

(86) Date de dépôt PCT/PCT Filing Date: 2010/09/20

(87) Date publication PCT/PCT Publication Date: 2011/03/24

(45) Date de délivrance/Issue Date: 2018/06/26

(85) Entrée phase nationale/National Entry: 2012/03/09

(86) N° demande PCT/PCT Application No.: US 2010/049471

(87) N° publication PCT/PCT Publication No.: 2011/035231

(30) **Priorité/Priority:** 2009/09/21 (US61/244,297)

(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01)

(72) Inventeurs/Inventors:

CHO, AESOP, US;
KIM, CHOUNG U., US;
METOBO, SAMUEL E., US;
RAY, ADRIAN S., US;
XU, JIE, US


(73) Propriétaire/Owner:

GILEAD SCIENCES, INC., US

(74) Agent: ROBIC

(54) Titre : ANALOGUES DE CARBANUCLEOSIDE 2'-FLUORO-SUBSTITUES POUR TRAITEMENT ANTIVIRAL

(54) Title: 2'-FLUORO SUBSTITUTED CARBA-NUCLEOSIDE ANALOGS FOR ANTIVIRAL TREATMENT



(57) **Abrégé/Abstract:**

Provided are pyrrolo[1,2-f][1,2,4]triazinyl, imidazo[1,5-f][1,2,4]triazinyl, imidazo(1,2-f)[1,2,4]triazinyl, and [1,2,4]triazolo[4,3-f][1,2,4]triazinyl nucleosides, nucleoside phosphates and prodrugs thereof, wherein the 2' position of the nucleoside sugar is substituted with halogen and carbon substituents. The compounds, compositions, and methods provided are useful for the treatment of Flaviviridae virus infections, particularly hepatitis C infections caused by both wild type and mutant strains of HCV.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
24 March 2011 (24.03.2011)

PCT

(10) International Publication Number
WO 2011/035231 A1(51) International Patent Classification:
C07D 487/04 (2006.01)(21) International Application Number:
PCT/US2010/049471(22) International Filing Date:
20 September 2010 (20.09.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/244,297 21 September 2009 (21.09.2009) US(71) Applicant (for all designated States except US):
GILEAD SCIENCES, INC. [US/US]; 333 Lakeside Drive, Foster City, CA 94404 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHO, Aesop** [US/US]; 1656 Notre Dame Drive, Mountain View, CA 94040 (US). **KIM, Choung, U.** [US/US]; 1750 Elizabeth Street, San Carlos, CA 94070 (US). **METOBO, Samuel, E.** [KE/US]; 35329 Severn Drive, Newark, CA 94560 (US). **RAY, Adrian, S.** [US/US]; 19 Pilot Circle, Redwood City, CA 94065 (US). **XU, Jie** [US/US]; 812 Rigel Lane, Foster City, CA 94404 (US).(74) Agents: **WARD, John** et al.; Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, CA 94404 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: 2' -FLUORO SUBSTITUTED CARBA-NUCLEOSIDE ANALOGS FOR ANTIVIRAL TREATMENT



(57) Abstract: Provided are pyrrolo[1,2-f][1,2,4]triazinyl, imidazo[1,5-f][1,2,4]triazinyl, imidazo(1,2-f)[1,2,4]triazinyl, and [1,2,4]triazolo[4,3-f][1,2,4]triazinyl nucleosides, nucleoside phosphates and prodrugs thereof, wherein the 2' position of the nucleoside sugar is substituted with halogen and carbon substituents. The compounds, compositions, and methods provided are useful for the treatment of Flaviviridae virus infections, particularly hepatitis C infections caused by both wild type and mutant strains of HCV.

WO 2011/035231 A1

5

2'-FLUORO SUBSTITUTED CARBA-NUCLEOSIDE ANALOGS FOR
ANTIVIRAL TREATMENT

FIELD OF THE INVENTION

10 The invention relates generally to compounds with antiviral activity, more particularly nucleosides active against *Flaviviridae* infections and most particularly to inhibitors of hepatitis C virus RNA-dependent RNA polymerase.

BACKGROUND OF THE INVENTION

Viruses comprising the *Flaviviridae* family comprise at least three
15 distinguishable genera including *pestiviruses*, *flaviviruses*, and *hepaciviruses* (Calisher, *et al.*, J. Gen. Virol., 1993, 70, 37-43). While *pestiviruses* cause many economically important animal diseases such as bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, hog cholera) and border disease of sheep (BDV), their importance in human disease is less well characterized (Moennig, V., *et al.*, Adv.
20 Vir. Res. 1992, 48, 53-98). *Flaviviruses* are responsible for important human diseases such as dengue fever and yellow fever while *hepaciviruses* cause hepatitis C virus infections in humans. Other important viral infections caused by the *Flaviviridae* family include West Nile virus (WNV) Japanese encephalitis virus (JEV), tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis, St Louis encephalitis,
25 Omsk hemorrhagic fever virus and Zika virus. Combined, infections from the *Flaviviridae* virus family cause significant mortality, morbidity and economic losses throughout the world. Therefore, there is a need to develop effective treatments for *Flaviviridae* virus infections.

The hepatitis C virus (HCV) is the leading cause of chronic liver disease
30 worldwide (Boyer, N. *et al.* J Hepatol. 32:98-112, 2000) so a significant focus of current antiviral research is directed toward the development of improved methods of treatment of chronic HCV infections in humans (Di Besceglie, A.M. and Bacon, B.

- 5 R., Scientific American, Oct.: 80-85, (1999); Gordon, C. P., et al., *J. Med. Chem.* **2005**, 48, 1-20; Maradpour, D.; et al., *Nat. Rev. Micro.* **2007**, 5(6), 453-463). A number of HCV treatments are reviewed by Bymock et al. in *Antiviral Chemistry & Chemotherapy*, 11:2; 79-95 (2000).

10 RNA-dependent RNA polymerase (RdRp) is one of the best studied targets for the development of novel HCV therapeutic agents. The NS5B polymerase is a target for inhibitors in early human clinical trials (Sommadossi, J., WO 01/90121 A2, US 2004/0006002 A1). These enzymes have been extensively characterized at the biochemical and structural level, with screening assays for identifying selective inhibitors (De Clercq, E. (2001) *J. Pharmacol. Exp. Ther.* 297:1-10; De Clercq, E. 15 (2001) *J. Clin. Virol.* 22:73-89). Biochemical targets such as NS5B are important in developing HCV therapies since HCV does not replicate in the laboratory and there are difficulties in developing cell-based assays and preclinical animal systems.

Currently, there are primarily two antiviral compounds, ribavirin, a nucleoside analog, and interferon-alpha (α) (IFN), that are used for the treatment of chronic HCV 20 infections in humans. Ribavirin alone is not effective in reducing viral RNA levels, has significant toxicity, and is known to induce anemia. The combination of IFN and ribavirin has been reported to be effective in the management of chronic hepatitis C (Scott, L. J., et al. *Drugs* **2002**, 62, 507-556) but less than half the patients infected with some genotypes show a persistent benefit when given this treatment. Other 25 patent applications disclosing the use of nucleoside analogs to treat hepatitis C virus include WO 01/32153, WO 01/60315, WO 02/057425, WO 02/057287, WO 02/032920, WO 02/18404, WO 04/046331, WO2008/089105 and WO2008/141079 but additional treatments for HCV infections have not yet become available for patients.

30 Virologic cures of patients with chronic HCV infection are difficult to achieve because of the prodigious amount of daily virus production in chronically infected patients and the high spontaneous mutability of HCV virus (Neumann, et al., *Science* **1998**, 282, 103-7; Fukimoto, et al., *Hepatology*, **1996**, 24, 1351-4; Domingo, et al., *Gene*, **1985**, 40, 1-8; Martell, et al., *J. Virol.* **1992**, 66, 3225-9. Experimental anti- 35 viral nucleoside analogs have been shown to induce viable mutations in the HCV

5 virus both *in vivo* and *in vitro* (Migliaccio, et al., *J. Biol. Chem.* **2003**, 926; Carroll, et al.,
Antimicrobial Agents Chemotherapy **2009**, 926; Brown, A. B., *Expert Opin. Investig. Drugs*
2009, 18, 709-725). Therefore, drugs having improved antiviral properties, particularly
enhanced activity against resistant strains of virus; improved oral bioavailability; fewer
undesirable side effects and extended effective half-life *in vivo* (De Francesco, R. et al. (2003)
10 Antiviral Research 58:1-16) are urgently needed.

Certain ribosides of the nucleobases pyrrolo[1,2-f][1,2,4]triazine, imidazo[1,5-f][1,2,4]triazine, imidazo[1,2-f][1,2,4]triazine, and [1,2,4]triazolo[4,3-f][1,2,4]triazine have
been disclosed in *Carbohydrate Research* 2001, 331(1), 77-82; *Nucleosides & Nucleotides*
(1996), 15(1-3), 793-807; *Tetrahedron Letters* (1994), 35(30), 5339-42; *Heterocycles* (1992),
15 34(3), 569-74; *J. Chem. Soc. Perkin Trans. 1* 1985, 3, 621-30; *J. Chem. Soc. Perkin Trans. 1*
1984, 2, 229-38; WO 2000056734; *Organic Letters* (2001), 3(6), 839-842; *J. Chem. Soc.*
Perkin Trans. 1 1999, 20, 2929-2936; and *J. Med. Chem.* 1986, 29(11), 2231-5. However,
these compounds have not been disclosed as useful for the treatment of HCV.


Ribosides of pyrrolo[1,2-f][1,2,4]triazinyl, imidazo[1,5-f][1,2,4]triazinyl, imidazo[1,2-f][1,2,4]triazinyl, and [1,2,4]triazolo[4,3-f][1,2,4]triazinyl nucleobases with antiviral, anti-
HCV, and anti-RdRp activity have been disclosed by Babu, Y. S., WO2008/089105 and
WO2008/141079; Cho, et al., WO2009/132123 and Francom, et al. WO2010/002877. Butler,
et al., WO2009/132135, has disclosed anti-viral pyrrolo[1,2-f][1,2,4]triazinyl, imidazo[1,5-f][1,2,4]triazinyl, imidazo[1,2-f][1,2,4]triazinyl, and [1,2,4]triazolo[4,3-f][1,2,4]triazinyl
25 nucleosides wherein the 1' position of the nucleoside sugar is substituted.

SUMMARY OF THE INVENTION

Provided are compounds that inhibit viruses of the *Flaviviridae* family. The
30 invention also comprises compounds of Formula I that inhibit viral nucleic acid
polymerases, particularly HCV RNA-dependent RNA polymerase (RdRp), rather than
cellular nucleic acid polymerases. The compounds of Formula I have been
discovered to be efficacious against both wild type and S282T mutant strains of HCV

5 virus. Therefore, a compound of Formula I is useful for treating *Flaviviridae* infections in humans and other animals.

In one embodiment, provided is a compound of Formula I:



Formula I

10 or a pharmaceutically acceptable salt or ester thereof;

wherein:

R¹ is (C₁-C₈)alkyl, (C₄-C₈)carbocyclalkyl, (C₁-C₈)substituted alkyl, (C₂-C₈)alkenyl, (C₂-C₈)substituted alkenyl, (C₂-C₈)alkynyl, (C₂-C₈)substituted alkynyl, or aryl(C₁-C₈)alkyl;

15 R² is halogen;

each R³, R⁴, and R⁵ is independently H, OR^a, N(R^a)₂, N₃, CN, NO₂, S(O)_nR^a, halogen, (C₁-C₈)alkyl, (C₄-C₈)carbocyclalkyl, (C₁-C₈)substituted alkyl, (C₂-C₈)alkenyl, (C₂-C₈)substituted alkenyl, (C₂-C₈)alkynyl, (C₂-C₈)substituted alkynyl, or aryl(C₁-C₈)alkyl;

or any two of R³, R⁴ or R⁵ on adjacent carbon atoms when taken together are
20 -O(CO)O- or when taken together with the ring carbon atoms to which they are attached form a double bond;


R⁶ is H, OR^a, N(R^a)₂, N₃, CN, NO₂, S(O)_nR^a, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹², -C(=O)SR¹¹, -S(O)R¹¹, -S(O)₂R¹¹, -S(O)(OR¹¹), -S(O)₂(OR¹¹), -SO₂NR¹¹R¹², halogen, (C₁-C₈)alkyl, (C₄-C₈)carbocyclalkyl, (C₁-C₈)substituted alkyl,

5 (C₂-C₈)alkenyl, (C₂-C₈)substituted alkenyl, (C₂-C₈)alkynyl, (C₂-C₈)substituted alkynyl, or aryl(C₁-C₈)alkyl;

each n is independently 0, 1, or 2;

each R^a is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, (C₄-C₈)carbocyclalkyl, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹², -C(=O)SR¹¹, -
 10 S(O)R¹¹, -S(O)₂R¹¹, -S(O)(OR¹¹), -S(O)₂(OR¹¹), or -SO₂NR¹¹R¹²;


R⁷ is H, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹², -C(=O)SR¹¹, -S(O)R¹¹, -S(O)₂R¹¹, -S(O)(OR¹¹), -S(O)₂(OR¹¹), -SO₂NR¹¹R¹², or



each Y and Y¹ is, independently, O, S, NR, ⁺N(O)(R), N(OR), ⁺N(O)(OR), or

15 N-NR₂;

W¹ and W², when taken together, are -Y³(C(R^y)₂)₃Y³-; or one of W¹ or W² together with either R³ or R⁴ is -Y³- and the other of W¹ or W² is Formula Ia; or W¹ and W² are each, independently, a group of the Formula Ia:



Formula Ia

20


wherein:

each Y² is independently a bond, O, CR₂, NR, ⁺N(O)(R), N(OR), ⁺N(O)(OR), N-NR₂, S, S-S, S(O), or S(O)₂;

each Y³ is independently O, S, or NR;

5 M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



wherein:

each M1a, M1c, and M1d is independently 0 or 1;

10 M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, R, $-\text{C}(=\text{Y}^1)\text{R}$, $-\text{C}(=\text{Y}^1)\text{OR}$, $-\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$, $-\text{N}(\text{R})_2$, $-\text{N}^+(\text{R})_3$, $-\text{SR}$, $-\text{S}(\text{O})\text{R}$, $-\text{S}(\text{O})_2\text{R}$, $-\text{S}(\text{O})(\text{OR})$, $-\text{S}(\text{O})_2(\text{OR})$, $-\text{OC}(=\text{Y}^1)\text{R}$, $-\text{OC}(=\text{Y}^1)\text{OR}$, $-\text{OC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{SC}(=\text{Y}^1)\text{R}$, $-\text{SC}(=\text{Y}^1)\text{OR}$, $-\text{SC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{R}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{OR}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$, $-\text{SO}_2\text{NR}_2$, $-\text{CN}$, $-\text{N}_3$, $-\text{NO}_2$, $-\text{OR}$, or W^3 ; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;


each R is independently H, $(\text{C}_1\text{--}\text{C}_8)$ alkyl, $(\text{C}_1\text{--}\text{C}_8)$ substituted alkyl, $(\text{C}_2\text{--}\text{C}_8)$ alkenyl, $(\text{C}_2\text{--}\text{C}_8)$ substituted alkenyl, $(\text{C}_2\text{--}\text{C}_8)$ alkynyl, $(\text{C}_2\text{--}\text{C}_8)$ substituted alkynyl, $\text{C}_6\text{--}\text{C}_{20}$ aryl, $\text{C}_6\text{--}\text{C}_{20}$ substituted aryl, $\text{C}_2\text{--}\text{C}_{20}$ heterocyclyl, $\text{C}_2\text{--}\text{C}_{20}$ substituted heterocyclyl, arylalkyl or substituted arylalkyl;

W^3 is W^4 or W^5 ; W^4 is R, $-\text{C}(\text{Y}^1)\text{R}^y$, $-\text{C}(\text{Y}^1)\text{W}^5$, $-\text{SO}_2\text{R}^y$, or $-\text{SO}_2\text{W}^5$; and W^5 is a carbocycle or a heterocycle wherein W^5 is independently substituted with 0 to 3 R^y groups;

each X^1 and X^2 is independently C--R^{10} or N;

each R^8 is halogen, $\text{NR}^{11}\text{R}^{12}$, $\text{N}(\text{R}^{11})\text{OR}^{11}$, $\text{NR}^{11}\text{NR}^{11}\text{R}^{12}$, N_3 , NO, NO_2 , CHO, CN, $-\text{CH}(=\text{NR}^{11})$, $-\text{CH}=\text{NNHR}^{11}$, $-\text{CH}=\text{N}(\text{OR}^{11})$, $-\text{CH}(\text{OR}^{11})_2$, $-\text{C}(=\text{O})\text{NR}^{11}\text{R}^{12}$, $-\text{C}(=\text{S})\text{NR}^{11}\text{R}^{12}$, $-\text{C}(=\text{O})\text{OR}^{11}$, $(\text{C}_1\text{--}\text{C}_8)$ alkyl, $(\text{C}_2\text{--}\text{C}_8)$ alkenyl, $(\text{C}_2\text{--}\text{C}_8)$ alkynyl, $(\text{C}_4\text{--}\text{C}_8)$ carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-\text{C}(=\text{O})(\text{C}_1\text{--}\text{C}_8)$ alkyl, $-\text{S}(\text{O})_n(\text{C}_1\text{--}\text{C}_8)$ alkyl, aryl $(\text{C}_1\text{--}\text{C}_8)$ alkyl, OR^{11} or SR^{11} ;


- 5 each R^9 and R^{10} is independently H, halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO, NO_2 , CHO, CN, $-CH(=NR^{11})$, $-CH=NHNR^{11}$, $-CH=N(OR^{11})$, $-CH(OR^{11})_2$, $-C(=O)NR^{11}R^{12}$, $-C(=S)NR^{11}R^{12}$, $-C(=O)OR^{11}$, R^{11} , OR^{11} or SR^{11} ;
- each R^{11} and R^{12} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl,
- 10 $-C(=O)(C_1-C_8)$ alkyl, $-S(O)_n(C_1-C_8)$ alkyl or aryl (C_1-C_8) alkyl; or R^{11} and R^{12} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or $-NR^a$;
- wherein each (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl or aryl (C_1-C_8) alkyl of
- 15 each R^1 , R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is, independently, optionally substituted with one or more halo, hydroxy, CN, N_3 , $N(R^a)_2$ or OR^a ; and wherein one or more of the non-terminal carbon atoms of each said (C_1-C_8) alkyl may be optionally replaced with -O-, -S- or $-NR^a$; wherein each (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl of each R^1 , R^6 , R^{11} , R^{12} , R^a and R is a hydrocarbon containing normal, secondary or tertiary carbon atoms.
- 20 In one embodiment, provided is a compound of Formula I:



Formula I

- or a pharmaceutically acceptable salt or ester thereof;
- wherein:
- 25 R^1 is (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;


- 5 R^2 is F;
each R^3 and R^5 is independently H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl(C_1-C_8)alkyl;
 R^4 is H, OR^b , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, halogen, (C_1-C_8) alkyl,
10 (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl(C_1-C_8)alkyl;
or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are $-O(CO)O-$ or when taken together with the ring carbon atoms to which they are attached form a double bond;
15 R^6 is H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl(C_1-C_8)alkyl;
20 each n is independently 0, 1, or 2;
each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;
each R^b is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;
25 R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



7a

5 each Y and Y¹ is, independently, O, S, NR, ⁺N(O)(R), N(OR), ⁺N(O)(OR), or N-NR₂;

W¹ and W², when taken together, are -Y³(C(R^y)₂)₃Y³-; or one of W¹ or W² together with either R³ or R⁴ is -Y³- and the other of W¹ or W² is Formula Ia; or W¹ and W² are each, independently, a group of the Formula Ia:



Formula Ia


wherein:

each Y² is independently a bond, O, CR₂, NR, ⁺N(O)(R), N(OR), ⁺N(O)(OR), N-NR₂, S, S-S, S(O), or S(O)₂;

15 each Y³ is independently O, S, or NR;

M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



wherein:

20 each M1a, M1c, and M1d is independently 0 or 1;


M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, R, -C(=Y¹)R, -C(=Y¹)OR, -C(=Y¹)N(R)₂, -N(R)₂, -⁺N(R)₃, -SR, -S(O)R, -S(O)₂R, -S(O)(OR), -S(O)₂(OR), -OC(=Y¹)R, -OC(=Y¹)OR, -OC(=Y¹)(N(R)₂), -SC(=Y¹)R, -SC(=Y¹)OR, -SC(=Y¹)(N(R)₂), -N(R)C(=Y¹)R,

- 5 -N(R)C(=Y¹)OR, -N(R)C(=Y¹)N(R)₂, -SO₂NR₂, -CN, -N₃, -NO₂, -OR, or W³; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;
- each R is independently H, (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl, (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl, (C₂-C₈) substituted alkynyl, C₆-C₂₀ aryl,
- 10 C₆-C₂₀ substituted aryl, C₂-C₂₀ heterocyclyl, C₂-C₂₀ substituted heterocyclyl, arylalkyl or substituted arylalkyl;
- W³ is W⁴ or W⁵; W⁴ is R, -C(Y¹)R^y, -C(Y¹)W⁵, -SO₂R^y, or -SO₂W⁵; and W⁵ is a carbocycle or a heterocycle wherein W⁵ is independently substituted with 0 to 3 R^y groups;
- each X¹ and X² is independently C-R¹⁰ or N;
- 15 each R⁸ is halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NNHR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl, aryl(C₁-C₈)alkyl, OR¹¹ or SR¹¹;
- 20 each R⁹ and R¹⁰ is independently H, halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NNHR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, R¹¹, OR¹¹ or SR¹¹;
- each R¹¹ and R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl,
- 25 -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl or aryl(C₁-C₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or -NR^a-;
- wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl or aryl(C₁-C₈)alkyl of
- 30 each R¹, R³, R⁴, R⁵, R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more halo, hydroxy, CN, N₃, N(R^a)₂ or OR^a; and wherein one or more of the non-terminal carbon atoms of each said (C₁-C₈)alkyl is optionally replaced with -O-, -S- or -NR^a-; wherein each

- 5 (C₁-C₈)alkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl of each R¹, R⁶, R¹¹, R¹², R^a and R is a hydrocarbon containing normal, secondary or tertiary carbon atoms.

In one embodiment, there is provided a compound of Formula II:




Formula II


- 10 or a pharmaceutically acceptable salt thereof;

wherein:

- R¹ is (C₁-C₈)alkyl;
 R³ and R⁵ are each H;
 R⁴ is OR^a;
 15 R⁶ is H, OR^a, CN or (C₁-C₈)alkyl;
 each R^a is independently H or (C₁-C₈)alkyl;
 R⁷ is H or



- each Y and Y¹ is O;
 20 one of W¹ or W² together with either R³ or R⁴ is -Y³- and the other of W¹ or W² is Formula Ia; or W¹ and W² are each, independently, a group of the Formula Ia:



Formula Ia


wherein:

each Y^2 is independently O or NR;

each Y^3 is O;

M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



wherein:

each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, R, $-\text{C}(=\text{Y}^1)\text{OR}$, $-\text{OC}(=\text{Y}^1)\text{OR}$, or $-\text{SC}(=\text{Y}^1)\text{R}$; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, (C₁-C₈)alkyl, (C₁-C₈) substituted alkyl, C₆-C₂₀ aryl,

C₆-C₂₀ substituted aryl, C₂-C₂₀ heterocyclyl, or arylalkyl;

X^1 is C-R¹⁰ or N;


X^2 is C-R¹⁰;


R^8 is halogen, NR¹¹R¹², or OR¹¹;

R^9 is H, NR¹¹R¹², or SR¹¹;


- 5 each R^{10} is H; and
- each R^{11} and R^{12} is independently H, or (C₁-C₈)alkyl, wherein each (C₁-C₈)alkyl of each R^1 , R^6 , R^{11} , R^{12} , R^a and R is a hydrocarbon containing normal, secondary or tertiary carbon atoms.

In another embodiment, there is provided a compound that is:






5




7h

5



5




10 or a pharmaceutically acceptable salt thereof.


7k

5

In another embodiment, there is provided a compound that is:




5




7m






5




7n




5




10






or a pharmaceutically acceptable salt thereof.

10 In another embodiment, there is provided a compound useful for the preparation of a compound of Formula I that is:



7q



In another embodiment, provided are compounds of Formula I and pharmaceutically acceptable salts thereof and all racemates, enantiomers, diastereomers, tautomers, polymorphs, pseudopolymorphs and amorphous forms thereof.


In another embodiment, provided are novel compounds of Formula I with activity against infectious *Flaviviridae* viruses. Without wishing to be bound by theory, the compounds of the invention may inhibit viral RNA-dependent RNA polymerase and thus inhibit the replication of the virus. They are useful for treating human patients infected with a human virus such as hepatitis C.

In another embodiment, provided are pharmaceutical compositions comprising an effective amount of a Formula I compound, or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable diluent or carrier.

5 In another embodiment, the present application provides for combination
pharmaceutical agent comprising:

 a) a first pharmaceutical composition comprising a compound of Formula I; or a
pharmaceutically acceptable salt, solvate, or ester thereof; and

 b) a second pharmaceutical composition comprising at least one additional
10 therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs,
HCV NS3 protease inhibitors, NS5a inhibitors, alpha-



5 glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers
10 and other drugs for treating HCV; or mixtures thereof.

In another embodiment, the present application provides for a method of inhibiting HCV polymerase, comprising contacting a cell infected with HCV with an effective amount of a compound of Formula I; or a pharmaceutically acceptable salts, solvate, and/or ester thereof.

15 In another embodiment, the present application provides for a method of inhibiting HCV polymerase, comprising contacting a cell infected with HCV with an effective amount of a compound of Formula I; or a pharmaceutically acceptable salts, solvate, and/or ester thereof; and at least one additional therapeutic agent.

In another embodiment, the present application provides for a method of
20 treating and/or preventing a disease caused by a viral infection wherein the viral infection is caused by a virus selected from the group consisting of dengue virus, yellow fever virus, West Nile virus, Japanese encephalitis virus, tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis virus, St Louis encephalitis virus, Omsk hemorrhagic fever virus, bovine viral diarrhea virus, Zika
25 virus and Hepatitis C virus; by administering to a subject in need thereof a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

In another embodiment, the present application provides for a method of treating HCV in a patient, comprising administering to said patient a therapeutically
30 effective amount of a compound of Formula I; or a pharmaceutically acceptable salt, solvate, and/or ester thereof.

In another embodiment, the present application provides for a method of treating HCV in a patient, comprising administering to said patient a therapeutically effective amount of a compound of Formula I; or a pharmaceutically acceptable salt,
35 solvate, and/or ester thereof; and at least one additional therapeutic agent.

5

Another aspect of the invention provides a method for the treatment or prevention of the symptoms or effects of an HCV infection in an infected animal which comprises administering to, i.e. treating, said animal with a pharmaceutical combination composition or formulation comprising an effective amount of a Formula I compound, and a second
10 compound having anti-HCV properties.

In another aspect, the invention also provides a method of inhibiting HCV, comprising administering to a mammal infected with HCV an amount of a Formula I compound, effective to inhibit the replication of HCV in infected cells in said mammal.

In another aspect, there is provided the use of therapeutically amount of a compound
15 or a pharmaceutically acceptable salt or ester thereof as defined herein or of a pharmaceutical composition as defined herein for treating or preventing a *Flaviviridae* virus infection in a mammal in need thereof.

In another aspect, there is provided the use of therapeutically amount of a compound or a pharmaceutically acceptable salt or ester thereof as defined herein or of a
20 pharmaceutical composition as defined herein for treating a *Flaviviridae* virus infection in a mammal in need thereof.

In another aspect, there is provided a compound or a pharmaceutically acceptable salt or ester thereof as defined herein or a pharmaceutical composition as defined herein for the treatment or prevention of a *Flaviviridae* virus infection.

25 In another aspect, there is provided a compound or a pharmaceutically acceptable salt or ester thereof as defined herein or a pharmaceutical composition as defined herein for the treatment of a *Flaviviridae* virus infection.

In another aspect, there is provided the use of a compound or a pharmaceutically acceptable salt or ester thereof as defined herein for the manufacture of a medicament for the
30 treatment or prevention of a *Flaviviridae* virus infection, Hepatitis C virus infection or S282T mutant of Hepatitis C virus infection.

5 In another aspect, there is provided the use of a compound or a pharmaceutically acceptable salt or ester thereof as defined herein for the manufacture of a medicament for the treatment of a *Flaviviridae* virus infection, Hepatitis C virus infection or S282T mutant of Hepatitis C virus infection.

10 In another aspect, there is provided the use of a therapeutically effective amount of the compound as defined herein or a pharmaceutically acceptable salt thereof, for treating or preventing a dengue virus infection in a mammal in need thereof.

 In another aspect, there is provided the use of a therapeutically effective amount of the compound as defined herein or a pharmaceutically acceptable salt thereof, for treating a dengue virus infection in a mammal in need thereof.

15 In another aspect, there is provided the use of a compound as defined herein or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a dengue virus infection.

 In another aspect, there is provided the use of a compound as defined herein or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a dengue virus infection.

 In another aspect, there is provided the use of a pharmaceutical composition as defined herein, for treating or preventing a dengue virus infection.

 In another aspect, there is provided the use of a pharmaceutical composition as defined herein, for treating a dengue virus infection.

25 In another aspect, there is provided the use of a compound or a pharmaceutically acceptable salt or ester thereof as defined herein for inhibiting HCV polymerase.

 In another aspect, provided is the use of a compound of Formula I for the manufacture of a medicament for the treatment of *Flaviviridae* viral infections. In another aspect, provided is a compound of Formula I for use in treating a *Flaviviridae* viral infection.

30 In one embodiment, the *Flaviviridae* viral infection is acute or chronic HCV infection. In one embodiment of each aspect of use and compound, the treatment results in the reduction of one or more of the viral loads or clearance of RNA in the patient.

5 In another aspect, the invention also provides processes and novel intermediates disclosed herein which are useful for preparing Formula I compounds of the invention.


 In other aspects, novel methods for synthesis, analysis, separation, isolation, purification, characterization, and testing of the compounds of this invention are provided.

DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

10


 Reference will now be made in detail to certain embodiments of the invention, examples of which are illustrated in the accompanying description, structures and formulas. While the invention will be described in conjunction with the enumerated embodiments, it will be understood that they are not intended to limit the invention to those embodiments.

15 On the contrary, the invention is intended to cover all



- 5 alternatives, modifications, and equivalents, which may be included within the scope of the present invention.

In another aspect, compounds of Formula I are represented by Formula II:



Formula II


- 10 or a pharmaceutically acceptable salt, thereof;

wherein:

- R^1 is (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;
- 15 each R^3 , R^4 , or R^5 is independently H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;
- or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are —
- 20 $O(CO)O-$ or when taken together with the ring carbon atoms to which they are attached form a double bond;
- R^6 is H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted
- 25 alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;
- each n is independently 0, 1, or 2;

5 each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;


R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



10

each Y or Y^1 is, independently, O, S, NR, $^+N(O)(R)$, N(OR), $^+N(O)(OR)$, or N-NR₂;

W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or W^1 and W^2 are each, independently, a group of the Formula Ia:



Formula Ia

wherein:


each Y^2 is independently a bond, O, CR₂, NR, $^+N(O)(R)$, N(OR), $^-N(O)(OR)$, N-NR₂, S, S-S, S(O), or S(O)₂;

20

each Y^3 is independently O, S, or NR;

M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



5

wherein:

each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, R, $-C(=Y^1)R$, $-C(=Y^1)OR$, $-C(=Y^1)N(R)_2$, $-N(R)_2$, $-N(R)_3$, $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-OC(=Y^1)R$, $-OC(=Y^1)OR$, $-OC(=Y^1)N(R)_2$, $-SC(=Y^1)R$, $-SC(=Y^1)OR$, $-SC(=Y^1)N(R)_2$, $-N(R)C(=Y^1)R$, $-N(R)C(=Y^1)OR$, $-N(R)C(=Y^1)N(R)_2$, $-SO_2NR_2$, $-CN$, $-N_3$, $-NO_2$, $-OR$, or W^3 ; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, (C_1-C_8) alkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, C_6-C_{20} aryl, C_6-C_{20} substituted aryl, C_2-C_{20} heterocyclyl, C_2-C_{20} substituted heterocyclyl, arylalkyl or substituted arylalkyl;

W^3 is W^4 or W^5 ; W^4 is R, $-C(Y^1)R^y$, $-C(Y^1)W^5$, $-SO_2R^y$, or $-SO_2W^5$; and W^5 is a carbocycle or a heterocycle wherein W^5 is independently substituted with 0 to 3 R^y groups;

each X^1 or X^2 is independently $C-R^{10}$ or N;

each R^8 is halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO, NO_2 , CHO, CN, $-CH(=NR^{11})$, $-CH=NNHR^{11}$, $-CH=N(OR^{11})$, $-CH(OR^{11})_2$, $-C(=O)NR^{11}R^{12}$, $-C(=S)NR^{11}R^{12}$, $-C(=O)OR^{11}$, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-C(=O)(C_1-C_8)$ alkyl, $-S(O)_n(C_1-C_8)$ alkyl, aryl (C_1-C_8) alkyl, OR^{11} or SR^{11} ;

each R^9 or R^{10} is independently H, halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, N_3 , NO, NO_2 , CHO, CN, $-CH(=NR^{11})$, $-CH=NNHR^{11}$, $-CH=N(OR^{11})$, $-CH(OR^{11})_2$, $-C(=O)NR^{11}R^{12}$, $-C(=S)NR^{11}R^{12}$, $-C(=O)OR^{11}$, R^{11} , OR^{11} or SR^{11} ;

5 each R^{11} or R^{12} is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl or aryl(C₁-C₈)alkyl; or R^{11} and R^{12} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring
10 can optionally be replaced with -O-, -S- or -NR^a-;

wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl or aryl(C₁-C₈)alkyl of each R^1 , R^3 , R^4 , R^5 , R^6 , R^{11} or R^{12} is, independently, optionally substituted with one or more halo, hydroxy, CN, N₃, N(R^a)₂ or OR^a; and wherein one or more of the non-terminal carbon atoms of each said (C₁-C₈)alkyl may be optionally replaced with -O-,
15 -S- or -NR^a-.

In one embodiment of the invention of Formula II, R^1 is (C₁-C₈)alkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl. In another aspect of this embodiment, R^1 is (C₁-C₈)alkyl. In another aspect of this embodiment, R^1 is methyl, CH₂F, or ethynyl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^1 is
20 (C₁-C₈)alkyl and R^6 is H. In another aspect of this embodiment, R^1 is (C₁-C₈)alkyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R^1 is (C₁-C₈)alkyl and R^6 is CN, OH, or CH₃.

In one embodiment of Formula II, R^3 is H, OR^a, N(R^a)₂, N₃, CN, SR^a, halogen, (C₁-C₈)alkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl. In one aspect of this embodiment,
25 R^3 is H. In another aspect of this embodiment, R^3 is H and R^1 is (C₁-C₈)alkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl. In another aspect of this embodiment, R^3 is H and R^1 is (C₁-C₈)alkyl. In another aspect of this embodiment, R^3 is H and R^1 is methyl, CH₂F, or ethynyl. In another aspect of this embodiment, R^3 is H and R^1 is methyl. In another aspect of this embodiment, R^3 is H, R^1 is (C₁-C₈)alkyl and at least one of X¹ or X² is N. In another aspect of this embodiment, R^3 is H, R^1 is methyl and at least
30 one of X¹ or X² is N. In another aspect of this embodiment, R^3 is H, R^1 is (C₁-C₈)alkyl and R^6 is CN, OH, or CH₃. In another aspect of this embodiment, R^3 is H, R^1 is methyl and R^6 is CN, OH, or CH₃. In another aspect of this embodiment, R^3 is H, R^1 is methyl and R^6 is H.

- 5 In one embodiment of Formula II, R^4 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.
- 25 In one embodiment of Formula II, R^5 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this

embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^5 is N_3 .


In another embodiment of Formula II, R^5 is H. In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

5 In another embodiment of Formula II, R^6 is H, CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.


20 In another embodiment of Formula II, R^6 is CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

5 aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 .

In one embodiment of Formula II, R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)SR^{11}$ or



. In a aspect of this embodiment, R^7 is H. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$ wherein R^{11} is (C_1-C_8) alkyl. In another aspect of this embodiment, R^7 is



. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment,

5 R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this

10 embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and

15 R^6 is H.

In one embodiment of Formula II, X^1 is N or $C-R^{10}$. In another aspect of this embodiment, X^1 is N. In another aspect of this embodiment, X^1 is $C-R^{10}$. In another aspect of this embodiment, X^2 is C-H. In another aspect of this embodiment, X^1 is N and X^2 is C-H. In another aspect of this embodiment, X^1 is $C-R^{10}$ and X^2 is CH. In

20 another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this

30 embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of

5 this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.


In another embodiment of Formula II, each R^8 is independently halogen,
 10 $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, OR^{11} or SR^{11} . In another aspect of this embodiment, R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$
 15 and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1
 20 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^8 is OH and R^9 is NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment R^6 is H. In another aspect of this
 25 embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this
 30 embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this
 35 embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In

5 another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is
 10 CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment of Formula II, each R^{10} is, independently, H, halogen, CN or optionally substituted heteroaryl. In another aspect of this embodiment, R^1 is
 15 methyl. In another aspect of this embodiment, R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another
 20 aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl, CH_2F or
 25 ethynyl. In another aspect of this embodiment, R^8 is OH and R^9 is NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6
 30 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is (C_1-C_8) alkyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In

5 another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is (C_1-C_8) alkyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment, compounds of Formula I or Formula II are represented by Formula III:



Formula III

or a pharmaceutically acceptable salt, thereof;

wherein:

R^1 is CH_3 , CH_2F , or ethynyl and all remaining variables are defined as for Formula I.

25 In one embodiment of Formula III, R^4 is H, OR^a , $N(R^a)_2$, N_3 , CN, SR^a , halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl. In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In


5 another aspect of this embodiment, R^4 is OR^a and R^1 is CH_3 , CH_2F , or ethynyl. In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

15 In another embodiment of Formula III, R^6 is H, CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.


In another embodiment of Formula III, R^6 is CN, OR^a or CH_3 . In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is

5 methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 .

In one embodiment of Formula III, R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)SR^{11}$ or



. In a aspect of this embodiment, R^7 is H. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$. In another aspect of this embodiment, R^7 is $-C(=O)R^{11}$ wherein R^{11} is (C_1-C_8) alkyl. In another aspect of this embodiment, R^7 is



. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In
 15 another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of
 20 this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

25

In one embodiment of Formula III, X^1 is N or $C-R^{10}$. In another aspect of this embodiment, X^1 is N. In another aspect of this embodiment, X^1 is $C-R^{10}$. In another aspect of this embodiment, X^2 is C-H. In another aspect of this embodiment, X^1 is N and X^2 is C-H. In another aspect of this embodiment, X^1 is $C-R^{10}$ and X^2 is CH. In

5 another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 .
15 In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.


In another embodiment of Formula III, each R^8 is independently halogen, $NR^{11}R^{12}$, $N(R^{11})OR^{11}$, $NR^{11}NR^{11}R^{12}$, OR^{11} or SR^{11} . In another aspect of this embodiment, R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl, CH_2F or ethynyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 is OH and R^9 is NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl, CH_2F , or ethynyl. In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is
35

5 methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In another embodiment of Formula III, each R^{10} is, independently, H, halogen, CN or optionally substituted heteroaryl. In another aspect of this embodiment, R^1 is methyl. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$. In another aspect of this embodiment, R^9 is H, halogen, or $NR^{11}R^{12}$ and R^1 is methyl. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen. In another aspect of this embodiment, R^8 is NH_2 and R^9 is H or halogen and R^1 is methyl. In another aspect of this embodiment, R^8 and R^9 are each NH_2 . In another aspect of this embodiment, R^8 and R^9 are each NH_2 and R^1 is methyl. In another aspect of this embodiment, R^8 is OH and R^9 is NH_2 . In another aspect of this embodiment, R^8 is OH, R^9 is NH_2 and R^1 is methyl. In another aspect of this embodiment R^6 is H. In another aspect of this embodiment R^6 is CN. In another aspect of this embodiment R^6 is OR^a . In another aspect of this embodiment R^6 is OH. In another aspect of this embodiment R^6 is CH_3 . In another aspect of this embodiment, R^4 is H or OR^a . In another aspect of this embodiment, R^4 is OR^a and R^1 is methyl. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OR^a , R^1 is methyl and R^6 is H. In another aspect of this embodiment, R^4 is OH and R^1 is methyl. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and at least one of X^1 or X^2 is N. In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is CN, OH, or CH_3 . In another aspect of this embodiment, R^4 is OH, R^1 is methyl and R^6 is H.

In one embodiment of Formulas I-III, R^{11} or R^{12} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_4-C_8) carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, $-C(=O)(C_1-C_8)$ alkyl, $-S(O)_n(C_1-$


- 5 C₈)alkyl or aryl(C₁-C₈)alkyl. In another embodiment, R¹¹ and R¹² taken together with a nitrogen to which they are both attached, form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or -NR^a-. Therefore, by way of example and not limitation, the moiety -NR¹¹R¹² can be represented by the heterocycles:




- In another embodiment of Formulas I-III, each R³, R⁴, R⁵, R⁶, R¹¹ or R¹² is, independently, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl or aryl(C₁-C₈)alkyl, wherein said (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl or aryl(C₁-C₈)alkyl are, independently, optionally substituted with one or more halo, hydroxy, CN, N₃, N(R^a)₂ or OR^a. Therefore, by way of example and not limitation, R³, R⁴, R⁵, R⁶, R¹¹ or R¹² could represent moieties such as -CH(NH₂)CH₃, -CH(OH)CH₂CH₃, -CH(NH₂)CH(CH₃)₂, -CH₂CF₃, -(CH₂)₂CH(N₃)CH₃, -(CH₂)₆NH₂ and the like.

- In another embodiment of Formula I-III, R³, R⁴, R⁵, R⁶, R¹¹ or R¹² is (C₁-C₈)alkyl wherein one or more of the non-terminal carbon atoms of each said (C₁-C₈)alkyl may be optionally replaced with -O-, -S- or -NR^a-. Therefore, by way of example and not limitation, R³, R⁴, R⁵, R⁶, R¹¹ or R¹² could represent moieties such as -CH₂OCH₃, -CH₂OCH₂CH₃, -CH₂OCH(CH₃)₂, -CH₂SCH₃, -(CH₂)₆OCH₃, -(CH₂)₆N(CH₃)₂ and the like.

- 25 In another embodiment, Formulas I-III is a compound selected from the group consisting of





5




Chemical structures of various nucleoside phosphonates and phosphates, including:

- Indinavir phosphonate (top left)
- Indinavir phosphonate (top right)
- Indinavir phosphonate (middle left)
- Indinavir phosphonate (middle right)
- Indinavir phosphonate (bottom left)
- Indinavir phosphonate (bottom right)
- Indinavir phosphonate (bottom left)
- Indinavir phosphonate (bottom right)











or a pharmaceutically acceptable salt or ester thereof.

10 In another embodiment, provided is a compound useful for the synthesis of the compounds of Formula I selected from the group consisting of





DEFINITIONS

10 Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings:

When trade names are used herein, applicants intend to independently include the tradename product and the active pharmaceutical ingredient(s) of the tradename product.

15 As used herein, "a compound of the invention" or "a compound of Formula I" means a compound of Formula I or a pharmaceutically acceptable salt, thereof. Similarly, with respect to isolatable intermediates, the phrase "a compound of Formula (number)" means a compound of that formula and pharmaceutically acceptable salts, thereof.

20 "Alkyl" is hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms. For example, an alkyl group can have 1 to 20 carbon atoms (*i.e.*, C₁-C₂₀ alkyl), 1 to 8 carbon atoms (*i.e.*, C₁-C₈ alkyl), or 1 to 6 carbon atoms (*i.e.*, C₁-C₆ alkyl). Examples of suitable alkyl groups include, but are not limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (n-Pr, n-propyl, -CH₂CH₂CH₃), 2-propyl (i-Pr, i-propyl, -CH(CH₃)CH₂CH₃), 1-butyl (n-Bu, n-butyl, -CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (i-Bu, i-

25

5 butyl, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, 2-butyl (s-Bu , s-butyl , $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 2-methyl-2-propyl (t-Bu , t-butyl , $-\text{C}(\text{CH}_3)_3$), 1-pentyl (n-pentyl , $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$), 3-pentyl ($-\text{CH}(\text{CH}_2\text{CH}_3)_2$), 2-methyl-2-butyl ($-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$), 3-methyl-2-butyl ($-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$), 3-methyl-1-butyl ($-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$), 2-methyl-1-butyl ($-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 1-hexyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-hexyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3-hexyl ($-\text{CH}(\text{CH}_2\text{CH}_3)(\text{CH}_2\text{CH}_2\text{CH}_3)$), 2-methyl-2-pentyl ($-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3-methyl-2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 4-methyl-2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$), 3-methyl-3-pentyl ($-\text{C}(\text{CH}_3)(\text{CH}_2\text{CH}_3)_2$), 2-methyl-3-pentyl ($-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)_2$), 2,3-dimethyl-2-butyl ($-\text{C}(\text{CH}_3)_2\text{CH}(\text{CH}_3)_2$), 3,3-dimethyl-2-butyl ($-\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)_3$), and octyl ($-(\text{CH}_2)_7\text{CH}_3$).

“Alkoxy” means a group having the formula $-\text{O-alkyl}$, in which an alkyl group, as defined above, is attached to the parent molecule via an oxygen atom. The alkyl portion of an alkoxy group can have 1 to 20 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{20}$ alkoxy), 1 to 12 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{12}$ alkoxy), or 1 to 6 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_6$ alkoxy).
 20 Examples of suitable alkoxy groups include, but are not limited to, methoxy ($-\text{O-CH}_3$ or $-\text{OMe}$), ethoxy ($-\text{OCH}_2\text{CH}_3$ or $-\text{OEt}$), t-butoxy ($-\text{O-C}(\text{CH}_3)_3$ or $-\text{OtBu}$) and the like.

“Haloalkyl” is an alkyl group, as defined above, in which one or more hydrogen atoms of the alkyl group is replaced with a halogen atom. The alkyl portion
 25 of a haloalkyl group can have 1 to 20 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{20}$ haloalkyl), 1 to 12 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_{12}$ haloalkyl), or 1 to 6 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_6$ alkyl). Examples of suitable haloalkyl groups include, but are not limited to, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CFH}_2$, $-\text{CH}_2\text{CF}_3$, and the like.

“Alkenyl” is a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, sp^2 double
 30 bond. For example, an alkenyl group can have 2 to 20 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_{20}$ alkenyl), 2 to 8 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_8$ alkenyl), or 2 to 6 carbon atoms (*i.e.*, $\text{C}_2\text{-C}_6$ alkenyl). Examples of suitable alkenyl groups include, but are not limited to, ethylene or vinyl ($-\text{CH}=\text{CH}_2$), allyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), cyclopentenyl ($-\text{C}_5\text{H}_7$), and 5-hexenyl
 35 ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$).

5 “Alkynyl” is a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp* triple bond. For example, an alkynyl group can have 2 to 20 carbon atoms (*i.e.*, C₂-C₂₀ alkynyl), 2 to 8 carbon atoms (*i.e.*, C₂-C₈ alkyne), or 2 to 6 carbon atoms (*i.e.*, C₂-C₆ alkynyl). Examples of suitable alkynyl groups include, but are not limited to,
10 acetylenic (-C≡CH), propargyl (-CH₂C≡CH), and the like.

“Alkylene” refers to a saturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. For example, an alkylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon
15 atoms. Typical alkylene radicals include, but are not limited to, methylene (-CH₂-), 1,1-ethyl (-CH(CH₃)-), 1,2-ethyl (-CH₂CH₂-), 1,1-propyl (-CH(CH₂CH₃)-), 1,2-propyl (-CH₂CH(CH₃)-), 1,3-propyl (-CH₂CH₂CH₂-), 1,4-butyl (-CH₂CH₂CH₂CH₂-), and the like.

“Alkenylene” refers to an unsaturated, branched or straight chain or cyclic
20 hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. For example, an alkenylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkenylene radicals include, but are not limited to, 1,2-ethylene (-CH=CH-).

25 “Alkynylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne. For example, an alkynylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkynylene radicals include, but are not limited to,
30 acetylene (-C≡C-), propargyl (-CH₂C≡C-), and 4-pentynyl (-CH₂CH₂CH₂C≡C-).

“Amino” refers generally to a nitrogen radical which can be considered a derivative of ammonia, having the formula -N(X)₂, where each “X” is independently H, substituted or unsubstituted alkyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, etc. The hybridization of the nitrogen is approximately sp³.
35 Nonlimiting types of amino include -NH₂, -N(alkyl)₂, -NH(alkyl), -N(carbocyclyl)₂, -

5 NH(carbocyclyl), -N(heterocyclyl)₂, -NH(heterocyclyl), -N(aryl)₂, -NH(aryl), -
 N(alkyl)(aryl), -N(alkyl)(heterocyclyl), -N(carbocyclyl)(heterocyclyl), -
 N(aryl)(heteroaryl), -N(alkyl)(heteroaryl), etc. The term “alkylamino” refers to an
 amino group substituted with at least one alkyl group. Nonlimiting examples of amino
 groups include -NH₂, -NH(CH₃), -N(CH₃)₂, -NH(CH₂CH₃), -N(CH₂CH₃)₂, -
 10 NH(phenyl), -N(phenyl)₂, -NH(benzyl), -N(benzyl)₂, etc. Substituted alkylamino refers
 generally to alkylamino groups, as defined above, in which at least one substituted alkyl,
 as defined herein, is attached to the amino nitrogen atom. Non-limiting examples of
 substituted alkylamino includes -NH(alkylene-C(O)-OH), -NH(alkylene-C(O)-O-alkyl),
 -N(alkylene-C(O)-OH)₂, -N(alkylene-C(O)-O-alkyl)₂, etc.

15 “Aryl” means an aromatic hydrocarbon radical derived by the removal of one
 hydrogen atom from a single carbon atom of a parent aromatic ring system. For
 example, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 10
 carbon atoms. Typical aryl groups include, but are not limited to, radicals derived from
 benzene (e.g., phenyl), substituted benzene, naphthalene, anthracene, biphenyl, and the
 20 like.

“Arylalkyl” refers to an acyclic alkyl radical in which one of the hydrogen
 atoms bonded to a carbon atom, typically a terminal or sp³ carbon atom, is replaced
 with an aryl radical. Typical arylalkyl groups include, but are not limited to, benzyl,
 2-phenylethan-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, naphthobenzyl,
 25 2-naphthophenylethan-1-yl and the like. The arylalkyl group can comprise 7 to 20
 carbon atoms, *e.g.*, the alkyl moiety is 1 to 6 carbon atoms and the aryl moiety is 6 to
 14 carbon atoms.

“Arylalkenyl” refers to an acyclic alkenyl radical in which one of the
 hydrogen atoms bonded to a carbon atom, typically a terminal or sp³ carbon atom, but
 30 also an sp² carbon atom, is replaced with an aryl radical. The aryl portion of the
 arylalkenyl can include, for example, any of the aryl groups disclosed herein, and the
 alkenyl portion of the arylalkenyl can include, for example, any of the alkenyl groups
 disclosed herein. The arylalkenyl group can comprise 8 to 20 carbon atoms, *e.g.*, the
 alkenyl moiety is 2 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

5 “Arylalkynyl” refers to an acyclic alkynyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also an sp carbon atom, is replaced with an aryl radical. The aryl portion of the arylalkynyl can include, for example, any of the aryl groups disclosed herein, and the alkynyl portion of the arylalkynyl can include, for example, any of the alkynyl groups disclosed herein. The arylalkynyl group can comprise 8 to 20 carbon atoms, *e.g.*, the
10 alkynyl moiety is 2 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

The term “substituted” in reference to alkyl, alkylene, aryl, arylalkyl, alkoxy, heterocyclyl, heteroaryl, carbocyclyl, etc., for example, “substituted alkyl”, “substituted alkylene”, “substituted aryl”, “substituted arylalkyl”, “substituted
15 heterocyclyl”, and “substituted carbocyclyl”, unless otherwise indicated, means alkyl, alkylene, aryl, arylalkyl, heterocyclyl, carbocyclyl respectively, in which one or more hydrogen atoms are each independently replaced with a non-hydrogen substituent. Typical substituents include, but are not limited to, $-X$, $-R^b$, $-O^-$, $=O$, $-OR^b$, $-SR^b$, $-S^-$, $-NR^b_2$, $-N^+R^b_3$, $=NR^b$, $-CX_3$, $-CN$, $-OCN$, $-SCN$, $-N=C=O$, $-NCS$, $-NO$, $-NO_2$, $=N_2$,
20 $-N_3$, $-NHC(=O)R^b$, $-OC(=O)R^b$, $-NHC(=O)NR^b_2$, $-S(=O)_2-$, $-S(=O)_2OH$, $-S(=O)_2R^b$, $-OS(=O)_2OR^b$, $-S(=O)_2NR^b_2$, $-S(=O)R^b$, $-OP(=O)(OR^b)_2$, $-P(=O)(OR^b)_2$, $-P(=O)(O^-)_2$, $-P(=O)(OH)_2$, $-P(O)(OR^b)(O^-)$, $-C(=O)R^b$, $-C(=O)X$, $-C(S)R^b$, $-C(O)OR^b$, $-C(O)O^-$, $-C(S)OR^b$, $-C(OSR^b)$, $-C(S)SR^b$, $-C(O)NR^b_2$, $-C(S)NR^b_2$, $-C(=NR^b)NR^b_2$, where each X is independently a halogen: F, Cl, Br, or I; and each R^b is independently H, alkyl,
25 aryl, arylalkyl, a heterocycle, or a protecting group or prodrug moiety. Alkylene, alkenylene, and alkynylene groups may also be similarly substituted. Unless otherwise indicated, when the term “substituted” is used in conjunction with groups such as arylalkyl, which have two or more moieties capable of substitution, the substituents can be attached to the aryl moiety, the alkyl moiety, or both.


30 The term “prodrug” as used herein refers to any compound that when administered to a biological system generates the drug substance, *i.e.*, active ingredient, as a result of spontaneous chemical reaction(s), enzyme catalyzed chemical reaction(s), photolysis, and/or metabolic chemical reaction(s). A prodrug is thus a covalently modified analog or latent form of a therapeutically active compound.

5 One skilled in the art will recognize that substituents and other moieties of the compounds of Formula I-III should be selected in order to provide a compound which is sufficiently stable to provide a pharmaceutically useful compound which can be formulated into an acceptably stable pharmaceutical composition. The definitions and substituents for various genus and subgenus of the present compounds are described and illustrated herein. It should be understood by one skilled in the art that any combination of the definitions and substituents described above should not result in an inoperable species or compound. "Inoperable species or compounds" means compound structures that violates relevant scientific principles (such as, for example, a carbon atom connecting to more than four covalent bonds) or compounds too unstable to permit isolation and formulation into pharmaceutically acceptable dosage forms.


"Heteroalkyl" refers to an alkyl group where one or more carbon atoms have been replaced with a heteroatom, such as, O, N, or S. For example, if the carbon atom of the alkyl group which is attached to the parent molecule is replaced with a heteroatom (e.g., O, N, or S) the resulting heteroalkyl groups are, respectively, an alkoxy group (e.g., -OCH₃, etc.), an amine (e.g., -NHCH₃, -N(CH₃)₂, etc.), or a thioalkyl group (e.g., -SCH₃). If a non-terminal carbon atom of the alkyl group which is not attached to the parent molecule is replaced with a heteroatom (e.g., O, N, or S) the resulting heteroalkyl groups are, respectively, an alkyl ether (e.g., -CH₂CH₂-O-CH₃, etc.), an alkyl amine (e.g., -CH₂NHCH₃, -CH₂N(CH₃)₂, etc.), or a thioalkyl ether (e.g., -CH₂-S-CH₃). If a terminal carbon atom of the alkyl group is replaced with a heteroatom (e.g., O, N, or S), the resulting heteroalkyl groups are, respectively, a hydroxyalkyl group (e.g., -CH₂CH₂-OH), an aminoalkyl group (e.g., -CH₂NH₂), or an alkyl thiol group (e.g., -CH₂CH₂-SH). A heteroalkyl group can have, for example, 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. A C₁-C₆ heteroalkyl group means a heteroalkyl group having 1 to 6 carbon atoms.

"Heterocycle" or "heterocyclyl" as used herein includes by way of example and not limitation those heterocycles described in Paquette, Leo A.; Principles of Modern Heterocyclic Chemistry (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; The Chemistry of Heterocyclic Compounds, A Series of

5 Monographs” (John Wiley & Sons, New York, 1950 to present), in particular
 Volumes 13, 14, 16, 19, and 28; and *J. Am. Chem. Soc.* (1960) 82:5566. In one
 specific embodiment of the invention “heterocycle” includes a “carbocycle” as
 defined herein, wherein one or more (*e.g.* 1, 2, 3, or 4) carbon atoms have been
 replaced with a heteroatom (*e.g.* O, N, or S). The terms “heterocycle” or
 10 “heterocyclyl” includes saturated rings, partially unsaturated rings, and aromatic rings
 (*i.e.*, heteroaromatic rings). Substituted heterocyclyls include, for example,
 heterocyclic rings substituted with any of the substituents disclosed herein including
 carbonyl groups. A non-limiting example of a carbonyl substituted heterocyclyl is:



15 Examples of heterocycles include by way of example and not limitation
 pyridyl, dihydropyridyl, tetrahydropyridyl (piperidyl), thiazolyl,
 tetrahydrothiophenyl, sulfur oxidized tetrahydrothiophenyl, pyrimidinyl, furanyl,
 thienyl, pyrrolyl, pyrazolyl, imidazolyl, tetrazolyl, benzofuranyl, thianaphthalenyl,
 indolyl, indolenyl, quinolinyl, isoquinolinyl, benzimidazolyl, piperidinyl, 4-
 20 piperidonyl, pyrrolidinyl, 2-pyrrolidonyl, pyrrolinyl, tetrahydrofuranyl,
 tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl,
 octahydroisoquinolinyl, azocinyl, triazinyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-
 dithiazinyl, thienyl, thianthrenyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl,
 phenoxathinyl, 2H-pyrrolyl, isothiazolyl, isoxazolyl, pyrazinyl, pyridazinyl,
 25 indolizinyl, isoindolyl, 3H-indolyl, 1H-indazolyl, purinyl, 4H-quinolizinyl,
 phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, 4aH-
 carbazolyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, pyrimidinyl,
 phenanthrolinyl, phenazinyl, phenothiazinyl, furazanyl, phenoxazinyl, isochromanyl,
 chromanyl, imidazolidinyl, imidazolyl, pyrazolidinyl, pyrazolyl, piperazinyl,
 30 indolinyl, isoindolinyl, quinuclidinyl, morpholinyl, oxazolidinyl, benzotriazolyl,
 benzisoxazolyl, oxindolyl, benzoxazolyl, isatinoyl, and bis-tetrahydrofuranyl:



5

By way of example and not limitation, carbon bonded heterocycles are bonded at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or isothiazole, position 2 or 3 of an aziridine, position 2, 3, or 4 of an azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline. Still more typically, carbon bonded heterocycles include 2-pyridyl, 3-pyridyl, 4-pyridyl, 5-pyridyl, 6-pyridyl, 3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl, 6-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 2-pyrazinyl, 3-pyrazinyl, 5-pyrazinyl, 6-pyrazinyl, 2-thiazolyl, 4-thiazolyl, or 5-thiazolyl.

By way of example and not limitation, nitrogen bonded heterocycles are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole, imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3-pyrazoline, piperidine, piperazine, indole, indoline, 1H-indazole, position 2 of a isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline. Still more typically, nitrogen bonded heterocycles include 1-aziridyl, 1-azetidyl, 1-pyrrolyl, 1-imidazolyl, 1-pyrazolyl, and 1-piperidinyl.

“Heterocyclalkyl” refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a heterocyclalkyl radical (*i.e.*, a heterocyclalkyl-alkylene- moiety). Typical heterocyclalkyl groups include, but are not limited to heterocyclalkyl- CH_2 -, 2-(heterocyclalkyl)ethan-1-yl, and the like, wherein the “heterocyclalkyl” portion includes any of the heterocyclalkyl groups described above, including those described in Principles of Modern Heterocyclic Chemistry. One skilled in the art will also understand that the

5 heterocyclyl group can be attached to the alkyl portion of the heterocyclyl alkyl by means of a carbon-carbon bond or a carbon-heteroatom bond, with the proviso that the resulting group is chemically stable. The heterocyclyl alkyl group comprises 3 to 20 carbon atoms, *e.g.*, the alkyl portion of the arylalkyl group is 1 to 6 carbon atoms and the heterocyclyl moiety is 2 to 14 carbon atoms. Examples of heterocyclylalkyls
10 include by way of example and not limitation 5-membered sulfur, oxygen, and/or nitrogen containing heterocycles such as thiazolylmethyl, 2-thiazolylethan-1-yl, imidazolylmethyl, oxazolylmethyl, thiadiazolylmethyl, etc., 6-membered sulfur, oxygen, and/or nitrogen containing heterocycles such as piperidylmethyl, piperazinylmethyl, morpholylmethyl, pyridylmethyl, pyridizylmethyl,
15 pyrimidylmethyl, pyrazinylmethyl, etc.

“Heterocyclylalkenyl” refers to an acyclic alkenyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also a sp^2 carbon atom, is replaced with a heterocyclyl radical (*i.e.*, a heterocyclyl-alkenylene- moiety). The heterocyclyl portion of the heterocyclyl alkenyl group
20 includes any of the heterocyclyl groups described herein, including those described in Principles of Modern Heterocyclic Chemistry, and the alkenyl portion of the heterocyclyl alkenyl group includes any of the alkenyl groups disclosed herein. One skilled in the art will also understand that the heterocyclyl group can be attached to the alkenyl portion of the heterocyclyl alkenyl by means of a carbon-carbon bond or a
25 carbon-heteroatom bond, with the proviso that the resulting group is chemically stable. The heterocyclyl alkenyl group comprises 4 to 20 carbon atoms, *e.g.*, the alkenyl portion of the heterocyclyl alkenyl group is 2 to 6 carbon atoms and the heterocyclyl moiety is 2 to 14 carbon atoms.

“Heterocyclylalkynyl” refers to an acyclic alkynyl radical in which one of the
30 hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, but also an sp carbon atom, is replaced with a heterocyclyl radical (*i.e.*, a heterocyclyl-alkynylene- moiety). The heterocyclyl portion of the heterocyclyl alkynyl group includes any of the heterocyclyl groups described herein, including those described in Principles of Modern Heterocyclic Chemistry, and the alkynyl portion of the
35 heterocyclyl alkynyl group includes any of the alkynyl groups disclosed herein. One

5 skilled in the art will also understand that the heterocyclyl group can be attached to the alkynyl portion of the heterocyclyl alkynyl by means of a carbon-carbon bond or a carbon-heteroatom bond, with the proviso that the resulting group is chemically stable. The heterocyclyl alkynyl group comprises 4 to 20 carbon atoms, *e.g.*, the alkynyl portion of the heterocyclyl alkynyl group is 2 to 6 carbon atoms and the
10 heterocyclyl moiety is 2 to 14 carbon atoms.

“Heteroaryl” refers to an aromatic heterocyclyl having at least one heteroatom in the ring. Non-limiting examples of suitable heteroatoms which can be included in the aromatic ring include oxygen, sulfur, and nitrogen. Non-limiting examples of heteroaryl rings include all of those aromatic rings listed in the definition of
15 “heterocyclyl”, including pyridinyl, pyrrolyl, oxazolyl, indolyl, isoindolyl, purinyl, furanyl, thienyl, benzofuranyl, benzothiophenyl, carbazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, quinolyl, isoquinolyl, pyridazyl, pyrimidyl, pyrazyl, etc.

“Carbocycle” or “carbocyclyl” refers to a saturated (*i.e.*, cycloalkyl), partially
20 unsaturated (*e.g.*, cycloakenyl, cycloalkadienyl, etc.) or aromatic ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to about 20 carbon atoms as a polycycle. Monocyclic carbocycles have 3 to 7 ring atoms, still more typically 5 or 6 ring atoms. Bicyclic carbocycles have 7 to 12 ring atoms, *e.g.*, arranged as a bicyclo [4,5], [5,5], [5,6] or [6,6] system, or 9 or 10 ring atoms arranged
25 as a bicyclo [5,6] or [6,6] system, or spiro-fused rings. Non-limiting examples of monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, and phenyl. Non-limiting examples of bicyclo carbocycles includes naphthyl, tetrahydronaphthalene, and decaline.

30 “Carbocyclylalkyl” refers to to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom is replaced with a carbocyclyl radical as described herein. Typical, but non-limiting, examples of carbocyclylalkyl groups include cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

35 “Arylheteroalkyl” refers to a heteroalkyl as defined herein, in which a

5 hydrogen atom (which may be attached either to a carbon atom or a heteroatom) has been replaced with an aryl group as defined herein. The aryl groups may be bonded to a carbon atom of the heteroalkyl group, or to a heteroatom of the heteroalkyl group, provided that the resulting arylheteroalkyl group provides a chemically stable moiety. For example, an arylheteroalkyl group can have the general formulae -alkylene-
 10 O-aryl, -alkylene-O-alkylene-aryl, -alkylene-NH-aryl, -alkylene-NH-alkylene-aryl, -alkylene-S-aryl, -alkylene-S-alkylene-aryl, etc. In addition, any of the alkylene moieties in the general formulae above can be further substituted with any of the substituents defined or exemplified herein.

“Heteroarylalkyl” refers to an alkyl group, as defined herein, in which a
 15 hydrogen atom has been replaced with a heteroaryl group as defined herein. Non-limiting examples of heteroaryl alkyl include -CH₂-pyridinyl, -CH₂-pyrrolyl, -CH₂-oxazolyl, -CH₂-indolyl, -CH₂-isoindolyl, -CH₂-purinyl, -CH₂-furanyl, -CH₂-thienyl, -CH₂-benzofuranyl, -CH₂-benzothiophenyl, -CH₂-carbazolyl, -CH₂-imidazolyl, -CH₂-thiazolyl, -CH₂-isoxazolyl, -CH₂-pyrazolyl, -CH₂-isothiazolyl,
 20 -CH₂-quinolyl, -CH₂-isoquinolyl, -CH₂-pyridazyl, -CH₂-pyrimidyl, -CH₂-pyrazyl, -CH(CH₃)-pyridinyl, -CH(CH₃)-pyrrolyl, -CH(CH₃)-oxazolyl, -CH(CH₃)-indolyl, -CH(CH₃)-isoindolyl, -CH(CH₃)-purinyl, -CH(CH₃)-furanyl, -CH(CH₃)-thienyl, -CH(CH₃)-benzofuranyl, -CH(CH₃)-benzothiophenyl, -CH(CH₃)-carbazolyl, -CH(CH₃)-imidazolyl, -CH(CH₃)-thiazolyl, -CH(CH₃)-isoxazolyl,
 25 -CH(CH₃)-pyrazolyl, -CH(CH₃)-isothiazolyl, -CH(CH₃)-quinolyl, -CH(CH₃)-isoquinolyl, -CH(CH₃)-pyridazyl, -CH(CH₃)-pyrimidyl, -CH(CH₃)-pyrazyl, etc.


The term “optionally substituted” in reference to a particular moiety of the compound of Formula I-III (e.g., an optionally substituted aryl group) refers to a
 30 moiety wherein all substituents are hydrogen or wherein one or more of the hydrogens of the moiety may be replaced by substituents such as those listed under the definition of “substituted” or as otherwise indicated.

The term “optionally replaced” in reference to a particular moiety of the compound of Formula I-III (e.g., the carbon atoms of said (C₁-C₈)alkyl may be
 35 optionally replaced by -O-, -S-, or -NR^a-) means that one or more of the methylene

5 groups of the (C₁-C₈)alkyl may be replaced by 0, 1, 2, or more of the groups specified (e.g., -O-, -S-, or -NR^a-).

The term “non-terminal carbon atom(s)” in reference to an alkyl, alkenyl, alkynyl, alkylene, alkenylene, or alkynylene moiety refers to the carbon atoms in the moiety that intervene between the first carbon atom of the moiety and the last carbon atom in the moiety. Therefore, by way of example and not limitation, in the alkyl
10 moiety -CH₂(C^{*})H₂(C^{*})H₂CH₃ or alkylene moiety -CH₂(C^{*})H₂(C^{*})H₂CH₂- the C^{*} atoms would be considered to be the non-terminal carbon atoms.

Certain Y and Y¹ alternatives are nitrogen oxides such as ⁺N(O)(R) or ⁺N(O)(OR). These nitrogen oxides, as shown here attached to a carbon atom, can also


15 be represented by charge separated groups such as , respectively, and are intended to be equivalent to the aforementioned representations for the purposes of describing this invention.

"Linker" or "link" means a chemical moiety comprising a covalent bond or a chain of atoms. Linkers include repeating units of alkyloxy (e.g. polyethyleneoxy, PEG, polymethyleneoxy) and alkylamino (e.g. polyethyleneamino, JeffamineTM); and
20 diacid ester and amides including succinate, succinamide, diglycolate, malonate, and caproamide.


The terms such as “oxygen-linked”, “nitrogen-linked”, “carbon-linked”, “sulfur-linked”, or “phosphorous-linked” mean that if a bond between two moieties
25 can be formed by using more than one type of atom in a moiety, then the bond formed between the moieties is through the atom specified. For example, a nitrogen-linked amino acid would be bonded through a nitrogen atom of the amino acid rather than through an oxygen or carbon atom of the amino acid.

Unless otherwise specified, the carbon atoms of the compounds of Formula I-
30 III are intended to have a valence of four. In some chemical structure representations where carbon atoms do not have a sufficient number of variables attached to produce a valence of four, the remaining carbon substituents needed to provide a valence of

5 four should be assumed to be hydrogen. For example,



has the same meaning as



“Protecting group” refers to a moiety of a compound that masks or alters the properties of a functional group or the properties of the compound as a whole. The chemical substructure of a protecting group varies widely. One function of a

10 protecting group is to serve as an intermediate in the synthesis of the parental drug substance. Chemical protecting groups and strategies for protection/deprotection are well known in the art. See: "Protective Groups in Organic Chemistry", Theodora W. Greene (John Wiley & Sons, Inc., New York, 1991). Protecting groups are often

15 utilized to mask the reactivity of certain functional groups, to assist in the efficiency of desired chemical reactions, e.g. making and breaking chemical bonds in an ordered and planned fashion. Protection of functional groups of a compound alters other physical properties besides the reactivity of the protected functional group, such as the polarity, lipophilicity (hydrophobicity), and other properties which can be measured

20 by common analytical tools. Chemically protected intermediates may themselves be

5 biologically active or inactive.

Protected compounds may also exhibit altered, and in some cases, optimized properties *in vitro* and *in vivo*, such as passage through cellular membranes and resistance to enzymatic degradation or sequestration. In this role, protected compounds with intended therapeutic effects may be referred to as prodrugs. Another
10 function of a protecting group is to convert the parental drug into a prodrug, whereby the parental drug is released upon conversion of the prodrug *in vivo*. Because active prodrugs may be absorbed more effectively than the parental drug, prodrugs may possess greater potency *in vivo* than the parental drug. Protecting groups are removed either *in vitro*, in the instance of chemical intermediates, or *in vivo*, in the case of
15 prodrugs. With chemical intermediates, it is not particularly important that the resulting products after deprotection, e.g. alcohols, be physiologically acceptable, although in general it is more desirable if the products are pharmacologically innocuous.

“Prodrug moiety” means a labile functional group which separates from the
20 active inhibitory compound during metabolism, systemically, inside a cell, by hydrolysis, enzymatic cleavage, or by some other process (Bundgaard, Hans, “Design and Application of Prodrugs” in Textbook of Drug Design and Development (1991), P. Krogsgaard-Larsen and H. Bundgaard, Eds. Harwood Academic Publishers, pp. 113-191). Enzymes which are capable of an enzymatic activation mechanism with the
25 phosphonate prodrug compounds of the invention include, but are not limited to, amidases, esterases, microbial enzymes, phospholipases, cholinesterases, and phosphatases. Prodrug moieties can serve to enhance solubility, absorption and lipophilicity to optimize drug delivery, bioavailability and efficacy.

A prodrug moiety may include an active metabolite or drug itself.

30 Exemplary prodrug moieties include the hydrolytically sensitive or labile acyloxymethyl esters $-\text{CH}_2\text{OC}(=\text{O})\text{R}^{30}$ and acyloxymethyl carbonates $-\text{CH}_2\text{OC}(=\text{O})\text{OR}^{30}$ where R^{30} is C_1 – C_6 alkyl, C_1 – C_6 substituted alkyl, C_6 – C_{20} aryl or C_6 – C_{20} substituted aryl. The acyloxyalkyl ester was used as a prodrug strategy for carboxylic acids and then applied to phosphates and phosphonates by Farquhar et al
35 (1983) *J. Pharm. Sci.* 72: 324; also US Patent Nos. 4816570, 4968788, 5663159 and

5 5792756. In certain compounds of the invention, a prodrug moiety is part of a phosphate group. The acyloxyalkyl ester may be used to deliver phosphoric acids across cell membranes and to enhance oral bioavailability. A close variant of the acyloxyalkyl ester, the alkoxycarbonyloxyalkyl ester (carbonate), may also enhance oral bioavailability as a prodrug moiety in the compounds of the combinations of the invention. An exemplary acyloxymethyl ester is pivaloyloxymethoxy, (POM) 10 $-\text{CH}_2\text{OC}(=\text{O})\text{C}(\text{CH}_3)_3$. An exemplary acyloxymethyl carbonate prodrug moiety is pivaloyloxymethylcarbonate (POC) $-\text{CH}_2\text{OC}(=\text{O})\text{OC}(\text{CH}_3)_3$.

The phosphate group may be a phosphate prodrug moiety. The prodrug moiety may be sensitive to hydrolysis, such as, but not limited to those comprising a 15 pivaloyloxymethyl carbonate (POC) or POM group. Alternatively, the prodrug moiety may be sensitive to enzymatic potentiated cleavage, such as a lactate ester or a phosphonamidate-ester group.

Aryl esters of phosphorus groups, especially phenyl esters, are reported to enhance oral bioavailability (DeLambert et al (1994) *J. Med. Chem.* 37: 498). Phenyl 20 esters containing a carboxylic ester ortho to the phosphate have also been described (Khamnei and Torrence, (1996) *J. Med. Chem.* 39:4109-4115). Benzyl esters are reported to generate the parent phosphonic acid. In some cases, substituents at the *ortho*-or *para*-position may accelerate the hydrolysis. Benzyl analogs with an acylated phenol or an alkylated phenol may generate the phenolic compound through 25 the action of enzymes, e.g. esterases, oxidases, etc., which in turn undergoes cleavage at the benzylic C–O bond to generate the phosphoric acid and the quinone methide intermediate. Examples of this class of prodrugs are described by Mitchell et al (1992) *J. Chem. Soc. Perkin Trans. I* 2345; Brook et al WO 91/19721. Still other benzylic prodrugs have been described containing a carboxylic ester-containing group 30 attached to the benzylic methylene (Glazier et al WO 91/19721). Thio-containing prodrugs are reported to be useful for the intracellular delivery of phosphonate drugs. These proesters contain an ethylthio group in which the thiol group is either esterified with an acyl group or combined with another thiol group to form a disulfide. Deesterification or reduction of the disulfide generates the free thio intermediate 35 which subsequently breaks down to the phosphoric acid and episulfide (Puech et al

5 (1993) *Antiviral Res.*, 22: 155-174; Benzaria et al (1996) *J. Med. Chem.* 39: 4958).
Cyclic phosphonate esters have also been described as prodrugs of phosphorus-
containing compounds (Erion et al, US Patent No. 6312662).

It is to be noted that all enantiomers, diastereomers, and racemic mixtures,
tautomers, polymorphs, pseudopolymorphs of compounds within the scope of
10 Formula I, Formula II, or Formula III and pharmaceutically acceptable salts thereof
are embraced by the present invention. All mixtures of such enantiomers and
diastereomers are within the scope of the present invention.

A compound of Formula I-III and its pharmaceutically acceptable salts may
exist as different polymorphs or pseudopolymorphs. As used herein, crystalline
15 polymorphism means the ability of a crystalline compound to exist in different crystal
structures. The crystalline polymorphism may result from differences in crystal
packing (packing polymorphism) or differences in packing between different
conformers of the same molecule (conformational polymorphism). As used herein,
crystalline pseudopolymorphism means the ability of a hydrate or solvate of a
20 compound to exist in different crystal structures. The pseudopolymorphs of the
instant invention may exist due to differences in crystal packing (packing
pseudopolymorphism) or due to differences in packing between different conformers
of the same molecule (conformational pseudopolymorphism). The instant invention
comprises all polymorphs and pseudopolymorphs of the compounds of Formula I-III
25 and their pharmaceutically acceptable salts.

A compound of Formula I-III and its pharmaceutically acceptable salts may
also exist as an amorphous solid. As used herein, an amorphous solid is a solid in
which there is no long-range order of the positions of the atoms in the solid. This
definition applies as well when the crystal size is two nanometers or less. Additives,
30 including solvents, may be used to create the amorphous forms of the instant
invention. The instant invention comprises all amorphous forms of the compounds of
Formula I-III and their pharmaceutically acceptable salts.

Selected substituents comprising the compounds of Formula I-III are present
to a recursive degree. In this context, "recursive substituent" means that a substituent
35 may recite another instance of itself. Because of the recursive nature of such


5 substituents, theoretically, a large number of compounds may be present in any given embodiment. For example, R^x comprises a R^y substituent. R^y can be R. R can be W^3 . W^3 can be W^4 and W^4 can be R or comprise substituents comprising R^y . One of ordinary skill in the art of medicinal chemistry understands that the total number of such substituents is reasonably limited by the desired properties of the compound intended. Such properties include, by way of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such as activity against the intended target, and practical properties such as ease of synthesis.

By way of example and not limitation, W^3 and R^y are recursive substituents in certain embodiments. Typically, each recursive substituent can independently occur 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0, times in a given embodiment. More typically, each recursive substituent can independently occur 12 or fewer times in a given embodiment. Even more typically, each recursive substituent can independently occur 3 or fewer times in a given embodiment. For example, W^3 will occur 0 to 8 times, R^y will occur 0 to 6 times in a given embodiment. Even more typically, W^3 will occur 0 to 6 times and R^y will occur 0 to 4 times in a given embodiment.

Recursive substituents are an intended aspect of the invention. One of ordinary skill in the art of medicinal chemistry understands the versatility of such substituents. To the degree that recursive substituents are present in an embodiment of the invention, the total number will be determined as set forth above.


The modifier "about" used in connection with a quantity is inclusive of the stated value and has the meaning dictated by the context (e.g., includes the degree of error associated with measurement of the particular quantity).

The compounds of the Formula I-III may comprise a phosphate group as R^7 ,



30 which may be a prodrug moiety wherein each Y or Y^1 is, independently, O, S, NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, or $N-NR_2$; W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or one of W^1 or W^2 together with either


- 5 R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or W^1 and W^2 are each, independently, a group of Formula Ia:



wherein:

- each Y^2 is independently a bond, O, CR_2 , NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$,
 10 $N-NR_2$, S, S-S, S(O), or S(O)₂;
 each Y^3 is independently O, S, or NR;
 M_2 is 0, 1 or 2;
 each R^y is independently H, F, Cl, Br, I, OH, R, $-C(=Y^1)R$, $-C(=Y^1)OR$, $-C(=Y^1)N(R)_2$, $-N(R)_2$, $^+N(R)_3$, $-SR$, $-S(O)R$, $-S(O)_2R$, $-S(O)(OR)$, $-S(O)_2(OR)$, $-OC(=Y^1)R$, $-OC(=Y^1)OR$, $-OC(=Y^1)N(R)_2$, $-SC(=Y^1)R$, $-SC(=Y^1)OR$, $-SC(=Y^1)N(R)_2$, $-N(R)C(=Y^1)R$, $-N(R)C(=Y^1)OR$, or $-N(R)C(=Y^1)N(R)_2$, $-SO_2NR_2$, $-CN$, $-N_3$, $-NO_2$, $-OR$, a protecting group or W^3 ; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R^x is independently R^y , a protecting group, or the formula:



20

wherein:

- M_{1a} , M_{1c} , and M_{1d} are independently 0 or 1;
 M_{12c} is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;
 each R is H, halogen, (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl,
 25 (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl, (C₂-C₈) substituted alkynyl, C₆-C₂₀ aryl,


5 C₆–C₂₀ substituted aryl, C₂–C₂₀ heterocycle, C₂–C₂₀ substituted heterocyclyl, arylalkyl, substituted arylalkyl or a protecting group;

W³ is W⁴ or W⁵; W⁴ is R, -C(Y¹)R^y, -C(Y¹)W⁵, -SO₂R^y, or -SO₂W⁵; and W⁵ is a carbocycle or a heterocycle wherein W⁵ is independently substituted with 0 to 3 R^y groups.


10 W⁵ carbocycles and W⁵ heterocycles may be independently substituted with 0 to 3 R^y groups. W⁵ may be a saturated, unsaturated or aromatic ring comprising a mono- or bicyclic carbocycle or heterocycle. W⁵ may have 3 to 10 ring atoms, e.g., 3 to 7 ring atoms. The W⁵ rings are saturated when containing 3 ring atoms, saturated or mono-unsaturated when containing 4 ring atoms, saturated, or mono- or di-
 15 unsaturated when containing 5 ring atoms, and saturated, mono- or di-unsaturated, or aromatic when containing 6 ring atoms.

A W⁵ heterocycle may be a monocycle having 3 to 7 ring members (2 to 6 carbon atoms and 1 to 3 heteroatoms selected from N, O, P, and S) or a bicycle having 7 to 10 ring members (4 to 9 carbon atoms and 1 to 3 heteroatoms selected from N, O, P, and S). W⁵ heterocyclic monocycles may have 3 to 6 ring atoms (2 to 5 carbon
 20 atoms and 1 to 2 heteroatoms selected from N, O, and S); or 5 or 6 ring atoms (3 to 5 carbon atoms and 1 to 2 heteroatoms selected from N and S). W⁵ heterocyclic bicycles have 7 to 10 ring atoms (6 to 9 carbon atoms and 1 to 2 heteroatoms selected from N, O, and S) arranged as a bicyclo [4,5], [5,5], [5,6], or [6,6] system; or 9 to 10
 25 ring atoms (8 to 9 carbon atoms and 1 to 2 hetero atoms selected from N and S) arranged as a bicyclo [5,6] or [6,6] system. The W⁵ heterocycle may be bonded to Y² through a carbon, nitrogen, sulfur or other atom by a stable covalent bond.

W⁵ heterocycles include for example, pyridyl, dihydropyridyl isomers, piperidine, pyridazinyl, pyrimidinyl, pyrazinyl, s-triazinyl, oxazolyl, imidazolyl,
 30 thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, furanyl, thiofuranyl, thienyl, and pyrrolyl. W⁵ also includes, but is not limited to, examples such as:





W^5 carbocycles and heterocycles may be independently substituted with 0 to 3 R groups, as defined above. For example, substituted W^5 carbocycles include:




10


Examples of substituted phenyl carbocycles include:



Embodiments of  of Formula I-III compounds include substructures such as:




wherein each Y^{2b} is, independently, O or N(R). In another aspect of this embodiment, each Y^{2b} is O and each R^x is independently:




wherein M12c is 1, 2 or 3 and each Y^2 is independently a bond, O, CR_2 , or S. In


another aspect of this embodiment, one $Y^{2b}-R^x$ is NH(R) and the other $Y^{2b}-R^x$ is O- R^x wherein R^x is:




wherein M12c is 2. In another aspect of this embodiment, each Y^{2b} is O and each R^x is independently:



10 wherein M12c is 2. In another aspect of this embodiment, each Y^{2b} is O and each R^x is independently:




wherein M12c is 1 and Y^2 is a bond, O, or CR_2 .




Other embodiments of of Formulas I-III compounds include


15 substructures such as:




- 5 wherein each Y^3 is, independently, O or N(R). In another aspect of this embodiment, each Y^3 is O. In another aspect of this embodiment, the substructure is:





wherein R^y is W^5 as defined herein.

- 10 Another embodiment of  of Formula I-III includes the substructures:






wherein each Y^{2c} is, independently, O, N(R^y) or S.




Another embodiment of  of Formula I-III compounds

- 15 includes the substructures wherein one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia. Such an embodiment is represented by a compound of Formula Ib selected from:


5






Formula Ib

In another aspect of the embodiment of Formula Ib, each Y and Y³ is O. In another aspect of the embodiment of Formula Ib, W¹ or W² is Y^{2b}-R^x; each Y, Y³ and Y^{2b} is O and R^x is:




M12c

wherein M12c is 1, 2 or 3 and each Y² is independently a bond, O, CR₂, or S. In another aspect of the embodiment of Formula Ib, W¹ or W² is Y^{2b}-R^x; each Y, Y³ and Y^{2b} is O and R^x is:




M12c

wherein M12c is 2. In another aspect of the embodiment of Formula Ib, W¹ or W² is Y^{2b}-R^x; each Y, Y³ and Y^{2b} is O and R^x is:



5


wherein M12c is 1 and Y² is a bond, O, or CR₂.




Another embodiment of

of Formula I-III compounds


10 includes a substructure:




wherein W⁵ is a carbocycle such as phenyl or substituted phenyl. In another aspect of this embodiment, the substructure is:




15 wherein Y^{2b} is O or N(R) and the phenyl carbocycle is substituted with 0 to 3 R groups. In another aspect of this embodiment of the substructure, R^x is:




wherein M12c is 1, 2 or 3 and each Y^2 is independently a bond, O, CR_2 , or S.




Another embodiment of of Formula I-III includes substructure:



The chiral carbon of the amino acid and lactate moieties may be either the *R* or *S* configuration or the racemic mixture.




Another embodiment of of Formula I-III is substructure




15 wherein each Y^2 is, independently, $-O-$ or $-NH-$. In another aspect of this embodiment, R^y is (C_1-C_8) alkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8)

- 5 substituted alkenyl, (C₂-C₈) alkynyl or (C₂-C₈) substituted alkynyl. In another aspect of this embodiment, R^y is (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl, (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl or (C₂-C₈) substituted alkynyl; and R is CH₃. In another aspect of this embodiment, R^y is (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl, (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl or (C₂-C₈) substituted alkynyl; R is CH₃; and each Y² is -NH-. In a aspect of this embodiment, W¹ and W² are, independently, nitrogen-linked, naturally occurring amino acids or naturally occurring amino acid esters. In another aspect of this embodiment, W¹ and W² are, independently, naturally-occurring 2-hydroxy carboxylic acids or naturally-occurring 2-hydroxy carboxylic acid esters wherein the acid or ester is linked to P through the 2-hydroxy group.




Another embodiment of of Formula I, Formula II, or Formula III is substructure:




- In one aspect of this embodiment, each R^x is, independently, (C₁-C₈) alkyl. In another aspect of this embodiment, each R^x is, independently, C₆-C₂₀ aryl or C₆-C₂₀ substituted aryl.


In a preferred embodiment,




is selected from



Another embodiment of



of Formulas I-III is substructure







- 10 wherein W^1 and W^2 are independently selected from one of the formulas in Tables 20.1-20.37 and Table 30.1 below. The variables used in Tables 20.1-20.37 (e.g., W^{23} , R^{21} , etc.) pertain only to Tables 20.1-20.37, unless otherwise indicated.



The variables used in Tables 20.1 to 20.37 have the following definitions:



- each R^{21} is independently H or (C_1-C_8) alkyl;
- 15 each R^{22} is independently H, R^{21} , R^{23} or R^{24} wherein each R^{24} is independently substituted with 0 to 3 R^{23} ;

- 5 each R^{23} is independently R^{23a} , R^{23b} , R^{23c} or R^{23d} , provided that when R^{23} is bound to a heteroatom, then R^{23} is R^{23c} or R^{23d} ;
- each R^{23a} is independently F, Cl, Br, I, -CN, N_3 or $-NO_2$;
- each R^{23b} is independently Y^{21} ;
- each R^{23c} is independently $-R^{2x}$, $-N(R^{2x})(R^{2x})$, $-SR^{2x}$, $-S(O)R^{2x}$, $-S(O)_2R^{2x}$, -
- 10 $S(O)(OR^{2x})$, $-S(O)_2(OR^{2x})$, $-OC(=Y^{21})R^{2x}$, $-OC(=Y^{21})OR^{2x}$, $-OC(=Y^{21})(N(R^{2x})(R^{2x}))$, $-SC(-Y^{21})R^{2x}$, $-SC(=Y^{21})OR^{2x}$, $-SC(=Y^{21})(N(R^{2x})(R^{2x}))$, $-N(R^{2x})C(=Y^{21})R^{2x}$, -
- $N(R^{2x})C(=Y^{21})OR^{2x}$, or $-N(R^{2x})C(=Y^{21})(N(R^{2x})(R^{2x}))$;
- each R^{23d} is independently $-C(=Y^{21})R^{2x}$, $-C(=Y^{21})OR^{2x}$ or -
- $C(=Y^{21})(N(R^{2x})(R^{2x}))$;
- 15 each R^{2x} is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl, heteroaryl; or two R^{2x} taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or $-NR^{21}-$; and wherein one or more of the non-terminal carbon atoms of each said (C_1-C_8) alkyl may be
- 20 optionally replaced with -O-, -S- or $-NR^{21}-$;
- each R^{24} is independently (C_1-C_8) alkyl, (C_2-C_8) alkenyl, or (C_2-C_8) alkynyl;
- each R^{25} is independently R^{24} wherein each R^{24} is substituted with 0 to 3 R^{23} groups;
- each R^{25a} is independently (C_1-C_8) alkylene, (C_2-C_8) alkenylene, or $(C_2-$
- 25 $C_8)$ alkynylene any one of which said (C_1-C_8) alkylene, (C_2-C_8) alkenylene, or (C_2-C_8) alkynylene is substituted with 0-3 R^{23} groups;
- each W^{23} is independently W^{24} or W^{25} ;
- each W^{24} is independently R^{25} , $-C(=Y^{21})R^{25}$, $-C(=Y^{21})W^{25}$, $-SO_2R^{25}$, or -
- SO_2W^{25} ;
- 30 each W^{25} is independently carbocycle or heterocycle wherein W^{25} is independently substituted with 0 to 3 R^{22} groups; and
- each Y^{21} is independently O or S.

5 Table 20.1Table 20.2


5 Table 20.3Table 20.4

5 Table 20.5Table 20.6


5 Table 20.7Table 20.8

5


Table 20.9




50




51




52




53




54



55




56




57


Table 20.10



58




59




60


10

5 Table 20.11


61




62




63




64




65




66




67




68

Table 20.12

69




70




71


002.1.1

5 Table 20.13


72




73




74




75




76




77




78




79

Table 20.14



80





81




82

5 Table 20.15Table 20.16

5 Table 20.17Table 20.18

10

Table 20.19

5

Table 20.20






Table 20.21





10


Table 20.22





15







5 Table 20.23Table 20.24


10

Table 20.25


15

5 Table 20.26Table 20.27


5 Table 20.28

5 Table 20.29


186




187




188




189




190




191




192




193

Table 20.30


194




195




196



197



198



199




5 Table 20.31

Table 20.32



5

Table 20.33

5

Table 20.34






















Table 20.35



10


5 Table 20.36Table 20.37

5 Table 30.1

10

5

Embodiments of R^x include esters, carbamates, carbonates, thioesters, amides, thioamides, and urea groups:



Any reference to the compounds of the invention described herein also includes a reference to a physiologically acceptable salt thereof. Examples of physiologically acceptable salts of the compounds of the invention include salts derived from an appropriate base, such as an alkali metal or an alkaline earth (for example, Na^+ , Li^+ , K^+ , Ca^{+2} and Mg^{+2}), ammonium and NR_4^+ (wherein R is defined herein). Physiologically acceptable salts of a nitrogen atom or an amino group include (a) acid addition salts formed with inorganic acids, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, sulfamic acids, phosphoric acid, nitric acid and the like; (b) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, isethionic acid, lactobionic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, malonic acid, sulfosalicylic acid, glycolic acid, 2-hydroxy-3-naphthoate, pamoate, salicylic acid, stearic acid, phthalic acid, mandelic acid, lactic acid, ethanesulfonic acid, lysine, arginine, glutamic acid, glycine, serine, threonine, alanine, isoleucine, leucine and the like; and (c) salts formed from elemental anions for example, chlorine, bromine, and iodine. Physiologically acceptable salts of a compound of a hydroxy group include the anion of said compound in combination with a suitable cation such as Na^+ and NR_4^+ .

For therapeutic use, salts of active ingredients of the compounds of the invention will be physiologically acceptable, i.e. they will be salts derived from a

5 physiologically acceptable acid or base. However, salts of acids or bases which are not physiologically acceptable may also find use, for example, in the preparation or purification of a physiologically acceptable compound. All salts, whether or not derived from a physiologically acceptable acid or base, are within the scope of the present invention.

10 Finally, it is to be understood that the compositions herein comprise compounds of the invention in their un-ionized, as well as zwitterionic form, and combinations with stoichiometric amounts of water as in hydrates.

The compounds of the invention, exemplified by Formula I-III may have chiral centers, e.g. chiral carbon or phosphorus atoms. The compounds of the
15 invention thus include racemic mixtures of all stereoisomers, including enantiomers, diastereomers, and atropisomers. In addition, the compounds of the invention include enriched or resolved optical isomers at any or all asymmetric, chiral atoms. In other words, the chiral centers apparent from the depictions are provided as the chiral isomers or racemic mixtures. Both racemic and diastereomeric mixtures, as well as
20 the individual optical isomers isolated or synthesized, substantially free of their enantiomeric or diastereomeric partners, are all within the scope of the invention. The racemic mixtures are separated into their individual, substantially optically pure isomers through well-known techniques such as, for example, the separation of diastereomeric salts formed with optically active adjuncts, e.g., acids or bases
25 followed by conversion back to the optically active substances. In most instances, the desired optical isomer is synthesized by means of stereospecific reactions, beginning with the appropriate stereoisomer of the desired starting material.

The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner, while the term "achiral" refers to
30 molecules which are superimposable on their mirror image partner.

The term "stereoisomers" refers to compounds which have identical chemical constitution, but differ with regard to the arrangement of the atoms or groups in space.

"Diastereomer" refers to a stereoisomer with two or more centers of chirality and whose molecules are not mirror images of one another. Diastereomers have
35 different physical properties, e.g. melting points, boiling points, spectral properties,

5 and reactivities. Mixtures of diastereomers may separate under high resolution analytical procedures such as electrophoresis and chromatography.

"Enantiomers" refer to two stereoisomers of a compound which are non-superimposable mirror images of one another.


Stereochemical definitions and conventions used herein generally follow S. P. Parker, Ed., McGraw-Hill Dictionary of Chemical Terms (1984) McGraw-Hill Book Company, New York; and Eliel, E. and Wilen, S., Stereochemistry of Organic Compounds (1994) John Wiley & Sons, Inc., New York. Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane-polarized light. In describing an optically active compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l, D and L, or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with S, (-), or l meaning that the compound is levorotatory while a compound prefixed with R, (+), or d is dextrorotatory. For a given chemical structure, these stereoisomers are identical except that they are mirror images of one another. A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric mixture. A 50:50 mixture of enantiomers is referred to as a racemic mixture or a racemate, which may occur where there has been no stereoselection or stereospecificity in a chemical reaction or process. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species, devoid of optical activity.

Whenever a compound described herein is substituted with more than one of the same designated group, e.g., "R" or "R¹", then it will be understood that the groups may be the same or different, i.e., each group is independently selected. Wavy lines, , indicate the site of covalent bond attachments to the adjoining substructures, groups, moieties, or atoms.

The compounds of the invention can also exist as tautomeric isomers in certain cases. Although only one delocalized resonance structure may be depicted, all such forms are contemplated within the scope of the invention. For example, ene-amine tautomers can exist for purine, pyrimidine, imidazole, guanidine, amidine, and

5 tetrazole systems and all their possible tautomeric forms are within the scope of the invention.

One skilled in the art will recognize that the pyrrolo[1,2-f][1,2,4]triazine, imidazo[1,5-f][1,2,4]triazine, imidazo[1,2-f][1,2,4]triazine, and [1,2,4]triazolo[4,3-f][1,2,4]triazine nucleosides can exist in tautomeric forms. For example, but not by
 10 way of limitation, structures (a) and (b) can have equivalent tautomeric forms as shown below:



15 All possible tautomeric forms of the heterocycles in all of the embodiments disclosed herein are within the scope of the invention.

Methods of Inhibition of HCV polymerase

Another aspect of the invention relates to methods of inhibiting the activity of HCV polymerase comprising the step of treating a sample suspected of containing HCV with a composition of the invention.

20 Compositions of the invention may act as inhibitors of HCV polymerase, as intermediates for such inhibitors or have other utilities as described below. The

5 inhibitors will bind to locations on the surface or in a cavity of HCV polymerase having a geometry unique to HCV polymerase. Compositions binding HCV polymerase may bind with varying degrees of reversibility. Those compounds binding substantially irreversibly are ideal candidates for use in this method of the invention. Once labeled, the substantially irreversibly binding compositions are useful as probes for the detection of HCV polymerase.

10 Accordingly, the invention relates to methods of detecting HCV polymerase in a sample suspected of containing HCV polymerase comprising the steps of: treating a sample suspected of containing HCV polymerase with a composition comprising a compound of the invention bound to a label; and observing the effect of the sample on the activity of the label. Suitable labels are well known in the diagnostics field and include stable free radicals,

15 fluorophores, radioisotopes, enzymes, chemiluminescent groups and chromogens. The compounds herein are labeled in conventional fashion using functional groups such as hydroxyl, carboxyl, sulfhydryl or amino.

Within the context of the invention, samples suspected of containing HCV polymerase include natural or man-made materials such as living organisms; tissue or cell

20 cultures; biological samples such as biological material samples (blood, serum, urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like. Typically the sample will be suspected of containing an organism which produces HCV polymerase, frequently a

25 pathogenic organism such as HCV. Samples can be contained in any medium including water and organic solvent\water mixtures. Samples include living organisms such as humans, and man made materials such as cell cultures.

The treating step of the invention comprises adding the composition of the invention to the sample or it comprises adding a precursor of the composition to the sample. The

30 addition step comprises any method of administration as described above.

If desired, the activity of HCV polymerase after application of the composition can be observed by any method including direct and indirect methods of detecting

5 HCV polymerase activity. Quantitative, qualitative, and semiquantitative methods of determining HCV polymerase activity are all contemplated. Typically one of the screening methods described above are applied, however, any other method such as observation of the physiological properties of a living organism are also applicable.

10 Organisms that contain HCV polymerase include the HCV virus. The compounds of this invention are useful in the treatment or prophylaxis of HCV infections in animals or in man.

However, in screening compounds capable of inhibiting human immunodeficiency viruses, it should be kept in mind that the results of enzyme assays may not correlate with cell culture assays. Thus, a cell based assay should be the primary screening tool.

15 Screens for HCV polymerase Inhibitors.

Compositions of the invention are screened for inhibitory activity against HCV polymerase by any of the conventional techniques for evaluating enzyme activity. Within the context of the invention, typically compositions are first screened for inhibition of HCV polymerase *in vitro* and compositions showing inhibitory activity are then screened for activity *in vivo*. Compositions having *in vitro* K_i (inhibitory constants) of less than about 5 $\times 10^{-6}$ M, typically less than about 1×10^{-7} M and preferably less than about 5×10^{-8} M are preferred for *in vivo* use.

Useful *in vitro* screens have been described in detail and will not be elaborated here. However, the examples describe suitable *in vitro* assays.

25 Pharmaceutical Formulations

The compounds of this invention are formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. All formulations will optionally contain excipients such as those set forth in the "Handbook of Pharmaceutical Excipients" (1986). Excipients include ascorbic acid and other antioxidants, chelating agents such

5 as EDTA, carbohydrates such as dextran, hydroxyalkylcellulose, hydroxyalkylmethylcellulose, stearic acid and the like. The pH of the formulations ranges from about 3 to about 11, but is ordinarily about 7 to 10.

While it is possible for the active ingredients to be administered alone it may be preferable to present them as pharmaceutical formulations. The formulations, both
10 for veterinary and for human use, of the invention comprise at least one active ingredient, as above defined, together with one or more acceptable carriers therefor and optionally other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and physiologically innocuous to the recipient thereof.

15 The formulations include those suitable for the foregoing administration routes. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Techniques and formulations generally are found in Remington's Pharmaceutical Sciences (Mack Publishing Co., Easton, PA). Such methods include the step of bringing into
20 association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be
25 presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste.

30 A tablet is made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be made by molding in a suitable
35 machine a mixture of the powdered active ingredient moistened with an inert liquid

5 diluent. The tablets may optionally be coated or scored and optionally are formulated so as to provide slow or controlled release of the active ingredient therefrom.

For infections of the eye or other external tissues e.g. mouth and skin, the formulations are preferably applied as a topical ointment or cream containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active
10 ingredient(s) in a range between 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc.), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base.

15 If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e. an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol (including PEG 400) and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or
20 penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethyl sulphoxide and related analogs.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an
25 emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and
30 the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.


Emulgents and emulsion stabilizers suitable for use in the formulation of the invention include Tween[®] 60, Span[®] 80, cetostearyl alcohol, benzyl alcohol, myristyl alcohol, glyceryl mono-stearate and sodium lauryl sulfate.

35 The choice of suitable oils or fats for the formulation is based on achieving the

5 desired cosmetic properties. The cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters
10 known as CrodamolTM CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils are used.

Pharmaceutical formulations according to the present invention comprise a
15 combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. Pharmaceutical formulations containing the active ingredient may be in any form suitable for the intended method of administration. When used for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft
20 capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents including sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic
25 pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as calcium or sodium carbonate, lactose, calcium or sodium phosphate; granulating and disintegrating agents, such as maize starch, or alginic acid; binding agents, such as starch, gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc. Tablets may be uncoated
30 or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a

- 5 longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.



5 Formulations for oral use may be also presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

 Aqueous suspensions of the invention contain the active materials in
10 admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally-occurring phosphatide (e.g., lecithin), a condensation
15 product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous suspension may also contain one or more preservatives
20 such as ethyl or n-propyl p-hydroxy-benzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose or saccharin.

 Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as arachis oil, olive oil, sesame oil or coconut oil, or in a mineral
25 oil such as liquid paraffin. The oral suspensions may contain a thickening agent, such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

30 Dispersible powders and granules of the invention suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening,
35 flavoring and coloring agents, may also be present.

5 The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as gum acacia and gum tragacanth, naturally-occurring phosphatides, such as soybean lecithin, esters or
10 partial esters derived from fatty acids and hexitol anhydrides, such as sorbitan monoolcate, and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also
15 contain a demulcent, a preservative, a flavoring or a coloring agent.

 The pharmaceutical compositions of the invention may be in the form of a sterile injectable preparation, such as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been
20 mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may
25 conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

 The amount of active ingredient that may be combined with the carrier
30 material to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a time-release formulation intended for oral administration to humans may contain approximately 1 to 1000 mg of active material compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95% of the total compositions
35 (weight:weight). The pharmaceutical composition can be prepared to provide easily

5 measurable amounts for administration. For example, an aqueous solution intended for intravenous infusion may contain from about 3 to 500 μg of the active ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about 30 mL/hr can occur.

Formulations suitable for topical administration to the eye also include eye
10 drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active ingredient. The active ingredient is preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10%, and particularly about 1.5% w/w.

Formulations suitable for topical administration in the mouth include lozenges
15 comprising the active ingredient in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with
20 a suitable base comprising for example cocoa butter or a salicylate.

Formulations suitable for intrapulmonary or nasal administration have a particle size for example in the range of 0.1 to 500 microns, such as 0.5, 1, 30, 35 etc., which is administered by rapid inhalation through the nasal passage or by inhalation through the mouth so as to reach the alveolar sacs. Suitable formulations include
25 aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol or dry powder administration may be prepared according to conventional methods and may be delivered with other therapeutic agents such as compounds heretofore used in the treatment or prophylaxis of HCV infections as described below.

Formulations suitable for vaginal administration may be presented as
30 pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers,
35 bacteriostats and solutes which render the formulation isotonic with the blood of the

5 intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

The formulations are presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection,
10 immediately prior to use. Extemporaneous injection solutions and suspensions are prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned
15 above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

The invention further provides veterinary compositions comprising at least one active ingredient as above defined together with a veterinary carrier therefor.

20 Veterinary carriers are materials useful for the purpose of administering the composition and may be solid, liquid or gaseous materials which are otherwise inert or acceptable in the veterinary art and are compatible with the active ingredient. These veterinary compositions may be administered orally, parenterally or by any other desired route.

25 Compounds of the invention are used to provide controlled release pharmaceutical formulations containing as active ingredient one or more compounds of the invention ("controlled release formulations") in which the release of the active ingredient are controlled and regulated to allow less frequency dosing or to improve the pharmacokinetic or toxicity profile of a given active ingredient.

30 Effective dose of active ingredient depends at least on the nature of the condition being treated, toxicity, whether the compound is being used prophylactically (lower doses) or against an active viral infection, the method of delivery, and the pharmaceutical formulation, and will be determined by the clinician using conventional dose escalation studies. It can be expected to be from about

5 0.0001 to about 100 mg/kg body weight per day; typically, from about 0.01 to about 10
mg/kg body weight per day; more typically, from about .01 to about 5 mg/kg body weight
per day; most typically, from about .05 to about 0.5 mg/kg body weight per day. For
example, the daily candidate dose for an adult human of approximately 70 kg body weight
will range from 1 mg to 1000 mg, preferably between 5 mg and 500 mg, and may take the
10 form of single or multiple doses.

Routes of Administration

One or more compounds of the invention (herein referred to as the active ingredients)
are administered by any route appropriate to the condition to be treated. Suitable routes
include oral, rectal, nasal, topical (including buccal and sublingual), vaginal and parenteral
15 (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural),
and the like. It will be appreciated that the preferred route may vary with for example the
condition of the recipient. An advantage of the compounds of this invention is that they are
orally bioavailable and can be dosed orally.

Combination Therapy

20 Combinations of the compounds of Formula I-III are typically selected based on the
condition to be treated, cross-reactivities of ingredients and pharmaco-properties of the
combination. For example, when treating an infection (e.g., HCV), the compositions of
the invention are combined with other active therapeutic agents (such as those described
herein).

25 Compositions of the invention are also used in combination with one or more other
active ingredients. Preferably, the other active therapeutic ingredients or agents are
interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-
glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists,
antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin
30 antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside
inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin
inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers or other drugs for treating
HCV; or mixtures thereof.

5

More specifically, one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of


- 1) interferons, *e.g.*, pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (PegasysTM), rIFN-alpha 2b (Intron ATM), rIFN-alpha 2a (Roferon-ATM), interferon alpha (MOR-22, OPC-18, Alfaferone, Alfaferone, Multiferon, subalin), interferon alfacon-1 (InfergenTM), interferon alpha-n1 (WellferonTM), interferon alpha-n3 (AlferonTM), interferon-beta (AvonexTM, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (AlbuferonTM), IFN alpha XL, BLX-883 (LocteronTM), DA-3021, glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon lambda (PEGylated IL-15 29), and belerofon,
- 2) ribavirin and its analogs, *e.g.*, ribavirin (RebetolTM, CopegusTM), and taribavirin (ViramidineTM),
- 3) HCV NS3 protease inhibitors, *e.g.*, boceprevir (SCH-503034 , SCH-7), telaprevir (VX-950), VX-813, TMC-435 (TMC435350), ABT-450, BI-201335, BI-1230, MK-7009, 20 SCH-900518, VBY-376, VX-500, GS-9256, GS-9451, BMS-790052, BMS-605339, PHX-1766, AS-101, YH-5258, YH5530, YH5531, and ITMN-191 (R-7227),
- 4) alpha-glucosidase 1 inhibitors, *e.g.*, celgosivir (MX-3253), Miglitol, and UT-231B,
- 5) hepatoprotectants, *e.g.*, emericasan (IDN-6556), ME-3738, GS-9450 (LB-84451), 25 silibinin, and MitoQ,
- 6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, *e.g.*, R1626, R7128 (R4048), IDX184, IDX-102, PSI-7851, BCX-4678, valopicitabine (NM-283), and MK-0608,
- 7) non-nucleoside inhibitors of HCV NS5B polymerase, *e.g.*, filibuvir (PF-868554), 30 ABT-333, ABT-072, BI-207127, VCH-759, VCH-916, JTK-652, MK-3281, VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773, A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, and GS-9190,

- 5 8) HCV NS5A inhibitors, *e.g.*, AZD-2836 (A-831), AZD-7295 (A-689), and BMS-790052,
- 9) TLR-7 agonists, *e.g.*, imiquimod, 852A, GS-9524, ANA-773, ANA-975, AZD-8848 (DSP-3025), PF-04878691, and SM-360320,
- 10) cyclophilin inhibitors, *e.g.*, DEBIO-025, SCY-635, and NIM811,
- 10 11) HCV IRES inhibitors, *e.g.*, MCI-067,
- 12) pharmacokinetic enhancers, *e.g.*, BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin,
- 13) other drugs for treating HCV, *e.g.*, thymosin alpha 1 (ZadaxinTM), nitazoxanide (AlineaTM, NTZ), BIVN-401 (virostatTM), PYN-17 (altirexTM), KPE02003002, actilon (CPG-15 10101), GS-9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225, PTX-111, ITX2865, TT-033i, ANA 971, NOV-205, tarvacin, EHC-18, VGX-410C, EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-4538), Oglufanide, FK-788, and VX-497 (merimepodib)
- 14) mevalonate decarboxylase antagonists, *e.g.*, statins, HMGCoA synthase inhibitors (*e.g.*, hymeglusin), squalene synthesis inhibitors (*e.g.*, zaragozic acid);
- 20 15) angiotensin II receptor antagonists, *e.g.*, losartan, irbesartan, olmesartan, candesartan, valsartan, telmisartan, eprosartan;
- 16) angiotensin-converting enzyme inhibitors, *e.g.*, captopril, zofenopril, enalapril, ramipril, quinapril, perindopril, lisinopril, benazepril, fosinopril;
- 25 17) other anti-fibrotic agents, *e.g.*, amiloride and
- 18) endothelin antagonists, *e.g.* bosentan and ambrisentan.

 In yet another embodiment, the present application discloses pharmaceutical compositions comprising a compound of the present invention, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, in combination with at least one additional

30 therapeutic agent, and a pharmaceutically acceptable carrier or excipient.

 According to the present invention, the therapeutic agent used in combination with the compound or composition of the present invention can be any agent having a therapeutic

- 5 effect when used in combination with the compound of the present invention. For example, the therapeutic agent used in combination with the compound or composition of the present invention can be interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of
- 

5 the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers or other drugs for treating HCV; or mixtures thereof.

10 More specifically, compositions of one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of

1) interferons, *e.g.*, pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon
15 alpha (MOR-22, OPC-18, Alfaferone, Alfaferone, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (Albuferon), IFN alpha XL, BLX-883 (Locteron), DA-3021, glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon
20 lambda (PEGylated IL-29), and belerofon,

2) ribavirin and its analogs, *e.g.*, ribavirin (Rebetol, Copegus), and taribavirin (Viramidine),

3) HCV NS3 protease inhibitors, *e.g.*, bocoprevir (SCH-503034, SCH-7), telaprevir (VX-950), VX-813, TMC-435 (TMC435350), ABT-450, BI-201335, BI-
25 1230, MK-7009, SCH-900518, VBY-376, VX-500, GS-9256, GS-9451, BMS-790052, BMS-605339, PHX-1766, AS-101, YH-5258, YH5530, YH5531, and ITMN-191 (R-7227),

4) alpha-glucosidase 1 inhibitors, *e.g.*, celgosivir (MX-3253), Miglitol, and UT-231B,

30 5) hepatoprotectants, *e.g.*, emericasan (IDN-6556), ME-3738, GS-9450 (LB-84451), silibinin, and MitoQ,

6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, *e.g.*, R1626, R7128 (R4048), IDX184, IDX-102, PSI-7851, BCX-4678, valopicitabine (NM-283), and MK-0608,

- 5 7) non-nucleoside inhibitors of HCV NS5B polymerase, *e.g.*, filibuvir (PF-868554), ABT-333, ABT-072, BI-207127, VCH-759, VCH-916, JTK-652, MK-3281, VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773, A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, and GS-9190,
- 10 8) HCV NS5A inhibitors, *e.g.*, AZD-2836 (A-831), AZD-7295 (A-689), and BMS-790052,
- 9) TLR-7 agonists, *e.g.*, imiquimod, 852A, GS-9524, ANA-773, ANA-975, AZD-8848 (DSP-3025), PF-04878691, and SM-360320,
- 10) cyclophilin inhibitors, *e.g.*, DEBIO-025, SCY-635, and NIM811,
- 15 11) HCV IRES inhibitors, *e.g.*, MCI-067,
- 12) pharmacokinetic enhancers, *e.g.*, BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin,
- 13) other drugs for treating HCV, *e.g.*, thymosin alpha 1 (Zadaxin), nitazoxanide (Alinea, NTZ), BIVN-401 (virostat), PYN-17 (altirex), KPE02003002, actilon (CPG-10101), GS-9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225, 20 PTX-111, ITX2865, TT-033i, ANA 971, NOV-205, tarvacin, EHC-18, VGX-410C, EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-4538), Oglufanide, FK-788, and VX-497 (merimepodib)
- 14) mevalonate decarboxylase antagonists, *e.g.*, statins, HMGCoA synthase inhibitors (*e.g.*, hymeglusin), squalene synthesis inhibitors (*e.g.*, zaragozic acid);
- 25 15) angiotensin II receptor antagonists, *e.g.*, losartan, irbesartan, olmesartan, candesartan, valsartan, telmisartan, eprosartan;
- 16) angiotensin-converting enzyme inhibitors, *e.g.*, captopril, zofenopril, enalapril, ramipril, quinapril, perindopril, lisinopril, benazepril, fosinopril;
- 30 17) other anti-fibrotic agents, *e.g.*, amiloride and
- 18) endothelin antagonists, *e.g.* bosentan and ambrisentan.

In yet another embodiment, the present application provides a combination pharmaceutical agent comprising:

- a) a first pharmaceutical composition comprising a compound of the 35 present invention, or a pharmaceutically acceptable salt, solvate, or ester thereof; and

5 b) a second pharmaceutical composition comprising at least one additional therapeutic agent selected from the group consisting of HIV protease inhibiting compounds, HIV non-nucleoside inhibitors of reverse transcriptase, HIV nucleoside inhibitors of reverse transcriptase, HIV nucleotide inhibitors of reverse transcriptase, HIV integrase inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120
10 inhibitors, CCR5 inhibitors, interferons, ribavirin analogs, NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, cyclophilin inhibitors, hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating HCV, and combinations thereof.

 Combinations of the compounds of Formula I-III and additional active
15 therapeutic agents may be selected to treat patients infected with HCV and other conditions such as HIV infections. Accordingly, the compounds of Formula I-III may be combined with one or more compounds useful in treating HIV, for example HIV protease inhibiting compounds, HIV non-nucleoside inhibitors of reverse transcriptase, HIV nucleoside inhibitors of reverse transcriptase, HIV nucleotide
20 inhibitors of reverse transcriptase, HIV integrase inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120 inhibitors, CCR5 inhibitors, interferons, ribavirin analogs, NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, cyclophilin inhibitors, hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating HCV.

25 More specifically, one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of 1) HIV protease inhibitors, *e.g.*, amprenavir, atazanavir, fosamprenavir, indinavir, lopinavir, ritonavir, lopinavir + ritonavir, nelfinavir, saquinavir, tipranavir, brexanavir, darunavir, TMC-126, TMC-114, mozenavir (DMP-450), JE-2147 (AG1776),
30 AG1859, DG35, L-756423, RO0334649, KNI-272, DPC-681, DPC-684, and GW640385X, DG17, PPL-100, 2) a HIV non-nucleoside inhibitor of reverse transcriptase, *e.g.*, capravirine, emivirine, delaviridine, efavirenz, nevirapine, (+) calanolide A, etravirine, GW5634, DPC-083, DPC-961, DPC-963, MIV-150, and TMC-120, TMC-278 (rilpivirine), efavirenz, BILR 355 BS, VRX 840773, UK-
35 453,061, RDEA806, 3) a HIV nucleoside inhibitor of reverse transcriptase, *e.g.*,

5 zidovudine, emtricitabine, didanosine, stavudine, zalcitabine, lamivudine, abacavir,
 amdoxovir, elvucitabine, alovudine, MIV-210, racivir ($\pm$ -FTC), D-d4FC,
 emtricitabine, phosphazide, fozivudine tidoxil, fosaltivudine tidoxil, apricitibine
 (AVX754), amdoxovir, KP-1461, abacavir + lamivudine, abacavir + lamivudine +
 zidovudine, zidovudine + lamivudine, 4) a HIV nucleotide inhibitor of reverse
 10 transcriptase, *e.g.*, tenofovir, tenofovir disoproxil fumarate + emtricitabine, tenofovir
 disoproxil fumarate + emtricitabine + efavirenz, and adefovir, 5) a HIV integrase
 inhibitor, *e.g.*, curcumin, derivatives of curcumin, chicoric acid, derivatives of
 chicoric acid, 3,5-dicaffeoylquinic acid, derivatives of 3,5-dicaffeoylquinic acid,
 aurintricarboxylic acid, derivatives of aurintricarboxylic acid, caffeic acid phenethyl
 15 ester, derivatives of caffeic acid phenethyl ester, tyrphostin, derivatives of tyrphostin,
 quercetin, derivatives of quercetin, S-1360, zintevir (AR-177), L-870812, and L-
 870810, MK-0518 (raltegravir), BMS-707035, MK-2048, BA-011, BMS-538158,
 GSK364735C, 6) a gp41 inhibitor, *e.g.*, enfuvirtide, sifuvirtide, FB006M, TRI-1144,
 SPC3, DES6, Locus gp41, CovX, and REP 9, 7) a CXCR4 inhibitor, *e.g.*, AMD-070,
 20 8) an entry inhibitor, *e.g.*, SP01A, TNX-355, 9) a gp120 inhibitor, *e.g.*, BMS-488043
 and BlockAide/CR, 10) a G6PD and NADH-oxidase inhibitor, *e.g.*, immunitin, 10) a
 CCR5 inhibitor, *e.g.*, aplaviroc, vicriviroc, INCB9471, PRO-140, INCB15050, PF-
 232798, CCR5mAb004, and maraviroc, 11) an interferon, *e.g.*, pegylated rIFN-alpha
 2b, pegylated rIFN-alpha 2a, rIFN-alpha 2b, IFN alpha-2b XL, rIFN-alpha 2a,
 25 consensus IFN alpha, infergen, rebif, locteron, AVI-005, PEG-infergen, pegylated
 IFN-beta, oral interferon alpha, feron, reaferon, intermax alpha, r-IFN-beta, infergen +
 actimmune, IFN-omega with DUROS, and albuferon, 12) ribavirin analogs, *e.g.*,
 rebetol, copegus, VX-497, and viramidine (taribavirin) 13) NS5a inhibitors, *e.g.*, A-
 831, A-689 and BMS-790052, 14) NS5b polymerase inhibitors, *e.g.*, NM-283,
 30 valopicitabine, R1626, PSI-6130 (R1656), IDX184, PSI-7851, HCV-796, BILB 1941,
 MK-0608, NM-107, R7128, VCH-759, PF-868554, GSK625433, and XTL-2125, 15)
 NS3 protease inhibitors, *e.g.*, SCH-503034 (SCH-7), VX-950 (Telaprevir), ITMN-
 191, and BILN-2065, 16) alpha-glucosidase 1 inhibitors, *e.g.*, MX-3253 (celgosivir)
 and UT-231B, 17) hepatoprotectants, *e.g.*, IDN-6556, ME 3738, MitoQ, and LB-
 35 84451, 18) non-nucleoside inhibitors of HCV, *e.g.*, benzimidazole derivatives, benzo-

5 1,2,4-thiadiazine derivatives, and phenylalanine derivatives, 19) other drugs for treating HCV, *e.g.*, zadaxin, nitazoxanide (alinea), BIVN-401 (virostat), DEBIO-025, VGX-410C, EMZ-702, AVI 4065, bavituximab, oglufanide, PYN-17, KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975, XTL-6865, ANA 971, NOV-205, tarvacin, EHC-18, and NIM811, 19) pharmacokinetic enhancers, *e.g.*,
10 BAS-100 and SPI452, 20) RNAse H inhibitors, *e.g.*, ODN-93 and ODN-112, 21) other anti-HIV agents, *e.g.*, VGV-1, PA-457 (bevirimat), ampligen, HRG214, cytolin, polymun, VGX-410, KD247, AMZ 0026, CYT 99007, A-221 HIV, BAY 50-4798, MDX010 (iplimumab), PBS119, ALG889, and PA-1050040.

It is also possible to combine any compound of the invention with one or more
15 other active therapeutic agents in a unitary dosage form for simultaneous or sequential administration to a patient. The combination therapy may be administered as a simultaneous or sequential regimen. When administered sequentially, the combination may be administered in two or more administrations.

Co-administration of a compound of the invention with one or more other
20 active therapeutic agents generally refers to simultaneous or sequential administration of a compound of the invention and one or more other active therapeutic agents, such that therapeutically effective amounts of the compound of the invention and one or more other active therapeutic agents are both present in the body of the patient.

Co-administration includes administration of unit dosages of the compounds
25 of the invention before or after administration of unit dosages of one or more other active therapeutic agents, for example, administration of the compounds of the invention within seconds, minutes, or hours of the administration of one or more other active therapeutic agents. For example, a unit dose of a compound of the invention can be administered first, followed within seconds or minutes by administration of a
30 unit dose of one or more other active therapeutic agents. Alternatively, a unit dose of one or more other therapeutic agents can be administered first, followed by administration of a unit dose of a compound of the invention within seconds or minutes. In some cases, it may be desirable to administer a unit dose of a compound of the invention first, followed, after a period of hours (*e.g.*, 1-12 hours), by
35 administration of a unit dose of one or more other active therapeutic agents. In other

5 cases, it may be desirable to administer a unit dose of one or more other active therapeutic agents first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of a compound of the invention.

The combination therapy may provide “synergy” and “synergistic”, i.e. the effect achieved when the active ingredients used together is greater than the sum of
10 the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined formulation; (2) delivered by alternation or in parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are
15 administered or delivered sequentially, e.g. in separate tablets, pills or capsules, or by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e. serially, whereas in combination therapy, effective dosages of two or more active ingredients are administered together. A synergistic anti-viral effect denotes an antiviral effect
20 which is greater than the predicted purely additive effects of the individual compounds of the combination.

In still yet another embodiment, the present application provides for methods of inhibiting HCV polymerase in a cell, comprising: contacting a cell infected with HCV with an effective amount of a compound of Formula I-III, or a pharmaceutically
25 acceptable salt, solvate, and/or ester thereof, whereby HCV polymerase is inhibited.

In still yet another embodiment, the present application provides for methods of inhibiting HCV polymerase in a cell, comprising: contacting a cell infected with HCV with an effective amount of a compound of Formula I-III, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active
30 therapeutic agent, whereby HCV polymerase is inhibited.

In still yet another embodiment, the present application provides for methods of inhibiting HCV polymerase in a cell, comprising: contacting a cell infected with HCV with an effective amount of a compound of Formula I-III, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active
35 therapeutic agent selected from the group consisting of one or more interferons,

5 ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7
10 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

In still yet another embodiment, the present application provides for methods of treating HCV in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of Formula I-III, or a pharmaceutically acceptable
15 salt, solvate, and/or ester thereof.

In still yet another embodiment, the present application provides for methods of treating HCV in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of Formula I-III, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent,
20 whereby HCV polymerase is inhibited.

In still yet another embodiment, the present application provides for methods of treating HCV in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of Formula I-III, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent
25 selected from the group consisting of one or more interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, endothelin antagonists, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV
30 NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

In still yet another embodiment, the present application provides for the use of a compound of the present invention, or a pharmaceutically acceptable salt, solvate,

5 and/or ester thereof, for the preparation of a medicament for treating an HCV infection in a patient.

Metabolites of the Compounds of the Invention

Also falling within the scope of this invention are the *in vivo* metabolic products of the compounds described herein, to the extent such products are novel and
10 unobvious over the prior art. Such products may result for example from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes novel and unobvious compounds produced by a process comprising contacting a compound of this invention with a mammal for a period of
15 time sufficient to yield a metabolic product thereof. Such products typically are identified by preparing a radiolabelled (e.g. ^{14}C or ^3H) compound of the invention, administering it parenterally in a detectable dose (e.g. greater than about 0.5 mg/kg) to an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time for metabolism to occur (typically about 30 seconds to 30 hours) and isolating its
20 conversion products from the urine, blood or other biological samples. These products are easily isolated since they are labeled (others are isolated by the use of antibodies capable of binding epitopes surviving in the metabolite). The metabolite structures are determined in conventional fashion, e.g. by MS or NMR analysis. In general, analysis of metabolites is done in the same way as conventional drug
25 metabolism studies well-known to those skilled in the art. The conversion products, so long as they are not otherwise found *in vivo*, are useful in diagnostic assays for therapeutic dosing of the compounds of the invention even if they possess no HCV polymerase inhibitory activity of their own.

Recipes and methods for determining stability of compounds in surrogate
30 gastrointestinal secretions are known. Compounds are defined herein as stable in the gastrointestinal tract where less than about 50 mole percent of the protected groups are deprotected in surrogate intestinal or gastric juice upon incubation for 1 hour at 37°C. Simply because the compounds are stable to the gastrointestinal tract does not mean that they cannot be hydrolyzed *in vivo*. The prodrugs of the invention typically

- 5 will be stable in the digestive system but may be substantially hydrolyzed to the parental drug in the digestive lumen, liver or other metabolic organ, or within cells in general.

Examples

10

Certain abbreviations and acronyms are used in describing the experimental details. Although most of these would be understood by one skilled in the art, Table 1 contains a list of many of these abbreviations and acronyms.

- 15 Table 1. List of abbreviations and acronyms.

Abbreviation	Meaning
Ac ₂ O	acetic anhydride
AIBN	2,2'-azobis(2-methylpropionitrile)
Bn	benzyl
BnBr	benzylbromide
BSA	bis(trimethylsilyl)acetamide
BzCl	benzoyl chloride
CDI	carbonyl diimidazole
DABCO	1,4-diazabicyclo[2.2.2]octane
DBN	1,5-diazabicyclo[4.3.0]non-5-ene
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
DBU	1,5-diazabicyclo[5.4.0]undec-5-ene
DCA	dichloroacetamide
DCC	dicyclohexylcarbodiimide
DCM	dichloromethane
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMTCI	dimethoxytrityl chloride
DMSO	dimethylsulfoxide
DMTr	4, 4'-dimethoxytrityl


DMF	dimethylformamide
EtOAc	ethyl acetate
ESI	electrospray ionization
HMDS	hexamethyldisilazane
HPLC	High pressure liquid chromatography
LDA	lithium diisopropylamide
LRMS	low resolution mass spectrum
MCPBA	meta-chloroperoxybenzoic acid
MeCN	acetonitrile
MeOH	methanol
MMTC	mono methoxytrityl chloride
m/z or m/e	mass to charge ratio
MH ⁺	mass plus 1
MH ⁻	mass minus 1
MsOH	methanesulfonic acid
MS or ms	mass spectrum
NBS	N-bromosuccinimide
Ph	phenyl
rt or r.t.	room temperature
TBAF	tetrabutylammonium fluoride
TMSCl	chlorotrimethylsilane
TMSBr	bromotrimethylsilane
TMSI	iodotrimethylsilane
TMSOTf	(trimethylsilyl)trifluoromethylsulfonate
TEA	triethylamine
TBA	tributylamine
TBAP	tributylammonium pyrophosphate
TBSCl	t-butyldimethylsilyl chloride
TEAB	triethylammonium bicarbonate
TFA	trifluoroacetic acid

TLC or tlc	thin layer chromatography
Tr	triphenylmethyl
Tol	4-methylbenzoyl
Turbo Grignard	1:1 mixture of isopropylmagnesium chloride and lithium chloride
δ	parts per million down field from tetramethylsilane

5

Preparation of Compounds


Compound 1




10 To a suspension of 7-bromo-2,4-bis-methylsulfanylimidazo[2,1-f][1,2,4]triazine (prepared according to WO2008116064, 500 mg, 1.72 mmol) in anhydrous THF (5 mL) was dropwise added BuLi (1.6 M in hexanes, 1.61 mL, 2.41 mmol) at -78 °C. The suspension became red brown solution after 5 min, and then a mixture of **1a** (prepared according to WO 200631725, 675 mg, 1.81 mmol) and boron trifluoride etherate (2.40 mL, 1.89 mmol) in THF (5 mL) was added dropwise to the mixture. After stirring for 2 h at -78 °C, saturated NH₄Cl was added to quench the reaction. The mixture was diluted with ethyl acetate; the organic layer was washed with brine and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (EtOAc / hexanes), affording **1b** as a rich yellow foam (650 mg, 67%). ¹H NMR (400 MHz, CDCl₃): δ 8.13 (d, 2H), 8.03 (d, 2H), 7.81 (d, 1H), 7.59 (t, 1H), 7.45 (m, 3H), 7.36 (t, 2H), 6.40 (brs, 1H), 6.01 (dd, 1H), 4.78 (m, 2H), 4.60 (dd, 1H), 2.68 (s, 3H), 2.45 (s, 3H), 1.62 (d, 3H). ¹⁹F NMR (376 MHz, CDCl₃): δ -167.5. MS = 585.1 (M + H⁺).

15

20




To a solution of **1b** (820 mg, 1.40 mmol) in dichloromethane (20 mL) were added boron trifluoride etherate (2 mL) and triethylsilane (2 mL), and stirred at room temperature for 16 h. Additional boron trifluoride etherate (1 mL) and triethylsilane (1 mL) were added, and stirred for 7 d. The mixture was diluted with dichloromethane and saturated sodium bicarbonate. The organic layer was washed sequentially with water, saturated ammonium chloride and brine, dried over magnesium sulfate, and concentrated. The residue was purified by silica gel column chromatography (EtOAc / hexanes), affording **1c** (605 mg, 76%). ¹H NMR (400 MHz, CDCl₃): δ 8.10 (d, *J* = 7.2 Hz, 2H), 8.00 (d, *J* = 7.2 Hz, 2H), 7.66 (s, 1H), 7.61 (t, *J* = 7.2 Hz, 1H), 7.53 (t, *J* = 7.2 Hz, 1H), 7.46 (t, *J* = 7.2 Hz, 2H), 7.38 (t, *J* = 7.2 Hz, 2H), 5.78 (m, 2H), 4.80 (dd, 1H), 4.68 (m, 1H), 4.60 (dd, 1H), 2.68 (s, 3H), 2.65 (s, 3H), 1.32 (d, 3H). ¹⁹F NMR (376 MHz, CDCl₃): δ -149.9. MS = 569.1 (M + H⁺).



Compound **1c** (635 mg, 1.12 mmol) was placed in a steel bomb reactor. Liquid ammonia (~30 mL) was charged and the bomb reactor was tightly sealed. The mixture was stirred at 50 °C for 16 h. After cooling to room temperature, ammonia was evaporated and the solid residue was dissolved in THF (10 mL) and MeOH (10 mL). Sodium ethoxide (25% wt. 0.63 mL) was added and stirred at 60 °C for 40 min. The mixture was neutralized with AcOH and concentrated. The residue was purified

5 by RP HPLC, affording the product **1d** (175 mg, 48%). ^1H NMR (400 MHz, DMSO- d_6): δ 8.21 (brs, 2H), 7.60 (s, 1H), 5.45 (brs, 1H), 5.43 (d, 1H), 4.91 (t, 1H), 3.92 (m, 1H), 3.76 (m, 2H), 3.57 (m, 1H), 2.44 (s, 3H), 1.09 (d, 3H). ^{19}F NMR (376 MHz, DMSO- d_6): δ -153.5. MS = 330.1 ($\text{M} + \text{H}^+$).




10

1d**1e**


To a solution of **1d** (175 mg, 0.53 mmol) in dichloromethane (11 mL) was added MCPBA (370 mg, ~ 1.5 mmol) and stirred at room temperature for 16 h. The mixture was concentrated, affording crude **1e** which was used for the next reaction without purification. MS = 362.0 ($\text{M} + \text{H}^+$).

15

**1e****1**


Compound **1e** (obtained from the previous reaction) was placed in a steel bomb reactor. Liquid ammonia (~ 30 mL) was charged, and the bomb reactor was tightly sealed. The mixture was stirred at 115 $^{\circ}\text{C}$ for 3 d. After cooling to room temperature, ammonia was evaporated. The solid residue was purified by RP HPLC, affording compound **1** (105 mg, 66% in two steps). ^1H NMR (400 MHz, D_2O): δ 7.31 (s, 1H), 5.43 (d, $J = 25.2$ Hz, 1H), 4.07 (dd, $J = 9.6, 23.2$, 1H), 3.89 (m, 1H), 3.83 (dd, $J = 2.4, 12.8$ Hz, 1H), 3.67 (dd, $J = 4.8, 12.8$ Hz, 1H), 1.05 (d, $J = 22.8$ Hz, 3H). ^{19}F NMR (376 MHz, D_2O): δ -153.5. MS = 299.2 ($\text{M} + \text{H}^+$).

25

5 **Compound 2**

To a solution of compound 1 (82 mg, 0.28 mmol) in water (340 mL) was added adenosine deaminase (A5168 bovine spleen type IX from Sigma-Aldrich, 0.125 Unit per mL of water) and stirred at 37 °C for 4 h. The mixture was concentrated and purified by RP HPLC, affording compound 2 (56 mg, 68%). ¹H NMR (400 MHz, D₂O): δ 7.35 (s, 1H), 5.46 (d, *J* = 25.2 Hz, 1H), 4.08 (dd, *J* = 9.6, 22.6, 1H), 3.93 (m, 1H), 3.87 (dd, *J* = 2.4, 12.8 Hz, 1H), 3.71 (dd, *J* = 4.8, 12.8 Hz, 1H), 1.12 (d, *J* = 23.2 Hz, 3H). ¹⁹F NMR (376 MHz, D₂O): δ -153.4. MS = 300.2 (M + H⁺).

15

Compound 3


To a suspension of 7-bromo-pyrrolo[2,1-f][1,2,4]triazin-4-ylamine (prepared according to WO2007056170, 2.13 g, 10 mmol) in THF (20 mL) was added TMSCl (2.66 mL, 21 mmol) and stirred at room temperature for 16 h under argon. After cooling to -78 °C, a solution of BuLi (1.6 M, 21 mL, 33 mmol) in hexanes was added dropwise. The mixture was stirred for 1 h at the same temperature. A solution of 1a (prepared according to WO 200631725, 4.46 g, 12 mmol) in THF (10 mL) was then added. After stirring for 2 h at -78 °C, saturated ammonium chloride was added to quench the reaction. The mixture was extracted with ethyl acetate. The organic

25

5 extract was concentrated *in vacuo*. The residue was purified by silica gel chromatography (ethyl acetate / hexanes), affording **3b** as a yellow solid (1.6 g, 32%). MS = 507.1 (M + H⁺).

10 **Alternative procedure for Compound 3b using 1,2-bis-[(chlorodimethyl)silanyl]ethane instead of chlorotrimethylsilane**


To a suspension of 7-bromo-pyrrolo[2,1-f][1,2,4]triazin-4-ylamine (500 mg, 2.35 mmol) in THF (6.5 mL) was added BuLi (1.6 M in hexanes, 1.6 mL) at -78 °C. After 30 min., a solution of 1,2-bis-[(chlorodimethyl)silanyl]ethane (538 mg, 2.4 mmol) in THF (1.2 mL) was added. After 45 min., BuLi (1.6 mL) was added. After an additional 30 min., BuLi (1.5 mL) was added. After 30 min., a solution of **1a** (610 mg, 1.64 mmol) in THF (2 mL) was then added dropwise. The resulting mixture was stirred at -78 °C for 2 h under argon. Acetic acid (0.7 mL) was added dropwise to quench the reaction, followed by addition of saturated ammonium chloride. The mixture was extracted with ethyl acetate. The organic extract was concentrated *in vacuo*. The residue was purified by silica gel chromatography (ethyl acetate / hexanes), affording **3b** (320 mg, 40%). The starting **1a** was also recovered (350 mg) from the chromatography.



To a solution of compound **3b** (50 mg, 0.1 mmol) and TMSCN (67 uL, 0.5 mmol) in acetonitrile (2.0 mL) at 0 °C was added TMSOTf (91 uL, 0.5 mmol). The reaction mixture was stirred at room temperature for 1 h, then at 65 °C for 3 d. The reaction was quenched with saturated NaHCO₃ at room temperature, and diluted with


30

- 5 $\text{CH}_3\text{CO}_2\text{Et}$. The organic phase was separated, washed with brine, dried over Na_2SO_4 , filtered and concentrated. The residue was purified by RP-HPLC (acetonitrile / water), to give the desired compound **3c** (28 mg, 54%). MS = 516.1 ($\text{M} + \text{H}^+$).




- 10 To a solution of **3c** (56 mg, 0.11 mmol) in methanol (1.2 mL) was added ammonium hydroxide (28% in water, 0.8 mL) and stirred at room temperature for 16 h. The mixture was concentrated and the residue was purified by RP HPLC (water / acetonitrile), affording compound **3** (20 mg, 60%). ^1H NMR (500 MHz, D_2O): δ 7.88 (s, 1H), 7.07 (d, 1H), 6.92 (d, 1H), 4.17 (m, 2H), 4.04 (dd, 1H), 3.87 (dd, 1H), 1.15 (d, 15 3H). MS = 308.1 ($\text{M} + \text{H}^+$).

Compound 4



- 20 To a solution of compound **3b** (60 mg, 0.12 mmol) in methanol (0.5 mL) was added ammonium hydroxide (28% in water, 0.5 mL) and stirred at room temperature for 16 h. The mixture was concentrated and the residue was purified by RP HPLC (water / acetonitrile), affording compound **4** (25 mg, 70%). MS = 299.1 ($\text{M} + \text{H}^+$).

25 Compound 5



5

Compound **3b** was converted to compound **5a** by a procedure similar to conversion of **1b** to **1c**. Compound **5a** was then converted to compound **5** by a procedure similar to conversion of **3c** to **3**. ^1H NMR (300 MHz, D_2O): δ 7.68 (s, 1H), 6.75 (d, $J = 4.5$ Hz, 1H), 6.65 (d, $J = 4.5$ Hz, 1H), 5.65 (d, $J = 25.2$ Hz, 1H), 3.95 (m, 3H), 3.74 (dd, 1H), 0.98 (d, $J = 22.8$ Hz, 3H). ^{19}F NMR (282 MHz, D_2O): δ -154.2. MS = 283.2 ($\text{M} + \text{H}^+$).

10

General procedure for preparation of a nucleoside triphosphate:


15

To a pear-shaped flask (5-15 mL) is charged with a nucleoside (~20 mg). Trimethyl phosphate (0.5-1.0 mL) is added. The solution is cooled with ice-water bath. POCl_3 (40-45 mg) is added and stirred at 0 °C until the reaction is complete (1 to 4 h; the reaction progress is monitored by ion-exchange HPLC; analytical samples are prepared by taking ~3 μL of the reaction mixture and diluting it with 1.0 M $\text{Et}_3\text{NH}_2\text{CO}_3$ (30-50 μL)). A solution of pyrophosphate- Bu_3N (250 mg) and Bu_3N (90-105 mg) in acetonitrile or DMF (1-1.5 mL) is then added. The mixture is stirred at 0 °C for 0.3 to 2.5 h, and then the reaction is quenched with 1.0 M $\text{Et}_3\text{NH}_2\text{CO}_3$ (~5 mL). The resulting mixture is stirred for additional 0.5-1 h while warming up to room

20


5 temperature. The mixture is concentrated to dryness, re-dissolved in water (4 mL), and purified by ion exchange HPLC. The fractions containing the desired product is concentrated to dryness, dissolved in water (~5 mL), concentrated to dryness, and again dissolved in water (~5 mL). NaHCO₃ (30-50 mg) is added and concentrated to dryness. The residue is dissolved in water and concentrated to dryness again. This
 10 process is repeated 2-5 times. The residue is then subjected to C-18 HPLC purification, affording the desired product as a sodium or salt. Alternatively, the crude reaction mixture is subjected to C-18 HPLC first and then ion exchange HPLC purification to afford the desired product as a triethylammonium salt.

15 Compound TP-1



Compound TP-1 was prepared by the general method using **Compound 2** as starting material. ¹H NMR (300 MHz, D₂O): δ 7.44 (s, 1H), 5.45 (d, *J* = 25.5 Hz, 1H), 4.0-4.4 (m, 4H), 3.05 (m, NCH₂CH₃), 1.10 (m, NCH₂CH₃ and 2'-C-CH₃). ³¹P NMR (121.4 MHz, D₂O): δ -9.5 (d, *J* = 22.1 Hz), -11.0 (d, *J* = 19.9 Hz), -23.2 (t, *J* = 23.0 Hz). ¹⁹F NMR (282 MHz, D₂O): δ -153.9.


Compound TP-2



25

5 **Compound TP-2** was prepared by the general method using **Compound 3** as starting material. ^1H NMR (300 MHz, D_2O): δ 7.82 (s, 1H), 7.03 (d, 1H), 6.90 (d, 1H), 4.1-4.4 (m, 4H), 3.05 (m, NCH_2CH_3), 1.10 (m, NCH_2CH_3 and 2'-C- CH_3). ^{31}P NMR (121.4 MHz, D_2O): δ -10.7 (d, $J = 19.5$ Hz), -11.3 (d, $J = 19.8$ Hz), -23.1 (t, $J = 19.8$ Hz).


10

Compound TP-3

15 **Compound TP-3** was prepared by the general method using **Compound 5** as starting material. ^1H NMR (300 MHz, D_2O): δ 7.73 (s, 1H), 6.87 (d, 1H), 6.82 (d, 1H), 5.71 (d, $J = 24.6$ Hz, 1H), 4.0-4.4 (m, 4H), 3.05 (m, NCH_2CH_3), 1.14 (m, NCH_2CH_3), 1.00 (d, $J = 22.8$ Hz, 3H, 2'-C- CH_3). ^{31}P NMR (121.4 MHz, D_2O): δ -8.1 (d, $J = 22.1$ Hz), -11.1 (d, $J = 19.9$ Hz), -22.7 (t, $J = 23.0$ Hz). ^{19}F NMR (282 MHz, D_2O): δ -155.6. MS = 520.9 ($\text{M} - \text{H}^+$).

20

General procedure for preparation of a nucleoside prodrug (Method A):




5

To a solution of a nucleoside (0.1 mmol) in trimethylphosphite (1.0 mL) are added 1*H*-tetrazole (42 mg, 0.6 mmol) followed by addition of 2,2-dimethylthiopropionic acid S-(2-{diisopropylamino-[2-(2,2-dimethyl-propionylsulfanyl)-ethoxy]-phosphanyloxy}-ethyl) ester (prepared according to *J. Med. Chem.*, **1985**, 38, 3941, 90 mg, 0.2 mmol) at 0 °C. After stirring for 2 h, 30% hydrogen peroxide in H₂O (140 µL) was added to the mixture. The mixture was then allowed to warm up to room temperature. After 30 min stirring, 1 M Na₂S₂O₃ in H₂O (5 mL) was added to quench the reaction. The organic layer was washed with saturated aqueous Na₂CO₃ (10 mL x 2), brine, concentrated *in vacuo*. The residue was purified by RP-HPLC (MeCN–H₂O gradient) to afford a prodrug **A**.

15

Compound A-1



5 Compound **A-1** was prepared by Method A using compound **1** as starting material. ^1H NMR (400 MHz, CDCl_3): δ 7.42 (s, 1H), 5.47 (d, $J = 26.4$ Hz, 1H), 4.95 (brs, 2H), 4.59 (m, 2H), 4.35 (m, 1H, 4'-H), 4.18 (m, 2H, 5'-H), 4.10 (m, 4H), 3.13 (m, 4H), 1.24 (d, 3H), 1.22 (s, 9H), 1.19 (d, 9H). ^{31}P NMR (161.9 MHz, CDCl_3): δ -1.26. MS = 667.1 ($\text{M} + \text{H}^+$).


10

General procedure for preparation of a nucleoside prodrug (Method B):

Non-limiting examples of mono-phosphoramidate prodrugs comprising the instant invention may be prepared according to general Scheme 1.

15


Scheme 1



20 The general procedure comprises the reaction of an amino acid ester salt **19b**, e.g., HCl salt, with an aryl dichlorophosphate **19a** in the presence of about two to ten equivalents of a suitable base to give the phosphoramidate **19c**. Suitable bases include, but are not limited to, imidazoles, pyridines such as lutidine and DMAP,

5 tertiary amines such as triethylamine and DABCO, and substituted amidines such as DBN and DBU. Tertiary amines are particularly preferred. Preferably, the product of each step is used directly in the subsequent steps without recrystallization or chromatography. Specific, but non-limiting, examples of **19a**, **19b**, and **19c** can be found in WO 2006/121820 that is hereby incorporated by reference in its entirety. A
10 nucleoside base **19d** reacts with the phosphoramidate **19c** in the presence of a suitable base. Suitable bases include, but are not limited to, imidazoles, pyridines such as lutidine and DMAP, tertiary amines such as triethylamine and DABCO, and substituted amidines such as DBN and DBU. The product **B** may be isolated by recrystallization and/or chromatography.


15

Compound B-1**B-1**


20 Phenyl ethoxyalaninyl phosphorochloridate (124 mg, 0.42 mmol; prepared according to McGuigan et al, *J. Med. Chem.* **1993**, 36, 1048–1052) was added to a mixture of **Compound 3** (20 mg, 0.065 mmol) and *N*-methylimidazole (42 μ L, 0.52 mmol) in anhydrous trimethyl phosphate (0.8 mL). The reaction mixture is stirred for 3 h at room temperature, and then methanol was added to quench the reaction. The
25 methanol solvent is removed under reduced pressure. The residue was purified by reverse-phase HPLC and then by silica gel column chromatography (100% ethyl acetate), affording compound **B-1** (10 mg, 27%). ^{31}P NMR (121.4 MHz, CDCl_3): δ - 3.42, 3.77. MS = 563.0 ($\text{M} + \text{H}^+$), 561.0 ($\text{M} - \text{H}^+$).

30 **Compound B-2**


5

**B-2**

About 3.1 mmol of 4-chlorophenyl 2-propyloxycarbonyl phosphorochloridate (prepared according to McGuigan et al, *J. Med. Chem.* **1993**, 36, 1048–1052) is added to a mixture of about 0.5 mmol of **Compound 3** and about 3.8 mmol of *N*-methylimidazole in about 3 mL anhydrous trimethyl phosphate. The reaction mixture is stirred for about one hour to 24 hours at room temperature and methanol is added to quench the reaction. The methanol solvent is removed under reduced pressure. The residue is purified by reverse-phase HPLC to give compound **B-2**.

15 Compound B-3**B-3**

Compound B-3 was obtained by a similar procedure used for compound **B-1**.
 20 ^{31}P NMR (121.4 MHz, CDCl_3): δ -3.50, 3.76. MS = 577.2 ($\text{M} + \text{H}^+$).


Compound B-4**B-4**

5

Compound B-4 was obtained by a similar procedure used for compound **B-1**.

^{31}P NMR (162 MHz, CD_3OD): δ 2.2. MS = 633.4 ($\text{M} + \text{H}^+$).

Compound B-5




10

B-5

Compound B-5 was obtained by a similar procedure used for compound **B-1**.

^{31}P NMR (162 MHz, CDCl_3): δ 4.15, 4.27. MS = 549.3 ($\text{M} + \text{H}^+$).


15 Compound B-6

**B-6**

Compound B-6 was obtained by a similar procedure used for compound **B-1**.

20 ^{31}P NMR (162 MHz, CDCl_3): δ 3.50, 4.07. MS = 613.1 ($\text{M} + \text{H}^+$).


Compound B-7

**B-7**

Compound B-7 was obtained by a similar procedure used for compound **B-1**, using compound **5** as parent nucleoside. ^{31}P NMR (162 MHz, CDCl_3): δ 3.37, 3.97.

MS = 538.1 ($\text{M} + \text{H}^+$).


Compound B-8

**B-8**

Compound B-8 was obtained by a similar procedure used for compound **B-1**, using compound **5** as parent nucleoside. ^{31}P NMR (162 MHz, CDCl_3): δ 3.69, 4.39.

MS = 588.1 ($\text{M} + \text{H}^+$).

Alternative procedure for preparation of a nucleoside prodrug (Method C):

**C-1a**

5 Into a flask containing ethyl *L*-valine hydrochloride (2.5 g, 13.8 mmol, 1 equiv.) was added CH₂Cl₂ (46 mL, 0.3 M) and phenyl dichlorophosphate (2.1 mL, 13.8 mmol, 1 equiv.) before being cooled to -10 °C. After 10 minutes, TEA (3.8 mL, 13.8 mmol, 1 equiv) was added slowly to the reaction mixture over five minutes. The reaction was allowed to proceed for an hour before *p*-nitrophenol (1.9 g, 13.8 mmol, 1 equiv.) was added to the reaction mixture followed by addition of more TEA (3.8 mL, 13.8 mmol, 1 equiv.) over five minutes. The reaction was allowed to warm up and proceed for another two hours. The reaction was concentrated *in vacuo* and taken up in diethyl ether (200 mL). The insoluble salts were filtered off and the filtrate concentrated *in vacuo*. Flash column chromatography was carried out using 4/1 Hex / EtOAc to furnish a clear oil as **C-1a**.


¹H NMR (400 MHz, CDCl₃): δ 8.21 (s, 2 H), 7.41 – 7.20 (m, 7 H), 4.22 -4.05 (m, 3 H), 2.46 (s, 2 H), 1.99 (dd, *J* = 23.0, 20.1 Hz, 2 H), 1.68 (s, 1 H), 1.20 -1.05 (m, 8 H).

³¹P NMR (162 MHz, CDCl₃): δ -2.79 (dd, *J* = 28.0, 4.2 Hz).

LC MS *m/z* 422.99 [M + H⁺].

20

Compound C-1



C-1

25 Into a flask containing **compound 3** (70 mg, 0.23 mmol, 1 equiv.) was added THF (1 mL, 0.2 M) and NMP (1 mL, 0.2 M) before cooling to 0 °C. *t*-BuMgCl (560 μL, 2.5 equiv., 1M THF) was added slowly and allowed to stir for 5 minutes before the above phenolate **C-1a** (207 mg, 0.46 mmol, 2 equiv. dissolved in 500 μL of THF) was added. The reaction mixture was warmed to 50 °C. The reaction was monitored by LCMS. Once the reaction was complete, the mixture was then concentrated *in vacuo*, and the residue was purified by HPLC, affording **Compound C-1**.


30

5 ^1H NMR (400 MHz, CDCl_3) δ 7.87 (s, 1 H), 7.24 – 7.10 (m, 4 H), 7.03 (t, $J = 7.2$ Hz, 1H), 6.81 (d, $J = 4.6$ Hz, 1 H), 6.52 (d, $J = 4.7$ Hz, 1 H), 5.61 (s, 2H), 4.46 (dd, $J = 24.0, 11.4$ Hz, 2 H), 4.33 – 4.14 (m, 2 H), 4.06 (dt, $J = 7.2, 4.2$ Hz, 2 H), 3.82 – 3.70 (m, 1 H), 3.63 (t, $J = 10.6$ Hz, 2 H), 1.98 (s, 1 H), 1.17 (dd, $J = 14.8, 7.6$ Hz, 3 H), 0.82 (dd, $J = 22.8, 6.8$ Hz, 6 H).

10 ^{31}P NMR (162 MHz, CDCl_3): δ 5.11.

^{19}F NMR (376 MHz, CDCl_3): δ -152.28.

LC MS m/z 591.21 $[\text{M} + \text{H}^+]$.



15 **C-2a**


Compound C-2a was obtained in a procedure similar to that exemplified for **Compound C-1a** but using the methionine ester.

^1H NMR (400 MHz, CDCl_3) δ 8.19 (s, 2 H), 7.44 – 7.03 (m, 7 H), 4.11 (s, 2 H), 3.81 (d, $J = 44.5$ Hz, 1H), 2.04 (s, 3 H), 1.61 (s, 2 H), 1.21 (d, $J = 6.1$ Hz, 2 H), 1.01 – 0.65 (m, 4 H).

^{31}P NMR (162 MHz, CDCl_3) δ -2.00 (d, $J = 12.9$ Hz).

LC MS m/z 455.03 $[\text{M} + \text{H}^+]$.

Compound C-2



5

C-2

Compound C-2 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-2a**.


¹H NMR (400 MHz, CDCl₃) δ 7.96 (d, *J* = 15.8 Hz, 1H), 7.40 – 7.06 (m, 13H), 6.93 (d, *J* = 6.7 Hz, 1H), 6.70 (s, 1H), 5.98 (s, 1H), 4.54 (dd, *J* = 21.6, 11.7 Hz, 2H), 4.32 (d, *J* = 12.0 Hz, 2H), 4.14 (dt, *J* = 13.0, 6.4 Hz, 4H), 2.44 (d, *J* = 7.5 Hz, 2H), 2.00 (d, *J* = 16.2 Hz, 5H), 1.89 (s, 2H), 1.35 – 1.13 (m, 7H).

³¹P NMR (162 MHz, CDCl₃) δ 4.12, 3.58.

¹⁹F NMR (376 MHz, CDCl₃) δ -152.28 (s).

LC MS *m/z* 623.27 [M + H⁺].

15

**C-3a**

Compound C-3a was obtained in a procedure similar to that exemplified for **Compound C-1a** but using a tryptophan ester.

¹H NMR (400 MHz, CDCl₃) δ 8.18 - 8.03 (m, 3 H), 7.29 – 7.08 (m, 8 H), 7.36 – 6.98 (m, 3 H), 4.41 - 4.11 (m, 1 H), 4.15 – 3.95 (m, 2 H), 3.68 – 3.80 (m, 1 H), 3.33 – 3.04 (m, 2 H), 1.06 - 1.17 (m, 3 H).


³¹P NMR (162 MHz, CDCl₃) δ -2.87, -2.99.

LC MS *m/z* 510.03 [M + H⁺].

25

Compound C-3

5

**C-3**


Compound C-3 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-3a**.

10 ^1H NMR (400 MHz, CDCl_3) δ 8.27 (s, 1H), 7.84 (s, 1H), 7.47 (s, 1H), 7.36 – 6.77 (m, 11 H), 6.57 (s, 1 H), 4.40 – 3.96 (m, 6 H), 3.20 (s, 4 H), 2.60 (s, 1H), 1.30 – 1.04 (m, 6 H).

^{31}P NMR (162 MHz, CDCl_3) δ 4.02, 3.75

^{19}F NMR (376 MHz, CDCl_3) δ -152.13.

15 LC MS m/z 678.32 $[\text{M} + \text{H}^+]$.

**C-4a**


Compound C-4a was obtained in a procedure similar to that exemplified for **Compound C-1a** by substituting the phenylalanine ester.

20 ^1H NMR (400 MHz, CDCl_3) δ 8.15 (t, $J = 8.7$ Hz, 2H), 7.43 – 7.11 (m, 10 H), 7.04 (ddd, $J = 11.4, 6.7, 2.9$ Hz, 2 H), 4.32 (ddd, $J = 15.3, 11.3, 6.1$ Hz, 4 H), 4.15 – 3.99 (m, 7 H), 3.74 (td, $J = 11.0, 5.0$ Hz, 8 H), 3.01 (d, $J = 5.7$ Hz, 2 H), 1.17 (td, $J = 7.1, 5.2$ Hz, 2 H).

25 ^{31}P NMR (162 MHz, CDCl_3) δ -2.97, -2.99.

5 LC MS m/z 471.03 [M + H⁺].

Compound C-4



10 **C-4**


Compound C-4 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-4a**.

¹H NMR (400 MHz, CDCl₃) δ 7.92 (d, *J* = 13.2 Hz, 1H), 7.46 – 6.97 (m, 17H), 6.91 (s, 1H), 6.75 (s, 1H), 4.10 (dd, *J* = 29.6, 19.2 Hz, 8H), 2.97 (s, 3H), 1.32 – 1.05 (m, 15 7H).

³¹P NMR (162 MHz, CDCl₃) δ 5.11.

¹⁹F NMR (376 MHz, CDCl₃) δ -152.34 (s).

LC MS m/z 639.24 [M + H⁺].



20 **C-5a**


Compound C-5a was obtained in a procedure similar to that exemplified for **Compound C-1a** but using the proline ester.

¹H NMR (400 MHz, CDCl₃) δ 8.20 (d, *J* = 7.8 Hz, 2 H), 7.45 – 7.08 (m, 7 H), 4.37 (td, *J* = 8.0, 3.8 Hz, 2 H), 4.17 – 3.98 (m, 2 H), 3.61 – 3.34 (m, 2 H), 2.21 – 1.77 (m, 3 25 H), 1.19 (td, *J* = 7.1, 3.8 Hz, 3 H).

5 ^{31}P NMR (162 MHz, CDCl_3) δ -3.92, -3.96.

LC MS m/z 420.98 $[\text{M} + \text{H}^+]$.

Compound C-5



C-5


Compound C-5 was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and **C-5a**.

15 ^1H NMR (400 MHz, CDCl_3) δ 7.95 (d, $J = 4.5$ Hz, 1 H), 7.39 – 7.10 (m, 4 H), 6.92 (dd, $J = 16.0, 4.6$ Hz, 1 H), 6.69 (s, 1H), 6.03 (bs, 2 H), 4.46 – 4.36 (m, 1 H), 4.36 – 3.96 (m, 4 H), 3.37 (d, $J = 58.9$ Hz, 2 H), 2.26 – 1.66 (m, 4 H), 1.39 – 1.12 (m, 8 H).

^{31}P NMR (162 MHz, CDCl_3) δ 3.47, 2.75.

^{19}F NMR (376 MHz, CDCl_3) δ -152.36.

LC MS m/z 589.14 $[\text{M} + \text{H}^+]$.




C-6

25 **Compound C-6** was obtained in a procedure similar to that exemplified for **Compound C-1** using **Compound 3** and the sulphone analog of **C-1a**.

- 5 ^1H NMR (400 MHz, CDCl_3) δ 7.93 (s, 1 H), 7.89 (s, 1 H), 7.35 – 7.01 (m, 5 H), 6.93 (d, $J = 2.8$ Hz, 1 H), 6.58 (d, $J = 2.8$ Hz, 1H), 5.79 (bs, 2 H), 4.30 (s, 6 H), 4.11 (d, $J = 7.0$ Hz, 6H), 3.10 – 2.84 (m, 3 H), 2.75 (s, 3 H), 2.54 (s, 6 H), 1.31 -1.15 (m, 6 H).
 ^{31}P NMR (162 MHz, CDCl_3) δ 3.39, 3.33.
 ^{19}F NMR (376 MHz, CDCl_3) δ -152.40
 10 LC MS m/z 655.24 $[\text{M} + \text{H}^+]$.

Compound 6




15

Compound 4 (about 0.04 mmol) and anhydrous MeOH (about 5 mL) is treated with acetic acid (about 5 mL) and the reaction is stirred overnight at room temperature. Saturated NaHCO_3 is added to neutralize the reaction mixture and the crude material is purified using a HPLC system (acetonitrile- H_2O) to give **6**.

20

Compound 7




25

To a dry, argon purged round bottom flask (50 mL) is added compound **3b** (about 0.39 mmol) and anhydrous dichloromethane (about 10 mL). The flask is


5 placed into a dry ice/acetone bath ($\sim -78^{\circ}\text{C}$) and the solution is stirred for about 10 min. $\text{BF}_3\cdot\text{Et}_2\text{O}$ (about 0.10 mL) is added dropwise and the reaction is stirred for about 10 min. AlMe_3 (about 1.16 mmol, 2.0 M in toluene) is then added. After a few minutes, the dry ice/acetone bath is removed and the reaction mixture is stirred at room temperature to about 45°C over about 4 h to about 4 d. A solution of pyridine
 10 (about 2 mL) in MeOH (about 10 mL) is added and the solvent is removed under reduced pressure. The crude material is purified by chromatography and is treated with ammonium hydroxide in methanol for about 16 h at about room temperature. The mixture is concentrated and the residue is purified by HPLC to give 7.

15 Compound 8



Raney Ni (about 500 mg) is neutralized by washing with H_2O , and added to a
 20 solution of **1d** (about 100 mg) in ethanol (about 10 mL). The mixture is then heated to about 35 to about 80°C until the reaction is complete. The catalyst is removed by filtration and the solution is concentrated *in vacuo*. The mixture is concentrated and the residue is purified by HPLC to give **8**.

25 Compound 9




5

9

Into a flask containing **Compound 3** (120 mg, 0.39 mmol, 1 equiv.) was added PO(OMe)₃ (1.5 mL, 0.25 M) and cooled to 0 °C before adding POCl₃ (125 µL, 1/37 mmol, 3.5 equiv.). The reaction mixture was allowed to stir for 5 hr before the reaction was quenched with water. It was directly purified by HPLC to furnish the

10 monophosphate **Compound 9**.
LC MS m/z 387.95 [M + H⁺].

Compound 10

15

10

Into a flask containing **Compound 9** (30 mg, 0.078 mmol, 1 equiv.) was added NMP (0.8 mL, 0.1 M) followed by addition of TEA (43 µL, 0.31 mmol, 4 equiv.), tetrabutylammonium bromide (25 mg, 0.078 mmol, 1 equiv.) before adding chloromethylisopropyl carbonate (60 µL, 0.38 mmol, 5 equiv.). The reaction

20 mixture was heated to 50 °C and allowed to stir overnight. It was purified directly by HPLC, affording **Compound 10**.


¹H NMR (400 MHz, CDCl₃) δ 7.98 (s, 1 H), 7.01 (d, *J* = 4.7 Hz, 1 H), 6.72 (d, *J* = 4.7 Hz, 1 H), 6.04 (bs, 2 H), 5.74 – 5.61 (m, 4 H), 4.91 (ddt, *J* = 12.6, 9.4, 6.3 Hz, 2 H), 4.64 – 4.28 (m, 4 H), 1.37 – 1.19 (m, 15 H).

25 ³¹P NMR (162 MHz, CDCl₃) δ -4.06.


¹⁹F NMR (376 MHz, CDCl₃) δ -76.58, -151.95 TFA salt.

LC MS m/z 620.03[M + H⁺].

Compound 11

**11**


A solution of **Compound B-2** in DMSO is treated with about 3 mole equivalents of potassium *t*-butoxide for about 15 min to 24 hours. The reaction is quenched with 1N HCl and **Compound 11** is isolated by reverse-phase HPLC.

Compound 12

Compound 1b (about 1 mmol) is placed in a steel bomb reactor. The reactor is charged with liquid ammonia (about 30 mL) and the mixture is stirred at about 0 to 50 °C for about 16 h. The ammonia is evaporated and the residue is purified to give **12a**. A solution of **12a** (about 100 mg) in ethanol (about 10 ml) is treated with Raney Ni (about 500 mg) that is neutralized by washing with H₂O. The mixture is then

5 heated to about 35 to about 80 °C until the reaction is complete. The catalyst is removed by filtration and the solution is concentrated *in vacuo*. The mixture is concentrated and the residue is purified by HPLC to give **12b**. To a solution of compound **12b** (about 50 mg) and TMSCN (about 0.5 mmol) in acetonitrile (about 2.0 mL) at about 0 °C is added TMSOTf (about 0.5 mmol). The reaction mixture is
10 stirred at room temperature for about 1 h, then at 65 °C for about 3 d. The reaction is quenched with saturated NaHCO₃ at room temperature, and diluted with CH₃CO₂Et. The organic phase was separated, washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue is purified by RP-HPLC then dissolved in methanol (about 1 mL). Ammonium hydroxide (28% in water, about 0.8 mL) is added and the
15 mixture is stirred at about room temperature for 16 h. The mixture is concentrated and the residue is purified by RP HPLC to give **12**.


Compound 13



20 **13**

Compound 13 is prepared in the same manner as **Compound 9** using **Compound 12** as a starting material.

Compound 14




25 **14**

Compound 14 is prepared by treating **Compound 13** with about one to about five equivalents of DCC in pyridine and heating the reaction to reflux for about one to

5 about 24 hours. **Compound 14** is isolated by conventional ion exchange and reverse-phase HPLC.

Compound 15




10

15

A solution of about 0.4 mmol of **Compound 14** in about 10 mL of DMF is treated with about 0.8 mmol of DIPEA and about 0.8 mmol of chloromethyl isopropyl carbonate (WO2007/027248). The reaction is heated to about 25 to about 80 °C for about 15 min to about 24 hours. The solvent is removed under vacuum and the residue is purified by HPLC to give **Compound 15**.

15

Compound 16




16

20 Compound **3** (about 0.22 mmol) is dissolved in anhydrous pyridine (about 2 mL) and chlorotrimethylsilane (about 0.17 mL) is added. The mixture is stirred at about 0 to about 25 °C for about one to about 24 hours. Additional chlorotrimethylsilane (about 0.1 mL) is added and the reaction is stirred for about one to about 24 hours. 4,4'-Dimethoxytrityl chloride (about 0.66 mmol) and DMAP

25 (about 0.11 to about 0.22 mmol) is sequentially added. The mixture is stirred for about one to about 24 hours. A solution of TBAF (1.0 M, about 0.22 mL) in THF is added and the reaction is stirred for about one to about 24 hours. The mixture is

5 partitioned between ethyl acetate and water. The ethyl acetate layer is dried and concentrated. The residue is purified chromatography to afford **Compound 16**.

Compound 17




17

10

A mixture of about 1.25 mmol of **Compound 16** and about 1.9 mmol of triethylammonium 2-(2,2-dimethyl-3-(trityloxy)propanoylthio)ethyl phosphinate (WO2008082601) is dissolved in anhydrous pyridine (about 19 mL). Pivaloyl chloride (about 2.5 mmol) is added dropwise at about -30 to about 0 °C and the solution is stirred at for about 30 min to about 24 hours. The reaction is diluted with methylene chloride and is neutralized with aqueous ammonium chloride (about 0.5M). The methylene chloride phase is evaporated and the residue is dried and is purified by chromatography to give **Compound 17**.

Compound 18




18

To a solution of about 0.49 mmol of **Compound 17** in anhydrous carbon tetrachloride (about 5 mL) is added dropwise benzylamine (about 2.45 mmol). The reaction mixture is stirred for about one to about 24 hours. The solvent is evaporated and the residue is purified by chromatography to give **Compound 18**.


25

5

Compound 20

10 A solution of about 2 mmol of **Compound 18** in methylene chloride (about 10 mL) is treated with an aqueous solution of trifluoroacetic acid (90%, about 10 mL). The reaction mixture is stirred at about 25 to about 60 °C for about one to about 24 hours. The reaction mixture is diluted with ethanol, the volatiles are evaporated and the residue is purified by chromatography to give **Compound 20**.

15


Compound 21

20 About 90 mM **Compound 2** in THF is cooled to about -78 °C and about 2.2 to about 5 equivalents of *t*-butylmagnesium chloride (about 1 M in THF) is added. The mixture is warmed to about 0 °C for about 30 min and is again cooled to about -78 °C. A solution of (2*S*)-2-{[chloro(1-phenoxy)phosphoryl]amino}propyl pivaloate (WO2008085508) (1 M in THF, about 2 equivalents) is added dropwise. The cooling
25 is removed and the reaction is stirred for about one to about 24 hours. The reaction is quenched with water and the mixture is extracted with ethyl acetate. The extracts are

5 dried and evaporated and the residue purified by chromatography to give **Compound 21**.

Compound 22

10



22a


Compound 22a was obtained in a procedure similar to that for preparation of **C-1a**.

¹H NMR (400 MHz, CDCl₃) δ 8.11 (d, *J* = 9.0 Hz, 2 H), 8.02 (s, 1 H), 7.48 (t, *J* = 7.5 Hz, 2 H), 7.42 – 7.25 (m, 4 H), 7.21 (dt, *J* = 14.9, 5.5 Hz, 2 H), 7.08 (t, *J* = 7.3 Hz, 2 H), 5.17 – 5.03 (m, 2 H), 4.99 (dd, *J* = 16.5, 9.7 Hz, 2 H), 3.44 (s, 1H), 3.35 – 3.21 (m, 2 H), 3.19 (d, *J* = 9.2 Hz, 1H), 3.00 – 2.80 (m, 2 H).

³¹P NMR (162 MHz, CDCl₃) δ 4.27.

LC MS *m/z* 452.09 [M + H⁺].

20



22b

Compound 22b was obtained in a procedure similar to that for preparation of **C-1** using **Compound 3** and **22a**.


¹H NMR (400 MHz, CD₃OD) δ 7.76 (d, *J* = 6.3 Hz, 1H), 7.38 (t, *J* = 8.2 Hz, 1 H), 7.27 – 7.12 (m, 4 H), 7.06 – 6.81 (m, 3 H), 6.74 (dd, *J* = 4.6, 3.5 Hz, 1 H), 4.95 – 4.79

5 (m, 1 H), 4.35 – 3.90 (m, 4 H), 3.23 (dt, $J = 3.2, 1.6$ Hz, 3H), 3.18 – 3.05 (m, 2 H),
2.82 (dt, $J = 14.7, 7.3$ Hz, 2 H), 1.15 (d, $J = 22.4$ Hz, 3 H).

^{31}P NMR (162 MHz, CD_3OD) δ 10.76, 10.71.

LC MS m/z 620.05 $[\text{M} + \text{H}^+]$.

10



Compound 22

15 Into a flask containing the **22b** (50 mg, 0.08 mmol, 1 equiv.) was added
ethanol (4 mL) followed by $\text{Pd}(\text{OH})_2$ (56 mg, 0.08 mmol, 1 equiv.) and ammonium
formate (42 mg, 0.64 mmol, 8 equiv.). The reaction was heated to 80 °C for about an
hour. The solid was filtered off and the material purified by HPLC.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.72 (s, 1H), 7.91 (s, 1 H), 7.95 - 7.89 (bs, 2 H),
7.41 (d, $J = 7.7$ Hz, 1 H), 7.26 (d, $J = 8.1$ Hz, 1 H), 7.19 – 6.66 (m, 3 H), 4.20 – 3.75
20 (m, 3 H), 2.99 (dd, $J = 16.5, 9.6$ Hz, 2 H), 2.89 – 2.70 (m, 2 H), 2.48 – 2.58 (m, 8 H),
1.10 (d, $J = 22.3$ Hz, 3 H).


^{31}P NMR (162 MHz, $\text{DMSO}-d_6$) δ 7.49.

^{19}F NMR (376 MHz, $\text{DMSO}-d_6$) δ -154.89.

LC MS m/z 530.21 $[\text{M} + \text{H}^+]$.

25

Compound 23



Compound **3** (250mg, 0.82mmol) was dissolved in PO(OMe)₃ (5 mL, 0.16M) and cooled to 0°C under argon. To this stirring solution was added POCl₃ (0.32 mL, 4.1 mmol) slowly dropwise, and the reaction mixture allowed to warm to room temperature for 16 h. The resulting solution was added dropwise to a rapidly stirring solution of acetonitrile (400 mL) and 0.08M aqueous KOH (300 mL). When addition was complete, the reaction progress was checked by LCMS. When the reaction was complete, solvents were removed under reduced pressure. The resulting solid residue was dissolved in water and purified by HPLC to give 140mg of **Compound 23** (yield 47%).


¹H-NMR (400MHz; CD₃OD) : d 8.15 (s, 1H), 7.40 (d, 1H; J = 4.8Hz), 7.09 (d, 1H; J = 4.8Hz), 4.64 (dd, 1H; J = 24Hz, 7.2Hz), 4.50-4.36 (m, 3H), 1.32 (d, 3H; J=22Hz).

¹⁹F-NMR (376MHz; CD₃OD) : d -153.11.

20 ³¹P-NMR (162MHz; CD₃OD) : d -2.20.


MS [M + H⁺] = 370.2.

Compound 24



- 5 A solution of **Compound 23** (7mg, 0.02 mmol) in DCM (2 mL) and PO(OMe)₃ (1 mL) was prepared and cooled to 0°C. To this solution was added oxalyl-Cl (10 uL) followed by DMF (2 uL). The mixture was allowed to stir for 1 min before an aliquot was taken out and quenched in MeOH and then checked by LCMS for activation. Successive amounts of oxalyl-Cl (10 uL) and DMF (2 uL) were
- 10 added until activation was complete. At this point, a large volume of 2-propanol (5 mL) was added to the reaction mixture and allowed to stir and warm to room temperature. Once the reaction was complete, the solvents were removed under reduced pressure, and the resulting crude material was purified by preparative HPLC to give 5.5 mg of **Compound 24** (yield 70%).
- 15 ¹H-NMR (400MHz; DMSO-d₆) : d 8.26 (br, 1H), 8.15 (br, 1H), 7.97 (s, 1H), 7.00 (d, 1H; J = 4.4Hz), 6.88 (d, 1H; J = 4.4Hz), 4.59-4.51 (m, 2H), 4.37-4.25 (m, 2H), 1.23 (d, 3H; J=22.8Hz).
- ¹⁹F-NMR (376MHz; CD₃OD) : d -151.72.
- ³¹P-NMR (162MHz; CD₃OD) : d -5.69.
- 20 MS [M + H⁺] = 412.0.

Compound 25



25

Compound 25 was prepared from **Compound 23** in a matter similar to that of **Compound 24** substituting the heptyl ester of alanine for 2-propanol (yield 5.3%).

¹H-NMR (400MHz; CD₃OD) : d 7.91 (s, 1H), 6.98 (d, 1H; J = 4.8Hz), 6.92 (d, 1H; J = 4.8Hz), 5.29 (dd, 1H; J = 24.4Hz, 8.8Hz), 4.66-4.60 (m, 2H), 4.48-4.40 (m, 1H),

5 4.15-4.11(m, 3H), 3.92(dd, 1H; J = 9.6Hz, 7.2Hz), 1.67-1.64 (m, 3H), 1.40-1.27 (m, 15H), 0.91-0.87(m, 6H).


^{19}F -NMR (376MHz; CD_3OD) : d -151.46.

^{31}P -NMR (162MHz; CD_3OD) : d 7.36.

MS $[\text{M} + \text{H}^+] = 539.4$.

10

Compound 26



Compound 26

15

Compound 26 is prepared from compound **22** in a matter similar to that for preparation of compound **10**.

Antiviral Activity

20

Another aspect of the invention relates to methods of inhibiting viral infections, comprising the step of treating a sample or subject suspected of needing such inhibition with a composition of the invention.

25

Within the context of the invention samples suspected of containing a virus include natural or man-made materials such as living organisms; tissue or cell cultures; biological samples such as biological material samples (blood, serum, urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like. Typically the sample will be suspected of containing an organism which induces a

5 viral infection, frequently a pathogenic organism such as a tumor virus. Samples can be contained in any medium including water and organic solvent/water mixtures. Samples include living organisms such as humans, and man made materials such as cell cultures.

If desired, the anti-virus activity of a compound of the invention after
10 application of the composition can be observed by any method including direct and indirect methods of detecting such activity. Quantitative, qualitative, and semiquantitative methods of determining such activity are all contemplated. Typically one of the screening methods described above are applied, however, any other method such as observation of the physiological properties of a living organism
15 are also applicable.

The antiviral activity of a compound of the invention can be measured using standard screening protocols that are known. For example, the antiviral activity of a compound can be measured using the following general protocols.

Cell-based Flavivirus Immunodetection assay

20 BHK21 or A549 cells are trypsinized, counted and diluted to 2×10^5 cells/mL in Hams F-12 media (A549 cells) or RPMI-1640 media (BHK21 cells) supplemented with 2% fetal bovine serum (FBS) and 1% penicillin/streptomycin. 2×10^4 cells are dispensed in a clear 96-well tissue culture plates per well and palced at 37° C, 5% CO₂ overnight. On the next day, the cells are infected with viruses at multiplicity of
25 infection (MOI) of 0.3 in the presence of varied concentrations of test compounds for 1 hour at 37° C and 5% CO₂ for another 48 hours. The cells are washed once with PBS and fixed with cold methanol for 10 min. After washing twice with PBS, the fixed cells are blocked with PBS containing 1% FBS and 0.05% Tween-20 for 1 hour at room temperature. The primary antibody solution (4G2) is then added at a
30 concentration of 1:20 to 1:100 in PBS containing 1% FBS and 0.05% Tween-20 for 3 hours. The cells are then washed three times with PBS followed by one hour incubation with horseradish peroxidase(HRP)-conjugated anti-mouse IgG (Sigma, 1:2000 dilution). After washing three times with PBS, 50 microliters of 3,3',5,5'-

5 tetramethylbenzidine (TMB) substrate solution (Sigma) is added to each well for two minutes. The reaction is stopped by addition of 0.5 M sulfuric acid. The plates are read at 450 nm absorbance for viral load quantification. After measurement, the cells are washed three times with PBS followed by incubation with propidium iodide for 5 min. The plate is read in a Tecan Safire™ reader (excitation 537 nm, emission
10 617 nm) for cell number quantification. Dose response curves are plotted from the mean absorbance versus the log of the concentration of test compounds. The EC₅₀ is calculated by non-linear regression analysis. A positive control such as N-nonyl-deoxynojirimycin may be used.

Cell-based Flavivirus cytopathic effect assay

15 For testing against West Nile virus or Japanese encephalitis virus, BHK21 cells are trypsinized and diluted to a concentration of 4×10^5 cells/mL in RPMI-1640 media supplemented with 2% FBS and 1% penicillin/streptomycin. For testing against dengue virus, Huh7 cells are trypsinized and diluted to a concentration of 4×10^5 cells/mL in DMEM media supplemented with 5% FBS and 1%
20 penicillin/streptomycin. A 50 microliter of cell suspension (2×10^4 cells) is dispensed per well in a 96-well optical bottom PIT polymer-based plates (Nunc). Cells are grown overnight in culture medium at 37°C, 5% CO₂, and then infected with West Nile virus (e.g. B956 strain) or Japanese encephalitis virus (e.g. Nakayama strain) at MOI = 0.3, or with dengue virus (e.g. DEN-2 NGC strain) at MOI = 1, in the presence
25 of different concentrations of test compounds. The plates containing the virus and the compounds are further incubated at 37°C, 5% CO₂ for 72 hours. At the end of incubation, 100 microliters of CellTiter-Glo™ reagent is added into each well. Contents are mixed for 2 minutes on an orbital shaker to induce cell lysis. The plates are incubated at room temperature for 10 minutes to stabilize luminescent signal.
30 Luminescence reading is recorded using a plate reader. A positive control such as N-nonyl-deoxynojirimycin may be used.

Antiviral Activity in a Mouse Model of Dengue Infection.

Compounds are tested *in vivo* in a mouse model of dengue virus infection

5 (Schul *et al.* J. Infectious Dis. 2007; 195:665-74). Six to ten week old AG129 mice (B&K Universal Ltd, Hill, UK) are housed in individually ventilated cages. Mice are injected intraperitoneally with 0.4 mL TSV01 dengue virus 2 suspension. Blood samples are taken by retro orbital puncture under isoflurane anaesthesia. Blood samples are collected in tubes containing sodium citrate to a final concentration of
10 0.4%, and immediately centrifuged for 3 minutes at 6000g to obtain plasma. Plasma (20 microliters) is diluted in 780 microliters RPMI-1640 medium and snap frozen in liquid nitrogen for plaque assay analysis. The remaining plasma is reserved for cytokine and NS1 protein level determination. Mice develop dengue viremia rising over several days, peaking on day 3 post-infection.

15 For testing of antiviral activity, a compound of the invention is dissolved in vehicle fluid, e.g. 10% ethanol, 30% PEG 300 and 60% D5W (5% dextrose in water; or 6N HCl (1.5 eq):1N NaOH (pH adjusted to 3.5): 100 mM citrate buffer pH 3.5 (0.9% v/v:2.5% v/v: 96.6% v/v). Thirty six 6-10 week old AG129 mice are divided into six groups of six mice each. All mice are infected with dengue virus as described
20 above (day 0). Group 1 is dosed by oral gavage of 200 μ L/mouse with 0.2 mg/kg of a compound of the invention twice a day (once early in the morning and once late in the afternoon) for three consecutive days starting on day 0 (first dose just before dengue infection). Groups 2, 3 and 4 are dosed the same way with 1 mg/kg, 5 mg/kg and 25 mg/kg of the compound, respectively. A positive control may be used, such as
25 (2R,3R,4R,5R)-2-(2-amino-6-hydroxy-purin-9-yl)-5-hydroxymethyl-3-methyl-tetrahydro-furan-3,4-diol, dosed by oral gavage of 200 microliters/mouse the same way as the previous groups. A further group is treated with only vehicle fluid.

On day 3 post-infection approximately 100 microliter blood samples (anti-coagulated with sodium citrate) are taken from the mice by retro-orbital puncture
30 under isoflurane anaesthesia. Plasma is obtained from each blood sample by centrifugation and snap frozen in liquid nitrogen for plaque assay analysis. The collected plasma samples are analyzed by plaque assay as described in Schul *et al.* Cytokines are also analysed as as described by Schul. NS1 protein levels are analysed using a PlateliaTM kit (BioRad Laboratories). An anti-viral effect is indicated by a

- 5 reduction in cytokine levels and/or NS1 protein levels.

Typically, reductions in viremia of about 5-100 fold, more typically 10-60 fold, most typically 20-30 fold, are obtained with 5-50 mg/kg bid dosages of the compounds of the invention.





10 HCV IC₅₀ Determination


Assay Protocol: Either wild type or S282T (Migliaccio, et al, *J. Biol. Chem.* **2003**, 49164-49170; Klumpp, et al., *J. Biol. Chem.* **2006**, 3793-3799) mutant polymerase enzyme was used in this assay. NS5b polymerase assay (40 μ L) was
15 assembled by adding 28 μ L polymerase mixture (final concentration: 50 mM Tris-HCl at pH 7.5, 10 mM KCL, 5 mM MgCl₂, 1 mM DTT, 10 mM EDTA, 4 ng/ μ L of RNA template, and 75 nM HCV Δ 21 NS5b polymerase) to assay plates followed by 4 μ L of compound dilution. The polymerase and compound were pre-incubated at 35 °C for 10 minute before the addition of 8 μ L of nucleotide substrate mixture (33P- α -
20 labeled competing nucleotide at K_M and 0.5 mM of the remaining three nucleotides). The assay plates were covered and incubated at 35 °C for 90 min. Reactions were then filtered through 96-well DEAE-81 filter plates via vacuum. The filter plates were then washed under vacuum with multiple volumes of 0.125 M NaHPO₄, water, and ethanol to remove unincorporated label. Plates were then counted on TopCount to assess the
25 level of product synthesis over background controls. The IC₅₀ value is determined using Prism fitting program.

Preferably, compounds described herein inhibited NS5b polymerase with an IC₅₀'s below 1000 μ M, more preferably below 100 μ M, and most preferably below 10 μ M. For example, compound **TP-1** has an IC₅₀ of 0.15 μ M against both wild type
30 HCV polymerase and the S282T mutant enzyme. Table II below shows the activity of **TP-1** and **TP-2** against both wild type and the S282T mutant enzyme compared to the activities obtained with the triphosphate of 2'-methyl guanidine and the triphosphate of (2R,3R,4R,5R)-2-(4-aminopyrrolo[1,2-f][1,2,4]triazin-7-yl)-3,4-

- 5 dihydroxy-5-(hydroxymethyl)-3-methyl-tetrahydrofuran-2-carbonitrile. This demonstrates that replacing the 2' OH of the pyrrolo[1,2-f][1,2,4]triazin-7-yl nucleosides with a 2' F unexpectedly confers activity against resistant S282T HCV mutant strains of virus.

10 Table II

Triphosphate	WT IC50(uM)	S282T IC50(uM)	Note
 2'-C-MeGTP	0.1	20	from <i>J. Bio. Chem.</i> , 2003 , 278, 49164 (200 fold shift)
 TP-1	0.15	0.15	(1 fold shift)
	0.525	111	WO/2009/132135 (242 fold shift)
 TP-2	0.24	1.60	(7 fold shift)



5

HCV EC₅₀ Determination

Replicon cells were seeded in 96-well plates at a density of 8×10^3 cells per well in 100 μ L of culture medium, excluding Geneticin. Compound was serially diluted in 100% DMSO and then added to the cells at a 1:200 dilution, achieving a final concentration of 0.5% DMSO and a total volume of 200 μ L. Plates were incubated at 37°C for 3 days, after which culture medium was removed and cells were lysed in lysis buffer provided by Promega's luciferase assay system. Following the manufacturer's instruction, 100 μ L of luciferase substrate was added to the lysed cells and luciferase activity was measured in a TopCount luminometer. Preferably, compounds described herein have EC₅₀'s below 1000 μ M, more preferably below 100 μ M, and most preferably below 10 μ M. The activities of representative compounds of Formula I are shown in the Table below.

20

Compound No.	EC ₅₀ , μ M
A-1	23
B-1	1.4-4.3
B-3	16-28
B-4	8.4-19
B-5	1.93-25.5
B-6	3.75-11.1
B-7	63-73
B-8	35-60

C-1	67-70
C-2	3.9-12
C-3	43-84
C-4	9.8-31
C-5	24-28
C-6	11
10	6.5-8
22	31-45
23	39.4-40.3
24	40.3-70.5
25	9.7-10

5

The cytotoxicity of a compound of the invention can be determined using the following general protocol.

Cytotoxicity Cell Culture Assay (Determination of CC50):

10 The assay is based on the evaluation of cytotoxic effect of tested compounds using a metabolic substrate.

Assay protocol for determination of CC50:

1. Maintain MT-2 cells in RPMI-1640 medium supplemented with 5% fetal bovine serum and antibiotics.
- 15 2. Distribute the cells into a 96-well plate (20,000 cell in 100 μ l media per well) and add various concentrations of the tested compound in triplicate (100 μ l/well). Include untreated control.
3. Incubate the cells for 5 days at 37 °C.
4. Prepare XTT solution (6 ml per assay plate) in dark at a concentration of 2mg/ml
- 20 in a phosphate-buffered saline pH 7.4. Heat the solution in a water-bath at 55°C for 5 min. Add 50 μ l of N-methylphenazonium methasulfate (5 μ g/ml) per 6 ml of XTT solution.


5

5. Remove 100 μ l media from each well on the assay plate and add 100 μ l of the XTT substrate solution per well. Incubate at 37 °C for 45 to 60 min in a CO₂ incubator.
6. Add 20 μ l of 2% Triton X-100 per well to stop the metabolic conversion of XTT.
7. Read the absorbance at 450 nm with subtracting off the background at 650 nm.
- 10 8. Plot the percentage absorbance relative to untreated control and estimate the CC50 value as drug concentration resulting in a 50% inhibition of the cell growth. Consider the absorbance being directly proportional to the cell growth.

The scope of the claims should not be limited by the embodiments set forth in the examples,
15 but should be given the broadest interpretation consistent with the description as a whole.

5 WHAT IS CLAIMED IS:

1. A compound of Formula II:




Formula II


- 10 or a pharmaceutically acceptable salt thereof;

wherein:

- R^1 is (C_1-C_8) alkyl;
 R^3 and R^5 are each H;
 R^4 is OR^a ;
 15 R^6 is H, OR^a , CN or (C_1-C_8) alkyl;
 each R^a is independently H or (C_1-C_8) alkyl;
 R^7 is H or



- each Y and Y^1 is O;
 20 one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is
 Formula Ia; or W^1 and W^2 are each, independently, a group of the Formula Ia:



Formula Ia


wherein:

each Y^2 is independently O or NR;

each Y^3 is O;

M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



wherein:

each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, R, $-\text{C}(=\text{Y}^1)\text{OR}$, $-\text{OC}(=\text{Y}^1)\text{OR}$, or $-\text{SC}(=\text{Y}^1)\text{R}$; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, $(\text{C}_1\text{--}\text{C}_8)$ alkyl, $(\text{C}_1\text{--}\text{C}_8)$ substituted alkyl, $\text{C}_6\text{--}\text{C}_{20}$ aryl,

$\text{C}_6\text{--}\text{C}_{20}$ substituted aryl, $\text{C}_2\text{--}\text{C}_{20}$ heterocyclyl, or arylalkyl;

X^1 is C-R^{10} or N;

X^2 is C-R^{10} ;

R^8 is halogen, $\text{NR}^{11}\text{R}^{12}$, or OR^{11} ;

R^9 is H, $\text{NR}^{11}\text{R}^{12}$, or SR^{11} ;

5 each R¹⁰ is H; and
 each R¹¹ and R¹² is independently H, or (C₁-C₈)alkyl, wherein each (C₁-C₈)alkyl of
each R¹, R⁶, R¹¹, R¹², R^a and R is a hydrocarbon containing normal, secondary or tertiary
carbon atoms.


10 2. The compound according to claim 1, wherein R⁶ is H, OR^a, CN, or methyl.

3. The compound according to claim 1 or 2, wherein R⁶ is H.

4. The compound according to claim 1 or 2, wherein R⁶ is CN.

15


5. The compound according to any one of claims 1-4, which is represented by
Formula III:



Formula III

20 wherein R¹ is methyl; or a pharmaceutically acceptable salt thereof.


6. The compound according to any one of claims 1-5, wherein R⁷ is



5

7. The compound according to any one of claims 1-4, wherein R^1 is CH_3 .

8. A compound of Formula I:



10

Formula I

or a pharmaceutically acceptable salt or ester thereof;


wherein:

R^1 is (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl,
 15 (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or
 aryl (C_1-C_8) alkyl;


R^2 is F;

each R^3 and R^5 is independently H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, halogen,
 (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl,
 20 (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl (C_1-C_8) alkyl;

- 5 R^4 is H, OR^b , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or aryl(C_1-C_8)alkyl;
- or any two of R^3 , R^4 or R^5 on adjacent carbon atoms when taken together are $-O(CO)O-$ or when taken together with the ring carbon atoms to which they are attached
- 10 form a double bond;
- R^6 is H, OR^a , $N(R^a)_2$, N_3 , CN, NO_2 , $S(O)_nR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_4-C_8) carbocyclalkyl, (C_1-C_8) substituted alkyl, (C_2-C_8) alkenyl, (C_2-C_8) substituted alkenyl, (C_2-C_8) alkynyl, (C_2-C_8) substituted alkynyl, or
- 15 aryl(C_1-C_8)alkyl;
- each n is independently 0, 1, or 2;
- each R^a is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;
- 20 each R^b is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, (C_4-C_8) carbocyclalkyl, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, or $-SO_2NR^{11}R^{12}$;
- R^7 is H, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)R^{11}$, $-S(O)_2R^{11}$, $-S(O)(OR^{11})$, $-S(O)_2(OR^{11})$, $-SO_2NR^{11}R^{12}$, or



- each Y and Y^1 is, independently, O, S, NR, $^+N(O)(R)$, $N(OR)$, $^+N(O)(OR)$, or $N-NR_2$;
- W^1 and W^2 , when taken together, are $-Y^3(C(R^y)_2)_3Y^3-$; or one of W^1 or W^2 together with either R^3 or R^4 is $-Y^3-$ and the other of W^1 or W^2 is Formula Ia; or W^1 and W^2 are each,
- 30 independently, a group of the Formula Ia:



Formula Ia


wherein:

each Y^2 is independently a bond, O, CR_2 , NR, $^+\text{N}(\text{O})(\text{R})$, $\text{N}(\text{OR})$, $^+\text{N}(\text{O})(\text{OR})$, $\text{N}-\text{NR}_2$, S, S-S, S(O), or S(O)₂;

each Y^3 is independently O, S, or NR;

M2 is 0, 1 or 2;

each R^x is independently R^y or the formula:



wherein:


each M1a, M1c, and M1d is independently 0 or 1;

M12c is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12;

each R^y is independently H, F, Cl, Br, I, OH, R, $-\text{C}(=\text{Y}^1)\text{R}$, $-\text{C}(=\text{Y}^1)\text{OR}$, $-\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$, $-\text{N}(\text{R})_2$, $^+\text{N}(\text{R})_3$, $-\text{SR}$, $-\text{S}(\text{O})\text{R}$, $-\text{S}(\text{O})_2\text{R}$, $-\text{S}(\text{O})(\text{OR})$, $-\text{S}(\text{O})_2(\text{OR})$, $-\text{OC}(=\text{Y}^1)\text{R}$, $-\text{OC}(=\text{Y}^1)\text{OR}$, $-\text{OC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{SC}(=\text{Y}^1)\text{R}$, $-\text{SC}(=\text{Y}^1)\text{OR}$, $-\text{SC}(=\text{Y}^1)(\text{N}(\text{R})_2)$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{R}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{OR}$, $-\text{N}(\text{R})\text{C}(=\text{Y}^1)\text{N}(\text{R})_2$, $-\text{SO}_2\text{NR}_2$, $-\text{CN}$, $-\text{N}_3$, $-\text{NO}_2$, $-\text{OR}$, or W^3 ; or when taken together, two R^y on the same carbon atom form a carbocyclic ring of 3 to 7 carbon atoms;

each R is independently H, (C₁-C₈) alkyl, (C₁-C₈) substituted alkyl, (C₂-C₈) alkenyl, (C₂-C₈) substituted alkenyl, (C₂-C₈) alkynyl, (C₂-C₈) substituted alkynyl, C₆-C₂₀ aryl,

- 5 C₆–C₂₀ substituted aryl, C₂–C₂₀ heterocyclyl, C₂–C₂₀ substituted heterocyclyl, arylalkyl or substituted arylalkyl;
- W³ is W⁴ or W⁵; W⁴ is R, -C(Y¹)R^y, -C(Y¹)W⁵, -SO₂R^y, or -SO₂W⁵; and W⁵ is a carbocycle or a heterocycle wherein W⁵ is independently substituted with 0 to 3 R^y groups;
- each X¹ and X² is independently C-R¹⁰ or N;
- 10 each R⁸ is halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NNHR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl, -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl, aryl(C₁-C₈)alkyl, OR¹¹ or SR¹¹;
- 15 each R⁹ and R¹⁰ is independently H, halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², N₃, NO, NO₂, CHO, CN, -CH(=NR¹¹), -CH=NNHR¹¹, -CH=N(OR¹¹), -CH(OR¹¹)₂, -C(=O)NR¹¹R¹², -C(=S)NR¹¹R¹², -C(=O)OR¹¹, R¹¹, OR¹¹ or SR¹¹;
- each R¹¹ and R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₄-C₈)carbocyclalkyl, optionally substituted aryl, optionally substituted heteroaryl,
- 20 -C(=O)(C₁-C₈)alkyl, -S(O)_n(C₁-C₈)alkyl or aryl(C₁-C₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S- or -NR^a;
- wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl or aryl(C₁-C₈)alkyl of
- 25 each R¹, R³, R⁴, R⁵, R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more halo, hydroxy, CN, N₃, N(R^a)₂ or OR^a; and wherein one or more of the non-terminal carbon atoms of each said (C₁-C₈)alkyl is optionally replaced with -O-, -S- or -NR^a; wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl of each R¹, R⁶, R¹¹, R¹², R^a and R is a hydrocarbon containing normal, secondary or tertiary carbon atoms.
- 30
9. A compound according to claim 8, represented by Formula II



Formula II

wherein each Y and Y¹ is O; X¹ is CR¹⁰ or N and X² is CH; or a pharmaceutically acceptable salt or ester thereof.

10

10. A compound according to claim 8 or 9, wherein R⁸ is halogen, NR¹¹R¹², N(R¹¹)OR¹¹, NR¹¹NR¹¹R¹², OR¹¹ or SR¹¹.

11. A compound according to any one of claims 8-10, wherein R⁹ is H, halogen, SR¹¹ or NR¹¹R¹².

15

12. A compound according to any one of claims 8-11, wherein R⁶ is H, OR^a, CN, methyl, ethenyl, or ethynyl.

13. A compound according to any one of claims 8-12, wherein R⁶ is H.


20

14. A compound according to any one of claims 8-12, wherein R⁶ is CN.

15. A compound according to any one of claims 8-14, wherein R⁴ is OR^b.

25


16. A compound according to any one of claims 8-15, represented by Formula III




Formula III

wherein R¹ is methyl or ethynyl; or a pharmaceutically acceptable salt or ester thereof.

17. A compound according to any one of claims 8-16, wherein R⁷ is H or








18. A compound according to any one of claims 8-17, wherein R⁷ is




19. A compound according to any one of claims 8-18, wherein R¹ is CH₃.

20. A compound that is:


5




10




5




5




10




5







5



or a pharmaceutically acceptable salt thereof.

10 22. A compound that is:





or a pharmaceutically acceptable salt thereof.

10 23. A pharmaceutical composition, comprising the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

24. Use of a therapeutically effective amount of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for treating or preventing a *Flaviviridae* virus
15 infection in a mammal in need thereof.

25. Use of a therapeutically effective amount of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for treating a *Flaviviridae* virus infection in a mammal in need thereof.

5

26. Use of a therapeutically effective amount of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for treating or preventing a dengue virus infection in a mammal in need thereof.

10 27. Use of a therapeutically effective amount of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for treating a dengue virus infection in a mammal in need thereof.

15 28. Use of a therapeutically effective amount of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for treating or preventing a Hepatitis C virus infection in a mammal in need thereof.

20 29. Use of a therapeutically effective amount of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for treating a Hepatitis C virus infection in a mammal in need thereof.

30. The use of claim 28 or 29, wherein the Hepatitis C virus infection is caused by a S282T mutant of Hepatitis C virus.

25 31. The use of any one of claims 28-30, further comprising using at least one additional therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, endothelin antagonists, other anti-fibrotic agents, nucleoside or
30 nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.

- 5 32. Use of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a *Flaviviridae* virus infection.
33. Use of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a *Flaviviridae* virus infection.
- 10 34. Use of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a dengue virus infection.
- 15 35. Use of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a dengue virus infection.
- 20 36. Use of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment or prevention of a Hepatitis C virus infection.
- 25 37. Use of the compound of any one of claims 1-22 or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of a Hepatitis C virus infection.
- 30 38. The use of claim 36 or 37, wherein the Hepatitis C virus infection is caused by a S282T mutant of Hepatitis C virus.
39. The use of any one of claims 36-38, wherein the compound is used in combination with at least one additional therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists,

- 5 antagonists of the renin-angiotensin system, endothelin antagonists, other anti-fibrotic agents, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof.
- 10
40. Use of the pharmaceutical composition of claim 23, for treating or preventing a *Flaviviridae* virus infection.
41. Use of the pharmaceutical composition of claim 23, for treating a *Flaviviridae* virus
- 15 infection.
42. Use of the pharmaceutical composition of claim 23, for treating or preventing a dengue virus infection.
- 20 43. Use of the pharmaceutical composition of claim 23, for treating a dengue virus infection.
44. Use of the pharmaceutical composition of claim 23, for treating or preventing a Hepatitis C virus infection.
- 25
45. Use of the pharmaceutical composition of claim 23, for treating a Hepatitis C virus infection.
46. The use of claim 44 or 45, wherein the Hepatitis C virus infection is caused by
- 30 a S282T mutant of Hepatitis C virus.
47. The use of any one of claims 44-46, wherein the pharmaceutical composition further comprises at least one additional therapeutic agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-

- 5 glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists,
antagonists of the renin-angiotensin system, endothelin antagonists, other anti-fibrotic
agents, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside
inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin
inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating
10 HCV; or mixtures thereof.

