



(86) Date de dépôt PCT/PCT Filing Date: 2002/05/17
(87) Date publication PCT/PCT Publication Date: 2002/11/28
(85) Entrée phase nationale/National Entry: 2003/10/02
(86) N° demande PCT/PCT Application No.: EP 2002/005467
(87) N° publication PCT/PCT Publication No.: 2002/094879
(30) Priorités/Priorities: 2001/05/18 (01112237.1) EP;
2001/09/26 (60/325,147) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07K 16/28, A61K 31/00, A61K 47/48,
A61K 51/10, A61K 47/10, A61P 35/04, A61P 35/00,
C07K 16/46, C12N 15/13, C12N 5/10, C12N 15/63,
C12N 15/62
(71) Demandeurs/Applicants:
BOEHRINGER INGELHEIM INTERNATIONAL GMBH,
DE;
BOEHRINGER INGELHEIM PHARMACEUTICALS INC.,
US
(72) Inventeurs/Inventors:
ADOLF, GUNTHER, AT;
OSTERMANN, ELINBORG, AT;
PATZELT, ERIK, AT;
...

(54) Titre : ANTICORPS SPECIFIQUES CONTRE CD44V6
(54) Title: ANTIBODIES SPECIFIC FOR CD44V6

(57) **Abrégé/Abstract:**

The present invention belongs to the field of oncology. The invention relates to antibodies with specified sequence which are specific for an epitope which is coded by the variant exon v6 of the CD44 gene and to derivatives of said antibody. The invention also provides nucleic acid molecules encoding said antibody proteins. The invention furthermore pertains to methods for producing said antibody proteins. The invention also provides pharmaceutical compositions comprising said antibody proteins. The invention furthermore is concerned with the use in the manufacture of a medicament for the treatment of cancer.



(72) **Inventeurs(suite)/Inventors(continued)**: SPROLL, MARLIES, DE; HEIDER, KARL-HEINZ, AT; MIGLIETTA, JOHN J., US;
VAN DONGEN, AUGUSTINUS ANTONIUS MARIA SILVESTE, NL

(74) **Agent**: FETHERSTONHAUGH & CO.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 November 2002 (28.11.2002)

PCT

(10) International Publication Number
WO 02/094879 A1

(51) International Patent Classification⁷: **C07K 16/28**,
16/46, C12N 5/10, 15/13, 15/62, A61K 47/48, 51/10,
31/00, 47/10, A61P 35/00, 35/04, C12N 15/63 // (A61K
51/10, 103:10) (A61K 51/10, 103:32) (A61K 51/10,
101:02) (A61K 51/10, 121:00)

(21) International Application Number: PCT/EP02/05467

(22) International Filing Date: 17 May 2002 (17.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01112237.1 18 May 2001 (18.05.2001) EP
60/325,147 26 September 2001 (26.09.2001) US

(71) Applicants (*for all designated States except US*):
**BOEHRINGER INGELHEIM INTERNATIONAL
GMBH** [DE/DE]; 55216 Ingelheim am Rhein (DE).
**BOEHRINGER INGELHEIM PHARMACEUTI-
CALS INC.** [US/US]; 900 Ridgebury Road, Ridgefield,
CT 06877 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ADOLF, Gün-
ther** [AT/AT]; Stiftgasse 15-17/10, A-1070 Wien (AT).
OSTERMANN, Elinborg [AT/AT]; Mauerbachstrasse
56/6, A-1140 Wien (AT). **PATZELT, Erik** [AT/AT];
Hans Buchmueller-Gasse 8, A-3002 Purkersdorf (AT).

SPROLL, Marlies [DE/DE]; Münchenerstrasse 32a,
82131 Gauting (DE). **HEIDER, Karl-Heinz** [DE/AT];
Johann-Strauss-Promenade 4/11, A-2000 Stockerau (AT).
MIGLIETTA, John, J. [US/US]; 34 Country Way,
Bethel, CT 06801-2936 (US). **VAN DONGEN, Augusti-
nus, Antonius, Maria, Silvester** [NL/NL]; Poortstraat 52,
NL-3572 HL Utrecht (NL).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: ANTIBODIES SPECIFIC FOR CD44V6

(57) **Abstract:** The present invention belongs to the field of oncology. The invention relates to antibodies with specified sequence which are specific for an epitope which is coded by the variant exon v6 of the CD44 gene and to derivatives of said antibody. The invention also provides nucleic acid molecules encoding said antibody proteins. The invention furthermore pertains to methods for producing said antibody proteins. The invention also provides pharmaceutical compositions comprising said antibody proteins. The invention furthermore is concerned with the use in the manufacture of a medicament for the treatment of cancer.



WO 02/094879 A1

5 **Antibodies specific for CD44v6**

Technical Field of the Invention

The present invention belongs to the field of oncology. The invention relates to antibodies with
10 specified sequence which are specific for an epitope which is coded by the variant exon v6 of
the CD44 gene and to derivatives of said antibody. The invention also provides nucleic acid
molecules encoding said antibody proteins. The invention furthermore pertains to methods for
producing said antibody proteins. The invention also provides pharmaceutical compositions
comprising said antibody proteins. The invention furthermore is concerned with the use in the
15 manufacture of a medicament for the treatment of cancer.

Background Art

Recently it has been shown that the expression of variants of the surface glycoprotein CD44 is
20 necessary and sufficient for causing so-called spontaneous metastatic behaviour of a non-
metastasizing rat pancreatic adenocarcinoma cell line as well as a non-metastasizing rat
fibrosarcoma cell line (Günthert *et al.*, 1991). While the smallest CD44 isoform, the standard
form CD44s (or CD44std), is ubiquitarily expressed in different tissues including epithelial cells,
certain CD44 splice variants (CD44v, CD44var) are expressed only on a subset of epithelial
25 cells. The CD44 variants are generated by alternative splicing in a way that the sequences of
ten exons (v1-v10) are completely excised in CD44s but can appear in the bigger variants in
different combinations (Screaton *et al.*, 1992; Tölg *et al.*, 1993; Hofmann *et al.*, 1991). The
variants differ in that different amino acid sequences are inserted at a certain site of the
extracellular part of the protein. Such variants can be detected in various human tumor cells as
30 well as in human tumor tissue. So, the expression of CD44 variants in the course of colorectal
carcinogenesis has recently been investigated (Heider *et al.*, 1993a). The expression of CD44
variants is absent in normal human colon epithelium, and only a weak expression is detectable
in the proliferating cells of the crypts. In later stages of the tumor progression, e.g. in
adenocarcinomas, all malignancies express variants of CD44. Tissue expression of variant

5 CD44 on a high level has also been shown in aggressive Non-Hodgkin lymphomas (Koopman *et al.*, 1993).

Exon v6 appears to play a special role especially in the course of metastatic spread (Rudy *et al.*, 1993). In an animal model, antibodies against v6 specific epitopes could prevent the
10 settlement of metastatic cells and the growth of metastases (Seiter *et al.*, 1993). In colon carcinomas, v6 expression correlates with tumor progression (Wielenga *et al.*, 1993). In gastric carcinomas, v6 expression is an important diagnostic marker to distinguish tumors of the intestinal type from those of the diffuse type (Heider *et al.*, 1993b). In the latter two publications, v6 expression has been determined using antibodies against v6 specific epitopes.

15 As CD44v6 has been shown to be a tumor-associated antigen with a favorable expression pattern in human tumors and normal tissues (Heider *et al.*, 1995; Heider *et al.*, 1996), it has been subject to antibody-based diagnostic and therapeutic approaches, (Heider *et al.*, 1996; WO 95/33771; WO 97/21104).

20 One serious problem that arises when using non-human antibodies for applications in humans is that they quickly raise a human anti-non-human response that reduces the efficacy of the antibody in patients and impairs continued administration. To overcome that problem, concepts of "humanising" non-human antibodies have been developed in the art. In the first approach,
25 humanization of non-human antibodies has been tried to achieve by constructing non-human/human chimeric antibodies, wherein the non-human variable regions are joined to human constant regions (Boulianne G. L., Hozumi N. and Shulman, M. J. (1984) Production of functional chimeric mouse/human antibody *Nature* 312: 643) The chimeric antibodies thus generated retain the binding specificity and affinity of the original non-human antibody.
30 However, chimeric antibodies, although significantly better than mouse antibodies, can still elicit an anti-chimeric response in humans (LoBuglio A. F., Wheeler R. H., Trang J., Haynes A., Rogers K., Harvey E. B., Sun L., Ghrayeb J. and Khazaeli M. B. (1989) Mouse/human chimeric monoclonal antibody in man: Kinetics and immune response. *Proc. Natl. Acad. Sci.* 86: 4220). This approach was later refined by further reducing the amount of non-human

5 sequences by grafting the complementarity determining regions (CDRs) from the non-human variable regions to human variable regions and then joining these "reshaped human" variable regions to human constant regions (Riechmann L., Clark M., Waldmann H. and Winter G. (1988) Reshaping human antibodies for therapy. *Nature* **332**: 323). CDR-grafted or reshaped human antibodies contain little or no protein sequences that can be identified as being derived
10 from mouse antibodies. Although an antibody humanised by CDR-grafting may still be able to elicit some immune reactions, such as an anti-allotype or an anti-idiotypic response, as seen even with natural human antibodies, the CDR-grafted antibody will be significantly less immunogenic than a mouse antibody thus enabling a more prolonged treatment of patients.

15 However, it soon turned out that CDR-grafting alone did not result in antibodies with sufficient binding affinity. CDR-grafted antibodies have relatively poor binding characteristics as compared to their parent non-human antibodies because more amino acids than those within the CDR's are involved in antigen binding. In consequence, CDR-grafted antibodies with poor binding affinity are not regarded to be useful in therapy. Therefore, attempts have been made
20 to create antibodies which combine the low immunogenicity of CDR-grafted antibodies with the good binding characteristics of the non-human parent antibodies. The concept was developed that, in addition to CDR-grafting, one to several amino acids in the humanized framework region have to be retained as residues of rodent donor origin for retaining binding affinity (Queen et al, (1989)*Proc. Natl. Acad. Sci.* **86**: 10029-10033).

25 Because of the high potential utility such antibodies could have in diagnosis and therapy, there is a need of antibodies with improved properties which are suitable for treatment of human cancer.

30 The problem underlying the present invention was to provide an antibody with significantly better properties as compared to the known CD44v6 specific antibodies.

Summary of the invention

5 The above-captioned technical problem is solved by the embodiments characterized in the claims and the description. The before-mentioned disadvantages in the art are overcome by the claims and the description of the present invention.

10 In order to solve the problems mentioned above, the present inventors have designed and generated a CD44v6 specific humanised antibody called BIWA8, which was both CDR-grafted and framework-mutated and had low immunogenicity combined with high affinity.

15 However, the inventors were able to create an antibody with even superior therapeutic utility, called BIWA4. Albeit this one has less binding affinity as compared to BIWA8, it surprisingly shows a much more favorable biodistribution and tumor uptake when administered in vivo.

20 The present invention belongs to the field of oncology. The invention relates to antibodies with specified sequence which are specific for an epitope which is coded by the variant exon v6 of the CD44 gene and to derivatives of said antibody. The invention also provides nucleic acid molecules encoding said antibody proteins. The invention furthermore pertains to methods for producing said antibody proteins. The invention also provides pharmaceutical compositions comprising said antibody proteins. The invention furthermore is concerned with the use in the manufacture of a medicament for the treatment of cancer.

25 **Brief description of the figures**
(see also example 1)

Figure 1: Evaluation of relative binding affinities tested in a competitive cell ELISA.

30 IC50: concentrations of cMAb and hMAbs at which binding of mMAb BIWA 1 to attached A431 cells is reduced by 50%. IC50 values relative to BIWA 2 are indicated.

Figure 2. Biodistributions of co-injected ^{125}I - and ^{131}I -labeled CD44v6-specific MAbs (10 μCi , 50 μg) in HNX-OE xenograft-bearing mice at 3 or 4 days p.i. Three groups of mice received either (A) ^{131}I - U36 (black bars) and ^{125}I -BIWA 1 (hatched bars) (n=5), (B) ^{131}I -BIWA 4 (black

bars) and ^{125}I -BIWA 2 (hatched bars) (n=6) or (C) ^{131}I -BIWA 4 (black bars) and ^{125}I -BIWA 8 (hatched bars) (n=6). At 3 (A) or 4 days (B,C) after injection mice were bled, sacrificed, dissected and the radioactivity levels (%ID/g $\pm$ s.e.m.) of tumor, blood and several organs were assessed. (Bld: blood, Tum: tumor, Liv: liver, Spl: spleen, Kid: kidney, Hrt: heart, Stm: stomach, Ilm: ileum, Cln: colon, Blr: bladder, Str: sternum, Msc: muscle, Lng: lung, Skn: skin, Tng: tongue).

Figure 3. Therapeutic efficacy of ^{186}Re -labeled CD44v6-specific MAbs in HNX-OE xenograft-bearing nude mice. Mice received 300 μCi ^{186}Re -U36 (- $\otimes$ -, Fig A), 300 μCi ^{186}Re -BIWA 1 (- $\otimes$ -, Fig A), 300 μCi ^{186}Re -BIWA 4 (- $\otimes$ -, Fig. B), 300 μCi ^{186}Re -BIWA 2 (- $\otimes$ -, Fig. B), 400 μCi ^{186}Re -BIWA 4 (- $\otimes$ -, Fig. C), 400 μCi ^{186}Re -BIWA 8 (- $\otimes$ -, Fig. C), or saline (- $\otimes$ -, Fig. A, B, C) as control. Control groups in Fig A and B are the same. The tumor size is expressed as the average tumor volume ($\pm$ s.e.m.) during treatment relative to the average tumor volume at the start of therapy.

Figure 4. Relationship between MAb dose administered and the AUC observed following BIWA 4 intravenous infusion to 10 patients in Part A of the study.

Figure 5. Relationship between MAb dose administered and the maximum plasma BIWA 4 concentration observed following BIWA 4 intravenous infusion to 10 patients in Part A of the study.

5 **Disclosure of the preferred embodiments of the invention**

Before the embodiments of the present invention it must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "an antibody" includes a
10 plurality of such antibodies, reference to the "cell" is a reference to one or more cells and equivalents thereof known to those skilled in the art, and so forth. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing
15 of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by
20 virtue of prior invention.

The terms "antibody molecule" or "antibody protein" or "antibody" as used herein shall be considered equivalent.

"Complementarity determining regions of a monoclonal antibody" are understood to be those amino acid sequences involved in specific antigen binding according to Kabat (Kabat E. A.,
25 Wu T. T., Perry H. M., Gottesman K. S. and Foeller C. (1991) *Sequences of Proteins of Immunological Interest* (5th Ed.). NIH Publication No. 91-3242. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, Bethesda, MD.) in connection with Chothia and Lesk (Chothia and Lesk (1987) *J. Mol. Biol.* 196:901-917).

30 As used herein, the term "framework modifications" refers to the exchange, deletion or addition of single or multiple amino acids in the variable regions surrounding the individual complementarity determining regions. Framework modifications may have an impact on the immunogenicity, producibility or binding specificity of an antibody protein.

5 The present invention provides an antibody molecule comprising a variable region of the heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 1 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof. Both antibodies BIWA4 and BIWA 8 comprise the variable region of the heavy chain as characterized in amino acid sequence SEQ ID No. 1.

10 A „fragment“ according to the invention is a shorter antibody molecule, i.e. any polypeptide subset, characterized in that it is encoded by a shorter nucleic acid molecule than disclosed below, however still retains its antibody binding activity.

15 A „functional variant“ of the antibody molecule according to the invention is an antibody molecule which possesses a biological activity (either functional or structural) that is substantially similar to the antibody molecule according to the invention, i.e. a substantially similar substrate specificity or cleavage of the substrate. The term „functional variant“ also includes „a fragment“, „an allelic variant“, „a functional variant“, „variant based on the degenerative nucleic acid code“ or „chemical derivatives“. Such a „functional variant“ e.g. may
20 carry one or several point mutations, one or several nucleic acid exchanges, deletions or insertions or one or several amino acid exchanges, deletions or insertions. Said functional variant is still retaining its biological activity such as antibody binding activity, at least in part or even going along with an improvement said biological activity.

25 A „functional variant“ of the antibody molecule according to the invention is a antibody molecule which possesses a biological activity (either functional or structural) that is substantially similar to the antibody molecule according to the invention, i.e. a substantially similar target molecule binding activity. The term „functional variant“ also includes „a fragment“, „an allelic variant“, „a functional variant“, „variant based on the degenerative nucleic acid code“ or „chemical derivatives“.

30 An „allelic variant“ is a variant due to the allelic variation, e.g. differences in the two alleles in humans. Said variant is still retaining its biological activity such as antibody target binding activity, at least in part or even going along with an improvement said biological activity.

A „variant based on the degenerative of the genetic code“ is a variant due to the fact that a certain amino acid may be encoded by several different nucleotide triplets. Said variant is still

5 retaining its biological activity such as antibody binding activity, at least in part or even going along with an improvement said biological activity.

A „fusion molecule“ may be the antibody molecule according to the invention fused to e.g. a reporter such as a radiolabel, a chemical molecule such as a toxin or a fluorescent label or any other molecule known in the art.

10 As used herein, a „chemical derivative“ according to the invention is an antibody molecule according to the invention chemically modified or containing additional chemical moieties not normally being part of the molecule. Such moieties may improve the molecule's activity such as target destruction (e.g. killing of tumor cells) or may improve its solubility, absorption, biological half life etc.

15 A molecule is „substantially similar“ to another molecule if both molecules have substantially similar structures or biological activity. Thus, provided that two molecules possess a similar activity, they are considered variants as that term is used herein even if the structure of one of the molecules is not found in the other, or if the sequence of amino acid residues is not identical.

20 For many uses of the antibodies according to the invention it is desirable to have the smallest possible antigen-binding, i.e. CD44v6-binding units. Therefore in another preferred embodiment an antibody protein according to the invention is a Fab fragment (Fragment antigen-binding = Fab). These CD44v6-specific antibody proteins according to the invention consist of the variable regions of both chains which are held together by the adjacent constant
25 region. These may be formed by protease digestion, e.g. with papain, from conventional antibodies, but similar Fab fragments may also be produced in the mean time by genetic engineering. In another preferred embodiment an antibody protein according to the invention is an F(ab')₂ fragment, which may be prepared by proteolytic cleaving with pepsin.

Using genetic engineering methods it is possible to produce shortened antibody fragments
30 which consist only of the variable regions of the heavy (VH) and of the light chain (VL). These are referred to as Fv fragments (Fragment variable = fragment of the variable part). In another preferred embodiment an CD44v6-specific antibody molecule according to the invention is such an Fv fragment. Since these Fv-fragments lack the covalent bonding of the two chains by the cysteines of the constant chains, the Fv fragments are often stabilised. It is advantageous to

5 link the variable regions of the heavy and of the light chain by a short peptide fragment, e.g. of 10 to 30 amino acids, preferably 15 amino acids. In this way a single peptide strand is obtained consisting of VH and VL, linked by a peptide linker. An antibody protein of this kind is known as a single-chain-Fv (scFv). Examples of scFv-antibody proteins of this kind known from the prior art are described in Huston et al. (1988, PNAS 16: 5879-5883). Therefore, in another
10 preferred embodiment an CD44v6-specific antibody protein according to the invention is a single-chain-Fv protein (scFv).

In recent years, various strategies have been developed for preparing scFv as a multimeric derivative. This is intended to lead, in particular, to recombinant antibodies with improved pharmacokinetic and biodistribution properties as well as with increased binding avidity. In
15 order to achieve multimerisation of the scFv, scFv were prepared as fusion proteins with multimerisation domains. The multimerisation domains may be, e.g. the CH3 region of an IgG or *coiled coil* structure (helix structures) such as *Leucin-zipper* domains. However, there are also strategies in which the interaction between the VH/VL regions of the scFv are used for the multimerisation (e.g. di-, tri- and pentabodies). Therefore in another preferred embodiment an
20 antibody protein according to the invention is an CD44v6-specific diabody antibody fragment. By diabody the skilled person means a bivalent homodimeric scFv derivative (Hu et al., 1996, PNAS 16: 5879-5883). The shortening of the *Linker* in an scFv molecule to 5- 10 amino acids leads to the formation of homodimers in which an inter-chain VH/VL-superimposition takes place. Diabodies may additionally be stabilised by the incorporation of disulphide bridges.
25 Examples of diabody-antibody proteins from the prior art can be found in Perisic et al. (1994, Structure 2: 1217-1226).

By minibody the skilled person means a bivalent, homodimeric scFv derivative. It consists of a fusion protein which contains the CH3 region of an immunoglobulin, preferably IgG, most preferably IgG1 as the dimerisation region which is connected to the scFv via a *Hinge region*
30 (e.g. also from IgG1) and a *Linker* region. The disulphide bridges in the *Hinge* region are mostly formed in higher cells and not in prokaryotes. In another preferred embodiment an antibody protein according to the invention is an CD44v6-specific minibody antibody fragment. Examples of minibody-antibody proteins from the prior art can be found in Hu et al. (1996, Cancer Res. 56: 3055-61).

5 By triabody the skilled person means a: trivalent homotrimeric scFv derivative (Kortt et al. 1997 Protein Engineering 10: 423-433). ScFv derivatives wherein VH-VL are fused directly without a linker sequence lead to the formation of trimers.

The skilled person will also be familiar with so-called miniantibodies which have a bi-, tri- or tetravalent structure and are derived from scFv. The multimerisation is carried out by di-, tri-
10 or tetrameric coiled coil structures (Pack et al., 1993 Biotechnology 11:, 1271-1277; Lovejoy et al. 1993 Science 259: 1288-1293; Pack et al., 1995 J. Mol. Biol. 246: 28-34).

Therefore in a preferred embodiment an antibody protein according to the invention is an CD44v6-specific multimerised molecule based on the abovementioned antibody fragments and may be, for example, a triabody, a tetravalent miniantibody or a pentabody.

15

In a more preferred embodiment, the invention relates to an antibody molecule wherein the variable region of the heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 1.

In another preferred embodiment, the invention relates to an antibody molecule comprising a
20 variable region of the light chain as characterized by the amino acid sequence as defined in SEQ ID No. 2 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof. Antibody BIWA4 as used herein comprises the variable region of the light chain as defined in amino acid sequence SEQ ID No. 2.

In another more preferred embodiment, the invention relates to an antibody molecule wherein
25 the variable region of the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 2.

In another preferred embodiment, the invention relates to an antibody molecule comprising a variable region of the light chain as characterized by the amino acid sequence as defined in SEQ ID No. 3 or a fragment, allelic variant, functional variant, glycosylation variant, fusion
30 molecule or a chemical derivative thereof. Antibody BIWA 8 comprises the variable region of the light chain as characterized in amino acid sequence SEQ ID No. 3.

In another more preferred embodiment, the invention relates to an antibody molecule wherein the variable region of the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 3.

5 In another more preferred embodiment, the invention relates to an antibody molecule according to the invention comprising a variable region of the heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 1 and comprising a variable region of light chain as characterized by the amino acid sequence as defined in SEQ ID No. 2 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical
10 derivative thereof. Antibody BIWA4 comprises the variable region of the heavy chain as characterized in amino acid sequence SEQ ID No. 1 and variable region of the light chain as defined in amino acid sequence SEQ ID No. 2.

In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the variable region of the heavy chain consists of the amino acids as
15 characterized by the amino acid sequence of SEQ ID No. 1 and wherein the variable region of the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 2.

In another more preferred embodiment, the invention relates to an antibody molecule according to the invention comprising a variable region of the heavy chain as characterized by
20 the amino acid sequence as defined in SEQ ID No. 1 and comprising a variable region of the light chain as characterized by the amino acid sequence as defined in SEQ ID No. 3 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof. Antibody BIWA8 comprises the variable region of the heavy chain as characterized in amino acid sequence SEQ ID No. 1 and variable region of the light chain as
25 defined in amino acid sequence SEQ ID No. 3.

In another most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the variable region of the heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 1 and wherein the variable region of the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ
30 ID No. 3.

In another preferred embodiment, the invention relates to an antibody molecule comprising a variable region of the heavy chain encoded by the nucleic acid sequence as defined in SEQ ID No. 4 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Both antibodies BIWA4

5 and BIWA 8 comprise the variable region of the heavy chain as characterized in nucleic acid sequence SEQ ID No. 4.

In another more preferred embodiment, the invention relates to an antibody molecule wherein the variable region of the heavy chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 4.

10 In another preferred embodiment, the invention relates to an antibody molecule comprising a variable region of the light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 5 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA4 as used herein comprises the variable region of the light chain as defined in nucleic acid sequence SEQ
15 ID No. 5.

In another more preferred embodiment, the invention relates to an antibody molecule wherein the variable region of the light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 5.

In another preferred embodiment, the invention relates to an antibody molecule comprising a
20 variable region of the light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 6 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA 8 comprises the variable region of the light chain as characterized in nucleic acid sequence SEQ ID No. 6.

25 In another more preferred embodiment, the invention relates to an antibody molecule wherein the variable region of the light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 6.

In another more preferred embodiment, the invention relates to an antibody molecule
30 according to the invention comprising a variable region of the heavy chain encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and comprising a variable region of the light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 5 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA4 comprises the variable region of the heavy

5 chain as characterized in nucleic acid sequence SEQ ID No. 4 and variable region of the light chain as defined in nucleic acid sequence SEQ ID No. 5.

In another most preferred embodiment, the invention relates to an antibody molecule according to to the invention wherein the variable region of the heavy chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and wherein the variable region of the light chain is
10 encoded by the nucleic acid sequence as defined in SEQ ID No. 5.

In another more preferred embodiment, the invention relates to an antibody molecule according to the invention comprising a variable region of the heavy chain encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and comprising a variable region of the light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 6 or a fragment, allelic
15 variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA8 comprises the variable region of the heavy chain as characterized in nucleic acid sequence SEQ ID No. 4 and variable region of the light chain as defined in nucleic acid sequence SEQ ID No. 6.

In another most preferred embodiment, the invention relates to an antibody molecule according
20 to the invention wherein the variable region of the heavy chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and wherein the variable region of the light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 6.

To generate humanised CD44v6-specific antibody proteins the disclosed nucleic acid sequences were expressed (see *infra* and examples) by molecular biology methods known in
25 the art.

The variable regions of the antibody proteins of the present invention are typically linked to at least a portion of the immunoglobulin constant region (F_c), typically that of a human immunoglobulin. Human constant region DNA sequences can be isolated in accordance with well-known procedures from a variety of human cells, but preferably immortalized B cells (see
30 Kabat et al., *supra*, and WO 87/02671). Hence the antibody proteins of the invention may contain all or only a portion of the constant region as long as they exhibit specific binding to the CD44v6 antigen. The choice of the type and extent of the constant region depends on whether effector functions like complement fixation or antibody dependent cellular toxicity are desired, and on the desired pharmacological properties of the antibody protein. The antibody

5 protein of the invention will typically be a tetramer consisting of two light chain/heavy chain pairs, but may also be dimeric, i.e. consisting of a light chain/heavy chain pair, e.g. a Fab or Fv fragment.

Therefore, in a further embodiment the invention relates to antibody proteins according to the
10 invention, characterised in that they have a variable light chain region and a variable heavy chain region, each joined to a human constant region. In particular, the variable region of the light chain was joined to a human kappa constant region and the variable region of the heavy chain was joined to a human gamma-1 constant region. Other human constant regions for chimerizing light and heavy chains are also available to the expert.

15 Humanization of the variable region of a murine antibody may be achieved employing methods known in the art. EP 0239400 discloses grafting of the CDRs of a murine variable region into the framework of a human variable region. WO 90/07861 discloses methods of reshaping a CDR-grafted variable region by introducing additional framework modifications. WO 92/11018 discloses methods of producing humanized Ig combining donor CDRs with an
20 acceptor framework that has a high homology to the donor framework. WO 92/05274 discloses the preparation of framework mutated antibodies starting from a murine antibody. Further prior art references related to humanization of murine monoclonal antibodies are EP 0368684; EP 0438310; WO 92/07075, or WO 92/22653.

In another preferred embodiment, the invention relates to an antibody molecule according to
25 the invention characterised in that each of said variable region of the light chain and said variable region of the heavy chain region is separately joined to a human constant region.

In another more preferred embodiment, the invention relates to an antibody molecule according to the invention, wherein said human constant region of the light chain is a human kappa constant region.

30 In another more preferred embodiment, the invention relates to an antibody protein according to the invention, wherein said human constant region of the heavy chain is a human IgG1 constant region.

5 Preferred are also antibodies comprising the heavy chain as characterized by the amino acid sequence of SEQ ID No. 7 and/or the light chain as characterized by the amino acid sequence of SEQ ID No. 8 or as characterized by the amino acid sequence of SEQ ID No. 9.

Thus, another important embodiment is an antibody molecule according to the invention comprising a heavy chain as characterized by the amino acid sequence as defined in SEQ ID
10 No. 7 and comprising a light chain as characterized by the amino acid sequence as defined in SEQ ID No. 8 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof. Antibody BIWA4 comprises the heavy chain as characterized in amino acid sequence SEQ ID No. 7 and variable region of the light chain as defined in amino acid sequence SEQ ID No. 8.

15 In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 7 and wherein the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 8. Antibody BIWA4 consists of the sequences as disclosed in amino acid sequence SEQ ID No. 7 (heavy chain) and amino acid
20 sequence SEQ ID No. 8 (light chain). BIWA4 is a CDR-grafted antibody without framework modifications. Surprisingly, this antibody has, despite lower binding affinity, superior therapeutic efficacy, better biodistribution and tumor uptake over the framework-mutated antibody BIWA8 (see example). It is a humanised version of antibody VFF-18 (=BIWA1) mentioned above, having the complementary determining regions of the murine monoclonal
25 antibody VFF-18 in a completely human framework, and human constant regions. It is therefore an antibody of very low immunogenicity in man, which is a favorable trait. However, as it has no murine framework residues to optimise antigen binding, it has a significantly lower antigen binding affinity as its parent antibody VFF-18, and therefore would not have been regarded as a good candidate for a therapeutic drug. Unexpectedly, it has been found that
30 BIWA4, despite its poor binding affinity, has a very favorable biodistribution and tumor uptake in vivo, making it superior to other humanised versions of VFF-18 with higher binding affinity.

Another important embodiment is an antibody molecule according to the invention comprising a heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 7 and

5 comprising a light chain as characterized by the amino acid sequence as defined in SEQ ID No. 9 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof. Antibody BIWA8 comprises the heavy chain as characterized in amino acid sequence SEQ ID No. 7 and variable region of the light chain as defined in amino acid sequence SEQ ID No. 9.

10 In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 7 and wherein the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 9. Antibody BIWA8 consists of the sequences as disclosed in amino acid sequence SEQ ID No. 7 (heavy chain) and amino acid
15 sequence SEQ ID No. 9 (light chain). BIWA8 is a CDR-grafted antibody with framework modifications. This antibody has significant higher binding affinity than BIWA4 (see example). Preferred are also antibodies comprising the heavy chain as encoded by the nucleic acid sequence of SEQ ID No. 10 and/or the light chain as characterized by the nucleic acid sequence of SEQ ID No. 11 or as characterized by the nucleic acid sequence of SEQ ID No.
20 12. Said sequences include non-translated sequences and the leader sequence as cloned in vector pAD-CMV1/pAD-CMV19.

Therefore, another important embodiment is an antibody molecule according to the invention comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 10 and comprising a light chain as characterized by the nucleic acid sequence as defined in
25 SEQ ID No. 11 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA4 comprises the heavy chain as encoded by nucleic acid sequence SEQ ID No. 10 and variable region of the light chain as encoded by nucleic acid sequence SEQ ID No. 11.

In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 10
30 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 11.

Another important embodiment is an antibody molecule according to the invention comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 10 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID

5 No. 12 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA8 comprises the heavy chain as encoded by nucleic acid sequence SEQ ID No. 10 and variable region of the light chain as encoded by nucleic acid sequence SEQ ID No. 12.

In a most preferred embodiment, the invention relates to an antibody molecule according to the
10 invention wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 10 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 12.

Preferred are also antibodies comprising the heavy chain as encoded by the nucleic acid sequence of SEQ ID No. 13 and/or the light chain as characterized by the nucleic acid sequence of SEQ ID No. 14 or as characterized by the nucleic acid sequence of SEQ ID No.
15 15. Said sequences include the leader sequence as cloned in vector N5KG1val.

Therefore, another important embodiment is an antibody molecule according to the invention comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 13 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID No. 14 or a fragment, allelic variant, functional variant, variant based on the
20 degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA4 comprises the heavy chain as encoded by nucleic acid sequence SEQ ID No. 13 and variable region of the light chain as encoded by nucleic acid sequence SEQ ID No. 14.

In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 13
25 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 14.

Another important embodiment is an antibody molecule according to the invention comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 13 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID No. 15 or a fragment, allelic variant, functional variant, variant based on the degenerative
30 nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody BIWA8 comprises the heavy chain as encoded by nucleic acid sequence SEQ ID No. 13 and variable region of the light chain as encoded by nucleic acid sequence SEQ ID No. 15.

5 In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 13 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 15.

Most preferred is the antibody protein comprising the heavy and light chain as encoded by the nucleic acid sequence of SEQ ID No. 16. Said sequence includes the leader sequence as cloned
10 in vector N5KG1val.

Therefore, another highly important embodiment is an antibody molecule according to the invention comprising a heavy and light chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 16 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof. Antibody
15 BIWA4 comprises the heavy and light chain as encoded by nucleic acid sequence SEQ ID No. 16.


In a most preferred embodiment, the invention relates to an antibody molecule according to the invention wherein the heavy and light chain is encoded by the nucleic acid sequence of SEQ ID No. 16. This sequence is encoding the entire antibody BIWA4.

20 The antibody proteins of the invention provide a highly specific tool for targeting therapeutic agents to the CD44v6 antigen. Therefore, in a further aspect, the invention relates to antibody proteins according to the invention, wherein said antibody protein is conjugated to a therapeutic agent. Of the many therapeutic agents known in the art, therapeutic agents selected
25 from the group consisting of radioisotopes, toxins, toxoids, inflammotogenic agents, enzymes, antisense molecules, peptides, cytokines, and chemotherapeutic agents are preferred. Among the radioisotopes, gamma, beta and alpha-emitting radioisotopes may be used as a therapeutic agent. β -emitting radioisotopes are preferred as therapeutic radioisotopes. $^{186}\text{Rhenium}$, $^{188}\text{Rhenium}$, $^{131}\text{Iodine}$ and $^{90}\text{Yttrium}$ have been proven to be particularly useful β -emitting
30 isotopes to achieve localized irradiation and destruction of malignant tumor cells. Therefore, radioisotopes selected from the group consisting of $^{186}\text{Rhenium}$, $^{188}\text{Rhenium}$, $^{131}\text{Iodine}$ and $^{90}\text{Yttrium}$ are particularly preferred as therapeutic agents conjugated to the antibody proteins of the invention. For example, for the radioiodination of an antibody of the invention, a method as disclosed in WO 93/05804 may be employed.

Thus, a more preferred aspect of the present invention is an antibody protein according to the invention, wherein said therapeutic agent is a therapeutic agent selected from the group consisting of radioisotopes, toxins, toxoids, pro-drugs and chemotherapeutic agents.

A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said therapeutic agent is linked to the antibody protein via a linker selected from the group of MAG-3 (US 5082930 A, EP 0247866 B1 (page 2 lines 55-56 – page 3 lines 1-23)); MAG-2 GABA (US 5681927 A, EP 0284071 B1 (page 6 lines 9-29)); and N2S2 ((=phenthioate) US 4897255 A, US 5242679 A, EP 0188256 B1 (page 2, lines 38- page 3, lines 18)), all herein incorporated by reference.

The formulae of said linkers are as follows:



A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said therapeutic agent is linked to the antibody protein via MAG-2 GABA.

A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said radioisotope is a β -emitting radioisotope.

A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said radioisotope is selected from the group consisting of ^{186}Re Rhenium, ^{188}Re Rhenium, ^{131}I Iodine and ^{90}Y Yttrium.

A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said radioisotope is ^{186}Re Rhenium.

5 A further aspect of the present invention pertains to antibody proteins according to the invention, characterised in that they are labelled. Such an CD44v6-specific labelled antibody allows for the localisation and/or detection of the CD44v6 antigen *in vitro* and/or *in vivo*. A label is defined as a marker that may be directly or indirectly detectable. An indirect marker is defined as a marker that cannot be detected by itself but needs a further directly detectable
10 marker specific for the indirect marker. Preferred labels for practicing the invention are detectable markers. From the large variety of detectable markers, a detectable marker selected from the group consisting of enzymes, dyes, radioisotopes, digoxigenin, and biotin is most preferred.

Thus, a more preferred aspect of the present invention is an antibody protein according to the
15 invention, characterised in that it is labelled. More preferred is the antibody protein according to the invention, wherein said label is a detectable marker. Also more preferred is the antibody protein according to the invention, wherein the detectable marker is a detectable marker selected from the group consisting of enzymes, dyes, radioisotopes, digoxigenin, and biotin.

A further aspect of the present invention relates to antibody proteins according to the
20 invention, characterised in that they are conjugated to an imageable agent. A large variety of imageable agents, especially radioisotopes, are available from the state of the art. For practising the invention gamma-emitting isotopes are more preferred. Most preferred is ¹²⁵Iodine.

Therefore, a more preferred aspect of the present invention is an antibody protein to the
25 invention conjugated to an imageable agent. A more preferred aspect of the present invention is an antibody protein according to the invention, wherein the imageable agent is a radioisotope. A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said radioisotope is a γ -emitting radioisotope. A more preferred aspect of the present invention is an antibody protein according to the invention, wherein said
30 radioisotope is ¹²⁵I.

Therefore, a more preferred aspect of the present invention is an antibody protein conjugated to a radioisotope as described above, wherein the antibody protein has specific activity of from about 0.5 to about 15 mCi/mg, or from about 0.5 to about 14 mCi/mg, preferably about 1 to

5 about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.

Another preferred embodiment of the present invention is a pharmaceutical composition containing an antibody according to the invention and a pharmaceutically acceptable carrier or excipient.

10 A pharmaceutically acceptable carrier can contain physiologically acceptable compounds that act, for example, to stabilize or to increase the absorption of an AMPA glutamate receptor agonist, antagonist or modulator. Such physiologically acceptable compounds include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins or other stabilizers or
15 excipients (see also e.g. Remington's Pharmaceutical Sciences (1990), 18th ed. Mack Publ., Easton). One skilled in the art would know that the choice of a pharmaceutically acceptable carrier, including a physiologically acceptable compound, depends, for example, on the route of administration of the composition.

In an animal or human body, it can prove advantageous to apply the pharmaceutical
20 compositions as described above via an intravenous or other route, e.g. systemically, locally or topically to the tissue or organ of interest, depending on the type and origin of the disease or problem treated, e.g. a tumor. For example, a systemic mode of action is desired when different organs or organ systems are in need of treatment as in e.g. systemic autoimmune diseases, or allergies, or transplantations of foreign organs or tissues, or tumors that are diffuse
25 or difficult to localise. A local mode of action would be considered when only local manifestations of neoplastic or immunologic action are expected, such as, for example local tumors.

The pharmaceutical compositions comprising antibody proteins of the present invention may be applied by different routes of application known to the expert, notably intravenous injection or
30 direct injection into target tissues. For systemic application, the intravenous, intravascular, intramuscular, intraarterial, intraperitoneal, oral, or intrathecal routes are preferred. A more local application can be effected subcutaneously, intracutaneously, intracardially, intralobally, intramedullary, intrapulmonarily or directly in or near the tissue to be treated (connective-, bone-, muscle-, nerve-, epithelial tissue). Depending on the desired duration and effectiveness

5 of the treatment, pharmaceutical antibody compositions may be administered once or several times, also intermittently, for instance on a daily basis for several days, weeks or months and in different dosages.

For preparing suitable pharmaceutical compositions comprising antibody preparations for the
10 applications described above, the expert may use known injectable, physiologically acceptable sterile solutions. For preparing a ready-to-use solution for parenteral injection or infusion, aqueous isotonic solutions, such as e.g. saline or corresponding plasma protein solutions are readily available. The pharmaceutical compositions may be present as lyophilisates or dry preparations, which can be reconstituted with a known injectable solution directly before use
15 under sterile conditions, e.g. as a kit of parts. The final preparation of the antibody compositions of the present invention are prepared for injection, infusion or perfusion by mixing purified antibodies according to the invention with a sterile physiologically acceptable solution, that may be supplemented with known carrier substances or/and additives (e.g. serum albumin, dextrose, sodium bisulfite, EDTA).

20 The amount of the antibody applied depends on the nature of the disease. In cancer patients, the applied dose of a 'naked' antibody which is comprised in the pharmaceutical composition according to the invention may be between 0.1 and 100 mg/m², preferably between 5 and 50 mg/m² per application, preferably 10 mg/m² to about 40 mg/m², preferably 10 mg/m² to about 30 mg/m², also preferably 20 mg/m² to about 30 mg/m², and most preferably about 25 mg/m²
25 body surface area. Also most preferred is an antibody protein dose of about 50 mg/m² body surface area.

The dose of radioactivity applied to the patient per administration has to be high enough to be effective, but must be below the dose limiting toxicity (DLT). For pharmaceutical compositions comprising radiolabeled antibodies, e.g. with ¹⁸⁶Rhenium, the maximally tolerated
30 dose (MTD) has to be determined which must not be exceeded in therapeutic settings. Application of radiolabeled antibody to cancer patients may then be carried out by repeated (monthly or weekly) intravenous infusion of a dose which is below the MTD (See e.g. Welt et al. (1994) *J. Clin. Oncol.* 12: 1193-1203). Multiple administrations are preferred, generally at weekly intervals; however, radiolabelled materials should be administered at longer intervals,

5 i.e., 4-24 weeks apart, preferable 12-20 weeks apart. The artisan may choose, however, to divide the administration into two or more applications, which may be applied shortly after each other, or at some other predetermined interval ranging, e.g. from 1 day to 1 week. Furthermore, the applied radioactivity dose will be in accordance with the guidelines outlined below. In general, the radioactivity dose per administration will be between 30 and 75 mCi/m² 10 body surface area (BSA). Thus, the amount of radiolabelled antibody in the pharmaceutical composition according to the invention, preferably labelled with ¹⁸⁶Rhenium, ¹⁸⁸Rhenium, ^{99m}Technetium, ¹³¹Iodine, or ⁹⁰Yttrium, most preferably labelled with ¹⁸⁶Rhenium, to be applied to a patient is 10, 20, 30, 40, 50 or 60 mCi/m², preferably 50 mCi/m². In a preferred embodiment, the invention relates to a pharmaceutical composition, wherein the dose of said 15 radiolabelled antibody according to the invention is MTD, preferably 50 mCi/m². This is extensively exemplified in clinical studies as set out in examples 3 to 6.

Preferred also is a pharmaceutical composition according to the invention comprising an antibody protein conjugated to a radioisotope according to the invention as defined *supra*, wherein the antibody protein has specific activity of from about 0.5 to about 15 mCi/mg, or 20 from about 0.5 to about 14 mCi/mg, preferably about 1 to about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.

Preferred also is a pharmaceutical composition according to the invention comprising an antibody protein conjugated to a radioisotope according to the invention as defined *supra*, wherein said antibody or antibody derivative is in an aqueous solution at pH of from about 7 to 25 about 8, and at a concentration of from about 0.5 to about 2.0 mg/ml.

A preferred embodiment is a pharmaceutical composition according to the invention, further comprising one or more radioprotectants selected from the group of ascorbic acid, gentisic acid, reductic acid, erythrorbic acid, p-aminobenzoic acid, 4-hydroxybenzoic acid, nicotinic acid, nicotinamide, 2-5-dihydroxy-1,4-benzenedisulfonic acid, povidone, inositol, and/or 30 citrate.

Preferred is a pharmaceutical composition according to the invention, wherein the radioprotectant is ascorbic acid.

Another preferred embodiment is a pharmaceutical composition according to the invention, wherein said antibody protein comprises an antibody molecule selected from the group of

5 antibody molecules BIWA4 or BIWA8 as described *supra* linked to $^{186}\text{Rhenium}$ via MAG-2 GABA further comprising the radioprotectant ascorbic acid.

Another preferred embodiment of the present invention is the use of an antibody protein according to the invention in the manufacture of a medicament for treatment of cancer. In a preferred embodiment the present invention relates to the use of antibody proteins according to
10 the invention conjugated to a therapeutic agent as described above for the treatment of cancer. Cancer includes any disease associated with malignant growth such as solid tumors, sarcomas and leukemias. A necessary precondition for such diseases is the expression of CD44v6. Cancer according to the invention includes, but is not limited to:

- 1) The treatment of epithelial carcinomas including breast, lung, colorectal, head and
15 neck, pancreatic, ovarian, bladder, gastric, skin, endometrial, ovarian, testicular, esophageal, prostatic and renal origin;
- 2) Bone and soft-tissue sarcomas: Osteosarcoma, chondrosarcoma, fibrosarcoma, malignant fibrous histiocytoma (MFH), leiomyosarcoma;
- 3) Hematopoietic malignancies: Hodgkin's and non-Hodgkin's lymphomas, leukemias;
- 20 4) Neuroectodermal tumors: Peripheral nerve tumors, astrocytomas, melanomas;
- 5) Mesotheliomas.

Examples for cancerous disease states associated with solid tumors include, but are not limited to: colorectal cancer, non-small cell lung cancer, breast cancer, head and neck cancer, ovarian
25 cancer, lung cancer, bladder cancer, pancreatic cancer and metastatic cancers of the brain.

Thus, a preferred embodiment is the use of an antibody protein according to the invention wherein said cancer is selected from the group consisting of colorectal cancer, non-small cell lung cancer, breast cancer, head and neck cancer, ovarian cancer, lung cancer, bladder cancer, pancreatic cancer and metastatic cancers of the brain.

30 Preferably also is the use of an antibody protein according to the invention as defined *supra* in the manufacture of a medicament for treatment of cancer, wherein the amount of antibody protein per application is between 0.1 and 100 mg/m², preferably between 5 and 50 mg/m², preferably 10 mg/m² to about 40 mg/m², preferably 10 mg/m² to about 30 mg/m², also

5 preferably 20 mg/m² to about 30 mg/m², and most preferably about 25 mg/m² body surface area. Also most preferred is an antibody protein dose of about 50 mg/m² body surface area.

Preferred also is the use of an antibody protein conjugated to a radioisotope according to the invention as defined *supra* in the manufacture of a medicament for treatment of cancer, wherein the radioactivity dose per administration is between 30 and 75 mCi/m² body surface
10 area (BSA). Preferred is the use of an antibody protein conjugated to a radioisotope according to the invention as defined *supra* in the manufacture of a medicament for treatment of cancer, wherein the antibody protein according to the invention is radiolabelled with ¹⁸⁶Rhenium, ¹⁸⁸Rhenium, ^{99m}Technetium, ¹³¹Iodine, or ⁹⁰Yttrium, and most preferably is labelled with ¹⁸⁶Rhenium. In yet another preferred embodiment the invention relates to the use of an
15 antibody protein conjugated to a radioisotope according to the invention as defined *supra* in the manufacture of a medicament for treatment of cancer, wherein the antibody dose is 10, 20, 30, 40, 50 or 60 mCi/m², most preferably 50 mCi/m². This is extensively exemplified in clinical studies as set out in examples 3 to 6.

Preferred also is the use of an antibody protein conjugated to a radioisotope according to the
20 invention as defined *supra* in the manufacture of a medicament for treatment of cancer, wherein the antibody protein has specific activity of from about 0.5 to about 15 mCi/mg, or from about 0.5 to about 14 mCi/mg, preferably about 1 to about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.

Preferred also is the use of an antibody protein conjugated to a radioisotope according to the
25 invention as defined *supra* in the manufacture of a medicament for treatment of cancer, wherein said antibody or antibody derivative is in an aqueous solution at pH of from about 7 to about 8, and at a concentration of from about 0.5 to about 2.0 mg/ml.

The invention further relates to a method of cancer treatment, wherein an antibody
protein according to the invention is administered once to several times to an individual in need
30 thereof, said antibody protein selectively binds to CD44v6, destroys tumor cells via the therapeutic agent linked to the antibody protein and the therapeutic success is monitored. Said antibody protein may be present as naked/unmodified antibody protein, modified antibody protein, such as e.g. fusion protein, or antibody protein conjugated to a therapeutic agent, which comprises contacting the tumor with an effective amount of said antibodies. The method

5 of treating tumors as described above may be effective *in vitro or in vivo*. Cancer is any cancer as described above.

The amount of the antibody applied depends on the nature of the disease. In cancer patients, the applied dose of a 'naked' antibody may be between 0.1 and 100 mg/m², preferably between 5 and 50 mg/m² per application, preferably 10 mg/m² to about 40 mg/m², preferably 10 mg/m²
10 to about 30 mg/m², also preferably 20 mg/m² to about 30 mg/m², and most preferably about 25 mg/m² body surface area. Also most preferred is an antibody protein dose of about 50 mg/m² body surface area.

The dose of radioactivity applied to the patient per administration has to be high enough to be effective, but must be below the dose limiting toxicity (DLT). For radiolabeled antibodies, e.g.
15 with ¹⁸⁶Rhenium, the maximally tolerated dose (MTD) has to be determined which must not be exceeded in therapeutic settings. Application of radiolabeled antibody to cancer patients may then be carried out by repeated (monthly or weekly) intravenous infusion of a dose which is below the MTD (See e.g. Welt et al. (1994) *J. Clin. Oncol.* 12: 1193-1203). Multiple administrations are preferred, generally at weekly intervals; however, radiolabelled materials
20 should be administered at longer intervals, i.e., 4-24 weeks apart, preferably 12-20 weeks apart. The artisan may choose, however, to divide the administration into two or more applications, which may be applied shortly after each other, or at some other predetermined interval ranging, e.g. from 1 day to 1 week.

Furthermore, the applied radioactivity dose will be in accordance with the guidelines outlined
25 below. In general, the radioactivity dose per administration will be between 30 and 75 mCi/m² body surface area (BSA). Thus, the amount of radiolabelled antibody, preferably labelled with ¹⁸⁶Rhenium, ¹⁸⁸Rhenium, ^{99m}Technetium, ¹³¹Iodine, or ⁹⁰Yttrium, most preferably labelled with ¹⁸⁶Rhenium, to be applied to a patient is 10, 20, 30, 40, 50 or 60 mCi/m², preferably 50 mCi/m². In a preferred embodiment, the invention relates to a method of treatment, wherein
30 the radiolabelled antibody as described above is administered to a patient suffering from cancer, wherein the dose of said radiolabelled antibody is MTD, preferably 50 mCi/m², whereby said cancer is prevented or treated. This is extensively exemplified in clinical studies as set out in examples 3 to 6.

5 Preferred also is a method of cancer treatment according to the invention (see above), wherein the antibody protein conjugated to a radioisotope according to the invention as defined *supra* has specific activity of from about 0.5 to about 15 mCi/mg, or from about 0.5 to about 14 mCi/mg, preferably about 1 to about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.

10 Preferred also is a method of cancer treatment according to the invention (see above), wherein the antibody protein conjugated to a radioisotope according to the invention as defined *supra* is in an aqueous solution at pH of from about 7 to about 8, and at a concentration of from about 0.5 to about 2.0 mg/ml.

Preferably, the invention relates to a method according to the invention, wherein the tumor is a
15 tumor selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.

A further aspect of the present invention is a nucleic acid, characterised in that it codes for an antibody protein according to the invention. Said nucleic acid may be RNA or preferably DNA.
20 Said DNA molecule may be chemically synthesized. First, suitable oligonucleotides can be synthesized with methods known in the art (e.g. Gait, M.J., 1984, *Oligonucleotide Synthesis. A Practical Approach*. IRL Press, Oxford, UK), which can be used to produce a synthetic gene. Methods to generate synthetic genes are known in the art (e.g. Stemmer et al. 1995, *Single-step assembly of a gene and entire plasmid from large numbers of oligodeoxyribonucleotides*,
25 Gene 164(1): 49-53; Ye et al. 1992, *Gene synthesis and expression in E. coli for pump, a human matrix metalloproteinase*, Biochem Biophys Res Commun 186(1):143-9; Hayden et Mandecki 1988, *Gene synthesis by serial cloning of oligonucleotides*, DNA 7(8): 571-7). These methods can be used to synthesize any DNA molecule disclosed in the present application, e.g. the DNA encoding BIWA4.

30 Preferably, too, a nucleic acid according to the invention is characterised in that it contains 5' or 3' or 5' and 3' untranslated regions. The nucleic acid according to the invention may contain other untranslated regions upstream and/or downstream. The untranslated region may contain a regulatory element, such as e.g. a transcription initiation unit (promoter) or enhancer. Said promoter may, for example, be a constitutive, inducible or development-controlled

5 promoter. Preferably, without ruling out other known promoters, the constitutive promoters of the human Cytomegalovirus (CMV) and Rous sarcoma virus (RSV), as well as the Simian virus 40 (SV40) and Herpes simplex promoter. Inducible promoters according to the invention comprise antibiotic-resistance promoters, heat-shock promoters, hormone-inducible „Mammary tumour virus promoter“ and the metallothioneine promoter. Preferably, too, a
10 nucleic acid according to the invention is characterised in that it codes for a fragment of the antibody protein according to the invention. This refers to part of the polypeptide according to the invention.

Preferably, a nucleic acid according to the invention is a nucleic acid as disclosed in SEQ ID
SEQ ID No. 4, 5, 6, 10, 11, 12, 13, 14, 15, and/or 16. Most preferred, said nucleic acid is a
15 nucleic acid of SEQ ID No. 16.

Another important aspect of the present invention is a recombinant DNA vector, characterised in that it contains a nucleic acid according to the invention. Preferably, said vector contains a nucleic acid as characterized in SEQ ID No. 4, 5, 6, 10, 11, 12, 13, 14, 15, and/or 16. Most preferred, said vector contains the nucleic acid as characterized in SEQ ID No. 16.

20 Examples are viral vectors such as e.g. Vaccinia, Semliki-Forest-Virus and Adenovirus. Vectors for use in COS-cells have the SV40 origin of replication and make it possible to achieve high copy numbers of the plasmids. Vectors for use in insect cells are, for example, *E. coli* transfer vectors and contain e.g. the DNA coding for polyhedrin as promoter.

Another preferred aspect of the present invention is a recombinant DNA vector according to
25 the invention, characterized in that it is an expression vector.

Another preferred aspect of the present invention is a recombinant DNA vector according to the invention, characterized in that it is vector pAD-CMV or a functional derivative thereof. Such derivatives are e.g. pAD-CMV1, pAD-CMV19 or pAD-CMV25.

Another preferred aspect of the present invention is a recombinant DNA vector according to
30 the invention, characterized in that it is the of SEQ ID No. 17 or a functional derivative thereof.

Another preferred aspect of the present invention is a recombinant DNA vector according to the invention, characterized in that it is the of SEQ ID No. 18 or a functional derivative thereof.

5 Preferably also, said vectors comprise one or several of the nucleic acid molecules as characterized in SEQ ID No. 4, 5, 6, 10, 11, 12, 13, 14, 15, and/or 16.

Preferred is also a vector as disclosed in US 5648267 A or US 5733779 A comprising a nucleotide sequence according to the invention. Preferably also, said vector comprises one or several of the nucleic acid molecules as characterized in SEQ ID No. 4, 5, 6, 10, 11, 12, 13,
10 14, 15, and/or 16. Another preferred aspect of the present invention is a recombinant DNA vector according to the invention, characterized in that it is vector N5KG1Val or a derivative thereof.

Another important aspect is a host, characterised in that it contains a vector according to the invention.

15 Another important aspect is a host according to the invention, characterised in that it is a eukaryotic host cell. The eukaryotic host cells according to the invention include fungi, such as e.g. *Pichia pastoris*, *Saccharomyces cerevisiae*, *Schizosaccharomyces*, *Trichoderma*, insect cells (e.g. from *Spodoptera frugiperda* Sf-9, with a Baculovirus expression system), plant cells, e.g. from *Nicotiana tabacum*, mammalian cells, e.g. COS cells, BHK, CHO or myeloma cells.

20 In descendants of the cells of the immune system in which antibody proteins are also formed in our body, the antibody proteins according to the invention are particularly well folded and glycosylated. Mammalian host cells, preferably CHO or COS cells are preferred, e.g. a CHO DG44 (Urlaub and Chasin, Proc. Natl. Acad. Sci. U.S.A. 77(7): 4216-20 (1980)), or CHO-K1 (ATCC CCL-61) cells. Thus, another preferred aspect is a host according to the invention
25 according to the invention, characterised in that it is a BHK, CHO or COS cell, most preferred CHO DG44 or CHO-K1 (ATCC CCL-61) cells.

Another preferred aspect is a host according to the invention, characterised in that it is a bacteriophage.

Another preferred aspect is a host according to the invention, characterised in that it is a
30 prokaryotic host cell. Examples of prokaryotic host cells are *Escherichia coli*, *Bacillus subtilis*, *Streptomyces* or *Proteus mirabilis*.

The invention further relates to a process for preparing an antibody protein according to the invention, characterized in that it comprises the following steps: a host according to the invention is cultivated under conditions in which said antibody protein is expressed by said host

5 cell and said antibody protein is isolated. The antibody according to the invention may be produced as follows. Nucleic acid molecules coding for the light chain and the heavy chain may be synthesised chemically and enzymatically by standard methods. First, suitable oligonucleotides can be synthesized with methods known in the art (details *supra*). Methods to generate synthetic genes from oligonucleotides are known in the art (details *supra*). These
10 nucleic acid molecules encoding the antibody heavy and light chains may be cloned into an expression vector (either both chains in one vector molecule, or each chain into a separate vector molecule), which then is introduced into a host cell. The host cell preferably is a mamalian host cell (details *supra*), e.g. a COS, CHO, or BHK cell, more preferably a chinese hamster ovary (CHO) cell, The host cell then is cultured in a suitable culture medium under
15 conditions where the antibody is produced, and the antibody is then isolated from the culture according to standard procedures. Procedures for production of antibodies from recombinant DNA in host cells and respective expression vectors are well-known in the art (see e.g. WO 94/11523, WO 97/9351, EP 0481790.)

The invention preferably relates to a process according to the invention, characterised in that
20 said host is a mammalian cell, preferably a CHO or COS cell.

The invention preferably relates to a process according to the invention, characterised in that said host cell is co-transfected with two plasmids which carry the expression units for the light or the heavy chain.

The following examples serve to further illustrate the present invention; but the same should
25 not be construed as limiting the scope of the invention disclosed herein.

Examples

Example 1 Radioimmunotherapy

30

MATERIALS AND METHODS.

Monoclonal antibodies. mMAb BIWA 1=VFF 18 (which is secreted by a hybridoma cell line which has been deposited on 7 June 1994 with the accession number DSM ACC2174 with the DSM-Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg

5 1b, D-38124 Braunschweig, Deutschland; see also WO 95/33771) was generated by immunizing BALB/c mice with glutathione S-transferase fusion protein containing the human CD44 domains v3-v10 (Heider et al., 1996). The epitope recognized by BIWA 1 has been mapped to amino acids 360-370 in domain v6 of CD44 (numbering according to Kugelman *et al.* (1992)). The batch used for the present studies was obtained after purification on protein-
10 G-Sepharose and dialysis against PBS.

MAb U36 (IgG1) was derived after immunization of mice with the HNSCC cell line UM-SCC-22B and recognised a different epitope within CD44v6 as BIWA 1. U36 was purified from a concentrated tissue culture supernatant by affinity chromatography on protein-A-Sepharose and further purified on Q-Sepharose.

15 **Generation of chimeric and humanized MAbs.** mRNA was isolated from the BIWA 1 hybridoma cell line by use of the QuickPrep mRNA Purification Kit (Pharmacia, Uppsala, Sweden). cDNA from the variable heavy (V_H) and variable light (V_L) chain was generated by RT-PCR.

20 The fragments were cloned into the TA cloning vector pCR II (Invitrogen, Groningen, The Netherlands) and sequenced. Two expression vectors derived from the plasmid pAD CMV1 (Himmeler et al., 1990) were constructed carrying the constant region of human gamma-1 and the constant region of the human kappa light chain, respectively. Subsequently, the V_H and V_L fragments of BIWA 1 were cloned into the corresponding expression vectors in front of the
25 constant regions. The chimeric antibody was named cMAb BIWA 2. Humanized versions of the BIWA 1 heavy and light chain variable regions (generated by CDR grafting) were cloned in front of the immunoglobulin constant regions of the above mentioned expression vectors. For the construction of humanized antibodies, the human variable regions used were derived for the heavy chain from the human immunoglobulin fragment accession number S31669 of
30 databank GenPept and for the light chain from the human immunoglobulin HUMIGKAX (rearranged anti-myelin kappa chain), Genbank accession number M29469. The resulting MAbs were named hMAb BIWA 4 and BIWA 8, respectively. BIWA 8 contained two amino acids of the murine parent antibody within the light chain framework 2 while BIWA 4 did not contain murine residues in the framework.

5 Recombinant MAbs were stably expressed in dihydrofolate reductase deficient Chinese hamster ovary cells by electroporation with heavy and light chain expression plasmids. Cells were seeded into 96 well microtiter plates at densities of 500 and 100 cells/well in selection medium (α -MEM with 10% dialyzed fetal calf serum). When colonies became visible (after ~ 14 days), culture supernatants were tested for their IgG content by ELISA, and the best producers were
10 expanded. Gene amplification was performed by culturing in the presence of increasing concentrations of methotrexate (20-500 nM).

Laboratory scale production of chimeric and humanized MAbs was performed in a standard culture medium containing 1% fetal calf serum. IgG fractions were purified from tissue culture supernatants by affinity chromatography on protein A sepharose. Purity was
15 tested by SDS-PAGE and high performance size exclusion chromatography.

Evaluation of antibody affinity. Measurement of kinetic and affinity constants using recombinant antigen was performed on a BIAcore 2000 system (BIAcore AB, Uppsala, Sweden). A glutathione-S-transferase fusion protein containing domains v3-v10 of human
20 CD44 (GST/CD44v3-v10; 20 μ g/ml) was immobilized on a CM5 sensor chip by the amine coupling method according to the manufacturer's instructions, using 10 mM sodium acetate pH 5.0 as coupling buffer. 35 μ l of MAb at various concentrations (8-67 nM) in HBS (10 mM HEPES, pH 7.4, 150 mM NaCl, 3.4 mM EDTA, 0.05% BIAcore surfactant P20) were injected over the antigen-coated surface at a flow rate of 5 μ L/min. Dissociation of the MAb
25 was assessed for 5 minutes in buffer flow (HBS). Between two analyses, the surface of the chip was regenerated with a single pulse of 15 μ l 30 mM HCl. Analysis of the data and calculation of the kinetic constants were performed with BIAcore's BIAevaluation software, version 2.1. Association rates (k_a), dissociation rates (k_d), and dissociation constants (K_d) were assessed for all antibodies.

30 Relative binding affinities were also evaluated by competitive cell ELISA. Human A431 cells, originating from an epidermoid carcinoma of the vulva and known to express high levels of CD44v6, were seeded in 96 well tissue culture plates in 200 μ l per well RPMI 1640 with 10% fetal calf serum at a density of $2.5 - 5 \times 10^5$ cells/ml. The plates were incubated overnight at 37°C in a humidified incubator with 5% CO₂ in air. After removal of the medium the cells were

5 washed once with PBS, fixed with 96% ethanol for 1 min, and washed again with PBS. cMAb BIWA 2, hMAb BIWA 4 and hMAb BIWA 8 (prediluted to 10 µg/ml) were applied in 1:2 serial dilutions (8 steps) in 100 µl/well in PBS/0.5%BSA/0.05% Tween 20 (assay buffer) and incubated for 30 min at room temperature. 100 µl prediluted mMAb BIWA 1 (20 ng/ml) was added and the plates were incubated for 2 h at room temperature on an orbital shaker. Control
10 samples contained prediluted samples only, without BIWA 1 (0% control) or BIWA 1 only without any competing antibodies (100% control). After washing three times with PBS/0.05% Tween 20 (washing buffer), 100 µl of the secondary antibody (peroxidase-conjugated goat anti-mouse Fc, diluted 1:15,000 in assay buffer, DAKO Copenhagen, Denmark) was added for detection of mMAb BIWA 1, and plates were incubated for 1 h at room temperature on an
15 orbital shaker. After washing three times with washing buffer, the plates were developed with 100 µl/well tetramethylbenzidine substrate solution (Kierkegaard and Perry Laboratories, Gaithersburg, USA). The reaction was stopped after 15 min with 50 µl/well 1 M phosphoric acid. Absorbance was measured in an ELISA plate reader at 450 nm (reference 610-690 nm).

20 **Radioiodination of Antibodies.** Iodination of MAbs was performed essentially as described by Haisma *et al.* (1986), using either ^{125}I (100 mCi/mL) or ^{131}I (200 mCi/mL), both purchased from Amersham, Aylesbury, England. One mg MAb IgG dissolved in 500 µl PBS, pH=7.4, and 1 mCi ^{125}I or ^{131}I were mixed in a vial coated with 75 µg Iodogen (Pierce, Oud Bijerland, The Netherlands). After 5 minutes of incubation at room temperature, free iodine was removed
25 by gelfiltration on a PD10-column (Pharmacia-LKB, Woerden, The Netherlands). After removal of unbound ^{125}I or ^{131}I the radiochemical purity always exceeded 97% as determined by TLC and HPLC procedures which have been described before (Van Gog *et al.*, 1997a). No aggregates or fragments were formed as assessed by HPLC analysis.

30 **Preparation of Rhenium-186-labeled MAbs.** ^{186}Re -labeled MAbs were prepared according to a multistep procedure using the chelate S-benzoylmercaptoacyltriglycine (S-benzoyl-MAG3) as previously described (Van Gog *et al.*, 1997a). In this procedure a solid-state synthesis for the preparation of ^{186}Re -MAG3 is followed by esterification with 2,3,5,6-tetrafluorophenol (TFP) and conjugation of the reactive ^{186}Re -MAG3-TFP ester to the MAb.

5 After conjugation the ^{186}Re -labeled MAb was purified on a PD10-column. After removal of unbound ^{186}Re the radiochemical purity always exceeded 98%.

Binding-assay for radiolabeled antibodies. *In vitro* binding characteristics of the labeled MAbs used in the biodistribution and therapy studies were determined in an immunoreactivity assay essentially as described previously (Van Gog et al., 1997a). To test the binding of iodinated or ^{186}Re -labeled MAbs, UM-SCC-11B cells fixed in 0.1% glutaraldehyde were used. UM-SCC-11B cells were kindly provided by Dr. T.E. Carey, University of Michigan, Ann Arbor, MI. Five serial dilutions (ranging from 5×10^6 cells per tube to 3.1×10^5 cells per tube) were prepared with 1% BSA in PBS. Excess of unlabeled MAb IgG was added to a second tube with the lowest concentration of cells to determine non-specific binding. IgG labeled with 10,000 cpm of ^{125}I , ^{131}I or ^{186}Re was added to each tube and the samples were incubated overnight at 4°C. Cells were spun down, radioactivity in the pellet and supernatant was measured in a gamma counter (LKB-Wallace 1282 CompuGamma, Kabi Pharmacia, Woerden, The Netherlands), and the percentage of bound and free radioactivity was calculated. Data were graphically analyzed in a modified Lineweaver Burk plot and the immunoreactivity was determined by linear extrapolation to conditions representing infinite antigen excess.

Biodistribution studies in HNSCC-bearing nude mice. For the biodistribution experiments nude mice bearing subcutaneously implanted human HNSCC xenografts (HNX-OE) were used as described previously (Van Gog et al., 1997a). Female mice (Hsd: Athymic nu/nu, 25-32 g, Harlan CPB, Zeist, The Netherlands) were 8-10 weeks old at the time of the experiments. Three biodistribution experiments were conducted with mice bearing 1 or 2 tumors ranging from 30 to 470 mm³. In the first experiment, 10 µCi (50 µg) ^{131}I -labeled mMAb U36 were injected simultaneously with 10 µCi (50 µg) ^{125}I -labeled mMAb BIWA 1 in mice bearing tumors of 133 ± 28 mm³ (n=20 mice, 37 tumors). In the second experiment, 10 µCi (50 µg) ^{131}I -labeled hMAb BIWA 4 and 10 µCi (50 µg) ^{125}I -labeled cMAb BIWA 2 were co-injected in mice bearing tumors of 167 ± 31 mm³ (n=21 mice, 32 tumors). In the third experiment, 10 µCi (50 µg) ^{131}I -labeled hMAb BIWA 4 and 10 µCi (50 µg) ^{125}I -labeled hMAb BIWA 8 were co-injected in mice with tumors of 130 ± 21 mm³ (n=23 mice, 40 tumors). Conjugates were

5 intravenously (i.v.) injected in a volume of 100 μ l after dilution in 0.9% NaCl. To obtain a comparable blood/body clearance of the co-injected MAbs, only MAbs with an identical murine or human isotype were combined. The antibody dose (total dose 100 μ g per mouse) was chosen high enough to prevent rapid isotype-related elimination of the MAb from the blood (Sharkey et al., 1991, Van Gog et al., 1997b), and low enough to prevent antigen
10 saturation in the tumor.

At indicated time points after injection, mice were anaesthetized, bled, killed and dissected. Besides the tumors, the following organs were removed: liver, spleen, kidney, heart, stomach, ileum, colon, bladder, sternum, muscle, lung, skin and tongue. After weighing, radioactivity in tumors, blood and organs was counted in a dual-isotope gamma counter (LKB-
15 Wallace 1282 CompuGamma), with automatic correction for the ^{131}I -comptons in the ^{125}I window setting. Radioactivity uptake in these tissues was calculated as the percentage of the injected dose per gram of tissue (%ID/g).

Until the day of MAb administration mice were routinely housed under specific-pathogen-free conditions, in sterile cages in a humidity- and temperature controlled clean room, classification
20 2000 according to the Federal Standard 209d. On the day of injection, mice were transported to a Radio Nuclide Center, and sterile radioimmunoconjugates were administered under aseptic conditions in a laminar flow hood.

Radioimmunotherapy studies in nude mice. Animal RIT studies were performed to
25 compare the therapeutic efficacy of the different MAbs labeled with ^{186}Re . The immunoreactive fractions of the conjugates always exceeded 75 %. Three therapy experiments were conducted with mice bearing 1 or 2 HNX-OE tumors ranging from 45 to 195 mm^3 . The ^{186}Re doses were chosen at the maximum tolerated dose (MTD) level (i.e. 400 μCi) or lower (300 μCi). The MTD level is defined as the dose resulting in 5-15% body weight loss. In the first experiment
30 mice were given a single i.v. injection with either 300 μCi (100 μg) ^{186}Re -labeled mMAb U36 or 300 μCi (100 μg) ^{186}Re -labeled mMAb BIWA 1. In the second experiment either 300 μCi (100 μg) ^{186}Re -labeled hMAb BIWA 4 or 300 μCi (100 μg) ^{186}Re -labeled cMAb BIWA 2 were administered, and in the third experiment either 400 μCi (100 μg) ^{186}Re -labeled hMAb BIWA 4 or 400 μCi (100 μg) ^{186}Re -labeled hMAb BIWA 8. Average tumor volumes were

5 similar for all experimental groups. Experiment 1: $95 \text{ mm}^3 \pm 34 \text{ mm}^3$ ($n = 7$ mice, 12 tumors) for the ^{186}Re -mMAb U36 treated group, $91 \text{ mm}^3 \pm 15 \text{ mm}^3$ ($n = 7$ mice, 12 tumors) for the ^{186}Re -mMAb BIWA 1 treated group, and $99 \text{ mm}^3 \pm 54 \text{ mm}^3$ ($n = 6$ mice, 11 tumors) for the control group. Experiment 2: $101 \text{ mm}^3 \pm 35 \text{ mm}^3$ ($n = 7$ mice, 12 tumors) for the ^{186}Re -hMAb BIWA 4 treated group, $92 \text{ mm}^3 \pm 43 \text{ mm}^3$ ($n = 7$ mice, 12 tumors) for the ^{186}Re -cMAb BIWA 2 treated group, while the control group was the same as in experiment 1. Experiment 3: $105 \text{ mm}^3 \pm 43 \text{ mm}^3$ ($n = 8$ mice, 13 tumors) for the ^{186}Re -hMAb BIWA 4 treated group, $100 \text{ mm}^3 \pm 42 \text{ mm}^3$ ($n = 8$ mice, 13 tumors) for the ^{186}Re -hMAb BIWA 8 treated group, and $110 \text{ mm}^3 \pm 46 \text{ mm}^3$ ($n = 7$ mice, 11 tumors) for the control group. During treatment tumors were measured twice weekly and tumor volumes relative to the volume at the start of treatment were calculated. Toxicity was monitored by measurement of the body weight twice weekly. Mice were sacrificed when one of the tumors exceeded 1000 mm^3 .

Statistics. Differences in tissue uptake between co-injected MAbs were statistically analyzed for each time point with the Student's *t*-test for paired data. Differences in average tumor volume between the various RIT treatment groups were statistically analyzed for each time point with the Student's *t*-test for independent samples.

RESULTS

***In vitro* binding characteristics of the CD44v6-specific MAbs.** The binding affinities of the five MAbs were analyzed using recombinant antigen as well as human tumor cell lines. Kinetic and affinity constants were evaluated by surface plasmon resonance using GST/CD44v3-v10 as immobilized antigen. Table 1 shows the association rates (k_a), dissociation rates (k_d) and dissociation constants (K_d). mMAb BIWA 1 and cMAb BIWA 2, containing identical variable regions, have similar k_a , k_d , and K_d and show the highest affinity. In contrast, mMAb U36 and hMAb BIWA 4 have lower k_a and higher k_d , resulting in markedly lower dissociation constants (factors 35.0 and 10.5, respectively). hMAb BIWA 8, containing murine residues in the light chain framework region 2, shows a marked decrease of k_d resulting in increased affinity.

Table 1

Kinetics and affinity constants of MAbs directed against CD44v6.

Antibody	k_a ($M^{-1}s^{-1}$)	k_d (s^{-1})	K_d (M)	K_d relative to murine BIWA 1
Murine BIWA 1	1.3×10^5	4.2×10^{-5}	3.2×10^{-10}	1.0
Murine U36	1.5×10^4	1.7×10^{-4}	1.1×10^{-8}	35.0
Chimeric BIWA 2	1.7×10^5	4.1×10^{-5}	2.4×10^{-10}	0.7
Humanized BIWA 4	6.5×10^4	2.2×10^{-4}	3.4×10^{-9}	10.5
Humanized BIWA 8	7.5×10^4	6.3×10^{-5}	8.4×10^{-10}	2.6

5

The relative binding affinities of the cMAb and the hMAbs were also evaluated in a competitive cell ELISA using human A431 tumor cells (Fig. 1). In accordance with the affinity measurements on recombinant antigen, cMAb BIWA 2 was the most effective competitor, followed by hMAb BIWA 8 and hMAb BIWA 4. Similar results (not shown) were obtained

10 with two other human HNSCC cell lines (FaDu and LICR-LON-HN5).

15

Biodistribution in HNSCC-bearing nude mice. Biodistribution studies were performed in HNX-OE xenograft bearing nude mice. Two MAbs with identical murine or human isotype were labeled with either ^{125}I or ^{131}I and injected simultaneously (50 μ g, 10 μ Ci each). Each pair

15 of MAbs was selected to provide a stepwise decrease in the difference in affinities: mMAB U36 has a 35.0 fold lower affinity than mMAB BIWA 1 (experiment 1), hMAb BIWA 4 has a 14.0-fold lower affinity than cMAb BIWA 2 (experiment 2) and hMAb BIWA 4 has a 4.0-fold lower affinity than hMAb BIWA 8 (experiment 3). The immunoreactive fractions of all iodinated MAbs were at least 74% after extrapolation (Table 2).

5 **Table 2** Immunoreactive fraction of iodinated MAbs determined by binding to UM-SCC-11B cells.

Experiment no.	Antibody	Label	Binding to 5x10 ⁶ cells (%)	Binding Extrapolated ^a (%)
1	Murine U36	¹³¹ I	59.7	87.4
	Murine BIWA 1	¹²⁵ I	91.1	91.1
2	Humanized BIWA 4	¹³¹ I	77.4	82.3
	Chimeric BIWA 2	¹²⁵ I	80.5	79.9
3	Humanized BIWA 4	¹³¹ I	77.3	74.5
	Humanized BIWA 8	¹²⁵ I	91.8	92.1

^a Immunoreactivity was determined by linear extrapolation to conditions representing infinite antigen excess: see Materials and Methods

10 The biodistributions in experiment 1 were determined at day 1, 2, 3 and 7 after injection; biodistributions in experiments 2 and 3 were determined at day 1, 2, 4 and 7 days after injection. The calculated average %ID/g of tumor and blood of all three experiments are given in Table 3.

15 **Table 3** Tumor and blood levels of iodinated CD44v6-specific MAbs with different affinity after co-injection to HNX-OE bearing mice

		Conjugate A (%ID/g)		Conjugate B (%ID/g)		Conj. A:B ratio			
Exp. No		Time after	Tumor	Blood	Tumor	Blood	Tumor		
	Blood	injection							
20	1	Conj. A: ¹³¹ I-mMAb U36	1 d	15.7	17.9	13.0	17.8	1.2	1.0
		Conj. B: ¹²⁵ I-mMAb BIWA 1	2 d	18.4	15.0	13.4	14.9	1.4	1.0
			3 d	20.6	11.8	13.7	11.0	1.5	1.1
			7 d	16.5	7.1	7.8	4.8	2.1	1.5
25									

5	2. Conj. A: ^{131}I -hMAb BIWA 4	1 d	10.8	10.2	9.1	10.6	1.2	1.0
	Conj. B: ^{125}I -cMAb BIWA 2	2 d	12.4	10.2	9.8	10.1	1.3	1.0
		4 d	12.9	7.2	8.9	7.2	1.5	1.0
		7 d	7.6	3.3	4.8	2.7	1.6	1.2
10	3. Conj. A: ^{131}I -hMAb BIWA 4	1 d	10.5	11.1	9.5	11.4	1.1	1.0
	Conj. B: ^{125}I -hMAb BIWA 8	2 d	10.9	10.0	9.6	9.8	1.1	1.0
		4 d	11.7	6.2	9.6	5.9	1.2	1.0
		7 d	10.1	4.2	7.8	3.9	1.3	1.1

15 For each pair of co-injected MAbs the uptake ratios for tumor and blood are provided. The average %ID/g and s.e.m. of tumor, blood and various organs at 3 (experiment 1) or 4 days p.i. (experiments 2 and 3) are shown in Figure 2.

In a direct comparison of the two murine MAbs, tumor uptake of low affinity U36 was significantly higher than uptake of high affinity BIWA 1 at all time points ($p < 0.001$) (Table 3). In contrast, no significant differences were found between the uptake values of these MAbs in blood and normal tissues at 1, 2 and 3 days p.i.. At day 7 p.i., BIWA 1 levels in blood and most of the organs were significantly lower ($p < 0.05$) than U36 levels, indicating more rapid clearance of BIWA 1 from the blood/body. A 50% higher tumor uptake of U36 in comparison with BIWA 1 at day 3 p.i. is illustrated by Figure 2A.

25 Similar relationships were found in the evaluation of the two other MAb pairs. hMAb BIWA 4, while having the lower affinity, showed a significantly higher tumor uptake ($p < 0.001$) than cMAb BIWA 2 and hMAb BIWA 8 at all time points (Table 3). In contrast, MAb levels in blood and normal tissues were similar for these pairs of MAbs at 1, 2, and 4 days p.i.. At 7 days p.i., BIWA 2 and BIWA 8 levels in blood and most of the organs were significantly lower ($p < 0.05$) than BIWA 4 levels, indicating more rapid clearance of these MAbs from the blood/body. A 45% higher tumor uptake of BIWA 4 in comparison with BIWA 2 is illustrated by Figure 2B, while a 20% higher tumor uptake of BIWA 4 in comparison with BIWA 8 is illustrated by Figure 2C, for the 4 days post injection time points.

35 Consistent results were obtained from an additional experiment (data not shown) in which the radiolabels were exchanged: ^{125}I -BIWA 4 *versus* ^{131}I -BIWA 8 instead of ^{131}I -BIWA 4 *versus*

5 ¹²⁵I-BIWA 8. Data from this latter experiment rule out the possibility that the type of radiolabel had influenced the pharmacokinetic behavior of the labeled MAb.

Radioimmunotherapy in HNSCC-bearing nude mice. From the three biodistribution experiments it appeared that the low affinity MAbs showed a higher and more selective tumor uptake than the high affinity MAbs, and thus might be better suited for RIT. To test this possibility the following treatment groups were compared in RIT studies with HNX-OE xenograft bearing mice:

Experiment 1: 300 μ Ci ¹⁸⁶Re-U36 or 300 μ Ci ¹⁸⁶Re-BIWA 1 or saline as control. Experiment 2: 300 μ Ci ¹⁸⁶Re-BIWA 4 or 300 μ Ci ¹⁸⁶Re-BIWA 2 or saline as control. Experiment 3: 400 μ Ci ¹⁸⁶Re-BIWA 4 or 400 μ Ci ¹⁸⁶Re-BIWA 8 or saline as control.

15 In Figure 3, the mean relative tumor volume (as a percentage of the tumor volume at day 0) for the control and treatment groups is plotted against time. Tumors of mice in the control group in all three experiments showed exponential growth with a tumor volume doubling time of about 7 days. In the groups treated with the ¹⁸⁶Re-labeled MAbs, tumors stopped growing, in some cases accompanied by tumor regression, shortly after injection of the conjugates. However, all tumors ultimately regrew.

In experiment 1, administration of 300 μ Ci ¹⁸⁶Re-BIWA 1 resulted in a decrease of the tumor growth rate, but not in a reduction of the mean tumor size. Administration of 300 μ Ci ¹⁸⁶Re-U36, however, caused a reduction of the mean tumor volume from 185 mm³ to 120 mm³ between day 7 and day 17 post injection, after which tumors started growing again. The mean relative tumor volume in the ¹⁸⁶Re-U36-treated group was significantly smaller ($p < 0.001$) than that of the ¹⁸⁶Re-BIWA 1-treated group from day 14 on.

In experiment 2, administration of either 300 μ Ci ¹⁸⁶Re-BIWA 4 or 300 μ Ci ¹⁸⁶Re-BIWA 2 resulted in tumor growth arrest at day 7 with start of regrowth at day 17 p.i.. BIWA 4 was more effective in RIT than BIWA 2 from day 14 on, but a significant difference between the mean relative tumor volumes was only found at day 14 p.i. ($p < 0.05$).

30 In experiment 3, mice were treated with either 400 μ Ci ¹⁸⁶Re-BIWA 4 or BIWA 8, which resulted in a decrease of the relative tumor volume to a minimum of $80 \pm 62\%$ and $98 \pm 81\%$, respectively, at day 19. Thereafter, tumors started regrowth.

5 These data indicate that the low affinity MAb BIWA 4 is more effective in RIT than the high affinity Mabs cBIWA 2 and BIWA 8.

References

- 10 Günthert, U., Hofmann, M., Rudy, W., Reber, S., Zöller, M., Haußmann, I., Matzku, S., Wenzel, A., Ponta, H., and Herrlich, P. A new variant of glycoprotein CD44 confers metastatic potential to rat carcinoma cells. *Cell* 65: 13-24 (1991).
- 15 Haisma, H.J., Hilgers, J., and Zurawski, V.R. Jr. Iodination of monoclonal antibodies for diagnosis and therapy using a convenient one vial method. *J. Nucl. Med.*, 27: 1890-1895, 1986.
- 20 Heider, K.-H., Hofmann, M., Horst, E., van den Berg, F., Ponta, H., Herrlich, P., and Pals, S.T. A human homologue of the rat metastasis-associated variant of CD44 is expressed in colorectal carcinomas and adenomatous polyps. *J. Cell Biol.* 120: 227-233 (1993a).
- 25 Heider, K.-H., Dämmrich, J., Skroch-Angel, P., Müller-Hermelink, H.-K., Vollmers, H.-P., Herrlich, P., and Ponta, H. Differential expression of CD44 splice variants in intestinal- and diffuse-type human gastric carcinomas and normal gastric mucosa. *Cancer Res.* 53: 4197-4203 (1993b).
- 30 Heider KH, Mulder JWR, Ostermann E, Susani S, Patzelt E, Pals ST, Adolf GRA. Splice variants of the cell surface glycoprotein CD44 associated with metastatic tumor cells are expressed in normal tissues of humans and cynomolgus monkeys. *Eur. J. Cancer* 31A: 2385-2391, 1995.
- 35 Heider KH, Sproll M, Susani S, Patzelt E, Beaumier P, Ostermann O, Ahorn H, Adolf GRA. Characterization of a high affinity monoclonal antibody specific for CD44v6 as candidate for immunotherapy of squamous cell carcinomas. *Cancer Immunology Immunotherapy* 43: 245-253, 1996.
- 40 Himmler, A., Maurer-Fogy, I., Kronke, M., Scheurich, P., Pfizenmaier, K., Lantz, M., Olsson, I., Hauptmann, R., Stratowa, C., and Adolf, G.R. Molecular cloning and expression of human and rat tumor necrosis factor receptor chain (p60) and its soluble derivative, tumor necrosis binding protein. *DNA & Cell Biol.*, 9: 705-715, 1990.
- 45 Hofmann, M., Rudy, W., Zöller, M., Tölg, C., Ponta, H., Herrlich P., and Günthert, U. CD44 splice variants confer metastatic behavior in rats: homologous sequences are expressed in human tumor cell lines. *Cancer Res.* 51: 5292-5297 (1991).

- 5 Kugelman, L.C., Gangluly, S., Haggerty, J.G., Weissman, S.M., and Milstone, L.M. The core protein of epican, a heparan sulfate proteoglycan on keratinocytes, is an alternative form of CD44. *J. Invest. Dermatol.*, 99: 886-891, 1992.
- 10 Koopman, G., Heider, K.-H., Horts, E., Adolf, G. R., van den Berg, F., Ponta, H., Herrlich, P., Pals, S. T. Activated human lymphocytes and aggressive Non-Hodgkin's lymphomas express a homologue of the rat metastasis-associated variant of CD44. *J. Exp. Med.* 177: 897-904 (1993).
- 15 Rudy, W., Hofmann, M., Schwartz-Albiez, R., Zöller, M., Heider, K.-H., Ponta, H., Herrlich, P. The two major CD44 proteins expressed on a metastatic rat tumor cell line are derived from different splice variants: Each one individually suffices to confer metastatic behaviour. *Cancer Res.* 53: 1262-1268 (1993).
- 20 Screaton, G.R., Bell, M.V., Jackson, D.G., Cornelis, F.B., Gerth, U., and Bell, J. I. Genomic structure of DNA encoding the lymphocyte homing receptor CD44 reveals at least 12 alternatively spliced exons. *Proc. Natl. Acad. Sci. U.S.A.* 89: 12160-12164 (1992).
- 25 Sharkey, R.M., Natale, A., Goldenberg, D.M., and Mattes, M.J. Rapid blood clearance of immunoglobulin G2a and immunoglobulin G2b in nude mice. *Cancer Res.*, 51: 3102-3107, 1991.
- Tölg, C., Hofmann, M., Herrlich, P., and Ponta, H. Splicing choice from ten variant exons establishes CD44 variability. *Nucleic Acids. Res.* 21: 1225-1229 (1993).
- 30 Van Gog, F.B., Visser, G.W.M., Stroomeer, J.W.G., Roos, J.C., Snow, G.B., and Van Dongen, G.A.M.S. High dose ¹⁸⁶Re-labeling of monoclonal antibodies for clinical application: pitfalls and solutions. *Cancer*, 80: 2360-2370, 1997a.
- 35 Van Gog, F.B., Brakenhoff, R.H., Snow, G.B., and Van Dongen, G.A.M.S. Rapid elimination of mouse/human chimeric monoclonal antibodies in nude mice. *Cancer Immunol. Immunother.*, 44: 103-111, 1997b.
- 40 Wielenga, V. J. M., Heider, K.-H., Offerhaus, G. J. A., Adolf, G. R., van den Berg, F. M., Ponta, H., Herrlich, P., Pals, S.T. Expression of CD44 variant proteins in human colorectal cancer is related to tumor progression. *Cancer Res.* 53: 4754-4756 (1993).

5 Example 2 Details of sequences

This example shows the details of sequences, e.g. the position of cloning sites, leaders and untranslated regions.

10 Abbreviations:

aa = amino acids

nt = nucleotide sequence

SEQ ID No. 1 VH BIWA 4/8 aa

15 EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYDMSWVRQAPGKGLEWVSTISSGGSY
TYYLDSIKGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARQGLDYWGRGTLVTVSS

SEQ ID No. 2 VL BIWA 4 aa

20 EIVLTQSPATLSLSPGERATLSCSASSSINYIYWYQQKPGQAPRLLIYLTSNLASGVPAR
FSGSGSGTDFTLTISSELPEDFAVYYCLQWSSNPLTFGGGGTKVEIK

SEQ ID No. 3 VL BIWA 8 aa

EIVLTQSPATLSLSPGERATLSCSASSSINYIYWLQQKPGQAPRILYLTSNLASGVPARF
SGSGSGTDFTLTISSELPEDFAVYYCLQWSSNPLTFGGGGTKVEIK

25

SEQ ID No. 4 VH BIWA 4/8 nt

30 GAAGTGCAGCTGGTGGAGTCTGGGGGAGGCTTAGTGAAGCCTGGAGGGTCCCTAAGACTCTCC
TGTGCAGCCTCTGGATTCACTTTCAGTAGCTATGACATGTCTTGGGTTCGCCAGGCTCCGGGG
AAGGGGCTGGAGTGGGTCTCAACCATTAGTAGTGGTGGTAGTTACACCTACTATCTAGACAGT
ATAAAGGGCCGATTCACCATCTCCAGAGACAATGCCAAGAACTCCCTGTACCTGCAAATGAAC
AGTCTGAGGGCTGAGGACACGGCCGTGTATTACTGTGCAAGACAGGGGTGGACTACTGGGGT
CGAGGAACCTTAGTCACCGTCTCCTCA

SEQ ID No. 5 VL BIWA 4 nt

35

GAAATTGTTCTCACCCAGTCTCCAGCAACCCTGTCTCTGTCTCCAGGGGAGAGGGCCACCCTG
TCCTGCAGTGCCAGCTCAAGTATAAATTACATATACTGGTACCAGCAGAAGCCAGGACAGGCT
CCTAGACTCTTGATTTATCTCACATCCAACCTGGCTTCTGGAGTCCCTGCGCGCTTCAGTGGC
AGTGGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTGAAGATTTTGCCGTT
40 TATTACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAGATT
AAA

SEQ ID No. 6 VL BIWA 8 nt

45 GAAATTGTTCTCACCCAGTCTCCAGCAACCCTGTCTCTGTCTCCAGGGGAGAGGGCCACCCTG
TCCTGCAGTGCCAGCTCAAGTATAAATTACATATACTGGCTCCAGCAGAAGCCAGGACAGGCT
CCTAGAATCTTGATTTATCTCACATCCAACCTGGCTTCTGGAGTCCCTGCGCGCTTCAGTGGC
AGTGGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTGAAGATTTTGCCGTT
TATTACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAGATT
50 AAA

5 SEQ ID No. 7 heavy chain (variable + constant) BIWA 4/8 aa

EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYDMSWVRQAPGKGLEWVSTISSGGSY
 TYYLDSIKGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARQGLDYWGRGTLVTVSS
 ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
 10 SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEL
 LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKP
 REEQYNSTYRVVSVLTVHLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV
 YTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFL
 YSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

15

SEQ ID No. 8 light chain (variable + constant) BIWA 4 aa

EIVLTQSPATLSLSPGERATLSCSASSSINYIYWYQQKPGQAPRLLIYLTSLNLA SGV PAR
 FSGSGSGTDFTLTISSELPEDFAVYYCLQWSSNPLTFGGGGTKVEIKRTVAAPSVFIFPPS
 20 DEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLST
 LTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID No. 9 light chain (variable + constant) BIWA 8 aa

25 EIVLTQSPATLSLSPGERATLSCSASSSINYIYWLQQKPGQAPRILIIYLTSLNLA SGV PARF
 SSGSGSGTDFTLTISSELPEDFAVYYCLQWSSNPLTFGGGGTKVEIKRTVAAPSVFIFPPSD
 EQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTL
 TLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC

30 SEQ ID No. 10 heavy chain (variable + constant) BIWA 4/8 nt ; insert in pAD-CMV1/pAD-
 CMV19, contains introns (lower case) between CH1-hinge, hinge-CH2, CH2-CH3, leader
 sequence underlined, non-translated sequences in italic, cloning sites bold

aagctttgacagacgcacaaccctggactcccaagtctttctcttcagtgacaaacacagacataggatatcaca
 35 *tttgcttctgacacaactgtgttcactagcagcctcaaacagacacc*ATGAAC TTTGGGCTCAGCTTGATTTTCC
TTGTCCTAATTTTAAAGGTGTCCAGTGTGAAGTGCAGCTGGTGGAGTCTGGGGGAGGCTTAGTGAAGCCTGGAG
GGTCCCTAAGACTCTCCTGTGCAGCCTCTGGATTCACTTTCAGTAGCTATGACATGTCTTGGGTTCGCCAGGCTC
 CGGGGAAGGGGCTGGAGTGGGTCTCAACCATTAGTAGTGGTGGTAGTTACACCTACTATCTAGACAGTATAAAGG
 GCCGATTCAACATCTCCAGAGACAATGCCAAGAACTCCCTGTACCTGCAAATGAACAGTCTGAGGGGCTGAGGACA
 40 CGGCCGTGTATTACTGTGCAAGACAGGGGTGGACTACTGGGGTCGAGGAACCTTAGTCACCGTCTCCTCAGCTA
 GCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCT
 GCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCCCTGACCAGCGGCGTGCACA
 CCTTCCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGG
 GCACCCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTggtgagaggc
 45 cagcacagggagggaggggtgtctgtctggaagcaggctcagcgctcctgcctggacgcacatccggctatgcagccc
 cagtccagggcagcaaggcaggccccgtctgcctcttcacccggagcctctgcccgccccactcatgctcaggga
 gaggtcttcttggtttttcccaggctctgggcaggcagggctaggtgcccctaaccaggccctgcacacaaa
 ggggcagggtgctgggctcagacctgccaagagccatatccgggaggaccctgcccctgacctaaagcccaccccaa
 aggccaaactctccactccctcagctcggacaccttctctcctcccagattccagtaactcccaatcttctctct
 50 gcaGAGCCCAAATCTTG TGACAAAAC TCACACATGCCACCGTGCCCAggtaagccagcccaggcctcgccctcc
 agctcaaggcgggacagggtgccctagagtagcctgcacccagggaagggcccggtgctgacacgtccac
 ctccatctcttccctcaGCACCTGAAC TCTGGGGGACCGTCAGTCTTCTCTTCCCCCAAACCCAAGGACAC
 CCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTT
 CAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTA

5 CCGGGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAA
 CAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAggtgggacccgtgggggtgagggggccaca
 tggacagagggccggctcgccccaccctctgccctgagagtaccgctgtaccaacctctgtcctacaGGGCAGCC
 CCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCT
 10 GGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGAC
 CACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCA
 GCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCT
 GTCTCCGGGTAAATGAgtgcgacggccgcgaattc

15 SEQ ID No. 11 light chain (variable + constant) BIWA 4 nt ; sequence in pAD-CMV1/pAD-
 CMV19, leader sequence underlined, non-translated sequences in italic, cloning sites bold

aagcttgatcttcaggatatcacattttgcttctgacacaactgtgttctactagcaacctcaaacagacaccATGG
 ATTTTCAGGTGCAGATTTTCAGCTTCCTGCTAATGAGTGCCTCAGTCATAATGTCCAGGGGAGAAATTGTTCTCA
 CCCAGTCTCCAGCAACCCTGTCTGTCTCCAGGGGAGAGGGCCACCCTGTCTGCAGTGCCAGCTCAAGTATAA
 20 ATTACATATACTGGTACCAGCAGAAGCCAGGACAGGCTCCTAGACTCTTGATTATCTCACATCCAACCTGGCTT
 CTGGAGTCCCTGCGCGCTTCAGTGGCAGTGGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTG
 AAGATTTTGCCGTTTATTACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAGA
 TTAAACGGACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAACCTGCTA
 GCGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAGGTGGATAACGCCCTCCAAT
 25 CGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGC
 TGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCA
 CAAAGAGCTTCAACAGGGGAGAGTGTTAGgaattc

30 SEQ ID No. 12 light chain (variable + constant) BIWA 8 nt ; sequence in pAD-CMV1/pAD-
 CMV19, leader sequence underlined, non-translated sequences in italic, cloning sites bold

aagcttgatcttcaggatatcacattttgcttctgacacaactgtgttctactagcaacctcaaacagacaccATGG
 ATTTTCAGGTGCAGATTTTCAGCTTCCTGCTAATGAGTGCCTCAGTCATAATGTCCAGGGGAGAAATTGTTCTCA
 CCCAGTCTCCAGCAACCCTGTCTGTCTCCAGGGGAGAGGGCCACCCTGTCTGCAGTGCCAGCTCAAGTATAA
 35 ATTACATATACTGGCTCCAGCAGAAGCCAGGACAGGCTCCTAGAATCTTGATTATCTCACATCCAACCTGGCTT
 CTGGAGTCCCTGCGCGCTTCAGTGGCAGTGGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTG
 AAGATTTTGCCGTTTATTACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAGA
 TTAAACGGACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAACCTGCTA
 GCGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAGGTGGATAACGCCCTCCAAT
 40 CGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGC
 TGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCA
 CAAAGAGCTTCAACAGGGGAGAGTGTTAGgaattc

45 SEQ ID No. 13 heavy chain (variable + constant) BIWA 4/8 nt ; sequence in N5KG1val, no
 introns contained, leader sequence underlined

ATGGAGTTTGGGCTGAGCTGGCTTTTTCTTGTGGCTATTTTAAAAGGTGTCCAGTGTGAAGTG
 CAGCTGGTGGAGTCTGGGGGAGGCTTAGTGAAGCCTGGAGGGTCCCTAAGACTCTCCTGTGCA
 GCCTCTGGATTCACTTTCAGTAGCTATGACATGTCTTGGGTTCGCCAGGCTCCGGGGGAAGGGG
 50 CTGGAGTGGGTCTCAACCATTAGTAGTGGTGGTAGTTACACCTACTATCTAGACAGTATAAAG
 GGCCGATTCACCATCTCCAGAGACAATGCCAAGAACTCCCTGTACCTGCAAATGAACAGTCTG
 AGGGCTGAGGACACGGCCGTGTATTACTGTGCAAGACAGGGGTGGACTACTGGGGTTCGAGGA
 ACCTTAGTCACCGTCTCCTCAGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCC
 TCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAA
 55 CCGGTGACGGTGTGCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTC
 CTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGC
 ACCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTT

5 GAGCCCAAATCTTGTGACAAAACCTCACACATGCCCACCGTGCCCAGCACCTGAACTCCTGGGG
GGACCGTCAGTCTTCCTCTTCCCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCT
GAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTAC
GTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACG
TACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAG
10 TGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGG
CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAG
GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGC
AATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTC
TTCCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGC
15 TCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGT
AAATGA

SEQ ID No. 14 light chain (variable + constant) BIWA 4 nt ; sequence in N5KG1val, leader
sequence underlined

20 ATGGAAGCCCCAGCTCAGCTTCTCTTCCTCCTGCTGCTCTGGCTCCCAGATACCACCGGAGAA
ATTGTTCTCACCCAGTCTCCAGCAACCCTGTCTCTGTCTCCAGGGGAGAGGGCCACCCTGTCC
TGCAGTGCCAGCTCAAGTATAAATTACATATACTGGTACCAGCAGAAGCCAGGACAGGCTCCT
AGACTCTTGATTTATCTCACATCCAACCTGGCTTCTGGAGTCCCTGCGCGCTTCAGTGGCAGT
25 GGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTGAAGATTTTGCCGTTTAT
TACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAGATTAAA
CGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGA
ACTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGGCCAAAGTACAGTGGAAG
GTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGAC
30 AGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTC
TACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGA
GAGTGTTGA

SEQ ID No. 15 light chain (variable + constant) BIWA 8 nt ; sequence in N5KG1val, leader
sequence underlined

35 ATGGAAGCCCCAGCTCAGCTTCTCTTCCTCCTGCTGCTCTGGCTCCCAGATACCACCGGAGAA
ATTGTTCTCACCCAGTCTCCAGCAACCCTGTCTCTGTCTCCAGGGGAGAGGGCCACCCTGTCC
TGCAGTGCCAGCTCAAGTATAAATTACATATACTGGCTCCAGCAGAAGCCAGGACAGGCTCCT
40 AGAATCTTGATTTATCTCACATCCAACCTGGCTTCTGGAGTCCCTGCGCGCTTCAGTGGCAGT
GGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTGAAGATTTTGCCGTTTAT
TACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAGATTAAA
CGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGA
ACTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGGCCAAAGTACAGTGGAAG
45 GTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGAC
AGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTC
TACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGA
GAGTGTTGA

50 SEQ ID No. 16 BIWA 4 in N5KG1val

CTGACGCGCCCTGTAGCGGCGCATTAAGCGCGGGCGGGTGTGGTGGTTACGCGCAGCGTGACCG
CTACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTCGCTTTCTTCCCTTCCTTTCTCGCCACGT

5 TCGCCGGCTTTCCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTT
TACGGCACCTCGACCCCAAAAACTTGATTAGGGTGATGGTTCACGTAGGGTCGCGACGTACC
GGGCCCCCCTCGATTAATTAATCGAGCTACTAGCTTTGCTTCTCAATTTCTTATTTGCATAA
TGAGAAAAAAGGAAAATTAATTTTAACACCAATTCAGTAGTTGATTGAGCAAATGCGTTGCC
10 AAAAAGGATGCTTTAGAGACAGTGTTCTCTGCACAGATAAGGACAAACATTATTCAGAGGGAG
TACCCAGAGCTGAGACTCCTAAGCCAGTGAGTGGCACAGCATTCTAGGGAGAAATATGCTTGT
CATCACCGAAGCCTGATTCCGTAGAGCCACACCTTGGTAAGGGCCAATCTGCTCACACAGGAT
AGAGAGGGCAGGAGCCAGGGCAGAGCATATAAGGTGAGGTAGGATCAGTTGCTCCTCACATTT
GCTTCTGACATAGTTGTGCCAGCATGGAGGAATCGATCCTCCATGCTTGAACAAGATGGATTG
CACGCAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTATTCCGGCTATGACTGGGCACAACAGACA
15 ATCGGCTGCTCTGATGCCGCCGTGTTCCGGCTGTGAGCGCAGGGGCGCCCGGTTCTTTTTGTC
AAGACCGACCTGTCCGGTGCCCTGAATGAACTGCAGGTAAGTGCGGCCGCTCTAGGCCTCCAA
AAAAGCCTCCTCACTACTTCTGGAATAGCTCAGAGGCCGAGGCGGCCTCGGCCTCTGCATAAA
TAAAAAAAATTAGTCAGCCATGCATGGGGCGGAGAATGGGCGGAACTGGGCGGAGTTAGGGGC
GGGATGGGCGGAGTTAGGGGCGGGACTATGGTTGCTGACTAATTGAGATGCATGCTTTGCATA
20 CTCTGCTGCTGGGGAGCCTGGGGACTTTCCACACCTGGTTGCTGACTAATTGAGATGCATG
CTTTGCATACTTCTGCCTGCTGGGGAGCCTGGGGACTTTCCACACCCTAACTGACACACATTC
CACAGAATTAATTCCCCTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCA
TATATGGAGTTCCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGAC
CCCCGCCCATTTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCAT
25 TGACGTCAATGGGTGGACTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCAT
ATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAG
TACATGACCTTATGGGACTTTCCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACC
ATGGTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTTGACTCACGGGGATTT
CCAAGTCTCCACCCCATTTGACGTCAATGGGAGTTTGTTTTTGGCACCAAATCAACGGGACTTT
30 CCAAATGTGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAG
GTCTATATAAGCAGAGCTGGGTACGTGAACCGTCAGATCGCCTGGAGACGCCATCACAGATCT
CTCACCATGGAAGCCCCAGCTCAGCTTCTCTTCCCTCCTGCTGCTCTGGCTCCCAGATAACCACC
GGAGAAATTGTTCTCACCCAGTCTCCAGCAACCCTGTCTCTGTCTCCAGGGGAGAGGGCCACC
CTGTCTGACGTGCCAGCTCAAGTATAAATTACATATACTGGTACCAGCAGAAGCCAGGACAG
35 GCTCCTAGACTCTTGATTTATCTCACATCCAACCTGGCTTCTGGAGTCCCTGCGCGCTTCAGT
GGCAGTGGGTCTGGAACCGACTTCACTCTCACAATCAGCAGCCTGGAGCCTGAAGATTTTGCC
GTTTATTACTGCCTGCAGTGGAGTAGTAACCCGCTCACATTCGGTGGTGGGACCAAGGTGGAG
ATTAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAA
TCTGGAACCTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAG
40 TGGAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTCACAGAGCAGGACAGC
AAGGACAGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACAC
AAAGTCTACGCCTGCGAAGTCAC
CCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTGAATTCAGAT
CCGTTAACGGTTACCAACTACCTAGACTGGATTCTGTGACAACATGCGGCCGTGATATCTACGT
45 ATGATCAGCCTCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCCCTCCCCCGTGCC
TTCCTTGACCCTGGAAGGTGCCACTCCCCTGTCCTTTCTTAATAAAATGAGGAAATTGCATC
GCATTGTCTGAGTAGGTGTCATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGA
GGATTGGGAAGACAATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGAACCAGCTGGGAC
TAGTAGCTTTGCTTCTCAATTTCTTATTTGCATAATGAGAAAAAAGGAAAATTAATTTTAAC
50 ACCAATTCAGTAGTTGATTGAGCAAATGCGTTGCCAAAAAGGATGCTTTAGAGACAGTGTTCT
CTGCACAGATAAGGACAAACATTATTCAGAGGGAGTACCCAGAGCTGAGACTCCTAAGCCAGT
GAGTGGCACAGCATTCTAGGGAGAAATATGCTTGTCTACCCGAAGCCTGATTCCGTAGAGCC
ACACCTTGGTAAGGGCCAATCTGCTCACACAGGATAGAGAGGGCAGGAGCCAGGGCAGAGCAT
ATAAGGTGAGGTAGGATCAGTTGCTCCTCACATTTGCTTCTGACATAGTTGTGTTGGGAGCTT

5 GGATAGCTTGGACAGCTCAGGGCTGCGATTTTCGCGCCAAACTTGACGGCAATCCTAGCGTGAA
GGCTGGTAGGATTTTATCCCCGCTGCCATCATGGTTCGACCATTGAACTGCATCGTCGCCGTG
TCCCAAATATGGGGATTGGCAAGAACGGAGACCTACCTGGCCTCCGCTCAGGAACGAGTTC
AAGTACTTCCAAAGAATGACCACAACCTCTTCAGTGGAAGGTAAACAGAATCTGGTGATTATG
GGTAGGAAAACCTGGTTCTCCATTCTTGAGAAGAATCGACCTTTAAAGGACAGAATTAATATA
10 GTTCTCAGTAGAGAACTCAAAGAACCACACGAGGAGCTCATTTTCTTGCCAAAAGTTTGGAT
GATGCCTTAAGACTTATTGAACAACCGGAATTGGCAAGTAAAGTAGACATGGTTTGGATAGTC
GGAGGCAGTTCTGTTTACCAGGAAGCCATGAATCAACCAGGCCACCTTAGACTCTTTGTGACA
AGGATCATGCAGGAATTTGAAAGTGACACGTTTTTCCCAGAAATTGATTTGGGGAAATATAAA
CTTCTCCCAGAAATACCCAGGCGTCCTCTCTGAGGTCCAGGAGGAAAAAGGCATCAAGTATAAG
15 TTTGAAGTCTACGAGAAGAAAGACTAACAGGAAGATGCTTTCAAGTTCTCTGCTCCCCTCCTA
AAGCTATGCATTTTTTATAAGACCATGGGACTTTTGCTGGCTTTAGATCAGCCTCGACTGTGCC
TTCTAGTTGCCAGCCATCTGTTGTTTGCCCCCTCCCCCGTGCCTTCCTTGACCCTGGAAGGTGC
CACTCCCCTGTCTTTTCTAATAAAAATGAGGAAATTGCATCGCATTTGTCTGAGTAGGTGTCA
TTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAG
20 GCATGCTGGGGATGCGGTGGGCTCTATGGAACCAGCTGGGGCTCGAAGCGGCCGCTCCGGATA
TGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGT
ACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCA
TGGTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGAATCACGGGGATTTC
CAAGTCTCCACCCCATTTGACGTCAATGGGAGTTTGTGTTTGGCACCAAAATCAACGGGACTTTC
25 CAAAATGTCGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGG
TCTATATAAGCAGAGCTGGGTACGTCCTCACATTCAGTGATCAGCACTGAACACAGACCCGTC
GACATGGAGTTTGGGCTGAGCTGGCTTTTTCTTGTGGCTATTTTAAAAGGTGTCCAGTGTGAA
GTGCAGCTGGTGGAGTCTGGGGGAGGCTTAGTGAAAGCCTGGAGGGTCCCTAAGACTCTCCTGT
GCAGCCTCTGGATTCACTTTCAGTAGCTATGACATGTCTTGGGTTCGCCAGGCTCCGGGGAAAG
30 GGGCTGGAGTGGGTCTCAACCATTAGTAGTGGTGGTAGTTACACCTACTATCTAGACAGTATA
AAGGGCCGATTCACCATCTCCAGAGACAATGCCAAGAACTCCCTGTACCTGCAAATGAACAGT
CTGAGGGCTGAGGACACGGCCGT
GTATTACTGTGCAAGACAGGGGTGGACTACTGGGGTCGAGGAACCTTAGTCACCGTCTCCTC
AGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCTCCTCCAAGAGCACCTCTGGGGG
35 CACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAA
CTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCTTACAGTCCTCAGGACTCTA
CTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAA
CGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTGACAA
AACTCACACATGCCACCGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCCTCTT
40 CCCCCCAAACCAAGGACACCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGT
GGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCA
TAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAAGCTCCT
CACCGTCTTGACACAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGC
CCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGT
45 GTACACCTTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTGACCTGCCTGGT
CAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA
CTACAAGACCACGCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCAC
CGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCT
GCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGAGGATCCGTTAACGG
50 TTACCAACTACCTAGACTGGATTCGTGACAACATGCGGCCGTGATATCTACGTATGATCAGCC
TCGACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGCCCCCTCCCCCGTGCCTTCCTTGACC
CTGGAAGGTGCCACTCCCCTGTCTTCTTAATAAAAATGAGGAAATTGCATCGCATTTGTCTG
AGTAGGTGTCAATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAA
GACAATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGAACCAGCTGGGGCTCGACAGCGC

5 TCGGATCGCCTCGAGGCCGCTACTAACTCTCTCCTCCCTCCTTTTTCTCCTGCAGGACGAGGCAG
CGCGGCTATCGTGGCTGGCCACGACGGGCGTTCCTTGCGCAGCTGTGCTCGACGTTGTCACTG
AAGCGGGAAGGGACTGGCTGCTATTGGGCGAAGTGCCGGGGCAGGATCTCCTGTCATCTCACC
TTGCTCCTGCCGAGAAAGTATCCATCATGGCTGATGCAATGCGGGCGGCTGCATACGCTTGATC
CGGCTACCTGCCCATTTCGACCACCAAGCGAAACATCGCATCGAGCGAGCACGTACTCGGATGG
10 AAGCCGGTCTTGTCGATCAGGATGATCTGGACGAAGAGCATCAGGGGCTCGCGCCAGCCGAAC
TGTTCCGCCAGGCTCAAGGCGCGCATGCCCGACGGCGAGGATCTCGTCTGACCCATGGCGATG
CCTGCTTGCCGAATATCATGGTGGAAAATGGCCGCTTTTCTGGATTCATCGACTGTGGCCGGC
TGGGTGTGGCGGACCGCTATCAGGACATAGCGTTGGCTACCCGTGATATTGCTGAAGAGCTTG
GCGGCGAATGGGCTGACCGCTTCCTCGTGCTTTACGGTATCGCCGCTCCCGATTTCGCAGCGCA
15 TCGCCTTCTATCGCCTTCTTGACGAGTTCTTCTGAGCGGGACTCTGGGGTTCGAAATGACCGA
CCAAGCGACGCCCAACCTGCCATCACGAGATTTTCGATTCCACCGCCGCTTCTATGAAAGGTT
GGGCTTCGGAATCGTTTTCCGGGACGCCGGCTGGATGATCCTCCAGCGCGGGGATCTCATGCT
GGAGTTCTTCGCCCACCCCAACTTGTTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAG
CATCACAATTTTACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAAACT
20 CATCAATCTATCTTATCATGTCTGGATCGCGGCCGGCCGCACCGCGGTGGAGCTTTAATTAAG
GCGCGCCAGCTCCAGCTTTTGTTCCTTTAGTGAGGGTTAATTTTCGAGCTTGGCGTAATCATG
GTCATAGCTGTTTCCTGTGTGAA
ATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGG
GTGCCATAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTTCCAGTCGG
25 GAAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCGTA
TTGGGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCGTTCCGGCTGCGGCGAG
CGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAA
AGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGT
TTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGC
30 GAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTC
CTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGC
TTTCTCATAGCTCACGCTGTAGGTATCTCAGTTCCGGTGTAGGTCGTTCCGCTCCAAGCTGGGCT
GTGTGCACGAACCCCCCGTTTACGCCCCGACCGCTGCGCCTTATCCGGTAACCTATCGTCTTGAGT
CCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAG
35 CGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAA
GGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCT
CTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTA
CGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTTTGATCTTTTCTACGGGGTCTGACGCTCAGT
GGAACGAAAACCTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGA
40 TCCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTG
ACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCA
TAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCA
GTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGC
CAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTA
45 ATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTTCGCAACGTTGTTGCCA
TTGCTACAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCCGGTTCCC
AACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTC
CTCCGATCGTTGTGAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGC
ATAATTCTCTTACTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCA
50 AGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATA
ATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTTGGAAAACGTTCTTCGGGGCGAA
AACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAAACCCACTCGTGCACCCAACT
GATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATG
CCGCAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAAT

5 ATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGA
AAAATAAACAAATAGGGGTTCGCGGCACATTTCCCCGAAAAGTGCCACA

SEQ ID No. 17 pAD-CMV1, cloning sites in bold

10 TCGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTT
CCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCATTGACGTCAATAAT
GACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGC
CCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGC
CTGGCATTATGCCCAGTACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTAT
15 TACCATGGTGTATGCGGTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCTAAGTCT
CCACCCCATTTGACGTCAATGGGAGTTTGTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACTC
CGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAACTAG
AGAACCCACTGCTTAACTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCC**AAGCTT**CTGCAGGTGAC
ATCGATGGATCCGGTACCTCGAGCGC**GAAATTC**CTAGAGGATCTTTGTGAAGGAACCTTACTTCTGTGGTGTGAC
20 ATAATTGGACAACTACCTACAGAGATTTAAAGCTCTAAGGTAAATATAAAATTTTAAAGTGTATAATGTGTTAA
ACTACTGATTCTAATTGTTTGTGTATTTTAGATTCCAACCTATGGAAGTGAATGGGAGCAGTGGTGGAAATGC
CTTTAATGAGGAAAACCTGTTTTGCTCAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGACTCTCAACA
TTCTACTCCTCCAAAAAAGAAGAGAAAGGTAGAAGACCCCAAGGACTTTCCTTCAGAATTGCTAAGTTTTTTGAG
TCATGCTGTGTTTAGTAATAGAAGTCTTGCTTGCTTTGCTATTTACACCACAAAGGAAAAAGCTGCACTGCTATA
25 CAAGAAAATTATGAAAAATATTTGATGTATAGTGCCTTGACTAGAGATCATAATCAGCCATACCACATTTGTAG
AGGTTTTACTTGCTTTAAAAAACCTCCACACCTCCCCCTGAACCTGAAACATAAAATGAATGCAATTGTTGTTG
TTAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTTACAAATAAAGCATT
TTTCACTGCATTCTAGTTGTGGTTTTGTCCAACTCATCAATGTATCTTATCATGTCTGGATCAATTCTGAGAAAC
TAGCCTTAAAGACAGACAGCTTTGTTCTAGTCAGCCAGGCAAGCATATGTAAATAAAGTTCCTCAGGGAAGTGA
30 GTTAAAGATGTATCCTGGACCTGCCAGACCTGGCCATTCACGTAAACAGAAGATTCCGCCTCAAGTTCCGGTTA
ACAACAGGAGGCAACGAGATCTCAAATCTATTACTTCTAATCGGGTAATTAAAACCTTTCACTAAAACACGGAC
CCACGGATGTCACCCACTTTTCTTCCCCGGCTCCGCCCTTCTCAGTACTCCCCACCATTAGGCTCGCTACTCCA
CCTCCACTTCCGGGCGCGACACCCACGTGCCCTCTCCCACCCGACGCTAACCCCGCCCCCTGCCCGTCTGACCCCG
CCCACCACCTGGCCCCGCCCCGTTGAGGACAGAAGAAACCCCGGCGAGCCGAGCCAAGGCGGACGGGTAGACGC
35 TGGGGGCGCTGAGGAGTCGTCTCTACCTTCTCTGCTGGCTCGGTGGGGGACGCGGTGGATCTCAGGCTTCCGGA
AGACTGGAAGAACC GGCTCAGAACC GGCTTGTCTCCGCGGGGCTTGGGCGGCGGAAGAATGGCCGCTAGACGCGGA
CTTGGTGCGAGGCATCGCAGGATGCAGAAGAGCAAGCCCGCGGGAGCGCGCGGCTGTACTACCCGCGCCTGGA
GCGGCCACGCGGACTGGGCGGGGCGGCTGGTGGAGGCGGAGTCTGACCTCGTGGAGGCGGGGCTCTGATGT
TCAAATAGGATGCTAGGCTTGTGAGGCGTGGCCTCCGATTCACAAGTGGGAAGCAGCGCCGGGCGACTGCAATT
40 TCGCGCCAACTTGGGGGAAGCACAGCGTACAGGCTGCCTAGGTGATCGCTGCTGCTGTATGGTTCGACCGCTG
AACTGCATCGTCGCCGTGTCCAGAATATGGGCATCGGCAAGAACGGAGACCTTCCCTGGCCAATGCTCAGGTAC
TGGCTGGATTGGGTAGGGAAACCGAGGCGGTTGCTGAATCGGGTTCGAGCACTTGGCGGAGACGCGCGGGCCAA
CTACTTAGGGACAGTCATGAGGGGTAGGCCCCGGGCTGCTGCCCTTGCCCATGCCCGCGGTGATCCCCATGCTG
TGCCAGCCTTTGCCAGAGGCGCTCTAGCTGGGAGCAAAGTCCGGTCACTGGGCAGCACACCCCCCGGACTTGC
45 ATGGGTAGCCGCTGAGATGGAGCCTGAGCACACGTGACAGGGTCCCTGTAAACGCAGTGTCTCTAACTTTTCAG
GAACGAGTTCAAGTACTTCCAAAGAATGACCACACCTCCTCAGTGGAAAGGTAAACAGAACCTGGTGATTATGGG
CCGAAAACCTGGTTCTCCATTCTGAGAAGAATCGACCTTTAAAGGACAGAATTAATATAGTTCTCAGTAGAGA
GCTCAAGGAACCACCAAGGAGCTCATTTTCTTGCCAAAAGTCTGGACCATGCCTTAAACTTATTGAACAACC
AGAGTTAGCAGATAAAGTGGACATGGTTTGGATAGTTGGAGGCGAGTTCGGTTTACAAGGAAGCCATGAATCAGCC
50 AGGCCATCTCAGACTCTTTGTGACAAGGATCATGCAGGAATTTGAAAGTGACACGTTCTTCCAGAAATTGATTT
GGAGAAATATAAACTTCTCCAGAGTACCCAGGGGTCTTTCTGAAGTCCAGGAGGAAAAAGGCATCAAGTATAA
ATTTGAAGTCTATGAGAAGAAAGGCTAACAGAAAGATACTTGCTGATTGACTTCAAGTTCTACTGCTTTCTCCT
AAAATTATGCATTTTACAAGACCATGGGACTTGTGTTGGCTTTAGATCCTGTGCATCCTGGGCAACTGTTGTAC
TCTAAGCCACTCCCCAAAGTCATGCCCCAGCCCCTGTATAATTCTAAACAATTAGAATTATTTTCATTTTCATTA
55 GTCTAACCAGGTATATTAATATACTTTAAGAAACACCATTTGCCATAAAGTTCTCAATGCCCTCCCATGCAG
CCTCAAGTGGCTCCCCAGCAGATGCATAGGGTAGTGTGTGTACAAGAGACCCCAAGACATAGAGCCCCCTGAGAG
CATGAGCTGATATGGGGGCTCATAGAGATAGGAGCTAGATGAATAAGTACAAAGGGCAGAAATGGGTTTTAACCA
GCAGAGCTAGAACTCAGACTTTAAAGAAAATTAGATCAAAGTAGAGACTGAATTATTCTGCACATCAGACTCTGA
GCAGAGTTCTGTTCACTCAGACAGAAATGGGTAAATTGAGAGCTGGCTCCATTGTGCTCCTTAGAGATGGGAGC
60 AGGTGGAGGATTATATAAGGTCTGGAACATTTAACTTCTCCGTTTCTCATCTTCAGTGAGATTCCAAGGGATACT
ACAATTCTGTGGAATGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAGAAGTATGCAAAGC

5 ATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAGAAGTATGCAAAGCATG
CATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCCATCCCGCCCCCTAACTCCGCCCCAGTTCCGCC
CATTCTCCGCCCCCATGGCTGACTAATTTTTTTTTTATTTATGCAGAGGCCGAGGCGCCTCTGAGCTATTCAGAAAGT
AGTGAGGAGGCTTTTTTGGAGGCCCTAGGCTTTTGCAAAAAAGCTAATTCAGCCTGAATGGCGAATGGGACGCGCC
10 CTGTAGCGGCGCATTAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGC
GCCCGCTCCTTTTCGCTTTCTTCCCTTCTTCTCGCCACGTTCCGCGGCTTTCCCCGTCAAGCTCTAAATCGGGG
GCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAAGTTGATTAGGGTGATGGTTCACGT
AGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTG
TTCCAAACTGGAACAACACTCAACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCC
15 TATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTTTAACAAAATATTAACGTTTACAATTTCA
GGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCG
CTCATGAGACAATAACCCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGT
GTCGCCCTTATTCCCTTTTTTTCGCGCATTTTGCCTTCTGTTTTTGCTCAGCCAGAAACGCTGGTGAAAGTAAAA
GATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGT
TTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATT
20 GACGCCGGGCAAGAGCAACTCGGTGCGCGCATACACTATTCTCAGAAATGACTTGGTTGAGTACTCACCAGTCACA
GAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAACACTGCG
GCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTA
ACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTA
GCAATGGCAACAACGTTGCGCAAACTATTAAGTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGAC
25 TGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAA
TCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTA
GTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTG
ATTAAGCATTGGTAACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAGCTTCATTTTTTAATTT
AAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAATCCCTTAACGTGAGTTTTCGTTCCACTGA
30 GCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAA
ACAAAAAACCACCGCTACCAGCGGTGGTTTGGTTGCCGGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAAGT
GGCTTCAGCAGAGCGCAGATACCAATACTGTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCT
GTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTT
ACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGGCTGAACGGGGGGTTTCGTGCACACAG
35 CCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCATTGAGAAAGCGCCACGCTTCCC
GAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGG
GGAAACGCCTGGTATCTTTATAGTCCTGTGCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCG
TCAGGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCC

40 SEQ ID No. 18 pAD-CMV19, cloning sites in bold

TCGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTT
CCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCGCCCATTGACGTCAATAAT
GACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGC
45 CCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGC
CTGGCATTATGCCAGTACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTAT
TACCATGGTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTTGACTCACGGGGATTTCCAAGTCT
CCACCCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCAAAATCAACGGGACTTTCCAAATGTGCTAACAACCTC
CGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCG
50 TCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCGCGG
CCGGGAACGGTGCAATTGGAACGCGGATTCCCCGTGCCAAGAGTCAGGTAAAGTACCGCCTATAGAGAAGACTCTTG
GGTTTCTGATAGGCACTGACTCTCTGCTTATTGGTCTATTTTCCCACCCCTTAGGCTGCTGGTGCTTAACTGGC
TTATCGAAATTAATACGACTCACTATAGGGAGACCCA**AGCTT**CTGCAGGTGACATCGATGGATCCGGTACCTCG
AGCGCG**GAATT**CTCTAGAGATATCTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATT
55 TCACAAATAAAGCATTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAAACTCATCAATGTATCTTATCATGTCT
GGATCAATTCTGAAAACTAGCCTTAAAGACAGACAGCTTTGTTCTAGTCAGCCAGGCAAGCATATGTAAATAAA
GTTCTCAGGGAACTGAGGTTAAAGATGTATCCTGGACCTGCCAGACCTGGCCATTACAGTAAACAGAAGATTC
CGCCTCAAGTTCCGGTTAACAACAGGAGGCAACGAGATCTCAAATCTATTACTTCTAATCGGGTAATTAAACCT
TTCAACTAAAACACGGACCCACGGATGTCACCCACTTTTCTTCCCCGGCTCCGCCCTTCTCAGTACTCCCCACC
60 ATTAGGCTCGCTACTCCACCTCCACTTCCGGGCGCGACACCCACGTGCCCTCTCCACCCGACGCTAACCCCGCC
CCTGCCCGTCTGACCCCGCCACCTGGCCCCGCCCGTTGAGGACAGAAGAAACCCGGGCAGCCGCAGCCA
AGGCGGACGGGTAGACGCTGGGGGCGCTGAGGAGTCGTCTCTACCTTCTCTGCTGGCTCGGTGGGGGACGCGGT

5 GGATCTCAGGCTTCCGGAAGACTGGAAGAACCGGCTCAGAACCGCTTGTCTCCGCGGGGCTTGGGCGGCGGAAGA
ATGGCCGCTAGACGCGGACTTGGTGCGAGGCATCGCAGGATGCAGAAGAGCAAGCCCGCGGGAGCGCGCGGCTG
TACTACCCCGCGCCTGGAGCGGCCACGCCGACTGGGCGGGGCGGCTGGTGGAGGCGGAGTCTGACCTCGTGG
AGGCGGGGCTCTGATGTTCAAATAGGATGCTAGGCTTGTGAGGCGTGGCCTCCGATTACAAAGTGGGAAGCAG
CGCCGGGCGACTGCAATTTGCGGCCAACTTGGGGGAAGCACAGCGTACAGGCTGCCTAGGTGATCGCTGCTGCT
10 GTCATGGTTCGACCGCTGAACTGCATCGTCGCCGTGTCCAGAATATGGGCATCGGCAAGAACGGAGACCTTCCC
TGGCCAATGCTCAGGTACTGGCTGGATTGGGTAGGGAAACCGAGGCGGTTGCTGAATCGGGTCGAGCACTTGG
CGGAGACGCGCGGGCCAACTACTTAGGGACAGTCATGAGGGGTAGGCCCGCGGCTGCTGCCCTTGCCCATGCCC
GCGGTGATCCCCATGCTGTGCCAGCCTTTGCCAGAGGCGCTCTAGCTGGGAGCAAAGTCCGGTCACTGGGCAGC
ACCACCCCCCGGACTTGCATGGGTAGCCGCTGAGATGGAGCCTGAGCACACGTGACAGGGTCCCTGTAAACGCAG
15 TGTTCCTCTAACTTTAGGAACGAGTTCAAGTACTTCCAAAGAATGACCACCACCTCCTCAGTGGAAAGGTAAACA
GAACCTGGTGATTATGGGCCGGAACCTGGTTCTCCATTCTGAGAAGAATCGACCTTTAAAGGACAGAATTAA
TATAGTTCTCAGTAGAGAGCTCAAGGAACCACCACAAGGAGCTCATTTTCTTGCCAAAAGTCTGGACCATGCCTT
AAAACCTTATTGAACAACCAGAGTTAGCAGATAAAGTGGACATGGTTTGGATAGTTGGAGGCAGTTCGTTTACAA
GGAAGCCATGAATCAGCCAGGCCATCTCAGACTCTTTGTGACAAGGATCATGCAGGAATTTGAAAGTGACACGTT
20 CTTCCCAGAAATTGATTTGGAGAAATATAAATCTCTCCAGAGTACCCAGGGGTCTTTCTGAAGTCCAGGAGGA
AAAAGGCATCAAGTATAAATTTGAAGTCTATGAGAAGAAAGGCTAACAGAAAGATACTTGCTGATTGACTTCAAG
TTCTACTGCTTTCTCTAAATTTATGCATTTTACAAGACCATGGGACTTGTGTTGGCTTTAGATCCTGTGCAT
CCTGGGCAACTGTTGTACTCTAAGCCACTCCCCAAAGTCATGCCCCAGCCCCGTGATAATTCTAAACAATTAGAA
TTATTTTCATTTTCATTAGTCTAACCAGGTTATATTAAATATACTTTAAGAAACACCATTTGCCATAAAGTTCTC
25 AATGCCCCCTCCCATGCAGCCTCAAGTGGCTCCCCAGCAGATGCATAGGGTAGTGTGTGTACAAGAGACCCCAAAG
ACATAGAGCCCCTGAGAGCATGAGCTGATATGGGGGCTCATAGAGATAGGAGCTAGATGAATAAGTACAAAGGGC
AGAAATGGGTTTTAACCAGCAGAGCTAGAACTCAGACTTTAAAGAAAATTAGATCAAAGTAGAGACTGAATTATT
CTGCACATCAGACTCTGAGCAGAGTTCTGTTCACTCAGACAGAAAATGGGTAAATTGAGAGCTGGCTCCATTGTG
CTCCTTAGAGATGGGAGCAGGTGGAGGATTATATAAGGTCTGGAACATTTAACTTCTCCGTTTCTCATCTTCAGT
30 GAGATTCCAAGGGATACTACAATTCTGTGGAATGTGTGTGTCAGTTAGGGTGTGGAAAGTCCCAGGCTCCCAGCA
GGCAGAAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCAGGCTCCCAGCAGGC
AGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCCCCCTA
ACTCCGCCCAGTTCCGCCCATTTCTCCGCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCGCCT
CTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTAATTCAGCCTGAA
35 TGGCGAATGGGAAATTGTAAACGTTAATATTTTGTAAATTCGCGTTAAATTTTGTAAATCAGCTCATTTTTT
TAACCAATAGGCCGAAATCGGCAAAATCCCTTATAAATCAAAGAATAGACCGAGATAGGGTTGAGTGTGTTCC
AGTTTGGAAACAAGAGTCCACTATTAAAGAACGTGGACTCCAACGTCAAAGGGCGAAAAACCGTCTATCAGGGCGA
TGGCCCACTACGTGAACCATCACCTAATCAAGTTTTTGGGGTGCAGGTGCCGTAAAGCACTAAATCGGAACCCCT
AAAGGGAGCCCCGATTTAGAGCTTGACGGGGAAAGCCGGCGAACGTGGCGAGAAAGGAAGGAAGAAAGCGAAA
40 GGAGCGGGCGCTAGGGCGCTGGCAAGTGTAGCGGTACGCTGCGCGTAACCACCACACCCGCGCGCTTAATGCG
CCGCTACAGGGCGCGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTCTAAATA
CATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAGGAAGAGTATG
AGTATTCACATTTCCGTGTGCCCCCTATTCCCTTTTTTGCGGCATTTTGCCTTCTGTTTTTGTCTACCCAGAA
ACGCTGGTGAAAGTAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAAGTGGATCTCAACAGC
45 GGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGC
GCGGTATTATCCCGTATTGACGCCGGGCAAGAGCAACTCGGTGCGCGCATACTATTCTCAGAATGACTTGGTT
GAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGTGCCATAACC
ATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCCTTTTTTGCAC
AACATGGGGGATCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGT
50 GACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAACCTATTAACCTGGCGAACTACTTACTCTAGCTTCC
CGGCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGC
TGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTGCGGTATCATTGCAGCACTGGGGCCAGATGGT
AAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCT
GAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTGACACCAAGTTTACTCATATATACTTTAGATTGATTAA
55 AAACCTTCATTTTTAATTTAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAATCCCTTAACGT
GAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGC
GTAATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTACCAACT
CTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCTTCTAGTGTAGCCGTAGTTAGGC
CACCCTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGT
60 GCGGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGCTGAACG
GGGGGTTGCTGCACACAGCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCATTGA
GAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGC

5 ACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTCGGGTTTCGCCACCTCTGACTTGAGCGT
CGATTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCAGCTGC

Example 3 Clinical study 1170.1

10 1. LIST OF ABBREVIATIONS AND DEFINITION OF TERMS

ADCC	Antibody Dependent Cell-mediated Cytotoxicity
AE	Adverse Event
ALT	Alanine Amino Transferase
a.p.	Anterior posterior
AP	Alkaline phosphatase
AST	Aspartate Amino Transferase
AUC	Area Under the concentration-time Curve
Bq	Becquerel, SI unit for radioactivity Becquerel (1Bq = one decay /s)
BSA	Bovine Serum Albumin
CD44v6	CD44 variant isoform v6
CDS	Corporate Drug Safety
cGy	Centi Gray measure for radioactivity
CHO	Chinese Hamster Ovary cells
Ci	Curie; unit for radioactivity; $1 \text{ Ci} = 37 \times 10^9 \text{ decays / s} = 37 \text{ GBq}$
CL	Total body clearance
cMAb	Chimeric Monoclonal Antibody
cm	Centimetre
$C_{\max}$	Maximum drug concentration observed
cpm	Count per minute
CRF	Case Report/Record Form
CT (scan)	Computed Tomography
CTC	Common Toxicity Criteria
CV	Coefficient of Variation
DLT	Dose Limiting Toxicity
ECG	Electrocardiogram
ELISA	Enzyme-Linked Immuno-Sorbent Assay
ENT	Ear Nose Throat
f	Female
FDA	Food and Drug Administration
g	Gram
GBq	SI unit for radioactivity Giga Becquerel ($1 \text{ GBq} = 10^9 \text{ decays / s}$)
GCP	Good Clinical Practice

GGT	Gamma Glutaryl Transpeptidase
GMP	Good Manufacturing Practice
Gy	Gray
HAHA	Human-Anti-Human-Antibody
Hb	Haemoglobin
HER2	Human Epidermal growth factor Receptor 2
hMAb	Humanised Monoclonal Antibody
HNSCC	Head and Neck Squamous Cell Carcinoma
HPLC	High Performance Liquid Chromatography
hr / h	Hour
hrs	Hours
Ht	Haematocrit
¹³¹ I	Iodine-131 (half-life 8.05 days)
ICH	International committee on harmonisation
ID	Injected Dose
IEC	Independent Ethics Committee
IgG	Immunoglobulin G
INN	International Non proprietary Name
IRB	Institutional Review Board
ITT	Intent-To-Treat
i.v.	Intravenous(ly)
KeV	Kilo electron volt
l / L	Litre
m	Male
m ²	Square metres
MAb	Monoclonal Antibody
MAG2GABA-TFP	Mercaptoacetylglcylglycyl-gamma-aminobutyrate-tetrafluorophenol ester (MAG2GABA-TFP), chelate used for coupling ¹⁸⁶ Re to monoclonal antibodies (future method)
MAG3	Mercaptoacetyltriglycine. Chelate used for coupling of ^{99m} Tc and ¹⁸⁶ Re to monoclonal antibodies
MBq	SI unit for radioactivity; Mega Becquerel (1 MBq = 10 ⁶ decays/s)
mCi	Milli Curie; unit for radioactivity; 1 mCi = 37 million decays / s = 37 MBq
MCV	Mean Corpuscular Volume
mGy	Milli Gray
µg	Microgram
mg	Milligram
min	Minute

ml / mL	Millilitre
mMAb	Murine Monoclonal Antibody
mmHg	Millimetre mercury
μmol	Micromol
mmol	Millimol
MRI	Magnetic Resonance Imaging
MRT	Mean Residence Time
mSv	Milli Sievert
MTD	Maximum Tolerated Dose
NA / n.a.	Not Applicable
NCI	US National Cancer Institute
ND	Not Done
ng	Nanogram
No / N	Number
nos	Not other specified
NSCLC	Non Small Cell Lung Cancer
n.y.r.	Not yet recovered
p.a.	Posterior anterior
PBS	Phosphate Buffered Saline
p.i.	Post infusion
p.o.	Per os
PP	Per Protocol
Pt. / Pat	Patient
Pts.	Patients
¹⁸⁶ Re	Rhenium-186, radionuclide; half-life of 3.7 days (tissue penetration of β-particles of about 1.2 mm)
Recov.	Recovered
RES	Reticulo Endothelial System
RIS	Radioimmunosintigraphy
RIT	Radioimmunotherapy
ROI	Region of Interest
SAE	Serious Adverse Event
SCC	Squamous Cell Carcinoma
sCD44v6	Soluble CD44v6
SD	Stable Disease
sGOT	Serum Glutamic Oxalacetic Transaminase
sGPT	Serum Glutamic Pyruvic Transaminase

SOC	System Organ Class
SOP	Standard Operating Procedure
SPECT	Single Photon Emission Computed Tomography
$t_{1/2}$	Elimination half life
^{99m}Tc	Technetium 99m (half-life 6 hours)
TLC	Thin Layer Chromatography
$T_{\max}$	Time point at which the maximum drug concentration is observed
TNM	Tumour Node Metastasis system for staging tumours
TSH	Thyroid Stimulating Hormone
UICC	Union Internationale Contre le Cancer (international union against cancer)
unk	Unknown
V_{ss}	Apparent volume of distribution under steady-state conditions
V_z	Apparent volume of distribution during the terminal phase
WBC	White Blood Count
WHO	World Health Organisation

5

2. Study Objectives

2.1 GENERAL AIM / CLINICAL OBJECTIVE

The general aim of the present study was to assess the safety and tolerability of intravenously administered ^{99m}Tc and ^{186}Re -labelled hMAb BIWA 4, to confirm preferential accumulation in the tumour of ^{99m}Tc -labelled hMAb BIWA 4, to determine the maximum tolerated radiation dose of ^{186}Re -labelled hMAb BIWA 4 and to propose a safe dose for phase II development. In order to reach this goal the study was divided into two parts:

Part A:

Objectives:

- To determine the safety and tolerability of a single infusion of intravenously administered ^{99m}Tc -labelled hMAb BIWA 4 in patients with advanced squamous cell carcinoma of the head and neck.
- To determine the biodistribution of a single infusion of ^{99m}Tc -labelled hMAb BIWA 4 at different BIWA 4 dose levels in patients with advanced squamous cell carcinoma of the head and neck.
- To study the pharmacokinetics of a single infusion of ^{99m}Tc -labelled hMAb BIWA 4.

Part B:

Objectives

- 5 • To determine the qualitative and quantitative toxic effects of ^{186}Re -labelled hMAb BIWA 4 and to study the predictability, onset, duration, intensity, reversibility and dose-relationship of the toxic side effects.
- To determine the maximum tolerated radiation dose of intravenously administered ^{186}Re -labelled hMAb BIWA 4 in head and neck cancer patients.
- 10 • To study the pharmacokinetics of intravenously administered ^{186}Re -labelled hMAb BIWA 4 in patients with squamous cell carcinoma of the head and neck.
- A secondary objective was to determine the preliminary therapeutic effects of ^{186}Re -labelled hMAb BIWA 4.

15 The following objectives outlined in the protocol could not be addressed during the performance of the trial. The development of the programme with the linker chelate mercaptoacetyltriglycine (MAG3), used for coupling of $^{99\text{m}}\text{Tc}$ and ^{186}Re to monoclonal antibodies was discontinued and thus the trial was finished. The further development with BIWA 4 will continue by using the linker mercaptoacetylglycylglycyl-gamma-aminobutyrate-tetrafluorophenol ester (MAG2GABA-TFP).

- 20 • The MTD for single dose treatment was to be identified first before more patients were entered to define the MTD for a second treatment with ^{186}Re -labelled hMAb BIWA 4.
- To propose a safe dose for first and consecutive infusions with ^{186}Re -labelled hMAb BIWA 4 for further studies.
- To obtain initial results on a dose schedule for repeated dosing.

25 2.2 PRIMARY VARIABLES

- Safety: clinical laboratory tests, human-anti-human-antibody (HAHA) assessments, vital signs measurements and adverse events.
- Efficacy: biopsy biodistribution data (Part A only) and radioimmunoscentigraphic images (Part A and B including dosimetry for Part B of the trial).
- 30 • Pharmacokinetic results.

2.3 SECONDARY VARIABLES

A secondary variable for the study was tumour response (Part B only).

3. INVESTIGATIONAL PLAN

3.1 OVERALL STUDY DESIGN AND PLAN – DESCRIPTION

The trial was performed in two parts. Part A evaluated the optimal dose of cold BIWA 4 and quantified tumour uptake while Part B investigated the maximum tolerated dose of ^{186}Re -BIWA 4.

3.1.1 Part A:

This part of the clinical trial was an uncontrolled, rising dose sequential group study. It was designed to provide initial data on the safety and tolerability of a single infusion of $^{99\text{m}}\text{Tc}$ -labelled hMAb BIWA 4, to investigate the pattern and level of biodistribution and to establish the pharmacokinetic profile of hMAb BIWA 4 in patients with head and neck cancer. Three protein doses of hMAb BIWA 4 were used in this Part A of the study with three patients planned to be treated at each dose level.

Patients were routinely investigated by the Department of Otolaryngology to determine the extent of the tumour. This included physical examination, computed tomography (CT) or MRI scans of the head and neck and panendoscopy (optional). During these procedures samples of tissues, suspect of tumour, were taken and investigated for the presence of squamous cell carcinoma. Based on the results of these investigations, the patients were destined to undergo surgery including neck dissection.

Radiolabelled antibody was injected and the patient was observed for occurrence of adverse events. Radioimmunoscintigraphic scans were performed 21 hours post infusion (p.i.) prior to surgery. Patients underwent surgery 48 hours (hrs) after the infusion of the radioactively labelled hMAb BIWA 4. The pathologist investigated the neck dissection specimen to determine the exact tumour load. Moreover, the amount of $^{99\text{m}}\text{Tc}$ in biopsies from tumour site(s) and normal tissues in the surgical specimen was measured. Tumour sites and tumour infiltrated nodes were examined for the presence of CD44v6 antigen by immunohistochemical techniques. The first three patients were administered 2 mg hMAb BIWA 4 labelled with 20 mCi $^{99\text{m}}\text{Tc}$ combined with 23 mg unlabelled hMAb BIWA 4. The second group of three evaluable patients was administered 2 mg hMAb BIWA 4 labelled with 20 mCi $^{99\text{m}}\text{Tc}$ combined with 48 mg unlabelled hMAb BIWA 4 and the third group was administered 2 mg of the labelled antibody combined with 98 mg of unlabelled antibody.

Pharmacokinetic assessments were done at the specified timepoints.

3.1.2 Part B:

This part of the clinical trial was an open uncontrolled, dose escalation study. It was designed to assess the safety and tolerability of ^{186}Re -labelled hMAb BIWA 4, to determine the maximum tolerated dose (MTD) of intravenously administered ^{186}Re -labelled hMAb BIWA 4 and to determine the preliminary therapeutic effects of ^{186}Re -labelled hMAb BIWA 4 in head

5 and neck cancer patients for whom no curative options were available. In addition the pharmacokinetic profile of ^{186}Re -labelled hMAb BIWA 4 was assessed.

All patients entering into this part of the trial received the dose of hMAb BIWA 4, which was selected basing on the results of Part A of the study. The hMAb BIWA 4 was labelled with escalating doses of ^{186}Re . At the lower dose levels (toxicity observed did not exceed grade 1) 10 two patients were entered per dose group and at the higher levels ($\geq$ grade 2 toxicity) a minimum of three patients were entered per dose group. All patients were evaluated to determine the safety of the administered hMAb BIWA 4.

15 Patients were routinely investigated by the Department of Otolaryngology to determine the extent of the tumour. This included physical examination and CT or MRI scanning of tumour locations.

^{186}Re -labelled antibody was injected with escalating radiation doses: patients in the first dose group received a radiation dose of 20 mCi/m^2 , after which the dose for patients in the subsequent dose groups was escalated by 10 mCi/m^2 increments until the MTD was reached. Patients were observed for occurrence of adverse events. Radioimmunoscentigraphic scans 20 were performed.

3.1.3 STUDY PROCEDURES AT EACH VISIT

3.1.3.1 Visit Schedule

a) Screening visit

Before entry into the study each patient was screened for eligibility. Demographics and 25 medically relevant history were recorded, concomitant therapy was recorded and a general physical examination was done and the body weight was measured. The Karnofsky performance score was determined. Written informed consent had to be obtained. A pregnancy test was required for women with childbearing potential. A 12-lead electrocardiogram (ECG) was made and blood as well as urine safety laboratory assessments were done. A blood sample 30 for HAHA-testing was also taken. A Chest X-ray was made and a full disease assessment (CT/MRI, ear-nose-throat [ENT] examination) was performed.

At the end of this visit the results of the required investigations were evaluated, the inclusion and exclusion criteria verified and arrangements for surgery (Part A only) were made.

b) Visit 2: Study days 1-7

35 Parts A and B:

On the first study day, which was not more than 3 weeks after the screening visit, the patient was admitted to the hospital. All required baseline laboratory assessments not obtained at the screening visit had to be done and evaluated prior to the infusion and the eligibility criteria had to be met. The body weight was measured. All concomitant therapy had to be recorded. Every

- 5 adverse event starting after the signing of the written informed consent had to be recorded in the patient file and the case report form (CRF).

5 On the day of treatment, before antibody administration a blood sample was collected for pharmacokinetics (including the assessment of soluble CD44v6) and the vital signs were recorded. Urine was collected from 0-4 hrs, 4-8 hrs, 8-12 hrs and 12-24 hrs during the first 24 hrs and then in 24 hrs intervals up to 48 and 96 hrs post infusion for Part A and B, respectively.

10 The antibody was administered at the department of Nuclear Medicine or in a designated room in the hospital. The antibody had to be administered through a peripheral upper extremity vein as proximally as possible. In all cases, the infusion was given over 5 minutes, after a 10-mL saline flush, through a freely flowing line. In order to obtain reproducible pharmacokinetic results a syringe pump had to be used. Therefore, dilution of the volume with NaCl 0.9 % was
15 required to 20 mL. This antibody infusion was followed by a 10 mL saline flush. Any suspected extravasation was documented in the CRF and the patient was imaged.

An emergency unit available with resuscitation equipment, anti-histamines, corticosteroids and epinephrine was within reach to counteract possible anaphylaxis. Adverse events and changes in the concomitant therapy were recorded daily.

20 Part A:

Vital signs were recorded at 10, 60 and 120 minutes p.i. Blood samples for safety and soluble CD44v6 were collected 21, 48 and 144 hrs post infusion.

Blood samples for pharmacokinetics were drawn at the end of the infusion and at 5, 10, 30 minutes and 1, 2, 4, 16, 21, 48, 72 hrs after the end of the infusion, and on day 7 (144-hrs p.i. including a serum sample for HAHA and soluble CD44v6 assessment).
25

Urine for pharmacokinetics was collected from 0-4 hrs, 4-8 hrs, 8-12 hrs and 12-24 hrs during the first 24 hrs and a 24-hr sample until 48 hrs p.i.

Whole body scintigraphic images were made directly after the infusion and 21 hours after the infusion. Twenty-one hours after infusion, in addition to the whole body image, single photon
30 emission computed tomography (SPECT) and planar images of the head and neck region were made.

The patient was operated at 48 hrs p.i. and stayed in the hospital for post-operative care.

On day 7 (144 hrs p.i.) safety urine samples were collected.

Adverse events and changes in the concomitant therapy were recorded daily.

35 Part B:

Vital signs were recorded at 10, 60, 120 and 240 minutes post infusion. Blood samples for safety and for soluble CD44v6 were collected 21, 48 and 144 hrs post infusion. A blood sample for HAHA assessment was collected 144 hrs after the infusion. A urine sample for safety was also collected 144 hrs p.i.

5 Blood samples for pharmacokinetics were drawn at the end of the infusion and at 5 and 30 minutes and 1, 2, 4, 16, 21, 48, 72 hours after the end of the infusion and on day 7 (144-hrs p.i.) after infusion. The originally planned sample at 10 minutes was omitted as per amendment 1.

10 Urine for pharmacokinetics was collected from 0-4 hrs, 4-8 hrs, 8-12 hrs and 12-24 hrs during the first 24 hrs and in 24-hr samples until 96 hrs p.i.

Whole body scintigraphic images were made directly after, at 21, 48, 72 and 144 hrs p.i. and optionally at two weeks p.i. if counting statistics permitted.

Planar images of the head and neck region were made at 21, 48 (optional), 72 and 144 hrs p.i. and optionally at two weeks p.i., if counting statistics permitted.

15 SPECT imaging was only performed seventy-two hours after the infusion.

Adverse events and changes in the concomitant therapy were recorded daily.

Patients were allowed to leave the hospital after three days.

c) Visit 3: Follow-up visits

Part A:

20 Patients visited the outpatient clinic six weeks after the infusion. A physical examination was performed, safety blood and urine samples collected, body weight and vital signs measured and adverse events recorded. Blood samples for pharmacokinetics, HAHA assessment and soluble CD44v6 were also obtained. A pregnancy test was required for women with childbearing potential. Follow-up of adverse events was recorded. Any changes in the concomitant therapy
25 were recorded.

Part B:

Patients visited the outpatient clinic weekly for at least six weeks for recording of adverse events and collection of safety blood samples. Blood samples for pharmacokinetics were drawn at 240 hours and 336 hours after infusion. The originally planned sample at 6 weeks p.i. was
30 omitted as per amendment 1. Any changes in the concomitant therapy were also recorded. The disease assessment performed at baseline was repeated six weeks after the infusion and thereafter if indicated (the assessment was done after six weeks rather than after four weeks as mentioned once in the protocol). Six weeks after the infusion blood samples for safety, HAHA assessment and soluble CD44v6 were collected, vital signs recorded and body weight
35 measured. A physical examination was done during this visit. A urine sample for safety was collected. A pregnancy test was required for women with childbearing potential.

5 d) Visits 4 and 5: second treatment (Part B)

A second treatment as outlined in the original protocol was not performed due to the premature termination of the trial because of the linker change. Instead patients who responded to the first dose of ^{186}Re -BIWA 4 were eligible for a second administration. They underwent the same visit schedule as for the first administration.

10 **3.2 DISCUSSION OF STUDY DESIGN, INCLUDING THE CHOICE OF CONTROL GROUPS**

15 The aim of the present study was to assess the preferential accumulation of $^{99\text{m}}\text{Tc}$ -labelled BIWA 4 in the tumour (Part A) and to evaluate the maximum tolerated dose of ^{186}Re -BIWA 4 (Part B) as well as the pharmacokinetics of BIWA 4 in patients suffering from advanced head and neck cancer (Part A and B).

20 An open design was used as is general practice in these types of Phase I trials in oncology. A minimum of two patients in the lower radiation dose tiers and three patients in the higher radiation dose tiers were included in Part B of the trial. In case of occurrence of drug-related Common Toxicity Criteria (CTC) grade 4 haematology and grade 3 non-haematology toxicity a further three patients were treated with the respective dose tier (for further dosing details refer to section 3.4.4. and 3.4.5).

The dose of BIWA 4 administered was based on previous results with mMAb BIWA 1 indicating that a dose of 50 mg yielded a high and selective uptake in tumour tissue with a low uptake in non-tumour tissues and the results from Part A of the trial (see also section 3.4.4.1).

25 The starting dose level for radioactivity chosen was based on previous data suggesting that a dose of 20 mCi/m² may be a safe dose (see also section 3.4.4.2).

The criteria for efficacy which applied as well as the criteria for assessing tolerability are well-established for this patient population and can also be evaluated in an open design.

3.3 SELECTION OF STUDY POPULATION

30 **3.3.1 Inclusion criteria**

- Patients with histological confirmation of squamous cell carcinoma in the head and neck.
- Patients destined for surgery by means of a neck dissection (Part A)

or:

- 35
- Patients with either local and/or regional recurrent disease for which curative treatment options were not available, or distant metastases. The tumour deposits had to be measurable either clinically or by one or more radiological technique(s) (CT, MRI, bone scintigraphy).

5 Because RIT was expected to be more effective in smaller size tumour deposits, patients with lesions measuring < 3 cm in greatest dimension were preferred (Part B).

- Patients over 18 years of age.
- Patients younger than 80 years of age.
- Patients who had given "written informed consent".
- 10 - Patients with a life expectancy of at least 3 months.
- Patients with a good performance status: Karnofsky > 60.

3.3.2 Exclusion criteria

- 15 - Life-threatening infection, allergic diathesis, organ failure (bilirubin > 30 $\mu\text{mol/l}$ and/or creatinine > 150 $\mu\text{mol/l}$) or evidence of a recent myocardial infarction on ECG or unstable angina pectoris.
- Pre-menopausal women (last menstruation $\leq$ 1 year prior to study start):
 - Not surgically sterile (hysterectomy, tubal ligation) and
 - Not practising acceptable means of birth control, (or not planned to be continued throughout the study). Acceptable methods of birth control include oral, implantable or
- 20 - injectable contraceptives.
- Women with a positive serum pregnancy test at baseline.
- Chemotherapy or radiotherapy within 4 weeks before inclusion in the study.
- White blood cell count < 3000/ mm^3 , granulocyte count < 1500/ mm^3 or platelet count < 100,000 / mm^3 .
- 25 - Haematological disorders, congestive heart failure, bronchial asthma, alimentary or contact allergy, severe atopy or allergy.

3.3.3 Removal of subjects from therapy or assessment

3.3.3.1 Criteria for Stopping Subject Treatment

30 The infusion had to be terminated immediately if the patient developed tachycardia (pulse rate greater than 120 per minute), hypotension (blood pressure less than 100 mm Hg systolic), respiratory distress, chest pain, or any symptoms intolerable to the patient.

3.3.3.2 Dropouts and Withdrawals

5 The subjects were free to discontinue their participation in this study at any time.

Evaluation of radiolabelled hMAb BIWA 4 was considered not to be feasible if the patient was prematurely removed from the study because of voluntary withdrawal. A case was considered not evaluable if adequate follow-up information was not available. Any unevaluable patients were planned to be replaced.

10 If the patient discontinued early from the study, the reason had to be documented on the CRF. If a patient did not return for the post-infusion blood samples for HAHA determination or pharmacokinetics, the reason had to be documented. If a patient developed a serious adverse event the study schedule had to be followed as closely as possible depending on the serious adverse events (SAE).

15 3.4 TREATMENTS

3.4.1 Treatments administered

Patients in Part A were administered ^{99m}Tc -BIWA 4 at a radioactivity dose of 20 mCi. The dose of BIWA 4 administered was 25 mg, 50 mg or 100 mg for three patients each. The drug was administered intravenously as a single dose.

20 Patients in Part B received 50 mg BIWA 4 labelled with Rhenium 186. The lowest radioactivity dose was 20 mCi/m² which was increased in dose tiers of 10 mCi/m². The trial drug was administered intravenously as a single dose.

3.4.2 Identity of investigational product

Part A:

25	Substance (INN):	BIWA 4 (bivatuzumab)
	Pharmaceutical form:	solution for injection
	Chiffre number:	BIWA 4 LOI 99 1D 1A
	Batch number:	B981101
	Source:	Boehringer Ingelheim Pharma KG
30	Unit strength:	5 mg / mL
	Daily dose:	25 mg
	Duration of use:	single dose
	Route of administration:	intravenous
	Posology:	infusion over five minutes

35	Substance (INN):	BIWA 4 (bivatuzumab)
	Pharmaceutical form:	solution for injection
	Chiffre number:	BIWA 4 LOI 99 1D 1A
	Batch number:	B981101
	Source:	Boehringer Ingelheim Pharma KG
40	Unit strength:	5 mg / mL

Daily dose:	50 mg
Duration of use:	single dose
Route of administration:	intravenous
Posology:	infusion over five minutes

Substance (INN):	BIWA 4 (bivatuzumab)
10 Pharmaceutical form:	solution for injection
Chiffre number:	BIWA 4 LOI 99 1D 1A
Batch number:	B981101
Source:	Boehringer Ingelheim Pharma KG
Unit strength:	5 mg / mL
15 Daily dose:	100 mg
Duration of use:	single dose
Route of administration:	intravenous
Posology:	infusion over five minutes

20 BIWA 4 was administered as radioconjugate linked with ^{99m}Tc . Linker molecule was MAG3. MAG3 was purchased from Mallinckrodt, Petten, The Netherlands.

^{99m}Tc was ordered locally via the laboratory where the radioimmunoconjugate was prepared (laboratory of Prof. Dr. van Dongen, Section Tumor Biology, Department of Otorhinolaryngology / Head and Neck Surgery, *Vrije Universiteit* University Medical Center, De Boelelaan 1117, 1081 HV Amsterdam, The Netherlands).

25 Part B:

Substance (INN):	BIWA 4 (bivatuzumab)
Pharmaceutical form:	solution for injection
Chiffre number:	BIWA 4 LOI 99 1D 1A
Batch number:	B981101
30 Source:	Boehringer Ingelheim Pharma KG
Unit strength:	5 mg / mL
Daily dose:	50 mg
Duration of use:	single dose
Route of administration:	intravenous
35 Posology:	infusion over five minutes

BIWA 4 was administered as radioconjugate linked with ^{186}Re . Linker molecule was MAG3. MAG3 was purchased from Mallinckrodt, Petten, The Netherlands.

¹⁸⁶Re was ordered locally via the laboratory where the radioimmunoconjugate was prepared (laboratory of Prof. Dr. van Dongen, Section Tumor Biology, Department of
40 Otorhinolaryngology / Head and Neck Surgery, *Vrije Universiteit* University Medical Center, De Boelelaan 1117, 1081 HV Amsterdam, The Netherlands).

3.4.2.1 Characteristics and quality of the trial drug

5 Antibody characteristics

The antigen recognised by hMAb BIWA 4 is a transmembrane glycoprotein located on the outer cell surface and is only to a small extent internalised (< 20 %). Further analysis revealed that hMAb BIWA 4 recognises an epitope encoded by variant exon v6 of CD44. The antigen was shown to be expressed by all primary head and neck tumours (n=54) and by the majority
10 of cells within these tumours. A comparable expression was observed for sixty-eight tumour infiltrated lymph nodes from neck dissection specimens (R97-2054). The reactivity pattern of hMAb BIWA 4 in human normal tissues is provided in Appendix III of the protocol. Reactivity of hMAb BIWA 4 was found to be essentially restricted to squamous epithelia. As demonstrated by the previous RIS study with murine monoclonal antibody (mMAb) BIWA 1,
15 reactivity with normal squamous epithelium was not a limiting factor for utility in tumour targeting with respect to tumour uptake.

Quality control of ^{99m}Tc - or ^{186}Re -labelled hMAb BIWA 4

The antibody was labelled with ^{99m}Tc or ^{186}Re according to a method described by Fritzberg *et al.* (R96-2106: See protocol Appendix IV) which was modified according to Visser *et al.*
20 (R96-2094) and Van Gog *et al.* (R96-2111).

The procedures for radiolabelling hMAb BIWA 4 with ^{99m}Tc and ^{186}Re had been validated with respect to the final quality of the prepared conjugate. In five independent labelling experiments, performed according to the procedure described in appendix IV of the protocol, the percentage of label bound to the antibody was found to be 96-99%.

25 In this clinical investigation the radiochemical purity of each ^{99m}Tc - or ^{186}Re -labelled antibody batch prepared was assessed by thin layer chromatography (TLC) or high performance liquid chromatography (HPLC) and passage through a PD-10 gel filtration column, and had to be more than 90% to allow administration to a patient.

The immunoreactive fraction of each $^{99m}\text{Tc}/^{186}\text{Re}$ -labelled antibody batch was checked after
30 administration by use of validated methods and had to be above 60%. In brief, analyses were essentially performed according to a procedure as described by Lindmo *et al.* (R96-2104). UM-SCC 11B cells (human larynx carcinoma), containing the CD44v6 antigen, were fixed in 0.1% glutaraldehyde. Six dilutions, ranging from 5×10^6 cells per tube to 3.1×10^5 cells per tube, were made with 1% bovine serum albumine (BSA) in phosphate buffered saline (PBS).

35 To the tubes, 80,000 counts per minute (cpm) of the ^{99m}Tc - or 10,000 of the ^{186}Re -labelled hMAb BIWA 4 were added and incubated overnight at room temperature.

To the last sample, excess unlabelled hMAb BIWA 4 was added to determine non-specific binding. Cells were spun down and the radioactivity in pellet and supernatant were determined in a gamma counter and the percentage bound and free radiolabelled MAb calculated (LKB-
40 Wallac 1218 CompuGamma).

5 Data were graphically analysed in a modified Lineweaver Burk plot and the immunoreactive fraction was determined by linear extrapolation to conditions representing infinite antigen excess.

In case the discharge level of 60% was not reached in this binding assay the preparation for infusion was re-evaluated in a second binding assay. For this purpose the antibody preparation
10 was labelled with ^{131}I . The immunoreactive fraction in at least one of both assays, had to be larger than 60% for each patient to be evaluable and for the study to continue with the next patient.

Antibody Safety

15 BIWA 4 used for radiolabelling with $^{99\text{m}}\text{Tc}$ and ^{186}Re in this study is a well-characterised monoclonal antibody product which was produced by chinese hamster ovary (CHO)-cell culture fermentation. A master cell bank was established under Good Manufacturing Practice (GMP) conditions and thoroughly examined for microbiological status (bacteria, fungi, mycoplasma) as well as viral status (adventitious viruses, retroviruses). With the exception of endogenous retroviruses, which were known to be present in most CHO cells, no contaminants
20 were detected.

In accordance with the Food and Drug Administration (FDA) 1994 "Points to consider" document on MAbs used in clinical phase I studies in cancer patients, a standardised downstream purification process, which has been validated for efficient virus removal, was applied to the purification of BIWA 4 material. Four model viruses were included in the
25 validation (MuLV, PsRV, REO-3, SV 40). The concentrated bulk harvest was examined for the retrovirus titer to ensure that the subsequent downstream purification process was capable of adequately removing the retrovirus. Data on removal of endotoxins were within the acceptance limits (≤ 0.01 EU/mg) and were available for the bulk product. Furthermore pyrogenicity testing according to European Pharmacopoea guidelines had been successfully
30 performed.

The final hMAb BIWA 4 to be used for further radiolabelling was analysed in great detail and was proven to be a highly pure (SDS page, isoelectric focusing), sterile solution containing minimal endotoxin levels (≤ 0.01 EU/mg). Results of testing for pyrogenicity were conform to European Pharmacopoea standard. The equality of the BIWA 4 product from the manufacture
35 for pre-clinical and clinical supplies was demonstrated by analytical results.

Radiolabelling of hMAb BIWA 4 with $^{99\text{m}}\text{Tc}$ or ^{186}Re was performed by the department of Otolaryngology and Head & Neck Surgery of the *Vrije Universiteit* University Medical Center according to Standard Operating Procedures (SOPs). Sterility of the final product was guaranteed. Absence of endotoxins was tested during validation runs. For drug administration
40 to a patient in the University Hospital Nijmegen the proper amount of radiolabelled hMAb BIWA 4 was transported in a special container directly after preparation to the study centre in Nijmegen. Twenty-four hours were allowed between labelling of the compound and administration to the patient.

5 See also appendix 16.1.6 for an allocation of the individual Rhenium batches to the patients.

3.4.2.2 Packaging, Labelling and Supply

10 BIWA 4 was supplied by Boehringer Ingelheim The Netherlands. It was produced by Boehringer Ingelheim, Germany using a GMP manufacturing and purification process and filled in vials as a sterile, non-pyrogenic solution containing 25 mg hMAb BIWA 4 in 5 mL isotonic PBS, pH 7.2. Examples of the vial labels of the native and also of the labelled antibody were included in the Clinical Trial Manual.

One batch with a total amount of 27 g of hMAb BIWA 4 had been prepared and filled into vials. Labelling of hMAb BIWA 4 with ^{99m}Tc or ^{186}Re was performed in a class B certified nuclear laboratory of the *Vrije Universiteit* University Medical Center, Amsterdam.

15 3.4.2.3 Storage Conditions

The unlabelled hMAb BIWA 4 had to be stored in the hospital pharmacy in a limited access area for study materials at a monitored temperature between +2 and +8°C.

3.4.3 Method of assigning subjects to treatment groups

20 No randomisation was used. Patients were assigned to the different dose groups according to the sequence of inclusion.

3.4.4 Selection of doses in the study

3.4.4.1 Selection of the BIWA 4 doses

Part A:

25 Since hMAb BIWA 4 had never been administered to patients before, it was essential to be informed about its safety and biodistribution before starting RIT trials. Its biodistribution might strongly depend on the MAb dose used for tumour targeting and would need careful consideration. On the basis of MAb protein dose escalation studies with the low affinity anti-CD44v6 mMAb U36 and the high affinity anti-CD44v6 mMAb BIWA 1, it was anticipated that the optimal dose was in the range of 25-100 mg, with 50 mg being optimal for mMAb
30 U36.

To confirm that this dose also was adequate for the intermediate affinity of hMAb BIWA 4 and to obtain initial information on the variability of the tumour uptake the biodistribution was assessed at 25, 50 and 100 mg total hMAb BIWA 4 and three evaluable patients were planned to be treated at each of these dose levels.

35 All evaluable patients received a single intravenous infusion of 2 mg hMAb BIWA 4 labelled with 20 mCi ^{99m}Tc , measured by a radiation calibration system just prior to administration. Patients scheduled to receive 25 mg, 50 mg or 100 mg received 23 mg, 48 mg or 98 mg,

5 respectively of unlabelled hMAb BIWA 4 administered together with this 2 mg hMAb BIWA 4 labelled with 20 mCi ^{99m}Tc .

Selection of the BIWA 4 dose for Part B:

10 On theoretical grounds a hMAb BIWA 4 dose of 50 mg was calculated as most suited for development. Part A of this study was performed to confirm the tumour preferential uptake of hMAb BIWA 4 at the three dose levels tested (25 mg, 50 mg and 100 mg). It was expected that the tumour uptake (expressed as percent injected dose per kilogram, % ID/kg) and tumour to non-tumour uptake ratio for these three dose levels would not differ much. If this was the case the hMAb BIWA 4 dose to be used for Part B of the study was 50 mg. However, if there was a clinically relevant difference between these dose levels, which favoured one over the other levels, the dose level with the best pattern of biodistribution would have been selected.

In this case a clinically relevant difference was defined as a difference of the tumour (mean of the tumour uptake) to bone marrow uptake ratio (mean of the bone marrow uptake = the cellular fraction and the supernatant) of more than 50 %.

Finally the dose selected was 50 mg hMAb BIWA 4 basing on the results of Part A.

20 3.4.4.2 Selection of the radiation starting dose

Part B:

The hMAb BIWA 4 dose selected in Part A of the study was labelled with escalating doses of ^{186}Re .

25 It has been reported that the maximum tolerated dose of ^{186}Re -labelled murine MAb NR-LU-10 was 90 mCi/m² for heavily pre-treated patients, while dose-limiting myelosuppression was observed at 120 mCi/m². For chimeric MAb NR-LU-13, recognising the same antigen as NR-LU-10, reversible myelosuppression occurred at 60 mCi/m². In a study with cMAb U36, performed at the *Vrije Universiteit* University Medical Center dose limiting myelosuppression was observed at 41 mCi/m².

30 In the light of the above-mentioned results a radiation dose of 20 mCi/m² administered as ^{186}Re -labelled hMAb BIWA 4 was considered a safe starting dose.

The dose was escalated with 10 mCi/m² increments. Two evaluable patients were entered at the lower dose levels. When $\geq$ grade 2 drug-related toxicity according to the CTC was observed a minimum of three patients were treated per dose level.

5 Before entering patients at a next higher dose level it had to be sure that the patients at the ongoing dose level did not experience dose limiting toxicity (DLT) defined as: drug-related CTC grade 3 non-haematologic toxicity or drug-related CTC grade 4 haematologic toxicity excluding nausea and vomiting without adequate antiemetic treatment. For this purpose all patients at such an ongoing dose level had to be observed long enough to ensure that possibly
10 induced toxicity was reversible.

When at the ongoing dose level one patient experienced DLT the number of patients treated at that dose level was increased to a total of six patients maximum. When 1 out of 6 patients experienced DLT the dose was escalated to the next level. When two or more of the patients experienced DLT, the next lower dose level was expanded (if not previously done) to a total of
15 six patients in order to establish MTD and a safe recommended dose for phase II.

It was originally planned that in case of acceptable toxicity (less than two patients with DLT) at that dose level additional patients could be entered who received this dose and an escalating, lower second dose. This plan was not pursued due to the change of the linker. MAG3 will be replaced by MAG2GABA-TFP, a comparable linker, but with a more convenient linkage
20 procedure. The present development programme with MAG3 has been finished.

Alternatively patients who responded to and tolerated the first administration of ^{186}Re -BIWA 4 were eligible for a second administration with a dose of 50 mCi/m² ^{186}Re -BIWA 4.

3.4.5 Selection and timing of dose for each subject

3.4.5.1 Dosage and Treatment Schedule

25 The investigators were allowed to administer the trial drug at any time. The time elapsed between radiolabelling and administration, however, should not exceed 24 hours.

3.4.6 Blinding

This was an open uncontrolled trial. No blinding was done.

3.4.7 Prior and concomitant therapy or procedures

30 3.4.7.1 Rescue Medication and Additional Treatment(s)

Anaphylaxis was considered to be the most serious potential side effect and would have mandated immediate cessation of antibody infusion and the institution of appropriate resuscitative measures. Precautions to be taken were: Resuscitation equipment, within reach: anti-histaminics, corticosteroids and epinephrine. Any patient experiencing this type of adverse
35 reaction was not allowed to receive additional monoclonal antibody.

Patients were allowed to receive other treatment(s) either for the study indication or for unrelated illnesses providing the inclusion and exclusion criteria were met. All concomitant therapy had to be recorded in the CRF.

- 5 In case of serious haematologic toxicity (CTC grade 4) cytokine intervention or other methods for alleviating bone marrow toxicity were allowed.

3.4.7.2 Restrictions

10 Additional chemotherapy or radiotherapy was not allowed and the last chemotherapy or radiotherapy should have stopped more than 4 weeks before inclusion in this study. All prior chemotherapy and radiotherapy was to be recorded in the CRF.

3.4.8 Treatment compliance

The study medication was given as a single intravenous infusion. The compliance was verified with pharmacokinetic assessments and with radioimmunoscentigraphic images.

15 3.5 EFFICACY / CLINICAL PHARMACOLOGY AND SAFETY VARIABLES

3.5.1 Efficacy / pharmacodynamics and safety measurements assessed and flow chart

Flow Charts:

Part A

- 20 The investigations performed in Part A and the respective times are detailed in the flow chart given below. A description of the investigations is given in the following sections:

5 TABLE 3.5.1: 1 FLOW CHART: Part A: Biodistribution study with ^{99m}Tc-labelled hMAb BIWA 4

	Visit 1	Visit 2						Visit 3
	screening visit*	Day 1*		Day 2	Day 3	Day 4	Day 7	Follow-Up Week 6
		pre-inj.	Post-inj.					
Signed informed consent	x							
Demographics	x							
Conc. Therapy	x	x	x	x	x	x	x	X
Medical History	x							
Physical Examination	x							x
ECG	x							
Chest X-ray	x							
Blood safety analysis	x ¹	x ¹		x ¹	x ¹		x ¹	x ¹
Urine safety analysis	x						x	x
Pregnancy test	x							x
HAHA assessment	x ²						x ²	x ²
ENT examination	x							
CT or MRI	x							
In-/exclusion criteria	x							
Adverse events		x	x	x	x	x	x	X
Vital signs	x ³	x ³	x ³					x ³
body weight	x	x						x
Immunoscintigraphy:								
- Whole Body Scan			x ⁴	x ⁴				
- Planar Scan				x ⁵				
- SPECT Scan				x ⁵				
Pharmacokinetics:								
Blood samples		x ⁶	x ⁶	x ⁶	x ⁶	x ⁶	x ⁶	6
Urine collection		x ⁶	x ⁶	x ⁶	x ⁶			
Serum soluble CD44v6		x ⁷		x ⁷	x ⁷		x ⁷	6
Surgery					x			

*: The pre-infusion assessments should occur no more than 3 weeks prior to the actual infusion.

10 x¹: Blood samples were tested for: Glucose, sodium, potassium, calcium, chloride, creatinine, total protein, albumin, serum glutamic oxalacetic transaminase (sGOT), serum glutamic pyruvic transaminase (sGPT), Alkaline Phosphatase, Gamma Glutaryl Transpeptidase (GGT), bilirubin, urea, uric-acid, thyroid stimulating hormone (TSH), haemoglobin (Hb), haematocrit (Ht), mean corpuscular volume (MCV), reticulocytes, leucocytes, neutrophils, bands, lymphocytes, basophils, eosinophils, monocytes and platelets at the screening visit or on day 1 pre-infusion at 21, 48 and 144 hrs p.i. and at six weeks post infusion (p.i.).

15 x²: HAHA was assessed in a serum sample obtained at the screening visit, after one week (144 hrs) and after six weeks p.i.

x³: Vital signs were assessed at the screening visit, pre-infusion, at 10, 60 and 120 minutes post infusion and after 6 weeks.

x⁴: Whole body scan was done immediately after infusion and 21 hrs p.i.

x⁵: The SPECT scan and planar scan was done once 21 hrs p.i.

20 x⁶: Blood samples for pharmacokinetics were obtained pre-infusion, at the end of the infusion, at 5, 10, 30 minutes; 1, 2, 4, 16, 21, 48, 72 and 144 hrs post end of infusion and six weeks p.i. Blood samples were drawn for enzyme-linked

- 5 immuno-sorbent assay (ELISA) and for measurement of radioactivity (for details see section 9.5.4.1). Urine was collected from 0-4 hrs, 4-8 hrs, 8-12 hrs, and 12-24 hrs during the first 24 hrs and in a 24-hr sample for the remaining time until 48 hrs p.i.
- x⁷: Soluble CD44v6 was measured from a serum sample obtained pre-infusion, at 21, 48 and 144 hrs p.i and six weeks p.i.
- 10 6 week six

Part B

The investigations performed and the respective times are given in the flow chart below. The individual investigations are described in more detail in the following sections.

15 TABLE 3.5.1: 2 FLOW CHART Part B: Dose escalation study with ¹⁸⁶Re-labelled hMAb BIWA 4

	Visit 1	Visit 2						Visit 3
	Screening visit*	Day 1*		Day 2	Day 3	Day 4	Day 7	Follow-Up Week No.
		pre-inj.	Post-inj.					
Signed informed consent	x							
Demographics	x							
Conc. Therapy	x	x	x	x	x	x	x	2, 3, 4, 5, 6
Medical History	x							
Physical Examination	x							6
ECG	x							
Chest X-ray	x							
Blood safety analysis	x ¹	x ¹		x ¹	x ¹		x ¹	2, 3, 4, 5, 6
Urine safety analysis	x						x	6
Pregnancy test	x							6
HAHA assessment	x ²						x ²	6
ENT examination	x							6 x ⁸
CT or MRI	x							6 x ⁸
CT thorax	x ⁸							
In-/exclusion criteria	x							
Adverse events		x	x	x	x	x	x	2, 3, 4, 5, 6
Vital signs	x ³	x ³	x ³					6
Body weight	x	x						6
Immunoscintigraphy:								
- Whole Body Scan			x ⁴	x ⁴	x ⁴	x ⁴	x ⁴	
- Planar Scan				x ⁵	(x ⁵)	x ⁵	x ⁵	
- SPECT Scan						x ⁵		

Pharmacokinetics:								
Blood samples		x ⁶	x ⁶	x ⁶	x ⁶	x ⁶	x ⁶	6
Urine collection		x ⁶	x ⁶	x ⁶	x ⁶	x ⁶		
Serum soluble CD44v6		x ⁷		x ⁷	x ⁷		x ⁷	6

5 *: The pre-infusion assessment had to occur no more than 3 weeks prior to the actual infusion.

x¹: Blood samples were tested for: Glucose, sodium, potassium, calcium, chloride, creatinine, total protein, albumin, sGOT, sGPT, Alkaline Phosphatase, GGT, bilirubin, urea, uric-acid, TSH, Hb, Ht, MCV, reticulocytes, leucocytes, neutrophils, bands, lymphocytes, basophils, eosinophils, monocytes and platelets at the screening visit or on day 1 pre-infusion, at 21 hrs p.i., at 48 hrs p.i. and at 144 hrs p.i.. During weeks 2-6 p.i. safety blood samples were
10 obtained at least weekly.

x²: HAHA was assessed in a serum sample obtained at the screening visit, at 144 hrs and six weeks post infusion.

x³: Vital signs were assessed once at the screening visit, pre-infusion, at 10, 60, 120 and 240 minutes post infusion and after 6 weeks.

x⁴: Whole body scan was done immediately p.i., at 21, 48, 72, 144 hrs p.i. and optionally at two weeks p.i.

15 x⁵: The planar scan was done at 21, 48 (optional), 72, 144 hrs p.i. and optionally at two weeks p.i. The SPECT scan was done at 72 hrs p.i.

x⁶: Blood samples for pharmacokinetics were obtained pre-infusion, at the end of the infusion, at 5, 30 minutes; 1, 2, 4, 16, 21, 48, 72, 144, 240 and 336 hrs post end of infusion. Blood samples were drawn for ELISA and for measurement of radioactivity. Urine was collected from 0-4 hrs, 4-8 hrs, 8-12 hrs, and 12-24 hrs during the first 24
20 hrs and in 24-hr samples for the remaining time until 96 hrs p.i.

x⁷: Soluble CD44v6 was measured from a serum sample obtained pre-infusion, at 21, 48 and 144 hrs p.i. and six weeks p.i.

x⁸: Disease assessment was to be repeated every 6 weeks until disease progression or loss to follow-up. CT thorax was done at baseline and repeated at follow-up if there were abnormalities.

25 3.5.1.1 Radioimmunoscintigraphy and dosimetry

The data obtained from radioimmunoscintigraphy (RIS) were presented qualitatively for both Part A and B (i.e. uptake in tumour, bone marrow, liver, lung, intestine, kidney and additional organs expressed as low, medium or high) and quantitatively for Part B.

30 For qualitative assessment the rating scale was transformed into numbers (0 being no uptake, 1 being low uptake, 2 being medium uptake and 3 being high uptake). Mean values were calculated and presented.

For Part B the quantitative presentation of RIS results used dosimetric calculations as outlined in section 5.1.4 of the protocol by including the following parameters:

- Actual organ activity at a given time point expressed in MBq.
- 35 • The residence time of radioactivity in organs expressed in hours.
- The absorbed (radiation) dose expressed in mGy/MBq.
- The effective (radiation) dose expressed in mSv.

More details concerning the analysis of dosimetry can be found in the dosimetry report (dated November 21, 2001).

40 e) Radioimmunoscintigraphic Procedures

- 5 A description of the methods used for Part A and B and the time points are given in the following section:

Part A

10 With a large field of view dual headed gamma camera equipped with a low energy collimator, digital whole body images (anterior posterior (a.p.) and posterior anterior (p.a.)) were obtained directly after and 21 hrs after infusion. At 21 hrs p.i. planar and SPECT images of the head and neck were acquired. A calibration source also was acquired. Data was stored to enable quantitative analysis.

Part B

With a large field of view dual headed gamma camera equipped with a low energy collimator, digital whole body images (a.p. and p.a.) were obtained directly after, at 21, 48, 72 and 144 hrs after administration of the radioimmunoconjugate. Additional planar images of the head and neck region were made at 21, 48 (optional), 72, and 144 hrs p.i. Optionally additional imaging (whole body and planar) were performed two weeks p.i. A calibration source (i.e. an aliquot with a known fraction of the injected dose in a 10 millilitre (ml) vial), inserted in an Adams phantom, had to be placed between the lower legs of the patient during the whole body scan. The initial activity of the calibration source had to be 100-200 MBq ^{186}Re and this source had to be used during all whole body imaging studies. The energy-window and peak settings, the scanspeed, the scanlength, the scanning date, the time of starting the scan and the scan duration had to be reported. The anterior-posterior thickness of the neck and of the abdomen had to be measured, while the patient was in supine position on the scanning table.

At each imaging session, anterior and planar static images had to be taken and just before or just after this image a static image had to be acquired from the calibration source in the Adams-phantom.

At 72 hr. p.i. a SPECT study of the neck had to be performed. The methods used for reconstruction and the filter functions with cut off frequencies had to be reported.

Single Photon Emission Computed Tomography (SPECT)

SPECT images were obtained using a double headed rotating gamma camera equipped with a low energy collimator. Acquisition required at least thirty minutes. Twenty percent symmetric windows were centred at the 137 keV photon peaks.

Planar and SPECT data acquisition parameters

Planar imaging included the following minimal requirements: matrix 128x128 (detail) or 256 x 256 (whole body) and a minimum of 400000 counts with a maximum acquisition time of 10 minutes for detail and 60 minutes for whole body.

SPECT imaging included the following minimal requirements: 64 images, matrix size 64 x 64, 360 degree circular orbit, 60 second acquisitions per angle.

Analysis of the data

At the 21 hr p.i. anterior whole body images, rectangular regions of interest (ROI's) had to be drawn around the whole body and the calibration source. Also irregular ROI's around the organs which accumulated ^{186}Re (e.g. liver, spleen and left kidney) and around the tumour had to be drawn. One or more representative background regions had to be drawn. These regions had to be mirrored to the posterior images. Originally it was planned to draw a ROI around the sacrum on the posterior image. This plan was not pursued during the trial. All regions had to be saved on the computer, in order to project these ROI's on images at other time points. The

- 5 number of pixels and the counts per pixel in each region had to be reported. The number of counts in the regions for all imaging time points had to be recorded digitally in a spreadsheet.

Calculation of the absorbed dose in the organs

- 10 The amount of activity in the organs, tumour and the total body was estimated from the geometric mean counts in the ROI's of the anterior and posterior views. Background and attenuation correction were applied when indicated. The activity in the urine was not used to estimate the absorbed dose in the bladder as originally planned. Instead the dynamic bladder model was used. The residence times in the organs and the rest of the body were calculated and imported in the MIRDOSE3 program.

Required quality control tests

15 Planar Imaging

Routine quality controls were performed at the department of nuclear medicine weekly:

- 1) 100M flood table.
- 2) Extrinsic ^{57}Co flood with the low energy collimator was obtained each week.

SPECT imaging

- 20 Quality control of the SPECT imaging system had to include center of rotation determination.

Tomographic processing

A filtered back projection algorithm was used for tomographic image reconstruction using a ramp filter.

Data storage

- 25 All planar images and tomographic data had to be stored permanently on magnetic tape or optical disc. For tomographic studies, original projecting images and reconstructed studies had to be written to a back up tape.

Copies of images and reports

- 30 Copies of all relevant images, pathology and surgery reports were required for all patients. Also copies of MRI and CT reports were required.

3.5.1.2 Biopsy biodistribution of $^{99\text{m}}\text{Tc}$ -labelled hMAb BIWA 4 (Part A)

- 35 Patients entered in Part A of the study underwent surgery 48 hrs after infusion of the radiolabelled hMAb BIWA 4. Biopsies from tumour site(s) and from normal tissues (as many as possible) in the surgical specimen were taken. Then also under general anaesthesia a bone biopsy and a bone marrow aspirate was taken. The bone marrow aspirates were centrifuged to assess the radioactivity in the supernatant (plasma) and in the sediment (cellular fraction).

- ^s All biopsies were weighed and the amount of ^{99m}Tc was measured. Specific uptake of radioactivity into tumour was evaluated by comparing %ID/kg tumour with %ID/kg normal tissue.

- 5 CD44v6 antigen expression was assessed by immunohistochemistry using cryostate sections, which were first incubated with mMAb BIWA 1, followed by anti-mouse Immunoglobulin G (IgG) secondary reagent. The surgical specimen and the biopsies were investigated histo/cytopathologically.

Evaluation of surgical specimen and biodistribution

10 After receiving the surgical specimen it was processed in the following chronological order.

1. Pictures were taken of the specimen. A Polaroid from the front and slides from the front and back.
2. The size of the surgical specimen was assessed.
- 15 3. Biopsies of primary tumour, suspect lymph node, and if possible from normal tissues in the surgical specimen like normal mucosa, normal lymph node, fat and muscle were taken. All biopsies were weighed and the amount of ^{99m}Tc was measured. All data were converted to percentages injected dose/kilogram tissue. Specific uptake of radioactivity into tumour was evaluated by comparing %ID/kg tumour with %ID/kg normal tissue.
- 20 4. Next the specimen was nailed to a board and fixed in formaldehyde 4% for at least 36 hours.
5. After dissecting the sternocleidomastoid muscle (a structure with a high radiodensity), a specimen radiograph was made to show the exact size and location of the lymph nodes involved. This radiograph was made while the specimen was being immersed in ethanol 96%, which has the same X-ray absorption as fat.
- 25 6. All the nodes visualised with the X-ray were indicated on the Polaroid and specimen radiograph.
7. All the nodes found by examining the surgical specimen and by X-ray were dissected from the specimen.
- 30 8. All macroscopically negative nodes were entirely processed for microscopy and one single section was evaluated. Of all macroscopically positive nodes two or more slices were made. Macroscopical evidence for the presence or absence of tumour necrosis were recorded.
- 35 9. Furthermore, the number of nodes enclosed and the number, localisation and lymph node level (according to the Memorial Sloan Kettering Cancer Center Classification) of tumour containing nodes were recorded.

5 3.5.1.3 Tumour Response (Part B)

The efficacy parameter for the radioimmunotherapy treatment was tumour response. Tumour response was assessed with tumour measurements as assessed clinically and/or with CT, MRI or bone scintigraphy investigations. Evaluation was done according to response criteria of the World Health Organisation (WHO). See Appendix VI of the protocol, (R96-0941).

10 3.5.1.4 Physical Examination

Before inclusion in the study the disease status of each patient was evaluated by ENT-examination (palpation included) and by CT or MRI of the tumour site(s).

All head and neck lesions were described per neck side.

Palpation

15 All patients were examined at baseline by an otolaryngologist/head and neck surgeon. This clinical investigator assessed the number, size, location and mobility of all palpable lymph nodes in the head and neck area. The character of the lymph nodes was described as: not suspected, suspected or tumour infiltrated. The status of the neck lymph nodes was classified according to the Tumour Node Metastasis system for staging tumours (TNM) of the Union
20 Internationale Contre le Cancer (UICC) at diagnosis.

Radiological examination

Depending on the localisation of the tumour site(s), one or more of the following examinations were required: CT, MRI, bone scintigraphy. For tumour involvement of the head and neck region MRI was generally preferred. In case there were bone lesions, CT was preferred. For
25 Part B all radiological disease assessment parameters obtained at baseline were repeated six weeks after the infusion and every six weeks until progression or lost to follow-up. For patients entered in Part B of the study a CT thorax was obtained at baseline, and was repeated at follow-up if there were tumour lesions in the thorax. Ultrasound might have been employed as additional technique of tumour imaging. Guidelines for the investigations are given below:

30 **Primary tumour / loco-regional recurrence**

For patients participating in Part A of the study and patients with a loco-regional recurrence (Part B) CT scan and/or MRI of the head and neck region had to be performed.

Computed Tomography

35 Computed tomography of the Head and Neck region: Dynamic CT was preferred over spiral CT. However, spiral CT might have been helpful in patients who were unable to cooperate.

The patient had to be examined in supine position, the neck slightly hyperextended, the head immobilised and the shoulders relaxed and pushed downwards. The patient had to breath quietly with use of abdominal rather than chest muscles. The area to be scanned was

5 determined from the initial overview made in the lateral projection. The plane of the scan had to be parallel to that of the vocal cords. Contiguous three millimetre-slices had to be used routinely. Imaging had to be performed from skull base to upper mediastinum.

Magnetic resonance imaging

10 The patients had to be examined in supine position, the neck slightly hyperextended and the head immobilised. Quiet breathing was mandatory during the examination with the use of abdominal rather than the chest muscles. The appropriate images to demonstrate the neck anatomy and to assess the extent of the primary lesion and the presence of lymph node spread was obtained. They had to be obtained from the skull base to the upper mediastinum.

Distant metastases

15 For all patients participating in Part B of the study CT scanning of the thorax was done at baseline. Additional investigations to visualise metastases, like bone scintigraphy or CT abdomen, were optional.

CT thorax

20 Spiral CT-scans were obtained. Patients were examined in supine position and had to raise their arms above their heads. Patients had to breath quietly with use of abdominal rather than chest muscles. The patient was scanned from just above the lungs to the level of the adrenal glands. Images had to be photographed in mediastinal and lung setting.

CT abdomen

25 Spiral scans were obtained. Patients were examined in supine position and had to raise their arms above their heads. Patients had to breath quietly. Oral contrast was used in all patients. The area to be scanned ran from just above the diaphragm to the symphysis. Images had to be photographed in abdominal and liver setting.

Bone scintigraphy

30 After intravenous infusion of maximally 600 MBq of ^{99m}Tc -labelled HDP/MDP, the patient was asked to consume enough fluids and to void frequently. Static views were obtained 3 hours after infusion: one whole body view and if necessary 3 more detailed views. Suspect lesions might have needed closer analysis by CT and/or MRI.

Reporting of response of bone metastases were planned to be conducted according to a separate set of response criteria. No data was provided.

35 3.5.1.5 Soluble CD44v6

Soluble CD44v6 (sCD44v6) had to be measured in serum. Concentrations were determined by means of a validated enzyme-linked immuno-sorbent assay (ELISA) that was based on a commercially available test kit and conducted in accordance with current international

5 guidelines at the Boehringer Ingelheim Department of Pharmacokinetics and Drug Metabolism, Biberach, Germany. Blood samples (to be processed to serum) of 5 mL were obtained pre-infusion, at 21, 48 and 144 hrs p.i. and six weeks after the infusion. Samples were allowed to clot and centrifuged to prepare serum. Storage and shipment conditions: The sample for ELISA measurement was put in cryotubes, labelled carefully to enable unique identification,
10 stored at -20°C until radioactivity has decreased (^{186}Re : 4 weeks, $^{99\text{m}}\text{Tc}$: 3 days) and sent to BI Department of Pharmacokinetics and Drug Metabolism, Biberach, Germany in batches every four weeks. The serum samples were sent on dry ice.

3.5.1.6 Safety

Protection of subjects

15 All patients were monitored carefully during and after administration of the radiolabelled monoclonal antibody. For this purpose at least one of the investigators was present.

Laboratory Assessments

Blood samples were collected for glucose, sodium, potassium, calcium, chloride, creatinine, total protein, albumin, serum glutamic oxalacetic transaminase (sGOT), serum glutamic
20 pyruvic transaminase (sGPT), alkaline phosphatase (AP), gamma glutaryl transpeptidase (GGT), bilirubin, urea, uric-acid, thyroid stimulating hormone (TSH), haemoglobin (Hb), haematocrit (Ht), median corpuscular volume (MCV), reticulocytes, leucocytes, neutrophils, bands, lymphocytes, basophils, eosinophils, monocytes and platelets at the following time points:

- 25
- At the screening visit or on day one pre-infusion and
 - At 21, 48 and 144 hrs post infusion.
 - For Part A at 6 weeks post infusion. For Part B: During weeks 2, 3, 4, 5 and 6 of the study at least once weekly (more often in case of toxicity).

30 Baseline laboratory assessments pre-infusion could have been obtained either at the screening visit or on day one of the study, provided the samples were obtained less than 21 days before the infusion of radiolabelled hMAb BIWA 4 and all required assessments were done. All required laboratory test results had to be reviewed and checked for eligibility by the responsible physician prior to the antibody infusion. The same also applied to a second hMAb BIWA 4 administration in Part B of the study.

35 Urine samples were collected for standard hospital screening (protein, blood, and glucose) at the time points:

- Screening visit
- One week post infusion (144 hrs p.i.)

- 5 • Six weeks post infusion

A pregnancy test was required for women with childbearing potential at the screening visit and at the study end.

Human-Anti-Human-Antibody Assessment

10 The presence and/or the development of HAHA was evaluated in serum samples. Therefore blood samples (to be processed to serum) of 5 mL were taken at the screening visit, at one week (144 hrs p.i.) and at six weeks after infusion of the antibody. For patients receiving two BIWA 4 administrations in Part B of the study an additional HAHA sample was collected before the second administration. Serum level was evaluated by means of validated ELISA methods.

15 Samples were allowed to clot and centrifuged to prepare serum.

Storage and shipment conditions: The samples for ELISA measurement were to be put in cryotubes, labelled carefully to enable unique identification, stored at -20°C until radioactivity had decreased (¹⁸⁶Re: 4 weeks, ^{99m}Tc: 3 days) and sent to BI Department of Pharmacokinetics and Drug Metabolism, Biberach, Germany in batches every four weeks. The serum samples
20 had to be sent on dry ice.

Any elevations in human anti-human antibody levels were compared casewise carefully with the patient's possible adverse events. This data was collected and analysed retrospectively during the course of the study.

Adverse Events

25 All adverse events were recorded in the CRF. The events were graded according to the US National Cancer Institute CTC (NCI-CTC) version 2.0 (a version of this document had to be downloaded from the Internet at the site: <http://ctep.info.nih.gov/CTC3/ctc.htm>). All SAEs had to be reported.

Immunogenicity and toxicity

30 Before performance of this trial no human data existed with respect to safety and immunogenicity of hMAb BIWA 4. In the previous study performed with the parental murine antibody BIWA 1 no clinically significant toxicity was encountered. No toxicity was expected from the naked hMAb BIWA 4 antibody alone (no antibody dependent cell-mediated cytotoxicity (ADCC) effector functions in vitro; no interference with proliferation of antigen-
35 expressing cells expected). It was demonstrated that hMAb BIWA 4 could be safely administered to mice (xenograft studies) and monkeys (toxicity studies). Good local tolerability was demonstrated in three local toxicity studies.

Murine and chimeric MAb U36 (anti-CD44v6-epitope with defined overlapping epitope specificity) was so far safe in HNSCC patients; no toxicities except for myelotoxicity (caused

5 by ^{186}Re -labels) were seen in a Phase I dose-escalation RIT study with cMAb U36. It was theoretically possible that hypersensitivity reactions to radiolabelled hMAb BIWA 4 might occur. Monoclonal antibodies have been administered to several thousands of patients for diagnostic applications.

10 Although unlikely, potential reactions to intravenously administered BIWA 4 might include hypotension, transient fever and chills, skin rashes, dyspnoea, itching, nausea, and anaphylaxis.

Vital signs (blood pressure, temperature, pulse rate and respiratory rate) were therefore recorded at the screening visit and at the following time points: pre-infusion and 10, 60, 120 minutes post-infusion and after 6 weeks. Additionally vital signs were recorded at 240 minutes post-infusion for patients participating in Part B only.

15 In this study radionuclides were used. The radiation burden associated with the gamma emitting radionuclide $^{99\text{m}}\text{Tc}$ (in the current study 20 mCi) was similar to that encountered in many routine nuclear medicine procedures, and was known to be small. To minimise radiation exposure to the bladder and kidneys the patients were well hydrated and asked to void at frequent intervals during the first 48 hours (96 hours after infusion with ^{186}Re -labelled
20 antibody).

For the beta- and gamma-emitting radionuclide ^{186}Re , Breitz *et al.* (R98-2459) found a maximum tolerated dose of ^{186}Re -labelled MAb IgG in heavily pretreated patients of 90 mCi/m². One of the aims of this study was to investigate the toxicity of ^{186}Re -labelled hMAb BIWA 4 in patients who either had local and/or regional recurrent disease for which curative
25 treatment options were not available, or distant metastases. Because bone marrow, thyroid, kidney and liver toxicity might occur, blood samples for organ function testing were taken as described in section 3.5.1.6.

To monitor the occurrence of weight loss or weight gain the body weight was recorded at the screening visit, before the infusion on day 1 and after 6 weeks.

30 Patients were monitored for the occurrence of local toxicity. Any sign of local toxicity was recorded as adverse event.

3.5.2 Appropriateness of measurements

The measurements used are well-accepted for this type of trial.

3.5.3 Primary efficacy / pharmacodynamic variables(s)

35 3.5.3.1 Part A of the trial

The primary efficacy variables determined in Part A of the trial were the biopsy distribution and the radioimmunosintigraphy. Description of the methods how to obtain the data is given in section 3.5.1.1 and 3.5.1.2.

- 5 The uptake in the normal tissue and tumour was measured in biopsies of the surgical specimen and the uptake expressed as percent of the injected dose per kg (% ID/kg). The time course of uptake in the tumour and other tissues was evaluated and compared concerning the different doses of BIWA 4.

3.5.3.2 Part B of the trial

- 10 The primary parameters for efficacy were the analysis of radioimmunoscinigraphy and dosimetry.

3.5.4 Drug concentration measurements / pharmacokinetics

Blood concentrations of radiolabelled BIWA 4 were determined in both Part A and Part B of the trial.

15 3.5.4.1 Methods and timing of sample collection

- Patient blood samples were collected and handled as follows: 7 mL of blood (3.5 mL in potassium EDTA containing tubes and 3.5 mL in coagulation tubes: amount of millilitres requested are given as per amendment 1) were sampled from a peripheral vein of the arm opposite to the infusion site at the designated time points (i.e. just prior to antibody
20 administration, at the end of the infusion and at 5, 10, 30 minutes; 1, 2, 4, 16, 21, 48, 72 hrs post-end of infusion and on day 7 p.i. (144 hrs) and 6 weeks after the infusion). End of infusion was regarded as $t = 0$.

The planned samples at 10 minutes and 6 week p.i. were omitted for Part B of the trial. Instead samples were collected at 240 and 366 hour p.i. as per amendment 1.

- 25 Urine was collected during 48 hrs after the ^{99m}Tc -labelled hMAb BIWA 4 infusion and during 96 hrs after the ^{186}Re -labelled hMAb BIWA 4 infusion. Urine was collected from 0-4 hrs, 4-8 hrs, 8-12 hrs, and 12-24 hrs during the first 24 hrs and in 24-hr samples for the remaining time. Radioactivity of urine samples was counted to determine the excretion of radioactivity.

- Blood samples were centrifuged to prepare serum. Before processing, the radioactivity in
30 whole blood was measured. Radioactivity was also measured in serum. Refrigerated storage of blood samples was obligatory, though samples had not to be frozen down. Using the aliquot retained from the conjugate preparation, a weighted dilution of the injected patient dose was prepared as standard. The counting of the patient samples and standards were performed according to local SOPs. Results were recorded in the appropriate sections of the CRF. The
35 radionuclide content was reported as a percentage of the injected doses (expressed as % ID/kg blood or serum). Apart from radioactivity counting all samples were analysed for presence of immune complexes by means of HPLC analysis.

- All sample tubes were labelled with the following information: Trial number, patient number, sample identification (i.e., serum, plasma, or blood), time relative to infusion, actual time,
40 actual date and isotope (i.e., ^{99m}Tc or ^{186}Re). The volume of each urine sample was measured

5 and recorded and all urine samples were labelled with the following information: Trial number, patient number, total sample volume, collection interval, actual time, date and isotope (i.e., ^{99m}Tc or ^{186}Re).

Dates, times and radioactivity measurement results of all pharmacokinetic samples were recorded in the CRF.

10 The plasma samples for ELISA measurement were transferred in cryotubes, labelled carefully to enable unique identification, stored at -20°C until radioactivity had decreased (^{186}Re : 4 weeks, ^{99m}Tc : 3 days) and sent to Boehringer Ingelheim Department of Pharmacokinetics and Drug Metabolism, Biberach, Germany in batches every four weeks. The plasma samples were sent on dry ice.

15 3.5.4.2 Analytical determinations

Plasma samples were measured by validated ELISA methods at the Boehringer Ingelheim Department of Pharmacokinetics and Drug Metabolism, Biberach, Germany. Radioactivity counting of the samples were performed in full blood, serum and urine by the investigator.

3.5.4.3 Parameters and evaluation

20 The following pharmacokinetic parameters using WinNonlin 3.1 Professional (Pharsight Corporation, Mountain View, CA) were determined from full blood, plasma, serum levels and imaging data as described below:

The primary pharmacokinetic parameters are the time point at which the maximum drug concentration is observed (T_{max}), maximum drug concentration observed (C_{max}), area under the concentration-time curve ($\text{AUC}_{0 \rightarrow \infty}$), terminal elimination half-life ($t_{1/2}$), volume of distribution (terminal phase and predicted steady state), total body clearance (CL), and mean residence time ($\text{MRT}_{0 \rightarrow \infty}$).

The primary objectives of pharmacokinetic analysis were:

30 To determine and compare the disposition of total radioactivity and immunoreactive hMAb BIWA 4 following administration of single doses of ^{99m}Tc -labelled (Part A) and ^{186}Re -labelled hMAb (Part B) BIWA 4.

To calculate, from imaging data and blood disposition data, the fraction of the injected radioactivity reaching the liver and spleen during the studies. To achieve this objective, the WinNonlin® software package was used to perform non-compartmental pharmacokinetic analysis on the plasma concentration vs. time profiles of immunoreactive hMAb BIWA 4 and of total blood radioactivity levels, as a function of time, following intravenous administration.

The results are reported as summary statistics.

5 3.5.5 Pharmacokinetic / pharmacodynamic relationship

A compartmental pharmacokinetic model was initially intended to describe the distribution and metabolism of ^{186}Re -labelled BIWA 4 in humans. It was intended to combine data from BIWA 4 plasma levels, full blood and serum radioactivity measurements, radioactivity from whole body images and specific regions of interest, the administered dose ^{186}Re -labelled BIWA 4, the dose of unlabelled BIWA 4, soluble CD44v6 levels and the assessment of the radiolabelled antibody prior to infusion.

The results were to be reported as summary statistics. However, variability was too high to be used for a reasonable model.

3.5.6 Primary Safety variable

15 Determination of maximum tolerated dose and assessment of safety were the primary endpoints in Part B of the trial. The measurements done are described in section 3.5.1.6.

3.5.6.1 Dose Limiting Toxicity and Maximum Tolerated Dose

DLT was defined as drug-related CTC grade 3 non-haematologic toxicity or drug-related CTC grade 4 haematologic toxicity, excluding nausea and vomiting without adequate antiemetic treatment.

The MTD was defined as the dose level at which less than two out of six patients developed drug-related DLT.

3.6 DATA QUALITY ASSURANCE

25 The trial was in general conducted according to the principles of Good Clinical Practice as specified in the appropriate regulations and in the company standard operating procedures reflecting these regulations.

Throughout the course of the study, a representative from Boehringer Ingelheim The Netherlands was contacting and/or visiting the study site to monitor the progress of the study. There were frequent contacts with the investigator and onsite visits for the purpose of data audits, including the comparison of source documents with Case Report Forms and drug accountability checks. The investigator or his/her designee had to be available to the Boehringer Ingelheim The Netherlands representative during these onsite visits.

Duly filled in CRFs were collected on a regular basis. Data were double-data entered in-house. Review of the data was done and implausibilities questioned to the investigator.

35 Serious adverse events were reported according to the guidelines defined in the protocol and CRF and according to Boehringer Ingelheim SOPs.

5 **3.7 STATISTICAL METHODS PLANNED IN THE PROTOCOL AND
 DETERMINATION OF SAMPLE SIZE**

3.7.1 Statistical and analytical plans

 3.7.1.1 Statistical design / model

The designs used in this trial are commonly used in Phase I oncology trials.

10 Part A

Part A of this phase I trial was an uncontrolled, rising dose sequential group study to determine the safety, tolerability, biodistribution and pharmacokinetics of a single infusion of ^{99m}Tc-labelled hMAb BIWA 4 in patients with advanced squamous cell carcinoma of the head and neck.

15 Three hMAb BIWA 4 dose levels were used in this part of the study with three patients planned at each dose level. All patients had a proven tumour of the head and neck and were destined for surgery.

Part B

20 Part B of this phase I trial was an open uncontrolled dose escalation study to determine the safety, tolerability, MTD, pharmacokinetics and preliminary therapeutic effects of a single infusion of ¹⁸⁶Re-labelled hMAb BIWA 4 in patients with advanced squamous cell carcinoma of the head and neck for whom no curative treatment options were available.

25 In this part of the study the radiation dose was escalated. Two patients were treated at dose levels where < grade 2 CTC toxicity was seen. If ≥ grade 2 toxicity was seen at least three patients were treated per dose group.

Safety and tolerability evaluation (Parts A + B)

Safety and tolerability of the study were assessed in terms of changes in laboratory parameters, vital signs, development of HAHA and the incidence of adverse experiences. The results were reported for each dose level separately as well as in terms of overall means.

30 Biodistribution evaluation (Part A)

The biodistribution of ^{99m}Tc-labelled hMAb BIWA 4 in solid tissue was assessed by radioimmunosintigraphy (see sections 3.5.1.1) and by measurements of radioactivity of biopsy specimens (see sections 3.5.1.1 and 3.5.1.2). Because only a maximum of 3 patients were entered per dose level only descriptive statistics were possible.

35 Immunoscintigraphic imaging evaluation (Parts A+B)

- ^s At given time-points after the administration of radio-labelled antibody immunoscintigraphic images were obtained (see section 9.5.1.1). Again only descriptive statistics were applicable.

5 Pharmacokinetic evaluation (Parts A + B)

The following pharmacokinetic parameters were determined from plasma or serum, urine levels and imaging data as described below:

Primary parameters (non-compartmental)

10 T_{max} , C_{max} , AUC (0-infinite time), terminal elimination half-life, volume of distribution (terminal phase V_z and predicted steady state V_{ss}), CL, and MRT (0-infinite time). Cumulative urinary excretion of radioactivity over time was determined from total urine output.

Secondary parameters (compartmental)

15 A compartmental pharmacokinetic model was planned to be developed to define the distribution and metabolism of ^{186}Re -labelled hMAb BIWA 4 in humans. The model collated data from serum and urine radioactivity measurements, radioactivity from whole body images and specific regions of interest, the administered dose ^{186}Re -labelled hMAb BIWA 4, the cold-loading dose of hMAb BIWA 4, soluble CD44v6 levels and the assessment of the radiolabelled antibody prior to infusion.

20 The results were planned to be reported for each dose level separately and by means of summary statistics.

Therapeutic efficacy evaluation (Part B)

25 Tumour response was the parameter for therapeutic efficacy. Tumour response was assessed according to the WHO guidelines. The sum of the products of the largest perpendicular diameters of all measured tumour lesions were the primary parameter for tumour response. There were separate criteria for the assessment of response of bone lesions.

3.7.1.2 Null and alternative hypotheses

30 All analyses in this trial were descriptive and exploratory by nature. Any statistical tests were performed only to provide a statistical framework from which to view the results and providing aid for planning further studies. No formal statistical inferences were foreseen and, accordingly, no statistical tests were performed.

3.7.1.3 Planned analyses

Two populations were distinguished:

- The intention to treat subset consists of all patients who received an infusion of radiolabelled hMAb BIWA 4 and for whom data after baseline were available.
- 35 • The per protocol subset consists of evaluable patients. A patient was evaluable if the following criteria were met:

5 Part A:

1. Fulfilling the criteria for the intention to treat subset.
2. The follow-up in terms of adverse events, vital signs and safety blood and urine samples was adequate.
3. The immunoreactive fraction in at least one of both performed assays was larger than 60%.
- 10 4. Adequate biopsies and scintigraphic images were available to assess the biodistribution.

Part B:

1. Fulfilling the criteria for the intention to treat subset.
2. The follow-up in terms of adverse events, vital signs and safety blood and urine samples was adequate.
- 15 3. The immunoreactive fraction in at least one of both performed assays was larger than 60%.
4. Response: Patients (Pts.) were evaluable for response if adequate tumour measurements were obtained at baseline and if patients were followed over at least a six week period. If disease progression was observed within this six weeks period, patients were also evaluable for response.

20 Primary Analyses

The primary analysis was performed for the per protocol subset.

Non evaluable patients were replaced. For Part A therefore the evaluable subset was planned to consist of 9 patients and the intent to treat subset consisted of at least 9 patients. For Part B the number of patients depended on the toxicity encountered. Patients who were evaluable for
25 toxicity but not for response were not replaced.

Pharmacokinetic data:

The analysis of pharmacokinetic data was performed by Dr. Thomas R. MacGregor, Boehringer Ingelheim Pharmaceuticals Inc. Ridgefield CT, USA.

The primary objectives of pharmacokinetic analysis were:

- 30 1. To determine and compare the disposition of total radioactivity and immunoreactive hMAb BIWA 4 following administration of single doses of ^{99m}Tc -labelled hMAb BIWA 4 and ^{186}Re -labelled hMAb BIWA 4.
2. To identify an appropriate cold-loading dose of hMAb BIWA 4.

- 5 3. To assess the disposition of a second dose of ^{186}Re -labelled hMAb BIWA 4.
4. To calculate, from imaging data and blood disposition data, the fraction of the injected radioactivity reaching the liver and spleen during the studies. To achieve this objective, the WinNonLin® software package was used to perform noncompartmental pharmacokinetic analysis on the plasma concentration vs. time profiles of immunoreactive hMAb BIWA 4 and of total blood radioactivity levels, as a function of time, following intravenous
10 administration. Any measurements involving radioactivity were normalised to the fraction of the radiolabel that was protein-bound and immunoreactive prior to infusion. The specific pharmacokinetic parameters that were generated include: $T_{\max}$, $C_{\max}$, AUC (0- infinite time), terminal elimination half-life, volume of distribution (terminal phase and predicted
15 steady state), total body clearance, and mean residence time (0- infinite time). Cumulative urinary excretion of radioactivity over time were determined from total urine output.

The secondary objective of the pharmacokinetic analysis to develop a pharmacokinetic model to describe the distribution and metabolism of ^{186}Re -labelled hMAb BIWA 4 in humans was not achieved due to the high variability of the data (especially the radioscintigraphic data).

20 Secondary Analyses

The secondary analyses were performed for the intention to treat subset of patients participating in Part B of the study.

The secondary analyses were restricted to key endpoints: e.g.: Response rate, duration of response, time to progression.

25 Interim Analysis

No interim analyses were made.

3.7.1.4 Handling of missing data

In this phase I study it was anticipated that most data were available for analysis. In case of missing data it was most likely that the reason for missing was not outcome related. No
30 missing data were imputed.

3.7.2 Determination of sample size

In Part A three patients per dose level provide some information on safety of the hMAb BIWA 4 and on the anticipated correct dose of 50 mg unlabelled hMAb BIWA 4.

35 In Part B six patients were regarded sufficient to determine DLT. TABLE 3.7.2:1 exhibits the probabilities of 2 or more out of 6 patients to be observed with DLT for some assumed underlying rates of DLT in the population of all patients.

5 TABLE 3.7.2: 1 Assumed Population Rates of DLT

	0.40	0.42	0.45
Probability of observing DLT in two or more out of six patients	0.77	0.80	0.84

Source: Probabilities calculated from the cumulative distribution function of the binomial distribution with n=6 and varying p's.

10 With the escalation scheme in this trial, there was a probability of at least 80% for two or more patients to exhibit DLT, if the underlying individual probability for a patient to reach DLT is 42 % or larger.

3.8 CHANGES IN THE CONDUCT OF THE STUDY OR PLANNED ANALYSES

The following changes from protocol were implemented via an amendment.

15 Time points for collection of blood samples were adjusted by implementation of amendment No 1 dated 1 September 1999. The samples at 10 minutes and 6 weeks after infusion were dropped while additional samples were collected after 240 and 336 hours post end of infusion in Part B of the trial. It had turned out that the original blood sampling schedule was not sufficient to adequately cover an assumed half-life of 50 hours which was observed in Part A of the trial. A total of 3.5 mL blood was collected in potassium EDTA tubes and 3.5 mL in coagulation tubes.

20 The protocol stated that patients could be administered a second dose of ^{186}Re -BIWA 4 after having determined the MTD. This procedure was modified by amendment 2. Patients were considered eligible for a second treatment cycle in case they responded to the first administration of ^{186}Re -BIWA 4, were free of or had recovered from adverse events and had no HAHA antibodies. Response was defined as stable disease (i.e. no change), partial
25 remission or complete remission. The second dose administered was always 50 mCi/m². Due to the change in the linker the trial was concluded before having investigated a second dose with the dosing regimen described in the protocol. To allow patients who responded to be treated anyhow, amendment 2 was issued.

30 The trial was completed and terminated after having reached the MTD due to the change of the linker. MAG3 will be replaced by MAG2GABA-TFP in the future development of the BIWA 4 project.

Samples were available for sCD44v6 determination for more time points than originally planned.

35 The following changes were made after having evaluated the data and after discussion within the trial team.

- 5 The ratio of %ID/kg tumour versus %ID/kg non-tumour tissue was performed for bone marrow only, due to the variability of the results.

The same holds true for the evaluation of the tumour size. Measurements were incomplete thus jeopardising a reasonable evaluation. Therefore, no formal analysis was done.

4. STUDY SUBJECTS

4.1 DISPOSITION OF SUBJECTS

Of the 33 patients screened three patients discontinued the trial due to adverse events before administration of the trial drug (TABLE 6.1: 1). These three patients are not included in the analyses. A total of 30 patients were entered and treated in this trial.

Part A

Ten patients of the 30 patients treated were included in Part A with 3 each receiving 25 mg and 100 mg ^{99m}Tc -BIWA 4, respectively, while four patients received 50 mg ^{99m}Tc -BIWA 4. All patients in Part A completed the trial.

Part B

Two out of the 20 patients treated in Part B discontinued due to adverse events (two patients died). Three patients were administered a second dose of 50 mCi/m².

TABLE 6.1: 1 Disposition of Patients

	Part A				Part B					Total
	^{99m} Tc-BIWA 4			20	¹⁸⁶ Re-BIWA 4 [mCi/m ²]					
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4		30	40	50	60		
Screened										33
Entered	3	4	3	2	4	3	6	5		30
Treated	3	4	3	2	4	3	6	5		30
Completed	3	4	3	2	3	2	6	5		28
Discontinued	0	0	0	0	1	1	0	0		2
due to AE	0	0	0	0	1	1	0	0		2
due to lack of efficacy	0	0	0	0	0	0	0	0		0
due to other reason	0	0	0	0	0	0	0	0		0

Source

Appendix 16.1.9.2 TABLE 1.1

4.2 PROTOCOL DEVIATIONS

The majority of protocol violations were deviations ($> \pm 10\%$) from the time windows provided in the protocol concerning blood and/or urine sampling. The patients affected were not excluded from the analyses but the actual times of collecting blood or urine were used for determination of blood, plasma, serum and urine concentrations.

5. EFFICACY / CLINICAL PHARMACOLOGY EVALUATION

Results from Part A confirmed a dose of 50 mg BIWA 4 as the optimal dose for treatment. Data from Part B indicate that patients may clinically benefit from ^{186}Re -BIWA 4 therapy at a radiation dose below that dose at which DLT was observed.

Concentrations of BIWA 4 measured were dose-proportional and a moderate amount of the dose administered was excreted via the kidneys.

5.1 DATA SETS ANALYSED

All patients who had received at least one dose of the trial drug were included in the intent-to-treat analysis. Seven patients from Part A were included in the per protocol analysis. No per protocol analysis was performed for Part B.

5.2 DEMOGRAPHICS AND OTHER BASELINE CHARACTERISTICS

The mean age of all patients included was 56 years (range 37 to 78). Nineteen patients were males and eleven were females (TABLE 5.2: 1).

Three patients treated in Part A never smoked while seven were current smokers (pack years range 19 - 47). Frequent alcohol consumption was reported in eight patients while two were non drinkers (TABLE 5.2: 1).

All patients in Part B were either ex-smokers or current smokers (TABLE 5.2: 1) with a mean pack-year of 35 (range 10 to 86). Six patients were non-drinkers while the remaining patients were frequent consumers of alcohol (TABLE 5.2: 1).

Stage of the disease was local operable for all patients included in Part A while patients from Part B had recurrent disease (15 patients) and/or metastases (3 patients). One patient had local inoperable disease. In one patient the information is missing.

Concomitant diseases or relevant medical history were reported in 28 patients (TABLE 5.2: 1). Concomitant therapy was required by all patients. The most often used medications were analgesics, sedatives, lactulose, H_2 -blockers, anti-emetics and antimicrobial agents.

5 TABLE 5.2: 1 Demographics and baseline characteristics. Given are means for age. All other figures denote numbers of patients

	Part A			Part B					Total
	^{99m} Tc-BIWA 4			¹⁸⁶ Re-BIWA 4 [mCi/m ²]					
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4	20	30	40	50	60	
n	3	4	3	2	4	3	6	5	30
age [mean]	45.7	58.5	58.7	57.0	61.0	53.0	57.0	60.4	56.4
male/female	2/1	2/2	1/2	1/1	3/1	2/1	5/1	3/2	19/11
conc.disease	3	3	2	2	4	3	6	5	28
concom. therapy	3	4	3	2	4	3	6	5	30
smoker	2	4	1	0	1	0	3	1	12
ex-smoker	0	0	0	2	3	3	3	4	15
Alcohol yes	3	3	2	0	3	3	6	2	22

10 Initial diagnosis of disease was known in patients from Part A about one month before inclusion while patients included in Part B were ill since two years on average (range 0.1 to 17.5 years; TABLE 5.2: 2). The Karnofsky score ranged from 70 to 100 in patients from Part A. Patients from Part B had Karnofsky scores of 70 to 90 (TABLE 5.2: 2).

Metastases at diagnosis were reported for one patient in Part B of the trial.

15 All patients in Part B had undergone previous radio- and / or chemotherapy while patients in Part A had neither previous chemo- nor radiotherapy reported (TABLE 5.2: 2). Cisplatin, methotrexate and fluorouracil were the most often systemic anticancer drugs used).

Previous surgery was radical in eleven of the 13 patients in Part B who had a previous surgery. Only one patient from Part A, had undergone previous surgery (TABLE 5.2: 2).

20 Lesion sizes (including metastases) of the tumours ranged from 14 mm² to 10304 mm². The primary tumour site was moderately differentiated in the majority of the patients. Lymph nodes were affected in 13 patients (TABLE 5.2: 2).

5 TABLE 5.2: 2 Demographics and baseline characteristics. Given are means for years of known disease and Karnofsky score. All other figures denote numbers of patients

	Part A			Part B				
	^{99m} Tc-BIWA 4			¹⁸⁶ Re-BIWA 4 [mCi/m ²]				
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4	20	30	40	50	60
n	3	4	3	2	4	3	6	5
years of known disease	0.11	0.08	0.07	9.30	2.13	1.17	0.80	1.16
Karnofsky score	90	90	100	80	75	76.7	85	80
radical surgery	0	0	0	2	2	1	2	4
lymph nodes affected	2	2	0	2	3	2	1	1
previous chemotherapy	0	0	0	1	3	1	4	1
previous radiotherapy	0	0	0	2	4	3	6	5

5.3 MEASUREMENTS OF TREATMENT COMPLIANCE

10 All medication was administered under the supervision of the investigator or his/her delegate. Blood was collected to determine the pharmacokinetics of BIWA 4. All patients had detectable blood levels of BIWA 4.

5.4 EFFICACY / CLINICAL PHARMACOLOGY RESULTS

15 Three out of six patients experienced stable disease at the maximum tolerated dose of 50 mCi/m². The dose of 50 mg BIWA 4 was confirmed in Part A of the trial to be the optimal dose concerning blood concentrations and selective tumour uptake. The plasma concentrations of BIWA 4 were dose-proportional in the range of 25 mg to 100 mg BIWA 4 in Part A and peaked at 0.9 hours with a terminal elimination half-life of 54 – 74 hours and 94 hours for Part A and Part B, respectively.

20 5.4.1 Analysis of efficacy / pharmacodynamics

5.4.1.1 Primary endpoints

Primary endpoint for efficacy was the biodistribution of ^{99m}Tc-BIWA 4 in Part A. Further primary endpoints were the analyses of the radioimmunoscentigraphic images and the

5 pharmacokinetics (described in section 5.4.2) for Part A and B of the trial, respectively. Dosimetry was done for Part B only.

Biodistribution of ^{99m}Tc -BIWA 4 in Part A

Tissue samples were obtained during surgery and the uptake of ^{99m}Tc -BIWA 4 expressed as % injected dose (ID)/kg tissue.

10 **Intent-to-treat (ITT) subset:** The relative biodistribution of ^{99m}Tc -BIWA 4 was highest in the tumour in all three dosing groups except for patient 1 (25 mg BIWA 4), patient 5 (50 mg BIWA 4, no tumour cells) and patient 9 (100 mg BIWA 4), respectively. The uptake in tumour ranged from 6 to 17 % ID/kg, 5 to 28 % ID/kg and 13 to 17 % ID/kg for the dose group 25 mg, 50 mg and 100 mg, respectively. The calculated mean ratio of tumour uptake versus
15 uptake in bone marrow was 1.7, 2.6 and 2 in the 25 mg, 50 mg and 100 mg dose group, respectively, (TABLES 7.2: 1 and 7.2: 2).

Per-protocol (PP) subset: The relative biodistribution of ^{99m}Tc -BIWA 4 was highest in the tumour in all three dosing groups except for patient 1 (25 mg BIWA 4). The uptake in tumour ranged from 6 to 17 % ID/kg, 23 to 28 % ID/kg and 16 % ID/kg for the dose group 25 mg,
20 50 mg and 100 mg, respectively. The calculated mean ratio of tumour uptake versus uptake in bone marrow was 1.7, 3.2 and 2.5 in the 25 mg, 50 mg and 100 mg dose group, respectively, (TABLES 7.2: 1 and 7.2: 3).

A trend to greater uptake with smaller tumour size appeared to be present for the 25 mg BIWA 4 dose group. A similar evaluation was not possible for the 50 mg and 100 mg BIWA 4
25 dose groups due to the limited number of patients (TABLE 7.2: 4).

High %ID/kg uptake was observed in bone marrow supernatant (up to 17 %ID/kg) and plasma or blood (up to 13 %ID/kg) which was always below the uptake in the tumour in the per protocol population except for patient 1 (25 mg BIWA 4). Uptake in skin and mucosa was always below the uptake in primary tumour in the per protocol subset.

Radioimmunoscentigraphic images of ^{99m}Tc -BIWA 4 in Part A

30 While no radioactivity uptake was found in the tumour directly after infusion, a medium uptake of 1.9 was recorded after 21 hours. Uptake of radioimmunoconjugate appeared to be low in bone marrow and kidney both directly after infusion and 21 hours after infusion. Uptake was low to medium in lung while liver revealed a higher uptake.

35 A similar pattern was observed for the per protocol population. The only difference appeared to be a higher uptake in kidney while the uptake was lower in lung as compared to the intent-to-treat population.

5 **Radioimmunoscintigraphic images of ^{186}Re -BIWA 4 in Part B**

Radioimmunoscintigraphic examinations were done immediately after infusion and 21, 48, 72, 144 and 336 hours after infusion. Hardly any uptake in tumour was observed directly after the infusion. Relative uptake in tumour appeared to be dose-dependent and increased over time reaching medium and high uptake after 72 to 144 hours with a decline after 336 hours.

10 Biodistribution of radioactivity was similar in bone marrow, lung, liver, kidney and intestine and did not reveal the same dose dependent effect (except for the intestine). Moreover, relative uptake of radioactivity appeared to be constant or modestly decreased over time and was similar for all doses of radioactivity. Uptake in bone marrow was lowest for most treatment groups.

15 **5.4.1.2 Secondary endpoint(s)**

Secondary endpoints of efficacy were tumour response to therapy in Part B and the level of CD44v6 expression in tumour in Part A, respectively. Soluble CD44v6 expression was determined for both Part A and Part B of the trial.

CD44v6 expression in tumour in Part A

20 Data of determination of CD44v6 antigen expression is available for seven patients. CD44v6 antigen expression was observed in more than 90% and 80% of the tumour cells in five and two patients, respectively, while CD44v6 antigen expression was detected in more than 90% of the cells of lymph node metastases in the four patients for which lymph node was available for evaluation. Homogenous CD44v6 antigen expression was observed in all patients in the
25 mucosa.

Tumour response in Part B

Patients treated with doses of up to 40 mCi/m^2 ^{186}Re -BIWA 4 did not experience a clinical tumour response. All patients experienced progressive disease. One patient was not evaluable due to intercurrent death.

30 Three of the six patients treated with 50 mCi/m^2 ^{186}Re -BIWA 4 developed stable disease or no change after the first cycle. Two of these three patients who were treated with another cycle with 50 mCi/m^2 ^{186}Re -BIWA 4 had progressive disease after the second cycle. Progressive disease was observed after a total of 148 days (patient 109) and 127 days (patient 116) after first administration of the trial drug, respectively.

35 One patient treated with 60 mCi/m^2 ^{186}Re -BIWA 4 experienced stable disease after the first cycle. This patient went into progressive disease after a second cycle with 50 mCi/m^2 ^{186}Re -BIWA 4 (173 days after the first administration of the trial drug).

The overall time to progression for patients who did not respond to therapy ranged from 0 to 55 days with a mean of about five to six weeks irrespective of the treatment group.

5 **Soluble CD44v6 in Part A and B**

Soluble CD44v6 was determined in all patients treated with BIWA 4 in the present trial.

The amount of soluble CD44v6 detected tended to increase for the ^{186}Re -BIWA 4 treated patients (TABLE 7.2: 6) during the trial in Part B while for patients treated with $^{99\text{m}}\text{Tc}$ in Part A no such trend could be observed.

10 **5.4.2 Drug dose, drug concentration, and relationship to response**

The pharmacokinetics of BIWA 4 will be presented separately for Parts A and B because the radiolabel was different (Part A: $^{99\text{m}}\text{Tc}$; Part B: ^{186}Re).

5.4.2.1 Analytical performance with regard to BIWA 4, HAHAAs and CD44v6

15 Validated assays were used for analysis of plasma samples. Analysis of quality control samples produced results of high accuracy and precision.

Analytical performance with regard to gamma counting of samples

20 The signal linearity of the scintillation crystal in the gamma counter was not tested. It can be assumed based on the type of instrument at least in the energy range that was used in the current study. Calibration samples showed typically 40000-200000 counts per minute vs less than 50-100 counts per minute in background samples. The precision of the seven measurements per calibration standard was regularly within 2%. The same applies for the precision of the individual triplicate measurements in blood, serum and urine. Blood and serum samples showed typically activities between 2000 and 100000 counts per minute.

25 There were no quality control samples for radioactivity counting. The precision of repeated measurements of calibration standards and samples derived from the current study indicate that the precision of triplicate measurements was at least better than 2 %.

5.4.2.1 Pharmacokinetics

Part A.

Part A consisted of 10 patients receiving a single BIWA 4 dose ranging from 25 to 100 mg with a constant radiolabelled dose of 20 mCi ^{99m}Tc. The pharmacokinetic parameters in plasma for BIWA 4 measured by ELISA following a short intravenous infusion are given in TABLE 5.4.2.2: 1.

TABLE 5.4.2.2: 1 Pharmacokinetic parameters for BIWA 4 in plasma from Part A of the study

Patient	BIWA 4 Dose (mg)	tmax (min)	Cmax (ng/mL)	Half- Life (h)	AUC _{inf} (μg•h/mL)	Vz (L)	CL (mL/min)	MRT (h)
1	25	0	6634	44.1	387.4	4.1	1.076	58.0
2	25	30	6640	67.0	534.7	4.5	0.779	94.8
3	25	10	9590	53.3	542.9	3.5	0.768	71.0
4	50	29	14900	64.9	1112.3	4.2	0.749	84.8
5	50	121	15500	102.8	1580.2	4.7	0.527	135.2
8	50	5	13900	53.0	831.0	4.6	1.003	71.7
10	50	41	16700	83.8	1461.6	4.1	0.570	110.0
6	100	4	30500	85.0	2548.9	4.8	0.654	112.0
7	100	17	30600	76.0	2709.6	4.0	0.615	110.4
9	100	30	24500	44.6	1708.1	3.8	0.976	61.1
Geometric mean								
	25 mg	.	7503	54.0	482.7	4.0	0.863	73.1
	50 mg	29	15216	73.8	1208.8	4.4	0.689	97.5
	100 mg	13	28383	66.0	2276.4	4.2	0.732	91.1

The increase in C_{max} BIWA 4 plasma concentrations observed and the area under the curve were proportional to the dose administered (FIGURES 4 and 5).

Serum concentrations of the radiolabel (^{99m}Tc) were also determined and the pharmacokinetic parameters are presented in TABLE 5.4.2.2: 2.

5 TABLE 5.4.2.2: 2 Pharmacokinetic parameters for ^{99m}Tc -BIWA 4 in serum from Part A of the study

Patient	^{99m}Tc -BIWA 4 Dose (mCi)	Tmax (min)	Cmax (%ID/kg)	Half-Life (h)	AUC _{inf} (%ID•h/kg)	Vz (kg)	CL (kg/min)	MRT (h)
1	19.0 mCi	117	52.66	44.6	3376.9	1.9	0.030	63.8
2	20.0 mCi	10	28.08	55.6	2151.7	3.7	0.048	79.1
3	21.4 mCi	10	39.11	27.5	1445.3	2.7	0.072	41.5
4	19.5 mCi	29	31.84	31.3	1432.8	3.2	0.072	43.8
5	19.1 mCi	121	32.29	39.6	1934.8	3.0	0.054	56.3
8	18.8 mCi	114	28.41	42.8	1620.1	3.8	0.060	60.8
10	19.3 mCi	41	40.31	43.1	2247.1	2.8	0.042	62.3
6	19.3 mCi	120	30.45	45.3	1946.5	3.4	0.054	65.2
7	19.8 mCi	77	35.79	36.0	1628.6	3.2	0.060	51.1
9	18.3 mCi	238	31.94	35.5	1805.6	2.8	0.054	50.6
Harmonic mean				38.7				
Geometric mean		57.9	34.5	39.4	1898.0	3.0	0.053	56.5

10 The geometric mean half-life for BIWA 4 measured by ELISA in plasma following intravenous infusion ranged from 54 to 73.8 hours for the three treatment groups studied in Part A. The geometric mean half-life for the ^{99m}Tc -radiolabeled BIWA 4 for the same samples was shorter at 39.4 hours in serum and 46.4 hours in blood. The discrepancy between the two mean estimates was attributed to longer sampling times possible with BIWA 4 due to radioactivity assay restrictions.

15 Radioactivity was excreted into the urine in the amounts collected in TABLE 5.4.2.2: 3. Due to incomplete collections over the 48 hours (i.e. cumulative data may not always be comparable because of varying duration of collection), no further assessment was made of this urinary excretion data. For six patients with complete 48-hour collections, the mean amount of radioactivity collected was 10% of the initial dose.

5 TABLE 5.4.2.2: 3 Amount of radioactivity (^{99m}Tc) excreted (as percent of initial dose) into urine in Part A

Patient	Collection Interval (h)					Cumulative Amount (%ID)
	0-4	4-8	8-12	12-24	24-48	
1	2.80	10.13	5.08	n.a.	n.a.	18.01
2	1.94	0.97	0.52	3.09	3.23	9.75
3	0.29	1.44	1.19	n.a.	n.a.	2.92
4	1.68	1.42	1.95	1.14	2.21	8.40
5	2.20	1.60	1.02	3.96	3.98	12.76
8	1.29	2.45	0.88	3.67	3.00	11.29
10	n.a.	n.a.	n.a.	n.a.	n.a.	
6	2.55	1.59	1.80	1.89	2.00	9.83
7	n.a.	3.14	n.a.	1.79	3.13	8.06
9	1.54	3.34	0.00	1.05	2.31	8.24

n.a. = not applicable, no data available

Part B:

10 Part B consisted of 20 patients receiving BIWA 4 doses of 50 mg with varying radiolabelled doses of ^{186}Re -BIWA 4 ranging from 20-60 mCi/m². Three patients received a second intravenous infusion while the other 17 patients received only a single course of therapy.

15 Graphically, there was consistency between plasma BIWA 4 concentrations and serum radioactivity concentrations for each patient and within the three patients that received ^{186}Re -BIWA 4 on two occasions. This consistency observed graphically contrasts with the perception of a longer half-life for radioactivity when the data was modelled (noncompartmental evaluation). The geometric mean half-life of 122 hours for the radioactive portion of the ^{186}Re -BIWA 4 was longer than the ELISA-determined plasma half-life of 94 hours for BIWA 4.

20 Grouping (TABLES 5.4.2.2: 6 and 5.4.2.2: 7) the pharmacokinetic parameters by amount of radioactivity administered showed some trends of longer exposure with increased radioactivity dosed, but the numbers of individuals in each dose group are too small to make any consistent conclusions.

5 TABLE 5.4.2.2: 6 Geometric means for plasma BIWA 4 pharmacokinetic parameters grouped by amount of radioactivity given.

¹⁸⁶ Re Dose (mCi/m ²)	n	Cmax (ng/mL)	Half-Life (h)	AUC _{inf} (μg•h/mL)	Vz (L)	CL (mL/h)	MRT (h)
20	2	12160	106.5	1217	6.3	41.1	130.3
30	4	15239	86.2	1399	4.4	35.7	116.2
40	3	11121	90.6	1201	5.4	41.7	125.1
50	9*	12982	92.4	1055	6.3	47.4	117.4
60	5	11927	99.8	939	7.7	53.2	115.9

* patients who received two doses of 50 mCi/m² are included

10 TABLE 5.4.2.2: 7 Geometric means for serum ¹⁸⁶Re-BIWA 4 pharmacokinetic parameters grouped by amount of radioactivity given.

¹⁸⁶ Re Dose (mCi/m ²)	n	Cmax (%ID/kg)	Half-Life (h)	AUC _{inf} (%ID•h/kg)	Vz (kg)	CL (kg/h)	MRT (h)
20	2	30.88	114.4	3296	5.01	0.030	154.6
30	4	40.09	98.8	4365	3.27	0.023	134.5
40	3	28.82	126.4	3856	4.73	0.026	172.4
50	9*	40.22	123.3	4182	4.25	0.024	162.5
60	4	32.10	145.2	3493	6.00	0.029	185.5

* patients who received two doses of 50 mCi/m² are included

5.4.2.2 Pharmacokinetic / pharmacodynamic analysis

A pharmacokinetic/pharmacodynamic analysis was not performed.

5 **5.4.3 Statistical / analytical issues**

The analysis was performed as originally planned. Details are described in section 3.7.1.3.

5.4.3.1 Primary endpoint(s)

Not applicable.

5.4.3.2 Secondary endpoint(s)

10 Not applicable.

5.4.4 Drug-drug and drug-disease interactions

No tests concerning drug-drug or drug-disease interactions were done.

5.4.5 By-subject displays

For details see section 6.4.2.2 and 6.4.2.3.

15 **5.5 EFFICACY / CLINICAL PHARMACOLOGY CONCLUSIONS**

Part A:

Results from Part A confirmed a dose of 50 mg BIWA 4 as the optimal dose for treatment based on blood concentrations and tissue uptake level. The distribution as assessed by
20 radioscintigraphy and biopsy measurements was in almost all cases highest in the tumour as compared to other tissues. Uptake of radioactivity increased in the tumour over time. CD44v6 expression was present in more than 80 % and 90% of the cells of primary tumour and lymph node metastases, respectively, and in all mucosa specimens obtained.

The amount of soluble CD44v6 appeared to be constant before and after ^{99m}Tc-BIWA 4 administration.

25 Concentrations of BIWA 4 measured were dose-proportional in the range of 25 mg to 100 mg BIWA 4. A moderate amount of the dose administered was excreted via the kidneys. The % ID excreted in urine was similar for all dosing groups.

Part B:

30 Data from Part B indicate that patients may clinically benefit from ¹⁸⁶Re-BIWA 4 therapy at MTD. One out of five patients treated with 60 mCi/m² had stable disease. Three out of six patients experienced stable disease at the maximum tolerated dose of 50 mCi/m². Time until progression ranged between 127 and 173 days in those patients who received a second dose of 50 mCi/m². Radioimmunoscintigraphy indicates uptake of radioactivity in tumour tissue. Biodistribution was comparable in other tissues irrespective of dose. Dosimetric analysis did
35 not reveal unexpected high absorbed doses in tissues other than the tumour except for the

5 testes. However, the relevance of the observed testes dose for the impairment of fertility is currently unclear.

The amount of soluble CD44v6 detected tended to increase for the ^{186}Re -BIWA 4 treated patients.

10 The plasma concentrations of BIWA 4 peaked at 0.92 hours and the antibody was eliminated with a geometric mean half-life of 94 hours for BIWA 4 determined by ELISA. C_{max} and AUC values (ELISA measurement) were similar to those obtained in Part A of the trial for the 50 mg BIWA 4 dose group.

Dose limiting toxicity occurred at a dose of 60 mCi/m² while a dose of 50 mCi/m² ¹⁸⁶Re-BIWA 4 turned out to be the maximum tolerated dose in the present trial. The dose limiting toxicity were adverse events from the bone marrow i.e. thrombocytopenia and leucopenia.

6.1 EXTENT OF EXPOSURE

All ten patients treated in Part A of the trial were administered one single dose of ^{99m}Tc-BIWA 4. Doses of BIWA 4 were either 25 mg, 50 mg or 100 mg while ^{99m}Tc was 20 mCi.

In Part B, 20 patients were administered one single dose of ¹⁸⁶Re-BIWA 4. Three patients received a second dose of 50 mCi/m² due to stable disease (no change). The dose of BIWA 4 was kept stable at 50 mg while the radioactivity of ¹⁸⁶Re was increased in 10 mCi/m² body surface increments.

The dose was calculated according to body surface area (TABLE 6.1: 1).

TABLE 6.1: 1 Mean dose of radioactivity administered

	Part A			Part B				
	^{99m} Tc-BIWA 4			¹⁸⁶ Re-BIWA 4 [mCi/m ²]				
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4	20	30	40	50	60
N	3	4	3	2	4	3	6	5
mCi ^{99m} Tc	20.1±1. 2	19.2±0. 3	19.1±0. 8	n.a	n.a	n.a	n.a	n.a
mCi ¹⁸⁶ Re	n.a.	n.a	n.a	33.1±1. 7	52.4±5.6	70.6±5. 2	87.2±8. 5	97.2±8. 3

n.a. = not applicable, mean values and standard deviation are given

The radiochemical purity of the drug was more than 95 % and the immunoreactive fraction was higher than 80 %.

An observation period of at least 6 weeks followed the administration of the trial drug.

6.2 ADVERSE EVENTS (AEs)

Adverse events were reported in all patients treated. Two patients discontinued the observation period in Part B due to an AE. No action was taken with the trial drug since treatment had been completed.

Part A:

The majority of patients in Part A experienced adverse events from body as a whole (50 %) which may be also due to the underlying surgery and the associated pain. No special pattern of adverse events was reported and none of the patients developed CTC criteria 3 or 4 which were judged to be drug-related (TABLES 6.2.2: 1 and 2).

Three patients experienced serious adverse events which are described in section 6.3.1.

Part B:

Adverse events of the system organ class (SOC) 'body as a whole' were reported in 65 % of the patients, followed by adverse events from the system organ class 'platelet, bleeding and clotting disorders' (60 % of the patients) and 'white cell and reticulo endothelial system disorders' (50 % of the patients). Drug-related thrombocytopenias and leucopenias occurred in 11 (55 %) and 10 (50 %) patients on average 21 days after start of radioimmunotherapy and were generally reversible after 6 weeks. A dose-response was observed for these events (TABLE 6.2.2: 3).

Haematological dose-limiting toxicity defined as drug-related CTC grade 4 was reported in three patients.

Non-haematological dose-limiting toxicity defined as drug-related CTC grade 3 or 4 non-haematological toxicity occurred in 4 patients (at 30 mCi/m² one patient experienced rash and Quincke's oedema, two patients experienced fever at 60 mCi/m² and one patient treated with 60 mCi/m² experienced fatigue).

Six of the nine patients who experienced serious adverse events died. Deaths and serious adverse events are described in section 12.3.1.

6.2.2 Display of adverse eventsPart A:

The following adverse events sorted by SOC, preferred term and the three dose groups of BIWA 4 were reported in more than one patient in Part A of the trial (TABLE 6.2.2: 1). More details as well as the preferred terms can be found in section 7.3.1.

TABLE 6.2.2: 1: Patients with adverse events in Part A (pre-surgical treatment with a single dose of 20 mCi ^{99m}Tc-BIWA 4)

	^{99m} Tc-BIWA 4			
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4	total
number of patients (% of patients treated)	3 (100)	4 (100)	3 (100)	10 (100)
number of patients with any adverse event	3 (100)	4 (100)	3 (100)	10 (100)
application site disorders	2 (66.7)	0 (0)	0 (0)	2 (20.0)
body as a whole	2 (66.7)	2 (50.0)	1 (33.3)	5 (50.0)
Pain	1 (33.3)	1 (25.0)	1 (33.3)	3 (30.0)

metabolic and nutritional disorders	1 (33.3)	2 (50.0)	1 (33.3)	4 (40.0)
weight decrease	1 (33.3)	1 (25.0)	0 (0)	2 (20.0)
resistance mechanism disorder	0 (0)	4 (100)	0 (0)	4 (40.0)
Infection	0 (0)	3 (75.0)	0 (0)	3 (30.0)
vascular disorders	1 (33.3)	1 (25.0)	1 (33.3)	3 (30.0)
Haemorrhage (not specified)	1 (33.3)	1 (25.0)	0 (0)	2 (20.0)

given are SOCs and preferred terms in case they were reported in more than one patient; figure in brackets denote percentage of patients treated

The events which were considered drug-related by the investigator graded according to CTC criteria are listed in TABLE 6.2.2: 2. The events were grade 1 according to CTC.

TABLE 6.2.2: 2 Severity of drug-related (as judged by the investigator) adverse events according to CTC criteria reported in Part A (pre-surgical treatment with a single dose of 20 mCi ^{99m}Tc -BIWA 4).

	^{99m}Tc -BIWA 4			
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4	total
number of patients	3	4	3	10
Number of adverse events considered drug-related	0	2	0	2
Liver and biliary system disorder	0	2	0	2
sGOT increased	0	1	0	1
sGPT increased	0	1	0	1

sGOT = serum glutamic oxalacetic transaminase ; sGPT = serum glutamic pyruvic transaminase

Part B:

TABLE 6.2.2: 3 presents the number of patients with adverse events sorted by SOC, preferred term and the radiation dose groups of ^{186}Re -BIWA 4.

¹⁸⁶Re-BIWA 4 first and second doses considered)

	¹⁸⁶ Re-BIWA 4 [mCi/m ²]					total
	20	30	40	50	60	
number of patients treated	2 (100)	4 (100)	3 (100)	6 (100)	5 (100)	20 (100)
number of patients with adverse events	2 (100)	4 (100)	3 (100)	6 (100)	5 (100)	20 (100)
body as a whole	2 (100)	3 (75.0)	2 (66.7)	2 (33.3)	4 (80.0)	13 (65.0)
allergic reaction	0 (0)	1 (25.0)	0 (0)	0 (0)	1 (20.0)	2 (10.0)
Fatigue	0 (0)	0 (0)	0 (0)	1 (16.7)	1 (20.0)	2 (10.0)
Fever	0 (0)	0 (0)	2 (66.7)	0 (0)	2 (40.0)	4 (20.0)
oedema mouth	1 (50.0)	0 (0)	0 (0)	1 (16.7)	0 (0)	2 (10.0)
Pain	0 (0)	2 (50.0)	0 (0)	1 (16.7)	1 (20.0)	4 (20.0)
central and peripheral nervous system disorder	0 (0)	1 (25.0)	2 (66.7)	1 (16.7)	0 (0)	4 (20.0)
gastro-intestinal system disorder	0 (0)	1 (25.0)	1 (33.3)	2 (33.3)	3 (60.0)	7 (35.0)
Mucositis	0 (0)	0 (0)	0 (0)	1 (16.7)	3 (60.0)	4 (20.0)
Stomatitis	0 (0)	0 (0)	0 (0)	1 (16.7)	1 (20.0)	2 (10.0)
liver and biliary system disorders	1 (50.0)	0 (0)	0 (0)	0 (0)	1 (20.0)	2 (10.0)
sGPT increased	1 (50.0)	0 (0)	0 (0)	0 (0)	1 (20.0)	2 (10.0)
metabolic and nutritional disorders	1 (50.0)	0 (0)	1 (33.3)	1 (16.7)	1 (20.0)	4 (20.0)
Neoplasm	0 (0)	0 (0)	2 (66.7)	0 (0)	2 (40.0)	4 (20.0)
neoplasm malignant aggravated	0 (0)	0 (0)	1 (33.3)	0 (0)	2 (40.0)	3 (15.0)
platelet, bleeding and clotting disorders	0 (0.0)	1 (25.0)	1 (33.3)	5 (83.3)	5 (100)	12 (60.0)
Thrombocytopenia	0 (0.0)	1 (25.0)	1 (33.3)	4 (66.7)	5 (100)	11 (55.0)
red blood cell disorder	0 (0)	0 (0)	0 (0)	2 (33.3)	3 (60.0)	5 (25.0)
Anaemia	0 (0)	0 (0)	0 (0)	2 (33.3)	3 (60.0)	5 (25.0)
respiratory system disorder	0 (0)	1 (25.0)	1 (33.3)	0 (0)	2 (40.0)	4 (20.0)
Bronchitis	0 (0)	1 (25.0)	0 (0)	0 (0)	1 (20.0)	2 (10.0)
Dyspnoea	0 (0)	0 (0)	0 (0)	0 (0)	2 (40.0)	2 (10.0)
Rhinitis	0 (0)	0 (0)	1 (33.3)	0 (0)	1 (20.0)	2 (10.0)
skin and appendages disorder	1 (50.0)	1 (25.0)	1 (33.3)	0 (0)	1 (20.0)	4 (20.0)
Rash	0 (0)	1 (25.0)	0 (0)	0 (0)	1 (20.0)	2 (10.0)
white cell and RES disorder	0 (0)	0 (0)	1 (33.3)	4 (66.7)	5 (100)	10 (50.0)
Granulocytopenia	0 (0)	0 (0)	0 (0)	1 (16.7)	2 (40.0)	3 (15.0)
Leucopenia	0 (0)	0 (0)	1 (33.3)	4 (66.7)	5 (100)	10 (50.0)

given are SOCs and preferred terms in case they were reported in more than one patient; sGPT = serum glutamic pyruvic transaminase ; RES = reticulo endothelial system, figure in brackets denote per centage of patients treated

5

Drug-related adverse events were reported in 75 % of the patients. Drug-related leucopenia was reported in 50 % and thrombocytopenia in 55 % of the patients, respectively.

Adverse events considered drug-related are provided in TABLE 6.2.2: 4 graded according to the CTC.

CTC criteria reported in Part B (first and second dose considered)

adverse event	¹⁸⁶ Re-BIWA 4 [mCi/m ²]					total
	20	30	40	50	60	
CTC grade 1	1	0	2	6	5	14
face oedema	1	n.a.	0	0	0	1
leucopenia	0	n.a.	1	2	0	3
moniliasis	0	n.a.	0	0	1	1
mucositis	0	n.a.	0	0	1	1
platelets abnormal	0	n.a.	0	1	0	1
purpura	0	n.a.	0	0	1	1
stomatitis	0	n.a.	0	1	0	1
thrombocytopenia	0	n.a.	1	2	2	5
CTC grade 2	0	1	0	7	7	15
anaemia	n.a.	0	n.a.	1	1	2
gout	n.a.	0	n.a.	1	0	1
leucopenia	n.a.	0	n.a.	1	2	3
mucositis	n.a.	0	n.a.	2	3	5
stomatitis	n.a.	0	n.a.	0	1	1
taste loss	n.a.	0	n.a.	1	0	1
thrombocytopenia	n.a.	1	n.a.	1	0	2
CTC grade 3	0	2	0	2	8	12
allergic reaction	n.a.	1	n.a.	0	0	1
fatigue	n.a.	0	n.a.	0	1	1
fever	n.a.	0	n.a.	0	2	2
leucopenia	n.a.	0	n.a.	1	1	2
rash	n.a.	1	n.a.	0	0	1
thrombocytopenia	n.a.	0	n.a.	1	4	5
CTC grade 4	0	0	0	3	5	8
anaemia	n.a.	n.a.	n.a.	0	1	1
granulocytopenia	n.a.	n.a.	n.a.	1	2	3
leucopenia	n.a.	n.a.	n.a.	1	2	3
thrombocytopenia	n.a.	n.a.	n.a.	1	0	1

given is the number of adverse events; n.a. = not applicable

6.2.3 Analysis of adverse events

Part A:

None of the adverse events occurring in Part A of the trial were considered drug-related except for one patient experiencing mild and reversible CTC grade 1 elevation of AST and ALT.

No difference in AE profile was observed when comparing the different BIWA 4 doses.

Part B:

Thrombocytopenia and leucopenia were dose-dependent (TABLES 6.2.2: 3 and 6.2.2: 4) and dose-limiting. Time course is provided in section 6.4.2.1.

causing dose-limiting toxicity (TABLES 6.2.2: 3 and 6.2.2: 4).

An allergic reaction was reported in two patients, one experiencing drug-related Quincke's oedema and rash (30 mCi/m²) and one patient (60 mCi/m²) had an allergic reaction after a thrombocyte concentrate transfusion, which was not considered drug-related. One patient 20 mCi/m²) had urticaria which also was not considered drug-related and one patient (20 mCi/m²) experienced drug-related face oedema.

HAHAs were detected in two patients (see section 12.4.3).

Discontinuations due to adverse events occurred in two patients. Both patients died.

Adverse events required initiation of concomitant therapy in 17 patients. These patients are shortly described in section 6.3.1.

6.3 DEATHS, OTHER SERIOUS ADVERSE EVENTS, AND OTHER SIGNIFICANT ADVERSE EVENTS

Serious adverse events were reported in 12 patients. Of those six patients died in the course of the trial or follow-up. All patients who died were treated in Part B of the trial. None of the patients treated in Part A died. All fatalities were due to disease progression.

Two patients in Part B discontinued the trial prematurely due to adverse events (death in both patient 112 and 105).

Adverse events requiring therapy were most often thrombocytopenia, leucopenia and fever.

INCLUDED IN THE TEXT

7.1 DEMOGRAPHIC DATA

Details of demographics are not included.

7.2 EFFICACY / PHARMACODYNAMIC DATA

TABLE 7.2: 1 Tumour to bone marrow uptake ratio – Part A of the trial

TABLE 7.2: 2 Tumour to bone marrow uptake ratio (ITT subset) – Part A of the trial

TABLE 7.2: 3 Tumour to bone marrow uptake ratio (PP subset) – Part A of the trial

TABLE 7.2: 4 Tumour size and % uptake of antibody – Part A of the trial (per protocol population)

TABLE 7.2: 5 CD44v6 antigen expression in tumour and other tissue – Part A of the trial

TABLE 7.2: 6 Soluble CD44v6 in serum. Mean values are given in ng/mL

Part A				
^{99m} Tc-BIWA 4				
	patient No	uptake tumour (%ID/kg; calculated)	uptake bone marrow (%ID/kg; calculated)	ratio tumour / bone marrow (calculated)
25 mg BIWA 4	1*	6.14	5.58	1.10
	2*	15.78	7.49	2.11
	3*	16.78	8.63	1.95
50 mg BIWA 4	4*	28.10	6.98	4.03
	8*	27.76	7.57	3.67
	10*	22.63	11.54	1.96
100 mg BIWA 4	6*	15.89	6.47	2.46
	7	16.99	7.70	2.21
	9	13.31	10.11	1.32

No = number; * = included in per protocol analysis; % ID = % injected dose; calculations were done by taking different weights of samples into account

TABLE 7.2: 2 Tumour to bone marrow uptake ratio (ITT subset) – Part A of the trial

Part A	
^{99m} Tc-BIWA 4	
	ratio tumour / bone marrow (calculated)
25 mg BIWA 4	1.72
50 mg BIWA 4	2.57
100 mg BIWA 4	1.99

mean values are given; ITT = intent-to-treat; calculations were done by taking different weights of samples into account

TABLE 7.2: 3 Tumour to bone marrow uptake ratio (PP subset) – Part A of the trial

Part A	
^{99m} Tc-BIWA 4	
	ratio tumour / bone marrow (calculated)
25 mg BIWA 4	1.72
50 mg BIWA 4	3.22
100 mg BIWA 4	2.46

mean values are given; PP = per protocol; calculations were done by taking different weights of samples into account

population)

Part A			
^{99m} Tc-BIWA 4			
	patient No	tumour size [mm ²]	uptake tumour (%ID/kg; calculated)
25 mg BIWA 4	1	1350	6.14
	2	900	15.78
	3	225	16.78
50 mg BIWA 4	4	n.a.	28.10
	8	1504	27.76
	10	1400	22.63
100 mg BIWA 4	6	800	15.89

No = patient number, n.a. = not applicable, only length or width given but not length and width; calculations were done by taking different weights of samples into account

TABLE 7.2: 5 CD44v6 antigen expression in tumour and other tissue – Part A of the trial

Part A				
^{99m} Tc-BIWA 4				
	patient No	CD44v6 antigen expression in tumour	CD44v6 expression in mucosa	CD44v6 expression in lymph node metastases
25 mg BIWA 4	1*	n.a.	n.a.	n.a.
	2*	++/+++; > 80 %	+++	++/+++; > 90 %
	3*	+++; > 90 %	+++	+++; > 90 %
50 mg BIWA 4	4*	+++; > 90 %	+++	+++; > 90 %
	8*	n.a.	n.a.	n.a.
	10*	++; > 90 %	+++	+++; > 90 %
100 mg BIWA 4	6*	+++; > 80 %	+++	n.a.
	7	+++; > 90 %	+++	n.a.
	9	+++; > 90 %	+++	n.a.

No = number; * = included in per protocol analysis, n.a. = not applicable

	Part A			Part B				
	^{99m} Tc-BIWA 4			¹⁸⁶ Re-BIWA 4 [mCi/m ²]				
	25 mg BIWA 4	50 mg BIWA 4	100 mg BIWA 4	20	30	40	50	60
N	3	4	3	2	4	3	6	5
pre-dose	203	337	174	408	262	346	171	160
21 hours	223	301	179	396	177	403	199	170
48 hours	200	275	157	384	198	423	199	174
72 hours	140	386	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
144 hours	188	282	182	444	284	516	233	195
240/336 hours	n.a.	n.a.	n.a.	445	337	n.a.	n.a.	n.a.
week 6	203	310	210	289	220	487	226	162

n.a. = not applicable, data include the second dosing

Efficacy results:**Part A:**

Results from Part A confirmed a dose of 50 mg BIWA 4 as the optimal dose for treatment based on blood concentrations and tissue uptake levels. The distribution as assessed by radioscintigraphy and biopsy measurements was in almost all cases highest in the tumour as compared to other tissues. Uptake of radioactivity increased in the tumour over time. CD44v6 expression was present in more than 80 % and 90% of the cells of primary tumour and lymph node metastases, respectively, and in all mucosa specimens obtained.

No correlation was observed between tumour size and uptake of radioactivity in the 50 mg BIWA 4 dose group. The amount of soluble CD44v6 appeared to be constant before and after ^{99m}Tc -BIWA 4 administration.

Concentrations of BIWA 4 measured were dose-proportional in the range of 25 mg to 100 mg BIWA 4. A moderate amount of the dose administered was excreted via the kidneys. The percent injected dose (% ID) excreted in urine was similar for all dosing groups.

Part B:

Data from Part B indicate that patients may clinically benefit from ^{186}Re -BIWA 4 therapy at maximum tolerated dose (MTD). One out of five patients treated with 60 mCi/m² had stable disease. Three out of six patients experienced stable disease at the maximum tolerated dose of 50 mCi/m². Time until progression ranged between 127 and 173 days in those patients who received a second dose of 50 mCi/m². Radioimmunosintigraphy indicates uptake of radioactivity in tumour tissue. Biodistribution was comparable in other tissues irrespective of dose. Dosimetric analysis did not reveal unexpected high absorbed doses in tissues other than the tumour except for the testes. The amount of soluble CD44v6 detected tended to increase for the ^{186}Re -BIWA 4 treated patients.

The plasma concentrations of BIWA 4 peaked at 0.92 hours and the antibody was eliminated with a geometric mean half-life of 94 hours for BIWA 4 determined by enzyme-linked immuno-sorbent assay (ELISA) measurement. C_{max} and AUC values (ELISA) were similar to those obtained in Part A of the trial for the 50 mg BIWA 4 dose group.

Tolerability of single dose BIWA 4 coupled to low radiation dose of Technetium-99 was acceptable. Two of the three serious adverse events were due to complications as a result of surgery.

Part B:

The maximum tolerated dose (MTD) was 50 mCi/m² ¹⁸⁶Re-BIWA 4.

Dose-limiting adverse events were dose dependent reversible reductions in thrombocyte and leucocyte count with subsequent fever in individual patients. Clinical symptoms of thrombocytopenia were mild petechiae.

Mucositis was observed in patients treated with higher radiation doses but was not dose-limiting.

No relevant changes in thyroid stimulating hormone (TSH) values were observed during the course of the trial.

Twelve patients experienced a serious adverse event. Of those, six patients died during the course of Part B of the trial mainly due to progression of the underlying disease.

Allergic reactions were observed rarely with one serious drug-related Quincke's oedema. No allergic reactions occurred during or shortly after infusion of the drug.

Two patients developed HAHAs. Repeated dosing did not induce HAHA development.

Conclusions:

Results indicate uptake of ¹⁸⁶Re-BIWA 4 in tumour tissue and clinical benefit in patients with advanced squamous cell carcinoma of the head and neck. The safety profile appears to be acceptable. BIWA 4 showed dose-proportional pharmacokinetics and tumour uptake did not change relevantly between doses of 50 mg and 100 mg BIWA4.

Example 4

Introduction. Patients with an advanced stage of head and neck squamous cell carcinoma (HNSCC) have an increased risk for development of locoregional recurrent tumors and/or distant metastases. For these patients, development of an effective adjuvant systemic treatment is needed. Knowing that HNSCC are intrinsically radiosensitive, targeting of radionuclides selectively to HNSCC by use of monoclonal antibodies as a form of radioimmunotherapy, might contribute to a more effective therapy.

Objective. To determine the safety, maximum tolerated dose (MTD), immunogenicity and efficacy of radioimmunotherapy with Rhenium-186 (^{186}Re)-labelled humanised monoclonal antibody BIWA4 in patients with HNSCC.

Patients and methods. A phase I dose escalation study was conducted in HNSCC patients for whom no curative therapeutic options were available. In a total of 20 patients ^{186}Re -labelled BIWA4 was administered intravenously in doses of 20, 30, 40, 50 or 60 mCi/m². Three patients received, at least 3 months after a dose of 50 or 60 mCi/m², a second dose of 50 mCi/m².

Results. All single as well as repeated administrations were well tolerated and no signs of acute adverse events were observed. The only significant manifestations of toxicity at the higher doses were oral mucositis and dose-limiting myelotoxicity consisting of thrombo- and leucocytopenia. The MTD was established at 50 mCi/m². One patient developed a human-anti-human response after one single administration. Stable disease, lasting 4 to 19 weeks, was observed in 5 patients treated at the highest dose levels.

Conclusion. Radioimmunotherapy with ^{186}Re -labelled BIWA4 in HNSCC patients seems to be safe and tumoricidal doses can be reached. Moreover, due to the low rate of immunogenicity, repeated administrations appear possible. The results of this phase I study encourage the further development of radioimmunotherapy with Rhenium-186 (^{186}Re)-labelled humanised monoclonal antibody BIWA4 towards an adjuvant therapy for head and neck cancer patients.

SEQUENCE LISTING

<110> Boehringer Ingelheim International GmbH

5 <120> Antibodies specific for CD44v6

<130> 1-1212 PCT

<140> not yet assigned

10 <141> 2002-05-16

<150> EP 01112237.1

<151> 2001-05-18

15 <150> US 60/325147

<151> 2001-09-26

<160> 18

20 <170> PatentIn Ver. 2.1

<210> 1

<211> 114

<212> PRT

25 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

30

<400> 1

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

35 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Asp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

40 Ser Thr Ile Ser Ser Gly Gly Ser Tyr Thr Tyr Tyr Leu Asp Ser Ile
50 55 60

45 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

50 Ala Arg Gln Gly Leu Asp Tyr Trp Gly Arg Gly Thr Leu Val Thr Val
100 105 110

Ser Ser

55

<210> 2

<211> 106

<212> PRT

60 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

<400> 2

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Ser Ala Ser Ser Ser Ile Asn Tyr Ile
 20 25 30

Tyr Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Tyr
 35 40 45

Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60

Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro Glu
 65 70 75 80

Asp Phe Ala Val Tyr Tyr Cys Leu Gln Trp Ser Ser Asn Pro Leu Thr
 85 90 95

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 3

<211> 106

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

<400> 3

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Ser Ala Ser Ser Ser Ile Asn Tyr Ile
 20 25 30

Tyr Trp Leu Gln Gln Lys Pro Gly Gln Ala Pro Arg Ile Leu Ile Tyr
 35 40 45

Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60

Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro Glu
 65 70 75 80

Asp Phe Ala Val Tyr Tyr Cys Leu Gln Trp Ser Ser Asn Pro Leu Thr
 85 90 95

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 4

<211> 342

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

<400> 4

```

gaagtgcagc tgggtggagtc tggggggaggc ttagtgaagc ctggaggggc cctaagactc 60
tcctgtgcag cctctggatt cactttcagt agctatgaca tgtcttgggt tcgccaggct 120
5 ccggggaagg ggctggagtg ggtctcaacc attagtagtg gtggtagtta cacctactat 180
ctagacagta taaagggccg attcaccatc tccagagaca atgccaagaa ctccctgtac 240
ctgcaaataa acagtctgag ggctgaggac acggccgtgt attactgtgc aagacagggg 300
ttggactact ggggtcgagg aaccttagtc accgtctcct ca 342

```

<210> 5

<211> 318

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized antibody sequence

<400> 5

```

gaaattgttc tcacccagtc tccagcaacc ctgtctctgt ctccagggga gagggccacc 60
ctgtcctgca gtgccagctc aagtataaat tacatatact ggtaccagca gaagccagga 120
caggctccta gactcttgat ttatctcaca tccaacctgg cttctggagt ccctgcgcgc 180
ttcagtggca gtgggtcttg aaccgacttc actctcacia tcagcagcct ggagcctgaa 240
25 gattttgccc tttattactg cctgcagtgg agtagtaacc cgctcacatt cggtggtggg 300
accaaggtgg agattaata 318

```

<210> 6

<211> 318

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized antibody sequence

<400> 6

```

gaaattgttc tcacccagtc tccagcaacc ctgtctctgt ctccagggga gagggccacc 60
40 ctgtcctgca gtgccagctc aagtataaat tacatatact ggctccagca gaagccagga 120
caggctccta gaatcttgat ttatctcaca tccaacctgg cttctggagt ccctgcgcgc 180
ttcagtggca gtgggtcttg aaccgacttc actctcacia tcagcagcct ggagcctgaa 240
gattttgccc tttattactg cctgcagtgg agtagtaacc cgctcacatt cggtggtggg 300
accaaggtgg agattaata 318

```

<210> 7

<211> 444

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized antibody sequence

<400> 7

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1          5          10          15
60 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20          25          30
Asp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35          40          45

```

Ser Thr Ile Ser Ser Gly Gly Ser Tyr Thr Tyr Tyr Leu Asp Ser Ile
 50 55 60
 5 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 10 Ala Arg Gln Gly Leu Asp Tyr Trp Gly Arg Gly Thr Leu Val Thr Val
 100 105 110
 Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser
 115 120 125
 Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys
 130 135 140
 20 Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu
 145 150 155 160
 Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu
 165 170 175
 25 Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr
 180 185 190
 Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val
 195 200 205
 Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro
 210 215 220
 35 Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe
 225 230 235 240
 Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 245 250 255
 40 Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe
 260 265 270
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 275 280 285
 Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr
 290 295 300
 50 Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
 305 310 315 320
 Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala
 325 330 335
 55 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 340 345 350
 Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 355 360 365
 60 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 370 375 380

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
385 390 395 400

Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
405 410 415

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
420 425 430

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
435 440

<210> 8

<211> 213

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

<400> 8

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Ser Ala Ser Ser Ser Ile Asn Tyr Ile
20 25 30

Tyr Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Tyr
35 40 45

Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro Glu
65 70 75 80

Asp Phe Ala Val Tyr Tyr Cys Leu Gln Trp Ser Ser Asn Pro Leu Thr
85 90 95

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130 135 140

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145 150 155 160

Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
165 170 175

Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
180 185 190

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
195 200 205

Asn Arg Gly Glu Cys
210

<210> 9
 <211> 213
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:humanized
 antibody sequence

<400> 9
 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Ser Ala Ser Ser Ser Ile Asn Tyr Ile
 20 25 30
 Tyr Trp Leu Gln Gln Lys Pro Gly Gln Ala Pro Arg Ile Leu Ile Tyr
 35 40 45
 Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60
 Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro Glu
 65 70 75 80
 Asp Phe Ala Val Tyr Tyr Cys Leu Gln Trp Ser Ser Asn Pro Leu Thr
 85 90 95
 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
 100 105 110
 Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
 115 120 125
 Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
 130 135 140
 Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
 145 150 155 160
 Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
 165 170 175
 Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
 180 185 190
 Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
 195 200 205
 Asn Arg Gly Glu Cys
 210

<210> 10
 <211> 2135
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:humanized
 antibody sequence

<400> 10

aagctttgac agacgcacaa ccctggactc ccaagtcttt ctcttcagtg acaaacacag 60
 acataggata tcacatttgc ttctgacaca actgtgttca ctagcagcct caaacagaca 120
 ccatgaactt tgggctcagc ttgattttcc ttgtcctaata tttaaaagggt gtccagtggtg 180
 5 aagtgcagct ggtggagtct gggggaggct tagtgaagcc tggaggggtcc ctaagactct 240
 cctgtgcagc ctctggattc actttcagta gctatgacat gtcttgggtt cgccaggctc 300
 cggggaagg gctggagtgg gtctcaacca ttagtagtgg tggtagttac acctactatc 360
 tagacagtat aaagggccga ttcaccatct ccagagacaa tgccaagaac tccctgtacc 420
 tgcaaatgaa cagtctgagg gctgaggaca cggccgtgta ttactgtgca agacaggggt 480
 10 tggactactg gggctcagga accttagtca ccgtctcctc agctagcacc aagggcccat 540
 cggctcttccc cctggcacc cctccaaga gcacctctgg gggcacagcg gccctgggct 600
 gcctggtcaa ggactacttc cccgaaccgg tgacgggtgc gtggaaactca ggcgccctga 660
 ccagcggcgt gcacaccttc ccggtgttcc tacagtcctc aggactctac tccctcagca 720
 gcgtggtgac cgtgccctcc agcagcttgg gcacccagac ctacatctgc aacgtgaatc 780
 15 acaagcccag caacaccaag gtggacaaga aagttggtga gaggccagca caggaggga 840
 ggggtgtctgc tgggaagcagg ctccagcgtc ctgcctggac gcatcccggc tatgcagccc 900
 cagtccaggc cagcaaggca ggccccgtct gcctcttcac ccggagcctc tgcccgcccc 960
 actcatgctc agggagaggg tcttctggct ttttcccagg ctctgggcag gcacaggcta 1020
 ggtgccccta acccaggccc tgcacacaaa ggggcagggt ctgggctcag acctgccaag 1080
 20 agccatatcc gggaggacc tggccctgac ctaagcccac cccaaaggcc aaactctcca 1140
 ctccctcagc tcggacacct tctctcctcc cagattccag taactcccaa tcttctctct 1200
 gcagagccca aatcttgtga caaaactcac acatgccac cgtgccagc taagccagcc 1260
 caggcctcgc cctccagctc aaggcgggac aggtgcccta gagtagcctg catccaggga 1320
 caggccccag ccgggtgctg acacgtccac ctccatctct tctcagcac ctgaactcct 1380
 25 ggggggaccg tcagtcttcc tcttcccccc aaaacccaag gacacctca tgatctccc 1440
 gacccctgag gtcacatgcg tgggtggtgga cgtgagccac gaagaccctg aggtcaagtt 1500
 caactggtac gtggacggcg tggaggtgca taatgccaag acaaagccgc gggaggagca 1560
 gtacaacagc acgtaccggg tggtcagcgt cctcaccgtc ctgcaccagg actggctgaa 1620
 tggcaaggag tacaagtgca aggtctccaa caaagccctc ccagccccca tcgagaaaac 1680
 30 catctccaaa gccaaagggt ggacccgtgg ggtgcgaggg ccacatggac agaggccggc 1740
 tcggcccacc ctctgccctg agagtgaacc ctgtaccaac ctctgtccta cagggcagcc 1800
 ccgagaacca caggtgtaca ccctgcccc atcccgggat gagctgacca agaaccaggt 1860
 cagcctgacc tgcctggtca aaggcttcta tcccagcgac atcgccgtgg agtgggagag 1920
 caatgggcag ccggagaaca actacaagac cagcctccc gtgctggact ccgacggctc 1980
 35 cttcttctc tacagcaagc tcaccgtgga caagagcagg tggcagcagg ggaacgtctt 2040
 ctcatgctcc gtgatgcatg aggtcttgca caaccactac acgcagaaga gcctctccct 2100
 gtctccgggt aaatgagtgc gacggccgcg aattc 2135

<210> 11

<211> 785

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
 antibody sequence

<400> 11

50 aagcttgatc ttcaggatat cacatttgct tctgacacaa ctgtgttcac tagcaacctc 60
 aaacagacac catggatttt cagggtgcaga ttttcagctt cctgctaata agtgcctcag 120
 tcataatgct caggggagaa attgtttctca ccagctctcc agcaaccctg tctctgtctc 180
 caggggagag ggccaccctg tctgagcagt ccagctcaag tataaattac atatactgg 240
 accagcagaa gccaggacag gctcctagac tcttgattta tctcacatcc aacctggctt 300
 55 ctggagtccc tgcgcgcttc agtggcagtg ggtctggaac cgacttcact ctcaaatca 360
 gcagcctgga gcctgaagat tttgccgttt attactgcct gcagtggagt agtaaccgc 420
 tcacattcgg tgggtgggacc aagggtggaga ttaaacggac tgtggctgca ccatctgtct 480
 tcatcttccc gccatctgat gagcagttga aatctggaac tgctagcgtt gtgtgcctgc 540
 tgaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 600
 60 cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc tacagcctca 660
 gcagcaccct gacgctgagc aaagcagact acgagaaaca caaagtctac gcctgcgaag 720
 tcacccatca gggcctgagc tcgcccgtca caaagagctt caacagggga gagtgtagg 780
 aattc 785

<210> 12

<211> 785

<212> DNA

5 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

10

<400> 12

```

aagcttgatc ttcaggatat cacatttgct tctgacacaa ctgtgtttcac tagcaacctc 60
aaacagacac catggattht caggtgcaga ttttcagctt cctgctaata agtgcctcag 120
tcataatgtc caggggagaa attgttctca cccagtctcc agcaaccctg tctctgtctc 180
15 caggggagag ggccaccctg tcctgcagtg ccagctcaag tataaattac atatactggc 240
tccagcagaa gccaggacag gctcctagaa tcttgattta tctcacatcc aacctggctt 300
ctggagtccc tgcgcgcttc agtggcagtg ggtctggaac cgacttcact ctcacaatca 360
gcagcctgga gcctgaagat tttgccgttt attactgcct gcagtggagt agtaaccgcg 420
tcacattcgg tgggtgggacc aaggtggaga ttaaaccggac tgtggctgca ccatctgtct 480
20 tcatcttccc gccatctgat gagcagttga aatctggaac tgctagcgtt gtgtgcctgc 540
tgaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac gccctccaat 600
cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc tacagcctca 660
gcagcaccct gacgctgagc aaagcagact acgagaaaca caaagtctac gcctgcgaag 720
tcacccatca gggcctgagc tcgcccgtca caaagagctt caacagggga gagtgttagg 780
25 aattc 785

```

<210> 13

<211> 1392

30 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:humanized
antibody sequence

35

<400> 13

```

atggagtttg ggctgagctg gctttttctt gtggctatth taaaagggtgt ccagtgtgaa 60
gtgcagctgg tggagtctgg gggaggctta gtgaagcctg gagggctcct aagactctcc 120
40 tgtgcagcct ctggattcac tttcagtagc tatgacatgt cttgggttcg ccaggctccg 180
gggaaggggc tggagtgggt ctcaaccatt agtagtggtg gtagttacac ctactatcta 240
gacagtataa agggccgatt caccatctcc agagacaatg ccaagaactc cctgtacctg 300
caaatagaaca gtctgagggc tgaggacacg gccgtgtatt actgtgcaag acaggggttg 360
gactactggg gtcgaggaac cttagtcacc gtctcctcag ctagcaccaa gggcccatcg 420
45 gtcttcccc tggcaccctc ctccaagagc acctctgggg gcacagcggc cctgggctgc 480
ctgggtcaagg actacttccc cgaaccgggtg acggtgtcgt ggaactcagg cgccctgacc 540
agcggcgtgc acaccttccc ggctgtccta cagtcctcag gactctactc cctcagcagc 600
gtgggtgaccg tgccctccag cagcttgggg acccagacct acatctgcaa cgtgaatcac 660
aagcccagca acaccaagggt ggacaagaaa gttgagccca aatcttgtga caaaactcac 720
50 acatgcccac cgtgcccagc acctgaactc ctgggggggac cgtcagtctt cctcttcccc 780
ccaaaaccca aggacaccct catgatctcc cggacccttg aggtcacatg cgtgggtggg 840
gacgtgagcc acgaagaccc tgagggtcaag ttcaactggg acgtggacgg cgtggaggtg 900
cataatgcca agacaaagcc gcgggaggag cagtacaaca gcacgtaccg tgtgggtcagc 960
gtcctcaccg tcctgcacca ggactggctg aatggcaagg agtacaagtg caaggtctcc 1020
55 aacaaagccc tcccagcccc catcgagaaa accatctcca aagccaaagg gcagccccga 1080
gaaccacagg tgtacaccct gcccccatcc cgggatgagc tgaccaagaa ccaggtcagc 1140
ctgacctgcc tgggtcaaagg cttctatccc agcgacatcg ccgtggagtg ggagagcaat 1200
gggcagccgg agaacaacta caagaccacg cctcccgtgc tggactccga cggctccttc 1260
ttcctctaca gcaagctcac cgtggacaag agcaggtggc agcaggggaa cgtcttctca 1320
60 tgctccgtga tgcattgaggc tctgcacaac cactacacgc agaagagcct ctccctgtct 1380
ccgggtaaat ga 1392

```

<210> 14

<211> 702
<212> DNA
<213> Artificial Sequence

5 <220>
<223> Description of Artificial Sequence:humanized
antibody sequence

<400> 14

10 atggaagccc cagctcagct tctcttcctc ctgctgctct ggctcccaga taccaccgga 60
gaaattgttc tcacccagtc tccagcaacc ctgtctctgt ctccagggga gagggccacc 120
ctgtcctgca gtgccagctc aagtataaat tacatatact ggtaccagca gaagccagga 180
caggctccta gactcttgat ttatctcaca tccaacctgg cttctggagt ccctgcgcgc 240
ttcagtggca gtgggtcttg aaccgacttc actctcaca tcagcagcct ggagcctgaa 300
15 gattttgccg tttattactg cctgcagtgg agtagtaacc cgctcacatt cgggtggtggg 360
accaaggtgg agattaaacg tacggtggct gcaccatctg tcttcatctt cccgccatct 420
gatgagcagt tgaaatctgg aactgcctct gttgtgtgcc tgctgaataa cttctatccc 480
agagaggcca aagtacagtg gaaggtggat aacgccctcc aatcgggtaa ctcccaggag 540
agtgtcacag agcaggacag caaggacagc acctacagcc tcagcagcac cctgacgctg 600
20 agcaaagcag actacgagaa acacaaagtc tacgcctgcg aagtcaccca tcagggcctg 660
agctcgcccg tcacaaagag cttcaacagg ggagagtgtt ga 702

<210> 15
25 <211> 702
<212> DNA
<213> Artificial Sequence

<220>
30 <223> Description of Artificial Sequence:humanized
antibody sequence

<400> 15

atggaagccc cagctcagct tctcttcctc ctgctgctct ggctcccaga taccaccgga 60
35 gaaattgttc tcacccagtc tccagcaacc ctgtctctgt ctccagggga gagggccacc 120
ctgtcctgca gtgccagctc aagtataaat tacatatact ggctccagca gaagccagga 180
caggctccta gaatcttgat ttatctcaca tccaacctgg cttctggagt ccctgcgcgc 240
ttcagtggca gtgggtcttg aaccgacttc actctcaca tcagcagcct ggagcctgaa 300
gattttgccg tttattactg cctgcagtgg agtagtaacc cgctcacatt cgggtggtggg 360
40 accaaggtgg agattaaacg tacggtggct gcaccatctg tcttcatctt cccgccatct 420
gatgagcagt tgaaatctgg aactgcctct gttgtgtgcc tgctgaataa cttctatccc 480
agagaggcca aagtacagtg gaaggtggat aacgccctcc aatcgggtaa ctcccaggag 540
agtgtcacag agcaggacag caaggacagc acctacagcc tcagcagcac cctgacgctg 600
agcaaagcag actacgagaa acacaaagtc tacgcctgcg aagtcaccca tcagggcctg 660
45 agctcgcccg tcacaaagag cttcaacagg ggagagtgtt ga 702

<210> 16
50 <211> 9568
<212> DNA
<213> Artificial Sequence

<220>
55 <223> Description of Artificial Sequence:humanized
antibody sequence

<400> 16

ctgacgcgcc ctgtagcggc gcattaagcg cggcgggtgt ggtggttacg cgcagcgtga 60
ccgctacact tgccagcgcc ctagcgcccg ctcttttcgc tttcttcctt tcctttctcg 120
60 ccacgttcgc cggctttccc cgtcaagctc taaatcgggg gctcccttta gggttccgat 180
ttagtgcttt acggcacctc gaccccaaaa aacttgatta ggggtgatgg tcacgtaggg 240
tcgcgacgta ccgggcccc cctcgattaa ttaatcgagc tactagcttt gcttctcaat 300
ttcttatatt cataatgaga aaaaaaggaa aattaatttt aacaccaatt cagtagttga 360
ttgagcaaat gcgttgccaa aaaggatgct ttagagacag tgttctctgc acagataagg 420

	acaaacatta	ttcagagggga	gtacccagag	ctgagactcc	taagccagtg	agtggcacag	480
	cattctaggg	agaaatatgc	ttgtcatcac	cgaagcctga	ttccgtagag	ccacaccttg	540
	gtaagggcca	atctgctcac	acaggataga	gagggcagga	gccagggcag	agcatataag	600
	gtgaggtagg	atcagttgct	cctcacattt	gcttctgaca	tagttgtgcc	agcatggagg	660
5	aatcgatcct	ccatgcttga	acaagatgga	ttgcacgcag	gttctccggc	cgcttgggtg	720
	gagaggctat	tcggctatga	ctgggcacaa	cagacaatcg	gctgctctga	tgccgccgtg	780
	ttccggctgt	cagcgcaggg	gcgcccgggt	ctttttgtca	agaccgacct	gtccgggtgcc	840
	ctgaatgaac	tgcaggtaag	tgcggccgct	ctaggcctcc	aaaaaagcct	cctcactact	900
	tctggaatag	ctcagaggcc	gaggcggcct	cggcctctgc	ataaataaaa	aaaattagtc	960
10	agccatgcat	ggggcggaga	atgggcggaa	ctgggcggag	ttagggggcg	gatgggcgga	1020
	gttagggggcg	ggactatggg	tgctgactaa	ttgagatgca	tgctttgcat	acttctgcct	1080
	gctggggagc	ctggggactt	tccacacctg	gttgctgact	aattgagatg	catgctttgc	1140
	atacttctgc	ctgctgggga	gcctggggac	tttccacacc	ctaactgaca	cacattccac	1200
	agaattaatt	cccctagtta	ttaatagtaa	tcaattacgg	ggtcattagt	tcatagccca	1260
15	tatatggagt	tcgcggttac	ataacttacg	gtaaatggcc	cgcctggctg	accgcccac	1320
	gacccccgcc	cattgacgtc	aataatgacg	tatgttccca	tagtaacgcc	aatagggact	1380
	ttccattgac	gtcaatgggt	ggactattta	cggtaaactg	cccacttggc	agtacatcaa	1440
	gtgtatcata	tgccaagtac	gccccctatt	gacgtcaatg	acggtaaattg	gcccgcctgg	1500
	cattatgccc	agtacatgac	cttatgggac	tttccactct	ggcagtacat	ctacgtatta	1560
20	gtcatcgcta	ttaccatggg	gatgcgggtt	tggcagtaca	tcaatgggag	tggatagcgg	1620
	tttgactcac	ggggattttcc	aagtctccac	cccattgacg	tcaatgggag	tttggttttg	1680
	cacaaaaatc	aacgggactt	tccaaaatgt	cgtaacaact	ccgccccatt	gacgcaaattg	1740
	ggcggtaggc	gtgtacgggtg	ggaggtctat	ataagcagag	ctgggtacgt	gaaccgtcag	1800
	atcgccctgga	gacgccatca	cagatctctc	accatggaag	ccccagctca	gcttctcttc	1860
25	ctcctgctgc	tctggctccc	agataccacc	ggagaaattg	ttctcaccac	gtctccagca	1920
	accctgtctc	tgtctccagg	ggagagggcc	accctgtcct	gcagtgccag	ctcaagtata	1980
	aattacatat	actggtacca	gcagaagcca	ggacaggctc	ctagactctt	gatttatctc	2040
	acatccaacc	tggcttcttg	agtccctgcg	cgcttcagtg	gcagtgggtc	tggaaaccgac	2100
	ttcactctca	caatcagcag	cctggagcct	gaagattttg	ccgtttatta	ctgcctgcag	2160
30	tggagtagta	acccgctcac	attcgggtgg	gggaccaagg	tggagattaa	acgtacgggtg	2220
	gctgcaccat	ctgtcttcat	cttcccgcga	tctgatgagc	agttgaaatc	tggaaactgcc	2280
	tctgttgtgt	gcctgctgaa	taacttctat	cccagagagg	ccaaagtaca	gtggaagggtg	2340
	gataacgccc	tccaatcggg	taactcccag	gagagtgtca	cagagcagga	cagcaaggac	2400
	agcacctaca	gcctcagcag	caccctgacg	ctgagcaaag	cagactacga	gaaacacaaa	2460
35	gtctacgcct	gcgaagtcac	ccatcagggc	ctgagctcgc	ccgtcacaaa	gagcttcaac	2520
	aggggagagt	gttgaattca	gatccgttaa	cggttaccaa	ctacctagac	tggattcgtg	2580
	acaacatgcg	gccgtgatat	ctacgtatga	tcagcctoga	ctgtgccttc	tagttgccag	2640
	ccatctgttg	tttgcccctc	ccccgtgcct	tccttgacct	tggaaagggtg	cactcccact	2700
	gtccttttct	aataaaaatga	ggaaattgca	tcgcattgtc	tgagtagggtg	tcattctatt	2760
40	ctgggggggtg	gggtggggca	ggacagcaag	ggggaggatt	gggaagacaa	tagcaggcat	2820
	gctgggggatg	cggtgggctc	tatggaacca	gctgggacta	gtagctttgc	ttctcaattt	2880
	cttattttgca	taatgagaaa	aaaaggaaaa	ttaattttta	caccaattca	gtagttgatt	2940
	gagcaaatgc	gttgccaaaa	aggatgcttt	agagacagtg	ttctctgcac	agataaggac	3000
	aaacattatt	cagagggagt	accagagact	gagactccta	agccagttag	tggcacagca	3060
45	ttctagggag	aaatatgctt	gtcatcaccg	aagcctgatt	ccgtagagcc	acaccttggt	3120
	aaggggccaat	ctgctcacac	aggatagaga	gggcaggagc	cagggcagag	catataaggt	3180
	gaggtaggat	cagttgctcc	tcacattttgc	ttctgacata	gttgtgttg	gagcttggat	3240
	agcttggaca	gctcagggct	gcgattttgc	gccaaacttg	acggcaatcc	tagcgtgaag	3300
	gctggtagga	ttttatcccc	gctgccatca	tggttcagacc	attgaactgc	atcgtcgccg	3360
50	tgtcccaaaa	tatggggatt	ggcaagaacg	gagacctacc	ctggcctccg	ctcaggaacg	3420
	agttcaagta	cttccaaaga	atgaccacaa	cctcttcagt	ggaaggtaaa	cagaatctgg	3480
	tgattatggg	taggaaaacc	tggttctcca	ttcctgagaa	gaatcgacct	ttaaaggaca	3540
	gaattaatat	agttctcagt	agagaactca	aagaaccacc	acgaggagct	cattttcttg	3600
	ccaaaagttt	ggatgatgcc	ttaagactta	ttgaacaacc	ggaattggca	agtaaagtag	3660
55	acatgggtttg	gatagtcgga	ggcagttctg	tttaccagga	agccatgaat	caaccaggcc	3720
	accttagact	ctttgtgaca	aggatcatgc	aggaatttga	aagtgcacag	tttttcccag	3780
	aaattgattt	ggggaaatat	aaacttctcc	cagaataccc	aggcgtcctc	tctgagggtcc	3840
	aggaggaaaa	aggcatcaag	tataagtttg	aagtctacga	gaagaaagac	taacaggaag	3900
	atgctttcaa	gttctctgct	cccctcctaa	agctatgcat	ttttataaga	ccatgggact	3960
60	tttgctggct	ttagatcagc	ctcgactgtg	ccttctagtt	gccagccatc	tgttgtttgc	4020
	ccctcccccg	tgcccttctt	gaccctggaa	ggtgccactc	ccactgtcct	ttcctaataa	4080
	aatgaggaaa	ttgcatcgca	ttgtctgagt	aggtgtcatt	ctatttctggg	gggtgggggtg	4140
	gggcaggaca	gcaaggggga	ggattgggaa	gacaatagca	ggcatgctgg	ggatgcgggtg	4200
	ggctctatgg	aaccagctgg	ggctcgaagc	ggccgctccg	gatatgccaa	gtacgcccc	4260

	tattgacgtc	aatgacggta	aatggcccg	ctggcattat	gccagtaga	tgaccttatg	4320
	ggactttcct	acttggcagt	acatctacgt	attagtcac	gctattacca	tggtgatgcg	4380
	gttttggcag	tacatcaatg	ggcgtggata	gcggtttgac	tcacggggat	ttccaagtct	4440
	ccacccatt	gacgtcaatg	ggagtttgtt	ttggcaccaa	aatcaacggg	actttccaaa	4500
5	atgtcgtaac	aactccgccc	cattgacgca	aatgggagg	aggcgtgtac	ggtgggaggt	4560
	ctatataagc	agagctgggt	acgtcctcac	attcagtgat	cagcactgaa	cacagacccg	4620
	tcgacatgga	gtttgggctg	agctggcttt	ttcttgtggc	tatttttaaaa	ggtgtccagt	4680
	gtgaagtgca	gctgggtggag	tctggggggag	gcttagtgaa	gcctggagg	tccttaagac	4740
	tctcctgtgc	agcctctgga	ttcactttca	gtagctatga	catgtcttgg	gttcgccagg	4800
10	ctccggggaa	ggggctggag	tgggtctcaa	ccattagtag	tggtggtagt	tacacctact	4860
	atctagacag	tataaagggc	cgattcacca	tctccagaga	caatgccaa	aactccctgt	4920
	acctgcaa	gaacagtctg	agggtgagg	acacggccgt	gtattactgt	gcaagacagg	4980
	ggttggacta	ctgggggtcga	ggaaccttag	tcaccgtctc	ctcagctagc	accaagggcc	5040
	catcgggtctt	ccccctggca	ccctcctcca	agagcacctc	tgggggcaca	gcggccctgg	5100
15	gctgcctgg	caaggactac	ttccccgaac	cggtgacgg	gtcgtggaac	tcaggcgccc	5160
	tgaccagcgg	cgtgcacacc	ttccccggctg	tcctacagtc	ctcaggactc	tactccctca	5220
	gcagcgtgg	gaccgtgccc	tccagcagct	tgggcaccca	gacctacatc	tgcaacgtga	5280
	atcacaagcc	cagcaacacc	aagggtggaca	agaaagttga	gcccacatct	tgtgacaaaa	5340
	ctcacacatg	cccaccgtgc	ccagcacctg	aactcctggg	gggaccgtca	gtcttcctct	5400
20	tccccccaaa	acccaaggac	accctcatga	tctcccggac	ccctgaggtc	acatgcgtgg	5460
	tggtggacgt	gagccacgaa	gaccctgagg	tcaagttcaa	ctggtacgtg	gacggcggtg	5520
	aggtgcataa	tgccaagaca	aagccgcggg	aggagcagta	caacagcacg	taccgtgtgg	5580
	tcagcgtcct	caccgtcctg	caccaggact	ggctgaatgg	caaggagtac	aagtgcagg	5640
	tctccaacaa	agccctccca	gcccccatcg	agaaaaccat	ctccaaagcc	aaagggcagc	5700
25	cccgagaacc	acaggtgtac	accctgcccc	catcccggga	tgagctgacc	aagaaccagg	5760
	tcagcctgac	ctgcctggtc	aaaggcttct	atcccagcga	catcgccgtg	gagtgggaga	5820
	gcaatgggca	gccgggagaac	aactacaaga	ccacgcctcc	cgtgctggac	tccgacggct	5880
	ccttcttcct	ctacagcaag	ctcaccgtgg	acaagagcag	gtggcagcag	gggaacgtct	5940
	tctcatgctc	cgtgatgcat	gaggctctgc	acaaccacta	cacgcagaag	agcctctccc	6000
30	tgtctccggg	taaatgagga	tccgttaacg	gttaccaact	acctagactg	gattcgtgac	6060
	aacatgcggc	cgtgatattct	acgtatgatc	agcctcgact	gtgccttcta	gttgccagcc	6120
	atctgttgtt	tgcccctccc	ccgtgccttc	cttgaccctg	gaagggtgcca	ctcccactgt	6180
	ccttttcctaa	taaaatgagg	aaattgcatc	gcattgtctg	agtaggtgtc	attctattct	6240
	gggggggtgg	gtggggcagg	acagcaagg	ggaggattgg	gaagacaata	gcaggcatgc	6300
35	tgggggatg	gtgggctcta	tgggaaccagc	tggggctcga	cagcgtgcg	atcgccctga	6360
	ggccgctact	aactctctcc	tccctccttt	ttcctgcagg	acgaggcagc	gcggctatcg	6420
	tggctggcca	cgacgggctg	tccttgcgca	gctgtgctcg	acgttgtcac	tgaagcggga	6480
	agggactggc	tgctattggg	cgaagtgccg	gggcaggatc	tcctgtcatc	tcaccttgct	6540
	cctgcccaga	aagtatccat	catggctgat	gcaatgcggc	ggctgcatac	gcttgatccg	6600
40	gctacctgcc	cattcgacca	ccaagcgaaa	catcgcatcg	agcgagcacg	tactcggatg	6660
	gaagccggtc	ttgtcgatca	ggatgatctg	gacgaagagc	atcaggggct	cgcgccagcc	6720
	gaactgttcg	ccaggctcaa	ggcgcgcag	cccagcggcg	aggatctcgt	cgtgacccat	6780
	ggcgatgcct	gcttgccgaa	tatcatgggtg	gaaaatggcc	gcttttctgg	attcatcgac	6840
	tgtggccggc	tgggtgtggc	ggaccgctat	caggacatag	cgttggctac	ccgtgatatt	6900
45	gctgaagagc	ttggcggcga	atgggctgac	cgcttcctcg	tgctttacgg	tatcgccgct	6960
	cccgattcgc	agcgcatcgc	cttctatcgc	cttcttgacg	agttcttctg	agcgggactc	7020
	tgggggttcga	aatgaccgac	caagcgacgc	ccaacctgcc	atcacgagat	ttcgattcca	7080
	ccgcgcctt	ctatgaaagg	ttgggcttcg	gaatcgtttt	ccgggacgcc	ggctggatga	7140
	tcctccagcg	cggggatctc	atgctggagt	tcttcgcccc	ccccaaacttg	tttattgcag	7200
50	cttataatgg	ttacaaataa	agcaatagca	tcacaaatth	cacaaataaa	gcattttttt	7260
	cactgcattc	tagttgtgg	ttgtccaaac	tcatcaatct	atcttatcat	gtctggatcg	7320
	cggccggccg	caccgcgggtg	gagctttta	taaggcgcgc	cagctccagc	ttttgttccc	7380
	tttagtgagg	gttaatttcg	agcttggcgt	aatcatgggtc	atagctgttt	cctgtgtgaa	7440
	attgttatcc	gtcacaatt	ccacacaaca	tacgagccgg	aagcataaag	tgtaaagcct	7500
55	gggggtgccta	atgagtgagc	taactcacat	taattgcgtt	gcgtcactg	cccgttttcc	7560
	agtcgggaaa	cctgtcgtgc	cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	7620
	gtttgcgtat	tgggcgctct	tcgccttcct	cgctcactga	ctcgtgcgcg	tcggtcgttc	7680
	ggctgcggcg	agcggtatca	gctcactcaa	aggcggta	acggttatcc	acagaatcag	7740
	gggataacgc	aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	7800
60	aggccgcgtt	gctggcggtt	ttccataggg	tccgcccccc	tgacgagcat	cacaaaaatc	7860
	gacgctcaag	tcagaggtgg	cgaaacccga	caggactata	aagataccag	gcgtttcccc	7920
	ctggaagctc	cctcgtgcgc	tctcctgttc	cgacctgcc	gcttaccgga	tacctgtccg	7980
	cctttctccc	ttcggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	8040
	cggtgtaggt	cgttcgctcc	aagctgggct	gtgtgcacga	acccccggtt	cagcccgacc	8100

```

gctgogcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc 8160
cactggcagc agccactggg aacaggatta gcagagcgag gtatgtaggc ggtgctacag 8220
agttcttgaa gtgggtggcct aactacggct acactagaag gacagtatct ggtatctgcg 8280
ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa 8340
5 ccaccgctgg tagcgggtgg ttttttgttt gcaagcagca gattacgcgc agaaaaaaag 8400
gatctcaaga agatcctttg atcttttcta cgggggtctga cgctcagtgg aacgaaaact 8460
cacgttaagg gatttttggtc atgagattat caaaaaggat cttcacctag atccttttta 8520
attaaaaatg aagtttttaa tcaatctaaa gtatatatga gtaaacttgg tctgacagtt 8580
accaatgctt aatcagttag gcacctatct cagcgatctg tctatttctg tcatccatag 8640
10 ttgcctgact ccccgctcgt tagataacta cgatacggga gggcttacca tctggcccca 8700
gtgctgcaat gataccgcga gaccacgct caccggctcc agatttatca gcaataaacc 8760
agccagccgg aagggccgag cgcagaagtg gtcctgcaac tttatccgcc tccatccagt 8820
ctattaattg ttgccgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg 8880
ttgttgccat tgctacaggc atcgtgggtg cacgctcgtc gtttggtatg gcttcattca 8940
15 gctccgggtt ccaacgatca aggcgagtta catgatcccc catgttgtgc aaaaaagcgg 9000
ttagctcctt cggtcctccg atcgttgtca gaagtaagt ggccgcagtg ttatcactca 9060
tggttatggc agcactgcat aattctctta ctgtcatgcc atccgtaaga tgcttttctg 9120
tgactgggtg gtactcaacc aagtcattct gagaatagt tatgcggcga ccgagttgct 9180
cttgcccggc gtcaatacgg gataataccg cgccacatag cagaacttta aaagtgctca 9240
20 tcattggaaa acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca 9300
gttcgatgta acccactcgt gcacccaact gatcttcagc atcttttact ttcaccagcg 9360
tttctgggtg agcaaaaaa ggaaggcaaa atgccgcaaa aaagggaata agggcgacac 9420
ggaaatgttg aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt 9480
attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacia ataggggttc 9540
25 cgcgcacatt tccccgaaaa gtgccaca 9568

```

<210> 17

<211> 6414

30 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: plasmid

35

<400> 17

```

tcgacattga ttattgacta gttattaata gtaatcaatt acgggggtcat tagttcatag 60
cccatatatg gagttccgcg ttacataact tacggtaaat ggcccgccctg gctgaccgcc 120
caacgacccc cgccattga cgtcaataat gacgtatgtt cccatagtaa cgccaatagg 180
40 gactttccat tgacgtcaat ggggtggagta tttacggtaa actgcccact tggcagtaca 240
tcaagtgtat catatgccaa gtacgccccc tattgacgtc aatgacggta aatggccgc 300
ctggcattat gccagtaga tgaccttatg ggactttcct acttggcagt acatctacgt 360
attagtcatt gctattacca tgggtgatgcg gttttggcag tacatcaatg ggcgtggata 420
gcggtttgac tcacggggat ttccaagtct ccacccatt gagtcaatg ggagtttgtt 480
45 ttggcaccaa aatcaacggg actttccaaa atgtcgtaac aactccgccc cattgacgca 540
aatgggcggg aggcgtgtac ggtgggaggt ctatataagc agagctctct ggctaactag 600
agaaccact gcttaactgg cttatcgaaa ttaatacgac tcactatagg gagaccaag 660
cttctgcagg tcgacatcga tggatccggt acctcgagcg cgaattctct agaggatctt 720
tgtgaaggaa ccttacttct gtggtgtgac ataattggac aaactaccta cagagattta 780
50 aagctctaag gtaaatataa aatttttaag tgtataatgt gttaaactac tgattctaata 840
tgtttggtga ttttagattc caacctatgg aactgatgaa tgggagcagt ggtggaatgc 900
ctttaatgag gaaaacctgt tttgctcaga agaatgcca tctagttagt atgaggctac 960
tgctgactct caacattcta ctctccaaa aaagaagaga aaggtagaag accccaagga 1020
ctttccttca gaattgctaa gttttttgag tcatgctgtg tttagtaata gaactcttgc 1080
55 ttgctttgct atttacacca caaaggaaaa agctgcactg ctatacaaga aaattatgga 1140
aaaatatatt atgtatagtg ccttgactag agatcataat cagccatacc acattttagt 1200
aggttttact tgcttttaaaa aacctccac acctccccc gaacctgaaa cataaaatga 1260
atgcaattgt tggtgttaac ttgtttattg cagcttataa tgggtacaaa taaagcaata 1320
gcatacaciaa tttcacaaat aaagcatttt tttcactgca ttctagttgt ggtttgtcca 1380
60 aactcatcaa tgtatcttat catgtctgga tcaattctga gaaactagcc ttaaagacag 1440
acagctttgt tctagttagc caggcaagca tatgtaaata aagttcctca gggaactgag 1500
gttaaaagat gtatcctgga cctgccagac ctggccattc acgtaaagag aagattccgc 1560
ctcaagttcc ggtaacaac aggaggcaac gagatctcaa atctattact tctaactcgg 1620
taattaaaac ctttcaacta aaacacggac ccacggatgt caccactttt tccttccccg 1680

```

	gctccgccct	tctcagtact	ccccaccatt	aggctcgcta	ctccacctcc	acttccgggc	1740
	gcgacaccca	cgtgccctct	cccacccgac	gctaaccccg	cccctgcccg	tctgaccccg	1800
	cccaccacct	ggccccgccc	cgttgaggac	agaagaaacc	ccgggcagcc	gcagccaagg	1860
	cggacgggta	gacgctgggg	gcgctgagga	gtcgtcctct	accttctctg	ctggctcggt	1920
5	ggggggacgcg	gtggatctca	ggcttccgga	agactggaag	aaccggctca	gaaccgcttg	1980
	tctccgcggg	gcttgggcgg	cggaagaatg	gccgctagac	gcggacttgg	tgcgaggcat	2040
	cgcaggatgc	agaagagcaa	gcccgcgggg	agcgcgcggc	tgtactaccc	cgcgcctgga	2100
	gcggccacgc	cggactgggc	ggggccggcc	tggtggaggc	ggagtctgac	ctcgtggagg	2160
	cggggcctct	gatgttcaaa	taggatgcta	ggcttgttga	ggcgtggcct	ccgattcaca	2220
10	agtgggaagc	agcgccgggc	gactgcaatt	tcgcgcctaa	cttgggggaa	gcacagcgta	2280
	caggctgcct	aggtgatcgc	tgctgctgtc	atggttcgac	cgctgaactg	catcgctcgc	2340
	gtgtcccaga	atatgggcat	cggcaagaac	ggagaccttc	cctggccaat	gctcaggtag	2400
	tggtctggatt	gggttaggga	aaccgaggcg	gttcgctgaa	tcgggtcgag	cacttggcgg	2460
	agacgcgcgg	gccaactact	tagggacagt	catgaggggt	aggcccgcgg	gctgctgccc	2520
15	ttgcccatgc	ccgcggtgat	ccccatgctg	tgccagcctt	tgccagagg	cgctctagct	2580
	gggagcaaag	tccggtcact	gggcagcacc	accccccgga	cttgcatggg	tagccgctga	2640
	gatggagcct	gagcacacgt	gacaggggtc	ctgttaacgc	agtgtttctc	taactttcag	2700
	gaacgagttc	aagtacttcc	aaagaatgac	caccacctcc	tcagtgggaag	gtaaacagaa	2760
	cctggtgatt	atgggcccga	aaacctgggt	ctccattcct	gagaagaatc	gacctttaaa	2820
20	ggacagaatt	aatatagttc	tcagtagaga	gctcaaggaa	ccaccacaag	gagctcattt	2880
	tcttgccaaa	agtctggacc	atgccttaaa	acttattgaa	caaccagagt	tagcagataa	2940
	agtggacatg	gtttggatag	ttggaggcag	ttccgtttac	aagggaagcca	tgaatcagcc	3000
	aggccatctc	agactctttg	tgacaaggat	catgcaggaa	tttgaaagtg	acacgttctt	3060
	cccagaaatt	gattttggaga	aatataaact	tctcccagag	taccagggg	tcctttctga	3120
25	agtccaggag	gaaaaaggca	tcaagtataa	atltgaagtc	tatgagaaga	aaggctaaca	3180
	gaaagatact	tgctgattga	cttcaagttc	tactgctttc	ctcctaaaat	tatgcatttt	3240
	tacaagacca	tgggacttgt	gttggcttta	gatcctgtgc	atcctgggca	actgttgtac	3300
	tctaagccac	tccccaaagt	catgccccag	cccctgtata	attctaaaca	attagaatta	3360
	ttttcatttt	cattagtcta	accaggttat	attaaatata	ctttaagaaa	caccatttgc	3420
30	cataaagttc	tcaatgcccc	tcccatgcag	cctcaagtgg	ctccccagca	gatgcatagg	3480
	gtagtgtgtg	tacaagagac	cccaaagaca	tagagcccct	gagagcatga	gctgatatgg	3540
	gggctcatag	agataggagc	tagatgaata	agtacaaagg	gcagaaatgg	gttttaacca	3600
	gcagagctag	aactcagact	ttaaagaaaa	ttagatcaaa	gtagagactg	aattattctg	3660
	cacatcagac	tctgagcaga	gttctgttca	ctcagacaga	aaatgggtaa	attgagagct	3720
35	ggctccattg	tgctccttag	agatgggagc	aggtaggagga	ttatataagg	tctggaacat	3780
	ttaaacttctc	cgttttctcat	cttcagttag	attccaaggg	atactacaat	tctgtggaat	3840
	gtgtgtcagt	taggggtgtg	aaagtcccca	ggctccccag	caggcagaag	tatgcaaagc	3900
	atgcatctca	attagtcagc	aaccagggtg	ggaaagtccc	caggctcccc	agcaggcaga	3960
	agtatgcaaa	gcatgcatct	caattagtc	gcaaccatag	tcccgcctct	aactccgccc	4020
40	atcccgcctc	taactccgcc	cagttccgcc	cattctccgc	cccatggctg	actaattttt	4080
	tttatattatg	cagaggccga	ggcgccctctg	agctattcca	gaagtagtga	ggaggctttt	4140
	ttggaggcct	aggctttttgc	aaaaaagcta	attcagcctg	aatggcggaat	gggacgcgcc	4200
	ctgtagcggc	gcattaaagc	cggcgggtgt	ggtggttacg	cgcagcgtga	ccgctacact	4260
	tgccagcgcc	ctagcgcccg	ctcctttcgc	tttcttccct	tcctttctcg	ccacgttcgc	4320
45	cggctttccc	cgtcaagctc	taaatcgggg	gctcccttta	gggttccgat	ttagtgcctt	4380
	acggcacctc	gaccccaaaa	acttgattag	ggtgatggtt	cacgtagtgg	gccatcgccc	4440
	tgatagacgg	tttttcgccc	tttgacgttg	gagtcacagt	tcctttaatag	tggactcttg	4500
	ttccaaactg	gaacaacact	caaccctatc	tcgggtctatt	cttttgattt	ataagggatt	4560
	ttgccgattt	cggcctattg	gttaaaaaat	gagctgattt	aacaaaaatt	taacgcgaat	4620
50	tttaacaaaa	tattaacgtt	tacaatttca	ggtggcactt	ttcgggggaa	tgtgcgcgga	4680
	accctatatt	gtttattttt	ctaaatacat	tcaaatatgt	atccgctcat	gagacaataa	4740
	ccctgataaa	tgcttcaata	atattgaaaa	aggaagagta	tgagtattca	acatttccgt	4800
	gtcgccctta	ttcccttttt	tgccgcattt	tgccctcctg	tttttgctca	cccagaaacg	4860
	ctgggtgaaag	taaaagatgc	tgaagatcag	ttgggtgcac	gagtgggtta	catcgaactg	4920
55	gatctcaaca	gcggttaagat	ccttgagagt	tttcgccccg	aagaacgttt	tccaatgatg	4980
	agcactttta	aagttctgct	atgtggcgcg	gtattatccc	gtattgacgc	cgggcaagag	5040
	caactcggtc	gccgcataca	ctattctcag	aatgacttgg	ttgagtactc	accagtcaca	5100
	gaaaagcatc	ttacggatgg	catgacagta	agagaattat	gcagtgcctg	cataaccatg	5160
	agtataaaca	ctgcggccaa	cttacttctg	acaacgatcg	gaggaccgaa	ggagctaacc	5220
60	gctttttttgc	acaacatggg	ggatcatgta	actcgccctg	atcgttggga	accggagctg	5280
	aatgaagcca	taccaaacga	cgagcgtgac	accacgatgc	ctgtagcaat	ggcaacaacg	5340
	ttgcgcaaac	tattaactgg	cgaactactt	actctagctt	cccggcaaca	attaatagac	5400
	tggatggagg	cggataaagt	tgcaggacca	cttctgcgct	cggcccttcc	ggctggctgg	5460
	tttatgtctg	ataaatctgg	agccgggtgag	cgtgggtctc	gcggtatcat	tgcagcactg	5520

```

5   gggccagatg gtaagccctc ccgtatcgta gttatctaca cgacggggag tcaggcaact 5580
    atggatgaac gaaatagaca gatcgctgag ataggtgcct cactgattaa gcattggtaa 5640
    ctgtcagacc aagtttactc atatatactt tagattgatt taaaacttca tttttaattt 5700
    aaaaggatct aggtgaagat cctttttgat aatctcatga ccaaaatccc ttaacgtgag 5760
10  ttttcgttcc actgagcgtc agaccccgtc gaaaagatca aaggatcttc ttgagatcct 5820
    ttttttctgc gcgtaatctg ctgcttgcaa acaaaaaaac caccgctacc agcgggtggtt 5880
    tgtttgccgg atcaagagct accaactctt tttccgaagg taactggctt cagcagagcg 5940
    cagataccaa atactgtcct tctagtgtag ccgtagttag gccaccactt caagaactct 6000
    gtagcaccgc ctacatacct cgctctgcta atcctgttac cagtggctgc tgccagtggc 6060
15  gataagtcgt gtcttaccgg gttggactca agacgatagt taccggataa ggcgcagcgg 6120
    tcgggctgaa cgggggggttc gtgcacacag cccagcttgg agcgaacgac ctacaccgaa 6180
    ctgagatacc tacagcgtga gcattgagaa agcggccacgc ttcccggaagg gagaaaggcg 6240
    gacaggtatc cggtaagcgg cagggtcggg acaggagagc gcacgaggga gcttccaggg 6300
    ggaaacgcct ggtatcttta tagtcctgtc gggtttcgcc acctctgact tgagcgtcga 6360
20  tttttgtgat gctcgtcagg ggggaggagc ctatggaaaa acgccagcaa cgcc 6414

```

<210> 18

<211> 6062

20 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: plasmid

25

<400> 18

```

    tcgacattga ttattgacta gttattaata gtaatcaatt acgggggtcat tagttcatag 60
    cccatataat gagttccgcg ttacataact tacggtaaata ggcccgcctg gctgaccgcc 120
    caacgacccc cgccattga cgtcaataat gacgtatgtt cccatagtaa cgccaatagg 180
30  gactttccat tgacgtcaat ggggtggagta ttacggtaa actgccact tggcagtaca 240
    tcaagtgtat catatgccaa gtacgcccc tattgacgtc aatgacggta aatggccgcg 300
    ctggcattat gccagtgaca tgacctatg ggactttcct acttggcagt acatctacgt 360
    attagtcata gctattacca tgggtgatgcg gttttggcag tacatcaatg ggcgtggata 420
    gcggtttgac tcacggggat ttccaagtct ccaccccatg gacgtcaatg ggagtttgtt 480
35  ttggcaccaa aatcaacggg actttccaaa atgtcgtaac aactccgcc cattgacgca 540
    aatgggagggt aggcgtgtac ggtgggagggt ctatataagc agagctcgtt tagtgaaccg 600
    tcagatcgcc tggagacgcc atccacgctg ttttgacctc catagaagac accgggaccg 660
    atccagcctc cgcggccggg aacgggtgcat tggaaacgcg attccccgtg ccaagagtca 720
    ggtaagtacc gcctatagag aagactcttg ggtttctgat aggactgac tctctctgcc 780
40  tattgggtcta ttttcccacc cttaggctgc tgggtgctta ctggcttata gaaattaata 840
    cgactcata tagggagacc caagcttctg caggtcgaca tcgatggatc cgggtacctc 900
    agcgcgaatt ctctagagat atcttgttta ttgcagctta taatgggttac aaataaagca 960
    atagcatcac aaatttcaca aataaagcat ttttttctact gcattctagt tgtggtttgt 1020
    ccaaactcat caatgtatct tatcatgtct ggatcaattc tgaaaaacta gccttaaaga 1080
45  cagacagctt tgttctagtc agccaggcaa gcatatgtaa ataaagtacc tcagggaact 1140
    gaggttaaaa gatgtatcct ggacctgcca gacctggcca ttcacgtaaa cagaagattc 1200
    cgcctcaagt tccggttaac aacaggaggc aacgagatct caaatctatt acttctaata 1260
    gggtaattaa aacctttcaa ctaaaacacg gaccacgga tgtcaccac ttttccttcc 1320
    cgggctccgc ctttctcagt actccccacc attaggctcg ctactccacc tccacttccg 1380
50  ggcgcgacac ccacgtgcc tctcccaccc gacgctaacc ccgccctgc ccgtctgacc 1440
    ccgccacca cctggccccg cccggttgag gacagaagaa accccgggca gccgcagcca 1500
    aggcggacgg gtagacgctg ggggcgctga ggagtcgtcc tctaccttct ctgctggctc 1560
    ggtgggggac gcggtggatc tcaggcttcc ggaagactgg aagaaccggc tcagaaccgc 1620
    ttgtctccgc ggggcttggg cggcggaaga atggccgcta gacgcggact tgggtgcgag 1680
55  catcgcagga tgcagaagag caagcccgcg gggagcgcgc ggctgtacta cccgcgcct 1740
    ggagcggcca cgcgggactg ggcggggccg gcctgggtga ggcggagtct gacctcgtgg 1800
    aggcggggcc tctgatgttc aaataggatg ctaggcttgt tgaggcgtgg cctccgattc 1860
    acaagtggga agcagcgcgc ggcgactgca atttcgcgcc aaacttgggg gaagcacagc 1920
    gtacaggctg cctaggtgat cgctgctgct gtcattggtt gaccgctgaa ctgcatcgtc 1980
60  gccgtgtccc agaatatggg catcggcaag aacggagacc ttccctggcc aatgctcagg 2040
    tactggctgg attgggttag ggaaaccgag gcggttcgct gaatcgggtc gagcacttgg 2100
    cggagacgcg cgggccaaact acttagggac agtcatgagg ggtaggcccg ccggctgctg 2160
    cccttgccca tgcccgcggt gatcccatg ctgtgccagc ctttgcccag aggcgtcta 2220
    gctgggagca aagtccggtc actgggcagc accaccccc ggacttgcatt gggtagccgc 2280

```

	tgagatggag	cctgagcaca	cgtgacaggg	tccctgttaa	cgcagtgttt	ctctaacttt	2340
	caggaacgag	ttcaagtact	tccaaagaat	gaccaccacc	tcctcagtgg	aaggtaaaca	2400
	gaacctggtg	attatgggcc	ggaaaacctg	gttctccatt	cctgagaaga	atcgaccttt	2460
	aaaggacaga	attaatatag	ttctcagtag	agagctcaag	gaaccaccac	aaggagctca	2520
5	ttttcttgcc	aaaagtctgg	accatgcctt	aaaacttatt	gaacaaccag	agtttagcaga	2580
	taaagtggac	atggtttgga	tagttggagg	cagttccggt	tacaaggaag	ccatgaatca	2640
	gccaggccat	ctcagactct	ttgtgacaag	gatcatgcag	gaatttgaaa	gtgacacggt	2700
	cttcccagaa	attgatttgg	agaaatataa	acttctccca	gagtaccacg	gggtcctttc	2760
	tgaagtccag	gaggaaaaag	gcatcaagta	taaatttgaa	gtctatgaga	agaaaggcta	2820
10	acagaaagat	acttgctgat	tgacttcaag	ttctactgct	ttcctcctaa	aattatgcat	2880
	ttttacaaga	ccatgggact	tgtgttggt	ttagatcctg	tgcattcctg	gcaactgttg	2940
	tactctaagc	cactcccca	agtcattgccc	cagccctgt	ataattctaa	acaattagaa	3000
	ttattttcat	tttcattagt	ctaaccagg	tataattaa	atactttaag	aaacaccatt	3060
	tgccataaag	ttctcaatgc	ccctcccatg	cagcctcaag	tggtcccca	gcagatgcat	3120
15	agggtagtgt	gtgtacaaga	gaccccaaag	acatagagcc	cctgagagca	tgagctgata	3180
	tgggggctca	tagagatagg	agctagatga	ataagtacaa	agggcagaaa	tgggttttaa	3240
	ccagcagagc	tagaactcag	actttaaa	aaattagatc	aaagtagaga	ctgaattatt	3300
	ctgcacatca	gactctgagc	agagtctgt	tcactcagac	agaaaatggg	taaattgaga	3360
	gctggctcca	ttgtgctcct	tagagatggg	agcagggtga	ggattatata	aggtctggaa	3420
20	catttaactt	ctccgtttct	catcttcagt	gagattccaa	gggatactac	aattctgtgg	3480
	aatgtgtgtc	agttaggggt	tggaaagtcc	ccagggtccc	cagcaggcag	aagtatgcaa	3540
	agcatgcac	tcaattagtc	agcaaccagg	tgtggaaagt	ccccaggctc	cccagcaggc	3600
	agaagtatgc	aaagcatgca	tctcaattag	tcagcaacca	tagtcccgc	cctaactccg	3660
	cccattccgc	ccctaactcc	gccagttcc	gccattctc	cgcccatgg	ctgactaatt	3720
25	ttttttattt	atgcagaggc	cgaggcgct	ctgagctatt	ccagaagtag	tgaggaggct	3780
	tttttgagg	cctaggcttt	tgcaaaaaag	ctaattcagc	ctgaatggcg	aatgggaaat	3840
	tgtaaacgtt	aatattttgt	taaaattcgc	gttaaatttt	tgttaaataca	gctcattttt	3900
	taaccaatag	gccgaaatcg	gcaaaatccc	ttataaatca	aaagaataga	ccgagatagg	3960
	gttgagtgtt	gttccagttt	ggaacaagag	tccactatta	aagaacgtgg	actccaacgt	4020
30	caaagggcga	aaaaccgtct	atcagggcga	tggcccacta	cgtgaaccat	caccctaatt	4080
	aagtttttgg	ggtcagagg	ccgtaaagca	ctaaatcgga	accctaagag	gagccccga	4140
	tttagagctt	gacggggaaa	gccggcgaa	gtggcgagaa	aggaagggaa	gaaagcgaaa	4200
	ggagcggg	ctagggcgct	ggcaagtgt	gcggtcacgc	tgcgcgtaac	caccacaccc	4260
	gccgcgctta	atgcgcgct	acagggcgcg	tcagggtggc	cttttcgggg	aatgtgctgc	4320
35	ggaaccctta	tttggtttatt	tttctaaata	cattcaaata	tgtatccgct	catgagacaa	4380
	taaccctgat	aaatgcttca	ataatattga	aaaaggaaga	gtatgagtat	tcaacatttc	4440
	cgtgtcgccc	ttattccctt	ttttgcggca	ttttgccttc	ctgtttttgc	tcaccacagaa	4500
	acgctggtga	aagtaaaaga	tgtctgaagat	cagttgggtg	cacgagtggg	ttacatcgaa	4560
	ctggatctca	acagcggtaa	gatccttgag	agttttcgcc	ccgaagaacg	ttttccaatg	4620
40	atgagcactt	ttaaagtctt	gctatgtggc	gcggtattat	cccgtattga	cgccgggcaa	4680
	gagcaactcg	gtcgcgcgat	acactattct	cagaatgact	tggttgagta	ctcaccagtc	4740
	acagaaaagc	atcttacgga	tggcatgaca	gtaagagaat	tatgcagtgc	tgccataacc	4800
	atgagtgata	acactgcggc	caacttactt	ctgacaacga	tcggaggacc	gaaggagcta	4860
	accgcttttt	tgcacaacat	gggggatcat	gtaactcgcc	ttgatcggtg	ggaaccggag	4920
45	ctgaatgaag	ccataccaaa	cgacgagcgt	gacaccacga	tgctgtagc	aatggcaaca	4980
	acgttgcgca	aactattaac	tggcgaacta	cttactctag	cttcccggca	acaattaata	5040
	gactggatgg	aggcggataa	agttgcagga	ccacttctgc	gctcggccct	tcgggctggc	5100
	tgggtttattg	ctgataaatc	tggagccggt	gagcgtgggt	ctcgcggtat	cattgcagca	5160
	ctggggccag	atggtaagcc	ctcccgatat	gtagttatct	acacgacggg	gagtcaggca	5220
50	actatggatg	aacgaaatag	acagatcgct	gagataggtg	cctcactgat	taagcattgg	5280
	taactgtcag	accaagttta	ctcatatata	cttttagattg	atttaaaaact	tcatTTTTTaa	5340
	tttaaaagga	tctaggtgaa	gatccttttt	gataatctca	tgaccaaaa	cccttaacgt	5400
	gagttttcgt	tccactgagc	gtcagacccc	gtagaaaaga	tcaaaggatc	ttcttgagat	5460
	ccttttttttc	tgcgcgtaat	ctgctgcttg	caaacaaaaa	aaccaccgct	accagcgggtg	5520
55	gtttgtttgc	cggatcaaga	gctaccaact	ctttttccga	aggtaactgg	cttcagcaga	5580
	gcgcagatac	caaatactgt	ccttctagt	tagccgtagt	taggccacca	cttcaagaac	5640
	tctgtagcac	cgcctacata	cctcgctctg	ctaactcctg	taccagtggc	tgctgccagt	5700
	ggcgataagt	cgtgtcttac	cgggttgagc	tcaagacgat	agttaccgga	taaggcgag	5760
	cggtcgggct	gaacgggggg	ttcgtgcaca	cagcccagct	tggagcgaac	gacctacacc	5820
60	gaactgagat	acctacagcg	tgagcattga	gaaagcgcca	cgcttcccga	aggagaaaag	5880
	gcggacaggt	atccggtaag	cggcagggtc	ggaacaggag	agcgcacgag	ggagcttcca	5940
	gggggaaacg	cctggtatct	ttatagtcct	gtcgggtttc	gccacctctg	acttgagcgt	6000
	cgatttttgt	gatgctcgtc	aggggggagg	agcctatgga	aaaacgccag	caacgcagct	6060
	gc						6062

claims

1. Antibody molecule comprising a variable region of the heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 1 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof.
- 5 2. Antibody molecule wherein the variable region heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 1.
3. Antibody molecule comprising a variable region light chain as characterized by the amino acid sequence as defined in SEQ ID No. 2 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof.
- 10 4. Antibody molecule wherein the variable region light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 2.
5. Antibody molecule comprising a variable region light chain as characterized by the amino acid sequence as defined in SEQ ID No. 3 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof.
- 15 6. Antibody molecule wherein the variable region light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 3.
7. Antibody molecule according to claim 1 and 3 comprising a variable region heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 1 and comprising a variable region light chain as characterized by the amino acid sequence as defined in SEQ ID
20 No. 2 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof.
8. Antibody molecule according to claim 2 and 4 wherein the variable region heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 1 and wherein the variable region light chain consists of the amino acids as characterized by the
25 amino acid sequence of SEQ ID No. 2.
9. Antibody molecule according to claim 1 and 5 comprising a variable region heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 1 and comprising a variable region light chain as characterized by the amino acid sequence as defined in SEQ ID No. 3 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule
30 or a chemical derivative thereof.
10. Antibody molecule according to claim 2 and 6 wherein the variable region heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 1 and

wherein the variable region light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 3.

11. Antibody molecule comprising a variable region heavy chain encoded by the nucleic acid sequence as defined in SEQ ID No. 4 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
12. Antibody molecule wherein the variable region heavy chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 4.
13. Antibody molecule comprising a variable region light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 5 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
14. Antibody molecule wherein the variable region light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 5.
15. Antibody molecule comprising a variable region light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 6 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
16. Antibody molecule wherein the variable region light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 6.
17. Antibody molecule according to claim 11 and 13 comprising a variable region heavy chain encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and comprising a variable region light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 5 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
18. Antibody molecule according to claim 12 and 14 wherein the variable region heavy chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and wherein the variable region light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 5.
19. Antibody molecule according to claim 11 and 15 comprising a variable region heavy chain encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and comprising a variable region light chain encoded by the nucleic acid sequence as defined in SEQ ID No. 6 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
20. Antibody molecule according to claim 12 and 16 wherein the variable region heavy chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 4 and wherein the variable region light chain is encoded by the nucleic acid sequence as defined in SEQ ID No. 6.

21. An antibody molecule according to any one of claims 1 to 20 characterised in that each of said variable light chain and said variable heavy chain region is separately joined to a human constant region.
22. The antibody molecule of claim 21, wherein said human constant region of the light chain is a human kappa constant region.
23. The antibody molecule of claim 21 or 22, wherein said human constant region of the heavy chain is a human IgG1 constant region.
24. Antibody molecule according to any one of claims 1 to 23 comprising a heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 7 and comprising a light chain as characterized by the amino acid sequence as defined in SEQ ID No. 8 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof.
25. The antibody molecule according to any one of claims 1 to 24 wherein the heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 7 and wherein the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 8.
26. The antibody molecule according to any one of claims 1 to 25 comprising a heavy chain as characterized by the amino acid sequence as defined in SEQ ID No. 7 and comprising a light chain as characterized by the amino acid sequence as defined in SEQ ID No. 9 or a fragment, allelic variant, functional variant, glycosylation variant, fusion molecule or a chemical derivative thereof.
27. The antibody molecule according to any one of claims 1 to 26 wherein the heavy chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 7 and wherein the light chain consists of the amino acids as characterized by the amino acid sequence of SEQ ID No. 9.
28. The antibody molecule according to any one of claims 1 to 27 comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 10 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID No. 11 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
29. The antibody molecule according to any one of claims 1 to 28 wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 10 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 11.

30. The antibody molecule according to any one of claims 1 to 29 comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 10 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID No. 12 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
31. The antibody molecule according to any one of claims 1 to 30 wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 10 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 12.
32. The antibody molecule according to any one of claims 1 to 31 comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 13 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID No. 14 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
33. The antibody molecule according to any one of claims 1 to 32 wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 13 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 14.
34. The antibody molecule according to any one of claims 1 to 33 comprising a heavy chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 13 and comprising a light chain as characterized by the nucleic acid sequence as defined in SEQ ID No. 15 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
35. The antibody molecule according to any one of claims 1 to 34 wherein the heavy chain is encoded by the nucleic acid sequence of SEQ ID No. 13 and wherein the light chain is encoded by the nucleic acid sequence of SEQ ID No. 15.
36. The antibody molecule according to any one of claims 1 to 35 comprising a heavy and light chain as encoded by the nucleic acid sequence as defined in SEQ ID No. 16 or a fragment, allelic variant, functional variant, variant based on the degenerative nucleic acid code, fusion molecule or a chemical derivative thereof.
37. The antibody molecule according to any one of claims 1 to 36 wherein the heavy and light chain is encoded by the nucleic acid sequence of SEQ ID No. 16.
38. An antibody protein according to any one of claims 1 to 37, wherein said antibody protein is conjugated to a therapeutic agent.

39. The antibody protein of claim 38, wherein said therapeutic agent is a therapeutic agent selected from the group consisting of radioisotopes, toxins, toxoids, inflammatory agents and chemotherapeutic agents.
40. The antibody protein of claim 39, wherein said therapeutic agent is linked to the antibody protein via a linker selected from the group of MAG-3 GABA, MAG-2 GABA and N2S2.
41. The antibody protein of claim 40, wherein said therapeutic agent is linked to the antibody protein via MAG-2 GABA.
42. The antibody protein of any one of claims 38 to 41, wherein said radioisotope is a β -emitting radioisotope.
43. The antibody protein of claim 42, wherein said radioisotope is selected from the group consisting of $^{186}\text{Rhenium}$, $^{188}\text{Rhenium}$, $^{131}\text{Iodine}$ and $^{90}\text{Yttrium}$.
44. The antibody protein of claim 43, wherein said radioisotope is $^{186}\text{Rhenium}$.
45. The antibody protein according to any one of claims 39 to 44, wherein the antibody protein has specific activity of from about 0.5 to about 15 mCi/mg, or from about 0.5 to about 14 mCi/mg, preferably about 1 to about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.
46. The antibody protein according to any one of claims 1 to 37, characterised in that it is labelled.
47. The antibody protein of claim 46, wherein said label is a detectable marker.
48. The antibody protein of claim 47, wherein the detectable marker is a detectable marker selected from the group consisting of enzymes, dyes, radioisotopes, digoxigenin, and biotin.
49. An antibody protein according to any one of claims 1 to 37 conjugated to an imageable agent.
50. The antibody protein of claim 49, wherein the imageable agent is a radioisotope.
51. The antibody protein of claim 50, wherein said radioisotope is a γ -emitting radioisotope.
52. The antibody protein of claim 51, wherein said radioisotope is ^{125}I .
53. A pharmaceutical composition containing an antibody protein according to any one of claims 1 to 45 and a pharmaceutically acceptable carrier or excipient.
54. Pharmaceutical composition according to claim 53, wherein the antibody protein is conjugated to a radioisotope according to any one of claims 39 to 44 and has specific activity of from about 0.5 to about 15 mCi/mg, or from about 0.5 to about 14 mCi/mg, preferably about 1 to about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.

55. Pharmaceutical composition according to claim 53 or 54, wherein the amount of radiolabelled antibody in the pharmaceutical composition to be applied to a patient is 10, 20, 30, 40, 50 or 60 mCi/m².
56. Pharmaceutical composition according to any one of claims 53 to 55, wherein the amount of radiolabelled antibody in the pharmaceutical composition to be applied to a patient is 50 mCi/m².
57. Pharmaceutical composition according to any one of claims 53 to 56, further comprising one or more radioprotectant selected from the group of ascorbic acid, gentisic acid, reductic acid, erythrorbic acid, p-aminobenzoic acid, 4-hydroxybenzoic acid, nicotinic acid, nicotinamide, 2-5-dihydroxy-1,4-benzenedisulfonic acid, povidone, inositol, and/or citrate.
58. Pharmaceutical composition according to any one of claims 53 to 57, wherein the radioprotectant is ascorbic acid.
59. Pharmaceutical composition according to any one of claims 53 to 58, wherein said antibody protein comprises an antibody molecule selected from the group of antibody molecules of claims 8, 10, 18, 20, 24, 25, 28, 29, 32, 33, 36, 37 linked to ¹⁸⁶Rhenium via MAG-2 GABA further comprising the radioprotectant ascorbic acid.
60. Use of an antibody protein according to any one of claims 1 to 45 in the manufacture of a medicament for treatment of cancer.
61. Use of an antibody protein according to claim 60 wherein said cancer is selected from the group consisting of colorectal cancer, non-small cell lung cancer, breast cancer, head and neck cancer, ovarian cancer, lung cancer, bladder cancer, pancreatic cancer and metastatic cancers of the brain.
62. Use of an antibody protein according to any one of claims 60 or 61, wherein the antibody protein is conjugated to a radioisotope according to any one of claims 39 to 44 and has specific activity of from about 0.5 to about 15 mCi/mg, or from about 0.5 to about 14 mCi/mg, preferably about 1 to about 10 mCi/mg, preferably about 1 to about 5 mCi/mg, and most preferably 2 to 6 mCi/mg or 1 to 3 mCi/mg.
63. Use of an antibody protein according to claim 62, wherein the amount of radiolabelled antibody in the pharmaceutical composition to be applied to a patient is 10, 20, 30, 40, 50 or 60 mCi/m².
64. Use of an antibody protein according to claim 62, wherein the amount of radiolabelled antibody in the pharmaceutical composition to be applied to a patient is 50 mCi/m².

65. Method of cancer treatment, wherein an antibody protein according to any one of claims 1 to 45 is administered once to several times to an individual in need thereof, said antibody protein selectively binds to CD44v6, destroys tumor cells via the therapeutic agent linked to the antibody protein and the therapeutic success is monitored.
- 5 66. The method of claim 65, wherein the tumor is a tumor selected from the cancer group consisting of colorectal cancers, non-small cell lung cancers, breast cancers, head and neck cancer, ovarian cancers, lung cancers, bladder cancers, pancreatic cancers and metastatic cancers of the brain.
67. Nucleic acid, characterised in that it codes for an antibody protein according to one of claims 10 1 to 37.
68. Recombinant DNA vector, characterised in that it contains a nucleic acid according to claim 67.
69. Recombinant DNA vector according to claim 68, characterized in that it is an expression vector.
- 15 70. Recombinant DNA vector according to claim 68 or 69, characterized in that it is vector pAD-CMV or a functional derivative thereof.
71. Recombinant DNA vector according to any one of claims 68 to 70, characterized in that it is vector N5KG1Val or a derivative thereof.
72. Host, characterised in that it contains a vector according to any one of claims 68 to 71.
- 20 73. Host according to claim 72, characterised in that it is a eukaryotic host cell.
74. Host according to claim 72 or 73, characterised in that it is a mammalian cell.
75. Host according to any one of claims 72 to 74, characterised in that it is a BHK, CHO or COS cell.
76. Host according to claim 72, characterised in that it is a bacteriophage.
- 25 77. Host according to claim 72, characterised in that it is a prokaryotic host cell.
78. Process for preparing an antibody protein according to any one of claims 1 to 37, characterized in that it comprises the following steps: a host according to one of claims 72 to 77 is cultivated under conditions in which said antibody protein is expressed by said host cell and said antibody protein is isolated.
- 30 79. Process according to claim 78, characterised in that said host is a mammalian cell, preferably a CHO or COS cell.
80. Process according to claim 78 or 79, characterised in that said host cell is co-transfected with two plasmids which carry the expression units for the light or the heavy chain.




Fig. 1




Fig. 2




Fig. 3

1170.1 Part A Cmax Proportionality Assessment





Fig. 4

1170.1 Part A AUC Proportionality Assessment

**Fig. 5**