



US007425444B2

(12) **United States Patent**
Jestin et al.

(10) **Patent No.:** **US 7,425,444 B2**
(45) **Date of Patent:** **Sep. 16, 2008**

- (54) **CIRCOVIRUS SEQUENCES ASSOCIATED WITH PIGLET WEIGHT LOSS DISEASE (PWD)**
- (75) Inventors: **André Jestin**, Saint-Brieuc (FR); **Emmanuel Albina**, Trégueux (FR); **Pierre Le Cann**, Plédran (FR); **Philippe Blanchard**, Plérin (FR); **Evelyne Hutet**, Plérin (FR); **Claire Arnauld**, Saint-Brieuc (FR); **Catherine Truong**, Saint-Brieuc (FR); **Dominique Mahe**, Saint-Carreuc (FR); **Roland Cariolet**, Ploufragan (FR); **François Madec**, Saint-Brieuc (FR)

(73) Assignee: **Wyeth**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 4 days.

(21) Appl. No.: **11/588,306**

(22) Filed: **Oct. 27, 2006**

(65) **Prior Publication Data**

US 2007/0041990 A1 Feb. 22, 2007

Related U.S. Application Data

- (60) Division of application No. 10/718,264, filed on Nov. 21, 2003, now Pat. No. 7,179,472, which is a division of application No. 09/514,245, filed on Feb. 28, 2000, now Pat. No. 6,703,023, which is a continuation-in-part of application No. PCT/FR98/02634, filed on Dec. 4, 1998.

(30) **Foreign Application Priority Data**

Dec. 5, 1997 (FR) 97 15396

(51) **Int. Cl.**
C12N 15/00 (2006.01)

(52) **U.S. Cl.** **435/320.1**; 435/235.1; 435/325

(58) **Field of Classification Search** 435/320.1, 435/235.1, 325; 514/44

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,946,787 A 8/1990 Eppstein et al.
5,106,733 A 4/1992 Baker et al.
5,382,425 A 1/1995 Cochran et al.
5,459,127 A 10/1995 Felgner et al.
5,498,413 A 3/1996 Casal Alvarez et al.
5,545,412 A 8/1996 Eppstein et al.
5,580,859 A 12/1996 Felgner et al.
5,587,164 A 12/1996 Sanderson et al.
5,703,055 A 12/1997 Felgner et al.
5,705,385 A 1/1998 Bally et al.
5,719,131 A 2/1998 Harris et al.
5,756,103 A 5/1998 Paoletti et al.
5,770,212 A 6/1998 Falkner et al.
5,795,872 A 8/1998 Ricigliano et al.
5,811,103 A 9/1998 Meyers et al.

5,820,869 A 10/1998 Wasmoen et al.
5,833,975 A 11/1998 Paoletti et al.
5,888,513 A 3/1999 Plana Duran et al.
5,910,488 A 6/1999 Nabel et al.
5,990,091 A 11/1999 Tartaglia et al.
6,004,777 A 12/1999 Tartaglia et al.
6,015,694 A 1/2000 Dubensky, Jr. et al.
6,019,980 A 2/2000 Li et al.
6,033,904 A 3/2000 Cochran et al.
6,143,333 A 11/2000 Gordon et al.
6,143,334 A 11/2000 Reinbold et al.
6,143,734 A 11/2000 Garvey et al.
6,165,493 A 12/2000 Neurath et al.
6,207,165 B1 3/2001 Audonnet et al.
6,217,883 B1 4/2001 Allan et al.
6,287,856 B1 9/2001 Poet et al.
6,368,601 B1 4/2002 Allan
6,391,314 B1 5/2002 Allan et al.
6,475,779 B2 11/2002 Mathiowitz et al.
6,497,883 B1 12/2002 Bublot et al.
6,517,843 B1 2/2003 Ellis et al.
6,573,081 B2 6/2003 Bernhardt et al.
6,660,272 B2 12/2003 Allan et al.
6,703,023 B1 3/2004 Jestin et al.
6,794,163 B2 9/2004 Liu et al.

(Continued)

FOREIGN PATENT DOCUMENTS

DE	10/044648	3/2002
EP	0 737 750	10/1996
FR	2 422 956	11/1979
FR	2518 755	6/1983
FR	2769322	4/1999

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 60/069,750, filed Dec. 16, 1997, Wang et al.
U.S. Appl. No. 60/069,233, filed Dec. 11, 1997, Wang et al.
Allan, G. M. et al., "Isolation of Procine Circovirus-like Viruses from Pigs with a Wasting Disease in the USA and Europe", *Journal of Veterinary Diagnostic Investigation*, vol. 10, pp. 3-10, Jan. 1998, XP 002068503.
Allan, G.M. et al., 1995, *Vet. Microbiol.*, 44: 49-64.

(Continued)

Primary Examiner—Ali R. Salimi
(74) *Attorney, Agent, or Firm*—Bingham McCutchen, LLP

(57) **ABSTRACT**

The genome sequences and the nucleotide sequences coding for the PWD circovirus polypeptides, such as the circovirus structural and non-structural polypeptides, vectors including the sequences, and cells and animals transformed by the vectors are provided. Methods for detecting the nucleic acids or polypeptides, and kits for diagnosing infection by a PWD circovirus, also are provided. Method for selecting compounds capable of modulating the viral infection are further provided. Pharmaceutical, including vaccine, compositions for preventing and/or treating viral infections caused by PWD circovirus and the use of vectors for preventing and/or treating diseases also are provided.

68 Claims, 29 Drawing Sheets

U.S. PATENT DOCUMENTS

7,148,015	B2	12/2006	Jestin et al.
7,179,472	B2	2/2007	Jestin et al.
7,223,407	B2	5/2007	Jestin et al.
7,223,594	B2	5/2007	Jestin et al.
7,244,433	B2	7/2007	Jestin et al.
7,258,865	B2	8/2007	Jestin et al.
7,261,898	B2	8/2007	Jestin et al.
7,297,537	B2	11/2007	Jestin et al.
7,314,628	B2	1/2008	Jestin et al.
2002/0055189	A1	5/2002	Bernhardt et al.
2002/0106639	A1	8/2002	Li et al.
2002/0177216	A1	11/2002	Phillip et al.
2003/0170616	A1	9/2003	Wang et al.
2005/0058653	A1	3/2005	Ellis et al.

FOREIGN PATENT DOCUMENTS

FR	2772047	6/1999
SU	1 538 305	12/1994
WO	WO 90/11092	10/1990
WO	92/05255	4/1992
WO	94/01133	1/1994
WO	94/21797	9/1994
WO	WO 94/27238	11/1994
WO	94/27435	12/1994
WO	WO 95/11307	4/1995
WO	96/34109	10/1996
WO	96/40931	12/1996
WO	96/40945	12/1996
WO	98/40499	9/1998
WO	WO 99/18214	4/1999
WO	99/29717	6/1999
WO	99/29871	6/1999
WO	00/01409	1/2000
WO	00/24428	5/2000
WO	00/77216	12/2000
WO	WO 00/77188	12/2000
WO	WO 01/96377	6/2002
WO	WO 02/102999	12/2002

OTHER PUBLICATIONS

Author Unknown, GenBank info on AF027217 (Revised Jul. 5, 2002).

Author Unknown, NCBI Sequence Revision History of AF027217 (Revised Jul. 5, 2002).

Barany, F., 1991, PNAS. USA, 88: 189-193.

Blanchard et al., 2003, "An ORF2 Protein-Based ELISA for Porcine Circovirus Type 2 Antibodies in Post-Weaning Multisystemic Wasting Syndrome", *Veterinary Microbiology*, 94: 183-184.

Blanchard et al., 2003, "Protection of Swine Against Post-Weaning Multisystemic Wasting Syndrome (PMWS) by Porcine Circovirus Type 2 (PCV2) Proteins", *Vaccine* 21: 4565-4575.

Buckholz, R.G., 1993, Yeast systems for the expression of heterologous gene products. *Curr. Op. Biotechnology* 4: 538-542.

Burg, J.L. et al., 1996, *Mol. and Cell. Probes*, 10: 257-271.

Chu, B.C.F. et al., 1986, *NAR*, 14: 5591-5603.

Chu, P.W.G. et al., 1993, *Virus Research*, 27: 161-171.

Chu, Te-hua Terina et al., 1997, "Toward Highly Efficient Cell-Type-Specific Gene Transfer with Retroviral Vectors Displaying Single-Chain Antibodies", *J. Virol.*, 71(1): 720-725.

Clark, E.G., 1997, American Association of Swine Practitioners, 499-501.

Cosset, Francois-Loic et al., 1995, "Retroviral by Envelopes Expressing an N-Terminal Binding Domain", *J. Virol.*, 69:6314-6322.

Daft, B. et al., 1996, American Association of Veterinary Laboratory Diagnosticians, 32.

Derse, D. et al., 1995, *J. Virol.*, 69(3): 1907-1912.

Derwent Abstract of French Patent Appl. No. 2422956, 1979.

Derwent Abstract of French Patent Appl. No. 2518755, 1983.

Dialog Inpadoc Family and Legal Status Search for German Patent Appl. No. 10044648, 2000.

Dialog Inpadoc Family and Legal Status Search for WO 01/96377, 2001.

Dialog Inpadoc Family and Legal Status Search for WO 02/102999, 2002.

Duck, P. et al., 1990, *Biotechniques*, 9: 142-147.

Dulac, G.C. et al., 1989, *Can. J. Vet. Res.*, 53: 431-433.

Edwards, C.P., and Aruffo, A., 1993, Current applications of COS cell based transient expression systems. *Curr. Op. Biotechnology* 4: 558-563.

Edwards, S. et al., 1994, *Vet. Rec.*, 134: 680-681.

Ellis, J. et al., "Isolation of Circovirus from legions of Pigs with Postweaning Multisystemic Wasting Syndrome", *Canadian Veterinary Journal*, col. 39, pp. 44-51, Jan. 1998, XP-002068502.

Erlich, H.A., 1989, In *PCR Technology. Principles and Applications for DNA Amplification*. New York: Stockton Press.

Felgner, et al., 1987, *Proc. Natl. Acad. Sci.*, 84:7413-7417.

Fontes, E.P.B. et al., 1994, *J. Biol. Chem.*, vol. 269, No. 11:8459-8465.

Fralely et al., 1980, *J. Biol. Chem.*, 255: 10431-10435.

Guateli, J.C. et al., 1990, *Proc. Nat'l Acad. Sci., USA*, 87: 1874-1878.

Hackland, A.F. et al., 1994, *Arch. Virol.*, 139: 1-22.

Hamel, A. et al., "Nucleotide Sequence of Porcine Circovirus Associated with Postweaning Multisystemic Wasting Syndrome in Pigs", *Journal of Virology*, vol. 72, No. 6, pp. 5262-5267, Jun. 1998, XP-002078783.

Hanson, S.F. et al., 1995, *Virology*, 211: 1-9.

Harding, J.C. and Clark, E.G., 1997, *Swine Health and Production*, vol. 5, No. 5: 201-203.

Harding, J.C., 1997, American Association of Swine Practitioners, 503.

Harding, R.M. et al., 1993, *Journal of General Virology*, 74: 323-328.

Heyraud-Nitschke, F. et al., 1995, *Nucleic Acids Research*, vol. 23, No. 6: 910-916.

Horner, G.W., 1991, *Surveillance* 18(5): 23.

Huygen, K. et al., 1996, *Nature Medicine*, 2(8): 893-898.

Innis, M.A. et al., 1990, in *PCR Protocols. A guide to Methods and Applications*, San Diego, Academic Press.

Kaneda, et al., 1989, *Science*, 243: 375-378.

Kasahara, Noriyuki et al., 1994, "Tissue-Specific Targeting of Retroviral Vectors Through Ligan-Receptor Interactions", *Science*, 266: 1373-1376.

Kessler, C., "Overview of Amplification on Systems" in *Non-radioactive Labeling and Detection of Biomolecules*, 1992, Springer Verlag, Berlin, New-York: 197-205.

Kievitis, T. et al., 1991, *J. Virol. Methods*, 35: 273-286.

Kohler, G. & Milstein, 1975, *Nature*, 256(5517): 495-497.

Kwoh, D.Y. et al., 1989, *Proc. Nat'l Acad. Sci., USA*, 86: 1173-1177.

Ladany, S. et al., 1989, *J. Clin. Microbiol.* 27: 2778-2783.

Lazarowitz, S.G. et al., 1989, *The EMBO Journal*, vol. 8 No. 4: 1023-1032.

Liu et al., 1997, *J. Gen. Virol.*, 78 (Pt 6), 1265-1270.

Luckow, V.A., 1993, Baculovirus systems for the expression of human gene products. *Curr. Op. Biotechnol.* 4: 564-572.

Mankertz, A. et al., 1997, *J. Virol.*, 71: 2562-2566.

Marglin, A. and Merrifield, R.B., 1966, *J. Am. Chem. Soc.*, 88(21): 5051-5052.

Matthews, J.A. et al., 1988, *Anal. Biochem.*, 169: 1-25.

McNeilly, F. et al., 1996, *Vet. Immunol. Immunopathol.*, 49: 295-306.

Meehan, B.M. et al., 1997, *J. Gen. Virol.* 78: 221-227.

Midoux, 1993, *Nucleic Acids Research*, 21: 871-878.

Miele, E.A. et al., 1983, *J. Mol. Biol.* 171: 281-295.

Morozov, I. et al., "Detection of a Novel Strain of Porcine Circovirus in Pigs with Postweaning Multisystemic Wasting Syndrome", *Journal of Clinical Microbiology*, col. 36, No. 9, pp. 2535-2541, Sep. 1998, XP-002090921.

Müller, 1974, in *Methode der Organischen Chemie*, E. Wunsch Ed., vol. 15-I and 15-II, Thieme, Stuttgart.

Nayar, G.P. et al., 1997, *Can. Vet. J.* 38(6): 385-386.

Neddleman, Saul B. et al., 1970, "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins", *J. Mol. Biol.*, 48(3): 443-453.

- Olins, P.O., and Lee, S.C., 1993, *Curr. Op. Biotechnol.* 4: 520-525.
- Pagano et al., 1967, *J. Virol.*, 1: 891-897.
- Pearson, William R. et al., 1988, "Improved Tools for Biological Sequence Comparison", *Proc. Nat'l Acad. Sci., USA*, 85: 2444-2448.
- Rolfs, A. et al., "Usage of Polymerase Chain Reaction in Genetic and Infectious Disease" in *PCR Topics*, 1991, Springer-Verlag, Berlin.
- Rose, N. et al., 2002, "Risk Factors for Porcine Post-Weaning Multisystemic Wasting Syndrome (PMWS) in 149 French Farrow-to-Finish Herds", *Preventive Veterinary Microbiology*, 61: 209-225.
- Sambrook, J. et al., 1989, *Molecular cloning: A Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Sanchez-Pescador, R., 1988, *J. Clin. Microbiol.*, 26(10): 1934-1938.
- Ségales, J. et al., "First Report of Post-Weaning Multisystemic Wasting Syndrome in Pigs in Spain", *Beterinary Record*, col. 141, No. 23, pp. 600-601, Dec. 1997, XP-002068504.
- Shiver, J.W., "Immune Responses to HIV gp 120 Elicited by DNA Vaccination," in *Vaccines*, 1995, eds Chanock, et al., pp. 95-98, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Smith, Temple F. et al., 1981, "Comparison of Biosequences", *Advances in Applied Mathematics*, 2: 482-489.
- Tascon, R.E. et al., 1996, *Nature Medicine*, 2(8): 888-892.
- Tischer, I. and Bohk, H.J., 1988, *Zentralbl. Bakteriologie Mikrobiol. Hyg. [A]* 270: 280-287.
- Tischer, I. et al., 1982, *Nature*, 295: 64-66.
- Tischer, I. et al., 1986, *Arch. Virol.*, 91: 271-276.
- Tischer, I. et al., 1995, *Arch. Virol.*, 140: 737-743.
- Urdea, M.S., 1988, *Nucleic Acids Research*, 11: 4937-4957.
- Valesia-Whittmann, Sandrine et al., 1996, "Improvement of Retroviral Retargeting by Using Amino Acid Spacers between an Additional Binding Domain and the N Terminus of Moloney Nurine Leukemia Virus SU", *J. Virol.* 70(3): 2059-2064.
- Virus Taxonomy, Murphy, F.A. et al., Ed. 1995, *Sixth Report of the International Committee on Taxonomy of Viruses*, Springer-Verlag Wien New York.
- Walker, G.T. et al., 1992, *Nucleic Acids Res.* 20: 1691-1696.
- Walker, G.T. et al., 1992, *Proc. Nat'l Acad. Sci., USA*, 89: 392-396.
- White, B.A. et al. Eds., "PCR Cloning Protocol" in *Methods in Molecular Biology*, 67, Humana Press, Towota, 1997.
- Young, John A. T. et al., "Efficient Incorporation of Human CD4 Protein into Avian Leukosis Virus Particles", *Reports*, 1421-1423, 1990.
- Zhao, T.M. et al., 1996, *Proc. Natl. Acad. Sci., USA* 96(13): 6653-6648.
- ABI-Prism, "Automated DNA Sequencing, Chemistry Guide", Applied Biosystems, product manual, applicable to automated sequencers ABI Prism 310, 377 and 373 (2000).
- Adams, Mark D. et al. "Complementary DNA Sequencing: Expressed Sequence Tags and Human Genome Project," *Science*, 1651-1656. (1991).
- Albina et al., "Premiers Résultats du CNEVA sur le dépérissement fatal du porcelet en fin de post-sevrage" *La Semaine Veterinaire des Filières*, 26:1-2, Nov. 30, 1996.
- Allan et al., "Porcine Circoviruses: A Review," *J Vet Diagn Invest.* 12: 3-14 (2000).
- Allan et al., "Immunostimulation, PCV-2 and PMWS," *Veterinary Record* (2000); 147(6):170-171.
- Allan, G.M., et al., "Production, preliminary characterisation and applications of monoclonal antibodies to porcine circovirus," *Veterinary Immunology and Immunopathology*, 43 (1994) 357-371.
- Altmann, Curtis R. et al. "Microarray-Based Analysis of Early Development in *Xenopus laevis*," *Developmental Biology*, 236:64-75 (2001).
- Bantle, John A. et al. "Phase III Interlaboratory Study of FETAX Part 3. FETAX Validation using 12 Compounds with and without an Exogenous Metabolic Activation System," *Journal of Applied Toxicology*, 19:447-472 (1999).
- Behr, J.P., "Gene Transfer with Synthetic Cationic Amphiphiles: Prospects for Gene Therapy" *Bioconjugate Chem.* 5(5):382-389 (1994).
- Bei, R. et al., "The Use of a Cationic Liposome Formulation (DOTAP) Mixed with a Recombinant Tumor-Associated Antigen to Induce Immune Responses and Protective Immunity in Mice," *Journal of Immunotherapy*, 21(3):159-169 (1998).
- Cho et al., "Enhanced cellular immunity of hepatitis C virus nonstructural proteins by codelivery of granulocyte macrophage-colony stimulating factor gene in intramuscular DNA immunization," *Vaccine* 17:1136-1144 (1999).
- Cruse et al., "The Illustrated Dictionary of Immunology," Boca Raton: CRC Press (1995) p. 156.
- Cruse et al., "The Illustrated Dictionary of Immunology," 2nd edition. Boca Raton: CRC Press (2003) p613.
- Dedet, V., "Seule certitude: un malade à part entière," *La Semaine Veterinaire* p. 54, May 24, 1997.
- DMRIE structure, Printed from PubChem Compound Summary 2008.
- Donnelly, John J. et al., "Immunization with DNA" *Journal of Immunological Methods*, 176: 145-152 (1994).
- Dorland's Illustrated Medical Dictionary, 28th edition. Philadelphia. WB Saunders p. 1787 (1994).
- Ellis et al., "Reproduction of lesions of Post weaning Multisystemic wasting syndrome in gnotobiotic piglets," *J. Vet. Diagn. Invest.* 11:3-14 (1999).
- Ellis et al., "Coinfection by porcine circoviruses and porcine parvovirus in pigs with naturally acquired postweaning multisystemic wasting syndrome" *J. Vet. Diagn. Invest.* 12(1):21-27 (2000).
- Ertl, H.C.J. et al., "Genetic Immunization" *Viral Immunology*, 9(1):1-9 (1996).
- Felgner J.H. et al., "Enhanced gene delivery and mechanism studies with a novel series of cationic lipid formulations," *The journal of biological chemistry*, 269(4):2550-2561 (1994).
- Fenaux, M. et al., "Genetic Characterization of Type 2 Porcine Circovirus (PCV-2) from Pigs with Postweaning Multisystemic Wasting Syndrome in Different Geographic Regions of North America and Development of a Differential PCR-Restriction Fragment Length Polymorphism Assay to Detect and Differentiate between Infections with PCV-1 and PCV-2," *Journal of Clinical Microbiology*, Jul. 2000, 38(7):2494-2503.
- Fort, Douglas J. et al. "Evaluation of the Developmental Toxicity of Thalidomide Using Frog Embryo Teratogenesis Assay--Xenopus (FETAX): Biotransformation and Detoxification," *Teratogenesis, Carcinogenesis, and Mutagenesis*, 20:35-47 (2000).
- GenBank Accession No. AF027217; May 14, 1998.
- Gregoriadis G. et al., "Liposome-mediated DNA vaccination" *FEBS Letters*, Feb. 3, 1997; 402:107-110.
- Gregoriadis, Gregory, "Genetic vaccines: strategies for optimization" *Pharmaceutical Research*, 15(5):661-670 (1998).
- Grierson, S.S., et al., "Genome sequence analysis of 10 Dutch porcine circovirus type 2 (PCV-2) isolates from a PMWS case-control study," *Research in Veterinary Science*, 77 (2004) pp. 265-268.
- Haddad, D. et al., "Comparative study of DNA-based immunization vectors: effect of secretion signals on the antibody responses in mice," *FEMS Immunology and Medical Microbiology*, 18(3):193-202 (1997).
- Hamel et al., "Porcine Circovirus," (Sequence Alignments for SEQ ID No:s 1-4 and 6) Genbank Acc No: 027217, Dec. 19, 1997.
- Inumaru, S. et al., "cDNA Cloning of Porcine Granulocyte-Macrophage Colony-Stimulating Factor" *Immunology and Cell Biology*, 73:474-476, XP-000946635 (1995).
- Ishii N. et al., "Cationic liposomes are a strong adjuvant for a DNA vaccine of human immunodeficiency virus type 1" *AIDS Res Hum Retroviruses*, Nov. 1, 1997 13(6):1421-1428.
- Izumida, et al., "Establishment of the Attenuated Strain of Porcine Parvovirus of the Live Vaccine and its Biological-Immunological Characteristics," *Japanese Veterinary Science*, 48(2):293-303 (1996).
- Klavinskis, Linda S. et al., "Intranasal Immunization with Plasmid DNA-Lipid Complexes Elicits Mucosal Immunity in the Female Genital and Rectal Tracts" *J. Immunol* 1999, pp. 254-262.
- Knell, S. et al., "Comparative genetic characterization of Porcine Circovirus type 2 samples from German wild boar populations," *Veterinary Microbiology*, 109 (2005) pp. 169-177.
- Krakowka et al., "Activation of the Immune System in the Pivotal Event in the Production of Wasting Disease in Pigs Infected with Porcine Circovirus-2 (PCV-2)," *Vet Pathology*. 38:31-42 (2001).

- Krakowka et al., "Immunologic Features of Porcine Circovirus Type-2 Infection," *Viral Immunology*, 15(4): 567-582 (2002).
- LeCann et al., "Piglet Wasting disease," *Veterinary Record*, p. 660, Dec. 20/27 1997.
- Lekcharoensuk, P. et al., "Epitope Mapping of the Major Capsid Protein of Type 2 Porcine Circovirus (PCV2) by Using Chimeric PCV1 and PCV2," *Journal of Virology*, vol. 78, No. 15, Aug. 2004, pp. 8135-8145.
- Mahé et al., "Differential Recognition of ORF 2 Protein from Type 1 and Type 2 Porcine circoviruses and Identification of Immunorelevant Epitopes", *Journal of General Virology* 81:1815-1824 (2000).
- Mankerts et al., "Porcine Circovirus Complete Genome", EMBL Sequence Database XP-002104869 May 22, 1997.
- McCluskie, Michael J. et al., "Route and Method of Delivery of DNA Vaccine Influence Immune Responses in Mice and Non-Human Primates" *Molecular Medicine* 5:287-300 (1999).
- McInnes, C.J. et al., "Cloning and Expression of a cDNA Encoding Ovine Granulocyte-Macrophage Colony-Stimulating Factor" *Gene*, 105: 275-279, XP-002148815 (1991).
- Meehan et al., "Putative PCV Replication-Associated Protein (REP)", EMBL Sequence Database XP 002104867 (1997).
- Meehan et al., "Porcine Circovirus Complete Genome", EMBL Sequence Database XP-002104868 (1997).
- Meehan, B.M. et al., "Characterization of novel circovirus DNAs associated with wasting syndromes in pigs." , *Journal of General Virology*, vol. 79, No. 9, pp. 2171-2179, Sep. 1998, XP-002090386.
- Miller, J.S. et al., "The nucleotide sequence of RNA-1 of Indian peanut clump furovirus" *Arch. Virol.* 141:2301-2312, (1996).
- Mumford J.A. et al., "Antigenicity and immunogenicity of equine influenza vaccine containing a Carbomer adjuvant," *Epidemiol. Infect.* 112: 421-437 (1994).
- Nakakura, Norihiko et al. "Synthesis of Heterogenous mRNA-like RNA and Low-Molecular-Weight RNA before the Midblastula Transition in Embryos of *Xenopus laevis*," *Developmental Biology*, 123:421-429 (1987).
- Nash RA et al., "Molecular cloning and in vivo evaluation of canine granulocyte-macrophage colony-stimulating factor" *Blood* 78(4):930-937, XP 002133949 (1991).
- Nawagtigul et al., "Open Reading Frame 2 of Porcine Circovirus Type 2 Encodes a Major Capsid Protein", *Journal of General Virology* 81:2281-2287 (2000).
- Nayer et al., "Detection and Characterization of Porcine Circovirus Associated With Postweaving Multisystemic Wasting Syndrome Pigs", *Can Vet. J.* 38:385-386 (1997).
- Newport, John et al., "Major Developmental Transition in Early *Xenopus* Embryos: I. Characterization and Timing of Cellular Changes at the Midblastula Stage," *Cell* 30:675-686 (1982).
- Newport, John et al. "A Major Developmental Transition in Early *Xenopus* Embryos: II. Control of the Onset of Transactions," *Cell* 30:687-696 (1982).
- Nieuwkoop and Faber "Normal Table of *Xenopus laevis* (Daudin) - A Systematic and chronological survey of the development from the fertilized egg to the end of metamorphosis, " Chapter VII, 162-188 Garland Publishing(1994).
- Norman, JA et al., "Development of Improved Vectors for DNA-Based Immunization and other Gene Therapy Applications," *Vaccine*, 15(8): 801-803 (1997).
- Nuwaysir, Emile F. et al. "Microarrays and Toxicology: The Advent of Toxicogenomics," *Molecular Carcinogenesis*, 24:153-159 (1999).
- Okada, E. et al., "Intranasal immunization of a DNA vaccine with IL-12 and Granulocyte-macrophage colony-stimulating Factor (GM-(SF)) - expressing plasmids in Liposomes Induces Strong Mucosal and Cell-Mediated immune responses against HIV-1 antigens," *The Journal of Immunology*, 159:3638-3647 (1997).
- Parker, SE et al., "Plasmid DNA gene therapy: Studies with the human interleukin-2 gene in tumor cells in vitro and in the murine B16 melanoma model in vivo," *Cancer Gene Therapy*, 3(3):175-185 (1996).
- Restifo, N P et al., "The promises of nucleic acid vaccines," *Gene Therapy*, 7:89-92 (2000).
- Ruitenber, K. M. et al., "DNA-Mediated Immunization with Glycoprotein D of Equine Herpesvirus 1 (EHV-1) in a Murine Model of EHV-1 Respiratory Infection," *Vaccine*, 17:237-244 (1999).
- Schena, Mark, et al., "Quantitative Monitoring of Gene Expression Patterns with a Complementary DNA Microarray," *Science*, 270: 467-470 (1995).
- Schena, Mark et al. "Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes," *Proc. Natl. Acad. Sci. USA*, 93:10614-10619 (1999).
- Schultz, J et al., "Update on Antiviral DNA Vaccine Research (1998-2000)," *Intervirology*, 43: 197-217 (2000).
- Sin, J-I et al., "Protective Immunity Against Heterologous Challenge with Encephalomyocarditis Virus by VP1 DNA Vaccination: Effect on Coinjection with a Granulocyte-Macrophage Colony Stimulating Factor Gene," *Vaccine* 15:1827-1833 (1997).
- Somasundaram, C. et al., "Enhanced Protective Response and Immuno-Adjuvant Effects of Porcine GM-CSF on DNA Vaccination of Pigs against Aujeszky's Disease Virus," *Veterinary Immunology and Immunopathology* 70:277-287 (1999).
- Terpestra et al., "Potency control of modified live viral vaccines for veterinary use," *Vaccine* 14(6):570-575 (1996).
- Todd et al., "Comparison of Three Animal Viruses with Circular Single-Stranded DNA Genomes," *Arch Virol.*, 117:129-135 (1991).
- Vannier, et al., "Study of the Efficacy of an Inactivated Virus Vaccine Against Porcine Parvovirus," *Ann. Rech. Vet.* 17(4):425-432 (1986).
- Vogel, et al., "Nucleic Acid Vaccines," *Clinical Microbiology Reviews* 8(3):406-410 (1995).
- Walker, G.T. et al., *Nucleic Acids Res.* (1992) 20: 1691-1696.
- Watanabe, Y et al., "Highly Efficient Transfection into Primary Cultured Mouse Hepatocytes by Use of Cation-Liposomes: An Application for Immunization," *J. Biochem.* 116:1220-1226 (1994).
- Wen, L. et al., "Genotyping of porcine circovirus type 2 from a variety of clinical conditions in China," *Veterinary Microbiology*, 110(2005) pp. 141-146.
- West et al., "Myocarditis and abortion associated with intrauterine infection of sows with porcine circovirus 2" *J Vet Diagn Invest.* 11: 530-532 (1999).
- Wheeler, C.J. et al., "Converting an alcohol to an amine in a cationic lipid dramatically alters the co-lipid requirement, cellular transfection activity and the ultrastructure of DNA-cytoflectin complexes," *Biochimica et Biophysica Acta* 1280:1-11, XP 002035803 (1996).
- Xiang Z. et al., "Manipulation of the immune response to a plasmid-encoded viral antigen by coinoculation with plasmids expressing cytokines," *Immunity*, 2:129-135 (1995).
- Yokoyama et al. "DNA immunization: Effects of vehicle and route of administration on the induction of protective antiviral immunity," *FEMS Immunology and Medical Microbiology*, 14:221-230 (1996).
- Young, "Fields Virology" 3rd ed. Philadelphia: Lippencott-Raven Publishers Chapter 70- Parvoviruses; vol. 2, p2199-2220 (1996).
- Zhang, Michael Q. "Large-Scale Gene Expression Data Analysis: A New Challenge to Computational Biologists," *Genome Research*, 9(8):681-688 (1999).




FIG. 1

Leu Ala Ser Arg Cys Arg Cys Cys Arg Pro Leu Thr Leu Ser Phe Ala Leu Cys
 Trp Arg Val Glu Ala Ala Ala Ala Gly Arg Cys Arg *** His Phe His Trp Ala
 Gly Ala Cys Lys Pro Leu Pro Leu Val Glu Ala Ala Asp Thr Phe Ile Gly Leu
 3' TGG TCG CGT GAA GCC GTC GCC GTC GTG GAG CCG TCG CAG TCA CTT TTA CGG TTC
 9 18 27 36 45 54
 5' ACC AGC GCA CTT CGG CAG CGG CAG CAC CTC GGC AGC GTC AGT GAA AAT GCC AAG
 Thr Ser Ala Leu Arg Gln Arg Gln His Leu Gly Ser Val Ser Glu Asn Ala Lys
 Pro Ala His Phe Gly Ser Gly Ser Thr Ser Ala Ala Ser Val Lys Met Pro Ser
 Gln Arg Thr Ser Ala Ala Ala Ala Pro Arg Gln Arg Gln *** Lys Cys Gln Ala

 Ser Phe Arg Gly Ala Val Gly Tyr Ser Thr Pro Thr *** Gly *** Tyr Asp Lys
 Leu Phe Ala Ala Arg Leu Gly Met Leu Pro Pro His Glu Gly Lys Ile Ile Arg
 Leu Phe Leu Pro Gly Cys Gly Trp Leu Leu His Thr Asn Val Arg Leu Leu Gly
 GTT CTT TTC GCC GGG CGT TGG GGT ATT CTC CAC CCA CAA GTG GGA ATT ATT AGG
 63 72 81 90 99 108
 CAA GAA AAG CGG CCC GCA ACC CCA TAA GAG GTG GGT GTT CAC CCT TAA TAA TCC
 Gln Glu Lys Arg Pro Ala Thr Pro *** Glu Val Gly Val His Pro *** *** Ser
 Lys Lys Ser Gly Pro Gln Pro His Lys Arg Trp Val Phe Thr Leu Asn Asn Pro
 Arg Lys Ala Ala Arg Asn Pro Ile Arg Gly Gly Cys Ser Pro Leu Ile Ile Leu

 Arg Pro Pro Ser Phe Cys Phe Val Pro Ala Glu Leu Arg Gly Lys Gln Asn Asn
 Gly Leu Leu Leu Phe Val Phe Tyr Pro Leu Lys Trp Asp Gly Lys Lys Ile Ile
 Glu Ser Ser Ser Phe Phe Leu Ile Arg Ser Ser Gly Ile Glu Arg Lys Ser ***
 AAG GCT CCT CCT CTT TTT GTT TTA TGC CCT CGA AGG TTA GAG GGA AAA ACT AAT
 117 126 135 144 153 162
 TTC CGA GGA GGA GAA AAA CAA AAT ACG GGA GCT TCC AAT CTC CCT TTT TGA TTA
 Phe Arg Gly Gly Glu Lys Gln Asn Thr Gly Ala Ser Asn Leu Pro Phe *** Leu
 Ser Glu Glu Glu Lys Asn Lys Ile Arg Glu Leu Pro Ile Ser Leu Phe Asp Tyr
 Pro Arg Arg Arg Lys Thr Lys Tyr Gly Ser Phe Gln Ser Pro Phe Leu Ile Ile

 Gln Lys His Arg Pro Leu Asn Pro Leu Pro Tyr Phe Glu Glu Gly Gly Pro Thr
 Lys Asn Thr Ala Leu Phe Thr Gln Phe Leu Thr Ser Ser Arg Val Glu Leu Pro
 Lys Thr Gln Pro Ser Ser Pro Lys Ser Ser Pro Leu Val Gly *** Arg Trp Pro
 AAA ACA AAC ACC CCT TCC AAA CCT TCT CCC ATC TTG AGG AGT GGA GGT CCC
 171 180 189 198 207 216
 TTT TGT TTG TGG CGA GGA AGG TTT GGA AGA GGG TAG AAC TCC TCA CCT CCA GGG
 Phe Cys Leu Trp Arg Gly Arg Phe Gly Arg Gly *** Asn Ser Ser Pro Pro Gly
 Phe Val Cys Gly Glu Glu Gly Leu Glu Glu Gly Arg Thr Pro His Leu Gln Gly
 Leu Phe Val Ala Arg Lys Val Trp Lys Arg Val Glu Leu Leu Thr Ser Arg Gly

 Gln Ser Asn Gln *** Ser Ala Ser Lys *** Cys Pro Ser Thr Thr Asn Gln His
 Lys Arg Ile Lys Ser Leu Leu Leu Ser Lys Val Leu His Leu Pro Ile Lys Thr
 Asn Ala Phe Lys Ala Leu Phe Cys Val Lys Leu Leu Thr Phe His Tyr Lys Pro
 CAA ACG CTT AAA ACG ATT CTT CGT CTG AAA ATT GTT CCA CTT CAC CAT AAA ACC
 225 234 243 252 261 270
 GTT TGC GAA TTT TGC TAA GAA GCA GAC TTT TAA CAA GGT GAA GTG CTA TTT TGG
 Val Cys Glu Phe Cys *** Glu Ala Asp Phe *** Gln Gly Glu Val Val Phe Trp
 Phe Ala Asn Phe Ala Lys Lys Gln Thr Phe Asn Lys Val Lys Trp Tyr Phe Gly
 Leu Arg Ile Leu Leu Arg Ser Arg Leu Leu Thr Arg *** Ser Gly Ile Leu Val

FIG. 2a

Gly Ser Gly Cys Arg Ser Leu Ser Leu Phe Arg Gly Ala Ser Tyr Leu Ile Ser
 Gly Ala Ala Val Asp Leu Phe Arg Phe Ser Gly Val Leu Leu Ile Phe Phe Val
 Ala Arg Gln Trp Met Ser Phe Ala Phe Pro Val Ser Trp Cys Phe Leu Ser Tyr

 ACG GGC GAC GGT GTA GCT CTT TCG CTT TCC TTG GCT GGT CGT CTT ATT TCT TAT
 279 288 297 306 315 324
 TGC CCG CTG CCA CAT CGA GAA AGC GAA AGG AAC CGA CCA GCA GAA TAA AGA ATA

 Cys Pro Leu Pro His Arg Glu Ser Glu Arg Asn Arg Pro Ala Glu *** Arg Ile
 Ala Arg Cys His Ile Glu Lys Ala Lys Gly Thr Asp Gln Gln Asn Lys Glu Tyr
 Pro Ala Ala Thr Ser Arg Lys Arg Lys Glu Pro Thr Ser Arg Ile Lys Asn Thr

 Cys Tyr Leu Leu Gly Cys Val *** Arg Thr His Leu Glu Ala Ser Gly Pro Ser
 Ala Thr Phe Phe Ala Val Tyr Lys Asp Leu Thr Ser Ser Arg Pro Val Leu Pro
 Gln Leu Leu Ser Pro Trp Met Ser Ile Ser His Pro Ala Gly Arg Phe Trp Pro

 GAC GTC ATT TCT TCC GGT GTA TGA ATA GCT CAC ACC TCG AGG CGC CTT GGT CCC
 333 342 351 360 369 378
 CTG CAG TAA AGA AGG CCA CAT ACT TAT CGA GTG TGG AGC TCC GCG GAA CCA GGG

 Leu Gln *** Arg Arg Pro His Thr Tyr Arg Val Trp Ser Ser Ala Glu Pro Gly
 Cys Ser Lys Glu Gly His Ile Leu Ile Glu Cys Gly Ala Pro Arg Asn Gln Gly
 Ala Val Lys Lys Ala Thr Tyr Leu Ser Ser Val Glu Leu Arg Gly Thr Arg Gly

 Ala Cys Arg Gly Thr *** Gln Gln Ser Tyr Gly Lys Pro Ser Pro Thr Lys Pro
 Leu Ala Ala Val Gln Arg Ser Ser His Thr Gly Lys Gln Leu Arg Pro Arg Gln
 Phe Arg Leu Ser Arg Asp Val Ala Thr Leu Val Arg Lys Ser Val Pro Asp Lys

 CTT CGC GTC GCT GGA CAG ATG ACG ACA CTC ATG GGA AAA CCT CTG CCC CAG AAA
 387 396 405 414 423 432
 GAA GCG CAG CGA CCT GTC TAC TGC TGT GAG TAC CCT TTT GGA GAC GGG GTC TTT

 Glu Ala Gln Arg Pro Val Tyr Cys Cys Glu Tyr Pro Phe Gly Asp Gly Val Phe
 Lys Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu Thr Gly Ser Leu
 Ser Ala Ala Thr Cys Leu Leu Leu *** Val Pro Phe Trp Arg Arg Gly Leu Trp

 Ser Gln Leu Arg Ala Thr Glu Gln Leu Thr His Ser Phe Asn Gly Arg Ala Pro
 His Ser Tyr Gly Leu Leu Lys Arg Tyr Arg Ile His Ser Ile Glu Ala Pro Gln
 Thr Val Thr Ala Ser Cys Asn Gly Thr Val Tyr Thr Leu Phe Lys Arg Pro Ser

 CCA CTG ACA TCG GCT CGT CAA AGG ACA TTG CAT ACA CTC TTT AAA GGC GCC CGA
 441 450 459 468 477 486
 GGT GAC TGT AGC CGA GCA GTT TCC TGT AAC GTA TGT GAG AAA TTT CCG CGG GCT

 Gly Asp Cys Ser Arg Ala Val Ser Cys Asn Val Cys Glu Lys Phe Pro Arg Ala
 Val Thr Val Ala Glu Gln Phe Pro Val Thr Tyr Val Arg Asn Phe Arg Gly Leu
 *** Leu *** Pro Ser Ser Phe Leu *** Arg Met *** Glu Ile Ser Ala Gly Trp

 Gln Val Lys Ser Leu Ser Arg Ser Ser Ala Ala Ala His Asn Ser Ser Leu Gln
 Ser Phe Lys Gln Phe His Ala Pro Leu His Leu Leu Thr Ile Pro Leu Cys Ser
 Ala Ser Ser Lys Phe Thr Leu Pro Phe Ile Cys Cys Arg Ser Gln Phe Val Ala

 CCG ACT TGA AAA CTT TCA CTC GCC CTT CTA CGT CGT CGC ACT AAC CTT CTG TCG
 495 504 513 522 531 540
 GGC TGA ACT TTT GAA AGT GAG CGG GAA GAT GCA GCA GCG TGA TTG GAA GAC AGC

 Gly *** Thr Phe Glu Ser Glu Arg Glu Asp Ala Ala Ala *** Leu Glu Asp Ser
 Ala Glu Leu Leu Lys Val Ser Gly Lys Met Gln Gln Arg Asp Trp Lys Thr Ala
 Leu Asn Phe *** Lys *** Ala Gly Arg Cys Ser Ser Val Ile Gly Arg Gln Leu

FIG. 2b

Val Arg *** Leu Pro Gly Ala Arg Asn His Ser Ser Gly Thr Pro Gly Tyr Asn
 Tyr Val Asp Tyr His Ala Arg Gly Thr Thr Pro Leu Ala Leu Pro Gly Thr Ile
 Thr Cys Thr Met Thr Pro Gly Gly Pro Gln Pro Phe Leu Trp His Ala Arg Leu

 ACA TGT GCA GTA TCA CCC GGG CGG GCC AAC ACC CTT CTC GGT CAC CCG GGC ATT
 549 558 567 576 585 594
 TGT ACA CGT CAT AGT GGG CCC GCC CGG TTG TGG GAA GAG CCA GTG GGC CCG TAA

 Cys Thr Arg His Ser Gly Pro Ala Arg Leu Trp Glu Glu Pro Val Gly Pro ***
 Val His Val Ile Val Gly Pro Pro Gly Cys Gly Lys Ser Gln Trp Ala Arg Asn
 Tyr Thr Ser *** Trp Ala Arg Pro Val Val Gly Arg Ala Ser Gly Pro Val Ile

 Gln Gln Ala *** Pro Cys Arg Ser Ser Ala *** Tyr Phe Tyr Thr Thr Pro His
 Lys Ser Leu Arg Pro Val Gly Val Pro Leu Arg Thr Ser Ile Leu Pro Pro Ile
 Lys Ala Ser Gly Leu Ser Val *** Gln Phe Gly Leu Leu Phe Leu His His Ser

 AAA ACG ACT CGG ATC CCT GTG GAT GAC CTT CGG ATC ATC TTT ATT CAC CAC CCT
 603 612 621 630 639 648
 TTT TGC TGA GCC TAG GGA CAC CTA CTG GAA GCC TAG TAG AAA TAA GTG GTG GGA

 Phe Cys *** Ala *** Gly His Leu Leu Glu Ala *** *** Lys *** Val Val Gly
 Phe Ala Glu Pro Arg Asp Thr Tyr Trp Lys Pro Ser Arg Asn Lys Trp Trp Asp
 Leu Leu Ser Leu Gly Thr Pro Thr Gly Ser Leu Val Glu Ile Ser Gly Gly Met

 Ile Asp His Leu Leu Leu Gln Gln Lys Pro His Asn Lys His Ser Thr Val Lys
 Ser Ile Met Ser Phe Phe Asn Asn Asn Gln Ile Ile Lys Ile Ala Pro *** Arg
 Pro Tyr *** Pro Ser Ser Thr Thr Thr Lys Ser Ser Lys *** Pro Gln Asn Gly

 ACC TAT AGT ACC TCT TCT TCA ACA ACA AAA CCT ACT AAA AAT ACC GAC CAA TGG
 657 666 675 684 693 702
 TGG ATA TCA TGG AGA AGA AGT TGT TGT TTT GGA TGA TTT TTA TGG CTG GTT ACC

 Trp Ile Ser Trp Arg Arg Ser Cys Cys Phe Gly *** Phe Leu Trp Leu Val Thr
 Gly Tyr His Gly Glu Glu Val Val Val Leu Asp Asp Phe Tyr Gly Trp Leu Pro
 Asp Ile Met Glu Lys Lys Leu Leu Phe Trp Met Ile Phe Met Ala Gly Tyr Leu

 Pro His Asp Val Ser Val Thr His Gly Thr Asp Met Ser Gln Leu Ser *** Leu
 Pro Ile Ile *** Gln Ser Gln Thr Val Pro Ile Trp Gln Ser Tyr Leu Ser Phe
 Gln Ser Ser Arg Ser Leu Ser His Ser Arg Tyr Gly Asn Val Thr Ser Val Leu

 AAC CCT ACT AGA TGA CTC TGA CAC ACT GGC CAT AGG TAA CTG ACA TCT CTG ATT
 711 720 729 738 747 756
 TTG GGA TGA TCT ACT GAG ACT GTG TGA CCG GTA TCC ATT GAC TGT AGA GAC TAA

 Leu Gly *** Ser Thr Glu Thr Val *** Pro Val Ser Ile Asp Cys Arg Asp ***
 Trp Asp Asp Leu Leu Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Glu Thr Lys
 Gly Met Ile Tyr *** Asp Cys Val Thr Gly Ile His *** Leu *** Arg Leu Lys

 Pro Tyr Gln Glu Lys Lys Pro Gly Cys Tyr Lys Ser *** Trp Cys Asp Pro Gly
 Pro Thr Ser Asn Arg Lys Gln Gly Ala Thr Asn Gln Asn Gly Ala Ile Leu Gly
 Pro Pro Val Thr Gly Lys Lys Ala Arg Leu Ile Lys Ile Val Leu Leu *** Ala

 TCC CCC ATG ACA AGG AAA AAA CCG GGC GTC ATA AAA CTA ATG GTC GTT AGT CCG
 765 774 783 792 801 810
 AGG GGG TAC TGT TCC TTT TTT GGC CCG CAG TAT TTT GAT TAC CAG CAA TCA GGC

 Arg Gly Tyr Cys Ser Phe Phe Gly Pro Gln Tyr Phe Asp Tyr Gln Gln Ser Gly
 Gly Gly Thr Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn Gln Ala
 Gly Val Leu Phe Leu Phe Trp Pro Ala Val Phe *** Leu Pro Ala Ile Arg Pro

FIG. 2c

Gly Pro Ile Thr Ser Arg Leu Gln Gln Gly Leu Gln Leu Leu Glu Arg Asp Ser
 Gly Leu Phe Pro Val Gly *** Ser Ser Asp Trp Ser Tyr Phe Ser Glu Ile Pro
 Gly Trp Ser His Tyr Glu Glu Val Ala Thr Gly Ala Thr Ser Ala Arg *** Arg
 GGG GGT CCT TAC CAT GAG GAG TTG ACG ACA GGG TCG ACA TCT TCG AGA GAT AGC
 819 828 837 846 855 864
 CCC CCA GGA ATG GTA CTC CTC AAC TGC TGT CCC AGC TGT AGA AGC TCT CTA TCG
 Pro Pro Gly Met Val Leu Leu Asn Cys Cys Pro Ser Cys Arg Ser Ser Leu Ser
 Pro Gln Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu Ala Leu Tyr Arg
 Pro Arg Asn Gly Thr Pro Gln Leu Leu Ser Gln Leu *** Lys Leu Ser Ile Gly

 Ser *** *** Lys Ala Ile Lys Ser Ser Gln Gln Leu Val Ile Trp Pro Pro Val
 Pro Asn Ser Ser Gln Leu Lys Pro Leu Ser Ser Ser Phe Leu Gly Arg Leu Tyr
 Leu Ile Val Val Lys Cys Asn Gln Phe Val Ala Pro Ser Cys Asp Val Ser Thr
 CTC CTA ATG ATG AAA CGT TAA AAC CTT CTG ACG ACC TCT TGT TAG GTG CCT CCA
 873 882 891 900 909 918
 GAG GAT TAC TAC TTT GCA ATT TTG GAA GAC TGC TGG AGA ACA ATC CAC GGA GGT
 Glu Asp Tyr Tyr Phe Ala Ile Leu Glu Asp Cys Trp Arg Thr Ile His Gly Gly
 Arg Ile Thr Thr Leu Gln Phe Trp Lys Thr Ala Gly Glu Gln Ser Thr Glu Val
 Gly Leu Leu Leu Cys Asn Phe Gly Arg Leu Leu Glu Asn Asn Pro Arg Arg Tyr

 Arg Leu Gly Ile Gln Leu Leu Pro Gly Val Arg His Gly Lys Gly Met Tyr Phe
 Gly Phe Ala Ser Lys Phe Cys His Val Trp Gly Thr Gly Lys Glu Trp Ile Phe
 Gly Ser Pro Arg Asn Ser Ala Thr Ser Gly Gly Gln Ala Arg Lys Gly Tyr Leu
 TGG GCT TCC GGC TAA ACT TCG TCA CCT GGG TGG GAC ACG GGA AAA GGG TAT ATT
 927 936 945 954 963 972
 ACC CGA AGG CCG ATT TGA AGC AGT GGA CCC ACC CTG TGC CCT TTT CCC ATA TAA
 Thr Arg Arg Pro Ile *** Ser Ser Gly Pro Thr Leu Cys Pro Phe Pro Ile ***
 Pro Glu Gly Arg Phe Glu Ala Val Asp Pro Pro Cys Ala Leu Phe Pro Tyr Lys
 Pro Lys Ala Asp Leu Lys Gln Trp Thr His Pro Val Pro Phe Ser His Ile Lys

 Leu Asn Ser Leu Arg Lys Gln *** *** Met Thr Ile Thr Lys Ile Lys Ile ***
 Tyr Ile Val Ser Asp Lys Lys Asn Asp Cys Arg Leu Pro Lys *** Lys *** Glu
 Ile Phe *** Gln Thr Lys Lys Thr Ile Val Asp Tyr His Asn Lys Asn Lys Asn
 TTA TTT AAT GAC TCA GAA AAA ACA ATA GTG TAG CAT TAC CAA AAA TAA AAA TAA
 981 990 999 1008 1017 1026
 AAT AAA TTA CTG AGT CTT TTT TGT TAT CAC ATC GTA ATG GTT TTT ATT TTT ATT
 Asn Lys Leu Leu Ser Leu Phe Cys Tyr His Ile Val Met Val Phe Ile Phe Ile
 Ile Asn Tyr *** Val Phe Phe Val Ile Thr Ser *** Trp Phe Leu Phe Leu Phe
 *** Ile Thr Glu Ser Phe Leu Leu Ser His Arg Asn Gly Phe Tyr Phe Tyr Ser

 Lys Ser Pro Arg Glu Pro Tyr Ile Arg Gln Ile Thr Cys Leu Tyr Asp Val Lys
 Asn Leu Pro Asp Lys Leu Ile Phe Glu Arg Phe Gln Val Tyr Ile Thr Leu Arg
 Met *** Leu Thr Lys *** Ser Leu Asn Glu Ser Asn Tyr Met Phe Leu *** Gly
 GTA AAT CTC CCA GAA AGT CCT ATT TAA GAG ACT TAA CAT GTA TTT ATC AGT TGG
 1035 1044 1053 1062 1071 1080
 CAT TTA GAG GGT CTT TCA GGA TAA ATT CTC TGA ATT GTA CAT AAA TAG TCA ACC
 His Leu Glu Gly Leu Ser Gly *** Ile Leu *** Ile Val His Lys *** Ser Thr
 Ile *** Arg Val Phe Gln Asp Lys Phe Ser Glu Leu Tyr Ile Asn Ser Gln Pro
 Phe Arg Gly Ser Phe Arg Ile Asn Ser Leu Asn Cys Thr *** Ile Val Asn Leu

FIG. 2d

Gly Cys Leu Lys Pro Ser His Asn Cys Lys Pro Ala Cys Leu Gly Pro Arg His
 Val Val Tyr Asn Gln Ala Thr Thr Ala Asn Gln Leu Ala Tyr Gly Leu Gly Thr
 *** Trp Met Ile Lys Pro Gln Pro Gln Met Lys Ser Arg Met Ala Trp Ala Gln

 AAT GGT GTA TTA AAA CCC GAC ACC AAC GTA AAA CCT CGC GTA TCG GGT CCG GAC
 1089 1098 1107 1116 1125 1134
 TTA CCA CAT AAT TTT GGG CTG TGG TTG CAT TTT GGA GCG CAT AGC CCA GGC CTG

 Leu Pro His Asn Phe Gly Leu Trp Leu His Phe Gly Ala His Ser Pro Gly Leu
 Tyr His Ile Ile Leu Gly Cys Gly Cys Ile Leu Glu Arg Ile Ala Gln Ala Cys
 Thr Thr *** Phe Trp Ala Val Val Ala Phe Ser Ala *** Pro Arg Pro Val

 Ala Arg Cys Gln His Pro Tyr Lys Phe Pro Ala Val Ala Pro Lys Lys *** ***
 His Glu Val Asn Thr His Thr Asn Leu His Leu Trp Leu Gln Asn Arg Lys Asn
 Thr Ser Ser Met Pro Thr Pro Ile *** Ile Ser Gly Cys Ser Thr Glu Lys Ile

 ACA CGA GCT GTA ACC ACA CCC ATA AAT TTA CCT CGG TGT CGA CCA AAG AAA ATA
 1143 1152 1161 1170 1179 1188
 TGT GCT CGA CAT TGG TGT GGG TAT TTA AAT GGA GCC ACA GCT GGT TTC TTT TAT

 Cys Ala Arg His Trp Cys Gly Tyr Leu Asn Gly Ala Thr Ala Gly Phe Phe Tyr
 Val Leu Asp Ile Gly Val Gly Ile *** Met Glu Pro Gln Leu Val Ser Phe Ile
 Cys Ser Thr Leu Val Trp Val Phe Lys Trp Ser His Ser Trp Phe Leu Leu Leu

 Lys Ala Pro Val Leu *** Asn Asn Pro Arg Ala Arg Thr Gln Pro His Leu Val
 Asn Pro Gln Phe Trp Asp Ile Thr Gln Asp Leu Glu Pro Lys Pro Thr Phe Tyr
 Ile Gln Ser Ser Gly Ile Leu Gln Lys Thr *** Ser Gln Asn Pro Pro Ser Thr

 ATA AAC CGA CCT TGG TTA GTT AAC AAA CCA GAT CGA GAC CAA ACC CCC ACT TCA
 1197 1206 1215 1224 1233 1242
 TAT TTG GCT GGA ACC AAT CAA TTG TTT GGT CTA GCT CTG GTT TGG GGG TGA AGT

 Tyr Leu Ala Gly Thr Asn Gln Leu Phe Gly Leu Ala Leu Val Trp Gly *** Ser
 Ile Trp Leu Glu Pro Ile Asn Cys Leu Val *** Leu Trp Phe Gly Gly Gly Val
 Phe Gly Trp Asn Gln Ser Ile Val Trp Ser Ser Ser Gly Leu Gly Val Lys Tyr

 Gln Leu Pro Leu Tyr Leu Ala Ala Lys His His Pro Pro Leu Leu Leu *** Tyr
 Arg Ser His Tyr Thr Phe Pro Gln Arg Ile Thr His Arg Ser Ser Tyr Asn Ile
 Gly Pro Thr Thr Pro Leu Pro Ser Gly *** Pro Thr Ala Pro Pro Thr Thr Leu

 TGG ACC TCA CCA TCC ATT TCC CGA CGG AAT ACC ACA CCG CCC TCC TCA TCA ATT
 1251 1260 1269 1278 1287 1296
 ACC TGG AGT GGT AGG TAA AGG GCT GCC TTA TGG TGT GGC GGG AGG AGT AGT TAA

 Thr Trp Ser Gly Arg *** Arg Ala Ala Leu Trp Cys Gly Gly Arg Ser Ser ***
 Pro Gly Val Val Gly Lys Gly Leu Pro Tyr Gly Val Ala Gly Gly Val Val Asn
 Leu Glu Trp *** Val Lys Gly Cys Leu Met Val Trp Arg Glu Glu *** Leu Ile

 Leu Pro *** Leu Gly Leu Gln His Leu Pro Asn Cys Leu Gln Cys Gly Leu Tyr
 Tyr Pro Asp Tyr Ala Leu Asn Thr Ser Pro Thr Val Phe Asn Ala Asp Leu Ile
 Ile Pro Thr Met Pro Trp Thr Pro Pro Pro Pro *** Leu Thr Pro Met Trp Ser

 ATA TCC CCA GTA TCC GGT TCA ACC ACC TCC CCC AAT GTT TCA ACC GTA GGT TCT
 1305 1314 1323 1332 1341 1350
 TAT AGG GGT CAT AGG CCA AGT TGG TGG AGG GGG TTA CAA AGT TGG CAT CCA AGA

 Tyr Arg Gly His Arg Pro Ser Trp Trp Arg Gly Leu Gln Ser Trp His Pro Arg
 Ile Gly Val Ile Gly Gln Val Gly Gly Gly Gly Tyr Lys Val Gly Ile Gln Asp
 *** Gly Ser *** Ala Lys Leu Val Glu Gly Val Thr Lys Leu Ala Ser Lys Ile

FIG. 2e

Cys Cys His Val Trp Cys Arg Lys Ser *** Leu His His Pro Arg Gln Pro Leu
 Val Val Thr Ser Gly Val Gly Arg Gln Asn Ser Thr Ile Pro Asp Arg Pro Tyr
 Leu Leu Leu Pro Gly Leu Val Glu Lys Ile Leu Pro Ser Pro Thr Glu Pro Thr

 ATT GTT GTC ACC TGG GTT GTG GAG AAA CTA ATC TCC ACT ACC CCA GAG ACC CCA
 1359 1368 1377 1386 1395 1404
 TAA CAA CAG TGG ACC CAA CAC CTC TTT GAT TAG AGG TGA TGG GGT CTC TGG GGT

 Gln Gln Trp Thr Gln His Leu Phe Asp *** Arg *** Trp Gly Leu Trp Gly
 Asn Asn Ser Gly Pro Asn Thr Ser Leu Ile Arg Gly Asp Gly Val Ser Gly Val
 Thr Thr Val Asp Pro Thr Pro Leu *** Leu Glu Val Met Gly Ser Leu Gly ***

Ile *** Ile *** Gly Lys *** Tyr Pro Leu Ile Pro Phe Thr Pro Thr Pro Pro
 Phe Glu Tyr Lys Ala Lys Arg Ile Arg Tyr Tyr Gln Phe Pro Leu Pro Leu Pro
 Phe Asn Met Asn Leu Arg Glu Leu Val Thr Thr Asn Ser Leu Tyr Pro Tyr Pro

 TTT TAA GTA TAA ATC GGA AAG ATT ATG CCA TCA TAA CCT TTC CAT CCC CAT CCC
 1413 1422 1431 1440 1449 1458
 AAA ATT CAT ATT TAG CCT TTC TAA TAC GGT AGT ATT GGA AAG GTA GGG GTA GGG

 Lys Ile His Ile *** Pro Phe *** Tyr Gly Ser Ile Gly Lys Val Gly Val Gly
 Lys Phe Ile Phe Ser Leu Ser Asn Thr Val Val Leu Glu Arg *** Gly *** Gly
 Asn Ser Tyr Leu Ala Phe Leu Ile Arg *** Tyr Trp Lys Gly Arg Gly Arg Gly

Gln His Arg Arg Leu Pro Pro Pro Val Pro Arg His Gln Ile Glu Ala Arg ***
 Asn Thr Gly Gly Ser Pro Pro Leu Phe Gln Gly Ile Asn Phe Arg Leu Glu Asn
 Thr Pro Ala Ala Gln Pro Pro Ser Ser Ser Ala Ser Thr Ser Asp *** Ser Thr

 CCA ACC ACG GCG GAC TCC CCC CCT CCT TGA CCG GCT ACA ACT TAG AGT CGA GCA
 1467 1476 1485 1494 1503 1512
 GGT TGG TGC CGC CTG AGG GGG GGA GGA ACT GGC CGA TGT TGA ATC TCA GCT CGT

 Gly Trp Cys Arg Leu Arg Gly Gly Gly Thr Gly Arg Cys *** Ile Ser Ala Arg
 Val Gly Ala Ala *** Gly Gly Glu Glu Leu Ala Asp Val Glu Ser Gln Leu Val
 Leu Val Pro Pro Glu Gly Gly Arg Asn Trp Pro Met Leu Asn Leu Ser Ser Leu

Cys Glu Leu Ile Ala Ala Leu Thr Arg Arg Lys His His Thr Cys Ile Arg ***
 Val Asn Trp Ser Pro Gln Ser His Gly Gly Arg Ile Thr Leu Val Phe Glu Arg
 Leu Met Gly Leu His Ser Arg Thr Asp Glu Glu *** Pro Ser Tyr Leu Asn Glu

 ATT GTA AGG TTC TAC CGA CGC TCA CAG GAG GAG AAT ACC ACT CAT GTT TAA GAG
 1521 1530 1539 1548 1557 1566
 TAA CAT TCC AAG ATG GCT GCG AGT GTC CTC CTC TTA TGG TGA GTA CAA ATT CTC

 *** His Ser Lys Met Ala Ala Ser Val Leu Leu Leu Trp *** Val Gln Ile Leu
 Asn Ile Pro Arg Trp Leu Arg Val Ser Ser Ser Tyr Gly Glu Tyr Lys Phe Ser
 Thr Phe Gln Asp Gly Cys Glu Cys Pro Pro Leu Met Val Ser Thr Asn Ser Leu

Phe Pro Pro Phe Gln Leu Tyr Gly Asp Lys Pro Ala Met Gln Leu Pro Lys Gln
 Ser Leu Arg Ser Asn Phe Ile Gly Thr Lys Arg Arg Trp Arg Tyr Arg Asn Arg
 Leu Phe Ala Pro Ile Ser Ser Val Arg Arg Glu Ala Gly Asp Thr Val Thr Glu

 ATC TTT CCG CCC TTA ACT TCT ATG GGC AGA AAG CCG CGG TAG ACA TTG CCA AAG
 1575 1584 1593 1602 1611 1620
 TAG AAA GGC GGG AAT TGA AGA TAC CCG TCT TTC GGC GCC ATC TGT AAC GGT TTC

 *** Lys Gly Gly Asn *** Arg Tyr Pro Ser Phe Gly Ala Ile Cys Asn Gly Phe
 Arg Lys Ala Gly Ile Glu Asp Thr Arg Leu Ser Ala Pro Ser Val Thr Val Ser
 Glu Arg Arg Glu Leu Lys Ile Pro Val Phe Arg Arg His Leu *** Arg Phe Leu

FIG. 2f

Leu Arg Pro Thr Gly Phe Ile Thr Lys Glu Pro Pro His Lys Trp Ser Pro Gln
 Phe Ala Pro His Val Leu Tyr Pro Arg Arg Arg Leu Ile Asn Gly Leu His Ser
 Ser Pro Pro Thr Tyr Trp Ile His Asp Glu Gly Ser Ser Thr Glu Leu Ile Ala

 ACT TCC GCC CCA CAT GGT TTA TAC CAG AAG AGG CCT CCT ACA AAG GTT CTA CCG
 1629 1638 1647 1656 1665 1674
 TGA AGG CCG GGT GTA CCA AAT ATG GTC TTC TCC GGA GGA TGT TTC CAA GAT GGC

 *** Arg Arg Gly Val Pro Asn Met Val Phe Ser Gly Gly Cys Phe Gln Asp Gly
 Glu Gly Gly Val Tyr Gln Ile Trp Ser Ser Pro Glu Asp Val Ser Lys Met Ala
 Lys Ala Gly Cys Thr Lys Tyr Gly Leu Leu Arg Arg Met Phe Pro Arg Trp Leu

 Pro Pro Pro Asp Thr Lys Gln Pro Leu Ala Glu Lys Ala Val Asp Asp *** Leu
 Arg Pro Arg Thr Arg Arg Arg Arg Tyr Arg Arg Arg Pro Trp Thr Met Arg Tyr
 Ala Pro Ala Pro Gly Asp Glu Ala Thr Val Gly Gly Gln Gly Arg *** Gly Ile

 ACG CCC CCG CCC AGG CAG AAG ACG CCA TTG CGG AGG AAC CCG TGC AGT AGG ATA
 1683 1692 1701 1710 1719 1728
 TGC GGG GGC GGG TCC GTC TTC TGC GGT AAC GCC TCC TTG GCC ACG TCA TCC TAT

 Cys Gly Gly Gly Ser Val Phe Cys Gly Asn Ala Ser Leu Ala Thr Ser Ser Tyr
 Ala Gly Ala Gly Pro Ser Ser Ala Val Thr Pro Pro Trp Pro Arg His Pro Ile
 Arg Gly Arg Val Arg Leu Leu Arg *** Arg Leu Leu Gly His Val Ile Leu ***

 Leu Ser Leu Leu Ala Ser Ser Tyr Tyr
 Phe His Phe Phe His Ala Ala Thr Thr Asn
 Phe Thr Phe Ser Thr Arg Gln Gln Leu Ile

 TTT TCA CTT TCT TCA CGC GAC GAC ATC ATA A 5'
 1737 1746 1755
 AAA AGT GAA AGA AGT GCG CTG CTG TAG TAT T 3'

 Lys Ser Glu Arg Ser Ala Leu Leu *** Tyr
 Lys Val Lys Glu Val Arg Cys Cys Ser Ile
 Lys *** Lys Lys Cys Cys Ala Ala Val Val

FIG. 2g

circopormank	1	10	20	30	40	50	50
circopormeeh	1	10	20	30	40	50	50
circopordfp	1	10	20	30	40	50	50
circopormank	51	60	70	80	90	100	100
circopormeeh	51	60	70	80	90	100	100
circopordfp	51	60	70	80	90	100	100
circopormank	101	110	120	130	140	150	150
circopormeeh	101	110	120	130	140	150	150
circopordfp	101	110	120	130	140	150	150
circopormank	151	160	170	180	190	200	200
circopormeeh	151	160	170	180	190	200	200
circopordfp	151	160	170	180	190	200	200
circopormank	201	210	220	230	240	250	250
circopormeeh	201	210	220	230	240	250	250
circopordfp	201	210	220	230	240	250	250
circopormank	251	260	270	280	290	300	300
circopormeeh	251	260	270	280	290	300	300
circopordfp	251	260	270	280	290	300	300
circopormank	301	310	320	330	340	350	350
circopormeeh	301	310	320	330	340	350	350
circopordfp	301	310	320	330	340	350	350
circopormank	351	360	370	380	390	400	400
circopormeeh	351	360	370	380	390	400	400
circopordfp	351	360	370	380	390	400	400
circopormank	401	410	420	430	440	450	450
circopormeeh	401	410	420	430	440	450	450
circopordfp	401	410	420	430	440	450	450
circopormank	451	460	470	480	490	500	500
circopormeeh	451	460	470	480	490	500	500
circopordfp	451	460	470	480	490	500	500
circopormank	501	510	520	530	540	550	550
circopormeeh	501	510	520	530	540	550	550
circopordfp	501	510	520	530	540	550	550
circopormank	551	560	570	580	590	600	600
circopormeeh	551	560	570	580	590	600	600
circopordfp	551	560	570	580	590	600	600

FIG. 3a

circopormank	601	610	620	630	640	650	
circopormeeh	601	610	620	630	640	650	650
circopordfp	601	610	620	630	640	650	650
		660	670	680	690	700	
circopormank	651	660	670	680	690	700	700
circopormeeh	651	660	670	680	690	700	700
circopordfp	651	660	670	680	690	700	700
		710	720	730	740	750	
circopormank	701	710	720	730	740	750	750
circopormeeh	701	710	720	730	740	750	750
circopordfp	701	710	720	730	740	750	750
		760	770	780	790	800	
circopormank	751	760	770	780	790	800	800
circopormeeh	751	760	770	780	790	800	800
circopordfp	751	760	770	780	790	800	800
		810	820	830	840	850	
circopormank	801	810	820	830	840	850	850
circopormeeh	801	810	820	830	840	850	850
circopordfp	801	810	820	830	840	850	850
		860	870	880	890	900	
circopormank	851	860	870	880	890	900	900
circopormeeh	851	860	870	880	890	900	900
circopordfp	851	860	870	880	890	900	900
		910	920	930	940	950	
circopormank	901	910	920	930	940	950	950
circopormeeh	901	910	920	930	940	950	950
circopordfp	901	910	920	930	940	950	950
		960	970	980	990	1000	
circopormank	951	960	970	980	990	1000	1000
circopormeeh	951	960	970	980	990	1000	1000
circopordfp	951	960	970	980	990	1000	1000
		1010	1020	1030	1040	1050	
circopormank	1001	1010	1020	1030	1040	1050	1050
circopormeeh	1001	1010	1020	1030	1040	1050	1050
circopordfp	1001	1010	1020	1030	1040	1050	1050
		1060	1070	1080	1090	1100	
circopormank	1051	1060	1070	1080	1090	1100	1100
circopormeeh	1051	1060	1070	1080	1090	1100	1100
circopordfp	1051	1060	1070	1080	1090	1100	1100
		1110	1120	1130	1140	1150	
circopormank	1101	1110	1120	1130	1140	1150	1150
circopormeeh	1101	1110	1120	1130	1140	1150	1150
circopordfp	1101	1110	1120	1130	1140	1150	1150
		1160	1170	1180	1190	1200	
circopormank	1151	1160	1170	1180	1190	1200	1200
circopormeeh	1151	1160	1170	1180	1190	1200	1200
circopordfp	1151	1160	1170	1180	1190	1200	1200

FIG. 3b

		1210	1220	1230	1240	1250	
circopormank	1201	ACCAATCAAT	TGTTTGGTCC	AGCTCAGGTT	TGGGGGTGAA	GTACCTGGAG	1250
circopormeeh	1201	ACCAATCAAT	TGTTTGGTCC	AGCTCAGGTT	TGGGGGTGAA	GTACCTGGAG	1250
circopordfp	1201	ACCAATCAAT	TGTTTGGTCT	AGCTCTGGTT	TGGGGGTGAA	GTACCTGGAG	1250
		1260	1270	1280	1290	1300	
circopormank	1251	TGGTAGGTAA	AGGGCTGCCT	TATGGTGTGG	CGGGAGGAGT	AGTTAATATA	1300
circopormeeh	1251	TGGTAGGTAA	AGGGCTGCCT	TATGGTGTGG	CGGGAGGAGT	AGTTAATATA	1300
circopordfp	1251	TGGTAGGTAA	AGGGCTGCCT	TATGGTGTGG	CGGGAGGAGT	AGTTAATATA	1300
		1310	1320	1330	1340	1350	
circopormank	1301	GGGTCATAG	GCCAAGTTGG	TGGAGGGGGT	TACAAAGTTG	GCATCCAAGA	1350
circopormeeh	1301	GGGTCATAG	GCCAAGTTGG	TGGAGGGGGT	TACAAAGTTG	GCATCCAAGA	1350
circopordfp	1301	GGGTCATAG	GCCAAGTTGG	TGGAGGGGGT	TACAAAGTTG	GCATCCAAGA	1350
		1360	1370	1380	1390	1400	
circopormank	1351	TAACAACAGT	GGACCCAACA	CCTCTTTGAT	TAGAGGTGAT	GGGGTCTCTG	1400
circopormeeh	1351	TAACAACAGT	GGACCCAACA	CCTCTTTGAT	TAGAGGTGAT	GGGGTCTCTG	1400
circopordfp	1351	TAACAACAGT	GGACCCAACA	CCTCTTTGAT	TAGAGGTGAT	GGGGTCTCTG	1400
		1410	1420	1430	1440	1450	
circopormank	1401	GGGTAAAATT	CATATTTAGC	CTTCTAATA	CGGTAGTATT	GGAAAGGTAG	1450
circopormeeh	1401	GGGTAAAATT	CATATTTAGC	CTTCTAATA	CGGTAGTATT	GGAAAGGTAG	1450
circopordfp	1401	GGGTAAAATT	CATATTTAGC	CTTCTAATA	CGGTAGTATT	GGAAAGGTAG	1450
		1460	1470	1480	1490	1500	
circopormank	1451	GGGTAGGGGG	TTGGTGCCCG	CTGAGGGGGG	GAGGAAC TGG	CCGATGTTGA	1500
circopormeeh	1451	GGGTAGGGGG	TTGGTGCCCG	CTGAGGGGGG	GAGGAAC TGG	CCGATGTTGA	1500
circopordfp	1451	GGGTAGGGGG	TTGGTGCCCG	CTGAGGGGGG	GAGGAAC TGG	CCGATGTTGA	1500
		1510	1520	1530	1540	1550	
circopormank	1501	ATCTGAGGTG	GTTAACATTC	CAAGATGGCT	GGGAGTATCC	TCCTTTTATG	1550
circopormeeh	1501	ATCTGAGGTG	GTTAACATTC	CAAGATGGCT	GGGAGTATCC	TCCTTTTATG	1550
circopordfp	1501	ATCTGAGGTC	GTTAACATTC	CAAGATGGCT	GGGAGTATCC	TCCTTTTATG	1550
		1560	1570	1580	1590	1600	
circopormank	1551	GTGAGTACAA	ATTCTCTAGA	AAGCGGGGAA	TTGAAGATAC	CCGTCTTTCC	1600
circopormeeh	1551	GTGAGTACAA	ATTCTCTAGA	AAGCGGGGAA	TTGAAGATAC	CCGTCTTTCC	1600
circopordfp	1551	GTGAGTACAA	ATTCTCTAGA	AAGCGGGGAA	TTGAAGATAC	CCGTCTTTCC	1600
		1610	1620	1630	1640	1650	
circopormank	1601	GGGCCATCTG	TAACGGTTTC	TGAAGCGGGG	GTGTGCCAAA	TATGGTCTTC	1650
circopormeeh	1601	GGGCCATCTG	TAACGGTTTC	TGAAGCGGGG	GTGTGCCAAA	TATGGTCTTC	1650
circopordfp	1601	GGGCCATCTG	TAACGGTTTC	TGAAGCGGGG	GTGTGCCAAA	TATGGTCTTC	1650
		1660	1670	1680	1690	1700	
circopormank	1651	TCCGGAGGAT	GTTTCCAAGA	TGGCTGCCGG	GGCGGGTCCG	TCTTCTGCGG	1700
circopormeeh	1651	TCCGGAGGAT	GTTTCCAAGA	TGGCTGCCGG	GGCGGGTCCG	TCTTCTGCGG	1700
circopordfp	1651	TCCGGAGGAT	GTTTCCAAGA	TGGCTGCCGG	GGCGGGTCCG	TCTTCTGCGG	1700
		1710	1720	1730	1740	1750	
circopormank	1701	TAACGGCTCC	TGGCCACGT	CATCCTATAA	AAGTCAAAGA	AGTGGCGTGC	1750
circopormeeh	1701	TAACGGCTCC	TGGCCACGT	CATCCTATAA	AAGTCAAAGA	AGTGGCGTGC	1750
circopordfp	1701	TAACGGCTCC	TGGCCACGT	CATCCTATAA	AAGTCAAAGA	AGTGGCGTGC	1750
		1760	1770	1780	1790	1800	
circopormank	1751	TGTAGTATT	1800
circopormeeh	1751	TGTAGTATT	1800
circopordfp	1751	TGTAGTATT	1800

FIG. 3c

		10	20	30	40	50	
circopormank	1	NPSKKSQP	HKRWVFTLNN	PSBEEKNKIR	ELPISLFDYF	VCGEEGLEEG	50
circopormeeh	1	NPSKKSQP	HKRWVFTLNN	PSBEEKNKIR	ELPISLFDYF	VCGEEGLEEG	50
circopordfp{	1	NPSKKSQP	HKRWVFTLNN	PSBEEKNKIR	ELPISLFDYF	VCGEEGLEEG	50
		60	70	80	90	100	
circopormank	51	RTPHLOGFAN	FAKKOTFNKV	KWYFGARCHI	EKAKGTDQON	KBYCSKEGHI	100
circopormeeh	51	RTPHLOGFAN	FAKKOTFNKV	KWYFGARCHI	EKAKGTDQON	KBYCSKEGHI	100
circopordfp{	51	RTPHLOGFAN	FAKKOTFNKV	KWYFGARCHI	EKAKGTDQON	KBYCSKEGHI	100
		110	120	130	140	150	
circopormank	101	LIECGAPRNO	GKRSDELSTAV	STLLETGSLV	TVAEQFPVY	VRNFRGLAEL	150
circopormeeh	101	LIECGAPRNO	GKRSDELSTAV	STLLETGSLV	TVAEQFPVY	VRNFRGLAEL	150
circopordfp{	101	LIECGAPRNO	GKRSDELSTAV	STLLETGSLV	TVAEQFPVY	VRNFRGLAEL	150
		160	170	180	190	200	
circopormank	151	LKVSCKMOOR	DWKTAVHVIV	GPPGCGKSOW	ARNFAEPROT	YWKPSRNKWH	200
circopormeeh	151	LKVSCKMOOR	DWKTAVHVIV	GPPGCGKSOW	ARNFAEPROT	YWKPSRNKWH	200
circopordfp{	151	LKVSCKMOOR	DWKTAVHVIV	GPPGCGKSOW	ARNFAEPROT	YWKPSRNKWH	200
		210	220	230	240	250	
circopormank	201	DGYHGEEVVV	LDDFYGWLFW	DDLLRLCDRY	PLTVETKGGT	VPFLARSILI	250
circopormeeh	201	DGYHGEEVVV	LDDFYGWLFW	DDLLRLCDRY	PLTVETKGGT	VPFLARSILI	250
circopordfp{	201	DGYHGEEVVV	LDDFYGWLFW	DDLLRLCDRY	PLTVETKGGT	VPFLARSILI	250
		260	270	280	290	300	
circopormank	251	TSNOAPQEWY	SSTAVPAVEA	LYRRITTLQF	WKTAGEQSTE	VPEGRFEAVD	300
circopormeeh	251	TSNOAPQEWY	SSTAVPAVEA	LYRRITTLQF	WKTAGEQSTE	VPEGRFEAVD	300
circopordfp{	251	TSNOAPQEWY	SSTAVPAVEA	LYRRITTLQF	WKTAGEQSTE	VPEGRFEAVD	300
		310	320	330	340	350	
circopormank	301	PPCALFPYKI	NY	350
circopormeeh	301	PPCALFPYKI	NY	350
circopordfp{	301	PPCALFPYKI	NY	350

FIG. 4

		10	20	30	40	50	
circopormank	1	MTWPRRRYRR	RRTRPRSHLG	NILRRRPYLA	HPAFRRRYRW	RRKTGIFNCR	50
circopormeeh	1	MTWPRRRYRR	RRTRPRSHLG	NILRRRPYLA	HPAFRRRYRW	RRKTGIFNSR	50
circopordfp[1	MTWPRRRYRR	RRTRPRSHLG	NILRRRPYLV	HPAFRRRYRW	RRKTGIFNSR	50
		60	70	80	90	100	
circopormank	51	LSKEFVLIFFK	GGYSQPSWIV	NILRFNIGOF	LPPSGGTNPE	PLPFOYYRIR	100
circopormeeh	51	LSKEFVLIFFK	GGYSQPSWIV	NILRFNIGOF	LPPSGGTNPE	PLPFOYYRIR	100
circopordfp[51	LSREFVLTIR	GGISQPSWIV	NILRFNIGOF	LPPSGGTNPE	PLPFOYYRIR	100
		110	120	130	140	150	
circopormank	101	KAKYEFYPRD	PITSNORGVG	STVVILDANE	VTPSTNLAYD	PYINYSSRHT	150
circopormeeh	101	KAKYEFYPRD	PITSNORGVG	STVVILDANE	VTPSTNLAYD	PYINYSSRHT	150
circopordfp[101	KAKYEFYPRD	PITSNORGVG	STVVILDANE	VTPSTNLAYD	PYINYSSRHT	150
		160	170	180	190	200	
circopormank	151	IRQPFTYHSR	YFTPKEPDDQ	TIDWFHPNPK	RNQLWLHLNT	HTNVEHTGLG	200
circopormeeh	151	IRQPFTYHSR	YFTPKEPDDQ	TIDWFHPNPK	RNQLWLHLNT	HTNVEHTGLG	200
circopordfp[151	IRQPFTYHSR	YFTPKEPDDQ	TIDWFQPNPK	RNQLWLHLNT	HTNVEHTGLG	200
		210	220	230	240	250	
circopormank	201	YALONAATAQ	NYVVRTIYV	QREFILKDP	LNK*	250
circopormeeh	201	YALONAATAQ	NYVVRTIYV	QREFILKDP	LNK*	250
circopordfp[201	YALONAATAQ	NYVVRTIYV	QREFILKDP	LNE*	250

FIG. 5

		10	20	30	40	50	
circopormank	1	MISIPPLIST	RLPVGVARELS	KITGPLEALPT	TGRAHYDVYS	CLPITLLHLP	50
circopormeeh	1	MISIPPLIST	RLPVGVPRLS	KITGPLEALPT	TGRAHYDVYS	CLPITLLHLP	50
circopordfp[1	MISIPPLIST	RLPVGVPRLS	KITGPLEALPT	TGRAHYDVYS	CLPITLLHLP	50
		60	70	80	90	100	
circopormank	51	AHFQKFSOPA	EISHIRYREL	LGYSHORPRE	OKGTHSSROV	AALPLVPRSS	100
circopormeeh	51	AHFQKFSOPA	EISHIRYREL	LGYSHORPRE	OKGTHSSROV	AAEPLVPRSS	100
circopordfp[51	AHFQKFSOPA	EISHIRYRKL	LGYSHORPRE	OKGTHSSROV	AALPLVPRSS	100
		110	120	130	140	150	
circopormank	101	TLDKYVAFFT	AVFFILLVGS	FRFLDVAAGT	KIPLHLVKSL	LLSKISKPLE	150
circopormeeh	101	TLDKYVAFFT	AVFFILLVGS	FRFLDVAAGT	KIPLHLVKSL	LLSKIRKPLE	150
circopordfp[101	TLDKYVAFFT	AVFFILLVGS	FRFLDVAAGT	KIPLHLVKSL	LLSKIRKPLE	150
		160	170	180	190	200	
circopormank	151	VRSSTLFQTF	LSANKLIKKG	DWKLPEYVFL	LLGRIIKGEH	PPLMGERAAF	200
circopormeeh	151	VRSSTLFQTF	LSANKLIKKG	DWKLPEYVFL	LLGRIIKGEH	PPLMGLRAAF	200
circopordfp[151	VRSSTLFQTF	LATNKLIKKG	DWKLPEYVFL	LLGRIIKGEH	PPLMGERAAF	200
		210	220	230	240	250	
circopormank	201	LAWHFF*	250
circopormeeh	201	LAWHFFH	250
circopordfp[201	LAWHFF-	250

FIG. 6




FIG. 7

Leu Ala Ser Arg Cys Arg Cys Cys Arg Pro Leu Val Glu Ala Ala Val His Gly
 Trp Arg Val Glu Ala Ala Ala Ala Gly Arg Cys Cys Arg Leu Leu Leu Met Gly
 Gly Ala Cys Lys Pro Leu Pro Leu Val Glu Ala Ala Gly *** Cys Cys Cys Ala
 3' TGG TCG CGT GAA GCC GTC GCC GTC GTG GAG CCG TCG TGG AGT CGT CGT TGT ACG
 9 18 27 36 45 54
 5' ACC AGC GCA CTT CGG CAG CGG CAG CAC CTC GGC AGC ACC TCA GCA GCA ACA TGC
 Thr Ser Ala Leu Arg Gln Arg Gln His Leu Gly Ser Thr Ser Ala Ala Thr Cys
 Pro Ala His Phe Gly Ser Gly Ser Thr Ser Ala Ala Pro Gln Gln Gln His Ala
 Gln Arg Thr Ser Ala Ala Ala Ala Pro Arg Gln His Leu Ser Ser Asn Met Pro
 Ala Leu Leu Ile Ser Ser Ala Ser Gly Leu Gly Met Phe Pro Pro His Glu Ser
 Leu Leu Phe Phe Pro Leu Leu Pro Gly Trp Gly Trp Leu Leu His Thr Asn Val
 Trp Cys Ser Ser His Phe Phe Arg Val Gly Val Gly Tyr Phe Thr Pro Thr ***
 GGT CGT TCT TCT TAC CTT CTT CGC CTG GGG TTG GGG TAT TTT CCA CCC ACA AGT
 63 72 81 90 99 108
 CCA GCA AGA AGA ATG GAA GAA GCG GAC CCC AAC CCC ATA AAA GGT GGG TGT TCA
 Pro Ala Arg Arg Met Glu Glu Ala Asp Pro Asn Pro Ile Lys Gly Gly Cys Ser
 Gln Gln Glu Glu Trp Lys Lys Arg Thr Pro Thr Pro *** Lys Val Gly Val His
 Ser Lys Lys Asn Gly Arg Ser Gly Pro Gln Pro His Lys Arg Trp Val Phe Thr
 Gln Ile Ile Arg Gly Phe Val Leu Ala Leu Phe Tyr Pro Ile Lys Trp Tyr Gly
 Arg Phe Leu Gly Glu Ser Ser Ser Arg Leu Phe Ile Arg Ser Arg Gly Ile Asp
 Glu Ser Tyr Asp Lys Arg Leu Arg Ala Cys Ser Phe Val Pro Asp Glu Leu Ile
 GAG ACT TAT TAG GAA GGC TTC TGC TCG CGT TCT TTT ATG CCC TAG AAG GTT ATA
 117 126 135 144 153 162
 CTC TGA ATA ATC CTT CCG AAG ACG AGC GCA AGA AAA TAC GGG ATC TTC CAA TAT
 Leu *** Ile Ile Leu Pro Lys Thr Ser Ala Arg Lys Tyr Gly Ile Phe Gln Tyr
 Ser Glu *** Ser Phe Arg Arg Arg Ala Gln Glu Asn Thr Gly Ser Ser Asn Ile
 Leu Asn Asn Pro Ser Glu Asp Glu Arg Lys Lys Ile Arg Asp Leu Pro Ile Ser
 *** Lys Ile Ile Lys Asn Asn Ala Leu Leu Thr Ile Leu Phe Ser Ser Cys Arg
 Arg Asn Ser *** Lys Ile Thr Pro Ser Ser Pro Leu Ser Ser Pro Arg Val Gly
 Gly Ile Gln Asn Asn *** Gln Gln Arg Pro Pro Tyr His Pro Leu Val Phe Val
 GGG ATA AAC TAA TAA AAT AAC AAC CGC TCC TCC CAT TAC TCC TTC CTG CTT GTG
 171 180 189 198 207 216
 CCC TAT TTG ATT ATT TTA TTG TTG GCG AGG AGG GTA ATG AGG AAG GAC GAA CAC
 Pro Tyr Leu Ile Ile Leu Leu Leu Ala Arg Arg Val Met Arg Lys Asp Glu His
 Pro Ile *** Leu Phe Tyr Cys Trp Arg Gly Gly *** *** Gly Arg Thr Asn Thr
 Leu Phe Asp Tyr Phe Ile Val Gly Glu Glu Gly Asn Glu Glu Gly Arg Thr Pro
 Val Glu Leu Pro Glu Ser Ile Lys His Leu Leu Leu Ser Lys Ile Phe His Leu
 *** Arg Trp Pro Asn Ala Leu Lys Thr Phe Phe Cys Val Lys Leu Leu Thr Phe
 Glu Gly Gly Pro Thr Arg *** Asn Gln Ser Ser Ala Ser Lys *** Tyr Leu Ser
 GAG TGG AGG TCC CCA AGC GAT TAA AAC ACT TCT TCG TCT GAA AAT TAT TTC ACT
 225 234 243 252 261 270
 CTC ACC TCC AGG GGT TCG CTA ATT TTG TGA AGA AGC AGA CTT TTA ATA AAG TGA
 Leu Thr Ser Arg Gly Ser Leu Ile Leu *** Arg Ser Arg Leu Leu Ile Lys ***
 Ser Pro Pro Gly Val Arg *** Phe Cys Glu Glu Ala Asp Phe *** *** Ser Glu
 His Leu Gln Gly Phe Ala Asn Phe Val Lys Lys Gln Thr Phe Asn Lys Val Lys

FIG. 8a

Pro Ile Gln Thr Gly Ala Ala Val Asp Leu Phe Arg Phe Ser Cys Ile Leu Leu
 His Tyr Lys Pro Ala Arg Gln Trp Met Ser Phe Ala Phe Pro Val Ser *** Cys
 Thr Thr Asn Pro His Gly Ser Gly Cys Arg Ser Leu Ser Leu Phe Leu Asp Ala
 TCA CCA TAA ACC CAC GGG CGA CGG TGT AGC TCT TTC GCT TTC CTT GTC TAG TCG
 279 288 297 306 315 324
 AGT GGT ATT TGG GTG CCC GCT GCC ACA TCG AGA AAG CGA AAG GAA CAG ATC AGC
 Ser Gly Ile Trp Val Pro Ala Ala Thr Ser Arg Lys Arg Lys Glu Gln Ile Ser
 Val Val Phe Gly Cys Pro Leu Pro His Arg Glu Ser Glu Arg Asn Arg Ser Ala
 Trp Tyr Leu Gly Ala Arg Cys His Ile Glu Lys Ala Lys Gly Thr Asp Gln Gln

Ile Phe Phe Val Ala Thr Phe Phe Ala Val *** Gln His Leu Thr Ser Ser Arg
 Phe Leu Ser Tyr Gln Leu Leu Ser Pro Leu Lys Ser His Ser His Pro Ala Gly
 Ser Tyr Leu Ile Ser Cys Tyr Leu Leu Cys Ser Val Ser Pro Thr His Leu Glu
 TCT TAT TTC TTA TGA CGT CAT TTC TTC CGT TGA ATG ACT ACC TCA CAC CTC GAG
 333 342 351 360 369 378
 AGA ATA AAG AAT ACT GCA GTA AAG AAG GCA ACT TAC TGA TGG AGT GTG GAG CTC
 Arg Ile Lys Asn Thr Ala Val Lys Lys Ala Thr Tyr *** Trp Ser Val Glu Leu
 Glu *** Arg Ile Leu Gln *** Arg Arg Gln Leu Thr Asp Gly Val Trp Ser Ser
 Asn Lys Glu Tyr Cys Ser Lys Glu Gly Asn Leu Leu Met Glu Cys Gly Ala Pro

Ser Arg Leu Ser Leu Pro Thr Val Gln Arg Ser Ser His Thr Gly Gln Gln Leu
 Leu Asp *** Pro Cys Arg Leu Ser Arg Asp Val Ala Thr Leu Val Lys Asn Ser
 *** Ile Glu Pro Val Val Ser His Gly Thr *** Gln Gln Ser Tyr Arg Thr Pro
 GAT CTA GAG TCC CTG TTG CCT CAC TGG ACA GAT GAC GAC ACT CAT GGA ACA ACC
 387 396 405 414 423 432
 CTA GAT CTC AGG GAC AAC GGA GTG ACC TGT CTA CTG CTG TGA GTA CCT TGT TGG
 Leu Asp Leu Arg Asp Asn Gly Val Thr Cys Leu Leu Leu *** Val Pro Cys Trp
 *** Ile Ser Gly Thr Thr Glu *** Pro Val Tyr Cys Cys Glu Tyr Leu Val Gly
 Arg Ser Gln Gly Gln Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu

Ala Pro Thr Gln His Gly Asn Cys Leu Leu Val Arg Tyr Arg Lys Asp Ser Ile
 Leu Pro Leu Arg Thr Val Thr Ala Ser Cys Cys Gly Thr Val Asn Thr Leu Phe
 Ser Arg Ser Asp Pro Ser Arg Gln Leu Ala Ala Gly Gln Leu Thr Gln *** Phe
 TCT CGC CCT CAG ACC ACT GGC AAC GTC TCG TCG TGG GAC ATT GCA AAC AGT CTT
 441 450 459 468 477 486
 AGA GCG GGA GTC TGG TGA CCG TTG CAG AGC AGC ACC CTG TAA CGT TTG TCA GAA
 Arg Ala Gly Val Trp *** Pro Leu Gln Ser Ser Thr Leu *** Arg Leu Ser Glu
 Glu Arg Glu Ser Gly Asp Arg Cys Arg Ala Ala Pro Cys Asn Val Cys Gln Lys
 Ser Gly Ser Leu Val Thr Val Ala Glu Gln His Pro Val Thr Phe Val Arg Asn

Glu Ala Pro Gln Ser Phe Lys Gln Phe His Ala Pro Phe His Leu Leu Thr Ile
 Lys Arg Pro Ser Ala Ser Ser Lys Phe Thr Leu Pro Phe Ile Cys Phe Arg Ser
 Asn Gly Arg Ala Pro Gln Val Lys Ser Leu Ser Arg Ser Phe Ala Ser Ala His
 TAA AGG CGC CCG ACC GAC TTG AAA ACT TTC ACT CGC CCT TTT ACG TCT TCG CAC
 495 504 513 522 531 540
 ATT TCC GCG GGC TGG CTG AAC TTT TGA AAG TGA GCG GGA AAA TGC AGA AGC GTG
 Ile Ser Ala Gly Trp Leu Asn Phe *** Lys *** Ala Gly Lys Cys Arg Ser Val
 Phe Pro Arg Ala Gly *** Thr Phe Glu Ser Glu Arg Glu Asn Ala Glu Ala ***
 Phe Arg Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met Gln Lys Arg Asp

FIG. 8b

Pro Leu Ser Ile Tyr Val Asp Asn His Pro Trp Arg Pro Thr Thr Phe Ala Phe
 Gln Phe Val Leu Thr Cys Thr Met Thr Pro Gly Gly Pro His Pro Leu Leu Leu
 Asn Ser Ser *** His Val Arg *** Gln Pro Ala Val Gln Thr His Tyr Phe Cys
 TAA CCT TCT GAT TAC ATG TGC AGT AAC ACC CCG GTG GAC CCA CAC CAT TTT CGT
 549 558 567 576 585 594
 ATT GGA AGA CTA ATG TAC ACG TCA TTG TGG GGC CAC CTG GGT GTG GTA AAA GCA
 Ile Gly Arg Leu Met Tyr Thr Ser Leu Trp Gly His Leu Gly Val Val Lys Ala
 Leu Glu Asp *** Cys Thr Arg His Cys Gly Ala Thr Trp Val Trp *** Lys Gln
 Trp Lys Thr Asn Val His Val Ile Val Gly Pro Pro Gly Cys Gly Lys Ser Lys
 Pro Ser Ser Ile Lys Cys Val Arg Phe Gly Cys Val Pro Phe Trp Arg Ser Val
 His Ala Ala Leu Lys Ala Ser Gly Ser Val Val Tyr Gln Phe Gly Gly Leu Phe
 Ile Pro Gln *** Asn Gln Leu Gly Pro Phe Trp Met Ser Ser Val Val *** Phe
 TTA CCC GAC GAT TAA AAC GTC TGG GCC TTT GGT GTA TGA CCT TTG GTG GAT CTT
 603 612 621 630 639 648
 AAT GGG CTG CTA ATT TTG CAG ACC CGG AAA CCA CAT ACT GGA AAC CAC CTA GAA
 Asn Gly Leu Leu Ile Leu Gln Thr Arg Lys Pro His Thr Gly Asn His Leu Glu
 Met Gly Cys *** Phe Cys Arg Pro Gly Asn His Ile Leu Glu Thr Thr *** Lys
 Trp Ala Ala Asn Phe Ala Asp Pro Glu Thr Thr Tyr Trp Lys Pro Pro Arg Asn
 Leu Pro Pro Ile Thr Val Met Thr Phe Phe His Asn Asn Asn Ile Val Lys Ile
 Leu His His Ser Pro *** Trp Pro Ser Ser Thr Thr Thr Ile Ser Ser Lys ***
 Cys Thr Thr Pro His Asn Gly His His Leu Leu Pro Gln *** Gln His Ser Lys
 TGT TCA CCA CCC TAC CAA TGG TAC CAC TTC TTC ACC AAC AAT AAC TAC TGA AAA
 657 666 675 684 693 702
 ACA AGT GGT GGG ATG GTT ACC ATG GTG AAG AAG TGG TTG TTA TTG ATG ACT TTT
 Thr Ser Gly Gly Met Val Thr Met Val Lys Lys Trp Leu Leu Leu Met Thr Phe
 Gln Val Val Gly Trp Leu Pro Trp *** Arg Ser Gly Cys Tyr *** *** Leu Leu
 Lys Trp Trp Asp Gly Tyr His Gly Glu Glu Val Val Val Ile Asp Asp Phe Tyr
 Ala Pro Gln Gly Pro Ile Ile *** Gln Ser Gln Thr Ile Ser Ile Trp Gln Ser
 Pro Gln Ser Gly Gln Ser Ser Arg Ser Leu Ser His Ser Arg Tyr Gly Asn Val
 His Ser Ala Ala Arg Pro His Asp Val Ser Val Thr His Asp Ile Asp Met Ser
 TAC CGA CCG ACG GGA CCC TAC TAG ATG ACT CTG ACA CAC TAG CTA TAG GTA ACT
 711 720 729 738 747 756
 ATG GCT GGC TGC CCT GGG ATG ATC TAC TGA GAC TGT GTG ATC GAT ATC CAT TGA
 Met Ala Gly Cys Pro Gly Met Ile Tyr *** Asp Cys Val Ile Asp Ile His ***
 Trp Leu Ala Ala Leu Gly *** Ser Thr Glu Thr Val *** Ser Ile Ser Ile Asp
 Gly Trp Leu Pro Trp Asp Asp Leu Leu Arg Leu Cys Asp Arg Tyr Pro Leu Thr
 Tyr Leu Ser Phe Thr Ser Ser Tyr Arg Lys Gln Gly Ala Thr Asn Gln Asn Gly
 Thr Ser Val Leu Pro Pro Val Thr Gly Lys Lys Ala Arg Leu Ile Arg Ile Val
 Gln Leu Ser *** Leu His Phe Gln Val Lys Lys Pro Gly Cys Tyr Glu Ser ***
 GAC ATC TCT GAT TTC CAC CTT GAC ATG GAA AAA ACC GGG CGT CAT AAG ACT AAT
 765 774 783 792 801 810
 CTG TAG AGA CTA AAG GTG GAA CTG TAC CTT TTT TGG CCC GCA GTA TTC TGA TTA
 Leu *** Arg Leu Lys Val Glu Leu Tyr Leu Phe Trp Pro Ala Val Phe *** Leu
 Cys Arg Asp *** Arg Trp Asn Cys Thr Phe Phe Gly Pro Gln Tyr Ser Asp Tyr
 Val Glu Thr Lys Gly Gly Thr Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr

FIG. 8c

Ala Ile Leu Gly Arg Gln Phe Pro Val Gly *** Ser Ser Asp Trp Ser Tyr Phe
 Leu Leu *** Val Gly Asn Ser His Tyr Glu Glu Val Ala Thr Gly Ala Thr Ser
 Trp Cys Asp Ser Gly Thr Pro Ile Thr Ser Arg Leu Gln Gln Gly Leu Gln Leu

 GGT CGT TAG TCT GGG GCA ACC TTA CCA TGA GGA GTT GAC GAC AGG GTC GAC ATC
 819 828 837 846 855 864
 CCA GCA ATC AGA CCC CGT TGG AAT GGT ACT CCT CAA CTG CTG TCC CAG CTG TAG

 Pro Ala Ile Arg Pro Arg Trp Asn Gly Thr Pro Gln Leu Leu Ser Gln Leu ***
 Gln Gln Ser Asp Pro Val Gly Met Val Leu Leu Asn Cys Cys Pro Ser Cys Arg
 Ser Asn Gln Thr Pro Leu Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu

Ser Lys Ile Pro Pro Asn Ser Gly Gln Tyr Lys Pro Leu Ile Ser Cys Phe Leu
 Ala Arg *** Arg Leu Ile Val Glu Lys Thr Asn Gln Phe Phe Ala Val Ser Cys
 Leu Glu Lys Asp Ser Ser *** Lys Arg Pro Ile Lys Ser Ser His *** Leu Val

 TTC GAG AAA TAG CCT CCT AAT GAA GGA ACC ATA AAA CCT TCT TAC GAT GTC TTG
 873 882 891 900 909 918
 AAG CTC TTT ATC GGA GGA TTA CTT CCT TGG TAT TTT GGA AGA ATG CTA CAG AAC

 Lys Leu Phe Ile Gly Gly Leu Leu Pro Trp Tyr Phe Gly Arg Met Leu Gln Asn
 Ser Ser Leu Ser Glu Asp Tyr Phe Leu Gly Ile Leu Glu Cys Tyr Arg Thr
 Ala Leu Tyr Arg Arg Ile Thr Ser Leu Val Phe Trp Lys Asn Ala Thr Glu Gln

Gly Arg Leu Phe Pro Ala Leu Glu Asp Gly Lys Gly Gly Trp Ala Arg Phe Lys
 Asp Val Ser Ser Pro Pro Trp Asn Thr Val Arg Glu Gly Gly His Gly Ser Asn
 Ile Trp Pro Pro Leu Pro Gly Thr Arg *** Gly Lys Gly Gly Met Gly Gln Ile

 TTA GGT GCC TCC TTC CCC CGG TCA AGC AGT GGG AAA GGG GGG GTA CGG GAC TTA
 927 936 945 954 963 972
 AAT CCA CGG AGG AAG GGG GCC AGT TCG TCA CCC TTT CCC CCC CAT GCC CTG AAT

 Asn Pro Arg Arg Lys Gly Ala Ser Ser Ser Pro Phe Pro Pro His Ala Leu Asn
 Ile His Gly Gly Arg Gly Pro Val Arg His Pro Phe Pro Pro Met Pro *** Ile
 Ser Thr Glu Glu Gly Gly Gln Phe Val Thr Leu Ser Pro Pro Cys Pro Glu Phe

Trp Ile Phe Tyr Ile Val Ser Asp Lys Lys Asp Ser Arg Leu Pro Lys *** ***
 Gly Tyr Ser Ile Phe *** Gln Thr Lys Lys Ile Val Glu Tyr His Asn Lys Asn
 Glu Met His Phe Leu Asn Ser Leu Arg Lys *** *** Lys Thr Ile Thr Lys Ile

 AAG GTA TAC TTT ATT TAA TGA CTC AGA AAA AAT AGT GAA GCA TTA CCA AAA ATA
 981 990 999 1008 1017 1026
 TTC CAT ATG AAA TAA ATT ACT GAG TCT TTT TTA TCA CTT CGT AAT GGT TTT TAT

 Phe His Met Lys *** Ile Thr Glu Ser Phe Leu Ser Leu Arg Asn Gly Phe Tyr
 Ser Ile *** Asn Lys Leu Leu Ser Leu Phe Tyr His Phe Val Met Val Phe Ile
 Pro Tyr Glu Ile Asn Tyr *** Val Phe Phe Ile Thr Ser *** Trp Phe Leu Leu

Glu Asn Leu Thr Leu His Pro Thr Lys Leu Ile Leu Asn Glu Ser Asn Tyr Met
 Asn Met Leu Pro *** Thr Pro Pro Arg *** Phe *** Ile Arg Gln Ile Thr Cys
 Ile *** *** Pro Asn Leu Pro Pro Asp Lys Phe Asn Phe Glu Arg Phe Gln Val

 ATA AGT AAT TCC CAA TTC ACC CCC CAG AAA TTT TAA TTT AAG AGA CTT AAC ATG
 1035 1044 1053 1062 1071 1080
 TAT TCA TTA AGG GTT AAG TGG GGG GTC TTT AAA ATT AAA TTC TCT GAA TTG TAC

 Tyr Ser Leu Arg Val Lys Trp Gly Val Phe Lys Ile Lys Phe Ser Glu Leu Tyr
 Ile His *** Gly Leu Ser Gly Gly Ser Leu Lys Leu Asn Ser Leu Asn Cys Thr
 Phe Ile Lys Gly *** Val Gly Gly Leu *** Asn *** Ile Leu *** Ile Val His

FIG. 8d

Cys Pro *** Val Ser Ile Thr Asn Arg Thr Thr Tyr Val Thr Lys Ser Arg Leu
 Val His Asn Cys Pro Tyr Gln Ile Gly Pro Arg Ile Tyr Gln Lys Arg Val Cys
 Tyr Met Thr Val Arg Ile Asn Tyr Glu Gln Asp Tyr Ile Ser Asn Glu Phe Ala

 TAT GTA CCA ATG TGC CTA TAA CAT AAG GAC CAG CAT ATA TGA CAA AAG CTT GCG
 1089 1098 1107 1116 1125 1134
 ATA CAT GGT TAC ACG GAT ATT GTA TTC CTG GTC GTA TAT ACT GTT TTC GAA CGC

 Ile His Gly Tyr Thr Asp Ile Val Phe Leu Val Val Tyr Thr Val Phe Glu Arg
 Tyr Met Val Thr Arg Ile Leu Tyr Ser Trp Ser Tyr Ile Leu Phe Ser Asn Ala
 Thr Trp Leu His Gly Tyr Cys Ile Pro Gly Arg Ile Tyr Cys Phe Arg Thr Gln

Ala Ser Ala *** Thr Thr *** Met Glu Leu Leu Lys Tyr Asp *** Gly Cys Ser
 His Arg Pro Arg Arg Pro Arg Cys Lys Trp Cys Asn Thr Thr Glu Ala Val Ala
 Thr Gly Leu Gly Val His Asp Val Asn Gly Ala Thr Gln Leu Arg Leu Trp Leu

 TCA CGG CTC CGG ATG CAC CAG ATG TAA AGG TCG TCA AAC ATC AGA GTC GGT GTC
 1143 1152 1161 1170 1179 1188
 AGT GCC GAG GCC TAC GTG GTC TAC ATT TCC AGC AGT TTG TAG TCT CAG CCA CAG

 Ser Ala Glu Ala Tyr Val Val Tyr Ile Ser Ser Ser Leu *** Ser Gln Pro Gln
 Val Pro Arg Pro Thr Trp Ser Thr Phe Pro Ala Val Cys Ser Leu Ser His Ser
 Cys Arg Gly Leu Arg Gly Leu His Phe Gln Gln Phe Val Val Ser Ala Thr Ala

Thr Glu Lys Thr Thr Gln Asn Ser Thr Ile Leu Leu Ser Ile *** Ser Leu Asn
 Pro Lys Lys Gln Gln Lys Thr Pro Leu Leu *** Tyr His Phe Arg Pro Cys Thr
 Gln Asn Arg Lys Asn Asn Pro Gln Phe Tyr Asp Ile Thr Phe Asp Leu Val Pro

 GAC CAA AGA AAA CAA CAA ACC AAC CTT CAT TAG TTA TCA CTT TAG ATC CTG TCC
 1197 1206 1215 1224 1233 1242
 CTG GTT TCT TTT GTT GTT TGG TTG GAA GTA ATC AAT AGT GAA ATC TAG GAC AGG

 Leu Val Ser Phe Val Val Trp Leu Glu Val Ile Asn Ser Glu Ile *** Asp Arg
 Trp Phe Leu Leu Leu Phe Gly Trp Lys *** Ser Ile Val Lys Ser Arg Thr Gly
 Gly Phe Phe Cys Cys Leu Val Gly Ser Asn Gln *** *** Asn Leu Gly Gln Val

Pro Pro Leu Thr Gly Pro Thr Thr Pro Ser Pro Ser Pro *** Pro Ile Ala Pro
 Gln Pro Tyr Leu Val Pro Leu Pro Leu Leu Leu Ala Pro Asn His Tyr Pro Pro
 Lys Pro Thr Phe Tyr Arg Ser His Tyr Ser Phe Pro Gln Thr Ile Thr His Arg

 AAA CCC CCA TTT CAT GGC CCT CAC CAT CCT CTT CCC GAC CCA ATA CCA TAC CGC
 1251 1260 1269 1278 1287 1296
 TTT GGG GGT AAA GTA CCG GGA GTG GTA GGA GAA GGG CTG GGT TAT GGT ATG GCG

 Phe Gly Gly Lys Val Pro Gly Val Val Gly Glu Gly Leu Gly Tyr Gly Met Ala
 Leu Gly Val Lys Tyr Arg Glu Trp *** Glu Lys Gly Trp Val Met Val Trp Arg
 Trp Gly *** Ser Thr Gly Ser Gly Arg Arg Arg Ala Gly Leu Trp Tyr Gly Gly

Pro Thr Thr *** Met Pro Thr Met Pro Ser Pro Gln Pro Arg Gln *** Leu Thr
 Leu Leu Leu Lys Cys Leu Pro *** Leu His Pro Ser His Gly Lys Asn Cys Leu
 Ser Ser Tyr Asn Val Tyr Pro Asp Tyr Thr Leu Ala Thr Ala Lys Thr Val Phe

 CCT CCT CAT CAA ATG TAT CCC CAG TAT CCA CTC CCG ACA CCG GAA ACA ATG TTT
 1305 1314 1323 1332 1341 1350
 GGA GGA GTA GTT TAC ATA GGG GTC ATA GGT GAG GGC TGT GGC CTT TGT TAC AAA

 Gly Gly Val Val Tyr Ile Gly Val Ile Gly Glu Gly Cys Gly Leu Cys Tyr Lys
 Glu Glu *** Phe Thr *** Gly Ser *** Val Arg Ala Val Ala Phe Val Thr Lys
 Arg Ser Ser Leu His Arg Gly His Arg *** Gly Leu Trp Pro Leu Leu Gln Ser

FIG. 8e

Ile Met *** Phe Leu Leu Val Pro Ala Trp Glu Gly Thr Val Arg Pro Ser Arg
 *** *** Arg Phe Tyr Cys Cys Gln Leu Gly Ser Gly Gln *** Gly Pro His Asp
 Asn Asp Asp Leu Ile Val Ala Ser Ser Gly Val Gly Arg Asp Gly Gln Thr Ile

 CAA TAG TAG ATT TTA TTG TCG TGA CCT CGG GTG AGG GGA CAG TGG GAC CCA CTA
 1359 1368 1377 1386 1395 1404
 GTT ATC ATC TAA AAT AAC AGC ACT GGA GCC CAC TCC CCT GTC ACC CTG GGT GAT

 Val Ile Ile *** Asn Asn Ser Thr Gly Ala His Ser Pro Val Thr Leu Gly Asp
 Leu Ser Ser Lys Ile Thr Ala Leu Glu Pro Thr Pro Leu Ser Pro Trp Val Ile
 Tyr His Leu Lys *** Gln His Trp Ser Pro Leu Pro Cys His Pro Gly *** Ser

 Pro Ala Pro Gly Ser Asn Leu Arg Leu Arg Glu *** Glu Thr Thr Asn Leu Pro
 Pro Leu Leu Ala Leu Ile *** Gly *** Gly Lys Lys Asn Gln Leu Ile *** Leu
 Pro Ser Cys Pro Trp Phe Glu Val Lys Val Lys Arg Ile Arg Tyr Tyr Glu Phe

 GCC CCT CGT CCC GGT CTT AAG TTG GAA TTG GAA AGA ATA AGA CAT CAT AAG TTT
 1413 1422 1431 1440 1449 1458
 CGG GGA GCA GGG CCA GAA TTC AAC CTT AAC CTT TCT TAT TCT GTA GTA TTC AAA

 Arg Gly Ala Gly Pro Glu Phe Asn Leu Asn Leu Ser Tyr Ser Val Val Phe Lys
 Gly Glu Gln Gly Gln Asn Ser Thr Leu Thr Phe Leu Ile Leu *** Tyr Ser Lys
 Gly Ser Arg Ala Arg Ile Gln Pro *** Pro Phe Leu Phe Cys Ser Ile Gln Arg

 Cys Leu Ala Pro Thr Gln Gly Gly Glu Gln Pro Phe Phe Thr Met Leu Ile Ser
 Ala Cys Leu Pro Pro Lys Val Gly Arg Arg Pro Ser Ser Leu *** *** Tyr Gln
 Pro Val Ser Arg Pro Asn Ser Gly Gly Gly Pro Pro Leu Phe Asp Asn Ile Asn

 CCC GTG TCT CGC CCC CAA ACT GGG GGG AGG ACC CCC TTC TTT CAG TAA TTA TAA
 1467 1476 1485 1494 1503 1512
 GGG CAC ACA GCG GGG GTT TGA CCC CCC TCC TGG GGG AGG AAA GTC ATT AAT ATT

 Gly His Arg Ala Gly Val *** Pro Pro Ser Trp Gly Lys Lys Val Ile Asn Ile
 Gly Thr Glu Arg Gly Phe Asp Pro Pro Pro Gly Gly Arg Lys Ser Leu Ile Leu
 Ala Gln Ser Gly Gly Leu Thr Pro Leu Leu Gly Glu Glu Ser His *** Tyr ***

 Asp *** *** Thr Trp Arg Gly Pro Pro Arg Glu Ser Gln Pro Glu Ser Ser Leu
 Ile Glu Asp His Gly Gly Gly Leu Leu Ala Asn Gln Ser His Asn Ala Gln Cys
 Phe Arg Met Met Asp Val Ala Trp Ser Pro Thr Arg Val Thr Thr Arg Lys Val

 CCT AGA GTA GTA CAG GTG GCG GGT CCT CCC GCA AGA CTG ACA CCA AGC GAA CTG
 1521 1530 1539 1548 1557 1566
 GAA TCT CAT CAT GTC CAC CGC CCA GGA GGG CGT TCT GAC TGT GGT TCG CTT GAC

 Glu Ser His His Val His Arg Pro Gly Gly Arg Ser Asp Cys Gly Ser Leu Asp
 Asn Leu Ile Met Ser Thr Ala Gln Glu Gly Val Leu Thr Val Val Arg Leu Thr
 Ile Ser Ser Cys Pro Pro Pro Arg Arg Ala Phe *** Leu Trp Phe Ala *** Gln

 Ile Asp Ser Pro Ala Pro Ser Ala Pro Thr Ser Ser Ala Met Lys Gly Glu Gly
 Tyr Ile Arg Leu His Pro Leu Pro Pro His Gln Leu His Trp Lys Glu Lys Glu
 Thr Tyr Gly Phe Thr Arg Ser Leu Arg Thr Asn Phe Ile Gly Asn Lys Arg Arg

 TCA TAT AGG CTT CCA CGC CCT CTC CGC CCA CAA CTT CTA CGG TAA AAA GGA AGA
 1575 1584 1593 1602 1611 1620
 AGT ATA TCC GAA GGT GCG GGA GAG GCG GGT GTT GAA GAT GCC ATT TTT CTT TCT

 Ser Ile Ser Glu Gly Ala Gly Glu Ala Gly Val Glu Asp Ala Ile Phe Pro Ser
 Val Tyr Pro Lys Val Arg Glu Arg Arg Val Leu Lys Met Pro Phe Phe Leu Leu
 Tyr Ile Arg Arg Cys Gly Arg Gly Gly Cys *** Arg Cys His Phe Ser Phe Ser

FIG. 8f

Ala Thr Val Thr Ala Pro Thr Ser Ser Gly Pro Ala Ala Ala Ser Ser Arg Ala
 Leu Pro Leu Pro Pro Pro Pro Pro Arg Ala Leu Pro Pro Pro Pro Asp Pro
 Trp Arg Tyr Arg His Arg Pro His Val Leu Trp Pro Arg Arg Arg Leu Ile Gln

 GGT CGC CAT TGC CAC CGC CCC CAC CTG CTC GGT CCC CGC CGC CGC CTC CTA GAC
 1629 1638 1647 1656 1665 1674
 CCA GCG GTA ACG GTG GCG GGG GTG GAC GAG CCA GGG GCG GCG GCG GAG GAT CTG

 Pro Ala Val Thr Val Ala Gly Val Asp Glu Pro Gly Ala Ala Ala Glu Asp Leu
 Gln Arg *** Arg Trp Arg Gly Trp Thr Ser Gln Gly Arg Arg Arg Ile Trp
 Ser Gly Asn Gly Gly Gly Gly Gly Arg Ala Arg Gly Gly Gly Gly Ser Gly

 Leu Ile Ala Ala Pro Ala Thr Asp Glu Glu Glu Thr Val Gly Gly Gln Ile Arg
 Trp Ser Pro Gln Pro Pro Pro Thr Lys Lys Lys Pro Leu Ala Glu Lys Ser Val
 Gly Leu His Ser Arg Pro Arg His Arg Arg Arg Arg Tyr Arg Arg Arg Pro Tyr

 CGG TTC TAC CGA CGC CCC CGC CAC AGA AGA AGA AGC CAT TGC GGA GGA ACC TAT
 1683 1692 1701 1710 1719 1728
 GCC AAG ATG GCT GCG GGG GCG GTG TCT TCT TCT TCG GTA ACG CCT CCT TGG ATA

 Ala Lys Met Ala Ala Gly Ala Val Ser Ser Ser Ser Val Thr Pro Pro Thr Ile
 Pro Arg Trp Leu Arg Gly Arg Cys Leu Leu Leu Arg *** Arg Leu Leu Gly Tyr
 Gln Asp Gly Cys Gly Gly Gly Val Phe Phe Phe Gly Asn Ala Ser Leu Asp Thr

 *** Ile Gln Phe Arg Phe Phe His Ala Thr Leu Ile
 Asp Tyr Arg Phe Val Phe Ser Thr Arg Gln Leu Tyr
 Thr Met Asp Ser Phe Ser Leu Leu Ala Ser Tyr Thr Asn

 GCA GTA TAG ACT TTT GCT TTC TTC ACG CGA CAT TCA TAA 5'
 1737 1746 1755 1764
 CGT CAT ATC TGA AAA CGA AAG AAG TGC GCT GTA AGT ATT 3'

 Arg His Ile *** Lys Arg Lys Lys Cys Ala Val Ser Ile
 Val Ile Ser Glu Asn Glu Arg Ser Ala Leu *** Val
 Ser Tyr Leu Lys Thr Lys Glu Val Arg Cys Lys Tyr

FIG. 8g




FIG. 9




FIG. 10





FIG. 11



TYPE B
SPOT NO. 104 TO 159

TYPE A
SPOT NO. 160 TO 215

FIG. 12




FIG. 14




FIG. 15




FIG. 16

1

**CIRCOVIRUS SEQUENCES ASSOCIATED
WITH PIGLET WEIGHT LOSS DISEASE
(PWD)**

INFORMATION ON RELATED APPLICATIONS

This application is a divisional of U.S. patent application Ser. No. 10/718,264, filed Nov. 21, 2003, now U.S. Pat. No. 7,179,472, which is a divisional of U.S. patent application Ser. No. 09/514,245, filed Feb. 28, 2000, now U.S. Pat. No. 6,703,023, which is a continuation-in-part of International Patent Application No. PCT/FR98/02634, filed Dec. 4, 1998, published in a non-English language, the specifications of which are incorporated herein by reference in their entireties.

BACKGROUND OF THE INVENTION

The invention relates to the genomic sequence and nucleotide sequences coding for polypeptides of PWD circovirus, such as the structural and nonstructural polypeptides of said circovirus, as well as vectors including said sequences and cells or animals transformed by these vectors. The invention likewise relates to methods for detecting these nucleic acids or polypeptides and kits for diagnosing infection by the PWD circovirus. The invention is also directed to a method for selecting compounds capable of modulating the viral infection. The invention further comprises pharmaceutical compositions, including vaccines, for the prevention and/or the treatment of viral infections by PWD circovirus as well as the use of a vector according to the invention for the prevention and/or the treatment of diseases by gene therapy.

Piglet weight loss disease (PWD), alternatively called fatal piglet wasting (FPW) has been widely described in North America (Harding, J. C., 1997), and authors have reported the existence of a relationship between this pathology and the presence of porcine circovirus (Daft, B. et al., 1996; Clark, E. G., 1997; Harding, J. C., 1997; Harding, J. C. and Clark, E. G., 1997; Nayar, G. P. et al., 1997). A porcine circovirus has already been demonstrated in established lines of cell cultures derived from pigs and chronically infected (Tischer, I., 1986, 1988, 1995; Dulac, G. C., 1989; Edwards, S., 1994; Allan, G. M., 1995 and McNeilly, F., 1996). This virus, during experimental infection of piglets, does not prove pathogenic for pigs (Tischer, I., 1986, Horner, G. W., 1991) and its nucleotide sequence has been determined and characterized (Tischer, I., 1982; Meehan, B. M. et al., 1997; Mankertz, A., 1997). The porcine circovirus, called PCV virus, is part of the circovirus genus of the circoviridae family (Murphy, F. A. et al., 1995) whose virion has a circular DNA of size between 1.7 and 2.3 kb, which DNA comprises three open reading frames (ORF1 to ORF3), coding for a replication protein REP involved in the initiation and termination phase of rolling circular replication (RCR) (Heyraud-Nitschke, F., et al., 1995; Harding, M. R. et al., 1993; Hanson, S. F. et al., 1995; Fontes, E. P. B. et al., 1994), coding for a capsid protein (Boulton, L. H. et al., 1997; Hackland, A. F. et al., 1994; Chu, P. W. G. et al., 1993) and coding for a nonstructural protein called a dissemination protein (Lazarowitz, S. G. et al., 1989).

The inventors of the present invention have noticed that the clinical signs perceptible in pigs and linked to infection by the PWD circovirus are very distinctive. These manifestations in general appear in pigs of 8 to 12 weeks of age, weaned for 4 to 8 weeks. The first signs are hypotonia without it being possible to speak of prostration. Rapidly (48 hours), the flanks hollow, the line of the spine becomes apparent, and the pigs "blanch." These signs are in general accompanied by hyperthermia, anorexia and most often by respiratory signs

2

(coughing, dyspnea, polypnea). Transitory diarrhea can likewise appear. The disease state phase lasts approximately one month at the end of which the rate of mortality varies from 5 to 20%. To these mortalities, it is expedient to add a variable proportion (5-10%) of cadaveric animals which are no longer able to present an economic future. It is to be noted that outside of this critical stage of the end of post-weaning, no anomaly appears on the farms. In particular, the reproductive function is totally maintained.

On the epidemiological level, the first signs of this pathology appeared at the start of 1995 in the east of the Côtes d'Armor region in France, and the farms affected are especially confined to this area of the region. In December 1996, the number of farms concerned could not be evaluated with precision because of the absence of a specific laboratory diagnostic method or of an epidemiological surveillance system of the livestock. Based on the clinical facts as well as on results of postmortem examinations supplied by veterinarians, it is possible to estimate this number as several dozen (80-100). The contagiousness of the disease is weak to moderate. Cases are being reported outside the initial area and for the majority are following the transfer of animals coming from farms familiar with the problem. On the other hand, a characteristic of the condition is its strong remanence. Thus, farms which have been affected for a year are still affected in spite of the massive administration of therapeutics. Farms with clinical expression are drawn from various categories of specialization (breeders/fatteners, post-weaners/fatteners) and different economic structures are concerned. In addition, the disorders appear even in farms where the rules of animal husbandry are respected.

Numerous postmortem examinations have been carried out either on farms or in the laboratory. The elements of the lesional table are disparate. The most constant macroscopic lesions are pneumonia which sometimes appears in patchy form as well as hypertrophy of the lymphatic ganglia. The other lesions above all affect the thoracic viscera including, especially, pericarditis and pleurisy. However, arthritis and gastric ulcers are also observed. The lesions revealed in the histological examination are essentially situated at the pulmonary level (interstitial pneumonia), ganglionic level (lymphoid depletion of the lymph nodes, giant cells) and renal level (glomerulonephritis, vasculitis). The infectious agents have been the subject of wide research. It has been possible to exclude the intervention of pestiviruses and Aujeszky's disease. The disorders appear in the seropositive PDRS (Porcine Dysgenic and Respiratory Syndrome, an infection linked to an arterivirus) herds, but it has not been possible to establish the role of the latter in the genesis of the disorders (the majority of the farms in Brittany are PDRS seropositive).

The inventors of the present invention, with the aim of identifying the etiological agent responsible for PWD, have carried out "contact" tests between piglets which are obviously "ill" and SPF pigs (specific pathogen-free) from CNEVA (Centre National d'Etudes Vétérinaires et Alimentaires, France). These tests allow the development of signs comparable to those observed on the farm to be observed in protected animal houses. The discrete signs such as moderate hyperthermia, anorexia and intermittent diarrhea appeared after one week of contact. It must be noted that the PDRS virus only diffused subsequent to the clinical signs. In addition, inoculations of organ homogenates of sick animals to healthy pigs allowed signs related to those observed on the farms to be reproduced, although with a lower incidence, linked to the favorable conditions of upkeep of the animals in the experimental installations.

Thus, the inventors of the present invention have been able to demonstrate that the pathological signs appear as a well-defined entity affecting the pig at a particular stage of its growth.

This pathology has never been described in France. However, sparse information, especially Canadian, relates to similar facts.

The disorders cannot be mastered with the existing therapeutics.

The data collected both on the farm and by experimentation have allowed the following points to be highlighted:

PWD is transmissible but its contagiousness is not very high,

its etiological origin is of infectious and probably viral nature,

PWD has a persistent character in the affected farms.

Considerable economic consequences ensue for the farms.

Thus, there is currently a significant need for a specific and sensitive diagnostic, whose production is practical and rapid, allowing the early detection of the infection.

A reliable, sensitive and practical test which allows the distinction between strains of porcine circovirus (PCV) is thus strongly desirable.

On the other hand, a need for efficient and well-tolerated treatment of infections with PWD circovirus likewise remains desirable, no vaccine currently being available against PWD circovirus.

Concerning PWD circovirus, it will probably be necessary to understand the role of the immune defense in the physiology and the pathology of the disease to develop satisfactory vaccines.

Fuller information concerning the biology of these strains, their interactions with their hosts, the associated infectivity phenomena and those of escape from the immune defenses of the host especially, and finally their implication in the development of associated pathologies, will allow a better understanding of these mechanisms. Taking into account the facts which have been mentioned above and which show in particular the limitations of combating infection by the PWD circovirus, it is thus essential today on the one hand to develop molecular tools, especially starting from a better genetic knowledge of the PWD circovirus, and likewise to perfect novel preventive and therapeutic treatments, novel methods of diagnosis and specific, efficacious and tolerated novel vaccine strategies. This is precisely the subject of the present invention.

SUMMARY OF THE INVENTION

The present invention relates to vaccines comprising a nucleotide sequence of the genome of Porcine circovirus type B, or a homologue or fragment thereof, and an acceptable pharmaceutical or veterinary vehicle. In one embodiment of the invention, the nucleotide sequence is selected from SEQ ID No. 15, SEQ ID No. 19, SEQ ID No. 23, or SEQ ID No. 25, or a homologue or fragment thereof. In another embodiment of the invention, the homologue has at least 80% sequence identity to SEQ ID No. 15, SEQ ID No. 19, SEQ ID No. 23 or SEQ ID No. 25. In yet another embodiment, the vaccines further comprising an adjuvant

The present invention also relates to vaccines comprising a polypeptide encoded by a nucleotide sequence of the genome of PCVB, or a homologue or fragment thereof, and an acceptable pharmaceutical or veterinary vehicle. In one embodiment, the homologue has at least 80% sequence identity to SEQ ID No. 15, SEQ ID No. 19, SEQ ID No. 23 or SEQ ID No. 25. In another embodiment of the invention, the nucle-

otide sequence is selected from SEQ ID No. 23 or SEQ ID No. 25, or a homologue or fragment thereof. In still another embodiment, the polypeptide has the amino acid sequence of SEQ ID No. 24 or SEQ ID No. 26. In yet another embodiment, the homologue has at least 80% sequence identity to SEQ ID No. 24 or SEQ ID No. 26. In another embodiment, the polypeptide has the amino acid sequence of SEQ ID No. 29, SEQ ID No. 30, SEQ ID No. 31, or SEQ ID No. 32.

A further aspect of the invention relates to vaccines comprising a vector and an acceptable pharmaceutical or veterinary vehicle, the vector comprising a nucleotide sequence of the genome of Porcine circovirus type B, or a homologue or fragment thereof. In one embodiment, the vaccine further comprises a gene coding for an expression product capable of inhibiting or retarding the establishment or development of a genetic or acquired disease.

The present invention also relates to vaccines comprising a cell and an acceptable pharmaceutical or veterinary vehicle, wherein the cell is transformed with a nucleotide sequence of the genome of Porcine circovirus type B, or a homologue or fragment thereof.

Still further, the present invention relates to vaccines comprising a pharmaceutically acceptable vehicle and a single polypeptide, wherein the single polypeptide consists of SEQ ID No. 26.

Additionally, the present invention relates to methods of immunizing a mammal against piglet weight loss disease comprising administering to a mammal an effective amount of the vaccines described above.

These and other aspects of the invention will become apparent to the skilled artisan in view of the teachings contained herein.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1: Experimental scheme which has made it possible to bring about the isolation and the identification of the circovirus associated with PWD of type A and B.

Test 1: experimental reproduction of the PWD by inoculation of pig organ homogenates from farms affected by PWD.

Test 2: experimental reproduction of PWD.

Test 3: experimental reproduction of PWD.

Test 4: no experimental reproduction of PWD.

FIG. 2: Organization of the genome of the circovirus associated with PWD of type A (PCVA)

strand of (+) polarity (SEQ ID No. 1);

strand of (-) polarity (SEQ ID No. 5, represented according to the orientation 3'→5');

sequences of amino acids of proteins encoded by the two DNA strands in the three possible reading frames SEQ ID NOS: 2-4 and 6-8 respectively.

FIG. 3: Alignment of the nucleotide sequence SEQ ID No. 1 of the PWD circovirus of type A (PCVA) and of the MEEHAN SEQ ID No. 163 strain and MANKERTZ SEQ ID No. 164 strain circoviruses of the porcine cell lines.

FIG. 4: Alignment of the sequence of amino acids SEQ ID No. 12 of a polypeptide encoded by the nucleotide sequence SEQ ID No. 9 (ORF1) of the PWD circovirus of type A (PCVA) and of corresponding nucleotide sequences of the MEEHAN SEQ ID No. 165 strain and MANKERTZ SEQ ID No. 166 strain circoviruses of the porcine cell lines.

FIG. 5: Alignment of the sequence of amino acids SEQ ID No. 12 of a polypeptide encoded by the nucleotide sequence SEQ ID No. 11 (ORF2) of the PWD circovirus of type A (PCVA) and of corresponding nucleotide sequences of the MEEHAN SEQ ID No. 167 strain and MANKERTZ SEQ ID No. 168 strain circoviruses of the porcine cell lines.

FIG. 6: Alignment of the sequence of amino acids SEQ ID No. 14 of a polypeptide encoded by the nucleotide sequence SEQ ID No. 13 (ORF3) of the PWD circovirus of type A (PCVA) and of corresponding nucleotide sequences of the MEEHAN SEQ ID No. 169 strain and MANKERTZ SEQ ID No. 170 strain circoviruses of the porcine cell lines.

FIG. 7: Western blot analysis of recombinant proteins of the PWD circovirus of type A (PCVA).

The analyses were carried out on cell extracts of Sf9 cells obtained after infection with recombinant baculovirus PCF ORF 1.

FIG. 8: Organization of the genome of the circovirus associated with the PWD of type B (PCVB)

strand of (+) polarity (SEQ ID No. 15);

strand of (-) polarity (SEQ ID No. 19, represented according to the orientation 3'→5');

sequence of amino acids of proteins encoded by the two DNA strands in the three possible reading frames SEQ ID NOS: 16-18 and 20-22 respectively.

FIG. 9: Evolution of the daily mean gain (DMG) of pig farms affected by piglet weight loss disease (PWD), placed under experimental conditions.

FIG. 10: DMG compared for the 3 batches of pigs (F1, F3 and F4) calculated over a period of 28 days, after vaccination test.

FIG. 11: Hyperthermia greater than 41° C., expressed as a percentage compared for the 3 batches of pigs (F1, F3 and F4) calculated per week over a period of 28 days, after vaccination test.

FIG. 12: Membranes of peptide spots corresponding to the ORF2s revealed with the aid of an infected pig serum, originating from a conventional farm.

The numbers of specific peptides of the circovirus of type B as well as their nonreactive homologs (type A) are indicated in bold.

The nonspecific immunogenic peptides are indicated in italics.

FIG. 13: Alignment of amino acid sequences of proteins encoded by the ORF2 of the PWD circovirus of type A SEQ ID No. 12 and by the ORF2 of the PWD circovirus of type B SEQ ID No. 26. The position of 4 peptides corresponding to specific epitopes of the PWD circovirus of type B is indicated on the corresponding sequence by a bold line, their homolog on the sequence of the PWD circovirus of type A is likewise indicated by an ordinary line.

FIG. 14: Charts the results of experiments that demonstrate, in terms of percent hyperthermia, that vaccination with ORF1 and ORF2 of PCV-B enhances the level of protection in swine challenged with PCV-B (Percent hyperthermia: >40.5 C, control: not vaccinated and not challenged, ORF1: vaccinated and challenged, ORF2: vaccinated and challenged, ORF: not vaccinated, challenged).

FIG. 15: Charts the results of experiments that demonstrate, in terms of animal growth, that vaccination with ORF1 and ORF2 of PCV-B enhances the level of protection in swine challenged with PCV-B (Control: not vaccinated, not challenged, ORF1: vaccinated and challenged, ORF2: vaccinated and challenged, ORF: not vaccinated, challenged).

FIG. 16: Immunoperoxidase staining of PK15 cells at 24 h post-transfection with the pcDNA3/ORF2 plasmid. Expression of PCVB ORF2 was confirmed by IPMA following incubation in the presence of the swine anti-PCVB monospecific serum.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to nucleotide sequences of the genome of PWD circovirus selected from the sequences SEQ ID No. 1, SEQ ID No. 5, SEQ ID No. 15, SEQ ID No. 19 or one of their fragments.

The nucleotide sequences of sequences SEQ ID No. 1 and SEQ ID No. 5 correspond respectively to the genome sequence of the strand of (+) polarity and of the strand of (-) polarity of the PWD circovirus of type A (or PCVA), the sequence SEQ ID No. 5 being represented according to the orientation 5'→3'.

The nucleotide sequences of sequences SEQ ID No. 15 and SEQ ID No. 19 correspond respectively to the genome sequence of the strand of (+) polarity and of the strand of (-) polarity of the PWD circovirus of type B (or PCVB), the sequence SEQ ID No. 19 being represented according to the orientation 5'→3'.

The present invention likewise relates to nucleotide sequences, characterized in that they are selected from:

- a) a nucleotide sequence of a specific fragment of the sequence SEQ ID No. 1, SEQ ID No. 5, SEQ ID No. 15, SEQ ID No. 19 or one of their fragments;
- b) a nucleotide sequence homologous to a nucleotide sequence such as defined in a);
- c) a nucleotide sequence complementary to a nucleotide sequence such as defined in a) or b), and a nucleotide sequence of their corresponding RNA;
- d) a nucleotide sequence capable of hybridizing under stringent conditions with a sequence such as defined in a), b) or c);
- e) a nucleotide sequence comprising a sequence such as defined in a), b), c) or d); and
- f) a nucleotide sequence modified by a nucleotide sequence such as defined in a), b), c), d) or e).

Nucleotide, polynucleotide or nucleic acid sequence will be understood according to the present invention as meaning both a double-stranded or single-stranded DNA in the monomeric and dimeric (so-called in tandem) forms and the transcription products of said DNAs.

It must be understood that the present invention does not relate to the genomic nucleotide sequences taken in their natural environment, that is to say in the natural state. It concerns sequences which it has been possible to isolate, purify or partially purify, starting from separation methods such as, for example, ion-exchange chromatography, by exclusion based on molecular size, or by affinity, or alternatively fractionation techniques based on solubility in different solvents, or starting from methods of genetic engineering such as amplification, cloning and subcloning, it being possible for the sequences of the invention to be carried by vectors.

The nucleotide sequences SEQ ID No. 1 and SEQ ID No. 15 were obtained by sequencing of the genome by the Sanger method.

Nucleotide sequence fragment according to the invention will be understood as designating any nucleotide fragment of the PWD circovirus, type A or B, of length of at least 8 nucleotides, preferably at least 12 nucleotides, and even more preferentially at least 20 consecutive nucleotides of the sequence from which it originates.

Specific fragment of a nucleotide sequence according to the invention will be understood as designating any nucleotide fragment of the PWD circovirus, type A or B, having, after alignment and comparison with the corresponding fragments of known porcine circoviruses, at least one nucleotide or base of different nature. For example, the specific nucleotide fragments of the PWD circovirus of type A can easily be determined by referring to FIG. 3 of the present invention in which the nucleotides or bases of the sequence SEQ ID No. 1 (circopordfp) are shown which are of different nature, after

alignment of said sequence SEQ ID No. 1 with the other two sequences of known porcine circovirus (circopormeeh and circopormank).

Homologous nucleotide sequence in the sense of the present invention is understood as meaning a nucleotide sequence having at least a percentage identity with the bases of a nucleotide sequence according to the invention of at least 80%, preferably 90% or 95%, this percentage being purely statistical and it being possible to distribute the differences between the two nucleotide sequences at random and over the whole of their length.

Specific homologous nucleotide sequence in the sense of the present invention is understood as meaning a homologous nucleotide sequence having at least one nucleotide sequence of a specific fragment, such as defined above. Said "specific" homologous sequences can comprise, for example, the sequences corresponding to the genomic sequence or to the sequences of its fragments representative of variants of PWD circovirus of type A or B. These specific homologous sequences can thus correspond to variations linked to mutations within strains of PWD circovirus of type A and B, and especially correspond to truncations, substitutions, deletions and/or additions of at least one nucleotide. Said homologous sequences can likewise correspond to variations linked to the degeneracy of the genetic code.

The term "degree or percentage of sequence homology" refers to "degree or percentage of sequence identity between two sequences after optimal alignment" as defined in the present application.

Two amino-acids or nucleotidic sequences are said to be "identical" if the sequence of amino-acids or nucleotidic residues, in the two sequences is the same when aligned for maximum correspondence as described below. Sequence comparisons between two (or more) peptides or polynucleotides are typically performed by comparing sequences of two optimally aligned sequences over a segment or "comparison window" to identify and compare local regions of sequence similarity. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman, *Ad. App. Math* 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970), by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. (U.S.A.)* 85: 2444 (1988), by computerized implementation of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wis.), or by visual inspection.

"Percentage of sequence identity" (or degree or identity) is determined by comparing two optimally aligned sequences over a comparison window, where the portion of the peptide or polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical amino-acid residue or nucleic acid base occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The definition of sequence identity given above is the definition that would use one of skill in the art. The definition by itself does not need the help of any algorithm, said algorithms being helpful only to achieve the optimal alignments of sequences, rather than the calculation of sequence identity.

From the definition given above, it follows that there is a well defined and only one value for the sequence identity between two compared sequences which value corresponds to the value obtained for the best or optimal alignment.

In the BLAST N or BLAST P "BLAST 2 sequence", software which is available in the web site <http://www.ncbi.nlm.nih.gov/gorf/b12.html>, and habitually used by the inventors and in general by the skilled man for comparing and determining the identity between two sequences, gap cost which depends on the sequence length to be compared is directly selected by the software (i.e. 11.2 for substitution matrix BLOSUM-62 for length >85).

In the present description, PWD circovirus will be understood as designating the circoviruses associated with piglet weight loss disease (PWD) of type A (PCVA) or type B (PCVB), defined below by their genomic sequence, as well as the circoviruses whose nucleic sequences are homologous to the sequences of PWD circoviruses of type A or B, such as in particular the circoviruses corresponding to variants of the type A or of the type B.

Complementary nucleotide sequence of a sequence of the invention is understood as meaning any DNA whose nucleotides are complementary to those of the sequence of the invention, and whose orientation is reversed (antiparallel sequence).

Hybridization under conditions of stringency with a nucleotide sequence according to the invention is understood as meaning a hybridization under conditions of temperature and ionic strength chosen in such a way that they allow the maintenance of the hybridization between two fragments of complementary DNA.

By way of illustration, conditions of great stringency of the hybridization step with the aim of defining the nucleotide fragments described above are advantageously the following.

The hybridization is carried out at a preferential temperature of 65° C. in the presence of SSC buffer, 1×SSC corresponding to 0.15 M NaCl and 0.05 M Na citrate. The washing steps, for example, can be the following:

2×SSC, at ambient temperature followed by two washes with 2×SSC, 0.5% SDS at 65° C.; 2×0.5×SSC, 0.5% SDS; at 65° C. for 10 minutes each.

The conditions of intermediate stringency, using, for example, a temperature of 42° C. in the presence of a 2×SSC buffer, or of less stringency, for example a temperature of 37° C. in the presence of a 2×SSC buffer, respectively require a globally less significant complementarity for the hybridization between the two sequences.

The stringent hybridization conditions described above for a polynucleotide with a size of approximately 350 bases will be adapted by the person skilled in the art for oligonucleotides of greater or smaller size, according to the teaching of Sambrook et al., 1989.

Among the nucleotide sequences according to the invention, those are likewise preferred which can be used as a primer or probe in methods allowing the homologous sequences according to the invention to be obtained, these methods, such as the polymerase chain reaction (PCR), nucleic acid cloning and sequencing, being well known to the person skilled in the art.

Among said nucleotide sequences according to the invention, those are again preferred which can be used as a primer or probe in methods allowing the presence of PWD circovirus or one of its variants such as defined below to be diagnosed.

The nucleotide sequences according to the invention capable of modulating, of inhibiting or of inducing the expression of PWD circovirus gene, and/or capable of modulating the replication cycle of PWD circovirus in the host cell

and/or organism are likewise preferred. Replication cycle will be understood as designating the invasion and the multiplication of PWD circovirus, and its propagation from host cell to host cell in the host organism.

Among said nucleotide sequences according to the invention, those corresponding to open reading frames, called ORF sequences, and coding for polypeptides, such as, for example, the sequences SEQ ID No. 9 (ORF1), SEQ ID No. 11 (ORF2) and SEQ ID No. 13 (ORF3) respectively corresponding to the nucleotide sequences between the positions 47 and 985 determined with respect to the position of the nucleotides on the sequence SEQ ID No. 1, the positions 1723 and 1022 and the positions 658 and 38 with respect to the position of the nucleotides on the sequence SEQ ID No. 5 (represented according to the orientation 3'→5'), the ends being included, or alternatively the sequences SEQ ID No. 23 (ORF'1), SEQ ID No. 25 (ORF'2) and SEQ ID No. 27 (ORF'3), respectively corresponding to the sequences between the positions 51 and 995 determined with respect to the position of the nucleotides on the sequence SEQ ID No. 15, the positions 1734 and 1033 and the positions 670 and 357, the positions being determined with respect to the position of the nucleotides on the sequence SEQ ID No. 19 (represented according to the orientation 3'→5'), the ends being included, are finally preferred.

The nucleotide sequence fragments according to the invention can be obtained, for example, by specific amplification, such as PCR, or after digestion with appropriate restriction enzymes of nucleotide sequences according to the invention, these methods in particular being described in the work of Sambrook et al., 1989. Said representative fragments can likewise be obtained by chemical synthesis when their size is not very large and according to methods well known to persons skilled in the art.

Modified nucleotide sequence will be understood as meaning any nucleotide sequence obtained by mutagenesis according to techniques well known to the person skilled in the art, and containing modifications with respect to the normal sequences according to the invention, for example mutations in the regulatory and/or promoter sequences of polypeptide expression, especially leading to a modification of the rate of expression of said polypeptide or to a modulation of the replicative cycle.

Modified nucleotide sequence will likewise be understood as meaning any nucleotide sequence coding for a modified polypeptide such as defined below.

The present invention relates to nucleotide sequences of PWD circovirus according to the invention, characterized in that they are selected from the sequences SEQ ID No. 9, SEQ ID No. 11, SEQ ID No. 13, SEQ ID No. 23, SEQ ID No. 25, SEQ ID No. 27 or one of their fragments.

The invention likewise relates to nucleotide sequences characterized in that they comprise a nucleotide sequence selected from:

- a) a nucleotide sequence SEQ ID No. 9, SEQ ID No. 11, SEQ ID No. 13, SEQ ID No. 23, SEQ ID No. 25, SEQ ID No. 27 or one of their fragments;
- b) a nucleotide sequence of a specific fragment of a sequence such as defined in a);
- c) a homologous nucleotide sequence having at least 80% identity with a sequence such as defined in a) or b);
- d) a complementary nucleotide sequence or sequence of RNA corresponding to a sequence such as defined in a), b) or c); and
- e) a nucleotide sequence modified by a sequence such as defined in a), b), c) or d).

As far as homology with the nucleotide sequences SEQ ID No. 9, SEQ ID No. 11, SEQ ID No. 13, SEQ ID No. 23, SEQ ID No. 25, SEQ ID No. 27 or one of their fragments is concerned, the homologous, especially specific, sequences having a percentage identity with one of the sequences SEQ

ID No. 9, SEQ ID No. 11, SEQ ID No. 13, SEQ ID No. 23, SEQ ID No. 25, SEQ ID No. 27 or one of their fragments of at least 80%, preferably 90% or 95%, are preferred. Said specific homologous sequences can comprise, for example, the sequences corresponding to the sequences ORF1, ORF2, ORF3, ORF'1, ORF'2 and ORF'3 of PWD circovirus variants of type A or of type B. In the same manner, these specific homologous sequences can correspond to variations linked to mutations within strains of PWD circovirus of type A or of type B and especially correspond to truncations, substitutions, deletions and/or additions of at least one nucleotide.

Among nucleotide sequences according to the invention, the sequence SEQ ID No. 23 which has a homology having more than 80% identity with the sequence SEQ ID No. 9, as well as the sequence SEQ ID No. 25, are especially preferred.

Preferably, the invention relates to the nucleotide sequences according to the invention, characterized in that they comprise a nucleotide sequence selected from the following sequences:

- a) SEQ ID No. 33 170 5' TGTGGCGA 3';
- b) SEQ ID No. 34 450 5' AGTTTCCT 3';
- c) SEQ ID No. 35 1026 5' TCATTTAGAGGGTCTTTCAG 3';
- d) SEQ ID No. 36 1074 5' GTCAACCT 3';
- e) SEQ ID No. 37 1101 5' GTGGTTGC 3';
- f) SEQ ID No. 38 1123 5' AGCCCAGG 3';
- g) SEQ ID No. 39 1192 5' TTGGCTGG 3';
- h) SEQ ID No. 40 1218 5' TCTAGCTCTGGT 3';
- i) SEQ ID No. 41 1501 5' ATCTCAGCTCTGT 3';
- j) SEQ ID No. 42 1536 5' TGTCCTCCTTCT 3';
- k) SEQ ID No. 43 1563 5' TCTCTAGA 3';
- l) SEQ ID No. 44 1623 5' TGTACCAA 3';
- m) SEQ ID No. 45 1686 5' TCCGTCTT 3';

and their complementary sequences.

In the list of nucleotide sequences a)-m) above, the underlined nucleotides are mutated with respect to the two known sequences of circovirus which are nonpathogenic to pigs. The number preceding the nucleotide sequence represents the position of the first nucleotide of said sequence in the sequence SEQ ID No. 1.

The invention comprises the polypeptides encoded by a nucleotide sequence according to the invention, preferably a polypeptide whose sequence is represented by a fragment, especially a specific fragment, of one of the six sequences of amino acids represented in FIG. 2, these six amino acid sequences corresponding to the polypeptides which can be encoded according to one of the three possible reading frames of the sequence SEQ ID No. 1 or of the sequence SEQ ID No. 5, or a polypeptide whose sequence is represented by a fragment, especially a specific fragment, of one of the six sequences of amino acids shown in FIG. 8, these six sequences of amino acids corresponding to the polypeptides which can be encoded according to one of the three possible reading frames of the sequence SEQ ID No. 15 or of the sequence SEQ ID No. 19.

The invention likewise relates to the polypeptides, characterized in that they comprise a polypeptide selected from the amino acid sequences SEQ ID No. 10, SEQ ID No. 12, SEQ ID No. 14, SEQ ID No. 24, SEQ ID No. 26, SEQ ID No. 28 or one of their fragments.

Among the polypeptides according to the invention, the polypeptide of amino acid sequence SEQ ID No. 24 which has a homology having more than 80% identity with the sequence SEQ ID No. 10, as well as the polypeptide of sequence SEQ ID No. 26, are especially preferred.

The invention also relates to the polypeptides, characterized in that they comprise a polypeptide selected from:

- a) a specific fragment of at least 5 amino acids of a polypeptide of an amino acid sequence according to the invention;
- b) a polypeptide homologous to a polypeptide such as defined in a);
- c) a specific biologically active fragment of a polypeptide such as defined in a) or b); and
- d) a polypeptide modified by a polypeptide such as defined in a), b) or c).

Among the polypeptides according to the invention, the polypeptides of amino acid sequences SEQ ID No. 29, SEQ ID No. 30, SEQ ID No. 31 and SEQ ID No. 32 are also preferred, these polypeptides being especially capable of specifically recognizing the antibodies produced during infection by the PWD circovirus of type B. These polypeptides thus have epitopes specific for the PWD circovirus of type B and can thus be used in particular in the diagnostic field or as immunogenic agent to confer protection in pigs against infection by PWD circovirus, especially of type B.

In the present description, the terms polypeptide, peptide and protein are interchangeable.

It must be understood that the invention does not relate to the polypeptides in natural form, that is to say that they are not taken in their natural environment but that they can be isolated or obtained by purification from natural sources, or else obtained by genetic recombination, or alternatively by chemical synthesis and that they can thus contain unnatural amino acids, as will be described below.

Polypeptide fragment according to the invention is understood as designating a polypeptide containing at least 5 consecutive amino acids, preferably 10 consecutive amino acids or 15 consecutive amino acids.

In the present invention, specific polypeptide fragment is understood as designating the consecutive polypeptide fragment encoded by a specific fragment nucleotide sequence according to the invention.

Homologous polypeptide will be understood as designating the polypeptides having, with respect to the natural polypeptide, certain modifications such as, in particular, a deletion, addition or substitution of at least one amino acid, a truncation, a prolongation, a chimeric fusion, and/or a mutation. Among the homologous polypeptides, those are preferred whose amino acid sequence has at least 80%, preferably 90%, homology with the sequences of amino acids of polypeptides according to the invention.

Specific homologous polypeptide will be understood as designating the homologous polypeptides such as defined above and having a specific fragment of polypeptide according to the invention.

In the case of a substitution, one or more consecutive or nonconsecutive amino acids are replaced by "equivalent" amino acids. The expression "equivalent" amino acid is directed here at designating any amino acid capable of being substituted by one of the amino acids of the base structure without, however, essentially modifying the biological activities of the corresponding peptides and such that they will be defined by the following.

These equivalent amino acids can be determined either by depending on their structural homology with the amino acids which they substitute, or on results of comparative tests of biological activity between the different polypeptides, which are capable of being carried out.

By way of example, the possibilities of substitutions capable of being carried out without resulting in an extensive modification of the biological activity of the corresponding modified polypeptides will be mentioned, the replacement, for example, of leucine by valine or isoleucine, of aspartic acid by glutamic acid, of glutamine by asparagine, of arginine by lysine etc., the reverse substitutions naturally being envisageable under the same conditions.

The specific homologous polypeptides likewise correspond to polypeptides encoded by the specific homologous nucleotide sequences such as defined above and thus comprise in the present definition the polypeptides which are mutated or correspond to variants which can exist in PWD circovirus, and which especially correspond to truncations, substitutions, deletions and/or additions of at least one amino acid residue.

Specific biologically active fragment of a polypeptide according to the invention will be understood in particular as designating a specific polypeptide fragment, such as defined above, having at least one of the characteristics of polypeptides according to the invention, especially in that it is:

- capable of inducing an immunogenic reaction directed against a PWD circovirus; and/or
- capable of being recognized by a specific antibody of a polypeptide according to the invention; and/or
- capable of linking to a polypeptide or to a nucleotide sequence of PWD circovirus; and/or
- capable of exerting a physiological activity, even partial, such as, for example, a dissemination or structural (capsid) activity; and/or
- capable of modulating, of inducing or of inhibiting the expression of PWD circovirus gene or one of its variants, and/or capable of modulating the replication cycle of PWD circovirus in the cell and/or the host organism.

The polypeptide fragments according to the invention can correspond to isolated or purified fragments naturally present in a PWD circovirus or correspond to fragments which can be obtained by cleavage of said polypeptide by a proteolytic enzyme, such as trypsin or chymotrypsin or collagenase, or by a chemical reagent, such as cyanogen bromide (CNBr) or alternatively by placing said polypeptide in a very acidic environment, for example at pH 2.5. Such polypeptide fragments can likewise just as easily be prepared by chemical synthesis, from hosts transformed by an expression vector according to the invention containing a nucleic acid allowing the expression of said fragments, placed under the control of appropriate regulation and/or expression elements.

"Modified polypeptide" of a polypeptide according to the invention is understood as designating a polypeptide obtained by genetic recombination or by chemical synthesis as will be described below, having at least one modification with respect to the normal sequence. These modifications will especially be able to bear on amino acids at the origin of a specificity, of pathogenicity and/or of virulence, or at the origin of the structural conformation, and of the capacity of membrane insertion of the polypeptide according to the invention. It will thus be possible to create polypeptides of equivalent, increased or decreased activity, and of equivalent, narrower, or wider specificity. Among the modified polypeptides, it is necessary to mention the polypeptides in which up to 5 amino acids can be modified, truncated at the N- or C-terminal end, or even deleted or added.

As is indicated, the modifications of the polypeptide will especially have as objective:

- to render it capable of modulating, of inhibiting or of inducing the expression of PWD circovirus gene and/or capable of modulating the replication cycle of PWD circovirus in the cell and/or the host organism,

of allowing its incorporation into vaccine compositions, of modifying its bioavailability as a compound for therapeutic use.

The methods allowing said modulations on eukaryotic or prokaryotic cells to be demonstrated are well known to the person skilled in the art. It is likewise well understood that it will be possible to use the nucleotide sequences coding for said modified polypeptides for said modulations, for example through vectors according to the invention and described below, in order, for example, to prevent or to treat the pathologies linked to the infection.

The preceding modified polypeptides can be obtained by using combinatorial chemistry, in which it is possible to systematically vary parts of the polypeptide before testing them on models, cell cultures or microorganisms for example, to select the compounds which are most active or have the properties sought.

Chemical synthesis likewise has the advantage of being able to use:

unnatural amino acids, or
nonpeptide bonds.

Thus, in order to improve the duration of life of the polypeptides according to the invention, it may be of interest to use unnatural amino acids, for example in D form, or else amino acid analogs, especially sulfur-containing forms, for example.

Finally, it will be possible to integrate the structure of the polypeptides according to the invention, its specific or modified homologous forms, into chemical structures of polypeptide type or others. Thus, it may be of interest to provide at the N- and C-terminal ends compounds not recognized by the proteases.

The nucleotide sequences coding for a polypeptide according to the invention are likewise part of the invention.

The invention likewise relates to nucleotide sequences utilizable as a primer or probe, characterized in that said sequences are selected from the nucleotide sequences according to the invention.

Among the pairs of nucleotide sequences utilizable as a pair of primers according to the invention, the pairs of primers selected from the following pairs are preferred:

a) SEQ ID No. 465' GTG TGC TCG ACA TTG GTG TG 3',
and

SEQ ID No. 475' TGG AAT GTT AAC GAG CTG AG 3';

b) SEQ ID No. 465' GTG TGC TCG ACA TTG GTG TG 3',
and

SEQ ID No. 485' CTC GCA GCC ATC TTG GAA TG 3';

c) SEQ ID No. 495' CGC GCG TAA TAC GAC TCA CT 3',
and

SEQ ID No. 465' GTG TGC TCG ACA TTG GTG TG 3';

d) SEQ ID No. 495' CGC GCG TAA TAC GAC TCA CT 3',
and

SEQ ID No. 485' CTC GCA GCC ATC TTG GAA TG 3';
and

e) SEQ ID No. 505' CCT GTC TAC TGC TGT GAG TAC CTT
GT 3',

and

SEQ ID No. 51 5' GCA GTA GAC AGG TCA CTC CGT TGT
CC 3'.

The cloning and the sequencing of the PWD circovirus, type A and B, has allowed it to be identified, after comparative analysis with the nucleotide sequences of other porcine circoviruses, that, among the sequences of fragments of these nucleic acids, were those which are strictly specific to the PWD circovirus of type A, of type B or of type A and B, and those which correspond to a consensus sequence of porcine circoviruses other than the PWD circoviruses of type A and/or B.

There is likewise a great need for nucleotide sequences utilizable as a primer or probe specific to the whole of the other known and nonpathogenic porcine circoviruses.

Said consensus nucleotide sequences specific to all circoviruses, other than PWD circovirus of type A and B, are easily identifiable from FIG. 3 and the sequence SEQ ID No. 15, and are part of the invention.

Among said consensus nucleotide sequences, that which is characterized in that it is part of the following pair of primers is preferred:

a) SEQ ID No. 465' GTG TGC TCG ACA TTG GTG TG 3',
and

SEQ ID No. 525' TGG AAT GTT AAC TAC CTC AA 3'.

The invention likewise comprises a nucleotide sequence according to the invention, characterized in that said sequence is a specific consensus sequence of porcine circovirus other than PWD circovirus of type B and in that it is one of the primers of the following pairs of primers:

a) SEQ ID No. 535' GGC GGC GCC ATC TGT AAC GGT
TT 3',

and

SEQ ID No. 545' GAT GGC GCC GAA AGA CGG GTA
TC 3'.

It is well understood that the present invention likewise relates to specific polypeptides of known porcine circoviruses other than PWD circovirus, encoded by said consensus nucleotide sequences, capable of being obtained by purification from natural polypeptides, by genetic recombination or by chemical synthesis by procedures well known to the person skilled in the art and such as described in particular below. In the same manner, the labeled or unlabeled mono- or polyclonal antibodies directed against said specific polypeptides encoded by said consensus nucleotide sequences are also part of the invention.

It will be possible to use said consensus nucleotide sequences, said corresponding polypeptides as well as said antibodies directed against said polypeptides in procedures or sets for detection and/or identification such as described below, in place of or in addition to nucleotide sequences, polypeptides or antibodies according to the invention, specific to PWD circovirus type A and/or B.

These protocols have been improved for the differential detection of the circular monomeric forms of specific replicative forms of the virion or of the DNA in replication and the dimeric forms found in so-called in-tandem molecular constructs.

The invention additionally relates to the use of a nucleotide sequence according to the invention as a primer or probe for the detection and/or the amplification of nucleic acid sequences.

The nucleotide sequences according to the invention can thus be used to amplify nucleotide sequences, especially by

the PCR technique (polymerase chain reaction) (Erich, 1989; Innis et al., 1990; Rolfs et al., 1991; and White et al., 1997).

These oligodeoxyribonucleotide or oligoribonucleotide primers advantageously have a length of at least 8 nucleotides, preferably of at least 12 nucleotides, and even more preferentially at least 20 nucleotides.

Other amplification techniques of the target nucleic acid can be advantageously employed as alternatives to PCR.

The nucleotide sequences of the invention, in particular the primers according to the invention, can likewise be employed in other procedures of amplification of a target nucleic acid, such as:

the TAS technique (Transcription-based Amplification System), described by Kwok et al. in 1989;

the 3SR technique (Self-Sustained Sequence Replication), described by Guatelli et al. in 1990;

the NASBA technique (Nucleic Acid Sequence Based Amplification), described by Kievitis et al. in 1991;

the SDA technique (Strand Displacement Amplification) (Walker et al., 1992);

the TMA technique (Transcription Mediated Amplification).

The polynucleotides of the invention can also be employed in techniques of amplification or of modification of the nucleic acid serving as a probe, such as:

the LCR technique (Ligase Chain Reaction), described by Landegren et al. in 1988 and improved by Barany et al. in 1991, which employs a thermostable ligase;

the RCR technique (Repair Chain Reaction), described by Segev in 1992;

the CPR technique (Cycling Probe Reaction), described by Duck et al. in 1990;

the amplification technique with Q-beta replicase, described by Miele et al. in 1983 and especially improved by Chu et al. in 1986, Lizardi et al. in 1988, then by Burg et al. as well as by Stone et al. in 1996.

In the case where the target polynucleotide to be detected is possibly an RNA, for example an mRNA, it will be possible to use, prior to the employment of an amplification reaction with the aid of at least one primer according to the invention or to the employment of a detection procedure with the aid of at least one probe of the invention, an enzyme of reverse transcriptase type in order to obtain a cDNA from the RNA contained in the biological sample. The cDNA obtained will thus serve as a target for the primer(s) or the probe(s) employed in the amplification or detection procedure according to the invention.

The detection probe will be chosen in such a manner that it hybridizes with the target sequence or the amplicon generated from the target sequence. By way of sequence, such a probe will advantageously have a sequence of at least 12 nucleotides, in particular of at least 20 nucleotides, and preferably of at least 100 nucleotides.

The invention also comprises the nucleotide sequences utilizable as a probe or primer according to the invention, characterized in that they are labeled with a radioactive compound or with a nonradioactive compound.

The unlabeled nucleotide sequences can be used directly as probes or primers, although the sequences are generally labeled with a radioactive element (^{32}P , ^{35}S , ^3H , ^{125}I) or with a nonradioactive molecule (biotin, acetylaminofluorene, digoxigenin, 5-bromodeoxyuridine, fluorescein) to obtain probes which are utilizable for numerous applications.

Examples of nonradioactive labeling of nucleotide sequences are described, for example, in French Patent No. 78.10975 or by Urdea et al. or by Sanchez-Pescador et al. in 1988.

In the latter case, it will also be possible to use one of the labeling methods described in patents FR-2 422 956 and FR-2 518 755.

The hybridization technique can be carried out in various manners (Matthews et al., 1988). The most general method consists in immobilizing the nucleic acid extract of cells on a support (such as nitrocellulose, nylon, polystyrene) and in incubating, under well-defined conditions, the immobilized target nucleic acid with the probe. After hybridization, the excess of probe is eliminated and the hybrid molecules formed are detected by the appropriate method (measurement of the radioactivity, of the fluorescence or of the enzymatic activity linked to the probe).

The invention likewise comprises the nucleotide sequences according to the invention, characterized in that they are immobilized on a support, covalently or noncovalently.

According to another advantageous mode of employing nucleotide sequences according to the invention, the latter can be used immobilized on a support and can thus serve to capture, by specific hybridization, the target nucleic acid obtained from the biological sample to be tested. If necessary, the solid support is separated from the sample and the hybridization complex formed between said capture probe and the target nucleic acid is then detected with the aid of a second probe, a so-called detection probe, labeled with an easily detectable element.

Another subject of the present invention is a vector for the cloning and/or expression of a sequence, characterized in that it contains a nucleotide sequence according to the invention.

The vectors according to the invention, characterized in that they contain the elements allowing the expression and/or the secretion of said nucleotide sequences in a determined host cell, are likewise part of the invention.

The vector must then contain a promoter, signals of initiation and termination of translation, as well as appropriate regions of regulation of transcription. It must be able to be maintained stably in the host cell and can optionally have particular signals specifying the secretion of the translated protein. These different elements are chosen as a function of the host cell used. To this end, the nucleotide sequences according to the invention can be inserted into autonomous replication vectors within the chosen host, or integrated vectors of the chosen host.

Such vectors will be prepared according to the methods currently used by the person skilled in the art, and it will be possible to introduce the clones resulting therefrom into an appropriate host by standard methods, such as, for example, lipofection, electroporation and thermal shock.

The vectors according to the invention are, for example, vectors of plasmid or viral origin.

A preferred vector for the expression of polypeptides of the invention is baculovirus.

The vector pBS KS in which is inserted the in-tandem DNA sequence of the PWD circovirus type A (or DFP) as deposited at the CNCM on 3 Jul. 1997, under the number I-1891, is likewise preferred.

These vectors are useful for transforming host cells in order to clone or to express the nucleotide sequences of the invention.

The invention likewise comprises the host cells transformed by a vector according to the invention.

These cells can be obtained by the introduction into host cells of a nucleotide sequence inserted into a vector such as

defined above, then the culturing of said cells under conditions allowing the replication and/or expression of the transfected nucleotide sequence.

The host cell can be selected from prokaryotic or eukaryotic systems, such as, for example, bacterial cells (Olins and Lee, 1993), but likewise yeast cells (Buckholz, 1993), as well as animal cells, in particular the cultures of mammalian cells (Edwards and Aruffo, 1993), and especially Chinese hamster ovary (CHO) cells, but likewise the cells of insects in which it is possible to use procedures employing baculoviruses, for example (Luckow, 1993).

A preferred host cell for the expression of the proteins of the invention is constituted by sf9 insect cells.

A more preferred host cell according to the invention is *E. coli*, such as deposited at the CNCM on 3 Jul. 1997, under the number I-1891.

The invention likewise relates to animals comprising one of said transformed cells according to the invention.

The obtainment of transgenic animals according to the invention overexpressing one or more of the genes of PWD circovirus or part of the genes will be preferably carried out in rats, mice or rabbits according to methods well known to the person skilled in the art, such as by viral or nonviral transfections. It will be possible to obtain the transgenic animals overexpressing one or more of said genes by transfection of multiple copies of said genes under the control of a strong promoter of ubiquitous nature, or selective for one type of tissue. It will likewise be possible to obtain the transgenic animals by homologous recombination in embryonic cell strains, transfer of these cell strains to embryos, selection of the affected chimeras at the level of the reproductive lines, and growth of said chimeras.

The transformed cells as well as the transgenic animals according to the invention are utilizable in procedures for preparation of recombinant polypeptides.

It is today possible to produce recombinant polypeptides in relatively large quantity by genetic engineering using the cells transformed by expression vectors according to the invention or using transgenic animals according to the invention.

The procedures for preparation of a polypeptide of the invention in recombinant form, characterized in that they employ a vector and/or a cell transformed by a vector according to the invention and/or a transgenic animal comprising one of said transformed cells according to the invention, are themselves comprised in the present invention.

Among said procedures for preparation of a polypeptide of the invention in recombinant form, the preparation procedures employing a vector, and/or a cell transformed by said vector and/or a transgenic animal comprising one of said transformed cells, containing a nucleotide sequence according to the invention coding for a polypeptide of PWD circovirus, are preferred.

The recombinant polypeptides obtained as indicated above can just as well be present in glycosylated form as in nonglycosylated form and can or cannot have the natural tertiary structure.

A preferred variant consists in producing a recombinant polypeptide used to a "carrier" protein (chimeric protein). The advantage of this system is that it allows a stabilization of and a decrease in the proteolysis of the recombinant product, an increase in the solubility in the course of renaturation in vitro and/or a simplification of the purification when the fusion partner has an affinity for a specific ligand.

More particularly, the invention relates to a procedure for preparation of a polypeptide of the invention comprising the following steps:

a) culture of transformed cells under conditions allowing the expression of a recombinant polypeptide of nucleotide sequence according to the invention;

b) if need be, recovery of said recombinant polypeptide.

When the procedure for preparation of a polypeptide of the invention employs a transgenic animal according to the invention, the recombinant polypeptide is then extracted from said animal.

The invention also relates to a polypeptide which is capable of being obtained by a procedure of the invention such as described previously.

The invention also comprises a procedure for preparation of a synthetic polypeptide, characterized in that it uses a sequence of amino acids of polypeptides according to the invention.

The invention likewise relates to a synthetic polypeptide obtained by a procedure according to the invention.

The polypeptides according to the invention can likewise be prepared by techniques which are conventional in the field of the synthesis of peptides. This synthesis can be carried out in homogeneous solution or in solid phase.

For example, recourse can be made to the technique of synthesis in homogeneous solution described by Houben-Weyl in 1974.

This method of synthesis consists in successively condensing, two by two, the successive amino acids in the order required, or in condensing amino acids and fragments formed previously and already containing several amino acids in the appropriate order, or alternatively several fragments previously prepared in this way, it being understood that it will be necessary to protect beforehand all the reactive functions carried by these amino acids or fragments, with the exception of amine functions of one and carboxyls of the other or vice-versa, which must normally be involved in the formation of peptide bonds, especially after activation of the carboxyl function, according to the methods well known in the synthesis of peptides.

According to another preferred technique of the invention, recourse will be made to the technique described by Merrifield.

To make a peptide chain according to the Merrifield procedure, recourse is made to a very porous polymeric resin, on which is immobilized the first C-terminal amino acid of the chain. This amino acid is immobilized on a resin through its carboxyl group and its amine function is protected. The amino acids which are going to form the peptide chain are thus immobilized, one after the other, on the amino group, which is deprotected beforehand each time, of the portion of the peptide chain already formed, and which is attached to the resin. When the whole of the desired peptide chain has been formed, the protective groups of the different amino acids forming the peptide chain are eliminated and the peptide is detached from the resin with the aid of an acid.

The invention additionally relates to hybrid polypeptides having at least one polypeptide according to the invention, and a sequence of a polypeptide capable of inducing an immune response in man or animals.

Advantageously, the antigenic determinant is such that it is capable of inducing a humoral and/or cellular response.

It will be possible for such a determinant to comprise a polypeptide according to the invention in glycosylated form used with a view to obtaining immunogenic compositions capable of inducing the synthesis of antibodies directed against multiple epitopes. Said polypeptides or their glycosylated fragments are likewise part of the invention.

These hybrid molecules can be formed, in part, of a polypeptide carrier molecule or of fragments thereof accord-

ing to the invention, associated with a possibly immunogenic part, in particular an epitope of the diphtheria toxin, the tetanus toxin, a surface antigen of the hepatitis B virus (patent FR 79 21811), the VP1 antigen of the poliomyelitis virus or any other viral or bacterial toxin or antigen.

The procedures for synthesis of hybrid molecules encompass the methods used in genetic engineering for constructing hybrid nucleotide sequences coding for the polypeptide sequences sought. It will be possible, for example, to refer advantageously to the technique for obtaining of genes coding for fusion proteins described by Minton in 1984.

Said hybrid nucleotide sequences coding for a hybrid polypeptide as well as the hybrid polypeptides according to the invention characterized in that they are recombinant polypeptides obtained by the expression of said hybrid nucleotide sequences are likewise part of the invention.

The invention likewise comprises the vectors characterized in that they contain one of said hybrid nucleotide sequences. The host cells transformed by said vectors, the transgenic animals comprising one of said transformed cells as well as the procedures for preparation of recombinant polypeptides using said vectors, said transformed cells and/or said transgenic animals are, of course, likewise part of the invention.

The polypeptides according to the invention, the antibodies according to the invention described below and the nucleotide sequences according to the invention can advantageously be employed in procedures for the detection and/or identification of PWD circovirus, or of porcine circovirus other than a PWD circovirus, in a biological sample (biological tissue or fluid) capable of containing them. These procedures, according to the specificity of the polypeptides, the antibodies and the nucleotide sequences according to the invention which will be used, will in particular be able to detect and/or to identify a PWD circovirus or a porcine circovirus other than a PWD circovirus or other than the PWD circovirus of type B.

The polypeptides according to the invention can advantageously be employed in a procedure for the detection and/or the identification of PWD circovirus of type A, of type B, of type A or B, or porcine circovirus other than the PWD circovirus of type B, or of porcine circovirus other than the PWD circovirus of type A or B, in a biological sample (biological tissue or fluid) capable of containing them, characterized in that it comprises the following steps:

- a) contacting of this biological sample with a polypeptide or one of its fragments according to the invention (under conditions allowing an immunological reaction between said polypeptide and the antibodies possibly present in the biological sample);
- b) demonstration of the antigen-antibody complexes possibly formed.

In the present description, PWD circovirus, except if a particular mention is indicated, will be understood as designating a PWD circovirus of type A or of type B, and porcine circovirus other than PWD, except if a particular mention is indicated, will be understood as designating a porcine circovirus other than a PWD circovirus of type A and B.

Preferably, the biological sample is formed by a fluid, for example a pig serum, whole blood or biopsies.

Any conventional procedure can be employed for carrying out such a detection of the antigen-antibody complexes possibly formed.

By way of example, a preferred method brings into play immunoenzymatic processes according to the ELISA technique, by immunofluorescence, or radioimmunological processes (RIA) or their equivalent.

Thus, the invention likewise relates to the polypeptides according to the invention, labeled with the aid of an adequate label such as of the enzymatic, fluorescent or radioactive type.

Such methods comprise, for example, the following steps: deposition of determined quantities of a polypeptide composition according to the invention in the wells of a microtiter plate,

introduction into said wells of increasing dilutions of serum, or of a biological sample other than that defined previously, having to be analyzed,

incubation of the microplate,

introduction into the wells of the microtiter plate of labeled antibodies directed against pig immunoglobulins, the labeling of these antibodies having been carried out with the aid of an enzyme selected from those which are capable of hydrolyzing a substrate by modifying the absorption of the radiation of the latter, at least at a determined wavelength, for example at 550 nm,

detection, by comparison with a control test, of the quantity of hydrolyzed substrate.

The invention likewise relates to a kit or set for the detection and/or identification of PWD circovirus, of porcine circovirus other than a PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, characterized in that it comprises the following elements:

a polypeptide according to the invention,

if need be, the reagents for the formation of the medium favorable to the immunological or specific reaction,

if need be, the reagents allowing the detection of the antigen-antibody complexes produced by the immunological reaction between the polypeptide(s) of the invention and the antibodies possibly present in the biological sample, these reagents likewise being able to carry a label, or to be recognized in their turn by a labeled reagent, more particularly in the case where the polypeptide according to the invention is not labeled,

if need be, a biological reference sample (negative control) devoid of antibodies recognized by a polypeptide according to the invention,

if need be, a biological reference sample (positive control) containing a predetermined quantity of antibodies recognized by a polypeptide according to the invention.

The polypeptides according to the invention allow monoclonal or polyclonal antibodies to be prepared which are characterized in that they specifically recognize the polypeptides according to the invention. It will advantageously be possible to prepare the monoclonal antibodies from hybridomas according to the technique described by Kohler and Milstein in 1975. It will be possible to prepare the polyclonal antibodies, for example, by immunization of an animal, in particular a mouse, with a polypeptide or a DNA, according to the invention, associated with an adjuvant of the immune response, and then purification of the specific antibodies contained in the serum of the immunized animals on an affinity column on which the polypeptide which has served as an antigen has previously been immobilized. The polyclonal antibodies according to the invention can also be prepared by purification, on an affinity column on which a polypeptide according to the invention has previously been immobilized, of the antibodies contained in the serum of pigs infected by a PWD circovirus.

The invention likewise relates to mono- or polyclonal antibodies or their fragments, or chimeric antibodies, characterized in that they are capable of specifically recognizing a polypeptide according to the invention.

It will likewise be possible for the antibodies of the invention to be labeled in the same manner as described previously

21

for the nucleic probes of the invention, such as a labeling of enzymatic, fluorescent or radioactive type.

The invention is additionally directed at a procedure for the detection and/or identification of PWD circovirus, of porcine circovirus other than a PWD circovirus, or other than the PWD circovirus of type B, in a biological sample, characterized in that it comprises the following steps:

- a) contacting of the biological sample (biological tissue or fluid) with a mono- or polyclonal antibody according to the invention (under conditions allowing an immunological reaction between said antibodies and the polypeptides of PWD circovirus, of porcine circovirus other than a PWD circovirus, of porcine circovirus other than the PWD circovirus of type B, possibly present in the biological sample);
- b) demonstration of the antigen-antibody complex possibly formed.

Likewise within the scope of the invention is a kit or set for the detection and/or the identification of PWD circovirus, of porcine circovirus other than a PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, characterized in that it comprises the following components:

- a) polyclonal or monoclonal antibody according to the invention, if need be labeled;
- if need be, a reagent for the formation of the medium favorable to the carrying out of the immunological reaction;
- if need be, a reagent allowing the detection of the antigen-antibody complexes produced by the immunological reaction, this reagent likewise being able to carry a label, or being capable of being recognized in its turn by a labeled reagent, more particularly in the case where said monoclonal or polyclonal antibody is not labeled;
- if need be, reagents for carrying out the lysis of cells of the sample tested.

The present invention likewise relates to a procedure for the detection and/or the identification of PWD, of porcine circovirus other than a PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, in a biological sample, characterized in that it employs a nucleotide sequence according to the invention.

More particularly, the invention relates to a procedure for the detection and/or the identification of PWD circovirus, of porcine circovirus other than a PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, in a biological sample, characterized in that it contains the following steps:

- a) if need be, isolation of the DNA from the biological sample to be analyzed;
- b) specific amplification of the DNA of the sample with the aid of at least one primer, or a pair of primers, according to the invention;
- c) demonstration of the amplification products.

These can be detected, for example, by the technique of molecular hybridization utilizing a nucleic probe according to the invention. This probe will advantageously be labeled with a nonradioactive (cold probe) or radioactive element.

For the purposes of the present invention, "DNA of the biological sample" or "DNA contained in the biological sample" will be understood as meaning either the DNA present in the biological sample considered, or possibly the cDNA obtained after the action of an enzyme of reverse transcriptase type on the RNA present in said biological sample.

Another aim of the present invention consists in a procedure according to the invention, characterized in that it comprises the following steps:

22

a) contacting of a nucleotide probe according to the invention with a biological sample, the DNA contained in the biological sample having, if need be, previously been made accessible to hybridization under conditions allowing the hybridization of the probe with the DNA of the sample;

b) demonstration of the hybrid formed between the nucleotide probe and the DNA of the biological sample.

The present invention also relates to a procedure according to the invention, characterized in that it comprises the following steps:

a) contacting of a nucleotide probe immobilized on a support according to the invention with a biological sample, the DNA of the sample having, if need be, previously been made accessible to hybridization, under conditions allowing the hybridization of the probe with the DNA of the sample;

b) contacting of the hybrid formed between the nucleotide probe immobilized on a support and the DNA contained in the biological sample, if need be after elimination of the DNA of the biological sample which has not hybridized with the probe, with a nucleotide probe labeled according to the invention;

c) demonstration of the novel hybrid formed in step b).

According to an advantageous embodiment of the procedure for detection and/or identification defined previously, this is characterized in that, prior to step a), the DNA of the biological sample is first amplified with the aid of at least one primer according to the invention.

The invention is additionally directed at a kit or set for the detection and/or the identification of PWD circovirus, of porcine circovirus other than the PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, characterized in that it comprises the following elements:

- a) a nucleotide probe according to the invention;
- b) if need be, the reagents necessary for the carrying out of a hybridization reaction;
- c) if need be, at least one primer according to the invention as well as the reagents necessary for an amplification reaction of the DNA.

The invention likewise relates to a kit or set for the detection and/or the identification of PWD circovirus, of porcine circovirus other than a PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, characterized in that it comprises the following components:

- a) a nucleotide probe, called a capture probe, according to the invention;
- b) an oligonucleotide probe, called a revealing probe, according to the invention,
- c) if need be, at least one primer according to the invention, as well as the reagents necessary for an amplification reaction of the DNA.

The invention also relates to a kit or set for the detection and/or identification of PWD circovirus, of porcine circovirus other than a PWD circovirus or of porcine circovirus other than the PWD circovirus of type B, characterized in that it comprises the following elements:

- a) at least one primer according to the invention;
- b) if need be, the reagents necessary for carrying out a DNA amplification reaction;
- c) if need be, a component allowing the sequence of the amplified fragment to be verified, more particularly an oligonucleotide probe according to the invention.

The invention additionally relates to the use of a nucleotide sequence according to the invention, of a polypeptide according to the invention, of an antibody according to the invention, of a cell according to the invention, and/or of an animal

transformed according to the invention, for the selection of an organic or inorganic compound capable of modulating, inducing or inhibiting the expression of genes, and/or of modifying the cellular replication of PWD circovirus or capable of inducing or of inhibiting the pathologies linked to an infection by a PWD circovirus.

The invention likewise comprises a method of selection of compounds capable of binding to a polypeptide or one of its fragments according to the invention, capable of binding to a nucleotide sequence according to the invention, or capable of recognizing an antibody according to the invention, and/or capable of modulating, inducing or inhibiting the expression of genes, and/or of modifying the cellular replication of PWD circovirus or capable of inducing or inhibiting the pathologies linked to an infection by a PWD circovirus, characterized in that it comprises the following steps:

- a) contacting of said compound with said polypeptide, said nucleotide sequence, or with a cell transformed according to the invention and/or administration of said compound to an animal transformed according to the invention;
- b) determination of the capacity of said compound to bind to said polypeptide or said nucleotide sequence, or to modulate, induce or inhibit the expression of genes, or to modulate the growth or the replication of PWD circovirus, or to induce or inhibit in said transformed animal the pathologies linked to an infection by PWD circovirus (designated activity of said compound).

The compounds capable of being selected can be organic compounds such as polypeptides or carbohydrates or any other organic or inorganic compounds already known, or novel organic compounds elaborated by molecular modeling techniques and obtained by chemical or biochemical synthesis, these techniques being known to the person skilled in the art.

It will be possible to use said selected compounds to modulate the cellular replication of PWD circovirus and thus to control infection by this virus, the methods allowing said modulations to be determined being well known to the person skilled in the art.

This modulation can be carried out, for example, by an agent capable of binding to a protein and thus of inhibiting or of potentiating its biological activity, or capable of binding to an envelope protein of the external surface of said virus and of blocking the penetration of said virus into the host cell or of favoring the action of the immune system of the infected organism directed against said virus. This modulation can likewise be carried out by an agent capable of binding to a nucleotide sequence of a DNA of said virus and of blocking, for example, the expression of a polypeptide whose biological or structural activity is necessary for the replication or for the proliferation of said virus host cells to host cells in the host animal.

The invention relates to the compounds capable of being selected by a selection method according to the invention.

The invention likewise relates to a pharmaceutical composition comprising a compound selected from the following compounds:

- a) a nucleotide sequence according to the invention;
- b) a polypeptide according to the invention;
- c) a vector, a viral particle or a cell transformed according to the invention;
- d) an antibody according to the invention;
- e) a compound capable of being selected by a selection method according to the invention;

possibly in combination with a pharmaceutically acceptable vehicle and, if need be, with one or more adjuvants of the appropriate immunity.

The invention also relates to an immunogenic and/or vaccine composition, characterized in that it comprises a compound selected from the following compounds:

- a) a nucleotide sequence according to the invention;
- b) a polypeptide according to the invention;
- c) a vector or a viral particle according to the invention; and
- d) a cell according to the invention.

In one embodiment, the vaccine composition according to the invention is characterized in that it comprises a mixture of at least two of said compounds a), b), c) and d) above and in that one of the two said compounds is related to the PWD circovirus of type A and the other is related to the PWD circovirus of type B.

In another embodiment of the invention, the vaccine composition is characterized in that it comprises at least one compound a), b), c), or d) above which is related to PWD circovirus of type B. In still another embodiment, the vaccine composition is characterized in that it comprises at least one compound a), b), c), or d) above which is related to PWD circovirus of type B ORF'2.

A compound related to the PWD circovirus of type A or of type B is understood here as respectively designating a compound obtained from the genomic sequence of the PWD circovirus of type A or of type B.

The invention is additionally aimed at an immunogenic and/or vaccine composition, characterized in that it comprises at least one of the following compounds:

- a nucleotide sequence SEQ ID No. 23, SEQ ID No. 25, or one of their fragments or homologues;
- a polypeptide of sequence SEQ ID No. 24, SEQ ID No. 26, or one of their fragments, or a modification thereof;
- a vector or a viral particle comprising a nucleotide sequence SEQ ID No. 23, SEQ ID No. 25, or one of their fragments or homologues;
- a transformed cell capable of expressing a polypeptide of sequence SEQ ID No. 24, SEQ ID No. 26, or one of their fragments, or a modification thereof; or
- a mixture of at least two of said compounds.

The invention also comprises an immunogenic and/or vaccine composition according to the invention, characterized in that it comprises said mixture of at least two of said compounds as a combination product for simultaneous, separate or protracted use for the prevention or the treatment of infection by a PWD circovirus, especially of type B.

In a preferred embodiment, the vaccine composition according to the invention comprises the mixture of the following compounds:

- a pDNA3 plasmid containing a nucleic acid of sequence SEQ ID No. 23;
- a pDNA3 plasmid containing a nucleic acid of sequence SEQ ID No. 25;
- a pDNA3 plasmid containing a nucleic acid coding for the GM-CSF protein;
- a recombinant baculovirus containing a nucleic acid of sequence SEQ ID No. 23;
- a recombinant baculovirus containing a nucleic acid of sequence SEQ ID No. 25; and
- if need be, an adjuvant of the appropriate immunity, especially the adjuvant AIF™.

The invention is likewise directed at a pharmaceutical composition according to the invention, for the prevention or the treatment of an infection by a PWD circovirus.

The invention is also directed at a pharmaceutical composition according to the invention for the prevention or the treatment of an infection by the PWD circovirus of type B.

The invention likewise concerns the use of a composition according to the invention, for the preparation of a medicament intended for the prevention or the treatment of infection by a PWD circovirus, preferably by the PWD circovirus of type B.

Under another aspect, the invention relates to a vector, a viral particle or a cell according to the invention, for the treatment and/or the prevention of a disease by gene therapy.

Finally, the invention comprises the use of a vector, of a viral particle or of a cell according to the invention for the preparation of a medicament intended for the treatment and/or the prevention of a disease by gene therapy.

The polypeptides of the invention entering into the immunogenic or vaccine compositions according to the invention can be selected by techniques known to the person skilled in the art such as, for example, depending on the capacity of said polypeptides to stimulate the T cells, which is translated, for example, by their proliferation or the secretion of interleukins, and which leads to the production of antibodies directed against said polypeptides.

In pigs, as in mice, in which a weight dose of the vaccine composition comparable to the dose used in man is administered, the antibody reaction is tested by taking of the serum followed by a study of the formation of a complex between the antibodies present in the serum and the antigen of the vaccine composition, according to the usual techniques.

The pharmaceutical compositions according to the invention will contain an effective quantity of the compounds of the invention, that is to say in sufficient quantity of said compound(s) allowing the desired effect to be obtained, such as, for example, the modulation of the cellular replication of PWD circovirus. The person skilled in the art will know how to determine this quantity, as a function, for example, of the age and of the weight of the individual to be treated, of the state of advancement of the pathology, of the possible secondary effects and by means of a test of evaluation of the effects obtained on a population range, these tests being known in these fields of application.

According to the invention, said vaccine combinations will preferably be combined with a pharmaceutically acceptable vehicle and, if need be, with one or more adjuvants of the appropriate immunity.

Today, various types of vaccines are available for protecting animals or man against infectious diseases: attenuated living microorganisms (*M. bovis*—BCG for tuberculosis), inactivated microorganisms (influenza virus), acellular extracts (*Bordetella pertussis* for whooping cough), recombinant proteins (surface antigen of the hepatitis B virus), polysaccharides (pneumococcal). Vaccines prepared from synthetic peptides or genetically modified microorganisms expressing heterologous antigens are in the course of experimentation. More recently still, recombinant plasmid DNAs carrying genes coding for protective antigens have been proposed as an alternative vaccine strategy. This type of vaccination is carried out with a particular plasmid originating from a plasmid of *E. coli* which does not replicate in vivo and which codes uniquely for the vaccinating protein. Animals have been immunized by simply injecting the naked plasmid DNA into the muscle. This technique leads to the expression of the vaccine protein in situ and to an immune response of cellular type (CTL) and of humoral type (antibody). This double induction of the immune response is one of the principal advantages of the vaccination technique with naked DNA.

The vaccine compositions comprising nucleotide sequences or vectors into which are inserted said sequences are especially described in the international application No. WO 90/11092 and likewise in the international application No. WO 95/11307.

The constitutive nucleotide sequence of the vaccine composition according to the invention can be injected into the host after having been coupled to compounds which favor the penetration of this polynucleotide into the interior of the cell or its transport to the cell nucleus. The resultant conjugates can be encapsulated in polymeric microparticles, as described in the international application No. WO 94/27238 (Medisorb Technologies International).

According to another embodiment of the vaccine composition according to the invention, the nucleotide sequence, preferably a DNA, is complexed with DEAE-dextran (Pagano et al., 1967) or with nuclear proteins (Kaneda et al., 1989), with lipids (Felgner et al., 1987) or encapsulated in liposomes (Fraley et al., 1980) or else introduced in the form of a gel facilitating its transfection into the cells (Midoux et al., 1993, Pastore et al., 1994). The polynucleotide or the vector according to the invention can also be in suspension in a buffer solution or be combined with liposomes.

Advantageously, such a vaccine will be prepared according to the technique described by Tacson et al. or Huygen et al. in 1996 or alternatively according to the technique described by Davis et al. in the international application No. WO 95/11307.

Such a vaccine can likewise be prepared in the form of a composition containing a vector according to the invention, placed under the control of regulation elements allowing its expression in man or animal. It will be possible, for example, to use, by way of in vivo expression vector of the polypeptide antigen of interest, the plasmid pcDNA3 or the plasmid pcDNA1/neo, both marketed by Invitrogen (R&D Systems, Abingdon, United Kingdom). It is also possible to use the plasmid V1Jns.tPA, described by Shiver et al. in 1995. Such a vaccine will advantageously comprise, apart from the recombinant vector, a saline solution, for example a sodium chloride solution.

Pharmaceutically acceptable vehicle is understood as designating a compound or a combination of compounds entering into a pharmaceutical composition or vaccine which does not provoke secondary reactions and which allows, for example, the facilitation of the administration of the active compound, an increase in its duration of life and/or its efficacy in the body, an increase in its solubility in solution or alternatively an improvement in its conservation. These pharmaceutically acceptable vehicles are well known and will be adapted by the person skilled in the art as a function of the nature and of the mode of administration of the chosen active compound.

As far as the vaccine formulations are concerned, these can comprise adjuvants of the appropriate immunity which are known to the person skilled in the art, such as, for example, aluminum hydroxide, a representative of the family of muramyl peptides such as one of the peptide derivatives of N-acetyl muramyl, a bacterial lysate, or alternatively Freund's incomplete adjuvant.

These compounds can be administered by the systemic route, in particular by the intravenous route, by the intramuscular, intradermal or subcutaneous route, or by the oral route. In a more preferred manner, the vaccine composition comprising polypeptides according to the invention will be administered by the intramuscular route, through the food or by nebulization several times, staggered over time.

Their administration modes, dosages and optimum pharmaceutical forms can be determined according to the criteria

generally taken into account in the establishment of a treatment adapted to an animal such as, for example, the age or the weight, the seriousness of its general condition, the tolerance to the treatment and the secondary effects noted. Preferably, the vaccine of the present invention is administered in an amount that is protective against piglet weight loss disease.

For example, in the case of a vaccine according to the present invention comprising a polypeptide encoded by a nucleotide sequence of the genome of PCV, or a homologue or fragment thereof, the polypeptide will be administered one time or several times, spread out over time, directly or by means of a transformed cell capable of expressing the polypeptide, in an amount of about 0.1 to 10 µg per kilogram weight of the animal, preferably about 0.2 to about 5 µg/kg, more preferably about 0.5 to about 2 µg/kg for a dose.

The present invention likewise relates to the use of nucleotide sequences of PWD circovirus according to the invention for the construction of autoreplicative retroviral vectors and the therapeutic applications of these, especially in the field of human gene therapy *in vivo*.

The feasibility of gene therapy applied to man no longer needs to be demonstrated and this relates to numerous therapeutic applications like genetic diseases, infectious diseases and cancers. Numerous documents of the prior art describe the means of employing gene therapy, especially through viral vectors. Generally speaking, the vectors are obtained by deletion of at least some of the viral genes which are replaced by the genes of therapeutic interest. Such vectors can be propagated in a complementation line which supplies in trans the deleted viral functions in order to generate a defective viral vector particle for replication but capable of infecting a host cell. To date, the retroviral vectors are amongst the most widely used and their mode of infection is widely described in the literature accessible to the person skilled in the art.

The principle of gene therapy is to deliver a functional gene, called a gene of interest, of which the RNA or the corresponding protein will produce the desired biochemical effect in the targeted cells or tissues. On the one hand, the insertion of genes allows the prolonged expression of complex and unstable molecules such as RNAs or proteins which can be extremely difficult or even impossible to obtain or to administer directly. On the other hand, the controlled insertion of the desired gene into the interior of targeted specific cells allows the expression product to be regulated in defined tissues. For this, it is necessary to be able to insert the desired therapeutic gene into the interior of chosen cells and thus to have available a method of insertion capable of specifically targeting the cells or the tissues chosen.

Among the methods of insertion of genes, such as, for example, microinjection, especially the injection of naked plasmid DNA (Derse, D. et al., 1995, and Zhao, T. M. et al., 1996), electroporation, homologous recombination, the use of viral particles, such as retroviruses, is widespread. However, applied *in vivo*, the gene transfer systems of recombinant retroviral type at the same time have a weak infectious power (insufficient concentration of viral particles) and a lack of specificity with regard to chosen target cells.

The production of cell-specific viral vectors, having a tissue-specific tropism, and whose gene of interest can be translated adequately by the target cells, is realizable, for example, by fusing a specific ligand of the target host cells to the N-terminal part of a surface protein of the envelope of PWD circovirus. It is possible to mention, for example, the construction of retroviral particles having the CD4 molecule on the surface of the envelope so as to target the human cells infected by the HIV virus (YOUNG, J. A. T. et al., Sciences 1990, 250, 1421-1423), viral particles having a peptide hor-

mone fused to an envelope protein to specifically infect the cells expressing the corresponding receptor (KASAHARA, N. et al., Sciences 1994, 266, 1373-1376) or else alternatively viral particles having a fused polypeptide capable of immobilizing on the receptor of the epidermal growth factor (EGF) (COSSET, F. L. et al., J. of Virology 1995, 69, 10, 6314-6322). In another approach, single-chain fragments of antibodies directed against surface antigens of the target cells are inserted by fusion with the N-terminal part of the envelope protein (VALSESIA-WITTMAN, S. et al., J. of Virology 1996, 70, 3, 2059-2064; TEARINA CHU, T. H. et al., J. of Virology 1997, 71, 1, 720-725).

For the purposes of the present invention, a gene of interest in use in the invention can be obtained from a eukaryotic or prokaryotic organism or from a virus by any conventional technique. It is, preferably, capable of producing an expression product having a therapeutic effect and it can be a product homologous to the cell host or, alternatively, heterologous. In the scope of the present invention, a gene of interest can code for an (i) intracellular or (ii) membrane product present on the surface of the host cell or (iii) secreted outside the host cell. It can therefore comprise appropriate additional elements such as, for example, a sequence coding for a secretion signal. These signals are known to the person skilled in the art.

In accordance with the aims pursued by the present invention, a gene of interest can code for a protein corresponding to all or part of a native protein as found in nature. It can likewise be a chimeric protein, for example arising from the fusion of polypeptides of various origins or from a mutant having improved and/or modified biological properties. Such a mutant can be obtained, by conventional biological techniques, by substitution, deletion and/or addition of one or more amino acid residues.

It is very particularly preferred to employ a gene of therapeutic interest coding for an expression product capable of inhibiting or retarding the establishment and/or the development of a genetic or acquired disease. A vector according to the invention is in particular intended for the prevention or for the treatment of cystic fibrosis, of hemophilia A or B, of Duchenne's or Becker's myopathy, of cancer, of AIDS and of other bacteria or infectious diseases due to a pathogenic organism: virus, bacteria, parasite or prion. The genes of interest utilizable in the present invention are those which code, for example, for the following proteins:

- a cytokine and especially an interleukin, an interferon, a tissue necrosis factor and a growth factor and especially a hematopoietic growth factor (G-CSF, GM-CSF),
- a factor or cofactor involved in clotting and especially factor VIII, von Willebrand's factor, antithrombin III, protein C, thrombin and hirudin,
- an enzyme or an enzyme inhibitor such as the inhibitors of viral proteases,
- an expression product of a suicide gene such as thymidine kinase of the HSV virus (herpesvirus) of type 1,
- an activator or an inhibitor of ion channels,
- a protein of which the absence, the modification or the deregulation of expression is responsible for a genetic disease, such as the CFTR protein, dystrophin or minidystrophin, insulin, ADA (adenosine diaminase), glucocerebrosidase and phenylhydroxylase,
- a protein capable of inhibiting the initiation or the progression of cancers, such as the expression products of tumor suppressor genes, for example the P53 and Rb genes,
- a protein capable of stimulating an immune or an antibody response, and

a protein capable of inhibiting a viral infection or its development, for example the antigenic epitopes of the virus in question or altered variants of viral proteins capable of entering into competition with the native viral proteins.

The invention thus relates to the vectors characterized in that they comprise a nucleotide sequence of PWD circovirus according to the invention, and in that they additionally comprise a gene of interest.

The present invention likewise relates to viral particles generated from said vector according to the invention. It additionally relates to methods for the preparation of viral particles according to the invention, characterized in that they employ a vector according to the invention, including viral pseudoparticles (VLP, virus-like particles).

The invention likewise relates to animal cells transfected by a vector according to the invention.

Likewise comprised in the invention are animal cells, especially mammalian, infected by a viral particle according to the invention.

The present invention likewise relates to a vector, a viral particle or a cell according to the invention, for the treatment and/or the prevention of a genetic disease or of an acquired disease such as cancer or an infectious disease. The invention is likewise directed at a pharmaceutical composition comprising, by way of therapeutic or prophylactic agent, a vector or a cell according to the invention, in combination with a vehicle acceptable from a pharmaceutical point of view.

Other characteristics and advantages of the invention appear in the examples and the figures.

The invention is described in more detail in the following illustrative examples. Although the examples may represent only selected embodiments of the invention, it should be understood that the following examples are illustrative and not limiting.

EXAMPLES

Example 1

Cloning, Sequencing and Characterization of the PWD Circovirus of Type A (PCVA)

1. Experimental Procedures

Experimental reproduction of the infection and its syndrome are provided (cf. FIG. 1).

A first test was carried out with pigs from a very well-kept farm, but affected by piglet weight loss disease (PWD), likewise called fatal piglet wasting (FPW). Tests carried out with SPF (specific pathogen-free) pigs showed a transfer of contaminant(s) finding expression in a complex pathology combining hyperthermia, retardation of growth, diarrhea and conjunctivitis. The PDRS (porcine dysgenic and respiratory syndrome) virus, an infectious disease due to an arteriovirus) was rapidly isolated from breeding pigs and contact pigs. It should have been possible to attribute all the clinical signs to the presence of the PDRS virus. However, two farm pigs presented signs of FPW without the PDRS virus being isolated. The histological analyses and blood formulas, however, showed that these pigs were suffering from an infectious process of viral origin.

In a second test, 8-week SPF pigs were inoculated by the intratracheal route with organ homogenates of two farm pigs suffering from FPW. The inoculated pigs exhibited hyperthermia 8 to 9 days post-infection, then their growth was retarded. Other SPF pigs, placed in contact, had similar, attenuated signs 30 days after the initial experiment. No sero-conversion with respect to a European or Canadian strain of PDRS virus was recorded in these animals.

A third test allowed the syndrome to be reproduced from samples taken from the pigs of the second test.

Conclusion

The syndrome is reproduced under the experimental conditions. It is determined by at least one infectious agent, which is transmittable by direct contact. The clinical constants are a sometimes high hyperthermia (greater than or equal to 41.5° C.) which develops 8 to 10 days after infection. Retardation of the growth can be observed. The other signs are a reversal of the blood formula (reversal of the lymphocyte/polynuclear ratio from 70/30 to 30/70) and frequent lesions on the ganglia, especially those draining the respiratory apparatus (ganglionic hypertrophy, loss of structure with necrosis and infiltration by mononucleated or plurinucleated giant cells).

2. Laboratory Studies

Various cell supports including primary pig kidney cells or cell lines, pig testicle cells, monkey kidney cells, pig lymphocytes, pig alveolar macrophages and circulating blood monocytes were used to demonstrate the possible presence of a virus. No cytopathic effect was demonstrated in these cells. On the other hand, the use of a serum of a pig sick after experimental infection allowed an intracellular antigen to be revealed in the monocytes, the macrophages and approximately 10% of pig kidney (PK) cells infected with organ homogenates. This indirect revealing was carried out kinetically at different culture times. It is evident from this that the antigen initially appears in the nucleus of the infected cells before spreading into the cytoplasm. The successive passages in cell culture did not allow the signal to be amplified.

Under electron microscopy on organ homogenates, spherical particles labeled specifically by the serum of sick pigs, infected under the experimental conditions, were visualized. The size of these particles is estimated at 20 nm.

After two passages of these organ homogenates over pig lymphocytes and then three passages over pig kidney or testicle cells, a cytopathic effect developed and was amplified. An adenovirus was visualized in the electron microscope, which, under the experimental conditions, did not reproduce FPW (only a hyperthermia peak was noted 24 to 48 hours after infection, and then nothing more).

It has been possible to demonstrate DNA bands in certain samples of pigs infected under the experimental conditions and having exhibited signs of the disease (results not shown). A certain connection exists between the samples giving a positive result in cell culture and those having a DNA band.

Conclusion

At least two types of virus were demonstrated in the organ homogenates from pigs suffering from FPW. One is an adenovirus, but by itself alone it does not reproduce the disease. The other type of virus is a circovirus and is associated with FPW. This circovirus, of which two types have been isolated and sequenced, designated below PWD circovirus type A (or PCVA) and PWD circovirus of type B (or PCVB) have mutations with respect to the known sequences of circovirus which are nonpathogenic for the pig.

3. Cloning and Sequencing of the DNA of the PWD Circovirus of Type A

Cloning and sequencing of the DNA of PHD circovirus Type A is accomplished by extraction of the replicative form (RF) DNA, followed by cleavage by the Kpn I enzyme and amplification by a pair of primers flanking the Kpn I restriction site. The two strands of DNA are sequenced at least twice by the Sanger method.

The nucleic sequence of the strand of (+) polarity of the genome of the PWD circovirus of type A (or PCVA), strain FPW, is represented by the sequence SEQ ID No. 1 in the list of sequences, the nucleic acid sequence of the strand of (-)

polarity of the genome of the PWD circovirus of type A (or PCVA) being represented by the nucleic acid sequence 3'→5' of FIG. 3 or by the sequence SEQ ID No. 5 (represented according to the orientation 5'→3') in the list of sequences.

The amino acid sequences SEQ ID No. 10, SEQ ID No. 12 and SEQ ID No. 14 of the list of sequences respectively represent the sequences of proteins encoded by the nucleic sequences of the 3 open reading frames SEQ ID No. 9 (ORF1), corresponding to the REP protein, SEQ ID No. 11 (ORF2) and SEQ ID No. 13 (ORF3), determined from the sequence SEQ ID No. 1 of the strand of (+) polarity or of the nucleic sequence SEQ ID No. 5 of the strand of (-) polarity of the genome of the PWD circovirus of type A.

4. Comparison of the Nucleotide Sequences and Amino Acids of the PWD Circovirus of Type A (or Associated with PWD) which are Obtained with the Corresponding Sequences of MEEHAN and MANKERTZ Circoviruses of Porcine Cell Lines.

DNA sequences are analyzed using, DNASIS software.

Sequences of Oligonucleotides used as Primers or Probes in the Detection and/or Identification Procedures

1. Specific Detection of the PWD Circovirus of Type A:

SEQ ID No. 46 primer PCV 5: 5' GTG TGC TCG ACA TTG GTG TG 3';

SEQ ID No. 47 primer PCV 10: 5' TGG AAT GTT AAC GAG CTG AG 3';

2. Specific Detection of the Circovirus of the Cell Lines:

SEQ ID No. 46 primer PCF 5: 5' GTG TGC TCG ACA TTG GTG TG 3';

SEQ ID No. 52 primer MEE 1: 5' TGG AAT GTT AAC TAC CTC AA 3';

3. Differential Detection:

the pairs of primers used are those described, for example, in the paragraphs 1 and 2 above;

4. Detection of the Monomeric Circular Replicative Forms RF:

SEQ ID No. 46 primer PCV 5: 5' GTG TGC TCG ACA TTG GTG TG 3';

SEQ ID No. 48 primer PCV 6: 5' CTC GCA GCC ATC TTG GAA TG 3';

5. Detection of the Vectors Carrying the Dimers in Tandem: Nar Dimer:

SEQ ID No. 49 primer KS 620: 5' CGC GCG TAA TAC GAC TCA CT 3';

SEQ ID No. 46 primer PCV 5: 5' GTG TGC TCG ACA TTG GTG TG 3';

Kpn Dimer:

SEQ ID No. 49 primer KS 620: 5' CGC GCG TAA TAC GAC TCA CT 3';

SEQ ID No. 48 primer PCV 6: 5' CTC GCA GCC ATC TTG GAA TG 3';

6. Differential Detection:

The pairs of primers used are those described, for example, in paragraphs 4 and 5 above.

The procedures using the pairs or primers described in paragraphs 4 and 5 are of particular interest for differentially detecting the circular monomeric forms of specific replicative forms of the virion or of the DNA in replication and the dimeric forms found in the so-called in-tandem molecular constructs.

The in-tandem constructs of the viral genome (dimers) such as the constructs used for the preparation of the pBS KS+tandem PCV Kpn I vector, deposited at the CNCM under the number I-1891, 3 Jul. 1997 (*E. coli* transformed by said vector) are very interesting for their use in methods of production of sufficient quantity of an inoculum formed of DNA, intended for the virus production, this in the absence of a satisfactory virus production protocol in a cell system. These said methods of production using in-tandem constructs of the viral genome will allow the virulence factors to be studied by mutation and by way of consequence will be able to be used for the production of a collection of viruses carrying the mutations indicated in the construction of vectors which will have the appropriate tropism and virulence. These vectors with autoreplicative structure have the sought gene transfer properties, especially for their applications in gene therapy, and in vaccinology.

Western-Blot Analysis of Recombinant Proteins of the PWD Circovirus of Type A

The results were obtained using a specific antiserum of the PWD circovirus produced during test 1 (cf. FIG. 1).

Type of Products Analyzed

The analyses were carried out on cell extracts of Sf9 cells obtained after infection by the recombinant baculovirus PCV ORF 1.

The culture of Sf9 cells was carried out in a 25 cm² Petri dish according to the standard culture methods for these cells. After centrifugation, the cell pellets are taken up with 300 µl of PBS buffer (phosphate saline buffer).

Electrophoresis (PAGE-SDS)

The electrophoresis is carried out on the cell extracts of Sf9 cells obtained previously on 5 samples (cf. Table 1 below) under the following conditions:

% polyacrylamide gel: 8%; conditions: denaturing
Voltage: 80 V; duration: 135 mn.

TABLE 1

	Nature of the samples subjected to electrophoresis				
	Well No.				
	1	2	3	4	5
Sample applied	PM Rainbow	Raoul 24 h	Raoul 48 h	Raoul 72 h	Raoul 96 h
µl of sample	10	15	15	15	15
µl of Laemmli 4x	0	5	5	5	5

Legends to Table 1:

Laemmli 4x: loading buffer
PM Rainbow: molecular-weight markers (35, 52, 77, 107, 160 and 250 kD)
Raoul 24 h, 48 h, 72 h and 96h: expression products of the ORF1 of the PWD circovirus of type A.

Western Blot

After electrophoresis, the bands obtained in the different wells are transferred to nitrocellulose membrane for 1 h at 100 v in a TGM buffer (tris-glycine-methanol).

The Western blot is carried out under the following conditions:

1) Saturation with a solution containing 5% of skimmed milk; 0.05% of Tween 20 in a TBS 1× buffer (tris buffer saline) for 30 min.

2) 1st Antibody:

10 ml of PWD anticircovirus antibody of type A are added diluted to 1/100, then the reaction mixture is incubated for one night at 4° C. Three washes of 10 min in TBS 1× are carried out.

3) 2nd Antibody:

10 ml of pig rabbit P164 antibody anti-immunoglobulins, coupled to peroxidase (Dakopath), are added diluted to 1/100, then the reaction medium is incubated for 3 hours at 37° C. Three washes of 10 min in TBS 1× are carried out.

4) Visualization

The substrate 4-chloro-1-naphthol in the presence of oxygenated water is used for visualization.

Results

The results are shown in FIG. 7.

Kinetics of Appearance of Antibodies Specific for the REP Recombinant Protein of the PWD Circovirus of Type A Expressed in Baculovirus After Infection of Pigs by the PWD Circovirus of Type A (Test 4, cf. FIG. 1)

After infection of the pigs, a sample of serum of each of the infected pigs is taken at different periods expressed in the table by the date of taking (carried out here in the same year) and is then analyzed by Western blot.

The visualization of the specific antibodies is carried out in the manner described previously.

The results obtained are shown by Table 2 below.

TABLE 2

Kinetics of appearance of specific antibodies								
Sample	Pigs	10/6	16/06	23/06	01/07	08/07	15/07	21/07
A3	1							Neg.
Control	2							Neg.
B2 Infec.	1	Neg.	Neg.	Neg.	+	+	++	+++
RP+	2	Neg.	Neg.	Neg.	Neg.	Neg.	Neg.	Neg.
	3	Neg.	Neg.	Neg.	Neg.	+	+	+
	4	Neg.	Neg.	Neg.	Neg.	Neg.	Neg.	++

Legends to Table 2:

A3 control: uninfected control animals;

B2 Infec. RP+: animals infected with pig kidney (PK) cells containing the circovirus;

Neg.: negative;

+, ++, +++: intensity scale of the positive reaction;

10/06, 16/06, 23/06, 01/07, 08/07, 15/07, 21/07: dates expressed in day/month on which the different withdrawals of serum were carried out.

Example 2

Cloning, Sequencing and Characterization of the Type B PWD Circovirus (PCVB)

The techniques used for cloning, sequencing and characterization of the type B PWD circovirus (PCVB) are those used in Example 1 above for the type A PWD circovirus (PCVA).

The nucleic acid sequence of the strand of (+) polarity of the genome of the PWD circovirus of type B (or PCVB) is represented by the sequence SEQ ID No. 15 in the sequence listing, the nucleic acid sequence of the strand of (-) polarity of the genome of the PWD circovirus of type B (or PCVB) being represented by the nucleic acid sequence 3'→5' of FIG.

8 or by the sequence SEQ ID No. 19 (represented according to the orientation 5'→3') in the sequence listing.

The amino acid sequences, SEQ ID No. 24, SEQ ID No. 26 and SEQ ID No. 28 of the sequence listing, respectively, represent the sequences of the proteins encoded by the nucleic sequences of the 3 open reading frames SEQ ID No. 23 (ORF'1), corresponding to the REP protein, SEQ ID No. 25 (ORF'2) and SEQ ID No. 27 (ORF'3), determined from the sequence SEQ ID No. 15 of the strand of (+) polarity or from the nucleic sequence SEQ ID No. 19 of the strand of (-) polarity of the genome of the PWD circovirus of type B.

Example 3

Comparative Analysis of Nucleotide Sequences (ORF1, ORF2 and Genomic) and Amino Acid Sequences Encoded by the ORF1 and the ORF2 of the PWD Circoviruses of Type A (PCVA) and of Type B (PCVB)

The results expressed in % of homology are shown in Tables 3 and 4 below.

TABLE 3

Compared analysis of the amino acid sequences		
% homology	ORF1	ORF2
PCVA/PCVB	80.4	56.2

TABLE 4

Compared analysis of the nucleotide sequences				
% homology	Genomic	ORF1	ORF2	The remainder
PCVA/PCVB	70.4	80.4	60.1	66.1

Example 4

Observation of the Disease and Reproduction of the Disease Under Experimental Conditions

a) Test No. 1: Observation of the Disease

The objective is to take breeding animals at the start of disease and to place them under experimental conditions to follow the progression of the pathology and describe all the clinical signs thereof. This first test was carried out on 3 breeding pigs aged 10 weeks of which 2 were already ill (suffering from wasting), and on 3 other pigs aged 13 weeks, not having signs of disease. The clinical observation was spread over a period of 37 days. Two pigs of 10 weeks wasted rapidly (pigs 1 and 2, FIG. 9) and had to be painlessly killed 5 and 6 days after their arrival. A single pig exhibited hyperthermia over 5 days and diarrhea. Two other pigs exhibited dyspnea and cough, of which one additionally had hyperthermia, greater than 41° C., for the two first days of its stay. Another pig had retarded growth in the second week (pig 6, FIG. 9), without any other clinical sign being recorded. On the lesional level, 5 pigs out of 6 exhibited macroscopic lesions of gray pneumonia, the sixth exhibited cicatricial lesions on the lung.

b) Test No. 2: Reproduction of the Disease from Inocula Prepared in Farm Pigs.

The two sick pigs in test 1 served to prepare inocula which were tested in test 2 on specific-pathogen-free (SPF) pigs. The SPF pigs were aged 9 weeks at the time of inoculation. The clinical and lesional results are shown in Table 5.

TABLE 5

	Test Measurement					
	2	3	4	5	6	7
Status of the pigs	SPF CNEVA	SPF field	SPF CNEVA	SPF CNEVA	Conventional	Conventional
Age	9 weeks	6 weeks	5 weeks	5 weeks	5 weeks	6-7 weeks
Number	4	6	12	8	8	8
Inoculation route	Intratracheal route	Intratracheal route	Intratracheal + intramuscular route	Intratracheal + intramuscular route	Intratracheal + intramuscular route	Intratracheal + intramuscular route
Inoculum titer per pig	ND*	ND*	10 ^{4.53} TCID ₅₀ per ml: 1 ml IM + 5 ml IT	10 ^{4.53} TCID ₅₀ per ml: 1 ml IM + 5 ml IT	10 ^{4.53} TCID ₅₀ per ml: 1 ml IM + 5 ml IT	10 ^{4.53} TCID ₅₀ per ml: 1 ml IM + 5 ml IT
Start of hyperthermia	10 days post-infection	9-13 days post-infection	12-13 days post-infection	9-14 days post-infection	8-12 days post-infection	12 days post-infection
% of pigs in hyperthermia**	100%	83%	92%	100%	75%	88%
Number of days per pig**	7	4.5	3.3	5.8	7.5	11.6
Maximum temperatures***	40.4 to 41.7° C.	40.6 to 42.3° C.	40.2 to 41.6° C.	40.3 to 40.8° C.	40.6 to 42° C.	40.2 to 41.9° C.
Hyperthermia**** % per week						
W1	3.5 (3.5)	17 (36)	7 (5)	37 (17)	16 (17)	20 (28)
W2	<u>42 (3.5)</u>	7 (13)	<u>13 (1)</u>	<u>21 (3)</u>	<u>52 (10)</u>	37 (28)
W3	<u>35 (3.5)</u>	<u>33 (10)</u>	<u>28 (7)</u>	<u>62 (2)</u>	<u>34 (12)</u>	<u>79 (17)</u>
W4	<u>21 (3.5)</u>	<u>28 (7)</u>	5 (0)	6 (3)	25 (22)	<u>55 (3)</u>
DMG:						
W1	928 (1053)	417 (357)	564 (620)	650 (589)	401 (407)	509 (512)
W2	<u>678 (1028)</u>	<u>428 (617)</u>	<u>503 (718)</u>	612 (584)	<u>294 (514)</u>	410 (310)
W3	<u>661 (1000)</u>	771 (642)	<u>381 (657)</u>	<u>520 (851)</u>	<u>375 (586)</u>	435 (440)
W4	<u>786 (1100)</u>	<u>550 (657)</u>	764 (778)	641 (696)	<u>473 (610)</u>	<u>451 (681)</u>
Contact pigs transmission	Yes to 100%	Yes to 75%	Not tested	Not tested	Not tested	Not tested
% of pulmonary lesions	25	75	0	25	25	12
% of ganglionic lesions	17	33	67	25	50	12

*ND: not determined,

**hyperthermia when the temperature is greater than 40° C.,

***range of maximum temperatures recorded at the individual level,

****the percentage corresponds to the number of temperature recordings greater than 40° C. divided by the total number of temperature recordings in the week on all of the pigs.

In this test, there was no wasting, at the very most a retardation of the growth in the second, third or fourth week after infection. These data illustrate that certain breeding conditions probably favor the expression of the disease.

c) Tests No. 3 to No. 7: Reproduction of the Experimental Tests

The increase in the number of the experimental tests on pigs had the mastering and better characterization of the experimental model as an objective. All of the results are presented in Table 5.

Under the experimental conditions, PWD is thus characterized by a long incubation, of 8 to 14 days, true hyperthermia over 2 to 8 days, a decrease in food consumption and a retardation of the increase in weight on the second, third or fourth week post-infection. The lesional table associated with this clinical expression includes, in the main, ganglionic hypertrophy and lesions of pneumonia.

Conclusion

The perfection of this experimental model allows the direct etiological role of the PWD circovirus in the disease to be indisputably demonstrated. In addition, this model is an indis-

45

pensable tool for the understanding of pathogenic mechanisms and the study of future vaccine candidates.

Example 5

Demonstration of the Vaccine Composition Protective Efficacy Produced from Nucleic Fragments of PWD Circovirus Sequence

1) Animals Used for the Study

Piglets having the PWD disease, reproduced under experimental conditions described in paragraph c) of Example 4, were used in a protocol for evaluating the vaccine composition efficacy, comprising nucleic fragments of PWD circovirus sequence.

2) Tested Vaccine Composition and Vaccination Protocol

a) Components Used for the Study

The plasmids were obtained from the pcDNA3 plasmid of INVITROGENE

pcDNA3ORF-plasmids

These plasmids are plasmids which do not carry a PWD circovirus nucleic acid insert and are used as a negative control plasmid.

65

pcDNA3ORF1+plasmid and pcDNA3ORF2+plasmid

The pcDNA3ORF1+ and pcDNA3ORF2+ plasmids are plasmids which carry a nucleic acid insert of the sequence of the PWD circovirus of TYPE B, and an insert comprising the nucleic acid fragment SEQ ID No. 23 (ORF'1) coding for the Rep protein of sequence SEQ ID No. 24 and an insert comprising the nucleic acid fragment SEQ ID No. 25 (ORF'2) coding for the protein of sequence SEQ ID No. 26, probably corresponding to the capsid protein, respectfully. These nucleic constructs further comprise the ATG initiation codon of the coding sequence of the corresponding protein.

GMCSF+Plasmid

GM-CSF (granulocyte/macrophage colony stimulating factor) is a cytokine which occurs in the development, the maturation and the activation of macrophages, granulocytes and dendritic cells which present an antigen. The beneficial contribution of the GM-CSF in vaccination is considered to be a cellular activation with, especially, the recruitment and the differentiation of cells which present an antigen.

This pcDNA3-GMCSF+ plasmid carries a nucleic acid insert coding for the granulocyte/macrophage colony stimulation factor, the GM-CSF protein.

The gene coding for this GM-CSF protein was cloned and sequenced by Inumaru et al. (*Immunol. Cell Biol.*, 1995, 73 (5), 474-476). The pcDNA3-GMCSF+ plasmid was obtained by Dr. B. Charley of INRA of Jouy-en-Josas (78, France).

Recombinant Baculoviruses

The so-called ORF-baculoviruses are viruses not carrying any insert comprising a nucleic acid fragment capable of expressing a PWD circovirus protein.

The so-called ORF1+(BAC ORF1+) or ORF2+(BAC ORF2+) baculoviruses are recombinant baculoviruses carrying an insert comprising a nucleic acid fragment SEQ ID No. 23 (ORF'1) and an insert comprising the nucleic acid fragment SEQ ID No. 25 (ORF'2), respectively.

Adjuvant

The adjuvant supplied by the Seppic Company, a subsidiary of AIR LIQUIDE, is the adjuvant corresponding to the reference AIF SEPPIC.

b) Vaccination Protocol

Weaned piglets aged 3 weeks are divided into four batches A, B, C and D each comprising 8 piglets.

Batches A, B and C, aged 3 weeks, each receive a first injection (injection M1) of 1 ml containing 200 micrograms of plasmids (naked DNA) in PBS, pH: 7.2, by the intramuscular route for each of the plasmids mentioned below for each batch, then, at the age of 5 weeks, a second injection (injection M2) comprising these same plasmids. A third injection is carried out simultaneously on the other side of the neck. This third injection comprises 1 ml of a suspension containing 5×10^8 cells infected by recombinant baculoviruses and 1 ml of AIF SEPPIC adjuvant.

Batch A (F1) (Control Batch):

First Injection

pcDNA3ORF1-plasmid, pcDNA3ORF2-plasmid and GMCSF+plasmid.

Second and Third Injection (Simultaneous)

pcDNA3ORF1-plasmid, pcDNA3ORF2-plasmid and GMCSF+plasmid;

Cells transformed by baculoviruses not containing any nucleic acid insert coding for a PWD circovirus protein;

AIF SEPPIC Adjuvant.

Batch B (F2) (Control Batch):

First Injection

pcDNA3ORF1-plasmid, pcDNA3ORF2-plasmid and GMCSF+plasmid;

Second and Third Injection (Simultaneous)

pcDNA3ORF1-plasmid, pcDNA3ORF2-plasmid and GMCSF+plasmid;

Cells transformed by baculoviruses not containing any nucleic acid insert coding for a PWD circovirus protein;

AIF SEPPIC Adjuvant.

Batch C (F3):

First Injection

pcDNA3ORF1+plasmid, pcDNA3ORF2+plasmid and GMCSF+plasmid;

Second and Third Injection (Simultaneous)

pcDNA3ORF1+plasmid, pcDNA3ORF2+plasmid and GMCSF+plasmid;

Cells transformed by BAC ORF1+ and BAC ORF2+ recombinant baculoviruses capable of respectively expressing the Rep protein of sequence SEQ ID No. 24 and the protein of sequence SEQ ID No. 26 of the PWD circovirus of TYPE B.

Batch D (F4) (Control Batch): No Injection

The batches of piglets B, C and D are infected (tested) at the age of 6 weeks although batch A is not subjected to the test.

3) Observation of the Batches

counting of coughing/sneezing: 15 minutes/batch/day;

consistency of fecal matter: every day;

regular recordings: weekly taking of blood, weighing;

weighing of food refuse: 3 times per week;

calculation of the daily mean gain in weight (dmg);

The daily mean gains were calculated for each of the batches over a period of 28 days following testing (cf. FIG. 10), an intermediate calculation of the dmG was likewise carried out for each of the batches over the first and second periods of 14 days. The results obtained are reported below in Table 6.

TABLE 6

	Daily mean gains			
	F1	F2	F3	F4
d0-d14	411 g	450 g	511 g	461 g
d14-d28	623 g	362 g	601 g	443 g
d0-d28	554 g	406 g	556 g	452 g

Measurement of Hyperthermia

The measurement of hyperthermia, of greater than 41° C. (cf. FIG. 11) and greater than 40.2° C., was carried out for each of the batches over a total period of 28 days following testing. The results obtained, corresponding to the ratio expressed as a percentage between the number of temperature recordings of greater than 41° C. (or greater than 40.2° C.) and the total number of temperature recordings carried out on all of the pigs per one-week period are reported below in Tables 7 and 8, respectively, for the hyperthermia measurements of greater than 41° C. and greater than 40.2° C.

TABLE 7

	Hyperthermia > 41° C.			
	F1	F2	F3	F4
W1	4.1	0	0	0
W2	10.7	16.	0	8.9
W3	4.7	27.	0	45.
W4	0	0	0	7.5

TABLE 8

Hyperthermia > 40.2				
	F1	F2	F3	F4
W1	29.1	10.41	29.1	20.8
W2	28.5	39.2	10.7	37.5
W3	14.3	68.7	25.0	81.2
W4	3.3	17.5	20.0	55

4) Conclusion

The recordings carried out clearly show that the animals which received the three injections of a vaccine composition comprising nucleic acid fragments of PWD circovirus according to the invention and/or capable of expressing recombinant proteins of PWD circovirus, in particular of type B, did not exhibit hyperthermia (cf. FIG. 10). These animals additionally did not experience a decline in their growth, the dmgs being comparable to those of uninfected control animals (cf. FIG. 9). They did not exhibit any particular clinical sign.

These results demonstrate the efficacious protection of the piglets against infection with a PWD circovirus of the invention, the primary agent responsible for PWD or FPW, provided by a vaccine composition prepared from a nucleic acid fragment of the nucleic sequence of PWD circovirus according to the invention, in particular of type B, and/or from recombinant proteins encoded by these nucleic acid fragments.

These results in particular show that the proteins encoded by the ORF1 and ORF2 of PWD circovirus according to the invention are immunogenic proteins inducing an efficacious protective response for the prevention of infection by a PWD circovirus.

Example 6

Serological Diagnosis of PWD Circovirus by Immunodetermination using Recombinant Proteins or Synthetic Peptides of PWD Circovirus

A. Serological Diagnosis with Recombinant Proteins

The identification and the sequencing of porcine PWD circovirus allow recombinant proteins of PWD circovirus to be produced by the techniques of genetic recombination well known to the person skilled in the art. Using these techniques, recombinant proteins encoded, in particular, by the ORF'2 of the PWD circovirus, type B, were expressed by transformed Sf9 insect cells and then isolated.

These recombinant proteins encoded by the ORF'2 are extracted, after culture of the transformed Sf9 cells, by thermal cell lysis by means of 3 cycles of freezing/thawing to $-70^{\circ}\text{C.}/+37^{\circ}\text{C.}$ Healthy Sf9 cells or nontransformed control Sf9 cells are also lysed.

Two antigenic fractions originating from nontransformed control Sf9 cells and Sf9 cells expressing the ORF'2 are precipitated at 4°C. by a 60% plus or minus 5% saturated ammonium sulfate solution. Determination of total proteins is carried out with the aid of the Biorad kit. 500 ng of control Sf9 proteins and of semipurified Sf9 proteins expressing the

ORF'2, in solution in 0.05 M bicarbonate buffer pH 9.6, are passively adsorbed at the bottom of 3 different wells of a Nunc Maxisorp microplate by incubation for one night at $+4^{\circ}\text{C.}$

The reactivity of pig sera with respect to each of these antigenic fractions is evaluated by an indirect ELISA reaction of which the experimental protocol is detailed below:

Saturation step: 200 $\mu\text{l/well}$ of PBS1 \times /3% semi-skimmed milk, 1 h 30 incubation at 37°C.

Washing: 200 $\mu\text{l/well}$ of PBS1 \times /Tween 20: 0.05%, 3 rapid washes.

Serum incubation step: 100 $\mu\text{l/well}$ of serum diluted to $1/100$ in PBS1 \times /semi-skimmed milk, 1%/Tween 20: 0.05%, 1 h incubation at 37°C.

Washing: 200 $\mu\text{l/well}$ of PBS1 \times /Tween 20: 0.05%, 2 rapid washes followed by 2 washes of 5 min.

Conjugate incubation step: 50 $\mu\text{l/well}$ of rabbit anti-pig conjugate diluted to $1/1000$ in PBS1 \times /semi-skimmed milk, 1%/Tween 20: 0.05%, 1 h incubation at 37°C.

Washing: 200 $\mu\text{l/well}$ of PBS1 \times /Tween 20: 0.05%, 2 rapid washes followed by 2 washes of 5 min.

Visualization step: 100 $\mu\text{l/well}$ of OPD substrate/citrate buffer/ H_2O_2 , 15 min incubation at 37°C.

Termination: 50 $\mu\text{l/well}$ of 1 N H_2SO_4 .

Read optical density in a spectrophotometer at 490 nm.

Results

The results obtained are shown below in Table 9.

TABLE 9

Antigens	Reactivity of Pig Serum not inoculated with Circovirus	Reactivity of Pig Serum inoculated with Circovirus
Purified Sf9 control	0.076	0.088
Sf9 expressing purified ORF'2	0.071	1.035

The results are expressed in optical density measured in a spectrophotometer at 490 nm during analysis by ELISA of the reactivity of pig sera which are or are not inoculated with the type B PWD circovirus according to the protocol indicated above.

B. Serological Diagnosis by Synthetic Peptide

The epitopic mapping of the proteins encoded, for example, by the nucleic sequences ORF1 and ORF2 of the two types of PWD circovirus (types A and B) additionally allowed immunogenic circoviral epitopes to be identified on the proteins encoded by the nucleic sequences ORF'1 and ORF'2 as well as the specific epitopes of the protein encoded by the nucleic acid sequence ORF'2 of the type B PWD circovirus. Four specific epitopes of the type B PWD circovirus and one epitope common to the two types of PWD circovirus situated on the protein encoded by the nucleic sequence ORF'2 were synthesized in peptide form. The equivalent peptides in the circovirus of type A were likewise synthesized. All peptides were evaluated as diagnostic antigens within the context of carrying out a serological test.

Results

The results obtained are shown in Table 10, below.

TABLE 10

Results of the evaluation as a diagnostic antigen of synthetic peptides encoded by the nucleic sequences ORF2 and ORF'2 of PWD circovirus of type A and B.									
SEQ ID NO:	Peptide	Type	PWD circovirus	Position	AA sequence	Infected pig serum reactivity			Epitopic specificity
						SPF D0/D54	Conventional 1 D0/D42	Conventional 2 D0/D42	
SEQ ID NO: 29	121	B		71-85	VDMMRFNINDFLPPG	+/-, +++	+/-, +++	-, +++	Circovirus B
SEQ ID NO: 55	177	B		70-84	NVNELRFNIGQFLPP	+/-, +	+/-, +/-	+/-, -	
SEQ ID NO: 30	131	B		115-129	QGDRGVGSSAVILDD	+/-, +/-	++, ++	+/-, +	Circovirus B
SEQ ID NO: 56	188	A		114-127	TSNQRGVGSTVVIL	+/-, -	-, +/-	+/-, +/-	
SEQ ID NO: 31	133	B		119-134	GVGSSAVILDDNVFTK	-, ++	++, +++	+/-, ++	
SEQ ID NO: 57	189	A		118-132	RGVGVSTVVILDANFV	+/-, -	-, +/-	+/-, +/-	
SEQ ID NO: 58	146	B		171-185	FTIDYFQPNNKRNQL	-, +/-	-, ++	-, ++	Circovirus A&B
SEQ ID NO: 59	202	A		170-184	DQTIDWFQPNKRNQ	+++ , +++	+/-, ++	+, ++	
SEQ ID NO: 32	152	B		195-209	VDHVGLGTAFENSIY	-, ++	+++ , +++	+/-, +	Circovirus B
SEQ ID NO: 60	208	A		194-208	NVEHTGLGYALQMAT	- , -	- , -	- , -	

+/-, +, ++, +++. Increasing intensities of the reactivities observed in Spot peptides on a nitrocellulose membrane. The porcine sera tested are from animals experimentally infected with the circovirus of type B within the animal houses of the CNEVA. Samples are taken from the animals before inoculation on d0 and 42 days or 54 days after inoculation, on d42, d54.

Example 7

Characterization of the Specific Epitopes of the PWD Circovirus of Type B

25

The proteins encoded by the ORF2 of the porcine circoviruses of type A and B were chosen for this study. For each of the ORF2s (types A and B), 56 peptides of 15 amino acids which overlap every 4 amino acids were synthesized, thus covering the whole of the protein (cf. Table 11 below).

30

TABLE 11

Sequence of amino acids of the 56 peptides of 15 amino acids synthesized from the nucleic sequence ORF2' (type B) and ORF2 (type A) of PWD circovirus with their corresponding spot number (cf. FIG. 12)					
Type B ORF'2			Type A ORF2		
Spot No.	Sequence		Spot No.	Sequence	
SEQ ID NO: 61	107	HRPRSHLGQILRRRP	SEQ ID NO: 84	163	TRPRSHLGNILRRRP
SEQ ID NO: 62	108	SHLGQILRRRPWLHVH	SEQ ID NO: 85	164	SHLGNILRRRPYLHVH
SEQ ID NO: 63	109	QILRRRPWLHVHPRHR	SEQ ID NO: 86	165	NILRRRPYLHVHPAPR
SEQ ID NO: 64	110	RRPWLHVHPRHRWR	SEQ ID NO: 87	166	RRPYLVHHPAFNRNRYR
SEQ ID NO: 65	111	LVHPRHRWRRRKNG	SEQ ID NO: 88	167	LVHHPAFNRNRYRWRK
SEQ ID NO: 66	112	RHRWRRRKNGIFNT	SEQ ID NO: 89	168	AFNRNRYRWRKNGIF
SEQ ID NO: 67	113	RWRKNGIFNTRLSR	SEQ ID NO: 90	169	RYRWRKNGIFNTRLSR
SEQ ID NO: 68	114	KNGIFNTRLSRTFGY	SEQ ID NO: 91	170	RRKNGIFNTRLSRREF
SEQ ID NO: 69	115	FNTRLSRTFGYTVKR	SEQ ID NO: 92	171	GIFNTRLSRREFVLT
SEQ ID NO: 70	116	LSRTFGYTVKTFVVR	SEQ ID NO: 93	172	SRLSREFVLTIRGGH
SEQ ID NO: 71	117	FGYTVKRTTVRTPSW	SEQ ID NO: 94	173	REFVLTIRGGHSQPS
SEQ ID NO: 72	118	VKRTTVRTPSWAVDM	SEQ ID NO: 95	174	LTIRGGHSOPSWNVN
SEQ ID NO: 73	119	TVRTPSWAVDMMRFN	SEQ ID NO: 96	175	GGHSQPSWNVNELRF
SEQ ID NO: 74	120	PSWAVDMMRFNINDF	SEQ ID NO: 97	176	QPSWNVNELRFNIGO
SEQ ID NO: 29	121	VDMMRFNINDFLPPG	SEQ ID NO: 98	177	NVNELRFNIGQFLPP

TABLE 11-continued

Sequence of amino acids of the 56 peptides of 15 amino acids synthesized from the nucleic sequence ORF2' (type B) and ORF2 (type A) of PWD circovirus with their corresponding spot number (cf. FIG. 12)					
Type B ORF'2			Type A ORF2		
SEQ ID NO:	Spot No.	Sequence	SEQ ID NO:	Spot No.	Sequence
75	122	RFNINDFLPPGGGSN	99	178	LRFNIGQFLPPSGGT
76	123	NDFLPPGGGSNPRSV	100	179	IGQFLPPSGGTNPLP
77	124	PPGGGSNPRSVPFY	101	180	LPPSGGTNPLPLPFQ
78	125	GSNPRSVPFYRIR	102	181	GGTNPLPLPFQYRI
79	126	RSVPFYRIRKVKV	103	182	PLPLPFQYRIRKAK
80	127	FYRIRKVKVEFWP	104	183	PFQYRIRKAKYEFY
81	128	RIRKVKVEFWPCSPI	105	184	YRIRKAKYEFYPRDP
82	129	VKVEFWPCSPITQGD	106	185	KAKYEFYPRDPITSN
83	130	FWPCSPITQDRGVG	107	186	EFYPRDPITSNQRGV
30	131	SPITQDRGVGSSAV	108	187	RDPITSNQRGVGSTV
31	132	QDRGVGSSAVILDD	109	188	TSNQRGVGSTVVILD
110	133	GVGSSAVILDDNFVT	136	189	RGVGSTVVILDANFV
111	134	SAVILDDNFVTKATA	137	190	STVVILDANFVTPST
112	135	LDDNFVTKATALTYD	138	191	ILDANFVTPSTNLAY
113	136	FVTKATALTYDPYVN	139	192	NFVTPSTNLAYDPYI
114	137	ATALTYDPYVNYSSR	140	193	PSTNLAYDPYINYSS
115	138	TYDPYVNYSSRIITIT	141	194	LAYDPYINYSSRHTI
116	139	VVNYSSRHTITQPFS	142	195	PYINYSSRHTIRQPF
117	140	SSRHTITQPFSYHSR	143	196	YSSRIITIRQPFTYHS
118	141	TITQPFSYHSRYFTP	144	197	HTIRQPFTYHSRYFT
119	142	PFSYHSRYFTP KPVL	145	198	QPFTYHSRYFTP KPPE
120	143	HSRYFTP KPVLDFTI	146	199	YHSRYFTP KPPELDQT
121	144	FTP KPVLDFTI DYFQ	147	200	YFTP KPPELDQTI DWF
122	145	PVLDFTI DYFQ PNNK	148	201	KPELDQTI DWFQ PNN
123	146	FTI DYFQ PNNKRNQL	149	202	DQTI DWFQ PNNKRNQ
124	147	YFQ PNNKRNQLWLR	150	203	DWFQ PNNKRNQLWLH
125	148	NNKRNQLWLR LQTAG	151	204	PNNKRNQLWLRHLNTH
126	149	NQLWLR LQTAGNVDH	152	205	RNQLWLRHLNTH TNVE
127	150	LRLQTAGNVDHVGLG	153	206	WLHLNTH TNVEHTGL
128	151	TAGNVDHVGLGTAFE	154	207	NTH TNVEHTGLGYAL
32	152	VDHVGLGTAFENSIY	155	208	NVEHTGLGYALQ NAT
129	153	GLGTAFENSIYDQEY	156	209	TGLGYALQ NATTAQN
130	154	AFENSIYDQEYNIRV	157	210	YALQ NATTAQ NYVVR
131	155	SIYDQEYNIRV TMYV	158	211	NATTAQ NYVVR LTIY
132	156	QEYNIRV TMYVQFRE	159	212	AQ NYVVR LTIYVQFR

TABLE 11-continued

Sequence of amino acids of the 56 peptides of 15 amino acids synthesized from the nucleic sequence ORF2' (type B) and ORF2 (type A) of PWD circovirus with their corresponding spot number (cf. FIG. 12)					
Type B ORF'2			Type A ORF2		
Spot No.	Sequence		Spot No.	Sequence	
SEQ ID NO:133	157	IRVTMYVQFREFNFK	SEQ ID NO:160	213	VVRLTIYVQFREFIL
SEQ ID NO:134	158	MYVQFREFNFKDPPL	SEQ ID NO:161	214	TIYVQFREFILKDPL
SEQ ID NO:135	159	VQFREFNFKDPLNP	SEQ ID NO:162	215	YVQFREFILKDPLNE

15

These peptides were synthesized according to the "spot" method which consists of simultaneous synthesis of a large number of peptides on a cellulose solid support, each site of synthesis of a peptide constituting a spot (Synt:em, NIMES). This method involves orientation of the peptides on the plate, these being fixed covalently by the carboxy-terminal end. A spot represents approximately 50 nmol of peptide.

The reference of the spots and corresponding peptide sequences is given in Table 11.

These membranes were used for immunoreactivity tests with respect to serum of SPF pigs which were or were not infected experimentally with the type B PWD circovirus strain as well as with respect to sera of infected pigs from conventional farms (conventional farms 1 or 2). This study allowed specific immunoreactive peptides of the circovirus of type B corresponding to the spots No. 121, No. 132, No. 133 and No. 152 (respectively of amino acid sequences SEQ ID No. 29, SEQ ID No. 30, SEQ ID No. 31 and SEQ ID No. 32) to be demonstrated. An illustration is shown in FIG. 12 where the membranes are visualized with an infected pig serum coming from a conventional farm. Nonspecific immunoreactive peptides of type [Iacuna] were likewise demonstrated, among which we shall keep the peptide No. 146 SEQ ID No. 123 which is strongly immunogenic.

A comparison between the peptide sequences of circoviruses of type A and B (FIG. 13) indicates a divergence ranging from 20 to 60% for the specific immunoreactive peptides of the type B, and a weaker divergence (13%) between the nonspecific peptides.

Example 8

Protection of Swine From Post-Weaning Multisystemic Wasting Syndrome (PMWS) Conferred by Procine Circovirus Type B (PCV-B) ORF'2 Protein

The ORF'1-encoded protein (REP) and ORF'2-encoded putative capsid protein of PCV-B were expressed, either in insect cells by recombinant baculovirus vectors, or in mammalian cell lines by transfection with plasmidic expression vectors. These two circovirus-derived proteins were detectable in both expression systems. As evaluated by weight gains, hyperthermia and absence of lesions following challenge, the pigs were protected against a virulent circovirus challenge after one first DNA immunization with plasmids directing ORF'2 protein and GM-CSF expression and a second injection, 15 days later, with the same plasmid preparation plus the ORF'2 recombinant protein. A lower level of protection was observed when the pigs were vaccinated with ORF'1 protein, as opposed to pigs vaccinated with ORF'2 protein.

A. Development of an Experimental Model of PMWS in Swine:

Eight 3 week-old SPF pigs were inoculated intratracheally (5 ml) and intramuscularly (1 ml).

B. Production and Control of PCV-B Plasmids:

PCV-B ORF'1 and ORF'2 genes, isolated from PCV-B challenge strain, was cloned into vector plasmid pcDNA3.1. All constructs were validated through a partial sequencing of the PCV-B genes in the final plasmids and expression control by immunoperoxidase on PK15 cells respectively transfected with each plasmid, using swine polyclonal antibodies.

Plasmid Encoding GM-CSF has Been Co-Administered.

C. Construction of Recombinant Baculoviruses:

ORF'1 and ORF'2 proteins were expressed under polyhedrin promoter control. Recombinant proteins were detected by western-blot using swine polyclonal antibodies.

D. Vaccination and Challenge:

Four groups of 7 pigs were vaccinated intramuscularly at day 0 (Do), two weeks later, they received the same plasmid preparation plus the recombinant baculovirus.

E. Monitoring:

All groups of pigs were housed in isolated experimental units with air filtration and low air pressure. Clinical observations and rectal temperatures were recorded every day. The pigs were weighed weekly.

F. Conclusions

Expression of PCV-B ORF'2 or PCV-B ORF'1 in swine resulted in a significantly enhanced level of protection as evaluated by weight evolution and body temperature evolution following challenge with PCV-B circovirus. These results are summarized in FIGS. 14 and 15.

The invention described herein may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The specific embodiments previously described are therefore to be considered as illustrative of, and not limiting, the scope of the invention. Additionally, the disclosure of all publications and patent applications cited above and below, including International Patent Application No. PCT/FR98/02634, filed Dec. 4, 1998, and published as International Publication No. WO 99/29871 on Jun. 17, 1999, are expressly incorporated herein by reference in their entireties to the same extent as if each were incorporated by reference individually.

BIBLIOGRAPHIC REFERENCES

- Allan, G. M. et al., 1995, *Vet. Microbiol.*, 44: 49-64.
 65 Barany, F., 1911, *PNAS. USA*, 88: 189-193.
 Boulton, L. H. et al., 1997, *J. Gen. Virol.*, 78 (Pt 6), 1265-1270.

- Buckholz, R. G., 1993, Yeast systems for the expression of heterologous gene products. *Curr. Op. Biotechnology* 4: 538-542.
- Burg, J. L. et al., 1996, *Mol. and Cell. Probes*, 10: 257-271.
- Chu, B. C. F. et al., 1986, *NAR*, 14: 5591-5603.
- Chu, P. W. G. et al., 1993, *Virus Research*, 27: 161-171.
- Clark, E. G., 1997, *American Association of Swine Practitioners*, 499-501.
- Daft, B. et al., 1996, *American Association of Veterinary Laboratory Diagnosticians*, 32.
- Derse, D. et al., 1995, *J. Virol.*, 69(3): 1907-1912.
- Duck, P. et al., 1990, *Biotechniques*, 9: 142-147.
- Dulac, G. C. et al., 1989, *Can. J. Vet. Res.*, 53: 431-433.
- Edwards, C. P., and Aruffo, A., 1993, Current applications of COS cell based transient expression systems. *Curr. Op. Biotechnology* 4: 558-563.
- Edwards, S. et al., 1994, *Vet. Rec.*, 134: 680-681.
- Erlich, H. A., 1989, In *PCR Technology. Principles and Applications for DNA Amplification*. New York: Stockton Press.
- Felgner, et al., 1987, *Proc. Natl. Acad. Sci.*, 84: 7413.
- Fontes, E. P. B. et al., 1994, *J. Biol. Chem.*, Vol. 269, No. 11: 8459-8465.
- Fraley et al., 1980, *J. Biol. Chem.*, 255: 10431.
- Guateli, J. C. et al., 1990, *PNAS. USA*, 87: 1874-1878.
- Hackland, A. F. et al., 1994, *Arch. Virol.*, 139: 1-22.
- Hanson, S. F. et al., 1995, *Virology*, 211: 1-9.
- Harding, J. C., 1997, *American Association of Swine Practitioners*, 503.
- Harding, R. M. et al., 1993, *Journal of General Virology*, 74: 323-328.
- Harding, J. C. and Clark, E. G., 1997, *Swine Health and Production*, Vol. 5, No. 5: 201-203.
- Heyraud-Nitschke, F. et al., 1995, *Nucleic Acids Research*, Vol. 23, No. 6.
- Homer, G. W., 1991, *Surveillance* 18(5): 23.
- Houben-Weyl, 1974, in *Methode der Organischen Chemie*, E. Wunsch Ed., Volume 15-I and 15-II, Thieme, Stuttgart.
- Huygen, K. et al., 1996, *Nature Medicine*, 2(8): 893-898.
- Innis, M. A. et al., 1990, in *PCR Protocols. A guide to Methods and Applications*, San Diego, Academic Press.
- Kaneda, et al., 1989, *Science*, 243: 375.
- Kievitis, T. et al., 1991, *J. Virol. Methods*, 35: 273-286.
- Kohler, G. et al., 1975, *Nature*, 256(5517): 495-497.
- Kwoh, D. Y. et al., 1989, *PNAS. USA*, 86: 1173-1177.
- Ladany, S. et al., 1989, *J. Clin. Microbiol.* 27: 2778-2783.
- Lazarowitz, S. G. et al., 1989, *The EMBO Journal*, Vol. 8 No. 4: 1023-1032.

- Luckow, V. A., 1993, Baculovirus systems for the expression of human gene products. *Curr. Op. Biotechnology* 4: 564-572.
- Mankertz, A. et al., 1997, *J. Virol.*, 71: 2562-2566.
- 5 Matthews, J. A. et al., 1988, *Anal. Biochem.*, 169: 1-25.
- McNeilly, F. et al., 1996, *Vet. Immunol. Immunopathol.*, 49: 295-306.
- Meehan, B. M. et al., 1997, *J. Gen. Virol.* 78: 221-227.
- 10 Merrifield, R. D., 1966, *J. Am. Chem. Soc.*, 88(21): 5051-5052.
- Midoux, 1993, *Nucleic Acids Research*, 21: 871-878.
- Miele, E. A. et al., 1983, *J. Mol. Biol.*, 171: 281-295.
- Murphy, F. A. et al., 1995, *Sixth Report of the International Committee on Taxonomy of Viruses*. Springer-Verlag Wien N.Y.
- 15 Nayar, G. P. et al., 1997, *Can. Vet. J.* 38(6): 385-386.
- Olins, P. O., and Lee, S. C., 1993, Recent advances in heterologous gene expression in *E. coli*. *Curr. Op. Biotechnology* 4: 520-525.
- 20 Pagano et al., 1967, *J. Virol.*, 1: 891.
- Rolfs, A. et al., 1991, In *PCR Topics. Usage of Polymerase Chain reaction in Genetic and Infectious Disease*. Berlin: Springer-Verlag.
- Sambrook, J. et al., 1989, In *Molecular cloning: A Laboratory Manual*. Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press.
- 25 Sanchez-Pescador, R., 1988, *J. Clin. Microbiol.*, 26(10): 1934-1938.
- Segev D., 1992, in "Non-radioactive Labeling and Detection of Biomolecules". Kessler C. Springer Verlag, Berlin, N.Y.: 197-205.
- Shiver, J. W., 1995, in *Vaccines 1995*, eds Chanock, R. M. Brown, F. Ginsberg, H. S. & Norrby, E., pp. 95-98, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- 35 Tascon, R. E. et al., 1996, *Nature Medicine*, 2(8): 888-892.
- Tischer, I. et al., 1982, *Nature*, 295: 64-66.
- Tischer, I. et al., 1986, *Arch. Virol.*, 91: 271-276.
- Tischer, I. et al., 1988, *Zentralbl Bakteriell Mikrobiol Hyg [A]* 270: 280-287.
- Tischer, I. et al., 1995, *Arch. Virol.*, 140: 737-743.
- Urdea, M. S., 1988, *Nucleic Acids Research*, II: 4937-4957.
- Walker, G. T. et al., 1992, *NAR* 20: 1691-1696.
- Walker, G. T. et al., 1992, *PNAS. USA*, 89: 392-396.
- 45 White, B. A. et al., 1997, *Methods in Molecular Biology*, 67, Humana Press, Towota.
- Zhao, T. M. et al., 1996, *Proc. Natl. Acad. Sci., USA* 93(13): 6653-6648.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 170

<210> SEQ ID NO 1
 <211> LENGTH: 1759
 <212> TYPE: DNA
 <213> ORGANISM: Type A PWD circovirus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(78)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (82)..(99)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (106)..(156)

-continued

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (160) .. (195)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (199) .. (231)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (235) .. (246)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (250) .. (315)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (319) .. (330)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (334) .. (489)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (493) .. (525)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (529) .. (591)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (595) .. (600)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (604) .. (606)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (610) .. (627)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (634) .. (636)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (640) .. (681)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (685) .. (708)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (712) .. (726)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (730) .. (753)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (757) .. (933)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (937) .. (969)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (973) .. (1047)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1051) .. (1056)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1060) .. (1071)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1075) .. (1236)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1240) .. (1257)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1261) .. (1293)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1297) .. (1350)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1354) .. (1380)

-continued

```

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1384)..(1386)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1390)..(1416)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1420)..(1425)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1429)..(1497)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1501)..(1512)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1516)..(1551)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1555)..(1566)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1570)..(1581)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1585)..(1620)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1624)..(1752)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1756)..(1758)

<400> SEQUENCE: 1

acc agc gca ctt cgg cag cgg cag cac ctc ggc agc gtc agt gaa aat      48
Thr Ser Ala Leu Arg Gln Arg Gln His Leu Gly Ser Val Ser Glu Asn
1          5          10         15

gcc aag caa gaa aag cgg ccc gca acc cca taa gag gtg ggt gtt cac      96
Ala Lys Gln Glu Lys Arg Pro Ala Thr Pro Glu Val Gly Val His
          20          25          30

cct taataa tcc ttc cga gga gga gaa aaa caa aat acg gga gct tcc      144
Pro          Ser Phe Arg Gly Gly Glu Lys Gln Asn Thr Gly Ala Ser
          35          40          45

aat ctc cct ttt tga tta ttt tgt ttg tgg cga gga agg ttt gga aga      192
Asn Leu Pro Phe          Leu Phe Cys Leu Trp Arg Gly Arg Phe Gly Arg
          50          55          60

ggg tag aac tcc tca cct cca ggg gtt tgc gaa ttt tgc taa gaa gca      240
Gly          Asn Ser Ser          Pro Pro Gly Val Cys Glu Phe Cys          Glu Ala
          65          70

gac ttt taa caa ggt gaa gtg gta ttt tgg tgc ccg ctg cca cat cga      288
Asp Phe          Gln Gly Glu Val Val Phe Trp Cys Pro Leu Pro His Arg
75          80          85

gaa agc gaa agg aac cga cca gca gaa taa aga ata ctg cag taa aga      336
Glu Ser Glu Arg Asn Arg Pro Ala Glu          Arg Ile Leu Gln          Arg
90          95          100

agg cca cat act tat cga gtg tgg agc tcc gcg gaa cca ggg gaa gcg      384
Arg Pro His Thr Tyr Arg Val Trp Ser Ser Ala Glu Gly Glu Ala
          105          110          115

cag cga cct gtc tac tgc tgt gag tac cct ttt gga gac ggg gtc ttt      432
Gln Arg Pro Val Tyr Cys Cys Glu Tyr Pro Phe Gly Asp Gly Val Phe
120          125          130          135

ggt gac tgt agc cga gca gtt tcc tgt aac gta tgt gag aaa ttt ccg      480
Gly Asp Cys Ser Arg Ala Val Ser Cys Asn Val Cys Glu Lys Phe Pro
          140          145          150

cgg gct ggc tga act ttt gaa agt gag cgg gaa gat gca gaa gcg tga      528
Arg Ala Gly          Thr Phe Glu Ser Glu Arg Glu Asp Ala Glu Ala
          155          160          165

```

-continued

ttg gaa gac agc tgt aca cgt cat agt ggg ccc gcc cgg ttg tgg gaa	576
Leu Glu Asp Ser Cys Thr Arg His Ser Gly Pro Ala Arg Leu Trp Glu	
170 175 180	
gag cca gtg ggc ccg taa ttt tgc tga gcc tag gga cac cta ctg gaa	624
Glu Pro Val Gly Pro Phe Cys Ala Gly His Leu Leu Glu	
185 190	
gcc tagtag aaa taa gtg gtg gga tgg ata tca tgg aga aga agt tgt	672
Ala Lys Val Val Gly Trp Ile Ser Trp Arg Arg Ser Cys	
195 200 205	
tgt ttt gga tga ttt tta tgg ctg gtt acc ttg gga tga tct act gag	720
Cys Phe Gly Phe Leu Trp Leu Val Thr Leu Gly Ser Thr Glu	
210 215 220	
act gtg tga ccg gta tcc att gac tgt aga gac taa agg ggg tac tgt	768
Thr Val Pro Val Ser Ile Asp Cys Arg Asp Arg Gly Tyr Cys	
225 230 235	
tcc ttt ttt ggc ccg cag tat ttt gat tac cag caa tca ggc ccc cca	816
Ser Phe Phe Gly Pro Gln Tyr Phe Asp Tyr Gln Gln Ser Gly Pro Pro	
240 245 250	
gga atg gta ctc ctc aac tgc tgt ccc agc tgt aga agc tct cta tcg	864
Gly Met Val Leu Leu Asn Cys Cys Pro Ser Cys Arg Ser Ser Leu Ser	
255 260 265	
gag gat tac tac ttt gca att ttg gaa gac tgc tgg aga aca atc cac	912
Glu Asp Tyr Tyr Phe Ala Ile Leu Glu Asp Cys Trp Arg Thr Ile His	
270 275 280	
gga ggt acc cga agg ccg att tga agc agt gga ccc acc ctg tgc cct	960
Gly Gly Thr Arg Arg Pro Ile Ser Ser Gly Pro Thr Leu Cys Pro	
285 290 295	
ttt ccc ata taa aat aaa tta ctg agt ctt ttt tgt tat cac atc gta	1008
Phe Pro Ile Asn Lys Leu Leu Ser Leu Phe Cys Tyr His Ile Val	
300 305 310	
atg gtt ttt att ttt att cat tta gag ggt ctt tca gga taa att ctc	1056
Met Val Phe Ile Phe Ile His Leu Glu Gly Leu Ser Gly Ile Leu	
315 320 325	
tga att gta cat aaa tag tca acc tta cca cat aat ttt ggg ctg tgg	1104
Ile Val His Lys Ser Thr Leu Pro His Asn Phe Gly Leu Trp	
330 335 340	
ttg cat ttt gga gcg cat agc cca ggc ctg tgt gct cga cat tgg tgt	1152
Leu His Phe Gly Ala His Ser Pro Gly Leu Cys Ala Arg His Trp Cys	
345 350 355	
ggg tat tta aat gga gcc aca gct ggt ttc ttt tat tat ttg gct gga	1200
Gly Tyr Leu Asn Gly Ala Thr Ala Gly Phe Phe Tyr Tyr Leu Ala Gly	
360 365 370	
acc aat caa ttg ttt ggt cta gct ctg gtt tgg ggg tga agt acc tgg	1248
Thr Asn Gln Leu Phe Gly Leu Ala Leu Val Trp Gly Ser Thr Trp	
375 380 385	
agt ggt agg taa agg gct gcc tta tgg tgt ggc ggg agg agt agt taa	1296
Ser Gly Arg Arg Ala Ala Leu Trp Cys Gly Gly Arg Ser Ser	
390 395 400	
tat agg ggt cat agg cca agt tgg tgg agg ggg tta caa agt tgg cat	1344
Tyr Arg Gly His Arg Pro Ser Trp Trp Arg Gly Leu Gln Ser Trp His	
405 410 415	
cca aga taa caa cag tgg acc caa cac ctc ttt gat tag agg tga tgg	1392
Pro Arg Gln Gln Trp Thr Gln His Leu Phe Asp Arg Trp	
420 425 430	
ggt ctc tgg ggt aaa att cat att tag cct ttc taa tac ggt agt att	1440
Gly Leu Trp Gly Lys Ile His Ile Pro Phe Tyr Gly Ser Ile	
435 440 445	
gga aag gta ggg gta ggg ggt tgg tgc cgc ctg agg ggg gga gga act	1488
Gly Lys Val Gly Val Gly Gly Trp Cys Arg Leu Arg Gly Gly Gly Thr	
450 455 460	

-continued

```

ggc cga tgt tga atc tca gct cgt taa cat tcc aag atg gct gcg agt 1536
Gly Arg Cys Ile Ser Ala Arg His Ser Lys Met Ala Ala Ser
465 470 475

gtc ctc ctc tta tgg tga gta caa att ctc tag aaa ggc ggg aat tga 1584
Val Leu Leu Leu Trp Val Gln Ile Leu Lys Gly Gly Asn
480 485

aga tac ccg tct ttc ggc gcc atc tgt aac ggt ttc tga agg cgg ggt 1632
Arg Tyr Pro Ser Phe Gly Ala Ile Cys Asn Gly Phe Arg Arg Gly
490 495 500

gta cca aat atg gtc ttc tcc gga gga tgt ttc caa gat ggc tgc ggg 1680
Val Pro Asn Met Val Phe Ser Gly Gly Cys Phe Gln Asp Gly Cys Gly
505 510 515 520

ggc ggg tcc gtc ttc tgc ggt aac gcc tcc ttg gcc acg tca tcc tat 1728
Gly Gly Ser Val Phe Cys Gly Asn Ala Ser Leu Ala Thr Ser Ser Tyr
525 530 535

aaa agt gaa aga agt gcg ctg ctg tag tat t 1759
Lys Ser Glu Arg Ser Ala Leu Leu Tyr
540 545

```

<210> SEQ ID NO 2

<211> LENGTH: 545

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 2

```

Thr Ser Ala Leu Arg Gln Arg Gln His Leu Gly Ser Val Ser Glu Asn
1 5 10 15

Ala Lys Gln Glu Lys Arg Pro Ala Thr Pro Glu Val Gly Val His Pro
20 25 30

Ser Phe Arg Gly Gly Glu Lys Gln Asn Thr Gly Ala Ser Asn Leu Pro
35 40 45

Phe Leu Phe Cys Leu Trp Arg Gly Arg Phe Gly Arg Gly Asn Ser Ser
50 55 60

Pro Pro Gly Val Cys Glu Phe Cys Glu Ala Asp Phe Gln Gly Glu Val
65 70 75 80

Val Phe Trp Cys Pro Leu Pro His Arg Glu Ser Glu Arg Asn Arg Pro
85 90 95

Ala Glu Arg Ile Leu Gln Arg Arg Pro His Thr Tyr Arg Val Trp Ser
100 105 110

Ser Ala Glu Pro Gly Glu Ala Gln Arg Pro Val Tyr Cys Cys Glu Tyr
115 120 125

Pro Phe Gly Asp Gly Val Phe Gly Asp Cys Ser Arg Ala Val Ser Cys
130 135 140

Asn Val Cys Glu Lys Phe Pro Arg Ala Gly Thr Phe Glu Ser Glu Arg
145 150 155 160

Glu Asp Ala Glu Ala Leu Glu Asp Ser Cys Thr Arg His Ser Gly Pro
165 170 175

Ala Arg Leu Trp Glu Glu Pro Val Gly Pro Phe Cys Ala Gly His Leu
180 185 190

Leu Glu Ala Lys Val Val Gly Trp Ile Ser Trp Arg Arg Ser Cys Cys
195 200 205

Phe Gly Phe Leu Trp Leu Val Thr Leu Gly Ser Thr Glu Thr Val Pro
210 215 220

Val Ser Ile Asp Cys Arg Asp Arg Gly Tyr Cys Ser Phe Phe Gly Pro
225 230 235 240

Gln Tyr Phe Asp Tyr Gln Gln Ser Gly Pro Pro Gly Met Val Leu Leu
245 250 255

```

-continued

Asn Cys Cys Pro Ser Cys Arg Ser Ser Leu Ser Glu Asp Tyr Tyr Phe
 260 265 270
 Ala Ile Leu Glu Asp Cys Trp Arg Thr Ile His Gly Gly Thr Arg Arg
 275 280 285
 Pro Ile Ser Ser Gly Pro Thr Leu Cys Pro Phe Pro Ile Asn Lys Leu
 290 295 300
 Leu Ser Leu Phe Cys Tyr His Ile Val Met Val Phe Ile Phe Ile His
 305 310 315 320
 Leu Glu Gly Leu Ser Gly Ile Leu Ile Val His Lys Ser Thr Leu Pro
 325 330 335
 His Asn Phe Gly Leu Trp Leu His Phe Gly Ala His Ser Pro Gly Leu
 340 345 350
 Cys Ala Arg His Trp Cys Gly Tyr Leu Asn Gly Ala Thr Ala Gly Phe
 355 360 365
 Phe Tyr Tyr Leu Ala Gly Thr Asn Gln Leu Phe Gly Leu Ala Leu Val
 370 375 380
 Trp Gly Ser Thr Trp Ser Gly Arg Arg Ala Ala Leu Trp Cys Gly Gly
 385 390 395 400
 Arg Ser Ser Tyr Arg Gly His Arg Pro Ser Trp Trp Arg Gly Leu Gln
 405 410 415
 Ser Trp His Pro Arg Gln Gln Trp Thr Gln His Leu Phe Asp Arg Trp
 420 425 430
 Gly Leu Trp Gly Lys Ile His Ile Pro Phe Tyr Gly Ser Ile Gly Lys
 435 440 445
 Val Gly Val Gly Gly Trp Cys Arg Leu Arg Gly Gly Thr Gly Arg
 450 455 460
 Cys Ile Ser Ala Arg His Ser Lys Met Ala Ala Ser Val Leu Leu Leu
 465 470 475 480
 Trp Val Gln Ile Leu Lys Gly Gly Asn Arg Tyr Pro Ser Phe Gly Ala
 485 490 495
 Ile Cys Asn Gly Phe Arg Arg Gly Val Pro Asn Met Val Phe Ser Gly
 500 505 510
 Gly Cys Phe Gln Asp Gly Cys Gly Gly Ser Val Phe Cys Gly Asn
 515 520 525
 Ala Ser Leu Ala Thr Ser Ser Tyr Lys Ser Glu Arg Ser Ala Leu Leu
 530 535 540
 Tyr
 545

<210> SEQ ID NO 3
 <211> LENGTH: 577
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus
 <400> SEQUENCE: 3

Pro Ala His Phe Gly Ser Gly Ser Thr Ser Ala Ala Ser Val Lys Met
 1 5 10 15
 Pro Ser Lys Lys Ser Gly Pro Gln Pro His Lys Arg Trp Val Phe Thr
 20 25 30
 Leu Asn Asn Pro Ser Glu Glu Glu Lys Asn Lys Ile Arg Glu Leu Pro
 35 40 45
 Ile Ser Leu Phe Asp Tyr Phe Val Cys Gly Glu Glu Gly Leu Glu Glu
 50 55 60
 Gly Arg Thr Pro His Leu Gln Gly Phe Ala Asn Phe Ala Lys Lys Gln

-continued

65	70	75	80
Thr Phe Asn Lys Val Lys Trp Tyr Phe Gly Ala Arg Cys His Ile Glu 85 90 95			
Lys Ala Lys Gly Thr Asp Gln Gln Asn Lys Glu Tyr Cys Ser Lys Glu 100 105 110			
Gly His Ile Leu Ile Glu Cys Gly Ala Pro Arg Asn Gln Gly Lys Arg 115 120 125			
Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu Thr Gly Ser Leu 130 135 140			
Val Thr Val Ala Glu Gln Phe Pro Val Thr Tyr Val Arg Asn Phe Arg 145 150 155 160			
Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met Gln Lys Arg Asp 165 170 175			
Trp Lys Thr Ala Val His Val Ile Val Gly Pro Pro Gly Cys Gly Lys 180 185 190			
Ser Gln Trp Ala Arg Asn Phe Ala Glu Pro Arg Asp Thr Tyr Trp Lys 195 200 205			
Pro Ser Arg Asn Lys Trp Trp Asp Gly Tyr His Gly Glu Glu Val Val 210 215 220			
Val Leu Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp Asp Leu Leu Arg 225 230 235 240			
Leu Cys Asp Arg Tyr Pro Leu Thr Val Glu Thr Lys Gly Gly Thr Val 245 250 255			
Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn Gln Ala Pro Gln 260 265 270			
Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu Ala Leu Tyr Arg 275 280 285			
Arg Ile Thr Thr Leu Gln Phe Trp Lys Thr Ala Gly Glu Gln Ser Thr 290 295 300			
Glu Val Pro Glu Gly Arg Phe Glu Ala Val Asp Pro Pro Cys Ala Leu 305 310 315 320			
Phe Pro Tyr Lys Ile Asn Tyr Val Phe Phe Val Ile Thr Ser Trp Phe 325 330 335			
Leu Phe Leu Phe Ile Arg Val Phe Gln Asp Lys Phe Ser Glu Leu Tyr 340 345 350			
Ile Asn Ser Gln Pro Tyr His Ile Ile Leu Gly Cys Gly Cys Ile Leu 355 360 365			
Glu Arg Ile Ala Gln Ala Cys Val Leu Asp Ile Gly Val Gly Ile Met 370 375 380			
Glu Pro Gln Leu Val Ser Phe Ile Ile Trp Leu Glu Pro Ile Asn Cys 385 390 395 400			
Leu Val Leu Trp Phe Gly Gly Glu Val Pro Gly Val Val Gly Lys Gly 405 410 415			
Leu Pro Tyr Gly Val Ala Gly Gly Val Val Asn Ile Gly Val Ile Gly 420 425 430			
Gln Val Gly Gly Gly Tyr Lys Val Gly Ile Gln Asp Asn Asn Ser 435 440 445			
Gly Pro Asn Thr Ser Leu Ile Arg Gly Asp Gly Val Ser Gly Val Lys 450 455 460			
Phe Ile Phe Ser Leu Ser Asn Thr Val Val Leu Glu Arg Gly Val Gly 465 470 475 480			
Ala Ala Gly Gly Glu Glu Leu Ala Asp Val Glu Ser Gln Leu Val Asn 485 490 495			

-continued

```

Ile Pro Arg Trp Leu Arg Val Ser Ser Ser Tyr Gly Glu Tyr Lys Phe
      500                               505                               510

Ser Arg Lys Ala Gly Ile Glu Asp Thr Arg Leu Ser Ala Pro Ser Val
      515                               520                               525

Thr Val Ser Glu Gly Gly Val Tyr Gln Ile Trp Ser Ser Pro Glu Asp
      530                               535                               540

Val Ser Lys Met Ala Ala Gly Ala Gly Pro Ser Ser Ala Val Thr Pro
      545                               550                               555                               560

Pro Trp Pro Arg His Pro Ile Lys Val Lys Glu Val Arg Cys Cys Ser
      565                               570                               575

```

Ile

```

<210> SEQ ID NO 4
<211> LENGTH: 553
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

```

<400> SEQUENCE: 4

```

Gln Arg Thr Ser Ala Ala Ala Ala Pro Arg Gln Arg Gln Lys Cys Gln
 1                               5                               10                               15

Ala Arg Lys Ala Ala Arg Asn Pro Ile Arg Gly Gly Cys Ser Pro Leu
 20                               25                               30

Leu Pro Arg Arg Arg Lys Thr Lys Tyr Gly Ser Phe Gln Ser Pro Phe
 35                               40                               45

Leu Ile Ile Leu Phe Val Ala Arg Lys Val Trp Lys Arg Val Glu Leu
 50                               55                               60

Leu Thr Ser Arg Gly Leu Arg Ile Leu Leu Arg Ser Arg Leu Leu Thr
 65                               70                               75                               80

Arg Ser Gly Ile Leu Val Pro Ala Ala Thr Ser Arg Lys Arg Lys Glu
 85                               90                               95

Pro Thr Ser Arg Ile Lys Asn Thr Ala Val Lys Lys Ala Thr Tyr Leu
100                               105                               110

Ser Ser Val Glu Leu Arg Gly Thr Arg Gly Ser Ala Ala Thr Cys Leu
115                               120                               125

Leu Leu Val Pro Phe Trp Arg Arg Gly Leu Trp Leu Pro Ser Ser Phe
130                               135                               140

Leu Arg Met Glu Ile Ser Ala Gly Trp Leu Asn Phe Lys Ala Gly Arg
145                               150                               155                               160

Cys Arg Ser Val Ile Gly Arg Gln Leu Tyr Thr Ser Trp Ala Arg Pro
165                               170                               175

Val Val Gly Arg Ala Ser Gly Pro Val Ile Leu Leu Ser Leu Gly Thr
180                               185                               190

Pro Thr Gly Ser Leu Val Glu Ile Ser Gly Gly Met Asp Ile Met Glu
195                               200                               205

Lys Lys Leu Leu Phe Trp Met Ile Phe Met Ala Gly Tyr Leu Gly Met
210                               215                               220

Ile Tyr Asp Cys Val Thr Gly Ile His Leu Arg Leu Lys Gly Val Leu
225                               230                               235                               240

Phe Leu Phe Trp Pro Ala Val Phe Leu Pro Ala Ile Arg Pro Pro Arg
245                               250                               255

Asn Gly Thr Pro Gln Leu Leu Ser Gln Leu Lys Leu Ser Ile Gly Gly
260                               265                               270

Leu Leu Leu Cys Asn Phe Gly Arg Leu Leu Glu Asn Asn Pro Arg Arg
275                               280                               285

```

-continued

Tyr Pro Lys Ala Asp Leu Lys Gln Trp Thr His Pro Val Pro Phe Ser
 290 295 300
 His Ile Lys Ile Thr Glu Ser Phe Leu Leu Ser His Arg Asn Gly Phe
 305 310 315 320
 Tyr Phe Tyr Ser Phe Arg Gly Ser Phe Arg Ile Asn Ser Leu Asn Cys
 325 330 335
 Thr Ile Val Asn Leu Thr Thr Phe Trp Ala Val Val Ala Phe Trp Ser
 340 345 350
 Ala Pro Arg Pro Val Cys Ser Thr Leu Val Trp Val Phe Lys Trp Ser
 355 360 365
 His Ser Trp Phe Leu Leu Leu Phe Gly Trp Asn Gln Ser Ile Val Trp
 370 375 380
 Ser Ser Ser Gly Leu Gly Val Lys Tyr Leu Glu Trp Val Lys Gly Cys
 385 390 395 400
 Leu Met Val Trp Arg Glu Glu Leu Ile Gly Ser Ala Lys Leu Val Glu
 405 410 415
 Gly Val Thr Lys Leu Ala Ser Lys Ile Thr Thr Val Asp Pro Thr Pro
 420 425 430
 Leu Leu Glu Val Met Gly Ser Leu Gly Asn Ser Tyr Leu Ala Phe Leu
 435 440 445
 Ile Arg Tyr Trp Lys Gly Arg Gly Arg Gly Leu Val Pro Pro Glu Gly
 450 455 460
 Gly Arg Asn Trp Pro Met Leu Asn Leu Ser Ser Leu Thr Phe Gln Asp
 465 470 475 480
 Gly Cys Glu Cys Pro Pro Leu Met Val Ser Thr Asn Ser Leu Glu Arg
 485 490 495
 Arg Glu Leu Lys Ile Pro Val Phe Arg Arg His Leu Arg Phe Leu Lys
 500 505 510
 Ala Gly Cys Thr Lys Tyr Gly Leu Leu Arg Arg Met Phe Pro Arg Trp
 515 520 525
 Leu Arg Gly Arg Val Arg Leu Leu Arg Arg Leu Leu Gly His Val Ile
 530 535 540
 Leu Lys Lys Lys Cys Ala Ala Val Val
 545 550

<210> SEQ ID NO 5

<211> LENGTH: 1759

<212> TYPE: DNA

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 5

aatactacag cagcgcactt ctttcacttt tataggatga cgtggccaag gaggcgttac 60
 cgcagaagac ggaccgcgcc ccgcagccat cttggaaaacg tcctccggag aagaccatat 120
 ttggtacacc ccgccttcag aaaccggttac agatggcgcc gaaagacggg tatcttcaat 180
 tcccgccttt ctagagaatt tgtactcacc ataagaggag gacactcgca gccatcttgg 240
 aatgttaacg agctgagatt caacatcggc cagttcctcc cccctcagg cggcaccaac 300
 ccctacccc tacctttcca atactaccgt attagaaagg ctaaatatga attttacc 360
 agagacccca tcacctctaa tcaaagaggt gttgggtcca ctgtgttat cttggatgcc 420
 aactttgtaa cccctccac caacttggcc tatgaccct atattaacta ctctcccg 480
 cacaccataa ggcagccctt tacctaccac tccaggtact tcaccccaaa accagagcta 540
 gaccaaacaa ttgattggtt ccagccaaat aataaaagaa accagctgtg gctccattta 600

-continued

```

aataccaca ccaatgtcga gcacacaggc ctgggctatg cgctccaaaa tgcaaccaca 660
gccccaaatt atgtggtaag gttgactatt tatgtacaat tcagagaatt tatectgaaa 720
gaccctctaa atgaataaaa ataaaaacca ttacgatgtg ataacaaaaa agactcagta 780
atattattta tatgggaaaa gggcacaggg tgggtccact gcttcaaatc ggccttcggg 840
tacctccgtg gattgttctc cagcagtctt ccaaaattgc aaagtagtaa tcttccgata 900
gagagcttct acagctggga cagcagttga ggagtacat tcttgggggg cctgattgct 960
ggtaatcaaa atactcgggg ccaaaaaagg aacagtacc cctttagtct ctacagtcaa 1020
tggataccgg tcacacagtc tcagtagatc atcccaaggt aaccagccat aaaaatcacc 1080
caaaacaaca acttcttctc catgatatcc atcccaccac ttatttctac taggcttcca 1140
gtaggtgtcc ctaggctcag caaaattacg ggcccactgg ctcttccacc aaccgggceg 1200
gcccactatg acgtgtacag ctgtcttcca atcacgctgc tgcattctcc cgctcacttt 1260
caaaagtcca gccagcccgc ggaaatttct cacatacgtt acaggaaact gctcggctac 1320
agtcacaaa gaccocgtct ccaaaaggtt actcacagca gtacacaggt cgctgcgctt 1380
cccttggttc cgcggagctc cacactcgat aagtatgtgg ccttctttac tgcagtatc 1440
tttattctgc tggtcgggtc ctttcgcttt ctcgatgtgg cagcgggcac caaaatacca 1500
cttcaccttg ttaaaagtct gcttcttagc aaaattcgca aaccctgga ggtgaggagt 1560
tctacctctt tccaaacctt cctcgcaca aacaaaataa tcaaaaaggg agattggaag 1620
ctcccgtatt ttgttttct cctcctcgga aggattatta aggggaaca cccacctctt 1680
atggggttgc gggccgcttt tcttgcttgg catttctact gacgctgcc aggtgctgcc 1740
gctgccgaag tgcgctggt 1759

```

```

<210> SEQ ID NO 6
<211> LENGTH: 567
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 6

```

```

Gly Ala Cys Lys Pro Leu Pro Leu Val Glu Ala Ala Asp Thr Phe Ile
1 5 10 15
Gly Leu Leu Phe Leu Pro Gly Cys Gly Trp Leu Leu His Thr Asn Val
20 25 30
Arg Leu Leu Gly Glu Ser Ser Ser Phe Phe Leu Ile Arg Ser Ser Gly
35 40 45
Ile Glu Arg Lys Ser Lys Thr Gln Pro Ser Ser Pro Lys Ser Ser Pro
50 55 60
Leu Val Gly Arg Trp Pro Asn Ala Phe Lys Ala Leu Phe Cys Val Lys
65 70 75 80
Leu Leu Thr Phe His Tyr Lys Pro Ala Arg Gln Trp Met Ser Phe Ala
85 90 95
Phe Pro Val Ser Trp Cys Phe Leu Ser Tyr Gln Leu Leu Ser Pro Trp
100 105 110
Met Ser Ile Ser His Pro Ala Gly Arg Phe Trp Pro Phe Arg Leu Ser
115 120 125
Arg Asp Val Ala Thr Leu Val Arg Lys Ser Val Pro Asp Lys Thr Val
130 135 140
Thr Ala Ser Cys Asn Gly Thr Val Tyr Thr Leu Phe Lys Arg Pro Ser
145 150 155 160

```


-continued

```

<210> SEQ ID NO 7
<211> LENGTH: 580
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 7

Trp Arg Val Glu Ala Ala Ala Ala Gly Arg Cys Arg His Phe His Trp
1          5          10          15
Ala Leu Phe Ala Ala Arg Leu Gly Met Leu Pro Pro His Glu Gly Lys
20          25          30
Ile Ile Arg Gly Leu Leu Leu Phe Val Phe Tyr Pro Leu Lys Trp Asp
35          40          45
Gly Lys Lys Ile Ile Lys Asn Thr Ala Leu Phe Thr Gln Phe Leu Thr
50          55          60
Ser Ser Arg Val Glu Leu Pro Lys Arg Ile Lys Ser Leu Leu Leu Ser
65          70          75          80
Lys Val Leu His Leu Pro Ile Lys Thr Gly Ala Ala Val Asp Leu Phe
85          90          95
Arg Phe Ser Gly Val Leu Leu Ile Phe Phe Val Ala Thr Phe Phe Ala
100         105         110
Val Tyr Lys Asp Leu Thr Ser Ser Arg Pro Val Leu Pro Leu Ala Ala
115         120         125
Val Gln Arg Ser Ser His Thr Gly Lys Gln Leu Arg Pro Arg Gln His
130         135         140
Ser Tyr Gly Leu Leu Lys Arg Tyr Arg Ile His Ser Ile Glu Ala Pro
145         150         155         160
Gln Ser Phe Lys Gln Phe His Ala Pro Leu His Leu Leu Thr Ile Pro
165         170         175
Leu Cys Ser Tyr Val Asp Tyr His Ala Arg Gly Thr Thr Pro Leu Ala
180         185         190
Leu Pro Gly Thr Ile Lys Ser Leu Arg Pro Val Gly Val Pro Leu Arg
195         200         205
Thr Ser Ile Leu Pro Pro Ile Ser Ile Met Ser Phe Phe Asn Asn Asn
210         215         220
Gln Ile Ile Lys Ile Ala Pro Arg Pro Ile Ile Gln Ser Gln Thr Val
225         230         235         240
Pro Ile Trp Gln Ser Tyr Leu Ser Phe Pro Thr Ser Asn Arg Lys Gln
245         250         255
Gly Ala Thr Asn Gln Asn Gly Ala Ile Leu Gly Gly Leu Phe Pro Val
260         265         270
Gly Ser Ser Asp Trp Ser Tyr Phe Ser Glu Ile Pro Pro Asn Ser Ser
275         280         285
Gln Leu Lys Pro Leu Ser Ser Ser Phe Leu Gly Arg Leu Tyr Gly Phe
290         295         300
Ala Ser Lys Phe Cys His Val Trp Gly Thr Gly Lys Glu Trp Ile Phe
305         310         315         320
Tyr Ile Val Ser Asp Lys Lys Asn Asp Cys Arg Leu Pro Lys Lys Glu
325         330         335
Asn Leu Pro Asp Lys Leu Ile Phe Glu Arg Phe Gln Val Tyr Ile Thr
340         345         350
Leu Arg Val Val Tyr Asn Gln Ala Thr Thr Ala Asn Gln Leu Ala Tyr
355         360         365
Gly Leu Gly Thr His Glu Val Asn Thr His Thr Asn Leu His Leu Trp
370         375         380

```

-continued

Leu Gln Asn Arg Lys Asn Asn Pro Gln Phe Trp Asp Ile Thr Gln Asp
 385 390 395 400
 Leu Glu Pro Lys Pro Thr Phe Tyr Arg Ser His Tyr Thr Phe Pro Gln
 405 410 415
 Arg Ile Thr His Arg Ser Ser Tyr Asn Ile His Pro Asp Tyr Ala Leu
 420 425 430
 Asn Thr Ser Pro Thr Val Phe Asn Ala Asp Leu Ile Val Val Thr Ser
 435 440 445
 Gly Val Gly Arg Gln Asn Ser Thr Ile Pro Asp Arg Pro Tyr Phe Glu
 450 455 460
 Tyr Lys Ala Lys Arg Ile Arg Tyr Tyr Gln Phe Pro Leu Pro Leu Pro
 465 470 475 480
 Asn Thr Gly Gly Ser Pro Pro Leu Phe Gln Gly Ile Asn Phe Arg Leu
 485 490 495
 Glu Asn Val Asn Trp Ser Pro Gln Ser His Gly Gly Arg Ile Thr Leu
 500 505 510
 Val Phe Glu Arg Ser Leu Arg Ser Asn Phe Ile Gly Thr Lys Arg Arg
 515 520 525
 Trp Arg Tyr Arg Asn Arg Phe Ala Pro His Val Leu Tyr Pro Arg Arg
 530 535 540
 Arg Leu Ile Asn Gly Leu His Ser Arg Pro Arg Thr Arg Arg Arg Arg
 545 550 555 560
 Tyr Arg Arg Arg Pro Trp Thr Met Arg Tyr Phe His Phe Phe His Ala
 565 570 575
 Ala Thr Thr Asn
 580

<210> SEQ ID NO 8
 <211> LENGTH: 557
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 8

Leu Ala Ser Arg Cys Arg Cys Cys Arg Pro Leu Thr Leu Ser Phe Ala
 1 5 10 15
 Leu Cys Ser Phe Arg Gly Ala Val Gly Tyr Ser Thr Pro Thr Gly Tyr
 20 25 30
 Asp Lys Arg Pro Pro Ser Phe Cys Phe Val Pro Ala Glu Leu Arg Gly
 35 40 45
 Lys Gln Asn Asn Gln Lys His Arg Pro Leu Asn Pro Leu Pro Tyr Phe
 50 55 60
 Glu Glu Gly Gly Pro Thr Gln Ser Asn Gln Ser Ala Ser Lys Cys Pro
 65 70 75 80
 Ser Thr Thr Asn Gly His Gly Ser Gly Cys Arg Ser Leu Ser Leu Phe
 85 90 95
 Arg Gly Ala Ser Tyr Leu Ile Ser Cys Tyr Leu Leu Gly Cys Val Arg
 100 105 110
 Thr His Leu Glu Ala Ser Gly Pro Ser Ala Cys Arg Gly Thr Gln Gln
 115 120 125
 Ser Tyr Gly Lys Pro Ser Pro Thr Lys Pro Ser Gln Leu Arg Ala Thr
 130 135 140
 Glu Gln Leu Thr His Ser Phe Asn Gly Arg Ala Pro Gln Val Lys Ser
 145 150 155 160
 Leu Ser Arg Ser Ser Ala Ala Ala His Asn Ser Ser Leu Gln Val Arg
 165 170 175

-continued

Leu Pro Gly Ala Arg Asn His Ser Ser Gly Thr Pro Gly Tyr Asn Gln
 180 185 190
 Gln Ala Pro Cys Arg Ser Ser Ala Tyr Phe Tyr Thr Thr Pro His Ile
 195 200 205
 Asp His Leu Leu Leu Gln Gln Lys Pro His Asn Lys His Ser Thr Val
 210 215 220
 Lys Pro His Asp Val Ser Val Thr His Gly Thr Asp Met Ser Gln Leu
 225 230 235 240
 Ser Leu Pro Tyr Gln Glu Lys Lys Pro Gly Cys Tyr Lys Ser Trp Cys
 245 250 255
 Asp Pro Gly Gly Pro Ile Thr Ser Arg Leu Gln Gln Gly Leu Gln Leu
 260 265 270
 Leu Glu Arg Asp Ser Ser Lys Ala Ile Lys Ser Ser Gln Gln Leu Val
 275 280 285
 Ile Trp Pro Pro Val Arg Leu Gly Ile Gln Leu Leu Pro Gly Val Arg
 290 295 300
 His Gly Lys Gly Met Tyr Phe Leu Asn Ser Leu Arg Lys Gln Met Thr
 305 310 315 320
 Ile Thr Lys Ile Lys Ile Lys Ser Pro Arg Glu Pro Tyr Ile Arg Gln
 325 330 335
 Ile Thr Cys Leu Tyr Asp Val Lys Gly Cys Leu Lys Pro Ser His Asn
 340 345 350
 Cys Lys Pro Ala Cys Leu Gly Pro Arg His Ala Arg Cys Gln His Pro
 355 360 365
 Tyr Lys Phe Pro Ala Val Val Pro Lys Lys Lys Ala Pro Val Leu Asn
 370 375 380
 Asn Pro Arg Ala Arg Thr Gln Pro His Leu Val Gln Leu Pro Leu Tyr
 385 390 395 400
 Leu Ala Ala Lys His His Pro Pro Leu Leu Leu Tyr Leu Pro Leu Gly
 405 410 415
 Leu Gln His Leu Pro Asn Cys Leu Gln Cys Gly Leu Tyr Cys Cys His
 420 425 430
 Val Trp Cys Arg Lys Ser Leu His His Pro Arg Gln Pro Leu Ile Ile
 435 440 445
 Gly Lys Tyr Pro Leu Ile Pro Phe Thr Pro Thr Pro Pro Gln His Arg
 450 455 460
 Arg Leu Pro Pro Pro Val Pro Arg His Gln Ile Glu Ala Arg Cys Glu
 465 470 475 480
 Leu Ile Ala Ala Leu Thr Arg Arg Lys His His Thr Cys Ile Arg Phe
 485 490 495
 Pro Pro Phe Gln Leu Tyr Gly Asp Lys Pro Ala Met Gln Leu Pro Lys
 500 505 510
 Gln Leu Arg Pro Thr Gly Phe Ile Thr Lys Glu Pro Pro His Lys Trp
 515 520 525
 Ser Pro Gln Pro Pro Pro Asp Thr Lys Gln Pro Leu Ala Glu Lys Ala
 530 535 540
 Val Asp Asp Leu Leu Ser Leu Leu Ala Ser Ser Tyr Tyr
 545 550 555

<210> SEQ ID NO 9
 <211> LENGTH: 939
 <212> TYPE: DNA
 <213> ORGANISM: Type A PWD circovirus
 <220> FEATURE:

-continued

<221> NAME/KEY: CDS

<222> LOCATION: (1) .. (936)

<400> SEQUENCE: 9

```

atg cca agc aag aaa agc ggc ccg caa ccc cat aag agg tgg gtg ttc      48
Met Pro Ser Lys Lys Ser Gly Pro Gln Pro His Lys Arg Trp Val Phe
1           5           10           15

acc ctt aat aat cct tcc gag gag gag aaa aac aaa ata cgg gag ctt      96
Thr Leu Asn Asn Pro Ser Glu Glu Glu Lys Asn Lys Ile Arg Glu Leu
20           25           30

cca atc tcc ctt ttt gat tat ttt gtt tgt ggc gag gaa ggt ttg gaa     144
Pro Ile Ser Leu Phe Asp Tyr Phe Val Cys Gly Glu Glu Gly Leu Glu
35           40           45

gag ggt aga act cct cac ctc cag ggg ttt gcg aat ttt gct aag aag     192
Glu Gly Arg Thr Pro His Leu Gln Gly Phe Ala Asn Phe Ala Lys Lys
50           55           60

cag act ttt aac aag gtg aag tgg tat ttt ggt gcc cgc tgc cac atc     240
Gln Thr Phe Asn Lys Val Lys Trp Tyr Phe Gly Ala Arg Cys His Ile
65           70           75           80

gag aaa gcg aaa gga acc gac cag cag aat aaa gaa tac tgc agt aaa     288
Glu Lys Ala Lys Gly Thr Asp Gln Gln Asn Lys Glu Tyr Cys Ser Lys
85           90           95

gaa ggc cac ata ctt atc gag tgt gga gct ccg cgg aac cag ggg aag     336
Glu Gly His Ile Leu Ile Glu Cys Gly Ala Pro Arg Asn Gln Gly Lys
100          105          110

cgc agc gac ctg tct act gct gtg agt acc ctt ttg gag acg ggg tct     384
Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu Thr Gly Ser
115          120          125

ttg gtg act gta gcc gag cag ttt cct gta acg tat gtg aga aat ttc     432
Leu Val Thr Val Ala Glu Gln Phe Pro Val Thr Tyr Val Arg Asn Phe
130          135          140

cgc ggg ctg gct gaa ctt ttg aaa gtg agc ggg aag atg cag cag cgt     480
Arg Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met Gln Gln Arg
145          150          155          160

gat tgg aag aca gct gta cac gtc ata gtg ggc ccg ccc ggt tgt ggg     528
Asp Trp Lys Thr Ala Val His Val Ile Val Gly Pro Pro Gly Cys Gly
165          170          175

aag agc cag tgg gcc cgt aat ttt gct gag cct agg gac acc tac tgg     576
Lys Ser Gln Trp Ala Arg Asn Phe Ala Glu Pro Arg Asp Thr Tyr Trp
180          185          190

aag cct agt aga aat aag tgg tgg gat gga tat cat gga gaa gaa gtt     624
Lys Pro Ser Arg Asn Lys Trp Trp Asp Gly Tyr His Gly Glu Glu Val
195          200          205

gtt gtt ttg gat gat ttt tat ggc tgg tta cct tgg gat gat cta ctg     672
Val Val Leu Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp Asp Leu Leu
210          215          220

aga ctg tgt gac cgg tat cca ttg act gta gag act aaa ggg ggt act     720
Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Glu Thr Lys Gly Gly Thr
225          230          235          240

gtt cct ttt ttg gcc cgc agt att ttg att acc agc aat cag gcc ccc     768
Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn Gln Ala Pro
245          250          255

cag gaa tgg tac tcc tca act gct gtc cca gct gta gaa gct ctc tat     816
Gln Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu Ala Leu Tyr
260          265          270

cgg agg att act act ttg caa ttt tgg aag act gct gga gaa caa tcc     864
Arg Arg Ile Thr Thr Leu Gln Phe Trp Lys Thr Ala Gly Glu Gln Ser
275          280          285

acg gag gta ccc gaa ggc cga ttt gaa gca gtg gac cca ccc tgt gcc     912
Thr Glu Val Pro Glu Gly Arg Phe Glu Ala Val Asp Pro Pro Cys Ala

```

-continued

290	295	300	
ctt ttc cca tat aaa ata aat tac tga			939
Leu Phe Pro Tyr Lys Ile Asn Tyr			
305	310		

<210> SEQ ID NO 10
 <211> LENGTH: 312
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 10

Met	Pro	Ser	Lys	Lys	Ser	Gly	Pro	Gln	Pro	His	Lys	Arg	Trp	Val	Phe
1			5					10						15	
Thr	Leu	Asn	Asn	Pro	Ser	Glu	Glu	Glu	Lys	Asn	Lys	Ile	Arg	Glu	Leu
		20					25						30		
Pro	Ile	Ser	Leu	Phe	Asp	Tyr	Phe	Val	Cys	Gly	Glu	Glu	Gly	Leu	Glu
		35					40					45			
Glu	Gly	Arg	Thr	Pro	His	Leu	Gln	Gly	Phe	Ala	Asn	Phe	Ala	Lys	Lys
	50					55					60				
Gln	Thr	Phe	Asn	Lys	Val	Lys	Trp	Tyr	Phe	Gly	Ala	Arg	Cys	His	Ile
65					70					75					80
Glu	Lys	Ala	Lys	Gly	Thr	Asp	Gln	Gln	Asn	Lys	Glu	Tyr	Cys	Ser	Lys
				85					90						95
Glu	Gly	His	Ile	Leu	Ile	Glu	Cys	Gly	Ala	Pro	Arg	Asn	Gln	Gly	Lys
			100					105					110		
Arg	Ser	Asp	Leu	Ser	Thr	Ala	Val	Ser	Thr	Leu	Leu	Glu	Thr	Gly	Ser
		115					120						125		
Leu	Val	Thr	Val	Ala	Glu	Gln	Phe	Pro	Val	Thr	Tyr	Val	Arg	Asn	Phe
	130					135					140				
Arg	Gly	Leu	Ala	Glu	Leu	Leu	Lys	Val	Ser	Gly	Lys	Met	Gln	Gln	Arg
145					150					155					160
Asp	Trp	Lys	Thr	Ala	Val	His	Val	Ile	Val	Gly	Pro	Pro	Gly	Cys	Gly
				165					170						175
Lys	Ser	Gln	Trp	Ala	Arg	Asn	Phe	Ala	Glu	Pro	Arg	Asp	Thr	Tyr	Trp
			180					185						190	
Lys	Pro	Ser	Arg	Asn	Lys	Trp	Trp	Asp	Gly	Tyr	His	Gly	Glu	Glu	Val
		195					200					205			
Val	Val	Leu	Asp	Asp	Phe	Tyr	Gly	Trp	Leu	Pro	Trp	Asp	Asp	Leu	Leu
	210						215					220			
Arg	Leu	Cys	Asp	Arg	Tyr	Pro	Leu	Thr	Val	Glu	Thr	Lys	Gly	Gly	Thr
225					230					235					240
Val	Pro	Phe	Leu	Ala	Arg	Ser	Ile	Leu	Ile	Thr	Ser	Asn	Gln	Ala	Pro
				245					250					255	
Gln	Glu	Trp	Tyr	Ser	Ser	Thr	Ala	Val	Pro	Ala	Val	Glu	Ala	Leu	Tyr
			260					265					270		
Arg	Arg	Ile	Thr	Thr	Leu	Gln	Phe	Trp	Lys	Thr	Ala	Gly	Glu	Gln	Ser
		275					280					285			
Thr	Glu	Val	Pro	Glu	Gly	Arg	Phe	Glu	Ala	Val	Asp	Pro	Pro	Cys	Ala
	290					295					300				
Leu	Phe	Pro	Tyr	Lys	Ile	Asn	Tyr								
305					310										

<210> SEQ ID NO 11
 <211> LENGTH: 702
 <212> TYPE: DNA
 <213> ORGANISM: Type A PWD circovirus

-continued

```

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (699)

<400> SEQUENCE: 11

atg acg tgg cca agg agg cgt tac cgc aga aga cgg acc cgc ccc cgc      48
Met Thr Trp Pro Arg Arg Arg Tyr Arg Arg Arg Arg Thr Arg Pro Arg
1           5           10           15

agc cat ctt gga aac atc ctc cgg aga aga cca tat ttg gta cac ccc      96
Ser His Leu Gly Asn Ile Leu Arg Arg Arg Pro Tyr Leu Val His Pro
           20           25           30

gcc ttc aga aac cgt tac aga tgg cgc cga aag acg ggt atc ttc aat     144
Ala Phe Arg Asn Arg Tyr Arg Trp Arg Arg Lys Thr Gly Ile Phe Asn
           35           40           45

tcc cgc ctt tct aga gaa ttt gta ctc acc ata aga gga gga cac tcg     192
Ser Arg Leu Ser Arg Glu Phe Val Leu Thr Ile Arg Gly Gly His Ser
           50           55           60

cag cca tct tgg aat gtt aac gag ctg aga ttc aac atc ggc cag ttc     240
Gln Pro Ser Trp Asn Val Asn Glu Leu Arg Phe Asn Ile Gly Gln Phe
           65           70           75           80

ctc ccc ccc tca ggc ggc acc aac ccc cta ccc cta cct ttc caa tac     288
Leu Pro Pro Ser Gly Gly Thr Asn Pro Leu Pro Leu Pro Phe Gln Tyr
           85           90           95

tac cgt att aga aag gct aaa tat gaa ttt tac ccc aga gac ccc atc     336
Tyr Arg Ile Arg Lys Ala Lys Tyr Glu Phe Tyr Pro Arg Asp Pro Ile
           100          105          110

acc tct aat caa aga ggt gtt ggg tcc act gtt gtt atc ttg gat gcc     384
Thr Ser Asn Gln Arg Gly Val Gly Ser Thr Val Val Ile Leu Asp Ala
           115          120          125

aac ttt gta acc ccc tcc acc aac ttg gcc tat gac ccc tat att aac     432
Asn Phe Val Thr Pro Ser Thr Asn Leu Ala Tyr Asp Pro Tyr Ile Asn
           130          135          140

tac tcc tcc cgc cac acc ata agg cag ccc ttt acc tac cac tcc agg     480
Tyr Ser Ser Arg His Thr Ile Arg Gln Pro Phe Thr Tyr His Ser Arg
           145          150          155          160

tac ttc acc ccc aaa cca gag cta gac caa aca att gat tgg ttc cag     528
Tyr Phe Thr Pro Lys Pro Glu Leu Asp Gln Thr Ile Asp Trp Phe Gln
           165          170          175

cca aat aat aaa aga aac cag ctg tgg ctc cat tta aat acc cac acc     576
Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu His Leu Asn Thr His Thr
           180          185          190

aat gtc gag cac aca ggc ctg ggc tat gcg ctc caa aat gca acc aca     624
Asn Val Glu His Thr Gly Leu Gly Tyr Ala Leu Gln Asn Ala Thr Thr
           195          200          205

gcc caa aat tat gtg gta agg ttg act att tat gta caa ttc aga gaa     672
Ala Gln Asn Tyr Val Val Arg Leu Thr Ile Tyr Val Gln Phe Arg Glu
           210          215          220

ttt atc ctg aaa gac cct cta aat gaa taa                               702
Phe Ile Leu Lys Asp Pro Leu Asn Glu
           225          230

<210> SEQ ID NO 12
<211> LENGTH: 233
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 12

Met Thr Trp Pro Arg Arg Arg Tyr Arg Arg Arg Arg Thr Arg Pro Arg
1           5           10           15

Ser His Leu Gly Asn Ile Leu Arg Arg Arg Pro Tyr Leu Val His Pro
           20           25           30

```

-continued

Ala Phe Arg Asn Arg Tyr Arg Trp Arg Arg Lys Thr Gly Ile Phe Asn
35 40 45

Ser Arg Leu Ser Arg Glu Phe Val Leu Thr Ile Arg Gly Gly His Ser
50 55 60

Gln Pro Ser Trp Asn Val Asn Glu Leu Arg Phe Asn Ile Gly Gln Phe
65 70 75 80

Leu Pro Pro Ser Gly Gly Thr Asn Pro Leu Pro Leu Pro Phe Gln Tyr
85 90 95

Tyr Arg Ile Arg Lys Ala Lys Tyr Glu Phe Tyr Pro Arg Asp Pro Ile
100 105 110

Thr Ser Asn Gln Arg Gly Val Gly Ser Thr Val Val Ile Leu Asp Ala
115 120 125

Asn Phe Val Thr Pro Ser Thr Asn Leu Ala Tyr Asp Pro Tyr Ile Asn
130 135 140

Tyr Ser Ser Arg His Thr Ile Arg Gln Pro Phe Thr Tyr His Ser Arg
145 150 155 160

Tyr Phe Thr Pro Lys Pro Glu Leu Asp Gln Thr Ile Asp Trp Phe Gln
165 170 175

Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu His Leu Asn Thr His Thr
180 185 190

Asn Val Glu His Thr Gly Leu Gly Tyr Ala Leu Gln Asn Ala Thr Thr
195 200 205

Ala Gln Asn Tyr Val Val Arg Leu Thr Ile Tyr Val Gln Phe Arg Glu
210 215 220

Phe Ile Leu Lys Asp Pro Leu Asn Glu
225 230

<210> SEQ ID NO 13
<211> LENGTH: 621
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(618)

<400> SEQUENCE: 13

atg ata tcc atc cca cca ctt att tct act agg ctt cca gta ggt gtc 48
Met Ile Ser Ile Pro Pro Leu Ile Ser Thr Arg Leu Pro Val Gly Val
1 5 10 15

cct agg ctc agc aaa att acg ggc cca ctg gct ctt ccc aca acc ggg 96
Pro Arg Leu Ser Lys Ile Thr Gly Pro Leu Ala Leu Pro Thr Thr Gly
20 25 30

cgg gcc cac tat gac gtg tac agc tgt ctt cca atc acg ctg ctg cat 144
Arg Ala His Tyr Asp Val Tyr Ser Cys Leu Pro Ile Thr Leu Leu His
35 40 45

ctt ccc gct cac ttt caa aag ttc agc cag ccc gcg gaa att tct cac 192
Leu Pro Ala His Phe Gln Lys Phe Ser Gln Pro Ala Glu Ile Ser His
50 55 60

ata cgt tac agg aaa ctg ctc ggc tac agt cac caa aga ccc cgt ctc 240
Ile Arg Tyr Arg Lys Leu Leu Gly Tyr Ser His Gln Arg Pro Arg Leu
65 70 75 80

caa aag ggt act cac agc agt aga cag gtc gct gcg ctt ccc ctg gtt 288
Gln Lys Gly Thr His Ser Ser Arg Gln Val Ala Ala Leu Pro Leu Val
85 90 95

ccg cgg agc tcc aca ctc gat aag tat gtg gcc ttc ttt act gca gta 336
Pro Arg Ser Ser Thr Leu Asp Lys Tyr Val Ala Phe Phe Thr Ala Val
100 105 110

-continued

ttc ttt att ctg ctg gtc ggt tcc ttt cgc ttt ctc gat gtg gca gcg	384
Phe Phe Ile Leu Leu Val Gly Ser Phe Arg Phe Leu Asp Val Ala Ala	
115 120 125	
ggc acc aaa ata cca ctt cac ctt gtt aaa agt ctg ctt ctt agc aaa	432
Gly Thr Lys Ile Pro Leu His Leu Val Lys Ser Leu Leu Leu Ser Lys	
130 135 140	
att cgc aaa ccc ctg gag gtg agg agt tct acc ctc ttc caa acc ttc	480
Ile Arg Lys Pro Leu Glu Val Arg Ser Ser Thr Leu Phe Gln Thr Phe	
145 150 155 160	
ctc gcc aca aac aaa ata atc aaa aag gga gat tgg aag ctc ccg tat	528
Leu Ala Thr Asn Lys Ile Ile Lys Lys Gly Asp Trp Lys Leu Pro Tyr	
165 170 175	
ttt gtt ttt ctc ctc ctc gga agg att att aag ggt gaa cac cca cct	576
Phe Val Phe Leu Leu Leu Gly Arg Ile Ile Lys Gly Glu His Pro Pro	
180 185 190	
ctt atg ggg ttg cgg gcc gct ttt ctt gct tgg cat ttt cac tga	621
Leu Met Gly Leu Arg Ala Ala Phe Leu Ala Trp His Phe His	
195 200 205	

<210> SEQ ID NO 14

<211> LENGTH: 206

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 14

Met Ile Ser Ile Pro Pro Leu Ile Ser Thr Arg Leu Pro Val Gly Val	
1 5 10 15	
Pro Arg Leu Ser Lys Ile Thr Gly Pro Leu Ala Leu Pro Thr Thr Gly	
20 25 30	
Arg Ala His Tyr Asp Val Tyr Ser Cys Leu Pro Ile Thr Leu Leu His	
35 40 45	
Leu Pro Ala His Phe Gln Lys Phe Ser Gln Pro Ala Glu Ile Ser His	
50 55 60	
Ile Arg Tyr Arg Lys Leu Leu Gly Tyr Ser His Gln Arg Pro Arg Leu	
65 70 75 80	
Gln Lys Gly Thr His Ser Ser Arg Gln Val Ala Ala Leu Pro Leu Val	
85 90 95	
Pro Arg Ser Ser Thr Leu Asp Lys Tyr Val Ala Phe Phe Thr Ala Val	
100 105 110	
Phe Phe Ile Leu Leu Val Gly Ser Phe Arg Phe Leu Asp Val Ala Ala	
115 120 125	
Gly Thr Lys Ile Pro Leu His Leu Val Lys Ser Leu Leu Leu Ser Lys	
130 135 140	
Ile Arg Lys Pro Leu Glu Val Arg Ser Ser Thr Leu Phe Gln Thr Phe	
145 150 155 160	
Leu Ala Thr Asn Lys Ile Ile Lys Lys Gly Asp Trp Lys Leu Pro Tyr	
165 170 175	
Phe Val Phe Leu Leu Leu Gly Arg Ile Ile Lys Gly Glu His Pro Pro	
180 185 190	
Leu Met Gly Leu Arg Ala Ala Phe Leu Ala Trp His Phe His	
195 200 205	

<210> SEQ ID NO 15

<211> LENGTH: 1767

<212> TYPE: DNA

<213> ORGANISM: Type B PWD circovirus

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(111)

-continued

```

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (115)..(243)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (247)..(267)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (271)..(360)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (364)..(417)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (421)..(447)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (451)..(471)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (475)..(510)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (514)..(516)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (520)..(729)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (733)..(753)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (757)..(759)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (763)..(804)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (808)..(861)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (865)..(984)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (988)..(1173)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1177)..(1233)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1237)..(1359)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1363)..(1476)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1480)..(1737)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1741)..(1767)

<400> SEQUENCE: 15

acc agc gca ctt cgg cag cgg cag cac ctc ggc agc acc tca gca gca      48
Thr Ser Ala Leu Arg Gln Arg Gln His Leu Gly Ser Thr Ser Ala Ala
1          5          10          15

aca tgc cca gca aga aga atg gaa gaa gcg gac ccc aac ccc ata aaa      96
Thr Cys Pro Ala Arg Arg Met Glu Glu Ala Asp Pro Asn Pro Ile Lys
          20          25          30

ggt ggg tgt tca ctc tga ata atc ctt ccg aag acg agc gca aga aaa      144
Gly Gly Cys Ser Leu Ile Ile Leu Pro Lys Thr Ser Ala Arg Lys
          35          40          45

tac ggg atc ttc caa tat ccc tat ttg att att tta ttg ttg gcg agg      192
Tyr Gly Ile Phe Gln Tyr Pro Tyr Leu Ile Ile Leu Leu Leu Ala Arg
          50          55          60

```

-continued

agg gta atg agg aag gac gaa cac ctc acc tcc agg ggt tcg cta att Arg Val Met Arg Lys Asp Glu His Leu Thr Ser Arg Gly Ser Leu Ile 65 70 75	240
ttg tga aga agc aga ctt tta ata aag tga agt ggt att tgg gtg ccc Leu Arg Ser Arg Leu Leu Ile Lys Ser Gly Ile Trp Val Pro 80 85 90	288
gct gcc aca tcg aga aag cga aag gaa cag atc agc aga ata aag aat Ala Ala Thr Ser Arg Lys Arg Lys Glu Gln Ile Ser Arg Ile Lys Asn 95 100 105	336
act gca gta aag aag gca act tac tga tgg agt gtg gag ctc cta gat Thr Ala Val Lys Lys Ala Thr Tyr Trp Ser Val Glu Leu Leu Asp 110 115 120	384
ctc agg gac aac gga gtg acc tgt cta ctg ctg tga gta cct tgt tgg Leu Arg Asp Asn Gly Val Thr Cys Leu Leu Leu Val Pro Cys Trp 125 130 135	432
aga gcg gga gtc tgg tga ccg ttg cag agc agc acc ctg taa cgt ttg Arg Ala Gly Val Trp Pro Leu Gln Ser Ser Thr Leu Arg Leu 140 145 150	480
tca gaa att tcc gcg ggc tgg ctg aac ttt tga aag tga gcg gga aaa Ser Glu Ile Ser Ala Gly Trp Leu Asn Phe Lys Ala Gly Lys 155 160 165	528
tgc aga agc gtg att gga aga cta atg tac acg tca ttg tgg ggc cac Cys Arg Ser Val Ile Gly Arg Leu Met Tyr Thr Ser Leu Trp Gly His 170 175 180	576
ctg ggt gtg gta aaa gca aat ggg ctg cta att ttg cag acc cgg aaa Leu Gly Val Val Lys Ala Asn Gly Leu Leu Ile Leu Gln Thr Arg Lys 185 190 195	624
cca cat act gga aac cac cta gaa aca agt ggt ggg atg gtt acc atg Pro His Thr Gly Asn His Leu Glu Thr Ser Gly Gly Met Val Thr Met 200 205 210 215	672
gtg aag aag tgg ttg tta ttg atg act ttt atg gct ggc tgc cct ggg Val Lys Lys Trp Leu Leu Leu Met Thr Phe Met Ala Gly Cys Pro Gly 220 225 230	720
atg atc tac tga gac tgt gtg atc gat atc cat tga ctg tag aga cta Met Ile Tyr Asp Cys Val Ile Asp Ile His Leu Arg Leu 235 240	768
aag gtg gaa ctg tac ctt ttt tgg ccc gca gta ttc tga tta cca gca Lys Val Glu Leu Tyr Leu Phe Trp Pro Ala Val Phe Leu Pro Ala 245 250 255	816
atc aga ccc cgt tgg aat ggt act cct caa ctg ctg tcc cag ctg tag Ile Arg Pro Arg Trp Asn Gly Thr Pro Gln Leu Leu Ser Gln Leu 260 265 270	864
aag ctc ttt atc gga gga tta ctt cct tgg tat ttt gga aga atg cta Lys Leu Phe Ile Gly Gly Leu Leu Pro Trp Tyr Phe Gly Arg Met Leu 275 280 285 290	912
cag aac aat cca cgg agg aag ggg gcc agt tcg tca ccc ttt ccc ccc Gln Asn Asn Pro Arg Arg Lys Gly Ala Ser Ser Ser Pro Phe Pro Pro 295 300 305	960
cat gcc ctg aat ttc cat atg aaa taa att act gag tct ttt tta tca His Ala Leu Asn Phe His Met Lys Ile Thr Glu Ser Phe Leu Ser 310 315 320	1008
ctt cgt aat ggt ttt tat tat tca tta agg gtt aag tgg ggg gtc ttt Leu Arg Asn Gly Phe Tyr Tyr Ser Leu Arg Val Lys Trp Gly Val Phe 325 330 335	1056
aaa att aaa ttc tct gaa ttg tac ata cat ggt tac acg gat att gta Lys Ile Lys Phe Ser Glu Leu Tyr Ile His Gly Tyr Thr Asp Ile Val 340 345 350	1104
ttc ctg gtc gta tat act gtt ttc gaa cgc agt gcc gag gcc tac gtg Phe Leu Val Val Tyr Thr Val Phe Glu Arg Ser Ala Glu Ala Tyr Val 355 360 365	1152

-continued

```

gtc tac att tcc agc agt ttg tag tct cag cca cag ctg gtt tct ttt 1200
Val Tyr Ile Ser Ser Ser Leu Ser Gln Pro Gln Leu Val Ser Phe
370 375 380

gtt gtt tgg ttg gaa gta atc aat agt gaa atc tag gac agg ttt ggg 1248
Val Val Trp Leu Glu Val Ile Asn Ser Glu Ile Asp Arg Phe Gly
385 390 395

ggt aaa gta ccg gga gtg gta gga gaa ggg ctg ggt tat ggt atg gcg 1296
Gly Lys Val Pro Gly Val Val Gly Glu Gly Leu Gly Tyr Gly Met Ala
400 405 410 415

gga gga gta gtt tac ata ggg gtc ata ggt gag ggc tgt ggc ctt tgt 1344
Gly Gly Val Val Tyr Ile Gly Val Ile Gly Glu Gly Cys Gly Leu Cys
420 425 430

tac aaa gtt atc atc taa aat aac agc act gga gcc cac tcc cct gtc 1392
Tyr Lys Val Ile Ile Asn Asn Ser Thr Gly Ala His Ser Pro Val
435 440 445

acc ctg ggt gat ccg gga gca ggg cca gaa ttc aac ctt aac ctt tct 1440
Thr Leu Gly Asp Arg Gly Ala Gly Pro Glu Phe Asn Leu Asn Leu Ser
450 455 460

tat tct gta gta ttc aaa ggg cac aga gcg ggg gtt tga ccc ccc tcc 1488
Tyr Ser Val Val Phe Lys Gly His Arg Ala Gly Val Pro Pro Ser
465 470 475

tgg ggg aag aaa gtc att aat att gaa tct cat cat gtc cac cgc cca 1536
Trp Gly Lys Lys Val Ile Asn Ile Glu Ser His His Val His Arg Pro
480 485 490

gga ggg cgt tct gac tgt ggt tcg ctt gac agt ata tcc gaa ggt gcg 1584
Gly Gly Arg Ser Asp Cys Gly Ser Leu Asp Ser Ile Ser Glu Gly Ala
495 500 505

gga gag gcg ggt gtt gaa gat gcc att ttt cct tct cca gcg gta acg 1632
Gly Glu Ala Gly Val Glu Asp Ala Ile Phe Pro Ser Pro Ala Val Thr
510 515 520 525

gtg gcg ggg gtg gac gag cca ggg gcg gcg gcg gag gat ctg gcc aag 1680
Val Ala Gly Val Asp Glu Pro Gly Ala Ala Ala Glu Asp Leu Ala Lys
530 535 540

atg gct gcg ggg gcg gtg tct tct tct tcg gta acg cct cct tgg ata 1728
Met Ala Ala Gly Ala Val Ser Ser Ser Ser Val Thr Pro Pro Trp Ile
545 550 555

cgt cat atc tga aaa cga aag aag tgc gct gta agt att 1767
Arg His Ile Lys Arg Lys Lys Cys Ala Val Ser Ile
560 565

```

```

<210> SEQ ID NO 16
<211> LENGTH: 569
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

```

<400> SEQUENCE: 16

```

```

Thr Ser Ala Leu Arg Gln Arg Gln His Leu Gly Ser Thr Ser Ala Ala
1 5 10 15

Thr Cys Pro Ala Arg Arg Met Glu Glu Ala Asp Pro Asn Pro Ile Lys
20 25 30

Gly Gly Cys Ser Leu Ile Ile Leu Pro Lys Thr Ser Ala Arg Lys Tyr
35 40 45

Gly Ile Phe Gln Tyr Pro Tyr Leu Ile Ile Leu Leu Leu Ala Arg Arg
50 55 60

Val Met Arg Lys Asp Glu His Leu Thr Ser Arg Gly Ser Leu Ile Leu
65 70 75 80

Arg Ser Arg Leu Leu Ile Lys Ser Gly Ile Trp Val Pro Ala Ala Thr
85 90 95

```

-continued

Ser Arg Lys Arg Lys Glu Gln Ile Ser Arg Ile Lys Asn Thr Ala Val
 100 105 110

Lys Lys Ala Thr Tyr Trp Ser Val Glu Leu Leu Asp Leu Arg Asp Asn
 115 120 125

Gly Val Thr Cys Leu Leu Leu Val Pro Cys Trp Arg Ala Gly Val Trp
 130 135 140

Pro Leu Gln Ser Ser Thr Leu Arg Leu Ser Glu Ile Ser Ala Gly Trp
 145 150 155 160

Leu Asn Phe Lys Ala Gly Lys Cys Arg Ser Val Ile Gly Arg Leu Met
 165 170 175

Tyr Thr Ser Leu Trp Gly His Leu Gly Val Val Lys Ala Asn Gly Leu
 180 185 190

Leu Ile Leu Gln Thr Arg Lys Pro His Thr Gly Asn His Leu Glu Thr
 195 200 205

Ser Gly Gly Met Val Thr Met Val Lys Lys Trp Leu Leu Leu Met Thr
 210 215 220

Phe Met Ala Gly Cys Pro Gly Met Ile Tyr Asp Cys Val Ile Asp Ile
 225 230 235 240

His Leu Arg Leu Lys Val Glu Leu Tyr Leu Phe Trp Pro Ala Val Phe
 245 250 255

Leu Pro Ala Ile Arg Pro Arg Trp Asn Gly Thr Pro Gln Leu Leu Ser
 260 265 270

Gln Leu Lys Leu Phe Ile Gly Gly Leu Leu Pro Trp Tyr Phe Gly Arg
 275 280 285

Met Leu Gln Asn Asn Pro Arg Arg Lys Gly Ala Ser Ser Ser Pro Phe
 290 295 300

Pro Pro His Ala Leu Asn Phe His Met Lys Ile Thr Glu Ser Phe Leu
 305 310 315 320

Ser Leu Arg Asn Gly Phe Tyr Tyr Ser Leu Arg Val Lys Trp Gly Val
 325 330 335

Phe Lys Ile Lys Phe Ser Glu Leu Tyr Ile His Gly Tyr Thr Asp Ile
 340 345 350

Val Phe Leu Val Val Tyr Thr Val Phe Glu Arg Ser Ala Glu Ala Tyr
 355 360 365

Val Val Tyr Ile Ser Ser Ser Leu Ser Gln Pro Gln Leu Val Ser Phe
 370 375 380

Val Val Trp Leu Glu Val Ile Asn Ser Glu Ile Asp Arg Phe Gly Gly
 385 390 395 400

Lys Val Pro Gly Val Val Gly Glu Gly Leu Gly Tyr Gly Met Ala Gly
 405 410 415

Gly Val Val Tyr Ile Gly Val Ile Gly Glu Gly Cys Gly Leu Cys Tyr
 420 425 430

Lys Val Ile Ile Asn Asn Ser Thr Gly Ala His Ser Pro Val Thr Leu
 435 440 445

Gly Asp Arg Gly Ala Gly Pro Glu Phe Asn Leu Asn Leu Ser Tyr Ser
 450 455 460

Val Val Phe Lys Gly His Arg Ala Gly Val Pro Pro Ser Trp Gly Lys
 465 470 475 480

Lys Val Ile Asn Ile Glu Ser His His Val His Arg Pro Gly Gly Arg
 485 490 495

Ser Asp Cys Gly Ser Leu Asp Ser Ile Ser Glu Gly Ala Gly Glu Ala
 500 505 510

Gly Val Glu Asp Ala Ile Phe Pro Ser Pro Ala Val Thr Val Ala Gly

-continued

Arg Asn Phe Arg Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met
 165 170 175

Gln Lys Arg Asp Trp Lys Thr Asn Val His Val Ile Val Gly Pro Pro
 180 185 190

Gly Cys Gly Lys Ser Lys Trp Ala Ala Asn Phe Ala Asp Pro Glu Thr
 195 200 205

Thr Tyr Trp Lys Pro Pro Arg Asn Lys Trp Trp Asp Gly Tyr His Gly
 210 215 220

Glu Glu Val Val Val Ile Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp
 225 230 235 240

Asp Leu Leu Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Glu Thr Lys
 245 250 255

Gly Gly Thr Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn
 260 265 270

Gln Thr Pro Leu Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu
 275 280 285

Ala Leu Tyr Arg Arg Ile Thr Ser Leu Val Phe Trp Lys Asn Ala Thr
 290 295 300

Glu Gln Ser Thr Glu Glu Gly Gly Gln Phe Val Thr Leu Ser Pro Pro
 305 310 315 320

Cys Pro Glu Phe Pro Tyr Glu Ile Asn Tyr Val Phe Phe Ile Thr Ser
 325 330 335

Trp Phe Leu Leu Phe Ile Lys Gly Val Gly Gly Leu Ile Val His Thr
 340 345 350

Trp Leu His Gly Tyr Cys Ile Pro Gly Arg Ile Tyr Cys Phe Arg Thr
 355 360 365

Gln Cys Arg Gly Leu Arg Gly Leu His Phe Gln Gln Phe Val Val Ser
 370 375 380

Ala Thr Ala Gly Phe Phe Cys Cys Leu Val Gly Ser Asn Gln Asn Leu
 385 390 395 400

Gly Gln Val Trp Gly Ser Thr Gly Ser Gly Arg Arg Arg Ala Gly Leu
 405 410 415

Trp Tyr Gly Gly Arg Ser Ser Leu His Arg Gly His Arg Gly Leu Trp
 420 425 430

Pro Leu Leu Gln Ser Tyr His Leu Lys Gln His Trp Ser Pro Leu Pro
 435 440 445

Cys His Pro Gly Ser Gly Ser Arg Ala Arg Ile Gln Pro Pro Phe Leu
 450 455 460

Phe Cys Ser Ile Gln Arg Ala Gln Ser Gly Gly Leu Thr Pro Leu Leu
 465 470 475 480

Gly Glu Glu Ser His Ile Ser Ser Cys Pro Pro Arg Arg Ala Phe
 485 490 495

Leu Trp Phe Ala Gln Tyr Ile Arg Arg Cys Gly Arg Gly Gly Cys Arg
 500 505 510

Cys His Phe Ser Phe Ser Ser Gly Asn Gly Gly Gly Gly Arg Ala
 515 520 525

Arg Gly Gly Gly Gly Ser Gly Gln Asp Gly Cys Gly Gly Gly Val
 530 535 540

Phe Phe Phe Gly Asn Ala Ser Leu Asp Thr Ser Tyr Leu Lys Thr Lys
 545 550 555 560

Glu Val Arg Cys Lys Tyr
 565

-continued

```

<210> SEQ ID NO 19
<211> LENGTH: 1767
<212> TYPE: DNA
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 19
aatacttaca ggcgacttct ttcgttttca gatatgacgt atccaaggag gcgttaccga      60
agaagaagac accgccccg cagccatctt ggccagatcc tccgccgccc cccctggctc      120
gtccaccccc gccaccgtta ccgctggaga aggaaaaatg gcactctcaa caccgcctc      180
tcccgcacct tcggatatac tgtcaagcga accacagtca gaacgccctc ctgggcggtg      240
gacatgatga gattcaatat taatgacttt ctccccccag gagggggggtc aaacccccgc      300
tctgtgccct ttgaatacta cagaataaga aaggttaagg ttgaattctg gccctgctcc      360
ccgatcacc cagggtgacag gggagtgggc tccagtgctg ttattttaga tgataacttt      420
gtaacaaagg ccacagccct cacctatgac ccctatgtaa actactctc cggccatacc      480
ataaccagc cttctccta ccaactcccg tactttacc ccaaacctgt cctagatttc      540
actattgatt acttccaacc aaacaacaaa agaaaccagc tgtggctgag actacaaact      600
gctggaaatg tagaccacgt aggcctcgcc actgcgttcg aaaacagtat atacgaccag      660
gaatacaata tccgtgtaac catgtatgta caattcagag aatttaattt taaagacccc      720
ccacttaacc cttaatgaat aataaaaacc attacgaagt gataaaaag actcagtaat      780
ttattcata tggaaattca gggcatgggg gggaaagggt gacgaactgg cccccttct      840
ccgtggattg ttctgtagca ttcttccaaa ataccaagga agtaatctc cgataaagag      900
cttctacagc tgggacagca gttgaggagt accattccaa cggggctga ttgctggtaa      960
tcagaatact gggggccaaa aaaggtacag ttccacctt agtctctaca gtcaatggat      1020
atcgatcaca cagtctcagt agatcatccc agggcagcca gccataaaag tcatcaataa      1080
caaccacttc ttcacatgg taaccatccc accacttggt tctaggtggt tccagtatg      1140
tggtttccgg gtctgcaaaa ttagcagccc atttgctttt accacacca ggtggcccca      1200
caatgacgtg tacattagtc ttccaatcac gttctgcat tttcccgctc actttcaaaa      1260
gttcagccag cccgcggaaa tttctgacaa acggttacagg gtgctgctct gcaacggta      1320
ccagactccc gctctccaac aaggactca cagcagtaga caggctactc cgttgctcct      1380
gagatctagg agctccacac tccatcagta agttgccttc tttactgcag tattctttat      1440
tctgctgac tgctcttctc gctttctcga tgggcagcg ggcacccaaa taccactca      1500
ctttattaaa agtctgcttc ttcacaaaat tagcgaaccc ctggagggtga ggtgttcgtc      1560
cttctcatt acctctctcg ccaacaataa aataatcaaa tagggatatt ggaagatccc      1620
gtattttctt gcgctcgtct tcggaagat tattcagagt gaacacccac cttttatggg      1680
gttggggtec gcttcttcca ttctcttgc tgggeatggt gctgctgagg tgetgcccag      1740
gtgctgccgc tgccgaagtg cgctggt      1767

```

```

<210> SEQ ID NO 20
<211> LENGTH: 567
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 20

```

```

Gly Ala Cys Lys Pro Leu Pro Leu Val Glu Ala Ala Gly Cys Cys Cys
1           5           10           15

```

-continued

Ala Trp Cys Ser Ser His Phe Phe Arg Val Gly Val Gly Tyr Phe Thr
20 25 30

Pro Thr Glu Ser Tyr Asp Lys Arg Leu Arg Ala Cys Ser Phe Val Pro
35 40 45

Asp Glu Leu Ile Gly Ile Gln Asn Asn Gln Gln Arg Pro Pro Tyr His
50 55 60

Pro Leu Val Phe Val Glu Gly Gly Pro Thr Arg Asn Gln Ser Ser Ala
65 70 75 80

Ser Lys Tyr Leu Ser Thr Thr Asn Pro His Gly Ser Gly Cys Arg Ser
85 90 95

Leu Ser Leu Phe Leu Asp Ala Ser Tyr Leu Ile Ser Cys Tyr Leu Leu
100 105 110

Cys Ser Val Ser Pro Thr His Leu Glu Ile Glu Pro Val Val Ser His
115 120 125

Gly Thr Gln Gln Ser Tyr Arg Thr Pro Ser Arg Ser Asp Pro Ser Arg
130 135 140

Gln Leu Ala Ala Gly Gln Leu Thr Gln Phe Asn Gly Arg Ala Pro Gln
145 150 155 160

Val Lys Ser Leu Ser Arg Ser Phe Ala Ser Ala His Asn Ser Ser His
165 170 175

Val Arg Gln Pro Ala Val Gln Thr His Tyr Phe Cys Ile Pro Gln Asn
180 185 190

Gln Leu Gly Pro Phe Trp Met Ser Ser Val Val Phe Cys Thr Thr Pro
195 200 205

His Asn Gly His His Leu Leu Pro Gln Gln His Ser Lys His Ser Ala
210 215 220

Ala Arg Pro His Asp Val Ser Val Thr His Asp Ile Asp Met Ser Gln
225 230 235 240

Leu Ser Leu His Phe Gln Val Lys Lys Pro Gly Cys Tyr Glu Ser Trp
245 250 255

Cys Asp Ser Gly Thr Pro Ile Thr Ser Arg Leu Gln Gln Gly Leu Gln
260 265 270

Leu Leu Glu Lys Asp Ser Ser Lys Arg Pro Ile Lys Ser Ser His Leu
275 280 285

Val Ile Trp Pro Pro Leu Pro Gly Thr Arg Gly Lys Gly Gly Met Gly
290 295 300

Gln Ile Glu Met His Phe Leu Asn Ser Leu Arg Lys Lys Thr Ile Thr
305 310 315 320

Lys Ile Ile Pro Asn Leu Pro Pro Asp Lys Phe Asn Phe Glu Arg Phe
325 330 335

Gln Val Tyr Met Thr Val Arg Ile Asn Tyr Glu Gln Asp Tyr Ile Ser
340 345 350

Asn Glu Phe Ala Thr Gly Leu Gly Val His Asp Val Asn Gly Ala Thr
355 360 365

Gln Leu Arg Leu Trp Leu Gln Asn Arg Lys Asn Asn Pro Gln Phe Tyr
370 375 380

Asp Ile Thr Phe Asp Leu Val Pro Lys Pro Thr Phe Tyr Arg Ser His
385 390 395 400

Tyr Ser Phe Pro Gln Thr Ile Thr His Arg Ser Ser Tyr Asn Val Tyr
405 410 415

Pro Asp Tyr Thr Leu Ala Thr Ala Lys Thr Val Pro Asn Asp Asp Leu
420 425 430

Ile Val Ala Ser Ser Gly Val Gly Arg Asp Gly Gln Thr Ile Pro Ser

-continued

Ser Arg Tyr Gly Asn Val Thr Ser Val Leu Pro Pro Val Thr Gly Lys
245 250 255

Lys Ala Arg Leu Ile Arg Ile Val Leu Leu Val Gly Asn Ser His Tyr
260 265 270

Glu Glu Val Ala Thr Gly Ala Thr Ser Ala Arg Arg Leu Ile Val Glu
275 280 285

Lys Thr Asn Gln Phe Phe Ala Val Ser Cys Asp Val Ser Ser Pro Pro
290 295 300

Trp Asn Thr Val Arg Glu Gly Gly His Gly Ser Asn Gly Tyr Ser Ile
305 310 315 320

Phe Gln Thr Lys Lys Ile Val Glu Tyr His Asn Lys Asn Asn Met Leu
325 330 335

Pro Thr Pro Pro Arg Phe Ile Arg Gln Ile Thr Cys Val His Asn Cys
340 345 350

Pro Tyr Gln Ile Gly Pro Arg Ile Tyr Gln Lys Arg Val Cys His Arg
355 360 365

Pro Arg Arg Pro Arg Cys Lys Trp Cys Asn Thr Thr Glu Ala Val Ala
370 375 380

Pro Lys Lys Gln Gln Lys Thr Pro Leu Leu Tyr His Phe Arg Pro Cys
385 390 395 400

Thr Gln Pro Tyr Leu Val Pro Leu Pro Leu Leu Leu Ala Pro Asn His
405 410 415

Tyr Pro Pro Leu Leu Leu Lys Cys Leu Pro Leu His Pro Ser His Gly
420 425 430

Lys Asn Cys Leu Arg Phe Tyr Cys Cys Gln Leu Gly Ser Gly Gln Gly
435 440 445

Pro His Asp Pro Leu Leu Ala Leu Ile Gly Gly Lys Lys Asn Gln Leu
450 455 460

Ile Leu Ala Cys Leu Pro Pro Lys Val Gly Arg Arg Pro Ser Ser Leu
465 470 475 480

Tyr Gln Ile Glu Asp His Gly Gly Gly Leu Leu Ala Asn Gln Ser His
485 490 495

Asn Ala Gln Cys Tyr Ile Arg Leu His Pro Leu Pro Pro His Gln Leu
500 505 510

His Trp Lys Glu Lys Glu Leu Pro Leu Pro Pro Pro Pro Arg Ala
515 520 525

Leu Pro Pro Pro Pro Pro Asp Pro Trp Ser Pro Gln Pro Pro Pro Thr
530 535 540

Lys Lys Lys Pro Leu Ala Glu Lys Ser Val Asp Tyr Arg Phe Val Phe
545 550 555 560

Ser Thr Arg Gln Leu Tyr
565

<210> SEQ ID NO 22

<211> LENGTH: 569

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 22

Leu Ala Ser Arg Cys Arg Cys Cys Arg Pro Leu Val Glu Ala Ala Val
1 5 10 15

His Gly Ala Leu Leu Ile Ser Ser Ala Ser Gly Leu Gly Met Phe Pro
20 25 30

Pro His Glu Ser Gln Ile Ile Arg Gly Phe Val Leu Ala Leu Phe Tyr
35 40 45

-continued

Pro Ile Lys Trp Tyr Gly Lys Ile Ile Lys Asn Asn Ala Leu Leu Thr
 50 55 60

Ile Leu Phe Ser Ser Cys Arg Val Glu Leu Pro Glu Ser Ile Lys His
 65 70 75 80

Leu Leu Leu Ser Lys Ile Phe His Leu Pro Ile Gln Thr Gly Ala Ala
 85 90 95

Val Asp Leu Phe Arg Phe Ser Cys Ile Leu Leu Ile Phe Phe Val Ala
 100 105 110

Thr Phe Phe Ala Val Gln His Leu Thr Ser Ser Arg Ser Arg Leu Ser
 115 120 125

Leu Pro Thr Val Gln Arg Ser Ser His Thr Gly Gln Gln Leu Ala Pro
 130 135 140

Thr Gln His Gly Asn Cys Leu Leu Val Arg Tyr Arg Lys Asp Ser Ile
 145 150 155 160

Glu Ala Pro Gln Ser Phe Lys Gln Phe His Ala Pro Phe His Leu Leu
 165 170 175

Thr Ile Pro Leu Ser Ile Tyr Val Asp Asn His Pro Trp Arg Pro Thr
 180 185 190

Thr Phe Ala Phe Pro Ser Ser Ile Lys Cys Val Arg Phe Gly Cys Val
 195 200 205

Pro Phe Trp Arg Ser Val Leu Pro Pro Ile Thr Val Met Thr Phe Phe
 210 215 220

His Asn Asn Asn Ile Val Lys Ile Ala Pro Gln Gly Pro Ile Ile Gln
 225 230 235 240

Ser Gln Thr Ser Ser Ile Trp Gln Ser Tyr Leu Ser Phe Thr Ser Ser
 245 250 255

Tyr Arg Lys Gln Gly Ala Thr Asn Gln Asn Gly Ala Ile Leu Gly Arg
 260 265 270

Gln Phe Pro Val Gly Ser Ser Asp Trp Ser Tyr Phe Ser Lys Ile Pro
 275 280 285

Pro Asn Ser Gly Gln Tyr Lys Pro Leu Ile Ser Cys Phe Leu Gly Arg
 290 295 300

Leu Phe Pro Ala Leu Glu Asp Gly Lys Gly Gly Trp Ala Arg Phe Lys
 305 310 315 320

Trp Ile Phe Trp Ile Val Ser Asp Lys Lys Asp Ser Arg Leu Pro Lys
 325 330 335

Glu Asn Leu Thr Leu His Pro Thr Lys Leu Ile Leu Asn Glu Ser Asn
 340 345 350

Tyr Met Cys Pro Val Ser Ile Thr Asn Arg Thr Thr Tyr Val Thr Lys
 355 360 365

Ser Arg Leu Ala Ser Ala Thr Thr Met Glu Leu Leu Lys Tyr Asp Gly
 370 375 380

Cys Ser Thr Glu Lys Thr Thr Gln Asn Ser Thr Ile Leu Leu Ser Ile
 385 390 395 400

Ser Leu Asn Pro Pro Leu Thr Gly Pro Thr Thr Pro Ser Pro Ser Pro
 405 410 415

Pro Ile Ala Pro Pro Thr Thr Met Pro Thr Met Pro Ser Pro Gln Pro
 420 425 430

Arg Gln Leu Thr Ile Met Phe Leu Leu Val Pro Ala Trp Glu Gly Thr
 435 440 445

Val Arg Pro Ser Arg Pro Ala Pro Gly Ser Asn Leu Arg Leu Arg Glu
 450 455 460

-continued

Glu Thr Thr Asn Leu Pro Cys Leu Ala Pro Thr Gln Gly Gly Glu Gln
 465 470 475 480

Pro Phe Phe Thr Met Leu Ile Ser Asp Thr Trp Arg Gly Pro Pro Arg
 485 490 495

Glu Ser Gln Pro Glu Ser Ser Leu Ile Asp Ser Pro Ala Pro Ser Ala
 500 505 510

Pro Thr Ser Ser Ala Met Lys Gly Glu Gly Ala Thr Val Thr Ala Pro
 515 520 525

Thr Ser Ser Gly Pro Ala Ala Ala Ser Ser Arg Ala Leu Ile Ala Ala
 530 535 540

Pro Ala Thr Asp Glu Glu Glu Thr Val Gly Gly Gln Ile Arg Ile Gln
 545 550 555 560

Phe Arg Phe Phe His Ala Thr Leu Ile
 565

<210> SEQ ID NO 23
 <211> LENGTH: 945
 <212> TYPE: DNA
 <213> ORGANISM: Type B PWD circovirus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(942)

<400> SEQUENCE: 23

atg ccc agc aag aag aat gga aga agc gga ccc caa ccc cat aaa agg 48
 Met Pro Ser Lys Lys Asn Gly Arg Ser Gly Pro Gln Pro His Lys Arg
 1 5 10 15

tgg gtg ttc act ctg aat aat cct tcc gaa gac gag cgc aag aaa ata 96
 Trp Val Phe Thr Leu Asn Asn Pro Ser Glu Asp Glu Arg Lys Lys Ile
 20 25 30

cgg gat ctt cca ata tcc cta ttt gat tat ttt att gtt ggc gag gag 144
 Arg Asp Leu Pro Ile Ser Leu Phe Asp Tyr Phe Ile Val Gly Glu Glu
 35 40 45

ggt aat gag gaa gga cga aca cct cac ctc cag ggg ttc gct aat ttt 192
 Gly Asn Glu Glu Gly Arg Thr Pro His Leu Gln Gly Phe Ala Asn Phe
 50 55 60

gtg aag aag cag act ttt aat aaa gtg aag tgg tat ttg ggt gcc cgc 240
 Val Lys Lys Gln Thr Phe Asn Lys Val Lys Trp Tyr Leu Gly Ala Arg
 65 70 75 80

tgc cac atc gag aaa gcg aaa gga aca gat cag cag aat aaa gaa tac 288
 Cys His Ile Glu Lys Ala Lys Gly Thr Asp Gln Gln Asn Lys Glu Tyr
 85 90 95

tgc agt aaa gaa ggc aac tta ctg atg gag tgt gga gct cct aga tct 336
 Cys Ser Lys Glu Gly Asn Leu Leu Met Glu Cys Gly Ala Pro Arg Ser
 100 105 110

cag gga caa cgg agt gac ctg tct act gct gtg agt acc ttg ttg gag 384
 Gln Gly Gln Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu
 115 120 125

agc ggg agt ctg gtg acc gtt gca gag cag cac cct gta acg ttt gtc 432
 Ser Gly Ser Leu Val Thr Val Ala Glu Gln His Pro Val Thr Phe Val
 130 135 140

aga aat ttc cgc ggg ctg gct gaa ctt ttg aaa gtg agc ggg aaa atg 480
 Arg Asn Phe Arg Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met
 145 150 155 160

cag aag cgt gat tgg aag act aat gta cac gtc att gtg ggg cca cct 528
 Gln Lys Arg Asp Trp Lys Thr Asn Val His Val Ile Val Gly Pro Pro
 165 170 175

ggg tgt ggt aaa agc aaa tgg gct gct aat ttt gca gac ccg gaa acc 576
 Gly Cys Gly Lys Ser Lys Trp Ala Ala Asn Phe Ala Asp Pro Glu Thr
 180 185 190

-continued

```

aca tac tgg aaa cca cct aga aac aag tgg tgg gat ggt tac cat ggt      624
Thr Tyr Trp Lys Pro Pro Arg Asn Lys Trp Trp Asp Gly Tyr His Gly
      195                200                205

gaa gaa gtg gtt gtt att gat gac ttt tat ggc tgg ctg ccc tgg gat      672
Glu Glu Val Val Val Ile Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp
      210                215                220

gat cta ctg aga ctg tgt gat cga tat cca ttg act gta gag act aaa      720
Asp Leu Leu Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Glu Thr Lys
      225                230                235                240

ggg gga act gta cct ttt ttg gcc cgc agt att ctg att acc agc aat      768
Gly Gly Thr Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn
      245                250                255

cag acc ccg ttg gaa tgg tac tcc tca act gct gtc cca gct gta gaa      816
Gln Thr Pro Leu Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu
      260                265                270

gct ctt tat cgg agg att act tcc ttg gta ttt tgg aag aat gct aca      864
Ala Leu Tyr Arg Arg Ile Thr Ser Leu Val Phe Trp Lys Asn Ala Thr
      275                280                285

gaa caa tcc acg gag gaa ggg ggc cag ttc gtc acc ctt tcc ccc cca      912
Glu Gln Ser Thr Glu Glu Gly Gly Gln Phe Val Thr Leu Ser Pro Pro
      290                295                300

tgc cct gaa ttt cca tat gaa ata aat tac tga      945
Cys Pro Glu Phe Pro Tyr Glu Ile Asn Tyr
      305                310

```

<210> SEQ ID NO 24

<211> LENGTH: 314

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 24

```

Met Pro Ser Lys Lys Asn Gly Arg Ser Gly Pro Gln Pro His Lys Arg
1      5      10      15

Trp Val Phe Thr Leu Asn Asn Pro Ser Glu Asp Glu Arg Lys Lys Ile
      20      25      30

Arg Asp Leu Pro Ile Ser Leu Phe Asp Tyr Phe Ile Val Gly Glu Glu
      35      40      45

Gly Asn Glu Glu Gly Arg Thr Pro His Leu Gln Gly Phe Ala Asn Phe
      50      55      60

Val Lys Lys Gln Thr Phe Asn Lys Val Lys Trp Tyr Leu Gly Ala Arg
      65      70      75      80

Cys His Ile Glu Lys Ala Lys Gly Thr Asp Gln Gln Asn Lys Glu Tyr
      85      90      95

Cys Ser Lys Glu Gly Asn Leu Leu Met Glu Cys Gly Ala Pro Arg Ser
      100     105     110

Gln Gly Gln Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu
      115     120     125

Ser Gly Ser Leu Val Thr Val Ala Glu Gln His Pro Val Thr Phe Val
      130     135     140

Arg Asn Phe Arg Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met
      145     150     155     160

Gln Lys Arg Asp Trp Lys Thr Asn Val His Val Ile Val Gly Pro Pro
      165     170     175

Gly Cys Gly Lys Ser Lys Trp Ala Ala Asn Phe Ala Asp Pro Glu Thr
      180     185     190

Thr Tyr Trp Lys Pro Pro Arg Asn Lys Trp Trp Asp Gly Tyr His Gly
      195     200     205

```

-continued

Glu Glu Val Val Val Ile Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp
 210 215 220
 Asp Leu Leu Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Glu Thr Lys
 225 230 235 240
 Gly Gly Thr Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn
 245 250 255
 Gln Thr Pro Leu Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu
 260 265 270
 Ala Leu Tyr Arg Arg Ile Thr Ser Leu Val Phe Trp Lys Asn Ala Thr
 275 280 285
 Glu Gln Ser Thr Glu Glu Gly Gly Gln Phe Val Thr Leu Ser Pro Pro
 290 295 300
 Cys Pro Glu Phe Pro Tyr Glu Ile Asn Tyr
 305 310

<210> SEQ ID NO 25
 <211> LENGTH: 702
 <212> TYPE: DNA
 <213> ORGANISM: Type B PWD circovirus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(699)

<400> SEQUENCE: 25

atg acg tat cca agg agg cgt tac cga aga aga cac cgc ccc cgc 48
 Met Thr Tyr Pro Arg Arg Arg Tyr Arg Arg Arg Arg His Arg Pro Arg
 1 5 10 15
 agc cat ctt ggc cag atc ctc cgc cgc cgc ccc tgg ctc gtc cac ccc 96
 Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro Trp Leu Val His Pro
 20 25 30
 cgc cac cgt tac cgc tgg aga agg aaa aat ggc atc ttc aac acc cgc 144
 Arg His Arg Tyr Arg Trp Arg Arg Lys Asn Gly Ile Phe Asn Thr Arg
 35 40 45
 ctc tcc cgc acc ttc gga tat act gtc aag cga acc aca gtc aga acg 192
 Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Arg Thr Thr Val Arg Thr
 50 55 60
 ccc tcc tgg gcg gtg gac atg atg aga ttc aat att aat gac ttt ctt 240
 Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asn Asp Phe Leu
 65 70 75 80
 ccc cca gga ggg ggg tca aac ccc cgc tct gtg ccc ttt gaa tac tac 288
 Pro Pro Gly Gly Gly Ser Asn Pro Arg Ser Val Pro Phe Glu Tyr Tyr
 85 90 95
 aga ata aga aag gtt aag gtt gaa ttc tgg ccc tgc tcc ccg atc acc 336
 Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr
 100 105 110
 cag ggt gac agg gga gtg ggc tcc agt gct gtt att tta gat gat aac 384
 Gln Gly Asp Arg Gly Val Gly Ser Ser Ala Val Ile Leu Asp Asp Asn
 115 120 125
 ttt gta aca aag gcc aca gcc ctc acc tat gac ccc tat gta aac tac 432
 Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr
 130 135 140
 tcc tcc cgc cat acc ata acc cag ccc ttc tcc tac cac tcc cgg tac 480
 Ser Ser Arg His Thr Ile Thr Gln Pro Phe Ser Tyr His Ser Arg Tyr
 145 150 155 160
 ttt acc ccc aaa cct gtc cta gat ttc act att gat tac ttc caa cca 528
 Phe Thr Pro Lys Pro Val Leu Asp Phe Thr Ile Asp Tyr Phe Gln Pro
 165 170 175
 aac aac aaa aga aac cag ctg tgg ctg aga cta caa act gct gga aat 576
 Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ala Gly Asn

-continued

180	185	190	
gta gac cac gta ggc ctc ggc act gcg ttc gaa aac agt ata tac gac			624
Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Ile Tyr Asp			
195	200	205	
cag gaa tac aat atc cgt gta acc atg tat gta caa ttc aga gaa ttt			672
Gln Glu Tyr Asn Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe			
210	215	220	
aat ttt aaa gac ccc cca ctt aac cct taa			702
Asn Phe Lys Asp Pro Pro Leu Asn Pro			
225	230		

<210> SEQ ID NO 26
 <211> LENGTH: 233
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 26

Met Thr Tyr Pro Arg Arg Arg Tyr Arg Arg Arg His Arg Pro Arg			
1	5	10	15
Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro Trp Leu Val His Pro			
20	25	30	
Arg His Arg Tyr Arg Trp Arg Arg Lys Asn Gly Ile Phe Asn Thr Arg			
35	40	45	
Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Arg Thr Thr Val Arg Thr			
50	55	60	
Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asn Asp Phe Leu			
65	70	75	80
Pro Pro Gly Gly Gly Ser Asn Pro Arg Ser Val Pro Phe Glu Tyr Tyr			
85	90	95	
Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr			
100	105	110	
Gln Gly Asp Arg Gly Val Gly Ser Ala Val Ile Leu Asp Asp Asn			
115	120	125	
Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr			
130	135	140	
Ser Ser Arg His Thr Ile Thr Gln Pro Phe Ser Tyr His Ser Arg Tyr			
145	150	155	160
Phe Thr Pro Lys Pro Val Leu Asp Phe Thr Ile Asp Tyr Phe Gln Pro			
165	170	175	
Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ala Gly Asn			
180	185	190	
Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Ile Tyr Asp			
195	200	205	
Gln Glu Tyr Asn Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe			
210	215	220	
Asn Phe Lys Asp Pro Pro Leu Asn Pro			
225	230		

<210> SEQ ID NO 27
 <211> LENGTH: 315
 <212> TYPE: DNA
 <213> ORGANISM: Type B PWD circovirus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(312)

<400> SEQUENCE: 27

atg gta acc atc cca cca ctt gtt tct agg tgg ttt cca gta tgt ggt	48
---	----

-continued

```

Met Val Thr Ile Pro Pro Leu Val Ser Arg Trp Phe Pro Val Cys Gly
1          5          10          15
ttc cgg gtc tgc aaa att agc agc cca ttt gct ttt acc aca ccc agg      96
Phe Arg Val Cys Lys Ile Ser Ser Pro Phe Ala Phe Thr Thr Pro Arg
          20          25          30
tgg ccc cac aat gac gtg tac att agt ctt cca atc acg ctt ctg cat      144
Trp Pro His Asn Asp Val Tyr Ile Ser Leu Pro Ile Thr Leu Leu His
          35          40          45
ttt ccc gct cac ttt caa aag ttc agc cag ccc gcg gaa att tct gac      192
Phe Pro Ala His Phe Gln Lys Phe Ser Gln Pro Ala Glu Ile Ser Asp
          50          55          60
aaa cgt tac agg gtg ctg ctc tgc aac ggt cac cag act ccc gct ctc      240
Lys Arg Tyr Arg Val Leu Leu Cys Asn Gly His Gln Thr Pro Ala Leu
65          70          75          80
caa caa ggt act cac agc agt aga cag gtc act ccg ttg tcc ctg aga      288
Gln Gln Gly Thr His Ser Ser Arg Gln Val Thr Pro Leu Ser Leu Arg
          85          90          95
tct agg agc tcc aca ctc cat cag taa      315
Ser Arg Ser Ser Thr Leu His Gln
          100

```

```

<210> SEQ ID NO 28
<211> LENGTH: 104
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

```

<400> SEQUENCE: 28

```

```

Met Val Thr Ile Pro Pro Leu Val Ser Arg Trp Phe Pro Val Cys Gly
1          5          10          15
Phe Arg Val Cys Lys Ile Ser Ser Pro Phe Ala Phe Thr Thr Pro Arg
          20          25          30
Trp Pro His Asn Asp Val Tyr Ile Ser Leu Pro Ile Thr Leu Leu His
          35          40          45
Phe Pro Ala His Phe Gln Lys Phe Ser Gln Pro Ala Glu Ile Ser Asp
          50          55          60
Lys Arg Tyr Arg Val Leu Leu Cys Asn Gly His Gln Thr Pro Ala Leu
65          70          75          80
Gln Gln Gly Thr His Ser Ser Arg Gln Val Thr Pro Leu Ser Leu Arg
          85          90          95
Ser Arg Ser Ser Thr Leu His Gln
          100

```

```

<210> SEQ ID NO 29
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

```

<400> SEQUENCE: 29

```

```

Val Asp Met Met Arg Phe Asn Ile Asn Asp Phe Leu Pro Pro Gly
1          5          10          15

```

```

<210> SEQ ID NO 30
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

```

<400> SEQUENCE: 30

```

```

Gln Gly Asp Arg Gly Val Gly Ser Ser Ala Val Ile Leu Asp Asp
1          5          10          15

```

-continued

```

<210> SEQ ID NO 31
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

```

<400> SEQUENCE: 31

```

```

Gly Val Gly Ser Ser Ala Val Ile Leu Asp Asp Asn Phe Val Thr
1           5           10          15

```

```

<210> SEQ ID NO 32
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

```

<400> SEQUENCE: 32

```

```

Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Ile Tyr
1           5           10          15

```

```

<210> SEQ ID NO 33
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 33

```

```

tgtggcga

```

8

```

<210> SEQ ID NO 34
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 34

```

```

agtttcct

```

8

```

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 35

```

```

tcatttagag ggtctttcag

```

20

```

<210> SEQ ID NO 36
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 36

```

```

gtcaacct

```

8

```

<210> SEQ ID NO 37
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 37

```

```

gtggttgc

```

8

```

<210> SEQ ID NO 38
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 38

```

-continued

agcccagg	8
<210> SEQ ID NO 39 <211> LENGTH: 8 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 39	
ttggctgg	8
<210> SEQ ID NO 40 <211> LENGTH: 12 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 40	
tctagetctg gt	12
<210> SEQ ID NO 41 <211> LENGTH: 12 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 41	
atctcagctc gt	12
<210> SEQ ID NO 42 <211> LENGTH: 12 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 42	
tgtectectc tt	12
<210> SEQ ID NO 43 <211> LENGTH: 8 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 43	
tctctaga	8
<210> SEQ ID NO 44 <211> LENGTH: 8 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 44	
tgtaccaa	8
<210> SEQ ID NO 45 <211> LENGTH: 8 <212> TYPE: DNA <213> ORGANISM: Type A PWD circovirus <400> SEQUENCE: 45	
tccgtett	8
<210> SEQ ID NO 46 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 46	

-continued

gtgtgctcga cattggtgtg	20
<210> SEQ ID NO 47 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 47	
tggaatgtta acgagctgag	20
<210> SEQ ID NO 48 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 48	
ctcgcagcca tcttggaatg	20
<210> SEQ ID NO 49 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 49	
cgcgcgtaat acgactcact	20
<210> SEQ ID NO 50 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 50	
cctgtctact gctgtgagta ccttgt	26
<210> SEQ ID NO 51 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 51	
gcagtagaca ggtcactccg ttgtcc	26
<210> SEQ ID NO 52 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 52	
tggaatgtta actacctcaa	20
<210> SEQ ID NO 53 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Primer <400> SEQUENCE: 53	
ggcggcgcca tetgtaacgg ttt	23
<210> SEQ ID NO 54 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Primer	

-continued

<400> SEQUENCE: 54

gatggcgccg aaagacgggt atc

23

<210> SEQ ID NO 55

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 55

Asn	Val	Asn	Glu	Leu	Arg	Phe	Asn	Ile	Gly	Gln	Phe	Leu	Pro	Pro
1			5						10				15	

<210> SEQ ID NO 56

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 56

Thr	Ser	Asn	Gln	Arg	Gly	Val	Gly	Ser	Thr	Val	Val	Ile	Leu
1			5						10				

<210> SEQ ID NO 57

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 57

Arg	Gly	Val	Gly	Ser	Thr	Val	Val	Ile	Leu	Asp	Ala	Asn	Phe	Val
1			5						10				15	

<210> SEQ ID NO 58

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 58

Phe	Thr	Ile	Asp	Tyr	Phe	Gln	Pro	Asn	Asn	Lys	Arg	Asn	Gln	Leu
1			5						10				15	

<210> SEQ ID NO 59

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 59

Asp	Gln	Thr	Ile	Asp	Trp	Phe	Gln	Pro	Asn	Asn	Lys	Arg	Asn	Gln
1			5						10				15	

<210> SEQ ID NO 60

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 60

Asn	Val	Glu	His	Thr	Gly	Leu	Gly	Tyr	Ala	Leu	Gln	Asn	Ala	Thr
1			5						10				15	

<210> SEQ ID NO 61

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 61

His Arg Pro Arg Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro

-continued

1	5	10	15
---	---	----	----

<210> SEQ ID NO 62
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 62

Ser His Leu Gly	Gln Ile Leu Arg Arg	Arg Pro Trp Leu Val His	
1	5	10	15

<210> SEQ ID NO 63
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 63

Gln Ile Leu Arg	Arg Arg Pro Trp Leu Val His	Pro Arg His Arg	
1	5	10	15

<210> SEQ ID NO 64
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 64

Arg Arg Pro Trp	Leu Val His Pro Arg His	Arg Tyr Arg Trp Arg	
1	5	10	15

<210> SEQ ID NO 65
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 65

Leu Val His Pro	Arg His Arg Tyr Arg	Trp Arg Arg Lys Asn Gly	
1	5	10	15

<210> SEQ ID NO 66
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 66

Arg His Arg Tyr	Arg Trp Arg Arg Lys Asn Gly	Ile Phe Asn Thr	
1	5	10	15

<210> SEQ ID NO 67
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 67

Arg Trp Arg Arg	Lys Asn Gly Ile Phe Asn Thr	Arg Leu Ser Arg	
1	5	10	15

<210> SEQ ID NO 68
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 68

Lys Asn Gly Ile	Phe Asn Thr Arg Leu Ser	Arg Thr Phe Gly Tyr	
1	5	10	15

-continued

<210> SEQ ID NO 69
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 69

Phe Asn Thr Arg Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Arg
 1 5 10 15

<210> SEQ ID NO 70
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 70

Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Arg Thr Thr Val Arg
 1 5 10 15

<210> SEQ ID NO 71
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 71

Phe Gly Tyr Thr Val Lys Arg Thr Thr Val Arg Thr Pro Ser Trp
 1 5 10 15

<210> SEQ ID NO 72
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 72

Val Lys Arg Thr Thr Val Arg Thr Pro Ser Trp Ala Val Asp Met
 1 5 10 15

<210> SEQ ID NO 73
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 73

Thr Val Arg Thr Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn
 1 5 10 15

<210> SEQ ID NO 74
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 74

Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asn Asp Phe
 1 5 10 15

<210> SEQ ID NO 75
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 75

Arg Phe Asn Ile Asn Asp Phe Leu Pro Pro Gly Gly Gly Ser Asn
 1 5 10 15

<210> SEQ ID NO 76

-continued

```

<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 76

Asn Asp Phe Leu Pro Pro Gly Gly Gly Ser Asn Pro Arg Ser Val
1           5           10           15

<210> SEQ ID NO 77
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 77

Pro Pro Gly Gly Gly Ser Asn Pro Arg Ser Val Pro Phe Glu Tyr
1           5           10           15

<210> SEQ ID NO 78
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 78

Gly Ser Asn Pro Arg Ser Val Pro Phe Glu Tyr Tyr Arg Ile Arg
1           5           10           15

<210> SEQ ID NO 79
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 79

Arg Ser Val Pro Phe Glu Tyr Tyr Arg Ile Arg Lys Val Lys Val
1           5           10           15

<210> SEQ ID NO 80
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 80

Phe Glu Tyr Tyr Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro
1           5           10           15

<210> SEQ ID NO 81
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 81

Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile
1           5           10           15

<210> SEQ ID NO 82
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 82

Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr Gln Gly Asp
1           5           10           15

<210> SEQ ID NO 83
<211> LENGTH: 15
<212> TYPE: PRT

```

-continued

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 83

Phe	Trp	Pro	Cys	Ser	Pro	Ile	Thr	Gln	Gly	Asp	Arg	Gly	Val	Gly
1			5					10					15	

<210> SEQ ID NO 84

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 84

Thr	Arg	Pro	Arg	Ser	His	Leu	Gly	Asn	Ile	Leu	Arg	Arg	Arg	Pro
1			5					10						15

<210> SEQ ID NO 85

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 85

Ser	His	Leu	Gly	Asn	Ile	Leu	Arg	Arg	Arg	Pro	Tyr	Leu	Val	His
1			5					10						15

<210> SEQ ID NO 86

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 86

Asn	Ile	Leu	Arg	Arg	Arg	Pro	Tyr	Leu	Val	His	Pro	Ala	Phe	Arg
1			5					10						15

<210> SEQ ID NO 87

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 87

Arg	Arg	Pro	Tyr	Leu	Val	His	Pro	Ala	Phe	Arg	Asn	Arg	Tyr	Arg
1			5						10					15

<210> SEQ ID NO 88

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 88

Leu	Val	His	Pro	Ala	Phe	Arg	Asn	Arg	Tyr	Arg	Trp	Arg	Arg	Lys
1			5						10					15

<210> SEQ ID NO 89

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 89

Ala	Phe	Arg	Asn	Arg	Tyr	Arg	Trp	Arg	Arg	Lys	Thr	Gly	Ile	Phe
1			5						10					15

<210> SEQ ID NO 90

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

-continued

<400> SEQUENCE: 90

Arg	Tyr	Arg	Trp	Arg	Arg	Lys	Thr	Gly	Ile	Phe	Asn	Ser	Arg	Leu
1				5					10					15

<210> SEQ ID NO 91

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 91

Arg	Arg	Lys	Thr	Gly	Ile	Phe	Asn	Ser	Arg	Leu	Ser	Arg	Glu	Phe
1				5					10					15

<210> SEQ ID NO 92

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 92

Gly	Ile	Phe	Asn	Ser	Arg	Leu	Ser	Arg	Glu	Phe	Val	Leu	Thr	Ile
1			5						10					15

<210> SEQ ID NO 93

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 93

Ser	Arg	Leu	Ser	Arg	Glu	Phe	Val	Leu	Thr	Ile	Arg	Gly	Gly	His
1			5						10					15

<210> SEQ ID NO 94

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 94

Arg	Glu	Phe	Val	Leu	Thr	Ile	Arg	Gly	Gly	His	Ser	Gln	Pro	Ser
1				5					10					15

<210> SEQ ID NO 95

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 95

Leu	Thr	Ile	Arg	Gly	Gly	His	Ser	Gln	Pro	Ser	Trp	Asn	Val	Asn
1				5					10					15

<210> SEQ ID NO 96

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 96

Gly	Gly	His	Ser	Gln	Pro	Ser	Trp	Asn	Val	Asn	Glu	Leu	Arg	Phe
1			5						10					15

<210> SEQ ID NO 97

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 97

-continued

Gln Pro Ser Trp Asn Val Asn Glu Leu Arg Phe Asn Ile Gly Gln
1 5 10 15

<210> SEQ ID NO 98
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 98

Asn Val Asn Glu Leu Arg Phe Asn Ile Gly Gln Phe Leu Pro Pro
1 5 10 15

<210> SEQ ID NO 99
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 99

Leu Arg Phe Asn Ile Gly Gln Phe Leu Pro Pro Ser Gly Gly Thr
1 5 10 15

<210> SEQ ID NO 100
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 100

Ile Gly Gln Phe Leu Pro Pro Ser Gly Gly Thr Asn Pro Leu Pro
1 5 10 15

<210> SEQ ID NO 101
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 101

Leu Pro Pro Ser Gly Gly Thr Asn Pro Leu Pro Leu Pro Phe Gln
1 5 10 15

<210> SEQ ID NO 102
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 102

Gly Gly Thr Asn Pro Leu Pro Leu Pro Phe Gln Tyr Tyr Arg Ile
1 5 10 15

<210> SEQ ID NO 103
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 103

Pro Leu Pro Leu Pro Phe Gln Tyr Tyr Arg Ile Arg Lys Ala Lys
1 5 10 15

<210> SEQ ID NO 104
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 104

Pro Phe Gln Tyr Tyr Arg Ile Arg Lys Ala Lys Tyr Glu Phe Tyr
1 5 10 15

-continued

<210> SEQ ID NO 105
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus
 <400> SEQUENCE: 105

Tyr Arg Ile Arg Lys Ala Lys Tyr Glu Phe Tyr Pro Arg Asp Pro
 1 5 10 15

<210> SEQ ID NO 106
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 106

Lys Ala Lys Tyr Glu Phe Tyr Pro Arg Asp Pro Ile Thr Ser Asn
 1 5 10 15

<210> SEQ ID NO 107
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 107

Glu Phe Tyr Pro Arg Asp Pro Ile Thr Ser Asn Gln Arg Gly Val
 1 5 10 15

<210> SEQ ID NO 108
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 108

Arg Asp Pro Ile Thr Ser Asn Gln Arg Gly Val Gly Ser Thr Val
 1 5 10 15

<210> SEQ ID NO 109
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 109

Thr Ser Asn Gln Arg Gly Val Gly Ser Thr Val Val Ile Leu Asp
 1 5 10 15

<210> SEQ ID NO 110
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 110

Gly Val Gly Ser Ser Ala Val Ile Leu Asp Asp Asn Phe Val Thr
 1 5 10 15

<210> SEQ ID NO 111
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 111

Ser Ala Val Ile Leu Asp Asp Asn Phe Val Thr Lys Ala Thr Ala
 1 5 10 15

-continued

```

<210> SEQ ID NO 112
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 112

Leu Asp Asp Asn Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp
1             5             10             15

```

```

<210> SEQ ID NO 113
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 113

```

```

Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn
1             5             10             15

```

```

<210> SEQ ID NO 114
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 114

```

```

Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr Ser Ser Arg
1             5             10             15

```

```

<210> SEQ ID NO 115
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 115

```

```

Thr Tyr Asp Pro Tyr Val Asn Tyr Ser Ser Arg His Thr Ile Thr
1             5             10             15

```

```

<210> SEQ ID NO 116
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 116

```

```

Tyr Val Asn Tyr Ser Ser Arg His Thr Ile Thr Gln Pro Phe Ser
1             5             10             15

```

```

<210> SEQ ID NO 117
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 117

```

```

Ser Ser Arg His Thr Ile Thr Gln Pro Phe Ser Tyr His Ser Arg
1             5             10             15

```

```

<210> SEQ ID NO 118
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 118

```

```

Thr Ile Thr Gln Pro Phe Ser Tyr His Ser Arg Tyr Phe Thr Pro
1             5             10             15

```

```

<210> SEQ ID NO 119
<211> LENGTH: 15

```

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 119

Pro Phe Ser Tyr His Ser Arg Tyr Phe Thr Pro Lys Pro Val Leu
1           5           10           15

<210> SEQ ID NO 120
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 120

His Ser Arg Tyr Phe Thr Pro Lys Pro Val Leu Asp Phe Thr Ile
1           5           10           15

<210> SEQ ID NO 121
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 121

Phe Thr Pro Lys Pro Val Leu Asp Phe Thr Ile Asp Tyr Phe Gln
1           5           10           15

<210> SEQ ID NO 122
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 122

Pro Val Leu Asp Phe Thr Ile Asp Tyr Phe Gln Pro Asn Asn Lys
1           5           10           15

<210> SEQ ID NO 123
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 123

Phe Thr Ile Asp Tyr Phe Gln Pro Asn Asn Lys Arg Asn Gln Leu
1           5           10           15

<210> SEQ ID NO 124
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 124

Tyr Phe Gln Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu
1           5           10           15

<210> SEQ ID NO 125
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 125

Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ala Gly
1           5           10           15

<210> SEQ ID NO 126
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

```

-continued

<400> SEQUENCE: 126

Asn	Gln	Leu	Trp	Leu	Arg	Leu	Gln	Thr	Ala	Gly	Asn	Val	Asp	His
1				5					10					15

<210> SEQ ID NO 127

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 127

Leu	Arg	Leu	Gln	Thr	Ala	Gly	Asn	Val	Asp	His	Val	Gly	Leu	Gly
1				5					10					15

<210> SEQ ID NO 128

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 128

Thr	Ala	Gly	Asn	Val	Asp	His	Val	Gly	Leu	Gly	Thr	Ala	Phe	Glu
1				5					10					15

<210> SEQ ID NO 129

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 129

Gly	Leu	Gly	Thr	Ala	Phe	Glu	Asn	Ser	Ile	Tyr	Asp	Gln	Glu	Tyr
1				5					10					15

<210> SEQ ID NO 130

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 130

Ala	Phe	Glu	Asn	Ser	Ile	Tyr	Asp	Gln	Glu	Tyr	Asn	Ile	Arg	Val
1				5					10					15

<210> SEQ ID NO 131

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 131

Ser	Ile	Tyr	Asp	Gln	Glu	Tyr	Asn	Ile	Arg	Val	Thr	Met	Tyr	Val
1				5					10					15

<210> SEQ ID NO 132

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 132

Gln	Glu	Tyr	Asn	Ile	Arg	Val	Thr	Met	Tyr	Val	Gln	Phe	Arg	Glu
1				5					10					15

<210> SEQ ID NO 133

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 133

-continued

```
Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe Asn Phe Lys
1           5           10           15
```

```
<210> SEQ ID NO 134
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 134
```

```
Met Tyr Val Gln Phe Arg Glu Phe Asn Phe Lys Asp Pro Pro Leu
1           5           10           15
```

```
<210> SEQ ID NO 135
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type B PWD circovirus

<400> SEQUENCE: 135
```

```
Val Gln Phe Arg Glu Phe Asn Phe Lys Asp Pro Pro Leu Asn Pro
1           5           10           15
```

```
<210> SEQ ID NO 136
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 136
```

```
Arg Gly Val Gly Ser Thr Val Val Ile Leu Asp Ala Asn Phe Val
1           5           10           15
```

```
<210> SEQ ID NO 137
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 137
```

```
Ser Thr Val Val Ile Leu Asp Ala Asn Phe Val Thr Pro Ser Thr
1           5           10           15
```

```
<210> SEQ ID NO 138
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 138
```

```
Ile Leu Asp Ala Asn Phe Val Thr Pro Ser Thr Asn Leu Ala Tyr
1           5           10           15
```

```
<210> SEQ ID NO 139
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 139
```

```
Asn Phe Val Thr Pro Ser Thr Asn Leu Ala Tyr Asp Pro Tyr Ile
1           5           10           15
```

```
<210> SEQ ID NO 140
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 140
```

```
Pro Ser Thr Asn Leu Ala Tyr Asp Pro Tyr Ile Asn Tyr Ser Ser
```

-continued

1 5 10 15

<210> SEQ ID NO 141
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 141

Leu Ala Tyr Asp Pro Tyr Ile Asn Tyr Ser Ser Arg His Thr Ile
 1 5 10 15

<210> SEQ ID NO 142
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 142

Pro Tyr Ile Asn Tyr Ser Ser Arg His Thr Ile Arg Gln Pro Phe
 1 5 10 15

<210> SEQ ID NO 143
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 143

Tyr Ser Ser Arg His Thr Ile Arg Gln Pro Phe Thr Tyr His Ser
 1 5 10 15

<210> SEQ ID NO 144
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 144

His Thr Ile Arg Gln Pro Phe Thr Tyr His Ser Arg Tyr Phe Thr
 1 5 10 15

<210> SEQ ID NO 145
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 145

Gln Pro Phe Thr Tyr His Ser Arg Tyr Phe Thr Pro Lys Pro Glu
 1 5 10 15

<210> SEQ ID NO 146
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 146

Tyr His Ser Arg Tyr Phe Thr Pro Lys Pro Glu Leu Asp Gln Thr
 1 5 10 15

<210> SEQ ID NO 147
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 147

Tyr Phe Thr Pro Lys Pro Glu Leu Asp Gln Thr Ile Asp Trp Phe
 1 5 10 15

-continued

<210> SEQ ID NO 148
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 148

Lys Pro Glu Leu Asp Gln Thr Ile Asp Trp Phe Gln Pro Asn Asn
 1 5 10 15

<210> SEQ ID NO 149
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 149

Asp Gln Thr Ile Asp Trp Phe Gln Pro Asn Asn Lys Arg Asn Gln
 1 5 10 15

<210> SEQ ID NO 150
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 150

Asp Trp Phe Gln Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu His
 1 5 10 15

<210> SEQ ID NO 151
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 151

Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu His Leu Asn Thr His
 1 5 10 15

<210> SEQ ID NO 152
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 152

Arg Asn Gln Leu Trp Leu His Leu Asn Thr His Thr Asn Val Glu
 1 5 10 15

<210> SEQ ID NO 153
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 153

Trp Leu His Leu Asn Thr His Thr Asn Val Glu His Thr Gly Leu
 1 5 10 15

<210> SEQ ID NO 154
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 154

Asn Thr His Thr Asn Val Glu His Thr Gly Leu Gly Tyr Ala Leu
 1 5 10 15

<210> SEQ ID NO 155

-continued

```

<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 155
Asn Val Glu His Thr Gly Leu Gly Tyr Ala Leu Gln Asn Ala Thr
1           5           10           15

<210> SEQ ID NO 156
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 156
Thr Gly Leu Gly Tyr Ala Leu Gln Asn Ala Thr Thr Ala Gln Asn
1           5           10           15

<210> SEQ ID NO 157
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 157
Tyr Ala Leu Gln Asn Ala Thr Thr Ala Gln Asn Tyr Val Val Arg
1           5           10           15

<210> SEQ ID NO 158
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 158
Asn Ala Thr Thr Ala Gln Asn Tyr Val Val Arg Leu Thr Ile Tyr
1           5           10           15

<210> SEQ ID NO 159
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 159
Ala Gln Asn Tyr Val Val Arg Leu Thr Ile Tyr Val Gln Phe Arg
1           5           10           15

<210> SEQ ID NO 160
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 160
Val Val Arg Leu Thr Ile Tyr Val Gln Phe Arg Glu Phe Ile Leu
1           5           10           15

<210> SEQ ID NO 161
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 161
Thr Ile Tyr Val Gln Phe Arg Glu Phe Ile Leu Lys Asp Pro Leu
1           5           10           15

<210> SEQ ID NO 162
<211> LENGTH: 15
<212> TYPE: PRT

```

-continued

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 162

Tyr Val Gln Phe Arg Glu Phe Ile Leu Lys Asp Pro Leu Asn Glu
 1 5 10 15

<210> SEQ ID NO 163

<211> LENGTH: 1759

<212> TYPE: DNA

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 163

```

accagcgcac ttcggcgcgc gcagcacctc ggcagcgtca gtgaaaatgc caagcaagaa    60
aagcggcccc caaccccata agaggtgggt gttcaccctt aataatcctt ccgaggagga    120
gaaaaacaaa atacgggagc ttccaatctc cctttttgat tttttggtt gccggagagga    180
aggtttgtaa gagggtagaa ctccctcacct ccaggggttt gcgaattttg ctaagaagca    240
gacttttaac aaggtgaagt ggtattttgg tgcccgcctg cacatcgaga aagcgaagg    300
aaccgaccag cagaataaag aatactgcag taaagaaggc cacatactta tcgagtgtgg    360
agctccggcg aaccagggga agcgcagcga cctgtctact gctgtgagta cccttttgga    420
gacggggtct ttggtgactg tagccgagca gttccctgta acgtatgtga gaaatttccg    480
cgggctggct gaacttttga aagtgagcgg gaagatgcag aagcgtgatt ggaagacagc    540
tgtcacagtc atagtgggcc cgcgggttg tgggaagagc cagtgggccc gtaattttgc    600
tgagcctagg gacacctact ggaagcctag tagaaataag tgggtgggatg gatatcatgg    660
agaagaagtt gttgttttgg atgattttta tggctgggta ccttgggatg atctactgag    720
actgtgtgac cggtatccat tgactgtaga gactaaaggg ggtactgttc cttttttggc    780
ccgcagtatt ttgattacca gcaatcaggc ccccagcaa tggtagctct caactgctgt    840
cccagctgta gaagctctct atcggaggat tactactttg caattttgga agactgctgg    900
agaacaatcc acggaggtac ccgaaggccg atttgaagca gtggaccac cctgtgccct    960
tttcccatat aaaataaatt actgagtctt ttttgttacc acatcgtaat ggtttttatt   1020
tttatttatt tagagggctt tttaggataa attctctgaa ttgtacataa atagtcagcc   1080
ttaccacata attttgggct gtggttgcac tttggagcgc atagcccagg cctgtgtgct   1140
cgacattggt gtgggtatth aaatggagcc acagctggtt tcttttatta tttgggtgga   1200
accaatcaat tgttttggtc agctcagggt tgggggtgaa gtacctggag tggtaggtaa   1260
agggctgcct tatggtgtgg cgggaggagt agttaatata ggggtcatag gccaaagtgg   1320
tggagggggg tacaaagttg gcatccaaga taacaacagt ggaccaaca cctctttgat   1380
tagaggtgat ggggtctctg gggtaaaatt catatttagc ctttctaata cggtagtatt   1440
ggaaaggtag gggtaggggg ttggtgccgc ctgagggggg gaggaactgg ccgatgttga   1500
atctcagcta gttaacatcc caagatggct gcgagtatcc tccttttatg gtgagtacaa   1560
attctgtaga aaggcgggaa ttgaagatac ccgtctttcg gcgccatctg taacgggttc   1620
tgaaggcggg gtgtgccaaa tatggtcttc tccggaggat gttccaaga tggctgcggg   1680
ggcgggtcct tcttctgcgg taacgcctcc ttggccactg catcctataa aagtgaagaa   1740
agtgcgctgc tgtagtatt                                     1759

```

<210> SEQ ID NO 164

<211> LENGTH: 1759

<212> TYPE: DNA

-continued

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 164

```

accagcgcac ttcggcagcg gcagcacctc ggcagcgtca gtgaaaatgc caagcaagaa    60
aagcggcccc caaccccata agaggtgggt gttcacccctt aataatcctt cagaggagga    120
gaaaaacaaa atacgggagc ttccaatctc cctttttgat tattttgttt gcggagagga    180
aggtttggaa gagggtagaa ctccctcacct ccaggggttt gctaattttg ctaagaagca    240
gacttttaac aaggtgaagt ggtatttttg tgcccgtgc cacatcgaga aagcgaagg    300
aaccgaccag cagaataaag aatactgcag taaagaaggc cacatactta tcgagtgtgg    360
agctccgcgg aaccagggga agcgcagcga cctgtctact gctgtgagta cccttttggg    420
gacgggtgct ttggtgactg tagccgagca gttccctgta acgtatgtga gaaatttccg    480
cgggctggct gaacttttga aagtgagcgg gaagatgcag aagcgtgatt ggaagacagc    540
tgtacacgtc atagtgggcc cgcccggttg tgggaagagc cagtggggcc gtaattttgc    600
tgagcctage gacacctact ggaagcctag tagaaataag tgggtgggatg gatatcatgg    660
agaagaagtt gttgttttgg atgattttta tggctgggta ccttgggatg atctactgag    720
actgtgtgac cggtatccat tgactgtaga gactaaaggc ggtactgttc cttttttggc    780
tcgcagtatt ttgattacca gcaatcaggc cccccaggaa tggctactcct caactgctgt    840
cccagctgta gaagctctct atcggaggat tactactttg caattttgga agactgctgg    900
agaacaatca acggaggtag ccgaaggcgg atttgaagca gtggaccacac cctgtgccct    960
tttcccatat aaaataaatt actgagtctt ttttgttatc acatcgtaat ggtttttatt   1020
tttatttatt tagagggctc tttaggataa attctctgaa ttgtacataa atagtcagcc   1080
ttaccacata attttgggct gtggttgcat tttggagcgc atagcccagg cctgtgtgct   1140
cgacattggt gtgggtattt aaatggagcc acagctggtt tcttttatta tttgggtgga   1200
accattcaat tgtttgttcc agctcaggtt tgggggtgaa gtacctggag tggtaggtaa   1260
agggctgcct tatggtgtgg cgggaggagt agttaatata ggggtcatag gccaaagtgg   1320
tggagggggt tacaaagttg gcatccaaga taacaacagt ggaccaaca cctctttcat   1380
tagaggtgat ggggtctctg gggtaaaatt catatntagc ctttctaata cggtagtatt   1440
ggaaaggtag gggtaggggg ttggtgccgc ctgagggggg gaggaactgg cggatgttga   1500
atctgaggtg gttaacatgc caagatggct gcgagtatcc tccttttatg gtgattacaa   1560
attctttaga aaggcggcaa ttgaagatc cgtcttttcg gcgccatctg taacgggttc   1620
tgaaggcggg gtgtgccaaa tatggtcttc tccggaggat gttccaaga tggctgcggg   1680
ggcgggtcct tcttctcggg taacgcctcc ttggccacgt catcctataa aagtgaagaa   1740
agtgcgctgc tgtagtatt                                     1759

```

<210> SEQ ID NO 165

<211> LENGTH: 312

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 165

```

Met Pro Ser Lys Lys Ser Gly Pro Gln Pro His Lys Arg Trp Val Phe
 1             5             10             15

Thr Leu Asn Asn Pro Ser Gly Gly Gly Lys Asn Lys Ile Arg Gly Leu
 20             25             30

Pro Ile Ser Leu Phe Asp Tyr Phe Val Cys Gly Gly Gly Gly Leu Gly

```

-continued

35					40					45					
Gly	Gly	Arg	Thr	Pro	His	Leu	Gln	Gly	Phe	Ala	Asn	Phe	Ala	Lys	Lys
	50					55					60				
Gln	Thr	Phe	Asn	Lys	Val	Lys	Trp	Tyr	Phe	Gly	Ala	Arg	Cys	His	Ile
65					70					75					80
Gly	Lys	Ala	Lys	Gly	Thr	Asp	Gln	Gln	Asn	Lys	Gly	Tyr	Cys	Ser	Lys
			85						90					95	
Gly	Gly	His	Ile	Leu	Ile	Gly	Cys	Gly	Ala	Pro	Arg	Asn	Gln	Gly	Lys
			100					105					110		
Arg	Ser	Asp	Leu	Ser	Thr	Ala	Val	Ser	Thr	Leu	Leu	Gly	Thr	Gly	Ser
		115					120					125			
Leu	Val	Thr	Val	Ala	Gly	Gln	Phe	Pro	Val	Thr	Tyr	Val	Arg	Asn	Phe
	130					135					140				
Arg	Gly	Leu	Ala	Gly	Leu	Leu	Lys	Val	Ser	Gly	Lys	Met	Gln	Gln	Arg
145					150					155					160
Asp	Trp	Lys	Thr	Ala	Val	His	Val	Ile	Val	Gly	Pro	Pro	Gly	Cys	Gly
				165					170					175	
Lys	Ser	Gln	Trp	Ala	Arg	Asn	Phe	Ala	Gly	Pro	Arg	Asp	Thr	Tyr	Trp
			180					185					190		
Lys	Pro	Ser	Arg	Asn	Lys	Trp	Trp	Asp	Gly	Tyr	His	Gly	Gly	Gly	Val
		195					200					205			
Val	Val	Leu	Asp	Asp	Phe	Tyr	Gly	Trp	Leu	Pro	Trp	Asp	Asp	Leu	Leu
	210					215					220				
Arg	Leu	Cys	Asp	Arg	Tyr	Pro	Leu	Thr	Val	Gly	Thr	Lys	Gly	Gly	Thr
225					230					235					240
Val	Pro	Phe	Leu	Ala	Arg	Ser	Ile	Leu	Ile	Thr	Ser	Asn	Gln	Ala	Pro
				245					250					255	
Gln	Gly	Trp	Tyr	Ser	Ser	Thr	Ala	Val	Pro	Ala	Val	Gly	Ala	Leu	Tyr
			260					265					270		
Arg	Arg	Ile	Thr	Thr	Leu	Gln	Phe	Trp	Lys	Thr	Ala	Gly	Gly	Gln	Ser
		275					280					285			
Thr	Gly	Val	Pro	Gly	Gly	Arg	Phe	Gly	Ala	Val	Asp	Pro	Pro	Cys	Ala
	290					295					300				
Leu	Phe	Pro	Tyr	Lys	Ile	Asn	Tyr								
305					310										

<210> SEQ ID NO 166

<211> LENGTH: 312

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 166

Met	Pro	Ser	Lys	Lys	Ser	Gly	Pro	Gln	Pro	His	Lys	Arg	Trp	Val	Phe
1				5					10					15	
Thr	Leu	Asn	Asn	Pro	Ser	Gly	Gly	Gly	Lys	Asn	Lys	Ile	Arg	Gly	Leu
			20					25					30		
Pro	Ile	Ser	Leu	Phe	Asp	Tyr	Phe	Val	Cys	Gly	Gly	Gly	Gly	Leu	Gly
		35					40					45			
Gly	Gly	Arg	Thr	Ala	His	Leu	Gln	Gly	Phe	Ala	Asn	Phe	Ala	Lys	Lys
	50					55					60				
Gln	Thr	Phe	Asn	Lys	Val	Lys	Trp	Tyr	Phe	Gly	Ala	Arg	Cys	His	Ile
65					70					75					80
Gly	Lys	Ala	Lys	Gly	Thr	Asp	Gln	Gln	Asn	Lys	Gly	Tyr	Cys	Ser	Lys
			85						90					95	

-continued

Gly Gly His Ile Leu Ile Gly Cys Gly Ala Pro Arg Asn Gln Gly Lys
 100 105 110
 Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Gly Thr Gly Ser
 115 120 125
 Leu Val Thr Val Ala Gly Gln Phe Pro Val Thr Tyr Val Arg Asn Phe
 130 135 140
 Arg Gly Leu Ala Gly Leu Leu Lys Val Ser Gly Lys Met Gln Gln Arg
 145 150 155 160
 Asp Trp Lys Thr Ala Val His Val Ile Val Gly Pro Pro Gly Cys Gly
 165 170 175
 Lys Ser Gln Trp Ala Arg Asn Phe Ala Gly Pro Ser Asp Thr Tyr Trp
 180 185 190
 Lys Pro Ser Arg Asn Lys Trp Trp Asp Gly Tyr His Gly Gly Gly Val
 195 200 205
 Val Val Leu Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp Asp Leu Leu
 210 215 220
 Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Gly Thr Lys Gly Gly Thr
 225 230 235 240
 Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn Gln Ala Pro
 245 250 255
 Gln Gly Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Gly Ala Leu Tyr
 260 265 270
 Arg Arg Ile Thr Thr Leu Gln Phe Trp Lys Thr Ala Gly Gly Gln Ser
 275 280 285
 Thr Gly Val Pro Gly Gly Arg Phe Gly Ala Val Asp Pro Pro Cys Ala
 290 295 300
 Leu Phe Pro Tyr Lys Ile Asn Tyr
 305 310

<210> SEQ ID NO 167

<211> LENGTH: 233

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 167

Met Thr Trp Pro Arg Arg Arg Tyr Arg Arg Arg Thr Arg Pro Arg
 1 5 10 15
 Ser His Leu Gly Asn Ile Leu Arg Arg Arg Pro Tyr Leu Ala His Pro
 20 25 30
 Ala Phe Arg Asn Arg Tyr Arg Trp Arg Arg Lys Thr Gly Ile Phe Asn
 35 40 45
 Ser Arg Leu Ser Thr Glu Phe Val Leu Thr Ile Arg Gly Gly His Ser
 50 55 60
 Gln Pro Ser Trp Asn Val Asn Tyr Leu Lys Phe Asn Ile Gly Gln Phe
 65 70 75 80
 Leu Pro Pro Ser Gly Gly Thr Asn Pro Leu Pro Leu Pro Phe Gln Tyr
 85 90 95
 Tyr Arg Ile Arg Lys Ala Lys Tyr Glu Phe Tyr Pro Arg Asp Pro Ile
 100 105 110
 Thr Ser Asn Gln Arg Gly Val Gly Ser Thr Val Val Ile Leu Asp Ala
 115 120 125
 Asn Phe Val Thr Pro Ser Thr Asn Leu Ala Tyr Asp Pro Tyr Ile Asn
 130 135 140
 Tyr Ser Ser Arg His Thr Ile Arg Gln Pro Phe Thr Tyr His Ser Arg
 145 150 155 160

-continued

```

Tyr Phe Thr Pro Lys Pro Glu Leu Asp Gln Thr Ile Asp Trp Phe His
      165                      170                      175

Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu His Leu Asn Thr His Thr
      180                      185                      190

Asn Val Glu His Thr Gly Leu Gly Tyr Ala Leu Gln Asn Ala Ala Thr
      195                      200                      205

Ala Gln Asn Tyr Val Val Arg Leu Thr Ile Tyr Val Gln Phe Arg Glu
      210                      215                      220

Phe Ile Leu Lys Asp Pro Leu Asn Lys
225                      230

```

```

<210> SEQ ID NO 168
<211> LENGTH: 233
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 168

```

```

Met Thr Trp Pro Arg Arg Arg Tyr Arg Arg Arg Thr Arg Pro Arg
 1                      5                      10                      15

Ser His Leu Gly Asn Ile Leu Arg Arg Arg Pro Tyr Leu Val His Pro
      20                      25                      30

Ala Phe Arg Asn Arg Tyr Arg Trp Arg Arg Lys Thr Gly Ile Phe Asn
      35                      40                      45

Cys Arg Leu Ser Lys Glu Phe Val Ile Thr Ile Arg Gly Gly His Ser
      50                      55                      60

Gln Pro Ser Trp Ile Val Asn Ile Leu Arg Phe Asn Ile Gly Gln Phe
      65                      70                      75                      80

Leu Pro Pro Ser Gly Gly Thr Asn Pro Leu Pro Leu Pro Phe Gln Tyr
      85                      90                      95

Tyr Arg Ile Arg Lys Ala Lys Tyr Glu Phe Tyr Pro Arg Asp Pro Ile
      100                      105                      110

Thr Ser Asn Glu Arg Gly Val Gly Ser Thr Val Val Ile Leu Asp Ala
      115                      120                      125

Asn Phe Val Thr Pro Ser Thr Asn Leu Ala Tyr Asp Pro Tyr Ile Asn
      130                      135                      140

Tyr Ser Ser Arg His Thr Ile Arg Gln Pro Phe Thr Tyr His Ser Arg
      145                      150                      155                      160

Tyr Phe Thr Pro Lys Pro Glu Leu Asp Gln Thr Ile Glu Trp Phe His
      165                      170                      175

Pro Asn Asn Lys Arg Asn Gln Leu Trp Leu His Leu Asn Thr His Thr
      180                      185                      190

Asn Val Glu His Thr Gly Leu Gly Tyr Ala Leu Gln Asn Ala Ala Thr
      195                      200                      205

Ala Gln Asn Tyr Val Val Arg Leu Thr Ile Tyr Val Gln Phe Arg Glu
      210                      215                      220

Phe Ile Leu Lys Asp Pro Leu Asn Lys
225                      230

```

```

<210> SEQ ID NO 169
<211> LENGTH: 206
<212> TYPE: PRT
<213> ORGANISM: Type A PWD circovirus

```

```

<400> SEQUENCE: 169

```

```

Met Ile Ser Ile Pro Pro Leu Ile Ser Thr Arg Leu Pro Val Gly Val
 1                      5                      10                      15

```

-continued

```

Pro Arg Leu Ser Lys Ile Thr Gly Pro Leu Ala Leu Pro Thr Thr Gly
      20                25                30

Arg Ala His Tyr Asp Val Tyr Ser Cys Leu Pro Ile Thr Leu Leu His
      35                40                45

Leu Pro Ala His Phe Gln Lys Phe Ser Gln Pro Ala Glu Ile Ser His
      50                55                60

Ile Arg Tyr Arg Glu Leu Leu Gly Tyr Ser His Gln Arg Pro Arg Leu
      65                70                75                80

Gln Lys Gly Thr His Ser Ser Arg Gln Val Ala Ala Leu Pro Leu Val
      85                90                95

Pro Arg Ser Ser Thr Leu Asp Lys Tyr Val Ala Phe Phe Thr Ala Val
      100               105               110

Phe Phe Ile Leu Leu Val Gly Ser Phe Arg Phe Leu Asp Val Ala Ala
      115               120               125

Gly Thr Lys Ile Pro Leu His Leu Val Lys Ser Leu Leu Leu Ser Lys
      130               135               140

Ile Arg Lys Pro Leu Glu Val Arg Ser Ser Thr Leu Phe Gln Thr Phe
      145               150               155               160

Leu Ser Ala Asn Lys Ile Ile Lys Lys Gly Asp Trp Lys Leu Pro Tyr
      165               170               175

Phe Val Phe Leu Leu Leu Gly Arg Ile Ile Lys Gly Glu His Pro Pro
      180               185               190

Leu Met Gly Leu Arg Ala Ala Phe Leu Ala Trp His Phe His
      195               200               205

```

<210> SEQ ID NO 170

<211> LENGTH: 206

<212> TYPE: PRT

<213> ORGANISM: Type A PWD circovirus

<400> SEQUENCE: 170

```

Met Ile Ser Ile Pro Pro Leu Ile Ser Thr Arg Leu Pro Val Gly Val
  1                5                10                15

Ala Arg Leu Ser Lys Ile Thr Gly Pro Leu Ala Leu Pro Thr Thr Gly
      20                25                30

Arg Ala His Tyr Asp Val Tyr Ser Cys Leu Pro Ile Thr Leu Leu His
      35                40                45

Leu Pro Ala His Phe Gln Lys Phe Ser Gln Pro Ala Glu Ile Ser His
      50                55                60

Ile Arg Tyr Arg Glu Leu Leu Gly Tyr Ser His Gln Arg Pro Arg Leu
      65                70                75                80

Gln Lys Gly Thr His Ser Ser Arg Gln Val Ala Ala Leu Pro Leu Val
      85                90                95

Pro Arg Ser Ser Thr Leu Asp Lys Tyr Val Ala Phe Phe Thr Ala Val
      100               105               110

Phe Phe Ile Leu Leu Val Gly Ser Phe Arg Phe Leu Asp Val Ala Ala
      115               120               125

Gly Thr Lys Ile Pro Leu His Leu Val Lys Ser Leu Leu Leu Ser Lys
      130               135               140

Ile Ser Lys Pro Leu Glu Val Ser Ser Ser Thr Leu Phe Gln Thr Phe
      145               150               155               160

Leu Ser Ala Asn Lys Ile Ile Lys Lys Gly Asp Trp Lys Leu Pro Tyr
      165               170               175

```

-continued

Phe	Val	Phe	Leu	Leu	Leu	Gly	Arg	Ile	Ile	Lys	Gly	Glu	His	Pro	Pro
			180					185					190		
Leu	Met	Gly	Leu	Arg	Ala	Ala	Phe	Leu	Ala	Trp	His	Phe	His		
		195					200					205			

We claim:

1. A vector capable of expressing at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 90% sequence identity to SEQ ID NO: 25 and at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 90% sequence identity to SEQ ID NO: 23.

2. The vector of claim 1, wherein the vector is a baculovirus vector.

3. A vector capable of expressing at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 95% sequence identity to SEQ ID NO: 25 and at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 90% sequence identity to SEQ ID NO: 23.

4. The vector of claim 3, wherein the vector is a baculovirus vector.

5. A vector capable of expressing at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 90% sequence identity to SEQ ID NO: 25 and at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 95% sequence identity to SEQ ID NO: 23.

6. The vector of claim 5, wherein the vector is a baculovirus vector.

7. A vector capable of expressing at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 95% sequence identity to SEQ ID NO: 25 and at least one polypeptide encoded by a nucleic acid sequence comprising a sequence having at least 95% sequence identity to SEQ ID NO: 23.

8. The vector of claim 7, wherein the vector is a baculovirus vector.

9. A vaccine composition, comprising the vector of claim 1 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

10. A vaccine composition, comprising the vector of claim 2 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

11. A vaccine composition, comprising the vector of claim 3 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

12. A vaccine composition, comprising the vector of claim 4 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

13. A vaccine composition, comprising the vector of claim 5 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic

14. A vaccine composition, comprising the vector of claim 6 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

15. A vaccine composition, comprising the vector of claim 7 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

16. A vaccine composition, comprising the vector of claim 8 and a pharmaceutically or veterinarily acceptable carrier, wherein said nucleic acid sequence encodes an immunogenic protein that induces a protective response effective against infection by a piglet weight loss disease virus.

17. An immunogenic composition, comprising an immunizing amount of the vector of claim 1 and a pharmaceutically or veterinarily acceptable carrier.

18. An immunogenic composition, comprising an immunizing amount of the vector of claim 2 and a pharmaceutically or veterinarily acceptable carrier.

19. An immunogenic composition, comprising an immunizing amount of the vector of claim 3 and a pharmaceutically or veterinarily acceptable carrier.

20. An immunogenic composition, comprising an immunizing amount of the vector of claim 4 and a pharmaceutically or veterinarily acceptable carrier.

21. An immunogenic composition, comprising an immunizing amount of the vector of claim 5 and a pharmaceutically or veterinarily acceptable carrier.

22. An immunogenic composition, comprising an immunizing amount of the vector of claim 6 and a pharmaceutically or veterinarily acceptable carrier.

23. An immunogenic composition, comprising an immunizing amount of the vector of claim 7 and a pharmaceutically or veterinarily acceptable carrier.

24. An immunogenic composition, comprising an immunizing amount of the vector of claim 8 and a pharmaceutically or veterinarily acceptable carrier.

25. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 9.

26. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 10.

27. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 11.

28. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 12.

29. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 13.

30. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 14.

31. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 15.

32. A method for treating or preventing porcine circovirus type B infection in a mammalian subject, comprising administering to said subject a therapeutically effective amount of a vaccine according to claim 16.

33. A recombinant polynucleotide comprising a sequence having at least 90% sequence identity to SEQ ID NO: 25 and at least 90% sequence identity to SEQ ID NO: 23.

34. A recombinant polynucleotide comprising a sequence having at least 95% sequence identity to SEQ ID NO: 25 and at least 90% sequence identity to SEQ ID NO: 23.

35. A recombinant polynucleotide comprising a sequence having at least 90% sequence identity to SEQ ID NO: 25 and at least 95% sequence identity to SEQ ID NO: 23.

36. A recombinant polynucleotide comprising a sequence having at least 95% sequence identity to SEQ ID NO: 25 and at least 95% sequence identity to SEQ ID NO: 23.

37. A method for detecting and quantifying at least one polynucleotide of a porcine circovirus type B (PCVB) by a hybridization method, wherein said polynucleotide comprises a sequence with at least 90% identity to SEQ ID NO: 25 and at least 90% sequence identity to SEQ ID NO: 23.

38. A method for detecting and quantifying at least one polynucleotide of a porcine circovirus type B (PCVB) by a hybridization method, wherein said polynucleotide comprises a sequence with at least 95% identity to SEQ ID NO: 25 and at least 90% sequence identity to SEQ ID NO: 23.

39. A method for detecting and quantifying at least one polynucleotide of a porcine circovirus type B (PCVB) by a hybridization method, wherein said polynucleotide comprises a sequence with at least 90% identity to SEQ ID NO: 25 and at least 95% sequence identity to SEQ ID NO: 23.

40. A method for detecting and quantifying at least one polynucleotide of a porcine circovirus type B (PCVB) by a hybridization method, wherein said polynucleotide comprises a sequence with at least 95% identity to SEQ ID NO: 25 and at least 95% sequence identity to SEQ ID NO: 23.

41. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 1.

42. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 2.

43. The method of claim 42, wherein the cells are SF9 cells.

44. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 3.

45. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 4.

46. The method of claim 45, wherein the cells are SF9 cells.

47. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 5.

48. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 6.

49. The method of claim 48, wherein the cells are SF9 cells.

50. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 7.

51. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vitro, comprising transfecting cells in culture with the vector of claim 8.

52. The method of claim 51, wherein the cells are SF9 cells.

53. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 1 to a mammal.

54. The method of claim 53, wherein the mammal is a pig.

55. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 2 to a mammal.

56. The method of claim 55, wherein the mammal is a pig.

57. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 3 to a mammal.

58. The method of claim 57, wherein the mammal is a pig.

59. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 4 to a mammal.

60. The method of claim 59, wherein the mammal is a pig.

61. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 5 to a mammal.

62. The method of claim 61, wherein the mammal is a pig.

63. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 6 to a mammal.

64. The method of claim 63, wherein the mammal is a pig.

65. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 7 to a mammal.

66. The method of claim 65, wherein the mammal is a pig.

67. A method for expressing ORF'1 and ORF'2 of a porcine circovirus type B (PCVB) in vivo, comprising administering the vector of claim 8 to a mammal.

68. The method of claim 67, wherein the mammal is a pig.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,425,444 B2
APPLICATION NO. : 11/588306
DATED : September 16, 2008
INVENTOR(S) : Andre Jestin et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page, Item [73] Assignee, after "Wyeth" please insert --Madison, New Jersey--.

Signed and Sealed this

Second Day of December, 2008

A handwritten signature in black ink that reads "Jon W. Dudas". The signature is written in a cursive style with a large, looped initial "J".

JON W. DUDAS
Director of the United States Patent and Trademark Office