




(86) Date de dépôt PCT/PCT Filing Date: 2003/11/21
(87) Date publication PCT/PCT Publication Date: 2004/06/03
(45) Date de délivrance/Issue Date: 2013/02/12
(85) Entrée phase nationale/National Entry: 2005/05/20
(86) N° demande PCT/PCT Application No.: IT 2003/000764
(87) N° publication PCT/PCT Publication No.: 2004/046192
(30) Priorité/Priority: 2002/11/21 (IT RM2002A000588)

(51) Cl.Int./Int.Cl. *C12N 15/10* (2006.01),
C07K 16/00 (2006.01), *C07K 16/18* (2006.01)
(72) Inventeurs/Inventors:
VISINTIN, MICHELA, IT;
CATTANEO, ANTONINO, IT
(73) Propriétaire/Owner:
ROTTAPHARM BIOTECH S.R.L., IT
(74) Agent: DENNISON ASSOCIATES

(54) Titre : METHODE D'ISOLEMENT D'ANTICORPS INTRACELLULAIRES VISANT A NEUTRALISER DES INTERACTIONS PROTEIQUES
(54) Title: METHOD FOR ISOLATING INTRACELLULAR ANTIBODIES ABLE TO NEUTRALIZE PROTEIN INTERACTIONS



(57) Abrégé/Abstract:

A method is described for isolating neutralizing intracellular antibodies of an interaction between a protein ligand x and a protein ligand y inside a cell. A method to identify a protein ligand x able to bind to a known y ligand using intracellular antibodies able to disrupt the interaction between x and y is also described. A method for the isolation of a set of antibody fragments against a significant proportion of the protein-protein interactions of a given cell (interactome) or against the protein interactions that constitute an intracellular pathway or network is also described.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
3 June 2004 (03.06.2004)

PCT

(10) International Publication Number
WO 2004/046192 A3

(51) International Patent Classification⁷: **C12N 15/10,**
C07K 16/00

(74) Agents: **CAPASSO, Olga** et al.; De Simone & Partners
S.p.A., Via V. Bellini, 20, I-00198 Roma (IT).

(21) International Application Number:
PCT/IT2003/000764

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
21 November 2003 (21.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
RM2002A000588
21 November 2002 (21.11.2002) IT

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **LAY LINE GENOMICS S.P.A.** [IT/IT]; Via di Castel Romano, 100, I-00128 Roma (IT).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **VISINTIN, Michela** [IT/IT]; c/o Lay Line Genomics S.P.A., Via di Castel Romano, 100, I-00128 Roma (IT). **CATTANEO, Antonino** [IT/IT]; c/o Lay Line Genomics S.p.A., Via di Castel Romano, 100, I-00128 Roma (IT).


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
23 December 2004

[Continued on next page]

(54) Title: METHOD FOR ISOLATING INTRACELLULAR ANTIBODIES ABLE TO NEUTRALIZE PROTEIN INTERACTIONS



(57) Abstract: A method is described for isolating neutralizing intracellular antibodies of an interaction between a protein ligand x and a protein ligand y inside a cell. A method to identify a protein ligand x able to bind to a known y ligand using intracellular antibodies able to disrupt the interaction between x and y is also described. A method for the isolation of a set of antibody fragments against a significant proportion of the protein-protein interactions of a given cell (interactome) or against the protein interactions that constitute an intracellular pathway or network is also described.

WO 2004/046192 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHOD FOR ISOLATING INTRACELLULAR ANTIBODIES ABLE TO NEUTRALIZE PROTEIN INTERACTIONS

5 The present invention concerns a method for the isolation of stable intracellular
antibodies, able to neutralize protein interactions. In particular, the invention concerns an
antibody fragment library for isolating and expressing functional antibodies in the
intracellular environment (intracellular antibodies), capable of interacting with specific
target proteins in the intracellular environment. The invention concerns a method for the
isolation of a set of antibody fragments against significant portion of the protein-protein
10 interactions of a given cell (interactome) or against the protein interactions that constitute
an intracellular pathway or network.

It is known that that antibodies can be used to interfere in a highly specific manner with
biological processes inside the cell (Cattaneo and Biocca 1997) (Cattaneo and Biocca
1999) (Biocca, Neuberger et al. 1990). A specific intracellular expression can result in the
15 inhibition of target protein function (protein knock-out). In this way, a phenotype of
interest can be conferred to a cell or to an organism by a suitable intracellular antibody.
In particular, protein knock-out represents an advantageous method to approach and
resolve the bottleneck in functional genomics (from genes to function). In fact, genomic
and proteomic programs provide a wealth of information for the identification of genes
20 and proteins but not of their function. The solution of the problem of previous art could
facilitate the identification and validation of target proteins for drug development as well.
Protein knock-out has remarkable advantages over alternative methods (gene knock-out
or RNA knock-out, antisense, ribozymes or RNAi) (Rossi 1999; Capecchi 1989; Fire
1999; Tavernarakis, Wang et al. 2000) since:

- 25
- the protein is directly correlated with its biological function;
 - it permits post-transduction modified versions of the protein to be knocked out or the
protein's compartmentalized subpopulations;
 - it allows modulation of knock-out intensity (gradual knock-out or knock down).

Since protein function very often depends on specific protein-protein interactions,
30 antibodies or antibody fragments able to interfere with protein-protein interactions will
allow selective and specific protein function neutralization, while allowing other
functions of the same protein, depending on other protein interactions.

Intracellular antibody technology can be extremely useful for functional genomics since it exploits:

- a virtually unlimited repertoire of antibodies against any protein;
- the possibility to direct these antibodies to sites where the protein exerts its function inside the cell, and
- the possibility to obtain a specific and localized protein knock-out.

However, its application to functional genomics and to global proteomics requires the technology be implemented in a high-throughput process. This is why intracellular antibody technology is presently not applicable to functional genomics.

10 Intracellular antibody technology uses genes for specific antibodies that are suitably modified and obtained from the natural immune system (mouse or human) or from antibody libraries (synthetic or natural) exposed on the surfaces of filamentous phages as antibody fragments (phage display technologies, ScFv).

15 In terms of affinity, stability and solubility, the quality of antibody fragments isolated with phage display technology is proportional to the size of the library used for in vitro selection (Vaughan, Williams et al. 1996), given that the probability of finding an antibody with the properties of interest is proportional to the diversity of the library. This is why it is necessary to have large antibody libraries in order to obtain high-affinity antibodies capable of recognizing any antigen (Griffiths, Malmqvist et al. 1993)
20 (Griffiths, Williams et al. 1994) (Sheets, Amersdorfer et al. 1998) (Vaughan, Williams et al. 1996) (Sblattero and Bradbury 2000). These libraries are tested with the antigens of interest, ligated in the solid phase; the antibodies are then selected by sequential cycles of absorption, washing, elution and growth. This selection procedure, however, does not ensure that selected antibodies are suitable for intracellular expression, since not all
25 antibodies are able to function in the intracellular environment. At the current state of the art, the performance of antibodies suited for intracellular expression is not predictable a priori (Biocca, Ruberti et al. 1995; Proba, Worn et al. 1998), since the intracellular physicochemical conditions (cytoplasm and nucleus) differ greatly from those in which the antibodies are naturally found and normally synthesized (endoplasmic reticulum and
30 secretory pathways).

Hence, an application of the technology to functional genomics requires a robust and predictable method to select a priori antibodies capable of functioning in the intracellular environment. An improvement has been achieved with the selection method based on a

double-hybrid system (IACT or Intracellular Antibody Capture Technology) (Visintin et al. 1999; Visintin et al. 2002) and described in international patent application PCT WO 00/54057. Antibodies in the IACT format are able to interact with the corresponding antigen and result to be effective and functional when expressed as ordinary intracellular antibodies in the final user system. The IACT selection strategy is effectively diagnostic and predictive for the identification of functional intracellular antibodies starting from large libraries. The IACT technology has also been used in a derived procedure to rapidly and effectively identify the epitopes recognized by antibodies selected on their respective antigens (in vivo epitope mapping, IVEM, patent application RM2000A00561) (Visintin et al. 2002).

However, the development of IACT does not yet allow intracellular antibody technology to be applied to functional genomics, which requires a parallel throughput of at least 500 to 1000 genes to several thousand genes. Moreover, the IACT procedure does not ensure that selected antibody fragments are able to neutralize the function of the target protein.

Only a subset of the selected antibodies, at most, will have the property of being neutralizing antibodies.

The solution to this problem constitutes the object of the invention, allowing a rapid, effective high-volume processing from genes to validated intracellular antibodies that are then expressed in appropriate cell models to study and neutralize the function of corresponding proteins.

In a specific implementation, the procedure described in this invention allows the direct isolation of antibody fragments able to specifically disrupt protein-protein interactions, thereby having the ability to selectively and specifically neutralize protein function.

Therefore, the procedure described allows to isolate antibody fragments being validated intracellularly and protein function neutralizing.

As already mentioned, the use of IACT to isolate validated intracellular antibodies is severely limited by low cell transformation efficacy (yeast or mammal) where double-hybrid selection is operated, which, in turn, does not allow the introduction of antibody libraries (of scFv or single domain antibodies) with a diversity greater than 10^5 antibodies, a range far from that required to ensure a significant probability of finding a clone with the desired properties (10^{10} — 10^{11}).

Presently, the IACT step is preceded by an enrichment of the library to reduce the diversity to levels compatible with the transformation in cells for the double-hybrid selection (Figure 1) by using:

- one or two absorption cycles on antigen columns in the solid phase in the case of a phage antibody library, or
- immunization of a mouse with the antigen if the antibody library is prepared from mouse lymphocytes.

These steps are extremely costly and slow. Furthermore, for the first step a good phage display library is needed. In both cases the enrichment step requires the availability of a protein coded by the gene of interest. This creates a highly significant bottleneck since the functional genomics programs provide the genes (cDNA, EST, ORF, etc.) but not the protein. The possible pipelines at the current state of the art are, therefore, limited by the fact that for each group of genes of interest an ad hoc preselection must be carried out each time.

The solution to this problem is offered by the construction of an antibody library expressed directly in the double-hybrid rather than in the phage display format. The entire procedure to obtain intracellular antibodies starting from the genes as input is facilitated, shortened, accelerated and rendered more parallel since the genes of interest could be directly inserted into the vector for IACT selection or its modifications, thus short-circuiting the need to express and purify the corresponding proteins.

A further implementation with respect to the IACT system is to ensure the isolation of neutralizing intracellular antibodies. In fact, the IACT system ensures only the isolation of biologically functional antibodies in an intracellular environment but does not select for neutralizing antibodies.

DESCRIPTION OF THE INVENTION

In the following, an intracellular antibody is a synthetic or recombinant antibody in a single chain format (scFv), FAb or single domain (dAb) antibody, able to fold and to be functional in an intracellular environment (i.e. cytoplasm, nucleus).

A library of intracellular antibodies shall comprise: a) library of intracellular antibodies in which the antibody genes are V regions derived from i) mouse lymphocytes; ii) lymphocytes derived from patients with autoimmune disease; iii) lymphocytes derived from healthy people; b) a library of intracellular antibodies of point a) obtained with the method of the genetic recombination by two homologous recombination sites (BIG-

SPLINT); c) a library of intracellular antibodies engineered on particular frameworks designed on the basis of the intracellular consensus sequences present in intracellular antibodies.

5 A specific object of the invention is a method to isolate intracellular antibodies able to neutralize an interaction between a known protein ligand x and a known protein ligand y inside a cell, comprising the steps of:

a) obtaining a recombinant yeast strain wherein:

- i) the protein ligand x and the protein ligand y are expressed, but
 - ii) at least one of x or y is expressed in an inducible manner, and
 - 10 iii) the interaction between the protein ligand x and the protein ligand would lead to the production either of a repressor of an essential function of the yeast cell or of a toxic agent, so that non viable yeast cells would be obtained;
- b) constructing an expression library of intracellular antibodies in the recombinant yeast strain of point a);
- 15 c) selecting transformed yeast clones by inducing the expression of both ligand x and ligand y; and
- d) isolating recipient yeast clones able to grow.

Another object of the invention is a method to isolate intracellular antibodies able to neutralize an interaction between a known protein ligand x and a known protein ligand y

20 inside a cell, comprising the steps of:

- a) obtaining an expression library of intracellular antibodies in a yeast strain with mat-a or mat-alpha sexual competence;
 - b) obtaining a recombinant yeast strain having a sexually competence opposite to that of the yeast strain in point a) wherein:
- 25 i) the protein ligand x and the protein ligand y are expressed, but
 - ii) at least one of x or y is expressed in an inducible manner, and
 - iii) the interaction between the protein ligand x and the protein ligand would lead to the production either of a repressor of an essential function of the yeast cell or of a toxic agent, so that non viable yeast cells would be obtained;
- 30 c) exposing yeasts of point a) and recombinant yeasts of point b) to conditions able to promote sexual mating and the expression of the ligands x and y;
- d) selecting transformed yeast clones by inducing the expression of both ligand x and ligand y; and

e) isolating recipient yeast clones able to grow.

In a preferred embodiment the recombinant yeast strain is obtained by:

- 5 a) cloning a first sequence encoding the protein ligand x and a second sequence encoding the protein ligand y into a yeast expression vector, each first and second sequence being under the control of each one of two portions of a bidirectional promoter, the first sequence being fused to a sequence encoding a first molecule, and the second sequence being fused to a sequence encoding a second molecule, so that when the protein ligand x and the protein ligand y interact said first molecule and said second molecule interact too in a way to induce the production either of a repressor of an essential function of the yeast
- 10 cell or of a toxic agent;
- b) transforming the recipient yeast strain with the vector of point a); and
- c) selecting transformed yeasts.

Preferably the expression vector in yeast comprises as a bidirectional promoter the promoters Gal 1 and Gal 10.

- 15 Preferably the first and the second molecule leads to the production of a repressor able to shut down the transcription of the yeast HIS3 gene.

Preferably the expression library of intracellular antibodies in yeast is obtained from mouse or human lymphocytes.

- 20 Preferably the expression library of intracellular antibodies in yeast is obtained in a scFv or in a single domain antibody format.

It is another object of the invention an intracellular antibody able to neutralize an interaction between a protein ligand x and a protein ligand y inside a cell obtainable according to the method described. Preferably the intracellular antibody neutralizes the interaction between p65 RelA and p65 RelA; alternatively between p65 RelA and p50

- 25 NF- κ B1; alternatively between A β 1-42 and A β 1-42; alternatively between JX-TK TrkA and PTB p66 Shc; alternatively between Synuclein and Synphilin.

It is another object of the invention a nucleotide sequence encoding the intracellular antibody as above defined.

- 30 It is another object of the invention a yeast expression vector for cloning a sequence encoding a protein ligand x and a sequence encoding a protein ligand y able to interact each other, comprising a bidirectional promoter wherein the first promoter portion is 5' to 3' followed by appropriate restriction cloning sites for the sequence encoding the protein ligand x and a first sequence encoding a first molecule; and the second promoter portion

is 5' to 3' followed by appropriate restriction cloning sites for the sequence encoding the protein ligand y and a second sequence encoding a second molecule, so that when the protein ligand x and the protein ligand y interact, said first molecule and said second molecule interact too in a way to induce the production either of a repressor of an essential function of the yeast cell or of a toxic agent.

It is another object of the invention a method to simultaneously identify both a protein ligand x able to bind to a known protein ligand y inside an eukaryotic cell, and an intracellular antibody able to neutralize said interaction between the protein ligand x and the known protein ligand y, comprising the steps of:

- 10 a) obtaining an expression library of intracellular antibodies in a yeast strain with mat-a or mat-alpha sexual competence;
- b) obtaining a recombinant yeast strain having a sexually competence opposite to that of the yeast strain in point a) wherein:
 - 15 i) the protein ligand y is expressed in an inducible manner, and
 - ii) the interaction between the protein ligand x and the protein ligand y would lead to the production either of a repressor of an essential function of the yeast cell or of a toxic agent, so that non viable yeast cells would be obtained;
- c) cloning into a suitable yeast expression vector a cDNA library containing the sequence encoding the protein ligand x;
- 20 d) transforming the yeast strain of point b) with the cDNA library of point c),
- e) selecting transformed yeasts;
- f) submitting the yeasts of point a) and the yeasts of point e) to such conditions to promote sexual reproduction and the expression of the bidirectional promoter in an environment able to induce the expression of the protein ligand x and protein ligand y;
- 25 g) selecting viable yeast clones in which the interaction between the protein ligand x and the protein ligand y is disrupted by the intracellular antibody.

It is another object of the invention a method to simultaneously identify both a protein ligand x able to bind to a known protein ligand y inside an eukaryotic cell, and an intracellular antibody able to neutralize said interaction between the protein ligand x and the known protein ligand y, comprising the steps of:

- 30 a) obtaining an expression library of intracellular antibodies in a yeast strain with mat-a or mat-alpha sexual competence;

- b) cloning a cDNA library containing the sequence encoding the protein ligand x and a sequence encoding the protein ligand y into a yeast expression vector, the cDNA library sequences being under the control of one of two portions of a bidirectional inducible promoter, the cDNA library sequences being fused to a sequence encoding a first molecule; and the sequence encoding the protein ligand y being under the control of the other of two portions of the bidirectional promoter, and being fused to a sequence encoding a second molecule, so that when the protein ligand x and the protein ligand y interact said first molecule and said second molecule interact too, in a way to induce the production either of a repressor of an essential function of the yeast cell or of a toxic agent;
- c) transforming a yeast strain having a sexually competence opposite to that of the yeast strain in point a) with the recombinant yeast expression vector of point b);
- d) selecting transformed yeasts;
- e) submitting the yeasts of point a) and the yeasts of point d) to such conditions to promote sexual reproduction and the expression of the bidirectional promoter in an environment able to induce the expression of the protein ligand x and protein ligand y;
- g) selecting viable yeast clones in which the interaction between the protein ligand x and the protein ligand y is disrupted by the intracellular antibody.

DETAILED DESCRIPTION OF THE INVENTION

- The authors have developed an intracellular antibody library according to a method comprising the following steps (Figure 2):
- a) recovering DNA encoding the immunoglobulin natural V regions of B cells from spleen cells or peripheral lymphocytes by isolation of RNA, synthesis of complement DNA and amplification of DNA encoding light and heavy chains of immunoglobulins;
- b) cloning said DNA using primers designed to contain restriction sites suitable for an expression vector comprising: 1) an antibiotic resistance gene different from the vector that expresses the protein x and y ligands, i.e. ampicillin; 2) a nutritional marker different from the vector that expresses the x and y protein ligands, i.e. LEU2; 3) replication origins for both E. coli (i.e. ColE1, fl) and for the yeast (i.e. 2 μ); 4) a polylinker suitable for the insertion of an antibody library; 5) one or more nuclear localization signals downstream from the sequence encoding the scFv, but also FAb and antibody domains;
- c) characterizing the library obtained by colony PCR-fingerprinting, and
- d) selecting the SPLINT by IACT (SPLINT-2HY) on a known panel of antigens.

The vector of item b) is the vector pVP16/mv1 (Fig.3).

To overcome the limitation posed by the low efficiency of yeast transformation the authors set up a technology, named BIG-SPLINT (Fig. 4a, Fig. 4b), that uses recombination events to obtain a larger intracellular library after creating smaller primary
5 libraries. A recombination strategy has been described as an alternative to create scFv fragment libraries expressed in phages (Sblattero and Bradbury, 2000; PCT WO 00311246), but it was never been applied to get intracellular antibody libraries in eukaryotic cells. The event of recombination is applied to the SPLINT intracellular library and is based on a modification of the pVP16 vector so as to clone two lox recombination
10 sites (lox P and lox P511) between VH and VL genes (pVP/16mv1-lox vector, Fig. 5). This system introduces a gene encoding the CRE recombinase induced by the GAL1 promoter only in the presence of galactose, integrated in the yeast genome. Initially, two mini-libraries are obtained in two pVP16/mv1-lox vectors that differ only in their nutritional and antibiotic markers. The VH and VL chains are then recombined after
15 transformation (or mating) in the yeast strain containing the CRE recombinase gene, whose activity is induced only after the introduction of the two scFv libraries in the yeast strain (Fig. 4a, Fig. 4b).

The invention overcomes the previous technical problems by completely avoiding protein manipulation steps in the process “gene - intracellular antibody”, thus allowing
20 the isolation of intracellular antibodies against any of proteins encoded by a genome, and permitting the isolation of intracellular antibodies against all protein-protein interactions in a cell.

The SPLINT technology of the invention is predictive for sequences preferred by antibodies in intracellular environments, thus augmenting the knowledge of the structure
25 adopted by intracellular antibodies in physiologically incompatible environments, and leading to the formulation of a family of consensus sequences of antibody scaffolds validated for intracellular expression.

A preferred embodiment of the invention makes use of antibody sequences already validated by the IACT procedure according to the following method:

30 **SUPER-SPLINT**

As the selection procedures for intracellular antibodies are implemented, information is acquired on validated intracellular antibody sequences. Therefore various SPLINT generations (SUPER-SPLINT) may be obtained by using validated antibody sequences

with the current IACT technology, as disclosed in the WO0054057 application, thus augmenting the intracellular stability and affinity of antibodies selected in vivo. The so called SUPER-SPLINT library is then obtained by grafting CDR (complementarity determining region) libraries on antibody scaffolds designed on the basis of information acquired from the study of validated intracellular antibodies. In this way, the size of the input library is smaller, thus permitting the antibody diversity to be increased in an antibody structure that is more targeted to intracellular expression.

The SPLINT libraries constitute an unlimited source of antibodies for a variety of uses in diagnosis, research, industry and therapy and, in general, for all those applications requiring more stable antibodies.

The SPLINT technology can be adapted to other cell types besides yeast, including *E. coli* and mammalian cells, so as to avoid the possible limitation of having antibodies against proteins expressed by the host cell.

The SPLINT technology offers a new field of application for intracellular antibodies not possible before, in the so-called inverse functional genomics or AntiGENomics (Fig. 6), that leads to the identification of proteins involved in a cellular function of interest. In this format, intracellular antibody libraries derived by SPLINT (SUB-SPLINT) are used in suitable cell lines, for example mammalian cell lines, to induce a phenotype of interest and to identify the proteins responsible for the underlying biological processes. The AntiGENomics method thus allows the identification of the proteins involved in a given biological function, and solves a typical bottleneck of proteomic screenings, namely to assign a function to the proteins identified.

The invention is now described according to following examples.

FIGURES

Figure 1. Diagram of IACT technology. The intracellular antibodies can be isolated after combination between the two-hybrid system and the phage display technology or after immunization of a mouse and the subsequent construction of a scFv or antibody domains (Tomlinson and Holliger, 2000) (Wu and Yazaki, 2000; Kortt et al., 2001; Todorovska et al., 2001) (Tanaka et al., 2003)) library on a vector that has expressed an activation domain. After in vivo selection in the yeast cell, the positive testing clones are isolated and the scFv can be used for in vitro and in vivo applications.

Figure 2. Diagram of the construction of a SPLINT library: the antibody variable regions can be cloned directly from a non-immunized mouse on a vector that expresses the

activation domain and two NLS fused in tandem after the scFv antibody fragment. The SPLINT library is then inserted in the yeast strain in an episomal form and can then be used for IACT screening immediately after transformation of the antigen to be used as bait.

5 **Figure 3.** Diagram of the plinker220 and pVP16mv1 vectors. The vectors presents 2 NLS fused at the 3' and 5' ends of scFv for pLinker220 or 2 NLS fused in tandem after the scFv in pVP16mv1. The scFv library is transcribed by the ADH1 promoter. Downstream from the library, there is an ADH term termination sequence. The vector also contains the sequence that encodes the ampicillinase gene (bla), the gene sequence LEU2 (nutritional
10 marker for selection in yeast), an E. coli origin of replication fl and an yeast origin of replication 2 μ .

Figure 4 a-b. Diagram of the technology used to create a BIG-SPLINT library. In this system, two SPLINT libraries cloned in two pVP16mv1 vectors are used that differ only in antibiotic resistance to E. coli. The vectors present lox sites for heterologous
15 recombination between the antibody chains. The gene encoding the CRE recombinase can be inserted or as a gene expressed by an inducible Gal1 promoter cloned in an episomal vector or can be present in the yeast strain previously constructed for this purpose.

Figure 5. Diagram of the pVP16mv1-lox and pVP16m2 kan vectors. The two vectors are conceptually identical to the pVP16mv1 vector. The only difference is that upstream and
20 downstream from the scFv-VP16 construct there are two lox511-loxPWT sequences that are recognized by the CRE enzyme that starts to recombine the variable chains directly in the intracellular environment.

Figure 6. Diagram of the AntiGENomics technology. The technology uses in vivo selection of a cDNA library created from a cell line and the SPLINT antibody library.
25 After isolation of a large panel of Ag-scFv pairs, the isolated genes are identified and the scFv they recognize are sequenced. These scFv are then expressed as a cell-specific library (SUB-SPLINT) in vivo, in the target cell of interest and used to recapture in vivo the proteins expressed in the cell, either directly or after a phenotypic selection. The intracellular scFv-protein complexes are then immunoprecipitated using the scFv
30 sequence tag and analyzed using classic proteomic techniques such as 2Degel and MS.

Figure 7. Diagram of the vectors pMIC-BD1 and pMIC-BD2. The following vectors were constructed to express the antigen used as bait downstream from a DNA-binding domain (lexA). The two vectors differ only in the lack of NLS in pMIC-BD2, which was

removed before mutagenesis. The ADH1 promoter promotes the transcription of the
lexA-Ag fusion protein. Downstream from this construct there is an ADH1 term
destruction sequence. The vector also has the sequence that encodes the chloramphenicol
(cam) gene, the TRP1 gene sequence (nutritional marker for selection in yeast), an origin
of replication fl in E. coli and an origin of replication 2 μ for the yeast.

Figure 8. Diagram of the construction of the anti-GFP library. A mouse is immunized
with the GFP protein. The mouse spleen is then used as a source of lymphocytes to
extract the RNA encoding the immunoglobulin variable chains. Once the cDNA has been
prepared, the variable chains are amplified with specific degenerated primers. The anti-
GFP library in the scFv format is then cloned in the pVP16mv1 vector and screened
against the GFP cloned in the pBMIC-BD1 vector.

Figure 9. X-gal assay and PCR-BSTNI fingerprinting on anti-CH2-Shc clones isolated
from the hyperimmune library obtained after immunization of mice with GFP protein and
selected with the IACT method.

Figure 10. The figure shows the various cloning steps of the SPLINT library: in the first
panel to the left one can see the analysis of total RNA extracted from the nonimmunized
mouse spleen. To the right, one can see the RT-PCR product reamplified with a mix of
primers specific for the variable chains. The purpose of this step is to evaluate the quality
of the cDNA before proceeding to amplification programmed with the various primer
pairs (see table I). Below, one can see the analysis of the amplifications for the VH and
VL with each degenerated primer pair. Lastly, after having quantified each single
amplification for the VH and for the VL, each amplified product is mixed in equal
concentrations; the final result is illustrated in the upper right photograph.

Figure 11. Photograph of the VH and VL chain pull-through.

Figure 12. a. Photograph of the SPLINT library assembly in the scFv format.**b.** PCR-
BstNI fingerprinting of the SPLINT library after electroporation in the DH5 α F' strain.

Figure 13. X-gal assay and PCR-BstNI fingerprinting on anti-SH2-Shc clones isolated
from the SPLINT library selected with the IACT method.

Figure 14. X-gal assay and PCR-BstNI fingerprinting on anti-K-RAS clones isolated
from the SPLINT library selected with the IACT method.

Figure 15. X-gal assay for secondary screening on anti-SH2-SHc and anti-K-RAS clones
with respective antigens used as baits.

Figure 16. X-gal assay and PCR-BstNI fingerprinting on anti-Syk clones isolated from the SLPINT library selected with the IACT method.

Figures 17. Rescue scFv plasmids are reintroduced into original selection strain and into different bait strains to verify specificity. The isolated plasmid expressing the scFvs (i.e. anti-LT β 4) are reintroduced into strains containing target antigen (LT β 4) and another unrelated antigen (lamin). As an example clones b and h result to be specific only for LT β 4 and fail to interact with lamin antigen, thus confirming their specificity for LT β 4.

Figure 18. Anti-LT β 4 scFv fragment specifically recognized LT β 4 and fail to recognize the UT β 4 bait. The thymosins are a family of small proteins entirely composed of a single WH2 domain, a ~35 residue actin monomer-binding motif, that is found in many different regulators of the actin cytoskeleton. The lymphoid-specific thymosin encodes for an extra-six-residue NH2-terminal extension. Five antibody fragments obtained from the LT β 4 screening were tested for their ability to bind UT β 4. As shown in figure, all the scFvs tested result to be specific only for the lymphoid form and not for the ubiquitous one.

Figures 19. Diagram of the method used with SPLINT in the three-hybrid system (see explanation in Results).

Figure 20. Diagram of the pBiDi3HY, pBiDi3HYlexA-VP16 and pSLINT vectors.

pBiDi3HYlexA-VP16 contains a bidirectional promoter for the expression of two cDNA libraries (1 and 2). Library cDNA1 is expressed as a fusion protein with lexA, whereas library cDNA2 is expressed as a fusion vector with VP16. Both libraries have two different stop sequences. The vector expresses as a nutritional marker for selection in the TRP1 yeast. The vector also has the sequence encoding the chloramphenicol gene (cam), an origin of replication fl in E. coli and an origin of replication 2 μ for the yeast.

pSPLINT is a vector identical to pVP16mv1 but without the VP16 activation domain.

This vector is used only to express scFv in the intracellular environment.

Figure 21. Western blot analysis of lexA and VP16 proteins expressed upon induction with galactose in pBiDi3HYlexA-VP16 vector. After blotting, lexA was detected with the anti-lexA polyclonal antibody (Invitrogen) and VP16 with the anti-VP16 polyclonal antibody (Clontech).

Figure 22. Western blot analysis of lexA-p65 and VP16-p65 fusion proteins expressed upon induction with galactose in pBiDi3HYlexA-VP16 vector. After blotting, lexA-p65 was detected with the anti-lexA polyclonal antibody (Invitrogen) and VP16-p65 with the anti-VP16 polyclonal antibody (Clontech).

Figure 23. p65 monomer-anti-p65 scFvs interactions in the yeast IACT system. L40 yeast cells were cotransformed with lexA-p65 bait vector and/or lamin bait with anti-p65 A1 or anti-p65 A2 –VP16 fusions in pLinker220 vector. Yeast colonies were grown on –his plates and scored for β -gal assay. Clone A2 display a positive interaction with p65 monomer, thus indicating its specificity with the target protein. On the contrary, clone A1 was only able to recognize DNA-binding domain lexA.

Figure 24. Immunofluorescence microscopy of the anti-p65 A2 scFv transiently transfected in HEK cells. Cells were reacted with the anti-SV5 tag monoclonal antibody followed by incubation with an anti-FITC-conjugated anti-mouse (Vector).

Figure 25. HEK2973T cells were transiently transfected with anti-p65 A2 cloned in scFvex cyto-SV5 vector. Cell lysates were assayed for expression of p65 and p50 proteins (controls, c) and immunoprecipitation analysis with anti-p65 A2 or an irrelevant antibody. p65 protein was detected with anti-p65 (upper lane) and anti-p50 (lower lane) polyclonal antibody (Santa Cruz).

Vectors

plinker220 (Fig. 3).

The vector for the expression of the scFvs or a library of scFv fragments was designed to contain a polylinker that can be used to clone antibody fragments that contain at their termini unique restriction sites identical to those present on the polylinker. To create this vector, the polylinker of the pDAN3 vector was cloned (Sblattero et al. 2000) in VP16* (Hollenberg et al. 1995). To obtain this vector, the VP16*vector was cut with the NcoI enzyme, treated with Calf Intestinal Phosphatase (CIP) and purified with a 0.75% agarose gel preparation. The pDAN3 polylinker (where a scFv was cloned beforehand) was amplified using the following oligonucleotides:

oligonucleotide POLY BACK:

TCGATTTCACCATGGCGCCGAAAAAAAAAGAGAAAAGTGGCCCAGCCGAGCG
CGCATGCCGATATTGTG

Oligonucleotide POLY FOR:

TTCTTGGGTGCCATGGCGGAGCTCGAAACGCTAGCTGAGGAGACGGTGACCG
TG GT

The amplified fragment was then digested with the NcoI restriction enzyme and purified with 1.5% agarose gel. The two fragments were ligated using T4 DNA ligase. To confirm the perfect orientation of the insert and to confirm cloning, the vector was sequenced. The

VP16 activation domain was then modified by removing several nonessential amino acids. To do this, the plinker200 was used as a template and VP16 was amplified using the following nucleotides:

VP16-220 back: TCGAGCGGTACCGCTAGCGTTTCGAGCTCCGCCATGG

5 MIC-AD2 for: CGGCCAGTGAATTCCTACTACCCACCGTACTCGTCAAT

The fragment containing the VP16 activation domain was then digested using the restriction enzymes NheI and EcoRI and cloned in the modified VP16 vector, cut in the same fashion, treated with Calf Intestinal Phosphatase (CIP) and purified with a 0.75% agarose gel preparation. The two fragments were ligated using T4 DNA ligase. The
10 resulting vector was then called plinker220.

pVP16/mv1 (Fig. 3)

The expression vector of the scFvs or a library of scFv fragments, pVP16/mv1, was designed to contain a polylinker that can be used to clone antibody fragments containing at their termini unique restriction sites identical to those present on the polylinker. The
15 two nuclear location signals were shifted to permit their tandem expression downstream from the scFv and immediately upstream from the VP16 activation domain. This shift yields a scFv without steric hindrance at its 5' end, thus augmenting the specificity and folding of the antibody fragment fused to the VP16 activation domain at its 3' end.

pVP16/mv1 was constructed in the following way:

20 oligonucleotide MIC-AD1 back:

CAAACCGAAAAGCTTATGAGCGCGCATGCCGATATC

and a mixture of mouse VL oligonucleotides for the mix (Orlandi et al., 1992) were used to amplify a light chain cloned in plinker220. MIC-AD1 back contains the ATG start transcription codon and the restriction sites HindIII and BssHIII, while the mix of VL
25 mouse oligonucleotides for the mixture contains the restriction site SalI. The DNA fragment amplified with these nucleotides was cut using the enzymes HindIII and SalI and purified with a 1.5% agarose gel preparation. The plinker220 vector was cut in the same fashion, treated with Calf Intestinal Phosphatase (CIP) and purified with a 0.75% agarose gel preparation. The two fragments were ligated using T4 DNA ligase. The
30 resulting vector was then called pMIC-AD1.

pMIC-AD1 was used to prepare the final pVP16/mv1 vector. The following oligos were used:

MIC-AD2 back

CAAACCGCTAGCGCACCCAAGAAGAAGCGGAAGGTCGCACCCAAGAAGAAG
CGGAAGGTCCCAGGTGTACCGAGCTCG.

This nucleotide contains two nuclear location sequences (NLS) and the NheI restriction site.

5 MIC-AD2 for:

CGGCCAGTGAATTCCTACTACCCACCGTACTCGTCAAT.

This nucleotide contains the EcoRI restriction site.

The following nucleotides were used to amplify the VP16 activation domain from the plinker220 vector and to add upstream from it the two NLS encoded in tandem. The
10 resulting DNA fragment was cut using NheI-EcoRI and purified with a 1.5% agarose gel preparation. The plinker220 vector was cut in the same fashion, treated with Calf Intestinal Phosphatase (CIP) and purified with a 0.75% agarose gel preparation. The two fragments were ligated using T4 DNA ligase to create the pVP16/mv1 vector.

pMIC-BD 1 (Fig. 7)

15 The expression vectors pVP16/mv and pMIC-BD were designed to selectively change the resistance for expression in *E. coli*. The introduction of two different antibiotic resistances (ampicillin and chloramphenicol, respectively) permits the removal of the bait plasmid from the cotransformed cells with the two plasmids, thus eliminating the need for classic segregation in yeast, which is very long and costly.

20 The vector for the antigen expression in yeast was designed to contain a prokaryotic marker selectable by the antibiotic chloramphenicol. The pMIC-BD1 vector was constructed so: the sequence containing the region encoding the DNA-binding domain *lexA*, the transcription stop sequence of the *ADH1* gene and a part of the *TRP1* yeast gene was isolated from the pBTM116 vector (Bartel, Chien et al. 1993) by digesting the
25 plasmid with HindIII. The fragment was purified with 1.5% agarose gel. The expression vector used as a backbone, pBD-GAL4 Cam (Stratagene), was also cut using HindIII, dephosphorylated with Calf Intestinal Phosphatase (CIP), purified with 0.75% agarose gel and ligated with the *lexA-ADH1-TRP1* fragment to create the pMIC-BD1. The correct orientation of the insert was confirmed by sequence analysis.

30 **pMIC-BD2** (Fig. 7)

The *lexA* protein cloned in pMIC-BD1 was changed to abolish the nuclear location signal present inside the protein.

wt-*lexA* GTT ACC GTT AAG CGC CTG AAA AAA CAG GGC AAT

V T V K R L K K Q G N

mut-lexA GTT ACC GTT AAG GGC CTG GAA AAA CAG GGC AAT

V T V K G L E K Q G N

(Rhee et al., 2000).

- 5 The CGC and AAA codons of the region encoding lexA were then mutated in the following way:

lexA-NLS back oligonucleotide

AAGATGAAAGCGTTAACGGCCAGGCAACAAGAGGTG

that contains the restriction site HpaI, and

- 10 lexA-NLS for

AACGACAATTGGTTTAAACTCGCTATTTTCTGGCAACAGTTCGACTTTATTGC
CCTGTTTTTCCAGGCCCTTAACGGT

that contains the two mutations and the PmeI restriction sites were used to amplify a part of lexA. The DNA fragment amplified with these oligonucleotides was cut with enzymes

- 15 HpaI and PmeI and purified with a 2% agarose gel preparation. The pMIC-BD1 vector was cut in the same fashion, treated with Calf Intestinal Phosphatase (CIP) and purified with a 0.75% agarose gel preparation. The two fragments were ligated using T4 DNA ligase. The resulting vector was then called pMIC-BD2.

pVP16mv1-lox and pVP16mv2-kan (Fig. 5)

- 20 As described in Sblattero D. et al 2000, the following recombination sequences were inserted in the pVP16mv1 vectors:

loxP511 ATA ACT TCG TAT AAT GTA TAC TAT ACG AAG TTA T

loxPWT ATA ACT TCG TAT AAT GTA TGC TAT ACG AAG TTA T

The polylinker of the vector is modified so:

- 25 loxP511 linker

TCC GGA GGG TCG ACC ATA ACT TCG TAT AAT GTA TAC TAT ACG AAG TTA TCC TCG AGC GGT ACC
BspEI SalI loxP511 XhoI KpNI

The first lox site (loxP511) is added by pull-through of the variable chains VL and VH.

- 30 The primers for variable chain pull-through were:

VH PTL lox back

GGA GGG TCG ACC ATA ACT TCG TAT AAT GTA TAC TAT ACG AAG TTA
TCC TCG AGC GGT A

VH PTMIC for

CCA GGC CCA GCA GTG GGT TTG GGA TTG GTT TGC CGC TA

VL PTL lox for

ACC GCT CGA GGA TAA CTT CGT ATA GTA TAC ATT ATA CGA AGT TAT

GGT CGA CCC TCC

5 VL PTMIC back

CGC TGG ATT GTT ATT ACT CGC AGC AAG CGG CGC GCA TGC C

The second lox site (loxPWT) is added directly on vector pVP16 at 3' of the VP16 activation domain. The VP16 gene is amplified from pVP16mv1 with the following primers:

10 VP16loxPWT for (cutting with BamHI)

CTA TCT ATC TAg gat ccA TAA CTT CGT ATA GCA TAC ATT ATA CGA AGT

TAT CCC ACC GTA CTC GTC AAT

VP16 back (cutting with HindIII)

GGC TAA AGC TTA TTT AGG TGA CAC TAT

15 The amplification products are then directly added as fragment Hind III-Bam HI on the vectors cut with the same enzymes pVP16mv1 and pVP16mv1-kan. The kan gene + ColE1 are inserted in vector pVP16* as an amplification product amplified by the plasmid pZero-2 (Invitrogen) and cloned in the sites DraIII and AaTII.

Kan DraIII back

20 Ccg atc gga cac CAG GTG CTG TCA GAA GAA CTC GTC AAG AAG GC

ColE1 AaTII for

AA TGA AAT GCA TCG GGA CGT CCT TCC GCT TCC TCG CTC ACT GAC

Having cloned colE1 and the gene for kanamycin resistance, the vector is changed as previously described to obtain the pVP16mv2-kan vector.

25 Recombination of the variable chains takes place in vivo in the yeast cell. The CRE gene, which encodes recombinase, is transcribed by the Gal1 inducible promoter. pGal1-CRE can be a plasmid integrated in the yeast strain or an episomal vector can be introduced with antibiotic resistance (neomycin) that expresses CRE under the Gal1 promoter.

CRE can, for example, be cloned in a pESC vector (Stratagene) controlled by the Gal1

30 promoter. In this vector, antibiotic resistance (G418, neomycin) can be cloned to promote selection of this plasmid in the yeast cell. Gene G418 is transcribed by the Gal10 promoter, which is a bidirectional promoter with respect to Gal1. Both promoters are

induced in the presence of galactose in a culture medium, thus permitting the control of CRE enzyme recombinase expression.

pBiDi3HY vector (Fig.18):

to construct pBiDi3HY vector 2 vectors were used. The pESC-TRP vector (Stratagene) was PvuII cut to isolate the bidirectional promoters Gal1-Gal10. The pMIC-BD1 vector was SphI cut and DNA polymerase I, Large (Klenow) fragment was subsequently used to remove the 3' overhangs to form blunt ends. The vector was also dephosphorylated with calf intestinal alkaline phosphatase to prevent recircularization of cloning vector. Cloning of Gal1-Gal10 PvuII digested into pMIC-BD1 SphI cut and blunt generated pBiDi3HY.

pBiDi3HYlexA-VP16 vector (Fig.18):

the DNA fragment VP16 was amplified from VP16* vector (Hollenberg et al., 1995) and cloned between the BamHI/ApaI of pBiDi3HY plasmid.

The DNA fragment lexA was amplified from pMIC-BD1 and cloned between the EcoRI/SpeI of pBiDi3HY-VP16 vector. The final vector pBiDi3HY-lexA-VP16 was sequenced and the expression of lexA and VP16 proteins was verified by western blot analysis of yeast crude extract after induction of promoters with galactose 20% (see figure 23).

pBiDi3HYlexAp65-VP16p65 vector: the member of NF-kB/Rel family p65 protein, was cloned as C-terminal fusion protein with lexA and VP16 domains. The DNA fragment p65 was amplified from pRSV NF-kB relA (p65) plasmid and cloned into SpeI to produce lexA-p65 fusion protein and between ApaI/SalI to produce VP16-p65 fusion protein. The expression of both p65 fusion proteins were verified by western blot analysis and sequences (see figure 24).

All clones were sequenced, using the Epicentre Sequitherm Excel II kit (Alsbyte, Mill Valley, CA), with a Li-Cor 4000L automatic sequencer (Lincoln, NE).

Construction of a library of an immunized mouse (anti-GFP) (Fig. 8)

In the preparatory work leading to this invention, within the development of IACT technology, antibody libraries were constructed from the spleens of mice immunized against various antigens of interest, including EGFP (green fluorescent protein) (Prasher et al., 1992), human protein A β 1-42 and others. The selection of these libraries against the immunized antigens does not constitute a novel aspect but is mentioned because it introduces the unexpected result described in the paragraph below.

Protein EGFP was expressed in *E. coli* and purified according to a standard protocol developed by (Baird et al., 1999). EGFP was then conjugated with BSA (bovine serum albumin) to augment its immunogenicity in so far as GFP is a highly compacted protein. 0.7 mg of the purified protein (both native and conjugated GFP) were used for the initial immunization of three mice. The subsequent boosts contained 1.0—1.7 mg of the purified protein (~200-500µg of purified protein per mouse).

After about 2 months, the immunized mouse sera was analyzed by ELISA. The results showed that the mice presented a high titer of antibodies against GFP (we can find specific anti-GFP antibodies in analyzed sera also at higher dilutions, i.e. 1:10000). The spleens were then removed and used for RNA extraction.

The immunoglobulin variable regions were amplified using V-region PCR (Orlandi et al., 1992), using the following primers (Table I):

Table I. Panel of degenerated primers used for the construction of SPLINT libraries

VH mouse BACK:

15	TTA TCC TCG AGC GGT ACC GAK GTR MAG CTT CAG GAG TC HB1
	TTA TCC TCG AGC GGT ACC GAG GTB CAG CTB CAG CAG TC HB2
	TTA TCC TCG AGC GGT ACC CAG GTG CAG CTG AAG SAS TC HB3
	TTA TCC TCG AGC GGT ACC GAG GTC CAR CTG CAA CAR TC HB4
	TTA TCC TCG AGC GGT ACC CAG GTY CAG CTB CAG CAR TC HB5
20	TTA TCC TCG AGC GGT ACC CAG GTY CAR CTG CAG CAG TC HB6
	TTA TCC TCG AGC GGT ACC CAG GTC CAC GTG AAG CAG HB7
	TTA TCC TCG AGC GGT ACC GAG GTG AAS STG GTG GAA T HB8
	TTA TCC TCG AGC GGT ACC GAV GTG AWG YTG GTG GAG TC HB9
	TTA TCC TCG AGC GGT ACC GAG GTG CAG SKG GTG GAG TC HB10
25	TTA TCC TCG AGC GGT ACC GAK GTG CAM CTG GTG GAG TC HB11
	TTA TCC TCG AGC GGT ACC GAG GTG AAG CTG ATG GAR TC HB12
	TTA TCC TCG AGC GGT ACC GAG GTG CAR CTT GTT GAG TC HB13
	TTA TCC TCG AGC GGT ACC GAR GTR AAG CTT CTC GAG TC HB14
	TTA TCC TCG AGC GGT ACC GAA GTG AAR STT GAG GAG TC HB15
30	TTA TCC TCG AGC GGT ACC CAG GTT ACT CTR AAA GWG TST G HB16
	TTA TCC TCG AGC GGT ACC CAG GTC CAA CTV CAG CAR CC HB17
	TTA TCC TCG AGC GGT ACC GAT GTG AAC TTG GAA GTG TC HB18
	TTA TCC TCG AGC GGT ACC GAG GTG AAG GTC ATC GAG TC HB19

VH mouse FOR:

5 GAT TGG TTT GCC GCT AGC TGA GGA GAC GGT GAC CGT GGT HF1
GAT TGG TTT GCC GCT AGC TGA GGA GAC TGT GAG AGT GGT HF2
GAT TGG TTT GCC GCT AGC TGC AGA GAC AGT GAC CAG AGT HF3
GAT TGG TTT GCC GCT AGC TGA GGA GAC GGT GAC TGA GGT HF4

VL mouse BACK:

AGC AAG CGG CGC GCA TGC CGA YAT CCA GCT GAC TCA GCC LB1
AGC AAG CGG CGC GCA TGC CGA YAT TGT TCT CWC CCA GTC LB2
AGC AAG CGG CGC GCA TGC CGA YAT TKT GMT VAC TCA GTC LB3
10 AGC AAG CGG CGC GCA TGC CGA YAT TGT GYT RAC ACA GTC LB4
AGC AAG CGG CGC GCA TGC CGA YAT TGT RAT GAC MCA GTC LB5
AGC AAG CGG CGC GCA TGC CGA YAT TMA GAT RAM CCA GTC LB6
AGC AAG CGG CGC GCA TGC CGA YAT TCA GAT GAY DCA GTC LB7
AGC AAG CGG CGC GCA TGC CGA YAT YCA GAT GAC ACA GA LB8
15 AGC AAG CGG CGC GCA TGC CGA YAT TGT TCT CAW CCA GTC LB9
AGC AAG CGG CGC GCA TGC CGA YAT TGW GCT SAC CCA ATC LB10
AGC AAG CGG CGC GCA TGC CGA YAT TST RAT GAC CCA RTC LB11
AGC AAG CGG CGC GCA TGC CGA YRT TKT GAT GAC CCA RAC LB12
AGC AAG CGG CGC GCA TGC CGA YAT TGT GAT GAC BCA GKC LB13
20 AGC AAG CGG CGC GCA TGC CGA YAT TGT GAT AAC YCA GGA LB14
AGC AAG CGG CGC GCA TGC CGA YAT TGT GAT GAC CCA GWT LB15
AGC AAG CGG CGC GCA TGC CGA YAT TGT GAT GAC ACA ACC LB16
AGC AAG CGG CGC GCA TGC CGA CAG GCT GTT GTG ACT CAG GAA TC LBL

VL mouse FOR:

25 GAA GTT ATG GTC GAC CCT CCG GAA CGT TTK ATT TCC AGC TTG G LF1/2
GAA GTT ATG GTC GAC CCT CCG GAA CGT TTT ATT TCC AAC TTT G LF4
GAA GTT ATG GTC GAC CCT CCG GAA CGT TTC AGC TCC AGC TTG G LF5
GAA GTT ATG GTC GAC CCT CCG GAA CCT AGG ACA GTC AGT TTG G LFL

To create a library that is as representative as possible, single amplifications were performed using each oligonucleotide described above according to the following schemes:

[illegible]

HF3	x	X	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
HF4	x	X	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x

	LB1	LB2	LB3	LB4	LB5	LB6	LB7	LB8	LB9	LB10	LB11	LB12	LB13	LB14	LB15	LB16	LBL
LF1/2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	
HF3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	
HF4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	
HF5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	
LFL																	x

This procedure allows verification of the library quality, without having to introduce a prevalence of determinate variable regions that are more easily amplified than others. In this specific case, 141 individual amplifications are carried out, starting from the cDNA obtained from the total RNA extracted from the immunized mouse spleen. The conditions used to amplify the variable chains were:

cDNA 0.5 μ l; H₂O 30.5 μ l; 10X PCR buffer 5 μ l; dNTP 2.5mM each 5 μ l; MgCl₂ 25mM 3 μ l; Oligo1 (10pmol/ μ l) 2.5 μ l; Oligo 2 (10pmol/ μ l); 2.5 μ lTaq polymerase1 μ l are mixed.

Mineral oil (50-100 μ l) is added to prevent evaporation in the test tube containing the reaction mixture. The tube is then placed in a thermocycler programmed as follows: 94°C -5', (94°C -1' 60°C -1' 72°C -1') for 30 cycles; 72°C -10'; 4°C -24h.

After the variable regions have been amplified, they are analyzed on a 1.5% agarose gel and quantified. The same quantity of DNA for each amplified products is then used for cloning. A total of 10 μ g of a variable chain is then used for each cloning.

An initial VL library is created.

The light chains were processed as follows:

VL DNA mix 10 μ g; BssHII (NEB 20 U/ μ l)1.5 μ l; Buffer 3 (NEB)10 μ l; H₂O Q.S. (final V =100 μ l).

The reaction mixture is incubated at 50° C for 8 h. The digested chains are then purified with 1.5% gel using a purification kit (Gel extraction kit, Qiagen). The purified DNA is then digested using the SalI enzyme according to the following scheme: VL DNA mix BssHII 100 μ l; SalI (NEB 20 U/ μ l)1.5 μ l; Buffer Sal (NEB) 2 μ l; H₂O 78.5 μ l.

The reaction mixture is incubated at 37° C for 8 h. The digested chains are then purified with a 1.5% gel using a purification kit (Gel extraction kit, Qiagen). After purification, the DNA is quantified on 1.5% agarose gel and analyzed under spectrophotometry at a wavelength of 260 nm.

10 μ g of the pVP16/mv1 vector are cut in the same fashion: pVP16/mv1 DNA 10 μ g; BssHII (NEB 20 U/ μ l) 1.5 μ l; Buffer 3 (NEB) 10 μ l; H₂O Q.S. (final V=100 μ l).

The reaction mixture is incubated at 50° C for 8 h. The digested chains are then purified with 0.75% gel using a purification kit (Gel extraction kit, Qiagen). The purified DNA is

5 then digested using the Sall enzyme according to the following scheme: pVP16/mv1 BssHII 100 μ l; Sall (NEB 20 U/ μ l) 1.5 μ l; Buffer Sal (NEB) 20 μ l; H₂O 78.5 μ l.

The reaction mixture is incubated at 37° C for 8 h. The digested vector is dephosphorylated with Calf Intestinal Phosphatase (CIF) and purified with 0.75% agarose gel. After purification, the DNA is quantified on 0.75% agarose gel and analyzed
10 under spectrophotometry at a wavelength of 260 nm. pVP16/mv1 (BssHII-Sall-CIP) is then tested in E. coli to verify total digestion and dephosphorylation by electroporation of a known quantity of digested DNA incubated overnight with ligase or only in H₂O. After the vector background from a partial digestion of the vector or its religation has been verified (the background must be equal to or as close as possible to 0), pVP16/mv1
15 (BssHII-Sall-CIP) is then ligated with the light variable regions purified in a proportion of insert: vector=5:1.

About 6 μ g of the vector is used in the ligase mixture. The ligase reaction is then left overnight at 16° C, the ligase is then inactivated for 10 min at 70° C; the ligase mix is then purified in the following way: 1 purification step with phenol (1:1); 1 purification
20 step with phenol-chloroform (1:1); 1 purification step with chloroform-isoamyl (25:24:1) DNA precipitation with 3 volumes of sodium acetate/ethanol (solution 1 ml NaAc pH 5.2 + 27.5 ml EtOH). Wash the precipitated DNA with 200 μ l of 70% EtOH. The DNA pellet is then resuspended in 20—30 μ l of distilled H₂O.

10—40 ng of the purified ligase is then used for each electroporation. DH5 α F' or TG1 is
25 then made electrocompetent with the following method (Engberg et al., 1996) about 3—4 E. coli colonies are inoculated in 100 ml 2YT and grown at 37° C until O.D. 0.8—1. Two 50-ml flacon tubes are filled and left on ice for about 10 min. Thereafter, all steps are carried out in the cold room at 4° C. The tubes are centrifuged for 8 min at 3500 rpm at 4° C. The supernatant is removed. Each pellet is resuspended in 2 ml of 10% iced glycerol
30 buffer (10% sterile glycerol in distilled sterile H₂O). The resuspended bacteria are then brought to a volume of 50 ml in a glycerol buffer. The tubes are centrifuged for 8 min at 3500 rpm at 4° C. The supernatant is eliminated. The pellets are resuspended in 2 ml of 10% iced glycerol buffer and transferred to a falcon tube. The bacteria are then brought to

a volume of 50 ml in a glycerol buffer. Two further washings are performed according to the scheme described from points 6 to 8. Lastly, the final pellet is resuspended in 1 ml and brought to 3 ml in 10% glycerol buffer. The tubes are centrifuged for 8 min at 3500 rpm at 4° C. The supernatant is removed. The final pellet is resuspended in 125 µl 10% glycerol buffer. The bacteria are then divided into aliquots (30 µl per tube) in 0.5 ml Eppendorf tubes and conserved for about 1 month at -80° C.

The electroporator is set to 1800 V and the electrocompetent cells, together with the cuvettes, are placed on ice and cooled for about 10--15 min.

For each electroporation, 30 µl of bacteria plus 1 µl of ligase mixture (10—40 ng DNA) are used. After transformation, the bacteria are resuspended in 1 ml of SOC medium + 10 mM MgCl₂ and incubated for 1 h at 37° C. The bacteria are then seeded on LB + ampicillin plates. The next day, the efficacy of transformation is evaluated. To control the diversity of the VL library, 100 colonies are sampled at random from the plate and analyzed by PCR-fingerprinting using the BstNI restriction enzyme. After diversity has been confirmed, the bacteria are collected from the plate and frozen at -80° C in 20% glycerol.

The assembly of the VH chains with the VL chains is performed directly on the VL library. About 10µl is removed from the VL library glycerolate and inoculated in 50 ml LB + Amp overnight. The next day, the DNA for VH chain cloning is prepared. The heavy variable chains are processed as follows: VH DNA mix 10µg; XhoI (NEB 20 U/µl) 1.5µl; NheI (NEB 20 U/µl) 1.5µl; Buffer 2 (NEB)10µl; BSA (10mg/ml) 1µl; H₂O Q.S. (final V=100µl). The reaction mixture is incubated at 37° C for 8 h. The digested chains are then purified with a 1.5% gel using a purification kit (Gel extraction kit, Qiagen). After purification, the DNA is quantified on 1.5% agarose gel and analyzed under spectrophotometry at a wavelength of 260 nm.

10µg of the VL library is cut in the same fashion: VH DNA mix10µg; XhoI (NEB 20 U/µl) 1.5µl; NheI (NEB 20 U/µl) 1.5µl; Buffer 2 (NEB) 10µl; BSA (10mg/ml) 1µl; H₂O Q.S. (final V=100µl). The reaction mixture is incubated at 37° C for 8 h. The digested vector is dephosphorylated with Calf Intestinal Phosphatase (CIF) and purified with 0.75% agarose gel. After purification, the DNA is quantified on 0.75% agarose gel and analyzed under spectrophotometry at a wavelength of 260 nm. The VL library (XhoI-NheI-CIP) is then tested in E. coli to verify total digestion and dephosphorylation by electroporation of a known quantity of digested DNA incubated overnight with ligases or

only in H₂O. After the vector background from a partial digestion of the vector or its rebinding is verified (the background must be equal to or as close as possible to 0), the library is then ligated with the heavy variable regions purified in a proportion of insert: vector=5:1.

- 5 Thereafter, the procedure is continued as described above by the preparation of the VL library. A good library should have about 10^6 — 10^7 different scFv/ μ g of the DNA used. The pVP16/mv1 vector library is then assayed with GFP bait (immunizing antigen). The GFP bait was amplified from pEGFP-N1. The amplification primers of the GFP gene were: GFP sense: CCG GTC GAA TTC ATG GTG AGC AAG GGC GAG GAG and
10 GFP antisense: TGA TCT GGA TCC GCG GCC GCT TTA AGT GAT CCC GGC. Since previous experiments showed that GFP alone was able to transactivate the transcription of the reporter genes in the yeast system, a mutant of the GFP deletion was used for this experiment. The literature reports that the removal of the last 8 amino acids of the GFP maintains the fluorescent activity of the protein and its conformation
15 unchanged (Siegel and Isacoff, 1997).

The DNA fragment amplified with these oligonucleotides was then cut with enzymes EcoRI and BamHI and purified with a 1.5% agarose gel preparation. The pMIC-BD1 vector was cut in the same fashion, treated with Calf Intestinal Phosphatase (CIP) and purified with a 0.75% agarose gel preparation. The two fragments were ligated using T4
20 DNA ligase.

To verify transactivation of the bait GFP, self-activation of reporter gene transcription was tested according to the transformation steps shown below:

bait	Prey	YC selection medium	HIS3 phenotype	LacZ phenotype
LexA-GFP		-W	/	white
LexA-GFP		-WHUK	/	
LexA-GFP	VP16/mv1	-WL		white
LexA-GFP	VP16/mv1	-WHULK	/	

25 Having confirmed the non transactivation of the bait GFP and its good expression in the yeast strain (a yeast extract transformed with bait GFP is tested by western blot analysis using a primary antibody that recognizes the lexA protein), selection of the library in yeast cells is performed. L40 expressing bait GFP is transformed with 100 μ g of scFv anti-GFP library according to the protocol described below:

Day 1: inoculation of L40 containing bait GFP in 5 ml of YC-UW overnight;

Day 2: inoculation of 100 ml of YC-UW with an aliquot of the overnight culture in order to have a dilution that allows the algorithmic growth phase to be reached the next day;

Day 3: the overnight culture is transferred in 1 l of heated YPAD to obtain a culture with an OD 600= 0.3; the yeast is grown at 30° C for 3 h, the cells are centrifuged at 1500 rpm for 5 min at room temperature; the yeast pellet is washed with 500 ml of 1 x TE and then centrifuged at 1500 rpm for 5 min at room temperature; the pellet is resuspended in 20 ml of 1 x LiAC, 0.5 x TE and transferred to a fresh flask; 100 µg of scFv anti-GFP library and 1 ml of denatured salmon sperm are added; 140 ml of 1 x LiAc, 40% PEG 3350, 1 x TE; the product is mixed and incubated in a low-speed agitator for 30 min at 30° C; 17.6 ml of DMSO are added and mixed. Thermal shock is performed for 10 min at 42° C while agitating occasionally. The product is quickly cooled by adding 400 ml of YPA. The yeast is pelleted by centrifugation and washed once with 500 ml of YPA. After centrifugation, the pellet is resuspended in 1 l of YPAD preheated to 30° C. The product is incubated for 1 h at 30° C; 1 ml of the culture is isolated and the pellet obtained by centrifugation of this ml is resuspended in 1 ml of YC-UWL. Dilutions of 1:10, 1:100, and 1:1000 are seeded on YC-UWL plates to calculate the efficiency of the transformation. The pellet obtained from the remaining culture is washed twice with YC-WHULK. The final pellet is resuspended in 10 ml of YC-WHULK. The aliquots are plated on YC-WHULK plates and after 3—4 days the colonies that have grown are analyzed to determine the interaction.

96 colonies grown on YC-WHULK and testing blue on beta-Gal assay were analyzed by PCR-fingerprinting analysis using the BstNI restriction enzyme. The analysis of digestion patterns and scFv sequences isolated from the HIS+lacZ+ colonies permit the isolation of about a dozen different scFv that recognize the bait (also further to a second screening in yeast cells, the same bait that expresses GFP with scFv isolated in L40 is transformed) for each selection. The isolated scFv were then cloned in a scFv expression vector (Persic, Righi et al. 1997) for expression in mammalian cells and simultaneously in pUC119CAT for expression and production of the protein in E. coli. The binding affinity of the proteins purified with the affinity column was analyzed using plasmonic surface resonance. On average, the scFv isolated with IACT have an affinity constant of 100—500 nM. The scFvs expressed in vivo in mammalian cells (COS, CHO, PC12, C6, etc.) bind the target protein and in most cases manage to dislocate the same from its cellular expression environment if they are expressed with localization signals differing from those of antigen

expression. Also, neutralizing scFv can be found that inhibit antigen function or the antigen can be neutralized by taking it to another cell compartment using the Ag-Ab specific binding.

5 The results of this selection demonstrate that the enrichment obtained by immunization allows the construction of a library with a size adequate for yeast transformation and with a diversity wide enough to ensure the isolation of a panel of numerous antibodies against the immunizing protein.

Similar results have been obtained with a library created from the spleens of mice immunized against protein A β 1-42, a human protein, from which numerous antibodies
10 against the immunizing protein were derived (see table 2).

At this point, a surprising and completely unexpected result emerged from the screening of the hyperimmune spleen-derived libraries with bait constructed from proteins not relevant to immunization.

15 **Selection of a hyperimmune SPLINT library against antigens different from that used for immunization**

The libraries obtained from immunized mouse spleens (mice immunized with GFP), as described in the paragraph above, were screened against protein Shc (a protein that regulates cell growth, apoptosis and lifespan) and human A β 1-42 (the highly amyloidogenic component accumulation linked to the pathogenesis of Alzheimer's
20 disease).

A functional domain of the Shc protein (CH2-Shc) and the human A β 1-42 were cloned in the pMIC-BD1 vector and expressed in yeast to control the capacity of the two antigens to transactivate the reporter genes.

Construction of the baits:

25 **CH2-Shc/MIC-BD1**: the DNA fragment CH2-Shc was PCR amplified from pGEX4T1-p66Shc and cloned between EcoRI/BamHI restriction sites of pMIC-BD1 plasmid.

A β 1-42/MIC-BD1: the DNA fragment of beta amyloid 1-42 domain was synthesized by primers annealing and then cloned into BamHI/PstI restriction sites of pMIC-BD1 plasmid.

30 The pMIC-BD vector was constructed to be used in a yeast strain that kept the reporter genes under the control of the operator site for protein lexA. This vector also permits a rapid extraction of positive clones, after verification of an antibody-antigen interaction in

a two-hybrid system that uses as a vector containing the activation domain an ampicillin-resistant plasmid in that it utilizes chloramphenicol as the phenotype marker in *E. coli*. All antigens were tested in vivo as described elsewhere (Visintin and Cattaneo 2001) (Visintin, Tse et al. 1999) and the expression of the fusion proteins was assayed after the proteins were extracted from the transformed yeast strain. All antigens were found to be expressed in optimum quantities in the yeast and not to transactivate the reporter genes (HIS3 and lacZ).

CH2-Shc was selected as the antigen for library screening. The efficiency of the transformation of the library was about 2.5×10^4 clones/ μ g DNA. Unexpectedly, hundreds of positive colonies (Table III) were also obtained after the library was plated after the first day of transformation (see protocol in Visintin and Cattaneo 2001). The protocol used here was the same as that described above (see Construction of a library of an anti-GFP immunized mouse)

After about 15 min., the colonies turned blue on X-Gal assay, an indication that the observed interactions were quite strong. Twenty blue colonies were isolated and the plasmids containing the scFv fragments to be analyzed were rapidly isolated after extraction of total DNA from the yeast and transformed in *E. coli*. Analysis by PCR-BstNI-fingerprinting and a second two-hybrid screening yielded the following results (Fig. 9): 19 scFv, 9 classes of different scFv. Of the 9 classes of scFv, 8 tested specific against CH2-Shc and not against an irrelevant antigen (laminin) used for the screening. Likewise, in a separate selection from a library derived from a spleen hyperimmune for an unrelated antigen, and challenged with the A β 1-42 bait, a high number of different antibody domains against the A β 1-42 protein were isolated (data not shown, table III).

Table III. Selection of ICABs with different antigen specificities.

Antigen	cfu/ μ g	n. of SPLINT-DNA in L40	n. of His3 ⁺ -lacZ ⁺ rescued clones	n. of His3 ⁺ -lacZ ⁺ analyzed clones	n. positive clones
CH2-Shc	2.4×10^6	430	20		8
SH2-Shc	3.7×10^6	189	20		5
K-RAS	4.4×10^6	27	27		2
Syk	1.4×10^6	3000	10		6

UTβ4	3e10 ⁷	300	96	3
LTβ4	2.1e10 ⁶	123	123	13
Gephyrin	2.6e10 ⁶	5824	50	5
TFII-I	1.29e10 ⁶	1000	96	4
Aβ 1-42	1.2e10 ⁶	900	66	6
ORF X5	6.8e10 ⁵	34	34	6
SARS-CoV				

These results were obtained in just 1 month from the beginning of bait cloning (vector controls and rescreening included) and demonstrated the rapidity and precision of the system in obtaining antibodies against any protein of interest. The antibodies were then
5 cloned in a phagemid vector to produce soluble scFv for the in vitro specificity tests ELISA and plasmon surface resonance.

The conclusion is that although the selection libraries were hyperimmune toward the respective immunizing antigens (and contained numerous antibodies against the immunizing proteins, as demonstrated in the paragraph above), they also represent an
10 optimum source of intracellular antibodies against generic proteins. Hence, they represent SPLINT libraries and demonstrate that the spleen-derived libraries have unexpected properties required for SPLINT libraries, i.e. a size adequate for yeast cell transformation, although with a diversity wide enough to permit the isolation of antibodies against generic antigens.

15 The unexpectedly positive outcome obtained from a hyperimmune library, which should be enriched only for antibodies against immunizing antigens, laid the foundation for the experiment described below, i.e. the construction of a SPLINT library starting from a pool of mouse spleens that were not expressly immunized.

SPLINT library from non-immune mouse lymphocytes (Fig. 2)

20 Construction of SPLINT library

The spleen is rich in B cells and is an optimum tissue from which lymphocytes can be relatively easily extracted without needing to purify them by long and laborious methods. Mouse spleens (extracted from freshly sacrificed 3—6-month-old mice) were washed and cleaned in PBS and the lymphocytes released by pressing the spleen with a hypodermic
25 needle and a sterile spatula. The cell suspension was then transferred to a sterile tube and left to deposit for 5 min. The supernatant was then removed and the cells washed twice in

PBS and H₂O to remove the red blood cells. The cells were then immediately used for total RNA extraction. The total RNA was extracted using a kit (Rneasy Mini Kit, Qiagen) according to the manufacturer's protocols. After extraction, the RNA was quantified (about 30 µg from 10 mg of original tissue) and used for cDNA synthesis using random primers (GIBCO Brl). The synthesis of the DNA complementary to individual RNA filaments was obtained by using Superscript II Rnase H- Reverse Transcriptase enzyme (GIBCO Brl).

The following protocol was used for the production of cDNA: Random primers (3 µg/µl); 1 µl (100 µg); RNA total 5 µl; dNTP mix (10mM each dATP, dGTP, dCTP, dTTP at physiologic pH) 1 µl; H₂O DEPC 5 µl. The mix is heated to 65° C for 5 min, then rapidly cooled on ice. The content of the tube is collected after brief centrifugation and First-strand buffer 5 x 4 µl, DTT 0.1M 2 µl are added. The tube content is mixed and then incubated for 2 min at 42° C, after which 1 µl (200U) of Superscript II is added and incubated for 50 min at 42° C. The reaction is inactivated at 70° C for 15 min. 1 µl (2U) of RNaseH is added to remove the residual RNA bound to the formed DNA and is left for 20 min at 37° C. The cDNA can then be used to amplify the variable chains.

To obtain cDNA from the immunoglobulin variable regions, the V regions were amplified by PCR using degenerated 5' and 3' primers (TableI) that permit the amplification of a greater number of variable chains.

The amplification primers of the variable chains are identical to those described in the section Construction of an anti-GFP immunized mouse library.

A total of 141 amplifications were performed to be certain that a major number of antibody families were expressed in the final library. The single antibody chain amplifications are shown in Figure 10.

After all the VL and the VH variable regions have been amplified from the cDNA, they are reamplified to increase the quantity of DNA for cloning and to add extra sites to each terminus of the variable region previously amplified with the degenerated primers. After the variable chains (PTVL and PTVH) have undergone pull through, the amplified products are purified with gels (see Figure 11). After purification, the PTVL and the PTVH are used for in vitro assembly. The pull through is used to insert a small polypeptide chain (called linker) that, after in vitro assembly, is in turn needed to reconstruct the single chain sequence (see Figure 12). The assembly was done in a single step rather than in two steps as described in the section on anti-GFP library cloning

because a problem was noted in sequential cloning of the variable chains in the VP16mv vector. A possible explanation for this is the poor transformability of the pLinker220 vector in *E. coli* and the possible partial amplification of several variable chain families that make cloning in this vector particularly difficult.

5 The primers used for pull through were:

VH PTL 220 BACK

GGA GGG TCG ACC AGC GGT TCT GGG AAA CCA GGT TCC GGT GAA GGC
TCG AGC GGTA

VH PTMIC FOR

10 CCA GGC CCA GCA GTG GGT TTG GGA TTG GTT TGC CGC TA

VL PTL 220 FOR

ACC GCT CGA GCC TTC ACC GGA ACC TGG TTT CCC AGA ACC GCT GGT
CGA CCC TCC

VL PTMIC BACK

15 CGC TGG ATT GTT ATT ACT CGC AGC AAG CGG CGC GCA TGC C

The pull-through protocol was: VL 5 μ l (~100 ng); VL PTL 220 FOR 5 μ l; VL PTMIC BACK 5 μ l; MgCl₂ 2 μ l; dNTPs 10 μ l; buffer 10X 10 μ l; Taq polymerase 2 μ l; H₂O 61 μ l; VH 5 μ l (~100 ng); VH PTL 220 BACK 5 μ l; VH PTMIC FOR 5 μ l; MgCl₂ 2 μ l; dNTPs 10 μ l; buffer 10X 10 μ l; Taq polymerase 2 μ l; H₂O 61 μ l.

20 Mineral oil (50-100 μ l) is added to prevent evaporation in the test tube containing the reaction mixture. The tube is then placed in a thermocycler programmed as follows: 94°C -5'; (94°C -30'', 60°C -30'', 72°C -30'') for 20 cycles; 72°C -10'; 4°C -24h.

After the pull-through band for the VH and VL is isolated, the chains are purified with gel and the amplified products are used for in vitro assembly. In all, about 200 ng of DNA is
25 used for the reaction (100 ng of PTVL and 100 ng of PTVH).

The assembly reaction primers are:

VH PT2 FOR

TGG TGA TGG TGA GTA CTA TCC AGG CCC AGC AGT GGG TTT G

VL PT2 BACK

30 TAC CTA TTG CCT ACG GCA GCC GCT GGA TTG TTA TTA CTC

The reaction is: PTVL 100 ng; PTVH 100 ng; VL PT2 BACK (100 μ M) 0.5 μ l; VH PT2 FOR (100 μ M) 0.5 μ l; dNTPs 10 μ l; Buffer 10X 10 μ l; MgCl₂ 2 μ l; Taq polymerase 2 μ l; add H₂O to 100 μ l. Mineral oil (50-100 μ l) is added to prevent evaporation in the test tube

containing the reaction mixture. The tube is then placed in a thermocycler programmed as follows: 94°C –5'; (94°C –30", 68°C –30", 72°C –30") for 8 cycles without primers and 12 cycles with primers; 72°C –10'; 4°C –24h.

After assembly of the VH and the VL, the assembled scFv library is purified with gel as described above. After purification, the DNA is digested with enzymes NheI and BssH2
5 in the following way: DNA mix 10µg; NheI (NEB 20 U/µl) 1.5µl; Buffer 2 (NEB)10µl BSA (10mg/ml)1µl; H₂O Q.S. (final V=100µl) , and left to digest for 4 h at 37°C.

After digestion and purification with gel, the DNA is cut with BssH2: DNA mix 10µg BssH2 (NEB 20 U/µl) 1.5µl; Buffer 3 (NEB) 10µl; H₂O Q.S. (final V=100µl).

10 The reaction mixture is incubated at 50°C for 4 h. The digested chains are then purified with 1.5% gel using a purification kit (Gel extraction kit, Qiagen). After purification, the DNA is quantified on 1.5% agarose using spectrophotometry at a wavelength of 260 nm. The library is then ligated in the vector pLinker220: about 200ng of vector were used for this step; ligation was performed in gradient to optimize the reaction. The reaction mix is:
15 pLinker220: 1 µl (200ng); VL-VH assembly: 2µl (250 ng); Ligase:1µl (400U/µl); Buffer: 1µl; H₂O: 5µl; The reaction mixture is placed in an Eppendorf tube and incubated overnight in a beaker filled with water at 18° C in the cold room. The next day, the reaction tube is removed from the cold room and 1 µl of ligase is added and left at room temperature for 2 h. After incubation, the reaction mix is purified with phenol-chloroform
20 (as described above), resuspended in 10 µl of H₂O, and then electroporated in the bacterial strain.

For each electroporation, 30 µl of bacteria plus 1 µl of ligase mix is used (max. 100 ng per electroporation). After transformation, the bacteria are resuspended in 1 ml of SOC medium + 10 nM MgCl₂ and incubated for 1 h at 37° C, after which they are seeded on
25 LB+ ampicillin plates. The next day, the efficiency of the transformation is evaluated. The library thus obtained was estimated to be about 4.10⁵ CFU.

At this point, the SPLINT library was used for the control test such as PCR-fingerprinting to test library diversity, antibody sequences taken randomly, western blot analysis to check the expression of the individual chains in yeast and screening on a panel of known
30 antigens.

The results demonstrate that even from this initial SPLINT library intracellular antibodies against a wide variety of antigens can be isolated.

Description and list of antigens (tableII):

SPLINT was selected against the following antigens: two domains of Shc (Fig.13, 15), K-RAS (Fig.14, 15), Syk (Fig.16), β 4-thymosin ubiquitous, β 4-thymosin lymphoid (Fig.17, 18), amyloid β 1-42, gephyrin, TFII-I and X5 ORF SARS-CoV proteins.

Construction of the baits:

5 **SH2-Shc/MIC-BD1:** the DNA fragment SH2-Shc was PCR amplified from pGEX4T1-p66Shc and cloned between EcoRI/BamHI restriction sites of pMIC-BD1 plasmid.

CH2-Shc/MIC-BD1: the DNA fragment CH2-Shc was PCR amplified from pGEX4T1-p66Shc and cloned between EcoRI/BamHI restriction sites of pMIC-BD1 plasmid.

10 **Syk/MIC-BD2:** the Syk EcoRI/BamHI fragment from Syk/BTM116 (Visintin et al., 1999) was subcloned between EcoRI/BamHI restriction sites of pMIC-BD2 plasmid.

K-RAS/MIC-BD1 and K-RAS/MIC-BD2: K-RAS BamHI/PstI from K-RAS/BTM116 (Visintin et al., 1999) was subcloned between BamHI/PstI restriction sites of pMIC-BD1 and pMIC-BD2 respectively.

15 **LT β 4/MIC-BD1:** the DNA fragment LT β 4 was amplified from LT β 4/pCI-neo vector and cloned between the EcoRI/BamHI restriction sites of pMIC-BD1 plasmid.

UT β 4/MIC-BD1: the DNA fragment LT β 4 was amplified from UT β 4/pCI-neo vector and cloned between the EcoRI/BamHI restriction sites of pMIC-BD1 plasmid.

Gephyrin/MIC-BD1: the DNA fragment of the 153-348 domain of gephyrin protein was amplified and cloned between the EcoRI/BamHI restriction sites of pMIC-BD1 plasmid.

20 Before starting selection of SPLINT library all baits were tested for transactivation of reporter genes as described (Visintin and Cattaneo, 2001). All baits resulted to be non transactivating and thus used for the selections.

25 **A β 1-42/MIC-BD1:** the DNA fragment of β amyloid 1-42 domain was synthesized by primers annealing and then cloned into BamHI/PstI restriction sites of pMIC-BD1 plasmid.

TFII-I 1-397/MIC-BD1: the DNA fragment of the 1-397 domain of TFII-I transcription factor was amplified from human TFII-I cDNA and cloned into Bam HI restriction site of pMIC-BD1 plasmid)

30 **X5 ORF SARS-CoV/MIC-BD1:** the DNA fragment of the X5 ORF of SARS-Coronavirus was synthesized by primers annealing and then cloned into EcoRI/SalI restriction sites of pMIC-BD1 plasmid.

Selection of scFv intracellular antibodies with different specificities by IACT.

To verify whether we could select from this library intracellular antibodies against signalling proteins, and other intracellular proteins, we screened the library against a panel of different baits (Table II). These include, two domains of Shc, K-RAS, Syk, β 4-thymosin ubiquitous, β 4-thymosin lymphoid, amyloid β 1-42, gephyrin, TFII-I and X5 ORF SARS-CoV proteins.

Table II. Panel of different baits used for SPLINT-2HY selection.

- Human p66Shc CH2 domain (CH2-Shc)
- Human p66Shc SH2 domain (SH2-Shc)
- Human K-RAS
- Human Syk
- Mouse β 4-Thymosin ubiquitous (TU β 4)
- Mouse β 4-Thymosin lymphoid (TL β 4)
- Human Amyloid β 1-42
- Rat Gephyrin
- Human TFII-I
- X5 ORF SARS-CoV

The baits chosen for the validation of the SPLINT library include proteins of different species, of viral origin and of very different function and cellular location:

Two different domains (CH2 and SH2) of human p66^{Shc} adaptor protein (Ventura et al., 2002), which contains a unique amino-terminal proline-rich region (CH2) an NH2-terminal phosphotyrosine binding domain (PTB) followed by a collagen homology domain (CH1) and a COOH-terminal SH2 domain. By virtue of its PTB and SH2 domains, Shc proteins link receptor and nonreceptor tyrosine kinase activation to downstream cytoplasmic can bind to activated receptors that have been tyrosine-phosphorylated. This protein is implicated in pathways activated by environmental stresses and regulation of life-span.

Human K-Ras is a monomeric membrane-localized G protein of 21 kd that functions as a molecular or nuclear events. The ras-signaling pathway is an attractive target for anticancer therapy because of its important role in carcinogenesis.

Syk protein-tyrosine kinase has been implicated in a variety of hematopoietic cell responses, in particular immunoreceptor signaling events that mediate diverse cellular responses including proliferation, differentiation, and phagocytosis. Syk appears also to

play a general physiological function in a wide variety of cells including neuronal cells in which it is suggested to play an important role in signaling steps to neurite extension (Yanagi et al., 2001).

5 Thymosin β 4 encodes both an ubiquitous actin-binding protein (UT β 4) with demonstrated capacity to inhibit neutrophilic infiltration, and a splice-variant limited to lymphoid tissue (LT β 4) that was shown to be activation-responsive expressed by dendritic epidermal T cells, and other intraepithelial lymphocyte, to down-regulate local inflammation (Girardi, 2003).

10 Gephyrin is a 93 kDa protein that copurified with the glycine receptor (GlyR,) and was found to be localized at the postsynaptic side of glycinergic synapses (Prior et al., 1992) (Sassoe-Pognetto and Fritschy, 2000; Craig and Lichtman, 2001).

The major constituent of senile plaques in Alzheimer's disease is a 42-aa peptide, referred to as beta-amyloid (Abeta). Abeta is generated from a family of differentially spliced, type-1 transmembrane domain (TM)-containing proteins, called APP, by endoproteolytic processing. The major, relatively ubiquitous pathway of APP metabolism in cell culture involves cleavage by alpha-secretase, which cleaves within the Abeta sequence, thus precluding Abeta formation and deposition. Amyloid beta-peptide (1-42) (Butterfield, 15 2002) may be central to the pathogenesis of AD (Sinha and Lieberburg, 1999).

20 TFII-I is an inducible multifunctional transcription factor that is activated in response to a variety of extracellular signals and translocates to the nucleus to turn on signal-induced genes (Roy, 2001).

Severe acute respiratory syndrome (SARS) is a life-threatening form of atypical pneumonia. SARS-CoV genome is ~29.7 kb long and contains 14 open reading frames (ORFs) flanked by 5' and 3'-untranslated regions. Coronaviruses encode a number of non-structural proteins. These non-structural proteins, which vary widely among the different coronavirus species, are of unknown function and seems to be dispensable for virus replication (Rota et al., 2003; Snijder et al., 2003; Zeng et al., 2003). 25

DNA sequences coding for each of these proteins, fused to lexA protein to create antigen-specific baits, were introduced by transformation into yeast cells expressing the SPLINT library and IACT selection was performed as described (Visintin et al., 2002). It should be noted that no manipulation whatsoever of the corresponding protein was performed. Double transformants were selected for histidine prototrophy and lacZ activity. In the search for antigen-binding scFv fragments, about 10-20 His⁺/lacZ⁺ different clones for each 30

antigen were analyzed in a second two-hybrid screening, except for UT β 4, LT β 4, Ab1-42, Gephyrin and TFII-I (TableII) whose screening was more exhaustive (more than 90 His⁺/lacZ⁺ different clones). Specifically binding clones were obtained in each case, demonstrated that bait-specific single chain antibodies could be expressed in the cytoplasm and in the nucleus of yeast cells (Table III). Subsequent secondary screening confirmed that true positives could be identified that interact specifically with the original bait, but not with other lexA fusions (lexA-lamin).

8 clones for CH2-Shc, 5 clones for SH2-Shc, 2 for K-RAS, 6 for Syk, 11 for LT β 4, 2 for UT β 4 and 6 for gephyrin, 6 for A β 1-42, 4 for TFII-I and 6 for X5 ORF SARS-CoV were identified that interact with their antigens but did not interact with lexA-Lamin .

A subset of antibody fragments obtained from LT β 4 screening were also analyzed for binding to UT β 4 (Fig.18).

The lymphoid-specific thymosin encodes for a six-residue NH₂-terminal extension relative to the ubiquitous form of thymosin. In order to verify if it would be possible to isolate scFv specific only for the lymphoid form we tested five antibody fragments obtained from the LT β 4 screening that were shown to have the most different aminoacid sequences among the scFv isolated, for their ability to bind UT β 4. All the scFvs tested result to be highly specific for the lymphoid form and failed to bind the ubiquitous one (Fig.18). The parallel selection of SPLINT on UT β 4 resulted to be very stringent; very few different fingerprinting-patterns were found among 96 analyzed clones. Moreover it was shown that the selection of SPLINT against the UT β 4 bait was able to isolate scFvs whose aminoacid sequences correspond to some scFvs previously isolated against LT β 4, thus indicating that the selection of SPLINT against LT β 4 was exhaustive enough for fishing out specific antibodies against both forms of the thymosin protein.

In conclusion, the results obtained confirmed that indeed the SPLINT library constructed represents a good source for validated intracellular antibodies, against a wide variety of different proteins. The procedure to isolate such antibodies, which depends directly from the format of the library, requires no manipulation whatsoever of the target protein, thus obviating the need for protein expression and purification, that represents a significant experimental bottleneck for increasing the throughput of antibody isolation. The procedure described, that depends on the construction of the SPLINT library, allows isolating stable validated antibodies directly from gene sequences.

Improving the stringency of selections.

The results summarized in Table III shows that for each target protein chosen, a number of different antigen-specific antibodies could be isolated directly from gene sequences. The procedure has been experimentally optimized and refined and is very robust, rapid and efficient.

In order to fish out very strong binders and to improve the stringency of selections, if required, we also tested our SPLINT library by using a modified bait, fused with a mutant *lexA* protein. The bacterial *LexA* protein of pMIC-BD1 vector was modified (mLexA) to abolish its intrinsic NLS, as previously described (Rhee et al., 2000) (Fig.7). In this way, only the antibody/antigen complex formed in the cytoplasm has the information to be targeted to the nucleus. To check whether it should be possible to screen SPLINT also with antigen expressed only in the yeast cytoplasm (as in the case with the pMIC-BD2 vector) we chose the protein K-RAS as target antigen. The choice of RAS was also due to the apparent difficulty of isolating intracellular antibodies in the standard selection procedure (Table III). In this modified selection procedure, the interaction between the antigen and the scFv must occur in the cytoplasm, because the antigen is expressed in yeast cells without the intrinsic nuclear localization signal of *lexA*. ScFv and antigen interact in the cell cytoplasm before the complex is translocated into the nucleus and activates transcription. We tried in parallel the selection of K-RAS as *lexA*-bait fusion protein with or without nuclear localization signal. In both cases we obtained the same number of functional intracellular antibodies.

This indicates that the technology was strong enough to allow the selection of scFv with such an affinity to bring the antibody-antigen complex into the nucleus and to activate the transcription of reporter genes.

As a general comment, the selection with K-RAS either with or without NLS, gave very few positive clones if compared with other antigens (Table II). It should be noted that very few positive clones were also obtained in other reported IACT selection strategies when RAS was used as a bait (Tanaka and Rabbitts, 2003). This might indicate that the antigen has difficulty entering into the nucleus, and this difficulty interferes with the normal selection procedure. Alternatively, interference with the endogenous RAS protein might explain this low number of colonies obtained from this selection.

In conclusion we established that functional antigen-specific ICABs were selected from the natural SPLINT library against a wide spectrum of very different protein antigens,

including viral proteins. It should be noted that the optimized selection procedure, ensure that the selected antibodies are all functional also when expressed in mammalian cells. Indeed IACT selection predicts the property of being a functional intrabody in all cases (Visintin et al., 2002) (Tse et al., 2002). This was confirmed for a number of the ICABs selected from SPLINT, which showed good solubility and half life when expressed in mammalian cells (data not shown). Thus, SPLINT libraries represent a rich and good source for antibodies to function in the intracellular environment.

BIG SPLINT: Increasing the size of SPLINT

Library size is a critical variable for the isolation of antibody fragments. However, the true diversity of a library is not easily determinable; this in fact depends not only on the number of independent counted clones but also on the number of clones that actually express scFv proteins.

Notwithstanding the fact that we demonstrated here that it is possible to select functional intracellular antibodies from a SPLINT library of 10^6 - 10^7 clones, functional ICABs could be more easily directly selected making a larger SPLINT library. This has been accomplished by directly scaling up (i.e. repetition a number of times of the procedure described above) or by recombination methods (figure 4 a-b), as has been done in bacteria (Sblattero and Bradbury, 2000) and in yeast for making large antibody libraries of surface displayed Fab fragments (Feldhaus et al., 2003). The BIG-SPLINT library obtained in this way allows not only to isolate stable intracellular antibodies but also to increase the number of possible antibody candidates for intracellular expression.

Consensus sequence framework-based library

A further refinement of the SPLINT library will be carried out with the study of scFv sequences isolated by IACT.

It has been noted that the set of antibodies selected with these technologies is characterized by a number of highly conservative residues. These critical residues represent a sort of trademark that identifies those antibodies capable of being expressed in the intracellular environment by a multitude of scFv which are otherwise unable to function there. IACT technology combined with SPLINT is also able to select the scFv that possess an optimum sequence very similar to a consensus sequence present in the database of immunoglobulins present in nature. As the database of intracellular antibodies selected with this technology grows, the number of conserved residues will decrease. It will therefore be possible to achieve a final generic framework on which an antibody

library for intracellular use may be engineered. This will allow the creation of a library compatible with the low transformability of yeast and, more generally, a phagmid or phagic library of highly stable scFv that are functional for expression in any cellular or noncellular environment. It can be concluded that both IACT and SPLINT are two technologies with the capability to surpass their intrinsic potential and to thus create ever more sophisticated and validated libraries.

SPLINT libraries as a source of super stable antibodies, that can be isolated rapidly, directly from gene sequences.

It has been demonstrated above that a SPLINT library, as described above, represents a good generic source of intracellular antibodies, that is of antibodies/antibody domains that are able to bind any antigen of interest under conditions of intracellular expression. In light of the fact that these validated intracellular antibodies fold in the intracellular environment despite the fact that they fail to form the intrachain disulfide bond that links the two cysteine residues present in all known antibody variable regions (Visintin et al 1999; Biocca et al 1995), and in light of the fact that this disulfide bond contributes approximately 4.5 kCal/mol to the folding stability of an antibody domain (Proba et al., 1997), it can be concluded that antibodies generated through the IACT procedure from SPLINT libraries are much more stable than average antibodies in any given repertoire. In fact, when SPLINT derived antibodies are expressed under the normal oxidising conditions, thus allowing their intrachain disulfide bonds to form, their folding stability under experimentally controlled denaturing conditions (such as urea, guanidinium or thermal denaturation) is much greater than that of average antibodies (data not shown). Thus, SPLINT libraries allow isolating antibodies of superior stability. This property, together with the possibility of isolating antibodies directly from gene sequences, makes the SPLINT libraries the tool of choice to solve the bottleneck for the high throughput generation of stable antibodies, in applications such as antibody arrays, to be used as advanced diagnostic tools and for protein expression profiling in proteomics.

Protein-protein interaction targeted antibodies from SPLINT libraries:

The unexpected demonstration that a SPLINT library as described above represents a good source for intracellular antibodies represents the essential prerequisite for the final development of this invention, namely the exploitation of the SPLINT library, after suitable engineering, for the isolation of antibodies that are intrinsically neutralizing. In fact, the antibodies isolated from SPLINT are not necessarily endowed with the property

of inhibiting the biological function of their target protein. Indeed, the neutralization of protein function (PKO) is an added-on property, that has to be engineered or implemented on an ad hoc case-by case basis, with strategies such as, for instance, traffic diversion. These methods are not general, and therefore only a subset of antibodies will be useable to neutralize protein function, notwithstanding their in built property of being validated intracellular antibodies.

We describe in the following the use of SPLINT libraries to generate rapidly (directly from gene sequences) antibodies that are intrinsically endowed with the ability to interfere with a given protein-protein interaction. These will be intrinsically neutralizing antibodies.

Protein interactions play a crucial role in the discovery of diverse biological functions. The interpretation of genes sequenced from the human genome is leading researchers to develop new technologies for the interpretation of unpredictable functions of yet undiscovered genes. Diverse interactomes have been explored (protein interactions in a single organism). It has been seen that the two-hybrid system can be used to create a vast network of connections that can be further validated by other biotechnology system such as 2D gels, mass spectrometry, etc. The currently available approaches to studying protein function are limited and poorly generic. In contrast, with the methodology adopted for DNA study, the proteins are so variable that the study of the genome by protein interaction cannot be undertaken using a single methodology.

For this reason, researchers are examining whether antibody intracellular technology may offer an approach suitable for various fields of functional genomics.

For example, one can imagine constructing an ad hoc library of antibody fragments selected on a certain proteome (Fig. 6). The technology would entail a first step in which a cDNA library is created from a cell type, a tissue, a virus or a bacterium, etc. The library would then be directly cloned in the two-hybrid vector fused to a protein that binds DNA (e.g. pMIC-BD2) and tested against a SPLINT library in vivo. After positive clones have been obtained, the scFv involved in an interaction are isolated and cloned (in pools in a library format) in a eukaryotic expression vector (e.g. scFvexpress). After the new sub-library is created, the scFv can be expressed in the cell type, tissue, virus or bacterium from which the cDNA library was originally constructed. Once the library is introduced, a protein extract is prepared and used in immunoprecipitation experiments. The immunoprecipitated proteins are separated with a bidirectional gel (2D gel) and the

isolated proteins can then be analyzed by mass spectrometry. With this technology (IntrAP), antibodies directed against the entire cell proteome can be obtained in a single step.

Another possible consideration is the interactions and networks of genetic regulation at the level of a complete genome. In this way, it would be possible to conceive of the genome as a complex of functions expressed by proteins programmed to stabilize specific interactions. In this light, the total number of genes in an organism would appear less important than the complete repertory of interactions that may be potentially encoded by the same genome (interactoma). For this reason, one can conceive of constructing a subset of scFv libraries (similar to that described in the paragraph above) by isolating only the scFv that recognize protein modules or peptide sequences recognized as being a functional part of an interaction. The library could then be used to directly interfere in a known interactome through a methodology based on the three-hybrid system (Fig. 19). The 3HY-SPLINT strategy draws on various publications from the field of three-hybrid systems (Vidal M, et al. 1996; Huang J, et al. 1997; Leanna CA, et al. 1996; Shi HM, et al. 1996; Vidal M, et al. 1999).

The strategy uses a genetic selection in which the dissociation of an interaction becomes a selective advantage for the yeast strain used for this purpose. To reiterate the concept, the binding between protein X-DNA-BD and protein Y-AD is lethal for a yeast cell because in this system a toxic gene is used as a reporter gene. The dissociation by an antibody that binds one of the two interaction domains between X and Y confers a selective advantage that may be conveniently identified.

In our 3-HY system, the expression of the tetracycline repressor represses the expression of the HIS3 gene. Under these conditions, the interaction between X and Y confers an auxotrophic phenotype for histidine. When an antibody disrupts the interaction between X and Y, the TetR repressor is not expressed and the strain can survive also in the absence of histidine since gene HIS3 transcription is no longer repressed.

A novelty of our system is the use of two vectors instead of the classical set of three vectors to express either the two interacting proteins and the library of neutralizing antibodies. This allow to simplify the methodology, avoiding the difficult engineering of three vectors with different nutritional markers and different antibiotic resistance and the need of a particular yeast strain to be also engineered in which more than 3 genes must be knocked-out. Moreover a highly optimized protocols for the transformation of the yeast

cells should be assessed thus resulting in very poor reproducibility of the experiments. Our three hybrid methodology allow to maintain the current methodology of IAC technology as little modified and thus accessible to non specialized researchers. The results obtained so far are:

5 **STEP 1:lexA-p65 and VP16-p65 interact in two-hybrid system**

Construction of the bait and the prey:

p65/MIC-BD1: the DNA fragment of p65 protein member of NF-kB/Rel family was amplified from pRSV NF-kB relA (p65) plasmid and cloned between BamHI/Sall restriction sites of pMIC-BD1.

10 **p65/VP16c-t:** the DNA fragment of p65 was amplified from pRSV NF-kB relA (p65) plasmid and cloned between BamHI/NotI restriction sites of VP16c-t.

Intracellular interaction of lexA-p65 and VP16-p65 fusion proteins: L40 yeast cells were cotransfected with bait and prey vectors by using lithium acetate transformation protocol. Positive interaction was verified by using auxotrophic markers for both
15 plasmids and for histidine prototrophy. Histidine-positive colonies and controls were lysed in liquid nitrogen and assayed for b-gal activity on filters as described (Visintin and Cattaneo, 2001).

STEP 2: lexA-p65 and VP16-p65 interact in two-hybrid system when cloned in pBiDi3HYlexA-VP16 vector

20 **Construction of pBiDi3HY vector:** to construct pBiDi3HY vector 2 vectors were used. The pESC-TRP vector was PvuII cutted to isolate the bidirectional promoters Gal1-Gal10. The pMIC-BD1 vector was SphI cut and DNA polymerase I, Large (Klenow) fragment was subsequently used to remove the 3' overhangs to form blunt ends. The vector was also dephosphorylated with calf intestinal alkaline phosphatase to prevent
25 recircularization of cloning vector. Cloning of Gal1-Gal10 PvuII digested into pMIC-BD1 SphI cut and blunt generated pBiDi3HY.

Construction of pBiDi3HYlexA-VP16 vector: the DNA fragment VP16 was amplified from VP16* vector (Hollenberg et al., 1995) and cloned between the BamHI/ApaI of pBiDi3HY plasmid.

30 The DNA fragment lexA was amplified from pMIC-BD1 and cloned between the EcoRI/SpeI of pBiDi3HY-VP16 vector. The final vector pBiDi3HY-lexA-VP16 was sequenced and the expression of lexA and VP16 proteins was verified by western blot

analysis of yeast crude extract after induction of promoters with galactose 20% (see figure 20).

Construction of pBiDi3HYlexAp65-VP16p65 vector: the member of NF-kB/Rel family p65 protein, was cloned as C-terminal fusion protein with lexA and VP16 domains. The DNA fragment p65 was amplified from pRSV NF-kB relA (p65) plasmid (gently provided by Dr. Francesca Demarchi) and cloned into SpeI to produce lexA-p65 fusion protein and between ApaI/SalI to produce VP16-p65 fusion protein. The expression of both p65 fusion proteins were verified by western blot analysis and sequences (see figure 21).

All clones were sequenced, using the Epicentre Sequitherm Excel II kit (Alsbyte, Mill Valley, CA), with a Li-Cor 4000L automatic sequencer (Lincoln, NE).

Construction of pSPLINT vector (Fig.20): For construction of plasmid pSPLINT, the following cloning strategy was applied. A PCR Vk and VH cloned with 220 linker from pLINKER220 was performed by using these oligonucleotides: sense

5'-CATCATCATAAGCTTATTTAGGTGACACTA-3' and antisense

5'ACCCGGGGATCCCTAATGGTGATGGTGATGGTGAGTACTATCCAGGCCAG CAGTGGGTTTGGGATTGGTTTGCCGACCTTCCGCTTCTT-3'. The resulting PCR product was digested with HindIII/BamHI and ligated to HindIII/BamHI -digested vector VP16* (Hollenberg et al., 1995) to give pSPLINT. The activation domain VP16 was removed while maintaining both nuclear localization signals between scFv. A tag signal SV5 was added at the C-terminal of scFv linker.

Construction of SPLINT-3HY library.

A pool of scFv assembled from SPLINT library was also cloned into BssHII/NheI-digested pSPLINT to obtain SPLINT/pSPLINT library of $1.5 \sim 10^5$ different clones in E.coli.

STEP 3: lexA-p65 and VP16-p65 interaction can be abrogated by a neutralizing antibody in a SPLINT-3HY selection strategy

Co-transformation of pBiDi3HYlexAp65-VP16p65 vector with SPLINT-3HY library in YI596 yeast strain.

The genotype of *Saccharomyces cerevisiae* reporter strain YI596 is MATa/ α , his3 Δ 200/his3 Δ 200, trp1-901 leu 2-3, 112/leu2-3, 112ade2/ade2, URA::lexAOp)₈-TetRLYS2::ADH(Tet Op)₂-HIS.

The constructed system is a positive selection method designed to identify antibody domains that disrupt protein-protein interaction in vivo. The proteins of interest (p65 homodimers) are expressed as fusion with the lexA DNA binding domain and the transcriptional activator, VP16 under the control of the inducible bidirectional promoters Gal1-Gal10. Interaction between lexA-p65 and VP16-p65 leads the expression of the tetracycline repressor protein (TetR) which prevents transcription of the HIS3 reporter gene leading to the inability of yeast cells to grow on media lacking histidine. Disruption of homodimerization of p65 protein by a neutralizing antibody restores the ability to grow on such media. LexA-p65 fusion protein was shown to display activation capability. In order to block histidine production suppressing background growth on media lacking histidine a modulation of the system with tetracycline/3-AT was performed. After the identification of growth conditions of lexAp65-VP16p65 homodimer this construct was used for the selection of SPLINT library in order to identify the neutralizing antibody that disrupt p65 homodimer formation in vivo.

SPLINT library is introduced into YI595 strain by a modified protocol of yeast transformation (Visintin and Cattaneo, 2001). Each transformant presumably contains an individual library plasmid encoding a scFv in addition to the p65 homodimer proteins. Cells containing the plasmid encoding a scFv that bind one or both p65 interacting proteins will be selected on SG-Raf-WHULK. The transformation procedures must be carefully followed to ensure a high transformation efficiency.

SPLINT-3HY transformation

The following transformation protocol is a modification of published method (Visintin and Cattaneo, 2001).

Materials: 500 µg SPLINT/pSPLINT library; 150 ml YC-UKW ; 2 l YPAD; 1 l YPA; 1.5 l YC-UKWL + 10 YC-UKWL plates (100mm); 1 l YC-WHULK ; 1 l SG-Raf-WHULK + 3AT 25mM (see recipe below); 20 SG-Raf-WHULK + 3AT 25mM plates 100 ml 10X TE; 20 ml 10X LiAc; 1 ml of 10 mg/ml denatured salmon sperm; 150 ml 50% PEG 4000; 20 ml DMSO.

Procedure:

Day1: grow YI595 yeast containing p65 homodimer plasmid in YC-UKW O/N

Day2: inoculate 100 ml of YC-UKW with an aliquot of the overnight culture in order to find a dilution that places the 100 ml culture to logarithmic phase the next day

- Day3:** transfer enough overnight culture in 1 l of prewarmed to 30°C YPAD to produce an $OD_{600} = 0.3$. Grow at 30°C for 3 hours. Centrifuge the cells at 1500 X g for 5 min at room temperature. Wash pellet in 500 ml of 1X TE then centrifuge again the cells at 1500 X g for 5 min at room temperature. Resuspend pellet in 20 ml 1X LiAc, 0.5X TE and
- 5 transfer to a new flask. Add 500 µg DNA library and 1 ml denatured salmon sperm. Add 140 ml of 1X LiAc, 40% PEG 3350, 1X TE; mix and incubate for 30 min at 30°C with gently shaking. Add 17.6 ml of DMSO; swirl to mix. Heat shock for 10 minutes at 42°C in a water bath swirl occasionally to mix. Rapidly cool at room temperature cells in a water bath diluting with 400 ml YPA. Pellet cells by centrifugation and wash with 500
- 10 ml YPA. After centrifugation resuspend pellet in 1 l of prewarmed YPAD. Incubate at for 1 hour at 30°C with gently shaking. Pellet cells from 1 ml: resuspend in 1 ml YC-UKWL; spread 100 µl of a 1:1000, 1:100, 1: 10 dilutions for transformation efficiency controls. Pellet cells from the remaining culture. Wash pellet with 500 ml YC-UWL. Resuspend in 1l of prewarmed YC-UWL and incubate O/N at 30°C with gently shaking.
- 15 **Day4:** pellet cells and wash with 500 ml of YC-WHULK. Pellet cells and wash with 500 ml of SG-Raf-WHULK + 3AT 25mM . Resuspend final pellet in 10 ml of SG-Raf-WHULK + 3AT 25mM . Spread dilutions of the total on YC-UKWL plates to compare to the number of primary transformants Spread the remaining transformation suspension on SG-Raf-WHULK + 3AT 25mM plates.
- 20 **Synthetic Galactose Minimal Medium (SG dropout media)**
- 1. YNB w/o aa & $(NH_4)_2SO_4$:** 1.2 g yeast nitrogen base, w/o amino acids and ammonium sulfate (Difco # 0335-15-9); 20 g bacto-agar; Add H_2O to 800 ml. Autoclave 121°C for 15 min.
- 2. Salts:** 5.4g NaOH; 10g succinic acid (Sigma # S-7501); 5g ammonium sulfate (Sigma
- 25 # A-3920).
- 3. Gal-Raf:** 20g galactose; 10g raffinose
- Add H_2O to 100 ml and dissolve all components one by one. Add galactose and raffinose and H_2O to make a final volume of 150 ml.
- 4. amino acids MIX:** 5.8 g NaOH; 1 g Adenine hemisulfate salts; 1 g L-Arginine HCl
- 30 (Sigma # A-5131); 1 g L-Cysteine (Sigma # C-8277); 1 g L-Threonine (Sigma # T-8625); 0.5 g L-Aspartic acid (Sigma # A-4534); 0.5 g L-Isoleucine (Sigma # I-7383) 0.5 g L-Methionine (Sigma # M-9625); 0.5 g L-Phenylalanine (Sigma # P-5030)

0.5 g L-Proline (Sigma # P-4655) ; 0.5 g L-Serine (Sigma # S-5511). Dissolve in 80 ml H₂O

5. L-Tyrosine: 0.5 g L-Tyrosine (Sigma # T-3754); 0.2 g NaOH; Dissolve in 10 ml by heating.

5 6. Omitted amino acid solutions:

L-Histidine (Sigma # H-9511): 5 g/l H₂O

Uracil (Sigma # U-0750) : 10 g/l H₂O (+2 pellets NaOH)

L-Leucine (Sigma # L-1512): 10 g/l H₂O

L-Lysine HCl (Sigma # L-1262): 10 g/l H₂O

10 L-Tryptophan (Sigma # T-0254): 10g/lH₂O

Add to aa MIX (4.) the L-Tyrosine solution (5.) and H₂O to make a final volume of 100 ml. Filter –sterilize, aliquot and store at -20°C for up to 1 year. Filter-sterilize and aliquot individually omitted amino acids (aa) solutions (H, W, L, K, U) and store at -20°C for up to 1 year. Before preparing SG-Raf plates or media by mixing appropriate solutions, media must be adjusted to pH 5.8 and sterilized by filtering salts + aa final mix.

Results of SPLINT-2HY selections

It was already demonstrated that IACT is a genetic selection method that allows the isolation of soluble intrabodies directly from antigen gene sequences (Visintin et al., 2002).

20 The major goal of IAC technology was to develop intracellular one-pot scFv libraries of sufficient size and diversity to facilitate isolation of antibodies of every conceivable specificity, including antibodies with high solubility, stability and good affinity for intracellular expression. An important determinant of the composition and diversity of this *alternative in vitro* immune system is the source of antibody genes used as building blocks to construct the library. In order to explore the probability of isolating intrabodies directly from primary mice antibody repertoire we have engineered a single pot antibody library in an activation domain vector and tested it in a two-hybrid selection strategy by using a wide panel of different antigens.

25 The results obtained from all the different selections have indicated that SPLINT library is a good source of antibody fragments. In fact, it was possible to select several scFv against each antigen challenged with the IAC technology. For some target antigens it was possible also to isolate several antibodies thus indicating that although the library is not very complex the diversity obtained is still sufficient to acquire immunological reagents

that recognize particular antigens as for example, proteins that are highly conserved between species.

We have demonstrated that a small library which has proved to be enriched in functional antibodies can still provide a good source of antibody for ectopic expression because the method for selecting the antibody fragments act as a filtering funnel. SPLINT has demonstrated to contain a source of stable superior antibodies, naturally present in naïve repertoires but difficult to find if the methodology used is not specialized.

In the original format, SPLINT-2HY guarantee that the antibodies selected are good intracellular binder but cannot assure for neutralizing antibodies.

In order to generate a method that easily provide neutralizing antibodies, IAC technology and SPLINT library were modified to generate a novel system named SPLINT-3HY.

Results of the 3-HY selections

In the original format of IAC technology, an antigen-antibody interaction brings the two hybrid proteins together and results in the activation of the reporter genes.

In the new generation of the technology, the three hybrid version, we can identify antibody domains that dissociate or abrogate defined macromolecular interactions.

In order to achieve the proof of principle of this technology we used SPLINT-3HY to undertake the de novo selection of scFv antibody fragments against NF-kB RelA p65 protein.

Mammalian NF-kB is a family of structurally related proteins, including p65/RelA, which form homo- and heterodimers. Regulation of gene expression by NF-kB/rel family of transcription factors is controlled mainly by the inhibitory IkB proteins. The active complex of NF-kB is composed of homodimers and heterodimers of p50, RelA, RelB, and c-Rel. These complexes are sequestered in the cytoplasm by IKBs. Extracellular signal-induced phosphorylation and subsequent degradation of IKBs is essential for nuclear translocation of NF-kB. NF-kB binds to the recognition elements in the promoter region of target genes and activate transcription. IkB proteins determine the duration of transactivation by NF-kB. IKBa is involved in transient activation of NF-kB because it is degraded rapidly upon stimulation and is resynthesized by activated NF-kB. Most of the inducers of NF-kB cause degradation of IKB α . The newly synthesized IKB α sequesters NF-kB in the cytoplasm and determines the signal.

p65 has a bipartite structure-a DNA-binding region and a C-terminal transcription activation domain. These two functional regions are linked by a segment that contains a nuclear transport signal (Chen et al., 2000).

I κ B degradation activates this PKAc that phosphorylates the NF- κ B p65 subunit at Ser 276 rendering NF- κ B that enters the nucleus transcriptionally active through recruitment of CREB binding protein (CBP)/p300 (Zhong et al., 1997; Zhong et al., 1998).

It is believed that p50 homodimers compete for κ B site binding with NF- κ B complexes containing subunits such as p65 that have transactivation domains; however, this is unlikely to be their primary mechanism of repression, as their affinity for classical κ B sites is lower than that of heterodimers such as p50:p65 (Kunsch et al., 1992). One possible mechanism that has been demonstrated for other repressive DNA binding proteins (Glass and Rosenfeld, 2000) is that nuclear p50 may interact with separate inhibitory proteins that recruit corepressor complexes containing HDACs to gene promoters.

The homodimer of p65 protein was cotransformed into yeast cells YI595 with SPLINT/pSPLINT scFv library following the large scale procedure described in (Visintin and Cattaneo, 2001). The co-transformed library was spread onto SG-Raf-WHULK +25mM plates to allow the activation of Gal 1-10 promoters and the selection of the plasmid encoding a scFv that bind one or both p65 interacting proteins due to repression of tetracycline repressor that restores the transcription of His3 gene. The high efficiency of transformation (2.6×10^7) obtained in this experiment ensured that the yield of primary transformants was three times the complexity of the library, thus the library complexity ensures that 95% of the clones represented in the library have been examined (Clarke and Carbon, 1976). After three days 4 clones that grew on -HIS plates were picked and grown on a fresh -HIS plates grid for a second verification of the transcriptional activation of his gene. After 4 days only 2 clones out of the initial 4 clones selected in the first round of selection were able to be rescued on SG-Raf-WHULK +25mM plates. Both clones from secondary screening were picked and grown for the isolation of DNA. Isolated scFv/pSPLINT plasmids were analysed by PCR-BstNI fingerprinting and sequences.

The conclusion from this first part of the experiments is that both scFvs selected from SPLINT-3HY were able to abrogate the interaction between p65-p65 dimer. In order to verify that the two isolated scFvs were also able to recognise p65 as a monomer both antibody fragments were cloned in fusion with VP16 into pLINKER220 vector.

STEP 4: are anti-p65 isolated scFv binders for p65 monomers?

Construction of anti-p65 (A1 and A2)/pLinker220 plasmid. Anti-p65 scFvs fragments were subcloned into pLinker220 between BssHII-NheI restriction sites.

As shown in figure 23 clone A2 display a positive interaction with p65 monomer, thus indicating its specificity with the target protein.

On the contrary, clone A1 was only able to recognized DNA-binding domain lexA.

Although A1 clone result to be specific for lexA protein the result obtained indicates that scFvA1 is probably able to inhibit p65-binding domain or p65-DNA-binding domain in a non conventional manner or by steric hindrance.

scFv A2 was subsequently cloned into mammalian expression vector scFvexcyto-SV5.

STEP 5: in vivo mammalian cells studies

Construction of anti-p65A2/scFvexcyto-SV5 plasmid. Anti-p65 scFvs fragments were subcloned into scFvexcyto-SV5 between BssHII-NheI restriction sites.

Anti-p65 A2 was expressed in HEK cell line as leaderless cytoplasmic protein.

Transfected cells display a diffuse intracellular staining, typical of soluble cytoplasmic protein (see figure 24).

To assess whether anti-p65 A2 inhibition interferes with the physical association between p65 dimer and p50, a coimmunoprecipitation experiment was carried out (Fig. 25). Fig.

25 shows controls (lane 1-3-5- p65 and p50 expressed by HEK in non transfected (c), transfected with anti-p65 (lane 3) and transfected with an irrelevant scFv (lane 5) immunostained with anti-p65 (upper part of the picture) and immunostained with an anti-p50 (lower part). Lanes 4 show the coimmunoprecipitated band corresponding to the p65 monomer (upper part) and p50 protein (lower part). The conclusion for this experiment is that anti-p65 is able to recognize endogenous p65 in HEK cell lines. In this experiment also p50 protein was pulled down with p65 thus indicating that the interaction between anti-p65 and p65 does not interfere with the natural binding sites of p65 and p50.

SPLINT-3HY was also challenge with a panel of interacting pairs (see Table IV). From each selection strategy several neutralizing antibodies were isolated (data not shown) thus indicating that the SPLINT-3HY strategy should provide a robust methodology to elucidate cellular mechanisms in numerous fields, of particular salience in the area of signal transduction. The assessment of gene function may be performed using the tools of protein knock-out including peptide, aptamers and antibodies. These approaches have

been superseded by intracellular antibody technology, which exhibits much more potency for the investigation of protein function than the techniques listed above. In fact they can block or stabilize macromolecular interactions, modulate enzyme function by occluding an active site, sequestering substrate or fixing the enzyme in an active or inactive conformation, they can divert proteins from their usual cellular compartments, target specifically a post-transcriptionally modified version of a given protein (including heterodimerization), they can specifically target the subcellular localization of a given protein, and finally they can selectively target non-protein antigens.

SPLINT-3HY not only provides a good source of stable antibodies but also provides in one step of selection a good source of intracellular and neutralizing antibodies, thus becoming the first antibody-based selection strategy engineered for this aim.

Table IV. Panel of interacting pairs used for SPLINT-3HY selection

<u>Protein ligand x</u>	<u>Protein ligand y</u>
p65 RelA	p65 RelA
p65 RelA	p50 NF- κ B1
Amyloid β aa. 1-42	Amyloid β aa.1-42
JX-TK TrkA	PTB p66 Shc
Synuclein	Synphilin

Consensus sequence framework-based library

A further refinement of the SPLINT library will be carried out with the study of scFv sequences isolated by IACT.

It has been noted that the set of antibodies selected with these technologies is characterized by a number of highly conservative residues. These critical residues represent a sort of trademark that identifies those antibodies capable of being expressed in the intracellular environment by a multitude of scFv which are otherwise unable to function there. IACT technology combined with SPLINT is also able to select the scFv that possess an optimum sequence very similar to a consensus sequence present in the database of immunoglobulins present in nature. As the database of intracellular antibodies selected with this technology grows, the number of conserved residues will decrease. It will therefore be possible to achieve a final generic framework on which an antibody library for intracellular use may be engineered. This will allow the creation of a library

compatible with the low transformability of yeast and, more generally, a phagemid or phagic library of highly stable scFv that are functional for expression in any cellular or noncellular environment. It can be concluded that both IACT and SPLINT are two technologies with the capability to surpass their intrinsic potential and to thus create ever more sophisticated and validated libraries.

Sequences:

URA3 (nutritional marker that encodes the enzyme orotidine-5'phosphate decarboxylase. This enzyme is active in the last step of uracil biosynthesis and is responsible for the conversion of compound C into uracil).

10 ATGTCGAAAGCTACATATAAGGAACGTGCTGCTACTCATCCTAGTCCTGTTGCTGCCAAGCTAT
TTAATATCATGCACGAAAAGCAAACAACTTGTGTGCTTCATTGGATGTTTCGTACCACCAAGG
AATTACTGGAGTTAGTTGAAGCATTAGGTCCCAAATTTGTTTACTAAAAACACATGTGGATA
TCTTGACTGATTTTTCCATGGAGGGCACAGTTAAGCCGCTAAAGGCATTATCCGCCAAGTACA
ATTTTTTACTCTTCGAAGACAGAAAATTTGCTGACATTGGTAATACAGTCAAATTGCAGTACTC
15 TGCGGGTGTATACAGAATAGCAGAATGGGCAGACATTACGAATGCACACGGTGTGGTGGGCC
CAGGTATTGTTAGCGGTTTGAAGCAGGCGGCAGAAGAAGTAACAAAGGAACCTAGAGGCCTT
TTGATGTTAGCAGAATTGTCATGCAAGGGCTCCCTATCTACTGGAGAATATACTAAGGGTACT
GTTGACATTGCGAAGAGCGACAAAGATTTTGTATCGGCTTTATTGCTCAAAGAGACATGGGT
GGAAGAGATGAAGGTTACGATTGGTTGATTATGACACCCGGTGTGGGTTTAGATGACAAGGG
20 AGACGCATTGGGTCAACAGTATAGAACCGTGGATGATGTGGTCTCTACAGGATCTGACATTAT
TATTGTTGGAAGAGGACTATTTGCAAAGGGAAGGGATGCTAAGGTAGAGGGTGAACGTTACA
GAAAAGCAGGCTGGGAAGCATATTTGAGAAGATGCGGCCAGCAAACTAA

TRP1 (nutritional marker that encodes the enzyme N-(5-phosphoribosyl) anthranilate isomerase. This enzyme is found in the next to last step of tryptophan).

25 ATGTCTGTTATTAATTTACAGGTAGTTCTGGTCCATTGGTGAAAGTTTGCGGCTTGCAGAGCA
CAGAGGCCGCAGAATGTGCTCTAGATTCCGATGCTGACTTGCTGGGTATTATATGTGTGCCCA
ATAGAAAGAGAACAATTGACCCGGTTATTGCAAGGAAAATTTCAAGTCTTGTAAGCATATA
AAAATAGTTCAGGCACTCCGAAATACTTGGTTGGCGTGTTTCGTAATCAACCTAAGGAGGATG
TTTTGGCTCTGGTCAATGATTACGGCATTGATATCGTCCAAGTGCATGGAGATGAGTCGTGGCA
30 AGAATACCAAGAGTTCCTCGGTTTGCCAGTTATTAAGAACTCGTATTTCCAAAAGACTGCAA
CATACTACTCAGTGCAGCTTCACAGAAACCTCATTCGTTTATTCCCTTGTTTGATTGAGAAGCA
GGTGGGACAGGTGAACTTTTGGATTGGAAGTCTGATTTCTGACTGGGTGGAAGGCAAGAGAGC
CCCGAAAGCTTACATTTTATGTTAGCTGGTGGACTGACGCCAGAAAATGTTGGTGATGCGCTT
AGATTAAATGGCGTTATTGGTGTGATGTAAGCGGAGGTGTGGAGACAAATGGTGTAAGAA
35 CTCTAACAAAATAGCAAATTTTCGTCAAAAATGCTAAGAAATAG

LEU2 (a nutritionally essential marker that encodes the enzyme β -isopropylamate dehydrogenase. Leu3p is a positive regulator of all 3 genes for the biosynthesis of leucine

and is thought to activate the expression of the genes for isoleucyl-valine ILV2 and ILV5).

ATGTCTGCCCCTATGTCTGCCCCTAAGAAGATCGTCGTTTTGCCAGGTGACCACGTTGGTCAAG
 AAATCACAGCCGAAGCCATTAAGGTTCTTAAAGCTATTTCTGATGTTTCGTTCCAATGTCAAGTT
 5 CGATTTTCGAAAATCATTTAATTGGTGGTGCTGCTATCGATGCTACAGGTGTCCCACTTCCAGAT
 GAGGCGCTGGAAGCCTCCAAGAAGGTTGATGCCGTTTTGTAGGTGCTGTGGCTGGTCCTAAA
 TGGGGTACCGGTAGTGTTAGACCTGAACAAGGTTTACTAAAAATCCGTAAAGAACTTCAATTG
 TACGCCAACTTAAGACCATGTAACCTTGCATCCGACTCTCTTTTAGACTTATCTCCAATCAAGC
 CACAATTTGCTAAAGGTACTGACTTCGTTGTTGTCAGAGAATTAGTGGGAGGTATTTACTTTGG
 10 TAAGAGAAAGGAAGACGATGGTGATGGTGTCGCTTGGGATAGTGAACAATACACCGTTCCAG
 AAGTGCAAAGAATCACAAGAATGGCCGCTTTCATGGCCCTACAACATGAGCCACCATTCGCTA
 TTTGGTCCTTGGATAAAGCTAATCTTTTGGCCTCTTCAAGATTATGGAGAAAACTGTGGAGG
 AAACCATCAAGAACGAATTCCCTACATTGAAGGTTCAACATCAATTGATTGATTCTGCCGCCA
 TGATCCTAGTTAAGAACCCAACCCACCTAAATGGTATTATAATCACCAGCAACATGTTTGGTG
 15 ATATCATCTCCGATGAAGCCTCCGTTATCCCAGGTTCTTGGGTTTGTGTCATCTGCGTCCTTG
 GCCTCTTTGCCAGACAAGAACACCGCATTTGGTTTGTACGAACCATGCCACGGTTCTGCTCCAG
 ATTTGCCAAAGAATAAGGTTGACCCTATCGCCACTATCTTGTCTGCTGCAATGATGTTGAAATT
 GTCATTGAACTTGCCTGAAGAAGGTAAGGCCATTGAAGATGCAGTTAAAAAGGTTTTGGATGC
 AGGTATCAGAACTGGTGATTTAGGTGGTTCCAACAGTACCACCGAAGTCGGTGATGCTGTCGC
 20 CGAAGAAGTTAAGAAAATCCTTGCTTAA

HIS3 (nutritional marker that encodes the enzyme imidazolglycerol phosphate dehydratase. This enzyme is involved in the regulation cycle of histidine biosynthesis).

ATGACAGAGCAGAAAGCCCTAGTAAAGCGTATTACAAATGAAACCAAGATTCAGATTGCGAT
 CTCTTTAAAGGGTGGTCCCCTAGCGATAGAGCACTCGATCTTCCCAGAAAGAGGCAGAAGCAG
 25 TAGCAGAACAGGCCACACAATCGCAAGTGATTAACGTCCACACAGGTATAGGGTTTCTGGACC
 ATATGATACATGCTCTGGCCAAGCATTCCGGCTGGTCGCTAATCGTTGAGTGCAATTGGTGACTT
 ACACATAGACGACCATCACACCACTGAAGACTGCGGGATTGCTCTCGGTCAAGCTTTTAAAGA
 GGCCCTAGGGGGCCGTGCGTGGAGTAAAAAGGTTTGGATCAGGATTTGCGCCTTTGGATGAGGC
 ACTTTCTAGAGCGGTGGTAGATCTTTCGAACAGGCCGTACGCAGTTGTCGAACTTGGTTTGCA
 30 AAGGGAGAAAGTAGGAGATCTCTCTTGCGAGATGATCCCGCATTTTCTTGAAAGCTTTGCAGA
 GGCTAGCAGAATTACCCTCCACGTTGATTGTCTGCGAGGCAAGAATGATCATCACCGTAGTGA
 GAGTGCGTTCAAGGCTCTTGCGGTTGCCATAAGAGAAGCCACCTCGCCCAATGGTACCAACGA
 TGTTCCCTCCACCAAAGGTGTTCTTATGTAG

LYS2 (nutritional marker that encodes the enzyme α -aminoadipic semialdehyde. It is
 35 thought that the product of the reaction controlled by LYS2 acts as a co-inductor together
 with product LYS14 to stimulate the expression of numerous genes involved in the same
 cycle).

ATGACTAACGAAAAGGTCTGGATAGAGAAGTTGGATAATCCAACCTCTTTCAGTGTTACCACAT
GACTTTTTACGCCCACAACAAGAACCTTATACGAAACAAGCTACATATTCGTTACAGCTACCT
CAGCTCGATGTGCCTCATGATAGTTTTTTCTAACAAATACGCTGTCGCTTTGAGTGTATGGGCTG
CATTGATATATAGAGTAACCGGTGACGATGATATTGTTCTTTATATTGCGAATAACAAAATCTT
5 AAGATTCAATATTCAACCAACGTGGTCATTTAATGAGCTGTATTCTACAATTAACAATGAGTTG
AACAAGCTCAATTCTATTGAGGCCAATTTTTCTTTGACGAGCTAGCTGAAAAAATTCAAAGTT
GCCAAGATCTGGAAAGGACCCCTCAGTTGTTCCGTTTGGCCTTTTTGGAAAACCAAGATTTC
AATTAGACGAGTTCAAGCATCATTTAGTGGACTTTGCTTTGAATTTGGATACCAGTAATAATGC
GCATGTTTTGAACTTAATTTATAACAGCTTACTGTATTGGAATGAAAGAGTAACCATTTGTTGCG
10 GACCAATTTACTCAATATTTGACTGCTGCGCTAAGCGATCCATCCAATTGCATAACTAAAATCT
CTCTGATCACCGCATCATCCAAGGATAGTTTACCTGATCCAATAAGAACTTGGGCTGGTGCG
ATTTTCGTGGGGTGTATTCACGACATTTTCCAGGACAATGCTGAAGCCTTCCCAGAGAGAACCT
GTGTTGTGGAGACTCCAACACTAAATTCCGACAAGTCCCGTTCTTTCACTTATCGCGACATCAA
CCGCACTTCTAACATAGTTGCCATTATTTGATTAAACAGGTATCAAAAGAGGTGATGTAGT
15 GATGATCTATTCTTCTAGGGGTGTGGATTTGATGGTATGTGTGATGGGTGTCTTGAAAGCCGGC
GCAACCTTTTCAGTTATCGACCCTGCATATCCCCCAGCCAGACAAACCATTACTTAGGTGTTG
CTAAACCACGTGGGTGATTGTTATTAGAGCTGCTGGACAATTGGATCAACTAGTAGAAGATT
ACATCAATGATGAATTGGAGATTGTTTCAAGAATCAATTCCATCGCTATTCAAGAAAATGGTA
CCATTGAAGGTGGCAAATTGGACAATGGCGAGGATGTTTTGGCTCCATATGATCACTACAAAG
20 ACACCAGAACAGGTGTTGTAGTTGGACCAGATTCCAACCCAACCCTATCTTTCACATCTGGTTC
CGAAGGTATTCCTAAGGGTGTCTTGGTAGACATTTTTCTTGGCTTATTATTTCAATTGGATGT
CCAAAAGGTTCAACTTAACAGAAAATGATAAATTCACAATGCTGAGCGGTATTGCACATGATC
CAATTCAAAGAGATATGTTTACACCATTATTTTTAGGTGCCCAATTGTATGTCCCTACTCAAGA
TGATATTGGTACACCGGGCCGTTTAGCGGAATGGATGAGTAAGTATGGTTGCACAGTTACCCA
25 TTTAACACCTGCCATGGGTCAATTACTTACTGCCCAAGCTACTACACCATTCCCTAAGTTACAT
CATGCGTTCTTTGTGGGTGACATTTTAACAAAACGTGATTGTCTGAGGTTACAAACCTTGGCAG
AAAATTGCCGTATTGTTAATATGTACGGTACCACTGAAACACAGCGTGCAGTTTCTTATTTCTGA
AGTTAAATCAAAAAATGACGATCCAAACTTTTTGAAAAAATTGAAAGATGTCATGCCTGCTGG
TAAAGGTATGTTGAACGTTTACGCTACTAGTTGTTAACAGGAACGATCGTACTCAAATATGTGG
30 TATTGGCGAAATAGGTGAGATTTATGTTTCGTGCAGGTGGTTTGGCCGAAGGTATAGAGGATT
ACCAGAATTGAATAAAGAAAAATTTGTGAACAACCTGGTTTGTGAAAAAGATCACTGGAATTA
TTTGGATAAGGATAATGGTGAACCTTGGAGACAATTCTGGTTAGGTCCAAGAGATAGATTGTA
CAGAACGGGTGATTTAGGTCGTTATCTACCAAACGGTGACTGTGAATGTTGCGGTAGGGCTGA
TGATCAAGTTAAAATTCGTGGGTTCAGAATCGAATTAGGAGAAATAGATACGCACATTTCCCA
35 ACATCCATTGGTAAGAGAAAACATTACTTTAGTTTCGCAAAAATGCCGACAATGAGCCAACATT
GATCACATTTATGGTCCCAAGATTTGACAAGCCAGATGACTTGTCTAAGTTCCAAAGTGATGTT
CCAAAGGAGGTTGAAACTGACCCTATAGTTAAGGGCTTAATCGGTACCATCTTTTATCCAAG
GACATCAGGACTTTCTTAAAGAAAAGATTGGCTAGCTATGCTATGCCTTCCTTGATTGTGGTTA
TGGATAAACTACCATTGAATCCAAATGGTAAAGTTGATAAGCCTAAACTTCAATTCCCAACTC
40 CCAAGCAATTAAATTTGGTAGCTGAAAATACAGTTTCTGAAACTGACGACTCTCAGTTTACCA

ATGTTGAGCGCGAGGTTAGAGACTTATGGTTAAGTATATTACCTACCAAGCCAGCATCTGTAT
 CACCAGATGATTCGTTTTTCGATTTAGGTGGTCATTCTATCTTGGCTACCAAAATGATTTTTACC
 TTAAAGAAAAAGCTGCAAGTTGATTTACCATTTGGGCACAATTTTCAAGTATCCAACGATAAAG
 GCCTTTGCCGCGGAAATTGACAGAATTAAATCATCGGGTGGATCATCTCAAGGTGAGGTCGTC
 5 GAAAATGTCACTGCAAATTATGCGGAAGACGCCAAGAAATTGGTTGAGACGCTACCAAGTTC
 GTACCCCTCTCGAGAATATTTTGTGTAACCTAATAGTGCCGAAGGAAAAACAACAATTAATGT
 GTTTGTTACCGGTGTCACAGGATTTCTGGGCTCCTACATCCTTGCAGATTTGTTAGGACGTTCT
 CCAAAGAAGTACAGTTTCAAAGTGTGTTGCCACGTCAGGGCCAAGGATGAAGAAGCTGCATTT
 GCAAGATTACAAAAGGCAGGTATCACCTATGGTACTTGGAACGAAAAATTTGCCTCAAATATT
 10 AAAGTTGTATTAGGCGATTTATCTAAAAGCCAATTTGGTCTTTCAGATGAGAAGTGGATGGAT
 TTGGCAAACACAGTTGATATAATTATCCATAATGGTGCGTTAGTTCAGTGGGTTTATCCATATG
 CCAAATTGAGGGATCCAAATGTTATTTCAACTATCAATGTTATGAGCTTAGCCGCCGTCGGCA
 AGCCAAAGTTCTTTGACTTTGTTTCCTCCACTTCTACTCTTGACACTGAATACTACTTTAATTTG
 TCAGATAAACTTGTTAGCGAAGGGAAGCCAGGCATTTTAGAATCAGACGATTTAATGAACTCT
 15 GCAAGCGGGCTCACTGGTGGATATGGTCAGTCCAAATGGGCTGCTGAGTACATCATTAGACGT
 GCAGGTGAAAGGGGCCTACGTGGGTGTATTGTCAGACCAGGTACGTAACAGGTGCCTCTGCC
 AATGGTTCTTCAAACACAGATGATTTCTTATTGAGATTTTTGAAAGGTTTCAGTCCAATTAGGTA
 AGATTCCAGATATCGAAAATTCCGTGAATATGGTTCCAGTAGATCATGTTGCTCGTGTTGTTGT
 TGCTACGTCTTTGAATCCTCCCAAAGAAAATGAATTGGCCGTTGCTCAAGTAACGGGTACCCC
 20 AAGAATATTATTCAAAGACTACTTGTATACTTTACACGATTATGGTTACGATGTCGAAATCGA
 AAGCTATTCTAAATGGAAGAAATCATTGGAGGCGTCTGTTATTGACAGGAATGAAGAAAATGC
 GTTGTATCCTTTGCTACACATGGTCTTAGACAACCTACCTGAAAGTACCAAAGCTCCGGAACCTA
 GACGATAGGAACGCCGTGGCATCTTTAAAGAAAGACACCGCATGGACAGGTGTTGATTGGTCT
 AATGGAATAGGTGTTACTCCAGAAGAGGTTGGTATATATATTGCATTTTTTAAACAAGGTTGGA
 25 TTTTACCTCCACCAACTCATAATGACAACTTCCACTGCCAAGTATAGAACTAACTCAAGCGC
 AAATAAGTCTAGTTGCTTCAGGTGCTGGTGCTCGTGGAAGCTCCGCAGCAGCTTA
KAN (antibiotic resistance: it is an aminoglycosidase that acts on protein synthesis by
 binding to the ribosome subunit 30S to prevent the start of the complex with 'mRNA. It
 can also cause a messenger misreading, leading to the formation of a nonsense peptide.
 30 Another important function of aminoglycosidase is to increase membrane permeability).
 ATGATTGAACAAGATGGATTGCACGCAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTATTCGGC
 TATGACTGGGCACAACAGACAATCGGCTGCTCTGATGCCGCCGTGTTCCGGCTGTCAGCGCAG
 GGGCGCCCGGTTCTTTTTGTCAAGACCGACCTGTCCGGTGCCCTGAATGAACTCCAAGACGAG
 GCAGCGCGGCTATCGTGGCTGGCCACGACGGGCGTTCCTTGCGCAGCTGTGCTCGACGTTGTC
 35 ACTGAAGCGGGAAGGGACTGGCTGCTATTGGGCGAAGTGCCGGGGCAGGATCTCCTGTCATCT
 CACCTTGCTCCTGCCGAGAAAGTATCCATCATGGCTGATGCAATGCGGCGGCTGCATACGCTT
 GATCCGGCTACCTGCCCATTCGACCACCAAGCGAAACATCGCATCGAGCGAGCACGTACTCGG
 ATGGAAGCCGGTCTTGTCGATCAGGATGATCTGGACGAAGAGCATCAGGGGCTCGCGCCAGC
 CGAACTGTTCCGCCAGGCTCAAGGCGCGGATGCCCGACGGCGAGGATCTCGTCGTGACCCATGG

CGATGCCTGCTTGCCGAATATCATGGTGGAAAATGGCCGCTTTTCTGGATTCATCGACTGTGGC
CGGCTGGGTGTGGCGGACCGCTATCAGGACATAGCGTTGGCTACCCGTGATATTGCTGAAGAG
CTTGGCGGCGAATGGGCTGACCGCTTCCTCGTGCTTTACGGTATCGCCGCTCCCGATTTCGCAGC
GCATCGCCTTCTATCGCCTTCTTGACGAGTTCTTCTGA

- 5 BLA (β -lactamase: antibiotic resistance. The product of this gene inhibits certain enzymes involved in bacterial cell wall synthesis. β -lactamase hydrolyzes the β -lactame contained in the antibiotic ampicillin).

ATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTGTTTTT
GCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGG
10 TTACATCGAGCTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTTCGCCCCGAAGAACGTTT
TCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGTGCGGTATTATCCCGTGTTGACGCCGGG
CAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTC
ACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCAT
GAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCG
15 CTTTTTTGCACAACATGGGGGATCATGTAACCCGCCTTGATCGTTGGGAACCGGAGCTGAATG
AAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTTGCGC
AACTATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAG
GCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATA
AATCTGGAGCCGGTGAGCGTGGATCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGC
20 CCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGAC
AGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAA

CAM (antibiotic resistance: the product of this gene acts by binding to the ribosome subunit 50S, thus blocking the formation of peptide binding and inhibiting the activity of peptidyl transferase. It is a potent inhibitor of protein synthesis in eukaryotic cells).

ATGGAGAAAAAATCACTGGATATACCACCGTTGATATATCCCAATGGCATCGTAAAGAACAT
TTTGAGGCATTTTCAGTCAGTTGCTCAATGTACCTATAACCAGACCGTTCAGCTGGATATTACGG
CCTTTTTAAAGACCGTAAAGAAAAATAAGCACAAGTTTTATCCGGCCTTTATTCACATTCTTGC
CCGCCTGATGAATGCTCATCCGGAATTTTCGTATGGCAATGAAAGACGGTGAGCTGGTGATATG
GGATAGTGTTACCCCTTGTTACACCGTTTTCCATGAGCAAACCTGAAACGTTTTTCATCGCTCTGG
30 AGTGAATACCACGACGATTTCCGGCAGTTTCTACACATATATTCGCAAGATGTGGCGTGTTAC
GGTGAAAACCTGGCCTATTTCCCTAAAGGGTTTATTGAGAATATGTTTTTCGTCTCAGCCAATC
CCTGGGTGAGTTTCACCAGTTTTGATTAAACGTGGCCAATATGGACAACTTCTTCGCCCCCGT
TTTACCATGGGCAAATATTATACGCAAGGCGACAAGGTGCTGATGCCGCTGGCGATTTCAGGT
TCATCATGCCGTTTGTGATGGCTTCCATGTCGGCAGAATGCTTAATGAATTACAACAGTACTGC
35 GATGAGTGGCAGGGCGGGGCGTAA

Tet R (antibiotic resistance: inhibits the binding of aminoacyl tRNA to the ribosome subunit 30S in bacteria).

ATAGTGATGGCACGACTAAGCTTGGACGACGTAATTTCAATGGCGCTCACCTGCTGGACAGC
 GAAGGGCTAGAGGGCTTGACTACGCGTAAGCTGGCGCAGTCCCTAAAAATTGAGCAACCGAC
 TCTGTATTGGCACGTGCGCAACAAGCAGACTCTTATGAACATGCTTTCAGAGGCAATACTGGC
 GAAGCATCACACCCGTTTCAGCACCGTTACCGACTGAGAGTTGGCAGCAGTTTCTCCAGGAAAA
 5 TGCTCTGAGTTTCCGTAAAGCATTACTGGTCCATCGTGATGGAGCCCGATTGCATATAGGGAC
 CTCTCCTACGCCCCCCCCAGTTTGAACAAGCAGAGGGCGCAACTACGCTGTCTATGCGATGCAGG
 GTTTTCGGTCGAGGAGGCTCTTTTCATTCTGCAATCTATCAGCCATTTTACGTTGGGTGCAGTA
 TTAGAGGAGCAAGCAACAAACCAGATAGAAAATAATCATGTGATAGACGCTGCACCACCATT
 ATTACAAGAGGCATTTAATATTCAGGCGAGAACCTCTGCTGAAATGGCCTTCCATTTTCGGGCT
 10 GAAATCATTAAATATTTGGATTTTCTGCACAGTTAGATGAAAAAAGCATACACCCATTGAGGA
 TGGTAATAAATGA

G418-Neomycin (antibiotic resistance. It is an aminoglycoside phosphotransferase. It is used in eukaryotic, mammalian and yeast vectors).

ATGAGCCATATTCAACGGGAAACGTCTTGCTCGAGGCCGCGATTAAATTCCAACATGATGCTG
 15 ATTTATATGGGTATAAATGGGCTCGCGATAATGTCGGGCAATCAGGTGCACAATCTATCGATT
 GTATGGGAAGCCCGATGCGCCAGAGTTGTTTCTGAAACATGGAAAGGTAGCGTTGCCAATGAT
 GTTACAGATGAGATGGTCAGACTAACTGGCTGACGAATTTATGCCTCTTCCGACCATCAAGC
 ATTTTATCCGTACTCCTGATGATGCATGGTACTCACCCTGCGATCCCCGGGAAAACAGCATTC
 CAGGTATTAGAAGAATATCCTGATCAGGTGAAAATATTGTTGATGCGCTGGCAGTGTTCCCTGC
 20 GCCGGTTGCATTCGATCCTGTTTGTAATTGTCCTTTTAAACAGCGATCGCGTATTTTCGTCTCGCTC
 AGGCGCAACACGAATGAATAACGGTTTGGTTGATGCGAGTGATTTTGATGACGAGCGTAATGG
 CGGCCTGTTGAACAAGTCTGGAAAGAAATGCATAAGCTTTTGCCATTCTCACCGGATTAGTCG
 TCACTCATGGTGATTTCTCACTTGATAACCTTATTTTTGACGAGGGGAAATTATAGGTTGTATT
 GATGTTGGACGAGTCGGAATCGCAGACCGATACCAGGATCTTGCCTCCTATGGAAGTGCCTCG
 25 GTGAGTTTTCTCCTTCATTACAGAAACGGCTTTTTCAAATATGGTATTGATAATCCTGATATG
 AATAAATTGCAGTTTCATTTGATGCTCGATGAGTTTTCTA

ColE1 (origin of replication in E.coli)

CTGTCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAACTTCATTTTTAATTTAAAA
 GGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCGTT
 30 CCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGC
 GTAATCTGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCAA
 GAGCTACCAACTCTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATACTGTC
 CTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACATACCTCG
 CTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGA
 35 CTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGCTGAACGGGGGGTTTCGTGCACAC
 AGCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAA
 AGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAAC
 AGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTGCGGGTT
 TCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGGCGGAGCCTATGGAA

AAACGCCAGCAACGCGGCCTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCACATGTTC
TTTCCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCG
CTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGA

2 μ (origin of replication in *S.cerevisiae*)

5 CATTTAAGCATAAACACGCACTATGCCGTTCTTCTCATGTATATATATACAGGCAACACGCA
GATATAGGTGCGACGTGAACAGTGAGCTGTATGTGCGCAGCTCGCGTTGCATTTTCGGAAGCG
CTCGTTTTTCGGAACGCTTTGAAGTTCCTATTCCGAAGTTCCTATTCTCTAGCTAGAAAGTATA
GGAAC TTCAGAGCGCTTTTGAAAACCAAAAGCGCTCTGAAGACGCACTTTCAAAAACCAAA
AACGCACCGGACTGTAACGAGCTACTAAAATATTGCGAATACCGCTTCCACAAACATTGCTCA
10 AAAGTATCTCTTTGCTATATATCTCTGTGCTATATCCCTATATAACCTACCCATCCACCTTTCGC
TCCTTGAAC TTGCATCTAAACTCGACCTCTACATTTTTTATGTTTATCTCTAGTATTACTCTTTA
GACAAAAAATTGTAGTAAGAACTATTCATAGAGTGAATCGAAAACAATACGAAAATGTAAA
CATTTCTTATACGTAGTATATAGAGACAAAATAGAAGAAACCGTTCATAATTTTCTGACCAAT
GAAGAATCATCAACGCTATCACTTTCTGTTCACAAAGTATGCGCAATCCACATCGGTATAGAA
15 TATAATCGGGGATGCCTTTATCTTGAAAAAATGCACCCGCAGCTTCGCTAGTAATCAGTAAAC
GCGGGAAGTGAGTCAGGCTTTTTTTATGGAAGAGAAAATAGACACCAAAGTAGCCTTCTTCT
AACCTTAACGGACCTACAGTGCAAAAAGTTATCAAGAGACTGCATTATAGAGCGCACAAAGG
AGAAAAAAGTAATCTAAGATGCTTTGTTAGAAAAATAGCGCTCTCGGGATGCATTTTGTAG
AACAAAAAAGAAGTATAGATTCTTTGTTGGTAAAATAGCGCTCTCGCGTTGCATTTCTGTTCTG
20 TAAAAATGCAGCTCAGATTCTTTGTTTGAAAAATTAGCGCTCTCGCGTTGCATTTTGTGTTTAC
AAAAATGAAGCACAGATTCTTCGTTGGTAAAATAGCGCTTTCGCGTTGCATTTCTGTTCTGTAA
AAATGCAGCTCAGATTCTTTGTTTGAAAAATTAGCGCTCTCGCGTTGCATTTTGTGTTCTACAAA
ATGAAGCACAGATGCTTCGTT

LexA (DNA-binding domain)

25 ATGAAAGCGTTAACGGCCAGGCAACAAGAGGTGTTTGATCTCATCCGTGATCACATCAGCCAG
ACAGGTATGCCGCCGACGCGTGCGGAAATCGCGCAGCGTTTGGGGTTCCGTTCCCCAAACGCG
GCTGAAGAACATCTGAAGGCGCTGGCACGCAAAGGCGTTATTGAAATTGTTTCCGGCGCATCA
CGCGGGATTCTGTTGTCAGGAAGAGGAAGAAGGGTTGCCGCTGGTAGGTCGTGTGGCTGCC
GGTGAACCACTTCTGGCGCAACAGCATATTGAAGGTCATTATCAGGTCGATCCTTCCTTATTCA
30 AGCCGAATGCTGATTTCTGCTGCGCGTCAGCGGGATGTCGATGAAAGATATCGGCATTATGG
ATGGTGACTTGCTGGCAGTGCATAAACTCAGGATGTACGTAACGGTCAGGTCGTTGTCGCAC
GTATTGATGACGAAGTTACCGTTAAGCGCCTGAAAAACAGGGGCAATAAAGTCGAACTGTTGC
CAGAAAATAGCGAGTTTAAACCAATTGTCGTTGACCTTCGTCAGCAGAGCTTCACCATTGAAG
GGCTGGCGGTTGGGGTTATTCGCAACGGCGACTGGCTG

35 VP16 (activation domain)

CGGCCCCCCCCGACCGATGTCAGCCTGGGGGACGAGCTCCACTTAGACGGCGAGGACGTGGCG
ATGGCGCATGCCGACGCGCTAGACGATTTTCGATCTGGACATGTTGGGGGACGGGGATTCCCCG
GGGCCGGGATTTACCCCCACGACTCCGCCCCCTACGGCGCTCTGGATATGGCCGACTTCGAG
TTTGAGCAGATGTTTACCGATGCCCTTGGAATTGACGAGTACGGTGGG

B42 (activation domain)

ATGGGTGCTCCTCCAAAAAGAAGAGAAAGGTAGCTGGTATCAATAAAGATATCGAGGAGTG
CAATGCCATCATTGAGCAGTTTATCGACTACCTGCGCACCGGACAGGAGATGCCGATGGAAAT
GGCGGATCAGGCGATTAACGTGGTGCCGGGCATGACGCCGAAAACCATTCCTTCACGCCGGGCC
5 GCCGATCCAGCCTGACTGGCTGAAATCGAATGGTTTTTCATGAAATTGAAGCGGATGTTAACGA
TACCAGCCTCTTGCTGAGTGGAGATGCCTCCTACCCTTATGA

ADH1 prom (promoter)

ATGAAAGCGTTAACGGCCAGGCAACAAGAGGTGTTTGATCTCATCCGTGATCACATCAGCCAG
ACAGGTATGCCGCCGACGCGTGCGGAAATCGCGCAGCGTTTGGGGTTCCGTTCGCCAAACGCG
10 GCTGAAGAACATCTGAAGGCGCTGGCACGCAAAGGCGTTATTGAAATTGTTTCCGGCGCATCA
CGCGGATTCGTCTGTTGCAGGAAGAGGAAGAAGGGTTGCCGCTGGTAGGTCGTGTGGCTGCCG
GTGAACCACTTCTGGCGCAACAGCATATTGAAGGTCATTATCAGGTCGACCTTCCTTATTCAAG
CCGAATGCTGATTTCTTGCTGCGCGTCAGCGGGATGTGATGAAAGATATCGGCATTATGGAT
GGTGACTTGCTGGCAGTGCATAAACTCAGGATGTACGTAACGGTCAGGTCGTTGTCGCACGT
15 ATTGATGACGAAGTTACCGTTAAGCGCCTGAAAAACAGGGCAATAAAGTCGAAGTGTGCC
AGAAAATAGCGAGTTTAAACCAATTGTCGTAGATCTTCGTCAGCAGAGCTTCACCATTGAAGG
GCTGGCGGTTGGGGTTATTCGCAACGGCGACTGGCTGGAATTCCCGGGGATCCGTCGACCATG
GCGGCCGCTCGAGTCGACCTGCAGCCAAGCTAATTCCGGGCGAATTTCTTATGATTTATGA

GAL1-10 bi-directional prom (inducible bi-directional promoter)

GGATCCCCTTTCAAAAATTCTTACTTTTTTTTTTGGATGGACGCAAAGAAGTTTAATAATCATAT
TACATGGCATTACCACCATATACATATCCATATACATATCCATATCTAATCTTACTTATATGTT
GTGGAAATGTAAAGAGCCCCATTATCTTAGCCTAAAAAAACCTTCTCTTTGGAACCTTTCAGTA
ATACGCTTAACTGCTCATTGCTATATTGAAGTACGGATTAGAAGCCGCCGAGCGGGTGACAGC
CCTCCGAAGGAAGACTCTCCTCCGTGCGTCCTCGTCTTCACCGGTCGCGTTCCTGAAACGCAGA
25 TGTGCCTCGCGCCGCACTGCTCCGAACAATAAAGATTCTACAATACTAGCTTTTATGGTTATGA
AGAGGAAAAATTGGCAGTAACCTGGCCCCACAAACCTTCAAATGAACGAATCAAATTAACAA
CCATAGGATGATAATGCGATTAGTTTTTTAGCCTTATTTCTGGGGTAATTAATCAGCGAAGCGA
TGATTTTTGATCTATTAACAGATATATAAATGCAAAAACCTGCATAACCACTTTAACTAATACTT
TCAACATTTTCGGTTTGTATTACTTCTTATTCAAATGTAATAAAAGTATCAACAAAAAATTGTT
30 AATATACCTCTATACTTTAACGTCAAGGAGAAAAAACCCGGG

GAL1 prom (inducible promoter)

ACGGATTAGAAGCCGCCGAGCGGGTGACAGCCCTCCGAAGGAAGACTCTCCTCCGTGCGTCCT
CGTCTTCACCGGTCGCGTTCCTGAAACGCAGATGTGCCTCGCGCCGCACTGCTCCGAACAATA
AAGATTCTACAATACTAGCTTTTATGGTTATGAAGAGGAAAAATTGGCAGTAACCTGGCCCCA
35 CAAACCTTCAAATGAACGAATCAAATTAACAACCATAGGATGATAATGCGATTAGTTTTTTAG
CCTTATTTCTGGGGTAATTAATCAGCGAAGCGATGATTTTTGATCTATTAACAGATATATAAAT
GCAAAAACCTGCATAACCACTTTAACTAATACTTTCAACATTTTCGGTTTGTATTACTTCTTATTC
AAATGTAATAAAAGTATCAACAAAAAATTGTTAATATACCTCTATACTTTAACGTCAAGGAGA
AAAAACC

ADH1 term (transcription stop sequence)

AAGCTTTTCATAGGGTAGGGGAATTTCTGGGCATGCCGGTAGAGGTGTGGTCAATAAGAGCGAC
CTCATGCTATACCTGAGAAAGCAACCTGACCTACAGGAAAGAGTTACTCAAGAACAAGAATTT
TCGTTTTAAACCTAAGAGTCACTTTAAAATTTGTATACACTTATTTTTTTTATAACTTATTTAA
5 TAATAAAAATCATAAATCATAAGAAATTCGCTTATTTAGAAAGTGTCAACAACGTATCTACCAA
CGATTTGACCCTTTTCCATCTTTTCGTAAATTTCTGGCAAGGTAGACAAGCCGACAACCTTGAT
TGGAGACTTGACCAAACCTCTGGCGAAGAA GTCCA

ADH2 term (transcription stop sequence)

GAATTCGACACTTCTAAATAAGCGGATCTCCTATGCCTTCACGATTTATAGTTTCCATTATCAA
10 GTATGCCTATATTAGTATATAGCATCTTTAGATGACAGTGTTCGAAGTTTCACGAATAAAAGAT
AATATTCTACTTTTTGCTCCCACCGCGTTTGCTAGCACGAGTGAACACCATCCCTCGCCTGTGA
GTTGTACCCATTCTCTAAACTGTAGACATGGTAGCTTCAGCAGTGTTCGTTATGTACGGCATC
CTCCAACAAACAGTCGGTTATAGTTTGTCTGCTCCTCTGAATCGTCTCCCTCGATATTTCTCAT
TTTCCTTCGCATGCCCTCGAG

15 CRE GENE

ATGTCCAATTTACTGACCGTACACCAAAATTTGCCTGCATTACCGGTCGATGCAACGAGTGAT
GAGGTTTCGCAAGAACCTGATGGACATGTTTCAGGGATCGCCAGGCGTTTTCTGAGCATACCTGG
AAAATGCTTCTGTCCGTTTGCCGGTCGTGGGCGGCATGGTGCAAGTTGAATAACCGGAAATGG
TTTCCCGCAGAACCTGAAGATGTTTCGCGATTATCTTCTATATCTTCAGGCGCGCGGTCTGGCAG
20 TAAAAACTATCCAGCAACATTTGGGCCAGCTAAACATGCTTCATCGTCGGTCCGGGCTGCCAC
GACCAAGTGACAGCAATGCTGTTTCACTGGTTATGCGGCGGATCCGAAAAGAAAACGTTGATG
CCGGTGAACGTGCAAAACAGGCTCTAGCGTTCGAACGCACTGATTTTCGACCAGGTTTCGTTTAC
TCATGGAAAATAGCGATCGCTGCCAGGATATACGTAATCTGGCATTCTGTTGGGGATTGCTTATA
ACACCCTGTTACGTATAGCCGAAATTGCCAGGATCAGGGTTAAAGATATCTCACGTACTGACG
25 GTGGGAGAATGTTAATCCATATTGGCAGAACGAAAACGCTGGTTAGCACCGCAGGTGTAGAG
AAGGCACTTAGCCTGGGGGTAACTAAACTGGTCGAGCGATGGATTTCCGTCTCTGGTGTAGCT
GATGATCCGAATAACTACCTGTTTTTGCCGGGTCAGAAAAAATGGTGTTGCCGCGCCATCTGCC
ACCAGCCAGCTATCAACTCGCGCCCTGGAAGGGATTTTTGAAGCAACTCATCGATTGATTAC
GGCGCTAAGGATGACTCTGGTCAGAGATACCTGGCCTGGTCTGGACACAGTGCCCGTGTCGGA
30 GCCGCGCGAGATATGGCCCGCGCTGGAGTTTCAATACCGGAGATCATGCAAGCTGGTGGCTGG
ACCAATGTAAATATTGTCATGAACTATATCCGTACCCTGGATAGTGAAACAGGGGCAATGGTG
CGCCTGCTGGAAGATGGCGATTAG

PolyLINKER pMIC-BD1 and pMIC-BD2 and pBTM116

GAATTCCCGGGGATCCGTCGACCTGCAG

35 POLYLINKER pVP16

GGCCCAGCCGGCCGTCGACGCGGCCGCTCGAG

POLYLINKERS plinker220 and pVP16mv1

GCGCGCATGCC....(VL)....TCCGGAGGGTTCGACCAGCGGTTCTGGGAAACCAGGTTCCGGTGAA
 GGCTCGAGCGGTACC....(VH)....GCTAGCGGCAAACCAATCCCAAACCCACTGCTGGGCCTGGA
 TAGTACT

REFERENCES

- 5 Baird, G.S., Zacharias, D.A. and Tsien, R.Y. (1999) Circular permutation and receptor insertion within green fluorescent proteins. *Proc Natl Acad Sci U S A* 96, 11241-6.
- Bartel, P.L., Chien, C., Sternglanz, R. and Fields, S. (1993) Using the two-hybrid system to detect protein-protein interaction. In: D.A. Hartley (Ed) *Cellular interaction in development: a practical approach*. IRL Press, Oxford, p. 153-179.
- 10 Biocca, S., Neuberger, M.S. and Cattaneo, A. (1990) Expression and targeting of intracellular antibodies in mammalian cells. *Embo J* 9, 101-8.
- Biocca, S., Ruberti, F., Tafani, M., Pierandrei-Amaldi, P. and Cattaneo, A. (1995) Redox state of single chain Fv fragments targeted to the endoplasmic reticulum, cytosol and mitochondria. *Biotechnology (N Y)* 13, 1110-5.
- 15 Capecchi, M.R. (1989) Altering the genome by homologous recombination. *Science* 244, 1288-92.
- Cattaneo, A. and Biocca, S. (1997) *Intracellular Antibodies: Development and Applications*.
- Cattaneo, A. and Biocca, S. (1999) The selection of intracellular antibodies. *Trends in*
- 20 *Biotechnology* 17, 115-121.
- Chen, Y.Q., Sengchanthalangsy, L.L., Hackett, A. and Ghosh, G. (2000) NF-kappaB p65 (RelA) homodimer uses distinct mechanisms to recognize DNA targets. *Structure Fold Des* 8, 419-28.
- Clarke, L. and Carbon, J. (1976) A colony bank containing synthetic Col El hybrid
- 25 plasmids representative of the entire E. coli genome. *Cell* 9, 91-9.
- Engberg, J., Andersen, P.S., Nielsen, L.K., Dziegiel, M., Johansen, L.K. and Albrechtsen, B. (1996) Phage-display libraries of murine and human antibody Fab fragments. *Mol Biotechnol* 6, 287-310.
- Feldhaus, M.J., Siegel, R.W., Opresko, L.K., Coleman, J.R., Feldhaus, J.M., Yeung, Y.A.,
- 30 Cochran, J.R., Heinzelman, P., Colby, D., Swers, J., Graff, C., Wiley, H.S. and Wittrup, K.D. (2003) Flow-cytometric isolation of human antibodies from a nonimmune *Saccharomyces cerevisiae* surface display library. *Nat Biotechnol* 21, 163-70.
- Fire, A. (1999) RNA-triggered gene silencing. *Trends Genet* 15, 358-63.

- Glass, C.K. and Rosenfeld, M.G. (2000) The coregulator exchange in transcriptional functions of nuclear receptors. *Genes Dev* 14, 121-41.
- Griffiths, A.D., Malmqvist, M., Marks, J.D., Bye, J.M., Embleton, M.J., McCafferty, J., Baier, M., Holliger, K.P., Gorick, B.D., Hughes-Jones, N.C. and et al. (1993) Human anti-self antibodies with high specificity from phage display libraries. *Embo J* 12, 725-34.
- Griffiths, A.D., Williams, S.C., Hartley, O., Tomlinson, I.M., Waterhouse, P., Crosby, W.L., Kontermann, R.E., Jones, P.T., Low, N.M., Allison, T.J. and et al. (1994) Isolation of high affinity human antibodies directly from large synthetic repertoires. *Embo J* 13, 3245-60.
- Hollenberg, S.M., Sternglanz, R., Cheng, P.F. and Weintraub, H. (1995) Identification of a new family of tissue -specific basic helix-loop-helix proteins with a two-hybrid system. *Molecular and Cellular Biology* 15, 3813-3822.
- Kortt, A.A., Dolezal, O., Power, B.E. and Hudson, P.J. (2001) Dimeric and trimeric antibodies: high avidity scFvs for cancer targeting. *Biomol Eng* 18, 95-108.
- Kunsch, C., Ruben, S.M. and Rosen, C.A. (1992) Selection of optimal kappa B/Rel DNA-binding motifs: interaction of both subunits of NF-kappa B with DNA is required for transcriptional activation. *Mol Cell Biol* 12, 4412-21.
- Orlandi, R., Gussow, D.H., Jones, P.T. and Winter, G. (1992) Cloning immunoglobulin variable domains for expression by the polymerase chain reaction. 1989. *Biotechnology* 24, 527-31.
- Persic, L., Righi, M., Roberts, A., Hoogenboom, H.R., Cattaneo, A. and Bradbury, A. (1997) Targeting vectors for intracellular immunisation. *Gene* 187, 1-8.
- Prasher, D.C., Eckenrode, V.K., Ward, W.W., Prendergast, F.G. and Cormier, M.J. (1992) Primary structure of the *Aequorea victoria* green-fluorescent protein. *Gene* 111, 229-33.
- Proba, K., Honegger, A. and Pluckthun, A. (1997) A natural antibody missing a cysteine in VH: consequences for thermodynamic stability and folding. *J Mol Biol* 265, 161-72.
- Proba, K., Worn, A., Honegger, A. and Pluckthun, A. (1998) Antibody scFv fragments without disulfide bonds made by molecular evolution. *J Mol Biol* 275, 245-53.
- Rhee, Y., Gurel, F., Gafni, Y., Dingwall, C. and Citovsky, V. (2000) A genetic system for detection of protein nuclear import and export. *Nat Biotechnol* 18, 433-7.
- Rossi, J.J. (1999) Ribozymes, genomics and therapeutics. *Chem Biol* 6, R33-7.

- Sblattero, D. and Bradbury, A. (2000) Exploiting recombination in single bacteria to make large phage antibody libraries. *Nat Biotechnol* 18, 75-80.
- Sheets, M.D., Amersdorfer, P., Finnern, R., Sargent, P., Lindquist, E., Schier, R., Hemingsen, G., Wong, C., Gerhart, J.C., Marks, J.D. and Lindqvist, E. (1998) Efficient construction of a large nonimmune phage antibody library: the production of high-affinity human single-chain antibodies to protein antigens (Sheets et al.). *Proc Natl Acad Sci U S A* 95, 6157-62.
- Siegel, M.S. and Isacoff, E.Y. (1997) A genetically encoded optical probe of membrane voltage. *Neuron* 19, 735-41.
- 10 Tanaka, T., Chung, G.T., Forster, A., Lobato, M.N. and Rabbitts, T.H. (2003) De novo production of diverse intracellular antibody libraries. *Nucleic Acids Res* 31, e23.
- Tavernarakis, N., Wang, S.L., Dorovkov, M., Ryazanov, A. and Driscoll, M. (2000) Heritable and inducible genetic interference by double-stranded RNA encoded by transgenes. *Nat Genet* 24, 180-3.
- 15 Todorovska, A., Roovers, R.C., Dolezal, O., Kortt, A.A., Hoogenboom, H.R. and Hudson, P.J. (2001) Design and application of diabodies, triabodies and tetrabodies for cancer targeting. *J Immunol Methods* 248, 47-66.
- Tomlinson, I. and Holliger, P. (2000) Methods for generating multivalent and bispecific antibody fragments. *Methods Enzymol* 326, 461-79.
- 20 Uetz, P., Giot, L., Cagney, G., Mansfield, T.A., Judson, R.S., Knight, J.R., Lockshon, D., Narayan, V., Srinivasan, M., Pochart, P., Qureshi-Emili, A., Li, Y., Godwin, B., Conover, D., Kalbfleisch, T., Vijayadamodar, G., Yang, M., Johnston, M., Fields, S. and Rothberg, J.M. (2000) A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*. *Nature* 403, 623-7.
- 25 Vaughan, T.J., Williams, A.J., Pritchard, K., Osbourn, J.K., Pope, A.R., Earnshaw, J.C., McCafferty, J., Hodits, R.A., Wilton, J. and Johnson, K.S. (1996) Human antibodies with sub-nanomolar affinities isolated from a large non-immunized phage display library (Uetz et al.). *Nat Biotechnol* 14, 309-14.
- 30 Visintin, M. and Cattaneo, A. (2001) Selecting intracellular antibodies using the two-hybrid system. In: R. Kontermann, & Dubel, S., eds (Ed) *Antibody Engineering*, Vol. 1, Springer Lab Manual. Springer, Heidelberg, Germany, p. 790.

- Visintin, M., Settanni, G., Maritan, A., Graziosi, S., Marks, J.D. and Cattaneo, A. (2002) The intracellular antibody capture technology (IACT): towards a consensus sequence for intracellular antibodies. *J Mol Biol* 317, 73-83.
- 5 Visintin, M., Tse, E., Axelson, H., Rabbitts, T.H. and Cattaneo, A. (1999) Selection of antibodies for intracellular function using a two-hybrid in vivo system. In: *Proc Natl Acad Sci U S A*, Vol. 96, p. 11723-8.
- Visintin, M., Tse, E., Axelson, H., Rabbitts, T.H. and Cattaneo, A. (1999) Selection of antibodies for intracellular function using a two-hybrid in vivo system. In: *Proc Natl Acad Sci U S A*, Vol. 96, p. 11723-8.
- 10 Wu, A.M. and Yazaki, P.J. (2000) Designer genes: recombinant antibody fragments for biological imaging. *Q J Nucl Med* 44, 268-83.
- Zhong, H., SuYang, H., Erdjument-Bromage, H., Tempst, P. and Ghosh, S. (1997) The transcriptional activity of NF-kappaB is regulated by the IkappaB-associated PKAc subunit through a cyclic AMP-independent mechanism. *Cell* 89, 413-24.
- 15 Zhong, H., Voll, R.E. and Ghosh, S. (1998) Phosphorylation of NF-kappa B p65 by PKA stimulates transcriptional activity by promoting a novel bivalent interaction with the coactivator CBP/p300. *Mol Cell* 1, 661-71.

CLAIMS

1. A method to isolate yeast clones expressing intracellular antibodies able to neutralize an interaction between a known protein ligand x and a known protein ligand y inside a cell, comprising the steps of:

a) obtaining a recombinant yeast strain obtained by cloning a first sequence encoding the protein ligand x and a second sequence encoding the protein ligand y into a yeast expression vector, each first and second sequence being under the control of each one of two portions of a bidirectional promoter, the first sequence being fused to a sequence encoding a first molecule, and the second sequence being fused to a sequence encoding a second molecule, so that when the protein ligand x and the protein ligand y interact said first molecule and said second molecule interact in a way to induce the production either of a repressor of an essential function of the yeast cell or of a toxic agent, transforming the recipient yeast strain with the vector of point a); and selecting transformed yeasts wherein:

- i) the first molecule is a DNA binding domains and the second molecule is an activating domains respectively, but
- ii) at least one of x or y is expressed in an inducible manner, and
- iii) the interaction between the protein ligand x and the protein ligand y would lead to the production either of a repressor of an essential function of the yeast cell or of a toxic agent, so that non viable yeast cells would be obtained;

b) constructing an expression library of intracellular antibodies in the recombinant yeast strain of point a);

c) selecting transformed yeast clones by inducing the expression of both ligand x and ligand y; and

d) isolating recipient yeast clones able to grow.

2. A method to isolate yeast clones expressing intracellular antibodies able to neutralize an interaction between a known protein ligand x and a known protein ligand y inside a cell, comprising the steps of:

a) obtaining an expression library of intracellular antibodies in a yeast strain with mat-a or mat-alpha sexual competence and carrying a repressor of an essential function of the yeast cell;

b) obtaining a recombinant yeast strain having a sexually competence opposite to that of the yeast strain in point a) obtained by cloning a first sequence encoding the protein ligand x and a second sequence encoding the protein ligand y into a yeast expression vector, each first and second sequence being under the control of each one of two portions of a bidirectional promoter, the first sequence being fused to a sequence encoding a first molecule, and the second sequence being fused to a sequence encoding a second molecule, so that when the protein

ligand x and the protein ligand y interact said first molecule and said second molecule interact in a way to induce the production either of a repressor of an essential function of the yeast cell or of a toxic agent, transforming the recipient yeast strain with the vector of point a); and selecting transformed yeasts wherein:

- i) the first molecule is a DNA binding domains and the second molecule is an activating domains respectively, but
 - ii) at least one of x or y is expressed in an inducible manner, and
 - iii) the interaction between the protein ligand x and the protein ligand y would lead to the production either of a repressor of an essential function of the yeast cell or of a toxic agent, so that non viable yeast cells would be obtained;
- c) exposing yeasts of point a) and recombinant yeasts of point b) to conditions able to promote sexual mating and the expression of the ligands x and y;
- d) selecting transformed yeast clones by inducing the expression of both ligand x and ligand y; and
- e) isolating recipient yeast clones able to grow.
3. The method according to claim 1 or 2 wherein the expression vector in yeast comprises as a bidirectional promoter the promoters Gal 1 and Gal 10.
4. The method according to claim 1 or 2 wherein the first and the second molecule leads to the production of a repressor able to shut down the transcription of the yeast HIS3 gene.
5. The method according to any of claims 1 to 4 in which the expression library of intracellular antibodies in yeast is obtained from mouse or human lymphocytes.
6. The method according to any of claims 1 to 5 in which the expression library of intracellular antibodies in yeast is obtained in a scFv or in a single domain antibody format.
7. A yeast expression vector for cloning both a sequence encoding a protein ligand x and a sequence encoding a protein ligand y able to interact each other, the protein ligand x and the protein ligand y being fused to a DNA binding domain and an activating domain respectively comprising a bidirectional promoter wherein the first promoter portion is 5' to 3' followed by appropriate restriction cloning sites for the sequence encoding the protein ligand x and a first sequence encoding a first molecule; and the second promoter portion is 5' to 3' followed by appropriate restriction cloning sites for the sequence encoding the protein ligand y and a second sequence encoding a second molecule, so that when the protein ligand x and the protein ligand y interact, said first molecule and said second molecule interact in a way to induce the production either of a repressor of an essential function of the yeast cell or of a toxic agent.

Fig.1




Fig.2




Fig.3






Fig.4a



es) yeast strain che abbia:

Matα, ura 3-52, leu 2-3112, his 3- 200, trp 1-901,
ade2-101, LYS::(lexAop)-HIS3, ura3::(lexAop)-LacZ,
trp::(Gal1)-CRE

mating




Fig.4b




Fig.5





Fig.6





Fig.7



WT-lexA (pMIC-BD1)	GTT ACC GTT AAG CGC CTG AAA AAA CAG GGC AAT
	V T V K R L K K Q G N
mut-lexA (pMIC-BD2)	GTT ACC GTT AAG GGC CTG GAA AAA CAG GGC AAT
	V T V K G L E K Q G N

Fig.8




Fig.9





Fig.10




Fig.11




Fig.12a




Fig.12b





Fig.13

Fingerprinting anti-SH2-Shc



ϕ X 1 16 45 50 61 79 126 137 140 155




 β -gal assay anti SH2-Shc


Fig.14

Fingerprinting anti-K-RAS


ϕ X 1a 1b 7a 7b 12a 12b




4a 4b 4c 4d 6a 6b 6c 6d 10a 10b 10c 10d ϕ X 15a 15b 15c 15d



Screening A




β -gal assay anti K-RAS





Screening B

Fig.15



II screening: in black SH2-MIC-BD1 + anti-SH2-Shc
in blue K-RAS-MIC-BD2 + anti-K-RAS

Fig.16

 β -gal assay anti Syk

PCR-BstNI fingerprinting anti-Syk

Fig.17




Fig.18




Fig.19




Fig.20





Fig.21




Fig.22




Fig.23






Fig.24

anti-p65
A2



scFv c





Fig.25

