

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 November 2001 (22.11.2001)

PCT

(10) International Publication Number  
**WO 01/88110 A1**

(51) International Patent Classification<sup>7</sup>: **C12N 9/00**,  
15/00, C07H 21/04

**Ioanna, P.** [GR/US]; 385 S. Catalina Avenue, Apt. 206,  
Pasadena, CA 91106 (US). **SUN, Lianhong** [CN/US]; 307  
South Wilson Avenue, Apt. 1, Pasadena, CA 91106 (US).

(21) International Application Number: PCT/US00/32345

(74) Agents: **SCHAFFER, Robert** et al.; Darby & Darby P.C.,  
805 Third Avenue, New York, NY 10022-7513 (US).

(22) International Filing Date:  
27 November 2000 (27.11.2000)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL,  
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL,  
PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ,  
UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:  
09/571,553 16 May 2000 (16.05.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier application:  
US 09/571,553 (CIP)  
Filed on 16 May 2000 (16.05.2000)

(71) Applicant (*for all designated States except US*): **CALI-  
FORNIA INSTITUTE OF TECHNOLOGY** [US/US];  
1200 East California Boulevard, Mail Code 210-85,  
Pasadena, CA 91125 (US).

**Published:**  
— with international search report  
— with amended claims

(71) Applicants and  
(72) Inventors: **ARNOLD, Frances, H.** [US/US]; 629 S.  
Grand Avenue, Pasadena, CA 91105 (US). **PETROUNIA,**

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 01/88110 A1**

(54) Title: DIRECTED EVOLUTION OF OXIDASE ENZYMES

(57) Abstract: This invention relates to the expression of improved polynucleotide and polypeptide sequences encoding for eukaryotic enzymes, particularly oxidase enzymes. The enzymes are advantageously produced in conventional or facile expression systems. Various methods for directed evolution of polynucleotide sequences can be used to obtain the improved sequences. The improved characteristics of the polypeptides or proteins generated in this manner include improved expression, enhanced activity toward one or more substrates, and increased thermal stability. In a particular embodiment, the invention relates to improved expression of the galactose oxidase gene and galactose oxidase enzymes. GAO mutants that are highly active and/or thermostable are disclosed.

## **DIRECTED EVOLUTION OF OXIDASE ENZYMES**

This invention is concerned with the production of modified enzymes, particularly oxidase enzymes, and more particularly galactose oxidase enzymes. Recombinant techniques such as directed evolution are used to obtain polynucleotide and polypeptide products having desirable properties. Galactose oxidase variants with increased activity and increased thermostability relative to the wild-type enzyme are described.

### **BACKGROUND OF THE INVENTION**

An "oxidation enzyme" is an enzyme that catalyzes one or more oxidation reactions, typically by adding, inserting, contributing or transferring oxygen from a source or donor to a substrate. Such enzymes are also called oxidoreductases or redox enzymes, and encompasses oxygenases, hydrogenases or reductases, oxidases and peroxidases. One such enzyme is galactose oxidase. This invention relates to the selection and production of polynucleotides that encode polypeptides or proteins with biological activity as oxidation enzymes, and in particular galactose oxidase enzymes. These enzymes are produced in facile expression systems such as robust prokaryotic cells (*e.g.* bacteria) and eukaryotic systems (*e.g.* fungi and yeast).

### **FIELD OF THE INVENTION**

The invention concerns the recombinant production of functional eukaryotic proteins by host cells, in high yield, with increased activity, and/or with increased stability, *e.g.* thermostability. Preferred proteins of the invention include oxidase enzymes (oxidases) such as polypeptides evolved from galactose oxidase (*D*-galactose:oxygen 6-oxidoreductase or GAO; EC 1.1.3.9). Polynucleotides which encode and express these proteins in recombinant host cell expression systems, and the resulting polypeptides, are encompassed by the invention.

The publications and reference materials noted herein and listed in the appended Bibliography are each incorporated by reference in their entirety. They are referenced numerically in the text and the Bibliography below.

### **Production of Enzyme Variants**

5 Many proteins of interest are produced by organisms having "eukaryotic" cells. These are cells having a nucleus surrounded by its own membrane and containing DNA on structures called chromosomes. All multicellular organisms, such as humans and animals, and many single-cell animals, have eukaryotic cells. Other single-cell organisms, such as bacteria have "prokaryotic" cells. These cells have a primitive nucleus with DNA in a defined  
10 structure, but without chromosomes and a nuclear membrane that is characteristic of eukaryotes. Prokaryotic organisms are generally much easier and less costly to grow, maintain and manipulate than eukaryotic cells.

Genetic engineering and recombinant DNA and RNA technologies have made it possible to produce proteins, hormones and enzymes that are native to one organism, by using  
15 the cells of a different organism as "factories" or host cell expression systems. In particular, it is often desirable to express a protein of eukaryotic origin in a prokaryotic host cell, because the prokaryotes can be grown in large quantities of identical cells, to produce large amounts of the desired foreign protein. For example, certain human proteins may be useful as drugs if they can be supplied in sufficient quantity to patients who have a protein deficiency. Such  
20 proteins may not easily or ethically be obtained by isolating them from human cells, nor can they easily be made by direct chemical synthesis or by growing them in isolated tissue cultures. Other proteins and enzymes are useful in industry. For example, certain enzymes can break down food products, and are useful in laundry detergent. However, commercial applications require large amounts of protein and a high degree of quality control. Desirable  
25 applications also require or would benefit from more active or more thermostable (heat resistant) proteins or enzymes.

To solve some of these problems, recombinant genetic engineering techniques have been developed to use genetic machinery of other cells, such as bacteria and yeast, to produce human or other proteins. Selected genetic material, such as a polynucleotide that encodes a  
30 desired protein, is "recombined" with genetic material in a host cell, so that the host cell

-3-

expresses the introduced foreign genetic material and produces the desired polypeptide or protein. Bacteria, fungi and yeast can be suitable host cells because they are easy and economical to grow and maintain in large quantities, and can be used to reliably and repeatably produce foreign proteins. Some proteins that are made by cells can be secreted or delivered outside the cell, which can improve the yield and the efficiency of subsequent isolation and purification steps.

Directed evolution has been successfully applied to improve a variety of enzyme properties, such as substrate specificity, activity in organic solvents, and stability at high temperatures, which are often critical for industrial applications (5). This evolutionary approach uses DNA shuffling, for simultaneous random mutagenesis and recombination, to generate a variant having an improved desirable property over the existing wild type protein. Point mutations are generated due to the intrinsic infidelity of Taq-based polymerase chain reactions (PCR) associated with reassembly of nucleic acid sequences. In one example, Stemmer and coworkers applied this technique to the gene encoding for green fluorescence protein (GFP), which resulted in a protein that folded better than the wild type in *E. coli* (10). Other examples are in the literature. (11-18, 21-25, 27-34, 47-58, 60-63, 65-75). Eukaryotic enzymes have a myriad of existing and potential applications, but improvement of these and other proteins by directed evolution is desirable. For example, the difficulty of expressing certain oxidase enzymes in a facile expression host has posed technical challenges. Efforts to modify these enzymes for industrial applications by protein engineering methods have been impeded. Directed evolution, for example, exploits expression in a host such as *E. coli* or *S. cerevisiae*, organisms in which large libraries of mutants or variants can be made. Also, the lack of efficient expression in an appropriate foreign (heterologous) host can prevent the mass production of some of these proteins on an economical scale. Thus, there continues to be a need for new ways to produce new proteins, and for new proteins and enzymes having new or enhanced biological properties.

### **Galactose Oxidase Enzymes**

One protein of interest is the oxidation enzyme galactose oxidase. Galactose oxidase (D-galactose: oxygen 6-oxidoreductase, GAO; EC 1.1.3.9) is an enzyme containing a single copper ion, and is secreted by a number of fungal species. *Fusarium* NRRL 2903, formerly

known as *Dactylium dendroides*, has been the most extensively studied (76). The enzyme is a glycoprotein with a carbohydrate content of about 1.7% and consists of a single polypeptide chain of 639 amino acid residues with molecular mass of 68,000 Da (77, 78). The reaction catalyzed by GAO is the oxidation of primary alcohols to the corresponding aldehydes, coupled to the two-electron reduction of O<sub>2</sub> to hydrogen peroxide (79).

The enzyme oxidizes an unusually broad range of substrates. It accepts D-galactose (FIG. 1), alpha- and beta-galactopyranosides, oligo- and polysaccharides and considerably smaller molecules, such as glycerol and allyl alcohol, as substrates (77, 80-82). GAO exhibits prochiral (only the *pro-S* hydrogen is abstracted) as well as enantiomeric specificity for galactose (only D-galactose is oxidized by the enzyme) (80, 83). Furthermore, GAO strictly discriminates against D-glucose, the C-4 epimer of D-galactose, as a substrate or ligand. D-glucose does not bind to GAO at concentrations as high as 1 M (80, 84). The kinetic parameters of GAO for the oxidation of galactose are:  $K_m = 67 \text{ mM}$ ,  $k_{cat} = 3,000 \text{ sec}^{-1}$ ,  $k_{cat}/K_m = 45 \times 10^3 \text{ M}^{-1}\text{sec}^{-1}$  (85).

The crystal structure of GAO has been reported (86). It consists of three predominantly beta-structure domains. The copper ion lies on the solvent-accessible surface of the second and largest domain (residues 156-532) (78, 87). Tyr-272, Tyr-495, His-496, His-581 and a water molecule are the copper ligands at pH 7.0. The crystal structure also reveals a novel thioether bond linking Cys-228 and Tyr-272 and supports the presence of a tyrosine free radical at the active site (79). The active site structure of GAO is shown in FIG. 2. Site-directed mutagenesis of Tyr-495 and Cys-228 have confirmed their involvement in catalysis (85, 88).

GAO is useful in a wide variety of applications, ranging from analytical and food chemistry to chemoenzymatic synthesis and clinical testing. For example, biological sensors based on GAO have been developed to determine the content of galactose (89), lactose and other GAO substrates (90). Such biosensors have also been used for quality control in dairy industries (91, 92), online bioprocess monitoring (93) and analysis of blood samples of patients with suspected galactosemia (94). The stereospecificity and broad substrate specificity of GAO have been exploited in the chemoenzymatic synthesis of L-sugars from polyols (95), which are usually difficult to prepare by chemical methods (96, 97), as well as sugar-containing polyamines (98) and 5-C-(hydroxymethyl)hexoses (99). GAO applications

in synthesis have been limited due to its relatively low activity toward a large number of primary alcohols (100). Additionally, GAO is also used for the detection of the disaccharide D-galactose-beta-(1->3)-N-acetylgalactosamine (Gal-GalNAc), a tumor marker in colonic cancer and precancer, and provides a cost-effective screening test for patients with neoplasia or at the risk of developing neoplasia (101, 102). GAO finds applications in food chemistry. For example, it has been used in oxidized guar manufacture (103) and to treat the oligosaccharide fraction contained in honey (104). Finally, GAO is used to oxidize the cell surface polysaccharides of membrane-bound glycoproteins containing terminal non-reducing galactose residues: this is an essential step in the successful radiolabeling of these glycoconjugates (105, 106).

Modified and particularly improved or optimized GAO enzymes are useful to improve and expand the use of the enzyme in practical applications. For example, enzymes of the invention include GAO variants that are more active, more thermostable, or both. Increased activity and/or expression as well as high thermostability may significantly decrease the cost of enzyme production, simplify its purification and handling, and prolong its shelf-life. Other properties of the enzyme may also be varied, for example to optimize activity towards particular substrates or toward other substrates such as polymeric materials and glucose.

Use of these evolved enzymes in biosensors and diagnostics can increase sensitivity, decrease the response time and enhance the detection range. In addition, a more stable enzyme will find applications in the construction of biosensors with prolonged stability. An evolved GAO with improved activity toward poor GAO substrates, such as allyl alcohol and glucose, will provide new and improved applications of the enzyme in organic synthesis and other sensor applications. For chemical synthesis applications, selective oxidation of alcohols to the corresponding aldehydes avoids the use of protecting groups, minimizes side reactions often observed in traditional chemical synthesis, and is an environmentally friendly process. Use of such GAO enzymes as a synthetic reagent would facilitate the use of more inexpensive, safe and biodegradable carbohydrate materials in industrial processes (107).

A more efficient enzyme is expected to be advantageous in the food chemistry applications of GAO, and, in particular in the selective modification of guar and other carbohydrate-based polymers. GAO variants according to the invention would also be useful for modification of carbohydrate-based (e.g. cellulosic) textiles and other materials. The

aldehyde function produced by the GAO can be used to couple other substances selectively at the modified position on the polymer.

Accordingly, there is a need to develop new and improved GAO enzymes, as well as methods for expressing such proteins. In particular, there is a need for protein expression methods which are well-suited for use in connection with directed evolution techniques.

This invention describes methods for screening libraries of GAO mutants produced by error-prone PCR and DNA shuffling, to identify mutations that are expressed in bacteria (e.g. *E. coli*) and with improved GAO function. Micro-plate and membrane screening techniques are disclosed. In one embodiment, the mutant is a functional and active galactose oxidase (GAO) that is expressed in *E. coli* at levels of about 65 times the activity of a parent recombinant wild type (for D-galactose). The activity for other substrates, such as allyl alcohol, is also about 65 times that of wild type. Mutants of the invention can have any fraction or multiple of the corresponding wild type activity, but preferably are more active, e.g. about 2 to 200 times as active. Mutants also are more thermostable. Enzyme yield is generally at least about 10 mg/l.

### **SUMMARY OF THE INVENTION**

The observed constraints on the use of native proteins are thought to be a consequence of evolution. Proteins have evolved in the context and environment of a living organism, to carry out specific biological functions under conditions conducive to life – not in the laboratory or under industrial conditions. In some cases, evolution may favor or even require less than optimally efficient enzymes. The output, efficiency, working conditions, stability and other properties of known expression systems are not thought to be unalterable, nor are they limitations which should be seen as intrinsic to the nature of cellular expression systems. It is possible that the proteins used in these systems can be evolved *in vitro*, or that analogous proteins can be otherwise developed, to alter or enhance the protein's properties, for example, to obtain much more efficient expression, activity and thermostability. Improved proteins can also be obtained by screening cultures of native organisms or expressed gene libraries (3).

The invention provides a method for improving the expression, thermostability, and/or the activity toward one or more substrates, of a polynucleotide encoding oxidase

enzymes by using directed evolution. The invention also provides polynucleotides encoding for variant oxidase enzymes which have improved properties in conventional expression systems. According to one embodiment of the invention, directed evolution or random mutagenesis is used to produce GAO variants which are more highly expressed, more active, and/or more thermostable in prokaryotic expression systems such as *E. coli*.

The above features and many other attendant advantages of the invention will become better understood by reference to the following detailed description when taken in conjunction with the accompanying drawings.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

**FIG. 1** shows a reaction scheme in which a D-galactose substrate is oxidized to produce a D-galactohexodialdose product, in the presence of galactose oxidase (GAO) enzyme.

**FIG. 2** shows the active site structure of GAO pH 7.0

**FIG. 3** is a graph showing the effect of metal ions (particularly copper ions) on the activity of a recombinant wild-type GAO, pGAO-010. Enzyme solutions with additives were kept at 4 °C for 1 hr before assay. Relative activity of enzyme solution with 1 mM copper (II) sulfate was estimated as 100 %.

**FIG. 4** is a graph showing GAO activity for various clones generated by error-prone PCR, with varying concentrations of MnCl<sub>2</sub>, using conditions A of TABLE 3.

**FIG. 5** is a graph showing GAO activity for various clones generated by error-prone PCR, with varying concentrations of MnCl<sub>2</sub>, using conditions C of TABLE 3.

**FIG. 6** shows the sequences of PCR primers used herein for amplification, *e.g.* of the whole galactose oxidase gene.

**FIG. 7** is a schematic representation of the construction of plasmid pUC18-EHL.

**FIG. 8** is a schematic representation of the construction of plasmid pGAO-010.

**FIG. 9** is a schematic representation of the construction of plasmids pGAO-027 and pGAO-036.

**FIG. 10** is a schematic representation of the construction of plasmids pGAO-006 and pGAO-011.



**FIG. 11** shows the structures and activities of representative plasmids encoding GAO according to the invention, with IPTG-induced expression in host *E. coli*. Permeable cells which were treated by freeze (-20 °C), thaw (4 °C) and 0.5 mg/l lysozyme for 30 minutes at 37 °C were used for assay. Activities given as \* indicates that cells did not grow in test tube culture; \*\* indicates that a transformant was not obtained.

**FIG. 12** shows a scheme for the design of plasmids according to the invention.

**FIG. 13** shows the structures and activities of additional plasmids encoding GAO according to the invention, with IPTG-induced expression in host *E. coli*.

**FIG. 14** is a graph comparing the GAO activities of GAO plasmids with and without random codon alternation.

**FIG. 15** shows substrate specificities for a wild type galactose oxidase and a recombinant galactose oxidase enzyme of the invention. Partially purified galactose oxidase from *D. dendroides* (Sigma) and cell-free extract from *E. coli* BL21(DE3)/pGAO-010 were used. Relative activities for D-galactose were estimated as 100 %. (+) indicates that oxidation was detected, but activities were too low to be estimated. n.d. indicates that activities were not distinguishable from background absorbance levels.

**FIG. 16** is a graph showing the thermal stability of selected GAO mutants.

**FIGS. 17A-C** show the sequence of representative mutant 9.16.8D2 of the invention [SEQ. ID NO. 10]

**FIGS. 18A-C** show the sequence of representative mutant 9.16.6C11 of the invention [SEQ. ID NO. 11]

**FIGS. 19A-C** show the sequence of representative mutant 9.16.16D12 of the invention [SEQ. ID NO. 12]

**FIGS. 20A-C** show the sequence of representative mutant 11.03.6D3 of the invention [SEQ. ID NO. 13]

**FIGS. 21A-C** show the sequence of representative mutant 11.03.10C3 of the invention [SEQ. ID NO. 14]

**FIGS. 22A-C** show the sequence of representative mutant 11.03.10D6 of the invention [SEQ. ID NO. 15]

**FIGS. 23A-C** show the sequence of representative mutant 11.03.13E12 of the invention [SEQ. ID NO. 16]

FIGS. 24A-C show the sequence of representative mutant 1.06.20E7 of the invention [SEQ. ID NO. 17]

FIGS. 25A-C show the sequence of representative mutant 1.D4 of the invention [SEQ. ID NO. 18]

5 FIGS. 26A-C show the sequence of representative mutant 2G4 of the invention [SEQ. ID NO. 19]

FIGS. 27A-C show the sequence of representative mutant 3.H7 of the invention [SEQ. ID NO. 20]

10 FIGS. 28A-C show the sequence of representative mutant 4.F12 of the invention [SEQ. ID NO. 21]

### **DETAILED DESCRIPTION OF THE INVENTION**

This invention concerns methods for improving the expression, activity and/or thermostability of proteins using facile or conventional expression systems.

#### **Definitions**

15 As used herein, "about" or "approximately" shall mean within 20 percent, preferably within 10 percent, and more preferably within 5 percent of a given value or range.

The term "substrate" means any substance or compound that is converted or meant to be converted into another compound by the action of an enzyme catalyst. The term  
20 includes aromatic and aliphatic compounds, and includes not only a single compound, but also combinations of compounds, such as solutions, mixtures and other materials which contain at least one substrate.

An "oxidation reaction" or "oxygenation reaction", as used herein, is a chemical or biochemical reaction involving the addition of oxygen to a substrate, to form an  
25 oxygenated or oxidized substrate or product. An oxidation reaction is typically accompanied by a reduction reaction (hence the term "redox" reaction, for oxidation and reduction). A compound is "oxidized" when it receives oxygen or loses electrons. A compound is "reduced" when it loses oxygen or gains electrons. GAO typically catalyzes the oxidation of a primary alcohol group to an aldehyde.

The term "enzyme" means any substance composed wholly or largely of protein or polypeptides that catalyzes or promotes, more or less specifically, one or more chemical or biochemical reactions.

5 A "polypeptide" (one or more peptides) is a chain of chemical building blocks called amino acids that are linked together by chemical bonds called peptide bonds. A protein or polypeptide, including an enzyme, may be "native" or "wild-type", meaning that it occurs in nature or has the amino acid sequence of a native protein, respectively. These terms are sometimes used interchangeably. A polypeptide may or may not be glycosylated. A "recombinant wild-type" typically means the wild type sequence in a recombinant host without glycosylation. Comparisons in the examples and figures of this application are generally with reference to a wild type that is a recombinant wild type. A polypeptide may also be a "mutant", "variant" or "modified", meaning that it has been made, altered, derived, or is in some way different or changed from a native protein, or from another mutant. A native wild type protein comprises the natural sequence of amino acids in the polypeptide and typically includes glycosylation. A "parent" polypeptide or enzyme is any polypeptide or enzyme from which any other polypeptide or enzyme is derived or made, using any methods, tools or techniques, and whether or not the parent is itself a native or mutant polypeptide or enzyme. A parent polynucleotide is one that encodes a parent polypeptide. A "test enzyme" is a protein-containing substance that is tested to determine whether it has properties of an enzyme. The term "enzyme" can also refer to a catalytic polynucleotide (e.g. RNA or DNA).

10  
15  
20

The "activity" of an enzyme is a measure of its ability to catalyze a reaction, and may be expressed as the rate at which the product of the reaction is produced. For example, enzyme activity can be represented as the amount of product produced per unit of time, per unit (*e.g.* concentration or weight) of enzyme. The "stability" of an enzyme means its ability to function, over time, in a particular environment or under particular conditions. One way to evaluate stability is to assess its ability to resist a loss of activity over time, under given conditions. Enzyme stability can also be evaluated in other ways, for example, by determining the relative degree to which the enzyme is in a folded or unfolded state. Thus, one enzyme is more stable than another, or has improved stability, when it is more resistant than the other enzyme to a loss of activity under the same

25  
30

conditions, is more resistant to unfolding, or is more durable by any suitable measure. For example, a more "thermally stable" or "thermostable" enzyme is one that is more resistant to loss of structure (unfolding) or function (enzyme activity) when exposed to heat or an elevated temperature. One way to evaluate this is to determine the "melting temperature" or  $T_m$  for the protein. The melting temperature, also called a midpoint, is the temperature at which half of the protein is unfolded from its fully folded state. This midpoint is typically determined by calculating the midpoint of a titration curve that plots protein unfolding as a function of temperature. Thus, a protein with a higher  $T_m$  requires more heat to cause unfolding and is more stable or more thermostable. Stated another way, a protein with a higher  $T_m$  indicates that fewer molecules of that protein are unfolded at the same temperature as a protein with a lower  $T_m$ , again meaning that the protein which is more resistant to unfolding is more stable (it has less unfolding at the same temperature). Another measure of stability is  $T_{1/2}$  or  $T_{50}$ , which is the transition midpoint of the inactivation curve of the protein as a function of temperature.  $T_{1/2}$  is the temperature at which the protein loses half of its activity. Thus, a protein with a higher  $T_{1/2}$  requires more heat to deactivate it, and is more stable or more thermostable. Stated another way, a protein with a higher  $T_{1/2}$  indicates that fewer molecules of that protein are inactive at the same temperature as a protein with a lower  $T_{1/2}$ , again meaning that the protein which is more resistant to deactivation is more stable (it has more activity at the same temperature). These assays are also called "thermal shift" assays, because the inactivation or unfolding curve, plotted against temperature, is "shifted" to higher or lower temperatures when stability increases or decreases. Thermostability can also be measured in other ways. For example, a longer half-life ( $t_{1/2}$ ) for the enzyme's activity at elevated temperature is an indication of thermostability.

An "oxidation enzyme" is an enzyme that catalyzes one or more oxidation reactions, typically by adding, inserting, contributing or transferring oxygen from a source or donor to a substrate. Such enzymes are also called oxidoreductases or redox enzymes, and encompasses oxygenases, hydrogenases or reductases, oxidases and peroxidases.

The terms "oxygen donor", "oxidizing agent" and "oxidant" mean a substance, molecule or compound which donates oxygen to a substrate in an oxidation reaction. Typically, the oxygen donor is reduced (accepts electrons). Exemplary oxygen donors,

which are not limiting, include molecular oxygen or dioxygen (O<sub>2</sub>) and peroxides, including alkyl peroxides such as t-butyl peroxide, and most preferably hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). A peroxide is any compound having two oxygen atoms bound to each other.

5 A "luminescent" substance means any substance which produces detectable electromagnetic radiation, or a change in electromagnetic radiation, most notably visible light, by any mechanism, including color change, UV absorbance, fluorescence and phosphorescence. Preferably, a luminescent substance according to the invention produces a detectable color, fluorescence or UV absorbance. The term "chemiluminescent agent" means any luminescent substance which enhances the detectability of a luminescent (*e.g.*,  
10 fluorescent) signal, for example by increasing the strength or lifetime of the signal. One exemplary and preferred chemiluminescent agent is azinobis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS). 5-amino-2,3-dihydro-1,4-phthalazinedione (luminol) and analogs. Others include 5-amino-2,3-dihydro-1,4-phthalazinedione (luminol) and analogs, 1,2-dioxetanes such as tetramethyl-1,2-dioxetane (TMD), 1,2-dioxetanones, and 1,2-  
15 dioxetanediones, *o*-anisidine, *o*-dianisidine, and *o*-tolidine. Another term for these kinds of materials is "chromogen."

The term "polymer" means any substance or compound that is composed of two or more building blocks ('mers') that are repetitively linked to each other. For example, a "dimer" is a compound in which two building blocks have been joined together.

20 The term "cofactor" means any non-protein substance that is necessary or beneficial to the activity of an enzyme. A "coenzyme" means a cofactor that interacts directly with and serves to promote a reaction catalyzed by an enzyme. Many coenzymes serve as carriers. For example, NAD<sup>+</sup> and NADP<sup>+</sup> carry hydrogen atoms from one enzyme to another. An "ancillary protein" means any protein substance that is necessary or  
25 beneficial to the activity of an enzyme.

The term "host cell" means any cell of any organism that is selected, modified, transformed, grown, or used or manipulated in any way, for the production of a substance by the cell, for example the expression by the cell of a gene, a DNA or RNA sequence, a protein or an enzyme.

30 "DNA" (deoxyribonucleic acid) means any chain or sequence of the chemical building blocks adenine (A), guanine (G), cytosine (C) and thymine (T), called nucleotide

bases, that are linked together on a deoxyribose sugar backbone. DNA can have one strand of nucleotide bases, or two complimentary strands which may form a double helix structure. "RNA" (ribonucleic acid) means any chain or sequence of the chemical building blocks adenine (A), guanine (G), cytosine (C) and uracil (U), called nucleotide bases, that are linked together on a ribose sugar backbone. RNA typically has one strand of nucleotide bases.

A "polynucleotide" or "nucleotide sequence" is a series of nucleotide bases (also called "nucleotides") in DNA and RNA, and means any chain of two or more nucleotides. A nucleotide sequence typically carries genetic information, including the information used by cellular machinery to make proteins and enzymes. These terms include double or single stranded genomic and cDNA, RNA, any synthetic and genetically manipulated polynucleotide, and both sense and anti-sense polynucleotide (although only sense stands are being represented herein). This includes single- and double-stranded molecules, *i.e.*, DNA-DNA, DNA-RNA and RNA-RNA hybrids, as well as "protein nucleic acids" (PNA) formed by conjugating bases to an amino acid backbone. This also includes nucleic acids containing modified bases, for example thio-uracil, thio-guanine and fluoro-uracil.

The polynucleotides herein may be flanked by natural regulatory sequences, or may be associated with heterologous sequences, including promoters, enhancers, response elements, signal sequences, polyadenylation sequences, introns, 5'- and 3'- non-coding regions, and the like. The nucleic acids may also be modified by many means known in the art. Non-limiting examples of such modifications include methylation, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, and internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.). Polynucleotides may contain one or more additional covalently linked moieties, such as, for example, proteins (*e.g.*, nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), intercalators (*e.g.*, acridine, psoralen, etc.), chelators (*e.g.*, metals, radioactive metals, iron, oxidative metals, etc.), and alkylators. The polynucleotides may be derivatized by formation of a methyl or ethyl phosphotriester or an alkyl phosphoramidate linkage. Furthermore, the polynucleotides herein may also be modified with a label capable of

providing a detectable signal, either directly or indirectly. Exemplary labels include radioisotopes, fluorescent molecules, biotin, and the like.

5 Proteins and enzymes are made in the host cell using instructions in DNA and RNA, according to the genetic code. Generally, a DNA sequence having instructions for a particular protein or enzyme is "transcribed" into a corresponding sequence of RNA. The RNA sequence in turn is "translated" into the sequence of amino acids which form the protein or enzyme. An "amino acid sequence" is any chain of two or more amino acids. Each amino acid is represented in DNA or RNA by one or more triplets of nucleotides. Each triplet forms a codon, corresponding to an amino acid. For example, the amino acid  
10 lysine (Lys) can be coded by the nucleotide triplet or codon AAA or by the codon AAG. (The genetic code has some redundancy, also called degeneracy, meaning that most amino acids have more than one corresponding codon.) Because the nucleotides in DNA and RNA sequences are read in groups of three for protein production, it is important to begin reading the sequence at the correct amino acid, so that the correct triplets are read. The  
15 way that a nucleotide sequence is grouped into codons is called the "reading frame."

The term "gene", also called a "structural gene" means a DNA sequence that codes for or corresponds to a particular sequence of amino acids which comprise all or part of one or more proteins or enzymes, and may or may not include regulatory DNA sequences, such as promoter sequences, which determine for example the conditions under which the  
20 gene is expressed. Some genes, which are not structural genes, may be transcribed from DNA to RNA, but are not translated into an amino acid sequence. Other genes may function as regulators of structural genes or as regulators of DNA transcription.

A "coding sequence" or a sequence "encoding" a polypeptide, protein or enzyme is a nucleotide sequence that, when expressed, results in the production of that polypeptide, protein or enzyme, *i.e.*, the nucleotide sequence encodes an amino acid sequence for that  
25 polypeptide, protein or enzyme. A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced and translated into the protein encoded by the coding sequence. Preferably, the coding  
30 sequence is a double-stranded DNA sequence which is transcribed and translated into a polypeptide in a cell *in vitro* or *in vivo* when placed under the control of appropriate

regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (*e.g.*, mammalian) DNA,  
5 and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, terminators, and the like, that provide for the expression of a  
10 coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining this invention, the promoter sequence is bounded at its  
15 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding  
20 of RNA polymerase. As described above, promoter DNA is a DNA sequence which initiates, regulates, or otherwise mediates or controls the expression of the coding DNA. A promoter may be "inducible", meaning that it is influenced by the presence or amount of another compound (an "inducer"). For example, an inducible promoter includes those which initiate or increase the expression of a downstream coding sequence in the presence  
25 of a particular inducer compound. A "leaky" inducible promoter is a promoter that provides a high expression level in the presence of an inducer compound and a comparatively very low expression level, and at minimum a detectable expression level, in the absence of the inducer.

A "signal sequence" is included at the beginning of the coding sequence of a  
30 protein to be expressed in the periplasmic space, or outside the cell. This sequence encodes a signal peptide, N-terminal to the mature polypeptide, that directs the host cell to



translocate the polypeptide. The term "translocation signal sequence" is also used to refer to a signal sequence. Translocation signal sequences can be found associated with a variety of proteins native to eukaryotes and prokaryotes, and are often functional in both types of organisms. Proteins of the invention may be further modified and improved by adding a sequence which directs the secretion of the protein outside the host cell. The addition of the signal sequence does not interfere with the folding of the secreted protein, and evidence thereof is easily tested for using techniques known in the art and depending on the protein (*e.g.*, tests for activity of a given protein after modification).

The terms "express" and "expression" mean allowing or causing the information in a gene or DNA sequence to become manifest, for example producing a protein by activating the cellular functions involved in transcription and translation of a corresponding gene or DNA sequence. A DNA sequence is expressed in or by a cell to form an "expression product" such as a protein. The expression product itself, *e.g.* the resulting protein, may also be said to be "expressed" by the cell. A polynucleotide or polypeptide is expressed recombinantly, for example, when it is expressed or produced in a foreign host cell under the control of a foreign or native promoter, or in a native host cell under the control of a foreign promoter.

A polynucleotide or polypeptide is "over-expressed" when it is expressed or produced in an amount or yield that is substantially higher than a given base-line yield, *e.g.* a yield that occurs in nature. For example, a polypeptide is over-expressed when the yield is substantially greater than the normal, average or base-line yield of the native polypeptide in native host cells under given conditions, for example conditions suitable to the life cycle of the native host cells. Over-expression of a polypeptide can be obtained, for example, by altering any one or more of: (a) the growth or living conditions of the host cells; (b) the polynucleotide encoding the polypeptide to be over-expressed; (c) the promoter used to control expression of the polynucleotide; and (d) the host cells themselves. This is a relative, and thus "over-expression" can also be used to compare or distinguish the expression level of one polypeptide to another, without regard for whether either polypeptide is a native polypeptide or is encoded by a native polynucleotide. Typically, over-expression means a yield that is at least about two times a normal, average or given base-line yield. Thus, a polypeptide is over-expressed when it is produced in an

amount or yield that is substantially higher than the amount or yield of a parent polypeptide or under parent conditions. Likewise, a polypeptide is “under-expressed” when it is produced in an amount or yield that is substantially lower than the amount or yield of a parent polypeptide or under parent conditions, *e.g.* at least half the base-line yield. In this context, the expression level or yield refers to the amount or concentration of polynucleotide that is expressed, or polypeptide that is produced (*i.e.* expression product), whether or not in an active or functional form. As one example, a polynucleotide or polypeptide may be said to be under-expressed when it is expressed in detectable amounts under the control of an inducible promoter, but without induction, *i.e.* in the absence of an inducer compound.

An expression product can be characterized as intracellular, extracellular or secreted. The term “intracellular” means something that is inside a cell. The term “extracellular” means something that is outside a cell. A substance is “secreted” by a cell if it delivered to the periplasm or outside the cell, from somewhere on or inside the cell.

As used herein, the terms “expression-resistant polypeptide” and “resistant to functional expression” are synonymous and refer to a polypeptide that is difficult to functionally express in selected host cells. For example, an expression-resistant polypeptide is not produced, or is produced in very low yield or in non-functional form, when a polynucleotide encoding that polypeptide is transformed or introduced into host cells, *e.g.* into a facile host cell expression system.

The term “transformation” means the introduction of a “foreign” (*i.e.* extrinsic or extracellular) gene, DNA or RNA sequence to a host cell, so that the host cell will express the introduced gene or sequence to produce a desired substance, typically a protein or enzyme coded by the introduced gene or sequence. The introduced gene or sequence may also be called a “cloned” or “foreign” gene or sequence, may include regulatory or control sequences, such as start, stop, promoter, signal, secretion, or other sequences used by a cell’s genetic machinery. The gene or sequence may include nonfunctional sequences or sequences with no known function. A host cell that receives and expresses introduced DNA or RNA has been “transformed” and is a “transformant” or a “clone.” The DNA or RNA introduced to a host cell can come from any source, including cells of the same genus or species as the host cell, or cells of a different genus or species.

The terms “vector”, “cloning vector” and “expression vector” mean the vehicle by which a DNA or RNA sequence (*e.g.* a foreign gene) can be introduced into a host cell, so as to transform the host and promote expression (*e.g.* transcription and translation) of the introduced sequence.

5            Vectors typically comprise the DNA of a transmissible agent, into which foreign DNA is inserted. A common way to insert one segment of DNA into another segment of DNA involves the use of enzymes called restriction enzymes that cleave DNA at specific sites (specific groups of nucleotides) called restriction sites. Generally, foreign DNA is inserted at one or more restriction sites of the vector DNA, and then is carried by the  
10            vector into a host cell along with the transmissible vector DNA. A segment or sequence of DNA having inserted or added DNA, such as an expression vector, can also be called a “DNA construct.”

          A common type of vector is a “plasmid”, which generally is a self-contained molecule of double-stranded DNA, that can readily accept additional (foreign) DNA and  
15            which can readily introduced into a suitable host cell. A plasmid vector often contains coding DNA and promoter DNA and has one or more restriction sites suitable for inserting foreign DNA. Promoter DNA and coding DNA may be from the same gene or from different genes, and may be from the same or different organisms. A large number of vectors, including plasmid and fungal vectors, have been described for replication and/or  
20            expression in a variety of eukaryotic and prokaryotic hosts. Non-limiting examples include pKK plasmids (Clonetech), pUC plasmids, pET plasmids (Novagen, Inc., Madison, WI), pRSET or pREP plasmids (Invitrogen, San Diego, CA), or pMAL plasmids (New England Biolabs, Beverly, MA), and many appropriate host cells, using methods disclosed or cited herein or otherwise known to those skilled in the relevant art.  
25            Recombinant cloning vectors will often include one or more replication systems for cloning or expression, one or more markers for selection in the host, *e.g.* antibiotic resistance, and one or more expression cassettes. Routine experimentation in biotechnology can be used to determine which vectors are best suited for used with the invention. In general, the choice of vector depends on the size of the polynucleotide  
30            sequence and the host cell to be employed in the methods of this invention.

A "cassette" refers to a segment of DNA that can be inserted into a vector at specific restriction sites. The segment of DNA encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure insertion of the cassette in the proper reading frame for transcription and translation.

5           The term "expression system" means a host cell and compatible vector under suitable conditions, *e.g.* for the expression of a protein coded for by foreign DNA carried by the vector and introduced to the host cell. Common expression systems include bacteria (*e.g.* *E. coli* and *B. subtilis*) or yeast (*e.g.* *S. cerevisiae*) host cells and plasmid vectors, and insect host cells and *Baculovirus* vectors. As used herein, a "facile expression  
10           system" means any expression system that is foreign or heterologous to a selected polynucleotide or polypeptide, and which employs host cells that can be grown or maintained more advantageously than cells that are native or heterologous to the selected polynucleotide or polypeptide, or which can produce the polypeptide more efficiently or in higher yield. For example, the use of robust prokaryotic cells to express a protein of  
15           eukaryotic origin would be a facile expression system. Preferred facile expression systems include *E. coli*, *B. subtilis* and *S. cerevisiae* host cells and any suitable vector.

          The terms "mutant" and "mutation" mean any detectable change in genetic material, *e.g.* DNA, or any process, mechanism, or result of such a change. This includes gene mutations, in which the structure (*e.g.* DNA sequence) of a gene is altered, any gene  
20           or DNA arising from any mutation process, and any expression product (*e.g.* protein or enzyme) expressed by a modified gene or DNA sequence. The term "variant" may also be used to indicate a modified or altered gene, DNA sequence, enzyme, cell, etc., *i.e.*, any kind of mutant. Such changes also include changes in the promoter, ribosome binding site, etc.

25           "Sequence-conservative variants" of a polynucleotide sequence are those in which a change of one or more nucleotides in a given codon position results in no alteration in the amino acid encoded at that position.

          "Function-conservative variants" are those in which a given amino acid residue in a protein or enzyme has been changed without altering the overall conformation and  
30           function of the polypeptide, including, but not limited to, replacement of an amino acid with one having similar properties (such as, for example, acidic, basic, hydrophobic, and

the like). Amino acids with similar properties are well known in the art. For example, arginine, histidine and lysine are hydrophilic-basic amino acids and may be interchangeable. Similarly, isoleucine, a hydrophobic amino acid, may be replaced with leucine, methionine or valine. Amino acids other than those indicated as conserved may differ in a protein or enzyme so that the percent protein or amino acid sequence similarity between any two proteins of similar function may vary and may be, for example, from 70% to 99% as determined according to an alignment scheme such as by the Cluster Method, wherein similarity is based on the MEGALIGN algorithm. A "function-conservative variant" also includes a polypeptide or enzyme which has at least 60 % amino acid identity as determined by BLAST or FASTA algorithms, preferably at least 75%, most preferably at least 85%, and even more preferably at least 90%, and which has the same or substantially similar properties or functions as the native or parent protein or enzyme to which it is compared.

The term "DNA reassembly" is used when recombination occurs between identical sequences. "DNA shuffling" refers to a group of *in vitro* or *in vivo* methods involving recombination of nucleic acid species. For example, homologous recombination of pools of nucleic acid fragments or polynucleotides can be employed to generate polynucleotide molecules having variant sequences of the invention. Such methods can be employed to generate polynucleotide molecules having variant sequences of the invention.

"Isolation" or "purification" of a polypeptide or enzyme refers to the derivation of the polypeptide by removing it from its original environment (for example, from its natural environment if it is naturally occurring, or from the host cell if it is produced by recombinant DNA methods). Methods for polypeptide purification are well-known in the art, including, without limitation, preparative disc-gel electrophoresis, isoelectric focusing, HPLC, reversed-phase HPLC, gel filtration, ion exchange and partition chromatography, and countercurrent distribution. For some purposes, it is preferable to produce the polypeptide in a recombinant system in which the protein contains an additional sequence tag that facilitates purification, such as, but not limited to, a polyhistidine sequence. The polypeptide can then be purified from a crude lysate of the host cell by chromatography on an appropriate solid-phase matrix. Alternatively, antibodies produced against the protein or against peptides derived therefrom can be used as purification reagents. Other

purification methods are possible. A purified polynucleotide or polypeptide may contain less than about 50%, preferably less than about 75%, and most preferably less than about 90%, of the cellular components with which it was originally associated. A "substantially pure" enzyme indicates the highest degree of purity which can be achieved using  
5 conventional purification techniques known in the art.

Polynucleotides are "hybridizable" to each other when at least one strand of one polynucleotide can anneal to another polynucleotide under defined stringency conditions. Stringency of hybridization is determined, e.g., by a) the temperature at which hybridization and/or washing is performed, and b) the ionic strength and polarity (e.g.,  
10 formamide) of the hybridization and washing solutions, as well as other parameters.

Hybridization requires that the two polynucleotides contain substantially complementary sequences; depending on the stringency of hybridization, however, mismatches may be tolerated. Typically, hybridization of two sequences at high stringency (such as, for example, in an aqueous solution of 0.5X SSC at 65°C) requires that the sequences exhibit  
15 some high degree of complementarity over their entire sequence. Conditions of intermediate stringency (such as, for example, an aqueous solution of 2X SSC at 65°C) and low stringency (such as, for example, an aqueous solution of 2X SSC at 55°C), require correspondingly less overall complementarity between the hybridizing sequences. (1X SSC is 0.15 M NaCl, 0.015 M Na citrate.) Polynucleotides that "hybridize" to the  
20 polynucleotides herein may be of any length. In one embodiment, such polynucleotides are at least 10, preferably at least 15 and most preferably at least 20 nucleotides long. In another embodiment, polynucleotides that hybridizes are of about the same length. In another embodiment, polynucleotides that hybridize include those which anneal under suitable stringency conditions and which encode polypeptides or enzymes having the  
25 same function, such as the ability to catalyze an oxidation, oxygenase, or coupling reaction of the invention.

The general genetic engineering tools and techniques discussed here, including transformation and expression, the use of host cells, vectors, expression systems, etc., are well known in the art.

### **Mutagenesis and Directed Evolution of Proteins**

To improve the expression and function of proteins using conventional expression systems, the invention makes the unexpected discovery that directed evolution can be used to generate mutant libraries of polynucleotides which, when expressed using conventional or facile expression systems, result in functional proteins having increased activity and/or thermostability.

According to the invention, proteins that are expressed in facile gene expression systems can be obtained by using directed evolution to generate mutant polynucleotides in a library format for selection. General methods for generating libraries and isolating and identifying improved proteins (also described as “variants”) according to the invention using directed evolution are described briefly below and more extensively, for example, in U.S. Patent Nos. 5,741,691 and 5,811,238. *See also*, International Applications WO 98/42832, WO 95/22625, WO 97/20078, and WO 95/ and U.S. Patents 5,605,793 and 5,830,721 (143, 149-156). It should be understood that any method for generating mutations in polynucleotide sequences to provide an evolved polynucleotide for use in expression systems can be employed. Proteins produced by directed evolution methods can then be screened for improved expression, activity, thermostability, folding, secretion, and other functions and properties according to conventional methods.

Any source of nucleic acid in purified form can be utilized as the starting nucleic acid. Thus the process may employ DNA or RNA including messenger RNA, which DNA or RNA may be single or double stranded. In addition, a DNA-RNA hybrid which contains one strand of each may be utilized. The nucleic acid sequence may be of various lengths depending on the size of the nucleic acid sequence to be mutated. Preferably the specific nucleic acid sequence is from 50 to 50,000 base pairs. It is contemplated that entire vectors containing the nucleic acid encoding the protein of interest may be used in the methods of this invention.

Any specific nucleic acid sequence can be used to produce the population of mutants by the present process. An initial population of the specific nucleic acid sequences having mutations may be created by a number of different known methods, some of which are set forth below.

Error-prone polymerase chain reaction (20,45,46) and cassette mutagenesis (38-44), in which the specific region optimized is replaced with a synthetically mutagenized oligonucleotide can be employed in the invention. Error-prone PCR can be used to mutagenize a mixture of fragments of unknown sequences. These techniques can also be employed under low-fidelity polymerization conditions to introduce a low level of point mutations randomly over a long sequence, or to mutagenize a mixture of fragments of unknown sequence.

Oligonucleotide-directed mutagenesis, which replaces a short sequence with a synthetically mutagenized oligonucleotide may also be employed to generate evolved polynucleotides having improved expression.

Alternatively, nucleic acid or DNA shuffling, which uses a method of *in vitro* or *in vivo*, generally homologous, recombination of pools of nucleic acid fragments or polynucleotides, can be employed to generate polynucleotide molecules having variant sequences of the invention.

Parallel PCR is another method that can be used to evolve polynucleotides for improved expression, function or properties in conventional expression systems, which uses a large number of different PCR reactions that occur in parallel in the same vessel, such that the product of one reaction primes the product of another reaction. Sequences can be randomly mutagenized at various levels by random fragmentation and reassembly of the fragments by mutual priming. Site-specific mutations can be introduced into long sequences by random fragmentation of the template followed by reassembly of the fragments in the presence of mutagenic oligonucleotides.

A particularly useful application of parallel PCR, which can be used in the invention, is called sexual PCR. In sexual PCR, also known as DNA shuffling, parallel PCR is used to perform *in vitro* recombination on a pool of DNA sequences. Sexual PCR can also be used to construct libraries of chimaeras of genes from different species.

The polynucleotide sequences for use in the invention can also be altered by chemical mutagenesis. Chemical mutagens include, for example, sodium bisulfite, nitrous acid, hydroxylamine, hydrazine or formic acid. Other agents which are analogues of nucleotide precursors include nitrosoguanidine, 5-bromouracil, 2-aminopurine, or acridine. Generally, these agents are added to the PCR reaction in place of the nucleotide precursor



thereby mutating the sequence. Intercalating agents such as proflavine, acriflavine, quinacrine and the like can also be used. Random mutagenesis of the polynucleotide sequence can also be achieved by irradiation with X-rays or ultraviolet light, or by  
5     subjecting the polynucleotide to propagation in a host (such as *E. coli*) that is deficient in the normal DNA damage repair function. Generally, plasmid DNA or DNA fragments so mutagenized are introduced into *E. coli* and propagated as a pool or library of mutant plasmids.

Alternatively a mixed population of specific nucleic acids may be found in nature in that they may consist of different alleles of the same gene or the same gene from  
10     different related species (*i.e.*, cognate genes). Alternatively, they may be related DNA sequences found within one species, for example, the oxidase class of genes. Once the mixed population of the specific nucleic acid sequences is generated, the polynucleotides can be used directly or inserted into an appropriate cloning vector, using techniques well-known in the art.

Once the evolved polynucleotide molecules are generated they can be cloned into a suitable vector selected by the skilled artisan according to methods well known in the art. If a mixed population of the specific nucleic acid sequence is cloned into a vector it can be  
15     clonally amplified by inserting each vector into a host cell and allowing the host cell to amplify the vector. The mixed population may be tested to identify the desired  
20     recombinant nucleic acid fragment. The method of selection will depend on the DNA fragment desired. For example, in this invention a DNA fragment which encodes for a protein with improved properties can be determined by tests for functional activity and/or stability of the protein. Such tests are well known in the art.

Using the methods of directed evolution, the invention provides a novel means for  
25     producing functional, and soluble proteins with improved activity toward one or more substrates. The mutants can be expressed in conventional or facile expression systems such as *E. coli*. Conventional tests can be used to determine whether a protein of interest produced from an expression system has improved expression, folding and/or functional properties. For example, to determine whether a polynucleotide subjected to directed  
30     evolution and expressed in a foreign host cell produces a protein with improved activity, one skilled in the art can perform experiments designed to test the functional activity of the

protein. Briefly, the evolved protein can be rapidly screened, and is readily isolated and purified from the expression system or media if secreted. It can then be subjected to assays designed to test functional activity of the particular protein in native form. Such experiments for various proteins are well known in the art, and are discussed in the Examples below.

In one embodiment, the invention contemplates the use polynucleotides encoding for variants of oxidase enzymes. The invention employs directed evolution to generate novel oxidase enzymes, such as GAO, which are expressed in host cells (*e.g. E. coli*) used in an expression system, and which exhibit increased functional activity and increased thermostability.

The invention can also be applied to select or optimize an expression system, including selection of host cells, promoters, and signal sequences. Expression conditions can also be optimized according to the invention.

#### **Directed Evolution of Galactose Oxidase**

Galactose oxidase (EC 1.1.3.9) is an alcohol oxidase enzyme. It oxidizes the hydroxyl group of the sixth carbon of D-galactose. It also oxidizes many other kinds of sugars and alcohols (77, 108, 114, 115, 118-120). Although many fungi produce galactose oxidase, no bacterium has been reported to produce the enzyme (109). There are many reports about galactose oxidase from *Fusarium* ssp. NRRL2903, which is identical to *Dactylium dendroides* ATCC46032 (76-78, 84-86, 88, 95, 99, 108, 110-128). **FIG. 1** The native enzyme is an extra-cellular monomer enzyme and has molecular weight as 67,000. It has one copper (II) ion associated with it active site and related to its oxidation properties. **FIG. 2.** Structure and amino acid residues related to catalysis have been characterized and reported (76, 78, 84-86, 88, 111-113, 116-119).

Galactose oxidase is currently used mainly for assays of D-galactose and D-galactosamine. The enzyme oxidizes the hydroxyl group in the substrate to an aldehyde, which is reactive. Therefore, the enzyme is implicated for use in production of non-natural sugars and derivatives of sugars (118, 119, 95, 99, 128). Hyper-production of galactose oxidase would be useful for a wide variety of applications. The gene of the galactose oxidase has been cloned (110) and expressed in *Escherichia coli* (127). This recombinant

-26-

galactose oxidase was produced as a fused protein with the N-terminal sequence of *LacZ*. However, the yield of the galactose oxidase by this recombinant *E. coli* was not satisfactory.

5 According to the invention, galactose oxidase enzyme (GAO) has been produced in high activity and with improved properties by recombinant techniques in *E. coli*.


The following Examples are understood to be exemplary only, and do not limit the scope of the invention or the appended claims. A person of ordinary skill in the art will appreciate that the invention can be practiced in many forms according to the claims and disclosures here.

10

### EXAMPLE 1

#### Activity Assays for Galactose Oxidase Expressed in *E. coli*

This Example describes assays used for evaluating galactose oxidase activity. Galactose oxidase generates equimolar amounts of hydrogen peroxide by oxidation of a substrate. Colorimetric detection of hydrogen peroxide was therefore used to assay  
15 galactose oxidase activity, employing the following reaction scheme:



This system can be used to assay for oxidation of various substrates, with a very high sensitivity. In the reaction scheme above, an alcohol group of a substrate R is oxidized to produce an aldehyde and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) is released. For example, D-galactose is converted to D-galactohexodialdose plus  $\text{H}_2\text{O}_2$ . The chromogen, in the  
25 presence of hydrogen peroxide and peroxidase enzyme, *e.g.* horseradish peroxidase (HRP), produces a detectable color change, indicating that the reaction catalyzed by GAO has occurred.

### **A. Test Tube Assay**

The activity of galactose oxidase produced in *E. coli* was investigated using fungal galactose oxidase (Sigma, partially purified) as a standard. For detecting hydrogen peroxide with peroxidase (Sigma, type I from horseradish), a chromogen was selected for the GAO assays (85).

#### ***1. Materials***

*Cells.* *E. coli* DH5aMCR (Life Technologies) was used for gene manipulation. *E. coli* BL21(DE3) (Novagen) was used as a host strain for expression of galactose oxidase gene. *E. coli* KY-14478 (SN0029, lacking catalase, Kyowa Hakko Kogyo, Co. Ltd.) was also used for manipulation and expression of genes (157). Competent cells for electroporation were prepared (147).

*Cultivation Media.* Luria-Bertani LB medium (10 g/l bacto tryptone, 5 g/l bacto yeast extract, 10 g/l NaCl, pH 7.5) was used mainly for cultivation of *E. coli* (19). LB plates contained 15 g/l agar in LB medium. Ampicillin (100 mg/l) was added to the medium when required.

*Buffers.* Solutions of sodium phosphate, potassium phosphate and Tris-HCl at various pHs were tested as buffer solution for the assay.

*Chromogens.* Many aromatic compounds can be used as a chromogen for the assay. Four chromogens showed particularly strong color formation; green, orange, red and red, respectively: (a) 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) (85); (b) *o*-anisidine; (c) *o*-dianisidine (127, 123, 121, 122) and (d) *o*-tolidine (114, 119). Their peaks of absorbance were 410 nm, 490 nm, 460 nm and 420 nm.

#### ***2. Methods***

*Cultivation.* Three steps of cultivation were performed for production of galactose oxidase. Recombinant *E. coli* strains were cultivated on LB plate containing ampicillin at 30 °C for 18 hours. The cells were inoculated to LB containing ampicillin. After cultivation at 30 °C for 12 hours, the culture was transferred to a new test tube containing 3 ml LB supplemented with ampicillin. The inoculation rate was 0.5 % of medium. Isopropyl beta-D-thiogalactopyranoside (IPTG) (1 mM) was added for induction after cultivation at 30 °C for 7 hours. Cultivation was continued at 30°C for 6 hours.

*Permeabilization.* Permeable cells were prepared by freezing (-20°C) - thawing

-28-

(4°C) and treatment with 0.5 mg/l lysozyme (Sigma, from chicken egg white) for 30 minutes at 37°C. This pre-treatment for permeabilization was used for assay in evaluation of recombinant galactose oxidase. (Example 3).

5           *Activity assay.* The extract was assayed for galactose oxidase activity. Copper (II) sulfate solution (0.4 mM) was added to the cell-free extract. The cell-free extract was diluted in the buffer solution. Peroxidase (Sigma, type I from horseradish) (10 units/ml) and azinobis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) (2 g/l) were added to the reaction solution. The reaction solution was pre-incubated at 37 °C for 5 minutes. Substrate was added to the solution to be 100 mM. The increase of absorbance (410 nm or 10 405 nm) was measured at 37 °C for 1 minute. Fungal galactose oxidase (Sigma, partially purified) was used as standard for estimation of the activity.

### 3. Results

From these experiments, ABTS was selected as a preferred chromogen for these types of assays, since ABTS formed its color most strongly and sensitively. Moreover, the 15 highest assay sensitivity and lowest background was achieved when using a 100 mM sodium phosphate buffer solution (pH 7.0) for the assay.

Minimum detectable activity of galactose oxidase for this assay system was 0.05 units/ml. Galactose oxidase activity between 0.1 and 1 units/ml was measured quantitatively by photometer at 410 nm or 405 nm.

20           Catalase produced by *E. coli* degrades hydrogen peroxide and may influence the assay. In practice, catalase was not observed to pose a problem, because the activity of the galactose oxidase was greatly higher than that of catalase.

Provided below are additional galactose oxidase screening techniques and/or activity assays, having the following advantages: high specificity for galactose oxidase, 25 high sensitivity, good reproducibility, quantitative measurements, simplicity, flexibility for many substrates, and low cost. One screening system utilizes microplates and the other utilizes membranes. Both systems applies horseradish peroxidase (type I, Sigma) together with a chromogen (ABTS).

## **B. Microplate Screening Method**

The following micro-plate assay has a high sensitivity. Moreover, the enzyme activity can be determined quantitatively. To increase throughput, the method can be automated, for example robotically. This method is particularly suitable as a second screen, after active clones are identified by a more rapid first screen, such as a membrane screen. In experiments using these procedures, the active cultures on the microplate had galactose oxidase activity as indicated by strong green color formation, where each positive well on the microplate was visible as a dark circle. GAO activity was screened in 96-well plates.

Briefly, single colonies were picked from LB-Ampicillin (LB-Ap) agar plates into deep-well plates and grown in LB-Ap. The master plates were duplicated into new deep-well plates containing LB-Ap-1 mM IPTG. Following cultivation at 30 °C, CuSO<sub>4</sub> was added and the cells were lysed with lysozyme and SDS. Cell extracts were reacted with galactose and allyl alcohol using the GAO-HRP coupled assay described above.

### ***1. Methods for Approach A***

Single colonies were picked from Luria-Bertani/100 µg/ml ampicillin (LB-Ap) agar plates into deep-well polypropylene plates (well depth: 2.4 cm; volume: 1 ml; from Beckton Dickinson Labware) and cells were grown for 10 h at 30 °C and 270 rpm in 200 µl LB-Ap. The master plates were duplicated by transferring a 10 µl aliquot to a new deep-well plate containing 300 µl LB-Ap and 1 mM isopropyl-beta-D-thiogalactopyranoside (IPTG) and grown for 12 h at 30 °C and 250 rpm. The cultures were then centrifuged for 10 min at 5000 rpm and the cell pellet was resuspended in 300 µl 100 mM sodium phosphate (NaPi) buffer, pH 7.0 containing 0.4 mM CuSO<sub>4</sub>. Following addition of 0.5 mg/ml lysozyme (35 min at 37 °C) and 2.5% (w/v) SDS (overnight at 4 °C), the GAO activity was assayed using the GAO-horseradish peroxidase (HRP) coupled assay (85). Aliquots of the cell extracts were reacted with galactose (50 mM for generation A1 or 25 mM for generations A2 and A3) and allyl alcohol (0.5 M for all generations) at pH 7.0. The initial rate of H<sub>2</sub>O<sub>2</sub> formation was followed by monitoring the HRP-catalyzed oxidation of 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) (ABTS) at 405 nm. To assay thermostability, the plates were heated at a given temperature for 10 min, cooled down on ice for 10 min, and allowed to reach room temperature for ca. 5 min before the activity toward galactose was measured. The thermostability index was determined from the ratio of the residual GAO activity to the initial activity. Mutants identified as

thermostable were then grown in test tubes (3 ml cultures) and the residual activity after heating at various temperatures was measured at room temperature.

### ***2. Methods for Approach B***

5 Single colonies were picked from LB-Ap agar plates into deep-well polypropylene plates (well depth: 4.4 cm; volume: 2.2 ml; from Qiagen) and cells were grown for 8 h at 30 °C and 270 rpm in 500  $\mu$ l LB-Ap. The master plates were duplicated by transferring a 10  $\mu$ l aliquot to a new deep-well plate containing 500  $\mu$ l LB-Ap-1 mM IPTG and grown overnight at 30 °C and 270 rpm. An aliquot of the culture was transferred to a microtiter plate. Following addition of 0.5 mg/ml (30 min at 37 °C) and 0.4% (w/v) SDS - 0.4 mM  
10  $\text{CuSO}_4$  in 100 mM NaPi buffer, pH 7.0 (4 h at 4 °C), the GAO activity was assayed using the GAO-HRP coupled assay as described above. The galactose concentration used was 25 mM (generations B1 and B2) or 10 mM (generations B3 and B4).

### **C. Membrane Screening Method**

15 Although the micro-plate screening system is highly sensitivity and quantitative, it is desirable to provide a method that contemporaneously assay many more, *e.g.* thousands more clones in a sensitive, accurate, practical and efficient manner. Methods for detection of galactose oxidase activities directly from colonies on agar-plate were examined, but were found to exhibit relatively low sensitivity, low reproducibility, and very slow color formation. Hence, to evaluate very large number of mutants, methods for detection of their  
20 activities directly from colonies on agar-plate or from colonies transferred onto a membrane were examined. These methods were based on colorimetric detection using chromogen and peroxidase, as in the micro-plate screening system.

A suitable screening method using membranes was developed, as is shown here in one optimized form. After transformants formed colonies on an LB-Ap plate (100 mg/l at  
25 30 °C for 18-24 hours), these colonies were transferred to a membrane, *i.e.* they were adsorbed onto the membrane and lifted. for cultivation, the membrane was placed on a new LB-Ap plate (100 mg/l) and was incubated at 30 °C till new colonies were formed on the membrane (6-12 hours). The membrane then was transferred to a new LB-Ap (100 mg/l) plate with 1 mM IPTG, at 30 °C for 6 hours, for induction. Then, the membrane  
30 was put on a filter paper at room temperature, containing lysozyme (0.5 mg/ml), D-galactose (100 mM), ABTS (2 mg/ml), peroxidase (10 units/ml) and  $\text{CuSO}_4$  (0.4 mM). In experiments using these procedures, colonies which had galactose oxidase activities showed as deep purple on the filter paper. This simple method has suitable sensitivity and can be used to evaluate several thousands colonies on one membrane at once.

Several thousands colonies can be evaluate by the screening method with one membrane. This method can be used with an image analyzer, for quantitative determination of activity of each colony. Although the sensitivity of this method is not as high as others, the method is fast and is suitable for a first or initial screening, because many thousands or even millions of colonies can be contemporaneously or rapidly evaluated.

In a preferred embodiment, galactose oxidase activities of colonies which were transferred on a membrane were estimated directly. Colonies, which were formed on LB-Apicillin plate at 30 °C for 24 hours, were transferred onto a membrane (Immobilon NC (HATF), surfactant-free, 45 mm, 82 mm, Millipore). The membrane was put on a new LB-Apicillin plate and was kept at 30 °C for 6~12 hours till colonies were re-formed. Then the membrane was transferred onto an LB-Apicillin plate containing 1 mM IPTG and was incubated for 6 hours at 30 °C. After the membrane was put on filter paper containing 0.5 mg/l lysozyme, 100 mM substrate, 2 mg/ml ABTS, 10 units/ml peroxidase and 0.4 mM CuSO<sub>4</sub> in 100 mM sodium phosphate buffer solution (pH 7.0), the membrane was kept at room temperature for one day, covered with a shield (ABTS is light sensitive). Active colonies showed deep purple color formations.

#### **D. Assay Reagents and Conditions**

Some of the assays herein use CuSO<sub>4</sub>, and/or SDS.

*Copper sulfate* is used to provide copper (II) ion to activate the recombinant (mutant or variant) enzyme. The activity of partially purified galactose oxidase from *D. dendroides* (Sigma) was detected well by using peroxidase and ABTS as described; the addition of copper (II) ion and other cofactors was not needed. (The Sigma enzyme already includes copper ions.) However, experiments with cell-free extracts of recombinant GAO enzymes of the invention showed that almost no activity was detected in the absence of copper (II) ions. Thus, the presence of copper (II) ion is preferred, and without being bound by any theory, is believed to be essential, to activate recombinant GAO enzymes produced by *E. coli* as described herein. Treatment with copper ions at 4 °C is preferred. Copper ion can be provided as copper sulfate (CuSO<sub>4</sub>). Experiments showed that 0.1 mM CuSO<sub>4</sub> is sufficient, whereas 10 mM CuSO<sub>4</sub> slightly inhibited GAO activity. Experiments under assay conditions showed that the preferred concentration of CuSO<sub>4</sub> for activating crude enzyme solution is 0.4 mM. The metal (II) ions of iron, cobalt, nickel, and manganese, and the metal chelator EDTA, did not affect activation of the



recombinant GAO in experiments under assay conditions. Experimental results are shown in FIG. 3. under assay conditions, with and without various metal (II) ions or EDTA.

*Detection enhancers.* In certain assay embodiments, sodium azide or sodium sulfide may be added, for example in an amount of from about 0.01 mM to less than 1 mM. These reagents may enhance detection of GAO activity in some circumstances.

*Detergents.* Addition of detergents to the assay solution also increased the observed activity. Pretreatment with SDS was most effective for increasing the galactose oxidase activity. Treatment with SDS for longer than 12 hours at 4 °C after treatment with lysozyme was suitable for the assay. The galactose oxidase activity did not change within the treatment for 12 to 24 hours at 4 °C. Cultivation, pre-treatment and assay were done as described above.

Other detergents may also be used, as shown in TABLE 1. In these experiments, approximately 0.1 units/ml culture of *E. coli* BL21(DE3)/pGAO-010 and 0.25 units of partially purified galactose oxidase (Sigma) were used. Cells were treated with 0.5 mg/ml lysozyme at 37 °C for 30 minutes. Enzyme and cells were treated with detergents at 4 °C for 1-12 hours. Galactose oxidase activities were assayed using the microplate method described above.

*Cultivation.* Activation on LB-Ap (100 mg/l) plate for 12~24 hours at 30 °C and seed-cultivation in LB-Ap (100 mg/l) 200-500  $\mu$ l/well for 8-10 hours at 30 °C provided uniform growth for cultivation. These conditions are suitable if not necessary for the assay, using the cells, reactants and reagents in these experiments.

The addition of IPTG as an inducer was observed to be necessary for the expression of galactose oxidase on microplate cultivation in these experiments. Initial addition of IPTG to the medium was preferred to the addition of IPTG during cultivation. A cultivation time of 12-16 hours was preferred, and provided superior results (overall higher activities) for almost all recombinant *E. coli* which had a plasmid for expression of galactose oxidase in these experiments. The growth of cells was stopped before 16 hours and the cell extracts had almost no activity at 37 °C. Cultivation at about 30 °C was the optimal temperature in these experiments.

-33-

**TABLE 1**

	SDS [% w/v]	Treatment	0	0.00001	0.000 1	0.001	0.01	0.1
	Relative activity of culture [%]	4°C, 12 hr	100	131	106	146	241	394
	Relative activity of culture [%]	4°C, 1 hr	100	96	118	134	146	189
5	Relative activity of GAO <sup>4</sup> [%]	4°C, 12 hr	100	99	95	103	99	101
	Triton X-100 [% w/v]	Treatment	0	0.00001	0.000 1	0.001	0.01	1
	Relative activity of culture [%]	4°C, 12 hr	100	133	145	190	220	250
	Relative activity of culture [%]	4°C, 1 hr	100	85	95	123	118	149
	Relative activity of GAO [%]	4°C, 12 hr	100	114	114	109	108	98
10	Tween 80 [% w/v]	Treatment	0	0.00001	0.000 1	0.001	0.01	1
	Relative activity of culture [%]	4°C, 12 hr	100	135	113	142	139	140
	Relative activity of culture [%]	4°C, 1 hr	100	159	125	144	122	139
	Relative activity of GAO [%]	4°C, 12 hr	100	120	113	106	114	102
	DMSO [% w/v]	Treatment	0	0.00001	0.000 1	0.001	0.01	1
15	Relative activity of culture [%]	4°C, 12 hr	100	152	140	150	155	152
	Relative activity of culture [%]	4°C, 1 hr	100	169	106	116	116	96
	Relative activity of GAO [%]	4°C, 12 hr	100	104	107	103	97	99

(<sup>1</sup>0.09 units/ml, <sup>2</sup>0.07 units/ml, <sup>3</sup>0.25 units/ml) <sup>4</sup>GAO obtained from SIGMA

**EXAMPLE 2**

20

**Construction of Galactose Oxidase Plasmids**

Plasmids were constructed to express galactose oxidase gene (*gao*) from *Fusarium* ssp. as described below. Several vectors were examined for high expression. Plasmids with different promoters and different sequences between the GAO gene and the ribosome binding site were constructed, as described. *Escherichia coli* strain BL21(DE3) and KY-

14478 were transformed with these plasmids. Permeable cells from test tube cultures were used for the assay.

## **A. Construction of Plasmids**

### ***1. Modified pUC18 Vector Plasmids***

Modified pUC18 plasmids were made to be used for constructing galactose oxidase expression plasmids. As shown in **FIG. 7**, vector pUC18 was digested with the restriction enzyme *Hind*III, blunted with T4 DNA polymerase and ligated with T4 DNA ligase to create vector pUC18-HL lacking the *Hind*III site. pUC18-HL was digested with *Eco*RI, blunted with T4 DNA polymerase and ligated with T4 DNA ligase to create vector pUC18-EHL lacking the *Eco*RI and *Hind*III sites. Similarly, pUC18-EHL was digested with *Pst*I, blunted with T4 DNA polymerase and ligated with T4 DNA ligase to create vector pUC18-EHPL, lacking the *Eco*RI, *Hind*III, and *Pst*I sites.

### ***2. GAO Vector Plasmids***

As shown in **FIG. 8**, plasmid pGAO-010 expressing GAO was made using plasmid pR3. Plasmid pR3 contains the gene for mature galactose oxidase (GAO) fused to the 5' end of the *lacZ* fragment, and was obtained from Dr. Howard K. Kuramitsu (Dept. of Oral Biology, State University of New York, Buffalo, NY). The GAO gene was amplified from pR3 by PCR using primers P-MY001 and P-MY002 in order to introduce a *Hind*III restriction site followed by an ATG initiation codon immediately upstream from the mature GAO sequence, and an *Xba*I site immediately downstream from the stop codon. (Primer sequences are shown in **FIG. 6**). The PCR product was digested with *Hind*III and *Xba*I and ligated into a similarly digested pUC18 vector to create pGAO-001. Plasmid pPLA-001 is a modified pUC18 vector containing a double *lac* promoter. The *lac* promoter from pUC18 was amplified using primers P-MY003 and P-MY004. The PCR product was digested with *Eco*RI and *Hind*III and ligated into a similarly digested pUC18 vector. Following digestion of pGAO-001 with *Hind*III and *Xba*I, pPLA-001 with *Eco*RI and *Hind*III, and pUC18-HL with *Eco*RI and *Xba*I, plasmid pGAO-010 was generated by ligation with T4-DNA ligase.

Another plasmid, pGAO-036, was made by amplifying pGAO-010 using primers P-MY036 and P-MY002. **FIG. 9**. The PCR product was digested with *Kpn*I and *Xba*I and

-35-

ligated with a similarly digested pUC18-EHL to create plasmid pGAO-027. Plasmid pGAO-027 was digested with *KpnI* and *XbaI* and ligated with a similarly digested pUC18-EHPL to create plasmid pGAO-036. This plasmid contains a unique *PstI* site. Plasmid pGAO-036 was used as a for directed evolution experiments described herein.

5 Another plasmid, pGAO-011, was made using similar techniques, as shown in FIG. 10.

### **B. Plasmids and transformation**

Plasmids for expression of galactose oxidase were constructed as described above. The galactose oxidase enzyme was amplified from pR3 (*Fusarium* spp.) by PCR. The *lac* promoter of pUC18 and *T7* promoter of pET-22b(+) (Novagen) were used for expression. 10 In addition to expression as mature sequence of galactose oxidase, expression of the gene as a fused protein with other peptides was examined. The N terminal sequence of *LacZ* was selected to express the galactose oxidase as a fused protein (127). *PelB* leader sequence was also used to produce galactose oxidase in periplasm. Furthermore, His-tag 15 which is useful for purification of recombinant proteins was examined as an additional sequence of the C-terminal of galactose oxidase. *T7* terminator sequence was used for stabilization of expression. Two different *oris* were chosen for replication of plasmid. The copy number of plasmid with *ori* from pUC series is higher than the plasmid with *ori* from pBR series.

20 In more detail, plasmids pUC18, pET-22b(+) (Novagen) and derivatives were used as vector plasmids. Galactose oxidase gene from *Fusarium* spp. was amplified from pR3 according to known techniques. (110, 127). Genes were manipulated according to conventional methods using kits from Qiagen (Valencia, CA). The QIAprep Spin Miniprep Kit, QIAquick Gel Extraction Kit and QIAEX II Gel Extraction Kit, were used 25 reselectively for purification of plasmids from cells, purification of DNA fragments and extraction of DNA fragments from agarose gel. *E. coli* DH5aMCR was transformed with plasmids by treatment with  $\text{CaCl}_2$  (19). Electroporation was used for transformation of *E. coli* BL21(DE3) with plasmids (147, 148).

30 pUC18 and pET-22b(+) (Navagen) were used as vector plasmids. The gene of galactose oxidase from pR3 (127) was used. *lac* promoter from pUC18, *tac* promoter

-36-

from pKK223-3 (Amersham Pharmacia Biotech) and *T7* promoter from pET-22b(+) were selected for expression of the gene. The N terminal sequence of *LacZ* from pUC18, *PelB* leader, His-tag and *T7* terminator sequences from pET-22b(+) were used for production of galactose oxidase. The gene and parts for expression were prepared by PCR. PCR was  
5 done in 100 µl of reaction solution containing PCR buffer (10 mM Tris-HCl, pH 8.5, 50 mM KCl, 2.5 mM MgCl<sub>2</sub>, 0.01 % gelatin), 1 ng of DNA as template, 50 p mole of each primers, 2.5 units of Taq DNA polymerase (Perkin Elmer) and 50 n mole of each dNTPs. DNA fragments were amplified in 30 cycles of 30 seconds at 94 °C, 30 seconds at 50 °C and 60 seconds at 72 °C. PCR products were purified by QIAquick PCR Purification Kit  
10 (Qiagen). Cutting and ligation of DNA by enzymes were according by "molecular cloning" (19). *E. coli* cells were transformed with plasmids by electroporation (Bio-Rad, gene Pulser). QIAprep Spin Miniprep Kit (Qiagen) was used for purification of plasmid from *E. coli* recombinant cells.

Using these strategies, plasmids were designed to produce the galactose oxidase  
15 gene. The plasmids were transformed to *E. coli* DH5αMCR, BL21(DE3) and KY-14478. Representative plasmids are shown diagrammatically in **FIG. 11**, according to the general scheme shown in **FIG. 12**.

Expression of the galactose oxidase gene in all constructed plasmids was controlled by the *lac* operator. Therefore, induction by isopropyl β-D-  
20 thiogalactopyranoside (IPTG) was necessary for production of the enzyme (**FIG. 11**). The expression of galactose oxidase was highest when IPTG (1 mM) was added after cultivation for 7 hours and cells were incubated for 6 more hours. Cultivation at 30 °C gave greatest activity of galactose oxidase per cultivation. Expression of the enzyme was remarkably decreased at 37 °C. Lower temperatures than 27 °C were not suitable in the  
25 experiments because the cells grew very slowly.

Incubation on LB plate at 30 °C for 18 hours and pre-cultivation in LB at 30°C for 12 hours stabilized the main cultivation. The optimal culture conditions were selected as shown above.

### C. Galactose oxidase activity

Galactose oxidase activities of the recombinant *E. coli* were measured (FIG. 11). Some recombinant strains showed much higher activities than the recombinant plasmid pR3. These recombinants hold plasmids which were constructed with *lac* promoter and *ori* from pUC series. Some recombinant *E. coli* with plasmids, pGAO-018 and pGAO-023, expressing the galactose oxidase gene by *T7* promoter did not grow well. Their galactose oxidase activities were not detected. Although some recombinants holding plasmid with *T7* promoter, pGAO-008 and pGAO-009, grow normally, they showed low galactose oxidase activity. From these results, *lac* promoter was suitable for expression of galactose oxidase gene. Furthermore, double *lac* promoter seemed to be stronger than single *lac* promoter in some but not all cases.

For example, plasmid pGAO-025 was designed to have double *lac* promoter and *lacZ-gao* fused gene (FIG. 13). However, galactose oxidase activity of a recombinant with pGAO-025 was almost the same as a recombinant with pGAO-011 which had a single *lac* promoter in KY-1447 cells but was more active than pGAO-011 in BL21(DE3) cells. Triple *lac* promoter was also examined to express the galactose oxidase gene. The effect of triple promoter was about the same as double promoter, e.g. in pGAO-028 and pGAO-010 (FIGS. 15 and 17).

Galactose oxidase which was fused with the N-terminal sequence of *LacZ* or *PelB* leader was produced, as well as non-fused proteins. The activity of galactose oxidase fused with *PelB* leader was not detected without a pre-treatment of cells. Detection of activity of the enzyme required same the pre-treatment of recombinant cells as others. In these experiments GAO was not secreted in the medium, although a secretion signal sequence was present.

Plasmids pGAO-003 and pGAO-005 were designed to produce galactose oxidase in fused form with His-tag at its C-terminal. No galactose oxidase activity was detected from recombinant strains with these plasmids.

Terminator sequence sometimes stabilizes gene expression. In these experiments, introduction of *T7* terminator sequence apparently did not increase GAO expression.

Compare pGAO-020 with pGAO-010 or pGAO-022 with pGAO-017.

*E. coli* DH5aMCR expressed the galactose oxidase gene with these plasmids. However their activities were lower than that of recombinant strains of *E. coli* BL21(DE3) and *E. coli* KY-14478 (data not shown). *E. coli* BL21(DE3) and *E. coli* KY-14478 with plasmid pGAO-010 or pGAO-027 successfully expressed galactose oxidase in high activity. These two plasmids have the same sequence except for one restriction endonuclease site in the vector sequence. Their structure is suitable to express the galactose oxidase in a mature fungal sequence. Consequently, *E. coli* BL21(DE3) and *E. coli* KY-14478 harvesting plasmid pGAO-010, pGAO-027 or their derivatives were used for continued experiments.

**D. Codon Alternation**

Codon alternation of the N-terminal sequence of a gene, without changing the peptide sequence, may cause higher expression of the gene in some cases. Codons of six N-terminal amino acid residues of galactose oxidase were exchanged randomly by PCR with a mixed primer, with the following alternations.

												<b><u>SEQ ID NO:</u></b>
	(M)	A	S	A	P	I	G	S	A			<b>26</b>
Wild-type sequence	ATG	GCC	TCA	GCA	CCT	ATC	GGA	AGC	GCC	...		<b>27</b>
Random Alternation	---	--N	--N	--N	--N	--A	--N	---	---	...		<b>28</b>
							T					
							C					

The galactose oxidase gene of pGAO-010 was replaced with PCR products comprising the galactose oxidase gene with random codon alternation. The plasmids of this library were named pGAO-010M. This random codon alternation of the N-terminal sequence did not cause higher expression (**FIG. 14**), and in many cases GAO activity was reduced. No significant difference was observed when *E. coli* KY-14478 was used as a host strain, compared with *E. coli* BL21(DE3).

**E. Optimization of upper sequence of gao**

The region between the Shine-Dalgarno (“SD”) sequence AGGA and the initiation codon, ATG, is sensitive for efficient RNA translation and has a significant influence on expression of gene. One to three bases were inserted between SD of the *lac* promoter and the ATG of the galactose oxidase gene in pGAO-027 to investigate the impact of altering

the distance between SD and ATG. A change in the length of the region between SD and ATG causes a decrease in galactose oxidase activity when *E. coli* BL21(DE3) was used as a host strain (TABLE 2; SEQ ID NOS: 29-36). The original sequence of pGAO-027 or the one-base extended sequence of pGAO-029 were preferred for expression of the gene.

5 When *E. coli* KY-14478 was used as a host strain, one or two bases extension of the sequence between SD and ATG were preferred to express the gene.

TABLE 2

Plasmid	Sequence between SD and ATG	Promoter	GAO Activity (units/ml)	
			BL21(DE3)	KY-14478
027	... <u>AGGAAAAGCTTATG</u> ...	<i>Plac</i>	19.0	12.5
029	... <u>AGGAAAAAGCTTATG</u> ...		19.1	15.7
030	... <u>AGGAAACAAGCTTATG</u> ...		16.3	15.9
031	... <u>AGGAACAAAGCTTATG</u> ...		14.3	13.1
032	... <u>AGGAAAAGCTTATG</u> ...	<i>Ptac</i>	30.6	52.4
033	... <u>AGGAAAAAGCTTATG</u> ...		25.7	56.2
034	... <u>AGGAAACAAGCTTATG</u> ...		34.6	49.8
035	... <u>AGGAACAAAGCTTATG</u> ...		22.1	38.7

\*Plasmids are designated pGAO-XXX, where XXX is 027 through 035

The *tac* promoter often if not usually expresses genes at higher levels than *lac* promoter. *tac* promoter was prepared from pKK223-3 (Amercham Pharmacia Biotech) by PCR. *lac* promoters of plasmids, pGAO-027, pGAO-29, pGAO-030 and pGAO-031 were replaced with *tac* promoter. Recombinant strains with plasmids using *tac* promoter for expression showed approximately twice as much activity than the recombinant strains using *lac* promoter (TABLE 3). The optimal distance between SD and ATG under the *tac* promoter was almost the same as that under the *lac* promoter in both *E. coli* strains.

20

25



Recombinant strains *E. coli* BL21(DE3)/pGAO-034 and *E. coli* KY-14478/pGAO-033 were considered to be good for expression of galactose oxidase. Optimal culture conditions for these strains were as described above.

#### **F. Properties of recombinant galactose oxidase**

5 Galactose oxidase from *Dactylium dendroides* (*Fusarium* ssp.) and the enzyme from recombinant *E. coli* BL21(DE3)/pGAO-010 differs only in glycosilation; their amino acid sequences are identical.

Substrate specificities of recombinant galactose oxidase from *E. coli* and the enzyme from fungi were compared. Cell-free extract of *E. coli* BL21(DE3)/pGAO-010 was used as a crude recombinant enzyme from *E. coli*. Partially purified galactose oxidase from *Dactylium dendroides* (Sigma, partially purified) was used as fungal enzyme. Substrate specificities of these two enzymes were almost same (**FIG. 15**).

### **EXAMPLE 3**

#### **Optimization of error-prone PCR conditions**

##### **A. General PCR Conditions**

15 Mutation of the galactose oxidase gene (*gao*) was induced by error-prone PCR and according to known techniques (66, 129-133, 136-139). Wild type *gao* on pGAO-027 was replaced by the PCR products which were mutant galactose oxidase genes. The resultant plasmids were named as pGAO-027M. *E. coli* BL21(DE3) was transformed with these plasmids. Almost all transformants carrying error prone PCR products instead of wild type *gao* lost their galactose oxidase activities (**FIG. 7**). Mutations were induced on the whole galactose oxidase gene by error-prone PCR, using conditions "A" of **TABLE 3**. 228 clones were selected randomly from each set of conditions with different manganese concentrations. These clones were cultivated and assayed with micro-plates. More than 25 65 % of transformants lost their galactose oxidase activity, even though manganese ions were not added to the PCR solution.

Various reaction conditions for error-prone PCR were compared, and in particular milder conditions were examined for mutation of the galactose oxidase gene. Conditions

"A" and "C" are the previous conditions of error-prone PCR (above) and normal PCR conditions, respectively. The use of a buffer solution for error-prone PCR (Buffer EP) increased the error rate. Non-uniform composition of dNTPs for error-prone PCR (dNTPs EP) induced mutations in a higher rate than uniform composition of dNTPs for normal PCR (dNTPs normal). *Taq* DNA polymerase from Promega Corporation showed a higher error rate than the enzyme from Perkin Elmer. Since the rate of inactivation was 31 % at most in condition "C" (**FIG. 5**), induction of mutation was not optimal, and may have been insufficient. In **FIG. 5**, mutations were induced in the whole galactose oxidase gene by error-prone PCR using conditions "C" of **TABLE 3**. Activities of 288 clones from each set of conditions with different manganese concentration were estimated using micro-plate screening.

From the alternatives examined in these experiments, Error-prone PCR condition "F" had a suitable frequency of error and was selected to induce mutation on the galactose oxidase gene in further experiments. The composition of buffer solution, the content of dNTPs and thermophilic DNA polymerase each affected the rate of mutation. For example, the difference between the buffer solution for normal PCR and the buffer solution for error-prone PCR was that the EP buffer contained gelatin. Since gelatin is not expected to influence the error rate of the PCR reaction, the observed rate difference may be due to a small difference in the final pH of reaction mixtures with these buffer solutions. More error was induced by non-uniform content of dNTPs for error-prone PCR than uniform content of dNTPs for normal PCR. Selection of the thermophilic DNA polymerase can be significant when optimizing an error-prone PCR experiment, as the particular polymerase may influence the mutation rate.

PCR conditions selected for mutation of the whole galactose oxidase gene in these experiments was milder than previously disclosed conditions (66, 129-133, 136-139). When the PCR conditions described previously were used for error-prone PCR of galactose oxidase gene, the mutation rate was too high, resulting in too many inactive or low activity clones. This result may be related to the fact that the galactose oxidase gene is as much as twice as large as genes previously used for error-prone PCR in the literature. Without being bound by any theory, deadly mutations may be induced more frequently as the target gene becomes larger.

In **TABLE 3**, 96 of 288 clones were selected randomly from each library. Their galactose oxidase activities were estimated by micro-plate screening method. Rates of clones which lost their galactose oxidase activities are show in the table.

**FIG. 4** and **FIG. 5** show the effect of varying amounts of  $MnCl_2$  in these experiments.

In the mutagenesis methods used herein, the error rate is from 1-6 mutations per polynucleotide, preferably 4-6, and most preferably 6. In certain embodiments with more than one round of directed evolution, the error rate may be different from one round to another. For example, the error rate may be about 1-2 mutations per polynucleotide in one round (e.g. a first round), and may be about 4-6 mutations per polynucleotide in another round (e.g. a second round).

**TABLE 3**

PCR conditions						Inactivated clones [%]					
	Buffer	$MgCl_2$	dNTPs	TaqDNA polymerase		$MnCl_2$ 0 mM	$MnCl_2$ 0.1mM	$MnCl_2$ 0.15mM	$MnCl_2$ 0.2mM	$MnCl_2$ 0.4mM	$MnCl_2$ 0.5mM
A	EP	7mM	EP	Promega	50 u/ml	60 (173/288)	69 (199/288)	77 (223/288)	76 (220/288)	90 (258/288)	94 (270/288)
B	EP	7mM	normal	Promega	50 u/ml	55 (53/96)	61 (59/96)				
C	normal	2.5mM	normal	Perkin Elmer	25 u/ml	3 (3/96)	10 (10/96)				
						5 (14/288)	9 (27/288)	10 (29/288)	11 (31/288)	28 (81/288)	31 (90/288)
D	EP	7mM	EP	Perkin Elmer	25 u/ml	45 (43/96)	61 (59/96)				
E	EP	7mM	EP	Perkin Elmer	50 u/ml	39 (37/96)	52 (50/96)				
F	normal	7mM	EP	Perkin Elmer	25 u/ml	23 (22/96)	41 (39/96)				
G	normal	7mM	EP	Promega	50 u/ml	41 (39/96)	52 (50/96)				
H	EP	7mM	normal	Promega	50 u/ml	51 (49/96)	61 (59/96)				

Buffer EP : (x10) 500 mM KCl, 100 mM Tris-HCl (pH 8.3), 0.1% (w/v) gelatin  
 Buffer (normal) : (x10) 500 mM KCl, 100 mM Tris-HCl (pH 8.3)  
 dNTPs EP : 0.2mM dGTP, 0.2 mM dATP, 1 mM dCTP, 1 mM dTTP  
 dNTPs (normal) : 0.5M dGTP, 0.5 mM dATP, 0.5 mM dCTP, 0.5 mM dTTP

**EXAMPLE 4****Production of Galactose Oxidase Mutants**

The directed evolution of galactose oxidase (GAO) is described. GAO variants with increased activity toward allyl alcohol and D-galactose and increased thermostability relative to wild-type have been identified.

**A. Construction of GAO Mutant Libraries**

Plasmid pGAO-036, expressing wild-type GAO, was used as the parent for the directed evolution of GAO (FIG. 9).

Two strategies have been followed for the directed evolution of the enzyme: (A) mutagenesis of the whole GAO gene (bases 1-1917) and (B) mutagenesis of part of the GAO gene (bases 518-1917). In Approach A, two rounds of error-prone PCR (45) have been performed (generations A1 and A2), followed by one round of StEP recombination (generation A3) (139) of four improved variants identified in library A2. In Approach B, four rounds of error-prone PCR (45) have been performed (generations B1 through B4). *E. coli* strain BL21(DE3) (Novagen) was used for the expression of GAO.

***1. Approach A***

Error-prone PCR was carried out in a 100  $\mu$ l reaction mixture containing about 0.3  $\mu$ g plasmid DNA as template, 30 pmol of each primer, 0.2 mM dGTP, 0.2 mM dATP, 1 mM dCTP, 1 mM dTTP, 7 mM MgCl<sub>2</sub>, 0.1 mM MnCl<sub>2</sub>, and 2.5 U *Taq* polymerase (Perkin Elmer) in 10 mM Tris-HCl, 50 mM KCl buffer, pH 8.5. PCR conditions were as follows: 30 cycles of 94 °C for 30 seconds, 50 °C for 30 seconds and 72 °C for 60 seconds. The percentage of inactive clones was between 30 and 50%.

StEP recombination of the four improved variants identified in generation A2 was performed in a 100  $\mu$ l reaction mixture containing about 0.3 mg (total) plasmid DNA as template (prepared by mixing equal amounts of all four plasmids), 10 pmol of each primer, 0.5 mM of each dNTP, 2.5 mM MgCl<sub>2</sub>, and 5 U *Taq* polymerase (Perkin Elmer) in 10 mM Tris-HCl, 50 mM KCl buffer, pH 8.5. PCR conditions were: 95 °C for 3 minutes and 100 cycles of 94 °C for 30 seconds and 58 °C for 10 seconds. The primers used for error-prone PCR and StEP were:

5'-AATTCGAAGCTTATGGCCTCAGCACCTATCGGAAGC-3' (forward) [SEQ. ID. NO. 1]

and 5'-CTTCCTTCTAGATTACTGAGTAACGCGAATCGT-3' (reverse) [SEQ. ID. NO. 2].

5

### **2. Approach B**

Error-prone PCR was carried out in a 100  $\mu$ l reaction mixture containing 10 ng plasmid DNA as template, 50 pmol of each primer, 0.2 mM of each dNTP, 7 mM (generations B1 and B2) or 4 mM MgCl<sub>2</sub> (generations B3 and B4), and 5 U *Taq* polymerase (Boehringer Mannheim) in 10 mM Tris-HCl, 50 mM KCl buffer, pH 8.3. PCR conditions were as follows: 94 °C for 2 minutes and 25 cycles of 94 °C for 30 seconds, 58 °C for 30 seconds and 72 °C for 60 seconds. The primers used were:

10

5'-TTGTTCTGCGGCTGCAGCAATTGAACCG-3' (forward) [SEQ. ID. NO. 8] and 5'-TGCCGGTCGACTCTAGATTACTGAGTAACG-3' (reverse) [SEQ. ID. NO. 9].

The percentage of inactive clones was between 30 and 40%.

15

### **B. Screening of GAO Libraries**

GAO activity was screened in 96-well plates, using the methods of Approaches A and B, respectively, as described in Example 1(D).

### **C. Laboratory Evolution of GAO**

20

The thermal stability curves of selected GAO variants are shown in FIG. 16. Variants were grown in test tubes (3 ml cultures). Following centrifugation and resuspension of the cell pellets in NaPi buffer, pH 7.0 containing CuSO<sub>4</sub>, the cells were lysed. Aliquots of the cell extracts were heated at each temperature for 10 min and then cooled down on ice for 10 min before the residual activity toward D-galactose was determined at room temperature.

25

Results of the laboratory evolution of GAO to increase activity and thermostability are listed in TABLE 4.  $T_{50}$  is an operational measure of stability and is defined as the temperature at which the enzyme loses 50% of its activity following incubation for a set time.

Wild type GAO (pGAO-036) was used as the parent for generation A1 of GAO variants. After screening about 1500 clones, three mutants, 9.16.8D2, 9.16.6C11 and 9.16.16D12, were identified as more active toward allyl alcohol and/or galactose. Clone 9.16.16D12, which was also more thermostable than wild-type GAO, was used to parent generation A2  
5 of GAO variants. Four improved mutants were identified in this library following screening of about 1500 clones: 11.03.6D3, 11.03.10C3, 11.03.10D6 and 11.03.13E12. These clones were more active than the parent toward allyl alcohol and galactose. Clone 11.03.10C3 was substantially more thermostable than the parent, as well. These four improved variants were recombined by StEP in generation A3. Screening of about 2000  
10 clones led to the identification of variant 1.06.20E7 which shows about a 200-fold increased activity toward allyl alcohol and D-galactose and exhibits about a 12 °C higher  $T_{50}$  with respect to wild-type GAO.

Wild-type GAO (pGAO-036) was used as the parent for generation B1 of GAO variants. After screening about 900 clones, variant 1.D4 was identified as more active  
15 toward galactose and used to parent generation B2. Mutant 2.G4 was identified as more active toward galactose in this library following screening of about 1500 clones. Library B3 of GAO variants was generated using 2.G4 as the parent, and clone 3.H7 was identified as an improved variant after screening about 1500 clones. Finally, library 4B was created using 3.H7 as the parent and about 1500 clones were screened. Variant 4.F12 was  
20 identified as about 15-fold more active toward galactose relative to wild-type GAO.

#### **D. Active and Thermostable Mutations**

Most beneficial mutations occur in domains II and III of the GAO gene (residues 156-532 and 533-639, respectively) (87). Mutation V494A, which was identified several times in the screen, is located at the bottom of the active site adjacent to the copper ligand  
25 Y495. Its presence increases the binding affinity for galactose approximately 3-fold. N535D is found in a solvent-exposed loop in domain III. The amino acid substitution G195E is largely responsible for the observed increase in thermostability of variant 1.06.20E7 relative to wild-type. *See FIG. 16 and TABLE 4.*

It should also be noted that a large number of mutations (five in these experiments)  
30 resulted from the substitution of a neutral residue by a negatively charged residue. This

tends to decrease the isoelectric point of GAO in the mutants (the pI of wild type GAO is 12). A decrease in pI is advantageous, in that it may lead to fewer interactions between the mutant GAO and other macromolecules, and lower adhesion to glass. It may also permit increased use of crude galactose oxidase preparations in organic synthesis (107).

5

**TABLE 4**  
Mutations identified in GAO variants and their effects on GAO properties.

10

15

20

GEN	GAO name	nucleotide base substitution	amino acid substitution	relative activity for allyl alcohol*	relative activity for D-galactose	T <sub>50</sub> (°C)
0	pGAO-036	N/A (WT)	N/A (WT)	1.0	1.0	42
A1	9.16.8D2	A1609G	N537D	2.6	4.6	
A1	9.16.6C11	T1481C T1543A	V494A C515S	2.8	1.3	
A1	9.16.16D12	T1481C T408C	V494A P136	3.0	4.9	44
A2	11.03.6D3	T1481C T408C T28C	V494A P136 S10P	6.4	11	
A2	11.03.10C3	T1481C T408C G584A A9C	V494A P136 G195E A3	3.8	9.6	54
A2	11.03.10D6	T1481C T408C A936G A1603G T654C	V494A P136 L312 N535D T218	5.4	11	
A2	11.03.13E12	T1481C T408C A208G	V494A P136 M70V	5.1	9.1	
A3	1.06.20E7	T1481C T28C T408C A208G G584A A1603G	V494A S10P P136 M70V G195E N535D	20	55	54
B1	1.D4	A1237G	N413D		2.4	
B2	2.G4	A1237G T1650A	N413D S550		4.0	
B3	3.H7	A1237G T1650A T1481C	N413D S550 V494A		8.6	
B4	4.F12	A1237G T1650A T1481C T1830A	N413D S550 V494A S610		15.2	

\*Allyl alcohol is oxidized by wild-type GAO at ca. 3% the rate of galactose oxidation.

-47-

Mutations identified at residues A3, L312, T218, P136, S550 and S610 are synonymous and, without being bound by theory, the observed increase in activity is probably due to higher expression of GAO in *E. coli*. Given the low expression level of recombinant wild-type GAO (less than 3% of total intracellular protein as determined by SDS-PAGE), this is a much needed improvement.

The variants identified also exhibit increased activity toward a variety of GAO substrates. Mutant 1.06.20E7 is about 200-fold more active toward 3-pyridylcarbinol and mutant 4.F12 is about 15-fold more active toward glycerol, xylitol, beta-D-lactose, and IPTG.

The sequences of representative mutants of the invention identified in **TABLE 4** are shown in **FIGS. 17-28**.

As shown in the above Examples, the galactose oxidase gene can be expressed in *E. coli* in relatively high yield, with an increased activity toward at least one substrate. In certain embodiments the activity is greatly increased toward several substrates. In certain embodiments the mutants exhibit thermostability.

The inducible promoters *Plac* or *Ptac* were effective for expression of the galactose oxidase gene and are preferred. Much higher expression may be possible when other strong promoters are used. However, some strong promoters may be counterproductive. For example, *E. coli* did not grow well when *T7* promoter, which is stronger than *lac* promoter, was used for expression of the galactose oxidase gene. Double promoters of two *Plac-Plac* or *Plac-Ptac* were selected to express the galactose oxidase gene. Double promoters express the gene stronger than single promoter as compared pGAO-025 and pGAO-011. Triple promoters expressed the gene as well as double promoters. Upper promoter of double promoters seemed to be less effective than lower promoter in the Examples. Therefore, double promoters of *Plac-Plac* or *Plac-Ptac* are preferred. Induction of gene by IPTG was necessary when *lac* promoter or *tac* promoter was used. Timing of induction and incubation time after that were optimized.

In these experiments the fused form of GAO (*i.e.* as a fusion protein with *lacZ*) was not found to provide advantages, and was not necessary to express the fungal gene.

Galactose oxidase generally had reduced activity or lost its activity when codons were alternated or when it was produced as fused enzyme with His-tag. Culture condition was also important for production of the enzyme.



-48-

Galactose oxidase was engineered by directed evolution to produce more active variants toward natural and additional substrates. Activity of the present mutants was as high as about 65 times that of wild-type GAO. Mutants of the invention also are more stable than wild-type, and in particular exhibit improved thermal stability.

**BIBLIOGRAPHY**

1. Cleland, J. L.; Wang, D. I. C. *Bio/Technology* **8**, 1274 (1990).
2. Bernarderz-Clark, E. D.; Georgiou, G. Inclusion Bodies and Recovery of Proteins from the Aggregated States. In *Protein Refolding*; Bernarderz-Clark, E. D., Georgiou, G., Eds.; ACS: Washington, D. C. p. 1-20 (1990)
3. Thatcher, D. R.; Hitchcock, A. Protein Folding in Biotechnology. In *Mechanisms of Protein Folding*; Pain, R. H., Ed.; IRL Press: Oxford p. 229-261 (1994).
4. Parekh, R.; Forrester, K.; Wittrup, D. *Protein Expres. Purif.* **6**, 537 (1995).
5. Arnold, F. H. *Accounts Chem. Res.* **31**, 125 (1998).
6. Mitraki, A.; King, J. *FEBS Lett.* **307**, 20 (1992).
7. Zhang, J. X.; Goldenberg, D. P. *Biochemistry* **32**, 14075 (1993).
8. Wetzel, R.; Perry, L. P.; Veilleux, C. *Bio/Technology* **9**, 731 (1991).
9. Knappik, A.; Pluckthun, A. *Protein Eng.* **8**, 81 (1995).
10. Crameri, A.; Whiteborn, E. A.; Tate, E.; Stemmer, W. P. C. *Nature Biotechnol.* **14**, 315 (1996).
11. Tams, J. W.; Welinder, K. G. *FEBS Lett.* **421**, 234 (1998).
12. Ortlepp, S. A.; Pollard-Knight, D.; Chiswell, D. J. *J. Biotechnol.* **11**, 353 (1989).
13. Smith, A. T. et al. *J. Biol. Chem.* **265**, 13335-13343 (1990).
14. Egorov, A. M.; Gazaryan, I. G.; Savelyev, S. V.; Fechina, V. A.; Veryovkin, A. N.; Kim, B. B. *Ann. N. Y. Acad. Sci.* **646**, 35 (1991).
15. Moore, J. C.; Arnold, F. H. *Nature Biotechnol.* **14**, 458 (1996).
16. Fitzgerald, M. M.; Churchill, M. J.; McRee, D. E.; Goodin, D. B. *Biochemistry* **33**, 3807 (1994).
17. Goodin, D. B.; Davidson, M. G.; Roe, J. A.; Mauk, A. G.; Smith, M. *Biochemistry* **30**, 4953 (1991)
18. De Sutter, K.; Hostens, K.; Vandekerckhove, J.; Fiers, W. *GENE* **141**, 163 (1994).
19. Sambrook, J.; Fritsch, E. F.; Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory: New York (1989).

20. Caldwell, R. C.; Joyce, G. F. *PCR Methods Applic.* **2**, 28 (1992).
21. Beckman, R. A.; Mildvan, A. S.; Loeb, L. A. *Biochemistry* **24**, 5810 (1994).
22. Shafikhani, S.; Siegel, R. A.; Ferrari, E.; Schellenberger, V. *Biotechniques* **23**, 304 (1997).
23. Stemmer, W. P. C. *Proc. Natl. Acad. Sci. USA* **91**, 10747-51 (1994).
24. Zhao, H. M.; Arnold, F. H. *Nucleic Acids Res.* **25**, 1307 (1997).
25. Carbon, J.; Clarke, L.; Ilgen, C.; Ratzkin, B. The Construction and Use of Hybrid Plasmid Gene Banks in *Escherichia coli*. In *Recombinant Molecules: Impact on Science and Society*; Beers, R. F. J., Bassett, E. G., Eds; Raven Press: New York, pp 355-378 (1977).
26. Shindler, J. S.; Childs, R. E.; Bardsley, W. G. *Eur. J. Biochem.* **65**, 325 (1976).
27. Lei, S. P.; Lin, H. C.; Wang, S. S.; Callaway, J.; Wilcox, G. *J. Bacteriol.* **169**, 4379 (1987).
28. Better, M.; Chang, C. P.; Robinson, R. R.; Horwitz, A. H. *Science* **240**, 1041 (1988).
29. Goshorn, S. C.; Svensson, H. R.; Kerr, D. E.; Somerville, J. E.; Senter, P. D.; Fell, H. P. *Cancer Res.* **53**, 2123 (1993).
30. Rathore, D.; Nayak, S. K.; Batra, J. K. *FEBS Lett.* **392**, 259 (1996).
31. Studier, F. W.; Rosenberg, A. H.; Dunn, J. J.; Dubendorff, J. W. *Meth. Enzymol.* **185**, 60 (1990).
32. Ostermeier, M.; Desutter, K.; Georgiou, G. *Eukaryotic J. Biol. Chem.* **271**, 10616 (1996).
33. Savenkova, M. I.; Kuo, J. M.; Ortiz de Montellano, P. R. *Biochemistry* **37**, 10828 (1998).
34. Gajhede, M.; Schuller, D. J.; Henriksen, A.; Smith, A. T.; Poulos, T. L. *Nature Struct. Biol.* **4**, 1032 (1997).
35. Anfinsen, C. B. *Science* **181**, 223 (1973).
36. Schein, C. H. *Bio/Technology* **8**, 308 (1990).
37. Martineau, P.; Jones, P.; Winter, G. *J. Mol. Biol.* **20**, 117 (1998).

38. Stemmer, W. P. C. et al., *Biotechniques* **14**, 256 (1992).
39. Arkin, A. and Youvan, D. C. *Proc. Natl. Acad. Sci. USA* **89**, 7811 (1992).
40. Oliphant, A. R. et al., *Gene* **44**, 177 (1986).
41. Hermes, J. D. et al., *Proc. Natl. Acad. Sci. USA* **87**, 696 (1990).
42. Delagrave et al. *Protein Engineering* **6**, 327 (1993).
43. Delagrave et al. *Bio/Technology* **11**, 1548 (1993)
44. Goldman, E. R. and Youvan D. C. *Bio/Technology* **10**, 1557 (1992).
45. Leung, D. W. et al., *Technique* **1**, (1989).
46. Gramm, H. et al., *Proc. Natl. Acad. Sci. USA* **89**, 3576 (1992).
47. Castelli, M. C. et al., *Gene* **142**, 113 (1994).
48. Gietz, D., Schiestl, R. H., Willems, A., Woods, R. A., *Yeast* **11**, 355 (1995).
49. Welinder, K. G., *Eur J. Biochem* **96**, 483-502 (1979).
50. Sirotkin, K. J. *Theor. Biol.* **123**, 261 (1986).
51. Glover, D.M. (ed.), 1985, DNA Cloning: A Practical Approach, MRL Press, Ltd., Oxford, U.K. Vol. I, II.
52. Schatz, P.J. Et al., *Annu. Rev. Genet.* **24**, 215-248 (1990).
53. Gussow, D. & Clackson, T. (1989) *Nucleic Acids Res.* , **17**, 4000-4000.
54. Gillam, E.M., Guo, Z., Martin, M.V., Jenkins, C.M. & Guengerich, F.P. (1995) *Arch. Biochem. Biophys.*, **319**, 540-550.
55. Guengerich, F.P., Martin, M.V., Guo, Z. & Chun, Y.J. (1996) *Meth. Enzymol.*, **272**, 35-44.
56. Joo, H., Lin, Z. & Arnold, F.H. (1999) *Nature* , 399, 670-673.
57. Joo, H., Arisawa, A., Lin, Z. & Arnold, F.H. (1999) *Chem. Biol.*, **6**, 669-706.
58. Khosla, C., Curtis, JE, Demodena, J., Rinas, U. & Bailey, JE (1990) *Bio/Technology*, **8**, 849-853

59. Bradford, M. (1978) *Anal. Biochem.*, **72**, 248-254
60. Dunford, H.B. (1991) *Peroxidases in Chemistry and Biology*, Vol 2. pp. 1-24
61. Miele, R.G., Prorok, M., Costa V.A. & Castellino F.J. (1999) *J.Biol.Chem.*, **274**, 7769-7776
62. Gazaryan, I.G. (1994) *LABPV Newsletters*, **4**, 8-15
63. Rodriguez-Lopez, J.N., Smith, A.T., and Thorneley, R.N.F. (1995) *J.Biol.Chem.*, **271**, 4023-4030
64. Haschke, R.H. & Friedhoff, J.M. (1978) *Biochim. Biophys. Res. Commun.*, **90**, 1039-1042
65. Smith, A.T., & Veitch, N.C. (1998) *Curr.Opin.Chem.Biol.*, **2**, 269-278
66. Chen, K. & Arnold, F.H. (1993) *Proc. Natl. Acad. Sci. USA*, **90**, 5618-5622
67. Giver, L., Gershenson, A., Freskgard, P-O. & Arnold, F.H. (1998) *Proc. Natl. Acad. Sci. USA*, **95**, 12809-12813
68. Yano, T., Oue, S. & Kagamiyama, H. (1998) *Proc. Natl. Acad. Sci. USA*, **95**, 5511-5515
69. Cherry, J.R., Lamsa, M.H., Schneider, P., Vind, J., Svendsen, A., Jones, A., & Pederson, A.H. (1999) *Nat. Biotechnol.*, **17**, 379-384
70. Romanos, M.A., Scorer, C.A. & Clare J.J. (1992) *Yeast*, **8**, 423-488
71. Fiedler, K. & Simons, K. (1995) *Cell*, **81**, 309-312
72. Nagayama, Y., Namba, H., Yokoyama, N., Yamashita, S. & Niwa, M. (1998) *J. Biol. Chem.*, **273**, 33423-33428
73. Helenius, A. (1994) *Mol. Biol. Cell.*, **5**, 253-265
74. Miyazaki, K., Wintrode, P.L., Grayling, R.A., Rubingh, D.N. & Arnold, F.H. (2000) *J.Mol.Biol.* in press.
75. Zhao, H.M. & Arnold, F.H. (1999) *Protein Eng.* **12**, 47-53
76. Borman, C. D., Saysell, C. G., and Sykes, A. G. (1997) *J. Biol. Inorg. Chem.* **2**, 480-487.

77. Mendonca, M. H., and Zancan, G. T. (1987) *Arch. Biochem. Biophys.* **272**, 507-514.
78. Ito, N., Phillips, S. E. V., Yadav, K. D. S., and Knowles, P. F. (1994) *J. Mol. Biol.* **238**, 794-814.
79. Whitaker, M. W., and Whitaker, J. W. (1988) *J. Biol. Chem.* **263**, 6074-6080.
80. Avigad, G., Amaral D., Asensio, C., and Horecker, B. L. (1962) *J. Biol. Chem.* **237**, 2736-2743.
81. Amaral, D., Kelly-Falcoz, F., and Horecker, B. L. (1966) *Methods Enzymol.* **9**, 87-2.
82. Klibanov, A. M., Alberti, B. N., and Marletta, M. A. (1982) *Biochem. Biophys. Res. Commun.* **108**, 804-808.
83. Maradufu, A., Cree, G. M., and Perlin, A. S. (1971) *Canad. J. Chem.* **49**, 3429-3436.
84. Wachter, R. M., and Branchaud, B. L. (1996) *J. Am. Chem. Soc.* **118**, 2782-2789.
85. Baron, A. J., Stevens, C., Wilmot, C., Seneviratne, K. D., Blakeley, V., Dooley, D. M., Phillips, S. E. V., Knowles, P. F., and McPherson, M. J. (1994) *J. Biol. Chem.* **269**, 25095-25105.
86. Ito, N., Phillips, S. E. V., Stevens, C., Ogel, Z. B., McPherson, M. J., Keen, J. N., Yadav, K. D. S., and Knowles, P. F. (1991) *Nature* **350**, 87-90.
87. Ito, N., Knowles, P. F., and Phillips, S. E. V. (1995) *Methods Enzymol.* **258**, 235-262.
88. Reynolds, M. P., Baron, A. J., Wilmot, C. M., Vinecombe, E., Stevens, C., Phillips, S. E. V., Knowles, P. F., and McPherson, M. J. (1997) *J. Biol. Inorg. Chem.* **2**, 327-335.
89. Tkac, J., Gemeiner, P., and Sturdik, E. (1999) *Biotechnology Techniques* **13**, 931-936.
90. Vega, F. A., Nunez, C. G., Weigel, B., Hitzmann, B., and Ricci, J. C. D. (1998) *Anal. Chim. Acta* **373**, 57-62.
91. Adanyi, N., Szabo, E. E., and Varadi, M. (1999) *European Food Research and Technology* **209**, 220-226.
92. Mannino, S., Cosio, M. S., and Buratti, S. (1999) *Italian Journal of Food Science* **11**, 57-65.

93. Szabo, E. E., Adanyi, N., and Varadi, M. (1996) *Biosensors & Bioelectronics* **11**, 1051-1058.
94. Vrbova, E., Peckova, J., and Marek, M. (1992) *Collection of Czechoslovak Chemical Communications* **57**, 2287-2294.
95. Root, R. L., Durrwachter, J. R., and Wong, C. H., (1985) *J. Am. Chem. Soc.* **107**, 2997-2999.
96. Dahlhoff, W. V., Idelmann, P., and Koster, R., (1980) *Angew. Chem. Int. Ed. Engl.* **19**, 546-547.
97. Koster, R., Idelmann, P., and Dahlhoff, W. V. (1982) *Synthesis*, 650-652.
98. Liu, X. C., and Dordick, J. S. (1999) *J. Am. Chem. Soc.* **121**, 466-467.
99. Mazur, A. W., and Hiler, G. D. (1997) *J. Org. Chem.* **62**, 4471-4475.
100. Arts, S. J. H. F., Mombarg, E. J. M., vanBekkum, H., and Sheldon, R. A. (1997) *Synthesis-Stuttgart* **6**, 597-610.
101. Yang, G. Y., and Shamsuddin, A. M. (1996) *Histol. Histopathol.* **11**, 801-806.
102. Said, I. T., Shamsuddin, A.M., Sherief, M. A., Taleb, S. G., Aref, W. F., and Kumar, D.(1999) *Histol. Histopathol.* **14**, 351-357.
103. Marrs, B. L. in *IBC's Fifth Annual World Congress on Enzyme Technologies* (2000) Las Vegas, NV.
104. Martin, I. G., Macias, E. M., Sanchez, J. S., and Rivera, B. G. (1998) *Food Chemistry* **61**, 281-286.
105. Calderhead, D. M., and Lienhard, G. E. (1988) *J. Biol. Chem.* **263**, 12171-12174.
106. Gahmberg, C. G., and Tolvanen, M. (1994) *Methods Enzymol.* **230**, 32-44.
107. Mazur, A. in *Enzymes in Carbohydrate Synthesis* (1991) Bednarski, M. D. and Simon, E. S. Eds, pp. 99-110.
108. Tressel, P. (1980) Ph.D. Thesis, State University of New York, Buffalo.
109. Aisaka, K., and Terada, O. (1981) Production of galactose oxidase by *Gibberella fujikuroi*., *Agric. Biol. Chem.*, 45(10), 2311-2316.
110. McPherson, M.J., Ogel, Z.B., Stevens, C., Yadav, K.D.S., Keen, J.N., and Knowles, P.F. (1992) Galactose oxidase of *Dactylium dendroides*., *J. Biol. Chem.*, 267(12), 8146-8152.

111. McPherson, M.J., Stevens, C., Baron, A.J., Ogel, Z.B., Sneviratne, K., Wilmot, C., Ito, N., Brocklebank, I., Phillips, S.E.V., and Knowles, P.F. (1993) Galactose oxidase: Molecular analysis and mutagenesis studies., *Biochem. Soc. Transact.*, 21, 752-756.
112. Saysell, C.G., Barna, T., Borman, C.D., Baron, A.J., McPherson, M.J., and Sykes, A.G. (1997) Properties of the Trp290His variant of *Fusarium* NRRL 2903 galactose oxidase: Interaction of GOasesemi state with different buffer, its redox activity and ability to bind azide., *JBIC*, 2, 702-709.
113. Whittaker, M.M., Ballou, D.P., and Whittaker, J.W. (1998) Kinetic isotope effects as probes of the mechanism of galactose oxidase., *Biochemistry*, 37, 8426-8436.
114. Maradufu, A., et al. (1974) A non-hydrogen-binding role for the 4-hydroxyl group of D-galactose in its reaction with D-galactose oxidase., *Carbohydr. Res.* 32, 93-99.
115. Schlegel, R.A., Gerbeck, C.M., and Montgomery, R. (1968) Substrate specificity of D-galactose oxidase., *Carbohydr. Res.*, 7, 193-199.
116. Hamilton, G.A., de Jersey, J., and Adolf, P.K. (1973) Galactose oxidase : The complexities of a simple enzyme., in King, T.E., et al. Eds., *Oxidases and related redox enzyme*, University Park Press, Baltimore, MD, 103-124.
117. Hamilton, F.A., Adolf, P.K., de Jersey, J., DuBois, G.C., Dyrkacz, G.R., and Libby, D. (1978) Trivalent copper, superoxide, and galactose oxidase., *J. Am. Chem. Soc.*, 100(6), 1899-1912.
118. Arts, S.J.H.F., Monbarg, E.J.M., van Bekkum, H., and Sheldon, R.A. (1997) Hydrogen peroxide and oxygen in catalytic oxidation of carbohydrates and related compounds., *Synthesis*, June, 1997, 597-613.
119. Kosman, D.J. (1984) Galactose oxidase., in Lontie, R., Eds., *Copper proteins and copper enzymes*. Vol. 2., CRC Press, Boca Raton, Fla., 1-26.
120. Mendonca, M.H., and Zancan, G.T. (1988) Role of carbohydrate content on the properties of galactose oxidase from *Dactylium dendroides*., *Arch. Biochem. Biophys.*, 266(2), 427-434.
121. Koroleva, O.V., Rabinovich, M.L., Buglova, T.T., and Yaropolov, A.I. (1983) Some properties of galactose oxidase from *Fusarium graminearum*., *Prikl. Biokhim. Mikrobiol.*, 19(5), 632-637.
122. Avigad, G. (1985) Oxidation rates of some desialylated glycoproteins by galactose oxidase., *Arch. Biochem. Biophys.*, 239(2), 531-537.



123. Tressel, P.S., and Kosman, D.J. (1989) Galactose oxidase from *Dactylium dendroides*., *Methods Enzymol.*, 89, 163-171.
124. Tressel, P., and Kosman, D.J. (1980) A simple purification procedure for galactose oxidase., *Anal. Biochem.*, 105, 150-153.
125. Avigad, G. (1978) An NADH coupled assay system for galactose oxidase., *Anal. Biochem.*, 86, 470-476.
126. Kiba, N., Shitara, K., and Furusawa, M. (1989) A post-column co-immobilized galactose oxidase/peroxidase reactor for fluorometric detection of saccharides in a liquid chromatographic system., *J. Chromatogr.*, 463, 183-187.
127. Lis, M., and Kuramitsu, H.K. (1997) Galactose oxidase-glucan binding domain fusion proteins as targeting inhibitors of dental plaque bacteria., *Antimicrob. Agents Chemother.*, 41(5), 999-1003.
128. Martin, B.D., Linhardt, R.J., and Dordick, J.S. (1998) Highly swelling hydrogels from ordered galactose-based polyacrylates., *Biomaterials*, 19(1-3), 69-76.
129. Arnold, F.H. (1993) Engineering proteins for nonnatural environments., *FEBS J.*, 7, 744-749.
130. Moore, J.C., and Arnold, F.H. (1996) Directed evolution of a para-nitrobenzyl esterase for aqueous-organic solvents., *Nature Biotechnol.*, 14, 458-467.
131. You, L., and Arnold, F.H. (1996) Directed evolution of subtilisin E in *Bacillus subtilis* to enhance total activity in aqueous dimethylformamide., *Protein Eng.*, 9, 77-83.
132. Moore, J.C., Jin, H.M., Kuchner, O., and Arnold, F.H. (1997) Strategies for the in vitro evolution of protein function : Enzyme evolution by random recombination of improved sequences., *J. Mol. Biol.*, 272, 336-347.
133. Arnold, F.H., and Moore, J.C. (1997) Optimization industrial enzymes by directed evolution., *Adv. Biochem. Eng. Biotechnol.*, 58, 1-14.
134. Zhao H., and Arnold, F.H. (1997) Optimization of DNA shuffling for high fidelity recombination., *Nucleic acids Res.*, 25(6), 1307-1308.
135. Zhao, H., and Arnold, F.H. (1997) Functional and nonfunctional mutation distinguished by random recombination of homologous gene., *PNAS. USA*, 94, 7997-800.
136. Giver, L., Gershenson, A., Freskgard, P.O., and Arnold, F.H. (1998) Directed evolution of a thermostable esterase., *Proc. Natl. Acad. Sci. USA*, 95, 1209-12813.

137. Giver, L., and Arnold, F.H. (1998) Combinatorial protein design by in vitro recombination., *Curr. Opin. Chem. Biol.*, 2, 335-338.
138. Arnold, F.H. (1998) Design by directed evolution., *Accounts Chem. Res.*, 31, 125-131.
139. Zhao, H.M., Giver, L., Shao, Z.X., Affholter, J.A., and Arnold, F.H. (1998) Molecular evolution by staggered extension process (StEP) in vitro recombination., *Nature Biotechnol.*, 16, 258-261.
140. Shao, Z.X., Zhao, H.M., Giver, L., and Arnold, F.H. (1998) Random-priming in vitro recombination, *Nucleic Acids Res.*, 26, 681-683.
141. Kuchner, O., and Arnold, F.H. (1997) Directed evolution of enzyme catalysts., *Trends Biotechnol.*, 15, 523-530.
142. Stemmer, W.P.C. (1994) Rapid evolution of a protein in vitro by DNA shuffling., *Nature*, 370, 389-391.
143. Stemmer, W.P.C. (1994) Methods for in vitro recombination., U.S. Patent 5,605,793.
144. Cramer, A., Whitehorn, E.A., Tate, E., and Stemmer, W.P.C. (1996) Improved green fluorescent protein by molecular evolution using DNA shuffling., *Nature Biotechnol.*, 14, 315-318.
145. Zhang, J.H., Dawes, G., and Stemmer, W.P.C. (1997) Directed evolution of fucosidase from a galactosidase by DNA shuffling and screening., *Proc. Natl. Acad. Sci. USA*, 94, 4504-4509.
146. Cramer, A., Dawes, G., Rodriguez, E. Jr., Silver, S., and Stemmer, W.P.C. (1997) Molecular evolution of an arsenate detoxification pathway by DNA shuffling., *Nature Biotechnol.*, 15, 436-438.
147. Dower, W.J., Miller, J.F., and Ragsdale, C.W. (1988) High Efficiency transformation of *E. coli* by high voltage electroporation., *Nucleic Acids Res.*, 16(13), 6127-6145.
148. Calvin, N.M., and Hanawalt, P.C. (1988) High-efficiency transformation of bacteria cells by electroporation., *J. Bacteriol.*, 170(6), 2796-2801.
149. U.S. Patent No. 5,741,691
150. U.S. Patent No. 5,811,238
151. U.S. Patent No. 5,830,721

-58-

152. International Application WO 98/42832
153. International Application WO 95/22625
154. International Application WO 97/20078
156. International Application WO 95/41653
157. Nakagawa, S., Ishino, S., and Teshiba, S. (1996) Construction of catalase deficient *Escherichia coli* strains for production of uricase. *Biosci. Biotech. Biochem.*, 60(3), 415-420.

CLAIMSWhat is claimed is:

- 1       **1.**       A method of obtaining and improving the production of a functional galactose oxidase  
2               polypeptide by a host cell comprising the steps of:  
3               (a)       providing at least one parent galactose oxidase polynucleotide encoding a  
4                       parent galactose oxidase polypeptide,  
5               (b)       altering the nucleotide sequence of the parent polynucleotide by random  
6                       mutagenesis to produce a population of mutant polypeptides;  
7               (c)       transforming host cells to express the mutant polypeptides;  
8               (d)       screening for first-generation functional mutants produced by the host cells  
9                       and having at least one modified property;  
10              (e)       selecting at least one polynucleotide encoding a first-generation mutant as a  
11                       parent polynucleotide; and  
12              (f)       repeating a round of altering, transforming and screening steps at least once  
13                       to obtain at least one other generation of one or more mutants.
  
- 1       **2.**       The method of claim 1 wherein the method of random mutagenesis comprises an  
2               error-prone polymerase chain reaction.
  
- 1       **3.**       The method of claim 2, wherein the error-prone polymerase chain reaction employs  
2               unbalanced nucleotide concentrations.
  
- 1       **4.**       The method of claim 2, wherein the error-prone polymerase chain reaction employs  
2               manganese ions in a concentration of about 0 to about 500  $\mu\text{M}$ .
  
- 1       **5.**       The method of claim 2, wherein the error-prone polymerase chain reaction employs  
2               manganese ions in a concentration of about 100  $\mu\text{M}$ .
  
- 1       **6.**       The method of claim 2, wherein the polymerase chain reaction generates an error rate  
2               of about 1-2 mutations per polynucleotide.

- 1 7. The method of claim 2, wherein the polymerase chain reaction generates an error rate  
2 of up to about six mutations per polynucleotide.
- 1 8. The method of claim 1, wherein at least one of the altering, transforming and  
2 screening steps are changed in at least one repeated round.
- 1 9. The method of claim 1, wherein the conditions for random mutagenesis in at least one  
2 repeated round of altering, transforming and screening are different from the  
3 conditions in any other round of altering, transforming and screening.
- 1 10. The method of claim 1, wherein the host cells in at least one repeated round of  
2 altering, transforming and screening are different from the host cells in any other  
3 round of altering, transforming and screening.
- 1 11. The method of claim 10, wherein the host cells in at least one round are bacterial cells.
- 1 12. The method of claim 11, wherein the bacterial cells are *E. coli* cells.
- 1 13. The method of claim 1, wherein at least one round of altering, transforming and  
2 screening comprises screening for a property of the polypeptide that was not screened  
3 for in another round of altering, transforming and screening.
- 1 14. The method of claim 13, wherein at least one property is selected from the group  
2 consisting of enzyme activity, enzyme selectivity, enzyme stability, and enzyme yield.
- 1 15. The method of claim 1, wherein each screening step comprises screening for one or  
2 more of the biological activity of the polypeptide, the selectivity of the polypeptide,  
3 the stability of the polypeptide, and the yield of expressed polypeptide.

- 1 16. The method of claim 2, wherein the error rate in the altering step of at least one round  
2 of altering, transforming and screening is about 1-2 mutations per polynucleotide, and  
3 the error rate in the altering step of at least one other round is about 4-6 mutations per  
4 polynucleotide.
- 1 17. The method of claim 2, wherein the polymerase chain reaction employs manganese  
2 ions in a concentration of about 0.35 mM.
- 1 18. The method of claim 1, wherein screening comprises pre-screening for mutant  
2 colonies using nitrocellulose membranes.
- 1 19. A polynucleotide evolved according to the method of claim 8
- 1 20. A polynucleotide encoding for a galactose oxidase which has a mutation in at least one  
2 amino acid selected from the group consisting of A3, S10, M70, P136, G195, T218,  
3 L312, N413, V494, C515, N535, N537, S550, and S610.
- 1 21. A polynucleotide encoding for a galactose oxidase which has at least one amino acid  
2 mutation selected from the group consisting of S10P, M70V, G195E, N413D, V494A,  
3 C515S, N535D, and N537D.
- 1 22. A polynucleotide encoding for a galactose oxidase which has the amino acid mutation  
2 N537D.
- 1 23. A polynucleotide encoding for a galactose oxidase which has the amino acid mutation  
2 V494A.
- 1 24. The polynucleotide of claim 23, further comprising the amino acid mutation C515S.
- 1 25. The polynucleotide of claim 23, further comprising the amino acid mutation S10P.

- 1 26. The polynucleotide of claim 23, further comprising a silent mutation at P136.
- 1 27. The polynucleotide of claim 25, further comprising a silent mutation at P136.
- 1 28. The polynucleotide of claim 23, further comprising the amino acid mutation G195E.
- 1 29. The polynucleotide of claim 28, further comprising a silent mutation in at least one of  
2 A3 and P136.
- 1 30. The polynucleotide of claim 23, further comprising the amino acid mutation N535D.
- 1 31. The polynucleotide of claim 30, further comprising a silent mutation in at least one of  
2 P136, L312, and T218.
- 1 32. The polynucleotide of claim 23, further comprising the amino acid mutation M70V.
- 1 33. The polynucleotide of claim 32, further comprising a silent mutation at P136.
- 1 34. A polynucleotide encoding for a galactose oxidase which has the amino acid  
2 mutations V494A, S10P, M70V, G195E and N535D.
- 1 35. The polynucleotide of claim 34, further comprising a silent mutation at P136.
- 1 36. A polynucleotide encoding for a galactose oxidase which has the amino acid mutation  
2 N413D.
- 1 37. The polynucleotide of claim 36, further comprising a silent mutation at S550.
- 1 38. The polynucleotide of claim 23, further comprising the amino acid mutation N413D.

- 1 39. The polynucleotide of claim 38, further comprising a silent mutation in at least one of  
2 S550 and S610.
- 1 40. A polynucleotide encoding for a galactose oxidase which has a nucleotide mutation  
2 in at least one of positions 9, 28, 208, 408, 584, 654, 830, 936, 1237, 1481, 1543,  
3 1603, 1609, 1650, and 1830.
- 1 41. The polynucleotide of claim 40, wherein the mutation at any of positions 9, 408, 654,  
2 936, 1650 and 1830 is a silent mutation.
- 1 42. The polynucleotide of claim 40 which has a mutation in at least one of nucleotide  
2 positions 28, 408, 654, and 1481, wherein a thymine is replaced by a cytosine.
- 1 43. The polynucleotide of claim 40, which has a mutation in at least one of nucleotide  
2 positions 1543, 1650 and 1830, wherein a thymine is replaced by an adenine.
- 1 44. The polynucleotide of claim 40, which has a mutation in at least one of nucleotide  
2 positions 206, 936, 1237, 1603, and 1609, wherein adenine is replaced by guanine.
- 1 45. A polynucleotide encoding for a galactose oxidase which has at least one nucleotide  
2 mutation in a region encompassed by nucleotides selected from the group consisting  
3 of:  
4 (a) 1 through 30;  
5 (b) 200 through 700;  
6 (c) 800 through 1000; and  
7 (d) 1200 through 1650.
- 1 46. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 encompassed by nucleotides 1-30, wherein a thymine is replaced by a cytosine.



- 1 47. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 encompassed by nucleotides 1450-1550, wherein a thymine is replaced by one of a  
3 cytosine and an adenine.
- 1 48. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 encompassed by nucleotides 1200-1250, wherein an adenine is replaced by a guanine.
- 1 49. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 encompassed by nucleotides 1600-1650, wherein an adenine is replaced by a guanine.
- 1 50. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 proximate to and encompassing nucleotide 208, wherein an adenine is replaced by a  
3 guanine.
- 1 51. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 proximate to and encompassing nucleotide 585, wherein a guanine is replaced by an  
3 adenine.
- 1 52. The polynucleotide of claim 45, which has a nucleotide mutation in a region  
2 proximate to and encompassing nucleotide 1543, wherein a thymine is replaced by an  
3 adenine.
- 1 53. A polynucleotide encoding for a galactose oxidase which has at least one of the  
2 nucleotide mutations A9C, T28C, A208G, T408C, G584A, T654C, A936G, A1237G,  
3 T1481C, T1543A, A1603G, A1609G, T1650A, and T1830A.
- 1 54. The polynucleotide of claim 53, which has the nucleotide mutation T1481C.
- 1 55. The polynucleotide of claim 54, further comprising the nucleotide mutation T1543A.
- 1 56. The polynucleotide of claim 54, further comprising the nucleotide mutation T408C.

- 1 57. The polynucleotide of claim 56, further comprising a nucleotide mutation selected  
2 from the group consisting of G584A, A1603G, and A208G.
- 1 58. The polynucleotide of claim 56, further comprising at least one of the nucleotide  
2 mutations A9C, A936G, and T654C.
- 1 59. The polynucleotide of claim 56, further comprising the nucleotide mutations T28C,  
2 A208G, G584A and A1603G.
- 1 60. The polynucleotide of claim 53 which has the nucleotide mutation A1237G.
- 1 61. The polynucleotide of claim 60, further comprising at least one of the nucleotide  
2 mutations selected from the group consisting of T1650A, T1830A, and T1481C.
- 1 62. The polynucleotide of claim 61, having the nucleotide mutations A1237G, T1650A,  
2 T1481C and T1830A.
- 1 63. A galactose oxidase which has a mutation in at least one amino acid selected from the  
2 group consisting of A3, S10, M70, P136, G195, T218, L312, N413, V494, C515,  
3 N535, N537, S550, and S610.
- 1 64. A galactose oxidase which has at least one of the amino acid mutations S10P, M70V,  
2 G195E, N413D, V494A, C515S, N535D, and N537D.
- 1 65. The galactose oxidase of claim 64, which has the amino acid mutation N537D.
- 1 66. The galactose oxidase of claim 64, which has the amino acid mutation V494A.
- 1 67. The galactose oxidase of claim 66, further comprising the amino acid mutation  
2 C515S.

- 1 68. The galactose oxidase claim 66, further comprising the amino acid mutation S10P.
- 1 69. The galactose oxidase of claim 66, further comprising a silent mutation at P136.
- 1 70. The galactose oxidase of claim 68, further comprising a silent mutation at P136.
- 1 71. The galactose oxidase of claim 66, further comprising the amino acid mutation  
2 G195E.
- 1 72. The galactose oxidase of claim 71, further comprising a silent mutation in at least one  
2 of A3 and P136.
- 1 73. The galactose oxidase of claim 66, further comprising the amino acid mutation  
2 N535D.
- 1 74. The galactose oxidase of claim 73, further comprising a silent mutation in at least one  
2 of P136, L312, and T218.
- 1 75. The galactose oxidase of claim 66, further comprising the amino acid mutation M70V.
- 1 76. The galactose oxidase of claim 75, further comprising a silent mutation at P136.
- 1 77. The galactose oxidase of claim 64, which has the amino acid mutations S10P, M70V,  
2 G195E, V494A and N535D.
- 1 78. The galactose oxidase of claim 77, further comprising a silent mutation at P136.
- 1 79. The galactose oxidase of claim 64, which has the amino acid mutation N413D.
- 1 80. The galactose oxidase of claim 80, further comprising a silent mutation at S550.

1 81. The galactose oxidase of claim 66, further comprising the amino acid mutation  
2 N413D.

1 82. The galactose oxidase of claim 81, further comprising a silent mutation in at least one  
2 of S550 and S610.

**AMENDED CLAIMS**

[received by the International Bureau on 15 October 2001 (15.10.01);  
original claims 1-82 replaced by new claims 1-130 (14 pages)]

- 1 1. A polynucleotide evolved according to a method, wherein the method comprises  
2 the steps of:  
3 (i) providing at least one parent galactose oxidase polynucleotide encoding  
4 a parent galactose oxidase polypeptide,  
5 (ii) altering the nucleotide sequence of the parent polynucleotide by random  
6 mutagenesis to produce a population of mutant polypeptides;  
7 (iii) transforming host cells to express the mutant polypeptides;  
8 (iv) screening for first-generation functional mutants produced by the host  
9 cells and having at least one modified property;  
10 (v) selecting at least one polynucleotide encoding a first-generation mutant  
11 as a parent polynucleotide; and  
12 (vi) repeating a round of altering, transforming and screening steps at least  
13 once to obtain at least one other generation of one or more mutants,  
14 wherein at least one of the altering, transforming and screening steps are changed  
15 in at least one repeated round.
- 1 2. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in at least one amino acid selected from the group consisting of  
3 A3, M70, P136, G195, T218, L312, N413, V494, C515, N535, N537, S550, S10,  
4 S610.
- 1 3. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in at least one amino acid selected from the group consisting of  
3 M70V, G195E, N413D, V494A, C515S, N535D, N537D, and S10P.
- 1 4. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has the amino acid mutation N537D.

- 1 5. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has the amino acid mutation V494A.
- 1 6. The polynucleotide of claim 5, wherein the galactose oxidase further comprises  
2 the amino acid mutation C515S.
- 1 7. The polynucleotide of claim 5, wherein the galactose oxidase further comprises  
2 the amino acid mutation S10P.
- 1 8. The polynucleotide of claim 5, wherein the galactose oxidase further comprises  
2 a silent mutation at P136.
- 1 9. The polynucleotide of claim 7, wherein the galactose oxidase further comprises  
2 a silent mutation at P136.
- 1 10. The polynucleotide of claim 5, wherein the galactose oxidase further comprises  
2 the amino acid mutation G195E.
- 1 11. The polynucleotide of claim 10, wherein the galactose oxidase further comprises  
2 a silent mutation in at least one of A3 and P136.
- 1 12. The polynucleotide of claim 5, wherein the galactose oxidase further comprises  
2 the amino acid mutation N535D.
- 1 13. The polynucleotide of claim 12, wherein the galactose oxidase further comprises  
2 a silent mutation in at least one of P136, L312, and T218.
- 1 14. The polynucleotide of claim 5, wherein the galactose oxidase further comprising  
2 the amino acid mutation M70V.

- 1 15. The polynucleotide of claim 14, wherein the galactose oxidase further comprises  
2 a silent mutation at P136.
- 1 16. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has the amino acid mutations V494A, S10P, M70V, G195E and N535D.
- 1 17. The polynucleotide of claim 16, wherein the galactose oxidase further comprises  
2 a silent mutation at P136.
- 1 18. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has the amino acid mutation N413D.
- 1 19. The polynucleotide of claim 18, wherein the galactose oxidase further comprises  
2 a silent mutation at S550.
- 1 20. The polynucleotide of claim 5, wherein the galactose oxidase further comprises  
2 the amino acid mutation N413D.
- 1 21. The polynucleotide of claim 20, wherein the galactose oxidase further comprises  
2 a silent mutation in at least one of S550 and S610.
- 1 22. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a nucleotide mutation in at least one position selected from the group  
3 consisting of 9, 28, 208, 408, 584, 654, 830, 936, 1237, 1481, 1543, 1603, 1609,  
4 1650, and 1830.
- 1 23. The polynucleotide of claim 22, wherein the mutation at any of positions 9, 408,  
2 654, 936, 1650 and 1830 is a silent mutation.

- 1 24. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a nucleotide mutation in at least one position of a region encompassed by  
3 nucleotides 1 through 30.
- 1 25. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a nucleotide mutation in at least one position of a region encompassed by  
3 nucleotides 200 through 700.
- 1 26. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a nucleotide mutation in at least one position of a region encompassed by  
3 nucleotides 800 through 1000.
- 1 27. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a nucleotide mutation in at least one position of a region encompassed by  
3 nucleotides 1200 through 1650.
- 1 28. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region encompassed by  
3 nucleotides 1-30, wherein a thymine is replaced by a cytosine.
- 1 29. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region encompassed by  
3 nucleotides 1450-1550, wherein thymine is replaced by one of cytosine and  
4 adenine.
- 1 30. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region encompassed by  
3 nucleotides 1200-1250, wherein adenine is replaced by guanine.



- 1 31. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region encompassed by  
3 nucleotides 1600-1650, wherein adenine is replaced by guanine.
- 1 32. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region proximate to and  
3 encompassing nucleotide 208, wherein adenine is replaced by guanine.
- 1 33. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region proximate to and  
3 encompassing nucleotide 584, wherein guanine is replaced by adenine.
- 1 34. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one position within a region proximate to and  
3 encompassing nucleotide 1543, wherein thymine is replaced by adenine.
- 1 35. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one of nucleotide positions 28, 408, 654, and 1481,  
3 wherein thymine is replaced by cytosine.
- 1 36. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one of nucleotide positions 1543, 1650 and 1830,  
3 wherein thymine is replaced by adenine.
- 1 37. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has a mutation in at least one of nucleotide positions 208, 936, 1237, 1603, and  
3 1609, wherein adenine is replaced by guanine.
- 1 38. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has at least one of the nucleotide mutations A1609G, T1481C, T1543A, T408C,

- 3 T28C, G584A, A9C, A936G, A1603G, T654C, A208G, A1237G, T1650A, and  
4 T1830A.
- 1 39. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has the nucleotide mutation A1609G.
- 1 40. A polynucleotide encoding for a galactose oxidase, wherein the polynucleotide  
2 has the nucleotide mutation T1481C.
- 1 41. The polynucleotide of claim 40, further comprising the nucleotide mutation  
2 T1543A.
- 1 42. The polynucleotide of claim 40, further comprising the nucleotide mutation  
2 T408C.
- 1 43. The polynucleotide of claim 42, further comprising the nucleotide mutation  
2 G584A.
- 1 44. The polynucleotide of claim 42, further comprising the nucleotide mutation  
2 A1603G.
- 1 45. The polynucleotide of claim 42, further comprising the nucleotide mutation  
2 A208G.
- 1 46. The polynucleotide of claim 42, further comprising at least one of the nucleotide  
2 mutations A9, A936G, and T654C.
- 1 47. The polynucleotide of claim 42, further comprising the nucleotide mutations  
2 T28C, A208G, G584A and A1603G.

- 1 48. A polynucleotide encoding a galactose oxidase, wherein the polynucleotide has  
2 the nucleotide mutation A1237G.
- 1 49. The polynucleotide of claim 48, further comprising at least one of the nucleotide  
2 mutations T1650A, and T1830A.
- 1 50. The polynucleotide of claim 48, further comprising the nucleotide mutation  
2 T1481C.
- 1 51. A polynucleotide encoding a galactose oxidase, wherein the polynucleotide has  
2 the nucleotide mutations A1237G, T1650A, T1481C and T1830A.
- 1 52. A polynucleotide encoding for a galactose oxidase which has a mutation in at  
2 least one amino acid selected from the group consisting of A3, S10, M70, P136,  
3 T218, L312, N413, C515, N535, N537, S550, and S610.
- 1 53. The polynucleotide encoding for a galactose oxidase of claim 52, wherein the  
2 galactose oxidase further comprises at least one amino acid mutation selected  
3 from the group consisting of G195 and V494.
- 1 54. The polynucleotide of claim 52, wherein the galactose oxidase has at least a  
2 mutation selected from the group consisting of S10P, M70V, N413D, C515S,  
3 N535D, and N537D.
- 1 55. The polynucleotide of claim 54, wherein the galactose oxidase further comprises  
2 at least one amino acid mutation selected from the group consisting of G195E and  
3 V494A.
- 1 56. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in amino acid N537.

- 1 57. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in amino acid N537.
- 1 58. The polynucleotide of claim 57, wherein the mutation is N537D.
- 1 59. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494 and C515.
- 1 60. The polynucleotide of claim 59, wherein the mutations are V494A and C515S.
- 1 61. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494 and P136.
- 1 62. The polynucleotide of claim 61, wherein the V494 mutation is V494A.
- 1 63. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494, P136, and S10.
- 1 64. The polynucleotide of claim 63, wherein the V494 mutation is V494A, and the  
2 S10 mutation is S10P.
- 1 65. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494, P136, G195, and A3.
- 1 66. The polynucleotide of claim 65, wherein the V494 mutation is V494A, and the  
2 G195 mutation is G195E.
- 1 67. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494, P136, L312, N535, and T218.

- 1 68. The polynucleotide of claim 67, wherein the V494 mutation is V494A, and the  
2 N535 mutation is N535D.
- 1 69. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494, P136, and M70.
- 1 70. The polynucleotide of claim 69, wherein the V494 mutation is V494A, and the  
2 M70 mutation is M70V.
- 1 71. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids V494, S10, P136, M70, G195, and N535.
- 1 72. The polynucleotide of claim 71, wherein the V494 mutation is V494A, the S10  
2 mutation is S10P, the M70 mutation is M70V, the G195 mutation is G195E, and  
3 the N535 mutation is N535D.
- 1 73. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in amino acid N413.
- 1 74. The polynucleotide of claim 73, wherein the mutation is N413D.
- 1 75. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in amino acids N413 and S550.
- 1 76. The polynucleotide of claim 75, wherein the N413 mutation is N413D.
- 1 77. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has a mutation in amino acids N413, S550, and V494.

- 1 78. The polynucleotide of claim 77, wherein the N413 mutation is N413D, and the  
2 V494 mutation is V494A.
- 1 79. A polynucleotide encoding for a galactose oxidase, wherein the galactose oxidase  
2 has mutations in amino acids N413, S550, V494, and S610.
- 1 80. The polynucleotide of claim 79, wherein the N413 mutation is N413D, and the  
2 V494 mutation is V494A.
- 1 81. A polynucleotide having a sequence selected from the group consisting of SEQ  
2 ID NOS:37-48.
- 1 82. A galactose oxidase which has a mutation in at least one amino acid selected from  
2 the group consisting of A3, S10, M70, P136, G195, T218, L312, N413, V494,  
3 C515, N535, N537, S550, and S610.
- 1 83. A galactose oxidase which has at least one of the amino acid mutations S10P,  
2 M70V, G195E, N413D, V494A, C515S, N535D, and N537D.
- 1 84. The galactose oxidase of claim 83, which has the amino acid mutation N537D.
- 1 85. The galactose oxidase of claim 83, which has the amino acid mutation V494A.
- 1 86. The galactose oxidase of claim 85, further comprising the amino acid mutation  
2 C515S.
- 1 87. The galactose oxidase claim 85, further comprising the amino acid mutation  
2 S10P.
- 1 88. The galactose oxidase of claim 85, further comprising a silent mutation at P136.

- 1 89. The galactose oxidase of claim 87, further comprising a silent mutation at P136.
- 1 90. The galactose oxidase of claim 85, further comprising the amino acid mutation  
2 G195E.
- 1 91. The galactose oxidase of claim 90, further comprising a silent mutation in at least  
2 one of A3 and P136.
- 1 92. The galactose oxidase of claim 85, further comprising the amino acid mutation  
2 N535D.
- 1 93. The galactose oxidase of claim 92, further comprising a silent mutation in at least  
2 one of P136, L312, and T218.
- 1 94. The galactose oxidase of claim 85, further comprising the amino acid mutation  
2 M70V.
- 1 95. The galactose oxidase of claim 94, further comprising a silent mutation at P136.
- 1 96. The galactose oxidase of claim 83, which has the amino acid mutations S10P,  
2 M70V, G195E, V494A and N535D.
- 1 97. The galactose oxidase of claim 96, further comprising a silent mutation at P136.
- 1 98. The galactose oxidase of claim 83, which has the amino acid mutation N413D.
- 1 99. The galactose oxidase of claim 98, further comprising a silent mutation at S550.

- 1 100. The galactose oxidase of claim 85, further comprising the amino acid mutation  
2 N413D.
- 1 101. The galactose oxidase of claim 100, further comprising a silent mutation in at  
2 least one of S550 and S610.
- 1 102. A galactose oxidase which has a mutation in at least one amino acid selected from  
2 the group consisting of A3, S10, M70, P136, T218, L312, N413, C515, N535,  
3 N537, S550, and S610.
- 1 103. The galactose oxidase of claim 102, further comprising at least one amino acid  
2 mutation selected from the group consisting of G195 and V494.
- 1 104. The galactose oxidase of claim 102, wherein the mutation is selected from the  
2 group consisting of S10P, M70V, N413D, C515S, N535D, and N537D.
- 1 105. The galactose oxidase of claim 104, further comprising at least one amino acid  
2 mutation selected from the group consisting of G195E and V494A.
- 1 106. A galactose oxidase which has a mutation in amino acid N537.
- 1 107. The galactose oxidase of claim 106, wherein the mutation is N537D.
- 1 108. A galactose oxidase which has mutations in amino acids V494 and C515.
- 1 109. The galactose oxidase of claim 108, wherein the mutations are V494A and  
2 C515S.
- 1 110. A galactose oxidase which has mutations in amino acids V494 and P136.



- 1 111. The galactose oxidase of claim 110, wherein the V494 mutation is V494A.
- 1 112. A galactose oxidase which has mutations in amino acids V494, P136, and S10.
- 1 113. The galactose oxidase of claim 112, wherein the V494 mutation is V494A, and  
2 the S10 mutation is S10P.
- 1 114. A galactose oxidase which has mutations in amino acids V494, P136, G195, and  
2 A3.
- 1 115. The galactose oxidase of claim 114, wherein the V494 mutation is V494A, and  
2 the G195 mutation is G195E.
- 1 116. A galactose oxidase which has mutations in amino acids V494, P136, L312,  
2 N535, and T218.
- 1 117. The galactose oxidase of claim 116, wherein the V494 mutation is V494A, and  
2 the N535 mutation is N535D.
- 1 118. A galactose oxidase which has mutations in amino acids V494, P136, and M70.
- 1 119. The galactose oxidase of claim 118, wherein the V494 mutation is V494A, and  
2 the M70 mutation is M70V.
- 1 120. A galactose oxidase which has mutations in amino acids V494, S10, P136, M70,  
2 G195, and N535.
- 1 121. The galactose oxidase of claim 120, wherein the V494 mutation is V494A, the  
2 S10 mutation is S10P, the M70 mutation is M70V, the G195 mutation is G195E,  
3 and the N535 mutation is N535D.

- 1 122. A galactose oxidase which has a mutation in amino acid N413.
- 1 123. The galactose oxidase of claim 122, wherein the mutation is N413D.
- 1 124. A galactose oxidase which has a mutation in amino acids N413 and S550.
- 1 125. The galactose oxidase of claim 124, wherein the N413 mutation is N413D.
- 1 126. A galactose oxidase which has a mutation in amino acids N413, S550, and V494.
- 1 127. The galactose oxidase of claim 126, wherein the N413 mutation is N413D, and  
2 the V494 mutation is V494A.
- 1 128. A galactose oxidase which has mutations in amino acids N413, S550, V494, and  
2 S610.
- 1 129. The galactose oxidase of claim 128, wherein the N413 mutation is N413D, and  
2 the V494 mutation is V494A.
- 1 130. A galactose oxidase having an amino acid sequence selected from the group  
2 consisting of SEQ ID NOS: 10-21.

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5




**FIG. 6**

PCR primers	Sequence	
MY		
001	5'-AAT TCG AAG CTT ATG GCC TCA GCA CCT ATC GGA AGC-3'	SEQ. ID NO. 1
002	5'-CTT CCT TCT AGA TTA CTG AGT AAC GCG AAT CGT-3'	SEQ. ID NO. 2
003	5'-GGA AGA GAA TTC AAT ACG CAA ACC GCC TCT-3'	SEQ. ID NO. 3
004	5'-GGT CAT AAG CTT TTC CTG TGT GAA ATT GTT AT-3'	SEQ. ID NO. 4
005	5'-ACC ATG ATT TCG ACG TCG GTA CCC TCA GCA-3'	SEQ. ID NO. 5
009	5'-CTT CCT AAG CTT TCA CTG AGT AAC GCG AAT-3'	SEQ. ID NO. 6
036	5'-GGA AGA GGT ACC AAT ACG CAA ACC GCC TCT-3'	SEQ. ID NO. 7

---

FIG. 7







FIG. 8








FIG. 10

FIG. 11

Plasmid	(vector)		GAO activities [units/ml-culture]					
			Host strain	DH5αMCR		BL21(DE3)		KY-14478
			Induction	-	IPTG	-	IPTG	IPTG
pR3	(pUC118)		0	0.01	0.01	0.03	0.31	
pGAO-003	(pET22b(+))		0	0	0	0	0	
pGAO-004	(pET22b(+))		0	0	0	0	0	
pGAO-005	(pET22b(+))		0	0	0	0	0	
pGAO-006	(pUC118)		1.22	1.72	0.08	1.35	0.87	
pGAO-007	(pET22b(+))		0.02	0	0.05	0	0	
pGAO-008	(pET22b(+))		0	0	0.03	0.01	0	
pGAO-009	(pET22b(+))		0	0	0.02	0.03	0	
pGAO-010	(pUC18)		0	0	0.67	1.43	0.40	
pGAO-011	(pUC18)		0.04	0.04	0.01	0.85	0.41	
pGAO-014	(pUC18)		0	0.01	- **	- **	0	
pGAO-015	(pUC18)		0	0	- **	- **	0	
pGAO-016	(pUC18)		0.19	0.15	0.03	0.04	0	
pGAO-017	(pUC18)		0	0	0.06	0.47	0.31	
pGAO-018	(pUC18)		- *	- *	- **	- **	- **	
pGAO-019	(pUC18)		- *	- *	- **	- **	- **	
pGAO-020	(pUC18)		0	0.01	0.97	2.21	0.22	
pGAO-021	(pUC18)		0.02	0	0.03	0.31	0.24	
pGAO-022	(pUC18)		0.03	0.08	0.12	0.93	0.14	
pGAO-023	(pUC18)		- *	- *	- **	- **	- *	
pGAO-024	(pUC18)		- *	- *	- **	- **	- *	

FIG. 12




FIG. 13

Plasmid	GAO activity [units/ml] (+ IPTG)	
	BL21(DE3)	KY-14478
pGAO-011 (pUC18)	0.28	0.74
pGAO-025 (pUC18)	0.91	0.88
pGAO-010 (pUC18)	3.32	1.20
pGAO-027 (pUC18)	3.64	1.57
pGAO-028 (pUC18)	2.97	1.77

FIG. 14








FIG. 15

Substrate (100 mM)	Relative activities of galactose oxidase [%]	
	<i>D. dendroides</i> (Sigma)	<i>E. coli</i> BL21(DE3)/pGAO-010
D-Galactose	100	100
D-Glucose	0	0
D-Sucrose	0	0
$\alpha$ -D-Lactose	20	17
$\beta$ -D-Lactose	42	32
D-Raffinose	114	110
D-Melibiose	75	75
Benzyl alcohol (25 % Methanol)	15	11
2-Hydroxybenzyl alcohol	(+)	(+)
2-Pyridylcarbinol	14	15
3-Pyridylcarbinol	50	46
4-Pyridylcarbinol	32	29
Cyclohexylmethanol (45 % Methanol) 1)	1.9	2.1
Tetrahydropyran-2-methanol 2)	0	0
Cyclopentamethanol (30 % Methanol) 3)	0.42	0.25
Tetrahydrofurfuryl alcohol 4)	n.d.	n.d.
Glycerol	4.1	3.4
Ethylene glycol	0.45	0.16
1-Propanol	0	0
1,2-Propanediol	(+)	(+)
Acetol	13	13
Allyl alcohol	4.6	3.6






FIG. 16

Date : 2000.04.10  
 Mutant ID : 9.16.8D2  
 Mutation : N537D(A1609G)  
 Sequence Size : 1917

FIG. 17A

```

      10      20      30      40      50      60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

      70      80      90      100      110      120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A Q S G N E C N K A I D G N K D T F W H

      130      140      150      160      170      180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

      190      200      210      220      230      240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

      250      260      270      280      290      300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

      310      320      330      340      350      360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

      370      380      390      400      410      420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCT TGG ACT AGC ATT
A R Y V R L V A I T E A N G Q P W T S I

      430      440      450      460      470      480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

      490      500      510      520      530      540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

      550      560      570      580      590      600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F G G S P G G

      610      620      630      640      650      660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

      670      680      690      700      710      720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

      730      740      750      760      770      780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

      790      800      810      820      830      840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```



FIG. 17B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E

910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M

970 980 990 1000 1010 1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W

1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S

1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D

1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G

1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G

1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T

1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R

1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q

1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GTC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R V Y H S I S L

1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T

1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC GAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S D G N L

1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I

1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T

1750 1760 1770 1780 1790 1800  
CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
H T V N T D Q R R I P L T L T N N G G N

1810 1820 1830 1840 1850 1860  
AGC TAT TCT TTC CAA GTT CCT AGC GAC TCT GGT GTT GCT TTG CCT GGC TAC TGG ATG TTG  
S Y S F Q V P S D S G V A L P G Y W M L

1870 1880 1890 1900 1910 1920  
TTC GTG ATG AAC TCG GCC GGT GTT CCT AGT GTG GCT TCG ACG ATT CGC GTT ACT CAG  
F V M N S A G V P S V A S T I R V T Q

FIG. 17C

Date : 2000.04.10  
 Mutant ID : 9.16.6C11  
 Mutation : V494A(T1481C), C515S(T1543A)  
 Sequence Size : 1917

FIG. 18A

```

      10          20          30          40          50          60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

      70          80          90          100         110         120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A Q S G N E C N K A I D G N K D T F W H

      130         140         150         160         170         180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

      190         200         210         220         230         240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

      250         260         270         280         290         300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

      310         320         330         340         350         360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

      370         380         390         400         410         420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCT TGG ACT AGC ATT
A R Y V R L V A I T E A N G Q P W T S I

      430         440         450         460         470         480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

      490         500         510         520         530         540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

      550         560         570         580         590         600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F G G S P G G

      610         620         630         640         650         660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

      670         680         690         700         710         720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

      730         740         750         760         770         780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

      790         800         810         820         830         840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```

FIG. 18B

850                    860                    870                    880                    890                    900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E

910                    920                    930                    940                    950                    960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M

970                    980                    990                    1000                    1010                    1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W

1030                    1040                    1050                    1060                    1070                    1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S

1090                    1100                    1110                    1120                    1130                    1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D

1150                    1160                    1170                    1180                    1190                    1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G

1210                    1220                    1230                    1240                    1250                    1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G

1270                    1280                    1290                    1300                    1310                    1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T

1330                    1340                    1350                    1360                    1370                    1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R

1390                    1400                    1410                    1420                    1430                    1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q

1450                    1460                    1470                    1480                    1490                    1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L

1510                    1520                    1530                    1540                    1550                    1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT AGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L S G D C T T

1570                    1580                    1590                    1600                    1610                    1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L

1630                    1640                    1650                    1660                    1670                    1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I

1690                    1700                    1710                    1720                    1730                    1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T

1750                    1760                    1770                    1780                    1790                    1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

1810 1820 1830 1840 1850 1860  
AGC TAT TCT TTC CAA GTT CCT AGC GAC TCT GGT GTT GCT TTG CCT GGC TAC TGG ATG TTG  
S Y S F Q V P S D S G V A L P G Y W M L

1870 1880 1890 1900 1910 1920  
TTC GTG ATG AAC TCG GCC GGT GTT CCT AGT GTG GCT TCG ACG ATT CGC GTT ACT CAG  
F V M N S A G V P S V A S T I R V T Q

FIG. 18C

Date : 2000.04.10  
 Mutant ID : 9.16.16D12  
 Mutation : P136(T408C), V494A(T1481C)  
 Sequence Size : 1917

FIG. 19A

```

      10      20      30      40      50      60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

      70      80      90      100     110     120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A Q S G N E C N K A I D G N K D T F W H

      130     140     150     160     170     180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

      190     200     210     220     230     240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

      250     260     270     280     290     300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

      310     320     330     340     350     360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

      370     380     390     400     410     420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCC TGG ACT AGC ATT
A R Y V R L V A I T E A N G Q P W T S I

      430     440     450     460     470     480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

      490     500     510     520     530     540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

      550     560     570     580     590     600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F G G S P G G

      610     620     630     640     650     660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

      670     680     690     700     710     720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

      730     740     750     760     770     780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

      790     800     810     820     830     840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```

FIG. 19B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E

910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M

970 980 990 1000 1010 1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W

1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S

1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D

1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G

1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G

1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T

1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R

1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q

1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L

1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T

1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L

1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I

1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T

1750 1760 1770 1780 1790 1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

	1810		1820		1830		1840		1850		1860								
AGC	TAT	TCT	TTC	CAA	GTT	CCT	AGC	GAC	TCT	GGT	GTT	GCT	TTG	CCT	GGC	TAC	TGG	ATG	TTG
S	Y	S	F	Q	V	P	S	D	S	G	V	A	L	P	G	Y	W	M	L
	1870		1880		1890		1900		1910		1920								
TTC	GTG	ATG	AAC	TCG	GCC	GGT	GTT	CCT	AGT	GTG	GCT	TCG	ACG	ATT	CGC	GTT	ACT	CAG	
F	V	M	N	S	A	G	V	P	S	V	A	S	T	I	R	V	T	Q	

FIG. 19C



FIG. 20A

Date : 2000.04.13  
 Mutant ID : 11.03.6D3  
 Mutation : S10P(T28C), P136(T408C), V494A(T1481C)  
 Sequence Size : 1917

```

      10          20          30          40          50          60
GCC TCA GCA CCT ATC GGA AGC GCC ATT CCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A  S  A  P  I  G  S  A  I  P  R  N  N  W  A  V  T  C  D  S

      70          80          90          100         110         120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A  Q  S  G  N  E  C  N  K  A  I  D  G  N  K  D  T  F  W  H

     130         140         150         160         170         180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T  F  Y  G  A  N  G  D  P  K  P  P  H  T  Y  T  I  D  M  K

     190         200         210         220         230         240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T  T  Q  N  V  N  G  L  S  M  L  P  R  Q  D  G  N  Q  N  G

     250         260         270         280         290         300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W  I  G  R  H  E  V  Y  L  S  S  D  G  T  N  W  G  S  P  V

     310         320         330         340         350         360
GCG TCA GGT GAT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A  S  G  S  W  F  A  D  S  T  T  K  Y  S  N  F  E  T  R  P

     370         380         390         400         410         420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCC TGG ACT AGC ATT
A  R  Y  V  R  L  V  A  I  T  E  A  N  G  Q  P  W  T  S  I

     430         440         450         460         470         480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A  E  I  N  V  F  Q  A  S  S  Y  T  A  P  Q  P  G  L  G  R

     490         500         510         520         530         540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W  G  P  T  I  D  L  P  I  V  P  A  A  A  A  I  E  P  T  S

     550         560         570         580         590         600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G  R  V  L  M  W  S  S  Y  R  N  D  A  F  G  G  S  P  G  G

     610         620         630         640         650         660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I  T  L  T  S  S  W  D  P  S  T  G  I  V  S  D  R  T  V  T

     670         680         690         700         710         720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V  T  K  H  D  M  F  C  P  G  I  S  M  D  G  N  G  Q  I  V

     730         740         750         760         770         780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V  T  G  G  N  D  A  K  K  T  S  L  Y  D  S  S  S  D  S  W

     790         800         810         820         830         840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I  P  G  P  D  M  Q  V  A  R  G  Y  Q  S  S  A  T  M  S  D
    
```

FIG. 20B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E  
  
 910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M  
  
 970 980 990 1000 1010 1020  
 TTG ACG GGT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W  
  
 1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S  
  
 1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D  
  
 1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G  
  
 1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G  
  
 1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T  
  
 1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R  
  
 1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q  
  
 1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L  
  
 1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T  
  
 1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L  
  
 1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I  
  
 1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T  
  
 1750 1760 1770 1780 1790 1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

		1810		1820		1830		1840		1850		1860							
AGC	TAT	TCT	TTC	CAA	GTT	CCT	AGC	GAC	TCT	GGT	GTT	GCT	TTG	CCT	GGC	TAC	TGG	ATG	TTG
S	Y	S	F	Q	V	P	S	D	S	G	V	A	L	P	G	Y	W	M	L
		1870		1880		1890		1900		1910		1920							
TTC	GTG	ATG	AAC	TCG	GCC	GGT	GTT	CCT	AGT	GTG	GCT	TCG	ACG	ATT	CGC	GTT	ACT	CAG	
F	V	M	N	S	A	G	V	P	S	V	A	S	T	I	R	V	T	Q	

FIG. 20C

Date : 2000.04.10  
 Mutant ID : 11.03.10C3  
 Mutation : A3(A9C),P136(T408C),G195E(G584A),V494A(T1481C)  
 Sequence Size : 1917

FIG. 21A

```

      10      20      30      40      50      60
GCC TCA GCC CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

      70      80      90      100      110      120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A Q S G N E C N K A I D G N K D T F W H

      130      140      150      160      170      180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

      190      200      210      220      230      240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

      250      260      270      280      290      300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

      310      320      330      340      350      360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

      370      380      390      400      410      420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCC TGG ACT AGC ATT
A R Y V R L V A I T E A N G Q P W T S I

      430      440      450      460      470      480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

      490      500      510      520      530      540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

      550      560      570      580      590      600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GAA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F E G S P G G

      610      620      630      640      650      660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

      670      680      690      700      710      720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

      730      740      750      760      770      780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

      790      800      810      820      830      840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```

FIG. 21B

850                    860                    870                    880                    890                    900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E  
  
 910                    920                    930                    940                    950                    960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M  
  
 970                    980                    990                    1000                    1010                    1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W  
  
 1030                    1040                    1050                    1060                    1070                    1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC ATG  
 K K G S V F Q A G P S T A M N W Y Y T S  
  
 1090                    1100                    1110                    1120                    1130                    1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D  
  
 1150                    1160                    1170                    1180                    1190                    1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G  
  
 1210                    1220                    1230                    1240                    1250                    1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G  
  
 1270                    1280                    1290                    1300                    1310                    1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACC  
 E P G T S P N T V F A S N G L Y F A R T  
  
 1330                    1340                    1350                    1360                    1370                    1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R  
  
 1390                    1400                    1410                    1420                    1430                    1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q  
  
 1450                    1460                    1470                    1480                    1490                    1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L  
  
 1510                    1520                    1530                    1540                    1550                    1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T  
  
 1570                    1580                    1590                    1600                    1610                    1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L  
  
 1630                    1640                    1650                    1660                    1670                    1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I  
  
 1690                    1700                    1710                    1720                    1730                    1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T  
  
 1750                    1760                    1770                    1780                    1790                    1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

1810 1820 1830 1840 1850 1860  
AGC TAT TCT TTC CAA GTT CCT AGC GAC TCT GGT GTT GCT TTG CCT GGC TAC TGG ATG TTG  
S Y S F Q V P S D S G V A L P G Y W M L

1870 1880 1890 1900 1910 1920  
TTC GTG ATG AAC TCG GCC GGT GTT CCT AGT GTG GCT TCG ACG ATT CGC GTT ACT CAG  
F V M N S A G V P S V A S T I R V T Q

FIG. 21C

FIG. 22A

Date : 2000.04.10  
 Mutant ID : 11.03.10D6  
 Mutation : P136(T408C), T218(T654C), L312(A936G), V494A(T1481C), N535D(A1603G)  
 Sequence Size : 1917

```

        10          20          30          40          50          60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
 A   S   A   P   I   G   S   A   I   S   R   N   N   W   A   V   T   C   D   S

        70          80          90          100         110         120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
 A   Q   S   G   N   E   C   N   K   A   I   D   G   N   K   D   T   F   W   H

       130          140          150          160          170          180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
 T   F   Y   G   A   N   G   D   P   K   P   P   H   T   Y   T   I   D   M   K

       190          200          210          220          230          240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
 T   T   Q   N   V   N   G   L   S   M   L   P   R   Q   D   G   N   Q   N   G

       250          260          270          280          290          300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
 W   I   G   R   H   E   V   Y   L   S   S   D   G   T   N   W   G   S   P   V

       310          320          330          340          350          360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
 A   S   G   S   W   F   A   D   S   T   T   K   Y   S   N   F   E   T   R   P

       370          380          390          400          410          420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCC TGG ACT AGC ATT
 A   R   Y   V   R   L   V   A   I   T   E   A   N   G   Q   P   W   T   S   I

       430          440          450          460          470          480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
 A   E   I   N   V   F   Q   A   S   S   Y   T   A   P   Q   P   G   L   G   R

       490          500          510          520          530          540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
 W   G   P   T   I   D   L   P   I   V   P   A   A   A   A   I   E   P   T   S

       550          560          570          580          590          600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
 G   R   V   L   M   W   S   S   Y   R   N   D   A   F   G   G   S   P   G   G

       610          620          630          640          650          660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACC GTG ACA
 I   T   L   T   S   S   W   D   P   S   T   G   I   V   S   D   R   T   V   T

       670          680          690          700          710          720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
 V   T   K   H   D   M   F   C   P   G   I   S   M   D   G   N   G   Q   I   V

       730          740          750          760          770          780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
 V   T   G   G   N   D   A   K   K   T   S   L   Y   D   S   S   S   D   S   W

       790          800          810          820          830          840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
 I   P   G   P   D   M   Q   V   A   R   G   Y   Q   S   S   A   T   M   S   D
    
```

FIG. 22B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E  
 910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTG CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M  
 970 980 990 1000 1010 1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W  
 1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S  
 1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D  
 1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G  
 1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G  
 1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T  
 1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R  
 1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q  
 1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L  
 1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T  
 1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC GAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y D S N G N L  
 1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I  
 1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T  
 1750 1760 1770 1780 1790 1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N



		1810		1820		1830		1840		1850		1860							
AGC	TAT	TCT	TTC	CAA	GTT	CCT	AGC	GAC	TCT	GGT	GTT	GCT	TTG	CCT	GGC	TAC	TGG	ATG	TTG
S	Y	S	F	Q	V	P	S	D	S	G	V	A	L	P	G	Y	W	M	L
		1870		1880		1890		1900		1910		1920							
TTC	GTG	ATG	AAC	TCG	GCC	GGT	GTT	CCT	AGT	GTG	GCT	TCG	ACG	ATT	CGC	GTT	ACT	CAG	
F	V	M	N	S	A	G	V	P	S	V	A	S	T	I	R	V	T	Q	

FIG. 22C

FIG. 23A

Date : 2000.04.10  
 Mutan ID : 11.03.13E12  
 Mutation : M70V(A208G), P136(T408C), V494A(T1481C)  
 Sequence Size : 1917

```

        10          20          30          40          50          60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A  S  A  P  I  G  S  A  I  S  R  N  N  W  A  V  T  C  D  S

        70          80          90          100         110         120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A  Q  S  G  N  E  C  N  K  A  I  D  G  N  K  D  T  F  W  H

       130         140         150         160         170         180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T  F  Y  G  A  N  G  D  P  K  P  P  H  T  Y  T  I  D  M  K

       190         200         210         220         230         240
ACA ACT CAG AAC GTC AAC GGC TTG TCT GTG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T  T  Q  N  V  N  G  L  S  V  L  P  R  Q  D  G  N  Q  N  G

       250         260         270         280         290         300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W  I  G  R  H  E  V  Y  L  S  S  D  G  T  N  W  G  S  P  V

       310         320         330         340         350         360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A  S  G  S  W  F  A  D  S  T  T  K  Y  S  N  F  E  T  R  P

       370         380         390         400         410         420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCC TGG ACT AGC ATT
A  R  Y  V  R  L  V  A  I  T  E  A  N  G  Q  P  W  T  S  I

       430         440         450         460         470         480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A  E  I  N  V  F  Q  A  S  S  Y  T  A  P  Q  P  G  L  G  R

       490         500         510         520         530         540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W  G  P  T  I  D  L  P  I  V  P  A  A  A  A  I  E  P  T  S

       550         560         570         580         590         600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G  R  V  L  M  W  S  S  Y  R  N  D  A  F  G  G  S  P  G  G

       610         620         630         640         650         660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I  T  L  T  S  S  W  D  P  S  T  G  I  V  S  D  R  T  V  T

       670         680         690         700         710         720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V  T  K  H  D  M  F  C  P  G  I  S  M  D  G  N  G  Q  I  V

       730         740         750         760         770         780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V  T  G  G  N  D  A  K  K  T  S  L  Y  D  S  S  S  D  S  W

       790         800         810         820         830         840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I  P  G  P  D  M  Q  V  A  R  G  Y  Q  S  S  A  T  M  S  D
    
```

FIG. 23B

850                    860                    870                    880                    890                    900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E  
  
 910                    920                    930                    940                    950                    960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M  
  
 970                    980                    990                    1000                    1010                    1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W  
  
 1030                    1040                    1050                    1060                    1070                    1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S  
  
 1090                    1100                    1110                    1120                    1130                    1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D  
  
 1150                    1160                    1170                    1180                    1190                    1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G  
  
 1210                    1220                    1230                    1240                    1250                    1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G  
  
 1270                    1280                    1290                    1300                    1310                    1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T  
  
 1330                    1340                    1350                    1360                    1370                    1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R  
  
 1390                    1400                    1410                    1420                    1430                    1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q  
  
 1450                    1460                    1470                    1480                    1490                    1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L  
  
 1510                    1520                    1530                    1540                    1550                    1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T  
  
 1570                    1580                    1590                    1600                    1610                    1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L  
  
 1630                    1640                    1650                    1660                    1670                    1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I  
  
 1690                    1700                    1710                    1720                    1730                    1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T  
  
 1750                    1760                    1770                    1780                    1790                    1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

1810 1820 1830 1840 1850 1860  
AGC TAT TCT TTC CAA GTT CCT AGC GAC TCT GGT GTT GCT TTG CCT GGC TAC TGG ATG TTG  
S Y S F Q V P S D S G V A L P G Y W M L

1870 1880 1890 1900 1910 1920  
TTC GTG ATG AAC TCG GCC GGT GTT CCT AGT GTG GCT TCG ACG ATT CGC GTT ACT CAG  
F V M N S A G V P S V A S T I R V T Q

FIG. 23C

FIG. 24A

Date : 2000.04.10  
 Filename : 1.06.20E7  
 Mutation : S10P(T28C),M70V(A208G),P136(T408C),G195E(G584A),V494A(T1481C)  
 N535D(A1603G)  
 Sequence Size : 1917

```

      10          20          30          40          50          60
GCC TCA GCA CCT ATC GGA AGC GCC ATT CCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A  S  A  P  I  G  S  A  I  P  R  N  N  W  A  V  T  C  D  S

      70          80          90          100         110         120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A  Q  S  G  N  E  C  N  K  A  I  D  G  N  K  D  T  F  W  H

     130         140         150         160         170         180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T  F  Y  G  A  N  G  D  P  K  P  P  H  T  Y  T  I  D  M  K

     190         200         210         220         230         240
ACA ACT CAG AAC GTC AAC GGC TTG TCT GTG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T  T  Q  N  V  N  G  L  S  V  L  P  R  Q  D  G  N  Q  N  G

     250         260         270         280         290         300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W  I  G  R  H  E  V  Y  L  S  S  D  G  T  N  W  G  S  P  V

     310         320         330         340         350         360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A  S  G  S  W  F  A  D  S  T  T  K  Y  S  N  F  E  T  R  P

     370         380         390         400         410         420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCC TGG ACT AGC ATT
A  R  Y  V  R  L  V  A  I  T  E  A  N  G  Q  P  W  T  S  I

     430         440         450         460         470         480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A  E  I  N  V  F  Q  A  S  S  Y  T  A  P  Q  P  G  L  G  R

     490         500         510         520         530         540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W  G  P  T  I  D  L  P  I  V  P  A  A  A  A  I  E  P  T  S

     550         560         570         580         590         600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GAA GGA TCC CCT GGT GGT
G  R  V  L  M  W  S  S  Y  R  N  D  A  F  E  G  S  P  G  G

     610         620         630         640         650         660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I  T  L  T  S  S  W  D  P  S  T  G  I  V  S  D  R  T  V  T

     670         680         690         700         710         720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V  T  K  H  D  M  F  C  P  G  I  S  M  D  G  N  G  Q  I  V

     730         740         750         760         770         780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V  T  G  G  N  D  A  K  K  T  S  L  Y  D  S  S  S  D  S  W

     790         800         810         820         830         840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I  P  G  P  D  M  Q  V  A  R  G  Y  Q  S  S  A  T  M  S  D
    
```

FIG. 24B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E

910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M

970 980 990 1000 1010 1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W

1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S

1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D

1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G

1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC AAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T N A H I I T L G

1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T

1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R

1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q

1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L

1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T

1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC GAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y D S N G N L

1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I

1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T

1750 1760 1770 1780 1790 1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

1810 1820 1830 1840 1850 1860  
AGC TAT TCT TTC CAA GTT CCT AGC GAC TCT GGT GTT GCT TTG CCT GGC TAC TGG ATG TTG  
S Y S F Q V P S D S G V A L P G Y W M L

1870 1880 1890 1900 1910 1920  
TTC GTG ATG AAC TCG GCC GGT GTT CCT AGT GTG GCT TCG ACG ATT CGC GTT ACT CAG  
F V M N S A G V P S V A S T I R V T Q

**FIG. 24C**

Date : 2000.04.11  
 Mutant ID : 1.D4  
 Mutation : N413D(A1237G)  
 Sequence Size : 1917

FIG. 25A

```

      10      20      30      40      50      60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

      70      80      90      100      110      120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAG
A Q S G N E C N K A I D G N K D T F W H

      130      140      150      160      170      180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

      190      200      210      220      230      240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

      250      260      270      280      290      300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

      310      320      330      340      350      360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

      370      380      390      400      410      420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCT TGG ACT AGC ATT
A R Y V R L V A I T E A N G Q P W T S I

      430      440      450      460      470      480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

      490      500      510      520      530      540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

      550      560      570      580      590      600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F G G S P G G

      610      620      630      640      650      660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

      670      680      690      700      710      720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

      730      740      750      760      770      780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

      790      800      810      820      830      840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```



FIG. 25B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E

910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M

970 980 990 1000 1010 1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W

1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S

1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D

1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G

1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC GAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T D A H I I T L G

1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T

1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R

1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q

1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GTC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R V Y H S I S L

1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T

1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L

1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCT ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I

1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T

1750 1760 1770 1780 1790 1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

1810 1820 1830 1840 1850 1860  
AGC TAT TCT TTC CAA GTT CCT AGC GAC TCT GGT GTT GCT TTG CCT GGC TAC TGG ATG TTG  
S Y S F Q V P S D S G V A L P G Y W M L

1870 1880 1890 1900 1910 1920  
TTC GTG ATG AAC TCG GCC GGT GTT CCT AGT GTG GCT TCG ACG ATT CGC GTT ACT CAG  
F V M N S A G V P S V A S T I R V T Q

FIG. 25C

Date : 2000.04.11  
 Mutant ID : 2.G4  
 Mutation : N413D(A1237G),S550(T1650A)  
 Sequence Size : 1917

FIG. 26A

```

      10      20      30      40      50      60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

      70      80      90      100      110      120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A Q S G N E C N K A I D G N K D T F W H

      130      140      150      160      170      180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

      190      200      210      220      230      240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

      250      260      270      280      290      300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

      310      320      330      340      350      360
GGC TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

      370      380      390      400      410      420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCT TGG ACT AGC ATT
A R Y V R L V A I T E A N G Q P W T S I

      430      440      450      460      470      480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

      490      500      510      520      530      540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

      550      560      570      580      590      600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F G G S P G G

      610      620      630      640      650      660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

      670      680      690      700      710      720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

      730      740      750      760      770      780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

      790      800      810      820      830      840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```

FIG. 26B

850 860 870 880 890 900  
GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
G R V F T I G G S W S G G V F E K N G E

910 920 930 940 950 960  
GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
V Y S P S S K T W T S L P N A K V N P M

970 980 990 1000 1010 1020  
TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
L T A D K Q G L Y R S D N H A W L F G W

1030 1040 1050 1060 1070 1080  
AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
K K G S V F Q A G P S T A M N W Y Y T S

1090 1100 1110 1120 1130 1140  
GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
G S G D V K S A G K R Q S N R G V A P D

1150 1160 1170 1180 1190 1200  
GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
A M C G N A V M Y D A V K G K I L T F G

1210 1220 1230 1240 1250 1260  
GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC GAC GCC CAC ATC ATC ACC CTC GGT  
G S P D Y Q D S D A T T D A H I I T L G

1270 1280 1290 1300 1310 1320  
GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
E P G T S P N T V F A S N G L Y F A R T

1330 1340 1350 1360 1370 1380  
TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
F H T S V V L P D G S T F I T G G Q R R

1390 1400 1410 1420 1430 1440  
GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
G I P F E D S T P V F T P E I Y V P E Q

1450 1460 1470 1480 1490 1500  
GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GTC TAC CAT AGC ATT TCC CTT  
D T F Y K Q N P N S I V R V Y H S I S L

1510 1520 1530 1540 1550 1560  
TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
L L P D G R V F N G G G G L C G D C T T

1570 1580 1590 1600 1610 1620  
AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
N H F D A Q I F T P N Y L Y N S N G N L

1630 1640 1650 1660 1670 1680  
GCG ACA CGT CCC AAG ATT ACC AGA ACC TCA ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
A T R P K I T R T S T Q S V K V G G R I

1690 1700 1710 1720 1730 1740  
ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
T I S T D S S I S K A S L I R Y G T A T

1750 1760 1770 1780 1790 1800  
CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
H T V N T D Q R R I P L T L T N N G G N

		1810		1820		1830		1840		1850		1860							
AGC	TAT	TCT	TTC	CAA	GTT	CCT	AGC	GAC	TCT	GGT	GTT	GCT	TTG	CCT	GGC	TAC	TGG	ATG	TTG
S	Y	S	F	Q	V	P	S	D	S	G	V	A	L	P	G	Y	W	M	L
		1870		1880		1890		1900		1910		1920							
TTC	GTG	ATG	AAC	TCG	GCC	GGT	GTT	CCT	AGT	GTG	GCT	TCG	ACG	ATT	CGC	GTT	ACT	CAG	
F	V	M	N	S	A	G	V	P	S	V	A	S	T	I	R	V	T	Q	

FIG. 26C

FIG. 27A

Date : 2000.04.11  
 Mutant ID : 3.H7  
 Mutation : N413D(A1237G), S550(T1650A), V494A(T1481C)  
 Sequence Size : 1917

```

10      20      30      40      50      60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A  S  A  P  I  G  S  A  I  S  R  N  N  W  A  V  T  C  D  S

70      80      90      100     110     120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A  Q  S  G  N  E  C  N  K  A  I  D  G  N  K  D  T  F  W  H

130     140     150     160     170     180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T  F  Y  G  A  N  G  D  P  K  P  P  H  T  Y  T  I  D  M  K

190     200     210     220     230     240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T  T  Q  N  V  N  G  L  S  M  L  P  R  Q  D  G  N  Q  N  G

250     260     270     280     290     300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W  I  G  R  H  E  V  Y  L  S  S  D  G  T  N  W  G  S  P  V

310     320     330     340     350     360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A  S  G  S  W  F  A  D  S  T  T  K  Y  S  N  F  E  T  R  P

370     380     390     400     410     420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCT TGG ACT AGC ATT
A  R  Y  V  R  L  V  A  I  T  E  A  N  G  Q  P  W  T  S  I

430     440     450     460     470     480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A  E  I  N  V  F  Q  A  S  S  Y  T  A  P  Q  P  G  L  G  R

490     500     510     520     530     540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W  G  P  T  I  D  L  P  I  V  P  A  A  A  A  I  E  P  T  S

550     560     570     580     590     600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G  R  V  L  M  W  S  S  Y  R  N  D  A  F  G  G  S  P  G  G

610     620     630     640     650     660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I  T  L  T  S  S  W  D  P  S  T  G  I  V  S  D  R  T  V  T

670     680     690     700     710     720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V  T  K  H  D  M  F  C  P  G  I  S  M  D  G  N  G  Q  I  V

730     740     750     760     770     780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT AGC GAT AGC TGG
V  T  G  G  N  D  A  K  K  T  S  L  Y  D  S  S  S  D  S  W

790     800     810     820     830     840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I  P  G  P  D  M  Q  V  A  R  G  Y  Q  S  S  A  T  M  S  D
    
```

FIG. 27B

850 860 870 880 890 900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E

910 920 930 940 950 960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M

970 980 990 1000 1010 1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W

1030 1040 1050 1060 1070 1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S

1090 1100 1110 1120 1130 1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D

1150 1160 1170 1180 1190 1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G

1210 1220 1230 1240 1250 1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC GAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T D A H I I T L G

1270 1280 1290 1300 1310 1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T

1330 1340 1350 1360 1370 1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R

1390 1400 1410 1420 1430 1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q

1450 1460 1470 1480 1490 1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L

1510 1520 1530 1540 1550 1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T

1570 1580 1590 1600 1610 1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L

1630 1640 1650 1660 1670 1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCA ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I

1690 1700 1710 1720 1730 1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T

1750 1760 1770 1780 1790 1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

	1810	1820	1830	1840	1850	1860													
AGC	TAT	TCT	TTC	CAA	GTT	CCT	AGC	GAC	TCT	GGT	GTT	GCT	TTG	CCT	GGC	TAC	TGG	ATG	TTG
S	Y	S	F	Q	V	P	S	D	S	G	V	A	L	P	G	Y	W	M	L
	1870	1880	1890	1900	1910	1920													
TTC	GTG	ATG	AAC	TCG	GCC	GGT	GTT	CCT	AGT	GTG	GCT	TCG	ACG	ATT	CGC	GTT	ACT	CAG	
F	V	M	N	S	A	G	V	P	S	V	A	S	T	I	R	V	T	Q	

FIG. 27C



FIG. 28A

Date : 2000.04.11  
 Mutant ID : 4.F12  
 Mutation : N413D(A1237G),S550(T1650A),V494A(T1481C),S610(T1830A)  
 Sequence Size : 1917

```

10      20      30      40      50      60
GCC TCA GCA CCT ATC GGA AGC GCC ATT TCT CGC AAC AAC TGG GCC GTC ACT TGC GAC AGT
A S A P I G S A I S R N N W A V T C D S

70      80      90      100      110      120
GCA CAG TCG GGA AAT GAA TGC AAC AAG GCC ATT GAT GGC AAC AAG GAT ACC TTT TGG CAC
A Q S G N E C N K A I D G N K D T F W H

130     140     150     160     170     180
ACA TTC TAT GGC GCC AAC GGG GAT CCA AAG CCC CCT CAC ACA TAC ACG ATT GAC ATG AAG
T F Y G A N G D P K P P H T Y T I D M K

190     200     210     220     230     240
ACA ACT CAG AAC GTC AAC GGC TTG TCT ATG CTG CCT CGA CAG GAT GGT AAC CAA AAC GGC
T T Q N V N G L S M L P R Q D G N Q N G

250     260     270     280     290     300
TGG ATC GGT CGC CAT GAG GTT TAT CTA AGC TCA GAT GGC ACA AAC TGG GGC AGC CCT GTT
W I G R H E V Y L S S D G T N W G S P V

310     320     330     340     350     360
GCG TCA GGT AGT TGG TTC GCC GAC TCT ACT ACA AAA TAC TCC AAC TTT GAA ACT CGC CCT
A S G S W F A D S T T K Y S N F E T R P

370     380     390     400     410     420
GCT CGC TAT GTT CGT CTT GTC GCT ATC ACT GAA GCG AAT GGC CAG CCT TGG ACT AGC ATT
P R Y V R L V A I T E A N G Q P W T S I

430     440     450     460     470     480
GCA GAG ATC AAC GTC TTC CAA GCT AGT TCT TAC ACA GCC CCC CAG CCT GGT CTT GGA CGC
A E I N V F Q A S S Y T A P Q P G L G R

490     500     510     520     530     540
TGG GGT CCG ACT ATT GAC TTA CCG ATT GTT CCT GCG GCT GCA GCA ATT GAA CCG ACA TCG
W G P T I D L P I V P A A A A I E P T S

550     560     570     580     590     600
GGA CGA GTC CTT ATG TGG TCT TCA TAT CGC AAT GAT GCA TTT GGA GGA TCC CCT GGT GGT
G R V L M W S S Y R N D A F G G S P G G

610     620     630     640     650     660
ATC ACT TTG ACG TCT TCC TGG GAT CCA TCC ACT GGT ATT GTT TCC GAC CGC ACT GTG ACA
I T L T S S W D P S T G I V S D R T V T

670     680     690     700     710     720
GTC ACC AAG CAT GAT ATG TTC TGC CCT GGT ATC TCC ATG GAT GGT AAC GGT CAG ATC GTA
V T K H D M F C P G I S M D G N G Q I V

730     740     750     760     770     780
GTC ACA GGT GGC AAC GAT GCC AAG AAG ACC AGT TTG TAT GAT TCA TCT ACG GAT AGC TGG
V T G G N D A K K T S L Y D S S S D S W

790     800     810     820     830     840
ATC CCG GGA CCT GAC ATG CAA GTG GCT CGT GGG TAT CAG TCA TCA GCT ACC ATG TCA GAC
I P G P D M Q V A R G Y Q S S A T M S D
    
```

FIG. 28B

850                    860                    870                    880                    890                    900  
 GGT CGT GTT TTT ACC ATT GGA GGC TCC TGG AGC GGT GGC GTA TTT GAG AAG AAT GGC GAA  
 G R V F T I G G S W S G G V F E K N G E  
  
 910                    920                    930                    940                    950                    960  
 GTC TAT AGC CCA TCT TCA AAG ACA TGG ACG TCC CTA CCC AAT GCC AAG GTC AAC CCA ATG  
 V Y S P S S K T W T S L P N A K V N P M  
  
 970                    980                    990                    1000                    1010                    1020  
 TTG ACG GCT GAC AAG CAA GGA TTG TAC CGT TCA GAC AAC CAC GCG TGG CTC TTT GGA TGG  
 L T A D K Q G L Y R S D N H A W L F G W  
  
 1030                    1040                    1050                    1060                    1070                    1080  
 AAG AAG GGT TCG GTG TTC CAA GCG GGA CCT AGC ACA GCC ATG AAC TGG TAC TAT ACC AGT  
 K K G S V F Q A G P S T A M N W Y Y T S  
  
 1090                    1100                    1110                    1120                    1130                    1140  
 GGA AGT GGT GAT GTG AAG TCA GCC GGA AAA CGC CAG TCT AAC CGT GGT GTA GCC CCT GAT  
 G S G D V K S A G K R Q S N R G V A P D  
  
 1150                    1160                    1170                    1180                    1190                    1200  
 GCC ATG TGC GGA AAC GCT GTC ATG TAC GAC GCC GTT AAA GGA AAG ATC CTG ACC TTT GGC  
 A M C G N A V M Y D A V K G K I L T F G  
  
 1210                    1220                    1230                    1240                    1250                    1260  
 GGC TCC CCA GAT TAT CAA GAC TCT GAC GCC ACA ACC GAC GCC CAC ATC ATC ACC CTC GGT  
 G S P D Y Q D S D A T T D A H I I T L G  
  
 1270                    1280                    1290                    1300                    1310                    1320  
 GAA CCC GGA ACA TCT CCC AAC ACT GTC TTT GCT AGC AAT GGG TTG TAC TTT GCC CGA ACG  
 E P G T S P N T V F A S N G L Y F A R T  
  
 1330                    1340                    1350                    1360                    1370                    1380  
 TTT CAC ACC TCT GTT GTT CTT CCA GAC GGA AGC ACG TTT ATT ACA GGA GGC CAA CGA CGT  
 F H T S V V L P D G S T F I T G G Q R R  
  
 1390                    1400                    1410                    1420                    1430                    1440  
 GGA ATT CCG TTC GAG GAT TCA ACC CCG GTA TTT ACA CCT GAG ATC TAC GTC CCT GAA CAA  
 G I P F E D S T P V F T P E I Y V P E Q  
  
 1450                    1460                    1470                    1480                    1490                    1500  
 GAC ACT TTC TAC AAG CAG AAC CCC AAC TCC ATT GTT CGC GCC TAC CAT AGC ATT TCC CTT  
 D T F Y K Q N P N S I V R A Y H S I S L  
  
 1510                    1520                    1530                    1540                    1550                    1560  
 TTG TTA CCT GAT GGC AGG GTA TTT AAC GGT GGT GGT GGT GGT CTT TGT GGC GAT TGT ACC ACG  
 L L P D G R V F N G G G G L C G D C T T  
  
 1570                    1580                    1590                    1600                    1610                    1620  
 AAT CAT TTC GAC GCG CAA ATC TTT ACG CCA AAC TAT CTT TAC AAT AGC AAC GGC AAT CTC  
 N H F D A Q I F T P N Y L Y N S N G N L  
  
 1630                    1640                    1650                    1660                    1670                    1680  
 GCG ACA CGT CCC AAG ATT ACC AGA ACC TCA ACA CAG AGC GTC AAG GTC GGT GGC AGA ATT  
 A T R P K I T R T S T Q S V K V G G R I  
  
 1690                    1700                    1710                    1720                    1730                    1740  
 ACA ATC TCG ACG GAT TCT TCG ATT AGC AAG GCG TCG TTG ATT CGC TAT GGT ACA GCG ACA  
 T I S T D S S I S K A S L I R Y G T A T  
  
 1750                    1760                    1770                    1780                    1790                    1800  
 CAC ACG GTT AAT ACT GAC CAG CGC CGC ATT CCC CTG ACT CTG ACA AAC AAT GGA GGA AAT  
 H T V N T D Q R R I P L T L T N N G G N

		1810		1820		1830		1840		1850		1860							
AGC	TAT	TCT	TTC	CAA	GTT	CCT	AGC	GAC	TCA	GGT	GTT	GCT	TTG	CCT	GGC	TAC	TGG	ATG	TTG
S	Y	S	F	Q	V	P	S	D	S	G	V	A	L	P	G	Y	W	M	L
		1870		1880		1890		1900		1910		1920							
TTC	GTG	ATG	AAC	TCG	GCC	GGT	GTT	CCT	AGT	GTG	GCT	TCG	ACG	ATT	CGC	GTT	ACT	CAG	
F	V	M	N	S	A	G	V	P	S	V	A	S	T	I	R	V	T	Q	

**FIG. 28C**

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/32345

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 9/00, 15/00; C07H 21/04

US CL : 435/183, 440; 536/23.2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/183, 440; 536/23.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROGERS, M.S. et al. Characterization fo the active site of galactose oxidase and its active site mutational variants Y495/F/H/K and W290H by circular dichroism spectroscopy. Inorganica Chimica Acta.	45
Y,P	US 6,090,604 A (GOLIGHTLY et al) 18 July 2000, entire document, especially col. 8, line 64-col. 10, line 16.	1-19
Y	US 5,811,238 A (STEMMER et al ) 22 September 1998, entire document.	1-19

 Further documents are listed in the continuation of Box C.  See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"B" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 16 JULY 2001	Date of mailing of the international search report 14.08.2001
---	--

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRONWEN M. LOEB

Telephone No. (703) 305-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/32345

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

MEDLINE, EMBASE, BIOSIS, CAPLUS, various sequence databases

search terms: galactose oxidase, evolving, directed evolution, SEQ ID Nos. 10 and 11