

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2018214554 C1**

(54) Title
Treatment of diuretic resistance

(51) International Patent Classification(s)
C07K 16/00 (2006.01) **A61P 7/10** (2006.01)
A61K 39/395 (2006.01) **A61P 13/12** (2006.01)

(21) Application No: **2018214554** (22) Date of Filing: **2018.02.01**

(87) WIPO No: **WO18/144773**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/453,257	2017.02.01	US

(43) Publication Date: **2018.08.09**

(44) Accepted Journal Date: **2022.06.02**

(44) Amended Journal Date: **2022.12.15**

(71) Applicant(s)
Novo Nordisk A/S; Yale University

(72) Inventor(s)
Lo, Chih-Hung; Testani, Jeffrey; Rao, Veena; Kakkar, Rahul; Devalaraja, Madhav N.

(74) Agent / Attorney
Phillips Ormonde Fitzpatrick, PO Box 323, COLLINS STREET WEST, VIC, 8007, AU

(56) Related Art
US 20070218063 A1
US 20150246092 A1
HANBERG J. S. et al., "Hypochloremia and Diuretic Resistance in Heart Failure: Mechanistic Insights", Circulation: Heart Failure, (2016-08), vol. 9, no. 8, pages 1 - 12
WO 2007106811 A2



(51) International Patent Classification:

A61K 39/395 (2006.01) A61P 7/10 (2006.01)
C07K 16/00 (2006.01) A61P 13/12 (2006.01)

(21) International Application Number:

PCT/US2018/016508

(22) International Filing Date:

01 February 2018 (01.02.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/453,257 01 February 2017 (01.02.2017) US

(71) Applicants: YALE UNIVERSITY [US/US]; Two Whitney Avenue, New Haven, CT 06511 (US). CORVIDIA THERAPEUTICS, INC. [US/US]; 35 Gatehouse Drive, Waltham, MA 02451 (US).

(72) Inventor; and

(71) Applicant: LO, Chih-Hung [US/US]; c/o Corvidia Therapeutics, Inc., 35 Gatehouse Drive, Waltham, MA 02451 (US).

(72) Inventors: TESTANI, Jeffrey; 532 Tanner Marsh Road, Guilford, CT 06437 (US). RAO, Veena; 844 Tall Timber Road, Orange, CT 06477 (US). KAKKAR, Rahul; c/o Corvidia Therapeutics, Inc., 35 Gatehouse Drive, Waltham, MA 02451 (US). DEVALARAJA, Madhav N.; c/o Corvidia Therapeutics, Inc., 35 Gatehouse Drive, Waltham, MA 02451 (US).

(74) Agent: SUN, Danqiong et al.; Fenwick & West LLP, Silicon Valley Center, 801 California Street, Mountain View, CA 94041 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(54) Title: TREATMENT OF DIURETIC RESISTANCE




FIG. 1B

(57) Abstract: The present disclosure provides methods of treating diuretic resistance by administering an IL-6 antagonist to patients who require diuresis. In typical embodiments, the patient has heart failure. Optionally, the patient has elevated urine levels of IL-6, plasma levels of IL-6, or both urine and plasma levels of IL-6.





(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

TREATMENT OF DIURETIC RESISTANCE

1. CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/453,257, filed February 1, 2017, which is hereby incorporated in its entirety by reference.

2. STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] This invention was made with government support under grants 5R01HL128973 and 4K23HL114868 awarded by the National Institutes of Health. The government has certain rights in the invention.

3. STATEMENT REGARDING JOINT RESEARCH AGREEMENT

[0003] This invention was made under a Joint Research Agreement by and among Yale University, MedImmune Ltd., AstraZeneca Pharmaceuticals LP, and Corvidia Therapeutics, Inc.

4. SEQUENCE LISTING

[0004] The instant application contains a Sequence Listing filed during international phase and is hereby incorporated herein by reference in its entirety.

5. BACKGROUND

[0005] Renal injury is often observed in heart failure, and heart failure is often observed in renal disease. The term cardiorenal syndrome (“CRS”) encompasses a variety of clinical conditions in which dysfunction of the heart, kidney, or both, leads to accelerated failure of both organs. Evidence of this adverse organ crosstalk portends a high degree of morbidity and mortality. Despite its dire clinical implications, the mechanistic underpinnings of CRS are only now being elucidated.

[0006] There is a need for new methods of treating renal injury and renal impairment in heart failure patients. There is also a need for new methods of detecting cardiorenal syndrome, for determining which cardiorenal patients will be responsive to treatment, and for monitoring efficacy of therapy.

[0007] Diuretics are a mainstay in the treatment of heart failure. However, certain patients are, or become, resistant to diuretics. There is a need for new methods of treating patients who require diuretics but are resistant to diuretics.

6. SUMMARY

[0008] In one aspect, the present invention provides a method of treating left ventricular heart failure, the method comprising: administering to a patient with left ventricular heart failure a therapeutically effective amount of an anti IL 6 antibody.

[0008a] In another aspect, the present invention provides a use of a therapeutically effective amount of an anti IL-6 antibody in the manufacture of a medicament for treating left ventricular heart failure in a patient.

[0008b] As further described below in Example 1, consecutive heart failure (“HF”) patients receiving high dose diuretic therapy at an outpatient treatment center were enrolled in a prospective observational study. Plasma levels of IL-6 were measured to query systemic associations of this pro-inflammatory cytokine with various disease parameters, and urine levels of IL-6 were measured to query IL-6 associations with local inflammation and neurohormonal activation at the level of renal tissues.

[0009] Urine IL-6 and plasma IL-6 levels were found to be only modestly correlated with one another.

[0010] Increases in urine levels of IL-6 were significantly correlated in these heart failure patients with measures of renal impairment, such as diuretic resistance, lower estimated glomerular filtration rate (“eGFR”), and increased tissue-level renin-angiotensin-aldosterone system (“RAAS”) activation.

[0011] Although an inverse association between diuretic efficiency and plasma IL-6 was also observed, upon adjustment for eGFR, only urine IL-6 remained significantly associated with risk of low diuretic efficiency in these patients. Furthermore, when urine IL-6 and plasma IL-6 were both entered into a logistic regression model, only urine IL-6 remained associated with risk of low diuretic efficiency while plasma IL-6 showed no such association.

[0012] These data demonstrate that urine IL-6 level is a useful biomarker for renal inflammation, and can be used to gauge renal dysfunction in the setting of heart failure (cardiorenal syndrome). The data further suggest that serial measurements of urine IL-6 can be used to measure the renal benefits of treatments administered to patients with heart failure, notably heart failure patients with cardiorenal syndrome.

[0013] The urine IL-6 data, and to some extent the plasma IL-6 data, also predict that treatment with an IL-6 antagonist should be effective to reduce renal inflammation in heart failure patients, that is, to treat renal symptoms of cardiorenal syndrome.

[0014] However, because infection is often a precipitating cause of acute decompensation in heart failure patients, it is important to limit anti-cytokine and other immunosuppressive therapies to those heart failure patients who are likely to respond with improved renal and/or cardiac function. The cost of chronic IL-6 antagonist therapy also militates for limiting treatment to those heart failure patients who are likely to respond with improved renal and/or cardiac function.

[0015] Analysis was expanded to 129 patients, and further assessed each patient's genotype at the rs855791 single nucleotide polymorphism ("SNP") in transmembrane protease serine 6 ("TMPRSS6").

[0016] Urine levels of IL-6 were inversely correlated with diuretic efficiency only in the patients having at least one copy of the major allele of the TMPRSS6 rs855791 SNP (AG and GG); urine levels of IL-6 were not significantly correlated with diuretic efficiency in patients homozygous for the minor allele (AA). Plasma levels of IL-6 correlated inversely with diuretic efficiency only in the patients having at least one copy of the major allele of the TMPRSS6 rs855791 SNP; plasma levels of IL-6 were not significantly correlated with diuretic efficiency in patients homozygous for the minor allele.

[0017] These data suggested that diuretic resistance (low diuretic efficiency) in heart failure patients could be treated with an IL-6 antagonist, but only in those having at least one copy of the TMPRSS6 rs855791 major allele.

[0018] In mouse M1 CCD cells, which are genotypically analogous to human cells homozygous for the *TMPRSS6* rs855791 major allele, the addition of IL-6 correlated with the expression of ion transporters, NKCC2, ENaC-beta, and NCC. Increased expression of these ion transporters provides a putative mechanism for IL-6 mediated diuretic resistance.

[0019] Because the IL-6 mediated increase in expression of ion transporters is not known to be linked to hepcidin expression, these data suggested that IL-6 antagonism could also be effective in treating diuretic resistance in patients homozygous for the *TMPRSS6* rs855791 minor allele.

[0020] Secondary analysis of data from two additional large heart failure clinical trials confirmed the association of diuretic resistance with IL-6 level (Example 5), independently of

TMPRSS6 rs855791 genotype (Example 6), providing evidence that IL-6 antagonism should also be effective in treating diuretic resistance in patients homozygous for the *TMPRSS6* rs855791 minor allele.

[0021] Accordingly, in a first aspect, methods are provided for treating a patient who requires diuresis but is resistant to diuretics. The methods comprise administering, in combination with a diuretic, a therapeutically effective amount of an IL-6 antagonist to the patient.

[0022] In some embodiments, the patient has elevated pre-treatment plasma IL-6 levels. In certain embodiments, the patient has a pre-treatment plasma IL-6 level of greater than 2 pg/mL. In certain embodiments, the patient has a pre-treatment IL-6 level of greater than 3 pg/mL. In certain embodiments, the patient has a pre-treatment IL-6 level of greater than 5 pg/mL. In certain embodiments, the patient has a pre-treatment IL-6 level of greater than 10 pg/mL.

[0023] In some embodiments, the patient has a diuretic efficiency of less than 500. In some embodiments, the patient has a diuretic efficiency of less than 200. In some embodiments, the patient has a diuretic efficiency of less than 150. In some embodiments, the patient has a diuretic efficiency of less than 100.

[0024] In some embodiments, the patient has diuretic resistant heart failure. In certain embodiments, the patient has acute heart failure. In certain embodiments, the patient has chronic heart failure.

[0025] In some embodiments, the patient has cardiorenal syndrome. In some of these embodiments, the patient has cardiorenal syndrome type 4.

[0026] In some embodiments, the patient has kidney disease. In certain embodiments, the patient has hepatorenal syndrome.

[0027] In some embodiments, the patient has at least one copy of the *TMPRSS6* rs855791 major allele.

[0028] In certain embodiments, the IL-6 antagonist is an anti-IL-6 antibody, or antigen-binding fragment or derivative thereof. In particular embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative has a K_D for binding human IL-6 of less than 100 nM, less than 50 nM, less than 10 nM, even less than 1 nM.

[0029] In certain embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative has an elimination half-life following intravenous administration of at least 7 days, at least 14 days, at least 21 days, or at least 30 days.

[0030] In certain embodiments, the IL-6 antagonist is a full-length monoclonal anti-IL-6 antibody. In particular embodiments, the antibody is an IgG1 or IgG4 antibody. In certain embodiments, the antibody is an IgG1 antibody.

[0031] In certain embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative is fully human. In certain embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative is humanized.

[0032] In certain embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative comprises all six variable region CDRs of MED5117. In specific embodiments, the antibody comprises the VH and VL of MED5117. In particular embodiments, the antibody is MED5117.

[0033] In certain embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative comprises all six variable region CDRs of an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, elsilimomab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb). In certain embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative comprises the heavy chain V region and light chain V region from an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb). In particular embodiments, the anti-IL-6 antibody or antigen-binding fragment or derivative is an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb).

[0034] In certain embodiments, the IL-6 antagonist is a single domain antibody, a V_{HH} Nanobody, an Fab, or a scFv.

[0035] In certain embodiments, the IL-6 antagonist is an anti-IL-6R antibody, or antigen-binding fragment or derivative thereof. In certain embodiments, the anti-IL-6R antibody, antigen-binding fragment, or derivative comprises the 6 CDRs of tocilizumab or vobarilizumab.

[0036] In certain embodiments, the IL-6 antagonist is a JAK inhibitor. In certain embodiments, the JAK inhibitor is selected from the group consisting of tofacitinib (Xeljanz), decemotinib, ruxolitinib, upadacitinib, baricitinib, filgotinib, lestaurtinib, pacritinib, peficitinib, INCB-039110, ABT-494, INCB-047986 and AC-410.

[0037] In certain embodiments, the IL-6 antagonist is a STAT3 inhibitor.

[0038] In certain embodiments, the IL-6 antagonist is administered parenterally. In particular embodiments, the IL-6 antagonist is administered subcutaneously.

[0039] In certain embodiments, the IL-6 antagonist is administered orally.

[0040] In certain embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase diuretic efficiency. In certain embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase diuretic efficiency to normal levels. In certain embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase eGFR. In particular embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase eGFR to normal levels.

[0041] In certain embodiments, the method further comprises the subsequent step of determining the level of IL-6 in urine, determining the level of IL-6 in plasma, or determining the level of IL-6 in urine and in plasma. In particular embodiments, the method further comprises a final step of adjusting the dose of IL-6 antagonist for subsequent administration based on IL-6 level determined in the immediately preceding step.

7. BRIEF DESCRIPTION OF THE DRAWINGS

[0042] The following detailed description of preferred embodiments of the invention will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, there are shown in the drawings embodiments which are presently preferred. It should be understood, however, that the invention is not limited to the precise arrangements and instrumentalities of the embodiments shown in the drawings.

[0043] **FIG. 1A** is a bar graph showing diuretic efficiency (“DE”) by tertiles of urine IL-6 in 129 heart failure (“HF”) patients receiving high dose diuretic therapy who were enrolled in the study described in Example 1 herein. Whiskers extend from the 10th to 90th percentile.

[0044] **FIG. 1B** is a bar graph showing diuretic efficiency (“DE”) by tertiles of plasma IL-6 in the 129 heart failure (“HF”) patients whose data is shown in FIG. 1A. Whiskers extend from the 10th to 90th percentile.

[0045] **FIG. 2A** plots DE by tertiles of urine IL-6 in the patients reported in FIGS. 1A and 1B further stratified by genotype, with the left panel showing results for patients homozygous for the *TMPRSS6* rs855791 SNP minor allele (2321G→A; A736V) and right panel showing results for patients having at least one copy of the *TMPRSS6* rs855791 SNP major allele (2321G; A736).

[0046] **FIG. 2B** plots DE by tertiles of plasma IL-6 in the patients reported in FIGS. 1A and 1B further stratified by genotype, with the left panel showing results for patients homozygous for the *TMPRSS6* rs855791 SNP minor allele (2321G→A; A736V) and right panel showing results for patients having at least one copy of the *TMPRSS6* rs855791 SNP major allele (2321G; A736).

[0047] **FIGS. 3A and 3B** plot the association in a 98 patient subset of the 129 patients reported in FIGS. 1 and 2 of (A) Urine IL-6 levels and (B) Plasma IL-6 levels with the following clinical determinations: Reduced kidney function, low diuretic efficiency (“DE”), increased neurohormonal activation, and risk of mortality. Whiskers represent 95% confidence interval (“CI”). All analyses adjusted for both urine and plasma levels of IL-6. Urine IL-6 levels are indexed to urinary creatinine. Due to the skewed distribution of urine and plasma IL-6 variables, a log transform was applied before performing logistic and Cox regressions. **Abbreviations:** OR=Odds ratio. HR=Hazards ratio. IL=interleukin.

SD=standard deviation. eGFR=estimated glomerular filtration rate. * = adjusted for use of angiotensin converting enzyme inhibitor (ACE-I) or angiotensin receptor blocker (ARB). ** = adjusted for baseline characteristics including age, race, amino terminal pro B-type natriuretic peptide (NT-proBNP), use of ACE-I or ARB, home loop diuretic dose, and eGFR.

[0048] **FIGS. 4A, 4B, and 4C** show the expression of NKCC2, ENaC-beta, and NCC in M1 CCD cells after treatment with IL-6 and/or Ruxolitinib, with **FIG. 4A** showing the expression of NKCC2 after the treatment with IL-6 and/or Ruxolitinib; **FIG. 4B** showing the expression

of ENaC-beta after the treatment with IL-6 and/or Ruxolitinib; and **FIG. 4C** showing the expression of NCC after the treatment with IL-6 and/or Ruxolitinib.

[0049] **FIGS. 5A, 5B, and 5C** show the expression of NKCC2, ENaC-beta, and NCC in M1 CCD cells after treatment with IL-6 and/or Spironolactone, with **FIG. 5A** showing the expression of NKCC2 after the treatment with IL-6 and/or Spironolactone; **FIG. 5B** showing the expression of ENaC-beta after the treatment with IL-6 and/or Spironolactone; and **FIG. 5C** showing the expression of NCC after the treatment with IL-6 and/or Spironolactone.

[0050] **FIG. 6** shows the association of baseline characteristics with higher levels of IL-6 in the PROTECT trial described in Example 5.

[0051] **FIGS. 7A and 7B** show the Kaplan-Meier survival curve by tertiles of IL-6, with **FIG. 7A** showing the all-cause mortality at 180 days and **FIG. 7B** showing all-cause mortality or cardiovascular related rehospitalization at 60 days.

[0052] **FIGS. 8A and 8B** show the Kaplan-Meier survival curve for change of IL-6 between baseline and day 7, with **FIG. 8A** showing the all-cause mortality at 180 days and **FIG. 8B** showing all-cause mortality or cardiovascular related rehospitalization at 60 days.

[0053] **FIG. 9** shows the association of baseline characteristics with higher levels of IL-6 in the BIOSTAT-CHF study described in Example 6.

[0054] **FIGS. 10A and 10B** show the Kaplan-Meier survival curve by tertiles of IL-6, with **FIG. 10A** showing the all-cause mortality and/or rehospitalization for heart failure at 2 years and **FIG. 10B** showing all-cause mortality at 2 years.

[0055] **FIG. 11** show the analysis between ferritin levels and tertiles of IL-6.

8. DETAILED DESCRIPTION

8.1. Definitions

[0056] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

[0057] By “transmembrane protease serine 6 (*TMPRSS6*) polypeptide” is meant a polypeptide or fragment thereof having at least about 85% or greater amino acid identity to the amino acid sequence provided at NCBI Accession No. NP_001275929 and having serine

proteinase activity. The *TMPRSS6* polypeptide, also known as Matriptase-2 (MT2), cleaves hemojuvelin and inhibits bone morphogenetic protein signaling. An exemplary *TMPRSS6* amino acid sequence having an alanine at position 736 (736A) is provided below:

```

1      MPVAEAPQVA GGQGDGGDGE EAEPEGMFKA CEDSKRKARG YLRLVPLFVL
LALLVLASAG
61     VLLWYFLGYK AEVMVSQVYS GSLRVLNRHF SQDLTRRESS AFRSETAKAQ
KMLKELITST
121    RLGTYYNSSS VYSFGEGPLT CFFWFILQIP EHRLMLSPV VVQALLVEEL
LSTVNSSAAV
181    PYRAEYVDP EGLVILEASV KDIAALNSTL GCYRYSYVGQ QQVLRLLKGPD
HLASSCLWHL
241    QGPKDLMLKL RLEWTLAECR DRLAMYDVAG PLEKRLITSV YGCSRQEPVW
EVLASGAIMA
301    VVWKKGLHSY YDPFVLSVQP VVFQACEVNL TLDNRDLSQG VLSTPYFPSY
YSPQTHCSWH
361    LTVPSLDYGL ALWFDAYALR RQKYDLPTQ GQWTIQNRRL CGLRILQPYA
ERIPVVATAG
421    ITINFTSQIS LTGPGVRVHY GLYNQSDPCP GEFLCSVNGL CVPACDGVKD
CPNGLDERNC
481    VCRATFQCKE DSTCISLPKV CDGQPDCLNG SDEEQCQEGV PCGTFTFQCE
DRSCVKKPNP
541    QCDGRPDGRD GSDEEHCDG LQGPSSRIVG GAVSSEGEWP WQASLQVRGR
HICGGALIAD
601    RWVITAAHCF QEDSMASVTL WTVFLGKVWQ NSRWPGEVSF KVSRLLLHPY
HEEDSHDYDV
661    ALLQLDHPVV RSAAVRPVCL PARSHFFEPG LHCWITGWA LREGALRADA
VALFYGWRNQ
721    GSETCCCPIS NALQKADVQL IPQDLCEVY RYQVTPRMLC AGYRKGGKDA
CQDSSGGPLV
781    CKALSGRWFL AGLVSWGLGC GRPNYFGVYT RITGVISWIQ QVVT (SEQ ID NO:1)

```

[0058] An exemplary *TMPRSS6* amino acid sequence having a valine at position 736 (736V) is provided below:

```

1      MPVAEAPQVA GGQGDGGDGE EAEPEGMFKA CEDSKRKARG YLRLVPLFVL
LALLVLASAG
61     VLLWYFLGYK AEVMVSQVYS GSLRVLNRHF SQDLTRRESS AFRSETAKAQ
KMLKELITST
121    RLGTYYNSSS VYSFGEGPLT CFFWFILQIP EHRLMLSPV VVQALLVEEL
LSTVNSSAAV

```

```

181 PYRAEYEVDP EGLVILEASV KDIAALNSTL GCYRYSYVGQ GQVLRRLKGPD
HLASSCLWHL
24 QGPKDLMLKL RLEWTLAECR DRLAMYDVAG PLEKRLITSV YGCSRQEPVV
EVLASGAIMA
30 VVWKKGLHSY YDPFVLSVQP VVFQACEVNL TLDNR LDSQG VLSTPYFPSY
YSPQTHCSWH
36 LTVPSLDYGL ALWFDAYALR RQKYDLPCTQ GQWTIQNRRL CGLRILQPYA
ERIPVVATAG
42 ITINFTSQIS LTGPGVRVHY GLYNQSDPCP GEFLCSVNGL CVPACDGVKD
CPNGLDERNC
48 VCRATFQCKE DSTCISLPKV CDGQPDCLNG SDEEQCQEGV PCGTFTFQCE
DRSCVKKPNP
54 QCDGRPD CRD GSDEEHCD CG LQGPSSRIVG GAVSSEGEWP WQASLQVRGR
HICGGALIAD
60 RWVITAAHCF QEDSMAS TVL WTVFLGKVWQ NSRWPGEVSF KVSRI LLHPY
HEEDSHDYDV
66 ALLQLDHPVV RSAAVRPVCL PARSHFFEPG LHCWITGWGA LREGALRADA
VALFYGWRNQ
72 GSETCCCPIS NALQKVDVQL IPQDLCSEVY RYQVTPRMLC AGYRKGKKDA
CQGDSSGGPLV
78 CKALSGRWFL AGLVSWGLGC GRPNYFGVYT RITGVISWIQ QVVT (SEQ ID NO:2)

```

[0059] By “*TMPRSS6* nucleic acid molecule” is meant a polynucleotide encoding a *TMPRSS6* polypeptide (Matriptase-2; MT2). An exemplary *TMPRSS6* nucleic acid molecule sequence is provided at NCBI Accession No. NM_001289000. A *TMPRSS6* nucleic acid sequence having a G at nucleotide position 2321 (“G allele”; “major allele”) is provided below:

```

1 GGACAAACAG AGGCTCCTGA GGCCTGTGTG CAGGCCCGGC ACCTATCTGC
CGCTCCCAAA
61 GGATGCCCCG GCGCGAGGCC CCCCAGGTGG CTGGCGGGCA GGGGGACGGA
GGTGATGGCG
121 AGGAAGCGGA GCCGGAGGGG ATGTTCAAGG CCTGTGAGGA CTCCAAGAGA
AAAGCCCGGG
181 GCTACCTCCG CCTGGTGCCC CTGTTTGTGC TGCTGGCCCT GCTCGTGCTG
GCTTCGGCGG
241 GGGTGCTACT CTGGTATTTC CTAGGGTACA AGGCGGAGGT GATGGTCAGC
CAGGTGTACT
301 CAGGCAGTCT GCGTGACTC AATCGCCACT TCTCCAGGA TCTTACCCGC
CGGGAATCTA

```

361 GTGCCTTCCG CAGTGAAACC GCCAAAGCCC AGAAGATGCT CAAGGAGCTC
ATCACCAGCA
421 CCCGCCTGGG AACTTACTAC AACTCCAGCT CCGTCTATTC CTTTGGGGAG
GGACCCCTCA
481 CCTGCTTCTT CTGGTTCATT CTCCAAATCC CCGAGCACCG CCGGCTGATG
CTGAGCCCCG
541 AGGTGGTGCA GGCACGCTG GTGGAGGAGC TGCTGTCCAC AGTCAACAGC
TCGGCTGCCG
601 TCCCCTACAG GGCCGAGTAC GAAGTGGACC CCGAGGGCCT AGTGATCCTG
GAAGCCAGTG
661 TGAAAGACAT AGCTGCATTG AATTCCACGC TGGGTTGTTA CCGCTACAGC
TACGTGGGCC
721 AGGGCCAGGT CCTCCGGCTG AAGGGGCCTG ACCACCTGGC CTCCAGCTGC
CTGTGGCACC
781 TGCAGGGCCC CAAGGACCTC ATGCTCAAAC TCCGGCTGGA GTGGACGCTG
GCAGAGTGCC
841 GGGACCGACT GGCCATGTAT GACGTGGCCG GGCCCCTGGA GAAGAGGCTC
ATCACCTCGG
901 TGTACGGCTG CAGCCGCCAG GAGCCCGTGG TGGAGTTCTT GCGTCGGGG
GCCATCATGG
961 CGGTCTCTG GAAGAAGGGC CTGCACAGCT ACTACGACCC CTTCGTGCTC
TCCGTGCAGC
1021 CGGTGGTCTT CCAGGCCTGT GAAGTGAACC TGACGCTGGA CAACAGGCTC
GACTCCCAGG
1081 GCGTCCTCAG CACCCCGTAC TTCCCCAGCT ACTACTCGCC CCAAACCCAC
TGCTCCTGGC
1141 ACCTCACGGT GCCCTCTCTG GACTACGGCT TGGCCCTCTG GTTTGATGCC
TATGCACTGA
1201 GGAGGCAGAA GTATGATTTG CCGTGCAACC AGGGCCAGTG GACGATCCAG
AACAGGAGGC
1261 TGTGTGGCTT GCGCATCCTG CAGCCCTACG CCGAGAGGAT CCCCGTGGTG
GCCACGGCCG
1321 GGATCACCAT CAACTTCACC TCCCAGATCT CCCTCACCAG GCCCGGTGTG
CGGGTGCACT
1381 ATGGCTTGTA CAACCAGTCG GACCCCTGCC CTGGAGAGTT CCTCTGTTCT
GTGAATGGAC
1441 TCTGTGTCCC TGCCTGTGAT GGGGTCAAGG ACTGCCCCAA CGGCCTGGAT
GAGAGAACT
1501 GCGTTTGACAG AGCCACATTC CAGTGCAAAG AGGACAGCAC ATGCATCTCA
CTGCCCAAGG

1561 TCTGTGATGG GCAGCCTGAT TGTCTCAACG GCAGCGACGA AGAGCAGTGC
CAGGAAGGGG

1621 TGCCATGTGG GACATTCACC TTCCAGTGTG AGGACCGGAG CTGCGTGAAG
AAGCCCAACC

1681 CGCAGTGTGA TGGGCGGCCC GACTGCAGGG ACGGCTCGGA TGAGGAGCAC
TGTGACTGTG

1741 GCCTCCAGGG CCCCTCCAGC CGCATTTGTTG GTGGAGCTGT GTCCTCCGAG
GGTGAGTGGC

1801 CATGGCAGGC CAGCCTCCAG GTTCGGGGTC GACACATCTG TGGGGGGGCC
CTCATCGCTG

1861 ACCGCTGGGT GATAACAGCT GCCCACTGCT TCCAGGAGGA CAGCATGGCC
TCCACGGTGC

1921 TGTGGACCGT GTTCCTGGGC AAGGTGTGGC AGAACTCGCG CTGGCCTGGA
GAGGTGTCTT

1981 TCAAGGTGAG CCGCCTGCTC CTGCACCCGT ACCACGAAGA GGACAGCCAT
GACTACGACG

2041 TGGCGCTGCT GCAGCTCGAC CACCCGGTGG TGCCTCGGC CGCCGTGCGC
CCCGTCTGCC

2101 TGCCCGCGCG CTCCCACTTC TTCGAGCCCG GCCTGCACTG CTGGATTACG
GGCTGGGGCG

2161 CCTTGCGCGA GGGCGCCCTA CGGGCGGATG CTGTGGCCCT ATTTTATGGA
TGGAGAAACC

2221 AAGGCTCAGA GACATGTTGC TGCCCCATCA GCAACGCTCT GCAGAAAGTG
GATGTGCAGT

2281 TGATCCACAC GGACCTGTGC AGCGAGGTCT ATCGCTACCA GGTGACGCCA
CGCATGCTGT

2341 GTGCCGGCTA CCGCAAGGGC AAGAAGGATG CCTGTCAGGG TGA CTGAGGT
GGTCCGCTGG

2401 TGTGCAAGGC ACTCAGTGGC CGCTGGTTCC TGGCGGGGCT GGTGAGCTGG
GGCCTGGGCT

2461 GTGGCCGGCC TAACTACTTC GCGTCTACA CCCGCATCAC AGGTGTGATC
AGCTGGATCC

2521 AGCAAGTGGT GACCTGAGGA ACTGCCCCC TGCAAAGCAG GGCCACCTC
CTGGA CTGAG

2581 AGAGCCCAGG GCAACTGCCA AGCAGGGGGA CAAGTATTCT GCGGGGGGT
GGGGGAGAGA

2641 GCAGGCCCTG TGGTGGCAGG AGGTGGCATC TTGTCTCGTC CCTGATGTCT
GCTCCAGTGA

2701 TGGCAGGAGG ATGGAGAAGT GCCAGCAGCT GGGGGTCAAG ACGTCCCCTG
AGGACCCAGG

2761 CCCACACCCA GCCCTTCTGC CTCCCAATTC TCTCTCCTCC GTCCCCCTTCC
 TCCACTGCTG
 2821 CCTAATGCAA GGCAGTGGCT CAGCAGCAAG AATGCTGGTT CTACATCCCG
 AGGAGTGTCT
 2881 GAGGTGCGCC CCACTCTGTA CAGAGGCTGT TTGGGCAGCC TTGCCTCCAG
 AGAGCAGATT
 2941 CCAGCTTCGG AAGCCCCTGG TCTAACTTGG GATCTGGGAA TGGAAGGTGC
 TCCCATCGGA
 3001 GGGGACCCCTC AGAGCCCTGG AGACTGCCAG GTGGGCCTGC TGCCACTGTA
 AGCCAAAAGG
 3061 TGGGGAAGTC CTGACTCCAG GGTCTTGCC CCACCCCTGC CTGCCACCTG
 GGCCCTCACA
 3121 GCCCAGACCC TCACTGGGAG GTGAGCTCAG CTGCCCTTTG GAATAAAGCT
 GCCTGATCCA
 3181 AAAAAAAAAA AAAAAA (SEQ ID NO:3)

[0060] A *TMPRSS6* nucleic acid sequence having an A at nucleotide position 2321 is provided below:

1 GGACAAACAG AGGCTCCTGA GGCCTGTGTG CAGGCCCGGC ACCTATCTGC
 CGCTCCCAA
 61 GGATGCCCCGT GGCCGAGGCC CCCCAGGTGG CTGGCGGGCA GGGGGACGGA
 GGTGATGGCG
 121 AGGAAGCGGA GCCGGAGGGG ATGTTCAAGG CCTGTGAGGA CTCCAAGAGA
 AAAGCCCGGG
 181 GCTACCTCCG CCTGGTGCCC CTGTTTGTGC TGCTGGCCCT GCTCGTGCTG
 GCTTCGGCGG
 241 GGGTGCTACT CTGGTATTTC CTAGGGTACA AGGCGGAGGT GATGGTCAGC
 CAGGTGTACT
 301 CAGGCAGTCT GCGTGACTC AATCGCCACT TCTCCAGGA TCTTACCCGC
 CGGGAATCTA
 361 GTGCCTTCCG CAGTGAAACC GCCAAAGCCC AGAAGATGCT CAAGGAGCTC
 ATCACCAGCA
 421 CCCGCTGGG AACTTACTAC AACTCCAGCT CCGTCTATTC CTTTGGGGAG
 GGACCCCTCA
 481 CCTGCTTCTT CTGGTTCATT CTCCAAATCC CCGAGCACCG CCGGCTGATG
 CTGAGCCCCG
 541 AGGTGGTGCA GGCAC TGCTG GTGAGGAGC TGCTGTCCAC AGTCAACAGC
 TCGGCTGCCG
 601 TCCCCTACAG GGCCGAGTAC GAAGTGGACC CCGAGGGCCT AGTGATCCTG
 GAAGCCAGTG

661 TGAAAGACAT AGCTGCATTG AATTCCACGC TGGGTTGTTA CCGCTACAGC
TACGTGGGCC
721 AGGGCCAGGT CCTCCGGCTG AAGGGGCGTG ACCACCTGGC CTCCAGCTGC
CTGTGGCACC
781 TGCAGGGCCC CAAGGACCTC ATGCTCAAAC TCCGGCTGGA GTGGACGCTG
GCAGAGTGCC
841 GGGACCGACT GGCCATGTAT GACGTGGCCG GGCCCCTGGA GAAGAGGCTC
ATCACCTCGG
901 TGTACGGCTG CAGCCGCCAG GAGCCCGTGG TGGAGGTTCT GCGTCGGGG
GCCATCATGG
961 CGGTCGTCTG GAAGAAGGGC CTGCACAGCT ACTACGACCC CTTCGTGCTC
TCCGTGCAGC
1021 CGGTGGTCTT CCAGGCCTGT GAAGTGAACC TGACGCTGGA CAACAGGCTC
GACTCCCAGG
1081 GCGTCCTCAG CACCCCGTAC TTCCCAGCT ACTACTCGCC CCAAACCCAC
TGCTCCTGGC
1141 ACCTCACGGT GCCCTCTCTG GACTACGGCT TGGCCCTCTG GTTTGATGCC
TATGCACTGA
1201 GGAGGCAGAA GTATGATTTG CCGTGCACCC AGGGCCAGTG GACGATCCAG
AACAGGAGGC
1261 TGTGTGGCTT GCGCATCCTG CAGCCCTACG CCGAGAGGAT CCCCCTGGTG
GCCACGGCCG
1321 GGATCACCAT CAACTTCACC TCCCAGATCT CCCTCACC GGCCCGGTGTG
CGGGTGCACT
1381 ATGGCTTGTA CAACCAGTCG GACCCCTGCC CTGGAGAGTT CCTCTGTTCT
GTGAATGGAC
1441 TCTGTGTCCC TGCCTGTGAT GGGGTCAAGG ACTGCCCCAA CGGCCTGGAT
GAGAGAAACT
1501 GCGTTTGCAG AGCCACATTC CAGTGCAAAG AGGACAGCAC ATGCATCTCA
CTGCCCAAGG
1561 TCTGTGATGG GCAGCCTGAT TGTCTCAACG GCAGCGACGA AGAGCAGTGC
CAGGAAGGGG
1621 TGCCATGTGG GACATTCACC TTCCAGTGTG AGGACCGGAG CTGCGTGAAG
AAGCCCAACC
1681 CGCAGTGTGA TGGGCGGCCC GACTGCAGGG ACGGCTCGGA TGAGGAGCAC
TGTGACTGTG
1741 GCCTCCAGGG CCCCTCCAGC CGCATTGTTG GTGGAGCTGT GTCCTCCGAG
GGTGAGTGGC
1801 CATGGCAGGC CAGCCTCCAG GTTCGGGGTC GACACATCTG TGGGGGGGCC
CTCATCGCTG

1861 ACCGCTGGGT GATAACAGCT GCCCACTGCT TCCAGGAGGA CAGCATGGCC
TCCACGGTGC
1921 TGTGGACCGT GTTCCTGGGC AAGGTGTGGC AGAACTCGCG CTGGCCTGGA
GAGGTGTCCT
1981 TCAAGGTGAG CCGCCTGCTC CTGCACCCGT ACCACGAAGA GGACAGCCAT
GACTACGACG
2041 TGGCGCTGCT GCAGCTCGAC CACCCGGTGG TGCCTCGGC CGCCGTGCGC
CCCGTCTGCC
2101 TGCCCGCGCG CTCCCACTTC TTCGAGCCCG GCCTGCACTG CTGGATTACG
GGCTGGGGCG
2161 CCTTGCGCGA GGGCGCCCTA CGGGCGGATG CTGTGGCCCT ATTTTATGGA
TGGAGAAACC
2221 AAGGCTCAGA GACATGTTGC TGCCCCATCA GCAACGCTCT GCAGAAAGTG
GATGTGCAGT
2281 TGATCCACA GGACCTGTGC AGCGAGGTCT ATCGCTACCA AGTGACCCA
CGCATGCTGT
2341 GTGCCGGCTA CCGCAAGGGC AAGAAGGATG CCTGTCAGGG TGA CTCAAGT
GGTCCGCTGG
2401 TGTGCAAGGC ACTCAGTGGC CGCTGGTTCC TGGCGGGGCT GGTCAAGTGG
GGCCTGGGCT
2461 GTGGCCGGCC TAACTACTTC GGCGTCTACA CCCGCATCAC AGGTGTGATC
AGCTGGATCC
2521 AGCAAGTGGT GACCTGAGGA ACTGCCCCC TGCAAAGCAG GGCCACCTC
CTGGACTCAG
2581 AGAGCCCAGG GCAACTGCCA AGCAGGGGGA CAAGTATTCT GGCGGGGGT
GGGGGAGAGA
2641 GCAGGCCCTG TGGTGGCAGG AGGTGGCATC TTGTCTCGTC CCTGATGTCT
GCTCCAGTGA
2701 TGGCAGGAGG ATGGAGAAGT GCCAGCAGCT GGGGGTCAAG ACGTCCCCTG
AGGACCCAGG
2761 CCCACACCCA GCCCTTCTGC CTCCCAATTC TCTCTCCTCC GTCCCCCTCC
TCCACTGCTG
2821 CCTAATGCAA GGCAGTGGCT CAGCAGCAAG AATGCTGGTT CTACATCCCG
AGGAGTGTCT
2881 GAGGTGCGCC CCACTCTGTA CAGAGGCTGT TTGGGCAGCC TTGCCTCCAG
AGAGCAGATT
2941 CCAGCTTCGG AAGCCCCTGG TCTAACTTGG GATCTGGGAA TGGAAGGTGC
TCCCATCGGA
3001 GGGGACCCTC AGAGCCCTGG AGACTGCCAG GTGGGCCTGC TGCCACTGTA
AGCCAAAAGG

```

3061 TGGGGAAGTC CTGACTCCAG GGTCTTGCC CCACCCCTGC CTGCCACCTG
GGCCCTCACA
3121 GCCCAGACCC TCACTGGGAG GTGAGCTCAG CTGCCCTTTG GAATAAAGCT
GCCTGATCCA
3181 AAAAAAAAAA AAAAAA (SEQ ID NO:4)

```

[0061] By “**variant**” is meant a polynucleotide or polypeptide sequence that differs from a reference sequence by one or more nucleotides or one or more amino acids. An exemplary *TMPRSS6* variant is *TMPRSS6* (A736V), resulting from SNP rs855791 (G→A).

[0062] By “**single nucleotide polymorphism**” or “**SNP**” is meant a naturally occurring DNA sequence variant in which a single nucleotide in the genome differs between members of a biological species or between paired chromosomes in an individual. SNPs can be used as genetic markers for variant alleles. In one embodiment, the *TMPRSS6* SNP is rs855791.

[0063] By “**rs855791**” is meant a single nucleotide polymorphism (SNP) in the human *TMPRSS6* gene, 2321G→A, resulting in an alanine to valine substitution (A736V) in the catalytic domain of Matriptase-2 (MT2), which is encoded by the *TMPRSS6* gene. The allele with highest frequency in the human population (the major allele) is 2321G, encoding 736A. The allele with lowest frequency in the human population (minor allele) is 2321A, encoding 736V.

[0064] By “**heterozygous**” is meant that a chromosomal locus has two different alleles. In one embodiment of the methods described herein, heterozygous refers to a genotype in which one allele has a *TMPRSS6* nucleic acid sequence encoding a *TMPRSS6* polypeptide having an alanine at amino acid position 736 (e.g., having a G or C at nucleotide position 2321 of a *TMPRSS6* nucleic acid molecule) (rs855791 major allele), and the other allele has a variant *TMPRSS6* nucleic acid sequence encoding a *TMPRSS6* polypeptide comprising a valine at amino acid position 736 (e.g., having an A or T at nucleotide position 2321 of a *TMPRSS6* nucleic acid molecule) (rs855791 minor allele).

[0065] By “**homozygous**” is meant that a chromosomal locus has two identical alleles. In certain embodiments of the methods described herein, homozygous refers to a genotype in which both alleles have a *TMPRSS6* nucleic acid sequence encoding a *TMPRSS6* polypeptide comprising an alanine at amino acid position 736 (e.g., having a G or C at nucleotide position 2321 of a *TMPRSS6* nucleic acid molecule) (rs855791 homozygous major allele). In certain embodiments, homozygous refers to a genotype in which both alleles have a *TMPRSS6* nucleic acid sequence encoding a *TMPRSS6* polypeptide comprising a valine at amino acid

position 736 (e.g., having an A or T at nucleotide position 2321 of a *TMPRSS6* nucleic acid molecule) (rs855791 homozygous minor allele).

[0066] “**Determining that a patient has at least one copy of the *TMPRSS6* rs855791 major allele**” includes, but is not limited to, performing an assay to determine that a patient has at least one copy of the *TMPRSS6* rs855791 major allele; ordering an assay to determine that a patient has at least one copy of the *TMPRSS6* rs855791 major allele; prescribing an assay to determine that a patient has at least one copy of the *TMPRSS6* rs855791 major allele; otherwise directing or controlling that an assay be performed to determine that a patient has at least one copy of the *TMPRSS6* rs855791 major allele; and reviewing *TMRSS6* genotype assay data or protein or nucleic acid sequence data to determine that a patient has at least one copy of the *TMPRSS6* rs855791 major allele.

[0067] By “**interleukin 6**” or “**IL-6**” or “**IL-6 polypeptide**” is meant a polypeptide or fragment thereof having at least about 85% or greater amino acid identity to the amino acid sequence provided at NCBI Accession No. NP_000591 and having IL-6 biological activity. IL-6 is a pleiotropic cytokine with multiple biologic functions. Exemplary IL-6 biological activities include immunostimulatory and pro-inflammatory activities. An exemplary IL-6 amino acid sequence is provided below:

```

1  MCVGARRLGR  GPCAALLLLG  LGLSTVTGLH  CVGDTYPSND  RCHECRPGN  GMVSRCSRSQ
61  NTVCRPCGPG  FYNDVVSSKP  CKPCTWCNLR  SGSERKQLCT  ATQDTVCRCR  AGTQPLDSYK
121 PGVDCAPCPF  GHFSPGDNQA  CKPWTNCTLA  GKHTLQPASN  SSDAICEDRD  PPATQPQETQ
181 GPPARPITVQ  PTEAWPRTSQ  GPSTRPVEVP  GGRAVAAILG  LGLVLGLLGP  LAILLALYLL
241 RRDQRLPPDA  HKPPGGGSFR  TPIQEEQADA  HSTLAKI  (SEQ ID NO:5)

```

[0068] By “**interleukin 6 (IL-6) nucleic acid**” is meant a polynucleotide encoding an interleukin 6 (IL-6) polypeptide. An exemplary interleukin 6 (IL-6) nucleic acid sequence is provided at NCBI Accession No. NM_000600. The exemplary sequence at NCBI Accession No. NM_000600 is provided below.

```

1  AATATTAGAG  TCTCAACCCC  CAATAAATAT  AGGACTGGAG  ATGTCTGAGG
CTCATTTCTGC
61  CCTCGAGCCC  ACCGGAACG  AAAGAGAAGC  TCTATCTCCC  CTCCAGGAGC
CCAGCTATGA
121 ACTCCTTCTC  CACAAGCGCC  TTCGGTCCAG  TTGCCTTCTC  CCTGGGGCTG
CTCCTGGTGT
181 TGCCTGCTGC  CTTCCCTGCC  CCAGTACCCC  CAGGAGAAGA  TTCCAAAGAT
GTAGCCGCCC

```

```

241 CACACAGACA GCCACTCACC TCTTCAGAAC GAATTGACAA ACAAATTCGG
TACATCCTCG
301 ACGGCATCTC AGCCCTGAGA AAGGAGACAT GTAACAAGAG TAACATGTGT
GAAAGCAGCA
361 AAGAGGCACT GGCAGAAAAC AACCTGAACC TTCCAAAGAT GGCTGAAAAA
GATGGATGCT
421 TCCAATCTGG ATTCAATGAG GAGACTTGCC TGGTGAAAAT CATCACTGGT
CTTTTGGAGT
481 TTGAGGTATA CCTAGAGTAC CTCCAGAACA GATTTGAGAG TAGTGAGGAA
CAAGCCAGAG
541 CTGTGCAGAT GAGTACAAAA GTCCTGATCC AGTTCCTGCA GAAAAAGGCA
AAGAATCTAG
601 ATGCAATAAC CACCCCTGAC CCAACCACAA ATGCCAGCCT GCTGACGAAG
CTGCAGGCAC
661 AGAACCAGTG GCTGCAGGAC ATGACAACTC ATCTCATTCT GCGCAGCTTT
AAGGAGTTCC
721 TGCAGTCCAG CCTGAGGGCT CTTCGGCAA TGTAGCATGG GCACCTCAGA
TTGTTGTTGT
781 TAATGGGCAT TCCTTCTTCT GGTGAGAAAC CTGTCCACTG GGCACAGAAC
TTATGTTGTT
841 CTCTATGGAG AACTAAAAGT ATGAGCGTTA GGACACTATT TTAATTATTT
TTAATTATTT
901 AATATTTTAAA TATGTGAAGC TGAGTTAATT TATGTAAGTC ATATTTATAT
TTTTAAGAAG
961 TACCACTTGA AACATTTTAT GTATTAGTTT TGAAATAATA ATGGAAAGTG
GCTATGCAGT
1021 TTGAATATCC TTTGTTTCAG AGCCAGATCA TTTCTTGGAA AGTGTAGGCT
TACCTCAAAT
1081 AAATGGCTAA CTTATACATA TTTTAAAGA AATATTTATA TTGTATTTAT
ATAATGTATA
1141 AATGGTTTTT ATACCAATAA ATGGCATT TTAAAAATTC GCAAAAAAAA
AAAAA
1201 A (SEQ ID NO:6)

```

[0069] By “**interleukin 6 receptor (IL-6R) complex**” is meant a protein complex comprising an IL-6 receptor subunit alpha (IL-6R α) and interleukin 6 signal transducer Glycoprotein 130, also termed interleukin 6 receptor subunit β (IL-6R β).

[0070] By “**interleukin 6 receptor subunit α (IL-6R α) polypeptide**” is meant a polypeptide or fragment thereof having at least about 85% or greater amino acid identity to the amino acid sequence provided at NCBI Accession No. NP_000556 or NP_852004 and having IL-6

receptor biological activity. Exemplary IL-6 α biological activities include binding to IL-6, binding to glycoprotein 130 (gp130), and regulation of cell growth and differentiation. An exemplary IL-6R sequence is provided below:

```

1 MLAVGCALLA ALLAAPGAAL APRRCPAQEV ARGVLTSLPG DSVTLTCEPGV EPEDNATVHW
61 VLRKPAAGSH PSRWAGMGRR LLLRSVQLHD SGNYSYRAG RPAGTVHLLV DVPPEEPQLS
121 CFRKSPLSNV VCEWGPRSTP SLTTKAVLLV RKFQNSPAED FQEPQYSQE SQKFSCQLAV
181 PEGDSSFYIV SMCVASSVGS KFSKTQTFQG CGILQPDPPA NITVTAVARN PRWLSVTWQD
241 PHSWNSSFYR LRFELRYRAE RSKTFTTMV KDLQHHCVIH DAWSGLRHVV QLRAQEEFGQ
301 GEWSEWSPEA MGTPWTESRS PPAENEVSTP MQALTTNKDD DNILFRDSAN ATSLPVQDSS
361 SVPLPTFLVA GGSLAFGTLL CIAIVLRFKK TWKLRALKEG KTSMHPPYSL GQLVPERPRP
421 TPVLVPLISP FVSPSSLGSD NTSSHNRDPA RDPRSPYDIS NTDYFFPR (SEQ ID
NO: 7)

```

[0071] By “interleukin 6 receptor subunit β (IL-6R β) polypeptide” is meant a polypeptide or fragment thereof having at least about 85% or greater amino acid identity to the amino acid sequence provided at NCBI Accession No. NP_002175, NP_786943, or NP_001177910 and having IL-6 receptor biological activity. Exemplary IL-6R β biological activities include binding to IL-6 α , IL-6 receptor signaling activity, and regulation of cell growth, differentiation, hepcidin expression etc. An exemplary IL-6R β sequence is provided below:

```

1 MLTLQTWLVQ ALFIFLTES TGEILDPCGY ISPEPVVQL HSNFTAVCVL KEKCMDYFHW
61 NANYIVWKTN HFTIPKEQYT IINRTASSVT FTDIASLNIQ LTCNLTFTGQ LEQNVYGITI
121 ISGLPPEKPK NLSCIVNEGK KMRCEWDGGR ETHLETNFTL KSEWATHKFA DCAKARDTPT
181 SCTVDYSTVY FVNIEVVEA ENALGKVTSD HINFDPVYKV KPNPPHNLV INSEELSSIL
241 KLTWTNPSIK SVIILKYNIQ YRTKASTWS QIPPEDTAST RSSFTVQDLK PFTBYVFRIR
301 CMKEDGKGYW SDWSEEASGI TYEDRPSKAP SFWYKIDPSH TQGYRTVQLV WKTLPPEAN
361 GKILDYEVTL TRWKSHLQNY TVNATKLTVN LTNDRYLATL TVRNLVGKSD AAVLTIPACD
421 FQATHPVMDL KAFPKDNMLW VEWTPRESV KKYILEWCVL SDKAPCITDW QQEDGTVHRT
481 YLRGNLAESK CYLITVTPVY ADGPGSPESI KAYLKQAPPS KGPTVTRTKV GKNEAVLEWD
541 QLPVDVQNGF IRNYTIFYRT IIGNETAVNV DSSHTYETLS SLTSDTLYMV RMAAYTDEGG
601 KDGPEFTFTT PKFAQGEIEA IVPVCLAFLL LTTLLGVLFV FNKRDLIKKH IWPNVDPDSK
661 SHIAQWSPHT PPRHNFNSKD QMYSDGNFTD VSVVEIEAND KKPFPEDLKS LDLFKKEKIN
721 TEGHSSGIGG SSCMSSSRPS ISSSDENESS QNTSSTVQYS TVVHSGYRHQ VPSVQVFSRS
781 ESTQPLLDSE ERPEDLQLVD HVDGGDGILP RQYFKQNC QHESPDISH FERSKQVSSV
841 NEEDFVRLKQ QISDHISQSC GSGQMKMFQE VSAADAFGPG TEGQVERFET VGMEAATDEG
901 MPKSYLPQTV RQGGYMPQ (SEQ ID NO: 8)

```

[0072] By “IL-6 antagonist” is meant an agent that is capable of decreasing the biological activity of IL-6. IL-6 antagonists include agents that decrease the level of IL-6 polypeptide in

serum, including agents that decrease the expression of an IL-6 polypeptide or nucleic acid; agents that decrease the ability of IL-6 to bind to the IL-6R; agents that decrease the expression of the IL-6R; and agents that decrease signal transduction by the IL-6R receptor when bound by IL-6. In preferred embodiments, the IL-6 antagonist decreases IL-6 biological activity by at least about 10%, 20%, 30%, 50%, 70%, 80%, 90%, 95%, or even 100%. As further described in Section 6.3.4 below, IL-6 antagonists include IL-6 binding polypeptides, such as anti-IL-6 antibodies and antigen binding fragments or derivatives thereof; IL-6R binding polypeptides, such as anti-IL-6R antibodies and antigen binding fragments or derivatives thereof; and synthetic chemical molecules, such as JAK1 and JAK3 inhibitors.

[0073] By “**IL-6 antibody**” or “**anti-IL-6 antibody**” is meant an antibody that specifically binds IL-6. Anti-IL-6 antibodies include monoclonal and polyclonal antibodies that are specific for IL-6, and antigen-binding fragments or derivatives thereof. IL-6 antibodies are described in greater detail in Section 8.3.6.1 below.

[0074] As used herein, “**diuretic efficiency**” is calculated as mmol urinary sodium per doubling of loop diuretic dose (mmol Na/doubling of loop diuretic dose) according to the methods described in Hanberg *et al.*, *Circ. Heart Fail.* 2016;9:e003180, the disclosure of which is incorporated herein by reference in its entirety.

[0075] By “**diuretic resistant heart failure**” is meant heart failure in which the patient’s diuretic efficiency is less than 100.

[0076] The terms “**biomarker**” or “**marker**,” as used herein, refers to a molecule that can be detected. Therefore, a biomarker according to the present invention includes, but is not limited to, a nucleic acid, a polypeptide, a carbohydrate, a lipid, an inorganic molecule, an organic molecule, each of which may vary widely in size and properties. A “biomarker” can be a bodily substance relating to a bodily condition or disease. A “biomarker” can be detected using any means known in the art or by a previously unknown means that only becomes apparent upon consideration of the marker by the skilled artisan.

[0077] As used herein, “**biomarker**” in the context of the present invention encompasses, without limitation, proteins, nucleic acids, and metabolites, together with their polymorphisms, mutations, variants, modifications, subunits, fragments, protein-ligand complexes, and degradation products, elements, related metabolites, and other analytes or sample-derived measures. Biomarkers can also include mutated proteins or mutated nucleic

acids. Biomarkers also encompass non-blood borne factors or non-analyte physiological markers of health status, such as clinical parameters, as well as traditional laboratory risk factors. As defined by the Food and Drug Administration (FDA), a biomarker is a characteristic (e.g. measurable DNA and/or RNA or a protein) that is “objectively measured and evaluated as an indicator of normal biologic processes, pathogenic processes, or pharmacologic responses to a therapeutic intervention or other interventions”. Biomarkers also include any calculated indices created mathematically or combinations of any one or more of the foregoing measurements, including temporal trends and differences. Biomarkers may be measured at any level spatial or temporal localization, including but not limited to within a tumor, within in a cell, or on the membrane of a cell.

[0078] By “**agent**” is meant any compound or composition suitable to be administered in therapy, and explicitly includes chemical compounds; proteins, including antibodies or antigen-binding fragments thereof; peptides; and nucleic acid molecules.

[0079] By “**subject**” or “**individual**” is meant a human or non-human mammal, including, but not limited to, bovine, equine, canine, ovine, feline, and rodent, including murine and rattus, subjects. A “**patient**” is a human subject.

[0080] As used herein, the terms “**treat**,” “**treating**,” “**treatment**,” and the like refer to reducing or ameliorating a disorder, and/or signs or symptoms associated therewith, or slowing or halting the progression thereof. It will be appreciated that, although not precluded, treating a disorder or condition does not require that the disorder, condition or symptoms associated therewith be completely eliminated.

[0081] “**Pre-treatment**” means prior to the first administration of an IL-6 antagonist according the methods described herein. Pre-treatment does not exclude, and often includes, the prior administration of treatments other than an IL-6 antagonist, such as treatment with a diuretic, such as a loop diuretic.

[0082] By “**biological sample**” is meant any tissue, cell, fluid, or other material derived from an organism (e.g., human subject). In certain embodiments, the biological sample is serum, plasma, urine, or whole blood.

[0083] As used herein, an “**instructional material**” includes a publication, a recording, a diagram, or any other medium of expression which can be used to communicate the usefulness of a component of the invention in a kit for detecting biomarkers disclosed herein. The instructional material of the kit of the invention can, for example, be affixed to a

container which contains the component of the invention or be shipped together with a container which contains the component. Alternatively, the instructional material can be shipped separately from the container with the intention that the instructional material and the component be used cooperatively by the recipient.

[0084] The “**level**” of one or more biomarkers means the absolute or relative amount or concentration of the biomarker in the sample as determined by measuring mRNA, cDNA, small organic molecules, nucleotides, ions or protein, or any portion thereof such as oligonucleotide or peptide. A level of a biomarker may refer, based on context, to a global level or a level within some subdivision of an organism or within a specific sample, by way of non-limiting example a level may refer to the amount or concentration of a biomarker in a urine sample or the level may refer to the amount or concentration of the same biomarker in a plasma sample.

[0085] “**Measuring**” or “**measurement,**” or alternatively “**detecting**” or “**detection,**” means determining the presence, absence, quantity or amount (which can be an effective amount) of either a given substance within a clinical or subject-derived sample, including the derivation of qualitative or quantitative concentration levels of such substances, or otherwise determining the values or categorization of a subject's clinical parameters.

[0086] A “**reference level**” of a biomarker means a level of a biomarker that is indicative of the presence or absence of a particular phenotype or characteristic. When the level of a biomarker in a subject is above the reference level of the biomarker it is indicative of the presence of, or relatively heightened level of, a particular phenotype or characteristic. When the level of a biomarker in a subject is below the reference level of the biomarker it is indicative of a lack of or relative lack of a particular phenotype or characteristic.

8.1. Other interpretational conventions

[0087] Unless otherwise specified, antibody constant region residue numbering is according to the EU index as in Kabat.

[0088] Ranges: throughout this disclosure, various aspects of the invention can be presented in a range format. Ranges include the recited endpoints. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible

subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. This applies regardless of the breadth of the range.

[0089] Unless specifically stated or apparent from context, as used herein the term “or” is understood to be inclusive.

[0090] Unless specifically stated or apparent from context, as used herein, the terms “a”, “an”, and “the” are understood to be singular or plural. That is, the articles “a” and “an” are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0091] In this disclosure, “comprises,” “comprising,” “containing,” “having,” “includes,” “including,” and linguistic variants thereof have the meaning ascribed to them in U.S. Patent law, permitting the presence of additional components beyond those explicitly recited.

[0092] Unless specifically stated or otherwise apparent from context, as used herein the term “about” is understood as within a range of normal tolerance in the art, for example within 2 standard deviations of the mean and is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, more preferably $\pm 5\%$, even more preferably $\pm 1\%$, and still more preferably $\pm 0.1\%$ from the stated value.

[0093] Where an antibody equilibrium dissociation constant (K_D) is reported, K_D is determined by surface plasmon resonance with the antibody (or antigen-binding fragment thereof) fixed to the chip surface with ligand flowed thereover.

8.2. Summary of experimental observations

[0094] As further described below in Example 1, consecutive heart failure (“HF”) patients receiving high dose diuretic therapy at an outpatient treatment center were enrolled in a prospective observational study. Plasma levels of IL-6 were measured to query systemic associations of this pro-inflammatory cytokine with various disease parameters, and urine levels of IL-6 were measured to query IL-6 associations with local inflammation and neurohormonal activation at the level of renal tissues.

[0095] Plasma and urine IL-6 levels were only modestly correlated with one another.

[0096] Increases in urine levels of IL-6 were significantly correlated in these heart failure patients with measures of renal impairment, such as diuretic resistance, lower estimated glomerular filtration rate (“eGFR”), and increased tissue-level renin-angiotensin-aldosterone system (“RAAS”) activation.

[0097] Although an inverse association between diuretic efficiency and plasma IL-6 was also observed, upon adjustment for eGFR only urine IL-6 remained significantly associated with risk of low diuretic efficiency in these patients. Furthermore, when urine IL-6 and plasma IL-6 were both entered into a logistic regression model, only urine IL-6 remained associated with risk of low diuretic efficiency while plasma IL-6 showed no such association.

[0098] These data demonstrate that urine IL-6 level is a useful biomarker for renal inflammation, and can be used to gauge renal dysfunction in the setting of heart failure (cardiorenal syndrome). The data further suggest that serial measurements of urine IL-6 can be used to measure the renal benefits of treatments administered to patients with heart failure, notably heart failure patients with cardiorenal syndrome.

[0099] The urine IL-6 data, and to some extent the plasma IL-6 data, predict that treatment with an IL-6 antagonist should be effective to reduce renal inflammation in heart failure patients, that is, to treat symptoms of cardiorenal syndrome. However, because infection is often a precipitating cause of acute decompensation in heart failure patients, it is important to limit anti-cytokine and other immunosuppressive therapies to those heart failure patients who are likely to respond with improved renal and/or cardiac function. The cost of chronic IL-6 antagonist therapy also militates for limiting treatment to those heart failure patients who are likely to respond with improved renal and/or cardiac function.

[0100] As detailed below in Example 2, the analysis conducted in Example 1 was expanded to 129 patients. FIG. 1A is a bar graph showing diuretic efficiency (“DE”) by tertiles of urine IL-6, confirming the inverse correlation of urinary IL-6 level with diuretic efficiency observed in the 98 patient subset. FIG. 1B is a bar graph showing diuretic efficiency (“DE”) by tertiles of plasma IL-6 in these 129 patients, confirming an inverse correlation of plasma IL-6 levels with diuretic efficiency.

[0101] The genotype of each of the 129 patients at the rs855791 single nucleotide polymorphism (“SNP”) in transmembrane protease serine 6 (“TMPRSS6”) was assessed. The TMPRSS6 polypeptide, also known as Matriptase-2 (MT2), cleaves hemojuvelin and inhibits bone morphogenetic protein signaling. The rs855791 (G2321A) SNP alters the

TMPRSS6 protein sequence: the allele with highest frequency in the human population (the major allele) is 2321G, encoding 736A; the allele with lowest frequency in the human population (minor allele) is 2321A, encoding 736V.

[0102] As shown in FIG. 2A, urine levels of IL-6 were inversely correlated with diuretic efficiency only in the patients having at least one copy of the major allele of the TMPRSS6 rs855791 SNP (FIG. 2A, right panel, “AG+GG”); urine levels of IL-6 are not significantly correlated with diuretic efficiency in patients homozygous for the minor allele (FIG. 2A, left panel, “AA”).

[0103] As shown in FIG. 2B, plasma levels of IL-6 correlated inversely with diuretic efficiency only in the patients having at least one copy of the major allele of the TMPRSS6 rs855791 SNP (FIG. 2B, right panel, “AG+GG”); plasma levels of IL-6 are not significantly correlated with diuretic efficiency in patients homozygous for the minor allele (FIG. 2B, left panel, “AA”).

[0104] These data suggested that diuretic resistance (low diuretic efficiency) in heart failure patients could be treated with an IL-6 antagonist, but only in those having at least one copy of the TMPRSS6 rs855791 major allele.

[0105] In mouse M1 CCD cells, which are genotypically analogous to human cells homozygous for the *TMPRSS6* rs855791 major allele, the addition of IL-6 correlated with the expression of ion transporters, NKCC2, ENaC-beta, and NCC. Increased expression of these ion transporters provides a putative mechanism for IL-6 mediated diuretic resistance. And because the increased expression is not known to be linked to hepcidin expression, these data suggested that IL-6 antagonism could be effective in treating diuretic resistance even in patients homozygous for the *TMPRSS6* rs855791 minor allele.

[0106] Secondary analysis of data from two large clinical trials in different heart failure patient populations confirmed the association of diuretic resistance with IL-6 level (Example 5), independently of *TMPRSS6* rs855791 genotype (Example 6), providing evidence that IL-6 antagonism should also be effective in treating diuretic resistance in patients homozygous for the *TMPRSS6* rs855791 minor allele.

8.3. Methods of treating diuretic resistance

[0107] Accordingly, in a first aspect, methods are provided for treating a patient who requires diuresis but is resistant to diuretics. The methods comprise administering, in combination with a diuretic, a therapeutically effective amount of an IL-6 antagonist to the patient.

[0108] In certain embodiments, the patient has elevated pre-treatment urine levels of IL-6. In some embodiments, the patient has elevated pre-treatment plasma IL-6 levels. In certain embodiments, the patient has elevated pre-treatment levels of IL-6 in urine and in plasma.

[0109] In some embodiments, the patient has diuretic-resistant heart failure. In various embodiments, the patient has cardiorenal syndrome.

[0110] In some embodiments, the patient has been determined to have at least one copy of the TMPRSS6 rs855791 major allele. In other embodiments, the patient is homozygous for the TMPRSS6 rs855791 minor allele.

8.3.1. Diuretic efficiency

[0111] In the methods described herein, the patient in need of the IL-6 antagonist treatment has a disease or condition that requires diuresis, and is diuretic resistant.

[0112] In certain embodiments, the patient has been treated or is being treated with a thiazide diuretic, such as chlorothiazide (Diuril®), chlorthalidone, hydrochlorothiazide (Microzide®), indapamide, or metolazone. In certain embodiments, the patient has been treated or is being treated with a loop diuretic, such as bumetanide (Bumex®), ethacrynic acid (Edecrin®), furosemide (Lasix®), or torsemide (Demadex®). In certain embodiments, the patient has been treated or is being treated with a potassium-sparing diuretic, such as amiloride, eplerenone (Inspra®), spironolactone (Aldactone®), or triamterene (Dyrenium®). In some embodiments, the patient has been treated or is being treated with more than one diuretic. In some embodiments, the patient has been treated or is being treated with a plurality of different types of diuretics.

[0113] By definition, the patient resistant to diuretics has a low diuretic efficiency. Diuretic efficiency is calculated as the increase in sodium output per doubling of the loop diuretic dose, centered on a dose of 40 mg of IV furosemide equivalents: $\text{diuretic efficiency} = (\text{mmol Na output}) / (\log_2(\text{administered loop diuretic dose}) - 4.32)$. See Hanberg *et al.*, *Circ. Heart Fail.* 2016;9:e003180, the disclosure of which is incorporated herein by reference in its entirety.

[0114] In some embodiments, the patient has a diuretic efficiency of less than 500, such as less than 450, 400, 350, 300, 250, or 200. In some embodiment, the patient has a diuretic efficiency of less than 200, such as less than 195, 190, 185, 180, 175, 170, 165, 160, 155, or 150. In some embodiments, the patient has a diuretic efficiency of less than 150, such as less than 145, 140, 135, 130, 125, 120, 110, 105, or 100. In some embodiments, the patient has a diuretic efficiency of less than 100, such as less than 95, 90, 85, 80, 75, 70, 65, 60, 55, or 50. In particular embodiments, the patient has a diuretic efficiency of less than 50, such as less than 45, 40, 35, 30, or even less than 25, 20, 15, or 10.

[0115] In some embodiments, the patient resistant to diuretics requires a diuretic treatment of no less than 40 mg of furosemide (or equivalent) daily. In some of these embodiments, the patient requires a diuretic treatment of no less than 80 mg of furosemide (or equivalent) daily. In some of these embodiments, the patient requires a diuretic treatment of no less than 120 mg of furosemide (or equivalent) daily.

8.3.2. Pre-treatment IL-6 levels

8.3.2.1. Pre-treatment levels of IL-6 in urine

[0116] In certain embodiments, the patient has pre-treatment urine levels of IL-6 of more than 5.0 pg IL-6/g creatinine, 6.0 pg IL-6/g creatinine, 7.0 pg IL-6/g creatinine, 8.0 pg IL-6/g creatinine, 9.0 pg IL-6/g creatinine, or 10.0 pg IL-6/g creatinine. In certain embodiments, the patient has pre-treatment IL-6 urine levels of more than 11.0 pg IL-6/g creatinine, 12.0 pg IL-6/g creatinine, 13.0 pg IL-6/g creatinine, 14.0 pg IL-6/g creatinine, or 15.0 pg IL-6/g creatinine. In further embodiments, the patient has pre-treatment IL-6 levels in urine of more than 16.0 pg IL-6/g creatinine, 17.0 pg IL-6/g creatinine, 18.0 pg IL-6/g creatinine, 19.0 pg IL-6/g creatinine, or 20.0 pg IL-6/g creatinine. In particular embodiments, the patient has pre-treatment IL-6 levels in urine of more than 21.0 pg IL-6/g creatinine, 22.0 pg IL-6/g creatinine, 23.0 pg IL-6/g creatinine, 24.0 pg IL-6/g creatinine, 25.0 pg IL-6/g creatinine, 26.0 pg IL-6/g creatinine, 27.0 pg IL-6/g creatinine, 28.0 pg IL-6/g creatinine, 29.0 pg IL-6/g creatinine, or even more than 30.0 pg IL-6/g creatinine. In certain embodiments, the patient has pre-treatment urine IL-6 levels of more than 35.0 pg/g creatinine.

[0117] In certain embodiments, the patient has pre-treatment urine levels of more than 14.2 pg IL-6/g creatinine (“elevated urine IL-6 levels”). In other embodiments, the patient has pre-treatment urine levels of less than 14.2 pg IL-6/g creatinine.

[0118] In certain embodiments, the patient has levels of IL-6 in urine prior to treatment with an IL-6 antagonist and prior to treatment with a loop diuretic of more than 5.0 pg IL-6/g creatinine, 6.0 pg IL-6/g creatinine, 7.0 pg IL-6/g creatinine, 8.0 pg IL-6/g creatinine, 9.0 pg IL-6/g creatinine, or 10.0 pg IL-6/g creatinine. In some of these embodiments, the patient has IL-6 urine levels of more than 11.0 pg IL-6/g creatinine, 12.0 pg IL-6/g creatinine, 13.0 pg IL-6/g creatinine, 14.0 pg IL-6/g creatinine, or 15.0 pg IL-6/g creatinine. In further embodiments, the patient has levels of IL-6 in urine prior to treatment with an IL-antagonist and prior to treatment with a loop diuretic of more than 16.0 pg IL-6/g creatinine, 17.0 pg IL-6/g creatinine, 18.0 pg IL-6/g creatinine, 19.0 pg IL-6/g creatinine, or 20.0 pg IL-6/g creatinine. In particular embodiments, the patient has IL-6 levels in urine of more than 21.0 pg IL-6/g creatinine, 22.0 pg IL-6/g creatinine, 23.0 pg IL-6/g creatinine, 24.0 pg IL-6/g creatinine, 25.0 pg IL-6/g creatinine, 26.0 pg IL-6/g creatinine, 27.0 pg IL-6/g creatinine, 28.0 pg IL-6/g creatinine, 29.0 pg IL-6/g creatinine, or even more than 30.0 pg IL-6/g creatinine. In certain embodiments, the patient has urine IL-6 levels prior to treatment with an IL-antagonist and prior to treatment with a loop diuretic of more than 35.0 pg/g creatinine.

[0119] In certain embodiments, the patient has levels of IL-6 in urine prior to treatment with an IL-6 antagonist and prior to treatment with a loop diuretic of more than 14.2 pg IL-6/g creatinine. In other embodiments, the patient has levels of IL-6 in urine prior to treatment with an IL-antagonist and prior to treatment with a loop diuretic of less than 14.2 pg IL-/g creatinine.

8.3.2.2. Pre-treatment levels of IL-6 in plasma

[0120] In various embodiments, the patient has elevated pre-treatment plasma IL-6 levels.

[0121] In certain embodiments, the patient has pre-treatment plasma levels of IL-6 of more than 2.0 pg/mL. In other embodiments, the patient has pre-treatment plasma levels of IL-6 of less than 2.0 pg/mL.

[0122] In certain embodiments, the patient has pre-treatment plasma levels of IL-6 of more than 1.0 pg/ml, 1.1 pg/ml, 1.2 pg/ml, 1.3 pg/ml, 1.4 pg/ml, 1.5 pg/ml, 1.6 pg/ml, 1.7 pg/ml, 1.8 pg/ml, 1.9 pg/ml, or 2.0 pg/ml. In certain embodiments, the patient has pre-treatment plasma levels of IL-6 of more than 2.1 pg/ml, 2.2 pg/ml, 2.3 pg/ml, 2.4 pg/ml, 2.5 pg/ml, 2.6 pg/ml, 2.7 pg/ml, 2.8 pg/ml, 2.9 pg/ml, or 3.0 pg/ml. In certain embodiments, the patient has pre-treatment plasma levels of IL-6 of more than 3.1 pg/ml, 3.2 pg/ml, 3.3 pg/ml, 3.4 pg/ml, 3.5 pg/ml, 3.6 pg/ml, 3.7 pg/ml, 3.8 pg/ml, or 3.9 pg/ml.

[0123] In some embodiments, the patient has a pre-treatment IL-6 level of greater than 2 pg/mL, such as greater than 3 pg/mL, 4 pg/mL, 5 pg/mL, 6 pg/mL, 8 pg/mL, 10 pg/mL, 15 pg/mL, or 20 pg/mL. In certain embodiments, the patient has a pre-treatment IL-6 level of greater than 3 pg/mL. In certain embodiments, the patient has a pre-treatment IL-6 level of greater than 5 pg/mL. In certain embodiments, the patient has a pre-treatment IL-6 level of greater than 10 pg/mL.

8.3.2.3. Measurement of pre-treatment IL-6 levels

[0124] Concentrations of IL-6 in urine, plasma, and serum can be determined using any standard assay known in the art. When IL-6 is measured in urine, the level may be indexed or normalized to another biomarker, in certain embodiments urinary creatinine.

[0125] In particular embodiments, concentrations are measured using the MesoScale Discovery (MSD) platform (Meso Scale diagnostics, Gaithersburg, MD, USA).

8.3.3. Heart failure

[0126] In typical embodiments of the methods described herein, the patient has heart failure.

[0127] In certain embodiments, the patient has NYHA functional class I heart failure. In certain embodiments, the patient has NYHA functional class II heart failure. In certain embodiments, the patient has NYHA functional class III heart failure. In certain embodiments, the patient has NYHA functional class IV heart failure.

[0128] In certain embodiments, the patient has acute heart failure. In certain embodiments, the patient has chronic heart failure.

[0129] In certain embodiments, the patient has a type of heart failure selected from Table 1 below.

Table 1	
ICD-10-CM	Description
I50	Heart failure
	Heart failure complicating abortion or ectopic or molar pregnancy, heart failure following surgery, heart failure due to hypertension, heart failure due to hypertension with chronic kidney disease, obstetric surgery and procedures, rheumatic heart failure
I50.9	Heart failure, unspecified

Table 1	
ICD-10-CM	Description
	Biventricular (heart) failure NOS
	Cardiac, heart or myocardial failure NOS
	Congestive heart disease
	Congestive heart failure
	Right ventricular failure (secondary to left heart failure)
I50.1	Left ventricular failure
	Cardiac asthma
	Edema of lung with heart disease NOS
	Edema of lung with heart failure
	Left heart failure
	Pulmonary edema with heart disease NOS
	Pulmonary edema with heart failure
I50.20	Unspecified systolic (congestive) heart failure
I50.21	Acute systolic (congestive) heart failure
I50.22	Chronic systolic (congestive) heart failure
I50.23	Acute on chronic systolic (congestive) heart failure
I50.30	Unspecified diastolic (congestive) heart failure
I50.31	Acute diastolic (congestive) heart failure
I50.32	Chronic diastolic (congestive) heart failure
I50.33	Acute on chronic diastolic (congestive) heart failure
I50.40	Unspecified combined systolic (congestive) and diastolic (congestive) heart failure
I50.41	Acute combined systolic (congestive) and diastolic (congestive) heart failure
I50.42	Chronic combined systolic (congestive) and diastolic (congestive) heart failure
I50.43	Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure
I50.1	Left ventricular failure
	Heart failure, unspecified
	Biventricular (heart) failure NOS

Table 1	
ICD-10-CM	Description
	Cardiac, heart or myocardial failure NOS
	Congestive heart disease
	Congestive heart failure
	Right ventricular failure (secondary to left heart failure)

[0130] In certain embodiments, the patient has cardiorenal syndrome. In particular embodiments, the patient has cardiorenal syndrome type 1. In particular embodiments, the patient has cardiorenal syndrome type 2. In particular embodiments, the patient has cardiorenal syndrome type 3. In particular embodiments, the patient has cardiorenal syndrome type 4.

[0131] In certain embodiments, the patient has diuretic resistant heart failure. In certain of these embodiments, the heart failure patient has a diuretic efficiency of less than 100, 95, 90, 85, 80, 75, 70, 65, 60, 55, or 50. In particular embodiments, the patient has a diuretic efficiency of less than 45, 40, 35, 30, or even less than 25, 20, 15, or 10.

8.3.4. Kidney disease

[0132] In some embodiments of methods described herein, the patient has kidney disease.

[0133] In certain embodiments, the patient has diuretic resistant kidney disease. In certain of these embodiments, the kidney disease patient has a diuretic efficiency of less than 100, 95, 90, 85, 80, 75, 70, 65, 60, 55, or 50. In particular embodiments, the patient has a diuretic efficiency of less than 45, 40, 35, 30, or even less than 25, 20, 15, or 10.

[0134] In particular embodiments, the patient has hepatorenal syndrome.

8.3.5. TMPRSS6 rs855791 genotype

[0135] In certain embodiments, the patient has previously been determined to have at least one copy of the TMPRSS6 rs855791 major allele. In other embodiments, the method further comprises the earlier step of determining that the patient has at least one copy of the TMPRSS6 rs855791 major allele.

[0136] Preferably, the genotype at both alleles is determined, thus permitting identification and discrimination of patients who are homozygous for the *TMPRSS6* rs855791 major allele, heterozygous for the major and minor *TMPRSS6* rs855791 alleles, and homozygous for the *TMPRSS6* rs855791 minor allele.

[0137] The absence (major allele) or presence (minor allele) of SNP rs855791 (2321G→A) in the *TMPRSS6* gene is determined using standard techniques.

[0138] Typically, PCR is used to amplify a biological sample obtained from the patient.

[0139] In certain embodiments, the absence or presence of polymorphism is detected concurrently with amplification using real-time PCR (RT-PCR). In certain embodiments, the RT-PCR assay employs 5' nuclease (TaqMan[®] probes), molecular beacons, and/or FRET hybridization probes. *Reviewed in Espy et al., Clin. Microbiol. Rev.* 2006 January; 19(1): 165-256, incorporated herein by reference in its entirety. In typical embodiments, a commercially available assay is used. In select embodiments, the commercially available assay is selected from the group consisting of TaqMan[™] SNP Genotyping Assays (ThermoFisher); PCR SNP Genotyping Assay (Qiagen); Novallele Genotyping Assays (Canon); and SNP Type[™] assays (formerly SNPtype) (Fluidigm).

[0140] In certain embodiments, the absence or presence of polymorphism is detected following amplification using hybridization with a probe specific for SNP rs855791, restriction endonuclease digestion, nucleic acid sequencing, primer extension, microarray or gene chip analysis, mass spectrometry, and/or a DNase protection assay. In certain embodiments, the allelic variants are called by sequencing. In certain embodiments, Sanger sequencing is used. In certain embodiments, one of a variety of next-generation sequencing techniques is used, including for example a sequencing technique selected from the group consisting of microarray sequencing, Solexa sequencing (Illumina), Ion Torrent (Life Technologies), SOLiD (Applied Biosystems), pyrosequencing, single-molecule real-time sequencing (Pacific Bio), nanopore sequencing and tunneling currents sequencing.

[0141] In certain embodiments, the absence or presence of polymorphisms is detected using the procedures set forth in Example 2 below.

8.3.6. IL-6 antagonists

[0142] The IL-6 antagonist used in the methods described herein is capable of decreasing the biological activity of IL-6.

8.3.6.1. Anti-IL-6 antibodies

[0143] In certain embodiments, the IL-6 antagonist is an anti-IL-6 antibody or antigen-binding fragment or derivative thereof.

[0144] In certain embodiments, the IL-6 antagonist is a full-length anti-IL-6 monoclonal antibody. In particular embodiments, the full-length monoclonal antibody is an IgG antibody. In certain embodiments, the full-length monoclonal antibody is an IgG1, IgG2, IgG3, or IgG4 antibody. In certain embodiments, the IL-6 antagonist is a polyclonal composition comprising a plurality of species of full-length anti-IL-6 antibodies, each of the plurality having unique CDRs. In certain embodiments, the IL-6 antagonist is an antibody fragment selected from Fab, Fab', and F(ab')₂ fragments. In certain embodiments, the IL-6 antagonist is a scFv, a disulfide-linked Fv (dsFv), or a single domain antibody, such as a camelid-derived VHH single domain Nanobody. In certain embodiments, the IL-6 antagonist is immunoconjugate or fusion comprising an IL-6 antigen-binding fragment. In certain embodiments, the antibody is bispecific or multispecific, with at least one of the antigen-binding portions having specificity for IL-6.

[0145] In certain embodiments, the antibody is fully human. In certain embodiments, the antibody is humanized. In certain embodiments, the antibody is chimeric and has non-human V regions and human C region domains. In certain embodiments, the antibody is murine.

[0146] In typical embodiments, the anti-IL-6 antibody has a K_D for binding human IL-6 of less than 100 nM. In certain embodiments, the anti-IL-6 antibody has a K_D for binding human IL-6 of less than 75 nM, 50 nM, 25 nM, 20 nM, 15 nM, or 10 nM. In particular embodiments, the anti-IL-6 antibody has a K_D for binding human IL-6 of less than 5 nM, 4 nM, 3 nM, or 2 nM. In selected embodiments, the anti-IL-6 antibody has a K_D for binding human IL-6 of less than 1 nM, 750 pM, or 500 pM. In specific embodiments, the anti-IL-6 antibody has a K_D for binding human IL-6 of no more than 500 pM, 400 pM, 300 pM, 200 pM, or 100 pM.

[0147] In typical embodiments, the anti-IL-6 antibody neutralizes the biological activity of IL-6. In certain embodiments, the neutralizing antibody prevents binding of IL-6 to the IL-6 receptor.

[0148] In typical embodiments, the anti-IL-6 antibody has an elimination half-life following intravenous administration of at least 7 days. In certain embodiments, the anti-IL-6 antibody has an elimination half-life of at least 14 days, at least 21 days, or at least 30 days.

[0149] In certain embodiments, the anti-IL-6 antibody has a human IgG constant region with at least one amino acid substitution that extends serum half-life as compared to the unsubstituted human IgG constant domain.

[0150] In certain embodiments, the IgG constant domain comprises substitutions at residues 252, 254, and 256, wherein the amino acid substitution at amino acid residue 252 is a substitution with tyrosine, the amino acid substitution at amino acid residue 254 is a substitution with threonine, and the amino acid substitution at amino acid residue 256 is a substitution with glutamic acid ("YTE"). See U.S. Pat. No. 7,083,784, incorporated herein by reference in its entirety. In certain extended half-life embodiments, the IgG constant domain comprises substitutions selected from T250Q/M428L (Hinton *et al.*, *J. Immunology* 176:346-356 (2006)); N434A (Yeung *et al.*, *J. Immunology* 182:7663-7671 (2009)); or T307A/E380A/N434A (Petkova *et al.*, *International Immunology*, 18: 1759-1769 (2006)).

[0151] In certain embodiments, the elimination half-life of the anti-IL-6 antibody is increased by utilizing the FcRN-binding properties of human serum albumin. In certain embodiments, the antibody is conjugated to albumin (Smith *et al.*, *Bioconjug. Chem.*, 12: 750-756 (2001)). In certain embodiments, the anti-IL-6 antibody is fused to bacterial albumin-binding domains (Stork *et al.*, *Prot. Eng. Design Science* 20: 569-76 (2007)). In certain embodiments, the anti-IL-6 antibody is fused to an albumin-binding peptide (Nguyen *et al.*, *Prot Eng Design Sel* 19: 291-297 (2006)). In certain embodiments, the anti-IL-antibody is bispecific, with one specificity being to IL-6, and one specificity being to human serum albumin (Ablynx, WO 2006/122825 (bispecific Nanobody)).

[0152] In certain embodiments, the elimination half-life of the anti-IL-6 antibody is increased by PEGylation (Melmed *et al.*, *Nature Reviews Drug Discovery* 7: 641-642 (2008)); by HPMA copolymer conjugation (Lu *et al.*, *Nature Biotechnology* 17: 1101-1104 (1999)); by dextran conjugation (*Nuclear Medicine Communications*, 16: 362-369 (1995)); by conjugation with homo-amino-acid polymers (HAPs; HAPylation) (Schlapschy *et al.*, *Prot Eng Design Sel* 20: 273-284 (2007)); or by polysialylation (Constantinou *et al.*, *Bioconjug. Chem.* 20: 924-931 (2009)).

8.3.6.1.1. MEDI5117 and derivatives

[0153] In certain embodiments, the anti-IL-6 antibody or antigen-binding portion thereof comprises all six CDRs of MEDI5117. In particular embodiments, the antibody or antigen-binding portion thereof comprises the MEDI5117 heavy chain V region and light chain V

region. In specific embodiments, the antibody is the full-length MEDI5117 antibody. The MEDI5117 antibody is described in WO 2010/088444 and US 2012/0034212, the disclosures of which are incorporated herein by reference in their entireties. The MEDI5117 antibody has the following CDR and heavy and light chain sequences:

MEDI5117 VH CDR1

SNYMI (SEQ ID NO:9)

MEDI5117 VH CDR2

DLYYYAGDITYYADSVKG (SEQ ID NO:10)

MEDI5117 VH CDR3

WADDHPPWIDL (SEQ ID NO:11)

MEDI5117 VL CDR1

RASQGISSWLA (SEQ ID NO:12)

MEDI5117 VL CDR2

KASTLES (SEQ ID NO:13)

MEDI5117 VL CDR3

QQSWLGGS (SEQ ID NO:14)

MEDI5117 Heavy chain

EVQLVESGGGLVQPGGSLRLSCAASGFTISSNYMIWVRQAPGKGLEWVSDLYYYAGDITYY
ADSVKGRFTMSRDISKNTVYLQMNSLR AEDTAVYYCARWADDHPPWIDLWGRGTLTVSS
ASTKGPSVFPLAPSSKSTSGGTAA LGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS
GLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRV EPKSCDKTHTCPPCPAPELLGG
PSVFLFPPKPKDTLYITREPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN
STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREE
MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO:15)

MEDI5117 Light chain

DIQMTQSPSTLSASVGDRTITCRASQGISSWLAWYQQKPGKAPKVLIIYKASTLESQVPS
RFGSGSGTEFTLTISSLQPD FATYYCQQSWLGGSFGQGTKLEIKRTVAAPSVFIFPPS
DEQLKSGTASVVCCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTTL
SKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:16)

[0154] In certain embodiments, the anti-IL-6 antibody is a derivative of MEDI5117.

[0155] In certain embodiments, the MED5117 derivative includes one or more amino acid substitutions in the MED5117 heavy and/or light chain V regions.

[0156] In certain embodiments, the derivative comprises fewer than 25 amino acid substitutions, fewer than 20 amino acid substitutions, fewer than 15 amino acid substitutions, fewer than 10 amino acid substitutions, fewer than 5 amino acid substitutions, fewer than 4 amino acid substitutions, fewer than 3 amino acid substitutions, fewer than 2 amino acid substitutions, or 1 amino acid substitution relative to the original V_H and/or V_L of the MEDI5117 anti-IL-6 antibody, while retaining specificity for human IL-6.

[0157] In certain embodiments, the MED5117 derivative comprises an amino acid sequence that is at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of the V_H and V_L domain of MEDI5117. The percent sequence identity is determined using BLAST algorithms using default parameters.

[0158] In certain embodiments, the MED5117 derivative comprises an amino acid sequence in which the CDRs comprise an amino acid sequence that is at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of the respective CDRs of MEDI5117. The percent sequence identity is determined using BLAST algorithms using default parameters.

[0159] In certain embodiments, the V_H and/or V_L CDR derivatives comprise conservative amino acid substitutions at one or more predicted nonessential amino acid residues (i.e., amino acid residues which are not critical for the antibody to specifically bind to human IL-6).

8.3.6.1.2. Other anti-IL-6 antibodies

[0160] In certain embodiments, the anti-IL-6 antibody comprises the six CDRs from an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, elsilimomab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb). In certain embodiments, the anti-IL-6 antibody comprises the heavy chain V region and light chain V region from an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029;

Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb). In particular embodiments, the anti-IL-6 antibody is an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb).

[0161] In certain embodiments, the anti-IL-6 antibody comprises the six CDRs from an antibody selected from those described in US 2016/0168243, US 2016/0130340, US 2015/0337036, US 2015/0203574, US 2015/0140011, US 2015/0125468, US 2014/0302058, US 2014/0141013, US 2013/0280266, US 2013/0017575, US 2010/0215654, US 2008/0075726, US Pat. No. 5,856,135, US 2006/0240012, US 2006/0257407, or U.S. Pat. No. 7291721, the disclosures of which are incorporated herein by reference in their entireties.

8.3.6.2. Anti-IL-6 receptor antibodies

[0162] In certain embodiments, the IL-6 antagonist is an anti-IL-6 receptor antibody or antigen-binding fragment or derivative thereof.

[0163] In certain embodiments, the IL-6 antagonist is a full-length anti-IL-6 receptor monoclonal antibody. In particular embodiments, the full-length monoclonal antibody is an IgG antibody. In certain embodiments, the full-length monoclonal antibody is an IgG1, IgG2, IgG3, or IgG4 antibody. In certain embodiments, the IL-6 antagonist is a polyclonal composition comprising a plurality of species of full-length anti-IL-6 receptor antibodies, each of the plurality having unique CDRs. In certain embodiments, the IL-6 antagonist is an antibody fragment selected from Fab and Fab' fragments. In certain embodiments, the IL-6 antagonist is a scFv, a single domain antibody, including a camelid-derived VHH single domain Nanobody. In certain embodiments, the antibody is bispecific or multispecific, with at least one of the antigen-binding portions having specificity for IL-6R.

[0164] In certain embodiments, the antibody is fully human. In certain embodiments, the antibody is humanized. In certain embodiments, the antibody is chimeric and has non-human V regions and human C region domains. In certain embodiments, the antibody is murine.

[0165] In typical embodiments, the anti-IL-6 receptor antibody has a K_D for binding human IL-6R of less than 100 nM. In certain embodiments, the anti-IL-6R antibody has a K_D for

binding human IL-6R of less than 75 nM, 50 nM, 25 nM, 20 nM, 15 nM, or 10 nM. In particular embodiments, the anti-IL-6 receptor antibody has a K_D for binding human IL-6R of less than 5 nM, 4 nM, 3 nM, or 2 nM. In selected embodiments, the anti-IL-6 receptor antibody has a K_D for binding human IL-6R of less than 1 nM, 750 pM, or 500 pM. In specific embodiments, the anti-IL-6 receptor antibody has a K_D for binding human IL-6R of no more than 500 pM, 400 pM, 300 pM, 200 pM, or 100 pM.

[0166] In typical embodiments, the anti-IL-6R reduces the biological activity of IL-6.

[0167] In typical embodiments, the anti-IL-6R antibody has an elimination half-life following intravenous administration of at least 7 days. In certain embodiments, the anti-IL-6R antibody has an elimination half-life of at least 14 days, at least 21 days, or at least 30 days.

[0168] In certain embodiments, the anti-IL-6R antibody has a human IgG constant region with at least one amino acid substitution that extends serum half-life as compared to the unsubstituted human IgG constant domain.

[0169] In certain embodiments, the IgG constant domain comprises substitutions at residues 252, 254, and 256, wherein the amino acid substitution at amino acid residue 252 is a substitution with tyrosine, the amino acid substitution at amino acid residue 254 is a substitution with threonine, and the amino acid substitution at amino acid residue 256 is a substitution with glutamic acid ("YTE"). See U.S. Pat. No. 7,083,784, incorporated herein by reference in its entirety. In certain extended half-life embodiments, the IgG constant domain comprises substitutions selected from T250Q/M428L (Hinton *et al.*, J. Immunology 176:346-356 (2006)); N434A (Yeung *et al.*, J. Immunology 182:7663-7671 (2009)); or T307A/E380A/N434A (Petkova *et al.*, International Immunology, 18: 1759-1769 (2006)).

[0170] In certain embodiments, the elimination half-life of the anti-IL-6R antibody is increased by utilizing the FcRN-binding properties of human serum albumin. In certain embodiments, the antibody is conjugated to albumin (Smith *et al.*, Bioconjug. Chem., 12: 750-756 (2001)). In certain embodiments, the anti-IL-6R antibody is fused to bacterial albumin-binding domains (Stork *et al.*, Prot. Eng. Design Science 20: 569-76 (2007)). In certain embodiments, the anti-IL-6 antibody is fused to an albumin-binding peptide (Nguyen *et al.*, Prot Eng Design Sel 19: 291-297 (2006)). In certain embodiments, the anti-IL-6 antibody is bispecific, with one specificity being to IL-6R, and one specificity being to human serum albumin (Ablynx, WO 2006/122825 (bispecific Nanobody)).

[0171] In certain embodiments, the elimination half-life of the anti-IL-6R antibody is increased by PEGylation (Melmed *et al.*, Nature Reviews Drug Discovery 7: 641-642 (2008)); by HPMa copolymer conjugation (Lu *et al.*, Nature Biotechnology 17: 1101-1104 (1999)); by dextran conjugation (Nuclear Medicine Communications, 16: 362-369 (1995)); by conjugation with homo-amino-acid polymers (HAPs; HAPylation) (Schlappschy *et al.*, Prot Eng Design Sel 20: 273-284 (2007)); or by polysialylation (Constantinou *et al.*, Bioconjug. Chem. 20: 924-931 (2009)).

[0172] In certain embodiments, the anti-IL-6R antibody or antigen-binding portion thereof comprises all six CDRs of tocilizumab. In particular embodiments, the antibody or antigen-binding portion thereof comprises the tocilizumab heavy chain V region and light chain V region. In specific embodiments, the antibody is the full-length tocilizumab antibody.

[0173] In certain embodiments, the anti-IL-6R antibody or antigen-binding portion thereof comprises all six CDRs of sarilumab. In particular embodiments, the antibody or antigen-binding portion thereof comprises the sarilumab heavy chain V region and light chain V region. In specific embodiments, the antibody is the full-length sarilumab antibody.

[0174] In certain embodiments, the anti-IL-6R antibody or antigen-binding portion thereof comprises all six CDRs of VX30 (Vaccinex), ARGX-109 (arGEN-X), FM101 (Formatech), SA237 (Roche), NI-1201 (NovImmune), or an antibody described in US 2012/0225060.

[0175] In certain embodiments, the anti-IL-6R antibody or antigen-binding portion thereof is a single domain antibody. In particular embodiments, the single domain antibody is a camelid VHH single domain antibody. In specific embodiments, the antibody is vobarilizumab (ALX-0061) (Ablynx NV).

8.3.6.3. Anti-IL-6:IL-6R complex antibodies

[0176] In certain embodiments, the IL-6 antagonist is an antibody specific for the complex of IL-6 and IL-6R. In certain embodiments, the antibody has the six CDRs of an antibody selected from those described in US 2011/0002936, which is incorporated herein by reference in its entirety.

8.3.6.4. JAK and STAT inhibitors

[0177] IL-6 is known to signal via the JAK-STAT pathway.

[0178] In certain embodiments, the IL-6 antagonist is an inhibitor of the JAK signaling pathway. In certain embodiments, the JAK inhibitor is a JAK1-specific inhibitor. In certain embodiments, the JAK inhibitor is a JAK3-specific inhibitor. In certain embodiments, the JAK inhibitor is a pan-JAK inhibitor.

[0179] In certain embodiments, the JAK inhibitor is selected from the group consisting of tofacitinib (Xeljanz), decernotinib, ruxolitinib, upadacitinib, baricitinib, filgotinib, lestaurtinib, pacritinib, peficitinib, INCB-039110, ABT-494, INCB-047986 and AC-410.

[0180] In certain embodiments, the IL-6 antagonist is a STAT3 inhibitor. In a specific embodiment, the inhibitor is AZD9150 (AstraZeneca, Isis Pharmaceuticals), a STAT3 antisense molecule.

8.3.6.5. Additional IL-6 antagonists

[0181] In certain embodiments, the IL-6 antagonist is an antagonist peptide.

[0182] In certain embodiments, the IL-6 antagonist is C326 (an IL-6 antagonist by Avidia, also known as AMG220), or FE301, a recombinant protein inhibitor of IL-6 (Ferring International Center S.A., Conaris Research Institute AG). In certain embodiments, the anti-IL-6 antagonist comprises soluble gp130, FE301 (Conaris/Ferring).

8.3.7. Dosage regimens

8.3.7.1. Antibodies, antigen-binding fragments, peptides

[0183] In typical embodiments, antibody, antigen-binding fragments, and peptide IL-6 antagonists are administered parenterally.

[0184] In some parenteral embodiments, the IL-6 antagonist is administered intravenously. In certain intravenous embodiments, the IL-6 antagonist is administered as a bolus. In certain intravenous embodiments, the IL-6 antagonist is administered as an infusion. In certain intravenous embodiments, the IL-6 antagonist is administered as a bolus followed by infusion. In some parenteral embodiments, the IL-6 antagonist is administered subcutaneously.

[0185] In certain embodiments, the antibody, antigen-binding fragment, or peptide IL-6 antagonist is administered in a dose that is independent of patient weight or surface area (flat dose).

[0186] In certain embodiments, the intravenous flat dose is 1 mg, 2 mg, 3 mg, 4 mg, 5 mg, 6 mg, 7 mg, 8 mg, 9 mg, or 10 mg. In certain embodiments, the intravenous flat dose is 11 mg, 12 mg, 13 mg, 14 mg, 15 mg, 16 mg, 17 mg, 18 mg, 19 mg, or 20 mg. In certain embodiments, the intravenous flat dose is 25 mg, 30 mg, 40 mg, or 50 mg. In certain embodiments, the intravenous flat dose is 60 mg, 70 mg, 80 mg, 90 mg, or 100 mg. In certain embodiments, the intravenous flat dose is 1 – 10 mg, 10 – 15 mg, 15 – 20 mg, 20 – 30 mg, 30 – 40 mg, or 40 – 50 mg. In certain embodiments, the intravenous flat dose is 1 – 40 mg, or 50 – 100 mg.

[0187] In certain embodiments, the subcutaneous flat dose is 10 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 80 mg, 90 mg, or 100 mg. In certain embodiments, the subcutaneous flat dose is 110 mg, 120 mg, 130 mg, 140 mg, 150 mg, 160 mg, 170 mg, 180 mg, 190 mg, or 200 mg. In certain embodiments, the subcutaneous flat dose is 210 mg, 220 mg, 230 mg, 240 mg, or 250 mg. In certain embodiments, the subcutaneous flat dose is 10 – 100 mg, 100 – 200 mg, or 200 – 250 mg. In certain embodiments, the subcutaneous flat dose is 10 – 20 mg, 20 – 30 mg, 30 – 40 mg, 40 – 50 mg, 50 – 60 mg, 60 – 70 mg, 70 – 80 mg, 80 – 90 mg, or 90 – 100 mg. In certain embodiments, the subcutaneous flat dose is 100 – 125 mg, 125 – 150 mg, 150 – 175 mg, 175 – 200 mg, or 200 – 250 mg.

[0188] In certain embodiments, the antibody, antigen-binding fragment, or peptide IL-6 antagonist is administered as a patient weight-based dose.

[0189] In certain embodiments, the antagonist is administered at an intravenous dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg or 1.0 mg/kg. In certain embodiments, the antagonist is administered at a dose of 1.5 mg/kg, 2 mg/kg, 2.5 mg/kg, 3 mg/kg, 3.5 mg/kg, 4 mg/kg, 4.5 mg/kg, or 5 mg/kg.

[0190] In certain embodiments, the subcutaneous weight-based dose is 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg or 1.0 mg/kg. In certain embodiments, the antagonist is administered at a dose of 1.5 mg/kg, 2 mg/kg, 2.5 mg/kg, 3 mg/kg, 3.5 mg/kg, 4 mg/kg, 4.5 mg/kg, or 5 mg/kg.

[0191] In various intravenous embodiments, the IL-6 antagonist is administered once every 7 days, once every 14 days, once every 21 days, once every 28 days, or once a month. In various subcutaneous embodiments, the IL-6 antagonist is administered once every 14 days, once every 28 days, once a month, once every two months (every other month), or once every three months.

[0192] In certain preferred embodiments, the IL-6 antagonist is the MEDI5117 antibody. In certain embodiments, MEDI5117 is administered in a flat dose of 1 – 30 mg IV once every week. In certain embodiments, the MEDI5117 antibody is administered in a flat dose of 1, 2, 3, 4, 5, 7.5, 10, 15, 20, 25, or 30 mg IV once every week. In certain embodiments, the MEDI5117 antibody is administered in a flat dose of 25 – 250 mg s.c. once every month to once every three months. In particular embodiments, MEDI5117 is administered at a dose of 30 mg, 45 mg, 60 mg, 75 mg, 100 mg, 120 mg, 125 mg, 150 mg, 175 mg, 200 mg, 225 mg, 240 mg, or 250 mg s.c. once every month, once every two months, or once every 3 months.

[0193] In certain embodiments, the IL-6 antagonist is tocilizumab. In certain embodiments, tocilizumab is administered s.c. in a starting dose for patients ≥ 100 kg of 162 mg once every week. In certain embodiments, tocilizumab is administered intravenously at a dose of 4 mg/kg once every 4 weeks followed by an increase to 8 mg/kg every 4 weeks based on clinical response.

8.3.7.2. JAK and STAT inhibitors

[0194] In typical embodiments, small molecule JAK inhibitors and STAT inhibitors are administered orally.

[0195] In certain embodiments, the inhibitor is administered once or twice a day at an oral dose of 1 – 10 mg, 10 – 20 mg, 20 – 30 mg, 30 – 40 mg, or 40 – 50 mg. In certain embodiments, the inhibitor is administered once or twice a day at a dose of 50 – 60 mg, 60 – 70 mg, 70 – 80 mg, 80 – 90 mg, or 90 – 100 mg. In certain embodiments, the inhibitor is administered at a dose of 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mg PO once or twice a day. In certain embodiments, the inhibitor is administered at a dose of 75 mg PO QD or BID, 100 mg PO QD or BID.

[0196] In certain embodiments, the JAK inhibitor is tofacitinib, and is administered at a dose of 5 mg PO BID or 11 mg PO qDay,

[0197] In certain embodiments, the JAK inhibitor is decernotinib, and is administered at a dose of 25 mg, 50 mg, 100 mg, or 150 mg PO BID.

[0198] In certain embodiments, the inhibitor is ruxolitinib, and is administered at dose of 25 mg PO BID, 20 mg PO BID, 15 mg PO BID, 10 mg PO BID, or 5 mg PO BID.

8.3.7.3. Treating to goal

[0199] In certain embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase diuretic efficiency. In certain of these embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase diuretic efficiency to normal levels.

[0200] In certain embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase eGFR, and in certain of these embodiments, the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase eGFR to normal levels.

8.3.7.4. IL-6 level monitoring

[0201] In certain embodiments, the impact of treatment with IL-6 antagonists on cardiorenal parameters may be monitored by measuring a level of IL-6 in a urine or plasma sample from the patient. It is specifically contemplated that the methods of the invention may be used to monitor the efficacy of treatments for cardiorenal syndrome and may motivate, for example, a change of dose or therapeutic.

[0202] Accordingly, in certain embodiments, the method further comprises the subsequent step of determining the level of IL-6 in urine, in plasma, or in both urine and plasma. In certain of these embodiments, the method further comprises a final step of adjusting the dose of at least one subsequent administration of IL-6 antagonist based on IL-6 level determined in the immediately preceding step.

8.3.8. Additional therapeutic agents

[0203] In certain embodiments of the methods described herein, the method further comprises administering at least one therapeutic agent additional to the IL-6 antagonist, wherein the additional therapeutic agent treats one or more cardiovascular or renal symptoms of heart failure. Specific treatment will be determined on a case by case basis by the attending physician.

8.3.8.1. Standard heart failure agents

[0204] In certain embodiments, the additional therapeutic agent is a diuretic.

[0205] In particular embodiments, the diuretic is a loop diuretic. In select embodiments, the loop diuretic is selected from the group consisting of furosemide, torsemide, bumetanide, and ethacrynic acid. In particular embodiments, the loop diuretic is furosemide. In certain embodiments, furosemide is administered orally. In certain embodiments, furosemide is administered intravenously. In certain embodiments, the diuretic is a thiazide diuretic. In particular embodiments, the thiazide diuretic is chlorothiazide, hydrochlorothiazide, chlorthalidone, indapamide, or metolazone. In certain embodiments, the diuretic is a potassium sparing diuretic.

[0206] In certain embodiments, the additional therapeutic agent is an ACE inhibitor. In certain embodiments, the ACE inhibitor is selected from the group consisting of benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril andtrandolapril.

[0207] In certain embodiments, the additional therapeutic agent is an angiotensin receptor blocker (“ARB”). In certain embodiments, the ARB is eprosartan, olmesartan, valsartan, telmisartan, losartan, azilsartan medoxomil, candesartan, or irbesartan.

[0208] In certain embodiments, the additional therapeutic agent is a β -blocker, a calcium antagonist, or a mineralocorticoid receptor antagonist.

[0209] In certain embodiments, the additional therapeutic agent is a natriuretic peptide, such as a B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide.

[0210] In certain embodiments, the additional therapeutic agent is an adenosine antagonist, such as rolofylline.

[0211] Specific treatment will be determined on a case-by-case basis by the attending physician.

8.3.8.2. Nitroxyl donors

[0212] In certain embodiments, the additional therapeutic agent is a nitroxyl donor, and the method further comprises administering a therapeutically effective amount of the nitroxyl donor.

[0213] In particular embodiments, the nitroxyl donor is selected from the compounds described in one or more of U.S. Pat. Nos. 9,499,511; 9,487,498; 9,464,061; 9,458,127; 9,221,780; 9,181,213; 9,156,804; 9,115,064; 9,018,411; 8,987,326; RE45,314; 8,674,132;

8,227,639; and 8,030,356, the disclosures of which are incorporated herein by reference in their entireties.

[0214] In selected embodiments, the nitroxyl donor is selected from the compounds described in U.S. Pat. No. RE45,314. In specific embodiments, the nitroxyl donor is selected from the compounds described in U.S. Pat. No. 9,156,804.

8.3.8.3. Sodium-free chloride supplementation

[0215] In certain embodiments, the additional therapeutic agent is a sodium free chloride salt. In certain embodiments, the agent is lysine chloride.

8.4. Methods of improving treatment of heart failure

[0216] In another aspect, methods are provided for improving treatment of heart failure by discontinuing therapy that is ineffective, thereby reducing side effects and reducing cost without loss of treatment efficacy. The methods comprise discontinuing administration of an IL-6 antagonist to a patient with heart failure, wherein the patient has been determined to be homozygous for the *TMPRSS6* rs855791 minor allele. In one series of embodiments, the patient has previously been determined to be homozygous for the *TMPRSS6* rs855791 minor allele. In another series of embodiments, the method further comprises the earlier step of determining that the patient is homozygous for the *TMPRSS6* rs855791 minor allele.

[0217] In certain embodiments, the patient has elevated pre-treatment urine levels of IL-6. In certain embodiments, the patient has elevated pre-treatment plasma levels of IL-6. In certain embodiments, the patient has elevated pre-treatment urine and elevated pre-treatment plasma levels of IL-6.

[0218] In particular embodiments, the patient has cardiorenal syndrome.

8.5. Diagnostic, prognostic, and treatment guidance methods

[0219] In another aspect, methods are provided for determining if a subject would benefit from IL-6 antagonist treatment for heart failure. The method comprises measuring a level of IL-6 in a urine sample or plasma sample from the subject, comparing the measured level of IL-6 to a predetermined reference level, and determining whether or not the measured level of IL-6 is greater than the corresponding reference level, wherein when the measured level of

IL-6 is greater than the corresponding reference level, IL-6 antagonist treatment is recommended.

[0220] IL-6 may be measured in plasma or in urine according to the methods set forth in Section 6.3.2.3 above. Reference levels for IL-6 in urine and plasma may be determined by measuring IL-6 levels in a reference population. A person of skill in the art is able to determine a reference level for the level of a biomarker in a population based on clinical experience and common levels of the biomarker in samples from the population.

[0221] In further aspects, methods are provided for determining if a subject is in need of IL-6 antagonist treatment for impaired glomerular filtration, low diuretic efficiency, high urine angiotensin, high plasma renin, or is at risk of mortality due to cardiorenal syndrome using the above described method. As shown in FIG. 3 and discussed in Example 1 hereinbelow, high levels of IL-6 in urine or plasma are associated with these parameters.

8.6. Kits

[0222] In a further aspect, a kit is provided. In general, kits will comprise detection reagents that are suitable for detecting the presence of biomarkers of interest and with instructions for use in accordance with the methods of the invention. The kit may comprise antibodies or other immunohistochemical reagents capable of binding to IL-6. The kit may contain capture and detection antibodies suitable for performing an ELISA for measuring IL-6 in urine or plasma. In certain embodiments the kit may contain tools and reagents for preparing urine and plasma samples for the ELISA or for indexing IL-6 in urine to the concentration of another biomarker, in certain embodiments IL-6 is indexed against creatinine.

8.7. Further embodiments

[0223] Further embodiments are provided in the following numbered embodiments.

1. A method of treating heart failure, comprising:
administering a therapeutically effective amount of an IL-6 antagonist to a patient with heart failure,
wherein the patient has been determined to have at least one copy of the *TMPRSS6* rs855791 major allele.

2. The method of embodiment 1, wherein the patient has previously been determined to have at least one copy of the *TMPRSS6* rs855791 major allele.
3. The method of embodiment 1, further comprising the earlier step of:
determining that the patient has at least one copy of the *TMPRSS6* rs855791 major allele.
4. The method of any one of embodiments 1-3, wherein the patient has elevated pre-treatment urine levels of IL-6.
5. The method of any one of embodiments 1-4, wherein the patient has elevated pre-treatment plasma levels of IL-6.
6. The method of any one of embodiments 1-5, wherein the patient has acute heart failure.
7. The method of any one of embodiments 1-5, wherein the patient has chronic heart failure.
8. The method of any one of embodiments 1-7, wherein the patient has cardiorenal syndrome.
9. The method of embodiment 8, wherein the patient has cardiorenal syndrome type 4.
10. The method of any one of embodiments 1-9, wherein the patient has diuretic resistant heart failure.
11. The method of embodiment 10, wherein the patient's diuretic efficiency is less than 95.
12. The method of embodiment 11, wherein the patient's diuretic efficiency is less than 90.
13. The method of embodiment 12, wherein the patient's diuretic efficiency is less than 85.

14. The method of embodiment 13, wherein the patient's diuretic efficiency is less than 80.
15. The method of embodiment 14, wherein the patient's diuretic efficiency is less than 75.
16. The method of embodiment 15, wherein the patient's diuretic efficiency is less than 70.
17. The method of embodiment 16, wherein the patient's diuretic efficiency is less than 65.
18. The method of any one of embodiments 1-17, wherein the IL-6 antagonist is an anti-IL-6 antibody, or antigen-binding fragment or derivative thereof.
19. The method of embodiment 18, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative has a K_D for binding human IL-6 of less than 100 nM.
20. The method of embodiment 19, wherein the antibody or antigen-binding fragment or derivative has a K_D for binding human IL-6 of less than 50 nM.
21. The method of embodiment 20, wherein the antibody or antigen-binding fragment or derivative has a K_D for binding human IL-6 of less than 10 nM.
22. The method of embodiment 21, wherein the antibody or antigen-binding fragment or derivative has a K_D for binding human IL-6 of less than 1 nM.
23. The method of any one of embodiments 18-22, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative has an elimination half-life following intravenous administration of at least 7 days.

24. The method of embodiment 23, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative has an elimination half-life following intravenous administration of at least 14 days.
25. The method of embodiment 24, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative has an elimination half-life following intravenous administration of at least 21 days.
26. The method of embodiment 25, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative has an elimination half-life following intravenous administration of at least 30 days.
27. The method of any one of embodiments 18 - 26, wherein the IL-6 antagonist is a full-length monoclonal anti-IL-6 antibody.
28. The method of embodiment 27, wherein the antibody is an IgG1 or IgG4 antibody.
29. The method of embodiment 28, wherein the antibody is an IgG1 antibody.
30. The method of any one of embodiments 18-29, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative is fully human.
31. The method of any one of embodiments 18-29, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative is humanized.
32. The method of any one of embodiments 18- 31, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative comprises all six variable region CDRs of MED5117.
33. The method of embodiment 32, wherein the antibody comprises the VH and VL of MED5117.
34. The method of embodiment 33, wherein the antibody is MED5117.

35. The method of any one of embodiments 18– 31, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative comprises all six variable region CDRs of an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, elsilimomab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb).

36. The method of embodiment 35, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative comprises the heavy chain V region and light chain V region from an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb).

37. The method of embodiment 36, wherein the anti-IL-6 antibody or antigen-binding fragment or derivative is an antibody selected from the group consisting of siltuximab, gerilimumab, sirukumab, clazakizumab, olokizumab, VX30 (VOP-R003; Vaccinex), EB-007 (EBI-029; Eleven Bio), ARGX-109 (ArGEN-X), FM101 (Femta Pharmaceuticals, Lonza) and ALD518/BMS-945429 (Alder Biopharmaceuticals, Bristol-Myers Squibb).

38. The method of any one of embodiments 18 - 26, wherein the IL-6 antagonist is a single domain antibody, a V_{HH} Nanobody, an Fab, or a scFv.

39. The method of any one of embodiments 1 – 17, wherein the IL-6 antagonist is an anti-IL-6R antibody, or antigen-binding fragment or derivative thereof.

40. The method of embodiment 39, wherein the anti-IL-6R antibody, antigen-binding fragment, or derivative comprises the 6 CDRs of tocilizumab.

41. The method of embodiment 39, wherein the anti-IL-6R antibody, antigen-binding fragment, or derivative comprises the 6 CDRs of vobarilizumab.

42. The method of any one of embodiments 1 – 17, wherein the IL-6 antagonist is a JAK inhibitor.

43. The method of embodiment 42, wherein the JAK inhibitor is selected from the group consisting of tofacitinib (Xeljanz), decernotinib, ruxolitinib, upadacitinib, baricitinib, filgotinib, lestaurtinib, pacritinib, peficitinib, INCB-039110, ABT-494, INCB-047986 and AC-410.

44. The method of any one of embodiments 1 – 17, wherein the IL-6 antagonist is a STAT3 inhibitor.

45. The method of any one of embodiments 18-41, wherein the IL-6 antagonist is administered parenterally.

46. The method of embodiment 45, wherein the IL-6 antagonist is administered subcutaneously.

47. The method of any one of embodiments 42 or 43, wherein the IL-6 antagonist is administered orally.

48. The method of any one of embodiments 1-47, wherein the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase diuretic efficiency.

49. The method of embodiment 48, wherein the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase diuretic efficiency to normal levels.

50. The method of any one of embodiments 1 – 49, wherein the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase eGFR.

51. The method of embodiment 50, wherein the IL-6 antagonist is administered at a dose, on a schedule, and for a period sufficient to increase eGFR to normal levels.

52. The method of any one of embodiments 1-51, further comprising the subsequent step of determining the level of IL-6 in urine.

53. The method of any one of embodiments 1-51, further comprising the subsequent step of determining the level of IL-6 in plasma.

54. The method of any one of embodiments 1-51, further comprising the subsequent step of determining the level of IL-6 in urine and in plasma.

55. The method of any one of embodiments 52-54, further comprising a final step of adjusting the dose of IL-6 antagonist for subsequent administration based on IL-6 level determined in the immediately preceding step.

56. A method of determining if a subject requires treatment for cardiorenal syndrome, the method comprising:

- a) measuring a level of IL-6 in a urine or plasma sample from the subject,
 - b) comparing the measured level of IL-6 to a predetermined reference level,
 - c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,
- wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject has cardiorenal syndrome, and treatment is recommended.

57. A method of determining if a subject requires treatment for impaired glomerular filtration, comprising:

- a) measuring a level of IL-6 in a urine sample from the subject,
 - b) comparing the measured level of IL-6 to a predetermined reference level,
 - c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,
- wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject has impaired glomerular filtration, and treatment is recommended.

58. A method of determining if a subject requires treatment for low diuretic efficiency, comprising:

- a) measuring a level of IL-6 in a urine sample from the subject,
- b) comparing the measured level of IL-6 to a predetermined reference level,

c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,
wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject has low diuretic efficiency, and treatment is recommended.

59. A method of determining if a subject requires treatment for high urine angiotensin, comprising:

a) measuring a level of IL-6 in a urine sample from the subject,
b) comparing the measured level of IL-6 to a predetermined reference level,
c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,
wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject has high urine angiotensin, and treatment is recommended.

60. A method of determining if a subject requires treatment for high plasma renin, comprising:

a) measuring a level of IL-6 in a plasma sample from the patient,
b) comparing the measured level of IL-6 to a predetermined reference level,
c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,
wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject has high plasma renin, and treatment is recommended.

61. A method of determining if a subject is at risk of mortality due to cardiorenal syndrome and requires treatment, comprising:

a) measuring a level of IL-6 in a plasma sample from the patient,
b) comparing the measured level of IL-6 to a predetermined reference level,
c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,
wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject is at risk of mortality due to cardiorenal syndrome, and treatment is recommended.

62. The method of any one of embodiments 56-61, wherein the patient is a heart failure patient.

63. The method of any of embodiments 56-62, wherein the level of IL-6 is measured using an enzyme-linked immunosorbent assay (ELISA).

64. A method of treating cardiorenal syndrome in a patient comprising:
a) measuring a level of IL-6 in a urine sample or a plasma sample from the patient,
b) comparing the measured level of IL-6 to a predetermined reference level,
c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level
wherein when the measured level of IL-6 is greater than the corresponding reference level, treatment for cardiorenal syndrome is administered.

65. The method of embodiment 64, wherein the patient is a heart failure patient.

66. The method of embodiment 64 or embodiment 65, wherein the level of IL-6 is measured using an enzyme-linked immunosorbent assay (ELISA).

67. The method of any one of embodiments 64-66, wherein the treatment administered for cardiorenal syndrome comprises administering one or more of at least one diuretic, angiotensin-converting-enzyme inhibitor, angiotensin receptor blocker, natriuretic peptide, adenosine antagonist, an IL-6 antagonist, or any combination thereof.

68. The method of embodiment 67, wherein treating cardiorenal syndrome comprises administering at least one anti-IL-6 antibody or anti-IL-6R antibody.

69. A kit comprising reagents for an assay for measuring a level of IL-6 and written instructions, the written instructions comprising:
a) measuring a level of IL-6 in a urine or plasma sample from the subject,
b) comparing the measured level of IL-6 to a predetermined reference level,
c) determining whether or not the measured level of IL-6 is greater than the corresponding reference level,

wherein when the measured level of IL-6 is greater than the corresponding reference level, the subject has cardiorenal syndrome, and treatment is recommended.

70. The kit of embodiment 69, wherein the instructions direct the kit for use on a heart failure patient.

71. The kit of embodiment 69 or 70, wherein the kit contains reagents for measuring the level of IL 6 using an enzyme-linked immunosorbent assay (ELISA).

8.8. Experimental examples

[0224] The invention is further described in detail by reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified. Thus, the invention should in no way be construed as being limited to the following examples, but rather, should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

[0225] Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, practice the claimed methods of the present invention. The following working examples therefore, specifically point out the preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

[0226] The materials and methods employed in the experiments disclosed herein are now described.

8.8.1. Example 1: Urine and plasma IL-6 associations in heart failure patients

[0227] **Methods.** Consecutive heart failure (“HF”) patients receiving high dose diuretic therapy at the Yale Transitional Care Clinic (YTCC) were enrolled in the prospective observational study.

[0228] The YTCC is an outpatient clinic with a focus on diuretic and fluid status management. Patients present for 4-8 hours of treatment, during which they receive 1-3 doses of loop diuretic. The dosing protocol is determined by patient fluid status; the choice of oral (PO) torsemide or intravenous (IV) bumetanide is at the discretion of the treating physician.

During the treatment period all urine produced is saved in a cumulative collection container and sent to the clinical laboratory for electrolyte measurements at the conclusion of the visit. A cumulative urine collection is conducted during the treatment period. Additional spot urine samples are saved both before and one hour after diuretic administration.

[0229] Assays. A Randox RxDaytona™ automated clinical chemistry analyzer was used to measure urine and serum electrolytes using ion selective electrodes. Urea, creatinine, bicarbonate and cystatin C were measured using Randox reagents per the manufacturer's instructions (Randox Laboratories™, UK). Concentrations of interleukin 10 and IL-6 were measured using the MesoScale Discovery (MSD) platform (Meso Scale diagnostics™, Gaithersburg, MD, USA). Levels of amino terminal pro B-type natriuretic peptide (NT-proBNP) were measured at the Yale clinical laboratory on a Roche Elecsys 120 analyzer (Roche Diagnostics, Indianapolis, USA). Plasma renin activity (PRA), angiotensinogen and active renin were measured using commercially available competitive ELISA kits from ALPCO™ per manufacturer's instructions (ALPCO™, Salem, NH, USA). Total renin was also analyzed using commercially available ELISA kits (R&D Systems™, Minneapolis, USA). The total renin immunoassay kit from R&D systems™ recognizes both active renin and prorenin. The assay's mean detectable limit is 4.43 pg/ml for total renin and 0.81 pg/ml for active renin. Liquid chromatography mass spectrometry was used to measure levels of bumetanide and torsemide in urine. Ultra-high performance liquid chromatography was performed on the Agilent Infinity 1290 UPLC system™. Chromatographic separation was achieved on the Zorbax Bonus RP™ 2.1x50 mm 1.8μ column with a flow rate of 0.6 ml/min. The mobile phase contained of 0.1% Formic acid (Buffer A) and 80% acetonitrile in 0.1% Formic acid (Buffer B). Mass spectrometry was performed on Agilent Q-TOF system™ (Agilent™, Santa Clara, CA, USA) in positive ion mode.

[0230] Calculations and Definitions. eGFR was calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formula. Doses of loop diuretics were converted to furosemide equivalents with 1 mg bumetanide = 20 mg torsemide = 40 mg intravenous furosemide = 80 mg oral furosemide. As published previously, diuretic efficiency was defined as the increase in urinary sodium output over the treatment period per each doubling of the loop diuretic dose, centered on 40 mg of furosemide equivalents; this scale was chosen to account for the sigmoidal dose-response curve of these drugs. Urinary diuretic excretion was calculated by multiplying the concentration of diuretic in the urine by the volume of urine produced in the first 3 hours after diuretic administration; this quantity of

estimated diuretic was then normalized to the administered diuretic dose in furosemide equivalents, taking into account the published urinary clearance of the particular diuretic received by the patient (bumetanide or torsemide). Fractional excretions of sodium and potassium were calculated using the formula: Fractional excretion of X (FEX) = $\frac{[X]_{\text{Urine}} * [Creatinine]_{\text{Serum}}}{([X]_{\text{Serum or plasma}} * [Creatinine]_{\text{Urine}})}$. Urine proteins including renin, angiotensinogen and IL-6 were indexed to urinary creatinine. A low urine IL-6 was defined as a value less than the median of 14.2 pg/g of urinary creatinine. A low plasma IL-6 was defined as a value less than the median of 2.0 pg/mL. High levels of urine and plasma renin and angiotensinogen were defined as values greater than or equal to the cohort medians of these variables.

[0231] Statistical Analysis. Values reported are mean +/- SD, median (quartile 1 – quartile 3) and percentage. Correlations between continuous variables are Spearman's rho with the exception of adjusted correlations. Pearson's chi-squared test was used to compare categorical variables between groups. To compare continuous variables between groups, either Student's t-test or the Wilcoxon Rank Sum test was used. A log transformation was applied to skewed variables including plasma IL-6, urine IL-6 and NT-pro-BNP before entering them into multivariable models and partial correlation analyses. Logistic regression was used to evaluate association between the odds of low diuretic efficiency, an eGFR < 60 mL/min/1.73m², or high levels of urine or plasma neurohormonal parameters with plasma and urine levels of IL-6, both on a univariate level and with adjustment for plasma or urine IL-6 and/or eGFR. Cox proportional hazards modeling was used to evaluate time-to-event associations with all-cause mortality. Statistical analysis was performed with IBM SPSS Statistics version 23 (IBM Corp., Armonk, NY) and Stata version 13 (StataCorp™, College Station, TX). Statistical significance was defined as 2-tailed $p < 0.05$ for all analyses.

[0232] The baseline characteristics of our population are described in Table 2, below. In this subset, 98 patients underwent determination of IL-6 levels in blood and urine. The median (IQR) pre-diuretic level of urine IL-6 was 14.2 pg/g of creatinine (5.6-36.2 pg/g) whereas the median level of plasma IL-6 was 2.0 pg/mL (1.2-3.9 pg/mL). Plasma and urine IL-6 levels were only modestly correlated ($r=0.40$, $p<0.001$). Notably, those with lower than median levels of urine IL-6 tended to be younger, Caucasian, more often on angiotensin receptor blocker (ARB) or angiotensin converting enzyme (ACE-I) therapy, more likely to have heart failure with reduced ejection fraction (HFrEF), a higher GFR and substantially lower plasma

levels of NT-proBNP levels. The profile of patients defined by a lower than median plasma IL-6 was somewhat different, but largely reflected the same trends.

Table 2: Baseline Characteristics

	Overall Cohort (n=98)	Low Urine IL-6 (n=49)	High Urine IL-6 (n=49)	p-value	Low Plasma IL-6 (n=49)	High Plasma IL-6 (n=49)	p-value
Demographics							
Age	67.9 ± 13.5	62.1 ± 13.2	73.8 ± 11.2	<0.001*	65.8 ± 15.1	70.1 ± 11.5	0.12
Black race, %	36	57	15	<0.001*	47	26	0.04*
Male sex, %	44	49	39	0.31	49	39	0.31
Past Medical History, %							
Hypertension	91	94	88	0.29	96	86	0.08
Diabetes	47	55	39	0.11	37	57	0.04*
Gout	13	10	16	0.37	10	16	0.37
Ischemic etiology	30	33	27	0.46	33	27	0.46
Post-discharge follow-up visit	63	71	55	0.09	61	65	0.68
Baseline Medications, %							
ACEI or ARB	55	65	45	0.04*	67	43	0.02*
Beta-blocker	77	82	71	0.23	82	71	0.23
Thiazide-type diuretic	14	12	16	0.56	6	22	0.02*
MRA	33	37	29	0.39	31	35	0.67
Digoxin	10	12	8	0.50	8	12	0.50
Home loop diuretic dose (mg furosemide equivalents)	80 (60-160)	80 (60-160)	100 (80-160)	0.60	80 (40-120)	160 (80-200)	0.002*
Physical Exam							
Weight (kg)	98.2 ± 33.4	106.0 ± 37.9	90.4 ± 26.4	0.02*	98.4 ± 31.5	98.0 ± 35.5	0.95
BMI	34.8 ± 10.8	36.7 ± 11.3	32.5 ± 9.9	0.09	33.5 ± 8.4	36.2 ± 13.0	0.28

Table 2: Baseline Characteristics

	Overall Cohort (n=98)	Low Urine IL-6 (n=49)	High Urine IL-6 (n=49)	p-value	Low Plasma IL-6 (n=49)	High Plasma IL-6 (n=49)	p-value
Systolic blood pressure (mmHg)	122.6 ± 18.9	120.6 ± 19.1	124.5 ± 18.7	0.32	123.4 ± 18.9	121.7 ± 19.1	0.65
Heart rate (beats per minute)	75.9 ± 13.2	75.6 ± 13.2	76.2 ± 13.4	0.82	74.9 ± 13.3	76.8 ± 13.2	0.48
Echocardiogram							
Left ventricular ejection fraction (%)	45 (27-56)	35 (25-50)	51 (31-59)	0.03*	39 (29-54)	45 (26-57)	0.56
Ejection fraction ≥ 40%	53%	43%	63%	0.04*	49%	57%	0.42
Laboratory Values							
Sodium (mmol/L)	136 (134-140)	137 (134-138)	139 (134-141)	0.06	138 (135-140)	136 (132-140.0)	0.12
Chloride (mmol/L)	96 (95-102)	100 (96-102)	98 (95-102)	0.50	101 (98-103)	96 (91-101)	<0.001*
Hypochloremia, %	34	27	42	0.13	14	55	<0.001*
Potassium (mmol/L)	4.1 (3.7-4.5)	4.2 (3.8-4.6)	4.1 (3.6-4.3)	0.053	4.1 (3.7-4.5)	4.1 (3.7-4.4)	0.64
Bicarbonate (mmol/L)	23.9 (21.7-26.4)	23.1 (21.6-25.3)	24.9 (22.2-27.9)	0.03*	23.1 (21.5-25.2)	25.0 (22.6-28.0)	0.03*
Blood urea nitrogen (mg/dL)	29.0 (20.0-47.0)	27.0 (17.5-44.0)	32.0 (24.0-48.0)	0.10	27.0 (19.0-39.0)	36.0 (23.0-65.0)	0.04*
Serum creatinine (mg/dL)	1.5 ± 0.8	1.4 ± 0.6	1.7 ± 0.9	0.046*	1.3 ± 0.5	1.8 ± 1.0	0.007*
BUN/Creatinine ratio	24.0 ± 8.3	24.1 ± 8.1	23.9 ± 8.5	0.92	23.6 ± 8.0	24.5 ± 8.5	0.58
Estimated glomerular filtration rate (eGFR) (mL/min/1.73m ²)	54.9 ± 28.2	64.3 ± 30.1	45.7 ± 22.8	<0.001*	61.4 ± 27.2	48.3 ± 27.8	0.02*
eGFR < 60 mL/min/1.73m ²	61%	48%	73%	0.01*	55%	67%	0.24
Albumin (g/dL)	3.8 ± 0.4	3.9 ± 0.4	3.8 ± 0.4	0.045*	4.0 ± 0.4	3.7 ± 0.4	<0.001*
Hemoglobin (g/dL)	12.2 ± 2.3	12.7 ± 2.2	11.7 ± 2.3	0.04*	12.8 ± 2.0	11.5 ± 2.4	0.004*

Table 2: Baseline Characteristics

	Overall Cohort (n=98)	Low Urine IL-6 (n=49)	High Urine IL-6 (n=49)	p-value	Low Plasma IL-6 (n=49)	High Plasma IL-6 (n=49)	p-value
Diuretic Parameters							
Diuretic dose administered (mg furosemide equivalents)	140 (40-240)	80 (40-160)	160 (80-280)	0.001*	80 (40-160)	160 (80-320)	<0.001*
Diuretic administered, %							
IV Bumetanide:	45	27	63	<0.001*	33	57	0.02*
Torsemide:	55	73	37		67	43	

ACE-I = angiotensin-converting enzyme inhibitor. ARB = angiotensin receptor blocker. MRA = mineralocorticoid receptor antagonist. IL = interleukin. BUN = blood urea nitrogen. BNP = brain natriuretic peptide. BMI = body mass index. * $p < 0.05$.

Table 3: Plasma and Urine Biomarkers at Baseline

	Overall Cohort (n=98)	Low Urine IL-6 (n=49)	High Urine IL-6 (n=49)	p-value	Low Plasma IL-6 (n=49)	High Plasma IL-6 (n=49)	p-value
Plasma Biomarkers							
IL-6 (pg/mL)	2.0 (1.2-3.9)	1.4 (0.9-2.9)	2.8 (1.5-6.1)	<0.001*	1.2 (0.9-1.4)	3.9 (2.9-6.8)	<0.001*
IL-10 (pg/mL)	0.3 (0.2-0.7)	0.3 (0.2-0.4)	0.5 (0.3-0.8)	0.001*	0.3 (0.2-0.5)	0.5 (0.3-0.8)	0.01*
IL-10/IL-6 ratio	0.2 (0.1-0.4)	0.2 (0.1-0.3)	0.2 (0.1-0.4)	0.73	0.3 (0.2-0.6)	0.1 (0.1-0.2)	<0.001*
Total renin (pg/mL)	1167.7 (734.7-2759.7)	1169.2 (734.7-3013.5)	1166.3 (816.0-2390.9)	0.97	909.6 (595.2-1517.9)	1784.4 (921.6-3868.7)	<0.001*
Active renin (pg/mL)	36.8 (8.4-162.8)	79.0 (7.7-184.2)	28.1 (12.0-139.4)	0.38	13.0 (4.4-160.7)	49.1 (15.7-169.2)	0.07
Angiotensinogen (ng/mL)	167.9 (54.1, 445.3)	193.9 (48.2, 518.1)	160.3 (58.3, 430.2)	0.81	178.8 (31.6, 481.2)	167.2 (75.5, 445.3)	0.50
NT pro-BNP (pg/mL)	1825.0 (681.0-5030.0)	840.0 (256.0-3115.0)	3115.0 (1440.0-6070.0)	<0.001*	1440.0 (412.0-3010.0)	3390.0 (792.0-6410.0)	0.01*
Urine Biomarkers							
IL-6 (pg/g of creatinine)	14.2 (5.6-36.2)	5.6 (2.5-7.7)	36.2 (20.7-68.9)	<0.001*	7.0 (3.2-19.1)	21.2 (9.2-45.3)	<0.001*
IL-6 (pg/mL)	1.2 (0.5-2.6)	0.5 (0.2-0.7)	2.5 (1.6-4.0)	<0.001*	0.8 (0.4-1.9)	1.7 (0.6-3.2)	0.03*
Renin (pg/mg of creatinine)	0.3 (0.1-0.8)	0.2 (0.1-0.5)	0.6 (0.3-1.2)	<0.001*	0.2 (0.1-0.6)	0.5 (0.2-1.5)	<0.001*
Renin (pg/mL)	28.2 (10.2-88.3)	19.9 (7.5-53.1)	37.0 (13.4-109.6)	0.01*	23.7 (7.5-53.1)	35.9 (12.3-109.6)	0.06
Angiotensinogen (pg/mg creatinine)	18.7 (5.4-126.3)	6.2 (3.3-19.0)	123.5 (17.3-549.7)	<0.001*	7.3 (3.7-39.7)	33.7 (12.3-318.9)	0.002*
Angiotensinogen (pg/mL)	1530 (624-10941)	966 (231-1567)	5850 (1518-53865)	<0.001*	1105 (463-5801)	2153 (966-23952)	0.02*
FE _{Na} (%)	0.5 (0.2-1.3)	0.4 (0.2-1.2)	0.5 (0.2-1.4)	0.22	0.4 (0.1-1.0)	0.7 (0.3-1.8)	0.02*
FE _K (%)	18.4 (11.7-39.3)	15.5 (9.9-32.5)	21.6 (15.2-44.6)	0.02*	15.8 (9.6-24.5)	24.5 (13.9-50.3)	0.001*
FE _{Urea} (%)	63.6 (44.8-81.3)	59.5 (44.4-80.5)	65.5 (44.9-81.6)	0.62	66.7 (47.5-82.8)	58.2 (44.6-76.0)	0.20

IL = interleukin. NT pro-BNP = N-terminal pro b-type natriuretic peptide. FE_{Na} = fractional excretion of sodium. FE_K = fractional excretion of potassium. FE_{Urea} = fractional excretion of urea.

8.8.1.1. Kidney function and IL-6

[0233] As shown in Table 2, eGFR was lower in patients with high urine or plasma IL-6, but this association was only significant between eGFR and urine IL-6 ($p=0.01$). A correlation between both plasma IL-6 and eGFR ($r=-0.26$, $p=0.01$) as well as urine IL-6 and eGFR ($r=-0.38$, $p<0.001$) was observed. However, on adjustment for urine IL-6, there was no longer a significant association between plasma IL-6 and eGFR ($p=0.20$), whereas a significant association remained between urine IL-6 and eGFR after adjustment for plasma IL-6 (partial $r=-0.32$, $p=0.002$). Similarly, the risk of reduced eGFR as defined by an eGFR $< 60\text{mL/min/1.73m}^2$ was increased with higher levels of urine IL-6 (OR=1.9 per SD increase, 95% CI=1.2, 3.1, $p=0.006$) and not with higher levels of plasma IL-6 (OR=1.3 per SD increase, 95% CI=0.8-2.0, $p=0.25$).

8.8.1.2. Diuretic response and IL-6

[0234] There was an inverse association between diuretic efficiency and both urine IL-6 ($r=-0.43$, $p<0.001$) and plasma IL-6 ($r=-0.31$, $p=0.002$; FIG. 3). Odds for a low diuretic efficiency increased with higher levels of either urine IL-6 (OR=2.3 per SD increase, 95% CI 1.4-3.8, $p=0.001$) or plasma IL-6 (OR=1.7 per SD increase, 95% CI=1.1-2.7, $p=0.02$). Upon adjustment for eGFR, only urine IL-6 remained significantly associated with risk of low diuretic efficiency (adjusted OR=1.8 per SD increase, 95% CI 1.1-3.1, $p=0.02$; FIG. 3). Furthermore, when urine IL-6 and plasma IL-6 were both entered into a logistic regression model, only urine IL-6 remained associated with risk of low diuretic efficiency (adjusted OR=2.1 per SD increase, 95% CI 1.3-3.5, $p=0.004$) while plasma IL-6 showed no such association (OR=1.4, 95% CI 0.9-2.2, $p=0.17$).

8.8.1.3. Neurohormonal activation

[0235] Plasma IL-6 was associated with higher levels of plasma renin (Table 3 and FIG. 3). Notably, higher levels of plasma IL-6 conferred additional risk of high plasma total renin (OR=1.9 per SD increase, 95% CI 1.2-3.0, $p=0.008$), and this persisted despite adjustment for use of ACE-I or ARB and urine IL-6 levels (adjusted OR=2.3 per SD increase, 95% CI 1.3-3.9, $p=0.003$). Urine IL-6 was not associated with increased risk of high plasma renin (OR=1.0, 95% CI 0.7-1.5, $p=0.98$).

[0236] Urine IL-6 was strongly associated with high levels of tissue level RAAS activation as measured by urine angiotensinogen (OR=4.2 per SD increase, 95% CI 2.2-7.9, $p<0.001$)

and urine renin (OR=2.1 per SD increase, 95% CI 1.3-3.4, $p=0.002$; FIG. 3). These associations persisted after adjustment for plasma IL-6 levels and ACE-I or ARB use (adjusted OR for high urine angiotensinogen=4.2 per SD increase, 95% CI 2.2-8.3, $p<0.001$; adjusted OR for high urine renin=2.0 per SD increase, 95% CI 1.2-3.3, $p=0.005$). Plasma IL-6 did not demonstrate a univariate association with risk of high urine renin (OR=1.3 per SD increase, 95% CI 0.9-2.0, $p=0.16$). Although a trend toward association with urine angiotensinogen (OR=1.5 per SD increase, 95% CI 0.98-2.3, $p=0.06$) was observed, it was eliminated with adjustment for urine IL-6 (adjusted OR=1.1, 95% CI 0.7-1.7, $p=0.76$).

8.8.1.4. Association with survival

[0237] Over a median follow-up time of 713 days, 32 deaths occurred. Consistent with previous reports, increases in plasma IL-6 were associated with a higher risk of mortality (univariate HR=2.8 per SD increase, 95% CI 2.0-4.0, $p<0.001$). Plasma IL-6 remained associated with mortality after multivariable adjustment for baseline characteristics including age, race, baseline NT-proBNP levels, use of ACE-I or ARB, home loop diuretic dose and eGFR (adjusted HR=2.3 per SD increase, 95% CI=1.5-3.7, $p<0.001$). In contrast, urine IL-6 was not associated the risk of mortality (univariate HR=1.3 per SD increase, 95% CI 0.9-1.8, $p=0.15$; adjusted HR=1.02, 95% CI=0.6-1.6, $p=0.93$).

8.8.1.5. Summary

[0238] Plasma and urine IL-6 levels capture distinctive aspects of the role of IL-6 in cardiorenal disease pathophysiology. Plasma IL-6 levels are associated with global measures of disease severity such as risk of mortality. In contrast, urine IL-6, likely a measure of kidney inflammation, is strongly correlated with multiple measures of cardiorenal syndrome ("CRS") in these heart failure patients, including diuretic resistance, renin angiotensin aldosterone and system (RAAS) activation, and lower estimated glomerular filtration rate (eGFR).

[0239] These data demonstrate that urine IL-6 level is a useful biomarker for renal inflammation, and is particularly useful in assessing renal symptoms in heart failure patients. In addition, serial urine IL-6 levels should prove useful in monitoring and assessing the renal benefit of therapeutic interventions in heart failure.

8.8.2. Example 2: TMPRSS6 genotype stratifies heart failure patients whose renal symptoms are predicted to be responsive to IL-6 antagonists

[0240] The data obtained in Example 1 also predict that treatment with an IL-6 antagonist should be effective in reducing renal inflammation in heart failure patients.

[0241] However, because infection is often a precipitating cause of acute decompensation in heart failure patients, it is important to limit anti-cytokine and other immunosuppressive therapies to those heart failure patients who are likely to respond with improved renal and/or cardiac function. The cost of chronic IL-6 antagonist therapy also militates for limiting treatment to those heart failure patients who are likely to respond with improved renal and/or cardiac function.

[0242] Analysis conducted in Example 1 was expanded to 129 patients. FIG. 1A is a bar graph showing diuretic efficiency (“DE”) by tertiles of urine IL-6, confirming the inverse correlation of urinary IL-6 with diuretic efficiency observed in the 98 patient subset. FIG. 1B is a bar graph showing diuretic efficiency (“DE”) by tertiles of plasma IL-6 in these 129 patients, confirming an inverse correlation of plasma IL-6 with diuretic efficiency (uncorrected by urinary IL-6 levels).

[0243] Each patient’s genotype at the rs855791 single nucleotide polymorphism (“SNP”) in transmembrane protease serine 6 (“TMPRSS6”) was further assessed. The TMPRSS6 polypeptide, also known as Matriptase-2 (MT2), cleaves hemojuvelin and inhibits bone morphogenetic protein signaling. The rs855791 (G2321A) SNP alters the TMPRSS6 protein sequence: the allele with highest frequency in the human population (the major allele) is 2321G, encoding 736A; the allele with lowest frequency in the human population (minor allele) is 2321A, encoding 736V.

[0244] Genomic DNA was isolated from buffy coats using ReliaPrep large volume HT gDNA isolation system on the HSM Instrument (Promega, Madison, USA). The purity of the isolated DNA was assessed by Nanodrop. Genotyping was carried out at the Yale Centre for Genome analysis. Whole genome genotyping was done using Infinium® Exome-24 v1.0 BeadChip Kit from Illumina using standard protocols suggested by the manufacturer (Illumina, Inc., San Diego, CA). The amplification, fragmentation, precipitation, resuspension and hybridization steps were done manually. The arrays were scanned on the Illumina HiScan instrument. The Illumina HiScan or iScan System scans the BeadChip, using a laser to excite the fluorophore of the single-base extension product on the beads. The

scanner records high resolution images of the light emitted from the fluorophores. The Illumina GenomeStudio Genotyping Module, included with the Illumina Infinium Assay system, was used for extracting genotyping data from intensity data files (*.idat files) collected from the Illumina HiScan System. The Infinium Exome-24 v1.0 BeadChip contains over 240,000 putative functional exonic variants selected from over 12,000 individual human exome and whole-genome sequences. The > 240,000 markers represent diverse populations, including European, African, Chinese, and Hispanic individuals, and a range of common conditions, such as type 2 diabetes, cancer, metabolic, and psychiatric disorders. Detailed Illumina genotyping protocol is available at http://support.illumina.com/infinium_hts_assay_protocol_user_guide_15045738_a.pdf. The details of the SNPs in this exome chip are available at <http://support.illumina.com/downloads/infinium-exome-24-v1-0-product-files.html>.

[0245] As shown in FIG. 2A, urine levels of IL-6 were inversely correlated with diuretic efficiency only in the patients having at least one copy of the major allele of the TMPRSS6 rs855791 SNP (FIG. 2A, right panel, “AG+GG”); urine levels of IL-6 are not significantly correlated with diuretic efficiency in patients homozygous for the minor allele (FIG. 2A, left panel, “AA”).

[0246] As shown in FIG. 2B, plasma levels of IL-6 correlated inversely with diuretic efficiency only in the patients having at least one copy of the major allele of the TMPRSS6 rs855791 SNP (FIG. 2B, right panel, “AG+GG”); plasma levels of IL-6 are not significantly correlated with diuretic efficiency in patients homozygous for the minor allele (FIG. 2B, left panel, “AA”).

[0247] These data indicate that treatment of heart failure with an IL-6 antagonist will only improve renal symptoms in heart failure patients having at least one copy of the TMPRSS6 rs855791 major allele.

8.8.3. Example 3: Correlation of IL-6 with the expression of NKCC2, ENaC-beta, and NCC in the absence or presence of Ruxolitinib

Methods

[0248] The mouse M1 CCD cell line (American Type Culture Collection (ATCC), Cat# CRL-2038) was maintained the cell culture medium containing equal amount of DMEM (Sigma-Aldrich, Cat#D6046) and Han F12 (Sigma-Aldrich, Cat#11765-047), supplemented

with 5% Fetal Bovine Serum (FBS), 1% Penicillin-Streptomycin (Thermo Fisher Scientific, Cat# 15140-122), 1% Insulin-Transferrin-Selenium (Thermo Fisher Scientific, Cat# 51500-056) and 100 nM Dexamethasone (Sigma-Aldrich, Cat# D4902-100MG).

[0249] M1 CCD cells were seeded at 1 million/well in 6-well plate in the cell culture medium, and incubated overnight in a 37°C, 5% CO₂ incubator on Day 0. The cell culture medium was changed to DMEM/F12 serum-free medium on Day 1 and the cells were incubated overnight in a 37°C, 5% CO₂ incubator. On Day 2, the serum-free medium was removed and the cell culture medium was added to each well. Ruxolitinib (Selleckchem, Cat# S1378) was added at the final concentration of 1 μM or 100 μM 10 min before the addition of IL-6 (Sigma-Aldrich, Cat# SRP3096-20UG) at the final concentration of 10 ng/mL, 100 ng/mL, or 500 ng/mL. A control well without Ruxolitinib or IL-6 was included. The cells were treated for 24 hours with IL-6 and/or Ruxolitinib. The cells of each well were washed with once with 1xPBS and collected in 250 μL 1xPIPA buffer (10x, Millipore, Cat# 20-188) supplemented with 1% protease inhibitor cocktail (100x, Thermo Fisher Scientific, Cat# 78430). The sample were analyzed by immunoblotting using anti-NKCC2 antibody (Millipore, Cat# AB3562P), anti-ENaC-beta antibody (Millipore, Cat# AB3532P), or anti-NCC antibody (Millipore, Cat# AB3553) and the protein expression was quantified. Each experiment was done in triplicate.

Results

[0250] We examined the expression of NKCC2 (Na-K-Cl cotransporter 2), ENaC-beta (epithelial sodium channel, beta subunit), and NCC (sodium-chloride symporter) proteins in mouse M1 CCD cell line after treatment of IL-6 in the absence or presence of a JAK inhibitor, Ruxolitinib. The mouse M1 CCD cells are genotypically analogous to human cells homozygous for the *TMPRSS6* rs855791 major allele. As shown in FIG. 4A, 4B, and 4C, the treatment of IL-6 increased the expression of NKCC2, ENaC-beta, and NCC. Ruxolitinib blocked the effect of IL-6 on the ion transporter proteins.

[0251] Increased expression of these ion transporters provides a putative mechanism for IL-6 mediated diuretic resistance.

[0252] Because the increased expression is not known to be linked to hepcidin expression, these data suggested that IL-6 antagonism could be effective in treating diuretic resistance even in patients homozygous for the *TMPRSS6* rs855791 minor allele.

8.8.4. Example 4: Correlation of IL-6 with the expression of NKCC2, ENaC-beta, and NCC in the absence or presence of Spironolactone

Methods

[0253] The mouse M1 CCD cell line (American Type Culture Collection (ATCC), Cat# CRL-2038) was maintained the cell culture medium containing equal amount of DMEM (Sigma-Aldrich, Cat#D6046) and Han F12 (Sigma-Aldrich, Cat#11765-047), supplemented with 5% Fetal Bovine Serum (FBS), 1% Penicillin-Streptomycin (Thermo Fisher Scientific, Cat# 15140-122), 1% Insulin-Transferrin-Selenium (Thermo Fisher Scientific, Cat# 51500-056) and 100 nM Dexamethasone (Sigma-Aldrich, Cat# D4902-100MG).

[0254] M1 CCD cells were seeded at 1 million/well in 6-well plate in the cell culture medium, and incubated overnight in a 37°C, 5% CO₂ incubator on Day 0. The cell culture medium was changed to DMEM/F12 serum-free medium on Day 1 and the cells were incubated overnight in a 37°C, 5% CO₂ incubator. On Day 2, the serum-free medium was removed and the cell culture medium was added to each well. Spironolactone (Selleckchem, Cat# S4054) was added at the final concentration of 1 µM or 100 µM 10 min before the addition of IL-6 (Sigma-Aldrich, Cat# SRP3096-20UG) at the final concentration of 10 ng/mL, 100 ng/mL, or 500 ng/mL. A control well without Spironolactone or IL-6 was included. The cells were treated for 24 hours with IL-6 and/or Spironolactone. The cells of each well were washed with once with 1xPBS and collected in 250 µL 1xPIPA buffer (10x, Millipore, Cat# 20-188) supplemented with 1% protease inhibitor cocktail (100x, Thermo Fisher Scientific, Cat# 78430). The sample were analyzed by immunoblotting using anti-NKCC2 antibody (Millipore, Cat# AB3562P), anti-ENaC-beta antibody (Millipore, Cat# AB3532P), or anti-NCC antibody (Millipore, Cat# AB3553) and the protein expression was quantified. Each experiment was done in triplicate.

Results

[0255] We examined the expression of NKCC2 (Na-K-Cl cotransporter 2), ENaC-beta (epithelial sodium channel, beta subunit), and NCC (sodium-chloride symporter) proteins in mouse M1 CCD cell line after treatment of IL-6 in the absence or presence of a potassium-sparing diuretic, Spironolactone. The mouse M1 CCD cells are genotypically analogous to human cells homozygous for the *TMPRSS6* rs855791 major allele. As shown in FIG. 5A, 5B, and 5C, the treatment of IL-6 increased the expression of NKCC2, ENaC-beta, and NCC. Spironolactone blocked the effect of IL-6 on the ion transporter proteins.

[0256] Increased expression of these ion transporters provides a putative mechanism for IL-6 mediated diuretic resistance.

[0257] Because the increased expression is not known to be linked to hepcidin expression, these data suggested that IL-6 antagonism could be effective in treating diuretic resistance even in patients homozygous for the *TMPRSS6* rs855791 minor allele.

8.8.5. Example 5: Association of IL-6 with diuretic response in patients hospitalized for acute heart failure (HF)

[0258] **Methods.** Data from the PROTECT trial (Weatherley *et al.*, 2010, *J. Card. Fail.* 16:25–35; Massie *et al.*, 2010, *N. Engl. J. Med.* 363:1419–1428) was analyzed according to tertiles of IL-6. The PROTECT trial was a randomized placebo-controlled trial testing the effect of Adenosine A1–Receptor Antagonist Rolofylline on dyspnea relief, risk of worsening renal function and clinical outcomes. The key inclusion and exclusion criteria of the trial are shown below.

TABLE A	
Inclusion	Exclusion
>18 years	Acute contrast-induced nephropathy
History of HF >14 days with diuretic therapy	Temp >38 or sepsis requiring IV antimicrobial treatment
Hospitalized for ADHF* requiring IV diuretic therapy	Serum potassium <3.5 mEq/L
Within 24hrs of presentation to the hospital	Ongoing or planned IV therapy for ADHF with positive inotropic agents, vasopressors, vasodilators
Anticipated need for IV furosemide >40mg/day for at least 24hr	BNP <500 pg/mL or NT-pro-BNP <2000 pg/mL
Impaired renal function defined as creatinine clearance of admission between 20 and 80 mL/min (Cockcroft-Gault)	Ongoing or planned treatment with ultrafiltration
Systolic blood pressure >95 mm Hg	Severe pulmonary disease
	Clinical evidence of acute coronary syndrome in the 2 weeks before screening
	Hgb <8 g/dL, or Hct <25%, or the need for a blood transfusion
	Systolic blood pressure ≤160 mm Hg at randomization

*ADHF: dyspnea at rest or with minimal exertion and signs of fluid overload manifested by at least one of the following at time of randomization: JVP >8 cm, or Pulmonary rales ≥1/3 up the lung fields, not clearing with cough, or ≥2+ peripheral edema, or presacral edema

[0259] In total, 2033 patients with ADHF were included in the PROTECT study. Of these patients, IL-6 was measured by Singulex in 1445 patients at admission (baseline), 1462 patients at day 2 (24hr after baseline) and 1445 patients at day 7. Diuretic response was defined as weight change on day 4 per 40 mg of furosemide (or equivalent doses) administered from baseline to day 3. The primary endpoint of this study was all-cause mortality at 180 days.

[0260] **Statistical analysis.** Baseline characteristics are presented according to tertiles of IL-6. Differences between tertiles of baseline characteristics were tested using one-way analysis of variance (ANOVA), Kruskal Wallis or chi2-test where appropriate. Univariable linear regression was performed using diuretic response as the dependent variable and (log-transformed) IL-6 at baseline as the independent variable correcting for clinically relevant variables associated with diuretic response. Survival analysis was performed using Cox regression analysis correcting for clinically relevant variables and the PROTECT risk model (O'Connor *et al.*, 2012, *Eur. J. Heart Fail.* 14:605–612). The PROTECT risk model includes: previous hospitalization for HF, edema, systolic blood pressure, sodium levels, BUN, creatinine and albumin at admission.

[0261] **Results.** The baseline characteristics of the population are described in Table 4, below. Higher levels of IL-6 at baseline are associated with higher levels of BNP, anemia, eGFR <60 and older age (FIG. 6).

Table 4: Baseline Characteristics				
Factor	1st tertile	2nd tertile	3rd tertile	p-value
N	531	530	530	
Demographics	0.66-7.8	7.9-16.1	16.2-274.2	
	pg/mL	pg/mL	pg/mL	
Age	69.2 (11.5)	70.8 (11.0)	72.5 (10.8)	<0.001
Sex	347 (65.3%)	349 (65.8%)	354 (66.8%)	0.88
BMI	28.3 (5.8)	28.8 (6.0)	29.1 (6.2)	0.093
eGFR	52.5 (20.2)	47.2 (19.3)	44.9 (17.4)	<0.001
NYHA class				
I/II	107 (21.0%)	71 (14.1%)	96 (19.2%)	0.008
III	267 (52.4%)	263 (52.2%)	237 (47.4%)	
IV	136 (26.7%)	170 (33.7%)	167 (33.4%)	
LVEF	30 (25, 40)	30 (20, 40)	30 (22, 40)	0.35
HFpEF	42 (15.6%)	39 (16.3%)	38 (14.9%)	0.92
Systolic BP	126.2 (17.4)	124.5 (17.5)	123.6 (17.4)	0.043
Diastolic BP	75.3 (11.2)	73.5 (12.1)	72.3 (11.9)	<0.001
Heart Rate	79.2 (14.6)	79.4 (15.8)	81.3 (16.0)	0.052
Respiratory rate	20.6 (4.3)	21.2 (4.1)	21.7 (4.9)	<0.001

Table 4: Baseline Characteristics				
Factor	1st tertile	2nd tertile	3rd tertile	p-value
Medical history				
Atrial fibrillation	95 (45.5%)	92 (43.6%)	78 (39.4%)	0.45
Valve disease	196 (37.0%)	195 (36.8%)	208 (39.5%)	0.61
Mitral regurgitation	174 (32.8%)	178 (33.6%)	184 (34.8%)	0.78
Aortic stenosis	18 (3.4%)	16 (3.0%)	32 (6.0%)	0.027
Aortic regurgitation	43 (8.1%)	18 (3.4%)	38 (7.2%)	0.004
Heart failure (HF)	510 (96.0%)	505 (95.3%)	501 (94.5%)	0.51
Hospitalization for HF previous year	258 (48.6%)	266 (50.2%)	259 (48.9%)	0.86
No. of HF hospitalizations, median (IQR)	1.0 (1.0, 2.0)	1.0 (1.0, 2.0)	1.0 (1.0, 2.0)	0.62
Ischemic heart disease	362 (68.4%)	396 (74.9%)	365 (68.9%)	0.037
Myocardial infarction	255 (48.3%)	296 (56.0%)	242 (45.7%)	0.003
Hypertension	424 (79.8%)	422 (79.6%)	428 (80.8%)	0.89
Stroke or PVD	78 (14.7%)	102 (19.3%)	117 (22.2%)	0.007
Anemia	174 (36.3%)	216 (47.2%)	238 (50.6%)	<0.001
Thyroid disease	60 (11.3%)	61 (11.5%)	57 (10.8%)	0.92
Depression	28 (5.3%)	41 (7.7%)	35 (6.6%)	0.27
Hyperlipidemia	298 (56.1%)	267 (50.5%)	244 (46.0%)	0.004
Current smoker	92 (17.4%)	114 (21.6%)	105 (19.8%)	0.23
COPD or asthma	106 (20.0%)	102 (19.2%)	112 (21.2%)	0.73
Diabetes mellitus	247 (46.5%)	229 (43.2%)	258 (48.7%)	0.2
History of Atrial Fibrillation/Flutter	260 (49.2%)	291 (55.0%)	310 (58.8%)	0.007
AICD	73 (13.7%)	106 (20.0%)	68 (12.8%)	0.002
Biventricular pacemaker	54 (10.2%)	60 (11.3%)	47 (8.9%)	0.41
Pacemaker	50 (9.5%)	66 (12.5%)	65 (12.3%)	0.22
Medication				
Beta-blockers	417 (78.7%)	396 (74.7%)	385 (72.6%)	0.068
ACEIs	353 (66.6%)	312 (58.9%)	330 (62.3%)	0.033
ARBs	85 (16.0%)	85 (16.0%)	71 (13.4%)	0.38
ACE-I/ARB	424 (80.0%)	390 (73.6%)	393 (74.2%)	0.026
MRA	264 (49.8%)	226 (42.6%)	229 (43.2%)	0.033
Digoxin	159 (30.0%)	153 (28.9%)	146 (27.5%)	0.68
Nitrates	136 (25.7%)	146 (27.6%)	135 (25.5%)	0.68
Hydralazine	17 (3.2%)	15 (2.8%)	13 (2.5%)	0.76
CCBs	74 (14.0%)	71 (13.4%)	81 (15.3%)	0.67
Signs and symptoms				
Orthopnea	445 (84.3%)	440 (84.5%)	435 (82.5%)	0.65
Dyspnea at rest (NYHA IV)	264 (50.6%)	304 (59.1%)	327 (64.8%)	<0.001
Angina pectoris	134 (25.2%)	117 (22.1%)	111 (21.0%)	0.24
CCS Class III & IV	48 (36.4%)	38 (33.0%)	29 (26.6%)	0.27
Edema	82 (15.4%)	151 (28.5%)	184 (34.8%)	<0.001
Jugular venous distension	190 (39.7%)	195 (40.5%)	199 (41.6%)	0.82
Edema & raised JVP	127 (26.5%)	148 (30.7%)	150 (31.4%)	0.2
Rales	37 (7.0%)	55 (10.4%)	62 (11.7%)	0.027

[0262] The association of IL-6 levels and diuretic response is shown in Table 5, below. The diuretic response was defined as weight change on day 4 per 40 mg of furosemide (or equivalent doses) administered from baseline to day 3.

Table 5: Diuretic Response		
	Beta	p-value
<i>Univariable</i>	0.06	0.027
<i>Model 1 (age, sex)</i>	0.06	0.030
<i>Model 2(model 1 + eGFR, BMI)</i>	0.06	0.035

[0263] Table 6 and FIGs. 7A and 7B show the association of IL-6 levels with all-cause mortality at 180 days and the association of IL-6 levels with all-cause mortality at 60 days and/or cardiovascular related rehospitalization (CV hosp).

Table 6: Cox Regression Results		
	All-cause mortality at 180 days	60 days mortality and/or CV hosp
<i>Univariable</i>	1.59 (1.43-1.76) <0.001	1.14 (1.04-1.26) 0.007
<i>Model 1</i>	1.57 (1.41-1.74) <0.001	1.14 (1.04-1.26) 0.007
<i>Model 2</i>	1.55 (1.39-1.72) <0.001	1.11 (1.00-1.22) 0.049
<i>Model 3</i>	1.50 (1.35-1.68) <0.001	1.09 (0.99-1.21) 0.087
<i>PROTECT model</i>	1.41 (1.26-1.58) <0.001	1.06 (0.95-1.17) 0.309

Model 1: age, sex

Model 2: model 1 + eGFR, BMI

Model 3: model 2 + BNP

PROTECT model: age, previous hospitalization for HF, edema at admission, sodium, bun (log), creatinine (log) and albumin

[0264] Table 7 and FIGs. 8A and 8B show that an increase in IL-6 levels at day 7 compared to the baseline predicts adverse outcomes.

Table 7: Cox Regression Results			
180 days all-cause mortality			
	>1 pg/mL decrease	<1.0 pg/mL increase/decrease	>1.0 pg/mL increase
<i>Univariable</i>	2.13 (1.16-3.91) 0.014	ref	3.01 (1.65-5.49) <0.001
<i>Model 1</i>	1.79 (0.96-3.32) 0.066	ref	2.80 (1.53-5.11) 0.001
<i>Model 2</i>	1.80 (0.95-3.44) 0.073	ref	2.65 (1.41-4.97) 0.002
<i>Model 3</i>	1.65 (0.87-3.16) 0.128	ref	2.35 (1.25-4.44) 0.008
<i>PROTECT</i>	1.71 (0.90-3.26) 0.101	ref	2.38 (1.30-4.47) 0.007
60 days mortality and/or CV hosp			
	>1 pg/mL decrease	<1.0 pg/mL increase/decrease	>1.0 pg/mL increase
<i>Univariable</i>	1.31 (0.91-1.92) 0.146	ref	1.83 (1.26-2.65) 0.001
<i>Model 1</i>	1.31 (0.89-1.92) 0.002	ref	1.81 (1.25-2.62) 0.002
<i>Model 2</i>	1.29 (0.88-1.92) 0.187	ref	1.70 (1.16-2.48) 0.006
<i>Model 3</i>	1.28 (0.86-1.89) 0.223	ref	1.67 (1.14-2.45) 0.008

<i>PROTECT</i>	1.20 (0.81-1.78) 0.353	ref	1.57 (1.07-2.30) 0.020
----------------	------------------------	-----	------------------------

Model 1: age, sex

Model 2: model 1 + eGFR, BMI

Model 3: model 2 + BNP

PROTECT model: age, previous hospitalization for HF, edema at admission, sodium, bun (log), creatinine (log) and albumin

8.8.6. Example 6: Association of IL-6 levels with outcomes in a worsening heart failure population

[0265] **Methods.** Data from the BIOlogy Study to TAilored Treatment in Chronic Heart Failure (BIOSTAT-CHF) study was analyzed to investigate the association of IL-6 with outcomes in patients with worsening heart failure. In brief, BIOSTAT-CHF was a multicenter, multinational, prospective, observational study including 2516 patients from 69 centers in 11 European countries (Voors *et al.*, 2016, *Eur. J. Heart Fail.* 18:716–726). We performed secondary analyses in the BIOSTAT-CHF study, excluding patients with ferritin <100 from subsequent analysis. Inclusion criteria for the index cohort include: patients with >18 years of age, having symptoms of new-onset or worsening HF, confirmed either by a LVEF of ≤40% or BNP and/or NT-proBNP plasma levels >400 pg/ml or >2,000 pg/ml, respectively. Furthermore, these patients had not been previously treated with an ACEi/ARBs and/or beta-blocker or they were receiving ≤50% of the target doses of these drugs at the time of inclusion and anticipated initiation or up-titration of ACEi/ARBs and beta-blockers. All patients needed to be treated with loop diuretics.

Table B
Inclusion criteria
Age >18
Diagnosed with HF
Previous documented admission with HF requiring diuretic treatment
Treated with furosemide >20 mg/day or equivalent
Not previously treated or receiving <50% of target doses of ACE inhibitors/ARBs and/or beta-blockers according to 2008 ESC guidelines
Anticipated uptitration of ACE inhibitors/ARBs and/or beta-blockers

[0266] In total, IL-6 was measured in 2329 patients with worsening HF from the BIOSTAT-CHF study. The primary outcome of this study was a composite outcome of all-cause mortality and hospitalization for HF.

[0267] **Statistical analyses.** Baseline characteristics are presented according to tertiles of IL-6. Differences between tertiles of baseline characteristics were tested using one-way analysis of variance (ANOVA), Kruskal Wallis or chi2-test where appropriate. Survival analysis was performed using Cox regression analysis correcting for clinically relevant variables and the BIOSTAT-CHF risk model. The BIOSTAT-CHF risk model for all-cause mortality and/or hospitalization for heart failure includes: age, N-terminal pro-B-type natriuretic peptide (NT-proBNP), hemoglobin (Hb), the use of a beta-blocker at time of inclusion, a HF-hospitalization in year before inclusion, peripheral edema, systolic blood pressure, high-density lipoprotein cholesterol and sodium (Voors *et al.*, 2017, *Eur. J. Heart Fail.* 19:627-634). We performed interaction analysis between ferritin levels and IL-6 for the primary outcome. To investigate the association of IL-6 with outcomes depending on the position of the TMPRS6 SNP (rs855791).

[0268] **Results.** The baseline characteristics of the population are described in Table 8, below. Higher levels of IL-6 at baseline are associated with higher levels of NTproBNP and Anemia (FIG. 9).

Table 8: Baseline Characteristics				
Factor	1st tertile	2nd tertile	3rd tertile	p-value
N	781	775	773	
Demographics	0.3-3.4 pg/mL	3.5-7.8 pg/mL	7.9-260.7 pg/mL	
Age	66.3 (12.2)	69.3 (11.7)	70.9 (11.7)	<0.001
Female	200 (25.6%)	200 (25.8%)	212 (27.4%)	0.67
HF status				
HF _r EF	607 (84.1%)	568 (81.6%)	524 (78.3%)	<0.001
HF _m rEF	92 (12.7%)	74 (10.6%)	87 (13.0%)	
HF _p EF	23 (3.2%)	54 (7.8%)	58 (8.7%)	
BMI	27.9 (5.1)	27.7 (5.6)	27.7 (5.6)	0.77
Ischemic etiology	329 (43.2%)	353 (46.4%)	357 (46.7%)	0.31
NYHA				
I	70 (9.0%)	82 (10.6%)	62 (8.0%)	0.005
II	400 (51.2%)	355 (45.8%)	320 (41.4%)	
III	199 (25.5%)	216 (27.9%)	251 (32.5%)	
IV	22 (2.8%)	25 (3.2%)	33 (4.3%)	
NA	90 (11.5%)	97 (12.5%)	107 (13.8%)	
Systolic BP	126.4 (20.0)	124.3 (22.4)	123.7 (23.5)	0.034
Diastolic BP	76.7 (12.7)	74.8 (13.4)	73.5 (14.0)	<0.001
LVEF	30.6 (8.9)	30.9 (11.1)	31.3 (11.7)	0.54
Heart Rate	76.4 (18.2)	80.1 (18.9)	83.6 (21.0)	<0.001
Signs and symptoms				
Peripheral edema				
Not Present	352 (57.2%)	253 (39.7%)	180 (26.4%)	<0.001
Ankle	170 (27.6%)	188 (29.5%)	221 (32.5%)	

Table 8: Baseline Characteristics				
Factor	1st tertile	2nd tertile	3rd tertile	p-value
Below Knee	83 (13.5%)	156 (24.5%)	194 (28.5%)	
Above Knee	10 (1.6%)	40 (6.3%)	86 (12.6%)	
Elevated JVP				
No	420 (76.5%)	318 (58.9%)	289 (53.7%)	<0.001
Yes	106 (19.3%)	190 (35.2%)	216 (40.1%)	
Uncertain	23 (4.2%)	32 (5.9%)	33 (6.1%)	
Hepatomegaly	75 (9.6%)	128 (16.6%)	132 (17.1%)	<0.001
Orthopnea	168 (21.6%)	273 (35.3%)	361 (46.8%)	<0.001
Medical history				
Anemia	158 (23.6%)	271 (38.1%)	348 (47.0%)	<0.001
Atrial fibrillation	293 (37.5%)	372 (48.0%)	387 (50.1%)	<0.001
Diabetes mellitus	208 (26.6%)	270 (34.8%)	276 (35.7%)	<0.001
COPD	113 (14.5%)	126 (16.3%)	163 (21.1%)	0.002
Renal disease	152 (19.5%)	229 (29.5%)	261 (33.8%)	<0.001
Hypertension	486 (62.2%)	496 (64.0%)	473 (61.2%)	0.51
Peripheral artery disease	60 (7.7%)	92 (11.9%)	105 (13.6%)	<0.001
Stroke	59 (7.6%)	84 (10.8%)	76 (9.8%)	0.075
PCI	163 (20.9%)	166 (21.4%)	173 (22.4%)	0.76
CABG	120 (15.4%)	122 (15.7%)	156 (20.2%)	0.02
Medication				
Loop diuretics	779 (99.7%)	769 (99.2%)	769 (99.5%)	0.36
ACE/ARB	601 (77.0%)	564 (72.8%)	518 (67.0%)	<0.001
Beta blocker	675 (86.4%)	654 (84.4%)	603 (78.0%)	<0.001
Aldosterone antagonist	446 (57.1%)	411 (53.0%)	382 (49.4%)	0.01
Laboratory				
Hemoglobin	13.7 (1.7)	13.2 (1.8)	12.7 (2.0)	<0.001
Total cholesterol	4.5 (3.7, 5.5)	4.1 (3.4, 4.9)	3.7 (3.1, 4.5)	<0.001
AST	24.0 (19.0, 32.0)	25.0 (19.0, 34.0)	27.0 (20.0, 39.0)	<0.001
ALT	25.0 (18.0, 37.0)	25.0 (17.0, 37.0)	23.5 (15.0, 40.0)	0.12
Sodium	140.0 (138.0, 142.0)	140.0 (137.0, 142.0)	139.0 (136.0, 141.0)	<0.001
Potassium	4.3 (4.0, 4.6)	4.2 (3.9, 4.6)	4.2 (3.8, 4.6)	<0.001
HbA1c	6.0 (5.6, 6.7)	6.5 (5.9, 7.5)	6.5 (5.9, 7.3)	<0.001
NT-proBNP	2661.0 (1445.0, 4820.0)	4344.0 (2517.0, 7837.0)	5734.5 (3141.0, 11452.0)	<0.001
Troponin-I	0.0 (0.0, 0.1)	0.0 (0.0, 0.1)	0.0 (0.0, 0.1)	<0.001

[0269] As shown in Table 9 and FIGs 10A and 10B, levels of IL-6 measured at baseline were associated with the combined outcome of all-cause mortality and/or a hospitalization for HF as well as all-cause mortality alone at two years.

Table 9: Cox Regression Analyses		
	<i>All-cause mort and/or HF hosp at 2 years</i>	<i>All-cause mortality at 2 years</i>
	HR (95%CI) p-value	HR (95%CI) p-value
<i>Univariable</i>	1.38 (1.31-1.46) <0.001	1.42 (1.32-1.53) <0.001
<i>Model 1</i>	1.34 (1.26-1.42) <0.001	1.48 (1.38-1.58) <0.001
<i>Model 2</i>	1.25 (1.17-1.33) <0.001	1.34 (1.24-1.44) <0.001
<i>Model 3</i>	1.24 (1.16-1.32) <0.001	1.33 (1.23-1.44) <0.001
<i>BIOSTAT model</i>	1.13 (1.04-1.19) 0.001	1.20 (1.11-1.31) <0.001

Model 1: Age, sex

Model 2: model 1+ BMI, country, hypertension (history of), diabetes (history of) and anemia

Model 3: model 2+ beta-blocker at baseline, ACEi/ARB at baseline and MRA at baseline

BIOSTAT model: age, N-terminal pro-B-type natriuretic peptide (NT-proBNP), hemoglobin (Hb), the use of a beta-blocker at time of inclusion, a HF-hospitalization in year before inclusion, peripheral edema, systolic blood pressure, high-density lipoprotein cholesterol and sodium

[0270] As shown in Table 10 and FIG. 11, there is no differential association of IL-6 with outcome depending on ferritin status.

Table 10: Cox Regression Analyses		
	<i>Ferritin>100 ug/L</i>	<i>Interaction ferritin*IL-6</i>
	HR (95%CI) p-value	
<i>Univariable</i>	1.39 (1.27-1.52) <0.001	0.610
<i>Model 1</i>	1.35 (1.23-1.48) <0.001	
<i>Model 2</i>	1.26 (1.14-1.40) <0.001	
<i>Model 3</i>	1.26 (1.14-1.39) <0.001	
<i>BIOSTAT model</i>	1.11 (0.99-1.22) 0.052	

Model 1: Age, sex

Model 2: model 1+ BMI, country, hypertension (history of), diabetes (history of) and anemia

Model 3: model 2+ beta-blocker at baseline, ACEi/ARB at baseline and MRA at baseline

BIOSTAT model: age, N-terminal pro-B-type natriuretic peptide (NT-proBNP), hemoglobin (Hb), the use of a beta-blocker at time of inclusion, a HF-hospitalization in year before inclusion, peripheral edema, systolic blood pressure, high-density lipoprotein cholesterol and sodium

[0271] As shown in Table 11, there was no differential association of IL-6 with outcome depending on TMPRSS6 genotype.

Table 11: IL-6 according to TMPRSS6 allele	
<i>TMPRSS6</i>	<i>HR (95%CI) p-value</i>
AA (n=191)	1.50 (1.19-1.89) 0.001
AG (n=579)	1.34 (1.17-1.54) <0.001
GG (n=409)	1.54 (1.32-1.79) <0.001

9. INCORPORATION BY REFERENCE

[0272] The disclosures of each and every patent, patent application, and publication cited herein are hereby incorporated herein by reference in their entirety.

10. EQUIVALENTS

[0273] While this invention has been disclosed with reference to specific embodiments, it is apparent that other embodiments and variations of this invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention. The appended claims are intended to be construed to include all such embodiments and equivalent variations.

[0274] Throughout the description and claims of the specification, the word “comprise” and variations of the word, such as “comprising” and “comprises”, is not intended to exclude other additives, components, integers or steps.

[0275] A reference herein to a patent document or other matter which is given as prior art is not to be taken as admission that the document or matter was known or that the information it contains was part of the common general knowledge as at the priority date of any of the claims.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method of treating left ventricular heart failure, the method comprising:
administering to a patient with left ventricular heart failure a therapeutically effective amount of an anti IL-6 antibody.
2. The method of claim 1, wherein the patient has elevated pre-treatment plasma IL-6 levels.
3. The method of claim 2, wherein the patient has a pre-treatment plasma IL-6 level of greater than 2 pg/mL.
4. The method of claim 1, wherein the patient has a diuretic efficiency of less than 500 mmol Na/doubling of loop diuretic dose.
5. The method of claim 4, wherein the patient has a diuretic efficiency of less than 200 mmol Na/doubling of loop diuretic dose.
6. The method of claim 1, wherein the patient has diuretic resistant heart failure.
7. The method of claim 1, wherein the patient has acute heart failure.
8. The method of claim 1, wherein the patient has chronic heart failure.
9. The method of claim 1, wherein the anti-IL-6 antibody comprises all six variable region CDRs of MEDI5117.
10. The method of claim 9, wherein the antibody comprises the VH and VL of MEDI5117.
11. The method of claim 10, wherein the antibody is MEDI5117.
12. The method of claim 1, further comprising administering a diuretic.
13. The method of claim 1, wherein a single species of antibody is administered, without administration of any other antibody species.
14. The method of claim 1, wherein the anti-IL-6 antibody is MEDI5117; and wherein MEDI5117 is administered subcutaneously in a flat dose of from 10 - 20 mg.
15. Use of a therapeutically effective amount of an anti IL-6 antibody in the manufacture of a medicament for treating left ventricular heart failure in a patient.
16. The use of claim 15, wherein the patient has elevated pre-treatment plasma IL-6 levels.

17. The use of claim 15, wherein the patient has a diuretic efficiency of less than 500 mmol Na/doubling of loop diuretic dose
18. The use of claim 15, wherein the anti-IL-6 antibody comprises all six variable region CDRs of MEDI5117.
19. The use of claim 18, wherein the antibody comprises the VH and VL of MEDI5117.
20. The use of claim 19, wherein the antibody is MEDI5117.




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 3




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 6




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B




FIG. 9




FIG. 10A




FIG. 10B




FIG. 11

SEQUENCE LISTING

<110> YALE UNIVERSITY
CORVIDIA THERAPEUTICS, INC.

<120> TREATMENT OF DIURETIC RESISTANCE

<130> 32601-35789/WO

<140> PCT/US2018/016508

<141> 2018-02-01

<150> 62/453,257

<151> 2017-02-01

<160> 16

<170> PatentIn version 3.5

<210> 1

<211> 824

<212> PRT

<213> Homo sapiens

<400> 1

Met Pro Val Ala Glu Ala Pro Gln Val Ala Gly Gly Gln Gly Asp Gly
1 5 10 15

Gly Asp Gly Glu Glu Ala Glu Pro Glu Gly Met Phe Lys Ala Cys Glu
20 25 30

Asp Ser Lys Arg Lys Ala Arg Gly Tyr Leu Arg Leu Val Pro Leu Phe
35 40 45

Val Leu Leu Ala Leu Leu Val Leu Ala Ser Ala Gly Val Leu Leu Trp
50 55 60

Tyr Phe Leu Gly Tyr Lys Ala Glu Val Met Val Ser Gln Val Tyr Ser
65 70 75 80

Gly Ser Leu Arg Val Leu Asn Arg His Phe Ser Gln Asp Leu Thr Arg
85 90 95

Arg Glu Ser Ser Ala Phe Arg Ser Glu Thr Ala Lys Ala Gln Lys Met
100 105 110

Leu Lys Glu Leu Ile Thr Ser Thr Arg Leu Gly Thr Tyr Tyr Asn Ser
115 120 125

Ser Ser Val Tyr Ser Phe Gly Glu Gly Pro Leu Thr Cys Phe Phe Trp
130 135 140

Phe Ile Leu Gln Ile Pro Glu His Arg Arg Leu Met Leu Ser Pro Glu
145 150 155 160

Val Val Gln Ala Leu Leu Val Glu Glu Leu Leu Ser Thr Val Asn Ser
165 170 175

Ser Ala Ala Val Pro Tyr Arg Ala Glu Tyr Glu Val Asp Pro Glu Gly
180 185 190

Leu Val Ile Leu Glu Ala Ser Val Lys Asp Ile Ala Ala Leu Asn Ser
195 200 205

Thr Leu Gly Cys Tyr Arg Tyr Ser Tyr Val Gly Gln Gly Gln Val Leu
210 215 220

Arg Leu Lys Gly Pro Asp His Leu Ala Ser Ser Cys Leu Trp His Leu
225 230 235 240

Gln Gly Pro Lys Asp Leu Met Leu Lys Leu Arg Leu Glu Trp Thr Leu
245 250 255

Ala Glu Cys Arg Asp Arg Leu Ala Met Tyr Asp Val Ala Gly Pro Leu
260 265 270

Glu Lys Arg Leu Ile Thr Ser Val Tyr Gly Cys Ser Arg Gln Glu Pro
275 280 285

Val Val Glu Val Leu Ala Ser Gly Ala Ile Met Ala Val Val Trp Lys
290 295 300

Lys Gly Leu His Ser Tyr Tyr Asp Pro Phe Val Leu Ser Val Gln Pro
305 310 315 320

Val Val Phe Gln Ala Cys Glu Val Asn Leu Thr Leu Asp Asn Arg Leu

30 Aug 2019

					325					330					335				
Asp	Ser	Gln	Gly	Val	Leu	Ser	Thr	Pro	Tyr	Phe	Pro	Ser	Tyr	Tyr	Ser				
			340				345						350						
Pro	Gln	Thr	His	Cys	Ser	Trp	His	Leu	Thr	Val	Pro	Ser	Leu	Asp	Tyr				
			355				360						365						
Gly	Leu	Ala	Leu	Trp	Phe	Asp	Ala	Tyr	Ala	Leu	Arg	Arg	Gln	Lys	Tyr				
			370				375						380						
Asp	Leu	Pro	Cys	Thr	Gln	Gly	Gln	Trp	Thr	Ile	Gln	Asn	Arg	Arg	Leu				
			385				390						395			400			
Cys	Gly	Leu	Arg	Ile	Leu	Gln	Pro	Tyr	Ala	Glu	Arg	Ile	Pro	Val	Val				
			405							410			415						
Ala	Thr	Ala	Gly	Ile	Thr	Ile	Asn	Phe	Thr	Ser	Gln	Ile	Ser	Leu	Thr				
			420							425			430						
Gly	Pro	Gly	Val	Arg	Val	His	Tyr	Gly	Leu	Tyr	Asn	Gln	Ser	Asp	Pro				
			435				440						445						
Cys	Pro	Gly	Glu	Phe	Leu	Cys	Ser	Val	Asn	Gly	Leu	Cys	Val	Pro	Ala				
			450				455						460						
Cys	Asp	Gly	Val	Lys	Asp	Cys	Pro	Asn	Gly	Leu	Asp	Glu	Arg	Asn	Cys				
			465							475			480						
Val	Cys	Arg	Ala	Thr	Phe	Gln	Cys	Lys	Glu	Asp	Ser	Thr	Cys	Ile	Ser				
			485							490			495						
Leu	Pro	Lys	Val	Cys	Asp	Gly	Gln	Pro	Asp	Cys	Leu	Asn	Gly	Ser	Asp				
			500				505						510						
Glu	Glu	Gln	Cys	Gln	Glu	Gly	Val	Pro	Cys	Gly	Thr	Phe	Thr	Phe	Gln				
			515				520						525						
Cys	Glu	Asp	Arg	Ser	Cys	Val	Lys	Lys	Pro	Asn	Pro	Gln	Cys	Asp	Gly				
			530				535						540						

Arg 545	Pro	Asp	Cys	Arg	Asp 550	Gly	Ser	Asp	Glu	Glu 555	His	Cys	Asp	Cys	Gly 560
Leu	Gln	Gly	Pro	Ser 565	Ser	Arg	Ile	Val	Gly 570	Gly	Ala	Val	Ser	Ser 575	Glu
Gly	Glu	Trp	Pro 580	Trp	Gln	Ala	Ser	Leu 585	Gln	Val	Arg	Gly	Arg 590	His	Ile
Cys	Gly	Gly 595	Ala	Leu	Ile	Ala	Asp 600	Arg	Trp	Val	Ile	Thr 605	Ala	Ala	His
Cys 610	Phe	Gln	Glu	Asp	Ser	Met 615	Ala	Ser	Thr	Val	Leu 620	Trp	Thr	Val	Phe
Leu 625	Gly	Lys	Val	Trp	Gln 630	Asn	Ser	Arg	Trp	Pro 635	Gly	Glu	Val	Ser	Phe 640
Lys	Val	Ser	Arg	Leu 645	Leu	Leu	His	Pro	Tyr 650	His	Glu	Glu	Asp	Ser 655	His
Asp	Tyr	Asp	Val 660	Ala	Leu	Leu	Gln	Leu 665	Asp	His	Pro	Val	Val 670	Arg	Ser
Ala	Ala	Val 675	Arg	Pro	Val	Cys	Leu 680	Pro	Ala	Arg	Ser	His 685	Phe	Phe	Glu
Pro 690	Gly	Leu	His	Cys	Trp	Ile 695	Thr	Gly	Trp	Gly	Ala 700	Leu	Arg	Glu	Gly
Ala 705	Leu	Arg	Ala	Asp	Ala 710	Val	Ala	Leu	Phe	Tyr 715	Gly	Trp	Arg	Asn	Gln 720
Gly	Ser	Glu	Thr	Cys 725	Cys	Cys	Pro	Ile	Ser 730	Asn	Ala	Leu	Gln	Lys 735	Ala
Asp	Val	Gln	Leu	Ile	Pro	Gln	Asp 740	Leu 745	Cys	Ser	Glu	Val	Tyr 750	Arg	Tyr

Gln Val Thr Pro Arg Met Leu Cys Ala Gly Tyr Arg Lys Gly Lys Lys
755 760 765

Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Lys Ala Leu
770 775 780

Ser Gly Arg Trp Phe Leu Ala Gly Leu Val Ser Trp Gly Leu Gly Cys
785 790 795 800

Gly Arg Pro Asn Tyr Phe Gly Val Tyr Thr Arg Ile Thr Gly Val Ile
805 810 815

Ser Trp Ile Gln Gln Val Val Thr
820

<210> 2
<211> 824
<212> PRT
<213> Homo sapiens

<400> 2
Met Pro Val Ala Glu Ala Pro Gln Val Ala Gly Gly Gln Gly Asp Gly
1 5 10 15

Gly Asp Gly Glu Glu Ala Glu Pro Glu Gly Met Phe Lys Ala Cys Glu
20 25 30

Asp Ser Lys Arg Lys Ala Arg Gly Tyr Leu Arg Leu Val Pro Leu Phe
35 40 45

Val Leu Leu Ala Leu Leu Val Leu Ala Ser Ala Gly Val Leu Leu Trp
50 55 60

Tyr Phe Leu Gly Tyr Lys Ala Glu Val Met Val Ser Gln Val Tyr Ser
65 70 75 80

Gly Ser Leu Arg Val Leu Asn Arg His Phe Ser Gln Asp Leu Thr Arg
85 90 95

Arg Glu Ser Ser Ala Phe Arg Ser Glu Thr Ala Lys Ala Gln Lys Met
100 105 110

30 Aug 2019

Leu Lys Glu Leu Ile Thr Ser Thr Arg Leu Gly Thr Tyr Tyr Asn Ser
115 120 125

Ser Ser Val Tyr Ser Phe Gly Glu Gly Pro Leu Thr Cys Phe Phe Trp
130 135 140

Phe Ile Leu Gln Ile Pro Glu His Arg Arg Leu Met Leu Ser Pro Glu
145 150 155 160

Val Val Gln Ala Leu Leu Val Glu Glu Leu Leu Ser Thr Val Asn Ser
165 170 175

Ser Ala Ala Val Pro Tyr Arg Ala Glu Tyr Glu Val Asp Pro Glu Gly
180 185 190

Leu Val Ile Leu Glu Ala Ser Val Lys Asp Ile Ala Ala Leu Asn Ser
195 200 205

Thr Leu Gly Cys Tyr Arg Tyr Ser Tyr Val Gly Gln Gly Gln Val Leu
210 215 220

Arg Leu Lys Gly Pro Asp His Leu Ala Ser Ser Cys Leu Trp His Leu
225 230 235 240

Gln Gly Pro Lys Asp Leu Met Leu Lys Leu Arg Leu Glu Trp Thr Leu
245 250 255

Ala Glu Cys Arg Asp Arg Leu Ala Met Tyr Asp Val Ala Gly Pro Leu
260 265 270

Glu Lys Arg Leu Ile Thr Ser Val Tyr Gly Cys Ser Arg Gln Glu Pro
275 280 285

Val Val Glu Val Leu Ala Ser Gly Ala Ile Met Ala Val Val Trp Lys
290 295 300

Lys Gly Leu His Ser Tyr Tyr Asp Pro Phe Val Leu Ser Val Gln Pro
305 310 315 320

Val	Val	Phe	Gln	Ala	Cys	Glu	Val	Asn	Leu	Thr	Leu	Asp	Asn	Arg	Leu	325	330	335	
Asp	Ser	Gln	Gly	Val	Leu	Ser	Thr	Pro	Tyr	Phe	Pro	Ser	Tyr	Tyr	Ser	340	345	350	
Pro	Gln	Thr	His	Cys	Ser	Trp	His	Leu	Thr	Val	Pro	Ser	Leu	Asp	Tyr	355	360	365	
Gly	Leu	Ala	Leu	Trp	Phe	Asp	Ala	Tyr	Ala	Leu	Arg	Arg	Gln	Lys	Tyr	370	375	380	
Asp	Leu	Pro	Cys	Thr	Gln	Gly	Gln	Trp	Thr	Ile	Gln	Asn	Arg	Arg	Leu	385	390	395	400
Cys	Gly	Leu	Arg	Ile	Leu	Gln	Pro	Tyr	Ala	Glu	Arg	Ile	Pro	Val	Val	405	410	415	
Ala	Thr	Ala	Gly	Ile	Thr	Ile	Asn	Phe	Thr	Ser	Gln	Ile	Ser	Leu	Thr	420	425	430	
Gly	Pro	Gly	Val	Arg	Val	His	Tyr	Gly	Leu	Tyr	Asn	Gln	Ser	Asp	Pro	435	440	445	
Cys	Pro	Gly	Glu	Phe	Leu	Cys	Ser	Val	Asn	Gly	Leu	Cys	Val	Pro	Ala	450	455	460	
Cys	Asp	Gly	Val	Lys	Asp	Cys	Pro	Asn	Gly	Leu	Asp	Glu	Arg	Asn	Cys	465	470	475	480
Val	Cys	Arg	Ala	Thr	Phe	Gln	Cys	Lys	Glu	Asp	Ser	Thr	Cys	Ile	Ser	485	490	495	
Leu	Pro	Lys	Val	Cys	Asp	Gly	Gln	Pro	Asp	Cys	Leu	Asn	Gly	Ser	Asp	500	505	510	
Glu	Glu	Gln	Cys	Gln	Glu	Gly	Val	Pro	Cys	Gly	Thr	Phe	Thr	Phe	Gln	515	520	525	

Cys Glu Asp Arg Ser Cys Val Lys Lys Pro Asn Pro Gln Cys Asp Gly
 530 535 540
 Arg Pro Asp Cys Arg Asp Gly Ser Asp Glu Glu His Cys Asp Cys Gly
 545 550 555 560
 Leu Gln Gly Pro Ser Ser Arg Ile Val Gly Gly Ala Val Ser Ser Glu
 565 570 575
 Gly Glu Trp Pro Trp Gln Ala Ser Leu Gln Val Arg Gly Arg His Ile
 580 585 590
 Cys Gly Gly Ala Leu Ile Ala Asp Arg Trp Val Ile Thr Ala Ala His
 595 600 605
 Cys Phe Gln Glu Asp Ser Met Ala Ser Thr Val Leu Trp Thr Val Phe
 610 615 620
 Leu Gly Lys Val Trp Gln Asn Ser Arg Trp Pro Gly Glu Val Ser Phe
 625 630 635 640
 Lys Val Ser Arg Leu Leu Leu His Pro Tyr His Glu Glu Asp Ser His
 645 650 655
 Asp Tyr Asp Val Ala Leu Leu Gln Leu Asp His Pro Val Val Arg Ser
 660 665 670
 Ala Ala Val Arg Pro Val Cys Leu Pro Ala Arg Ser His Phe Phe Glu
 675 680 685
 Pro Gly Leu His Cys Trp Ile Thr Gly Trp Gly Ala Leu Arg Glu Gly
 690 695 700
 Ala Leu Arg Ala Asp Ala Val Ala Leu Phe Tyr Gly Trp Arg Asn Gln
 705 710 715 720
 Gly Ser Glu Thr Cys Cys Cys Pro Ile Ser Asn Ala Leu Gln Lys Val
 725 730 735
 Asp Val Gln Leu Ile Pro Gln Asp Leu Cys Ser Glu Val Tyr Arg Tyr

	740		745		750	
Gln Val Thr Pro Arg Met Leu Cys Ala Gly Tyr Arg Lys Gly Lys Lys	755		760		765	
Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Lys Ala Leu	770		775		780	
Ser Gly Arg Trp Phe Leu Ala Gly Leu Val Ser Trp Gly Leu Gly Cys	785		790		795	800
Gly Arg Pro Asn Tyr Phe Gly Val Tyr Thr Arg Ile Thr Gly Val Ile		805		810		815
Ser Trp Ile Gln Gln Val Val Thr	820					

<210> 3
 <211> 3196
 <212> DNA
 <213> Homo sapiens

<400> 3
 ggacaaacag aggctcctga ggcctgtgtg caggcccggc acctatctgc cgctcccaaa 60
 ggatgcccgt ggccgaggcc ccccaggtgg ctggcgggca gggggacgga ggtgatggcg 120
 aggaagcgga gccggagggg atgttcaagg cctgtgagga ctccaagaga aaagcccggg 180
 gctacctccg cctggtgccc ctgtttgtgc tgctggccct gctcgtgctg gcttcggcgg 240
 ggggtgctact ctggtatttc ctagggtaca aggcggaggt gatggtcagc caggtgtact 300
 caggcagtct gcgtgtactc aatcgccact tctcccagga tcttaccgc cggaatcta 360
 gtgccttccg cagtgaacc gccaaagccc agaagatgct caaggagctc atcaccagca 420
 cccgcctggg aacttactac aactccagct ccgtctattc ctttggggag ggacccctca 480
 cctgcttctt ctggttcatt ctccaaatcc ccgagcaccg ccggctgatg ctgagccccg 540
 aggtggtgca ggcactgctg gtggaggagc tgctgtccac agtcaacagc tcggctgccg 600
 tcccctacag ggccgagtac gaagtggacc ccgagggcct agtgatcctg gaagccagtg 660
 tgaaagacat agctgcattg aattccacgc tgggttggtta ccgctacagc tacgtgggcc 720

agggccaggt cctccggctg aaggggcctg accacctggc ctccagctgc ctgtggcacc	780
tgcagggccc caaggacctc atgctcaaac tccggctgga gtggacgctg gcagagtggc	840
gggaccgact ggccatgtat gacgtggccg ggcccctgga gaagaggctc atcacctcgg	900
tgtacggctg cagccgccag gagcccgtgg tggaggttct ggcgtcgggg gccatcatgg	960
cggtcgtctg gaagaagggc ctgcacagct actacgacct cttcgtgctc tccgtgcagc	1020
cgggtggtctt ccaggcctgt gaagtgaacc tgacgctgga caacaggctc gactcccagg	1080
gcgtcctcag caccctgtac ttccccagct actactcgcc ccaaaccac tgctcctggc	1140
acctcacggg gccctctctg gactacggct tggccctctg gtttgatgcc tatgcactga	1200
ggaggcagaa gtatgatttg ccgtgcacct agggccagtg gacgatccag aacaggaggc	1260
tgtgtggctt gcgcatcctg cagccctacg ccgagaggat ccccggtgtg gccacggccg	1320
ggatcaccat caacttcacc tcccagatct ccctcaccgg gcccggtgtg cgggtgcact	1380
atggcttgta caaccagtcg gaccctgcc ctggagagtt cctctgttct gtgaatggac	1440
tctgtgtccc tgcctgtgat ggggtcaagg actgccccaa cggcctggat gagagaaact	1500
gcgtttgcag agccacattc cagtgcaaag aggacagcac atgcatctca ctgccaagg	1560
tctgtgatgg gcagcctgat tgtctcaacg gcagcgacga agagcagtgc caggaagggg	1620
tgccatgtgg gacattcacc ttccagtgtg aggaccggag ctgcgtgaag aagcccaacc	1680
cgcagtgtga tgggcggccc gactgcaggg acggctcgga tgaggagcac tgtgactgtg	1740
gcctccaggg cccctccagc cgcatgttg gtggagctgt gtcctccgag ggtgagtggc	1800
catggcaggc cagcctccag gttcggggtc gacacatctg tggggggggc ctcacgctg	1860
accgctgggt gataacagct gccactgct tccaggagga cagcatggcc tccacggtgc	1920
tgtggaccgt gttcctgggc aaggtgtggc agaactcgcg ctggcctgga gaggtgtcct	1980
tcaaggtgag ccgcctgctc ctgcacctgt accacgaaga ggacagccat gactacgacg	2040
tggcgctgct gcagctcgac caccgggtgg tgcgctcggc cgccgtgcgc cccgtctgcc	2100
tgcccgcgcg ctcccacttc ttcgagcccg gcctgcactg ctggattacg ggctggggcg	2160
ccttgcgca gggcgcccta cgggcggatg ctgtggccct attttatgga tggagaaacc	2220
aaggctcaga gacatgttgc tgcccatca gcaacgctct gcagaaagtg gatgtgcagt	2280
tgatcccaca ggacctgtgc agcgaggtct atcgctacca ggtgacgcca cgcatgctgt	2340

gtgccggcta	ccgcaagggc	aagaaggatg	cctgtcaggg	tgactcaggt	ggtccgctgg	2400
tgtgcaaggc	actcagtggc	cgctggttcc	tggcggggct	ggtcagctgg	ggcctgggct	2460
gtggccggcc	taactacttc	ggcgtctaca	cccgcacac	aggtgtgatc	agctggatcc	2520
agcaagtggg	gacctgagga	actgcccccc	tgcaaagcag	ggcccacctc	ctggactcag	2580
agagcccagg	gcaactgcca	agcaggggga	caagtattct	ggcggggggg	gggggagaga	2640
gcaggccctg	tggtggcagg	aggtggcatc	ttgtctcgtc	cctgatgtct	gctccagtga	2700
tggcaggagg	atggagaagt	gccagcagct	gggggtcaag	acgtcccctg	aggacccagg	2760
cccacaccca	gcccttctgc	ctcccaattc	tctctcctcc	gtccccttcc	tccactgctg	2820
cctaattgcaa	ggcagtggct	cagcagcaag	aatgctgggt	ctacatcccg	aggagtgtct	2880
gaggtgcgcc	ccactctgta	cagaggctgt	ttgggcagcc	ttgcctccag	agagcagatt	2940
ccagcttcgg	aagcccctgg	tctaacttgg	gatctgggaa	tggaaggtgc	tcccatcgga	3000
ggggaccctc	agagcccctg	agactgccag	gtgggcctgc	tgccactgta	agccaaaagg	3060
tggggaagtc	ctgactccag	ggtccttgcc	ccaccctgc	ctgccacctg	ggccctcaca	3120
gcccagaccc	tcactgggag	gtgagctcag	ctgccctttg	gaataaagct	gcctgatcca	3180
aaaaaaaaaa	aaaaaa					3196

<210> 4
 <211> 3196
 <212> DNA
 <213> Homo sapiens

<400> 4						
ggacaaacag	aggctcctga	ggcctgtgtg	caggcccggc	acctatctgc	cgctcccaaa	60
ggatgcccgt	ggccgaggcc	ccccagggtg	ctggcgggca	gggggacgga	ggtgatggcg	120
aggaagcgga	gccggagggg	atgttcaagg	cctgtgagga	ctccaagaga	aaagcccggg	180
gctacctccg	cctggtgccc	ctgtttgtgc	tgctggccct	gctcgtgctg	gcttcggcgg	240
gggtgctact	ctggtatttc	ctagggtaca	aggcggaggt	gatggtcagc	caggtgtact	300
caggcagtct	gcgtgtactc	aatcgccact	tctcccagga	tcttaccgcg	cgggaatcta	360
gtgccttcgg	cagtgaacc	gcaaagccc	agaagatgct	caaggagctc	atcaccagca	420
cccgcctggg	aacttactac	aactccagct	ccgtctattc	ctttggggag	ggaccctca	480

cctgcttctt ctggttcatt ctccaaatcc ccgagcaccg ccggctgatg ctgagccccg	540
aggtgggtgca ggcactgctg gtggaggagc tgctgtccac agtcaacagc tcggctgccg	600
tcccctacag ggccgagtac gaagtggacc ccgagggcct agtgatcctg gaagccagtg	660
tgaaagacat agctgcattg aattccacgc tgggttggtta ccgctacagc tacgtgggcc	720
agggccaggt cctccggctg aaggggcctg accacctggc ctccagctgc ctgtggcacc	780
tgcagggccc caaggacctc atgctcaaac tccggctgga gtggacgctg gcagagtgcc	840
gggaccgact ggccatgtat gacgtggccg ggcccctgga gaagaggctc atcacctcgg	900
tgtacggctg cagccgccag gagcccgtgg tggaggttct ggcgtcgggg gccatcatgg	960
cggtcgtctg gaagaagggc ctgcacagct actacgacc cttcgtgctc tccgtgcagc	1020
cgggtggtctt ccaggcctgt gaagtgaacc tgacgtgga caacaggctc gactcccagg	1080
gcgtcctcag caccctgtac ttcccagct actactgcc ccaaaccac tgctcctggc	1140
acctcacggg gccctctctg gactacggct tggccctctg gtttgatgcc tatgcactga	1200
ggaggcagaa gtatgatttg ccgtgcaccc agggccagtg gacgatccag aacaggaggc	1260
tgtgtggctt gcgcatcctg cagccctacg ccgagaggat ccccggtgtg gccacggccg	1320
ggatcaccat caacttcacc tcccagatct ccctcaccgg gcccggtgtg cgggtgcact	1380
atggccttga caaccagtcg gaccctgcc ctggagagtt cctctgttct gtgaatggac	1440
tctgtgtccc tgcctgtgat ggggtcaagg actgccccaa cggcctggat gagagaaact	1500
gcgtttgcag agccacattc cagtgc aaag aggacagcac atgcatctca ctgccc aagg	1560
tctgtgatgg gcagcctgat tgtctcaacg gcagcgacga agagcagtgc caggaagggg	1620
tgccatgtgg gacattcacc ttccagtgtg aggaccggag ctgcgtgaag aagcccaacc	1680
cgcagtgtga tgggcggccc gactgcaggg acggctcgga tgaggagcac tgtgactgtg	1740
gcctccaggg cccctccagc cgcatgttg gtggagctgt gtcctccgag ggtgagtggc	1800
catggcaggc cagcctccag gttcggggtc gacacatctg tggggggggc ctcacgctg	1860
accgctgggt gataacagct gccactgct tccaggagga cagcatggcc tccacggtgc	1920
tgtggaccgt gttcctgggc aagggtgtggc agaactcgcg ctggcctgga gaggtgtcct	1980
tcaaggtgag ccgcctgctc ctgcaccgt accacgaaga ggacagccat gactacgacg	2040

tggcgctgct gcagctcgac cacccggtgg tgcgctcggc cgccgtgcgc cccgtctgcc	2100
tgcccgcgcg ctcccacttc ttcgagcccg gcctgcactg ctggattacg ggctggggcg	2160
ccttgcgcgga gggcgcccta cgggcgggatg ctgtggccct attttatgga tggagaaacc	2220
aaggctcaga gacatgttgc tgcccatca gcaacgctct gcagaaagtg gatgtgcagt	2280
tgatcccaca ggacctgtgc agcgaggtct atcgctacca agtgacgcca cgcattgtgt	2340
gtgccggcta ccgcaagggc aagaaggatg cctgtcaggg tgactcaggt ggtccgctgg	2400
tgtgcaaggc actcagtggc cgctggttcc tggcggggct ggtcagctgg ggcctgggct	2460
gtggccggcc taactacttc ggcgtctaca cccgcatcac aggtgtgatc agctggatcc	2520
agcaagtggg gacctgagga actgcccccc tgcaaagcag ggcccacctc ctggactcag	2580
agagcccagg gcaactgcca agcaggggga caagtattct ggcgggggggt gggggagaga	2640
gcaggccctg tgggtggcagg aggtggcatc ttgtctcgtc cctgatgtct gctccagtga	2700
tggcaggagg atggagaagt gccagcagct ggggggtcaag acgtcccctg aggacccagg	2760
cccacacca gcccttctgc ctccaattc tctctcctcc gtccccttcc tccactgctg	2820
cctaattgcaa ggcagtggct cagcagcaag aatgctgggt ctacatcccg aggagtgtct	2880
gaggtgcgcc ccactctgta cagaggctgt ttgggcagcc ttgcctccag agagcagatt	2940
ccagcttcgg aagcccctgg tctaacttgg gatctgggaa tggaagggtgc tcccatcgga	3000
ggggaccctc agagccctgg agactgccag gtgggcctgc tgccactgta agccaaaagg	3060
tgggggaagtc ctgactccag ggtccttgcc ccaccctgc ctgccacctg ggccctcaca	3120
gcccagaccc tctactgggag gtgagctcag ctgccctttg gaataaagct gcctgatcca	3180
aaaaaaaaaa aaaaaa	3196

<210> 5
 <211> 274
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Cys Val Gly Ala Arg Arg Leu Gly Arg Gly Pro Cys Ala Ala Leu
 1 5 10 15

Leu Leu Leu Gly Leu Gly Leu Ser Thr Val Thr Gly Leu His Cys Val
 20 25 30

30 Aug 2019

Gly Asp Thr Tyr Pro Ser Asn Asp Arg Cys Cys His Glu Cys Arg Pro
35 40 45

Gly Asn Gly Met Val Ser Arg Cys Ser Arg Ser Gln Asn Thr Val Cys
50 55 60

Arg Pro Cys Gly Pro Gly Phe Tyr Asn Asp Val Val Ser Ser Lys Pro
65 70 75 80

Cys Lys Pro Cys Thr Trp Cys Asn Leu Arg Ser Gly Ser Glu Arg Lys
85 90 95

Gln Leu Cys Thr Ala Thr Gln Asp Thr Val Cys Arg Cys Arg Ala Gly
100 105 110

Thr Gln Pro Leu Asp Ser Tyr Lys Pro Gly Val Asp Cys Ala Pro Cys
115 120 125

Pro 130 Gly 135 Asp 140 Asn Gln Ala Cys Lys Pro Trp

Thr Asn Cys Thr Leu Ala Gly Lys His Thr Leu Gln Pro Ala Ser Asn
145 150 155 160

Ser Ser Asp Ala Ile Cys Glu Asp Arg Asp Pro Pro Ala Thr Gln Pro
165 170 175

Gln Glu Thr Gln Gly Pro Pro Ala Arg Pro Ile Thr Val Gln Pro Thr
180 185 190

Glu Ala Trp Pro Arg Thr Ser Gln Gly Pro Ser Thr Arg Pro Val Glu
195 200 205

Val Pro Gly Gly Arg Ala Val Ala Ala Ile Leu Gly Leu Gly Leu Val
210 215 220

Leu Gly Leu Leu Gly Pro Leu Ala Ile Leu Leu Ala Leu Tyr Leu Leu
225 230 235 240

Arg Arg Asp Gln Arg Leu Pro Pro Asp Ala His Lys Pro Pro Ser Phe
245 250 255

Arg Thr Pro Ile Gln Glu Glu Gln Ala Asp Ala His Ser Thr Leu Ala
260 265 270

Lys Ile

<210> 6
<211> 1201
<212> DNA
<213> Homo sapiens

<400> 6
aatattagag tctcaacccc caataaatat aggactggag atgtctgagg ctcatcttgc 60
cctcgagccc accgggaacg aaagagaagc tctatctccc ctccaggagc ccagctatga 120
actccttctc cacaagcgcc ttcgggtccag ttgccttctc cctggggctg ctcttggtgt 180
tgcctgctgc cttccctgcc ccagtacccc caggagaaga ttccaaagat gtagccgccc 240
cacacagaca gccactcacc tcttcagaac gaattgacaa acaaattcgg tacatcctcg 300
acggcatctc agccctgaga aaggagacat gtaacaagag taacatgtgt gaaagcagca 360
aagaggcact ggcagaaaac aacctgaacc ttccaaagat ggctgaaaaa gatggatgct 420
tccaatctgg attcaatgag gagacttgcc tggtgaaaat catcactggt cttttggagt 480
ttgaggtata cctagagtac ctccagaaca gatttgagag tagtgaggaa caagccagag 540
ctgtgcagat gagtacaaaa gtcctgatcc agttcctgca gaaaaaggca aagaatctag 600
atgcaataac caccctgac ccaaccacaa atgccagcct gctgacgaag ctgcaggcac 660
agaaccagtg gctgcaggac atgacaactc atctcattct gcgcagcttt aaggagttcc 720
tgcagtccag cctgagggct cttcggcaaa tgtagcatgg gcacctcaga ttgtttgtgt 780
taatgggcat tccttcttct ggtcagaaac ctgtccactg ggcacagaac ttatgtttgt 840
ctctatggag aactaaaagt atgagcgtaa ggacactatt ttaattattt ttaatttatt 900
aatatttaaa tatgtgaagc tgagttaatt tatgtaagtc atatttatat ttttaagaag 960
taccacttga aacattttat gtattagttt tgaaataata atggaaagtg gctatgcagt 1020

ttgaatatcc tttgtttcag agccagatca tttcttggaa agtgtaggct tacctcaaat 1080
 aaatggctaa cttatacata tttttaaaga aatatttata ttgtatttat ataatgtata 1140
 aatggttttt ataccaataa atggcatttt aaaaaattca gcaaaaaaaaa aaaaaaaaaa 1200
 a 1201

<210> 7
 <211> 468
 <212> PRT
 <213> Homo sapiens

<400> 7
 Met Leu Ala Val Gly Cys Ala Leu Leu Ala Ala Leu Leu Ala Ala Pro
 1 5 10 15

Gly Ala Ala Leu Ala Pro Arg Arg Cys Pro Ala Gln Glu Val Ala Arg
 20 25 30

Gly Val Leu Thr Ser Leu Pro Gly Asp Ser Val Thr Leu Thr Cys Pro
 35 40 45

Gly Val Glu Pro Glu Asp Asn Ala Thr Val His Trp Val Leu Arg Lys
 50 55 60

Pro Ala Ala Gly Ser His Pro Ser Arg Trp Ala Gly Met Gly Arg Arg
 65 70 75 80

Leu Leu Leu Arg Ser Val Gln Leu His Asp Ser Gly Asn Tyr Ser Cys
 85 90 95

Tyr Arg Ala Gly Arg Pro Ala Gly Thr Val His Leu Leu Val Asp Val
 100 105 110

Pro Pro Glu Glu Pro Gln Leu Ser Cys Phe Arg Lys Ser Pro Leu Ser
 115 120 125

Asn Val Val Cys Glu Trp Gly Pro Arg Ser Thr Pro Ser Leu Thr Thr
 130 135 140

Lys Ala Val Leu Leu Val Arg Lys Phe Gln Asn Ser Pro Ala Glu Asp
 145 150 155 160

Phe	Gln	Glu	Pro	Cys	Gln	Tyr	Ser	Gln	Glu	Ser	Gln	Lys	Phe	Ser	Cys	165	170	175	
Gln	Leu	Ala	Val	Pro	Glu	Gly	Asp	Ser	Ser	Phe	Tyr	Ile	Val	Ser	Met	180	185	190	
Cys	Val	Ala	Ser	Ser	Val	Gly	Ser	Lys	Phe	Ser	Lys	Thr	Gln	Thr	Phe	195	200	205	
Gln	Gly	Cys	Gly	Ile	Leu	Gln	Pro	Asp	Pro	Pro	Ala	Asn	Ile	Thr	Val	210	215	220	
Thr	Ala	Val	Ala	Arg	Asn	Pro	Arg	Trp	Leu	Ser	Val	Thr	Trp	Gln	Asp	225	230	235	240
Pro	His	Ser	Trp	Asn	Ser	Ser	Phe	Tyr	Arg	Leu	Arg	Phe	Glu	Leu	Arg	245	250	255	
Tyr	Arg	Ala	Glu	Arg	Ser	Lys	Thr	Phe	Thr	Thr	Trp	Met	Val	Lys	Asp	260	265	270	
Leu	Gln	His	His	Cys	Val	Ile	His	Asp	Ala	Trp	Ser	Gly	Leu	Arg	His	275	280	285	
Val	Val	Gln	Leu	Arg	Ala	Gln	Glu	Glu	Phe	Gly	Gln	Gly	Glu	Trp	Ser	290	295	300	
Glu	Trp	Ser	Pro	Glu	Ala	Met	Gly	Thr	Pro	Trp	Thr	Glu	Ser	Arg	Ser	305	310	315	320
Pro	Pro	Ala	Glu	Asn	Glu	Val	Ser	Thr	Pro	Met	Gln	Ala	Leu	Thr	Thr	325	330	335	
Asn	Lys	Asp	Asp	Asp	Asn	Ile	Leu	Phe	Arg	Asp	Ser	Ala	Asn	Ala	Thr	340	345	350	
Ser	Leu	Pro	Val	Gln	Asp	Ser	Ser	Ser	Val	Pro	Leu	Pro	Thr	Phe	Leu	355	360	365	

Val Ala Gly Gly Ser Leu Ala Phe Gly Thr Leu Leu Cys Ile Ala Ile
370 375 380

Val Leu Arg Phe Lys Lys Thr Trp Lys Leu Arg Ala Leu Lys Glu Gly
385 390 395 400

Lys Thr Ser Met His Pro Pro Tyr Ser Leu Gly Gln Leu Val Pro Glu
405 410 415

Arg Pro Arg Pro Thr Pro Val Leu Val Pro Leu Ile Ser Pro Pro Val
420 425 430

Ser Pro Ser Ser Leu Gly Ser Asp Asn Thr Ser Ser His Asn Arg Pro
435 440 445

Asp Ala Arg Asp Pro Arg Ser Pro Tyr Asp Ile Ser Asn Thr Asp Tyr
450 455 460

Phe Phe Pro Arg
465

<210> 8
<211> 918
<212> PRT
<213> Homo sapiens

<400> 8
Met Leu Thr Leu Gln Thr Trp Leu Val Gln Ala Leu Phe Ile Phe Leu
1 5 10 15

Thr Thr Glu Ser Thr Gly Glu Leu Leu Asp Pro Cys Gly Tyr Ile Ser
20 25 30

Pro Glu Ser Pro Val Val Gln Leu His Ser Asn Phe Thr Ala Val Cys
35 40 45

Val Leu Lys Glu Lys Cys Met Asp Tyr Phe His Val Asn Ala Asn Tyr
50 55 60

Ile Val Trp Lys Thr Asn His Phe Thr Ile Pro Lys Glu Gln Tyr Thr
65 70 75 80

Ile	Ile	Asn	Arg	Thr	Ala	Ser	Ser	Val	Thr	Phe	Thr	Asp	Ile	Ala	Ser	
				85					90					95		
Leu	Asn	Ile	Gln	Leu	Thr	Cys	Asn	Ile	Leu	Thr	Phe	Gly	Gln	Leu	Glu	
			100					105					110			
Gln	Asn	Val	Tyr	Gly	Ile	Thr	Ile	Ile	Ser	Gly	Leu	Pro	Pro	Glu	Lys	
		115					120					125				
Pro	Lys	Asn	Leu	Ser	Cys	Ile	Val	Asn	Glu	Gly	Lys	Lys	Met	Arg	Cys	
	130					135					140					
Glu	Trp	Asp	Gly	Gly	Arg	Glu	Thr	His	Leu	Glu	Thr	Asn	Phe	Thr	Leu	
145					150					155					160	
Lys	Ser	Glu	Trp	Ala	Thr	His	Lys	Phe	Ala	Asp	Cys	Lys	Ala	Lys	Arg	
				165					170					175		
Asp	Thr	Pro	Thr	Ser	Cys	Thr	Val	Asp	Tyr	Ser	Thr	Val	Tyr	Phe	Val	
			180					185					190			
Asn	Ile	Glu	Val	Trp	Val	Glu	Ala	Glu	Asn	Ala	Leu	Gly	Lys	Val	Thr	
		195					200					205				
Ser	Asp	His	Ile	Asn	Phe	Asp	Pro	Val	Tyr	Lys	Val	Lys	Pro	Asn	Pro	
	210					215					220					
Pro	His	Asn	Leu	Ser	Val	Ile	Asn	Ser	Glu	Glu	Leu	Ser	Ser	Ile	Leu	
225					230					235					240	
Lys	Leu	Thr	Trp	Thr	Asn	Pro	Ser	Ile	Lys	Ser	Val	Ile	Ile	Leu	Lys	
				245					250					255		
Tyr	Asn	Ile	Gln	Tyr	Arg	Thr	Lys	Asp	Ala	Ser	Thr	Trp	Ser	Gln	Ile	
			260					265					270			
Pro	Pro	Glu	Asp	Thr	Ala	Ser	Thr	Arg	Ser	Ser	Phe	Thr	Val	Gln	Asp	
		275					280					285				

Leu Lys Pro Phe Thr Glu Tyr Val Phe Arg Ile Arg Cys Met Lys Glu
 290 295 300
 Asp Gly Lys Gly Tyr Trp Ser Asp Trp Ser Glu Glu Ala Ser Gly Ile
 305 310 315 320
 Thr Tyr Glu Asp Arg Pro Ser Lys Ala Pro Ser Phe Trp Tyr Lys Ile
 325 330 335
 Asp Pro Ser His Thr Gln Gly Tyr Arg Thr Val Gln Leu Val Trp Lys
 340 345 350
 Thr Leu Pro Pro Phe Glu Ala Asn Gly Lys Ile Leu Asp Tyr Glu Val
 355 360 365
 Thr Leu Thr Arg Trp Lys Ser His Leu Gln Asn Tyr Thr Val Asn Ala
 370 375 380
 Thr Lys Leu Thr Val Asn Leu Thr Asn Asp Arg Tyr Leu Ala Thr Leu
 385 390 395 400
 Thr Val Arg Asn Leu Val Gly Lys Ser Asp Ala Ala Val Leu Thr Ile
 405 410 415
 Pro Ala Cys Asp Phe Gln Ala Thr His Pro Val Met Asp Leu Lys Ala
 420 425 430
 Phe Pro Lys Asp Asn Met Leu Trp Val Glu Trp Thr Thr Pro Arg Glu
 435 440 445
 Ser Val Lys Lys Tyr Ile Leu Glu Trp Cys Val Leu Ser Asp Lys Ala
 450 455 460
 Pro Cys Ile Thr Asp Trp Gln Gln Glu Asp Gly Thr Val His Arg Thr
 465 470 475 480
 Tyr Leu Arg Gly Asn Leu Ala Glu Ser Lys Cys Tyr Leu Ile Thr Val
 485 490 495

Thr	Pro	Val	Tyr	Ala	Asp	Gly	Pro	Gly	Ser	Pro	Glu	Ser	Ile	Lys	Ala	
			500					505					510			
Tyr	Leu	Lys	Gln	Ala	Pro	Pro	Ser	Lys	Gly	Pro	Thr	Val	Arg	Thr	Lys	
		515					520					525				
Lys	Val	Gly	Lys	Asn	Glu	Ala	Val	Leu	Glu	Trp	Asp	Gln	Leu	Pro	Val	
	530					535					540					
Asp	Val	Gln	Asn	Gly	Phe	Ile	Arg	Asn	Tyr	Thr	Ile	Phe	Tyr	Arg	Thr	
545					550					555					560	
Ile	Ile	Gly	Asn	Glu	Thr	Ala	Val	Asn	Val	Asp	Ser	Ser	His	Thr	Glu	
				565					570					575		
Tyr	Thr	Leu	Ser	Ser	Leu	Thr	Ser	Asp	Thr	Leu	Tyr	Met	Val	Arg	Met	
			580					585					590			
Ala	Ala	Tyr	Thr	Asp	Glu	Gly	Gly	Lys	Asp	Gly	Pro	Glu	Phe	Thr	Phe	
		595					600					605				
Thr	Thr	Pro	Lys	Phe	Ala	Gln	Gly	Glu	Ile	Glu	Ala	Ile	Val	Val	Pro	
	610					615					620					
Val	Cys	Leu	Ala	Phe	Leu	Leu	Thr	Thr	Leu	Leu	Gly	Val	Leu	Phe	Cys	
625					630					635					640	
Phe	Asn	Lys	Arg	Asp	Leu	Ile	Lys	Lys	His	Ile	Trp	Pro	Asn	Val	Pro	
				645					650					655		
Asp	Pro	Ser	Lys	Ser	His	Ile	Ala	Gln	Trp	Ser	Pro	His	Thr	Pro	Pro	
			660					665					670			
Arg	His	Asn	Phe	Asn	Ser	Lys	Asp	Gln	Met	Tyr	Ser	Asp	Gly	Asn	Phe	
		675					680					685				
Thr	Asp	Val	Ser	Val	Val	Glu	Ile	Glu	Ala	Asn	Asp	Lys	Lys	Pro	Phe	
	690					695					700					
Pro	Glu	Asp	Leu	Lys	Ser	Leu	Asp	Leu	Phe	Lys	Lys	Glu	Lys	Ile	Asn	

2018214554 30 Aug 2019

705					710						715					720
Thr	Glu	Gly	His	Ser	Ser	Gly	Ile	Gly	Gly	Ser	Ser	Cys	Met	Ser	Ser	
				725					730					735		
Ser	Arg	Pro	Ser	Ile	Ser	Ser	Ser	Asp	Glu	Asn	Glu	Ser	Ser	Gln	Asn	
			740					745					750			
Thr	Ser	Ser	Thr	Val	Gln	Tyr	Ser	Thr	Val	Val	His	Ser	Gly	Tyr	Arg	
		755					760					765				
His	Gln	Val	Pro	Ser	Val	Gln	Val	Phe	Ser	Arg	Ser	Glu	Ser	Thr	Gln	
	770					775					780					
Pro	Leu	Leu	Asp	Ser	Glu	Glu	Arg	Pro	Glu	Asp	Leu	Gln	Leu	Val	Asp	
785					790					795					800	
His	Val	Asp	Gly	Gly	Asp	Gly	Ile	Leu	Pro	Arg	Gln	Gln	Tyr	Phe	Lys	
				805					810					815		
Gln	Asn	Cys	Ser	Gln	His	Glu	Ser	Ser	Pro	Asp	Ile	Ser	His	Phe	Glu	
			820					825					830			
Arg	Ser	Lys	Gln	Val	Ser	Ser	Val	Asn	Glu	Glu	Asp	Phe	Val	Arg	Leu	
		835					840					845				
Lys	Gln	Gln	Ile	Ser	Asp	His	Ile	Ser	Gln	Ser	Cys	Gly	Ser	Gly	Gln	
	850					855					860					
Met	Lys	Met	Phe	Gln	Glu	Val	Ser	Ala	Ala	Asp	Ala	Phe	Gly	Pro	Gly	
865					870					875					880	
Thr	Glu	Gly	Gln	Val	Glu	Arg	Phe	Glu	Thr	Val	Gly	Met	Glu	Ala	Ala	
			885						890					895		
Thr	Asp	Glu	Gly	Met	Pro	Lys	Ser	Tyr	Leu	Pro	Gln	Thr	Val	Arg	Gln	
			900					905					910			
Gly	Gly	Tyr	Met	Pro	Gln											
			915													

<210> 9
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 9
 Ser Asn Tyr Met Ile
 1 5

<210> 10
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 10
 Asp Leu Tyr Tyr Tyr Ala Gly Asp Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 11
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 11
 Trp Ala Asp Asp His Pro Pro Trp Ile Asp Leu
 1 5 10

<210> 12
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 12

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
1 5 10

<210> 13

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 13

Lys Ala Ser Thr Leu Glu Ser
1 5

<210> 14

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 14

Gln Gln Ser Trp Leu Gly Gly Ser
1 5

<210> 15

<211> 450

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 15

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Ile Ser Ser Asn
20 25 30

Tyr	Met	Ile	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
		35					40					45				
Ser	Asp	Leu	Tyr	Tyr	Tyr	Ala	Gly	Asp	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	
	50					55					60					
Lys	Gly	Arg	Phe	Thr	Met	Ser	Arg	Asp	Ile	Ser	Lys	Asn	Thr	Val	Tyr	
65					70					75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
				85					90					95		
Ala	Arg	Trp	Ala	Asp	Asp	His	Pro	Pro	Trp	Ile	Asp	Leu	Trp	Gly	Arg	
			100					105					110			
Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	
		115					120					125				
Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	
	130					135					140					
Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	
145					150					155					160	
Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	
				165					170					175		
Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	
			180					185					190			
Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	
		195					200					205				
Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	
	210					215					220					
Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	
225					230					235					240	

Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Tyr	Ile	245	250	255	
Thr	Arg	Glu	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	260	265	270	
Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	275	280	285	
Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	290	295	300	
Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	305	310	315	320
Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	325	330	335	
Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	340	345	350	
Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	355	360	365	
Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	370	375	380	
Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	385	390	395	400
Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	405	410	415	
Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	420	425	430	
Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	435	440	445	

Gly Lys
450

<210> 16
<211> 213
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 16
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Val Leu Ile
35 40 45

Tyr Lys Ala Ser Thr Leu Glu Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Asp Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Trp Leu Gly Gly Ser
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130 135 140

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145 150 155 160

Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
165 170 175

Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
180 185 190

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
195 200 205

Asn Arg Gly Glu Cys
210