



(51) International Patent Classification:

A61K 47/48 (2006.01) A61P 29/00 (2006.01)
A61K 49/00 (2006.01) A61K 9/127 (2006.01)
A61K 49/18 (2006.01) C07D 405/12 (2006.01)
A61P 35/00 (2006.01) C07K 7/02 (2006.01)

(21) International Application Number:

PCT/EP2012/002652

(22) International Filing Date:

22 June 2012 (22.06.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11005110.9 22 June 2011 (22.06.2011) EP

(71) Applicant (for all designated States except US): J. STEFAN INSTITUTE [SI/SI]; Jamova 39, 1000 Ljubljana (SI).

(72) Inventor; and

(71) Applicant : SCHASCHKE, Norbert [DE/DE]; Hügelstrasse 1, 33613 Bielefeld (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): VASILJEVA, Olga [SI/SI]; Preloška cesta 16, Prelog, 1230 Domzale (SI). MIKHAYLOV, Georgy [KZ/KZ]; ul Aleksandra Nevskogo 11a, 050040 Almaty (KZ). TURK, Boris [SI/SI]; Gradišče IV 069, 1291 Škofljica (SI).

(74) Agent: BÖSL, Raphael; Isenbruck Bösl Hörschler LLP, Prinzregentenstrasse 68, 81675 München (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: CATHEPSIN-BINDING COMPOUNDS BOUND TO A CARRIER AND THEIR DIAGNOSTIC USE

(57) Abstract: The invention relates to Cathepsin-binding compounds bound to a carrier comprising a diagnostic moiety, for use in the diagnosis of inflammatory diseases, and/or for use in the diagnosis of neoplastic diseases, wherein the Cathepsin-binding compound binds to inflammatory cells of the tumour stroma. The invention also relates to Cathepsin B- targeting compounds and Cathepsin B-binding and liposome-binding compounds.



WO 2012/175223 A1

Cathepsin-binding Compounds bound to a Carrier and their Diagnostic Use

The invention relates to Cathepsin-binding compounds bound to a carrier comprising a
5 therapeutic and/or diagnostic moiety, for use in the diagnosis and/or treatment of
inflammatory diseases, and/or for use in the diagnosis and/or treatment of neoplastic
diseases, wherein the Cathepsin-binding compound binds to inflammatory cells of the
tumour stroma. The invention also relates to Cathepsin B-targeting compounds and
Cathepsin B-binding and liposome-binding compounds.

10

Background

Inflammation is an essential localized host response to invading microorganisms or tissue
injury which involves cells of the immune system. The classic signs of inflammation
include redness (erythema), swelling (edema), pain and increased heat production (pyrema)
15 at the site of injury. The inflammatory response allows the body to specifically recognize
and eliminate an invading organism and/or repair tissue injury. Inflammatory disease
occurs when an inflammatory response is initiated that is inappropriate and/or does not
resolve in the normal manner but rather persists and results in a chronic inflammatory
state. Inflammatory disease may be systemic (e.g. lupus) or localized to particular tissues
20 or organs and exerts an enormous personal and economic burden to the society. Examples
of some of the most common and problematic inflammatory diseases are rheumatoid
arthritis, inflammatory bowel disease, psoriasis, asthma, chronic obstructive pulmonary
disease, emphysema, colitis and ischemia-reperfusion injury.

25 Proteases, and particularly lysosomal cysteine proteases, have recently been considered as
pharmacological targets for immune disorders and inflammatory diseases, including
rheumatoid arthritis, atherosclerosis, and myasthenia gravis (Link et al., 2006; Riese, R. J.
et al., Immunity 1996, 4, 357; Thurmond et al. (1), 2004; Vasiljeva et al., 2007). An
inflammatory cell infiltrate is an important component of the tumour microenvironment,
30 and many clinical studies have established a positive correlation between inflammation and
tumour progression (Mantovani et al., 2008). Also, stromal cells that infiltrate the tumour
mass can secrete a variety of tumour-promoting factors including growth factors,
chemokines, and proteases (Murdoch et al., 2008).

35 Proteases can be very harmful if not strictly controlled. There is a growing body of
information that cysteine cathepsins are involved in numerous pathologies, with at least

some of them representing suitable targets for therapy. One of the areas, where cysteine cathepsins have gathered a lot of attraction recently is cancer biology, especially after the failure of the MMP inhibitors in clinical trials (Coussens et al., 2002; Turk et al., 2004). The other emerging areas include viral infections, cardiovascular diseases and osteoarthritis, although the major focus is still on osteoporosis, rheumatoid arthritis and other diseases linked with the immune system.

There is increasing evidence that cysteine proteases, mostly cathepsins B and L, and to a lesser extent cathepsins H, S, X, and K, contribute to the proteolytic events during tumour progression. Cysteine cathepsins upregulation has been reported for many human tumours, including breast, lung, brain, gastrointestinal, prostate cancer, and melanoma (Jedesko et al., 2004). The expression of lysosomal proteases, e.g. cathepsins B and L, have often been positively correlated with a poor prognosis for patients with a variety of malignancies, and antigen levels of cysteine cathepsins were measured as potential prognostic markers for several types of cancer.

Cysteine proteases have been implicated in the progression of tumours from a premalignant to a malignant state and in various critical tumourbiological processes, including tumour cell hyperproliferation and apoptosis; tumour induced angiogenesis, as well as invasion of surrounding tissues and metastasis by malignant cells, suggesting that they are relevant drug targets for treating cancer (Turk et al., 2004).

Rheumatoid arthritis is a chronic inflammatory joint disorder involving autoimmune processes that result in destruction of joint cartilage eventually leading to loss of joint function. In RA, the synovial membrane proliferates to form a pannus that destroys adjacent bone and cartilage. Among the cells mainly responsible for cartilage damage are fibroblast-like synoviocytes (synovial fibroblasts) and activated macrophages that accumulate in the interface between pannus and cartilage.

Cathepsin B and, to a lesser extent, cathepsin L were found in the synovial fluids (Turk et al., 1988; Huet et al., 1993) and synovial lining tissues of patients with rheumatoid arthritis (Esser et al., 1994). Both enzymes have been detected at high levels in membranes of patients with rheumatoid arthritis, even at very early stages, while their levels were very low in normal synovium (Hansen et al., 2000). Keyszer and co-authors proposed post-transcriptional up-regulation of these enzymes in rheumatoid arthritis (Keyszer et al., 1998). In addition, selective secretion of cathepsins B and L by synovial fibroblast-like cells upon cytokine induction was observed (Lemaire et al., 1997). In experimental animal models increased levels of cathepsin B have been detected in joint tissues during the course

of experimental arthritis (Esser et al., 1994; Etherington et al., 1988), while elevated levels of cathepsin L have been detected in the synovial lining of rabbits with antigen-induced arthritis (Etherington et al., 1988). Furthermore, involvement of cathepsins B and L in bone degradation was demonstrated by inhibition of bone resorption by selective
5 inactivators of cysteine proteases (Hill et al., 1994). Using retroviral gene transfer, ribozymes cathepsin L mRNA, specifically inhibiting the synthesis of cathepsin L, reduced cartilage destruction *in vitro* and *in vivo* using a mouse co-implantation model of rheumatoid arthritis (Schedel et al., 2004).

10 Innate immune cells can constitute a substantial proportion of the cells within the tumour microenvironment and have been associated with tumour malignancy in patients and animal models of cancer (Mantovani et al., 2008; Murdoch et al., 2008); however, the mechanisms by which they modulate tumour progression are incompletely understood. It was shown that high levels of Cathepsin protease activity are induced in the majority of
15 macrophages in the microenvironment of pancreatic islet cancers, mammary tumours, and lung metastases during malignant progression (Gocheva et al., 2010). It was also shown that tumour-associated macrophage (TAM)-supplied Cathepsins B and S are critical for promoting pancreatic tumour growth, angiogenesis, and invasion *in vivo*, and markedly enhance the invasiveness of cancer cells in culture. It was demonstrated that interleukin-4
20 (IL-4) is responsible for inducing Cathepsin activity in macrophages *in vitro* and *in vivo*, hence being an important regulator, and Cathepsin proteases as critical mediators, of the tumour-promoting functions of TAMs (Gocheva et al., 2010).

There are 11 human cysteine Cathepsin proteases (Cts) (B, C, H, F, K, L, O, S, V, W, and
25 X/Z), that share a conserved active site formed by cysteine and histidine residues. Cathepsins not only mediate terminal protein degradation in the lysosomes, but they also process and activate proteins including growth factors, transcription factors, and other proteases (Vasiljeva et al., 2007). For many years cysteine Cathepsins were believed to be localized exclusively in lysosomes, however, during pathological processes Cathepsins are
30 found in other cellular compartments or extracellularly (Jane et al., 2006), and at the plasma membrane (Mohamed and Sloane, 2006; Jane et al., 2006; Lechner et al., 2006). Besides their main function as lysosomal protein recycling machinery, the Cathepsins are known to participate in numerous physiological processes such as activation of proenzymes and prohormones (Friedrichs et al., 2003; Wolters et al., 2001), MHC-II
35 antigen processing and presentation (Honey et al., 2002), bone remodeling (Potts et al., 2004), keratinocyte differentiation (Roth et al., 2000; Reinheckel et al., 2005) and several others (Vasiljeva et al., 2007). However, proteases can be very harmful if not strictly controlled. There is a growing body of information that cysteine Cathepsins are involved in

numerous pathologies, like in particular cancer biology, but also viral infections, cardiovascular diseases and osteoarthritis, although the major focus is still on osteoporosis, rheumatoid arthritis, atherosclerosis and other diseases linked with the immune system (reviewed in Vasiljeva et al., 2007).

5

Interestingly, clinical studies in various tumour types have shown that certain Cathepsins, including Cts B, L, and S, are supplied to the tumour by stromal cells (Iacobuzio-Donahue et al. 1997; Hazen et al. 2000; Lindahl et al. 2009; Sullivan et al. 2009). This interplay between different cell types in the tumour microenvironment is highly complex, and therapeutic implications are thus difficult to predict.

10

A major role for cysteine Cathepsin proteases, produced by leukocytes and epithelial cells, as important mediators of tumour development, has also been recently appreciated (Turk et al., 2004). While many cysteine Cathepsins are lysosomal proteases, they are known to be involved in remodelling of ECM, to regulate cellular proliferation and death, to activate tumour angiogenesis, to promote invasion and metastasis of tumour cells, and to regulate inflammatory and immune responses in tissues (Chapman et al., 1997). Joyce and colleagues recently demonstrated the association of increased Cathepsin activity with angiogenic vasculature and invasive fronts of carcinomas during tumorigenesis in transgenic mouse models of Islet cell and cervical carcinogenesis using activity-based chemical probes and *in vivo* imaging (Joyce et al., 2004).

15

20

Cathepsins inhibitors were proposed for use in treatment of diseases associated with Cathepsins overexpression and/or secretion such as rheumatoid arthritis, osteoarthritis, atherosclerosis, neoplastic diseases and others. WO 2004/033445 describes compounds which are useful for treating diseases in which Cathepsin-dependent bone resorption was indicated. WO 2004/033445 relates to a method of treating or preventing diseases such as osteoporosis, Paget's disease, periodontal disease, tooth loss, bone fractures, rheumatoid arthritis, osteoarthritis, periprosthetic osteolysis, and some others using 4-amino-azepan-3-one cysteine proteases inhibitor. The use of specific inhibitors of Cathepsins L, B, S for treatment and prevention of diseases and disorders associated with these Cathepsins is described in patent WO 2009/136997.

25

30

WO 2009/133545 reports a drug targeting strategy for selective delivery of the anticancer drugs to endothelial cells by polymer-drug conjugates modified with a carbohydrate ligand. The use of such specific ligand is limited to the targeting of endothelial cells only. There is therefore a need for the identification and validation of novel surface antigens expressed on tumour cells.

35

Cathepsin cleavable peptide linkers are e.g. shown in (Jaffer et al., 2009). Those linkers could be cleaved by cathepsins. The cleavage will be used in that case for “liberating fluorochromes from the backbone and consequently generating strong fluorescence”.

5

The system developed in (Cegnar et al., 2004) was designed to be used for the inhibition of intracellular cathepsins. Cathepsin inhibitor, cystatin C, was “incorporated” as a drug, thus have been used as treatment compound transported being entrapped in the system.

10

Notably, the tumour cell-centered view of the metastatic process has been recently revised and the gained information on crosstalk between tumour cells and its surrounding tissue, or tumour stroma (Liotta and Kohn, 2001), supported the notion that the microenvironment is at least as determinative for tumour progression as the intrinsic features of tumour cells (Mueller and Fusenig, 2004). Thus, Cathepsins have been recently reported to be mainly overexpressed by tumour stromal cells (Vasiljeva et al., 2006; Gocheva et al., 2010). There is a need for novel approaches for the selective treatment of inflammatory and neoplastic diseases, with limited side effects. Therefore, a targeted approach is desirable. Moreover, there is a need for a reliable diagnosis of inflammatory and neoplastic diseases.

15

Although there are hints for a role of Cathepsins, in particular Cathepsin B, in the development and/or progression of various diseases, abundance of the Cathepsins in the human body and the complexity of the potential role of extracellular or membrane-localized Cathepsins appeared to make this target unsuitable for a targeting approach for delivering diagnostic and/or therapeutic moieties to a disease site. This is in particular true, as extracellularly localized Cathepsin is expected to be insufficiently close to cells to be targeted.

25

Compounds that will target cathepsins will target the site of disease (e.g. inflammation or neoplastic). Depends on the disease of interest where cathepsins are participated such a targeting system would enable targeted drug delivery to the site of the disease. Cysteine cathepsins are normally intracellular enzymes, thus their appearance extracellular could occur only at the site of the disease, thus enabling the fine targeting of medicine avoiding any possible toxic side effects.

30

Surprisingly, it could be shown in the Examples, that a Cathepsin-binding compound NS-629 bound to liposomes can surprisingly be used for binding to inflammatory cells. Moreover, it could surprisingly be shown in the Examples, that NS-629 bound to liposomes, wherein the liposomes carry a diagnostic label, like luciferine or the MRI agent Gd-DTPA, respectively, can be used for *in vivo* diagnosing inflammatory diseases, and/or

35

neoplastic diseases, by binding to cells of the tumour stroma, and that the compound surprisingly targets to the disease site *in vivo*.

The present invention relates to a Cathepsin-binding compound bound to a carrier comprising a therapeutic and/or diagnostic moiety,

- (i) for use in the diagnosis and/or treatment of inflammatory diseases, and/or
- (ii) for use in the diagnosis and/or treatment of neoplastic disease, wherein the Cathepsin-binding compound binds to inflammatory cells of the tumour stroma.

A Cathepsin-binding compound bound to a carrier preferably has following structure of formula (V):

Cathepsin-binding compound - linker - carrier (V)

It is understood, that in case of covalent linkage of the Cathepsin-binding compound to a linker, a Cathepsin-binding moiety is linked via at least one chemical bond to the linker. For example, MeO-Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH is described as Cathepsin-binding compound. In NS-629, the Cathepsin-binding moiety of this compound linked to the linker is -Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH.

In a further embodiment, a Cathepsin-binding compound bound to a carrier has following structure of formula (VI):

Cathepsin-binding compound - carrier (VI)

“Cathepsin” (Cts) according to the present invention is understood as a mammalian cysteine protease, in particular a human, rat, monkey, dog, cat, pig, cow or horse Cathepsin protease (Cts), more preferably a human cysteine protease. Preferably, the Cathepsin is Cts B, C, H, F, K, L, O, S, V, W, or X/Z, more preferably the Cathepsin is Cts B, even more preferably human Cts B, most preferably human Cts B having the NCBI Accession number AAC37547.1, AAH95408.1, AAH10240.1, or CAA77178.1.

Thus, in a preferred embodiment, the Cathepsin bound by the compound is selected from Cathepsin (Cts) B, C, H, F, K, L, O, S, V, W, and X/Z. In particular, the Cathepsin bound by the compound is overexpressed in an inflammatory and/or neoplastic disease, more preferably it is excessively located in the extracellular matrix of affected tissue in an

inflammatory and/or neoplastic disease. Preferably, the Cathepsin bound by the compound is selected from Cts B, and/or S, more preferably the compound is selected from Cts B.

Thus, in a further preferred embodiment, the Cathepsin is mammalian Cathepsin B, in particular human, rat, monkey, dog, cat, pig, cow or horse Cathepsin B, more preferably human Cathepsin B.

“Cathepsin-binding compound” is understood as compound having a K_D value with respect to at least one Cathepsin of less than about 100 μM , preferably less than about 10 μM , more preferably less than about 1 μM , even more preferably less than about 1 μM , especially preferably less than about 100 nM, most preferably less than about 10 nM, as determined in an assay described in Bieth, 1995, or by surface plasmon resonance described in Mirkovic et al., 2009. In one embodiment, the compound has a K_D value with respect to one Cathepsin of less than about 10 nM, and the K_D value with respect to at least one other Cathepsins is at least 10-fold, more preferably at least 100-fold higher than the K_D value with respect to one Cathepsin. Such compounds bind specifically to one Cathepsin. In a preferred embodiment, the compound binding specifically to one Cathepsin is an antibody or antibody fragment. The Cathepsin-binding compound may be any kind of compound, in particular, it is selected from an antibody or antibody fragment specifically binding to Cathepsin, a synthetic inhibitor of Cathepsin, a peptide, a protein and an antibody mimetic. In a further embodiment, the Cathepsin-binding compound is a protein or compound specifically binding to Cathepsin. In a preferred embodiment, the compound is a synthetic inhibitor of Cathepsin. In a further preferred embodiment, the compound is an inhibitor of the protease activity of a Cathepsin. The use of an inhibitor of the protease activity of a Cathepsin as a Cathepsin-binding compound enables to combine both the targeted delivery of therapeutically active compounds and the therapeutic effect of Cathepsin protease inhibition itself.

Herein, the term “antibody or antibody fragment” is understood as meaning antibodies, antibody fragments, or antigen-binding parts thereof, in particular complementarity-determining regions (CDRs) of antibodies, which have been prepared recombinantly and, where appropriate, modified, such as chimaeric antibodies, human antibodies, humanized antibodies, multifunctional antibodies, bispecific or oligospecific antibodies, single-chain antibodies and Fab, Fab' or $F(ab)_2$ fragments, or single domain VHH, VH or VL single domains, camelid antibodies and lama antibodies (see for example, Harmsen und De Haard, 2007; Holliger and Hudson, 2005; EP-B1-0 368 684, US 4,816,567, US 4,816,397, WO 88/01649, WO 93/06213 or WO 98/24884), fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the

above. The antibody is preferably a human IgG1 antibody. However, other human antibody isotypes are also encompassed by the invention, including IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgAsec, IgD and IgE. Also animal-derived antibodies of various isotypes can be used according to the invention.

5

An “amino-compound” according to the present invention is a compound of the general formula $NR^1R^2R^3$, wherein R^1 , R^2 and R^3 are selected from the group consisting of H, aliphatic and aromatic organic substituents.

10 “Antibody mimetics” according to the present invention are protein scaffolds different from antibodies and antibody fragments, e.g. DARPINs or anticalins which are based on lipocalin (Beste et al., 1999). The natural ligand-binding sites of the lipocalins, for example the retinol-binding protein or the bilin-binding protein, can be altered, for example by means of a “combinatorial protein design” approach, in such a way that they bind to
15 selected haptens (Skerra (1), 2000). Other known protein scaffolds are known as being alternatives to antibodies for molecular recognition (Skerra (2), 2000). Designed ankyrin repeat proteins (DARPin) are a promising class of non-immunoglobulin proteins that can offer advantages over antibodies for target binding in drug discovery and drug development. DARPins specifically targeting the tumour marker HER2 have also been
20 generated and were shown to function in both *in vitro* diagnostics and *in vivo* tumour targeting. DARPins are ideally suited for *in vivo* imaging or delivery of toxins or other therapeutic payloads because of their favorable molecular properties, including small size and high stability. The low-cost production in bacteria and the rapid generation of many target-specific DARPins make the DARPins approach useful for drug discovery.
25 Additionally, DARPins can be easily generated in multispecific formats, offering the potential to target an effector DARPins to a specific organ or to target multiple receptors with one molecule composed of several DARPins.

30 “About” according to the present invention is understood as meaning the experimental error range, in particular $\pm 5\%$ or $\pm 10\%$.

An “inhibitor of the protease activity of a Cathepsin” is understood as compound having a second order rate constant with respect to at least one Cathepsin of at least about 10^3 , preferably at least about 10^4 , more preferably at least about 10^5 , even more preferably at
35 least about 10^6 k_2/k_1 ($M^{-1} s^{-1}$) as determined in an assay described in Schaschke et al. (2), 2000. Preferably, the second order rate constant with respect to papain is at least about 10-fold, more preferably at least about 100-fold lower, than the second order rate constant for at least one Cathepsin. Examples for inhibitors of inhibitor of the protease activity of a

Cathepsin B are Ac-Leu-Leu-NLe-H described in Schaschke et al., 1996, and 4-amino-azepan-3-one compounds disclosed in WO 2004/033445, the inhibitors summarized in Yasuda et al., 2005, in particular cathepsin inhibitor AAE581 (Balicatib), compound 462795 (relacatib), azepanones SB 331750 and SB 357114, inhibitor of cathepsin S MV57471, reversible aldehyde type inhibitors as described in Saegusa et al., 2002, Katunuma et al., 1999, Yasuma et al., 1998, and Katunuma et al., 2000; cyclic hydrazide compounds as disclosed in Duffy et al., 1999; reversible cyclic ketone type inhibitors as described in Marquis et al. (1), 2001; Marquis et al., 1999, Marquis et al. (2), 2001, Marquis et al., 1998; reversible ketoamide type inhibitors as disclosed in Tavares et al. (1), 2004, Tavares et al. (2), 2004, Tavares et al. (3), 2004; reversible aminoethyl amide type inhibitors as disclosed in Altmann et al., 2002; reversible nitrile type inhibitors as disclosed in Greenspan et al., 2001, WO 01/58886, WO 02/20485, WO 01/09110, WO 01/19816, Robichaud et al., 2003, Ward et al., 2002; reversible cyanamide type inhibitors as disclosed in Falguyret et al., 2001; beta-Lactam 6-substituted oxapenam compounds as disclosed in Zhou et al., 2002, and US 5,905,076, beta-Lactam penam compounds as disclosed in US 6,232,305 B1 and Zhou et al., 2002; beta-Lactam monobactam compounds as disclosed in Zhou et al., 2003, and US 5,986,108; acrylamide compounds as disclosed in US 6,635,621; pyrazole compounds as disclosed in Thurmond et al. (2), 2004, Thurmond et al. (3), 2004; epoxysuccinate compounds as disclosed in Katunuma et al., 2002; vinyl sulfone compounds as disclosed in Palmer et al., 1995, Brömme et al., 1996, McKerrow et al., 1999; diacyl hydroxamate compounds as disclosed in Bromme et al., 1989, Bromme et al., 1994,, bis-hydrazides compounds as disclosed in Thompson et al., 1997, and Thompson et al., 1998,, acyclic ketone compounds as disclosed in Marquis et al., 1999, Huang et al., 2002, Shaw, 1994, Krantz, 1994, Smith et al., 2001, Seyfried et al., 2001; and beta-Lactam 2-substituted oxapenam compounds as disclosed in Zhou et al., 2001, and in US 5,925, 633; and the cathepsin inhibitors as summarized in Deaton and Kumar, 2004, in particular inhibitors of formula (1) to (273) disclosed therein. Preferred inhibitors of the protease activity of a Cathepsin are cathepsin inhibitors AAE581 (balicatib; N-[1-(cyanomethylcarbamoyl)cyclohexyl]-4-(4-propylpiperazin-1-yl)benzamide) and compound 462795 (relacatib; N-[(2S)-4-methyl-1-[[[(4S,7R)-7-methyl-3-oxo-1-pyridin-2-ylsulfonyl-azepan-4-yl]amino]-1-oxopentan-2-yl]-1-benzofuran-2-carboxamide).

An “inhibitor of the protease activity of Cathepsin B” is understood as compound having a second order rate constant with respect to at least one Cathepsin of at least about 10^3 , preferably at least about 10^4 , more preferably at least about 10^5 , even more preferably at least about 10^6 k_2/k_1 ($M^{-1} s^{-1}$) as determined in an assay described in Schaschke et al. (2), 2000. Preferably, the second order rate constant with respect to papain is at least about 10-fold, more preferably at least about 100-fold lower, than the second order rate for at least

one other Cathepsin. Preferably, the second order rate with respect to at least one other Cathepsin, in particular Cathepsin L, is at least about 10-fold, more preferably at least about 100-fold lower than the second order rate constant for Cathepsin B. In a preferred embodiment, the Cathepsin B is human Cathepsin B. In a preferred embodiment, the inhibitor of the protease activity of Cathepsin B is selected from NS-629, MeO-Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH and H₂N-(CH₂)₆-NH-Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH.

The “tumour stroma” is understood as cells and tissue components surrounding and interwoven with the tumour mass.

A “carrier” according to the present invention is understood a chemical entity which allows covalent or non-covalent binding of further moieties. In particular, the “carrier” according to the present invention is a nanodevice with a length in at least one dimension exceeding at least about 1 nm, preferably at least about 10 nm, more preferably at least about 100 nm, and which allows covalent or non-covalent binding of further moieties. In a preferred embodiment, the carrier is selected from a nanotube, a liposome, a lipoplex, a polymersome, a micell, a nanogel, a nanoparticle, a mesoporous silica particle, a dendrimer, and a nanoshell, in particular the carrier is a liposome, or other chemical entity that could be linked to the Cathepsin-binding compound. In a further preferred embodiment of the present invention, the carrier is selected from a nanotube, a liposome, a lipoplex, a polymersome, a micell, a nanogel, a nanoparticle, a mesoporous silica nanoparticle, a dendrimer, and a nanoshell, in particular the carrier is a liposome. In the case of a liposome, the therapeutic and/or diagnostic moiety may be encapsulated within the liposome, or the moiety may be bound via a lipid moiety to the membrane of the liposome. The generation of liposomes may be performed by methods known to a skilled person. In particular, the liposomes may be generated by extrusion as described in Example 3, or by sonification as described in Example 4. The Cathepsin-binding compound may be bound covalently or non-covalently to the carrier. As shown in Example 5, the Cathepsin-binding compound is in a preferred embodiment bound to the liposome non-covalently, in particular via a lipid moiety. In other embodiment, like e.g. dendrimers, the Cathepsin-binding compound and the therapeutic and/or diagnostic moiety are bound covalently to the dendrimer. The synthesis and use of a nanotube, a liposome, a lipoplex, a polymersome, a micell, a nanogel, a mesoporous silica nanoparticle, a dendrimer, and a nanoshell are for example described in Peer et al., 2007; Orive et al., 2009; Fahmy et al., 2007; Sun et al., 2008; and Ferrari, 2005.

Cathepsin-binding compounds may initially be chemically modified to enable further covalent or non-covalent coupling, in particular by introducing a linker. In one embodiment, the Cathepsin-binding compound is lipidated by introducing a lipid moiety, preferably via a linker, as for example shown in Example 1 and Figure 2. Lipidation of a
5 Cathepsin-binding compound allows easier coupling of such ligand to liposomes, micelles or lipoplexes, because of the lipid part a lipidated Cathepsin-binding compound can be added to the primary mixture in a carrier preparation. This method not only simplifies the coupling procedure but also excludes any manipulation of encapsulated compound.

10 Lipid vesicles are highly compatible with biological membranes in both composition and structure, and thus their use as delivery systems is well established. In this respect, liposomes loaded with at least one diagnostic or/and therapeutic moiety and bound to a Cathepsin-binding compound can be effective “drug carriers” for targeted delivery. This approach is very promising because, unlike magnetic targeting, it allows active targeting of
15 the bionanocomposite to the site of choice, in particular tumour stroma and/or inflammation site, without the need for additional devices such as a source of the external magnetic field.

Suitable linkers are described in Quasba et al., 2008; Cruz et al., 2010; Rossin et al., 2005;
20 Shuvaev et al., 2004; Liang et al., 2002 and Hylarides et al., 2001. In particular a functionalized PEG moiety and/or a diaminoalkyl moiety may be used as linker, as in NS-629.

Labelling of liposomes by use of a lipidated inhibitor is advantageous over other methods
25 such as direct coupling of ligand to liposome surface by covalent coupling (US Patent No. 5,013,556) or attaching to the free ends of the polymer chains forming the surface coat on the liposomes (Allen et al., 1995; Blume et al., 1993). The lipid part of the developed inhibitor, hence, allows its incorporation in the lipid bilayer by adding the compound to the primary lipid mixture. Incorporation into the bilayer can in particular be performed by
30 extrusion or sonification. Extrusion is exemplified in Example 3. Sonification is exemplified in Example 4.

A “liposome-binding compound” according to the present invention is understood as compound which is able to associate with liposomes covalently or non-covalently,
35 preferably covalently. In particular, a liposome-binding compound comprises at least one lipid moiety which allows incorporation in the lipid bilayer of a liposome.

A “therapeutic moiety” according to the present invention is a chemical moiety, which is capable of exhibiting a therapeutic effect when administered to a patient in need thereof in an effective amount. The therapeutic moiety is preferably selected from:

a toxin,

5 a chemotherapeutic agent,

in particular an alkylating agent and/or an anti-metabolite and/or a plant alkaloid and/or a taxane and/or a topoisomerase inhibitor and/or an antineoplastic agent

a radioactive moiety,

10 an apoptosis-inducing agent, and

an anti-inflammatory agent,

in particular a non-steroidal anti-inflammatory agents,

preferably selected from a salicylate, propionic acid

derivative, acetic acid derivative, enolic acid derivative, and

15 fenamic acid derivative, a selective COX-2 inhibitor, and a sulphonanilide, or

in particular a steroidal anti-inflammatory agents,

preferably a glucocorticoid.

20 It is understood, that the selection of the therapeutic moiety depends on the disease to be treated.

A “diagnostic moiety” according to the present invention is a chemical moiety, which can be detected. Preferably, the chemical moiety can be detected *in vitro*, *ex vivo*, or *in vivo*,
25 preferably *in vivo*. The diagnostic moiety is preferably selected from:

a radioactive label,

in particular technetium-99m, iodine-123, iodine-131, rhenium-186,
rhenium-188, gallium-67, yttrium-90, and lutetium-177,

a paramagnetic agent,

30 in particular a Gadolinium-complex, more preferably Gd-DTPA,

a super paramagnetic nanoparticle,

in particular iron oxide based nanoparticles,

a PET-imageable agent,

an MRI-imageable agent,

35 in particular a Gadolinium-complex, more preferably Gd-DTPA,

a fluorophore,

in particular Alexa Fluor,

a chromophore,

a phosphorescing agent,
a chemiluminescent agent, and
a bioluminescent agent.

5 In some embodiments, a chemical moiety is both a diagnostic and a therapeutic moiety, as e.g. certain radioactive isotopes like iodine-123 and iodine-131.

In vivo detection and imaging can be performed for example with a magnetic resonance imaging (MRI) apparatus, in case an MRI or paramagnetic agent is to be detected, or with
10 SPECT or PET cameras after accumulation of the radioactively labeled cathepsin-binding compound in the diseased tissue.

In one embodiment, the Cathepsin-binding compounds bound to a carrier comprising a therapeutic and/or diagnostic moiety and the Cathepsin-targeting compounds comprising a
15 therapeutic and/or diagnostic moiety, may be administered as the only treatment, or may be used simultaneously with, before, or after different types of treatment, e.g. surgery, radiation; or may be administered to patients with, before, or after conventional chemotherapeutic, radiotherapeutic, anti-inflammatory or anti-angiogenic agents, or targeted immunotoxins or coagulgands. The compounds of a combinatory treatment may
20 be administered to the patient simultaneously, either in a single composition, or as two distinct compositions using the same or different administration routes.

As shown in Example 9 and Figure 7, the Cathepsin-binding compound of the invention surprisingly localizes *in vivo* at an inflammation area, namely a paw edema. Therefore, in
25 one embodiment of the present invention, the invention relates to a Cathepsin-binding compound for use as described herein, wherein the use is a diagnostic use, and wherein the detection of a Cathepsin *in vivo* is indicative of an inflammatory site.

Cathepsin represents an early but dynamic target in inflammatory diseases and neoplastic
30 diseases. A Cathepsin-binding compound as described herein can therefore be applied for example at multiple points in the treatment cascade of neoplastic diseases, from primary chemoprevention to treatment of relapsed and disseminated disease.

In a preferred embodiment, the neoplastic disease is selected from bladder cancer, breast
35 cancer, colorectal cancer, endometrial cancer, head and neck cancer, leukemia, lung cancer, lymphoma, melanoma, non-small-cell lung cancer, ovarian cancer, prostate cancer, testicular cancer, uterine cancer, cervical cancer, thyroid cancer, gastric cancer, brain stem glioma, cerebellar astrocytoma, cerebral astrocytoma, glioblastoma, ependymoma, Ewing's

sarcoma family of tumours, germ cell tumour, extracranial cancer, Hodgkin's disease, acute lymphoblastic leukemia, acute myeloid leukemia, liver cancer, medulloblastoma, neuroblastoma, brain tumours, non-Hodgkin's lymphoma, mantle cell lymphoma, osteosarcoma, malignant fibrous histiocytoma of bone, retinoblastoma, rhabdomyosarcoma, soft tissue sarcomas, supratentorial primitive neuroectodermal pineal tumours, visual pathway and hypothalamic glioma, Wilms' tumour, acute lymphocytic leukemia, adult acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, esophageal cancer, hairy cell leukemia, kidney cancer, multiple myeloma, oral cancer, pancreatic cancer, primary central nervous system lymphoma, skin cancer, and small-cell lung cancer, in particular lung cancer, non-small-cell lung cancer, prostate cancer, ovarian cancer, colorectal cancer, pancreatic cancer, breast cancer, glioblastoma, and metastasis thereof, more preferably breast cancer and metastasis thereof.

In a preferred embodiment, the inflammatory disease is selected from psoriasis, contact dermatitis, atopic dermatitis, alopecia areata, erythema multiforme, dermatitis herpetiformis, scleroderma, vitiligo, hypersensitivity angiitis, urticaria, bullous pemphigoid, lupus erythematosus, pemphigus, epidermolysis bullosa acquisita, inflammatory bowel disease, rheumatoid arthritis, juvenile rheumatoid arthritis, psoriatic arthritis, osteoarthritis, refractory rheumatoid arthritis, chronic non-rheumatoid arthritis, osteoporosis/bone resorption, Crohn's disease, septic shock, endotoxic shock, atherosclerosis, ischaemia-reperfusion injury, coronary heart disease, vasculitis, amyloidosis, multiple sclerosis, sepsis, chronic recurrent uveitis, hepatitis C virus infection, malaria, ulcerative colitis, cachexia, plasmocytoma, endometriosis, Behcet's disease, Wegener's granulomatosis, AIDS, HIV infection, autoimmune disease, immune deficiency, common variable immunodeficiency (CVID), chronic graft-versus-host disease, trauma and transplant rejection, adult respiratory distress syndrome, pulmonary fibrosis, recurrent ovarian cancer, lymphoproliferative disease, refractory multiple myeloma, myeloproliferative disorder, diabetes, juvenile diabetes, meningitis, viral infections, cardiovascular diseases, ankylosing spondylitis, skin delayed type hypersensitivity disorders, Alzheimer's disease, systemic lupus erythematosus and allergic asthma, preferably selected from rheumatoid arthritis, Crohn's disease, ulcerative colitis, septic shock, psoriasis and atherosclerosis.

In another embodiment, the present invention relates to the use of a Cathepsin-binding compound bound to a carrier comprising a therapeutic and/or diagnostic moiety,

- (i) for the preparation of a diagnostic and/or medicament for the diagnosis and/or treatment of inflammatory diseases, and/or

- (ii) for the preparation of a diagnostic and/or medicament for the diagnosis and/or treatment of neoplastic diseases, wherein the Cathepsin-binding compound binds to inflammatory cells of the tumour stroma.

5 In a further embodiment, the present invention relates to a method for diagnosing and/or treating an inflammatory disease, comprising administering an effective amount of a Cathepsin-binding compound bound to a carrier comprising a therapeutic and/or diagnostic moiety to a patient in need thereof.

10 In a further embodiment, the present invention relates to a method for diagnosing and/or treating neoplastic diseases, comprising administering an effective amount of a Cathepsin-binding compound bound to a carrier comprising a therapeutic and/or diagnostic moiety to a patient in need thereof, wherein the Cathepsin-binding compound binds to inflammatory cells of the tumour stroma.

15

Surprisingly, it could be shown in Examples 6 and 7, that the Cathepsin-binding compound NS-629 non-covalently bound to liposomes can surprisingly be used for binding to inflammatory cells. Moreover, it could surprisingly be shown in the Examples 8 and 9, that NS-629 bound to liposomes, wherein the liposomes carry a diagnostic label, like luciferine or the MRI agent Gd-DTPA, respectively, can be used for *in vivo* diagnosing inflammatory diseases, and/or neoplastic diseases, by binding to tumour stromal cells, and that the compound surprisingly targets to disease site *in vivo* (see also Figure 7 and 8). NS-629 is obtained by reacting $\text{H}_2\text{N}-(\text{CH}_2)_6-\text{NH}-\text{Gly}-\text{Gly}-\text{Leu}(2\text{S},3\text{S})-t\text{Eps}-\text{Leu}-\text{Pro}-\text{OH}\times\text{TFA}$ with DSPE-PEG(2000). The hexane-1,6-diamine moiety and the PEG moiety are part of the linker between the Cathepsin B-inhibitor moiety and the lipid moiety.

20
25

Therefore, in another embodiment, the present invention relates to a Cathepsin -binding and liposome-binding compound, comprising:

- 30 (i) a Cathepsin -binding moiety,
(ii) optionally a linker, and
(iii) a liposome-binding moiety.

In a preferred embodiment, the Cathepsin-binding and liposome-binding compound is a Cathepsin B-binding and liposome-binding compound and comprises:

- 35 (i) a Cathepsin B-binding moiety,
(ii) optionally a linker, and
(iii) a liposome-binding moiety.

In a further preferred embodiment, the Cathepsin-binding and liposome-binding compound is a Cathepsin S-binding and liposome-binding compound and comprises:

- 5 (i) a Cathepsin S-binding moiety,
 (ii) optionally a linker, and
 (iii) a liposome-binding moiety.

In a preferred embodiment, a Cathepsin -binding and liposome-binding compound, preferably a Cathepsin B-binding and liposome-binding compound of the invention or Cathepsin S-binding and liposome-binding compound of the invention has the structure of
 10 formula (IV):

Cathepsin -binding moiety - linker - liposome-binding moiety (IV),

and in the case of a Cathepsin B-binding and liposome-binding compound,

15

Cathepsin B-binding moiety - linker - liposome-binding moiety (IV)',

and in the case of a Cathepsin S-binding and liposome-binding compound,

20

Cathepsin S-binding moiety - linker - liposome-binding moiety (IV)''.

In a preferred embodiment, the Cathepsin-binding and liposome-binding compound, preferably the Cathepsin B-binding and liposome-binding compound comprises a linker, even more preferably the linker comprises a functionalized PEG moiety and/or a
 25 diaminoalkyl moiety. The diaminoalkyl moiety is preferably a moiety having the structure -HN-(CH₂)_l-NH- wherein l is an integer from n=2 to 30, preferably 3 to 20, most preferably n=6.

In a further preferred embodiment, the linker in the Cathepsin-binding and liposome-
 30 binding compound, preferably the Cathepsin B-binding and liposome-binding compound has the structure of formulas II to V:


- NH-(CH₂)_{l+1}-(NH)-C(O)-(CH₂)_m-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (II),

35

-NH-(CH₂)_{m+1}-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (III),

-C(O)-(CH₂)_l-(NH)-C(O)-(CH₂)_m-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (IV),

and



wherein

l is an integer from n=1 to 30, preferably 3 to 20, most preferably n=5,


5 m is an integer from n=1 to 30, more preferably 1 to 10, most preferably 1,

p is an integer from n=1 to 200, more preferably 15 to 100, most preferably 38 to 53,

q is an integer from n=2 to 30, more preferably 2 to 10, most preferably 2.

10 In a preferred embodiment, the liposome-binding moiety of the Cathepsin binding and liposome-binding compound, preferably the Cathepsin B-binding and liposome-binding compound of the invention is a phospholipid, in particular a phospholipid comprising 1, 2 or more stearyl moieties, 1, 2 or more oleyl moieties, 1, 2 or more lauroyl moieties, 1, 2 or more myristoyl moieties, 1, 2 or more palmitoyl moieties, and/or 1, 2 or more arachinoyl moieties, more preferably a phospholipid comprising 2 stearyl moieties.

In a more preferred embodiment, the Cathepsin B-binding and liposome-binding compound of the invention has the structure of formula (XIII):



20 wherein n=38 to 53.

It is understood, that in a preparation, various lengths of the PEG moiety may be present. Preferably, the most abundant PEG species has n=45 or n=46. Thus, in a further preferred embodiment, the invention relates to a mixture Cathepsin B-binding and liposome-binding compounds of the invention of formula (XIII), wherein the n in the compounds are 2 or more different integers from 38 to 53.

In a further preferred embodiment, the Cathepsin-binding and liposome-binding compound of the invention comprises a Cathepsin-binding moiety, wherein the Cathepsin-binding

moiety is an inhibitor of the protease activity of Cathepsin. The Cathepsin-binding moiety is in particular a peptide inhibitor.

The Cathepsin-binding moiety is preferably selected from the group consisting of:

-Gly-Aaa-Bbb-X-Ccc-Ddd (VI),

-Gly-Aaa-Bbb-X-Ccc (VII)

and

-Gly-Aaa-Bbb-Y (VIII),

wherein


X is an electrophilic unit, in particular an epoxide-derivative or a semicarbazide-derivative, preferably *t*Eps, -(NH)-(NH)-C(O)-, most preferably (2S,3S)-*t*Eps;

Y is an electrophilic unit, in particular an epoxide-derivative, a semicarbazide-derivative or a nitrile-derivative, preferably -*t*Eps, -(NH)-(NH)-C(O)-NH₂, -(NH)-CH₂-CN, more preferably (2S,3S)-*t*Eps;


Aaa is selected from the group consisting of Gly, Phe, Asp, Glu, 1-aminoadipic acid, in particular Gly;

Bbb is selected from the group consisting of aliphatic or aromatic hydrophobic amino


acids, in particular Leu, Ile, Phe, Tyr, Val,




(XI),



(XII),



(IX),



(X),

2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, preferably amino acids (IX), (X), (XI), (XII), 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, Leu, Ile, Phe, Tyr, more preferably amino acids (IX), (X), (XI), (XII), 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, Leu and Ile;

Ccc is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Phe, Tyr, Val, Trp, Nle, preferably Nle and Leu, more preferably Leu;

5 Ddd is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids and amino compounds, in particular Trp, Val, Ile, Phe, Tyr, Ala, Ser, Thr, Pro, $-\text{NH}(\text{CH}_2)_{n=0-4}-\text{CH}_3$, $-\text{NH}-\text{CH}(\text{CH}_3)_2$, $-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$, $-\text{NH}-(\text{CH}_2)_{n=1-2}\text{Ph}$, preferably, Trp, Val, Ile, Phe, Tyr, more preferably Trp.

10 In preferred embodiments, the electrophilic units X and Y are epoxide-derivatives. A preferred epoxide-derivative according to the invention is *tEps*, wherein *tEps* is available in several stereoisomers. Preferable stereoisomers are (2S,3S)-*tEps* and (2R,3R)-*tEps*, wherein (2S,3S)-*tEps* is more preferred.

15 In a preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-\text{Gly}-\text{Aaa}-\text{Bbb}-\text{X}-\text{Ccc}-\text{Ddd}$, wherein X is (2S,3S)-*tEps* and Aaa, Bbb, Ccc and Ddd are selected from the group consisting of aliphatic amino acids, preferably from the group consisting of aliphatic
20 hydrophobic amino acids.

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-\text{Gly}-\text{Aaa}-\text{Bbb}-\text{X}-\text{Ccc}-\text{Ddd}$,
25 wherein X is $-(\text{NH})-(\text{NH})-\text{C}(\text{O})-$ and Aaa, Bbb, Ccc and Ddd are selected from the group consisting of aliphatic amino acids, preferably from the group consisting of aliphatic hydrophobic amino acids.

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin
30 B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-\text{Gly}-\text{Aaa}-\text{Bbb}-\text{X}-\text{Ccc}-\text{Ddd}$, wherein X is (2S,3S)-*tEps* and Aaa, Bbb, Ccc and Ddd are selected from the group consisting of aromatic amino acids, preferably from the group consisting of aromatic hydrophobic amino acids.

35

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-\text{Gly}-\text{Aaa}-\text{Bbb}-\text{X}-\text{Ccc}-\text{Ddd}$,

wherein X is $-(NH)-(NH)-C(O)-$ and Aaa, Bbb, Ccc and Ddd are selected from the group consisting of aromatic amino acids, preferably from the group consisting of aromatic hydrophobic amino acids.

5 In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-Gly-Aaa-Bbb-Y$, wherein Y is (2S,3S)-*t*Eps and Aaa, Bbb are selected from the group consisting of aliphatic amino acids, preferably from the group consisting of aliphatic hydrophobic amino acids.

10

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-Gly-Aaa-Bbb-Y$, wherein Y is $-(NH)-(NH)-C(O)-NH_2$ and Aaa, Bbb are selected from the group consisting of aliphatic amino acids, preferably from the group consisting of aliphatic hydrophobic amino acids.

15

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-Gly-Aaa-Bbb-Y$, wherein Y is (2S,3S)-*t*Eps and Aaa, Bbb are selected from the group consisting of aromatic amino acids, preferably from the group consisting of aromatic hydrophobic amino acids.

20

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-Gly-Aaa-Bbb-Y$, wherein Y is $-(NH)-(NH)-C(O)-NH_2$ and Aaa, Bbb are selected from the group consisting of aromatic amino acids, preferably from the group consisting of aromatic hydrophobic amino acids.

25

In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-Gly-Aaa-Bbb-Y$, wherein Y is $-(NH)-CH_2-CN$ and Aaa, Bbb are selected from the group consisting of aliphatic amino acids, preferably from the group consisting of aliphatic hydrophobic amino acids.

30

35 In a further preferred embodiment, the Cathepsin-binding moiety, preferably the Cathepsin B-binding moiety or Cathepsin S-binding moiety, most preferably the Cathepsin B-binding moiety, is a peptide inhibitor represented by the formula $-Gly-Aaa-Bbb-Y$, wherein Y is $-(NH)-CH_2-CN$ and Aaa, Bbb are selected from the group consisting of aliphatic amino acids, preferably from the group consisting of aliphatic hydrophobic amino acids.

(NH)-CH₂-CN and Aaa, Bbb are selected from the group consisting of aromatic amino acids, preferably from the group consisting of aromatic hydrophobic amino acids.

In a further embodiment according to the invention, the Cathepsin-binding and liposome-binding compound is a Cathepsin B-binding moiety and is an inhibitor of the protease activity of Cathepsin B. The Cathepsin B-binding moiety is in particular a peptide inhibitor, preferably a peptide inhibitor selected from the group consisting of



10 and



wherein X, Aaa, Bbb, Ccc and Ddd have the meanings as described below.

15 In a preferred embodiment X is an electrophilic unit, in particular an epoxide-derivative or a semicarbazide-derivative, preferably *t*Eps or -(NH)-(NH)-C(O)-, most preferably (2S,3S)-*t*Eps.

In a further preferred embodiment Aaa is an aliphatic amino acid of the general formula
20 NH₂-CHR-COOH, wherein R is preferably selected from the group consisting of hydrogen, methyl or isopropyl. In a preferred embodiment R is hydrogen and therefore Aaa is Gly.

In a further preferred embodiment Bbb is selected from the group consisting of aliphatic or
25 aromatic hydrophobic amino acids, in particular Leu, Ile, Phe, Tyr, Val, preferably Leu, Ile, Phe, Tyr, more preferably Leu and Ile;

According to the invention Ccc is preferably selected from the group consisting of
30 aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Val, Nle, preferably Nle and Leu, more preferably Leu;

In another preferred embodiment Ddd is selected from the group consisting of aliphatic or
aromatic hydrophobic amino acids and amino compounds, in particular Trp, Val, Ile, Phe,
Tyr, Ala, Ser, Thr, Pro, preferably, Trp, Val, Ile, Phe, Tyr, more preferably Trp.

35

Preferred embodiments of the Cathepsin-binding moiety, in particular of the Cathepsin B-binding moiety are -Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-Pro-OH, -Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-Trp-OH, -Gly-Gly-Leu-(2S,3S)-*t*Eps-Nle-Pro-OH, -Gly-Gly-Ile-(2S,3S)-*t*Eps-Leu-

Pro-OH, –Gly-Gly-Ile-(2S,3S)-*t*Eps-Leu-Trp-OH, –Gly-Gly-Ile-(2S,3S)-*t*Eps-Nle-Trp-OH, more preferably are the moieties –Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-Pro-OH and –Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-Trp-OH, most preferably –Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-Pro-OH. According to the invention, -OH represents that the amino acid is present as free acid.

5

In a further embodiment according to the invention, the Cathepsin-binding and liposome-binding compound is a Cathepsin S-binding moiety and is an inhibitor of the protease activity of Cathepsin S. The Cathepsin S-binding moiety is in particular a peptide inhibitor, preferably a peptide inhibitor selected from the group consisting of

10

-Gly-Aaa-Bbb-X-Ccc-Ddd (VI),

-Gly-Aaa-Bbb-X-Ccc (VII)

15 and

-Gly-Aaa-Bbb-Y (VIII),

wherein X, Y, Aaa, Bbb, Ccc and Ddd have the meanings as described below.

20 In a preferred embodiment X is an electrophilic unit, in particular an epoxide-derivative or a semicarbazide-derivative, preferably *t*Eps or -(NH)-(NH)-C(O)-, most preferably (2S,3S)-*t*Eps.


In a further preferred embodiment Y is an electrophilic unit, in particular an epoxide-derivative, a semicarbazide-derivative or a nitrile-derivative. Preferably, Y is *t*Eps, -(NH)-(NH)-C(O)-NH₂, -(NH)-CH₂-CN, more preferably (2S,3S)-*t*Eps.

In a further preferred embodiment Aaa is an aliphatic, aromatic or acidic α -amino acid. Aaa is preferably selected from the group consisting of Gly, Phe, Asp, Glu, 1-aminoadipic acid, wherein Gly is most preferred.

30

In a further preferred embodiment Bbb is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids. In particular Bbb is selected from the group of proteinogenic amino acids consisting of Leu, Ile, Phe, Tyr, Val or selected from the group of non-proteinogenic amino acids consisting of 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine or selected from the group of non-proteinogenic amino acids with the formulas:

35




Bbb is more preferably selected from the group of 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, Leu, Ile, Phe, Tyr and amino acids (IX), (X), (XI) or (XII). Most preferably, Bbb is selected from the group of amino acids (IX), (X), (XI) or (XII), 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine and Leu.

In a further preferred embodiment, Ccc is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Phe, Tyr, Val, Trp, preferably Leu and Ile, more preferably Leu.

In another preferred embodiment Ddd is an amino compound, in particular an alkyl amino compound, an alkaryl amino compound or an aryl amino compound, in particular -NH(CH₂)_{n=0-4}-CH₃, -NH-CH(CH₃)₂, -NH-CH₂-CH(CH₃)₂, -NH-(CH₂)_{n=1-2}Ph.

Preferred embodiments of the Cathepsin-binding moiety, in particular of the Cathepsin S-moiety are -Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-OH, -Gly-Glu-Leu-(2S,3S)-*t*Eps-Leu-OH, -Gly-Glu-Leu-(2S,3S)-*t*Eps-Ile-OH, -Gly-Gly-Leu-(2S,3S)-*t*Eps-Leu-NH-CH₂-CH₃, more preferably is the moiety -Gly-Glu-Leu-(2S,3S)-*t*Eps-Leu-NH-CH₂-CH₃, -Gly-Gly-Xxx - (2S,3S)-*t*Eps-Leu-NH-CH₂-CH₃, where Xxx represents an amino acid with the formula (IX) or (X). According to the invention, -OH represents that the amino acid is present as free acid.

Further examples of preferred Cathepsin S-moieties are:




In formulas (XIII), (XIV) and (XV) the phenyl-ring can be substituted in ortho-, meta or para-position.

5 In preferred embodiments of the invention, the Cathepsin-binding and liposome-binding compound, in particular the Cathepsin B-binding and liposome-binding compound and Cathepsin S-binding and liposome-binding compound, more preferably the Cathepsin B-binding and liposome-binding compound comprises next to the Cathepsin-binding moiety a linker.


10

In embodiments of the present invention in which the electrophilic unit X and/or Y of the Cathepsin-binding moiety is an epoxide-derivative, the linker is preferably selected from



15

and/or



20

wherein

l is an integer from n=1 to 30, preferably 3 to 20, most preferably n=5,

m is an integer from n=1 to 30, more preferably 1 to 10, most preferably 1,

p is an integer from n=1 to 200, more preferably 15 to 100, most preferably 38 to 53,

25


q is an integer from n=2 to 30, more preferably 2 to 10, most preferably 2.

In embodiments of the present invention in which the electrophilic unit X and/or Y of the Cathepsin-binding moiety is a semicarbazide-derivative or nitrile-derivative, the linker is preferably selected from

30



and/or



35

wherein

l is an integer from n=1 to 30, preferably 3 to 20, most preferably n=5,

m is an integer from n=1 to 30, more preferably 1 to 10, most preferably 1,

- p is an integer from n=1 to 200, more preferably 15 to 100, most preferably 38 to 53,
- q is an integer from n=2 to 30, more preferably 21 to 10, most preferably 2.

5 The invention further relates to a Cathepsin-targeting, in particular to a Cathepsin B-targeting compound or Cathepsin-S-targeting compound, comprising a Cathepsin-binding and liposome-binding compound of the invention, in particular a Cathepsin B-binding and liposome-binding compound of the invention or Cathepsin S-binding and liposome-binding compound of the invention, bound to a liposome.

10

The invention further relates to a Cathepsin targeting compound, further comprising a therapeutic and/or diagnostic moiety.

15 The invention further relates to a Cathepsin B-targeting compound, further comprising a therapeutic and/or diagnostic moiety.

In a preferred embodiment, the Cathepsin targeting compound of the invention, or the Cathepsin binding and liposome-binding compound of the invention comprise a cathepsin-binding compound, which is an inhibitor of the protease activity of Cathepsin, in particular
20 a peptide inhibitor, more preferably the moiety -Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH.

In a preferred embodiment, the Cathepsin B-targeting compound of the invention, or the Cathepsin B-binding and liposome-binding compound of the invention comprise a cathepsin-binding compound, which is an inhibitor of the protease activity of Cathepsin B,
25 in particular a peptide inhibitor, more preferably the moiety -Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH.

In a preferred embodiment, the Cathepsin targeting compound of the invention, in particular the Cathepsin B-targeting compound of the invention or the Cathepsin S-targeting compound of the invention, comprises a diagnostic moiety, wherein the
30 diagnostic moiety is preferably selected from:

- a radioactive label,
- a paramagnetic agent,
 - in particular a Gadolinium-complex, more preferably Gd-DTPA,
- 35 a super paramagnetic nanoparticle,
 - in particular iron oxide based nanoparticles,
- a PET-imageable agent,
- an MRI-imageable agent,

in particular a Gadolinium-complex, more preferably Gd-DTPA,
a fluorophore,
in particular Alexa Fluor,
a chromophore,
5 a phosphorescing agent,
a chemiluminescent agent, and
a bioluminescent agent.

10 In a preferred embodiment, the Cathepsin targeting compound of the invention comprises a therapeutic moiety,

wherein the therapeutic moiety is preferably selected from:

a toxin,
a chemotherapeutic agent,
15 in particular an alkylating agent or/and an anti-metabolite or/and a plant alkaloid or/and a taxane or/and a topoisomerase inhibitor or/and a antineoplastic agent
a radioactive moiety,
an apoptosis-inducing agent, and
an anti-inflammatory agent,

20 in particular a non-steroidal anti-inflammatory agents,
preferably selected from a salicylate, propionic acid derivative, acetic acid derivative, enolic acid derivative, and fenamic acid derivative, a selective COX-2 inhibitor, and a sulphonanilide, or
25 in particular a steroidal anti-inflammatory agents,
preferably a glucocorticoids.

30 The invention further relates to a Cathepsin targeting compound of the invention, in particular to a Cathepsin B-targeting compound of the invention or a Cathepsin S-targeting compound of the invention,

- (i) for use in the diagnosis of inflammatory diseases,
and/or
(ii) for use in the diagnosis of neoplastic diseases, wherein the Cathepsin-targeting moiety binds to inflammatory cells of the tumor stroma, in
35 particular in breast neoplastic disease.

“bound to” according to the present invention is understood as being covalently or non-covalently bound. In the case of non-covalent binding, e.g. hydrophobic forces may ensure

stable binding, for example by a lipid anchor inserted into a lipid bilayer, as shown in the Examples.

5 In a preferred embodiment, the Cathepsin -binding and liposome-binding compound of the invention is non-covalently bound to a liposome.

In a preferred embodiment, the Cathepsin B-binding and liposome-binding compound of the invention is non-covalently bound to a liposome.

10 In a further preferred embodiment, the Cathepsin binding moiety of the invention is covalently coupled to a lipid or lipid derivative via a linker. As in Figure 1, the linker may comprise ϵ -aminohexanoic acid and a functionalized PEG moiety.

15 In a further preferred embodiment, the Cathepsin B-binding moiety of the invention is covalently coupled to a lipid or lipid derivative via a linker. As in Figure 1, the linker may comprise ϵ -aminohexanoic acid and a functionalized PEG moiety.

In a further preferred embodiment, the a Cathepsin binding moiety is an inhibitor of the protease activity of Cathepsin. As shown in the examples, NS-629 may be used. In a
20 further preferred embodiment, the Cathepsin -targeting compound comprises NS-629 bound to a liposome.

In a further preferred embodiment, the a Cathepsin B-binding moiety is an inhibitor of the protease activity of Cathepsin B. As shown in the examples, NS-629 may be used. In a
25 further preferred embodiment, the Cathepsin B-targeting compound comprises NS-629 bound to a liposome.

30 In a further embodiment, the present invention relates to a method for diagnosing and/or treating an inflammatory disease, comprising administering an effective amount of a Cathepsin-targeting compound of the invention comprising a therapeutic and/or diagnostic moiety to a patient in need thereof.

35 In a further embodiment, the present invention relates to a method for diagnosing and/or treating neoplastic diseases, comprising administering an effective amount of a Cathepsin-targeting compound to a patient in need thereof, wherein the Cathepsin-binding moiety binds to inflammatory cells of the tumour stroma.

In a preferred embodiment, the compounds and compounds for use are administered as a pharmaceutical composition.

5 The pharmaceutical composition of the present invention contains therapeutically and/or diagnostically effective amounts of the individual compounds of the invention and generally an acceptable pharmaceutical "drug carrier", diluent or excipient, e.g. sterile water, physiological saline, bacteriostatic saline, i.e. saline containing about 0.9% mg/ml benzyl alcohol, phosphate-buffered saline, Hank's solution, Ringer's-lactate, lactose, dextrose, sucrose, trehalose, sorbitol, mannitol, and the like. The composition is generally a
10 solution or suspension. It can be administered systemically, intravenously, orally, subcutaneously, intramuscularly, pulmonary, by inhalation and/or through sustained release administrations. Preferably, the composition is administered systemically, in particular intravenously.

15 The term "therapeutically effective amount" generally means the quantity of a compound of the invention which results in the desired therapeutic effect. A typical dosage range is from about 0.01 mg 1000 mg per application.

The term "diagnostically effective amount" generally means the quantity of a compound of
20 the invention which results in the desired diagnostic effect without causing unacceptable side-effects. A typical dosage range is from about 0.01 mg 1000 mg per application. An advantage of the invention is that low dosages can be used and therefore compounds of the invention e.g. can hardly be detected 24 hours after administration.

25 Generally, the application of the pharmaceutical composition to a patient is one or several times per day, or one or several times a week, or even during longer time periods as the case may be.

The invention further relates to a method for preparing a Cathepsin-binding and liposome-binding
30 compound, comprising providing a Cathepsin-binding moiety and associating a liposome-binding moiety, in particular associating a linker and a liposome-binding moiety with the Cathepsin-binding moiety. Preferably, the Cathepsin-binding moiety and/or the linker and/or the liposome-binding moiety are connected by chemical synthesis known to a person skilled in the art. In a further embodiment the Cathepsin-binding moiety and/or the
35 linker and/or the liposome-binding moiety are protected by appropriate protecting groups. Examples of protecting groups are for amines BOC, Cbz or Fmoc, for alcohols are MOM or THP and for carboxygroups are Bn or methyl. Further protecting groups are possible, too, and are known to a person skilled in the art.

Figures

Fig. 1: Preferred embodiment of the present invention. **A**, cathepsins are normally
5 localized in the intracellular organelles: endosomes or lysosomes. However, in
several pathological conditions cathepsins could be translocated to the extracellular
milieu. **B**, An example of the cathepsin-binding compounds bound to a carrier,
represented as LNP-NS-629. NS-629 represents the preferred Cathepsin-binding
and liposome-binding compound according to formula (III), wherein $n=38$ to 53 . In
10 exemplary embodiments, the cathepsin-binding compounds bound to a carrier are
 $90-110$ nm. **C**, Targeting system based on cathepsin-binding compounds binding to
extracellular cathepsins enabling active targeting of encapsulated drug.

Fig. 2: Chemical structure of the lipidated inhibitor NS-629 (length distribution of the PEG
15 segment: $n=38-53$).

Fig. 3: Titration of Cathepsin B with free NS-629 (a) and liposome-coupled lipidated NS-
629 (b) at pH 6.0 and 25° C. The solid lines were generated by linear regression
analysis.

20 Fig. 4: Size distribution of liposomes functionalized with lipidated inhibitor as determined
by dynamic light scattering (DLS). Average size is 85.66 nm.

Fig. 5: Validation of targeting of liposomes with lipidated Cathepsin inhibitor to the
25 immune cells. Liposomes functionalized (NS-Lip-Alx) and not functionalized (Lip-
Alx) with NS-629 were loaded with fluorescence marker (Alexa Fluor 546TM;
(Invitrogen)) and incubated with mouse bone marrow-derived macrophages for 15
minutes at 4° C. Fluorescence of accumulated marker was measured with TECAN
plate reader. As a control liposomes without labeling were used.

30 Fig. 6: Representation of liposomes with lipidated Cathepsin inhibitor targeting efficiency
in primary mouse immune cells.

Non-functionalized liposomes (Lip-Alx) and liposomes functionalized with NS-629
(NS-Lip-Alx) were loaded with fluorescence marker (Alexa Fluor 555TM

(Invitrogen)) and incubated with mouse bone marrow-derived macrophages for 15 minutes at 4° C. Fluorescence of accumulated marker was examined with an Olympus fluorescence microscope (Olympus IX 81) with Imaging Software for Life Science Microscopy Cell^f.


- 5 Fig. 7: Targeted delivery of liposomes labeled by lipidated inhibitor carrying D-luciferin into transgenic mouse expressing luciferase (FVB.luc^{tg/+}). The high-intensity luciferase signal associated with the induced paw edema demonstrates selective accumulation of labelled liposomes in the inflammation area. The scale is in photons/sec/sm²/sr.
- 10 Fig. 8: T₁-weighted MR images (TE=8.5 ms, TR=400 ms) of an orthotopic transplanted breast cancer mouse before, 1 and 24 hours after (T₁) intraperitoneal injection of 200 µl NS-629 labeled liposomes containing Magnevist® (Bayer HealthCare Pharmaceuticals). The tumour tissue possess negative MR signal on T₁-weighted images. The bright signal at 1 and 24 hours after injection in T₁-weighted MR
- 15 image indicates successful targeting of Magnevist® (Bayer HealthCare Pharmaceuticals) loaded NS-629 labelled liposomes.

Examples

20 Example 1: Synthesis of the lipidated inhibitor NS-629

DSPE-PEG(2000) refers to 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[carboxy(polyethylene glycol)-2000] (ammonium salt) (CAS Reg. 474922-20-8). DSPE-PEG(2000) carboxylic acid with PEG unit length of n=45 (PEG lengths of n=38 to 53 are present) has the following formula I:

25



- 30 DSPE-PEG(2000) carboxylic acid (Avanti Polar Lipids, Inc.) (30.0 mg, 10.5 µmol) was suspended in MeCN (1 mL) and CHCl₃ was added dropwise until a clear solution was obtained. Then at room temperature under stirring a solution of DSC (Fluka, Buchs) in

MeCN (c=0.0105 mol/L, 1 mL) and a solution of DIPEA in MeCN (c=0.105 mol/L, 0.1 mL) were added. In parallel, H₂N-(CH₂)₆-NH-Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH×TFA (16.5 mg, 21 μmol; prepared by treating H₂N-(CH₂)₆-NH-Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-*O**t*Bu (Schaschke et al. (1), 2000) with TFA/H₂O (95:5, v/v)) was suspended in MeCN (1 mL) and H₂O was added dropwise until a clear solution was obtained. H₂N-(CH₂)₆-NH-Gly-Gly-Leu-(2*S*,3*S*)-*t*Eps-Leu-Pro-OH×TFA and its synthesis are described in Schaschke et al. (2), 2000, and in Schaschke et al., 1998. Then to this solution a solution of DIPEA in MeCN (c=0.105 mol/L, 0.4 mL) was added. After 60 min, this solution was added to the formed active ester of DSPE-PEG(2000) carboxylic acid and stirring was continued for 48 h. The solvent was evaporated under reduced pressure and the resulting material was dissolved in CHCl₃ (50 mL). The organic phase was washed with 5% aq KHSO₄ (3×20 mL), brine (1×20 mL), dried (Na₂SO₄), and the solvent evaporated. The obtained crude product was dissolved in MeCN/H₂O (1:3, v/v; 10 mL), lyophilized, and purified by thick-layer chromatography using glass plates from Merck, Darmstadt (type: PLC Silica gel 60 F₂₅₄, 1mm). Prior to use the plates were developed with MeOH twice. Upon development with CHCl₃/MeOH (2:1, v/v containing 1% AcOH) as solvent system, the area containing homogeneous product was scraped from the plate, the product extracted from the collected silica gel with MeOH, and the solvent evaporated. The lipidated inhibitor (NS-629) was obtained upon lyophilization as colorless powder; yield: 2.8 mg (8%); TLC (CHCl₃/MeOH/AcOH 13:5:0.18, v/v/v) R_f 0.62; ESI-MS: *m/z* = 1762.0 [M-2H]²⁻; calcd for C₁₆₇H₃₁₉N₈O₆₆P: 1761.6 (most abundant signal for n=46). The chemical structure of NS-629 for is shown in Figure 2.

Example 2: Titration of Cathepsin B active site with lipidated inhibitor NS-629

Powdered lipidated inhibitor of Cathepsin B, NS-629, was dissolved in 0.1 M phosphate buffer, pH 6.0, containing 1 mM EDTA and 0.1% (v/v) PEG for final concentration of 0.05 μM. The kinetic reaction between Cathepsin B and its lipidated inhibitor was analyzed by continuous measurements of the loss of enzymatic activity at different concentration of inhibitor in the presence of fluorogenic substrate Z-Arg-Arg-AMC (AMC=7-amido-4-methylcoumarin) (Bachem). Inhibitor NS-629 in increasing concentrations (0.01–0.06 mM concentration), recombinant Cathepsin B (0.05 mM) and the dithiothreitol (DTT) (0.5 mM) were mixed in a plate with 0.1 M phosphate buffer, pH 6.0, containing 1 mM EDTA and 0.1% (v/v) PEG. After 15 minutes incubation at 37° C the inhibition kinetics of

Cathepsin B and NS-629 were determined. The reaction was started by the addition of 150 μ l of Cathepsin substrate Z-Arg-Arg-AMC solution and the kinetics of substrate hydrolysis was monitored continuously during 10 min by a TECAN plate reader at excitation and emission wavelengths of 370 and 460 nm, respectively. As can be seen in Figure 3a, the inhibitor bound to Cathepsin B with an apparent 1:1.5 stoichiometry, suggesting that Cathepsin B efficiently binds NS-629.

Example 3: Lipid vesicles linked with lipidated inhibitor preparation by extrusion

Aliquots of lipids (2.6 mM of egg phosphatidylcholine (Avanti Polar Lipids, Inc.) and 0.1 mM of lipidated inhibitor) supplied as chloroform solutions are placed into vials to form thin films by removing chloroform by evaporation under vacuum. Dry films are then hydrated by adding of the 0.1 mM phosphate buffer buffer pH 6.0. Dispersions are homogenized with vortex mixing and extruded under pressure through polycarbonate filters of decreasing pore diameter 0.1 μ m using extruder. Figure 4 shows the size distribution of liposomes functionalized with lipidated inhibitor as determined by dynamic light scattering (DLS). The average size was shown to be 85.66 nm.

Example 4: Lipid vesicles linked with lipidated inhibitor preparation by sonification

Aliquots of lipids (2.6 mM of egg phosphatidylcholine (Avanti Polar Lipids, Inc.) and 0.1 mM of lipidated inhibitor) supplied as chloroform solutions are placed into vials to form thin films by removing chloroform by evaporation under vacuum. Dry films are then hydrated by adding of the 0.1 mM phosphate buffer pH 6.0. Dispersions are homogenized with vortex mixing and then emulsified by sonication in bath sonicator during 20 min.

Example 5: Titration of Cathepsin B active site with liposomes labelled by lipidated inhibitor NS-629

The liposomes labelled by lipidated inhibitor were prepared as described above in Example 3. The kinetic reaction between Cathepsin B and liposomes labelled by lipidated Cathepsin inhibitor NS-629 was analyzed by continuous measurements of the loss of enzymatic activity at different concentration of inhibitor in the presence of fluorogenic substrate Z-Arg-Arg-AMC. Liposomes labelled by Cathepsin inhibitor NS-629 in increasing inhibitor concentrations (0.01–0.09 mM concentration) in increasing concentrations (0.01–0.06 mM concentration), recombinant Cathepsin B (0.05 mM) and the DTT (0.5 mM) were mixed in

a plate with 0.1 M phosphate buffer, pH 6.0, containing 1 mM EDTA and 0.1% (v/v) PEG. After 15 minutes incubation at 37° C the inhibition kinetics of Cathepsin B and NS-629 were determined. The reaction was started by the addition of 150 µl of Cathepsin substrate Z-Arg-Arg-AMC solution and the kinetics of substrate hydrolysis was monitored continuously during 10 min by a TECAN plate reader at excitation and emission wavelengths of 370 and 460 nm, respectively. As can be seen in Figure 3a, the liposomes labelled by Cathepsin inhibitor NS-629 bound to Cathepsin B with an apparent 1 : 2.5 stoichiometry, suggesting that Cathepsin B could efficiently bind liposomes labelled by Cathepsin inhibitor NS-629.

10

Example 6: *Ex vivo* binding of Alexa Fluor 555™ (Invitrogen) loaded liposomes labeled by lipidated inhibitor to mouse bone marrow-derived macrophages

Liposomes were prepared as: aliquots of lipids (2,6 mM of egg phosphatidylcholine (Avanti Polar Lipids, Inc.) and 0,1 mM of lipidated inhibitor) supplied as chloroform solutions are placed into vials to form thin films by removing chloroform by evaporation under vacuum. Dry films are then hydrated by adding of 0.1 mg Alexa Fluor 555™ (Invitrogen) containing 0.01 M phosphate buffer, pH 7.4. Active endocytosis of macrophages was stopped by incubation at 4° C during 15 minutes. Next, 200 µl of liposomes were placed on the cells and incubated for 15 minutes at 4° C. After incubation cells were washed by PBS and fluorescence intensity was examined with TECAN plate reader. Figure 4 shows that this experiment proves targeting of liposomes with lipidated Cathepsin inhibitor to the immune cells. Liposomes functionalized (NS-Lip-Alx) and not functionalized (Lip-Alx) with NS-629 were loaded with fluorescence marker (Alexa Fluor 555™ (Invitrogen)) and incubated with mouse bone marrow-derived macrophages for 15 minutes at 4° C. Fluorescence of accumulated marker was measured with TECAN plate reader. As a control liposomes without labeling were used.

15

20

25

Example 7: *Ex vivo* binding of Alexa Fluor 555™ (Invitrogen) loaded liposomes labeled by lipidated inhibitor to mouse bone marrow-derived macrophages

Liposomes were prepared as: aliquots of lipids (2.6 mM of egg phosphatidylcholine (Avanti Polar Lipids, Inc.) and 0,1 mM of lipidated inhibitor) supplied as chloroform solutions are placed into vials to form thin films by removing chloroform by evaporation under vacuum. Dry films are then hydrated by adding of 0.1 mg Alexa Fluor 555™

30

(Invitrogen) containing 0.01 M phosphate buffer, pH 7.4. Active endocytosis of macrophages was stopped by incubation at 4° C during 15 minutes. Next, 200 µl of liposomes were placed on the cells and incubated for 15 minutes at 4° C. After incubation cells were washed by PBS and examined with an Olympus fluorescent microscope (Olympus IX 81, Olympus) with Imaging Software for Life Science Microscopy Cell. Non-functionalized liposomes (Lip-Alx) and liposomes functionalized with NS-629 (NS-Lip-Alx) were examined with an Olympus fluorescence microscope (Olympus IX 81, Olympus) with Imaging Software for Life Science Microscopy Cell. Figure 6 shows representation of liposomes with lipidated Cathepsin inhibitor targeting efficiency in primary mouse immune cells.

Example 8: Encapsulation of Magnevist® (Bayer HealthCare Pharmaceuticals) (dimeglumine salt of Gd-DTPA; Bayer AG) into the liposome with following extrusion

Aliquots of lipids (2.6 mM of egg phosphatidylcholine (Avanti Polar Lipids, Inc.) and 0.1 mM of lipidated inhibitor) supplied as chloroform solutions are placed into vials to form thin films by removing chloroform by evaporation under vacuum. Dry films are then hydrated by adding of Magnevist® (Bayer HealthCare Pharmaceuticals). Dispersions are homogenized with vortex mixing and extruded under pressure through polycarbonate filters of decreasing pore diameter 0.1 µm using extruder. T₁-weighted MR images (TE=8.5 ms, TR=400 ms) of an orthotopic transplanted breast cancer mouse before, 1 and 24 hours after (T₁) intraperitoneal injection of 200 µl NS-629 labeled liposomes containing Magnevist® (Bayer HealthCare Pharmaceuticals) are shown in Figure 8. The tumour tissue possess negative MR signal on T₁-weighted images. The bright signal at 1 and 24 hours after injection in T₁-weighted MR image shows successful targeting of Magnevist® (Bayer HealthCare Pharmaceuticals) loaded NS-629 labelled liposomes.

Example 9: Encapsulation of D-luciferine into the liposome with following extrusion

Aliquots of lipids (2,6 mM of egg phosphatidylcholine (Avanti Polar Lipids, Inc.) and 0,1 mM of lipidated inhibitor) supplied as chloroform solutions are placed into vials to form thin films by removing chloroform by evaporation under vacuum. Dry films are then hydrated by adding of D-luciferine in PBS (15 mg/ml). Dispersions are homogenized with vortex mixing and extruded under pressure through polycarbonate filters of decreasing pore diameter 0.1 µm using extruder. Targeted delivery of liposomes labeled by lipidated

inhibitor carrying D-luciferin into transgenic mouse expressing luciferase (FVB.luc^{tg/+}) is shown in Figure 7. The high-intensity luciferase signal associated with the induced paw edema demonstrates selective accumulation of labelled liposomes in the inflammation area.

5

References

- Allen, T. M. et al., 1995, *Biochim Biophys Acta* 1237(2): 99-108.
- Altmann et al., 2002, *J. Med. Chem.* 45: 2352– 2354.
- Beste et al., 1999, *Proc. Natl. Acad. Sci. USA*, 96: 1898-1903.
- 10 Bieth J.G., 1995, *Methods Enzymol.*, 248: 59-84.
- Blume, G. et al., 1993, *Biochim Biophys Acta* 1149(1): 180-184.
- Bromme et al., 1989, *Biochem. J.* 263: 861– 866.
- Bromme et al., 1994, *Methods Enzymol.* 244: 671– 685.
- Brömme et al., 1996, *Biochem. J.* 315: 85– 89.
- 15 Cegnar M. et al., 2004, *Experimental Cell Research*, 301(2), 223-231.
- Chapman H.A. et al., 1997, *Annu Rev Physiol*, 59: 63–88.
- Coussens L.M. et al., 2002, *Science*; 295: 2387-92.
- Cruz et al., 2010, *Molecular Pharmaceutics* 8: 104-116.
- Deaton & Kumar, 2004, *Progress in Medicinal Chemistry*, 42: 245-375.
- 20 Duffy et al., 1999, *Bioorg. Med. Chem. Lett.* 9: 1907– 1910.
- Esser R.E. et al., 1994, *Arthritis Rheum.* 37: 236-47.
- Etherington D.J. et al., 1988, *Br J Exp Pathol*, 69: 281-9.
- Fahmy et al., 2007, *AAPS Journal* 9(2): E171-E180.
- Falgueyret et al., 2001, *J. Med. Chem.* 44: 94–104.
- 25 Ferrari, 2005, *Nature Reviews* 5: 161-171.
- Friedrichs, B. et al., 2003, *J Clin Invest* 111(11): 1733-1745.
- Gocheva, V. et al., 2010, *Genes Dev* 24(3): 241-255.
- Greenspan et al., 2001, *J. Med. Chem.* 44: 4524– 4534.
- Hansen T. et al., 2000, *J Rheumatol*, 27: 859-65.
- 30 Harmsen & De Haard, 2007, *Applied Microbiology and Biotechnology*, 77(1), 13-22.
- Holliger & Hudson, *Nature Biotech.*, 2005, 23(9): 1126-1136.
- Honey, K. et al., 2002, *Nat Immunol* 3(11): 1069-1074.
- Hill P.A. et al., 1994, *J Cell Biochem*, 56: 118-30.
- Huang et al., 2002, *J. Med. Chem.* 45: 676– 684.
- 35 Huet G. et al., 1993, *Arthritis Rheum*, 36: 772-80.

- Hylarides et al., 2001, *Bioconjugate Chemistry* 12, 421-427.
- Jaffer Farouc A. et al., 2009, *Arteriosclerosis Thrombosis and Vascular Biology*, 29(7), 1017.
- Jane, D. T. et al., 2006, *Biol Chem* 387(2): 223-234.
- 5 Jedeszko C. & Sloane B.F., 2004, *Biol Chem*, 385: 1017-27.
- Joyce J.A. et al., 2004, *Cancer Cell*, 5:443-53.
- Katunuma et al., 1999, *FEBS Lett.* 458: 6-10.
- Katunuma et al., 2000, *Biochem. Biophys. Res. Commun.* 267: 850-854.
- Katunuma et al., 2002, *Arch. Biochem. Biophys.* 397: 305-311.
- 10 Keyszer G. et al., 1998, *Arthritis Rheum*, 41: 1378-87.
- Lechner, A. M. et al., 2006, *J Biol Chem* 281(51): 39588-39597.
- Lemaire R. et al., 1997, *Br J Rheumatol*, 36: 735-43.
- Lenar et al., 1998, *Biol Chem Hoppe Seyler*, 369 Suppl: 257-61.
- Krantz in: A.J. Barrett (Ed.), *Methods Enzymol.*, vol. 244, Academic Press, New York, 15 1994, pp. 656-671.
- Liang et al., 2002, *Journal of Controlled Release* 78, 67-79.
- Link, J. O. et al., 2006, *Curr. Opin. Drug Discov. Devel.*, 9, 471.
- Liotta, L. A. & Kohn, E. C., 2001, *Nature* 411: 375-379.
- Mantovani A. et al., 2008, *Cancer-related inflammation. Nature* 454: 436-444.
- 20 Marquis et al., 1998, *J. Med. Chem.* 41: 3563-3567.
- Marquis et al., 1999, *Bioorg. Med. Chem.* 7: 581-588.
- Marquis et al. (1), 2001, *J. Med. Chem.* 44: 1380-1395.
- Marquis et al. (2), 2001, *J. Med. Chem.* 44: 725-736.
- McKerrow et al., 1999, *Bioorg. Med. Chem.* 7: 639-644.
- 25 Mirković B. et al., 2009, *FEBS J.*, ;276(17): 4739-51.
- Mohamed, M. M. & Sloane, B. F., 2006, *Nat Rev Cancer* 6(10): 764-775.
- Mueller, M. M. & Fusenig, N. E., 2004, *Nat. Rev. Cancer* 4(11): 839-849.
- Murdoch C, et al., 2008, *Nat Rev Cancer* 8: 618-63.
- Orive et al., 2009, *Nature Reviews Neuroscience* 10: 682-692.
- 30 Peer et al., 2007, *Nature Nanotechnology* 2: 751 - 760.
- Palmer et al., 1995, *J. Med. Chem.* 38: 3193-3196.
- Potts, W. et al., 2004, *Int J Exp Pathol* 85(2): 85-96.
- Quasba et al., 2008, *Biotechnology Progress* 24 (3), 520-526.
- Reinheckel, T. et al., 2005, *J Cell Sci* 118(Pt 15): 3387-3395.
- 35 Robichaud et al., 2003, *J. Med. Chem.* 46: 3709-3727.
- Rossi, A. et al., A., 2004, *Biol Chem* 385(5): 363-372.

- Rossin et al., 2005, *The Journal of Nuclear Medicine* 46: 1210-1218.
- Roth, W. et al., 2000, *FASEB J* 14(13): 2075-2086.
- Saegusa et al., 2002, *J. Clin. Invest.* 110: 361– 369.
- Schaschke et al., 1996, *FEBS Lett.*, 391: 297-301.
- 5 Schaschke et al., 1998, *J. Am. Chem. Soc.*, 1998: 7030-7038.
- Schaschke et al. (1), 2000, *FEBS Lett.* 482: 91.
- Schaschke et al. (2), 2000, *Bioorganic & Medicinal Chemistry Letters*, 10: 677-680.
- Schedel J. et al., 2004, *Gene Ther.*, 11: 1040-7.
- Seyfried et al., 2001, *Brain Res.* 901: 94– 101.
- 10 Shaw, 1994, *Methods Enzymol.* 244: 649– 656.
- Shuvaev et al., 2004, *Methods in Molecular Biology* 283: 3-19.
- Skerra (1), 2000, *Biochim. Biophys. Acta*, 1482: 337-50.
- Skerra (2), 2000, *J. Mol. Recognit.*, 13: 167-187.
- Smith et al., 2001, *Bioorg. Med. Chem. Lett.*, 11: 2951–2954.
- 15 Sun et al., 2008, *Advanced Drug Delivery Reviews* 60, 1252-1265.
- Tavares et al. (1), 2004, *J. Med. Chem.* 47: 588– 599.
- Tavares et al. (2), 2004, *J. Med. Chem.* 47: 5049–5056.
- Tavares et al. (3), 2004, *J. Med. Chem.* 47: 5057– 5068.
- Thompson et al., 1997, *Proc. Natl. Acad. Sci. U. S. A.* 94: 14249–14254.
- 20 Thompson et al., 1998, *J. Med. Chem.* 41: 3923–3927.
- Thurmond et al. (1), 2004, *Pharmacol. Exp. Ther.*, 308: 268.
- Thurmond et al. (2), 2004, *J. Pharmacol. Exp. Ther.* 308: 268– 276.
- Thurmond et al. (3), 2004, *J. Med. Chem.* 47: 4799– 4801.
- Turk, B. et al., 2000, *Biochim Biophys Acta* 1477(1-2): 98-111.
- 25 Turk V., et al., 2004, *Cancer Cell*, 5: 409–10.
- Vasiljeva, O. et al., 2006, *Cancer Res* 66(10): 5242-5250.
- Vasiljeva, O. et al., 2007, *Current Pharmaceutical Design* 13: 385-401.
- Vasiljeva, O. et al., *Curr. Pharm. Des.*, 2007, 13: 387
- Ward et al., 2002, *J. Med. Chem.* 45: 5471– 5482.
- 30 Wolters, P. J. et al., 2001, *J Biol Chem* 276(21): 18551-18556.
- Yasuda et al., 2005, *Advanced Drug Delivery Reviews*, 57: 973– 993.
- Yasuma et al., 1998, *J. Med. Chem.* 41: 4301– 4308.
- Zhou et al., 2001, *Arkivoc VI*, 116– 121.
- Zhou et al., 2002, *Bioorg. Med. Chem. Lett.* 12: 3417–3419.
- 35 Zhou et al., 2003, *Bioorg. Med. Chem. Lett.* 12: 139-141.

Claims

1. A Cathepsin-binding compound bound to a carrier comprising a diagnostic moiety,
 - (i) for use in the diagnosis of inflammatory diseases,and/or
 - (ii) for use in the diagnosis of neoplastic diseases,wherein the Cathepsin-binding compound binds to inflammatory cells of the tumour stroma.
2. The Cathepsin-binding compound for use according to claim 1, wherein
 - (a) the Cathepsin bound by the compound is selected from Cathepsin (Cts) B, C, H, F, K, L, O, S, V, W, and X/Z, in particular Cts B, and/or S, more preferably Cts B,
 - (b) wherein the Cathepsin is mammalian Cathepsin, in particular human Cathepsin.
3. The Cathepsin-binding compound for use according to claim 1 or 2, wherein the compound specifically binds to Cts B.
4. The Cathepsin-binding compound for use according to any of the previous claims, wherein the carrier is selected from a nanotube, a liposome, a lipoplex, a polymersome, a micell, a nanogel, a nanoparticle, a mesoporous silica particle, a dendrimer, and a nanoshell, in particular the carrier is a liposome, or other chemical entity that could be linked to the Cathepsin-binding compound.
5. The Cathepsin-binding compound for use according to any of the preceding claims, wherein the Cathepsin-binding compound
 - (i) is selected from an antibody or antibody fragment or protein or compound specifically binding to Cathepsin, a synthetic inhibitor of Cathepsin, a peptide, and an antibody mimetic, and/or
 - (ii) an inhibitor of the protease activity of a Cathepsin.
6. The Cathepsin-binding compound for use any of claims, wherein the use is a diagnostic use, and wherein the detection of a Cathepsin *in vivo* is indicative of an inflammatory site.
7. The Cathepsin-binding compound for use according to claim 6, wherein the diagnostic moiety is selected from


- a radioactive label,
in particular technetium-99m, iodine-123, iodine-131, rhenium-186,
rhenium-188, gallium-67, yttrium-90, and lutetium-177,
a paramagnetic agent,
5 in particular a Gadolinium-complex, more preferably Gd-DTPA,
a super paramagnetic nanoparticle,
in particular iron oxide based nanoparticles,
a PET-imageable agent,
an MRI-imageable agent,
10 in particular a Gadolinium-complex, more preferably Gd-DTPA,
a fluorophore,
in particular Alexa Fluor,
a chromophore,
a phosphorescing agent,
15 a chemiluminescent agent, and
a bioluminescent agent.

8. The Cathepsin-binding compound for use according to any of the preceding claims,
wherein the neoplastic disease is breast cancer and metastasis thereof.
20
9. The Cathepsin-binding compound for use according to any of claims 1 to 7,
wherein the inflammatory disease is selected from psoriasis, contact dermatitis,
atopic dermatitis, alopecia areata, erythema multiforme, dermatitis herpetiformis,
scleroderma, vitiligo, hypersensitivity angiitis, urticaria, bullous pemphigoid, lupus
25 erythematosus, pemphigus, epidermolysis bullosa acquisita, inflammatory bowel
disease, rheumatoid arthritis, juvenile rheumatoid arthritis, psoriatic arthritis,
osteoarthritis, refractory rheumatoid arthritis, chronic non-rheumatoid arthritis,
osteoporosis/bone resorption, Crohn's disease, septic shock, endotoxic shock,
atherosclerosis, ischaemia-reperfusion injury, coronary heart disease, vasculitis,
30 amyloidosis, multiple sclerosis, sepsis, chronic recurrent uveitis, hepatitis C virus
infection, malaria, ulcerative colitis, cachexia, plasmocytoma, endometriosis,
Behcet's disease, Wegener's granulomatosis, AIDS, HIV infection, autoimmune
disease, immune deficiency, common variable immunodeficiency (CVID), chronic
graft-versus-host disease, trauma and transplant rejection, adult respiratory distress
35 syndrome, pulmonary fibrosis, recurrent ovarian cancer, lymphoproliferative
disease, refractory multiple myeloma, myeloproliferative disorder, diabetes,
juvenile diabetes, meningitis, viral infections, cardiovascular diseases, ankylosing
spondylitis, skin delayed type hypersensitivity disorders, Alzheimer's disease,

systemic lupus erythematosus and allergic asthma, preferably selected from rheumatoid arthritis, Crohn's disease, ulcerative colitis, septic shock, psoriasis and atherosclerosis.

- 5 10. A Cathepsin-binding and liposome-binding compound, in particular a Cathepsin B-binding and liposome-binding compound or Cathepsin S-binding and liposome-binding compound, more preferably a Cathepsin B-binding and liposome-binding compound, comprising:
- 10 (i) a Cathepsin-binding moiety, in particular a Cathepsin B-binding moiety or Cathepsin S-binding moiety, more preferably a Cathepsin B-binding compound,
- (ii) optionally a linker, and
- (iii) a liposome-binding moiety.
- 15 11. The Cathepsin-binding and liposome-binding compound of claim 10, comprising a linker, wherein the linker preferably comprises a functionalized PEG moiety and/or a diaminoalkyl moiety.
- 20 12. The Cathepsin-binding and liposome-binding compound of claim 10 or 11, wherein the linker has the structure selected of formulas II-V:
- NH-(CH₂)_{l+1}-(NH)-C(O)-(CH₂)_m-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (II)
- NH-(CH₂)_{m+1}-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (III),
- C(O)-(CH₂)_l-(NH)-C(O)-(CH₂)_m-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (IV),
- and/or
- 25 -C(O)-(CH₂)_m-O-[CH₂-CH₂-O]_p-C(O)-NH-(CH₂)_q- (V)
- wherein
- l is an integer from n= 1 to 30, preferably 3 to 20, most preferably n=5,
- m is an integer from n=1 to 30, more preferably 1 to 10, most preferably 1,
- p is an integer from n=1 to 200, more preferably 15 to 100, most preferably 38
- 30 to 53,
- q is an integer from n=2 to 30, more preferably 2 to 10, most preferably 2.
- 35 13. The Cathepsin binding and liposome-binding compound of any of claims 10 to 12, wherein the liposome-binding moiety, is a phospholipid, in particular a phospholipid comprising 1, 2 or more stearyl moieties, 1, 2 or more oleyl moieties, 1, 2 or more lauroyl moieties, 1, 2 or more myristoyl moieties, 1, 2 or more palmitoyl moieties, and/or 1, 2 or more arachinoyl moieties, more preferably a phospholipid comprising 2 stearyl moieties.

14. The Cathepsin binding and liposome-binding compound of any of claims 10 to 13, having the structure of formula (XIII):



5 wherein $n=38$ to 53 .

15. The Cathepsin-binding and liposome-binding compound of any of claims 10 to 14, wherein the Cathepsin-binding moiety is an inhibitor of the protease activity of Cathepsin, in particular a peptide inhibitor.

10

16. The Cathepsin-binding and liposome-binding compound of any of claims 10 to 15, wherein the Cathepsin-binding moiety is selected from the group consisting of

-Gly-Aaa-Bbb-X-Ccc-Ddd (VI)

-Gly-Aaa-Bbb-X-Ccc (VII)

15

and

-Gly-Aaa-Bbb-Y (VIII),

wherein

X is an electrophilic unit, in particular an epoxide-derivative or a semicarbazide-derivative, preferably tEps, $-(NH)-(NH)-C(O)-$, most preferably (2S,3S)-tEps;

20


Y is an electrophilic unit, in particular an epoxide-derivative, a semicarbazide-derivative or a nitrile-derivative, preferably -tEps, $-(NH)-(NH)-C(O)-NH_2$, $-(NH)-CH_2-CN$, more preferably (2S,3S)-tEps;

Aaa is selected from the group consisting of Gly, Phe, Asp, Glu, 1-aminoadipic acid, in particular Gly;


25

Bbb is selected from the group consisting of aliphatic or aromatic hydrophobic


amino acids, in particular Leu, Ile, Phe, Tyr, Val,




(IX),



(X),



(XI),



(XII),

2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, preferably amino acids (IX), (X), (XI), (XII), 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, Leu, Ile, Phe, Tyr, more preferably amino acids (IX), (X), (XI), (XII), 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, Leu and Ile;

Ccc is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Phe, Tyr, Val, Trp, Nle, preferably Nle and Leu, more preferably Leu;

Ddd is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids and amino compounds, in particular Trp, Val, Ile, Phe, Tyr, Ala, Ser, Thr, Pro, $-\text{NH}(\text{CH}_2)_n=0-4-\text{CH}_3$, $-\text{NH}-\text{CH}(\text{CH}_3)_2$, $-\text{NH}-\text{CH}_2-\text{CH}(\text{CH}_3)_2$, $-\text{NH}-(\text{CH}_2)_n=1-2\text{Ph}$, preferably, Trp, Val, Ile, Phe, Tyr, more preferably Trp.

17. The Cathepsin-binding and liposome-binding compound according to any of claims 10 to 16, wherein the Cathepsin-binding moiety is a Cathepsin B-binding moiety and is an inhibitor of the protease activity of Cathepsin B, in particular a peptide inhibitor, preferably a peptide inhibitor selected from the group consisting of

-Gly-Aaa-Bbb-X-Ccc-Ddd (VI),

and

-Gly-Aaa-Bbb-X-Ccc (VII)

wherein

X is an electrophilic unit, in particular an epoxide-derivative or a semicarbazide-derivative, preferably tEps, $-(\text{NH})-(\text{NH})-\text{C}(\text{O})-$, most preferably (2S,3S)-tEps;

Aaa is Gly;

Bbb is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Phe, Tyr, Val, preferably Leu, Ile, Phe, Tyr, more preferably Leu and Ile;

Ccc is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Val, Nle, preferably Nle and Leu, more preferably Leu;

Ddd is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids and amino compounds, in particular Trp, Val, Ile, Phe, Tyr, Ala, Ser, Thr, Pro, preferably, Trp, Val, Ile, Phe, Tyr, more preferably Trp.

18. The Cathepsin-binding and liposome-binding compound according to claim 17, wherein the Cathepsin B-binding moiety is -Gly-Gly-Leu-(2S,3S)-tEps-Leu-Pro-OH.

19. The Cathepsin-binding and liposome-binding compound according to any of claims 10 to 16, wherein the Cathepsin-binding moiety is a Cathepsin S-binding moiety and is an inhibitor of the protease activity of Cathepsin S, in particular a peptide inhibitor, preferably a peptide inhibitor selected from the group consisting of

-Gly-Aaa-Bbb-X-Ccc-Ddd (VI),

-Gly-Aaa-Bbb-X-Ccc (VII)

and

-Gly-Aaa-Bbb-Y (VIII),

wherein


X is an electrophilic unit, in particular an epoxide-derivative or a semicarbazide-derivative, preferably tEps, -(NH)-(NH)-C(O)-, most preferably (2S,3S)-tEps;

Y is an electrophilic unit, in particular an epoxide-derivative, a semicarbazide-derivative or a nitrile-derivative, preferably tEps, -(NH)-(NH)-C(O)-NH₂, -(NH)-CH₂-CN, more preferably (2S,3S)-tEps;


Aaa is selected from the group consisting of Gly, Phe, Asp, Glu, 1-aminoadipic acid, in particular Gly;

Bbb is selected from the group consisting of aliphatic or aromatic hydrophobic


amino acids, in particular Leu, Ile, Phe, Tyr, Val,




(IX),



(X),



(XI),



(XII),

2-naphthyl alanine, cyclohexyl alanine, 4-

bromophenyl alanine, preferably amino acids (IX), (X), (XI), (XII), 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine, Leu, Ile, Phe, Tyr, more preferably amino acids 1-4, 2-naphthyl alanine, cyclohexyl alanine, 4-bromophenyl alanine and Leu;

5 Ccc is selected from the group consisting of aliphatic or aromatic hydrophobic amino acids, in particular Leu, Ile, Phe, Tyr, Val, Trp, preferably Leu and Ile, more preferably Leu;

10 Ddd is an amino compound, in particular an alkyl amino compound, an alkaryl amino compound or an aryl amino compound, preferably -NH(CH₂)_{n=0-4}-CH₃, -NH-CH(CH₃)₂, -NH-CH₂-CH(CH₃)₂, -NH-(CH₂)_{n=1-2}Ph.

20. A Cathepsin targeting compound, in particular a Cathepsin B-Targeting compound or Cathepsin S-targeting compound, comprising a Cathepsin binding and liposome-binding compound of any of claims 10 to 18, bound to a liposome.

15

21. The Cathepsin targeting compound according to claim 20, further comprising a diagnostic moiety.

20

22. The Cathepsin targeting compound according to any of claims 20 to 21, comprising a diagnostic moiety,

wherein the diagnostic moiety is preferably selected from:

a radioactive label,

a paramagnetic agent,

in particular a Gadolinium-complex, more preferably Gd-DTPA,

25

a super paramagnetic nanoparticle,

in particular iron oxide based nanoparticles,

a PET-imageable agent,

an MRI-imageable agent,

in particular a Gadolinium-complex, more preferably Gd-DTPA,

30

a fluorophore,

in particular Alexa Fluor,

a chromophore,

a phosphorescing agent,

a chemiluminescent agent, and

35

a bioluminescent agent.

23. The Cathepsin targeting compound according to any of claims 20 to 22,

(i) for use in the diagnosis of inflammatory diseases,

and/or

- (ii) for use in the diagnosis of neoplastic diseases, wherein the Cathepsin-targeting compound binds to inflammatory cells of the tumour stroma, in particular in breast cancer.

5

- 24. A method for preparing a Cathepsin-binding and liposome-binding compound according to any of claims 10 to 19, comprising providing a Cathepsin-binding moiety and associating a liposome-binding moiety, in particular associating a linker and a liposome-binding moiety with the Cathepsin-binding moiety.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7





Figure 8



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/002652

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A61K47/48 A61K49/00 A61K49/18 A61P35/00 A61P29/00
 ADD. A61K9/127 C07D405/12 C07K7/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 A61K A61P C07K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHASCHKE N N ET AL: "beta-Cyclodextrin/epoxysuccinyl peptide conjugates: a new drug targeting system for tumor cells", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 10, no. 7, 1 April 2000 (2000-04-01), pages 677-680, XP004191853, PERGAMON, ELSEVIER SCIENCE, GB ISSN: 0960-894X, DOI: 10.1016/S0960-894X(00)00078-0	10,11, 15-19,24
Y	page 677, right-hand column, lines 2-6 page 678; figure 1 page 679, right-hand column, paragraph 2 ----- -/--	14,18

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search 18 July 2012	Date of mailing of the international search report 30/07/2012
---	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Villard, Anne-Laure
--	---

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/002652

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHASCHKE N ET AL: "Epoxy succinyl peptide-derived affinity labels for cathepsin B", FEBS LETTERS, vol. 482, no. 1-2, 29 September 2000 (2000-09-29), pages 91-96, XP004337627, ELSEVIER, AMSTERDAM, NL ISSN: 0014-5793, DOI: 10.1016/S0014-5793(00)02047-0	1-11, 15-19,24
Y	page 91, left-hand column, paragraph 1 page 93; compounds 6,7 page 95, section "Conclusions"	14,18
X	JAFFER FAROUC A ET AL: "Optical and Multimodality Molecular Imaging Insights Into Atherosclerosis", ARTERIOSCLEROSIS THROMBOSIS AND VASCULAR BIOLOGY, vol. 29, no. 7, July 2009 (2009-07), pages 1017-1024,1-12, XP002661291, ISSN: 1079-5642 figure 1 pages 1017-1018, section "Imaging of Protease Activity" Supplement material: 1st page - 2nd page, 1st paragraph	1-11,24
X	CEGNAR M ET AL: "Poly(lactide-co-glycolide) nanoparticles as a carrier system for delivering cysteine protease inhibitor cystatin into tumor cells", EXPERIMENTAL CELL RESEARCH, vol. 301, no. 2, 10 December 2004 (2004-12-10), pages 223-231, XP004626708, ACADEMIC PRESS, US ISSN: 0014-4827, DOI: 10.1016/J.YEXCR.2004.07.021 abstract	1-11,15, 24
X	US 2005/002865 A1 (KLAIVENESS JO [NO] ET AL) 6 January 2005 (2005-01-06) paragraphs [0014], [0028], [0029] page 14, line referring to antibodies targeting Cathepsin B	1-13, 15-17, 19,24
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/002652

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/096367 A2 (TARGESOME INC [US]) 5 December 2002 (2002-12-05)	1-13, 15-17, 19-24
Y	page 14, lines 20-21 page 16, lines 1-18 page 18, lines 1-7 page 31, line 31 - page 32, line 21 page 38, lines 5-8 examples 44,45 figure 33 claims 1,21,34	14,18
X	----- GREENBAUM D ET AL: "Epoxide electrophiles as activity-dependent cysteine protease profiling and discovery tools.", CHEMISTRY & BIOLOGY AUG 2000 LNKD- PUBMED:11048948, vol. 7, no. 8, August 2000 (2000-08), pages 569-581, XP002680233, ISSN: 1074-5521 figures 1,2,6	1-11
A,P	----- MIKHAYLOV G ET AL: "Ferri-liposomes as an MRI-visible drug-delivery system for targeting tumours and their microenvironment", NATURE NANOTECHNOLOGY 2011 NATURE PUBLISHING GROUP GBR LNKD- DOI:10.1038/NNANO.2011.112, vol. 6, no. 9, September 2011 (2011-09), pages 594-602, XP002680234, ISSN: 1748-3387 abstract supplementary methods, "preparation of ferri-liposomes" -----	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/002652

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2005002865	A1	06-01-2005	
		US 6331289 B1	18-12-2001
		US 2002102217 A1	01-08-2002
		US 2005002865 A1	06-01-2005

WO 02096367	A2	05-12-2002	
		AU 2002314861 A1	09-12-2002
		EP 1401818 A2	31-03-2004
		JP 2004535407 A	25-11-2004
		WO 02096367 A2	05-12-2002
