



(72) URBAN, ROBERT G., US

(72) CHICZ, ROMAN M., US

(72) COLLINS, EDWARD J., US

(72) HEDLEY, MARY LYNNE, US

(71) ZYCOS INC., US

(51) Int.Cl.⁶ A61K 39/12, C07H 21/04, C12N 15/00, C12N 7/00, A61K 39/00

(30) 1997/10/09 (08/948,378) US

(54) **PEPTIDES IMMUNOGENES ISSUS DE LA PROTEINE HPV E7**

(54) **IMMUNOGENIC PEPTIDES FROM THE HPV E7 PROTEIN**

(57) L'invention concerne des peptides immunogènes issus de la protéine HPV E7 du type 16 qui comprennent des épitopes chevauchants restreints de la cellule T de classe I. L'invention concerne également des procédés qui permettent d'administrer à un mammifère hôte des molécules d'ADN codant ces peptides.

(57) The invention provides immunogenic peptides from the HPV type 16 E7 protein that comprise overlapping class I restricted T cell epitopes. Also disclosed are methods of administering DNA molecules encoding these peptides to a host mammal.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 39/12, 39/00, C12N 15/00, 7/00, C07H 21/04</p>	A1	<p>(11) International Publication Number: WO 99/18995</p> <p>(43) International Publication Date: 22 April 1999 (22.04.99)</p>
<p>(21) International Application Number: PCT/US98/21456</p> <p>(22) International Filing Date: 9 October 1998 (09.10.98)</p> <p>(30) Priority Data: 08/948,378 9 October 1997 (09.10.97) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/948,378 (CON) Filed on 9 October 1997 (09.10.97)</p> <p>(71) Applicant (for all designated States except US): PANGAEA PHARMACEUTICALS, INC. [US/US]; 763 East Concord Avenue, Cambridge, MA 02138 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): URBAN, Robert, G. [US/US]; 51 Follen Road, Lexington, MA 02173 (US). CHICZ, Roman, M. [US/US]; 4 Cottage Street, Belmont, MA 02178 (US). COLLINS, Edward, J. [US/US]; 103 John Martin Court, Carrboro, NC 27510 (US). HEDLEY, Mary, Lynne [US/US]; 51 Follen Road, Lexington, MA 02173 (US).</p>	<p>(74) Agent: FRASER, Janis, K.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: IMMUNOGENIC PEPTIDES FROM THE HPV E7 PROTEIN</p> <p>(57) Abstract</p> <p>The invention provides immunogenic peptides from the HPV type 16 E7 protein that comprise overlapping class I restricted T cell epitopes. Also disclosed are methods of administering DNA molecules encoding these peptides to a host mammal.</p>		

- 1 -

IMMUNOGENIC PEPTIDES FROM THE HPV E7 PROTEINBackground of the Invention

This invention relates to treatment of human
5 papilloma virus (HPV) infection.

Papilloma viruses are non-enveloped DNA viruses
with a double stranded circular genome of approximately
8,000 bp. Over 75 types of human papilloma viruses (HPV)
have been typed at the DNA level, and these can be
10 broadly grouped into families on the basis of their
tissue tropism.

Histologic, molecular, and epidemiologic evidence
have implicated some HPV strains in cervical dysplasia
and cervical cancer. Many studies support the view that
15 most moderate and severe cervical intraepithelial
neoplasias (CIN) contain HPV DNA which is exclusively
detected in the histologically abnormal epithelium of
these lesions. Persistent infection with HPV is
believed to be the predominant risk factor for
20 development of cervical carcinoma. HPV DNA is readily
found in episomal form within cells exhibiting a
cytopathic effect, while the HPV DNA is found integrated
within the chromosomes of cells associated with most high
grade precancerous lesions and cancer. Approximately 23
25 HPV types are commonly found in anogenital screening
programs, but only 10-15 are associated with progressive
disease. Type 16 is the type most commonly found in
cervical cancer tissue.

Papillomaviruses contain nine open reading frames.
30 HPV genes with transforming properties have been mapped
to open reading frames E6 and E7. Substantial
biochemical work has demonstrated that the HPV E6 protein
inactivates the protein p53, whereas the E7 protein
interferes with retinoblastoma (Rb) protein function.
35 Since p53 and Rb are tumor-suppressor proteins which
function as cell division inhibitors, their inactivation

- 2 -

by E6 and E7 leads the cell to enter into S phase of the cell cycle. Expression of E6 and E7 is sufficient to immortalize some primary cell lines, and blocking E6 or E7 function has been shown to reverse the transformed
5 state.

Summary of the Invention

The invention is based on the discovery that a 13 amino acid peptide from the HPV strain 16 E7 protein that contains overlapping class I HLA binding, T cell
10 epitopes can induce a CTL response in an animal. Accordingly, the invention includes an immunogenic peptide having within its sequence multiple class I MHC-binding epitopes from a human papillomavirus (HPV) protein, and which has a length of less than 19 amino
15 acids and includes the sequence of Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:16) (hereinafter "immunogenic peptide"). The immunogenic peptide can optionally include sequences in addition to those derived from the E7 protein.

20 The immunogenic peptide can have the sequence of Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:3) or Xaa Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys, Xaa being Met, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu (SEQ ID NO:19), e.g., Ala Leu Met Gly Thr Leu
25 Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:4).

The invention also includes the peptides Thr Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:20) and Gly Thr Leu Gly Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:21), as well as Xaa Thr Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:27) and
30 Gly Thr Leu Gly Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:28), Xaa being Met, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu.

In addition, all of the peptides discussed herein may include additional amino acids to facilitate

- 3 -

expression, e.g., an amino terminal methionine to facilitate translation.

The invention also includes a polypeptide having the sequence of a first peptide linked to a second peptide by a peptide bond. The first peptide (which can be at the carboxy terminus or the amino terminus of the second peptide, so long as it functions in that site) is a peptide which controls intracellular trafficking of a peptide to which it is attached, and the second peptide is the immunogenic peptide described above. The polypeptide may optionally be modified to introduce an amino acid substitution at the junction between the first and second peptides to promote cleavage of the first and second peptides by a signal peptidase.

The trafficking peptides can be any recognized signal sequence, e.g. a signal sequence from the adenovirus E3 protein. A preferred trafficking peptide is the signal peptide of HLA-DR α , Met Ala Ile Ser Gly Val Pro Val Leu Gly Phe Phe Ile Ile Ala Val Leu Met Ser Ala Gln Glu Ser Trp Ala (SEQ ID NO:18).

The invention in addition includes a therapeutic composition containing the immunogenic peptide described above, and a pharmaceutically acceptable carrier. The polypeptide can optionally be formulated in a microparticle, a liposome or an immune-stimulating complex (ISCOM) (which may contain saponin alone as the active ingredient), or any other vehicle suitable for delivering into subjects the immunogenic peptides of the invention. When a microparticle is used, it preferably has a polymeric matrix that is a copolymer such as poly-lactic-co-glycolic acid (PLGA).

An immune response (e.g., a cellular immune response, including an MHC class I-mediated or class II-mediated immune response) in a mammal can be elicited by administering the immunogenic peptide to a mammal, e.g.,

- 4 -

a human, non-human primate, dog, cat, rabbit, cow, mouse, rat, guinea pig, or hamster, that has an MHC molecule that binds to the immunogenic peptide. The immunogenic peptide can be administered as part of a microparticle, liposome, or ISCOM, or in solution.

Another way to administer the peptide utilizes a nucleic acid, e.g., an expression vector, comprising a coding sequence encoding the immunogenic peptide. The nucleic acid can optionally encode a signal sequence linked to the immunogenic peptide, as described above. When the nucleic acid encodes such a signal sequence, it is preferred that it encodes the signal sequence from HLA-DR α (SEQ ID NO:18). In such a case, the immunogenic peptide can have the sequence, for example, of SEQ ID NO:4 or SEQ ID NO:3. Preferably, the nucleic acid does not include sequences from a viral genome that would render the nucleic acid infectious, and does not encode an intact E7 protein.

The nucleic acid described above can be included in a plasmid, optionally provided in a microparticle that also includes a polymeric matrix. In preferred embodiments, the polymeric matrix consists essentially of a copolymer of PLGA. The microparticle preferably has a diameter of, e.g., 0.02 to 20 microns, or less than about 11 microns. A plurality of microparticles preferably has diameter of, e.g., 0.02 to 20 microns, or less than about 11 microns

Also within the invention is a cell containing the plasmid of the invention. The cell can, e.g., be a B cell or other antigen presenting cell (APC). The cell may be cultured or otherwise maintained under conditions permitting expression of the peptide from the plasmid encoding it.

The nucleic acid and plasmid of the invention are useful in a method of inducing an immune response in a

- 5 -

mammal, e.g., a human, by administering the above-described plasmid to the mammal, e.g., as "naked DNA". The mammal may be at risk for, or suffer from, HPV infection, cervical dysplasia, and/or cervical cancer.

5 The nucleic acids and plasmids of the invention can also be incorporated into microparticles, liposomes, ISCOMS, or any other suitable delivery vehicle as described above.

The invention further includes a plasmid having a
10 sequence essentially identical to that of pBIOTOPE_{HPV} (SEQ ID NO:7), or a microparticle consisting essentially of a PLGA polymeric matrix and the pBIOTOPE_{HPV} plasmid, as well as methods of inducing an immune response in a mammal by
15 administering either the plasmid alone, or the plasmid incorporated into such a microparticle, to the mammal.

By a "substantially pure polypeptide" is meant a polypeptide which is separated from those components (proteins and other naturally-occurring organic
20 molecules) which naturally accompany it. Typically, the polypeptide is substantially pure when it constitutes at least 60%, by weight, of the protein in the preparation. Preferably, the protein in the preparation consists of at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, of an immunogenic
25 peptide.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The preferred methods
30 and materials for practicing the invention are described below, although other methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention. All publications, patent applications, patents, and other
35 references mentioned herein are incorporated by reference

- 6 -

in their entirety. In case of conflict, the present application, including definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting.

5 Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

Brief Description of the Drawings

Fig. 1 is a schematic drawing of the pBIOTOPE_{HPV} plasmid.
10

Fig. 2 is a graph showing the results of subjecting cells to triple color flow cytometry staining for CD8, CD16, and interferon-gamma.

Fig. 3 is a graph showing CTL lysis of an HLA-A2⁺, HPV16⁺ cell line with T cells from an HLA-A2⁺ donor stimulated with an influenza peptide (-■-), the A2.1 peptide (-●-), or the A2.4-C peptide (-▲-).
15

Fig. 4 is a graph showing CTL lysis of an HLA-A2⁺, HPV16⁺ cell line with T cells from a second HLA-A2⁺ donor stimulated with an influenza peptide (-■-), the A2.1 peptide (-●-), or the A2.4 peptide (-▲-).
20

Detailed Description

The peptides disclosed herein, and the nucleic acids encoding the peptides, can be used to elicit an immune response against the HPV E7 protein. The peptides were identified in part based on their binding affinity with the MHC class I HLA-A2 allele. Thus, the immune response elicited by these peptides is likely to be class I-mediated but may also involve class II mediated responses, B cell responses, or NK cell responses. The immune response can thus involve, e.g., cells expressing MHC class I molecules or cells expressing MHC class II molecules. The immune response can also include immune
25
30

- 7 -

cells such as macrophages, polymorphonuclear monocytes (PMN), natural killer cells, and B cells.

Five immunogenic peptides derived from the HPV type 16 E7 protein are shown in Table I. Peptide A2.1/4, 5 Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:3), corresponds to amino acids 82-94 in the HPV Type 16 E7 protein and includes the overlapping sequences of peptides A2.1, Leu Leu Met Gly Thr Leu Gly Ile Val (SEQ ID NO:1), A2.4, Thr Leu Gly Ile Val Cys Pro Ile Cys 10 (SEQ ID NO:2) A2.4-C, Thr Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:20), and A2.5, Gly Thr Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:21). Thus, peptide A2.1/4 has at least four overlapping epitopes potentially recognized by class I MHC restricted T cells.

15 **Table I. Amino acid sequences of conserved, class I-MHC binding, TCR binding HPV strain 16 E7 peptides**

	A2.1	LLMGTLGIV	(SEQ ID NO:1)
	A2.4	TLGIVCPIC	(SEQ ID NO:2)
20	A2.1/4	LLMGTLGIVCPIC	(SEQ ID NO:3)
	A2.4-C	TLGIVCPI	(SEQ ID NO:20)
	A2.5	GTLGIVCPI	(SEQ ID NO:21)

A peptide of the invention may optionally include 25 one having the amino acids SQK added to the carboxy terminus of the A2.1/4 peptide sequence ("the extended peptide"). Processing of the extended peptide can generate the peptide IVCPICSQK (SEQ ID NO:22), which has been reported as binding the MHC class I molecules HLA-A3 30 and HLA-A11 (Kast et al., J. Immunol. 152:3904-11, 1994). This region of the HPV E7 protein has several peptides that can be processed into MHC binding peptides. Additional extensions to the amino or carboxy terminus of

- 8 -

the A2.1/4 peptide may further increase the number of peptides that can be generated from this region of the E7 protein.

The peptides of the invention can be linked to a
5 trafficking sequence that directs the peptides to a
desired intracellular compartment. A trafficking
sequence is an amino acid sequence which functions to
control intracellular trafficking (directed movement from
organelle to organelle or to the cell surface) of a
10 polypeptide to which it is attached. Such trafficking
sequences might traffic the polypeptide to ER, a
lysosome, or an endosome, and include signal peptides
(the amino terminal sequences which direct proteins into
the ER during translation), ER retention peptides such as
15 KDEL (SEQ ID NO:20), and lysosome-targeting peptides such
as KFERQ (SEQ ID NO:21), QREFK (SEQ ID NO:22), and other
pentapeptides having Q flanked on one side by four
residues selected from K, R, D, E, F, I, V, and L.

Short amino acid sequences can act as signals to
20 target proteins to specific intracellular compartments.
For example, hydrophobic signal peptides are found at the
amino terminus of proteins destined for the ER, while the
sequence KFERQ (SEQ ID NO:21) (and other closely related
sequences) is known to target intracellular polypeptides
25 to lysosomes, while other sequences target polypeptides
to endosomes.

One such trafficking sequence is the HLA-DR α
leader sequence, Met Ala Ile Ser Gly Val Pro Val Leu Gly
Phe Phe Ile Ile Ala Val Leu Met Ser Ala Gln Glu Ser Trp
30 Ala (SEQ ID NO:18). The signal peptide may include only
a portion (e.g., at least ten amino acid residues) of the
specified 25 residue sequence, provided that portion is
sufficient to cause trafficking of the polypeptide to the
ER.

- 9 -

In some cases it is desirable to modify the portion of the peptide spanning the trafficking sequence and the sequence encoding the HPV E7 antigenic peptide to facilitate processing, i.e., cleavage, by the signal
5 peptidase. Recognition sequences for signal peptides are described in Von Heijne, NAR 14:4683, 1986.

Standard techniques can be used to construct a DNA encoding the antigenic peptide (see, e.g., the techniques described in WO 94/04171). The construct may include
10 additional sequences for enhancing expression in human cells, e.g., appropriate promoters, RNA stabilization sequences 5' and 3' to the coding sequence, introns (which can be placed at any location 5' or 3' within encoded sequence), and poly(A) addition sites, as well as
15 an origin of replication and selectable markers enabling the constructs to replicate and be selected for in prokaryotic and/or eukaryotic hosts.

An example of a DNA sequence encoding an immunogenic HPV E7 antigen is the BIOTOPE_{HPV} construct
20 (SEQ ID NO:7), which is shown schematically in Fig. 1. This plasmid contains a minigene (SEQ ID NO: 5) at positions 3290-3413. The minigene encodes the HLA-DR α trafficking peptide linked to 12 residues of the A2.1/4 peptide. In the peptide encoded by the minigene, an
25 alanine has been substituted for the amino terminal leucine in the A2.1/4 peptide in order to facilitate cleaving of the trafficking peptide from the immunogenic peptide by a signal peptidase. The BIOTOPE_{HPV} construct also carries the immediate early promoter of human
30 cytomegalovirus (CMV) at positions 2619-3315, and RNA stabilization sequences (RST) derived from the *Xenopus laevis* β -globin gene flanking the minigene (positions 3219-3279 and 3426-3624). To maximize export from the nucleus, the pre-mRNA expressed from the plasmid contains
35 a chimeric intron between the coding sequence of the

- 10 -

minigene and the SV40 polyadenylation site. The intron can also function if located between the promoter and the coding region.

Once in the cytoplasm of the cell, the mRNA
5 transcribed from the minigene is translated to produce a
40 amino acid hybrid peptide. The first two amino acids
are methionine and aspartic acid (derived from vector
sequences), and the next 38 amino acids correspond to Met
Ala Ile Ser Gly Val Pro Val Leu Gly Phe Phe Ile Ile Ala
10 Val Leu Met Ser Ala Gln Glu Ser Trp Ala Ala Leu Met Gly
Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:6). The
amino-terminal 25 amino acids of the 38-residue portion
are identical in sequence to the non-polymorphic HLA-DR α
chain gene leader sequence (SEQ ID NO:18). The last 13
15 amino acids have the sequence Ala Leu Met Gly Thr Leu Gly
Ile Val Cys Pro Ile Cys (SEQ ID NO:4), which is the
A2.1/4 peptide described above, but with an alanine
residue substituted for the amino terminal leucine
residue. While the translated peptide is 40 amino acids
20 in this example, it is understood that a longer peptide
would be generated if the plasmid encodes, for example,
an immunogenic peptide with the amino acid sequence of
SEQ ID NO:23.

Also within the plasmid is a kanamycin resistance
25 gene (positions 519-1313), which is driven by the SV40
early promoter (positions 131-484) and which has a
thymidine kinase (TK) polyadenylation site (positions
1314-1758). The kanamycin resistance gene and
accompanying regulatory sequences are for selection
30 purposes only and can be removed from the plasmid if
selection is not required or desired.

Once expressed in a cell, the encoded peptide can
be processed into one of several HLA MHC class I binding
epitopes. At least some of these are included in Table
35 1. These peptides can bind the HLA-A2 allele and may

- 11 -

also bind other alleles, such as HLA-A1, HLA-A3, HLA-A11, HLA-A24. The MHC molecule, upon binding to the peptide, can activate a T cell response. MHC class II binding peptides may also be generated from the encoded peptide.

5 These peptides would be expected to activate T helper cells or CTL upon presentation by the MHC class II expressing cells. Other receptors may also bind the encoded peptide or its processed fragments to activate immune cells such as NK or B cells. These cells may also

10 be activated by cytokines elicited in response to the peptides of the invention.

The peptides and nucleic acids of the invention can be used as vaccines prophylactically or therapeutically in subjects known to be infected by HPV, suspected of being infected by HPV, or likely to become

15 infected by HPV. Other suitable subjects include those displaying symptoms of, or likely to develop, HPV-associated conditions. The immunogenic peptides, and nucleic acids encoding these peptides, can be used as

20 vaccines in preventing or treating conditions associated with infections of HPV strain 16, e.g., bowenoid papulosis, anal dysplasia, respiratory or conjunctival papillomas, cervical dysplasia, cervical cancer, vulval cancer, or prostate cancer. They can also be used to

25 treat conditions associated with other HPV strains, especially those associated with HPV strains 18, 45, 6, 11, 35 and 31, which have regions of homology to the peptide of SEQ ID NO:3. These conditions include, e.g., exophytic condyloma (HPV strains 6 and 11), flat

30 condyloma, especially of the cervix (HPV strains 6, 11, 16, 18, and 31), giant condyloma (HPV strains 6 and 11), cervical cancer (HPV strains 18, 31, and 33, in addition to HPV strain 16), respiratory and conjunctival papillomas (HPV 6 and 11), and infection with genital-

35 tract HPVs (HPV 6, 11, and 16).

- 12 -

The immunogenic peptides or nucleic acids encoding the peptides can administered alone or in combination with other therapies known in the art, e.g., chemotherapeutic regimens, radiation, and surgery, to
5 treat HPV infections, or diseases associated with HPV infections. In addition, the peptides and nucleic acids of the invention can be administered in combination with other treatments designed to enhance immune responses, e.g., by co-administration with adjuvants or cytokines
10 (or nucleic acids encoding cytokines) as is well known in the art.

The peptides or nucleic acids of the invention can also be used in manufacture of a medicament for the prevention or treatment of HPV infection, or conditions
15 associated with HPV infection.

*Delivery of Immunogenic Peptides and Nucleic Acids
Encoding Immunogenic Peptides*

The delivery systems of the invention may be used to deliver, into appropriate cells, peptides, or DNA
20 constructs which express peptides, intended to stimulate an immune response against HPV. An advantage of DNA delivery is that the antigenic peptide is produced inside the target cell itself, where the interaction with a class I or class II MHC molecule to which the immunogenic
25 peptide binds is kinetically favored. This is in contrast to standard vaccine protocols which do not specifically direct antigenic peptides to MHC molecules. In addition, the immune response directly stimulated by DNA vaccines of the invention is likely to be limited to
30 a T cell mediated response, in contrast to standard vaccine protocols which result in a more generalized immune response, although it is possible that an antibody response may be indirectly induced when cells bearing viral particles are killed, or by other mechanisms.

- 13 -

The immunogenic peptides, or nucleic acids encoding the peptides, can be administered using standard methods, e.g., those described in Donnelly et al., J. Imm. Methods 176:145, 1994, and Vitiello et al., J. Clin. Invest. 95:341, 1995. Peptides and nucleic acids of the invention can be injected into subjects in any manner known in the art, e.g., intramuscularly, intravenously, intraarterially, intradermally, intraperitoneally, intranasally, intravaginally, intrarectally or subcutaneously, or they can be introduced into the gastrointestinal tract, the mucosa, or the respiratory tract, e.g., by inhalation of a solution or powder containing the microparticles. Administration can be local (e.g., at the cervix or other site of infection) or systemic.

The immunogenic peptides and nucleic acids encoding immunogenic peptides can be delivered in a pharmaceutically acceptable carrier such as saline, lipids, liposomes, microspheres, nanospheres, as colloidal suspensions, or as powders. They can be naked or associated or complexed with delivery vehicles and delivered using delivery systems known in the art, such as lipids, liposomes, microparticles, gold, nanoparticles, polymers, condensing agents, polysaccharides, polyamino acids, dendrimers, saponins, adsorption enhancing materials, or fatty acids.

It is expected that a dosage of approximately 0.1 to 100 μ moles of the polypeptide, or of about 1 to 200 μ g of DNA, would be administered per kg of body weight per dose. Where the patient is an adult human, vaccination regimens can include, e.g., intramuscular, intravenous, oral, or subcutaneous administrations of 10-1000 μ g of pBIOTOPE_{HPV} DNA when delivered in a microparticle, or of about 100-1000 μ g of naked pBIOTOPE_{HPV} DNA delivered intramuscularly or

- 14 -

intradermally, repeated 3-6 times. Of course, as is well known in the medical arts, dosage for any given patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Determination of optimal dosage is well within the abilities of a pharmacologist of ordinary skill.

10 Other standard delivery methods, e.g., biolistic transfer, or ex vivo treatment, can also be used. In ex vivo treatment, e.g., antigen presenting cells (APCs), dendritic cells, peripheral blood mononuclear cells, or bone marrow cells, can be obtained from a patient or an appropriate donor and activated ex vivo with the immunogenic compositions, and then returned to the patient.

Microparticle Delivery of Synthetic Immunogenic Peptides or Plasmids Encoding Immunogenic Peptides

20 Microparticles, including those described in U. S. Patent No. 5,783,567, can be used as vehicles for delivering macromolecules such as DNA, RNA, or polypeptides into cells. They contain macromolecules embedded in a polymeric matrix or enclosed in a shell of polymer. Microparticles act to maintain the integrity of the macromolecule e.g., by maintaining enclosed DNA in a nondegraded state. Microparticles can also be used for pulsed delivery of the macromolecule, and for delivery at a specific site or to a specific cell or target cell population.

30 The polymeric matrix can be a biodegradable copolymer such as poly-lactic-co-glycolic acid, starch, gelatin, or chitin. Microparticles can be used in particular to maximize delivery of DNA molecules into a

- 15 -

subject's phagocytotic cells. Alternatively, the microparticles can be injected or implanted in a tissue, where they form a deposit. As the deposit breaks down, the nucleic acid is released gradually over time and
5 taken up by neighboring cells (including APCs) as free DNA.

Liposomal Delivery of Synthetic Immunogenic Peptides or Plasmids Encoding Immunogenic Peptides

10 The immunogenic peptides of the invention can be administered into subjects via lipids, dendrimers, or liposomes using techniques that are well known in the art. For example, liposomes carrying immunogenic polypeptides or nucleic acids encoding immunogenic
15 peptides are known to elicit CTL responses *in vivo* (Reddy et al., J. Immunol. 148:1585, 1992; Collins et al., J. Immunol. 148:3336-3341, 1992; Fries et al., Proc. Natl. Acad. Sci. USA 89:358, 1992; Nabel et al., Proc. Nat. Acad. Sci. (USA) 89:5157, 1992).

20 *Delivery of Synthetic Immunogenic Peptides or Plasmids Encoding Immunogenic Peptides Using Saponin*

The peptides and nucleic acids of the invention can be administered by using Immune Stimulating Complexes (ISCOMS), which are negatively charged cage-like
25 structures of 30-40nm in size formed spontaneously on mixing cholesterol and Quil A (saponin), or saponin alone. The peptides and nucleic acids of the invention can be co-administered with the ISCOMS, or can be administered separately.

30 Protective immunity has been generated in a variety of experimental models of infection, including toxoplasmosis and Epstein-Barr virus-induced tumors, using ISCOMS as the delivery vehicle for antigens (Mowat et al., Immunology Today 12:383-385, 1991). Doses of

- 16 -

antigen as low as 1 μ g encapsulated in ISCOMS have been found to produce class I mediated CTL responses, where either purified intact HIV-1-IIIB gp 160 envelope glycoprotein or influenza hemagglutinin is the antigen
5 (Takahashi et al., Nature 344:873-875, 1990).

Measuring Responses of the Immune System and of HPV Virus Infections to the Immunogenic Peptides or Nucleic Acids Encoding the Immunogenic Peptides

The ability of immunogenic peptides, or nucleic
10 acids encoding the same, to elicit an immune response can be assayed by using methods for measuring immune responses that are well known in the art. For example, the generation of cytotoxic T cells can be demonstrated in a standard ⁵¹Cr release assay, by measuring
15 intracellular cytokine expression, or by using MHC tetramers. Standard assays, such as ELISA or ELISPOT, can also be used to measure cytokine profiles attributable to T cell activation. T cell proliferation can also be measured using assays such as ³H-thymidine
20 uptake and other assays known in the art. B cell responses can be measured using art recognized assays such as ELISA.

Other methodologies, e.g., digital imaging, cytologic, colposcopic and histological evaluations, can
25 also be used to evaluate the effects of immunogenic peptides, and of nucleic acids encoding the immunogenic peptides, on papilloma virus-associated lesions, or on papilloma virus levels generally.

The following are examples of the practice of the
30 invention. They are not to be construed as limiting the scope of the invention in any way.

- 17 -

EXAMPLES

As described in the Examples below, experimental models were chosen to demonstrate the generation of vigorous CTL responses to plasmids encoding the immunogenic peptides of the invention, e.g., pBIOTOPE_{HPV}.

Initial screening of HPV peptide sequences was performed by assessing binding affinity to the human class I HLA-A2 molecule. This was done by measuring the changes in circular dichroism (CD) as the receptor/ligand complex "melted". Examples of this type of screening are shown in Example 1. Of particular interest in Example 1 was the hybrid peptide A2.1/4, which contains at least two known epitopes.

Using a murine transgenic model, plasmids containing minigenes encoding these peptides were evaluated for their ability to generate HLA-A2 restricted CTLs (Examples 2 and 3). CTL activity, as measured using human target cells labeled with HPV peptides, was significantly increased over control targets for both the plasmids encoding A2.4 and A2.1/4, including the pA2.4 plasmid delivered in a PLGA microparticle.

Example 1. Peptides derived from HPV strain 16 E7 protein bind purified HLA-A*0201 with high affinity

To determine if peptides A2.1 (SEQ ID NO:1), A2.2 (SEQ ID NO:17), A2.4 (SEQ ID NO:2) A2.1/4 (SEQ ID NO:3), and A2.1/4 SWQ (SEQ ID NO:23) bind with biological affinity to the human class I molecule HLA-A2 (for the peptides A2.1, A2.2, A2.4 and A2.1/4) or HLA-A3 (for the A2.1/4-SQK peptide), recombinant HLA-A2 or HLA-A3 was produced in *E. coli* and refolded in the presence of the HPV-derived peptides and purified human β_2 -microglobulin. The resulting peptide-HLA complexes were then further purified by HPLC. To determine the precise thermokinetic interaction energy between receptor and ligand, each

- 18 -

complex was "melted" while its structure was monitored by circular dichroism. The temperature required to "melt" the complex is an accurate indication of the affinity between receptor and ligand.

5 The results of the binding studies are shown in Table II.

Table II. Peptides binding HLA-A molecules

TABLE II

10	NAME	Amino Acid Sequence	IC ₅₀ [□]	Tm [◆]
	A2.1	SEQ ID NO:1	8	47.8
	A2.2	SEQ ID NO:17	49	52.5
	A2.4	SEQ ID NO:2	153	41.5
15	A2.1/4	SEQ ID NO:3	ND	41.0
	A2.1/4SWQ	SEQ ID NO:23	ND	47.8

20 [□]IC₅₀ represents the amount (nM) of peptide required for 50% inhibition of binding of a radiolabeled standard peptide to HLA-A*0201 or HLA-A*0301 measured in a molecular binding assay.

25 *Values represent the temperature in degrees Celsius at which 50% of the refolded complexes are melted. HLA-A2 and HLA-A3 will not refold in the absence of a peptide ligand of sufficient affinity.

30 Of particular interest is a hybrid peptide A2.1/4, which contains at least two known overlapping epitopes, A2.1 and A2.4, each of which is presented by HLA-A2 positive human cervical tumor cells expressing the HPV 16 E7 protein (Ressing et al., J. Immunology 154:5934, 1995). Of the peptides studied, A2.4 is predicted to be the most capable of eliciting cross reactive immune responses between HPV strains. Moreover, the hybrid peptide generates both the A2.1 and A2.4

- 19 -

peptides; administration of pBIOTOPE_{HPV} to mice was found to generate T cell responses to both immunogenic peptides.

Example 2. Induction of HPV-specific CTL in HLA-transgenic mice immunized with intramuscular injections of a plasmid encoding the HPV strain 16 derived A2.4 peptide.

To demonstrate that a plasmid encoding the A2.4 peptide (SEQ ID NO:2) produced HPV peptides *in vivo* and that CTL to these peptides were generated, a transgenic animal model was employed. The HLA-A2/K^b mouse line produces a hybrid MHC class I molecule. In this hybrid, the peptide binding domains ($\alpha 1$ and $\alpha 2$) are derived from the human class I molecule HLA-A*0201, whereas the domain ($\alpha 3$) which interacts with the CD8 co-receptor on CTLs is derived from the murine class I molecule K^b. The resulting animal is capable of responding to immunogens which contain HLA-A2 restricted epitopes and of generating murine CTLs that recognize human target cells expressing HLA-A2 (Vitiello et al., J. Exp. Med. 173:1007, 1991).

6-8 week old HLA-A2/K^b females were immunized with either a plasmid encoding the A2.4 peptide having the amino terminal leucine replaced with an alanine residue, or with a null vector. Injections were performed with 50 μ g of plasmid DNA injected as "naked DNA" (that is, with no liposome, microparticle, or other carrier) into each anterior tibialis muscle. A booster immunization was performed 14 days after the first immunization, and a second booster immunization was performed 14 days after the first boost. Ten days following the third immunization, splenocytes were harvested and stimulated *in vitro* with syngeneic lipopolysaccharide (LPS) blasts which had been incubated with the synthetic A2.4 peptide.

- 20 -

After 4 days of co-culture, CTL activity was measured on human targets labeled with HPV peptides (Table III).

5 **Table III. Lysis of Human Cells Labeled with HPV-derived Peptides by Murine CTL from HLA-Transgenic Mice Immunized with Plasmid Encoding an A2.4 peptide.**

	IMMUNOGEN	% LYSIS OF TARGET CELLS*
	pVA2.4	28.7 ± 0.7*
10	Vector	6.8 ± 2.9*

15 *Data are reported as the mean lysis values at 100:1 effector to target ratio. Error is reported as the standard deviation; p=0.05 by Students t-test.

Mice immunized with a plasmid encoding the A2.4 peptide generate CTL that lyse human targets expressing HLA-A2 and the appropriate HPV peptide. This response is significantly greater than that achieved by immunization with null vector DNA alone.

Example 3. Plasmid DNA encoding the A2.1/4 peptide delivered to mice in PLGA microparticles elicits CTL responses

25 6-8 week old HLA-A2/K^b females were immunized intraperitoneally one time with 2-5µg of PLGA microparticles containing plasmid pBIOTOPE_{HPV}. Seven days following the immunization, splenocytes were harvested and *in vitro* stimulated with IL-2. After 2 days, CTL activity was measured on human targets labeled with HPV peptides (HPV(+)), or lacking HPV peptide (HPV(-)), at an E:T ratio of 50:1 (Table IV).

- 21 -

Table IV. Lysis of Human Cells Labeled with HPV-derived Peptides by Murine Splenocytes from HLA-Transgenic Mice Immunized with PLGA Microparticles Containing pBIOTOPE_{HPV}

IMMUNOGEN	% LYSIS OF TARGET CELLS	
	HPV(+)	HPV(-)
pBIOTOPE _{HPV}	17.4 ± 2.8*	3.9 ± 4.2*

10

Data are reported as the mean lysis values from three individual measurements.

*Error is reported as the standard deviation; p value <0.05 as determined by the Students t-test.

15

Thus, mice immunized with PLGA microparticles containing pBIOTOPE_{HPV} generate CTL that lyse human targets expressing HLA-A2 and A2.1/4 peptide.

Example 4. Synthetic peptides derived from HPV type 16 activate human CTL

20

Peripheral blood mononuclear cells (PBMC) from an HLA-A2* donor were cultured *in vitro* for two rounds of stimulation in the presence of 300 units of IL-2 and peptide A2.1/4 (LLMGTLGIVCPIC) (SEQ ID NO:3) or an immunodominant peptide having the amino acid sequence
 25 GILGFVFTL (SEQ ID NO:24) from influenza virus, which was used as a positive control.

Seven days after the second stimulation, each culture was subdivided into two subgroups. One subgroup of each culture was stimulated for an additional 7 hours
 30 with the respective peptide ("the third peptide stimulation"), while the other subgroups were cultured without the peptide. All samples were pretreated with brefeldin A to prevent cytokine secretion. The cells

- 22 -

were then subjected to triple color flow cytometry staining for CD8, CD16, and interferon- γ .

The results of the experiments are shown in Fig. 2. For cells treated with the A2.1/4 peptide, 28% of the cells subjected to the third peptide stimulation stained positive for interferon- γ , compared to 1.7% of the cells that did not receive a final stimulation. The percentage of CD8⁺ cells in cells receiving a third stimulation with peptide was 14.4%, while 19.2% of the cells which did not received a third stimulation were CD8⁺. Overall, 3.1% of the PBMC receiving a final pulse of the A2.1/4 peptide were activated CTL, i.e., were CD8⁺ CD16⁻ IFN- γ ⁺, compared to 0.5% of the cells receiving no final pulse of HPV-derived peptide.

For cells treated with the influenza peptide, 11.5% of the cells receiving a third stimulation with the influenza peptide were positive for interferon γ , compared to 1.7% of the cells that did not receive a third stimulation. For cells cultured with influenza peptide, 11.5% of the cells given a final pulse of influenza peptide were activated, compared to 1.49% of cells which were not given a final pulse of influenza.

Figs. 3 and 4 demonstrate that CTL specific for the A2.1 or A2.4-C peptides can recognize and lyse HPV 16-infected cells. Fig. 3 shows CTL-mediated lysis of an HLA-A2⁺, HPV-16⁺ transformed line (Caski) by T cell populations exposed to peptide A2.1, A2.4-C, or a peptide derived from influenza virus ("Flu"). Effector/target (E/T) ratios ranging from 25 to 1.5 were used. The peptide A2.4-C was highly effective at inducing lysis, with nearly 35% release detected at an E/T ratio of 25:1. The A2.1 peptide was less effective, but nevertheless caused much higher percentages of lysis at E/T ratios of 25:1 and 12:1 than did the influenza peptide.

- 23 -

Results with PBL isolated from a second HLA-A2⁺ individual and subjected to two rounds of stimulation with peptide A2.1, peptide A2.4, or the influenza peptide ("Flu") are shown in Fig. 4. Both the A2.1 and A2.4
5 peptides induced higher levels of lysis than did the influenza peptide.

These observations demonstrate that the A2.1/4 peptide, or peptides derived therefrom, can activate and expand PBL from humans, and that these peptides can cause
10 CTL-mediated lysis of target cells transformed with HPV16.

Other Embodiments

It is to be understood that while the invention has been described in conjunction with the detailed
15 description thereof, that the foregoing description is intended to illustrate and not limit the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

- 24 -

What is claimed is:

1. A peptide less than 19 amino acids in length, wherein the peptide comprises the amino sequence Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:16).
- 5 2. The peptide of claim 1, wherein the peptide's amino acid sequence comprises Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:3).
3. The peptide of claim 1, wherein the peptide's sequence comprises Xaa Leu Met Gly Thr Leu Gly Ile Val
10 Cys Pro Ile Cys, Xaa being Met, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu (SEQ ID NO:19).
4. The peptide of claim 3, wherein Xaa is Ala or Met.
5. The peptide of claim 1, wherein the peptide's
15 sequence comprises Leu Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys Ser Gln Lys (SEQ ID NO:25).
6. A peptide less than 19 amino acids in length, wherein the peptide comprises the amino acid sequence Gly Thr Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:21).
- 20 7. The peptide of claim 6, wherein the peptide's sequence comprises Xaa Gly Thr Leu Gly Ile Val Cys Pro Ile Cys, Xaa being Met, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu (SEQ ID NO:25).
8. The peptide of claim 6, wherein the peptide's
25 sequence comprises Met Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:26).

- 25 -

9. The peptide of claim 7, wherein the peptide's sequence consists of Xaa Gly Thr Leu Gly Ile Val Cys Pro Ile Cys, Xaa being Met, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu.
- 5 10. The peptide of claim 8, wherein the peptide's sequence consists of Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys Ser Gln Lys (SEQ ID NO: 26).
11. A peptide consisting of the amino acid sequence Thr Leu Gly Ile Val Cys Pro Ile (SEQ ID NO:20).
- 10 12. A polypeptide comprising a first peptide and a second peptide linked by a peptide bond, the first peptide being a peptide which controls intracellular trafficking of a peptide to which it is attached, and the second peptide consisting of a sequence 12-18 amino acids
15 in length comprising the sequence Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:16).
13. The polypeptide of claim 12, wherein the sequence of the first peptide comprises the amino acid sequence Met Ala Ile Ser Gly Val Pro Val Leu Gly Phe Phe
20 Ile Ile Ala Val Leu Met Ser Ala Gln Glu Ser Trp Ala (SEQ ID NO:18).
14. The polypeptide of claim 12, wherein the amino acid sequence of the second peptide is Xaa Leu Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys, Xaa being Met,
25 Leu, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu (SEQ ID NO:19).
15. The polypeptide of claim 12, wherein the amino acid sequence of the second polypeptide is Ala Leu

- 26 -

Met Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID
NO:4).

16. The polypeptide of claim 13, wherein the
amino acid sequence of the second peptide is Xaa Leu Met
5 Gly Thr Leu Gly Ile Val Cys Pro Ile Cys, Xaa being Met,
Leu, Ala, Ser, Arg, Lys, Gly, Gln, Asp, or Glu (SEQ ID
NO:19).

17. The polypeptide of claim 13, wherein the
amino acid sequence of the second peptide is Ala Leu Met
10 Gly Thr Leu Gly Ile Val Cys Pro Ile Cys (SEQ ID NO:4).

18. A polypeptide comprising a first peptide and
a second peptide linked by a peptide bond, the first
peptide being a peptide which controls intracellular
trafficking of a peptide to which it is attached, and the
15 second peptide consisting of a sequence 8-18 amino acids
in length comprising the sequence Thr Leu Gly Ile Val Cys
Pro Ile (SEQ ID NO:20).

19. A therapeutic composition comprising
(a) the peptide of claim 1, and
20 (b) a pharmaceutically acceptable carrier.

20. A therapeutic composition comprising
(a) the peptide of claim 6, and
(b) a pharmaceutically acceptable carrier.

21. A microparticle comprising a polymeric
25 matrix and the peptide of claim 1.

22. A microparticle comprising a polymeric
matrix and the peptide of claim 6.

- 27 -

23. A microparticle comprising a polymeric matrix and the polypeptide of claim 1.

24. A microparticle comprising a polymeric matrix and the polypeptide of claim 18.

5 25. A liposome or immune-stimulating complex (ISCOM) containing the peptide of claim 1.

26. A liposome or immune-stimulating complex (ISCOM) containing the peptide of claim 6.

10 27. A method of eliciting an MHC class I-mediated immune response in a mammal, which method comprises administering a purified preparation of the peptide of claim 1 to a mammal.

15 28. A method of eliciting an MHC class I-mediated immune response in a mammal, which method comprises administering a purified preparation of the peptide of claim 6 to a mammal.

20 29. A method of eliciting an MHC class I-mediated immune response in a mammal, which method comprises administering the microparticle of claim 21 to a mammal.

30. The method of claim 29, wherein the polymeric matrix of said microparticle consists essentially of a copolymer of poly-lactic-co-glycolic acid (PLGA).

25 31. A method of eliciting an MHC class I-mediated immune response in a mammal, which method

- 28 -

comprises administering the microparticle of claim 22 to a mammal.

32. The method of claim 31, wherein the polymeric matrix of said microparticle consists essentially of a
5 copolymer of poly-lactic-co-glycolic acid (PLGA).

33. A nucleic acid comprising a coding sequence coding for expression of the peptide of claim 1.

34. A nucleic acid comprising a coding sequence coding for expression of the peptide of claim 6.

10 35. A nucleic acid comprising a coding sequence coding for expression of the polypeptide of claim 12.

36. A nucleic acid comprising a coding sequence coding for expression of the polypeptide of claim 18.

15 37. A plasmid comprising a coding sequence coding for expression of the polypeptide of claim 12.

38. A microparticle comprising a polymeric matrix and the plasmid of claim 37.

39. The microparticle of claim 38, wherein the polymeric matrix of the microparticle consists
20 essentially of a copolymer of PLGA.

40. The microparticle of claim 38, wherein the microparticle has a diameter of 0.02 to 20 microns.

41. The microparticle of claim 38, wherein the microparticle has a diameter of less than about 11
25 microns.

- 29 -

42. A cell comprising the plasmid of claim 37.

43. The cell of claim 42, wherein the cell is a mammalian B cell or APC.

44. A method of making a polypeptide, which
5 method comprises maintaining the cell of claim 42 under conditions permitting expression of said polypeptide.

45. A plasmid comprising a coding sequence coding for expression of the polypeptide of claim 18.

46. A microparticle comprising a polymeric
10 matrix and the plasmid of claim 45.

47. The microparticle of claim 46, wherein the polymeric matrix of said microparticle consists essentially of a copolymer of PLGA.

48. The microparticle of claim 46, wherein the
15 microparticle has a diameter of 0.02 to 20 microns.

49. The microparticle of claim 46, wherein the microparticle has a diameter of less than about 11 microns.

50. A cell comprising the plasmid of claim 45.

20 51. The cell of claim 50, wherein the cell is a mammalian B cell or APC.

52. A method of making a peptide, which method comprises maintaining the cell of claim 50 under conditions permitting expression of said polypeptide.

- 30 -

53. A method of inducing an immune response in a mammal, which method comprises administering the nucleic acid of claim 35 to a mammal.

54. A method of inducing an immune response in a mammal, which method comprises administering the nucleic acid of claim 36 to a mammal.

55. A method of inducing an immune response in a mammal, which method comprises administering the plasmid of claim 37 to a mammal.

56. A method of inducing an immune response in a mammal, which method comprises administering the plasmid of claim 45 to a mammal.

57. A method of inducing an immune response in a mammal, which method comprises administering the microparticle of claim 38 to a mammal.

58. The method of claim 57, wherein the mammal is a human.

59. The method of claim 58, wherein the human suffers from, or is at risk of a condition selected from the group consisting of exophytic condyloma, flat condyloma, cervical cancer, respiratory papilloma, conjunctival papilloma, genital-tract HPV infection, and cervical dysplasia.

60. A method of inducing an immune response in a mammal, which method comprises administering the microparticle of claim 46 to a mammal.

- 31 -

61. The method of claim 60, wherein the mammal is a human.

62. The method of claim 61, wherein the human suffers from, or is at risk of, a condition selected from the group consisting of exophytic condyloma, flat condyloma, cervical cancer, respiratory papilloma, conjunctival papilloma, genital-tract HPV infection, and cervical dysplasia.

63. A plasmid DNA comprising the sequence of SEQ ID NO:7.

64. A microparticle comprising a polymeric matrix and a nucleic acid, wherein the polymeric matrix consists essentially of PLGA and the nucleic acid comprises the sequence of SEQ ID NO:7.

65. A method of inducing a cell mediated, anti-HPV immune response in a mammal, which method comprises administering to the mammal a DNA comprising the sequence of SEQ ID NO:7.


66. A method of inducing an immune response in a patient, which method comprises administering to the patient a microparticle having a diameter of less than 20 microns and consisting essentially of a polymeric matrix and a nucleic acid molecule, wherein the polymeric matrix consists essentially of PLGA and the nucleic acid molecule comprises the sequence of SEQ ID NO:7.

67. A DNA comprising the sequence of SEQ ID NO:5.

- 32 -

68. A DNA comprising the sequence of nucleotides 3219-3624 of SEQ ID NO:7.

69. A DNA comprising the sequence of nucleotides 3290-3413 of SEQ ID NO:7.






FIG. 2

3/3




FIG. 3




FIG. 4