



(86) Date de dépôt PCT/PCT Filing Date: 2001/07/26
(87) Date publication PCT/PCT Publication Date: 2002/02/07
(45) Date de délivrance/Issue Date: 2011/08/02
(85) Entrée phase nationale/National Entry: 2003/01/24
(86) N° demande PCT/PCT Application No.: JP 2001/006467
(87) N° publication PCT/PCT Publication No.: 2002/009754
(30) Priorité/Priority: 2000/07/28 (JP2000-228239)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),
A61K 31/704 (2006.01), *A61P 35/00* (2006.01),
C07K 16/32 (2006.01)
(72) Inventeurs/Inventors:
NOGUCHI, TOSHIHIRO, JP;
BABA, AKEMI, JP;
HANADA, MITSU HARU, JP
(73) Propriétaire/Owner:
DAINIPPON SUMITOMO PHARMA CO., LTD., JP
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : MEDICAMENT COMPRENANT DE L'AMRUBICINE ET UN ANTICORPS ANTI-HER2 POUR LE TRAITEMENT
DU CANCER DU SEIN
(54) Title: MEDICAMENT COMPRISING AMRUBICIN AND AN ANTI-HER2 ANTIBODY FOR TREATING BREAST
CANCER

(57) **Abrégé/Abstract:**

The present invention is directed to a medicament for treating breast cancer for use in combination therapy. The medicament comprises an anti-HER2 antibody and amrubicin or a pharmaceutically acceptable salt thereof as an active ingredient.

ABSTRACT

The present invention is directed to a medicament for treating breast cancer for use in combination therapy. The medicament comprises an anti-HER2 antibody and amrubicin or a pharmaceutically acceptable salt thereof as an active ingredient.

DESCRIPTION

MEDICAMENT COMPRISING AMRUBICIN AND AN ANTI-HER2 ANTIBODY FOR
TREATING BREAST CANCERTechnical Field

The present invention relates to medicament for treating cancer for use in combined therapy with an anti-HER2 antibody, which comprises amrubicin or a pharmaceutically acceptable salt thereof as an active ingredient.

Background Art

Amrubicin ((+)-(7S,9S)-9-acetyl-9-amino-7-[(2-deoxy- β -D-erythro-pentopyranosyl)oxy]-7,8,9,10-tetrahydro-6,11-dihydroxy-5,12-naphthacenedione) that is an anthracycline compound is known as an anti-cancer agent exhibiting less side effects in comparison with known anthracycline compounds such as doxorubicin and daunorubicin (Japanese Examined Patent Publication JP-B 3-5397 published April 27, 1985).

Amrubicin is characterized by a stronger anti-cancer activity of its metabolite (amrubicinol: hydroxylated product at 13-position of the anthracycline skeleton) obtained by being readily reduced in vivo, than that of amrubicin itself. To the contrary, doxorubicin and daunorubicin do not have such characteristics (Cancer

Chemother. Pharmacol., 30, 51-57 (1992)). In addition, cardiotoxicity of amrubicin was far less than that of doxorubicin in a rabbit chronic experimental model (Invest. New Drug, 15, 219-225 (1997)).

Although anthracycline anti-cancer agents have similar structures, diseases which are susceptible to treatment, modes of the action and the like have been known to vary as described below.

Daunorubicin and idarubicin are efficacious against leukemia, however, they are not efficacious against solid cancer. Doxorubicin, epirubicin, pirarubicin and aclarubicin have efficacies toward solid cancer (Nihon Iyakuhinshuu 23th. Ed., 2000, Yakugyou Jiho Co., Ltd.). Daunorubicin and doxorubicin inhibit DNA synthesis and RNA synthesis to the same extent, however, aclarubicin and marceromycin inhibit RNA synthesis to a greater extent than DNA synthesis, with mechanisms to exert anti-cancer effects being entirely different each other (JJSHP 27, 1087-1110 (1991)). Although an inhibitory action for cell proliferation of daunorubicin and doxorubicin is attenuated upon reduction of the ketone at 13-position, idarubicin and its reduced form exhibit the action to a similar extent (Cancer Chemother. Pharmacol. 30, 51-57 (1992)). To the contrary, in regard to amrubicin, its reduced form exhibits the action which is stronger than amrubicin (Jpn. J. Cancer

Res. 89, 1067-1073 (1998)). Tumor selectivity cannot be achieved by intravenous injection of doxorubicin, however, it was found that intravenous injection of amrubicin can result in the achievement of tumor selectivity in experiments on mouse (Jpn. J. Cancer Res. 89, 1061-1066 (1998)). An intercalation activity of doxorubicin is about ten times stronger than those of amrubicin and amrubicinol (Jpn. J. Cancer Res. 89, 1229-1238 (1998)). Although doxorubicin and daunorubicin largely distribute in cell nuclei, amrubicin and idarubicin largely distribute in cytoplasms (Ann. Haematol. 69, S13-S17 (1994); Jpn. J. Cancer Res. 89, 1229-1238 (1998); and Urol. Res. 25, 125-130 (1997)). Amrubicin and amrubicinol sufficiently stabilize a cleavable complex via topoisomerase II at a concentration which leads to 50% inhibition of cell proliferation, however, doxorubicin does not stabilize the cleavable complex at a concentration which leads to 50% inhibition of cell proliferation (Jpn. J. Cancer Res. 89, 1229-1238 (1998)).

Members in the anthracycline compounds thus vary in terms of their modes of action, and the indication therefor also vary. Among them, doxorubicin, and amrubicin and amrubicinol are extremely closely related in respect of chemical structures, however, the mechanisms of anti-cancer action are widely different.

Recently, an anti-cancer action by combined use of

doxorubicin that is an anthracycline compound with HerceptinTM was reported, finding that the combination of doxorubicin with HerceptinTM showed additive effects in an in vitro experiment (M. Pegram et al., Oncogene, 18, 2241-2251 (1999)). However, when doxorubicin and HerceptinTM were used in combination, it was also reported that cardiotoxicity is developed at a high frequency in clinical use. Therefore, such a combination therapy was not always recognized as being effective. HerceptinTM is one of the antibodies directed against a human epidermal cell growth factor receptor 2 protein (hereinafter, referred to as HER2), which has been used as a remedy for breast cancer.

Disclosure of the Invention

An object of the present invention is to provide a remedy for cancer by using amrubicin or a pharmaceutically acceptable salt thereof with an anti-HER2 antibody (for example, HerceptinTM) in combination.

Aiming at improvement of an anti-cancer action of amrubicin, the present inventors have carried out investigation for combination effects with various medicines, and could achieve the desired effects by using in combination with HerceptinTM. In other words, the present invention was accomplished through the finding that synergistic effects

on a cancer cell strain can be achieved, accompanied with small influences on cardiotoxicity by using amrubicin with HerceptinTM in combination. When the present invention was compared with a combined therapy of doxorubicin with HerceptinTM, merely additive effects have been found for doxorubicin as described above, and to the contrary, synergistic effects have been found according to the present invention. Moreover, doxorubicin results in cardiotoxicity at a high frequency through the use in combination with HerceptinTM, and thus application in a clinical field is difficult. However, in accordance with the present invention, small influences on cardiotoxicity are exerted, leading to expectations as new a remedy for cancer. Difference of these effects between the present invention and doxorubicin is believed to be responsible for difference of modes of action of amrubicin and doxorubicin.

Summary of the present invention is as described below.

(1) A medicament for treating cancer for use in combination therapy with an anti-HER2 antibody, which comprises amrubicin or a pharmaceutically acceptable salt thereof as an active ingredient.

(2) The medicament according to (1) described above wherein said cancer is breast cancer, gastric cancer, renal cancer, salivary gland cancer, ovarian cancer, uterine body cancer or lung cancer.

(3) The medicament according to (1) or (2) described above wherein said cancer is a cancer in which HER2 is highly expressed.

(4) The medicament according to any one of (1) to (3) described above for administering amrubicin or a pharmaceutically acceptable salt thereof at 0.5 - 3.0 mg/kg/day when an anti-HER2 antibody is administered at 0.1 - 4.5 mg/kg/day.

(5) The medicament according to (4) described above for administering amrubicin or a pharmaceutically acceptable salt thereof on successive three days in every three weeks, while administering an anti-HER2 antibody every one week.

(6) The medicament according to any one of (1) to (3) described above for administering amrubicin or a pharmaceutically acceptable salt thereof at 1.5 - 9.0 mg/kg/day when an anti-HER2 antibody is administered at 0.1 - 4.5 mg/kg/day.

(7) The medicament according to (6) described above for administering amrubicin or a pharmaceutically acceptable salt thereof every three weeks, while administering an anti-HER2 antibody every one week.

(8) The medicament according to any one of (1) to (7) wherein a dosage ratio of amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody is 2 : 1 - 8 : 1 at one week after initiating the administration

of both agents.

(9) Use of amrubicin or a pharmaceutically acceptable salt thereof for manufacturing a medicament for treating cancer which is administered in combination with an anti-HER2 antibody.

(10) The use according to (9) described above wherein said cancer is breast cancer, gastric cancer, renal cancer, salivary gland cancer, ovarian cancer, uterine body cancer or lung cancer.

(11) The use according to (9) or (10) described above wherein said cancer is a cancer in which HER2 is highly expressed.

(12) The use according to any one of (9) to (11) described above for administering amrubicin or a pharmaceutically acceptable salt thereof at 0.5 - 3.0 mg/kg/day when an anti-HER2 antibody is administered at 0.1 - 4.5 mg/kg/day.

(13) The use according to (12) described above for administering amrubicin or a pharmaceutically acceptable salt thereof on successive three days in every three weeks, while administering an anti-HER2 antibody every one week.

(14) The use according to any one of (9) to (11) described above for administering amrubicin or a pharmaceutically acceptable salt thereof at 1.5 - 9.0 mg/kg/day when an anti-HER2 antibody is administered at 0.1

- 4.5 mg/kg/day.

(15) The use according to (14) described above for administering amrubicin or a pharmaceutically acceptable salt thereof every three weeks, whilst administering an anti-HER2 antibody every one week.

(16) The use according to any one of (9) to (15) wherein a dosage ratio of amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody is 2 : 1 - 8 : 1 at one week after initiating the administration of both agents.

(17) A method of treating cancer comprising using an effective amount of an anti-HER2 antibody and an effective amount of amrubicin or a pharmaceutically acceptable salt thereof in combination.

(18) The method according to (17) described above wherein said cancer is breast cancer, gastric cancer, renal cancer, salivary gland cancer, ovarian cancer, uterine body cancer or lung cancer.

(19) The method according to (17) or (18) described above wherein said cancer is a cancer in which HER2 is highly expressed.

(20) The method according to any one of (17) to (19) described above comprising administering amrubicin or a pharmaceutically acceptable salt thereof at 0.5 - 3.0 mg/kg/day while administering an anti-HER2 antibody at 0.1 - 4.5 mg/kg/day.

(21) The method according to (20) described above comprising administering amrubicin or a pharmaceutically acceptable salt thereof on successive three days in every three weeks, while administering an anti-HER2 antibody every one week.

(22) The method according to any one of (17) to (19) described above comprising administering amrubicin or a pharmaceutically acceptable salt thereof at 1.5 - 9.0 mg/kg/day while administering an anti-HER2 antibody at 0.1 - 4.5 mg/kg/day.

(23) The method according to (22) described above comprising administering amrubicin or a pharmaceutically acceptable salt thereof every three weeks, while administering an anti-HER2 antibody every one week.

(24) The method according to any one of (17) to (23) wherein a dosage ratio of amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody is 2 : 1 - 8 : 1 at one week after initiating the administration of both agents.

(25) A kit for treating cancer comprising a first composition including amrubicin or a pharmaceutically acceptable salt thereof and a second composition including an anti-HER2 antibody.

(26) A medicament, use, method, kit or the like according to any one of (1) to (25) described above wherein

Herceptin is used as said anti-HER2 antibody.

In one particular embodiment, there is provided a medicament for treating breast cancer, which comprises amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody.

The medicament for treating cancer of the present invention comprises an effective amount of amrubicin, amrubicinol or a pharmaceutically acceptable salt thereof as an active ingredient, and is for use in combination with an effective amount of an anti-HER2 antibody.

The medicament for treating cancer of the present invention comprising an effective amount of amrubicin, amrubicinol or a pharmaceutically acceptable salt thereof as an active ingredient can enhance therapeutic effects on cancer of an anti-HER2 antibody by using it with anti-HER2 antibody in combination.

Exemplary pharmaceutically acceptable salt of amrubicin or amrubicinol may include acid addition salts and base addition salts. The acid addition salts include for example, inorganic acid salts such as hydrochloride, hydrobromide, sulfate, hydroiodide, nitrate, phosphate and the like; organic acid salts such as citrate, oxalate, acetate, formate, propionate, benzoate, trifluoroacetate, fumarate, maleate, tartrate, aspartate, glutamate, methanesulfonate, benzenesulfonate, camphor-sulfonate and the like. The base addition salts include inorganic base salts such as sodium salt, potassium salt, calcium salt, magnesium salt, ammonium salt and the like; organic base salts such as triethylammonium salt, triethanolammonium salt, pyridinium salt,

diisopropylammonium salt and the like.

The anti-HER2 antibody means an antibody against HER2 (human epidermal cell growth factor receptor 2 protein), which has a function inhibiting proliferation of cancer cells those are highly expressing HER2. The antibody may include a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a humanized antibody and the like, which can be produced in accordance with a known process. For example, an anti-HER2 monoclonal antibody can be produced in accordance with a process described in WO89/06692. Further, an anti-HER2 humanized antibody can be produced in accordance with a process described in WO94/04679. Preferable examples of the anti-HER2 antibody include a humanized antibody, and specifically, Trastuzumab (HerceptinTM, Genentech, Ltd.) is included.

Cancer which may be a therapeutic target in the present invention is not particularly limited, however cancer in which HER2 is highly expressed is preferred. For example, breast cancer, gastric cancer, renal cancer, salivary gland cancer, ovarian cancer, uterine body cancer, lung cancer and the like are included, and in particular, breast cancer is a preferable therapeutic target.

To a cancer patient, may be administered amrubicin or a pharmaceutically acceptable salt thereof in advance, may be administered an anti-HER2 antibody in advance, or may

be administered both agent concurrently. When both are not concurrently administered, an interval between the administrations of both may be determined ad libitum.

In accordance with the present invention, since a synergistic effect is observed by using amrubicin with an anti-HER2 antibody in combination, a therapeutic effect on cancer can be achieved at a lower dosage than usual dosages of amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody. Therefore, in accordance with the present invention, side effects resulting from increased dosages of amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody can be minimized.

Amrubicin or a pharmaceutically acceptable salt thereof according to the present invention can be administered parenterally (for example, via intravenous, subcutaneous or intramuscular injection, or intravesically, topically, transrectally, transdermally or transnasally) in general. Preferably, intravenous injection can be exemplified. In addition, it can be also used for the oral administration, and forms for the oral administration may include for example, tablet, capsule, pill, granule, powder, liquid, syrup, suspension and the like.

Dose and administration frequency of amrubicin or a pharmaceutically acceptable salt thereof according to the present invention vary depending upon symptoms, age and body

weight of the patient, mode of administration, and dose and administration frequency of the anti-HER2 which is used in combination. When administered as an injectable agent, it can be usually administered to an adult in the range of from about 10 mg to about 1000 mg, preferably from about 30 mg to about 500 mg per day, once or in divided doses for several times. When administered orally, it can be usually administered to an adult in the range of from about 100 mg to about 3000 mg, preferably from about 300 mg to about 1500 mg per day, once or in divided doses for several times. Furthermore, the administration can be repeated by providing intervals of from about one day to about 30 days.

The anti-HER2 antibody according to the present invention is usually administered to a targeted cancer cell site if possible, or alternatively, administered parenterally by an intravenous administration. In instances of the intravenous administration, it is preferably infused intravenously over about 30 minutes to about 90 minutes. For the parenteral administration, an anti-HER2 antibody is formulated into an injectable dosage form (solution, suspension or emulsion) together with a pharmaceutically acceptable excipient for the parenteral use. Such an excipient is essentially nontoxic and non-therapeutic. Examples of such an excipient include water, saline, Ringer solution, dextrose solution and 5%

human serum albumin. Nonaqueous excipients such as nonvolatile oil, ethyl oleate and the like can be also used. The excipient can contain a substance for increasing isotonicity and chemical stability, for example buffer agents and stabilizers.

Dose and administration frequency of the anti-HER2 antibody according to the present invention depend upon symptoms, age and body weight of the patient, mode of administration, and dose and administration frequency of the amrubicin which is used in combination, however, it can be usually administered to an adult in the range of from about 5 mg to about 1000 mg, preferably from about 10 mg to about 300 mg per day, once or in divided doses for several times. Furthermore, the administration can be repeated by providing intervals of from about 5 days to about 10 days.

Preferred embodiments for exhibiting suppressive effects on cancer according to the present invention are explained below. For example, dosage ratio of amrubicin hydrochloride and HerceptinTM used in combination is preferably in the range of 2 : 1 - 8 : 1. Since amrubicin is readily converted into its active metabolite amrubicinol in vivo as demonstrated in Example 1, it is believed that the ratio of the concentration 2 : 1-8 : 1, which demonstrated synergistic effects when amrubicin and HerceptinTM were used in combination, reflects the dosage ratio of amrubicin

hydrochloride and HerceptinTM.

Also in the case of treating a cancer patient, the dosage ratio in the range of 2 : 1 - 8 : 1 is similarly preferred as a dosage ratio for the one week on the initiation of administration of amrubicin hydrochloride and HerceptinTM.

Amrubicin hydrochloride is administered at a dosage of from about 1.5 to about 9.0 mg/kg for a single administration, or at a dosage of from about 0.5 to about 3.0 mg/kg on successive three days with repeating the administrations every three weeks. A route of administration is preferably intravenous administration. HerceptinTM is intravenously administered in the dosage range of from about 0.1 to about 4.5 mg/kg, which is administered every one week.

The administration of amrubicin hydrochloride and HerceptinTM follows one cycle of three weeks. The initiation of administration in each cycle is preferably executed on the same day. More specifically, in instances when amrubicin hydrochloride is administered on successive three days, it is preferred that amrubicin hydrochloride is administered on the first, second and third days of each cycle, while HerceptinTM is administered on the first, eighth and fifteenth days of the cycle.

The present invention also involves a kit for use in combination therapy of cancer comprising (a) a first

composition including amrubicin or a pharmaceutically acceptable salt thereof as an active ingredient, and (b) a second composition including an anti-HER2 antibody as an active ingredient.

The fact that small influences on cardiotoxicity are exerted in combination use of amrubicin hydrochloride with HerceptinTM can be verified by a test in which a monkey that responds to HerceptinTM is used. More specifically, it can be verified by the observation that cardiotoxicity is strongly presented in a rhesus monkey when doxorubicin hydrochloride and HerceptinTM are intravenously administered, although cardiotoxicity is not presented even when the rhesus monkey is administered with amrubicin hydrochloride at a dosage of two times under the same condition. The administration of amrubicin hydrochloride at a dosage of two times of doxorubicin hydrochloride is carried out on the basis of therapeutic dosages of both agents.

Examples

The present invention is explained in more detail by way of examples below, and the present invention is not limited thereto.

[Experimental Process]

Method for evaluation of combination effects in vitro

Combination effects of amrubicin hydrochloride, amrubicinol hydrochloride and doxorubicin hydrochloride with HerceptinTM in vitro were examined using human breast cancer cells. The effects of the combined use were evaluated by calculating a Combination Index (CI) following a method of Pegram et al. (M. Pegram et al., Oncogene, 18, 2241-2251 (1999)).

Because amrubicinol is an active metabolite of amrubicin, it was added to the test agent. Amrubicinol can be produced by a method described in a literature (Ishizumi et al., J. Org. Chem., 52, 4477-4485 (1987)).

Human breast cancer cell line BT-474 (a cell line in which HER2 is highly expressed) was obtained from ATCC (American Type Culture Collection). In a medium comprising equal amount of D-MEM (Dulbecco's Modified Eagle Medium) medium and F-12 (Nutrient Mixture F-12) medium which further contains 2.5 mM L-glutamine, 10% bovine fetal serum (FCS) and 2.5 µg/ml amphotericin B, human breast cancer cell line BT-474 was subcultured. The culture was carried out in an incubator with 5% CO₂ at 37°C. This medium was also used in the following experiments.

Test agents were prepared as below.

HerceptinTM was dissolved in a dissolution solution attached thereto to prepare a 21 mg/ml stock solution. Thereafter, the solution was used after 2 fold serial

dilutions with the medium.

Amrubicin hydrochloride and amrubicinol hydrochloride were dissolved in distilled water to give 1 mg/ml in use, which was used after sterilization by filtration followed by 2 fold serial dilutions with the medium.

Doxorubicin hydrochloride was obtained from Kyowa Hakko Kogyo Co., Ltd. A 2 mg/ml solution was prepared by adding 5 ml of sterile water to a 10 mg vial, and the solution was used.

Human breast cancer cell line BT-474 on subculture were subjected to a trypsin treatment, suspended in the medium, and seeded on a 96-well plate. The seeding density was set to be 1×10^4 cells/0.1 ml/well. After seeding, cells were cultured overnight in an incubator with 5% CO₂ at 37°C (Day 0).

To the group for evaluating a single agent, were added 0.05 ml/well of a diluted solution of a test agent and 0.05 ml/well of the medium; and to the group for evaluating the combination effect, were added 0.05 ml/well of a diluted solution of a test agent and 0.05 ml/well of the diluted solution of HerceptinTM (Day 1). Final concentration of each agent was: 0.0625 - 16 µg/ml for amrubicin hydrochloride; 0.00625 - 1.6 µg/ml for amrubicinol hydrochloride; 0.00625 - 1.6 µg/ml for doxorubicin hydrochloride; 0.000781 - 0.4 µg/ml for HerceptinTM. The experiments were conducted with

n=3 for the agent-treated group, and with n=6 for the non-treated (control) group.

Until Day 4, culture was conducted in an incubator with 5% CO₂ at 37° C.

On Day 4, 0.02 ml of WST solution (a solution containing 1.3 mg/ml WST-1 (2-(4-iodophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium, sodium salt) and 0.14 mg/ml 1-Methoxy PMS (1-Methoxy-5-methylphenazinium methylsulfate) in a phosphate buffer solution (PBS)) was added to each of the wells. Thereafter, culture was conducted in an incubator with 5% CO₂ at 37° C for 2 - 4 hours. Absorbance of the culture was then measured with MICROPLATE READER Model 3550-UV (BioRad), and living cell number was measured.

Concentration D of amrubicin hydrochloride, amrubicinol hydrochloride, doxorubicin hydrochloride and HerceptinTM required for achieving a particular inhibition rate of proliferation (fa) was determined by the following formula (1). The concentration for the single agent was defined as Df₁ and Df₂, while the concentration when used in combination was defined as D₁ and D₂, and thus a Combination Index (CI) was calculated by the formula (2) (M. Pegram et al., Oncogene, 18, 2241-2251 (1999)).

$$fa = 1 - f$$

$$D = D_m \times \{fa / (1 - fa)\}^{1/m} \dots (1)$$

$$CI = D_1/Df_1 + D_2/Df_2 + (D_1 \times D_2) / (Df_1 \times Df_2) \dots (2)$$

For a reference, f , D_m and m were calculated as follows.

$f = (\text{mean value of absorbance at each of the agent concentrations}) / (\text{mean value of absorbance at the agent concentration of 0 mg/ml})$

absorbance: $A_{420} - A_{630}$

Common logarithm of $(1/f) - 1$ was plotted versus common logarithm of the agent concentration, and then a regression line was drawn in accordance with a least square method to find the slope and X intercept. The value of X intercept represents a common logarithm of an IC_{50} value (concentration of the agent required for 50% inhibition of the cell proliferation). (As described above, IC_{50} value was represented as D_m , and amrubicin hydrochloride, amrubicinol hydrochloride or doxorubicin hydrochloride is represented with the subscript 1 while HerceptinTM is represented with the subscript 2).

CI and inhibition rate of proliferation fa were calculated as described above. A CI value is obtained when the value of fa is 0.9. Evaluation is made as an additive effect at $CI = 1$, as a synergistic effect at $CI < 1$, and as an antagonistic effect at $C > 1$.

Example 1

Combination effects of amrubicinol hydrochloride and doxorubicin hydrochloride with HerceptinTM

Using a method for evaluation of combination effects as described above, the combination effects were measured when a dosage ratio of amrubicinol hydrochloride and HerceptinTM is 2 : 1, and when a dosage ratio of doxorubicin hydrochloride and HerceptinTM is 2 : 1. The CI value was obtained when the value of fa is 0.9, which is shown in Table 1 below.

Table 1

	CI value	D ₁	D ₂	Df ₁	Df ₂
Present invention (Amrubicinol hydrochloride)	0.38	0.239	0.119	0.671	6.312
Control compound (Doxorubicin hydrochloride)	0.92	0.373	0.187	0.434	6.312

Units for D₁, D₂, Df₁ and Df₂ are (μg/ml).

The results described above revealed that the combination effect with HerceptinTM was a synergistic effect in the case of the present amrubicinol hydrochloride because the CI value indicated a considerably lower value than 1.0. However, in the case of doxorubicin hydrochloride, an additive effect was suggested because the CI value indicated approximately 1.0, which did not suggest a synergistic effect. Thus, it was demonstrated that the effects by combined use with HerceptinTM greatly differ between amrubicinol hydrochloride and doxorubicin hydrochloride.

Example 2

Combination effects and dosage ratio of amrubicinol
hydrochloride with HerceptinTM

Using a method for evaluation of combined use as described above, variation of the dosage ratio of amrubicinol hydrochloride and HerceptinTM was brought, and the evaluation test by the combined use in the range of 2 : 1 - 8 : 1 was conducted. A CI value was obtained when the value of fa is 0.9, which is shown in Table 2 below.

Table 2

Dosage Ratio of Amrubicinol hydrochloride and Herceptin	CI value	D ₁	D ₂	Df ₁	Df ₂
2 : 1	0.38	0.239	0.119	0.671	6.312
4 : 1	0.33	0.212	0.053	0.671	6.312
8 : 1	0.42	0.279	0.035	0.671	6.312

Units for D₁, D₂, Df₁ and Df₂ are (μg/ml).

As shown in Table 2 above, it was revealed that the combination of amrubicinol hydrochloride with HerceptinTM showed a synergistic effect, because the CI value showed lower values than 1.0 in either case in the range of 2 : 1 - 8 : 1 of the dosage ratio.

Example 3

Combination effects of amrubicin hydrochloride with
HerceptinTM

Using a method for evaluation of combination as

described above, the effect by combination of amrubicin hydrochloride and HerceptinTM was evaluated. The test for the effect by combination was conducted at a dosage ratio of 40 : 1 of amrubicin hydrochloride and HerceptinTM. A CI value was obtained when the value of f_a is 0.9, which is shown in Table 3 below.

Table 3

	CI value	D ₁	D ₂	Df ₁	Df ₂
Present invention (Amrubicin hydrochloride)	0.17	1.267	0.032	7.542	6.312

Units for D₁, D₂, Df₁ and Df₂ are (μ g/ml).

As shown in Table 3 above, it was revealed that in the case of combination of amrubicin hydrochloride with HerceptinTM, a synergistic effect was showed because the CI value showed a lower value than 1.0.

As described herein above, evaluation of the effect by combination was carried out by means of a Combination Index (CI) with a method for evaluation of combination by the Pegram method, and thus marked synergistic effects were found in combination of amrubicin hydrochloride with HerceptinTM and amrubicinol hydrochloride with HerceptinTM.

Example 4

Cardiotoxicity test using monkey

Presence of influences on cardiotoxicity is evaluated

in an experiment using a rhesus monkey in the case of the combination of amrubicin hydrochloride with HerceptinTM. More specifically, amrubicin hydrochloride and HerceptinTM are intravenously injected, and then the presence of cardiotoxicity is determined in a rhesus monkey administered with two times dose of amrubicin hydrochloride instead of doxorubicin hydrochloride under the following conditions in which cardiotoxicity is strongly presented.

Into vein of a rhesus monkey, is administered doxorubicin at a dose of 2.92 mg/kg (35 mg/m²), 7 - 8 times with three weeks intervals. On 140 - 150 days post initiation of the administration, vacuolation of myocardial cells are found upon histopathological observation of the heart (E. J. Gralla et al., Toxicology, 13, 263-273, 1979). Moreover, the administration of HerceptinTM to a human is carried out using a schedule of: administration at 4 mg/kg, followed by 2 mg/kg every week.

Accordingly, the agent-administered groups are set to be as follows:

- 1) agent non-administered group
- 2) doxorubicin hydrochloride 2 - 4 mg/kg, iv, 8q3W
- 3) doxorubicin hydrochloride 2 - 4 mg/kg, iv, 8q3W
HerceptinTM 4 mg/kg, followed by 2 mg/kg every week,
iv, administered 24 times
- 4) amrubicin hydrochloride 4 - 8 mg/kg, iv, 8q3W

5) amrubicin hydrochloride 4 - 8 mg/kg, iv, 8q3W
HerceptinTM 4 mg/kg, followed by 2 mg/kg every week,
iv, administered 24 times

On 140 - 150 days post initiation of the administration,
the animal is dissected, and histopathological diagnosis
of the heart is carried out. Judging from the comparison
of impairment of the myocardial cells, cardiotoxicity is
evaluated.

Industrial applicability

According to the present invention, a medicament for
combination therapy for cancer which is useful in treating
a cancer patient is provided. More specifically, by using
amrubicin hydrochloride with HerceptinTM in combination,
synergistic anti-cancer effects can be achieved. Therefore,
anti-cancer effects of amrubicin hydrochloride can be
improved by using HerceptinTM in combination. Furthermore,
the effects have come to be exerted also to a kind of cancer
cells in which the effects of amrubicin hydrochloride are
hard to be exerted, by using with HerceptinTM in combination.
In addition, it becomes possible to suppress the dose of
amrubicin hydrochloride on behalf of the synergistic effects
by combination, thereby enabling cancer therapy with
alleviated side effects to more extent.

CLAIMS

1. A medicament for treating breast cancer, which comprises amrubicin or a pharmaceutically acceptable salt thereof and an anti-HER2 antibody.
2. The medicament according to claim 1, wherein the breast cancer is a breast cancer in which HER2 is highly expressed.
3. The medicament according to claim 1, which is formulated for administration of amrubicin or a pharmaceutically acceptable salt thereof at an amount of 0.5 to 3.0 mg/kg/day and for administration of an anti-HER2 antibody at an amount of 0.1 to 4.5 mg/kg/day.
4. The medicament according to claim 1, which is formulated for administration of amrubicin or a pharmaceutically acceptable salt thereof at an amount of 1.5 to 9.0 mg/kg/day and for administration of an anti-HER2 antibody at an amount of 0.1 to 4.5 mg/kg/day.
5. The medicament according to any one of claims 1 to 4, which is formulated to provide a dosage ratio of amrubicin or a pharmaceutically acceptable salt thereof to an anti-HER2 antibody of 2:1 to 8:1.
6. Use of amrubicin, or a pharmaceutically acceptable salt thereof, in the manufacture of a first medicament for use in combination therapy with a second medicament comprising an anti-HER2 antibody for the treatment of breast cancer.
7. The use according to claim 6, wherein the breast cancer is a breast cancer in which HER2 is highly expressed.

8. The use according to claim 6 or 7, wherein the first medicament is formulated for administration of amrubicin or a pharmaceutically acceptable salt thereof at an amount of 0.5 - 3.0 mg/kg/day.

9. The use according to claim 8, wherein the second medicament is formulated for administration of an anti-HER2 antibody at an amount of 0.1 - 4.5 mg/kg/day.

10. The use according to claim 8 or 9, wherein the first medicament is formulated for administration of amrubicin or a pharmaceutically acceptable salt thereof on successive three days in every three weeks.

11. The use according to claim 10, wherein the second medicament is formulated for administration of an anti-HER2 antibody every one week.

12. The use according to claim 6 or 7, wherein the first medicament is formulated for administration of amrubicin or a pharmaceutically acceptable salt thereof at an amount of 1.5 to 9.0 mg/kg/day.

13. The use according to claim 12, wherein the second medicament is formulated for administration of anti-HER2 antibody at an amount of 0.1 - 4.5 mg/kg/day.

14. The use according to claim 12 or 13, wherein the first medicament is formulated for administration of amrubicin or a pharmaceutically acceptable salt thereof every three weeks.

15. The use according to claim 14, wherein the second medicament is formulated for administration of an anti-HER2 antibody every one week.

16. The use according to any one of claims 6 to 15, wherein the first and second medicaments are formulated to provide a dosage ratio of amrubicin or a pharmaceutically acceptable salt thereof to an anti-HER2 antibody of from 2:1 to 8:1 for one week after initiation of the administration.

17. A kit for treating breast cancer comprising: (1) a first composition including amrubicin, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient; and (2) a second composition including an anti-HER2 antibody and a pharmaceutically acceptable carrier or excipient.

18. A medicament for treating breast cancer comprising an effective amount of amrubicin, or a pharmaceutically acceptable salt thereof, for use in combination with an effective amount of an anti-HER2 antibody.

19. The medicament according to claim 18, wherein the effective amount of amrubicin, or a pharmaceutically acceptable salt thereof is 0.5 to 3.0 mg/kg/day.

20. The medicament according to claim 18, wherein the effective amount of amrubicin, or a pharmaceutically acceptable salt thereof is 1.5 to 9.0 mg/kg/day.

21. The medicament according to claim 18, wherein the effective amount of an anti-HER2 antibody is 0.1 to 4.5 mg/kg/day.