



US010604765B2

(12) **United States Patent**
Cole et al.

(10) **Patent No.:** **US 10,604,765 B2**

(45) **Date of Patent:** **Mar. 31, 2020**

(54) **SOYBEAN PLANT AND SEED
CORRESPONDING TO TRANSGENIC EVEN
MON87712 AND METHODS FOR
DETECTION THEREOF**

2004/0078852 A1 4/2004 Thomashow et al.
2007/0128225 A1 6/2007 Prior et al.
2014/0380520 A1 12/2014 Khanna et al.
2019/0002907 A1 1/2019 Cole et al.

(71) Applicant: **Monsanto Technology LLC**, St. Louis,
MO (US)

FOREIGN PATENT DOCUMENTS

WO WO 2008/005210 1/2008
WO WO 2009/055597 4/2009
WO WO 2009/064652 5/2009
WO WO 2009/102873 8/2009

(72) Inventors: **Robert H. Cole**, Florissant, MO (US);
John A. Korte, Westerly, RI (US);
John R. LeDeaux, St. Louis, MO (US);
Melissa Compton Spears, St. Louis,
MO (US); **Kunsheng Wu**, Ballwin, MO
(US)

OTHER PUBLICATIONS

(73) Assignee: **Monsanto Technology LLC**, St. Louis,
MO (US)

U.S. Appl. No. 16/033,088, filed Jul. 11, 2018, Cole et al.
U.S. Appl. No. 15/802,397, filed Nov. 2, 2017, Khanna et al.
Anklam et al., "Analytical methods for detection and determination
of genetically modified organisms in agricultural crops and plant-
derived food products", *Eur. Food Res. Technol.*, 214:3-26, 2011.
Axelos et al., "The gene family encoding the *Arabidopsis thaliana*
translation elongation factor EF-1 alpha: molecular cloning, char-
acterization and expression," *Mol. Gen. Genet.*, 219:106-112, 1989.
Barker et al., "Nucleotide sequence of the T-DNA region from the
Agrobacterium tumefaciens octopine Ti plasmid pTi15955," *Plant
Mol. Biol.*, 2:335-350, 1983.
Coruzzi et al., "Tissue-specific and light-regulated expression of a
pea nuclear gene encoding the small subunit of ribulose-1,5-
bisphosphate carboxylase," *EMBO J.*, 3:1671-1679, 1984.
Depicker et al., "Nopaline synthase: transcript mapping and DNA
sequence," *J. Mol. Appl. Genet.*, 1:561-573, 1982.
Fling et al., "Nucleotide sequence of the transposon Tn7 gene
encoding an aminoglycoside-modifying enzyme, 3'(9)-O-
nucleotidyltransferase," *Nucl. Acids Res.*, 13(19):7095-7106, 1985.
Giza et al., "A self-inducing runaway-replication plasmid expres-
sion system utilizing the Rop protein," *Gene*, 78:73-84, 1989.
Holtan et al., "BBX32, an *Arabidopsis* B-Box protein, functions in
light signaling by suppressing HY5-regulated gene expression and
interacting with STH2/BBX21," *Plant Physiol.*, 156(4):2109-2123,
2011.
John, "Structural characterization of genes corresponding to cotton
fiber mRNA, E6: reduced E6 protein in transgenic plants by
antisense gene," *Plant Mol. Biol.*, 30:297-306, 1996.

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/033,091**

(22) Filed: **Jul. 11, 2018**

(65) **Prior Publication Data**

US 2019/0002908 A1 Jan. 3, 2019

Related U.S. Application Data

(62) Division of application No. 15/343,049, filed on Nov.
3, 2016, now Pat. No. 10,053,704, which is a division
of application No. 13/879,238, filed as application
No. PCT/US2011/055799 on Oct. 11, 2011, now Pat.
No. 9,493,786.

(60) Provisional application No. 61/393,448, filed on Oct.
15, 2010, provisional application No. 61/392,267,
filed on Oct. 12, 2010.

(51) **Int. Cl.**

C12N 15/82 (2006.01)
A01H 1/02 (2006.01)
C12Q 1/68 (2018.01)
A01H 5/10 (2018.01)
C12Q 1/6895 (2018.01)

(52) **U.S. Cl.**

CPC **C12N 15/8261** (2013.01); **A01H 1/02**
(2013.01); **A01H 5/10** (2013.01); **C12Q**
1/6895 (2013.01); **C12Q 2600/13** (2013.01);
C12Q 2600/156 (2013.01); **Y02A 40/146**
(2018.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,067,627 B2 6/2006 Cohen et al.
7,692,067 B2 4/2010 Creelman et al.
7,960,607 B2 6/2011 Molinero
8,088,976 B2 1/2012 Boukharov et al.

(Continued)

Primary Examiner — Bratislav Stankovic

(74) *Attorney, Agent, or Firm* — Dentons US LLP; David
Lanzotti, Esq.

(57) **ABSTRACT**

The present invention provides a transgenic soybean com-
prising event MON87712 that exhibits increased yield. The
invention also provides cells, plant parts, seeds, plants,
commodity products related to the event, and DNA mol-
ecules that are unique to the event and were created by the
insertion of transgenic DNA into the genome of a soybean
plant. The invention further provides methods for detecting
the presence of said soybean event nucleotide sequences in
a sample, probes and primers for use in detecting nucleotide
sequences that are diagnostic for the presence of said
soybean event.

10 Claims, 2 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

Kay et al., "Duplication of CaMV 35S promoter sequences creates a strong enhancer for plant genes," *Science*, 236:1299-1302, 1987.

Khanna et al., "The *Arabidopsis* B-Box Zinc Finger Family," *Plant Cell*, 21:3416-3420, 2009.

Koyejo et al., "Petition for the Determination of Nonregulated Status for MON 87712 Soybean", Jul. 19, 2011.

Klee et al., "Cloning of an *Arabidopsis thaliana* gene encoding 5-enolpyruvylshikimate-3-phosphate synthase: sequence analysis and manipulation to obtain glyphosate-tolerant plants," *Molec. Gen. Genet.*, 210:437-442, 1987.

LW et al., "An Accurate and Rapid PCR-Based Zygosity Testing Method for Genetically Modified Maize," *GMO Biosafety Research*, 1(1):1-4, 2010.

Odell et al., "Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter," *Nature*, 313:810-812, 1985.

Padgett et al., "New Weed Control Opportunities: Development of Soybeans with a Roundup Ready™ Gene," in *Herbicide-Resistant Crops* Ch. 4, pp. 53-84, CRC Press, 1996.

Richins et al., "Sequence of figwort mosaic virus DNA (caulimovirus group)," *Nucl. Acids Res.*, 15:8451-8466, 1987.

Stalker et al., "Nucleotide sequence of the region of the origin of replication of the broad host range plasmid RK2," *Molec. Gen. Genet.*, 181:8-12, 1981.

Sutcliffe, "Complete nucleotide sequence of the *Escherichia coli* plasmid pBR322," *Cold Spring Harbor Symposia on Quantitative Biology* 43:77-90, Cold Spring Harbor Laboratory Press, 1979.

Zambryski et al., "Tumor induction by *Agrobacterium tumefaciens*: analysis of the boundaries of T-DNA," *J Mol. Appl. Genet.*, 1:361-370, 1982.

Arabidopsis thaliana At3g21150 mRNA, complete cds, GenBank, published Mar. 14, 2003.

Office Action regarding Indonesian Application No. W00201302014, dated Jun. 27, 2016.

Response to Office Action regarding U.S. Appl. No. 16/033,088, filed Jan. 27, 2020.




Figure 1





Figure 2



**SOYBEAN PLANT AND SEED
CORRESPONDING TO TRANSGENIC EVEN
MON87712 AND METHODS FOR
DETECTION THEREOF**

CROSS REFERENCE TO RELATED
APPLICATIONS

This application is a Division of U.S. application Ser. No. 15/343,049 filed Nov. 3, 2016, which is a Division of U.S. application Ser. No. 13/879,238, filed Sep. 9, 2013, now U.S. Pat. No. 9,493,786, which is a 371 National Stage Entry of International Application No. PCT/US2011/055799, filed Oct. 11, 2011, which claims the benefit of U.S. Provisional Application Nos. 61/392,267 and 61/393,448, filed Oct. 12, 2010 and Oct. 15, 2010, respectively, all of which are incorporated herein by reference in their entireties.

INCORPORATION OF SEQUENCE LISTING

The sequence listing that is contained in the file named "MONS299WO.txt", which is 26.3 kilobytes as measured in Microsoft Windows operating system and was created on 5 Oct. 2011, is filed electronically herewith and incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to transgenic soybean event MON87712 and plants comprising the event that exhibit increased yield. The invention also provides cells, plant parts, seeds, plants, commodity products related to the event, and DNA molecules that are unique to the event and were created by the insertion of transgenic DNA into the genome of a soybean plant. The invention further provides methods for detecting the presence of said soybean event nucleotide sequences in a sample, probes and primers for use in detecting nucleotide sequences that are diagnostic for the presence of said soybean event.

BACKGROUND OF THE INVENTION

Soybean (*Glycine max*) is an important crop and is a primary food source in many areas of the world. The methods of biotechnology have been applied to soybean for improvement of agronomic traits and the quality of the product. One such agronomic trait is increased yield.

Increased yield may be achieved in transgenic plants by the expression of a transgene capable of providing such increased yield. The expression of foreign genes in plants is known to be influenced by many factors, such as the regulatory elements used in the transgene cassette, the chromosomal location of the transgene insert, the proximity of any endogenous regulatory elements close to the transgene insertion site, and environmental factors such as light and temperature. For example, it has been observed that there may be a wide variation in the overall level of transgene expression or in the spatial or temporal pattern of transgene expression between similarly-produced events. For this reason, it is often necessary to screen hundreds of independent transformation events in order to ultimately identify one event useful for commercial agricultural purposes. Such an event, once identified as having the desired transgene expression, molecular characteristics and the improved trait, may then be used for introgressing the improved trait into other genetic backgrounds using plant breeding methods. The resulting progeny would contain the

transgenic event and would therefore have the transgene expression characteristics for that trait of the original transformant. This may be used to produce a number of different crop varieties that comprise the improved trait and are suitably adapted to specific local growing conditions.

SUMMARY OF THE INVENTION

The present invention provides transgenic soybean plants and seeds comprising event MON87712, a representative seed sample of which have been deposited with American Type Culture Collection (ATCC) under the Accession No. PTA-10296. Plants comprising the event exhibit increased yield.

The invention provides a plant, seed, cell, progeny plant, or plant part comprising the event and commodity products derived from a plant, cell, plant part, or seed comprising event MON87712. The invention thus provides a plant, seed, cell, progeny plant, plant part, or commodity product comprising a DNA molecule having a nucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 7, SEQ ID NO: 8 and SEQ ID NO: 6, and complements thereof. The invention provides a plant, seed, cell, progeny plant, or plant part comprising a recombinant DNA molecule that produces an amplicon comprising a DNA molecule of the invention, for instance in a DNA amplification method. The plant parts of the soybean comprising event MON87712 of the present invention include, but are not limited to pollen, ovule, flower, shoot, root, stem, leaf, pod and seed. Novel genetic compositions contained in the genome of a plant comprising event MON87712 and products made from a plant comprising event MON87712 such as whole or processed seed, animal feed, oil, meal, flour, food products, protein supplements, fuel products and biomass from which soybean plants comprising MON87712 have been harvested are aspects of this invention.

The invention provides DNA molecules related to event MON87712. These DNA molecules may comprise nucleotide sequences representing or derived from the junction of the transgene insertion and flanking genomic DNA of event MON87712, and/or a region of the genomic DNA flanking the inserted DNA, and/or a region of the integrated transgenic DNA flanking the insertion site, and/or a region of the integrated transgenic expression cassette, and/or a contiguous sequence of any of these regions.

In one embodiment, the invention provides a DNA molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 7, SEQ ID NO: 8 and SEQ ID NO: 6, and complements thereof. In another embodiment, the DNA molecule may comprise a polynucleotide having a sequence comprising from at least about 51-100 consecutive nucleotides of a sequence selected from the group consisting of SEQ ID NO: 7 and SEQ ID NO: 8 or a polynucleotide molecule having a sequence with at least 90% identity to SEQ ID NO: 6. In yet another embodiment, the invention provides a DNA molecule comprising SEQ ID NO: 1 and SEQ ID NO: 2. According to still another embodiment, there are no more than 5000 consecutive nucleotides between the 3' end of SEQ ID NO: 1 and the 5' end of SEQ ID NO: 2, for instance, no more than 4500, 4000, 3500, 3000, 2500, 2250, 2100, 2050, 2000, or 1992 consecutive nucleotides between the 3' end of SEQ ID NO: 1 and the 5' end of SEQ ID NO: 2. According to yet another embodiment, there are no more than 5000 consecutive nucleotides between the 3' end of SEQ ID NO: 7 and the 5'

end of SEQ ID NO: 8, for instance, no more than 4500, 4000, 3500, 3000, 2500, 2250, 2100, 2050, 2000, or 1914 consecutive nucleotides between the 3' end of SEQ ID NO: 7 and the 5' end of SEQ ID NO: 8. In a further embodiment, the invention provides a DNA molecule comprising SEQ ID NO: 1 or SEQ ID NO: 2 and further comprising a BBX32 coding sequence. In some embodiments, the BBX32 coding sequence is operably or genetically linked to SEQ ID NO: 1 or SEQ ID NO: 2. In other embodiments, the BBX32 coding sequence is flanked by SEQ ID NO: 1 and SEQ ID NO: 2.

The invention also provides DNA molecules useful as primers and probes diagnostic for the event. Plants, cells, plant parts, commodity products, progeny, and seeds comprising these molecules are provided.

According to one aspect of the invention, compositions and methods are provided for detecting the presence of the transgene/genomic insertion region from a novel soybean plant comprising MON87712. DNA sequences are provided that comprise at least one junction sequence of MON87712 selected from the group consisting of SEQ ID NO: 1 (corresponding to positions 3495 through 3516 of SEQ ID NO: 6, FIG. 1 [F]), SEQ ID NO: 2 (corresponding to positions 5509 through 5530 of SEQ ID NO: 6, FIG. 1 [F]), SEQ ID NO: 7 (corresponding to positions 3456 through 3555 of SEQ ID NO: 6), SEQ ID NO: 8 (corresponding to positions 5470 through 5569 of SEQ ID NO: 6), and complements thereof; wherein a junction sequence is a nucleotide sequence that spans the point at which heterologous DNA inserted into the genome is linked to the soybean cell genomic DNA and detection of this sequence in a biological sample containing soybean DNA is diagnostic for the presence of the soybean event MON87712 DNA in said sample (FIG. 1). A soybean event MON87712 and soybean seed comprising these DNA molecules is an aspect of this invention.

According to another aspect of the invention, two DNA molecules are provided for use in a DNA detection method, wherein the DNA molecules function as DNA primers when used together in an amplification reaction with a template derived from event MON87712 to produce an amplicon diagnostic for event MON87712 DNA in a sample. In one embodiment, a first and second DNA molecules comprise a polynucleotide molecule having a nucleotide sequence of sufficient length of consecutive nucleotides of SEQ ID NO: 6, or a complement thereof.

In another embodiment, the first DNA molecule comprises a polynucleotide having a nucleotide sequence of sufficient length of consecutive polynucleotide of any portion of the transgene region of the DNA molecule of SEQ ID NO: 3 or SEQ ID NO: 5 and the second DNA molecule comprises a polynucleotide of similar length of any portion of a 5' flanking soybean genomic DNA region of SEQ ID NO: 3. Any amplicon produced by DNA primers homologous or complementary to any portion of SEQ ID NO: 3 and SEQ ID NO: 5, and any amplicon that comprises SEQ ID NO: 1 or at least 51 consecutive nucleotides of SEQ ID NO: 7 is an aspect of the invention.

According to another aspect of the invention, the first DNA molecule comprises a polynucleotide having a nucleotide sequence of sufficient length of consecutive polynucleotide of any portion of the transgene region of the DNA molecule of SEQ ID NO: 4 or SEQ ID NO: 5 and the second DNA molecule comprises a polynucleotide of similar length of any portion of a 3' flanking soybean genomic DNA of SEQ ID NO: 4. Any amplicons produced by DNA primers homologous or complementary to any portion of SEQ ID NO: 4 and SEQ ID NO: 5, and any amplicon that comprises

SEQ ID NO: 2 or at least 51 consecutive nucleotides of SEQ ID NO: 8 is an aspect of the invention.

The invention provides methods, compositions, and kits useful for detecting the presence of DNA derived from soybean event MON87712. Certain methods comprise (a) contacting a sample comprising DNA with a primer set that when used in a nucleic acid amplification reaction with genomic DNA from a soybean comprising event MON87712 produces an amplicon diagnostic for the event; (b) performing a nucleic acid amplification reaction thereby producing the amplicon; and (c) detecting the amplicon, wherein said amplicon comprises SEQ ID NO: 1 and/or SEQ ID NO: 2, or at least 51 consecutive nucleotides of SEQ ID NO: 7 or at least 51 consecutive nucleotides of SEQ ID NO: 8. The invention also provides a method for detection of the event by (a) contacting a sample comprising DNA with a probe that when used in a hybridization reaction with genomic DNA from the event hybridizes to a DNA molecule specific for the event; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting the hybridization of the probe to the DNA molecule. Kits comprising the methods and compositions of the invention useful for detecting the presence of DNA derived from the event are also provided.

Another aspect of the invention is a method of determining the zygosity of a soybean plant genome comprising soybean event MON87712 DNA in a sample comprising: a) contacting the sample with three different primers that i) when used together in a nucleic acid amplification reaction with soybean event MON87712 DNA, produces a first amplicon that is diagnostic for soybean event MON87712; and ii) when used together in a nucleic acid amplification reaction with soybean genomic DNA other than MON87712 DNA, produces a second amplicon that is diagnostic for soybean wild type genomic DNA other than event MON87712 DNA; b) performing a nucleic acid amplification reaction; and c) detecting said first amplicon and said second amplicon, wherein the presence of said first and second amplicons is diagnostic of a heterozygous genome in said sample, and wherein the presence of only said first amplicon is diagnostic of a homozygous genome in said sample. In one embodiment the primers comprise SEQ ID NO: 10, SEQ ID NO: 11 or 15, and SEQ ID NO: 13.

In one embodiment the method of determining the zygosity of a soybean comprising event MON87712, comprises (a) contacting the sample comprising soybean DNA with the primer set SEQ ID NO: 10, SEQ ID NO: 11 or SEQ ID NO: 15, SEQ ID NO: 13, and the probe set 6FAMTM-labeled SEQ ID NO: 12 and VICTM-labeled SEQ ID NO: 14 that when used in a nucleic acid amplification reaction with genomic DNA from a soybean comprising event MON87712, produces a first amplicon, releasing a fluorescent signal from the combination of primers SEQ ID NO: 10 and SEQ ID NO: 11 or 15 and the 6FAMTM-labeled probe (SEQ ID NO: 12) that is diagnostic for soybean event MON87712; (b) performing a nucleic acid amplification reaction, thereby producing the first amplicon; and (c) detecting said first amplicon; and (d) contacting the sample comprising soybean DNA with the primer set, SEQ ID NO: 10 and SEQ ID NO: 13, and the VICTM-labeled probe (SEQ ID NO: 14) that when used in a nucleic acid amplification reaction with genomic DNA from soybean plants produces a second amplicon, releasing a fluorescent signal that is diagnostic of wild-type soybean genomic DNA homologous to the soybean genomic region of a transgene insertion identified as soybean event MON87712; (e) performing a nucleic acid amplification reaction, thereby producing the second ampli-

con; and (f) detecting said second amplicon; and (g) comparing the first and the second amplicons in a sample, wherein the presence of both amplicons indicates the sample is heterozygous for the transgene insertion.

The invention further provides a method of producing a soybean plant with increased yield comprising: (a) selfing a MON87712 comprising soybean plant, thereby producing a seed; (b) growing said seed to produce a plurality of progeny plant; and (c) selecting a progeny plant that comprises MON87712 or a progeny plant with increased yield. Another aspect of the invention provides a method of producing a soybean plant with increased yield comprising: (a) crossing a MON87712 comprising soybean plant with a second soybean plant, thereby producing a seed; (b) growing said seed to produce a plurality of progeny plants; and (c) selecting a progeny plant that comprises MON87712 or a progeny plant with increased yield.

In yet another aspect, the invention provides a method of increasing yield in a crop comprising a) planting a crop plant or seed comprising event MON87712; and b) growing said crop plant or seed.

The foregoing and other aspects of the invention will become more apparent from the following detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Diagrammatical representation of the transgenic insert in the genome of a soybean comprising event MON87712; [A] corresponds to the relative positions of SEQ ID NO: 1 and SEQ ID NO: 7, both forming the junction between the 5' portion of the transgenic insert and the 3' portion of the flanking genomic DNA; [B] corresponds to the relative positions of SEQ ID NO: 2 and SEQ ID NO: 8, both forming the junction between the 3' portion of the transgenic insert and the 5' portion of the flanking genomic DNA; [C] corresponds to the relative position of SEQ ID NO: 3, which contains the soybean genomic flanking region and a portion of the arbitrarily designated 5' end of the transgenic DNA insert; [D] corresponds to the relative position of SEQ ID NO: 4, which contains the soybean genome flanking region and a portion of the arbitrarily designated 3' end of the transgenic DNA insert; [E] represents SEQ ID NO: 5, which is the sequence of the transgenic DNA insert including the BBX32 expression cassette integrated into the genome of a soybean plant comprising event MON87712; [F] represents SEQ ID NO: 6, which is the contiguous sequence comprising the 5' flanking genomic sequence, the transgenic insert and the 3' flanking genomic sequence, comprising as represented in the figure from left to right, SEQ ID NO: 3, SEQ ID NO: 5 and SEQ ID NO: 4, in which SEQ ID NOs: 1 and 7, and SEQ ID NOs: 2 and 8 are incorporated as set forth above, as these sequences are present in the genome of a plant comprising event MON87712; [G] and [H] represent forward primers (SEQ ID NO: 11 or SEQ ID NO: 15, and SEQ ID NO: 13) for event-specific zygosity endpoint TAQMAN® PCR for identification of MON87712 and wild-type allele, respectively; [I] represents reverse primer (SEQ ID NO: 10) for identification of MON87712 and wild-type alleles; [J] and [K] represent probes (SEQ ID NO: 12 and SEQ ID NO: 14) used for event-specific zygosity endpoint TAQMAN® PCR for identification of MON87712 and wild-type allele, respectively. Arrows indicate the direction of 5' to 3'. LB: refers to the left border of T-DNA; RB: refers to the right border of T-DNA.

FIG. 2. Meta-analysis of yield data for 9 transgenic events from 4 field seasons of field testing in the United States or South America, showing event 1 comprising MON87712 DNA with the highest yield. Panel A: average yield increases of transgenic events over their corresponding negative isolines (bushel/acre). Panel B: average yield increases of transgenic events over wild type control (bushel/acre). The numbers above each bar represent % increase.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO: 1—A 22 bp nucleotide sequence representing the left border junction between the soybean genomic DNA and the integrated DNA insert. This sequence corresponds to positions 3495 to 3516 of SEQ ID NO: 6, and to positions 3495 through 3516 of SEQ ID NO: 3 ([C] of FIG. 1). In addition, SEQ ID NO: 1 corresponds to the integrated left border of the expression cassette at positions 1 through 11 of SEQ ID NO: 5 ([E] of FIG. 1).

SEQ ID NO: 2—A 22 bp nucleotide sequence representing the right border junction between the integrated DNA insert and the soybean genomic DNA. This sequence corresponds to positions 5509 to 5530 of SEQ ID NO: 6, and to positions 90 through 111 of SEQ ID NO: 4 ([D] of FIG. 1). In addition, SEQ ID NO: 2 corresponds to positions 2004 through 2014 SEQ ID NO: 5 ([E] of FIG. 1).

SEQ ID NO: 3—A 3605 bp nucleotide sequence including the 5' soybean genomic sequence (3505 bp) flanking the inserted DNA of event MON87712 plus a region (100 bp) of the integrated DNA. This sequence corresponds to positions 1 through 3605 of SEQ ID NO: 6. It is believed that the sequence at position 3451 to 3463 of SEQ ID NO: 3 may contain anywhere from 13 to 100 individual c residues in a row.

SEQ ID NO: 4—A 2065 bp nucleotide sequence including the 3' soybean genomic sequence (1965 bp) flanking the inserted DNA of event MON87712 plus a region (100 bp) of the integrated DNA. This sequence corresponds to positions 5420 through 7484 of SEQ ID NO: 6.

SEQ ID NO: 5—The sequence of the integrated expression cassette imparting increased yield, including the left and the right border sequences after integration. SEQ ID NO: 5 corresponds to nucleotide positions 3506 through 5519 of SEQ ID NO: 6.

SEQ ID NO: 6—A 7484 bp nucleotide sequence representing the contig of the 5' sequence flanking the inserted DNA of MON87712 (SEQ ID NO: 3), the sequence of the integrated DNA insert (SEQ ID NO: 5) and the 3' sequence flanking the inserted DNA of MON87712 (SEQ ID NO: 4). It is believed that the sequence at position 3451 to 3463 of SEQ ID NO: 6 may contain anywhere from 13 to 100 individual c residues in a row.

SEQ ID NO: 7: A 100 bp nucleotide sequence representing the left border junction between soybean genomic DNA and the integrated DNA insert. This sequence corresponds to positions 3456 to 3555 of SEQ ID NO: 6, and to positions 3456 through 3555 of SEQ ID NO: 3 ([C] of FIG. 1). In addition, SEQ ID NO: 7 corresponds to the integrated left border of the expression cassette at positions 1 through 50 of SEQ ID NO: 5 ([E] of FIG. 1).

SEQ ID NO: 8—A 100 bp nucleotide sequence representing the right border junction between the integrated DNA insert and the soybean genomic DNA. This sequence corresponds to positions 5470 to 5569 of SEQ ID NO: 6, and to positions 51 through 150 of SEQ ID NO: 4 ([D] of FIG. 1). In addition, SEQ ID NO: 8 corresponds to positions 1965 through 2014 SEQ ID NO: 5 ([E] of FIG. 1).

SEQ ID NO: 9—The nucleotide sequence of the BBX32 expression cassette of pMON83132.

SEQ ID NO: 10—The reverse primer sequence used in event-specific zygosity endpoint TAQMAN PCR assay for identification of both MON87712 DNA/allele and wild-type DNA/allele. SEQ ID NO: 10 corresponds to nucleotide positions 5517 to 5546 of SEQ ID NO: 6. Production of a PCR amplicon using the combination of primers SEQ ID NO: 10 and SEQ ID NO: 11 or 15 is a positive result for the presence of event MON87712.

SEQ ID NO: 11—The forward primer sequence used to identify MON87712 event and the zygosity of MON87712 event. SEQ ID NO: 11 corresponds to nucleotide positions 5435 to 5458 of SEQ ID NO: 6.

SEQ ID NO: 12—The probe sequence used to identify MON87712 event and the zygosity of MON87712 event. SEQ ID NO: 12 is a 6FAMTM-labeled synthetic oligonucleotide. Release of a fluorescent signal in an amplification reaction using primers SEQ ID NO: 10 and SEQ ID NO: 11 or 15 in combination with the 6FAMTM-labeled probe in a TAQMAN[®] assay is diagnostic of event MON87712.

SEQ ID NO: 13—The forward primer sequence used in event-specific zygosity endpoint TAQMAN[®] PCR for identification of wild-type DNA/allele. Production of a PCR amplicon using primers SEQ ID NO: 13 and SEQ ID NO: 10 is diagnostic of wild-type DNA.

SEQ ID NO: 14—The probe sequence used to determine the presence of soybean wild-type DNA/allele. SEQ ID NO: 14 is a VICTM-labeled synthetic oligonucleotide. Release of a fluorescent signal in an amplification reaction using primers SEQ ID NO: 10 and SEQ ID NO: 13 in combination with the VICTM-labeled probe in a TAQMAN[®] assay is diagnostic of the wild-type allele in a zygosity assay.

SEQ ID NO: 15—An alternative forward primer sequence used to identify MON87712 event and the zygosity of MON87712 event. SEQ ID NO: 15 corresponds to nucleotide positions 5432 to 5455 of SEQ ID NO: 6.

SEQ ID NO: 16—The sequence of primer SQ3983 used to identify MON87712 event. Production of a 97 bp PCR amplicon using the combination of primers SQ3983 and SQ22982 (SEQ ID NO: 17) is a positive result for the presence of event MON87712.

SEQ ID NO: 17—The sequence of primer SQ22982 used to identify MON87712 event.

SEQ ID NO: 18—The sequence of probe PB10453 used to identify MON87712 event. It is a 6FAMTM-labeled synthetic oligonucleotide.

SEQ ID NO: 19—The sequence of primer SQ1532 used as an internal control in end-point TAQMAN[®] assays.

SEQ ID NO: 20—The sequence of primer SQ1533 used as an internal control in end-point TAQMAN[®] assays.

SEQ ID NO: 21—The sequence of a VICTM-labeled synthetic oligonucleotide probe PB0359 used as an internal control in end-point TAQMAN[®] assays.

DETAILED DESCRIPTION

The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger et al., *Glossary of Genetics: Classical and Molecular*, 5th edition, Springer-Verlag: New York, 1991; and Lewin, *Genes V*, Oxford University Press: New York, 1994.

The present invention provides transgenic soybean event MON87712. The term “event” as used herein refers to DNA molecules produced as a result of inserting transgenic DNA into a plant’s genome at a particular location on a chromosome. Event MON87712 refers to the DNA molecules produced as a result of the insertion of transgenic DNA having a sequence provided herein as SEQ ID NO: 5 into a particular chromosomal location in the *Glycine max* genome. Plants, seeds, progeny, cells, and plant parts thereof comprising event MON87712 are also provided in the present invention. A sample of seed comprising MON87712 has been deposited with American Type Culture Collection (ATCC) under Accession No. PTA-10296 on Aug. 20, 2009. Plants comprising MON87712 exhibit increased yield.

As used herein, the term “soybean” means *Glycine max* and includes all plant varieties that can be bred with soybean, including wild soybean species as well as those plants belonging to *Glycine soja* that permit breeding between species.

A transgenic “event” is produced by transformation of plant cells with heterologous DNA, i.e., a nucleic acid construct that comprises a transgene of interest, regeneration of a population of independently transformed transgenic plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant with desirable molecular characteristics, such as insertion of a single copy of the transgene into a particular genome location, integrity of the transgenic DNA, and an enhanced trait such as increased yield. A plant comprising the event can refer to the original transformant that includes the transgene inserted into the particular location in the plant’s genome. A plant comprising the event can also refer to progeny of the original transformant that retain the transgene at the same particular location in the plant’s genome. Such progeny may be produced by selfing or by a sexual outcross between the transformant, or its progeny, and another plant. Such another plant may be a transgenic plant comprising the same or a different transgene; or may be a non-transgenic plant, such as one from a different variety. The resulting progeny may be homozygous or heterozygous for event MON87712 DNA (inserted DNA and flanking DNA). Even after repeated back-crossing to a recurrent parent, the event DNA from the transformed parent is present in the progeny of the cross at the same genomic location.

A DNA molecule comprising event MON87712 refers to a DNA molecule comprising at least a portion of the inserted transgenic DNA (provided as SEQ ID NO: 5) and at least a portion of the flanking genomic DNA immediately adjacent to the inserted DNA. As such, a DNA molecule comprising event MON87712 has a nucleotide sequence representing at least a portion of the transgenic DNA insert and at least a portion of the particular region of the genome of the plant into which the transgenic DNA was inserted. The arrangement of the inserted DNA in event MON87712 in relation to the surrounding plant genome is specific and unique to event MON87712 and as such the nucleotide sequence of such a DNA molecule is diagnostic and identifying for event MON87712. Examples of the sequence of such a DNA molecule are provided herein as SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 7, SEQ ID NO: 8 and SEQ ID NO: 6. Such a DNA molecule is also an integral part of the chromosome of a plant that comprises event MON87712 and may be passed on to progenies of the plant.

As used herein, a “recombinant DNA molecule” is a DNA molecule comprising a combination of DNA molecules that would not naturally occur together and is the result of human

intervention, e.g., a DNA molecule that is comprised of a combination of at least two DNA molecules heterologous to each other, and/or a DNA molecule that is artificially synthesized and comprises a polynucleotide sequence that deviates from the polynucleotide sequence that would normally exist in nature, and/or a DNA molecule that comprises a transgene artificially incorporated into a host cell's genomic DNA and the associated flanking DNA of the host cell's genome. An example of a recombinant DNA molecule is a DNA molecule described herein resulting from the insertion of the transgene into the *Glycine max* genome, which may ultimately result in the expression of a recombinant RNA and/or protein molecule in that organism. The nucleotide sequence or any fragment derived therefrom would also be considered a recombinant DNA molecule if the DNA molecule can be extracted from cells, or tissues, or homogenate from a plant or seed or plant tissue; or can be produced as an amplicon from extracted DNA or RNA from cells, or tissues, or homogenate from a plant or seed or plant tissue, any of which is derived from such materials derived from a plant comprising event MON87712. For that matter, the junction sequences as set forth at SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7 and SEQ ID NO: 8, and nucleotide sequences derived from event MON87712 that also contain these junction sequences are considered to be recombinant DNA, whether these sequences are present within the genome of the cells comprising event MON87712 or present in detectable amounts in tissues, progeny, biological samples or commodity products derived from plants comprising event MON87712. As used herein, the term "transgene" refers to a polynucleotide molecule artificially incorporated into a host cell's genome. Such transgene may be heterologous to the host cell. The term "transgenic plant" refers to a plant comprising such a transgene. A "transgenic plant" includes a plant, plant part, a plant cell or seed whose genome has been altered by the stable integration of recombinant DNA. A transgenic plant includes a plant regenerated from an originally-transformed plant cell and progeny transgenic plants from later generations or crosses of a transformed plant. As a result of such genomic alteration, the transgenic plant is distinctly different from the related wild type plant. An example of a transgenic plant is a plant described herein as comprising event MON87712.

As used herein, the term "heterologous" refers to a sequence which is not normally present in a given host genome in the genetic context in which the sequence is currently found. In this respect, the sequence may be native to the host genome, but be rearranged with respect to other genetic sequences within the host sequence.

The present invention provides DNA molecules and their corresponding nucleotide sequences. As used herein, the terms "DNA sequence", "nucleotide sequence" and "polynucleotide sequence" refer to the sequence of nucleotides of a DNA molecule, usually presented from the 5' (upstream) end to the 3' (downstream) end. The nomenclature used herein is that required by Title 37 of the United States Code of Federal Regulations § 1.822 and set forth in the tables in WIPO Standard ST.25 (1998), Appendix 2, Tables 1 and 3. The present invention is disclosed with reference to only one strand of the two nucleotide sequence strands that are provided in transgenic event MON87712. Therefore, by implication and derivation, the complementary sequences, also referred to in the art as the complete complement or the reverse complementary sequences, are within the scope of the present invention and are therefore also intended to be within the scope of the subject matter claimed.

The nucleotide sequence corresponding to the complete nucleotide sequence of the inserted transgenic DNA and substantial segments of the *Glycine max* genomic DNA flanking either end of the inserted transgenic DNA is provided herein as SEQ ID NO: 6. A subsection of this is the inserted transgenic DNA provided as SEQ ID NO: 5. The nucleotide sequence of the genomic DNA flanking the 5' end of the inserted transgenic DNA and a portion of the 5' end of the inserted DNA is provided herein as SEQ ID NO: 3. The nucleotide sequence of the genomic DNA flanking the 3' end of the inserted transgenic DNA and a portion of the 3' end of the inserted DNA is provided herein as SEQ ID NO: 4. The region spanning the location where the transgenic DNA connects to and is linked to the genomic DNA is referred to herein as the junction. A "junction sequence" or "junction region" refers to a DNA sequence and/or corresponding DNA molecule that spans the inserted transgenic DNA and the adjacent flanking genomic DNA. Examples of a junction sequence of event MON87712 are provided herein as SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7, and SEQ ID NO: 8. The identification of one of these junction sequences in a nucleotide molecule derived from a soybean plant or seed is conclusive that the DNA was obtained from event MON87712 and is diagnostic for the presence of DNA from event MON87712. SEQ ID NO: 1 is a 22 bp nucleotide sequence spanning the junction between the genomic DNA and the 5' end of the inserted DNA. SEQ ID NO: 7 is a 100 bp nucleotide sequence spanning the junction between the genomic DNA and the 5' end of the inserted DNA. SEQ ID NO: 2 is a 22 bp nucleotide sequence spanning the junction between the genomic DNA and the 3' end of the inserted DNA. SEQ ID NO: 8 is a 100 bp nucleotide sequence spanning the junction between the genomic DNA and the 3' end of the inserted DNA. Any segment of DNA derived from transgenic event MON87712 that includes the consecutive nucleotides of SEQ ID NO: 1 or 51 consecutive nucleotides, 55 consecutive nucleotides, 60 consecutive nucleotides, 65 consecutive nucleotides, 70 consecutive nucleotides, 75 consecutive nucleotides, 80 consecutive nucleotides, 85 consecutive nucleotides, 90 consecutive nucleotides, 95 consecutive nucleotides, or all of the nucleotides of SEQ ID NO: 7 is within the scope of the present invention. Any segment of DNA derived from transgenic event MON87712 that includes the consecutive nucleotides of SEQ ID NO: 2 or 51 consecutive nucleotides, 55 consecutive nucleotides, 60 consecutive nucleotides, 65 consecutive nucleotides, 70 consecutive nucleotides, 75 consecutive nucleotides, 80 consecutive nucleotides, 85 consecutive nucleotides, 90 consecutive nucleotides, 95 consecutive nucleotides, or all of the nucleotides of SEQ ID NO: 8 is within the scope of the present invention. In addition, any polynucleotide molecule comprising a sequence complementary to any of the sequences described within this paragraph is within the scope of the present invention. FIG. 1 is an illustration of the transgenic DNA insert in the genome of a soybean comprising event MON87712, and the relative positions of SEQ ID NOs: 1-8 arranged 5' to 3'. The present invention also provides a nucleic acid molecule comprising a polynucleotide having a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the full-length of SEQ ID NO: 6.

The present invention further provides exemplary DNA molecules that can be used either as primers or probes for diagnosing the presence of DNA derived from event MON87712 in a sample. Such primers or probes are specific for a target nucleic acid sequence and as such are useful for

the identification of event MON87712 nucleic acid sequence by the methods of the invention described herein.

A "probe" is an isolated nucleic acid to which is attached a detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of genomic DNA from a soybean comprising event MON87712 whether from a soybean plant or from a sample that comprises DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that bind specifically to a target DNA sequence and the detection of such binding can be used to diagnose/determine/confirm the presence of that target DNA sequence in a particular sample.

A "primer" is typically an isolated polynucleotide that is designed for use in specific annealing or hybridization methods to hybridize to a complementary target DNA strand to form a hybrid between the primer and the target DNA strand, and then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. A pair of primers may be used with template DNA, such as a sample of *Glycine max* genomic DNA, in a thermal or isothermal amplification, such as polymerase chain reaction (PCR), or other nucleic acid amplification methods, to produce an amplicon, where the amplicon produced from such reaction would have a DNA sequence corresponding to sequence of the template DNA located between the two sites where the primers hybridized to the template. As used herein, an "amplicon" is a piece or fragment of DNA that has been synthesized using amplification techniques, i.e. the product of an amplification reaction. In one embodiment of the invention, an amplicon diagnostic for event MON87712 comprises a sequence not naturally found in the *Glycine max* genome. Primer pairs, as used in the present invention, are intended to refer to use of two primers binding opposite strands of a double stranded nucleotide segment for the purpose of amplifying linearly the polynucleotide segment between the positions targeted for binding by the individual members of the primer pair, typically in a thermal or isothermal amplification reaction or other nucleic acid amplification methods. Exemplary DNA molecules useful as primers are provided as SEQ ID NOs: 10-11, SEQ ID NO: 13 and SEQ ID NO: 15. The primer pair provided as SEQ ID NO: 10 and SEQ ID NO: 11 or 15 may be used as a first DNA molecule and a second DNA molecule that is different from the first DNA molecule, and both molecules are each of sufficient length of consecutive nucleotides of either SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6 or the complements thereof to function as DNA primers so that, when used together in an amplification reaction with template DNA derived from event MON87712, an amplicon that is specific and unique to transgenic event MON87712 would be produced. The use of the term "amplicon" specifically excludes primer-dimers that may be formed in the DNA amplification reaction.

Probes and primers according to the present invention may have complete sequence identity to the target sequence, although primers and probes differing from the target sequence that retain the ability to hybridize preferentially to target sequences may be designed by conventional methods. In order for a nucleic acid molecule to serve as a primer or probe it needs only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed. Any nucleic acid hybridization or amplification method can be used to identify the presence of transgenic DNA from event MON87712 in a sample. Probes and

primers are generally at least about 11 nucleotides, at least about 18 nucleotides, at least about 24 nucleotides, and at least about 30 nucleotides or more in length. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions.

Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989 (hereinafter, "Sambrook et al., 1989"); *Current Protocols in Molecular Biology*, ed. Ausubel et al., Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, "Ausubel et al., 1992"); and Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, 1990. PCR-primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, © 1991, Whitehead Institute for Biomedical Research, Cambridge, Mass.).

Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm the disclosed sequences by known methods, e.g., by re-cloning and sequencing such sequences.

The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure. A nucleic acid molecule is said to be the "complement" of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit "complete complementarity" when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be "minimally complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least "low-stringency" conditions. Similarly, the molecules are said to be "complementary" if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under "high-stringency" conditions. Stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985). Departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

As used herein, a substantially homologous sequence is a nucleic acid sequence that will specifically hybridize to the complement of the nucleic acid sequence to which it is being compared under high stringency conditions. Appropriate stringency conditions that promote DNA hybridization, for example, 6.0x sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2.0xSSC at 50° C., are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in

the wash step can be selected from a low stringency of about 2.0×SSC at 50° C. to a high stringency of about 0.2×SSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22° C., to high stringency conditions at about 65° C. Both temperature and salt may be varied, or either the temperature or the salt concentration may be held constant while the other variable is changed. In one embodiment, a nucleic acid of the present invention will specifically hybridize to one or more of the nucleic acid molecules set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 7 and SEQ ID NO: 8, or complements or fragments thereof under high stringency conditions. The hybridization of the probe to the target DNA molecule can be detected by any number of methods known to those skilled in the art. These can include, but are not limited to, fluorescent tags, radioactive tags, antibody based tags, and chemiluminescent tags.

Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, “stringent conditions” are conditions that permit the primer pair to hybridize only to the target nucleic acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA amplification reaction. Examples of DNA amplification methods include PCR, Recombinase Polymerase Amplification (RPA) (see for example U.S. Pat. No. 7,485,428), Strand Displacement Amplification (SDA) (see for example, U.S. Pat. Nos. 5,455,166 and 5,470,723), Transcription-Mediated Amplification (TMA) (see for example, Guatelli et al., *Proc. Natl. Acad. Sci. USA* 87:1874-1878 (1990)), Rolling Circle Amplification (RCA) (see for example, Fire and Xu, *Proc. Natl. Acad. Sci. USA* 92:4641-4645 (1995); Lui, et al., *J. Am. Chem. Soc.* 118:1587-1594 (1996); Lizardi, et al., *Nature Genetics* 19:225-232 (1998), U.S. Pat. Nos. 5,714,320 and 6,235,502)), Helicase Dependant Amplification (HDA) (see for example Vincent et al., *EMBO Reports* 5 (8): 795-800 (2004); U.S. Pat. No. 7,282,328), and Multiple Displacement Amplification (MDA) (see for example Dean et al., *Proc. Natl. Acad. Sci. USA* 99:5261-5266 (2002)).

The term “specific for (a target sequence)” indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

As used herein, the term “isolated” refers to at least partially separating a molecule from other molecules normally associated with it in its native or natural state. In one embodiment, the term “isolated” refers to a DNA molecule that is at least partially separated from the nucleic acids that normally flank the DNA molecule in its native or natural state. Thus, DNA molecules fused to regulatory or coding sequences with which they are not normally associated, for example as the result of recombinant techniques, are considered isolated herein. Such molecules are considered isolated even when integrated into the chromosome of a host cell or present in a nucleic acid solution with other DNA molecules.

Any number of methods well known to those skilled in the art can be used to isolate and manipulate a DNA molecule, or fragment thereof, disclosed in the present invention. For example, PCR (polymerase chain reaction) technology can be used to amplify a particular starting DNA molecule and/or to produce variants of the original molecule. DNA molecules, or fragments thereof, can also be obtained by other techniques such as by directly synthesizing the frag-

ment by chemical means, as is commonly practiced by using an automated oligonucleotide synthesizer.

It would be advantageous to be able to detect the presence of transgene/genomic DNA of a particular plant in order to determine whether progeny of a sexual cross contain the transgene/genomic DNA of interest. In addition, a method for detecting a particular plant would be helpful when complying with regulations requiring the pre-market approval and labeling of foods derived from the transgenic crop plants.

The presence of a transgene may be detected by any well known nucleic acid detection method such as the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, etc. As a result, such methods may not be useful for discriminating between different transformation events, particularly those produced using the same DNA construct unless the sequence of chromosomal DNA adjacent to the inserted DNA (“flanking DNA”) is known. An event-specific PCR assay is discussed, for example, by Taverniers et al. (*J. Agric. Food Chem.*, 53: 3041-3052, 2005) in which an event-specific tracing system for transgenic maize lines Bt11, Bt176, and GA21 and for canola event GT73 was demonstrated. In this study, event-specific primers and probes were designed based upon the sequences of the genome/transgene junctions for each event. Transgenic plant event specific DNA detection methods have also been described in U.S. Pat. Nos. 6,893,826; 6,825,400; 6,740,488; 6,733,974; 6,689,880; 6,900,014 and 6,818,807.

The DNA molecules and corresponding nucleotide sequences provided herein are therefore useful for, among other things, identifying event MON87712, selecting plant varieties or hybrids comprising event MON87712, detecting the presence of DNA derived from event MON87712 in a sample, and monitoring samples for the presence and/or absence of event MON87712 or plants and plant parts comprising event MON87712.

The present invention provides plants, progeny, seeds, plant cells, plant parts (such as pollen, ovule, pod, flower, root or stem tissue, fibers, and leaf), and commodity products. These plants, progeny, seeds, plant cells, plant parts, and commodity products contain a detectable amount of a polynucleotide of the present invention, i.e., such as a polynucleotide comprising at least one of the sequences provided as the consecutive nucleotides of SEQ ID NO: 1, the consecutive nucleotides of SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, or at least 51 consecutive nucleotides of SEQ ID NO: 8. Plants, progeny, seeds, plant cells, plant parts and commodity products of the present invention may also contain one or more additional transgenes. Such transgene may be any nucleotide sequence encoding a protein or RNA molecule conferring a desirable trait including but not limited to increased insect resistance, increased water use efficiency or drought tolerance, increased yield performance, increased nitrogen use efficiency or increase tolerance to nitrogen stress such as high or low nitrogen supply, increased seed quality, increased disease resistance, improved nutritional quality, and/or increased herbicide tolerance, such as glyphosate or dicamba tolerance, in which the desirable trait is measured with respect to a comparable plant lacking such additional transgene.

The present invention provides plants, progeny, seeds, plant cells, and plant part such as pollen, ovule, pod, flower, root or stem tissue, and leaf derived from a transgenic plant

comprising event MON87712. A representative sample of seed comprising event MON87712 has been deposited according to the Budapest Treaty for the purpose of enabling the present invention. The repository selected for receiving the deposit is the American Type Culture Collection (ATCC) having an address at 10801 University Boulevard, Manassas, Va. USA, Zip Code 20110. The ATCC repository has assigned the accession No. PTA-10296 to event MON87712 containing seed.

The present invention provides a microorganism comprising a DNA molecule having a nucleotide sequence selected from the group consisting of the consecutive nucleotides of SEQ ID NO: 1, the consecutive nucleotides of SEQ ID NO: 2, or at least 51 consecutive nucleotides of SEQ ID NO: 7, or of SEQ ID NO: 8 present in its genome. An example of such a microorganism is a transgenic plant cell. Microorganisms, such as a plant cell of the present invention, are useful in many industrial applications, including but not limited to: (i) use as research tool for scientific inquiry or industrial research; (ii) use in culture for producing endogenous or recombinant carbohydrate, lipid, nucleic acid, enzymes or protein products or small molecules that may be used for subsequent scientific research or as industrial products; and (iii) use with modern plant tissue culture techniques to produce transgenic plants or plant tissue cultures that may then be used for agricultural research or production. The production and use of microorganisms such as transgenic plant cells utilizes modern microbiological techniques and human intervention to produce a man-made, unique microorganism. In this process, recombinant DNA is inserted into a plant cell's genome to create a transgenic plant cell that is separate and unique from naturally occurring plant cells. This transgenic plant cell can then be cultured much like bacteria and yeast cells using modern microbiology techniques and may exist in an undifferentiated, unicellular state. The new plant cell's genetic composition and phenotype is a technical effect created by the integration of the heterologous DNA into the genome of the cell. Another aspect of the present invention is a method of using a microorganism of the present invention. Methods of using microorganisms of the present invention, such as transgenic plant cells, include (i) methods of producing transgenic cells by integrating recombinant DNA into genome of the cell and then using this cell to derive additional cells possessing the same heterologous DNA; (ii) methods of culturing cells that contain recombinant DNA using modern microbiology techniques; (iii) methods of producing and purifying endogenous or recombinant carbohydrate, lipid, nucleic acid, enzymes or protein products from cultured cells; and (iv) methods of using modern plant tissue culture techniques with transgenic plant cells to produce transgenic plants or transgenic plant tissue cultures.

As used herein, "progeny" includes any plant, seed, plant cell, and/or regenerable plant part comprising the event DNA derived from an ancestor plant and/or a polynucleotide having at least one of the sequences provided as the consecutive nucleotides of SEQ ID NO: 1, the consecutive nucleotides of SEQ ID NO: 2, or at least 51 consecutive nucleotides of SEQ ID NO: 7, or at least 51 consecutive nucleotides of SEQ ID NO: 8. Plants, progeny, and seeds may be homozygous or heterozygous for the transgene. Progeny may be grown from seeds produced by a plant comprising event MON87712 and/or from seeds produced by a plant fertilized with pollen from a plant comprising event MON87712.

Progeny plants may be self-pollinated (also known as "selfing") to generate a true breeding line of plants, i.e.,

plants homozygous for the transgene. Alternatively, progeny plants may be outcrossed, e.g., bred with another plant, to produce a varietal or a hybrid seed or plant. The other plant may be transgenic or nontransgenic. A varietal or hybrid seed or plant of the present invention may thus be derived by crossing a first parent that lacks the specific and unique DNA of event MON87712 with a second parent comprising event MON87712, resulting in a hybrid comprising the specific and unique DNA of event MON87712. Each parent can be a hybrid or an inbred/variety, so long as the cross or breeding results in a plant or seed of the present invention, i.e., a seed having at least one allele comprising the specific and unique DNA of event MON87712 and/or the consecutive nucleotides of SEQ ID NO: 1 or SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7 or at least 51 consecutive nucleotides of SEQ ID NO: 8. Two different transgenic plants may thus be mated to produce hybrid offspring that contain two independently segregating, added, exogenous genes. For example, a plant comprising event MON87712 with increased yield can be crossed with another transgenic plant, such as one tolerant to glyphosate, to produce a plant having the characteristics of both transgenic parents. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcox J. ed., American Society of Agronomy, Madison Wis. (1987). Sexually crossing one plant with another plant, i.e., cross-pollinating, may be accomplished or facilitated by human intervention, for example: by human hands collecting the pollen of one plant and contacting this pollen with the style or stigma of a second plant; by human hands and/or human actions removing, destroying, or covering the stamen or anthers of a plant (e.g., by manual intervention or by application of a chemical gametocide) so that natural self-pollination is prevented and cross-pollination would have to take place in order for fertilization to occur; by human placement of pollinating insects in a position for "directed pollination" (e.g., by placing beehives in orchards or fields or by caging plants with pollinating insects); by human opening or removing of parts of the flower to allow for placement or contact of foreign pollen on the style or stigma; by selective placement of plants (e.g., intentionally planting plants in pollinating proximity); and/or by application of chemicals to precipitate flowering or to foster receptivity (of the stigma for pollen).

In practicing this method, the step of sexually crossing one plant with itself, i.e., self-pollinating or selfing, may be accomplished or facilitated by human intervention, for example: by human hands collecting the pollen of the plant and contacting this pollen with the style or stigma of the same plant and then optionally preventing further fertilization of the plant; by human hands and/or actions removing, destroying, or covering the stamen or anthers of other nearby plants (e.g., by detasseling or by application of a chemical gametocide) so that natural cross-pollination is prevented and self-pollination would have to take place in order for fertilization to occur; by human placement of pollinating insects in a position for "directed pollination" (e.g., by caging a plant alone with pollinating insects); by human manipulation of the flower or its parts to allow for self-pollination; by selective placement of plants (e.g., intentionally planting plants beyond pollinating proximity); and/or by

application of chemicals to precipitate flowering or to foster receptivity (of the stigma for pollen).

The present invention provides a plant part that is derived from a plant comprising event MON87712. As used herein, a "plant part" refers to any part of a plant that is comprised of material derived from a plant comprising event MON87712. Plant parts include but are not limited to cells, pollen, ovule, pod, flower, root or stem tissue, fibers, and leaf. Plant parts may be viable, nonviable, regenerable, and/or non-regenerable.

The present invention provides a commodity product that is derived from a plant comprising event MON87712. As used herein, a "commodity product" refers to any composition or product that is comprised of material derived from a plant, seed, plant cell, or plant part comprising event MON87712. Commodity products may be sold to consumers and may be viable or nonviable. Nonviable commodity products include but are not limited to nonviable seeds and grains; processed seeds, seed parts, and plant parts; dehydrated plant tissue, frozen plant tissue, and processed plant tissue; seeds and plant parts processed for animal feed for terrestrial and/or aquatic animal consumption, oil, meal, flour, flakes, bran, fiber, and any other food for human consumption; and biomasses and fuel products. Processed soybeans are the largest source of protein feed and vegetable oil in the world. Soybeans and soybean oils from plants comprising MON87712 can be used in the manufacture of many different products, not limited to, nontoxic plastics, printing inks, lubricants, waxes, hydraulic fluids, electric transformer fluids, solvents, cosmetics, and hair care products. Soybeans and oils of plants comprising MON87712 can be suitable for use in a variety of soy foods made from whole soybeans, such as soymilk, soy nut butter, natto, and tempeh, and soy foods made from processed soybeans and soybean oil, including soybean meal, soy flour, soy protein concentrate, soy protein isolates, texturized soy protein concentrate, hydrolyzed soy protein, whipped topping, cooking oil, salad oil, shortening, and lecithin. Whole soybeans are also edible, and are typically sold to consumers raw, roasted, or as edamame. Soymilk, which is typically produced by soaking and grinding whole soybeans, may be consumed without other processing, spray-dried, or processed to form soy yogurt, soy cheese, tofu, or yuba. Viable commodity products include but are not limited to seeds and plant cells. A plant comprising event MON87712 can thus be used to manufacture any commodity product typically acquired from a soybean plant. Any such commodity product that is derived from the plants comprising event MON87712 may contain at least a detectable amount of the specific and unique DNA corresponding to event MON87712, and specifically may contain a detectable amount of a polynucleotide having a nucleotide sequence of the consecutive nucleotides of SEQ ID NO: 1, the consecutive nucleotides of SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7 or at least 51 consecutive nucleotides of SEQ ID NO: 8. Any standard method of detection for polynucleotide molecules may be used, including methods of detection disclosed herein. A commodity product is within the scope of the present invention if there is any detectable amount of the consecutive nucleotides of SEQ ID NO: 1, the consecutive nucleotides of SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7 or at least 51 consecutive nucleotides of SEQ ID NO: 8 in the commodity product.

The plant, progeny, seed, plant cell, plant part (such as pollen, ovule, pod, flower, root or stem tissue, and leaf), and commodity products of the present invention are therefore

useful for, among other things, growing plants for the purpose of producing seed and/or plant parts comprising event MON87712 for agricultural purposes, producing progeny comprising event MON87712 for plant breeding and research purposes, use with microbiological techniques for industrial and research applications, and sale to consumers.

The present invention provides methods for producing plants with increased yield and plants comprising event MON87712. An event MON87712 containing plant was produced by an *Agrobacterium* mediated transformation method similar to that described in U.S. Pat. No. 5,914,451, using an inbred soybean line with the construct pMON83132 (Table 1). Construct pMON83132 contains a plant expression cassette for expression of the BBX32 protein in soybean plant cells. Transgenic soybean cells were regenerated into intact soybean plants and individual plants were selected from the population of independently transformed transgenic plants that showed desirable molecular characteristics, such as single copy of the transgene cassette at a single locus, the integrity of the transgene cassette, absence of the construct backbone sequence, loss of the unlinked glyphosate resistance selection cassette. Furthermore, inverse PCR and DNA sequence analyses were performed to determine the 5' and 3' insert-to-plant genome junctions, to confirm the organization of the elements within the insert (FIG. 1), and to determine the complete DNA sequence of the insert in soybean event MON87712 (SEQ ID NO: 5). In addition, transgenic plants were screened and selected for increased yield under field conditions. A soybean plant that contains in its genome the BBX32 expression cassette of pMON83132 is an aspect of the present invention.

Increased yield of a transgenic plant of the present invention can be measured in a number of ways, including test weight, seed number per plant, seed weight, seed number per unit area (i.e. seeds, or weight of seeds, per acre), bushels per acre, tons per acre, or kilo per hectare. Expression of the BBX32 protein in plants comprising event MON87712 leads to an increase in yield brought about by the increased capacity for growth and reproductive development. Increased yield results from the partitioning of growth in one or more of the yield parameters including more seeds per plot, larger seeds, more seeds per plant, and/or more plants per plot. Recombinant DNA used in this invention provides increased yield through an increase in source compounds when compared to non-transgenic soybean with the same genetic background. These biochemical and metabolic changes are linked to diurnally-regulated pathways that are impacted by the activity of BBX32 and include carbon metabolism and nitrogen metabolism.

Methods for producing a plant with increased yield comprising transgenic event MON87712 are provided. Transgenic plants used in these methods may be homozygous or heterozygous for the transgene. Progeny plants produced by these methods may be varietal or hybrid plants; may be grown from seeds produced by a plant and/or from seed comprising event MON87712 produced by a plant fertilized with pollen from a plant comprising event MON87712; and may be homozygous or heterozygous for the transgene. Progeny plants may be subsequently self-pollinated to generate a true breeding line of plants, i.e., plants homozygous for the transgene, or alternatively may be outcrossed, e.g., bred with another unrelated plant, to produce a varietal or a hybrid seed or plant. As used herein, the term "zygosity" refers to the similarity of DNA at a specific chromosomal location (locus) in a plant. In the present invention, the DNA specifically refers to the transgene insert along with the

junction sequence (event DNA). A plant is homozygous if the transgene insert with the junction sequence is present at the same location on each chromosome of a chromosome pair (2 alleles). A plant is considered heterozygous if the transgene insert with the junction sequence is present on only one chromosome of a chromosome pair (1 allele). A wild-type plant is null for the event DNA.

A plant with increased yield may be produced by sexually crossing a plant comprising event MON87712 comprising a polynucleotide having the nucleotide sequence of consecutive nucleotides of SEQ ID NO: 1, a polynucleotide having the nucleotide sequence of consecutive nucleotides of SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8 with another plant and thereby producing seed, which is then grown into progeny plants. These progeny plants may be analyzed using diagnostic methods to select for progeny plants that contain event MON87712 DNA or for progeny plants with increased yield. The other plant used may or may not be transgenic. The progeny plant and/or seed produced may be varietal or hybrid seed.

A plant with increased yield may be produced by selfing a plant comprising event MON87712 comprising a polynucleotide having the nucleotide sequence of consecutive nucleotides of SEQ ID NO: 1, a polynucleotide having the nucleotide sequence of consecutive nucleotides of SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8 and thereby producing seed, which is then grown into progeny plants. These progeny plants may then be analyzed using diagnostic methods to select for progeny plants that comprise event MON87712 DNA.

Progeny plants and seeds encompassed by these methods and produced by using these methods are distinct from other plants, for example, because the progeny plants and seeds are recombinant and as such created by human intervention; contain at least one allele that consists of the transgenic DNA of the present invention; and/or contain a detectable amount of a polynucleotide sequence selected from the group consisting of consecutive nucleotides of SEQ ID NO: 1, consecutive nucleotides of SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and at least 51 consecutive nucleotides of SEQ ID NO: 8. A seed may be selected from an individual progeny plant, and so long as the seed comprises SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8, it will be within the scope of the present invention.

The plants, progeny, seeds, plant cells, plant parts (such as pollen, ovule, pod, flower, root or stem tissue, and leaves), and commodity products of the present invention may be evaluated for DNA composition, gene expression, and/or protein expression. Such evaluation may be done by using various methods such as PCR, sequencing, northern blotting, southern analysis, western blotting, immuno-precipitation, and ELISA or by using the methods of detection and/or the detection kits provided herein.

Methods of detecting the presence of compositions specific to event MON87712 in a sample are provided. One method consists of detecting the presence of DNA specific to and derived from a cell, a tissue, a seed, a plant or plant parts comprising event MON87712. The method provides for a template DNA sample to be contacted with a primer pair that is capable of producing an amplicon from event MON87712 DNA upon being subjected to conditions appropriate for amplification, particularly an amplicon that comprises SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive

nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complements thereof. The amplicon is produced from a template DNA molecule derived from event MON87712, so long as the template DNA molecule incorporates the specific and unique nucleotide sequences of SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8. The amplicon may be single or double stranded DNA or RNA, depending on the polymerase selected for use in the production of the amplicon. The method provides for detecting the amplicon molecule produced in any such amplification reaction, and confirming within the sequence of the amplicon the presence of the nucleotides corresponding to SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complements thereof. The detection of the nucleotides corresponding to SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, and/or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complements thereof in the amplicon are determinative and/or diagnostic for the presence of event MON87712 specific DNA and thus biological material comprising event MON87712 in the sample.

Another method is provided for detecting the presence of a DNA molecule corresponding to SEQ ID NO: 3 or SEQ ID NO: 4 in a sample consisting of material derived from plant or plant tissue. The method consists of (i) obtaining a DNA sample from a plant, or from a group of different plants, (ii) contacting the DNA sample with a DNA probe molecule comprising the nucleotides as set forth in either SEQ ID NO: 1 or SEQ ID NO: 2, (iii) allowing the probe and the DNA sample to hybridize under stringent hybridization conditions, and then (iv) detecting a hybridization event between the probe and the target DNA sample. Detection of the hybrid composition is diagnostic for the presence of SEQ ID NO: 3 or SEQ ID NO: 4, as the case may be, in the DNA sample. Absence of hybridization is alternatively diagnostic of the absence of the transgenic event in the sample if the appropriate positive controls are run concurrently. Alternatively, determining that a particular plant contains either or both of the sequences corresponding to SEQ ID NO: 1 or SEQ ID NO: 2, or the complements thereof, is determinative that the plant contains at least one allele corresponding to event MON87712.

It is thus possible to detect the presence of a nucleic acid molecule of the present invention by any well known nucleic acid amplification and detection methods such as polymerase chain reaction (PCR), recombinase polymerase amplification (RPA), or DNA hybridization using nucleic acid probes. An event-specific PCR assay is discussed, for example, by Taverniers et al. (J. Agric. Food Chem., 53: 3041-3052, 2005) in which an event-specific tracing system for transgenic maize lines Bt11, Bt176, and GA21 and for transgenic event RT73 is demonstrated. In this study, event-specific primers and probes were designed based upon the sequences of the genome/transgene junctions for each event. Transgenic plant event specific DNA detection methods have also been described in U.S. Pat. Nos. 6,893,826; 6,825,400; 6,740,488; 6,733,974; 6,689,880; 6,900,014 and 6,818,807.

DNA detection kits are provided. One type of kit contains at least one DNA molecule of sufficient length of contiguous nucleotides of SEQ ID NO: 3, SEQ ID NO: 5, or SEQ ID NO: 6 to function as a DNA primer or probe specific for detecting the presence of DNA derived from transgenic event MON87712 in a sample. The DNA molecule being

detected with the kit comprises contiguous nucleotides of the sequence as set forth in SEQ ID NO: 1 or at least 51 consecutive nucleotides of SEQ ID NO: 7. Alternatively, the kit may contain at least one DNA molecule of sufficient length of contiguous nucleotides of SEQ ID NO: 4, SEQ ID NO: 5, or SEQ ID NO: 6 to function as a DNA primer or probe specific for detecting the presence of DNA derived from transgenic event MON87712 in a sample. The DNA molecule being detected with the kit comprises contiguous nucleotides as set forth in SEQ ID NO: 2 or at least 51 consecutive nucleotides of SEQ ID NO: 8.

An alternative kit employs a method in which the target DNA sample is contacted with a primer pair as described above, then performing a nucleic acid amplification reaction sufficient to produce an amplicon comprising the consecutive nucleotides of SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, or at least 51 consecutive nucleotides of SEQ ID NO: 8. Detection of the amplicon and determining the presence of the consecutive nucleotides of SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complements thereof within the sequence of the amplicon is diagnostic for the presence of event MON87712 specific DNA in a DNA sample.

A DNA molecule sufficient for use as a DNA probe is provided that is useful for determining, detecting, or for diagnosing the presence or even the absence of DNA specific and unique to event MON87712 DNA in a sample. The DNA molecule contains the consecutive nucleotides of SEQ ID NO: 1, or the complement thereof, the consecutive nucleotides of SEQ ID NO: 2, or the complement thereof, at least 51 consecutive nucleotides of SEQ ID NO: 7, or the complement thereof, or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complement thereof.

Nucleic acid amplification can be accomplished by any of the various nucleic acid amplification methods known in the art, including thermal and isothermal amplification methods. The sequence of the heterologous DNA insert, junction sequences, or flanking sequences from event MON87712 (with representative seed samples comprising event MON87712 deposited as ATCC PTA-10296) can be verified by amplifying such sequences from the event using primers derived from the sequences provided herein followed by standard DNA sequencing of the amplicon or of the cloned DNA.

The amplicon produced by these methods may be detected by a plurality of techniques. One such method is Genetic Bit Analysis (Nikiforov, et al. *Nucleic Acid Res.* 22:4167-4175, 1994) where a DNA oligonucleotide is designed which overlaps both the adjacent flanking genomic DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following thermal amplification of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking genomic sequence), a single-stranded amplicon can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labelled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. Detection of a fluorescent or other signal indicates the presence of the insert/flanking sequence due to successful amplification, hybridization, and single base extension.

Another method is the Pyrosequencing technique as described by Winge (*Innov. Pharma. Tech.* 00:18-24, 2000). In this method an oligonucleotide is designed that overlaps

the adjacent genomic DNA and insert DNA junction. The oligonucleotide is hybridized to a single-stranded amplicon from the region of interest (one primer in the inserted sequence and one in the flanking genomic sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. ddNTPs are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

Fluorescence Polarization as described by Chen, et al., (*Genome Res.* 9:492-498, 1999) is a method that can be used to detect the amplicon. Using this method an oligonucleotide is designed which overlaps the genomic flanking and inserted DNA junction. The oligonucleotide is hybridized to single-stranded amplicon from the region of interest (one primer in the inserted DNA and one in the flanking genomic DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single base extension.

TAQMAN® (PE Applied Biosystems, Foster City, Calif.) may also be used to detect and/or quantifying the presence of a DNA sequence using the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the genomic flanking and insert DNA junction. The FRET probe and amplification primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

Molecular Beacons have been described for use in sequence detection as described in Tyangi, et al. (*Nature Biotech.* 14:303-308, 1996). Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking genomic and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and amplification primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties, which leads to the production of a fluorescent signal. The fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

Other described methods, such as, microfluidics (US Patent Publication No. 2006068398, U.S. Pat. No. 6,544,734) provide methods and devices to separate and amplify DNA samples. Optical dyes are used to detect and measure specific DNA molecules (WO/05017181). Nanotube devices (WO/06024023) that comprise an electronic sensor for the detection of DNA molecules or nanobeads that bind specific DNA molecules and can then be detected.

DNA detection kits can be developed using the compositions disclosed herein and the methods well known in the

art of DNA detection. The kits are useful for the identification of event MON87712 in a sample and can be applied to methods for breeding plants containing the appropriate event DNA. The kits may contain DNA primers or probes that are similar or complementary to SEQ ID NO: 1-6, or fragments or complements thereof.

The kits and detection methods of the present invention are therefore useful for, among other things, identifying event MON87712, selecting plant varieties or hybrids comprising event MON87712, detecting the presence of DNA derived from event MON87712 in a sample, and monitoring samples for the presence and/or absence of event MON87712 or plants, plant parts or commodity products comprising event MON87712.

The following examples are included to demonstrate examples of certain embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLES

Example 1: Transformation of Soybean with pMON83132 and Event Selection

This example describes transformation and generation of transgenic soybean events, and selection of event MON87712.

An *Agrobacterium*-mediated transformation method was used to transform soybean cells with a binary construct designated as pMON83132. The construct contains two plant transformation cassettes or T-DNAs. Each cassette is flanked by right border and left border sequences. The transgenic insert as set forth in SEQ ID NO: 5 comprises a transformation cassette containing an enhanced 35S promoter from cauliflower mosaic virus (CaMV), operably

linked to a DNA molecule encoding BBX32 protein, operably linked to the 3' untranslated region from fiber protein E6 gene of *Gossypium barbadense* (cotton). The second transformation cassette contains a chimeric promoter consisting of enhancer sequences from the promoter of the figwort mosaic virus (FMV) 35S RNA combined with the promoter of the EF-1 α gene from *Arabidopsis thaliana* that encodes elongation factor EF-1 alpha. It also contains a leader (exon 1) and intron with flanking splice sites of the *Arabidopsis* EF-1 α gene, operably linked to a DNA molecule encoding a chloroplast transit peptide (CTP2) from *Arabidopsis* EPSP synthase, fused to a codon modified coding sequence of the *aroA* gene from the *Agrobacterium* sp. strain CP4 encoding the CP4 EPSPS protein, operably linked to a 3' untranslated region of the *RbcS2* gene from *Pisum sativum*. The CP4 *aroA* gene confers tolerance to glyphosate, and was used as a selectable marker. Table 1 is a summary of the genetic elements in pMON83132.

After transformation with construct pMON83132, transformed cells were allowed to grow and multiply on media containing glyphosate. Plants were regenerated from surviving cells. A total of 4944 independent R0 transformation events were produced and characterized by detailed molecular analyses. Events were screened by quantitative PCR and RT-PCR (TAQMAN[®]) for insert number (number of integration sites within the soybean genome), copy number (the number of copies of the T-DNA within one locus), the integrity of the inserted cassette and the absence of backbone sequence, and expression of the AtBBX32 transgene transcript. Events with undesirable molecular characteristics, such as presence of multiple copies of the transgene and/or molecular complexity of the insert, the presence of the transformation vector backbone sequence were eliminated. Furthermore, linkage Southern analysis was done to remove events where the BBX32 expression cassette was linked to the CP4 selectable marker cassette. A total of 284 events met the stringent molecular selection criteria based on the analyses described above. Three additional events were further eliminated due to unavailability of homozygous seed stocks, resulting in 281 events being advanced to R1 generation.

TABLE 1

Summary of the genetic elements in pMON83132.		
Genetic Element	Location in Construct pMON83132	Function (Reference)
T-DNA I		
B ¹ -Left Border	1-442	DNA region from <i>Agrobacterium tumefaciens</i> containing the left border sequence used for transfer of the T-DNA (Barker <i>et al.</i> , Plant Mol. Biol. 2: 335-350, 1983).
P ² -e35S	511-1123	Promoter for the cauliflower mosaic virus (CaMV) 35S RNA (Odell <i>et al.</i> , Nature 313: 810-812, 1985) containing the duplicated enhancer region (Kay <i>et al.</i> , Science 236: 1299-1302, 1987) that directs transcription in plant cells.
CS ³ -BBX32	1148-1825	Coding sequence of the BBX32 gene from <i>Arabidopsis thaliana</i> encoding a zinc finger protein (B-box type) (Khanna <i>et al.</i> , Plant Cell 21: 3416-3420, 2009) which modulates aspects of diurnal biology (Holtan <i>et al.</i> , submitted).
T ⁴ -E6	1840-2154	3' UTR region of the E6 gene of <i>Gossypium barbadense</i> (cotton) that encodes a fiber protein involved in early fiber development (John, Plant Mol Biol. 30: 297-306, 1996) which functions to direct polyadenylation of mRNA.

TABLE 1-continued

Summary of the genetic elements in pMON83132.		
Genetic Element	Location in Construct pMON83132	Function (Reference)
B-Right Border	2193-2549	DNA region from <i>Agrobacterium tumefaciens</i> containing the right border sequence used for transfer of the T-DNA (Depicker <i>et al.</i> , J. of Mol. and Applied Genetics 1: 561-573, 1982; Zambryski <i>et al.</i> , J. of Mol. and Applied Genetics 1: 361-370, 1982). T-DNA II
B-Right Border	2721-3077	DNA region from <i>Agrobacterium tumefaciens</i> containing the right border sequence used for transfer of the T-DNA (Depicker <i>et al.</i> , J. of Mol. and Applied Genetics 1: 561-573, 1982; Zambryski <i>et al.</i> , J. of Mol. and Applied Genetics 1: 361-370, 1982).
P-FMV/EF-1 α	3100-4139	Chimeric promoter consisting of enhancer sequences from the promoter of the Figwort Mosaic virus (FMV) 35S RNA (Richins <i>et al.</i> , Nucleic Acids Research 15: 8451-8466, 1987) combined with the promoter from the EF-1 α gene of <i>Arabidopsis thaliana</i> that encodes elongation factor EF-1 alpha (Axelos <i>et al.</i> , Molecular and General Genetics 219: 106-112, 1989). It is associated with constitutive expression of the gene.
L ⁵ -EF-1 α	4140-4185	Leader (exon 1) of the EF-1 α gene from <i>Arabidopsis thaliana</i> that encodes elongation factor EF-1 alpha (Axelos <i>et al.</i> , Molecular and General Genetics 219: 106-112, 1989), which enhances gene expression.
I ⁶ -EF-1 α	4186-4807	Intron with flanking splice sites of the EF-1 α gene from <i>Arabidopsis thaliana</i> that encodes elongation factor EF-1 alpha (Axelos <i>et al.</i> , Molecular and General Genetics 219: 106-112, 1989), that enhances gene expression.
TS ⁷ -CTP2	4817-5044	Targeting sequence of the <i>ShkG</i> gene from <i>Arabidopsis thaliana</i> encoding EPSPS containing the transit peptide region) that directs transport of the EPSPS protein to the chloroplast (Klee <i>et al.</i> , Molecular and General Genetics 210: 437-442, 1987).
CS-modified cp4 <i>epsps</i>	5045-6412	Codon modified coding sequence of the <i>aroA</i> gene from the <i>Agrobacterium</i> sp. strain CP4 encoding the CP4 EPSPS protein (Padgett <i>et al.</i> , 1996, in <i>Herbicide-Resistant Crops: Agricultural, Economic, Environmental, Regulatory, and Technological Aspects</i> , S. O. Duke, (ed.), Pages 53-84, CRC Press, Boca Raton, FL).
T-E9	6419-7061	3' nontranslated region of the <i>RbcS2</i> gene from <i>Pisum sativum</i> (pea) encoding the Rubisco small subunit, which functions to direct polyadenylation of the mRNA (Coruzzi <i>et al.</i> , EMBO J. 3: 1671-1679, 1984).
B-Left Border	7076-7486	DNA region from <i>Agrobacterium tumefaciens</i> containing the left border sequence used for transfer of the T-DNA (Barker <i>et al.</i> , Plant Mol. Biol. 2: 335-350, 1983). Vector Backbone
<i>aadA</i>	7582-8470	Bacterial promoter, coding sequence, and 3' UTR for an aminoglycoside-modifying enzyme, 3'-(9)-O-nucleotidyltransferase from the transposon Tn7 (Fling <i>et al.</i> , Nucleic Acids Research 13: 7095-7106, 1985) that confers spectinomycin and streptomycin resistance.
OR ⁸ -ori-pBR322	9001-9589	Origin of replication from pBR322 for maintenance of plasmid in <i>E. coli</i> (Sutcliffe, 1979, Complete nucleotide sequence of the <i>Escherichia coli</i> plasmid pBR322. Pages 77-90 in <i>Cold Spring Harbor Symposia on Quantitative Biology</i> , Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press).
CS- <i>rop</i>	10017-10208	Coding sequence for repressor of primer protein derived from the <i>ColE1</i> plasmid for maintenance of plasmid copy number in <i>E. coli</i> (Giza and Huang, Gene 78: 73-84, 1989).

TABLE 1-continued

Summary of the genetic elements in pMON83132.		
Genetic Element	Location in Construct	pMON83132 Function (Reference)
OR-ori V	10946-11342	Origin of replication from the broad host range plasmid RK2 for maintenance of plasmid in <i>Agrobacterium</i> (Stalker <i>et al.</i> , Molecular and General Genetics 181: 8-12, 1981).

- ¹B, Border
- ²P, Promoter
- ³CS, Coding Sequence
- ⁴T, Transcription Termination Sequence
- ⁵L, Leader
- ⁶I, Intron
- ⁷TS, Targeting Sequence
- ⁸OR, Origin of Replication

At the R1 generation, events underwent segregation analysis to select only homozygous and heterozygous plants going forward. In addition, CP4 linkage analysis, TAQ-MAN® analysis and BBX32 expression analysis were performed to remove events where CP4 was linked to the BBX32 transgene, or where the sequence of the OriV origin of replication (backbone) was detected, or where BBX32 transgene expression level was undesirable, resulting in 121 events remaining. Southern hybridization analysis using different probes further eliminated 42 events that were not single copy, or single insert or not backbone-free. Probes included the intact BBX32 coding region and its respective promoter, the polyadenylation sequence and the plasmid backbone. An additional 2 events were stopped due to lack of sufficient seeds to advance, leaving 77 events to carry forward to the R2 generation.

At the R2 generation, an additional 9 events were dropped due to 1) lack of sufficient seed for the R2 nursery (1 event), or 2) undesirable agronomic phenotypes (2 events), or 3) BBX32 expression level or the absence of the representative homozygous lines as determined by Invader® (Third Wave Technologies, Inc., Madison, Wis.) (6 events). As a result, 68 events were carried forward for year 1 yield field test in the US. Based upon yield performance, 16 events were advanced to the second season field testing in South America, from which, 9 events were selected for further yield testing. Meta-analysis of yield data from 4 seasons of field trials in the United States or South America, each with multiple locations (Table 2), showed that one progeny line designated as comprising event MON87712 was the highest yielding event (FIG. 2).

TABLE 2

Details of the four season field testing for yield.		
Year	Geography	# Locations
2007	United States	18
2007-2008	South America	14
2008	United States	24
2008-2009	South America	14

Example 2: Isolation of Flanking Sequences Using Inverse PCR And Identification of Flanking Sequences by Sequencing

This example describes isolation of the soybean genomic DNA sequences flanking the transgenic DNA insert using

inverse PCR for event MON87712, and identification of the flanking genomic sequences by sequencing.

Sequences flanking the T-DNA insertion in event MON87712 were determined using inverse PCR as described in Ochman *et al.*, 1990 (PCR Protocols: A guide to Methods and Applications, Academic Press, Inc.). Plant genomic DNA was isolated from both wild-type A3525 and the transgenic line from tissue grown under greenhouse conditions. Frozen leaf tissue was ground with a mortar and a pestle in liquid nitrogen or by mechanical grinding, followed by DNA extraction using methods known in the art. This method can be modified by one skilled in the art to extract DNA from any tissue, including, but not limited to seed.

An aliquot of DNA from each sample was digested with restriction endonucleases selected based on restriction analysis of the transgenic DNA. After self-ligation of the restriction fragments, PCR amplification was performed using primers designed from the transgenic sequence that would amplify sequences extending away from the 5' and 3' ends of the transgenic DNA. A variety of Taq polymerases and amplification systems may be used. Table 3 shows an example of PCR amplification for flanking sequence isolation using Phusion High Fidelity DNA Polymerase (Cat. No. F531S or F531L, New England Biolabs), and Thermalcyclers Applied Biosystems GeneAmp 9700, ABI 9800 Fast Thermal Cycler and MJ Opticon.

TABLE 3

An example of inverse PCR amplification for flanking sequence isolation.		
	Volume	Component
PCR master mix (per reaction)	2.9 µl	Water
	0.05 µl	Primer 1 (100 µM)
	0.05 µl	Primer 1 (100 µM)
	5.0 µl	2X Phusion Taq
	2.0 µl	ligated DNA
	10 µl	Total
	Step	Condition
DNA amplification in a fast thermocycler	1	98° C. 30 sec
	2	98° C. 5 sec
	3	60° C. 10 sec

TABLE 3-continued

An example of inverse PCR amplification for flanking sequence isolation.		
4	72° C.	2 min
5	Go to step 2	30 times
6	72° C.	4 min
7	10° C.	forever
8	End	

PCR products were separated by agarose gel electrophoresis and purified using a QIAGEN gel purification kit (Qiagen, Valencia, Calif.). The subsequent products were sequenced directly using standard sequencing protocols. Using these two methods, the 5' flanking sequence, which extends into the left border sequence of the integrated DNA insert including the BBX32 expression cassette, was identified and is presented as SEQ ID NO: 3 ([C] of FIG. 1). The 3' flanking sequence, which extends into the right border sequence of the integrated DNA insert including the BBX32 expression cassette, was identified and is presented as SEQ ID NO: 4 ([D] of FIG. 1). The transgenic DNA integrated into the soybean genomic DNA is presented as SEQ ID NO: 5 ([E] of FIG. 1).

The isolated sequences were compared to the T-DNA sequence to identify the flanking sequences and the co-isolated T-DNA fragments. Confirmation of the presence of the expression cassette was achieved by PCR with primers designed based upon the deduced flanking sequence data and the known T-DNA sequence. The wild type sequence corresponding to the same region in which the T-DNA was integrated in the transformed line was isolated using primers designed from the flanking sequences in MON87712. The flanking sequences in MON87712 and the wild type sequence were analyzed against multiple nucleotide and protein databases. This information was used to examine the relationship of the transgene to the plant genome and to look

at the insertion site integrity. The flanking sequence and wild type sequences were used to design primers for TAQMAN® endpoint assays used to identify the events and determine zygosity as described in Example 3.

Example 3: Event-Specific Endpoint TAQMAN® And Zygosity Assays

This example describes an event-specific endpoint TAQMAN® thermal amplification method for identification of event MON87712 DNA in a sample.

Examples of conditions useful with the event MON87712-specific endpoint TAQMAN® method are described in Table 4 and Table 5. The DNA primers used in the endpoint assay are primers SQ3983 (SEQ ID NO: 16) and SQ22982 (SEQ ID NO: 17) and the 6-FAM™ labeled oligonucleotide probe is PB10453 (SEQ ID NO: 18). 6FAM™ is a fluorescent dye product of Applied Biosystems (Foster City, Calif.) attached to the DNA probe. For TAQMAN® MGB (Minor Groove Binding) probes, the 5' exonuclease activity of Taq DNA polymerase cleaves the probe from the 5'-end, between the fluorophore and quencher. When hybridized to the target DNA strand, quencher and fluorophore are separated enough to produce a fluorescent signal.

Primers SQ3983 (SEQ ID NO: 16) and SQ22982 (SEQ ID NO: 17), when used as described with probe PB10453 (SEQ ID NO: 18), produce an amplicon that is diagnostic for event MON87712 DNA. The analysis includes a positive control from soybean known to contain event MON87712 DNA, a negative control from non-transgenic soybean and a negative control that contains no template DNA.

These assays are optimized for use with Applied Biosystems GeneAmp PCR System 9700, ABI 9800 Fast Thermal Cycler and MJ Opticon. Other methods and apparatus known to those skilled in the art may be used to produce amplicons that identify the event MON87712 DNA.

TABLE 4

Soybean MON87712 Event-Specific Endpoint TAQMAN® PCR Conditions			
Step	Reagent	Volume	Comments
1	18 megohm water	adjust for final volume of 10 µl	
2	2X Universal Master Mix (dNTPs, enzyme and buffer)	5.0 µl	1X final concentration of dNTPs, enzyme and buffer
3	Event Primers SQ3983 and SQ22982 Mix (resuspended in 18 megohm water to a concentration of 20 µM for each primer) Example: In a microcentrifuge tube, the following are added to achieve 500 µl at a final concentration of 20 µM: 100 µl of Primer SQ22982 at a concentration of 100 µM 100 µl of Primer SQ3983 at a concentration of 100 µM 300 µl of 18 megohm water	0.5 µl	1.0 µM final concentration
4	Event 6-FAM™ MGB Probe PB10453 (resuspended in 18 megohm water to a concentration of 10 µM)	0.2 µl	0.2 µM final concentration
5	Internal control Primer-1 and internal control Primer-2 Mix (resuspended in 18 megohm	0.5 µl	1 µM final concentration

TABLE 4-continued

Soybean MON87712 Event-Specific Endpoint TAQMAN® PCR Conditions			
Step	Reagent	Volume	Comments
	water to a concentration of 20 µM for each primer)		
6	Internal control VIC™ probe (resuspended in 18 megohm water to a concentration of 10 µM)	0.2 µl	0.2 µM final concentration
7	Extracted DNA (template): 1. Leaf or seed samples to be analyzed 2. Negative control (non-transgenic DNA) 3. Negative water control (no template control) 4. Positive control (MON87712DNA)	3.0 µl	

TABLE 5

Endpoint TAQMAN® thermocycler conditions		
Cycle No.	Settings	
1	50° C.	2 minutes
1	95° C.	10 minutes
10	95° C.	15 seconds
	64° C.	1 minute (-1° C./cycle)
30	95° C.	15 seconds
	54° C.	1 minute
1	10° C.	Forever

The following example describes an event-specific endpoint TAQMAN® thermal amplification method developed to determine the zygosity of event MON87712 in a sample. A zygosity assay is useful for determining if a plant comprising an event is homozygous for the event DNA, that is comprising the exogenous DNA at the same location on each chromosome of a chromosomal pair; or heterozygous for an event DNA, that is comprising the exogenous DNA on only one chromosome of a chromosomal pair; or is null for the event DNA, that is wild type. A set of three primers (SEQ ID NO: 10, 13 and 11 or 15), a 6FAM™ labeled probe (SEQ ID NO: 12) and a VIC™ labeled probe (SEQ ID NO: 14) were used in the assays (Table 8). 6FAM™ and VIC™ are fluorescent dye products of Applied Biosystems (Foster City, Calif.) attached to the DNA probes. Primer SEQ ID NO: 11 or 15 hybridizes and extends specifically from the inserted exogenous DNA; primer SEQ ID NO: 10 hybridizes and extends specifically from the DNA flanking the 3' side of the inserted exogenous DNA; primer SEQ ID NO: 13 hybridizes and extends specifically from the wild-type DNA corresponding to a region that was deleted during T-DNA insertion in the transgenic event MON87712. These primers are diagnostic. In this example, primers SEQ ID NO: 10 and SEQ ID NO: 11 or 15 and the 6FAM™-labeled oligonucleotide probe SEQ ID NO: 12 produce a DNA amplicon revealed by the liberation of a fluorescent signal from probe SEQ ID NO: 12, which is diagnostic for event MON87712 DNA, indicating a copy of the inserted transgenic DNA present in the genomic DNA. Primers SEQ ID NO: 10 and SEQ ID NO: 13 and the VIC™-labeled oligonucleotide probe SEQ ID NO: 14 produce an amplicon revealed by the liberation of a fluorescent signal from probe SEQ ID NO: 14, which is diagnostic for the wild type allele, indicating no copy of the inserted exogenous DNA present in the genomic DNA. When the three primers and two probes are mixed

together in a PCR reaction with DNA extracted from a plant, release of a fluorescent signal only from the 6FAM™-labeled oligonucleotide probe (SEQ ID NO: 12) is indicative of and diagnostic of a plant homozygous for event MON87712. When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant, release of a fluorescent signal from both the 6FAM™-labeled oligonucleotide probe SEQ ID NO: 12 and the VIC™-labeled oligonucleotide probe SEQ ID NO: 14 is indicative of and diagnostic of a plant heterozygous for event MON87712. When the three primers and two probes are mixed together in a PCR reaction with DNA extracted from a plant, release of a fluorescent signal from only the VIC™-labeled oligonucleotide probe SEQ ID NO: 14 is indicative of and diagnostic of a plant null for event MON87712, i.e. wild type. The assays also include a positive control from soybean containing event MON87712 DNA, a negative control from non-transgenic soybean and a negative control that contains no template DNA.

Table 6 and Table 7 provide examples of conditions used for zygosity assay for event MON87712 in a sample. These assays are optimized for use with an Applied Biosystems GeneAmp PCR System 9700. The allelic detection was optimized with a 7900HT Sequence Detection System by Applied Biosystems, or a PHERAstar by BMG Labtech. Other methods and apparatus known to those skilled in the art that produce amplicons that identify the event MON87712 DNA is within the skill of the art.

TABLE 6

Soybean MON87712 Event-Specific Zygosity Endpoint TAQMAN® PCR		
Step	Details	
1	Prepare the following:	
	200 µM Primers	
	100 µM Event 6-FAM™ MGB* probe	
	100 µM Wild type VIC™ MGB probe	
	50:50 2X ABI universal master mix:2X AbGene universal master mix	
	0.1X TE	
2	Prepare the following 80X Primer + probe stock:	
	100 µM Event 6-FAM™ MGB Probe	16 µl
	100 µM Wild type VIC™ MGB probe	16 µl
	200 µM Wild type forward primer	8.5 µl
	200 µM Event forward primer	8.5 µl
	200 µM reverse primer	17 µl

TABLE 6-continued

Soybean MON87712 Event-Specific Zygosity Endpoint TAQMAN® PCR			
Step	Details		
3	Dilute the 80X Primer + Probe stock to 10X working dilution with 0.1X TE		
4	Prepare the following reaction solution (per reaction):		
	50:50 2X ABI universal master mix:2X AbGene universal master mix	3 µl	
	10X Primer + Probe working dilution	0.6 µl	
	Extracted DNA (template):	2.4 µl	
	1) Samples to be analyzed		
	2) Negative control (non-transgenic DNA)		
	3) Negative water control (no template control)		
	4) Positive control homozygous MON87712 DNA		
	5) Positive control heterozygous MON87712 DNA		

*MGB: Minor Groove Binding

TABLE 7

Zygosity Endpoint TAQMAN® Thermocycler Conditions			
Cycle No.	Settings		
1	95° C.	10 minutes	
40	92° C.	15 seconds	
	64° C.	1 Minute	
1	10° C.	Forever	

TABLE 8

Examples of Primer And Probe Combinations Used for Zygosity Assays					
Combination	Type	Direction	SEQ ID NO	Sequences	
1	Primers	Reverse	10	GTTTTACAATTACCTCG TTTAAGTAAATCA	
		Forward	13	CTATTATTTGCTATAAG TATTTGATGTAAGAA	
	Probe	Wild type allele		14	VIC- TTGTATTAATAACAAA AAATTG
				10	GTTTTACAATTACCTCG TTTAAGTAAATCA
	Primers	Reverse	10	GTTTTACAATTACCTCG TTTAAGTAAATCA	
		Forward	11	CATTCTCGAGCAGGAC CTGCAGAA	
Probe	MON87712 allele		12	6FAM- AACACTGATAGTTTAA ACTGAAG	
2	Primers	Reverse	10	GTTTTACAATTACCTCG TTTAAGTAAATCA	
		Forward	13	CTATTATTTGCTATAAG TATTTGATGTAAGAA	
	Probe	Wild type allele		14	VIC- TTGTATTAATAACAAA AAATTG
				10	GTTTTACAATTACCTCG TTTAAGTAAATCA
	Primers	Reverse	10	GTTTTACAATTACCTCG TTTAAGTAAATCA	
		Forward	15	GTACATTCTCGAGCAG GACCTGCA	
Probe	MON87712 allele		12	6FAM- AACACTGATAGTTTAA ACTGAAG	

Example 4: Identification of Event MON87712 in any MON87712 Containing Breeding Event

This example describes how one may identify the MON87712 event within progeny of any breeding event containing MON87712 soybean.

Event DNA primer pairs are used to produce an amplicon diagnostic for soybean event MON87712. An amplicon diagnostic for MON87712 comprises at least one junction sequence, provided herein as SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complements thereof ([A] or [B], respectively as illustrated in FIG. 1). SEQ ID NO: 1 ([A] of FIG. 1) is a nucleotide sequence corresponding to the junction of the 5' flanking sequence (positions 3495 through 3516 of SEQ ID NO: 3 [C], see FIG. 1) and the left border of the integrated DNA insert (positions 1 through 11 of SEQ ID NO: 5 [E], see FIG. 1). SEQ ID NO: 2 ([B] of FIG. 1) is a nucleotide sequence corresponding to the junction of the right border of the integrated DNA insert (positions 2004 through 2014 of SEQ ID NO: 5 [E], see FIG. 1) and the 3' flanking sequence (positions 90 through 111 of SEQ ID NO: 4 [D], see FIG. 1). SEQ ID NO: 7 ([A] of FIG. 1) is a nucleotide sequence corresponding to the junction of the 5' flanking sequence (positions 3456 through 3555 of SEQ ID NO: 3 [C], see FIG. 1) and the left border of the integrated DNA insert (positions 1 through 50 of SEQ ID NO: 5 [E], see FIG. 1). SEQ ID NO: 8 ([B] of FIG. 1) is a nucleotide sequence corresponding to the junction of the right border of the integrated DNA insert (positions 1965 through 2014 of SEQ ID NO: 5 [E], see FIG. 1) and the 3' flanking sequence (positions 51 through 150 of SEQ ID NO: 4 [D], see FIG. 1).

Event primer pairs that produce an amplicon diagnostic for MON87712 include primer pairs designed using the flanking sequences (SEQ ID NO: 3 and SEQ ID NO: 4) and the integrated transgenic DNA sequence (SEQ ID NO: 5). To acquire a diagnostic amplicon in which at least 15 nucleotides of SEQ ID NO: 1 is found, one would design a forward primer based upon SEQ ID NO: 3 from bases 1 through 3494 and a reverse primer based upon the inserted expression cassette DNA sequence, SEQ ID NO: 5 from positions 12 through 2014, in which the primers are of sufficient length of contiguous nucleotides to specifically hybridize to SEQ ID NO: 3 and SEQ ID NO: 5. To acquire a diagnostic amplicon in which at least 15 nucleotides of SEQ ID NO: 2 is found, one would design a forward primer based upon the inserted expression cassette DNA sequence, SEQ ID NO: 5, from positions 1 through 2003 and a reverse primer based upon the 3' flanking sequence, SEQ ID NO: 4, from bases 12 through 2065, in which the primers are of sufficient length of contiguous nucleotides to specifically hybridize to SEQ ID NO: 4 and SEQ ID NO: 5. For practical purposes, one should design primers that produce amplicons of a limited size range, for example, between 100 to 1000 bases. Smaller sized (shorter nucleotide length) amplicons in general may be more reliably produced in PCR reactions, allow for shorter cycle times and be easily separated and visualized on agarose gels or adapted for use in endpoint TAQMAN®-like assays. In addition, amplicons produced using primer pairs can be cloned into vectors, propagated, isolated and sequenced or can be sequenced directly with methods well established in the art. Any primer pair derived from the combination of SEQ ID NO: 3 and SEQ ID NO: 5 or the combination of SEQ ID NO: 4 and SEQ ID NO: 5 that are useful in a DNA amplification method to produce an amplicon diagnostic for MON87712, plants comprising MON87712 or progeny thereof is an aspect of the present invention. Any single isolated DNA primer molecule comprising at least 11 contiguous nucleotides of SEQ ID NO: 3, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for MON87712, plants comprising MON87712 or progeny thereof is an

35

aspect of the present invention. Any single isolated DNA primer molecule comprising at least 11 contiguous nucleotides of SEQ ID NO: 4, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for MON87712, plants comprising MON87712 or progeny thereof is an aspect of the present invention. Any single isolated DNA primer molecule comprising at least 11 contiguous nucleotides of SEQ ID NO: 5, or its complement that is useful in a DNA amplification method to produce an amplicon diagnostic for MON87712, plants comprising MON87712 or progeny thereof is an aspect of the present invention.

An example of the amplification conditions for this analysis is illustrated in Table 4 and Table 5 (Example 3). However, any modification of these methods or the use of DNA primers homologous or complementary to SEQ ID NO: 3, SEQ ID NO: 4 or DNA sequences of the transgenic insert (SEQ ID NO: 5) of event MON87712 that produce an amplicon diagnostic for MON87712 is within the scope of the present disclosure. A diagnostic amplicon comprises a DNA molecule homologous or complementary to at least one transgene/genomic junction DNA (SEQ ID NO: 1, SEQ ID NO: 2, at least 51 consecutive nucleotides of SEQ ID NO: 7, or at least 51 consecutive nucleotides of SEQ ID NO: 8, or the complements thereof), or a substantial portion thereof.

An analysis for event MON87712 DNA in a sample should include a positive control from event MON87712, a negative control from a soybean plant that does not contain event MON87712, for example, but not limited to wild type control, and a negative control that contains no soybean genomic DNA. A primer pair that will amplify an endogenous soybean DNA molecule will serve as an internal control for the DNA amplification conditions. Additional primer sequences can be selected from SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5 by those skilled in the art of DNA amplification methods, and conditions selected for the production of an amplicon by the methods shown in Table 4 and Table 5 may differ, but result in an amplicon diagnostic

36

for event MON87712 DNA. The use of these DNA primer sequences with modifications to the methods of Table 4 and Table 5 are within the scope of the invention. The amplicon produced by at least one DNA primer sequence derived from SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5 that is diagnostic for MON87712 is an aspect of the invention.

DNA detection kits that contain at least one DNA primer derived from SEQ ID NO: 3, SEQ ID NO: 4, or SEQ ID NO: 5, that when used in a DNA amplification method, produces a diagnostic amplicon for MON87712, plants comprising MON87712 or progeny thereof is an aspect of the invention. A soybean plant or seed, wherein its genomic DNA produces an amplicon diagnostic for MON87712 when tested in a DNA amplification method is an aspect of the invention. The assay for the MON87712 amplicon can be performed by using an Applied Biosystems GeneAmp PCR System 9700, ABI 9800 Fast Thermal Cycler and MJ Opticon, Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700 or Eppendorf Mastercycler Gradient thermocycler or any other amplification system that can be used to produce an amplicon diagnostic of MON87712.

A deposit of the soybean event MON87712 comprising seed disclosed above and recited in the claims, has been made under the Budapest Treaty with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110. The date of deposit is Aug. 20, 2009 and the ATCC accession number is PTA-10296. Upon issuance of a patent, all restrictions upon the deposit will be removed, and the deposit is intended to meet all of the requirements of 37 C.F.R. §§ 1.801-1.809. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced as necessary during that period.

Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 21

<210> SEQ ID NO 1
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 5' junction

<400> SEQUENCE: 1

ctataagtat ttgatggtca at

22

<210> SEQ ID NO 2
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: 3' junction

<400> SEQUENCE: 2

tcagtgtttg atttacttaa ac

22

<210> SEQ ID NO 3
 <211> LENGTH: 3605

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' flank + 100 bp insert
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (3451)..(3463)
<223> OTHER INFORMATION: The sequence at position 3451 to 3463 may
    contain additional c residues including, for example, from 13 to
    100 or more total consecutive c residues.

<400> SEQUENCE: 3

gtcgtcgaag gtgtgtcgtt gaagctcctt tttttaagta gtggtagaaa tactagaaat    60
ataatatgca tgggataatc attaacttac cttaaagagt tggaagaagg catccttttc    120
atggtcctttc attgctcctg atgtaggcca tggatgaaga aactattgct tgatggctcgt    180
tgagatagct tgtgacgcaa tctgcttgg ttcaaaccta aatacaaaaag aagggtacac    240
aaaattaaaa gacaatgaat gagttataat atatatgaa aaagtttga attctaattt    300
actcttacc cccaccctg ggtataatca tagggcgatc attgggtaag ggttctttca    360
tttccccgtc taaatcatgt tcattgatag acaatttata taatgatata ttatattaat    420
acatattdaa attttcaact ttaaataat ctaaaaaatc ttttaacaa taataagtat    480
aacaatttaa tatatatata tatcattata ttaacaacg ttaaccatta aaatttcctt    540
atggtgttaa ctatttaagt agtcattttt tattatctca cgtgtttaa aacattcaat    600
gggtcctgtt tcttttttg accgattttg tttattatcg atgaattata ttaccgacaa    660
aacatttgtc ggaaagctct gtcgataatg ctaacttttc tgacaaaat tttatccatc    720
aaaaagttcc atcacaata aataatttaa aaattcatca gaaaaaatc atatttcccta    780
catactataa tccgtcagaa atttccattg gaactaaca catttttatg gacatatatt    840
cgtcggaaca ctctatcgtt aattttaatg tttatgata attagatct gtcaattata    900
tttcgccgaa aatcgacgct tttttgtag tgattaatc aatgatgaaa tgaattatat    960
taatgatgat atatgaatgt attaatgatg agatatgaca ttgtttgaaa tgattataat   1020
atacatgcat ctatgaatga cattgcatat gagttggcac gtaagttgag actacttaga   1080
aggccctcaa cctagtgttg gaggttgaca tgggtgctat agaatgaata ggtgggtgaa   1140
tccttgataa ggttaagggt tttgcaagcc ttatcgaggc aagccattat ccatactcat   1200
caattttaga tacaaccaca cttccttcat agggttaatt ggagtctccc tgagtggcca   1260
agggtgact atgttaaggg atgtactagg ttaatcgctc aaaaagtaa atctccttgg   1320
agtaaagtgt caacatgtca tatcataata catagacatg tgcatttacc tttatgaata   1380
aagaagaatg acatgttact atatgagtta tggaaatgct ttgatgatg atgattgaat   1440
ctttatgtaa tgcttgatga actaacgatt ataagattga ccttaaaaca acaagatta   1500
atacaattg ctataaaaca aacttgattt ttatctcgt tacctttctt aataatgaaa   1560
aataatttga aaaaagcgag tatggccata aggcacatag ttttcaatt gtatatacat   1620
atcatatgat gtcttcttat cttatcatgt gttttggctt tttagcttgg aaagcgtgaa   1680
aaatttataa gtattttcac aattaaatta atagaaggtt tttaaaagaa ttaaaataaa   1740
gtcttccgct tgagtgttc ttagtggat ataaccatga caaatcatta caattacaag   1800
taagaaagac atatttaagg caaaaaatag tcttcgacta tcggttaagg acattattag   1860
tgcattcaaa aaagaaaaaa acaaaaaaca aaattttttt attgtagcca actagaatat   1920
aaagatgttg aagcaggtta ttgtacaaaa gtttaattatt ggcaaatatg acaggtattt   1980

```

-continued

```

tgtatTTTT atTTTgCcat aacattgata caaaatattt acatacttta ttgatgacta 2040
atattTTTT agaaacctaa ctaaactatc tataataaag tctctcattt taattacatt 2100
tttgTTTTca ttgatgagac ttgaacccaaa acctttgtca agggatcaaa cacaattctc 2160
cacacaaaoc aataaatcaa catgTTTgtt taaagttgag attagagata ctaattcata 2220
cttataacat agcataaaat atttatatta ttgaaaaat aaaataagtc agaagaaat 2280
aatataagat tctTTTgttt aattaaatga ttagaagaa aataaaatag tataaaataa 2340
TTTTagTta attatttatt ttaactctat agttatctcc TTTTaaGat ttagTTTTgt 2400
acgaaaaaaa tataaattgt aattcttata tatacataaa agTTaaattt taccaaaagc 2460
attaaacttt ttaatctatt attatttttc taacatgaaa ctattaaata ttaattttaa 2520
TTTTtattgt tataaagaaa tttatttaac TTTTaaGaa taaaattaa aaaatgaaaa 2580
gagattttat tctctttatt tatgccacag gcagggacag aggcagaaat aaacattag 2640
gggccaattt tgaatttaaa atttatctaa ttaaaaatg tatgagtatt aattttaaC 2700
aaactactgc taatatgttt tttccaaaa aaattcaatg acatataata gaaatatatt 2760
cacataaaag acaattaaa catttaaaaa atctattaca aatgtcattt gacatgtac 2820
catgatTTag tttctaatat ttactttgtt tttccaaat ctagtcaata gtgtgcctag 2880
accatgcact caatgttaaa agcatgtagg agcatcaca tgacaataaa atcaagataa 2940
tattttatat aatgttcata attaagcttg tcccatagac aaagattact aaataattt 3000
aacttctcaa ataaattata atcaaatcaa tataaaggaa gaataataat gtcttttaT 3060
cacaatgca ataatttata acctttaaaa tagtataaaa aattataaaa ataaaattta 3120
tgaaaaaaat aaatgagact taaattttt taaaagaca aaatgtaaaa tttttacaat 3180
aaaatctact aaaaagaaat agaataattg ttttaaaaa aataaaaatg tcattatagt 3240
tatgcaaaac tataaaggat aattgattat tttattaaa aataaaaat aacattttaa 3300
tataattatt tttataatta aacttttaaa aaataattat ctaaaattta aaatattatg 3360
tatattaaaa aataaaaact aaaaaaaaaag ttaggggggc ctcgcacctc cgtatctggc 3420
cgcaggagaa gttatcacat gagtttgcg ccccccccc cccgcgcc cctattttt 3480
ttttctatta tttgctataa gtatttgatg gtcaatatgg agaaaaagaa agagtaatta 3540
ccaattttt tccaattcaa aaatgtagat gtccgcagcg ttattataaa atgaaagtac 3600
atTTT 3605

```

```

<210> SEQ ID NO 4
<211> LENGTH: 2065
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' flank + 100 bp insert

<400> SEQUENCE: 4

```

```

ataaaaatgt gagtacattc tcgagcagga cctgcagaag ctagcttgat ggggatcaga 60
ttgtcgtttc ccgccttcag tttaaactat cagtgttga tttacttaa cgaggtaatt 120
gtaaaactta aacttgatat agttaaattg tatatatata tatatataat acttaactta 180
tttaattatc aaaatttatt ttgaacccaaa cttttactaa tttaatatta atttattttt 240
ttgttaaaaa ggcaagtaaa gtaataaaaa aattcataaa atgattttaa attatagagt 300
gataaaacta tcttttttag tacataaaat ttcaaaaacg gagaccatga tgtctttagg 360
attaaaaagt aattaaagga aaaatgtaaa aaaaaaaaa ttaaggaact aaaaataagg 420

```

-continued

```

ataaaaagt aaagaatggt gatcaataaa aaaaagagt tcaataacta aaaacaaaa 480
tcctaaaaat attagtgact agtaaatact aatcttgaac aacgttctat caataattca 540
attaagatc gagtgaagga agaatgcttc tttaaaaata aactaataaa gagatatata 600
catttcatac ttttttttat cacgtataca tttctctctt aaaaaacat atatacattt 660
ctaaagaaaa tattgcacat ttctactttc aaaatacatt tgacggtaac atgtagatgg 720
agggaaaaat tatggttttt cattttttatt ttagtgtaa tttattttta ttatttttt 780
ttttatctta ttgatagcc tgacattttt ttgatttgtt tttttattaa ttgtgtgtaa 840
agacaaagta ttagatttgt gcaataaac taggtcaaga ctaaagtta ttaattttta 900
aagactagac tttagatcat ttaaaaaaaa ttttaagtct aacatgtcaa cctatttaag 960
taaatgtaag tatttaattt ttgttgttat tttttataat ggataggat tcagttgggt 1020
caagaacccc ataaccatcc ctactaaata ggtatccagt cttaccccca attggcccc 1080
acaaacaggt gattaaaccc ataataaaaa aagtgaattg attaaggga aagagaagt 1140
taggccactt aaacccaat tatggaagaa ttacctagac ggtaatccaa gcacaaaggt 1200
ccatggttca aggccacaa atgtttgtat aaataggggg aaaaccctaa gtaatgacca 1260
ttgttcgttt gatgtgcatt ttattgctt gcttactact cacaattgaa ctctgactta 1320
agcgtttgag tatcttttgc aagtatcccc ccttgggtc atcttatttg gcacgtgaag 1380
aagacaaagt ccaaggcag gacccttcta ttagaggac agtgtccgac cattatttgg 1440
taggaatatt ttggcacccc aaagtaagat agtccttgac cctaagaatt atggcaatca 1500
cacgcacaag aaccaacata gcagcaaaaa ccaacttggg aatcattagc ctgaagaagt 1560
ccatgtagag ggctagattg acttatoctt aatggtgagg aggctacaag cacaaattt 1620
agatagcgt caacatcatg tcgaagaggt cattaccctt tgacaagaaa atgcacaact 1680
acaagagggg aaccgttctc agaaacccga aatgaaaca cctgttggag cggaggtaga 1740
gtcaaccaat tttgagatgc caaaagccac caatgttagt gtttggcagg atacacccc 1800
tcagttcata aatagttag ggaagcaact acagtagttc aagaacacct tttcctcgg 1860
tgtcatggaa gccacctcc tcccacta gaagaacctc accatcgaga agtactatgg 1920
taccactaac ccatatgagc accttgacgt ttataccacc caagtcagcc tgtacactat 1980
ggataatggt gtcttatgcc aggtgttccc tacatctttg aaggggtaga tcctccatta 2040
gttcaccac ctcctcccc caaca 2065

```

```

<210> SEQ ID NO 5
<211> LENGTH: 2014
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: insert

```

```

<400> SEQUENCE: 5

```

```

tgatgggtcaa tatggagaaa aagaaagagt aattaccaat ttttttcaa ttcaaaaatg 60
tagatgtccg cagcgttatt ataaaaatgaa agtacatttt gataaaacga caaattacga 120
tccgtcgat ttataggcga aagcaataaa caaattattc taattcggaa atctttattt 180
cgacgtgtct acattcacgt ccaaatgggg gcttagatga gaaacttcac gatttggcgc 240
gccaagcctt gatatcgaat tcctgcagcc ccttaaagg ggggatccg gtccgatgtg 300
agacttttca acaaagggtg ataccggaa acctcctcgg attccattgc ccagctatct 360

```

-continued

```

gtcactttat tgtgaagata gtggaaaagg aaggtggctc ctacaaatgc catcattgcy 420
ataaaggaaa ggccatcggt gaagatgcct ctgccgacag tggtoceaaa gatggacccc 480
caccacgag gagcatcggt gaaaaagaag acgttccaac cacgtcttca aagcaagtgg 540
attgatgtga tggtocegatt gagacttttc aacaaagggt aatatccgga aacctcctcg 600
gattccattg cccagctatc tgtcacttta ttgtgaagat agtggaaaag gaagtggtgt 660
cctacaaatg ccatcattgc gataaaggaa aggccatcgt tgaagatgcc tctgccgaca 720
gtggtcccaa agatggaccc ccaccacga ggagcatcgt ggaaaaagaa gacgttccaa 780
ccacgtcttc aaagcaagtg gattgatgtg atatctccac tgacgtaagg gatgacgcac 840
aatcccacta tccttcgcaa gacccttctt ctatataagg aagttcattt catttgagga 900
ggaccaggtg gtaccggcgc gccaccatgg tgagcttttg cgagctttgt ggtgccgaag 960
ctgatctcca ttgtgccgcy gactctgcct tectctgcgc ttcttgtgac gctaagtctc 1020
atgctcaaaa tttctcttc gctcgtcatt tccggcgtgt catctgccc aattgcaaat 1080
ctcttactca aaatttcggt tctggtcttc ttcttcttg gcctccacga acaacatggt 1140
gttcagaatc gtcgtctctt tcttgtgct cgtctcttga ctgtgtctca agctccgagc 1200
tatcgtcaac gacgcgtgac gtaaacagag cgcgaggag ggaaaacaga gtgaatgcca 1260
aggccgttgc ggttacggtg gcggatggca tttttgtaa ttggtgtggt aagttaggac 1320
taaacaggga ttaacaaac gctgctggtt catatgcctc tttggcttg gctgtggaga 1380
cgaggccaag agcgcgaag agagtgttct tagcggcggc gttttggttc ggcgtaaga 1440
acacgacgac gtggcagaat ttaaagaaag tagaagatgt gactggagtt tcagctggga 1500
tgattcgagc ggttgaagc aaattggcgc gtgcaatgac gcagcagctt agacggtggc 1560
gcygtggatc ggaggaagga tggcgtgaaa acgacaacct ttaggcgac gcyggccctg 1620
atcacctgct gtacagtatt tctacatttg atgtgtgatt tgtgaagaac atcaacaaa 1680
acaagcactg gctttaatat gatgataagt attatggtaa ttaattaatt ggcaaaaaca 1740
acaatgaagc taaaatttta tttattgagc cttgcgggta atttcttgg atgatctttt 1800
ttttatttt ctaattatat atagtttctt ttgctttgaa atgctaaagg tttgagagag 1860
ttatgctctt tttttcttcc tctttctttt ttaactttat catacaaatt ttgaataaaa 1920
atgtgagtac attctcgagc aggacctgca gaagctagct tgatggggat cagattgtcg 1980
tttccgcct tcagtttaaa ctatcagtgt ttga 2014

```

<210> SEQ ID NO 6

<211> LENGTH: 7484

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: flanks + insert

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (3451)..(3463)

<223> OTHER INFORMATION: The sequence at position 3451 to 3463 may contain additional c residues including, for example, from 13 to 100 or more total consecutive c residues.

<400> SEQUENCE: 6

```

gtcgtcgaag gtgtgctggt gaagctcctt ttttaagta gtgtagaaa tactagaat 60
ataatagca tgggataatc attaacttac cttaaagagt tggagaagg catcctttt 120
atggtcttcc attgctccgt atgtaggcca tggatgaaga aactattgct tgatggtcgt 180
tgagatagct tgtgacgcaa tctgcttgg ttcaaaccta aatacaaaag aagggtacac 240

```

-continued

aaaattaaaa	gacaatgaat	gagttataat	atatattgaa	aaagttttga	attctaattt	300
actcttacc	tccaccctg	ggtataatca	tagggcgatc	attgggtaag	ggttctttca	360
tttccccg	tcaatcatgt	tcattgatag	acaatttata	taatgatata	ttatattaat	420
acataattaa	attttcaott	ttaatatata	ctaaaaaatc	ttttaatcaa	taataagtat	480
aacaatttaa	tatatatata	tatcattata	tttaacaacg	ttaaccatta	aaatttctct	540
atgtgtttaa	ctatttfaat	agtcattttt	tattatctca	cggtgttfaat	aacattcaat	600
gggtcttg	tctttttttg	accgattttg	tttattatcg	atgaattata	ttaccgacaa	660
aacatttg	ggaaagctct	gtcgataatg	ctaacttttc	tgacaaatat	ttatccatcg	720
aaaaagttcc	atcacaata	aataatttaa	aaattcatca	gaaaaaatc	atatttctca	780
catactataa	tccgctagaa	atttccattg	gaactaacia	catttttatg	gacatatatt	840
cgctcgaa	ctctatcgtt	aatttfaat	tttatgataa	atttagatct	gtcaattata	900
tttcccgaa	aatcgacgct	ttttttgtag	tgattaattc	aatgatgaaa	tgaattatat	960
taatgatgat	atatgaatgt	attaatgatg	agatatgaca	ttgtttgaaa	tgattataat	1020
atacatgcat	ctatgaatga	cattgcatat	gagttggcac	gtaagttgag	actacttaga	1080
aggccctcaa	cctagtgttg	gaggttgaca	tggtggctat	agaatgaata	ggtgggtgaa	1140
tccttgata	ggttaagggt	tttgcagcc	ttatcgaggc	aagccattat	ccatactcat	1200
caattttaga	tacaaccaca	cttcttcat	agggttaatt	ggagtctccc	tgagtggcca	1260
agggtgact	atgttaaggg	atgtactagg	ttaatcgctc	aaaaagtaa	atctccttgg	1320
agtaaagtgt	caacatgtca	tatcataata	catagacatg	tgcatttacc	tttatgaata	1380
aagaagaatg	acatgttact	atatgagtta	tggaaatgct	ttgatgtatg	atgattgaat	1440
ctttatgtaa	tgcttgatga	actaacgatt	ataagattga	ccttaaaaca	acaaagatta	1500
ataacaattg	ctataaaaca	aacttgtatt	ttatcttctg	taccttctt	aataatgaaa	1560
aataatttga	aaaaagcgag	tatggccata	aggcacatag	tttttcaatt	gtatatacat	1620
atcatatgat	gtcttcttat	cttatcatgt	gttttggctt	tttagcttgg	aaagcgtgaa	1680
aaatttataa	gtattttcac	aattaaatta	atagaagggg	tttaaaagaa	ttaaaataaa	1740
gtcttccgct	tgagtgttc	ttagtggat	ataaccatga	caaatcatta	caattacaag	1800
taagaaagac	atatttfaat	caaaaaatag	tcttcgacta	tcggttfaat	acattattag	1860
tgcatcacia	aaagaaaaaa	aacaaaaaca	aaattttttt	attgtagcca	actagaatat	1920
aaagatgtg	aagcaggtta	ttgtacaaaa	gttaattatt	ggcaaatatg	acaggtattt	1980
tgtatttttt	attttgccat	aacattgata	caaaatattt	acatacttta	ttgatgacta	2040
atatttttta	agaaacctaa	ctaaactatc	tataataaag	tctctcattt	taattacatt	2100
tttgttttca	ttgatgagac	ttgaacaaa	acctttgtca	agggatcaaa	cacaattctc	2160
cacacaaacc	aataaatcaa	catgtttgtt	taaagttgag	attagagata	ctaattcata	2220
cttataacat	agcataaaat	atttatatta	tttgaaaaat	aaaataagtc	agaaagaaat	2280
aatataagat	tcttttgttt	aattfaatga	ttagaaagaa	aaataaatag	tataaaataa	2340
ttttaggtta	attatttatt	ttaactctat	agttatctcc	tttttaagat	ttagttttgt	2400
acgaaaaaaa	tataaattgt	aattcttata	tatacataaa	agttaaattt	tacaaaaagc	2460
atataacttt	ttaatctatt	attatttttc	taacatgaaa	ctattaaata	ttaattttaa	2520
ttttttatgt	tataaagaaa	tttatttaac	tttttaagaa	taaaaattaa	aaatgaaaa	2580

-continued

gagattttat tctctttatt tatgccacag gcagggacag aggcagaaat aaacattagg	2640
gggccaattht tgaattataa atttatctaa ttataaaaatg tatgagtatt aatttttaaac	2700
aaactactgc taatatgttt tttccaaaaa aaattcaatg acatataata gaaatatatt	2760
cacataaaaag acaaatataa catttaaaaa atctattaca aatgtcattt gacatgctac	2820
catgatthtag tttctaataat ttactttgtt tttccaaaaa ctagtcaata gtgtgcctag	2880
accatgcact caatgttaaa agcatgtagg agcatcacia tgacaataaa atcaagataa	2940
tattttatat aatgttcata attaagcttg tcccatagac aaagattact aaataattht	3000
aacttctcaa ataaattata atcaaatcaa tataaaggaa gaataataat gtctthtaat	3060
cacaaatgca ataatttata acctttaaaa tagtataaaa aattataaaa ataaaattht	3120
tgaaaaaaat aatgagact taaatthttt taaaaagaca aatgtaaaa thtttacaat	3180
aaaatctact aaaaagaat agaataatg thtttaaaaa aataaaaatg tcattatagt	3240
tatgcaaaac tataaaggat aatgattat thttataaaa aaataaaaat aacattthta	3300
tataattatt thttataatta aactthtaaa aaataattat ctaaatttht aaatattatg	3360
tatattaaaa aataaaacta aaaaaaaaag ttaggggggc ctgcacctc cgtatctggc	3420
cgcaggagaa gttatcacat gagthtcgag ccccccccc cccgcgccc cctatthttt	3480
thttctatta thtgctataa gtatthgatg gtcaatatgg agaaaaagaa agagtaatta	3540
ccaatthttt thcaatcaa aaatgtagat gtccgcagag thttataaaa atgaaagtac	3600
atthtgataa aacgacaaat tacgatccgt cgtatthta ggcaaaaagca ataaacaaat	3660
tattctaatt cggaaatctt thttcgagc tgtctacatt cacgtccaaa tgggggctta	3720
gatgaaaaac thcacgattt ggccgcgcaa agcttgatat cgaattcctg cagccccctt	3780
aaaggggggg atccggctcg atgtgagact thtcaacaaa gggtaatatc cggaaaacctc	3840
ctcggattcc attgccagc tatctgtcac thttatgtga agatagtgga aaaggaaggt	3900
ggctcctaca aatgccatca thgcgataaa ggaaaggcca tcgttgaaga tgctctgccc	3960
gacagtggtc ccaaatgtag acccccaccc acgaggagca tcgtggaaaa agaagacgtt	4020
ccaaccacgt cttcaagca agtgattga tgtgatggtc cgattgagac thttcaacaa	4080
agggtaatat ccgaaaacct cctcggattc cattgccagc ctatctgtca cthttatgtg	4140
aagatagtgg aaaaaggaag tggctcctac aaatgccatc attgcgataa aggaaaggcc	4200
atcgttgaag atgcctctgc cgacagtggc cccaaagatg gacccccacc cacgaggagc	4260
atcgtgaaa aagaagacgt thcaaccacg thttcaagc aatggattg atgtgatatc	4320
thcaactgag taagggatga cgcacaatcc cactatcctt cgcaagaccc thctctata	4380
taaggaaatt catttcattt ggagaggacc aggtggtaac ggccgcccac catggtgagc	4440
thttgcgagc thttgtgtgc cgaagctgat thccattgtg ccgcggactc tgctctcctc	4500
tgccgttctt gtgacgctaa gthccatgccc thaaatthtt thttcgctcg thatttcggg	4560
cgtgtcatct gcccatttg caaatctctt actcaaaatt tcgtthctgg thctctctt	4620
ccttggcctc cacgaacaac atgttgthta gaatcgtctt thttctcttg ctgctcgtct	4680
cttgactgtg thtcaagctc cgagctatcg thcaacgagc gtgacgtaaa cagagcgcga	4740
gggagggaaa acagagtga thccaaggcc gthtgggtht cgggtggcgga tggcatttht	4800
gtaaatgggt gtggtaagtt aggactaaac agggatttht caaacgctgt cgtthcatat	4860
gcgtctthgg cthttggctgt ggagacgagc ccaagagcga cgaagagagt gthcttagcg	4920
ggcgctthtt gthtccggct taagaacacg acgacgtggc agaattthaa gaaagtagaa	4980

-continued

gatgtgactg gagtttcagc tgggatgatt cgagcggttg aaagcaaatt ggcgcgtgca	5040
atgacgcagc agcttagacg gtggcgcgtg gattcggagg aaggatgggc tgaaaaacgac	5100
aacgttttagc cgatcgcggg cccatgatcac ctgtcgtaca gtatttctac atttgatgtg	5160
tgatttgtga agaacaatcaa acaaaaacaag cactggcctt aatatgatga taagtattat	5220
ggtaattaat taattggcaa aaacaacaat gaagctaaaa ttttatttat tgagccttgc	5280
ggtaatttc ttgtgatgat cttttttttt attttctaata tatatatagt ttcccttgc	5340
ttgaaatgct aaaggttga gagagttagc ctcttttttt cttcctcttt cttttttaac	5400
tttatcatac aaattttgaa taaaaatgtg agtacattct cgagcaggac ctgcagaagc	5460
tagcttgatg gggatcagat tgcctgttcc cgccttcagt ttaaactatc agtggttgat	5520
ttacttaaac gaggtaattg taaaacttaa acttgatata gttaaatggt atatatatat	5580
atatataata ctttaacttat ttaattatca aaatttattt tgaaccaaac ttttactaat	5640
ttaattatca tttatttttt tgttaaaaag gcaagtaaaag taataaaaaa attcataaaa	5700
tgatttaaaa ttatagagtg ataaacttat ctttttttagt acataaaatt tcaaaaacgg	5760
agaccatgat gtcttttagg taaaaagta attaaaggaa aaatgtaaaa aaaaaaagt	5820
taaggaacta aaaataagga taaaaagtta aagaatgttg atcaataaaa aaaaagagt	5880
caataactaa aaacaaaaat cctaaaaata ttagtgacta gtaataacta atcttgaaca	5940
acgttctatc aataattcaa ttaaagatcg agtgaaggaa gaatgcttct ttaaaaaata	6000
actaataaag agatatatac atttcatact tttttttatc acgtatacat ttctctctta	6060
aaaaaccata tatacatttc taaagaaaat attgcacatt tctactttca aaatacattt	6120
gacgtaaca ttagatgga gggaaaaatt atggtttttc atttttattt tagtgtaaat	6180
ttatttttat tattattttt tttatcttat ttgatagcct gacatttttt tgatttgttt	6240
ttttattaat tgtgtgtaaa gacaaaagtat tagatttgtyg caaataaaact aggtcaagac	6300
taaagtttat ttaattttta agactagact ttagatcatt taaaaaaat ttaagtcta	6360
acatgtcaac ctattttaagt aaatgtaagt atttaatttt tggttgtatt ttttataatg	6420
gataggtatt cagttgggtc aagaacccca taacctccc tactaaatag gtatccagtc	6480
ttaccccaaa ttggcccca caaacaggtg attaaaccca taataaaaaa agtgaattga	6540
ttaagggaaa agagaagttt aggccactta aacccaatt atggaagaat tacctagacg	6600
gtaatccaag cacaaaggtc catggttcaa ggcccacaaa tgtttgtata aataggggga	6660
aaaccctaag taatgacct tgttcgtttg atgtgcattt tattgccttg cttactactc	6720
acaattgaac tctgacttaa gcgtttgagt atcttttgca agtatcccc ccttgggtca	6780
tcttatttgg cacgtgaaga agacaaagtc caaaggcagg acccttctat tagagggaca	6840
gtgtccgacc attatttggg aggaatattt tggcaccoca aagtaagata gtccttgacc	6900
ctaagaatta tggcaatcac acgcacaaga accaacatag cagcaaaaac caacttggga	6960
atcattagcc tgaagaagtc catgtagagg gctagattga cttatcctta atggtaggga	7020
ggctacaagc acaagtatta gatatgcgtc aacatcatgt cgaagaggtc attacccttt	7080
gacaagaaaa tgcacaacta caagagggtg accgttccta gaaacccgaa aatgaaacac	7140
ctgttgagc ggaggtagag tcaaccaatt ttgagatgcc aaaagccacc aatgtagtg	7200
ttggcagga tacacacct cagttcataa atatgttagg gaagcaacta cagtagttca	7260
agaacacett tttcctcggg gtcattggaag ccacctcct tcccaactag aagaacctca	7320

-continued

```

ccatcgagaa gtactatggt accactaacc catatgagca ccttgacggt tatatcaccc 7380
aagtcagcct gtacacatag gataatggtg tcttatgcca ggtgttcctt acatctttga 7440
aggggtagat cctccattag ttcacccacc tcctccccc aaca 7484

```

```

<210> SEQ ID NO 7
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' longer junction

```

```

<400> SEQUENCE: 7

```

```

cccccccgcg cgcccctat ttttttttc tattatttgc tataagtatt tgatggtcaa 60
tatggagaaa aagaaagagt aattaccaat tttttttcaa 100

```

```

<210> SEQ ID NO 8
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' longer junction

```

```

<400> SEQUENCE: 8

```

```

gggatcaga ttgtcgttcc cgccttcag tttaaactat cagtgtttga tttacttaaa 60
cgaggtaatt gtaaaactta aacttgatat agttaaagt 100

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1644
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: BBX32 cassette

```

```

<400> SEQUENCE: 9

```

```

ggtccgatgt gagacttttc aacaaagggt aatatccgga aacctcctcg gattccattg 60
cccagctatc tgtcacttta ttgtgaagat agtggaaaag gaaggtggct cctacaaatg 120
ccatcattgc gataaaggaa aggccatcgt tgaagatgcc tctgccgaca gtggtcccaa 180
agatggaccc ccaccacga ggagcatcgt ggaaaaagaa gacgttccaa ccacgtcttc 240
aaagcaagtg gattgatgtg atgggtccgat tgagactttt caacaaaggg taatatccgg 300
aaacctcttc ggattccatt gccagctat ctgtcacttt attgtgaaga tagtgaaaa 360
ggaaggtggc tcctacaat gccatcattg cgataaagga aaggccatcg ttgaagatgc 420
ctctgccgac agtgggtcca aagatggacc cccaccacg aggagcatcg tggaaaaaga 480
agacgttcca accacgtctt caaagcaagt ggattgatgt gatatctcca ctgacgtaag 540
ggatgacgca caatcccact atccttcgca agacccttc tctatataag gaagttcatt 600
tcattggag aggaccaggt ggtaccggcg cgccaocatg gtgagctttt gcgagctttg 660
tggtgccgaa gctgatctcc attgtgccgc ggactctgcc tcctctgcc gttctgtgta 720
cgctaagttc catgcctcaa atttctctt cgctcgtcat tccggcgtg tcatctgccc 780
aaattgaaa tctcttactc aaaatttctg ttctggctct cttctctctt ggctccacg 840
aacaacatgt tgttcagaat cgtcgtcttc ttctgtctgc tcgtctcttg actgtgtctc 900
aagctccgag ctatcgtcaa cgacgcgtga cgtaaacaga gcgcgagggga gggaaaaacg 960
agtgaatgcc aaggccgttg cggttacggt ggccgatggc atttttgtaa attggtgtgg 1020
taagttagga ctaaacaggg atttaacaaa cgctgtcgtt tcatatgcgt ctttggcttt 1080

```

-continued

```

ggctgtggag acgaggccaa gagcgacgaa gagagtgttc ttagcggcgg cgttttggtt 1140
cggcgtaag aacacgacga cgtggcagaa tttaaagaaa gtagaagatg tgactggagt 1200
ttcagctggg atgattcgag cggttgaaag caaattggcg cgtgcaatga cgcagcagct 1260
tagacggtgg cgcgtggatt cggaggaagg atgggctgaa aacgacaacg tttagcggat 1320
cgcgggccct gatcacctgt cgtacagtat ttctacattt gatgtgtgat ttgtgaagaa 1380
catcaacaa aacaagcact ggctttaata tgatgataag tattatggta attaattaat 1440
tggcaaaaac aacaatgaag ctaaaatttt atttattgag ccttgcgggtt aatttcttgt 1500
gatgatcttt tttttatatt tctaattata tatagtttcc tttgctttga aatgctaaag 1560
gtttgagaga gttatgtctt tttttcttcc ctctttcttt ttaacttta tcatacaaat 1620
tttgaataaa aatgtgagta catt 1644

```

```

<210> SEQ ID NO 10
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: reverse primer for event zygosity

```

```

<400> SEQUENCE: 10

```

```

gttttacaat tacctcgttt aagtaaatca 30

```

```

<210> SEQ ID NO 11
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: forward primer for event zygosity

```

```

<400> SEQUENCE: 11

```

```

cattctcgag caggacctgc agaa 24

```

```

<210> SEQ ID NO 12
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 6FAM-labeled probe for event zygosity

```

```

<400> SEQUENCE: 12

```

```

aacactgata gtttaaactg aag 23

```

```

<210> SEQ ID NO 13
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: forward primer for wild type allele

```

```

<400> SEQUENCE: 13

```

```

ctattatttg ctataagtat ttgatgtaag aa 32

```

```

<210> SEQ ID NO 14
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: VIC-labeled probe for wild type allele

```

```

<400> SEQUENCE: 14

```

-continued

ttgtattaat aacaaaaaat tg	22
<210> SEQ ID NO 15 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: event-specific primer for zygoty	
<400> SEQUENCE: 15	
gtacattctc gagcaggacc tgca	24
<210> SEQ ID NO 16 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: event-specific primer SQ3983	
<400> SEQUENCE: 16	
tgatgggat cagattgtcg t	21
<210> SEQ ID NO 17 <211> LENGTH: 34 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: event-specific primer SQ22982	
<400> SEQUENCE: 17	
actatatcaa gtttaagttt tacaattacc tcgt	34
<210> SEQ ID NO 18 <211> LENGTH: 22 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: event-specific probe PB10453	
<400> SEQUENCE: 18	
ccttcagttt aaactatcag tg	22
<210> SEQ ID NO 19 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: primer SQ1532 for internal control	
<400> SEQUENCE: 19	
ggtatccctc cagaccagca	20
<210> SEQ ID NO 20 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: primer SQ1533 for internal control	
<400> SEQUENCE: 20	
gtggactcct tctggatggt gtaa	24
<210> SEQ ID NO 21 <211> LENGTH: 29 <212> TYPE: DNA <213> ORGANISM: Artificial sequence	

-continued

<220> FEATURE:

<223> OTHER INFORMATION: VIC-labeled probe PB0359 for internal control

<400> SEQUENCE: 21

atatttgctg gaaagcagct tgaggatgg

29

The invention claimed is:

1. A method of determining the zygosity of a soybean plant genome comprising soybean event MON87712 DNA in a sample, the method comprising:

a) contacting the sample with three different primers that

i) when used together in a nucleic acid amplification reaction with soybean event MON87712 DNA, produces a first amplicon that is diagnostic for soybean event MON87712; and

ii) when used together in a nucleic acid amplification reaction with soybean genomic DNA other than event MON87712 DNA, produces a second amplicon that is diagnostic for soybean wild type genomic DNA other than event MON87712 DNA;

b) performing a nucleic acid amplification reaction; and

c) detecting said first amplicon and said second amplicon; wherein the presence of said first and second amplicons is diagnostic of a genome heterozygous for event MON87712 in said sample, and wherein the presence of only said first amplicon is diagnostic of a genome homozygous for event MON87712 in said sample.

2. The method of claim 1, wherein the three different primers comprise (i) SEQ ID NO:10 or 17, (ii) SEQ ID NO:11, 15, or 16, and (iii) SEQ ID NO:13.

3. The method of claim 1, wherein the first amplicon comprises SEQ ID NO: 2, 12, or 18.

4. The method of claim 1, wherein the second amplicon comprises SEQ ID NO: 14.

5. A method of determining the zygosity of a soybean plant genome comprising soybean event MON87712 DNA in a sample, the method comprising:

a) contacting the sample with three different primers and a probe set that comprises a first probe and a second probe, wherein said first probe comprises a first fluorescent label and said second probe comprises a second fluorescent label that

i) when used together in a nucleic acid amplification reaction with soybean event MON87712 DNA, produces a first amplicon and releases a first fluorescent signal from said first probe that is diagnostic for soybean event MON87712 DNA; and

ii) when used together in a nucleic acid amplification reaction with soybean genomic DNA other than event MON87712 DNA, produces a second amplicon and releases a second fluorescent signal from said second probe that is diagnostic for soybean wild type genomic DNA other than event MON87712 DNA;

b) performing a nucleic acid amplification reaction; and
c) detecting said first fluorescent signal and said second fluorescent signal;

wherein the presence of said first and second fluorescent signals is diagnostic of a genome heterozygous for event MON87712 in said sample, and wherein the presence of only said first fluorescent signal is diagnostic of a genome homozygous for event MON87712 in said sample.

6. The method of claim 5, wherein the three different primers comprise (i) SEQ ID NO:10 or 17, (ii) SEQ ID NO:11, 15, or 16, and (iii) SEQ ID NO:13.

7. The method of claim 5, wherein said first probe comprises SEQ ID NO: 12 or 18 and said second probe comprises SEQ ID NO: 14.

8. The method of claim 5, wherein said first fluorescent label comprises 6FAM™ and said second fluorescent label comprises VIC™.

9. The method of claim 5, wherein said first amplicon comprises SEQ ID NO: 2, 12, or 18.

10. The method of claim 5, wherein said second amplicon comprises SEQ ID NO: 14.

* * * * *