



(86) **Date de dépôt PCT/PCT Filing Date:** 2013/06/04
(87) **Date publication PCT/PCT Publication Date:** 2013/12/27
(85) **Entrée phase nationale/National Entry:** 2014/12/12
(86) **N° demande PCT/PCT Application No.:** KR 2013/004913
(87) **N° publication PCT/PCT Publication No.:** 2013/191396
(30) **Priorité/Priority:** 2012/06/20 (KR10-2012-0066333)

(51) **Cl.Int./Int.Cl.** **C07C 311/12** (2006.01),
A61K 31/18 (2006.01), **A61P 3/10** (2006.01)
(71) **Demands/ Applicants:**
AHN-GOOK PHARMACEUTICAL CO., LTD., KR;
BAMICHEM CO., LTD., KR;
INCHEON UNIVERSITY INDUSTRY ACADEMIC
COOPERATION FOUNDATION, KR
(72) **Inventeurs/Inventors:**
HAN, CHEOL KYU, KR;
AUH, JIN, KR;
CHOI, NAM SONG, KR;
HAN, CHANG KYUN, KR;
...
(74) **Agent:** NORTON ROSE FULBRIGHT CANADA

(54) **Titre : NOUVEAU COMPOSE AYANT LA CAPACITE A INHIBER L'ENZYME 11B-HSD1 OU UN SEL PHARMACEUTIQUEMENT
ACCEPTABLE DE CELUI-CI, PROCEDE DE PRODUCTION ASSOCIE, ET COMPOSITION PHARMACEUTIQUE CONTENANT
CELUI-CI EN TANT QUE PRINCIPE ACTIF**
(54) **Title: NOVEL COMPOUND HAVING ABILITY TO INHIBIT 11B-HSDI ENZYME OR PHARMACEUTICALLY ACCEPTABLE SALT
THEREOF, METHOD FOR PRODUCING SAME, AND PHARMACEUTICAL COMPOSITION CONTAINING SAME AS ACTIVE
INGREDIENT**

(57) **Abrégé/Abstract:**

The present invention relates to a novel compound having the ability to inhibit the 11 β -HSD1 enzyme or to a pharmaceutically acceptable salt thereof, to a method for producing same, and to a pharmaceutical composition containing same as an active ingredient. The compound according to the present invention selectively inhibits the activity of 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1), and can thus be effectively used as a therapeutic agent for treating diseases caused by the excessive activity of 11 β -HSD1, e.g. non-insulin dependent (type 2) diabetes, insulin resistance, obesity, lipid disorders, metabolic syndrome, and diseases and conditions brought on by the excessive activity of glucocorticoids.



(72) **Inventeurs(suite)/Inventors(continued)**: KIM, TAE-JEONG, KR; PAE, KAMSA, KR; SHIN, YOUNG JUNE, KR;
HAN, DONG-OH, KR; AHN, SOON KIL, KR

(74) **Agent(suite/continued)**: LLP/S.E.N.C.R.L., S.R.L.

ABSTRACT

The present invention relates to a novel compound or a pharmaceutically acceptable salt thereof
5 inhibiting 11 β -HSD1 enzyme activity, a preparation method of the same, and a pharmaceutical composition comprising the same as an active ingredient. Since the compound of the present invention selectively inhibits the activity of 11 β -HSD1 (11 β -Hydroxysteroid
10 dehydrogenase type 1), the compound of the invention can be effectively used as a therapeutic agent for the treatment of diseases caused by the over-activation of 11 β -HSD1 such as non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder,
15 metabolic syndrome, and other diseases or condition mediated by the excessive activity of glucocorticoid.

NOVEL COMPOUND HAVING ABILITY TO INHIBIT 11B-HSD1
ENZYME OR PHARMACEUTICALLY ACCEPTABLE SALT THEREOF,
METHOD FOR PRODUCING SAME, AND PHARMACEUTICAL
COMPOSITION CONTAINING SAME AS ACTIVE INGREDIENT

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a novel compound
10 or a pharmaceutically acceptable salt thereof
inhibiting 11 β -HSD1 enzyme activity, a preparation
method of the same, and a pharmaceutical composition
comprising the same as an active ingredient.

15 2. Description of the Related Art

Metabolic syndrome is the disease in increase
particularly in advanced countries, in Asia, and in
developing countries. This disease is characterized
by obesity, type II diabetes, hyperlipidemia,
20 hypertension, arteriosclerosis, coronary heart
disease, and chronic renal failure (C. T. Montague et
al.(2000), Diabetes, 49,883-888).

Glucocorticoid (cortisol in human, and
corticosterone in rat and mouse) binding to
25 glucocorticoid receptor is a steroid hormone found in

almost every vertebrate (Dallman MF, Strack AM, Akana SF et al. 1993; Front Neuroendocrinol 14, 303-347).

This hormone regulates the expressions of liver enzymes involved in gluconeogenesis so as to release
5 glycerol from adipocytes and amino acid from muscle, resulting in the increase of substrate supplement. It has also been reported that glucocorticoid plays an important role in the differentiation of adipocyte precursors into mature adipocytes that can store
10 triglyceride (Bujalska IJ et al. 1999; Endocrinology 140, 3188-3196). This implies that the glucocorticoid induced by "stress" is deeply involved in the disease caused by abdominal obesity which is a major risk factor causing type II diabetes, hypertension, and
15 coronary artery disease (Bjorntorp P & Rosmond R 2000; Int. J. Obesity 24, S80-S85).

It was experimentally confirmed that the activity of glucocorticoid could be controlled in tissue level not only by the secretion of cortisol but also by the
20 intracellular interconversion between active cortisol and inactive cortisol mediated by 11 β -HSD1 (11 β -Hydroxysteroid dehydrogenase type 1) and 11 β -HSD2 (11 β -Hydroxysteroid dehydrogenase type 2) (Sandeep TC & Walker BR 2001 Trends in Endocrinol & Metab. 12,
25 446-453).

Glucocorticoid and 11β -HSD1 are known as important factors for the differentiation of adipocytes. The 11β -HSD1 mRNA level is increased in the visceral fat tissue of an obesity patient, 5 compared with that in the subcutaneous tissue. The over-expression of 11β -HSD1 in the fat tissue of a transgenic mouse is related to the up-regulation of corticosterone in the fat tissue, visceral obesity, insulin sensitivity, type II diabetes, hyperlipidemia, 10 and hyperphagia (H. Masuzaki et al(2001), Science, 294, 2166-2170). Thus, it can be said that 11β -HSD1 is mainly involved in visceral obesity and metabolic syndrome.

At this time, 11β -HSD1 converts the inactive 15 glucocorticoid to the active form, suggesting that it plays an important role in activating glucocorticoid receptor in target tissues and in regulating the concentrations of glucocorticoid in there.

It has been confirmed that the above mechanism 20 can be favorably used for the treatment of diabetes and obesity. Particularly, the treatment effect was confirmed when the antiulcerative agent carbenoxolone that can inhibit both 11β -HSD1 and 11β -HSD2 was used for the treatment. This treatment increased insulin 25 sensitivity, and the suppressed 11β -HSD1 reduced

intracellular cortisol level, suggesting that the insulin effect could be controlled (Walker BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159). The inhibition of 11 β -HSD1 in the 11 β -HSD1 knock-out mouse
5 weakened the activity of gluconeogenesis enzyme to induce glucocorticoid and reduced the level of plasma glucose responding stress or obesity (Kotelevtsev Y. et al., Proc Natl Acad Sci USA. 1997 Dec 23; 94(26):14924-14929), indicating that the inhibition of
10 11 β -HSD1 was effective in lowering plasma glucose and hepatic glucose in type II diabetes. The inhibition of 11 β -HSD1 not only reduces the typical diabetes related symptoms but also does not carry any significant side effect.

15 In the course of study using the non-specific inhibitor carbenoxolone however, a side effect such as blood pressure increase was observed when 11 β -HSD2 was inhibited. So, it is necessary to develop an inhibitor that has selectivity to 11 β -HSD1.

20 Osteological development and bone function are also regulated by glucocorticoid. 11 β -HSD1 exists in human osteoclasts and osteoblasts. When a healthy volunteer was treated with carbenoxolone, the bone formation marker showed no changes but the bone
25 resorption marker was reduced (Cooper MS et al 2000;

Bone 27, 375-381). The inhibition of 11 β -HSD1 activity in bone can be used as a protective mechanism in the treatment of osteoporosis.

Further, glucocorticoid is also involved in eye
5 disease like glaucoma. 11 β -HSD1 has been known to affect the intraocular pressure in human, and thus the inhibition of 11 β -HSD1 is expected to alleviate the increased intraocular pressure in relation to glaucoma (Rauz S et al. 2001; Investigative Ophthalmology &
10 Visual Science 42, 2037-2042).

In the course of study to prepare compounds that can selectively inhibit 11 β -HSD1, the present inventors succeeded in synthesizing novel compounds
15 which showed excellent activity to inhibit 11 β -HSD1 and to have plasma glucose lowering effect dose-dependently, which were confirmed by *in vivo* animal study. The present inventors completed this invention by confirming that the novel compounds synthesized by
20 the inventors could be effectively used in the treatment of the following diseases or conditions; non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder, metabolic syndrome, and other diseases mediated by the excessive
25 activity of glucocorticoid.

SUMMARY OF THE INVENTION


It is an object of the present invention to
 5 provide a compound or a pharmaceutically acceptable
 salt thereof having the activity of inhibiting 11 β -
 HSD1 enzyme.

It is another object of the present invention to
 provide a preparation method of the said compound.

10 It is also an object of the present invention to
 provide a pharmaceutical composition comprising the
 said compound or the pharmaceutically acceptable salt
 thereof as an active ingredient.

15 To achieve the above objects, the present
 invention provides a compound represented by formula 1
 having the activity of inhibiting 11 β -HSD1 enzyme or a
 pharmaceutically acceptable salt thereof:

[Formula 1]






20

In formula 1,


X is carbonyl or sulfonyl;

R¹ is C₁₋₄ straight or branched alkyl, C₃₋₇ cycloalkyl, C₅₋₁₂ aryl, C₅₋₁₂ arylamine, 5-8 membered monocyclic or 8-12 membered bicyclic heterocycloalkyl, or 5-8 membered monocyclic or 8-12 membered bicyclic heteroaryl;

R² and R³ are independently hydrogen, C₁₋₄ straight or branched alkyl, C₅₋₈ cycloalkyl, C₅₋₈ bicycloalkyl, C₅₋₈ aryl, 1-adamantyl, or 2-adamantyl, wherein the 1-adamantyl or 2-adamantyl is unsubstituted or substituted with one or more substituents selected from the group consisting of hydroxy, amino, aldehyde, hydroxy C₁₋₄ straight or branched alkyl, hydroxycarbonyl, aminocarbonyl, N-hydroxyaminocarbonyl, nitrile, hydroxycarbonyl C₁₋₄ straight or branched alkyl, aminocarbonyl C₁₋₄ straight

or branched alkyl, , , and ;

R⁴ is hydrogen, C₁₋₄ straight or branched alkyl, hydroxy C₁₋₄ straight or branched alkyl, or sodium;



wherein the said $R^1 - R^3$, the aryl is
 5 unsubstituted or substituted with one or more
 substituents selected from the group consisting of
 hydroxy, halogen, amino, C_{1-4} straight or branched
 alkyl, C_{1-4} straight or branched haloalkyl, C_{1-4}
 straight or branched haloalkyloxy, C_{1-4} straight or
 10 branched alkyloxy, $-O-Z^1-NZ^2Z^3$, $-O-NZ^2Z^3$, and $-Z^1-NZ^2Z^3$,
 wherein the Z_1 is C_{1-4} straight or branched alkyl, and
 the Z^2 and Z^3 are independently hydrogen or C_{1-4}
 straight or branched alkyl;

wherein the said $R^1 - R^3$, the heterocycloalkyl or
 15 heteroaryl is monocyclic or bicyclic including one or
 more heteroatoms selected from the group consisting of
 oxygen, nitrogen and sulfur, wherein the
 heterocycloalkyl or heteroaryl is unsubstituted or
 substituted with one or more substituents selected
 20 from the group consisting of trifluoro C_{1-4} straight or
 branched alkyl, pyrimidinyl, C_{1-4} straight or branched


alkyl, and C₁₋₄ straight or branched alkyloxycarbonyl,
 and also the heterocycloalkyl can be fused-bicyclic
 with C₅₋₁₂ aryl or heteroaryl, and also the heteroaryl
 can be fused-bicyclic with C₃₋₇ cycloalkyl or
 5 heterocycloalkyl;

wherein the R¹ and R⁴ can form 5-8 membered
 monocyclic or 8-12 membered bicyclic heterocycloalkyl
 with X and N which are attached to the R¹ and R⁴;

wherein the R⁴ can form 5-8 membered monocyclic
 10 heterocycloalkyl with N and A which are attached to
 the R⁴.

The present invention also provides a preparation
 method of the compound represented by formula 1
 15 containing the step of reacting the compound
 represented by formula 2 with the compound represented
 by formula 3 to prepare the compound of formula 1, as
 represented in the following reaction formula 1.

20 [Reaction Formula 1]



In reaction formula 1,

$R^1 - R^4$, X and A are as defined in formula 1.

In addition, the present invention provides a pharmaceutical composition for the prevention or
5 treatment of diseases caused by the over-activation of 11β -HSD1 enzyme which comprises the compound represented by formula 1 or the pharmaceutically acceptable salt thereof as an active ingredient.

10 **ADVANTAGEOUS EFFECT**


As explained hereinbefore, the compounds of the present invention can be efficiently used as a therapeutic agent for the treatment of the disease or
15 condition mediated by the over-activation of 11β -HSD1, such as non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder, metabolic syndrome, and other diseases mediated by the excessive activity of glucocorticoid by selectively inhibiting
20 the activity of 11β -HSD1 (11β -Hydroxysteroid dehydrogenase type 1).

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Hereinafter, the present invention is described in detail.

The present invention provides a compound represented by formula 1 having the activity of inhibiting 11 β -HSD1 enzyme or a pharmaceutically acceptable salt thereof.

[Formula 1]






In formula 1,

10 X is carbonyl or sulfonyl;








R¹ is C₁₋₄ straight or branched alkyl, C₃₋₇ cycloalkyl, C₅₋₁₂ aryl, C₅₋₁₂ arylamine, 5-8 membered monocyclic or 8-12 membered bicyclic heterocycloalkyl, or 5-8 membered monocyclic or 8-12 membered bicyclic
15 heteroaryl;

R² and R³ are independently hydrogen, C₁₋₄ straight or branched alkyl, C₅₋₈ cycloalkyl, C₅₋₈ bicycloalkyl, C₅₋₈ aryl, 1-adamantyl, or 2-adamantyl, wherein the 1-adamantyl or 2-adamantyl is unsubstituted or
20 substituted with one or more substituents selected from the group consisting of hydroxy, amino, aldehyde, hydroxy C₁₋₄ straight or branched alkyl,

hydroxycarbonyl, aminocarbonyl, N-hydroxyaminocarbonyl, nitrile, hydroxycarbonyl C₁₋₄ straight or branched alkyl, aminocarbonyl C₁₋₄ straight

or branched alkyl, , , and ;

5 R⁴ is hydrogen, C₁₋₄ straight or branched alkyl, hydroxy C₁₋₄ straight or branched alkyl, or sodium;

A is , , , , , , or , wherein Y is unsubstituted or halogen;

10 wherein the said R¹ - R³, the aryl is unsubstituted or substituted with one or more substituents selected from the group consisting of hydroxy, halogen, amino, C₁₋₄ straight or branched alkyl, C₁₋₄ straight or branched haloalkyl, C₁₋₄ straight or branched haloalkyloxy, C₁₋₄ straight or branched alkyloxy, -O-Z¹-NZ²Z³, -O-NZ²Z³, and -Z¹-NZ²Z³,
15 wherein the Z₁ is C₁₋₄ straight or branched alkyl, and the Z² and Z³ are independently hydrogen or C₁₋₄ straight or branched alkyl;

wherein the said $R^1 - R^3$, the heterocycloalkyl or heteroaryl is monocyclic or bicyclic including one or more heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, wherein the
 5 heterocycloalkyl or heteroaryl is unsubstituted or substituted with one or more substituents selected from the group consisting of trifluoro C_{1-4} straight or branched alkyl, pyrimidinyl, C_{1-4} straight or branched alkyl, and C_{1-4} straight or branched alkyloxycarbonyl,
 10 and also the heterocycloalkyl can be fused-bicyclic with C_{5-12} aryl or heteroaryl, and also the heteroaryl can be fused-bicyclic with C_{3-7} cycloalkyl or heterocycloalkyl;

wherein the R^1 and R^4 can form 5-8 membered
 15 monocyclic or 8-12 membered bicyclic heterocycloalkyl with X and N which are attached to the R^1 and R^4 ;

wherein the R^4 can form 5-8 membered monocyclic heterocycloalkyl with N and A which are attached to the R^4 .

20


Preferably,

X is carbonyl or sulfonyl;


R^1 is C_{1-4} straight or branched alkyl, C_{3-7} cycloalkyl, C_{5-6} aryl, C_{5-6} arylamine, 5-6 membered
 25 monocyclic or 9-10 membered bicyclic heterocycloalkyl,

or 5-6 membered monocyclic or 9-10 membered bicyclic heteroaryl;

R^2 and R^3 are independently hydrogen, methyl, ethyl, propyl, C_{6-7} cycloalkyl, C_{6-8} bicycloalkyl, C_{5-6} aryl, 1-adamantyl, or 2-adamantyl, wherein the 1-adamantyl or 2-adamantyl is unsubstituted or substituted with one or more substituents selected from the group consisting of hydroxy, amino, aldehyde, hydroxymethyl, hydroxyethyl, hydroxycarbonyl, aminocarbonyl, N-hydroxyaminocarbonyl, nitrile, hydroxycarbonylmethyl, hydroxycarbonylethyl,



R^4 is hydrogen, methyl, ethyl, propyl, hydroxy methyl, hydroxy ethyl, or sodium;




wherein the said $R^1 - R^3$, the aryl is unsubstituted or substituted with one or more substituents selected from the group consisting of halogen, trifluoromethyl, trifluoroethyl, trifluoromethoxy, trifluoroethoxy, methoxy, ethoxy, propoxy, methyl, ethyl, 1,1,1-trifluoro-2-ol-isopropyl, $-O(CH_2)_nNZ^2Z^3$, and $-(CH_2)_nNZ^2Z^3$, the n is 1 or 2, and the Z^2 and Z^3 are independently hydrogen, methyl or ethyl;

10 wherein the said $R^1 - R^3$, the heterocycloalkyl or heteroaryl contains one or more heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, wherein the heterocycloalkyl or heteroaryl is unsubstituted or substituted with one or more substituents selected from the group consisting of trifluoromethyl, trifluoroethyl, pyrimidinyl, methyl, ethyl, and isobutyloxycarbonyl, and also the heterocycloalkyl can be fused-bicyclic with C_{5-12} aryl or heteroaryl, and also the heteroaryl can be fused-bicyclic with C_{3-7} cycloalkyl or heterocycloalkyl;

20 wherein the R^1 and R^4 can form 5-6 membered monocyclic or 8-9 membered bicyclic heterocycloalkyl with X and N which are attached to the R^1 and R^4 ;


R^4 can form 5-6 membered monocyclic heterocycloalkyl with N and A which are attached to the R^4 .

- 5 More preferably,
X is carbonyl or sulfonyl,
 R^1 is




R^2 is $-H$ or $-(CH_2)_2CH_3$,

 $\mathbb{R}^3$ is



R^4 is $-H$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2OH$, or $-Na$,


A is



5

More preferably,

R^4 is -H or -Na,



A is .

More preferably,


5 X is sulfonyl;

R^1 is phenyl, wherein the phenyl is unsubstituted or substituted with one or more substituents selected from the group consisting of halogen and trifluoromethyl, and the halogen can be
10 substituted with one or more;

R^2 is -H;

R^3 is  or  ;

R^4 is -H or -Na,

A is .

15

Specific examples of the compound represented by formula 1 of the present invention are as follows:

(1) N-(adamantan-2-yl)-1-[(3-chloro-2-methylbenzenesulfonylamino)methyl]cyclopropanecarboxam

20 ide;

- (2) (1-[(2-fluoro-benzenesulfonylamino)methyl]-N-(5-hydroxyadamantan-2-yl)cyclopropanecarboxamide;
- (3) E-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (4) Z-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (5) E-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (6) Z-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (7) E-4-[1-((3-chloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (8) E-4-[1-((3-chloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (9) E-4-[1-((3,5-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (10) E-4-[1-((2-fluoro-6-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (11) E-4-[1-((2,3-difluoro-
5 benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (12) E-4-[1-((2,4,6-trifluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (13) E-4-[1-((2-fluoro-*N*,6-dimethyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (14) E-4-[1-((2,4-dichloro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
15 adamantan-1-carboxylic acid amide;
- (15) E-4-[1-((4-chloro-2-fluoro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (16) E-4-[1-((4,5-dichloro-2-fluoro-
20 benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (17) E-4-[1-((furan-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (18) E-4-[1-(3,5-dichloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (19) E-4-[1-((thiophen-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (20) E-4-[1-((2-(trifluoromethyl)-4-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (21) E-4-[1-((3,4-difluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (22) E-4-[1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (23) E-4-[1-((4-trifluoromethoxybenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (24) E-4-[1-((2,3-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (25) E-4-[1-((3,4-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(26) E-4-[1-((1-methyl-1H-indole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(27) E-4-[1-((1-methyl-1H-pyrazole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(28) E-4-[1-((benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

10 (29) E-4-[1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid;

(30) *N*-(bicyclo[2.2.1]heptan-2-yl)-1-((2-fluoro-*N*-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;

(31) *N*-(adamantan-1-yl)-1-((2-fluoro-*N*-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;

20 (32) E-4-[1-((*N*-ethyl-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(33) E-3-(4-(1((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido)adamantan-1-yl)propanoic acid;

(34) E-N- (5- (3-amino-3-oxopropyl) adamantan-2-yl) -
1- ((2-fluoro-
benzenesulfonylamino)methyl) cyclopropanecarboxamide;

(35) E-N- (5-aminoadamantan-2-yl) -1- ((2-fluoro-
5 benzenesulfonylamino)methyl) cyclopropanecarboxamide
hydrochloride;

(36) E-4- [1- ((2,4,5-trifluoro-
benzenesulfonylamino)methyl) cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

10 (37) E-4- [1- ((4-chloro-
benzenesulfonylamino)methyl) cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(38) E-4- [1- (3-chloro-
benzenesulfonylamino) cyclopropanecarboxamido] -
15 adamantan-1-carboxylic acid amide;

(39) E-4- [1- (2-fluoro-
benzenesulfonylamino) cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(40) E-4- [1- (2-fluoro-6-methyl-
20 benzenesulfonylamino) cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(41) E-4- [1- (3-chloro-2-methyl-
benzenesulfonylamino) cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

- (42) E-4-[1-(4-fluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (43) E-4-[1-(2,4-dichloro-5-methyl-
5 benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (44) E-4-[1-(2,4-difluorochloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (45) E-4-[1-(2-fluoro-4,5-dichloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (46) E-4-[1-(2-fluoro-4-chloro-5-methyl-
15 benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (47) E-4-[1-(2,3,4-trifluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (48) E-4-[1-(thiophen-2-
20 sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (49) E-4-[3-(6-trifluoromethyl-pyridin-2-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(50) E-4-[1-(1-methyl-1H-indole-7-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(51) 1-(3-chloro-benzenesulfonylamino)-N-(4-fluoro-2-(trifluoromethyl)phenyl)cyclopropanecarboxamide;

(52) E-4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;

10 (53) 1-(3-chloro-benzenesulfonylamino)-N-[4-(1,1,1-trifluoro-2-hydroxypropan-2-yl)phenyl]cyclopropanecarboxamide;

(54) E-4-[1-(3-chloro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(55) N-(bicyclo[2.2.1]heptan-2-yl)-1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamide;

(56) E-4-[1-(1,1-dioxido benzo[d]isothiazol-2(3H)-yl)cyclopropanecarboxamido]adamantan-1-carboxylic acid amide;

(57) E-4-[1-(3,4-difluoro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(58) E-4-[1-(1-methyl-1H-pyrazole-5-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(59) E-4-[1-(2,3-difluorobenzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(60) E-4-[1-(benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;

10 (61) E-4-[1-(2-fluoro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(62) E-4-[1-(3-chloro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(63) E-4-[1-(2-fluorobenzamido)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(64) E-N-[1-(5-carbamoyladamantan-2-yl)carbamoyl)cyclopropyl]-5-(trifluoromethyl)pyrrolineamide;

(65) E-4-(1-(benzo[d][1,3]dioxol-5-sulfonylamino)cyclopropanecarboxamino)adamantan-1-carboxylic acid amide;

- (66) 1-(3-chloro-benzenesulfonylamino)-*N*-cycloheptyl-*N*-propylcyclopropanecarboxamide;
- (67) E-4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid amide;
- 5 (68) E-4-[3-(2-fluoro-benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
- (69) E-4-[3-(benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
- (70) E-4-[3-(3-chloro-2-methyl-
10 benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
- (71) E-4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid amide;
- (72) E-4-[2-(2-fluoro-*N*-methyl-
15 benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (73) E-4-[2-(3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (74) E-*N*-(5-cyanoadamantan-2-yl)-2-(2-fluoro-
20 benzenesulfonylamino)-2-methylpropanamide;
- (75) E-2-(2-fluoro-benzenesulfonylamino)-*N*-[5-(*N'*-carbamimidoyl)adamantan-2-yl]-2-methylpropanamide;
- (76) E-2-(2-fluoro-benzenesulfonylamino)-*N*-[5-(hydroxymethyl)adamantan-2-yl]-2-methylpropanamide;

- (77) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-formyladamantan-2-yl)-2-methylpropanamide;
- (78) E-[4-(2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]methyl-4-methylbenzenesulfonate;
- (79) E-2-[4-(2-(2-fluorobenzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]acetic acid;
- (80) E-N-[5-(2-amino-2-oxoethyl)adamantan-2-yl]-2-(2-fluorobenzenesulfonylamino)-2-methylpropanamide;
- 10 (81) E-4-[2-methyl-2-(benzenesulfonylamino)propanamido]adamantan-1-carboxylic acid amide;
- (82) E-4-[2-(2-fluoro-3-chlorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- 15 (83) E-4-[2-(3,5-difluoro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (84) E-4-[2-(2,6-difluoro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- 20 (85) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide;
- (86) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide;

(87) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide;

(88) E-4-[2-(4-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;

(89) E-4-[2-(2,5-dichloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;

(90) 2-(3-chloro-benzenesulfonylamino)-N-cycloheptyl-2-methyl-N-propylpropanamide;

(91) E-4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(92) E-4-[2-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(93) E-4-[4-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(94) E-4-[3-(4-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(95) E-4-[3-(3-chloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(96) E-4-[3-(3-chloro-2-methyl-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(97) E-4-{3-[(3-chloro-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;

- (98) E-4-[3-N-(2-hydroxyethyl)-2-(trifluoromethyl-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide;
- (99) E-4-{3-[(2-trifluoromethyl-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;
- (100) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl](2-(trifluoromethyl)benzenesulfonylamide);
- 10 (101) E-N-(5-carbamoyladamantan-2-yl)-5-[(N-methyl-2-(trifluoromethyl)benzenesulfonylamino)nicotinamide];
- (102) E-4-[3-(thiophen-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 15 (103) E-4-[3-(furan-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (104) E-4-[3-(pyridin-3-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (105) E-4-(3-(benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;
- 20 (106) E-4-[3-[(2-chloro-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide;

- (107) E-4-[3-[(2,4-dimethyl-thiazol-5-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (108) E-4-[3-(3,5-dimethyl-1*H*-pyrazole-4-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (109) E-*N*-(5-hydroxy-adamantan-2-yl)-3-benzenesulfonylamino-benzamide;
- (110) E-*N*-cycloheptyl-3-phenylsulfamoyl-benzamide;
- (111) E-*N*-(5-hydroxyadamantan-2-yl)-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (112) E-4-(3-(*N*-phenylsulfamoylbenzamino)-adamantan-1-carboxylic acid amide;
- (113) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl]-2-fluoro-3-chloro-benzenesulfonylamide;
- (114) E-4-[3-(3-chloro-4-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (115) E-4-[3-[(2-(trifluoromethyl)benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

- (116) E-4-[3-(2-chloro-4-bromo-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (117) E-4-[3-(2,4,6-trichloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (118) E-4-[3-(3-chloro-5-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 10 (119) E-4-[3-(3,5-dichloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (120) E-4-[3-(3-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (121) E-4-[3-(2,4-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 15 (122) E-4-[3-(2,5-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (123) E-4-[3-(2,6-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 20 (124) E-N-cycloheptyl-N-propyl-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (125) E-4-[2-fluoro-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

- (126) E-4-[2-chloro-5-(3-chloro-benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (127) E-4-[3-(3-chlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (128) E-4-[4-chloro-3-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (129) E-4-[2-chloro-5-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (130) E-4-[3-(3,5-dichlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (131) E-4-[5-(3,5-dichlorobenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (132) E-4-[2-fluoro-3-(3-chlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (133) E-4-[2-chloro-5-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (134) E-4-[4-chloro-3-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

- (135) E-4-[5-(3-chloro-4-methoxybenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (136) E-4-(3-(4-chloro-benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;
- (137) E-4-(3-(4-chloro-benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;
- (138) E-4-(3-(cyclopropanesulfonamido)benzamido)adamantan-1-carboxylic acid amide;
- (139) E-4-(3-(1-methylethylsulfonamido)benzamido)adamantan-1-carboxylic acid amide;
- (140) E-[3,4-dihydro-1H-isoquinolin-2-carboxylic acid-1-[(5-carbamoyl-adamantan-2-ylcarbamoyl)cyclopropylmethyl]]-amide;
- (141) E-3,4-dihydro-2H-quinolin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;
- (142) E-piperidin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;
- (143) E-4-{[1-5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-carbamoyl}-3,4-dihydro-2H-quinolin-1-carboxylic acid-butylester;

(144) E-4-pyrimidin-2-yl-piperazin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;

(145) E-4-({1[(3-phenyl-ureido)methyl]-cyclopropanecarbonyl}amino)-adamantan-1carboxylic acid amide;

(146) E-3,4-dihydro-2H-quinoxalin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;

10 (147) E-3,4-dihydro-1H-isoquinolin-2-carboxylic acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide;

(148) E-3,4-dihydro-2H-quinolin-1-carboxylic acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide; and

15 (149) E-2-(2-fluoro-benzamido)-thiazol-4-carboxylic acid(5-carbamoyl-adamantan-2-yl)amide.

Preferable examples of the compound represented by formula 1 of the present invention are as follows:

(1) N-(adamantan-2-yl)-1-[(3-chloro-2-methylbenzenesulfonylamino)methyl]cyclopropanecarboxamide;

(2) (1-[(2-fluoro-benzenesulfonylamino)methyl]-N-(5-hydroxyadamantan-2-yl)cyclopropanecarboxamide;

- (3) E-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (4) Z-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (5) E-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (6) Z-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (7) E-4-[1-((3-chloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (8) E-4-[1-((3-chloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (9) E-4-[1-((3,5-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (10) E-4-[1-((2-fluoro-6-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (11) E-4-[1-((2,3-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (12) E-4-[1-((2,4,6-trifluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (13) E-4-[1-((2-fluoro-N,6-dimethyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (14) E-4-[1-((2,4-dichloro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (15) E-4-[1-((4-chloro-2-fluoro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (16) E-4-[1-((4,5-dichloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (17) E-4-[1-((furan-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (18) E-4-[1-(3,5-dichloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (19) E-4-[1-((thiophen-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (20) E-4-[1-((2-(trifluoromethyl)-4-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (21) E-4-[1-((3,4-difluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (22) E-4-[1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (23) E-4-[1-((4-trifluoromethoxybenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (24) E-4-[1-((2,3-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (25) E-4-[1-((3,4-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (26) E-4-[1-((1-methyl-1H-indole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(27) E-4-[1-((1-methyl-1H-pyrazole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(28) E-4-[1-((benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(29) E-4-[1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid;

10 (30) N-(bicyclo[2.2.1]heptan-2-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;

(31) N-(adamantan-1-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;

(32) E-4-[1-((N-ethyl-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

20 (33) E-3-(4-(1((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido)adamantan-1-yl)propanoic acid;

(34) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamide;

(35) E-N- (5-aminoadamantan-2-yl) -1- ((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamide hydrochloride; and

(36) E-4- [1- ((2,4,5-trifluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
5 adamantan-1-carboxylic acid amide.

Other preferable examples of the compound represented by formula 1 of the present invention are
10 as follows:

(37) E-4- [1- ((4-chloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(38) E-4- [1- (3-chloro-benzenesulfonylamino)cyclopropanecarboxamido] -
15 adamantan-1-carboxylic acid amide;

(39) E-4- [1- (2-fluoro-benzenesulfonylamino)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

20 (40) E-4- [1- (2-fluoro-6-methyl-benzenesulfonylamino)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(41) E-4- [1- (3-chloro-2-methyl-benzenesulfonylamino)cyclopropanecarboxamido] -
25 adamantan-1-carboxylic acid amide;

- (42) E-4-[1-(4-fluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (43) E-4-[1-(2,4-dichloro-5-methyl-
5 benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (44) E-4-[1-(2,4-difluorochloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (45) E-4-[1-(2-fluoro-4,5-dichloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (46) E-4-[1-(2-fluoro-4-chloro-5-methyl-benzenesulfonylamino)cyclopropanecarboxamido]-
15 adamantan-1-carboxylic acid amide;
- (47) E-4-[1-(2,3,4-trifluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (48) E-4-[1-(thiophen-2-
20 sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (49) E-4-[3-(6-trifluoromethyl-pyridin-2-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(50) E-4-[1-(1-methyl-1H-indole-7-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(51) 1-(3-chloro-benzenesulfonylamino)-N-(4-fluoro-2-(trifluoromethyl)phenyl)cyclopropanecarboxamide;

(52) E-4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;

10 (53) 1-(3-chloro-benzenesulfonylamino)-N-[4-(1,1,1-trifluoro-2-hydroxypropan-2-yl)phenyl]cyclopropanecarboxamide;

(54) E-4-[1-(3-chloro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(55) N-(bicyclo[2.2.1]heptan-2-yl)-1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamide;

(56) E-4-[1-(1,1-dioxido benzo[d]isothiazol-2(3H)-yl)cyclopropanecarboxamido]adamantan-1-carboxylic acid amide;

(57) E-4-[1-(3,4-difluoro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(58) E-4-[1-(1-methyl-1H-pyrazole-5-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(59) E-4-[1-(2,3-difluorobenzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(60) E-4-[1-(benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;

10 (61) E-4-[1-(2-fluoro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(62) E-4-[1-(3-chloro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(63) E-4-[1-(2-fluorobenzamido)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(64) E-N-[1-(5-carbamoyladamantan-2-yl)carbamoyl)cyclopropyl]-5-(trifluoromethyl)pyrrolineamide;

(65) E-4-(1-(benzo[d][1,3]dioxol-5-sulfonylamino)cyclopropanecarboxamino)adamantan-1-carboxylic acid amide; and

(66) 1-(3-chloro-benzenesulfonylamino)-*N*-cycloheptyl-*N*-propylcyclopropanecarboxamide.

5 Preferable examples of the compound represented by formula 1 of the present invention are as follows:

(67) E-4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid amide;

(68) E-4-[3-(2-fluoro-benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;

10 (69) E-4-[3-(benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
and

(70) E-4-[3-(3-chloro-2-methyl-
15 benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide.

Preferable examples of the compound represented by formula 1 of the present invention are as follows:

20 (71) E-4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid amide;

(72) E-4-[2-(2-fluoro-*N*-methyl-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;

- (73) E-4-[2-(3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (74) E-N-(5-cyanoadamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide;
- 5 (75) E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(N'-carbamimidoyl)adamantan-2-yl]-2-methylpropanamide;
- (76) E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(hydroxymethyl)adamantan-2-yl]-2-methylpropanamide;
- (77) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-
10 formyladamantan-2-yl)-2-methylpropanamide;
- (78) E-[4-(2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]methyl-4-methylbenzenesulfonate;
- (79) E-2-[4-(2-(2-fluorobenzenesulfonylamino)-2-
15 methylpropanamido)adamantan-1-yl]acetic acid;
- (80) E-N-[5-(2-amino-2-oxoethyl)adamantan-2-yl]-2-(2-fluorobenzenesulfonylamino)-2-methylpropanamide;
- (81) E-4-[2-methyl-2-(benzenesulfonylamino)propanamido]adamantan-1-
20 carboxylic acid amide;
- (82) E-4-[2-(2-fluoro-3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (83) E-4-[2-(3,5-difluoro-benzenesulfonylamino)-
25 2-methylpropanamido]adamantan-1-carboxylic acid amide;

(84) E-4-[2-(2,6-difluoro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;

(85) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-

5 methylpropanamide;

(86) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide;

(87) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-

10 methylpropanamide;

(88) E-4-[2-(4-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;

(89) E-4-[2-(2,5-dichloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;

15 and

(90) 2-(3-chloro-benzenesulfonylamino)-N-cycloheptyl-2-methyl-N-propylpropanamide.

Preferable examples of the compound represented
20 by formula 1 of the present invention are as follows:

(91) E-4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(92) E-4-[2-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

- (93) E-4-[4-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (94) E-4-[3-(4-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 5 (95) E-4-[3-(3-chloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (96) E-4-[3-(3-chloro-2-methyl-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 10 (97) E-4-{3-[(3-chloro-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;
- (98) E-4-[3-N-(2-hydroxyethyl)-2-(trifluoromethyl-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide;
- 15 (99) E-4-{3-[(2-trifluoromethyl-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;
- (100) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl](2-(trifluoromethyl)benzenesulfonylamide;
- 20 (101) E-N-(5-carbamoyladamantan-2-yl)-5-[(N-methyl-2-(trifluoromethyl)benzenesulfonylamino]nicotinamide;

- (102) E-4-[3-(thiophen-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (103) E-4-[3-(furan-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 5 (104) E-4-[3-(pyridin-3-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (105) E-4-(3-(benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;
- (106) E-4-[3-[(2-chloro-10 benzenesulfonylamino]benzamido]-adamantan-1-carboxylic acid amide;
- (107) E-4-[3-[(2,4-dimethyl-thiazol-5-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 15 (108) E-4-[3-(3,5-dimethyl-1*H*-pyrazole-4-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (109) E-*N*-(5-hydroxy-adamantan-2-yl)-3-benzenesulfonylamino-benzamide;
- 20 (110) E-*N*-cycloheptyl-3-phenylsulfamoyl-benzamide;
- (111) E-*N*-(5-hydroxyadamantan-2-yl)-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (112) E-4-(3-(*N*-phenylsulfamoylbenzamino)-25 adamantan-1-carboxylic acid amide;

(113) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl]-2-fluoro-3-chloro-benzenesulfonylamide;

(114) E-4-[3-(3-chloro-4-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

(115) E-4-[3-[(2-(trifluoromethyl)benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

10 (116) E-4-[3-(2-chloro-4-bromobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(117) E-4-[3-(2,4,6-trichlorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(118) E-4-[3-(3-chloro-5-fluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(119) E-4-[3-(3,5-dichlorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(120) E-4-[3-(3-fluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(121) E-4-[3-(2,4-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(122) E-4-[3-(2,5-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(123) E-4-[3-(2,6-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

5 (124) E-N-cycloheptyl-N-propyl-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;

(125) E-4-[2-fluoro-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

10 (126) E-4-[2-chloro-5-(3-chloro-benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

(127) E-4-[3-(3-chlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;

15 (128) E-4-[4-chloro-3-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

(129) E-4-[2-chloro-5-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

20

(130) E-4-[3-(3,5-dichlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;

(131) E-4-[5-(3,5-dichlorobenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;

- (132) E-4-[2-fluoro-3-(3-chloro-benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (133) E-4-[2-chloro-5-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (134) E-4-[4-chloro-3-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- 10 (135) E-4-[5-(3-chloro-4-methoxybenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (136) E-4-(3-(4-chloro-benzenesulfonylamino)benzamido)-adamantan-1-carboxylic acid amide;
- 15 (137) E-4-(3-(4-chloro-benzenesulfonylamino)benzamido)-adamantan-1-carboxylic acid amide;
- (138) E-4-(3-(cyclopropanesulfonamido)benzamido)adamantan-1-carboxylic acid amide; and
- 20 (139) E-4-(3-(1-methylethylsulfonamido)benzamido)adamantan-1-carboxylic acid amide.

Preferable examples of the compound represented
 25 by formula 1 of the present invention are as follows:

(140) E-[3,4-dihydro-1H-isoquinolin-2-carboxylic
acid-1-[(5-carbamoyl-adamantan-2-
ylcarbamoyl)cyclopropylmethyl]-amide;

(141) E-3,4-dihydro-2H-quinolin-1-carboxylic
5 acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-
cyclopropylmethyl]-amide;

(142) E-piperidin-1-carboxylic acid[1-(5-
carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-
amide;

10 (143) E-4-{[1-(5-carbamoyl-adamantan-2-
ylcarbamoyl)-cyclopropylmethyl]-carbamoyl}-3,4-
dihydro-2H-quinolin-1-carboxylic acid-butylester;

(144) E-4-pyrimidin-2-yl-piperazin-1-carboxylic
acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-
15 cyclopropylmethyl]-amide;

(145) E-4-({1[(3-phenyl-ureido)methyl]-
cyclopropanecarbonyl}amino)-adamantan-1-carboxylic acid
amide;

(146) E-3,4-dihydro-2H-quinoxalin-1-carboxylic
20 acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-
cyclopropylmethyl]-amide;

(147) E-3,4-dihydro-1H-isoquinolin-2-carboxylic
acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-
2-yl]-amide;

(148) E-3,4-dihydro-2H-quinolin-1-carboxylic acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide; and

(149) E-2-(2-fluoro-benzamido)-thiazol-4-carboxylic acid(5-carbamoyl-adamantan-2-yl)amide.

The compound represented by formula 1 of the present invention can be used as a form of a pharmaceutically acceptable salt, in which the salt is preferably acid addition salt formed by pharmaceutically acceptable free acids. Herein, the pharmaceutically acceptable salt indicates any organic or inorganic addition salt of the base compound represented by formula 1 that is relatively nontoxic to a patient and has non-harmful activity whose side effect cannot reduce any positive effect of the said base compound represented by formula 1. The acid addition salt herein can be obtained from inorganic acids such as hydrochloric acid, bromic acid, nitric acid, sulfuric acid, perchloric acid, and phosphoric acid; or organic acids such as citric acid, acetic acid, lactic acid, maleic acid, fumaric acid, gluconic acid, methanesulfonic acid, glyconic acid, succinic acid, tartaric acid, galacturonic acid, embonic acid, glutamic acid, aspartic acid, oxalic acid, (D) or (L)

malic acid, methanesulfonic acid, ethanesulfonic acid, 4-toluenesulfonic acid, salicylic acid, citric acid, benzoic acid, and malonic acid. The salt herein includes alkali metal salt (sodium salt, potassium salt, etc) and alkali earth metal salt (calcium salt, magnesium salt, etc). For example, the acid addition salt is exemplified by acetate, aspartate, benzoate, besylate, bicarbonate/carbonate, bisulfate/sulfate, borate, camcilate, citrate, edisylate, ethylate, formate, fumarate, gluceptate, gluconate, glucuronate, hexafluorophosphate, hibenzate, hydrochloride/chloride, hydrobromide/bromide, hydroiodide/iodide, isethionate, lactate, malate, maleate, malonate, methylate, methylsulfate, naphthalate, 2-naphthylate, nicotinate, nitrate, orotate, oxalate, palmitate, pamoate, phosphate/hydrogen phosphate/dihydrogen phosphate, saccharate, stearate, succinate, tartrate, tosylate, trifluoroacetate, aluminum, arginine, benzathine, calcium, choline, diethylamine, diolamine, glycine, lysine, magnesium, meglumine, olamine, potassium, sodium, tromethamine, and zinc salt. Among them, hydrochloride or trifluoroacetate is preferred.

The compound represented by formula 1 of the present invention not only includes the pharmaceutically acceptable salts but also includes every possible salts, isomers, hydrates, and solvates
5 constructed from the same by the conventional method.

The acid addition salt in this invention can be prepared by the conventional method known to those in the art. For example, the compound of formula 1 of the present invention is dissolved in a water-miscible
10 organic solvent such as acetone, methanol, ethanol, or acetonitrile, to which excessive organic acid or acid aqueous solution of inorganic acid is added to induce precipitation or crystallization. Then, the solvent or the excessive acid is evaporated from the mixture,
15 followed by drying the mixture to give addition salt or suction-filtering the precipitated salt to give the same.

Some compounds among those represented by formula
20 1 contain chiral center or geometrical isomer center (E and Z isomers). It is understood that the present invention includes all the optical isomers, diastereomers, and geometrical isomers having the activity of inhibiting 11 β -HSD1.

The present invention also provides a preparation method of the compound represented by formula 1 containing the step of reacting the compound represented by formula 2 with the compound represented by formula 3 to prepare the compound of formula 1, as represented in the following reaction formula 1.

$$\begin{array}{c} \text{R}^1\text{-X-Cl} \\ \text{2} \end{array} + \begin{array}{c} \text{H-N-A-C(=O)-N(R}^2\text{)(R}^3\text{)} \\ | \\ \text{R}^4 \\ \text{3} \end{array} \longrightarrow \begin{array}{c} \text{R}^1\text{-X-N(R}^4\text{)-A-C(=O)-N(R}^2\text{)(R}^3\text{)} \\ \text{1} \end{array}$$

In reaction formula 1,

58

In the preparation method of the present invention, the compound represented by formula 2 can be a sulfonylhalide derivative or an acetylhalide derivative. The compounds of formula 2 and formula 3
5 can be the commercial compounds on the market or can be synthesized by those of knowledge in the art, considering the substituents therein.

Particularly, the compound of formula 1 of the present invention can be easily prepared by reacting
10 the amine derivative compound represented by formula 3 with the sulfonylhalide derivative or the acetylhalide derivative represented by formula 2 in the presence of a general organic solvent like dichloromethane with the addition of a proper amount of
15 diisopropylethylamine. At this time, the reaction temperature and the reaction time can vary with the chemical reactivity of the sulfonylhalide derivative or the acetylhalide derivative represented by formula 2, but the reaction for 1 - 24 hours at room
20 temperature is preferred, but not always limited thereto.

After the compound of formula 1 was prepared by the method described above, the molecular structure of the compound was confirmed by infrared spectroscopy
25 (IR), nuclear magnetic resonance spectrum (NMR), mass

spectrometry (Mass), liquid chromatography, X-ray crystallography, polarimetry, and the comparison of the estimated data and the analyzed data of the representative compound.

5

In addition, the present invention provides a pharmaceutical composition for the prevention or treatment of diseases caused by the over-activation of 11 β -HSD1 enzyme which comprises the compound
10 represented by formula 1 or the pharmaceutically acceptable salt thereof as an active ingredient.

The compound represented by formula 1 was confirmed to have excellent 11 β -HSD1 inhibiting activity. Precisely, IC₅₀ of the compound, which was
15 the concentration of the compound to inhibit the 11 β -HSD1 activity 50%, was 0.02 - 672 nM.

Therefore, since the compound of formula 1 of the present invention has excellent 11 β -HSD1 inhibiting activity, it can be effectively used for the
20 prevention or treatment of diseases mediated by abnormally activated 11 β -HSD1, such as non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder, metabolic syndrome, and other diseases mediated by the excessive activity of
25 glucocorticoid.

The pharmaceutical composition of the present invention comprising the compound represented by formula 1 or the pharmaceutically acceptable salt thereof as an active ingredient can be administered orally or parenterally and be used in general forms of pharmaceutical formulation, but not always limited thereto

The formulations for oral administration are exemplified by tablets, pills, hard/soft capsules, solutions, suspensions, emulsions, syrups, granules, and elixirs, etc. These formulations can include diluents (for example, lactose, dextrose, sucrose, mannitol, sorbitol, cellulose, and/or glycine) and lubricants (for example, silica, talc, stearate and its magnesium or calcium salt, and/or polyethylene glycol) in addition to the active ingredient. Tablets can include binding agents such as magnesium aluminum silicate, starch paste, gelatin, methylcellulose, sodium carboxymethylcellulose and/or polyvinylpyrrolidone, and if necessary disintegrating agents such as starch, agarose, alginic acid or its sodium salt or azeotropic mixtures and/or absorbents, coloring agents, flavors, and sweeteners can be additionally included thereto.


The effective dosage of the compound of the present invention can be determined according to age, weight, gender, administration method, health condition, and severity of disease. For example, the
 5 dose for an adult in 70 kg of body weight is 0.1 - 1,000 mg/day and preferably 1 - 500 mg/day. This administration can be performed once a day - several times a day according to the decision of a doctor or a pharmacist.

10

Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those
 15 skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.


Preparational Example 1: Preparation of 1-
 20 (aminomethyl)-cyclopropylacetic acid hydrochloride



200 mg (1.75 mmol) of 1-cyano-1-cyclopropylacetic acid was dissolved in 10 ml of methanol, to which 40 mg (20 wt%) of platinum oxide was added. 0.3 ml of concentrated hydrochloric acid solution was loaded thereto, and the mixture was stirred at room temperature for 4 hours in hydrogen stream. Upon completion of the reaction, the mixture was filtered to eliminate platinum oxide. The solvent was evaporated under reduced pressure to give 256 mg of 1-(aminomethyl)-cyclopropylacetic acid hydrochloride (yield: 97%).

^1H NMR (400 MHz, MeOD) δ 3.13 (s, 2H), 1.36-1.38 (m, 2H), 1.16-1.18 (m, 2H)

15 Preparational Example 2: Preparation of 1-(aminomethyl)-cyclopropylcarboxylic acid methylester hydrochloride




273 mg (1.8 mmol) of the 1-(aminomethyl)-cyclopropylacetic acid hydrochloride prepared in Preparational Example 1 was dissolved in 6 ml of methanol, to which 0.26 ml (3.6 mmol) of thionyl

chloride was added, followed by stirring at room temperature for 3 hours. Upon completion of the reaction, the solvent was evaporated under reduced pressure to give 291 mg of 1-(aminomethyl)-
 5 cyclopropylcarboxylic acid methylester hydrochloride (yield: 98%).

^1H NMR (400 MHz, D_2O) δ 3.60 (s, 3H), 3.06 (s, 2H), 1.28-1.31 (m, 2H), 0.97-0.99 (m, 2H)

10 Preparational Example 3: Preparation of 1-[(2-fluoro-benzenesulfonylamino)methyl]-cyclopropylcarboxylic acid methyl ester




150 (0.91 mmol) mg of the 1-(aminomethyl)-
 15 cyclopropylcarboxylic acid methylester hydrochloride prepared in Preparational Example 2 was dissolved in 3 ml of methylene chloride, to which 0.25 ml of TEA was added, followed by stirring at room temperature for 5 minutes. 194 mg (1.0 mmol) of 2-
 20 fluorosulfonylchloride was added to the reactant, followed by stirring at room temperature for 3 hours, followed by evaporation under reduced pressure to

eliminate the reaction solvent. The residue was dissolved in 5 ml of water, followed by extraction with 310 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 215 mg of 1-[(2-fluorobenzenesulfonylamino)methyl]-cyclopropylcarboxylic acid methyl ester (yield: 83%).

^1H NMR (400 MHz, CDCl_3) δ 7.86-7.92 (m, 1H), 7.56-7.60 (m, 1H), 7.20-7.32 (m, 2H), 5.67 (d, $J = 5.2$ Hz, 1H), 3.64-3.72 (m, 3H), 3.14-3.21 (m, 2H), 1.17-1.21 (m, 2H), 0.84-0.91 (m, 2H)

Preparational Example 4: Preparation of 1-[(2-fluorobenzenesulfonylamino)methyl]-cyclopropylcarboxylic acid




215 mg (0.75 mmol) of the 1-[(2-fluorobenzenesulfonylamino)methyl]-cyclopropylcarboxylic acid methyl ester prepared in Preparational Example 3 was dissolved in 7.5 ml of the mixed solvent of distilled water and THF (1:2), to which 94 mg (2.25 mmol) of lithium hydroxide monohydrate was added,

followed by stirring at room temperature for 16 hours. Upon completion of the reaction, the reaction solvent was eliminated by evaporation under reduced pressure and the residue was acidized with 1 N HCl to be pH 3, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 200 mg of 1-[(2-fluorobenzenesulfonylamino)methyl]-cyclopropylcarboxylic acid (yield: 98%).

^1H NMR (400 MHz, CDCl_3) δ 7.85-7.90 (m, 1H), 7.56-7.61 (m, 1H), 6.98-7.31 (m, 2H), 5.90 (t, $J = 6.2$ Hz, 1H), 3.15-3.17 (m, 2H), 1.29-1.32 (m, 2H), 0.93-0.97 (m, 2H)

15

Preparational Example 5: Preparation of 1-[(t-butoxycarbonylamino)methyl]cyclopropyl acetic acid




256 mg (1.69 mmol) of the 1-(aminomethyl)-cyclopropylacetic acid hydrochloride prepared in Preparational Example 1 was dissolved in 3 ml of methylene chloride, to which 0.26 ml (1.86 mmol) of TEA was added, followed by stirring for 10 minutes.

Methylene chloride was eliminated by evaporation under reduced pressure. The reactant was dissolved in 2 ml of 1.0 N NaOH and 6 ml of 1,4-dioxane, and then 443 mg (2.0 mmol) of Boc_2O was added thereto. After stirring
 5 the reaction mixture at room temperature for 16 hours, 1,4-dioxane was eliminated by evaporation under reduced pressure. The residue was acidized with 1 N HCl to be pH 3, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous
 10 magnesium sulfate, followed by evaporation under reduced pressure to give 345 mg of 1-[(t-butoxycarbonylamino)methyl]cyclopropyl acetic acid (yield: 95%).

^1H NMR (400 MHz, MeOD) δ 1.51 (s, 2H), 1.43 (s, 9H), 1.13-1.16 (m, 2H), 0.89-0.91 (m, 2H)
 15

Preparational Example 6: Preparation of 4-oxo-adamantan-1-carboxylic acid methyl ester




20 30% fuming sulfuric acid solution was heated at 60°C, to which 1 g (6.02 mmol) of 5-hydroxy-2-adamantanone dissolved in 6 ml of 99% formic acid was slowly added over 1 hour. After adding 6 ml of 99%

formic acid slowly for 1 hour, the mixture was stirred at 60°C for 1 hour. The reaction solution was slowly added to 50 ml of methanol cooled down to 0°C, followed by stirring at room temperature for 2 hours. Then, the reaction solution was evaporated under reduced pressure. 15 g of ice and 50 ml of methylene chloride were added thereto, followed by extraction with methylene chloride twice. After washing with brine, the extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 1.09 g of 4-oxo-adamantan-1-carboxylic acid methyl ester (yield: 87%).

^1H NMR (400 MHz, CDCl_3) δ 3.69 (s, 3H), 2.19 (m, 2H), 1.98-2.24 (m, 11H)

15

Preparational Example 7: Preparation of 4-amino-adamantan-1-carboxylic acid methyl ester




1.09 g (5.234 mmol) of the 4-oxo-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 6 and 0.5 g of 4Å molecular sieves were dissolved in 9.3 ml of methanolic ammonia (7N). The mixture was stirred at room temperature overnight,

followed by cooling down to 0°C. NaBH₄ was slowly
 added thereto, followed by stirring at room
 temperature for 2 hours. The floating materials
 generated therein were eliminated by filtering. The
 5 solvent was eliminated by evaporation under reduced
 pressure. The residue was dissolved in 100 ml of
 methylene chloride, followed by acidization with 10%
 citric acid. Then, the reactant was neutralized with
 NaHCO₃ solution, followed by washing with brine. The
 10 reactant was extracted twice with methylene chloride.
 The extract was dried over anhydrous magnesium
 sulfate, followed by evaporation under reduced
 pressure to give 800 mg of 4-amino-adamantan-1-
 carboxylic acid methyl ester (yield: 73%).

15 ¹H NMR (400 MHz, CDCl₃) δ 3.67 (s, 3H), 3.03 (m,
 1H), 1.48-2.02 (m, 13H)

Preparational Example 8: Preparation of t-butyl[(1-
 (adamantan-2-ylcarbamoyl)cyclopropyl)methyl]carbamate




20

33 mg (0.15 mmol) of the 1-[(t-
 butoxycarbonylamino)methyl]cyclopropyl acetic acid
 prepared in Preparational Example 5 and 32 mg (0.17

mmol) of 2-admantaneamine hydrochloride were dissolved in 1.5 ml of methylene chloride, to which 44 mg (0.17 mmol) of BOP-Cl was added. 0.04 ml (0.31 mmol) of TEA was also added thereto. The reaction mixture was stirred at room temperature for 16 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 5 ml of water, followed by extraction with 310 ml of methylene chloride. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 34 mg of t-butyl[(1-(adamantan-2-ylcarbamoyl)cyclopropyl)methyl]carbamate (yield: 64%).

^1H NMR (400 MHz, CDCl_3) δ 7.02 (brs, 1H), 4.89 (s, 1H), 4.03 (d, J = 7.2 Hz, 1H), 3.37 (d, J = 6.8 Hz, 2H), 1.49-2.05 (m, 14 H), 1.44 (s, 9H), 1.23-1.26 (m, 2H), 0.60-0.66 (m, 2H)


Preparational Example 9: Preparation of 4-[1-(t-butoxycarbonylamino)methyl-cyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester



391 mg (1.82 mmol) of the 1-[(t-butoxycarbonylamino)methyl]cyclopropyl acetic acid prepared in Preparational Example 5 and 492 mg (2.00 mmol) of the 4-amino-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 7 were dissolved in 6.0 ml of methylene chloride, to which 525 mg (2.00 mmol) of BOP-Cl was added. 0.5 ml (3.64 mmol) of TEA was also added thereto. The reaction mixture was stirred at room temperature for 16 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 400 mg of 4-[1-(t-butoxycarbonylamino)methylcyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester (yield: 54%).

^1H NMR (400 MHz, CDCl_3) δ 7.16 (brs, 1H), 4.87 (t, $J = 5.8$ Hz, 1H), 4.01 (d, $J = 6.8$ Hz, 1H), 3.65 (s, 3H), 3.37 (d, $J = 6.4$ Hz, 2H), 1.94-2.05 (m, 9H), 1.89 (s, 2H), 1.54 (s, $J = 12.4$ Hz, 2H), 1.42 (s, 9H), 1.25-1.28 (m, 2H), 0.63-0.65 (m, 2H)


Preparational Example 10: Preparation of 4-[1-(aminomethyl)cyclopropanylcarboxamido]-adamantan-1-carboxylic acid methyl ester hydrochloride



5 4.0 g (9.8 mmol) of the 4-[1-(t-butoxycarbonylamino)methyl-cyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 9 was dissolved in 33 ml of ethyl acetate, to which 25 ml of 4 M HCl 1,4-dioxane
10 solution was added. The reaction mixture was stirred at room temperature for 6 hours. Upon completion of the reaction, the reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in 50 ml of ethylene acetate and the
15 precipitated solid compound was filtered. As a result, 3.2 g of 4-[1-(aminomethyl)cyclopropanylcarboxamido]-adamantan-1-carboxylic acid methyl ester hydrochloride was obtained (yield: 92%).

20 ¹H NMR (400 MHz, MeOD) δ 3.90 (s, 1H), 3.64 (s, 3H), 3.08 (s, 2H), 1.90-2.05 (m, 11H), 1.55 (s, J=12.4 Hz, 2H), 1.35-1.38 (s, 2H), 1.05-1.11 (m, 2H)

Preparational Example 11: Preparation of 4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester



5


1.0 g (3.2 mmol) of the 4-[1-(aminomethyl)cyclopropanylcarboxamido]-adamantan-1-carboxylic acid methyl ester hydrochloride prepared in Preparational Example 10 was dissolved in 10 ml of methylene chloride, to which 0.8 ml of TEA was added, followed by stirring at room temperature for 5 minutes. 624 mg (3.2 mmol) of 2-fluorosulfonylchloride was added to the reaction mixture, followed by stirring at room temperature for 6 hours. The reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in 20 ml of water, followed by extraction with 350 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 1.5 g of 4-[1-((2-fluoro-

benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid methyl ester (yield: 99%).

^1H NMR (400 MHz, CDCl_3) δ 7.90 (dt, J = 7.6, 1.7 Hz, 1H), 7.59-7.64 (m, 1H), 7.32 (dt, J = 7.6, 1.0 Hz, 1H), 7.21-7.24 (m, 1H), 6.53 (d, J = 7.6 Hz, 1H), 5.28 (t, J = 6.6 Hz, 1H), 4.05 (d, J = 7.2 Hz, 1H), 3.66 (s, 3H), 3.12 (d, J = 6.4 Hz, 2H), 1.91-2.05 (m, 11H), 1.47 (d, J = 12.0 Hz, 2H), 1.21-1.23 (m, 2H), 0.67-0.70 (m, 2H)

10

Preparational Example 12: Preparation of 4-[1-((2-fluoro-
benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid




15

1.5 g (3.23 mmol) of the 4-[1-((2-fluoro-
benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid methyl ester prepared in
Preparational Example 11 was dissolved in 20 ml of
20 THF/ethanol mixed solution (1:1), to which 8.0 ml of
2N NaOH solution was added, followed by stirring at
room temperature overnight. The residue was acidized

with 1 N HCl, followed by extraction with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 1.44 g of 4-[1-((2-fluoro-
 5 benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid (yield: 99%).

^1H NMR (400 MHz, MeOD) δ 7.89 (dt, J = 7.4, 1.6 Hz, 1H), 7.65-7.70 (m, 1H), 7.46 (d, J = 7.2 Hz, 1H), 7.31-7.39 (m, 2H), 4.00 (d, J = 7.6 Hz, 1H), 3.15 (s, 2H),
 10 2H), 2.16 (d, J = 13.2 Hz, 2H), 1.98-2.03 (m, 7H), 1.94 (s, 2H), 1.60 (d, J = 13.2 Hz, 2H), 1.13-1.16 (m, 2H), 0.69-0.72 (m, 2H)

Example 1: Preparation of N-(adamantan-2-yl)-1-[(3-
 15 chloro-2-methylbenzenesulfonylamino)methyl]cyclopropanecarboxamide




34 mg (0.09 mmol) of the t-butyl[(1-(adamantan-2-ylcarbamoyl)cyclopropyl)methyl]carbamate prepared in
 20 Preparational Example 8 was dissolved in 1.0 ml of ethyl acetate, to which 0.12 ml of 4 M HCl 1,4-dioxane solution was added, followed by stirring at room

temperature for 3 hours. Upon completion of the reaction, the reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in 10 ml of ethylene acetate and the precipitated solid compound was filtered and dried. This solid compound was dissolved in 2 ml of methylene chloride, to which 0.03 ml of TEA was added, followed by stirring at room temperature for 5 minutes. 24 mg (0.10 mmol) of 3-chloro-2-methylbenzenesulfonylchloride was added to the reactant, which was then stirred at room temperature for 4 hours. The reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 16 mg of *N*-(adamantan-2-yl)-1-[(3-chloro-2-methylbenzenesulfonylamino)methyl]cyclopropanecarboxamide (yield: 39%).

^1H NMR (400 MHz, CDCl_3) δ 7.90 (dd, J = 8.0, 1.2 Hz, 1H), 7.58 (dd, J = 8.0, 0.8 Hz, 1H), 7.26 (t, J = 8.0 Hz, 1H), 5.96 (d, J = 7.6 Hz, 1H), 5.62 (t, J = 6.2 Hz, 1H), 4.01 (d, J = 7.6 Hz, 1H), 3.04 (d, J =

6.0 Hz, 2H), 2.72 (s, 3H), 1.66-1.89 (m, 14H), 1.07-1.09 (m, 2H), 0.86-0.88 (m, 2H)


Example 2: Preparation of 1-[(2-fluorobenzenesulfonylamino)methyl]-N-(5-hydroxyadamantan-2-yl)cyclopropanecarboxamide



43 mg (0.16 mmol) of the 1-[(2-fluorobenzenesulfonylamino)methyl]-cyclopropylcarboxylic acid prepared in Preparational Example 4 was dissolved in 1.6 ml of methylene chloride, to which 44 mg (0.32 mmol) of BOP-Cl, 29 mg (0.17 mmol) of 4-aminoadamantane-1-ol and 0.04 ml (0.32 mmol) of TEA were added. The reaction mixture was stirred at room temperature for 4 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 5 ml of water, followed by extraction with 310 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 9 mg of 1-[(2-fluorobenzenesulfonylamino)methyl]-N-(5-hydroxyadamantan-2-yl)cyclopropanecarboxamide (yield: 14%).

^1H NMR (400 MHz, CDCl_3) δ 7.88-7.92 (m, 1H), 7.59-7.64 (m, 1H), 7.21-7.33 (m, 2H), 6.48 (d, J = 6.8 Hz, 1H), 5.29 (t, J = 6.4 Hz, 1H), 4.04 (t, J = 4.0 Hz, 1H), 3.12 (s, J = 6.4 Hz, 2H), 2.12-2.19 (m, 3H),
 5 1.71-1.89 (m, 9H), 1.53 (d, J = 12.8 Hz, 2H), 1.21-1.23 (m, 2H), 0.67-0.69 (m, 2H)

Example 3: Preparation of E-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
 10 adamantan-1-carboxylic acid amide




1.44 g (3.20 mmol) of the 4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
 adamantan-1-carboxylic acid prepared in Preparational
 15 Example 12 was dissolved in 107 ml of methylene
 chloride, to which 533 mg (3.84 mmol) of HOBt and 736
 mg (3.84 mmol) of EDCI were added. The mixture was
 stirred at room temperature for 10 minutes and then
 107 ml of 35% ammonia water was added thereto. The
 20 mixture was stirred at room temperature for 20 hours.
 Upon completion of the reaction, extraction was
 performed with methylene chloride. After washing with




brine, the extract was dried over anhydrous magnesium sulfate, followed by column chromatography to give 1.09 g of E-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide (yield: 76%).




^1H NMR (400 MHz, CDCl_3) δ 7.88-7.92 (m, 1H), 7.59-7.64 (m, 1H), 7.31-7.33 (m, 1H), 7.24 (t, J = 8.8 Hz, 1H), 6.61 (d, J = 7.2 Hz, 1H), 5.61 (brs, 1H), 5.45 (t, J = 6.6 Hz, 1H), 5.33 (brs, 1H), 4.06 (t, J = 3.8 Hz, 1H), 3.13 (d, J = 6.4 Hz, 2H), 1.91-2.09 (m, 11H), 1.63 (d, J = 12.0 Hz, 2H), 1.22-1.24 (m, 2H), 0.67-0.69 (m, 2H)




The compounds of Example 4 - Example 37 were prepared by the preparation method described in Example 3.



【Table 1】




Ex am pl e	Compound Name	Structure	NMR data
4	Z-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-		^1H NMR (400 MHz, CDCl_3) δ 7.87-7.89 (m, 1H), 7.60-7.66 (m, 1H), 7.31-7.35 (m, 1H), 7.22-7.25 (m, 2H), 7.01 (d, J = 7.6 Hz, 1H), 6.57 (t, J = 6.2 Hz, 1H), 6.28 (brs, 1H), 5.56 (brs, 1H), 4.08 (d, J = 7.6




	adamantan-1-carboxylic acid amide		Hz, 1H), 3.17 (d, J = 6.4 Hz, 2H), 2.04-2.12 (m, 4H), 1.92 (s, 2H), 1.74-7.84 (m, 7H), 1.27-1.30 (m, 2H), 0.65-0.67 (m, 2H)
5	E-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropane carboxamidol]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.90 (d, J = 8.0 Hz, 1H), 7.59 (d, J = 8.0 Hz, 1H), 7.27 (t, J = 8.0 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.69 (t, J = 6.4 Hz, 1H), 5.62 (brs, 1H), 5.37 (brs, 1H), 4.02 (t, J = 3.4 Hz, 1H), 3.06 (d, J = 6.4 Hz, 2H), 2.70 (s, 3H), 1.98-2.06 (m, 7H), 1.90 (s, 2H), 1.86 (d, J = 13.6 Hz, 2H), 1.61 (d, J = 13.6 Hz, 2H), 1.14-1.16 (m, 2H), 0.71-0.73 (m, 2H)
6	Z-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropane carboxamidol]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.89 (t, J = 6.2 Hz, 1H), 7.62 (t, J = 6.6 Hz, 1H), 7.26-7.31 (m, 2H), 6.85 (s, 1H), 6.23 (brs, 1H), 5.48 (brs, 1H), 4.08 (s, 1H), 3.07-3.08 (m, 2H), 2.70-2.75 (m, 3H), 1.54-2.17 (m, 13H), 1.26-1.27 (m, 2H), 0.67-0.72 (m, 2H)
7	E-4-[1-((3-chloro-benzenesulfonylamino)methyl)cyclopropane carboxamidol]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.86 (t, J = 1.8 Hz, 1H), 7.75-7.77 (m, 1H), 7.58-7.60 (m, 1H), 7.50 (t, J = 8.0 Hz, 1H),




	)methyl)cyclopropane carboxamid o] - adamantan-1-carboxylic acid amide		6.62 (d, $J = 7.6$ Hz, 1H), 5.96 (t, $J = 6.4$ Hz, 1H), 5.72 (brs, 1H), 5.51 (brs, 1H), 4.06 (d, $J = 6.8$ Hz, 1H), 3.11 (d, $J = 6.4$ Hz, 2H), 1.92-2.09 (m, 11H), 1.64 (d, $J = 12.8$ Hz, 2H), 1.22-1.24 (m, 2H), 0.71-0.73 (m, 2H)
8	E-4-[1-((3-chloro-2-fluorobenzenesulfonylamino)methyl)cyclopropane carboxamid o] - adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.78-7.82 (m, 1H), 7.63-7.67 (m, 1H), 7.23-7.28 (m, 1H), 6.39 (d, $J = 7.2$ Hz, 1H), 5.61-5.63 (m, 2H), 5.30 (brs, 1H), 4.05 (d, $J = 7.2$ Hz, 1H), 3.19 (d, $J = 5.2$ Hz, 2H), 1.88-2.08 (m, 11H), 1.62 (d, $J = 13.6$ Hz, 2H), 1.20-1.22 (m, 2H), 0.73-0.75 (m, 2H)
9	E-4-[1-((3,5-difluorobenzenesulfonylamino)methyl)cyclopropane carboxamid o] - adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.40-7.42 (m, 2H), 7.02-7.07 (m, 1H), 6.67 (t, $J = 6.2$ Hz, 1H), 6.61 (d, $J = 7.6$ Hz, 1H), 5.82 (brs, 1H), 5.73 (brs, 1H), 4.05 (d, $J = 7.2$ Hz, 1H), 3.09 (d, $J = 6.4$ Hz, 2H), 1.19-2.07 (m, 11H), 1.62 (d, $J = 12.8$ Hz, 2H), 1.20-1.23 (m, 2H), 0.72-0.74 (m, 2H)
10	E-4-[1-((2-fluoro-6-methylbenzenesulfonylamino)methyl)cyclopropane carboxamid o] - adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.40-7.45 (m, 1H), 7.13 (d, $J = 8.0$ Hz, 1H), 7.06 (dd, $J = 10.8, 8.4$ Hz, 1H), 6.79 (d, J




	fonylamino)methyl)cyclopropane carboxamid o]- adamantan- 1- carboxylic acid amide		= 7.6 Hz, 1H), 5.63 (brs, 1H), 5.58 (s, 1H), 5.50 (brs, 1H), 4.07 (d, $J = 7.2$ Hz, 1H), 3.14 (d, $J = 6.8$ Hz, 2H), 2.69 (s, 3H), 2.10 (s, 2H), 1.82-2.06 (m, 9H), 1.63 (d, $J = 12.8$ Hz, 2H), 1.25-1.27 (m, 2H), 0.66-0.68 (m, 2H)
11	E-4-[1- ((2,3- difluoro- benzenesul fonylamino)methyl)cyclopropane carboxamid o]- adamantan- 1- carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.64-7.68 (m, 1H), 7.39-7.47 (m, 1H), 7.23- 7.28 (m, 1H), 6.51 (d, J $= 7.2$ Hz, 1H), 6.06 (t, $J = 6.2$ Hz, 1H), 5.75 (brs, 1H), 5.63 (brs, 1H), 4.05 (d, $J = 7.2$ Hz, 1H), 3.19 (d, $J =$ 6.4 Hz, 2H), 1.90-2.07 (m, 11H), 1.61 (d, $J =$ 12.8 Hz, 2H), 1.20-1.23 (m, 2H), 0.73-0.76 (m, 2H)
12	E-4-[1- ((2,4,6- trifluoro- benzenesul fonylamino)methyl)cyclopropane carboxamid o]- adamantan- 1- carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 6.80-6.85 (m, 2H), 6.26 (d, $J = 7.6$ Hz, 1H), 5.81 (t, $J = 6.2$ Hz, 1H), 5.59 (brs, 1H), 5.30 (brs, 1H), 4.04 (d, $J = 7.6$ Hz, 1H), 3.23 (d, $J = 6.0$ Hz, 2H), 1.85-2.07 (m, 11H), 1.62 (d, $J = 13.2$ Hz, 2H), 1.20-1.23 (m, 2H), 0.78-0.81 (m, 2H)
13	E-4-[1- ((2- fluoro- N,6- dimethyl-		^1H NMR (400 MHz, CDCl_3) d 7.30-7.44 (m, 1H), 7.23-7.25 (m, 1H), 7.13 (d, $J = 7.6$ Hz, 1H), 7.02-7.06 (m, 1H), 5.93




	benzenesulfonylamino)methyl)cyclopropane carboxamidate]-adamantan-1-carboxylic acid amide		(brs, 1H), 5.64 (brs, 1H), 4.10 (d, $J = 7.2$ Hz, 1H), 3.40 (s, 2H), 2.94 (d, $J = 2.4$ Hz, 3H), 2.71 (s, 3H), 2.08-2.13 (m, 4H), 1.98-2.04 (m, 5H), 1.90-1.95 (m, 2H), 1.60 (d, $J = 11.2$ Hz, 2H), 1.38-1.41 (m, 2H), 0.60-0.63 (m, 2H)
14	E-4-[1-((2,4-dichloro-5-methyl-benzenesulfonylamino)methyl)cyclopropane carboxamidate]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.94 (s, 1H), 7.54 (s, 1H), 6.58 (d, $J = 8$ Hz, 1H), 5.59 (t, $J = 6.4$ Hz, 2H), 5.46 (s, 1H), 4.07-4.06 (m, 1H), 3.00 (d, $J = 6.4$ Hz, 2H), 2.42 (s, 3H), 2.09-1.92 (m, 11H), 1.64 (d, $J = 6.6$ Hz, 2H), 1.25-1.22 (m, 2H), 0.68-0.66 (m, 2H)
15	E-4-[1-((4-chloro-2-fluoro-5-methyl-benzenesulfonylamino)methyl)cyclopropane carboxamidate]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.74 (d, $J = 7.6$ Hz, 1H), 7.26 (d, $J = 9.6$ Hz, 1H), 6.51 (d, $J = 7.6$ Hz, 1H), 5.62 (s, 1H), 5.51 (t, $J = 6.4$ Hz, 1H), 5.34 (s, 1H), 4.06-4.06 (m, 1H), 3.12 (d, $J = 6.4$ Hz, 2H), 2.40 (s, 3H), 2.08-1.91 (m, 11H), 1.63 (d, $J = 12$ Hz, 2H), 1.24-1.21 (m, 2H), 0.72-0.69 (m, 2H)





16	E-4-[1- (4,5-dichloro-2-fluorobenzenesulfonylamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.98 (d, J = 6.8 Hz, 1H), 7.38 (d, J = 9.2 Hz, 1H), 6.20 (d, J = 7.2 Hz, 1H), 5.68 (t, J = 6.4 Hz, 1H), 5.60 (s, 1H), 5.46 (s, 1H), 4.04-4.03 (m, 1H), 3.16 (d, J = 6.4 Hz, 2H), 2.07-1.83 (m, 11H), 1.64-1.59 (d, J = 18.4 Hz, 2H), 1.21-1.18 (m, 2H), 0.80-0.77 (m, 2H)
17	E-4-[1- (furan-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.58 (dd, J = 1.8, 0.8 Hz, 1H), 7.07 (dd, J = 3.6, 0.8 Hz, 1H), 6.53 (dd, J = 3.2, 1.6 Hz, 1H), 6.47 (d, J = 7.6 Hz, 1H), 5.60 (s, 1H), 5.41 (t, J = 6.4 Hz, 1H), 5.28 (s, 1H), 4.06-4.04 (m, 1H), 3.20 (d, J = 6.8 Hz, 2H), 2.07-1.90 (m, 11H), 1.61 (d, J = 18.8 Hz, 2H), 1.25-1.20 (m, 2H), 0.75-0.72 (m, 2H)
18	E-4-[1- (3,5-dichlorobenzenesulfonylamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 8.04 (s, 1H), 7.87 (s, 1H), 7.87 (s, 1H), 7.15-7.06 (m, 2H), 6.81 (s, 1H), 3.88-3.87 (m, 1H), 3.23 (d, J = 4.4 Hz, 1H), 3.16 (s, 2H), 2.06-1.82 (m, 11H), 1.50 (d, J = 12.4 Hz, 2H), 1.04-0.98 (m, 2H), 0.75-0.70 (m, 2H)




19	E-4-[1- (thiophen-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.64-7.62 (m, 2H), 7.14-7.12 (m, 1H), 6.54 (d, <i>J</i> = 7.6 Hz, 1H), 5.56 (s, 1H), 5.19 (s, 1H), 5.03 (t, <i>J</i> = 6.6 Hz, 1H), 4.06-4.03 (m, 1H), 3.16 (d, <i>J</i> = 6.8 Hz, 2H), 2.08-1.91 (m, 11H), 1.62 (d, <i>J</i> = 13.2 Hz, 2H), 1.25-1.23 (m, 2H), 0.71-0.70 (m, 2H)
20	E-4-[1- (2-(trifluoromethyl)-4-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 8.25 (dd, <i>J</i> = 8.8, 5.2 Hz, 1H), 7.61 (dd, <i>J</i> = 8.8, 2.4 Hz, 1H), 7.43-7.39 (m, 1H), 6.47 (d, <i>J</i> = 7.6 Hz, 1H), 5.64 (s, 1H), 5.54 (t, <i>J</i> = 6.2 Hz, 1H), 5.49 (s, 1H), 4.06-4.04 (m, 1H), 3.09 (d, <i>J</i> = 6 Hz, 2H), 2.08-1.90 (m, 11H), 1.67-1.61 (m, 2H), 1.24-1.21 (m, 2H), 0.71-0.68 (m, 2H)
21	E-4-[1- (3,4-difluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.73-7.68 (m, 1H), 7.66-7.63 (m, 1H), 7.37-7.30 (m, 1H), 6.42 (d, <i>J</i> = 5.6 Hz, 1H), 5.76 (s, 1H), 5.66 (s, 1H), 5.42 (s, 1H), 4.04 (s, 1H), 3.08 (d, <i>J</i> = 6 Hz, 2H), 2.07-1.91 (m, 11H), 1.71-1.61 (m, 2H), 1.22-1.19 (m, 2H), 0.76-0.72 (m, 2H)

22	E-4-[1- (2-fluoro-N-methyl-benzenesulfonylamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.86-7.90 (m, 1H), 7.61-7.66 (m, 1H), 7.34 (t, <i>J</i> = 7.6 Hz, 1H), 7.23-7.26 (m, 2H), 6.05 (brs, 1H), 6.83 (brs, 1H), 4.10 (d, <i>J</i> = 7.2 Hz, 1H), 3.31 (s, 2H), 2.90 (d, <i>J</i> = 2.0 Hz, 3H), 1.90-2.12 (m, 11H), 1.61 (d, <i>J</i> = 12.0 Hz, 2H), 1.36-1.38 (m, 2H), 0.57-0.60 (m, 2H)
23	E-4-[1- (4-trifluoromethoxybenzenesulfonylamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 8.14 (brs, 1H), 7.96 (m, 2H), 7.61 (d, <i>J</i> = 8.4 Hz, 2H), 7.10 (d, <i>J</i> = 6.8 Hz, 1H), 7.01 (brs, 1H), 6.74 (brs, 1H), 3.85 (m, 1H), 3.04 (s, 2H), 2.00 (d, <i>J</i> = 12.4 Hz, 1H), 1.91-1.76 (m, 9H), 1.45 (d, <i>J</i> = 12.4 Hz, 2H), 0.95 (m, 2H), 0.64 (m, 2H)
24	E-4-[1- (2,3-difluorobenzeneamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 8.95 (t, <i>J</i> = 6 Hz, 1H), 7.71 (d, <i>J</i> = 6.8 Hz, 1H), 7.62-7.56 (m, 1H), 7.38-7.29 (m, 2H), 7.02 (s, 1H), 6.72 (s, 1H), 4.18 (s, 1H), 3.17 (d, <i>J</i> = 4.4 Hz, 2H), 2.02-1.73 (m, 11H), 1.40 (d, <i>J</i> = 11.6 Hz, 2H), 1.03-1.01 (m, 2H), 0.84-0.82 (m, 2H)

25	E-4-[1-((3,4-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 8.96 (t, <i>J</i> = 6 Hz, 1H), 7.93 (t, <i>J</i> = 9.6 Hz, 1H), 7.85 (d, <i>J</i> = 6.8 Hz, 1H), 7.64-7.58 (m, 1H), 6.97 (s, 1H), 6.70 (s, 1H), 3.81-3.79 (m, 1H), 3.60 (d, <i>J</i> = 6.4 Hz, 2H), 2.01-1.73 (m, 11H), 1.39 (d, <i>J</i> = 12.4 Hz, 2H), 1.04-1.02 (m, 2H), 0.87-0.85 (m, 2H)
26	E-4-[1-((1-methyl-1H-indole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 8.08 (d, <i>J</i> = 1.6 Hz, 1H), 7.82 (t, <i>J</i> = 6.2 Hz, 1H), 7.65 (d, <i>J</i> = 8.8 Hz, 1H), 7.59 (dd, <i>J</i> = 8.8, 1.6 Hz, 1H), 7.53 (d, <i>J</i> = 3.2 Hz, 1H), 7.27 (d, <i>J</i> = 7.6 Hz, 1H), 7.01 (s, 1H), 6.73 (s, 1H), 6.64 (d, <i>J</i> = 2.8 Hz, 1H), 3.85 (s, 3H), 2.90 (d, <i>J</i> = 6 Hz, 2H), 2.08-1.78 (m, 11H), 1.48 (d, <i>J</i> = 12.8 Hz, 2H), 0.94 (m, 2H), 0.60-0.57 (m, 2H)
27	E-4-[1-((1-methyl-1H-pyrazole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 7.55 (d, <i>J</i> = 2 Hz, 1H), 7.16 (d, <i>J</i> = 8 Hz, 1H), 7.03 (s, 1H), 6.74 (d, <i>J</i> = 2.4 Hz, 2H), 3.95 (s, 3H), 3.72 (d, <i>J</i> = 7.2 Hz, 1H), 1.93 (s, 1H), 1.83-1.81 (m, 8H), 1.76-1.75 (m, 2H), 1.49 (d, <i>J</i> = 12.4 Hz, 2H), 1.20-1.17 (m, 2H), 0.88-0.85 (m, 2H)


28	E-4-[1-((benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 8.05 (s, 1H), 7.83 (d, <i>J</i> = 8 Hz, 2H), 7.69-7.60 (m, 3H), 7.17 (s, 1H), 7.01 (s, 1H), 6.73 (s, 1H), 3.87-3.86 (m, 1H), 2.99 (s, 2H), 2.03 (d, <i>J</i> = 13.2 Hz, 2H), 1.91 (s, 3H), 1.83 (s, 4H) 1.77 (s, 2H), 1.46 (d, <i>J</i> = 13.6 Hz, 2H), 0.96-0.94 (m, 2H), 0.64-0.62 (m, 2H)
29	E-4-[1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid		¹ H NMR (400 MHz, MeOD) δ 7.91-7.87 (m, 1H), 7.70-7.66 (m, 1H), 7.48-7.43 (m, 1H), 7.39-7.30 (m, 2H), 4.00 (s, 1H), 3.16 (s, 2H), 2.17-1.85 (m, 11H), 1.61 (d, <i>J</i> = 12.8 Hz, 2H), 1.15-1.13 (m, 2H), 0.72-0.70 (m, 2H)
30	N-(bicyclo[2.2.1]heptan-2-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido		¹ H NMR (400 MHz, CDCl ₃) δ 7.90 (t, <i>J</i> = 7.0 Hz, 1H), 7.61 (d, <i>J</i> = 5.6 Hz, 1H), 7.32 (t, <i>J</i> = 7.4 Hz, 1H), 7.24 (t, <i>J</i> = 10.0 Hz, 1H), 7.08 (s, 1H), 4.15 (d, <i>J</i> = 4.0 Hz, 1H), 3.44 (d, <i>J</i> = 14.4 Hz, 1H), 3.11 (d, <i>J</i> = 14.4 Hz, 1H), 2.93 (s, 3H), 2.48 (s, 1H), 2.22 (s, 1H), 2.02-2.08 (m, 1H), 1.60-1.74 (m, 3H), 1.26-1.45 (m, 5H), 1.00 (d, <i>J</i> = 12.4 Hz, 1H), 0.56 (s, 2H)

31	N-(adamantan-1-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido		¹ H NMR (400 MHz, CDCl ₃) d 7.88 (t, J = 7.4 Hz, 1H), 7.58-7.63 (m, 1H), 7.31 (t, J = 7.7 Hz, 1H), 7.23 (t, J = 9.2 Hz, 1H), 6.62 (s, 1H), 3.24 (s, 2H), 2.90 (d, J = 2.0 Hz, 3H), 2.07 (s, 9H), 1.65-1.73 (m, 6H), 1.28-1.31 (m, 2H), 0.51-0.53 (m, 2H)
32	E-4-[1-((N-ethyl-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.92 (m, 1H), 7.59 (m, 1H), 7.32 (td, J ₁ = 1.2 Hz, J ₂ = 6.4 Hz, 1H), 7.22 (m, 2H), 5.63 (brs, 1H), 5.43 (brs, 1H), 4.12 (d, J = 5.6 Hz, 1H), 3.52 (s, 2H), 3.32 (m, 2H), 2.18-1.86 (m, 13H), 1.39 (m, 2H), 1.12 (t, J = 6.8 Hz, 3H), 0.65 (m, 2H)
33	E-3-(4-(1((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido)adamantan-1-yl)propanoic acid		¹ H NMR (400 MHz, MeOD) d 7.86-7.94 (m, 1H), 7.65-7.70 (m, 1H), 7.30-7.43 (m, 2H), 3.94 (s, 1H), 3.65 (s, 1H), 2.23-2.32 (m, 2H), 2.13 (d, J = 13.2 Hz, 2H), 1.92-2.01 (m, 2H), 1.53-1.61 (m, 9H), 1.42-1.46 (m, 2H), 1.12-1.15 (m, 2H), 0.68-0.71 (m, 2H)
34	E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-1-((2-		¹ H NMR (400 MHz, DMSO) d 7.88-7.92 (m, 1H), 7.58-7.64 (m, 1H), 7.31 (t, J = 7.6 Hz, 1H), 7.21-7.25 (m, 1H), 6.47 (d, J = 7.6 Hz, 1H), 5.62 (brs,

	fluoro-benzenesulfonylamino)methyl)cyclopropane carboxamide		1H), 5.45-5.48 (m, 2H), 3.98 (d, $J = 7.2$ Hz, 1H), 3.12 (t, $J = 6.4$ Hz, 2H), 2.16-2.20 (m, 2H), 1.98 (s, 2H), 1.87 (d, $J = 13.6$ Hz, 2H), 1.46-1.62 (m, 11H), 1.18-1.21 (m, 2H), 0.68-0.71 (m, 2H)
35	E-N-(5-aminoadamantan-2-yl)-1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropane carboxamide hydrochloride		^1H NMR (400 MHz, D_2O) d 7.79 (dt, $J = 7.6, 1.6$ Hz, 1H), 7.64-7.70 (m, 1H), 7.27-7.35 (m, 2H), 3.75 (s, 1H), 3.15 (s, 2H), 1.88-2.10 (m, 11H), 1.48 (d, $J = 13.6$ Hz, 2H), 0.99-1.02 (m, 2H), 0.66-0.69 (m, 2H)
36	E-4-[1-((2,4,5-trifluoro-benzenesulfonylamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.73-7.79 (m, 1H), 7.08-7.14 (m, 1H), 6.18 (d, $J = 7.6$ Hz, 1H), 5.55 (t, $J = 6.4$ Hz, 1H), 5.21 (brs, 1H), 4.03 (d, $J = 7.2$ Hz, 1H), 3.15 (d, $J = 6.4$ Hz, 2H), 1.99-2.07 (m, 7H), 1.91 (s, 2H), 1.85 (d, $J = 13.2$ Hz, 2H), 1.63 (d, $J = 13.2$ Hz, 2H), 1.19-1.22 (m, 2H), 0.77-0.80 (m, 2H)
37	E-4-[1-((4-chloro-benzenesulfonylamino)methyl)cyclopropane carboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 8.11 (t, $J = 6.4$ Hz, 1H), 7.85-7.81 (m, 2H), 7.71-7.68 (m, 2H), 7.08 (d, $J = 7.6$ Hz, 1H), 7.00 (s, 1H), 6.73 (s,

clopropane carboxamid o]- adamantan- 1- carboxylic acid amide		1H), 3.86-3.84 (m, 1H), 3.01 (d, $J = 6.4$ Hz, 2H), 2.02-1.76 (m, 11H), 1.46 (d, $J = 12.8$ Hz, 2H), 0.97-0.94 (m, 2H), 0.65-0.63 (m, 2H)
---	--	--

Preparational Example 13: Preparation of 1-aminocyclopropyl acetic acid hydrochloride



5 After reducing pressure in the reaction vessel, the vessel was filled with nitrogen gas, to which 6.8 ml of 7 N NH₃ in MeOH was loaded. 1.0 g (6.30 mmol) of dimethylcyclopropane-1,1-dicarboxylate was dissolved in 63 ml of MeOH, which was also loaded into


10 the reaction vessel. If the starting material still remained, NH₃ gas could be provided for bubbling for 15 minutes. Upon completion of the reaction, the reaction solvent was eliminated by evaporation under reduced pressure and the precipitated solid compound

15 was filtered and washed with 0°C methanol, followed by vacuum drying. The obtained reaction product was dissolved in 5 ml of 7.4% NaOH/H₂O, followed by stirring at 40°C for 20 minutes. Then, the temperature was lowered to room temperature (Reactant

A). 7 ml of 12.3% NaOCl and 2 ml of 30% NaOH/H₂O were stirred together at room temperature for 1 hour (Reactant B). These reactants A and B were mixed and stirred at 80°C for 4 minutes. After lowering the
 5 temperature thereof, 4 ml of hydrochloric acid was slowly added thereto carefully not to increase the temperature more than 60°C. The solvent was eliminated by evaporation under reduced pressure and the reactant was dissolved in ethanol. The
 10 precipitated solid compound was filtered. The remaining solution was evaporated under reduced pressure and then dissolved in hot acetone. The generated solid compound was filtered and dried to give 0.5 g of 1-aminocyclopropyl acetic acid
 15 hydrochloride (yield: 58%).

¹H NMR (400 MHz, D₂O) δ 1.40-1.43 (m, 2H), 1.22-1.25 (m, 2H)

Preparational Example 14: Preparation of 1-(t-butoxycarbonylamino)cyclopropyl acetic acid
 20




4.18 g (30.37 mmol) of the 1-aminocyclopropyl acetic acid hydrochloride prepared in Preparational

Example 13 was dissolved in 61 ml of methylene chloride, to which 4.7 ml (33.40 mmol) of TEA was added. The mixture was stirred for 10 minutes and then methylene chloride was evaporated under reduced pressure. The reactant was dissolved in 36.5 ml of 1N NaOH solution and 101 ml of 1,4-dioxane, to which 8.4 ml (36.44 mmol) of Boc_2O was added. The reaction mixture was stirred at room temperature for 16 hours, and then 1,4-dioxane was evaporated under reduced pressure. The residue was acidized with 1 N HCl to be pH 3, followed by extraction with 3100 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 3.59 g of 1-(t-butoxycarbonylamino)cyclopropyl acetic acid (yield: 59%).

^1H NMR (400 MHz, MeOD) δ 1.51 (s, 2H), 1.51 (s, 9H), 1.44 (s, 4H)

20 Preparational Example 15: Preparation of 4-[1-(t-butoxycarbonylamino)-cyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester




3.59 g (17.85 mmol) of the 1-(t-butoxycarbonylamino)cyclopropyl acetic acid prepared in Preparational Example 14 and 4.83 g (19.64 mmol) of the 4-amino-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 7 were dissolved in 60 ml of methylene chloride, to which 5.15 g (19.64 mmol) of BOP-Cl was added. 4.98 ml (35.70 mmol) of TEA was also added thereto. The mixture was stirred at room temperature for 16 hours and methylene chloride was evaporated under reduced pressure. The residue was dissolved in 50 ml of water, followed by extraction with 3100 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 4.95 g of 4-[1-(t-butoxycarbonylamino)-cyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester (yield: 71%).

^1H NMR (400 MHz, CDCl_3) δ 7.27 (brs, 1H), 5.10 (brs, 1H), 4.00 (d, J = 6.8 Hz, 1H), 3.66 (s, 3H), 1.90-2.05 (m, 9H), 1.80 (d, J = 12.8 Hz, 2H), 1.60 (d,

$J = 13.6$ Hz, 2H), 1.53-1.56 (m, 2H), 1.47 (s, 9H).
0.98-1.01 (m, 2H)


Preparational Example 16: Preparation of 4-(1-
5 aminocyclopropanylcarboxamido)-adamantan-1-carboxylic
acid methyl ester hydrochloride



4.95 g (14.42 mmol) of the 4-[1-(t-butoxycarbonylamino)-cyclopropanecarboxamido]-
10 adamantan-1-carboxylic acid methyl ester prepared in
Preparational Example 15 was dissolved in 48 ml of
ethyl acetate, to which 36 ml of 4M HCl 1,4-dioxane
was added, followed by stirring at room temperature
for 6 hours. Upon completion of the reaction, the
15 reaction solvent was eliminated by evaporation under
reduced pressure. The residue was dissolved in 50 ml
of ethylene acetate and the precipitated solid
compound was filtered. As a result, 4.1 g of 4-(1-
aminocyclopropanylcarboxamido)-adamantan-1-carboxylic
20 acid methyl ester hydrochloride was obtained (yield:
86%).

^1H NMR (400 MHz, D_2O) δ 3.79 (s, 1H), 3.56 (s, 3H), 3.08 (s, 2H), 1.86-1.95 (m, 7H), 1.71 (d, J = 13.6 Hz, 2H), 1.40-1.46 (m, 4H), 1.30-1.33 (m, 2H)

5 Preparational Example 17: Preparation of 4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid methyl ester




200 mg (0.61 mmol) of the 4-(1-aminocyclopropanylcarboxamido)-adamantan-1-carboxylic acid methyl ester hydrochloride prepared in Preparational Example 16 was dissolved in 2 ml of methylene chloride, to which 0.17 ml of TEA was added, followed by stirring at room temperature for 5 minutes. 142 mg (0.67 mmol) of 2-chlorosulfonylchloride was added to the reactant, which was then stirred at room temperature for 6 hours. The reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in 10 ml of water, followed by extraction with 330 ml of ethyl acetate. The extract was dried

over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 108 g of 4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-

5 adamantan-1-carboxylic acid methyl ester (yield: 38%).

^1H NMR (400 MHz, CDCl_3) δ 7.85 (t, J = 1.8 Hz, 1H), 7.75 (d, J = 7.8 Hz, 1H), 7.60 (d, J = 8.0 Hz, 1H), 7.49 (t, J = 8.0 Hz, 1H), 7.18 (d, J = 7.6 Hz, 1H), 5.44 (d, J = 10.4 Hz, 1H), 3.98 (d, J = 7.6 Hz, 1H), 3.66 (s, 3H), 1.89-2.05 (m, 10H), 1.64 (s, 1H), 1.59 (d, J = 10.4 Hz, 2H), 1.44-1.48 (m, 2H), 0.82-0.85 (m, 2H)

Preparational Example 18: Preparation of 4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-
 15 adamantan-1-carboxylic acid



100 mg (0.21 mmol) of the 4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-


20 adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 17 was dissolved in 1.5 ml of

THF/ethanol mixed solution, to which 1.5 ml of 2N NaOH solution was added, followed by stirring at room temperature overnight. The residue was acidized with 1 N HCl, followed by extraction with ethyl acetate.
 5 The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 96 mg of 4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid (yield: 98%).

10 ^1H NMR (400 MHz, MeOD) δ 8.98 (s, 1H), 7.72-7.78 (m, 3H), 7.62-7.66 (m, 1H), 7.14 (d, J = 7.6 Hz, 1H), 3.61 (d, J = 6.8 Hz, 1H), 1.75-2.01 (m, 11H), 1.47 (d, J = 12.8 Hz, 2H), 1.16-1.19 (m, 2H), 0.88-0.91 (m, 11H)

15

Example 38: Preparation of E-4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide





20 96 mg (0.21 mmol) of the 4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid prepared in Preparational





Example 18 was dissolved in 1 ml of acetonitrile, to which 34 mg (0.25 mmol) of HOBt, 48 mg (0.25 mmol) of EDCI, and 60 ml (0.42 mmol) of TEA were added. The mixture was stirred at room temperature for 10
 5 minutes, to which 13 mg (0.25 mmol) of ammonium chloride was added, followed by stirring at room temperature for 20 hours. Upon completion of the reaction, the reaction solvent was eliminated by evaporation under reduced pressure and the reactant
 10 was dissolved in 60 ml of methylene chloride. The organic layer was washed with 1 N HCl (2 x 2 ml) and then washed again with brine. The reaction product was dried over anhydrous magnesium sulfate, followed by column chromatography to give 41 mg of E-4-[1-(3-
 15 chloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide (yield: 43%).




^1H NMR (400 MHz, CDCl_3) δ 7.84 (t, J = 2.0 Hz, 1H), 7.75 (dt, J = 7.2, 1.2 Hz, 1H), 7.59 (m, 1H), 7.49 (t, J = 8.0 Hz, 1H), 7.22 (d, J = 8.0 Hz, 1H),
 20 5.75 (s, 1H), 5.62 (brs, 1H), 5.35 (brs, 1H), 3.99 (t, J = 7.6 Hz, 1H), 2.08 (s, 3H), 1.92-1.99 (m, 8H), 1.63 (d, J = 12.8 Hz, 2H), 1.45-1.48 (m, 2H), 0.82-0.85 (m, 2H)




The compounds of Example 39 - Example 66 were prepared by the preparation method described in Example 38.





【Table 2】




Ex am pl e	Compound Name	Structure	NMR data
39	E-4-[1-(2-fluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.92 (dt, $J = 1.6, 7.6$ Hz, 1H), 7.63-7.65 (m, 1H), 7.34 (t, $J = 7.6$ Hz, 2H), 7.22-7.26 (m, 1H), 5.58 (brs, 1H), 5.52 (s, 1H), 5.22 (brs, 1H), 4.02 (d, $J = 7.6$ Hz, 1H), 3.65-3.66 (m, 2H), 1.93-2.11 (m, 11H), 1.65 (d, $J = 13.6$ Hz, 2H), 1.44-1.47 (m, 2H), 0.77-0.80 (m, 2H)
40	E-4-[1-(2-fluoro-6-methyl-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.90 (d, $J = 8.0$ Hz, 1H), 7.59 (d, $J = 8.0$ Hz, 1H), 7.27 (t, $J = 8.0$ Hz, 1H), 6.27 (d, $J = 7.6$ Hz, 1H), 5.69 (t, $J = 6.4$ Hz, 1H), 5.62 (brs, 1H), 5.37 (brs, 1H), 4.02 (t, $J = 3.4$ Hz, 1H), 3.06 (d, $J = 6.4$ Hz, 2H), 2.70 (s, 3H), 1.98-2.06 (m, 7H), 1.90 (s, 2H), 1.86 (d, $J = 13.6$ Hz, 2H), 1.61 (d, $J = 13.6$ Hz, 2H), 1.14-1.16 (m, 2H), 0.71-0.73 (m, 2H)




41	E-4-[1-(3-chloro-2-methyl-benzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.93 (d, <i>J</i> = 7.6 Hz, 1H), 7.63 (d, <i>J</i> = 7.6 Hz, 1H), 7.37-7.40 (m, 1H), 7.30 (t, <i>J</i> = 8.0 Hz, 1H), 6.11 (brs, 2H), 5.77 (brs, 1H), 4.00 (d, <i>J</i> = 4.8 Hz, 1H), 2.63 (s, 3H), 1.91-2.09 (m, 11H), 1.63 (d, <i>J</i> = 12.8 Hz, 2H), 1.37-1.40 (m, 1H), 0.71-0.74 (m, 2H)
42	E-4-[1-(4-fluorobenzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 7.82-7.85 (m, 2H), 7.45 (t, <i>J</i> = 8.8 Hz, 2H), 7.19 (d, <i>J</i> = 7.6 Hz, 1H), 7.02 (brs, 1H), 6.74 (brs, 1H), 3.66 (d, <i>J</i> = 7.2 Hz, 1H), 1.91 (s, 1H), 1.75-1.84 (m, 9H), 1.46 (d, <i>J</i> = 12.0 Hz, 2H), 1.14-1.17 (m, 2H), 0.83-0.86 (m, 2H)
43	E-4-[1-(2,4-dichloro-5-methyl-benzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.96 (s, 1H), 7.54 (s, 1H), 7.34 (d, <i>J</i> = 8.0 Hz, 1H), 6.20 (s, 1H), 5.66 (brs, 1H), 5.54 (brs, 1H), 4.02 (d, <i>J</i> = 7.6 Hz, 1H), 2.43 (s, 3H), 1.92-2.10 (m, 10H), 1.63-1.70 (m, 3H), 1.42-1.45 (m, 2H), 0.75-0.78 (m, 2H)
44	E-4-[1-(2,4-difluorochlorobenzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.91-7.96 (m, 1H), 7.26 (d, <i>J</i> = 6.4 Hz, 1H), 7.03-7.08 (m, 1H), 6.96-7.01 (m, 1H), 5.79 (s, 1H), 5.62 (brs, 1H),





	)cycloprop anecarboxa mido]- adamantan- 1- carboxylic acid amide		5.48 (brs, 1H), 4.01 (d, $J = 8.0$ Hz, 1H), 2.10 (s, 2H), 1.92-2.05 (m, 9H), 1.65 (d, $J = 13.2$ Hz, 2H), 1.45-1.48 (m, 2H), 0.73-0.82 (m, 2H)
45	E-4-[1-(2-fluoro-4,5-dichloro-benzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 9.38 (brs, 1H), 8.06 (d, $J = 9.6$ Hz, 1H), 7.64 (d, $J = 7.2$ Hz, 1H), 7.09 (d, $J = 7.6$ Hz, 1H), 7.03 (s, 1H), 6.75 (s, 1H), 3.66 (d, $J = 7.2$ Hz, 1H), 1.90 (s, 1H), 1.74-1.79 (m, 10H), 1.45 (d, $J = 12.4$ Hz, 2H), 1.19-1.22 (m, 2H), 0.94-0.97 (m, 2H)
46	E-4-[1-(2-fluoro-4-chloro-5-methyl-benzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.76 (d, $J = 7.6$ Hz, 1H), 7.24-7.30 (m, 2H), 6.32 (s, 1H), 5.69 (brs, 1H), 5.61 (brs, 1H), 4.00 (d, $J = 7.6$ Hz, 1H), 2.41 (s, 3H), 2.07-2.09 (m, 3H), 1.92-1.00 (m, 8H), 1.63 (d, $J = 13.2$ Hz, 2H), 1.44-1.47 (m, 2H), 0.82-0.85 (m, 2H)
47	E-4-[1-(2,3,4-trifluoro-benzenesulfonylamino)cyclopropyl]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.67-7.73 (m, 1H), 7.13-7.20 (m, 2H), 6.24 (brs, 1H), 5.64 (brs, 1H), 5.42 (brs, 1H), 4.01 (d, $J = 8.0$ Hz, 1H), 2.09 (s, 3H), 1.93-2.04 (m, 8H), 1.64 (d, $J = 14.0$ Hz, 2H), 1.47-



	1-carboxylic acid amide		1.50 (m, 2H), 0.82-0.86 (m, 2H)
48	E-4-[1-(thiophen-2-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 9.11 (brs, 1H), 7.97 (dd, <i>J</i> = 4.8, 1.2 Hz, 1H), 7.59 (dd, <i>J</i> = 3.8, 1.4 Hz, 1H), 7.17-7.20 (m, 2H), 7.04 (brs, 1H), 6.75 (brs, 1H), 3.68 (d, <i>J</i> = 7.2 Hz, 1H), 1.74-1.91 (m, 11H), 1.45 (d, <i>J</i> = 11.6 Hz, 2H), 1.14-1.21 (m, 2H), 0.92-0.95 (m, 2H)
49	E-4-[3-(6-trifluoromethylpyridin-2-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) δ 8.36 (d, <i>J</i> = 8.4 Hz, 1H), 7.88 (d, <i>J</i> = 8.0 Hz, 1H), 6.82-6.84 (m, 2H), 5.78 (s, 1H), 5.58 (brs, 1H), 5.25 (brs, 1H), 3.93-3.95 (m, 1H), 2.08 (s, 1H), 1.83-2.04 (m, 12H), 1.48-1.51 (m, 2H), 0.93-0.96 (m, 2H)
50	E-4-[1-(1-methyl-1H-indole-7-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, MeOD) δ 7.85 (dd, <i>J</i> = 7.8, 1.0 Hz, 1H), 7.68 (d, <i>J</i> = 4 Hz, 1H), 7.64 (dd, <i>J</i> = 7.7, 0.8 Hz, 1H), 7.27 (d, <i>J</i> = 3.2 Hz, 1H), 7.13 (t, <i>J</i> = 7.8 Hz, 1H), 6.62 (d, <i>J</i> = 3.2 Hz, 1H), 4.19 (s, 3H), 3.95-3.93 (m, 1H), 2.09-1.92 (m, 22H), 1.65 (d, <i>J</i> = 11.2 Hz, 2H), 1.32-1.29 (m, 2H), 0.94-0.91 (m, 2H)

51	1-(3-chloro-benzenesulfonylamino)-N-(4-fluoro-2-(trifluoromethyl)phenyl)cyclopropanecarboxamide		¹ H NMR (400 MHz, CDCl ₃) d 8.84 (s, 1H), 7.98-8.01 (m, 1H), 7.84 (s, 1H), 7.75 (d, J = 7.6 Hz, 1H), 7.55 (d, J = 8.0 Hz, 1H), 7.45 (t, J = 8.0 Hz, 1H), 7.32 (dd, J = 8.4, 2.8 Hz, 1H), 7.22-7.26 (m, 1H), 6.57 (s, 1H), 1.50-1.53 (m, 2H), 0.94-0.97 (m, 2H)
52	E-4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 10.30 (brs, 1H), 8.58 (brs, 1H), 7.73-7.77 (m, 3H), 7.62-7.66 (m, 1H), 7.15 (d, J = 7.6 Hz, 1H), 3.60 (d, J = 7.2 Hz, 1H), 1.74-1.90 (m, 11H), 1.45 (d, J = 12.0 Hz, 2H), 1.15-1.17 (m, 2H), 0.88-0.91 (m, 2H)
53	1-(3-chloro-benzenesulfonylamino)-N-[4-(1,1,1-trifluoro-2-hydroxyprop-2-yl)phenyl]cyclopropanecarboxamide		¹ H NMR (400 MHz, DMSO) d 9.44 (s, 1H), 7.77 (s, 1H), 7.72-7.74 (m, 1H), 7.52-7.53 (m, 2H), 7.45-7.47 (m, 2H), 7.39-7.41 (m, 2H), 1.66 (s, 3H), 1.25-1.28 (m, 2H), 0.99-1.02 (m, 2H)
54	E-4-[1-(3-chloro-benzeneamino)cyclopropanecarboxamido]-		¹ H NMR (400 MHz, DMSO) d 9.09 (s, 1H) 7.92 (t, J = 1.6 Hz, 1H), 7.82 (d, J = 8 Hz, 1H), 7.65-7.63 (m, 1H), 7.53 (t, J = 8 Hz, 1H), 7.07 (d, J =


	adamantan-1-carboxylic acid amide		7.2 Hz, 1H), 7.01 (s, 1H), 6.72 (s, 1H), 3.76-3.74 (m, 1H), 1.91-1.65 (m, 11H), 1.40 (d, $J = 12.8$ Hz, 2H), 1.34-1.31 (m, 2H), 1.04-1.01 (m, 2H)
55	N-(bicyclo[2.2.1]heptan-2-yl)-1-(3-chlorobenzenesulfonylamino)cyclopropanecarboxamide		^1H NMR (400 MHz, CDCl_3) d 7.85 (t, $J = 1.8$ Hz, 1H), 7.75 (dt, $J = 1.2$, 7.6 Hz, 1H), 7.56-7.59 (m, 1H), 7.47 (t, $J = 8.0$ Hz, 1H), 6.90 (d, $J = 7.2$ Hz, 1H), 6.56 (s, 1H), 3.97-4.03 (m, 1H), 2.37 (s, 1H), 2.22 (s, 1H), 2.00-2.07 (m, 1H), 1.53-1.63 (m, 2H), 1.38-1.47 (m, 4H), 1.31-1.34 (m, 1H), 1.23-1.26 (m, 1H), 0.85-0.94 (m, 2H), 0.73-0.78 (m, 1H)
56	E-4-[1-(1,1-dioxido[1,2,3]benzodisothiazol-2(3H)-yl)cyclopropanecarboxamido]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.83 (d, $J = 8.0$ Hz, 1H), 7.65-7.72 (m, 2H), 7.59 (t, $J = 7.6$ Hz, 1H), 7.45 (t, $J = 7.6$ Hz, 1H), 5.58 (brs, 1H), 5.37 (brs, 1H), 4.53 (s, 2H), 4.02 (d, $J = 7.2$ Hz, 1H), 1.94-2.03 (m, 10 H), 1.84 (s, 2H), 1.66-1.71 (m, 3H), 1.48 (t, $J = 12.4$ Hz, 2H)
57	E-4-[1-(3,4-difluorobenzeneamino)cyclopropanecarboxamido]-adamantan-		^1H NMR (400 MHz, DMSO) d 9.07 (s, 1H), 7.95 (t, $J = 9.6$ Hz, 1H), 7.79-7.76 (m, 1H), 7.62-7.55 (m, 1H), 7.06 (d, $J = 7.6$ Hz, 1H), 6.99 (s, 1H), 6.72 (s, 1H), 3.77-3.75 (m, 1H), 1.89-1.66 (m,

	1-carboxylic acid amide		11H), 1.40 (d, J = 12.4 Hz, 2H), 1.34-1.32 (m, 2H), 1.03-1.00 (m, 2H)
58	E-4-[1-(1-methyl-1H-pyrazole-5-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 7.55 (d, J = 2 Hz, 1H), 7.16 (d, J = 8 Hz, 1H), 7.03 (s, 1H), 6.74 (d, J = 2.4 Hz, 2H), 3.95 (s, 3H), 3.72 (d, J = 7.2 Hz, 1H), 1.93 (s, 1H), 1.83-1.81 (m, 8H), 1.76-1.75 (m, 2H), 1.49 (d, J = 12.4 Hz, 2H), 1.20-1.17 (m, 2H), 0.88-0.85 (m, 2H)
59	E-4-[1-(2,3-difluorobenzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 9.15 (s, 1H), 7.64-7.57 (m, 1H), 7.42-7.39 (m, 1H), 7.35-7.30 (m, 1H), 7.03-7.01 (m, 2H), 6.73 (s, 1H), 3.81-3.79 (m, 1H), 1.90-1.74 (m, 11H), 1.47 (d, J = 12.4 Hz, 2H), 1.35-1.33 (m, 2H), 1.04-1.01 (m, 2H)
60	E-4-[1-(benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 8.87 (s, 1H), 7.78 (d, J = 7.6 Hz, 2H), 7.69-7.65 (m, 1H), 7.62-7.59 (m, 2H), 7.21 (d, J = 7.6 Hz, 1H), 7.01 (s, 1H), 6.75 (s, 1H), 3.68-3.66 (m, 1H), 1.92 (s, 1H), 1.82-1.75 (m, 10H), 1.46 (d, J = 12 Hz, 2H), 1.15-1.12 (m, 2H), 0.83-0.80 (m, 2H)

61	E-4-[1-(2-fluoro- <i>N</i> -methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 7.84 (td, <i>J</i> = 7.6, 1.6 Hz, 1H), 7.82-7.76 (m, 1H), 7.52-7.48 (m, 1H), 7.43 (td, <i>J</i> = 8, 0.8 Hz, 1H), 7.02 (s, 1H), 7.00 (d, <i>J</i> = 7.6 Hz, 1H), 6.75 (s, 1H), 3.82-3.80 (m, 1H), 2.99 (d, <i>J</i> = 1.2 Hz, 3H), 1.93-1.77 (m, 11H), 1.53 (d, <i>J</i> = 12.8 Hz, 2H), 1.39-1.36 (m, 2H), 1.18-1.16 (m, 2H)
62	E-4-[1-(3-chloro- <i>N</i> -methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 7.80-7.76 (m, 3H), 7.68-7.64 (m, 1H), 7.02 (s, 1H), 6.96 (d, <i>J</i> = 7.2 Hz, 1H), 6.74 (s, 1H), 3.79-3.77 (m, 1H), 2.95 (s, 3H), 1.93-1.77 (m, 11H), 1.51 (d, <i>J</i> = 12.8 Hz, 2H), 1.41-1.38 (m, 2H), 1.21-1.19 (m, 2H)
63	E-4-[1-(2-fluorobenzamido)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 9.06 (s, 1H), 7.56 (m, 2H), 7.32 (m, 2H), 7.03 (m, 2H), 6.74 (brs, 1H), 3.78 (m, 1H), 1.88-1.74 (m, 10H), 1.45 (d, <i>J</i> = 12.4 Hz, 2H), 1.32 (m, 2H), 1.24 (s, 1H), 1.01 (m, 2H)
64	E- <i>N</i> -[1-(5-carbamoyladamantan-2-yl)carbamoyl]cyclopropanecarboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 9.62 (s, 1H), 9.08 (s, 1H), 8.44 (d, <i>J</i> = 7.6 Hz, 1H), 8.21 (d, <i>J</i> = 8.4 Hz, 1H), 7.07 (d, <i>J</i> = 7.2 Hz, 1H), 6.99 (s, 1H)

	opyl)-5-(trifluoromethyl)pyrrolineamide		1H), 6.73 (s, 1H), 3.76 (m, 1H), 1.97-1.64 (m, 11H), 1.34 (m, 4H), 1.02 (m, 2H)
65	E-4-(1-(benzo[d][1,3]dioxol-5-sulfonylamino)cyclopropanecarboxamino)adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 8.70 (brs, 1H), 7.31 (dd, J = 6.4, 2.0 Hz, 1H), 7.19 (m, 2H), 7.08 (d, J = 8.4 Hz, 1H), 7.01 (brs, 1H), 6.75 (brs, 1H), 6.75 (s, 1H), 3.68 (m, 1H), 1.91-1.75 (m, 11H), 1.45 (d, J = 12.0 Hz, 2H), 1.13 (m, 2H), 0.87 (m, 2H)
66	1-(3-chlorobenzenesulfonylamino)-N-cycloheptyl-N-propylcyclopropanecarboxamide		¹ H NMR (400 MHz, CDCl ₃) δ 7.88 (t, J = 2.0 Hz, 1H), 7.84 (m, 1H), 7.55 (m, 1H), 7.46 (d, J = 7.6 Hz, 1H), 5.07 (brs, 1H), 4.06/2.98 (m, 2H), 1.73-1.23 (m, 21H)

Preparational Example 19: Preparation of 3-azido-2,2-dimethylpropanoic acid




5 1 g (7.32 mmol) of 3-chloro-2,2-dimethylpropanoic acid was dissolved in 15.0 ml of water, to which 4.8 g (73.22 mmol) of NaN₃ was added, followed by stirring for 18 hours with high temperature reflux. The

reaction mixture was acidized with conc. HCl to be pH 3, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 1 g of 3-azido-2,2-dimethylpropanoic acid (yield: 95%).

^1H NMR (400 MHz, CDCl_3) δ 3.48 (s, 2H), 1.31 (s, 6H)


10 Preparational Example 20: Preparation of 3-amino-2,2-dimethylpropanoic acid



1 g (6.99 mmol) of the 3-azido-2,2-dimethylpropanoic acid prepared in Preparational Example 19 was dissolved in 100 ml of MeOH, to which 695 mg of 10% Pd/C was added, followed by stirring for 3 hours in the presence of hydrogen. 100 ml of water was added to the reaction mixture, followed by stirring for 30 minutes. The mixture was filtered by using celite filter and the resultant solution was evaporated under reduced pressure to give 753 mg of 3-amino-2,2-dimethylpropanoic acid (yield: 92%).

^1H NMR (400 MHz, DMSO) δ 2.65 (s, 2H), 1.30 (s, 2H), 1.01 (s, 6H)


Preparational Example 21: Preparation of 3-(t-butoxycarbonylamino)-2,2-dimethylpropanoic acid



256 mg (2.22 mmol) of the -amino-2,2-dimethylpropanoic acid prepared in Preparational Example 20 was dissolved in 2.0 ml of water and 3.0 ml of t-BuOH, to which 4.0 ml of 1.0 N NaOH solution and 725 mg (3.32 mmol) of Boc_2O were added. The mixture was stirred at room temperature for 16 hours, followed by evaporation under reduced pressure to eliminate the solvent. The residue was acidized with 1 N HCl to be pH 3, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 305 mg of 3-(t-butoxycarbonylamino)-2,2-dimethylpropanoic acid (yield: 63%).

^1H NMR (400 MHz, CDCl_3) δ 3.25 (s, 2H), 1.44 (s, 9H), 1.18 (s, 6H)

Preparational Example 22: Preparation of 4-(3-(t-butoxycarbonyl)amido)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid methyl ester



5


305 mg (1.40 mmol) of the 3-(t-butoxycarbonylamino)-2,2-dimethylpropanoic acid prepared in Preparational Example 21 and 379 mg (1.54 mmol) of the 4-amino-adamantane-1-carboxylic acid methyl ester prepared in Preparational Example 7 were dissolved in 5.0 ml of methylene chloride, to which 323 mg (1.38 mmol) of EDCI and 227 mg (1.68 mmol) of HOBT were added. 0.6 ml (4.21 mmol) of TEA was also added thereto. The reaction mixture was stirred at room temperature for 16 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 466 mg of 4-(3-(t-butoxycarbonyl)amido)-2,2-

20

dimethylpropanamido)adamantan-1-carboxylic acid methyl ester (yield: 95%).

^1H NMR (400 MHz, CDCl_3) δ 6.11 (brs, 1H), 5.12 (brs, 1H), 3.98 (m, 1H), 3.67 (s, 3H), 3.24 (d, J = 6.4 Hz, 1H), 2.05-1.74 (m, 10H), 1.63 (s, 3H), 1.43 (s, 9H), 1.21 (s, 6H)

Preparational Example 23: Preparation of 4-(3-amido-2,2-dimethylpropanamido)adamantan-1-carboxylic acid methyl ester hydrochloride



466 mg (1.33 mmol) of the 4-(3-(t-butoxycarbonyl)amido)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 22 was dissolved in 6 ml of ethyl acetate, to which 3.32 ml of 4 M HCl 1,4-dioxane solution was added, followed by stirring at room temperature for 16 hours. Upon completion of the reaction the reaction solvent was eliminated by evaporation under reduced pressure to give 422 mg of 4-(3-amido-2,2-


dimethylpropanamido)adamantan-1-carboxylic acid methyl ester hydrochloride (yield: 92%).

^1H NMR (400 MHz, CDCl_3) δ 8.73 (brs, 2H), 5.98 (m, 1H), 3.96 (m, 1H), 3.72 (s, 2H), 3.66 (s, 3H), 2.04-
5 1.91 (m, 9H), 1.66 (s, 4H), 1.39 (s, 6H)

Preparational Example 24: Preparation of 4-(3-(3-chloro-benzenesulfonylamino)-2,2-

dimethylpropanamido)adamantan-1-carboxylic acid methyl

10 ester




80 mg (0.23 mmol) of the 4-(3-amido-2,2-dimethylpropanamido)adamantan-1-carboxylic acid methyl ester hydrochloride prepared in Preparational Example
15 23 was dissolved in 1 ml of methylene chloride, to which 0.1 ml of TEA was added, followed by stirring at room temperature for 5 minutes. 59 mg (0.28 mmol) of 3-chlorosulfonylchloride was added to the reactant, followed by stirring at room temperature for 16 hours.
20 The reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in 20 ml of water, followed by extraction with 350 ml of

ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 101 g of 4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid methyl ester (yield: 91%).

^1H NMR (400 MHz, CDCl_3) δ 7.83 (s, 1H), 7.72 (d, J = 7.6 Hz, 1H), 7.53 (d, J = 8.4 Hz, 1H), 7.45 (t, J = 8.0 Hz, 1H), 5.88 (m, 1H), 5.57 (t, J = 6.4 Hz, 1H), 3.92 (m, 1H), 3.66 (s, 3H), 2.95 (d, J = 6.4 Hz, 2H), 1.99 (s, 8H), 1.85 (s, 2H), 1.65 (m, 3H), 1.26 (s, 6H)

Preparational Example 25: Preparation of 4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid




100 mg (0.21 mmol) of the 4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 24 was dissolved in 1.0 ml of THF/ethanol mixed solution (1:1), to which 1.0 ml of 2 N NaOH solution was added,

followed by stirring at room temperature overnight. The reactant was acidized with 1 N HCl, followed by extraction with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by
 5 evaporation under reduced pressure to give 98 mg of 4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid (yield: 99%).

^1H NMR (400 MHz, CDCl_3) δ 7.85 (t, J = 1.6 Hz, 1H), 7.73 (m, 1H), 7.54 (m, 1H), 7.43 (m, 1H), 6.00 (m, 2H), 3.92 (d, J = 6.8 Hz, 1H), 2.95 (d, J = 6.0 Hz, 1H), 2.00 (s, 7H), 1.93 (s, 2H), 1.66 (m, 4H), 1.26 (s, 6H)

15 Example 67: Preparation of E-4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid amide



98 mg (0.21 mmol) of the 4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid
 20 prepared in Preparational Example 25 was dissolved in




1.0 ml of methylene chloride, to which 44 mg (0.23 mmol) of HOBt and 31 mg (0.23 mmol) of EDCI were added. The mixture was stirred at room temperature for 10 minutes and then 1.0 ml of 35% ammonia water
 5 was added thereto. The mixture was stirred at room temperature for 20 hours. Upon completion of the reaction, extraction was performed with methylene chloride. After washing with brine, the extract was dried over anhydrous magnesium sulfate, followed by
 10 column chromatography to give 20.7 mg of 4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid amide (yield: 21%).

¹H NMR (400 MHz, CDCl₃) δ 7.84 (s, 1H), 7.73 (d, *J*
 15 = 7.6 Hz, 1H), 7.55 (d, *J* = 9.2 Hz, 1H), 7.45 (m, 1H), 5.96 (d, *J* = 7.2 Hz, 1H), 5.80 (m, 1H), 5.68 (s, 1H), 3.95 (s, 1H), 2.95 (d, *J* = 6.4 Hz, 1H), 1.62-2.04 (m, 11H), 1.26 (s, 6H)


20 The compounds of Example 68 - Example 70 were prepared by the preparation method described in Example 67.

【Table 3】

Ex am pl	Compound Name	Structure	NMR data

e			
68	E-4-[3-(2-fluorobenzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) δ 7.79 (m, 1H), 7.68 (m, 2H), 7.41 (m, 2H), 6.99 (s, 1H), 6.82 (s, 1H), 6.72 (s, 1H), 3.71 (m, 1H), 2.98 (s, 1H), 1.91-1.38 (m, 13H), 1.11 (s, 6H)
69	E-4-[3-(benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) δ 7.84 (m, 2H), 7.57 (m, 3H), 6.16 (brs, 1H), 5.98 (d, J = 7.6 Hz, 1H), 5.69 (brs, 1H), 5.59 (m, 1H), 3.93 (d, J = 4.8 Hz, 1H), 2.94 (d, J = 6.8 Hz, 1H), 2.04-1.80 (m, 9H), 1.73-1.61 (m, 4H), 1.24 (s, 6H)
70	E-4-[3-(3-chloro-2-methylbenzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) δ 7.84 (dd, J = 6.8, 0.8 Hz, 1H), 7.56 (d, J = 4.8 Hz, 1H), 7.25 (m, 1H), 5.86 (m, 4H), 3.95 (d, J = 7.6 Hz, 1H), 2.91 (d, J = 6.4 Hz, 2H), 2.75 (s, 3H), 1.59-2.05 (m, 11H), 1.23 (s, 6H)


Preparational Example 26: Preparation of 2-(t-butoxycarbonylamino)-2-methylpropanoic acid



200 mg (1.94 mmol) of 2-amino-2-methylpropanoic acid was dissolved in 8.0 ml of 1.0 N NaOH solution and 8.0 ml of 1,4-dioxane solution, to which 846 mg
 5 (3.88 mmol) of Boc_2O was added. The mixture was stirred at room temperature for 16 hours, and then 1,4-dioxane was eliminated by evaporation under reduced pressure. The residue was acidized with 1 N HCl to be pH 3, and followed by extraction with 320 ml
 10 of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 315 mg of 2-(tert-butoxycarbonylamino)-2-methylpropanoic acid (yield: 80%).

15 ^1H NMR (400 MHz, CDCl_3) δ 5.07 (brs, 1H), 1.54 (s, 6H), 1.45 (s, 9H)


Preparational Example 27: Preparation of 4-(2-((tert-butoxycarbonyl)amino)-2-methylpropanamido)adamantan-1-
 20 carboxylic acid methyl ester



380 mg (1.87 mmol) of the 2-(t-butoxycarbonylamino)-2-methylpropanoic acid prepared in Preparational Example 26 and 505 mg (2.06 mmol) of the 4-amino-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 7 were dissolved in 10.0 ml of methylene chloride, to which 430 mg (2.24 mmol) of EDCI and 303 mg (2.24 mmol) of HOBt were added. 0.8 ml (5.61 mmol) of TEA was also added thereto. The reaction mixture was stirred at room temperature for 16 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 580 mg of 4-(2-((t-butoxycarbonyl)amino)-2-methylpropanamido)adamantan-1-carboxylic acid methyl ester (yield: 92%).

^1H NMR (400 MHz, CDCl_3) δ 5.30 (s, 1H), 4.84 (brs, 1H), 3.98 (brs, 1H), 3.66 (s, 3H), 2.02-1.44 (m, 28H)

Preparational Example 28: Preparation of 4-(2-amino-2-methylpropanamido)adamantan-1-carboxylic acid methyl ester hydrochloride




5
580 mg (1.72 mmol) of the 4-(2-((t-butoxycarbonyl)amino)-2-methylpropanamido)adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 27 was dissolved in 6 ml of ethyl acetate, to
10 which 4.31 ml of 4 M HCl 1,4-dioxane was added. The mixture was stirred at room temperature for 16 hours. Upon completion of the reaction, the reaction solvent was eliminated by evaporation under reduced pressure to give 495 mg of 4-(2-amino-2-methylpropanamido)adamantan-1-carboxylic acid methyl
15 ester hydrochloride (yield: 87%).

^1H NMR (400 MHz, CDCl_3) δ 8.80 (brs, 2H), 6.48 (brs, 1H), 4.01 (brs, 1H), 3.66 (s, 3H), 2.09-1.60 (m, 19H)

20

Preparational Example 29: Preparation of 4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid methyl ester



5 100 mg (0.3 mmol) of the 4-(2-amino-2-methylpropanamido)adamantan-1-carboxylic acid methyl ester hydrochloride prepared in Preparational Example 28 was dissolved in 1 ml of methylene chloride, to which 0.13 ml of TEA was added, followed by stirring

10 at room temperature for 5 minutes. 70 mg (0.36 mmol) of 3-chlorosulfonylchloride was added to the reactant, followed by stirring at room temperature for 16 hours. The reaction solvent was eliminated by evaporation under reduced pressure. The residue was dissolved in


15 20 ml of water, followed by extraction with 350 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give

107 g of 4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid methyl

20 ester (yield: 78%).

^1H NMR (400 MHz, CDCl_3) δ 7.89 (td, $J_1 = 1.6$ Hz, $J_2 = 6.0$ Hz, 1H), 7.60 (m, 1H), 7.28 (m, 2H), 7.00 (d, $J = 9.2$ Hz, 1H), 3.97 (m, 1H), 3.66 (s, 3H), 2.03 (m, 7H), 1.88 (m, 4H), 1.62 (d, $J = 13.2$ Hz, 2H), 1.39 (s, 6H)

Preparational Example 30: Preparation of 4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid



10 105 mg (0.23 mmol) of the 4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 29 was dissolved in 1.0 ml of THF/ethanol mixed solution (1:1), to which 1.0 ml of 2 N NaOH solution was added, followed by stirring at room temperature overnight. The reactant was acidized with 1 N HCl, followed by extraction with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give 100 mg of 4-[2-(2-fluoro-


15

20

benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid (yield: 99%).

^1H NMR (400 MHz, CDCl_3) δ 7.89 (td, $J_1 = 2.0$ Hz, $J_2 = 6.0$ Hz, 1H), 7.60 (m, 1H), 7.28 (m, 2H), 7.03 (d, $J = 8.0$ Hz, 1H), 3.98 (d, $J = 8.0$ Hz, 1H), 2.05 (m, 7H), 1.95 (s, 2H), 1.88 (d, $J = 12.8$ Hz, 2H), 1.45 (d, $J = 12.0$ Hz, 2H), 1.39 (s, 6H)

Example 71: Preparation of E-4-[2-(2-fluorobenzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid amide




100 mg (0.23 mmol) of the 4-[2-(2-fluorobenzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid prepared in Preparational Example 30 was dissolved in 7.0 ml of methylene chloride, to which 34 mg (0.25 mmol) of HOBt and 48 mg (0.25 mmol) of EDCI were added. The mixture was stirred at room temperature for 10 minutes and then 7.0 ml of 35% ammonia water was added thereto. The mixture was stirred at room temperature for 20 hours. Upon completion of the reaction, extraction was performed




with methylene chloride. After washing with brine, the extract was dried over anhydrous magnesium sulfate, followed by column chromatography to give 15 mg of 4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid amide (yield: 15%).




^1H NMR (400 MHz, CDCl_3) δ 7.89 (td, $J_1 = 1.6$ Hz, $J_2 = 6.0$ Hz, 1H), 7.61 (m, 1H), 7.36 (d, $J = 4.8$ Hz, 1H), 7.30 (m, 1H), 7.24 (m, 1H), 7.03 (d, $J = 7.6$ Hz, 1H), 5.57 (brs, 1H), 3.97 (d, $J = 7.6$ Hz, 1H), 2.09 (s, 3H), 2.00 (m, 4H), 1.89 (m, 4H), 1.63 (d, $J = 12.4$ Hz, 2H), 1.39 (s, 6H)




The compounds of Example 72 - Example 90 were prepared by the preparation method described in Example 71.




【Table 4】




Ex am pl e	Compound Name	Structure	NMR data
72	E-4-[2-(2-fluoro- <i>N</i> -methyl-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-		^1H NMR (400 MHz, CDCl_3) δ 7.93 (m, 1H), 7.61 (m, 1H), 7.31 (td, $J_1 = 0.8$ Hz, $J_2 = 7.6$ Hz, 1H), 7.27 (m, 2H), 7.06 (d, $J = 8.0$ Hz, 1H), 5.62 (brs, 2H), 4.00 (d, $J =$




	carboxylic acid amide		8.4 Hz, 1H), 3.05 (d, $J = 1.6$ Hz, 3H), 2.18-1.50 (m, 13H), 1.46 (s, 6H)
73	E-4-[2-(3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.87 (t, $J = 1.6$ Hz, 1H), 7.76 (m, 1H), 7.55 (m, 1H), 6.74 (d, $J = 7.2$ Hz, 1H), 5.56 (brs, 1H), 5.19 (brs, 1H), 3.95 (d, $J = 7.2$ Hz, 1H), 2.07 (s, 3H), 2.00 (s, 4H), 1.92 (s, 2H), 1.82 (d, $J = 15.2$ Hz, 2H), 1.63 (d, $J = 14.0$ Hz, 2H), 1.44 (s, 6H)
74	E-N-(5-cyanoadamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide		^1H NMR (400 MHz, CDCl_3) d 7.88 (td, $J_1 = 1.6$ Hz, $J_2 = 4.0$ Hz, 1H), 7.62 (m, 1H), 7.32 (m, 2H), 7.10 (d, $J = 7.6$ Hz, 1H), 5.28 (s, 1H), 3.99 (d, $J = 7.6$ Hz, 1H), 2.16 (s, 4H), 2.09 (m, 5H), 1.91 (d, $J = 13.2$ Hz, 2H), 1.64 (d, $J = 13.6$ Hz, 2H), 1.34 (s, 6H)
75	E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(N'-carbamimidoyl)adamanta		^1H NMR (400 MHz, CDCl_3) d 7.89 (td, $J_1 = 1.6$ Hz, $J_2 = 6.0$ Hz, 1H), 7.61 (m, 1H), 7.30 (td, $J_1 = 0.8$ Hz, $J_2 = 6.8$ Hz, 1H), 7.25

	n-2-yl]-2-methylpropa namide		(m, 1H), 7.03 (d, $J = 8.0$ Hz, 1H), 5.39 (s, 1H), 4.55 (s, 2H), 3.97 (m, 1H), 2.10-1.84 (m, 11H), 1.61 (d, $J = 13.2$ Hz, 2H), 1.39 (s, 6H)
76	E-2-(2-fluoro-benzenesulfonfylamino)-N-[5-(hydroxymethyl)adamantan-2-yl]-2-methylpropa namide		^1H NMR (400 MHz, CDCl_3) d 7.89 (t, $J = 7.2$ Hz, 1H), 7.61 (m, 1H), 7.26 (m, 2H), 7.05 (d, $J = 8.0$ Hz, 1H), 5.74 (s, 1H), 3.92 (m, 1H), 3.24 (s, 2H), 2.02 (s, 3H), 1.86 (d, $J = 13.2$ Hz, 2H), 1.61 (m, 8H), 1.39 (s, 6H)
77	E-2-(2-fluoro-benzenesulfonfylamino)-N-(5-formyladama ntan-2-yl)-2-methylpropa namide		^1H NMR (400 MHz, CDCl_3) d 9.37 (s, 1H), 7.90 (td, $J_1 = 2.0$ Hz, $J_2 = 5.6$ Hz, 1H), 7.61 (m, 1H), 7.30 (m, 2H), 7.11 (d, $J = 7.6$ Hz, 1H), 5.55 (s, 1H), 3.97 (m, 1H), 2.13-1.61 (m, 13H), 1.39 (s, 6H)
78	E-[4-(2-(2-fluoro-benzenesulfonfylamino)-2-methylpropa namido)adam antan-1-yl]methyl-4-methylbenze		^1H NMR (400 MHz, CDCl_3) d 7.88 (td, $J_1 = 1.6$ Hz, $J_2 = 6.0$ Hz, 1H), 7.77 (d, $J = 8.4$ Hz, 2H), 7.59 (m, 1H), 7.33 (d, $J = 8.0$ Hz, 2H), 7.30 (m, 2H), 6.97 (d, $J = 8.0$ Hz, 1H), 5.34 (s, 1H), 3.86 (d,


	nesulfonate		$J = 7.6 \text{ Hz, 1H), 3.60 (s, 2H), 2.46 (s, 3H), 1.99 (s, 3H), 1.89 (d, } J = 12.0 \text{ Hz, 2H), 1.55 (m, 8H), 1.37 (s, 6H)}$
79	E-2-[4-(2-(2-fluorobenzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]acetic acid		$^1\text{H NMR (400 MHz, MeOD) d 7.75 (m, 1H), 7.65 (d, } J = 4.4 \text{ Hz, 1H), 7.46 (m, 1H), 7.00 (m, 2H), 3.86 (d, } J = 4.8 \text{ Hz, 1H), 2.11-1.16 (m, 13H), 1.34 (s, 6H)}$
80	E-N-[5-(2-amino-2-oxoethyl)adamantan-2-yl]-2-(2-fluorobenzenesulfonylamino)-2-methylpropanamide		$^1\text{H NMR (400 MHz, DMSO) d 8.43 (s, 1H), 7.81 (m, 1H), 7.71 (m, 1H), 7.41 (m, 2H), 7.18 (m, 2H), 6.69 (s, 1H), 3.61 (d, } J = 7.6 \text{ Hz, 1H), 3.32 (s, 2H), 1.89-1.34 (m, 13H), 1.24 (s, 6H)}$
81	E-4-[2-methyl-2-(benzenesulfonylamino)propanamido]adamantan-1-carboxylic acid amide		$^1\text{H NMR (400 MHz, CDCl}_3\text{) d 7.89 (m, 2H), 7.56 (m, 1H), 7.51 (m, 2H), 7.01 (d, } J = 7.6 \text{ Hz, 1H), 5.71 (s, 1H), 5.64 (brs, 1H), 5.54 (brs, 1H), 3.96 (m, 1H), 2.08 (s, 3H), 1.99 (s, 4H), 1.90 (m, 4H), 1.61 (d, } J = 14.4 \text{ Hz, 2H), 1.40 (s, 6H)}$

82	E-4-[2-(2-fluoro-3-chlorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.80-7.77 (m, 1H), 7.67-7.63 (m, 1H), 7.25-7.23 (m, 1H), 6.94 (d, <i>J</i> = 8.0 Hz, 1H), 5.72 (s, 1H), 5.62 (s, 1H), 5.46 (s, 1H), 3.99-3.93 (m, 1H), 2.09-1.87 (m, 11H), 1.66 (d, <i>J</i> = 13.2 Hz, 3H), 1.42 (s, 6H)
83	E-4-[2-(3,5-difluorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.45-7.40 (m, 2H), 7.06-7.00 (m, 1H), 6.65 (d, <i>J</i> = 8.0 Hz, 1H), 5.66 (s, 1H), 5.59 (s, 1H), 5.32 (s, 1H), 3.98 (<i>J</i> = 8.0 Hz, 1H), 2.08 (s, 3H), 2.00 (s, 4H), 1.92 (d, <i>J</i> = 2.0 Hz, 2H), 1.84 (d, <i>J</i> = 13.2 Hz, 2H), 1.66 (d, <i>J</i> = 13.2 Hz, 2H), 1.47 (s, 6H)
84	E-4-[2-(2,6-difluorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 7.58-7.51 (m, 1H), 7.06 (t, 2H), 6.95 (d, <i>J</i> = 8.0 Hz, 1H), 5.88 (s, 1H), 5.62 (s, 1H), 5.42 (s, 1H), 3.98 (d, <i>J</i> = 8.0 Hz, 1H), 2.09 (s, 3H), 1.09-1.86 (m, 8H), 1.65 (d, <i>J</i> = 11.6 Hz, 3H), 1.53 (s, 3H)

85	E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide		¹ H NMR (400 MHz, DMSO) δ 8.77 (s, 1H), 7.54 (dd, <i>J</i> = 6.4, 1.4 Hz, 1H), 7.44-7.36 (m, 2H), 7.16 (d, <i>J</i> = 7.6 Hz, 1H), 6.93 (s, 1H), 6.69 (t, <i>J</i> = 7.4 Hz, 1H), 4.34 (s, 2H), 4.15 (s, 1H), 3.68-3.66 (m, 1H), 1.93-1.77 (m, 11H), 1.49 (d, <i>J</i> = 12 Hz, 2H), 1.21 (s, 6H)
86	E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide		¹ H NMR (400 MHz, CDCl ₃) δ 7.84-7.91 (m, 1H), 7.57-7.62 (m, 1H), 7.25-7.31 (m, 1H), 7.21-7.33 (m, 1H), 7.01 (d, <i>J</i> = 7.6 Hz, 1H), 5.92 (s, 1H), 5.63 (brs, 1H), 5.61 (brs, 1H), 3.90 (t, <i>J</i> = 4.0 Hz, 1H), 2.17-2.21 (m, 2H), 1.99 (s, 3H), 1.85 (d, <i>J</i> = 13.2 Hz, 2H), 1.46-1.58 (m, 10H), 1.39 (s, 6H)
87	E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide		¹ H NMR (400 MHz, DMSO) δ 8.42 (s, 1H), 8.12 (s, 1H), 7.79-7.83 (m, 1H), 7.69-7.74 (m, 1H), 7.43-7.48 (m, 1H), 7.36-7.43 (m, 1H), 7.20 (d, <i>J</i> = 7.2 Hz, 1H), 3.64 (d, <i>J</i> = 6.8 Hz, 1H), 1.82-2.09 (m, 11H),

			1.44 (d, $J = 12.8$ Hz, 2H), 1.24 (s, 6H)
88	E-4-[2-(4-chlorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 8.26 (s, 1H), 7.82-7.79 (m, 2H), 7.67-7.65 (m, 2H), 7.07 (d, $J = 7.6$ Hz, 1H), 7.02 (s, 1H), 6.75 (s, 1H), 3.57-3.55 (m, 1H), 1.90-1.75 (m, 11H), 1.46 (d, $J = 12$ Hz, 2H), 1.25 (s, 6H)
89	E-4-[2-(2,5-dichlorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 8.50 (s, 1H), 7.95 (d, $J = 2$ Hz, 1H), 7.77-7.72 (m, 2H), 7.18-7.14 (m, 1H), 7.01 (s, 1H), 6.34 (s, 1H), 3.75-3.73 (m, 1H), 1.93-1.77 (m, 11H), 1.49 (d, $J = 9.6$ Hz, 2H), 1.25 (s, 6H)
90	2-(3-chlorobenzenesulfonylamino)-N-cycloheptyl-2-methyl-N-propylpropanamide		^1H NMR (400 MHz, CDCl_3) d 7.89 (s, 1H), 7.81 (d, $J = 8.0$ Hz, 1H), 7.50-7.48 (m, 1H), 7.43 (t, 1H), 6.73 (s, 1H), 3.05-3.00 (m, 1H), 2.79 (t, 2H), 2.08-2.01 (m, 2H), 1.78-1.60 (m, 6H), 1.59-1.42 (m, 5H), 0.94 (t, 3H)

Preparational Example 31: Preparation of 4-(3-nitro-benzamido)-adamantan-1-carboxylic acid methyl ester




33 mg (0.2 mmol) of 3-nitrobenzoic acid and 50 mg
 5 (0.2 mmol) of 4-amino-adamantane-1-carboxylic acid
 methyl ester were dissolved in 5.0 ml of methylene
 chloride, to which 60 mg (0.3 mmol) of EDC and 40 mg
 (0.3 mmol) of HOBt were added. 0.3 ml of TEA was also
 added thereto. The reaction mixture was stirred at
 10 room temperature for 12 hours, followed by evaporation
 under reduced pressure to eliminate methylene
 chloride. The residue was dissolved in 10 ml of
 water, followed by extraction with 320 ml of ethyl
 acetate. The extract was dried over anhydrous
 15 magnesium sulfate, and evaporated under reduced
 pressure, followed by column chromatography to give 70
 mg of 4-(3-nitro-benzamido)-adamantan-1-carboxylic
 acid methyl ester (yield: 97%).

^1H NMR (400 MHz, CDCl_3) δ 8.56 (s, 1H), 8.37-8.35
 20 (m, 1H), 8.16-8.14 (m, 1H) 7.66 (t, J = 8.0 Hz, 1H),
 6.46 (d, J = 4.0 Hz, 1H), 4.27 -4.26 (m, 1H), 3.96 (s,

3H), 2.21 (s, 2H), 2.12-2.03 (m, 5H), 1.96 (s, 2H),
1.89-1.85 (m, 2H), 1.68-1.66 (m, 2H).


Preparational Example 32: Preparation of 4-(3-amino-
5 benzamido)-adamantan-1-carboxylic acid methyl ester



30 mg (0.08 mmol) of the 4-(3-nitro-benzamido)-
adamantan-1-carboxylic acid methyl ester prepared in
Preparational Example 31 was dissolved in 10 ml of
10 methanol, to which 3 mg of Pd/C was added. The
mixture was stirred at room temperature for 12 hours
in the presence of hydrogen, which was then filtered.
Methanol was eliminated by evaporation under reduced
pressure. The obtained compound proceeded to the next
15 reaction without purification.

^1H NMR (400 MHz, CDCl_3) δ 7.40 (t, J = 4.0 Hz,
1H), 7.15 (s, 1H), 7.09-7.07 (m, 2H), 6.84-6.81 (m,
1H), 6.36-6.35 (m, 1H), 4.24 (s, 1H), 3.70 (s, 1H),
2.18-1.85 (m, 9H), 1.88-1.85 (m, 2H) 1.69-1.61 (m,
20 2H).

Preparational Example 33: Preparation of 4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid methyl ester




5 30 mg (0.09 mmol) of the 4-(3-amino-benzamido)-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 32 was dissolved in 5.0 ml of methylene chloride, to which 14 mg (0.18 mmol) of pyridine was added. 20 mg (0.11 mmol) of 2-fluoro-
 10 benzenesulfonylchloride was also added thereto. The mixture was stirred at room temperature for 12 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 10 ml of water, followed by extraction
 15 with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 30 mg of 4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid methyl ester
 20 (yield: 70%).

^1H NMR (400 MHz, CDCl_3) δ 7.85 (s, 1H), 7.60-7.46 (m, 3H), 7.22 -7.09 (m, 2H) 7.01 (s, 1H), 6.33-6.28

(m, 2H), 4.19 (s, 1H), 3.68 (s, 3H), 2.14- 1.93 (m 9H), 1.78-1.65 (m, 4H).

Preparational Example 34: Preparation of 4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid




30 mg (0.06 mmol) of the 4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 33 was dissolved in 1.0 ml of methanol and 1.0 ml of water, to which 24 mg (0.6 mmol) of sodium hydroxide was added. The mixture was stirred at room temperature for 12 hours, followed by evaporation under reduced pressure to eliminate methanol. The residue was acidized with 1 N HCl, followed by extraction with 320 ml of chloroform. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure. The obtained compound proceeded to the next reaction without purification.

^1H NMR (400 MHz, CDCl_3) δ 7.85-7.80 (m, 1H), 7.60 (s, 1H), 7.50 -7.45 (m, 1H) 7.35-7.21 (m, 4H), 7.11 (s, 1H), 6.30 (d, $J = 4.0$ Hz, 1H), 4.20 (s, 1H), 2.20-2.00 (m 9H), 1.89-1.82 (m, 2H), 1.72-1.69 (m, 2H).

5

Example 91: Preparation of E-4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide




10 15 mg (0.03 mmol) of the 4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid prepared in Preparational Example 34 was dissolved in 5.0 ml of methylene chloride, to which 9 mg (0.045 mmol) of EDC and 6 mg (0.045 mmol) of HOBT were added. Then, ammonia water was also added thereto. The mixture was stirred at room temperature for 12 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of methylene chloride. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced





pressure, followed by column chromatography to give 6 mg of 4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide (yield: 43%).




¹H NMR (400 MHz, CDCl₃) δ 8.53 (s, 1H), 7.89-7.80 (m, 2H), 7.50 (s, 1H) 7.46-7.32 (m, 2H), 7.28-7.12 (m, 3H), 6.42 (s, 1H), 5.75 (s, 1H), 5.30 (s, 1H), 4.12 (s, 1H), 2.17-1.94 (m, 9H), 1.84-1.81 (m, 2H), 1.69-1.66 (m, 2H).





10 The compounds of Example 92 - Example 139 were prepared by the preparation method described in Example 91.




【Table 5】




Ex am pl e	Compound Name	Structure	NMR data
92	E-4-[2-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) δ 11.75 (brs, 1H), 7.95 (brs, 1H), 7.63-7.61 (m, 1H), 7.42 (s, 1H), 7.38-7.23 (m, 2H), 7.17-7.05 (m, 3H) 6.37 (s, 1H), 5.39 (s, 1H), 5.31 (s, 1H), 4.20 (s, 1H), 2.30-1.81 (m, 9H), 1.67-1.63 (m, 4H)





93	E-4-[4-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, MeOD) d 7.92 (s, 1H), 7.90-7.80 (m, 3H), 7.63-7.16 (m, 6H), 6.61 (s, 1H), 4.12 (s, 1H), 2.29-1.83 (m, 1H), 1.65-1.50 (m, 2H)
94	E-4-[3-(4-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 8.70 (s, 1H), 7.84-7.76 (m, 3H), 7.55-7.50 (m, 1H), 7.41-7.32 (m, 2H) 7.11-7.05 9m, 2H0, 6.40 (d, J = 8.0 Hz, 1H), 5.44 (brs, 1H), 5.30 (brs, 1H), 4.40 (s, 1H), 2.18 -2.09 (m, 5H), 2.01-1.95 (m, 4H), 1.85-1.82 (m, 2H), 1.70-1.67 (m, 2H)
95	E-4-[3-(3-chloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, CDCl ₃) d 8.30 (brs, 1H), 7.80 (s, 1H), 7.50-7.30 (m, 5H), 6.45 (d, J = 8.0 Hz, 1H), 5.68 (brs, 1H), 5.30 (brs, 1H), 4.39 (s, 1H), 2.19-2.10 (m, 5H), 2.02-1.95 (m, 4H), 1.85-1.82 (m, 2H), 1.71-1.68 (m, 2H)
96	E-4-[3-(3-chloro-2-methyl-benzenesulfonylamino)-benzamido]-adamantan-1-		¹ H NMR (400 MHz, CDCl ₃) d 8.80 (s, 1H), 7.97 (d, J = 8.0 Hz, 1H), 7.84 (s, 1H), 7.54 (d, J = 4.0 Hz, 1H), 7.84 (s, 1H), 7.54 (d, J = 4.0 Hz, 1H), 7.44-7.19 (m, 3H), 6.46 (d, J =




	carboxylic acid amide		8.0 Hz, 1H), 5.70 (brs, 1H), 5.30 (s, 3H), 4.42 (d, $J = 8.0$ Hz, 1H), 2.72 (s, 3H), 2.16-2.09 (m, 5H), 1.99-1.94 (m, 4H), 1.83-1.80 (m, 2H), 1.70-1.66 (m, 2H)
97	E-4-{3-[(3-chloro-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.69 (d, $J = 4.0$ Hz, 1H), 7.60-7.51 (m, 3H), 7.44-7.28 (m, 3H), 6.32 (d, $J = 8.0$ Hz, 1H), 5.60 (brs, 1H), 5.26 (brs, 1H), 4.22 (d, $J = 8.0$ Hz, 1H), 3.22 (s, 3H), 2.20 (s, 2H), 2.10-1.95 (m, 7H), 1.87-1.83 (m, 2H), 1.71-1.67 (m, 2H)
98	E-4-[3-N-(2-hydroxyethyl)-2-(trifluoromethyl-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.89 (d, $J = 4.0$ Hz, 1H), 7.83 (d, $J = 4.0$ Hz, 1H), 7.69-7.55 (m, 4H), 7.43-7.36 (m, 3H), 6.31 (d, $J = 8.0$ Hz, 1H), 5.62 (brs, 1H), 5.40 (brs, 1H), 4.18 (s, 1H), 3.92 (t, $J = 8.0$ Hz, 2H), 3.68 (s, 2H), 2.19 (s, 1H), 2.10-1.94 (m, 7H), 1.85-1.82 (m, 2H), 1.70-1.66 (m, 2H)
99	E-4-{3-[(2-trifluoromethyl-benzenesulfonyl)-methylamino]-		^1H NMR (400 MHz, CDCl_3) d 7.90 (d, $J = 8.0$ Hz, 1H), 7.79 (d, $J = 4.0$ Hz, 1H), 7.67-7.42 (m, 4H), 7.40-7.33 (m, 2H), 6.31 (d, $J = 8.0$ Hz, 1H), 5.59 (brs,




	benzamido}-adamantan-1-carboxylic acid amide		1H), 5.23 (brs, 1H), 4.21 (d, $J = 8.0$ Hz, 1H), 3.34 (s, 3H), 2.19-2.18 (m, 2H), 2.09-2.00 (m, 5H), 1.95-1.83 (m, 2H), 1.70-1.67 (m, 2H)
100	E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl](2-(trifluoromethyl)benzenesulfonylamide		¹ H NMR (400 MHz, DMSO) d 8.02 (d, $J = 8.0$ Hz, 1H), 7.75 (d, $J = 4.0$ Hz, 1H), 7.59-7.51 (m, 2H), 7.46 (d, $J = 8.0$ Hz, 1H), 7.12 (s, 1H), 7.06 (s, 1H), 7.00-6.72 (m, 3H), 6.70 (s, 1H), 4.20 (s, 1H), 1.98-1.74 (m, 1H), 1.42-1.39 (m, 2)
101	E-N-(5-carbamoyladamantan-2-yl)-5-[(N-methyl-2-(trifluoromethyl)benzenesulfonylamino]nicotinamide		¹ H NMR (400 MHz, CDCl ₃) d 8.85 (s, 1H), 8.52 (s, 1H), 8.01-7.92 (m, 3H), 7.81-7.71 (m, 2H), 4.17 (s, 1H), 2.22 (s, 3H), 2.10-1.93 (m, 1H), 1.68-1.60 (s, 2H)
102	E-4-[3-(thiophen-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 10.59 (brs, 1H), 7.93-7.84 (m, 2H), 7.53-7.29 (m, 5H), 7.10 (s, 1H), 7.02 (s, 1H), 6.74 (s, 1H), 3.93 (s, 1H), 2.09-2.02 (m, 4H), 1.91-1.77 (m, 7H), 1.44-1.41 (m, 2H)
103	E-4-[3-(furan-2-sulfonylamino)-		¹ H NMR (400 MHz, DMSO) d 8.22 (s, 1H), 7.78 (s, 1H), 7.52-7.50 (m, 2H), 7.49-7.44 (m,




	benzamido]-adamantan-1-carboxylic acid amide		2H), 7.04-7.03 (m, 1H), 6.46-6.42 (m, 2H), 5.70 (s, 1H), 5.30 (s, 1H), 4.37-4.36 (m, 1H), 2.19-2.13 (m, 5H), 2.09-1.95 (m, 4H), 1.86-1.82 (m, 2H), 1.71-1.67 (m, 2H)
104	E-4-[3-(pyridin-3-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 8.89-8.87 (m, 1H), 8.76 (d, <i>J</i> = 4.8 Hz, 1H), 8.13 (d, <i>J</i> = 8.0 Hz, 1H), 7.98 (s, 1H), 7.62-7.58 (m, 1H), 7.48-7.47 (m, 2H), 7.32 (t, <i>J</i> = 8.0 Hz, 1H), 7.26-7.24 (m, 1H), 7.06 (s, 1H), 6.75 (s, 1H), 3.91 (s, 1H), 2.08-1.97 (m, 4H), 1.90-1.67 (m, 7H), 1.43-1.40 (m, 2H)
105	E-4-(3-benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 10.47 (brs, 1H), 7.97-7.92 (m, 1H), 7.77 (d, <i>J</i> = 8.0 Hz, 1H), 7.61-7.46 (m, 5H), 7.31-7.23 (m, 2H), 7.02 (s, 1H), 6.58 (s, 1H), 3.92 (s, 1H), 1.91-1.89 (m, 4H), 1.88-1.70 (m, 7H), 1.43-1.40 (m, 2H)
106	E-4-[3-[(2-chloro-benzenesulfonylamino]benzamido]-adamantan-1-carboxylic acid		¹ H NMR (400 MHz, DMSO) d 10.90 (brs, 1H), 8.04-7.90 (m, 2H), 7.60-7.40 (m, 5H), 7.30-7.20 (m, 2H), 7.05 (s, 1H), 6.70 (s, 1H), 4.00 (s, 1H), 2.00-1.95 (m, 4H),




	acid amide		1.90-1.80 (m, 5H), 1.75-1.70 (m, 2H), 1.45-1.40 (m, 2H)
107	E-4-[3-(2,4-dimethylthiazol-5-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 10.70 (brs, 1H), 7.93 (d, <i>J</i> = 6.4 Hz, 1H), 7.48-7.45 (m, 2H), 7.31 (t, <i>J</i> = 8.0 Hz, 1H), 7.30-7.24 (m, 2H), 7.06 (s, 1H), 6.74 (s, 1H), 2.57 (s, 3H), 2.40 (s, 1H), 2.08-2.00 (m, 4H), 1.90-1.75 (m, 7H), 1.48-1.40 (m, 2H)
108	E-4-[3-(3,5-dimethyl-1H-pyrazole-4-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 12.98 (brs, 1H), 10.33 (brs, 1H), 7.93 (d, <i>J</i> = 6.4 Hz, 1H), 7.48-7.45 (m, 2H), 7.31 (t, <i>J</i> = 8.0 Hz, 1H), 7.17-7.15 (m, 1H), 7.06 (s, 1H), 6.74 (s, 1H), 4.26-4.24 (m, 1H), 2.24 (s, 6H), 2.08-2.00 (m, 4H), 1.88-1.75 (m, 7H), 1.43-1.41 (m, 2H)
109	E-N-(5-hydroxyadamantan-2-yl)-3-benzenesulfonylamino-benzamide		¹ H NMR (400 MHz, CDCl ₃) d 8.02 (s, 1H), 7.88 (d, <i>J</i> = 8.0 Hz, 1H), 7.79 (d, <i>J</i> = 8.0 Hz, 1H), 7.46 (t, <i>J</i> = 8.0 Hz, 1H), 7.20-7.16 (m, 2H), 7.07-7.03 (m, 3H), 6.66 (s, 1H), 4.07 (s, 1H), 2.17-2.14 (m, 3H), 1.88-1.85 (m, 2H), 1.77-1.70 (m, 6H), 1.53-1.50 (m, 2H)





110	E- <i>N</i> -cycloheptyl-3-phenylsulfonylbenzamide		¹ H NMR (400 MHz, CDCl ₃) d 8.12 (s, 1H), 7.99 (d, <i>J</i> = 8.0 Hz, 1H), 7.83 (d, <i>J</i> = 8.0 Hz, 1H), 7.52 (t, <i>J</i> = 8.0 Hz, 1H), 7.20-7.11 (m, 3H), 6.97 (s, 1H), 6.03 (s, 1H), 4.15-4.12 (m, 1H), 2.07-1.98 (m, 2H), 1.67-1.40 (m, 10H)
111	E- <i>N</i> -(5-hydroxyadamantan-2-yl)-3-(2-(trifluoromethyl)benzenesulfonyl)benzamide		¹ H NMR (400 MHz, CDCl ₃) d 8.07 (d, <i>J</i> = 8.0 Hz, 1H), 7.92 (d, <i>J</i> = 8.0 Hz, 1H), 7.70-7.52 (m, 2H), 7.51-7.49 (m, 2H), 7.36-7.31 (m, 2H), 6.87 (brs, 1H), 6.22 (d, <i>J</i> = 6.4 Hz, 1H), 4.20-4.18 (m, 1H), 2.24-2.20 (m, m, 3H), 1.97-1.94 (m, 2H), 1.84-1.74 (m, 6H), 1.61-1.60 (m, 2H)
112	E-4-(3-(<i>N</i> -phenylsulfonylbenzamin)-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 10.40 (brs, 1H), 8.24 (d, <i>J</i> = 6.4 Hz, 1H), 8.16 (s, 1H), 8.03 (d, <i>J</i> = 6.4 Hz, 1H), 7.86 (d, <i>J</i> = 8.8 Hz, 1H), 7.61 (t, <i>J</i> = 8.0 Hz, 1H), 7.38-7.20 (m, 2H), 7.09-7.01 (m, 4H), 6.75 (s, 1H), 3.97-3.96 (m, 4H), 2.09-2.05 (m, 4H), 1.91-1.76 (m, 17H), 1.45-1.42 (m, 2H)
113	E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl		¹ H NMR (400 MHz, DMSO) d 7.69-7.65 (m, 1H), 7.54-7.51 (m, 2H), 7.17-7.13 (m, 2H), 7.03 (brs, 1H), 6.97-




	1)phenyl]- 2-fluoro-3- chloro- benzenesulf onylamide		6.92 (m, 3H), 6.73 (brs, 1H), 3.90 (brs, 1H) 2.09-2.01 (m, 4H), 1.99-1.73 (m, 7H), 1.43-1.40 (m, 2H)
11 4	E-4-[3-(3- chloro-4- (trifluorom ethyl)benze nesulfonyla mino)benzam ido]adamant an-1- carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 11.07 (s, 1H), 8.25 (d, J = 4.2 Hz, 1H), 8.15 (s, 1H), 7.93 (t, J = 4 Hz, 2H), 7.50- 7.48 (m, 1H), 7.32 (t, J = 4 Hz, 1H), 7.24 (d, J = 6 Hz, 1H), 7.02 (s, 1H), 6.75 (s, 1H), 3.91 (d, J = 3 Hz, 1H), 2.01-1.89 (m, 4H), 1.85-1.80 (m, 5H), 1.76 (s, 2H), 1.41 (d, J = 6.2 Hz, 2H)
11 5	E-4-[3-[(2- (trifluorom ethyl)benze nesulfonyla mino]- benzamido]- adamantan- 1- carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 10.86 (s, 1H), 8.09 (d, J = 7.2 Hz, 1H), 8.00 (d, J = 7.2 Hz, 1H), 7.92 (d, J = 6.4 Hz, 1H), 7.88-7.83 (d, 2H), 7.48 (d, J = 11.6 Hz, 2H), 7.32 (t, J = 7.6 Hz, 1H), 7.22 (d, J = 8 Hz, 1H), 7.01 (s, 1H), 6.74 (s, 1H), 3.92 (d, J = 5.6 Hz, 1H), 2.01 (s, 4H), 1.91-1.76 (m, 7H), 1.42 (d, J = 12 Hz, 2H)
11 6	E-4-[3-(2- chloro-4- bromo- benzenesulf onylamino)- benzamido]-		¹ H NMR (400 MHz, DMSO) d 10.91 (s, 1H), 7.94 (d, J = 15.6 Hz, 3H), 7.74 (d, J = 8.6 Hz, 1H), 7.45 (s, 2H), 7.25 (t, J = 8 Hz,




	adamantan-1-carboxylic acid amide		1H), 7.19 (d, $J = 7.6$ Hz, 1H), 7.02 (s, 1H), 6.75 (s, 1H), 4.12 (d, $J = 5.2$ Hz, 1H), 1.20 (d, $J = 5.4$ Hz, 4H), 1.91 (s, 2H), 1.84 (d, $J = 6.4$ Hz, 3H), 1.76 (s, 2H), 1.42 (d, $J = 12$ Hz, 2H)
117	E-4-[3-(2,4,6-trichlorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 11.09 (s, 1H), 7.99-7.90 (m, 2H), 7.46 (s, 2H), 7.33 (t, $J = 8$ Hz, 1H), 7.20 (d, $J = 8.4$ Hz, 2H), 7.02 (s, 1H), 6.75 (s, 1H), 3.91 (d, $J = 5.6$ Hz, 1H), 2.00 (d, $J = 7.6$ Hz, 5H), 1.87 (d, $J = 14.4$ Hz, 3H), 1.76 (s, 2H), 1.42 (d, $J = 12$ Hz, 2H)
118	E-4-[3-(3-chloro-5-fluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 10.60-10.53 (m, 1H), 7.981-7.949 (m, 2H), 7.94 (m, 1H), 7.63 (t, $J = 4.4$ Hz, 1H), 7.54 (m, 2H), 7.26 (d, $J = 2$ Hz, 1H), 7.03 (s, 1H), 6.76 (s, 1H), 3.93 (t, $J = 2$ Hz, 1H), 2.00 (d, $J = 3.4$ Hz, 4H), 1.91-1.76 (m, 7H), 1.42 (d, $J = 6.4$ Hz, 2H)
119	E-4-[3-(3,5-dichlorobenzenesulfonylamino)-benzamido]-adamantan-		^1H NMR (400 MHz, DMSO) d 7.89 (s, 2H), 7.70 (s, 2H), 7.41 (s, 2H), 7.31 (s, 1H), 7.21 (d, $J = 7.2$ Hz, 1H), 7.03 (s, 1H), 6.74 (s, 1H), 4.14




	1-carboxylic acid amide		(d, $J = 6.4$ Hz, 1H), 2.06 (d, $J = 27.6$ Hz, 7H), 1.91 (s 3H), 1.43 (d, $J = 11.6$ Hz, 3H)
120	E-4-[3-(3-fluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 10.74 (s, 1H), 7.99 (d, $J = 6.4$ Hz, 1H), 7.66 (d, $J = 7.6$ Hz, 2H), 7.59 (d, $J = 8.4$ Hz, 1H), 7.53 (d, $J = 7.6$ Hz, 3H), 7.37-7.27 (m, 2H), 7.06 (s, 1H), 6.78 (s, 1H), 3.96 (s, 1H), 2.04 (s, 4H), 1.98-1.78 (m, 8H), 1.47 (d, $J = 12.4$ Hz, 2H)
121	E-4-[3-(2,4-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 8.00 (s, 1H), 7.91-7.86 (m, 1H), 7.73 (t, 1H), 7.43-7.41 (m, 2H), 7.33 (t, 1H), 7.00-6.88 (m, 2H), 6.40 (d, $J = 7.6$ Hz, 1H), 5.67 (s, 1H), 5.35 (s, 1H), 4.34 (d, $J = 7.6$ Hz, 1H), 2.14 (t, 6H), 2.03 (t, 6H), 1.85 (d, $J = 14.4$ Hz, 2H), 1.70 (d, $J = 12.4$ Hz, 2H)
122	E-4-[3-(2,5-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 10.96 (s, 1H), 8.01 (d, $J = 6.4$ Hz, 1H), 7.68-7.61 (m, 2H), 7.58-7.54 (m, 3H), 7.39-7.29 (m, 2H), 7.05 (s, 1H), 6.78 (s, 1H), 3.95 (s, 1H), 2.04 (s, 1H), 1.92-1.79 (m, 8H), 1.47 (d, $J = 12.0$ Hz, 2H)


12 3	E-4-[3-(2,6-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide		¹ H NMR (400 MHz, DMSO) d 11.12 (s, 1H), 7.97 (s, 1H), 7.71 (s, 1H), 7.54 (d, <i>J</i> = 22.8 Hz, 2H), 7.35-7.29 (m, 4H), 7.06 (s, 1H), 6.78 (s, 1H), 3.96 (s, 1H), 2.04 (s, 4H), 1.94-1.79 (m, 7H), 1.47 (d, <i>J</i> = 12.0 Hz, 2H)
12 4	E-N-cycloheptyl-N-propyl-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide		¹ H NMR (400 MHz, CDCl ₃) d 8.03 (d, <i>J</i> = 8.0 Hz, 1H), 7.87 (d, <i>J</i> = 8.0 Hz, 1H), 7.64 (t, 1H), 7.55 (t, 1H), 7.28-7.22 (m, 2H), 7.17-7.13 (m, 1H), 7.05 (t, 2H), 3.47 (s, 1H), 3.24 (t, 1H), 2.98 (s, 1H), 1.86-1.60 (m, 10H), 1.40 (s, 4H), 1.28 (d, <i>J</i> = 7.6 Hz, 2H), 1.14 (s, 2H), 1.02-0.69 (m, 3H), 0.59 (s, 1H)
12 5	E-4-[2-fluoro-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 8.03 (t, <i>J</i> = 8.0 Hz, 1H), 7.91 (d, <i>J</i> = 7.6 Hz, 1H), 7.77 (td, <i>J</i> ₁ = 1.6 Hz, <i>J</i> ₂ = 6.4 Hz, 1H), 7.73 (m, 1H), 7.62 (t, <i>J</i> = 7.6 Hz, 1H), 6.97 (s, 1H), 6.63 (t, <i>J</i> = 7.2 Hz, 1H), 5.57 (brs, 1H), 5.21 (brs, 1H), 4.20 (d, <i>J</i> = 4.8 Hz, 1H), 2.22-1.64 (m, 13H)

12 6	E-4-[2-chloro-5-(3-chlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 9.04 (s, 1H), 7.79 (t, 1H), 7.69-7.66 (m, 2H), 7.52-7.47 (m, 2H), 7.39-7.33 (m, 2H), 7.08 (d, <i>J</i> = 7.6 Hz, 1H), 5.75 (s, 1H), 5.37 (s, 1H), 4.43 (d, <i>J</i> = 8.0 Hz, 1H), 2.36-2.15 (m, 4H), 2.09 (s, 1H), 1.98 (t, 4H), 1.87-1.79 (m, 3H), 1.69 (d, <i>J</i> = 12.4 Hz, 2H)
12 7	E-4-[3-(3-chlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, DMSO) d 10.50 (s, 1H), 8.14 (s, 1H), 7.93-7.56 (m, 6H), 7.22 (t, 1H), 7.02 (s, 1H), 6.74 (s, 1H), 3.94 (d, <i>J</i> = 5.2 Hz, 1H), 2.02 (s, 4H), 1.92-1.69 (m, 8H), 1.44 (d, <i>J</i> = 12.4 Hz, 2H)
12 8	E-4-[4-chloro-3-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 7.38-7.26 (m, 1H), 7.23 (d, <i>J</i> = 2 Hz, 1H), 7.00-6.97 (m, 1H), 6.31 (d, <i>J</i> = 7.2 Hz, 1H), 5.64 (s, 1H), 5.01 (s, 1H), 2.33-1.42 (m, 11H)
12 9	E-4-[2-chloro-5-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-		¹ H NMR (400 MHz, CDCl ₃) d 7.67-7.63 (m, 3H), 7.55-7.47 (m, 2H), 7.38 (d, <i>J</i> = 8.8 Hz, 1H), 7.15 (d, <i>J</i> = 8.0 Hz, 1H), 5.80 (s, 1H), 5.62 (s, 1H), 4.43 (d, <i>J</i> = 8.0 Hz, 1H),


	carboxylic acidamide		2.21-1.60 (m, 14H)
130	E-4-[3-(3,5-dichlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 7.26-7.21 (m, 1H), 7.03-6.99 (m, 2H), 6.27 (d, <i>J</i> = 7.2 Hz, 1H), 5.60 (s, 1H), 5.33 (s, 1H), 4.22 (d, <i>J</i> = 7.2 Hz, 1H), 3.87 (s, 2H), 3.75-3.61 (m, 1H), 2.37-1.56 (m, 15H)
131	E-4-[5-(3,5-dichlorobenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 8.90 (s, 1H), 7.95-7.92 (m, 1H), 7.71-7.67 (m, 1H), 7.63 (d, <i>J</i> = 2.0 Hz, 2H), 7.46 (t, 1H), 7.18-7.13 (m, 1H), 5.68 (s, 1H), 5.30 (s, 1H), 4.55 (d, <i>J</i> = 8.0 Hz, 1H), 2.32 (t, 4H), 2.19 (d, <i>J</i> = 6.0 Hz, 1H), 2.01-1.95 (t, 4H), 1.86 (d, <i>J</i> = 13.2 Hz, 2H), 1.71 (d, <i>J</i> = 13.6 Hz, 1H)
132	E-4-[2-fluoro-3-(3-chlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 7.84 (m, 2H), 7.68 (m, 2H), 7.56 (m, 1H), 7.43 (d, <i>J</i> = 8.0 Hz, 1H), 7.26 (m, 1H), 6.68 (m, 1H), 5.66 (brs, 1H), 5.50 (brs, 1H), 4.23 (d, <i>J</i> = 4.8 Hz, 1H), 2.19 (s, 2H), 2.07 (m, 5H), 1.94 (m, 2H), 1.76 (d, <i>J</i> = 13.6 Hz, 1H), 1.66 (d, <i>J</i> = 13.2 Hz, 1H)

13 3	E-4-[2-chloro-5-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 7.80 (d, <i>J</i> = 2.4 Hz, 1H), 7.68 (dd, <i>J</i> = 8.8, 2.4 Hz, 1H), 7.63 (d, <i>J</i> = 2.8, 1H), 7.46 (dd, <i>J</i> = 6.4, 1.6 Hz, 1H), 7.30 (d, <i>J</i> = 8.4 Hz, 1H), 7.01 (d, <i>J</i> = 8.0 Hz, 1H), 6.90 (d, <i>J</i> = 8.8 Hz, 1H), 5.92 (brs, 1H), 5.73 (brs, 1H), 4.37 (d, <i>J</i> = 7.6 Hz, 1H), 3.91 (s, 3H), 2.17 (m, 4H), 2.07 (s, 1H), 1.97 (d, <i>J</i> = 12.4 Hz, 1H), 1.93 (s, 2H), 1.84 (d, <i>J</i> = 13.2 Hz, 2H), 1.65 (d, <i>J</i> = 13.2 Hz, 2H)
13 4	E-4-[4-chloro-3-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acidamide		¹ H NMR (400 MHz, CDCl ₃) d 7.94 (d, <i>J</i> = 2.0 Hz, 1H), 7.80 (m, 1H), 7.66 (td, <i>J</i> = 2.4, 4.1 Hz, 1H), 7.53 (dd, <i>J</i> = 6.4, 2.0 Hz, 1H), 7.37 (d, <i>J</i> = 8.0 Hz, 1H), 6.92 (d, <i>J</i> = 8.8 Hz, 1H), 6.32 (d, <i>J</i> = 7.2 Hz, 1H), 5.56 (d, <i>J</i> = 33.2 Hz, 2H), 4.22 (m, 1H), 3.94 (s, 3H), 2.22-1.69 (m, 13H)
13 5	E-4-[5-(3-chloro-4-methoxybenzenesulfonylamino)-2-fluorobenza mido]adamantan-1-		¹ H NMR (400 MHz, CDCl ₃) d 8.47 (brs, 1H), 7.90 (dd, <i>J</i> = 4.0, 2.8 Hz, 1H), 7.71 (d, <i>J</i> = 2.4 Hz, 1H), 7.65 (m, 2H), 7.32 (m, 1H), 7.11 (m, 1H), 6.88 (d, <i>J</i> = 8.8 Hz, 1H), 5.71

	carboxylic acidamide		(brs, 1H), 5.30 (brs, 1H), 4.52 (d, $J = 7.6$ Hz, 1H), 3.91 (s, 3H), 2.17 (m, 4H), 2.05 (s, 1H), 1.97 (m, 4H), 1.82 (d, $J = 13.2$ Hz, 2H), 1.66 (d, $J = 13.2$ Hz, 2H)
136	E-4-(3-(4-chlorobenzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 10.54 (s, 1H), 7.94 (d, $J = 6.4$ Hz, 1H), 7.78-7.75 (m, 2H), 7.66-7.63 (m, 2H), 7.52-7.49 (m, 2H), 7.32 (t, $J = 7.8$ Hz, 1H), 7.24 (dd, $J = 8, 1.2$ Hz, 1H), 7.01 (s, 1H), 6.73 (s, 1H), 3.94-3.93 (m, 1H), 2.05-1.76 (m, 11H), 1.42 (d, $J = 12.8$ Hz, 2H)
137	E-4-(3-(2,4,5-trifluorobenzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, DMSO) d 10.96 (s, 1H), 7.84-7.98 (m, 3H), 7.48-7.53 (m, 2H), 7.35 (t, $J = 8.0$ Hz, 1H), 7.26 (dd, $J = 8.4, 1.2$ Hz, 1H), 7.02 (brs, 1H), 6.75 (brs, 1H), 3.78 (d, $J = 6.0$ Hz, 1H), 1.84-2.02 (m, 9H), 1.76 (s, 2H), 1.42 (d, $J = 12.8$ Hz, 2H)
138	E-4-(3-(cyclopropylsulfonylamino-benzamido)-adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 7.93 (s, 1H), 7.86 (brs, 1H), 7.62 (d, $J = 7.6$ Hz, 1H), 7.40-7.48 (m, 2H), 6.51 (d, $J = 8.0$ Hz, 1H), 5.77 (brs, 1H), 5.51 (brs, 1H), 4.35 (d, $J = 8.0$

			Hz, 1H), 2.47-2.54 (m, 1H), 2.09-2.19 (m, 4H), 2.00-2.03 (m, 2H), 1.95 (s, 2H), 1.86 (d, $J = 13.2$ Hz, 2H), 1.69 (d, $J = 12.8$ Hz, 2H), 1.16-1.20 (m, 2H), 0.93-0.99 (m, 2H)
13 9	E-4-(3-(1-methylethylsulfonamido)benzamido)adamantan-1-carboxylic acid amide		^1H NMR (400 MHz, CDCl_3) d 8.54 (s, 1H), 7.98 (s, 1H), 7.66-7.69 (m, 1H), 7.37-7.41 (m, 1H), 6.56 (d, $J = 8.0$ Hz, 1H), 5.91 (brs, 1H), 5.73 (brs, 1H), 4.39 (d, $J = 7.6$ Hz, 1H), 3.26-3.32 (m, 1H), 2.19 (d, $J = 12.0$ Hz, 4H), 1.92-2.04 (m, 5H), 1.86 (d, $J = 13.2$ Hz, 2H), 1.69 (d, $J = 12.8$ Hz, 2H), 1.38 (d, $J = 6.8$ Hz, 6H)

Preparational Example 35: Preparation of 4-[(1-[3,4-dihydro-1H-isoquinolin-2-carbonyl)-amino]-methyl-cyclopropanecarbonyl)-amino]-adamantan-1-carboxylic acid methyl ester




30 mg (0.1 mmol) of 4-[(1-aminomethyl-cyclopropanecarbonyl)-amino]-adamantane-1-carboxylic

acid methyl ester was dissolved in 1.0 ml of chloroform, to which 16 mg (0.1 mmol) of carbodiimide and TEA were added. Then, 14 mg (0.1 mmol) of 1,2,3,4-tetrahydroisoquinoline was also added thereto. The mixture was stirred for 12 hours with reflux, followed by evaporation under reduced pressure to eliminate chloroform. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and evaporated under reduced pressure, followed by column chromatography to give 30 mg of 4-[(1-[3,4-dihydro-1H-isoquinolin-2-carbonyl)-amino]-methyl-cyclopropanecarbonyl)-amino]-adamantan-1-carboxylic acid methyl ester (yield: 65%).


¹H NMR (400 MHz, CDCl₃) δ 7.63 (d, *J* = 7.2 Hz, 1H), 7.29-7.10 (m, 4H), 4.94-4.91 (m, 1H), 4.55 (s, 2H), 4.00-3.82 (m, 1H), 3.65 (s, 3H), 3.62-3.53 (m, 4H), 2.87 (t, *J* = 6.4 Hz, 2H), 2.19-1.74 (m, 11H), 1.48-1.42 (m, 2H), 0.90-0.87 (m, 2H), 0.71-0.68 (m, 2H).

Preparational Example 36: Preparation of 4-[(1-[3,4-dihydro-1H-isoquinolin-2-carbonyl)-amino]-methyl-cyclopropanecarbonyl)-amino]-adamantan-1-carboxylic acid



25 mg (0.05 mmol) of the 4-[(1-[3,4-dihydro-1H-isoquinolin-2-carbonyl)-amino]-methyl-cyclopropanecarbonyl)-amino]-adamantan-1-carboxylic acid methyl ester prepared in Preparational Example 35 was dissolved in 1.0 ml of methanol and 1.0 ml of water, to which 20 mg (0.5 mmol) of sodium hydroxide was added. The mixture was stirred at room temperature for 12 hours, followed by evaporation under reduced pressure to eliminate methanol. The residue was acidized with 1 N HCl, followed by extraction with 320 ml of chloroform. The extract was dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure. The obtained compound proceeded to the next reaction without purification.

Example 140: Preparation of E-[3,4-dihydro-1H-isoquinolin-2-carboxylic acid-1-[(5-carbamoyl-adamantan-2-yl)carbamoyl]cyclopropylmethyl]-amide



15 mg (0.03 mmol) of the 4-[(1-[3,4-dihydro-1H-isoquinolin-2-carbonyl)-amino]-methyl-cyclopropanecarbonyl)-amino]-adamantan-1-carboxylic
 5 acid prepared in Preparational Example 36 was dissolved in 5.0 ml of methylene chloride, to which 11 mg (0.06 mmol) of EDC and 8 mg (0.06 mmol) of HOBT were added, and then ammonia water was also added thereto. The mixture was stirred at room temperature
 10 for 12 hours, followed by evaporation under reduced pressure to eliminate methylene chloride. The residue was dissolved in 10 ml of water, followed by extraction with 320 ml of methylene chloride. The extract was dried over anhydrous magnesium sulfate,
 15 and evaporated under reduced pressure, followed by column chromatography to give 7 mg of [3,4-dihydro-1H-isoquinolin-2-carboxylic acid-1-[(5-carbamoyl-adamantan-2-ylcarbamoyl)cyclopropylmethyl]-amide (yield: 50%).



20 ^1H NMR (400 MHz, CDCl_3) δ 7.73 (d, J = 7.2 Hz, 1H), 7.22-7.11 (m, 4H), 5.63 (brs, 1H), 5.41 (brs, 1H), 4.97 (t, J = 7.2 Hz, 1H), 4.55 (s, 1H), 4.13-4.11




(m, 1H), 3.64 (t, $J = 6.4$ Hz, 1H), 3.55 (d, $J = 5.2$ Hz, 2H), 2.87 (t, $J = 6.4$ Hz, 1H), 2.05-1.86 (m, 1H), 1.48-1.45 (m, 2H), 1.31-1.20 (m, 2H), 0.71-0.68 (m, 2H).




5


The compounds of Example 141 - Example 149 were prepared by the preparation method described in Example 140.

【Table 6】

Ex am pl e	Compound Name	Structure	NMR data
14 1	E-3,4-dihydro-2H-quinolin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide		^1H NMR (400 MHz, CDCl_3) d 7.74 (d, $J = 7.2$ Hz, 1H), 7.27-7.16 (m, 3H), 7.11-7.07 (m, 1H), 5.64 (brs, 1H), 5.53-5.50 (m, 2H), 4.06-4.04 (m, 1H), 3.76 (t, $J = 6.4$ Hz, 2H), 3.55 (t, $J = 6.4$ Hz, 2H), 2.75 (t, $J = 6.4$ Hz, 2H), 2.11-1.89 (m, 11H), 1.56-1.53 (m, 2H), 1.28-1.21 (m, 2H), 0.61-0.59 (m, 2H)
14 2	E-piperidin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-		^1H NMR (400 MHz, CDCl_3) d 7.87 (d, $J = 7.2$ Hz, 1H), 5.66 (brs, 1H), 5.46 (brs, 1H), 4.86 (t, $J = 6.4$ Hz, 1H), 4.02-4.01 (m, 1H), 3.56-3.48 (m, 2H), 3.35-3.26 (m, 4H),

	ylcarbamoyl)- cyclopropylmethyl]- amide		2.08-1.76 (m, 11H), 1.59-1.50 (m, 8H), 1.30-1.24 (m, 2H), 0.65-0.63 (m, 2H)
14 3	E-4-{[1-5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-carbamoyl}-3,4-dihydroquinolin-1-carboxylic acid-butylester		¹ H NMR (400 MHz, CDCl ₃) d 7.98 (d, <i>J</i> = 8.0 Hz, 1H), 7.59 (d, <i>J</i> = 8.0 Hz, 1H), 7.20-7.14 (m, 2H), 7.10-7.03 (m, 1H), 5.59 (brs, 1H), 5.51 (t, <i>J</i> = 6.4 Hz, 1H), 5.28 (brs, 1H), 4.07-4.05 (m, 1H), 3.86-3.78 (m, 4H), 3.54 (d, <i>J</i> = 3.2 Hz, 1H), 2.19-1.90 (m, 11H), 1.60-1.57 (m, 2H), 1.54 (s, 9H), 1.27-1.25 (m, 2H), 0.63-0.60 (m, 2H)
14 4	E-4-pyrimidin-2-yl-piperazin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide		¹ H NMR (400 MHz, CDCl ₃) d 8.30 (d, <i>J</i> = 8.8 Hz, 2H), 6.65 (d, <i>J</i> = 4.8 Hz, 1H), 5.62 (brs, 1H), 5.37 (brs, 1H), 4.93 (t, <i>J</i> = 6.4 Hz, 1H), 4.02-4.01 (m, 1H), 3.90-3.84 (m, 4H), 3.57 (d, <i>J</i> = 8.8 Hz, 1H), 3.50-3.48 (m, 4H), 2.02-1.87 (m, 1H), 1.59-1.51 (m, 2H), 1.31-1.22 (m, 2H), 0.69-0.67 (m, 2H)
14 5	E-4-((1-(3-phenylureido)methyl)-cyclopropa		¹ H NMR (400 MHz, CDCl ₃) d 7.63-7.62 (m, 1H), 7.37-7.26 (m, 3H), 7.06-7.03 (m, 1H), 5.75-5.70 (brs, 2H), 5.42 (brs, 1H), 4.00

	necarbonyl }amino)- adamantan- 1carboxyli c acid amide		(brs, 1H), 3.50 (s, 2H), 2.07-1.85 (m, 11H), 1.53-1.48 (m, 2H), 1.26-1.22 (m, 2H), 0.66 (brs, 2H)
14 6	E-3,4- dihydro- 2H- quinoxalin -1- carboxylic acid[1-(5- carbamoyl- adamantan- 2- ylcarbamoy l)- cyclopropy lmethyl]- amide		¹ H NMR (400 MHz, MeOD) d 7.80 (brs, 1H), 7.41-7.24 (m, 3H), 4.04-3.89 (m, 3H), 3.74-3.60 (m, 2H), 3.33 (s, 1H), 2.18-1.84 (m, 9H), 1.58-1.55 (m, 2H), 1.20 (brs, 2H), 0.91 (brs, 2H)
14 7	E-3,4- dihydro- 1H- isoquinoli n-2- carboxylic acid[4-(5- carbamoyl- adamantan- 2- ylcarbamoy l)- thiazol-2- yl]-amide		¹ H NMR (400 MHz, CDCl ₃) d 7.80-7.60 (m, 2H), 7.40-7.20 (m, 6H), 5.70 (brs, 1H), 5.60 (brs, 1H), 4.70 (s, 2H), 4.20-4.10 (m, 1H), 3.80 (t, J = 6.4 Hz, 2H), 3.00 (t, J = 6.4 Hz, 2H), 2.10-1.80 (m, 9H), 1.65-1.55 (m, 2H)
14 8	E-3,4- dihydro- 2H- quinolin- 1- carboxylic acid[4-(5- carbamoyl- adamantan- 2- ylcarbamoy l)- thiazol-2- yl]-amide		¹ H NMR (400 MHz, CDCl ₃) d 7.49-7.44 (m, 1H), 7.19-7.17 (m, 2H), 7.13-7.05 (m, 4H), 5.74 (s, 1H), 5.67 (brs, 1H), 5.45 (brs, 1H), 3.97-3.95 (m, 1H), 3.77

	carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide		(t, $J = 6.4$ Hz, 2H), 2.78 (t, $J = 6.4$ Hz, 2H), 2.05-1.80 9m, 13H), 1.62-1.58 (m, 2H), 1.54-1.51 9m, 2H), 1.01-0.97 (m, 2H)
14 9	E-2-(2-fluoro-benzamido)-thiazol-4-carboxylic acid(5-carbamoyl-adamantan-2-yl)amide		^1H NMR (400 MHz, CDCl_3) d 8.08-8.04 (m, 1H), 7.52-7.42 (m, 1H), 7.34-7.29 (m, 2H), 7.20 (t, $J = 4.0$ Hz, 1H), 7.10 (t, $J = 4.4$ Hz, 1H), 5.67 (brs, 1H), 5.42 (brs, 1H), 4.16-4.14 (m, 1H), 2.12-1.92 (m, 9H), 1.80-1.60 (m, 4H)

Experimental Example 1: Examination of 11β -HSD1 inhibiting effect

To investigate 11β -HSD1 inhibiting activity of the compounds prepared in the examples of the present invention, the following experiment was performed.

The inhibiting activity of 11β -HSD1 derived from microsomal fractions was measured using the HTRF assay (62CO2PEB, Cisbio). Different concentrations of compounds were added to 96-well plates, followed by the addition of TE buffer (20 mM Tris buffer and 5 mM EDTA, pH 6.0) containing 200 μM NADPH(N1630, Sigma) and 160 nM cortisone (C2755, Sigma). The reactions were initiated by the addition of human microsomal

fractions (M0317, Sigma), and were allowed to incubate for 2 h at 37 °C. Europium (Eu³⁺) cryptate and XL665-conjugated cortisol were then added to each well and incubated for an additional 2 h at room temperature.

5 The cortisol concentration was calculated using a calibration curve. The IC₅₀ values were calculated from dose response curves using GraphPad Prism software. IC₅₀ of the example compound to 11β-HSD1 was obtained. The results are shown in Table 7.

10

【Table 7】

Example	human IC ₅₀ (nM)	Example	human IC ₅₀ (nM)
1	14.2	2	26.6
3	6.2	4	-
5	2.7	6	-
7	2.6	8	0.3
9	608.0	10	2.5
11	10.4	12	9.5
13	2.8	14	13.3
15	28.0	16	5.0
17	132.0	18	23.9
19	17.8	20	102.0
21	30.8	22	0.02
23	-	24	156.0
25	9.6	26	1.3
27	-	28	1.3
29	-	30	-
31	1.2	32	1.4
33	1.5	34	28.9
35	-	36	73.0
37	17.1	-	-
38	6.7	39	24.7

40	1.7	41	10.0
42	6.4	43	29.7
44	79.1	45	32.5
46	34.7	47	495.0
48	62.8	49	-
50	0.4	51	-
52	-	53	-
54	-	55	-
56	8.3	57	9.3
58	17.3	59	11.2
60	12.6	61	2.8
62	0.12	63	672.0
64	111.0	65	-
66	-	67	7.1
68	8.9	69	28.6
70	6.2	71	0.6
72	1.2	73	0.83
74	2.6	75	54.6
76	2.4	77	-
78	-	79	-
80	14.8	81	2.1
82	-	83	9.9
84	0.36	85	-
86	-	87	-
88	33.8	89	-
90	-	-	-
91	-	92	148.0
93	-	94	75.7
95	13.1	96	15.4
97	15.6	98	58.6
99	1.9	100	15.3
101	-	102	1.1
103	23.8	104	-
105	42.7	106	4.8
107	60.3	108	51.2
109	-	110	-
111	-	112	-
113	2.7	114	30.7
115	19.1	116	14.5
117	5.0	118	20.4

119	0.9	120	27.2
121	7.8	122	15.1
123	12.8	124	-
125	11.6	126	67.5
127	78.3	128	-
129	41.8	130	-
131	-	132	148.0
133	14.8	134	-
135	-	136	-
137	-	138	62.1
139	-	140	97.8
141	52.5	142	-
143	-	144	-
145	-	146	128.0
147	-	148	-
149	-	-	-

As shown in Table 7, IC₅₀ of the example compounds to 11 β -HSD1 was 0.02-672.0 nM, indicating the compounds had a significant 11 β -HSD1 inhibiting activity. In particular, IC₅₀ of those compounds prepared in Examples 8, 22, 50, 62, 71, 73, 84, and 119 were all less than 1.0 nM, indicating they had an excellent 11 β -HSD1 inhibiting activity.

Therefore, the compound represented by formula 1 of the present invention can be effectively used for the prevention or treatment of diseases caused by abnormal activation of 11 β -HSD1 such as non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder, metabolic syndrome, and other

diseases mediated by the excessive activity of glucocorticoid.

5 Experimental Example 2: In vivo study of the lowering effect on plasma glucose in KKAY mouse model

The inhibitory effect on plasma glucose of the compounds prepared in Examples of the present invention and showed an excellent 11 β -HSD1 activity was investigated *in vivo*.

10

<2-1> In vivo study

Ay gene was introduced into KK mice to induce severe obesity and hyperglycemia. The male KKAY mice (9-week old, DaehanBioLink, Korea) having severe
15 obesity and hyperglycemia prepared thereby were administered orally with the compounds prepared in Examples 3, 22, 71, 84, 113, and 119 at the concentrations of 50 mg/kg and 100 mg/kg. After 4 hour-fasting, the levels of plasma glucose and insulin
20 were measured. The test continued for 3 weeks.

<2-2> Plasma glucose lowering effect

KKAY mice were administered orally with the compounds prepared in Examples 3, 22, 71, and 113 at
25 the concentration of 100 mg/kg for 3 weeks.

Thereafter, the level of plasma glucose after fasting was measured. As a result, the plasma glucose lowering effect was approximately 9% in the group treated with the compound of Example 3. In the group
 5 treated with the compound of Example 22, the level of plasma glucose was decreased approximately 32%. In the groups treated with the compounds of Examples 71 and 113, the plasma glucose lowering effect was respectively 27% and 28%.

10 In the group treated with the compound of Example 84 at the concentration of 50 mg/kg, the plasma glucose lowering effect was 19%. In the group treated with the compound of Example 119 at the concentration of 50 mg/kg, the plasma glucose lowering effect was
 15 47%. The results are shown in Table 8.

【Table 8】

	Plasma glucose lowering effect					
	Example 3	Example 22	Example 71	Example 84	Example 113	Example 119
100 mg/kg	9%	32%	27%		28%	
50 mg/kg				19%		47%

<2-3> Plasma insulin lowering effect

20 KKAY mice were administered orally with the compounds prepared in Examples 22, 71, and 113 at the

concentration of 100 mg/kg for 3 weeks. Thereafter, the level of plasma insulin after fasting was measured. As a result, the plasma insulin lowering effect was approximately 53% in the group treated with the compound of Example 22. In the groups treated with the compounds of Examples 71 and 113, the plasma insulin lowering effect was respectively 40% and 60%.

In the group treated with the compound of Example 84 at the concentration of 50 mg/kg, the plasma insulin lowering effect was 43%. In the group treated with the compound of Example 119 at the concentration of 50 mg/kg, the plasma insulin lowering effect was 52%. The results are shown in Table 9.

15 【Table 9】

	Plasma insulin lowering effect				
	Example 22	Example 71	Example 84	Example 113	Example 119
100 mg/kg	53%	40%		60%	
50 mg/kg			43%		52%

As shown in Table 8 and Table 9, when the compounds prepared in Examples 3, 22, 71, 84, 113, and 119 were treated to KKAY mice at the concentration of either 100 mg/kg or 50 mg/kg, the levels of plasma glucose and plasma insulin were significantly reduced.

These results suggest that the compounds have an excellent anti-diabetic effect.

Therefore, the compound represented by formula 1 of the present invention can be effectively used for the prevention or treatment of diseases caused by the abnormal activation of 11β -HSD1 such as non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder, metabolic syndrome, and other diseases mediated by the excessive activity of glucocorticoid.

The compound represented by formula 1 of the present invention can be formulated in various forms according to the purpose of use. The followings are the examples of formulations comprising the compound of formula 1 of the present invention as an active ingredient, but these cannot limit the present invention thereto.

20 Manufacturing Example 1: Preparation of powders

The compound of formula 1	2 g
---------------------------	-----

Lactose	1 g
---------	-----

Powders were prepared by mixing all the above components, which were filled in airtight packs

according to the conventional method for preparing powders.

Manufacturing Example 2: preparation of tablets

5	The compound of formula 1	100 mg
	Corn starch	100 mg
	Lactose	100 mg
	Magnesium stearate	2 mg

Tablets were prepared by mixing all the above
 10 components by the conventional method for preparing tablets.

Manufacturing Example 3: Preparation of capsules

	The compound of formula 1	100 mg
15	Corn starch	100 mg
	Lactose	100 mg
	Magnesium stearate	2 mg

Capsules were prepared by mixing all the above
 components, which were filled in gelatin capsules
 20 according to the conventional method for preparing capsules.

Manufacturing Example 4: Preparation of injectable solutions

25	The compound of formula 1	100 mg
----	---------------------------	--------

Mannitol	180 mg
$\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$	26 mg
Distilled water	2974 mg

Injectable solutions were prepared by mixing all
5 the above components by the conventional method for
preparing injectable solutions.

INDUSTRIAL APPLICABILITY

10 Since the compound of the present invention
selectively inhibits the activity of 11β -HSD1 (11β -
Hydroxysteroid dehydrogenase type 1), the compound of
the invention can be effectively used as a therapeutic
agent for the treatment of diseases caused by the
15 over-activation of 11β -HSD1 such as non-insulin
dependent type II diabetes, insulin resistance,
obesity, lipid disorder, metabolic syndrome, and other
diseases or condition mediated by the excessive
activity of glucocorticoid.

20

Those skilled in the art will appreciate that
the conceptions and specific embodiments disclosed in
the foregoing description may be readily utilized as
a basis for modifying or designing other embodiments
25 for carrying out the same purposes of the present


invention. Those skilled in the art will also appreciate that such equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended Claims.

5

WHAT IS CLAIMED IS:

1. A compound represented by formula 1 having the activity of inhibiting 11 β -HSD1 enzyme or a
 5 pharmaceutically acceptable salt thereof:

[Formula 1]






(In formula 1,

X is carbonyl or sulfonyl;






10 R^1 is C_{1-4} straight or branched alkyl, C_{3-7} cycloalkyl, C_{5-12} aryl, C_{5-12} arylamine, 5-8 membered monocyclic or 8-12 membered bicyclic heterocycloalkyl, or 5-8 membered monocyclic or 8-12 membered bicyclic heteroaryl;



15 R^2 and R^3 are independently hydrogen, C_{1-4} straight or branched alkyl, C_{5-8} cycloalkyl, C_{5-8} bicycloalkyl, C_{5-8} aryl, 1-adamantyl, or 2-adamantyl, wherein the 1-adamantyl or 2-adamantyl is unsubstituted or substituted with one or more substituents selected
 20 from the group consisting of hydroxy, amino, aldehyde, hydroxy C_{1-4} straight or branched alkyl, hydroxycarbonyl, aminocarbonyl, N-

hydroxyaminocarbonyl, nitrile, hydroxycarbonyl C₁₋₄
straight or branched alkyl, aminocarbonyl C₁₋₄ straight

or branched alkyl, , , and ;

R⁴ is hydrogen, C₁₋₄ straight or branched alkyl,
5 hydroxy C₁₋₄ straight or branched alkyl, or sodium;

A is , , , , ,

, or , wherein Y is unsubstituted or
halogen;

wherein the said R¹ - R³, the aryl is
10 unsubstituted or substituted with one or more
substituents selected from the group consisting of
hydroxy, halogen, amino, C₁₋₄ straight or branched
alkyl, C₁₋₄ straight or branched haloalkyl, C₁₋₄
straight or branched haloalkyloxy, C₁₋₄ straight or
15 branched alkyloxy, -O-Z¹-NZ²Z³, -O-NZ²Z³, and -Z¹-NZ²Z³,
wherein the Z₁ is C₁₋₄ straight or branched alkyl, and
the Z² and Z³ are independently hydrogen or C₁₋₄
straight or branched alkyl;

wherein the said $R^1 - R^3$, the heterocycloalkyl or heteroaryl is monocyclic or bicyclic including one or more heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, wherein the
 5 heterocycloalkyl or heteroaryl is unsubstituted or substituted with one or more substituents selected from the group consisting of trifluoro C_{1-4} straight or branched alkyl, pyrimidinyl, C_{1-4} straight or branched alkyl, and C_{1-4} straight or branched alkyloxycarbonyl,
 10 and also the heterocycloalkyl can be fused-bicyclic with C_{5-12} aryl or heteroaryl, and also the heteroaryl can be fused-bicyclic with C_{3-7} cycloalkyl or heterocycloalkyl;

wherein the R^1 and R^4 can form 5-8 membered
 15 monocyclic or 8-12 membered bicyclic heterocycloalkyl with X and N which are attached to the R^1 and R^4 ;

wherein the R^4 can form 5-8 membered monocyclic heterocycloalkyl with N and A which are attached to the R^4).

20


2. The compound according to claim 1,

X is carbonyl or sulfonyl;


R^1 is C_{1-4} straight or branched alkyl, C_{3-7} cycloalkyl, C_{5-6} aryl, C_{5-6} arylamine, 5-6 membered
 25 monocyclic or 9-10 membered bicyclic heterocycloalkyl,

or 5-6 membered monocyclic or 9-10 membered bicyclic heteroaryl;

R^2 and R^3 are independently hydrogen, methyl, ethyl, propyl, C_{6-7} cycloalkyl, C_{6-8} bicycloalkyl, C_{5-6} aryl, 1-adamantyl, or 2-adamantyl, wherein the 1-adamantyl or 2-adamantyl is unsubstituted or substituted with one or more substituents selected from the group consisting of hydroxy, amino, aldehyde, hydroxymethyl, hydroxyethyl, hydroxycarbonyl, aminocarbonyl, N-hydroxyaminocarbonyl, nitrile, hydroxycarbonylmethyl, hydroxycarbonylethyl,



R^4 is hydrogen, methyl, ethyl, propyl, hydroxymethyl, hydroxyethyl, or sodium;




wherein the said $R^1 - R^3$, the aryl is unsubstituted or substituted with one or more substituents selected from the group consisting of halogen, trifluoromethyl, trifluoroethyl, trifluoromethoxy, trifluoroethoxy, methoxy, ethoxy, propoxy, methyl, ethyl, 1,1,1-trifluoro-2-ol-isopropyl, $-O(CH_2)_nNZ^2Z^3$, and $-(CH_2)_nNZ^2Z^3$, the n is 1 or 2, and the Z^2 and Z^3 are independently hydrogen, methyl or ethyl;

wherein the said $R^1 - R^3$, the heterocycloalkyl or heteroaryl contains one or more heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, wherein the heterocycloalkyl or heteroaryl is unsubstituted or substituted with one or more substituents selected from the group consisting of trifluoromethyl, trifluoroethyl, pyrimidinyl, methyl, ethyl, and isobutyloxycarbonyl, and also the heterocycloalkyl can be fused-bicyclic with C_{5-12} aryl or heteroaryl, and also the heteroaryl can be fused-bicyclic with C_{3-7} cycloalkyl or heterocycloalkyl;

wherein the R^1 and R^4 can form 5-6 membered monocyclic or 8-9 membered bicyclic heterocycloalkyl with X and N which are attached to the R^1 and R^4 ;


R^4 can form 5-6 membered monocyclic heterocycloalkyl with N and A which are attached to the R^4 .

- 5 3. The compound according to claim 1,
X is carbonyl or sulfonyl,
 R^1 is




R^2 is $-H$ or $-(CH_2)_2CH_3$,

 R^3 is



R^4 is -H, -CH₃, -CH₂CH₃, -CH₂CH₂OH, or -Na,

A is




5

4. The compound according to claim 1,


X is sulfonyl;

R¹ is phenyl, wherein the phenyl is unsubstituted or substituted with one or more substituents selected from the group consisting of
 5 halogen and trifluoromethyl, and the halogen can be substituted with one or more;

R² is -H or -CH₃;



R⁴ is -H,




5. The compound according to claim 1,


X is sulfonyl;

R¹ is phenyl, wherein the phenyl is
 15 unsubstituted or substituted with one or more substituents selected from the group consisting of halogen and trifluoromethyl, and the halogen can be substituted with one or more;

R² is -H;



R^4 is -H or -Na,




6. The compound according to claim 1,


5 X is sulfonyl;

R^1 is phenyl, wherein the phenyl is unsubstituted or substituted with one or more substituents selected from the group consisting of halogen and trifluoromethyl, and the halogen can be
10 substituted with one or more;

R^2 is -H;



R^4 is -H or -Na,



15

7. The compound according to claim 1, wherein the compound represented by formula 1 is selected from the group consisting of the following compounds:

(1) N-(adamantan-2-yl)-1-[(3-chloro-2-methylbenzenesulfonylamino)methyl]cyclopropanecarboxamide;
20

(2) (1-[(2-fluoro-benzenesulfonylamino)methyl]-N-(5-hydroxyadamantan-2-yl)cyclopropanecarboxamide;

(3) E-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
5 adamantan-1-carboxylic acid amide;

(4) Z-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

(5) E-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
10 adamantan-1-carboxylic acid amide;

(6) Z-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

15 (7) E-4-[1-((3-chloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

(8) E-4-[1-((3-chloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
20 adamantan-1-carboxylic acid amide;

(9) E-4-[1-((3,5-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

- (10) E-4-[1-((2-fluoro-6-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (11) E-4-[1-((2,3-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (12) E-4-[1-((2,4,6-trifluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (13) E-4-[1-((2-fluoro-*N*,6-dimethyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (14) E-4-[1-((2,4-dichloro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (15) E-4-[1-((4-chloro-2-fluoro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (16) E-4-[1-((4,5-dichloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (17) E-4-[1-((furan-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (18) E-4-[1-(3,5-dichloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (19) E-4-[1-((thiophen-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (20) E-4-[1-((2-(trifluoromethyl)-4-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (21) E-4-[1-((3,4-difluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (22) E-4-[1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (23) E-4-[1-((4-trifluoromethoxybenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (24) E-4-[1-((2,3-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (25) E-4-[1-((3,4-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (26) E-4-[1-((1-methyl-1H-indole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (27) E-4-[1-((1-methyl-1H-pyrazole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (28) E-4-[1-((benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (29) E-4-[1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid;
- (30) N-(bicyclo[2.2.1]heptan-2-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;
- 15 (31) N-(adamantan-1-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;
- 20 (32) E-4-[1-((N-ethyl-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (33) E-3-(4-(1((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido)adamantan-1-yl)propanoic acid;
- 25

(34) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-
1-((2-fluoro-
benzenesulfonylamino)methyl)cyclopropanecarboxamide;

(35) E-N-(5-aminoadamantan-2-yl)-1-((2-fluoro-
5 benzenesulfonylamino)methyl)cyclopropanecarboxamide
hydrochloride;

(36) E-4-[1-((2,4,5-trifluoro-
benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

10 (37) E-4-[1-((4-chloro-
benzenesulfonylamino)methyl)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

(38) E-4-[1-(3-chloro-
benzenesulfonylamino)cyclopropanecarboxamido]-
15 adamantan-1-carboxylic acid amide;

(39) E-4-[1-(2-fluoro-
benzenesulfonylamino)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

(40) E-4-[1-(2-fluoro-6-methyl-
20 benzenesulfonylamino)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

(41) E-4-[1-(3-chloro-2-methyl-
benzenesulfonylamino)cyclopropanecarboxamido]-
adamantan-1-carboxylic acid amide;

- (42) E-4-[1-(4-fluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (43) E-4-[1-(2,4-dichloro-5-methyl-
5 benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (44) E-4-[1-(2,4-difluorochloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (45) E-4-[1-(2-fluoro-4,5-dichloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (46) E-4-[1-(2-fluoro-4-chloro-5-methyl-benzenesulfonylamino)cyclopropanecarboxamido]-
15 adamantan-1-carboxylic acid amide;
- (47) E-4-[1-(2,3,4-trifluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (48) E-4-[1-(thiophen-2-
20 sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (49) E-4-[3-(6-trifluoromethyl-pyridin-2-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(50) E-4-[1-(1-methyl-1H-indole-7-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(51) 1-(3-chloro-benzenesulfonylamino)-N-(4-fluoro-2-(trifluoromethyl)phenyl)cyclopropanecarboxamide;

(52) E-4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;

(53) 1-(3-chloro-benzenesulfonylamino)-N-[4-(1,1,1-trifluoro-2-hydroxypropan-2-yl)phenyl]cyclopropanecarboxamide;

(54) E-4-[1-(3-chloro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(55) N-(bicyclo[2.2.1]heptan-2-yl)-1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamide;

(56) E-4-[1-(1,1-dioxido benzo[d]isothiazol-2(3H)-yl)cyclopropanecarboxamido]adamantan-1-carboxylic acid amide;

(57) E-4-[1-(3,4-difluoro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(58) E-4-[1-(1-methyl-1H-pyrazole-5-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(59) E-4-[1-(2,3-difluoro-5-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(60) E-4-[1-(benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;

10 (61) E-4-[1-(2-fluoro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(62) E-4-[1-(3-chloro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(63) E-4-[1-(2-fluorobenzamido)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(64) E-N-[1-(5-carbamoyladamantan-2-yl)carbamoyl)cyclopropyl]-5-(trifluoromethyl)pyrrolidineamide;

(65) E-4-(1-(benzo[d][1,3]dioxol-5-sulfonylamino)cyclopropanecarboxamino)adamantan-1-carboxylic acid amide;

- (66) 1-(3-chloro-benzenesulfonylamino)-N-cycloheptyl-N-propylcyclopropanecarboxamide;
- (67) E-4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid amide;
- 5 (68) E-4-[3-(2-fluoro-benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
- (69) E-4-[3-(benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
- (70) E-4-[3-(3-chloro-2-methyl-
10 benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
- (71) E-4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid amide;
- (72) E-4-[2-(2-fluoro-N-methyl-
15 benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (73) E-4-[2-(3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (74) E-N-(5-cyanoadamantan-2-yl)-2-(2-fluoro-
20 benzenesulfonylamino)-2-methylpropanamide;
- (75) E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(N'-carbamimidoyl)adamantan-2-yl]-2-methylpropanamide;
- (76) E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(hydroxymethyl)adamantan-2-yl]-2-methylpropanamide;

- (77) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-formyladamantan-2-yl)-2-methylpropanamide;
- (78) E-[4-(2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]methyl-4-methylbenzenesulfonate;
- (79) E-2-[4-(2-(2-fluorobenzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]acetic acid;
- (80) E-N-[5-(2-amino-2-oxoethyl)adamantan-2-yl]-2-(2-fluorobenzenesulfonylamino)-2-methylpropanamide;
- 10 (81) E-4-[2-methyl-2-(benzenesulfonylamino)propanamido]adamantan-1-carboxylic acid amide;
- (82) E-4-[2-(2-fluoro-3-chlorobenzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- 15 (83) E-4-[2-(3,5-difluoro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (84) E-4-[2-(2,6-difluoro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- 20 (85) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide;
- (86) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide;

- (87) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide;
- (88) E-4-[2-(4-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (89) E-4-[2-(2,5-dichloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (90) 2-(3-chloro-benzenesulfonylamino)-N-cycloheptyl-2-methyl-N-propylpropanamide;
- 10 (91) E-4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (92) E-4-[2-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (93) E-4-[4-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 15 (94) E-4-[3-(4-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (95) E-4-[3-(3-chloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 20 (96) E-4-[3-(3-chloro-2-methyl-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (97) E-4-{3-[(3-chloro-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid
- 25 amide;

(98) E-4-[3-N-(2-hydroxyethyl)-2-(trifluoromethyl-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide;

(99) E-4-{3-[(2-trifluoromethyl-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;

(100) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl](2-(trifluoromethyl)benzenesulfonylamide);

10 (101) E-N-(5-carbamoyladamantan-2-yl)-5-[(N-methyl-2-(trifluoromethyl)benzenesulfonylamino)nicotinamide];

(102) E-4-[3-(thiophen-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

15 (103) E-4-[3-(furan-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(104) E-4-[3-(pyridin-3-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

(105) E-4-(3-(benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;

20 (106) E-4-[3-[(2-chloro-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide;

- (107) E-4-[3-[(2,4-dimethyl-thiazol-5-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (108) E-4-[3-(3,5-dimethyl-1*H*-pyrazole-4-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (109) E-*N*-(5-hydroxy-adamantan-2-yl)-3-benzenesulfonylamino-benzamide;
- (111) E-*N*-(5-hydroxyadamantan-2-yl)-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (113) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl]-2-fluoro-3-chloro-benzenesulfonylamide;
- (114) E-4-[3-(3-chloro-4-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (115) E-4-[3-[(2-(trifluoromethyl)benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (116) E-4-[3-(2-chloro-4-bromo-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (117) E-4-[3-(2,4,6-trichloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

- (118) E-4-[3-(3-chloro-5-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (119) E-4-[3-(3,5-dichloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (120) E-4-[3-(3-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (121) E-4-[3-(2,4-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (122) E-4-[3-(2,5-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (123) E-4-[3-(2,6-difluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (124) E-N-cycloheptyl-N-propyl-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (125) E-4-[2-fluoro-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (126) E-4-[2-chloro-5-(3-chloro-benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (127) E-4-[3-(3-chlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;

- (128) E-4-[4-chloro-3-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (129) E-4-[2-chloro-5-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (130) E-4-[3-(3,5-dichlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (131) E-4-[5-(3,5-dichlorobenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (132) E-4-[2-fluoro-3-(3-chlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (133) E-4-[2-chloro-5-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (134) E-4-[4-chloro-3-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (135) E-4-[5-(3-chloro-4-methoxybenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (136) E-4-(3-(4-chloro-benzenesulfonylamino)benzamido)-adamantan-1-carboxylic acid amide;

- (137) E-4-(3-(4-chloro-benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;
- (138) E-4-(3-(cyclopropanesulfonamido)benzamido)adamantan-1-carboxylic acid amide;
- (139) E-4-(3-(1-methylethylsulfonamido)benzamido)adamantan-1-carboxylic acid amide;
- (140) E-[3,4-dihydro-1H-isoquinolin-2-carboxylic acid-1-[(5-carbamoyl-adamantan-2-ylcarbamoyl)cyclopropylmethyl]-amide;
- (141) E-3,4-dihydro-2H-quinolin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;
- (142) E-piperidin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;
- (143) E-4-{[1-5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-carbamoyl}-3,4-dihydro-2H-quinolin-1-carboxylic acid-butylester;
- (144) E-4-pyrimidin-2-yl-piperazin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;

(145) E-4-({1[(3-phenyl-ureido)methyl]-cyclopropanecarbonyl}amino)-adamantan-1-carboxylic acid amide;

(146) E-3,4-dihydro-2H-quinoxalin-1-carboxylic acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-amide;

(147) E-3,4-dihydro-1H-isoquinolin-2-carboxylic acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide;

(148) E-3,4-dihydro-2H-quinolin-1-carboxylic acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide; and

(149) E-2-(2-fluoro-benzamido)-thiazol-4-carboxylic acid(5-carbamoyl-adamantan-2-yl)amide.

15

8. The compound according to claim 1, wherein the compound represented by formula 1 is selected from the group consisting of the following compounds:

(1) N-(adamantan-2-yl)-1-[(3-chloro-2-methylbenzenesulfonylamino)methyl]cyclopropanecarboxamide;

(2) (1-[(2-fluoro-benzenesulfonylamino)methyl]-N-(5-hydroxyadamantan-2-yl)cyclopropanecarboxamide;

- (3) E-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (4) Z-4-[1-((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (5) E-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (6) Z-4-[1-((3-chloro-2-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (7) E-4-[1-((3-chloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (8) E-4-[1-((3-chloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (9) E-4-[1-((3,5-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (10) E-4-[1-((2-fluoro-6-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (11) E-4-[1-((2,3-difluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (12) E-4-[1-((2,4,6-trifluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (13) E-4-[1-((2-fluoro-N,6-dimethyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (14) E-4-[1-((2,4-dichloro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (15) E-4-[1-((4-chloro-2-fluoro-5-methyl-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (16) E-4-[1-((4,5-dichloro-2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (17) E-4-[1-((furan-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (18) E-4-[1-(3,5-dichloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (19) E-4-[1-((thiophen-2-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (20) E-4-[1-((2-(trifluoromethyl)-4-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (21) E-4-[1-((3,4-difluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (22) E-4-[1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (23) E-4-[1-((4-trifluoromethoxybenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (24) E-4-[1-((2,3-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (25) E-4-[1-((3,4-difluorobenzeneamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 20 (26) E-4-[1-((1-methyl-1H-indole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(27) E-4-[1-((1-methyl-1H-pyrazole-5-sulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(28) E-4-[1-((benzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

(29) E-4-[1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid;

10 (30) N-(bicyclo[2.2.1]heptan-2-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;

(31) N-(adamantan-1-yl)-1-((2-fluoro-N-methylbenzenesulfonylamino)methyl)cyclopropanecarboxamido;

(32) E-4-[1-((N-ethyl-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

20 (33) E-3-(4-(1((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamido)adamantan-1-yl)propanoic acid;

(34) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-1-((2-fluorobenzenesulfonylamino)methyl)cyclopropanecarboxamide;

(35) E-N- (5-aminoadamantan-2-yl) -1- ((2-fluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamide hydrochloride; and

(36) E-4- [1- ((2,4,5-trifluoro-benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
5 adamantan-1-carboxylic acid amide.

9. The compound according to claim 1, wherein the compound represented by formula 1 is selected from the
10 group consisting of the following compounds:

(37) E-4- [1- ((4-chloro-benzenesulfonylamino)methyl)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(38) E-4- [1- (3-chloro-benzenesulfonylamino)cyclopropanecarboxamido] -
15 adamantan-1-carboxylic acid amide;

(39) E-4- [1- (2-fluoro-benzenesulfonylamino)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

20 (40) E-4- [1- (2-fluoro-6-methyl-benzenesulfonylamino)cyclopropanecarboxamido] -
adamantan-1-carboxylic acid amide;

(41) E-4- [1- (3-chloro-2-methyl-benzenesulfonylamino)cyclopropanecarboxamido] -
25 adamantan-1-carboxylic acid amide;

- (42) E-4-[1-(4-fluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (43) E-4-[1-(2,4-dichloro-5-methyl-
5 benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (44) E-4-[1-(2,4-difluorochloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 10 (45) E-4-[1-(2-fluoro-4,5-dichloro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (46) E-4-[1-(2-fluoro-4-chloro-5-methyl-benzenesulfonylamino)cyclopropanecarboxamido]-
15 adamantan-1-carboxylic acid amide;
- (47) E-4-[1-(2,3,4-trifluoro-benzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (48) E-4-[1-(thiophen-2-
20 sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (49) E-4-[3-(6-trifluoromethyl-pyridin-2-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (50) E-4-[1-(1-methyl-1H-indole-7-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (51) 1-(3-chloro-benzenesulfonylamino)-N-(4-fluoro-2-(trifluoromethyl)phenyl)cyclopropanecarboxamide;
- (52) E-4-[1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;
- 10 (53) 1-(3-chloro-benzenesulfonylamino)-N-[4-(1,1,1-trifluoro-2-hydroxypropan-2-yl)phenyl]cyclopropanecarboxamide;
- (54) E-4-[1-(3-chloro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- 15 (55) N-(bicyclo[2.2.1]heptan-2-yl)-1-(3-chloro-benzenesulfonylamino)cyclopropanecarboxamide;
- (56) E-4-[1-(1,1-dioxido[benzo[d]isothiazol-2(3H)-yl)cyclopropanecarboxamido]adamantan-1-carboxylic acid
- 20 amide;
- (57) E-4-[1-(3,4-difluoro-benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;

- (58) E-4-[1-(1-methyl-1H-pyrazole-5-sulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (59) E-4-[1-(2,3-difluoro-5 benzeneamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (60) E-4-[1-(benzenesulfonylamino)cyclopropanecarboxamido]-N-hydroxyadamantan-1-carboxylic acid amide;
- 10 (61) E-4-[1-(2-fluoro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (62) E-4-[1-(3-chloro-N-methylbenzenesulfonylamino)cyclopropanecarboxamido]-15 adamantan-1-carboxylic acid amide;
- (63) E-4-[1-(2-fluorobenzamido)cyclopropanecarboxamido]-adamantan-1-carboxylic acid amide;
- (64) E-N-[1-(5-carbamoyladamantan-2-yl)carbamoyl)cyclopropyl]-5-20 (trifluoromethyl)pyrrolidineamide;
- (65) E-4-(1-(benzo[d][1,3]dioxol-5-sulfonylamino)cyclopropanecarboxamino)adamantan-1-carboxylic acid amide; and

(66) 1-(3-chloro-benzenesulfonylamino)-N-cycloheptyl-N-propylcyclopropanecarboxamide.

10. The compound according to claim 1, wherein
 5 the compound represented by formula 1 is selected from the group consisting of the following compounds:

(67) E-4-(3-(3-chloro-benzenesulfonylamino)-2,2-dimethylpropanamido)adamantan-1-carboxylic acid amide;

(68) E-4-[3-(2-fluoro-benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;

(69) E-4-[3-(benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide;
 and

(70) E-4-[3-(3-chloro-2-methyl-
 15 benzenesulfonylamino)-2,2-dimethylpropanamido]adamantan-1-carboxylic acid amide.

11. The compound according to claim 1, wherein
 the compound represented by formula 1 is selected from
 20 the group consisting of the following compounds:

(71) E-4-[2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido]-adamantan-1-carboxylic acid amide;

(72) E-4-[2-(2-fluoro-N-methyl-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-
 25 carboxylic acid amide;

- (73) E-4-[2-(3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (74) E-N-(5-cyanoadamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide;
- 5 (75) E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(N'-carbamimidoyl)adamantan-2-yl]-2-methylpropanamide;
- (76) E-2-(2-fluoro-benzenesulfonylamino)-N-[5-(hydroxymethyl)adamantan-2-yl]-2-methylpropanamide;
- (77) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-
10 formyladamantan-2-yl)-2-methylpropanamide;
- (78) E-[4-(2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamido)adamantan-1-yl]methyl-4-methylbenzenesulfonate;
- (79) E-2-[4-(2-(2-fluorobenzenesulfonylamino)-2-
15 methylpropanamido)adamantan-1-yl]acetic acid;
- (80) E-N-[5-(2-amino-2-oxoethyl)adamantan-2-yl]-2-(2-fluorobenzenesulfonylamino)-2-methylpropanamide;
- (81) E-4-[2-methyl-2-(benzenesulfonylamino)propanamido]adamantan-1-
20 carboxylic acid amide;
- (82) E-4-[2-(2-fluoro-3-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (83) E-4-[2-(3,5-difluoro-benzenesulfonylamino)-
25 2-methylpropanamido]adamantan-1-carboxylic acid amide;

- (84) E-4-[2-(2,6-difluoro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (85) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide;
- (86) E-N-(5-(3-amino-3-oxopropyl)adamantan-2-yl)-2-(2-fluoro-benzenesulfonylamino)-2-methylpropanamide;
- (87) E-2-(2-fluoro-benzenesulfonylamino)-N-(5-(hydrazinecarbonyl)adamantan-2-yl)-2-methylpropanamide;
- (88) E-4-[2-(4-chloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- (89) E-4-[2-(2,5-dichloro-benzenesulfonylamino)-2-methylpropanamido]adamantan-1-carboxylic acid amide;
- and
- (90) 2-(3-chloro-benzenesulfonylamino)-N-cycloheptyl-2-methyl-N-propylpropanamide.

12. The compound according to claim 1, wherein the compound represented by formula 1 is selected from the group consisting of the following compounds:

- (91) E-4-[3-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (92) E-4-[2-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;

- (93) E-4-[4-(2-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (94) E-4-[3-(4-fluoro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 5 (95) E-4-[3-(3-chloro-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (96) E-4-[3-(3-chloro-2-methyl-benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 10 (97) E-4-{3-[(3-chloro-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;
- (98) E-4-[3-N-(2-hydroxyethyl)-2-(trifluoromethyl-benzenesulfonylamino)benzamido]-adamantan-1-carboxylic acid amide;
- 15 (99) E-4-{3-[(2-trifluoromethyl-benzenesulfonyl)-methylamino]-benzamido}-adamantan-1-carboxylic acid amide;
- (100) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl](2-(trifluoromethyl)benzenesulfonylamide;
- 20 (101) E-N-(5-carbamoyladamantan-2-yl)-5-[(N-methyl-2-(trifluoromethyl)benzenesulfonylamino]nicotinamide;

- (102) E-4-[3-(thiophen-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (103) E-4-[3-(furan-2-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 5 (104) E-4-[3-(pyridin-3-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (105) E-4-(3-(benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;
- (106) E-4-[3-[(2-chloro-10 benzenesulfonylamino]benzamido]-adamantan-1-carboxylic acid amide;
- (107) E-4-[3-[(2,4-dimethyl-thiazol-5-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 15 (108) E-4-[3-(3,5-dimethyl-1H-pyrazole-4-sulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (109) E-N-(5-hydroxy-adamantan-2-yl)-3-benzenesulfonylamino-benzamide;
- 20 (111) E-N-(5-hydroxyadamantan-2-yl)-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (113) E-sodium[3-((5-carbamoyladamantan-2-yl)carbamoyl)phenyl]-2-fluoro-3-chloro-benzenesulfonylamide;

- (114) E-4-[3-(3-chloro-4-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (115) E-4-[3-[(2-(trifluoromethyl)benzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (116) E-4-[3-(2-chloro-4-bromobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 10 (117) E-4-[3-(2,4,6-trichlorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (118) E-4-[3-(3-chloro-5-fluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 15 (119) E-4-[3-(3,5-dichlorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (120) E-4-[3-(3-fluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 20 (121) E-4-[3-(2,4-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (122) E-4-[3-(2,5-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- (123) E-4-[3-(2,6-difluorobenzenesulfonylamino)-benzamido]-adamantan-1-carboxylic acid amide;
- 25

- (124) E-*N*-cycloheptyl-*N*-propyl-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamide;
- (125) E-4-[2-fluoro-3-(2-(trifluoromethyl)benzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (126) E-4-[2-chloro-5-(3-chlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (127) E-4-[3-(3-chlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (128) E-4-[4-chloro-3-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (129) E-4-[2-chloro-5-(3,5-dichlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;
- (130) E-4-[3-(3,5-dichlorobenzenesulfonylamino)-4-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (131) E-4-[5-(3,5-dichlorobenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;
- (132) E-4-[2-fluoro-3-(3-chlorobenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

(133) E-4-[2-chloro-5-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

(134) E-4-[4-chloro-3-(3-chloro-4-methoxybenzenesulfonylamino)benzamido]adamantan-1-carboxylic acid amide;

(135) E-4-[5-(3-chloro-4-methoxybenzenesulfonylamino)-2-fluorobenzamido]adamantan-1-carboxylic acid amide;

10 (136) E-4-(3-(4-chloro-benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;

(137) E-4-(3-(4-chloro-benzenesulfonylamino-benzamido)-adamantan-1-carboxylic acid amide;

(138) E-4-(3-(cyclopropanesulfonamido)benzamido)adamantan-1-carboxylic acid amide; and

(139) E-4-(3-(1-methylethylsulfonamido)benzamido)adamantan-1-carboxylic acid amide.

20

13. The compound according to claim 1, wherein the compound represented by formula 1 is selected from the group consisting of the following compounds:

(140) E-[3,4-dihydro-1H-isoquinolin-2-carboxylic
acid-1-[(5-carbamoyl-adamantan-2-
ylcarbamoyl)cyclopropylmethyl]-amide;

(141) E-3,4-dihydro-2H-quinolin-1-carboxylic
5 acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-
cyclopropylmethyl]-amide;

(142) E-piperidin-1-carboxylic acid[1-(5-
carbamoyl-adamantan-2-ylcarbamoyl)-cyclopropylmethyl]-
amide;

10 (143) E-4-{[1-(5-carbamoyl-adamantan-2-
ylcarbamoyl)-cyclopropylmethyl]-carbamoyl}-3,4-
dihydro-2H-quinolin-1-carboxylic acid-butylester;

(144) E-4-pyrimidin-2-yl-piperazin-1-carboxylic
acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-
15 cyclopropylmethyl]-amide;

(145) E-4-({1[(3-phenyl-ureido)methyl]-
cyclopropanecarbonyl}amino)-adamantan-1-carboxylic acid
amide;

(146) E-3,4-dihydro-2H-quinoxalin-1-carboxylic
20 acid[1-(5-carbamoyl-adamantan-2-ylcarbamoyl)-
cyclopropylmethyl]-amide;


(147) E-3,4-dihydro-1H-isoquinolin-2-carboxylic
acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-
2-yl]-amide;

(148) E-3,4-dihydro-2H-quinolin-1-carboxylic acid[4-(5-carbamoyl-adamantan-2-ylcarbamoyl)-thiazol-2-yl]-amide; and

(149) E-2-(2-fluoro-benzamido)-thiazol-4-carboxylic acid(5-carbamoyl-adamantan-2-yl)amide.

14. A preparation method of the compound of claim 1 containing the step of reacting the compound represented by formula 2 with the compound represented by formula 3 in the presence of an organic solvent to prepare the compound of formula 1, as represented in the following reaction formula 1:

[Reaction Formula 1]



(In reaction formula 1,

R¹ - R⁴, X and A are as defined in formula 1 of claim 1).

15. A pharmaceutical composition for the prevention or treatment of diseases caused by the over-activation of 11β-HSD1 enzyme which comprises the

compound of claim 1 or a pharmaceutically acceptable salt thereof as an active ingredient.

16. The pharmaceutical composition according to
5 claim 15, wherein the disease caused by the over-activation of 11 β -HSD1 enzyme is selected from the group consisting of non-insulin dependent type II diabetes, insulin resistance, obesity, lipid disorder, metabolic syndrome, and other diseases mediated by the
10 excessive activity of glucocorticoid.