

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
30 September 2010 (30.09.2010)(10) International Publication Number
WO 2010/108856 A1

(51) International Patent Classification:

C07C 59/68 (2006.01) **C07D 277/24** (2006.01)
C07C 59/90 (2006.01) **C07D 317/54** (2006.01)
A61P 11/08 (2006.01) **C07D 317/58** (2006.01)
C07D 209/08 (2006.01) **C07D 319/18** (2006.01)
C07D 213/38 (2006.01) **C07D 333/16** (2006.01)
C07D 213/61 (2006.01) **C07D 333/20** (2006.01)
C07D 239/26 (2006.01) **C07D 333/24** (2006.01)

(21) International Application Number:

PCT/EP2010/053580

(22) International Filing Date:

19 March 2010 (19.03.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/162,352 23 March 2009 (23.03.2009) US

(71) Applicant (for all designated States except US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DOMINIQUE, Romyr** [CA/US]; 1124 Crossing Way, Wayne, New Jersey 07470 (US). **GOODNOW, Robert Alan** [US/US]; 799 Long Hill Road, Morris County, Gillette, New Jersey 07933 (US). **KOWALCZYK, Agnieszka** [US/US]; 14 Autumn Drive, Mine Hill, New Jersey 07803 (US). **QIAO, Qi** [CA/US]; 21 D Alfred Lane, Bloomfield, New Jersey 07003 (US). **SIDDURI, Achyutharao** [US/US];

22 Washington Court, Livingston, New Jersey 07039 (US). **TILLEY, Jefferson Wright** [US/US]; 19 Evergreen Drive, North Caldwell, New Jersey 07006 (US).


(74) Agent: **HEIROTH, Ulrike**; Grenzacherstrasse 124, CH-4070 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report (Art. 21(3))

(54) Title: LEUKOTRIENE B₄ INHIBITORS

(57) Abstract: Provided herein are compounds of the formula (I) as well as pharmaceutically acceptable salts thereof, wherein the substituents are as those disclosed in the specification. These compounds, and the pharmaceutical compositions containing them, are useful for the treatment of diseases such as, for example, COPD.

LEUKOTRIENE B₄ INHIBITORS

The invention is directed to compounds of formula (I):



- 5 or pharmaceutically acceptable salts thereof, a pharmaceutical preparation comprising the same and their use for the preparation of medicaments. These compounds inhibit the interaction of leukotriene B₄ (LTB₄) pro-inflammatory lipid mediator binding to BLT-1 and BLT-2 receptors resulting in amelioration of disease states resulting from excessive inflammatory response, such as, for
10 example, severe asthma and chronic obstructive pulmonary disease (COPD).

All documents cited or relied upon below are expressly incorporated herein by reference.

- LTB₄ is a potent pro-inflammatory lipid mediator derived from arachidonic acid
15 through the 5-lipoxygenase signaling pathway. LTB₄ is produced by multiple cell types such as neutrophils, monocytes, macrophages, keratinocytes, lymphocytes and mast cells. It functions as a chemoattractant and as an activator of neutrophil cells. It has been shown that LTB₄ effects its action through the agonism of G-protein coupled receptors BLT-1 and BLT-2. (Prostaglandins, Leukotrienes and
20 Essential Fatty Acids 69, 2003, 123-134.)


LTB₄ is considered to be an important mediator of acute and chronic inflammatory diseases. Increased levels of LTB₄ have been detected in the lungs of patients with severe asthma and COPD. Thus, it is anticipated that an effective inhibitor of

- 2 -

the action of LTB₄ and BLT-1 and -2 would provide effective therapy for the treatment of inflammatory conditions such as asthma and COPD.

A need exists in the art for LTB₄ inhibitors that have efficacy for the treatment of
5 diseases such as COPD.


The present invention pertains to inhibitors of LTB₄. In a preferred embodiment, the invention provides for pharmaceutical compounds of the formula (I):



10

as well as pharmaceutically acceptable salts thereof, that are useful as inhibitors of LTB₄.

In an embodiment of the present invention, provided is a compound of formula (I):



15

wherein:

R1 is lower alkyl, alkoxy, alkoyl, dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl; and


20 R2 is benzo[1,3]dioxol, dihydro-benzo[1,4]dioxin, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted

- 3 -

with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl,

or a pharmaceutically acceptable salt thereof.

- 5 In another embodiment of the present invention, provided is a compound of formula (I):



wherein:

- 10 R1 is lower alkyl, alkoxy, alkoyl, dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl; and

- 15 R2 is benzo[1,3]dioxol, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl,

or a pharmaceutically acceptable salt thereof.

- 20 In another embodiment of the present invention, provided is a pharmaceutical composition, comprising a therapeutically effective amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

- 25 In a further preferred embodiment, R1 is lower alkyl, alkoxy or alkoyl and R2 is benzo[1,3]dioxol, dihydrobenzo[1,4]dioxin, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-

- 4 -

substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl.

5

In a further preferred embodiment, R1 is lower alkyl, alkoxy or alkoyl and R2 is benzo[1,3]dioxol, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or
10 mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl.

In an another preferred embodiment, R1 is dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl
15 and R2 is benzo[1,3]dioxol, dihydro-benzo[1,4]dioxin, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri- substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with
20 lower alkyl.

In a still another preferred embodiment, R1 is dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl and R2 is benzo[1,3]dioxol, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted
25 with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl.

In certain embodiment of formula (I), R1 is hydroxymethyl, methoxymethyl, ethoxy, ethoxymethyl, benzyloxy, acetyl or propionyl.
30

- 5 -

In an another embodiment of formula (I), R1 is dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl.

In certain embodiment of formula (I), R2 is phenyl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or

5 methanesulfonyl.

In certain embodiment of formula(I), R2 is thiophene, pyrimidine or pyridine, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy.

10

It is to be understood that the terminology employed herein is for the purpose of describing particular embodiments, and is not intended to be limiting. Further, although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the

15 preferred methods, devices and materials are now described.

As used herein, the term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more

20 preferably one to ten carbon atoms.

The term "cycloalkyl" refers to a monovalent carbocyclic radical of three to seven, preferably three to six carbon atoms. This term is further exemplified by radicals such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. In a preferred

25 embodiment, the "cycloalkyl" moieties can optionally be substituted with one, two, three or four substituents, wherein each substituent is independently, for example, hydroxy, alkyl, alkoxy, halogen or amino, unless otherwise specifically indicated.

Examples of cycloalkyl moieties include, but are not limited to, optionally

substituted cyclopropyl, optionally substituted cyclobutyl, optionally substituted

30 cyclopentyl, optionally substituted cyclopentenyl, optionally substituted cyclohexyl, optionally substituted cyclohexylene, optionally substituted cycloheptyl, and the like or those which are specifically exemplified herein.

The term "heterocycloalkyl" denotes a cyclic alkyl ring, wherein one, two or three of the carbon ring atoms is replaced by a heteroatom such as N, O or S. Examples of heterocycloalkyl groups include, but are not limited to, morpholine, thiomorpholine, piperazine, piperidine and the like. The heterocycloalkyl groups may be unsubstituted or substituted.

The term "lower alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent alkyl radical of one to six carbon atoms, preferably one to four carbon atoms. This term is further exemplified by radicals such as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, isobutyl, t-butyl, n-pentyl, 3-methylbutyl, n-hexyl, 2-ethylbutyl and the like.

The term "aryl" refers to an aromatic monovalent mono- or polycarbocyclic radical, such as phenyl or naphthyl, preferably phenyl.

The term "heteroaryl," alone or in combination with other groups, means a monocyclic or bicyclic radical of 5 to 12 ring atoms having at least one aromatic ring containing one, two, or three ring heteroatoms selected from N, O, and S, the remaining ring atoms being C. More specifically the term "heteroaryl" includes, but is not limited to, pyridinyl, furanyl, thienyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrimidinyl, pyrazolyl, pyrazinyl, pyrimidinyl, benzofuranyl, tetrahydrobenzofuranyl, isobenzofuranyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolyl, isoindolyl, benzoxazolyl, quinolyl, tetrahydroquinolyl, isoquinolyl, benzimidazolyl, benzisoxazolyl or benzothienyl, imidazo[1,2-a]-pyridinyl, imidazo[2,1-b]thiazolyl, and the derivatives thereof. Preferred heterocycle groups include, but are not limited to, thiophene, pyrimidine and pyridine. One or two ring carbon atoms of the heteroaryl group may be replaced with a carbonyl group. The heteroaryl group described above may be substituted independently with one, two, or three substituents, preferably one or two substituents such as, for example, halogen, hydroxy, C₁₋₆ alkyl, halo C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl sulfonyl, C₁₋₆ alkyl sulfinyl, C₁₋₆ alkylthio, amino, amino C₁₋₆ alkyl, mono- or di-substituted amino-C₁₋₆ alkyl, nitro, cyano, acyl, carbamoyl, mono- or di-substituted amino,

- 7 -

aminocarbonyl, mono- or di-substituted amino-carbonyl, aminocarbonyl C₁₋₆ alkoxy, mono- or di-substituted amino-carbonyl-C₁₋₆ alkoxy, hydroxy- C₁₋₆ alkyl, carboxyl, C₁₋₆ alkoxy carbonyl, aryl C₁₋₆ alkoxy, heteroaryl C₁₋₆ alkoxy, heterocyclyl C₁₋₆ alkoxy, C₁₋₆ alkoxycarbonyl C₁₋₆ alkoxy, carbamoyl C₁₋₆ alkoxy and carboxyl C₁₋₆ alkoxy, preferably halogen, hydroxy, C₁₋₆ alkyl, halo C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl sulfonyl, C₁₋₆ alkyl sulfinyl, C₁₋₆ alkylthio, amino, mono-C₁₋₆ alkyl substituted amino, di-C₁₋₆ alkyl substituted amino, amino C₁₋₆ alkyl, mono-C₁₋₆ alkyl substituted amino-C₁₋₆ alkyl, di-C₁₋₆ alkyl substituted amino-C₁₋₆ alkyl, nitro, carbamoyl, mono- or di-substituted amino-carbonyl, hydroxy- C₁₋₆ alkyl, carboxyl, C₁₋₆ alkoxy carbonyl and cyano.

The alkyl and aryl groups may be substituted or unsubstituted. Where substituted, there will generally be, for example, 1 to 3 substituents present, preferably 1 substituent. Substituents may include, for example: carbon-containing groups such as alkyl, aryl, arylalkyl (e.g. substituted and unsubstituted phenyl, substituted and unsubstituted benzyl); halogen atoms and halogen-containing groups such as haloalkyl (e.g. trifluoromethyl); oxygen-containing groups such as alcohols (e.g. hydroxyl, hydroxyalkyl, aryl(hydroxyl)alkyl), ethers (e.g. alkoxy, aryloxy, alkoxyalkyl, aryloxyalkyl), aldehydes (e.g. carboxaldehyde), ketones (e.g. alkylcarbonyl, alkylcarbonylalkyl, arylcarbonyl, arylalkylcarbonyl, arylcarbonylalkyl), acids (e.g. carboxy, carboxyalkyl), acid derivatives such as esters (e.g. alkoxycarbonyl, alkoxycarbonylalkyl, alkylcarbonyloxy, alkylcarbonyloxyalkyl), amides (e.g. aminocarbonyl, mono- or di-alkylaminocarbonyl, aminocarbonylalkyl, mono- or di-alkylaminocarbonylalkyl, arylaminocarbonyl), carbamates (e.g. alkoxycarbonylamino, arloxy carbonylamino, aminocarbonyloxy, mono- or di-alkylaminocarbonyloxy, arylaminocarbonyloxy) and ureas (e.g. mono- or di-alkylaminocarbonylamino or arylaminocarbonylamino); nitrogen-containing groups such as amines (e.g. amino, mono- or di-alkylamino, aminoalkyl, mono- or di-alkylaminoalkyl), azides, nitriles (e.g. cyano, cyanoalkyl), nitro; sulfur-containing groups such as thiols, thioethers, sulfoxides and sulfones (e.g. alkylthio, alkylsulfinyl, alkylsulfonyl, alkylthioalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, arylthio, arylsulfinyl, arylsulfonyl, arylthioalkyl, arylsulfinylalkyl, arylsulfonylalkyl); and heterocyclic groups containing one or more, preferably one, heteroatom, (e.g.

- 8 -

- thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, oxadiazolyl, thiadiazolyl, aziridinyl, azetidiny, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, tetrahydrofuranyl, pyranyl, pyronyl, pyridyl, pyrazinyl, pyridazinyl, piperidyl, hexahydroazepinyl, piperazinyl, morpholinyl, thianaphthyl, benzofuranyl, isobenzofuranyl, indolyl, oxyindolyl, isoindolyl, indazolyl, indolinyl, 7-azaindolyl, benzopyranyl, coumarinyl, isocoumarinyl, quinolinyl, isoquinolinyl, naphthridinyl, cinnolinyl, quinazolinyl, pyridopyridyl, benzoxazinyl, quinoxalinyl, chromenyl, chromanyl, isochromanyl, phthalazinyl and carbolinyl).
- 10 The lower alkyl groups may be substituted or unsubstituted. Where substituted, there will generally be, for example, 1 to 3 substituents present, preferably 1 substituent. The substituents include those that are noted for the alkyl group, *supra*, such as, for example, hydroxyl.
- 15 As used herein, the term "alkoxy" means alkyl-O-; and "alkoyl" means alkyl-CO-. Alkoxy or alkoyl groups may be substituted by, for example, one or more alkyl groups.

As used herein, the term "halogen" means a fluorine, chlorine, bromine or iodine radical, preferably a fluorine, chlorine or bromine radical, and more preferably a fluorine or chlorine radical.

Compounds of formula I can have one or more asymmetric carbon atoms and can exist in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates. The optically active forms can be obtained for example by resolution of the racemates, by asymmetric synthesis or asymmetric chromatography (chromatography with a chiral adsorbents or eluant). The invention embraces all of these forms.

30

As used herein, the term "pharmaceutically acceptable salt" means any pharmaceutically acceptable salt of the compound of formula (I). Salts may be prepared from pharmaceutically acceptable non-toxic acids and bases including

- 9 -

inorganic and organic acids and bases. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethenesulfonic, dichloroacetic, formic, fumaric, gluconic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, oxalic, pamoic, 5 pantothenic, phosphoric, succinic, sulfuric, tartaric, oxalic, p-toluenesulfonic and the like. Particularly preferred are fumaric, hydrochloric, hydrobromic, phosphoric, succinic, sulfuric and methanesulfonic acids. Acceptable base salts include alkali metal (e.g. sodium, potassium), alkaline earth metal (e.g. calcium, magnesium) and aluminium salts.

10

In the practice of the method of the present invention, an effective amount of any one of the compounds of this invention or a combination of any of the compounds of this invention or a pharmaceutically acceptable salt thereof, is administered via any of the usual and acceptable methods known in the art, either singly or in 15 combination. The compounds or compositions can thus be administered orally (e.g., buccal cavity), sublingually, parenterally (e.g., intramuscularly, intravenously, or subcutaneously), rectally (e.g., by suppositories or washings), transdermally (e.g., skin electroporation) or by inhalation (e.g., by aerosol), and in the form of solid, liquid or gaseous dosages, including tablets and suspensions. The administration 20 can be conducted in a single unit dosage form with continuous therapy or in a single dose therapy ad libitum. The therapeutic composition can also be in the form of an oil emulsion or dispersion in conjunction with a lipophilic salt such as pamoic acid, or in the form of a biodegradable sustained-release composition for subcutaneous or intramuscular administration.

25

Useful pharmaceutical carriers for the preparation of the compositions hereof, can be solids, liquids or gases; thus, the compositions can take the form of tablets, pills, capsules, suppositories, powders, enterically coated or other protected formulations (e.g. binding on ion-exchange resins or packaging in lipid-protein 30 vesicles), sustained release formulations, solutions, suspensions, elixirs, aerosols, and the like. The carrier can be selected from the various oils including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, and the like. Water, saline, aqueous dextrose, and glycols

- 10 -

are preferred liquid carriers, particularly (when isotonic with the blood) for injectable solutions. For example, formulations for intravenous administration comprise sterile aqueous solutions of the active ingredient(s) which are prepared by dissolving solid active ingredient(s) in water to produce an aqueous solution, and rendering the

5 solution sterile. Suitable pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, talc, gelatin, malt, rice, flour, chalk, silica, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol, and the like. The compositions may be subjected to conventional pharmaceutical additives such as preservatives, stabilizing agents,

10 wetting or emulsifying agents, salts for adjusting osmotic pressure, buffers and the like. Suitable pharmaceutical carriers and their formulation are described in Remington's Pharmaceutical Sciences by E. W. Martin. Such compositions will, in any event, contain an effective amount of the active compound together with a suitable carrier so as to prepare the proper dosage form for proper administration

15 to the recipient.

The dose of a compound of the present invention depends on a number of factors, such as, for example, the manner of administration, the age and the body weight of the subject, and the condition of the subject to be treated, and ultimately will be

20 decided by the attending physician or veterinarian. Such an amount of the active compound as determined by the attending physician or veterinarian is referred to herein, and in the claims, as a "therapeutically effective amount". For example, the dose of a compound of the present invention is typically in the range of about 1 to about 1000 mg per day. Preferably, the therapeutically effective amount is in an

25 amount of from about 1 mg to about 500 mg per day

It will be appreciated, that the compounds of general formula I in this invention may be derivatized at functional groups to provide derivatives which are capable of conversion back to the parent compound in vivo. Physiologically acceptable and


30 metabolically labile derivatives, which are capable of producing the parent compounds of general formula I in vivo are also within the scope of this invention.

The compounds of the present invention can be prepared by any conventional

- 11 -

means. Suitable processes for synthesizing these compounds are provided in the Examples. Generally, compounds of formula I can be prepared according to the Schemes described below. The sources of the starting materials for these reactions are also described.

5


Scheme 1

The compounds contained within this invention can be synthesized according to the following general synthetic strategies as shown below in Scheme 1. The synthesis of **1** may be effected by condensation of **3**, 4-[2-(2-carboxy-ethyl)-3-(6-E-hexyl)-phenoxy]-butyric acid, protected as a di-ester for R₀ = lower alkyl, preferably as a di-ethyl ester (R₀ = ethyl), and E is a leaving group, such as a halogen or mesylate with the fragment **2** wherein D is a nucleophile such as a hydroxyl group under standard conditions employed for the alkylation of phenols with primary halides or mesylates. Functional groups represented by symbol R₁ being halogen, nitro, and amino group and can be transformed to an aryl, aryl ether, or amine moiety before or after coupling to **3** according to chemistry described in this invention. Functional groups represented by symbol R₂ being hydroxy methyl, alkylated hydroxy methyl, cyano methyl, dimethyl amino methyl, ethers or ketones.

A synthesis of **3** for E = Br and R₀ = Et has been described in *Bioorganic & Medicinal Chemistry Letters* (1994), 4(24), 2883-8. A synthesis of **3** for E = Br and R₀ = Et is also shown below in Schemes 2 and 3.

Scheme 2

- 12 -




In Scheme 2, 2,3-dimethylphenol **4** is reacted with 4-bromo butyric acid ester **5** in presence of a base, preferably lithium hydride in aprotic solvent, preferably dimethylsulfoxide to obtain the dimethyl intermediate, **6**. Then, the more reactive methyl group at 2-position of **6** is selectively oxidized to the corresponding aldehyde **7** using an oxidizing conditions, such as copper(II)sulfate pentahydrate and potassium persulfate in a mixed solvent systems, preferably water and acetonitrile. The two carbon chain ester moiety can be selectively introduced by a modified Horner-Emmons condensation conditions from aldehyde **7** and triethylphosphonoacetate **8** in the presence of a base such as sodium ethoxide in a protic solvents, preferably ethanol. Then, the benzylic bromination of **9** is effected with N-bromosuccinimide in the presence of 2,2'-azobisisobutyronitrile (AIBN) in an aprotic solvents such as carbon tetrachloride or chlorobenzene or benzene. The hydroxy protected 5-carbon chain aldehyde **12** can be obtained by oxidation of a mono protected pentane-1,5-diol with any suitable oxidation conditions such as Swern oxidation or TEMPO oxidation, reactions well known to those skilled in the art. The protecting group on **11** and **12** can be any suitable protecting group for primary alcohols, for example t-butyldimethylsilyl group. Use and removal of protecting groups is well presented in the literature. For a leading reference, see

- 13 -

P.G. M. Wuts and T. W. Greene in Green's Protective Groups in Organic Synthesis, Wiley and Sons, 2007.


Scheme 3



- 5 As shown in Scheme 3, a one-pot Wittig condensation reaction is conducted first by making *in situ* Wittig salt from the benzylic bromide **10** and triphenylphosphine in acetonitrile and then the reaction of the resulting Wittig salt with the protected aldehyde **12** in 1,2-epoxybutane to obtain the olefinic intermediate **13** in a *cis* to *trans* ratio of ~1:3. The mixture of *cis* and *trans* compounds can be converted to
- 10 the corresponding alkyl bromide intermediate **16** by removal of the protecting group, using for example tetrabutyl ammonium fluoride for the case wherein the protecting group is a t-butyldimethylsilyl group, hydrogenation of the double bonds, and

- 14 -

conversion of the hydroxyl group to the bromide. These transformation are routine and well known to those skilled in the art.

Scheme 4: Method A

5

Target molecule (I) where R_2 is CH_2OH can be synthesized using the Method A described in Scheme 4 starting from commercially available 3-bromo-5-iodobenzoic acid. The first step consists in the selective displacement of the iodo group by the hydroxide ion using aqueous sodium hydroxide in the presence of catalytic amount of cuprous oxide at a temperature about 100°C (as described in *Organic Process Research & Development* **2002**, 6, 591-596). Reduction of the carboxylic group to benzylic alcohol **19** could be accomplished using a common reducing agent such as lithium aluminium hydride. The coupling reaction between **19** and **16**

- 15 -

is regioselective and only the most nucleophilic alcohol functionality reacts to form the desired intermediate **20**. The coupling is achieved in a mixture of acetone and N,N-dimethylformamide at a temperature about 75 °C in the presence of a base such as potassium carbonate or cesium carbonate. The ester hydrolysis can be

5 conveniently effected by treating the compound **20** with several equivalents of an alkali metal hydroxide, such as sodium hydroxide, potassium hydroxide or lithium hydroxide, in a suitable solvent such as a mixture of alcohol and water or tetrahydrofuran and water. The reaction can be carried out at a temperature ranging from 0 °C to 70 °C. The Suzuki coupling reaction is carried out between

10 aryl bromide **21** and various boronic acids to produce the derivatives of target molecule (**I**) where R₂ = CH₂OH. The conditions of this method are disclosed in many publications which have been reviewed by A. Suzuki in an article entitled "The Suzuki reaction with arylboron compounds in arene chemistry" in *Modern Arene Chemistry* **2002**, 53-106 and in a recent review: *Tetrahedron* **2002**, 58,


15 9633-9695. Generally Suzuki coupling reactions are carried out in the presence of a transition metal catalyst such as a palladium catalyst utilizing and a weak inorganic base. Among the preferred organic solvents are dimethoxyethane, ethanol and toluene. The weak inorganic base can be a carbonate or bicarbonate, such as potassium carbonate or cesium carbonate or phosphate such as

20 potassium phosphate. The reaction can also be done in anhydrous conditions or an aqueous solution of the base can be used. The source of palladium can be palladium(0) complex (e.g. tetrakis(triphenylphosphine)palladium(0)) or a compound which can be reduced *in situ* to give palladium(0) (for example palladium acetate(II) or bis(triphenylphosphine)palladium(II) chloride or

25 [1,1'-bis(diphenylphosphino) ferrocene] dichloropalladium(II).

Scheme 5: Method B

- 16 -




Target molecule (I) where R₂ equals CH₂OMe and CH₂OEt can be prepared as described in Scheme 5 starting from intermediate **20**. Using a strong base such as sodium hydride, alkylation of the benzylic alcohol could be accomplished in DMF

5 using alkyl halides such as methyl iodide and ethyl iodide. This reaction also known as the *Williamson reaction* is a very general method allowing the preparation of various unsymmetrical ethers. All the remaining steps leading to preparation of compounds of formula (I) have been described in Methods A.

- 17 -


Scheme 6: Method C



As shown in Scheme 6, 3-N,N-dimethylaminomethyl-5-aryl analogues and 3-cyanomethyl-5-aryl analogues having a structure such as (I) can be prepared using Method C. The desired methanesulfonate **25** can be obtained from 3-hydroxymethyl analogue **20** by treatment with methanesulfonyl chloride, triethylamine in dichloromethane and using a catalytic amount of N,N-dimethylaminopyridine. Then, the 3-N,N-dimethylaminomethyl-5-bromo analogue **26** was accomplished via a nucleophilic substitution reaction by treatment of **25** with N,N-dimethylamine in acetonitrile at room temperature. Alternatively, treatment of **25** with KCN in DMF at 60°C afforded the desired 3-cyanomethyl-5-bromo analogue **27**. The remaining two steps, Suzuki coupling with arylboronic acids and hydrolysis of the ethyl ester, were followed as described in Scheme 4 to obtain the final compounds such as (I).

- 18 -

Scheme 7: Method D




- As shown in Scheme 7, 3-alkoxy-5-aryl analogues having a structure such as (I) can be prepared using Method D. The desired diol **29** can be obtained from commercially available 3,5-dimethoxy bromobenzene **28** by treatment with any dealkylating agent, such as trimethylchlorosilane and sodium iodide or tribromoborane in a solvent like acetonitrile or dichloromethane at reflux or room temperature. Then, the monoalkylation of one hydroxy of the diol **29** was accomplished by using lithium hydride and one equivalent of **16** in DMSO to give the desired compound **30**. The second alkylation was accomplished by treatment of **30** with a lower alkyl iodide in the presence of a base such as potassium carbonate or cesium carbonate in a mixture of acetone and N,N-

- 19 -

dimethylformamide at a temperature of about 75 °C. The remaining two steps, Suzuki coupling with arylboronic acids and hydrolysis of the ethyl ester, were followed as described in the previous schemes to obtain the final compounds such as (I).

5

Scheme 8: Method E

The target 3-acyl-5-aryl analogues (I) can be synthesized according to the Scheme 8, Method E. The desired 3-acyl-5-bromophenol **33** can be prepared from


10 commercially available 3-bromo-5-iodobenzoic acid **17** in three steps. First, the 3-bromo-5-iodobenzoic acid **17** was converted to the corresponding acid chloride by treatment with oxalyl chloride in dichloromethane in the presence of a catalytic amount of N,N-dimethylformamide at 0 °C to room temperature. Then, the resulting acid chloride of **17** was treated with an *in situ* generated methylcopper reagent

15 from methylmagnesium chloride and a soluble copper cyanide and lithium chloride mixture in tetrahydrofuran at -70 to -30 °C for 15 h to obtain **32** (Paul Knochel and

- 20 -

Robert D. Singer, *Chem. Rev.* **1993**, 93, 2117-2188. Finally, the iodide of **32** was converted to the corresponding phenol by treatment of **32** with sodium hydroxide in the presence of a catalytic amount of copper(I) oxide in water and pyridine at 110 °C to afford the desired **33**. The remaining three steps, alkylation of phenol with **16**, Suzuki coupling with arylboronic acids and hydrolysis of ethyl ester, were followed as described in the previous schemes to obtain the final 3-acyl-5-aryl analogues (**I**).


Scheme 9: Method F



Compounds of Formula **40** can be conveniently prepared starting from 3,5-dibromobenzonitrile **35**. The hydroxyl group can be introduced via a nucleophilic displacement of a bromine by an alkoxide, preferably *para*-methoxybenzyloxy, followed by ether cleavage to give a phenol. For example, *para*-methoxybenzyl group can be removed by refluxing in glacial acetic acid. Aryl group can be

- 21 -

incorporated in a Suzuki coupling reaction between an appropriately substituted haloarene, preferably bromo- or iodoarene, and a boronic acid or a boronic acid equivalent (ester or anhydride) using a palladium catalyst. Nitrile group can be easily converted to ketone using a Grignard reagent and catalytic amount of cuprous bromide as described by Weiberth in *J. Org. Chem.* **1987**, 52, 3901-3904.

Scheme 10: Method G


- Compounds of Formula **47** can be prepared starting from 3,5-dimethoxybenzonitrile **42**. Nitrile group can be easily converted to ketone using a Grignard reagent and catalytic amount of cuprous bromide as described by Weiberth in *J. Org. Chem.* **1987**, 52, 3901-3904. The methoxy ether can be cleaved by methods known to those skilled in the art, preferably using boron tribromide yielding intermediate **44**. Dihydroxy derivative **44** can be converted to

- 22 -

monotriflate **46** in a two-step procedure: (i) conversion to the corresponding ditriflate **45** (Tf₂O, DIEA, CH₂Cl₂) and (ii) cleavage of one of the two sulfonyl groups by treatment with cesium carbonate in 1,2-dimethoxyethane as described by Hamura in *Helvetica Chimica Acta* **2002**, 85, 3589-3605. Aryl group can be

5 incorporated in a Suzuki coupling reaction between an appropriately substituted aromatic triflate and a boronic acid or a boronic acid equivalent using a palladium catalyst.

Scheme 11



- 10 Substituted phenyl boronic acids (**49**, R = H) and boronic esters such as 4,4,5,5-tetramethyl -[1,3,2]dioxaborolane (**49**, R = -(C(CH₃)₂)₂-) useful in the preparation of compounds of this invention may be commercially available or they can be made by reactions that are well known in the field of organic synthesis. Aryl boronic acids and aryl boronic esters are formed by treatment of aryl halides **48** with an
- 15 organometallic reagent such as *n*-butyl lithium followed by treatment with boron triisopropoxide or 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane followed by acidic work-up as is well known to those skilled in the art (For example in the literature, see *Org. Process Res. Dev.* **2004**, 8, 201).

Commercially available boronic acids used in this procedure are listed below. The

- 20 Available Chemicals Database (ACD) indicates the availability of greater than seven hundred commercially available aryl boronic acids. Some boronic acids useful for the preparation of compounds of the invention are listed below.

Table 1: Commercially available boronic acids

Boronic acid
3-CHLORO-PHENYLBORONIC ACID
PHENYLBORONIC ACID
2-CHLOROPHENYLBORONIC ACID
3-CHLORO-4-FLUOROPHENYLBORONIC ACID
3-METHYLPHENYLBORONIC ACID
1,4-BENZODIOXANEBORONIC ACID
3,4-DIFLUOROPHENYLBORONIC ACID
4-CHLOROPHENYLBORONIC ACID
4-CHLORO-3-FLUOROPHENYLBORONIC ACID
THIOPHENE-2-BORONIC ACID
2-FLUOROPHENYLBORONIC ACID
4-FLUORO-3-METHYLPHENYLBORONIC ACID
3-FLUOROPHENYLBORONIC ACID
THIOPHENE-3-BORONIC ACID
4-METHYL-3-THIOPHENE-BORONIC ACID
4-METHOXYPHENYLBORONIC ACID
4-ETHOXYPHENYLBORONIC ACID
3-FLUORO-4-METHYLPHENYLBORONIC ACID
4-ETHYLPHENYLBORONIC ACID
2,3-DIHYDROBENZOFURAN-5-BORONIC ACID
3-METHOXYPHENYLBORONIC ACID
2-TRIFLUOROMETHOXYPHENYLBORONIC ACID
4-TRIFLUOROMETHOXYPHENYLBORONIC ACID
4-FLUOROPHENYLBORONIC ACID
3-FLUORO-4-METHOXYPHENYLBORONIC ACID
3,5-DIFLUOROPHENYLBORONIC ACID
1-CYCLOHEXEN-1-YL-BORONIC ACID
5-BENZO[1,3]DIOXOLEBORONIC ACID
1H-PYRAZOLE-4-BORONIC ACID

- 24 -

1H-INDOLE-5-BORONIC ACID
4-PYRIDYL-BORONIC ACID
3-PYRIDYL-BORONIC ACID
4-METHANESULFONYLPHENYLBORONIC ACID
1-H-PYRAZOL-3-YLBORONIC ACID
4-FLUORO-3-HYDROXYPHENYLBORONIC ACID

Table 2: These boronic acids are also available from other suppliers that may not necessarily be listed in the ACD.

3-Fluoro-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)pyridine	ATLANTIC SCIENTIFIC CO., INC., JERSEY CITY, NJ,	791819-04-0
Quinoline-2-boronic acid	LANCASTER	745784-12-7
3-Chloro-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)pyridine	ATLANTIC SCIENTIFIC CO., INC., JERSEY CITY, NJ,	652148-93-1
6-Chloropyridine-2-boronic acid pinacol ester	INTERCHIM, MONTLUCON, FRANCE	652148-92-0
Boronic acid, (2-methyl-4-pyrimidinyl)-	CHEMSTEP, TALENCE, FRANCE	647853-31-4
Boronic acid, (3-methoxy-2-pyridinyl)-	CHEMSTEP, TALENCE, FRANCE	500707-34-6
Boronic acid, (6-methoxy-2-pyridinyl)-	CHEMSTEP, TALENCE, FRANCE	372963-51-4
Boronic acid, (6-methyl-2-pyridinyl)-	CHEMSTEP, TALENCE, FRANCE	372963-50-3
Boronic acid, (5-methyl-2-pyridinyl)-	CHEMSTEP, TALENCE, FRANCE	372963-49-0
Boronic acid, (4-methyl-2-	CHEMSTEP, TALENCE,	372963-48-9

- 25 -

pyridinyl)-	FRANCE	
Boronic acid, 2-pyridinyl-	CHEMSTEP, TALENCE, FRANCE	197958-29-5

In the practice of the method of the present invention, an effective amount of any one of the compounds of this invention or a combination of any of the compounds of this invention or a pharmaceutically acceptable salt thereof, is administered via any of the usual and acceptable methods known in the art, either singly or in

5 combination. The compounds or compositions can thus be administered orally (e.g., buccal cavity), sublingually, parenterally (e.g., intramuscularly, intravenously, or subcutaneously), rectally (e.g., by suppositories or washings), transdermally (e.g., skin electroporation) or by inhalation (e.g., by aerosol), and in the form of solid, liquid or gaseous dosages, including tablets and suspensions. The administration
10 can be conducted in a single unit dosage form with continuous therapy or in a single dose therapy ad lithium. The therapeutic composition can also be in the form of an oil emulsion or dispersion in conjunction with a lipophilic salt such as pamoic acid, or in the form of a biodegradable sustained-release composition for subcutaneous or intramuscular administration.

15

Useful pharmaceutical carriers for the preparation of the compositions hereof, can be solids, liquids or gases; thus, the compositions can take the form of tablets, pills, capsules, suppositories, powders, enterically coated or other protected formulations (e.g. binding on ion-exchange resins or packaging in lipid-protein

20 vesicles), sustained release formulations, solutions, suspensions, elixirs, aerosols, and the like. The carrier can be selected from the various oils including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, and the like. Water, saline, aqueous dextrose, and glycols are preferred liquid carriers, particularly (when isotonic with the blood) for injectable
25 solutions. For example, formulations for intravenous administration comprise sterile aqueous solutions of the active ingredient(s) which are prepared by dissolving solid active ingredient(s) in water to produce an aqueous solution, and rendering the solution sterile. Suitable pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, gelatin, malt, rice, flour, chalk, silica, magnesium stearate, sodium
30 stearate, glycerol monostearate, sodium chloride, dried skim milk, glycerol,

- 26 -

propylene glycol, water, ethanol, and the like. The compositions may be subjected to conventional pharmaceutical additives such as preservatives, stabilizing agents, wetting or emulsifying agents, salts for adjusting osmotic pressure, buffers and the like. Suitable pharmaceutical carriers and their formulation are described in

- 5 Remington's Pharmaceutical Sciences by E. W. Martin. Such compositions will, in any event, contain an effective amount of the active compound together with a suitable carrier so as to prepare the proper dosage form for proper administration to the recipient.
- 10 The dose of a compound of the present invention depends on a number of factors, such as, for example, the manner of administration, the age and the body weight of the subject, and the condition of the subject to be treated, and ultimately will be decided by the attending physician or veterinarian. Such an amount of the active compound as determined by the attending physician or veterinarian is referred to
- 15 herein, and in the claims, as an "effective amount". For example, the dose of a compound of the present invention is typically in the range of about 1 to about 1000 mg per day.

- The invention will now be further described in the Examples below, which are
- 20 intended as an illustration only and do not limit the scope of the invention.

EXAMPLES

- Reagents were purchased from Aldrich, Sigma, Maybridge, Advanced ChemTech, and Lancaster or other suppliers as indicated below and used without further
- 25 purification. Reactions using microwave irradiation for heating were conducted using either a Personal Chemistry Emrys Optimizer System or a CEM Discovery System. The purification of multi-milligram to multi-gram scale was conducted by methods known to those skilled in the art such as elution of silica gel flash column; preparative flash column purifications were also effected in some cases by
- 30 use of disposal pre-packed multigram silica gel columns (RediSep) eluted with a CombiFlash system. Biotage™ and ISCO™ are also flash column instruments that may have been used in this invention for purification of intermediates.

- 27 -

For the purpose of judging compound identity and purity, LC/MS (liquid chromatography/mass spectroscopy) spectra were recorded using the following system. For measurement of mass spectra, the system consists of a Micromass Platform II spectrometer: ES Ionization in positive mode (mass range: 150 -1200 amu). The simultaneous chromatographic separation was achieved with the following HPLC system: ES Industries Chromegabond WR C-18 3u 120Å (3.2 x 30mm) column cartridge; Mobile Phase A: Water (0.02% TFA) and Phase B: Acetonitrile (0.02% TFA); gradient 10% B to 90% B in 3 minutes; equilibration time of 1 minute; flow rate of 2 mL/minute. In some cases, ammonium acetate at 20 millimolar concentration was used as a modifier for effective ionization during preparative HPLC. In such cases, the ammonium salt was isolated.

For some separations, the use of super critical fluid chromatography may also be useful. Super critical fluid chromatography separations were performed using a Mettler-Toledo Minigram system with the following typical conditions: 100 bar, 30 °C, 2.0 mL/min eluting a 12 mm AD column with 40% MeOH in super critical fluid CO₂. In the case of analytes with basic amino groups, 0.2% isopropyl amine was added to the methanol modifier.

Many compounds of Formula 1 were also purified by reversed phased HPLC, using methods well known to those skilled in the art. In some cases, preparative HPLC purification was conducted using PE Sciex 150 EX Mass Spec controlling a Gilson 215 collector attached to a Shimadzu preparative HPLC system and a Leap autoinjector. Compounds were collected from the elution stream using LC/MS detection in the positive ion detection: The elution of compounds from C-18 columns (2.0 X 10 cm eluting at 20 ml/min) was effected using appropriate linear gradation mode over 10 minutes of Solvent (A) 0.05% TFA/H₂O and Solvent (B) 0.035% TFA/acetonitrile. For injection on to HPLC systems, the crude samples were dissolved in mixtures of methanol, acetonitrile and DMSO

30

H-Cube™ (produced by Thales Nanotechnology) is a continuous-flow hydrogenation reactor equipped with *in situ* hydrogen generation and a disposable catalyst cartridge CatCart™. The reaction mixture can be heated and pressurized

- 28 -

up to 100°C and 100 bar (1450 psi) respectively. Reaction scale can be varied from 10 mg to 100 g.

Compounds were characterized either by ¹H-NMR using a Varian Inova 400 MHz
5 NMR Spectrometer or a Varian Mercury 300 MHz NMR Spectrometer as well as by high resolution mass spectrometry using a Bruker Apex-II high-resolution 4.7T FT-Mass Spectrometer.

LIST OF ABBREVIATIONS:

10	AIBN	2,2'-azobisisobutyronitrile,
	Bu	butyl,
	DCE	1,2-dichloroethane,
	DCM	dichloromethane,
	DIPEA	diisopropylethylamine,
15	DMF	N,N-dimethylformamide,
	DMSO	dimethylsulfoxide,
	EtOAc	ethyl acetate,
	EtOH	ethyl alcohol,
	FCC	flash column chromatography,
20	h	hour,
	HPLC	high pressure liquid chromatography,
	HRMS	high resolution mass spectra,
	LRMS	low resolution mass spectra,
	LC	liquid chromatography,
25	L-Pro	L-proline,
	MCPBA	meta-chloroperoxybenzoic acid,
	MeOH	methyl alcohol,
	MW	microwave,
	NIS	N-iodosuccinimide,
30	NBS	N-bromosuccinimide,
	NMP	1-methyl-2-pyrrolidinone,
	PdCl ₂ (dppf)	[1,1'-Bis(diphenylphosphino)ferrocene] dichloropalladium(II),
	PG	protecting group,


- 29 -

PyBroP	bromo-tris-pyrrolidino-phosphonium hexafluorophosphate,
rt	room temperature,
TBAF	tetrabutylammonium fluoride,
TBDMS	tert-butyl-dimethylsilyl,
5 TBTU	2-(1 <i>H</i> -benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate,
TMS	trimethylsilyl,
TMSSMe	(methylthio)trimethylsilane,
TEA	triethylamine,
10 TEMPO	2,2,6,6-tetramethylpiperidine-1-oxyl,
TFA	trifluoroacetic acid,
THF	tetrahydrofuran.

I. PREPARATION OF PREFERRED INTERMEDIATES


15 Preparation of 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester

Preparation of 4-(2,3-dimethyl-phenoxy)-butyric acid ethyl ester




- 20 To a solution of 2,3-dimethylphenol (25 g, 204 mmol) in DMSO (205 mL) was added 4-bromo-butyric acid ethyl ester (40.96 g, 210 mmol) and lithium hydride (2.0 g, 250 mmol) at room temperature. The resulting light brown solution was stirred for 2 days. Then, the reaction mixture was cooled to 0 °C and water (200 mL) was added slowly. The organic compound was extracted into hexanes (2 x
- 25 200 mL). The combined organic extracts were washed with brine solution (150 mL) and the organic solution was dried over anhydrous magnesium sulfate. Filtration of the drying agent and the removal of the solvent gave light brown oil. The crude mixture was purified by using a Biotage™ (40L) column chromatography eluting with 5% ethyl acetate in hexanes to isolate 4-(2,3-dimethyl-phenoxy)-butyric acid
- 30 ethyl ester (45.32 g, 94%) as a colorless oil: ES(+)-HRMS m/e calculated for C₁₄H₂₀O₃ (M⁺)⁺ 236.1412, found 236.1419.

Preparation of 4-(2-formyl-3-methyl-phenoxy)-butyric acid ethyl ester



- A mixture of copper(II)sulfate pentahydrate (21.98 g, 88.06 mmol) and potassium persulfate (71.42 g, 264 mmol) in water (396 mL) was heated to 63-65 °C to obtain a blue colored solution. Then, a solution of 4-(2,3-dimethyl-phenoxy)-butyric acid ethyl ester (20.81 g, 88.06 mmol) in acetonitrile (220 mL) was added at the above temperature. The resulting light green solution was refluxed for 40 minutes. Then, the reaction mixture was cooled to ~5 °C in order to precipitate most of the inorganic solids. The resulting solids were collected by filtration and the solid cake was washed with dichloromethane (1.0 L). The two layers of filtrate were separated and the aqueous layer was extracted with dichloromethane (200 mL). The combined organic extracts were washed with brine solution (150 mL) and the organic solution was dried over anhydrous magnesium sulfate. Filtration of the drying agent and the removal of the solvent gave a brown oil. The crude mixture was purified by using a Biotage™ (40L) column chromatography eluting with 5-10% ethyl acetate in hexanes to obtain 4-(2-formyl-3-methyl-phenoxy)-butyric acid ethyl ester (45.32 g, 94%) as a colorless oil: EI(+)-HRMS m/e calculated for C₁₄H₁₈O₄ (M⁺)⁺ 250.1205, found 250.1202.
- Preparation of 4-[2-((E)-2-ethoxycarbonyl-vinyl)-3-methyl-phenoxy]-butyric acid ethyl ester.




- Sodium metal spheres (1.6 g, 69.6 mmol) were added to ethanol (100 mL) with stirring at room temperature under nitrogen atmosphere over 15 min. An exothermic reaction occurred and the mixture was stirred for another 15 min to form sodium ethoxide. After cooling to room temperature, triethylphosphonoacetate (14.7 mL, 73.4 mmol) and 4-(2-formyl-3-methyl-phenoxy)-butyric acid ethyl ester (13.25 g, 52.9 mmol) were added sequentially. During the addition of 4-(2-formyl-3-

- 31 -


methyl-phenoxy)-butyric acid ethyl ester, the color of the solution turned brown and the temperature increased to ~55 °C. The resulting brown solution was stirred for 2 days at room temperature. Then, the reaction mixture was diluted with water (150 mL) and stirred for 1 h. Then, the organic compound was extracted into
5 hexanes (3 x 100 mL). The combined organic extracts were washed with brine solution (150 mL) and the organic solution was dried over anhydrous magnesium sulfate. Filtration of the drying agent and the removal of the solvent gave a light yellow oil. The crude oil was dissolved in hexanes (~50 mL) and treated with charcoal and heated gently with a heat gun. After cooling to room temperature, the
10 charcoal was filtered-off and the filtrate was removed under vacuum to give 4-[2-((E)-2-ethoxycarbonyl-vinyl)-3-methyl-phenoxy]-butyric acid ethyl ester (13.25 g, 78%) as colorless oil: EI(+)-HRMS m/e calculated for C₁₈H₂₄O₅ (M+)⁺ 320.1624, found 320.1626.

Preparation of 4-[3-bromomethyl-2-((E)-2-ethoxycarbonyl-vinyl)-phenoxy]-butyric
15 acid ethyl ester.




To a solution of 4-[2-((E)-2-ethoxycarbonyl-vinyl)-3-methyl-phenoxy]-butyric acid ethyl ester (8.0 g, 25.0 mmol) in chlorobenzene (190 mL) were added N-bromosuccinimide (6.67 g, 37.5 mmol) and 2,2'-azobisisobutyronitrile (AIBN) (591
20 mg, 3.6 mmol) at room temperature. Then, the solution was heated to 85 °C and stirred for 1 h. Then, the reaction mixture was cooled to room temperature and diluted with water (100 mL). Then, the organic compound was extracted into hexanes (3 x 100 mL). The combined organic extracts were washed with brine solution (150 mL) and the organic solution was dried over anhydrous magnesium
25 sulfate. Filtration of the drying agent and the removal of the solvent gave a crude oil. The crude oil was purified by using a Biotage (40L) column eluting with 15-25% ethyl acetate in hexanes to isolate 4-[3-bromomethyl-2-((E)-2-ethoxycarbonyl-vinyl)-phenoxy]-butyric acid ethyl ester (7.11 g, 71%) as a low melting solid: ES(+)-HRMS m/e calculated for C₁₈H₂₃BrO₅ (M+Na)⁺ 421.0621, found 421.0621.
30 Preparation of 5-(*tert*-butyl-dimethyl-silanyloxy)-pentanal .

- 32 -



To a solution of 5-(*tert*-butyl-dimethyl-silyloxy)-pentanol (16.8 mmol, 3.66 g) in dichloromethane (30 mL) were added water (5.6 mL), potassium bromide (1.7 mmol, 202 mg), *n*-tetrabutylammonium hydrogensulfate (0.84 mmol, 290 mg), and TEMPO (30 mg) at room temperature. The resulting light brown solution was cooled to ~5 °C and a solution of sodium hypochlorite (19.3 mmol, 30 mL, 5%) was added dropwise at this temperature. After addition of half of the sodium hypochlorite solution, solid potassium carbonate (300 mg) was added to maintain the reaction mixture basic. Then, the remaining sodium hypochlorite solution was added at 5-10 °C. By this point, a precipitate had formed and the reaction mixture was stirred for another 1 h at ~10-15 °C. Then, water (100 mL) was added and the resulting solution was extracted into diethyl ether (2x100 mL). The combined organic extracts were washed with brine solution (150 mL) and the organic layer was dried over anhydrous magnesium sulfate. Filtration of the drying agent and the removal of the solvent gave 5-(*tert*-butyl-dimethyl-silyloxy)-pentanal (3.32 g, 91%) as a light brown oil: ES(+)-HRMS *m/e* calculated for C₁₁H₂₄O₂Si (M+H)⁺ 217.1619, found 217.1619.

Preparation of 4-[3-[6-(*tert*-butyl-dimethyl-silyloxy)-hex-1-enyl]-2-((*E*)-2-ethoxycarbonyl-vinyl)-phenoxy]-butyric acid ethyl ester.



A solution of 4-[3-bromomethyl-2-((*E*)-2-ethoxycarbonyl-vinyl)-phenoxy]-butyric acid ethyl ester (2.0 mmol, 798 mg) and triphenylphosphine (2.2 mmol, 577 mg) in acetonitrile (12 mL) was heated to reflux for 1 h under nitrogen atmosphere. Then, it was cooled to room temperature and a solution of 5-(*tert*-butyl-dimethyl-silyloxy)-pentanal (2.8 mmol, 606 mg) in 1,2-epoxybutane (22 mL) was added at room temperature and the mixture was again heated to reflux for 15 h. During this period, the mixture first turned to a brick red color and at the end of the reaction it had become a pale yellow solution. Then, the reaction mixture was cooled to room temperature and the solvent was removed under vacuum. The residue was

- 33 -

dissolved in a solution of ethyl acetate and hexanes (1:3, 150 mL) and the resulting cloudy solution was washed with a mixture of methanol and water (2:1, 225 mL).

The aqueous layer was extracted one more time with ethyl acetate and hexanes (1:3, 50 mL). The combined organic extracts were washed with brine solution (150


5 mL) and the organic solution was dried over anhydrous magnesium sulfate.

Filtration of the drying agent and the removal of the solvent gave light brown oil.

The crude mixture was purified by using a Biotage™ (40L) column chromatograph eluting with 5 and 15% ethyl acetate in hexanes to obtain the desired 4-[3-[6-(*tert*-butyl-dimethyl-silanyloxy)-hex-1-enyl]-2-((*E*)-2-ethoxycarbonyl-vinyl)-phenoxy]-

10 butyric acid ethyl ester (760 mg, 74%) as a colorless oil: ES(+)-HRMS *m/e* calculated for C₂₉H₄₆O₆Si (M+Na)⁺ 541.2956, found 541.2953.

7) Preparation of 4-[3-[6-(*tert*-butyl-dimethyl-silanyloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester.




15 To a solution of 4-[3-[6-(*tert*-butyl-dimethyl-silanyloxy)-hex-1-enyl]-2-((*E*)-2-ethoxycarbonyl-vinyl)-phenoxy]-butyric acid ethyl ester (0.977 mmol, 507 mg) in ethyl acetate (10 mL) was added 10% palladium on carbon (350 mg) at room temperature. The resulting black mixture was stirred in the presence of atmospheric hydrogen gas in a balloon for 36 h at room temperature. Then, the

20 catalyst was removed by filtration using a filter paper and the residue was washed with hot ethyl acetate (~60 mL). The filtrate was concentrated *in vacuo* and the resulting residue was dried under high vacuum to obtain 4-[3-[6-(*tert*-butyl-dimethyl-silanyloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (438 mg, 86%) as a colorless oil: ES(+)-HRMS *m/e* calculated for C₂₉H₅₀O₆Si (M+Na)⁺ 545.3269, found 545.3267.


8) Preparation of 4-[2-(2-ethoxycarbonyl-ethyl)-3-(6-hydroxy-hexyl)-phenoxy]-butyric acid ethyl ester.

- 34 -



To a solution of 4-[3-[6-(*tert*-butyl-dimethyl-silanyloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (0.837 mmol, 438 mg) in THF (12 mL) was added a solution of *n*-tetrabutyl ammonium fluoride (1.25 mmol, 1.25 mL, 1.0M) in THF at 0 °C. Then, the resulting colorless solution was allowed to warm to room temperature in 2 h and the mixture was stirred for another 2 h at room temperature before being diluted with water (~50 mL). The organic compound was extracted into ethyl acetate (2x50 mL) and the combined extracts were washed with brine solution (100 mL). The organic solution was dried over anhydrous magnesium sulfate and the filtrate was removed under vacuum after filtration of the drying agent. The crude residue was dried further under high vacuum and the desired 4-[2-(2-ethoxycarbonyl-ethyl)-3-(6-hydroxy-hexyl)-phenoxy]-butyric acid ethyl ester (342 mg, 99%) was isolated as a colorless oil: ES(+)-HRMS *m/e* calculated for C₂₃H₃₆O₆ (M+Na)⁺ 431.2404, found 431.2404.

9) Preparation of 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester.



To a solution of 4-[2-(2-ethoxycarbonyl-ethyl)-3-(6-hydroxy-hexyl)-phenoxy]-butyric acid ethyl ester (0.85 mmol, 349 mg) and carbon tetrabromide (1.26 mmol, 423 mg) in dichloromethane (10 mL) was added triphenylphosphine (1.07 mmol, 281 mg) at ~0 °C. The resulting colorless solution was stirred for 3 h at 5-10 °C. Then, the solvent was removed under vacuum and the crude was tried to dissolve in a mixture of ethyl acetate and hexanes (1:3, 50 mL). As a result, a cloudy solution containing some precipitate was formed and the cloudy solution was transferred into a separatory funnel and was washed with a mixture of methanol and water (2:1, 150 mL). The aqueous layer was extracted one more time with ethyl acetate and hexanes (1:3, 50 mL). The combined organic extracts were washed with brine


- 35 -

solution (100 mL) and the organic solution was dried over anhydrous magnesium sulfate. Filtration of the drying agent and the removal of the solvent gave a colorless oil which was purified by using a Biotage™ (40M) column chromatography eluting with 10% ethyl acetate in hexanes to obtain the desired 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (350 mg, 87.5%) as a colorless oil: ES(+)-HRMS m/e calculated for $C_{23}H_{35}BrO_5$ ($M+Na$)⁺ 493.1560, found 493.1560.

II. PREPARATION OF PREFERRED COMPOUNDS


Method A

Step 1: 3-Bromo-5-hydroxy-benzoic acid



To a solution of NaOH (12.2 g; 305.8 mmol) in 300 mL of water was added 3-bromo-5-iodo-benzoic acid (20g, 61.2 mmol) and Cu_2O (866 mg, 6.1 mmol). The reaction mixture was heated at 100 °C for 24 h. After complete consumption of the starting material, the reaction mixture was cooled to room temperature and filtered through Celite™. The filtrate was then acidified with 10% aq. HCl and extracted into ethyl acetate. The organic extract was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford the title compound as a tan solid (12.3g, 93% yield). HR-ES (-) calcd for $C_7H_5O_3Br$ [$M-H$]⁻ 214.9349, observed 214.9350

Step 2: 3-Bromo-5-hydroxymethyl-phenol




To a solution of 3-bromo-5-hydroxy-benzoic acid (5.0 g, 23.4 mmol) in 40 mL of THF was added a solution of 1M $LiAlH_4$ (30 mL), and the resulting solution was heated to reflux for 2 h. After cooling to room temperature, the reaction mixture was poured on ice and acidified with 10% aq. HCl. The solution was then filtered under vacuum and the filtrate was extracted with EtOAc. The organic extract was dried

- 36 -

over anhydrous sodium sulfate and concentrated under reduced pressure. The crude material was purified by column chromatography (Isco™ 120 g) using 50% ethyl acetate/hexanes as eluting solvents to afford the title compound as a white solid (1.7 g, 36% yield). HR-ES(-) calcd for C₇H₇O₂Br [M-H]⁻ 200.9556, observed

5 200.9557.

Step 3: 4-[3-[6-(3-Bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



10

To a solution of 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (4.2 g, 8.9 mmol), 3-Bromo-5-hydroxymethyl-phenol (1.5 g, 7.4 mmol) in *N,N*-dimethylformamide (40 mL) and acetone (80 mL) was added potassium carbonate (10.2 g, 74.2 mmol) at room temperature. The resulting

15 suspension was heated to 70 ° C for 24 h. Then, the reaction mixture was cooled to room temperature and diluted with water and 10% aq. HCl. The organic compound was extracted into ethyl acetate and the combined organic extracts were washed with water and brine solution. The organic layers were dried over anhydrous sodium sulfate and the solvent was removed under reduced pressure.


20

The crude material was purified by column chromatography (Isco™ 120 g) with 0-50% ethyl acetate/hexanes as eluting solvents to afford the title compound (4.3 g, 99%) as a colorless oil. HR-ES(+) calcd for C₃₀H₄₁O₇Br [M+Na]⁺ 615.1928, observed 615.1926.

25

Step 4 : 4-[3-[6-(3-Bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

- 37 -



To a solution of 4-[3-[6-(3-bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (1.0 g, 1.68 mmol) in hot ethanol (40 mL) was added aqueous 2.0 N sodium hydroxide (10 mL). The resulting solution was heated to 60 °C and stirred for 4 h. Then, the reaction mixture was cooled down and diluted with EtOAc, washed with 10% aq. HCl and brine. The combined ethyl acetate extracts were dried over anhydrous sodium sulfate and the solvent was removed under reduced pressure to afford 805 mg (89%) of title compound as a light yellow oil. HR-ES(+) calcd for C₂₆H₃₃O₇Br [M+Na]⁺ 559.1302, observed 559.1299.


Step 5: General procedure for Suzuki coupling

To a solution of 4-[3-[6-(3-bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (100 mg, 0.19 mmol), potassium carbonate (105 mg, 0.76 mmol), boronic acid (0.38 mmol) in EtOH (4 mL)/H₂O (1 mL) was added Pd(PPh₃)₄ (11 mg, 5 mol%). The reaction mixture was heated at 78 °C for 12 h, cooled down to room temperature then diluted with EtOAc. The resulting solution was washed with 10% aq. HCl and brine. The organic layer was then dried over anhydrous sodium sulfate and concentrated under vacuo to afford the crude material. The desired products were isolated by preparative HPLC.

- 38 -

Example 1:


4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



- 5 The title compound was prepared by following the general procedure for Suzuki coupling described above in Step 5 with 3,4-methylenedioxyphenyl boronic acid. HR-ES(+) calcd for $C_{33}H_{38}O_9$ ($M+Na$)⁺ 601.2408, found 601.2407.

Example 2:


- 10 **4-[2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid**



- The title compound was prepared by following the general procedure for Suzuki coupling described above in Step 5 with phenylboronic acid. HR-ES(+) calcd for $C_{32}H_{38}O_7$ ($M+H$)⁺ 535.2689, found 535.2689.
- 15

Example 3:

4-[2-(2-Carboxy-ethyl)-3-[6-(3-hydroxymethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid




- 39 -

The title compound was prepared by following the general procedure for Suzuki coupling described above in Step 5 with 3-thiopheneboronic acid. HR-ES(+) calcd for $C_{30}H_{36}O_7S$ ($M+Na$)⁺ 563.2074, found 563.2076.

5

Example 4:

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-fluoro-5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




The title compound was prepared by following the general procedure for Suzuki coupling described above in Step 5 with 4-fluorophenylboronic acid. HR-ES(+) calcd for $C_{32}H_{37}O_7F$ ($M+Na$)⁺ 575.2415, found 575.2413.

10

Example 5:

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid

15




The title compound was prepared by following the general procedure for Suzuki coupling described above in Step 5 with 4-chlororophenylboronic acid. HR-ES(+) calcd for $C_{32}H_{37}O_7Cl$ ($M+Na$)⁺ 591.2120, found 591.2121.

20

Example 6:


4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-hydroxymethyl -phenoxy]-hexyl}-phenoxy)-butyric acid

- 40 -




The title compound was prepared by following the general procedure for Suzuki coupling described above in Step 5 with 1,4-benzodioxane-6-boronic acid. HR-ES(+) calcd for $C_{34}H_{40}O_9Cl$ ($M+Na$)⁺ 615.2564, found 615.2563.

5

Example 7:**4-[2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid**

10 **Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester**



To a mixture of 4-[3-[6-(3-bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.0 g, 3.34 mmol), 4-methanesulfonylphenylboronic acid (1.35 g, 6.74 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (369.8 mg, 0.51 mmol), and cesium carbonate (2.22 g, 6.74 mmol) was added dimethoxyethane (100 mL) at room temperature under nitrogen atmosphere. The resulting brown reaction mixture was heated to 96 °C and stirred for 15 h at which time the TLC analysis of the reaction mixture indicated the absence of starting material. Then, the reaction

15


20

- 41 -


mixture was cooled to room temperature and diluted with water and ethyl acetate. The two layers were separated and the aqueous layer was extracted with ethyl acetate and the combined organic extracts were washed with water and brine solution. The organic layer was dried over anhydrous magnesium sulfate and
5 filtration of the drying agent and removal of the solvent under vacuum gave the colored residue which was purified by using an ISCO 120 g column to obtain 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (1.23 g, 55%) as a light brown oil: ES(+)-HRMS m/e calcd for $C_{37}H_{48}O_9S$ ($M+Na$)⁺ 691.2911, found 691.2913.

10

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid




To a solution of the 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (250 mg, 0.37 mmol) in THF (5 mL) and ethanol (5 mL) was added aqueous 1.0 N sodium hydroxide (4 mL) at room temperature. The resulting suspension was stirred for 5 h at room temperature at which time TLC analysis of the mixture indicated the absence of starting material. Then, the reaction mixture was concentrated and the
15 residue was diluted with water (20 mL) and extracted with diethyl ether (30 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N hydrochloric acid and the precipitated white organic compound was extracted into ethyl acetate (2 x 30 mL). The combined ethyl acetate extracts were washed with brine solution (50 mL) and the organic layers were dried over anhydrous
20 magnesium sulfate. Filtration of the drying agent and removal of the solvent afforded 4-[2-(2-carboxy-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (65 mg, 37%) as a white solid: ES(+)-HRMS m/e calcd for $C_{33}H_{40}O_9S$ ($M+Na$)⁺ 635.2285, found 635.2284.

Method B**Step 1: 4-[3-[6-(3-Bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester**

5 To a solution of 4-[3-[6-(3-bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (1.25 g, 2.1 mmol) in DMF (25 mL) were added sodium hydride (211 mg, 5.26 mmol) and iodomethane (747 mg, 5.26 mmol) at room temperature. The resulting suspension was stirred for 15 h
10 and then the excess sodium hydride was quenched by slow addition of water (5 mL). The mixture was diluted with 1.0N hydrochloric acid (50 mL) and the organic compound was extracted into ethyl acetate (2 x 50 mL). The combined extracts were washed with brine solution (100 mL) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and removal of the solvent under vacuum
15 afforded the crude residue which was purified by using an ISCO 40 g column, eluting with 0-30% ethyl acetate in hexanes to obtain 4-[3-[6-(3-bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (1.03 g, 81%) as a colorless viscous oil: EI(+)-HRMS m/e calcd for $C_{31}H_{43}BrO_7$ (M+Na)⁺ 629.2084, found 629.2085.

20

Step 2 : 4-[3-[6-(3-Bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

25 To a solution of 4-[3-[6-(3-Bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (500 mg, 0.82 mmol) in hot

- 43 -

ethanol (20 mL) was added aqueous 2.0 N sodium hydroxide (5 mL). The resulting solution was heated to 60 °C and stirred for 4 h. Then, the reaction mixture was cooled down and diluted with EtOAc, washed with 10% aq. HCl and brine. The combined ethyl acetate extracts were dried over anhydrous sodium sulfate and the solvent was removed under reduced pressure to afford 400 mg (89%) of title compound as a light yellow oil. HR-ES(+) calcd for C₂₇H₃₅O₇Br [M+Na]⁺ 573.1458, observed 573.1459.

Step 3: General procedure for Suzuki coupling

10 Method A:

To a solution of 4-[3-[6-(3-Bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (100 mg, 0.19 mmol), potassium carbonate (105 mg, 0.76 mmol), boronic acid (0.38 mmol) in EtOH (4 mL)/H₂O (1 mL) was added Pd(PPh₃)₄ (11 mg, 5 mol%). The reaction mixture was heated at 78 °C for 12 h, cooled down to room temperature then diluted with EtOAc. The resulting solution was washed with 10% aq. HCl and brine. The organic layer was then dried over anhydrous sodium sulfate and concentrated under vacuo to afford the crude material. The desired products were isolated by preparative HPLC.

20 Method B:


To a mixture of 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester, boronic acid [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), and cesium carbonate was added dimethoxyethane (10 mL) at room temperature under nitrogen atmosphere. The resulting brown reaction mixture was heated to 96 °C and stirred for 15 h at which time the TLC analysis of the reaction mixture indicated the absence of starting material. Then, the reaction mixture was cooled to room temperature and diluted with water (50 mL) and ethyl acetate (50 mL). The two layers were separated and the aqueous layer was extracted with ethyl acetate (50 mL) and the combined organic extracts were washed with water (50 mL) and brine solution (50 mL). The organic layer was dried over anhydrous magnesium sulfate and filtration of the drying agent and removal of the solvent under vacuum gave the colored

- 44 -

residue which was purified by using an ISCO 40 column. eluting with 0-25% ethyl acetate in hexanes to obtain the desired product.

Example 8:


- 5 **4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid**



- The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 3,4-methylenedioxyphenyl
10 boronic acid. HR-ES(+) calcd for $C_{34}H_{40}O_9$ ($M+Na$)⁺ 615.2564, found 615.2564.

Example 9:

- 4-[2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid**




- 15 The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with phenylboronic acid. HR-ES(+) calcd for $C_{33}H_{40}O_7$ ($M+Na$)⁺ 571.2666, found 571.2665.

- 20 **Example 10:**

- 4-[2-(2-Carboxy-ethyl)-3-[6-(4'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid**

- 45 -




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 4-fluorophenylboronic acid. HR-ES(+) calcd for $C_{33}H_{39}O_7F$ ($M+Na$)⁺ 589.2572, found 589.2574.

5

Example 11:


4-(2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy)-butyric acid



- 10 The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 4-chlorophenylboronic acid. HR-ES(+) calcd for $C_{33}H_{39}O_7Cl$ ($M+Na$)⁺ 605.2276, found 605.2279.

Example 12:

- 15 **4-(2-(2-Carboxy-ethyl)-3-[6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-methoxy methyl-phenoxy]-hexyl]-phenoxy)-butyric acid**




- 46 -

The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 1,4-benzodioxane-6-boronic acid. HR-ES(+) calcd for $C_{35}H_{42}O_9$ ($M+Na$)⁺ 629.2721, found 629.2719.

5

Example 13:


4-(2-(2-Carboxy-ethyl)-3-{6-[3-methoxymethyl-5-(4-methyl-thiophen-3-yl)-phenoxy]-hexyl}-phenoxy)-butyric acid



The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 4-methyl-3-thiopheneboronic acid. HR-ES(+) calcd for $C_{32}H_{40}O_7S$ ($M+Na$)⁺ 591.2387, found 591.2387.

Example 14:

4-{2-(2-Carboxy-ethyl)-3-[6-(3'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 3-fluorophenylboronic acid. HR-ES(+) calcd for $C_{33}H_{39}O_7F$ ($M+Na$)⁺ 589.2572, found 589.2571.

Example 15:

4-{2-(2-Carboxy-ethyl)-3-[6-(2'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid

- 47 -




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 2-fluorophenylboronic acid. HR-ES(+) calcd for $C_{33}H_{39}O_7F$ ($M+Na$)⁺ 589.2572, found 589.2573.

5

Example 16:


4-{2-(2-Carboxy-ethyl)-3-[6-(3',5'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid



- 10 The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 3,5-difluorophenylboronic acid. HR-ES(+) calcd for $C_{33}H_{38}O_7F_2$ ($M+Na$)⁺ 607.2478, found 607.2474.

Example 17:

- 15 **4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 4-

- 48 -

trifluoromethylphenylboronic acid. HR-ES(+) calcd for $C_{34}H_{39}O_7F_3$ (M+Na)⁺ 639.2540, found 639.2537.

Example 18:


- 5 **4-{2-(2-Carboxy-ethyl)-3-[6-(3'-chloro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**



- The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 3-chlorophenylboronic acid. HR-ES(+) calcd for $C_{33}H_{39}O_7Cl$ (M+Na)⁺ 605.2276, found 605.2279.
- 10

Example 19:

- 4-{2-(2-Carboxy-ethyl)-3-[6-(3'-methoxy-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**




- The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 3-methoxyphenylboronic acid. HR-ES(+) calcd for $C_{34}H_{42}O_8$ (M+Na)⁺ 601.2772, found 601.2774.
- 15

- 20 **Example 20:**

- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-4'-trifluoromethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**

- 49 -




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 3 using method A with 4-

trifluoromethoxyphenylboronic acid. HR-ES(+) calcd for $C_{34}H_{39}O_8F_3$ ($M+Na$)⁺

5 655.2489, found 655.2489.

Example 21:

4-[2-(2-Carboxy-ethyl)-3-[6-(3',4'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid



10


A similar procedure as described in step 3, method B was used, starting from 4-[3-[6-(3-bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (184 mg, 0.33 mmol), 3,4-difluorophenylboronic acid (106 mg, 0.66 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (44 mg, 0.06 mmol), and cesium carbonate (434 mg, 1.33 mmol) to obtain 4-[2-(2-carboxy-ethyl)-3-[6-(3',4'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (180 mg, 93%) as a light yellow paste: ES(+)-HRMS m/e calcd for $C_{33}H_{38}F_2O_7$ ($M+Na$)⁺ 607.2478, found 607.2478.

20

Example 22:

4-[2-(2-Carboxy-ethyl)-3-[6-(3-methoxymethyl-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid

- 50 -




A similar procedure as described in step 3, method B was used, starting from 4-[3-[6-(3-bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (209 mg, 0.38 mmol), 5-pyrimidinylboronic acid (94 mg, 0.76 mmol),
 5 [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (51 mg, 0.07 mmol), and cesium carbonate (492 mg, 1.51 mmol) to obtain 4-[2-(2-carboxy-ethyl)-3-[6-(3-methoxymethyl-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid (146 mg, 70%) as a light brown paste: ES(+)-HRMS m/e calcd for $C_{31}H_{38}N_2O_7$ (M+H)⁺ 551.2752, found 551.2748.

10

Example 23:

4-[2-(2-Carboxy-ethyl)-3-{6-[3-(1*H*-indol-5-yl)-5-methoxymethyl-phenoxy]-hexyl}-phenoxy]-butyric acid




15 A similar procedure as described in step 3, method B was used, starting from 4-[3-[6-(3-bromo-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (175 mg, 0.31 mmol), 5-indoleboronic acid (104 mg, 0.64 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (51 mg, 0.07 mmol), and cesium carbonate (417 mg, 1.28 mmol) to obtain 4-[2-(2-carboxy-ethyl)-3-{6-[3-(1*H*-indol-5-yl)-5-methoxymethyl-phenoxy]-hexyl}-phenoxy]-butyric acid (125 mg, 69%) as a light brown solid: ES(+)-HRMS m/e calcd for $C_{35}H_{41}NO_7$ (M+Na)⁺ 610.2775, found 610.2774.

20


Example 24:

- 51 -

4-[2-(2-Carboxy-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl -biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid




Step 1: 4-{2-(2-Ethoxycarbonyl-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester



To a solution of 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (250 mg, 0.37 mmol) in DMF (5 mL) were added sodium hydride (29.9 mg, 0.75 mmol), and iodomethane (106.2 mg, 0.75 mmol) at room temperature. The resulting suspension was stirred for 15 h and then the excess sodium hydride was quenched by slow addition of water (5 mL). The mixture was diluted with 1.0N hydrochloric acid (25 mL) and the organic compound was extracted into ethyl acetate (2 x 25 mL). The combined extracts were washed with brine solution (50 mL) and dried over anhydrous magnesium sulfate. Filtration of the drying agent and removal of the solvent under vacuum afforded the crude residue which was purified by using an ISCO 40 g column to obtain 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (200 mg, 78%) as a light brown oil: ES(+)-HRMS m/e calcd for C₃₈H₅₀O₉S (M+Na)⁺ 705.3068, found 705.3069.

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid


- 52 -



To a solution of the 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (195 mg, 0.29 mmol) in THF (5 mL) and ethanol (5 mL) was added aqueous 1.0 N sodium hydroxide (3 mL) at room temperature. The resulting suspension was stirred for 5 h at room temperature at which time TLC analysis of the mixture indicated the absence of starting material. Then, the reaction mixture was concentrated and the residue was diluted with water (20 mL) and extracted with diethyl ether (30 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N hydrochloric acid and the precipitated white organic compound was extracted into ethyl acetate (2 x 30 mL). The combined ethyl acetate extracts were washed with brine solution (50 mL) and the organic layers were dried over anhydrous magnesium sulfate. Filtration of the drying agent and removal of the solvent afforded 4-[2-(2-carboxy-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (168 mg, 94%) as a white solid: ES(+)-HRMS m/e calcd for C₃₄H₄₂O₉S (M+Na)⁺ 649.2442, found 649.2440.


Example 25 :

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



Step 1: 4-[3-[6-(3-Bromo-5-ethoxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester

- 53 -



To a solution of 4-[3-[6-(3-bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (1.0 g, 1.68 mmol) in *N,N*-dimethylformamide (10 mL) was added iodoethane (403 μ L, 5.04 mmol) and sodium hydride, 60% dispersion in mineral oil (202 mg, 5.04 mmol) at room temperature. The resulting suspension was stirred for 5 h. Then, the reaction mixture was diluted with water and brine. The organic compound was extracted into ethyl acetate and the combined organic extracts were washed with brine solution. The organic layers were dried over anhydrous sodium sulfate and the solvent was removed under reduced pressure. The crude material was purified by column chromatography (Isco™ 40 g) with 0- 30% ethyl acetate/hexanes as eluting solvents to afford the title compound (550 mg, 53%) as a colorless oil. HR-ES(+) calcd for $C_{32}H_{45}O_7Br$ $[M+Na]^+$ 643.2241, observed 643.2240.

15

Step 2 : General Procedure for Suzuki reaction and saponification


To a solution of 4-[3-[6-(3-Bromo-5-ethoxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (100 mg, 0.16 mmol), cesium carbonate (207 mg, 0.64 mmol) and boronic acid (0.32 mmol) in DME (4 mL) was added $Pd(PPh_3)_4$ (9 mg, 5 mol%). The reaction mixture was heated at 90 °C for 12 h, cooled down to room temperature then diluted with EtOAc. The resulting solution was washed with 10% aq. HCl and brine. The organic layer was then dried over anhydrous sodium sulfate and concentrated under vacuo to afford the crude material. Without further purification, the crude sample was dissolved in hot ethanol (4 mL) and 2.0N sodium hydroxide (1 mL) was added. The resulting solution was heated to 60 °C and stirred for 4 h. Then, the reaction mixture was cooled down and diluted with EtOAc, washed with 10% aq. HCl and brine. The combined ethyl acetate extracts were dried over anhydrous sodium sulfate and the

25

- 54 -

solvent was removed under reduced pressure. The desired products were isolated by preparative HPLC.

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 2 with 3,4-methylenedioxyphenyl boronic acid. HR-ES(+) calcd for $C_{35}H_{42}O_9$ ($M+Na$)⁺ 629.2721, found 629.2724.

10

Example 26:

4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




15 The title compound was prepared by following the general procedure for Suzuki coupling described above in step 2 with 4-fluorophenylboronic acid. HR-ES(+) calcd for $C_{34}H_{41}O_7F$ ($M+Na$)⁺ 603.2728, found 603.2733.

Example 27:

20 **4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**

- 55 -




The title compound was prepared by following the general procedure for Suzuki coupling described above in step 2 with 4-trifluoromethylphenylboronic acid. HR-ES(+) calcd for $C_{35}H_{41}O_7F_3$ ($M+Na$)⁺ 653.2696, found 653.2700.

5

Example 28:


4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-chloro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid



- 10 The title compound was prepared by following the general procedure for Suzuki coupling described above in step 2 with 4-chlorophenylboronic acid. HR-ES(+) calcd for $C_{34}H_{41}O_7Cl$ ($M+Na$)⁺ 619.2433, found 619.2434.


Example 29:

- 15 **4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid**




Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester

- 56 -



A similar procedure as described in **Example 24, step 1** was used, starting from 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (250 mg, 0.37 mmol), sodium hydride (29.9 mg, 0.75 mmol), and iodoethane (116.7 mg, 0.75 mmol) to obtain 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (110 mg, 42%) as a light brown oil: ES(+)-HRMS m/e calcd for C₃₉H₅₂O₉S (M+Na)⁺ 719.3224, found 719.3227.

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid




A similar procedure as described in **Example 24, step 2** was used, starting from 4-{2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid ethyl ester (101 mg, 0.145 mmol) and 1.0 N aqueous sodium hydroxide (1.5 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (90 mg, 97%) as a white solid: ES(+)-HRMS m/e calcd for C₃₅H₄₄O₉S (M+Na)⁺ 663.2598, found 663.2594.

Method C


Step 1: 4-[3-[6-(3-Bromo-5-methanesulfonyloxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester

- 57 -



To a solution of 4-[3-[6-(3-Bromo-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.99g, 5.05 mmol) in DCM (100 mL), TEA (3.52 mL, 25.26 mmol), DMAP (62 mg, 0.505 mmol) were added, followed by MsCl (0.60 mL, 7.58 mmol), and stirred at room temperature for 2.5 h. MeOH (4 mL) was added to quench the reaction. The solvent was removed under reduced pressure. Flash column chromatography (40% EtOAc/Hex) provided a colorless oil (2.51g, 74% yield). ¹H NMR (CDCl₃): δ 7.11-7.05 (m, 3H), 6.86 (s, 1H), 6.77 (d, 1H), 6.68 (d, 1H), 5.14 (s, 2H), 4.17-4.08 (m, 4H), 4.01-3.91 (m, 4H), 2.99-2.93 (m, 5H), 2.63 (dd, 2H), 2.56-2.46 (m, 4H), 2.17-0.89 (m, 16H).

Step 2: 4-[3-[6-(3-Bromo-5-dimethylaminomethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




The a solution of 4-[3-[6-(3-Bromo-5-methanesulfonyloxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.5 g, 3.72 mmol) in MeCN (40 mL), N,N-dimethylamine 2M/THF (3.72 mL, 7.44 mmol) was added, and stirred at room temperature overnight. The solvent was removed under reduced pressure. Flash column chromatography (50% EtOAc/Hex) provided a light yellow oil (1.47g, 64% yield). ¹H NMR (CDCl₃): δ 7.08 (t, 1H), 7.03 (s, 1H), 6.93 (s, 1H), 6.79 (s, 1H), 6.77 (d, 1H), 6.68 (d, 1H), 4.17-4.06 (m, 4H), 3.99 (dd, 2H), 3.92 (dd,

- 58 -

2H), 3.34 (s, 2H), 2.97 (dd, 2H), 2.63 (dd, 2H), 2.56-2.46 (m, 4H), 2.23 (s, 6H), 2.17-0.89 (m, 16H).

Step 3: 4-[3-[6-(3-Bromo-5-cyanomethyl-phenoxy)-hexyl]-2-(2-ethoxy carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



To a solution of 4-[3-[6-(3-bromo-5-methanesulfonyloxymethyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.51 g, 3.74 mmol) in DMF (40 mL), KCN (292 mg, 4.48 mmol) was added and stirred at 60 °C overnight. The reaction was diluted with EtOAc (250 mL) washed with water (100 mL X2), brine (50 mL), dried with Na₂SO₄. Concentration under reduced pressure provided an oil. Flash column chromatography (30% EtOAc/Hex) provided a light yellow oil (1.19 g, 53% yield). ¹H NMR (CDCl₃): δ 7.08 (t, 1H), 7.04 (s, 1H), 6.99 (s, 1H), 6.79 (s, 1H), 6.77 (d, 1H), 6.68 (d, 1H), 4.17-4.06 (m, 4H), 4.01-3.91 (m, 4H), 3.68 (s, 2H), 2.97 (dd, 2H), 2.63 (dd, 2H), 2.56-2.46 (m, 4H), 2.17-0.89 (m, 16H).


Step 4 : General Procedure for Suzuki reaction and saponification

In a sealable tube, an aryl bromide derivative (100 mg, 0.157 mmol) in DME (3 mL) was added, followed by Boronic acid (100 mg), Cs₂CO₃ (100 mg), and PdCl₂(dppf) (20 mg). The reaction mixture was shaken at 90 °C overnight. The reaction was diluted with EtOAc (5 mL), washed with water (3 mL). The organic layer was separated, concentration under reduced pressure gave an oil, which was used in the next step without further purification. The residue oil was dissolved in EtOH (5 mL), 3N NaOH (0.5 mL) was added and stirred at 60 °C for 3 h. 3N HCl (0.55 mL) was added to neutralize the reactions. Concentration under reduced pressure gave an oil which was purified by preparative HPLC to provide the following compounds.

- 59 -

Example 30:


4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-dimethylaminomethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



- 5 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 3,4-methylenedioxyphenylboronic Acid. LC / MS indicated a purity of 93% as measured by UV 214 nM. HR-ES(+):calculated for $C_{35}H_{44}NO_8$ ($M+H$)¹⁺ 606.3062, found 606.3060.
- 10

Example 31:


4-[2-(2-Carboxy-ethyl)-3-[6-(3-dimethylaminomethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid



15


- The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with thiopheneboronic acid.
- 20 LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{32}H_{42}NO_6S$ ($M+H$)¹⁺ 568.2728, found 568.2726.

- 60 -

Example 32:**4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**

- 5 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with phenylboronic acid. LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{34}H_{44}NO_6$ (M+H)¹⁺ 562.3163, found 562.3161.

10

Example 33:**4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**


- 15 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 4-fluorophenylboronic acid. LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{34}H_{43}NO_6F$ (M+H)¹⁺ 580.3069, found 580.3068.

20

- 61 -

Example 34:

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-dimethylaminomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




- 5 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 4-chlorophenylboronic acid. LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{34}H_{43}NO_6Cl$ ($M+H$)¹⁺ 596.2774, found 596.2774.

10

Example 35:

4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-fluoro-4'-methyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




- 15 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 3-fluor-4-methylphenylboronic acid. LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{35}H_{45}NO_6F$ ($M+H$)¹⁺ 594.3226, found 594.3225.

20

- 62 -

Example 36:

4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




- 5 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 3-fluorophenylboronic acid. LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{34}H_{43}NO_6F$ (M+H) $^{1+}$ 580.3069, found 580.3065.

10

Example 37:


4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid




- 15 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 4-trifluoromethylphenylboronic acid. LC / MS indicated a purity of 86% as measured by UV 214 nM. LC/MS: calculated for $C_{35}H_{43}F_3NO_6$ (M+H) $^{1+}$ 630.30, found 630.30.

20

- 63 -

Example 38:**4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**

- 5 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 4-methanesulfonylphenylboronic acid. LC / MS indicated a purity of 85% as measured by UV 214 nM. HR-ES(+):calculated for $C_{35}H_{46}NO_8S$ (M+H) $^{1+}$ 640.2939, found 640.2940.

Example 39:**4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid**


- 15 The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 3-methanesulfonylphenylboronic acid. LC / MS indicated a purity of 85% as

- 64 -

measured by UV 214 nM. HR-ES(+):calculated for $C_{35}H_{46}NO_8S$ (M+H)¹⁺ 640.2939, found 640.2937.

Example 40:


- 5 **4-{2-(2-Carboxy-ethyl)-3-[6-(3-dimethylaminomethyl-5-pyridin-4-yl-phenoxy)-hexyl]-phenoxy}-butyric acid**



- The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dimethylaminomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxy-carbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with 4-pyridineboronic acid.
 10 LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-ES(+):calculated for $C_{33}H_{43}N_2O_6$ (M+H)¹⁺ 563.3116, found 563.3113.

Example 41:


- 15 **4-{2-(2-Carboxy-ethyl)-3-[6-(3-cyanomethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid**



- The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-cyanomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with thiophene-3-boronic acid.
 20 LC / MS indicated a purity of 100% as measured by UV 214 nM. Calculated for $C_{31}H_{36}NO_6S$ (M+H)¹⁺ 550.22, found 550.20.

Example 42:

4-{2-(2-Carboxy-ethyl)-3-[6-(5-cyanomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid



5

The title compound was prepared according to general method described in step 4 by reaction of 4-[3-[6-(3-dyanomethyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester with phenylboronic acid.



LC / MS indicated a purity of 100% as measured by UV 214 nM. HR-

10 ES(+):calculated for $C_{33}H_{38}NO_6$ ($M+H$)¹⁺ 566.2513, found 566.2513.

Method D**Example 43:**

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

15


**Step 1: 5-Bromo-benzene-1,3-diol**

To a suspension of 5-bromo-1,3-dimethoxy-benzene (15 g, 67.02 mmol) and
20 sodium iodide (100.5 g, 670.2 mmol) in acetonitrile (577 mL) was added

- 66 -

trimethylsilyl chloride (42.39 mL, 335.1 mmol) at room temperature. Then, the resulting light yellow suspension was heated to reflux for 48 h. Then, it was cooled to room temperature and diluted with water (150 mL). The organic compound was extracted into ethyl acetate (2 x 150 mL) and the combined ethyl acetate extracts were washed with saturated sodium thiosulfate solution (300 mL) to remove the iodine color and was also washed with brine solution (300 mL). Then, the organic layer was dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. The crude residue was purified by using an ISCO 330 g column, eluting with 5-30% ethyl acetate in hexanes to obtain 5-bromo-benzene-1,3-diol (9.24 g, 73%) as a light brown solid: ES(+)-HRMS *m/e* calcd for C₆H₅BrO₂ (M+H)⁺ 189.2341, found 189.2346.


Step 2: 4-[3-[6-(3-Bromo-5-hydroxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



To a solution of 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (4.48 g, 9.5 mmol) and 5-bromo-benzene-1,3-diol (1.79 g, 9.47 mmol) in dimethylsulfoxide (30 mL) was added lithium hydride (160 mg, 20 mmol) at room temperature. The resulting suspension was stirred for 48 h and then the reaction mixture was diluted with water (100 mL). The organic compound was extracted into ethyl acetate (3 x 50 mL) and the combined organic extracts were washed with water (100 mL) and brine solution (100 mL). The organic layers were dried over anhydrous magnesium sulfate and filtration of the drying agent and concentration of the solvent under vacuum gave the crude product which was purified by using an ISCO 120 g column, eluting with 0-30% ethyl acetate in hexanes to afford 4-[3-[6-(3-bromo-5-hydroxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.73 g, 74%) as a white solid: ES(+)-HRMS *m/e* calcd for C₂₉H₃₉BrO₇ (M+Na)⁺ 601.1771, found 601.1773.


- 67 -

Step 3: 4-[3-[6-(3-Bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



To a mixture of 4-[3-[6-(3-bromo-5-hydroxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (163 mg, 0.28 mmol) and potassium carbonate (117 mg, 0.84 mmol) were added dimethylformamide (4 mL), acetone (8 mL), and iodoethane (219 mg, 1.4 mmol) at room temperature. The resulting suspension was heated to reflux for 2 days. Then, the reaction mixture was cooled to room temperature and diluted with water (20 mL). The organic compound was extracted into ethyl acetate (2 x 25 mL) and the combined organic extracts were washed with water (50 mL) and brine solution (50 mL). The organic layers were dried over anhydrous magnesium sulfate and filtration of the drying agent and concentration of the solvent gave the crude product which was purified by using an ISCO 40 g column, eluting with 0-30% ethyl acetate in hexanes to afford 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (166 mg, 98%) as a colorless oil: ES(+)-HRMS m/e calcd for $C_{31}H_{43}BrO_7$ (M+H)⁺ 607.2265, found 607.2270.

Step 4: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




To a mixture of 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (155 mg, 0.255 mmol), 3,4-methylenedioxybenzeneboronic acid (85 mg, 0.51 mmol), [1,1'-

- 68 -

bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (166 mg, 0.51 mmol) was added dimethoxyethane (10 mL) at room temperature under nitrogen atmosphere. The resulting brown reaction mixture was heated to 96 °C and stirred for 15 h at which time the TLC analysis of the reaction mixture indicated the absence of starting material. Then, the reaction mixture was cooled to room temperature and diluted with water (50 mL) and ethyl acetate (50 mL). The two layers were separated and the aqueous layer was extracted with ethyl acetate (50 mL) and the combined organic extracts were washed with water (50 mL) and brine solution (50 mL). The organic layer was dried over anhydrous magnesium sulfate and filtration of the drying agent and removal of the solvent under vacuum gave the colored residue which was purified by using an ISCO 40 column, eluting with 0-25% ethyl acetate in hexanes to obtain 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (165 mg, 99%) as a colorless viscous oil: ES(+)-LRMS m/e calcd for C₃₈H₄₈O₉ (M+Na)⁺ 671.3190, found 671.3185.

Step 5: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



To a solution of the 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (152 mg, 0.23 mmol) in THF (5 mL) and ethanol (5 mL) was added aqueous 1.0 N sodium hydroxide (5 mL) at room temperature. The resulting suspension was stirred for 5 h at room temperature at which time TLC analysis of the mixture indicated the absence of starting material. Then, the reaction mixture was concentrated and the residue was diluted with water (20 mL) and extracted with diethyl ether (30 mL) to remove any neutral impurities. The aqueous layer was acidified with 1.0 N hydrochloric acid and the precipitated white organic compound was extracted into ethyl acetate (2 x

- 69 -

30 mL). The combined ethyl acetate extracts were washed with brine solution (50 mL) and the organic layers were dried over anhydrous magnesium sulfate.


Filtration of the drying agent and removal of the solvent afforded 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-

- 5 butyric acid (129 mg, 95%) as a off-white waxy solid: ES(+)-HRMS m/e calcd for $C_{34}H_{40}O_9$ (M+H)⁺ 593.2745, found 593.2744.


Example 44:

4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-

- 10 **butyric acid**




Step 1: 3-Bromo-5-ethoxy-phenol



- A similar procedure as described in **Example 43, step 2** was used, starting from 5-bromo-benzene-1,3-diol (9.2 g, 48.67 mmol), iodoethane (8.13 g, 51.1 mmol), and lithium hydride (720 mg, 97.34 mmol) to afford 3-bromo-5-ethoxy-phenol (3.38 g, 32%) as a light yellow oil: ES(+)-HRMS m/e calcd for $C_8H_9BrO_2$ (M+Na)⁺ 217, found 217.


- 20 **Step 2: 4-[3-[6-(3-Bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester**

- 70 -



A similar procedure as described in **Example 43, step 3** was used, starting from 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (8.07 g, 17.13 mmol), 3-bromo-5-ethoxy-phenol (3.38 g, 15.57 mmol), and
 5 potassium carbonate (4.3 g, 31.14 mmol) to obtain 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (7.77 g, 82%) as a colorless oil: ES(+)-HRMS m/e calcd for $C_{31}H_{43}BrO_7$ (M+H)⁺ 607.2265, found 607.2270.

10 **Step 3: 4-[3-[6-(5-Ethoxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester**




To a mixture of 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), phenylboronic acid
 15 (125 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) in a microwave tube was added dimethoxyethane (5.9 mL) under nitrogen atmosphere at room temperature. The resulting brown reaction mixture was heated to 120 °C in a microwave for 30 minutes. The reaction mixture was diluted with ethyl acetate (50
 20 mL) and the undissolved residue was filtered off and washed with water (50 mL). Then, the two layers of filtrate were separated and the aqueous layer was extracted with ethyl acetate (20 mL). The combined organic extracts were washed with water (50 mL) and brine solution (50 mL). The organic layer was dried over anhydrous magnesium sulfate and filtration of the drying agent and removal of the

- 71 -

solvent under vacuum gave the colored residue which was purified by using an ISCO 40 column, eluting with 2-25% ethyl acetate in hexanes to obtain 4-[3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (255 mg, 84%) as a colorless viscous oil: ES(+)-LRMS m/e calcd for

5 $C_{37}H_{48}O_7$ (M+Na)⁺ 627.3292, found 627.3292.

Step 4: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid




10 A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (240 mg, 0.4 mmol) and 1.0 N aqueous sodium hydroxide (4 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (210 mg, 96%) as a light brown waxy solid: ES(+)-HRMS m/e

15 calcd for $C_{33}H_{40}O_7$ (M+Na)⁺ 571.2666, found 571.2665.

Example 45:


4-[2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid



20


Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid ethyl ester

- 72 -



- A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 3-thiophenylboronic acid (131 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid ethyl ester (160 mg, 52%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{35}H_{46}O_7S$ (M+Na)⁺ 633.2856, found 633.2860.

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid




- A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid ethyl ester (150 mg, 0.25 mmol) and 1.0 N aqueous sodium hydroxide (2.5 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid (136 mg, 99%) as a light brown waxy solid: ES(+)-HRMS m/e calcd for $C_{31}H_{38}O_7S$ (M+H)⁺ 555.2411, found 555.2410.


Example 46:

4-[2-(2-Carboxy-ethyl)-3-[6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy]-butyric acid

- 73 -




Step 1: 4-[3-{6-[3-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



- 5 A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 1,4-benzodioxane-6-boronic acid (184 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[3-{6-[3-(2,3-
- 10 dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (76 mg, 23%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for C₃₉H₅₀O₉ (M+Na)⁺ 685.3347, found 685.3353.

- Step 2: 4-[2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl}-phenoxy]-butyric acid**
- 15





A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (70 mg, 0.1 mmol) and 1.0

- 74 -

N aqueous sodium hydroxide (1 mL) to afford 4-[2-(2-carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy]-butyric acid (62 mg, 99%) as a light brown waxy solid: ES(+)-HRMS m/e calcd for $C_{35}H_{42}O_9$ ($M+Na$)⁺ 629.2721, found 629.2721.

5


Example 47:**4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid****Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester**

A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 3-fluoro-phenylboronic acid (142 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (220 mg, 70%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{37}H_{47}FO_7$ ($M+Na$)⁺ 645.3198, found 645.3198.

20

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid

- 75 -




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (210 mg, 0.34 mmol) and 1.0 N aqueous sodium hydroxide (3.4 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (190 mg, 99%) as a light brown waxy solid: ES(+)-HRMS m/e calcd for $C_{33}H_{39}FO_7$ ($M+Na$)⁺ 589.2572, found 589.2573.

10


Example 48:

4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid



Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester

15




A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 4-fluoro-phenylboronic acid (141 mg,

- 76 -

1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (161 mg, 52%) as a colorless viscous oil: ES(+)-HRMS m/e

5 calcd for $C_{37}H_{47}FO_7$ ($M+Na$)⁺ 645.3198, found 645.3196.

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid




10 A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (150 mg, 0.24 mmol) and 1.0 N aqueous sodium hydroxide (2.4 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (135 mg, 99%) as a light brown

15 waxy solid: ES(+)-HRMS m/e calcd for $C_{33}H_{39}FO_7$ ($M+Na$)⁺ 589.2572, found 589.2573.

Example 49:


4-[2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid

20



Step 1: 4-[3-[6-(4'-Chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester


- 77 -



A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 4-chloro-phenylboronic acid (165 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (231 mg, 72%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{37}H_{47}ClO_7$ ($M+Na$)⁺ 661.2902, found 661.2905.

10

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (150 mg, 0.24 mmol) and 1.0 N aqueous sodium hydroxide (2.4 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (200 mg, 98%) as a light brown waxy solid: ES(+)-HRMS m/e calcd for $C_{33}H_{39}ClO_7$ ($M+Na$)⁺ 605.2276, found 605.2278.

20


Example 50:

4-[2-(2-Carboxy-ethyl)-3-[6-(3',4'-difluoro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid

- 78 -




Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(3',4'-difluoro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester



- 5 A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 3,4-difluoro-phenylboronic acid (164 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(3',4'-difluoro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (75 mg, 23%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{37}H_{46}F_2O_7$ ($M+Na$)⁺ 663.3104, found 663.3102.

- Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(3',4'-difluoro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid**




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(3',4'-difluoro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (65 mg, 0.1 mmol) and 1.0 N aqueous sodium


- 79 -

hydroxide (1.0 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(3',4'-difluoro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (58 mg, 98%) as a light brown waxy solid: ES(+)-HRMS m/e calcd for $C_{33}H_{38}F_2O_7$ ($M+Na$)⁺ 607.2478, found 607.2481.

5

Example 51:**4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid****Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester**

10




A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 4-methanesulfonyl-phenylboronic acid (207 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-phenoxy]-butyric acid ethyl ester (57 mg, 17%) as a light brown oil: ES(+)-HRMS m/e calcd for $C_{38}H_{50}O_9S$ ($M+Na$)⁺ 705.3068, found 705.3065.

15

20

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid

- 80 -




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid ethyl ester (52 mg, 0.076 mmol) and 1.0 N aqueous sodium hydroxide (1.0 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (44 mg, 91%) as a light brown solid: ES(+)-HRMS m/e calcd for C₃₄H₄₂O₉S (M+Na)⁺ 649.2442, found 649.2442.

10


Example 52:

4-[2-(2-Carboxy-ethyl)-3-[6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy]-butyric acid



Step 1: 4-[3-[6-[3-(2-Chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester

15




A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 2-chloro-pyridin-4-ylboronic acid (161

- 81 -

mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[3-{6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (115 mg, 36%) as a light yellow oil: ES(+)-HRMS m/e calcd for $C_{36}H_{46}ClNO_7$ (M+H)⁺ 640.3036, found 640.3032.


Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy]-butyric acid



A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-{6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (105 mg, 0.16 mmol) and 1.0 N aqueous sodium hydroxide (1.6 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy]-butyric acid (92 mg, 96%) as a light brown solid: ES(+)-HRMS m/e calcd for $C_{32}H_{38}ClNO_7$ (M+H)⁺ 584.2410, found 584.2412.


Example 53:

4-[2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid




Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid ethyl ester

- 82 -



A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), pyrimidin-5-ylboronic acid (124 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid ethyl ester (94 mg, 31%) as a light brown oil: ES(+)-HRMS m/e calcd for $C_{35}H_{46}N_2O_7$ (M+H)⁺ 607.3378, found 607.3379.

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid ethyl ester (76 mg, 0.13 mmol) and 1.0 N aqueous sodium hydroxide (1.3 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy]-butyric acid (63 mg, 92%) as a light brown solid: ES(+)-HRMS m/e calcd for $C_{31}H_{38}N_2O_7$ (M+H)⁺ 551.2752, found 551.2750.

20


Example 54:

4-[2-(2-Carboxy-ethyl)-3-[6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl]-phenoxy]-butyric acid

- 83 -




Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid ethyl ester



- 5 A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 5-indolineboronic acid (170 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid ethyl ester (25 mg, 8%) as a light yellow oil: ES(+)-HRMS m/e calcd for $C_{39}H_{49}NO_7$ ($M+Na$)⁺ 666.3401, found 666.3399.

Step 2: 4-[2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid ethyl ester (24 mg, 0.04 mmol) and 1.0 N aqueous sodium


- 84 -

hydroxide (0.4 mL) to afford 4-[2-(2-carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid (20 mg, 91%) as a light brown solid: ES(+)-HRMS *m/e* calcd for C₃₅H₄₁NO₇ (M+Na)⁺ 610.2775, found 610.2773.

5

Example 55:**4-[2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid****Step 1: 4-[2-(2-Ethoxycarbonyl-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid ethyl ester**

10




A similar procedure as described in **Example 44, step 3** was used, starting from 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (305 mg, 0.5 mmol), 6-indolineboronic acid (170 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (55 mg, 0.075 mmol), and cesium carbonate (331 mg, 1.0 mmol) to obtain 4-[2-(2-ethoxycarbonyl-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid ethyl ester (125 mg, 39%) as a light yellow oil: ES(+)-HRMS *m/e* calcd for C₃₉H₄₉NO₇ (M+Na)⁺ 666.3401, found 666.3396.

20

Step 2: 4-[2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid


- 85 -




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[2-(2-ethoxycarbonyl-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid ethyl ester (111 mg, 0.17 mmol) and 1.0 N aqueous sodium hydroxide (1.7 mL) to afford 4-[2-(2-carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy]-butyric acid (79 mg, 78%) as a light brown solid: ES(+)-HRMS *m/e* calcd for C₃₅H₄₁NO₇ (M+Na)⁺ 610.2775, found 610.2774.

Example 56:

10 4-[3-{6-[3-Benzothiazol-5-yl-5-ethoxy-phenoxy]-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



Step 1: 4-[3-{6-(3-Benzothiazol-5-yl-5-ethoxy-phenoxy)-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




To a suspension of palladium(II) acetate (11.32 mg, 0.05 mmol), triphenylphosphine (26.2 mg, 0.1 mmol), and benzothiazol-5-yl pinacolboronate (258 mg, 0.99 mmol) in dioxane (7.5 mL) were added 4-[3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (300 mg, 0.49 mmol),

- 86 -

potassium phosphate (216 mg, 0.99 mmol), and pure water (15 mg, 0.99 mmol) at room temperature. Then, the resulting mixture was heated to 100 °C for 2 h. Then, the reaction mixture was cooled to room temperature and diluted with water (50 mL) and ethyl acetate (50 mL). The two layers were separated and the aqueous layer was extracted with ethyl acetate (50 mL) and the combined organic extracts were washed with water (50 mL) and brine solution (50 mL). The organic layer was dried over anhydrous magnesium sulfate and filtration of the drying agent and removal of the solvent under vacuum gave the colored residue which was purified by using an ISCO 40 g column, eluting with 2-25% ethyl acetate in hexanes to obtain 4-[3-{6-(3-benzothiazol-5-yl-5-ethoxy-phenoxy)-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (61 mg, 99%) as a off-white solid: ES(+)-LRMS m/e calcd for C₃₈H₄₇NO₇S (M+H)⁺ 662.3146, found 662.3144.

Step 2: 4-[3-{6-[3-Benzothiazol-5-yl-5-ethoxy-phenoxy]-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid





A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-{6-(3-benzothiazol-5-yl-5-ethoxy-phenoxy)-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (61 mg, 0.17 mmol) and 1.0 N aqueous sodium hydroxide (1.0 mL) to afford 4-[3-{6-[3-benzothiazol-5-yl-5-ethoxy-phenoxy]-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (9.2 mg, 17%) as a light brown solid: ES(+)-HRMS m/e calcd for C₃₄H₃₉NO₇S (M+H)⁺ 606.2520, found 606.2523.


Example 57:

4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-phenoxy)-butyric acid

- 87 -

**Step 1: 2,2-Difluoro-benzo[1,3]dioxole-5-boronic acid:**

5 *n*-BuLi in hexanes (2.5 M solution, 46 mL) was added to anhydrous THF (250 mL)
 cooled in acetone-dry ice bath under the flow of nitrogen. After 10 min 5-bromo-
 2,2-difluoro-1,3-benzodioxole (25.0 g) was added dropwise and the resulting
 solution was stirred for 30 min. Then, boron isopropoxide (30 mL) was added
 dropwise and the reaction mixture was allowed to reach room temperature over a
 period of 3 h. Then acetic acid (30 mL) was added in one portion and the resulting
 10 mixture was stirred at room temperature for 10 min. The reaction mixture was then
 diluted with ethyl acetate and washed with water and brine. The organic extract
 was dried over anhydrous sodium sulfate and concentrated under reduced
 pressure. The title compound was obtained by trituration with hexanes as a white
 solid (14.2 g, 73% yield). HRMS calcd for $C_7H_5O_4BF_2$ $[2M-H-H_2O]^-$ 385.0324,
 15 observed 385.0319.


Step 2: 4-[3-{6-[3-(2,2-Difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester


20 A similar procedure as described in **Example 44, step 3** was used, starting from 4-
 [3-[6-(3-bromo-5-ethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-
 butyric acid ethyl ester (577 mg, 0.95 mmol), 2,2-difluoro-benzo[1,3]dioxo-5-

- 88 -

ylboronic acid trimer (524 mg, 0.95 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (104 mg, 0.14 mmol), and cesium carbonate (625 mg, 1.9 mmol) to obtain 4-[3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (410 mg, 63%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{38}H_{46}F_2O_9$ (M+H)⁺ 685.3183, found 685.3180.

Step 3: 4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-phenoxy)-butyric acid




10

A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (395 mg, 0.58 mmol) and 1.0 N aqueous sodium hydroxide (5.8 mL) to afford 4-(2-(2-carboxy-ethyl)-3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-phenoxy)-butyric acid (260 mg, 72%) as a light yellow solid: ES(+)-HRMS m/e calcd for $C_{34}H_{38}F_2O_9$ (M+Na)⁺ 651.2376, found 651.2381.

15

Example 58:


4-[3-{6-(5-Benzyloxy-biphenyl-3-yloxy)-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



20


- 89 -

Step 1: 4-[3-[6-(3-Benzyloxy-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



A similar procedure as described in **Example 43, step 3** was used, starting from 4-[3-[6-(3-bromo-5-hydroxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (930 mg, 1.6 mmol), benzyl bromide (343 mg, 2.0 mmol), and potassium carbonate (553 mg, 4.0 mmol) to afford 4-[3-[6-(3-benzyloxy-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (1.06 g, 98%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{36}H_{45}BrO_7$ (M+Na)⁺ 691.2241, found 691.2238.

Step 2: 4-[3-[6-(5-Benzyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-benzyloxy-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (260 mg, 0.39 mmol), phenylboronic acid (95 mg, 0.78 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1

- 90 -

mmol), and cesium carbonate (253 mg, 0.78 mmol) to obtain 4-[3-[6-(5-benzyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (230 mg, 89%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{42}H_{50}O_7$ ($M+Na$)⁺ 689.3449, found 689.3447.

5

Step 3: 4-[3-[6-(5-Benzyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(5-benzyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (224 mg, 0.34 mmol) and 1.0 N aqueous sodium hydroxide (5 mL) to afford 4-[3-[6-(5-benzyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (165 mg, 80%) as an amorphous white solid: ES(+)-HRMS m/e calcd for $C_{38}H_{42}O_7$ ($M+Na$)⁺ 633.2823, found 633.2820.

15


Example 59:

4-[3-[6-(5-Benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




- 91 -

Step 1: 4-[3-[6-(5-Benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-benzyloxy-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (260 mg, 0.39 mmol), 3-fluoro-phenylboronic acid (109 mg, 0.78 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (253 mg, 0.78 mmol) to obtain 4-[3-[6-(5-benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (244 mg, 92%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for C₄₂H₄₉FO₇ (M+Na)⁺ 707.3354, found 707.3355.

Step 2: 4-[3-[6-(5-Benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(5-benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (229 mg, 0.34 mmol) and 1.0 N aqueous sodium hydroxide (5 mL) to afford 4-[3-[6-(5-benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid.

- 92 -


(2-carboxy-ethyl)-phenoxy]-butyric acid (163 mg, 78%) as an amorphous white solid: ES(+)-HRMS m/e calcd for $C_{38}H_{41}FO_7$ ($M+Na$)⁺ 651.2728, found 651.2729.

Example 60:

5 **4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid**



Step 1: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




10

A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-benzyloxy-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (260 mg, 0.39 mmol), 3,4-methylenedioxyphenylboronic acid (127 mg, 0.78 mmol), [1,1'-

15 bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (253 mg, 0.78 mmol) to obtain 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (260 mg, 94%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{43}H_{50}O_9$ ($M+Na$)⁺ 733.3347, found 733.3345.


Step 2: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



- 5 A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (254 mg, 0.36 mmol) and 1.0 N aqueous sodium hydroxide (5 mL) to afford 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (107 mg, 46%) as an
- 10 amorphous off-white solid: ES(+)-HRMS m/e calcd for $C_{39}H_{42}O_9$ ($M+Na$)⁺ 677.2721, found 677.2718.


Example 61:

- 4-[3-[6-(3-Benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid**
- 15



Step 1: 4-[3-[6-(3-Benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester


- 94 -



A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-benzyloxy-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (260 mg, 0.39 mmol), 3-thiophenylboronic acid (99 mg, 0.78 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (253 mg, 0.78 mmol) to obtain 4-[3-[6-(3-benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (165 mg, 67%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{40}H_{48}O_7S$ ($M+Na$)⁺ 695.3013, found 695.3008.

10

Step 2: 4-[3-[6-(3-Benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(3-benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (160 mg, 0.24 mmol) and 1.0 N aqueous sodium hydroxide (5 mL) to afford 4-[3-[6-(3-benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (129 mg, 88%) as a light brown waxy solid: ES(+)-HRMS m/e calcd for $C_{36}H_{40}O_7S$ ($M+Na$)⁺ 639.2387, found 639.2386.

15


Example 62:

4-[2-(2-Carboxy-ethyl)-3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid



5

Step 1: 4-[3-[6-(3-Bromo-5-cyclopentylmethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




A similar procedure as described in **Example 43, step 3** was used, starting from 4-[3-[6-(3-bromo-5-hydroxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (600 mg, 1.03 mmol), iodomethylcyclopentane (260 mg, 1.24 mmol), and potassium carbonate (356 mg, 2.57 mmol) to afford 4-[3-[6-(3-bromo-5-cyclopentylmethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (487g, 73%) as a colorless viscous oil: EI(+)-LRMS m/e calcd for C₃₅H₄₉BrO₇ (M+H)⁺ 663.45, found 663.5.

15

Step 2: 4-[3-[6-(3-Cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester


- 96 -



A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-bromo-5-cyclopentylmethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (136 mg, 0.2 mmol), 3-thiophenylboronic acid (53 mg, 0.41 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (134 mg, 0.41 mmol) to obtain 4-[3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (129 mg, 95%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for $C_{39}H_{52}O_7S$ ($M+Na$)⁺ 687.3326, found 687.3328.

10

Step 3: 4-[2-(2-Carboxy-ethyl)-3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (117 mg, 0.17 mmol) and 1.0 N aqueous sodium hydroxide (5 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy]-butyric acid (104 mg, 98%) as a off-white waxy solid: ES(+)-HRMS m/e calcd for $C_{35}H_{44}O_7S$ ($M+H$)⁺ 609.2881, found 609.2881.

20


Example 63:

4-[2-(2-Carboxy-ethyl)-3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid



5

Step 1: 4-[3-[6-(5-Cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-bromo-5-cyclopentylmethoxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (152 mg, 0.23 mmol), 3-fluorophenylboronic acid (65 mg, 0.46 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (150 mg, 0.46 mmol) to obtain 4-[3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (114 mg, 74%) as a colorless viscous oil: ES(+)-HRMS m/e calcd for C₄₁H₅₃FO₇ (M+H)⁺ 677.3848, found 677.3850.

15

Step 2: 4-[2-(2-Carboxy-ethyl)-3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid

- 98 -




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (104 mg, 0.15 mmol) and 1.0 N aqueous sodium hydroxide (5 mL) to afford 4-[2-(2-carboxy-ethyl)-3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy]-butyric acid (92 mg, 97%) as a off-white waxy solid: ES(+)-HRMS m/e calcd for C₃₇H₄₅FO₇ (M+H)⁺ 621.3222, found 621.3222.


10

Method E**Example 64:**

4-[3-[6-(3-Acetyl-5-benzo[1,3]dioxol-5-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




15

Step 1: 1-(3-Bromo-5-iodo-phenyl)-ethanone

To a solution of 3-bromo-5-iodobenzoic acid (4.9 g, 15.0 mmol) in dichloromethane (50 mL) were added dimethylformamide (250 μ L) and oxalyl chloride (2.54 mL, 30

- 99 -

mmol) at 0 °C. Then, the reaction mixture was warmed to room temperature in 1 h and was stirred for 15 h. Then, the solvent was removed under vacuum and dried under high vacuum to obtain the corresponding benzoyl chloride. In another 3-necked RB flask, lithium chloride (1.69 g, 40 mmol, predried at 130 °C under high vacuum for 2 h) and copper cyanide (1.79 g, 20 mmol) was charged and tetrahydrofuran (40 mL) was added at room temperature and it was stirred for 10 min to obtain a clear solution. Then, the reaction mixture was cooled to -70 °C and slowly treated with a 3M solution of methylmagnesium chloride in tetrahydrofuran (6.66 mL, 20 mmol). After addition, the reaction mixture was allowed to warm to -30 °C where it was stirred for 5 min. The resulting clear reaction mixture was again cooled back to -70 °C and then slowly treated with a solution of the above prepared substituted benzoyl chloride (15 mmol) in tetrahydrofuran (40 mL). After addition, the reaction mixture was allowed to warm to -20 °C and stirred for 15 h. The reaction mixture was then poured into a solution consisting of a saturated aqueous ammonium chloride solution (100 mL) and ammonium hydroxide (20 mL) and the organic compound was extracted into ethyl acetate (2 x 100 mL) and the combined ethyl acetate extracts were washed with brine solution (300 mL). Then, the organic layer was dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. The crude residue was purified by using an ISCO 120 g column, eluting with 0-20% ethyl acetate in hexanes to obtain 1-(3-bromo-5-iodo-phenyl)-ethanone (3.15 g, 65%) as a white solid: ES(+)-HRMS m/e calcd for C₈H₆BrIO (M)⁺ 323.8646, found 323.8642.

Step 2: 1-(3-Bromo-5-hydroxy-phenyl)-ethanone

25

To a mixture of 1-(3-bromo-5-iodo-phenyl)-ethanone (2.28 g, 7.0 mmol), sodium hydroxide (1.4 g, 35.1 mmol), and Cu₂O (143 mg, 1.0 mmol) were added pyridine (20 mL) and water (40 mL) at room temperature. The resulting suspension was heated to 110 °C and stirred for 24 h. Then the reaction mixture was cooled to room temperature and diluted with 1N hydrochloric acid (200 mL). Then, the


30

- 100 -

organic compound was extracted into ethyl acetate (3 x 100 mL) and the combined organic extracts were washed with 1N hydrochloric acid (200 mL) and brine solution (200 mL). The organic layers were dried over anhydrous magnesium sulfate and filtration of the drying agent and concentration of the solvent under vacuum gave the crude dark brown solid which was tried to dissolve in acetonitrile (50 mL) at reflux, but it did not dissolve completely. Then, it was diluted with ethyl acetate (~10 mL) and hexanes (~15 mL). The dark suspension was stored in the refrigerator. Only few solids were formed and they were collected by filtration and washed with acetonitrile. Then, the mother liquor was removed under vacuum and the dark brown solids were now dissolved in acetonitrile (50 mL) at reflux and it was treated with charcoal to remove some color. It was then filtered and the solvent was removed under vacuum to afford reasonably pure 1-(3-bromo-5-hydroxy-phenyl)-ethanone (965 mg, 64%) as a brown solid: ES(-)-HRMS m/e calcd for C₈H₇BrO₂ (M-H)⁻ 212.9556, found 212.9556.

15

Step 3: 4-[3-[6-(3-Acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




A similar procedure as described in **Example 43, step 3** was used, starting from 4-[3-[6-bromo-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.36 g, 5.0 mmol), 1-(3-bromo-5-hydroxy-phenyl)-ethanone (965 mg, 4.48 mmol), and potassium carbonate (1.38 g, 10 mmol) to afford 4-[3-[6-(3-acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (2.15 g, 79%) as a colorless viscous oil: EI(+)-HRMS m/e calcd for C₃₁H₄₁BrO₇ (M+Na)⁺ 627.1928, found 627.1925.

25


Step 4: 4-[3-[6-(3-Acetyl-5-benzo[1,3]-dioxol-5-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester

- 101 -



- A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (303 mg, 0.5 mmol), 3,4-methylenedioxyphenylboronic acid (166 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (326 mg, 1.0 mmol) to obtain 4-[3-[6-(3-acetyl-5-benzo[1,3]-dioxol-5-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (235 mg, 73%) as a colorless oil: ES(+)-HRMS m/e calcd for $C_{38}H_{46}O_9$ ($M+Na$)⁺ 669.3034, found 669.3031.

Step 5: 4-[3-[6-(3-Acetyl-5-benzo[1,3]dioxol-5-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




- A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(3-acetyl-5-benzo[1,3]-dioxol-5-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (230 mg, 0.36 mmol) and 1.0 N aqueous sodium hydroxide (8 mL) to afford 4-[3-[6-(3-acetyl-5-benzo[1,3]dioxol-5-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (192 mg, 92%) as a white solid: ES(+)-HRMS m/e calcd for $C_{34}H_{38}O_9$ ($M+Na$)⁺ 613.2408, found 613.2407.


Example 65:

4-[3-[6-(3-Acetyl-5-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

- 102 -




Step 1: 4-[3-[6-(3-Acetyl-5-(2,3-dihydro-benzo[1,4]-dioxin-6-yl)-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



5

A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (303 mg, 0.5 mmol), 1,4-benzodioxane-6-boronic acid (180 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (326 mg, 1.0 mmol) to obtain 4-[3-[6-(3-acetyl-5-(2,3-dihydro-benzo[1,4]-dioxin-6-yl)-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (330 mg, 99%) as a colorless oil: ES(+)-HRMS m/e calcd for C₃₉H₄₈O₉ (M+Na)⁺ 683.3190, found 683.3194.



Step 2: 4-[3-[6-(3-Acetyl-5-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(3-acetyl-5-(2,3-dihydro-benzo[1,4]-dioxin-6-yl)-phenoxy)-hexyl]-2-(2-

- 103 -


ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (330 mg, 0.5 mmol) and 1.0 N aqueous sodium hydroxide (10 mL) to afford 4-[3-[6-(3-acetyl-5-(2,3-dihydrobenzo[1,4]dioxin-6-yl)-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (200 mg, 61%) as a off-white solid: ES(+)-HRMS m/e calcd for $C_{35}H_{40}O_9$ ($M+Na$)⁺ 627.2564, found 627.2560.

Example 66:**4-[3-[6-(5-Acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid****Step 1: 4-[3-[6-(5-Acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester**

A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (303 mg, 0.5 mmol), phenylboronic acid (122 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (326 mg, 1.0 mmol) to obtain 4-[3-[6-(5-acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (288 mg, 96%) as a colorless oil: ES(+)-HRMS m/e calcd for $C_{37}H_{46}O_7$ ($M+Na$)⁺ 625.3136, found 625.3133.

Step 2: 4-[3-[6-(5-Acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid


- 104 -




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(5-acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (282 mg, 0.47 mmol) and 1.0 N aqueous sodium hydroxide (10 mL) to afford 4-[3-[6-(5-acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (168 mg, 66%) as a white solid: ES(+)-HRMS m/e calcd for C₃₃H₃₈O₇ (M+Na)⁺ 569.2510, found 569.2508.

Example 67:

10 4-[3-[6-(5-Acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



Step 1: 4-[3-[6-(5-Acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (303 mg, 0.5 mmol), 3-fluoro-phenylboronic acid (140 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1

- 105 -

mmol), and cesium carbonate (326 mg, 1.0 mmol) to obtain 4-[3-[6-(5-acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (288 mg, 93%) as a colorless oil: ES(+)-HRMS m/e calcd for $C_{37}H_{45}FO_7$ ($M+Na$)⁺ 643.3041, found 643.3039.

5

Step 2: 4-[3-[6-(5-Acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(5-acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (275 mg, 0.44 mmol) and 1.0 N aqueous sodium hydroxide (10 mL) to afford 4-[3-[6-(5-acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (165 mg, 66%) as a white solid: ES(+)-HRMS m/e calcd for $C_{33}H_{37}FO_7$ ($M+Na$)⁺ 587.2415, found 587.2415.

15

Example 68:


4-[3-[6-(3-Acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



Step 1: 4-[3-[6-(3-Acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester

20


- 106 -



A similar procedure as described in **Example 43, step 4** was used, starting from 4-[3-[6-(3-acetyl-5-bromo-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (303 mg, 0.5 mmol), 3-thiopheneboronic acid (128 mg, 1.0 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (73 mg, 0.1 mmol), and cesium carbonate (326 mg, 1.0 mmol) to obtain 4-[3-[6-(3-acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (285 mg, 94%) as a light brown oil: ES(+)-HRMS m/e calcd for $C_{35}H_{44}O_7S$ ($M+Na$)⁺ 631.2700, found 631.2704.

10

Step 2: 4-[3-[6-(3-Acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




A similar procedure as described in **Example 43, step 5** was used, starting from 4-[3-[6-(3-acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (272 mg, 0.44 mmol) and 1.0 N aqueous sodium hydroxide (10 mL) to afford 4-[3-[6-(3-acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid (163 mg, 66%) as a light brown solid: ES(+)-HRMS m/e calcd for $C_{31}H_{36}O_7S$ ($M+Na$)⁺ 575.2074, found 575.2074.

20


Example 69:

4-[3-[6-[3-Acetyl-5-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenoxy]-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

- 107 -




Step 1: Trifluoro-methanesulfonic acid 3-acetyl-5-trifluoromethanesulfonyloxy-phenyl ester



- 5 3,5-Dihydroxyacetophenone (5.0 g) was suspended in 100 mL of dichloromethane and then pyridine (10.6 mL) was added. The resulting clear solution was cooled to 0 °C and 11.6 mL of triflic anhydride were added dropwise. After addition was completed, the cooling bath was removed and the reaction mixture was stirred at room temperature for 1 h and then quenched with 1 N hydrochloric acid. The
- 10 organic solution was separated and washed with saturated sodium bicarbonate solution and brine and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure and the title compound was obtained by trituration with diethyl ether as white solid (9.9 g, 72% yield). HRMS calcd for $C_{10}H_6O_7S_2F_6$ M^+ 415.9459, observed 415.9459

15

Step 2: Trifluoro-methanesulfonic acid 3-acetyl-5-hydroxy-phenyl ester




- Trifluoro-methanesulfonic acid 3-acetyl-5-trifluoromethanesulfonyloxy-phenyl ester
- 20 (9.9 g) was dissolved in 1,2-dimethoxyethane (80 mL), followed by addition of cesium carbonate (11.6 g). The resulting suspension was stirred at 80 °C for 3 h and then quenched with saturated solution of ammonium chloride. The solution

- 108 -


was then extracted with diethyl ether and combined etherate extracts were washed with brine, dried over anhydrous sodium sulfate, and filtered through a silica pad. The filtrate was concentrated under reduced pressure and purified on a silica gel column using ethyl acetate and hexanes to afford 3.5 g of the title compound (52% yield). HRMS calcd for $C_{19}H_{17}O_5S_1F_3$ $[M+H]^+$ 285.0039, observed 285.0039.

Step 3: 4-[3-[6-(3-Acetyl-5-trifluoromethanesulfonyloxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



- To a solution of trifluoro-methanesulfonic acid 3-acetyl-5-hydroxy-phenyl ester (1.0 g) in a mixture of acetone and DMF (2:1, 80 mL) were added potassium carbonate (2.4 g) and 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (1.8 g). The resulting mixture was stirred at 75 °C for 3 h. Then the insoluble material was filtered out and the filtrate was diluted with ethyl acetate and washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 1.7 g of the title compound (72% yield). HRMS calcd for $C_{32}H_{41}O_{10}S_1F_3$ $[M+H]^+$ 675.2446, observed 675.2440.

Step 4: 4-[3-{6-[3-Acetyl-5-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenoxy]hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




To a solution of 4-[3-[6-(3-acetyl-5-trifluoromethanesulfonyloxy-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (500 mg) in 1,2-

- 109 -

dimethoxyethane (5 mL) were added 2,2-difluoro-benzo[1,3]dioxole-5-boronic acid (273 mg), 2 M aq. sodium carbonate solution (1.0 mL) and Pd(PPh₃)₄ (8 mg) and the resulting mixture was stirred at 90 °C for 3 h. Then the reaction mixture was diluted with ethyl acetate, filtered through Celite and washed with water and brine.

- 5 The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 300 mg of the title compound (59% yield). HRMS calcd for C₃₈H₄₄O₉F₂ [M+Na]⁺ 705.2845, observed 705.2843.

10 **Step 5: 4-[3-{6-[3-Acetyl-5-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenoxy]-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid**





- To a solution of 4-[3-{6-[3-acetyl-5-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenoxy]-hexyl}-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (300 mg) in ethanol (5 mL) was added a solution of NaOH (176 mg) in water (2 mL) and the resulting mixture was stirred at room temperature for 5 h. Then the reaction mixture was diluted with water, acidified with 3 N HCl and extracted into ethyl acetate. The organic extract was washed with brine, dried over anhydrous sodium sulfate, concentrated under reduced pressure and purified on a reverse-phase HPLC to
- 15 yield 137 mg of the title compound (50% yield). HRMS calcd for C₃₄H₃₆O₉F₂ [M+Na]⁺ 649.2219, observed 649.2222.
- 20


Example 70:

- 25 **4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-isobutyryl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid**

- 110 -

**Step 1: 3-Bromo-5-(4-methoxy-benzyloxy)-benzonitrile**

p-Methoxybenzyl alcohol (2.6 mL) was dissolved in anhydrous DMF (50 mL) and
 5 NaH (808 mg, 60% dispersion in oil) was added in several portions. After gas
 evolution ceased, 3,5-dibromobenzonitrile (5.0 g) was added and the reaction
 mixture was stirred at 70 °C. After 2 h additional amount of NaH (147 mg) was
 added and heating was continued for 6 h. The reaction mixture was diluted with
 ethyl acetate and washed with water and brine. The organic extract was dried over
 10 anhydrous sodium sulfate, concentrated and purified on a silica gel column using
 ethyl acetate and hexanes to yield 3.1 g of the title compound (51% yield). HRMS
 calcd for C₁₅H₁₂O₂BrN M⁺ 371.0051, observed 371.0048.

Step 2: 3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-benzonitrile

15


To a solution of 3-bromo-5-(4-methoxy-benzyloxy)-benzonitrile (1.1 g) in 1,2-
 dimethoxyethane (10 mL) were added 3,4-methylenedioxyphenylboronic acid (860
 mg), 2 M aq. sodium carbonate solution (3.5 mL) and Pd(PPh₃)₄ (40 mg) and the
 resulting mixture was stirred at 90 °C for 4 h. Then the reaction mixture was diluted

- 111 -

with ethyl acetate, filtered through Celite and washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, filtered through a silica pad, concentrated and triturated with diethyl ether to yield 1.0 g of the title compound (89% yield). HRMS calcd for $C_{22}H_{17}O_4N$ M^+ 359.1158, observed

5 359.1158.


Step 3: 1-[3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-phenyl]-2-methyl-propan-1-one



10 To a solution of 3-benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-benzonitrile (500 mg) in anhydrous THF (10 mL) were added a solution of isopropylmagnesium chloride (2.1 mL of 2 M solution in diethyl ether) and catalytic amount of cuprous bromide (4 mg). The resulting reaction mixture was refluxed for 2 h. After it was cooled down, 10% sulfuric acid (5 mL) was added and stirring was continued for 30

15 min at room temperature. Then the reaction mixture was diluted with diethyl ether, washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 358 mg of the title compound (64% yield). HRMS calcd for $C_{25}H_{24}O_5$ $[M+H]^+$ 405.1697, observed 405.1698.

20 **Step 4: 1-(3-Benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-2-methyl-propan-1-one**




1-[3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-phenyl]-2-methyl-propan-1-one (355 mg) was dissolved in glacial acetic acid (3 mL) and heated at 100 °C overnight. The solvent was removed under reduced pressure and the residue was

25 purified on a silica gel column using ethyl acetate and hexanes to yield 205 mg of

- 112 -


orange solid, which was dissolved in small amount of ethyl acetate and triturated with hexanes to yield 112 mg of the title compound (45% yield). HRMS calcd for $C_{17}H_{16}O_4$ $[M+H]^+$ 281.1122, observed 281.1121.

5 **Step 5: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-isobutyryl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester**



To a solution of 1-(3-benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-2-methyl-propan-1-one (109 mg) in a mixture of acetone and DMF (2:1, 3 mL) were added cesium carbonate (624 mg) and 4-[3-(6-bromo-hexyl)-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (190 mg). The resulting mixture was stirred at 75 °C overnight. Then the insoluble material was filtered out and the filtrate was diluted with ethyl acetate and washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 194 mg of the title compound (75% yield). HRMS calcd for $C_{40}H_{50}O_9$ $[M+Na]^+$ 697.3347, observed 697.3344.



15 **Step 6: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-isobutyryl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid**



20 To a solution of 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-isobutyryl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester (191 mg) in ethanol (2 mL) was added 10 M NaOH (566 μ L) and the resulting mixture was stirred at room

- 113 -

temperature for 5 h. Then the reaction mixture was diluted with water, acidified with 3 N HCl and extracted into ethyl acetate. The organic extract was washed with brine, dried over anhydrous sodium sulfate, concentrated under reduced pressure and purified on a reverse-phase HPLC to yield 50 mg of the title compound (29% yield). HRMS calcd for C₃₆H₄₂O₉ [M+Na]⁺ 641.2721, observed 641.2720.


Example 71:**4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-propionyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid****Step 1: 1-[3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-phenyl]-propan-1-one**

To a solution of 3-benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-benzonitrile (250 mg) in anhydrous THF (3 mL) were added a solution of ethylmagnesium chloride (517 μ L of 2 M solution in diethyl ether) and catalytic amount of cuprous bromide (2 mg). The resulting reaction mixture was refluxed for 1 h and then additional amount of ethylmagnesium chloride (517 μ L of 2 M solution in diethyl ether) was added and heating was continued for 30 min. After it was cooled down, 10% sulfuric acid (1.5 mL) was added and stirring was continued for 30 min at room temperature. Then the reaction mixture was diluted with diethyl ether, washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 136 mg

- 114 -

of the title compound (50% yield). HRMS calcd for $C_{24}H_{22}O_5$ $[M+H]^+$ 391.1540, observed 391.1541.

Step 2: 1-(3-Benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-propan-1-one




5

1-[3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-phenyl]-propan-1-one (135 mg) was dissolved in glacial acetic acid (2 mL) and heated at 100 °C for 2 days. The solvent was removed under reduced pressure and the residue was purified on a silica gel column using ethyl acetate and hexanes to yield 28.5 mg of the title compound (30% yield). HRMS calcd for $C_{16}H_{14}O_4$ $[M+H]^+$ 271.0965, observed 271.0964.

10

Step 3: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-propionyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester




15

The title compound was prepared by the same method described in **Example 70, Step 5**. Starting with 28.5 mg of 1-(3-benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-propan-1-one, 44 mg (66% yield) of the title compound were obtained. HRMS calcd for $C_{39}H_{48}O_9$ $[M+Na]^+$ 683.3190, observed 683.3191.

20

Step 4: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-propionyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

- 115 -



The title compound was prepared by the same method described in **Example 70**,


Step 6. Starting with 41 mg of 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-propionyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester, 34

5 mg (94% yield) of the title compound were obtained. HRMS calcd for $C_{35}H_{40}O_9$ $[M+Na]^+$ 627.2564, observed 627.2566.


Example 72:

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopropanecarbonyl-phenoxy)-hexyl]-2-

10 **(2-carboxy-ethyl)-phenoxy]-butyric acid**



Step 1: [3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-phenyl]-cyclopropyl-methanone




15 To a solution of 3-benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-benzonitrile (250 mg) in anhydrous THF (3 mL) were added a solution of cyclopropylmagnesium chloride (4.1 mL of 0.5 M solution in THF) and catalytic amount of cuprous bromide (2 mg). The resulting reaction mixture was refluxed for 1.5 h. After it was cooled

- 116 -


down, 10% sulfuric acid (1.5 mL) was added and stirring was continued for 30 min at room temperature. Then the reaction mixture was diluted with diethyl ether, washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 253 mg of the title compound (91% yield). HRMS calcd for $C_{25}H_{22}O_5$ $[M+H]^+$ 403.1540, observed 403.1537.

Step 2: (3-Benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-cyclopropyl-methanone



- 10 [3-Benzo[1,3]dioxol-5-yl-5-(4-methoxy-benzyloxy)-phenyl]-cyclopropyl-methanone (250 mg) was dissolved in glacial acetic acid (3 mL) and heated at 100 °C overnight. The solvent was removed under reduced pressure and the residue was purified on a silica gel column using ethyl acetate and hexanes to yield 138 mg of the title compound (79% yield). HRMS calcd for $C_{17}H_{14}O_4$ $[M+H]^+$ 283.0965, 15 observed 283.0964.


Step 3: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopropanecarbonyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



- 20 The title compound was prepared by the same method described in **Example 70**, **Step 5**. Starting with 35 mg of (3-benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-cyclopropyl-methanone, 74 mg (92% yield) of the title compound were obtained. HRMS calcd for $C_{40}H_{48}O_9$ $[M+Na]^+$ 695.3190, observed 695.3191.

- 117 -

Step 4: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopropanecarbonyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid




The title compound was prepared by the same method described in **Example 70**,

- 5 **Step 6.** Starting with 70 mg of 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-cyclopropanecarbonyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester, 48 mg (71% yield) of the title compound were obtained. HRMS calcd for $C_{36}H_{40}O_9$ $[M+H]^+$ 617.2745, observed 617.2745.


10

Example 73:

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopentanecarbonyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid



Step 1: Cyclopentyl-(3,5-dimethoxy-phenyl)-methanone




15


To a solution of 3,5-dimethoxybenzonitrile (5.0 g) in anhydrous THF (60 mL) were added a solution of cyclopentylmagnesium chloride (20 mL of 2 M solution in diethyl ether) and cuprous bromide (88 mg). The resulting reaction mixture was refluxed for 1 h. Then water (10 mL) and 10% sulfuric acid (20 mL) were added

- 118 -

and the resulting mixture was stirred at room temperature overnight. Then the reaction mixture was diluted with diethyl ether and washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 5.9 of the title compound (82% yield). HRMS calcd for $C_{14}H_{18}O_3$ $[M+H]^+$ 253.1329, observed 253.1328.

Step 2: Cyclopentyl-(3,5-dihydroxy-phenyl)-methanone

To a solution of cyclopentyl-(3,5-dimethoxy-phenyl)-methanone (234 mg) in anhydrous dichloromethane (5 mL) cooled in dry ice / acetone bath was added dropwise a solution of boron tribromide (4 mL of 1 M solution in dichloromethane). After the addition was completed, the cooling bath was removed and the reaction mixture was stirred at room temperature overnight. Unreacted boron tribromide was quenched with methanol. Then the reaction mixture was diluted with ethyl acetate and washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 200 mg of the title compound (97% yield). Note: the title compound decomposes on air. HRMS calcd for $C_{12}H_{14}O_3$ $[M+H]^+$ 207.1016, observed 207.1016.

Step 3: Trifluoro-methanesulfonic acid 3-cyclopentanecarbonyl-5-trifluoromethanesulfonyloxy-phenyl ester


Cyclopentyl-(3,5-dihydroxy-phenyl)-methanone (1.9 g) was suspended in 40 mL of dichloromethane and then pyridine (3.0 mL) was added. The resulting clear

- 119 -

solution was cooled to 0 °C and 3.2 mL of triflic anhydride were added dropwise. After addition was completed, the cooling bath was removed and the reaction mixture was stirred at room temperature for 1 h and then quenched with 1 N hydrochloric acid. The organic solution was separated and washed with saturated sodium bicarbonate solution and brine and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure and the residue was purified on a silica gel column using ethyl acetate and hexanes to yield 1.5 g of the title compound (34% yield). HRMS calcd for C₁₄H₁₂O₇S₂F₆ [M-H]⁻ 468.9856, observed 468.9854.

10

Step 4: Trifluoro-methanesulfonic acid 3-cyclopentanecarbonyl-5-hydroxy-phenyl ester




Trifluoro-methanesulfonic acid 3-cyclopentanecarbonyl-5-

15 trifluoromethanesulfonyloxy-phenyl ester (1.5 g) was dissolved in 1,2-dimethoxyethane (30 mL), followed by addition of cesium carbonate (1.5 g). The resulting suspension was stirred at 80 °C for 3 h and then quenched with saturated solution of ammonium chloride. The solution was then extracted with diethyl ether and combined etherate extracts were washed with brine and dried over anhydrous sodium sulfate. The filtrate was concentrated under reduced pressure and purified on a silica gel column using ethyl acetate and hexanes to yield 300 mg of the title compound (28% yield). HRMS calcd for C₁₃H₁₃O₅S₁F₃ [M+H]⁺ 339.0509, observed 339.0509.


25 **Step 5: (3-Benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-cyclopentyl-methanone**

- 120 -



To a solution of trifluoro-methanesulfonic acid 3-cyclopentanecarbonyl-5-hydroxy-phenyl ester (300 mg) in 1,2-dimethoxyethane (4 mL) were added 3,4-metylenedioxyphenylboronic acid (295 mg), 2 M aq. sodium carbonate solution (0.9 mL) and Pd(PPh₃)₄ (10 mg) and the resulting mixture was stirred at 90 °C for 3 h. Then the reaction mixture was diluted with ethyl acetate, filtered through Celite and washed with water and brine. The organic extract was dried over anhydrous sodium sulfate, concentrated and purified on a silica gel column using ethyl acetate and hexanes to yield 100 mg of the title compound (36% yield). HRMS calcd for C₁₉H₁₈O₄ [M+H]⁺ 311.1278, observed 311.1277.


Step 6: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopentanecarbonyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester



The title compound was prepared by the same method described in **Example 70, Step 5**. Starting with 94 mg of (3-benzo[1,3]dioxol-5-yl-5-hydroxy-phenyl)-cyclopentyl-methanone, 98 mg (47% yield) of the title compound were obtained. HRMS calcd for C₄₂H₅₂O₉ [M+Na]⁺ 723.3503, observed 723.3509.

Step 7: 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopentanecarbonyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid

- 121 -



The title compound was prepared by the same method described in **Example 70, Step 6**. Starting with 97 mg of 4-[3-[6-(3-benzo[1,3]dioxol-5-yl-5-cyclopentanecarbonyl-phenoxy)-hexyl]-2-(2-ethoxycarbonyl-ethyl)-phenoxy]-butyric acid ethyl ester, 3.9 mg (4% yield) of the title compound were obtained. HRMS calcd for $C_{38}H_{44}O_9$ $[M+H]^+$ 645.3058, observed 645.3056.

Example 74

Assay of compounds for inhibition of LTB_4 activity.

Ca^{2+} Flux Assay for LTB_4 Antagonist Assay

10 Cell Culture Conditions:

Human leukemia HL-60 cells endogenously expressing BLT1 and BLT2 receptors were cultured in RPMI-1640 medium supplemented with 20% fetal bovine serum, 2 mM glutamine, 100U/ml penicillin and 100ug/ml streptomycin.

Seventy two hours prior to experiment cells are counted using ViaCount reagent, centrifuged and resuspended at 2.0×10^5 cells /ml density with the complete growth media containing 1 μ M Retinoic Acid (Sigma).

Dye Loading and Assay:

On a day of the experiment loading buffer (Calcium-3 Assay Kit, Molecular Devices) was prepared by dissolving the contents of one vial (Express Kit) into 500 ml Hank's Balanced Salt Solution containing 20 mM HEPES and 5 mM probenecid. Equal volume of the loading buffer was mixed with the replacement buffer (Hank's Balanced Salt Solution containing 20 mM HEPES, 0.05%BSA and 5mM probenecid). Retinoic Acid induced HL-60 cells were counted using ViaCount reagent, centrifuged and resuspended at 2.0×10^6 cells /ml density with the loading

- 122 -

buffer/replacement buffer, dispensed into 384 well black/clear microplates (Falcon)(25µl/well) and placed in a 37°C/5% CO₂ incubator for 1 hour.

During the incubation, test compounds were prepared at 6X the desired

- 5 concentration in HBSS/20mM HEPES/0.05%BSA as well as LTB₄ (Biomol) was prepared at 2.2X concentration in HBSS/20mM HEPES/0.5%BSA buffer.

After the incubation, both the cell and compound plates were brought to the FLIPR and 5µl of the diluted compounds were transferred to the cell plates by the FLIPR.

- 10 Plates were then incubated for 30 min at room temperature. After the ½ hour incubation, plates were returned to the FLIPR and 25µl of 2.2X LTB₄ was added to the cell plates. During the assay, fluorescence readings were taken simultaneously from all 384 wells of the cell plate every 1.5 seconds. Five readings were taken to establish a stable baseline, then 25µl (LTB₄) of sample was rapidly and
- 15 simultaneously added to each well of the cell plate. The fluorescence was continuously monitored before, during and after sample addition for a total elapsed time of 100 seconds. Responses (increase in peak fluorescence) in each well following agonist addition was determined. The initial fluorescence reading from each well, prior to ligand stimulation, was used as a zero baseline value for the data
- 20 from that well. The responses are expressed as % inhibition of the neutral control (neural control: wells that received buffer plus DMSO but no test compound).

Assay results

Example #	Compound name	LTB ₄ antagonism (HL-60 FLIPR) (nM)
1	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	3.1
2	4-{2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	24.68

- 123 -

3	4-{2-(2-Carboxy-ethyl)-3-[6-(3-hydroxymethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	62.46
4	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-fluoro-5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	29.63
5	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	43.08
6	4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-hydroxymethyl-phenoxy]-hexyl}-phenoxy)-butyric acid	7.39
7	4-{2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	83.21
8	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl)-5-methoxymethyl-phenoxy]-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.14
9	4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	1.93
10	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	2.52

11	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	1.37
12	4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-methoxymethyl-phenoxy]-hexyl}-phenoxy)-butyric acid	1.38
13	4-(2-(2-Carboxy-ethyl)-3-{6-[3-methoxymethyl-5-(4-methyl-thiophen-3-yl)-phenoxy]-hexyl}-phenoxy)-butyric acid	140.16
14	4-{2-(2-Carboxy-ethyl)-3-[6-(3'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	1.47
15	4-{2-(2-Carboxy-ethyl)-3-[6-(2'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	5.98
16	4-{2-(2-Carboxy-ethyl)-3-[6-(3',5'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	14.85
17	4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	1.62
18	4-{2-(2-Carboxy-ethyl)-3-[6-(3'-chloro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-	13.38

	butyric acid	
19	4-{2-(2-Carboxy-ethyl)-3-[6-(3'-methoxy-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	9.44
20	4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-4'-trifluoromethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	15.14
21	4-{2-(2-Carboxy-ethyl)-3-[6-(3',4'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	2.66
22	4-{2-(2-Carboxy-ethyl)-3-[6-(3-methoxymethyl-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	339.3
23	4-(2-(2-Carboxy-ethyl)-3-{6-[3-(1 <i>H</i> -indol-5-yl)-5-methoxymethyl-phenoxy]-hexyl}-phenoxy)-butyric acid	30.19
24	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	6.83
25	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	1.06
26	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	11.88

- 126 -

27	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	6.24
28	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	6.51
29	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	15.96
30	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-dimethylaminomethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.1
31	4-{2-(2-Carboxy-ethyl)-3-[6-(3-dimethylaminomethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	0.11
32	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	0.1
33	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	0.11
34	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-dimethylaminomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	0.24

- 127 -

	acid	
35	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-fluoro-4'-methyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	0.16
36	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	0.14
37	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	0.37
38	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	2.22
39	4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	70.81
40	4-{2-(2-Carboxy-ethyl)-3-[6-(3-dimethylaminomethyl-5-pyridin-4-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	0.42
41	4-{2-(2-Carboxy-ethyl)-3-[6-(3-cyanomethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	15.19
42	4-{2-(2-Carboxy-ethyl)-3-[6-(5-cyanomethyl-biphenyl-3-yloxy)-	22.98

	hexyl]-phenoxy)- butyric acid	
43	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.22
44	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	1.26
45	4-{2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	0.4
46	4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy)-butyric acid	1.11
47	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	2.32
48	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	5.2
49	4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	6.05
50	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-3',4'-difluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	8.27
51	4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-	2.1

- 129 -

	biphenyl-3-yloxy)-hexyl]-phenoxy)-butyric acid	
52	4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy)-butyric acid	6.94
53	4-{2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy)-butyric acid	79.31
54	4-(2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1 <i>H</i> -indol-5-yl)-phenoxy]-hexyl]-phenoxy)-butyric acid	34.16
55	4-(2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1 <i>H</i> -indol-6-yl)-phenoxy]-hexyl]-phenoxy)-butyric acid	6.69
56	4-[3-[6-(3-Benzothiazol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.32
57	4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl]-phenoxy)-butyric acid	1.2
58	4-[3-[6-(5-Benzyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	74.58
59	4-[3-[6-(5-Benzyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	131.84

- 130 -

	acid	
60	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-benzyloxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	4.59
61	4-[3-[6-(3-Benzyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	77.67
62	4-{2-(2-Carboxy-ethyl)-3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid	7.98
63	4-{2-(2-Carboxy-ethyl)-3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid	32.52
64	4-[3-[6-(3-Acetyl-5-benzo[1,3]dioxol-5-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	2.95
65	4-[3-{6-[3-Acetyl-5-(2,3-dihydrobenzo[1,4]dioxin-6-yl)-phenoxy]-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	2.07
66	4-[3-[6-(5-Acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	3.33
67	4-[3-[6-(5-Acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	5.1
68	4-[3-[6-(3-Acetyl-5-thiophen-3-yl-	2

- 131 -


	phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	
69	4-[3-{6-[3-Acetyl-5-(2,2-difluorobenzo[1,3]dioxol-5-yl)-phenoxy]-hexyl}-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.9
70	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-isobutyryl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.31
71	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-propionyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.4
72	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopropanecarboxyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.16
73	4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopentanecarboxyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid	0.77

It is to be understood that the invention is not limited to the particular embodiments of the invention described above, as variations of the particular embodiments may be made and still fall within the scope of the appended claims.

- 132 -

Claims

1. A compound of formula (I)



5 wherein:

R1 is lower alkyl, alkoxy, alkoyl, dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl; and

10 R2 is benzo[1,3]dioxol, dihydro-benzo[1,4]dioxin, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl,

15 or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, wherein:

R1 is lower alkyl, alkoxy, alkoyl, dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl; and

20 R2 is benzo[1,3]dioxol, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl,

25 or a pharmaceutically acceptable salt thereof.

- 133 -

3. The compound according to claim 1, wherein:

R1 is lower alkyl, alkoxy or alkoyl; and

R2 is benzo[1,3]dioxol, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl.

4. The compound according to claim 1, wherein:

R1 is dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl; and

R2 is benzo[1,3]dioxol, benzo[1,4]dioxin, benzothiazole, difluoro-benzo[1,3]dioxole, cycloalkyl, aryl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl, heteroaryl, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy, N-aryl or indole, unsubstituted or substituted with lower alkyl.

5. The compound according to claim 1, wherein R1 is hydroxymethyl, methoxymethyl, ethoxy, ethoxymethyl, benzyloxy, acetyl or propionyl.

6. The compound according to claim 1, wherein R1 is dimethylaminomethyl, cyanomethyl, benzyloxy, cyclopentylmethoxy, cyclopropanecarbonyl or cyclopentanecarbonyl.

7. The compound according to claim 1, wherein said aryl at R2 is phenyl, unsubstituted or mono-, di- or tri-substituted with halogen, hydroxy, alkoxy, lower alkyl, -CF₃, -OCF₃ or methanesulfonyl.

8. The compound according to claim 1, wherein said heteroaryl at R2 is thiophene, pyrimidine or pyridine, unsubstituted or mono-, di- or tri-substituted with lower alkyl, halogen or hydroxy.

9. The compound according to claim 1, wherein said compound is:

- 134 -

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-hydroxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,

5 4-{2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3-hydroxymethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid,

10 4-{2-(2-Carboxy-ethyl)-3-[6-(4'-fluoro-5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-hydroxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

15

4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-hydroxymethyl-phenoxy]-hexyl}-phenoxy)-butyric acid,

20 4-{2-(2-Carboxy-ethyl)-3-[6-(5-hydroxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-methoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,

25 4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

30

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

- 135 -

4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-methoxymethyl-phenoxy]-hexyl}-phenoxy)-butyric acid,

4-(2-(2-Carboxy-ethyl)-3-{6-[3-methoxymethyl-5-(4-methyl-thiophen-3-yl)-
5 phenoxy]-hexyl}-phenoxy)-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-
phenoxy}-butyric acid,

10 4-{2-(2-Carboxy-ethyl)-3-[6-(2'-fluoro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-
phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3',5'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-
hexyl]-phenoxy}-butyric acid,

15

4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-
hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3'-chloro-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-
20 phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3'-methoxy-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-
phenoxy}-butyric acid or

25 4-{2-(2-Carboxy-ethyl)-3-[6-(5-methoxymethyl-4'-trifluoromethoxy-biphenyl-3-
yloxy)-hexyl]-phenoxy}-butyric acid.

10. The compound according to claim 1, wherein said compound is:

4-{2-(2-Carboxy-ethyl)-3-[6-(3',4'-difluoro-5-methoxymethyl-biphenyl-3-yloxy)-
30 hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3-methoxymethyl-5-pyrimidin-5-yl-phenoxy)-hexyl]-
phenoxy}-butyric acid,

- 4-(2-(2-Carboxy-ethyl)-3-{6-[3-(1*H*-indol-5-yl)-5-methoxymethyl-phenoxy]-hexyl}-phenoxy)-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(4'-methanesulfonyl-5-methoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 5 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxymethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-6-(5-ethoxymethyl-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 10 4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxymethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxymethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 15 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-dimethylaminomethyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(3-dimethylaminomethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 20 4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-dimethylaminomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 25 4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-fluoro-4'-methyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-trifluoromethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 30 4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(5-dimethylaminomethyl-3'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid or

4-{2-(2-Carboxy-ethyl)-3-[6-(3-dimethylaminomethyl-5-pyridin-4-yl-phenoxy)-hexyl]-phenoxy}-butyric acid.

5

11. The compound according to claim 1, wherein said compound is:

4-{2-(2-Carboxy-ethyl)-3-[6-(3-cyanomethyl-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(5-cyanomethyl-biphenyl-3-yloxy)-hexyl]-phenoxy}-
10 butyric acid,

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric
acid,

15 4-{2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-
butyric acid,

4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-5-ethoxy-phenoxy]-hexyl}-phenoxy)-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-
20 butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-
butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(4'-chloro-5-ethoxy-biphenyl-3-yloxy)-hexyl]-phenoxy}-
butyric acid,

25 4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-3',4'-difluoro-biphenyl-3-yloxy)-hexyl]-
phenoxy}-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(5-ethoxy-4'-methanesulfonyl-biphenyl-3-yloxy)-hexyl]-
phenoxy}-butyric acid,

4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2-chloro-pyridin-4-yl)-5-ethoxy-phenoxy]-hexyl}-
30 phenoxy)-butyric acid,

4-{2-(2-Carboxy-ethyl)-3-[6-(3-ethoxy-5-pyrimidin-5-yl-phenoxy)-hexyl]-phenoxy}-
butyric acid,

- 4-(2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-5-yl)-phenoxy]-hexyl}-phenoxy)-butyric acid,
- 4-(2-(2-Carboxy-ethyl)-3-{6-[3-ethoxy-5-(1*H*-indol-6-yl)-phenoxy]-hexyl}-phenoxy)-butyric acid,
- 5 4-[3-[6-(3-Benzothiazol-5-yl-5-ethoxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-(2-(2-Carboxy-ethyl)-3-{6-[3-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-5-ethoxy-phenoxy]-hexyl}-phenoxy)-butyric acid,
- 4-[3-[6-(5-Benzoyloxy-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric
- 10 acid,
- 4-[3-[6-(5-Benzoyloxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-benzoyloxy-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 15 4-[3-[6-(3-Benzoyloxy-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(3-cyclopentylmethoxy-5-thiophen-3-yl-phenoxy)-hexyl]-phenoxy}-butyric acid,
- 4-{2-(2-Carboxy-ethyl)-3-[6-(5-cyclopentylmethoxy-3'-fluoro-biphenyl-3-yloxy)-hexyl]-phenoxy}-butyric acid,
- 20 hexyl]-phenoxy}-butyric acid,
- 4-[3-[6-(3-Acetyl-5-benzo[1,3]dioxol-5-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-[3-[6-[3-Acetyl-5-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-phenoxy]-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 25 4-[3-[6-(5-Acetyl-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-[3-[6-(5-Acetyl-3'-fluoro-biphenyl-3-yloxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-[3-[6-(3-Acetyl-5-thiophen-3-yl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 30 4-[3-[6-[3-Acetyl-5-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-phenoxy]-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,
- 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-isobutyryl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,

- 139 -

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-propionyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,

4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopentanecarbonyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid,

- 5 4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopropanecarbonyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid or
4-[3-[6-(3-Benzo[1,3]dioxol-5-yl-5-cyclopentanecarbonyl-phenoxy)-hexyl]-2-(2-carboxy-ethyl)-phenoxy]-butyric acid.

- 10 12. A pharmaceutical composition, comprising a therapeutically effective amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

- 15 13. The use of a compound according to any one of the claims 1 to 11 for the preparation of a medicament for the treatment of severe asthma or chronic obstructive pulmonary disease.

14. Compound according to any one of the claims 1 to 11 for use in the treatment of severe asthma or chronic obstructive pulmonary disease.

20

15. The invention as described herein above.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/053580

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07C59/68 C07C59/90 A61P11/08 C07D209/08 C07D213/38
C07D213/61 C07D239/26 C07D277/24 C07D317/54 C07D317/58
C07D319/18 C07D333/16 C07D333/20 C07D333/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07C A61K A61P C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2009/024492 A1 (HOFFMANN LA ROCHE [CH]; DOMINIQUE ROMYR [US]; FOTOUHI NADER [US]; GILL) 26 February 2009 (2009-02-26) the whole document -----	1-6, 10-14

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 June 2010

Date of mailing of the international search report

24/06/2010

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Delanghe, Patrick

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2010/053580

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 15
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.2

Claims Nos.: 15

Claim 15 contains no technical features but a general reference to the description expressed as "The invention as described herein above". As a result, the claim is completely unclear and a meaningful search can not be done.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.2), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2010/053580

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009024492 A1	26-02-2009	CA 2696967 A1	26-02-2009
		EP 2190808 A1	02-06-2010
		US 2009054466 A1	26-02-2009