

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
14 January 2021 (14.01.2021)



(10) International Publication Number
WO 2021/007545 A1

(51) International Patent Classification:

A01N 43/22 (2006.01) *A01N 47/40* (2006.01)
A01N 43/40 (2006.01) *A01N 51/00* (2006.01)
A01N 43/56 (2006.01) *A01N 53/00* (2006.01)
A01N 47/06 (2006.01) *A01N 63/00* (2020.01)
A01N 47/38 (2006.01)

(21) International Application Number:

PCT/US2020/041668

(22) International Filing Date:

10 July 2020 (10.07.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/873,100 11 July 2019 (11.07.2019) US
62/873,302 12 July 2019 (12.07.2019) US

(71) Applicant: **FMC CORPORATION** [US/US]; Patent Dept., 2929 Walnut Street, Philadelphia, PA 19104 (US).

(72) Inventor: **ZHANG, Wenming**; FMC Corporation, 2929 Walnut Street, Philadelphia, PA 19104 (US).

(74) Agent: **EPELBAUM, Sabine U.**; FMC Corporation, Patent Dept., 2929 Walnut Street, Philadelphia, PA 19104 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO,


NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: MIXTURES COMPRISING INDAZOLE PESTICIDES



Formula (I)



Formula (II)

(57) Abstract: Disclosed are compositions comprising compounds of Formula (I) and compounds of Formula (II). Also disclosed are compositions containing compounds of Formula (I) or compounds of Formula (II), or combinations thereof and methods for controlling an invertebrate pest comprising contacting the invertebrate pest or its environment with a biologically effective amount of a compound or a composition of the disclosure.

WO 2021/007545 A1

TITLE

MIXTURES COMPRISING INDAZOLE PESTICIDES

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of U.S. provisional application number 62/873,100, filed July 11, 2019, and U.S. provisional application number 62/873,302, filed July 12, 2019, the disclosures of which are hereby incorporated by reference in their entireties.

FIELD


10 This disclosure relates to pesticidal mixtures and compositions comprising certain substituted indazole compounds, and at least one other invertebrate pest control agent, suitable for agronomic, nonagronomic and uses, and methods of their use for controlling invertebrate pests such as arthropods in both agronomic and nonagronomic environments.

BACKGROUND

15 The control of invertebrate pests is extremely important in achieving high crop efficiency. Damage by invertebrate pests to growing and stored agronomic crops can cause significant reduction in productivity and thereby result in increased costs to the consumer. The control of invertebrate pests in forestry, greenhouse crops, ornamentals, nursery crops, stored food and fiber products, livestock, household, turf, wood products, and public and animal health is also important. Many products are commercially available for these purposes,
20 but the need continues for new compounds that are more effective, less costly, less toxic, environmentally safer or have different sites of action. PCT Application Publication WO 2015/038503 A1 discloses related indazole compounds.


SUMMARY

25 This disclosure is directed to a composition comprising (a) at least one compound of Formula I or of Formula II, or a combination of the foregoing compounds, compositions containing them, and their use for controlling invertebrate pests:



Formula I

wherein n is 0, 1, or 2; or



Formula II

wherein R is CH₃, or CF₂H;

5 or a combination of the foregoing compounds; and

(b) at least one additional pest control agent different from the compound of Formula I or the compound of Formula II.

In some embodiments the additional pest control agent is an invertebrate pest control agent.

10 In some embodiments the additional pest control agent is selected from an insecticide, fungicide, nematocide, and bactericide.

In some embodiments the additional pest control agent is selected from abamectin, acephate, acequinocyl, acetamiprid, acrinathrin, acynonapyr, afidopyropen

15 [(3*S*,4*R*,4*aR*,6*S*,6*aS*,12*R*,12*aS*,12*bS*)-3-[(cyclopropylcarbonyl)oxy]-1,3,4,4*a*,5,6,6*a*,12,12*a*,12*b*-decahydro-6,12-dihydroxy-4,6*a*,12*b*-trimethyl-11-oxo-9-(3-pyridinyl)-2*H*,11*H*-naphtho[2,1-*b*]pyrano[3,4-*e*]pyran-4-yl)methyl

cyclopropanecarboxylate), amidoflumet, amitraz, avermectin, azadirachtin, azinphos-methyl, benfuracarb, bensultap, benzpyrimoxan, bifenthrin, kappa-bifenthrin, bifenazate, bistrifluron, borate, broflanilide, buprofezin, cadusafos, carbaryl, carbofuran, cartap, carzol, chlorantraniliprole, chlorfenapyr, chlorfluazuron, chlorprallethrin, chlorpyrifos, chlorpyrifos-*e*, chlorpyrifos-methyl, chromafenozide, clofentezin, chlorprallethrin, clothianidin, cyantraniliprole (CyazypyrTM) (3-bromo-1-(3-chloro-2-pyridinyl)-*N*-[4-cyano-2-methyl-6-[(methylamino)carbonyl]phenyl]-1*H*-pyrazole-5-carboxamide), cyclaniliprole (3-bromo-*N*-[2-bromo-4-chloro-6-[(1-cyclopropylethyl)amino]carbonyl]phenyl]-1-(3-

25 chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxamide), cyclobutrifluram, cycloprothrin, cycloxaprid ((5*S*,8*R*)-1-[(6-chloro-3-pyridinyl)methyl]-2,3,5,6,7,8-hexahydro-9-nitro-5,8-Epoxy-1*H*-imidazo[1,2-*a*]azepine), cyenopyrafen, cyetpyrafen, cyflumetofen, cyfluthrin, beta-cyfluthrin, cyhalodiamide, cyhalothrin, gamma-cyhalothrin, lambda-cyhalothrin, cypermethrin, alpha-cypermethrin, zeta-cypermethrin, cyromazine, deltamethrin,

diafenthiuron, diazinon, dicloromezotiaz, dieldrin, diflubenzuron, dimefluthrin, dimehypo, dimethoate, dimpropridaz, dinotefuran, diofenolan, DiPel®emamectin, emamectin benzoate, endosulfan, esfenvalerate, ethiprole, etofenprox, epsilon-metofluthrin, etoxazole, fenbutatin oxide, fenitrothion, fenothiocarb, fenoxycarb, fenpropathrin, fenvalerate, fipronil,

5 flometoquin (2-ethyl-3,7-dimethyl-6-[4-(trifluoromethoxy)phenoxy]-4-quinolinyl methyl carbonate), flonicamid, fluazaindolizine, flubendiamide, flucythrinate, flufenerim, flufenoxuron, flufenoxystrobin (methyl (αE)-2-[[2-chloro-4-(trifluoromethyl)phenoxy]methyl]- α -(methoxymethylene)benzeneacetate), fluensulfone (5-chloro-2-[(3,4,4-trifluoro-3-buten-1-yl)sulfonyl]thiazole), fluhexafon, fluopyram,

10 flupentiofenox, flupiprole (1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-5-[(2-methyl-2-propen-1-yl)amino]-4-[(trifluoromethyl)sulfinyl]-1*H*-pyrazole-3-carbonitrile), flupyradifurone (4-[[6-chloro-3-pyridinyl)methyl](2,2-difluoroethyl)amino]-2(5*H*)-furanone), flupyrimin, fluvalinate, tau-fluvalinate, fluxametamide, fonophos, formetanate, fosthiazate, gamma-cyhalothrin, halofenozide, heptafluthrin ([2,3,5,6-tetrafluoro-4-

15 (methoxymethyl)phenyl]methyl 2,2-dimethyl-3-[(1*Z*)-3,3,3-trifluoro-1-propen-1-yl]cyclopropanecarboxylate), hexaflumuron, hexythiazox, hydramethylnon, imidacloprid, indoxacarb, insecticidal soaps, isofenphos, isocycloseram, kappa-tefluthrin, lambda-cyhalothrin, lufenuron, malathion, meperfluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl

20 (1*R*,3*S*)-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate), metaflumizone, metaldehyde, methamidophos, methidathion, methiocarb, methomyl, methoprene, methoxychlor, metofluthrin, methoxyfenozide, epsilon-metofluthrin, metronidazole epsilon-momfluorothrin, monocrotophos, monofluorothrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 3-(2-cyano-1-propen-1-yl)-2,2-dimethylcyclopropanecarboxylate), nicofluprole, nicotine,

25 nitenpyram, nithiazine, novaluron, noviflumuron, oxamyl, oxazosulfil, parathion, parathion-methyl, permethrin, phorate, phosalone, phosmet, phosphamidon, pirimicarb, profenofos, profluthrin, propargite, protrifenbute, pyflubumide (1,3,5-trimethyl-*N*-(2-methyl-1-oxopropyl)-*N*-[3-(2-methylpropyl)-4-[2,2,2-trifluoro-1-methoxy-1-(trifluoromethyl)ethyl]phenyl]-1*H*-pyrazole-4-carboxamide), pymetrozine, pyrafluprole,

30 pyrethrin, pyridaben, pyridalyl, pyrifluquinazon, pyriminostrobin (methyl (αE)-2-[[[2-[(2,4-dichlorophenyl)amino]-6-(trifluoromethyl)-4-pyrimidinyl]oxy]methyl]- α -(methoxymethylene)benzeneacetate), pydiflumetofen, pyriprole, pyriproxyfen, rotenone, ryanodine, silafluofen, spinetoram, spinosad, spirodiclofen, spiromesifen, spiropidion, spirotetramat, sulprofos, sulfoxaflo (N-[methyloxido[1-[6-(trifluoromethyl)-3-pyridinyl]ethyl]- λ^4 -sulfanylidene]cyanamide), tebufenozide, tebufenpyrad, teflubenzuron,

35 tefluthrin, kappa-tefluthrin, terbufos, tetrachlorantraniliprole, tetrachlorvinphos, tetramethrin, tetramethylfluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 2,2,3,3-tetramethylcyclopropanecarboxylate), tetraniliprole, thiacloprid, thiamethoxam, thiodicarb,

thiosultap-sodium, tiozazafen (3-phenyl-5-(2-thienyl)-1,2,4-oxadiazole), tolfenpyrad, tralomethrin, triazamate, trichlorfon, triflumezopyrim (2,4-dioxo-1-(5-pyrimidinylmethyl)-3-[3-(trifluoromethyl)phenyl]-2*H*-pyrido[1,2-*a*]pyrimidinium inner salt), triflumuron, tyclopyrazoflor, zeta-cypermethrin, *Bacillus thuringiensis* delta-endotoxins, entomopathogenic bacteria, entomopathogenic viruses, and entomopathogenic fungi..

In some embodiments the additional pest control agent is selected from cyantraniliprole, acetamiprid, imidacloprid, spirotetramat, spirodiclofen, chlorantraniliprole, bifenthrin, indoxacarb, avermectin, *Bacillus* spp. and any active crystal proteins, buprofezin, carbofuran, chlorfenapyr, chlorpyrifos, clothianidin, cyromazine, diafenthiuron, dinotefuran, emamectin benzoate, fipronil, flonicamid, fluhexafon, flupyradifurone, methomyl, methoxyfenozide, metronidazole, novaluron, permethrin, pyriproxifen, sulfoxaflor, thiamethoxam, γ -cyhalothrin, and ζ -cypermethrin.

In some embodiments the additional pest control agent is selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

In some embodiments, the disclosure is directed to aforesaid composition wherein component (a) is at least one compound of Formula I, or at least one compound of Formula II, or a combination of the foregoing compounds of Formula I and Formula II.

In some embodiments, the disclosure provides a composition comprising any of the compositions described above and at least one additional component selected from surfactants, solid diluent, and liquid diluents.

In one embodiment, this disclosure also provides a composition for controlling an invertebrate pest comprising the compositions disclosed herein and at least one additional component selected from surfactants, solid diluents, and liquid diluents, said composition further comprising at least one additional biologically active compound or agent.

In one embodiment, this disclosure is also directed to compositions comprising at least one compound of Formula I or at least one compound of Formula II selected from:

N-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, and *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

In one embodiment, this disclosure is also directed to compositions comprising at least one compound of Formula I selected from:




N-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, and *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.



5 In one embodiment, this disclosure is also directed to compositions comprising at least one compound of Formula **II** selected from:

N-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, or *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

10 In one embodiment, the compounds of Formula **I** and Formula **II** are those wherein the compound is selected from the compounds in index Table 1.

Table 1.

Compound	Compound Structure	Chemical name
1		<i>N</i> -[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2 <i>H</i> -indazole-4-carboxamide
2		<i>N</i> -[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2 <i>H</i> -indazole-4-carboxamide
3		<i>N</i> -[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2 <i>H</i> -indazole-4-carboxamide

4		N-(1-methylcyclopropyl)-2-(3-pyridinyl)-2H-indazole-4-carboxamide
5		N-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2H-indazole-4-carboxamide

or combinations of compounds 1-5.

In one embodiment, the present disclosure also provides a composition comprising a compound of Formula **I** or a compound of Formula **II**, or a combination of any of the foregoing compounds.

In one embodiment, this disclosure also provides a composition comprising a compound of Formula **I** or a compound of Formula **II**, and at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents.

In one embodiment, this disclosure also provides a composition for controlling an invertebrate pest comprising compound of Formula **I** or a compound of Formula **II**, and at least one additional component selected from surfactants, solid diluents and liquid diluents, said composition optionally further comprising at least one additional biologically active compound or agent.

In one embodiment, this disclosure also provides a composition for controlling an invertebrate pest comprising compound of Formula **I** or a compound of Formula **II**, or a combination of the foregoing compounds, and at least one additional biologically active compound or pest control agent.

In one embodiment, this disclosure also provides a composition for controlling an invertebrate pest comprising compound of Formula **I**, or a combination of the foregoing compounds and at least one additional biologically active compound or pest control agent.

In one embodiment, this disclosure also provides a composition for controlling an invertebrate pest comprising compound of Formula **II** and at least one additional biologically active compound or pest control agent.

In one embodiment, this disclosure provides a method for controlling an invertebrate pest comprising contacting the invertebrate pest or its environment with a biologically effective amount of a compound of Formula **I** or a compound of Formula **II** (e.g., as a composition described herein). This disclosure also relates to such method wherein the invertebrate pest or its environment is contacted with a composition comprising a biologically effective amount of a compound of Formula **I** or a compound of Formula **II**, and at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents, said composition optionally further comprising a biologically effective amount of at least one additional biologically active compound or agent.

In one embodiment, this disclosure also relates to such method wherein the invertebrate pest or its environment is contacted with a composition comprising a biologically effective amount of a compound of Formula **I** or a compound of Formula **II**, and at least one additional biologically active compound or pest control agent.

In one embodiment, this disclosure also relates to such method wherein the invertebrate pest or its environment is contacted with a composition comprising a biologically effective amount of a compound of Formula **I** or a compound of Formula **II**, at least one additional biologically active compound or pest control agent and at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents. In some embodiments, the environment is soil or plant foliage.

In one embodiment, this disclosure also provides a method for protecting a seed from an invertebrate pest comprising contacting the seed with a biologically effective amount of a compound of Formula **I** or a compound of Formula **II** or a combination of the foregoing compounds.

In one embodiment, this disclosure also provides a method for protecting a seed from an invertebrate pest comprising contacting the seed with a biologically effective amount of a compound of Formula **I** or a compound of Formula **II**, or a combination of the foregoing compounds and at least one additional biologically active compound or pest control agent.

In one embodiment, this disclosure also provides a method for protecting a seed from an invertebrate pest comprising contacting the seed with a biologically effective amount of a compound of Formula **I** or a compound of Formula **II**, or a combination of the foregoing compounds, at least one additional biologically active compound or pest control agent, and at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents.

In some embodiments, the invertebrate pest is selected from the order Lepidoptera, Blattodea, Coleoptera, Dermaptera, Hemiptera, Homoptera, Acari, Orthoptera, Diptera, Thysanoptera, Hymenoptera, Isoptera, Thysanura, Mallophaga, Siphonoptera, Araneae, and Scutigromorpha.

In some embodiments, the invertebrate pest is selected from the order Lepidoptera, Thrips and Hemiptera.

In some embodiments, the invertebrate pest is selected from silverleaf whitefly, corn plant hopper, western flower thrips, potato leafhopper, cotton melon aphid, root knot
5 nematode, fall army worm and diamondback moth.

In one embodiment, the compositions as disclosed herein further comprise liquid fertilizer. In some embodiments, the liquid fertilizer is aqueous-based.

In one embodiment, this disclosure provides a soil drench formulation comprising the compositions disclosed herein.

10 In one embodiment, this disclosure provides a spray composition comprising the compositions disclosed herein. In some embodiments the spray composition further comprises a propellant.

In one embodiment, this disclosure provides a bait composition comprising the composition disclosed herein. In one embodiment, the bait composition further comprises one
15 or more food materials. In one embodiment, the bait composition further comprises an attractant. In one embodiment, the bait composition further comprises a humectant.

In one embodiment, the compositions disclosed herein are solid compositions, such as dusts, powders, granules, pellets, prills, pastilles, tablets, or filled films. In some
20 embodiments, the compositions disclosed herein are solid compositions and are water-dispersible or water-soluble.

In one embodiment, a liquid or dry formulation comprising the compositions as disclosed herein for use in a drip irrigation system, furrow during planting, handheld sprayer, backpack sprayer, boom sprayer, ground sprayer, aerial application, unmanned
aerial vehicle, or a seed treatment.

25 In one embodiment, the compositions as disclosed herein for use in a drip irrigation system, furrow during planting, handheld sprayer, backpack sprayer, boom sprayer, ground sprayer, aerial application, unmanned aerial vehicle, or a seed treatment wherein said formulation is sprayed at an ultra-low volume.

In one embodiment, this disclosure also relates to the treated seed.
30

DETAILED DESCRIPTION

As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains”, “containing,” “characterized by” or any other variation thereof, are intended to cover a non-exclusive inclusion, subject to any limitation explicitly indicated. For
35 example, a composition, mixture, process or method that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such composition, mixture, process or method.

The transitional phrase “consisting of” excludes any element, step, or ingredient not specified. If in the claim, such would close the claim to the inclusion of materials other than those recited except for impurities ordinarily associated therewith. When the phrase “consisting of” appears in a clause of the body of a claim, rather than immediately following the preamble, it limits only the element set forth in that clause; other elements are not excluded from the claim as a whole.

The transitional phrase “consisting essentially of” is used to define a composition or method that includes materials, steps, features, components, or elements, in addition to those literally disclosed, provided that these additional materials, steps, features, components, or elements do not materially affect the basic and novel characteristic(s) of the claims. The term “consisting essentially of” occupies a middle ground between “comprising” and “consisting of”.

Where applicants have defined an embodiment or a portion thereof with an open-ended term such as “comprising,” it should be readily understood that (unless otherwise stated) the description should be interpreted to also describe such an embodiment using the terms “consisting essentially of” or “consisting of.”

Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

Also, the indefinite articles “a” and “an” preceding an element or component of the disclosure are intended to be nonrestrictive regarding the number of instances (i.e. occurrences) of the element or component. Therefore “a” or “an” should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

As referred to in this disclosure, the term “invertebrate pest” includes arthropods, gastropods, nematodes and helminths of economic importance as pests. The term “arthropod” includes insects, mites, spiders, scorpions, centipedes, millipedes, pill bugs and symphylans. The term “gastropod” includes snails, slugs and other Stylommatophora. The term “nematode” includes members of the phylum Nematoda, such as phytophagous nematodes and helminth nematodes parasitizing animals. The term “helminth” includes all of the parasitic worms, such as roundworms (phylum Nematoda), heartworms (phylum Nematoda, class Secernentea), flukes (phylum Platyhelminthes, class Tematoda), acanthocephalans (phylum Acanthocephala), and tapeworms (phylum Platyhelminthes, class Cestoda).

In the context of this disclosure “invertebrate pest control” means inhibition of invertebrate pest development (including mortality, feeding reduction, and/or mating disruption), and related expressions are defined analogously.

The term “agronomic” refers to the production of field crops such as for food and fiber and includes the growth of maize or corn, soybeans and other legumes, rice, cereal (e.g., wheat, oats, barley, rye and rice), leafy vegetables (e.g., lettuce, cabbage, and other cole crops), fruiting vegetables (e.g., tomatoes, pepper, eggplant, crucifers and cucurbits), potatoes, sweet potatoes, grapes, cotton, tree fruits (e.g., pome, stone and citrus), small fruit (e.g., berries and cherries) and other specialty crops (e.g., canola, sunflower and olives).

The term “nonagronomic” refers to other than field crops, such as horticultural crops (e.g., greenhouse, nursery or ornamental plants not grown in a field), residential, agricultural, commercial and industrial structures, turf (e.g., sod farm, pasture, golf course, lawn, sports field, etc.), wood products, stored product, agro-forestry and vegetation management, public health (i.e. human) and animal health (e.g., domesticated animals such as pets, livestock and poultry, undomesticated animals such as wildlife) applications.

The term “crop vigor” refers to rate of growth or biomass accumulation of a crop plant. An “increase in vigor” refers to an increase in growth or biomass accumulation in a crop plant relative to an untreated control crop plant. The term “crop yield” refers to the return on crop material, in terms of both quantity and quality, obtained after harvesting a crop plant. An “increase in crop yield” refers to an increase in crop yield relative to an untreated control crop plant.

The term “biologically effective amount” refers to the amount of a biologically active compound (e.g., a compound of Formula **I** or a compound of Formula **II** and at least one additional biologically active compound or pest control agent) sufficient to produce the desired biological effect when applied to (i.e. contacted with) an invertebrate pest to be controlled or its environment, or to a plant, the seed from which the plant is grown, or the locus of the plant (e.g., growth medium) to protect the plant from injury by the invertebrate pest or for other desired effect (e.g., increasing plant vigor).

Embodiments of the present disclosure as described in the Summary include, but are not limited to those described below.

Embodiment 1. A compound of Formula **I**.

Embodiment 2. A compound of Formula **I** wherein n is 0, 1, or 2

Embodiment 3. A compound of Formula **I** wherein n is 0.

Embodiment 4. A compound of Formula **I** wherein n is 1.

Embodiment 5. A compound of Formula **I** wherein n is 2.

Embodiment 6. A compound of Formula **II** wherein R is CH₃, or CF₂H

Embodiment 7. A compound of Formula **II** wherein R is CH₃.

Embodiment 8. A compound of Formula **II** wherein R is CF₂H.

Embodiment 9. A composition comprising at least one compound according to any one of Embodiments 1-8, or a combination of the foregoing compounds.

Embodiment 10. A composition according to any one of Embodiments 1-9, and at least

one additional biologically active compound or pest control agent different from the compound of Formula I or Formula II.

Embodiment 11. A composition according to any one of Embodiments 1-10, and at least one additional component selected from surfactants, solid diluents and liquid diluents.

5 Embodiment 12. A composition according to any one of Embodiments 1-11, at least one additional biologically active compound or pest control agent, and at least one additional component selected from surfactants, solid diluents and liquid diluents.

Embodiment 13. A composition according to any one of Embodiments 9-12 wherein the compound of Formula I or II is selected from *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, and *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide; or a combination of any of the foregoing compounds.

Embodiment 14. A composition according to any one of Embodiments 9-12 wherein the compound of Formula I or II is selected from *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, and *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

Embodiment 15. A composition according to any one of Embodiments 9-12 wherein the compound of Formula I or II is selected from *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, and *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

25 Embodiment 16. A composition according to any one of Embodiments 10-15 wherein the pest control compound or agent is selected from an insecticide, fungicide, nematocide, and bactericide, or a combination of the foregoing.

Embodiment 17. A composition according to any one of Embodiments 10-16 wherein the pest control compound or agent is an insecticide.

30 Embodiment 18. A composition according to any one of Embodiments 10-16 wherein the pest control compound or agent is a fungicide.

Embodiment 17. A composition according to any one of Embodiments 10-16 wherein the pest control compound or agent is a nematocide.

35 Embodiment 18. A composition according to any one of Embodiments 10-16 wherein the pest control compound or agent is a bactericide.

Embodiment 19. A composition according to any one of Embodiments 10-18 wherein the at least one additional biologically active compound or agent is selected from abamectin, acephate, acequinocyl, acetamiprid, acrinathrin, acynonapyr, afidopyropen

([*(3S,4R,4aR,6S,6aS,12R,12aS,12bS)*]-3-[(cyclopropylcarbonyl)oxy]-
 1,3,4,4a,5,6,6a,12,12a,12b-decahydro-6,12-dihydroxy-4,6a,12b-trimethyl-11-oxo-9-(3-
 pyridinyl)-2*H*,11*H*-naphtho[2,1-*b*]pyrano[3,4-*e*]pyran-4-yl]methyl
 cyclopropanecarboxylate), amidoflumet, amitraz, avermectin, azadirachtin, azinphos-methyl,
 5 benfuracarb, bensultap, benzpyrimoxan, bifenthrin, kappa-bifenthrin, bifenazate, bistrifluron,
 borate, broflanilide, buprofezin, cadusafos, carbaryl, carbofuran, cartap, carzol,
 chlorantraniliprole, chlorfenapyr, chlorfluazuron, chlorprallethrin, chlorpyrifos,
 chlorpyrifos-*e*, chlorpyrifos-methyl, chromafenozide, clofentezin, chlorprallethrin,
 clothianidin, cyantraniliprole (CyazypyrTM) (3-bromo-1-(3-chloro-2-pyridinyl)-*N*-[4-cyano-
 10 2-methyl-6-[(methylamino)carbonyl]phenyl]-1*H*-pyrazole-5-carboxamide), cyclaniliprole
 (3-bromo-*N*-[2-bromo-4-chloro-6-[[*(1-cyclopropylethyl)*]amino]carbonyl]phenyl]-1-(3-
 chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxamide), cyclobutrifluram, cycloprothrin,
 cycloxaprid (*(5S,8R)*-1-[(6-chloro-3-pyridinyl)methyl]-2,3,5,6,7,8-hexahydro-9-nitro-5,8-
 Epoxy-1*H*-imidazo[1,2-*a*]azepine), cyenopyrafen, cyetpyrafen, cyflumetofen, cyfluthrin,
 15 beta-cyfluthrin, cyhalodiamide, cyhalothrin, gamma-cyhalothrin, lambda-cyhalothrin,
 cypermethrin, alpha-cypermethrin, zeta-cypermethrin, cyromazine, deltamethrin,
 diafenthiuron, diazinon, dicloromezotiaz, dieldrin, diflubenzuron, dimefluthrin, dimehypo,
 dimethoate, dimpropridaz, dinotefuran, diofenolan, DiPel®emamectin, emamectin
 benzoate, endosulfan, esfenvalerate, ethiprole, etofenprox, epsilon-metofluthrin, etoxazole,
 20 fenbutatin oxide, fenitrothion, fenothiocarb, fenoxycarb, fenpropathrin, fenvalerate, fipronil,
 flometoquin (2-ethyl-3,7-dimethyl-6-[4-(trifluoromethoxy)phenoxy]-4-quinolinyl methyl
 carbonate), flonicamid, fluazaindolizine, flubendiamide, flucythrinate, flufenerim,
 flufenoxuron, flufenoxystrobin (methyl (*αE*)-2-[[2-chloro-4-
 (trifluoromethyl)phenoxy]methyl]-*α*-(methoxymethylene)benzeneacetate), fluensulfone (5-
 25 chloro-2-[(3,4,4-trifluoro-3-buten-1-yl)sulfonyl]thiazole), fluhexafon, fluopyram,
 flupentiofenox, flupiprole (1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-5-[(2-methyl-2-
 propen-1-yl)amino]-4-[(trifluoromethyl)sulfinyl]-1*H*-pyrazole-3-carbonitrile),
 flupyradifurone (4-[[*(6-chloro-3-pyridinyl)*]methyl](2,2-difluoroethyl)amino]-2(*5H*)-
 furanone), flupyrimin, fluvalinate, tau-fluvalinate, fluxametamide, fonophos, formetanate,
 30 fosthiazate, gamma-cyhalothrin, halofenozide, heptafluthrin ([2,3,5,6-tetrafluoro-4-
 (methoxymethyl)phenyl]methyl 2,2-dimethyl-3-[(1*Z*)-3,3,3-trifluoro-1-propen-1-
 yl]cyclopropanecarboxylate), hexaflumuron, hexythiazox, hydramethylnon, imidacloprid,
 indoxacarb, insecticidal soaps, isofenphos, isocycloseram, kappa-tefluthrin, lambda-
 cyhalothrin, lufenuron, malathion, meperfluthrin ([2,3,5,6-tetrafluoro-4-
 35 (methoxymethyl)phenyl]methyl (*1R,3S*)-3-(2,2-dichloroethenyl)-2,2-
 dimethylcyclopropanecarboxylate), metaflumizone, metaldehyde, methamidophos,
 methidathion, methiocarb, methomyl, methoprene, methoxychlor, metofluthrin,
 methoxyfenozone, epsilon-metofluthrin, metronidazole epsilon-momfluorothrin,

monocrotophos, monofluorothrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 3-(2-cyano-1-propen-1-yl)-2,2-dimethylcyclopropanecarboxylate), nicofluprole, nicotine, nitenpyram, nithiazine, novaluron, noviflumuron, oxamyl, oxazosulfyl, parathion, parathion-methyl, permethrin, phorate, phosalone, phosmet, phosphamidon, pirimicarb, profenofos, profluthrin, propargite, protrifenbute, pyflubumide (1,3,5-trimethyl-*N*-(2-methyl-1-oxopropyl)-*N*-[3-(2-methylpropyl)-4-[2,2,2-trifluoro-1-methoxy-1-(trifluoromethyl)ethyl]phenyl]-1*H*-pyrazole-4-carboxamide), pymetrozine, pyrafluprole, pyrethrin, pyridaben, pyridalyl, pyrifluquinazon, pyriminostrobin (methyl (αE)-2-[[[2-[(2,4-dichlorophenyl)amino]-6-(trifluoromethyl)-4-pyrimidinyl]oxy]methyl]- α -(methoxymethylene)benzeneacetate), pydiflumetofen, pyriprole, pyriproxifen, rotenone, ryanodine, silafluofen, spinetoram, spinosad, spiroadiclofen, spiromesifen, spiropidion, spirotetramat, sulprofos, sulfoxaflor (*N*-[methyloxido[1-[6-(trifluoromethyl)-3-pyridinyl]ethyl]- λ^4 -sulfanylidene]cyanamide), tebufenozide, tebufenpyrad, teflubenzuron, tefluthrin, kappa-tefluthrin, terbufos, tetrachlorantraniliprole, tetrachlorvinphos, tetramethrin, tetramethylfluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 2,2,3,3-tetramethylcyclopropanecarboxylate), tetraniliprole, thiacloprid, thiamethoxam, thiodicarb, thiosultap-sodium, tioazafen (3-phenyl-5-(2-thienyl)-1,2,4-oxadiazole), tolfenpyrad, tralomethrin, triazamate, trichlorfon, triflumezopyrim (2,4-dioxo-1-(5-pyrimidinylmethyl)-3-[3-(trifluoromethyl)phenyl]-2*H*-pyrido[1,2-*a*]pyrimidinium inner salt), triflumuron, tylocyprazoflor, zeta-cypermethrin, *Bacillus thuringiensis* delta-endotoxins, entomopathogenic bacteria, entomopathogenic viruses, and entomopathogenic fungi.

Embodiment 20. A composition according to any one of Embodiments 10-19 wherein the at least one additional biologically active compound or pest control agent is selected from additional biologically active pest control compound or agent, wherein the at least one additional biologically active compound or agent is selected from cyantraniliprole, acetamiprid, imidacloprid, spirotetramat, spiroadiclofen, chlorantraniliprole, bifenthrin, or indoxacarb. Avermectin, *Bacillus* spp., any active crystal proteins thereof, Buprofezin, Carbofuran, Chlorfenapyr, Chlorpyrifos, Clothianidin, Cyromazine, Diafenthiuron, Dinotefuran, Emamectin Benzoate, Fipronil, Flonicamid, Flupyradifurone, methomyl (Lannate®), Methoxyfenozide, Novaluron, Permethrin, Pyriproxifen,

Sulfoxaflor, Thiamethoxam, γ -Cyhalothrin, or ζ -cypermethrin, broflanilide, dimpropridaz, isocycloseram, tetrachlorantraniliprole, oxazosulfyl, tylocyprazoflor, flupyrimin, spiropidion, acynonapyr, benzpyrimoxan, chloroprallethrin, epsilon-metofluthrin, kappa-bifenthrin, dicloromezotiaz, and kappa-tefluthrin.

Embodiment 21. A composition according to any one of Embodiments 10-20 wherein the at least one additional biologically active compound or agent is selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid,

methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

Embodiment 22. A composition according to any one of Embodiments 10-21 wherein the compound of Formula I or Formula II is *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and wherein the at least one additional biologically active compound or pest control agent is at least one selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

Embodiment 23. A composition according to any one of Embodiments 10-21 wherein the compound of Formula I or Formula II is *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and wherein the at least one additional biologically active compound or pest control agent is at least one selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

Embodiment 24. A composition according to any one of Embodiments 10-21 wherein the compound of Formula I or Formula II is *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and wherein the at least one additional biologically active compound or pest control agent is at least one selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

Embodiment 25. A composition according to any one of Embodiments 10-21 wherein the compound of Formula I or Formula II is *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and wherein the at least one additional biologically active compound or pest control agent is at least one selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

Embodiment 26. A composition according to any one of Embodiments 10-21 wherein the compound of Formula I or Formula II is *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and wherein the at least one additional biologically active compound or pest control agent is at least one selected from cyantraniliprole,

pest to pass through the opening so the invertebrate pest can gain access to said bait composition from a location outside the housing, and wherein the housing is further adapted to be placed in or near a locus of potential or known activity for the invertebrate pest.

5 Embodiment 40. A composition according to any one of Embodiments 9-30 wherein the composition is a solid composition selected from dusts, powders, granules, pellets, prills, pastilles, tablets, and filled films.

Embodiment 41. A composition according to Embodiment 40 wherein the solid composition is water-dispersible or water-soluble.

10 Embodiment 42. A liquid or dry formulation comprising the composition of any one of Embodiments 9-32, or 40-41 for use in a drip irrigation system or furrow during planting.

Embodiment 42. A method for controlling an invertebrate pest comprising contacting the invertebrate pest or its environment with a biologically effective amount of a composition according to any one of Embodiments 9-32, or 40-41.

15 Embodiment 43. A method according to Embodiment 42 wherein the environment is soil or plant foliage.

Embodiment 44. A method according to Embodiment 42 or 43 wherein the invertebrate pest is selected from the order Lepidoptera, Blattodea, Coleoptera, Dermaptera, Hemiptera, Homoptera, Acari, Orthoptera, Diptera, Thysanoptera, Hymenoptera, Isoptera, Thysanura, Mallophaga, Siphonoptera, Araneae, and Scutigromorpha.

20 Embodiment 45. A method according to Embodiment 44 wherein the invertebrate pest is selected from the order Lepidoptera, Thrips, and Hemiptera.

25 Embodiment 46. A method according to any one of Embodiments 42-45 wherein the invertebrate pest is selected from silverleaf whitefly, corn plant hopper, western flower thrips, potato leafhopper, cotton melon aphid, root knot nematode, fall army worm and diamondback moth.

Embodiment 47. A treated seed comprising the composition of any one of Embodiments 9-32, or 40-41 in an amount of from about 0.0001 to 1 % by weight of the seed before treatment.

30 In all embodiments of this disclosure, component (b) is different from component (a) in the compositions disclosed herein.

Of note is that compositions of this disclosure are characterized by favorable metabolic and/or soil residual patterns and exhibit activity controlling a spectrum of agronomic and nonagronomic invertebrate pests.

35 Of particular note, for reasons of invertebrate pest control spectrum and economic importance, protection of agronomic crops from damage or injury caused by invertebrate pests by controlling invertebrate pests are embodiments of the disclosure. Compounds and compositions of this disclosure because of their favorable translocation properties or systemicity in plants also protect foliar or other plant parts which are not directly contacted

with a compound of Formula **I** or a compound of Formula **II** or a composition comprising the compound.

Also noteworthy as embodiments of the present disclosure are compositions comprising components (a) and (b) (i.e. in biologically effective amounts) as described in any of the preceding Embodiments, as well as any other embodiments described herein, and any combinations thereof, further comprising at least one additional component selected from the group consisting of a surfactant, a solid diluent, and a liquid diluent, said compositions optionally further comprising at least one additional biologically active compound, and agent (i.e. in a biologically effective amount).

Embodiments of the disclosure also include a composition for protecting an animal comprising components (a) and (b) (i.e. in parasitically effective amounts) of any of the preceding Embodiments and a carrier.

Embodiments of the disclosure further include methods for controlling an invertebrate pest comprising contacting the invertebrate pest or its environment with a biologically effective amount of a composition of any of the preceding Embodiments. Of particular note is a method for protecting an animal comprising administering to the animal a parasitically effective amount of a composition of any of the preceding Embodiments.

Embodiments of the disclosure also include a composition comprising of any of the preceding Embodiments, in the form of a soil drench liquid formulation. Embodiments of the disclosure further include methods for controlling an invertebrate pest comprising contacting the soil with a liquid composition as a soil drench comprising a biologically effective amount of a composition of any of the preceding Embodiments.

Embodiments of the disclosure also include a spray composition for controlling an invertebrate pest comprising a biologically effective amount of a composition of any of the preceding Embodiments and a propellant. Embodiments of the disclosure further include a bait composition for controlling an invertebrate pest comprising a biologically effective amount of a composition of any of the preceding Embodiments, one or more food materials, optionally an attractant, and optionally a humectant. Embodiments of the disclosure also include a device for controlling an invertebrate pest comprising said bait composition and a housing adapted to receive said bait composition, wherein the housing has at least one opening sized to permit the invertebrate pest to pass through the opening so the invertebrate pest can gain access to said bait composition from a location outside the housing, and wherein the housing is further adapted to be placed in or near a locus of potential or known activity for the invertebrate pest.

Embodiments of the disclosure also include a method for protecting a seed from an invertebrate pest comprising contacting the seed with a biologically effective amount of a composition of any of the preceding Embodiments.

Embodiments of the disclosure also include methods for protecting an animal from an invertebrate parasitic pest comprising administering to the animal a parasitically effective amount of a composition of any of the preceding Embodiments.

Embodiments of the disclosure also include methods wherein the invertebrate pest or its environment is contacted with a biologically effective amount of a composition of any of the preceding Embodiments, and at least one additional component selected from the group consisting of surfactants, solid diluents and liquid diluents, said composition optionally further comprising a biologically effective amount of at least one additional biologically active compound or agent, provided that the methods are not methods of medical treatment of a human or animal body by therapy.

Embodiments of this disclosure also include use of an unmanned aerial vehicle (UAV) for the dispensation of the compositions disclosed herein over a planted area. In some embodiments the planted area is a crop-containing area. In some embodiments, the crop is selected from a monocot or dicot. In some embodiments, the crop is selected from rice, corn, barley, soybean, wheat, vegetable, tobacco, tea tree, fruit tree and sugar cane. In some embodiments, the compositions disclosed herein are formulated for spraying at an ultra-low volume. Products applied by drones may use water or oil as the spray carrier. Typical spray volume (including product) used for drone applications globally. 5.0 liters/ha – 100 liters/ha (approximately 0.5-10 gpa). This includes the range of ultra low spray volume (ULV) to low spray volume (LV). Although not common there may be situations where even lower spray volumes could be used as low as 1.0 liter/ha (0.1 gpa)

The term “or combinations thereof” as used herein refers to all permutations and combinations of the listed items preceding the term. For example, “A, B, C, or combinations thereof” is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or more item or term, such as BB, AAA, AB, BBC, AAABCCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

Embodiments of the disclosure also include a composition comprising a compound of any of the preceding Embodiments, in the form of a soil drench liquid formulation. Embodiments of the disclosure further include methods for controlling an invertebrate pest comprising contacting the soil with a liquid composition as a soil drench comprising a biologically effective amount of a compound of any of the preceding Embodiments.

Embodiments of the disclosure also include a spray composition for controlling an invertebrate pest comprising a biologically effective amount of a compound or composition of any of the preceding Embodiments and a propellant. Embodiments of the disclosure further include a bait composition for controlling an invertebrate pest comprising a biologically

effective amount of a compound or composition of any of the preceding Embodiments, one or more food materials, optionally an attractant, and optionally a humectant. Embodiments of the disclosure also include a device for controlling an invertebrate pest comprising said bait composition and a housing adapted to receive said bait composition, wherein the housing has at least one opening sized to permit the invertebrate pest to pass through the opening so the invertebrate pest can gain access to said bait composition from a location outside the housing, and wherein the housing is further adapted to be placed in or near a locus of potential or known activity for the invertebrate pest.

Embodiments of the disclosure also include methods for protecting a seed from an invertebrate pest comprising contacting the seed with a biologically effective amount of a compound or composition of any of the preceding Embodiments.

Embodiments of the disclosure also include methods for protecting an animal from an invertebrate parasitic pest comprising administering to the animal a parasitically effective amount of a compound or composition of any of the preceding Embodiments.


Embodiments of the disclosure also include methods for controlling an invertebrate pest comprising contacting the invertebrate pest or its environment with a biologically effective amount of a composition comprising at least one compound of Formula **I** or at least one compound of Formula **II** and at least one biologically active compound or pest control agent, provided that the methods are not methods of medical treatment of a human or animal body by therapy.

Embodiments of this disclosure also relates to such methods wherein the invertebrate pest or its environment is contacted with a composition comprising a biologically effective amount of compound of Formula **I** or at least one compound of Formula **II**, at least one biologically active compound or pest control agent, and at least one additional component selected from surfactants, solid diluents, and liquid diluents.

Compounds of Formula **I** or compounds of Formula **II** can be prepared by the following methods and variations as described in Schemes 1 and 2. The following abbreviations are used: DMF is *N,N*-dimethylformamide, DCC is *N,N'*-dicyclohexylcarbodiimide, and HATU is 1-[bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate.

The compound of Formula **I** can be prepared from Compound 6 by the method shown in Scheme 1. In this method, the compound of Formula **I** is prepared by an amide-bond-forming reaction of the carboxylic group of Compound 6 with the amine group of Compound 7a (when *n* is 0), 7b (when *n* is 1), or 7c (when *n* is 2), in the presence of an amide coupling reagent such as DCC or HATU.

20

Scheme 1

5


The compounds of Formula I wherein n is 1 or 2 (i.e. the sulfoxide, Compound 2, or the sulfone, Compound 3, respectively) can be prepared by oxidation of the sulfide (Compound 1). A variety of methods and reagents are known in the art for oxidizing sulfides to sulfoxides and sulfones. Examples of such oxidizing reagents include *meta*-chloroperoxybenzoic acid and sodium periodate.

10

Compounds according to Formula II can be prepared from Compound 8 by the method shown in Scheme 2. In this method, Compound 4 is prepared by an amide forming reaction of the carboxyl group of Compound 8 with the amine group of Compound 9 in the presence of an amide coupling reagent such as DCC or HATU. For representative reagents and reaction conditions, see Jones, J. *The Chemical Synthesis of Peptides*, International Series of Monographs on Chemistry, Oxford University: Oxford, 1994.

15

Scheme 2




20

25

Similar to Compound 4, Compound 5 can be prepared from Compound 8 by the method shown in Scheme 3. In this method, Compound 5 is prepared by an amide-bond-forming reaction of the carboxyl group of Compound 8 with the amine group of Compound 10 in the presence of an amide coupling reagent such as DCC or HATU.

21

Scheme 3



Compound 8 (CAS Registry Number 2001277-98-9) is known in the art.

5 It is recognized that some reagents and reaction conditions described above for preparing compounds of Formula I or compounds of Formula II may not be compatible with certain functionalities present in the intermediates. In these instances, the incorporation of protection/deprotection sequences or functional group interconversions into the synthesis will aid in obtaining the desired products. The use and choice of the protecting groups will be apparent to one skilled in chemical synthesis (see, for example, Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). One skilled in the art will recognize that, in some cases, after introduction of the reagents depicted in the individual schemes, additional routine synthetic steps not described in detail may be needed to complete the synthesis of compounds of Formula I or compounds of Formula II. One skilled in the art will also recognize that it may be necessary to perform a combination of the steps illustrated in the above schemes in an order other than that implied by the particular sequence presented to prepare compounds of Formula I or compounds of Formula II.

One skilled in the art will also recognize that compounds of Formula I or compounds of Formula II and the intermediates described herein can be subjected to various electrophilic, nucleophilic, radical, organometallic, oxidation, and reduction reactions to add substituents or modify existing substituents.

Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present disclosure to its fullest extent. The following Synthesis Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Steps in the following Synthesis Examples illustrate a procedure for each step in an overall synthetic transformation, and the starting material for each step may not have necessarily been prepared by a particular preparative run whose procedure is described in other Examples or Steps. Percentages are by weight except for chromatographic solvent mixtures or where otherwise indicated. Parts and percentages for chromatographic solvent mixtures are by volume unless otherwise indicated. ¹H NMR spectra are reported in ppm downfield from tetramethylsilane; “s” means singlet, “d” means doublet, “t” means triplet, “q” means quartet, “m” means multiplet, “dd” means doublet of

doublets, “dt” means doublet of triplets, “br s” means broad singlet. DMF means *N,N*-dimethylformamide. Compound numbers refer to Index Table 1.

SYNTHESIS EXAMPLE 1

5 Preparation of *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide (Compound 1) (*N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide) (Compound 2), and (*N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H* indazole-4-carboxamide (Compound 3)

10 Step A: Preparation of *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide (Compound 1)

A solution of 7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxylic acid (10 g, 39 mmol), HATU (17.7 g, 47 mmol), 2-methyl-1-methylsulfonyl-propan-2-amine (7 g, 58 mmol) in DMF (100 mL) was treated with triethylamine (16 mL, 117 mmol). The reaction mixture was stirred for 4 hours at room temperature. The reaction mixture was diluted with EtOAc (300 mL) and washed with water (6 x 100 mL). The organic layer was separated and concentrated *in vacuo*. The resulting crude solid was purified via normal phase chromatography (silica gel eluted with 70-100% EtOAc in hexane) to yield the title compound, a compound of this disclosure (9.9 g, 71% yield). ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 9.37-9.43 (m, 2H), 8.69-8.72 (m, 1H), 8.56-8.60 (m, 1H), 8.00 (s, 1H), 7.64-7.70 (m, 2 H), 7.22-7.27 (m, 1H), 3.11 (s, 2H), 2.09 (s, 3H), 1.47 (s, 6H).

20 Compound 2 (*N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide), and Compound 3 (*N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H* indazole-4-carboxamide, can each be prepared via oxidation of
25 Compound 1.

SYNTHESIS EXAMPLE 2

Preparation of *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide (Compound 4) and *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide (Compound 5)

30

Step A: Preparation of *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide (Compound 5)

35 A solution of 2-(3-pyridinyl)-2*H*-indazole-4-carboxylic acid (100 mg, 0.42 mmol, CAS Registry Number 2001277-98-9), HATU (190 mg, 0.5 mmol), 1-(difluoromethyl)cyclopropanamine hydrochloride (71 mg, 0.5 mmol) in DMF (2 mL) was treated with triethylamine (174 mL, 1.25 mmol). The reaction mixture was stirred overnight

at room temperature. The reaction mixture was then purified directly via reverse phase column chromatography [C18 column eluted with 10%-100% MeCN/MeOH (1:1) to water] to yield the title compound, a compound of this disclosure (105 mg, 76% yield). ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 9.35-9.39 (m, 2H) 9.14 (s, 1H) 8.67-8.68 (m, 1H) 8.55-8.58 (m, 1H) 7.94-7.97 (m, 1H), 7.73-7.75 (m, 1H), 7.63-7.67 (m, 1H), 7.41-7.45 (m, 1H), 6.21 (t, 1H), 1.13-1.17 (m, 2H), 1.02-1.06 (m, 2H). Compound 4 was prepared as described for Compound 5, but 1-(difluoromethyl)cyclopropanamine hydrochloride was replaced with 1-methylcyclopropanamine hydrochloride.

10 A composition of this disclosure will generally be used as an invertebrate pest control active ingredient in a composition, i.e. formulation, with at least one additional component selected from the group consisting of surfactants, solid diluents, and liquid diluents, which serves as a carrier. The formulation or composition ingredients are selected to be consistent with the physical properties of the active ingredient, mode of application and environmental factors such as soil type, moisture and temperature.

15 Useful formulations include both liquid and solid compositions. Liquid compositions include solutions (including emulsifiable concentrates), suspensions, emulsions (including microemulsions, oil in water emulsions, flowable concentrates and/or suspoemulsions) and the like, which optionally can be thickened into gels. The general types of aqueous liquid compositions are soluble concentrate, suspension concentrate, capsule suspension, 20 concentrated emulsion, microemulsion, oil in water emulsion, flowable concentrate and suspoemulsion. The general types of nonaqueous liquid compositions are emulsifiable concentrate, microemulsifiable concentrate, dispersible concentrate and oil dispersion.

25 The general types of solid compositions are dusts, powders, granules, pellets, prills, pastilles, tablets, filled films (including seed coatings) and the like, which can be water-dispersible (“wettable”) or water-soluble. Films and coatings formed from film-forming solutions or flowable suspensions are particularly useful for seed treatment. Active ingredient can be (micro)encapsulated and further formed into a suspension or solid formulation; alternatively the entire formulation of active ingredient can be encapsulated (or “overcoated”). 30 Encapsulation can control or delay release of the active ingredient. An emulsifiable granule combines the advantages of both an emulsifiable concentrate formulation and a dry granular formulation. High-strength compositions are primarily used as intermediates for further formulation.

35 Sprayable formulations are typically extended in a suitable medium before spraying. Such liquid and solid formulations are formulated to be readily diluted in the spray medium, usually water, but occasionally another suitable medium like an aromatic or paraffinic hydrocarbon or vegetable oil. Spray volumes can range from about one to several thousand liters per hectare, but more typically are in the range from about ten to several hundred liters

per hectare. Sprayable formulations can be tank mixed with water or another suitable medium for foliar treatment by aerial or ground application, or for application to the growing medium of the plant. Liquid and dry formulations can be metered directly into drip irrigation systems or metered into the furrow during planting. Liquid and solid formulations can be applied onto
 5 seeds of crops and other desirable vegetation as seed treatments before planting to protect developing roots and other subterranean plant parts and/or foliage through systemic uptake.

One way of dispensing the compositions disclosed herein over a target area, such as, but not limited to a crop-containing field, is by using drones. Use of drones or unmanned aerial vehicles (UAVs) in agricultural applications, such as for treating fields with chemical
 10 products, is rapidly expanding. A container of chemical products is coupled to the UAV and a material dispensing system mounted to the UAV, and the UAV is piloted above the area to be treated while the chemical product is dispensed.

Formulations will typically contain effective amounts of active ingredient, diluent and surfactant within the following approximate ranges which add up to 100 percent by weight.

	Weight Percent		
	<u>Active Ingredient</u>	<u>Diluent</u>	<u>Surfactant</u>
Water-Dispersible and Water-soluble Granules, Tablets and Powders	0.001–90	0–99.999	0–15
Oil Dispersions, Suspensions, Emulsions, Solutions (including Emulsifiable Concentrates)	1–50	40–99	0–50
Dusts	1–25	70–99	0–5
Granules and Pellets	0.001–95	5–99.999	0–15
High Strength Compositions	90–99	0–10	0–2

15 Solid diluents include, for example, clays such as bentonite, montmorillonite, attapulgite and kaolin, gypsum, cellulose, titanium dioxide, zinc oxide, starch, dextrin, sugars (e.g., lactose, sucrose), silica, talc, mica, diatomaceous earth, urea, calcium carbonate, sodium carbonate and bicarbonate, and sodium sulfate. Typical solid diluents are described in Watkins et al., *Handbook of Insecticide Dust Diluents and Carriers*, 2nd Ed., Dorland Books,
 20 Caldwell, New Jersey.

Liquid diluents include, for example, water, *N,N*-dimethylalkanamides (e.g., *N,N*-dimethylformamide), limonene, dimethyl sulfoxide, *N*-alkylpyrrolidones (e.g., *N*-methylpyrrolidinone), alkyl phosphates (e.g., triethylphosphate), ethylene glycol, triethylene glycol, propylene glycol, dipropylene glycol, polypropylene glycol, propylene
 25 carbonate, butylene carbonate, paraffins (e.g., white mineral oils, normal paraffins,

isoparaffins), alkylbenzenes, alkylnaphthalenes, glycerine, glycerol triacetate, sorbitol, aromatic hydrocarbons, dearomatized aliphatics, alkylbenzenes, alkylnaphthalenes, ketones such as cyclohexanone, 2-heptanone, isophorone and 4-hydroxy-4-methyl-2-pentanone, acetates such as isoamyl acetate, hexyl acetate, heptyl acetate, octyl acetate, nonyl acetate, 5 tridecyl acetate and isobornyl acetate, other esters such as alkylated lactate esters, dibasic esters alkyl and aryl benzoates, γ -butyrolactone, and alcohols, which can be linear, branched, saturated or unsaturated, such as methanol, ethanol, *n*-propanol, isopropyl alcohol, *n*-butanol, isobutyl alcohol, *n*-hexanol, 2-ethylhexanol, *n*-octanol, decanol, isodecyl alcohol, isooctadecanol, cetyl alcohol, lauryl alcohol, tridecyl alcohol, oleyl alcohol, cyclohexanol, 10 tetrahydrofurfuryl alcohol, diacetone alcohol, cresol and benzyl alcohol. Liquid diluents also include glycerol esters of saturated and unsaturated fatty acids (typically C₆-C₂₂), such as plant seed and fruit oils (e.g., oils of olive, castor, linseed, sesame, corn (maize), peanut, sunflower, grapeseed, safflower, cottonseed, soybean, rapeseed, coconut and palm kernel), animal-sourced fats (e.g., beef tallow, pork tallow, lard, cod liver oil, fish oil), 15 and mixtures thereof. Liquid diluents also include alkylated fatty acids (e.g., methylated, ethylated, butylated) wherein the fatty acids may be obtained by hydrolysis of glycerol esters from plant and animal sources, and can be purified by distillation. Typical liquid diluents are described in Marsden, *Solvents Guide*, 2nd Ed., Interscience, New York, 1950.

The solid and liquid compositions of the present disclosure often include one or more 20 surfactants. When added to a liquid, surfactants (also known as “surface-active agents”) generally modify, most often reduce, the surface tension of the liquid. Depending on the nature of the hydrophilic and lipophilic groups in a surfactant molecule, surfactants can be useful as wetting agents, dispersants, emulsifiers or defoaming agents.

Surfactants can be classified as nonionic, anionic or cationic. Nonionic surfactants 25 useful for the present compositions include, but are not limited to: alcohol alkoxyates such as alcohol alkoxyates based on natural and synthetic alcohols (which may be branched or linear) and prepared from the alcohols and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof; amine ethoxyates, alkanolamides and ethoxylated alkanolamides; alkoxyated triglycerides such as ethoxylated soybean, castor and rapeseed oils; alkylphenol 30 alkoxyates such as octylphenol ethoxyates, nonylphenol ethoxyates, dinonyl phenol ethoxyates and dodecyl phenol ethoxyates (prepared from the phenols and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); block polymers prepared from ethylene oxide or propylene oxide and reverse block polymers where the terminal blocks are prepared from propylene oxide; ethoxylated fatty acids; ethoxylated fatty esters and oils; ethoxylated 35 methyl esters; ethoxylated tristyrylphenol (including those prepared from ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); fatty acid esters, glycerol esters, lanolin-based derivatives, polyethoxylate esters such as polyethoxylated sorbitan fatty acid esters, polyethoxylated sorbitol fatty acid esters and polyethoxylated glycerol fatty acid esters; other

sorbitan derivatives such as sorbitan esters; polymeric surfactants such as random copolymers, block copolymers, alkyd peg (polyethylene glycol) resins, graft or comb polymers and star polymers; polyethylene glycols (pegs); polyethylene glycol fatty acid esters; silicone-based surfactants; and sugar-derivatives such as sucrose esters, alkyl polyglycosides and alkyl polysaccharides.

Useful anionic surfactants include, but are not limited to: alkylaryl sulfonic acids and their salts; carboxylated alcohol or alkylphenol ethoxylates; diphenyl sulfonate derivatives; lignin and lignin derivatives such as lignosulfonates; maleic or succinic acids or their anhydrides; olefin sulfonates; phosphate esters such as phosphate esters of alcohol alkoxyates, phosphate esters of alkylphenol alkoxyates and phosphate esters of styryl phenol ethoxylates; protein-based surfactants; sarcosine derivatives; styryl phenol ether sulfate; sulfates and sulfonates of oils and fatty acids; sulfates and sulfonates of ethoxylated alkylphenols; sulfates of alcohols; sulfates of ethoxylated alcohols; sulfonates of amines and amides such as *N,N*-alkyltaurates; sulfonates of benzene, cumene, toluene, xylene, and dodecyl and tridecylbenzenes; sulfonates of condensed naphthalenes; sulfonates of naphthalene and alkyl naphthalene; sulfonates of fractionated petroleum; sulfosuccinamates; and sulfosuccinates and their derivatives such as dialkyl sulfosuccinate salts.

Useful cationic surfactants include, but are not limited to: amides and ethoxylated amides; amines such as *N*-alkyl propanediamines, tripropylenetriamines and dipropylenetetramines, and ethoxylated amines, ethoxylated diamines and propoxylated amines (prepared from the amines and ethylene oxide, propylene oxide, butylene oxide or mixtures thereof); amine salts such as amine acetates and diamine salts; quaternary ammonium salts such as quaternary salts, ethoxylated quaternary salts and diquaternary salts; and amine oxides such as alkyldimethylamine oxides and bis-(2-hydroxyethyl)-alkylamine oxides.

Also useful for the present compositions are mixtures of nonionic and anionic surfactants or mixtures of nonionic and cationic surfactants. Nonionic, anionic and cationic surfactants and their recommended uses are disclosed in a variety of published references including *McCutcheon's Emulsifiers and Detergents*, annual American and International Editions published by McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; Sisely and Wood, *Encyclopedia of Surface Active Agents*, Chemical Publ. Co., Inc., New York, 1964; and A. S. Davidson and B. Milwidsky, *Synthetic Detergents*, Seventh Edition, John Wiley and Sons, New York, 1987.

Compositions of this disclosure may also contain formulation auxiliaries and additives, known to those skilled in the art as formulation aids (some of which may be considered to also function as solid diluents, liquid diluents or surfactants). Such formulation auxiliaries and additives may control: pH (buffers), foaming during processing (antifoams such polyorganosiloxanes), sedimentation of active ingredients (suspending agents), viscosity

(thixotropic thickeners), in-container microbial growth (antimicrobials), product freezing (antifreezes), color (dyes/pigment dispersions), wash-off (film formers or stickers), evaporation (evaporation retardants), and other formulation attributes. Film formers include, for example, polyvinyl acetates, polyvinyl acetate copolymers, polyvinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohols, polyvinyl alcohol copolymers and waxes. Examples of formulation auxiliaries and additives include those listed in *McCutcheon's Volume 2: Functional Materials*, annual International and North American editions published by McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; and PCT Publication WO 03/024222.

Compositions disclosed herein comprising compounds of Formula I or compounds of Formula II and any other active ingredients are typically incorporated into the present compositions by dissolving the active ingredient in a solvent or by grinding in a liquid or dry diluent. Solutions, including emulsifiable concentrates, can be prepared by simply mixing the ingredients. If the solvent of a liquid composition intended for use as an emulsifiable concentrate is water-immiscible, an emulsifier is typically added to emulsify the active-containing solvent upon dilution with water. Active ingredient slurries, with particle diameters of up to 2,000 μm can be wet milled using media mills to obtain particles with average diameters below 3 μm . Aqueous slurries can be made into finished suspension concentrates (see, for example, U.S. 3,060,084) or further processed by spray drying to form water-dispersible granules. Dry formulations usually require dry milling processes, which produce average particle diameters in the 2 to 10 μm range. Dusts and powders can be prepared by blending and usually grinding (such as with a hammer mill or fluid-energy mill). Granules and pellets can be prepared by spraying the active material upon preformed granular carriers or by agglomeration techniques. See Browning, "Agglomeration", *Chemical Engineering*, December 4, 1967, pp 147–48, *Perry's Chemical Engineer's Handbook*, 4th Ed., McGraw-Hill, New York, 1963, pages 8–57 and following, and WO 91/13546. Pellets can be prepared as described in U.S. 4,172,714. Water-dispersible and water-soluble granules can be prepared as taught in U.S. 4,144,050, U.S. 3,920,442 and DE 3,246,493. Tablets can be prepared as taught in U.S. 5,180,587, U.S. 5,232,701 and U.S. 5,208,030. Films can be prepared as taught in GB 2,095,558 and U.S. 3,299,566.

For further information regarding the art of formulation, see T. S. Woods, "The Formulator's Toolbox – Product Forms for Modern Agriculture" in *Pesticide Chemistry and Bioscience, The Food–Environment Challenge*, T. Brooks and T. R. Roberts, Eds., Proceedings of the 9th International Congress on Pesticide Chemistry, The Royal Society of Chemistry, Cambridge, 1999, pp. 120–133. See also U.S. 3,235,361, Col. 6, line 16 through Col. 7, line 19 and Examples 10–41; U.S. 3,309,192, Col. 5, line 43 through Col. 7, line 62 and Examples 8, 12, 15, 39, 41, 52, 53, 58, 132, 138–140, 162–164, 166, 167 and 169–182; U.S. 2,891,855, Col. 3, line 66 through Col. 5, line 17 and Examples 1–4; Klingman, *Weed*

Control as a Science, John Wiley and Sons, Inc., New York, 1961, pp 81–96; Hance et al., *Weed Control Handbook*, 8th Ed., Blackwell Scientific Publications, Oxford, 1989; and *Developments in formulation technology*, PJB Publications, Richmond, UK, 2000.

In the following Examples, all formulations are prepared in conventional ways. “Active ingredients” refers to the aggregate of biologically active compounds or agents consisting of invertebrate pest control agents selected from the at least one additional pest control agent or compound of (b) in combination with at least one compound of Formula I or Formula II, or a combination of one or more compounds of each of Formula I and II. “Active ingredients” can also refer to at least one compound of Formula I or Formula II, or a combination of one or more compounds of each of Formula I and II, in addition to at least one additional biologically active compound, such as an insecticide, fungicide, nematocide or bactericide. Without further elaboration, it is believed that one skilled in the art using the preceding description can utilize the present disclosure to its fullest extent. The following Examples are, therefore, to be construed as merely illustrative, and not limiting of the disclosure in any way whatsoever. Percentages are by weight except where otherwise indicated.

Example A

High Strength Concentrate

active ingredients	98.5%
silica aerogel	0.5%
synthetic amorphous fine silica	1.0%

Example B

Wettable Powder

active ingredients	65.0%
dodecylphenol polyethylene glycol ether	2.0%
sodium ligninsulfonate	4.0%
sodium silicoaluminate	6.0%
montmorillonite (calcined)	23.0%

20

Example C

Granule

active ingredients	10.0%
attapulgite granules (low volatile matter, 0.71/0.30 mm; U.S.S. No. 25–50 sieves)	90.0%

Example D

Extruded Pellet

active ingredients	25.0%
anhydrous sodium sulfate	10.0%

crude calcium ligninsulfonate	5.0%
sodium alkylphthalenesulfonate	1.0%
calcium/magnesium bentonite	59.0%

Example EEmulsifiable Concentrate

active ingredients	10.0%
polyoxyethylene sorbitol hexoleate	20.0%
C ₆ -C ₁₀ fatty acid methyl ester	70.0%

Example FMicroemulsion

active ingredients	5.0%
polyvinylpyrrolidone-vinyl acetate copolymer	30.0%
alkylpolyglycoside	30.0%
glyceryl monooleate	15.0%
water	20.0%

Example GSeed Treatment

active ingredients	20.00%
polyvinylpyrrolidone-vinyl acetate copolymer	5.00%
montan acid wax	5.00%
calcium ligninsulfonate	1.00%
polyoxyethylene/polyoxypropylene block copolymers	1.00%
stearyl alcohol (POE 20)	2.00%
polyorganosilane	0.20%
colorant red dye	0.05%
water	65.75%

Example HFertilizer Stick

active ingredients	2.5%
pyrrolidone-styrene copolymer	4.8%
tristyrylphenyl 16-ethoxylate	2.3%
talc	0.8%
corn starch	5.0%
slow-release fertilizer	36.0%
kaolin	38.0%
water	10.6%

30

Example ISuspension Concentrate

active ingredients	35%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
water	53.7%

Example JEmulsion in Water

active ingredients	10.0%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%
propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
aromatic petroleum based hydrocarbon	20.0
water	58.7%

Example KOil Dispersion

active ingredients	25%
polyoxyethylene sorbitol hexaoleate	15%
organically modified bentonite clay	2.5%
fatty acid methyl ester	57.5%

Example LSuspoemulsion

active ingredients	10.0%
imidacloprid	5.0%
butyl polyoxyethylene/polypropylene block copolymer	4.0%
stearic acid/polyethylene glycol copolymer	1.0%
styrene acrylic polymer	1.0%
xanthan gum	0.1%

propylene glycol	5.0%
silicone based defoamer	0.1%
1,2-benzisothiazolin-3-one	0.1%
aromatic petroleum based hydrocarbon	20.0%
water	53.7%

Compositions of this disclosure exhibit activity against a wide spectrum of invertebrate pests. These pests include invertebrates inhabiting a variety of environments such as, for example, plant foliage, roots, soil, harvested crops or other foodstuffs, building structures or animal integuments. These pests include, for example, invertebrates feeding on foliage (including leaves, stems, flowers and fruits), seeds, wood, textile fibers or animal blood or tissues, and thereby causing injury or damage to, for example, growing or stored agronomic crops, forests, greenhouse crops, ornamentals, nursery crops, stored foodstuffs or fiber products, or houses or other structures or their contents, or being harmful to animal health or public health. Those skilled in the art will appreciate that not all compounds are equally effective against all growth stages of all pests.

These present compositions are thus useful agronomically for protecting field crops from phytophagous invertebrate pests, and also nonagronomically for protecting other horticultural crops and plants from phytophagous invertebrate pests. This utility includes protecting crops and other plants (i.e. both agronomic and nonagronomic) that contain genetic material introduced by genetic engineering (i.e. transgenic) or modified by mutagenesis to provide advantageous traits. Examples of such traits include tolerance to herbicides, resistance to phytophagous pests (e.g., insects, mites, aphids, spiders, nematodes, snails, plant-pathogenic fungi, bacteria and viruses), improved plant growth, increased tolerance of adverse growing conditions such as high or low temperatures, low or high soil moisture, and high salinity, increased flowering or fruiting, greater harvest yields, more rapid maturation, higher quality and/or nutritional value of the harvested product, or improved storage or process properties of the harvested products. Transgenic plants can be modified to express multiple traits. Examples of plants containing traits provided by genetic engineering or mutagenesis include varieties of corn, cotton, soybean and potato expressing an insecticidal *Bacillus thuringiensis* toxin such as YIELD GARD[®], KNOCKOUT[®], STARLINK[®], BOLLGARD[®], NuCOTN[®] and NEWLEAF[®], INVICTA RR2 PRO[™], and herbicide-tolerant varieties of corn, cotton, soybean and rapeseed such as ROUNDUP READY[®], LIBERTY LINK[®], IMI[®], STS[®] and CLEARFIELD[®], as well as crops expressing *N*-acetyltransferase (GAT) to provide resistance to glyphosate herbicide, or crops containing the HRA gene providing resistance to herbicides inhibiting acetolactate synthase (ALS). The present compositions may interact synergistically with traits introduced by genetic engineering or modified by mutagenesis, thus enhancing phenotypic expression or effectiveness of the traits or increasing the invertebrate pest control effectiveness of the present compounds and compositions. In particular, the

present compositions may interact synergistically with the phenotypic expression of proteins or other natural products toxic to invertebrate pests to provide greater-than-additive control of these pests, i.e. produce a combined effect greater than the sum of their separate effects.

5 Compositions of this disclosure can also optionally comprise plant nutrients, e.g., a fertilizer composition comprising at least one plant nutrient selected from nitrogen, phosphorus, potassium, sulfur, calcium, magnesium, iron, copper, boron, manganese, zinc, and molybdenum. Of note are compositions comprising at least one fertilizer composition comprising at least one plant nutrient selected from nitrogen, phosphorus, potassium, sulfur, calcium and magnesium. Compositions of the present disclosure which further comprise at
10 least one plant nutrient can be in the form of liquids or solids. Of note are solid formulations in the form of granules, small sticks or tablets. Solid formulations comprising a fertilizer composition can be prepared by mixing the compound or composition of the present disclosure with the fertilizer composition together with formulating ingredients and then preparing the formulation by methods such as granulation or extrusion. Alternatively solid
15 formulations can be prepared by spraying a solution or suspension of a compound or composition of the present disclosure in a volatile solvent onto a previous prepared fertilizer composition in the form of dimensionally stable mixtures, e.g., granules, small sticks or tablets, and then evaporating the solvent.

Nonagronomic uses refer to invertebrate pest control in the areas other than fields of
20 crop plants. Nonagronomic uses of the present compositions include control of invertebrate pests in stored grains, beans and other foodstuffs, and in textiles such as clothing and carpets. Nonagronomic uses of the present compositions also include invertebrate pest control in ornamental plants, forests, in yards, along roadsides and railroad rights of way, and on turf such as lawns, golf courses and pastures. Nonagronomic uses of the present compositions
25 also include invertebrate pest control in houses and other buildings which may be occupied by humans and/or companion, farm, ranch, zoo or other animals. Nonagronomic uses of the present compositions also include the control of pests such as termites that can damage wood or other structural materials used in buildings.

Nonagronomic uses of the present compositions also include protecting human and
30 animal health by controlling invertebrate pests that are parasitic or transmit infectious diseases. The controlling of animal parasites includes controlling external parasites that are parasitic to the surface of the body of the host animal (e.g., shoulders, armpits, abdomen, inner part of the thighs) and internal parasites that are parasitic to the inside of the body of the host animal (e.g., stomach, intestine, lung, veins, under the skin, lymphatic tissue). External
35 parasitic or disease transmitting pests include, for example, chiggers, ticks, lice, mosquitoes, flies, mites and fleas. Internal parasites include heartworms, hookworms and helminths. of the present disclosure are suitable for systemic and/or non-systemic control of infestation or infection by parasites on animals. Compositions of the present disclosure are particularly

suitable for combating external parasitic or disease transmitting pests. Compositions of the present disclosure are suitable for combating parasites that infest agricultural working animals, such as cattle, sheep, goats, horses, pigs, donkeys, camels, buffalos, rabbits, hens, turkeys, ducks, geese and bees; pet animals and domestic animals such as dogs, cats, pet birds and aquarium fish; as well as so-called experimental animals, such as hamsters, guinea pigs, rats and mice. By combating these parasites, fatalities and performance reduction (in terms of meat, milk, wool, skins, eggs, honey, etc.) are reduced, so that applying a composition of the present disclosure allows more economic and simple husbandry of animals.

Examples of agronomic or nonagronomic invertebrate pests include eggs, larvae and adults of the order Lepidoptera, such as armyworms, cutworms, loopers, and heliothines in the family Noctuidae (e.g., pink stem borer (*Sesamia inferens* Walker), corn stalk borer (*Sesamia nonagrioides* Lefebvre), southern armyworm (*Spodoptera eridania* Cramer), fall armyworm (*Spodoptera frugiperda* J. E. Smith), beet armyworm (*Spodoptera exigua* Hübner), cotton leafworm (*Spodoptera littoralis* Boisduval), yellowstriped armyworm (*Spodoptera ornithogalli* Guenée), black cutworm (*Agrotis ipsilon* Hufnagel), velvetbean caterpillar (*Anticarsia gemmatalis* Hübner), green fruitworm (*Lithophane antennata* Walker), cabbage armyworm (*Barathra brassicae* Linnaeus), soybean looper (*Pseudoplusia includens* Walker), cabbage looper (*Trichoplusia ni* Hübner), tobacco budworm (*Heliothis virescens* Fabricius)); borers, casebearers, webworms, coneworms, cabbageworms and skeletonizers from the family Pyralidae (e.g., European corn borer (*Ostrinia nubilalis* Hübner), navel orangeworm (*Amyelois transitella* Walker), corn root webworm (*Crambus caliginosellus* Clemens), sod webworms (Pyralidae: *Crambinae*) such as sod worm (*Herpetogramma licarsisalis* Walker), sugarcane stem borer (*Chilo infuscatellus* Snellen), tomato small borer (*Neoleucinodes elegantalis* Guenée), green leafroller (*Cnaphalocrocis medinalis*), grape leaf folder (*Desmia funeralis* Hübner), melon worm (*Diaphania nitidalis* Stoll), cabbage center grub (*Helluata hydralis* Guenée), yellow stem borer (*Scirpophaga incertulas* Walker), early shoot borer (*Scirpophaga infuscatellus* Snellen), white stem borer (*Scirpophaga innotata* Walker), top shoot borer (*Scirpophaga nivella* Fabricius), dark-headed rice borer (*Chilo polychrysus* Meyrick), striped riceborer (*Chilo suppressalis* Walker), cabbage cluster caterpillar (*Crocidolomia binotalis* English)); leafrollers, budworms, seed worms, and fruit worms in the family Tortricidae (e.g., codling moth (*Cydia pomonella* Linnaeus), grape berry moth (*Endopiza viteana* Clemens), oriental fruit moth (*Grapholita molesta* Busck), citrus false codling moth (*Cryptophlebia leucotreta* Meyrick), citrus borer (*Ecdytolopha aurantiana* Lima), redbanded leafroller (*Argyrotaenia velutinana* Walker), obliquebanded leafroller (*Choristoneura rosaceana* Harris), light brown apple moth (*Epiphyas postvittana* Walker), European grape berry moth (*Eupoecilia ambiguella* Hübner), apple bud moth (*Pandemis pyrusana* Kearfott), omnivorous leafroller (*Platynota stultana* Walsingham), barred fruit-tree tortrix (*Pandemis cerasana* Hübner), apple brown tortrix (*Pandemis heparana* Denis &

Schiffermüller)); and many other economically important lepidoptera (e.g., diamondback moth (*Plutella xylostella* Linnaeus), pink bollworm (*Pectinophora gossypiella* Saunders), gypsy moth (*Lymantria dispar* Linnaeus), peach fruit borer (*Carposina niponensis* Walsingham), peach twig borer (*Anarsia lineatella* Zeller), potato tuberworm (*Phthorimaea operculella* Zeller), spotted teniform leafminer (*Lithocolletis blancardella* Fabricius), Asiatic apple leafminer (*Lithocolletis ringoniella* Matsumura), rice leaf folder (*Lerodea eufala* Edwards), apple leafminer (*Leucoptera scitella* Zeller)); eggs, nymphs and adults of the order Blattodea including cockroaches from the families Blattellidae and Blattidae (e.g., oriental cockroach (*Blatta orientalis* Linnaeus), Asian cockroach (*Blattella asahinai* Mizukubo), German cockroach (*Blattella germanica* Linnaeus), brownbanded cockroach (*Supella longipalpa* Fabricius), American cockroach (*Periplaneta americana* Linnaeus), brown cockroach (*Periplaneta brunnea* Burmeister), Madeira cockroach (*Leucophaea maderae* Fabricius)), smoky brown cockroach (*Periplaneta fuliginosa* Service), Australian Cockroach (*Periplaneta australasiae* Fabr.), lobster cockroach (*Nauphoeta cinerea* Olivier) and smooth cockroach (*Symploce pallens* Stephens)); eggs, foliar feeding, fruit feeding, root feeding, seed feeding and vesicular tissue feeding larvae and adults of the order Coleoptera including weevils from the families Anthribidae, Bruchidae, and Curculionidae (e.g., boll weevil (*Anthonomus grandis* Boheman), rice water weevil (*Lissorhoptrus oryzophilus* Kuschel), granary weevil (*Sitophilus granarius* Linnaeus), rice weevil (*Sitophilus oryzae* Linnaeus)), annual bluegrass weevil (*Listronotus maculicollis* Dietz), bluegrass billbug (*Sphenophorus parvulus* Gyllenhal), hunting billbug (*Sphenophorus venatus vestitus*), Denver billbug (*Sphenophorus cicatristriatus* Fahraeus)); flea beetles, cucumber beetles, rootworms, leaf beetles, potato beetles, and leafminers in the family Chrysomelidae (e.g., Colorado potato beetle (*Leptinotarsa decemlineata* Say), western corn rootworm (*Diabrotica virgifera* LeConte)); chafers and other beetles from the family Scarabaeidae (e.g., Japanese beetle (*Popillia japonica* Newman), oriental beetle (*Anomala orientalis* Waterhouse, *Exomala orientalis* (Waterhouse) Baraud), northern masked chafer (*Cyclocephala borealis* Arrow), southern masked chafer (*Cyclocephala immaculata* Olivier or *C. lurida* Bland), dung beetle and white grub (*Aphodius* spp.), black turfgrass ataenius (*Ataenius spretulus* Haldeman), green June beetle (*Cotinis nitida* Linnaeus), Asiatic garden beetle (*Maladera castanea* Arrow), May/June beetles (*Phyllophaga* spp.) and European chafer (*Rhizotrogus majalis* Razoumowsky)); carpet beetles from the family Dermestidae; wireworms from the family Elateridae; bark beetles from the family Scolytidae and flour beetles from the family Tenebrionidae.

35 In addition, agronomic and nonagronomic pests include: eggs, adults and larvae of the order Dermaptera including earwigs from the family Forficulidae (e.g., European earwig (*Forficula auricularia* Linnaeus), black earwig (*Chelisoches morio* Fabricius)); eggs, immatures, adults and nymphs of the orders Hemiptera and Homoptera such as, plant bugs

from the family Miridae, cicadas from the family Cicadidae, leafhoppers (e.g. *Empoasca* spp.) from the family Cicadellidae, potato leafhoppers, bed bugs (e.g., *Cimex lectularius* Linnaeus) from the family Cimicidae, planthoppers from the families Fulgoroidae and Delphacidae, treehoppers from the family Membracidae, psyllids from the family Psyllidae, whiteflies from the family Aleyrodidae, aphids from the family Aphididae, phylloxera from the family Phylloxeridae, mealybugs from the family Pseudococcidae, scales from the families Coccidae, Diaspididae and Margarodidae, lace bugs from the family Tingidae, stink bugs from the family Pentatomidae, chinch bugs (e.g., hairy chinch bug (*Blissus leucopterus hirtus* Montandon) and southern chinch bug (*Blissus insularis* Barber)) and other seed bugs from the family Lygaeidae, spittlebugs from the family Cercopidae squash bugs from the family Coreidae, and red bugs and cotton stainers from the family Pyrrhocoridae.

Agronomic and nonagronomic pests also include : eggs, larvae, nymphs and adults of the order Acari (mites) such as spider mites and red mites in the family Tetranychidae (e.g., European red mite (*Panonychus ulmi* Koch), two spotted spider mite (*Tetranychus urticae* Koch), McDaniel mite (*Tetranychus mcdanieli* McGregor)); flat mites in the family Tenuipalpidae (e.g., citrus flat mite (*Brevipalpus lewisi* McGregor)); rust and bud mites in the family Eriophyidae and other foliar feeding mites and mites important in human and animal health, i.e. dust mites in the family Epidermoptidae, follicle mites in the family Demodicidae, grain mites in the family Glycyphagidae; ticks in the family Ixodidae, commonly known as hard ticks (e.g., deer tick (*Ixodes scapularis* Say), Australian paralysis tick (*Ixodes holocyclus* Neumann), American dog tick (*Dermacentor variabilis* Say), lone star tick (*Amblyomma americanum* Linnaeus)) and ticks in the family Argasidae, commonly known as soft ticks (e.g., relapsing fever tick (*Ornithodoros turicata*), common fowl tick (*Argas radiatus*)); scab and itch mites in the families Psoroptidae, Pyemotidae, and Sarcoptidae; eggs, adults and immatures of the order Orthoptera including grasshoppers, locusts and crickets (e.g., migratory grasshoppers (e.g., *Melanoplus sanguinipes* Fabricius, *M. differentialis* Thomas), American grasshoppers (e.g., *Schistocerca americana* Drury), desert locust (*Schistocerca gregaria* Forskal), migratory locust (*Locusta migratoria* Linnaeus), bush locust (*Zonocerus* spp.), house cricket (*Acheta domesticus* Linnaeus), mole crickets (e.g., tawny mole cricket (*Scapteriscus vicinus* Scudder) and southern mole cricket (*Scapteriscus borellii* Giglio-Tos)); eggs, adults and immatures of the order Diptera including leafminers (e.g., *Liriomyza* spp. such as serpentine vegetable leafminer (*Liriomyza sativae* Blanchard)), midges, fruit flies (Tephritidae), frit flies (e.g., *Oscinella frit* Linnaeus), soil maggots, house flies (e.g., *Musca domestica* Linnaeus), lesser house flies (e.g., *Fannia canicularis* Linnaeus, *F. femoralis* Stein), stable flies (e.g., *Stomoxys calcitrans* Linnaeus), face flies, horn flies, blow flies (e.g., *Chrysomya* spp., *Phormia* spp.), and other muscoid fly pests, horse flies (e.g., *Tabanus* spp.), bot flies (e.g., *Gastrophilus* spp., *Oestrus* spp.), cattle grubs (e.g., *Hypoderma* spp.), deer flies (e.g., *Chrysops* spp.), keds (e.g., *Melophagus ovinus* Linnaeus) and other Brachycera,

mosquitoes (e.g., *Aedes* spp., *Anopheles* spp., *Culex* spp.), black flies (e.g., *Prosimulium* spp., *Simulium* spp.), biting midges, sand flies, sciarids, and other Nematocera; eggs, adults and immatures of the order Thysanoptera including onion thrips (*Thrips tabaci* Lindeman), flower thrips (*Frankliniella* spp.), and other foliar feeding thrips; insect pests of the order

5 Hymenoptera including ants of the Family Formicidae including the Florida carpenter ant (*Camponotus floridanus* Buckley), red carpenter ant (*Camponotus ferrugineus* Fabricius), black carpenter ant (*Camponotus pennsylvanicus* De Geer), white-footed ant (*Technomyrmex albipes* fr. Smith), big headed ants (*Pheidole* sp.), ghost ant (*Tapinoma melanocephalum* Fabricius); Pharaoh ant (*Monomorium pharaonis* Linnaeus), little fire ant (*Wasmannia*

10 *auropunctata* Roger), fire ant (*Solenopsis geminata* Fabricius), red imported fire ant (*Solenopsis invicta* Buren), Argentine ant (*Iridomyrmex humilis* Mayr), crazy ant (*Paratrechina longicornis* Latreille), pavement ant (*Tetramorium caespitum* Linnaeus), cornfield ant (*Lasius alienus* Förster) and odorous house ant (*Tapinoma sessile* Say). Other Hymenoptera including bees (including carpenter bees), hornets, yellow jackets, wasps, and

15 sawflies (*Neodiprion* spp.; *Cephus* spp.); insect pests of the order Isoptera including termites in the Termitidae (e.g., *Macrotermes* sp., *Odontotermes obesus* Rambur), Kalotermitidae (e.g., *Cryptotermes* sp.), and Rhinotermitidae (e.g., *Reticulitermes* sp., *Coptotermes* sp., *Heterotermes tenuis* Hagen) families, the eastern subterranean termite (*Reticulitermes flavipes* Kollar), western subterranean termite (*Reticulitermes hesperus* Banks), Formosan

20 subterranean termite (*Coptotermes formosanus* Shiraki), West Indian drywood termite (*Incisitermes immigrans* Snyder), powder post termite (*Cryptotermes brevis* Walker), drywood termite (*Incisitermes snyderi* Light), southeastern subterranean termite (*Reticulitermes virginicus* Banks), western drywood termite (*Incisitermes minor* Hagen), arboreal termites such as *Nasutitermes* sp. and other termites of economic importance; insect

25 pests of the order Thysanura such as silverfish (*Lepisma saccharina* Linnaeus) and firebrat (*Thermobia domestica* Packard); insect pests of the order Mallophaga and including the head louse (*Pediculus humanus capitis* De Geer), body louse (*Pediculus humanus* Linnaeus), chicken body louse (*Menacanthus stramineus* Nitzsch), dog biting louse (*Trichodectes canis* De Geer), fluff louse (*Goniocotes gallinae* De Geer), sheep body louse (*Bovicola ovis*

30 Schrank), short-nosed cattle louse (*Haematopinus eurysternus* Nitzsch), long-nosed cattle louse (*Linognathus vituli* Linnaeus) and other sucking and chewing parasitic lice that attack man and animals; insect pests of the order Siphonoptera including the oriental rat flea (*Xenopsylla cheopis* Rothschild), cat flea (*Ctenocephalides felis* Bouche), dog flea (*Ctenocephalides canis* Curtis), hen flea (*Ceratophyllus gallinae* Schrank), sticktight flea

35 (*Echidnophaga gallinacea* Westwood), human flea (*Pulex irritans* Linnaeus) and other fleas afflicting mammals and birds. Additional arthropod pests covered include: spiders in the order Araneae such as the brown recluse spider (*Loxosceles reclusa* Gertsch & Mulaik) and

the black widow spider (*Latrodectus mactans* Fabricius), and centipedes in the order Scutigermorpha such as the house centipede (*Scutigera coleoptrata* Linnaeus).

5 Examples of invertebrate pests of stored grain include larger grain borer (*Prostephanus truncatus*), lesser grain borer (*Rhyzopertha dominica*), rice weevil (*Stiophilus oryzae*), maize weevil (*Stiophilus zeamais*), cowpea weevil (*Callosobruchus maculatus*), red flour beetle (*Tribolium castaneum*), granary weevil (*Stiophilus granarius*), Indian meal moth (*Plodia interpunctella*), Mediterranean flour beetle (*Ephestia kuhniella*) and flat or rusty grain beetle (*Cryptolestis ferrugineus*).

10 Compositions of the present disclosure may have activity on members of the Classes Nematoda, Cestoda, Trematoda, and Acanthocephala including economically important members of the orders Strongylida, Ascaridida, Oxyurida, Rhabditida, Spirurida, and Enoplida such as but not limited to economically important agricultural pests (i.e. root knot nematodes in the genus *Meloidogyne*, lesion nematodes in the genus *Pratylenchus*, stubby root nematodes in the genus *Trichodorus*, etc.) and animal and human health pests (i.e. all
15 economically important flukes, tapeworms, and roundworms, such as *Strongylus vulgaris* in horses, *Toxocara canis* in dogs, *Haemonchus contortus* in sheep, *Dirofilaria immitis* Leidy in dogs, *Anoplocephala perfoliata* in horses, *Fasciola hepatica* Linnaeus in ruminants, etc.).

20 Compositions of the disclosure may have activity against pests in the order Lepidoptera (e.g., *Alabama argillacea* Hübner (cotton leaf worm), *Archips argyrospila* Walker (fruit tree leaf roller), *A. rosana* Linnaeus (European leaf roller) and other *Archips* species, *Chilo suppressalis* Walker (rice stem borer), *Cnaphalocrosis medinalis* Guenée (rice leaf roller), *Crambus caliginosellus* Clemens (corn root webworm), *Crambus teterrellus* Zincken (bluegrass webworm), *Cydia pomonella* Linnaeus (codling moth), *Earias insulana* Boisduval (spiny bollworm), *Earias vittella* Fabricius (spotted bollworm), *Helicoverpa armigera*
25 Hübner (American bollworm), *Helicoverpa zea* Boddie (corn earworm), *Heliothis virescens* Fabricius (tobacco budworm), *Herpetogramma licarsisalis* Walker (sod webworm), *Lobesia botrana* Denis & Schiffermüller (grape berry moth), *Pectinophora gossypiella* Saunders (pink bollworm), *Phyllocnistis citrella* Stainton (citrus leafminer), *Pieris brassicae* Linnaeus (large white butterfly), *Pieris rapae* Linnaeus (small white butterfly), *Plutella xylostella* Linnaeus
30 (diamondback moth), *Spodoptera exigua* Hübner (beet armyworm), *Spodoptera litura* Fabricius (tobacco cutworm, cluster caterpillar), *Spodoptera frugiperda* J. E. Smith (fall armyworm), *Trichoplusia ni* Hübner (cabbage looper) and *Tuta absoluta* Meyrick (tomato leafminer)).

35 Compositions of the disclosure have significant activity on members from the order Homoptera including: *Acyrtosiphon pisum* Harris (pea aphid), *Aphis craccivora* Koch (cowpea aphid), *Aphis fabae* Scopoli (black bean aphid), *Aphis gossypii* Glover (cotton aphid, melon aphid), *Aphis pomi* De Geer (apple aphid), *Aphis spiraeicola* Patch (spirea aphid), *Aulacorthum solani* Kaltenbach (foxglove aphid), *Chaetosiphon fragaefolii* Cockerell

(strawberry aphid), *Diuraphis noxia* Kurdjumov/Mordvilko (Russian wheat aphid), *Dysaphis plantaginea* Paaserini (rosy apple aphid), *Eriosoma lanigerum* Hausmann (woolly apple aphid), *Hyalopterus pruni* Geoffroy (mealy plum aphid), *Lipaphis erysimi* Kaltentbach (turnip aphid), *Metopolophium dirrhodum* Walker (cereal aphid), *Macrosiphum euphorbiae* Thomas
 5 (potato aphid), *Myzus persicae* Sulzer (peach-potato aphid, green peach aphid), *Nasonovia ribisnigri* Mosley (lettuce aphid), *Pemphigus* spp. (root aphids and gall aphids), *Rhopalosiphum maidis* Fitch (corn leaf aphid), *Rhopalosiphum padi* Linnaeus (bird cherry-oat aphid), *Schizaphis graminum* Rondani (greenbug), *Sitobion avenae* Fabricius (English grain aphid), *Therioaphis maculata* Buckton (spotted alfalfa aphid), *Toxoptera aurantii* Boyer
 10 de Fonscolombe (black citrus aphid), and *Toxoptera citricida* Kirkaldy (brown citrus aphid); *Adelges* spp. (adelgids); *Phylloxera devastatrix* Pergande (pecan phylloxera); *Bemisia tabaci* Gennadius (tobacco whitefly, sweetpotato whitefly), *Bemisia argentifolii* Bellows & Perring (silverleaf whitefly), *Dialeurodes citri* Ashmead (citrus whitefly) and *Trialeurodes vaporariorum* Westwood (greenhouse whitefly); *Empoasca fabae* Harris (potato leafhopper),
 15 *Laodelphax striatellus* Fallen (smaller brown planthopper), *Macrolestes quadrilineatus* Forbes (aster leafhopper), *Nephotettix cincticeps* Uhler (green leafhopper), *Nephotettix nigropictus* Stål (rice leafhopper), *Nilaparvata lugens* Stål (brown planthopper), *Peregrinus maidis* Ashmead (corn planthopper), *Sogatella furcifera* Horvath (white-backed planthopper), *Sogatodes orizicola* Muir (rice delphacid), *Typhlocyba pomaria* McAtee white apple
 20 leafhopper, *Erythroneoura* spp. (grape leafhoppers); *Magicidada septendecim* Linnaeus (periodical cicada); *Icerya purchasi* Maskell (cottony cushion scale), *Quadraspidotus perniciosus* Comstock (San Jose scale); *Planococcus citri* Risso (citrus mealybug); *Pseudococcus* spp. (other mealybug complex); *Cacopsylla pyricola* Foerster (pear psylla), *Trioza diospyri* Ashmead (persimmon psylla).

25 Compositions of this disclosure also have activity on members from the order Hemiptera including: *Acrosternum hilare* Say (green stink bug), *Anasa tristis* De Geer (squash bug), *Blissus leucopterus* Say (chinch bug), *Cimex lectularius* Linnaeus (bed bug) *Corythuca gossypii* Fabricius (cotton lace bug), *Cyrtopeltis modesta* Distant (tomato bug), *Dysdercus suturellus* Herrich-Schäffer (cotton stainer), *Euchistus servus* Say (brown stink bug),
 30 *Euchistus variolarius* Palisot de Beauvois (one-spotted stink bug), *Graptosthetus* spp. (complex of seed bugs), *Halymorpha halys* Stål (brown marmorated stink bug), *Leptoglossus corculus* Say (leaf-footed pine seed bug), *Lygus lineolaris* Palisot de Beauvois (tarnished plant bug), *Nezara viridula* Linnaeus (southern green stink bug), *Oebalus pugnax* Fabricius (rice stink bug), *Oncopeltus fasciatus* Dallas (large milkweed bug), *Pseudatomoscelis seriatus*
 35 Reuter (cotton fleahopper). Other insect orders controlled by compounds of the disclosure include Thysanoptera (e.g., *Frankliniella occidentalis* Pergande (western flower thrips), *Scirtothrips citri* Moulton (citrus thrips), *Sericothrips variabilis* Beach (soybean thrips), and *Thrips tabaci* Lindeman (onion thrips); and the order Coleoptera (e.g., *Leptinotarsa*

decemlineata Say (Colorado potato beetle), *Epilachna varivestis* Mulsant (Mexican bean beetle) and wireworms of the genera *Agriotes*, *Athous* or *Limonius*).

Note that some contemporary classification systems place Homoptera as a suborder within the order Hemiptera.

5 Of note is use of compositions of this disclosure for controlling western flower thrips (*Frankliniella occidentalis*). Of note is use of compounds of this disclosure for controlling potato leafhopper (*Empoasca fabae*). Of note is use of compounds of this disclosure for controlling cotton melon aphid (*Aphis gossypii*). Of note is use of compounds of this disclosure for controlling diamondback moth (*Plutella xylostella* L.). Of note is use of
10 compounds of this disclosure for controlling silverleaf whitefly (*Bemisia argentifolii* Bellows & Perring).

Compounds of the present disclosure may also be useful for increasing vigor of a crop plant. This method comprises contacting the crop plant (e.g., foliage, flowers, fruit or roots) or the seed from which the crop plant is grown with compounds of Formula I or Formula II
15 in amount sufficient to achieve the desired plant vigor effect (i.e. biologically effective amount). Typically, compounds of Formula I or Formula II is applied in a formulated composition. Although compounds of Formula I or Formula II is often applied directly to the crop plant or its seed, it can also be applied to the locus of the crop plant, i.e. the environment of the crop plant, particularly the portion of the environment in close enough proximity to
20 allow compounds of Formula I or Formula II to migrate to the crop plant. The locus relevant to this method most commonly comprises the growth medium (i.e. medium providing nutrients to the plant), typically soil in which the plant is grown. Treatment of a crop plant to increase vigor of the crop plant thus comprises contacting the crop plant, the seed from which the crop plant is grown or the locus of the crop plant with a biologically effective amount of
25 compounds of Formula I or Formula II.

Increased crop vigor can result in one or more of the following observed effects: (a) optimal crop establishment as demonstrated by excellent seed germination, crop emergence and crop stand; (b) enhanced crop growth as demonstrated by rapid and robust leaf growth (e.g., measured by leaf area index), plant height, number of tillers (e.g., for rice), root mass and overall dry weight of vegetative mass of the crop; (c) improved crop yields, as
30 demonstrated by time to flowering, duration of flowering, number of flowers, total biomass accumulation (i.e. yield quantity) and/or fruit or grain grade marketability of produce (i.e. yield quality); (d) enhanced ability of the crop to withstand or prevent plant disease infections and arthropod, nematode or mollusk pest infestations; and (e) increased ability of the crop to
35 withstand environmental stresses such as exposure to thermal extremes, suboptimal moisture or phytotoxic chemicals.

The compounds of the present disclosure may increase the vigor of treated plants compared to untreated plants by killing or otherwise preventing feeding of phytophagous

invertebrate pests in the environment of the plants. In the absence of such control of phytophagous invertebrate pests, the pests reduce plant vigor by consuming plant tissues or sap, or transmitting plant pathogens such as viruses. Even in the absence of phytophagous invertebrate pests, the compounds of the disclosure may increase plant vigor by modifying metabolism of plants. Generally, the vigor of a crop plant will be most significantly increased by treating the plant with a compound of the disclosure if the plant is grown in a nonideal environment, i.e. an environment comprising one or more aspects adverse to the plant achieving the full genetic potential it would exhibit in an ideal environment.

Of note is a method for increasing vigor of a crop plant wherein the crop plant is grown in an environment comprising phytophagous invertebrate pests. Also of note is a method for increasing vigor of a crop plant wherein the crop plant is grown in an environment not comprising phytophagous invertebrate pests. Also of note is a method for increasing vigor of a crop plant wherein the crop plant is grown in an environment comprising an amount of moisture less than ideal for supporting growth of the crop plant. Of note is a method for increasing vigor of a crop plant wherein the crop is rice. Also of note is a method for increasing vigor of a crop plant wherein the crop is maize (corn). Also of note is a method for increasing vigor of a crop plant wherein the crop is soybean.

Compositions of this disclosure can also be mixed with one or more other biologically active compounds or agents including insecticides, fungicides, nematocides, bactericides, acaricides, herbicides, herbicide safeners, growth regulators such as insect molting inhibitors and rooting stimulants, chemosterilants, semiochemicals, repellents, attractants, pheromones, feeding stimulants, other biologically active compounds or entomopathogenic bacteria, virus or fungi to form a multi-component pesticide giving an even broader spectrum of agronomic and nonagronomic utility. Thus, the present disclosure also pertains to a composition comprising a biologically effective amount of at least one compound of Formula I or a compound of Formula II, or a combination of the foregoing compounds, at least one additional component selected from the group consisting of surfactants, solid diluents, and liquid diluents, and at least one additional biologically active compound or agent. For compositions of the present disclosure, the other biologically active compounds or agents can be formulated together with the present compounds, including compounds of Formula I or compounds of Formula II, or a combination of the foregoing compounds, to form a premix, or the other biologically active compounds or agents can be formulated separately from the present compounds, including compounds of Formula I or compounds of Formula II, and the two formulations combined together before application (e.g., in a spray tank) or, alternatively, applied in succession.

Examples of such biologically active compounds or agents with which compounds of this disclosure can be formulated are insecticides such as abamectin, acephate, acequinocyl, acetamiprid, acrinathrin, acynonapyr, afidopyropen ((1*S*,4*R*,4*aR*,6*S*,6*aS*,12*R*,12*aS*,12*bS*)-3-

[(cyclopropylcarbonyl)oxy]-1,3,4,4a,5,6,6a,12,12a,12b-decahydro-6,12-dihydroxy-
 4,6a,12b-trimethyl-11-oxo-9-(3-pyridinyl)-2*H*,11*H*-naphtho[2,1-*b*]pyrano[3,4-*e*]pyran-4-
 yl)methyl cyclopropanecarboxylate), amidoflumet, amitraz, avermectin, azadirachtin,
 azinphos-methyl, benfuracarb, bensultap, benzpyrimoxan, bifenthrin, kappa-bifenthrin,
 5 bifenazate, bistrifluron, borate, broflanilide, buprofezin, cadusafos, carbaryl, carbofuran,
 cartap, carzol, chlorantraniliprole, chlorfenapyr, chlorfluazuron, chlorprallethrin,
 chlorpyrifos, chlorpyrifos-*e*, chlorpyrifos-methyl, chromafenozide, clofentezin,
 chlorprallethrin, clothianidin, cyantraniliprole (Cyazypyr™) (3-bromo-1-(3-chloro-2-
 pyridinyl)-*N*-[4-cyano-2-methyl-6-[(methylamino)carbonyl]phenyl]-1*H*-pyrazole-5-
 10 carboxamide), cyclaniliprole (3-bromo-*N*-[2-bromo-4-chloro-6-[[1-(1-
 cyclopropylethyl)amino]carbonyl]phenyl]-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-
 carboxamide), cycloprothrin, cycloxaprid ((5*S*,8*R*)-1-[(6-chloro-3-pyridinyl)methyl]-
 2,3,5,6,7,8-hexahydro-9-nitro-5,8-Epoxy-1*H*-imidazo[1,2-*a*]azepine), cyenopyrafen,
 cyflumetofen, cyfluthrin, beta-cyfluthrin, cyhalodiamide, cyhalothrin, gamma-cyhalothrin,
 15 lambda-cyhalothrin, cypermethrin, alpha-cypermethrin, zeta-cypermethrin, cyromazine,
 deltamethrin, diafenthiuron, diazinon, dicloromesotiaz, dieldrin, diflubenzuron, dimefluthrin,
 dimehypo, dimethoate, dimpropridaz, dinotefuran, diofenolan, DiPel®, emamectin,
 emamectin benzoate, endosulfan, esfenvalerate, ethiprole, etofenprox, epsilon-metofluthrin,
 etoxazole, fenbutatin oxide, fenitrothion, fenothiocarb, fenoxycarb, fenpropathrin,
 20 fenvalerate, fipronil, flometoquin (2-ethyl-3,7-dimethyl-6-[4-(trifluoromethoxy)phenoxy]-4-
 quinolinyl methyl carbonate), flonicamid, fluazaindolizine, flubendiamide, flucythrinate,
 flufenerim, flufenoxuron, flufenoxystrobin (methyl (α *E*)-2-[[2-chloro-4-
 (trifluoromethyl)phenoxy]methyl]- α -(methoxymethylene)benzeneacetate), fluensulfone (5-
 chloro-2-[(3,4,4-trifluoro-3-buten-1-yl)sulfonyl]thiazole), fluhexafon, fluopyram, flupiprole
 25 (1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-5-[(2-methyl-2-propen-1-yl)amino]-4-
 [(trifluoromethyl)sulfinyl]-1*H*-pyrazole-3-carbonitrile), flupyradifurone (4-[[6-chloro-3-
 pyridinyl)methyl](2,2-difluoroethyl)amino]-2(5*H*)-furanone), flupyrimin, fluvalinate,
 tau-fluvalinate, fluxametamide, fonophos, formetanate, fosthiazate, gamma-cyhalothrin,
 halofenozide, heptafluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 2,2-
 30 dimethyl-3-[(1*Z*)-3,3,3-trifluoro-1-propen-1-yl]cyclopropanecarboxylate), hexaflumuron,
 hexythiazox, hydramethylnon, imidacloprid, indoxacarb, insecticidal soaps, isofenphos,
 isocycloseram, kappa-tefluthrin, lambda-cyhalothrin, lufenuron, malathion, meperfluthrin
 ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl (1*R*,3*S*)-3-(2,2-dichloroethenyl)-2,2-
 dimethylcyclopropanecarboxylate), metaflumizone, metaldehyde, methamidophos,
 35 methidathion, methiocarb, methomyl, methoprene, methoxychlor, metofluthrin,
 methoxyfenozide, epsilon-metofluthrin, epsilon-momfluorothrin, monocrotophos,
 monofluorothrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 3-(2-cyano-1-
 propen-1-yl)-2,2-dimethylcyclopropanecarboxylate), nicotine, nitenpyram, nithiazine,

novaluron, noviflumuron, oxamyl, oxazosulfyl, parathion, parathion-methyl, permethrin,
 phorate, phosalone, phosmet, phosphamidon, pirimicarb, profenofos, profluthrin, propargite,
 protrifenbute, pyflubumide (1,3,5-trimethyl-*N*-(2-methyl-1-oxopropyl)-*N*-[3-(2-
 5 4-methylpropyl)-4-[2,2,2-trifluoro-1-methoxy-1-(trifluoromethyl)ethyl]phenyl]-1*H*-pyrazole-
 4-carboxamide), pymetrozine, pyrafluprole, pyrethrin, pyridaben, pyridalyl, pyrfluquinazon,
 pyriminostrobin (methyl (αE)-2-[[[2-[(2,4-dichlorophenyl)amino]-6-(trifluoromethyl)-4-
 pyrimidinyl]oxy]methyl]- α -(methoxymethylene)benzeneacetate), pyriprole, pyriproxyfen,
 rotenone, ryanodine, silafluofen, spinetoram, spinosad, spirodiclofen, spiromesifen,
 spiropidion, spirotetramat, sulprofos, sulfoxaflor (*N*-[methyloxido[1-[6-(trifluoromethyl)-3-
 10 pyridinyl]ethyl]- λ^4 -sulfanylidene]cyanamide), tebufenozide, tebufenpyrad, teflubenzuron,
 tefluthrin, kappa-tefluthrin, terbufos, tetrachlorantraniliprole, tetrachlorvinphos, tetramethrin,
 tetramethylfluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 2,2,3,3-
 tetramethylcyclopropanecarboxylate), tetraniliprole, thiacloprid, thiamethoxam, thiodicarb,
 thiosultap-sodium, tioazafen (3-phenyl-5-(2-thienyl)-1,2,4-oxadiazole), tolfenpyrad,
 15 tralomethrin, triazamate, trichlorfon, triflumezopyrim (2,4-dioxo-1-(5-pyrimidinylmethyl)-3-
 [3-(trifluoromethyl)phenyl]-2*H*-pyrido[1,2-*a*]pyrimidinium inner salt), triflumuron,
 tyclopyrazoflor, zeta-cypermethrin, *Bacillus thuringiensis* delta-endotoxins,
 entomopathogenic bacteria, entomopathogenic viruses or entomopathogenic fungi.

Of note are insecticides such as abamectin, acetamiprid, acrinathrin, acynonapyr,
 20 afidopyropen, amitraz, avermectin, azadirachtin, benfuracarb, bensultap, bifenthrin,
 buprofezin, broflanilide, cadusafos, carbaryl, cartap, chlorantraniliprole, chloroprallethrin,
 chlorfenapyr, chlorpyrifos, clothianidin, cyantraniliprole, cyclaniliprole, cycloprothrin,
 cyfluthrin, beta-cyfluthrin, cyhalothrin, DiPel®, gamma-cyhalothrin, lambda-cyhalothrin,
 cypermethrin, alpha-cypermethrin, zeta-cypermethrin, cyromazine, deltamethrin, dieldrin,
 25 dinotefuran, diofenolan, emamectin, endosulfan, epsilon-metofluthrin, esfenvaterate,
 ethiprole, etofenprox, etoxazole, fenitrothion, fenothiocarb, fenoxycarb, fenvalerate, fipronil,
 flometoquin, fluxametamide, flonicamid, flubendiamide, fluensulfone, flufenoxuron,
 flufenoxystrobin, flufensulfone, flupiprole, flupyrimin, flupyradifurone, fluvalinate,
 formetanate, fosthiazate, gamma-cyhalothrin, heptafluthrin, hexaflumuron, hydramethylnon,
 30 imidacloprid, indoxacarb, isocycloseram, kappa-tefluthrin, lambda-cyhalothrin, lufenuron,
 meperfluthrin, metaflumizone, methiodicarb, methomyl, methoprene, methoxyfenozide,
 metofluthrin, monofluorothrin, nitenpyram, nithiazine, novaluron, oxamyl, pyflubumide,
 pymetrozine, pyrethrin, pyridaben, pyridalyl, pyriminostrobin, pyriproxyfen, ryanodine,
 spinetoram, spinosad, spirodiclofen, spiromesifen, spirotetramat, sulfoxaflor, tebufenozide,
 35 tetramethrin, tetramethylfluthrin, thiacloprid, thiamethoxam, thiodicarb, thiosultap-sodium,
 tralomethrin, triazamate, triflumezopyrim, triflumuron, tyclopyrazoflor, zeta-cypermethrin,
Bacillus thuringiensis delta-endotoxins, all strains of *Bacillus thuringiensis* and all strains of
 nucleo polyhedrosis viruses.

One embodiment of biological agents for mixing with compounds of this disclosure include entomopathogenic bacteria such as *Bacillus thuringiensis*, and the encapsulated delta-endotoxins of *Bacillus thuringiensis* such as MVP[®] and MVPII[®] bioinsecticides prepared by the CellCap[®] process (CellCap[®], MVP[®] and MVPII[®] are trademarks of Mycogen Corporation, Indianapolis, Indiana, USA); entomopathogenic fungi such as green muscardine fungus; and entomopathogenic (both naturally occurring and genetically modified) viruses including baculovirus, nucleopolyhedro virus (NPV) such as *Helicoverpa zea* nucleopolyhedrovirus (HzNPV), *Anagrapha falcifera* nucleopolyhedrovirus (AfNPV); and granulosis virus (GV) such as *Cydia pomonella* granulosis virus (CpGV).

One embodiment of biological agents for mixing with compounds of this disclosure include one or a combination of (i) a bacterium of the genus *Actinomycetes*, *Agrobacterium*, *Arthrobacter*, *Alcaligenes*, *Aureobacterium*, *Azobacter*, *Bacillus*, *Beijerinckia*, *Bradyrhizobium*, *Brevibacillus*, *Burkholderia*, *Chromobacterium*, *Clostridium*, *Clavibacter*, *Comamonas*, *Corynebacterium*, *Curtobacterium*, *Enterobacter*, *Flavobacterium*, *Gluconobacter*, *Hydrogenophaga*, *Klebsiella*, *Methylobacterium*, *Paenibacillus*, *Pasteuria*, *Photorhabdus*, *Phyllobacterium*, *Pseudomonas*, *Rhizobium*, *Serratia*, *Sphingobacterium*, *Stenotrophomonas*, *Streptomyces*, *Variovorax*, or *Xenorhabdus*, for example a bacterium of *Bacillus amyloliquefaciens*, *Bacillus cereus*, *Bacillus firmus*, *Bacillus licheniformis*, *Bacillus pumilus*, *Bacillus sphaericus*, *Bacillus subtilis*, *Bacillus thuringiensis*, *Bradyrhizobium japonicum*, *Chromobacterium subtsugae*, *Pasteuria nishizawae*, *Pasteuria penetrans*, *Pasteuria usage*, *Pseudomonas fluorescens*, and *Streptomyces lydicus*; (ii) a fungus such as green muscardine fungus; (iii) a virus including baculovirus, nucleopolyhedro virus such as *Helicoverpa zea* nucleopolyhedrovirus, *Anagrapha falcifera* nucleopolyhedrovirus; granulosis virus such as *Cydia pomonella* granulosis virus.

Of particular note is such a combination where the other invertebrate pest control active ingredient belongs to a different chemical class or has a different site of action than compounds of Formula I or Formula II. In certain instances, a combination with at least one other invertebrate pest control active ingredient having a similar spectrum of control but a different site of action will be particularly advantageous for resistance management. Thus, a composition of the present disclosure can further comprise a biologically effective amount of at least one additional invertebrate pest control active ingredient having a similar spectrum of control but belonging to a different chemical class or having a different site of action. These additional biologically active compounds or agents include, but are not limited to, acetylcholinesterase (AChE) inhibitors such as the carbamates methomyl, oxamyl, thiodicarb, triazamate, and the organophosphates chlorpyrifos; GABA-gated chloride channel antagonists such as the cyclodienes dieldrin and endosulfan, and the phenylpyrazoles ethiprole and fipronil; sodium channel modulators such as the pyrethroids bifenthrin, cyfluthrin, *beta*-cyfluthrin, cyhalothrin, *lambda*-cyhalothrin, cypermethrin, deltamethrin, dimefluthrin,

esfenvalerate, metofluthrin and profluthrin; nicotinic acetylcholinereceptor (nAChR) agonists such as the neonicotinoids acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, nithiazine, thiacloprid, and thiamethoxam, and sulfoxaflor; nicotinic acetylcholine receptor (nAChR) allosteric activators such as the spinosyns spinetoram and spinosad; chloride channel activators such as the avermectins abamectin and emamectin; juvenile hormone mimics such as diofenolan, methoprene, fenoxycarb and pyriproxyfen; selective homopteran feeding blockers such as pymetrozine and flonicamid; mite growth inhibitors such as etoxazole; inhibitors of mitochondrial ATP synthase such as propargite; uncouplers of oxidative phosphorylation via disruption of the proton gradient such as chlorfenapyr; nicotinic acetylcholine receptor (nAChR) channel blockers such as the nereistoxin analogs cartap; inhibitors of chitin biosynthesis such as the benzoylureas flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron and triflumuron, and buprofezin; dipteran moulting disrupters such as cyromazine; ecdysone receptor agonists such as the diacylhydrazines methoxyfenozide and tebufenozide; octopamine receptor agonists such as amitraz; mitochondrial complex III electron transport inhibitors such as hydramethylnon; mitochondrial complex I electron transport inhibitors such as pyridaben; voltage-dependent sodium channel blockers such as indoxacarb; inhibitors of acetyl CoA carboxylase such as the tetronic and tetramic acids spiroadiclofen, spiromesifen and spirotetramat; mitochondrial complex II electron transport inhibitors such as the β -ketonitriles cyenopyrafen and cyflumetofen; ryanidine receptor modulators such as the anthranilic diamides chlorantraniliprole, cyantraniliprole and cyantraniliprole, diamides such as flubendiamide, and ryanodine receptor ligands such as ryanodine; compounds wherein the target site responsible for biological activity is unknown or uncharacterized such as azadirachtin, bifenazate, pyridalyl, pyrifluquinazon and triflumezopyrim; microbial disrupters of insect midgut membranes such as *Bacillus thuringiensis* and the delta-endotoxins they produce and *Bacillus sphaericus*; and biological agents including nucleopolyhedroviruses (NPV) and other naturally occurring or genetically modified insecticidal viruses.

Further examples of biologically active compounds or agents with which compounds of this disclosure can be formulated are: fungicides such as acibenzolar-S-methyl, aldimorph, ametocradin, aminopyrifin, amisulbrom, anilazine, azaconazole, azoxystrobin, benalaxyl (including benalaxyl-M), benodanil, benomyl, bentiavalicarb (including bentiavalicarb-isopropyl), benzovindiflupyr, bethoxazin, binapacryl, biphenyl, bitertanol, bixafen, blastocidin-S, boscalid, bromuconazole, bupirimate, buthiobate, carboxin, carpropamid, captafol, captan, carbendazim, chloroneb, chlorothalonil, chlozolinate, copper hydroxide, copper oxychloride, copper sulfate, coumoxystrobin, cyazofamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, dichlobentiaxox, dichlofluanid, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, diflumerimol, dimethirimol, dimethomorph, dimoxystrobin, diniconazole (including diniconazole-M), dinocap, dipymetitrone, dithianon, dithiolanes,

dodemorph, dodine, econazole, etaconazole, edifenphos, enoxastrobin (also known as enestroburin), epoxiconazole, ethaboxam, ethirimol, etridiazole, famoxadone, fenamidone, fenaminstrobin, fenarimol, fenbuconazole, fenfuram, fenhexamide, fenoxanil, fencpiclonil, fencpicoxamid, fenpropidin, fenpropimorph, fenpyrazamine, fentin acetate, fentin hydroxide, 5 ferbam, ferimzone, flometoquin, florylpicoxamid, fluopimomide, fluazinam, flubeneteram, fludioxonil, flufenoxystrobin, fluindapyr, flumorph, fluopicolide, fluopyram, fluoxapiprolin, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutianil, flutolanil, flutriafol, fluxapyroxad, folpet, fthalide (also known as phthalide), fuberidazole, furalaxyl, furametpyr, hexaconazole, hymexazole, guazatine, imazalil, imibenconazole, iminoctadine albesilate, 10 iminoctadine triacetate, inpyrfluxam, iodcarb, ipconazole, ipfentrifluconazole, ipflufenquin, isofetamid, iprobenfos, iprodione, iprovalicarb, isoflucypram, isoprothiolane, isopyrazam, isotianil, kasugamycin, kresoxim-methyl, lancotrione, mancozeb, mandipropamid, mandestrobin, maneb, mapanipyrin, mefentrifluconazole, mepronil, meptyldinocap, metalaxyl (including metalaxyl-M/mefenoxam), metconazole, methasulfocarb, metiram, 15 metominostrobin, metyltetraprole, metrafenone, myclobutanil, naftitine, neo-asozin (ferric methanearsonate), nuarimol, octhilinone, ofurace, oryastrobin, oxadixyl, oxathiapiprolin, oxolinic acid, oxpoconazole, oxycarboxin, oxytetracycline, penconazole, pencycuron, penflufen, penthiopyrad, perfurazoate, phosphorous acid (including salts thereof, e.g., fosetyl-aluminm), picoxystrobin, piperalin, polyoxin, probenazole, prochloraz, procymidone, 20 propamocarb, propiconazole, propineb, proquinazid, prothiocarb, prothioconazole, pydiflumetofen (Adepidyn®), pyraclostrobin, pyrametostrobin, pyrapropoyne, pyraoxystrobin, pyraziflumid, pyrazophos, pyribencarb, pyributacarb, pyridachlometyl, pyrifenox, pyriofenone, perisoxazole, pyrimethanil, pyrifenox, pyrrolnitrin, pyroquilon, quinconazole, quinmethionate, quinofumelin, quinoxifen, quintozone, silthiofam, sedaxane, 25 simeconazole, spiroxamine, streptomycin, sulfur, tebuconazole, tebufloquin, teclofthalam, tecloftalam, tecnazene, terbinafine, tetraconazole, thiabendazole, thifluzamide, thiophanate, thiophanate-methyl, thiram, tiadinil, tolclofos-methyl, tolprocarb, tolyfluanid, triadimefon, triadimenol, triarimol, triazoxide, tribasic copper sulfate, triclopyricarb, tridemorph, trifloxystrobin, triflumizole, trimoprhamide, tricyclazole, trifloxystrobin, triforine, 30 triticonazole, uniconazole, validamycin, valifenalate (also known as valifenal), vinclozolin, zineb, ziram, zoxamide and 1-[4-[4-[5-(2,6-difluorophenyl)-4,5-dihydro-3-isoxazolyl]-2-thiazolyl]-1-piperidinyl]-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone; nematocides such as fluopyram, spirotetramat, thiodicarb, fosthiazate, abamectin, iprodione, fluensulfone, dimethyl disulfide, tioazafen, 1,3-dichloropropene (1,3-D), metam (sodium and potassium), dazomet, chloropicrin, fenamiphos, ethoprophos, cadusaphos, terbufos, 35 imicyafos, oxamyl, carbofuran, tioazafen, *Bacillus firmus* and *Pasteuria nishizawae*; bactericides such as streptomycin; acaricides such as amitraz, chinomethionat,

chlorobenzilate, cyhexatin, dicofol, dienochlor, etoxazole, fenazaquin, fenbutatin oxide, fenpropathrin, fenpyroximate, hexythiazox, propargite, pyridaben and tebufenpyrad.

In certain instances, combinations of a compound of this disclosure with other biologically active (particularly invertebrate pest control) compounds or agents (i.e. active ingredients) can result in a greater-than-additive (i.e. synergistic) effect. Reducing the quantity of active ingredients released in the environment while ensuring effective pest control is always desirable. When synergism of invertebrate pest control active ingredients occurs at application rates giving agronomically satisfactory levels of invertebrate pest control, such combinations can be advantageous for reducing crop production cost and decreasing environmental load.

Compounds of this disclosure and compositions thereof can be applied to plants genetically transformed to express proteins toxic to invertebrate pests (such as *Bacillus thuringiensis* delta-endotoxins). Such an application may provide a broader spectrum of plant protection and be advantageous for resistance management. The effect of the exogenously applied invertebrate pest control compounds of this disclosure may be synergistic with the expressed toxin proteins.

General references for these agricultural protectants (i.e. insecticides, fungicides, nematocides, acaricides, herbicides and biological agents) include *The Pesticide Manual, 13th Edition*, C. D. S. Tomlin, Ed., British Crop Protection Council, Farnham, Surrey, U.K., 2003 and *The BioPesticide Manual, 2nd Edition*, L. G. Copping, Ed., British Crop Protection Council, Farnham, Surrey, U.K., 2001.

Invertebrate pests are controlled in agronomic and nonagronomic applications by applying one or more compounds of this disclosure, typically in the form of a composition, in a biologically effective amount, to the environment of the pests, including the agronomic and/or nonagronomic locus of infestation, to the area to be protected, or directly on the pests to be controlled.

Thus, the present disclosure comprises a method for controlling an invertebrate pest in agronomic and/or nonagronomic applications, comprising contacting the invertebrate pest or its environment with a biologically effective amount of one or more of the compounds of the disclosure, or with a composition comprising at least one such compound or a composition comprising at least one such compound and a biologically effective amount of at least one additional biologically active compound or agent. Examples of suitable compositions comprising a compound of the disclosure and a biologically effective amount of at least one additional biologically active compound or agent include granular compositions wherein the additional active compound is present on the same granule as the compound of the disclosure or on granules separate from those of the compound of the disclosure.

To achieve contact with a compound or composition of the disclosure to protect a field crop from invertebrate pests, the compound or composition is typically applied to the seed of

the crop before planting, to the foliage (e.g., leaves, stems, flowers, fruits) of crop plants, or to the soil or other growth medium before or after the crop is planted.

One embodiment of a method of contact is by spraying. Alternatively, a granular composition comprising a compound of the disclosure can be applied to the plant foliage or the soil. Compounds of this disclosure can also be effectively delivered through plant uptake by contacting the plant with a composition comprising a compound of this disclosure applied as a soil drench of a liquid formulation, a granular formulation to the soil, a nursery box treatment or a dip of transplants. Of note is a composition of the present disclosure in the form of a soil drench liquid formulation. Also of note is a method for controlling an invertebrate pest comprising contacting the invertebrate pest or its environment with a biologically effective amount of a compound of the present disclosure or with a composition comprising a biologically effective amount of a compound of the present disclosure. Of further note is this method wherein the environment is soil and the composition is applied to the soil as a soil drench formulation. Of further note is that compounds of this disclosure are also effective by localized application to the locus of infestation. Other methods of contact include application of a compound or a composition of the disclosure by direct and residual sprays, aerial sprays, gels, seed coatings, microencapsulations, systemic uptake, baits, ear tags, boluses, foggers, fumigants, aerosols, dusts and many others. One embodiment of a method of contact is a dimensionally stable fertilizer granule, stick or tablet comprising a compound or composition of the disclosure. The compounds of this disclosure can also be impregnated into materials for fabricating invertebrate control devices (e.g., insect netting).

Compounds of the disclosure are useful in treating all plants, plant parts and seeds. Plant and seed varieties and cultivars can be obtained by conventional propagation and breeding methods or by genetic engineering methods. Genetically modified plants or seeds (transgenic plants or seeds) are those in which a heterologous gene (transgene) has been stably integrated into the plant's or seed's genome. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

Genetically modified plant and seed cultivars which can be treated according to the disclosure include those that are resistant against one or more biotic stresses (pests such as nematodes, insects, mites, fungi, etc.) or abiotic stresses (drought, cold temperature, soil salinity, etc.), or that contain other desirable characteristics. Plants and seeds can be genetically modified to exhibit traits of, for example, herbicide tolerance, insect-resistance, modified oil profiles or drought tolerance.

Treatment of genetically modified plants and seeds with compounds of the disclosure may result in super-additive or synergistic effects. For example, reduction in application rates, broadening of the activity spectrum, increased tolerance to biotic/abiotic stresses or enhanced storage stability may be greater than expected from just simple additive effects of the application of compounds of the disclosure on genetically modified plants and seeds.

Compounds of this disclosure are also useful in seed treatments for protecting seeds from invertebrate pests. In the context of the present disclosure and claims, treating a seed means contacting the seed with a biologically effective amount of a compound of this disclosure, which is typically formulated as a composition of the disclosure. This seed treatment protects the seed from invertebrate soil pests and generally can also protect roots and other plant parts in contact with the soil of the seedling developing from the germinating seed. The seed treatment may also provide protection of foliage by translocation of the compound of this disclosure or a second active ingredient within the developing plant. Seed treatments can be applied to all types of seeds, including those from which plants genetically transformed to express specialized traits will germinate. Representative examples include those expressing proteins toxic to invertebrate pests, such as *Bacillus thuringiensis* toxin or those expressing herbicide resistance such as glyphosate acetyltransferase, which provides resistance to glyphosate. Seed treatments with compounds of this disclosure can also increase vigor of plants growing from the seed.

One method of seed treatment is by spraying or dusting the seed with a compound of the disclosure (i.e. as a formulated composition) before sowing the seeds. Compositions formulated for seed treatment generally comprise a film former or adhesive agent. Therefore, typically a seed coating composition of the present disclosure comprises a biologically effective amount of compounds of Formula I or Formula II, and a film former or adhesive agent. Seed can be coated by spraying a flowable suspension concentrate directly into a tumbling bed of seeds and then drying the seeds. Alternatively, other formulation types such as wetted powders, solutions, suspoemulsions, emulsifiable concentrates and emulsions in water can be sprayed on the seed. This process is particularly useful for applying film coatings on seeds. Various coating machines and processes are available to one skilled in the art. Suitable processes include those listed in P. Kosters et al., *Seed Treatment: Progress and Prospects*, 1994 BCPC Mongraph No. 57, and references listed therein.

Compounds of Formula I or Formula II and their compositions, both alone and in combination with other insecticides, nematicides, and fungicides, are particularly useful in seed treatment for crops including, but not limited to, maize or corn, soybeans, cotton, cereal (e.g., wheat, oats, barley, rye and rice), potatoes, vegetables and oilseed rape.

Other insecticides with which compounds of Formula I or Formula II can be formulated to provide mixtures useful in seed treatment include abamectin, acetamiprid, acrinathrin, amitraz, avermectin, azadirachtin, bensultap, bifenthrin, buprofezin, cadusafos, carbaryl, carbofuran, cartap, chlorantraniliprole, chlorfenapyr, chlorpyrifos, clothianidin, cyantraniliprole, cyfluthrin, beta-cyfluthrin, cyhalothrin, gamma-cyhalothrin, lambda-cyhalothrin, cypermethrin, alpha-cypermethrin, zeta-cypermethrin, cyromazine, deltamethrin, dieldrin, dinotefuran, diofenolan, DiPel®, emamectin, endosulfan, esfenvalerate, ethiprole, etofenprox, etoxazole, fenothiocarb, fenoxycarb, fenvalerate, fipronil, flonicamid,

flubendiamide, flufenoxuron, fluvalinate, formetanate, fosthiazate, hexaflumuron, hydramethylnon, imidacloprid, indoxacarb, lufenuron, metaflumizone, methiocarb, methomyl, methoprene, methoxyfenozide, nitenpyram, nithiazine, novaluron, oxamyl, pymetrozine, pyrethrin, pyridaben, pyridalyl, pyriproxyfen, ryanodine, spinetoram, spinosad, 5 spirodiclofen, spiromesifen, spirotetramat, sulfoxaflor, tebufenozide, tetramethrin, thiacloprid, thiamethoxam, thiodicarb, thiosultap-sodium, tralomethrin, triazamate, triflumuron, *Bacillus thuringiensis* delta-endotoxins, all strains of *Bacillus thuringiensis* and all strains of nucleopolyhedrovirus viruses.

Fungicides with which compounds of Formula I or Formula II can be formulated to 10 provide mixtures useful in seed treatment include amisulbrom, azoxystrobin, boscalid, carbendazim, carboxin, cymoxanil, cyproconazole, difenoconazole, dimethomorph, fluazinam, fludioxonil, fluquinconazole, fluopicolide, fluoxastrobin, flutriafol, fluxapyroxad, ipconazole, iprodione, metalaxyl, mefenoxam, metconazole, myclobutanil, paclobutrazole, penflufen, picoxystrobin, prothioconazole, pyraclostrobin, sedaxane, silthiofam, 15 tebuconazole, thiabendazole, thiophanate-methyl, thiram, trifloxystrobin and triticonazole.

Compositions comprising compounds of Formula I or Formula II useful for seed treatment can further comprise bacteria and fungi that have the ability to provide protection from the harmful effects of plant pathogenic fungi or bacteria and/or soil born animals such as nematodes. Bacteria exhibiting nematicidal properties may include but are not limited to 20 *Bacillus firmus*, *Bacillus cereus*, *Bacillus subtilis* and *Pasteuria penitans*. A suitable *Bacillus firmus* strain is strain CNCM I-1582 (GB-126) which is commercially available as BioNem™. A suitable *Bacillus cereus* strain is strain NCMM I-1592. Both *Bacillus* strains are disclosed in US 6,406,690. Other suitable bacteria exhibiting nematicidal activity are *B. amyloliquefaciens* IN937a and *B. subtilis* strain GB03. Bacteria exhibiting fungicidal 25 properties may include but are not limited to *B. pumilus* strain GB34. Fungal species exhibiting nematicidal properties may include but are not limited to *Myrothecium verrucaria*, *Paecilomyces lilacinus* and *Purpureocillium lilacinum*.

For embodiments where one or more of these various mixing partners are used, the weight ratio of these various mixing partners (in total) to the compound of Formula I or II, is 30 typically between about 1: 10000 and about 10000:1. 1:3000 and about 3000:1. Of note are weight ratios between about 1:300 and about 300:1 (for example ratios between about 1:30 and about 30:1).

Useful weight ratios of the at least one additional biological agent or pest control of the compound of Formula I or the compound of Formula II, or a salt thereof to component (b) in 35 the mixtures, compositions and methods of the present disclosure are typically from 10000:1 to 1:1000, from 1000:1 to 1: 500, from 100:1 to 1:100, from 20:1 to 1:20, from 10:1 to 1:10.

Of note are mixtures, compositions and methods wherein the weight ratio of component (a) the compound of Formula **I** or **II** to component (b) is from 1:150 to 200:1, from 1:150 to 50:1, from 1:50 to 10:1 or from 1:5 to 5:1.

It also is understood that any numerical range recited herein includes all values from the lower value to the upper value. For example, if a weight ratio range is stated as 1 : 50, it is intended that values such as 2 : 40, 10 : 30, or 1 : 3, etc., are expressly enumerated in this specification. These are only examples of what is specifically intended, and all possible combinations of numerical values between and including the lowest value and the highest value enumerated are to be considered to be expressly stated in this application.

One skilled in the art can easily determine through simple experimentation the biologically effective amounts of active ingredients necessary for the desired spectrum of biological activity. It will be evident that including these additional components can expand the spectrum of invertebrate pests controlled beyond the spectrum controlled by the compound of Formula **I** or **II** alone.

Table A lists specific combinations of a compound of Formula **I** or **II** with other invertebrate pest control agents illustrative of the mixtures, compositions and methods disclosed herein. The first column of Table A lists the specific invertebrate pest control agents (e.g., “Acetamiprid” in the first line). The second column of Table A lists embodiments of ranges of weight ratios of rates at which a compound of Formula **I** or **II** can be applied relative to an invertebrate pest control agent (e.g., “50:1 to 1:50” of compound of Formula **I** or **II** relative to acetamiprid by weight).

Thus, for example, the first line of Table A specifically discloses the combination of a compound of Formula **I** or **II** with acetamiprid can be applied in a weight ratio between 50:1 to 1:50. The remaining lines of Table A are to be construed similarly. Of further note Table A lists specific combinations of a compound of Formula **I** or **II** with other invertebrate pest control agents illustrative of the mixtures, compositions and methods of the present disclosure and includes additional embodiments of weight ratio ranges for application rates.

Table A

Invertebrate Pest Control Agent Component (b)	Typical Weight Ratio	Invertebrate Pest Control Agent Component (b)	Typical Weight Ratio
Acetamiprid	150:1 to 1:200	Flupyrimin	50:1 to 1:500
Acynonapyr	100:1 to 1:400	Indoxacarb	200:1 to 1:50
Avermectin	50:1 to 1:50	Imidacloprid	1000:1 to 1:1000
<i>Bacillus</i> spp. and any active crystal proteins	50:1 to 1:10	Isocycloseram	50:1 to 1:100

Invertebrate Pest Control Agent Component (b)	Typical Weight Ratio	Invertebrate Pest Control Agent Component (b)	Typical Weight Ratio
Benzpyrimoxan	150:1 to 1:200	Methomyl	250:1 to 1:100
Bifenthrin	100:1 to 1:10	Methoxyfenozide	500:1 to 1:100
kappa-bifenthrin	100:1 to 1:250	epsilon-Metofluthrin	200:1 to 1:100
Broflanilide,	150:1 to 1:500	Novaluron	100:1 to 1:200
Buprofezin,	500:1 to 1:50	Oxazosulflor	100:1 to 1:200
Carbofuran	200:1 to 1:100	Permethrin	100:1 to 1:120
Chlorantraniliprole	100:1 to 1:120	Pyriproxifen	250:1 to 1:100
Chlorfenapyr	100:1 to 1:10	Spiropidion	1200:1 to 1:200
Chlorprallethrin	50:1 to 1:500	Spirotetramat	150:1 to 1:100
Chlorpyrifos,	500:1 to 1:200	Sulfoxaflor	200:1 to 1:100
Clothianidin,	100:1 to 1:400	kappa-Tefluthrin	100:1 to 1:1000
Cyantraniliprole	100:1 to 1:120	Tetrachlorantraniliprole	200:1 to 1:100
γ -Cyhalothrin	50:1 to 1:250	Thiamethoxam	1250:1 to 1:1000
ζ -Cypermethrin	150:1 to 1:200	Tylocyprazoflor	200:1 to 1:500
Cyromazine	400:1 to 1:50	<i>Bacillus thuringiensis</i>	50:1 to 1:10
Diafenthiuron	200:1 to 1:150	Dimpropridaz	250:1 to 1:150
Dicloromezotiaz	200:1 to 1:150	Dinotefuran	150:1 to 1:200
Emamectin Benzoate	500:1 to 1:100	Flonicamid	200:1 to 1:100
Fipronil	150:1 to 1:100	Flupyradifurone	200:1 to 1:200

Embodiments of the disclosure include the composition of the present disclosure wherein the at least one additional biologically active compound or agent is selected from the Invertebrate Pest Control Agents listed in Table A above.

5 The weight ratios of a compound, including a compound of Formula I or II, to the additional invertebrate pest control agent typically are between 10000:1 and 1:1000, with one embodiment between 1000:1 and 1:500, with one embodiment being between 500:1 and 1:500, another embodiment being between 250:1 and 1:200 and another embodiment being between 100:1 and 1:50.

10 Listed below in Tables A-1 to A-5 are embodiments of specific compositions comprising a compound of Formula I or II (compound numbers (Cmp No.) refer to compounds in Table 1 and Tests A-E.

Table A-1

Mixture No.	Cmp No.	Invertebrate Pest Control Agent
A1-1	1	Acetamiprid
A1-2	1	Acynonapyr

A1-3	1	Avermectin
A1-4	1	<i>Bacillus</i> spp.
A1-5	1	Benzpyrimoxan
A1-6	1	Bifenthrin
A1-7	1	kappa-bifenthrin
A1-8	1	Broflanilide
A1-9	1	Buprofezin
A1-10	1	Carbofuran
A1-11	1	Chlorantraniliprole
A1-12	1	Chlorfenapyr
A1-13	1	Chloroprallethrin
A1-14	1	Chlorpyrifos,
A1-15	1	Clothianidin,
A1-16	1	Cyantraniliprole
A1-17	1	γ -Cyhalothrin

Mixture No.	Cmp No.	Invertebrate Pest Control Agent
A1-18	1	ζ-Cypermethrin
A1-19	1	Cyromazine
A1-20	1	Diafenthiuron
A1-21	1	Dicloromezotiaz
A1-22	1	Dimpropridaz
A1-23	1	Dinotefuran
A1-24	1	Emamectin Benzoate
A1-25	1	Fipronil
A1-26	1	Flonicamid
A1-27	1	Flupyradifurone
A1-28	1	Flupyrimin
A1-29	1	Indoxacarb
A1-30	1	Imidacloprid
A1-31	1	Isocycloseram
A1-32	1	Methomyl
A1-33	1	Methoxyfenozide
A1-34	1	epsilon-Metofluthrin
A1-35	1	Novaluron
A1-36	1	Oxazosulfyl
A1-37	1	Permethrin
A1-38	1	Pyriproxifen
A1-39	1	Spiropidion
A1-40	1	Spirotetramat
A1-41	1	Sulfoxaflor
A1-42	1	kappa-Tefluthrin
A1-43	1	Tetrachlorantraniliprole
A1-44	1	Thiamethoxam
A1-45	1	Tyclopyrazoflor
A1-46	1	<i>Bacillus thuringiensis</i>

Table A2

Table A2 is identical to Table A1, except that each reference to compound 1 in the column headed "Cmpd. No." is replaced by a reference to compound 2. For example, the first mixture in Table 4 is designated A2-1 and is a mixture of compound 2 and the additional
5 invertebrate pest control agent acetamiprid.

Table A3

Table A3 is identical to Table A1, except that each reference to compound 1 in the column headed "Cmpd. No." is replaced by a reference to compound 3. For example, the first mixture in Table 5 is designated A3-1 and is a mixture of compound 3 and the additional
10 invertebrate pest control agent acetamiprid.

Table A4

Table A4 is identical to Table A1, except that each reference to compound 1 in the column headed "Cmpd. No." is replaced by a reference to compound 4. For example, the first mixture in Table 6 is designated A4-1 and is a mixture of compound 4 and the additional
15 invertebrate pest control agent acetamiprid.

Table A5

Table A5 is identical to Table A1, except that each reference to compound 1 in the column headed "Cmpd. No." is replaced by a reference to compound 5. For example, the first mixture in Table 7 is designated A5-1 and is a mixture of compound 5 and the additional
20 invertebrate pest control agent acetamiprid.

Seed treatments can also include one or more nematicidal agents of natural origin such as the elicitor protein called harpin which is isolated from certain bacterial plant pathogens such as *Erwinia amylovora*. An example is the Harpin-N-Tek seed treatment technology
25 available as N-HibitTM Gold CST.

Seed treatments can also include one or more species of legume-root nodulating bacteria such as the microsymbiotic nitrogen-fixing bacteria *Bradyrhizobium japonicum*. These inoculants can optionally include one or more lipo-chitooligosaccharides (LCOs), which are nodulation (Nod) factors produced by rhizobia bacteria during the initiation of nodule
30 formation on the roots of legumes. For example, the Optimize® brand seed treatment technology incorporates LCO Promoter TechnologyTM in combination with an inoculant.

Seed treatments can also include one or more isoflavones which can increase the level of root colonization by mycorrhizal fungi. Mycorrhizal fungi improve plant growth by enhancing the root uptake of nutrients such as water, sulfates, nitrates, phosphates and metals.
35 Examples of isoflavones include, but are not limited to, genistein, biochanin A, formononetin, daidzein, glycitein, hesperetin, naringenin and pratensein. Formononetin is

available as an active ingredient in mycorrhizal inoculant products such as PHC Colonize® AG.

Seed treatments can also include one or more plant activators that induce systemic acquired resistance in plants following contact by a pathogen. An example of a plant activator which induces such protective mechanisms is acibenzolar-*S*-methyl.

The treated seed typically comprises a compound of the present disclosure in an amount from about 0.1 g to 1 kg per 100 kg of seed (i.e. from about 0.0001 to 1% by weight of the seed before treatment). A flowable suspension formulated for seed treatment typically comprises from about 0.5 to about 70% of the active ingredient, from about 0.5 to about 30% of a film-forming adhesive, from about 0.5 to about 20% of a dispersing agent, from 0 to about 5% of a thickener, from 0 to about 5% of a pigment and/or dye, from 0 to about 2% of an antifoaming agent, from 0 to about 1% of a preservative, and from 0 to about 75% of a volatile liquid diluent.

The compounds of this disclosure can be incorporated into a bait composition that is consumed by an invertebrate pest or used within a device such as a trap, bait station, and the like. Such a bait composition can be in the form of granules which comprise (a) active ingredients, namely a biologically effective amount of compounds of Formula I or Formula II (b) one or more food materials; optionally (c) an attractant, and optionally (d) one or more humectants. Of note are granules or bait compositions which comprise between about 0.001-5% active ingredients, about 40-99% food material and/or attractant; and optionally about 0.05-10% humectants, which are effective in controlling soil invertebrate pests at very low application rates, particularly at doses of active ingredient that are lethal by ingestion rather than by direct contact. Some food materials can function both as a food source and an attractant. Food materials include carbohydrates, proteins and lipids. Examples of food materials are vegetable flour, sugar, starches, animal fat, vegetable oil, yeast extracts and milk solids. Examples of attractants are odorants and flavorants, such as fruit or plant extracts, perfume, or other animal or plant component, pheromones or other agents known to attract a target invertebrate pest. Examples of humectants, i.e. moisture retaining agents, are glycols and other polyols, glycerine and sorbitol. Of note is a bait composition (and a method utilizing such a bait composition) used to control at least one invertebrate pest selected from the group consisting of ants, termites and cockroaches. A device for controlling an invertebrate pest can comprise the present bait composition and a housing adapted to receive the bait composition, wherein the housing has at least one opening sized to permit the invertebrate pest to pass through the opening so the invertebrate pest can gain access to the bait composition from a location outside the housing, and wherein the housing is further adapted to be placed in or near a locus of potential or known activity for the invertebrate pest.

One embodiment of the present disclosure relates to a method for controlling invertebrate pests, comprising diluting the pesticidal composition of the present disclosure

compounds of Formula **I** or Formula **II** formulated with surfactants, solid diluents and liquid diluents or a formulated mixture of compounds of Formula **I** or Formula **II** and at least one other pesticide) with water, and optionally adding an adjuvant to form a diluted composition, and contacting the invertebrate pest or its environment with an effective amount of said diluted composition.

Although a spray composition formed by diluting with water a sufficient concentration of the present pesticidal composition can provide sufficient efficacy for controlling invertebrate pests, separately formulated adjuvant products can also be added to spray tank mixtures. These additional adjuvants are commonly known as “spray adjuvants” or “tank-mix adjuvants”, and include any substance mixed in a spray tank to improve the performance of a pesticide or alter the physical properties of the spray mixture. Adjuvants can be surfactants, emulsifying agents, petroleum-based crop oils, crop-derived seed oils, acidifiers, buffers, thickeners or defoaming agents. Adjuvants are used to enhancing efficacy (e.g., biological availability, adhesion, penetration, uniformity of coverage and durability of protection), or minimizing or eliminating spray application problems associated with incompatibility, foaming, drift, evaporation, volatilization and degradation. To obtain optimal performance, adjuvants are selected with regard to the properties of the active ingredient, formulation and target (e.g., crops, insect pests).

Among the spray adjuvants, oils including crop oils, crop oil concentrates, vegetable oil concentrates and methylated seed oil concentrates are most commonly used to improve the efficacy of pesticides, possibly by means of promoting more even and uniform spray deposits. In situations where phytotoxicity potentially caused by oils or other water-immiscible liquids are of concern, spray compositions prepared from the composition of the present disclosure will generally not contain oil-based spray adjuvants. However, in situations where phytotoxicity caused by oil-based spray adjuvants is commercially insignificant, spray compositions prepared from the composition of the present composition can also contain oil-based spray adjuvants, which can potentially further increase control of invertebrate pests, as well as rainfastness.

Products identified as “crop oil” typically contain 95 to 98% paraffin or naphtha-based petroleum oil and 1 to 2% of one or more surfactants functioning as emulsifiers. Products identified as “crop oil concentrates” typically consist of 80 to 85% of emulsifiable petroleum-based oil and 15 to 20% of nonionic surfactants. Products correctly identified as “vegetable oil concentrates” typically consist of 80 to 85% of vegetable oil (i.e. seed or fruit oil, most commonly from cotton, linseed, soybean or sunflower) and 15 to 20% of nonionic surfactants. Adjuvant performance can be improved by replacing the vegetable oil with methyl esters of fatty acids that are typically derived from vegetable oils. Examples of methylated seed oil concentrates include MSO[®] Concentrate (UAP-Loveland Products, Inc.) and Premium MSO Methylated Spray Oil (Helena Chemical Company).

The amount of adjuvants added to spray mixtures generally does not exceed about 2.5% by volume, and more typically the amount is from about 0.1 to about 1% by volume. The application rates of adjuvants added to spray mixtures are typically between about 1 to 5 L per hectare. Representative examples of spray adjuvants include: Adigor[®] (Syngenta) 47% methylated rapeseed oil in liquid hydrocarbons, Silwet[®] (Helena Chemical Company) polyalkyleneoxide modified heptamethyltrisiloxane and Assist[®] (BASF) 17% surfactant blend in 83% paraffin based mineral oil.

The compounds of this disclosure can be applied without other adjuvants, but most often application will be of a formulation comprising one or more active ingredients with suitable carriers, diluents, and surfactants and possibly in combination with a food depending on the contemplated end use. One method of application involves spraying a water dispersion or refined oil solution of a compound of the present disclosure. Combinations with spray oils, spray oil concentrations, spreader stickers, adjuvants, other solvents, and synergists such as piperonyl butoxide often enhance compound efficacy. For nonagronomic uses such sprays can be applied from spray containers such as a can, a bottle or other container, either by means of a pump or by releasing it from a pressurized container, e.g., a pressurized aerosol spray can. Such spray compositions can take various forms, for example, sprays, mists, foams, fumes or fog. Such spray compositions thus can further comprise propellants, foaming agents, etc. as the case may be. Of note is a spray composition comprising a biologically effective amount of a compound or a composition of the present disclosure and a carrier. One embodiment of such a spray composition comprises a biologically effective amount of a compound or a composition of the present disclosure and a propellant. Representative propellants include, but are not limited to, methane, ethane, propane, butane, isobutane, butene, pentane, isopentane, neopentane, pentene, hydrofluorocarbons, chlorofluorocarbons, dimethyl ether, and mixtures of the foregoing. Of note is a spray composition (and a method utilizing such a spray composition dispensed from a spray container) used to control at least one invertebrate pest selected from the group consisting of mosquitoes, black flies, stable flies, deer flies, horse flies, wasps, yellow jackets, hornets, ticks, spiders, ants, gnats, and the like, including individually or in combinations.

The rate of application required for effective control (i.e. “biologically effective amount”) will depend on such factors as the species of invertebrate to be controlled, the pest’s life cycle, life stage, its size, location, time of year, host crop or animal, feeding behavior, mating behavior, ambient moisture, temperature, and the like. Under normal circumstances, application rates of about 0.01 to 2 kg of active ingredients per hectare are sufficient to control pests in agronomic ecosystems, but as little as 0.0001 kg/hectare may be sufficient or as much as 8 kg/hectare may be required. For non-agronomic applications, effective use rates will range from about 1.0 to 50 mg/square meter but as little as 0.1 mg/square meter may be sufficient or as much as 150 mg/square meter may be required. One skilled in the art can easily

determine the biologically effective amount necessary for the desired level of invertebrate pest control.

Enhanced activity has been described as “the cooperative action of two components (e.g., component (a) and component (b)) in a mixture, such that the total effect is greater or more prolonged than the sum of the effects of the two (or more) taken independently” (see 5 P. M. L. Tames, *Neth. J. Plant Pathology* 1964, 70, 73–80). Mixtures containing the compounds of Formula I together with other invertebrate pest control agents are found to exhibit enhanced effects against certain important invertebrate pests.

The presence of an enhanced effect between two active ingredients is established with 10 the aid of the Colby equation (see S. R. Colby, “Calculating Synergistic and Antagonistic Responses of Herbicide Combinations”, *Weeds*, 1967, 15, 20–22):

$$p = A + B - \left[\frac{A \times B}{100} \right]$$

15 Using the method of Colby, the presence of an enhanced interaction between two active ingredients is established by first calculating the predicted activity, p, of the mixture based on activities of the two components applied alone. If p is lower than the experimentally established effect, an enhanced interaction has occurred. If p is equal or higher than the experimentally established effect, the interaction between the two components is 20 characterized to be only additive or antagonism. In the equation above, A is the observed result of one component applied alone at rate x. The B term is the observed result of the second component applied at rate y. The equation estimates p, the expected result of the mixture of A at rate x with B at rate y. To use the Colby equation the active ingredients of the mixture are applied in the test separately as well as in combination.

25 All patents and patent applications mentioned in this application are incorporated by reference herein in their entireties for all purposes. In case of conflict between the present disclosure and that of a patent or publication incorporated by reference, the present disclosure controls.

The following non-limiting examples are purely illustrative.

30 BIOLOGICAL EXAMPLES

The following tests demonstrate the control efficacy of mixtures or compositions of this disclosure on specific pests. The pest control protection afforded by the mixtures or compositions is not limited, however, to these species. The analysis of enhanced activity between the mixtures or compositions was determined using Colby’s equation. The average 35 % mortality data for the test compounds alone were inserted into the Colby’s equation. If the observed (obs) average % mortality was higher than “p”, the expected (exp) % mortality, the mixture or composition had enhanced effects. If the observed average % mortality was equal

to or lower than the expected mortality, the mixture or composition either had no enhanced effect. For bioassays where insect feeding damage was evaluated, enhanced activity is identified when the observed plant damage is less than the expected plant damage rating; ratings ranged between 0 (no damage) and 10 (dead plant) and each value was converted to a
5 % plant protection.

$$100 - (\text{Obs Plant Damage} * 10)$$

Thus, a plant given a plant damage rating of 3 out of 10 is equivalent to 70% plant protection. In all of these tests, compounds of Formula I are Compounds 1 (Cpd 1), 2 (Cpd2), and 3 (Cpd3) and compounds of Formula II are compounds 4 (Cpd4) and 5 (Cpd 5).

10

TEST A

For evaluating control of cyantraniliprole, acetamiprid, imidacloprid, spirotetramat, spiroidiclofen, chlorantraniliprole, bifenthrin, indoxacarb, Avermectin, *Bacillus* spp. and any active crystal proteins, Buprofezin, Carbofuran, Chlorfenapyr, Chlorpyrifos, Clothianidin, Cyromazine, Diafenthiuron, Dinotefuran, Emamectin Benzoate, Fipronil, Flonicamid,
15 Flupyradifurone, methomyl (Lannate®), Methoxyfenozide, Novaluron, Permethrin, Pyriproxifen, Sulfoxaflor, Thiamethoxam, γ -Cyhalothrin, or ζ -cypermethrin, broflanilide, dimpropyridaz, isocycloseram, tetrachlorantraniliprole, oxazosulfonyl, tyclopyrazoflor, flupyrimin, spiropidion, acynonapyr, benzpyrimoxan, chloroprallethrin, epsilon-metofluthrin, kappa-bifenthrin, dicloromezotiaz, and kappa-tefluthrin.

20 For evaluating control of silverleaf whitefly (*Bemisia argentifolii* Bellows and Perring) through contact and/or systemic means, each test unit consisted of a small open container with a 12- to 14-day-old cotton plant or 5-7 day old soybean plant inside. This was pre-infested by placing test units into cages containing adult whiteflies so that oviposition on the cotton leaves could occur. The adults were removed from the plants with an air-blast nozzle, and the
25 test units were capped. The test units were then stored 2 to 3 days before spraying.

Test compounds were formulated using a solution containing 10% acetone, 90% water and 300 ppm Activator 90® Spreader Lo-Foam Formula non-ionic surfactant containing alkylaryl polyoxyethylene, free fatty acids, glycols and 2-propanol (Loveland Industries, Inc.) to provide the desired concentration in ppm. Formulated test solutions were then applied in
30 1 mL volumes through a SUJ2 atomizer nozzle with 1/8 JJ custom body (Spraying Systems Co.) positioned 1.27 cm (0.5 inches) above the top of each test unit.

The results for all experimental compositions in this test were replicated three times. After spraying of the formulated test composition, each test unit was allowed to dry for 1 hour and the cap removed. The test units were held for 13 days in a growth chamber at 28°C and
35 50–70% relative humidity. Each test unit was then assessed for insect mortality using a binocular microscope; the results are listed in Tables 2a-i.

Table 2a – Mixtures comprising Cyantraniliprole and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		5.9	
Cpd 4	250		11.1	
Cpd 2	50		10.9	
Cpd 2	250		74.5	
Cpd 1	50		0	
Cpd 1	250		18.8	
Cpd 3	50		25	
Cpd 3	250		1.6	
Cpd 5	50		17	
Cpd 5	250		13.5	
Cyantraniliprole	3		29.1	
Cyantraniliprole	5.5		61.1	
Cpd 4 + Cyantraniliprole	50 + 3	1:0.06	25	62.7
Cpd 4 + Cyantraniliprole	50 + 5.5	1:0.11	69.4	82.2
Cpd 4 + Cyantraniliprole	250 + 3	1:0.012	21.5	31.4
Cpd 4 + Cyantraniliprole	250 + 5.5	1:0.022	67.9	73.3
Cpd 2 + Cyantraniliprole	50 + 3	1:0.06	38.2*	26.5
Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 2 + Cyantraniliprole	50 + 5.5	1:0.11	66.7	76
Cpd 2 + Cyantraniliprole	250 + 3	1:0.012	94.8*	42.9
Cpd 2 + Cyantraniliprole	250 + 5.5	1:0.022	100.0*	84
Cpd 1 + Cyantraniliprole	50 + 3	1:0.06	27.9	37.3
Cpd 1 + Cyantraniliprole	50 + 5.5	1:0.11	89.8*	76.3
Cpd 1 + Cyantraniliprole	250 + 3	1:0.012	76.1*	16.2
Cpd 1 + Cyantraniliprole	250 + 5.5	1:0.022	77.1	77.4
Cpd 3 + Cyantraniliprole	50 + 3	1:0.06	24.6*	10.9
Cpd 3 + Cyantraniliprole	50 + 5.5	1:0.11	70.5	79.3
Cpd 3 + Cyantraniliprole	250 + 3	1:0.012	39.2*	25
Cpd 3 + Cyantraniliprole	250 + 5.5	1:0.022	71.6*	20.1
Cpd 5 + Cyantraniliprole	50 + 3	1:0.06	39.2*	37.7
Cpd 5 + Cyantraniliprole	50 + 5.5	1:0.11	45.8*	14.9
Cpd 5 + Cyantraniliprole	250 + 3	1:0.012	59.2*	41.2
Cpd 5 + Cyantraniliprole	250 + 5.5	1:0.022	68.2*	66.3

*Denotes enhanced efficacy based on Colby formula

Table 2b – Mixtures comprising Acetamiprid and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		46.7	
Cpd 4	250		20	
Cpd 2	50		80	
Cpd 2	250		53.3	
Cpd 1	50		80	
Cpd 1	250		0	
Cpd 3	50		0	
Cpd 3	250		0	
Cpd 5	50		49	
Cpd 5	250		19.2	
Acetamiprid	2.52		6.5	
Acetamiprid	6.88		25.4	
Cpd 4 + Acetamiprid	50 + 2.52	1:0.0504	65.7*	50.1
Cpd 4+ Acetamiprid	50 + 6.88	1:0.1376	43.6	60.2
Cpd 4 + Acetamiprid	250 + 2.52	1:0.01008	40.3*	25.2
Cpd 4 + Acetamiprid	250 + 6.88	1:0.02752	81.9*	40.3
Cpd 2 + Acetamiprid	50 + 2.52	1:0.0504	28	81.3
Cpd 2 + Acetamiprid	50 + 6.88	1:0.1376	53.1	85.1
Cpd 2 + Acetamiprid	250 + 2.52	1:0.01008	60.0*	56.3
Cpd 2 + Acetamiprid	250 + 6.88	1:0.02752	100.0*	65.2
Cpd1 + Acetamiprid	50 + 2.52	1:0.0504	34.1	81.3
Cpd 1 + Acetamiprid	50 + 6.88	1:0.1376	44.7*	25.4
Cpd 1 + Acetamiprid	250 + 2.52	1:0.01008	23.3*	6.5
Cpd1 + Acetamiprid	250 + 6.88	1:0.02752	62.3*	25.4
Cpd 3 + Acetamiprid	50 + 2.52	1:0.0504	20.5*	6.5
Cpd 3+ Acetamiprid	50 + 6.88	1:0.1376	40.4*	25.4
Cpd 3 + Acetamiprid	250 + 2.52	1:0.01008	21.6*	6.5
Cpd 3 + Acetamiprid	250 + 6.88	1:0.02752	62.2*	25.4
Cpd 5 + Acetamiprid	50 + 2.52	1:0.0504	45.3	52.3
Cpd 5 + Acetamiprid	50 + 6.88	1:0.1376	83.7*	61.9
Cpd 5 + Acetamiprid	250 + 2.52	1:0.01008	56.4*	24.4
Cpd 5 + Acetamiprid	250 + 6.88	1:0.02752	64.4*	39.7

*Denotes enhanced efficacy based on Colby formula

Table 2c – Mixtures comprising Imidacloprid and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		16.1	
Cpd 4	250		20.5	
Cpd 2	50		11.8	
Cpd 2	250		75.4	
Cpd 1	50		0	
Cpd 1	250		3.1	
Cpd 3	50		16.1	
Cpd 3	250		26.7	
Cpd 5	50		56.5	
Cpd 5	250		41.1	
Imidacloprid	15.1		0	
Imidacloprid	21.15		24.7	
Cpd 4 + Imidacloprid	50 + 15.1	1:0.302	50.0*	16.1
Cpd 4 + Imidacloprid	50 + 21.15	1:0.423	39.7*	36.8
Cpd 4 + Imidacloprid	250 + 15.1	1:0.0604	0	20.5
Cpd 4 + Imidacloprid	250 + 21.15	1:0.0846	13	40.1
Cpd 2 + Imidacloprid	50 + 15.1	1:0.302	26.3*	11.8
Cpd 2 + Imidacloprid	50 + 21.15	1:0.423	58.1*	33.5
Cpd 2 + Imidacloprid	250 + 15.1	1:0.0604	96.7*	75.4
Cpd 2 + Imidacloprid	250 + 21.15	1:0.0846	100.0*	81.5
Cpd 1 + Imidacloprid	50 + 15.1	1:0.302	34.3*	0
Cpd 1 + Imidacloprid	50 + 21.15	1:0.423	41.3*	24.7
Cpd 1 + Imidacloprid	250 + 15.1	1:0.0604	23.8*	3.1
Cpd 1 + Imidacloprid	250 + 21.15	1:0.0846	34.7	40.1
Cpd 3 + Imidacloprid	50 + 15.1	1:0.302	0	16.1
Cpd 3 + Imidacloprid	50 + 21.15	1:0.423	13.7	36.8
Cpd 3 + Imidacloprid	250 + 15.1	1:0.0604	0	26.7
Cpd 3 + Imidacloprid	250 + 21.15	1:0.0846	36.0*	27
Cpd 5 + Imidacloprid	50 + 15.1	1:0.302	25.2*	16.1
Cpd 5 + Imidacloprid	50 + 21.15	1:0.423	54.4	67.3
Cpd 5 + Imidacloprid	250 + 15.1	1:0.0604	54.4	55.6
Cpd 5 + Imidacloprid	250 + 21.15	1:0.0846	23.4	55.6

*Denotes enhanced efficacy based on Colby formula

Table 2d – Mixtures comprising Spirotetramat and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		4.9	
Cpd 4	250		8.3	
Cpd 2	50		7.5	
Cpd 2	250		28	
Cpd 1	50		0	
Cpd 1	250		1.3	
Cpd 3	50		1.2	
Cpd 3	250		1.1	
Cpd 5	50		8.9	
Cpd 5	250		11.7	
Spirotetramat	50		2.1	
Spirotetramat	250		34.4	
Cpd 4 + Spirotetramat	50 + 50	1:1	0	6.8
Cpd 4 + Spirotetramat	50 + 250	1:5	9.1	37.6
Cpd 4 + Spirotetramat	250 + 50	1:0.2	0	10.2
Cpd 4 + Spirotetramat	250 + 250	1:1	1.9	39.9
Cpd 2 + Spirotetramat	50 + 50	1:1	0	9.4
Cpd 2 + Spirotetramat	50 + 250	1:5	9.4	39.3
Cpd 2 + Spirotetramat	250 + 50	1:0.2	25.3	29.5
Cpd 2 + Spirotetramat	250 + 250	1:1	53.0*	52.8
Cpd 1 + Spirotetramat	50 + 50	1:1	2.4*	2.1
Cpd 1 + Spirotetramat	50 + 250	1:5	20.5	34.4
Cpd 1 + Spirotetramat	250 + 50	1:0.2	0	2.1
Cpd 1 + Spirotetramat	250 + 250	1:1	31.4	34.4
Cpd 3 + Spirotetramat	50 + 50	1:1	0	2.1
Cpd 3 + Spirotetramat	50 + 250	1:5	28.1	34.4
Cpd 3 + Spirotetramat	250 + 50	1:0.2	0	2.1
Cpd 3 + Spirotetramat	250 + 250	1:1	61.0*	34.4
Cpd 5 + Spirotetramat	50 + 50	1:1	32.8*	10.8
Cpd 5 + Spirotetramat	50 + 250	1:5	35.7	40.2
Cpd 5 + Spirotetramat	250 + 50	1:0.2	46.4*	13.5
Cpd 5 + Spirotetramat	250 + 250	1:1	28.8	42.1

*Denotes enhanced efficacy based on Colby formula

Table 2e – Mixtures comprising Chlorantraniliprole and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		29	
Cpd 4	250		38.3	
Cpd 2	50		17.8	
Cpd 2	250		90.2	
Cpd 1	50		0	
Cpd 1	250		2.2	
Cpd 3	50		1.3	
Cpd 3	250		1.2	
Cpd 5	50		56.3	
Cpd 5	250		11.2	
Chlorantraniliprole	0.54		0	
Chlorantraniliprole	189.1		59.2	
Cpd 4 + Chlorantraniliprole	50 + 0.54	1:0.0108	27.6	29
Cpd 4 + Chlorantraniliprole	50 + 189.1	1:3.782	97.1*	71.1
Cpd 4 + Chlorantraniliprole	250 + 0.54	1:0.0022	20.6	38.3
Cpd 4 + Chlorantraniliprole	250 + 189.1	1:0.7564	69.1	74.9
Cpd 2 + Chlorantraniliprole	50 + 0.54	1:0.0108	1.2	17.8
Cpd 2 + Chlorantraniliprole	50 + 189.1	1:3.782	59.7	66.5
Cpd 2 + Chlorantraniliprole	250 + 0.54	1:0.0022	70.5	90.2
Cpd 2 + Chlorantraniliprole	250 + 189.1	1:0.7564	78.1	96
Cpd 1 + Chlorantraniliprole	50 + 0.54	1:0.0108	0	0
Cpd 1 + Chlorantraniliprole	50 + 189.1	1:3.782	55.1	59.2
Cpd 1 + Chlorantraniliprole	250 + 0.54	1:0.0022	1.3	2.2
Cpd 1 + Chlorantraniliprole	250 + 189.1	1:0.7564	75.0*	60.2
Cpd 3 + Chlorantraniliprole	50 + 0.54	1:0.0108	3.7*	1.3
Cpd 3 + Chlorantraniliprole	50 + 189.1	1:3.782	39.7	59.8
Cpd 3 + Chlorantraniliprole	250 + 0.54	1:0.0022	12.0*	1.2
Cpd 3 + Chlorantraniliprole	250 + 189.1	1:0.7564	69.0*	59.7
Cpd 5 + Chlorantraniliprole	50 + 0.54	1:0.0108	7.9	56.3
Cpd 5 + Chlorantraniliprole	50 + 189.1	1:3.782	48.9	82.2
Cpd 5 + Chlorantraniliprole	250 + 0.54	1:0.0022	13.5*	11.2
Cpd 5 + Chlorantraniliprole	250 + 189.1	1:0.7564	59.7	63.8

*Denotes enhanced efficacy based on Colby formula

Table 2f – Mixtures comprising Bifenthrin and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		15	
Cpd 4	250		17.9	
Cpd 2	50		15.7	
Cpd 2	250		90.6	
Cpd 1	50		9.3	
Cpd 1	250		0	
Cpd 3	50		21.2	
Cpd 3	250		25.3	
Cpd 5	50		84.4	
Cpd 5	250		55.1	
Bifenthrin	50		0	
Bifenthrin	250		0	
Cpd 4 + Bifenthrin	50 + 50	1:1	17.9*	15
Cpd 4 + Bifenthrin	50 + 250	1:5	78.3*	15
Cpd 4 + Bifenthrin	250 + 50	1:0.2	14.3	17.9
Cpd 4 + Bifenthrin	250 + 250	1:1	60.3*	17.9
Cpd 2 + Bifenthrin	50 + 50	1:1	22.5*	15.7
Cpd 2 + Bifenthrin	50 + 250	1:5	75.0*	15.7
Cpd 2 + Bifenthrin	250 + 50	1:0.2	100*	90.6
Cpd 2 + Bifenthrin	250 + 250	1:1	100*	90.6
Cpd 1 + Bifenthrin	50 + 50	1:1	0	9.3
Cpd 1 + Bifenthrin	50 + 250	1:5	4.8*	0
Cpd 1 + Bifenthrin	250 + 50	1:0.2	3.6*	0
Cpd 1 + Bifenthrin	250 + 250	1:1	11.9*	0
Cpd 3 + Bifenthrin	50 + 50	1:1	0	0
Cpd 3 + Bifenthrin	50 + 250	1:5	24.2*	0
Cpd 3 + Bifenthrin	250 + 50	1:0.2	0	0
Cpd 3 + Bifenthrin	250 + 250	1:1	27.6*	0
Cpd 5 + Bifenthrin	50 + 50	1:1	26.7	84.4
Cpd 5 + Bifenthrin	50 + 250	1:5	95.2*	84.4
Cpd 5 + Bifenthrin	250 + 50	1:0.2	27.3	55.1
Cpd 5 + Bifenthrin	250 + 250	1:1	74.6*	55.1

*Denotes enhanced efficacy based on Colby formula

Table 2g – Mixtures comprising Indoxacarb and their activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		30.6	
Cpd 4	250		23.1	
Cpd 2	50		31.1	
Cpd 2	250		91.7	
Cpd 1	50		2.6	
Cpd 1	250		2.1	
Cpd 3	50		18	
Cpd 3	250		27.7	
Cpd 5	50		30	
Cpd 5	250		73.4	
Indoxacarb	50		1.2	
Indoxacarb	250		0	
Cpd 4 + Indoxacarb	50 + 50	1:1	38.2*	31.5
Cpd 4 + Indoxacarb	50 + 250	1:5	90.3*	30.6
Cpd 4 + Indoxacarb	250 + 50	1:0.2	11.1	24
Cpd 4 + Indoxacarb	250 + 250	1:1	35.5*	23.1
Cpd 2 + Indoxacarb	50 + 50	1:1	1.7	32
Cpd 2 + Indoxacarb	50 + 250	1:5	2.5	31.1
Cpd 2 + Indoxacarb	250 + 50	1:0.2	100.0*	91.8
Cpd 2 + Indoxacarb	250 + 250	1:1	97.4*	91.7
Cpd 1 + Indoxacarb	50 + 50	1:1	8.0*	3.8
Cpd 1 + Indoxacarb	50 + 250	1:5	1.5*	0
Cpd 1 + Indoxacarb	250 + 50	1:0.2	0	1.2
Cpd 1 + Indoxacarb	250 + 250	1:1	6.7*	0
Cpd 3 + Indoxacarb	50 + 50	1:1	20.4*	1.2
Cpd 3 + Indoxacarb	50 + 250	1:5	8.8*	0
Cpd 3 + Indoxacarb	250 + 50	1:0.2	1.6*	1.2
Cpd 3 + Indoxacarb	250 + 250	1:1	0	0
Cpd 5 + Indoxacarb	50 + 50	1:1	46.0*	30.9
Cpd 5 + Indoxacarb	50 + 250	1:5	83.8*	30
Cpd 5 + Indoxacarb	250 + 50	1:0.2	50	73.8
Cpd 5 + Indoxacarb	250 + 250	1:1	73.9*	73.4

*Denotes enhanced efficacy based on Colby formula

Table 2h – Mixtures with Spinetoram and activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		0.9	
Cpd 4	250		9.2	
Cpd 2	50		0	
Cpd 2	250		2.9	
Cpd 1	50		2.4	
Cpd 1	250		7.6	
Cpd 3	50		0	
Cpd 3	250		0	
Cpd 5	50		0	
Cpd 5	250		6.9	
Spinetoram	5		0	
Spinetoram	29.5		64.4	
Cpd 4 + Spinetoram	50 + 5	1:0.1	20.5	0.9
Cpd 4 + Spinetoram	50 + 29.5	1:0.59	88.9*	64.7
Cpd 4 + Spinetoram	250 + 5	1:0.02	44.6	67.7
Cpd 4 + Spinetoram	250 + 29.5	1:0.118	77.6*	67.7
Cpd 2 + Spinetoram	50 + 5	1:0.1	23.9	64.7
Cpd 2 + Spinetoram	50 + 29.5	1:0.59	85.5*	64.7
Cpd 2 + Spinetoram	250 + 5	1:0.02	21.1*	9.2
Cpd 2 + Spinetoram	250 + 29.5	1:0.118	92.2*	67.7
Cpd 1 + Spinetoram	50 + 5	1:0.1	20.5*	0.9
Cpd 1 + Spinetoram	50 + 29.5	1:0.59	83.8*	64.7
Cpd 1 + Spinetoram	250 + 5	1:0.02	16.7*	9.2
Cpd 1 + Spinetoram	250 + 29.5	1:0.118	98.0*	67.7
Cpd 3 + Spinetoram	50 + 5	1:0.1	8.5*	0.9
Cpd 3 + Spinetoram	50 + 29.5	1:0.59	62.0	64.7
Cpd 3 + Spinetoram	250 + 5	1:0.02	19.1*	9.2
Cpd 3 + Spinetoram	250 + 29.5	1:0.118	48.0	67.7
Cpd 5 + Spinetoram	50 + 5	1:0.1	20.0*	0.9
Cpd 5 + Spinetoram	50 + 29.5	1:0.59	100*	64.7
Cpd 5 + Spinetoram	250 + 5	1:0.02	22.6*	9.2
Cpd 5 + Spinetoram	250 + 29.5	1:0.118	70.7*	67.7

*Denotes enhanced efficacy based on Colby formula

Table 2i – Mixtures with Pyriproxyfen and activity on Silverleaf Whitefly

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		7	
Cpd 4	250		8	
Cpd 5	50		0	
Cpd 5	250		0	
Pyriproxyfen	0.053		68	
Cpd 4 + Pyriproxyfen	50 + 0.053	1:0.00106	46	70
Cpd 4 + Pyriproxyfen	250 + 0.053	1:0.000212	66	70
Cpd 5 + Pyriproxyfen	50 + 0.053	1:0.00106	64	68
Cpd 5 + Pyriproxyfen	250 + 0.053	1:0.000212	65	68

TEST B

5 For evaluating control of the Western Flower Thrips (*Frankliniella occidentalis* Pergande) through contact and/or systemic means, each test unit consisted of a small open container with a 5- to 7-day-old bean (var. Soleil) plant inside.

10 Test solutions were formulated and sprayed with 3 replications as described for Test A. After spraying, the test units were allowed to dry for 1 hour, 22 to 27 adult thrips were added to each unit and then a black, screened cap was placed on top. The test units were held for 7 days at 25 °C and 45-55% relative humidity. To evaluate the level of plant protection provided by each treatment, each test unit was then visually assessed for plant damage inflicted by insect feeding and rated 0-10, where 0 = undamaged and 10 = plant is dead; the results are listed in Tables 3a-g.

15

20

25

Table 3a – Mixtures comprising Cyantraniliprole and their activity on Western Flower
Thrips

Treatment	Rate [ppm]	Ratio	Plant Damage (obs)	Plant Damage (exp)
Cpd 4	11.1		43.3	
Cpd 4	36.8		86.7	
Cpd 2	1.5		13.3	
Cpd 2	4.5		56.7	
Cpd 1	3.18		33.3	
Cpd 1	9.75		73.3	
Cpd 3	2.93		13.3	
Cpd 3	16.1		83.3	
Cpd 5	0.4		20.0	
Cpd 5	2		40.0	
Cyantraniliprole	0.08		6.7	
Cyantraniliprole	0.28		43.3	
Cpd 4 + Cyantraniliprole	11.1 + 0.08	1:0.0072	46.7	47.1
Cpd 4 + Cyantraniliprole	11.1 + 0.28	1:0.0252	83.3*	67.9
Cpd 4 + Cyantraniliprole	36.8 + 0.08	1:0.0022	73.3	87.6
Cpd 4 + Cyantraniliprole	36.8 + 0.28	1:0.0076	86.7*	50.9
Cpd 2 + Cyantraniliprole	1.5 + 0.08	1:0.0533	56.7*	19.1
Cpd 2 + Cyantraniliprole	1.5 + 0.28	1:0.1867	66.7*	59.6
Cpd 2 + Cyantraniliprole	4.5 + 0.08	1:0.0178	76.7	76.9
Cpd 2 + Cyantraniliprole	4.5 + 0.28	1:0.0622	86.7*	75.4
Cpd 1 + Cyantraniliprole	3.18 + 0.08	1:0.0252	70.0*	37.8
Cpd 1 + Cyantraniliprole	3.18 + 0.28	1:0.0881	70.0*	62.2
Cpd 1 + Cyantraniliprole	9.75 + 0.08	1:0.0082	36.7	75.1
Cpd 1 + Cyantraniliprole	9.75 + 0.28	1:0.0287	90.0*	84.9
Cpd 3 + Cyantraniliprole	2.93 + 0.08	1:0.0273	36.7*	19.1
Cpd 3 + Cyantraniliprole	2.93 + 0.28	1:0.0956	63.3*	50.9
Cpd 3 + Cyantraniliprole	16.1 + 0.08	1:0.005	70.0	90.6
Cpd 3 + Cyantraniliprole	16.1 + 0.28	1:0.0174	96.7*	90.6
Cpd 5 + Cyantraniliprole	0.4 + 0.08	1:0.2	20.0	25.3
Cpd 5 + Cyantraniliprole	0.4 + 0.28	1:0.7	23.3	54.7
Cpd 5 + Cyantraniliprole	2 + 0.08	1:0.04	36.7	44.0
Cpd 5 + Cyantraniliprole	2 + 0.28	1:0.14	60.0	66.0

*Denotes enhanced plant protection based on Colby formula

Table 3b – Mixtures comprising Acetamiprid and their activity on Western Flower Thrips

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	11.1		40.0	
Cpd 4	36.8		86.7	
Cpd 2	1.5		16.7	
Cpd 2	4.5		83.3	
Cpd 1	3.18		73.3	
Cpd 1	9.75		76.7	
Cpd 3	2.93		40.0	
Cpd 3	16.1		76.7	
Cpd 5	0.4		23.3	
Cpd 5	2		36.7	
Acetamiprid	5.49		30.0	
Acetamiprid	7.04		70.0	
Cpd 4 + Acetamiprid	11.1 + 5.49	1:0.4946	90.0*	58.0
Cpd 4 + Acetamiprid	11.1 + 7.04	1:0.6342	93.3*	82.0
Cpd 4 + Acetamiprid	36.8 + 5.49	1:0.1492	93.3*	90.7
Cpd 4 + Acetamiprid	36.8 + 7.04	1:0.1913	96.7*	75.0
Cpd 2 + Acetamiprid	1.5 + 5.49	1:3.66	86.7*	41.7
Cpd 2 + Acetamiprid	1.5 + 7.04	1:4.6933	86.7	88.3
Cpd 2 + Acetamiprid	4.5 + 5.49	1:1.22	80.0	98.3
Cpd 2 + Acetamiprid	4.5 + 7.04	1:1.5644	90.0	95.0
Cpd 1 + Acetamiprid	3.18 + 5.49	1:1.7264	86.7*	81.3
Cpd 1 + Acetamiprid	3.18 + 7.04	1:2.2138	90.0	92.0
Cpd 1 + Acetamiprid	9.75 + 5.49	1:0.5631	93.3*	83.7
Cpd 1 + Acetamiprid	9.75 + 7.04	1:0.7221	100.0*	93.0
Cpd 3 + Acetamiprid	2.93 + 5.49	1:1.8737	80.0*	58.0
Cpd 3 + Acetamiprid	2.93 + 7.04	1:2.4027	90.0*	82.0
Cpd 3+ Acetamiprid	16.1 + 5.49	1:0.341	93.3*	93.0
Cpd 3 + Acetamiprid	16.1 + 7.04	1:0.4373	100.0*	93.0
Cpd 5 + Acetamiprid	0.4 + 5.49	1:13.725	30.0	46.3
Cpd 5 + Acetamiprid	0.4 + 7.04	1:17.6	63.3	77.0
Cpd 5 + Acetamiprid	2 + 5.49	1:2.745	63.3*	55.7
Cpd 5 + Acetamiprid	2 + 7.04	1:3.52	80.0	81.0

*Denotes enhanced plant protection based on Colby formula

Table 3c – Mixtures comprising Imidacloprid and their activity on Western Flower Thrips

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	11.1		36.7	
Cpd 4	36.8		70.0	
Cpd 2	1.5		43.3	
Cpd 2	4.5		80.0	
Cpd 1	3.18		63.3	
Cpd 1	9.75		83.3	
Cpd 3	2.93		33.3	
Cpd 3	16.1		80.0	
Cpd 5	0.4		16.7	
Cpd 5	2		40.0	
Imidacloprid	6.4		66.7	
Imidacloprid	10.9		86.7	
Cpd 4 + Imidacloprid	11.1 + 6.4	1:0.5766	90.0*	78.9
Cpd 4 + Imidacloprid	11.1 + 10.9	1:0.982	90.0	91.6
Cpd 4 + Imidacloprid	36.8 + 6.4	1:0.1739	83.3	90.0
Cpd 4 + Imidacloprid	36.8 + 10.9	1:0.2962	96.7*	92.4
Cpd 2 + Imidacloprid	1.5 + 6.4	1:4.2667	73.3	81.1
Cpd 2 + Imidacloprid	1.5 + 10.9	1:7.2667	90.0	93.3
Cpd 2 + Imidacloprid	4.5 + 6.4	1:1.4222	83.3	98.0
Cpd 2 + Imidacloprid	4.5 + 10.9	1:2.4222	93.3	97.3
Cpd 1 + Imidacloprid	3.18 + 6.4	1:2.0126	86.7	87.8
Cpd 1 + Imidacloprid	3.18 + 10.9	1:3.4277	93.3	95.1
Cpd 1 + Imidacloprid	9.75 + 6.4	1:0.6564	90.0	94.4
Cpd 1 + Imidacloprid	9.75 + 10.9	1:1.1179	90.0	97.8
Cpd 3 + Imidacloprid	2.93 + 6.4	1:2.1843	83.3*	77.8
Cpd 3 + Imidacloprid	2.93 + 10.9	1:3.7201	93.3*	91.1
Cpd 3 + Imidacloprid	16.1 + 6.4	1:0.3975	80.0	97.3
Cpd 3 + Imidacloprid	16.1 + 10.9	1:0.677	100.0*	97.3
Cpd 5 + Imidacloprid	0.4 + 6.4	1:16	80.0*	72.2
Cpd 5 + Imidacloprid	0.4 + 10.9	1:27.25	80.0	88.9
Cpd 5 + Imidacloprid	2 + 6.4	1:3.2	46.7	80.0
Cpd 5 + Imidacloprid	2 + 10.9	1:5.45	86.7	92.0

*Denotes enhanced plant protection based on Colby formula

Table 3d – Mixtures comprising Spirotetramat and their activity on Western Flower Thrips

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	11.1		60.0	
Cpd 4	36.8		93.3	
Cpd 2	1.5		36.7	
Cpd 2	4.5		76.7	
Cpd 1	3.18		63.3	
Cpd 1	9.75		90.0	
Cpd 4	2.93		50.0	
Cpd 4	16.1		86.7	
Cpd 5	0.4		43.3	
Cpd 5	2		63.3	
Spirotetramat	100		53.3	
Spirotetramat	250		73.3	
Cpd 4 + Spirotetramat	11.1 + 100	1:9.009	66.7	81.3
Cpd 4 + Spirotetramat	11.1 + 250	1:22.5225	83.3	89.3
Cpd 4 + Spirotetramat	36.8 + 100	1:2.7174	93.3	96.9
Cpd 4 + Spirotetramat	36.8 + 250	1:6.7935	86.7*	83.1
Cpd 2 + Spirotetramat	1.5 + 100	1:66.6667	60.0	70.4
Cpd 2 + Spirotetramat	1.5 + 250	1:166.6667	86.7	89.1
Cpd 2 + Spirotetramat	4.5 + 100	1:22.2222	90.0	92.2
Cpd 2 + Spirotetramat	4.5 + 250	1:55.5556	96.7*	93.8
Cpd 1 + Spirotetramat	3.18 + 100	1:31.4465	86.7*	82.9
Cpd 1 + Spirotetramat	3.18 + 250	1:78.6164	86.7	90.2
Cpd 1 + Spirotetramat	9.75 + 100	1:10.2564	86.7	95.3
Cpd 1 + Spirotetramat	9.75 + 250	1:25.641	96.7	97.3
Cpd 3 + Spirotetramat	2.93 + 100	1:34.1297	56.7	76.7
Cpd 3 + Spirotetramat	2.93 + 250	1:85.3242	86.7	86.7
Cpd 3 + Spirotetramat	16.1 + 100	1:6.2112	93.3	96.4
Cpd 3 + Spirotetramat	16.1 + 250	1:15.528	96.7*	96.4
Cpd 5 + Spirotetramat	0.4 + 100	1:250	33.3	73.6
Cpd 5 + Spirotetramat	0.4 + 250	1:625	80.0	84.9
Cpd 5 + Spirotetramat	2 + 100	1:50	33.3	82.9
Cpd 5 + Spirotetramat	2 + 250	1:125	73.3	90.2

*Denotes enhanced plant protection based on Colby formula

Table 3e – Mixtures comprising Chlorantraniliprole and their activity on Western Flower
Thrips

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	11.1		60.0	
Cpd 4	36.8		76.7	
Cpd 2	1.5		66.7	
Cpd 2	4.5		43.3	
Cpd1	3.18		76.7	
Cpd 1	9.75		66.7	
Cpd 3	2.93		83.3	
Cpd 3	16.1		80.0	
Cpd 5	0.4		70.0	
Cpd 5	2		26.7	
Chlorantraniliprole	2.08		63.3	
Chlorantraniliprole	28.7		60.0	
Cpd 4 + Chlorantraniliprole	11.1 + 2.08	1:0.1874	76.7	85.3
Cpd 4 + Chlorantraniliprole	11.1 + 28.7	1:2.5856	90.0*	84.0
Cpd 4 + Chlorantraniliprole	36.8 + 2.08	1:0.0565	83.3	91.4
Cpd 4 + Chlorantraniliprole	36.8 + 28.7	1:0.7799	93.3*	86.7
Cpd 2 + Chlorantraniliprole	1.5 + 2.08	1:1.3867	63.3	87.8
Cpd 2 + Chlorantraniliprole	1.5 + 28.7	1:19.1333	90.0*	79.2
Cpd 2 + Chlorantraniliprole	4.5 + 2.08	1:0.4622	80.0	86.8
Cpd 2 + Chlorantraniliprole	4.5 + 28.7	1:6.3778	93.3*	77.3
Cpd 1 + Chlorantraniliprole	3.18 + 2.08	1:0.6541	66.7	91.4
Cpd 1 + Chlorantraniliprole	3.18 + 28.7	1:9.0252	90.0	90.7
Cpd 1 + Chlorantraniliprole	9.75 + 2.08	1:0.2133	36.7	87.8
Cpd 1 + Chlorantraniliprole	9.75 + 28.7	1:2.9436	100.0*	86.7
Cpd 3 + Chlorantraniliprole	2.93 + 2.08	1:0.7099	53.3	93.9
Cpd 3 + Chlorantraniliprole	2.93 + 28.7	1:9.7952	86.7	93.3
Cpd 3 + Chlorantraniliprole	16.1 + 2.08	1:0.1292	86.7	92.0
Cpd 3 + Chlorantraniliprole	16.1 + 28.7	1:1.7826	90.0	92.0
Cpd 5 + Chlorantraniliprole	0.4 + 2.08	1:5.2	33.3	89.0
Cpd 5 + Chlorantraniliprole	0.4 + 28.7	1:71.75	76.7	88.0
Cpd 5 + Chlorantraniliprole	2 + 2.08	1:1.04	76.7*	73.1
Cpd 5 + Chlorantraniliprole	2 + 28.7	1:14.35	93.3*	70.7

*Denotes enhanced plant protection based on Colby formula

Table 3f – Mixtures comprising Bifenthrin and their activity on Western Flower

Thrips

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	11.1		73.3	
Cpd 4	36.8		90.0	
Cpd 2	1.5		43.3	
Cpd 2	4.5		73.3	
Cpd 1	3.18		33.3	
Cpd 1	9.75		80.0	
Cpd 3	2.93		40.0	
Cpd 3	16.1		86.7	
Cpd 5	0.4		33.3	
Cpd 5	2		60.0	
Bifenthrin	50		36.7	
Bifenthrin	250		70.0	
Cpd 4 + Bifenthrin	11.1 + 50	1:4.5045	86.7	83.1
Cpd 4 + Bifenthrin	11.1 + 250	1:22.5225	96.7*	92.0
Cpd 4 + Bifenthrin	36.8 + 50	1:1.3587	90.0	93.7
Cpd 4 + Bifenthrin	36.8 + 250	1:6.7935	100.0*	83.0
Cpd 2 + Bifenthrin	1.5 + 50	1:33.3333	56.7	64.1
Cpd 2 + Bifenthrin	1.5 + 250	1:166.6667	90.0*	83.1
Cpd 2 + Bifenthrin	4.5 + 50	1:11.1111	66.7	96.4
Cpd 2 + Bifenthrin	4.5 + 250	1:55.5556	93.38	92.0
Cpd 1 + Bifenthrin	3.18 + 50	1:15.7233	83.3*	57.8
Cpd 1 + Bifenthrin	3.18 + 250	1:78.6164	90.0*	80.0
Cpd 1 + Bifenthrin	9.75 + 50	1:5.1282	86.7	87.3
Cpd 1 + Bifenthrin	9.75 + 250	1:25.641	100.0*	94.0
Cpd 3 + Bifenthrin	2.93 + 50	1:17.0648	73.3*	62.0
Cpd 3 + Bifenthrin	2.93 + 250	1:85.3242	83.3*	82.0
Cpd 3 + Bifenthrin	16.1 + 50	1:3.1056	73.3	96.0
Cpd 3 + Bifenthrin	16.1 + 250	1:15.528	93.3	96.0
Cpd 5 + Bifenthrin	0.4 + 50	1:125	33.3	57.8
Cpd 5 + Bifenthrin	0.4 + 250	1:625	73.3	80.0
Cpd 5 + Bifenthrin	2 + 50	1:25	66.7	74.7
Cpd 5 + Bifenthrin	2 + 250	1:125	90.0*	88.0

*Denotes enhanced plant protection based on Colby formula

Table 3g – Mixtures comprising Indoxacarb and their activity on Western Flower
Thrips

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	11.1		20.0	
Cpd 4	36.8		73.3	
Cpd 2	1.5		16.7	
Cpd 2	4.5		33.3	
Cpd 1	3.18		10.0	
Cpd 1	9.75		76.7	
Cpd 3	2.93		43.3	
Cpd 3	16.1		70.0	
Cpd 5	0.4		13.3	
Cpd 5	2		26.7	
Indoxacarb	50		10.0	
Indoxacarb	250		36.7	
Cpd 4 + Indoxacarb	11.1 + 50	1:4.5045	33.3*	28.0
Cpd 4 + Indoxacarb	11.1 + 250	1:22.5225	90.0*	49.3
Cpd 4 + Indoxacarb	36.8 + 50	1:1.3587	86.7*	76.0
Cpd 4 + Indoxacarb	36.8 + 250	1:6.7935	100.0*	47.2
Cpd 2 + Indoxacarb	1.5 + 50	1:33.3333	33.3*	25.0
Cpd 2 + Indoxacarb	1.5 + 250	1:166.6667	80.0*	40.0
Cpd 2 + Indoxacarb	4.5 + 50	1:11.1111	33.3	55.6
Cpd 2 + Indoxacarb	4.5 + 250	1:55.5556	80.0*	57.8
Cpd 1 + Indoxacarb	3.18 + 50	1:15.7233	86.7*	19.0
Cpd 1 + Indoxacarb	3.18 + 250	1:78.6164	90.0*	43.0
Cpd 1 + Indoxacarb	9.75 + 50	1:5.1282	86.7*	79.0
Cpd 1 + Indoxacarb	9.75 + 250	1:25.641	100.0*	85.2
Cpd 3 + Indoxacarb	2.93 + 50	1:17.0648	76.7*	49.0
Cpd 3 + Indoxacarb	2.93 + 250	1:85.3242	96.7*	64.1
Cpd 3 + Indoxacarb	16.1 + 50	1:3.1056	76.7	81.0
Cpd 3 + Indoxacarb	16.1 + 250	1:15.528	93.3*	81.0
Cpd 5 + Indoxacarb	0.4 + 50	1:125	40.0*	22.0
Cpd 5 + Indoxacarb	0.4 + 250	1:625	70.0*	45.1
Cpd 5 + Indoxacarb	2 + 50	1:25	70.0*	34.0
Cpd 5 + Indoxacarb	2 + 250	1:125	86.7*	53.6

5 *Denotes enhanced plant protection based on Colby formula

TEST C

For evaluating control of Potato Leafhopper (*Empoasca fabae* Harris) through contact and/or systemic means, each test unit consisted of a small open container with a 5- to 6-day-old Longio bean plant (primary leaves emerged) inside. White sand was added to the top of the soil, and one of the primary leaves was excised prior to application. Test compounds were formulated and sprayed with 3 replications as described for Test A. After spraying, the test units were allowed to dry for 1 hour before they were infested with 5 potato leafhoppers (18- to 21-day-old adults). A black, screened cap was placed on the top of each container. The test units were held for 6 days in a growth chamber at 19–21 °C and 50–70% relative humidity. Each test unit was then visually assessed for insect mortality; the results are listed in Tables 4a-c.

For evaluating control of corn planthopper (*Peregrinus maidis* (Ashmead)) through contact and/or systemic means, the test unit consisted of a small open container with a 3–4-day-old corn (maize) plant inside. White sand was added to the top of the soil prior to application of the test compound.

Test compounds were formulated and sprayed at 250 and/or 50 ppm and/or 10ppm. After spraying of the formulated test compound, the test units were allowed to dry for 1 h before they were post-infested with ~15–20 nymphs (18-to-21-day-old). A black, screened cap was placed on the top of each test unit, and the test units were held for 6 days in a growth chamber at 19-21 °C and 50–70% relative humidity. Each test unit was then visually assessed for insect mortality; the results are listed in Tables 4d-g.

25

30

35

40

Table 4a – Mixtures comprising Cyantraniliprole and their activity on Potato Leafhopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		7.1	
Cpd 4	250		6.7	
Cpd 2	50		46.7	
Cpd 2	250		86.7	
Cpd 1	50		60	
Cpd 1	250		35.7	
Cpd 3	50		0	
Cpd 3	250		0	
Cpd 5	50		0	
Cpd 5	250		26.7	
Cyantraniliprole	1.75		13.3	
Cyantraniliprole	3.3		28.6	
Cpd 4 + Cyantraniliprole	50 + 1.75	1:0.035	33.3*	19.5
Cpd 4 + Cyantraniliprole	50 + 3.3	1:0.066	53.3*	33.7
Cpd 4 + Cyantraniliprole	250 + 1.75	1:0.007	13.3	19.1
Cpd 4 + Cyantraniliprole	250 + 3.3	1:0.0132	46.7*	33.3
Cpd 2 + Cyantraniliprole	50 + 1.75	1:0.035	26.7	53.8
Cpd 2 + Cyantraniliprole	50 + 3.3	1:0.066	35.7	61.9
Cpd 2 + Cyantraniliprole	250 + 1.75	1:0.007	80	88.4
Cpd 2 + Cyantraniliprole	250 + 3.3	1:0.0132	80	90.5
Cpd 1 + Cyantraniliprole	50 + 1.75	1:0.035	53.3	72.9
Cpd 1 + Cyantraniliprole	50 + 3.3	1:0.066	60	71.4
Cpd 1 + Cyantraniliprole	250 + 1.75	1:0.007	53.3*	44.3
Cpd 1 + Cyantraniliprole	250 + 3.3	1:0.0132	73.3*	54.1
Cpd 3 + Cyantraniliprole	50 + 1.75	1:0.035	20.0*	13.3
Cpd 3 + Cyantraniliprole	50 + 3.3	1:0.066	33.3*	28.6
Cpd 3 + Cyantraniliprole	250 + 1.75	1:0.007	14.3*	13.3
Cpd 3 + Cyantraniliprole	250 + 3.3	1:0.0132	60.0*	28.6
Cpd 5 + Cyantraniliprole	50 + 1.75	1:0.035	14.3*	13.3
Cpd 5 + Cyantraniliprole	50 + 3.3	1:0.066	40.0*	28.6
Cpd 5 + Cyantraniliprole	250 + 1.75	1:0.007	33.3	36.4
Cpd 5 + Cyantraniliprole	250 + 3.3	1:0.0132	73.3*	47.6

*Denotes enhanced efficacy based on Colby formula

Table 4b – Mixtures comprising Imidacloprid and their activity on Potato Leafhopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		0	
Cpd 4	250		14.3	
Cpd 2	50		6.7	
Cpd 2	250		85.7	
Cpd 1	50		20	
Cpd 1	250		6.7	
Cpd 3	50		13.3	
Cpd 3	250		0	
Cpd 5	50		0	
Cpd 5	250		6.7	
Imidacloprid	0.82		53.3	
Imidacloprid	2.25		40	
Cpd 4 + Imidacloprid	50 + 0.82	1:0.0164	33.3	53.3
Cpd 4 + Imidacloprid	50 + 2.25	1:0.045	21.4	40
Cpd 4 + Imidacloprid	250 + 0.82	1:0.00328	33.3	60
Cpd 4 + Imidacloprid	250 + 2.25	1:0.009	60.0*	48.6
Cpd 2 + Imidacloprid	50 + 0.82	1:0.0164	33.3	56.4
Cpd 2 + Imidacloprid	50 + 2.25	1:0.045	66.7*	44
Cpd 2 + Imidacloprid	250 + 0.82	1:0.00328	86.7	93.3
Cpd 2 + Imidacloprid	250 + 2.25	1:0.009	100.0*	66.7
Cpd 1 + Imidacloprid	50 + 0.82	1:0.0164	40	62.7
Cpd 1 + Imidacloprid	50 + 2.25	1:0.045	73.3*	52
Cpd 1 + Imidacloprid	250 + 0.82	1:0.00328	66.7*	56.4
Cpd 1 + Imidacloprid	250 + 2.25	1:0.009	73.3*	44
Cpd 3 + Imidacloprid	50 + 0.82	1:0.0164	42.9	59.6
Cpd 3 + Imidacloprid	50 + 2.25	1:0.045	33.3	48
Cpd 3 + Imidacloprid	250 + 0.82	1:0.00328	20	40
Cpd 3 + Imidacloprid	250 + 2.25	1:0.009	85.7*	53.3
Cpd 5 + Imidacloprid	50 + 0.82	1:0.0164	33.3	40
Cpd 5 + Imidacloprid	50 + 2.25	1:0.045	73.3*	40
Cpd 5 + Imidacloprid	250 + 0.82	1:0.00328	50	56.4
Cpd 5 + Imidacloprid	250 + 2.25	1:0.009	35.7	44

*Denotes enhanced efficacy based on Colby formula

Table 4c – Mixtures comprising Imidacloprid and their activity on Potato Leafhopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		26.7	
Cpd 4	250		0	
Cpd 2	50		20	
Cpd 2	250		73.3	
Cpd 1	50		6.7	
Cpd 1	250		33.3	
Cpd 3	50		6.7	
Cpd 3	250		0	
Cpd 5	50		6.7	
Cpd 5	250		26.7	
Spirotetramat	50		28.6	
Spirotetramat	250		73.3	
Cpd 4 + Spirotetramat	50 + 50		53.3*	47.6
Cpd 4 + Spirotetramat	50 + 250		50	80.4
Cpd 4 + Spirotetramat	250 + 50		33.3*	28.6
Cpd 4 + Spirotetramat	250 + 250		66.7	73.3
Cpd 2 + Spirotetramat	50 + 50		46.7*	42.9
Cpd 2 + Spirotetramat	50 + 250		86.7*	78.7
Cpd 2 + Spirotetramat	250 + 50		66.7	81
Cpd 2 + Spirotetramat	250 + 250		93.3*	92.9
Cpd 1 + Spirotetramat	50 + 50		40.0*	35.5
Cpd 1 + Spirotetramat	50 + 250		73.3	75.1
Cpd 1 + Spirotetramat	250 + 50		26.7	52.4
Cpd 1 + Spirotetramat	250 + 250		73.3	82.2
Cpd 3 + Spirotetramat	50 + 50		33.3	33.3
Cpd 3 + Spirotetramat	50 + 250		50	75.1
Cpd 3 + Spirotetramat	250 + 50		46.7*	28.6
Cpd 3 + Spirotetramat	250 + 250		46.7	73.3
Cpd 5 + Spirotetramat	50 + 50		20	33.3
Cpd 5 + Spirotetramat	50 + 250		80.0*	75.1
Cpd 5 + Spirotetramat	250 + 50		53.3*	47.6
Cpd 5 + Spirotetramat	250 + 250		80	81

*Denotes enhanced efficacy based on Colby formula

Table 4d – Mixtures with Imidacloprid and activity on Corn Planthopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		0.0	
Cpd 4	250		9.1	
Cpd 2	50		0.0	
Cpd 2	250		54.1	
Cpd 1	50		14.8	
Cpd 1	250		45.8	
Cpd 3	50		8.3	
Cpd 3	250		0.0	
Cpd 5	50		5.3	
Cpd 5	250		75.0	
Imidacloprid	0.09		29.2	
Imidacloprid	0.16		20.0	
Cpd 4 + Imidacloprid	50 + 0.09	1:0.0018	0.0	29.2
Cpd 4 + Imidacloprid	50 + 0.16	1:0.0032	4.2	20.0
Cpd 4 + Imidacloprid	250 + 0.09	1:0.00036	12.0	27.3
Cpd 4 + Imidacloprid	250 + 0.16	1:0.00064	12.5	27.3
Cpd 2 + Imidacloprid	50 + 0.09	1:0.0018	81.8*	20.0
Cpd 2 + Imidacloprid	50 + 0.16	1:0.0032	36.4*	20.0
Cpd 2 + Imidacloprid	250 + 0.09	1:0.00036	48.8*	35.6
Cpd 2 + Imidacloprid	250 + 0.16	1:0.00064	23.1	27.3
Cpd 1 + Imidacloprid	50 + 0.09	1:0.0018	22.2	29.2
Cpd 1 + Imidacloprid	50 + 0.16	1:0.0032	100*	20.0
Cpd 1 + Imidacloprid	250 + 0.09	1:0.00036	100*	35.6
Cpd 1 + Imidacloprid	250 + 0.16	1:0.00064	88.9*	27.3
Cpd 3 + Imidacloprid	50 + 0.09	1:0.0018	66.7*	29.2
Cpd 3 + Imidacloprid	50 + 0.16	1:0.0032	22.6*	20.0
Cpd 3 + Imidacloprid	250 + 0.09	1:0.00036	25.0	35.6
Cpd 3 + Imidacloprid	250 + 0.16	1:0.00064	43.5*	27.3
Cpd 5 + Imidacloprid	50 + 0.09	1:0.0018	63.2*	29.2
Cpd 5 + Imidacloprid	50 + 0.16	1:0.0032	50.0*	20.0
Cpd 5 + Imidacloprid	250 + 0.09	1:0.00036	38.1*	35.6
Cpd 5 + Imidacloprid	250 + 0.16	1:0.00064	0	27.3

*Denotes enhanced efficacy based on Colby formula

Table 4e – Mixtures with Spinetoram and activity on Corn Planthopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		38.9	
Cpd 4	250		34.8	
Cpd 2	50		7.3	
Cpd 2	250		22.4	
Cpd 1	50		18.9	
Cpd 1	250		7.6	
Cpd 3	50		5.0	
Cpd 3	250		1.2	
Cpd 5	50		21.6	
Cpd 5	250		13.1	
Spinetoram	8.46		7.5	
Spinetoram	20.79		14.1	
Cpd 4 + Spinetoram	50 + 8.46	1:0.1692	13.2	43.4
Cpd 4 + Spinetoram	50 + 20.79	1:0.4158	21.0	47.5
Cpd 4 + Spinetoram	250 + 8.46	1:0.03384	16.7	43.4
Cpd 4 + Spinetoram	250 + 20.79	1:0.08316	34.9	44.0
Cpd 2 + Spinetoram	50 + 8.46	1:0.1692	21.4	47.5
Cpd 2 + Spinetoram	50 + 20.79	1:0.4158	30.1	47.5
Cpd 2 + Spinetoram	250 + 8.46	1:0.03384	4.1	39.7
Cpd 2 + Spinetoram	250 + 20.79	1:0.08316	27.9	44.0
Cpd 1 + Spinetoram	50 + 8.46	1:0.1692	16.4	43.4
Cpd 1 + Spinetoram	50 + 20.79	1:0.4158	38.7	47.5
Cpd 1 + Spinetoram	250 + 8.46	1:0.03384	4.9	39.7
Cpd 1 + Spinetoram	250 + 20.79	1:0.08316	26.9	44.0
Cpd 3 + Spinetoram	50 + 8.46	1:0.1692	1.4	43.4
Cpd 3 + Spinetoram	50 + 20.79	1:0.4158	14.1	47.5
Cpd 3 + Spinetoram	250 + 8.46	1:0.03384	10.4	39.7
Cpd 3 + Spinetoram	250 + 20.79	1:0.08316	27.3	44.0
Cpd 5 + Spinetoram	50 + 8.46	1:0.1692	6.0	43.4
Cpd 5 + Spinetoram	50 + 20.79	1:0.4158	13.2	47.5
Cpd 5 + Spinetoram	250 + 8.46	1:0.03384	4.2	39.7
Cpd 5 + Spinetoram	250 + 20.79	1:0.08316	22.6	44.0

Table 4f – Mixtures with Buprofezin and activity on Corn Planthopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		1.8	
Cpd 4	250		10.8	
Cpd 5	50		15.3	
Cpd 5	250		22.4	
Buprofezin	0.524		83.7	
Buprofezin	0.984		94.6	
Cpd 4 + Buprofezin	50 + 0.524	1:0.01048	78.9	84.0
Cpd 4 + Buprofezin	50 + 0.984	1:0.01968	98.4*	94.7
Cpd 4 + Buprofezin	250 + 0.524	1:0.002096	75.9	85.5
Cpd 4 + Buprofezin	250 + 0.984	1:0.003936	92.9	95.2
Cpd 5 + Buprofezin	50 + 0.524	1:0.01048	90.0*	86.2
Cpd 5 + Buprofezin	50 + 0.984	1:0.01968	81.8	95.5
Cpd 5 + Buprofezin	250 + 0.524	1:0.002096	70.3	87.4
Cpd 5 + Buprofezin	250 + 0.984	1:0.003936	89.1	95.8

*Denotes enhanced efficacy based on Colby formula

5

Table 4g – Mixtures with Fipronil and activity on Corn Planthopper

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		1.8	
Cpd 4	250		10.8	
Cpd 5	50		15.3	
Cpd 5	250		22.4	
Fipronil	0.78		65.2	
Fipronil	1.197		85.5	
Cpd 4 + Fipronil	50 + 0.78	1:0.0156	61.7	65.8
Cpd 4 + Fipronil	50 + 1.197	1:0.02394	85.7	85.7
Cpd 4 + Fipronil	250 + 0.78	1:0.00312	40.8	68.9
Cpd 4 + Fipronil	250 + 1.197	1:0.004788	87.0	87.1
Cpd 5 + Fipronil	50 + 0.78	1:0.0156	57.5	70.5
Cpd 5 + Fipronil	50 + 1.197	1:0.02394	82.2	87.7
Cpd 5 + Fipronil	250 + 0.78	1:0.00312	41.1	73.0
Cpd 5 + Fipronil	250 + 1.197	1:0.004788	90.9*	88.7

*Denotes enhanced efficacy based on Colby formula

TEST D

For evaluating control of cotton melon aphid (*Aphis gossypii* Glover) through contact and/or systemic means, each test unit consisted of a small open container with a 6- to 7-day-old okra plant inside. This was pre-infested by placing on a leaf of the test plant 30 to 40 aphids on a piece of leaf excised from a culture plant (cut-leaf method). The larvae moved onto the test plant as the leaf piece desiccated. After pre-infestation, the soil of the test unit was covered with a layer of sand.

Test compounds were formulated and sprayed as described for Test A. The applications were replicated three times. After spraying of the formulated test compounds, each test unit was allowed to dry for 1 hour and then a black, screened cap was placed on top. The test units were held for 6 days in a growth chamber at 19–21 °C and 50–70% relative humidity. Each test unit was then visually assessed for insect mortality; the results are listed in Tables 5a-n.

Table 5a – Mixtures comprising Cyantraniliprole and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		48.3	
Cpd 4	0.5		58.9	
Cpd 2	0.05		40.1	
Cpd 2	0.18		31.4	
Cpd 1	0.08		23.8	
Cpd 1	0.12		49.8	
Cpd 3	0.02		20.0	
Cpd 3	0.08		43.2	
Cpd 5	0.12		34.0	
Cpd 5	0.18		62.7	
Cyantraniliprole	0.06		21.1	
Cyantraniliprole	0.12		45.1	
Cpd 4 + Cyantraniliprole	0.2 + 0.06	1:0.3	84.0*	59.0
Cpd 4 + Cyantraniliprole	0.2 + 0.12	1:0.6	82.0*	72.0
Cpd 4 + Cyantraniliprole	0.5 + 0.06	1:0.12	85.0*	68.0
Cpd 4 + Cyantraniliprole	0.5 + 0.12	1:0.24	93.0*	77.0
Cpd 2 + Cyantraniliprole	0.05 + 0.06	1:1.2	36.0	53.0
Cpd 2 + Cyantraniliprole	0.05 + 0.12	1:2.4	42.0	67.0
Cpd 2 + Cyantraniliprole	0.18 + 0.06	1:0.3333	56.0*	46.0
Cpd 2 + Cyantraniliprole	0.18 + 0.12	1:0.6667	68.0*	62.0
Cpd 1 + Cyantraniliprole	0.08 + 0.06	1:0.75	41.0*	40.0
Cpd 1 + Cyantraniliprole	0.08 + 0.12	1:1.5	64.0*	58.0
Cpd 1 + Cyantraniliprole	0.12 + 0.06	1:0.5	72.0*	60.0
Cpd 1 + Cyantraniliprole	0.12 + 0.12	1:1	61.0	72.0
Cpd 3 + Cyantraniliprole	0.02 + 0.06	1:3	19.0	37.0
Cpd 3 + Cyantraniliprole	0.02 + 0.12	1:6	39.0	56.0
Cpd 3 + Cyantraniliprole	0.08 + 0.06	1:0.75	36.0	55.0
Cpd 3 + Cyantraniliprole	0.08 + 0.12	1:1.5	56.0	69.0
Cpd 5 + Cyantraniliprole	0.12 + 0.06	1:0.5	63.0*	48.0
Cpd 5 + Cyantraniliprole	0.12 + 0.12	1:1	73.0*	64.0
Cpd 5 + Cyantraniliprole	0.18 + 0.06	1:0.3333	72.0*	71.0
Cpd 5 + Cyantraniliprole	0.18 + 0.12	1:0.6667	81.0*	80.0

*Denotes enhanced efficacy based on Colby formula

Table 5b – Mixtures comprising Acetamiprid and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		48.3	
Cpd 4	0.5		58.9	
Cpd 2	0.05		40.1	
Cpd 2	0.18		31.4	
Cpd 1	0.08		23.8	
Cpd 1	0.12		49.8	
Cpd 3	0.02		20	
Cpd 3	0.08		43.2	
Cpd 5	0.12		34	
Cpd 5	0.18		62.7	
Acetamiprid	0.03		21.1	
Acetamiprid	0.06		45.1	
Cpd 4 + Acetamiprid	0.2 + 0.03	1:0.15	59.1	59.2
Cpd 4 + Acetamiprid	0.2 + 0.06	1:0.3	71.9*	71.6
Cpd 4 + Acetamiprid	0.5 + 0.03	1:0.06	75.9*	67.6
Cpd 4 + Acetamiprid	0.5 + 0.06	1:0.12	93.3*	77.4
Cpd 2 + Acetamiprid	0.05 + 0.03	1:0.6	31.7	52.8
Cpd 2 + Acetamiprid	0.05 + 0.06	1:1.2	44.7	67.1
Cpd 2 + Acetamiprid	0.18 + 0.03	1:0.1667	58.3*	45.9
Cpd 2 + Acetamiprid	0.18 + 0.06	1:0.3333	76.4*	62.3
Cpd 1 + Acetamiprid	0.08 + 0.03	1:0.375	34	39.9
Cpd 1 + Acetamiprid	0.08 + 0.06	1:0.75	67.2*	58.2
Cpd 1 + Acetamiprid	0.12 + 0.03	1:0.25	63.5*	60.4
Cpd 1 + Acetamiprid	0.12 + 0.06	1:0.5	68.8	72.4
Cpd 3 + Acetamiprid	0.02 + 0.03	1:1.5	16.3	36.9
Cpd 3 + Acetamiprid	0.02 + 0.06	1:3	39.5	56.1
Cpd 3 + Acetamiprid	0.08 + 0.03	1:0.375	18.5	55.2
Cpd 3 + Acetamiprid	0.08 + 0.06	1:0.75	41.2	68.8
Cpd 5 + Acetamiprid	0.12 + 0.03	1:0.25	28.9	47.9
Cpd 5 + Acetamiprid	0.12 + 0.06	1:0.5	56.3	63.7
Cpd 5 + Acetamiprid	0.18 + 0.03	1:0.1667	55.8	70.6
Cpd 5 + Acetamiprid	0.18 + 0.06	1:0.3333	79.8*	79.5

*Denotes enhanced efficacy based on Colby formula

Table 5c – Mixtures comprising Imidacloprid and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		6.7	
Cpd 4	0.5		6.7	
Cpd 2	0.05		0	
Cpd 2	0.18		80	
Cpd 1	0.08		100	
Cpd 1	0.12		26.7	
Cpd 3	0.02		0	
Cpd 3	0.08		20	
Cpd 5	0.12		73.3	
Cpd 5	0.18		6.7	
Imidacloprid	0.05		33.3	
Imidacloprid	0.1		6.7	
Cpd 4 + Imidacloprid	0.2 + 0.05	1:0.25	53.7*	52.6
Cpd 4 + Imidacloprid	0.2 + 0.1	1:0.5	67.0*	66.9
Cpd 4 + Imidacloprid	0.5 + 0.05	1:0.1	78.4*	73.4
Cpd 4 + Imidacloprid	0.5 + 0.1	1:0.2	86.0*	81.4
Cpd 2 + Imidacloprid	0.05 + 0.05	1:1	41.9	52.6
Cpd 2 + Imidacloprid	0.05 + 0.1	1:2	54	66.9
Cpd 2 + Imidacloprid	0.18 + 0.05	1:0.2778	61.6	78.7
Cpd 2 + Imidacloprid	0.18 + 0.1	1:0.5556	75	85.1
Cpd 1 + Imidacloprid	0.08 + 0.05	1:0.625	51.7	52.2
Cpd 1 + Imidacloprid	0.08 + 0.1	1:1.25	59.6	66.6
Cpd 1 + Imidacloprid	0.12 + 0.05	1:0.4167	44.7	61.7
Cpd 1 + Imidacloprid	0.12 + 0.1	1:0.8333	61.3	73.2
Cpd 3 + Imidacloprid	0.02 + 0.05	1:2.5	45.2*	41.4
Cpd 3 + Imidacloprid	0.02 + 0.1	1:5	47.6	59.1
Cpd 3 + Imidacloprid	0.08 + 0.05	1:0.625	35.8	58.1
Cpd 3 + Imidacloprid	0.08 + 0.1	1:1.25	56.4	70.7
Cpd 5 + Imidacloprid	0.12 + 0.05	1:0.4167	44.1	48.3
Cpd 5 + Imidacloprid	0.12 + 0.1	1:0.8333	56	63.8
Cpd 5 + Imidacloprid	0.18 + 0.05	1:0.2778	49.6	74.4
Cpd 5 + Imidacloprid	0.18 + 0.1	1:0.5556	61	82.1

*Denotes enhanced efficacy based on Colby formula

Table 5d – Mixtures comprising Spirotetramat and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		48.3	
Cpd 4	0.5		58.9	
Cpd 2	0.05		40.1	
Cpd 2	0.18		31.4	
Cpd 1	0.08		23.8	
Cpd 1	0.12		49.8	
Cpd 3	0.02		20	
Cpd 3	0.08		43.2	
Cpd 5	0.12		34	
Cpd 5	0.18		62.7	
Spirotetramat	0.25		21.1	
Spirotetramat	0.85		45.1	
Cpd 4 + Spirotetramat	0.2 + 0.25	1:1.25	59.1	59.2
Cpd 4 + Spirotetramat	0.2 + 0.85	1:4.25	71.9*	71.6
Cpd 4 + Spirotetramat	0.5 + 0.25	1:0.5	75.9*	67.6
Cpd 4 + Spirotetramat	0.5 + 0.85	1:1.7	93.3*	77.4
Cpd 2 + Spirotetramat	0.05 + 0.25	1:5	31.7	52.8
Cpd 2 + Spirotetramat	0.05 + 0.85	1:17	44.7	67.1
Cpd 2 + Spirotetramat	0.18 + 0.25	1:1.3889	58.3*	45.9
Cpd 2 + Spirotetramat	0.18 + 0.85	1:4.7222	76.4*	62.3
Cpd 1 + Spirotetramat	0.08 + 0.25	1:3.125	34	39.9
Cpd 1 + Spirotetramat	0.08 + 0.85	1:10.625	67.2*	58.2
Cpd 1 + Spirotetramat	0.12 + 0.25	1:2.0833	63.5*	60.4
Cpd 1 + Spirotetramat	0.12 + 0.85	1:7.0833	68.8	72.4
Cpd 3 + Spirotetramat	0.02 + 0.25	1:12.5	16.3	36.9
Cpd 3 + Spirotetramat	0.02 + 0.85	1:42.5	39.5	56.1
Cpd 3 + Spirotetramat	0.08 + 0.25	1:3.125	18.5	55.2
Cpd 3 + Spirotetramat	0.08 + 0.85	1:10.625	41.2	68.8
Cpd 5 + Spirotetramat	0.12 + 0.25	1:2.0833	28.9	47.9
Cpd 5 + Spirotetramat	0.12 + 0.85	1:7.0833	56.3	63.7
Cpd 5 + Spirotetramat	0.18 + 0.25	1:1.3889	55.8	70.6
Cpd 5 + Spirotetramat	0.18 + 0.85	1:4.7222	79.8*	79.5

*Denotes enhanced efficacy based on Colby formula

Table 5e – Mixtures comprising Chlorantraniliprole and their activity on Cotton Melon
Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		30.8	
Cpd 4	0.5		69.7	
Cpd 2	0.05		16.3	
Cpd 2	0.18		37	
Cpd 1	0.08		21.3	
Cpd 1	0.12		35.3	
Cpd 3	0.02		19.9	
Cpd 3	0.08		31.1	
Cpd 5	0.12		32.2	
Cpd 5	0.18		61.6	
Chlorantraniliprole	0.23		21	
Chlorantraniliprole	0.63		42.3	
Cpd 4 + Chlorantraniliprole	0.2 + 0.23	1:1.15	65.3*	45.3
Cpd 4 + Chlorantraniliprole	0.2 + 0.63	1:3.15	73.3*	60.1
Cpd 4 + Chlorantraniliprole	0.5 + 0.23	1:0.46	85.9*	76
Cpd 4 + Chlorantraniliprole	0.5 + 0.63	1:1.26	99.0*	82.5
Cpd 2 + Chlorantraniliprole	0.05 + 0.23	1:4.6	17.4	33.8
Cpd 2 + Chlorantraniliprole	0.05 + 0.63	1:12.6	36	51.7
Cpd 2 + Chlorantraniliprole	0.18 + 0.23	1:1.2778	42.8	50.2
Cpd 2 + Chlorantraniliprole	0.18 + 0.63	1:3.5	63.8*	63.6
Cpd 1 + Chlorantraniliprole	0.08 + 0.23	1:2.875	21.8	37.8
Cpd 1 + Chlorantraniliprole	0.08 + 0.63	1:7.875	43.3	54.6
Cpd 1 + Chlorantraniliprole	0.12 + 0.23	1:1.9167	26.9	48.8
Cpd 1 + Chlorantraniliprole	0.12 + 0.63	1:5.25	48.9	62.6
Cpd 3 + Chlorantraniliprole	0.02 + 0.23	1:11.5	19.8	36.7
Cpd 3 + Chlorantraniliprole	0.02 + 0.63	1:31.5	31.1	53.8
Cpd 3 + Chlorantraniliprole	0.08 + 0.23	1:2.875	26	45.5
Cpd 3 + Chlorantraniliprole	0.08 + 0.63	1:7.875	39.6	60.2
Cpd 5 + Chlorantraniliprole	0.12 + 0.23	1:1.9167	49.2*	46.5
Cpd 5 + Chlorantraniliprole	0.12 + 0.63	1:5.25	76.3*	60.9
Cpd 5 + Chlorantraniliprole	0.18 + 0.23	1:1.2778	60.3	69.7
Cpd 5 + Chlorantraniliprole	0.18 + 0.63	1:3.5	71.9	77.9

*Denotes enhanced efficacy based on Colby formula

Table 5f – Mixtures comprising Spinetoram and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		40.7	
Cpd 4	0.5		73.4	
Cpd 2	0.05		17.1	
Cpd 2	0.18		37.2	
Cpd 1	0.08		20.8	
Cpd 1	0.12		39.4	
Cpd 3	0.02		17.6	
Cpd 3	0.08		39	
Cpd 5	0.12		44.9	
Cpd 5	0.18		58.9	
Spinetoram	50		51.2	
Spinetoram	250		73.9	
Cpd 4 + Spinetoram	0.2 + 50	1:250	75.2*	71.1
Cpd 4 + Spinetoram	0.2 + 250	1:1250	93.0*	84.5
Cpd 4 + Spinetoram	0.5 + 50	1:100	82.7	87
Cpd 4 + Spinetoram	0.5 + 250	1:500	94.2*	93.1
Cpd 2 + Spinetoram	0.05 + 50	1:1000	46.7	59.5
Cpd 2 + Spinetoram	0.05 + 250	1:5000	79.1*	78.4
Cpd 2 + Spinetoram	0.18 + 50	1:277.7778	62.4	69.3
Cpd 2 + Spinetoram	0.18 + 250	1:1388.8889	81.4	83.6
Cpd 1 + Spinetoram	0.08 + 50	1:625	57.5	61.3
Cpd 1 + Spinetoram	0.08 + 250	1:3125	86.9*	79.3
Cpd 1 + Spinetoram	0.12 + 50	1:416.6667	63	70.4
Cpd 1 + Spinetoram	0.12 + 250	1:2083.3333	79.3	84.2
Cpd 3 + Spinetoram	0.02 + 50	1:2500	40.5	59.8
Cpd 3 + Spinetoram	0.02 + 250	1:12500	58.3	78.5
Cpd 3 + Spinetoram	0.08 + 50	1:625	48.9	70.2
Cpd 3 + Spinetoram	0.08 + 250	1:3125	78.1	84.1
Cpd 5 + Spinetoram	0.12 + 50	1:416.6667	66.9	73.1
Cpd 5 + Spinetoram	0.12 + 250	1:2083.3333	79.0	85.6
Cpd 5 + Spinetoram	0.18 + 50	1:277.7778	86.2*	79.9
Cpd 5 + Spinetoram	0.18 + 250	1:1388.8889	87.5	89.3

*Denotes enhanced efficacy based on Colby formula

Table 5g – Mixtures comprising Bifenthrin and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		21.0	
Cpd 4	0.5		45.1	
Cpd 2	0.05		23.4	
Cpd 2	0.18		38.3	
Cpd 1	0.08		29.1	
Cpd 1	0.12		35.9	
Cpd 3	0.02		15.8	
Cpd 3	0.08		30.6	
Cpd 5	0.12		25.8	
Cpd 5	0.18		44.3	
Bifenthrin	50		28.7	
Bifenthrin	250		58.2	
Cpd 4 + Bifenthrin	0.2 + 50	1:250	70.3*	43.6
Cpd 4 + Bifenthrin	0.2 + 250	1:1250	76.3*	66.9
Cpd 4 + Bifenthrin	0.5 + 50	1:100	84.2*	60.8
Cpd 4 + Bifenthrin	0.5 + 250	1:500	97.1*	77.0
Cpd 2 + Bifenthrin	0.05 + 50	1:1000	26.4	45.3
Cpd 2 + Bifenthrin	0.05 + 250	1:5000	52.8	67.9
Cpd 2 + Bifenthrin	0.18 + 50	1:277.7778	41.7	56.0
Cpd 2 + Bifenthrin	0.18 + 250	1:1388.8889	65.5	74.2
Cpd 1 + Bifenthrin	0.08 + 50	1:625	35.7	49.4
Cpd 1 + Bifenthrin	0.08 + 250	1:3125	69.0	70.3
Cpd 1 + Bifenthrin	0.12 + 50	1:416.6667	54.3	54.3
Cpd 1 + Bifenthrin	0.12 + 250	1:2083.3333	54.3	73.2
Cpd 3 + Bifenthrin	0.02 + 50	1:2500	22.8	40.0
Cpd 3 + Bifenthrin	0.02 + 250	1:12500	27.4	64.8
Cpd 3 + Bifenthrin	0.08 + 50	1:625	38.1	50.5
Cpd 3 + Bifenthrin	0.08 + 250	1:3125	63.8	71.0
Cpd 5 + Bifenthrin	0.12 + 50	1:416.6667	61.9*	47.1
Cpd 5 + Bifenthrin	0.12 + 250	1:2083.3333	83.8*	68.9
Cpd 5 + Bifenthrin	0.18 + 50	1:277.7778	81.9*	60.3
Cpd 5 + Bifenthrin	0.18 + 250	1:1388.8889	86.7*	76.7

*Denotes enhanced efficacy based on Colby formula

Table 5h – Mixtures comprising Indoxacarb and their activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		44.6	
Cpd 4	0.5		68.7	
Cpd 2	0.05		43.9	
Cpd 2	0.18		57.5	
Cpd 1	0.08		37.3	
Cpd 1	0.12		50.5	
Cpd 3	0.02		37.4	
Cpd 3	0.08		43.5	
Cpd 5	0.12		40.0	
Cpd 5	0.18		68.4	
Indoxacarb	50		23.4	
Indoxacarb	250		37.5	
Cpd 4 + Indoxacarb	0.2 + 50	1:250	79.6*	57.6
Cpd 4 + Indoxacarb	0.2 + 250	1:1250	55.6	65.3
Cpd 4 + Indoxacarb	0.5 + 50	1:100	94.2*	76.0
Cpd 4 + Indoxacarb	0.5 + 250	1:500	92.7*	80.4
Cpd 2 + Indoxacarb	0.05 + 50	1:1000	41.7	57.0
Cpd 2 + Indoxacarb	0.05 + 250	1:5000	38.8	64.9
Cpd 2 + Indoxacarb	0.18 + 50	1:277.7778	67.5*	67.5
Cpd 2 + Indoxacarb	0.18 + 250	1:1388.8889	74.0*	73.5
Cpd 1 + Indoxacarb	0.08 + 50	1:625	39.0	52.0
Cpd 1 + Indoxacarb	0.08 + 250	1:3125	68.8*	60.8
Cpd 1 + Indoxacarb	0.12 + 50	1:416.6667	61.8	62.1
Cpd 1 + Indoxacarb	0.12 + 250	1:2083.3333	79.3*	69.1
Cpd 3 + Indoxacarb	0.02 + 50	1:2500	71.3*	52.1
Cpd 3 + Indoxacarb	0.02 + 250	1:12500	60.8	60.9
Cpd 3 + Indoxacarb	0.08 + 50	1:625	57.0*	56.8
Cpd 3 + Indoxacarb	0.08 + 250	1:3125	56.1	64.7
Cpd 5 + Indoxacarb	0.12 + 50	1:416.6667	76.4*	54.1
Cpd 5 + Indoxacarb	0.12 + 250	1:2083.3333	89.0*	62.5
Cpd 5 + Indoxacarb	0.18 + 50	1:277.7778	80.3*	75.8
Cpd 5 + Indoxacarb	0.18 + 250	1:1388.8889	87.5*	80.3

*Denotes enhanced efficacy based on Colby formula

Table 5i – Mixtures with Sulfoxaflor and activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		33.5	
Cpd 4	0.5		80.3	
Cpd 5	0.12		41.6	
Cpd 5	0.18		70.3	
Sulfoxaflor	0.016		23.4	
Sulfoxaflor	0.04		48.0	
Cpd 4 + Sulfoxaflor	0.2 + 0.016	1:0.08	61.3*	49.1
Cpd 4 + Sulfoxaflor	0.2 + 0.04	1:0.2	77.9*	65.4
Cpd 4 + Sulfoxaflor	0.5 + 0.016	1:0.032	87.0*	84.9
Cpd 4 + Sulfoxaflor	0.5 + 0.04	1:0.08	97.1*	89.7
Cpd 5 + Sulfoxaflor	0.12 + 0.016	1:0.133	56.3*	55.2
Cpd 5 + Sulfoxaflor	0.12 + 0.04	1:0.333	85.3*	69.6
Cpd 5 + Sulfoxaflor	0.18 + 0.016	1:0.088	89.6*	77.2
Cpd 5 + Sulfoxaflor	0.18 + 0.04	1:0.222	97.9*	84.5

*Denotes enhanced efficacy based on Colby formula

5

10

15

Table 5j – Mixtures with Flupyradifurone and activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		33.5	
Cpd 4	0.5		80.3	
Cpd 5	0.12		41.6	
Cpd 5	0.18		70.3	
Flupyradifurone	0.02		23.4	
Flupyradifurone	0.075		48.0	
Cpd 4 + Flupyradifurone	0.2 + 0.02	1:0.1	61.3*	49.1
Cpd 4 + Flupyradifurone	0.2 + 0.075	1:0.375	77.9*	65.4
Cpd 4 + Flupyradifurone	0.5 + 0.02	1:0.04	87*	84.9
Cpd 4 + Flupyradifurone	0.5 + 0.075	1:0.15	97.1*	89.7
Cpd 5 + Flupyradifurone	0.12 + 0.02	1:0.166	56.3*	55.2
Cpd 5 + Flupyradifurone	0.12 + 0.075	1:0.625	85.3*	69.6
Cpd 5 + Flupyradifurone	0.18 + 0.02	1:0.111	89.6*	77.2
Cpd 5 + Flupyradifurone	0.18 + 0.075	1:0.417	97.9*	84.5

*Denotes enhanced efficacy based on Colby formula

5

Table 5k – Mixtures with Flonicamid and activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		40.2	
Cpd 4	0.5		69.3	
Cpd 5	0.12		35.9	
Cpd 5	0.18		67.2	
Flonicamid	0.027		24.2	
Flonicamid	0.267		49.3	
Cpd 4 + Flonicamid	0.2 + 0.027	1:0.135	69.4*	54.7
Cpd 4 + Flonicamid	0.2 + 0.267	1:1.335	78.6*	69.7
Cpd 4 + Flonicamid	0.5 + 0.027	1:0.054	80.1*	76.7
Cpd 4 + Flonicamid	0.5 + 0.267	1:0.534	93.8*	84.4
Cpd 5 + Flonicamid	0.12 + 0.027	1:0.225	62.9*	51.4
Cpd 5 + Flonicamid	0.12 + 0.267	1:2.225	78.8*	67.5
Cpd 5 + Flonicamid	0.18 + 0.027	1:0.15	77.7*	75.2
Cpd 5 + Flonicamid	0.18 + 0.267	1:1.483	72.7	83.4

*Denotes enhanced efficacy based on Colby formul

Table 51 – Mixtures with Chlorpyrifos and activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		40.2	
Cpd 4	0.5		69.3	
Cpd 5	0.12		35.9	
Cpd 5	0.18		67.2	
Chlorpyrifos	0.173		17.7	
Chlorpyrifos	0.624		37.9	
Cpd 4 + Chlorpyrifos	0.2 + 0.173	1:0.865	78.9*	50.8
Cpd 4 + Chlorpyrifos	0.2 + 0.624	1:3.12	83.7*	62.9
Cpd 4 + Chlorpyrifos	0.5 + 0.173	1:0.346	86.8*	74.7
Cpd 4 + Chlorpyrifos	0.5 + 0.624	1:1.248	94.5*	80.9
Cpd 5 + Chlorpyrifos	0.12 + 0.173	1:1.442	81.3*	47.2
Cpd 5 + Chlorpyrifos	0.12 + 0.624	1:5.2	92.5*	60.2
Cpd 5 + Chlorpyrifos	0.18 + 0.173	1:0.961	78.3*	73
Cpd 5 + Chlorpyrifos	0.18 + 0.624	1:3.467	93.2*	79.7

*Denotes enhanced efficacy based on Colby formula

5

10

15

Table 5m – Mixtures with Permethrin and activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		42.5	
Cpd 4	0.5		65	
Cpd 5	0.12		37.3	
Cpd 5	0.18		61.7	
Permethrin	0.0004		15.6	
Permethrin	0.04		32.1	
Cpd 4 + Permethrin	0.2 + 0.0004	1:0.002	65*	51.5
Cpd 4 + Permethrin	0.5 + 0.04	1:0.2	62.3*	61
Cpd 4 + Permethrin	0.12 + 0.0004	1:0.0008	84.2	87.8
Cpd 4 + Permethrin	0.18 + 0.04	1:0.08	86.5*	76.3
Cpd 5 + Permethrin	0.2 + 0.0004	1:0.003	51.1	51.5
Cpd 5 + Permethrin	0.5 + 0.04	1:0.333	65.1*	61
Cpd 5 + Permethrin	0.12 + 0.0004	1:0.002	72.1*	70.5
Cpd 5 + Permethrin	0.18 + 0.04	1:0.222	69.9	76.3

*Denotes enhanced efficacy based on Colby formula

5

Table 5n – Mixtures with Tolfenpyrad and activity on Cotton Melon Aphid

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	0.2		57.1	
Cpd 4	0.5		73.9	
Cpd 5	0.12		48.9	
Cpd 5	0.18		77.7	
Tolfenpyrad	0.192		29	
Tolfenpyrad	0.753		61.9	
Cpd 4 + Tolfenpyrad	0.2 + 0.192	1:0.96	78.2*	69.6
Cpd 4 + Tolfenpyrad	0.2 + 0.753	1:3.765	87.1*	83.7
Cpd 4 + Tolfenpyrad	0.5 + 0.192	1:0.384	92.6*	81.5
Cpd 4 + Tolfenpyrad	0.5 + 0.753	1:1.506	98.4*	90.1
Cpd 5 + Tolfenpyrad	0.12 + 0.192	1:1.6	84*	63.7
Cpd 5 + Tolfenpyrad	0.12 + 0.753	1:6.275	85.8*	80.5
Cpd 5+ Tolfenpyrad	0.18 + 0.192	1:1.067	89.1*	84.2
Cpd 5 + Tolfenpyrad	0.18 + 0.753	1:4.183	90.8	91.5

*Denotes enhanced efficacy based on Colby formula

TEST E

For evaluating control of diamondback moth (*Plutella xylostella* L.) through contact and/or systemic means, each test unit consisted of a small open container with a 10- to 12-day-old mustard plant inside. Test compounds were formulated and sprayed with 3 replications as described for Test A. After spraying, the test units were allowed to dry for 1 hour before they were infested with 30-50 neonate larvae. A black, screened cap was placed on the top of each container. The test units were held for 6 days in a growth chamber at 24–25 °C and 70% relative humidity. Each test unit was then visually assessed for plant damage and rated 0-10, where 0 = undamaged and 10 = plant is dead; the results are listed in Tables 6a-b.

For evaluating control of diamondback moth (*Plutella xylostella* L.) through ingestion only, the test substance Dipel® (active ingredient *Bacillus thuringiensis kurstaki*) was formulated by diluting in water and then adding an MSO based adjuvant at 0.1% v/v. Cabbage plants were sprayed with Dipel® to runoff and allowed to dry for three hours. Once dry, plants leaf tissue was excised and one piece was placed into one well in an eight well tray. Test units were maintained in a growth chamber at 24–25 °C and 70% relative humidity. Each well was then inspected for mortality after 4 days; the result is listed in Table 6c.

For evaluating control of fall armyworm (*Spodoptera frugiperda* (J.E. Smith)) the test unit consisted of a small open container with a 4–5-day-old corn (maize) plant inside. This was pre-infested with 10–15 1-day-old larvae on a piece of insect diet. Test compounds were formulated and sprayed with 3 replications as described for Test A. After spraying of the formulated test compound, the test units were maintained in a growth chamber for 6 days at 25 °C and 70% relative humidity. Plant feeding damage was then visually assessed based on foliage consumed, and larvae were assessed for mortality. Each test unit was then visually assessed for plant damage and rated 0-10, where 0 = undamaged and 10 = plant is dead; the results are listed in Tables 6d-i.

30

35

Table 6a – Mixtures comprising Cyantraniliprole and their activity on Diamondback

<u>Moth</u>				
Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0.0	
Cpd 4	250		0.0	
Cpd 2	50		0.0	
Cpd 2	250		0.0	
Cpd 1	50		0.0	
Cpd 1	250		0.0	
Cpd 3	50		0.0	
Cpd 3	250		0.0	
Cpd 5	50		0.0	
Cpd 5	250		0.0	
Cyantraniliprole	0.02		20.0	
Cyantraniliprole	0.013		73.3	
Cpd 4 + Cyantraniliprole	50 + 0.02	1:0.0004	50.0*	20.0
Cpd 4 + Cyantraniliprole	50 + 0.013	1:0.00026	63.3	73.3
Cpd 4 + Cyantraniliprole	250 + 0.02	1:0.00008	50.0*	20.0
Cpd 4 + Cyantraniliprole	250 + 0.013	1:0.000052	90.0*	73.3
Cpd 2 + Cyantraniliprole	50 + 0.02	1:0.0004	46.7*	20.0
Cpd 2 + Cyantraniliprole	50 + 0.013	1:0.00026	86.7*	73.3
Cpd 2 + Cyantraniliprole	250 + 0.02	1:0.00008	23.3*	20.0
Cpd 2 + Cyantraniliprole	250 + 0.013	1:0.000052	63.3	73.3
Cpd 1 + Cyantraniliprole	50 + 0.02	1:0.0004	33.3*	20.0
Cpd 1 + Cyantraniliprole	50 + 0.013	1:0.00026	73.3	73.3
Cpd 1 + Cyantraniliprole	250 + 0.02	1:0.00008	63.3*	20.0
Cpd 1 + Cyantraniliprole	250 + 0.013	1:0.000052	66.7	73.3
Cpd 3 + Cyantraniliprole	50 + 0.02	1:0.0004	26.7*	20.0
Cpd 3 + Cyantraniliprole	50 + 0.013	1:0.00026	76.7*	73.3
Cpd 3 + Cyantraniliprole	250 + 0.02	1:0.00008	56.7*	20.0
Cpd 3 + Cyantraniliprole	250 + 0.013	1:0.000052	66.7	73.3
Cpd 5 + Cyantraniliprole	50 + 0.02	1:0.0004	6.7	20.0
Cpd 5 + Cyantraniliprole	50 + 0.013	1:0.00026	70.0	73.3
Cpd 5 + Cyantraniliprole	250 + 0.02	1:0.00008	46.7*	20.0
Cpd 5 + Cyantraniliprole	250 + 0.013	1:0.000052	70.0	73.3

*Denotes enhanced plant protection based on Colby formula

Table 6b – Mixtures comprising Spirotetramat and their activity on Diamondback Moth

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0.0	
Cpd 4	250		0.0	
Cpd 2	50		0.0	
Cpd 2	250		0.0	
Cpd 1	50		0.0	
Cpd 1	250		0.0	
Cpd 3	50		0.0	
Cpd 3	250		0.0	
Cpd 5	50		0.0	
Cpd 5	250		0.0	
Spirotetramat	83.2		73.3	
Spirotetramat	150		80.0	
Cpd 4 + Spirotetramat	50 + 83.23	1:2	76.7*	73.3
Cpd 4 + Spirotetramat	50 + 150	1:5	96.7*	80.0
Cpd 4 + Spirotetramat	250 + 83.23	1:0.4	86.7*	73.3
Cpd 4 + Spirotetramat	250 + 150	1:1	93.3*	80.0
Cpd 2 + Spirotetramat	50 + 83.23	1:2	93.3*	73.3
Cpd 2 + Spirotetramat	50 + 150	1:5	86.7*	80.0
Cpd 2 + Spirotetramat	250 + 83.23	1:0.4	90.0*	73.3
Cpd 2 + Spirotetramat	250 + 150	1:1	80.0	80.0
Cpd 1 + Spirotetramat	50 + 83.23	1:2	93.3*	73.3
Cpd 1 + Spirotetramat	50 + 150	1:5	93.3*	80.0
Cpd 1 + Spirotetramat	250 + 83.23	1:0.4	90.0*	73.3
Cpd 1 + Spirotetramat	250 + 150	1:1	96.7*	80.0
Cpd 3 + Spirotetramat	50 + 83.23	1:2	83.3*	73.3
Cpd 3 + Spirotetramat	50 + 150	1:5	93.3*	80.0
Cpd 3 + Spirotetramat	250 + 83.23	1:0.4	83.3*	73.3
Cpd 3 + Spirotetramat	250 + 150	1:1	86.7*	80.0
Cpd 5 + Spirotetramat	50 + 83.23	1:2	90.0*	73.3
Cpd 5 + Spirotetramat	50 + 150	1:5	93.3*	80.0
Cpd 5 + Spirotetramat	250 + 83.23	1:0.4	73.3	73.3
Cpd 5 + Spirotetramat	250 + 150	1:1	93.3*	80.0

*Denotes enhanced plant protection based on Colby formula

Table 6c – Mixtures with Dipel®(*Bacillus thuringiensis kurstaki*) and activity on
Diamondback Moth

Treatment	Rate [ppm]	Ratio	% Mortality (obs)	% Mortality (exp)
Cpd 4	50		12.5	
Cpd 4	250		17	
Cpd 5	50		21	
Cpd 5	250		25	
Dipel®	0.15		25	
Dipel®	0.43		50	
Cpd 4 + Dipel®	50 + 0.15	1:0.003	25	34.4
Cpd 4 + Dipel®	50 + 0.43	1:0.0086	70.8*	56.3
Cpd 4 + Dipel®	250 + 0.15	1:0.0006	45.8*	37.5
Cpd 4 + Dipel®	250 + 0.43	1:0.00172	62.5*	58.3
Cpd 5 + Dipel®	50 + 0.15	1:0.003	29.2	34.4
Cpd 5 + Dipel®	50 + 0.43	1:0.0086	41.7	56.3
Cpd 5 + Dipel®	250 + 0.15	1:0.0006	16.7	37.5
Cpd 5 + Dipel®	250 + 0.43	1:0.00172	100*	58.3

*Denotes enhanced plant protection based on Colby formula

5

Table 6d – Mixtures with Methomyl and activity on Fall Armyworm

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0	
Cpd 4	250		0	
Cpd 5	50		0	
Cpd 5	250		13.3	
Methomyl	8.5		43.3	
Cpd 4 + Methomyl	50 + 8.5	1:0.17	36.7	43.3
Cpd 4 + Methomyl	250 + 8.5	1:0.034	80*	43.3
Cpd 5 + Methomyl	50 + 8.5	1:0.17	100*	43.3
Cpd 5 + Methomyl	250 + 8.5	1:0.034	76.7*	50.9

*Denotes enhanced plant protection based on Colby formula

10

Table 6e – Mixtures with Indoxacarb and activity on Fall Armyworm

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0	
Cpd 4	250		0	
Cpd 2	50		0	
Cpd 2	250		0	
Cpd 1	50		0	
Cpd 1	250		0	
Cpd 3	50		0	
Cpd 3	250		0	
Cpd 5	50		0	
Cpd 5	250		0	
Indoxacarb	0.2		70	
Indoxacarb	0.5		70	
Cpd 4 + Indoxacarb	50 + 0.2	1:2	60	70
Cpd 4 + Indoxacarb	50 + 0.5	1:5	93.3*	70
Cpd 4 + Indoxacarb	250+ 0.2	1:0.4	66.7	70
Cpd 4 + Indoxacarb	250 + 0.5	1:1	96.7*	70
Cpd 2 + Indoxacarb	50 + 0.2	1:2	50	70
Cpd 2 + Indoxacarb	50 + 0.5	1:5	93.3*	70
Cpd 2 + Indoxacarb	250+ 0.2	1:0.4	70	70
Cpd 2+ Indoxacarb	250 + 0.5	1:1	100*	70
Cpd 1 + Indoxacarb	50 + 0.2	1:2	30.0	70
Cpd 1 + Indoxacarb	50 + 0.5	1:5	86.7*	70
Cpd 1 + Indoxacarb	250+ 0.2	1:0.4	46.7	70
Cpd 1 + Indoxacarb	250 + 0.5	1:1	90*	70
Cpd 3 + Indoxacarb	50 + 0.2	1:2	43.3	70
Cpd 3 + Indoxacarb	50 + 0.5	1:5	73.3*	70
Cpd 3 + Indoxacarb	250+ 0.2	1:0.4	73.3*	70
Cpd 3 + Indoxacarb	250 + 0.5	1:1	86.7*	70
Cpd 5 + Indoxacarb	50 + 0.2	1:2	56.7	70
Cpd 5 + Indoxacarb	50 + 0.5	1:5	93.3*	70
Cpd 5 + Indoxacarb	250+ 0.2	1:0.4	60	70
Cpd 5 + Indoxacarb	250 + 0.5	1:1	93.3*	70

*Denotes enhanced plant protection based on Colby formula

Table 6f – Mixtures with Emamectin Benzoate and activity on Fall Armyworm

5

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0	
Cpd 4	250		0	
Cpd 5	50		0	
Cpd 5	250		0	
Emamectin Benzoate	0.03		43.3	
Emamectin Benzoate	0.06		23.3	
Cpd 4 + Emamectin Benzoate	50 + 0.03	1:0.0006	73.3	43.3
Cpd 4 + Emamectin Benzoate	50 + 0.06	1:0.0012	96.7	23.3
Cpd 4 + Emamectin Benzoate	250 + 0.03	1:0.00012	96.7	43.3
Cpd 4 + Emamectin Benzoate	250 + 0.06	1:0.00024	96.7	23.3
Cpd 5 + Emamectin Benzoate	50 + 0.03	1:0.0006	70	43.3
Cpd 5 + Emamectin Benzoate	50 + 0.06	1:0.0012	60	23.3
Cpd 5 + Emamectin Benzoate	250 + 0.03	1:0.00012	43.3	43.3
Cpd 5 + Emamectin Benzoate	250 + 0.06	1:0.00024	96.7	23.3

*Denotes enhanced plant protection based on Colby formula

10

15

20

Table 6g – Mixtures with Chlorfenapyr and activity on Fall Armyworm

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0	
Cpd 4	250		20	
Cpd 5	50		0	
Cpd 5	250		0	
Chlorfenapyr	3.7		66.7	
Chlorfenapyr	5.0		53.3	
Cpd 4 + Chlorfenapyr	50 + 3.7	1:0.074	30	66.7
Cpd 4 + Chlorfenapyr	50 + 5	1:0.1	60*	53.3
Cpd 4 + Chlorfenapyr	250 + 3.7	1:0.0148	80*	73.3
Cpd 4 + Chlorfenapyr	250 + 5	1:0.02	93.3*	62.7
Cpd 5 + Chlorfenapyr	50 + 3.7	1:0.074	70*	66.7
Cpd 5 + Chlorfenapyr	50 + 5	1:0.1	93.3*	53.3
Cpd 5 + Chlorfenapyr	250 + 3.7	1:0.0148	76.7*	66.7
Cpd 5 + Chlorfenapyr	250 + 5	1:0.02	100*	53.3

*Denotes enhanced plant protection based on Colby formula

5

Table 6h – Mixtures with Methoxyfenozide and activity on Fall Armyworm

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0	
Cpd 4	250		20	
Cpd 5	50		0	
Cpd 5	250		0	
Methoxyfenozide	2.8		36.7	
Cpd 4 + Methoxyfenozide	50 + 2.8	1:0.056	90*	36.7
Cpd 4 + Methoxyfenozide	250 + 2.8	1:0.011	86.7*	49.3
Cpd 5 + Methoxyfenozide	50 + 2.8	1:0.056	80*	36.7
Cpd 5 + Methoxyfenozide	250 + 2.8	1:0.011	66.7*	36.7

*Denotes enhanced plant protection based on Colby formula

10

Table 6i – Mixtures with Novaluron and activity on Fall Armyworm

Treatment	Rate [ppm]	Ratio	Plant Protection (obs)	Plant Protection (exp)
Cpd 4	50		0.0	
Cpd 4	250		0.0	
Cpd 5	50		0.0	
Cpd 5	250		13.3	
Novaluron	0.21		33.3	
Novaluron	0.36		80.0	
Cpd 4 + Novaluron	50 + 0.21	1:0.004	60.0*	33.3
Cpd 4 + Novaluron	50 + 0.36	1:0.007	83.3*	80.0
Cpd 4 + Novaluron	250 + 0.21	1:0.0008	80.0*	33.3
Cpd 4 + Novaluron	250 + 0.36	1:0.001	76.7	80.0
Cpd 5 + Novaluron	50 + 0.21	1:0.004	53.3*	33.3
Cpd 5 + Novaluron	50 + 0.36	1:0.007	80.0	80.0
Cpd 5 + Novaluron	250 + 0.21	1:0.0008	33.3	42.2
Cpd 5 + Novaluron	250 + 0.36	1:0.001	73.3	82.7

*Denotes enhanced plant protection based on Colby formula

5

TEST F

Tomato seeds (cv Tiny Tim) were germinated in plugs in Sunshine mix for 21 days. Seedlings with 3-4 true leaves were transplanted into 4" pots filled with pasteurized loamy sand soil (82.6% sand, 11.6% silt, and 5.6% clay, organic matter 1%, pH 6.6). The soil was treated at transplanting according to Table 1. 10% (v/v) acetone or water were included as negative controls and abamectin at 10 ppm was included as a positive control. Second application was made 7 days after transplanting and plants were inoculated with 7500 eggs of root knot nematode (*Meloidogyne incognita*). Each treatment had four replications. The plants were watered and fertilized as needed. The temperature in the greenhouse ranged from 25 to 30 °C. The assay was terminated after 6 weeks and severity of nematode infection was evaluated using 0-10 galling index scale (Bridge and Page 1980).

The observed % reduction of root galling (X) was calculated using the formula:

$$X = (UTC - T) / UTC * 100\%$$

where X – observed % reduction of root galling, UTC - galling in non-treated control, T – galling in treatment X.

Results are shown in Table 7a.

5 Table 7a – Mixtures with Biologicals and activity on Root Knot Nematode

Treatment	Rate [ppm]	Observed Reduction (%)	Expected Reduction (%)
Cpd 2 + CH200&CH201 ^{*1}	250 ppm + 5.1 E+09 CFU/pot	0	12
Cpd 3 + CH200&CH201	250 ppm + 5.1 E+09 CFU/pot	4	4
Cpd 4 + CH200&CH201	250 ppm + 5.1 E+09 CFU/pot	4	23
Cpd 5 + CH200&CH201	250 ppm + 5.1 E+09 CFU/pot	4	12
Cpd 1 + CH200&CH201	250 ppm + 5.1 E+09 CFU/pot	4	12
Cpd 2 + RTI545 ^{*2}	250 ppm + 2.78E+09 CFU/pot	0	4
Cpd 3 + RTI545	250 ppm + 2.78E+09 CFU/pot	4	-4
Cpd 4 + RTI545	250 ppm + 2.78E+09 CFU/pot	4	16
Cpd 5 + RTI545	250 ppm + 2.78E+09 CFU/pot	4	4
Cpd 1 + RTI545	250 ppm + 2.78E+09 CFU/pot	8	4
Cpd 2 + Nortica ^{*3}	250 ppm + 3.00E+10 CFU/pot	4	16
Cpd 3 + Nortica	250 ppm + 3.00E+10 CFU/pot	0	9
Cpd 4 + Nortica	250 ppm + 3.00E+10 CFU/pot	4	26
Cpd 5 + Nortica	250 ppm + 3.00E+10 CFU/pot	8	16
Cpd 1 + Nortica	250 ppm + 3.00E+10 CFU/pot	-4	16
Cpd 2 + FMC proprietary <i>Brevibacillus laterosporus</i> strain	250 ppm, (2.5 mg/pot) + 50% (v/v)	5	9
Cpd 3 + FMC proprietary <i>Brevibacillus laterosporus</i> strain	250 ppm, (2.5 mg/pot) + 50% (v/v)	5	9
Cpd 4 + FMC proprietary <i>Brevibacillus laterosporus</i> strain	250 ppm, (2.5 mg/pot) + 50% (v/v)	0	9
Cpd 5 + FMC proprietary <i>Brevibacillus laterosporus</i> strain	250 ppm, (2.5 mg/pot) + 50% (v/v)	14	17
Cpd 1 + FMC proprietary <i>Brevibacillus laterosporus</i> strain	250 ppm, (2.5 mg/pot) + 50% (v/v)	9	9

^{*1}see, WO 2018/045063 A1; ^{*2}see, WO 2018/067815 A1; ^{*3}Metronidazole

Tables 2a through 7a show mixtures and compositions of the present disclosure demonstrating control on a wide range of invertebrate pests, some with more than additive activity. As the % of mortality cannot exceed 100%, the unexpected increase in insecticidal activity can be greatest only when the separate active ingredient components alone are at application rates providing considerably less than 100% control. Likewise, plant damage cannot exceed a rating of 10 in these tests, and the unexpected increase in insecticidal activity can be greatest only when the separate active ingredient components alone are at application rates providing considerably less than 100% control. Enhanced activity may not be evident at low application rates where the individual active ingredient components alone have little activity. However, in some instances high activity was observed for combinations wherein individual active ingredient alone at the same application rate had essentially no activity. It is worth noting that the five experimental compounds are mixed with 25 representative active ingredients known to have various mode of action classifications; all were shown to enhance activity across the seven representative insect species. Mixtures were evaluated for both contact and/or systemic activity and the results indicate that both foliar and soil applied products and mixtures demonstrate enhanced activity.

A non-synthetic insecticide derived from *Bacillus* spp., was also evaluated for its ability to provide enhanced efficacy when mixed with cpds 1-5. Enhanced activity was also demonstrated in these mixtures.

Accordingly, this invention provides not only improved compositions but also methods of their use for control of invertebrate pests such as arthropods and nematodes in both agronomic and non-agronomic environments. The compositions of this invention demonstrate high controlling effect of invertebrate and nematode pests; consequently, their use as arthropodicides and nematicides can reduce crop production cost and environmental load.


CLAIMS

What is claimed is:

1. A composition comprising:

(a) at least one compound of


5 Formula I:



Formula I

wherein n is 0, 1, or 2; or

10 at least one compound of Formula II :



Formula II

wherein R is CH₃, or CF₂H; or

a combination of the foregoing compounds; and

15 (b) at least one additional invertebrate pest control agent different from the compound of Formula I, or the compound of Formula II.

2. The composition of claim 1 wherein component (a) is a compound of Formula **I** wherein n is 0.

5 3. The composition of claim 1 wherein component (a) is a compound of Formula **I** wherein n is 1.

4. The composition of claim 1 wherein component (a) is a compound of Formula **I** wherein n is 2.

10 5. The composition of claim 1 wherein component (a) is a compound of Formula **II** wherein R is CH₃.

6. The composition of claim 1 wherein component (a) is a compound of Formula **II** wherein R is CF₂H.

15 7. The composition of claim 1 wherein component (a) is a compound selected from *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

25 8. The composition of claim 1 or 7 wherein component (a) is a compound selected from *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide, and *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

30 9. The composition of claim 1 or 7 wherein component (a) is a compound selected from *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide.

10. The composition of any one of claims 1-9 wherein component (b) is selected from an insecticide, fungicide, nematocide, and bactericide.

11. The composition of any one of claims 1-10 wherein component (b) is an insecticide.

12. The composition of any one of claim 1-11 wherein component (b) is selected from
 5 abamectin, acephate, acequinocyl, acetamiprid, acrinathrin, acynonapyr, afidopyropen
 ([*(3S,4R,4aR,6S,6aS,12R,12aS,12bS)*-3-[(cyclopropylcarbonyl)oxy]-
 1,3,4,4a,5,6,6a,12,12a,12b-decahydro-6,12-dihydroxy-4,6a,12b-trimethyl-11-oxo-9-(3-
 pyridinyl)-2*H*,11*H*-naphtho[2,1-*b*]pyrano[3,4-*e*]pyran-4-yl)methyl
 cyclopropanecarboxylate), amidoflumet, amitraz, avermectin, azadirachtin, azinphos-methyl,
 10 benfuracarb, bensultap, benzpyrimoxan, bifenthrin, kappa-bifenthrin, bifenazate, bistrifluron,
 borate, broflanilide, buprofezin, cadusafos, carbaryl, carbofuran, cartap, carzol,
 chlorantraniliprole, chlorfenapyr, chlorfluazuron, chlorprallethrin, chlorpyrifos,
 chlorpyrifos-*e*, chlorpyrifos-methyl, chromafenozide, clofentezin, chlorprallethrin,
 clothianidin, cyantraniliprole (Cyazypyr™) (3-bromo-1-(3-chloro-2-pyridinyl)-*N*-[4-cyano-
 15 2-methyl-6-[(methylamino)carbonyl]phenyl]-1*H*-pyrazole-5-carboxamide), cyclaniliprole (3-
 bromo-*N*-[2-bromo-4-chloro-6-[[*(1-cyclopropylethyl)*amino]carbonyl]phenyl]-1-(3-chloro-2-
 pyridinyl)-1*H*-pyrazole-5-carboxamide), cyclobutrifluram, cycloprothrin, cycloxaprid
 ((*5S,8R*)-1-[(6-chloro-3-pyridinyl)methyl]-2,3,5,6,7,8-hexahydro-9-nitro-5,8-Epoxy-1*H*-
 imidazo[1,2-*a*]azepine), cyenopyrafen, cyetpyrafen, cyflumetofen, cyfluthrin,
 20 beta-cyfluthrin, cyhalodiamide, cyhalothrin, gamma-cyhalothrin, lambda-cyhalothrin,
 cypermethrin, alpha-cypermethrin, zeta-cypermethrin, cyromazine, deltamethrin,
 diafenthiuron, diazinon, dicloromezotiaz, dieldrin, diflubenzuron, dimefluthrin, dimehypo,
 dimethoate, dimpropyridaz, dinotefuran, diofenolan, DiPel®emamectin, emamectin benzoate,
 endosulfan, esfenvalerate, ethiprole, etofenprox, epsilon-metofluthrin, etoxazole, fenbutatin
 25 oxide, fenitrothion, fenothiocarb, fenoxycarb, fenpropathrin, fenvalerate, fipronil,
 flometoquin (2-ethyl-3,7-dimethyl-6-[4-(trifluoromethoxy)phenoxy]-4-quinolinyl methyl
 carbonate), flonicamid, fluazaindolizine, flubendiamide, flucythrinate, flufenimer,
 flufenoxuron, flufenoxystrobin (methyl (*αE*)-2-[[2-chloro-4-
 (trifluoromethyl)phenoxy]methyl]-*α*-(methoxymethylene)benzeneacetate), fluensulfone (5-
 30 chloro-2-[(3,4,4-trifluoro-3-buten-1-yl)sulfonyl]thiazole), fluhexafon, fluopyram,
 flupentiofenox, flupiprole (1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-5-[(2-methyl-2-
 propen-1-yl)amino]-4-[(trifluoromethyl)sulfinyl]-1*H*-pyrazole-3-carbonitrile),
 flupyradifurone (4-[[*(6-chloro-3-pyridinyl)*methyl](2,2-difluoroethyl)amino]-2(*5H*)-
 furanone), flupyrimin, fluvalinate, tau-fluvalinate, fluxametamide, fonophos, formetanate,
 35 fosthiazate, gamma-cyhalothrin, halofenozide, heptafluthrin ([2,3,5,6-tetrafluoro-4-
 (methoxymethyl)phenyl)methyl 2,2-dimethyl-3-[(1*Z*)-3,3,3-trifluoro-1-propen-1-

yl]cyclopropanecarboxylate), hexaflumuron, hexythiazox, hydramethylnon, imidacloprid, indoxacarb, insecticidal soaps, isofenphos, isocycloseram, kappa-tefluthrin, lambda-cyhalothrin, lufenuron, malathion, meperfluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl (1*R*,3*S*)-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate), metaflumizone, metaldehyde, methamidophos, methidathion, methiocarb, methomyl, methoprene, methoxychlor, metofluthrin, methoxyfenozide, epsilon-metofluthrin, metronidazole epsilon-momfluorothrin, monocrotophos, monofluorothrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 3-(2-cyano-1-propen-1-yl)-2,2-dimethylcyclopropanecarboxylate), nicofluprole, nicotine, nitenpyram, nithiazine, novaluron, noviflumuron, oxamyl, oxazosulfyl, parathion, parathion-methyl, permethrin, phorate, phosalone, phosmet, phosphamidon, pirimicarb, profenofos, profluthrin, propargite, protrifenbute, pyflubumide (1,3,5-trimethyl-*N*-(2-methyl-1-oxopropyl)-*N*-[3-(2-methylpropyl)-4-[2,2,2-trifluoro-1-methoxy-1-(trifluoromethyl)ethyl]phenyl]-1*H*-pyrazole-4-carboxamide), pymetrozine, pyrafluprole, pyrethrin, pyridaben, pyridalyl, pyrifluquinazon, pyriminostrobin (methyl (αE)-2-[[[2-[(2,4-dichlorophenyl)amino]-6-(trifluoromethyl)-4-pyrimidinyl]oxy]methyl]- α -(methoxymethylene)benzeneacetate), pydiflumetofen, pyriprole, pyriproxifen, rotenone, ryanodine, silafluofen, spinetoram, spinosad, spirodiclofen, spiromesifen, spiropidion, spirotetramat, sulprofos, sulfoxaflor (*N*-[methyloxido[1-[6-(trifluoromethyl)-3-pyridinyl]ethyl]- λ^4 -sulfanylidene]cyanamide), tebufenozide, tebufenpyrad, teflubenzuron, tefluthrin, kappa-tefluthrin, terbufos, tetrachlorantraniliprole, tetrachlorvinphos, tetramethrin, tetramethylfluthrin ([2,3,5,6-tetrafluoro-4-(methoxymethyl)phenyl]methyl 2,2,3,3-tetramethylcyclopropanecarboxylate), tetraniliprole, thiacloprid, thiamethoxam, thiodicarb, thiosultap-sodium, tioazafen (3-phenyl-5-(2-thienyl)-1,2,4-oxadiazole), tolfenpyrad, tralomethrin, triazamate, trichlorfon, triflumezopyrim (2,4-dioxo-1-(5-pyrimidinylmethyl)-3-[3-(trifluoromethyl)phenyl]-2*H*-pyrido[1,2-*a*]pyrimidinium inner salt), triflumuron, tyclopyrazoflor, zeta-cypermethrin, *Bacillus thuringiensis* delta-endotoxins, entomopathogenic bacteria, entomopathogenic viruses, and entomopathogenic fungi.

13. The composition of any one of claims 1-12 wherein the additional pest control agent (b) is selected from cyantraniliprole, acetamiprid, imidacloprid, spirotetramat, spirodiclofen, chlorantraniliprole, bifenthrin, indoxacarb, avermectin, *Bacillus* spp. and any active crystal proteins, buprofezin, carbofuran, chlorfenapyr, chlorpyrifos, clothianidin, cyromazine, diafenthiuron, dinotefuran, DiPel® emamectin benzoate, fipronil, flonicamid, fluhexafon, flupyradifurone, methomyl (Lannate®), methoxyfenozide, metronidazole, novaluron, permethrin, pyriproxifen, sulfoxaflor, thiamethoxam, permethrin, γ -cyhalothrin,

or ζ -cypermethrin, broflanilide, dimpropyridaz, isocycloseram, tetrachlorantraniliprole, tolfenpyrad, oxazosulfyl, tyclopyrazoflor, flupyrimin, spiropidion, acynonapyr, benzpyrimoxan, chloroprallethrin, epsilon-metofluthrin, kappa-bifenthrin, dicloromezotiaz, and kappa-tefluthrin.

5

14. The composition of any one of claims 1-13 wherein the additional pest control agent (b) is selected from cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spiroidiclofen, sulfoxaflo, tolfenpyrad, chlorantraniliprole, or indoxacarb.

10

15. The composition of claim 1 wherein component (a) is *N*-[1,1-dimethyl-2-(methylthio)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and component (b) is cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spiroidiclofen, sulfoxaflo, tolfenpyrad, chlorantraniliprole, or indoxacarb.

15

16. The composition of claim 1 wherein component (a) is *N*-[1,1-dimethyl-2-(methylsulfinyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and component (b) is cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spiroidiclofen, sulfoxaflo, tolfenpyrad, chlorantraniliprole, or indoxacarb.

20

25

17. The composition of any one of claim 1 wherein component (a) is *N*-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-7-fluoro-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and component (b) cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spiroidiclofen, sulfoxaflo, tolfenpyrad, chlorantraniliprole, or indoxacarb.

30

18. The composition of claim 1 wherein component (a) is *N*-(1-methylcyclopropyl)-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and component (b) is cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr,

35

emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram, spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

5 19. The composition of claim 1 wherein component (a) is *N*-[1-(difluoromethyl)cyclopropyl]-2-(3-pyridinyl)-2*H*-indazole-4-carboxamide and component
10 (b) is cyantraniliprole, chlorpyrifos, DiPel®, acetamiprid, *Bacillus* spp, bifenthrin, buprofezin, chlorfenapyr, emamectin benzoate, fipronil, flonicamid, flupyradifurone, imidacloprid, methomyl, methoxyfenozide, novaluron, permethrin, spinetoram,
10 spirotetramat, spirodiclofen, sulfoxaflor, tolfenpyrad, chlorantraniliprole, or indoxacarb.

 20. The composition of any one of claims 1-19 wherein the weight ratio of the
15 compound of Formula **I** or the compound of Formula **II** of component (b), is from 1:10 to 50:1.

 21. The composition of any one of claims 1-19 wherein the weight ratio of the
15 compound of Formula **I** or the compound of Formula **II** to component (b), is from 3000:1 to 1:5.

20 22. The composition of any one of claims 1-19 wherein the weight ratio of the
20 compound of Formula **I** or the compound of Formula **II** to component (b), is from 100:1 to 5:1.

25 23. The composition of any one of claims 1-22 further comprising at least one
25 additional component selected from the group consisting of surfactants, solid diluents, and liquid diluents, said composition further comprising at least one additional biologically
25 active compound or agent.

 24. The composition of any one of claims 1-23 further comprising a liquid fertilizer.
30

 25. The composition of claim 24, wherein said liquid fertilizer is aqueous-based.

 26. A soil drench formulation comprising the composition of any one claims 1-25.

27. A spray composition comprising the composition of any one claims 1-25 and a propellant.

28. A bait composition, comprising: the composition of Claim any one claims 1-23,
5 one or more food materials, optionally an attractant, and optionally a humectant.

29. A trap device for controlling an invertebrate pest, comprising: the bait
composition of claim 28 and a housing adapted to receive said bait composition, wherein the
housing has at least one opening sized to permit the invertebrate pest to pass through the
10 opening so the invertebrate pest can gain access to said bait composition from a location
outside the housing, and wherein the housing is further adapted to be placed in or near a
locus of potential or known activity for the invertebrate pest.

30. A composition comprising the composition of any of claims 1-23 wherein the
15 composition is a solid composition selected from dusts, powders, granules, pellets, prills,
pastilles, tablets, and filled films.

31. The composition of claim 30 wherein the composition is water-dispersible or
water-soluble.
20

32. A liquid or dry formulation comprising the composition of any one of claims 1-
28, and 30-31 for use in a drip irrigation system, furrow during planting, handheld sprayer,
backpack sprayer, boom sprayer, ground sprayer, aerial application, unmanned aerial
vehicle, or a seed treatment.
25

33. The liquid or dry formulation of claim 32 wherein said formulation is sprayed at
an ultra-low volume.

34. A method for controlling an invertebrate pest comprising contacting the
30 invertebrate pest or its environment with a biologically effective amount of a composition or
formulation of any one of claims 1-33.

35. The method of claim 34 wherein the environment is soil or plant foliage.

36. The method of claim 34 or 35 wherein the invertebrate pest is selected from the
35 order Lepidoptera, Blattodea, Coleoptera, Dermaptera, Hemiptera, Homoptera, Acari,

Orthoptera, Diptera, Thysanoptera, Hymenoptera, Isoptera, Thysanura, Mallophaga, Siphonoptera, Araneae, and Scutigeraomorpha.

5 37. The method of any one of claims 34- 36 wherein the invertebrate pest is selected from the order Lepidoptera, Thrips, and Hemiptera.

10 38. The method of any one of claims 34- 37 wherein the invertebrate pest is selected from silverleaf whitefly, corn plant hopper, western flower thrips, potato leafhopper, cotton melon sphid, root knot nematode, fall army worm and diamondback moth.

39. The method of any one claims 34-38 wherein the composition or formulation is dispensed by a drip irrigation system, furrow during planting, handheld sprayer, backpack sprayer, boom sprayer, ground sprayer, aerial application, or an unmanned aerial vehicle.

15 40. The method of claim 39 comprising spraying said composition or formulation at an ultra-low volume.

20 41. A treated seed comprising the composition or formulation of any one of claims 1-34 in an amount of from about 0.0001 to 1 % by weight of the seed before treatment.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2020/041668

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A01N43/22 A01N43/40 A01N43/56 A01N47/06 A01N47/38
 A01N47/40 A01N51/00 A01N53/00 A01N63/00
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 2020/018362 A1 (FMC CORP [US]) 23 January 2020 (2020-01-23) page 4, line 32 - page 6, line 10 page 7, line 3 - line 4 page 7, line 8 - line 18 page 11, line 6 - line 29 page 19, line 30 - line 33 claims 8-9 ----- -/--	1-4,7,8, 10-14, 18,19, 23,24, 26-33

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search 16 September 2020	Date of mailing of the international search report 18/11/2020
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Zanobini, Alessandra
--	--

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2020/041668

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2016/144351 A1 (DU PONT [US]) 15 September 2016 (2016-09-15) page 8, line 2 - line 8 last paragraph; page 31 emtry 502 and 696; table 1 -----	1-4,7,8, 10-14, 18-41
X	WO 2015/038503 A1 (DU PONT [US]) 19 March 2015 (2015-03-19) page 7, line 22 - line 31 page 24, line 2 - line 5 page 138; tables B2-B3 page 142; tables C2-C3 page 144; tables D2-D3 -----	1-4,7,8, 10-14, 18-41

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2020/041668

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

2-4, 8, 18, 19(completely); 1, 7, 10-14, 20-41(partially)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 2-4, 8, 18, 19(completely); 1, 7, 10-14, 20-41(partially)

Compound of Formula I (a) and at least one additional invertebrate pest control agent (b) different from it, various formulation forms of such composition and a method to control invertebrate pest with such composition

2. claims: 5, 6, 9, 15-17(completely); 1, 7, 10-14, 20-41(partially)

Compounds of Formula II (a), and at least one additional invertebrate pest control agent (b) different from it, various formulation forms of such composition and a method to control invertebrate pest with such composition

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2020/041668

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2020018362	A1	23-01-2020	TW 202005533 A UY 38300 A WO 2020018362 A1

WO 2016144351	A1	15-09-2016	CN 107635984 A WO 2016144351 A1

WO 2015038503	A1	19-03-2015	AR 097633 A1 AU 2014318984 A1 BR 112016005389 A2 CA 2923814 A1 CL 2016000581 A1 CR 20160127 A EA 201690584 A1 EP 3044220 A1 EP 3424919 A1 EP 3567036 A1 ES 2707398 T3 HU E042520 T2 IL 244480 A JP 6513682 B2 JP 2016530310 A KR 20160054002 A MD 20160038 A2 NI 201600036 A PE 20160676 A1 PH 12016500494 A1 PL 3044220 T3 PT 3044220 T SG 10201801587X A SG 11201601786P A SI 3044220 T1 TR 201901556 T4 TW 201601631 A US 2017174707 A1 US 2018009827 A1 US 2019055265 A1 UY 35737 A WO 2015038503 A1 ZA 201601782 B
