

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978**PUBLICATION PARTICULARS AND ABSTRACT**

(Section 32(3)(a) – Regulation 22(1)(g) and 31)

OFFICIAL APPLICATION NO.

21	01	2005/04898
----	----	------------

LODGING DATE

22	19 DEC 2003
----	-------------

ACCEPTANCE DATE

43	06-09-2006
----	------------

INTERNATIONAL CLASSIFICATION

51	C07D
----	------

NOT FOR PUBLICATION

CLASSIFIED BY: WIPO

FULL NAMES OF APPLICANT

71	PHARMACIA CORPORATION
----	-----------------------

FULL NAMES OF INVENTORS

72	1. HANAU, CATHLEEN E 2. MERSHON, SERENA MARIE 3. GRANETO, MATTHEW J 4. MEYERS, MARVIN J 5. HEGDE, SHRIDHAR G 6. BUCHLER, INGRID P 7. WU, KUN K 8. LIU, SHUANG 9. NACRO, KASSOUM
----	---

EARLIEST PRIORITY CLAIMED

COUNTRY

33	US
----	----

NUMBER

31	60/434,962
----	------------

DATE

32	20 DEC 2002
----	-------------

TITLE OF INVENTION

54	ACYCLIC PYRAZOLE COMPOUNDS
----	----------------------------

57

ABSTRACT (NOT MORE THAN 150 WORDS)

NUMBER OF SHEETS

269

If no classification is finished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

ABSTRACT

Compounds are described which inhibit mitogen activated protein kinase-activated protein kinase-2 (MK-2). Methods of making such compounds are described, as well as a method of using them for the inhibition of MK-2, and for the prevention or treatment of a disease or disorder that is mediated by TNF α , where the method involves administering to the subject an MK-2 inhibiting compound of the present invention. Pharmaceutical compositions and kits which contain the present MK-2 inhibiting compounds are also described.

**ACYCLIC PYRAZOLE COMPOUNDS FOR THE INHIBITION OF
MITOGEN ACTIVATED PROTEIN KINASE-ACTIVATED PROTEIN
KINASE-2**

**CROSS REFERENCE TO RELATED PATENTS AND PATENT
APPLICATIONS**

5

[0001] This application is related to and claims the benefit of U.S. Provisional Patent Application Serial No. 60/434,962, filed December 20, 2002, which is incorporated by reference herein in its entirety.

BACKGROUND OF THE INVENTION

10

(1) Field of the Invention:

[0002] The present invention relates to certain cyclic and heterocyclic compounds which inhibit mitogen-activated protein kinase-activated protein kinase-2 (MAPKAP kinase-2, or MK-2), and also to methods of using such compounds to inhibit MK-2 and for the prevention and treatment of TNF α mediated diseases or disorders in subjects that are in need of such prevention and/or treatment.

15

(2) Description of the Related Art:

[0003] Mitogen-activated protein kinases (MAPKs) are members of conserved signal transduction pathways that activate transcription factors, translation factors and other target molecules in response to a variety of extracellular signals. MAPKs are activated by phosphorylation at a dual phosphorylation motif with the sequence Thr-X-Tyr by mitogen-activated protein kinase kinases (MAPKKs). In higher eukaryotes, the physiological role of MAPK signaling has been correlated with cellular events such as proliferation, oncogenesis, development and differentiation. Accordingly, the ability to regulate signal transduction via these pathways could lead to the development of treatments and preventive therapies for human diseases associated with MAPK signaling, such as inflammatory diseases, autoimmune diseases and cancer.

20

25

30

[0004] In mammalian cells, three parallel MAPK pathways have been described. The best characterized pathway leads to the activation of the extracellular-signal-regulated kinase (ERK). Less well understood are the

signal transduction pathways leading to the activation of the cJun N-terminal kinase (JNK) and the p38 MAPK. See, e.g., Davis, *Trends Biochem. Sci.* 19:470-473 (1994); Cano, *et al.*, *Trends Biochem. Sci.* 20:117-122(1995).

5 [0005] The p38 MAPK pathway is potentially activated by a wide variety of stresses and cellular insults. These stresses and cellular insults include heat shock, UV irradiation, inflammatory cytokines (such as TNF and IL-1), tunicamycin, chemotherapeutic drugs (*i.e.*, cisplatinum), anisomycin, sorbitol/hyperosmolarity, gamma irradiation, sodium arsenite, and ischaemia. See, Ono, K., *et al.*, *Cellular Signalling* 12, 1 - 13 (2000).

10 Activation of the p38 pathway is involved in (1) production of proinflammatory cytokines, such as TNF- α ; (2) induction of enzymes, such as Cox-2; (3) expression of an intracellular enzyme, such as iNOS, which plays an important role in the regulation of oxidation; (4) induction of
15 adherent proteins, such as VCAM-1 and many other inflammatory-related molecules. Furthermore, the p38 pathway functions as a regulator in the proliferation and differentiation of cells of the immune system. See, Ono, K., *et al.*, *Id.* at 7.

[0006] The p38 kinase is an upstream kinase of mitogen-activated protein kinase-activated protein kinase-2 (MAPKAP kinase-2 or MK-2).
20 (See, Freshney, N. W., *et al.*, *J. Cell*, 78:1039-1049 (1994)). MK-2 is a protein that appears to be predominantly regulated by p38 in cells. Indeed, MK-2 was the first substrate of p38 α to be identified. For example, *in vitro* phosphorylation of MK-2 by p38 α activates MK-2. The
25 substrates that MK-2 acts upon, in turn, include heat shock protein 27, lymphocyte-specific protein 1 (LAP1), cAMP response element-binding protein (CREB), ATF1, serum response factor (SRF), and tyrosine hydroxylase. The substrate of MK-2 that has been best characterized is small heat shock protein 27 (hsp27).

30 [0007] The role of the p38 pathway in inflammatory-related diseases has been studied in several animal models. The pyridinyl imidazole compound SB203580 has been shown to be a specific inhibitor of p38 *in*

vivo, and also has been shown to inhibit activation of MK-2, (See, Rouse, J., *et al*, *Cell*, 78:1027-1037 (1994); Cuenda, A., *et al*, *Biochem. J.*, 333:11-15 (1998)), as well as a MAP kinase homologue termed reactivating kinase (RK). (See, Cuenda, A., *et al*, *FEBS Lett.*, 364(2):229 - 233 (1995)). Inhibition of p38 by SB203580 can reduce mortality in a murine model of endotoxin-induced shock and inhibit the development of mouse collagen-induced arthritis and rat adjuvant arthritis. See, *e.g.*, Badger, A. M., *et al*, *J. Pharmacol Exp. Ther.*, 279:1453 - 1461 (1996). Another p38 inhibitor that has been utilized in an animal model that is believed to be more potent than SB203580 in its inhibitory effect on p38 is SB 220025. A recent animal study has demonstrated that SB 220025 caused a significant dose-dependent decrease in vascular density of granulomas in laboratory rats. (See, Jackson, J. R., *et al*, *J. Pharmacol. Exp. Ther.*, 284:687 - 692 (1998)). The results of these animal studies indicated that p38, or the components of the p38 pathway, can be useful therapeutic targets for the prevention or treatment of inflammatory disease.

[0008] Due to its integral role in the p38 signaling pathway, MK-2 has been used as a monitor for measuring the level of activation in the pathway. Because of its downstream location in the pathway, relative to p38, MK-2 has been measured as a more convenient, albeit indirect, method of assessing p38 activation. However, so far, research efforts exploring therapeutic strategies associated with the modulation of this pathway have focused mainly on the inhibition of p38 kinase.

[0009] Several compounds that inhibit the activity of p38 kinase have been described in U.S. Patent Nos. 6,046,208, 6,251,914, and 6,335,340. These compounds have been suggested to be useful for the treatment of CSBP/RK/p38 kinase mediated disease. Commercial efforts to apply p38 inhibitors have centered around two p38 inhibitors, the pyridinylimidazole inhibitor SKF 86002, and the 2,4,5 triaryl imidazole inhibitor SB203580. See, Lee, J. C., *et al*, *Immunopharmacology* 47, 185-192 (2000). Compounds possessing a similar structure have also been investigated as

potential p38 inhibitors. Indeed, p38 MSP kinase's role in various disease states has been elucidated through the use of inhibitors.

[00010] Kotlyarov, A. *et al*, in *Nat. Cell Biol.*, 1(2):94 - 97 (1999)

introduced a targeted mutation into a mouse MK-2 gene, resulting in MK-

2-deficient mice. It was shown that mice lacking MK-2 possessed increased stress resistance and survived LPS-induced endotoxic shock

better than MK-2⁺ mice. The authors concluded that MK-2 was an essential component in the inflammatory response that regulates

biosynthesis of TNF α at a post-transcriptional level. More recently,

Lehner, M.D., *et al*, in *J. Immunol.*, 168(9):4667-4673 (2002), reported that MK-2-deficient mice showed increased susceptibility to *Listeria*

monocytogenes infection, and concluded that MK-2 had an essential role in host defense against intracellular bacteria, probably via regulation of

TNF and IFN-gamma production required for activation of antibacterial effector mechanisms.

[00011] The location of MK-2 in the p38 signaling pathway at a point that is downstream of p38 offers the potential that MK-2 could act as a focal point for modulating the pathway without affecting as many


substrates as would the regulation of an enzyme further upstream in the signaling cascade -- such as p38 MAP kinase.

[00012] Accordingly, it would be useful to provide compounds and methods that could serve to modulate the activity of MK-2 -- in particular, to act as inhibitors of MK-2 activity. Such compounds and methods would be useful for the provision of benefits similar to p38 MAP kinase inhibitors, which benefits include the prevention and treatment of diseases and disorders that are mediated by TNF α . It would be even more useful to provide MK-2 inhibitors having improved potency and reduced undesirable side effects, relative to p38 inhibitors.

SUMMARY OF THE INVENTION

[00013] Briefly therefore, the present invention is directed to a novel compound having the structure of formula I:

Formula I:




wherein:

Z^2 and Z^3 are nitrogen, Z^1 , Z^4 and Z^5 are carbon, and join with Z^2 and Z^3 to form a pyrazole ring, or optionally, Z^4 and Z^5 are nitrogen, Z^1 , Z^2 and Z^3 are carbon and join with Z^4 and Z^5 to form a pyrazole ring;

5


R^a is selected from:

1)




10

2)



3)



where dashed lines indicate optional single or double bonds;

when R^a is ring M and ring M is aromatic, M^1 is carbon and is substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon and nitrogen and is unsubstituted or substituted with $(L)_nR^1$;

when ring M is partially saturated, M^1 is carbon and is mono- or di-substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon, nitrogen, oxygen and sulfur, and when M^2 , M^3 , M^4 , or M^6 is oxygen or sulfur, it is unsubstituted, and when M^2 , M^3 , M^4 or M^6 is carbon or nitrogen, it is optionally unsubstituted; or mono- or di-substituted with $(L)_nR^1$;

when R^a is ring Q and ring Q is aromatic, Q^1 is selected from carbon and nitrogen, and when Q^1 is carbon, it is substituted with $(L)_nR^1$, and when Q^1 is nitrogen, it is unsubstituted, Q^4 is selected from nitrogen and carbon, and each of Q^2 , Q^3 and Q^5 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

optionally when ring Q is aromatic, Q^1 is carbon and is substituted with $(L)_nR^1$, Q^4 is carbon, and one of Q^2 , Q^3 and Q^5 is optionally oxygen or sulfur, and the remainder of Q^2 , Q^3 and Q^5 are independently selected from nitrogen and carbon, and if carbon, are substituted with $(L)_nR^1$;

when ring Q is partially saturated, Q^1 is selected from carbon and nitrogen, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$, Q^4 is selected from carbon and nitrogen, but only one of Q^1 and Q^4 can be nitrogen, each of Q^2 , Q^3 and Q^5 is independently selected from carbon, nitrogen, oxygen and sulfur, and if oxygen or sulfur, it is unsubstituted, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$;

when R^a is structure 3, it is fully conjugated, X^2 is selected from oxygen or nitrogen substituted with $(L)_nR^1$, X^1 is carbon and is substituted with $(L)_nR^1$, and each of X^5 and X^6 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkyl- R^{11} , C_2 - C_6 alkenyl- R^{11} , C_2 - C_6 alkynyl- R^{11} , C_1 - C_6 alkyl- $(R^{11})_2$, C_2 - C_6 alkenyl- $(R^{11})_2$, CSR^{11} , amino, $CONHR^{11}$, NHR^7 , NR^8R^9 , $N(R^7)-N(R^8)(R^9)$, $C(R^{11})=N-N(R^8)(R^9)$, $N=N(R^7)$, $N(R^7)-N=C(R^8)$, $C(R^{11})=N-O(R^{10})$, $ON=C(R^{11})$, C_1 - C_6 alkyl- NHR^7 , C_1 - C_6 alkyl- NR^8R^9 , (C_1-C_4) alkyl- $N(R^7)-N(R^8)(R^9)$, (C_1-C_4) alkyl- $C(R^{11})=N-N(R^8)(R^9)$, (C_1-C_4) alkyl- $N=N(R^7)$, (C_1-C_4) alkyl- $N(R^7)-N=C(R^8)$, nitro, cyano, CO_2R^{11} , $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , COR^{11} , SR^{10} , SSR^{10} , SOR^{11} , SO_2R^{11} , C_1 - C_6 alkyl- COR^{11} , C_1 - C_6 alkyl- SR^{10} , C_1 - C_6 alkyl- SOR^{11} , C_1 - C_6 alkyl- SO_2R^{11} , halo, $Si(R^{11})_3$, halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ;

R^7 , R^8 and R^9 are each independently selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_4 alkyl- R^{11} , C_1 - C_6 alkyl- NHR^{13} , C_1 - C_6 alkyl- $NR^{13}R^{14}$, $O-R^{15}$, C_1 - C_4 alkyl- OR^{15} , CO_2R^{15} , $C(S)OR^{15}$, $C(O)SR^{15}$, $C(O)R^{17}$, $C(S)R^{17}$, $CONHR^{16}$, $C(S)NHR^{16}$, $CON(R^{16})_2$, $C(S)N(R^{16})_2$, SR^{15} , SOR^{17} , SO_2R^{17} , C_1 - C_6 alkyl- CO_2R^{15} , C_1 - C_6 alkyl- $C(S)OR^{15}$, C_1 - C_6 alkyl- $C(O)SR^{15}$, C_1 - C_6 alkyl- COR^{17} , C_1 - C_6 alkyl- $C(S)R^{17}$, C_1 - C_6 alkyl- $CONHR^{16}$, C_1 - C_6 alkyl- $C(S)NHR^{16}$, C_1 - C_6 alkyl- $CON(R^{16})_2$, C_1 - C_6 alkyl- $C(S)N(R^{16})_2$, C_1 - C_6 alkyl- SR^{15} , C_1 - C_6 alkyl- SOR^{17} , C_1 - C_6 alkyl- SO_2R^{17} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{18} ;

R^{10} is selected from -H, C_1 - C_{10} alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkyl- NHR^{13} , C_1 - C_6 alkyl- $NR^{13}R^{14}$, C_1 - C_4 alkyl- OR^{15} , CSR^{11} , CO_2R^{15} , $C(S)OR^{15}$, $C(O)SR^{15}$, COR^{17} , $C(S)R^{17}$, $CONHR^{16}$, C_1 - C_4 alkyl- R^{11} , C_1 - C_4

alkyl-NH₂R¹³, C(S)NHR¹⁶, O-R¹⁵, CON(R¹⁶)₂, C(S)N(R¹⁶)₂, SOR¹⁷, SO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, Si(R¹³)₂R¹⁷, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹¹ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₂-C₆ alkenyl, C₂-C₆ alkynyl, amino, NHR¹³, NR¹³R¹⁴, N=NR¹³, C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, O-R¹⁵, C₁-C₄ alkyl-OR¹⁵, SR¹⁵, COR¹³, CO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹² is selected from -H, OH, oxo, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R¹¹, C₂-C₁₀ alkenyl-R¹¹, C₂-C₁₀ alkynyl-R¹¹, C₁-C₁₀ alkyl-(R¹¹)₂, C₂-C₁₀ alkenyl-(R¹¹)₂, CSR¹¹, hydroxyl C₁-C₆ alkyl-R¹¹, amino C₁-C₄ alkyl-R⁷, amino, NHR⁷, NR⁸R⁹, N(R⁷)-N(R⁸)(R⁹), C(R¹¹)=N-N(R⁸)(R⁹), N=N(R⁷), N(R⁷)-N=C(R⁸), C(R¹¹)=N-O(R¹⁰), ON=C(R¹¹), C₁-C₁₀ alkyl-NHR⁷, C₁-C₁₀ alkyl-NR⁸R⁹, (C₁-C₁₀)alkyl-N(R⁷)-N(R⁸)(R⁹), (C₁-C₁₀)alkylC(R¹¹)=N-N(R⁸)(R⁹), (C₁-C₁₀)alkyl-N=N(R⁷), (C₁-C₁₀)alkyl-N(R⁷)-N=C(R⁸), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-

R^{10} , C_1 - C_{10} alkyl- OR^{10} , COR^{11} , CO_2R^{11} , SR^{10} , SSR^{10} , SOR^{11} , SO_2R^{11} , C_1 - C_{10} alkyl- COR^{11} , C_1 - C_{10} alkyl- SR^{10} , C_1 - C_{10} alkyl- SOR^{11} , C_1 - C_{10} alkyl- SO_2R^{11} , halo, $Si(R^{11})_3$, halo C_1 - C_{10} alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{18} ;

R^{13} and R^{14} are each independently selected from -H, oxo, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_4 alkyl- R^{23} , C_1 - C_6 alkyl- NHR^{19} , C_1 - C_6 alkyl- $NR^{19}R^{20}$, $O-R^{21}$, C_1 - C_4 alkyl- OR^{21} , CO_2R^{21} , COR^{21} , $C(S)OR^{21}$, $C(O)SR^{21}$, $C(O)R^{23}$, $C(S)R^{23}$, $CONHR^{22}$, $C(S)NHR^{22}$, $CON(R^{22})_2$, $C(S)N(R^{22})_2$, SR^{21} , SOR^{23} , SO_2R^{23} , C_1 - C_6 alkyl- CO_2R^{21} , C_1 - C_6 alkyl- $C(S)OR^{21}$, C_1 - C_6 alkyl- $C(O)SR^{21}$, C_1 - C_6 alkyl- COR^{23} , C_1 - C_6 alkyl- $C(S)R^{23}$, C_1 - C_6 alkyl- $CONHR^{22}$, C_1 - C_6 alkyl- $C(S)NHR^{22}$, C_1 - C_6 alkyl- $CON(R^{22})_2$, C_1 - C_6 alkyl- $C(S)N(R^{22})_2$, C_1 - C_6 alkyl- SR^{21} , C_1 - C_6 alkyl- SOR^{23} , C_1 - C_6 alkyl- SO_2R^{23} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{24} ;

R^{15} and R^{16} are independently selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkyl- NHR^{19} , C_1 - C_6 alkyl- $NR^{19}R^{20}$, C_1 - C_4 alkyl- OR^{21} , CSR^{11} , CO_2R^{22} , COR^{23} , $CONHR^{22}$, $CON(R^{22})_2$, SOR^{23} , SO_2R^{23} , C_1 - C_6 alkyl- CO_2R^{22} , C_1 - C_6 alkyl- COR^{23} , C_1 - C_6 alkyl- $CONHR^{22}$, C_1 - C_6 alkyl- $CON(R^{22})_2$, C_1 - C_6 alkyl- SR^{21} , C_1 - C_6 alkyl- SOR^{23} , C_1 - C_6 alkyl- SO_2R^{23} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl,

heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

5 R¹⁷ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkenyl-R¹⁹, C₁-C₆ alkyl-R¹⁹, C₂-C₆ alkynyl, amino, NHR¹⁹, NR¹⁹R²⁰, C₁-C₆ alkyl-NHR¹⁹, C₁-C₆ alkyl-NR¹⁹R²⁰, O-R²¹, C₁-C₄ alkyl-OR²¹, SR²¹, C₁-C₆ alkyl-CO₂R²¹, C₁-C₆ alkyl-C(S)OR²¹, C₁-C₆ alkyl-C(O)SR²¹, C₁-C₆ alkyl-COR²³, C₁-C₆ alkyl-C(S)R²³, C₁-C₆ alkyl-CONHR²², C₁-C₆ alkyl-C(S)NHR²², C₁-C₆ alkyl-CON(R²²)₂, C₁-C₆ alkyl-C(S)N(R²²)₂, C₁-C₆ alkyl-SR²¹, C₁-C₆ alkyl-SOR²³, C₁-C₆ alkyl-SO₂R²³, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

15 R¹⁸ is selected from -H, oxo, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R²³, C₂-C₁₀ alkenyl-R²³, C₂-C₁₀ alkynyl-R²³, C₁-C₁₀ alkyl-(R²³)₂, C₂-C₁₀ alkenyl-(R²³)₂, CSR²³, amino, NHR¹⁹, NR²⁰R²⁰, N(R¹⁹)-N(R²⁰)(R²⁰), C(R²³)=N-N(R²⁰)(R²⁰), N=N(R¹⁹), N(R¹⁹)-N=C(R²⁰), C(R²³)=N-O(R²¹), ON=C(R²³), C₁-C₁₀ alkyl-NHR¹⁹, C₁-C₁₀ alkyl-NR²⁰R²⁰, (C₁-C₁₀)alkyl-N(R¹⁹)-N(R²⁰)(R²⁰), (C₁-C₁₀)alkylC(R²³)=N-N(R²⁰)(R²⁰), (C₁-C₁₀)alkyl-N=N(R¹⁹), (C₁-C₁₀)alkyl-N(R¹⁹)-N=C(R²⁰), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R²¹, C₁-C₁₀ alkyl-OR²¹, COR²³, CO₂R²³, SR²¹, SSR²¹, SOR²³, SO₂R²³, C₁-C₁₀ alkyl-COR²³, C₁-C₁₀ alkyl-SR²¹, C₁-C₁₀ alkyl-SOR²³, C₁-C₁₀ alkyl-SO₂R²³, halo, Si(R²³)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁹ and R²⁰ are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R²⁹, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, O-R²⁷, C₁-C₄ alkyl-OR²⁷, CO₂R²⁷, C(S)OR²⁷, C(O)SR²⁷, C(O)R²⁹, C(S)R²⁹, CONHR²⁸, C(S)NHR²⁸, CON(R²⁸)₂, C(S)N(R²⁸)₂, SR²⁷, SOR²⁹, SO₂R²⁹, C₁-C₆ alkyl-CO₂R²⁷, C₁-C₆ alkyl-C(S)OR²⁷, C₁-C₆ alkyl-C(O)SR²⁷, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-C(S)R²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-C(S)NHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-C(S)N(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclalkyl, alkylaryl, alkylheterocyclalkyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclalkyl, alkylaryl, alkylheterocyclalkyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

R²¹ and R²² are independently selected from -H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, C₁-C₄ alkyl-OR²⁷, CSR¹¹, CO₂R²⁸, COR²⁹, CONHR²⁸, CON(R²⁸)₂, SOR²⁹, SO₂R²⁹, C₁-C₆ alkyl-CO₂R²⁸, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclalkyl, alkylaryl, alkylheterocyclalkyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclalkyl, alkylaryl, alkylheterocyclalkyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

R²³ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R²⁵, C₁-C₆ alkyl-R²⁵, C₂-C₆ alkynyl, amino, NHR²⁵, NR²⁵R²⁶, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, O-R²⁷, C₁-C₄ alkyl-OR²⁷, SR²⁷, C₁-C₆ alkyl-CO₂R²⁷, C₁-C₆ alkyl-C(S)OR²⁷, C₁-C₆ alkyl-C(O)SR²⁷, C₁-C₆ alkyl-COR²⁹,

C₁-C₆ alkyl-C(S)R²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-C(S)NHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-C(S)N(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

R²⁴ is selected from -H, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R²⁹, C₂-C₁₀ alkenyl-R²⁹, C₂-C₁₀ alkynyl-R²⁹, C₁-C₁₀ alkyl-(R²⁹)₂, C₂-C₁₀ alkenyl-(R²⁹)₂, CSR²⁹, amino, NHR²⁵, NR²⁶R²⁶, N(R²⁵)-N(R²⁶)(R²⁶), C(R²⁹)=N-N(R²⁶)(R²⁶), N=N(R²⁵), N(R²⁵)-N=C(R²⁶), C(R²⁹)=N-O(R²⁷), ON=C(R²⁹), C₁-C₁₀ alkyl-NHR²⁵, C₁-C₁₀ alkyl-NR²⁶R²⁶, (C₁-C₁₀)alkyl-N(R²⁵)-N(R²⁶)(R²⁶), (C₁-C₁₀)alkylC(R²⁹)=N-N(R²⁶)(R²⁶), (C₁-C₁₀)alkyl-N=N(R²⁵), (C₁-C₁₀)alkyl-N(R²⁵)-N=C(R²⁶), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R²⁷, C₁-C₁₀ alkyl-OR²⁷, CO₂R²⁹, COR²⁹, SR²⁷, SSR²⁷, SOR²⁹, SO₂R²⁹, C₁-C₁₀ alkyl-COR²⁹, C₁-C₁₀ alkyl-SR²⁷, C₁-C₁₀ alkyl-SOR²⁹, C₁-C₁₀ alkyl-SO₂R²⁹, halo, Si(R²⁹)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

R²⁵ and R²⁶ are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R³⁵, C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹R³², O-R³³, C₁-C₄ alkyl-OR³³, CO₂R³³, C(S)OR³³, C(O)SR³³, C(O)R³⁵, C(S)R³⁵, CONHR³⁴, C(S)NHR³⁴, CON(R³⁴)₂, C(S)N(R³⁴)₂, SR³³, SOR³⁵, SO₂R³⁵, C₁-C₆ alkyl-CO₂R³³, C₁-C₆ alkyl-C(S)OR³³, C₁-C₆ alkyl-C(O)SR³³, C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-C(S)R³⁵, C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-C(S)NHR³⁴, C₁-C₆ alkyl-CON(R³⁴)₂, C₁-C₆ alkyl-C(S)N(R³⁴)₂,

C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-SOR³⁵, C₁-C₆ alkyl-SO₂R³⁵, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R²⁷ and R²⁸ are independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹R³², C₁-C₄ alkyl-OR³³, CSR¹¹, CO₂R³⁴, COR³⁵, CONHR³⁴, CON(R³⁴)₂, SOR³⁵, SO₂R³⁵, C₁-C₆ alkyl-CO₂R³⁴, C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-CON(R³⁴)₂, C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-SOR³⁵, C₁-C₆ alkyl-SO₂R³⁵, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R²⁹ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkenyl-R³¹, C₁-C₆ alkyl-R³¹, C₂-C₆ alkynyl, amino, NHR³¹, NR³¹R³², C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹R³², O-R³³, C₁-C₄ alkyl-OR³³, SR³³, C₁-C₆ alkyl-CO₂R³³, C₁-C₆ alkyl-C(S)OR³³, C₁-C₆ alkyl-C(O)SR³³, C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-C(S)R³⁵, C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-C(S)NHR³⁴, C₁-C₆ alkyl-CON(R³⁴)₂, C₁-C₆ alkyl-C(S)N(R³⁴)₂, C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-SOR³⁵, C₁-C₆ alkyl-SO₂R³⁵, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R^{30} is selected from -H, OH, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} alkyl- R^{35} , C_2 - C_{10} alkenyl- R^{35} , C_2 - C_{10} alkynyl- R^{35} , C_1 - C_{10} alkyl- $(R^{35})_2$, C_2 - C_{10} alkenyl- $(R^{35})_2$, CSR^{35} , amino, NHR^{31} , $NR^{32}R^{32}$, $N(R^{31})-N(R^{32})(R^{32})$, $C(R^{35})=N-N(R^{32})(R^{32})$, $N=N(R^{31})$, $N(R^{31})-N=C(R^{32})$, $C(R^{35})=N-O(R^{33})$, $ON=C(R^{35})$, C_1 - C_{10} alkyl- NHR^{31} , C_1 - C_{10} alkyl- $NR^{32}R^{32}$, $(C_1$ - $C_{10})$ alkyl- $N(R^{31})-N(R^{32})(R^{32})$, $(C_1$ - $C_{10})$ alkyl- $C(R^{35})=N-N(R^{32})(R^{32})$, $(C_1$ - $C_{10})$ alkyl- $N=N(R^{31})$, $(C_1$ - $C_{10})$ alkyl- $N(R^{31})-N=C(R^{32})$, SCN , NCS , C_1 - C_{10} alkyl SCN , C_1 - C_{10} alkyl NCS , nitro, cyano, $O-R^{33}$, C_1 - C_{10} alkyl- OR^{33} , COR^{35} , SR^{33} , SSR^{33} , SOR^{35} , SO_2R^{35} , C_1 - C_{10} alkyl- COR^{35} , C_1 - C_{10} alkyl- SR^{33} , C_1 - C_{10} alkyl- SOR^{35} , C_1 - C_{10} alkyl- SO_2R^{35} , halo, $Si(R^{35})_3$, halo C_1 - C_{10} alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{36} ;

R^{31} , R^{32} , R^{33} and R^{34} are each independently selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{36} ;

R^{35} is selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, OH, alkoxy, amino, alkylamino, dialkylamino, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic

cycloalkyl are optionally substituted with one or more of the groups defined by R^{36} ;

R^{36} is selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, OH, alkoxy, amino, nitro, cyano, halo, alkylamino, dialkylamino, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heterocyclylalkyl, and heteroarylalkyl;

L is selected from $C(R^{37})_2$, O, S, NR^{37} , C=O, C=S, $C=C(R^{37})_2$, SO, SO_2 , N=NO, $CR^{37}=CR^{37}$, $CR^{37}=N$, $N=CR^{37}$, N=N, NO=N, $C=ONR^{37}$, $C=SR^{37}$, $NR^{37}C=O$, $NR^{37}C=S$, C=OO, C=OS, C=SO, C=SS, OC=O, SC=O, OC=S, SC=S, $S(O)_m-(O,S,NR^{37})$, $(O,S,NR^{37})-S(O)_m$, $C=(O,S)-C=(O,S)$, aryl, heteroaryl, heterocyclyl, cycloalkyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heterocyclylalkyl, heteroarylalkyl, or C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ,

n is an integer from 0 to 10;


m is an integer from 1 to 2; and

R^2 , R^3 , R^4 , R^5 and R^{37} are each independently absent, or selected from an R^1 group; and

R^3 and R^4 optionally join to form a ring of 5, 6, 7, or 8 atoms, where the atoms in the ring are independently selected from Z^3 , Z^4 , O, S, C=O, C=S, S=O, SO_2 , C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

[00014] The present invention is also directed to a novel compound having the structure of formula II:

Formula II.






5

wherein:

Z^2 and Z^3 are nitrogen, Z^1 , Z^4 and Z^5 are carbon, and join with Z^2 and Z^3 to form a pyrazole ring, or optionally, Z^4 and Z^5 are nitrogen, Z^1 , Z^2 and Z^3 are carbon and join with Z^4 and Z^5 to form a pyrazole ring;

10

R^a is selected from:



where dashed lines indicate optional single or double bonds;

10 when R^a is ring M and ring M is aromatic, M^1 is carbon and is substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon and nitrogen and is unsubstituted or substituted with $(L)_nR^1$;

15 when ring M is partially saturated, M^1 is carbon and is mono- or di-substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon, nitrogen, oxygen and sulfur, and when M^2 , M^3 , M^4 , or M^6 is oxygen or sulfur, it is unsubstituted, and when M^2 , M^3 , M^4 or M^6 is carbon or nitrogen, it is optionally unsubstituted; or mono- or di-substituted with $(L)_nR^1$;

when R^a is ring Q and ring Q is aromatic, Q^1 is selected from carbon and nitrogen, and when Q^1 is carbon, it is substituted with $(L)_nR^1$, and when Q^1 is nitrogen, it is unsubstituted, Q^4 is selected from nitrogen and carbon, and each of Q^2 , Q^3 and Q^5 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

optionally when ring Q is aromatic, Q^1 is carbon and is substituted with $(L)_nR^1$, Q^4 is carbon, and one of Q^2 , Q^3 and Q^5 is optionally oxygen or sulfur, and the remainder of Q^2 , Q^3 and Q^5 are independently selected from nitrogen and carbon, and if carbon, are substituted with $(L)_nR^1$;

when ring Q is partially saturated, Q^1 is selected from carbon and nitrogen, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$, Q^4 is selected from carbon and nitrogen, but only one of Q^1 and Q^4 can be nitrogen, each of Q^2 , Q^3 and Q^5 is independently selected from carbon, nitrogen, oxygen and sulfur, and if oxygen or sulfur, it is unsubstituted, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$;

when R^a is structure 3, it is fully conjugated, X^2 is selected from oxygen or nitrogen substituted with $(L)_nR^1$, X^1 is carbon and is substituted with $(L)_nR^1$, and each of X^5 and X^6 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl- R^{11} , C_1 - C_6 alkoxy- R^{11} , COR^{17} , CO_2R^7 , $CONHR^7$, $N(R^8)_2$, amino C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkyl, amino, amino C_1 - C_4 alkyl- R^7 , halo C_1 - C_4 alkyl, C_1 - C_6 alkyl-NHR⁷, carbonitrile, SR^{10} , halo, NHR⁷, NR^8R^9 , NHR⁷- C_1 - C_6 alkyl, NR^8R^9 - C_1 - C_6 alkyl, nitro, cyano, $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , C_1 - C_6 alkyl-COR¹¹, halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, or C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ;

R^7 and R^8 are each independently selected from -H, C_1 - C_6 alkyl, C_1 - C_4 alkyl- R^{11} , C_1 - C_6 alkyl- $N(R^{13})_2$, CO_2R^{16} , COR^{17} , aryl, and arylalkyl, wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

5 R^9 and R^{10} are each independently selected from -H, hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{17} , C_1 - C_6 alkyl- NH_2R^{13} , CO_2R^{16} , COR^{17} , C_1 - C_6 alkyl- CO_2R^{16} , C_1 - C_6 alkyl- $CONH-R^{16}$, C_1 - C_6 alkyl- $CON(R^{16})_2$, hydroxy C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, halo C_1 - C_4 alkyl, $Si(R^{13})_2R^{17}$, aryl, heteroaryl, heterocyclyl, arylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein
10 aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

R^{11} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxyl, halo, amino, NHR^{13} , $N(R^{13})_2$, COR^{13} , CO_2R^{17} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl,
15 heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

R^{12} is selected from -H, hydroxyl, oxo, C_1 - C_6 alkyl, hydroxyl C_1 - C_6 alkyl- R^{11} , C_1 - C_{10} alkoxy, amino, amino C_1 - C_4 alkyl- R^7 , NHR^7 , $N(R^7)_2$, C_1 - C_6 alkyl- NHR^7 , C_1 - C_6 alkyl- NHR^8R^9 , C_1 - C_6 alkyl- $N(R^8)_2$, C_1 - C_6 alkyl- R^{11} ,
20 C_1 - C_6 alkyl- $CO_2R^7R^{11}$, C_1 - C_6 alkoxy- R^{11} , nitro, $O-R^{10}$, $C=O$, COR^{11} , CO_2R^{11} , SR^{10} , SOR^{11} , SO_2R^{11} , $NHSO_2R^{11}$, C_1 - C_6 alkyl- SR^{10} , halo, halo C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, hydroxy C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkoxy, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl,
25 heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{18} ;

R^{13} and R^{14} are each independently selected from -H, oxo, C_1 - C_6 alkyl, COR^{23} , and aryl;

30 R^{15} and R^{16} are each independently selected from -H, aryl, arylalkyl, wherein aryl, arylalkyl, are optionally substituted with one or more of the groups defined by R^{24} ;

R¹⁷ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkyl-R¹⁹, NHR¹⁹, aryl, heteroarylalkyl, and heterocyclylalkyl, wherein aryl is optionally substituted with one or more of the groups defined by R²⁴;

R¹⁸ is selected from -H, oxo, hydroxyl, C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, amino, amino C₁-C₆ alkyl, N(R¹⁹)₂, C₁-C₆ alkyl-N(R¹⁹)₂, CO₂R²³, SR²¹, halo, halo C₁-C₄ alkyl, aryl, heteroaryl, and heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁹ and R²⁰ are each independently selected from -H, C₁-C₆ alkyl, heteroaryl, heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R³⁰;

R²¹ and R²² are each independently selected from -H and C₁-C₆ alkyl;

R²³ is selected from -H and C₁-C₆ alkyl;

R²⁴ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, CO₂R²⁹, halo, and halo C₁-C₄ alkyl;

R²⁹ is selected from -H, and C₁-C₆ alkyl;

R³⁰ is selected from -H, aryl, heteroaryl, heterocyclyl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R³⁶;

R³⁶ is selected from -H and halo;

L is selected from:

- 1) carboxyamino,
- 2) carboxyaminoalkyl,
- 3) alkenyl,
- 4) alkynyl,
- 5) alkyl,
- 6) hydrazoalkyl,
- 7) arylcarbamyl,
- 8) aryl,
- 9) heteroaryl,
- 10) arylalkyl,
- 11) arylalkylamino, or

12) alkylaryl,

n is an integer that is selected from 0, or 1;

R², R³ and R⁴ are each independently selected from an R¹ group;

5 and

R³ and R⁴ optionally join to form a ring of 5, 6, 7, or 8 atoms, where the atoms in the ring are independently selected from Z³, Z⁴, O, S, C=O, C=S, S=O, SO₂, C that is mono or di-substituted with an R¹ group, and N that is unsubstituted or substituted with an R¹ group.

10 **[00015]** The present invention is also directed to a novel MK-2 inhibiting compound that is listed in Table I or Table II, below.

[00016] The present invention is also directed to a novel method of inhibiting MK-2, the method comprising contacting MK-2 with at least one compound that is described in Table I or Table II, below.

5 [00017] The present invention is also directed to a novel method of preventing or treating a $TNF\alpha$ mediated disease or disorder in a subject, the method comprising administering to the subject an effective amount of an MK-2 inhibiting compound having the structure described in formula I.

10 [00018] The present invention is also directed to a novel method of preventing or treating a $TNF\alpha$ mediated disease or disorder in a subject, the method comprising administering to the subject at least one MK-2 inhibiting compound that is described in Table I or Table II, below.

[00019] The present invention is also directed to a novel therapeutic composition comprising a compound having the structure described in formula I.

15 [00020] The present invention is also directed to a novel therapeutic composition comprising at least one MK-2 inhibitory compound that is described in Table I or Table II.

20 [00021] The present invention is also directed to a novel pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one MK-2 inhibitory compound having the structure described in formula I.

[00022] The present invention is also directed to a novel comprising a dosage form that includes a therapeutically effective amount of at least one MK-2 inhibitory compound having a structure described in formula I.

25 [00023] Among the several advantages found to be achieved by the present invention, therefore, may be noted the provision of a method that could serve to modulate the activity of MK-2 -- in particular, to inhibit MK-2 activity -- and the provision of a method for the prevention and treatment of diseases and disorders that are mediated by $TNF\alpha$.

BRIEF DESCRIPTION OF THE DRAWINGS

30 [00024] Figure 1 is a graph showing paw thickness as a function of time from day 0 to day 7 for MK2 (+/+) and MK2 (-/-) mice, which have received serum injection; and

[00025] Figure 2 is a bar chart showing paw thickness at seven days after injection for normal mice, MK2 (+/+) mice receiving serum, MK2 (-/-) mice receiving serum, and MK2 (+/+) mice receiving serum and anti-TNF antibody;

5 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[00026] In accordance with the present invention, it has been discovered that certain compounds can inhibit the activity of MAPKAP kinase-2. Many of these compounds exhibit their inhibitory effect at low concentrations -- having *in vitro* MK-2 inhibition IC_{50} values of under 1.0 μ M, and with some having IC_{50} values of under about 0.1 μ M, and even as low as about 0.02 μ M. Accordingly, these compounds can be potent and effective drugs for use in the inhibition of MK-2, and of special value in subjects where such inhibition would be useful. In particular, these compounds would be useful in methods to prevent or treat diseases and disorders that are mediated by $TNF\alpha$. For example, they can be used for the prevention or treatment of arthritis.

[00027] Compounds that have a high degree of MK-2 inhibiting activity offer advantages in therapeutic uses, because therapeutic benefits can be obtained by the administration of lower amounts of the present compounds than with less active compounds. Such highly active compounds also result in fewer side effects, and in some embodiments, demonstrate a selectivity for MK-2 inhibition over the inhibition of other related kinases.

[00028] The present MK-2 inhibitory compounds inhibit the activity of the MK-2 enzyme. When it is said that a subject compound inhibits MK-2, it is meant that the MK-2 enzymatic activity is lower in the presence of the compound than it is under the same conditions in the absence of such compound. One method of expressing the potency of a compound as an MK-2 inhibitor is to measure the " IC_{50} " value of the compound. The IC_{50} value of an MK-2 inhibitor is the concentration of the compound that is required to decrease the MK-2 enzymatic activity by one-half.


Accordingly, a compound having a lower IC_{50} value is considered to be a more potent inhibitor than a compound having a higher IC_{50} value. As

used herein, compounds that inhibit MK-2 can be referred to as MK-2 inhibitors, or MK-2 inhibiting compounds or MK-2 inhibiting agents.

[00029] In practice, the selectivity of an MK-2 inhibitor varies depending upon the condition under which the test is performed and on the inhibitors being tested. However, for the purposes of this specification, the selectivity of an MK-2 inhibitor can be measured as a ratio of the *in vitro* or *in vivo* IC_{50} value for inhibition of MK-3, divided by the IC_{50} value for inhibition of MK-2 ($IC_{50\text{ MK-3}}/IC_{50\text{ MK-2}}$). As used herein, the term " IC_{50} " refers to the concentration of a compound that is required to produce 50% inhibition of MK-2 or MK-3 activity. An MK-2 selective inhibitor is any inhibitor for which the ratio of $IC_{50\text{ MK-3}}$ to $IC_{50\text{ MK-2}}$ is greater than 1. In preferred embodiments, this ratio is greater than 2, more preferably greater than 5, yet more preferably greater than 10, still more preferably greater than 50, and more preferably still, is greater than 100. Such preferred selectivity may indicate an ability to reduce the incidence of side effects incident to the administration of an MK-2 inhibitor to a subject.

[00030] Compounds that are useful in the present method include those having the structure shown in formula I:




Formula I:



wherein:

Z^2 and Z^3 are nitrogen, Z^1 , Z^4 and Z^5 are carbon, and join with Z^2 and Z^3 to form a pyrazole ring, or optionally, Z^4 and Z^5 are nitrogen, Z^1 , Z^2 and Z^3 are carbon and join with Z^4 and Z^5 to form a pyrazole ring;

R^a is selected from:



10 where dashed lines indicate optional single or double bonds;
 when R^a is ring M and ring M is aromatic, M^1 is carbon and is substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon and nitrogen and is unsubstituted or substituted with $(L)_nR^1$;

15 when ring M is partially saturated, M^1 is carbon and is mono- or di-substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon, nitrogen, oxygen and sulfur, and when M^2 , M^3 , M^4 , or M^6 is oxygen or sulfur, it is unsubstituted, and when M^2 , M^3 , M^4 or M^6 is carbon or nitrogen, it is optionally unsubstituted; or
 20 mono- or di-substituted with $(L)_nR^1$;

when R^a is ring Q and ring Q is aromatic, Q^1 is selected from carbon and nitrogen, and when Q^1 is carbon, it is substituted with $(L)_nR^1$, and when Q^1 is nitrogen, it is unsubstituted, Q^4 is selected from nitrogen and carbon, and each of Q^2 , Q^3 and Q^5 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

optionally when ring Q is aromatic, Q^1 is carbon and is substituted with $(L)_nR^1$, Q^4 is carbon, and one of Q^2 , Q^3 and Q^5 is optionally oxygen or sulfur, and the remainder of Q^2 , Q^3 and Q^5 are independently selected from nitrogen and carbon, and if carbon, are substituted with $(L)_nR^1$;

when ring Q is partially saturated, Q^1 is selected from carbon and nitrogen, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$, Q^4 is selected from carbon and nitrogen, but only one of Q^1 and Q^4 can be nitrogen, each of Q^2 , Q^3 and Q^5 is independently selected from carbon, nitrogen, oxygen and sulfur, and if oxygen or sulfur, it is unsubstituted, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$;

when R^a is structure 3, it is fully conjugated, X^2 is selected from oxygen or nitrogen substituted with $(L)_nR^1$, X^1 is carbon and is substituted with $(L)_nR^1$, and each of X^5 and X^6 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkyl- R^{11} , C_2 - C_6 alkenyl- R^{11} , C_2 - C_6 alkynyl- R^{11} , C_1 - C_6 alkyl- $(R^{11})_2$, C_2 - C_6 alkenyl- $(R^{11})_2$, CSR^{11} , amino, $CONHR^{11}$, NHR^7 , NR^8R^9 , $N(R^7)$ - $N(R^8)(R^9)$, $C(R^{11})=N-N(R^8)(R^9)$, $N=N(R^7)$, $N(R^7)-N=C(R^8)$, $C(R^{11})=N-O(R^{10})$, $ON=C(R^{11})$, C_1 - C_6 alkyl- NHR^7 , C_1 - C_6 alkyl- NR^8R^9 , (C_1-C_4) alkyl- $N(R^7)-N(R^8)(R^9)$, (C_1-C_4) alkyl- $C(R^{11})=N-N(R^8)(R^9)$, (C_1-C_4) alkyl- $N=N(R^7)$, (C_1-C_4) alkyl- $N(R^7)-N=C(R^8)$, nitro, cyano, CO_2R^{11} , $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , COR^{11} , SR^{10} , SSR^{10} , SOR^{11} , SO_2R^{11} , C_1 - C_6 alkyl- COR^{11} , C_1 - C_6 alkyl- SR^{10} , C_1 - C_6 alkyl- SOR^{11} , C_1 - C_6 alkyl- SO_2R^{11} , halo, $Si(R^{11})_3$, halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic

cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹²;

5 R⁷, R⁸ and R⁹ are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R¹¹, C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, O-R¹⁵, C₁-C₄ alkyl-OR¹⁵, CO₂R¹⁵, C(S)OR¹⁵, C(O)SR¹⁵, C(O)R¹⁷, C(S)R¹⁷, CONHR¹⁶, C(S)NHR¹⁶, CON(R¹⁶)₂, C(S)N(R¹⁶)₂, SR¹⁵, SOR¹⁷, SO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

15 R¹⁰ is selected from -H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, C₁-C₄ alkyl-OR¹⁵, CSR¹¹, CO₂R¹⁵, C(S)OR¹⁵, C(O)SR¹⁵, COR¹⁷, C(S)R¹⁷, CONHR¹⁶, C₁-C₄ alkyl-R¹¹, C₁-C₄ alkyl-NH₂R¹³, C(S)NHR¹⁶, O-R¹⁵, CON(R¹⁶)₂, C(S)N(R¹⁶)₂, SOR¹⁷, SO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, Si(R¹³)₂R¹⁷, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R^{11} is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₂-C₆ alkenyl, C₂-C₆ alkynyl, amino, NHR¹³, NR¹³R¹⁴, N=NR¹³, C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, O-R¹⁵, C₁-C₄ alkyl-OR¹⁵, SR¹⁵, COR¹³, CO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R^{12} is selected from -H, OH, oxo, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R¹¹, C₂-C₁₀ alkenyl-R¹¹, C₂-C₁₀ alkynyl-R¹¹, C₁-C₁₀ alkyl-(R¹¹)₂, C₂-C₁₀ alkenyl-(R¹¹)₂, CSR¹¹, hydroxyl C₁-C₆ alkyl-R¹¹, amino C₁-C₄ alkyl-R⁷, amino, NHR⁷, NR⁸R⁹, N(R⁷)-N(R⁸)(R⁹), C(R¹¹)=N-N(R⁸)(R⁹), N=N(R⁷), N(R⁷)-N=C(R⁸), C(R¹¹)=N-O(R¹⁰), ON=C(R¹¹), C₁-C₁₀ alkyl-NHR⁷, C₁-C₁₀ alkyl-NR⁸R⁹, (C₁-C₁₀)alkyl-N(R⁷)-N(R⁸)(R⁹), (C₁-C₁₀)alkylC(R¹¹)=N-N(R⁸)(R⁹), (C₁-C₁₀)alkyl-N=N(R⁷), (C₁-C₁₀)alkyl-N(R⁷)-N=C(R⁸), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R¹⁰, C₁-C₁₀ alkyl-OR¹⁰, COR¹¹, CO₂R¹¹, SR¹⁰, SSR¹⁰, SOR¹¹, SO₂R¹¹, C₁-C₁₀ alkyl-COR¹¹, C₁-C₁₀ alkyl-SR¹⁰, C₁-C₁₀ alkyl-SOR¹¹, C₁-C₁₀ alkyl-SO₂R¹¹, halo, Si(R¹¹)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R^{13} and R^{14} are each independently selected from -H, oxo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R²³, C₁-C₆ alkyl-NHR¹⁹, C₁-

C_6 alkyl- $NR^{19}R^{20}$, $O-R^{21}$, C_1-C_4 alkyl- OR^{21} , CO_2R^{21} , COR^{21} , $C(S)OR^{21}$,
 $C(O)SR^{21}$, $C(O)R^{23}$, $C(S)R^{23}$, $CONHR^{22}$, $C(S)NHR^{22}$, $CON(R^{22})_2$,
 $C(S)N(R^{22})_2$, SR^{21} , SOR^{23} , SO_2R^{23} , C_1-C_6 alkyl- CO_2R^{21} , C_1-C_6 alkyl-
 $C(S)OR^{21}$, C_1-C_6 alkyl- $C(O)SR^{21}$, C_1-C_6 alkyl- COR^{23} , C_1-C_6 alkyl- $C(S)R^{23}$,
 C_1-C_6 alkyl- $CONHR^{22}$, C_1-C_6 alkyl- $C(S)NHR^{22}$, C_1-C_6 alkyl- $CON(R^{22})_2$, C_1-
 C_6 alkyl- $C(S)N(R^{22})_2$, C_1-C_6 alkyl- SR^{21} , C_1-C_6 alkyl- SOR^{23} , C_1-C_6 alkyl-
 SO_2R^{23} , halo C_1-C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl,
 alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl,
 heterocyclylalkyl, and C_1-C_{10} mono- and bicyclic cycloalkyl, wherein aryl,
 heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl,
 arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1-C_{10} mono- and bicyclic
 cycloalkyl are optionally substituted with one or more of the groups defined
 by R^{24} ;

R^{15} and R^{16} are independently selected from $-H$, C_1-C_6 alkyl, C_2-C_6
 alkenyl, C_2-C_6 alkynyl, C_1-C_6 alkyl- NHR^{19} , C_1-C_6 alkyl- $NR^{19}R^{20}$, C_1-C_4
 alkyl- OR^{21} , CSR^{11} , CO_2R^{22} , COR^{23} , $CONHR^{22}$, $CON(R^{22})_2$, SOR^{23} ,
 SO_2R^{23} , C_1-C_6 alkyl- CO_2R^{22} , C_1-C_6 alkyl- COR^{23} , C_1-C_6 alkyl- $CONHR^{22}$, C_1-
 C_6 alkyl- $CON(R^{22})_2$, C_1-C_6 alkyl- SR^{21} , C_1-C_6 alkyl- SOR^{23} , C_1-C_6 alkyl-
 SO_2R^{23} , halo C_1-C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl,
 alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl,
 heterocyclylalkyl, and C_1-C_{10} mono- and bicyclic cycloalkyl, wherein aryl,
 heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl,
 arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1-C_{10} mono- and bicyclic
 cycloalkyl are optionally substituted with one or more of the groups defined
 by R^{24} ;

R^{17} is selected from $-H$, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkenyl-
 R^{19} , C_1-C_6 alkyl- R^{19} , C_2-C_6 alkynyl, amino, NHR^{19} , $NR^{19}R^{20}$, C_1-C_6 alkyl-
 NHR^{19} , C_1-C_6 alkyl- $NR^{19}R^{20}$, $O-R^{21}$, C_1-C_4 alkyl- OR^{21} , SR^{21} , C_1-C_6 alkyl-
 CO_2R^{21} , C_1-C_6 alkyl- $C(S)OR^{21}$, C_1-C_6 alkyl- $C(O)SR^{21}$, C_1-C_6 alkyl- COR^{23} ,
 C_1-C_6 alkyl- $C(S)R^{23}$, C_1-C_6 alkyl- $CONHR^{22}$, C_1-C_6 alkyl- $C(S)NHR^{22}$, C_1-C_6
 alkyl- $CON(R^{22})_2$, C_1-C_6 alkyl- $C(S)N(R^{22})_2$, C_1-C_6 alkyl- SR^{21} , C_1-C_6 alkyl-
 SOR^{23} , C_1-C_6 alkyl- SO_2R^{23} , halo C_1-C_4 alkyl, aryl, heteroaryl, heterocyclyl,

alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁸ is selected from -H, oxo, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R²³, C₂-C₁₀ alkenyl-R²³, C₂-C₁₀ alkynyl-R²³, C₁-C₁₀ alkyl-(R²³)₂, C₂-C₁₀ alkenyl-(R²³)₂, CSR²³, amino, NHR¹⁹, NR²⁰R²⁰, N(R¹⁹)-N(R²⁰)(R²⁰), C(R²³)=N-N(R²⁰)(R²⁰), N=N(R¹⁹), N(R¹⁹)-N=C(R²⁰), C(R²³)=N-O(R²¹), ON=C(R²³), C₁-C₁₀ alkyl-NHR¹⁹, C₁-C₁₀ alkyl-NR²⁰R²⁰, (C₁-C₁₀)alkyl-N(R¹⁹)-N(R²⁰)(R²⁰), (C₁-C₁₀)alkylC(R²³)=N-N(R²⁰)(R²⁰), (C₁-C₁₀)alkyl-N=N(R¹⁹), (C₁-C₁₀)alkyl-N(R¹⁹)-N=C(R²⁰), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R²¹, C₁-C₁₀ alkyl-OR²¹, COR²³, CO₂R²³, SR²¹, SSR²¹, SOR²³, SO₂R²³, C₁-C₁₀ alkyl-COR²³, C₁-C₁₀ alkyl-SR²¹, C₁-C₁₀ alkyl-SOR²³, C₁-C₁₀ alkyl-SO₂R²³, halo, Si(R²³)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁹ and R²⁰ are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R²⁹, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, O-R²⁷, C₁-C₄ alkyl-OR²⁷, CO₂R²⁷, C(S)OR²⁷, C(O)SR²⁷, C(O)R²⁹, C(S)R²⁹, CONHR²⁸, C(S)NHR²⁸, CON(R²⁸)₂, C(S)N(R²⁸)₂, SR²⁷, SOR²⁹, SO₂R²⁹, C₁-C₆ alkyl-CO₂R²⁷, C₁-C₆ alkyl-C(S)OR²⁷, C₁-C₆ alkyl-C(O)SR²⁷, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-C(S)R²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-C(S)NHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-C(S)N(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic

cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

5 R²¹ and R²² are independently selected from -H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, C₁-C₄ alkyl-OR²⁷, CSR¹¹, CO₂R²⁸, COR²⁹, CONHR²⁸, CON(R²⁸)₂, SOR²⁹, SO₂R²⁹, C₁-C₆ alkyl-CO₂R²⁸, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

10 R²³ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkenyl-R²⁵, C₁-C₆ alkyl-R²⁵, C₂-C₆ alkynyl, amino, NHR²⁵, NR²⁵R²⁶, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, O-R²⁷, C₁-C₄ alkyl-OR²⁷, SR²⁷, C₁-C₆ alkyl-CO₂R²⁷, C₁-C₆ alkyl-C(S)OR²⁷, C₁-C₆ alkyl-C(O)SR²⁷, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-C(S)R²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-C(S)NHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-C(S)N(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

25 R²⁴ is selected from -H, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R²⁹, C₂-C₁₀ alkenyl-R²⁹, C₂-C₁₀ alkynyl-R²⁹, C₁-C₁₀ alkyl-(R²⁹)₂, C₂-C₁₀ alkenyl-(R²⁹)₂, CSR²⁹, amino, NHR²⁵, NR²⁶R²⁶, N(R²⁵)-

$N(R^{26})(R^{26})$, $C(R^{29})=N-N(R^{26})(R^{26})$, $N=N(R^{25})$, $N(R^{25})-N=C(R^{26})$, $C(R^{29})=N-O(R^{27})$, $ON=C(R^{29})$, C_1-C_{10} alkyl-NHR²⁵, C_1-C_{10} alkyl-NR²⁶R²⁶, (C_1-C_{10}) alkyl-N(R²⁵)-N(R²⁶)(R²⁶), (C_1-C_{10}) alkylC(R²⁹)=N-N(R²⁶)(R²⁶), (C_1-C_{10}) alkyl-N=N(R²⁵), (C_1-C_{10}) alkyl-N(R²⁵)-N=C(R²⁶), SCN, NCS, C_1-C_{10} alkyl SCN, C_1-C_{10} alkyl NCS, nitro, cyano, O-R²⁷, C_1-C_{10} alkyl-OR²⁷, CO₂R²⁹, COR²⁹, SR²⁷, SSR²⁷, SOR²⁹, SO₂R²⁹, C_1-C_{10} alkyl-COR²⁹, C_1-C_{10} alkyl-SR²⁷, C_1-C_{10} alkyl-SOR²⁹, C_1-C_{10} alkyl-SO₂R²⁹, halo, Si(R²⁹)₃, halo C_1-C_{10} alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C_1-C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C_1-C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

R²⁵ and R²⁶ are each independently selected from -H, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_4 alkyl-R³⁵, C_1-C_6 alkyl-NHR³¹, C_1-C_6 alkyl-NR³¹R³², O-R³³, C_1-C_4 alkyl-OR³³, CO₂R³³, C(S)OR³³, C(O)SR³³, C(O)R³⁵, C(S)R³⁵, CONHR³⁴, C(S)NHR³⁴, CON(R³⁴)₂, C(S)N(R³⁴)₂, SR³³, SOR³⁵, SO₂R³⁵, C_1-C_6 alkyl-CO₂R³³, C_1-C_6 alkyl-C(S)OR³³, C_1-C_6 alkyl-C(O)SR³³, C_1-C_6 alkyl-COR³⁵, C_1-C_6 alkyl-C(S)R³⁵, C_1-C_6 alkyl-CONHR³⁴, C_1-C_6 alkyl-C(S)NHR³⁴, C_1-C_6 alkyl-CON(R³⁴)₂, C_1-C_6 alkyl-C(S)N(R³⁴)₂, C_1-C_6 alkyl-SR³³, C_1-C_6 alkyl-SOR³⁵, C_1-C_6 alkyl-SO₂R³⁵, halo C_1-C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C_1-C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C_1-C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R²⁷ and R²⁸ are independently selected from -H, C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_6 alkyl-NHR³¹, C_1-C_6 alkyl-NR³¹R³², C_1-C_4 alkyl-OR³³, CSR¹¹, CO₂R³⁴, COR³⁵, CONHR³⁴, CON(R³⁴)₂, SOR³⁵, SO₂R³⁵, C_1-C_6 alkyl-CO₂R³⁴, C_1-C_6 alkyl-COR³⁵, C_1-C_6 alkyl-CONHR³⁴, C_1-C_6 alkyl-CON(R³⁴)₂, C_1-C_6 alkyl-SR³³, C_1-C_6 alkyl-SOR³⁵, C_1-C_6 alkyl-

SO₂R³⁵, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R²⁹ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkenyl-R³¹, C₁-C₆ alkyl-R³¹, C₂-C₆ alkynyl, amino, NHR³¹, NR³¹R³², C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹R³², O-R³³, C₁-C₄ alkyl-OR³³, SR³³, C₁-C₆ alkyl-CO₂R³³, C₁-C₆ alkyl-C(S)OR³³, C₁-C₆ alkyl-C(O)SR³³, C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-C(S)R³⁵, C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-C(S)NHR³⁴, C₁-C₆ alkyl-CON(R³⁴)₂, C₁-C₆ alkyl-C(S)N(R³⁴)₂, C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-SOR³⁵, C₁-C₆ alkyl-SO₂R³⁵, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R³⁰ is selected from -H, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R³⁵, C₂-C₁₀ alkenyl-R³⁵, C₂-C₁₀ alkynyl-R³⁵, C₁-C₁₀ alkyl-(R³⁵)₂, C₂-C₁₀ alkenyl-(R³⁵)₂, CSR³⁵, amino, NHR³¹, NR³²R³², N(R³¹)-N(R³²)(R³²), C(R³⁵)=N-N(R³²)(R³²), N=N(R³¹), N(R³¹)-N=C(R³²), C(R³⁵)=N-O(R³³), ON=C(R³⁵), C₁-C₁₀ alkyl-NHR³¹, C₁-C₁₀ alkyl-NR³²R³², (C₁-C₁₀)alkyl-N(R³¹)-N(R³²)(R³²), (C₁-C₁₀)alkylC(R³⁵)=N-N(R³²)(R³²), (C₁-C₁₀)alkyl-N=N(R³¹), (C₁-C₁₀)alkyl-N(R³¹)-N=C(R³²), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R³³, C₁-C₁₀ alkyl-OR³³, COR³⁵, SR³³, SSR³³, SOR³⁵, SO₂R³⁵, C₁-C₁₀ alkyl-COR³⁵, C₁-C₁₀ alkyl-SR³³, C₁-C₁₀ alkyl-SOR³⁵, C₁-C₁₀ alkyl-SO₂R³⁵, halo, Si(R³⁵)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀

mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

5 R³¹, R³², R³³ and R³⁴ and each independently selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

15 R³⁵ is selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, OH, alkoxy, amino, alkylamino, dialkylamino, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

25 R³⁶ is selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, OH, alkoxy, amino, nitro, cyano, halo, alkylamino, dialkylamino, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heterocyclylalkyl, and heteroarylalkyl;

30 L is selected from C(R³⁷)₂, O, S, NR³⁷, C=O, C=S, C=C(R³⁷)₂, SO, SO₂, N=NO, CR³⁷=CR³⁷, CR³⁷=N, N=CR³⁷, N=N, NO=N, C=ONR³⁷, C=SR³⁷, NR³⁷C=O, NR³⁷C=S, C=OO, C=OS, C=SO, C=SS, OC=O, SC=O, OC=S, SC=S, S(O)_m-(O,S,NR³⁷), (O,S,NR³⁷-S(O)_m, C=(O,S)-C=(O,S), aryl, heteroaryl, heterocyclyl, cycloalkyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heterocyclylalkyl, heteroarylalkyl, or C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and

bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹²,

n is an integer from 0 to 10;

m is an integer from 1 to 2; and

- 5 R², R³, R⁴, R⁵ and R³⁷ are each independently absent, or selected from an R¹ group; and

R³ and R⁴ optionally join to form a ring of 5, 6, 7, or 8 atoms, where the atoms in the ring are independently selected from Z³, Z⁴, O, S, C=O,

C=S, S=O, SO₂, C that is mono or di-substituted with an R¹ group, and N that is unsubstituted or substituted with an R¹ group.

[00031] The "M" ring and the "Q" ring of the structure of formula I can have any number of R¹-L_n- substituent groups, ranging from zero to one or more per ring atom, and such substituent groups can be located on any atom of the ring having a valence suitable for the addition of a substituent group(s). Each such substituent group can have any number of R¹ groups per L group, ranging from zero to 5. A preferred structure is the presence of either 0 or 1 R¹-L_n- substituent groups on the ring. It is also preferred that the R¹-L_n- substituent group is attached to the ring at the M¹ or the Q¹ location, respectively.

[00032] The meaning of any substituent at any one occurrence in Formula I, or any other general chemical formula herein, is independent of its meaning, or any other substituent's meaning, at any other occurrence, unless specified otherwise.

[00033] The term "alkyl" is used, either alone or within other terms such as "haloalkyl" and "alkylsulfonyl"; it embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about five carbon atoms. The number of carbon atoms can also be expressed as "C₁-C₅", for example. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, hexyl, octyl and the like. The term "alkenyl" refers to an unsaturated, acyclic hydrocarbon radical, linear or branched, in so much as it contains at least one double bond. Unless otherwise noted, such radicals preferably contain from 2 to about 6 carbon atoms, preferably from 2 to about 4 carbon atoms, more preferably from 2 to about 3 carbon atoms. The alkenyl radicals may be optionally substituted with groups as defined below. Examples of suitable alkenyl radicals include propenyl, 2-chloropropenyl, buten-1-yl, isobutenyl, penten-1-yl, 2-methylbuten-1-yl, 3-methylbuten-1-yl, hexen-1-yl, 3-

hydroxyhexen-1-yl, hepten-1-yl, octen-1-yl, and the like. The term "alkynyl" refers to an unsaturated, acyclic hydrocarbon radical, linear or branched, in so much as it contains one or more triple bonds, such radicals preferably containing 2 to about 6 carbon atoms, more preferably from 2 to about 3 carbon atoms. The alkynyl radicals may be optionally substituted with groups as described below. Examples of suitable alkynyl radicals include ethynyl, propynyl, hydroxypropynyl, butyn-1-yl, butyn-2-yl, pentyn-1-yl, pentyn-2-yl, 4-methoxypentyn-2-yl, 3-methylbutyn-1-yl, hexyl-1-yl, hexyn-2-yl, hexyn-3-yl, 3,3-dimethylbutyn-1-yl radicals, and the like.

The term "oxo" means a single double-bonded oxygen. The terms "hydrido", "-H", or "hydrogen", denote a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical, or two hydrido radicals may be attached to a carbon atom to form a methylene ($-\text{CH}_2-$) radical. The term "halo" means halogens


such as fluorine, chlorine, and bromine or iodine atoms. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl, and polyhaloalkyl radicals. A

monohaloalkyl radical, for one example, may have a bromo, chloro, or a fluoro atom within the radical. Dihalo radicals may have two or more of the same halo atoms or a combination of different halo radicals and polyhaloalkyl radicals may have more than two of the same halo atoms or a combination of different halo radicals. Likewise, the term "halo", when it is appended to alkenyl, alkynyl, alkoxy, aryl, cycloalkyl, heteroalkyl,

heteroaryl, and the like, includes radicals having mono-, di-, or tri-, halo substitution on one or more of the atoms of the radical. The term


"hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl radicals. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms, such as methoxy radical. The term "alkoxyalkyl" also embraces alkyl radicals having two or more alkoxy

radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and diaikoxyalkyl radicals. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro, or bromo, to provide "haloalkoxy" or "haloalkoxyalkyl" radicals. Examples of "alkoxy" radicals include methoxy, butoxy, and trifluoromethoxy. Terms such as "alkoxy(halo)alkyl", indicate a molecule having a terminal alkoxy that is bound to an alkyl, which is bonded to the parent molecule, while the alkyl also has a substituent halo group in a non-terminal location. In other words, both the alkoxy and the halo group are substituents of the alkyl chain. The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two, or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane, and biphenyl. The term "heterocyclyl" means a saturated or unsaturated mono- or multi-ring carbocycle wherein one or more carbon atoms is replaced by N, S, P, or O. This includes, for example, structures such as:



where Z, Z¹, Z², or Z³ is C, S, P, O, or N, with the proviso that one of Z, Z¹, Z², or Z³ is other than carbon, but is not O or S when attached to another Z atom by a double bond or when attached to another O or S atom. Furthermore, the optional substituents are understood to be attached to Z, Z¹, Z², or Z³ only when each is C. The term "heterocycle" also includes fully saturated ring structures, such as piperazinyl, dioxanyl, tetrahydrofuranyl, oxiranyl, aziridinyl, morpholinyl, pyrrolidinyl, piperidinyl, thiazolidinyl, and others. The term "heteroaryl" embraces unsaturated heterocyclic radicals. Examples of unsaturated heterocyclic radicals, also

termed "heteroaryl" radicals include thienyl, pyrrolyl, furyl, pyridyl, pyrimidyl, pyrazinyl, pyrazolyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, pyranyl, and tetrazolyl. The term also embraces radicals where heterocyclic radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like. The terms aryl or heteroaryl, as appropriate, include the following structures:



where:

when $n=1$, $m=1$ and A_1 - A_8 are each CR^x or N, A_9 and A_{10} are carbon;

when $n=0$, or 1, and $m=0$, or 1, one of A_2 - A_4 and/or A_5 - A_7 is optionally S, O, or NR^x , and other ring members are CR^x or N, with the proviso that oxygen cannot be adjacent to sulfur in a ring. A_9 and A_{10} are carbon;

when n is greater than or equal to 0, and m is greater than or equal to 0, 1 or more sets of 2 or more adjacent atoms A_1 - A_{10} are sp^3 O, S, NR^x , CR^xR^y , or $C=(O \text{ or } S)$, with the proviso that oxygen and sulfur cannot be adjacent. The remaining A_1 - A_8 are CR^x or N, and A_9 and A_{10} are carbon;

when n is greater than or equal to 0, and m greater than or equal to 0, atoms separated by 2 atoms (*i.e.*, A_1 and A_4) are sp^3 O, S, NR^x , CR^xR^y , and remaining A_1 - A_8 are independently CR^x or N, and A_9 and A_{10} are carbon.

[00034] The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals $-SO_2-$.


"Alkylsulfonyl", embraces alkyl radicals attached to a sulfonyl radical,

where alkyl is defined as above. The term "arylsulfonyl" embraces sulfonyl radicals substituted with an aryl radical. The terms "sulfamyl" or "sulfonamidyl", whether alone or used with terms such as "N-alkylsulfamyl", "N-arylsulfamyl", "N,N-dialkylsulfamyl" and "N-alkyl-N-arylsulfamyl", denotes a sulfonyl radical substituted with an amine radical, forming a sulfonamide ($-\text{SO}_2\text{-NH}_2$), which may also be termed an "aminosulfonyl". The terms "N-alkylsulfamyl" and "N,N-dialkylsulfamyl" denote sulfamyl radicals substituted, respectively, with one alkyl radical, a cycloalkyl ring, or two alkyl radicals. The terms "N-arylsulfamyl" and "N-alkyl-N-arylsulfamyl" denote sulfamyl radicals substituted, respectively, with one aryl radical, and one alkyl and one aryl radical. The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes $-\text{CO}_2\text{-H}$. The term "carboxyalkyl" embraces radicals having a carboxyl radical as defined above, attached to an alkyl radical. The term "carbonyl", whether used alone or with other terms, such as "alkylcarbonyl", denotes $-(\text{C}=\text{O})-$. The term "alkylcarbonyl" embraces radicals having a carbonyl radical substituted with an alkyl radical. An example of an "alkylcarbonyl" radical is $\text{CH}_3-(\text{CO})-$. The term "alkylcarbonylalkyl" denotes an alkyl radical substituted with an "alkylcarbonyl" radical. The term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl ($\text{C}=\text{O}$) radical. Examples of such "alkoxycarbonyl" radicals include $(\text{CH}_3)_3\text{C-O-C}=\text{O}-$ and $-(\text{O}=\text{C})\text{-OCH}_3$. The term "alkoxycarbonylalkyl" embraces radicals having "alkoxycarbonyl", as defined above substituted to an alkyl radical. Examples of such "alkoxycarbonylalkyl" radicals include $(\text{CH}_3)_3\text{C-OC(=O)-(CH}_2)_2-$ and $(\text{CH}_2)_2(-\text{O})\text{COCH}_3$. The terms "amido", or "carbamyl", when used alone or with other terms such as "amidoalkyl", "N-monoalkylamido", "N-monoarylamido", "N,N-dialkylamido", "N-alkyl-N-arylamido", "N-alkyl-N-hydroxyamido" and "N-alkyl-N-hydroxyamidoalkyl", embraces a carbonyl radical substituted with an amino radical. The terms "N-alkylamido" and "N,N-dialkylamido" denote amido groups which have been substituted with

one alkylradical and with two alkyl radicals, respectively. The terms "N-monoarylamido" and "N-alkyl-N-arylamido" denote amido radicals substituted, respectively, with one aryl radical, and one alkyl and one aryl radical. The term "N-alkyl-N-hydroxyamido" embraces amido radicals substituted with a hydroxyl radical and with an alkyl radical. The term "N-alkyl-N-hydroxyamidoalkyl" embraces alkylradicals substituted with an N-alkyl-N-hydroxyamido radical. The term "amidoalkyl" embraces alkyl radicals substituted with amido radicals. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. The term "alkylaminoalkyl" embraces aminoalkyl radicals having the nitrogen atom substituted with an alkyl radical. The term "amidino" denotes an $-C(-NH)-NH_2$ radical. The term "cyanoamidin" denotes an $-C(-N-CN)-NH_2$ radical. The term "heterocycloalkyl" embraces heterocyclic-substituted alkyl radicals such as pyridylmethyl and thienylmethyl. The terms "aralkyl", or "arylalkyl" embrace aryl-substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenethyl, and diphenethyl. The terms benzyl and phenylmethyl are interchangeable. The term "cycloalkyl" embraces radicals having three to ten carbon atoms, such as cyclopropyl cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. The term "cycloalkenyl" embraces unsaturated radicals having three to ten carbon atoms, such as cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and cycloheptenyl. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent sulfur atom. An example of "alkylthio" is methylthio, (CH_3-S-) . The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent $-S(-O)-$ atom. The terms "N-alkylamino" and "N, N-dialkylamino" denote amino groups which have been substituted with one alkyl radical and with two alkyl radicals, respectively. The term "acyl", whether used alone, or within a term such as "acylamino", denotes a radical provided by the residue after removal of hydroxyl from an organic acid. The term "acylamino"

embraces an amino radical substituted with an acyl group. An examples of an "acylamino" radical is acetylamino ($\text{CH}_3\text{-C(=O)-NH-}$).

[00035] In the naming of substituent groups for general chemical structures, the naming of the chemical components of the group is typically from the terminal group-toward the parent compound unless otherwise noted, as discussed below. In other words, the outermost chemical structure is named first, followed by the next structure in line, followed by the next, etc. until the structure that is connected to the parent structure is named. For example, a substituent group having a structure such as:




may be referred to generally as a "haloarylalkylaminocarboxylalkyl". An example of one such group would be fluorophenylmethylcarbamylylpentyl. The bonds having wavy lines through them represent the parent structure to which the alkyl is attached.

[00036] Substituent groups may also be named by reference to one or more "R" groups. The structure shown above would be included in a description, such as, " $\text{-C}_1\text{-C}_6\text{-alkyl-COR}^u$ ", where R^u is defined to include $\text{-NH-C}_1\text{-C}_4\text{-alkylaryl-R}^v$, and where R^v is defined to include halo. In this scheme, atoms having an "R" group are shown with the "R" group being the terminal group (*i.e.*, furthest from the parent). In a term such as " $\text{C(R}^x)_2$ ", it should be understood that the two R^x groups can be the same, or they can be different if R^x is defined as having more than one possible identity.

[00037] The present invention also comprises MK-2 inhibiting compounds having the structure shown in formula II:

Formula II.




wherein:

- 5 Z^2 and Z^3 are nitrogen, Z^1 , Z^4 and Z^5 are carbon, and join with Z^2 and Z^3 to form a pyrazole ring, or optionally, Z^4 and Z^5 are nitrogen, Z^1 , Z^2 and Z^3 are carbon and join with Z^4 and Z^5 to form a pyrazole ring;


 R^a is selected from:

1)




10

2)



3)



where dashed lines indicate optional single or double bonds;

when R^a is ring M and ring M is aromatic, M^1 is carbon and is substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon and nitrogen and is unsubstituted or substituted with $(L)_nR^1$;

5 when ring M is partially saturated, M^1 is carbon and is mono- or di-substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon, nitrogen, oxygen and sulfur, and when M^2 , M^3 , M^4 , or M^6 is oxygen or sulfur, it is unsubstituted, and when M^2 , M^3 , M^4 or M^6 is carbon or nitrogen, it is optionally unsubstituted; or
10 mono- or di-substituted with $(L)_nR^1$;

 when R^a is ring Q and ring Q is aromatic, Q^1 is selected from carbon and nitrogen, and when Q^1 is carbon, it is substituted with $(L)_nR^1$, and when Q^1 is nitrogen, it is unsubstituted, Q^4 is selected from nitrogen and carbon, and each of Q^2 , Q^3 and Q^5 is independently selected from
15 nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

 optionally when ring Q is aromatic, Q^1 is carbon and is substituted with $(L)_nR^1$, Q^4 is carbon, and one of Q^2 , Q^3 and Q^5 is optionally oxygen or sulfur, and the remainder of Q^2 , Q^3 and Q^5 are independently selected from nitrogen and carbon, and if carbon, are substituted with $(L)_nR^1$;

20 when ring Q is partially saturated, Q^1 is selected from carbon and nitrogen, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$, Q^4 is selected from carbon and nitrogen, but only one of Q^1 and Q^4 can be nitrogen, each of Q^2 , Q^3 and Q^5 is independently selected from carbon, nitrogen, oxygen
25 and sulfur, and if oxygen or sulfur, it is unsubstituted, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$;

 when R^a is structure 3, it is fully conjugated, X^2 is selected from oxygen or nitrogen substituted with $(L)_nR^1$, X^1 is carbon and is substituted with $(L)_nR^1$, and each of X^5 and X^6 is independently selected from nitrogen
30 and carbon, and if carbon, it is substituted with $(L)_nR^1$;

5 R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl- R^{11} , C_1 - C_6 alkoxy- R^{11} , COR^{17} , CO_2R^7 , $CONHR^7$, $N(R^8)_2$, amino C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkyl, amino, amino C_1 - C_4 alkyl- R^7 , C_1 - C_6 alkyl-NHR⁷, carbonitrile, SR^{10} , halo, NHR⁷, NR^8R^9 , NHR⁷- C_1 - C_6 alkyl, NR^8R^9 - C_1 - C_6 alkyl, nitro, cyano, $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , C_1 - C_6 alkyl- COR^{11} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, or C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ;

R^7 and R^8 are each independently selected from -H, C_1 - C_6 alkyl, C_1 - C_4 alkyl- R^{11} , C_1 - C_6 alkyl- $N(R^{13})_2$, CO_2R^{16} , COR^{17} , aryl, and arylalkyl, wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

15 R^9 and R^{10} are each independently selected from -H, hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{17} , C_1 - C_6 alkyl- NH_2R^{13} , CO_2R^{16} , COR^{17} , C_1 - C_6 alkyl- CO_2R^{16} , C_1 - C_6 alkyl- $CONH-R^{16}$, C_1 - C_6 alkyl- $CON(R^{16})_2$, hydroxy C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, halo C_1 - C_4 alkyl, $Si(R^{13})_2R^{17}$, aryl, heteroaryl, heterocyclyl, arylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

20 R^{11} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxyl, halo, amino, NHR¹³, $N(R^{13})_2$, COR^{13} , CO_2R^{17} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

25 R^{12} is selected from -H, hydroxyl, oxo, C_1 - C_6 alkyl, hydroxyl C_1 - C_6 alkyl- R^{11} , C_1 - C_{10} alkoxy, amino, amino C_1 - C_4 alkyl- R^7 , NHR⁷, $N(R^7)_2$, C_1 - C_6 alkyl-NHR⁷, C_1 - C_6 alkyl-NHR⁸ R^9 , C_1 - C_6 alkyl- $N(R^8)_2$, C_1 - C_6 alkyl- R^{11} , C_1 - C_6 alkyl- $CO_2R^7R^{11}$, C_1 - C_6 alkoxy- R^{11} , nitro, $O-R^{10}$, $C=O$, COR^{11} , CO_2R^{11} , SR^{10} , SOR^{11} , SO_2R^{11} , $NHSO_2R^{11}$, C_1 - C_6 alkyl- SR^{10} , halo, halo C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, hydroxy C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkoxy,

aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹³ and R¹⁴ are each independently selected from -H, oxo, C₁-C₆ alkyl, COR²³, and aryl;

R¹⁵ and R¹⁶ are each independently selected from -H, aryl, arylalkyl, wherein aryl, arylalkyl, are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁷ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkyl-R¹⁹, NHR¹⁹, aryl, heteroarylalkyl, and heterocyclalkyl, wherein aryl is optionally substituted with one or more of the groups defined by R²⁴;

R¹⁸ is selected from -H, oxo, hydroxyl, C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, amino, amino C₁-C₆ alkyl, N(R¹⁹)₂, C₁-C₆ alkyl-N(R¹⁹)₂, CO₂R²³, SR²¹, halo, halo C₁-C₄ alkyl, aryl, heteroaryl, and heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁹ and R²⁰ are each independently selected from -H, C₁-C₆ alkyl, heteroaryl, heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R³⁰;

R²¹ and R²² are each independently selected from -H and C₁-C₆ alkyl;

R²³ is selected from -H and C₁-C₆ alkyl;

R²⁴ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, CO₂R²⁹, halo, and halo C₁-C₄ alkyl;

R²⁹ is selected from -H, and C₁-C₆ alkyl;

R³⁰ is selected from -H, aryl, heteroaryl, heterocyclyl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R³⁶;

R³⁶ is selected from -H and halo;

L is selected from:

- 1) carboxyamino,
- 2) carboxyaminoalkyl,

- 5 3) alkenyl,
 4) alkynyl,
 5) alkyl,
 6) hydrazoalkyl,
 7) arylcarbamyl,
 8) aryl,
 9) heteroaryl,
 10) arylalkyl,
10 11) arylalkylamino, or
 12) alkylaryl,








n is an integer that is selected from 0, or 1;








R², R³ and R⁴ are each independently selected from an R¹ group; and








- 15 R³ and R⁴ optionally join to form a ring of 5, 6, 7, or 8 atoms, where the atoms in the ring are independently selected from Z³, Z⁴, O, S, C=O, C=S, S=O, SO₂, C that is mono or di-substituted with an R¹ group, and N that is unsubstituted or substituted with an R¹ group.








- 20 **[00038]** Table I and Table II show examples of MK-2 inhibiting compounds of the present invention, and also shows the chemical name and, where available, the IC₅₀ value of the compound for MK-2 inhibition.








- It is believed that any of the compounds that are listed in Table I and Table II are MK-2 inhibiting compounds that can be used in the method of the present invention. However, neither the novel MK-2 inhibiting compounds,
25 nor the uses of an MK-2 inhibiting compound that are described herein are intended to be limited to the compounds that are presented in the following Tables.








Table I: MK-2 Inhibiting Compounds			
No.	Structure ^a	Compound Name(s) ^b	MK-2 Avg. IC ₅₀ (uM)
1	 1.25 <chem>HO-C(=O)-CF3</chem>	1-(2-aminoethyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.0269
2	 H-Cl H-Cl	1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride	0.0397
3		6-(aminomethyl)-2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.0477
4	 <chem>HO-C(=O)-CF3</chem>	1-(2-aminoethyl)-3-{2-[(E)-2-phenylethenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.0505
5	 H-Cl H-Cl	1-(2-aminoethyl)-3-[2-(4-(hydroxymethyl)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride	0.0533
6		6-(hydroxymethyl)-2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.0615
7	 H-Cl H-Cl	1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylic acid dihydrochloride	0.0686








8	 2 HCl	1-(3-aminopropyl)-3-[2-(4-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride	0.102
9	 H-Cl H-Cl	1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride	0.109
10	 HO-CF₃	1-(2-aminoethyl)-3-[2-(4-(dimethylamino)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.117
11	 HO-CF₃	1-(2-aminoethyl)-3-[2-(4-(trifluoromethoxy)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.161
12	 HO-CF₃	1-(2-aminoethyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.168
13	 HO-CF₃	2-[2-(3-nitrophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.171
14		2-(2-isoquinolin-7-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.194








15	 2 HCl	1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride	0.196
16	 1.5 TFA	2-[2-(1H-indazol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.199
17	 2 HCl	1-(3-aminopropyl)-3-[2-(4-(dimethylamino)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride	0.203
18		2-[2-[(E)-2-pyridin-3-ylethenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.212
19		2-[2-(quinolin-3-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.216
20	 HO-CF₃	1-(2-aminoethyl)-3-[2-(3-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.217
21	 1.25 TFA	2-[2-(1H-pyrrolo[2,3-b]pyridin-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.228

22	 1.5 TFA	6-(hydroxymethyl)-2-[2-(1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.25
23		2-(2-quinolin-6-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.298
24	 2 HCl	1-(3-[[2-(4-bromophenyl)ethyl]amino]propyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid hydrochloride	0.3
25	 HO-CF₃	1-(3-[[2-(4-bromophenyl)ethyl]amino]propyl)-3-{2-[(E)-2-phenylethenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.315
26	 HO-CF₃	2-[2-(4-methoxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.318
27	 2.25 TFA	2-[2-(1-methyl-1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.322
28	 HO-CF₃	2-[2-[(E)-2-phenylvinyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.371

29		2-[2-(5-chlorothiophen-2-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.444
30		2-[2-(3-fluoro-4-methoxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.471
31		2-[2-(1-oxo-2,3-dihydro-1H-inden-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.478
32		2-[2-(1-methyl-1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.479
33		2-[2-(3-fluorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.505
34		methyl 4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]benzoate	0.511
35		2-[2-(3-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.52








36		2-[2-(2,3-dihydro-1-benzofuran-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.521
37		2-[2-(1-benzofuran-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.543
38		2-[2-(2-oxo-2,3-dihydro-1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.554
39		2-[2-(2-quinolin-3-ylpyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	0.577
40		2-[2-[4-(methylthio)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.581
41		2-[2-(2,3'-bipyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.586
42		2-[2-(4-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.611








43		2-[2-[4-(2-morpholin-4-ylethoxy)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.63
44		2-[2-(3-chlorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.638
45		1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate	0.735
46		2-[2-[4-(2-(dimethylamino)ethoxy)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.766
47		2-[2-(1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.808
48		2-[2-[4-(methylsulfonyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.821
49		2-[2-(1,2,3,4-tetrahydroquinolin-6-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.835








50		2-(6'-methoxy-2,3'-bipyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.866
51		2-[2-(3-methoxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.906
52		2-[2-(4-aminophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	0.914
53		5-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]-1H-indole-3-carbaldehyde	0.923
54		2-[2-[4-(hydroxymethyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.929
55		2-[2-(1H-indol-6-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.93
56		2-[2-(1-benzothien-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	0.965








57		4-oxo-2-(2-(quinolin-3-yl)pyridin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazine-6-carboxamide	1.01
58		ethyl 1-(2-(2-(quinolin-3-yl)pyridin-4-yl)-1H-pyrazole-5-carboxylate) dihydrochloride	1.03
59		2-[2-(2,3-dihydro-1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.07
60		2-[2-(4-methoxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	1.09
61		3-chloro-2-[2-(3-chlorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.13
62		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]benzaldehyde trifluoroacetate	1.15
63		1-(3-aminopropyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid	1.16

64		13012 or 6-(chloromethyl)-2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.17
65		2-[2-(2-furyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.2
66		2-[2-(3-nitrophenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride	1.26
67		2-[2-(1,3-thiazol-2-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.29
68		2-[2-(3-aminophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.3
69		1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxamide trifluoroacetate	1.32
70		2-[2-(1H-imidazol-1-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	1.36








71		2-[2-(4-hydroxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate	1.37
72		3-bromo-2-[2-(3-chlorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.39
73		2-[2-(3-furyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.46
74	 2.25 TFA	2-[2-[4-(dimethylamino)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	1.53
75		2-(2-thien-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.55
76		2-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.56
77		2-(5'-methoxy-2,3'-bipyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.69








78		2-(2-{4-[(isopropylamino)methyl]phenyl}pyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.73
79		2-[2-{4-(hydroxymethyl)phenyl}pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	1.75
80		2-(2-{4-[2-(dimethylamino)ethyl]phenyl}pyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	1.98
81		2-(2-pyrimidin-5-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	1.99
82		2-[2-(1,2,3,4-tetrahydroisoquinolin-7-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.08
83		2-[2-{3-(methylsulfonyl)phenyl}pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.1
84		2-[2-{3-(2-piperazin-1-ylethoxy)phenyl}pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrobromide	2.13








85		N-[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]phenyl]methanesulfonamide	2.29
86		2-[2-[4-(dimethylamino)quinazolin-6-yl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.4
87		2-[2-[3-[2-(dimethylamino)ethoxy]phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.4
88		2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	2.6
89		2-[2-(2,4-difluorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.62
90		2-[2-(4-methyl-1H-imidazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride	2.76
91		2-[2-(1H-pyrrol-2-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.79

92		2-[2-(2-methyl-1H-indol-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.83
93		2-[2-(2-fluorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.95
94		2-[2-[4-(morpholin-4-ylmethyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	2.99
95		ethyl 4-(2-[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]phenoxy]ethyl)piperazine-1-carboxylate	3.02
96		ethyl 1-(2-aminoethyl)-3-[2-[4-(hydroxymethyl)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	3.25
97		2-[2-[3-(aminomethyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	3.25
98		2-[2-[3-(2-pyrrolidin-1-ylethoxy)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	3.27








99		N-[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]phenyl]acetamide	3.31
100		2-[2-[4-(dimethylamino)phenyl]pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate	3.44
101		2-[2-[4-(morpholin-4-ylcarbonyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	3.65
102		3-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]phenyl]propanoic acid	4.19
103		2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	4.2
104		2-[2-(1,3-benzothiazol-2-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	4.27
105		2-[2-(4-phenyl-1H-imidazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate	4.31








106		2-[2-[3-(hydroxymethyl)phenyl]pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate	4.54
107		6-(hydroxymethyl)-2-[2-(1,2,3,4-tetrahydroquinolin-3-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	4.9
108		ethyl 1-(2-aminoethyl)-3-[2-(3-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	4.98
109		2-[2-(2H-1,2,3-triazol-2-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one compound with 2-[2-(1H-1,2,3-triazol-1-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one (1:1)	5.21
110		ethyl 1-(2-aminoethyl)-3-[2-(3-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	5.23
111		2-(2,2'-bipyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	5.34
112		2-[2-[4-[(dimethylamino)methyl]phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	5.35

113		2-[2-(1H-imidazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride	5.66
114		2-[2-(3-[(2-hydroxyethyl)amino]methyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	5.78
115		2-[2-(3-[(isopropylamino)methyl]phenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	6.27
116		2-[2-(3-(2-morpholin-4-ylethoxy)phenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	6.73
117		2-[2-(4-phenyl-1H-imidazol-1-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	7.04
118		1-(2-aminoethyl)-N-hydroxy-3-[2-(4-(hydroxymethyl)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxamide trifluoroacetate	7.91
119		1-(3-aminopropyl)-3-(4-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid dihydrochloride	8.11








120		2-[2-(1H-benzimidazol-2-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	8.92
121		2-[2-(1H-1,2,3-triazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride	9.05
122		2-pyridin-4-yl-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	10.1
123		2-[2-(1H-pyrrol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate	10.6
124		1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxamide dihydrochloride	10.7
125		2-[2-[3-(morpholin-4-ylmethyl)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	11.4
126		4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridine-2-carbonitrile	13.8








127		3-[2-(1H-imidazol-1-yl)pyridin-4-yl]-1-propyl-1H-pyrazole-5-carboxylic acid	14.3
128		2-(2-{3-[(dimethylamino)methyl]phenyl}pyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	15.4
129		2-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	15.5
130		2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride	16.2
131		ethyl 1-(2-aminoethyl)-3-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride	20
132		2-[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyridin-2-yl]benzyl]-1H-isoindole-1,3(2H)-dione	20
133		ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride	24.5








134		1-(2-((3-(5-methyl-2-furyl)butyl)amino)ethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate	27.2
135		2-(2-(1H-1,2,4-triazol-1-yl)pyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride	28.2
136		ethyl 1-(2-aminoethyl)-3-[2-(4-((tert-butyl(dimethyl)silyl)oxy)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	28.3
137		ethyl 1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	31.6
138		2-(2-(dimethylamino)pyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	37.2
139		ethyl 1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride	38.6
140		2-(2-chloropyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	44.8

141	 2 HCl	ethyl 1-(3-aminopropyl)-3-[2-(4-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride	47.5
142	 HO-C(=O)-CF₃	2-[2-(4-methyl-1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate	53.4
143	 HCl	ethyl 1-(3-aminopropyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate hydrochloride	60.8
144	 HO-C(=O)-CF₃	2-[2-(3,4-dichlorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	65.5
145		2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	80.4
146	 HO-C(=O)-CF₃	2-[2-(3,4-difluorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	88.9
147		2-(4-hydroxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	118

148		2-(2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	124
149		2-(2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	131
150		1-(3-aminopropyl)-3-(3-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid trifluoroacetate	153
151		1-[3-[(tert-butoxycarbonyl)amino]propyl]-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylic acid	155
152		ethyl 3-(1-oxidopyridin-4-yl)-1H-pyrazole-5-carboxylate	200
153		ethyl 1-(3-aminopropyl)-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate hydrochloride	200
154		1-[3-[(tert-butoxycarbonyl)amino]propyl]-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylic acid	200

155		ethyl 3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate	200
156		ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate	200
157		2-[2-(1H-1,2,4-triazol-1-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	200
158		ethyl 1-[2-[(tert-butoxycarbonyl)amino]ethyl]-3-[2-(4-[(tert-butyl(dimethyl)silyl]oxy)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate	200
159		ethyl 1-[2-[(tert-butoxycarbonyl)amino]ethyl]-3-[2-[4-(hydroxymethyl)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylate	200
160		2-[2-(4-methyl-1H-pyrazol-1-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	200
161		2-[2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate	200

162		ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate	200
163		2-(3-methoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	200
164		2-(3-hydroxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	200
165		ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylate	200
166		2-(4-methoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one	200
167		ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate	
168		ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate	

169	 H-Cl H-Cl	ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	
170	 H-Cl H-Cl	ethyl 1-(2-aminoethyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	
171	 H-Cl H-Cl	ethyl 1-(2-aminoethyl)-3-[2-(4-(trifluoromethoxy)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	
172	 H-Cl H-Cl	ethyl 1-(2-aminoethyl)-3-[2-((E)-2-phenylethenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride	
173	 HO-CF₃	ethyl 1-(2-aminoethyl)-3-[2-(4-(dimethylamino)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate trifluoroacetate	
174		ethyl 1-(3-((tert-butoxycarbonyl)amino)propyl)-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate	
175	 2 HCl	1-(3-aminopropyl)-3-[2-(3-(hydroxymethyl)phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride	






























176	 2 HCl	1-(3-aminopropyl)-3-[2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride	
177		2-(2-pyrimidin-5-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
178		2-[2-(2,4-dimethoxypyrimidin-5-yl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
179		4-[4-{4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl}pyridin-2-yl]benzoic acid	
180		2-[[4-oxo-2-(2-quinolin-3-ylpyridin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-6-yl]methyl]-1H-isoindole-1,3(2H)-dione	

Table II: Additional MK-2 Inhibiting Compounds			
No.	Structure ^a	Compound Name(s) ^b	MK-2 Avg. IC ₅₀ (uM)
181		2-[2-{3,5-bis(trifluoromethyl)phenyl}pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
182		2-(2-phenylpyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
183		2-[2-(2-bromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
184		2-[2-(2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
185		2-[2-(pentafluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
186		2-[2-(2,5-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
187		2-[2-(2,6-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
188		2-[2-(2-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









189		2-[2-(2-chloro-6-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
190		2-[2-(2,6-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
191		2-[2-(2-chloro-6-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
192		2-[2-(2-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
193		2-[2-(2,3-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
194		2-[2-(2,3,4-trimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
195		2-[2-(2,4-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
196		2-[2-(2,4,6-trimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

197		2-[2-(2,6-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
198		2-[2-(2-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
199		2-[2-(2-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
200		2-[2-(2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
201		2-[2-(2,5-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
202		2-[2-(3-bromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
203		2-[2-(3-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
204		2-[2-(3-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









205		2-[2-(3-chloro-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
206		2-[2-(3,5-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
207		2-[2-(3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
208		2-[2-(3,4-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
209		2-[2-(3,4,5-trimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
210		2-[2-(3,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
211		2-[2-(3-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
212		2-[2-(3-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

213		2-[2-(3-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
214		2-[2-(4-bromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
215		2-[2-(4-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
216		2-[2-(4-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
217		N-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
218		2-[2-[4-(dimethylamino)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
219		2-[2-(4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
220		2-[2-(4-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









221		2-[2-(4-hydroxy-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
222		2-[2-(1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
223		methyl 4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoate	
224		ethyl 4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoate	
225		2-[2-[4-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
226		2-[2-(4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
227		2-[2-(2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
228		2-[2-(3,5-dibromo-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









229		2-[2-(3,5-dichloro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
230		2-[2-(3-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
231		2-[2-(4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
232		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
233		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
234		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzaldehyde	
235		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzaldehyde	
236		2-[2-(1,3-benzodioxol-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









237		2-(2-quinolin-3-ylpyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
238		2-[2-(2-aminophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
239		2-[2-(4-amino-2,3,5,6-tetrafluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
240		2-[2-(3-aminophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
241		2-[2-(4-aminophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
242		2-[2-(6-methoxy-2-naphthyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
243		2-[2-(2,4-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
244		2-[2-(2,3-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

245		2-[2-(3,5-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
246		2-[2-[4-(methylthio)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
247		2-[2-(3,4-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
248		2-[2-(2,3,5,6-tetrafluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
249		2-[2-(2,3,6-trifluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
250		2-[2-(2,3,4-trifluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
251		2-[2-(2,4,5-trifluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
252		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzamide	









253		2-[2-(3-chloro-4-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
254		2-[2-(pentamethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
255		2-[2-(2-amino-6-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
256		2-[2-(2-amino-6-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
257		2-[2-[2-(methylthio)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
258		2-[2-(2,3-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
259		2-[2-(2,4-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
260		2-[2-(2,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









261		methly 2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoate	
262		2-[2-(2,4-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
263		2-(2-mesitylpyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
264		2-[2-(3,4-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
265		2-[2-(3,4-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
266		2-[2-[4-(methylamino)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
267		2-[2-(4-benzoylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
268		2-[2-(4-ethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

269		2-[2-(3,4-dihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
270		2-[2-(3,5-dihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
271		2-[2-(2-naphthyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
272		2-[2-(3-amino-4-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
273		2-[2-(2-amino-5-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
274		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzamide	
275		2-[2-(3,5-dibromo-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
276		2-[2-(4-amino-3,5-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









277		2-[2-(4-phenoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
278		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzaldehyde	
279		2-[2-(2,4,5-trimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
280		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzamide	
281		2-[2-(5-amino-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
282		2-[2-(2,5-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
283		2-[2-[2-chloro-5-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
284		2-[2-(2-chloro-6-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









285		2-[2-(2,3-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
286		2-[2-[4-(diethylamino)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
287		2-[2-(4-bromo-3-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
288		2-[2-(4-amino-2-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
289		2-[2-(2-amino-4-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
290		2-[2-(4'-pentyl-1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
291		2-[2-(2,6-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
292		2-[2-[4-(trifluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









293		2-[2-(2,3,4,5-tetrafluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
294		2-[2-(2-piperazin-1-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
295		2-[2-(4-bromo-2-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
296		2-[2-(3-(trifluoromethoxy)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
297		2-[2-(4-amino-3-ethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
298		2-[2-(4-bromo-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
299		2-[2-(4-amino-3-chloro-5-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
300		2-[2-(4-amino-2-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

301		2-[2-(3-chloro-2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
302		2-[2-(2-fluoro-6-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
303		2-[2-(5-fluoro-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
304		2-[2-(2,4,6-trifluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
305		2-[2-[2-(trifluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
306		2-[2-(2-chloro-4-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
307		2-[2-(4-butoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
308		2-[2-[4-(octyloxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

309		2-[2-(2-chloro-5-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
310		2-[2-(2-ethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
311		2-[2-(3-chloro-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
312		2-[2-(5-chloro-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
313		2-[2-(3-ethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
314		2-[2-(4-chloro-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
315		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
316		2-[2-(2-piperidin-1-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









317		2-[2-(4-amino-2,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrochloride	
318		2-[2-(2-chloro-6-phenoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
319		2-[2-(4-tert-butyl-2,6-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
320		2-[2-(2-chloro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
321		2-[2-(2-(dimethylamino)-6-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
322		2-[2-(4-butylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
323		N-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-3-(trifluoromethyl)phenyl]acetamide	
324		2-[2-[2-(2,4-dichlorophenoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

325		2-[2-[4-(2-hydroxyethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
326		2-[2-(2,4,6-trichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
327		N-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-2-(trifluoromethyl)phenyl]acetamide	
328		2-[2-(2-ethyl-6-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
329		2-[2-(2,4-dichloro-6-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
330		2-[2-(5-chloro-2-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
331		2-[2-(4-amino-3-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
332		methyl 4-methoxy-3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoate	









333		2-[2-(4-isopropylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
334		2-[2-(4-methoxy-2,5-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
335		2-[2-(3-bromo-4-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
336		2-[2-(4-amino-3-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
337		2-[2-(2-amino-4,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
338		2-[2-(2-amino-6-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
339		2-[2-(4'-hydroxy-1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
340		2-[2-(1,1'-biphenyl-2-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









341		2-[2-(3,5-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
342		2-[2-(4-chloro-2-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
343		2-[2-(2-fluoro-3-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
344		2-[2-(2-fluoro-6-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
345		2-[2-(2,4-bis(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
346		2-[2-(2,6-bis(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
347		2-[2-(2-fluoro-4-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
348		2-[2-(2-fluoro-5-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









349		2-(2-{3-fluoro-5-(trifluoromethyl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
350		2-(2-{4-fluoro-2-(trifluoromethyl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
351		2-(2-{4-fluoro-3-(trifluoromethyl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
352		2-(2-{2-fluoro-6-phenoxyphenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
353		2-(2-{2-fluoro-6-(4-fluorophenoxy)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
354		2-(2-{2-fluoro-6-(4-methylphenoxy)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
355		2-(2-{2-fluoro-6-[(4-methylbenzyl)oxy]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
356		2-(2-{2-[(4-chlorobenzyl)oxy]-6-fluorophenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









357		2-(2-[2-fluoro-6-[(4-methylphenyl)thio]phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
358		2-(2-[2-[(4-chlorophenyl)thio]-6-fluorophenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
359		2-(2-[2-fluoro-6-(2,2,2-trifluoroethoxy)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
360		2-(2-[2-(benzyloxy)-6-methoxyphenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
361		2-(2-[2-[(2-chlorobenzyl)oxy]-6-methoxyphenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
362		2-(2-[2-methoxy-6-(2,2,2-trifluoroethoxy)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
363		2-(2-[2-ethoxy-6-(2,2,2-trifluoroethoxy)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
364		2-(2-[2-isopropoxy-6-(2,2,2-trifluoroethoxy)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









365		2-[2-[2-(phenylthio)-5-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
366		2-[2-[4'-(pentyloxy)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
367		2-[2-[4'-(heptyl)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
368		2-[2-(3,4,5-trifluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
369		2-[2-[4'-(hexyl)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
370		2-[2-[4'-(octyloxy)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
371		2-[2-[4'-(octyl)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
372		2-[2-(4-bromo-2,3,5,6-tetrafluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









373		2-((2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl)thio)benzoic acid	
374		2-[2-(10,10-dioxido-9-oxo-9H-thioxanthen-3-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
375		2-[2-[4-(4-pentylcyclohexyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
376		2-[2-(4-tert-butylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
377		2-[2-[4-(benzyloxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
378		2-[2-(2,3,5-trifluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
379		2-[2-[2-chloro-4-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
380		2-[2-[4-(1,3-oxazol-5-yl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









381		2-[2-[4-(difluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
382		2-[2-(4-hydroxy-7-methyl-2,3-dihydro-1H-inden-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
383		2-[2-[2,6-bis(2,2,2-trifluoroethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
384		5-methoxy-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
385		2-[2-(2-[[3-(dimethylamino)propyl]amino]-6-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
386		2-[2-(2-fluoro-6-piperidin-1-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
387		2-[2-[3-bromo-2,6-bis(2,2,2-trifluoroethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
388		2-[2-[4'-(hexyloxy)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

389		2-{2-[4'-(heptyloxy)-1,1'-biphenyl-4-yl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
390		2-{2-(2-fluoro-5-methylphenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
391		2-{2-(2-bromo-5-fluorophenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
392		2-{2-(2-fluoro-5-methoxyphenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
393		2-{2-(3-bromo-4-hydroxyphenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
394		2-{2-(3-bromo-4-methoxyphenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
395		2-{2-(2-fluoro-4,5-dimethoxyphenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
396		2-{2-(4-bromo-2-fluorophenyl)pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

397		2-[2-(4-chloro-2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
398		2-[2-(4-chloro-3-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
399		2-[2-(9-oxo-9H-fluoren-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
400		2-[2-(9H-fluoren-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
401		2-[2-(4-(heptyloxy)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
402		2-[2-(4-(hexyloxy)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
403		2-[2-(4-heptylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
404		2-[2-(4-(pentyloxy)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









405		2-[2-(5-bromo-2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
406		2-[2-(5-bromo-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
407		2-[2-(5-bromo-2-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
408		2-[2-(4'-methyl-1,1'-biphenyl-2-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
409		2-[2-(3-fluoro-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
410		2-[2-(2-chloro-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
411		2-[2-(3,5-di-tert-butyl-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
412		2-[2-(2-amino-3,5-dichlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









413		2-[2-(2-bromo-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
414		1-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]-2-phenylethane-1,2-dione	
415		2-[2-[2-(phenylthio)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
416		N-butyl-N'-[4-methyl-3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]urea	
417		2-morpholin-4-yl-N-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
418		2-[2-(4-[[4-bromo-3-methylphenyl]amino]methyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
419		2-(2-[2-fluoro-6-[3-(trifluoromethyl)phenoxy]phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
420		4-tert-butyl-N-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]benzenesulfonamide	









421		2-[2-[2-(4-chloro-2-methylphenoxy)-6-fluorophenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
422		2-[2-(pentachlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
423		2-[2-(4-methoxy-7-methyl-2,3-dihydro-1H-inden-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
424		2-[2-(2-amino-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
425		2-[2-(4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
426		2-[2-(2-chloro-3,4-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
427		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzenesulfonamide	
428		2-[2-(3,5-dibromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









429		2-[2-(2-amino-3,5-dibromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
430		2-[2-(4-amino-2,5-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
431		2-[2-(4-chloro-3-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
432		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenylalanine	
433		2-[2-(4'-amino-1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
434		2-[2-(2-fluoro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
435		2-[2-(4'-[(2S)-2-methylbutyl]oxy)-1,1'-biphenyl-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
436		N-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-2-(trifluoromethoxy)phenyl]acetamide	









437		2-[2-[4-amino-3-(trifluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
438		2-(2-{3-[(2-oxo-1,3-benzoxazol-3(2H)-yl)methyl]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
439		N-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]methanesulfonamide	
440		2-(2-{2-amino-5-[1-hydroxy-2-(isopropylamino)ethyl]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
441		2-[2-[4-(4-mercaptobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
442		2-(2-{10-[3-(4-hydroxypiperidin-1-yl)propyl]-10H-phenothiazin-2-yl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
443		2-[2-(3-fluoro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
444		2-[2-(3-fluoro-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









445		2-[2-[4-(methylsulfonyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
446		2-[2-(4-pentylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
447		2-[2-(5-chloro-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
448		2-[2-[2,6-dichloro-4-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
449		N-[2-ethyl-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
450		2-[2-(2,3-dihydro-1,4-benzodioxin-6-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
451		2-[2-[2-(difluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
452		2-[2-(4-hydroxy-3,5-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









453		N-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
454		2-[2-(4-chloro-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
455		2-[2-(3-amino-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
456		2-[2-[3-(difluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
457		2-[2-[2,3-difluoro-4-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
458		2-[2-[3,4-difluoro-5-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
459		2-[2-[3-fluoro-4-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
460		2-[2-[5-fluoro-2-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









461		2-[2-(2,2,3,3-tetrafluoro-2,3-dihydro-1,4-benzodioxin-6-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
462		2-[(2-(2-(trifluoromethyl)thio)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
463		2-[(2-(3-(trifluoromethyl)thio)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
464		2-[2-(1,1'-biphenyl-3-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
465		2-[2-(4-hydroxy-2,6-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
466		2-[2-(2,3-difluoro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
467		2-[2-(2-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
468	CHIRAL 	4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	









469	CHIRAL 	4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-D-phenylalanine	
470		2-[2-(4-mercaptophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
471		2-[2-(4-amino-3-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
472		2-[2-(3-(cyclopentyloxy)-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
473		2-[2-(4-butyl-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
474		2-[2-(5-amino-2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
475		4'-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1,1'-biphenyl-4-carboxylic acid	
476		2-[2-(4'-propyl-1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









477		2-[2-(4'-butyl-1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
478		2-[(4-{4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl}phenyl)amino]carbonyl]benzoic acid	
479		2-[2-(3,5-dibromo-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
480	 CHIRAL HCl	2-(2-{3-[(3S)-1-propylpiperidin-3-yl]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrochloride	
481		2-[2-[4-(2-bromobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
482		2-[2-[4-(3-bromobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
483		2-[2-[4-(4-bromobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
484		2-[2-[2-(2-bromobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

485		2-[2-[2-(3-bromobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
486		2-[2-[2-(4-bromobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
487		2-[2-(2-benzoylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
488		N-acetyl-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
489		2-[2-(2-bromo-4-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
490		2-[2-(5-bromo-2-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
491		2-[2-(2,5-dibromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
492		2-[2-(3,5-dibromo-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









493		2-[2-[2-(octyloxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
494		2-(2-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
495		2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
496		2-[2-(2,2-difluoro-1,3-benzodioxol-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
497		2-[2-(10H-phenothiazin-3-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
498		2-[2-(2,5-dihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
499		2-[2-(4-propylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
500		2-[2-(4-hexylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









501		2-[2-(4-octylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
502		2-[2-(4'-ethyl-1,1'-biphenyl-4-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
503		2-[2-[4-(4-butylcyclohexyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
504		methyl 2-hydroxy-5-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoate	
505		2-[2-(2,3-dihydro-1-benzofuran-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
506		2-[2-(2-chloro-4-methylquinolin-3-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
507		2-[2-[4-(hydroxymethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
508		2-[(3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl)amino]carbonyl]benzoic acid	









509		{[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzyl}thioacetic acid	
510		N-[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
511		2-[2-(1-oxo-1,3-dihydro-2-benzofuran-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
512		N-acetyl-3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenylalanine	
513		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
514		2-[2-[4-(benzyloxy)-3-methoxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
515		2-[2-(3-ethoxy-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
516		2-[2-[4-(benzyloxy)-3-ethoxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

517		2-[2-(4-bromo-2,6-difluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
518		2-[2-[2-(2-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
519		2-[2-[3-(2-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
520		2-[2-[4-(2-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
521		2-[2-[2-(3-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
522		2-[2-[3-(3-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
523		2-[2-[4-(3-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
524		2-[2-[2-(4-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









525		2-[2-[3-(4-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
526		2-[2-[4-(4-methoxybenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
527		2-[2-[2-(dimethylamino)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
528		2-[2-(2-morpholin-4-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
529		2-[2-(2-azepan-1-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
530		2-[2-[2-(cyclopropylmethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
531		2-[2-(4'-bromo-1,1'-biphenyl-2-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
532		2-[2-(4'-bromo-1,1'-biphenyl-3-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









533		2-(2-[4-(1,3-benzothiazol-2-yl)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
534		ethyl 4'-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1,1'-biphenyl-4-carboxylate	
535		ethyl 4'-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1,1'-biphenyl-3-carboxylate	
536		ethyl 2'-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1,1'-biphenyl-3-carboxylate	
537		1-[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]-2-phenylethane-1,2-dione	
538		1-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]-2-phenylethane-1,2-dione	
539		ethyl oxo[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetate	
540		ethyl oxo[3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetate	









541		ethyl oxo{4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)phenyl]acetate	
542		2-[2-[2-fluoro-6-(2-fluorobenzoyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
543		2-[2-[2-(trifluoroacetyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
544		2-[2-[3-(trifluoroacetyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
545		2-[2-[4-(trifluoroacetyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
546		2-[2-(4-piperazin-1-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
547		2-[2-[3-(hydroxymethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
548		2-[2-[2-(benzyloxy)-5-bromophenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









549		2-[2-(3-chloro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
550		2-[2-(4-fluoro-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
551		2-[2-[4-(1H-pyrazol-3-yl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
552		2-[2-(5-chloro-2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
553		2-[2-[2-methyl-4-(trifluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
554		2-[2-[3-chloro-4-(trifluoromethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
555		2-[2-[3-fluoro-2-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
556		2-[2-[4-methoxy-3-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









557		2-{2-[2-methyl-3-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
558		2-{2-[2-methyl-5-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
559		2-{2-[3-methyl-5-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
560		2-{2-[4-methyl-2-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
561		2-{2-[4-methyl-3-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
562		2-{2-[2,4-difluoro-5-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
563		2-{2-[2,5-difluoro-4-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
564		2-{2-[3,5-difluoro-4-(trifluoromethyl)phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









565		2-[2-(4,5-difluoro-2-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
566		2-(2-[4-((trifluoromethyl)sulfonyl)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
567		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-D-phenylalanine	
568		2-[2-(5-amino-2-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrochloride	
569		2-[2-(3-chloro-4-hydroxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
570		2-(2-[3-[(dimethylamino)methyl]-1H-indol-5-yl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
571		2-[2-[4-(4-chloro-3-fluorophenoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
572		2-[2-[4-(2,4-dimethylphenoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









573		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-D-phenylalanine	
574		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
575		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
576		2-[2-(2-bromo-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
577		2-[2-(4-amino-3-bromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
578		N-[2-chloro-5-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
579		N-[2-chloro-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
580		2-[2-[4-(aminomethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrochloride	









581		2-[2-[3,4-bis(benzyloxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
582		N-acetyl-3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
583		4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenylalanine	
584		2-[2-(6-hydroxy-2-naphthyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
585		2-(2-[4-(3-(trifluoromethyl)phenoxy)phenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
586		2-(2-[4-(4-chlorobenzyl)oxy]phenyl)pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
587		2-[2-[2-(4-methoxyphenoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
588		2-[2-(4-piperazin-1-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrochloride	

589		2-[2-(4-amino-3-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
590		N-(4-methylphenyl)-2-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
591		2-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-phenylacetamide	
592		2-[2-[4'-(aminomethyl)-1,1'-biphenyl-2-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one hydrochloride	
593		2-[2-[2-(2-phenylethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
594		2-[2-(3-chloro-5-ethoxy-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
595		2-[2-[2-(benzyloxy)-3-methoxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
596		2-[4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	









597		2-[2-(4-ethoxy-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
598		2-[2-(3-methoxy-4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
599		2-[2-(4-butoxy-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
600		2-[2-[4-(2-hydroxyethoxy)-3-methoxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
601		2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl benzoate	
602		2-[2-(4-isopropoxy-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
603		2-(2-[4-((2,4-dichlorobenzyl)oxy)-3-methoxyphenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
604		2-(2-[4-((2-chlorobenzyl)oxy)-3-methoxyphenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

605		4-((2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)methyl)benzoic acid	
606		2-[2-(3-ethoxy-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
607		2-[2-(3,4-diethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
608		2-[2-(3-ethoxy-4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
609		2-[2-(4-butoxy-3-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
610		2-[2-[3-ethoxy-4-(2-hydroxyethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
611		2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl benzoate	
612		2-[2-(3-ethoxy-4-isopropoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

613		2-(2-(4-((2,4-dichlorobenzyl)oxy)-3-ethoxyphenyl)pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
614		2-(2-(4-((2-chlorobenzyl)oxy)-3-ethoxyphenyl)pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
615		2-(2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-phenylacetamide	
616		N-benzyl-2-(2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetamide	
617		2-(2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(2-phenylethyl)acetamide	
618		2-(2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(2-methylphenyl)acetamide	
619		2-(2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(4-methylphenyl)acetamide	
620		2-(2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(2-methoxyphenyl)acetamide	









621		ethyl 4-(((2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetyl)amino)benzoate	
622		2-[2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N,N-diphenylacetamide	
623		2-[2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-1-naphthylacetamide	
624		2-[2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-[3-(trifluoromethyl)phenyl]acetamide	
625		4-(((2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetyl)amino)benzoic acid	
626		2-[2-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
627		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-phenylacetamide	
628		N-benzyl-2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	









629		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-phenylethyl)acetamide	
630		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-methylphenyl)acetamide	
631		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(4-methylphenyl)acetamide	
632		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-methoxyphenyl)acetamide	
633		ethyl 4-[[[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetyl]amino]benzoate	
634		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-1-naphthylacetamide	
635		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-[3-(trifluoromethyl)phenyl]acetamide	
636		2-[2-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	

637		2-[2-(3-chloro-4,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
638		2-[2-(3-chloro-4-ethoxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
639		2-[2-(3-chloro-5-methoxy-4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
640		2-[2-(4-butoxy-3-chloro-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
641		2-chloro-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl benzoate	
642		2-[2-(3-chloro-4-isopropoxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
643		2-[2-[4-(benzyloxy)-3-chloro-5-methoxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
644		2-(2-[3-chloro-4-[(2-chlorobenzyl)oxy]-5-methoxyphenyl]pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









645		2-[2-(3-chloro-5-ethoxy-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
646		2-[2-(3-chloro-4,5-diethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
647		2-[2-(3-chloro-5-ethoxy-4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
648		2-[2-(4-butoxy-3-chloro-5-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
649		2-[2-[3-chloro-5-ethoxy-4-(2-hydroxyethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
650		2-chloro-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl benzoate	
651		2-[2-(3-chloro-5-ethoxy-4-isopropoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
652		2-[2-[4-(benzyloxy)-3-chloro-5-ethoxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

653		2-(2-{3-chloro-4-[(2,4-dichlorobenzyl)oxy]-5-ethoxyphenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
654		2-(2-{3-chloro-4-[(2-chlorobenzyl)oxy]-5-ethoxyphenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
655		4-[(2-chloro-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)methyl]benzoic acid	
656		2-[2-(3-bromo-4,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
657		2-[2-(3-bromo-4-ethoxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
658		2-[2-(3-bromo-5-methoxy-4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
659		2-[2-(3-bromo-4-butoxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
660		2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl benzoate	









661		2-[2-(3-bromo-4-isopropoxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
662		2-[2-(3-bromo-4-[(2,4-dichlorobenzyl)oxy]-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
663		2-[2-(3-bromo-4-[(2-chlorobenzyl)oxy]-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
664		2-[2-chloro-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-phenylacetamide	
665		2-[2-chloro-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
666		2-[2-chloro-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
667		2-[2-(3-bromo-5-ethoxy-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
668		2-[2-(3-bromo-4,5-diethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









669		2-[2-(3-bromo-5-ethoxy-4-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
670		2-[2-(3-bromo-4-butoxy-5-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
671		2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl benzoate	
672		2-[2-(3-bromo-5-ethoxy-4-isopropoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
673		2-[2-(3-bromo-4-[(2,4-dichlorobenzyl)oxy]-5-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
674		2-[2-(3-bromo-4-[(2-chlorobenzyl)oxy]-5-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
675		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-phenylacetamide	
676		N-benzyl-2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	

677		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-phenylethyl)acetamide	
678		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-methylphenyl)acetamide	
679		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(4-methylphenyl)acetamide	
680		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-methoxyphenyl)acetamide	
681		ethyl 4-[(2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetyl]amino]benzoate	
682		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-1-naphthylacetamide	
683		2-[2-bromo-6-methoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
684		2-[2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-phenylacetamide	









685		N-benzyl-2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetamide	
686		2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(2-phenylethyl)acetamide	
687		2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(2-methylphenyl)acetamide	
688		2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(4-methylphenyl)acetamide	
689		2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-(2-methoxyphenyl)acetamide	
690		ethyl 4-(((2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetyl)amino)benzoate	
691		2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)-N-1-naphthylacetamide	
692		2-(2-bromo-6-ethoxy-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetamide	









693		2-[2-(2-bromo-4,5-dimethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
694		2-[2-(2-bromo-4-ethoxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
695		2-[2-(2-butoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
696		2-[2-[2-(benzyloxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
697		2-[2-[2-(2-chlorobenzyl)oxy]phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
698		2-[2-[2-(2,4-dichlorobenzyl)oxy]phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
699		2-[2-(2-isopropoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
700		2-[2-(5-bromo-2-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









701		2-[2-(5-bromo-2-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
702		2-[2-(5-bromo-2-butoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
703		2-(2-{5-bromo-2-[(2-chlorobenzyl)oxy]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
704		2-(2-{5-bromo-2-[(2,4-dichlorobenzyl)oxy]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
705		2-[2-(5-bromo-2-isopropoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
706		2-[2-(3,5-dibromo-2-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
707		2-[2-(3,5-dibromo-2-propoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
708		2-[2-(3,5-dibromo-2-isopropoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









709		2-{2-[2-(benzyloxy)-3,5-dibromophenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
710		2-{2-[3,5-dibromo-2-[(2-chlorobenzyl)oxy]phenyl]pyrimidin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
711		N-(2-methoxyphenyl)-2-{2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy}acetamide	
712		ethyl 4-[(2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetyl]amino]benzoate	
713		N-benzyl-2-{2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy}acetamide	
714		N-(2-methylphenyl)-2-{2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy}acetamide	
715		2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-phenylacetamide	
716		2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(4-methylphenyl)acetamide	









717		2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-methoxyphenyl)acetamide	
718		N-benzyl-2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]acetamide	
719		2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-phenylethyl)acetamide	
720		ethyl 4-[(4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy)acetyl]amino]benzoate	
721		2-[4-bromo-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenoxy]-N-(2-methylphenyl)acetamide	
722		2-[2-[5-chloro-2-(methylamino)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
723		2-[2-(3-hydroxy-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
724		N-[3-methyl-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	









725		2-[2-(2,3-dihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
726		2-[2-(2,4-dihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
727		2-[2-(2,6-dihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
728		2-[2-(2,3,4-trihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
729		2-[2-(2,4,6-trihydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
730		2-[2-(2-hydroxy-5-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
731		2-[2-(2-hydroxy-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
732		2-[2-(5-bromo-2-hydroxy-3-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









733		2-[2-(3,5-dichloro-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
734		2-[2-[4-(diethylamino)-2-hydroxyphenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
735		2-[2-(2-hydroxy-6-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
736		2-[2-(2-hydroxy-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
737		2-[2-(3-amino-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
738		2-[2-(6-bromo-1,3-benzodioxol-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
739		2-[2-[3-(2-aminoethyl)-2-methyl-1H-indol-5-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
740		2-(4-benzylpiperazin-1-yl)-N-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	

741		2-[4-(4-fluorophenyl)piperazin-1-yl]-N-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)phenyl]acetamide	
742		N-[2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]-2-(4-pyridin-2-yl)piperazin-1-yl]acetamide	
743		2-(2-{4-[(2-fluorobenzyl)oxy]phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
744		2-[2-(3-hydroxy-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
745		2-[2-(5-fluoro-2-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
746		2-[2-(2,4,5-trimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
747		2-[2-(2-amino-5-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
748		2-[2-[3-(2-aminoethyl)-1H-indol-5-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









749		5-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]tryptophan	
750		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
751		3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-L-phenylalanine	
752		2-[2-[4-(5-propyl-1,3-dioxan-2-yl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
753		2-[2-(3-chloro-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
754		2-[2-(4-chloroquinolin-3-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
755		2-[2-(2-chloro-5-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
756		2-[2-(2-fluoro-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









757		2-[2-(3-fluoro-5-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
758		5-fluoro-2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
759		2-fluoro-5-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
760		3-fluoro-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
761		4-fluoro-3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	
762		2-[2-(2-fluoro-5-(hydroxymethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
763		2-[2-(2,5-bis(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
764		2-fluoro-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoic acid	









765		2-[2-(2,6-difluoro-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
766		2-[2-(2,6-difluoro-4-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
767		2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzenesulfonamide	
768		2-[2-(2-chloro-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
769		N-[2-bromo-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
770		N-[2,6-dibromo-4-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]acetamide	
771		2-[2-(4-amino-3,5-dibromophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
772		2-[2-(1,2,3,4-tetrahydroisoquinolin-7-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









773		2-[2-(5-fluoro-1H-indol-6-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
774		5-fluoro-6-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1H-indole-2-carboxylic acid	
775		2-[2-[3-(2-aminoethyl)-5-fluoro-1H-indol-6-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
776		2-[2-(5-fluoro-3-methyl-1H-indol-6-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
777		ethyl 5-fluoro-6-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1H-indole-2-carboxylate	
778		2-[2-(4-bromo-3-ethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
779		2-[2-(4-fluoro-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
780		2-[2-[4-bromo-3-(trifluoromethyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	









781		2-(2-{3-amino-4-(methylamino)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
782		2-(2-{4-(2-aminopyrimidin-4-yl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
783		2-(2-{2-amino-5-(trifluoromethyl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
784		2-(2-{4-amino-2,6-difluorophenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
785		2-(2-{4-amino-3,5-difluorophenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
786		2-(2-{4-(2-methylpyrimidin-4-yl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
787		2-(2-{4-(2-phenylpyrimidin-4-yl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
788		2-(2-{4-(2-pyridin-2-ylpyrimidin-4-yl)phenyl}pyrimidin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

789		2-[2-(2-chloro-3-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
790		2-[2-(3-chloro-5-fluorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
791		2-[2-(3-(1H-pyrazol-3-yl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
792		2-[2-(2,3,6-trifluoro-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
793		2-[2-(4-chloro-3,5-bis(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
794		2-[2-(2-bromo-5-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
795		2-[2-(2-chloro-6-(trifluoromethyl)phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
796		2-[2-(4-[chloro(difluoro)methoxy]phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

797		2-[2-[4'-(trifluoromethyl)-1,1'-biphenyl-2-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
798		2-[2-[4-(1-amino-3-hydroxypropyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
799		2-[2-[3-(1-amino-3-hydroxypropyl)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
800		1-[4-methyl-3-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]phenyl]pyrrolidine-2,5-dione	
801		ethyl 2-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]benzoate	
802		2-[2-[4-(cyclohexylmethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
803		2-[2-[4-[(1-methylheptyl)oxy]phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
804		2-[2-[7-methyl-2,3-dihydro-1H-inden-4-yl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

805		2-[2-(4-methoxy-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
806		2-[2-(2-methoxy-5-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
807		2-[2-(2-amino-3-chlorophenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
808		2-[2-(3-benzyl-2,3,4,5-tetrahydro-1H-1,4-benzodiazepin-7-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
809		2-[2-(4-piperidin-4-ylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
810		2-[2-(4-methoxy-3,5-dimethylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
811		2-[2-(3-oxo-1,3-dihydro-2-benzofuran-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
812		2-[2-(3,4-dimethoxy-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

813		2-[2-(3-methoxy-4-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
814		2-[2-(2-fluoro-6-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
815		2-[2-(3,4-difluoro-2-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
816		2-[2-(3,4-difluoro-2-hydroxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
817		2-[2-(2,3-difluoro-4-methoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
818		2-[2-(4-chloro-3-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
819		2-[2-(4-[[2-oxo-5-(trifluoromethyl)pyridin-1(2H)-yl]methyl]phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
820		2-[2-(4-[[5-(4-methylphenyl)pyrimidin-2-yl]oxy]phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

821		2-[2-(4-[[5-(4-methoxyphenyl)pyrimidin-2-yl]oxy]phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
822		2-[2-(4-[[5-(4-fluorophenyl)pyrimidin-2-yl]oxy]phenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
823		2-[2-(6-amino-1,3-benzodioxol-5-yl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
824		2-[2-[5-bromo-2-(2-hydroxyethoxy)phenyl]pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
825		2-[2-(4-fluoro-3-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
826		2-[2-(3-fluoro-2-methylphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	
827		methyl 4'-[4-(4-oxo-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-2-yl)pyrimidin-2-yl]-1,1'-biphenyl-4-carboxylate	
828		2-[2-(2-amino-4,5-diethoxyphenyl)pyrimidin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one	

Notes:

a) Chemical names were generated by ACD/Name software.

b) The MK-2 inhibiting compound may be shown with a solvent, such as, for example, trifluoroacetate, with which it can form a salt. Both the salt and base forms of each compound are included in the present invention.

[00039] In one embodiment of the present invention, the MK-2 inhibiting compound is one that is listed in Table 1.

[00040] In another embodiment of the present invention, the MK-2 inhibiting compound is one that is listed in Table 1 or in Table 2.

5 [00041] In yet another embodiment of the present invention, the MK-2 inhibiting compound is one that is listed in Table 2.

[00042] It is preferred that the MK-2 inhibiting compound is one that has an IC_{50} value for the inhibition of MK-2 that is lower than 1. By way of example, this would include the compounds in Table I numbered 1 – 56.

10 An MK-2 IC_{50} value that is lower than 0.5 is still more preferred (examples of these compounds include the compounds in Table I numbered 1 – 32), lower than 0.1 is even more preferred yet (examples of these compound include the compounds in Table I numbered 1 – 7).

15 [00043] In one embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except when Z^2 and Z^3 are both nitrogen, R^4 is other than pyrrole, or optionally when Z^4 and Z^5 are both nitrogen and R^a is ring Q, Q^2 is other than nitrogen.

[00044] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is selected from an M-ring or a Q-ring.

20 [00045] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring.

25 [00046] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring wherein ring M is an aromatic pyridine or pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are independently selected from carbon and nitrogen and if M^2 and/or M^6 is carbon, the carbon is substituted with $(L)_nR^1$.

30 [00047] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring wherein ring M is an aromatic pyridine or pyrimidine ring,

wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are independently selected from carbon and nitrogen and if M^2 and/or M^6 is carbon, the carbon is substituted with $(L)_nR^1$; and
[00048] R^3 and R^4 optionally join to form a ring of 5, 6, 7, or 8 atoms,
5 where the atoms in the ring are independently selected from Z^3 , Z^4 , O, S, C=O, C=S, S=O, SO₂, C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

[00049] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a
10 is an M-ring wherein ring M is an aromatic pyridine or pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$; and

[00050] R^3 and R^4 optionally join to form a ring of 6 or 7 atoms, where
15 the atoms in the ring are independently selected from Z^3 , Z^4 , C=O, C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.


[00051] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a
20 is an M-ring, wherein ring M is an aromatic pyridine or pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$; and

[00052] R^3 and R^4 optionally join to form a ring of 6 atoms, where the
25 atoms in the ring are independently selected from Z^3 , Z^4 , C=O, C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

[00053] In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a
30 is an M-ring wherein ring M is an aromatic pyridine or pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is

carbon, M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$; and

R^3 and R^4 optionally join to form a ring that is selected from:



5 **[00054]** In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring, wherein ring M is an aromatic pyridine ring, wherein M^1 , M^3 , M^4 and M^6 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 is nitrogen.

10 **[00055]** In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring, wherein ring M is an aromatic pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are nitrogen.

15 **[00056]** In another embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring wherein ring M is an aromatic pyridine ring, wherein:

M^1 , M^3 , M^4 and M^6 are carbon and are substituted with $(L)_nR^1$;

M^5 is carbon;

20 M^2 is nitrogen;

R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl- R^{11} , C_1 - C_6 alkoxy- R^{11} , COR^{17} , COR_6 , CO_2R_6 , $CONHR_6$, $N(R^8)_2$, amino C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkyl, amino, amino C_1 - C_4 alkyl- R^7 , C_1 - C_6 alkyl-NHR⁷, carbonitrile, SR^{10} , halo, NHR⁷, NR^8R^9 , NHR⁷- C_1 - C_6 alkyl, NR^8R^9 - C_1 - C_6 alkyl, nitro, cyano, $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , C_1 - C_6 alkyl- COR^{11} , halo C_1 - C_4 alkyl, aryl, heteroaryl,

heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, or C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹²;

5 R⁷, R⁸, are each independently selected from -H, C₁-C₆ alkyl, C₁-C₄ alkyl-R¹¹, C₁-C₆ alkyl-N(R¹³)₂, CO₂R¹⁶, COR¹⁷, aryl, and arylalkyl, wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R¹⁸;

10 R⁹, R¹⁰ are each independently selected from -H, hydroxyl, C₁-C₆ alkyl, C₁-C₆ alkyl-R¹⁷, C₁-C₆ alkyl-NH₂R¹³, CO₂R¹⁶, COR¹⁷, C₁-C₆ alkyl-CO₂R¹⁶, C₁-C₆ alkyl-CONH-R¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, hydroxy C₁-C₄ alkyl, halo C₁-C₄ alkoxy, halo C₁-C₄ alkyl, Si(R¹³)₂R¹⁷, aryl, heteroaryl, heterocyclyl, arylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with
15 one or more of the groups defined by R¹⁸;

 R¹¹ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxyl, halo, amino, NHR¹³, N(R¹³)₂, COR¹³, CO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or
20 more of the groups defined by R¹⁸;

 R¹² is selected from -H, hydroxyl, oxo, C₁-C₆ alkyl, hydroxyl C₁-C₆ alkyl-R¹¹, C₁-C₁₀ alkoxy, amino, amino C₁-C₄ alkyl-R⁷, NHR⁷, N(R⁷)₂, C₁-C₆ alkyl-NHR⁷, C₁-C₆ alkyl-NHR⁸R⁹, C₁-C₆ alkyl-N(R⁸)₂, C₁-C₆ alkyl-R¹¹, C₁-C₆ alkyl-CO₂R⁷R¹¹, C₁-C₆ alkoxy-R¹¹, nitro, O-R¹⁰, C=O, COR¹¹,
25 CO₂R¹¹, SR¹⁰, SOR¹¹, SO₂R¹¹, NHSO₂R¹¹, C₁-C₆ alkyl-SR¹⁰, halo, halo C₁-C₄ alkyl, halo C₁-C₄ alkoxy, hydroxy C₁-C₄ alkyl, hydroxy C₁-C₄ alkoxy, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclylalkyl, and C₁-C₁₀
30 mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R^{13} and R^{14} are each independently selected from -H, oxo, C_1 - C_6 alkyl, COR^{23} , and aryl;

5 R^{15} , R^{16} are each independently selected from -H, aryl, arylalkyl, wherein aryl, arylalkyl, are optionally substituted with one or more of the groups defined by R^{24} ;

R^{17} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{19} , NHR^{19} , aryl, heteroarylalkyl, and heterocyclalkyl, wherein aryl is optionally substituted with one or more of the groups defined by R^{24} ;

10 R^{18} is selected from -H, oxo, hydroxyl, C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, amino, amino C_1 - C_6 alkyl, $N(R^{19})_2$, C_1 - C_6 alkyl- $N(R^{19})_2$, CO_2R^{23} , SR^{21} , halo, halo C_1 - C_4 alkyl, aryl, heteroaryl, and heterocycl, wherein aryl, heteroaryl, and heterocycl, are optionally substituted with one or more of the groups defined by R^{24} ;

15 R^{19} and R^{20} are each independently selected from -H, C_1 - C_6 alkyl, heteroaryl, heterocycl, wherein aryl, heteroaryl, and heterocycl, are optionally substituted with one or more of the groups defined by R^{30} ;

R^{21} and R^{22} are each independently selected from -H and C_1 - C_6 alkyl;

R^{23} is selected from -H and C_1 - C_6 alkyl;

20 R^{24} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, CO_2R^{29} , halo, and halo C_1 - C_4 alkyl;

R^{29} is selected from -H, and C_1 - C_6 alkyl;

25 R^{30} is selected from -H, aryl, heteroaryl, heterocycl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocycl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{36} ;

R^{36} is selected from -H and halo; and

R^2 , R^3 , R^4 , R^{37} and R^{38} are each independently selected from an R^1 group.

30 **[00057]** In an embodiment of the present invention, the MK-2 inhibiting compound is one having the structure of formula I, except that R^a is an M-ring wherein ring M is an aromatic pyrimidine ring, wherein:

M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$;

M^5 is carbon;

M^2 and M^6 are nitrogen;

R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl- R^{11} , C_1 - C_6 alkoxy- R^{11} , COR^{17} , COR_6 , CO_2R_6 , $CONHR_6$, $N(R^8)_2$, amino C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkyl, amino, amino C_1 - C_4 alkyl- R^7 , halo C_1 - C_4 alkyl, C_1 - C_6 alkyl- NHR^7 , carbonitrile, SR^{10} , halo, NHR^7 , NR^8R^9 , NHR^7 - C_1 - C_6 alkyl, NR^8R^9 - C_1 - C_6 alkyl, nitro, cyano, $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , C_1 - C_6 alkyl- COR^{11} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, or C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ;

R^7 , R^8 , are each independently selected from -H, C_1 - C_6 alkyl, C_1 - C_4 alkyl- R^{11} , C_1 - C_6 alkyl- $N(R^{13})_2$, CO_2R^{16} , COR^{17} , aryl, and arylalkyl, wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

R^9 , R^{10} are each independently selected from -H, hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{17} , C_1 - C_6 alkyl- NH_2R^{13} , CO_2R^{16} , COR^{17} , C_1 - C_6 alkyl- CO_2R^{16} , C_1 - C_6 alkyl- $CONH-R^{16}$, C_1 - C_6 alkyl- $CON(R^{16})_2$, hydroxy C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, halo C_1 - C_4 alkyl, $Si(R^{13})_2R^{17}$, aryl, heteroaryl, heterocyclyl, arylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

R^{11} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxyl, halo, amino, NHR^{13} , $N(R^{13})_2$, COR^{13} , CO_2R^{17} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

R^{12} is selected from -H, hydroxyl, oxo, C_1 - C_6 alkyl, hydroxyl C_1 - C_6 alkyl- R^{11} , C_1 - C_{10} alkoxy, amino, amino C_1 - C_4 alkyl- R^7 , NHR^7 , $N(R^7)_2$, C_1 - C_6 alkyl- NHR^7 , C_1 - C_6 alkyl- NHR^8R^9 , C_1 - C_6 alkyl- $N(R^8)_2$, C_1 - C_6 alkyl- R^{11} ,

C_1-C_6 alkyl- $CO_2R^7R^{11}$, C_1-C_6 alkoxy- R^{11} , nitro, $O-R^{10}$, $C=O$, COR^{11} ,
 CO_2R^{11} , SR^{10} , SOR^{11} , SO_2R^{11} , $NHSO_2R^{11}$, C_1-C_6 alkyl- SR^{10} , halo, halo C_1-
 C_4 alkyl, halo C_1-C_4 alkoxy, hydroxy C_1-C_4 alkyl, hydroxy C_1-C_4 alkoxy,
 aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclalkyl,
 5 and C_1-C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl,
 heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclalkyl, and C_1-C_{10}
 mono- and bicyclic cycloalkyl are optionally substituted with one or more of
 the groups defined by R^{18} ;

R^{13} and R^{14} are each independently selected from -H, oxo, C_1-C_6
 10 alkyl, COR^{23} , and aryl;

R^{15} , R^{16} are each independently selected from -H, aryl, arylalkyl,
 wherein aryl, arylalkyl, are optionally substituted with one or more of the
 groups defined by R^{24} ;

R^{17} is selected from -H, C_1-C_6 alkyl, C_1-C_6 alkyl- R^{19} , NHR^{19} , aryl,
 15 heteroarylalkyl, and heterocyclalkyl, wherein aryl is optionally substituted
 with one or more of the groups defined by R^{24} ;

R^{18} is selected from -H, oxo, hydroxyl, C_1-C_{10} alkyl, C_1-C_{10} alkoxy,
 amino, amino C_1-C_6 alkyl, $N(R^{19})_2$, C_1-C_6 alkyl- $N(R^{19})_2$, CO_2R^{23} , SR^{21} ,
 halo, halo C_1-C_4 alkyl, aryl, heteroaryl, and heterocyclyl, wherein aryl,
 20 heteroaryl, and heterocyclyl, are optionally substituted with one or more of
 the groups defined by R^{24} ;

R^{19} and R^{20} are each independently selected from -H, C_1-C_6 alkyl,
 heteroaryl, heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are
 optionally substituted with one or more of the groups defined by R^{30} ;

R^{21} and R^{22} are each independently selected from -H and C_1-C_6
 25 alkyl;

R^{23} is selected from -H and C_1-C_6 alkyl;

R^{24} is selected from -H, C_1-C_6 alkyl, C_1-C_6 alkoxy, CO_2R^{29} , halo,
 and halo C_1-C_4 alkyl;

R^{29} is selected from -H, and C_1-C_6 alkyl;

R^{30} is selected from -H, aryl, heteroaryl, heterocyclyl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{36} ;


R^{36} is selected from -H and halo; and

5 R^2 , R^3 , R^4 , R^{37} and R^{38} are each independently selected from an R^1 group.

[00058] In another embodiment, the present MK-2 inhibiting compound has the structure shown in formula III:

Formula III:

10



wherein:

dashed lines indicate optional single or double bonds;

15 Z^2 and Z^3 are nitrogen, Z^1 , Z^4 and Z^5 are carbon, and join with Z^2 and Z^3 to form a pyrazole ring;

where dashed lines indicate optional single or double bonds;

M^1 , M^3 and M^4 is carbon and is substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 and M^6 is independently selected from nitrogen and carbon, and if carbon, it is unsubstituted or substituted with $(L)_nR^1$;

20 R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl- R^{11} , C_1 - C_6 alkoxy- R^{11} , COR^{17} , CO_2R^7 , $CONHR^7$, $N(R^8)_2$, amino C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkyl, amino, amino C_1 - C_4 alkyl- R^7 , halo C_1 - C_4 alkyl, C_1 - C_6 alkyl-NHR⁷, C_1 - C_6 alkyl- $N(R^7)_2$, carbonitrile, SR^{10} , halo, NHR^7 , NR^8R^9 , NHR^7 - C_1 - C_6 alkyl, NR^8R^9 -
25 C_1 - C_6 alkyl, nitro, cyano, $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , C_1 - C_6 alkyl- COR^{11} , aryl,

heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, or C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹²;

R⁷ and R⁸ are each independently selected from -H, C₁-C₆ alkyl, C₁-C₄ alkyl-R¹¹, C₁-C₆ alkyl-N(R¹³)₂, CO₂R¹⁶, COR¹⁷, aryl, and arylalkyl, wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R¹⁸;

R⁹ and R¹⁰ are each independently selected from -H, hydroxyl, C₁-C₆ alkyl, C₁-C₆ alkyl-R¹⁷, C₁-C₆ alkyl-NH₂R¹³, CO₂R¹⁶, COR¹⁷, C₁-C₆ alkyl-CO₂R¹⁶, C₁-C₆ alkyl-CONH-R¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, hydroxy C₁-C₄ alkyl, halo C₁-C₄ alkoxy, halo C₁-C₄ alkyl, Si(R¹³)₂R¹⁷, aryl, heteroaryl, heterocyclyl, arylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with one or more of the groups defined by R¹⁸;

R¹¹ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, hydroxyl, halo, amino, NHR¹³, N(R¹³)₂, COR¹³, CO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or more of the groups defined by R¹⁸;

R¹² is selected from -H, hydroxyl, oxo, C₁-C₆ alkyl, hydroxyl C₁-C₆ alkyl-R¹¹, C₁-C₁₀ alkoxy, amino, amino C₁-C₄ alkyl-R⁷, NHR⁷, N(R⁷)₂, C₁-C₆ alkyl-NHR⁷, C₁-C₆ alkyl-NHR⁸R⁹, C₁-C₆ alkyl-N(R⁸)₂, C₁-C₆ alkyl-R¹¹, C₁-C₆ alkyl-CO₂R⁷R¹¹, C₁-C₆ alkoxy-R¹¹, nitro, O-R¹⁰, C=O, COR¹¹, CO₂R¹¹, SR¹⁰, SOR¹¹, SO₂R¹¹, NHSO₂R¹¹, C₁-C₆ alkyl-SR¹⁰, halo, halo C₁-C₄ alkyl, halo C₁-C₄ alkoxy, hydroxy C₁-C₄ alkyl, hydroxy C₁-C₄ alkoxy, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R^{13} and R^{14} are each independently selected from -H, oxo, C_1 - C_6 alkyl, COR^{23} , and aryl;

5 R^{15} and R^{16} are each independently selected from -H, aryl, arylalkyl, wherein aryl, arylalkyl, are optionally substituted with one or more of the groups defined by R^{24} ;

R^{17} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{19} , NHR^{19} , aryl, heteroarylalkyl, and heterocyclalkyl, wherein aryl is optionally substituted with one or more of the groups defined by R^{24} ;

10 R^{18} is selected from -H, oxo, hydroxyl, C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, amino, amino C_1 - C_6 alkyl, $N(R^{19})_2$, C_1 - C_6 alkyl- $N(R^{19})_2$, CO_2R^{23} , SR^{21} , halo, halo C_1 - C_4 alkyl, aryl, heteroaryl, and heterocycl, wherein aryl, heteroaryl, and heterocycl, are optionally substituted with one or more of the groups defined by R^{24} ;

15 R^{19} and R^{20} are each independently selected from -H, C_1 - C_6 alkyl, heteroaryl, heterocycl, wherein aryl, heteroaryl, and heterocycl, are optionally substituted with one or more of the groups defined by R^{30} ;

R^{21} and R^{22} are each independently selected from -H and C_1 - C_6 alkyl;

R^{23} is selected from -H and C_1 - C_6 alkyl;

20 R^{24} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, CO_2R^{29} , halo, and halo C_1 - C_4 alkyl;

R^{29} is selected from -H, and C_1 - C_6 alkyl;


25 R^{30} is selected from -H, aryl, heteroaryl, heterocycl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocycl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{36} ;

R^{36} is selected from -H and halo;

R^2 , R^3 , R^4 , R^{37} and R^{38} are each independently selected from an R^1 group;

n is 0; and

30 R^3 and R^4 optionally join to form a ring structure that is selected from:



[00059] The MK-2 inhibiting compounds that are described in formulas I-III, and in Tables I and II can be made by the methods that are described in the Examples below. Compounds that are not described specifically in the Examples can be made by reference to the methods used in the Examples, but with the substitution of starting compounds that are suitable for the compound that is desired.

[00060] The present invention also includes a method of inhibiting mitogen activated protein kinase-activated protein kinase-2, the method comprising contacting a mitogen activated protein kinase-activated protein kinase-2 with one or more of any of the MK-2 inhibiting compounds described herein. In one embodiment, the contacting of MK-2 with an MK-2 inhibitory compound takes place inside a cell. The cell can be one of any type of organism, but is preferably an animal cell. Contacting can occur *in vitro* or *in vivo*, and the cell can be a living cell, or it can be non-living. When the contacting is carried out *in vitro*, the cell can be attached to other cells, or it can be a single cell, or clump of cells in suspension or on a solid medium. When the contacting is carried out *in vivo*, the MK-2 inhibitory compound can be administered as described below.

[00061] In one embodiment, the present invention provides a method for treating or preventing an MK-2 modulated disease or disorder in a subject, the method comprises contacting a mitogen activated protein kinase-activated protein kinase-2 in a subject with one or more of the MK-2 inhibiting compounds that are described herein. A preferred MK-2 inhibiting compound for the present method is one having the structure

described by formula I. In another preferred embodiment, the MK-2 inhibiting compound is one having the structure described by formula II.

[00062] The present invention also includes a method of inhibiting mitogen activated protein kinase-activated protein kinase-2 in a subject in need of such inhibition, the method comprising administering to the subject one or more of the MK-2 inhibiting compounds described herein.

[00063] The present invention also includes a method of preventing or treating a TNF α mediated disease or disorder in a subject, the method comprising administering to the subject an effective amount of one or more of the MK-2 inhibiting compounds described herein. In a preferred embodiment, the subject is one that is in need of such prevention or treatment.

[00064] The present methods can be practiced by the administration of any one or more of the present MK-2 inhibiting compounds. It is preferred that the MK-2 inhibiting compound is one having an MK-2 IC₅₀ of less than about 10 μ M, in an *in vitro* assay of MK-2 inhibitory activity, more preferred is a compound having an MK-2 IC₅₀ of less than about 1.0 μ M, yet more preferred is a compound having an MK-2 IC₅₀ of less than about 0.5 μ M.

[00065] It should be understood that the base forms, salts, pharmaceutically acceptable salts, and prodrugs of the compounds that are described herein, as well as isomeric forms, tautomers, racemic mixtures of the compounds, and the like, which have the same or similar activity as the compounds that are described, are to be considered to be included within the description of the compound.

[00066] The MK-2 inhibiting activity of any of the compounds described herein can be determined by any one of several methods that are well known to those having skill in the art of enzyme activity testing. One such method is described in detail in the general methods section of the examples. In addition, the efficacy of any one of the present MK-2 inhibiting compounds in therapeutic applications can be determined by

testing for inhibition of TNF α production in cell culture and in animal model assays. In general, it is preferred that the MK-2 inhibiting compounds of the present invention be capable of inhibiting the production and/or the release of TNF α in cell cultures and in animal models.

5 **[00067]** In the present method, the MK-2 inhibiting compounds that are described herein can be used as inhibitors of MAPKAP kinase-2. When this inhibition is for a therapeutic purpose, one or more of the present MK-2 inhibitory compounds can be administered to a subject that is in need of MK-2 inhibition. As used herein, a "subject in need of MK-2 inhibition" is a
10 subject who has, or who is at risk of contracting a TNF α mediated disease or disorder. TNF α mediated diseases and disorders are described in more detail below.

15 **[00068]** As described above, in an embodiment of the present method, a subject in need of prevention or treatment of a TNF α mediated disease or disorder is treated with one or more of the present MK-2 inhibiting compounds. In one embodiment, the subject is treated with an effective amount of the MK-2 inhibiting compound. The effective amount can be an amount that is sufficient for preventing or treating the TNF α mediated disease or disorder.

20 **[00069]** The MK-2 inhibiting compound that is used in the subject method can be any MK-2 inhibiting compound that is described herein.

25 **[00070]** In the subject method, the MK-2 inhibiting compound can be used in any amount that is an effective amount. It is preferred, however, that the amount of the MK-2 inhibiting compound that is administered is within a range of about 0.1 mg/day per kilogram (kg) of the subject to about 1500 mg/day/kg. It is more preferred that the amount of the compound is within a range of about 1 mg/day/kg to about 500 mg/day/kg. An amount that is within a range of about 10 mg/day/kg to about 400 mg/day/kg, is even more preferred.

30 **[00071]** When the term "about" is used herein in relation to a dosage amount of the MK-2 inhibiting compound, it is to be understood to mean an

amount that is within $\pm 10\%$ by weight of the amount or range that is described. By way of example, "about 0.1 - 10 mg/day" includes all dosages within 0.9 to 11 mg/day.

5 [00072] In an embodiment of the present invention, a therapeutic composition is provided that contains at least one of the MK-2 inhibiting compounds that are described herein. A preferred therapeutic composition contains a therapeutically effective amount of a compound that is described by formula I. In another embodiment, a preferred therapeutic composition is one having an MK-2 inhibiting compound that is described by formula II.

10 [00073] In another embodiment of the present invention, a pharmaceutical composition that contains one or more of the present MK-2 inhibitors can be administered to a subject for the prevention or treatment of a TNF α mediated disease or disorder. The pharmaceutical composition includes an MK-2 inhibitor of the present invention and a pharmaceutically acceptable carrier. A preferred MK-2 inhibitor for use in the pharmaceutical composition is described by formula I. In another embodiment, a preferred pharmaceutical composition is one having an MK-2 inhibiting compound that is described by formula II.

20 [00074] In another embodiment, a kit can be produced that is suitable for use in the prevention or treatment of a TNF α mediated disease or disorder. The kit comprises a dosage form comprising at least one of the MK-2 inhibitors that is described herein in an amount which comprises a therapeutically effective amount.

25 [00075] As used herein, an "effective amount" means the dose or effective amount to be administered to a patient and the frequency of administration to the subject which is readily determined by one or ordinary skill in the art, by the use of known techniques and by observing results obtained under analogous circumstances. The dose or effective amount to be administered to a patient and the frequency of administration to the subject can be readily determined by one of ordinary skill in the art by the use of known techniques and by observing results obtained under

30

analogous circumstances. In determining the effective amount or dose, a number of factors are considered by the attending diagnostician, including but not limited to, the potency and duration of action of the compounds used, the nature and severity of the illness to be treated, as well as the sex, age, weight, general health and individual responsiveness of the patient to be treated, and other relevant circumstances.

[00076] The phrase "therapeutically-effective" indicates the capability of an agent to prevent, or improve the severity of, the disorder, while avoiding adverse side effects typically associated with alternative therapies. The phrase "therapeutically-effective" is to be understood to be equivalent to the phrase "effective for the treatment, prevention, or inhibition", and both are intended to qualify the amount of the MK-2 inhibitory compound for use in therapy which will achieve the goal of improvement in the severity of pain and inflammation and the frequency of incidence over treatment, while avoiding adverse side effects typically associated with alternative therapies.

[00077] Those skilled in the art will appreciate that dosages may also be determined with guidance from Goodman & Goldman's The Pharmacological Basis of Therapeutics, Ninth Edition (1996), Appendix II, pp. 1707-1711.

[00078] The frequency of dose will depend upon the half-life of the active components of the composition. If the active molecules have a short half life (e.g. from about 2 to 10 hours) it may be necessary to give one or more doses per day. Alternatively, if the active molecules have a long half-life (e.g. from about 2 to about 15 days) it may only be necessary to give a dosage once per day, per week, or even once every 1 or 2 months. A preferred dosage rate is to administer the dosage amounts described above to a subject once per day.

[00079] Daily dosages can vary within wide limits and will be adjusted to the individual requirements in each particular case. In general, for administration to adults, an appropriate daily dosage has been described above, although the limits that were identified as being preferred may be

exceeded if expedient. The daily dosage can be administered as a single dosage or in divided dosages.

[00080] For the purposes of calculating and expressing a dosage rate, all dosages that are expressed herein are calculated on an average amount-per-day basis irrespective of the dosage rate. For example, one 100 mg dosage of an MK-2 inhibitor taken once every two days would be expressed as a dosage rate of 50 mg/day. Similarly, the dosage rate of an ingredient where 50 mg is taken twice per day would be expressed as a dosage rate of 100 mg/day.

[00081] For purposes of calculation of dosage amounts, the weight of a normal adult human will be assumed to be 70 kg.

[00082] When the MK-2 inhibitor is supplied along with a pharmaceutically acceptable carrier, the pharmaceutical compositions that are described above can be formed. Pharmaceutically acceptable carriers include, but are not limited to, physiological saline, Ringer's, phosphate solution or buffer, buffered saline, and other carriers known in the art. Pharmaceutical compositions may also include stabilizers, anti-oxidants, colorants, and diluents. Pharmaceutically acceptable carriers and additives are chosen such that side effects from the pharmaceutical compound are minimized and the performance of the compound is not canceled or inhibited to such an extent that treatment is ineffective.

[00083] The term "pharmacologically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by a researcher or clinician. This amount can be a therapeutically effective amount.

[00084] The term "pharmaceutically acceptable" is used herein to mean that the modified noun is appropriate for use in a pharmaceutical product. Pharmaceutically acceptable cations include metallic ions and organic ions. More preferred metallic ions include, but are not limited to, appropriate alkali metal salts, alkaline earth metal salts and other physiological acceptable metal ions. Exemplary ions include aluminum,

calcium, lithium, magnesium, potassium, sodium and zinc in their usual valences. Preferred organic ions include protonated tertiary amines and quaternary ammonium cations, including in part, trimethylamine, diethylamine, *N,N*-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (*N*-methylglucamine) and procaine. Exemplary pharmaceutically acceptable acids include, without limitation, hydrochloric acid, hydroiodic acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulfonic acid, acetic acid, formic acid, tartaric acid, maleic acid, malic acid, citric acid, isocitric acid, succinic acid, lactic acid, gluconic acid, glucuronic acid, pyruvic acid, oxalacetic acid, fumaric acid, propionic acid, aspartic acid, glutamic acid, benzoic acid, and the like.

[00085] Also included in the compounds and compositions of the invention are the isomeric forms and tautomers and the pharmaceutically-acceptable salts of the present MK-2 inhibitors. Illustrative pharmaceutically acceptable salts are prepared from formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, stearic, salicylic, *p*-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, cyclohexylaminosulfonic, algenic, β -hydroxybutyric, galactaric and galacturonic acids.

[00086] Suitable pharmaceutically-acceptable base addition salts of compounds of the present invention include metallic ion salts and organic ion salts. More preferred metallic ion salts include, but are not limited to, appropriate alkali metal (Group IA) salts, alkaline earth metal (Group IIA) salts and other physiological acceptable metal ions. Such salts can be made from the ions of aluminum, calcium, lithium, magnesium, potassium, sodium and zinc. Preferred organic salts can be made from tertiary amines and quaternary ammonium salts, including in part, trifluoroacetate, trimethylamine, diethylamine, *N,N*-dibenzylethylenediamine,

chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (*N*-methylglucamine) and procaine. All of the above salts can be prepared by those skilled in the art by conventional means from the corresponding compound of the present invention.

5 **[00087]** The method of the present invention is useful for, but not limited to, the prevention and/or treatment of diseases and disorders that are mediated by TNF α and/or mediated by MK-2, including pain, inflammation and/or arthritis. For example, the compounds described herein would be useful for the treatment of any inflammation-related disorder described
10 below, such as an analgesic in the treatment of pain and headaches, or as an antipyretic for the treatment of fever. The compounds described herein would also be useful for the treatment of an inflammation-related disorder in a subject suffering from such an inflammation-associated disorder.

15 **[00088]** As used herein, the terms "treating", "treatment", "treated", or "to treat," mean to alleviate symptoms, eliminate the causation either on a temporary or permanent basis. The term "treatment" includes alleviation, elimination of causation of pain and/or inflammation associated with, but not limited to, any of the diseases or disorders described herein. The terms "prevent", "prevention", "prevented", or "to prevent," mean to prevent
20 or to slow the appearance of symptoms associated with, but not limited to, any of the diseases or disorders described herein.

25 **[00089]** In preferred embodiments, the methods and compositions of the present invention encompass the prevention and/or treatment of pain, inflammation and inflammation-related disorders.

30 **[00090]** In other preferred embodiments, the methods and compositions of the present invention encompass the treatment of any one or more of the disorders selected from the group consisting of connective tissue and joint disorders, neoplasia disorders, cardiovascular disorders, otic disorders, ophthalmic disorders, respiratory disorders, gastrointestinal disorders, angiogenesis-related disorders, immunological disorders, allergic disorders, nutritional disorders, infectious diseases and disorders, endocrine disorders, metabolic disorders, neurological and

neurodegenerative disorders, psychiatric disorders, hepatic and biliary disorders, musculoskeletal disorders, genitourinary disorders, gynecologic and obstetric disorders, injury and trauma disorders, surgical disorders, dental and oral disorders, sexual dysfunction disorders, dermatologic disorders, hematological disorders, and poisoning disorders.

[00091] As used herein, the terms "neoplasia" and "neoplasia disorder", used interchangeably herein, refer to new cell growth that results from a loss of responsiveness to normal growth controls, *e.g.* to "neoplastic" cell growth. Neoplasia is also used interchangeably herein with the term "cancer" and for purposes of the present invention; cancer is one subtype of neoplasia. As used herein, the term "neoplasia disorder" also encompasses other cellular abnormalities, such as hyperplasia, metaplasia and dysplasia. The terms neoplasia, metaplasia, dysplasia and hyperplasia can be used interchangeably herein and refer generally to cells experiencing abnormal cell growth.

[00092] Both of the terms, "neoplasia" and "neoplasia disorder", refer to a "neoplasm" or tumor, which may be benign, premalignant, metastatic, or malignant. Also encompassed by the present invention are benign, premalignant, metastatic, or malignant neoplasias. Also encompassed by the present invention are benign, premalignant, metastatic, or malignant tumors. Thus, all of benign, premalignant, metastatic, or malignant neoplasia or tumors are encompassed by the present invention and may be referred to interchangeably, as neoplasia, neoplasms or neoplasia-related disorders. Tumors are generally known in the art to be a mass of neoplasia or "neoplastic" cells. Although, it is to be understood that even one neoplastic cell is considered, for purposes of the present invention to be a neoplasm or alternatively, neoplasia.

[00093] In still other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the connective tissue and joint disorders selected from the group consisting of arthritis, rheumatoid arthritis, spondyloarthropathies, gouty arthritis, lumbar spondylarthrosis, carpal tunnel syndrome, canine

hip dysplasia, systemic lupus erythematosus, juvenile arthritis, osteoarthritis, tendonitis and bursitis.

[00094] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the neoplasia disorders selected from the group consisting of acral lentiginous melanoma, actinic keratoses, adenocarcinoma, adenoid cystic carcinoma, adenomas, familial adenomatous polyposis, familial polyps, colon polyps, polyps, adenosarcoma, adenosquamous carcinoma, adrenocortical carcinoma, AIDS-related lymphoma, anal cancer, astrocytic tumors, bartholin gland carcinoma, basal cell carcinoma, bile duct cancer, bladder cancer, brain stem glioma, brain tumors, breast cancer, bronchial gland carcinomas, capillary carcinoma, carcinoids, carcinoma, carcinosarcoma, cavernous, central nervous system lymphoma, cerebral astrocytoma, cholangiocarcinoma, chondrosarcoma, choroid plexus papilloma/carcinoma, clear cell carcinoma, skin cancer, brain cancer, colon cancer, colorectal cancer, cutaneous T-cell lymphoma, cystadenoma, endodermal sinus tumor, endometrial hyperplasia, endometrial stromal sarcoma, endometrioid adenocarcinoma, ependymal, epithelioid, esophageal cancer, Ewing's sarcoma, extragonadal germ cell tumor, fibrolamellar, focal nodular hyperplasia, gallbladder cancer, gastrinoma, germ cell tumors, gestational trophoblastic tumor, glioblastoma, glioma, glucagonoma, hemangioblastomas, hemangioendothelioma, hemangiomas, hepatic adenoma, hepatic adenomatosis, hepatocellular carcinoma, Hodgkin's lymphoma, hypopharyngeal cancer, hypothalamic and visual pathway glioma, insulinoma, intraepithelial neoplasia, interepithelial squamous cell neoplasia, intraocular melanoma, invasive squamous cell carcinoma, large cell carcinoma, islet cell carcinoma, Kaposi's sarcoma, kidney cancer, laryngeal cancer, leiomyosarcoma, lentigo maligna melanomas, leukemia-related disorders, lip and oral cavity cancer, liver cancer, lung cancer, lymphoma, malignant mesothelial tumors, malignant thymoma, medulloblastoma, medulloepithelioma, melanoma, meningeal, merkel cell

carcinoma, mesothelial, metastatic carcinoma, mucoepidermoid carcinoma, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndrome, myeloproliferative disorders, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, neuroepithelial adenocarcinoma nodular melanoma, non-Hodgkin's lymphoma, oat cell carcinoma, oligodendroglial, oral cancer, oropharyngeal cancer, osteosarcoma, pancreatic polypeptide, ovarian cancer, ovarian germ cell tumor, pancreatic cancer, papillary serous adenocarcinoma, pineal cell, pituitary tumors, plasmacytoma, pseudosarcoma, pulmonary blastoma, parathyroid cancer, penile cancer, pheochromocytoma, pineal and supratentorial primitive neuroectodermal tumors, pituitary tumor, plasma cell neoplasm, pleuropulmonary blastoma, prostate cancer, rectal cancer, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, serous carcinoma, small cell carcinoma, small intestine cancer, soft tissue carcinomas, somatostatin-secreting tumor, squamous carcinoma, squamous cell carcinoma, submesothelial, superficial spreading melanoma, supratentorial primitive neuroectodermal tumors, thyroid cancer, undifferentiated carcinoma, urethral cancer, uterine sarcoma, uveal melanoma, verrucous carcinoma, vaginal cancer, vipoma, vulvar cancer, Waldenstrom's macroglobulinemia, well differentiated carcinoma, and Wilm's tumor.

[00095] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the cardiovascular disorders selected from the group consisting of myocardial ischemia, hypertension, hypotension, heart arrhythmias, pulmonary hypertension, hypokalemia, cardiac ischemia, myocardial infarction, cardiac remodeling, cardiac fibrosis, myocardial necrosis, aneurysm, arterial fibrosis, embolism, vascular plaque inflammation, vascular plaque rupture, bacterial-induced inflammation and viral induced inflammation, edema, swelling, fluid accumulation, cirrhosis of the liver, Bartter's syndrome, myocarditis, arteriosclerosis, atherosclerosis, calcification (such as vascular calcification and valvar calcification), coronary artery disease,

heart failure, congestive heart failure, shock, arrhythmia, left ventricular hypertrophy, angina, diabetic nephropathy, kidney failure, eye damage, vascular diseases, migraine headaches, aplastic anemia, cardiac damage, diabetic cardiac myopathy, renal insufficiency, renal injury, renal arteriopathy, peripheral vascular disease, left ventricular hypertrophy, cognitive dysfunction, stroke, and headache.

[00096] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the metabolic disorders selected from the group consisting of obesity, overweight, type I and type II diabetes, hypothyroidism, and hyperthyroidism.

[00097] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the respiratory disorders selected from the group consisting of asthma, bronchitis, chronic obstructive pulmonary disease (COPD), cystic fibrosis, pulmonary edema, pulmonary embolism, pneumonia, pulmonary sarcoisosis, silicosis, pulmonary fibrosis, respiratory failure, acute respiratory distress syndrome and emphysema.

[00098] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the angiogenesis-related disorders selected from the group consisting of angiofibroma, neovascular glaucoma, arteriovenous malformations, arthritis, osler-weber syndrome, atherosclerotic plaques, psoriasis, corneal graft neovascularization, pyogenic granuloma, delayed wound healing, retrolental fibroplasias, diabetic retinopathy, scleroderma, granulations, solid tumors, hemangioma, trachoma, hemophilic joints, vascular adhesions, hypertrophic scars, age-related macular degeneration, coronary artery disease, stroke, cancer, AIDS complications, ulcers and infertility.

[00099] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the infectious diseases and disorders selected from the group consisting of

viral infections, bacterial infections, prion infections, spirochetes infections, mycobacterial infections, rickettsial infections, chlamydial infections, parasitic infections and fungal infections.

5 **[000100]** In still further embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the infectious diseases and disorders selected from the group consisting of hepatitis, HIV (AIDS), small pox, chicken pox, common cold, bacterial influenza, viral influenza, warts, oral herpes, genital herpes, herpes simplex infections, herpes zoster, bovine spongiform encephalopathy, 10 septicemia, streptococcus infections, staphylococcus infections, anthrax, severe acquired respiratory syndrome (SARS), malaria, African sleeping sickness, yellow fever, chlamydia, botulism, canine heartworm, rocky mountain spotted fever, lyme disease, cholera, syphilis, gonorrhea, encephalitis, pneumonia, conjunctivitis, yeast infections, rabies, dengue 15 fever, Ebola, measles, mumps, rubella, West Nile virus, meningitis, gastroenteritis, tuberculosis, hepatitis, and scarlet fever.

[000101] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the neurological and neurodegenerative disorders selected from the group 20 consisting of headaches, migraine headaches, Alzheimer's disease, Parkinson's disease, dementia, memory loss, senility, amyotrophy, ALS, amnesia, seizures, multiple sclerosis, muscular dystrophies, epilepsy, schizophrenia, depression, anxiety, attention deficit disorder, hyperactivity, bulimia, anorexia nervosa, anxiety, autism, phobias, spongiform 25 encephalopathies, Creutzfeldt-Jakob disease, Huntington's Chorea, ischemia, obsessive-compulsive disorder, manic depression, bipolar disorders, drug addiction, alcoholism and smoking addiction.

[000102] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the 30 dermatological disorders selected from the group consisting of acne, psoriasis, eczema, burns, poison ivy, poison oak and dermatitis.

[000103] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the surgical disorders selected from the group consisting of pain and swelling following surgery, infection following surgery and inflammation following surgery.

[000104] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the gastrointestinal disorders selected from the group consisting of inflammatory bowel disease, irritable bowel syndrome, Crohn's disease, gastritis, irritable bowel syndrome, diarrhea, constipation, dysentery, ulcerative colitis, gastric esophageal reflux, gastric ulcers, gastric varices, ulcers, and heartburn.

[000105] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the otic disorders selected from the group consisting of otic pain, inflammation, otorrhea, otalgia, fever, otic bleeding, Lermoyez's syndrome, Meniere's disease, vestibular neuronitis, benign paroxysmal positional vertigo, herpes zoster oticus, Ramsay Hunt's syndrome, viral neuronitis, ganglionitis, geniculate herpes, labyrinthitis, purulent labyrinthitis, viral endolymphatic labyrinthitis, perilymph fistulas, noise-induced hearing loss, presbycusis, drug-induced ototoxicity, acoustic neuromas, aerotitis media, infectious myringitis, bullous myringitis, otitis media, otitis media with effusion, acute otitis media, secretory otitis media, serous otitis media, acute mastoiditis, chronic otitis media, otitis externa, otosclerosis, squamous cell carcinoma, basal cell carcinoma, nonchromaffin paragangliomas, chemodectomas, globus jugulare tumors, globus tympanicum tumors, external otitis, perichondritis, aural eczematoid dermatitis, malignant external otitis, subperichondrial hematoma, ceruminomas, impacted cerumen, sebaceous cysts, osteomas, keloids, otalgia, tinnitus, vertigo, tympanic membrane infection, typanitis, otic furuncles, otorrhea, acute mastoiditis, petrositis, conductive and sensorineural hearing loss, epidural abscess, lateral sinus thrombosis,

subdural empyema, otitic hydrocephalus, Dandy's syndrome, bullous myringitis, cerumen-impacted, diffuse external otitis, foreign bodies, keratosis obturans, otic neoplasm, otomycosis, trauma, acute barotitis media, acute eustachian tube obstruction, post-otic surgery, postsurgical otalgia, cholesteatoma, conductive and sensorineural hearing loss, epidural abscess, lateral sinus thrombosis, subdural empyema and otitic hydrocephalus.

[000106] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of the ophthalmic disorders selected from the group consisting of retinopathies, uveitis, ocular photophobia, acute injury to the eye tissue, conjunctivitis, age-related macular degeneration diabetic retinopathy, detached retina, glaucoma, vitelliform macular dystrophy type 2, gyrate atrophy of the choroid and retina, conjunctivitis, corneal infection, fuchs' dystrophy, iridocorneal endothelial syndrome, keratoconus, lattice dystrophy, map-dot-fingerprint dystrophy, ocular herpes, pterygium, myopia, hyperopia, and cataracts.

[000107] In other preferred embodiments, the methods and compositions of the present invention encompass the prevention and treatment of menstrual cramps, kidney stones, minor injuries, wound healing, vaginitis, candidiasis, sinus headaches, tension headaches, dental pain, periarteritis nodosa, thyroiditis, myasthenia gravis, multiple sclerosis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, swelling occurring after injury, closed head injury, liver disease, and endometriosis.

[000108] As used herein, the terms "TNF α mediated disease or disorder" are meant to include, without limitation, each of the symptoms or diseases that are mentioned above.

[000109] The term "subject" for purposes of treatment includes any human or animal subject who is in need of the prevention of or treatment of any one of the TNF α mediated diseases or disorders. The subject is typically a mammal. "Mammal", as that term is used herein, refers to any

animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cattle, etc., Preferably, the mammal is a human.

5 **[000110]** For methods of prevention, the subject is any human or animal subject, and preferably is a subject that is in need of prevention and/or treatment of a $\text{TNF}\alpha$ mediated diseases or disorders. The subject may be a human subject who is at risk of obtaining a $\text{TNF}\alpha$ mediated disease or disorder, such as those described above. The subject may be at risk due to genetic predisposition, sedentary lifestyle, diet, exposure to disorder-
10 causing agents, exposure to pathogenic agents and the like.

[000111] The subject pharmaceutical compositions may be administered enterally and parenterally. Parenteral administration includes subcutaneous, intramuscular, intradermal, intramammary, intravenous, and other administrative methods known in the art. Enteral administration
15 includes solution, tablets, sustained release capsules, enteric coated capsules, and syrups. When administered, the pharmaceutical composition may be at or near body temperature.

[000112] In particular, the pharmaceutical compositions of the present invention can be administered orally, for example, as tablets, coated
20 tablets, dragees, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or
25 more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These
30 excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch, or alginic

acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

[000113] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredients are mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredients are present as such, or mixed with water or an oil medium, for example, peanut oil, liquid paraffin, or olive oil.

[000114] Aqueous suspensions can be produced that contain the MK-2 inhibitors in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinylpyrrolidone gum tragacanth and gum acacia; dispersing or wetting agents may be naturally-occurring phosphatides, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate.

[000115] The aqueous suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, or one or more sweetening agents, such as sucrose or saccharin.

[000116] Oily suspensions may be formulated by suspending the active ingredients in an omega-3 fatty acid, a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol.

[000117] Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

[000118] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above.

Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

[000119] Syrups and elixirs containing the novel MK-2 inhibitory compounds may be formulated with sweetening agents, for example glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

[000120] The subject compositions can also be administered parenterally, either subcutaneously, or intravenously, or intramuscularly, or intrasternally, or by infusion techniques, in the form of sterile injectable aqueous or olagenous suspensions. Such suspensions may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above or other acceptable agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending

medium. For this purpose, any bland fixed oil may be employed including synthetic mono-, or di-, glycerides. In addition, n-3 polyunsaturated fatty acids may find use in the preparation of injectables.

5 [000121] The subject compositions can also be administered by inhalation, in the form of aerosols or solutions for nebulizers, or rectally, in the form of suppositories prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperature but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and poly-ethylene glycols.

10 [000122] The novel compositions can also be administered topically, in the form of creams, ointments, jellies, collyriums, solutions or suspensions.

[000123] Various delivery systems include capsules, tablets, and gelatin capsules, for example.

15 [000124] The following examples describe preferred embodiments of the invention. Other embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered to be exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples. In the examples all percentages are
20 given on a weight basis unless otherwise indicated.

GENERAL INFORMATION FOR PREPARATION METHODS:

[000125] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers.

25 [000126] NMR analysis:

[000127] Proton nuclear magnetic resonance spectra were obtained on a Varian Unity Innova 400, a Varian Unity Innova 300 a Varian Unity 300, a Bruker AMX 500 or a Bruker AV-300 spectrometer. Chemical shifts are given in ppm (δ) and coupling constants, J , are reported in Hertz.

30 Tetramethylsilane was used as an internal standard for proton spectra and the solvent peak was used as the reference peak for carbon spectra.

Mass spectra were obtained on a Perkin Elmer Sciex 100 atmospheric

pressure ionization (APCI) mass spectrometer, a Finnigan LCQ Duo LCMS ion trap electrospray ionization (ESI) mass spectrometer, a PerSeptive Biosystems Mariner TOF HPLC-MS (ESI), or a Waters ZQ mass spectrometer (ESI).

5 **[000128]** Determination of MK-2 IC₅₀:

[000129] Recombinant MAPKAPK2 was phosphorylated at a concentration of 42-78 μ M by incubation with 0.23 μ M of active p38 α in 50 mM HEPES, 0.1 mM EDTA, 10 mM magnesium acetate, and 0.25 mM ATP, pH 7.5 for one hour at 30°C.

10 **[000130]** The phosphorylation of HSP-peptide (KKKALSRQLSVAA) by MAPKAPK2 was measured using an anion exchange resin capture assay method. The reaction was carried out in 50 mM β -glycerolphosphate, 0.04 % BSA, 10 mM magnesium acetate, 2% DMSO and 0.8 mM dithiotheritol, pH 7.5 in the presence of the HSP-peptide with 0.2 μ Ci [γ ³³P]ATP and
15 0.03mM ATP. The reaction was initiated by the addition of 15 nM MAPKAPK2 and was allowed to incubate at 30°C for 30 min. The reaction was terminated and [γ ³³P]ATP was removed from solution by the addition of 150 μ l of AG 1X8 ion exchange resin in 900 mM sodium formate pH 3.0. A 50 μ l aliquot of head volume was removed from the quenched reaction
20 mixture and added to a 96-well plate, 150 μ l of Microscint-40 (Packard) was added and the amount of phosphorylated-peptide was determined. Allow the Microscint to sit in the plates for 60 minutes prior to counting.

[000131] Compounds are evaluated as potential inhibitors of the MK2 kinase by measuring their effects on MK2 phosphorylation of the peptide
25 substrate. Compounds may be screened initially at two concentrations prior to determination of IC₅₀ values. Screening results are expressed as percent inhibition at the concentrations of compound tested. For IC₅₀ value determinations, compounds are tested at six concentrations in ten-fold serial dilutions with each concentration tested in triplicate. Results are
30 expressed as IC₅₀ values in micromolar. The assay is performed at a final concentration of 2% DMSO.

[000132] U937 Cell TNF α release assay

[000133] The human monocyte-like cell line, U937 (ATCC #CRL-1593.2), is cultured in RPMI 1640 media with 10% heat-inactivated fetal calf serum (GIBCO), glutamine and pen/strep at 37°C and 5% CO₂. Differentiation of U937 to monocytic/macrophage-like cells is induced by the addition of phorbol 12-myristate 13-acetate (Sigma) at final concentration of 20 ng/ml to a culture of U937 cells at ~0.5 million cells/ml and incubated for 24 hrs. The cells are centrifuged, washed with PBS and resuspended in fresh media without PMA and incubated for 24 hrs. Cells adherent to the culture flask are harvested by scraping, centrifugation, and resuspended in fresh media to 2 million cells/ml, and 0.2 ml is aliquoted to each of 96 wells in flat-bottom plate. Cells are then incubated for an additional 24 hrs to allow for recovery. The media is removed from the cells, and 0.1 ml of fresh media is added per well. 0.05 ml of serially diluted compound or control vehicle (Media with DMSO) is added to the cells. The final DMSO concentration does not exceed 1%. After 1hr incubation, 0.05 ml of 400ng/ml LPS (E Coli serotype 0111:B4, Sigma) in media is added for final concentration of 100 ng/ml. Cells are incubated at 37°C for 4 hrs. After 4hrs incubation, supernatants are harvest and assayed by ELISA for the presence of TNF α .

[000134] U937 cell TNF α ELISA

[000135] ELISA plates (NUNC-Immuno™ Plate Maxisorb™ Surface) were coated with purified mouse monoclonal IgG1 anti-human TNF α antibody (R&D Systems #MAB610; 1.25 ug/ml in sodium bicarbonate pH 8.0, 0.1 ml/well) and incubated at 4°C. Coating solution was aspirated the following day and wells were blocked with 1 mg/ml gelatin in PBS (plus 1x thimerasol) for 2 days at 4°C. Prior to using, wells were washed 3x with wash buffer (PBS with 0.05% Tween). Cultured media samples were diluted in EIA buffer (5 mg/ml bovine γ -globulin, 1 mg/ml gelatin, 1 ml/l Tween-20, 1 mg/ml thimerasol in PBS), added to wells (0.1 ml/well) in triplicate and allowed to incubate for 1.5 hr at 37°C in a humidified

chamber. Plates were again washed and 0.1 ml/well of a mixture of rabbit anti-human TNF α polyclonal antibodies in EIA buffer (1:400 dilution of Sigma #T8300, and 1:400 dilution of Calbiochem #654250) was added for 1 hr at 37°C. Plates were washed as before and peroxidase-conjugated goat anti-rabbit IgG (H+L) antibody (Jackson ImmunoResearch #111-035-144, 1 ug/ml in EIA buffer, 0.1 ml/well) was added for 45 min. After final washing, plates were developed with peroxidase-ABTS solution (Kirkegaard/Perry #50-66-01, 0.1 ml/well). Enzymatic conversion of ABTS to colored product was measured after 5-30 minutes using a SpectroMax 340 spectrophotometer (Molecular Devices) at 405 nm. TNF levels were quantitated from a recombinant human TNF α (R&D Systems #210-TA-010) standard curve using a quadratic parameter fit generated by SoftMaxPRO software. ELISA sensitivity was approximately 30 pg TNF/ml. IC₅₀ values for compounds were generated using BioAssay Solver.

[000136] Lipopolysaccharide (LPS)-Induced TNF α Production.

[000137] Adult male 225-250 gram Lewis rats (Harlan Sprague-Dawley) were used. Rats were fasted 18 hr prior to oral dosing, and allowed free access to water throughout the experiment. Each treatment group consisted of 5 animals.

[000138] Compounds were prepared as a suspension in a vehicle consisting of 0.5% methylcellulose, 0.025% Tween-20 in PBS. Compounds or vehicle were orally administered in a volume of 1 ml using an 18 gauge gavage needle. LPS (*E. coli* serotype 0111:B4, Lot #39H4103, Cat. # L-2630, Sigma) was administered 1-4 hr later by injection into the penile vein at a dose of 1 mg/kg in 0.5 ml sterile saline. Blood was collected in serum separator tubes via cardiac puncture 1.5 hr after LPS injection, a time point corresponding to maximal TNF α production. After clotting, serum was withdrawn and stored at -20°C until assay by ELISA (described below).

[000139] Rat LPS TNF α ELISA

[000140] ELISA plates (NUNC-Immuno™ Plate Maxisorb™ Surface) were coated with 0.1 ml per well of a Protein G purified fraction of a 2.5 ug/ml of hamster anti-mouse/rat TNF α monoclonal antibody TN19.12 (2.5 ug/ml in PBS, 0.1 ml/well). The hybridoma cell line was kindly provided by Dr. Robert Schreiber, Washington University. Wells were blocked the following day with 1 mg/ml gelatin in PBS. Serum samples were diluted in a buffer consisting of 5 mg/ml bovine γ -globulin, 1 mg/ml gelatin, 1 ml/l Tween-20, 1 mg/ml thimerasol in PBS, and 0.1 ml of diluted serum was added wells in duplicate and allowed to incubate for 2 hr at 37°C. Plates were washed with PBS-Tween, and 0.1 ml per well of a 1:300 dilution of rabbit anti-mouse/rat TNF α antibody (BioSource International, Cat. #AMC3012) was added for 1.5 hr at 37°C. Plates were washed, and a 1:1000 fold dilution of peroxidase-conjugated donkey anti-rabbit IgG antibody (Jackson ImmunoResearch, Cat. #711-035-152) was added for 45 min. After washing, plates were developed with 0.1 ml of ABTS-peroxide solution (Kirkegaard/Perry, Cat. #50-66-01). Enzymatic conversion of ABTS to colored product was measured after ~30 minutes using a SpectroMax 340 spectrophotometer (Molecular Devices Corp.) at 405 nm. TNF levels in serum were quantitated from a recombinant rat TNF α (BioSource International, Cat. #PRC3014) standard curve using a quadratic parameter fit generated by SoftMaxPRO software. ELISA sensitivity was approximately 30 pg TNF/ml. Results are expressed in percent inhibition of the production of TNF α as compared to blood collected from control animals dosed only with vehicle.

[000141] NMR analysis:

[000142] Proton nuclear magnetic resonance spectra were obtained on a Varian Unity Innova 400, a Varian Unity Innova 300 a Varian Unity 300, a Bruker AMX 500 or a Bruker AV-300 spectrometer. Chemical shifts are given in ppm (δ) and coupling constants, J , are reported in Hertz. Tetramethylsilane was used as an internal standard for proton spectra and

the solvent peak was used as the reference peak for carbon spectra. Mass spectra were obtained on a Perkin Elmer Sciex 100 atmospheric pressure ionization (APCI) mass spectrometer, a Finnigan LCQ Duo LCMS ion trap electrospray ionization (ESI) mass spectrometer, a
5 PerSeptive Biosystems Mariner TOF HPLC-MS (ESI), or a Waters ZQ mass spectrometer (ESI).

[000143] Determination of MK-2 IC₅₀:

[000144] Recombinant MAPKAPK2 was phosphorylated at a concentration of 42-78 μM by incubation with 0.23 μM of active p38 α in 50
10 mM HEPES, 0.1 mM EDTA, 10 mM magnesium acetate, and 0.25 mM ATP, pH 7.5 for one hour at 30°C.

[000145] The phosphorylation of HSP-peptide (KKKALSRQLSVAA) by MAPKAPK2 was measured using an anion exchange resin capture assay method. The reaction was carried out in 50 mM β -glycerolphosphate, 0.04
15 % BSA, 10 mM magnesium acetate, 2% DMSO and 0.8 mM dithiotheritol, pH 7.5 in the presence of the HSP-peptide with 0.2 μCi [γ ³³P]ATP and 0.03mM ATP. The reaction was initiated by the addition of 15 nM MAPKAPK2 and was allowed to incubate at 30°C for 30 min. The reaction was terminated and [γ ³³P]ATP was removed from solution by the addition
20 of 150 μl of AG 1X8 ion exchange resin in 900 mM sodium formate pH 3.0. A 50 μl aliquot of head volume was removed from the quenched reaction mixture and added to a 96-well plate, 150 μl of Microscint-40 (Packard) was added and the amount of phosphorylated-peptide was determined. Allow the Microscint to sit in the plates for 60 minutes prior to counting.

[000146] Compounds are evaluated as potential inhibitors of the MK-2 kinase by measuring their effects on MK-2 phosphorylation of the peptide substrate. Compounds may be screened initially at two concentrations prior to determination of IC₅₀ values. Screening results are expressed as percent inhibition at the concentrations of compound tested. For IC₅₀
25 value determinations, compounds are tested at six concentrations in ten-fold serial dilutions with each concentration tested in triplicate. Results are
30

expressed as IC₅₀ values in micromolar. The assay is performed at a final concentration of 2% DMSO.

[000147] U937 Cell TNF α release assay

[000148] The human monocyte-like cell line, U937 (ATCC #CRL-1593.2), is cultured in RPMI1640 media with 10% heat-inactivated fetal calf serum (GIBCO), glutamine and pen/strep at 37°C and 5% CO₂. Differentiation of U937 to monocytic/macrophage-like cells is induced by the addition of

phorbol12-myristate 13-acetate (Sigma) at final concentration of 20 ng/ml to a culture of U937 cells at ~0.5 million cells/ml and incubated for 24 hrs. The cells are centrifuged, washed with PBS and resuspended in fresh media without PMA and incubated for 24 hrs. Cells adherent to the culture flask are harvested by scraping, centrifugation, and resuspended in fresh media to 2 million cells/ml, and 0.2 ml is aliquoted to each of 96 wells in flat-bottom plate. Cells are then incubated for an additional 24 hrs to allow for recovery. The media is removed from the cells, and 0.1 ml of fresh media is added per well. 0.05 ml of serially diluted compound or control vehicle (Media with DMSO) is added to the cells. The final DMSO concentration does not exceed 1%. After 1hr incubation, 0.05 ml of 400ng/ml LPS (E Coli serotype 0111:B4, Sigma) in media is added for final concentration of 100 ng/ml. Cells are incubated at 37°C for 4 hrs. After 4hrs incubation, supernatants are harvest and assayed by ELISA for the presence of TNF α .

[000149] U937 cell TNF α ELISA

[000150] ELISA plates (NUNC-Immuno™ Plate Maxisorb™ Surface)

were coated with purified mouse monoclonal IgG1 anti-human TNF α antibody (R&D Systems #MAB610; 1.25 μ g/ml in sodium bicarbonate pH 8.0, 0.1 ml/well) and incubated at 4°C. Coating solution was aspirated the following day and wells were blocked with 1 mg/ml gelatin in PBS (plus 1x thimerasol) for 2 days at 4°C. Prior to using, wells were washed 3x with wash buffer (PBS with 0.05% Tween). Cultured media samples were diluted in EIA buffer (5 mg/ml bovine gamma-globulin, 1 mg/ml gelatin, 1 ml/l Tween-20, 1 mg/ml thimerasol in PBS), added to wells (0.1 ml/well) in

triplicate and allowed to incubate for 1.5 hr at 37°C in a humidified chamber. Plates were again washed and 0.1 ml/well of a mixture of rabbit anti-human TNF α polyclonal antibodies in EIA buffer (1:400 dilution of Sigma #T8300, and 1:400 dilution of Calbiochem #654250) was added for 1 hr at 37°C. Plates were washed as before and peroxidase-conjugated goat anti-rabbit IgG (H+L) antibody (Jackson ImmunoResearch #111-035-144, 1 ug/ml in EIA buffer, 0.1 ml/well) was added for 45 min. After final washing, plates were developed with peroxidase-ABTS solution (Kirkegaard/Perry #50-66-01, 0.1 ml/well). Enzymatic conversion of ABTS to colored product was measured after 5-30 minutes using a SpectroMax 340 spectrophotometer (Molecular Devices) at 405 nm. TNF levels were quantitated from a recombinant human TNF α (R&D Systems #210-TA-010) standard curve using a quadratic parameter fit generated by SoftMaxPRO software. ELISA sensitivity was approximately 30 pg TNF/ml. IC₅₀ values for compounds were generated using BioAssay Solver.

[000151] Lipopolysaccharide (LPS)-Induced TNF α Production.

[000152] Adult male 225-250 gram Lewis rats (Harlan Sprague-Dawley) were used. Rats were fasted 18 hr prior to oral dosing, and allowed free access to water throughout the experiment. Each treatment group consisted of 5 animals.

[000153] Compounds were prepared as a suspension in a vehicle consisting of 0.5% methylcellulose, 0.025% Tween-20 in PBS. Compounds or vehicle were orally administered in a volume of 1 ml using an 18 gauge gavage needle. LPS (E. coli serotype 0111:B4, Lot #39H4103, Cat. # L-2630, Sigma) was administered 1-4 hr later by injection into the penile vein at a dose of 1 mg/kg in 0.5 ml sterile saline. Blood was collected in serum separator tubes via cardiac puncture 1.5 hr after LPS injection, a time point corresponding to maximal TNF α production. After clotting, serum was withdrawn and stored at -20°C until assay by ELISA (described below).

[000154] Rat LPS TNF α ELISA

[000155] ELISA plates (NUNC-Immuno™ Plate Maxisorb™ Surface) were coated with 0.1 ml per well of a Protein G purified fraction of a 2.5 ug/ml of hamster anti-mouse/rat TNF α monoclonal antibody TN19.12 (2.5 ug/ml in PBS, 0.1 ml/well). The hybridoma cell line was provided by Dr. Robert Schreiber, Washington University. Wells were blocked the following day with 1 mg/ml gelatin in PBS. Serum samples were diluted in a buffer consisting of 5 mg/ml bovine gamma-globulin, 1 mg/ml gelatin, 1 ml/l Tween-20, 1 mg/ml thimerasol in PBS, and 0.1 ml of diluted serum was added wells in duplicate and allowed to incubate for 2 hr at 37°C. Plates were washed with PBS-Tween, and 0.1 ml per well of a 1:300 dilution of rabbit anti-mouse/rat TNF α antibody (BioSource International, Cat. #AMC3012) was added for 1.5 hr at 37°C. Plates were washed, and a 1:1000 fold dilution of peroxidase-conjugated donkey anti-rabbit IgG antibody (Jackson ImmunoResearch, Cat. #711-035-152) was added for 45 min. After washing, plates were developed with 0.1 ml of ABTS-peroxide solution (Kirkegaard/Perry, Cat. #50-66-01). Enzymatic conversion of ABTS to colored product was measured after ~30 minutes using a SpectroMax 340 spectrophotometer (Molecular Devices Corp.) at 405 nm. TNF levels in serum were quantitated from a recombinant rat TNF α (BioSource International, Cat. #PRC3014.) standard curve using a quadratic parameter fit generated by SoftMaxPRO software. ELISA sensitivity was approximately 30 pg TNF/ml. Results are expressed in percent inhibition of the production of TNF α as compared to blood collected from control animals dosed only with vehicle.

[000156] Synthesis of MK-2 inhibiting compounds of the present invention:

[000157] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Proton nuclear magnetic resonance spectra were obtained on a Varian-300, Bruker AMX 500 or a Bruker AV-300 spectrometer. Spectra are given in ppm (δ) and coupling constants, J ,

are reported in Hertz. Tetramethylsilane was used as an internal standard for proton spectra and the solvent peak was used as the reference peak for carbon spectra. 3-Bromobenzotrifluoride was used as an internal standard in ^{19}F nuclear magnetic resonance spectra to calibrate the amount of TFA for TFA salts. Mass spectra were obtained on a Mariner electrospray ionization (ESI), Perkin Elmer Sciex 100 atmospheric pressure ionization (APCI) mass spectrometer, Hewlett Packard G1947A LCMS ion trap ionization (ESI), or a Finnigan LCQ Duo LCMS ion trap electrospray ionization (ESI) mass spectrometer. Thin-layer chromatography (TLC) was performed using Analtech silica gel plates and visualized by ultraviolet (UV) light. HPLC analyses were obtained using a YMC CombiScreen ODS-A (50 x 4.6 mm) with UV with diode array detection, using aqueous TFA in acetonitrile as the eluent on a Hewlett Packard 1100, or a Phenomenex C18 Luna column (150 x 4.6 mm) with UV detection at 254 nm, using aqueous TFA in acetonitrile as the eluent on a Varian Prostar. Purification using preparative HPLC was performed using a Phenomenex C18 Luna column (250 x 22 mm) with UV detection at 254 nm, using aqueous TFA in acetonitrile as the eluent on a Varian Prostar, or using a Waters Deltapack C 18 column (47 x 300 mm) with UV detection at 254 nm, using aqueous TFA in acetonitrile as the eluent on a Gilson. All reactions were carried out under nitrogen unless specified.

EXAMPLE 1

[000158] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride.

[000159] Step 1. The preparation of lithium (1Z)-4-ethoxy-3,4-dioxo-1-pyridin-4-ylbut-1-en-1-olate.

[000160] 4-Acetylpyridine (960.4 g, 7.93 mole) and diethyl oxylate (1170.8 g, 8.08 mole) were dissolved in 8 L toluene in a 22 L reactor. The reaction was cooled to -78°C under N_2 , and a 1 M solution of Lithium bis(trimetnylsilyl)amide in THF (8.08 L, 8.08 mole) was added in a medium stream over 45 minutes, keeping the temperature near 10°C . When the addition was complete, the reaction was allowed to warm to

room temperature with stirring for 18 hours. The reaction mixture was filtered, and the solid washed with toluene followed by diethyl ether. The product was air dried and desiccated to afford 1562.5 g of the lithium salt of the diketoester (87% yield).

5 **[000161]** Step 2. Lithium (1Z)-4-ethoxy-3,4-dioxo-1-pyridin-4-ylbut-1-en-1-olate (1562.5 g, 6.88 mole) was placed in a 22 L reactor and slurried in 7 L of ethanol. The slurry was heated to 63°C under N₂ with mixing. The heating mantle was removed, and the temperature stabilized at about 63°C. To this mixture hydrazine monohydrochloride (476.3 g, 6.95) was

10 added. After a few minutes a slow exotherm started. An ice/water bath was applied when the reaction reached 80°C. The ice bath was removed, and the temperature held steady at 80°C. The heating mantle was re-installed, and the reaction was maintained at 80.5°C (reflux) for one hour. The reaction was cooled and stirred at room temperature for 18 hours.

15 The mixture was filtered, and the solids washed with ethanol followed by diethyl ether. The product was air dried and desiccated to afford 1365.6 g (91% yield) of the desired pyrazole as a tan solid. LCMS showed a single peak with m/z 218 (M+H). ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.61 (d, 2H), 7.83 (d, 2H), 7.44 (s, 1H), 4.31 (q, 2H), 1.30 (t, 3H).

20 **[000162]** Step 3. To a cooled (0°C) solution of ethyl 3-pyridin-4-yl-1H-pyrazole-5-carboxylate (5.57 g, 25.6 mmol) in anhydrous DMF (140 mL) was added lithium *t*-butoxide (1M in THF, 38.5 mL) dropwise. The reaction stirred for 30 min, then a solution of *tert*-butyl 2-bromoethylcarbamate (8.62 g, 38.5 mmol) and sodium iodide (5.77 g, 38.5

25 mmol) in anhydrous DMF (25 mL) was added dropwise. The reaction was allowed to stir and warm to room temperature for 20 h. The reaction was poured into water and brine, extracted with ethyl acetate, dried over MgSO₄, and concentrated to an orange solid. The solid was rinsed with diethyl ether to afford an off-white solid (5.49 g, 59.5% yield): ¹H NMR

30 (DMSO-d₆ / 300 MHz) δ 8.59 (d, 2H), 7.82 (d, 2H), 7.51 (s, 1H), 6.90 (t, 1H), 4.58 (t, 2H), 4.33 (q, 2H), 3.38-3.33 (m, 2H), 1.34 (t, 3H), 1.27 (s, 9H); HRMS calculated for (M + H) 361.1870, found 361.1890.

EXAMPLE 2

[000163] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride.

5 [000164] To a flask charged with ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-pyridin-4-yl-1H-pyrazole-5-carboxylate (5.39 g, 15.0 mmol) was added 4N HCl/dioxane (20 mL). After 1 h the reaction mixture was filtered and rinsed with diethyl ether to afford an off-white solid (5.02 g, 100%
10 yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 8.93 (d, 2H), 8.48-8.43 (m, 5H), 7.95 (s, 1H), 4.89 (t, 2H), 4.37 (q, 2H), 3.41-3.38 (m, 2H), 1.35 (t, 3H); HRMS calculated for (M + H) 261.1346, found 261.1317.

EXAMPLE 3

[000165] This example illustrates the production of 2-pyridin-4-yl-6,7-dihydro-pyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

15 [000166] A flask was charged with ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride (1.13 g, 3.39 mmol), NH_4OH (30 mL), and ethanol (15 mL). After stirring for 1 h, the reaction mixture was purified by Gilson RP HPLC (5-95% acetonitrile/water). The appropriate fractions were concentrated to a pale yellow solid (0.725 g,
20 65.4% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 8.81 (d, 2H), 8.40 (br s, 1H), 8.22 (d, 2H), 7.67 (s, 1H), 4.43 (t, 2H), 3.70-3.64 (m, 2H); HRMS calculated for (M + H) 215.0927, found 215.0885.

EXAMPLE 4

[000167] This example illustrates the production of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate.

25 [000168] A solution of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride (0.794 g, 2.38 mmol) and $\text{LiOH}\cdot\text{H}_2\text{O}$ (0.310 g, 7.40 mmol) in 30 mL of THF/ H_2O (1:1) was heated to 80°C with stirring. After 2 h the reaction mixture was purified by Gilson RP HPLC (5-95% acetonitrile/water). The appropriate fractions were concentrated to a white
30 solid (0.442 g, 53.7% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 8.96 (br s,

2H), 8.48 (br s, 2H), 8.07-7.91 (m, 4H), 4.87 (br s, 2H), 3.41 (br s, 2H); HRMS calculated for (M + H) 233.1033, found 233.1025.

EXAMPLE 5

5 [000169] This example illustrates the production of 1-(2-([3-(5-methyl-2-furyl)butyl]amino)ethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate.

[000170] Ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride (0.806 g, 2.42 mmol) was neutralized by stirring with morpholino-methylpolystyrene resin (4.15 g, ~3.5 mmol base/g resin) in 10 dichloromethane/methanol (20:1) for 1 h. The resin was filtered, rinsed with methanol, and the filtrates concentrated. The resulting white residue was dissolved in dichloromethane/methanol (20:1). Six drops of glacial acetic acid were added with stirring, followed by 3-(5-methyl-2-furyl)butyraldehyde (0.422 g, 6.60 mmol. After 5 min, sodium 15 triacetoxyborohydride (1.04 g, 4.90 mmol) was added. The reaction stirred for 1 h, both the mono- and dialkylated products formed. The reaction was quenched with water and extracted with dichloromethane. The organic layers were concentrated to an oil, which was subjected to hydrolysis conditions [3 eq LiOH•H₂O, THF/H₂O (1:1)] for 3 h. The resulting product 20 mixture was purified by Gilson RP HPLC (5-95% acetonitrile/water) and the fractions corresponding to the monoalkylated product were concentrated to a mauve solid (0.106 g, 9.0% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.77-8.75 (m, 4H), 8.08 (d, 2H), 7.72 (s, 1H), 5.97-5.93 (m, 2H), 4.87 (t, 2H), 3.51-3.47 (m, 2H), 2.96 (br s, 2H), 2.82 (q, 1H), 2.19 (s, 25 3H), 1.92-1.72 (m, 2H), 1.16 (d, 3H); HRMS calculated for (M + H) 233.1033, found 233.1025.

EXAMPLE 6

[000171] This example illustrates the production of ethyl 3-(1-oxidopyridin-4-yl)-1H-pyrazole-5-carboxylate.

30 [000172] A flask was charged with ethyl 3-pyridin-4-yl-1H-pyrazole-5-carboxylate (5.04 g, 23.2 mmol) and 3-chloroperoxybenzoic acid (7.06g, 27.8 mmol) in 70 mL of dichloromethane. After 2 h the reaction was

concentrated *in vacuo* to remove most of the dichloromethane. To the residue was added 5% ethyl acetate/hexanes and the suspension filtered. The solid was slurried in NaHCO₃ (sat.), filtered, and rinsed with NaHCO₃ (sat.). The solid was then slurried in NaS₂O₃, filtered, and then rinsed with water. The solid was slurried up in ethanol and concentrated to afford an orange-tan solid (4.25 g, 78.0% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.24 (d, 2H), 7.86 (d, 2H), 7.42 (s, 1H), 4.31 (q, 2H), 1.31 (t, 3H); HRMS calculated for (M + H) 234.0873, found 234.0877.

EXAMPLE 7

[000173] This example illustrates the production of ethyl 3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate.

[000174] A suspension of ethyl 3-(1-oxidopyridin-4-yl)-1H-pyrazole-5-carboxylate (11.16 g, 14.9 mmol) and phosphorus oxychloride (110 mL, 1.2 mol) was heated to 106°C for 48 h. The reaction mixture was concentrated, and then dissolved in chloroform (300 mL) and ice water (300 mL). Solid NaHCO₃ was added until foaming was no longer observed, then the heterogeneous solution was extracted multiple times with chloroform. The organic layers were dried over MgSO₄, filtered, and concentrated. The residue was triturated with dichloromethane. The solid was filtered and rinsed with ethyl acetate, and then with diethyl ether to afford a pale yellow solid (3.59 g, 29.8% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 14.41 (br s, 1H), 8.44 (d, 1H), 7.98 (s, 1H), 7.88 (d, 1H), 7.61 (s, 1H), 4.33 (q, 2H), 1.32 (t, 3H); HRMS calculated for (M + H) 252.0534, found 252.0508.

EXAMPLE 8

[000175] This example illustrates the production of ethyl 1-{3-[(tert-butoxycarbonyl)-amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate.

[000176] Synthesis conducted as in the preparation of ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-pyridin-4-yl-1H-pyrazole-5-carboxylate using ethyl 3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (3.36 g, 13.3 mmol), lithium *t*-butoxide (1M in THF, 17.0 mL), *tert*-butyl 3-

bromopropylcarbamate (3.81 g, 16.0 mmol) and sodium iodide (2.39 g, 16.0 mmol). Chromatographic purification (25% ethyl acetate/hexane) afforded a white solid (3.29 g, 60.5% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 8.42 (d, 1H), 7.95 (s, 1H), 7.86 (dd, 1H), 7.67 (s, 1H), 6.85 (dd, 1H), 4.54 (t, 2H), 4.35 (q, 2H), 2.98-2.92 (m, 2H), 1.98-1.88 (m, 2H), 1.36-1.31 (m, 12H); HRMS calculated for (M + H) 409.1637, found 409.1634.

EXAMPLE 9

[000177] This example illustrates the production of ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate.

[000178] Synthesis was conducted as in the preparation of ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-pyridin-4-yl-1H-pyrazole-5-carboxylate, using ethyl 3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (6.00 g, 23.8 mmol), lithium *t*-butoxide (1M in THF, 31.0 mL), *tert*-butyl 2-bromoethylcarbamate (6.59 g, 29.4 mmol), and sodium iodide (4.41 g, 29.4 mmol). Flash chromatography (25% ethyl acetate/hexane) afforded a white solid (6.07 g, 64.6% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 8.43 (d, 1H), 7.94 (s, 1H), 7.85 (dd, 1H), 7.66 (s, 1H), 6.89 (t, 1H), 4.59 (t, 2H), 4.33 (q, 2H), 3.39-3.32 (m, 2H), 1.36-1.23 (m, 12H); HRMS calculated for (M + H) 395.1481, found 395.1512.

EXAMPLE 10

[000179] This example illustrates the production of 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000180] A flask was charged with ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (6.50 g, 15.9 mmol) and 4N HCl/dioxane (45 mL). The reaction mixture stirred for 1 h, then the suspension was filtered and the solid rinsed with diethyl ether. This solid was taken up in ethanol (20 mL) and NH_4OH (40 mL). After stirring for 20 h, the suspension was filtered and rinsed with ethanol and diethyl ether to afford a white solid (3.64 g, 87.4% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 8.41 (d, 1H), 8.35 (t, 1H), 7.90 (s, 1H), 7.83 (d, 1H), 7.48

(s, 1H), 4.49 (t, 2H), 3.24-3.18 (m, 2H), 2.20-2.14 (m, 2H), HRMS calculated for (M + H) 263.0694, found 263.0689.

EXAMPLE 11

5 [000181] This example illustrates the production of ethyl 1-{3-[(tert-butoxycarbonyl) amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate.

10 [000182] A flask was charged with ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.875 g, 2.14 mmol), 3-nitrophenylboronic acid (0.536 g, 3.21 mmol), 2M Na₂CO₃ (9 mL), toluene (25 mL), and [1, 1' bis(diphenylphosphino)ferrocene]dichloropalladium(II), 1:1 complex with dichloromethane [Pd(dppf)Cl₂·CH₂Cl₂ (0.140 g, 0.170 mmol)], then purged with N₂ and heated at 90°C for 20 h. The reaction mixture was diluted with ethyl acetate and washed with water and brine. The organic layer was dried
15 over MgSO₄, filtered, and concentrated. Chromatographic purification (30% ethyl acetate/hexane) afforded a white solid (0.767 g, 72.4% yield):
1H NMR (DMSO-d₆ / 300 MHz) δ 9.00 (dd, 1H), 8.75 (d, 1H), 8.68 (d, 1H), 8.54 (s, 1H), 8.30 (m, 1H), 7.91 (dd, 1H), 7.84-7.70 (m, 2H), 6.87 (dd, 1H), 4.58 (t, 2H), 4.36 (q, 2H), 3.00-2.95 (m, 2H), 1.98-1.92 (m, 2H), 1.38-1.31
20 (m, 12H); HRMS calculated for (M + H) 496.2191, found 496.2175.

EXAMPLE 12

[000183] This example illustrates the production of 1-(3-aminopropyl)-3-[2-(3 nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxamide trifluoroacetate.

25 [000184] Ammonia gas was bubbled into a pressure tube charged with ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate (0.084 g, 0.17 mmol) in 10 mL ethanol and cooled to -78 °C. The tube was then sealed and heated to 70°C for 6 days. The reaction mixture was concentrate *in vacuo*. To this residue was added 4N HCl/ dioxane (6 mL). After 1.5 h the solution was purified by
30 Gilson RP HPLC (5-95% acetonitrile/water). The appropriate fractions were concentrated to a light tan solid (0.048 g, 60% yield): 1H NMR

(DMSO-d₆ / 300 MHz) δ 8.94 (dd, 1H), 8.81 (d, 1H), 8.60 (d, 1H), 8.43 (s, 1H), 8.36 (dd, 1H), 8.20-8.09 (m, 4H), 7.87-7.74 (m, 3H), 4.66 (t, 2H), 2.86-2.79 (m, 2H), 2.20-2.15 (m, 2H); HRMS calculated for (M + H) 367.1513, found 367.1529.

5

EXAMPLE 13

[000185] This example illustrates the production of 1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxamide dihydrochloride.

[000186] Synthesis was conducted as for the production of 1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxamide trifluoroacetate using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate (0.127 g, 0.254 mmol). The TFA salt residue was taken up in methanol and 4N HCl/dioxane. After 20 h the suspension was concentrated to an off-white solid (0.069 g, 0.16 mmol): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.90 (d, 1H), 9.78 (s, 1H), 8.87 (d, 1H), 8.69 (s, 1H), 8.46-8.39 (m, 2H), 8.22-8.08 (m, 5H), 7.96-7.89 (m, 2H), 7.82-7.78 (m, 2H), 4.67 (t, 2H), 2.86-2.77 (m, 2H), 2.25-2.16 (m, 2H); HRMS calculated for (M + H) 373.1771, found 373.1769.

10

15

EXAMPLE 14

[000187] This example illustrates the production of ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate.

20

[000188] Synthesis was conducted as in the preparation of ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.732 g, 1.85 mmol), 3-quinolinylboronic acid (0.481 g, 2.78 mmol), 2M Na₂CO₃ (7 mL), toluene (20 mL), and Pd[dppf]Cl₂·CH₂Cl₂ (0.121 g, 0.148 mmol). The black residue obtained from the aqueous work-up was triturated with dichloromethane to afford an off-white solid (0.615 g, 68.2% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.72 (s, 1H), 9.14 (s, 1H), 8.78 (d, 1H), 8.62 (s, 1H), 8.15-8.08 (m,

25

30

2H), 7.90-7.82 (m, 3H), 7.71-7.66 (m, 1H), 6.94 (t, 1H), 4.64 (t, 2H), 4.37 (q, 2H), 3.39-3.32 (m, 2H), 1.39-1.28 (m, 12H); HRMS calculated for (M + H) 488.2292, found 488.2296.

EXAMPLE 15

5 **[000189]** This example illustrates the production of 2-{2-[4-(trifluoromethoxy)-phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

[000190] Synthesis was conducted as in the preparation of ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.954 g, 2.33 mmol), 3-quinolinyboronic acid (0.520 g, 3.00 mmol), 2M Na₂CO₃ (10 mL), toluene (25 mL), and Pd[dppf]Cl₂·CH₂Cl₂ (0.152 g, 0.186 mmol). Flash chromatography (25% ethyl acetate/hexane) afforded a beige solid (0.593g, 50.7% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.72 (d, 1H), 9.14 (d, 1H), 8.77 (d, 1H), 8.63 (s, 1H), 8.15-8.07 (m, 2H), 7.85-7.82 (m, 3H), 7.69 (dd, 1H), 6.88 (t, 1H), 4.60 (t, 2H), 4.37 (q, 2H), 3.02-2.96 (m, 2H), 2.01-1.93 (m, 2H), 1.39-1.34 (m, 12H); HRMS calculated for (M + H) 502.2449, found 502.2419.

20 **EXAMPLE 16**

[000191] This example illustrates the production of ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-{2-[4-(hydroxymethyl) phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate.

[000192] Synthesis was conducted as in the preparation of ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (1.100 g, 2.79 mmol), 4-hydroxymethyl-phenylboronic acid (0.550 g, 3.62 mmol), 2M Na₂CO₃ (12 mL), toluene (32 mL), and Pd[dppf]Cl₂·CH₂Cl₂ (0.182 g, 0.223 mmol). The black residue obtained from the aqueous work-up was triturated with dichloromethane to yield an off-white solid (0.567 g, 43.6% yield): ¹H NMR

(DMSO- d_6 / 300 MHz) δ 8.67 (d, 1H), 8.32 (s, 1H), 8.15 (d, 2H), 7.78-7.76 (m, 2H), 7.45 (d, 2H), 6.92 (t, 1H), 5.26 (t, 1H), 4.62-4.56 (m, 4H), 4.35 (q, 2H), 3.39-3.35 (m, 2H), 1.36-1.27 (m, 12H); HRMS calculated for (M + H) 467.2289, found 467.2259.

5

EXAMPLE 17

[000193] This example illustrates the production ethyl 1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride.

[000194] Synthesis was conducted as in the preparation of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate (0.763 g, 1.53 mmol) and 4N HCl/dioxane (10 mL). A light yellow solid was obtained (0.685 g, 95.4% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 9.00 (dd, 1H), 8.80 (d, 1H), 8.68 (d, 1H), 8.63 (s, 1H), 8.35 (dd, 1H), 8.13 (br s, 3H), 8.02 (dd, 1H), 7.93 (s, 2H), 7.85 (dd, 1H), 4.68 (t, 2H), 4.37 (q, 2H), 2.87-2.80 (m, 2H), 2.21-2.16 (m, 2H), 1.36 (t, 3H); HRMS calculated for (M + H) 396.1666, found 396.1660.

15

EXAMPLE 18

[000195] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride.

20

[000196] Synthesis was conducted as in the preparation of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate (0.588 g, 1.21 mmol) and 4N HCl/dioxane (12 mL). A white solid was obtained (0.557 g, 100% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 9.97 (s, 1H), 9.86 (s, 1H), 8.87-8.85 (m, 2H), 8.62 (s, 1H), 8.48-8.41 (m, 5H), 8.15-8.08 (m, 2H), 7.98-7.90 (m, 2H), 4.89 (t, 2H), 4.39 (q, 2H), 3.43-3.36 (m, 2H), 1.37 (t, 3H); HRMS calculated for (M + H) 388.1768, found 388.1754.

25

EXAMPLE 19

[000197] This example illustrates the production of ethyl 1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride.

5 [000198] Synthesis was conducted as for the preparation of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride, using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate (0.404 g, 0.805 mmol) and 4N HCl/dioxane (10 mL). A yellow solid was obtained (0.357g, 93.5% yield):
10 ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.92 (d, 1H), 9.72 (d, 1H), 8.84 (d, 1H), 8.79 (s, 1H), 8.40-8.36 (m, 2H), 8.16 (br s, 3H), 8.07-8.00 (m, 2H), 7.93-7.88 (m, 2H), 4.66 (t, 2H), 4.38 (q, 2H), 2.90-2.82 (m, 2H), 2.25-2.16 (m, 2H), 1.37 (t, 3H); HRMS calculated for (M + H) 402.1925, found 402.1937.

EXAMPLE 20

15 [000199] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride.

[000200] Synthesis was conducted as for the preparation of ethyl 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylate dihydrochloride,
20 using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate (0.498 g, 1.07 mmol) and 4N HCl/dioxane (10 mL). A white solid was obtained (0.474g, 100% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.83 (d, 1H), 8.74 (s, 1H), 8.38 (br s, 3H), 8.29-8.20 (m, 3H), 8.08 (s, 1H), 7.57 (d, 2H), 4.90 (t, 2H), 4.62 (m, 2H), 4.39 (q, 2H), 3.40-3.35 (m, 2H), 1.36 (t, 3H); HRMS
25 calculated for (M + H) 367.1765, found 367.1751.

EXAMPLE 21

[000201] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitro-phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate
30 dihydrochloride.

[000202] A flask was charged with ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.700 g, 1.78 mmol), 3-nitrophenylboronic acid (0.446 g, 2.67 mmol), 2M Na₂CO₃ (7 mL), toluene (20 mL), and [1, 1' bis(diphenylphosphino)ferrocene]dichloropalladium (II), 1:1 complex with dichloromethane [Pd(dppf)Cl₂·CH₂Cl₂ (0.116 g, 0.142 mmol)], then purged with N₂ and heated at 90°C for 20 h. The reaction mixture was filtered through Celite and the organic layer was concentrated. To this residue was added 4N HCl/dioxane (10.0 mL). After 2 h the reaction mixture was purified by Gilson RP HPLC (5-95% acetonitrile/water). The appropriate fractions were concentrated to a tacky residue, which was converted to the HCl salt using 4N HCl/dioxane and methanol. After stirring 1 h the reaction was concentrated to a brown solid (0.688 g, 85.1% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.01 (s, 1H), 8.80 (dd, 1H), 8.70-8.64 (m, 2H), 8.35-8.28 (m, 4H), 8.03 (d, 1H), 7.93 (s, 1H), 7.84 (dd, 1H), 4.87 (t, 2H), 4.38 (q, 2H), 3.39-3.33 (m, 2H), 1.36 (t, 3H); HRMS calculated for (M + H) 382.1510, found 382.1525.

EXAMPLE 22

[000203] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride.

[000204] Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.700 g, 1.78 mmol), 4-methoxyphenylboronic acid (0.406 g, 2.67 mmol), 2M Na₂CO₃ (7 mL), toluene (20 mL), Pd(dppf)Cl₂·CH₂Cl₂ (0.116 g, 0.142 mmol), and 4N HCl/dioxane (10.0 mL). The product was isolated as a yellow solid (0.709 g, 90.7% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.77 (d, 1H), 8.65 (s, 1H), 8.35 (br s, 3H), 8.24 (d, 2H), 8.16 (d, 1H), 8.04 (s, 1H), 7.18 (d, 2H),

4.89 (t, 2H), 4.38 (q, 2H), 3.87 (s, 3H), 3.42-3.38 (m, 2H), 1.36 (t, 3H),
HRMS calculated for (M + H) 367.1765, found 367.1790.

EXAMPLE 23

5 **[000205]** This example illustrates the production of Ethyl 1-(2-aminoethyl)-3-{2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride.

10 **[000206]** Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.700 g, 1.78 mmol), 4-trifluoromethoxyphenylboronic acid (0.550 g, 2.67 mmol), 2M Na₂CO₃ (7 mL), toluene (20 mL), Pd[dppf]Cl₂·CH₂Cl₂ (0.116 g, 0.142 mmol), and 4N HCl/dioxane (10.0 mL). The product was isolated as a reddish-tan solid (0.821 g, 93.5% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.79 (d, 1H), 8.59 (s, 1H), 8.36-8.33 (m, 5H), 8.07 (d, 1H), 7.94 (s, 1H), 7.57 (d, 2H), 4.87 (t, 2H), 4.38 (q, 2H), 3.42-3.38 (m, 2H), 1.36 (t, 3H); HRMS calculated for (M + H) 421.1482, found 421.1482.

EXAMPLE 24

20 **[000207]** This example illustrates the production of ethyl 1-(2-aminoethyl)-3-{2-[(E)-2-phenylethenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride.

25 **[000208]** Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride in using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (1.044 g, 2.64 mmol), *trans*-2-phenylvinylboronic acid (0.587 g, 3.97 mmol), 2M Na₂CO₃ (10 mL), toluene (30 mL), Pd[dppf]Cl₂·CH₂Cl₂ (0.172 g, 0.211 mmol), and 4N HCl/dioxane (10.0 mL). The product was isolated as a yellow solid (1.213 g, >100% yield, 73.0% pure): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.79-8.72 (m, 2H), 8.36 (br s, 3H), 8.24-8.19 (m, 2H), 7.96 (s, 1H), 7.71 (d, 2H),

30

7.62-7.45 (m, 5H), 4.89 (t, 2H), 4.39 (q, 2H), 3.41-3.39 (m, 2H), 1.37 (t, 3H); HRMS calculated for (M + H) 363.1816, found 363.1807.

EXAMPLE 25

5 [000209] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-[2-[4-(dimethylamino)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylate trifluoroacetate.

10 [000210] Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride in using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.758 g, 1.92 mmol), 4-dimethylaminophenylboronic acid (0.475 g, 2.88 mmol), 2M Na₂CO₃ (8 mL), toluene (23 mL), Pd[dppf]Cl₂·CH₂Cl₂ (0.126 g, 0.154 mmol), and 4N HCl/dioxane (10.0 mL). The TFA salt was isolated as a yellow solid (0.666 g, 70.3% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.64 (d, 1H), 8.46 (s, 1H), 8.07-8.04 (m, 5H), 7.95 (s, 1H), 7.88 (d, 1H), 6.86 (d, 2H), 4.85 (t, 2H), 4.38 (q, 2H), 3.47-3.37 (m, 2H), 3.03 (s, 6H), 1.36 (t, 3H); HRMS calculated for (M + H) 380.2081, found 380.2098.

EXAMPLE 26

20 [000211] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-[2-(3-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride.

25 [000212] Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride, using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.752g, 1.90 mmol), 3-methoxyphenylboronic acid (0.434 g, 2.86 mmol), 2M Na₂CO₃ (7 mL), toluene (20 mL), Pd[dppf]Cl₂·CH₂Cl₂ (0.124 g, 0.152 mmol), and 4N HCl/dioxane (10.0 mL). The product was isolated as an off-white solid (0.500 g, 60.0% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.81 (d, 1H), 8.67 (s, 1H), 8.37 (br s, 3H), 8.22 (d, 1H), 8.04 (s, 1H), 7.82-7.79 (m, 2H), 7.53 (dd, 1H), 7.18 (d, 1H), 4.89 (t, 2H), 4.38 (q, 2H), 3.89 (s, 3H), 3.42-3.36

(m, 2H), 1.36 (t, 3H); HRMS calculated for (M + H) 367.1765, found 367.1755.

EXAMPLE 27

5 [000213] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-[2-(3-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride.

[000214] Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride using ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (0.752g, 1.90 mmol), 3-hydroxyphenylboronic acid (0.629 g, 2.86 mmol), 2M Na₂CO₃ (7 mL), toluene (20 mL), Pd[dppf]Cl₂·CH₂Cl₂ (0.124 g, 0.152 mmol), and 4N HCl/dioxane (10.0 mL). The product was isolated as an off-white solid (0.381 g, 46.8% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.80 (d, 1H), 8.62 (s, 1H), 8.37 (br s, 3H), 8.22 (d, 1H), 8.04 (s, 1H), 7.62-7.58 (m, 2H), 7.41 (dd, 1H), 7.05 (dd, 1H), 4.89 (t, 2H), 4.38 (q, 2H), 3.89 (s, 3H), 3.42-3.36 (m, 2H), 1.36 (t, 3H); HRMS calculated for (M + H) 353.1608, found 353.1630.

EXAMPLE 28

20 [000215] This example illustrates the production of 2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

[000216] A flask was charged with ethyl 1-(2-aminoethyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride (0.303 g, 0.659 mmol), NH₄OH (6 mL) and ethanol (3 mL). After stirring for 20 h, the reaction was filtered, and the solid rinsed with water, ethanol, and diethyl ether to obtain a white solid (0.178 g, 76.8% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.73 (s, 1H), 9.16 (s, 1H), 8.79 (s, 1H), 8.64 (s, 1H), 8.35 (s, 1H), 8.16-8.08 (m, 2H), 7.90-7.69 (m, 4H), 4.45 (br s, 2H), 3.69 (br s, 2H); HRMS calculated for (M + H) 342.1349, found 342.1365.

EXAMPLE 29

[000217] This example illustrates the production of 2-(2-quinolin-3-ylpyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000218] Synthesis was conducted as it was for the production of 2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one using ethyl 1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride (0.144 g, 0.303 mmol), NH₄OH (6mL), and ethanol (3 mL). A white solid was obtained (0.046 g, 43% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.72 (d, 1H), 9.16 (d, 1H), 8.77 (d, 1H), 8.61 (s, 1H), 8.35 (br s, 1H), 8.15-8.07 (m, 2H), 7.88-7.79 (m, 2H), 7.71-7.66 (m, 2H), 4.55 (t, 2H), 3.29-3.23 (m, 2H), 2.22-2.18 (m, 2H); HRMS calculated for (M + H) 356.1506, found 356.1525.

EXAMPLE 30

[000219] This example illustrates the production of 2-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

[000220] Synthesis was conducted as it was for the production of 2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one using ethyl 1-(2-aminoethyl)-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride (0.202 g, 0.459 mmol), NH₄OH (8 mL), and ethanol (4 mL). A white solid was obtained (0.098 g, 66% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.67 (d, 1H), 8.36-8.32 (m, 2H), 8.15 (d, 2H), 7.78 (d, 1H), 7.64 (s, 1H), 7.45 (d, 2H), 5.27 (t, 1H), 4.57 (d, 2H), 4.42 (t, 2H), 3.70-3.66 (m, 2H); HRMS calculated for (M + H) 321.1346, found 321.1333.

EXAMPLE 31

[000221] This example illustrates the production of 2-[2-(3-methoxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

[000222] Synthesis was conducted as it was for the production of 2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one using ethyl 1-(2-aminoethyl)-3-[2-(3-methoxyphenyl)pyridin-4-yl]-1H-

pyrazole-5-carboxylate dihydrochloride (0.253 g, 0.575 mmol), NH_4OH (8 mL), and ethanol (4 mL). An off-white solid was obtained (0.160 g, 86.8% yield): ^1H NMR (DMSO-d_6 / 300 MHz) δ 8.69 (s, 1H), 8.37-8.32 (m, 2H), 7.80-7.68 (m, 4H), 7.42 (dd, 1H), 7.03 (d, 1H), 4.42 (t, 2H), 3.86 (s, 3H), 3.69-3.65 (m, 2H); HRMS calculated for (M + H) 321.1346, found 321.1344.

EXAMPLE 32

[000223] This example illustrates the production of 2-[2-(3-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

10 [000224] Synthesis was conducted as it was for the production of 2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one using ethyl 1-(2-aminoethyl)-3-[2-(3-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.304 g, 0.715 mmol), NH_4OH (10 mL), and ethanol (5 mL). An off-white solid was obtained (0.178 g, 81.1% yield): ^1H NMR (DMSO-d_6 / 300 MHz) δ 9.56 (br s, 1H), 8.66 (d, 1H), 8.32-8.28 (m, 2H), 7.79 (s, 1H), 7.63-7.59 (m, 3H), 7.30 (dd, 1H), 6.85 (d, 1H), 4.42 (t, 2H), 3.69-3.64 (m, 2H); HRMS calculated for (M + H) 307.1190, found 307.1214.

EXAMPLE 33

20 [000225] This example illustrates the production of 2-[2-(3-nitrophenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride.

[000226] Synthesis was conducted as it was for the production of 2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate using ethyl 1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.253 g, 0.540 mmol), NH_4OH (8 mL), and ethanol (4 mL). The isolated TFA salt was converted to the HCl salt using methanol and 4N HCl/dioxane. After 0.5 h, the suspension was concentrated and the solid rinsed with diethyl ether to afford a white solid
25
30 (0.140 g, 65.2% yield): ^1H NMR (DMSO-d_6 / 300 MHz) δ 9.00 (dd, 1H), 8.78 (d, 1H), 8.65 (d, 1H), 8.60 (s, 1H), 8.39-8.33 (m, 2H), 8.00 (dd, 1H),

7.84 (dd, 1H), 7.74 (s, 1H), 4.54 (t, 2H), 3.27-3.21 (m, 2H), 2.20-2.15 (m, 2H); HRMS calculated for (M + H) 342.1349, found 342.1365.

EXAMPLE 34

5 [000227] This example illustrates the production of 2-{2-[4-(trifluoromethoxy)-phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

[000228] Synthesis was conducted as it was for the production of 2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate using ethyl 1-(2-aminoethyl)-3-{2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride (0.439 g, 0.889 mmol), NH₄OH (8 mL), and ethanol (4 mL). An off-white solid was obtained (0.181 g, 43.0% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.72 (d, 1H), 8.45 (s, 1H), 8.33-8.30 (m, 3H), 7.88 (d, 1H), 7.69 (s, 1H), 7.51 (d, 2H), 4.43 (q, 2H), 3.70-3.64 (m, 2H); HRMS calculated for (M + H) 375.1063, found 375.1068.

EXAMPLE 35

[000229] This example illustrates the production of 2-{2-[(E)-2-phenylvinyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

20 [000230] Synthesis was conducted as it was for the production of 2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate using ethyl 1-(2-aminoethyl)-3-{2-[(E)-2-phenylvinyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride (0.610 g, 1.40 mmol), NH₄OH (10 mL), and ethanol (5 mL). A yellow solid was obtained (0.352 g, 58.4% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.69 (d, 1H), 8.40 (br s, 2H), 7.97-7.92 (m, 2H), 7.71-7.68 (m, 3H), 7.49-7.37 (m, 4H), 4.45 (t, 2H), 3.70-3.66 (m, 2H); HRMS calculated for (M + H) 317.1397, found 317.1405.

EXAMPLE 36

30 [000231] This example illustrates the production of 2-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

[000232] Synthesis was conducted as it was for the production of 2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate using ethyl 1-(2-aminoethyl)-3-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate trifluoroacetate (0.350 g, 708 μ mol), NH_4OH (8 mL), and ethanol (4 mL). A yellow solid was obtained (0.204 g, 64.5% yield): ^1H NMR ($\text{DMSO}-d_6$ / 300 MHz) δ 8.61 (d, 1H), 8.51 (s, 1H), 8.40 (s, 1H), 8.03 (d, 2H), 7.96 (d, 1H), 7.86 (s, 1H), 6.88 (d, 2H), 4.46 (t, 2H), 3.71-3.66 (m, 2H), 3.05 (s, 6H); HRMS calculated for (M + H) 334.1662, found 334.1673.

EXAMPLE 37

[000233] This example illustrates the production of 2-[2-(4-methoxyphenyl)-pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

[000234] Synthesis was conducted as it was for the production of 2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate using ethyl 1-(2-aminoethyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.368 g, 0.838 μ mol), NH_4OH (8 mL), and ethanol (4 mL). A white solid was obtained (0.225 g, 61.8% yield): ^1H NMR ($\text{DMSO}-d_6$ / 300 MHz) δ 8.69 (d, 1H), 8.45 (s, 1H), 8.36 (s, 1H), 8.13 (d, 2H), 7.93 (d, 1H), 7.76 (s, 1H), 7.12 (d, 2H), 4.44 (t, 2H), 3.85 (s, 3H), 3.71-3.66 (m, 2H); HRMS calculated for (M + H) 321.1346, found 321.1359.

EXAMPLE 38

[000235] This example illustrates the production of 2-[2-(3-nitrophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

[000236] Synthesis was conducted as it was for the production of 2-pyridin-4-yl-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate using ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.350 g, 0.771 μ mol), NH_4OH (8 mL), and ethanol (4 mL). A beige solid was obtained (0.152 g, 43.8% yield): ^1H

NMR (DMSO- d_6 / 300 MHz) δ 9.00 (s, 1H), 8.75 (d, 1H), 8.66 (d, 1H), 8.32-8.28 (m, 2H), 7.89 (d, 1H), 7.81 (dd, 1H), 7.73 (s, 1H), 4.43 (t, 2H), 3.70-3.65 (m, 2H); HRMS calculated for (M + H) 336.1091, found 336.1068.

EXAMPLE 39

5 **[000237]** This example illustrates the production of 2-[2-(3,4-difluorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

10 **[000238]** Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride using 2-(2-chloropyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one (0.175 g, 0.702 mmol), 3,4-difluorophenylboronic acid (0.167 g, 1.05 mmol), 2M Na₂CO₃ (2 mL), toluene (7 mL), and Pd[dppf]Cl₂·CH₂Cl₂ (0.046 g, 0.057 mmol). The TFA salt was isolated as a white solid (0.076 g, 25% yield): ¹H NMR (DMSO- d_6 / 300 MHz) δ 8.68 (d, 15 1H), 8.42 (s, 1H), 8.32-8.23 (m, 2H), 8.09 (d, 1H), 7.82 (s, 1H), 7.69 (s, 1H), 7.55 (dd, 1H), 4.41 (t, 2H), 3.69-3.64 (m, 2H); HRMS calculated for (M + H) 327.1052, found 327.1078.

EXAMPLE 40

20 **[000239]** This example illustrates the production of 2-[2-(3,4-dichlorophenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

25 **[000240]** Synthesis was conducted as for the production of ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride using 2-(2-chloropyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one (0.175 g, 0.702 mmol), 3,4-dichlorophenylboronic acid (0.200 g, 1.05 mmol), 2M Na₂CO₃ (2 mL), toluene (7 mL), and Pd[dppf]Cl₂·CH₂Cl₂ (0.046 g, 0.057 mmol). The TFA salt was obtained as a gray solid (0.016 g, 4.9% yield): ¹H NMR (DMSO- d_6 , TFA / 300 MHz) δ 8.81 (d, 1H), 8.66 (s, 1H), 8.42-8.34 (m, 2H), 8.28-8.18 (m, 2H), 7.87-7.85 (m, 2H), 4.45 (t, 2H), 3.71-3.64 (m, 2H), HRMS calculated for (M + H) 359.0461, found 359.0476.

30

EXAMPLE 41

[000241] This example illustrates the production of 1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride.

5 [000242] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate (0.253 g, 0.539 mmol) and LiOH•H₂O (0.0679 g, 1.62 mmol). The isolated TFA salt was a tacky
10 solid, and thus converted to the HCl salt using methanol and 4N HCl/dioxane. After 0.5 h, the suspension was concentrated and the solid rinsed with diethyl ether to afford a white solid (0.223 g, 94.6% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.00 (dd, 1H), 8.79 (d, 1H), 8.68 (d, 1H), 8.60 (s, 1H), 8.34 (dd, 1H), 8.11 (br s, 3H), 7.99 (dd, 1H), 7.88-7.81 (m, 2H), 4.68 (t, 2H), 2.86-2.79 (m, 2H), 2.20-2.15 (m, 2H); HRMS calculated
15 for (M + H) 368.1353, found 368.1336.

EXAMPLE 42

[000243] This example illustrates the production of 1-(2-aminoethyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylic acid trifluoroacetate.

20 [000244] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride (0.211 g, 0.457 mmol) and LiOH•H₂O (0.077 g, 1.8 mmol). A pink solid was obtained (0.150 g, 69.4% yield): ¹H NMR
25 (DMSO-d₆ / 300 MHz) δ 9.75 (s, 1H), 9.23 (d, 1H), 8.82 (d, 1H), 8.69 (s, 1H), 8.17-8.10 (m, 2H), 8.01-7.94 (m, 3H), 7.89-7.83 (m, 2H), 7.72 (dd, 1H), 4.86 (t, 2H), 3.44-3.38 (m, 2H); HRMS calculated for (M + H) 360.1455, found 360.1466.

EXAMPLE 43

30 [000245] This example illustrates the production of 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride.

[000246] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.262 g, 0.516 mmol) and
5 LiOH•H₂O (0.097 g, 2.3 mmol). The isolated TFA salt converted to the HCl salt using methanol and 4N HCl/dioxane. After 0.5 h, the suspension was concentrated and the solid rinsed with diethyl ether to afford a yellow solid (0.102 g, 46.5% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.01 (s, 1H), 8.80 (d, 1H), 8.70-8.64 (m, 2H), 8.35-8.28 (m, 4H), 8.02 (d, 1H), 7.89-7.82
10 (m, 2H), 3.39-3.33 (m, 2H); HRMS calculated for (M + H) 354.1197, found 354.1176.

EXAMPLE 44

[000247] This example illustrates the production of 1-(2-aminoethyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid
15 trifluoroacetate.

[000248] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.260 g, 0.591 mmol) and
20 LiOH•H₂O (0.099 g, 2.4 mmol). A pale yellow solid was obtained (0.212 g, 79.4% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.69 (d, 1H), 8.42 (s, 2H), 8.15 (d, 1H), 8.00 (br s, 3H), 7.88-7.84 (m, 2H), 7.09 (d, 2H), 4.84 (t, 2H), 3.84 (s, 3H), 3.38-3.42 (m, 2H); HRMS calcd for (M + H) 339.1452, found 339.1472.

EXAMPLE 45

[000249] This example illustrates the production of 1-(2-aminoethyl)-3-[2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylic acid
25 trifluoroacetate.

[000250] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-[2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl]-
30

1H-pyrazole-5-carboxylate dihydrochloride (0.257 g, 0.521 mmol) and LiOH•H₂O (0.097 g, 2.3 mmol). A beige solid was obtained (0.088 g, 34% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.73 (d, 1H), 8.50 (s, 1H), 8.32 (d, 2H), 8.01 (br s, 3H), 7.88 (d, 1H), 7.81 (s, 1H), 7.51 (d, 2H), 4.84 (t, 2H), 3.42-3.38 (m, 2H); HRMS calculated for (M + H) 393.1169, found 393.1189.

EXAMPLE 46

[000251] This example illustrates the production of 1-(2-aminoethyl)-3-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid trifluoroacetate.

[000252] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate using ethyl 1-(2-aminoethyl)-3-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate trifluoroacetate (0.238 g, 0.482 mmol) and LiOH•H₂O (0.088 g, 2.1 mmol). A neon orange solid was obtained (0.150 g, 67.0% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.64 (d, 1H), 8.46 (s, 1H), 8.07-8.04 (m, 5H), 7.91-7.88 (m, 2H), 6.86 (d, 2H), 4.86 (t, 2H), 3.43-3.37 (m, 2H), 3.03 (s, 6H); HRMS calcd for (M + H) 352.1768, found 352.1770.

EXAMPLE 47

[000253] This example illustrates the production of 1-(2-aminoethyl)-3-{2-[(E)-2-phenylethenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid trifluoroacetate.

[000254] Synthesis was conducted as in the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-{2-[(E)-2-phenylvinyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride (0.311 g, 0.714 mmol) and LiOH•H₂O (0.120 g, 2.86 mmol). The TFA salt was isolated as a light yellow solid (0.220 g, 68.7% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.68 (d, 1H), 8.26 (s, 1H), 8.03 (br s, 3H), 7.89-7.84 (m, 2H), 7.73-7.68 (m, 3H),

7.45-7.36 (m, 4H), 4.89 (t, 2H), 3.41-3.39 (m, 2H); HRMS calculated for (M + H) 335.1503, found 335.1496.

EXAMPLE 48

[000255] This example illustrates the production of 1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylic acid dihydrochloride.

[000256] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylate dihydrochloride (0.134 g, 0.282 mmol) and LiOH•H₂O (0.047 g, 1.1 mmol). The TFA salt was converted to the HCl salt using methanol and 4N HCl/dioxane. After 0.5 h, the suspension was concentrated and the solid rinsed with diethyl ether to afford a yellow solid (0.086 g, 68% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.92 (d, 1H), 9.95 (s, 1H), 9.83 (s, 1H), 8.44-8.41 (m, 2H), 8.17-8.09 (m, 4H), 8.03 (d, 1H), 7.94 (dd, 1H), 7.86 (s, 1H), δ 4.69 (t, 2H), 2.90-2.82 (m, 2H), 2.25-2.16 (m, 2H); HRMS calculated for (M + H) 374.1612, found 374.1622.

EXAMPLE 49

[000257] This example illustrates the production of 1-(2-aminoethyl)-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid dihydrochloride.

[000258] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride (0.167 g, 0.379 mmol) and LiOH•H₂O (0.064 g, 1.5 mmol). The TFA salt was converted to the HCl salt using methanol and 4N HCl/dioxane. After 0.5 h, the suspension was concentrated and the solid rinsed with diethyl ether to afford a pale pink solid (0.069 g, 70% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.80 (d, 1H), 8.67 (s, 1H), 8.33 (br s, 3H), 8.22-8.19 (m, 3H), 8.00 (s, 1H), 7.55 (d, 2H), 4.89 (t, 2H), 4.62 (m, 2H), 3.40-3.35 (m, 2H); HRMS calculated for (M + H) 339.1452, found 339.1435.

EXAMPLE 50

[000259] This example illustrates the production of 1-(2-aminoethyl)-3-[2-(3-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid trifluoroacetate.

5 [000260] Synthesis was conducted as for the preparation of 1-(2-aminoethyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid trifluoroacetate, using ethyl 1-(2-aminoethyl)-3-[2-(3-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (0.179 g, 0.407 mmol) and LiOH•H₂O (0.105 g, 2.50 mmol). An off-white solid was obtained (0.168 g,
10 91.2% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.72 (d, 1H), 8.43 (s, 1H), 8.00 (br s, 3H), 7.88 (d, 1H), 7.83 (s, 1H), 7.78-7.74 (m, 2H), 7.44 (dd, 1H), 7.06 (dd, 1H), 4.84 (t, 2H), 3.85 (s, 3H), 3.44-3.38 (m, 2H); HRMS calculated for (M + H) 339.1452, found 339.1469.

EXAMPLE 51

15 [000261] This example illustrates the production of 1-(2-aminoethyl)-N-hydroxy-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxamide trifluoroacetate.

[000262] Freshly made sodium methoxide (3.48 M, 0.27 mL) was added dropwise to a stirring solution of hydroxylamine hydrochloride (0.039 mL,
20 0.94 mmol) in methanol (1 mL) maintained at 40°C. The white slurry was cooled to room temperature, and then a solution of ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate (0.400 g, 0.857 mmol) in methanol (5 mL) was added. After stirring for 48 h, 7 mL of 4N HCl (aq) was added and the
25 reaction stirred for 20 h. Purification by Gilson RP HPLC (5-95% acetonitrile/water) afforded a pink solid (0.129 g, 32.1% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 11.53 (br s, 1H), 8.73 (d, 1H), 8.31 (s, 1H), 8.11-8.01 (m, 5H), 7.75 (d, 1H), 7.52-7.46 (m, 3H), 4.77 (t, 2H), 4.58 (s, 2H), 3.42-3.35 (m, 2H); HRMS calculated for (M + H) 354.1561, found
30 354.1538.

EXAMPLE 52

[000263] This example illustrates the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride.

5 [000264] A flask was charged with 1. 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol), pyrazole (0.325 g, 4.78 mmol), and sodium hydride (0.229 g, 5.73 mmol) in 6 mL anhydrous DMF and stirred under N₂ for 60 h at 138°C. The reaction was then quenched with 1N HCl (aq) and purified by Gilson RP
10 HPLC (5-95% acetonitrile/water). The appropriate fractions were concentrated and converted to the HCl salt using methanol and 4N HCl/dioxane. The mixture was concentrated to a light yellow solid (0.028 g, 8.2% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.64 (d, 1H), 8.48 (d, 1H), 8.35 (br s, 2H), 7.85 (s, 1H), 7.77 (d, 1H), 7.45 (s, 1H), 6.59 (s, 1H), 4.53
15 (t, 2H), 3.26-3.20 (m, 2H), 2.22-2.16 (m, 2H); HRMS calculated for (M + H) 295.1302, found 295.1285.

EXAMPLE 53

[000265] This example illustrates the production of 2-[2-(1H-imidazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride.
20

[000266] Synthesis was conducted as for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol)
25 and imidazole, sodium derivative (0.431 g, 4.78 mmol). A white solid was obtained (0.074 g, 23% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 10.0 (s, 1H), 8.62 (d, 1H), 8.55 (s, 1H), 8.48 (s, 1H), 8.41 (br s, 1H), 7.99 (d, 1H), 7.90 (s, 1H), 7.65 (s, 1H), 4.53 (t, 2H), 3.28-3.22 (m, 2H), 2.22-2.16 (m, 2H); HRMS calculated for (M + H) 295.1302, found 295.1290.

EXAMPLE 54

[000267] This example illustrates the production of 2-[2-(1H-pyrrol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate.

5 [000268] Synthesis was conducted as for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride, using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol), pyrrole (0.334 mL, 4.78 mmol), and sodium hydride (0.229 g, 5.73 mmol).

10 The TFA salt was isolated as a black solid (0.128 g, 32.7% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.42 (d, 1H), 8.34 (br s, 1H), 8.05 (s, 1H), 7.79 (br s, 2H), 7.67-7.64 (m, 2H), 6.30 (br s, 2H), 4.52 (t, 2H), 3.28-3.22 (m, 2H), 2.22-2.16 (m, 2H); HRMS calculated for (M + H) 294.1349, found 294.1348.

15

EXAMPLE 55

[000269] This example illustrates the production of 2-[2-(4-methyl-1H-imidazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride.

20 [000270] Synthesis was conducted as for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride, using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol), 4-methylimidazole (0.393 g, 4.78 mmol), and sodium hydride (0.229 g, 5.73 mmol). A brown solid was obtained (0.053 g, 16% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.88 (s, 1H), 8.60 (d, 1H), 8.40 (br s, 2H), 8.28 (s, 1H), 7.97 (d, 1H), 7.63 (s, 1H), 4.53 (t, 2H), 3.29-3.22 (m, 2H), 2.36 (s, 3H), 2.22-2.16 (m, 2H); HRMS calculated for (M + H) 309.1458, found 309.1462.

25

EXAMPLE 56

[000271] This example illustrates the production of 2-[2-(4-phenyl-1H-imidazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate.

5 [000272] Synthesis was conducted as for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride, using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol),
4-phenylimidazole (0.690 g, 4.78 mmol), and sodium hydride (0.229 g,
10 5.73 mmol). The TFA salt was isolated as a white solid (0.060 g, 13% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.17 (s, 1H), 8.76 (s, 1H), 8.57 (d, 1H), 8.37-8.32 (m, 2H), 7.93-7.88 (m, 3H), 7.65 (s, 1H), 7.47 (dd, 2H), 7.36-7.34 (m, 1H), 4.54 (t, 2H), 3.29-3.22 (m, 2H), 2.22-2.16 (m, 2H); HRMS calculated for (M + H) 371.1615, found 371.1626.

15

EXAMPLE 57

[000273] This example illustrates the production of 2-[2-(4-methyl-1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate.

[000274] Synthesis was conducted as it was for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride, using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol),
20 4-methylpyrazole (0.40 mL, 4.8 mmol), and sodium hydride (0.229 g, 5.73 mmol). The TFA salt was isolated as a white solid (0.068 g, 16% yield): ¹H
25 NMR (DMSO-d₆ / 300 MHz) δ 8.45-8.29 (m, 4H), 7.72 (m, 1H), 7.66 (s, 1H), 7.42 (s, 1H), 4.54 (t, 2H), 3.25-3.20 (m, 2H), 2.20-2.11 (m, 5H); HRMS calculated for (M + H) 309.1458, found 309.1448.

EXAMPLE 58

[000275] This example illustrates the production of 2-[2-(1H-1,2,4-triazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride.

5 [000276] Synthesis was conducted as it was for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride, using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol) and 1,2,4-triazole, sodium derivative (0.435 g, 4.78 mmol). A white solid
10 was obtained (0.157 g, 48.4% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 9.40 (s, 1H), 8.55 (d, 1H), 8.37-8.33 (m, 2H), 8.27 (s, 1H), 7.90 (dd, 1H), 7.50 (s, 1H), 4.53 (t, 2H), 3.25-3.20 (m, 2H), 2.22-2.14 (m, 2H); HRMS calcd for (M + H) 296.1254, found 296.1244.

EXAMPLE 59

15 [000277] This example illustrates the production of 2-[2-(1H-1,2,3-triazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride.

[000278] Synthesis was conducted as it was for the production of 2-[2-(1H-pyrazol-1-yl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one hydrochloride, using 2-(2-chloropyridin-4-yl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.251 g, 0.960 mmol),
20 1,2,3-triazole (0.28 mL, 4.8 mmol), and sodium hydride (0.229 g, 5.73 mmol). A white solid was obtained (0.118 g, 36.0% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.89 (s, 1H), 8.61 (d, 1H), 8.52 (s, 1H), 8.37 (br s, 1H), 8.01-7.96 (m, 2H), 7.55 (s, 1H), 4.54 (t, 2H), 3.29-3.22 (m, 2H), 2.22-2.16 (m, 2H); HRMS calcd for (M + H) 296.1254, found 296.1243.
25

EXAMPLE 60

[000279] This example illustrates the production of ethyl 1-{3-[(tert-butoxycarbonyl)-amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate.
30

[000280] To a cooled (0°C) solution of ethyl 3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate in anhydrous DMF (35 mL) was added lithium *t*-butoxide (1M in THF, 6.6 mL) dropwise. The reaction stirred for 30 min, and then a solution of *tert*-butyl 3-bromopropylcarbamate (1.57 g, 6.60 mmol) and sodium iodide (0.989 g, 6.60 mmol) in anhydrous DMF (10 mL) was added dropwise. The reaction was allowed to stir and warm to room temperature for 4 h. The reaction was then poured into water and brine and extracted with ethyl acetate. The organic layers were combined, dried over MgSO₄, filtered, and concentrated. Chromatographic purification (15% ethyl acetate/hexane) afforded a yellow oil (1.13 g, 63.7% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 7.81 (d, 2H), 7.27 (s, 1H), 6.87 (t, 1H), 6.79 (d, 2H), 4.52 (t, 2H), 4.37 (q, 2H), 3.80 (s, 3H), 3.30-2.94 (m, 2H), 1.96-1.91 (m, 2H), 1.39-1.33 (m, 12H); HRMS calculated for (M + H) 404.2180, found 404.2190.

EXAMPLE 61

[000281] This example illustrates the production of 1-{3-[(*tert*-butoxycarbonyl)-amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylic acid.

[000282] A solution of ethyl 1-{3-[(*tert*-butoxycarbonyl)amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate (0.467 g, 1.16 mmol) and LiOH•H₂O (0.097 g, 2.32 mmol) in 12 mL of THF/H₂O (1:1) stirred at room temperature for 3 h. The reaction mixture was concentrated to the aqueous phase, then diluted with 0.1N HCl and extracted three times with ethyl acetate. The organic layers were combined, dried over MgSO₄, filtered, and concentrated to a pale yellow solid (0.359 g, 82.3% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 7.71 (d, 2H), 7.01-6.94 (m, 3H), 6.88 (s, 1H), 4.62 (t, 2H), 3.80 (s, 3H), 2.94-2.90 (m, 2H), 1.91-1.84 (m, 2H), 1.39 (s, 9H); HRMS calculated for (M + H) 376.1867, found 376.1906.

EXAMPLE 61

[000283] This example illustrates the production of 1-(3-aminopropyl)-3-(4-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid dihydrochloride.

[000284] To a cooled (-78 °C) solution of 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylic acid (0.262 g, 0.70 mmol) in anhydrous dichloromethane (7 mL) was added boron tribromide (1.0M in CH₂Cl₂, 7.0 mL) dropwise. After 1 h the reaction was carefully quenched with water, then concentrated to the aqueous layer and purified by Gilson RP HPLC (5-95% acetonitrile/water). The appropriate fractions were concentrated. The residue was converted to the HCl salt using 4N HCl/dioxane and methanol. After stirring 1 h, the mixture was concentrated to a white solid (0.238 g, 100% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.07 (br s, 3H), 7.67 (d, 2H), 7.18 (s, 1H), 6.83 (d, 2H), 4.59 (t, 2H), 2.84-2.78 (m, 2H), 2.18-2.08 (m, 2H); HRMS calculated for (M + H) 262.1186, found 262.1195.

EXAMPLE 62

[000285] This example illustrates the production of ethyl 1-(3-aminopropyl)-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate hydrochloride.

[000286] To a flask charged with ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate (0.588 g, 1.46 mmol) was added 4N HCl/dioxane (5 mL). After 1 h the reaction mixture was filtered and rinsed with diethyl ether to yield a white solid (0.443 g, 89.4% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 8.06 (br s, 3H), 7.81 (d, 2H), 7.32 (s, 1H), 7.00 (d, 2H), 4.61 (t, 2H), 4.37 (q, 2H), 3.80 (s, 3H), 2.87-2.80 (m, 2H), 2.20-2.11 (m, 2H), 1.36 (t, 3H), HRMS calcd for (M + H) 304.1656, found 304.1665.

EXAMPLE 63

[000287] This example illustrates the production of 2-(4-methoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000288] To a flask charged with ethyl 1-(3-aminopropyl)-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate hydrochloride (0.418 g, 1.23 mmol) was added NH₄OH (20 mL) and ethanol (10 mL). After stirring for 18h, the reaction was filtered, and the white solid rinsed with diethyl ether to afford a white solid (0.243 g, 76.8% yield): ¹H NMR (DMSO-d₆ / 300

MHz) δ 8.28 (br s, 1H), 7.77 (d, 2H), 7.10 (s, 1H), 6.98 (d, 2H), 4.45 (t, 2H), 3.80 (s, 3H), 3.26-3.21 (m, 2H), 2.20-2.13 (m, 2H); HRMS calculated for (M + H) 258.1242, found 258.1237.

EXAMPLE 64

5 [000289] This example illustrates the production of 2-(4-hydroxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000290] Synthesis was conducted as it was for the preparation of 1-(3-amino-propyl)-3-(4-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid dihydrochloride using ethyl 1-(3-aminopropyl)-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate hydrochloride (0.182 g, 0.70 mmol) and boron
10 tribromide (7.0 mL). Purification by Gilson RP HPLC (5-95% acetonitrile/water) afforded a white solid (0.078 g, 46% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 9.55 (br s, 1H), 8.25 (br s, 1H), 7.65 (d, 2H), 7.03 (s, 1H), 6.80 (d, 2H), 4.44 (t, 2H), 3.26-3.21 (m, 2H), 2.18-2.13 (m, 2H); HRMS
15 calculated for (M + H) 244.1081, found 244.1049.

EXAMPLE 65

[000291] This example illustrates the production of ethyl 1-{3-[(tert-butoxy-carbonyl)amino]propyl}-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylate.

20 [000292] Synthesis was conducted as it was for the preparation of ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate, using ethyl 3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylate (2.01 g, 8.17 mmol), lithium *t*-butoxide (12.3 mL), *tert*-butyl 3-bromopropylcarbamate (2.92 g, 12.3 mmol) and sodium iodide (1.84 g,
25 12.3 mmol). Flash chromatography (12% ethyl acetate/hexane) afforded a yellow oil, which was triturated with diethyl ether to yield a pale yellow solid (1.47 g, 45.0% yield): HRMS calculated for (M + H) 404.2180, found 404.2206.

EXAMPLE 66

30 [000293] This example illustrates the production of 1-{3-[(tert-butoxycarbonyl)-amino]propyl}-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylic acid.

[000294] Synthesis was conducted as it was for the preparation of 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylic acid using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylate (0.610 g, 1.51 mmol) and
5 LiOH•H₂O (0.127 g, 3.02 mmol). An off-white solid was obtained (0.565g, 100% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 13.47 (br s, 1H), 7.46-7.32 (m, 4H), 6.93-6.86 (m, 2H), 4.56 (t, 2H), 3.83 (s, 3H), 3.01-2.95 (m, 2H), 1.96-1.90 (m, 2H), 1.39 (s, 9H); HRMS calculated for (M + H) 376.1873, found 376.1896.

10

EXAMPLE 67

[000295] This example illustrates the production of 1-(3-aminopropyl)-3-(3-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid trifluoroacetate.

[000296] Synthesis was conducted as it was for the preparation of 1-(3-amino-propyl)-3-(4-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid
15 dihydrochloride, using 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylic acid (0.496 g, 1.32 mmol) and boron tribromide (1.0M in CH₂Cl₂, 13.0 mL). The TFA salt was isolated as an off-white solid (0.353 g, 71.3% yield): ¹H NMR (DMSO-d₆ / 300 MHz) δ 13.62 (br s, 1H), 9.56 (br s, 1H), 7.80 (br s, 3H), 7.27-7.23 (m, 4H), 6.77
20 (d, 1H), 4.63 (t, 2H), 2.89-2.81 (m, 2H), 2.17-2.07 (m, 2H); HRMS calculated for (M + H) 262.1192, found 262.1223.

EXAMPLE 68

[000297] This example illustrates the production of 2-(3-methoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000298] Synthesis was conducted as it was for the preparation of ethyl 1-(3-aminopropyl)-3-(4-methoxyphenyl)-1H-pyrazole-5-carboxylate
25 hydrochloride, using ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(3-methoxyphenyl)-1H-pyrazole-5-carboxylate (0.737 g, 1.83 mmol) and 4N HCl/dioxane (10 mL). The resulting colorless oil was subjected to
30 conditions described for the production of 2-(4-methoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one, using NH₄OH (12 mL) and ethanol (6 mL). An off-white solid was obtained: ¹H NMR (DMSO-d₆ /

300 MHz) δ 8.29 (br s, 1H), 7.34-7.45 (m, 3H), 7.23 (s, 1H), 6.90 (dd, 1H), 4.48 (t, 2H), 3.83 (s, 3H), 3.26-3.21 (m, 2H), 2.22-2.14 (m, 2H); HRMS calculated for (M + H) 258.1242, found 258.1232.

EXAMPLE 69

5 **[000299]** This example illustrates the production of 2-(3-hydroxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000300] Synthesis was conducted as it was for the preparation of 1-(3-amino-propyl)-3-(4-hydroxyphenyl)-1H-pyrazole-5-carboxylic acid dihydrochloride, using 2-(3-methoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one (0.325 g 1.26 mmol) and boron tribromide (13.0 mL). Purification by Gilson RP HPLC (5-95% acetonitrile/water) afforded a white solid (0.194 g, 63.3% yield): ^1H NMR (DMSO- d_6 / 300 MHz) δ 9.45 (br s, 1H), 8.28 (br s, 1H), 7.28-7.19 (m, 3H), 7.08 (s, 1H), 6.75-6.72 (m, 1H), 4.47 (t, 2H), 3.26-3.21 (m, 2H), 2.20-2.16 (m, 2H); HRMS calculated for (M + H) 244.1086, found 244.1115.

EXAMPLE 70

[000301] This example illustrates the production of 1-(3-[[2-(4-bromophenyl)ethyl]-amino]propyl)-3-pyridin-4-yl-1H-pyrazole-5-carboxylic acid hydrochloride.

20 **[000302]** A single neck round bottom flask was charged with ethyl 3-pyridin-4-yl-1H-pyrazole-5-carboxylate (1.0 g, 4.6 mmol) and 30 mL DMF. The solution was cooled to -40°C in a dry ice/ CH_3CN bath. A 1 M solution of lithium *t*-butoxide in THF (6.9 mL, 6.9 mmol) was added dropwise over 5 minutes. After stirring at -40°C for 30 minutes, a solution of 3-[[2-(4-bromophenyl)ethyl](*tert*-butoxycarbonyl)amino]propyl methanesulfonate

25 (3.0 g, 6.9 mmol) in 10 mL DMF was added dropwise over 5 minutes. After 1 hour the reaction was allowed to warmed to ambient temperature and stirred for 18 hours. The reaction mixture was concentrated in vacuo, and the residue taken up in ethyl acetate. This was washed 3 times with

30 brine, dried over magnesium sulfate, and concentrated to a brown oil. The oil was treated with 20 mL of a 4 N HCl in dioxane solution, and stirred for 30 minutes to remove the protecting group. After concentrating in vacuo

the crude mixture was treated with 20 mL of a 2.5 N sodium hydroxide solution. The mixture was heated to 100 °C for 1 hour to hydrolyze the ester. The resulting product mixture was concentrated in vacuo to half the volume, and then chromatographed on a Gilson reverse phase HPLC
5 eluting with an acetonitrile/water gradient (5-70% CH₃CN over 15 minutes) to provide the desired product as a TFA salt. The salt was taken up in methanol, and treated with 10 mL of a 4 N solution of HCl in dioxane to convert to the HCl salt. Crystallization from diethyl ether afforded 1.30g (56%) of the title compound as a tan solid. LCMS showed a single peak
10 with m/z 429 (M+H). ¹H nmr (DMSO-d₆/300 MHz) δ 9.48 (broad s, 2H), 8.94 (d, 2H), 8.46 (d, 2H), 7.90 (s, 1H), 7.50 (d, 2H), 7.22 (d, 2H), 4.71 (t, 2H), 3.20-2.88 (m, 6H), 2.30 (m, 2H), ES⁺ HRMS calculated for M+H 429.0921, observed 429.0934.

EXAMPLE 71

15 **[000303]** This example illustrates the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride.

[000304] Step 1. Preparation of ethyl 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride.

20 **[000305]** This compound was prepared as part of a parallel library. A reaction tube was charged with ethyl 1-{3-[(tert-butoxycarbonyl)amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (550 mg, 1.35 mmol) and 4-methoxybenzene boronic acid (307 mg, 2.02 mmol). The mixture was purged with N₂, and then 16 mL
25 toluene and 6 mL of a 2M sodium carbonate solution were added. The reaction mixture was again purged with N₂. To this stirring mixture [1,1'-bis(diphenyl phosphino)ferrocene] dichloropalladium II·CH₂Cl₂ (88 mg, 0.11 mmol) was added, and the reaction heated to 80°C for 18 hours. The water layer was decanted, and the organic layer filtered through celite, and
30 washed with CH₂Cl₂. The filtrate was concentrated *in vacuo*, and the residue treated with 10 mL of a 4 N HCl in dioxane solution for 1 hour. The product was concentrated *in vacuo*, washed with diethyl ether, and air

dried to afford 673 mg (quantitative) of the desired ester. LCMS showed one major peak with m/z 381 ($M+H$).

[000306] Step 2. Preparation of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride.

5 **[000307]** This compound was prepared as part of a parallel library. A reaction tube was charged with ethyl 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride (200 mg, 0.44 mmol) and 10 mL of a 2.5 N NaOH solution. The mixture was heated to 100°C for 30 minutes, and concentrated *in vacuo*. The product
10 mixture was chromatographed on a Gilson reverse phase HPLC eluting with an acetonitrile/water gradient (5-70% CH_3CN over 15 minutes). The fractions containing the desired product were combined and concentrated. The oil was taken up in methanol and treated with 5 mL of a 4 N HCl in dioxane solution to obtain the HCl salt. The solution was concentrated to
15 dryness, washed with diethyl ether, and air dried to afford 196 mg (quantitative) of the title carboxylic acid. LCMS showed a single peak with m/z 353 ($M+H$). ES^+ HR MS calculated for $M+H$ 353.1608, observed 353.1640.

EXAMPLE 72

20 **[000308]** This example illustrates the production of 1-(3-aminopropyl)-3-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid hydrochloride.

[000309] The preparation of 1-(3-aminopropyl)-3-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid
25 hydrochloride was carried out in the same manner as that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride, using 4-dimethyl(amino)phenyl boronic acid. Purification afforded the title carboxylic acid as an orange solid. 6% yield over 2 steps. LCMS showed a
30 single peak with m/z 366 ($M+H$). ES^+ HR MS calculated for $M+H$ 366.1925, observed 366.1918.

EXAMPLE 73

[000310] This example illustrates the production of 1-(3-aminopropyl)-3-{2-[3-(hydroxymethyl)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid hydrochloride.

5 [000311] The preparation of 1-(3-aminopropyl)-3-{2-[3-(hydroxymethyl)phenyl]-pyridin-4-yl}-1H-pyrazole-5-carboxylic acid hydrochloride was carried out in the same manner as that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride, using 3-hydroxymethylphenyl boronic acid. Purification afforded the title carboxylic acid as a brown solid.
10 5% yield over 2 steps. LCMS showed a single peak with m/z 353 (M+H). ES⁺ HR MS calculated for M+H 353.1608, observed 353.1608.

EXAMPLE 74

15 [000312] This example illustrates the production of 1-(3-aminopropyl)-3-{2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid hydrochloride.

[000313] The preparation of 1-(3-aminopropyl)-3-{2-[4-(trifluoromethoxy)-phenyl] pyridin-4-yl}-1H-pyrazole-5-carboxylic acid hydrochloride was carried out in the same manner as that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride using 4-trifluoromethoxy-benzene boronic acid. Purification afforded the title carboxylic acid as a brown solid. 3%
20 yield over 2 steps. LCMS showed a single peak with m/z 407 (M+H). ES⁺ HR MS calculated for M+H 407.1326, observed 407.1358.

EXAMPLE 75

25 [000314] This example illustrates the production of 2-[2-(4-methoxyphenyl)-pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000315] This compound was prepared as part of a parallel library. A
30 reaction tube was charged with ethyl 1-(3-aminopropyl)-3-[2-(4-

methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride (250 mg, 0.55 mmol), 20 mL ammonium hydroxide, and 10 mL ethanol. The reaction mixture was stirred at room temperature for 18 hours. The mixture was then concentrated *in vacuo*, washed with water, filtered, and vacuum dried to afford 150 mg (81%) of the title lactam as a tan solid. LCMS showed a single peak with *m/z* 335 (*M*+*H*). ¹H nmr (DMSO-*d*₆ + TFA/300 MHz) δ 8.78 (d, 1H), 8.68 (s, 1H), 8.43 (br s, 1H), 8.26 (d, 1H), 8.09 (d, 2H), 7.93 (s, 1H), 7.23 (d, 2H), 4.59 (m, 2H), 3.89 (s, 3H), 3.26 (m, 2H), 2.21 (m, 2H). ES⁺ HR MS calculated for *M*+*H* 335.1503, observed 335.1490.

EXAMPLE 76

[000316] This example illustrates the production of 2-{2-[4-(dimethylamino)-phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-*a*][1,4]diazepin-4-one trifluoroacetate.

[000317] The preparation of 2-{2-[4-(dimethylamino)phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-*a*][1,4]diazepin-4-one trifluoroacetate was carried out in a manner similar to that described for the preparation of 2-[2-(4-methoxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-*a*][1,4]diazepin-4-one, except that it was chromatographed on a Gilson reverse phase HPLC eluting with an acetonitrile/water gradient (5-70% CH₃CN over 15 minutes). The pure fractions were combined and concentrated to afford the title lactam as a yellow solid (44% yield). LCMS showed a single peak with *m/z* 348 (*M*+*H*). ¹H nmr (DMSO-*d*₆/300 MHz) δ 8.61 (d, 1H), 8.52 (s, 1H), 8.41 (br s, 1H), 8.03 ((d, 2H), 7.98 (d, 1H), 7.84 (s, 1H), 6.89 (d, 2H), 4.57 (t, 2H), 3.25 (m, 2H), 3.05 (s, 6H), 2.21 (m, 2H). mp=223.0-226.7°C ES⁺ HR MS calculated for *M*+*H* 348.1819, observed 348.1790.

EXAMPLE 77

[000318] This example illustrates the production of 2-{2-[3-(hydroxymethyl)-phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-*a*][1,4]diazepin-4-one trifluoroacetate.

5 [000319] The preparation of 2-{2-[3-(hydroxymethyl)phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate was carried out in a manner similar to that described for the preparation of 2-[2-(4-methoxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one, except that it was chromatographed on a Gilson reverse phase HPLC eluting with an acetonitrile/water gradient (5-70% CH₃CN over 15 minutes). The pure fractions were combined and concentrated to afford the lactam as a white solid (26% yield). LCMS showed a single peak with *m/z* 335 (M+H). ¹H nmr (DMSO-d₆/300 MHz) δ 8.71 (d, 1H), 8.40 (s, 1H), 8.35 (br s, 1H), 8.11 (s, 1H), 8.03 (d, 1H), 7.86 (d, 1H), 7.63 (s, 1H), 7.53-7.41 (m, 2H), 4.61 (s, 2H), 4.54 (t, 2H), 3.24 (m, 2H), 2.19 (m, 2H). ES⁺ HR MS calculated for M+H 335.1503, observed 335.1520.

EXAMPLE 78

15 [000320] This example illustrates the production of 2-{2-[4-(trifluoromethoxy)-phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000321] The preparation of 2-{2-[4-(trifluoromethoxy)phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one was carried out in a manner similar to that described for the preparation of 2-[2-(4-methoxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one. The mixture was then concentrated *in vacuo*, washed with water, filtered, and vacuum dried to afford the title lactam as an off-white solid (65%). LCMS showed a single peak with *m/z* 389 (M+H). ¹H nmr (DMSO-d₆/300 MHz) δ 8.70 (d, 1H), 8.40 (s, 1H), 8.33 (m, 3H), 7.82 (d, 1H), 7.62 (s, 1H), 7.49 (d, 2H), 4.53 (m, 2H), 3.24 (m, 2H), 2.19 (m, 2H). ES⁺ HR MS calculated for M+H 389.1220, observed 389.1225.

EXAMPLE 79

30 [000322] This example illustrates the production of 2-{2-[4-(hydroxymethyl)phenyl]pyridin-4-yl}-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one.

[000323] Step 1. The preparation of ethyl 1-(3-aminopropyl)-3-[2-[4-(hydroxymethyl)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride was carried out in a manner similar to that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride, step 1 using 4-hydroxymethylphenyl boronic acid. Upon completion of the reaction, ethyl acetate and water were added. The layers were separated, and the organic layer was washed with water, dried over magnesium sulfate, filtered, and concentrated. The resulting residue was treated with excess 4 N HCl/dioxane for 15 minutes at room temperature. The reaction mixture was then chromatographed on a Gilson reverse phase HPLC eluting with an acetonitrile/water gradient (5-70% CH₃CN over 15 minutes). The pure fractions were combined, taken up in MeOH, and treated with 4 N HCl/dioxane to afford the HCl salt of the amine ester as a red solid (19 % yield). LCMS showed one major peak with *m/z* 381 (M+H).

[000324] Step 2. The preparation of 2-[2-[4-(hydroxymethyl)phenyl]pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one was carried out in a manner similar to that described for the preparation of 2-[2-(4-methoxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one. The product mixture was concentrated and filtered to afford the lactam as a tan solid (44% yield). LCMS showed a single peak with *m/z* 335 (M+H). ES⁺ HR MS calculated for M+H 335.1503, observed 335.1520.

EXAMPLE 80

[000325] This example illustrates the production of ethyl 1-(3-aminopropyl)-3-[2-(4 hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride.

[000326] The preparation of ethyl 1-(3-aminopropyl)-3-[2-(4-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate hydrochloride was carried out in a manner similar to that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic

acid hydrochloride, step 1 using 4-hydroxyphenyl boronic acid THP ether. Upon completion of the reaction in step 1, ethyl acetate and water were added. The layers were separated, and the organic layer was washed with water, dried over magnesium sulfate, filtered, and concentrated. The residue was treated with excess 4 N HCl/dioxane for 15 minutes. The mixture was concentrated, triturated with ether, and filtered to afford the title amine ester as a tan solid (95% yield). LCMS showed one major peak with m/z 367 ($M+H$). $mp = 241.6-244.4\text{ }^{\circ}\text{C}$.

EXAMPLE 81

[000327] This example illustrates the production of 1-(3-aminopropyl)-3-[2-(4-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride.

[000328] The preparation of 1-(3-aminopropyl)-3-[2-(4-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride was carried out in a manner similar to that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)-pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride, step 2. The title carboxylic acid was obtained as an off-white solid (51% yield). LCMS showed a single peak with m/z 339 ($M+H$). ES^+ HR MS calculated for $M+H$ 339.1452, observed 339.1455.

EXAMPLE 82

[000329] This example illustrates the production of 2-[2-(4-hydroxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate.

[000330] The preparation of 2-[2-(4-hydroxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one trifluoroacetate was carried out in a manner similar to that described for the preparation of 2-[2-(4-methoxyphenyl)pyridin-4-yl]-5,6,7,8-tetrahydro-4H-pyrazolo[1,5-a][1,4]diazepin-4-one. Upon completion of the reaction, the product mixture was concentrated, washed with water, and filtered. The mixture was then chromatographed on a Gilson reverse phase HPLC eluting with

an acetonitrile/water gradient (5-70% CH₃CN over 15 minutes). The pure fractions were combined and concentrated to afford the title compound as an off-white solid (41% yield). LCMS showed a single peak with m/z 321 (M+H). ¹H NMR (DMSO-d₆/300 MHz) δ 8.72 (d, 1H), 8.63 (s, 1H), 8.42 (br s, 1H), 8.22 (d, 1H), 8.02 (d, 2H), 7.81 (s, 1H), 7.03 (d, 2H), 4.57 (m, 2H), 3.25 (m, 2H), 2.20 (m, 2H). ES⁺ HR MS calculated for M+H 321.1346, observed 321.1368.

EXAMPLE 83

[000331] This examples illustrates the production of 1-(3-[[2-(4-bromophenyl)ethyl]-amino]propyl)-3-{2-[(E)-2-phenylvinyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid trifluoroacetate.

[000332] Step 1. Preparation of ethyl 1-{3-[[2-(4-bromophenyl)ethyl](tert-butoxycarbonyl)-amino]propyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate. A single neck roundbottom flask was charged with ethyl 3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (1.0 g, 4.0 mmol) and 30 mL dimethyl formamide under N₂. The solution was cooled to -40 °C in a dry ice/acetonitrile bath. A 1 M solution of lithium *t*-butoxide in THF (4.8 mL, 4.8 mmol) was added dropwise over 5 minutes. The reaction was stirred for 1 hour at -40 °C. A solution of 3-[[2-(4-bromophenyl)ethyl](tert-butoxycarbonyl)amino]propyl methanesulfonate (2.1 g, 4.8 mmol) in 10 mL dimethylformamide was added dropwise over 5 minutes. The resulting reaction mixture was stirred for 1 hour at -40 °C, and then stirred for 18 hours at room temperature. The mixture was then concentrated, and the residue taken up in diethyl ether. The solid impurities were filtered off, and washed with diethyl ether. The filtrate was concentrated to a brown oil. LCMS showed a mixture of the ethyl ester and hydrolyzed carboxylic acid. This product mixture was taken on to the next step without further purification.

[000333] Step 2. The preparation of ethyl 1-(3-[[2-(4-bromophenyl)ethyl]amino]-propyl)-3-{2-[(E)-2-phenylvinyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate. hydrochloride was carried out in a manner similar

to that described in example 2, step 1, using 2-phenylvinyl boronic acid. Upon completion of the reaction, ethyl acetate and water were added. The layers were separated, and the organic layer was washed with water, dried over magnesium sulfated, filtered and concentrated. The residue was treated with excess 4N HCl/dioxane for 30 minutes. The mixture was concentrated, washed with ether, and filtered to afford the ethyl ester as a brown solid (29% yield over 2 steps). LCMS showed one major peak with m/z 559 (M+H). ES⁺ HR MS calculated for M+H 559.1703, observed 559.1679.

[000334] Step 3. The preparation of 1-(3-{[2-(4-bromophenyl)ethyl]amino}propyl)-3-{2-[(E)-2-phenylvinyl]pyridin-4-yl}-1H-pyrazole-5-carboxylic acid trifluoroacetate was carried out in a manner similar to that described for the production of 1-(3-aminopropyl)-3-[2-(4-methoxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid hydrochloride, step 2. Upon completion of the reaction, the mixture was concentrated. The residue was then chromatographed on a Gilson reverse phase HPLC eluting with an acetonitrile/water gradient (5-70% CH₃CN over 15 minutes). The pure fractions were combined and concentrated to afford the title compound as an off-white solid (30% yield). LCMS showed a single peak with m/z 531 (M+H). ES⁺ HR MS calculated for M+H 531.1390, observed 531.1411.

EXAMPLE 84

[000335] This examples illustrates the production of ethyl 1-{2-[(tert-butoxy-carbonyl)amino]ethyl}-3-[2-(4-{[tert-butyl(dimethyl)silyl]oxy}phenyl)-pyridin-4-yl]-1H-pyrazole-5-carboxylate.

[000336] Ethyl 1-{2-[(tert-butoxycarbonyl)amino]ethyl}-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate (5.0 g, 0.013 mol), 4-(t-butyl dimethylsilyloxy)phenyl boronic acid (6.4 g, 0.025 mol), sodium carbonate (2.7 g, 0.025 mol), water (12.5 mL), and [1,1'-bis(diphenylphosphino)ferrocene]dichloro palladium (II) complex with dichloromethane (1:1) (1.0 g, 0.0013 mol) in toluene (100 mL) were

refluxed for 4 hours. Contents were allowed to cool and partitioned between EtOAc and water. The EtOAc layer was washed with brine, dried over MgSO_4 and concentrated *in vacuo*. The residue was filtered through a pad of silica gel, eluting with 25% EtOAc/hexanes to give a light amber oil. The oil crystallized under hexanes and was filtered to give the desired product as a white solid, 3.77 g (51% yield). FABHRMS m/z 567.2983 ($M+H$, $\text{C}_{30}\text{H}_{43}\text{N}_4\text{O}_5\text{Si}$ requires 567.2997). ^1H NMR (CDCl_3 /300 MHz): 8.70 (s, 1H); 8.12 (s, 1H); 7.97 (d, 2H); 7.60 (s, 1H); 7.29 (d, 1H); 6.95 (d, 2H); 4.97 (br, 1H); 4.77 (t, 2H); 4.40 (q, 2H); 3.63 (br, 2H); 1.40 (t, 3H); 1.38 (s, 9H); 1.00 (s, 9H); 0.25 (s, 6H).

[000337] Anal. Calculated for $\text{C}_{30}\text{H}_{42}\text{N}_4\text{O}_5\text{Si}$: C, 63.58; H, 7.47; N, 9.89. Found: C, 63.51; H, 7.58; N, 9.73.

EXAMPLE 85

[000338] This example illustrates the preparation of ethyl 1-(2-aminoethyl)-3-[2-(4-[[*tert*-butyl(dimethyl)silyl]oxy}phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride.

[000339] Ethyl 1-{2-[(*tert*-butoxycarbonyl)amino]ethyl}-3-[2-(4-[[*tert*-butyl(dimethyl)-silyl]oxy}phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate (1.0 g, 0.0018 mol) and 4N HCl in dioxane were mixed for 2 hours and filtered to give the desired product as a white solid, 875 mg (90% yield). FABHRMS m/z 467.2450 ($M+H$, $\text{C}_{25}\text{H}_{35}\text{N}_4\text{O}_3\text{Si}$ requires 467.2473). ^1H NMR ($\text{DMSO}-d_6$ + TFA/300 MHz): 8.80 (d, 1H); 8.75 (s, 1H); 8.35 (d, 1H); 8.22 (br, 3H); 8.20-8.10 (m, 3H); 7.13 (d, 2H); 4.90 (t, 2H); 4.40 (q, 2H); 3.40 (q, 2H); 1.39 (t, 3H); 0.95 (s, 9H); 0.23 (s, 6H).

[000340] Anal. Calculated for $\text{C}_{25}\text{H}_{34}\text{N}_4\text{O}_3\text{Si}$ (2 HCl, 1.1 H_2O): C, 53.68; H, 6.88; N, 10.02. Found: C, 53.34; H, 6.98; N, 10.18.

EXAMPLE 86

[000341] This example illustrates the production of 2-[2-(4-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

[000342] Ethyl 1-(2-aminoethyl)-3-[2-(4-[[*tert*-butyl(dimethyl)silyl]oxy}phenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride (770 mg, 0.0014 mol), conc ammonium hydroxide (2 mL)

and methanol (20 mL) were stirred overnight. Contents were concentrated *in vacuo* and the residue was slurried in water and filtered to give the desired product as a white solid, 373 mg, (87% yield). FABHRMS *m/z* 307.1222 (*M*+*H*, C₁₇H₁₅N₄O₂ requires 307.1190). ¹H NMR (DMSO-*d*₆ + TFA/300 MHz): 8.75 (d, 1H); 8.63 (s, 1H); 8.40 (s, 1H); 8.21 (d, 1H); 8.00-7.95 (m, 3H); 7.00 (d, 2H); 4.43 (t, 2H); 3.65 (br, 2H).

[000343] Anal. Calculated for C₁₇H₁₄N₄O₂ (1.3 H₂O): C, 61.92; H, 5.07; N, 16.99. Found: C, 61.79; H, 5.11; N, 16.85.

EXAMPLE 87

[000344] This example illustrates the production of 2-{2-[4-(2-morpholin-4-ylethoxy)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-*a*]pyrazin-4(5H)-one.

[000345] 2-[2-(4-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-*a*]pyrazin-4(5H)-one (500 mg, 0.0016 mol), 4-(2-chloroethyl)morpholine hydrochloride (372 mg (0.002 mol) and potassium carbonate (600 mg, 0.004 mol) were heated in DMF (20 mL) at 80°C for 3 hours. Contents were allowed to cool, diluted with water (50 mL), cooled to 0°C and filtered to give the desired product as a white solid, 515 mg (77% yield).

FABHRMS *m/z* 420.2004 (*M*+*H*, C₂₃H₂₆N₅O₃ requires 420.2030). ¹H NMR (DMSO-*d*₆ + TFA/300 MHz): 8.81 (d, 1H); 8.66 (s, 1H); 8.43 (s, 1H); 8.23 (d, 1H); 8.15 (d, 2H); 7.94 (s, 1H); 7.28 (d, 2H); 4.55-4.40 (m, 4H); 4.05-3.95 (m, 2H); 3.80-3.50 (m, 8H); 3.30-3.18 (m, 2H).

[000346] Anal. Calculated for C₂₃H₂₅N₅O₃ (0.8 H₂O): C, 63.60; H, 5.96; N, 16.25. Found: C, 63.67; H, 6.18; N, 16.14.

EXAMPLE 88

[000347] This example illustrates the production of 2-(2-{4-[2-(dimethylamino)ethoxy]-phenyl}pyridin-4-yl)-6,7-dihydropyrazolo[1,5-*a*]pyrazin-4(5H)-one.

[000348] 2-(2-{4-[2-(dimethylamino)ethoxy]phenyl}pyridin-4-yl)-6,7-dihydropyrazolo[1,5-*a*]pyrazin-4(5H)-one was prepared according to the procedure of 2-{2-[4-(2-morpholin-4-ylethoxy)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-*a*]pyrazin-4(5H)-one to give the desired product as a

white solid (72% yield). FABHRMS m/z 378.1901 ($M+H$, $C_{21}H_{24}N_5O_2$ requires 378.1925). 1H NMR (DMSO- d_6 + TFA/300 MHz): 9.80 (br, 1H); 8.80 (d, 1H); 8.70 (s, 1H); 8.42 (s, 1H); 8.28 (d of d, 1H); 8.12 (d, 2H); 7.96 (s, 1H); 7.28 (d, 2H); 4.50-4.40 (m, 4H); 3.70 (br, 2H); 3.60 (br, 2H); 2.90 (s, 6H). Anal. Calculated for $C_{21}H_{23}N_5O_2$ (0.6 H_2O): C, 64.91; H, 6.15; N, 18.03. Found: C, 64.97; H, 6.28; N, 18.04.

EXAMPLE 89

[000349] This example illustrates the production of Ethyl 1-(2-aminoethyl)-3-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride.

[000350] Ethyl 1-(2-[(tert-butoxycarbonyl)amino]ethyl)-3-(2-chloropyridin-4-yl)-1H-pyrazole-5-carboxylate and 3-benzyloxyboronic acid were reacted according to the procedure described for the preparation of ethyl 1-(2-[(tert-butoxycarbonyl)-amino]ethyl)-3-[2-(4-[(tert-butyl(dimethyl)silyl]oxy)phenyl]pyridin-4-yl)-1H-pyrazole-5-carboxylate, to give ethyl 1-(2-t-butoxycarbonylaminoethyl)-3-{2-[3-(benzyloxy-phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate as a light amber oil. Ethyl 1-(2-t-butoxycarbonylaminoethyl)-3-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate and 4N HCl in dioxane were stirred 3 hours and filtered to give the desired product as a pale yellow solid, 8.0 g (81% yield). FABHRMS m/z 443.2053 ($M+H$, $C_{26}H_{27}N_4O_3$ requires 443.2078). 1H NMR (DMSO- d_6 + TFA/300 MHz): 8.88 (d, 1H); 8.80 (s, 1H); 8.10 (s, 1H); 8.05 (br, 3H); 7.78 (s, 1H); 7.65 (d, 1H); 7.55 (t, 1H); 7.45 (d, 1H); 7.40-7.30 (m, 6H); 5.21 (s, 2H); 4.85 (t, 2H); 4.35 (q, 2H); 3.40 (q, 2H); 1.35 (t, 3H).

[000351] Anal. Calculated for $C_{26}H_{26}N_4O_3$ (3HCl, H_2O): C, 54.80; H, 5.48; N, 9.83. Found: C, 55.01; H, 5.84; N, 10.75.

EXAMPLE 90

[000352] This example illustrates the production of 2-{2-[3-(benzyloxy)phenyl]-pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

[000353] 2-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one was prepared according to the procedure for the preparation of 2-[2-(4-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one, using ethyl 1-(2-aminoethyl)-3-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride, to give the desired product as a white solid (63% yield). FABHRMS m/z 397.1634 ($M+H$, $C_{24}H_{21}N_4O_2$ requires 397.1659). 1H NMR ($DMSO-d_6$ + TFA/300 MHz): 8.81 (d, 1H); 8.62 (s, 1H); 8.40 (s, 1H); 8.21 (d, 1H); 7.92 (s, 1H); 7.80 (s, 1H); 7.70 (d, 1H); 7.63 – 7.25 (m, 6H); 5.23 (s, 2H); 4.51-4.40 (m, 2H); 3.78-3.60 (m, 2H). Anal. Calculated for $C_{24}H_{20}N_4O_2$ (H_2O): C, 69.55; H, 5.35; N, 13.52. Found: C, 69.65; H, 5.11; N, 14.50.

EXAMPLE 91

[000354] This example illustrates the production of ethyl 1-(2-aminoethyl)-3-[2-(3-hydroxyphenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylate dihydrochloride.

[000355] Ethyl 1-(2-*t*-butoxycarbonylaminoethyl)-3-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate (9.1 g, 0.017 mol), prepared as for ethyl 1-(2-aminoethyl)-3-{2-[3-(benzyloxy)phenyl]pyridin-4-yl}-1H-pyrazole-5-carboxylate dihydrochloride and 10% palladium/carbon (2.0 g) in ethanol (150 mL) were shaken at 55 psi H_2 on a Parr hydrogenator for three and a half days. Contents were filtered through clay and the filtrate was concentrated *in vacuo* leaving a pale yellow solid (6.4 g). The solid and 4N HCl in dioxane were stirred overnight and filtered to give the desired product as a white solid, 6.0 g (83% yield). HRMS calculated for ($M + H$) 353.1608, found 353.1630. 1H NMR ($DMSO-d_6$ + TFA/300 MHz): 8.86 (d, 1H); 8.77 (s, 1H); 8.40 (d, 1H); 8.20 (br s, 3H); 8.12 (s, 1H); 7.59-7.40 (m, 3H); 7.09 (d, 1H); 4.90 (t, 2H); 4.39 (q, 2H); 3.40 (q, 2H); 1.35 (t, 3H).

EXAMPLE 92

[000356] This example illustrates the production of 2-{2-[3-(2-morpholin-4-ylethoxy)-phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one.

[000357] 2-{2-[3-(2-morpholin-4-ylethoxy)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo-[1,5-a]pyrazin-4(5H)-one was prepared according to the procedure for 2-{2-[4-(2-morpholin-4-ylethoxy)phenyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one, using 2-[2-(3-hydroxyphenyl)pyridin-4-yl]-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one and chloroethylmorpholine hydrochloride to give the desired as a white solid (55% yield). FABHRMS m/z 420.1996 ($M+H$, $C_{23}H_{26}N_5O_3$ requires 420.2030). 1H NMR ($DMSO-d_6 + TFA/300$ MHz): 8.85 (d, 1H); 8.61 (s, 1H); 8.40 (s, 1H); 8.21 (d, 1H); 7.89 (s, 1H); 7.78 (s, 2H); 7.59 (t, 1H); 7.28 (d, 1H); 4.58-4.40 (m, 4H); 4.08-3.91 (m 2H); 3.84-3.48 (m, 8H); 3.45-3.13 (m, 2H).

[000358] Anal. Calculated for $C_{23}H_{25}N_5O_3$ ($0.7 H_2O$): C, 63.93; H, 6.16; N, 16.21. Found: C, 63.93; H, 5.96; N, 16.42.

EXAMPLE 93

[000359] This example illustrates the preparation of 2-(2-Chloropyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one.

[000360] Step 1. Preparation of 2-Chloro-4-hydrazinopyridine hydrochloride. A solution of 4-amino-2-chloropyridine (1.0 g, 7.78 mmol) in 20% sulfuric acid (20 mL) was cooled to 0 °C and treated with a solution of sodium nitrite (564 mg, 8.17 mmol) in water (3 mL) at a rate such that the reaction temperature did not exceed 10 °C. After 15 minutes, the solution was added to a 0°C suspension of tin(II) chloride in 20% sulfuric acid (20 mL). The frothy suspension was stirred for 15 minutes at 0 °C and then warmed to room temperature over 15 minutes. The mixture was poured into 100 mL of ice water and made basic with concentrated ammonium hydroxide. The product was extracted with diethyl ether and ethyl acetate repeatedly. The organic layers were dried (sodium sulfate) and concentrated to give crude 2-chloro-4-hydrazinopyridine as a yellow solid (830 mg, 5.78 mmol). The solid was dissolved in tetrahydrofuran (5 mL) and diluted with diethyl ether (15 mL). The solution was treated with 1 N HCl in diethyl ether (5.8 mL, 5.8 mmol). The white precipitate was filtered and washed with ether to give 2-chloro-4-hydrazinopyridine

hydrochloride as a white solid (995 mg, 5.53 mmol, 71% yield). LC-MS (ES+) $MH^+ = 144$. 1H NMR (300 MHz, DMSO- d_6) δ 10.0-9.40 (br s, 4H), 8.07 (d, $J = 6.1$, 1H), 6.95 (d, $J = 1.9$, 1H), 6.86 (dd, $J = 5.8$, 2.0, 1H).

[000361] Step 2. Preparation of Piperidine-2,4-dione 4-[(2-chloropyridin-4-yl)hydrazone]. A mixture of 2-chloro-4-hydrazinopyridine hydrochloride (961 mg, 5.33 mmol), piperdiene-2,4-dione (Example 1, step 3) (604 mg, 5.33 mmol) and ethanol (20 mL) was refluxed overnight. The reaction mixture was cooled to room temperature, diluted with diethyl ether (20 mL), and filtered. The precipitate was washed with 50% ethanol/diethyl ether and dried to give piperidine-2,4-dione 4-[(2-chloropyridin-4-yl)hydrazone] as an off-white solid (940 mg, 3.94 mmol, 74% yield). LC-MS (ES+) $MH^+ = 239$.

[000362] Step 3. Preparation of 2-(2-Chloropyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one.

[000363] A mixture of piperidine-2,4-dione 4-[(2-chloropyridin-4-yl)hydrazone] (863 mg, 3.62 mmol) in dimethylformamide dimethyl acetal (16 mL) was refluxed for 1 hour. The solvent was removed under reduced pressure. The residue was suspended in ethanol/diethyl ether and filtered. The precipitate was washed with 50% ethanol/diethyl ether to give 2-(2-chloropyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one as an off-white solid (378 mg, 1.52 mmol, 42% yield). The mother liquor was concentrated and purified by flash chromatography (0 $\rightarrow$ 10% methanol/ethyl acetate). The resultant oil was triturated with methanol/ether to give another 58 mg (0.23 mmol, 6%) of 2-(2-chloropyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one. 1H NMR (300 MHz, DMSO- d_6) δ 9.17 (s, 1H), 8.47 (d, $J = 5.7$, 1H), 8.04 (d, $J = 1.6$, 1H), 7.94 (dd, $J = 5.5$, 1.8, 1H), 7.71 (br s, 1H), 3.44 (td, $J = 6.5$, 2.6, 2H), 2.89 (t, $J = 6.6$, 2H). HRMS calculated for $C_{11}H_{10}ClN_4O$ (MH^+) 249.0538, found 249.0545.

EXAMPLE 94

[000364] This example illustrates the preparation of 2-(2-quinolin-3-ylpyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one bis(trifluoroacetate).

5 **[000365]** The title compound was prepared from 2-(2-chloropyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one (Example 508) and 3-quinolinylboronic acid by the procedure described for Example 2. ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.76 (d, *J* = 2.2, 1H), 9.38 (s, 1H), 9.26 (d, *J* = 2.0, 1H), 8.82 (d, *J* = 5.4, 1H), 8.71 (d, *J* = 1.6, 1H), 8.16 (d, *J* = 7.7, 1H), 8.12 (d, *J* = 8.3, 1H), 7.96 (dd, *J* = 5.6, 2.0, 1H), 7.87 (td, *J* = 7.7, 1.3, 1H), 7.76-7.68 (m, 2H), 3.48 (td, *J* = 6.6, 2.4, 2H), 2.95 (t, *J* = 6.6, 2H). HRMS calculated for C₂₀H₁₆N₅O (MH⁺) 342.1349, found 342.1334.

[000366] The following examples were made by the same method.

Example No.	Compound Name(s)	Calculated (m+H)	Found (m+H)
95	2-[2-(2-fluorophenyl)pyridin-4-yl]-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one trifluoroacetate	309.1146	309.119
96	2-(2-phenylpyridin-4-yl)-2,5,6,7-tetrahydro-4H-pyrazolo[4,3-c]pyridin-4-one trifluoroacetate	291.124	291.1252

EXAMPLE 97

[000367] This example illustrates the preparation of 2-{2-[(E)-2-(4-morpholin-4-ylphenyl)vinyl]pyridin-4-yl}-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one trifluoroacetate.

[000368] mp 290°C (decomposition); ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.62 (d, *J* = 6.5 Hz, 1H), 8.39 (2 x s, 2H), 7.97-7.83 (m, 2H), 7.70 (s, 1H), 7.57 (d, *J* = 8.6 Hz, 2H), 7.15 (d, *J* = 16.2 Hz, 1H), 7.03 (d, *J* = 16.2 Hz, 2H), 4.50-4.38 (m, 2H), 3.80-3.60 (m, 6H), 3.28-3.19 (m, 4H); *m/z* 402 [M+H]⁺.

EXAMPLE 98

[000369] This example illustrates that MK2 knock-out mice (MK2 (-/-)) are resistant to the formation of K/BN serum-induced arthritis and that compounds that inhibit MK-2 should be effective for the prevention and treatment of TNF α -mediated diseases or disorders.

A strain of mice has been reported that develops symptoms similar to human rheumatoid arthritis. The mice were designated K/BxN mice. See, Wipke, B. T. and P. M. Allen, *J. of Immunology*, 167:1601 - 1608 (2001). Serum from the mice can be injected into host animals to provoke a typical RA response. The progression of the RA symptoms in the mice is measured by measuring paw thickness as a function of time.

[000370] In the present example, host mice having normal MK-2 production (MK2 (+/+)) were genetically altered by disabling the gene encoding MK-2 to produce mice having no capability of endogenous synthesis of active MK-2 (MK2 (-/-)). Normal host mice (MK2 (+/+)) and MK-2 knock-out mice (MK2 (-/-)), were separated into four groups with each group containing both male and female mice. All groups of mice were treated similarly, except that one group (Normal), composed of MK2 (+/+) mice that served as the control group, was not injected with serum from K/BxN mice, while the other three groups were injected with K/BxN serum at day 0. The other three groups of mice were MK2 (+/+), MK2 (-/-), and Anti-TNF. The Anti-TNF group was composed of MK2 (+/+) mice which were also injected at day) with anti-TNF antibody. The paw thickness of all mice was measured immediately after the injections on day 0, and then on each successive day thereafter for 7 days.

[000371] Figure 1 is a graph that shows paw thickness as a function of time from day 0 to day 7 for MK2 (+/+) and MK2 (-/-) mice, which have received serum injection. It can be seen that paw thickness increased significantly for MK2(+/+) mice, whereas there was substantially no increase in paw thickness for MK2 knock-out mice. This indicated the requirement for a functioning MK2 regulatory system to the inflammatory response caused by the serum challenge. When anti-TNF antibody was

administered to the MK2 (+/+) mice along with the serum injection, the swelling response was significantly reduced. This can be seen in Figure 2, which is a bar chart showing paw thickness at seven days after injection for normal mice, MK2 (+/+) mice receiving serum, MK2 (-/-) mice receiving serum, and MK2 (+/+) mice receiving serum and anti-TNF antibody.

[000372] This data shows that the MK2 knock-out mice show no arthritic response to a serum challenge, whereas MK2 (+/+) mice show a normal response. Treatment of MK2 (+/+) mice that receive a serum challenge with anti-TNF antibody reduces the response back to near-normal levels.

This illustrates the utility of the MK2 regulatory system as a potential control point for the modulation of TNF production, and indicates that such regulation could serve as a treatment for inflammation -- such as that caused by arthritis, for example. It further shows that MK2 inhibition can have a beneficial effect on inflammation, and indicates that administration of an MK2 inhibitor can be an effective method of preventing or treating TNF modulated diseases or disorders.


[000373] All references cited in this specification, including without limitation all papers, publications, patents, patent applications, presentations, texts, reports, manuscripts, brochures, books, internet postings, journal articles, periodicals, and the like, are hereby incorporated by reference into this specification in their entireties. The discussion of the references herein is intended merely to summarize the assertions made by their authors and no admission is made that any reference constitutes prior art. Applicants reserve the right to challenge the accuracy and pertinency of the cited references.

[000374] In view of the above, it will be seen that the several advantages of the invention are achieved and other advantageous results obtained.

[000375] As various changes could be made in the above methods and compositions without departing from the scope of the invention, it is intended that all matter contained in the above description shall be interpreted as illustrative and not in a limiting sense.

WHAT IS CLAIMED IS:

1. A compound having the structure:




5 wherein:

Z^2 and Z^3 are nitrogen, Z^1 , Z^4 and Z^5 are carbon, and join with Z^2 and Z^3 to form a pyrazole ring, or optionally, Z^4 and Z^5 are nitrogen, Z^1 , Z^2 and Z^3 are carbon and join with Z^4 and Z^5 to form a pyrazole ring;

R^a is selected from:


10

1)




2)

15



3)



where dashed lines indicate optional single or double bonds;

when R^a is ring M and ring M is aromatic, M^1 is carbon and is substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon and nitrogen and is unsubstituted or substituted with $(L)_nR^1$;

when ring M is partially saturated, M^1 is carbon and is mono- or di-substituted with $(L)_nR^1$, M^5 is carbon, and each of M^2 , M^3 , M^4 and M^6 is independently selected from carbon, nitrogen, oxygen and sulfur, and when M^2 , M^3 , M^4 , or M^6 is oxygen or sulfur, it is unsubstituted, and when M^2 , M^3 , M^4 or M^6 is carbon or nitrogen, it is optionally unsubstituted; or mono- or di-substituted with $(L)_nR^1$;

when R^a is ring Q and ring Q is aromatic, Q^1 is selected from carbon and nitrogen, and when Q^1 is carbon, it is substituted with $(L)_nR^1$, and when Q^1 is nitrogen, it is unsubstituted, Q^4 is selected from nitrogen and carbon, and each of Q^2 , Q^3 and Q^5 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

optionally when ring Q is aromatic, Q^1 is carbon and is substituted with $(L)_nR^1$, Q^4 is carbon, and one of Q^2 , Q^3 and Q^5 is optionally oxygen or sulfur, and the remainder of Q^2 , Q^3 and Q^5 are independently selected from nitrogen and carbon, and if carbon, are substituted with $(L)_nR^1$;

when ring Q is partially saturated, Q^1 is selected from carbon and nitrogen, and if carbon, it is mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$, Q^4 is selected from carbon and nitrogen, but only one of Q^1 and Q^4 can be nitrogen, each of Q^2 , Q^3 and Q^5 is independently selected from carbon, nitrogen, oxygen and sulfur, and if oxygen or sulfur, it is unsubstituted, and if carbon, it is

mono- or di-substituted with $(L)_nR^1$, and if nitrogen, it is unsubstituted or substituted with $(L)_nR^1$;

when R^a is structure 3, it is fully conjugated, X^2 is selected from oxygen or nitrogen substituted with $(L)_nR^1$, X^1 is carbon and is substituted with $(L)_nR^1$, and each of X^5 and X^6 is independently selected from nitrogen and carbon, and if carbon, it is substituted with $(L)_nR^1$;

R^1 is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl- R^{11} , C₂-C₆ alkenyl- R^{11} , C₂-C₆ alkynyl- R^{11} , C₁-C₆ alkyl- $(R^{11})_2$, C₂-C₆ alkenyl- $(R^{11})_2$, CSR¹¹, amino, CONHR¹¹, NHR⁷, NR⁸R⁹, N(R⁷)-N(R⁸)(R⁹), C(R¹¹)=N-N(R⁸)(R⁹), N=N(R⁷), N(R⁷)-N=C(R⁸), C(R¹¹)=N-O(R¹⁰), ON=C(R¹¹), C₁-C₆ alkyl-NHR⁷, C₁-C₆ alkyl-NR⁸R⁹, (C₁-C₄)alkyl-N(R⁷)-N(R⁸)(R⁹), (C₁-C₄)alkylC(R¹¹)=N-N(R⁸)(R⁹), (C₁-C₄)alkyl-N=N(R⁷), (C₁-C₄)alkyl-N(R⁷)-N=C(R⁸), nitro, cyano, CO₂R¹¹, O-R¹⁰, C₁-C₄ alkyl-OR¹⁰, COR¹¹, SR¹⁰, SSR¹⁰, SOR¹¹, SO₂R¹¹, C₁-C₆ alkyl-COR¹¹, C₁-C₆ alkyl-SR¹⁰, C₁-C₆ alkyl-SOR¹¹, C₁-C₆ alkyl-SO₂R¹¹, halo, Si(R¹¹)₃, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ;

R^7 , R^8 and R^9 are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl- R^{11} , C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, O-R¹⁵, C₁-C₄ alkyl-OR¹⁵, CO₂R¹⁵, C(S)OR¹⁵, C(O)SR¹⁵, C(O)R¹⁷, C(S)R¹⁷, CONHR¹⁶, C(S)NHR¹⁶, CON(R¹⁶)₂, C(S)N(R¹⁶)₂, SR¹⁵, SOR¹⁷, SO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl,

alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹⁰ is selected from -H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, C₁-C₄ alkyl-OR¹⁵, CSR¹¹, CO₂R¹⁵, C(S)OR¹⁵, C(O)SR¹⁵, COR¹⁷, C(S)R¹⁷, CONHR¹⁶, C₁-C₄ alkyl-R¹¹, C₁-C₄ alkyl-NH₂R¹³, C(S)NHR¹⁶, O-R¹⁵, CON(R¹⁶)₂, C(S)N(R¹⁶)₂, SOR¹⁷, SO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, Si(R¹³)₂R¹⁷, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹¹ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₂-C₆ alkenyl, C₂-C₆ alkynyl, amino, NHR¹³, NR¹³R¹⁴, N=NR¹³, C₁-C₆ alkyl-NHR¹³, C₁-C₆ alkyl-NR¹³R¹⁴, O-R¹⁵, C₁-C₄ alkyl-OR¹⁵, SR¹⁵, COR¹³, CO₂R¹⁷, C₁-C₆ alkyl-CO₂R¹⁵, C₁-C₆ alkyl-C(S)OR¹⁵, C₁-C₆ alkyl-C(O)SR¹⁵, C₁-C₆ alkyl-COR¹⁷, C₁-C₆ alkyl-C(S)R¹⁷, C₁-C₆ alkyl-CONHR¹⁶, C₁-C₆ alkyl-C(S)NHR¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, C₁-C₆ alkyl-C(S)N(R¹⁶)₂, C₁-C₆ alkyl-SR¹⁵, C₁-C₆ alkyl-SOR¹⁷, C₁-C₆ alkyl-SO₂R¹⁷, halo, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹² is selected from -H, OH, oxo, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R¹¹, C₂-C₁₀ alkenyl-R¹¹, C₂-C₁₀ alkynyl-R¹¹, C₁-C₁₀

alkyl-(R¹¹)₂, C₂-C₁₀ alkenyl-(R¹¹)₂, CSR¹¹, hydroxyl C₁-C₆ alkyl-R¹¹, amino C₁-C₄ alkyl-R⁷, amino, NHR⁷, NR⁸R⁹, N(R⁷)-N(R⁸)(R⁹), C(R¹¹)=N-N(R⁸)(R⁹), N=N(R⁷), N(R⁷)-N=C(R⁸), C(R¹¹)=N-O(R¹⁰), ON=C(R¹¹), C₁-C₁₀ alkyl-NHR⁷, C₁-C₁₀ alkyl-NR⁸R⁹, (C₁-C₁₀)alkyl-N(R⁷)-N(R⁸)(R⁹), (C₁-C₁₀)alkylC(R¹¹)=N-N(R⁸)(R⁹), (C₁-C₁₀)alkyl-N=N(R⁷), (C₁-C₁₀)alkyl-N(R⁷)-N=C(R⁸), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R¹⁰, C₁-C₁₀ alkyl-OR¹⁰, COR¹¹, CO₂R¹¹, SR¹⁰, SSR¹⁰, SOR¹¹, SO₂R¹¹, C₁-C₁₀ alkyl-COR¹¹, C₁-C₁₀ alkyl-SR¹⁰, C₁-C₁₀ alkyl-SOR¹¹, C₁-C₁₀ alkyl-SO₂R¹¹, halo, Si(R¹¹)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹⁸;

R¹³ and R¹⁴ are each independently selected from -H, oxo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R²³, C₁-C₆ alkyl-NHR¹⁹, C₁-C₆ alkyl-NR¹⁹R²⁰, O-R²¹, C₁-C₄ alkyl-OR²¹, CO₂R²¹, COR²¹, C(S)OR²¹, C(O)SR²¹, C(O)R²³, C(S)R²³, CONHR²², C(S)NHR²², CON(R²²)₂, C(S)N(R²²)₂, SR²¹, SOR²³, SO₂R²³, C₁-C₆ alkyl-CO₂R²¹, C₁-C₆ alkyl-C(S)OR²¹, C₁-C₆ alkyl-C(O)SR²¹, C₁-C₆ alkyl-COR²³, C₁-C₆ alkyl-C(S)R²³, C₁-C₆ alkyl-CONHR²², C₁-C₆ alkyl-C(S)NHR²², C₁-C₆ alkyl-CON(R²²)₂, C₁-C₆ alkyl-C(S)N(R²²)₂, C₁-C₆ alkyl-SR²¹, C₁-C₆ alkyl-SOR²³, C₁-C₆ alkyl-SO₂R²³, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁵ and R¹⁶ are independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR¹⁹, C₁-C₆ alkyl-NR¹⁹R²⁰, C₁-C₄

alkyl-OR²¹, CSR¹¹, CO₂R²², COR²³, CONHR²², CON(R²²)₂, SOR²³,
 SO₂R²³, C₁-C₆ alkyl-CO₂R²², C₁-C₆ alkyl-COR²³, C₁-C₆ alkyl-CONHR²², C₁-
 C₆ alkyl-CON(R²²)₂, C₁-C₆ alkyl-SR²¹, C₁-C₆ alkyl-SOR²³, C₁-C₆ alkyl-
 SO₂R²³, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl,
 5 alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl,
 heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl,
 heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl,
 arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic
 cycloalkyl are optionally substituted with one or more of the groups defined
 10 by R²⁴;

R¹⁷ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkenyl-
 R¹⁹, C₁-C₆ alkyl-R¹⁹, C₂-C₆ alkynyl, amino, NHR¹⁹, NR¹⁹R²⁰, C₁-C₆ alkyl-
 NHR¹⁹, C₁-C₆ alkyl-NR¹⁹R²⁰, O-R²¹, C₁-C₄ alkyl-OR²¹, SR²¹, C₁-C₆ alkyl-
 CO₂R²¹, C₁-C₆ alkyl-C(S)OR²¹, C₁-C₆ alkyl-C(O)SR²¹, C₁-C₆ alkyl-COR²³,
 15 C₁-C₆ alkyl-C(S)R²³, C₁-C₆ alkyl-CONHR²², C₁-C₆ alkyl-C(S)NHR²², C₁-C₆
 alkyl-CON(R²²)₂, C₁-C₆ alkyl-C(S)N(R²²)₂, C₁-C₆ alkyl-SR²¹, C₁-C₆ alkyl-
 SOR²³, C₁-C₆ alkyl-SO₂R²³, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl,
 alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl,
 heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl,
 20 heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl,
 arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic
 cycloalkyl are optionally substituted with one or more of the groups defined
 by R²⁴;

R¹⁸ is selected from -H, oxo, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-
 25 C₁₀ alkynyl, C₁-C₁₀ alkyl-R²³, C₂-C₁₀ alkenyl-R²³, C₂-C₁₀ alkynyl-R²³, C₁-C₁₀
 alkyl-(R²³)₂, C₂-C₁₀ alkenyl-(R²³)₂, CSR²³, amino, NHR¹⁹, NR²⁰R²⁰, N(R¹⁹)-
 N(R²⁰)(R²⁰), C(R²³)=N-N(R²⁰)(R²⁰), N=N(R¹⁹), N(R¹⁹)-N=C(R²⁰), C(R²³)=N-
 O(R²¹), ON=C(R²³), C₁-C₁₀ alkyl-NHR¹⁹, C₁-C₁₀ alkyl-NR²⁰R²⁰, (C₁-
 C₁₀)alkyl-N(R¹⁹)-N(R²⁰)(R²⁰), (C₁-C₁₀)alkylC(R²³)=N-N(R²⁰)(R²⁰), (C₁-
 30 C₁₀)alkyl-N=N(R¹⁹), (C₁-C₁₀)alkyl-N(R¹⁹)-N=C(R²⁰), SCN, NCS, C₁-C₁₀
 alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R²¹, C₁-C₁₀ alkyl-OR²¹,
 COR²³, CO₂R²³, SR²¹, SSR²¹, SOR²³, SO₂R²³, C₁-C₁₀ alkyl-COR²³, C₁-C₁₀

alkyl-SR²¹, C₁-C₁₀ alkyl-SOR²³, C₁-C₁₀ alkyl-SO₂R²³, halo, Si(R²³)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R²⁴;

R¹⁹ and R²⁰ are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl-R²⁹, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, O-R²⁷, C₁-C₄ alkyl-OR²⁷, CO₂R²⁷, C(S)OR²⁷, C(O)SR²⁷, C(O)R²⁹, C(S)R²⁹, CONHR²⁸, C(S)NHR²⁸, CON(R²⁸)₂, C(S)N(R²⁸)₂, SR²⁷, SOR²⁹, SO₂R²⁹, C₁-C₆ alkyl-CO₂R²⁷, C₁-C₆ alkyl-C(S)OR²⁷, C₁-C₆ alkyl-C(O)SR²⁷, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-C(S)R²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-C(S)NHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-C(S)N(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

R²¹ and R²² are independently selected from -H, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR²⁵, C₁-C₆ alkyl-NR²⁵R²⁶, C₁-C₄ alkyl-OR²⁷, CSR¹¹, CO₂R²⁸, COR²⁹, CONHR²⁸, CON(R²⁸)₂, SOR²⁹, SO₂R²⁹, C₁-C₆ alkyl-CO₂R²⁸, C₁-C₆ alkyl-COR²⁹, C₁-C₆ alkyl-CONHR²⁸, C₁-C₆ alkyl-CON(R²⁸)₂, C₁-C₆ alkyl-SR²⁷, C₁-C₆ alkyl-SOR²⁹, C₁-C₆ alkyl-SO₂R²⁹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁰;

cycloalkyl are optionally substituted with one or more of the groups defined by R^{30} ;

R^{23} is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkenyl- R^{25} , C_1 - C_6 alkyl- R^{25} , C_2 - C_6 alkynyl, amino, NHR^{25} , $NR^{25}R^{26}$, C_1 - C_6 alkyl- NHR^{25} , C_1 - C_6 alkyl- $NR^{25}R^{26}$, $O-R^{27}$, C_1 - C_4 alkyl- OR^{27} , SR^{27} , C_1 - C_6 alkyl- CO_2R^{27} , C_1 - C_6 alkyl- $C(S)OR^{27}$, C_1 - C_6 alkyl- $C(O)SR^{27}$, C_1 - C_6 alkyl- COR^{29} , C_1 - C_6 alkyl- $C(S)R^{29}$, C_1 - C_6 alkyl- $CONHR^{28}$, C_1 - C_6 alkyl- $C(S)NHR^{28}$, C_1 - C_6 alkyl- $CON(R^{28})_2$, C_1 - C_6 alkyl- $C(S)N(R^{28})_2$, C_1 - C_6 alkyl- SR^{27} , C_1 - C_6 alkyl- SOR^{29} , C_1 - C_6 alkyl- SO_2R^{29} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{30} ;

R^{24} is selected from -H, OH, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} alkyl- R^{29} , C_2 - C_{10} alkenyl- R^{29} , C_2 - C_{10} alkynyl- R^{29} , C_1 - C_{10} alkyl- $(R^{29})_2$, C_2 - C_{10} alkenyl- $(R^{29})_2$, CSR^{29} , amino, NHR^{25} , $NR^{26}R^{26}$, $N(R^{25})-N(R^{26})(R^{26})$, $C(R^{29})=N-N(R^{26})(R^{26})$, $N=N(R^{25})$, $N(R^{25})-N=C(R^{26})$, $C(R^{29})=N-O(R^{27})$, $ON=C(R^{29})$, C_1 - C_{10} alkyl- NHR^{25} , C_1 - C_{10} alkyl- $NR^{26}R^{26}$, (C_1-C_{10}) alkyl- $N(R^{25})-N(R^{26})(R^{26})$, (C_1-C_{10}) alkyl- $C(R^{29})=N-N(R^{26})(R^{26})$, (C_1-C_{10}) alkyl- $N=N(R^{25})$, (C_1-C_{10}) alkyl- $N(R^{25})-N=C(R^{26})$, SCN , NCS , C_1 - C_{10} alkyl SCN , C_1 - C_{10} alkyl NCS , nitro, cyano, $O-R^{27}$, C_1 - C_{10} alkyl- OR^{27} , CO_2R^{29} , COR^{29} , SR^{27} , SSR^{27} , SOR^{29} , SO_2R^{29} , C_1 - C_{10} alkyl- COR^{29} , C_1 - C_{10} alkyl- SR^{27} , C_1 - C_{10} alkyl- SOR^{29} , C_1 - C_{10} alkyl- SO_2R^{29} , halo, $Si(R^{29})_3$, halo C_1 - C_{10} alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{30} ;

R^{25} and R^{26} are each independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₄ alkyl- R^{35} , C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹ R^{32} , O- R^{33} , C₁-C₄ alkyl-OR³³, CO₂ R^{33} , C(S)OR³³, C(O)SR³³, C(O) R^{35} , C(S) R^{35} , CONHR³⁴, C(S)NHR³⁴, CON(R^{34})₂, C(S)N(R^{34})₂, SR³³, SOR³⁵, SO₂ R^{35} , C₁-C₆ alkyl-CO₂ R^{33} , C₁-C₆ alkyl-C(S)OR³³, C₁-C₆ alkyl-C(O)SR³³, C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-C(S) R^{35} , C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-C(S)NHR³⁴, C₁-C₆ alkyl-CON(R^{34})₂, C₁-C₆ alkyl-C(S)N(R^{34})₂, C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-SOR³⁵, C₁-C₆ alkyl-SO₂ R^{35} , halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{36} ;

R^{27} and R^{28} are independently selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹ R^{32} , C₁-C₄ alkyl-OR³³, CSR¹¹, CO₂ R^{34} , COR³⁵, CONHR³⁴, CON(R^{34})₂, SOR³⁵, SO₂ R^{35} , C₁-C₆ alkyl-CO₂ R^{34} , C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-CON(R^{34})₂, C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-SOR³⁵, C₁-C₆ alkyl-SO₂ R^{35} , halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{36} ;

R^{29} is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, R^{31} , C₁-C₆ alkyl- R^{31} , C₂-C₆ alkynyl, amino, NHR³¹, NR³¹ R^{32} , C₁-C₆ alkyl-NHR³¹, C₁-C₆ alkyl-NR³¹ R^{32} , O- R^{33} , C₁-C₄ alkyl-OR³³, SR³³, C₁-C₆ alkyl-CO₂ R^{33} , C₁-C₆ alkyl-C(S)OR³³, C₁-C₆ alkyl-C(O)SR³³, C₁-C₆ alkyl-COR³⁵, C₁-C₆ alkyl-C(S) R^{35} , C₁-C₆ alkyl-CONHR³⁴, C₁-C₆ alkyl-C(S)NHR³⁴, C₁-C₆ alkyl-CON(R^{34})₂, C₁-C₆ alkyl-C(S)N(R^{34})₂, C₁-C₆ alkyl-SR³³, C₁-C₆ alkyl-

SOR³⁵, C₁-C₆ alkyl-SO₂R³⁵, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R³⁰ is selected from -H, OH, C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkyl-R³⁵, C₂-C₁₀ alkenyl-R³⁵, C₂-C₁₀ alkynyl-R³⁵, C₁-C₁₀ alkyl-(R³⁵)₂, C₂-C₁₀ alkenyl-(R³⁵)₂, CSR³⁵, N=NR³¹, amino, NHR³¹, NR³²R³², N(R³¹)-N(R³²)(R³²), C(R³⁵)=N-N(R³²)(R³²), N=N(R³¹), N(R³¹)-N=C(R³²), C(R³⁵)=N-O(R³³), ON=C(R³⁵), C₁-C₁₀ alkyl-NHR³¹, C₁-C₁₀ alkyl-NR³²R³², (C₁-C₁₀)alkyl-N(R³¹)-N(R³²)(R³²), (C₁-C₁₀)alkylC(R³⁵)=N-N(R³²)(R³²), (C₁-C₁₀)alkyl-N=N(R³¹), (C₁-C₁₀)alkyl-N(R³¹)-N=C(R³²), SCN, NCS, C₁-C₁₀ alkyl SCN, C₁-C₁₀ alkyl NCS, nitro, cyano, O-R³³, C₁-C₁₀ alkyl-OR³³, COR³⁵, SR³³, SSR³³, SOR³⁵, SO₂R³⁵, C₁-C₁₀ alkyl-COR³⁵, C₁-C₁₀ alkyl-SR³³, C₁-C₁₀ alkyl-SOR³⁵, C₁-C₁₀ alkyl-SO₂R³⁵, halo, Si(R³⁵)₃, halo C₁-C₁₀ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

R³¹, R³², R³³ and R³⁴ are each independently selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R³⁶;

- R^{35} is selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, OH, alkoxy, amino, alkylamino, dialkylamino, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{36} ;
- R^{36} is selected from -H, alkyl, alkenyl, alkynyl, aminoalkyl, OH, alkoxy, amino, nitro, cyano, halo, alkylamino, dialkylamino, hydroxyalkyl, alkylamino alkyl, dialkylaminoalkyl, alkoxyalkyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heterocyclylalkyl, and heteroarylalkyl;
- L is selected from $C(R^{37})_2$, O, S, NR^{37} , $C=O$, $C=S$, $C=C(R^{37})_2$, SO , SO_2 , $N=NO$, $CR^{37}=CR^{37}$, $CR^{37}=N$, $N=CR^{37}$, $N=N$, $NO=N$, $C=ONR^{37}$, $C=SR^{37}$, $NR^{37}C=O$, $NR^{37}C=S$, $C=OO$, $C=OS$, $C=SO$, $C=SS$, $OC=O$, $SC=O$, $OC=S$, $SC=S$, $S(O)_m-(O,S,NR^{37})$, $(O,S,NR^{37})-S(O)_m$, $C=(O,S)-C=(O,S)$, aryl, heteroaryl, heterocyclyl, cycloalkyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heterocyclylalkyl, heteroarylalkyl, or C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ,
- n is an integer from 0 to 10;
- m is an integer from 1 to 2; and
- R^2 , R^3 , R^4 , R^5 and R^{37} are each independently absent, or selected from an R^1 group; and
- R^3 and R^4 optionally join to form a ring of 5, 6, 7, or 8 atoms, where the atoms in the ring are independently selected from Z^3 , Z^4 , O, S, $C=O$, $C=S$, $S=O$, SO_2 , C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

2. The compound according to claim 1, wherein when Z^2 and Z^3 are both nitrogen, R^4 is other than pyrrole, or optionally when Z^4 and Z^5 are both nitrogen and R^a is ring Q, Q^2 is other than nitrogen.

3. The compound according to claim 1, wherein R^a is selected
5 from an M-ring and a Q-ring.

4. The compound according to claim 1, wherein R^a is an M-ring.

5. The compound according to claim 4, wherein ring M is an aromatic pyridine or pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are independently
10 selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$.

6. The compound according to claim 1, wherein R^a is an M-ring, wherein ring M is an aromatic pyridine or pyrimidine ring, wherein M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$, M^5 is carbon, M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$.

7. The compound according to claim 1, wherein:
 R^a is an M-ring that is an aromatic pyridine or pyrimidine ring;
 M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$;
 M^5 is carbon;
 M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$; and
 R^3 and R^4 optionally join to form a ring of 5, 6, 7, or 8 atoms, where the atoms in the ring are independently selected from Z^3 , Z^4 , O, S, C=O, C=S, S=O, SO₂, C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

8. The compound according to claim 1, wherein:
 R^a is an M-ring that is an aromatic pyridine or pyrimidine;
 M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$;
 M^5 is carbon;
 M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$; and
 R^3 and R^4 optionally join to form a ring of 6 or 7 atoms, where the atoms in the ring are independently selected from Z^3 , Z^4 , C=O, C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

9. The compound according to claim 1, wherein:
 R^a is an M-ring that is an aromatic pyridine or pyrimidine;
 M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$;
 M^5 is carbon;
 M^2 and M^6 are independently selected from carbon and nitrogen and if carbon, the carbon is substituted with $(L)_nR^1$; and R^3 and R^4 optionally join to form a ring of 6 atoms, where the atoms in the ring are

independently selected from Z^3 , Z^4 , $C=O$, C that is mono or di-substituted with an R^1 group, and N that is unsubstituted or substituted with an R^1 group.

10. The compound according to claim 1, wherein:

5

R^a is an M-ring that is an aromatic pyridine or pyrimidine ring;

M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$;


M^5 is carbon;

M^2 and M^6 are independently selected from carbon and nitrogen


and if carbon, the carbon is substituted with $(L)_nR^1$; and

10

R^3 and R^4 optionally join to form a ring that is selected from:



, and



15

11. The compound according to claim 1, wherein:

R^a is an M-ring that is an aromatic pyridine;

M^1 , M^3 , M^4 and M^6 are carbon and are substituted with

$(L)_nR^1$;

M^5 is carbon; and

20

M^2 is nitrogen.

12. The compound according to claim 1, wherein:

R^a is an M-ring that is an aromatic pyrimidine:

M^1 , M^3 and M^4 are carbon and are substituted with $(L)_nR^1$;

M^5 is carbon; and

25

M^2 and M^6 are nitrogen.

13. The compound according to claim 1, wherein:

R^a is an M-ring that is an aromatic pyridine:

M^1 , M^3 , M^4 and M^6 are carbon and are substituted with $(L)_nR^1$;

M^5 is carbon;

5

M^2 is nitrogen;

R^1 is selected from -H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyl- R^{11} , C_1 - C_6 alkoxy- R^{11} , COR^{17} , CO_2R^7 , $CONHR^7$, $N(R^8)_2$, amino C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkyl, amino, amino C_1 - C_4 alkyl- R^7 , halo C_1 - C_4 alkyl, C_1 - C_6 alkyl- NHR^7 , carbonitrile, SR^{10} , halo, NHR^7 , NR^8R^9 , NHR^7 - C_1 - C_6 alkyl, NR^8R^9 - C_1 - C_6 alkyl, nitro, cyano, $O-R^{10}$, C_1 - C_4 alkyl- OR^{10} , C_1 - C_6 alkyl- COR^{11} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, or C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{12} ;

10

15

R^7 and R^8 are each independently selected from -H, C_1 - C_6 alkyl, C_1 - C_4 alkyl- R^{11} , C_1 - C_6 alkyl- $N(R^{13})_2$, CO_2R^{16} , COR^{17} , aryl, and arylalkyl, wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

20

R^9 and R^{10} are each independently selected from -H, hydroxyl, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{17} , C_1 - C_6 alkyl- NH_2R^{13} , CO_2R^{16} , COR^{17} , C_1 - C_6 alkyl- CO_2R^{16} , C_1 - C_6 alkyl- $CONH-R^{16}$, C_1 - C_6 alkyl- $CON(R^{16})_2$, hydroxy C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, halo C_1 - C_4 alkyl, $Si(R^{13})_2R^{17}$, aryl, heteroaryl, heterocyclyl, arylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

25

R^{11} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxyl, halo, amino, NHR^{13} , $N(R^{13})_2$, COR^{13} , CO_2R^{17} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

30

R^{12} is selected from -H, hydroxyl, oxo, C_1 - C_6 alkyl, hydroxyl C_1 - C_6 alkyl- R^{11} , C_1 - C_{10} alkoxy, amino, amino C_1 - C_4 alkyl- R^7 , NHR^7 , $N(R^7)_2$, C_1 - C_6 alkyl- NHR^7 , C_1 - C_6 alkyl- NHR^8R^9 , C_1 - C_6 alkyl- $N(R^8)_2$, C_1 - C_6 alkyl- R^{11} , C_1 - C_6 alkyl- $CO_2R^7R^{11}$, C_1 - C_6 alkoxy- R^{11} , nitro, $O-R^{10}$, $C=O$, COR^{11} , CO_2R^{11} , SR^{10} , SOR^{11} , SO_2R^{11} , $NHSO_2R^{11}$, C_1 - C_6 alkyl- SR^{10} , halo, halo C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, hydroxy C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkoxy, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{18} ;

R^{13} and R^{14} are each independently selected from -H, oxo, C_1 - C_6 alkyl, COR^{23} , and aryl;

R^{15} and R^{16} are each independently selected from -H, aryl, arylalkyl, wherein aryl, arylalkyl, are optionally substituted with one or more of the groups defined by R^{24} ;

R^{17} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{19} , NHR^{19} , aryl, heteroarylalkyl, and heterocyclalkyl, wherein aryl is optionally substituted with one or more of the groups defined by R^{24} ;

R^{18} is selected from -H, oxo, hydroxyl, C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, amino, amino C_1 - C_6 alkyl, $N(R^{19})_2$, C_1 - C_6 alkyl- $N(R^{19})_2$, CO_2R^{23} , SR^{21} , halo, halo C_1 - C_4 alkyl, aryl, heteroaryl, and heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R^{24} ;

R^{19} and R^{20} are each independently selected from -H, C_1 - C_6 alkyl, heteroaryl, heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R^{30} ;

R^{21} and R^{22} are each independently selected from -H and C_1 - C_6 alkyl;

R^{23} is selected from -H and C_1 - C_6 alkyl;

R^{24} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, CO_2R^{29} , halo, and halo C_1 - C_4 alkyl;

R²⁹ is selected from -H, and C₁-C₆ alkyl ;

R³⁰ is selected from -H, aryl, heteroaryl, heterocyclyl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R³⁶;

5 R³⁶ is selected from -H and halo; and

R², R³, R⁴ and R³⁷ are each independently selected from an R¹ group.

14. The compound according to claim 1, wherein:

R^a is an M-ring that is an aromatic pyrimidine;

10 M¹, M³ and M⁴ are carbon and are substituted with (L)_nR¹;

M⁵ is carbon;

M² and M⁶ are nitrogen;

R¹ is selected from -H, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, hydroxyl, C₁-C₆ alkoxy, C₂-C₆ alkenyl-R¹¹, C₁-C₆ alkoxy-R¹¹, COR¹⁷, CO₂R⁷,
15 CONHR⁷, N(R⁸)₂, amino C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, amino, amino C₁-C₄ alkyl-R⁷, halo C₁-C₄ alkyl, C₁-C₆ alkyl-NHR⁷, carbonitrile, SR¹⁰, halo, NHR⁷, NR⁸R⁹, NHR⁷-C₁-C₆ alkyl, NR⁸R⁹-C₁-C₆ alkyl, nitro, cyano, O-R¹⁰, C₁-C₄ alkyl-OR¹⁰, C₁-C₆ alkyl-COR¹¹, halo C₁-C₄ alkyl, aryl, heteroaryl, heterocyclyl, alkylaryl, alkylheterocyclyl, alkylheteroaryl, arylalkyl,
20 heteroarylalkyl, heterocyclylalkyl, or C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R¹²;

R⁷ and R⁸, are each independently selected from -H, C₁-C₆ alkyl, C₁-C₄ alkyl-R¹¹, C₁-C₆ alkyl-N(R¹³)₂, CO₂R¹⁶, COR¹⁷, aryl, and arylalkyl,
25 wherein aryl and arylalkyl, are optionally substituted with one or more of the groups defined by R¹⁸;

R⁹ and R¹⁰ are each independently selected from -H, hydroxyl, C₁-C₆ alkyl, C₁-C₆ alkyl-R¹⁷, C₁-C₆ alkyl-NH₂R¹³, CO₂R¹⁶, COR¹⁷, C₁-C₆ alkyl-CO₂R¹⁶, C₁-C₆ alkyl-CONH-R¹⁶, C₁-C₆ alkyl-CON(R¹⁶)₂, hydroxy C₁-C₄ alkyl,
30 halo C₁-C₄ alkoxy, halo C₁-C₄ alkyl, Si(R¹³)₂R¹⁷, aryl, heteroaryl, heterocyclyl, arylalkyl, and C₁-C₁₀ mono- and bicyclic cycloalkyl, wherein

aryl, heteroaryl, heterocyclyl, and arylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

5 R^{11} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, hydroxyl, halo, amino, NHR^{13} , $N(R^{13})_2$, COR^{13} , CO_2R^{17} , halo C_1 - C_4 alkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, wherein heterocyclyl, heteroarylalkyl, and heterocyclylalkyl, are optionally substituted with one or more of the groups defined by R^{18} ;

10 R^{12} is selected from -H, hydroxyl, oxo, C_1 - C_6 alkyl, hydroxyl C_1 - C_6 alkyl- R^{11} , C_1 - C_{10} alkoxy, amino, amino C_1 - C_4 alkyl- R^7 , NHR^7 , $N(R^7)_2$, C_1 - C_6 alkyl- NHR^7 , C_1 - C_6 alkyl- NHR^8R^9 , C_1 - C_6 alkyl- $N(R^8)_2$, C_1 - C_6 alkyl- R^{11} , C_1 - C_6 alkyl- $CO_2R^7R^{11}$, C_1 - C_6 alkoxy- R^{11} , nitro, $O-R^{10}$, $C=O$, COR^{11} , CO_2R^{11} , SR^{10} , SOR^{11} , SO_2R^{11} , $NHSO_2R^{11}$, C_1 - C_6 alkyl- SR^{10} , halo, halo C_1 - C_4 alkyl, halo C_1 - C_4 alkoxy, hydroxy C_1 - C_4 alkyl, hydroxy C_1 - C_4 alkoxy, aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, 15 and C_1 - C_{10} mono- and bicyclic cycloalkyl, wherein aryl, heteroaryl, heterocyclyl, arylalkyl, heteroarylalkyl, and heterocyclylalkyl, and C_1 - C_{10} mono- and bicyclic cycloalkyl are optionally substituted with one or more of the groups defined by R^{18} ;

20 R^{13} and R^{14} are each independently selected from -H, oxo, C_1 - C_6 alkyl, COR^{23} , and aryl;

R^{15} and R^{16} are each independently selected from -H, aryl, arylalkyl, wherein aryl, arylalkyl, are optionally substituted with one or more of the groups defined by R^{24} ;

25 R^{17} is selected from -H, C_1 - C_6 alkyl, C_1 - C_6 alkyl- R^{19} , NHR^{19} , aryl, heteroarylalkyl, and heterocyclylalkyl, wherein aryl is optionally substituted with one or more of the groups defined by R^{24} ;

30 R^{18} is selected from -H, oxo, hydroxyl, C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, amino, amino C_1 - C_6 alkyl, $N(R^{19})_2$, C_1 - C_6 alkyl- $N(R^{19})_2$, CO_2R^{23} , SR^{21} , halo, halo C_1 - C_4 alkyl, aryl, heteroaryl, and heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R^{24} ;

R¹⁹ and R²⁰ are each independently selected from -H, C₁-C₆ alkyl, heteroaryl, heterocyclyl, wherein aryl, heteroaryl, and heterocyclyl, are optionally substituted with one or more of the groups defined by R³⁰;

5 R²¹ and R²² are each independently selected from -H and C₁-C₆ alkyl;

R²³ is selected from -H and C₁-C₆ alkyl;

R²⁴ is selected from -H, C₁-C₆ alkyl, C₁-C₆ alkoxy, CO₂R²⁹, halo, and halo C₁-C₄ alkyl;

R²⁹ is selected from -H, and C₁-C₆ alkyl;

10 R³⁰ is selected from -H, aryl, heteroaryl, heterocyclyl, alkylaryl, arylalkyl, wherein aryl, heteroaryl, heterocyclyl, alkylaryl, and arylalkyl, are optionally substituted with one or more of the groups defined by R³⁶;

R³⁶ is selected from -H and halo; and

15 R², R³, R⁴ and R³⁷ are each independently selected from an R¹ group.

15. An MK-2 inhibiting compound that is listed in Table I or Table II.

16. The compound according to claim 15, wherein the compound is selected from the group consisting of:

20 1-(2-aminoethyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylic acid trifluoroacetate,

1-(3-aminopropyl)-3-[2-(3-nitrophenyl)pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride,

25 6-(aminomethyl)-2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one,

1-(2-aminoethyl)-3-[2-[(E)-2-phenylethenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylic acid trifluoroacetate,

1-(2-aminoethyl)-3-[2-[4-(hydroxymethyl)phenyl]pyridin-4-yl]-1H-pyrazole-5-carboxylic acid dihydrochloride,

30 6-(hydroxymethyl)-2-(2-quinolin-3-ylpyridin-4-yl)-6,7-dihydropyrazolo[1,5-a]pyrazin-4(5H)-one, and

1-(3-aminopropyl)-3-(2-quinolin-3-ylpyridin-4-yl)-1H-pyrazole-5-carboxylic acid dihydrochloride, and mixtures thereof.

17. A compound having the structure described in claim 1, for use in a method of inhibiting MK-2.

5 18. A compound that is selected from the compounds described in claim 15, for use in a method of inhibiting MK-2.

19. A compound having the structure described in claim 1, for use in a method of preventing or treating a TNF α mediated disease or disorder in a subject, the method comprising administering to the subject an effective amount of the compound.

20. The compound according to claim 19, wherein the subject is one that is in need of such prevention or treatment.

21. The compound according to claim 19, wherein the subject is a mammal.

15 22. The compound according to claim 19, wherein the subject is a human.

23. The compound according to claim 19, wherein the TNF α mediated disease or disorder is one that is selected from the group consisting of connective tissue and joint disorders, neoplasia disorders, cardiovascular disorders, otic disorders, ophthalmic disorders, respiratory disorders, gastrointestinal disorders, angiogenesis-related disorders, immunological disorders, allergic disorders, nutritional disorders, infectious diseases and disorders, endocrine disorders, metabolic disorders, neurological and neurodegenerative disorders, psychiatric disorders, hepatic and biliary disorders, musculoskeletal disorders, genitourinary disorders, gynecologic and obstetric disorders, injury and trauma disorders, surgical disorders, dental and oral disorders, sexual dysfunction disorders, dermatologic disorders, hematological disorders, and poisoning disorders.

24. The compound according to claim 19, wherein the TNF α mediated disease or disorder is selected from the group consisting of: arthritis, rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus, juvenile arthritis, asthma,
5 bronchitis, menstrual cramps, tendinitis, bursitis, connective tissue injuries or disorders, skin related conditions, psoriasis, eczema, burns, dermatitis, gastrointestinal conditions, inflammatory bowel disease, gastric ulcer, gastric varices, Crohn's disease, gastritis, irritable bowel syndrome, ulcerative colitis, cancer, colorectal cancer, herpes simplex infections, HIV,
10 pulmonary edema, kidney stones, minor injuries, wound healing, vaginitis, candidiasis, lumbar spondylanhrosis, lumbar spondylarthrosis, vascular diseases, migraine headaches, sinus headaches, tension headaches, dental pain, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis,
15 multiple sclerosis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, swelling occurring after injury, myocardial ischemia, ophthalmic diseases, retinitis, retinopathies, conjunctivitis, uveitis, ocular photophobia, acute injury to the eye tissue, pulmonary inflammation, viral infections, cystic fibrosis, central nervous
20 system disorders, cortical dementias, and Alzheimer's disease.

25. A compound that is selected from the group consisting of the compounds described in claim 15, for use in a method of preventing or treating a TNF α mediated disease or disorder in a subject, the method comprising administering to the subject the compound.

25 26. A therapeutic composition comprising a compound having the structure described in claim 1.

27. A therapeutic composition comprising at least one MK-2 inhibitory compound that is described in claim 15.

28. A pharmaceutical composition comprising a
30 pharmaceutically acceptable carrier and at least one MK-2 inhibitory compound having the structure described in claim 1.

29. The pharmaceutical composition according to claim 28, wherein the MK-2 inhibitory compound has an IC_{50} for MK-2 of not over 0.1 mM.

30. A kit comprising a dosage form that includes a
5 therapeutically effective amount of at least one MK-2 inhibitory compound having a structure described in claim 1.

31. The use of a compound having the structure described in claim 1 in the manufacture of a medicament for use in a method of inhibiting MK-2.

10 32. The use of a compound that is selected from the compounds described in claim 15 in the manufacture of a medicament for use in a method of inhibiting MK-2.

33. The use of a compound having the structure described in claim 1 in the manufacture of a medicament for use in a method of
15 preventing or treating a TNF α mediated disease or disorder in a subject, the method comprising administering to the subject an effective amount of the medicament.

34. The use according to claim 33, wherein the subject is one that is in need of such prevention or treatment.

20 35. The use according to claim 33, wherein the subject is a mammal.

36. The use according to claim 33, wherein the subject is a human.

37. The use according to claim 33, wherein the TNF α mediated
25 disease or disorder is one that is selected from a group consisting of connective tissue and joint disorders, neoplasia disorders, cardiovascular disorders, otic disorders, ophthalmic disorders, respiratory disorders, gastrointestinal disorders, angiogenesis-related disorders, immunological disorders, allergic disorders, nutritional disorders, infectious diseases and
30 disorders, endocrine disorders, metabolic disorders, neurological and neurodegenerative disorders, psychiatric disorders, hepatic and biliary disorders, musculoskeletal disorders, genitourinary disorders, gynecologic and obstetric disorders, injury and trauma disorders, surgical disorders,

dental and oral disorders, sexual dysfunction disorders, dermatologic disorders, hematological disorders, and poisoning disorders.

38. The use according to claim 33, wherein the TNF α mediated disease or disorder is selected from the group consisting of:

- 5 arthritis, rheumatoid arthritis, spondyloarthopathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus, juvenile arthritis, asthma, bronchitis, menstrual cramps, tendinitis, bursitis, connective tissue injuries or disorders, skin related conditions, psoriasis, eczema, burns, dermatitis, gastrointestinal conditions, inflammatory bowel disease, gastric ulcer,
10 gastric varices, Crohn's disease, gastritis, irritable bowel syndrome, ulcerative colitis, cancer, colorectal cancer, herpes simplex infections, HIV, pulmonary edema, kidney stones, minor injuries, wound healing, vaginitis, candidiasis, lumbar spondylanhrosis, lumbar spondylarthrosis, vascular diseases, migraine headaches, sinus headaches, tension headaches,
15 dental pain, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis, multiple sclerosis, sarcoidosis, nephritic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, swelling occurring after injury, myocardial ischemia, ophthalmic diseases, retinitis, retinopathies,
20 conjunctivitis, uveitis, ocular photophobia, acute injury to the eye tissue, pulmonary inflammation, viral infections, cystic fibrosis, central nervous system disorders, cortical dementias, and Alzheimer's disease.

39. The use of a compound that is selected from the group consisting of the compounds described in claim 15 in the manufacture of a
25 medicament for use in a method of preventing or treating a TNF α mediated disease or disorder in a subject, the method comprising administering to the subject the medicament.





FIGURE 1.

**FIGURE 2.**