

(19) **DANMARK**

(10) **DK/EP 2950786 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 9/16 (2006.01)** **A 61 K 9/20 (2006.01)** **A 61 K 9/24 (2006.01)**
A 61 K 31/381 (2006.01) **A 61 K 31/4025 (2006.01)** **A 61 K 31/4184 (2006.01)**
A 61 K 31/439 (2006.01) **A 61 K 31/4709 (2006.01)** **A 61 K 31/497 (2006.01)**
A 61 K 31/4985 (2006.01) **A 61 K 31/501 (2006.01)** **A 61 K 31/5025 (2006.01)**
A 61 K 31/513 (2006.01) **A 61 K 31/5377 (2006.01)** **A 61 K 31/675 (2006.01)**
A 61 K 31/7056 (2006.01) **A 61 K 31/7072 (2006.01)** **A 61 K 31/7076 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-02-17**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-11-27**
- (86) Europæisk ansøgning nr.: **14704502.5**
- (86) Europæisk indleveringsdag: **2014-01-30**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-12-09**
- (86) International ansøgning nr.: **US2014013953**
- (87) Internationalt publikationsnr.: **WO2014120981**
- (30) Prioritet: **2013-01-31 US 201361759320 P** **2013-03-04 US 201361772292 P**
2013-05-30 US 201361828899 P **2013-08-27 US 201361870729 P**
2013-10-30 US 201361897793 P **2013-11-21 US 201361907332 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Gilead Pharmasset LLC, 333 Lakeside Drive, Foster City, CA 94404, USA**
- (72) Opfinder: **CHAL, Ben, c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404, USA**
MOGALIAN, Erik, c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404, USA
PAKDAMAN, Rowchanak, c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404, USA
OLIYAI, Reza, c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404, USA
STEFANIDIS, Dimitrios, c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404, USA
ZIA, Vahid, c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404, USA
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **KOMBINATIONSFORMULERING AF TO ANTIVIRALE FORBINDELSER**
- (56) Fremdragne publikationer:
WO-A1-2011/156757
GERMAN P ET AL: "Lack of clinically significant pharmacokinetic drug-drug interaction between sofosbuvir (GS-7977) and GS-5885 or GS-9669 in healthy volunteers", 63RD ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR THE STUDY OF LIVER DISEASES,, 1 November 2012 (2012-11-01), XP009177074,
CHIOU D ET AL: "Crystallization of Amorphous Components in Spray-Dried Powders", DRYING TECHNOLOGY, TAYLOR & FRANCIS, PHILADELPHIA, PA, US, vol. 25, 1 January 2007 (2007-01-01), pages 1427-1435, XP009108943, ISSN: 0737-3937, DOI: 10.1080/07373930701536718

Fortsættes ...

HILFIKER R ET AL: "Relevance of Solid-state Properties for Pharmaceutical products", 1 January 2006 (2006-01-01), 20060101, PAGE(S) 1 - 19, XP002525043, ISBN: 978-3-527-31146-0 the whole document
GUILLORY J K ED - BRITTAIN H G: "GENERATION OF POLYMORPHS, HYDRATES, SOLVATES, AND AMORPHOUS SOLIDS", 1 January 1999 (1999-01-01), POLYMORPHISM IN PHARMACEUTICAL SOLIDS; [DRUGS AND THE PHARMACEUTICAL SCIENCES ; 95], MARCEL DEKKER INC, NEW YORK * BASEL, PAGE(S) I - II,183, XP002350313, ISBN: 978-0-8247-0237-3 the whole document page 213

DESCRIPTION

BACKGROUND

[0001] Hepatitis C is recognized as a chronic viral disease of the liver which is characterized by liver disease. Although drugs targeting the liver are in wide use and have shown effectiveness, toxicity and other side effects have limited their usefulness. Inhibitors of hepatitis C virus (HCV) are useful to limit the establishment and progression of infection by HCV as well as in diagnostic assays for HCV.


[0002] Ledipasvir is a selective inhibitor of non-structural protein 5A (NS5A), which has been described previously (see, for example, WO 2010/132601). The chemical name of ledipasvir is (1-{3-[6-(9,9-difluoro-7-{2-[5-(2-methoxycarbonylamino-3-methyl-butyl)-5-aza-spiro[2.4]hept-6-yl]-3H-imidazol-4-yl]-9H-fluoren-2-yl)-1H-benzimidazol-2-yl]-2-aza-bicyclo[2.2.1]heptane-2-carbonyl}-2-methyl-propyl)-carbamic acid methyl ester.

[0003] Sofosbuvir (SOF) is a selective inhibitor of non-structural protein 5B (NS5B) (see, for example, WO 2010/132601 and U.S. Patent 7,964,580). The chemical name of sofosbuvir is (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-yl)methoxy)(phenoxy)phosphoryl)amino) propanoate.

SUMMARY


[0004] The present disclosure provides a pharmaceutical composition comprising a solid dispersion comprising amorphous ledipasvir dispersed within a polymer matrix formed by copovidone, and sofosbuvir in a crystalline form, and the composition for use in the treatment of HCV, as defined in claims 1 to 13.

[0005] Ledipasvir has the chemical name of (1-{3-[6-(9,9-difluoro-7-{2-[5-(2-methoxycarbonylamino-3-methyl-butyl)-5-aza-spiro[2.4]hept-6-yl]-3H-imidazol-4-yl]-9H-fluoren-2-yl)-1H-benzimidazol-2-yl]-2-aza-bicyclo[2.2.1]heptane-2-carbonyl}-2-methyl-propyl)-carbamic acid methyl ester, and has the following chemical formula:



[0006] Sofosbuvir (SOF) has the chemical name of (S)-isopropyl 2-(((S)-(((2R,3R,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-4-fluoro-3-hydroxy-4-methyltetrahydrofuran-2-

yl)methoxy)(phenoxy)phosphoryl)amino)propanoate and has the following chemical formula:



[

[0007] Herein disclosed also are pharmaceutical dosage forms and tablets, and methods for using the combination in the treatment of hepatitis C.

BRIEF DESCRIPTION OF THE DRAWINGS

[0008]

FIG. 1 is a XRPD pattern of the solid dispersion formulation of ledipasvir comprising copovidone in a drug:polymer ratio of 1:1. As shown by the XRPD, the solid dispersion is in the amorphous state.

FIG. 2 is a modulated differential scanning calorimetry (DSC) curve of the solid dispersion of ledipasvir comprising copovidone in a drug:polymer ratio of 1:1. The glass transition temperature of the solid dispersion is about 140 °C.

FIG. 3 shows a solid state characterization of the solid dispersion formulation of ledipasvir comprising copovidone in a drug:polymer ratio of 1:1 by solid state nuclear magnetic resonance (SS-NMR).

FIG. 4 is a Fourier-transformed Raman spectra of the solid dispersion of ledipasvir comprising copovidone in a drug:polymer ratio of 1:1.

FIG. 5 shows the dissolution of sofosbuvir in the sofosbuvir (400 mg)/ledipasvir (90 mg) combination described in Example 7.

FIG. 6 shows the dissolution of ledipasvir in the sofosbuvir (400 mg)/ledipasvir (90 mg) combination formulation described in Example 3.

FIG. 7A-D shows the HCV RNA levels during 12 weeks of treatment and 24 weeks post-treatment for treatment naive (FIG. 7A) or null responder (FIG. 7B) patients treated with sofosbuvir (SOF) and ribavirin (RBV) and for treatment naive (FIG. 7C) or null responder (FIG. 7D) patients treated with sofosbuvir (SOF), ledipasvir and ribavirin (RBV). This data and experimental method are further described in Example 5.

FIG. 8A-B present charts to show that all three formulations had comparable dissolution

performance, similar to that of the single-agent controls. This is more thoroughly described in Example 7.

FIG. 9 presents the pH-solubility profile of ledipasvir at room temperature (RT). The line is the nonlinear least-square regression fit using equation $S_T = S_0[(1+10^{(pK_{a1}-pH)})+10^{(pK_{a1}+pK_{a2}-2\cdot pH)}]$ with an intrinsic solubility (S_0) of 0.04 $\mu\text{g/mL}$ and a weakly basic pK_{a1} and pK_{a2} values of 5.0 and 4.0, respectively. This is more thoroughly described in Example 8.

FIG. 10 shows the study design for treatment naive (non-cirrhotic) and for null responders (50% cirrhotic) for patients treated with a fixed dose combination of sofosbuvir (SOF) and ledipasvir, with and without ribavirin (RBV) for 8 and 12 weeks. The data and experimental method are described in Example 9.

FIG. 11 shows the results for treatment naive (non-cirrhotic) and for null responders (50% cirrhotic) for patients treated with a fixed dose combination of sofosbuvir (SOF) and ledipasvir, with and without ribavirin (RBV) for 8 and 12 weeks. This data and experimental method are further described in Example 9.

DETAILED DESCRIPTION

1. Definitions

[0009] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise.

[0010] As used herein, the term "about" used in the context of quantitative measurements means the indicated amount $\pm 10\%$, or alternatively $\pm 5\%$, or $\pm 1\%$. For example, with a $\pm 10\%$ range, "about 2:8" can mean 1.8-2.2:7.2-8.8.

[0011] The term "amorphous" refers to a state in which the material lacks long range order at the molecular level and, depending upon temperature, may exhibit the physical properties of a solid or a liquid. Typically such materials do not give distinctive X-ray diffraction patterns and, while exhibiting the properties of a solid, are more formally described as a liquid. Upon heating, a change from solid to liquid properties occurs which is characterized by a change of state, typically second order (glass transition).

[0012] The term "crystalline" refers to a solid phase in which the material has a regular ordered internal structure at the molecular level and gives a distinctive X-ray diffraction pattern with defined peaks. Such materials when heated sufficiently will also exhibit the properties of a

liquid, but the change from solid to liquid is characterized by a phase change, typically first order (melting point).

[0013] The term "substantially amorphous" as used herein is intended to mean that greater than 70%; or greater than 75%; or greater than 80%; or greater than 85%; or greater than 90%; or greater than 95%, or greater than 99% of the compound present in a composition is in amorphous form. "Substantially amorphous" can also refer to material which has no more than about 20% crystallinity, or no more than about 10% crystallinity, or no more than about 5% crystallinity, or no more than about 2% crystallinity.

[0014] The term "substantially crystalline" as used herein is intended to mean that greater than 70%; or greater than 75%; or greater than 80%; or greater than 85%; or greater than 90%; or greater than 95%, or greater than 99% of the compound present in a composition is in crystalline form. "Substantially crystalline" can also refer to material which has no more than about 20%, or no more than about 10%, or no more than about 5%, or no more than about 2% in the amorphous form.

[0015] The term "polymer" refers to a chemical compound or mixture of compounds consisting of repeating structural units created through a process of polymerization. Suitable polymers useful in this invention are described throughout.

[0016] The term "polymer matrix" as used herein is defined to mean compositions comprising one or more polymers in which the active agent is dispersed or included within the matrix.

[0017] The term "solid dispersion" refers to the dispersion of one or more active agents in a polymer matrix at solid state prepared by a variety of methods, including spray drying, the melting (fusion), solvent, or the melting-solvent method.

[0018] The term "amorphous solid dispersion" as used herein, refers to stable solid dispersions comprising an amorphous active agent and a polymer. By "amorphous active agent," it is meant that the amorphous solid dispersion contains active agent in a substantially amorphous solid state form. In some aspects, as shown by the XRPD in **FIG. 1**, the solid dispersion is in the amorphous state, and the glass transition temperature of the solid dispersion is about 140 °C (see **FIG. 2**).

[0019] The term "pharmaceutically acceptable" indicates that the material does not have properties that would cause a reasonably prudent medical practitioner to avoid administration of the material to a patient, taking into consideration the disease or conditions to be treated and the respective route of administration. For example, it is commonly required that such a material be essentially sterile, e.g., for injectibles.

[0020] The term "pharmaceutically acceptable polymer" refers to a polymer that does not have properties that would cause a reasonably prudent medical practitioner to avoid administration of the material to a patient, taking into consideration the disease or conditions to be treated

and the respective route of administration.

[0021] The term "carrier" refers to a glidant, diluent, adjuvant, excipient, or vehicle etc with which the compound is administered, without limitation. Examples of carriers are described herein and also in "Remington's Pharmaceutical Sciences" by E.W. Martin.

[0022] The term "diluent" refers to chemical compounds that are used to dilute the compound of interest prior to delivery. Diluents can also serve to stabilize compounds. Non-limiting examples of diluents include starch, saccharides, disaccharides, sucrose, lactose, polysaccharides, cellulose, cellulose ethers, hydroxypropyl cellulose, sugar alcohols, xylitol, sorbitol, maltitol, microcrystalline cellulose, calcium or sodium carbonate, lactose, lactose monohydrate, dicalcium phosphate, cellulose, compressible sugars, dibasic calcium phosphate dehydrate, mannitol, microcrystalline cellulose, and tribasic calcium phosphate.

[0023] The term "binder" when used herein relates to any pharmaceutically acceptable film which can be used to bind together the active and inert components of the carrier together to maintain cohesive and discrete portions. Non-limiting examples of binders include hydroxypropylcellulose, hydroxypropylmethylcellulose, povidone, copovidone, and ethyl cellulose.

[0024] The term "disintegrant" refers to a substance which, upon addition to a solid preparation, facilitates its break-up or disintegration after administration and permits the release of an active ingredient as efficiently as possible to allow for its rapid dissolution. Non-limiting examples of disintegrants include maize starch, sodium starch glycolate, croscarmellose sodium, crospovidone, microcrystalline cellulose, modified corn starch, sodium carboxymethyl starch, povidone, pregelatinized starch, and alginic acid.

[0025] The term "lubricant" refers to an excipient which is added to a powder blend to prevent the compacted powder mass from sticking to the equipment during the tableting or encapsulation process. It aids the ejection of the tablet from the dies, and can improve powder flow. Non-limiting examples of lubricants include magnesium stearate, stearic acid, silica, fats, calcium stearate, polyethylene glycol, sodium stearyl fumarate, or talc; and solubilizers such as fatty acids including lauric acid, oleic acid, and C₈/C₁₀ fatty acid.

[0026] The term "film coating" refers to a thin, uniform, film on the surface of a substrate (e.g. tablet). Film coatings are particularly useful for protecting the active ingredient from photolytic degradation. Non-limiting examples of film coatings include polyvinylalcohol based, hydroxyethylcellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate film coatings.

[0027] The term "glidant" as used herein is intended to mean agents used in tablet and capsule formulations to improve flow-properties during tablet compression and to produce an anti-caking effect. Non-limiting examples of glidants include colloidal silicon dioxide, talc, fumed silica, starch, starch derivatives, and bentonite.

[0028] The term "effective amount" refers to an amount that is sufficient to effect treatment, as defined below, when administered to a mammal in need of such treatment. The therapeutically effective amount will vary depending upon the patient being treated, the weight and age of the patient, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art.

[0029] The term "treatment" or "treating," to the extent it relates to a disease or condition includes preventing the disease or condition from occurring, inhibiting the disease or condition, eliminating the disease or condition, and/or relieving one or more symptoms of the disease or condition.

[0030] The term "sustained virologic response" refers to the absence of detectable RNA (or wherein the RNA is below the limit of detection) of a virus (i.e. HCV) in a patient sample (i.e. blood sample) for a specific period of time after discontinuation of a treatment. For example, a SVR at 4 weeks indicates that RNA was not detected or was below the limit of detection in the patient at 4 weeks after discontinuing HCV therapy.

[0031] The term "% w/w" as used herein refers to the weight of a component based on the total weight of a composition comprising the component. For example, if component A is present in an amount of 50% w/w in a 100 mg composition, component A is present in an amount of 50 mg.

2. Pharmaceutical Compositions

[0032] The pharmaceutical compositions comprise a combination of an effective amount of ledipasvir, wherein ledipasvir is substantially amorphous, and an effective amount of sofosbuvir, wherein sofosbuvir is substantially crystalline.

[0033] Such a combination composition, as the experimental examples demonstrate, exhibit unexpected properties. Both sofosbuvir and ledipasvir have previously been demonstrated to act as effective anti-HCV agents. Ledipasvir, when administered alone in a conventional formulation, however, exhibited a negative food effect as evidenced by a roughly 2-fold decrease in exposure when given with a high-fat meal relative to dosing in the fasted state (see, e.g., **Tables 10 and 11, Example 3**). When ledipasvir is administered in a solid dispersion formulation and in the combination with sofosbuvir, no such negative food effect occurs (**Table 12, Example 3**).

[0034] In the combination composition, ledipasvir is present in a substantially amorphous form. Compared to crystalline agents, amorphous agents are expected to be unstable and have nonlinear solubility and exposure profiles. The data presented herein, however, show that ledipasvir in the combination composition is stable under various conditions, both short-term and long-term, and maintains high and consistent solubility and exposure profiles (**Example 6**).

[0035] Further, according the conventional wisdom, it is not advisable to co-formulate an amorphous agent with a crystalline agent, because the crystals can serve as seeds to induce crystallization of the amorphous agent, leading to instability of the amorphous agent. The current data show that, however, whether co-granulated or co-blended with sofosbuvir in the same layer or integrated as separate layers, ledipasvir stays stable and does not form crystals in the composition (**Example 6**).

[0036] It is also been discovered that, in tablet formations of the combination composition where sofosbuvir and ledipasvir are either co-granulated or co-blended, drug-drug interaction does not occur (**Example 7**).

A. Ledipasvir

[0037] Ledipasvir has previously been described (see, for example, WO 2010/132601) and can be prepared by methods described therein. In one embodiment, the pharmaceutical composition comprises ledipasvir formulated as a solid dispersion dispersed within a polymer matrix formed by a pharmaceutically acceptable polymer. The starting material of the solid dispersion can be a variety of forms of ledipasvir including crystalline forms, amorphous form, salts thereof, solvates thereof and the free base. For example, the acetone solvate, D-tartrate salt, anhydrous crystalline free base, amorphous free base, solvates or desolvates of ledipasvir can be used. Solvates of ledipasvir include, for example, those described in U.S. Publication No. 2013/0324740 such as, for example, the monoacetone solvate, diacetone solvate, ethyl acetone solvate, isopropyl acetate solvate, methyl acetate solvate, ethyl formate solvate, acetonitrile solvate, tetrahydrofuran solvate, methyl ethyl ketone solvate, tetrahydrofuran solvate, methyl ethyl ketone solvate, and methyl *tert*-butyl ether solvate. Particular starting materials contemplated to be useful are the monoacetone solvate, diacetone solvate, anhydrous crystalline free base, D-tartrate salt, anhydrous crystalline free base, and amorphous free base. These forms are characterized and described in U.S. Publication No. 2013/0324496.

[0038] After dispersion with the polymer, the solid dispersion is in the amorphous form. **FIGs. 1-4** characterize the amorphous solid dispersion comprising ledipasvir. As shown by the XRPD in **FIG. 1**, the solid dispersion is in the amorphous state, and the glass transition temperature of the solid dispersion is about 140 °C.

[0039] Various techniques are well known in the art for preparing solid dispersions including, but not limited to melt-extrusion, spray-drying, lyophilization, and solution-evaporation.

[0040] Melt-extrusion is the process of embedding a compound in a thermoplastic carrier. The mixture is processed at elevated temperatures and pressures, which disperses the compound in the matrix at a molecular level to form a solid solution. Extruded material can be further processed into a variety of dosage forms, including capsules, tablets and transmucosal

systems.

[0041] For the solution-evaporation method, the solid dispersion can be prepared by dissolving the compound in a suitable liquid solvent and then incorporating the solution directly into the melt of a polymer, which is then evaporated until a clear, solvent free film is left. The film is further dried to constant weight.

[0042] For the lyophilization technique, the compound and carrier can be co-dissolved in a common solvent, frozen and sublimed to obtain a lyophilized molecular dispersion.

[0043] For spray dried solid dispersions, the solid dispersion can be made by a) mixing the compound and polymer in a solvent to provide a feed solution; and b) spray drying the feed solution to provide the solid dispersion.

[0044] Spray dried solid dispersions of ledipasvir provide improved *in vivo* and *in vitro* performance and manufacturability/scalability relative to the other formulation approaches, such as wet and dry granulation formulations. Ledipasvir can be provided either as the free base, D-tartrate salt, crystalline acetone solvate, or other solvate as described herein.

[0045] The selection of the polymer for the solid dispersion is based on the stability and physical characteristics of the ledipasvir in the solution. Hypromellose and copovidone solid dispersions both showed adequate stability and physical characteristics. The polymer used in the solid dispersion is copovidone. Furthermore, the copovidone-based dispersion increased in bioavailability more than the equivalent hypromellose-based formulation (F = 30% and 22%, respectively) when prepared at 2:1 API:polymer ratio. Bioavailability of the copovidone-based formulation was further enhanced by increasing the fraction of polymer to a 1:1 ratio, resulting in a bioavailability of 35% in famotidine pretreated dogs.

[0046] The polymer (copovidone) is non-ionic. Non-ionic polymers showed benefits in screening solubility experiments.

[0047] Hypromellose and copovidone solid dispersions both showed adequate stability and physical characteristics. A copovidone-based dispersion increased bioavailability more than the equivalent hypromellose-based formulation (F = 30% and 22%, respectively) when spray dried at 2:1 ledipasvir:polymer ratio (data not shown). Accordingly, the polymer is copovidone.

[0048] The weight ratio of ledipasvir to polymer is about 1:1. A 1:1 ratio of ledipasvir:copovidone resulted in increased bioavailability (F = 35%) in famotidine pretreated dogs.

[0049] The pharmaceutical compositions comprises from about 10% to about 25% w/w, or from about 15% to about 20% w/w of the solid dispersion of ledipasvir. In further embodiments, the pharmaceutical composition comprises about 10% w/w, about 20% w/w, or about 25% w/w of the solid dispersion of ledipasvir. In a specific embodiment, the pharmaceutical composition

comprises about 18% w/w of the solid dispersion of ledipasvir.

[0050] Ledipasvir may be present in the pharmaceutical composition in a therapeutically effective amount. In some embodiments, the pharmaceutical composition comprises from about 7% to about 12% w/w of ledipasvir. In further embodiments, the pharmaceutical composition comprises about 1% w/w, about 3% w/w, about 5% w/w, about 7% w/w, about 11% w/w, or about 13% w/w, of ledipasvir. In a specific embodiment, the pharmaceutical composition comprises about 9% w/w of ledipasvir.

[0051] As noted above, after the ledipasvir is mixed with the polymer, the mixture can then be solubilized in a solvent. It is within the skill of those in the art to select an appropriate solvent based on the drug and/or polymer properties such as solubility, glass transition temperature, viscosity, and molecular weight. Acceptable solvents include but are not limited to, water, acetone, methyl acetate, ethyl acetate, chlorinated solvents, ethanol, dichloromethane, and methanol. In one embodiment, the solvent is selected from the group consisting of ethanol, dichloromethane, and methanol. In a further embodiment, the solvent is ethanol or methanol. In a specific embodiment, the solvent is ethanol.

[0052] Upon solubilization of the compound and polymer mixture with the solvent, the mixture may then be spray dried. Spray drying is a well known process wherein a liquid feedstock is dispersed into droplets into a drying chamber along with a heated process gas stream to aid in solvent removal and to produce a powder product. Suitable spray drying parameters are known in the art, and it is within the knowledge of a skilled artisan in the field to select appropriate parameters for spray drying. The target feed concentration is generally about 10 to about 50% with a target of about 20% and a viscosity of about 15 to about 300 cP. The inlet temperature of the spray dry apparatus is typically about 50-190° C, while the outlet temperature is about 30-90° C. The two fluid nozzle and hydrolic pressure nozzle can be used to spray dry ledipasvir. The two fluid nozzle gas flow can be about 1-10 kg/hr, the hydrolic pressure nozzle flow can be about 15-300 kg/hr, and the chamber gas flow may be about 25-2500 kg/hr. The spray-dried material typically has particle size (D_{90}) under 80 μm . In some instances, a milling step may be used, if desired to further reduce the particle size. Further descriptions of spray drying methods and other techniques for forming amorphous dispersions are provided in U.S. Pat. No. 6,763,607 and U.S. Pat. Pub. No. 2006-0189633, the entirety of each of which is incorporated herein by reference.

[0053] Spray drying out of ethanol resulted in high yields (88, 90, 92, 95, 97, 98, 99%) across a wide range of spray-drying outlet temperatures (30-90 °C) with no material accumulation on the spray dry chamber, and the yields obtained from spray drying out of DCM were 60%, 78%, and 44%. Furthermore, ledipasvir demonstrated good chemical stability in the ethanolic feed solution.

B. Sofosbuvir

[0054] Sofosbuvir has previously been described in U.S. Patent 7,964,580 and U.S. Publication Nos: 2010/0016251, 2010/0298257, 2011/0251152 and 2012/0107278. Sofosbuvir is provided as substantially crystalline in the pharmaceutical compositions described herein. Examples of preparing crystalline forms of sofosbuvir are disclosed in U.S. Publication Nos: 2010/0298257 and 2011/0251152. Crystalline forms, Forms 1-6, of sofosbuvir are described in U.S. Publication Nos.: 2010/0298257 and 2011/0251152, both of which are incorporated by reference. Forms 1-6 of sofosbuvir have the following characteristic X-ray powder diffraction (XRPD) pattern 2 θ -values measured according to the XRPD methods disclosed therein:

1. (1) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 7.5, 9.6, and 18.3 (Form 1);
2. (2) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 5.0, 7.3, and 18.1 (Form 1);
3. (3) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 6.9, 24.7, and 25.1 (Form 2);
4. (4) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 19.7, 20.6, and 24.6 (Form 3);
5. (5) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 5.0, 6.8, and 24.9 (Form 4);
6. (6) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 5.2, 6.6, and 19.1 (Form 5); and
7. (7) 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 6.1, 20.1, and 20.8 (Form 6).

[0055] Form 6, as described in the patent publications above, may be referred to as Form 2, such for example, by the Food and Drug Administration. Forms 1 and 6 are alternatively characterized by the following characteristic XRPD pattern 2 θ -values as measured according to the methods disclosed in U.S. Pat. Pub. Nos.: 2010/0298257 and 2011/0251152:

1. (1) 2 θ -reflections ($^{\circ}$) at about: 5.0 and 7.3 (Form 1); and
2. (2) 2 θ -reflections ($^{\circ}$) at about: 6.1 and 12.7 (Form 6).

[0056] In the pharmaceutical composition, the crystalline sofosbuvir has XRPD 2 θ -reflections ($^{\circ} \pm 0.2\theta$) at about: 6.1 and 12.7.

[0057] The pharmaceutical composition comprises from about 35% to about 45% w/w of sofosbuvir. In further embodiments, the pharmaceutical composition comprises about 35% w/w, or about 45% w/w. In a specific embodiment, the pharmaceutical composition comprises about 40% w/w of sofosbuvir.

C. Excipients

[0058] The pharmaceutical compositions provided in accordance with the present disclosure are usually administered orally. This disclosure therefore provides pharmaceutical compositions that comprise a solid dispersion comprising ledipasvir as described herein and one or more pharmaceutically acceptable excipients or carriers including but not limited to, inert solid diluents and fillers, diluents, including sterile aqueous solution and various organic

solvents, permeation enhancers, solubilizers, disintegrants, lubricants, binders, glidants, adjuvants, and combinations thereof. Such compositions are prepared in a manner well known in the pharmaceutical art (see, e.g., Remington's Pharmaceutical Sciences, Mace Publishing Co., Philadelphia, PA 17th Ed. (1985); and Modern Pharmaceutics, Marcel Dekker, Inc. 3rd Ed. (G.S. Banker & C.T. Rhodes, Eds.).

[0059] The pharmaceutical compositions may be administered in either single or multiple doses by oral administration. Administration may be via capsule, tablet, or the like. In one embodiment, the ledipasvir is in the form of a tablet. In a further embodiment, the tablet is a compressed tablet. In making the pharmaceutical compositions that include the solid described herein, the active ingredient is usually diluted by an excipient and/or enclosed within such a carrier that can be in the form of a capsule, tablet, sachet, paper or other container. When the excipient serves as a diluent, it can be in the form of a solid, semi-solid or liquid material (as above), which acts as a vehicle, carrier or medium for the active ingredient.

[0060] The pharmaceutical composition may be formulated for immediate release or sustained release. A "sustained release formulation" is a formulation which is designed to slowly release a therapeutic agent in the body over an extended period of time, whereas an "immediate release formulation" is an formulation which is designed to quickly release a therapeutic agent in the body over a shortened period of time. In some cases the immediate release formulation may be coated such that the therapeutic agent is only released once it reached the desired target in the body (e.g. the stomach). In a specific embodiment, the pharmaceutical composition is formulated for immediate release.

[0061] The pharmaceutical composition may further comprise pharmaceutical excipients such as diluents, binders, fillers, glidants, disintegrants, lubricants, solubilizers, and combinations thereof. Some examples of suitable excipients are described herein. When the pharmaceutical composition is formulated into a tablet, the tablet may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

[0062] In one embodiment, the pharmaceutical composition comprises a diluent selected from the group consisting of dicalcium phosphate, cellulose, compressible sugars, dibasic calcium phosphate dehydrate, lactose, lactose monohydrate, mannitol, microcrystalline cellulose, starch, tribasic calcium phosphate, and combinations thereof.

[0063] In further embodiments, the pharmaceutical composition comprises lactose monohydrate in an amount from about 5 to about 25% w/w, or from about 10 to about 20% w/w. In specific embodiments, the lactose monohydrate is present at about 5% w/w, at about 10% w/w, at about 15% w/w, at about 20% w/w, or at about 25% w/w. In a further specific embodiment, the lactose monohydrate is in an amount of about 16.5% w/w.

[0064] In yet further embodiments, the pharmaceutical composition comprises microcrystalline cellulose in an amount from about 5 to about 25% w/w, or from about 10 to about 25% w/w, or from about 15 to about 20% w/w. In specific embodiments, the microcrystalline cellulose is present in an amount of about 5%, or about 10%, or about 15%, or about 20%, or about 25%. In a further specific embodiment, the microcrystalline cellulose is in an amount of about 18% w/w.

[0065] In other embodiments, the pharmaceutical composition comprises a disintegrant, croscarmellose sodium.

[0066] In certain embodiments, the pharmaceutical composition comprises croscarmellose sodium in an amount from about 1 to about 10% w/w, or from about 1 to about 8% w/w, or from about 2 to about 8% w/w. In specific embodiments, the croscarmellose sodium is present in an amount of about 1%, or about 3%, or about 6%, or about 8%, or about 10%. In a further specific embodiment, the croscarmellose sodium is in an amount of about 5% w/w.

[0067] In other embodiments, the pharmaceutical composition comprises a glidant, colloidal silicon dioxide.

[0068] In further embodiments, the pharmaceutical composition comprises colloidal silicon dioxide in an amount from about 0.5 to about 3% w/w, or from about 0.5 to about 2% w/w, or from about 0.5 to about 1.5% w/w. In specific embodiments, the colloidal silicon dioxide is present in an amount of 0.5% w/w, 0.75% w/w, 1.25% w/w, 1.5% w/w, or 2% w/w. In a further specific embodiment, the colloidal silicon dioxide is present in an amount of about 1% w/w.

[0069] In other embodiments, the pharmaceutical composition comprises a lubricant, magnesium stearate.

[0070] In further embodiments, the pharmaceutical composition comprises magnesium stearate in an amount from about 0.1 to about 3% w/w, or from about 0.1 to about 2.5% w/w, or from about 0.5 to about 3% w/w, or from about 0.5 to about 2.5% w/w, or from about 0.5 to about 2% w/w, or from about 1 to about 3% w/w, or from about 1 to about 2% w/w. In specific embodiments, the magnesium stearate is present in an amount of about 0.1%, or about 0.5, or about 1%, or about 2%, or about 2.5%, or about 3% w/w. In a further specific embodiment, the magnesium stearate is in an amount of about 1.5% w/w.

[0071] In one embodiment, the pharmaceutical composition comprises a) about 40% w/w of sofosbuvir and b) about 18% w/w of the solid dispersion comprising ledipasvir. In yet a further related embodiment, the composition further comprises a) about 5 to about 25% w/w lactose monohydrate, b) about 5 to about 25% w/w microcrystalline cellulose, c) about 1 to about 10% w/w croscarmellose sodium, d) about 0.5 to about 3% w/w colloidal silicon dioxide, and e) about 0.1 to about 3% w/w magnesium stearate. In a further embodiment, the pharmaceutical composition comprises a) about 40% w/w of sofosbuvir, b) about 18 %w/w of the solid dispersion comprising ledipasvir, c) about 16.5% w/w lactose monohydrate, d) about 18% w/w

microcrystalline cellulose, e) about 5% w/w croscarmellose sodium, f) about 1% w/w colloidal silicon dioxide, and g) about 1.5% w/w magnesium stearate.

3. Pharmaceutical Dosage Forms

[0072] The disclosure provides for tablets, pills, and the like, comprising the pharmaceutical compositions or dosage forms described herein. The tablets or pills of the present disclosure may be coated to provide a dosage form affording the advantage of prolonged action or to protect from the acid conditions of the stomach. The tablets may also be formulated for immediate release as previously described. In certain embodiments, the tablet comprises a film coating. A film coating is useful for limiting photolytic degradation. Suitable film coatings are selected by routine screening of commercially available preparations. In one embodiment, the film coating is a polyvinylalcohol-based coating.

[0073] The tablets may be formulated into a monolayer or bilayer tablet. Typically, monolayer tablets comprise the active ingredients (i.e., ledipasvir and sofosbuvir) co-mixed in a single uniform layer. For making monolayer tablets, exemplary methods include, but are not limited to coblend (or bi-granulation) and codry granulation. Coblend granulation is a multi-step process consisting of separate dry granulations for each active ingredient with excipients followed by the blending of the two granulations together. Codry granulation consisted of dry granulating both active ingredients and excipients together.

[0074] Bilayer tablets comprise the active ingredients (i.e., ledipasvir and sofosbuvir) in separate layers and can be made by making a blend comprising excipients and one active ingredient (i.e., ledipasvir), and making a separate blend comprising the second active ingredient (i.e., sofosbuvir) and excipients. One blend may then be precompressed, and the second blend may then be added on top of the first precompressed blends. The resulting tablet comprises two separate layers, each layer comprising a different active ingredient.

[0075] In one embodiment, the tablet comprises a) about 30 to about 50% w/w of sofosbuvir and b) about 10 to about 40 %w/w of the solid dispersion comprising ledipasvir. In a related embodiment, the tablet comprises a) about 40% w/w of sofosbuvir and b) about 18% w/w of the solid dispersion comprising ledipasvir. In a further embodiment, the tablet comprises a) about 300 to about 500 mg of sofosbuvir and b) about 50 to about 130 mg of ledipasvir. In a yet further embodiment, the tablet comprises a) about 400 mg of sofosbuvir and b) about 90 mg of ledipasvir. In related embodiment, the tablet further comprises a) about 5 to about 25% w/w lactose monohydrate, b) about 5 to about 25% w/w microcrystalline cellulose, c) about 1 to about 10% w/w croscarmellose sodium, d) about 0.5 to about 3% w/w colloidal silicon dioxide, and e) about 0.1 to about 3% w/w magnesium stearate.

[0076] In some embodiments, the pharmaceutical compositions as described herein are formulated in a unit dosage or pharmaceutical dosage form. The term "unit dosage forms" or "pharmaceutical dosage forms" refers to physically discrete units suitable as unitary dosages

for human patients and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient (e.g., a tablet or capsule). The compounds are generally administered in a pharmaceutically effective amount. In some embodiments, each dosage unit contains from 3 mg to 2 g of ledipasvir. In other embodiments, the pharmaceutical dosage form comprises from about 3 to about 360 mg, or about 10 to about 200 mg, or about 10 to about 50 mg, or about 20 to about 40 mg, or about 25 to about 35 mg, or about 40 to about 140 mg, or about 50 to about 130 mg, or about 60 to about 120 mg, or about 70 to about 110 mg, or about 80 to about 100 mg. In specific embodiments, the pharmaceutical dosage form comprises about 40, or about 45, or about 50, or about 55, or about 60, or about 70, or about 80, or about 100, or about 120, or about 140, or about 160, or about 180, or about 200, or about 220 mg of ledipasvir. In a further specific embodiment, the pharmaceutical dosage form comprises about 90 mg of ledipasvir. In yet a further specific embodiment, the pharmaceutical dosage form comprises about 30 mg of ledipasvir.

[0077] In other embodiments, the pharmaceutical dosage form comprises from about 1 mg to about 3g of sofosbuvir. In other embodiments, the pharmaceutical dosage form comprises from about 1 to about 800 mg, or about 100 to about 700 mg, or about 200 to about 600 mg, or about 300 to about 500 mg, or about 350 to about 450 mg, of sofosbuvir. In specific embodiments, the pharmaceutical dosage form comprises about 50, or about 100, or about 150, or about 200, or about 250, or about 300, or about 350, or about 450, or about 500, or about 550, or about 600, or about 650, or about 700, or about 750, or about 800 mg of sofosbuvir. In a further specific embodiment, the pharmaceutical dosage form comprises about 400 mg of sofosbuvir. It will be understood, however, that the amount of ledipasvir and/or sofosbuvir actually administered usually will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight and response of the individual patient, the severity of the patient's symptoms, and the like.

[0078] In a specific embodiment, the pharmaceutical dosage form comprises about 400 mg of sofosbuvir and about 90 mg of ledipasvir.

[0079] In one embodiment, the pharmaceutical composition, or alternatively, the pharmaceutical dosage form or tablet comprises about 90 mg of amorphous ledipasvir formulated in a solid dispersion comprising a polymer: ledipasvir ratio of 1:1, about 400 mg crystalline sofosbuvir, lactose monohydrate in an amount from about 5 to about 25% w/w, microcrystalline cellulose in an amount from about 5 to about 25% w/w, croscarmellose sodium in an amount from about 1 to about 10% w/w, colloidal silicon dioxide in an amount from about 0.5 to about 3% w/w, and magnesium stearate in an amount from about 0.1 to about 3% w/w. In one embodiment, the polymer is copovidone.

[0080] In further embodiments, the pharmaceutical composition, pharmaceutical dosage form, or tablet as described herein is free of negative drug-drug interactions. In a related embodiment, the pharmaceutical composition, pharmaceutical dosage form, or tablet is free of

negative drug-drug interactions with acid suppressive therapies. In a further embodiment, the pharmaceutical composition, pharmaceutical dosage form, or tablet as described herein is administrable without regard to food and with or without regard to the patient being on an acid-suppressive therapy.

4. Methods of Use

[0081] The solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein may be administered to a patient suffering from hepatitis C virus (HCV) in a daily dose by oral administration. In one embodiment, the patient is human.

[0082] Previously, ledipasvir had been demonstrated to have a negative food effect when administered alone. Unexpectedly, the combination treatment of ledipasvir and sofosbuvir does not exhibit a negative food effect. Accordingly, the administration of the pharmaceutical composition comprising sofosbuvir and ledipasvir can be taken without regard to food.

[0083] In some embodiments, the combination composition achieved a reduced food effect. In some aspects, the composition achieves a first exposure, when administered to a patient following a meal, that is no more than 25%, or alternatively not more than 20%, 15% or 10%, lower than a second exposure when administered to the patient not following a meal. The exposures can be measured as C_{max} , AUC_{last} or AUC_{inf} . In some aspects, the administration is carried out within four, three, two or one hours following the meal.

[0084] In one embodiment, the solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating one or more of genotype 1 HCV infected patients, genotype 2 HCV infected patients, genotype 3 HCV infected patients, genotype 4 HCV infected patients, genotype 5 HCV infected patients, and/or genotype 6 HCV infected patients. In one embodiment, the solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating genotype 1 HCV infected patients, including genotype 1a and/or genotype 1b. In another embodiment, the solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating genotype 2 HCV infected patients, including genotype 2a, genotype 2b, genotype 2c and/or genotype 2d. In another embodiment, the solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating genotype 3 HCV infected patients, including genotype 3a, genotype 3b, genotype 3c, genotype 3d, genotype 3e and/or genotype 3f. In another embodiment, the solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating genotype 4 HCV infected patients, including genotype 4a, genotype 4b, genotype 4c, genotype 4d, genotype 4e, genotype 4f, genotype 4g, genotype 4h, genotype 4i and/or genotype 4j. In another embodiment, the solid

dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating genotype 5 HCV infected patients, including genotype 5a. In another embodiment, the solid dispersions, pharmaceutical compositions, pharmaceutical dosage forms, and tablets of ledipasvir and sofosbuvir as described herein are effective in treating genotype 6 HCV infected patients, including genotype 6a. In one embodiment, the compositions are pangenotypic, meaning they are useful across all genotypes and drug resistant mutants thereof.

[0085] In some embodiments, the pharmaceutical composition, pharmaceutical dosage form, or tablet of ledipasvir and sofosbuvir as described herein may be administered, either alone or in combination with one or more therapeutic agent(s) for treating HCV (such as a HCV NS3 protease inhibitor or an inhibitor of HCV NS5B polymerase), for about 24 weeks, for about 16 weeks, or for about 12 weeks, or less. In further embodiments, the pharmaceutical composition, pharmaceutical dosage form, or tablet of ledipasvir and sofosbuvir is administered, either alone or in combination with one or more therapeutic agent(s) for treating HCV (such as a HCV NS3 protease inhibitor or an inhibitor of HCV NS5B polymerase), for about 24 weeks or less, about 22 weeks or less, about 20 weeks or less, about 18 weeks or less, about 16 weeks or less, about 12 weeks or less, about 10 weeks or less, about 8 weeks or less, or about 6 weeks or less or about 4 weeks or less. The pharmaceutical composition, pharmaceutical dosage form, or tablet may be administered once daily, twice daily, once every other day, two times a week, three times a week, four times a week, or five times a week.

[0086] In further embodiments, a sustained virologic response is achieved at about 4 weeks, 6 weeks, 8 weeks, 12 weeks, or 16 weeks, or at about 20 weeks, or at about 24 weeks, or at about 4 months, or at about 5 months, or at about 6 months, or at about 1 year, or at about 2 years.

[0087] In one embodiment, the daily dose is 90 mg of ledipasvir and 400 mg of sofosbuvir administered in the form of a tablet. In a further embodiment, the daily dose is a tablet comprising a) about 30 to about 50% w/w of sofosbuvir, b) about 10 to about 40 %w/w of the solid dispersion comprising ledipasvir, c) about 5 to about 25% w/w lactose monohydrate, d) about 5 to about 25% w/w microcrystalline cellulose, e) about 1 to about 10% w/w croscarmellose sodium, f) about 0.5 to about 3% w/w colloidal silicon dioxide, and g) about 0.1 to about 3% w/w magnesium stearate.

[0088] In further embodiments, the patient is also suffering from cirrhosis. In yet a further embodiment, the patient is not suffering from cirrhosis.

5. Combination Therapy

[0089] In the methods described herein, the method can further comprise the administration of another therapeutic agent for treating HCV and other conditions such as HIV infections. In one embodiment, non-limiting examples of suitable additional therapeutic agents include one or

more interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers, and other drugs or therapeutic agents for treating HCV.

[0090] More specifically, the additional therapeutic agent may be selected from the group consisting of:


1. 1) interferons, e.g., pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon alpha (MOR-22, OPC-18, Alfaferone, Alfaferone, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (Albupharm), IFN alpha-2b XL, BLX-883 (Locteron), DA-3021, glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon lambda-1 (PEGylated IL-29), and belerofon;
2. 2) ribavirin and its analogs, e.g., ribavirin (Rebetol, Copegus), and taribavirin (Viramidine);
3. 3) HCV NS3 protease inhibitors, e.g., boceprevir (SCH-503034, SCH-7), telaprevir (VX-950), TMC435350, BI-1335, BI-1230, MK-7009, VBY-376, VX-500, GS-9256, GS-9451, BMS-605339, PHX-1766, AS-101, YH-5258, YH5530, YH5531, ABT-450, ACH-1625, ITMN-191, MK5172, MK6325, and MK2748;
4. 4) alpha-glucosidase 1 inhibitors, e.g., celgosivir (MX-3253), Miglitrol, and UT-231B;
5. 5) hepatoprotectants, e.g., emericasan (IDN-6556), ME-3738, GS-9450 (LB-84451), silibinin, and MitoQ;
6. 6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, e.g., R1626, R7128 (R4048), IDX184, IDX-102, BCX-4678, valopicitabine (NM-283), MK-0608, and INX-189 (now BMS986094);
7. 7) non-nucleoside inhibitors of HCV NS5B polymerase, e.g., PF-868554, VCH-759, VCH-916, JTK-652, MK-3281, GS-9190, VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773, A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, ABT-072, ABT-333, GS-9669, PSI-7792, and GS-9190;
8. 8) HCV NS5A inhibitors, e.g., AZD-2836 (A-831), BMS-790052, ACH-3102, ACH-2928, MK8325, MK4882, MK8742, PSI-461, IDX719, ABT-267, and A-689;
9. 9) TLR-7 agonists, e.g., imiquimod, 852A, GS-9524, ANA-773, ANA-975, AZD-8848 (DSP-3025), and SM-360320;
10. 10) cyclophilin inhibitors, e.g., DEBIO-025, SCY-635, and NIM81 1;
11. 11) HCV IRES inhibitors, e.g., MCI-067;
12. 12) pharmacokinetic enhancers, e.g., BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin; and
13. 13) other drugs for treating HCV, e.g., thymosin alpha 1 (Zadaxin), nitazoxanide (Alinia, NTZ), BIVN-401 (virostat), PYN-17 (altirex), KPE02003002, actilon (CPG-10101), GS-9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225, PTX-111, ITX2865, TT-033i, ANA


971, NOV-205, tarvacin, EHC-18, VGX-410C, EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-4538), Oglufanide, and VX-497 (merimepodib).

[0091] More specifically, the additional therapeutic agent may be combined with one or more compounds selected from the group consisting of non-nucleoside inhibitors of HCV NS5B polymerase (ABT-072 and ABT-333), HCV NS5A inhibitors (ACH-3102 and ACH-2928) and HCV NS3 protease inhibitors (ABT-450 and ACH-125).

[0092] In another embodiment, the therapeutic agent used in combination with the pharmaceutical compositions as described herein can be any agent having a therapeutic effect when used in combination with the pharmaceutical compositions as described herein. For example, the therapeutic agent used in combination with the pharmaceutical compositions as described herein can be interferons, ribavirin analogs, NS3 protease inhibitors, NS5B polymerase inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating HCV.

[0093] In another embodiment, the additional therapeutic agent used in combination with the pharmaceutical compositions as described herein is a cyclophilin inhibitor, including for example, a cyclophilin inhibitor disclosed in WO2013/185093. Non-limiting examples include one or more compounds selected from the group consisting of:





and stereoisomers and mixtures of stereoisomers thereof.

[0094] In another embodiment, the additional therapeutic agent used in combination with the pharmaceutical compositions as described herein is a non-nucleoside inhibitor of HCV NS5B polymerase. A non-limiting example includes Compound E (as described below).


[0095] Examples of additional anti-HCV agents which can be combined with the compositions provided herein include, without limitation, the following:

1. A. interferons, for example, pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon alpha (MOR-22, OPC-18, Alfaferone, Alfanative, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510), albinterferon alpha-2b (Albupharm), IFN alpha XL, BLX-883 (Locteron), DA-3021, glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon lambda (PEGylated IL-29), or belerofon, IFN alpha-2b XL, rIFN-alpha 2a, consensus IFN alpha, infergen, rebif, pegylated IFN-beta, oral interferon alpha, feron, reaferon, intermax alpha, r-IFN-beta, and infergen + actimmune/bavirin and ribavirin analogs, e.g., rebetol, copegus, VX-497, and viramidine (taribavirin);
2. B. NS5A inhibitors, for example, Compound B (described below), Compound C (described below), ABT-267, Compound D (described below), JNJ-47910382, daclatasvir (BMS-790052), ABT-267, MK-8742, EDP-239, IDX-719, PPI-668, GSK-2336805, ACH-3102, A-831, A-689, AZD-2836 (A-831), AZD-7295 (A-689), and BMS-790052;
3. C. NS5B polymerase inhibitors, for example, Compound E (described below), Compound F (described below), ABT-333, Compound G (described below), ABT-072, Compound H (described below), tegobuvir (GS-9190), GS-9669, TMC647055, setrobuvir (ANA-598), filibuvir (PF-868554), VX-222, IDX-375, IDX-184, IDX-102, BI-207127, valopicitabine (NM-283), PSI-6130 (R1656), PSI-7851, BCX-4678, nesbuvir (HCV-796), BILB 1941, MK-0608, NM-107, R7128, VCH-759, GSK625433, XTL-2125, VCH-916, JTK-652, MK-3281, VBY-708, A848837, GL59728, A-63890, A-48773, A-48547, BC-2329, BMS-791325, and BILB-1941;
4. D. NS3 protease inhibitors, for example, Compound I, Compound J, Compound K, ABT-450, Compound L (described below), simeprevir (TMC-435), boceprevir (SCH-503034), narlaprevir (SCH-900518), vaniprevir (MK-7009), MK-5172, danoprevir (ITMN-191), sovalprevir (ACH-1625), neceprevir (ACH-2684), Telaprevir (VX-950), VX-813, VX-500, faldaprevir (BI-201335), asunaprevir (BMS-650032), BMS-605339, VBY-376, PHX-1766, YH5531, BILN-2065, and BILN-2061;
5. E. alpha-glucosidase 1 inhibitors, for example, celgosivir (MX-3253), Miglitol, and UT-


231B;

6. F. hepatoprotectants, e.g., IDN-6556, ME 3738, MitoQ, and LB-84451;
7. G. non-nucleoside inhibitors of HCV, e.g., benzimidazole derivatives, benzo-1,2,4-thiadiazine derivatives, and phenylalanine derivatives; and
8. H. other anti-HCV agents, e.g., zidaxin, nitazoxanide (alinea), BIVN-401 (virostat), DEBIO-025, VGX-410C, EMZ-702, AVI 4065, bavituximab, oglufanide, PYN-17, KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975, XTL-6865, ANA 971, NOV-205, tarvacin, EHC-18, and NIM811.


[0096] Compound B is an NS5A inhibitor and is represented by the following chemical structure:



[0097] Compound C is an NS5A inhibitor and is represented by the following chemical structure:





[0098] Compound D is an NS5A inhibitor and is represented by the following chemical structure:




[0099] See U.S. Publication No. 2013/0102525 and references cited therein.

[0100] Compound E is an NS5B Thumb II polymerase inhibitor and is represented by the following chemical structure:






[0101] Compound F is a nucleotide inhibitor prodrug designed to inhibit replication of viral RNA by the HCV NS5B polymerase, and is represented by the following chemical structure:




[0102] Compound G is an HCV polymerase inhibitor and is represented by the following structure:




[0103] See U.S. Publication No. 2013/0102525 and references therein.


[0104] Compound H is an HCV polymerase inhibitor and is represented by the following structure:



[0105] See U.S. Publication No. 2013/0102525 and references therein.


[0106] Compound I is an HCV protease inhibitor and is represented by the following chemical structure:






[0107] See U.S. Publication No. 2014/0017198 and references therein.

[0108] Compound J is an HCV protease inhibitor and is represented by the following chemical structure:




[0109] See U.S. Patent No. 8,178,491 and references therein.

[0110] Compound K is an HCV protease inhibitor and is represented by the following chemical structure:




[0111] Compound L is an HCV protease inhibitor and is represented by the following chemical structure:



[0112] See U.S. Publication No. 2013/0102525 and references therein.

[0113] In one embodiment, the additional therapeutic agent used in combination with the pharmaceutical compositions as described herein is a HCV NS3 protease inhibitor. Non-limiting examples include one or more compounds selected from the group consisting of:



[0114] In another embodiment, the present application is provided a method of treating hepatitis C in a human patient in need thereof comprising administering to the patient a therapeutically effective amount of a pharmaceutical composition as described herein and an additional therapeutic selected from the group consisting of pegylated rIFN-alpha 2b, pegylated rIFN-alpha 2a, rIFN-alpha 2b, IFN alpha-2b XL, rIFN-alpha 2a, consensus IFN alpha, interferon, rebif, locteron, AVI-005, PEG-interferon, pegylated IFN-beta, oral interferon alpha, feron, reaferon, intermax alpha, r-IFN-beta, interferon + actimmune, IFN-omega with DUROS, albuferon, rebetol, copegus, levovirin, VX-497, viramidine (taribavirin), A-831, A-689, NM-283, valopicitabine, R1626, PSI-6130 (R1656), HCV-796, BILB 1941, MK-0608, NM-107, R7128, VCH-759, PF-868554, GSK625433, XTL-2125, SCH-503034 (SCH-7), VX-950 (Telaprevir), ITMN-191, and BILN-2065, MX-3253 (celgosivir), UT-231B, IDN-6556, ME 3738, MitoQ, and LB-84451, benzimidazole derivatives, benzo-1,2,4-thiadiazine derivatives, and phenylalanine derivatives, zadaxin, nitazoxanide (alinea), BIVN-401 (virostat), DEBIO-025, VGX-410C, EMZ-702, AVI 4065, bavituximab, oglufanide, PYN-17, KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975 (isatoribine), XTL-6865, ANA 971, NOV-205, tarvacin, EHC-18, and NIM811 and a pharmaceutically acceptable carrier or excipient.

[0115] In yet another embodiment, the present application provides a combination pharmaceutical agent comprising:

1. a) a first pharmaceutical composition comprising an effective amount of wherein ledipasvir is substantially amorphous; and an effective amount of sofosbuvir wherein

sofosbuvir is substantially crystalline as described herein and

2. b) a second pharmaceutical composition comprising at least one additional therapeutic agent selected from the group consisting of HIV protease inhibiting compounds, HIV non-nucleoside inhibitors of reverse transcriptase, HIV nucleoside inhibitors of reverse transcriptase, HIV nucleotide inhibitors of reverse transcriptase, HIV integrase inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120 inhibitors, CCR5 inhibitors, interferons, ribavirin analogs, NS3 protease inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating HCV, and combinations thereof.

[0116] The additional therapeutic agent may be one that treats other conditions such as HIV infections. Accordingly, the additional therapeutic agent may be a compound useful in treating HIV, for example HIV protease inhibiting compounds, non-nucleoside inhibitors of HIV reverse transcriptase, HIV nucleoside inhibitors of reverse transcriptase, HIV nucleotide inhibitors of reverse transcriptase, HIV integrase inhibitors, gp41 inhibitors, CXCR4 inhibitors, gp120 inhibitors, CCR5 inhibitors, interferons, ribavirin analogs, NS3 protease inhibitors, NS5b polymerase inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, non-nucleoside inhibitors of HCV, and other drugs for treating HCV.

[0117] More specifically, the additional therapeutic agent may be selected from the group consisting of

1. 1) HIV protease inhibitors, e.g., amprenavir, atazanavir, fosamprenavir, indinavir, lopinavir, ritonavir, lopinavir + ritonavir, nelfinavir, saquinavir, tipranavir, brexanavir, darunavir, TMC-126, TMC-114, mozenavir (DMP-450), JE-2147 (AG1776), AG1859, DG35, L-756423, RO0334649, KNI-272, DPC-681, DPC-684, and GW640385X, DG17, PPL-100,
2. 2) a HIV non-nucleoside inhibitor of reverse transcriptase, e.g., capravirine, emivirine, delaviridine, efavirenz, nevirapine, (+) calanolide A, etravirine, GW5634, DPC-083, DPC-961, DPC-963, MIV-150, and TMC-120, TMC-278 (rilpivirine), efavirenz, BILR 355 BS, VRX 840773, UK-453,061, RDEA806,
3. 3) a HIV nucleoside inhibitor of reverse transcriptase, e.g., zidovudine, emtricitabine, didanosine, stavudine, zalcitabine, lamivudine, abacavir, amdoxovir, elvucitabine, alovudine, MIV-210, racivir ($\pm$ -FTC), D-d4FC, emtricitabine, phosphazide, fozivudine tidoxil, fosaltivudine tidoxil, apricitabine (AVX754), amdoxovir, KP-1461, abacavir + lamivudine, abacavir + lamivudine + zidovudine, zidovudine + lamivudine,
4. 4) a HIV nucleotide inhibitor of reverse transcriptase, e.g., tenofovir, tenofovir disoproxil fumarate + emtricitabine, tenofovir disoproxil fumarate + emtricitabine + efavirenz, and adefovir,
5. 5) a HIV integrase inhibitor, e.g., curcumin, derivatives of curcumin, chicoric acid, derivatives of chicoric acid, 3,5-dicaffeoylquinic acid, derivatives of 3,5-dicaffeoylquinic acid, aurintricarboxylic acid, derivatives of aurintricarboxylic acid, caffeic acid phenethyl ester, derivatives of caffeic acid phenethyl ester, tyrphostin, derivatives of tyrphostin,

- quercetin, derivatives of quercetin, S-1360, zintevir (AR-177), L-870812, and L-870810, MK-0518 (raltegravir), BMS-707035, MK-2048, BA-011, BMS-538158, GSK364735C,
6. 6) a gp41 inhibitor, e.g., enfuvirtide, sifuvirtide, FB006M, TRI-1144, SPC3, DES6, Locus gp41, CovX, and REP 9,
 7. 7) a CXCR4 inhibitor, e.g., AMD-070,
 8. 8) an entry inhibitor, e.g., SP01A, TNX-355,
 9. 9) a gp120 inhibitor, e.g., BMS-488043 and BlockAide/CR,
 10. 10) a G6PD and NADH-oxidase inhibitor, e.g., immunitin, 10) a CCR5 inhibitor, e.g., aplaviroc, vicriviroc, INCB9471, PRO-140, INCB15050, PF-232798, CCR5mAb004, and maraviroc,
 11. 11) an interferon, e.g., pegylated rIFN-alpha 2b, pegylated rIFN-alpha 2a, rIFN-alpha 2b, IFN alpha-2b XL, rIFN-alpha 2a, consensus IFN alpha, infergen, rebif, locteron, AVI-005, PEG-infergen, pegylated IFN-beta, oral interferon alpha, feron, reaferon, intermax alpha, r-IFN-beta, infergen + actimmune, IFN-omega with DUROS, and albuferon,
 12. 12) ribavirin analogs, e.g., rebetol, copegus, levovirin, VX-497, and viramidine (taribavirin)
 13. 13) NS5a inhibitors, e.g., A-831, A-689, and BMS-790052,
 14. 14) NS5b polymerase inhibitors, e.g., NM-283, valopicitabine, R1626, PSI-6130 (R1656), HCV-796, BILB 1941, MK-0608, NM-107, R7128, VCH-759, PF-868554, GSK625433, and XTL-2125,
 15. 15) NS3 protease inhibitors, e.g., SCH-503034 (SCH-7), VX-950 (Telaprevir), ITMN-191, and BILN-2065,
 16. 16) alpha-glucosidase 1 inhibitors, e.g., MX-3253 (celgosivir) and UT-231B,
 17. 17) hepatoprotectants, e.g., IDN-6556, ME 3738, MitoQ, and LB-84451,
 18. 18) non-nucleoside inhibitors of HCV, e.g., benzimidazole derivatives, benzo-1,2,4-thiadiazine derivatives, and phenylalanine derivatives,
 19. 19) other drugs for treating Hepatitis C, e.g., zadaxin, nitazoxanide (alinea), BIVN-401 (virostat), DEBIO-025, VGX-410C, EMZ-702, AVI 4065, bavituximab, oglufanide, PYN-17, KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975 (isatoribine), XTL-6865, ANA 971, NOV-205, tarvacin, EHC-18, and NIM811,
 20. 20) pharmacokinetic enhancers, e.g., BAS-100 and SPI452, 20) RNase H inhibitors, e.g., ODN-93 and ODN-112, and
 21. 21) other anti-HIV agents, e.g., VGV-1, PA-457 (bevirimat), ampligen, HRG214, cytolin, polymun, VGX-410, KD247, AMZ 0026, CYT 99007, A-221 HIV, BAY 50-4798, MDX010 (iplimumab), PBS119, ALG889, and PA-1050040.

[0118] In one embodiment, the additional therapeutic agent is ribavirin. Accordingly, methods described herein include a method of treating hepatitis C in a human patient in need thereof comprising administering to the patient a therapeutically effective amount of ribavirin and a therapeutically effective amount of a pharmaceutical composition, pharmaceutical dosage form, or tablet as described herein. In a further embodiment, the ribavirin and pharmaceutical composition, pharmaceutical dosage form, or tablet comprising sofosbuvir and ledipasvir is

administered for about 12 weeks or less. In further embodiments, the ribavirin and pharmaceutical composition, pharmaceutical dosage form, or tablet comprising sofosbuvir and ledipasvir is administered for about 8 weeks or less, for about 6 weeks or less, or for about 4 weeks or less.

[0119] It is contemplated that the additional therapeutic agent will be administered in a manner that is known in the art and the dosage may be selected by someone of skill in the art. For example, the additional agent may be administered in a dose from about 0.01 milligrams to about 2 grams per day.

EXAMPLES

[0120] In the following examples and throughout this disclosure, abbreviations as used herein have respective meanings as follows:

ACN	Acetonitrile
AE	Adverse Event
API	Active Pharmaceutical Ingredient
AUC	Area Under the Curve
AUC _{inf}	Area under the concentration versus time curve extrapolated to infinite time, calculated as AUC _{0-last} + (C _{last} /λ _z)
AUC _{last}	Area under the concentration versus time curve from time zero to the last quantifiable concentration
BMI	Body Mass Index
BT	Breakthrough Rate
CI	Confidence Interval
CL/F	Apparent oral clearance after administration of the drug: CL/F = Dose/AUC
C _{last}	Last observed quantifiable concentration of the drug
cm	Centimeter
C _{max}	Maximum Concentration
cP	Centipoise
cP	Centipoise
CV	Coefficient of Variation
D ₉₀	Particle Size
DCF	Drug Content Factor
DCF	Drug Content Factor
DCM	Dichloromethane
dL	Deciliter

DRM	Drug Related Material
DSC	Differential Scanning Calorimetry
E _{max}	Maximum Effect
F%	Percent Bioavailability
FaSSIF	Fasted State Simulated Intestinal Fluids
FB	Free Base
FDC	Fixed-Dose Combination
FeSSIF	Fed State Simulated Intestinal Fluid
FT	Fourier Transform
g	Gram
GLSM	Geometric Least Squares Mean
GMR	Geometric Mean Ratio
GT	Genotype
h or hr	Hour
HCV	Hepatitis C virus
HDPE	High Density Polyethylene
HPC	Hydroxypropylcellulose
HPLC	High-performance Liquid Chromatography
HPMC	Hydroxymethylcellulose
ICH	International Conference on Harmonisation; Impurities guidelines
IFN	Interferon
IU	International Unit
KF	Karl Fischer
kg	Kilogram
L	Liter
LCT	Long Chain Triglycerides
LDV	Compound I, GS-5885, Ledipasvir
LLOD	Lower limit of detection
LLOQ	Lower Limit of Quantification
LOD	Limit of Detection
M	Molar
mg	Milligram
min	Minute
mL	Milliliter
mm	Millimeter

mM	Millimolar
N	Population Size
n	Number of Patients
ng	Nanogram
nM	Nanomolar
nm	Nanometer
°C	Degrees Celsius
PD	Pharmacodynamic(s)
PEG or PG	Polyethylene Glycol
P-gp or Pgp	P-glycoprotein
PI	Protease- Inhibitor
PK	Pharmacokinetic
PLS	Partial Least Squares
PPI	Proton-Pump Inhibitors
PS	Particle Size
PVP	Povidone
PVP/VA	Copovidone
QS	Quantum Satis
RAV	Resistance Associated Variants
RBV	Ribavirin
RH	Relative Humidity
RNA	Ribonucleic Acid
RSD	Relative Standard Deviation
RT	Room Temperature
S ₀	Intrinsic Solubility
SAE	Serious Adverse Event
SCT	Short Chain Triglycerides
SIBLM	Simulated Intestinal Bile Salt and Lecithin Mixture
SIF	Simulated Intestinal Fluids
SLS	Sodium Lauryl Sulfate
SOF	Sofosbuvir (GS-7977, formerly PSI-7977)
SS-NMR	Solid State Nuclear Magnetic Resonance
SVR	Sustained Virologic Response
t	Time
t _{1/2}	Half-life (h)

TFA	Trifluoroacetic acid
T _{max}	Time (observed time point) of C _{max}
UPLC	Ultra Performance Liquid Chromatography
Upper Resp Tract Infx	Upper Respiratory Tract Infection
USP	Uniform Standards and Procedures
UV	Ultraviolet
VL	Viral Load
vRVR	Very Rapid Viral Response
V _z /F	Apparent volume of distribution
wt or w	Weight
XRPD	Xray Powder Diffraction
µg	Microgram
µL	Microliter
µm	Micrometer

Example 1: Synthesis of Amorphous Ledipasvir

[0121] Methods for making various forms of ledipasvir may be found in U.S. Publication Nos. 2013/0324740, and 2013/0324496. Both of these applications are incorporated herein by reference. Following is a method for isolating amorphous free base of ledipasvir.

[0122] Combine ledipasvir acetone solvate (191.4 g) and acetonitrile (1356 g) in a reaction vessel and mix contents until a solution is achieved. Add this ledipasvir in acetonitrile solution slowly to another reaction vessel containing vigorously agitated water (7870 g). Agitate contents at about 23 °C for about 30 minutes. Filter the contents and dry at about 40-45 °C until constant weight is achieved to afford ledipasvir amorphous solid (146.4 g, 82% yield).

Example 2: Tablet Preparation and Formulation

A. Dose Selection of Tablets

i. Sofosbuvir

[0123] The sofosbuvir dose selected for the tablet formulation is 400 mg once daily. Support for the 400 mg sofosbuvir dose can be derived from $E_{\max}$ PK/PD modeling with early virological and human exposure data which also supports the selection of a 400 mg sofosbuvir dose over others tested.

[0124] The mean sofosbuvir major metabolite AUC_{τ} for the 400 mg sofosbuvir dose is associated with approximately 77% of the maximal HCV RNA change from baseline achievable as determined by this model, a value which is on the cusp of the plateau of the exposure-response sigmoidal curve. In a sigmoidal $E_{\max}$ model, there is a relatively linear exposure-response relationship in the 20 to 80% maximal effect range. Therefore, given that sofosbuvir exposure with 200 mg tablets appears dose-proportional with single doses up to 1200 mg, doses below 400 mg are expected to yield considerable reductions in the magnitude of HCV RNA change from baseline. Similarly, in order to improve upon an efficacy prediction of 77% in the plateau of the exposure-response curve, substantial increases in exposure (and hence dose) would be needed for an appreciable increase in antiviral effect.

[0125] The sofosbuvir dose of 400 mg once daily was associated with higher SVR rates in genotype 1 HCV infected patients as compared to the 200 mg once daily dose when given in conjunction with additional HCV therapeutics for 24 weeks. Safety and tolerability appeared similar across both dose levels. In addition, when sofosbuvir 400 mg once daily plus other HCV therapeutics were given to genotype 2 or 3 HCV infected patients, 100% SVR24 was observed.

ii. Ledipasvir

[0126] The maximum median HCV RNA log10 reduction was 3 or greater for all cohorts dosed with ≥ 3 mg of ledipasvir. An $E_{\max}$ PK/PD model indicates that the exposures achieved following administration of the 30 mg dose provides $> 95\%$ of maximal antiviral response in genotype 1a HCV infected patients. It was also observed that 30 mg or greater of ledipasvir likely provided coverage of some drug related mutations that doses less than 30 mg did not, based on an analysis of NS5A mutants that arose in response to exposure to ledipasvir. Therefore, 30 mg and 90 mg of ledipasvir were selected as the dose for the formulations described herein.

[0127] Further studies suggest that, when ledipasvir is administered in combination with other therapeutic agents, the breakthrough (BT) rate (number of patients with HCV RNA $>$ lower limit of quantification (LLOQ) after having achieved vRVR/total number of patients who achieved vRVR), is higher with doses of 30 mg (BT = 33%, 11/33; 30 mg ledipasvir), than with doses of 90 mg (BT = 12%, 9/74; 90 mg ledipasvir). Therefore, the 90 mg dose of ledipasvir may confer a greater antiviral coverage that prevents viral breakthrough.

B. Solid Dispersion Comprising Ledipasvir

[0128] To make the tablets comprising the combination of sofosbuvir and ledipasvir as described herein, a solid dispersion comprising ledipasvir was co-formulated with crystalline sofosbuvir. The starting material of the solid dispersion can be a variety of forms of ledipasvir including crystalline forms, amorphous form, salts thereof, solvates and free base, as described herein. Because of the high solubility in organic solvents and excipients and the ability to isolate the ledipasvir free base crystalline acetone solvate, this form was used in the amorphous solid dispersion of ledipasvir.

[0129] The spray dried solid dispersion approach achieved the most desirable characteristics relative to the other formulation approaches, which included improved *in vivo* and *in vitro* performance and manufacturability/scalability.

[0130] The spray dry feed solution was prepared by solubilizing ledipasvir acetone solvate and polymer in the feed solvent. Aggressive mixing or homogenization was used to avoid clumping of the composition.

[0131] Different polymers were tested for preferred characteristics in the solid dispersions. Non-ionic polymers such as hypromellose and copovidone solid dispersions both showed adequate stability and physical characteristics.

[0132] The feed solution was initially evaluated for appropriate solvent with regard to solubility, stability, and viscosity. Ethanol, methanol, and dichloromethane (DCM) all demonstrated excellent solubility (ledipasvir solubility >500 mg/mL). Ethanolic and DCM-based feed stocks were assessed for preparation ease and spray dried at a range of inlet and outlet temperatures to assess the robustness of the spray dry process. Both solvents gave rapid dissolution of ledipasvir and copovidone.

[0133] Spray drying out of ethanol resulted in high yields (88, 90, 92, 94, 95, 97, 98, 99%) across a wide range of spray-drying outlet temperatures (49-70 °C) with no material accumulation on the spray dry chamber. Spray drying out of DCM resulted in yields of 60%, 78%, and 44%. Overall, the ledipasvir Solid Dispersion (50% w/w) in a ledipasvir to copovidone ratio of 1:1 demonstrated good chemical stability in the ethanolic feed solution.

[0134] An ethanolic solution of 10% ledipasvir acetone solvate and 10% copovidone was prepared using homogenization. Viscosity of ethanolic solutions of ledipasvir:copovidone were low, measured through 30% solids content (~65 cP).

[0135] Spray drying was conducted using two fluid nozzle or a hydrolytic pressure nozzle. **Table 1** presents the spray dry process parameters evaluated at 100 g - 4000 g of total feed solution using the Anhydro MS35 spray dryer and **Table 2** shows the spray dry process parameters using the hydrolytic pressure nozzle. Particle size data suggested sufficiently large particle size (10-14 µm mean PS) and was minimally affected by using higher spray rates or a larger diameter spray nozzle. Nozzle gas flow was not modulated to increase particle size.

Table 1. Ledipasvir Spray Dry Parameters on Anhydro MS35 Spray Dryer Using a Two Fluid Nozzle

Parameter	Trial 1	Trial 2	Trial 3	Trial 4
Batch Size (g)	100	250	250	4000
Solids %	20	20	20	20
Feed Rate (mL/min)	30	40	40	40
Spray Nozzle (mm)	1.0	1.0	1.2	1.2
Nozzle Gas Flow (kg/hr)	6.0	6.0	6.0	6.0
Chamber Gas Flow (kg/hr)	35.0	35.0	35.0	35.0
Inlet Temp (°C)	125	165	165	165
Outlet Temp (°C)	70	73	72	76
PS d ₁₀ /d ₅₀ /d ₉₀ /mean (µm)	4/9/18/10	5/10/20/12	5/10/19/11	6/12/22/14
Post Spray LOD (%)	5.56	4.86	4.29	3.42

Table 2. Example of Ledipasvir Spray Dry Parameters Using a Hydrolytic Pressure Nozzle

Parameter	Trial 1
Batch Size (kg)	200
Solids %	20
Feed Rate (kg/hr)	178
Pressure Feed (bar)	52
Inlet Temp (°C)	158
Outlet Temp (°C)	65
PS d ₁₀ /d ₅₀ /d ₉₀ /mean (µm)	3/14/34
Post Spray LOD (%)	0.6

[0136] Organic volatile impurities, including the spray dry solvent ethanol and residual acetone from ledipasvir acetone solvate are rapidly removed during secondary drying at 60 °C. Smaller scale production can be tray dried. On larger scale batches, a double cone dryer or an agitated dryer can be used. Loss on drying (LOD) was proportionately slower and is attributable to water, which was later confirmed by Karl Fischer titration.

[0137] Residual ethanol was reduced below ICH guidelines of 0.5% w/w by 6 hours of drying (or 8 hours for larger scale). Ethanol content upon completion of drying was 0.08% w/w, and residual acetone was 0.002%, indicating that the secondary drying process is adequate for removal of residual solvent.

C. Tablet Preparation

i. Monolayer Tablet

[0138] Ledipasvir:copovidone solid dispersion (1:1) was made by dissolving ledipasvir and copovidone into ethanol, and then spray drying the mixture. The spray dried ledipasvir:copovidone solid dispersion is further dried in a secondary dryer. The amorphous solid dispersion comprising ledipasvir was blended with sofosbuvir and excipients and milled to facilitate mixing and blend uniformity. Either a coblend or codry granulation process can be used. Coblend granulation is a multi-step process consisting of separate dry granulations for each active ingredient with excipients followed by the blending of the two granulations together. Codry granulation consisted of dry granulating both active ingredients and excipients together. The coblend and codry processes demonstrated comparable physical and chemical tablet properties. Exemplary coblend and codry formulations are provided in **Table 3** and **Table 4** shown below.

Table 3. Representative Example Composition of Sofosbuvir/Ledipasvir Codry (Co-granulated) Tablets at Various Fill Weights

Intra-granular	% w/w Tablet			
Sofosbuvir	50.00	40.00	36.36	33.33
Ledipasvir:Copovidone Solid Dispersion (1:1)	22.50	18.00	16.36	15.00
Lactose Monohydrate	6.67	16.33	23.19	26.11
Microcrystalline cellulose	3.33	8.17	11.60	13.05
Croscarmellose Sodium	2.50	2.50	2.50	2.50
Silicon Dioxide	1.00	1.00	1.00	1.00
Magnesium stearate	0.75	0.75	0.75	0.75
Extra-granular				
Microcrystalline cellulose	10.00	10.00	5.00	5.00
Croscarmellose Sodium	2.50	2.50	2.50	2.50
Magnesium stearate	0.75	0.75	0.75	0.75
Fill wt (mg)	800	1000	1100	1200

Table 4. Representative Example Composition of Sofosbuvir/Ledipasvir Coblend (Bi-granulated) Tablets

	Composition	% w/w Intra-granular Blend	% w/w Tablet	mg/Tablet
	Sofosbuvir	80	40	400
	Microcrystalline cellulose	6	3	30

	Composition	% w/w Intra-granular Blend	% w/w Tablet	mg/Tablet
Sofosbuvir Intra-granular Blend	Lactose Monohydrate	6	3	30
	Croscarmellose Sodium	4	2	20
	Silicon Dioxide	3	1.5	15
	Magnesium stearate	1	0.5	5
	Intra-granular Subtotal	100	50	500
Ledipasvir Intra-granular Blend	Ledip asvir: Copovidone Solid Dispersion	42.4	18	180
	Microcrystalline cellulose	43.5	18.5	185
	Croscarmellose Sodium	9.4	4	40
	Silicon Dioxide	3.5	1.5	15
	Magnesium stearate	1.2	0.5	5
	Intra-granular Subtotal	100	42.5	425
Extra-granular	Microcrystalline cellulose	--	5	50
	Croscarmellose Sodium	--	2	20
	Magnesium stearate	--	0.5	5
	Total	--	100	1000
Coating	Film Coat	--	3	30
	Purified water	--	--	--

[0139] The granules were then mixed with a lubricant prior to tablet compression. The total resulting core tablet weight was 1000 mg.

[0140] Film-coating of the tablets is provided to reduce photolytic degradation. Tablets were coated to a target 3% weight gain. The film-coating material was a polyvinylalcohol-based coating. Exemplary tablet formulation is provided in **Table 5**.

Table 5. Representative Example of the Composition of Tablets Comprising the Solid Dispersion of Ledipasvir and Sofosbuvir

Ingredient	% w/w	Component Weight (mg/tablet)
Sofosbuvir	40.00	400
Ledipasvir Solid Dispersion	18.00	180.0
Lactose Monohydrate	16.50	165.0
Microcrystalline Cellulose	18.00	180.0
Croscarmellose Sodium	5.00	50.0
Colloidal Silicon Dioxide	1.00	10.0
Magnesium Stearate	1.50	15
Total Tablet Core Weight	100.0	1000.0
Film coating	3.00	30.0
Purified Water	--	--
Total Coated Tablet Weight		1030.0

ii. Bilayer Tablet

[0141] Tablets comprising the co-formulation of a solid dispersion comprising ledipasvir and crystalline sofosbuvir can also be made as a bilayer tablet wherein each active ingredient is in a separate layer. To make the bilayer tablet, a ledipasvir:copovidone (1:1) solid dispersion is made by dissolving ledipasvir and copovidone into ethanol, and then spray drying the mixture. The spray dried ledipasvir:copovidone solid dispersion is further dried in a secondary dryer. Next, the spray dried ledipasvir:copovidone solid dispersion is then blended with excipients. The mixture is milled and then blended with lubricant prior to dry granulation. The ledipasvir granules are blended with extragranular lubricant. Separately, the sofosbuvir drug substance is blended with excipients, and then the mixture is milled and then blended with lubricant prior to dry granulation. The sofosbuvir granules are then blended with extragranular lubricant. Finally, the sofosbuvir powder blend and ledipasvir powder blend are compressed into bilayer tablet cores. The bilayer tablet cores are then film-coated prior to packaging. A representative example composition of a bilayer tablet comprising the solid dispersion of ledipasvir and sofosbuvir is shown in Table 6. In this table, the solid dispersion comprises ledipasvir:copovidone in a 1:1 ratio.

Table 6. Representative Example of Composition of Bilayer Tablets Comprising the Solid Dispersion of Ledipasvir and Sofosbuvir

Ingredient	% w/w	Component Weight (mg/tablet)
Layer 1		
Sofosbuvir	33.34	400.0
Lactose Monohydrate	5.66	68.0

Ingredient	% w/w	Component Weight (mg/tablet)
Layer 1		
Microcrystalline Cellulose	7.50	90.0
Croscarmellose Sodium	2.00	24.0
Colloidal Silicon Dioxide	0.50	50.0
Magnesium Stearate	1.00	12.0
Layer 2		
Ledipasvir Solid Dispersion	15.00	180.0
Lactose Monohydrate	15.00	180.0
Microcrystalline Cellulose	17.00	204.0
Croscarmellose Sodium	2.50	30.0
Magnesium Stearate	0.50	6.0
Total Tablet Core	100.00	1200

Example 3: PK, Stability and Dissolution Properties of Ledipasvir Single-Agent Tablets and Ledipasvir/Sofosbuvir Tablets and Reduction of Food-Effect and Effects of Gastric Acid Suppressants

A. Ledipasvir Single-Agent Tablets Bioavailability

[0142] A series of *in vivo* experiments were conducted to evaluate the potential benefit of the solid dispersion approach relative to conventional formulations, as well as to optimize the solid dispersion by identifying the most beneficial polymer type and relative polymer concentration within the dispersion.

[0143] Equivalent bioavailability was achieved between formulations comprising the free base amorphous form (4% w/w, 10 mg amorphous free base tablet) and formulations comprising the D-tartrate salt of ledipasvir (5.85% w/w, 10 mg D-tartrate salt tablet), both using conventional formulations, in the pentagastrin pretreated dog model, as shown in **Table 7**. Pentagastrin is a synthetic polypeptide that stimulates the secretion of gastric acid, pepsin, and intrinsic factor.

Table 7. Mean (RSD) Pharmacokinetic Parameters of Ledipasvir Following Oral Administration of Tablets, 25 mg, in Beagle Dogs (n=6)

Drug Substance Form	Pretreatment	C _{max} (nM)	AUC ₀₋₂₄ (nM*hr)	F(%)
Amorphous Free base	Pentagastrin	743 (17)	8028 (22)	71

Drug Substance Form	Pretreatment	C _{max} (nM)	AUC ₀₋₂₄ (nM*hr)	F(%)
Crystalline D-tartrate	Pentagastrin	665 (38)	7623 (44)	67

[0144] Because these formulations displayed similar PK properties and the isolation properties of the D-tartrate salt were preferable to the free base amorphous form, the crystalline D-tartrate salt formulation was chosen to compare to the amorphous solid dispersion compositions. For these studies, 30 mg tablets comprising the crystalline D-tartrate salt of ledipasvir and 30 mg or 90 mg tablets comprising the amorphous solid dispersion of ledipasvir were used. Dog pharmacokinetic results for select immediate release ledipasvir tablets comprising ledipasvir solid dispersions are shown in Table 8.

Table 8. Mean (RSD) Pharmacokinetic Parameters of Ledipasvir after Oral Administration of Ledipasvir Tablets, Fasted Beagle Dogs (n=6)

Polymer	Ledipasvir: polymer Ratio	Dose (mg)	Pretreatment	C _{max} (nM)	AUC ₀₋₂₄ (nM*hr)	F (%)
Crystalline D-tartrate Ledipasvir Tablets	N/A	30	Pentagastrin	665 (38)	7623 (44)	67
			Famotidine	154 (44)	1038 (41)	9
		90	Pentagastrin	1831 (28)	18086 (36)	54
			Famotidine	349 (37)	3322 (40)	10
Amorphous Solid Dispersion Ledipasvir Tablet: HPMC	2:1	30	Famotidine	251 (51)	2553 (54)	22
Amorphous Solid Dispersion Ledipasvir Tablet: Copovidone	2:1	30	Famotidine	369 (26)	3383 (36)	30
	1:1		Pentagastrin	983 (22)	10541 (24)	93
	1:1		Famotidine	393 (30)	3930 (20)	35
	1:1	90	Pentagastrin	1644 (38)	20908 (41)	62
	1:1		Famotidine	740 (24)	7722 (28)	23

[0145] Compared to the crystalline D-tartrate ledipasvir formulations, the amorphous solid

dispersion tablets displayed higher bioavailability with lower variability. In pentagastrin pretreated animals, an approximate 40% increase in exposure and a 2-fold decrease in variability were noted. More importantly in famotidine pretreated animals, up to a 3.5-fold increase in bioavailability was observed compared to the D-tartrate salt tablet formulations.

[0146] A copovidone-based dispersion increased bioavailability more than the equivalent hypromellose-based formulation (F = 30% and 22%, respectively) when spray dried at 2:1 API:polymer ratio. Bioavailability of the copovidone-based formulation was further enhanced by increasing the fraction of polymer to a 1:1 ratio, resulting in a bioavailability of 35% in famotidine pretreated dogs.

[0147] Because of the improved *in vivo* performance and acceptable stability and physical properties, a 1:1 mixture of ledipasvir:copovidone was chosen as the spray-dried material.

[0148] Formulations comprising the amorphous solid dispersions proved to be advantageous over formulations comprising either the amorphous free base or the D-tartrate salt. It was observed that the bioavailability of amorphous free base formulations was similar to D-tartrate salt formulations. Additional data showed a decrease in bioavailability when ledipasvir was dosed with gastric acid suppressing agents (famotidine), indicating an unfavorable drug-drug interaction in free base amorphous and D-tartrate salt formulations of ledipasvir. A solid dispersion using spray drying with a hydrophilic polymer was identified to have acceptable stability, physical characteristics, and *in vivo* performance. A rapidly disintegrating tablet was developed using a dry granulation process and commonly used excipients. A bioavailability study comparing formulations comprising the D-tartrate salt with formulations comprising the amorphous solid dispersion showed improved biopharmaceutical performance and overcame much of the negative drug-drug interactions with acid suppressive therapies seen in the D-tartrate salt formulations.

B. Ledipasvir+Sofosbuvir Tablets Bioavailability

[0149] PK results for the combination of sofosbuvir with ledipasvir (wherein the ledipasvir is in solid dispersion with copovidone in a 1:1) are shown in **Table 9**, and demonstrate lack of a significant interaction between sofosbuvir and ledipasvir.

Table 9: Pharmacokinetic Data for Sofosbuvir and Ledipasvir on Administration of Sofosbuvir and Ledipasvir Alone or in Combination

Sofosbuvir (n=17)			
Mean (%CV)	Sofosbuvir alone	Sofosbuvir + Ledipasvir	%GMR (90%CI)
AUC _{inf} (ng.hr/mL)	794(36.4)	1750 (27.8)	229(191,276)
AUC _{last} (ng.hr/mL)	788 (36.6)	1740 (27.8)	230 (191, 277)
C _{max} (ng/mL)	929(52.3)	1870 (27.9)	221 (176, 278)

Metabolite I (n=17)			
Mean (%CV)	Sofosbuvir alone	Sofosbuvir + Ledipasvir	%GMR (90%CI)
AUC _{inf} (ng.hr/mL)	1110 (31.6)	1950 (22.8)	182 (157, 210)
AUC _{last} (ng.hr/mL)	1060 (32.7)	1890 (22.8)	179 (155, 207)
C _{max} (ng/mL)	312 (38.7)	553 (26.6)	182 (154, 216)
Metabolite II (n=17)			
Mean (%CV)	Sofosbuvir alone	Sofosbuvir + Ledipasvir	%GMR (90%CI)
AUC _{inf} (ng.hr/mL)	10900 (17.5)	13000 (16.7)	119 (113, 125)
AUC _{last} (ng.hr/mL)	10200 (17.9)	12100 (15.5)	119 (113, 126)
C _{max} (ng/mL)	1060 (17.3)	864 (20.1)	81.2 (76.9, 85.8)
Ledipasvir (n=17)			
Mean (%CV)	Ledipasvir alone	Sofosbuvir + Ledipasvir	%GMR (90%CI)
AUC _{inf} (ng.hr/mL)	11900 (26.2)	11400 (27.0)	95.7 (92.1, 99.5)
AUC _{last} (ng.hr/mL)	755 (24.7)	734 (27.0)	96.5 (89.9, 104)
C _{max} (ng/mL)	375 (28.8)	360 (31.2)	95.5 (91.9, 99.1)

[0150] Sofosbuvir plasma exposure was increased by ~ 2.3-fold by ledipasvir. The effect of ledipasvir on sofosbuvir is likely due to inhibition of P-gp, of which Sofosbuvir is a known substrate. The increase in sofosbuvir was not considered significant due to its very low and transient exposure relative to total drug related material (DRM) exposure (DRM, calculated as the sum of the AUCs for each of the analytes, corrected for molecular weight). Based on this calculation, the AUC of sofosbuvir with ledipasvir is only ~ 5.7% of DRM AUC. The exposure of metabolite II, the major circulating sofosbuvir metabolite, was not impacted by the administration of ledipasvir, and demonstrates the lack of significant interaction between sofosbuvir and ledipasvir.

C. Reduction of Food Effect in Solid Dispersions of Ledipasvir and Ledipasvir/Sofosbuvir Tablets

[0151] Ledipasvir alone in a conventional formulation (not the solid dispersion) has been demonstrated to have a negative food effect. **Table 10** summarizes PK parameters of ledipasvir following a single dose of ledipasvir, 30 mg, under fasted and fed conditions. The ledipasvir PK profile was altered in the presence of food. Specifically, the high-fat meal appeared to delay ledipasvir absorption, prolong T_{max} (median T_{max} of 8 hours), and

decreased ledipasvir plasma exposure (approximately 45% decrease each in mean $C_{\max}$, AUC_{last} , and AUC_{inf} , respectively).

Table 10: Plasma Ledipasvir PK Parameters Following Single-dose Administration of Ledipasvir by Concomitant Food Intake Status

PK Parameter	Mean (%CV)	
	Ledipasvir 30 mg (N=8)	Ledipasvir 30 mg Fed (N=8)
$C_{\max}$ (ng/mL)	73.1 (50.8)	36.5 (22.6)
$T_{\max}$ (h)	6.00 (5.00, 6.00)	8.00 (7.00, 8.00)
AUC_{last} (ng·h/mL)	1988.2 (58.2)	996.5 (21.6)
AUC_{inf} (ng·h/mL)	2415.9 (60.3)	1175.0 (25.3)
$t_{1/2}$ (h)	39.82 (33.15, 41.65)	36.83 (22.19, 49.08)
CL/F (mL/h)	17,034.5 (58.6)	26,917.9 (23.6)
V_z/F (mL)	876,546.3 (44.2)	1,386,469 (24.9)
C_{last} (ng/mL)	6.8 (68.0)	3.1 (42.2)

[0152] Table 11 presents the ratio of the GLSMs (ledipasvir 30 mg under fasted conditions/ledipasvir 30 mg under fed conditions) for each of the primary PK parameters.

Table 11: Statistical Evaluations of Ledipasvir PK Parameters for Food Effect

	Geometric Least Squares Mean (GLSM)		GLSM Ratio (Fed/Fasted) %	90% Confidence Interval
	Ledipasvir 30 mg Fed (N=8)	Ledipasvir 30 mg Fasted (N=8)		
$C_{\max}$ (ng/mL)	35.87	65.33	54.90	39.10, 77.08
AUC_{last} (ng·hr/mL)	977.76	1724.28	56.71	38.87, 82.73
AUC_{inf} (ng·hr/mL)	1143.64	2058.78	55.55	36.88, 83.67

[0153] Similar median half-lives of ledipasvir were observed independent of administration under fasted or fed conditions ($t_{1/2}$ of 39.82 hours under fasted conditions vs 36.83 hours under fed conditions) indicating that food decreased the bioavailability of ledipasvir by reducing its solubility and/or absorption.

[0154] Because ledipasvir has been demonstrated to have a negative food effect, the composition comprising both sofosbuvir and ledipasvir (as solid dispersion in copovidone (1:1))

was tested for a food effect. These results are shown in **Table 12**. Food slowed the rate of absorption of sofosbuvir (median T_{max} : 1.00 vs 2.00 hours) with only modest alteration in the bioavailability, as evidenced by increases of 2-fold or less in sofosbuvir and sofosbuvir metabolite I plasma exposure. For sofosbuvir metabolite II, an approximately 20-30% lower C_{max} was observed upon sofosbuvir administration with food with no change in AUC. The %GMR and associated 90% CI (fed/fasted treatments) for AUC of sofosbuvir metabolite II were within the equivalence bounds of 70% to 143%. Since the decrease in sofosbuvir metabolite II C_{max} was modest and the AUC parameters met the equivalence criteria, the effect of food on sofosbuvir metabolite II was not considered significant.

[0155] Similar ledipasvir plasma exposures (AUC and C_{max}) were achieved upon administration of ledipasvir under fasted or fed conditions. The %GMR and associated 90% CIs (fed/fasted treatments) were within the equivalence bounds of 70-143%. While a "negative" food effect was previously observed on ledipasvir when administered alone (as the amorphous free base, not solid dispersion), the pharmacokinetics of ledipasvir (amorphous solid dispersion; copovidone (1:1)) administered in combination with sofosbuvir does not appear to be altered by food. As such, the combination of sofosbuvir and ledipasvir may be administered without regard to food.

Table 12: Pharmacokinetic Data for Sofosbuvir, Sofosbuvir Metabolites I and II, and Ledipasvir on Administration of Sofosbuvir/Ledipasvir Tablets Fasted or with a Moderate-Fat Meal or with a High-Calorie/High Fat Meal

Sofosbuvir (n=29)			
Mean (%CV)	Sofosbuvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet Moderate-Fat Meal	% GMR (90% CI) [Moderate- Fat/Fasted]
AUC_{inf} (ng.hr/mL)	1520 (39.5)	2860 (33.4)	195 (176,216)
AUC_{last} (ng.hr/mL)	1520 (39.7)	2850 (33.5)	195 (176,216)
C_{max} (ng/mL)	1240 (49.6)	1520 (39.8)	126 (109, 147)
	Sofos buvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet High- Calorie/High-Fat Meal	GMR (90% CI) [High- Fat/Fasted]
AUC_{inf} (ng.hr/mL)	1520 (39.5)	2570 (34.0)	179 (162, 198)
AUC_{last} (ng.hr/mL)	1520 (39.7)	2550 (34.6)	178 (161, 198)
C_{max} (ng/mL)	1240 (49.6)	1350 (42.5)	115 (99.0, 134)

Sofosbuvir Metabolite I (n=29)			
Mean (%CV)	Sofosbuvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet Moderate-Fat Meal	% GMR (90% CI) [Moderate- Fat/Fasted]
AUC _{inf} (ng.hr/mL)	1520 (42.0)	2520 (21.4)	177 (163, 192)
AUC _{last} (ng.hr/mL)	1470 (43.3)	2460 (21.8)	180 (164, 196)
C _{max} (ng/mL)	352 (42.7)	495 (22.2)	151 (136, 167)
	Sofos buvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet High- Calorie/High-Fat Meal	GMR (90% CI) [High- Fat/Fasted]
AUC _{inf} (ng.hr/mL)	1520 (42.0)	2550 (22.2)	181 (166, 196)
AUC _{last} (ng.hr/mL)	1470 (43.3)	2500 (22.5)	184 (168, 201)
C _{max} (ng/mL)	352 (42.7)	501 (26.8)	154 (139, 171)
Sofosbuvir Metabolite II (n=29)			
Mean (%CV)	Sofosbuvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet Moderate-Fat Meal	% GMR (90% CI) [Moderate- Fat/Fasted]
AUC _{inf} (ng.hr/mL)	11800 (23.0)	13800 (17.7)	117 (112, 123)
AUC _{last} (ng.hr/mL)	11300 (23.4)	12900 (18.2)	114 (108, 121)
C _{max} (ng/mL)	865 (26.6)	700(19.5)	81.5 (75.6, 87.9)
	Sofos buvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet High- Calorie/High-Fat Meal	GMR (90% CI) [High- Fat/Fasted]
AUC _{inf} (ng.hr/mL)	11800 (23.0)	12900 (18.5)	112 (107, 118)
AUC _{last} (ng.hr/mL)	11300 (23.4)	12100 (20.1)	110 (103, 116)
C _{max} (ng/mL)	865 (26.6)	600 (22.9)	70.2 (65.0, 75.8)

Ledipasvir (n=29)			
Mean (%CV)	Sofosbuvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet Moderate-Fat Meal	% GMR (90% CI) [Moderate- Fat/Fasted]
AUC _{inf} (ng.hr/mL)	10600 (57.2)	10600 (35.6)	115 (99.4, 134)
AUC _{last} (ng.hr/mL)	8600 (53.8)	8650 (32.1)	114 (98.0, 133)
C _{max} (ng/mL)	324 (44.8)	319 (24.8)	109 (93.5, 126)
	Sofosbuvir/Ledipasvir Tablet Fasted	Sofosbuvir/Ledipasvir Tablet High- Calorie/High-Fat Meal	GMR (90% CI) [High- Fat/Fasted]
AUC _{inf} (ng.hr/mL)	10600 (57.2)	9220 (36.1)	103 (88.5, 119)
AUC _{last} (ng.hr/mL)	8600 (53.8)	7550 (33.9)	104 (88.8, 121)
C _{max} (ng/mL)	324 (44.8)	255 (25.9)	88.2 (75.8, 103)

D. Reduction of Effects of Gastric Acid Suppressants in Ledipasvir/Sofosbuvir Tablets

[0156] Ledipasvir, 30 mg, alone in both a conventional formulation (as the D-tartrate salt) and as the solid dispersion has been demonstrated to have a decrease in bioavailability when administered with some gastric acid suppressants; most significantly, proton-pump inhibitors (PPI's, e.g., omeprazole), but also including histamine-2 antagonists (H2RA's, e.g., famotidine, data not included). **Table 12A** summarizes PK parameters of ledipasvir following administration of ledipasvir conventional single agent tablets, 30 mg, ledipasvir tablets as solid dispersion (ledipasvir:copovidone 1:1), 30 mg, and sofosbuvir/ledipasvir FDC tablets (90 mg of ledipasvir solid dispersion comprising copovidone 1:1) with and without omeprazole. The bioavailability of ledipasvir as single agent tablets was reduced approximately 2-fold when administered with omeprazole; however, administration of ledipasvir as part of the sofosbuvir/ledipasvir FDC tablet with omeprazole resulted in no significant decrease in ledipasvir exposure (AUC and C_{max}) compared to sofosbuvir/ledipasvir FDC tablet administration in absence of omeprazole.

Table 12A. Pharmacokinetic Data for Ledipasvir on Administration of Ledipasvir Single Agent Tablets or Sofosbuvir/Ledipasvir Tablets with and without Omeprazole

Ledipasvir, Conventional Formulation (N = 10)			
Mean (%CV)	Ledipasvir alone	Ledipasvir + Omeprazole	%GMR (90%CI)
AUC _{tau} (ng.hr/mL)	1640 (18.5)	865 (37.7)	50.7 (43.4, 59.3)

Ledipasvir, Conventional Formulation (N = 10)			
Mean (%CV)	Ledipasvir alone	Ledipasvir + Omeprazole	%GMR (90%CI)
C _{max} (ng/mL)	99.0 (20.1)	51.2 (39.2)	49.7 (41.7, 59.1)
C _{tau} (ng/mL)	52.2 (22.1)	28.3 (36.0)	52.4 (44.3, 61.9)
Ledipasvir, Solid Dispersion (N = 17)			
Mean (%CV)	Ledipasvir alone	Ledipasvir + Omeprazole	%GMR (90%CI)
AUC _{inf} (ng.hr/mL)	2140 (38.8)	1300 (50.7)	58.5 (48.3, 70.8)
AUC _{last} (ng.hr/mL)	1850 (33.5)	1070 (45.5)	56.3 (46.4, 68.3)
C _{max} (ng/mL)	64.8 (32.9)	36.2 (55.9)	52.2 (41.4, 65.9)
Ledipasvir, SOF/LDV FDC (N = 16)			
Mean (%CV)	SOF/LDV FDC Alone	SOF/LDV FDC + Omeprazole	%GMR (90%CI)
AUC _{inf} (ng.hr/mL)	7990 (66.2)	6660 (51.8)	96.0 (66.5, 139)
AUC _{last} (ng.hr/mL)	7160 (65.8)	5700 (51.8)	92.5 (64.8, 132)
C _{max} (ng/mL)	242(68.6)	176(51.1)	89.1 (60.9, 130)

E. Dissolution of Ledipasvir/Sofosbuvir Tablets

[0157] Dissolution studies were conducted comparing the sofosbuvir 400 mg/ledipasvir 90 mg tablets (ledipasvir:copovidone (1:1)). The sofosbuvir/ledipasvir tablets (LOT 1-5) display greater than 85% sofosbuvir (FIG. 5) and ledipasvir (FIG. 6) dissolved in 30 minutes for both tablet formulations. These results are shown in FIGs. 5 and 6.

Example 4: Stability of Sofosbuvir/Ledipasvir Co-formulation

[0158] The compatibility of sofosbuvir anhydrous crystalline drug substance was evaluated with the ledipasvir:copovidone solid dispersion. A blend of the sofosbuvir and ledipasvir:copovidone (1:1) solid dispersion was prepared at a ratio representative of the final 400 mg sofosbuvir/90 mg ledipasvir tablets. The blend was compressed into pellets and placed in stability chambers at 40 °C/75% RH and 60 °C/ambient humidity and tested after two and four weeks of storage in open glass vials. The results summarized in Table 13 show that no degradation was observed for either sofosbuvir or ledipasvir, demonstrating the chemical compatibility of sofosbuvir and the ledipasvir:copovidone solid dispersion with each other.

Table 13: Strength and Impurity Content of Sofosbuvir and Ledipasvir:Copovidone

Solid Dispersion Blend Stored at 40 °C/75% RH and 60 °C

Condition	Time (weeks)	Ledipasvir		Sofosbuvir	
		Strength (%)	Total Impurity Content (%)	Strength (%)	Total Impurity Content (%)
45 °C/75 % RH	0	98.8	0.0	102.9	0.4
	2	96.9	0.0	101.6	0.3
	4	97.1	0.0	100.5	0.2
60 °C	0	98.8	0.0	102.9	0.4
	1	99.2	0.0	102.4	0.3
	2	99.6	0.0	103.2	0.3
	4	98.9	0.0	102.8	0.2

Example 5: Efficacy of Sofosbuvir/Ledipasvir/Ribavirin Treatment in Patients with HCV Infections

[0159] Patients with HCV infections were treated with either the combination of sofosbuvir, ledipasvir, and ribavirin or the combination of sofosbuvir and ribavirin. Patients used in the study included those that were treatment naïve, i.e. had not previously been treated for HCV and those that were null responders, i.e. had previously been treated for HCV but failed to respond to the treatment. Standard doses (90 mg of ledipasvir, 400 mg of sofosbuvir, and 1000 mg of ribavirin, for example) were given of each drug to the patients for a duration of 12 weeks. Throughout treatment, HCV RNA was measured, and the Sustained Virologic Response (SVR) was measured after treatment was discontinued. By four weeks of treatment, almost all patients had achieved an HCV RNA measurement below the limit of detection (LOD of 15 IU/mL), and by the end of treatment, 100% of patients achieved an HCV RNA level below the LOD (Table 14).

Table 14: Patients with HCV RNA below the limit of detection over time.

	Sofosbuvir + Ribavirin		Sofosbuvir + Ribavirin + Ledipasvir	
	Treatment-naïve (n = 25)	Null responder (n = 10)	Treatment-naïve (n = 25)	Null responder (n = 9)
Week 1	32%	10%	44%	0%
Week 2	68%	70%	88%	44%
Week 4	100%	100%	100%	89%
End of Treatment	100%	100%	100%	100%

[0160] Surprisingly, 100% of patients receiving the combination of sofosbuvir, ledipasvir, and

ribavirin achieved a sustained virologic response at four and twelve weeks post treatment. In contrast, only 88% of treatment naïve and 10% of null responder patients treated with the combination of sofosbuvir and ribavirin achieved a SVR at four weeks post treatment, and only 84% of treatment naïve and 10% of null responder patients treated with the combination of sofosbuvir and ribavirin achieved SVR at twelve weeks post treatment (**Table 15**).

Table 15: Sustained Virologic Response

	Sofosbuvir + Ribavirin		Sofosbuvir + Ribavirin + Ledipasvir	
	Treatment-naïve (n = 25)	Null responder (n = 10)	Treatment-naïve (n = 25)	Null responder (n = 9)
SVR4	88%	10%	100%	100%
SVR12	84%	10%	100%	100%

[0161] These results are graphically depicted as **FIG. 7A-D** and demonstrate that the addition of ledipasvir in the treatment regimen gave 100% SVR at weeks 4 and 12. Example 9, below, shows similar results are obtained with treatment regimens of less than twelve weeks (i.e. treatment regimens of about 8 or 6 weeks), and that similar results are obtained with treatment regimens of sofosbuvir and ledipasvir without the addition of ribavirin.

Example 6. Stability of SOF 400 mg/ Ledipasvir 90 mg Fixed-Dose Combination Tablets

[0162] This example summarizes the physicochemical stability of packaged Sofosbuvir (SOF) 400 mg/ ledipasvir 90 mg blue film-coated fixed-dose combination (FDC) tablets at 25 °C/60% relative humidity (RH) and 40 °C/75% RH as a function of desiccant. The ledipasvir portion of the table comprised ledipasvir:copovidone in a 1:1 ratio. In addition the chemical and physical stability of SOF/ledipasvir FDC tablets were evaluated at 40°C/75% RH under open condition for up to 4 weeks.

[0163] The physico-chemical properties that were evaluated included appearance, potency, degradant formation, dissolution rate and water content. Physical stability of the tablets in the absence of desiccant was evaluated after 24 weeks using FT-Raman spectroscopy and modulated differential scanning calorimetry (mDSC).

[0164] SOF 400 mg/ledipasvir 90 mg blue film-coated FDC tablets exhibited satisfactory stability at 25°C/60% RH and 40°C/75%RH for up to 24 weeks in the presence of 0, 1, and 3 g of desiccant. No significant changes were observed in potency, impurity content or dissolution rate. However, a ledipasvir photodegradant was present at 0.1% for all conditions. FT-Raman analysis for the tablets stored in the absence of desiccant showed no detectable crystallization after 24-weeks.

Methods and Materials

Materials

[0165] Table 16 lists the physicochemical properties for SOF drug substance and ledipasvir solid dispersion used to produce tablets. The quantities of SOF drug substance and ledipasvir solid dispersion were adjusted based on their respective drug content factor (DCF) with concomitant adjustment in the quantity of lactose monohydrate. The DCF used for SOF and ledipasvir solid dispersion powder, 50% w/w were 0.997 and 0.497 (0.994 when adjusted for the amount of copovidone), respectively.

Table 16. Physicochemical Properties of SOF Drug Substance and Ledipasvir Solid Dispersion, 50% w/w, Bulk Powder Used to Produce SOF 400 mg/Ledipasvir 90 mg Film-Coated FDC Tablets

Active Ingredient	Crystal Form	Assay by HPLC (%)	Impurities (%)	Drug Content Factor	Water Content by Karl Fischer (%)	Particle Size (µm)		
						d ₁₀	d ₅₀	d ₉₀
SOF	Anhydrous Form II	99.8	0.1	0.996	0.1	3	10	29
Ledipasvir Solid Dispersion, 50% w/w, Bulk Powder	Amorphous	49.7	0.2	0.497	1.09	5	22	44

Equipment

[0166] The primary equipment used to manufacture SOF 400 mg/Ledipasvir 90 mg film-coated FDC tablets included an 12 qt. V- Blender, a screening mill (Comil 197S, Quadro, Waterloo, Canada) equipped with a 0.094 in grated screen, a roller compactor/granulator (MiniPactor, Gerteis, Jona, Switzerland) equipped equipped with a 1.0 mm milling screen and a smooth/smooth roller configuration, a 12-station instrumented rotary tablet press (XM-12, Korsch, Berlin, Germany), and a tablet coater (LabCoat, O'Hara Technologies Inc., Ontario, Canada). The diamond-shaped tablet tooling (Elizabeth Carbide Die Co., Inc., McKeesport, PA, USA) consisted of diamond, standard concave D-type punches with dimensions of 0.7650 in x 0.4014 in (19.43 mm x 10.20 mm). A 15 inch perforated pan film coater was used to coat the tablet cores.

Container Closure

[0167] Sofosbuvir/Ledipasvir FDC tablets are packaged in 100 mL white, high density polyethylene (HDPE) bottles. Each bottle contained 30 tablets and 0, 1 or 3 g silica gel desiccant canister or sachet and polyester packing material. Each bottle was enclosed with a white, continuous thread, child-resistant screw cap with an induction-sealed, aluminum-faced liner.

[0168] A selected number of bottles were left open and packaged without desiccant to evaluate the physical and chemical stability at 40°C/75% RH under accelerated heat and humidity conditions.

General Study Design

[0169] The solid state and chemical stability of the packaged lot were evaluated in the following configurations:

1. 1) At 25 °C/60% RH and 40 °C/75% RH as a function of desiccant. The samples were stored under closed condition for a minimum of 24 weeks.
2. 2) At 40°C/75% RH under open condition for up to 4 weeks.

[0170] Samples were pulled at predetermined time points. Chemical stability testing for appearance, potency, degradant formation, dissolution rate and water content was conducted. Additional physical stability assays to monitor potential crystallization and phase separation were conducted.

Physical Stability Evaluation

[0171] Physical stability tests included appearance and FT-Raman. The visual inspection was performed on stressed film-coated tablets to identify changes in tablet color and coating integrity. FT-Raman spectroscopy was used to detect potential crystalline ledipasvir (Form III) in the film-coated tablets.

[0172] Tablets were visually inspected for changes in appearance at all time points and storage conditions. In contrast, FT-Raman was only performed on tablets with 0 g desiccant at 24 weeks (25°C/60% RH and 40°C/75% RH).

Appearance

[0173] At all time points tablets were examined for physical integrity (i.e. color, shape, coating

integrity and debossing).

FT-Raman

[0174] FT-Raman experiments were conducted. The 24-week SOF/ledipasvir film-coated FDC tablets stored in closed containers at 25°C/60% RH and 40°C/75% RH were analyzed using FT-Raman spectroscopy to detect the formation of crystalline ledipasvir (Form III). Briefly, the coating from the tablets was carefully removed using an Xacto™ knife followed by grinding of the tablet in a mortar and pestle. Tablet powder was then packed into cups and spectra were collected using a backscattering geometry.

Chemical Stability Evaluation

[0175] Chemical stability assays included measuring water content by Karl Fischer (KF), potency, formation of impurity/degradation products and dissolution rate were conducted.

KF Water Content

[0176] The water content was reported for SOF 400 mg/ledipasvir 90 mg film-coated FDC tablets following USP <921>.

Potency and Impurity/Degradant Formation by UPLC

[0177] The potency and degradation product formation of SOF/ledipasvir film-coated FDC tablets were evaluated by analysis of composite sample solution of 10 tablets according to STM-2542 [5]. The reference standard concentration for SOF and ledipasvir is 2.0 mg/mL and 0.45 mg/mL, respectively. The strength and degradation product content of SOF and ledipasvir was determined by UPLC using external reference standard and area normalization at wavelengths of 262 nm and 325 nm, respectively.

Dissolution Methodology

[0178] Dissolution testing was performed on SOF/ledipasvir film-coated FDC tablets. A USP type 2 dissolution apparatus with 900 mL of dissolution medium and a paddle speed of 75 rpm was used. The medium was 1.5% polysorbate 80 in 10 mM potassium phosphate buffer at pH 6.0 and the temperature was maintained at 37 °C for the duration of the assay. The extent of SOF and ledipasvir released as a function of time was monitored by UPLC using area

normalization and an external reference standard at a wavelength of 250 nm.

RESULTS

A. Physical Stability

A1. Appearance

[0179] Samples at all stability conditions and desiccant levels were visually inspected for all time points and found to resemble blue, diamond-shaped film-coated tablets.

A2. FT-Raman

[0180] The FT-Raman analysis was performed on powder extracted from tablets stored in the absence of desiccant after 24 weeks. Calculations of % crystallinity using the PLS model did not show signs of crystalline ledipasvir (Form III) above the LOD of 3% at either storage condition. This was consistent with the original sample ($t = 0$) in which ledipasvir (Form III) was also below the LOD. Spectra from selected samples were included in a chart, from 1577 cm^{-1} to 1514 cm^{-1} with the baselines artificially adjusted for clarity. This region is in one of the four spectral regions used to estimate the % ledipasvir (Form III) in tablets by PLS model.

[0181] The top two spectra (used as standards in the PLS model), in the chart, were from tablets spiked with 10% w/w and 3% w/w, of crystalline ledipasvir (Form III). The next two spectra represent stressed tablets stored for 24 weeks at $40\text{ }^{\circ}\text{C}/75\%\text{ RH}$ and $25\text{ }^{\circ}\text{C}/60\%\text{ RH}$. The last spectrum represents the initial time point ($t = 0$). Ledipasvir (Form III) has a distinct peak at 1552 cm^{-1} , which can clearly be seen in the spiked tablets with increasing intensity from 3% to 10%. The intensity in this region for the stressed samples stored for 24 weeks does not increase from the $t = 0$ sample, indicating no change in crystallinity. Ledipasvir (Form III) in the $t = 0$ sample and the 24 week samples is below that present in the tablets spiked with 3% Form III ledipasvir, the current limit of detection for this analytical technique.

B. Chemical Stability

B.1 KF Water Content

[0182] The water content of stressed samples stored for 4 weeks under open condition increased from 2.28% to 5.23%. The amount of water content of stressed samples stored at 25 °C/60%RH decreased to 1.91%, 1.58%, and 1.65% for tablets with no desiccant, with 1g desiccant, and 3g desiccant, respectively. At 40 °C/75%RH, the amount of water content decreased to 2.03%, 1.79%, and 1.46% for tablets without desiccant, with 1g desiccant, and 3g desiccant, respectively.

B.2 Potency and Impurity/Degradation Product Formation

[0183] The potency and impurity/degradation content for SOF 400 mg/ledipasvir 90 mg film-coated FDC tablets were determined at 25 °C/60%RH and 40 °C/75%RH. Representative chromatograms of stability samples stored at 40 °C/75%RH were obtained. The data showed that SOF and ledipasvir remained chemically stable in SOF 400 mg/ledipasvir 90 mg film-coated FDC tablets stored for 24 weeks at 25 °C/60%RH and 40 °C/75%RH. The label strength for SOF and ledipasvir remains unchanged at 25°C/60%RH and 40°C/75%RH.

Dissolution

[0184] The dissolution profiles of SOF and ledipasvir in SOF 400 mg/ledipasvir 90 mg film-coated FDC tablets were obtained. At the 24 week time point, the tablets ranged between 99% and 100% dissolution at 45 minutes for SOF, and between 99% and 98% for ledipasvir at both 25°C/60%RH and 40°C/75% RH for all desiccant levels tested.

[0185] From the foregoing, this example shows that SOF 400 mg/ledipasvir 90 mg Film-Coated FDC tablets exhibited satisfactory stability at 25°C/60% RH and 40 °C/75%RH for up to 24 weeks in the presence of 0, 1, and 3 g of desiccant. In addition, crystalline ledipasvir (Form III) was not detected by FT-Raman analysis after 24 weeks of storage.

Example 7. Formulation Development of a Fixed Dose Combination (FDC) Tablet SOF 400 mg/Ledipasvir 90 mg

[0186] This example shows the development of a SOF 400 mg/ledipasvir 90 mg fixed dose combination (FDC) tablet comprising ledipasvir: copovidone (1:1). There were expected difficulties with such a development, one of which was the expected poor powder flow and the other relates to non-homogenous blend, given the existing formulations of each individual agent.

[0187] Three tablet formulations were tested, including (1) a monolayer co-granulated tablet formulation, (2) a monolayer co-blended tablet formulation and (3) a bilayer tablet formulation. In all of these formulations, SOF was in anhydrous crystalline form II and ledipasvir was in

amorphous solid dispersion (ledipasvir:copovidone (1:1)).

[0188] Formulation (1) is typically associated with the highest risk of drug-drug interaction but is the most cost-effective during manufacturing. The bilayer formation of (3), by contrast, is perceived to have the lowest drug-drug interaction risk.

[0189] The dissolution performance of the formulations were tested in a dissolution media that included 10 mM phosphate buffer at pH 6.0 (1.5% Tween® 80). As shown in **FIG. 8A-B**, all three formulations had comparable dissolution performance, similar to that of the single-agent controls.

[0190] The pharmacokinetic (PK) performance of each formulation was also tested. Plasma Concentration of SOF/ledipasvir after oral administration of SOF/ledipasvir FDC and control tablets in fasted dogs (100 mg/22.5 mg fixed/dog). **Table 17** below shows the PK results.

Table 17. Pharmacokinetic performance of the formulations in famotidine pretreated dogs

Total Tablet weight /Formulation	Treatment	SOF		Ledipasvir	
		AUC _{0-t} (ng*hr/mL)	C _{max} (ng/mL)	AUC _{0-last} (ng*hr/mL)	C _{max} (ng/mL)
Control SOF tablet + Ledipasvir SD tablet	Famotidine	314 ± 207	503 ± 363	3260 ± 1312	345 ± 132
Monolayer, Co-granulated	Famotidine	501 ± 249	729 ± 434	3236 ± 730	333 ± 56
Monolayer, Co-blended	Famotidine	483 ± 406	652 ± 527	4208 ± 2216	444 ± 215
Bilayer	Famotidine	283 ± 193	288 ± 201	4,712 ± 2,270	421.7 ± 203.7

[0191] Based on these results, the monolayer co-granulated tablet was selected for further analysis. The composition of this formulation is provided in **Table 18**.

Table 18. Composition of SOF 400 mg/Ledipasvir 90 mg FDC Tablets

Composition	% w/w
Intra-granular	
SOF	40.00%
Ledipasvir SD	18.00%
Lactose Fast Flow 316	16.50%
MCC 101	8.00%
Croscarmellose	2.50%
Silicon Dioxide	1.00%
Magnesium Stearate	0.75%

Extragranular	
MCC 101	10.00%
Croscarmellose	2.50%
Magnesium stearate	0.75%
Total Fill weight Core Tablet (mg)	1000
Coating	
Opadry II Orange 85F13912	3.0%
Water	QS

[0192] A bioavailability clinical study was carried out with this formulation, with single agent tablets as control, in 24 healthy patients under fasted conditions. The results are shown in Table 19.

Table 19. Bioavailability of SOF/Ledipasvir fixed dose combination and single agent tablets

Total Tablet weight /Formulation	Dose (mg)	SOF		Ledipasvir	
		AUC _{inf} (ng*hr/mL)	C _{max} (ng/mL)	AUC _{inf} (ng*hr/mL)	C _{max} (ng/mL)
Single Agent Control SOF tablet + Ledipasvir SD tablet	SOF 400 mg Ledipasvir 90 mg	11900 (23.5)	764 (27.3)	9620 (45.6)	314 (40.5)
SOF/Ledipasvir FDC Tablet		12500 (23.1)	784 (36.2)	9570 (46.6)	314 (45.2)

[0193] These results, therefore, show that SOF/ledipasvir fixed dose combination (co-granulated) and single agent tablets are bioequivalent.

Example 8. Solubility Studies for Amorphous Ledipasvir

[0194] This example examines the physicochemical properties different ledipasvir forms, including amorphous and crystalline free base, solvates, and salts, with respect to solubility.

A. Materials and Methods

pH-Solubility Profile

[0195] The aqueous solubility of ledipasvir amorphous free base was determined across the pH range of 1 to 10. Excess solid ledipasvir was added to a range of pH-adjusted aqueous solutions (titrated with HCl or NaOH) and stirred for 48 hours at room temperature. The suspensions were then filtered through regenerated cellulose syringe filters. The pH value of the supernatant was measured, and the supernatant was diluted as appropriate with 50:50 H₂O+0.1% TFA:ACN and assayed for ledipasvir content by the HPLC-UV method.

Solubility in Simulated Intestinal Media

[0196] Solubility of ledipasvir amorphous free base was assessed in three types of simulated intestinal fluids at pH 6.5 or pH 5.0; and simulated intestinal bile salt and lecithin mixture (SIBLM), pH 6.4. Excess solid ledipasvir was added to the respective SIFs and stirred for 48 hours at room temperature. The resulting suspensions were then filtered through regenerated cellulose syringe filters. The supernatant was diluted as appropriate with 50:50 H₂O+0.1% TFA: ACN and assayed for ledipasvir content by the HPLC-UV method.

Excipient Solubility

[0197] Solubility of ledipasvir amorphous free base and ledipasvir crystalline D-tartrate was measured in a wide range of pharmaceutically acceptable solvents, including cosolvents, surfactants, fatty acids, triglycerides, or blends thereof. Material was weighed into scintillation vials and stirred for up to 48 hours at room temperature. In many cases, solubility was higher than the amount of solid used in the sample, thus many results are reported as 'greater than' or 'greater than or equal to' if the concentration was not quantitatively determined by HPLC-UV.

[0198] Additionally, aqueous solubility was measured as a function of time in the presence of 0.1% w/w surfactants and polymers in pH 2 (50 mM citrate) and pH 5 (50 mM citrate). ledipasvir crystalline forms (acetone solvate Form II; anhydrous FB Form III; D-tartrate) and amorphous form were evaluated to identify differences in dissolution behavior. Excess solid was added to aqueous buffered solutions; samples were withdrawn at predetermined intervals (2, 5, 8, 10, 15, 20, 30, 45, 60 minutes, and 24 hours), filtered through regenerated cellulose filters, and diluted for concentration measurement by the HPLC-UV method.

B. Results

Solubility and Dissolution Rate

[0199] The pH-solubility profiles of all available ledipasvir forms were determined at room

temperature and are graphically shown in **FIG. 9**. The flat portion of the solubility profile ($\text{pH} > 5$) represents the intrinsic aqueous solubility of the free base. The aqueous solubility of ledipasvir significantly increases as the pH of the solution is lowered below the pK_a of the ionizable groups. All forms lose crystallinity, reverting to the amorphous free base in aqueous solution, and thus show similar aqueous solubility properties at steady-state. However, dissolution properties are form dependent and are described in further detail below.

Ledipasvir Amorphous Free Base

[0200] The intrinsic solubility of ledipasvir amorphous free base (FB) is approximately 0.04 $\mu\text{g/mL}$. Under acidic conditions, the solubility increases to 1 mg/mL at pH 2.3, and peaks at about 7 mg/mL at pH 1.6, as shown in **Table 20** and **FIG. 9**. Solubility of ledipasvir in simulated intestinal fluids is governed by both the pH of the medium and the presence of bile salts and lecithin. In fasted state simulated intestinal fluids (FaSSIF) at pH 6.5 and room temperature, the solubility is 0.025 mg/mL , and this is increased approximately 10-fold to 0.232 mg/mL in simulated bile and lecithin mixture (SIBLM, pH 6.5) due to the increased concentration of bile salts and lecithin. A similar solubility enhancement to 0.230 mg/mL is observed in fed state simulated intestinal fluid (FeSSIF, pH 5), containing lower bile salt and lecithin mixtures than SIBLM. The solubility increase in this mixture is predominantly attributed to the ionization state of the molecule at pH 5.

Table 20. Solubility of Ledipasvir amorphous free base as a function of pH at room temperature

Aqueous Media	Solubility (mg/mL)
Aqueous, pH 1.6 (HCl)	6.855
Aqueous, pH 2.3 (HCl)	1.096
Aqueous, pH 3.1 (HCl)	0.0132
Aqueous, pH 4.1 (HCl)	0.00011
Aqueous, pH 5.5 (HCl)	0.00003
Aqueous, pH 6.2 (unaltered)	0.00003
Aqueous, pH 7.2 (NaOH)	0.00001
FaSSIF ¹ pH = 6.5	0.025
FeSSIF ² pH = 5.0	0.230
SIBLM ³ pH = 6.4	0.232

¹ FaSSIF is water with 3mM sodium taurocholate and 0.75 mM lecithin, pH adjusted to 6.5 with phosphate buffer, ionic strength adjusted to 0.15M with NaCl.

² FeSSIF is water with 15 mM sodium taurocholate and 3.75 mM lecithin, pH adjusted to 6.5 with phosphate buffer, ionic strength adjusted to 0.15M with NaCl.

³ SIBLM is water with 30 mM sodium glycocholate, 30 mM sodium glycochenodesoxycholate, 15 mM sodium glycodesoxycholate, 10 mM sodium

taurocholate, 10 mM sodium taurochenodesoxycholate, 5 mM sodium taurodesoxycholate, 50 mM sodium chloride, and 11 mM lecithin, pH adjusted to 6.4 with phosphate buffer, ionic strength adjusted to 0.15M with NaCl.

[0201] The dissolution rate of ledipasvir amorphous free base at pH 3 and 6 was also tested. At pH 3, the dissolution of the amorphous free base form is faster than that of the crystalline free base and acetone solvate forms. However, at pH 6, all free base forms show similar dissolution rate profiles.

[0202] As shown in **Table 21**, ledipasvir amorphous free base is freely soluble (>500 mg/mL) in ethanol and other organic solvents such as propylene glycol and PEG 400. Its solubility is greater than 200 mg/mL in surfactants (e.g., polysorbate 80, Cremophor EL, Labrasol) and lipid blends. Its solubility in oleic and octanoic acids is greater than 500 mg/mL. Solubility of ledipasvir in short-chain triglycerides (SCTs, tributyrin) is limited to 20 mg/mL, and decreases to less than 1 mg/mL in long-chain triglycerides (LCTs, soybean oil). It has a solubility of 25 mg/mL in the vehicle chosen for toxicological studies: 45% propylene glycol, 15% caprylocaproyl macrogol-8 glycerides (Solutol HS 15®), and 40% water (pH 2.5 by HCl).

Table 21. Solubility of Ledipasvir free base forms and Ledipasvir D-tartrate in organic solvents and excipients at room temperature

	Solubility (mg/mL)			
	Amorphous Free Base	Crystalline Acetone Solvate	Anhydrous Crystalline Free Base	Crystalline D-tartrate Salt
	(Ledipasvir)	(Ledipasvir-03)	(Ledipasvir Form III)	(Ledipasvir-02)
Acetone	5	5	--	<1
Acetonitrile	>500	--	--	12
Methanol	>500	>500	--	23
95% Methanol + 5% water	--	--	--	19
Ethanol	>500	>500	>500	4
95% Ethanol + 5% water	--	--	--	5
PEG 400	>500	>500	>500	4
Propylene glycol	>500	>500	>500	6
Octanoic acid	>500	>500	-	<1
Oleic acid	>500	>500	>500	<1
Polyoxyl 35 Castor Oil (Cremophor EL)	>200	--		--

	Solubility (mg/mL)			
	Amorphous Free Base	Crystalline Acetone Solvate	Anhydrous Crystalline Free Base	Crystalline D-tartrate Salt
	(Ledipasvir)	(Ledipasvir-03)	(Ledipasvir Form III)	(Ledipasvir-02)
Polysorbate 80 (Tween 80)	>200	>200	>200	3
Caprylocaproyl macrogolglycerides (Labrasol)	>300	>300	>300	3
Tributyrin	9	--	--	--
Soybean oil	2	2	2	--
RSSEDDS ¹	>500	--	--	10
¹ RSSEDDS: 10% Ethanol, 10% PG, 40% Solutol HS-15, 40% Labrasol				

[0203] Dilute nonionic surfactants generally increase ledipasvir solubility at both pH 2 and 5, as presented in **Table 22**. Similar effects were observed with nonionic polymers, though to a lesser extent. Sodium lauryl sulfate (SLS), an anionic surfactant, improves the solubility of ledipasvir at pH 5. However, a significant decrease in solubility is noted in presence of SLS under acidic conditions (pH 2). This observation is consistent with weakly basic compounds that have low intrinsic aqueous solubility, presumably forming an insoluble estolate salt.

Table 22. Solubility of Ledipasvir amorphous free base in surfactant or polymeric excipients (0.1% w/w) diluted into aqueous media at pH 2 and 5 at room temperature

Excipient (0.1% w/w in aqueous media)	Solubility (mg/mL)	
	pH 2	pH 5
No excipient	4.94	0.0001
Sodium lauryl sulfate	0.05	0.243
Labrasol	7.44	--
Cremaphor EL	9.11	0.0699
Polysorbate 80	9.27	0.0624
Poloxamer 188	7.19	0.0005
HPC (hydroxypropylcellulose)	4.67	0.0001
HPMC (hydroxymethylcellulose)	5.27	0.0003
PVP (povidone)	5.47	0.0004
PVP/VA (copovidone)	6.73	0.0010

Ledipasvir Crystalline Acetone Solvate (Ledipasvir-03)

[0204] The ledipasvir acetone solvate (ledipasvir-03) showed similar steady-state solubility as the other forms. Ledipasvir-03 has the slowest dissolution of all forms tested. Its dissolution at pH 6 was indistinguishable from that of other forms due to poor intrinsic solubility (<0.1 ng/mL).

[0205] Ledipasvir-03 is soluble in many organic solvents and pharmaceutically acceptable solvents, and the solubilities are comparable to those listed for ledipasvir amorphous free base, as also shown in **Table 21**.

Ledipasvir Crystalline Free Base (Form III)

[0206] Ledipasvir crystalline free base Form III showed similar steady-state solubility as the other forms (**FIG. 9**). This form dissolves more slowly than the amorphous free base, but faster than ledipasvir-03. Dissolution at pH 6 was indistinguishable from that of the other forms due to poor intrinsic solubility (<0.1 µg/mL). Solubility in a wider range of organic vehicles has not been explored, though is anticipated to be similar to other free base forms.

Ledipasvir Crystalline D-tartrate Salt (Ledipasvir-02)

[0207] Ledipasvir crystalline D-tartrate salt (ledipasvir-02) showed similar steady-state solubility as the other forms (**FIG. 9**). Dissolution behavior ledipasvir-02 is improved relative to all free base forms. At pH 3, ledipasvir-02 shows a roughly 5- to 10-fold faster initial dissolution rate than the free base forms, and roughly doubled the amount of ledipasvir in solution through 60 minutes compared to the amorphous form. At pH 6, the increased dissolution rate was also apparent. However, rapid dissociation of the salt at this pH resulted in equivalent solubility values to other forms within minutes.

[0208] Ledipasvir-02 is not soluble in various organic media, as shown in **Table 21**. Maximal solubility of ledipasvir-02 in any organic vehicle is 20 mg/mL in methanol; this limits the use of ledipasvir-02 in solubilized formulations or processes that require solubilization in organic media.

[0209] Ledipasvir has low aqueous solubility and high permeability, and is considered a BCS Class 2 compound. The data presented in this example indicate that in water, all forms of ledipasvir: the amorphous free base, crystalline free base acetone solvate (ledipasvir-03), crystalline anhydrous free base (Form III), and crystalline D-tartrate salt (ledipasvir-02), convert to the amorphous free base, and have similar aqueous solubility at steady state. The aqueous solubility of ledipasvir is less than 0.1 µg/mL in its neutral form (pH > 5), but substantially increases under acidic conditions due to protonation of two basic moieties. The aqueous dissolution rate of ledipasvir amorphous free base is faster than that of crystalline free

base forms. However, all free base forms have slower dissolution rates than the crystalline D-tartrate salt (ledipasvir-02). Ledipasvir-02 also shows improved wetting in aqueous media. ledipasvir free base forms, crystalline and amorphous, are highly soluble in a range of cosolvents and surfactants. In contrast, ledipasvir-02 is poorly soluble in organic excipients, and this property potentially limits its utility.

[0210] Ledipasvir amorphous free base was used in Phase 1 clinical studies, but drug substance manufacturing was identified as a critical limitation of the form. Ledipasvir crystalline D-tartrate salt (ledipasvir-02) was then identified as part of a more extensive salt and form screen and was used in Phase 2, however, poor solubility in organic excipients limits its utility in non-conventional formulations. Crystalline ledipasvir acetone solvate (ledipasvir-03) is used to develop a spray dried dispersion formulation to support future clinical studies due to its solubility in organic solvents and excipients relative to crystalline ledipasvir D-tartrate salt and improved manufacturability over the other free base forms.

Example 9: Efficacy of a Fixed Dose Combination of Sofosbuvir and Ledipasvir With and Without Ribavirin in Patients with HCV Infections

[0211] Patients with HCV infections were treated with the fixed dose combination of sofosbuvir and ledipasvir, with and without ribavirin. Patients used in the studies include those that were treatment naive (non-cirrhotic), i.e. had not previously been treated for HCV, and those that were prior protease-inhibitor (PI) failures and null responders (with and without cirrhosis), i.e. had previously been treated for HCV but failed to respond to the treatment. The treatment naive patients were treated for 6, 8, and 12 weeks and the null responders were treated for 12 weeks.

Study 1

[0212] Cohort 1 of study 1 included treatment-naive, Genotype-1 patients without cirrhosis. The patients were randomized 1:1:1 into three groups to receive 1) SOF/ledipasvir fixed dose combination for 8 weeks, 2) SOF/ledipasvir fixed dose combination with ribavirin for 8 weeks, or 3) SOF/ledipasvir fixed dose combination for 12 weeks (**FIG. 10**).

[0213] Cohort 2 of study 1 included Protease-Inhibitor treatment-experienced, Genotype -1 patients (Prior Protease-Inhibitor treatment failures, 50% of whom had compensated cirrhosis). The patients were randomized to receive 12 weeks of: 1) SOF/ledipasvir fixed dose combination or 2) SOF/ledipasvir fixed dose combination with ribavirin (**FIG. 10**). In Cohort 2, the patients must not have discontinued prior therapy due to an adverse event.

[0214] In study 1, there was a broad inclusion criteria, namely, there was no upper limit to age or BMI. Platelets were $\geq 50,000/\text{mm}^3$. The demographics of study 1 are shown in **Table 23**,

below.

Table 23. Demographics

	SOF/Ledipasvir fixed dose combination ± ribavirin
	(Cohort 1 and 2) N=100
Mean age, y (range)	50 (21-73)
Male, n (%)	66 (66)
Black, n (%)	9 (9)
Hispanic, n (%)	40 (40)
Mean BMI, kg/m ² (range)	29.9 (18-48)
IL28B CC, n (%)	15(15)
GT 1a, n (%)	87(87)
Mean baseline HCV RNA, log ₁₀ IU/mL (range)	6.1 (3.7-7.2)
	Cohort 2 (N=40)
Cirrhosis, n (%)	22/40 (55)
Mean Platelet Count (x10 ³ / µL)	107
Mean Albumin (g/dL)	3.8

[0215] Of 100 patients enrolled in study 1, 97% achieved sustained viral response. Of the failures, two patients relapsed (one from Group 1 (i.e., SOF/Ledipasvir x 8 Weeks) and one from Group 4 (i.e., SOF/ledipasvir x 12 Weeks), and one patient was lost to follow up from Group 3 (i.e., SOF/ledipasvir x 12 Weeks). However, the patient lost to follow up had achieved SVR at week 8 and declined further return visits.

[0216] In Cohort 1 of study 1 (i.e. Treatment Naive, Non-Cirrhotic Patients), 58 out of the 60 patients treated for 8 or 12 Weeks achieved SVR. In cohort 2 of study 1 (i.e. Treatment Experienced, PI failure Patients), 39 out of the 40 patients treated for 12 Weeks achieved SVR12. 21 out of the 21 patients with cirrhosis achieved SVR12 (**FIG. 11**).

[0217] In study 1, seven out of the nine patients with NS5A Resistance Associated Variants (RAVs) achieved sustained viral response. In addition, all patients with NS3/4A Resistance Associated Variants achieved sustained viral response. Interestingly, S282T mutation and multiple NS5A RAVs were detected at relapse in the patient who failed from the Group 1 (**Table 24**). The safety summary and a breakdown of the adverse effects are shown in **Tables 25** and **26**, respectively.

Table 24. Resistance analysis

	SOF/Ledipasvir fixed dose combination ± ribavirin
NS5A RAVs, n %	9/100 (9)

	SOF/Ledipasvir fixed dose combination ± ribavirin
NS3/4A RAVs, n %	29/40 (73)*
*number of patients in Cohort 2 with prior exposure to a protease inhibitor	

Table 25. Safety Summary

		SOF/Ledipasvir fixed dose combination	SOF/Ledipasvir fixed dose combination + ribavirin
	Patients, n (%)	N=58	N=42
Overall safety	AEs	24 (41%)	24 (57%)
	Grade 3-4 AEs	0	6 (14%)
	Serious AEs	2* (3%)	2** (5%)
	Treatment discontinuation due to AEs	0	0
Laboratory abnormalities	Grade 3-4 laboratory abnormality	4 (7%)	6 (14%)
	Hemoglobin <10 g/dL	0	8 (19%)
	Hemoglobin <8.5 g/dL	0	2 (5%)
*peptic ulcer, spinal compression fracture			
**delerium, suicidal ideation			

Table 26. Adverse Events (≥ 5% of patients overall)

	SOF/Ledipasvir fixed dose combination	SOF/Ledipasvir fixed dose combination + ribavirin
Preferred term, n (%)	N=58	N=42
Any adverse event	24 (41%)	24 (57%)
Nausea	3 (5%)	6 (14%)
Anemia	0	8 (19%)
Upper Resp Tract Infx	4 (7%)	4 (10%)
Headache	3 (5%)	4 (10%)

Study 2

[0218] In study 2, the treatment-naïve patients received SOF/ledipasvir fixed dose combination with ribavirin and prior null responders, all of whom had cirrhosis, were randomized to receive twelve weeks of: 1) SOF/ledipasvir fixed dose combination or 2) SOF/ledipasvir fixed dose combination with ribavirin.

Results

[0219] Of the 144 patients treated in both studies 1 and 2, 136 out of 144 (94%) achieved SVR at four weeks post treatment. Of the 85 treatment-naïve patients in these two studies, three of 25 patients failed to achieve SVR after 6 weeks of SOF/ledipasvir fixed dose combination with ribavirin therapy, whereas 100% (60/60) patients achieved SVR after 8 or 12 weeks of SOF/ledipasvir fixed dose combination with and without ribavirin therapy. Of the 59 treatment-experienced patients in these two studies, three cirrhotic patients relapsed after receiving 12 weeks of SOF/ledipasvir fixed dose combination without ribavirin. Conversely, no virologic failures were observed in the SOF/ledipasvir fixed dose combination with ribavirin treatment groups, but two patients in these groups were lost to follow-up. SOF/ledipasvir fixed dose combination with and without ribavirin was well tolerated, with few SAEs and minimal adverse events.

Conclusion

[0220] SOF/ledipasvir fixed dose combination +/- ribavirin may be given for as little as 8 weeks to treatment-naïve non-cirrhotic patients. Treatment-experienced patients, even those with cirrhosis, achieved high SVR rates with 12 weeks of the of SOF/ledipasvir fixed dose combination with and without ribavirin therapy.

Example 10: Efficacy of Multiple Anti-HCV Combination Therapy in Chronically Infected Hepatitis C Patients

[0221] To evaluate the safety, tolerability, and efficacy of 4 to 12 weeks of SOF with ledipasvir, alone or in combination with Compound E and/or Compound J in patients with HCV, patients with HCV will be dosed as shown in Table 27.

Table 27. Dosing

Group	Treatment	Dosing	Patient description
Group A	12 weeks of SOF/ledipasvir	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) administered orally for 12 weeks	Patients (n=20) monoinfected with HCV genotype 1 who are HCV treatment naïve
Group	6 weeks of	SOF with ledipasvir (400	Patients (n=20)

Group	Treatment	Dosing	Patient description
B	SOF/ledipasvir/ Compound E	mg/90 mg respectively once a day in a fixed dose combination) in combination with Compound E (500 mg once daily) for 6 weeks	monoinfected with HCV genotype 1 who are HCV treatment naive
Group C	6 weeks of SOF/ledipasvir/ Compound J	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) in combination with Compound J (80 mg once daily) for 6 weeks.	Patients (n=20) monoinfected with HCV genotype 1 who are HCV treatment naive
Group D	12 weeks of SOF/ledipasvir	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) administered orally for 12 weeks	Patients (n=up to 25) monoinfected with HCV genotype 1 who were previously treated in Group B, C, F, G or H of this study or a similar study
Group E	12 weeks of SOF/ledipasvir	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) administered orally for 12 weeks	Patients (n=20) monoinfected with HCV genotype 4 who are HCV treatment naive or treatment experienced
Group F	6 weeks of SOF/ledipasvir/ Compound J	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) in combination with Compound J (80 mg once daily) for 6 weeks.	Patients (n=50) monoinfected with HCV genotype 1 with advanced liver disease who are HCV treatment naive (n=25) or treatment experienced (n=25)
Group G	4 weeks of SOF/ledipasvir/ Compound J	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) in combination with Compound J (80 mg once daily) for 4 weeks	Patients (n=25) monoinfected with HCV genotype 1 who are HCV treatment naive, Stage 0-2 liver disease
Group H	4 weeks of SOF/ledipasvir/ Compound J/Compound E	SOF with ledipasvir (400 mg/90 mg respectively once a day in a fixed dose combination) in combination with Compound J (80 mg once daily) and Compound E (250 mg once daily) for 4 weeks	Patients (n=25) monoinfected with HCV genotype 1 who are HCV treatment naive, Stage 0-2 liver disease

[0222] The primary analysis set for safety analyses will include patients who received at least

one dose of study drug. On treatment data will be analyzed and defined as data collected from the first dose of study drug through the date of last dose of study drug plus 30 days. Patients who receive study drug other than that to which they were assigned will be analyzed according to the study drug received.

[0223] The analysis set for antiviral activity analyses will include patients who were enrolled into the study and received at least one dose of study drug.

[0224] The pharmacokinetic analysis set will include all patients who are enrolled and have received at least one dose of study medication.

[0225] The patient will be started on study treatment after confirming eligibility on Day 0 and after being informed fully about the remainder of the study, and then signing the specific consent for the treatment group (if not done previously). Blood will be drawn for HCV viral loads, study drug levels, lipid levels for research if not already drawn during screening, immunologic studies, and for storage prior to dosing as part of the screening consent. A pregnancy test will be done for females with childbearing potential and the pregnancy test must be negative on Day 0 prior to dosing with study drugs. Patients may be asked to fill out a baseline adherence questionnaire and an electronic pill bottle cap, which records pill bottle openings will be placed on all study drug bottles. Assistance will be provided filling out the questionnaire as needed. Patients on Arms B and H, will also be provided with a diary at Day 0, Week 2, Week 4 (Arm B only) on which to record gastrointestinal side effects.

[0226] On arrival at the clinic for scheduled study visits, patients will have their vital signs obtained, females will undergo a pregnancy test (if appropriate per schedule and of childbearing potential), clinical laboratories drawn, and a review of the study restrictions.

[0227] At each scheduled study visit (does not include Day 1, 3, 5, 10, Week 2, Week 3, Week 6 (not applicable for arm F, G or H) or post-treatment week 2 and 8 which are for lab collection only), patients will be asked about their state of health and use of any concomitant medication since the previous study visit. They will also be questioned about adverse events and their adherence with study restrictions. Vital signs, weight and examination will be performed as per the study flow. A complete list of study procedures and lab tests to be performed is in the Schedule of Tests, below. In addition, patients may be seen at unscheduled visits for a grade 3 or 4 adverse event or any unexpected adverse event or potential toxicity.

[0228] Patients may be asked to fill out a follow-up adherence questionnaire and pill bottle openings may be recorded from the electronic bottle cap at Day 7 (Group A), Week 4 (Group A), Week 6 (Groups B and C), Week 8 (Group A), and Week 12 (Group A). Assistance will be provided filling out the questionnaire as needed.

[0229] Patients in Groups B and H will be asked to bring their side-effect diaries to visits on Week 2, 4, 6 (B only).

[0230] Some of the visits have a small amount of flexibility regarding when they need to occur. Visits occurring during the interval when the patient is receiving study drug have limited flexibility since they occur so frequently, so a visit skipped during this period may be considered a missed visit. The window period for visit schedules is as shown in **Table 28**.

Table 28. Window Period for Visit Schedule

For 12 week regimen, Group A:			
Days 0, 1, 3 (no window)	Days 5, 7, 10, 14 (+/- 2 days)	Weeks 3, 4, 6 (+/- 3 days)	Weeks 8, 12 (+/- 5 days) Optional Week 12 Research Liver Biopsy (+/-14 days)
For 6 week regimens, Groups B & C :			
Days 0, 1, 3 (no window)	Days 5, 7, 10, 14 (+/- 2 days)	Weeks 3, 4, 6 (+/- 3 days) Optional Week 6 Research Liver Biopsy (+/-14 days)	
For 12 week regimens, Groups D and E:			
Day 0 (no window)	Week 4 (+/- 3 days)	Weeks 8, 12 (+/- 7 days)	
For 6 week regimen, Group F:			
Day 0 (no window)	Weeks 2, 4 (+/-3 days)	Week 6 (+/- 5 days) Optional Week 6 Research Liver Biopsy (+/-14 days)	
For 4 week regimen, Group G or H:			
Day 0 (no window)	Day 7 (+/- 2 days) Group H only	Weeks 2, 4 (+/- 3 days) Optional Week 4 Research Liver Biopsy (+/-14 days)	

[0231] During the four week visit, HCV RNA may be obtained to determine if virologic-response based treatment stopping criteria have been met. Patients who fail to achieve >2 log 10 HCV RNA drop at this time (unless > 2 log drop would be below LLOQ) should be discontinued from therapy unless a review by the PI/LAI/Sponsor Medical Monitor determines otherwise (see 9.3.1).

[0232] At the end of treatment duration as determined by the study group, patients may discontinue dosing of SOF and ledipasvir, Compound E, and/or Compound J. In addition, if a patient's participation terminates prior to completion of pre-specified study drug duration, the End of Treatment assessments may be performed at any end-of-treatment visit. An optional research liver biopsy for research purposes may be performed at this time in up to 10 patients in each study group. The additional liver biopsy data will serve to explore hepatic HCV RNA sequence analysis. If patients are undergoing the optional research liver biopsy, they may have safety labs completed prior to the procedure and imaging as medically indicated. Patients who have a HCV VL< LLOQ may receive education about how to prevent re-infection with HCV.

[0233] All patients may be assessed for sustained virologic response at the 12 Weeks Post End of Treatment visit. Patients who have HCV VL < LLOD may be provided with education about how to prevent re-infection with HCV.

[0234] After discontinuation of the study drug, patients may be followed at 2, 4, 8, 12, 24, 36, and 48 weeks post-end of treatment. A serum pregnancy test may be done with each visit, as appropriate. Week 2 and 8 Post-End of Treatment may include only collection of labs.

[0235] Subjects (n=18) received single doses of sofosbuvir (400 mg) alone or in combination with Compound E (500 mg QD) under fed conditions. Preliminary PK results for the combination of sofosbuvir with Compound E are presented in **Table 29** and demonstrate lack of a clinically significant interaction between sofosbuvir and Compound E.

Table 29: Pharmacokinetic Data for SOF, Compound E, and ledipasvir alone and upon co-administration

SOF (n=18)			
Mean (%CV)	SOF alone	SOF + Compound E	%GMR(90%CI)
AUC _{inf} (ng.hr/ml)	921 (61.2)	1150(40.2)	135 (116, 159)
		SOF + Compound E+ ledipasvir	%GMR(90%CI)
		2560 (42.9)	297 (253, 348)
AUC _{last} (ng.hr/ml)	908 (62.2)	SOF + Compound E	%GMR(90%CI)
		1140(41.4)	135 (115, 159)
		SOF + Compound E+ ledipasvir	%GMR(90%CI)
C _{max} (ng/ml)	515(78.3)	2550 (43.1)	301 (255, 354)
		SOF + Compound E	%GMR(90%CI)
		587(51.1)	130 (97.1, 175)
		SOF + Compound E+ ledipasvir	%GMR(90%CI)
		1260(55.1)	283 (219.366)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [WO2010132601A](#) [\[0002\]](#) [\[0003\]](#) [\[0037\]](#)
- [US7964580B](#) [\[0003\]](#) [\[0054\]](#)
- [US20130324740A](#) [\[0037\]](#) [\[0121\]](#)
- [US20130324496A](#) [\[0037\]](#) [\[0121\]](#)
- [US6763607B](#) [\[0052\]](#)
- [US20060189633A](#) [\[0052\]](#)
- [US20100016251A](#) [\[0054\]](#)
- [US20100298257A](#) [\[0054\]](#) [\[0054\]](#) [\[0054\]](#) [\[0055\]](#)
- [US20110251152A](#) [\[0054\]](#) [\[0054\]](#) [\[0054\]](#) [\[0055\]](#)
- [US20120107278A](#) [\[0054\]](#)
- [WO2013185093A](#) [\[0093\]](#)
- [US20130102525A](#) [\[0099\]](#) [\[0103\]](#) [\[0105\]](#) [\[0112\]](#)
- [US20140017198A](#) [\[0107\]](#)
- [US8178491B](#) [\[0109\]](#)

Non-patent literature cited in the description

- Remington's Pharmaceutical SciencesMace Publishing Co.19850000 [\[0058\]](#)
- Modern PharmaceuticsMarcel Dekker, Inc. [\[0058\]](#)


Patentkrav**1. Farmaceutisk sammensætning omfattende:**

- 5 a) fra ca. 10% til ca. 25% vægt/vægt af en fast dispersion omfattende ledipasvir dispergeret i en polymermatriks dannet af copovidon, hvor vægtforholdet mellem ledipasvir og copovidon i den faste dispersion er ca. 1:1 og hvor ledipasvir har formelen:



og er amorf;

- 10 b) fra ca. 35% til ca. 45% vægt/vægt sofosbuvir med formelen:



hvor sofosbuvir er krystallinsk og det krystallinske sofosbuvir har XRPD 2θ-reflektioner ($^{\circ} \pm 0,2\theta$) ved: 6,1 og 12,7;

- c) fra ca. 5% til ca. 25% vægt/vægt lactosemonohydrat;
- 15 d) fra ca. 5% til ca. 25% vægt/vægt mikrokrySTALLinsk cellulose;
- e) fra ca. 1% til ca. 10% vægt/vægt croscarmellose-natrium;
- f) fra ca. 0,5% til ca. 3% vægt/vægt kolloidalt siliciumdioxid; og
- g) fra ca. 0,1% til ca. 3% vægt/vægt magnesiumstearat.

2. Farmaceutisk sammensætning ifølge krav 1, omfattende

- 20 a) ca. 40% vægt/vægt sofosbuvir og

b) ca. 18% vægt/vægt af en fast dispersion omfattende ledipasvir.

3. Farmaceutisk sammensætning ifølge krav 1 eller 2 til anvendelse i behandlingen af en patient inficeret med hepatitis C-virus.

5 **4.** Farmaceutisk sammensætning til anvendelse ifølge krav 3, hvor den farmaceutiske sammensætning eller doseringsform administreres i ca. 24 uger eller derunder, fortrinsvis i ca. 12 uger eller derunder, mere fortrinsvis i ca. 8 uger eller derunder, og især i ca. 6 uger eller derunder.

10 **5.** Farmaceutisk sammensætning til anvendelse ifølge krav 3, hvor den farmaceutiske sammensætning eller doseringsform administreres én gang dagligt i ca. 12 uger eller derunder, fortrinsvis én gang dagligt i ca. 8 uger eller derunder, and more fortrinsvis én gang dagligt i ca. 6 uger eller derunder, og hvor hepatitis C-viruset er genotype 1, 2, 3, 4, 5, eller 6.

15

6. Farmaceutisk sammensætning til anvendelse ifølge krav 5, hvor hepatitis C-viruset er genotype 1a eller 1b.

7. Farmaceutisk sammensætning til anvendelse ifølge krav 5, hvor hepatitis
20 C-viruset er genotype 2.

8. Farmaceutisk sammensætning til anvendelse ifølge krav 5, hvor hepatitis C-viruset er genotype 3.

25 **9.** Farmaceutisk sammensætning til anvendelse ifølge krav 3, hvor den farmaceutiske sammensætning administreres én gang dagligt i ca. 12 uger, fortrinsvis én gang dagligt i ca. 8 uger, og hvor hepatitis C-viruset er genotype 1a, 1b, 2a, 2b, 2c, 2d, 3a, 3b, 3c, 3d, 3e, 3f, 4a, 4b, 4c, 4d, 4e, 4f, 4g, 4h, 4i, 4j, 5a, eller 6a.

30

10. Farmaceutisk sammensætning til anvendelse ifølge et hvilket som helst af kravene 3 til 9, hvor behandlingen omfatter administrationen af et yderligere terapeutisk middel.

11. Farmaceutisk sammensætning til anvendelse ifølge krav 10, hvor det yderligere terapeutiske middel er ribavirin.

12. Farmaceutisk sammensætning til anvendelse ifølge krav 10, hvor det
5 yderligere terapeutiske middel er en NS3-proteaseinhibitor.

13. Farmaceutisk sammensætning til anvendelse ifølge krav 10, hvor det yderligere terapeutiske middel er simeprevir.

DRAWINGS




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11