

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 3641795 T3**

-
- (51) Int.Cl.: **A 61 K 35/50 (2015.01)** **A 61 L 27/38 (2006.01)** **A 61 L 29/08 (2006.01)**
A 61 L 29/16 (2006.01) **A 61 M 25/10 (2013.01)**
- (45) Oversættelsen bekendtgjort den: **2023-02-27**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2022-11-30**
- (86) Europæisk ansøgning nr.: **18819951.7**
- (86) Europæisk indleveringsdag: **2018-06-21**
- (87) Den europæiske ansøgnings publiceringsdag: **2020-04-29**
- (86) International ansøgning nr.: **US2018038815**
- (87) Internationalt publikationsnr.: **WO2018237179**
- (30) Prioritet: **2017-06-22 US 201762523366 P** **2017-09-19 US 201715708406**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Petrucci, Gary M., 2975 County Road 24, Long Lake, MN 55356, USA**
- (72) Opfinder: **Petrucci, Gary M., 2975 County Road 24, Long Lake, MN 55356, USA**
- (74) Fuldmægtig i Danmark: **Potter Clarkson A/S, Havnegade 39, 1058 København K, Danmark**
- (54) Benævnelse: **METHODS AND MATERIALS FOR TREATING BLOOD VESSELS**
- (56) Fremdragne publikationer:
WO-A1-2016/007554
WO-A1-2016/198670
WO-A2-2012/112410
US-A- 5 674 192
US-A1- 2009 125 044
US-A1- 2014 236 161
US-A1- 2016 136 334
US-B2- 8 524 462
US-B2- 9 205 177
ROBERTSON L. et al.: "Angioplasty and stenting for peripheral arterial disease of the lower limbs: an overview of Cochrane Reviews (Protocol)", Cochrane Database of Systematic Reviews, 1 February 2017 (2017-02-01), XP055560230,
"The crucial, controversial carotid artery Part I: The artery in health and disease", HARVARD HEALTH PUBLISHING, vol. 16, no. 1 1 August 2011 (2011-08-01), pages 1-5, XP009518468, ISSN: 1089-1102 Retrieved from the Internet: URL:https://www.health.harvard.edu/heart-health/the-crucial-controversial-carotid-artery-part-i-the-artery-in-health-and-disease

DESCRIPTION

CLAIM OF PRIORITY

[0001] This application claims priority to U.S. Application No. 15/708,406, filed September 19, 2017, which claims the benefit of U.S. Provisional Application Serial No. 62/523,366, filed on June 22, 2017.

BACKGROUND

1. Technical Field

[0002] This document relates to amnion coated angioplasty balloon catheters for treating a mammal (e.g., a human) having one or more stenotic blood vessels.

2. Background Information

[0003] Angioplasty is frequently used to treat heart disease and peripheral artery disease (PAD). Heart disease is the leading cause of death for both men and women, and PAD affects at least 8 to 12 million Americans with prevalence increasing with age (CDC, *Heart Disease Facts*. Available online at cdc.gov/heartdisease/facts.htm). Each year, more than 1 million patients in the United States have angioplasty (American Heart Association. Heart Disease and Stroke Statistics-2007 Update. Dallas, TX: American Heart Association; 2007), but up to one-quarter of people who have angioplasty must have it repeated, or, in the case of heart disease, have bypass surgery within a few years (Harvard Health Publications, "Angioplasty or bypass surgery?" Harvard Heart Letter April 2008).

SUMMARY

[0004] This invention relates to an angioplasty balloon catheter comprising an angioplasty balloon, wherein said angioplasty balloon comprises a coating comprising as sole active ingredient, an amnion tissue preparation; or as sole active ingredients, an amnion tissue preparation and one or more therapeutic agents selected from the group consisting of a corticosteroid, a glucocorticoid, a non-steroidal anti-inflammatory drug, dexamethasone, an antibiotic, an antifungal agent, an antiproliferative drug, or an immunosuppressive drug.

[0005] This document provides methods and materials for treating a mammal (e.g., a human)

having one or more stenotic blood vessels, which are not part of the present invention. For example, amnion coated balloons provided herein (e.g., an angioplasty balloon coated with an amnion tissue preparation) can be used in balloon angioplasty. In some cases, amnion coated balloons provided herein can be used to treat a mammal having one or more stenotic blood vessels. Typically, a stent is placed into a blood vessel following balloon angioplasty to decrease recoil (e.g., restenosis) of the blood vessel. In some cases, amnion coated balloons provided herein can be used in stent-less balloon angioplasty (e.g., a balloon angioplasty in which a stent is not implanted).

[0006] The amnion tissue preparation can have viable cells or the amnion tissue preparation can lack viable cells. The amnion tissue preparation can be a dried amnion tissue preparation. A dried amnion tissue preparation can have a water content less than about 8 percent. A dried amnion tissue preparation can have a particle size ranging from about 0.1 μm to about 25 μm . The amnion tissue preparation can be the sole active ingredient on the amnion coated balloon or the amnion coated balloon also can be coated with one or more therapeutic agents. The angioplasty balloon catheter can be a stent-less angioplasty balloon catheter.

[0007] In another aspect, this document features methods for treating stenotic blood vessel in a mammal. The methods can include, or consist essentially of, inserting an angioplasty balloon into an obstructed blood vessel, where the angioplasty balloon has a coating including an amnion tissue preparation, and inflating the amnion coated balloon to perform angioplasty in the obstructed vessel. In some cases, one or more symptoms associated with said stenotic blood vessel can be reduced. When the stenotic blood vessel is a peripheral artery, the symptom can be claudication, cramps, tingling, pain, numbness, wounds, reduced body temperature, and/or poor nail growth, in the arms and/or legs. When the stenotic blood vessel is a stenotic coronary artery, the symptom can be stroke, heart attack, chest pain, shortness of breath, sweating, nausea, dizziness or light-headedness, palpitations, and/or arrhythmias. When the stenotic blood vessel is a stenotic carotid artery, the symptom can be weakness, confusion, difficulty speaking, dizziness, difficulty walking, difficulty standing, blurred vision, numbness, severe headache, and/or losing consciousness. When the stenotic blood vessel is a stenotic renal artery stenosis, the symptom can be decreased kidney blood flow, and/or chronic kidney disease. In some cases, the obstructed blood vessel is widened. The obstructed blood vessel can be widened to be about 20% to about 100% open. In some cases, the obstruction in said obstructed blood vessel is reduced. The obstruction can be reduced by about 20% volume to about 100% volume. In some cases, the blood flow in the obstructed blood vessel can be increased. The blood flow can be increased by about 20% to about 100%. The blood flow can be increased to between about 50 cm/s and about 250 cm/s. The mammal can be a human. The amnion coated balloon can be inflated to from about 5.1 bar (5 atmospheres (atm)) to about 35.5 bar (35 atm) of pressure. The amnion coated balloon can be inflated to less than about 10.1 bar (10 atm) of pressure. The angioplasty can have less than about 50% recoil. The angioplasty balloon can be inserted through a percutaneous puncture in the arm or groin. The obstruction can include an atherosclerotic plaque (e.g., a calcified atherosclerotic plaque). The method also can include (e.g., prior to said angioplasty) performing atherectomy in the obstructed vessel. The method can be a stent-less balloon

angioplasty.

[0008] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used to practice the invention, suitable methods and materials are described below. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

[0009] The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DETAILED DESCRIPTION

[0010] This document provides angioplasty balloons coated with an amnion tissue preparation, as well as methods and materials for making and using amnion coated balloons. For example, amnion coated balloons provided herein (e.g., an angioplasty balloon coated with an amnion tissue preparation) can be used in balloon angioplasty. In some cases, amnion coated balloons provided herein can be used to treat a mammal (e.g., a human) having one or more stenotic blood vessels. For example, an amnion coated balloon can be used in balloon angioplasty to treat a mammal having one or more (e.g., two, three, four, or more) stenotic blood vessels. In some cases, amnion coated balloons provided herein can be used in balloon angioplasty to reduce the symptoms associated with a stenotic blood vessel, widen a stenotic blood vessel, reduce or remove an obstruction from a stenotic blood vessel, and/or increase blood flow in a stenotic blood vessel. In some cases, a balloon catheter having an amnion coated balloon can be used to deliver an amnion tissue preparation to a stenotic blood vessel (e.g., to promote healing in the stenotic blood vessel).

[0011] An amnion coated balloon provided herein is coated with an amnion tissue preparation (e.g., human amnion tissue preparation). In some cases, an amnion coated balloon can be completely coated with an amnion tissue preparation. In some cases, an amnion coated balloon can be partially coated with an amnion tissue preparation. The term "amnion tissue preparation" as used herein refers to a preparation of amnion tissue or amnion material. In some cases, an amnion tissue preparation can be a liquid preparation (e.g., solution or suspension) that is prepared from a dried amnion tissue preparation. The term "dried amnion tissue preparation" as used herein refers to a preparation of amnion tissue or amnion material that is dried to have a water content that is less than about 8 percent (e.g., less than about 7 percent, less than about 6 percent, less than about 5 percent, less than about 4 percent, less than about 3 percent, less than about 2 percent, or less than about 1 percent). In some cases, a dried amnion tissue preparation can have a water content that is between about 0.1 percent and about 8 percent (e.g., between about 0.5 percent and about 8 percent, between about 1 percent and about 8 percent, between about 0.1 percent and about 5 percent, between about

0.1 percent and about 4 percent, between about 0.1 percent and about 3 percent, between about 0.5 percent and about 5 percent, or between about 1 percent and about 4 percent).

[0012] An amnion tissue preparation can be dried using any appropriate technique such as micronization, vacuum drying, spray drying, freeze drying, or combinations thereof. In some cases, an amnion tissue preparation can be dried as described elsewhere (e.g., U.S. Patent No. 5,656,498). A dried amnion tissue preparation can have any appropriate particle size. For example, a dried amnion tissue preparation can have a particle size ranging from about 0.1 μm to about 25 μm (e.g., from about 0.5 μm to about 25 μm , from about 0.75 μm to about 25 μm , from about 1 μm to about 25 μm , from about 0.1 μm to about 15 μm , from about 0.1 μm to about 10 μm , from about 0.1 μm to about 7.5 μm , from about 0.1 μm to about 5 μm , from about 0.75 μm to about 7.5 μm , or from about 1 μm to about 5 μm).

[0013] An amnion tissue preparation or a dried amnion tissue preparation can contain viable cells, non-viable cells, or a combination thereof. For example, an amnion tissue preparation or a dried amnion tissue preparation can be a preparation of amnion tissue or amnion material having viable cells. In some cases, an amnion tissue preparation can be a solution or suspension of amnion tissue or amnion material having viable cells.

[0014] In some cases, an amnion tissue preparation or a dried amnion tissue preparation can be a preparation of amnion tissue or amnion material where all the cells were removed, killed, or lysed such that the amnion tissue preparation or the dried amnion tissue preparation lacks viable cells. In some cases, an amnion tissue preparation or a dried amnion tissue preparation can be a preparation of amnion tissue or amnion material that was exposed to one or more physical and/or chemical treatments that killed, fixed, or lysed the cells of the amnion tissue or amnion material such that the amnion tissue preparation or the dried amnion tissue preparation lacks viable cells. For example, temperature (e.g., rapid freezing or rapid freezing-thawing), force and pressure, and/or electrical disruption can be used to kill or lyse cells within amnion tissue or amnion material to produce an amnion tissue preparation or a dried amnion tissue preparation that lacks viable cells.

[0015] In some cases, amnion tissue or amnion material can be obtained and then treated in a manner designed to lyse all the cells within the amnion tissue or amnion material. In these cases, the resulting material (e.g., matrix material and cellular remnants from lysed cells) can be used as an amnion tissue preparation that lacks viable cells or dried to form a dried amnion tissue preparation that lacks viable cells.

[0016] In some cases, an amnion tissue preparation or a dried amnion tissue preparation can be prepared from human amnion tissue. For example, human amnion tissue can be harvested, processed to maintain cell viability with or without removing blood, and used as an amnion tissue preparation or dried to form a dried amnion tissue preparation.

[0017] In some cases, human amnion tissue can be processed to remove blood prior to being used as an amnion tissue preparation or prior to being dried to form a dried amnion tissue

preparation. In some cases, human amnion tissue can be processed without removing cells or blood prior to forming an amnion tissue preparation or a dried amnion tissue preparation.

[0018] An example of an amnion tissue preparation includes, without limitation, a human amnion tissue preparation that includes viable cells. In some cases, an amnion tissue preparation can be obtained from MiMedX[®] or a tissue bank (e.g., a human tissue bank).

[0019] In some cases, an amnion tissue preparation also can include one or more therapeutic agents (e.g., a therapeutic agent that can be used to treat calcium deposits), one or more immunosuppressant agents (e.g., corticosteroids such as glucocorticoids), one or more anti-inflammatory agents (e.g., non-steroidal anti-inflammatory drugs, dexamethasone or other type of glucocorticoid steroids), one or more growth factors (e.g., vascular endothelial growth factor (VEGF), epithelial growth factor (EGF), fibroblast growth factor-2 (FGF2), or stem cell factor (SCF)), and/or one or more antimicrobial agents (e.g., antibiotics such as kanamycin, neomycin, streptomycin, or gentamicin, or an antifungal agent).

[0020] In some cases, an amnion coated balloon provided herein can be coated with an amnion tissue preparation (e.g., human amnion tissue preparation) as the sole active ingredient.

[0021] In some cases, an amnion coated balloon provided herein can be coated with an amnion tissue preparation (e.g., human amnion tissue preparation) and one or more therapeutic agents selected from the group consisting of a corticosteroid, a glucocorticoid, a non-steroidal anti-inflammatory drug, dexamethasone, an antibiotic, an antifungal agent, an antiproliferative drug, or an immunosuppressive drug.

[0022] An amnion tissue preparation can be coated onto a balloon to produce an amnion coated balloon provided herein using any appropriate method. For example, in cases where an amnion tissue preparation is a liquid preparation (e.g., solution or suspension), a balloon can be dipped into or sprayed with the liquid preparation.

[0023] In some cases, amnion coated balloons provided herein can be used to treat a mammal (e.g., a human) having one or more stenotic blood vessels. For example, an amnion coated balloon can be used for treating a mammal (e.g., a human) having one or more stenotic blood vessels. In some cases, an amnion coated balloon can be used in balloon angioplasty to reduce the symptoms associated with a stenotic blood vessel, widen a stenotic blood vessel, reduce or remove an obstruction from a stenotic blood vessel, and/or increase blood flow in a stenotic blood vessel. In some cases, an amnion coated balloon can be used in balloon angioplasty to deliver an amnion tissue preparation (e.g., an effective amount of an amnion tissue preparation described herein) to a stenotic blood vessel (e.g., to promote healing in the stenotic blood vessel). Effective amounts of amnion tissue preparations described herein can be determined by a physician, taking into account various factors such as overall health status, body weight, sex, diet, time and route of administration, other medications, and any other relevant clinical factors. As used herein, an "effective amount" or

"therapeutically effective amount" of a composition provided herein is the amount that is sufficient to provide a beneficial effect to the subject to which the composition or preparations are delivered. The effective amount can be the amount effective to achieve an improvement or elimination of one or more symptoms (e.g., symptoms associated with a stenotic blood vessel), widen a stenotic blood vessel, reduce or remove an obstruction from a stenotic blood vessel, and/or increase blood flow in a stenotic blood vessel.

[0024] In some cases, amnion coated balloons provided herein can be used in balloon angioplasty (e.g., coronary angioplasty, peripheral angioplasty, carotid angioplasty, venous angioplasty, femoral angioplasty, popliteal angioplasty, and tibial angioplasty). In some cases, amnion coated balloons provided herein can be used in stent-less balloon angioplasty (e.g., a balloon angioplasty in which a stent is not implanted). In some cases, amnion coated balloons provided herein can be used in balloon angioplasty that includes implanting stent. In cases where angioplasty with an amnion coated balloon includes implanting a stent, the stent can be implanted during the angioplasty procedure. For example, the stent can be provided on the same balloon catheter having an amnion coated balloon provided herein. In cases where angioplasty with an amnion coated balloon includes implanting a stent, the stent can be implanted after the angioplasty procedure. For example, the stent can be provided independently of an amnion coated balloon provided herein. In cases where angioplasty with an amnion coated balloon includes implanting a stent, the stent can be any appropriate stent. A stent can be made of any appropriate material (e.g., metals (such as nickel (e.g., nickel titanium), iron, magnesium, zinc, and/or alloys thereof) and/or polymers (such as poly(L-lactide) (PLLA); poly-D, L-lactide (PDLLA); tyrosine poly carbonate, and/or poly(anhydride-esters)-salicylates)). Examples of types of stents include, without limitation, bare stents, drug-eluting stents, bioabsorbable stents, coated stents, and combinations thereof. In cases where angioplasty with an amnion coated balloon includes implanting a drug-eluting stent, the stent can elute any appropriate drug (e.g., immunosuppressive drugs (such as rapamycin and everolimus) and antiproliferative drugs (such as paclitaxel)). In cases where angioplasty with an amnion coated balloon includes implanting a coated stent, the stent can be coated (e.g., completely coated or partially coated) with an amnion preparation described herein.

[0025] Methods of using amnion coated balloons provided herein can include inserting an amnion coated balloon provided herein into a stenotic blood vessel and inflating the amnion coated balloon. For example, an amnion coated balloon provided herein can be inflated to a pressure of from about 4 atm to about 35 atm (e.g., from about 4 atm to about 30 atm, from about 4 atm to about 25 atm, from about 4 atm to about 20 atm, from about 4 atm to about 15 atm, from about 4 atm to about 10 atm, from about 6 atm to about 35 atm, from about 8 atm to about 35 atm, from about 10 atm to about 35 atm, from about 15 atm to about 35 atm, from about 20 atm to about 35 atm, from about 5 atm to about 30 atm, from about 6 atm to about 25 atm, from about 8 atm to about 23 atm, or from about 10 atm to about 20 atm), wherein 1 bar = 1.01325 atm.

[0026] In some cases, an amnion coated balloon can be inflated at low pressure (e.g., lower pressure than is typically used with balloon angioplasty that includes implanting stent). For

example, an amnion coated balloon provided herein inflated at low pressure can be inflated to a pressure of less than about 10 atm (e.g., less than about 9 atm, less than about 8 atm, less than about 7 atm, less than about 6 atm, or less than about 5 atm). Methods of using amnion coated balloons provided herein also can include deflating and removing the amnion coated balloons.

[0027] An amnion coated balloon provided herein can be inserted into the stenotic blood vessel using any appropriate technique. For example, an amnion coated balloon can be inserted through a percutaneous puncture in the arm or groin. In some cases, methods of using amnion coated balloons provided herein can include using a guidewire to guide an amnion coated balloon to a stenotic blood vessel. In some cases, methods of using amnion coated balloons provided herein can include using optical imaging (e.g., X-ray, fluoroscopic guidance, and/or radiopaque contrast dye) to guide an amnion coated balloon to a stenotic blood vessel.

[0028] In some cases, a mammal can be identified as having one or more stenotic blood vessels using any appropriate technique. Examples of techniques that can be used to identify a mammal as having one or more stenotic blood vessels include, without limitation, angiography (e.g., coronary angiography), microphotography, stress testing, ultrasound (e.g., Doppler ultrasound and intravascular ultrasound), coronary calcium scoring (e.g., by CT), carotid measurement (e.g., carotid intimal media thickness (IMT) measurement by ultrasound), physiologic measurement (e.g., lipoprotein subclass analysis, HbA1c, hs-CRP, and homocysteine), and/or nuclear imaging techniques (e.g., PET and SPECT).

[0029] A stenotic blood vessel can be in any appropriate location of the body. Examples of body locations that can have stenotic blood vessels include, without limitation, the legs, the neck, the heart, the kidneys, the aorta, the chest, and the abdomen.

[0030] When a stenotic blood vessel is an obstructed blood vessel, the obstruction can be any appropriate type of obstruction. Examples of obstructions that can narrow a blood vessel include, without limitation, calcification (e.g., calcium deposits that accumulate and/or crystallize in soft tissues such as arteries), and plaques (e.g., atherosclerotic plaques such as atheromatous plaques). In some cases, plaques can include calcification (e.g., a calcified plaque). For example, a human having stenosis can have calcification in a carotid artery.

[0031] Any type of mammal having one or more stenotic blood vessels can be treated as described herein. Examples of mammals that can be treated with an amnion coated balloon provided herein include, without limitation, humans, non-human primates (e.g., monkeys), dogs, cats, horses, cows, pigs, sheep, rabbits, mice, and rats. For example, humans having one or more stenotic blood vessels can be treated with an amnion coated balloon provided herein.

[0032] When treating a mammal (e.g., a human) having one or more stenotic blood vessels, any appropriate blood vessel (e.g., an artery or a vein) can be narrowed and/or obstructed. In

cases where a stenotic blood vessel is an artery, the artery can be a peripheral artery (e.g., a leg artery such as a popliteal artery, a tibial artery (e.g., a posterior or anterior tibial artery), a peroneal artery, a plantar artery (e.g., a lateral, medial, or deep plantar artery), or a dorsalis pedis artery), a coronary artery (e.g., a carotid artery, a distal left main coronary artery (LMCA), a left circumflex artery (LCX), or a left anterior descending (LAD) artery), or a renal artery. In cases where a stenotic blood vessel is a vein, the vein can be a subclavian vein. For example, a human having a stenotic carotid artery can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure).

[0033] In some cases, a mammal (e.g., a human) having one or more stenotic blood vessels can have a condition associated with one or more stenotic blood vessels. The condition associated with one or more stenotic blood vessels can be any appropriate condition. Examples of conditions that can include a stenotic (e.g., narrowed and/or obstructed) blood vessel include, without limitation, heart disease, cardiovascular disease, arteriosclerosis (e.g., atherosclerosis), peripheral artery disease (PAD; e.g., PAD associated with diabetes), renal vascular hypertension, carotid artery disease, and coronary artery disease (CAD). For example, a human having atherosclerosis can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure). For example, a human having CAD can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure).

[0034] In some cases, the methods and materials provided herein can be used to reduce the symptoms associated with one or more stenotic blood vessels. In some cases, a balloon catheter having an amnion coated balloon can be used to reduce one or more (e.g., two, three, four, or more) symptoms associated with stenotic blood vessel. It will be appreciated that symptoms associated with stenosis will differ according to which blood vessel is stenotic and according to which organ(s) the stenotic blood vessel supplies blood. For example, symptoms associated with a stenotic coronary artery can include, without limitation, cardiovascular disorders (e.g., stroke or heart attack), chest pain (e.g., angina), shortness of breath, sweating, nausea, dizziness or light-headedness, palpitations, and arrhythmias. For example, symptoms associated with a stenotic carotid artery can include, without limitation, weakness, confusion, difficulty speaking, dizziness, difficulty walking, difficulty standing, blurred vision, numbness (e.g., of the face, arms, and/or legs), severe headache, and losing consciousness. For example, symptoms associated with a stenotic peripheral artery can include, without limitation, claudication, cramps, tingling, pain, numbness, wounds, reduced body temperature, and poor nail growth in the arms and/or legs. For example, symptoms associated with a stenotic renal artery can include, without limitation, decreased kidney blood flow, and chronic kidney disease. In some cases, a balloon catheter having an amnion coated balloon can be used to eliminate one or more symptoms associated with a stenotic blood vessel. For example, a human having one or more stenotic blood vessels can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure) to reduce or eliminate claudication, cramps, tingling, pain, numbness, wounds, reduced body

temperature, and/or poor nail growth in the hands and/or feet. For example, a human having one or more stenotic blood vessels can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure) to reduce or eliminate chest pain.

[0035] In some cases, the methods and materials provided herein can be used to widen (e.g., re-open) a stenotic blood vessel. In some cases, an amnion coated balloon (e.g., a balloon catheter having an amnion coated balloon) described herein can be inflated within a stenotic blood vessel to widen the blood vessel. For example, a human having one or more stenotic blood vessels can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure) to widen a stenotic blood vessel. In some cases, the methods provided herein can include widening a stenotic blood vessel to be from about 20% to about 100% open (e.g., about 25% to about 95%, about 30% to about 90%, about 50% to about 80%, about 60% to about 75%, about 20% to about 80%, about 20% to about 50%, about 25% to about 50%, about 30% to about 100%, about 40% to about 100%, about 50% to about 100%, or about 75% to about 100% open). For example, an amnion coated balloon provided herein can be inflated within a stenotic blood vessel to completely (e.g., 100%) open the blood vessel. In some cases, the methods provided herein (e.g., following the angioplasty) can have less than about 50% (e.g., less than about 45%, less than about 40%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, or less than about 5%) recoil.

[0036] In some cases, the methods and materials provided herein can be used to reduce or remove an obstruction from a stenotic blood vessel. In some cases, an amnion coated balloon (e.g., a balloon catheter having an amnion coated balloon) described herein can be inflated within a stenotic blood vessel to reduce or remove an obstruction from the blood vessel. For example, a human having one or more stenotic blood vessels can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure) to reduce or remove an obstruction from a stenotic blood vessel. In some cases, the methods provided herein can include reducing the size of an obstruction in a blood vessel by from about 20% volume to about 100% volume (e.g., about 25% to about 95%, about 30% to about 90%, about 50% to about 80%, about 60% to about 75%, about 20% to about 80%, about 20% to about 50%, about 25% to about 50%, about 30% to about 100%, about 40% to about 100%, about 50% to about 100%, or about 75% to about 100% volume). For example, an amnion coated balloon provided herein can be inflated within a stenotic blood vessel to completely remove (e.g., reduce by 100%) an obstruction in a blood vessel.

[0037] In some cases, the methods and materials provided herein can be used to increase blood flow in a blood vessel. In some cases, an amnion coated balloon (e.g., a balloon catheter having an amnion coated balloon) described herein can be inflated within a stenotic blood vessel to increase blood flow in the blood vessel. For example, a human having one or more stenotic blood vessels can be treated using an amnion coated balloon provided herein in an angioplasty procedure described herein (e.g., a stent-less angioplasty procedure) to increase blood flow in a blood vessel. In some cases, the methods provided herein can include

increasing blood flow in a blood vessel by from about 20% to about 100% (e.g., about 25% to about 95%, about 30% to about 90%, about 50% to about 80%, about 60% to about 75%, about 20% to about 80%, about 20% to about 50%, about 25% to about 50%, about 30% to about 100%, about 40% to about 100%, about 50% to about 100%, or about 75% to about 100%). In some cases, the methods provided herein can be used to increase blood flow in an artery to between about 50 cm/s and about 250 cm/s (e.g., between about 55 cm/s and about 250 cm/s, between about 60 cm/s and about 250 cm/s, between about 75 cm/s and about 250 cm/s, between about 100 cm/s and about 250 cm/s, between about 50 cm/s and about 225 cm/s, between about 50 cm/s and about 200 cm/s, between about 50 cm/s and about 175 cm/s, between about 50 cm/s and about 150 cm/s, between about 65 cm/s and about 225 cm/s, between about 75 cm/s and about 200 cm/s, or between about 100 cm/s and about 150 cm/s). For example, an amnion coated balloon provided herein can be inflated within a stenotic artery to restore a normal blood flow.

[0038] In some cases, the methods and materials provided herein can be used to deliver an amnion tissue preparation to a stenotic blood vessel. In some cases, an amnion coated balloon (e.g., a balloon catheter having an amnion coated balloon) described herein can be inflated within a stenotic blood vessel to reduce or remove an obstruction from the blood vessel. For example, an amnion coated balloon (e.g., a balloon catheter having an amnion coated balloon) described herein can be inflated within a stenotic blood vessel to deliver an amnion tissue preparation to a stenotic blood vessel. In some cases, the methods provided herein can include delivering, to mammal (e.g., a human), an amnion tissue preparation made with from about 0.01 mg to about 10 g (e.g., from about 0.01 mg to about 10 g, from about 0.1 mg to about 10 g, from about 1 mg to about 10 g, from about 10 mg to about 10 g, from about 100 mg to about 10 g, from about 1 g to about 10 g, from about 0.01 mg to about 5 g, from about 0.01 mg to about 1 g, from about 0.01 mg to about 100 mg, from about 10 mg to about 5 g, from about 100 mg to about 1 g, or from about 1 g to about 5 g) of amnion tissue per kg body weight of the subject being treated.

[0039] Methods for using amnion coated balloons provided herein (e.g., for treating a mammal (e.g., a human) having one or more stenotic blood vessels can be used as a combination therapy with one or more additional agents/therapies used to treat heart disease (e.g., heart disease involving one or more stenotic blood vessels). For example, a combination therapy used to treat stenotic blood vessels can include angioplasty with an amnion coated balloon provided herein and administering to the mammal one or more agents for treating stenotic blood vessels such as beta blockers (e.g., metoprolol, atenolol, and bisoprolol), angiotensin II receptor blockers (e.g., losartan, valsartan, and olmesartan), statins (e.g., atorvastatin, lovastatin, simvastatin), and blood thinners (e.g., anticoagulants such as heparin or warfarin, and antiplatelet drugs such as aspirin). In cases where a stenotic blood vessel includes calcification (e.g., a calcified plaque), the methods provided herein can include one or more agents and/or therapies to treat calcification. Examples of agents that can be used to treat calcification include, without limitation, vitamin K (e.g., vitamin K2), and vitamin D (e.g., if vitamin K is normal). Examples of therapies that can be used to treat calcification include, without limitation, atherectomy (e.g., orbital, rotational, laser, or directional atherectomy). In

cases where amnion coated balloons described herein are used in combination with one or more agents/therapies for treating stenotic blood vessels, the one or more agents/therapies for treating stenotic blood vessels can be administered/performed at the same time or independently. For example, angioplasty with an amnion coated balloon provided herein can be performed first, and the one or more agents/therapies for treating stenotic blood vessels can be administered second, or vice versa.

[0040] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US7084061Z **[0001]**
- US62523366 **[0001]**
- US5656498A **[0012]**

Non-patent literature cited in the description

- Heart Disease and Stroke Statistics-2007 Update American Heart Association American Heart Association 2007 0000 **[0003]**
- Harvard Heart Letter Harvard Health Publications, "Angioplasty or bypass surgery, 2008, **[0003]**

Titel; Metoder og materialer til behandling af blodkar.

1. Et angioplastikballonkateter omfattende en angioplastikballon, hvor angioplastikballonen omfatter en belægning omfattende
som eneste aktive ingrediens, et amnionvævspræparat; eller
som eneste aktive ingredienser, et amnionvævspræparat;
og et eller flere terapeutiske midler valgt fra gruppen bestående af et kortikosteroid, et glukokortikoid, et ikke-steroidt antiinflammatorisk lægemiddel, dexamethason, et antibiotikum, et svampedræbende middel, et antiproliferativt lægemiddel eller et immunsuppressivt lægemiddel.
2. Angioplastikballonkateteret ifølge krav 1, hvor amnionvævspræparatet har levedygtige celler.
3. Angioplastikballonkateteret ifølge krav 1, hvor amnionvævspræparatet mangler levedygtige celler.
4. Angioplastikballonkateteret ifølge et hvilket som helst af kravene 1-3, hvor amnionvævspræparatet er et tørret amnionvævspræparat.
5. Angioplastikballonkateteret ifølge krav 4, hvor det tørrede amnionvævspræparat har et vandindhold, der er mindre end omkring 8 procent.
6. Angioplastikballonkateteret ifølge krav 4, hvor det tørrede amnionvævspræparat har en partikelstørrelse i området fra omkring 0,1 μm til omkring 25 μm .
7. Angioplastikballonkateteret ifølge krav 1, hvor angioplastikballonkateteret er et stentløst angioplastikballonkateter.