

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 May 2006 (04.05.2006)

PCT

(10) International Publication Number
WO 2006/047702 A2

(51) International Patent Classification:
A61K 31/405 (2006.01)

(21) International Application Number:
PCT/US2005/038834

(22) International Filing Date: 27 October 2005 (27.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/622,022 27 October 2004 (27.10.2004) US

(71) Applicant (for all designated States except US): **SCI-CLONE PHARMACEUTICALS, INC.** [US/US]; 901 Mariner's Island Boulevard, Suite 205, San Mateo, California 94404 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MOSSEL, Eric, C.** [US/US]; 2213 Waters Edge Lane, League City, Texas 77573 (US). **TUTHILL, Cynthia, W.** [US/US]; 645 Woodland Avenue, Menlo Park, California 94025 (US). **RUDOLPH, Alfred, R.** [US/US]; 14142 Liddicoat Drive, Los Altos Hills, California 94022 (US).

(74) Agents: **REPPER, George, R.** et al.; 1425 K Street, N.w., Suite 800, Washington, District Of Columbia 20005 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2006/047702 A2

(54) Title: TREATMENT OR PREVENTION OF HEMORRHAGIC VIRAL INFECTIONS WITH IMMUNOMODULATOR COMPOUNDS

(57) Abstract: An immunomodulatory compound is administered to a patient having, or at risk of a hemorrhagic viral infection.

TREATMENT OR PREVENTION OF HEMORRHAGIC VIRAL INFECTIONS WITH IMMUNOMODULATOR COMPOUNDS

BACKGROUND OF THE INVENTION

5 CROSS-REFERENCE TO RELATED APPLICATION

[001] This application claims the benefit of U.S. Provisional Application Serial No. 60/622,022, filed October 27, 2004, which is incorporated herein.

Field of the Invention

[002] The present invention relates to the field of treatment or
10 prevention of hemorrhagic viral infections.

Description of the Background Art

[003] Hemorrhagic fever viruses (HFVs) are viruses classified in several taxonomic families. HFVs cause a variety of disease syndromes with similar clinical characteristics, referred to as acute hemorrhagic fever
15 syndromes. The pathophysiologic hallmarks of HFV infection are microvascular damage and increased vascular permeability. HFVs that are RNA viruses include Arenaviridae such as Lassa, as well as South American hemorrhagic fevers including Junin, Machupo, Guanarito, and Sabia viruses, which are the causative agents of Lassa fever and
20 Argentine, Bolivian, Venezuelan, and Brazilian hemorrhagic fevers, respectively, Whitewater Arroyo virus and Flexal virus; Filoviridae (Ebola and Marburg viruses); Bunyaviridae: Crimean-Congo hemorrhagic fever virus (CCHFV), Rift Valley fever virus, hemorrhagic fever with renal
25 syndrome-associated hantaviruses, including Hantaan virus, Seoul virus, Dobrava virus (also referred to as Dobrava-Belgrade virus), and Puumala virus, and hantavirus pulmonary syndrome-associated hantaviruses, including Bayou virus, Black Creek Canal virus, New York virus, Sin Nombre virus, Andes virus, Oran virus, Juquitiba virus, Laguna Negra

virus, and Lechiguanas virus; and Flaviviridae (dengue, dengue fever, dengue hemorrhagic fever, dengue shock syndrome, Kyasanur Forest disease, Omsk hemorrhagic fever, yellow fever).

[004] Under natural conditions, humans are infected through the bite of an infected arthropod or through contact with infected animal reservoirs. Hemorrhagic fever viruses are highly infectious by aerosol; are associated with high morbidity and, in some cases, high mortality; and are thought to pose a serious risk as biologic weapons.

[005] The exact pathogenesis for HFVs varies according to the etiologic agent. The major target organ is the vascular endothelium. Immunologic and inflammatory mediators are thought to play an important role in the pathogenesis of HFVs. All HFVs can produce thrombocytopenia, and some also cause platelet dysfunction. Infection with Ebola and Marburg viruses, Rift Valley fever virus, and yellow fever virus causes destruction of infected cells. Disseminated intravascular coagulation (DIC) is characteristic of infection with Filoviridae. Ebola and Marburg viruses may cause a hemorrhagic diathesis and tissue necrosis through direct damage to vascular endothelial cells and platelets with impairment of the microcirculation, as well as cytopathic effects on parenchymal cells, with release of immunologic and inflammatory mediators. Arenaviridae, on the other hand, appear to mediate hemorrhage via the stimulation of inflammatory mediators by macro-phages, thrombocytopenia, and the inhibition of platelet aggregation.

[006] The incubation period of HFVs ranges from 2 to 21 days. The clinical presentations of these diseases are nonspecific and variable, making diagnosis difficult. It is noteworthy that not all patients will develop hemorrhagic manifestations. Even a significant proportion of patients with Ebola virus infections may not demonstrate clinical signs of hemorrhage.

[007] Initial symptoms of the acute HFV syndrome may include fever, headache, myalgia, rash, nausea, vomiting, diarrhea, abdominal pain,

arthralgias, myalgias, and malaise. Illness caused by Ebola, Marburg, Rift Valley fever virus, yellow fever virus, Omsk hemorrhagic fever virus, and Kyasanur Forest disease virus are characterized by an abrupt onset, whereas Lassa fever and the diseases caused by the Machupo, Junin, 5 Guarinito, and Sabia viruses have a more insidious onset. Initial signs may include fever, tachypnea, relative bradycardia, hypotension (which may progress to circulatory shock), conjunctival injection, pharyngitis, and lymphadenopathy. Encephalitis may occur, with delirium, seizures, cerebellar signs, and coma. Most HFVs cause cutaneous flushing or a 10 macular skin rash, although the rash may be difficult to appreciate in dark-skinned persons and varies according to the causative virus. Hemorrhagic symptoms, when they occur, develop later in the course of illness and include petechiae, purpura, bleeding into mucous membranes and conjunctiva, hematuria, hematemesis, and melena. Hepatic involvement is 15 common, and renal involvement is proportional to cardiovascular compromise.


[008] Laboratory abnormalities include leukopenia (except in some cases of Lassa fever), anemia or hemoconcentration, and elevated liver enzymes; DIC with associated coagulation abnormalities and 20 thrombocytopenia are common. Mortality ranges from less than 1% for Rift Valley fever to 70% to 90% for Ebola and Marburg virus infections

[009] The nonspecific and variable clinical presentation of the HFVs presents a considerable diagnostic challenge. Clinical diagnostic criteria based on WHO surveillance standards for acute hemorrhagic fever 25 syndrome include temperature greater than 101 F (38.3 C) of less than 3 weeks' duration; severe illness and no predisposing factors for hemorrhagic manifestations; and at least two of the following hemorrhagic symptoms: hemorrhagic or purple rash, epistaxis, hematemesis, hematuria, hemoptysis, blood in stools, or other hemorrhagic symptom 30 with no established alternative diagnosis. Laboratory techniques for the diagnosis of HFVs include antigen detection, IgM antibody detection,

asymptomatic.

[0017] Immunomodulator compounds in accordance with the present invention, comprise immunomodulators of Formula A:

5



10

(A)

[0018] In Formula A, n is 1 or 2, R is hydrogen, acyl, alkyl or a peptide fragment, and X is an aromatic or heterocyclic amino acid or a derivative thereof. Preferably, X is L-tryptophan or D-tryptophan.

[0019] Appropriate derivatives of the aromatic or heterocyclic amino acids for "X" are: amides, mono- or di-(C₁-C₆) alkyl substituted amides, arylamides, and (C₁-C₆) alkyl or aryl esters. Appropriate acyl or alkyl moieties for "R" are: branched or unbranched alkyl groups of 1 to about 6 carbons, acyl groups from 2 to about 10 carbon atoms, and blocking groups such as carbobenzyloxy and t-butyloxycarbonyl. Preferably the carbon of the CH group shown in Formula A has a stereoconfiguration, when n is 2, that is different from the stereoconfiguration of X.

[0020] Preferred embodiments utilize compounds such as γ -D-glutamyl-L-tryptophan, γ -L-glutamyl-L-tryptophan, γ -L-glutamyl-N_m-formyl-L-tryptophan, N-methyl- γ -L-glutamyl-L-tryptophan, N-acetyl- γ -L-glutamyl-L-tryptophan, γ -L-glutamyl-D-tryptophan, β -L-aspartyl-L-tryptophan, and β -D-aspartyl-L-tryptophan. Particularly preferred embodiments utilize γ -D-glutamyl-L-tryptophan, sometimes referred to as SCV-07. These compounds, methods for preparing these compounds, pharmaceutically acceptable salts of these compounds and pharmaceutical formulations

thereof are disclosed in U.S. Patent No. 5,916,878, incorporated herein by reference.

[0021] The invention is applicable to prevention and/or treatment of hemorrhagic viral infections, including but not limited to Arenaviridae such as Lassa, as well as South American hemorrhagic fevers including 5 Junin, Machupo, Guanarito, and Sabia viruses, which are the causative agents of Lassa fever and Argentine, Bolivian, Venezuelan, and Brazilian hemorrhagic fevers, respectively, Whitewater Arroyo virus and Flexal virus; Filoviridae (Ebola and Marburg viruses); Bunyaviridae: Crimean- 10 Congo hemorrhagic fever virus (CCHFV), Rift Valley fever virus, hemorrhagic fever with renal syndrome-associated hantaviruses, including Hantaan virus, Seoul virus, Dobrava virus (also referred to as Dobrava-Belgrade virus), and Puumala virus, and hantavirus pulmonary syndrome-associated hantaviruses, including Bayou virus, Black Creek Canal virus, 15 New York virus, Sin Nombre virus, Andes virus, Oran virus, Juquitiba virus, Laguna Negra virus, and Lechiguanas virus; and Flaviviridae (dengue, dengue fever, dengue hemorrhagic fever, dengue shock syndrome, Kyasanur Forest disease, Omsk hemorrhagic fever, yellow fever).

20 [0022] Pichinde virus is used as an established model for Lassa fever.

[0023] The Formula A compounds may be administered as dosages in the range of about 0.001-10 mg. Dosages may be administered one or more times per week, preferably on a daily basis, with dosages administered one or more times per day. The dosages may be 25 administered by intramuscular injection, although other forms of injection and infusion may be utilized, and other forms of administration such as oral or nasal inhalation or oral ingestion may be employed.

[0024] In preferred embodiments, the compounds of Formula A are administered at a dosage within a range of about 0.01-10 mg, more 30 preferably at a dosage of about 0.1-1 mg.

[0025] Dosages may also be measured in micrograms per kilogram subject body weight, with dosages in the range of about 0.01-100 micrograms per kilogram, more preferably within the range of about 0.1-10 micrograms per kilogram, and most preferably at about 1 microgram per kilogram.

[0026] Included are biologically active analogs having substituted, deleted, elongated, replaced, or otherwise modified portions which possess bioactivity substantially similar to that of SCV-07, e.g., an SCV-07 derived peptide having sufficient homology with SVC-07 such that it functions in substantially the same way with substantially the same activity as SCV-07.

[0027] Administration can be by any suitable method, including orally, by injection, periodic infusion, continuous infusion, and the like.

[0028] In some embodiments, the Formula A compound is present in a pharmaceutically acceptable liquid carrier, such as water for injection, saline in physiological concentrations, or similar.

[0029] Effective amounts of Formula A compound can be determined by routine dose-titration experiments.

[0030] The Formula A compound also can be administered with other immune stimulators or antiviral agents.

[0031] Arenaviruses pose an ongoing public health concern both from naturally acquired infections and the potential for use as a biothreat agent. The family Arenaviridae is comprised of enveloped viruses with an ambisense, bi-segmented genome. Its 22 members are divided into 2 serocomplexes. The lymphocytic choriomeningitis-Lassa complex, contains the five known 'Old World' viruses while the Tacaribe complex is made up of 17 'New World' viruses of North and South America. Each virus is associated with a single rodent species or with a couple of closely related rodent species reservoir (the possible exception is Tacaribe, which has only been isolated from two species of bats). This close association with the rodent host limits the geographic range of the virus to that of the

reservoir. LCMV, being associated with the ubiquitous *Mus musculus*, is the only arenavirus found worldwide.

[0032] Asymptomatically infected rodents come into contact with humans, who become infected through the inhalation of virus in aerosolized excreta. Most arenaviruses have not been associated with human infection and not all arenaviruses known to infect humans cause disease. However, several arenaviruses do cause disease in humans that ranges from mild to very severe. The most important are LCMV, which causes a rarely fatal CNS syndrome, Lassa virus, the etiological agent of Lassa fever, and Junin, Machupo, Guanarito, and Sabia, the causative agents of the South American hemorrhagic fevers.

[0033] Lassa fever, like the Lassa virus reservoir, *Mastomys* sp., is endemic to West Africa. The disease is characterized by fever, weakness, malaise, headache and sore throat, with an onset 7-18 days after infection.

Other common symptoms include joint and back pain, cough, vomiting, and diarrhea. Unlike the hemorrhagic fevers, there is no associated skin rash, ecchymoses, or petechiae. Case fatality among hospitalized patients is 15-20%, but among all Lassa infections may be as low as 2-3%.

Aerosolized person-to-person transmission has not been shown to occur.

Contact with the secretions of an infected person is a significant risk factor, however.

[0034] Junin, Machupo, Guanarito, and Sabia viruses are the etiological agents of the South American hemorrhagic fevers: Argentine, Bolivian, Venezuelan, and Brazilian hemorrhagic fever, respectively. Though

distinct, all have similar presentations that include fever, malaise, and myalgia beginning 1-2 weeks after infection. As disease worsens, additional symptoms develop: gastrointestinal distress, dizziness, headache, photophobia, retroorbital pain, tachycardia, petechiae, and conjunctival injection. Machupo infection appears to have the greatest

propensity for person-to-person spread, with nosocomial and intrafamilial spread documented. Unlike the other hemorrhagic fever arenaviruses, but

similar to Lassa fever infection, deafness has been observed following Guanarito virus infection. Case fatality rate for the South American hemorrhagic fever viruses is around 20%.

[0035] Because of their high mortality, aerosol infectivity, and potential to
5 cause a public health crisis, Lassa, Junin, Machupo, Guanarito, and Sabia have been designated category A pathogens by the Center for Disease Control (CDC). Work with these agents must be done using BSL-4 precautions.

[0036] Pichinde virus (PIC) is a nonpathogenic Tacaribe complex virus.
10 PIC virulence in guinea pigs was dramatically increased through serial spleen passage in strain 13 guinea pigs. Infection of guinea pigs with the adapted virus produces a disease quite similar to Lassa fever in monkeys and humans. Features include fever, thrombocytopenia, platelet
15 dysfunction, vascular leakage, respiratory distress, viral replication in most extraneural tissues, and minimal histologic changes in infected
tissues. Because Pichinde is not pathogenic to man, this model has been extensively used as a low-containment proxy for the study of both Lassa
fever and arenavirus hemorrhagic fever.

[0037] The nucleoside analogue Ribavirin is the primary antiviral
20 therapeutic for Lassa fever and is a promising therapy for arenavirus hemorrhagic fever. However, while ribavirin therapy may prevent or relieve acute hemorrhagic fever, its use in arenavirus infection has
sometimes led to late neurologic disease. Several immunologic treatment modalities have also been tried in animal models of arenavirus infection
25 and disease without success. In vitro, arenaviruses are 10^{-3} - 10^{-4} -fold less sensitive to IFN β than is vesicular stomatitis virus. This insensitivity is supported by studies in Machupo-infected Rhesus monkeys with the IFN-inducing poly (ICLC). Overall mortality and time to death were not
improved with therapy and in some groups viremia was significantly
30 higher with treatment. Lucia et al. examined the use of IFN α , ribavirin, and the immunomodulatory drug C246,783 using the PIC-infected guinea

pig model. Among the effects of C246,783 are IFN induction and activation of NK cells and macrophages. No significant improvement in mortality was observed among animals treated with IFN α (up to 1.7×10^5 U) or with C246,783. An improvement in mortality was observed in this study with ribavirin administration if given daily for 28 days after infection; time to death was prolonged with 14 days of treatment. The relative insensitivity of arenaviruses to the type I IFNs probably contributed significantly to the virtual failure of these studies.

[0038] Despite these earlier failures of immunotherapy for the arenaviruses, T-cell stimulation is still an unexplored mode of treatment. A pilot study was therefore designed to examine the effect of the immunomodulatory drug SCV-07 on PIC-infected guinea pigs. SCV-07 is known to enhance the Th1 response to immunological stimuli.

SCV-07

[0039] SCV-07, γ -D-glutamyl-L-tryptophan, is a member of a class of immunomodulatory drugs that possess γ -glutamyl or β -aspartyl moieties, which was discovered by Russian scientists and is being examined for efficacy in several indications in the U.S. by SciClone Pharmaceuticals, Inc. SCV-07 possesses a number of immunomodulatory activities in vivo and in vitro. SCV-07 increases Con-A-induced thymocyte and lymphocyte proliferation, increases Con-A-induced interleukin-2 (IL-2) production and IL-2 receptor expression by spleen lymphocytes, and stimulates expression of Thy-1.2 on bone marrow cells. In vivo, SCV-07 has a strong immunostimulatory effect on 5-FU-immune-suppressed animals and in a model of immunization with sheep red blood cells.

[0040] The discrete mechanism of SCV-07 action has not been rigorously determined. However, based on the above data, it was suggested that SCV-07 acts via its influence on differentiation of pluripotent stem cells to thymocytes and/or differentiation of thymocytes to T-cells. Further, since SCV-07 administration results in increases of the Th1 cytokines IL-2 and

IFN γ , but not in Th2 cytokines, it was proposed that SCV-07 acts through a preferential activation of Th1 cells.

[0041] Increasing evidence suggests that CD4+ Th1 cells and associated cytokines, including IL-2 and IFN γ , are important in generating a

5 protective immune response to mycobacterial infection. In studies in a murine tuberculosis model, SCV-07 was shown to significantly decrease lung damage. IFN γ production by Con-A-induced spleen cells from SCV-07-treated animals was higher while IL-4 production was lower than in cells from untreated animals. Phagocytic activity by peritoneal

10 macrophages was markedly increased over untreated, infected controls. These pre-clinical data supported the commencement of clinical trials for the use of SCV-07. A phase II clinical trial performed in Russia found that SCV-07 administration increased the incidence of mycobacteria-negative sputum cultures and had a positive effect on cavity healing and symptoms
15 of tuberculosis without any drug-induced adverse effects. SCV-07's efficacy in human tuberculosis cases supports the hypothesis for its efficacy in other diseases in which Th1 immunity has been shown to be important, including arenaviral disease.

[0042] Based on the above, two pilot studies were undertaken to
20 examine the efficacy of SCV-07 in treating PIC-infected guinea pigs.

[0043] The invention is further illustrated by the following examples, which are not intended to be limiting.

Example 1:

[0044] 20 male Hartley guinea pigs (400-450g) were divided into groups
25 of five.

Group 0 received no SCV-07

Group 1 received 1 μ g/kg SCV-07 i.p. each of the five days prior to infection

Group 2 received 1 μ g/kg SCV-07 i.p. once on the same day as infection

30 Group 3 received 1 μ g/kg SCV-07 i.p. once on the day following infection

[0045] Infections: animals were inoculated i.p. with 100 pfu guinea pig-adapted PIC.

[0046] Body weight and temperature, and surrogate markers for disease progression were measured at least three times per week. Obviously ill
5 animals were assessed more often. Previous work showed that a 25% weight loss represented a terminally ill animal. However, the University of Texas Medical Branch IACUC required that a 20% weight loss would result in classification as terminal and sacrifice of the animal.

[0047] Group 0 animals are pooled from two separate experiments; the
10 one described here with treated groups 1 – 3, and the one described below with treated groups 4 – 6.

[0048] While there is a trend toward efficacy as shown by decreased mortality in the treated groups, Kaplan-Meier survival analysis by the log-rank statistic method supports the visual notion that none of the survival
15 curves of treated animals is significantly different from the survival curve of the control animals ($p = 0.66, 0.77,$ and 0.40 for groups 1, 2, and 3, respectively). Weight gain through 11 days post-infection (dpi) among treated animals was greatest in group 1 (Table 1), but for no group was weight gain different from controls. Neither was the temperature,
20 assessed 11 dpi, different between the treated and control animals (Table 2). Not evident from Tables 1 and 2 is the fact that the single control survivor came from the second control group. This animal dramatically skewed the control group temperature to the downside and weight gain to the upside. Therefore, the results from the initial study were more
25 suggestive of efficacy than they appear in the pooled data.

Table 1: Average percent weight gain by PIC-infected guinea pigs 10-11 dpi.

Group	Mean \pm Std Dev	<i>P</i> value*
0	1.79 \pm 13.50	
1	0.59 \pm 10.71	0.48
2	-3.75 \pm 12.48	0.92
3	-1.15 \pm 12.26	0.65
4	-0.35 \pm 20.07	0.64
5	6.73 \pm 19.02	0.21
6	5.62 \pm 16.78	0.23

*Student's *t*-test comparison of treated group to group 0.

Table 2: Average temperature ($^{\circ}$ C) of PIC-infected guinea pigs 10-11 dpi.

Group	Mean \pm Std Dev	<i>P</i> value*
0	40.35 \pm 0.63	
1	40.40 \pm 1.00	0.72
2	40.48 \pm 0.72	0.82
3	40.40 \pm 0.69	0.68
4	40.03 \pm 0.59	0.23
5	39.38 \pm 0.75	0.17
6	40.28 \pm 0.63	0.46

*Student's *t*-test comparison of treated group to group 0.

[0049] Though not significant, the first trial was suggestive enough that a second study was initiated.

Example 2:

5 [0050] 20 animals were again divided into four groups of five.

Group 0 received no SCV-07

Group 4 received 1 $\mu\text{g}/\text{kg}$ SCV-07 i.p. on each of the first five days after infection

10 Group 5 received 10 $\mu\text{g}/\text{kg}$ SCV-07 i.p. on each of the first five days after infection

Group 6 received 100 $\mu\text{g}/\text{kg}$ SCV-07 i.p. on each of the first five days after infection

[0051] Infections: animals were inoculated i.p. with 100 pfu guinea pig-adapted PIC.


15 [0052] Body weight and temperature, and surrogate markers for disease progression were measured as described above.

[0053] Mortality was decreased in groups 5 and 6, but only group 5 had a survival curve significantly different from group 0 ($p = 0.77, 0.03, \text{ and } 0.27$ for groups 4, 5, and 6, respectively). Weight gain through 11 dpi was
20 greater in treated groups 5 and 6 than among controls, but the difference was not significant (Table 1). Body temperature was lower in all three treated groups than in controls, but again, the difference was not significant (Table 2). Serum was collected from animals in both
25 experiments. Analysis of these samples is ongoing, but preliminary results suggest undetectable viremia until ~ 48 hours before acute illness and death.

[0054] These results suggest that SCV-07 administration was therapeutic in PIC-infected guinea pigs, as evidenced by a trend towards decreased mortality in several treatment groups, and with significantly decreased
30 mortality at a dose of 10 $\mu\text{g}/\text{kg}$ given i.p. for 5 days post infection.

CLAIMS

1. A method of treatment or prevention of a hemorrhagic viral infection in a subject comprising administering to said subject an effective amount of an immunomodulator compound of Formula A



(A)

- 15 wherein, n is 1 or 2, R is hydrogen, acyl, alkyl or a peptide fragment, and X is an aromatic or heterocyclic amino acid or a derivative thereof.

2. The method of claim 1, wherein X is L-tryptophan or D-tryptophan.

3. The method of claim 1 wherein said compound is SCV-07.

4. The method of claim 1 wherein said compound is administered at a dosage within a range of about 0.1-10 mg.

5. The method of claim 1 wherein said compound is administered at a dosage within a range of about 0.1-1 mg.

6. The method of claim 1 wherein said compound is administered at a dosage within a range of about 0.01-100 micrograms per kilogram subject body weight.

7. The method of claim 1 wherein said compound is administered at a dosage within a range of about 0.1-10 micrograms per kilogram subject body weight.

8. The method of claim 7 wherein said compound is SCV-07.

9. The method of claim 1 for treatment or prevention of infection by Arenaviridae including Lassa fever virus, Junin virus, Machupo

- virus, Guanarito virus, Sabia virus, Whitewater Arroyo virus or Flexal virus; Filoviridae including Ebola virus or Marburg virus; Bunyaviridae including Crimean-Congo hemorrhagic fever virus (CCHFV), Rift Valley fever virus, hemorrhagic fever with renal syndrome-associated hantaviruses including Hantaan virus, Seoul virus, Dobrava virus (also referred to as Dobrava-Belgrade virus), or Puumala virus, hantavirus pulmonary syndrome-associated hantaviruses including Bayou virus, Black Creek Canal virus, New York virus, Sin Nombre virus, Andes virus, Oran virus, Juquitiba virus, Laguna Negra virus, or Lechiguanas virus; or Flaviviridae including dengue virus, Kyasanur Forest disease virus, Omsk hemorrhagic fever virus or yellow fever virus.
- 5
- 10
10. The method of claim 1 for treatment or prevention of Lassa fever, South American hemorrhagic fevers including Argentine hemorrhagic fever, Bolivian hemorrhagic fever, Venezuelan hemorrhagic fever or Brazilian hemorrhagic fever, Whitewater Arroyo virus fever, Flexal virus fever, Ebola hemorrhagic fever, Marburg hemorrhagic fever, Crimean-Congo hemorrhagic fever, Rift Valley fever, hemorrhagic fevers with renal syndrome including Hantaan virus hemorrhagic fever, Seoul virus hemorrhagic fever, Dobrava virus hemorrhagic fever, Puumala virus hemorrhagic fever, hantavirus pulmonary syndrome-associated hemorrhagic fevers, including Bayou virus hemorrhagic fever, Black Creek Canal virus hemorrhagic fever, New York virus hemorrhagic fever, Sin Nombre virus hemorrhagic fever, Andes virus hemorrhagic fever, Oran virus hemorrhagic fever, Juquitiba virus hemorrhagic fever, Laguna Negra virus hemorrhagic fever, Lechiguanas virus hemorrhagic fever, dengue hemorrhagic fever, dengue shock syndrome, Kyasanur Forest disease, Omsk hemorrhagic fever or yellow fever.
- 15
- 20
- 25