




- (51) **International Patent Classification:**
C12Q 1/6806 (2018.01) C12N 15/10 (2006.01)
- (21) **International Application Number:** PCT/US2019/024107
- (22) **International Filing Date:** 26 March 2019 (26.03.2019)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/648,174 26 March 2018 (26.03.2018) US
- (71) **Applicant: QIAGEN SCIENCES, LLC [US/US];** 19300 Germantown Rd, Germantown, Maryland 20874 (US).
- (72) **Inventors; and**
(71) **Applicants: WANG, Yexun [CN/US];** 4442 Cross Country Drive, Ellicott City, Maryland 21042 (US). **PENG, Quan [CN/US];** 12010 Grey Squirrel St., Clarksburg, Maryland 20871 (US). **KIM, Daniel [US/US];** 524 W C St., Brunswick, Maryland 21716 (US).
- (74) **Agent: KIM, Judith;** Rubin and Rudman, 800 Connecticut Ave, NW, Suite 400, Washington, District of Columbia 20006 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

(54) **Title:** INTEGRATIVE DNA AND RNA LIBRARY PREPARATIONS AND USES THEREOF

FIG. 2



(57) **Abstract:** The invention relates to integrated DNA and RNA library preparations and methods of making and uses thereof, wherein the DNA molecules are labelled with a tag identifying the molecule as being a DNA molecule and the RNA molecules are labelled with a tag identifying the molecule as being a RNA molecule. The methods do not require physical separation of DNA and RNA. The methods output two separate libraries from DNA and RNA, respectively, which helps flexible manipulation on downstream sequencing platforms. The application also claims the DNA library and the cDNA library produced by the method and the DNA tag and the RNA tag used in the method.



UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

INTEGRATIVE DNA AND RNA LIBRARY PREPARATIONS AND USES THEREOF

BACKGROUND OF THE INVENTION

[0001] Paired DNA and RNA profiling enables researchers to gain more biological insights regarding the correlation between genotype and phenotype using samples from the same set of cell population.

[0002] To fully understand the complex biological systems, researchers are getting more and more interested in multi-omic information, a full view of genomic, transcriptomic and proteomic data and their interactions. As the first step, paired DNA and RNA profiling are made possible with the advancement of sequencing technology. Traditional approaches, however, usually required preparing DNA and RNA samples in parallel, which meant the data obtained were not necessarily from the same set of cell population. Thus, these separate workflows might yield less correlative DNA and RNA data due to the heterogeneity of the sample. In addition, they added extra time and effort because of doubled workload.

[0003] In order to circumvent the disadvantage of separate workflow, researchers tried to integrate them. DR-seq (Dey SS et al., *Nat. Biotechnol.* 33:285-289 (2015)) and G&T-seq (Macaulay IC et al., *Nat. Methods.* 12:519-22 (2015)) were among the first few attempts of integrative analysis of genomic DNA and mRNA from a single cell. However, these methods were designed specifically for single cell applications.

[0004] Another group developed an integrated DNA and RNA sequencing workflow, named Simul-seq. Reuter JA et al., *Nat. Methods* 13:953-958 (2016). It is a streamlined approach to profiling whole genome and transcriptome from the same set of cell population. Simul-seq is designed for whole genome and transcriptome sequencing.

[0005] There remains a need for improved, integrated DNA and RNA preparations amenable for sequencing analysis.

BRIEF SUMMARY OF THE INVENTION

[0006] Disclosed herein are methods for preparing DNA and cDNA libraries from a sample, comprising: ligating a DNA tag to an end of a DNA molecule in a sample, wherein the DNA tag comprises a unique molecular identifier (UMI) and a DNA identifier; and performing reverse transcription of a RNA molecule in the sample in the presence of a RNA tag, wherein the RNA tag comprises a RNA identifier, a UMI, and a poly(T).

[0007] In some embodiments, the reverse transcription is performed in the presence of a second RNA tag, wherein the second RNA tag comprises a RNA identifier, a UMI, and a template switching oligonucleotide (TSO).

[0008] In some embodiments, the methods further comprise amplifying the tagged DNA and tagged cDNA for enrichment with a set of gene specific primers. In some embodiments, the methods further comprise separating the amplified sample into first and second samples.

[0009] In some embodiments, the DNA and RNA molecules are obtained from a biological sample. In some embodiments, the DNA and RNA molecules are fragmented DNA and RNA from the biological sample.

[0010] In some embodiments, the DNA molecule contains polished ends for ligation. In some embodiments, the RNA molecule is polyadenylated.

[0011] In some embodiments, the methods do not require ribosomal depletion.

[0012] In further embodiments, the methods further comprise further amplifying the first sample with primers specific for the DNA tag. The amplification can generate a DNA library corresponding to the DNA molecules in the sample.

[0013] In further embodiments, the methods further comprise further amplifying the second sample with primers specific for the RNA tag. The amplification generates a cDNA library corresponding to the RNA molecules in a sample.

[0014] In some embodiments, the methods further comprise sequencing the DNA or cDNA library. The DNA library can be used for, but not limited to, DNA variant detection, copy number analysis, fusion gene detection, or structural variant detection. The cDNA library can be used for, but not limited to, RNA variant detection, gene expression analysis, or fusion gene detection. The libraries can be also used for paired DNA and RNA profiling.

[0015] Also disclosed herein are DNA libraries made by the methods disclosed herein. Further disclosed are cDNA libraries made by the methods disclosed herein.

[0016] Also disclosed herein are DNA tags comprising a unique molecular identifier (UMI) and a DNA identifier. In some embodiments, in the DNA tags, the UMI and the DNA identifier can be positioned in a 5' to 3' direction.

[0017] Also disclosed herein are RNA tags comprising a RNA identifier, a UMI, and a poly(T). In some embodiments, in the RNA tags, the RNA identifier, the UMI, and the poly(T) are positioned in a 5' to 3' direction. Also disclosed herein are RNA tags comprising a RNA identifier, a UMI, and a template switching oligonucleotide (TSO). In some

embodiments, in the RNA tags, the RNA identifier, the UMI, and the TSO are positioned in a 5' to 3' direction.

[0018] Disclosed herein are compositions comprising at least 2 of the above described tags. Also disclosed herein are compositions comprising the DNA tag and the 2 different RNA tags as described above.

[0019] Further disclosed herein are methods for preparing targeted DNA and cDNA libraries, comprising:

- (a) obtaining purified DNA and RNA from a biological sample;
- (b) fragmenting the DNA and RNA;
- (c) polishing the ends of the double stranded DNA fragments for ligation;
- (d) polishing the RNA fragments by polyadenylation;
- (e) ligating a DNA tag to a 3' end of the polished DNA fragments, wherein the DNA tag comprises in a 5' to 3' direction a unique molecular identifier (UMI) and a DNA identifier;
- (f) performing reverse transcription of the polished RNA fragments in the presence of a first RNA tag, wherein the first RNA tag comprises in a 5' to 3' direction a RNA identifier, a UMI, and a poly(T), and a second RNA tag, wherein the second RNA tag comprises in a 5' to 3' direction a RNA identifier, a UMI, and a template switching oligonucleotide (TSO);
- (g) amplifying the tagged DNA and tagged cDNA for enrichment with a set of gene specific primers;
- (h) separating the amplified sample into first and second samples;
- (i) amplifying the first sample with primers specific for the DNA tag; and
- (j) amplifying the second sample with primers specific for the RNA tag.

BRIEF DESCRIPTION OF THE DRAWINGS/FIGURES

[0020] FIG. 1. Exemplary DNA and RNA tag molecules.

[0021] FIG. 2. Exemplary process for generating DNA and cDNA libraries.

DETAILED DESCRIPTION OF THE INVENTION

[0022] Disclosed herein are innovative approaches for integrative DNA and cDNA library preparations for analysis, such as by next-generation sequencing (NGS) analysis,

without physical separation of DNA and RNA. These approaches integrate UMI (unique molecular index) technology and optionally, targeted enrichment technology, seamlessly into the workflow, which improve utilization of sequencing capacity and accuracy of the results. In addition, these methods output two separate DNA and cDNA libraries from DNA and RNA, respectively, which allow flexible manipulation on downstream sequencing platform. Compared to standalone DNA library and cDNA library methods, these approaches reduce sample consumption, simplify the experimental process, and can help researchers gain biological insights in genotype and phenotype correlations and molecular mechanisms of diseases.

[0023] Methods are described herein to prepare targeted DNA and cDNA libraries without the necessity of physical separation of genomic DNA (gDNA) and mRNA. The process involves three modules: (1) assign different DNA and RNA tag molecules to each individual DNA and RNA fragment, respectively, without separating them in the system; optionally, (2) amplify and enrich a subset of the tagged DNA and RNA fragments (target enrichment); and (3) differentially PCR amplify the tagged DNA and tagged cDNA in the (enriched) product to output two libraries corresponding to the original DNA and RNA, respectively.

[0024] The DNA and RNA tag molecules used in the first module are oligonucleotides comprising at least 1) an identifying sequence to distinguish a DNA library or RNA library, and 2) a UMI sequence for identifying each individual nucleic acid molecule.

[0025] The DNA and RNA tags are essential for the final separation of DNA and cDNA libraries in module 3, where they can serve as specific amplification primer sites for DNA and RNA. The UMI sequence helps improve accuracy for both DNA and RNA NGS analysis. Exemplary tag molecules are illustrated in FIG. 1.

[0026] Two types of RNA tag molecules can be used in order to sequence the single stranded RNA from both directions, and thus, two different mechanisms can be used to attach the RNA specific sequence. Only one type of DNA tag molecule is needed because the DNA tag molecule can be ligated to both ends of the double stranded DNA.

[0027] The targeted enrichment reaction (module 2) enables focused view on relevant regions of interest, and provides economic utilization of NGS sequencing capacity. It also mitigates the necessity for extra treatment of the sample associated with whole genome or transcriptome workflow, such as ribosomal RNA depletion. The enrichment is done in the same reaction for both DNA and RNA. Depending on the applications, the enrichment

primer pool can be the same if the target DNA and RNA regions are the same. If different regions are of interest for the DNA and RNA, users can simply mix the corresponding enrichment primer pools, and put them into the same reaction.

[0028] Module 3 enables separated output of DNA and cDNA libraries. The sequencing depth requirements for DNA and cDNA are usually quite different, and they vary depending on the applications. The output from the methods disclosed herein gives users flexibility so that sequencing capacity can be allocated individually according to specific needs. In addition, since the samples have already been partially amplified in module 2, the separation has negligible effect on sample loss.

[0029] FIG. 2 illustrates one exemplary, optimized way to utilize the methods disclosed herein. It starts with purified (not necessarily separated) gDNA and RNA from a biological sample (step 1). The total nucleic acids are fragmented by enzymatic digestion (for DNA) and by heat hydrolysis (for RNA). The double stranded DNA fragments are end polished so that they are ready for ligation (step 2). The fragmented RNAs are end polished by polyadenylation (step 3). In the next few steps, DNA fragments are ligated to DNA tag molecules (step 4), and the RNA fragments are attached with RNA tag molecules (on both ends) by template switching reverse transcription (step 5). With both DNA and RNA tags in place, the sample is subjected to targeted enrichment reaction by a set of gene specific primers, in which the regions of interest are amplified and enriched (step 6). Finally, the sample is split into two samples, and further amplified by primers specific for the DNA tag and RNA tag, respectively, and with proper NGS adapter sequences compatible with, e.g., Illumina NGS platform (step 7). The final products are two separate DNA and cDNA libraries resulted from the original DNA and RNA material, respectively, and are ready for sequencing.

[0030] Disclosed herein are methods for preparing DNA and cDNA libraries from a sample, comprising: ligating a DNA tag to an end of a DNA molecule in a sample, wherein the DNA tag comprises a unique molecular identifier (UMI) and a DNA identifier; and performing reverse transcription of a RNA molecule in the sample in the presence of a RNA tag, wherein the RNA tag comprises a RNA identifier, a UMI, and a poly(T). The methods do not require physical separation of the DNA and RNA from the sample.

[0031] In some embodiments, the reverse transcription is performed in the presence of a second RNA tag, wherein the second RNA tag comprises a RNA identifier, a UMI, and a template switching oligonucleotide (TSO).

[0032] In some embodiments, the methods can include ribosomal depletion. Alternatively, in some embodiments, the methods do not require ribosomal depletion. Methods for ribosomal depletion are known in the art, e.g., using RiboZero gold (Illumina: MRZG126).

[0033] The term “sample” can include RNA, DNA, a single cell, multiple cells, fragments of cells, or an aliquot of body fluid, taken from a subject (e.g., a mammalian subject, an animal subject, a human subject, or a non-human animal subject). Samples can be selected by one of skill in the art using any known means known including but not limited to centrifugation, venipuncture, blood draw, excretion, swabbing, biopsy, needle aspirate, lavage sample, scraping, surgical incision, laser capture microdissection, gradient separation, or intervention or other means known in the art. The term “mammal” or “mammalian” as used herein includes both humans and non-humans and include but is not limited to humans, non-human primates, canines, felines, murines, bovines, equines, and porcines.

[0034] As used herein, the term “biological sample” is intended to include, but is not limited to, tissues, cells, biological fluids and isolates thereof, isolated from a subject, as well as tissues, cells, and fluids present within a subject.

[0035] As used herein, a “single cell” refers to one cell. Single cells useful in the methods described herein can be obtained from a tissue of interest, or from a biopsy, blood sample, or cell culture. Additionally, cells from specific organs, tissues, tumors, neoplasms, or the like can be obtained and used in the methods described herein. In general, cells from any population can be used in the methods, such as a population of prokaryotic or eukaryotic organisms, including bacteria or yeast.

[0036] A single cell suspension can be obtained using standard methods known in the art including, for example, enzymatically using trypsin or papain to digest proteins connecting cells in tissue samples or releasing adherent cells in culture, or mechanically separating cells in a sample. Samples can also be selected by one of skill in the art using one or more markers known to be associated with a sample of interest.

[0037] Methods for manipulating single cells are known in the art and include fluorescence activated cell sorting (FACS), micromanipulation and the use of semi-automated cell pickers (e.g., the Quixell™ cell transfer system from Stoelting Co.). Individual cells can, for example, be individually selected based on features detectable by microscopic observation, such as location, morphology, or reporter gene expression.

[0038] Once a desired sample has been identified, the sample is prepared and the cell(s) are lysed to release cellular contents including DNA and RNA, such as gDNA and mRNA, using methods known to those of skill in the art. Lysis can be achieved by, for example, heating the cells, or by the use of detergents or other chemical methods, or by a combination of these. Any suitable lysis method known in the art can be used.

[0039] Nucleic acids from a cell such as DNA or RNA are isolated using methods known to those of skill in the art.

[0040] The term “polynucleotide(s)” or “oligonucleotide(s)” refers to nucleic acids such as DNA molecules and RNA molecules and analogs thereof (e.g., DNA or RNA generated using nucleotide analogs or using nucleic acid chemistry). As desired, the polynucleotides can be made synthetically, e.g., using art-recognized nucleic acid chemistry or enzymatically using, e.g., a polymerase, and, if desired, can be modified. Typical modifications include methylation, biotinylation, and other art-known modifications. In addition, a polynucleotide can be single-stranded or double-stranded and, where desired, linked to a detectable moiety. In some aspects, a polynucleotide can include hybrid molecules, e.g., comprising DNA and RNA.

[0041] “G,” “C,” “A,” “T” and “U” each generally stands for a nucleotide that contains guanine, cytosine, adenine, thymidine and uracil as a base, respectively. However, it will be understood that the term “ribonucleotide” or “nucleotide” can also refer to a modified nucleotide or a surrogate replacement moiety. The skilled person is well aware that guanine, cytosine, adenine, and uracil can be replaced by other moieties without substantially altering the base pairing properties of an oligonucleotide comprising a nucleotide bearing such replacement moiety. For example, without limitation, a nucleotide comprising inosine as its base can base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine can be replaced in nucleotide sequences by a nucleotide containing, for example, inosine. In another example, adenine and cytosine anywhere in the oligonucleotide can be replaced with guanine and uracil, respectively, to form G-U Wobble base pairing with the target mRNA. Sequences containing such replacement moieties are suitable for the compositions and methods described herein.

[0042] The term “DNA” refers to chromosomal DNA, plasmid DNA, phage DNA, or viral DNA that is single stranded or double stranded. DNA can be obtained from prokaryotes or eukaryotes.

[0043] The term “genomic DNA” or gDNA” refers to chromosomal DNA.

[0044] The term “messenger RNA” or “mRNA” refers to an RNA that is without introns and that can be translated into a polypeptide.

[0045] The term “cDNA” refers to a DNA that is complementary or identical to an mRNA, in either single stranded or double stranded form.

[0046] As used herein, “polymerase” and its derivatives, generally refers to any enzyme that can catalyze the polymerization of nucleotides (including analogs thereof) into a nucleic acid strand. Typically but not necessarily, such nucleotide polymerization can occur in a template-dependent fashion. Such polymerases can include without limitation naturally occurring polymerases and any subunits and truncations thereof, mutant polymerases, variant polymerases, recombinant, fusion or otherwise engineered polymerases, chemically modified polymerases, synthetic molecules or assemblies, and any analogs, derivatives or fragments thereof that retain the ability to catalyze such polymerization. Optionally, the polymerase can be a mutant polymerase comprising one or more mutations involving the replacement of one or more amino acids with other amino acids, the insertion or deletion of one or more amino acids from the polymerase, or the linkage of parts of two or more polymerases. Typically, the polymerase comprises one or more active sites at which nucleotide binding and/or catalysis of nucleotide polymerization can occur. Some exemplary polymerases include without limitation DNA polymerases and RNA polymerases. The term “polymerase” and its variants, as used herein, also refers to fusion proteins comprising at least two portions linked to each other, where the first portion comprises a peptide that can catalyze the polymerization of nucleotides into a nucleic acid strand and is linked to a second portion that comprises a second polypeptide. In some embodiments, the second polypeptide can include a reporter enzyme or a processivity-enhancing domain. Optionally, the polymerase can possess 5' exonuclease activity or terminal transferase activity. In some embodiments, the polymerase can be optionally reactivated, for example through the use of heat, chemicals or re-addition of new amounts of polymerase into a reaction mixture. In some embodiments, the polymerase can include a hot-start polymerase or an aptamer based polymerase that optionally can be reactivated.

[0047] The term “extension” and its variants, as used herein, when used in reference to a given primer, comprises any in vivo or in vitro enzymatic activity characteristic of a given polymerase that relates to polymerization of one or more nucleotides onto an end of an existing nucleic acid molecule. Typically but not necessarily such primer extension occurs in a template-dependent fashion; during template-dependent extension, the order and selection

of bases is driven by established base pairing rules, which can include Watson-Crick type base pairing rules or alternatively (and especially in the case of extension reactions involving nucleotide analogs) by some other type of base pairing paradigm. In one non-limiting example, extension occurs via polymerization of nucleotides on the 3'OH end of the nucleic acid molecule by the polymerase.

[0048] As used herein, the terms “ligating,” “ligation,” and their derivatives refer generally to the act or process for covalently linking two or more molecules together, for example, covalently linking two or more nucleic acid molecules to each other. In some embodiments, ligation includes joining nicks between adjacent nucleotides of nucleic acids. In some embodiments, ligation includes forming a covalent bond between an end of a first and an end of a second nucleic acid molecule. In some embodiments, for example embodiments wherein the nucleic acid molecules to be ligated include conventional nucleotide residues, the ligation can include forming a covalent bond between a 5' phosphate group of one nucleic acid and a 3' hydroxyl group of a second nucleic acid thereby forming a ligated nucleic acid molecule. In some embodiments, any means for joining nicks or bonding a 5' phosphate to a 3' hydroxyl between adjacent nucleotides can be employed. In an exemplary embodiment, an enzyme such as a ligase can be used. Generally for the purposes of this disclosure, an amplified target sequence can be ligated to an adapter to generate an adapter-ligated amplified target sequence.

[0049] As used herein, “ligase” and its derivatives, refers generally to any agent capable of catalyzing the ligation of two substrate molecules. In some embodiments, the ligase includes an enzyme capable of catalyzing the joining of nicks between adjacent nucleotides of a nucleic acid. In some embodiments, the ligase includes an enzyme capable of catalyzing the formation of a covalent bond between a 5' phosphate of one nucleic acid molecule to a 3' hydroxyl of another nucleic acid molecule thereby forming a ligated nucleic acid molecule. Suitable ligases can include, but not limited to, T4 DNA ligase, T4 RNA ligase, and *E. coli* DNA ligase.

[0050] As used herein, “ligation conditions” and its derivatives, generally refers to conditions suitable for ligating two molecules to each other. In some embodiments, the ligation conditions are suitable for sealing nicks or gaps between nucleic acids. As defined herein, a “nick” or “gap” refers to a nucleic acid molecule that lacks a directly bound 5' phosphate of a mononucleotide pentose ring to a 3' hydroxyl of a neighboring mononucleotide pentose ring within internal nucleotides of a nucleic acid sequence. As used

herein, the term nick or gap is consistent with the use of the term in the art. Typically, a nick or gap can be ligated in the presence of an enzyme, such as ligase at an appropriate temperature and pH. In some embodiments, T4 DNA ligase can join a nick between nucleic acids at a temperature of about 70°C-72°C.

[0051] As used herein, “blunt-end ligation” and its derivatives, refers generally to ligation of two blunt-end double-stranded nucleic acid molecules to each other. A “blunt end” refers to an end of a double-stranded nucleic acid molecule wherein substantially all of the nucleotides in the end of one strand of the nucleic acid molecule are base paired with opposing nucleotides in the other strand of the same nucleic acid molecule. A nucleic acid molecule is not blunt ended if it has an end that includes a single-stranded portion greater than two nucleotides in length, referred to herein as an “overhang.” In some embodiments, the end of nucleic acid molecule does not include any single stranded portion, such that every nucleotide in one strand of the end is based paired with opposing nucleotides in the other strand of the same nucleic acid molecule. In some embodiments, the ends of the two blunt ended nucleic acid molecules that become ligated to each other do not include any overlapping, shared or complementary sequence. Typically, blunted-end ligation excludes the use of additional oligonucleotide adapters to assist in the ligation of the double-stranded amplified target sequence to the double-stranded adapter, such as patch oligonucleotides as described in Mitra and Varley, US2010/0129874. In some embodiments, blunt-ended ligation includes a nick translation reaction to seal a nick created during the ligation process.

[0052] The term “amplicon” refers to the amplified product of a nucleic acid amplification reaction, e.g., RT-PCR.

[0053] The terms “reverse-transcriptase PCR” and “RT-PCR” refer to a type of PCR where the starting material is mRNA. The starting mRNA is enzymatically converted to complementary DNA or “cDNA” using a reverse transcriptase enzyme. The cDNA is then used as a template for a PCR reaction.

[0054] The terms “PCR product,” “PCR fragment,” and “amplification product” refer to the resultant mixture of compounds after two or more cycles of the PCR steps of denaturation, annealing and extension are complete. These terms encompass the case where there has been amplification of one or more segments of one or more target sequences.

[0055] The term “amplification reagents” refers to those reagents (deoxyribonucleotide triphosphates, buffer, etc.), needed for amplification except for primers, nucleic acid template, and the amplification enzyme. Typically, amplification reagents along

with other reaction components are placed and contained in a reaction vessel (test tube, microwell, etc.). Amplification methods include PCR methods known to those of skill in the art and also include rolling circle amplification (Blanco et al., *J. Biol. Chem.*, 264, 8935-8940, 1989), hyperbranched rolling circle amplification (Lizard et al., *Nat. Genetics*, 19, 225-232, 1998), and loop-mediated isothermal amplification (Notomi et al., *Nucl. Acids Res.*, 28, e63, 2000), each of which is hereby incorporated by reference in its entirety.

[0056] The term “hybridize” refers to a sequence specific non-covalent binding interaction with a complementary nucleic acid. Hybridization can occur to all or a portion of a nucleic acid sequence. Those skilled in the art will recognize that the stability of a nucleic acid duplex, or hybrids, can be determined by the T_m . Additional guidance regarding hybridization conditions can be found in: *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y., 1989, 6.3.1-6.3.6 and in: Sambrook et al., *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Vol. 3, 1989.

[0057] As used herein, “incorporating” a sequence into a polynucleotide refers to covalently linking a series of nucleotides with the rest of the polynucleotide, for example at the 3' or 5' end of the polynucleotide, by phosphodiester bonds, wherein the nucleotides are linked in the order prescribed by the sequence. A sequence has been “incorporated” into a polynucleotide, or equivalently the polynucleotide “incorporates” the sequence, if the polynucleotide contains the sequence or a complement thereof. Incorporation of a sequence into a polynucleotide can occur enzymatically (e.g., by ligation or polymerization) or using chemical synthesis (e.g., by phosphoramidite chemistry).

[0058] As used herein, the terms “amplify” and “amplification” refer to enzymatically copying the sequence of a polynucleotide, in whole or in part, so as to generate more polynucleotides that also contain the sequence or a complement thereof. The sequence being copied is referred to as the template sequence. Examples of amplification include DNA-templated RNA synthesis by RNA polymerase, RNA-templated first-strand cDNA synthesis by reverse transcriptase, and DNA-templated PCR amplification using a thermostable DNA polymerase. Amplification includes all primer-extension reactions. Amplification includes methods such as PCR, ligation amplification (or ligase chain reaction, LCR) and amplification methods. These methods are known and widely practiced in the art. See, e.g., U.S. Pat. Nos. 4,683,195 and 4,683,202 and Innis et al., “PCR protocols: a guide to method and applications” Academic Press, Incorporated (1990) (for PCR); and Wu et al. (1989) *Genomics* 4:560-569 (for LCR). In general, the PCR procedure describes a method of gene

amplification which is comprised of (i) sequence-specific hybridization of primers to specific genes within a DNA sample (or library), (ii) subsequent amplification involving multiple rounds of annealing, elongation, and denaturation using a DNA polymerase, and (iii) screening the PCR products for a band of the correct size. The primers used are oligonucleotides of sufficient length and appropriate sequence to provide initiation of polymerization, i.e. each primer is specifically designed to be complementary to each strand of the genomic locus to be amplified.

[0059] Reagents and hardware for conducting amplification reaction are commercially available. Primers useful to amplify sequences from a particular gene region are preferably complementary to, and hybridize specifically to sequences in the target region or in its flanking regions and can be prepared using the polynucleotide sequences provided herein. Nucleic acid sequences generated by amplification can be sequenced directly.

[0060] The term “associated” is used herein to refer to the relationship between a sample and the DNA molecules, RNA molecules, or other polynucleotides originating from or derived from that sample. A polynucleotide is associated with a sample if it is an endogenous polynucleotide, i.e., it occurs in the sample at the time the sample is selected, or is derived from an endogenous polynucleotide. For example, the mRNAs endogenous to a cell are associated with that cell. cDNAs resulting from reverse transcription of these mRNAs, and DNA amplicons resulting from PCR amplification of the cDNAs, contain the sequences of the mRNAs and are also associated with the cell. The polynucleotides associated with a sample need not be located or synthesized in the sample, and are considered associated with the sample even after the sample has been destroyed (for example, after a cell has been lysed). Molecular barcoding or other techniques can be used to determine which polynucleotides in a mixture are associated with a particular sample.

[0061] When hybridization occurs in an antiparallel configuration between two single-stranded polynucleotides, the reaction is called “annealing” and those polynucleotides are described as “complementary”. As used herein, and unless otherwise indicated, the term “complementary,” when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of a polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with a polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Such conditions can, for example, be stringent conditions, where stringent conditions can include: 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C

for 12-16 hours followed by washing. Other conditions, such as physiologically relevant conditions as can be encountered inside an organism, can apply. The skilled person will be able to determine the set of conditions most appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

[0062] Complementary sequences include base-pairing of a region of a polynucleotide comprising a first nucleotide sequence to a region of a polynucleotide comprising a second nucleotide sequence over the length or a portion of the length of one or both nucleotide sequences. Such sequences can be referred to as “complementary” with respect to each other herein. However, where a first sequence is referred to as “substantially complementary” with respect to a second sequence herein, the two sequences can be complementary, or they can include one or more, but generally not more than about 5, 4, 3, or 2 mismatched base pairs within regions that are base-paired. For two sequences with mismatched base pairs, the sequences will be considered “substantially complementary” as long as the two nucleotide sequences bind to each other via base-pairing.

[0063] Conventional notation is used herein to describe nucleotide sequences: the left-hand end of a single-stranded nucleotide sequence is the 5'-end; the left-hand direction of a double-stranded nucleotide sequence is referred to as the 5'-direction. The direction of 5' to 3' addition of nucleotides to nascent RNA transcripts is referred to as the transcription direction. The DNA strand having the same sequence as an mRNA is referred to as the “coding strand”; sequences on the DNA strand having the same sequence as an mRNA transcribed from that DNA and which are located 5' to the 5'-end of the RNA transcript are referred to as “upstream sequences”; sequences on the DNA strand having the same sequence as the RNA and which are 3' to the 3' end of the coding RNA transcript are referred to as “downstream sequences.”

[0064] In some embodiments, the double stranded DNA fragments can be end polished so that they are amenable for ligation. For example, the ends of the DNA fragments can be polished to have blunt ends. As known in the art, this can be achieved with enzymes that can either fill in or remove the protruding strand. Another method is to perform the ligation in the presence of short synthetic oligonucleotides, called “adaptors,” which have been prepared in such a way as to eventually ligate with one terminus to the fragment and make the fragment amenable for ligation with polynucleotides of interest such as DNA or RNA tags. As such, the DNA fragments can be ligated to DNA tags.

[0065] In some embodiments, the RNA fragments are end polished by polyadenylation. The RNA fragments can be attached to RNA tags, e.g., on both ends, by template switching reverse transcription.

[0066] A “DNA tag” or “DNA tag molecule” is a polynucleotide comprising a DNA identifier and a UMI. A DNA tag can be a deoxyribopolynucleotide. A “DNA identifier” is a polynucleotide sequence assigned to distinguish a gDNA molecule from a RNA molecule. A DNA tag can be ligated to the 5’ or 3’ end of double stranded DNA fragments.

[0067] A “RNA tag” or “RNA tag molecule” is a polynucleotide comprising a RNA identifier and a UMI. A RNA tag can be a deoxyribopolynucleotide. A “RNA identifier” is a polynucleotide sequence assigned to distinguish a cDNA molecule from a gDNA molecule. A RNA tag can further comprise poly(T). Alternatively, a RNA tag can further comprise a template switching oligonucleotide (TSO). A RNA tag can be used to add a 5’ tag to RNA-derived cDNA fragments through reverse transcription. In some embodiments, a RNA tag can be used to add a 3’ tag to RNA-derived cDNA through template switching in reverse transcription.

[0068] Two types of RNA tags are helpful because in order to sequence the single stranded RNA from both directions, two different mechanisms can be used to attach the RNA specific sequence. Only one type of DNA tag is needed because the DNA tag can be ligated to both ends of the double stranded DNA.

[0069] A composition can comprise at least 2 of the tags described above, e.g., a DNA tag and a RNA tag. A composition can also comprise the 3 tags described above, e.g., a DNA tag and the 2 types of RNA tags.

[0070] Unique molecular indices or identifiers (UMIs; also called Random Molecular Tags (RMTs)) are short sequences or “barcodes” of bases used to tag each DNA or RNA molecule (fragment) prior to library amplification, thereby aiding in the identification of each individual nucleic acid molecule, or PCR duplicates. Kivioja, T. *et al.*, *Nat. Methods* 9:72-74 (2012), and Suppl. If two reads align to the same location and have the same UMI, it is highly likely that they are PCR duplicates originating from the same fragment prior to amplification. UMIs can also be used to detect and quantify unique mRNA transcripts. In some embodiments, DNA tags containing the same DNA identifier sequence contain different UMI sequences. In some embodiments, RNA tags containing the same RNA identifier sequence contain different UMI sequences.

[0071] The concept of UMIs is that prior to any amplification, each original target molecule is ‘tagged’ by a unique barcode sequence. This DNA sequence must be long enough to provide sufficient permutations to assign each founder molecule a unique barcode. In some embodiments, a UMI sequence contains randomized nucleotides and is incorporated into the DNA or RNA tag. For example, a 12-base random sequence provides 4^{12} or 16,777,216 UMI’s for each target molecule in the sample.

[0072] In some embodiments, the RNA tag is a single-stranded DNA molecule and serves as a primer for reverse transcription. The RNA tag can be generated using a DNA polymerase (DNAP). Here, the binding site of the RNA tag is an RNA binding site (e.g., an mRNA binding site) and contains a sequence region complementary to a sequence region in one or more RNAs. In some embodiments, the binding site is complementary to a sequence region common to all RNAs in the sample to which the barcode adapter is added. For example, the binding site can be a poly(T) tract, which is complementary to the poly(A) tails of eukaryotic mRNAs. Alternatively or in addition, the binding site can include a random sequence tract. Upon adding the RNA tag to the RNAs associated with a sample, reverse transcription can occur and first strands of cDNA can be synthesized, such that the RNA identifier sequence is incorporated into the first strands of cDNA. It will be recognized that reverse transcription requires appropriate conditions, for example the presence of an appropriate buffer and reverse transcriptase enzyme, and temperatures appropriate for annealing of the barcode adapter to RNAs and the activity of the enzyme. It will also be recognized that reverse transcription, involving a DNA primer and an RNA template, is most efficient when the 3’ end of the primer is complementary to the template and can anneal directly to the template. Accordingly, the RNA tag can be designed so that the binding site occurs at the 3’ end of the adapter molecule.

[0073] As described above, the present methods can employ a reverse transcriptase enzyme that adds one or more non-templated nucleotides (such as Cs) to the end of a nascent cDNA strand upon reaching the 5’ end of the template RNA. These nucleotides form a 3’ DNA overhang at one end of the RNA/DNA duplex. If a second RNA molecule contains a sequence region, for example, a poly-G tract at its 3’ end that is complementary to the non-templated nucleotides, and binds to the non-templated nucleotides, the reverse transcriptase can switch templates and continue extending the cDNA, now using the second RNA molecule as a template. Such a second RNA molecule is referred to herein and known in the art as a template-switching oligo (TSO).

[0074] In embodiments of the present methods, a second RNA tag comprising a RNA identifier, UMI, and TSO can serve as a template-switching oligonucleotide for reverse transcription. Thus, the RNA identifier sequence is incorporated into the first strand of cDNA after template switching, and is present in DNA molecules resulting from amplification (for example, by PCR) of the first strand of cDNA. In these embodiments, any reverse transcriptase that has template switching activity can be used. The binding site of the first RNA tag is a cDNA binding site and preferably occurs at the 3' end of the adapter molecule. The binding site can include a G-tract (comprising one or more G nucleotides), or any other sequence that is at least partially complementary to that of the 3' overhang generated by the reverse transcriptase. It will be recognized that the overhang sequence, and thus an appropriate sequence for the binding site of the barcode adapter, can depend on the choice of reverse transcriptase used in the method.

[0075] Methods for reverse transcription and template switching are well known in the art. A procedure frequently referred to as "SMART" (switching mechanism at the 5' end of the RNA transcript) can generate full-length cDNA libraries, even from single-cell-derived RNA samples. This strategy relies on the intrinsic properties of Moloney murine leukemia virus (MMLV) reverse transcriptase and the use of a unique template switching oligonucleotide (TS oligo, or TSO). Moloney Murine Leukemia Virus Reverse Transcriptase (M-MLV RT) is an RNA-dependent DNA polymerase that can be used in cDNA synthesis with long messenger RNA templates (>5kb). The enzyme is a product of the *pol* gene of M-MLV and consists of a single subunit with a molecular weight of 71kDa. During first-strand synthesis, upon reaching the 5' end of the RNA template, the terminal transferase activity of the MMLV reverse transcriptase adds a few additional nucleotides (mostly deoxycytidine) to the 3' end of the newly synthesized cDNA strand. These bases function as a TS oligo-anchoring site. Upon base pairing between the TS oligo and the appended deoxycytidine stretch, the reverse transcriptase "switches" template strands, from cellular RNA to the TS oligo, and continues replication to the 5' end of the TS oligo. By doing so, the resulting cDNA contains the complete 5' end of the transcript, and universal sequences of choice can be added to the reverse transcription product. Along with tagging of the cDNA 3' end by oligo dT primers, this approach makes it possible to efficiently amplify the entire full-length transcript pool in a completely sequence-independent manner.

[0076] A TS oligo can be a DNA oligo sequence that carries 3 riboguanosines (rGrGrG) at its 3' end. The complementarity between these consecutive rG bases and the 3'

dC extension of the cDNA molecule allows the subsequent template switching. The 3' most rG can also be replaced with a locked nucleic acid base (LNA) to enhance thermostability of the LNA monomer, which would be advantageous for base pairing.

[0077] The TSO can include a 3' portion comprising a plurality of guanosines or guanosine analogues that base pair with cytosine. Non-limiting examples of guanosines or guanosine analogues useful in the methods described herein include, but are not limited to, deoxyriboguanosine, riboguanosine, locked nucleic acid-guanosine, and peptide nucleic acid-guanosine. The guanosines can be ribonucleosides or locked nucleic acid monomers.

[0078] The TSO can include a 3' portion including at least 2, at least 3, at least 4, at least 5, or 2, 3, 4, or 5, or 2-5 guanosines, or guanosine analogues that base pair with cytosine. The presence of a plurality of guanosines (or guanosine analogues that base pair with cytosine) allows the TSO to anneal transiently to the exposed cytosines at the 3' end of the first strand of cDNA. This causes the reverse transcriptase to switch template and continue to synthesis a strand complementary to the TSO. In one aspect of the invention, the 3' end of the TSO can be blocked, for example by a 3' phosphate group, to prevent the TSO from functioning as a primer during cDNA synthesis.

[0079] Before the tagged cDNA samples are pooled, synthesis of cDNA can be stopped, for example by removing or inactivating the reverse transcriptase. This prevents cDNA synthesis by reverse transcription from continuing in the pooled samples.

[0080] As used herein, "amplified target sequences" and its derivatives, refers generally to a nucleic acid sequence produced by the amplification of/amplifying the target sequences using target-specific primers and the methods provided herein. The amplified target sequences can be either of the same sense (the positive strand produced in the second round and subsequent even-numbered rounds of amplification) or antisense (i.e., the negative strand produced during the first and subsequent odd-numbered rounds of amplification) with respect to the target sequences. For the purposes of this disclosure, the amplified target sequences are typically less than 50% complementary to any portion of another amplified target sequence in the reaction.

[0081] The term "polymerase chain reaction" ("PCR") of Mullis (U.S. Pat. Nos. 4,683,195, 4,683,202, and 4,965,188) refers to a method for increasing the concentration of a segment of a target sequence in a mixture of nucleic acid sequences without cloning or purification. This process for amplifying the target sequence consists of introducing a large excess of two oligonucleotide primers to the nucleic acid sequence mixture containing the

desired target sequence, followed by a precise sequence of thermal cycling in the presence of a polymerase (e.g., DNA polymerase). The two primers are complementary to their respective strands of the double stranded target sequence. To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. The steps of denaturation, primer annealing, and polymerase extension can be repeated many times (i.e., denaturation, annealing and extension constitute one “cycle;” there can be numerous “cycles”) to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the “polymerase chain reaction” (hereinafter “PCR”). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be “PCR amplified.”

[0082] The methods disclosed herein can further comprise amplifying the tagged DNA the tagged cDNA for enrichment with a set of gene specific primers. Target enrichment can be achieved with, e.g., an SPE primer pool, DNA boosting primer, and RNA boosting primer. Amplicon-based next-generation sequencing (NGS) assays offer many advantages for targeted enrichment. For example, QIAseq NGS panels employ unique molecular indices (UMI's) to correct for PCR amplification bias and use single primer extension (SPE) technology which provides design flexibility and highly-specific target enrichment. The concept of UMIs is that prior to any amplification, each original target molecule is ‘tagged’ by a unique barcode sequence. This DNA sequence must be long enough to provide sufficient permutations to assign each founder molecule a unique barcode. In its current form, a 12-base random sequence provides 4^{12} or 16,777,216 UMI's for each target molecule in the sample.

[0083] As used herein, the term “primer” includes an oligonucleotide, either natural or synthetic, that is capable, upon forming a duplex with a polynucleotide template, of acting as a point of initiation of nucleic acid synthesis and being extended from its 3' end along the template so that an extended duplex is formed. The sequence of nucleotides added during the extension process is determined by the sequence of the template polynucleotide. Usually primers are extended by a DNA polymerase. Primers usually have a length in the range of

between 3 to 36 nucleotides, also 5 to 24 nucleotides, also from 14 to 36 nucleotides. Primers within the scope of the invention include orthogonal primers, amplification primers, constructions primers and the like. Pairs of primers can flank a sequence of interest or a set of sequences of interest. Primers and probes can be degenerate in sequence. Primers within the scope of the present invention bind adjacent to a target sequence. A “primer” can be considered a short polynucleotide, generally with a free 3'-OH group that binds to a target or template potentially present in a sample of interest by hybridizing with the target, and thereafter promoting polymerization of a polynucleotide complementary to the target. Primers of the instant invention are comprised of nucleotides ranging from 17 to 30 nucleotides. In some embodiments, the primer is at least 17 nucleotides, or alternatively, at least 18 nucleotides, or alternatively, at least 19 nucleotides, or alternatively, at least 20 nucleotides, or alternatively, at least 21 nucleotides, or alternatively, at least 22 nucleotides, or alternatively, at least 23 nucleotides, or alternatively, at least 24 nucleotides, or alternatively, at least 25 nucleotides, or alternatively, at least 26 nucleotides, or alternatively, at least 27 nucleotides, or alternatively, at least 28 nucleotides, or alternatively, at least 29 nucleotides, or alternatively, at least 30 nucleotides, or alternatively at least 50 nucleotides, or alternatively at least 75 nucleotides or alternatively at least 100 nucleotides.

[0084] As used herein, “target-specific primer” and its derivatives, refers generally to a single stranded or double-stranded polynucleotide, typically an oligonucleotide, that includes at least one sequence that is at least 50% complementary, typically at least 75% complementary or at least 85% complementary, more typically at least 90% complementary, more typically at least 95% complementary, more typically at least 98% or at least 99% complementary, or 100% identical, to at least a portion of a nucleic acid molecule that includes a target sequence. In such instances, the target-specific primer and target sequence are described as “corresponding” to each other. In some embodiments, the target-specific primer is capable of hybridizing to at least a portion of its corresponding target sequence (or to a complement of the target sequence); such hybridization can optionally be performed under standard hybridization conditions or under stringent hybridization conditions. In some embodiments, the target-specific primer is not capable of hybridizing to the target sequence, or to its complement, but is capable of hybridizing to a portion of a nucleic acid strand including the target sequence, or to its complement. In some embodiments, the target-specific primer includes at least one sequence that is at least 75% complementary, typically at least 85% complementary, more typically at least 90% complementary, more typically at least

95% complementary, more typically at least 98% complementary, or more typically at least 99% complementary, to at least a portion of the target sequence itself; in other embodiments, the target-specific primer includes at least one sequence that is at least 75% complementary, typically at least 85% complementary, more typically at least 90% complementary, more typically at least 95% complementary, more typically at least 98% complementary, or more typically at least 99% complementary, to at least a portion of the nucleic acid molecule other than the target sequence. In some embodiments, the target-specific primer is substantially non-complementary to other target sequences present in the sample; optionally, the target-specific primer is substantially non-complementary to other nucleic acid molecules present in the sample. In some embodiments, nucleic acid molecules present in the sample that do not include or correspond to a target sequence (or to a complement of the target sequence) are referred to as “non-specific” sequences or “non-specific nucleic acids”. In some embodiments, the target-specific primer is designed to include a nucleotide sequence that is substantially complementary to at least a portion of its corresponding target sequence. In some embodiments, a target-specific primer is at least 95% complementary, or at least 99% complementary, or 100% identical, across its entire length to at least a portion of a nucleic acid molecule that includes its corresponding target sequence. In some embodiments, a target-specific primer can be at least 90%, at least 95% complementary, at least 98% complementary or at least 99% complementary, or 100% identical, across its entire length to at least a portion of its corresponding target sequence. In some embodiments, a forward target-specific primer and a reverse target-specific primer define a target-specific primer pair that can be used to amplify the target sequence via template-dependent primer extension. Typically, each primer of a target-specific primer pair includes at least one sequence that is substantially complementary to at least a portion of a nucleic acid molecule including a corresponding target sequence but that is less than 50% complementary to at least one other target sequence in the sample. In some embodiments, amplification can be performed using multiple target-specific primer pairs in a single amplification reaction, wherein each primer pair includes a forward target-specific primer and a reverse target-specific primer, each including at least one sequence that substantially complementary or substantially identical to a corresponding target sequence in the sample, and each primer pair having a different corresponding target sequence. In some embodiments, the target-specific primer can be substantially non-complementary at its 3' end or its 5' end to any other target-specific primer present in an amplification reaction. In some embodiments, the target-specific primer can

include minimal cross hybridization to other target-specific primers in the amplification reaction. In some embodiments, target-specific primers include minimal cross-hybridization to non-specific sequences in the amplification reaction mixture. In some embodiments, the target-specific primers include minimal self-complementarity. In some embodiments, the target-specific primers can include one or more cleavable groups located at the 3' end. In some embodiments, the target-specific primers can include one or more cleavable groups located near or about a central nucleotide of the target-specific primer. In some embodiments, one of more targets-specific primers includes only non-cleavable nucleotides at the 5' end of the target-specific primer. In some embodiments, a target specific primer includes minimal nucleotide sequence overlap at the 3' end or the 5' end of the primer as compared to one or more different target-specific primers, optionally in the same amplification reaction. In some embodiments 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more, target-specific primers in a single reaction mixture include one or more of the above embodiments. In some embodiments, substantially all of the plurality of target-specific primers in a single reaction mixture includes one or more of the above embodiments.

[0085] Primer design is based on single primer extension, in which each genomic target is enriched by one target-specific primer and one universal primer – a strategy that removes conventional two target-specific primer design restriction and reduces the amount of required primers. All primers required for a panel are pooled into an individual primer pool to reduce panel handling and the number of pools required for enrichment and library construction.

[0086] The booster panel is a pool of up to 100 primers that can be used to boost the performance of certain primers in any panel (cataloged, extended, or custom), or to extend the contents of an existing custom panel. The primers are delivered as a single pool that can be spiked into the existing panel.

[0087] After removing unused adapters, a limited number of PCR cycles can be conducted using an adapter primer and a pool of single primers, each carrying a gene specific sequence and a 5' universal sequence. During this process, each single primer repeatedly samples the same target locus from different DNA templates. Afterwards, additional PCR cycles can be conducted using universal primers to attach complete adapter sequences and to amplify the library to the desired quantity.

[0088] Compared to existing targeted enrichment approaches, the SPE method relies on single end adapter ligation, which inherently has a much higher efficiency than requiring

adapters to ligate to both ends of the dsDNA fragment. More DNA molecules will be available for the downstream PCR enrichment step. PCR enrichment efficiency using one primer is also better than conventional two primer approach, due to the absence of an efficiency constraint from a second primer. During the initial PCR cycles, primers have repeated opportunities to convert (i.e. capture) maximal amount of original DNA molecules into amplicons.

[0089] All three features help to increase the efficiency of capturing rare mutations in the sample. In addition, incorporated UMI's within the amplicon are the key to estimating the number of DNA molecules captured and to greatly reduce sequencing errors in downstream analysis. Single primer extension also permits discovery of unknown structural variants, such as gene fusions.

[0090] The targeted enriched sample of DNA (e.g., gDNA) and cDNA are split into 2 separate samples. A first sample can be amplified by polymerase chain reaction (PCR) using primers specific for the DNA tag to generate a DNA library corresponding to the DNA in the sample. A second sample can be amplified by PCR using primers specific for the RNA tag to generate a cDNA library corresponding to the RNA in the sample.

[0091] A real-time polymerase chain reaction (Real-Time PCR), also known as quantitative polymerase chain reaction (qPCR), is a laboratory technique of molecular biology based on the polymerase chain reaction (PCR). It monitors the amplification of a targeted DNA molecule during the PCR, i.e. in real-time, and not at its end, as in conventional PCR. Real-time PCR can be used quantitatively (quantitative real-time PCR), and semi-quantitatively, i.e. above/below a certain amount of DNA molecules (semi quantitative real-time PCR). Other types of PCRs include but are not limited to nested PCR (used to analyze DNA sequences coming from different organisms of the same species but that can differ for a single nucleotide (SNIPS) and to ensure amplification of the sequence of interest in each of the organism analyzed) and Inverse-PCR (usually used to clone a region flanking an insert or a transposable element).

[0092] Two common methods for the detection of PCR products in real-time PCR are: (1) non-specific fluorescent dyes that intercalate with any double-stranded DNA, and (2) sequence-specific DNA probes consisting of oligonucleotides that are labeled with a fluorescent reporter which permits detection only after hybridization of the probe with its complementary sequence.

[0093] Methods and kits for performing PCR are well known in the art. PCR is a reaction in which replicate copies are made of a target polynucleotide using a pair of primers or a set of primers consisting of an upstream and a downstream primer, and a catalyst of polymerization, such as a DNA polymerase, and typically a thermally-stable polymerase enzyme. Methods for PCR are well known in the art, and taught, for example in MacPherson et al. (1991) PCR 1: A Practical Approach (IRL Press at Oxford University Press).

[0094] Embodiments of the invention provide 2 separate libraries for flexible manipulation downstream: a DNA library based on the original DNA and a cDNA library based on the original RNA produced by any of the methods described herein. The DNA library or cDNA library can be sequenced to provide an analysis of gene expression in single cells or in a plurality of single cells.

[0095] The amplified DNA or cDNA library can be sequenced and analyzed using methods known to those of skill in the art, e.g., by next-generation sequencing (NGS). In certain exemplary embodiments, RNA expression profiles are determined using any sequencing methods known in the art. Determination of the sequence of a nucleic acid sequence of interest can be performed using a variety of sequencing methods known in the art including, but not limited to, sequencing by synthesis (SBS), sequencing by hybridization (SBH), sequencing by ligation (SBL) (Shendure et al. (2005) Science 309:1728), quantitative incremental fluorescent nucleotide addition sequencing (QIFNAS), stepwise ligation and cleavage, fluorescence resonance energy transfer (FRET), molecular beacons, TaqMan reporter probe digestion, pyrosequencing, fluorescent in situ sequencing (FISSEQ), FISSEQ beads (U.S. Pat. No. 7,425,431), wobble sequencing (PCT/US05/27695), multiplex sequencing (U.S. Ser. No. 12/027,039, filed Feb. 6, 2008; Porreca et al (2007) Nat. Methods 4:931), polymerized colony (POLONY) sequencing (U.S. Pat. Nos. 6,432,360, 6,485,944 and 6,511,803, and PCT/US05/06425); nanogrid rolling circle sequencing (ROLONY) (US2009/0018024), allele-specific oligo ligation assays (e.g., oligo ligation assay (OLA), single template molecule OLA using a ligated linear probe and a rolling circle amplification (RCA) readout, ligated padlock probes, and/or single template molecule OLA using a ligated circular padlock probe and a rolling circle amplification (RCA) readout) and the like. High-throughput sequencing methods, e.g., using platforms such as Roche 454, Illumina Solexa, AB-SOLiD, Helicos, Complete Genomics, Polonator platforms and the like, can also be utilized. A variety of light-based sequencing technologies are known in the art (Landegren et

al. (1998) *Genome Res.* 8:769-76; Kwok (2000) *Pharmacogenomics* 1:95-100; and Shi (2001) *Clin. Chem.* 47:164-172).

[0096] Embodiments of the invention also provide methods for analyzing gene expression in a plurality of single cells, the method comprising the steps of preparing a cDNA library using the method described herein and sequencing the cDNA library. A “gene” refers to a polynucleotide containing at least one open reading frame (ORF) that is capable of encoding a particular polypeptide or protein after being transcribed and translated. Any of the polynucleotide sequences described herein can be used to identify larger fragments or full-length coding sequences of the gene with which they are associated. Methods of isolating larger fragment sequences are known to those of skill in the art.

[0097] As used herein, “expression” refers to the process by which polynucleotides are transcribed into mRNA and/or the process by which the transcribed mRNA is subsequently being translated into peptides, polypeptides, or proteins. If the polynucleotide is derived from genomic DNA, expression can include splicing of the mRNA in a eukaryotic cell.

[0098] The cDNA library can be sequenced by any suitable screening method. In particular, the cDNA library can be sequenced using a high-throughput screening method, such as Applied Biosystems’ SOLiD sequencing technology, or Illumina’s Genome Analyzer. In one aspect of the invention, the cDNA library can be shotgun sequenced. The number of reads can be at least 10,000, at least 1 million, at least 10 million, at least 100 million, or at least 1000 million. In another aspect, the number of reads can be from 10,000 to 100,000, or alternatively from 100,000 to 1 million, or alternatively from 1 million to 10 million, or alternatively from 10 million to 100 million, or alternatively from 100 million to 1000 million. A “read” is a length of continuous nucleic acid sequence obtained by a sequencing reaction.

[0099] The DNA or gDNA library generated by the methods disclosed herein can be useful for, but not limited to, DNA variant detection, copy number analysis, fusion gene detection and structural variant detection. The cDNA library generated by the methods disclosed herein can be useful for, but not limited to, RNA variant detection, gene expression analysis, and fusion gene detection. The DNA and cDNA libraries can also be used for paired DNA and RNA profiling.

[0100] The expression profiles described herein are useful in the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring

clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, some embodiments relate to diagnostic assays for determining the expression profile of nucleic acid sequences (e.g., RNAs), in order to determine whether an individual is at risk of developing a disorder and/or disease. Such assays can be used for prognostic or predictive purposes to thereby prophylactically treat an individual prior to the onset of the disorder and/or disease. Accordingly, in certain exemplary embodiments, methods of diagnosing and/or prognosing one or more diseases and/or disorders using one or more of expression profiling methods described herein are provided.

[0101] Some embodiments pertain to monitoring the influence of agents (e.g., drugs or other compounds administered either to inhibit or to treat or prevent a disorder and/or disease) on the expression profile of nucleic acid sequences (e.g., RNAs) in clinical trials. Accordingly, in certain exemplary embodiments, methods of monitoring one or more diseases and/or disorders before, during and/or subsequent to treatment with one or more agents using one or more of expression profiling methods described herein are provided.

[0102] Monitoring the influence of agents (e.g., drug compounds) on the level of expression of a marker of the invention can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent to affect an expression profile can be monitored in clinical trials of subjects receiving treatment for a disease and/or disorder associated with the expression profile. In certain exemplary embodiments, the methods for monitoring the effectiveness of treatment of a subject with an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting one or more expression profiled in the pre-administration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting one or more expression profiles in the post-administration samples; (v) comparing the one or more expression profiled in the pre-administration sample with the one or more expression profiles in the post-administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly.

[0103] The expression profiling methods described herein allow the quantitation of gene expression. Thus, not only tissue specificity, but also the level of expression of a variety of genes in the tissue is ascertainable. Thus, genes can be grouped on the basis of their tissue expression per se and level of expression in that tissue. This is useful, for example, in ascertaining the relationship of gene expression between or among tissues. Thus,

one tissue can be perturbed and the effect on gene expression in a second tissue can be determined. In this context, the effect of one cell type on another cell type in response to a biological stimulus can be determined. Such a determination is useful, for example, to know the effect of cell-cell interaction at the level of gene expression. If an agent is administered therapeutically to treat one cell type but has an undesirable effect on another cell type, the invention provides an assay to determine the molecular basis of the undesirable effect and thus provides the opportunity to co-administer a counteracting agent or otherwise treat the undesired effect. Similarly, even within a single cell type, undesirable biological effects can be determined at the molecular level. Thus, the effects of an agent on expression of other than the target gene can be ascertained and counteracted.

[0104] In another embodiment, the time course of expression of one or more nucleic acid sequences (e.g., genes, mRNAs and the like) in an expression profile can be monitored. This can occur in various biological contexts, as disclosed herein, for example development of a disease and/or disorder, progression of a disease and/or disorder, and processes, such a cellular alterations associated with the disease and/or disorder.

[0105] The expression profiling methods described herein are also useful for ascertaining the effect of the expression of one or more nucleic acid sequences (e.g., genes, mRNAs and the like) on the expression of other nucleic acid sequences (e.g., genes, mRNAs and the like) in the same cell or in different cells. This provides, for example, for a selection of alternate molecular targets for therapeutic intervention if the ultimate or downstream target cannot be regulated.

[0106] The expression profiling methods described herein are also useful for ascertaining differential expression patterns of one or more nucleic acid sequences (e.g., genes, mRNAs and the like) in normal and abnormal cells. This provides a battery of nucleic acid sequences (e.g., genes, mRNAs and the like) that could serve as a molecular target for diagnosis or therapeutic intervention.

EXAMPLES

[0107] Starting Material: Purified genomic DNA and total RNA. For example, 50ng gDNA and 50ng total RNA was purified from THP-1 cell line. Ideally, the relative amount of gDNA and RNA should represent the content in the sample.

[0108] DNA/RNA Fragmentation:

		uL	final conc.
DNA/RNA sample		X	
H ₂ O		11.8 - x	
10x Fragmentation Buffer		2	1x
100 mM dATP		0.6	3 mM
Exonuclease I (20 U/uL)		1.6	1.6 U/uL
5x Fragmentation Enzyme Mix		4	1x
		Total volume: 20 uL	
Incubate in thermocycler with heated lid on for 4°C 1min → 32°C 15min → 75°C 10min → 80°C 20min → 4°C hold			

[0109] RNA Polyadenylation:

		uL	final conc.
Sample from previous step		20	
H ₂ O		0.5	
10 mM ATP		1.25	0.5 mM
10 mM 3'-dATP (blocker)		1.25	0.5 mM
T4 Polynucleotide Kinase (10 U/uL)		1	0.4 U/uL
E. coli Poly(A) Polymerase (5 U/uL)		1	0.2 U/uL
		Total volume: 25 uL	
Incubate in thermocycler with heated lid on for 4°C 1min → 30°C 10min → 4°C hold			

[0110] DNA Ligation:

		uL	final conc.
Sample from previous step		25	
5x Ligation Buffer		10	1x
50 uM DNA ligation Adaptor		2.8	2.8 uM
50% PEG-6000		7.2	7.2%
T4 DNA ligase (600 U/uL)		5	60 U/uL
		Total volume: 50 uL	
Incubate in thermocycler with heated lid OFF for 4C 1min → 20C 15min → 4C hold			

[0111] Purification: Add 50 uL of ice cold water to the 50uL sample from previous step to make 100 uL total. Do 2 rounds of 1.2x Ampure XP beads purification following manufacturer's manual with the following exceptions: 1st round elution in 52uL water; and 2nd round elution in 13uL water.

[0112] Reverse Transcription:

	uL	final conc.
Sample from previous step	12.87	
7.5 uM TSON10T18NV oligo	1	300 nM
25 uM TSON10forTS oligo	1	1 uM
5x SuperScript II Buffer	5	1x
25 mM each dNTP mix	1	1 mM each
0.1 M DTT	1.25	5 mM
RNase Inhibitor (40 U/uL)	0.63	1 U/uL
300 mM MgCl ₂	0.5	6 mM
150 mM MnCl ₂	0.5	3 mM
MMLV Reverse Transcriptase RNase H- (200 U/uL)	1.25	10 U/uL
	Total volume: 25 uL	
Incubate in thermocycler with heated lid on for 4°C 1min → 25°C 10min → 42°C 45min → 70°C 15min → 4°C hold		

[0113] Purification: Add 75 uL of ice cold water to the 25uL sample from previous step to make 100uL total. Do 1 round of 1.2x Ampure XP beads purification following manufacturer's manual, and elute in 16.8 uL water.

[0114] Target Enrichment:

	uL	final conc.
Sample from previous step	16.8	
5x V2 Buffer	8	1x
2 mM each dNTP mix	4	0.2 mM each
100 nM each SPE primer pool	8	20 nM each
10 uM DNA boosting primer	0.8	400 nM
10 uM RNA boosting primer	0.8	400 nM
Hot-Star Taq Polymerase (6 U/uL)	1.6	0.24 U/uL
	Total volume: 40 uL	
Incubate in thermocycler with heated lid on for 95°C 13min → 98°C 2min → 8 cycles of (98°C 15sec → 68°C 10min) → 72°C 5min → 4°C hold		

[0115] Purification: Add 60 uL of ice cold water to the 40uL sample from previous step to make 100uL total. Do double size selection 0.5x/0.5x with Ampure XP beads following manufacturer's manual, and elute in 22uL water.

[0116] qPCR (real-time) to determine final amplification cycles:

	For DNA library		For RNA library	
	<u>uL</u>	<u>final conc.</u>	<u>uL</u>	<u>final conc.</u>
Sample from previous step	2		2	
5x V2 Buffer	2	1x	2	1x
2 mM each dNTP mix	1	0.2 mM each	1	0.2 mM each
H ₂ O	2.1		2.1	
20x EveGreen Dye	0.5	1x	0.5	1x
4 uM IL2N5RS2 Universal primer	1	400 nM	1	400 nM
4 uM DNA Universal Primer	1	400 nM	0	0
4 uM RNA Universal Primer	0	0	1	400 nM
Hot-Star Taq Polymerase (6 U/uL)	0.4	0.24 U/uL	0.4	0.24 U/uL
	Total volume: 10 uL		Total volume: 10 uL	
Run on ABI 7900 real time instrument: 95°C 13min → 98°C 2min → 30 cycles of (98°C 15sec → 62°C 2min).				
Record the counts for both samples				

[0117] Universal PCR:

	For DNA library		For RNA library	
	<u>uL</u>	<u>final conc.</u>	<u>uL</u>	<u>final conc.</u>
Sample from Target Enrichment	9		9	
5x V2 Buffer	5	1x	5	1x
2mM each dNTP mix	2.5	0.2 mM each	2.5	0.2 mM each
4uM IL2N5RS2 Universal primer	2.5	400 nM	2.5	400 nM
4uM DNA Universal Primer	2.5	400 nM	0	0
4uM RNA Universal Primer	0	0	2.5	400 nM
H ₂ O	2.5		2.5	
Hot-Star Taq Polymerase (6U/uL)	1	0.24 U/uL	1	0.24 U/uL
	Total volume: 25 uL		Total volume: 25 uL	
Incubate in thermocycler with heated lid on for 95°C 13min → 98°C 2min → "X" cycles of (98°C 15sec → 62°C 2min) → 72°C 5min → 4°C hold				
(X= Ct + 4) for DNA sample and RNA sample respectively. For example, if Ct = 19 for DNA, and 15 for RNA, then run 23 cycles for DNA, and 19 cycles for RNA				

[0118] Purification: Add 75 uL of ice cold water to each of the 25uL sample from previous step to make 100uL total. Do 1 round of 1.2x Ampure XP beads purification following manufacturer's manual, and elute in 20uL water.

[0119] Library Quantification using Agilent Bioanalyzer High Sensitivity DNA chip: Dilute the purified libraries to 2 ng/uL. Load 1 uL of this diluted sample on the bioanalyzer.

Obtain molar concentration of the libraries based on bionalyzer's electropherogram. The libraries are ready for sequencing.

[0120] Following the workflow, with 50 ng gDNA and 50 ng total RNA input, we obtained 675 ng of DNA library and 455 ng of RNA library. The same amount of 50 ng total RNA was also used with QIAseq Targeted RNAscan Panels system from QIAGEN for comparison purpose. The same amount of 50 ng gDNA was also used with QIAseq Targeted DNA Panels system from QIAGEN for comparison purpose. The samples were then put on Illumina's MiSeq machine for sequencing.

Results:

[0121] As shown in Table 1, compared to the standalone RNA library prep workflow (QIAseq Targeted RNAscan Panels system from QIAGEN), our method achieved around 24% of its enrichment efficiency on the 1st strand cDNA, and around 40% of its enrichment efficiency on the 2nd strand cDNA. Since RNAscan workflow had strand bias toward the 1st strand, our method had less bias and improved strand balance. The effect of enrichment efficiency on RNA analysis deserves further exploration.

[0122] Table 1.

Workflow	RNAScan	Ours
Average UMIs/primer 1 st strand	11061	2681
Average UMIs/primer 2 nd strand	5279	2077
Ratio 2 nd /1 st	0.48	0.77

[0123] UMI per SPE primer for RNA sample: Primers were divided into two groups based on the RNA strand they detected. As shown in Table 2, compared to the standalone DNA library prep workflow (QIAseq Targeted DNA Panels system from QIAGEN), our method achieved slightly better enrichment efficiency. Both of the methods had comparable sequencing specificity and uniformity.

[0124] Table 2.

Workflow	Targeted DNA Panels	Ours
Average UMIs/primer	1471	1701
Average reads/UMI	3.4	3.0
Overall specificity (on- target reads/all reads)	87%	90%
Coverage uniformity (T50)	24.9	21.6

[0125] Sequencing specs for DNA sample in both methods: Sequence coverage uniformity was measured by T50, the percentage of total sequence throughput captured by the bottom 50% of a target region. In the perfect uniform scenario, the T50 value equals to 50.

[0126] Cross talk between DNA and RNA was also evaluated since they remained in the same reaction. Using the same 50ng of DNA and RNA from THP-1 cell line, the effective leaking signal from RNA to DNA was only 0.75% of the real DNA signal, as measured by the total UMIs of the primers detecting both RNA and DNA. In this case, only the extremely highly expressed genes might have an effect on corresponding DNA copy number analysis. However, if DNA copy number analysis was limited on intron regions, this effect should disappear. The effective leaking signal from DNA to RNA was around 3% on average by the same measurement. Since there were only a few copies of genome DNA in each cell in most cases, this kind of leaking could only affect those extremely low expressing genes (less than 0.1 copy per cell), which might be lower than the background noise level. In conclusion, our method demonstrated minimal cross talk between DNA and RNA samples which might not have any significant effect in real cases.

[0127] The DNA library prepared by our method can be used for DNA variant detection, and copy number analysis. The RNA library prepared by our method is suitable for gene expression analysis, fusion gene detection, and RNA variant detection. Multi-modal NGS panels can be developed based on our proposed method, and be used for biomarker screening, or targeted eQTL analysis.

[0128] Adaptor for ligation:

Equal molar mix and annealing of the following 2 oligos to make double strand adaptor (DNA ligation Adaptor)		
SEQ ID NO:1	/5Phos/GGACTCCAATNNNNNNNNNNNNNACGCTAA GAAAGATCGGAAGAGCACACGTCTG/3ddC/	PAGE Purified
SEQ ID NO:2	ATT+GGAG+TCC*T/3Phos/	STD desalt

[0129] Reverse Transcription Oligos:

SEQ ID NO:3 TSON10T18 NV oligo	CGACTCACTATAGGGCTGGAATTCTGACGNNNNNNN NNNACGTTTTTTTTTTTTTTTTTTT	PAGE Purified
SEQ ID NO:4 TSON10for TS oligo	/5Me-isodC//iisodG//iisodG/TAATACGACTCACTATAG GGCTGGAATTCTGACGNNNNNNNNNNNATCTGCrGrGrG	PAGE Purified

[0130] Target Enrichment Oligos:

SEQ ID NO:5 DNA boosting primer	<u>AGCAGTGGTATCAACGCAGAGTCAAGCAGAAGACG</u> GCATACGAGATTCCGAAACGTGACTGGAGTTCAGAC GTGTGCTCTTCCGATCTTCTTAGCGT	STD desalt
SEQ ID NO:6 RNA boosting primer	<u>GTGAGTGATGGTTGAGGATGTGTGCAAGCAGAAGAC</u> GGCATAACGAGATTACGTACGGTGACTGGAGTTCAGA CGTGTGCTCTTCCGATCTCGACTCACTATAGGGCTGG <u>AATTCT</u>	STD desalt
For each primer, the first set of underlined nucleotides is priming site for PCR amplification in Universal PCR reactions, the second set of underlined nucleotides in the middle is the sample idx (index) region, which can be replaced with respective sample index sequences, and the third set of underlined nucleotides is part of DNA or RNA identifier used for PCR amplification in target enrichment reactions.		

[0131] uPCR Primers:

SEQ ID NO:7 IL2N5RS2 Universal primer	AATGATACGGCGACCACCGAGATCTACACT CTTCCCTACACGACGCTCTTCCGATCTNNN NNAATGTACAGTATTGCGTTTTG	PAGE Purified
SEQ ID NO:8 DNA Universal Primer	AAGCAGTGGTATCAACGCAGAGT	STD desalt
SEQ ID NO:9 RNA Universal Primer	GTGAGTGATGGTTGAGGATGTGTG	STD desalt

[0132] SPE Primer Pool (equal molar mix of the following oligos):

SEQ ID NO:10
AATGTACAGTATTGCGTTTTGAGCCCCAAGTCCTATGAGAACCTCTG

SEQ ID NO:11
AATGTACAGTATTGCGTTTTGTGGCACCAGCGATCAGGTCCTTTAT

SEQ ID NO:12
AATGTACAGTATTGCGTTTTGCTGAGTGGAGTCACAGCGGAGATAGT

SEQ ID NO:13
AATGTACAGTATTGCGTTTTGTGTTCCACCAGTAACAACAGTTGAATGTCC

SEQ ID NO:14
AATGTACAGTATTGCGTTTTGGTGTGAGGAACATACTAGTGCTTTGCAAGT

SEQ ID NO:15
AATGTACAGTATTGCGTTTTGTTCAAAGTTGGGTCTGCTTCAGTCCAAAG

SEQ ID NO:16
AATGTACAGTATTGCGTTTTGCCCCAGCTTCTTCTCTGCACTAAG

SEQ ID NO:17

AATGTACAGTATTGCGTTTTGGCCTTCCCAACATGCATTCTAACTTCTTCC

SEQ ID NO:18

AATGTACAGTATTGCGTTTTGCCAGCTACTCTCAAATCAGCATCCTTTGG

SEQ ID NO:19

AATGTACAGTATTGCGTTTTGCCAGTCCTTCTGTGAGTCTATCCTCAGTTC

SEQ ID NO:20

AATGTACAGTATTGCGTTTTGAGAGCGAACCAAGAATGCCTGTTTACAG

SEQ ID NO:21

AATGTACAGTATTGCGTTTTGGAGAGGCACGAGAACACACATCTATTCTG

SEQ ID NO:22

AATGTACAGTATTGCGTTTTGTTCTCTTCAGAAGTTCCTTCGTCATCCTT

SEQ ID NO:23

AATGTACAGTATTGCGTTTTGTGATGACATGCCCATCACTAAAACAC

SEQ ID NO:24

AATGTACAGTATTGCGTTTTGTGATAGAGACATGATGTAACCGTGGGAATTTCTTC

SEQ ID NO:25

AATGTACAGTATTGCGTTTTGCGTTCTAAGAGAGTGACAGAAAGGTAAAGAGGAG

SEQ ID NO:26

AATGTACAGTATTGCGTTTTGATCACAAAGTATCTTTTTCTGTGGCTTAGAAATCTT

SEQ ID NO:27

AATGTACAGTATTGCGTTTTGTCAAATGTTAGCTCATTTTTGTTAATGGTGGCTTTT

SEQ ID NO:28

AATGTACAGTATTGCGTTTTGTGTCACATTATAAAGATTCAGGCAATGTTTGTTAGT

SEQ ID NO:29

AATGTACAGTATTGCGTTTTGAGTTTGTATGCAACATTTCTAAAGTTACCTACTTGT

SEQ ID NO:30

AATGTACAGTATTGCGTTTTGAAAATCTGTTTTCCAATAAATTCTCAGATCCAGGAA

SEQ ID NO:31

AATGTACAGTATTGCGTTTTGCGACCCAGTTACCATAGCAATTTAGTGAAATAACTA

SEQ ID NO:32

AATGTACAGTATTGCGTTTTGAGAGGCGCTATGTGTATTATTATAGCTACCTGTTAA

SEQ ID NO:33

AATGTACAGTATTGCGTTTTGCGTTTTGACAGTTTACAGTTAAAGGCATTTCC

SEQ ID NO:34

AATGTACAGTATTGCGTTTTGCTGTCCTTATTTTTGGATATTTCTCCCAATGAAAGTA

SEQ ID NO:35
AATGTACAGTATTGCGTTTTGGACTTTTTGCAAATGTTAACATAGGTGACAGATT

SEQ ID NO:36
AATGTACAGTATTGCGTTTTGAAGTAGAAAATGGAAGTCTATGTGATCAAGAAATCG

SEQ ID NO:37
AATGTACAGTATTGCGTTTTGGGCCTCTTAAAGATCATGTTTGTTACAGTGCTTA

SEQ ID NO:38
AATGTACAGTATTGCGTTTTGACAAGATTGGTCAGGAAAAGAGAATTGTTCTATAA

SEQ ID NO:39
AATGTACAGTATTGCGTTTTGAGACCCTGTCTCAAAGTAAAAGTAAGTTAACATG

SEQ ID NO:40
AATGTACAGTATTGCGTTTTGTCAGTGTCTTCCAAATCCTTATGTATAGCAGCAAT

SEQ ID NO:41
AATGTACAGTATTGCGTTTTGAGGGTCGAGGAAGCCAGTTTACATCAA

SEQ ID NO:42
AATGTACAGTATTGCGTTTTGAACAAAAAGATATTTTCAATATTTCTGCGCAGGTTT

SEQ ID NO:43
AATGTACAGTATTGCGTTTTGGTCTCGACTTGAATTGCAAAAAGATGTTAGAAAAGC

SEQ ID NO:44
AATGTACAGTATTGCGTTTTGAAAATGTTGGCAGTCATAACATTTGAAACTAATGGA

SEQ ID NO:45
AATGTACAGTATTGCGTTTTGAGCCTCAAACAGGTTGGTTTTAAATTTGAAGTCT

SEQ ID NO:46
AATGTACAGTATTGCGTTTTGCCTCTGTGTGTATGTTTTAACTACAAAGCGAAACA

SEQ ID NO:47
AATGTACAGTATTGCGTTTTGGATTCACCTGGTAATGAGGAAAACAGCTTTAAAATC

SEQ ID NO:48
AATGTACAGTATTGCGTTTTGAGATCTGCTGAAAAGAAATTTGTTAAAGCACAATT

SEQ ID NO:49
AATGTACAGTATTGCGTTTTGCGGCATCCCCTACATCGAGACCTC

SEQ ID NO:50
AATGTACAGTATTGCGTTTTGCAGGGAGCAGATCAAACGGGTGAAG

SEQ ID NO:51
AATGTACAGTATTGCGTTTTGCAAGTCTTTTGAGGACATCCACCAGTACAG

SEQ ID NO:52
AATGTACAGTATTGCGTTTTGACGTGCCTGTTGGACATCCTGGATA

SEQ ID NO:53

AATGTACAGTATTGCGTTTTGCCTGTACTGGTGGATGTCCTCAAAGACT

SEQ ID NO:54

AATGTACAGTATTGCGTTTTGCCCTGAGGAGCGATGACGGAATATAAGC

SEQ ID NO:55

AATGTACAGTATTGCGTTTTGGTCGTATTCGTCCACAAAATGGTTCTGGATC

SEQ ID NO:56

AATGTACAGTATTGCGTTTTGTGACTGGCAATTGTGTCAACAGGTGAAAA

SEQ ID NO:57

AATGTACAGTATTGCGTTTTGCGCCAGCTGGAGTTTGGTCATGTTT

SEQ ID NO:58

AATGTACAGTATTGCGTTTTGAATCCCTCTCATCACAATTTTCATTCCACAATAGTTT

SEQ ID NO:59

AATGTACAGTATTGCGTTTTGTCAACAACAAAGAGAATCATGAAATCAACCCTAGC

SEQ ID NO:60

AATGTACAGTATTGCGTTTTGGATATGGAGCCAGCGTGTTCGATT

SEQ ID NO:61

AATGTACAGTATTGCGTTTTGGGCGCGGAAAGTCCTCACTCTC

SEQ ID NO:62

AATGTACAGTATTGCGTTTTGTATGGTGAGGTTTCGGCGTGTTTAAACG

SEQ ID NO:63

AATGTACAGTATTGCGTTTTGTGGTGACAAAGTTAGAAGGGTCCATGG

SEQ ID NO:64

AATGTACAGTATTGCGTTTTGCTTCTTTACCACCCAGATACGACGACTA

SEQ ID NO:65

AATGTACAGTATTGCGTTTTGCGCTCGTGGTGGTAGTCGTTCGTAT

SEQ ID NO:66

AATGTACAGTATTGCGTTTTGCCAGGAGGCCCTTTCTGTTTACAACC

SEQ ID NO:67

AATGTACAGTATTGCGTTTTGCCACAAGCCCAAATATTCTACTCACTTTGC

SEQ ID NO:68

AATGTACAGTATTGCGTTTTGATCGCCTGCATCAAGGAAAAGGTAATGG

SEQ ID NO:69

AATGTACAGTATTGCGTTTTGCGCGTAAGGATAGCAACTGAGGTTATCAC

SEQ ID NO:70

AATGTACAGTATTGCGTTTTGCGACCTGACGTAACCCCTTGCTTATC

SEQ ID NO:71

AATGTACAGTATTGCGTTTTGGGAAATGCTCTCACGTAGTCTCTCATGTCT

SEQ ID NO:72

AATGTACAGTATTGCGTTTTGGTCATAACCCGAAGAACAATGTTGCCACTA

SEQ ID NO:73

AATGTACAGTATTGCGTTTTGGTCAGCTCAGGATAAAGCACGGATGGATA

SEQ ID NO:74

AATGTACAGTATTGCGTTTTGCTCAGGATAAAAGCTTCCTTCTTAACAAGTTTTTCC

SEQ ID NO:75

AATGTACAGTATTGCGTTTTGAGAGATTGTTCCCTTGCATTGACCTCTTTTTTC

SEQ ID NO:76

AATGTACAGTATTGCGTTTTGCCCTCACCTTTGGAATTTACAGTCTGAA

SEQ ID NO:77

AATGTACAGTATTGCGTTTTGTAGGTTCTTCAGGTCTCTACACTCTCCTTTAAACT

SEQ ID NO:78

AATGTACAGTATTGCGTTTTGGAGAAGGAGTGCAATGCCAAGATTATGATCC

SEQ ID NO:79

AATGTACAGTATTGCGTTTTGGACGTTCTCCATTGTATTGGCAGTAACCA

SEQ ID NO:80

AATGTACAGTATTGCGTTTTGCACATCTCACAGGCTCTAAAGGAATTCTATATCCTA

SEQ ID NO:81

AATGTACAGTATTGCGTTTTGGAGGCAAGAGGTGAGTAGTACCAATACTGTC

SEQ ID NO:82

AATGTACAGTATTGCGTTTTGGAGCCCCTCCGCTTACTTGTAATCTG

SEQ ID NO:83

AATGTACAGTATTGCGTTTTGCCAGTAAAACGTATTGAGAAAAAGGTAAAAGCGTTA

SEQ ID NO:84

AATGTACAGTATTGCGTTTTGGCTCAGAATAAATCGTAACAATCTCAAAGTGCATTT

SEQ ID NO:85

AATGTACAGTATTGCGTTTTGTGAGGTGTCCACAGGGCTCAATCTTTAC

SEQ ID NO:86

AATGTACAGTATTGCGTTTTGCCCTTGTATCAGTAAAGGCTATATAATACCGAATT

SEQ ID NO:87

AATGTACAGTATTGCGTTTTGTCATGAAGAGAGTATCATCAGCTCGTTCATCATC

SEQ ID NO:88

AATGTACAGTATTGCGTTTTGTGTCCTTTCTGCCGATGTGAAATTAAGGTAC

SEQ ID NO:89

AATGTACAGTATTGCGTTTTGTGCGCCCAAATAATTCCTGCGAACA

SEQ ID NO:90

AATGTACAGTATTGCGTTTTGCTCATACCTCCATTCCAAGCTTTCATTGTCTC

SEQ ID NO:91

AATGTACAGTATTGCGTTTTGCCTGCCCTTATTTTTAACAGCAGGAACGAAT

SEQ ID NO:92

AATGTACAGTATTGCGTTTTGTGATAGCGAAAGTCCTCTTTGGTCAG

SEQ ID NO:93

AATGTACAGTATTGCGTTTTGGTTAAAGACCAACCACTAACTAAGAGACTTCCAAG

SEQ ID NO:94

AATGTACAGTATTGCGTTTTGAAACCTCTTCCAGTACCTTCTTCATGGTCT

SEQ ID NO:95

AATGTACAGTATTGCGTTTTGTTTCCAGGTGATGTGCTCTATGAACTCCTT

SEQ ID NO:96

AATGTACAGTATTGCGTTTTGGGAGCGGTGCAACAGTTCAATGGT

SEQ ID NO:97

AATGTACAGTATTGCGTTTTGCATCCGTGGATAATGTGCACCATAACC

SEQ ID NO:98

AATGTACAGTATTGCGTTTTGTGCGGAGAGCCTGGACTGTTTGAAATC

SEQ ID NO:99

AATGTACAGTATTGCGTTTTGAAGCCAGGTCTTCCCGATGAGAGAG

SEQ ID NO:100

AATGTACAGTATTGCGTTTTGGGCACTCCGTGGATTTCAAACAGTC

SEQ ID NO:101

AATGTACAGTATTGCGTTTTGCAGATATCTGCTGCCCTTTACCTTATGGTTT

SEQ ID NO:102

AATGTACAGTATTGCGTTTTGTGTAGACTGCTTTGGGATTACGTCTATCAGTTG

SEQ ID NO:103

AATGTACAGTATTGCGTTTTGGGAAAGGAGAAAAGGAAGTGCTACCTGAAC

SEQ ID NO:104

AATGTACAGTATTGCGTTTTGTTTTTCTCCCTTCTCCTTTGAACAAACAG

SEQ ID NO:105

AATGTACAGTATTGCGTTTTGACAGCTTTAGGAAAATGGAATCTCTTACCTCCTC

SEQ ID NO:106

AATGTACAGTATTGCGTTTTGGGGTGTATGGTCGCGTTGGATTTCTG

SEQ ID NO:107

AATGTACAGTATTGCGTTTTGGCTACGGCGTGCAACTCACAGAAC

SEQ ID NO:108

AATGTACAGTATTGCGTTTTGACCGACCTCTTCCAGCGCTACTT

SEQ ID NO:109

AATGTACAGTATTGCGTTTTGCGGGCAGGGCTTACTTACCTTGG

SEQ ID NO:110

AATGTACAGTATTGCGTTTTGTAGCTACTGCCTGCCTTCGAAGAACGAT

SEQ ID NO:111

AATGTACAGTATTGCGTTTTGTGTGGGTGGAAAAAGATGTGGTTAAGAAACAAC

SEQ ID NO:112

AATGTACAGTATTGCGTTTTGCCCCATATAGCTTAATCTGATGGGCATC

SEQ ID NO:113

AATGTACAGTATTGCGTTTTGGAAAGAGCATCAGGAACAAGCCTTGAGTAC

SEQ ID NO:114

AATGTACAGTATTGCGTTTTGTTGAGATGCCTGACAACCTTTACACCTTTG

SEQ ID NO:115

AATGTACAGTATTGCGTTTTGCTCTAGGGCTGAGGGAATATGCATCTCT

SEQ ID NO:116

AATGTACAGTATTGCGTTTTGCGTACCCAGAAGACAATGGCCTAGCTAT

SEQ ID NO:117

AATGTACAGTATTGCGTTTTGGGGCAGCACAGATTCCCTTAACCA

SEQ ID NO:118

AATGTACAGTATTGCGTTTTGCCATACCTTGGCTATCCCCTGAAAGTTG

SEQ ID NO:119

AATGTACAGTATTGCGTTTTGGCCCTGATGCTCATGGAGTGTTCTT

SEQ ID NO:120

AATGTACAGTATTGCGTTTTGCCTGGTGGTTGGGAGACGACTAC

SEQ ID NO:121

AATGTACAGTATTGCGTTTTGTGCTGACAGGACACAGAACAAGATACCT

SEQ ID NO:122

AATGTACAGTATTGCGTTTTGGGTACAGGTATCTTGTTCTGTGTCCTGTCAG

SEQ ID NO:123

AATGTACAGTATTGCGTTTTGGAGTCCCGGGCTCGATTACAG

SEQ ID NO:124

AATGTACAGTATTGCGTTTTGCTGGTCAGAGAGGTGTGTA CTGATTGTCT

SEQ ID NO:125
AATGTACAGTATTGCGTTTTGAGGAAAGATCAATTACATTCACAAGTTCACACTTCT

SEQ ID NO:126
AATGTACAGTATTGCGTTTTGCTGCACAGTTCAGAGGATATTTAAGCTCAATGAC

SEQ ID NO:127
AATGTACAGTATTGCGTTTTGCACAGACCGTCATGCATTTCTGACACTC

SEQ ID NO:128
AATGTACAGTATTGCGTTTTGAGGCTGGTACCTGCTCTTCTTCAATC

SEQ ID NO:129
AATGTACAGTATTGCGTTTTGCGAAATCAAACAGTTGTCTATCAGAGCCTGTC

SEQ ID NO:130
AATGTACAGTATTGCGTTTTGACAAAAGAAAAGAAGTCATGTCTGTATGTGGAAA

SEQ ID NO:131
AATGTACAGTATTGCGTTTTGTCCAGGATAATACACATCACAGTAAATAACACTCTG

SEQ ID NO:132
AATGTACAGTATTGCGTTTTGCATCCTCTTTGTCATCAAGCTACAGTCTTTTTGA

SEQ ID NO:133
AATGTACAGTATTGCGTTTTGCTCCCATTTTTGTGCATCTTTGTTGCTGTC

SEQ ID NO:134
AATGTACAGTATTGCGTTTTGCAGAACTGCCTATTCCTAACTGACTCATCATTC

SEQ ID NO:135
AATGTACAGTATTGCGTTTTGGAATTCTGTTTCATCGCTGAGTGACACTCTTTT

SEQ ID NO:136
AATGTACAGTATTGCGTTTTGTTTTTACCTTTGCTTTTACCTTTTTGTACTIONGTGAC

SEQ ID NO:137
AATGTACAGTATTGCGTTTTGAGAAGGAGTCTGGAATAGAAAGGCTAACAGAA

SEQ ID NO:138
AATGTACAGTATTGCGTTTTGCACAAGATGTGCCAAGGGAATTGTATGC

SEQ ID NO:139
AATGTACAGTATTGCGTTTTGAAGAGTCAATAGGTCAGAGAGTTTTATGTTCTTCCA

SEQ ID NO:140
AATGTACAGTATTGCGTTTTGACTGATCTTCTCAAAGTCGTCATCCTTCAGT

SEQ ID NO:141
AATGTACAGTATTGCGTTTTGACCCTGAGAAATAATCCAATTACCTGTTAATCAAGG

SEQ ID NO:142
AATGTACAGTATTGCGTTTTGAAAAGGTATTGAGTAAAATCAGTCTTCCTTCTACCC

SEQ ID NO:143
AATGTACAGTATTGCGTTTTGCCTTCCTCCCTCTTTCTTTCATAAAACCTCTCTT

SEQ ID NO:144
AATGTACAGTATTGCGTTTTGGCCAGAGCCACCCAACCTCTTAAGG

SEQ ID NO:145
AATGTACAGTATTGCGTTTTGTGGAAGAGGAATTTAATAACGAACGTTTTAAGAGGA

SEQ ID NO:146
AATGTACAGTATTGCGTTTTGGCATCTACTGCCGAGGATGTTCCAAG

SEQ ID NO:147
AATGTACAGTATTGCGTTTTGCACAGTGAGCTCAAGTGCGACATCA

SEQ ID NO:148
AATGTACAGTATTGCGTTTTGCCGACTGGCCATCTCCTCGTAG

SEQ ID NO:149
AATGTACAGTATTGCGTTTTGGTACCAGCGCGACTACGAGGAGAT

SEQ ID NO:150
AATGTACAGTATTGCGTTTTGTCTTTTCTGTCAAATGGAGATGATCTCTTCTGACTC

SEQ ID NO:151
AATGTACAGTATTGCGTTTTGGGGAGCCCATCATCTGCAAAAACATCC

SEQ ID NO:152
AATGTACAGTATTGCGTTTTGAAGCTGAAGAAGATGTGGAAAAGTCCCAATG

SEQ ID NO:153
AATGTACAGTATTGCGTTTTGGCGTGGGATGTTTTTGCAGATGATGG

SEQ ID NO:154
AATGTACAGTATTGCGTTTTGCGACGCTGAGGACGCTATGGATG

SEQ ID NO:155
AATGTACAGTATTGCGTTTTGGCTGAGGCGCGTCTTCGAGAAG

SEQ ID NO:156
AATGTACAGTATTGCGTTTTGGCGCTTGTCGTGAAAGCGAACGA

SEQ ID NO:157
AATGTACAGTATTGCGTTTTGGCTGCCCGCCCAGTTGTTACT

SEQ ID NO:158
AATGTACAGTATTGCGTTTTGAGACTCTGGACTGATGAAGCAATTCTGAGT

SEQ ID NO:159
AATGTACAGTATTGCGTTTTGTCACCGGTGACACCTTAAAACCAAAGC

SEQ ID NO:160
AATGTACAGTATTGCGTTTTGGGCTCCTTTGTACCTCCTCCATCTTGATC

SEQ ID NO:161
AATGTACAGTATTGCGTTTTGGTCAGTTGTCTAACAATAACAAAGATCTGCTCTTGG

SEQ ID NO:162
AATGTACAGTATTGCGTTTTGGGTGGGCAGCAAGAAAAAGTCCAGTAAA

SEQ ID NO:163
AATGTACAGTATTGCGTTTTGGCCAAGGCTTCTCTGGCATGATCTTTT

SEQ ID NO:164
AATGTACAGTATTGCGTTTTGGGATAACTTTCTCAGCATTCCACCAGTTTCAAG

SEQ ID NO:165
AATGTACAGTATTGCGTTTTGTGTCCCTAAGTTGAGTAAAATGATAGAGAATGAGTC

SEQ ID NO:166
AATGTACAGTATTGCGTTTTGGCTGCCAGAAATCCAGCATCCAAAATTTG

SEQ ID NO:167
AATGTACAGTATTGCGTTTTGGTCGCTTCTTTTCTTAGTGCCAGGAACT

SEQ ID NO:168
AATGTACAGTATTGCGTTTTGACAGTCGAGACGATTCATGAGGGAACTTC

SEQ ID NO:169
AATGTACAGTATTGCGTTTTGGGAAAGCTCGGCGTGTGGATAAGAAG

SEQ ID NO:170
AATGTACAGTATTGCGTTTTGACGCCACAAGTGACTGAAAGTTGGAAG

SEQ ID NO:171
AATGTACAGTATTGCGTTTTGTGATGGGCTGGAGATTTGGCATAGTTTTTC

SEQ ID NO:172
AATGTACAGTATTGCGTTTTGCTATGCACCCACTTTCAACACAGTTAGGT

SEQ ID NO:173
AATGTACAGTATTGCGTTTTGGCTTGGTCAGAAGTGCTGTTGTTGTC

SEQ ID NO:174
AATGTACAGTATTGCGTTTTGCGTGGGCCAGAAAGTTGTCCACAATG

SEQ ID NO:175
AATGTACAGTATTGCGTTTTGGGGATATGGATTCTCGTGGTAGAAGGTGTAA

SEQ ID NO:176
AATGTACAGTATTGCGTTTTGCTAATCACCAAGTTCCAAGTGTTGAGAATCTCC

SEQ ID NO:177
AATGTACAGTATTGCGTTTTGACCGTAATAACCAAGTTTCATCATAGGCATTGAT

SEQ ID NO:178
AATGTACAGTATTGCGTTTTGTCCCAGTGGAAGTTACTATGCACCCTAT

SEQ ID NO:179

AATGTACAGTATTGCGTTTTGTGCTTATGCTTGTGTTTGTGTTTCCTCTTATGG

SEQ ID NO:180

AATGTACAGTATTGCGTTTTGGCTTCTGTTTCTCCTTATGCTTGTTCCTTCTCAC

SEQ ID NO:181

AATGTACAGTATTGCGTTTTGCCTGAGTGGTCTTTTTGCAGGCAAAG

SEQ ID NO:182

AATGTACAGTATTGCGTTTTGCCGGCCACAAAGCTTCTAAGAACAAC

SEQ ID NO:183

AATGTACAGTATTGCGTTTTGGCGGTTTCATCTTGAAGGCTTGGATGT

SEQ ID NO:184

AATGTACAGTATTGCGTTTTGTTTCAGTGAAATGAACCCTTCGAATGACAAG

SEQ ID NO:185

AATGTACAGTATTGCGTTTTGCTCCTCCTCCTCTTTGCGTTTCTTGTC

SEQ ID NO:186

AATGTACAGTATTGCGTTTTGGCAGCAGAGAAACAAATGAAGGACAAACAG

SEQ ID NO:187

AATGTACAGTATTGCGTTTTGTAAGGAGGAGGAAGAAGACAAGAAACGCAA

SEQ ID NO:188

AATGTACAGTATTGCGTTTTGTAAGGCAGGTCTGTGAGCACAAAATTTGG

SEQ ID NO:189

AATGTACAGTATTGCGTTTTGTGGAGCTGACCAGTGACAATGACC

SEQ ID NO:190

AATGTACAGTATTGCGTTTTGGGCCAAGAAGTCGGTGGACAAGAAC

SEQ ID NO:191

AATGTACAGTATTGCGTTTTGGCGCAGGCGGTCATTGTCACTG

SEQ ID NO:192

AATGTACAGTATTGCGTTTTGTTGCTGTTCTTGTCCACCGACTTCTTG

SEQ ID NO:193

AATGTACAGTATTGCGTTTTGGCAGTGCGCGATCTGGAAGTCTG

SEQ ID NO:194

AATGTACAGTATTGCGTTTTGCGGCGGCGACTTTGACTACCC

SEQ ID NO:195

AATGTACAGTATTGCGTTTTGGAGCACGAGACGTCCATCGACATC

SEQ ID NO:196

AATGTACAGTATTGCGTTTTGCGGCCAGGAACTCGTCGTTGAA

SEQ ID NO:197

AATGTACAGTATTGCGTTTTGGCCATGCCGGGAGAACTCTAACTC

SEQ ID NO:198

AATGTACAGTATTGCGTTTTGTGTAACCCTCCTAAGTGTTACATACGTTGTCTTG

SEQ ID NO:199

AATGTACAGTATTGCGTTTTGGTCTTGGTCTCTGTTATATCTTGAGTCTAGAACAGT

SEQ ID NO:200

AATGTACAGTATTGCGTTTTGCAGGAGAACATGGAGGCGAGAAGAAAAT

SEQ ID NO:201

AATGTACAGTATTGCGTTTTGGGGAAAGATTGGATGCCGGGAATCAAC

SEQ ID NO:202

AATGTACAGTATTGCGTTTTGCGGAGGCTTGATTAGGTAGGAGGTG

SEQ ID NO:203

AATGTACAGTATTGCGTTTTGGCGGCAGCTCAACGAGAATAAACA

SEQ ID NO:204

AATGTACAGTATTGCGTTTTGGCCCGCATCCTTACTCCGCTTATC

SEQ ID NO:205

AATGTACAGTATTGCGTTTTGGCTGGTTTCAAGGTAAGTGGACTCTTCC

SEQ ID NO:206

AATGTACAGTATTGCGTTTTGGGGAATGACTGACGGAGAATCCCAAC

SEQ ID NO:207

AATGTACAGTATTGCGTTTTGCTAAGACCGAGAGCCTGTAGGAGCTTT

SEQ ID NO:208

AATGTACAGTATTGCGTTTTGGCCGGGCTTGTCTGGTCATCT

SEQ ID NO:209

AATGTACAGTATTGCGTTTTGCAGCTCACCTCCAAAAAGGCAAATCTTG

SEQ ID NO:210

AATGTACAGTATTGCGTTTTGGCAGGAGGCCATGATGGATTTCTTCAA

SEQ ID NO:211

AATGTACAGTATTGCGTTTTGCATGAGTGAAAGGAAAGAGGAAATCCCAATCC

SEQ ID NO:212

AATGTACAGTATTGCGTTTTGCCTATCTTCCACAGTACTTACACAACCTCCTAAGC

SEQ ID NO:213

AATGTACAGTATTGCGTTTTGCTCGCCGTAGACTGTCCAGGTTTT

SEQ ID NO:214

AATGTACAGTATTGCGTTTTGCTCACCTGATCCGTGACGTTGATGTC

SEQ ID NO:215

AATGTACAGTATTGCGTTTTGGCCCTGATGGACTCTCGGCTACT

SEQ ID NO:216

AATGTACAGTATTGCGTTTTGGAGAAAGATCAGGAACACTTGTCCCCTACTAG

SEQ ID NO:217

AATGTACAGTATTGCGTTTTGGTCCTCCACGATCTCCTCATACTCCTC

SEQ ID NO:218

AATGTACAGTATTGCGTTTTGTCGATGGACTTGACAAGCCCGTACTT

SEQ ID NO:219

AATGTACAGTATTGCGTTTTGCTGGACGACGAGGAGTATGAGGAGATC

SEQ ID NO:220

AATGTACAGTATTGCGTTTTGTACCAGAAGTCCCGGCGGTGATAAG

SEQ ID NO:221

AATGTACAGTATTGCGTTTTGGTTCACCTCTGTGTTTGACTGCCAGAAA

SEQ ID NO:222

AATGTACAGTATTGCGTTTTGCAATGAGTATTCTCTTCATTCAGGTCAGTTGATTT

SEQ ID NO:223

AATGTACAGTATTGCGTTTTGGGCTGCTTTCTTGAAGGCTATTGGGTAT

SEQ ID NO:224

AATGTACAGTATTGCGTTTTGAGGAGACTGGAATTCTCGAATAAGGATTAACA

SEQ ID NO:225

AATGTACAGTATTGCGTTTTGGCATAGTTAAAACCTGTGTTTGGTTTTGTAGGTCTT

SEQ ID NO:226

AATGTACAGTATTGCGTTTTGCTCTGTGTTGGCGGATACCCTTCCATA

SEQ ID NO:227

AATGTACAGTATTGCGTTTTGGGCATTCCTTCTTTATTGCCCTTCTTAAAAGC

SEQ ID NO:228

AATGTACAGTATTGCGTTTTGGCTGCTGGTCTGGCTACTATGATCTCTAC

SEQ ID NO:229

AATGTACAGTATTGCGTTTTGGCACACAGCTTTTAAGAAGGGCAATAAAGAAG

SEQ ID NO:230

AATGTACAGTATTGCGTTTTGTGTATGTTTAAATTCTGTACATGAGCATTTCATCAGT

SEQ ID NO:231

AATGTACAGTATTGCGTTTTGATTCATACCTTGCTTAATGGGTGTAGATACCAAAA

SEQ ID NO:232

AATGTACAGTATTGCGTTTTGTTGGCGTCAAATGTGCCACTATCACTC

SEQ ID NO:233
AATGTACAGTATTGCGTTTTGTTCTCTTTCAAGCTATGATTTAGGCATAGAGAATCG

SEQ ID NO:234
AATGTACAGTATTGCGTTTTGCTGCAGTTGTAGGTTATAACTATCCATTTGTCTGAA

SEQ ID NO:235
AATGTACAGTATTGCGTTTTGCCCTAGGTCAGATCACCCAGTCAGTTAAAAC

SEQ ID NO:236
AATGTACAGTATTGCGTTTTGTGGTTAAAGGTCAGCCCACCTTACCAGATATG

SEQ ID NO:237
AATGTACAGTATTGCGTTTTGGGGTATGCTCCCCATTTAGAGGATAAGG

SEQ ID NO:238
AATGTACAGTATTGCGTTTTGACGTCAGATCTACAGCGAACACA ACTACT

SEQ ID NO:239
AATGTACAGTATTGCGTTTTGAGTGGTGCCAGACTCACATTCAGTTCTAA

SEQ ID NO:240
AATGTACAGTATTGCGTTTTGCTTGCCAGTTCCTTTCTCTAATGTATCATCTC

SEQ ID NO:241
AATGTACAGTATTGCGTTTTGAAGTTTTCTTGTCTAGTATCACTTTCCCTCATAGG

SEQ ID NO:242
AATGTACAGTATTGCGTTTTGGGGCTCAACAGATGGTATGTGTTCTCTG

SEQ ID NO:243
AATGTACAGTATTGCGTTTTGGCTCTCGTTTCTAACAGTTCTTTGCATTGGATA

SEQ ID NO:244
AATGTACAGTATTGCGTTTTGGAGGTGACCTTCAAAGTCAGAGGCTGTAT

SEQ ID NO:245
AATGTACAGTATTGCGTTTTGGAGCAACCATCCCATCTGTCCTTGTAAC

SEQ ID NO:246
AATGTACAGTATTGCGTTTTGGGACAAGGATGAGAAACCCAATTGGAACC

SEQ ID NO:247
AATGTACAGTATTGCGTTTTGCGGTCCGCCAAAAGATCCCAGATTC

SEQ ID NO:248
AATGTACAGTATTGCGTTTTGGGAGGCCACTAACCCTTGTGATG

SEQ ID NO:249
AATGTACAGTATTGCGTTTTGTCCAGTTTCTAGAGGATGTAATGGGATTTGTC

SEQ ID NO:250
AATGTACAGTATTGCGTTTTGTCACATTTGGAGATGAGAAACGAGGTGTTCT

SEQ ID NO:251

AATGTACAGTATTGCGTTTTGCCCCTTGGCCTGTAACATTGCTCTGATC

SEQ ID NO:252

AATGTACAGTATTGCGTTTTGCACCTCGTTTCTCATCTCCAAATGTGATCTC

SEQ ID NO:253

AATGTACAGTATTGCGTTTTGCCAGTAGCTTTCCTGTTCTCGGCATT

SEQ ID NO:254

AATGTACAGTATTGCGTTTTGGCAGCGTCAAGAATGAGAAGACTTTTGTG

SEQ ID NO:255

AATGTACAGTATTGCGTTTTGTTGCCCTTCTGGAAATTACCCCGAGA

SEQ ID NO:256

AATGTACAGTATTGCGTTTTGAGTTCCACCAGCTTTAATTATTCTCTAGCTCTC

SEQ ID NO:257

AATGTACAGTATTGCGTTTTGGTTTCCCATGGCCATAATTTATTATCTCACCACAA

SEQ ID NO:258

AATGTACAGTATTGCGTTTTGGTCACGATGACTGTATTGGACCCTCAA

SEQ ID NO:259

AATGTACAGTATTGCGTTTTGTCCAGACCTTTGCTTTAGATTGGCAATTATTACTG

SEQ ID NO:260

AATGTACAGTATTGCGTTTTGCCCTAACAAACACAGAAGCAAAGCGTTCCTT

SEQ ID NO:261

AATGTACAGTATTGCGTTTTGCGCCCTCCTACCACCTGTACTACG

SEQ ID NO:262

AATGTACAGTATTGCGTTTTGACTATCCAGGCGCCTTCACCTACTC

SEQ ID NO:263

AATGTACAGTATTGCGTTTTGCTCCTAGGCGGTATCATCCTGGGTAG

SEQ ID NO:264

AATGTACAGTATTGCGTTTTGTCTGATTCTCTTCAGATACAAGGCAGATCC

SEQ ID NO:265

AATGTACAGTATTGCGTTTTGGCAGATACTTGGACTTGAGTAGGCTTATTAAACC

SEQ ID NO:266

AATGTACAGTATTGCGTTTTGGCGGCTCTATAAAGAATTGTCCTTATTTTCGAACTT

SEQ ID NO:267

AATGTACAGTATTGCGTTTTGGTTCGAGGCCTTTCTCTGAGCATCAAG

SEQ ID NO:268

AATGTACAGTATTGCGTTTTGACATCGGCAGAACTAGATGATCAGACCAA

SEQ ID NO:269
AATGTACAGTATTGCGTTTTGTTTAGGAAATCCACAATACTTTTTCTGATCTCTCC

SEQ ID NO:270
AATGTACAGTATTGCGTTTTGGCCACCAACCTCATTCTGTTTTGTTCTCTATC

SEQ ID NO:271
AATGTACAGTATTGCGTTTTGCTGCATTTGTCCTTTGACTGGTGTTAGGT

SEQ ID NO:272
AATGTACAGTATTGCGTTTTGCTTCGACCGACAAACCTGAGGTCATTAATC

SEQ ID NO:273
AATGTACAGTATTGCGTTTTGCCCCACATCCCAAGCTAGGAAGACC

SEQ ID NO:274
AATGTACAGTATTGCGTTTTGCGGGCCAGTACCTTGAAAGCGATG

SEQ ID NO:275
AATGTACAGTATTGCGTTTTGCTAACTCAATCGGCTTGTGTGATGCGTAT

SEQ ID NO:276
AATGTACAGTATTGCGTTTTGCCCTCCTGGACTGTTAGTAACTTAGTCTCC

SEQ ID NO:277
AATGTACAGTATTGCGTTTTGCCCTCCGAGCTCCGCGAAAAT

SEQ ID NO:278
AATGTACAGTATTGCGTTTTGGTGCTAAAAGTGTAAGAAGAAATGAGCTAGCAAAA

SEQ ID NO:279
AATGTACAGTATTGCGTTTTGCATATGCCTCAGTTTGAATCCTCTCACAAACAA

SEQ ID NO:280
AATGTACAGTATTGCGTTTTGGGGAGAAGAAAGAGAGATGTAGGGCTAGAG

SEQ ID NO:281
AATGTACAGTATTGCGTTTTGGCAAGCACTTCTGTTTTGTCTTTTCAGTTTCG

SEQ ID NO:282
AATGTACAGTATTGCGTTTTGTCTCTGATATACTTGGATTGGTAATTGAGAAAGTCT

SEQ ID NO:283
AATGTACAGTATTGCGTTTTGGTTTGATATCTTCCCAGCAAATAATCAGCTCTCAT

SEQ ID NO:284
AATGTACAGTATTGCGTTTTGTAGCCAACCTTTTTCGATGAGCTCACTAG

SEQ ID NO:285
AATGTACAGTATTGCGTTTTGTGGAACAGACAAACTATCGACTGAAGTTGT

SEQ ID NO:286
AATGTACAGTATTGCGTTTTGGAGGCTGAGTGCAAATTTGGTCTGGAA

SEQ ID NO:287

AATGTACAGTATTGCGTTTTGGATGGTGGTGGTTGTCTCTGATGATTACC

SEQ ID NO:288

AATGTACAGTATTGCGTTTTGGCAAGGCGAGTCCAGAACCAAGATT

SEQ ID NO:289

AATGTACAGTATTGCGTTTTGTCAGAAGCGACTGATCCCCATCAAGT

SEQ ID NO:290

AATGTACAGTATTGCGTTTTGCATATGGTCACATCACCTTAACTAAACCCATGTTT

SEQ ID NO:291

AATGTACAGTATTGCGTTTTGTTTCTCGGTACTGTTTATTTTGAACAAAACCAATCC

SEQ ID NO:292

AATGTACAGTATTGCGTTTTGCCTCCTCCCCAAATTCCAGGAACAATATGA

SEQ ID NO:293

AATGTACAGTATTGCGTTTTGTGTGCGTCATTTTATTTGGGAAAATTTGATACTAAC

SEQ ID NO:294

AATGTACAGTATTGCGTTTTGCATGCAGGAGAAGTCATCCCCCTTC

SEQ ID NO:295

AATGTACAGTATTGCGTTTTGTCTGAAAACCTGGTGGTTGCCTCTAGGTAA

SEQ ID NO:296

AATGTACAGTATTGCGTTTTGGCCCCTTCTTGCTCTTCTTGGACTIONG

SEQ ID NO:297

AATGTACAGTATTGCGTTTTGCCAAGCCAAGCCAAGCTGGATATTGTG

SEQ ID NO:298

AATGTACAGTATTGCGTTTTGCACTCACATTGTGCAGCTTGTAGTAGAG

SEQ ID NO:299

AATGTACAGTATTGCGTTTTGGCAAAGCGTCTGCATTTGAAGGAGTTT

SEQ ID NO:300

AATGTACAGTATTGCGTTTTGCCCTCCCGAGAACTTGCCGGTTAA

SEQ ID NO:301

AATGTACAGTATTGCGTTTTGGCTCCCCACCACAAAAACGCAAATG

SEQ ID NO:302

AATGTACAGTATTGCGTTTTGGTGTCACTGACGGAGAGCATGAAGATG

SEQ ID NO:303

AATGTACAGTATTGCGTTTTGCCACCCAAAGAAGTGTCTCCTGACC

SEQ ID NO:304

AATGTACAGTATTGCGTTTTGTCCGTCAGTGACACCTGGTACTTGAC

SEQ ID NO:305

AATGTACAGTATTGCGTTTTGCCCCTAGCTCTGCCTACCCTGATCTTTC

SEQ ID NO:306

AATGTACAGTATTGCGTTTTGACGAGGTGGACGTCTTCTTCAATCAC

SEQ ID NO:307

AATGTACAGTATTGCGTTTTGGCCCTGCGAGTCGAGGTGATTG

SEQ ID NO:308

AATGTACAGTATTGCGTTTTGCCATGACTCTCAGGAATTGGCCCTATACTTAG

SEQ ID NO:309

AATGTACAGTATTGCGTTTTGCTTGGGACCTTCATTTCTATATAACCCCTATCTGG

SEQ ID NO:310

AATGTACAGTATTGCGTTTTGTGCCAGGAACTTTTCATTGTGCCTCTC

SEQ ID NO:311

AATGTACAGTATTGCGTTTTGGTTACCCCATGGAACCTTACCAAGCACTAG

SEQ ID NO:312

AATGTACAGTATTGCGTTTTGGTATGAAATTCGCTGGAGGGTCATTGAATCAAT

SEQ ID NO:313

AATGTACAGTATTGCGTTTTGCAGGAAGGAGCACTTACGTTTTAGCATCTTC

SEQ ID NO:314

AATGTACAGTATTGCGTTTTGGATTTTGAGAAATTCCCTTAATATCCCCATGCTCAA

SEQ ID NO:315

AATGTACAGTATTGCGTTTTGCACAACCACATGTGTCCAGTGAAAATCC

SEQ ID NO:316

AATGTACAGTATTGCGTTTTGTGCTTTCATCAGCAGGGTTCAATCCAAA

SEQ ID NO:317

AATGTACAGTATTGCGTTTTGCATTTACATCATCACAGAGTATTGCTTCTATGGAGA

SEQ ID NO:318

AATGTACAGTATTGCGTTTTGGTGATCTCTGGATGTCGGAATATTTAGAAACCTCT

SEQ ID NO:319

AATGTACAGTATTGCGTTTTGATCTTTTGAAAACAATGGTGACTACATGGACATGAA

SEQ ID NO:320

AATGTACAGTATTGCGTTTTGGGTCTAAAAGGTCTGTGTTCCCTGAACTTACA

SEQ ID NO:321

AATGTACAGTATTGCGTTTTGCCAGCACCAATACATTTAATTTCTTTTCTGCAGAC

SEQ ID NO:322

AATGTACAGTATTGCGTTTTGGCTACAGATGGCTTGATCCTGAGTCATTTTC

SEQ ID NO:323

AATGTACAGTATTGCGTTTTGGTCAGGCCCATACCAAGGGAAAAGATC

SEQ ID NO:324

AATGTACAGTATTGCGTTTTGACACTGAGTGATGTCTGGTCTTATGGCATT

SEQ ID NO:325

AATGTACAGTATTGCGTTTTGCACTGAGCGTTTGTTAGTCCTGGTGTTTT

SEQ ID NO:326

AATGTACAGTATTGCGTTTTGCAGATTCTCCACAATCTCACTCAGGTGGTAAA

SEQ ID NO:327

AATGTACAGTATTGCGTTTTGCCCCACAGCTACGAGATCATGGTGAAAT

SEQ ID NO:328

AATGTACAGTATTGCGTTTTGTCTCTATTCATTTTTGAGGTTTGGTTGTTAACTT

SEQ ID NO:329

AATGTACAGTATTGCGTTTTGGGGAGTGCACCATTATCGGGAAAATGG

SEQ ID NO:330

AATGTACAGTATTGCGTTTTGGCTTATTCTCATTCGTTTCATCCAGGATCTCAAAA

SEQ ID NO:331

AATGTACAGTATTGCGTTTTGGGGCGACGAGATTAGGCTGTTATGC

SEQ ID NO:332

AATGTACAGTATTGCGTTTTGCCCTCTGCATTATAAGCAGTGCCAAAA

SEQ ID NO:333

AATGTACAGTATTGCGTTTTGGCCACATCGTTGTAAGCCTTACATTCAA

SEQ ID NO:334

AATGTACAGTATTGCGTTTTGCCGTTTGAAAGCTAGTGGTTCAGAGTTC

SEQ ID NO:335

AATGTACAGTATTGCGTTTTGGAGATCCCATCCTGCCAAAGTTTGTGATT

SEQ ID NO:336

AATGTACAGTATTGCGTTTTGGGAAAGCCCCTGTTTCATACTGACCAAAA

SEQ ID NO:337

AATGTACAGTATTGCGTTTTGCTTTCTCCCCACAGAAACCCATGTATGAAG

SEQ ID NO:338

AATGTACAGTATTGCGTTTTGGTTTGCCAGTTGTGCTTTTTGCTAAAATGC

SEQ ID NO:339

AATGTACAGTATTGCGTTTTGCCCTCCCACCCTCAGGACTATACCAAT

SEQ ID NO:340

AATGTACAGTATTGCGTTTTGTGCTCGGCAGATTGGTATAGTCCTG

SEQ ID NO:341

AATGTACAGTATTGCGTTTTGGGCATCCTCTGTCCTATCTCCCAGATACA

SEQ ID NO:342

AATGTACAGTATTGCGTTTTGAGGTTTTATACTAAACTTACTTTGACTGGGTTTGG

SEQ ID NO:343

AATGTACAGTATTGCGTTTTGCCCCAGAGGTAAGCGTCATATGG

SEQ ID NO:344

AATGTACAGTATTGCGTTTTGGCACAGGGAAGTAGGTACTGGGAGATTG

SEQ ID NO:345

AATGTACAGTATTGCGTTTTGAGGCCTGCAAGGTTTTAACTGGACCTA

SEQ ID NO:346

AATGTACAGTATTGCGTTTTGCGGGAGCTGATAAGTGGTACCTGTATGT

SEQ ID NO:347

AATGTACAGTATTGCGTTTTGGAAAAGGGTCCCAGGTAGGTCCAGTTAA

SEQ ID NO:348

AATGTACAGTATTGCGTTTTGCTCTCGGTGTATTTCTCTACTTACCTGTAATAATGC

SEQ ID NO:349

AATGTACAGTATTGCGTTTTGTTTATTGATGTCTATGAAGTGTGTGGTTCCTAAC

SEQ ID NO:350

AATGTACAGTATTGCGTTTTGCAGAAAACAAGCTGCCGCAAAGTTCTAC

SEQ ID NO:351

AATGTACAGTATTGCGTTTTGCAGGTGTTGCGATGATGTCACTGTACG

SEQ ID NO:352

AATGTACAGTATTGCGTTTTGTCATTTTTTCATTGGACTTGTTTTGTCAGCTTTTTGG

SEQ ID NO:353

AATGTACAGTATTGCGTTTTGGTTAGCCCCAATATGAAAAATAAAGCTGGTTGGA

SEQ ID NO:354

AATGTACAGTATTGCGTTTTGCTGGTTGGAGGTTTTTGCTAAATCTGGAATGA

SEQ ID NO:355

AATGTACAGTATTGCGTTTTGTTCTTTTTGACTAGAAAACCTCAGCCACTGTGTATT

SEQ ID NO:356

AATGTACAGTATTGCGTTTTGCATATGACCAATTGCAGATGAGCCCATTATTGAA

SEQ ID NO:357

AATGTACAGTATTGCGTTTTGAGGCATAGCTGACTCATCTATGTTTGTCT

SEQ ID NO:358

AATGTACAGTATTGCGTTTTGTTCCCTCATTTCTTTCACTCTGACAGTATAAAGGTAA

SEQ ID NO:359
AATGTACAGTATTGCGTTTTGGAAC TATTCCAACAGAACAAACCGATAACATCA

SEQ ID NO:360
AATGTACAGTATTGCGTTTTGTGGATAGCAAGACAATTAGAGCCCAACTTAGT

SEQ ID NO:361
AATGTACAGTATTGCGTTTTGCTACTCCTCCTGTCTCTTTCCACATCATCAATT

SEQ ID NO:362
AATGTACAGTATTGCGTTTTGAGGACCTTATGTTGTATGCTGTATAAATCTAAAGGT

SEQ ID NO:363
AATGTACAGTATTGCGTTTTGGTTTGT CATCTTCTATGGTAAGTATCTTTCTGGATG

SEQ ID NO:364
AATGTACAGTATTGCGTTTTGTGGAGGAGAAACAGATAAAAGTTGAGTATACGTTTA

SEQ ID NO:365
AATGTACAGTATTGCGTTTTGGAGGATGACGACATGTTAGTAAGCACTACTACT

SEQ ID NO:366
AATGTACAGTATTGCGTTTTGATTCCACCATCATTTCTTCTCCAAAATTATCATCC

SEQ ID NO:367
AATGTACAGTATTGCGTTTTGCTCAA AAGCACTGCCTTCTCTCATTATCTCAC

SEQ ID NO:368
AATGTACAGTATTGCGTTTTGAATGTATTTGACCTTCTTTTAAAGTGACATCGATGT

SEQ ID NO:369
AATGTACAGTATTGCGTTTTGTGATGTTCCCAACTTCTTCTCTCATGGTTATCTC

SEQ ID NO:370
AATGTACAGTATTGCGTTTTGCCCTCTGATCCCTAGATAATTTATGGGTAGCTAGA

SEQ ID NO:371
AATGTACAGTATTGCGTTTTGCACGAAATGCAGGTTTTGGAATATGATTAATGTT

SEQ ID NO:372
AATGTACAGTATTGCGTTTTGGAACAATGTTCTACGCACATTTTGTCTCAGTAAA

SEQ ID NO:373
AATGTACAGTATTGCGTTTTGTCCACGCTGCTCTCTAAATTACACTCGAA

SEQ ID NO:374
AATGTACAGTATTGCGTTTTGACGTAGAACACATTTTACTCCTCTTTGG

SEQ ID NO:375
AATGTACAGTATTGCGTTTTGGTCACATGAATGTAAATCAAGAAAACAGATGTTGTT

SEQ ID NO:376
AATGTACAGTATTGCGTTTTGTTCTGAACTATTTATGGACAAACAGTCAAACAACAAT

SEQ ID NO:377

AATGTACAGTATTGCGTTTTGTGAAGCCATTGCGAGAACTTTATCCATAAGTATTTTC

SEQ ID NO:378

AATGTACAGTATTGCGTTTTGGCCAGAGCACATGAATAAATGAGCATCCAT

SEQ ID NO:379

AATGTACAGTATTGCGTTTTGGGAAGCTCTCAGGGTACAAATTCTCAGATCAT

SEQ ID NO:380

AATGTACAGTATTGCGTTTTGCTCAGGGTACAAATTCTCAGATCATCAGTCCTC

SEQ ID NO:381

AATGTACAGTATTGCGTTTTGCTCTACACAAGCTTCCTTTCCGTCATGC

SEQ ID NO:382

AATGTACAGTATTGCGTTTTGCCCTTCAGATCTTCTCAGCATTGAGAGATC

SEQ ID NO:383

AATGTACAGTATTGCGTTTTGAATCGAAGCGCTACCTGATTCCAATTCC

SEQ ID NO:384

AATGTACAGTATTGCGTTTTGCCGACCGTAACTATTCGGTGCGTTG

SEQ ID NO:385

AATGTACAGTATTGCGTTTTGACATTCTATCCAAGCTGTGTTCTATCTTGAGAACT

SEQ ID NO:386

AATGTACAGTATTGCGTTTTGCGAGTGAGGGTTTTCGTGGTTCACATC

SEQ ID NO:387

AATGTACAGTATTGCGTTTTGCGTGGGTCCCAGTCTGCAGTTAAG

SEQ ID NO:388

AATGTACAGTATTGCGTTTTGGCTCAGAGCCGTTCCGAGATCTT

SEQ ID NO:389

AATGTACAGTATTGCGTTTTGGCGTTCCATCTCCCACTTGTCGTAGTT

SEQ ID NO:390

AATGTACAGTATTGCGTTTTGCTGGCCGAGTTGGTTCATCATCATTCAA

SEQ ID NO:391

AATGTACAGTATTGCGTTTTGTATGGTGTGTCCCCCAACTACGACAAG

SEQ ID NO:392

AATGTACAGTATTGCGTTTTGTGAAAAGCACTTCCTGAAATAATTCACCTTCGTTT

SEQ ID NO:393

AATGTACAGTATTGCGTTTTGAGGTACTCCATGGCTGACGAGATCTG

SEQ ID NO:394

AATGTACAGTATTGCGTTTTGTTGCCTTTGTTCCAAGGTCCAATGTGT

SEQ ID NO:395

AATGTACAGTATTGCGTTTTTGGCGCCCCGCATTCCAACGTCTC

SEQ ID NO:396

AATGTACAGTATTGCGTTTTTGGGCGCGCCGTTTACTTGAAGG

SEQ ID NO:397

AATGTACAGTATTGCGTTTTTGGCCTGGCGGTGCACACTATTCTG

SEQ ID NO:398

AATGTACAGTATTGCGTTTTTGGAGGTGCAGCCACAAAACCTTACAGATGC

SEQ ID NO:399

AATGTACAGTATTGCGTTTTTGGTGCCGAACCAATACAACCCTCTG

SEQ ID NO:400

AATGTACAGTATTGCGTTTTTGGGGCGGGTCCACCAGTTTGAAT

SEQ ID NO:401

AATGTACAGTATTGCGTTTTTGCCGCAGAGGGTTGTATTGGTTTCG

SEQ ID NO:402

AATGTACAGTATTGCGTTTTTGGAGCCACTCGCATTGACCATTCAAACCT

SEQ ID NO:403

AATGTACAGTATTGCGTTTTTGGCACGTCTGACAGGTAGCCATGG

SEQ ID NO:404

AATGTACAGTATTGCGTTTTTGGTGAGGCTGCTGGACGAGTACAAC

SEQ ID NO:405

AATGTACAGTATTGCGTTTTTGGCGACCAGGTTGTAAGTCTGTTCA

SEQ ID NO:406

AATGTACAGTATTGCGTTTTTGGCGCCTTTGTGCTTCTGTTCTTCGT

SEQ ID NO:407

AATGTACAGTATTGCGTTTTTGGCTGATTAATCGCGTAGAAAATGACCTTATTTGGAG

SEQ ID NO:408

AATGTACAGTATTGCGTTTTTGGCTCCATCGTCTACCTGGAGATTGACAA

SEQ ID NO:409

AATGTACAGTATTGCGTTTTTGTCTGCACGGCCTCGATCTTGTAGG

SEQ ID NO:410

AATGTACAGTATTGCGTTTTTGGCCAGCAGATGATCTTCCCCTACTACG

SEQ ID NO:411

AATGTACAGTATTGCGTTTTTGGCTCACGCTTGAAGACCACGTTG

SEQ ID NO:412

AATGTACAGTATTGCGTTTTTGGCCAGCATGCAGTTCTAAGGCTCT

SEQ ID NO:413

AATGTACAGTATTGCGTTTTGGTGCCCGTCTCGACTCTTAGGC

SEQ ID NO:414

AATGTACAGTATTGCGTTTTGTGTAGCCGCTGATCGTCGTGTATATGTC

SEQ ID NO:415

AATGTACAGTATTGCGTTTTGGACTGGTACTGGTTAGTAAAGGTTGATAATATTCCA

SEQ ID NO:416

AATGTACAGTATTGCGTTTTGGGTGAAGTAATCAGTTTGTTCAGTTACTAGTTACGTGATT

SEQ ID NO:417

AATGTACAGTATTGCGTTTTGCTGACATGCCTACTGATTATTCTTCAAACATCATCAC

SEQ ID NO:418

AATGTACAGTATTGCGTTTTGTGTGTGTTTTAATTGTTCCACTTGAGATTCTTAACC

SEQ ID NO:419

AATGTACAGTATTGCGTTTTGCGTCAGCATTTTGAATCACTTCATTCTGACATGATA

SEQ ID NO:420

AATGTACAGTATTGCGTTTTGAGTAATTTTCAACTATTGGCCTAGTGAATTTAAGCT

SEQ ID NO:421

AATGTACAGTATTGCGTTTTGAGAAAGAGGGAAGTCACATTTATAGAGTGCTAGC

SEQ ID NO:422

AATGTACAGTATTGCGTTTTGCATCAACAGAAACAGAAACAACAACTGTGACAAATC

SEQ ID NO:423

AATGTACAGTATTGCGTTTTGCCAAAGAATATCCCTTTATATAGCAGTGGAACAATT

SEQ ID NO:424

AATGTACAGTATTGCGTTTTGCAGAATATGCAGTGATAAGTGCTGTTTCATCACT

SEQ ID NO:425

AATGTACAGTATTGCGTTTTGTTCCCCCTGTGACGACTACTTTTCCTC

SEQ ID NO:426

AATGTACAGTATTGCGTTTTGCGGTCCCTATTTCTTCTCTGCTTCGT

SEQ ID NO:427

AATGTACAGTATTGCGTTTTGCTGAACAGTTCTGTCTCTATTACCCGACCTC

SEQ ID NO:428

AATGTACAGTATTGCGTTTTGCGTTCATAGCCTTCTATCCGAGTATGTAGCA

SEQ ID NO:429

AATGTACAGTATTGCGTTTTGCCCTTCTGTCTCGCAGGTTAATCC

SEQ ID NO:430

AATGTACAGTATTGCGTTTTGGCTTCCAGCCATTTCTGAGATATCCTCACAGT

SEQ ID NO:431

AATGTACAGTATTGCGTTTTGACCAGGAGGAACAAAGACACATGAAGATCAT

SEQ ID NO:432

AATGTACAGTATTGCGTTTTGGCGCCCCCGAGTTTCTTACGAATC

SEQ ID NO:433

AATGTACAGTATTGCGTTTTGTTTATACACAGTTTGGAGTTTGAGAATCAGAAGACT

SEQ ID NO:434

AATGTACAGTATTGCGTTTTGGGTTATCTCTGGCTGATGAGATTATGAGTGATTCTC

SEQ ID NO:435

AATGTACAGTATTGCGTTTTGGCCAAGCTAGTGATTGATGTGATTTCGCTAT

SEQ ID NO:436

AATGTACAGTATTGCGTTTTGCCCTCCTCTAGTACTCCCTGTTTGT

SEQ ID NO:437

AATGTACAGTATTGCGTTTTGCTCCTTCTGTCCCAATCAACTAGTCTAGC

SEQ ID NO:438

AATGTACAGTATTGCGTTTTGGCCTCGTCCCTCTTCCCTTAGGTAA

SEQ ID NO:439

AATGTACAGTATTGCGTTTTGTCTCTTCCCATTAGTCTGAGTACTGAGTGATT

SEQ ID NO:440

AATGTACAGTATTGCGTTTTGAGCATTCTTGAGACTTAAAGTGGCATTCTAAAGG

SEQ ID NO:441

AATGTACAGTATTGCGTTTTGATTTTTATTCTCAAGAGGCAGAAATACCAACTTACC

SEQ ID NO:442

AATGTACAGTATTGCGTTTTGAATTTATAGCTCTTTTCATCTGCTTTGGTATCATCA

SEQ ID NO:443

AATGTACAGTATTGCGTTTTGGCCTCTAATCTGATATACAGCCTTAGAAAGTCACA

SEQ ID NO:444

AATGTACAGTATTGCGTTTTGTGTGCCATTGTCCTGGAGCAACAATT

SEQ ID NO:445

AATGTACAGTATTGCGTTTTGAGTGTACTGCTCGTTTTCTTAATTTGAAAAGTGAGT

SEQ ID NO:446

AATGTACAGTATTGCGTTTTGACCCATGAACTAATACTTATTTGAGATTGGTCCAT

SEQ ID NO:447

AATGTACAGTATTGCGTTTTGCATGGTGCAACAAAAGTAAGAATCCAACAGTTTT

SEQ ID NO:448

AATGTACAGTATTGCGTTTTGTTGAAATGTTAAGTAAGCTTGAAATACCGATAGCAT

SEQ ID NO:449

AATGTACAGTATTGCGTTTTGGGGAGGAAGAAAATGAAGCACGAGGAAAAC

SEQ ID NO:450

AATGTACAGTATTGCGTTTTGATTTGGGATGTACTCTAAATTTAAAGCAGCAAATCA

SEQ ID NO:451

AATGTACAGTATTGCGTTTTGTCAAGAGCAGAATTTGGAGACTTTGATATTA AAACT

SEQ ID NO:452

AATGTACAGTATTGCGTTTTGCGGTTACTAACATGTTTAGGGAAATAGACA ACTGTT

SEQ ID NO:453

AATGTACAGTATTGCGTTTTGCCTGACAACAGATCCCATATAATTA ACTTTCATACC

SEQ ID NO:454

AATGTACAGTATTGCGTTTTGAGATGAAGAAGATGAGGAACGAGAGAGTAAAAGC

[0133] The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art, readily modify and/or adapt for various applications, without departing from the general concept of the invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.

[0134] The breadth and scope of the present invention should not be limited by any of the above-described exemplary embodiments, but should be defined only in accordance with the following claims and their equivalents.

[0135] All of the various aspects, embodiments, and options described herein can be combined in any and all variations.

[0136] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be herein incorporated by reference. U.S. Appl. No. 62/648,174, filed March 26, 2018, is incorporated herein by reference in its entirety.

WHAT IS CLAIMED IS:

1. A method for preparing DNA and cDNA libraries from a sample, comprising:
ligating a DNA tag to an end of a DNA molecule in a sample, wherein the DNA tag comprises a unique molecular identifier (UMI) and a DNA identifier; and
performing reverse transcription of a RNA molecule in the sample in the presence of a RNA tag, wherein the RNA tag comprises a RNA identifier, a UMI, and a poly(T).
2. The method of claim 1, wherein the reverse transcription is performed in the presence of a second RNA tag, wherein the second RNA tag comprises a RNA identifier, a UMI, and a template switching oligonucleotide (TSO).
3. The method of claim 1 or 2, further comprising amplifying the tagged DNA and the tagged cDNA for enrichment with a set of gene specific primers.
4. The method claim 3, further comprising separating the amplified sample into first and second samples.
5. The method of any one of claims 1-4, wherein the DNA and RNA molecules are obtained from a biological sample.
6. The method of any one of claims 1-5, wherein the DNA and RNA molecules are fragmented DNA and RNA from the biological sample.
7. The method of any one of claims 1-6, wherein the DNA molecule contains polished ends for ligation.
8. The method of any one of claims 1-7, wherein the RNA molecule is polyadenylated.
9. The method of any one of claims 1-8, wherein the method does not require ribosomal depletion.

10. The method of any one of claims 1-9, further comprising amplifying the first sample with primers specific for the DNA tag.

11. The method of claim 10, wherein the amplification generates a DNA library corresponding to the DNA in the sample.

12. The method of any one of claims 1-9, further comprising amplifying the second sample with primers specific for the RNA tag.

13. The method of claim 12, wherein the amplification generates a cDNA library corresponding to the RNA in a sample.

14. The method of any one of claims 1-13, further comprising sequencing the DNA or cDNA library.

15. The method of any one of claims 1-14, wherein the DNA molecule is genomic DNA.

16. The method of any one of claims 1-11, 14, and 15, wherein the DNA library can be used for DNA variant detection, copy number analysis, fusion gene detection, or structural variant detection.

17. The method of any one of claims 1-9 and 12-14, wherein the cDNA library can be used for RNA variant detection, gene expression analysis, or fusion gene detection.

18. The method of any one of claims 1-15, wherein the library can be used for paired DNA and RNA profiling.


19. A DNA library made by the method of claim 10.

20. A cDNA library made by the method of claim 12.

21. A DNA tag comprising a unique molecular identifier (UMI) and a DNA identifier.
22. The DNA tag of claim 21, wherein the UMI and the DNA identifier are positioned in a 5' to 3' direction.
23. A RNA tag comprising a RNA identifier, a UMI, and a poly(T).
24. The RNA tag of claim 23, wherein the RNA identifier, the UMI, and the poly(T) are positioned in a 5' to 3' direction.
25. A RNA tag comprising a RNA identifier, a UMI, and a template switching oligonucleotide (TSO).
26. The RNA tag of claim 25, wherein the RNA identifier, the UMI, and the TSO are positioned in a 5' to 3' direction.
27. A composition comprising at least 2 tags of claims 21-26.
28. A composition comprising the tags of claims 21-26.
29. A method for preparing targeted DNA and cDNA libraries, comprising:
 - (a) obtaining purified DNA and RNA from a biological sample;
 - (b) fragmenting the DNA and RNA;
 - (c) polishing the ends of the double stranded DNA fragments for ligation;
 - (d) polishing the RNA fragments by polyadenylation;
 - (e) ligating a DNA tag to a 3' end of the polished DNA fragments, wherein the DNA tag comprises in a 5' to 3' direction a unique molecular identifier (UMI) and a DNA identifier;
 - (f) performing reverse transcription of the polished RNA fragments in the presence of a first RNA tag, wherein the first RNA tag comprises in a 5' to 3' direction a RNA identifier, a UMI, and a poly(T), and a second RNA tag, wherein the second RNA tag comprises in a 5' to 3' direction a RNA identifier, a UMI, and a template switching oligonucleotide (TSO);

- (g) amplifying the tagged DNA and tagged cDNA for enrichment with a set of gene specific primers;
- (h) separating the amplified sample into first and second samples;
- (i) amplifying the first sample with primers specific for the DNA tag; and
- (j) amplifying the second sample with primers specific for the RNA tag.

FIG. 1



DNA tag molecule that can be ligated to 3' end of double strand DNA fragments.




RNA tag molecule that can be used to add a 5' tag to RNA-derived cDNA fragments through reverse transcription.



RNA tag molecule that can be used to add a 3' tag to RNA-derived cDNA through template switching in RT.

FIG. 2



INTERNATIONAL SEARCH REPORT

International application No PCT/US2019/024107
--

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C12Q1/6806 C12N15/10
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 C12Q C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2018/005811 A1 (GRAIL INC [US]) 4 January 2018 (2018-01-04)	1,2,4-28
Y	paragraph [0029]; claim 1 -----	3,29
X	WO 2014/164486 A1 (ZHANG YILIN [US]) 9 October 2014 (2014-10-09)	1
Y	paragraphs [0096], [0098] -----	3,29
A	WO 2014/201273 A1 (BROAD INST INC [US]; HARVARD COLLEGE [US] ET AL.) 18 December 2014 (2014-12-18) claims 10,47 ----- -/--	1-29

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 June 2019

Date of mailing of the international search report

02/07/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Knudsen, Henrik

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2019/024107

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JASON A REUTER ET AL: "Simul-seq: combined DNA and RNA sequencing for whole-genome and transcriptome profiling", NATURE METHODS, vol. 13, no. 11, 1 November 2016 (2016-11-01), pages 953-958, XP55597677, New York ISSN: 1548-7091, DOI: 10.1038/nmeth.4028 figure 1A	1-29
A	----- FLORIAN MERTES ET AL: "Combined ultra-low input mRNA and whole-genome sequencing of human embryonic stem cells", BMC GENOMICS, vol. 16, no. 1, 12 November 2015 (2015-11-12), XP055403401, DOI: 10.1186/s12864-015-2025-z page 8, left-hand column, paragraph 1	1-29
X,P	----- WO 2018/057820 A1 (PREDICINE INC [US]) 29 March 2018 (2018-03-29) claim 17	1
X,P	----- WO 2018/126278 A2 (EXOSOME DIAGNOSTICS INC [US]; ENDERLE DANIEL [DE] ET AL.) 5 July 2018 (2018-07-05) paragraphs [0118], [0121]; claim 1	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2019/024107

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2018005811 A1	04-01-2018	AU 2017290237 A1 CA 3024984 A1 CN 109563541 A EP 3478856 A1 US 2018002749 A1 US 2019062831 A1 WO 2018005811 A1	06-12-2018 04-01-2018 02-04-2019 08-05-2019 04-01-2018 28-02-2019 04-01-2018

WO 2014164486 A1	09-10-2014	AU 2014249273 A1 CA 2904899 A1 CN 105102633 A EP 2971186 A1 HK 1217734 A1 JP 2016510992 A US 2016024556 A1 WO 2014164486 A1	01-10-2015 09-10-2014 25-11-2015 20-01-2016 20-01-2017 14-04-2016 28-01-2016 09-10-2014

WO 2014201273 A1	18-12-2014	US 2016122753 A1 WO 2014201273 A1	05-05-2016 18-12-2014

WO 2018057820 A1	29-03-2018	NONE	

WO 2018126278 A2	05-07-2018	NONE	
