



US 20030163845A1

(19) **United States**

(12) **Patent Application Publication**
Mukerji et al.

(10) **Pub. No.: US 2003/0163845 A1**

(43) **Pub. Date: Aug. 28, 2003**

(54) **ELONGASE GENES AND USES THEREOF**

Publication Classification

(76) Inventors: **Pradip Mukerji**, Gahanna, OH (US);
Amanda Eun-Yeong Leonard,
Gahanna, OH (US); **Yung-Sheng**
Huang, Upper Arlington, OH (US);
Suzette L. Pereira, Westerville, OH
(US)

(51) **Int. Cl.⁷** **A01H 5/00**; C07H 21/04;
C12N 9/10; C12N 5/04; C12P 21/02;
C12N 1/12
(52) **U.S. Cl.** **800/281**; 435/69.1; 435/193;
435/419; 435/320.1; 536/23.2;
435/257.2

Correspondence Address:

STEVEN F. WEINSTOCK; ABBOTT
LABORATORIES
100 ABBOTT PARK ROAD
DEPT. 377/AP6A
ABBOTT PARK, IL 60064-6008 (US)

(57)

ABSTRACT

(21) Appl. No.: **10/156,911**

(22) Filed: **May 29, 2002**

Related U.S. Application Data

(63) Continuation-in-part of application No. 09/903,456, filed on Jul. 11, 2001, which is a continuation-in-part of application No. 09/624,670, filed on Jul. 24, 2000, which is a continuation-in-part of application No. 09/379,095, filed on Aug. 23, 1999, which is a continuation-in-part of application No. 09/145,828, filed on Sep. 2, 1998, now Pat. No. 6,403,349.

The subject invention relates to the identification of several genes involved in the elongation of polyunsaturated acids (i.e., "elongases") and to uses thereof. At least two of these genes are also involved in the elongation of monounsaturated fatty acids. In particular, elongase is utilized in the conversion of gamma linolenic acid (GLA) to dihomogamma linolenic acid (DGLA) and in the conversion of AA to adrenic acid (ADA), or eicosapentaenoic acid (EPA) to ω3-docosapentaenoic acid (DPA). DGLA may be utilized in the production of polyunsaturated fatty acids, such as arachidonic acid (AA), docosahexaenoic acid (DHA), EPA, adrenic acid, ω6-docosapentaenoic acid or ω3-docosapentaenoic acid which may be added to pharmaceutical compositions, nutritional compositions, animal feeds, as well as other products such as cosmetics.




FIG.1

Gap Weight: 6 Average Match: 2.912
 Length Weight: 4 Average Mismatch: -2.003

Quality: 50 Length: 84
 Ratio: 0.625 Gaps: 4
 Percent Similarity: 43.038 Percent Identity: 29.114

Match display thresholds for the alignment(s):

| = IDENTITY
 : = 2
 . = 1

jojobakcs x EL02 June 4, 1998 08:23 ..

```
jojobakcs 24 ATLPNFKSSINLHHVKL.GYHYLISNALFLVFIPLLGLASAHLSFSAHD 72
          .||| .| ::| :| :|:| . | || | ..
EL02      66 STLPPVLYAITAYYVIIFGGRFLLSKS..KPF.KLNGLFQLHNLVLTSL 112
```

```
jojobakcs 73 LSLLFDLLRRNLLPVVCSFLFVLLATLHFLTRP 106
          |.|| |:|.|::| |: : :|.|
EL02     113 LTLLL.LMVEQLVP IIVQHGLYFAICNIGAWTQP 145
```

FIG.2

S. cerevisiae EL02 (AA66-145) with *M. alpina* codon bias

S	T	L	P	P	V	L	Y	A	I	T	A	Y	Y	V	I	I	F	G	G	R	F	L	
TCC	ACC	CTC	CCC	CCC	GTC	CTC	TAC	GCC	ATC	ACC	GCC	TAC	TAC	GTC	ATC	ATC	TTC	GGT	GGT	CGC	TTC	CTC	
66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	
																						R0339	
L	S	K	S	K	P	F	K	L	N	G	L	F	Q	L	H	N	L	V	L	T	S	L	
CTC	TCC	AAG	TCC	AAG	CCC	TTC	AAG	CTC	AAC	GGT	CTC	TTC	CAG	CTC	CAC	AAC	CTC	GTC	CTC	ACC	TCC	CTC	
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	
																						R0339	
S	L	T	L	L	L	L	M	V	E	Q	L	V	P	I	I	V	Q	H	G	L	Y	F	
TCC	CTC	ACC	CTC	CTC	CTC	CTC	ATG	GTC	GAG	CAG	CTC	GTC	CCC	ATC	ATC	GTC	CAG	CAC	GGT	CTC	TAC	TTC	
112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	
																						R0339	
A	I	C	N	I	G	A	W	T	Q	P												R0339	
GCC	ATC	TGC	AAC	ATC	GGT	GCC	TGG	ACC	CAG	CCC													
135	136	137	138	139	140	141	142	143	144	145													

FIG.3




FIG.4A




FIG.4B

pRAE-5	GAATTCAGG	*	*	*	*	*	*	*	CATGGCCGCCGCAATCTTGGACAA
pRAE-6	GAATTCAGGCATCTCATGGATCCGCCATGGCCGCCGCAATCTTGGACAA								
	EcoRI						BamHI		NcoI

FIG.5

```

1  ATGGCCGCCG CAATCTTGGA CAAGGTCAAC TTCGGCATTG ATCAGCCCTT
51  CGGAATCAAG CTCGACACCT ACTTTGCTCA GGCCTATGAA CTCGTCACCG
101 GAAAGTCCAT CGACTCCTTC GTCTTCCAGG AGGGCGTCAC GCCTCTCTCG
151 ACCCAGAGAG AGGTCGCCAT GTGGACTATC ACTTACTTCG TCGTCATCTT
201 TGGTGGTCGC CAGATCATGA AGAGCCAGGA CGCCTTCAAG CTCAAGCCCC
251 TCTTCATCCT CCACAACTTC CTCCTGACGA TCGCGTCCGG ATCGCTGTTG
301 CTCCTGTTCA TCGAGAACCT GGTCCCCATC CTCGCCAGAA ACGGACTTTT
351 CTACGCCATC TGCGACGACG GTGCCTGGAC CCAGCGCCTC GAGCTCCTCT
401 ACTACCTCAA CTACCTGGTC AAGTACTGGG AGTTGGCCGA CACCGTCTTT
451 TTGGTCCTCA AGAAGAAGCC TCTTGAGTTC CTGCACTACT TCCACCACTC
501 GATGACCATG GTTCTCTGCT TTGTCCAGCT TGGAGGATAC ACTTCAGTGT
551 CCTGGGTCCC TATTACCCTC AACTTGACTG TCCACGTCTT CATGTACTAC
601 TACTACATGC GCTCCGCTGC CGGTGTTTCG ATCTGGTGGA AGCAGTACTT
651 GACCACTCTC CAGATCGTCC AGTTCGTTCT TGACCTCGGA TTCATCTACT
701 TCTGCGCCTA CACCTACTTC GCCTTCACCT ACTTCCCCTG GGCTCCCAAC
751 GTCGGCAAGT GCGCCGGTAC CGAGGGTGCT GCTCTCTTTG GCTGCGGACT
801 CCTCTCCAGC TATCTCTTGC TCTTTATCAA CTTCTACCGC ATTACCTACA
851 ATGCCAAGGC CAAGGCAGCC AAGGAGCGTG GAAGCAACTT TACCCCAAG
901 ACTGTCAAGT CCGGCGGATC GCCCAAGAAG CCCTCCAAGA GCAAGCACAT
951 CTAA

```

FIG.6

1 MAAAILDKVN FGIDQPFGIK LDTYFAQAYE LVTGKSIDSF VFQEGVTPLS
51 TQREVAMWTI TYFVVIFGGR QIMKSQDAFK LKPLFILHNF LLTIASGSLL
101 LLFIENLVPI LARNGLFYAI CDDGAWTQRL ELLYYLNYLV KYWELADTVF
151 LVLKKKPLEF LHYFHHSMTM VLCFVQLGGY TSVSWVPITL NLTVHVFMYY
201 YYMRSAAGVR IWMKQYLTTL QIVQFVLDLG FIYFCAYTYF AFTYFPWAPN
251 VGKCAGTEGA ALFGCGLLSS YLLLFINFYR ITYNAKAKAA KERGSNFTP
301 TVKSGGSPKK PSKSKHI*

FIG.7

1	1	~ ~ ~ ~ ~ M N S L V T Q Y A A P L F E R Y P Q L H D Y L P T L E R P F F N I S L W E H F D D V	50
GNS1			
SUR4			
MAELO			
	51	V T R V I N G R F V P S E F Q F I A G E L P L S T L P P V L Y A I T A Y V V I I F G G R F L L S K S	100
GNS1			
SUR4			
MAELO			
	101	K P . . F K L N G L F Q L H N L V I L T S L S L T L L L M V E I Q L V P I I V Q H G L Y F I A I G N I G	150
GNS1			
SUR4			
MAELO			
	151	A W T Q P L V T L Y Y M N Y I V K I F I E F I D T F F L V L K H K K L T F L H T Y H H G A T A L L G Y	200
GNS1			
SUR4			
MAELO			
	201	T Q L M G T T S T S W V P I S L N L G V H V V M Y Y F L A I A R G I I R V W W K E W V T R F I I I Q	250
GNS1			
SUR4			
MAELO			
	251	F V L D I I G F I Y F A V I Q K A V H L Y F P I L . P H C G D C V G S T T A T E A G C A I I S S Y L V	300
GNS1			
SUR4			
MAELO			

FIG.8A




FIG.8B

SCORES Init1: 153 Initn: 199 Opt: 495
57.4% identity in 549 bp overlap

MAELO	150	160	170	180	190	200
	TCTCGACCCAGAGAGAGGTCGCCATGTGGACTATCACTTACTTCGTCGTCATCTTTGGTG					
S78624	CATTAAAGCACTTTGCCCCCTGTGCTATACGCCATCACTGCCTATTACGTTATTATTTTGG					
	5990	6000	6010	6020	6030	6040
MAELO	210	220	230	240	250	260
	GTCGCCAGATCATGAAGAGCCAG--GACGCC-TTCAAGCTCAAGCCCCTCTTCATCCTCC					
S78624	GTGGCAGGTTTTTGTAAAGTAAGTCGAAACCATTAAATTAAATGGCCTTTTCCAATTGC					
	6050	6060	6070	6080	6090	6100
MAELO	270	280	290	300	310	320
	ACAACTTCCTCCTGACGATCGCGTCC--GGATCGCTGTTGCTCCTGTTTCATCGAGAACCT					
S78624	ATAATTTGGTTTTAAC-TTCACTTTTCATTGA-CGCTTTTATTGCTTATGGTTGAACAATT					
	6110	6120	6130	6140	6150	6160
MAELO	330	340	350	360	370	380
	GGTCCCCATCCTCGCCAGAAACGGACTTTTCTACGCCATCTGCGACGACGGTGCCTGGAC					
S78624	AGTGCCAATTATTGTTTCAGCAGGGTTATACTTCGCTATCTGTAATATTGGTGCTTGGAC					
	6170	6180	6190	6200	6210	6220
MAELO	390	400	410	420	430	440
	CCAGCGCCTCGAGCTCCTCTACTACCTCAACTACCTGGTCAAGTACTGGGAGTTGGCCGA					
S78624	TCAACCGCTCGTTACATTATATTACATGAATTACATTGTCAAGTTTATTGAATTTATAGA					
	6230	6240	6250	6260	6270	6280
MAELO	450	460	470	480	490	500
	CACCGTCTTTTTGGTCCTCAAGAAGAAGCCTCTTGAGTTCTGCACTACTTCCACCACTC					
S78624	CACCTTTTCTTGGTGCTAAAACATAAAAAAATTGACATTTTTCGA-TACTT--ATCA--C					
	6290	6300	6310	6320	6330	6340
MAELO	510	520	530	540	550	
	GATGACCATGGTCTCTGCTTTGT---CCAGCTTGGAGGATA-CACTTCAGTGTCTCTGG					
S78624	CATGGCGCTACTGCCTTATTATGTTACACCCAATTGATGGGCACCACATCTATTTCTTGG					
	6350	6360	6370	6380	6390	6400
MAELO	560	570	580	590	600	610
	GTCCCTATTACCCTCAACTTGACTGTCCACGTCTTCATGTACTACTACTACATGCGCTCC					
S78624	GTCCCTATTTTCATTGAACCTTGGTGTTACGTGGTTATGTATTGGTACTATTT---CTTG					
	6410	6420	6430	6440	6450	
MAELO	620	630	640	650	660	670
	GCTGCC---GGTGTTTCGCATCTGGTGGAAGCAGTACTTGACCACTCTCCAGATCGTCCAG					
S78624	GCTGCCAGAGGCATCAGGGTCTGGTGGAAGGAATGGGTTACCAGATTTCAAATTATCCAA					
	6460	6470	6480	6490	6500	6510

FIG.9A




FIG.9B

HOST(PLASMID)	334(pCGN7875) 25 μ M OA	334(pYES2) 25 μ M OA	334(pYX242) 25 μ M GLA	334(pRAE-5) 25 μ M GLA	334(pRAE-6) 25 μ M GLA	334(pYX242) NO SUBSTRATE	334(pRAE-5) NO SUBSTRATE
ADDED SUBSTRATE							
FATTY ACID	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)
C16:0	11.948	23.601	35.123	92.011	85.160	16.294	25.34
C16:1	30.665	71.217	32.789	315.464	115.456	56.183	113.913
C18:0	6.185	9.704	10.515	22.628	18.879	5.535	11.092
C18:1n-9	35.340	57.429	33.989	154.386	106.881	28.388	51.538
C18:3n-6			48.856	58.084	12.434		
C20:0			0.474	0.710	0.244		
C20:1n-9	(0.375%)* 0.352	(0.309%)* 0.527		1.405	0.867		0.516
C20:3n-6	ND	ND	(0.092%)* 0.226	(0.324%)* 2.504	(0.269%)* 1.006	ND	ND
C22:0				0.460			
C22:1n-9				0.321	0.315		
C24:0					1.825		0.999
TOTAL LIPID	93.760	170.490	245.090	771.690	374.420	112.99	256.52
ND = NOT DETECTED							
*% TOTAL FATTY ACID							

FIG. 10A

HOST(PLASMID)	334(pYX242)	334(pYX242)	334(pRAE-5)	334(pRAE-5)	334(pRAE-6)
ADDED SUBSTRATE	25 μM GLA	25 μM GLA	25 μM GLA	25 μM GLA	25 μM GLA
FATTY ACID	LIPID (μg)	LIPID (μg)	LIPID (μg)	LIPID (μg)	LIPID (μg)
C16:0	60.683	61.487	100.998	96.193	66.761
C16:1	79.838	79.586	359.754	220.440	87.359
C18:0	9.784	10.106	15.317	15.165	16.744
C18:1n-9	38.536	39.936	108.472	89.637	71.631
C18:3n-6	17.974	17.833	82.866	56.596	17.766
C20:0			0.510	0.570	
C20:1n-9					
C20:3n-6	(0.136%)* 0.389	(0.130%)* 0.374	(0.336%)* 3.035	(0.401%)* 2.689	(0.353%)* 1.185
C22:0			0.414		
C22:1n-9				0.383	
C24:0			1.513	1.626	
TOTAL LIPID	285.560	288.045	902.560	671.113	335.496
*% TOTAL FATTY ACID					

FIG. 10B

HOST(PLASMID)	334(pRAE-5/pCGR4) 25 μM GLA	334(pYY242/pYES2) 25 μM GLA	HOST(PLASMID) ADDED SUBSTRATE	334(pRAE-5/pCGR4) 25 μM GLA	334(pYY242/pYES2) 25 μM GLA
ADDED SUBSTRATE					
FATTY ACID	LIPID (μg)	LIPID (μg)		LIPID (μg)	LIPID (μg)
C16:0	41.050	37.169	C16:0	96.986	32.221
C16:1	99.393	100.552	C16:1n-7	209.667	62.757
C18:0	34.432	27.852	C18:0	80.418	14.027
C18:1	110.631	92.786	C18:1n-9	207.104	28.701
C18:3n-6	15.004	7.924	C18:3n-6	25.264	10.543
C20:0	0.643	0.574	C20:0	2.038	
C20:1	1.996	1.684	C20:1n-9	3.591	
C20:3n-6	0.542	0.607	C20:3n-6	1.284	0.326
C20:4n-6	0.579		C20:4n-6	1.394	
C22:0	1.242	2.604	C22:0	1.124	
C24:0	4.754	4.563	C24:0	3.952	
TOTAL LIPID	334	300	TOTAL LIPID	756	197

FIG. 11

HOST(PLASMID)	334(pYX242)	334(pRAE-5)	334(pRELO-1)	334(pRELO-2)
ADDED SUBSTRATE	25 μ M GLA	25 μ M GLA	25 μ M GLA	25 μ M GLA
	25°C/48HRS	25°C/48HRS	25°C/48HRS	25°C/48HRS
FATTY ACID	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)	LIPID (μ g)
C16:0	28.7	76.707	84.424	77.445
C16:1	0.729	2.513	1.532	1.056
C18:0	7.432	15.761	27.17	21.32
C18:1n-9	28.9	77.323	109.419	82.844
C18:3n-6	9.729	29.236	19.085	18.804
C20:0		0.643	0.522	0.537
C20:1n-9		0.77	0.426	0.299
C20:3n-6	(0.185%)* 0.374	(0.279%)* 1.472	(0.153%)* 0.748	(0.200%)* 0.832
C22:0		0.451		
C22:1n-9			0.224	
C24:0		0.918		
TOTAL LIPID	202	527	490	416
% TOTAL FATTY ACID				

FIG. 12

SCORES Init1: 156 Initn: 215 Opt: 296

Smith-Waterman score: 296; 28.8% identity in 264 aa overlap

	10	20	30	40	50	60	
U61954	RTFKMMDQILGTNFTYEGAKEVARGLEGFSAKLAVGYIATIFGLKYMKDRKAFDLSTPL						
				:: :: : :: :			
MAELO	AQAYELVTGKSIDSFVFQEGVTPNSTQREVAMWTITYFVVIFGGRQIMKSQDAFKLK-PL						
	30	40	50	60	70	80	
	70	80	90	100	110	119	120
U61954	NIWNGILSTFSLGFLTF-PTLLSVIRKDGFSHTYSHVSELYTDSTSGYWI-----F						
	:: :: :: : :: :: :: : : :						
MAELO	FILHNFLTIIASGSLLLFIENLVPILARNGL-----FYAICDDGAWTQRLELLYY						
	90	100	110		120	130	
	130	140	150	160	170		
U61954	LWVISKIPELLDTVFIVLRKRPLIFMHWYHHAALTGYALVCYHE--DAVHMVWV-VWMNY						
	: : : : : : :						
MAELO	LNLVVKYWELADTVFLVLKKKPLEFLHYFHHSMT---MVLCFVQLGGYTSVSWVPITLNL						
	140	150	160	170	180	190	
	180	190	200	210	220	230	
U61954	IIHAFMYGYLLKSLKVPIPPSVAQAITTSQMVFQFA----VAIFAQVHVSYKHYVEGVE						
	: : : : : : : : : : :						
MAELO	TVHVFMYYYMRSAAGVRI--WVKQYLTTLQIVQFVLDLGFIFYFCAYTYFAFTYFPWAPN						
	200	210	220	230	240	250	
	240	250	260	270	280		
U61954	-GLAYSFRGTAI-GFFMLTTYFYLWIQFYKEHYLKNGGKKYNLAKDQAKTQTKKAN						
	: : : : : : : : : : : :						
MAELO	VGKCAEGEAAALFGCGLLSYLLLFIFYRITY----NAKAKAAKERGSNFTPKTVKSGG						
	260	270	280	290	300		
MAELO	SPKKPSKSKHIX						
	310						

FIG.13

SCORES	Init1:	178	Initn:	178	Opt:	318
Smith-Waterman score:	318;	33.0% identity in 188 aa overlap				
	50	60	70	80	90	100
Z68749	SLLTNQDEVFPHIRARRFIQEHFGLFVQMAIAYVILVFSIKRFMRDREPFQLTTALRLWN					
MAELO	ELVTGKSIDSFVFQEGVTPLTSTQREVAMWTITYFVVIFGGRQIMKSQDAFKLPLFILHN					
	30	40	50	60	70	80
	110	120	130	140	150	160
Z68749	FFLSVFSIYGSWTMFPF--MVQQIRLYGLYGCCEALSNLPSQAEYWLFLTILSKAVEFV					
MAELO	FLLTIAS--GSLLLLFIEIENLVPILARNGLFYAICDD-GAWTQRLELLYYLNYLVKYWELA					
	90	100	110	120	130	140
	170	180	190	200	210	220
Z68749	DTFFLVLRKKPLIFLHWYHHMATVFVFFCSNYPTPSSQSRVGIVNLFVHAFMYPYYFTRS					
MAELO	DTVFLVLKKKPLEFLHYFHHSMTMVLCFVQLGGYTSVSWVPITLNLTVHVFMYYYMRSA					
	150	160	170	180	190	200
	230	240	250	260	270	
Z68749	MNIKVPAKISMAVTVLQLTQF--MCFIYGCTLMYYSLATNQARYPSNTPATLQCLSYTL					
MAELO	AGVRIWWK--QYLTTLQIVQFVLDLGFIFYFCAYTYFAFTYFPWAPNVGKCAGTEGAALFG					
	210	220	230	240	250	260
	280					
Z68749	HLL					
MAELO	CGLLSSYLLLFINFYRITYNAKAKAAKERGSNFTP KTVKSGGSPKKPSKSKHIX					
	270	280	290	300	310	

FIG.14




FIG.15

Mouse

SCORES Initl: 161 Initn: 191 Opt: 325

Smith-Waterman score: 325; 28.8% identity in 285 aa overlap

```

U97107      MDTSMNFSRGLKMD--LMQPYDFETFDLRPFLEEYWSSF-----LIVV
               ||| | : | :: | ::: | : | : | : | :
MAELO       MAAILDKVNFGIDQPFGIKLDTYFAQAYELVTGKSIDSFVFQEGVTP LSTQREVAMWTI
              10        20        30        40        50        60

                50        60        70        80        90        100
U97107      VYLLLIVVGQTYMRTRKSFSLSQRPLILWSFFLAIFSILGTLRMVKFMATVMFTVGLKQTV
             :|::|| | :|:::||::| :|::| | :|::| | :|::| :|::| :|::| :|::|
MAELO       TYFVVIFGGRRQIMKSQDAFKLKPLFILHNFLTITAS--GSL-LLLFIEINLV-PILARNGL
              70        80        90        100       110

                110       120       130       140       150
U97107      CFAIYTDDAVVRFWSFLLSKVV---ELGDTAFIILRKRLIFVHWYHHST--VLLFTS
             :|| | | :| :| :| :| :| :| :| :| :| :| :| :| :| :| :| :|
MAELO       FYAICDDGAWTORLELLYYLNVLKYVELADTVFLVLKKKPFLHLYFHHSMTMVLCFVQ
              120       130       140       150       160       170

            160         170         180         190         200         210
U97107      FGYKNKVPSGGWF-MTMNFVGHVSVMYTTYTMMKAALKHPNLPMVITSLQILQMVLG---
             :| :| :| :| :| :| :| :| :| :| :| :| :| :| :| :| :|
MAELO       LGGYTSV---SWVPITLNLTVHVFMYYYYYM RSAAGVR--IW WKQLTTLTQIVQFVLDLGF
              180         190         200         210         220         230

                  220         230         240         250         260
U97107      -----TI FGILNYIWRQEKG-CH TTTEHFFWSFMLYGTYFILFAHFFHRAYLRPKGKVA
               | |:: : | : | | | ::: | ::||| :|::| :|::| :|::| :|::|
MAELO       IYFCAYTYFAFTYFPWAPNVGKCAGTEGAALFGCGLLSSYLLLFINFYRITY-NAKAHAA
                 240         250         260         270         280         290

    270
U97107     SKSQX
           ::
MAELO      KERGSNFTP KTVKSGGSPKKPSKSHIX
          300        310
```

Human

SCORES Initl: 147 Initn: 147 Opt: 211

Smith-Waterman score: 211; 28.7% identity in 150 aa overlap

```

110      120      130      140      150      160
MAELO    NLVPILARNGLFYAICDDGAWTQRLELLYYLNVLVKYWELADTVFLVLKKKPLEFLHYFH
AC004050      SLLVVKDLTYLLPLCLPGDTIFIILRKQKLIFLHWYH
                        10      20      30

170      180      190      200      210      220
MAELO    HSMTMVLCFVQLGGYTSVSWVPITLNLTVHVFMYYYYMRSAAGVRIWWK--QYLTTLQIV
AC004050    HITVLLYSWYSYKDMVAGGGWFMTMNYGVHAVMYSYYALRAAGFRVSRKFAMFITLSQIT
                40      50      60      70      80      90

230      240      250      260      270      280
MAELO    QFVLDLGFIFYFCAYTYFAFTYFPWAPNVGKCGTEGAALFGCGLLSSYLLLFINFYRITY
AC004050    QMLMG-----CVVNYLVFC--WMQH-DQCHSHFQNIWFSSLMYLSYLVLFCHFFFEAY
                100      110      120      130      140

```

FIG. 16

SCORES Initl: 87 Initn: 218 Opt: 232
 Smith-Waterman score: 272; 29.7% identity in 232 aa overlap

	40	50	60	70	80	90
MAELO	SFVFQEGVTPLSTQREVAMWTITYFVVIFGGRQIMKSQDAFKLKPLFILHNFLLTIASGS					
				: : : : : :		
I05465	PRYKSQRMVPPGQLHPYVCLFCYLLTHCMAGTKTTEEPAAVLLPSILQLYNLGLTLLS--					
	20	30	40	50	60	70
	100	110	120	130	140	150
MAELO	LLLLFIENLVPILARNGLFYAICDDGAWTQRLELLYYL--NYLVKYWELADTVFLVLKKK					
	: : : : : : : : : : : : :					
I05465	-LYMFYELVTGVWEGKYNFFCQGTRSAGESDMKIIRVLWVWYFSLKIEFMDTFFFILRKN					
	80	90	100	110	120	
	160	170	180	190	200	210
MAELO	--PLEFLHYFHH-SMTMVLCFVQLGGYTSVSWVPITLNLTVHVFMYYYY-MRSAAGVR--					
	: : : : : : : : : : :					
I05465	NHQITVLHVYHHATMLNIWWFVMNWVPCGHSYFGATLNSFIHVLMSYYGLSSIPSMRPY					
	130	140	150	160	170	180
	220	230	240	250	260	270
MAELO	IWWKQYLTTLQIVQFVLDLGFYFCAYTYFAFTYFPWAPNVGKCAGTEGAALFGCGLLSS					
	: : : : : : : : : : : :					
I05465	LWWKKYITQGQLVQFVLTIIQTTTCG-----VFWP-----CSFPLGWLFFQIGYMIS					
	190	200	210	220	230	
	280	290	300	310		
MAELO	YLLLFINFYRITYNAKAKAAKERGSNFTPKTVKSGGSPKKPSKSKHIX					
	: : : : :					
I05465	LIALFTNFYIQTYNKKGASRRKEHLKGHQNGSVAAVNGHTNSFPSLENSVKPRKQRKDXQ					
	240	250	260	270	280	290

FIG.17

1 MGTDQGKTFT WEELAAHNTK DDLLLAIRGR VYDVTKFLSR HPGGVDTLLL
51 GAGRDVTPVF EMYHAFGAAD AIMKKYYVGT LVSNELPIFP EPTVFHKTIK
101 TRVEGYFTDR NIDPKNRPEI WGRYALIFGS LIASYQAQLF VPFVVERTWL
151 QVVFAIIMGF ACAQVGLNPL HDASHFSVTH NPTVWKILGA THDFFNGASY
201 LVWMYQHMLG HHPYTNIAGA DPDVSTSEPD VVRIKPNQKW FVNHINQHMF
251 VPFLYGLLAF KVRIQDINIL YFVKTNDAIR VNPISTWHTV MFWGGKAFFV
301 WYRLIVPLQY LPLGKVLLLF TVADMVSSYW LALTFQANHV VEEVQWPLPD
351 ENGIIQKDW A MQVETTQDY AHDShLWTSI TGSLNYQAVH HLFPNVSQHH
401 YPDILAIKN TCSEYKVPYL VKDTFWQAFA SHLEHLRVLG LRPKEE*

FIG.18




FIG. 19

HOST(PLASMID)	334(MAD708-2)	334(MAD708-10)	334(MAD708-18)	334(MAD708-19)	334(MAD708-30)	334(pRAE5)
ADDED SUBSTRATE	25 μM GLA	25 μM GLA	25 μM GLA	25 μM GLA	25 μM GLA	25 μM GLA
FATTY ACID	% TOTAL LIPID					
C16:0	14.1	14.68	14.38	15.45	14.13	13.59
C16:1	42.84	43.42	42.57	38.03	43.58	43.98
C18:0	3.19	3.28	3.63	4.08	3.37	2.04
C18:1n-9	17.66	19.39	19.6	20.8	20.06	10.88
C18:3n-6	6.65	5.58	10.24	9.46	3.56	11.14
C20:0	0.26	0.3	0.32	0.4	0.46	0.57
C20:3n-6	(47.5%) 6.03	(41.2%) 3.92	(8.0%) 0.91	(21.5%) 2.59	(49%) 3.43	(3.4%) 0.24
TOTAL LIPID (μg)	238.47	307.86	188.51	167.31	207.47	466.65
(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)						

FIG.20




FIG.21

```

1  ATGGAGTCGA TTGCGCCATT CCTCCCATCA AAGATGCCGC AAGATCTGTT
51 TATGGACCTT GCCACCGCTA TCGGTGTCCG GGCCGCGCCC TATGTCGATC
101 CTCTCGAGGC CGCGCTGGTG GCCCAGGCCG AGAAGTACAT CCCCACGATT
151 GTCCATCACA CGCGTGGGTT CCTGGTCGCG GTGGAGTCGC CTTTGGCCCG
201 TGAGCTGCCG TTGATGAACC CGTTCCACGT GCTGTTGATC GTGCTCGCTT
251 ATTTGGTCAC GGTCTTTGTG GGCATGCAGA TCATGAAGAA CTTTGAGCGG
301 TTCGAGGTCA AGACGTTTTT GCTCCTGCAC AACTTTTGTG TGGTCTCGAT
351 CAGCGCCTAC ATGTGCGGTG GGATCCTGTA CGAGGCTTAT CAGGCCAACT
401 ATGGA CTGTT TGAGAACGCT GCTGATCATA CCTTCAAGGG TCTTCCTATG
451 GCCAAGATGA TCTGGCTCTT CTACTTCTCC AAGATCATGG AGTTTGTCTGA
501 CACCATGATC ATGGTCCTCA AGAAGAACAA CCGCCAGATC TCCTTCTTGC
551 ACGTTTACCA CCACAGCTCC ATCTTCACCA TCTGGTGGTT GGTACCTTT
601 GTTGCACCCA ACGGTGAAGC CTACTTCTCT GCTGCGTTGA ACTCGTTCAT
651 CCATGTGATC ATGTACGGCT ACTACTTCTT GTCGGCCTTG GGCTTCAAGC
701 AGGTGTCGTT CATCAAGTTC TACATCACGC GCTCGCAGAT GACACAGTTC
751 TGCATGATGT CGGTCCAGTC TTCCTGGGAC ATGTACGCCA TGAAGGTCCT
801 TGGCCGCCCC GGATACCCCT TCTTCATCAC GGCTCTGCTT TGGTTCTACA
851 TGTGGACCAT GCTCGGTCTC TTCTACAACT TTTACAGAAA GAACGCCAAG
901 TTGGCCAAGC AGGCCAAGGC CGACGCTGCC AAGGAGAAGG CAAGGAAGTT
951 GCAGTAA

```

FIG.22

1 MESIAPFLPS KMPQDLFMDL ATAIGVRAAP YVDPLEAALV AQAKEYIPTI
51 VHHTRGFLVA VESPLARELP LMNPFHVLLI VLAYLVTVFV GMQIMKNFER
101 FEVKTFSLLE NFCLVSISAY MCGGILYEAY QANYGLFENA ADHTFKGLPM
151 AKMIWLFYFS KIMEFVDIMI MVLKKNNRQI SFLHVVHHSS IFTIWWLVTF
201 VAPNGEAYFS AALNSFIHVI MYGYYFLSAL GFKQVSFIKF YITRSQMTQF
251 CMMSVQSSWD MYAMKVLGRP GYPFFITALL WFYMWTLGL FYNFYRKNAK
301 LAKQAKADAA KEKARKLQ*

FIG.23

HOST(PLASMID)	334(pRPB2)	334(pYES2)
ADDED SUBSTRATE	25 μM GLA	25 μM GLA
	(n=4)	
FATTY ACID	% TOTAL LIPID	
C16:0	15.65	15.23
C16:1	35.2	38.59
C18:0	5.68	5.55
C18:1n-9	25.55	25.27
C18:3n-6	3.1	6.75
C20:0	0.36	0.14
C20:3n-6	(62.0%) 5.06	(2.6%) 0.18
TOTAL LIPID (μg)	314	247
(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)		

FIG.24

HOST(PLASMID)	334(pRPB2)	334(pRPB2)	334(pRPB2)	334(pRPB2)	334(pRPB2)	334(pRPB2)
ADDED SUBSTRATE	25 mM SA	25 mM OA	25 mM LA	25 mM DGLA	25 mM AA	25 mM ADRENIC
	C18:0	C18:1n-9	C18:2n-6	C20:3n-6	C20:4n-6	C22:4n-6
FATTY ACID	% TOTAL LIPID					
C16:0	15.07	14.52	15.74	15.69	16.06	15.15
C16:1	33.7	32.37	32.23	25.65	33.65	33.39
C18:0	*9.78	5.83	5.61	8.33	4.52	5.35
C18:1n-9	31.2	*37.25	26.05	20.15	24.54	28.54
C18:2n-6			*10.4			
C18:3n-6						
C20:2n-6			0.29			
C20:3n-6				*16.5		
C20:4n-6				0.27	*11.7	
C22:4n-6						*7.46
TOTAL LIPID (mg)	132	130	171	55	225	163

FIG.25A

HOST(PLASMID)	334(pRPB2)	334(pRPB2)	334(pRPB2)
ADDED SUBSTRATE	25 μM ALA	25 μM STA	25 μM EPA
	C18:3n-3	C18:4n-3	C20:5n-3
FATTY ACID	% TOTAL LIPID		
C16:0	17.32	16.01	20.67
C16:1	27.68	34.31	50.7
C18:0	6.75	5.39	6.14
C18:1n-9	28.4	28.54	
C18:3n-3	*8.39		
C18:4n-3		*1.95	
C20:4n-3		(73.2%) 5.33	
C20:5n-3			*10.33
C22:5n-3			0.25
TOTAL LIPID (μg)	114	199	201

*INDICATES SUBSTRATE ADDED
(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.25B

HOST(PLASMID)	334(pRPB2+PRPE31)	334(pYES2+pYX242)
ADDED SUBSTRATE	25 μM GLA	25 μM GLA
FATTY ACID	% TOTAL LIPID	
C16:0	15.54	18.26
C16:1	30.16	33.51
C18:0	8.76	5.58
C18:1n-9	27	27.37
C18:3n-6	*2.6	*5.6
C20:0	0.4	0.32
C20:3n-6	(57.4%) 3.55	(2.9%) 0.17
C20:4n-6	(27.6%) 1.32	ND
TOTAL LIPID (μg)	254	258

*INDICATES SUBSTRATE ADDED
(% CONVERSION)=RODUCT/(SUBSTRATE+PRODUCT)

FIG.26A

HOST(PLASMID)	334(pRPB2+PRPE31)	334(pYES2+pYX242)
ADDED SUBSTRATE	25 μM STA	25 μM STA
FATTY ACID	% TOTAL LIPID	
C16:0	18	16.4
C16:1	28.37	34.78
C18:0	7.42	5.71
C18:1n-9	26.44	30.15
C18:4n-3	*2.93	*4.57
C20:0	0.25	0.17
C20:4n-3	4.13	0.32
C20:5n-3	(39%) 1.87	(2.1%) .10
TOTAL LIPID (μg)	257	304

*INDICATES SUBSTRATE ADDED
(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.26B

SCORES Initl: 114 Initn: 278 Opt: 278
Smith-Waterman score: 308; 30.9% identity in 259 aa overlap

	40	50	60	70	80	90	99
GLELO	VAQAEKYIPTIVHHTRGFLVAVESPLARELPLMNPFHVLLIVLAYLVTVFVGMQIMKNFE						
					: : : :: : :: : :		
MAELO	GIKLDTYFAQAYELVTGKSIDSFVFEQGVTPPLSTQREVAMWTITYFVVI FGGRQIMKSQD						
	20	30	40	50	60	70	
	100	110	120	130	140	150	
GLELO	RFEVKTFSLHNFLVSI SAYMCGGILYE--AYQANYGLFENAADHTFKGLPMAKMIWLF						
	:: : : : : : :: : : : :						
MAELO	AFKCLKPLFILHNFLT IASGSLLLFIENLVPILARNGLFYAICDDGAWTQRLELLYYLN						
	80	90	100	110	120	130	
	160	170	180	190	200	210	
GLELO	YFSKIMEFVDTMIMVLKKNRQISFLHVYHHSSI FTIWWLVT FVAPNGEAYFSAALNSFI						
	: :: : : : : : : : : : : : : : : : :						
MAELO	YLVKYWELADTVFLVLKK--KPLEFLHYFHHS-MTMVLCFVQLGGYTSVSWVPITLNLTV						
	140	150	160	170	180	190	
	220	230	240	250		260	
GLELO	HVIMYGYGYFLSALGFKQVSFIKFYITRSQMTQF-----CMMSVQS----SWDMYAM						
	: : : : : : : : : : : :						
MAELO	HVFMYYYMRSAAGVRI--WWKQYLTTLQIVQFVLDLGFIFYCAYTYFAFTYFPWAPNVG						
	200	210	220	230	240	250	
	270	280	290	300	310		
GLELO	KVLGRPGYPFFITALLWFYMWMTMLGLFYNFYRKNAKLAKQAKADAAKEKARKLQ						
	: : : : : : :::						
MAELO	KCAGTEGAALFGCGLSSYLL----LFINFYR----ITYNAKAKAAKERGSNFTPKTVKS						
	260	270	280	290	300		
MAELO	GGSPKKPSKSKHIX						
	310						

FIG.27

GLELO	1	-	M	E	S	I	A	P	F	L	P	S	K	M	P	Q	D	L	F	M	D	L	A	T	A	I	G	V	R	A	A	P	Y	V	D	P	L	E	A	A	L	V	A	Q	42		
MAELO	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	20		
GNS1	1	-	-	-	-	-	-	-	-	-	M	N	S	L	V	T	Q	Y	A	A	P	L	F	E	R	Y	P	Q	L	H	D	Y	L	P	T	L	E	R	P	F	F	N	I	S	L	W	36
SUR4	1	M	N	T	T	T	S	T	V	I	A	A	V	A	D	Q	F	Q	S	L	N	S	S	S	C	F	L	K	V	H	V	P	S	I	E	N	P	-	F	G	I	E	L	42			
GLELO	43	A	E	K	V	I	P	T	I	V	H	H	T	R	G	F	L	V	A	V	E	S	P	L	A	R	E	L	P	L	M	N	P	F	H	V	L	L	I	V	L	A	Y	L	85		
MAELO	21	L	D	T	V	F	A	Q	A	Y	E	L	V	T	G	K	S	I	D	S	F	V	F	Q	E	G	V	I	P	L	S	I	Q	R	E	V	A	M	W	T	I	T	V	F	63		
GNS1	37	E	H	F	D	D	V	V	T	R	V	T	N	G	R	F	V	P	S	E	F	Q	F	I	A	G	E	L	P	L	S	I	L	P	P	V	L	Y	A	I	T	A	Y	Y	79		
SUR4	43	W	P	I	F	S	K	V	F	E	Y	F	S	G	-	Y	P	A	E	Q	F	E	F	I	H	N	K	I	F	L	A	N	G	Y	H	A	V	S	I	I	I	V	Y	Y	84		
GLELO	86	V	T	V	F	V	G	M	Q	I	M	K	N	F	E	R	F	E	V	K	T	F	S	L	L	H	N	F	C	L	V	S	I	S	A	Y	M	C	G	G	I	L	Y	E	128		
MAELO	64	V	V	I	F	G	G	R	Q	I	M	K	S	Q	D	A	-	-	F	K	L	K	P	L	F	I	L	H	N	F	L	I	A	S	G	S	L	L	L	L	F	I	104				
GNS1	80	V	I	I	F	G	G	R	F	L	S	-	-	K	S	K	P	F	K	L	N	G	L	F	Q	L	H	N	L	V	L	T	S	L	S	L	T	L	L	L	L	M	V	120			
SUR4	85	I	I	I	F	G	G	Q	A	I	L	R	A	L	N	A	S	P	L	K	F	K	L	L	F	E	I	H	N	L	F	L	T	S	I	S	L	V	L	W	L	L	M	L	127		
GLELO	129	-	-	A	Y	Q	-	-	A	N	Y	G	L	F	E	N	A	A	D	H	T	F	K	G	L	P	M	A	K	M	I	W	L	F	Y	F	S	K	I	M	E	F	V	D	167		
MAELO	105	E	N	L	V	P	I	L	A	R	N	G	L	F	Y	A	I	C	D	D	G	A	W	T	Q	R	L	E	L	Y	Y	L	N	Y	L	V	K	Y	W	E	L	A	D	147			
GNS1	121	E	Q	L	V	P	I	I	V	Q	H	G	L	Y	F	A	I	C	N	I	G	A	W	T	Q	P	L	V	T	L	Y	Y	M	N	Y	I	V	K	F	I	E	F	I	D	163		
SUR4	128	E	Q	L	V	P	M	V	Y	H	N	G	L	F	W	S	I	C	S	K	E	A	F	A	P	K	L	V	T	L	Y	L	N	Y	L	T	K	F	V	E	L	I	D	170			
GLELO	168	I	M	I	M	V	L	K	K	N	N	R	Q	I	S	F	L	H	V	Y	H	H	S	S	I	F	T	I	W	W	L	V	T	F	V	A	P	N	G	E	A	Y	F	S	210		
MAELO	148	T	V	F	L	V	L	K	K	-	-	K	P	L	E	F	L	H	Y	F	H	H	S	M	I	M	V	L	C	F	-	V	Q	L	G	G	Y	I	S	V	S	W	V	P	187		
GNS1	164	I	F	F	L	V	L	K	H	-	-	K	K	L	T	F	L	H	T	Y	H	H	G	A	T	A	L	L	C	Y	-	T	Q	L	M	G	T	I	S	I	S	W	V	P	203		
SUR4	171	T	V	F	L	V	L	R	R	-	-	K	K	L	L	F	L	H	T	Y	H	H	G	A	T	A	L	L	C	Y	-	T	Q	L	I	G	R	T	S	V	E	W	V	V	210		

FIG.28A

GLELO	211	A A L N S F I H V I M Y G Y Y F L S A L G F K Q V S F I K F Y I I T R S Q M T Q F C M M	253
MAELO	188	U T L N L T V H V F M Y Y Y Y M R S A A G V R - - I W W K Q Y L I T T L Q I V Q F V L D	228
GNS1	204	U S L N L G V H V V M Y W Y Y F L A A R G I R - - V W W K E W V I R F Q I I Q F V L D	244
SUR4	211	I L L N L G V H V I M Y W Y Y F L S C G I R - - V W W K Q W V I R F Q I I Q F L I D	251
GLELO	254	S V Q S S W D M Y A M K V L G R P G Y P F F I T A L L W F Y M W T M L G L F Y N F Y R	296
MAELO	229	L G F I Y F C A Y I Y F A F T Y F P W - A P N V G K C A G I E G A A L F G C G L L S S	270
GNS1	245	I G F I Y F A V M Q K A V H L Y F P - I L P H C G D C V G S T T A T F A G C A I I S S	286
SUR4	252	L V F V Y F A T Y I F Y A H K Y L D G I L P N K G T C Y G I Q A A A Y G Y L I L T S	294
GLELO	297	K N A K L A K Q A K A D A A K E K A R K L Q	318
MAELO	271	Y L L L F I N F Y R I T Y N A K A K A A K E R G S N F T P K T V K S G G S P K K P S K	313
GNS1	287	Y L V L F I S F Y I N V Y K R K G T K I S R V V K R A H G G V A A K V N E Y V N V D L	329
SUR4	295	Y L L L F I S F Y I Q S Y K K G G K K I V K K E S E V S G S V A S G S S T G V K T S N	337
MAELO	314	S K H I	317
GNS1	330	K N V T P S P S P K P Q H R R K R	347
SUR4	338	T K V S S R K A	345

FIG.28B

SCORES Init1: 83 Initn: 186 Opt: 271

Smith-Waterman score: 297; 28.5% identity in 242 aa overlap

	30	40	50	60	70	80
MAELO	YELVTGKSIDSFVFQEGVTPNSTQREVAMWTITYFVVIFGGRQIMKSQDAFKLKPLFILH					
				:: ::: : ::: : : ::::		
HS1	STYFKALLGPRDTRVKGWFLLDNYIPTFICSVIYLLIVWLGPKYMRNKQPFSCRGILVVY					
	10	20	30	40	50	60
	90	100	110	120	130	140
MAELO	NFLTIIASGSLLLLFIENLVPILARNGLFYAICDDGAWTQRLELLYYL--NYLVKYWELA					
	: :	:	: :	: :	: ::::	: :
HS1	NLGLTLLS---LYMFCELVTVGWEGKYNFFCQGTRTAGESDMKIIRVLWYYYFSKLIEFM					
	70	80	90	100	110	120
	150	160	170	180	190	200
MAELO	DTVFLVLKK--KPLEFLHYFHH-SMTMVLFCFVQLGGYTSVSWVPITLNLTVHVFMYYYY-					
	:: :	: :	:	: :	: :	: :
HS1	DTFFFILRKNNHQITVLHVYHHASMLNIWWFVMNWVPCGHSYFGATLNSFIHVLMYSSYG					
	130	140	150	160	170	180
	210	220	230	240	250	260
MAELO	MRSAGVR--IWWKQYLTTLQIVQFVLDLGFIFYCAYTYFAFTYFPWAPNVGKAGTEGA					
	: : ::	: : :	::	: : :		:
HS1	LSSVPSMRPYLWWKKYITQGQLLQFVLT-IQTSCGVI-----W-P----CTFPLGW					
	190	200	210	220		230
	270	280	290	300	310	
MAELO	ALFGCGLLSSYLLLFINFYRITYNAKAKAAKERGSNFTPKTVKSGGSPKKPSKSKI					
	: :		:	: ::		
HS1	LYFQIGYMISLIALFTNFYIQTYNKKGASRRKDHLDHQNGSMAAVNGHTNSFSPLENNV					
	240	250	260	270	280	290
HS1	KPRKLRKDX					
	300					

FIG.29

SCORES Init1: 88 Initn: 208 Opt: 272
Smith-Waterman score: 279; 28.2% identity in 266 aa overlap




FIG.30

SCORES Init1: 88 Initn: 207 Opt: 223
Smith-Waterman score: 236; 30.4% identity in 191 aa overlap

	100	110	120	130	140	150
MAELO	LLLLFIENLVPILARNGLFYAICDDGAWTQRLELLYYLNYLVKYWELADTVFLVLKKKP-					
				: : : : ::: :		
MM2	IVYEFLMSGWLSTYTWRCDPIDFSNSPEALRMVRVAWLFMLSKEIEMDTVIFILRKKDG					
	20	30	40	50	60	70
	160	170	180	190	200	209
MAELO	-LEFLHYFHHSMTMVLCF----VQLGGYTSVSWVPITLNLTVHVFMYYYMRSAAGV---					
	: :	:	: :	:	: :	
MM2	QVTFLHVFHHSVLPWSWWGIKIAPGGMGSFHAM---INSSVHVVMYLYYGLSALGPVAQ					
	80	90	100	110	120	130
	210	220	230	240	250	260
MAELO	-RIWWKQYLTTLQIVQFVL---DLGFIYF---CAYTYFAFTYFPWAPNVGKCAGTEGAAL					
	: ::: :: ::	:	:	: : :	:	:
MM2	PYLWWKKHMTAIQLIQFVLVSLHISQYYFMPSCNYQYPVIIHLIW-----M					
	140	150	160	170		
	270	280	290	300	310	
MAELO	FGCGLLSSYLLLFINFYRITYNAKAKAAKERGSNFTPVTKSGGSPKKPSKSKI					
	:	: :::	: :	:	: : :	
MM2	YG----TIFFILFSNFWYHSYTKGKRLPRAVQQNGAPATTKVKAN					
	180	190	200	210		

FIG.31

SCORES Init1: 51 Initn: 115 Opt: 168

Smith-Waterman score: 168; 30.4% identity in 115 aa overlap

	30	40	50	60	70	80
MAELO	YELVTGKSIDSFVFQEGVTP LSTQREVAMWTITYFVVIFGGRQIMKSQDAFKLKPLFILH					
				:: : : :: :: : :: :		
AI225632	NAFLDNMFGPRDSRVRGWFL LDSYLP TFI LTITYLLSIWLG NKYMKNRPALSLRGILTLY					
	20	30	40	50	60	70
	90	100	110	120	130	140
MAELO	NFL LTIASGSL LLLFIENLVPILARNGLFYAICDD---GAWTORLELLYYLNYLVKYWE					
	: : : : : :	: : : :	::	: :	:	
AI225632	NLAITLLSAYMLVELI-----LSSWEGGYNLQCQN LDSAGEGDVRVAKVLVWYYFSKLVE					
	80	90	100	110	120	
	150	160	170	180	190	200
MAELO	LADTVFLVLKKK--PLEFLHYFHHSMTMVL CFVQLGGYTSVSWVPITLNLTVHVFMYYYY					
	: : : :	: : :				
AI225632	FLDTIFFVLRKKANQITFLHVYH HASMFNI					
	130	140	150			

FIG.32

SCORES Frame: (3) Initl: 332 Initn: 332 Opt: 384
40.3% identity in 144 aa overlap

	80	90	100	110	120	130
GLELO	LIVLAYLVTVFVGMQIMKNFERFEVKTFSLHNFLVSI SAYMCGGILYEAYQANYGL-F					
					: : :: : : :: :::: :	
AI815960				LYNLGITLLSAYMLAELILSTWEGGYNLQC		
				10	20	30
	140	150	160	170	180	190
GLELO	ENAADHTFKGLPMAKMIWLFYFSKIMEFVDTMIMVLKKNRQISFLHVYHHSSIFTIWWL					
	:: ::	: : ::	:	: : :: : : : : : : :	: : : : : : : : : :	: : :
AI815960	QDLTSAGEADIRVAKVLWYYFSKSVEFLDTIFFVLRKQTSQITFLHVYHHASMFNIWWC					
	40	50	60	70	80	90
	200	210	220	230	240	250
GLELO	VTFVAPNGEAYFSAALNSFIHVIMYGYFLSAL-GFKQVSFIKFYITRSQMTQFCMMSVQ					
		: : : : : : : :	: :	: : : : : : : :	: : : : : :	: : : : :
AI815960	VLNWIPCGQSFFGPTLNSFIHILMYSYYGLSVFPSMHKYLWKKYLTQAQLVQF					
	100	110	120	130	140	
	260	270	280	290	300	310
GLELO	SSWDMYAMKVLGRPGYPFFITALLWFYMWTLGLFYNFYRKNALAKQAKADAAKEKARK					

FIG.33

SCORES Init1: 316 Initn: 384 Opt: 477
Smith-Waterman score: 477; 34.2% identity in 240 aa overlap

	50	60	70	80	90	100	
GLELO	AQAEKYIPTIVHHTRGFLVAVESPLARELPLMNPFHVLLIVLAYLVTVFVGMQIMKNFER						
				: :: : : :: : : :			
HS1	MEHFDASLSTYFKALLGPRDTRVKGWFLDNYIPTFICSVIYLLIVWLGPKYMRNKQP						
	10	20	30	40	50		
	110	120	130	140	150	159	
GLELO	FEVKTFSLHNFCLVVISAYMCGGILYEAYQANYGLF-ENAAHTFKGLPMAKMIWLFYF						
	: : :: : : :: : :: : :: : : : : : :						
HS1	FSCRGILVVYNLGLTLLSLYMFCELVGTGWEGKYNFFCQGRTRTAGESDMKIIRVLWYYF						
	60	70	80	90	100	110	
	160	170	180	190	200	210	219
GLELO	SKIMEFVDTMIMVLKKNRQISFLHVYHSSIFTIWWLTVFVAPNGEAYFSAALNSFIHV						
	:: : :: : : : : : : : : : :						
HS1	SKLIEFMDTFFFI LRKNNHQITVLHVYHHASMLNIWWFVMNWVPCGHSYFGATLNSFIHV						
	120	130	140	150	160	170	
	220	230	240	250	260	270	
GLELO	IMYGYFLSAL-GFKQVSFIKFYITRSQMTQFCMMSVQSSWDMYAMKVLGRPGYPFFITA						
	: : :: : : : : : : : : : : : :						
HS1	LMYSYGLSSVPSMRPYLWKKYITQGQLLQFVLTIIQTS-----CGVIWPCTFPLGWLY						
	180	190	200	210	220	230	
	280	290	300	310			
GLELO	LLWFYMWMTMLGLFYNFYRK--NAKLAKQAKADAAKEKARKLQ						
	: : :: : :						
HS1	FQIGYMISLIALFTNFYIQTYNKKGASRRKDHLKDHQNGSMAAVNGHTNSFSPLENNVKP						
	240	250	260	270	280	290	

FIG.34

SCORES Init1: 80 Initn: 114 Opt: 178
Smith-Waterman score: 178; 28.8% identity in 146 aa overlap

	140	150	160	170	180	190
GLELO	FEN	AAD	HTF	KGL	PM	AKMIWLFYFSKIMEFVDTMIMVLKKNRQISFLHVYHHSSIFTIWW
						::: : :: ::
AC004050						DTIFIILRK--QKLIFLHWYHHITVLLYSW
					10	20
	200	210	220	230	240	250
GLELO	LVTFVAPNGEAYFSAALNSFIHVIMYGYFLSALGFKQVSFIKFYITRSQMTQFCMMSVQ					
		:: :: : :: : :			: :: : :	
AC004050	YSYKDMVAGGGWF-MTMNYGVHAVMYSYYALRAAGFRVSRKFAMFITLSQITQMLMGCVV					
	30	40	50	60	70	80
	260	270	280	290	300	310
GLELO	SSWDMYAMKVLGRPGYPFFITALLW--FYMWTMLGLFYNFYRKN--AKLAKQAKADAAKE					
	: : : : :: ::: :: : : : : : : :					
AC004050	NYLVFCWMQ--HDQCHSHF-QNIFWSSLMYLSYLVLFCHFFFEAYIGKMRKTTKAEX					
	90	100	110	120	130	140
GLELO	KARKLQ					

FIG.35

SCORES Init1: 288 Initn: 288 Opt: 399
Smith-Waterman score: 399; 34.6% identity in 211 aa overlap

	80	90	100	110	120	130
GLELO	LLIVLAYLVTVFVGMQIMKNFERFEVKTFSLHNFCLVSISAYMCGGILYEAYQANYGLF					
				:::	:	:
MM2				IVYNFSLVILSLYIVYEFLMSGWLSTYTW		
				10	20	30
	140	150	160	170	180	190
GLELO	ENAAD--HTFKGLPMAKMIWLFYFSKIMEFVDTMIMVLKKNRQISFLHVYHHSSIFTIW					
	:		::	::		::
MM2	CDPIDFSNSPEALRMVRVAWLFLSKVIELMDTVIFILRKKDGQVTFHVFHHSVLPWSW					
	40	50	60	70	80	90
	200	210	220	230	240	250
GLELO	WLVTfVAPNGEAYFSAALNSFIHVIMYGYFLSALGFKQVSFI--KFYITRSQMTQFCMM					
		:	::	:	:	:
MM2	WWGIKIAPGGMGSFHAMINSSVHVVMYLYGLSALGPVAQPYLWKKHMTAIQLIQFVLV					
	100	110	120	130	140	150
	260	270	280	290	300	309
GLELO	SVQSSWDMYAMKVLGRPGYPFFITALLWFYMWTLGLFYNF----YRKNAKLAKQAKADA					
	::	::		:		:
MM2	SLHIS-QYYFMPSCNYQ-YPVIIH-LIWMYGTIFFILFSNFWYHSYTKGKRLPRAVQQNG					
	160	170	180	190	200	
	310					
GLELO	AKEKARKLQ					
MM2	APATTKVKAN					
	210					

FIG.36




FIG.37

SCORES Init1: 64 Initn: 129 Opt: 233

Smith-Waterman score: 239; 23.7% identity in 279 aa overlap

	20	30	40	50	60	70
GLELO	FMDLATAIGVRAAPYVDPLEAALVAQAEKYIPTIVHHTRGFLVAVESPLAREL-----PL					
				:		:
U97107				MDTSMNFSRGLKMDLMQPYDFETFQDLRPF		
				10	20	30
	80	90	100	110	120	129
GLELO	MNPFHV--LLIVLAYLVTVFVGMQIMKNFERFEVKTFSLLNHFCVLSISAYMCGGILYEA					
	:	:	:	:	:	:
U97107	LEEYWVSSFLIVVVYLLLVGGQTYMRTRKSFSLQRPLILWSFFLAIFS--ILGTLRMWK					
	40	50	60	70	80	
	130	140	150	160	170	180
GLELO	YQAN---YGLFENAADHTFKGLPMAMKIW--LFYFSKIMEFVDTMIMVLKKNRQISFL					
	:	:	:	:	:	:
U97107	FMATVMFTVGLKQTVCFAIYTDDAVVRF-WSFLFLLSKVVELGDTAFIILRK--RPLIFV					
	90	100	110	120	130	140
	190	200	210	220	230	240
GLELO	HVYHHSSI--FTIWWLVTFVAPNGEAYFSAALNSFIHVI MYGYFLSALGFKQVSFIKFY					
U97107	HWYHHSTVLLFTSFGYKNKV-PSGGWFMT--MNFGVHSV MYTYT MKA AKL KHPNLLPMV					
	150	160	170	180	190	200
	250	260	270	280	290	
GLELO	ITRSQMTQFCMMSVQSSWDMYAMKVLG--RPGYPFFITALLWFYMWMTMLGLFYN--FYRK					
U97107	ITSLQILQMVLGTIFGILNYIWRQEKGCCHTTTEHFFWSFMLYGT YFILFAHFFH RAYLRP					
	210	220	230	240	250	260
	300	310				
GLELO	NAKLAKQAKADAAKEKARKLQ					
	:	:	:	:	:	:
U97107	KGKVASKSQ					
	270					

FIG.38

SCORES Init1: 100 Initn: 205 Opt: 271

Smith-Waterman score: 271; 30.7% identity in 218 aa overlap

	60	70	80	90	100	110
GLELO	TRGFLVAVESPLARELPLMNP	FHVLLIVLAYLVT	FVGMQIMKNFERFEV	KTFSL	LLHNFC	
			: :	: :	: :	:
U68749	ATHGPKNFPDAEGRKFFADHFDVTIQASILY	MVVVFGTKWFM	RNRQPFQLTIPLNIWNFI			
(F56H11.4)	30	40	50	60	70	80
	120	130	140	150	160	
GLELO	LVSISAYMCGGILYEAYQ--ANYGL---	FENAADHTFKGLPM	AKMIWLFYFSKIMEF	VDT		
	::	: :	:	:::		: :: : : : :
U68749	LAAFSIAGAVKMTPEFFGTIAN	KGIVASYCKVDF	T-KG-ENGYWWLF	MASKLFEL	VDT	
(F56H11.4)	90	100	110	120	130	140
	170	180	190	200	210	220
GLELO	MIMVLKKNRQISFLHVVYHSSIFTIWWL	VT	FVAPNGEAYFSAALNSFIHVIMYGY	FLS		
	::: :	:		:	:	: : : : :
U68749	IFLVLRK--RPLMFLHWYHHIL	TMIYAWYSHPL	T-P-GFNRYGIYLN	FVVHAFMYSY	FLR	
(F56H11.4)	150	160	170	180	190	
	230	240	250	260	270	280
GLELO	ALGFKQVSFIKFYITRSQMTQFCMMSVQSSW	DMYAMKVLGRP-	GYPFFITALLWFY	MWTM		
	:::	:		: :	:	: : : : :
U68749	SMKIRVPGFIAQAITS	LQIVQFIISCAVLAHLGYLMHFT	NANCD	FEP	SFKLAVFMD	TTY
(F56H11.4)	200	210	220	230	240	250
	290	300	310			
GLELO	LGLFYNFYRKN	LAKQAKADA	AKEKARKLQ			
	:	:				
U68749	LALFVNFFLQSYVLR	GGKDKYKAV	PKKKNN			
(F56H11.4)	260	270	280			

FIG.39

SCORES Init1: 77 Initn: 155 Opt: 264
 Smith-Waterman score: 264; 27.2% identity in 206 aa overlap

	40	50	60	70	80	90
GLELO	AALVAQAEKYIPTIVHHTRGFLVAVESPLARELPLMNPFHVLLIVLAYLVTVFVGMQIMK					
				: : ::: :: :		
DM1	PTKMINMDISVTPNYSYIFDFENDFIHQRTWKWLENWTWVFYCGIYMLVIFGGQHFMQ					
	10	20	30	40	50	60
	100	110	120	130	140	150
GLELO	NFERFEVKTFSLHNFCLVSISAYMCGGILYEAYQA--NYGLFENAADHTF--KGLPMAK					
	::: :: : : : : :: :: ::: :: : :					
DM1	NRPRFQLRGPLIIWNTLLAMFSIMGAARTAPELIHVLRYHGLFHSVCVPSYIEQDRVCGF					
	70	80	90	100	110	120
	160	170	180	190	200	210
GLELO	MIWLFYFSKIMEFVDTMIMVLKKNRQISFLHVVYHSSIFTIWWLVTFVAPNGEAYFSAA					
	: : : :: : : : :: ::: :: : :					
DM1	WTWLFVLSKLPGLDTIFIVLRK--QPLIFLHWYHHITVLIYSWF-SYTEYTSSARWFIV					
	130	140	150	160	170	180
	220	230	240	250	260	270
GLELO	LNSFIHVIMYGYFLSALGFKQVSFIKFYITRSQMTQFCMMSVQSSWDMYAMKVLGRPGY					
	: : : : : ::: :: : : : : : :					
DM1	MNYCVHSMYSYYALKAAARNPPRFISMIITSLQLAQMIIGCAINWANGFLKTHGTXS					
	190	200	210	220	230	240
	280	290	300	310		
GLELO	PFFITALLWFYMTMLGLFYNFYRKNALAKQAKADAAKEKARKLQ					
DM1	HISQRNINLSIAMYSSYFVLFARFFYKAYLAPGGHKSRRMA					
	250	260	270	280		

FIG.41

SCORES Init1: 181 Initn: 279 Opt: 328
 Smith-Waterman score: 328; 30.0% identity in 237 aa overlap

	40	50	60	70	80	90
MAELO	VTGKSIDSFVFQEGVTPSTQREVAMWTITYFVVIFGGRQIMKSQDAFKLKPLFILHNFL					
				::	::	::
DM1	IFDFENDFIHQRTKRWMLNWTWVFYCGIYMLVIFGGQHFMQNRPRFQLRGPLIIWNTL					
	30	40	50	60	70	80
	100	110	120	130	140	149
MAELO	LTIASGSLLLFIENLVPILARNGLFYAICDDGAWTQ-RLELLY-YLNYLVKYWELADTV					
	::	:	:	:	:	:
DM1	LAMFSINGAARTAPELIHVLRYGLFHSVCVPSYIEQDRVCGFWTLFVLSKLPGLGDTI					
	90	100	110	120	130	140
	150	160	170	180	190	200
MAELO	FLVLKKKPLEFLHYFHSMTMVLCFVQLGGYTS-VSWVPITLNLTVHVFMYYYMRSAAG					
	::	::	::	::	::	::
DM1	FIVLRKQPLIFLHWYHHITVLIYSWFSYTEYTSSARWF-IVMNYCVHSMYSYYALKAAAR					
	150	160	170	180	190	200
	210	220	230	240	250	260
MAELO	VRI--WWKQYLTLQIVQFVLDLGFIFYFCAYTYFAFTYFPWAPNVGKCAGTEGAALFGCG					
	:	:	:	:	:	:
DM1	FNPPRFISMIITSLQLAQMIIG-----CAINVWANGFLK-THGTXSCHISQRNINLSIA					
	210	220	230	240	250	
	270	280	290	300	310	
MAELO	LLSSYLLLFINFYRITYNAKAKAAKERGSNFTPKTVKSGGSPKKPSKSKHI					
	:	:	:	:	:	:
DM1	MYSSYFVLFAFFYKAYLAPGGHKSRRMA					
	260	270	280			

FIG.42

```
1  ATGGAACATT TTGATGCATC ACTTAGTACC TATTTCAAGG CATTGCTAGG
51 CCCTCGAGAT ACTAGAGTAA AAGGATGGTT TCTTCTGGAC AATTATATAC
101 CCACATTTAT CTGCTCTGTC ATATATTTAC TAATTGTATG GCTGGGACCA
151 AAATACATGA GGAATAAACA GCCATTCTCT TGCCGGGGGA TTTTAGTGGT
201 GTATAACCTT GGA CTCACAC TGCTGTCTCT GTATATGTTT TGTGAGTTAG
251 TAACAGGAGT ATGGGAAGGC AAATACAAC TCTTCTGTCA GGGCACACGC
301 ACCGCAGGAG AATCAGATAT GAAGATTATC CGTGTCTCT GGTGGTACTA
351 CTTCTCCAAA CTCATAGAAT TTATGGACAC TTTCTTCTTC ATCCTGCGCA
401 AGAACAACCA CCAGATCACG GTCCTGCACG TCTACCACCA TGCCTCGATG
451 CTGAACATCT GGTGGTTTGT GATGAACTGG GTCCCCTGCG GCCACTCTTA
501 TTTTGGTGCC AACTTAATA GCTTCATCCA CGTCTCATG TACTCTTACT
551 ATGGTTTGTC GTCAGTCCCT TCCATGCGTC CATACTCTG GTGGAAGAAG
601 TACATCACTC AGGGGCAGCT GCTTCAGTTT GTGCTGACAA TCATCCAGAC
651 CAGCTGCGGG GTCATCTGGC CGTGCACATT CCCTCTTGGT TGGTTGTATT
701 TCCAGATTGG ATACATTATT TCCCTGATTG CTCTCTTCAC AAACCTCTAC
751 ATTCAGACCT ACAACAAGAA AGGGGCCTCC CGAAGGAAAG ACCACCTGAA
801 GGACCACCAG AATGGGTCCG TGGCTGCTGT GAATGGACAC ACCAACAGCT
851 TTTCACCCCT GGAAAACAAT GTGAAGCCAA GGAAGCTGCG GAAGGATTGA
901 AGTCAAAGAA TTGA
```

FIG.43

1 MEHFDASLST YFKALLGPRD TRVKGWFLD NYIPTFICSV IYLLIVWLGP
51 KYMRNKQPFS CRGILVVYNL GLTLLSLYMF CELVTGVWEG KYNFFCQGTR
101 TAGESDMKII RVLWWYYFSK LIEFMDTFFF ILRKNNHQIT VLHVYHHASM
151 LNIWWFVMNW VPCGHSYFGA TLNSFIHVLM YSYYGLSSVP SMRPYLWWKK
201 YITQGQLLQF VLTIIQTSCG VIWPCTFPLG WLYFQIGYII SLIALFTNFI
251 IQTYNKKGAS RRKDHLKDHQ NGSVAAVNGH TNSFSPLENN VKPRKLRKD*

FIG.44

HOST(PLASMID)	334(pYX242)	334(pRAE-58-A1)	334(pYX242)	334(pRAE-58-A1)
ADDED SUBSTRATE	25 μM GLA	25 μM GLA	25 μM AA	25 μM AA
FATTY ACID	% TOTAL FATTY ACID	% TOTAL FATTY ACID	% TOTAL FATTY ACID	% TOTAL FATTY ACID
C18:3n-6	4.40	2.71	0.03	0.04
C20:3n-6	0.09	(50.34%)* 2.75	0.02	0.02
C20:4n-6			7.48	3.97
C22:4n-6			ND	(23.37%)* 1.21
C16:1n-7	41.11	34.72	41.49	35.07
C18:1n-7	1.85	11.33	2.01	11.57
C20:1n-7	0.04	1.48	0.04	1.62
C18:1n-9	15.60	15.66	15.16	14.57
C20:1n-9	0.06	0.22	0.06	0.23
C18:1n-5	0.11	0.62	0.12	0.58
TOTAL LIPID	370	969	359	514
*% CONVERSION=PRODUCT/(SUBSTRATE+PRODUCT)				

FIG.45

```
1  ATGGCTCAGC ATCCGCTCGT TCAACGGCTT CTCGATGTCA AATTCGACAC
51  GAAACGATTT GTGGCTATTG CTAATCATGG GCCAAAGAAT TTCCCTGACG
101 CAGAAGGTCG CAAGTTCTTT GCTGATCACT TTGATGTTAC TATTCAGGCT
151 TCAATCCTGT ACATGGTCGT TGTGTTCCGA ACAAATGGT TCATGCGTAA
201 TCGTCAACCA TTCCAATTGA CTATTCCACT CAACATCTGG AATTTTCATCC
251 TCGCCGCATT TTCCATCGCA GGAGCTGTCA AAATGACCCC AGAGTTCTTT
301 GGAACCATTG CCAACAAAGG AATTGTCGCA TCCTACTGCA AAGTGTTTGA
351 TTTCACGAAA GGAGAGAATG GATACTGGGT GTGGCTCTTC ATGGCTTCCA
401 AACTTTTCGA ACTTGTTGAC ACCATCTTCT TGGTTCTCCG TAAACGTCCA
451 CTCATGTTCC TTCCTGGTA TCACCATATT CTCACCATGA TCTACGCCTG
501 GTAATCTCAT CCATTGACCC CAGGATTCAA CAGATACGGA ATTTATCTTA
551 ACTTTGTCGT CCACGCCTTC ATGTAATCTT ACTACTTCCT TCGCTCGATG
601 AAGATTCGCG TGCCAGGATT CATCGCCCAA GCTATCACAT CTCTTCAAAT
651 CGTTCAATTC ATCATCTCTT GCGCCGTTCT TGCTCATCTT GGTTATCTCA
701 TGCACTTCAC CAATGCCAAC TGTGATTTCT AGCCATCAGT ATTCAAGCTC
751 GCAGTTTTCA TGGACACAAC ATACTTGGCT CTTTTCGTCA ACTTCTTCCT
801 CCAATCATAT GTTCTCCGCG GAGGAAAAGA CAAGTACAAG GCAGTGCCAA
851 AGAAGAAGAA CAACTAA
```

FIG.46

1 MAQHPLVQRL LDVKFDTKRF VAIATHGPKN FPDAEGRKFF ADHFDVTIQA
51 SILYMVVVFG TKWFMRNRQP FQLTIPLNIW NFILAAFSIA GAVKMTPEFF
101 GTIANKGIVA SYCKVFDFTK GENGYVWVLF MASKLFELVD TIFLVLRKRP
151 LMFLHWYHHI LTMIYAWYSH PLTPGFNRYG IYLNFFVHAF MYSYYFLRSM
201 KIRVPGFIAQ AITSLQIVQF IISCAVLAHL GYLMHFTNAN CDFEPSVFKL
251 AVFMDTTYLA LFVNFFLQSY VLRGGKDKYK AVPKKKNN

FIG.47

HOST(PLASMID)	334(pYX242)	334(pRET-21)	334(pRET-22)
ADDED SUBSTRATES	50 μ M GLA + 50 μ M AA	50 μ M GLA + 50 μ M AA	50 μ M GLA + 50 μ M AA
FATTY ACID	% TOTAL FATTY ACID	% TOTAL FATTY ACID	% TOTAL FATTY ACID
C16:0	9.22	12.46	9.9
C16:1	0.09	0.18	0.13
C18:0	1.46	2.41	1.49
C18:1n-9	4.03	4.92	3.91
C18:3n-6	10.02	11.89	8.69
C20:3n-6	(1.28%)* 0.13	(11.1%)* 1.48	(19.4%)* 2.09
C20:4n-6	46.98	28.87	35.25
C20:4n-6	0	0	0
TOTAL LIPID (mg)	212	174	187
*% CONVERSION=PRODUCT/(SUBSTRATE+PRODUCT)			

FIG.48

1 ATGAACATGT CAGTGTTGAC TTTACAAGAA TATGAATTCG AAAAGCAGTT
51 CAACGAGAAT GAAGCCATCC AATGGATGCA GGAAAACTGG AAGAAATCTT
101 TCCTGTTTTT TGCTCTGTAT GCTGCCTTTA TATTCGGTGG TCGGCACCTA
151 ATGAATAAAC GAGCAAAGTT TGAAGTGAAG AAGCCATTAG TGCTCTGGTC
201 TCTGACCCTT GCAGTCTTCA GTATATTCGG TGCTCTTCGA ACTGGTGCTT
251 ATATGGTGTA CATTTTGATG ACCAAAGGCC TGAAGCAGTC AGTTTGTGAC
301 CAGGGTTTTT ACAATGGACC TGTCAGCAAA TTCTGGGCTT ATGCATTTGT
351 GCTAAGCAAA GCACCCGAAC TAGGAGATAC AATATTCATT ATTCTGAGGA
401 AGCAGAAGCT GATCTTCCTG CACTGGTATC ACCACATCAC TGTGCTCCTG
451 TACTCTTGGT ACTCCTACAA AGACATGGTT GCCGGGGGAG GTTGGTTCAT
501 GACTATGAAC TATGGCGTGC ACGCCGTGAT GTACTCTTAC TATGCCTTGC
551 GGGCGGCAGG TTTCCGAGTC TCCCGGAAGT TTGCCATGTT CATCACCTTG
601 TCCCAGATCA CTCAGATGCT GATGGGCTGT GTGGTTAACT ACCTGGTCTT
651 CTGCTGGATG CAGCATGACC AGTGTCAC TCACTTTCAG AACATCTTCT
701 GGTCTCACT CATGTACCTC AGCTACCTTG TGCTCTTCTG CCATTTCTTC
751 TTTGAGGCCT ACATCGGCAA AATGAGGAAA ACAACGAAAG CTGAATAG

FIG.49

1 MNMSVLTQE YEFKQFNEN EAIQWMQENW KKSFLFSALY AAFIFGGRHL
51 MNKRAKFELR KPLVLWSLTL AVFSIFGALR TGAYMVYILM TKGLKQSVCD
101 QGFYNGPVSF FWAYAFVLSK APELGDTIFI ILRKQKLIFL HWYHHITVLL
151 YSWYSYKDMV AGGGWFMTMN YGVHAVMYSY YALRAAGFRV SRKFAMFITL
201 SQITQMLMGC VVNYLVFCWM QHDQCHSHFQ NIFWSSLMYL SYLVLFCHFF
251 FEAYIGKMRK TTKAE*

FIG.50

HOST	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334
PLASMD	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pRAE-58
SUBSTRATE	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA
CONCENTRATION	25 μM	25 μM	100 μM	100 μM	25 μM	25 μM	25 μM	25 μM	100 μM	100 μM	25 μM	25 μM	25 μM	25 μM	100 μM	100 μM	100 μM
% TOTAL LIPID																	
C18:1n-9	23.82	21.49	18.49	17.41	22.09	19.23	17.45	18.44	24.78	21.28	19.42	18.85					
C18:1n-7	2.52	18.35	1.71	11.82	2.54	18.77	1.78	12.67	2.64	19.48	1.79	12.40					
C18:1n-5	0.15	1.13	0.10	0.54	0.15	1.23	0.10	0.63	0.15	1.18	0.09	0.62					
C18:3n-6	6.10	2.38	23.30	14.46	0.04	0.02	0.04	0.02	0.04	0.02	0.01	0.01					
C20:1n-9	0.08	0.83	0.05	0.48	0.10	1.18	0.04	0.56	0.10	1.30	0.06	0.63					
C20:1n-7	0.10	5.75	0.07	3.09	0.11	9.49	0.05	3.62	0.10	9.94	0.08	4.07					
C20:3n-6	0.15	(62.4%)3.95	0.31	(39.8%)9.56	0.02	ND	ND	0.04	ND	0.02	0.01	0.01					
C20:4n-6	ND	ND	0.01	ND	11.76	7.68	28.39	21.02	0.02	0.02	ND	0.01					
C20:5n-3	ND	ND	ND	ND	0.03	0.02	0.10	0.07	4.79	2.04	26.47	13.69					
C22:4n-6	ND	ND	ND	ND	ND	(27.5%)2.91	0.01	(15.7%)3.90	ND	ND	0.00	ND					
C22:5n-3	ND	ND	ND	ND	ND	ND	ND	0.03	0.02	(70.3%)4.82	0.04	(45.7%)11.50					
TOTAL LIPID	230	419	590	576	249	332	1014	961	372	390	1323	1065					

(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)
ND = NOT DETECTED

FIG.52

HOST	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334
PLASMID	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58
SUBSTRATE	PA	PA	SA	SA	ARA	ARA	BA	BA	PTA	PTA	OA	OA	OA	OA	EA	EA
CONCENTRATION	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM
% TOTAL LIPID																
C16:0	24.17	17.23	11.22	7.90	7.74	7.98	7.62	7.11	17.28	11.04	16.06	12.76	14.37	11.98		
C16:1n-7	39.83	33.83	30.62	20.56	21.61	19.81	21.34	22.89	50.06	39.43	40.95	30.06	43.34	29.51		
C16:1n-5	0.30	0.74	0.29	0.58	0.17	0.47	0.18	0.59	0.38	0.80	0.34	0.68	0.37	0.71		
C18:0	1.90	1.50	35.82	38.10	1.12	0.89	1.03	0.88	1.90	1.44	1.82	1.43	1.51	1.23		
C18:1n-9	15.36	14.11	11.52	10.88	8.29	10.03	8.09	10.25	14.55	13.86	20.12	21.37	14.12	15.15		
C18:1n-7	1.36	11.44	0.90	8.72	0.69	8.51	0.69	8.58	1.30	12.76	1.30	13.79	1.21	12.66		
C18:1n-5	0.11	0.78	0.08	0.69	0.08	0.54	0.06	0.61	0.19	0.76	0.10	0.90	0.15	0.84		
C20:0	0.15	0.17	0.09	0.12	52.07	41.48	ND	ND	ND	ND	ND	ND	0.17	0.23		
C20:1n-9	0.09	0.45	0.05	0.30	0.03	ND	0.06	0.28	0.05	0.38	0.18	0.58	7.47	10.97		
C20:1n-7	0.20	2.84	ND	1.52	0.05	1.43	0.14	1.60	0.07	2.76	0.12	2.08	ND	2.30		
C22:0	0.43	0.56	0.29	0.22	0.31	0.19	52.91	38.43	ND	ND	ND	ND	ND	0.32		
C24:0	0.59	1.39	0.36	0.85	0.45	0.71	0.53	1.14	0.45	1.63	0.66	1.02	0.56	0.79		
TOTAL LIPID	297	272	573	542	558	846	585	519	464	295	306	448	309	648		

ND = NOT DETECTED

FIG.53A

HOST	334	334	334	334	334	334	334	334	334	334	334
PLASMID	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pRAE-58	pYX242	pYX242	pRAE-58
SUBSTRATE	LA	LA	GLA	GLA	DGLA	DGLA	AA	AA	ADA	ADA	ADA
CONCENTRATION	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM
	% TOTAL LIPID										
C18:1n-9	15.27	16.83	14.85	15.58	13.62	16.24	15.08	15.64	16.18	13.98	
C18:1n-7	1.21	13.53	1.22	11.80	1.16	12.63	1.18	11.70	1.30	10.67	
C18:1n-5	0.13	0.95	0.20	0.73	0.12	0.72	0.14	0.59	0.12	0.70	
C18:2n-6	4.09	4.85	0.09	0.07	0.07	0.04	0.04	0.04	0.03	0.07	
C18:3n-6	ND	ND	4.66	2.33	ND	ND	ND	ND	ND	ND	
C20:1n-9	0.07	2.60	0.07	0.33	0.07	0.33	0.04	0.27	0.08	0.33	
C20:1n-7	0.10	0.18	0.14	1.65	0.08	1.68	0.12	1.58	0.12	1.85	
C20:2n-6	ND (13.2%)	0.74	ND	ND	ND	ND	ND	ND	ND	ND	
C20:3n-6	ND	ND	ND (51.4%)	2.46	6.37	7.86	ND	0.03	ND	ND	
C20:4n-6	ND	ND	ND	ND	ND	0.09	6.49	5.77	ND	ND	
C22:4n-6	ND	ND	ND	ND	ND	ND	ND (27.1%)	2.14	10.91	15.57	
C24:0	0.59	1.61	0.64	1.12	0.69	0.79	0.52	0.77	0.54	1.26	
TOTAL LIPID	333	373	260	392	260	672	553	690	706	440	

(%CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)
ND = NOT DETECTED

FIG.53B

HOST	334	334	334	334	334	334	334	334	334	334	334	334
PLASMID	pYX242	pRAE-58	pYX242	pYX242	pRAE-58	pYX242	pYX242	pRAE-58	pYX242	pYX242	pYX242	pRAE-58
SUBSTRATE	ALA	ALA	STA	STA	STA	EPA	DPA	DPA	DPA	DPA	DPA	pRAE-58
CONCENTRATION	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	
% TOTAL LIPID												
C18:1n-9	17.21	17.36	16.85	17.71	16.45	16.93	17.08	16.68	18.36	18.36	18.36	18.77
C18:1n-7	1.29	12.20	1.15	11.38	1.23	11.48	1.33	11.61	1.46	1.46	1.46	13.72
C18:1n-5	0.14	0.68	0.12	0.57	0.12	0.54	0.12	0.63	0.13	0.13	0.13	0.79
C18:3n-3	4.42	3.61	ND	0.03	ND	0.03	ND	0.03	ND	ND	ND	0.03
C18:4n-3	ND	0.13	3.04	1.38	ND	0.13	ND	0.13	ND	ND	ND	0.17
C20:1n-9	0.09	0.33	0.11	0.34	0.05	0.31	0.09	0.30	0.13	0.13	0.13	0.34
C20:1n-7	0.13	1.55	0.05	1.38	0.23	1.89	0.18	1.73	0.15	0.15	0.15	1.76
C20:3n-3	0.06	(22.2%)1.03	ND	ND	ND	0.11	ND	ND	ND	ND	ND	ND
C20:4n-3	ND	ND	0.06	(61.9%)2.24	ND	ND	ND	ND	ND	ND	ND	ND
C20:5n-3	ND	ND	0.05	0.05	7.43	4.88	ND	ND	0.07	0.07	0.07	ND
C22:4n-3	ND	ND	ND	0.39	ND	ND	ND	ND	ND	ND	ND	ND
C22:5n-6	ND	ND	ND	ND	ND	ND	0.28	0.41	ND	ND	ND	ND
C22:5n-3	ND	ND	ND	ND	ND	(39.5%)3.19	3.99	5.94	ND	ND	ND	ND
C24:0	0.43	0.73	0.33	0.73	0.45	0.84	0.64	1.07	0.68	0.68	0.68	0.77
C24:5n-3	ND	ND	ND	ND	ND	0.08	ND	0.06	ND	ND	ND	ND
TOTAL LIPID	696	729	911	710	719	703	602	642	397	397	397	684

(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)
ND=NOT DETECTED

FIG.53C

```
1  ATGGAGCAGC TGAAGGCCTT TGATAATGAA GTCAATGCTT TCTTGGACAA
51 CATGTTTGGA CCACGAGATT CTCGAGTTCG CGGGTGGTTC CTGCTGGACT
101 CTTACCTTCC CACCTTCATC CTCACCATCA CGTACCTGCT CTCGATATGG
151 CTGGGTAACA AGTACATGAA GAACAGGCCT GCTCTGTCTC TCAGGGGCAT
201 CCTCACCTTG TATAACCTCG CAATCACACT TCTTTCTGCG TATATGCTGG
251 TGGAGCTCAT CCTCTCCAGC TGGGAAGGAG GTTACAACCT GCAGTGTGAG
301 AATCTCGACA GTGCAGGAGA AGGTGATGTC CGGGTAGCCA AGGTCTTGTG
351 GTGGTACTAC TTCTCCAAAC TAGTGGAGTT CCTGGACACG ATTTTCTTTG
401 TTCTACGAAA AAAGACCAAT CAGATCACCT TCCTTCATGT CTATCACCAC
451 GCGTCCATGT TCAACATCTG GTGGTGTGTT TTGAACTGGA TACCTTGTGG
501 TCAAAGCTTC TTTGGACCCA CCCTGAACAG CTTTATCCAC ATTCTCATGT
551 ACTCCTACTA CGGCCTGTCT GTGTTCCCGT CCATGCACAA GTACCTTTGG
601 TGGAAGAAGT ACCTCACACA GGCTCAGCTG GTGCAGTTCG TACTCACCAT
651 CACGCACACG CTGAGTGCCG TGGTGAAGCC CTGTGGCTTC CCCTTTGGCT
701 GTCTCATCTT CCAGTCTTCC TATATGATGA CGCTGGTCAT CCTGTTCTTA
751 AACTTCTATA TTCAGACATA CCGGAAAAAG CCAGTGAAGA AAGAGCTGCA
801 AGAGAAAGAA GTGAAGAATG GTTTCCCCAA AGCCCACTTA ATTGTGGCTA
851 ATGGCATGAC GGACAAGAAG GCTCAATAA
```

FIG.54

1 MEQLKAFDNE VNAFLDNMFG PRDSRVRGWF LLDSYLPTEI LTITYLLSIW
51 LGNKYMKNRP ALSLRGILT YNLAITLLSA YMLVELILSS WEGGYNLQCQ
101 NLDSAGEGDV RVAKVLWYY FSKLVEFLDT IFFVLRKKTN QITFLHVVHH
151 ASMFNIWWCV LNWIPCGQSF FGPTLNSFIH ILMYSYYGLS VFPSMHKYLW
201 WKKYLTQAQL VQFVLTITHT LSAVVKPCGF PFGCLIFQSS YMTLVILFL
251 NFYIQTYRKK PVKKELQEKE VKNGFPKAHL IVANGMTDKK AQ*

FIG.55

HOST	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334
PLASMIID	pX242	pRAE-84	pX242	pRAE-84	pX242	pRAE-84	pX242	pRAE-84	pX242	pRAE-84	pX242	pX242	pRAE-84	pX242	pRAE-84
SUBSTRATE	GIA	GIA	AA	AA	ADA	ADA	ADA	ADA	ADA	ADA	ADA	ADA	EPA	DPA	DPA
CONCENTRATION	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM
% TOTAL LIPID															
C18:1n-9	15.94	14.16	12.30	15.67	11.77	11.41	14.81	17.92	15.91	16.33	15.04	14.63			
C18:1n-7	1.25	1.21	1.10	1.50	1.13	1.18	1.19	1.38	1.33	1.49	1.37	1.38			
C18:3n-6	4.53	4.21	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND			
C18:4n-3	ND	ND	ND	ND	ND	ND	2.78	2.70	ND	ND	ND	ND			
C20:1n-7	ND	ND	ND	ND	ND	0.32	ND	0.03	ND	0.05	ND	ND			
C20:3n-6	0.10	0.37	ND	ND	ND	ND	ND	0.05	ND	ND	ND	ND			
C20:4n-6	ND	ND	11.44	5.55	ND	ND	ND	ND	ND	ND	ND	ND			
C20:4n-3	ND	ND	ND	ND	ND	ND	ND	(14%)0.44	ND	ND	ND	ND			
C20:5n-3	ND	ND	ND	ND	ND	23.61	ND	ND	9.68	3.02	ND	ND			
C22:4n-6	ND	ND	ND	(10.4%)0.64	20.41	ND	ND	ND	ND	ND	ND	ND			
C22:4n-3	ND	ND	ND	ND	ND	ND	ND	(42.3%)0.33	ND	ND	0.57	0.57			
C22:5n-3	ND	ND	ND	ND	ND	ND	ND	ND	ND	(32.7%)1.47	7.87	4.88			
C24:4n-6	ND	ND	ND	(62.6%)1.07	ND	(9.2%)2.4	ND	ND	ND	ND	ND	ND			
C24:5n-3	ND	ND	ND	ND	ND	ND	ND	ND	ND	(82.8%)7.06	ND	(43.9%)3.82			
TOTAL LIPID	208	126	115	189	158	149	124	433	221	271	127	126			

(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)
ND = NOT DETECTED

FIG. 56

HOST	334	334	334	334	334	334	334	334	334	334	334	334	334	334	334
PLASMID	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242
SUBSTRATE	PA	SA	SA	SA	BA	BA	BA	BA	PTA	PTA	OA	OA	EA	EA	EA
CONCENTRATION	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM	25µM
C16:0	36.30	39.95	7.12	8.31	5.78	4.42	4.17	5.76	18.69	18.85	14.69	18.91	15.25	18.88	18.88
C16:1n-7	26.22	23.52	11.77	15.25	10.23	6.29	7.01	10.10	38.48	41.23	20.55	31.48	25.89	40.32	40.32
C16:1n-5	0.23	0.28	0.16	0.20	0.13	0.07	0.09	0.13	0.38	0.38	0.26	0.43	0.35	0.41	0.41
C18:0	2.26	2.14	64.90	58.73	0.94	1.01	0.64	0.85	2.17	2.29	3.02	2.73	2.71	2.15	2.15
C18:1n-9	14.83	11.27	6.35	7.22	5.20	4.33	3.84	5.12	14.25	14.27	18.44	22.20	14.62	16.91	16.91
C18:1n-7	1.44	1.36	0.57	0.73	0.54	0.51	0.41	0.56	1.57	1.68	1.53	1.67	1.65	1.84	1.84
C18:1n-5	0.10	ND	ND	0.06	ND	ND	ND	0.06	0.17	0.15	ND	0.18	ND	0.16	0.16
C20:0	0.59	0.24	0.09	0.08	66.40	74.78	0.10	0.05	0.17	0.17	0.24	0.20	0.33	0.04	0.04
C20:1n-9	0.06	0.10	ND	0.04	0.05	0.06	ND	ND	ND	ND	0.25	0.16	13.15	7.07	7.07
C20:1n-7	0.07	ND	ND	ND	ND	0.12	ND	ND	ND	ND	0.40	ND	ND	0.04	0.04
C22:0	0.45	0.75	0.29	0.30	0.43	0.31	77.35	70.71	0.74	0.80	0.98	0.74	0.83	0.44	0.44
C24:0	0.55	1.09	0.38	0.41	0.69	0.62	0.50	0.45	0.94	0.92	1.67	0.96	ND	0.53	0.53
TOTAL LIPID	158	104	144	112	324	209	178	94	148	87	243	315	70	529	529

ND = NOT DETECTED

FIG. 57A

HOST	334	334	334	334	334	334	334	334	334	334	334	334
PLASMID	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84	pYX242	pRAE-84
SUBSTRATE	LA	LA	GLA	GLA	DGLA	DGLA	DGLA	DGLA	AA	AA	ADA	ADA
CONCENTRATION	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM
% TOTAL LIPID												
C18:1n-9	12.30	16.12	15.63	16.28	14.28	13.77	16.21	15.04	15.38	15.38	15.38	12.94
C18:1n-7	1.34	1.87	1.69	1.90	1.41	1.61	1.61	1.62	1.51	1.51	1.51	1.47
C18:2n-6	2.67	3.61	0.17	0.20	0.24	0.21	0.09	0.09	0.06	0.06	0.06	0.14
C18:3n-6	ND	ND	2.03	2.49	ND	ND	ND	ND	ND	ND	ND	ND
C20:3n-6	ND	ND	ND	(14.7%)0.43	10.59	10.73	ND	ND	ND	ND	ND	ND
C20:4n-6	ND	ND	ND	ND	ND	ND	14.03	5.27	ND	ND	ND	ND
C22:4n-6	ND	ND	ND	ND	ND	ND	ND	(8.7%)0.5	11.44	11.44	11.44	16.60
C24:0	0.79	1.00	1.08	1.16	1.30	0.87	0.87	0.72	0.77	0.77	0.77	1.18
C24:4n-6	ND	ND	ND	ND	ND	ND	ND	(43.8%)0.39	ND	ND	ND	(7.3%)1.3
C24:5n-6	ND	ND	ND	ND	ND	ND	ND	0.38	ND	ND	ND	ND
TOTAL LIPID	85	87	88	79	107	98	208	212	304	304	304	122

(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)
ND = NOT DETECTED

FIG.57B

HOST	334	334	334	334	334	334	334	334	334	334	334	334
PLASMID	pYX242	pRAE-84	pYX242	pYX242	pRAE-84	pYX242	pYX242	pRAE-84	pYX242	pYX242	pRAE-84	pRAE-84
SUBSTRATE	ALA	ALA	ALA	ALA	ALA	ALA	ALA	ALA	ALA	ALA	ALA	ALA
CONCENTRATION	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM	25μM
% TOTAL LIPID												
C18:1n-9	16.69	16.38	18.24	15.95	14.07	15.16	16.05	15.06	17.47	17.15	17.15	17.15
C18:1n-7	1.37	1.43	1.71	1.40	1.37	1.47	1.67	1.51	1.75	1.73	1.73	1.73
C18:2n-6	0.08	0.08	0.12	0.04	0.13	0.06	0.11	0.18	0.13	0.15	0.15	0.15
C18:3n-3	4.47	4.28	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
C18:4n-3	ND	ND	2.28	2.39	ND	ND	ND	ND	ND	ND	ND	ND
C20:3n-3	(1.3%)0.06	(3.6%)0.16	ND	ND	ND	0.26	ND	ND	ND	0.12	ND	ND
C20:4n-3	ND	ND	ND	(11.1%)0.3	ND	ND	ND	ND	ND	ND	ND	ND
C20:5n-3	ND	0.07	ND	ND	9.97	3.84	ND	ND	ND	ND	ND	ND
C22:4n-3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
C22:5n-6	ND	ND	ND	(43.4%)0.23	ND	ND	0.64	0.55	ND	ND	ND	ND
C22:5n-3	ND	ND	ND	ND	ND	(24.0%)1.21	8.79	3.57	ND	ND	ND	ND
C24:0	0.65	0.43	1.41	0.58	1.38	0.78	1.45	1.35	0.89	0.67	0.67	0.67
C24:5n-3	ND	ND	ND	ND	ND	(73.6%)3.38	ND	(46.4%)3.09	ND	ND	ND	ND
TOTAL LIPID	362	384	173	393	124	280	137	151	190	200	200	200

(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)
ND = NOT DETECTED

FIG.57C

```
1  ATGGAACATT TCGATGCGTC ACTCAGTACC TATTTCAAGG CCTTCCTGGG
51  CCCCCGAGAT ACAAGAGTCA AAGGATGGTT CCTCCTGGAC AATTACATCC
101 CTACGTTTGT CTGTTCTGTT ATTTACTTAC TCATTGTATG GCTGGGACCA
151 AAATACATGA AGAACC GGCA GCCGTTCTCT TGCCGAGGCA TCCTGCAGTT
201 GTATAACCTT GGA CTACCC TGCTGTCTCT CTACATGTTC TATGAGTTGG
251 TGACAGGTGT GTGGGAGGGC AAATACA ACT TTTTCTGCCA GGAACACGC
301 AGCGCGGGAG AATCCGATAT GAAGATCATC CGCGTCCTCT GGTGGTACTA
351 CTTCTCCAAA CTCATCGAAT TCATGGACAC CTTTTTCTTC ATCCTTCGCA
401 AGAACAACCA CCAGATCACC GTGCTCCATG TCTACCACCA CGCTACCATG
451 CTCAACATCT GGTGGTTTGT GATGAACTGG GTTCCCTGCG GCCATTCATA
501 TTTTGGTGCG AACTCAACA GCTTCATCCA TGTCCTCATG TACTCGTACT
551 ATGGTCTGTC CTCCATCCCG TCCATGCGTC CCTACCTCTG GTGGAAAAAG
601 TACATCACTC AAGGGCAGCT GGTCCAGTTT GTGCTGACAA TCATCCAGAC
651 GACCTGCGGG GTCTTCTGGC CATGCTCCTT CCCTCTCGGG TGGCTGTTCT
701 TCCAGATTGG ATACATGATT TCCCTGATTG CTCTCTTCAC AAAGTTCTAC
751 ATTCAGACTT ACAACAAGAA AGGGGCTCT CGGAGGAAAG ACCACCTGAA
801 GGGCCACCAG AACGGGTCTG TGGCCGCCGT CAACGGACAC ACCAACAGCT
851 TCCCTTCCCT GGAAAACAGC GTGAAGCCCA GGAAGCAGCG AAAGGATTGA
```

FIG.58

1 MEHFDASLST YKAFGLGPRD TRVKGWFLD NYIPTFVCSV IYLLIVWLGP
51 KYMKNRQPFS CRGILQLYNL GLTLLSLYMF YELVTGVWEG KYNFFCQGTR
101 SAGESDMKII RVLWWYYFSK LIEFMDTFFF ILRKNNHQIT VLHVYHHATM
151 LNIWWFVMNW VPCGHSYFGA TLNSFIHVLM YSYYGLSSIP SMRPYLWKK
201 YITQGQLVQF VLTIIQTTCG VFWPCSFPLG WLFFQIGYMI SLIALFTNFY
251 IQTYNKKGAS RRKDHLKGHQ NGSVAAVNGH TNSFPSLENS VKPRKQRKD*

FIG.59

HOST(PLASMID)	334(pYX242)	334(pRET-22)	334(pYX242)	334(pRET-22)	334(pYX242)	334(pRET-22)
ADDED SUBSTRATES	50 mM GLA	50 mM GLA	50 mM AA	50 mM AA	NO SUBSTRATE	NO SUBSTRATE
	% TOTAL LIPID					
FATTY ACID						
C16:0	19.8	18.59	13.8	6.23	13.62	13.63
C16:1n-7	20.92	17.74	26.62	13.01	40.1	47.67
C18:0	5.79	4.94	3.62	2	4.86	5.031
C18:1n-7	(3.9%) 0.85	(9.12%) 1.78	(3.5%) 0.97	(12.54%) 1.18	(3.6%) 1.5	(7.53%) 3.88
C18:1n-9	8.46	7.45	10.27	5.36	13.7	16.93
C18:3n-6	*26.62	*22.03	0.03	0.01		
C20:3n-6	(1.1%) 0.3	(38.2%) 13.61				
C20:4n-6			*27.36	*65.38		
C22:4n-6						
TOTAL LIPID (μg)	36	42	85	280	55	79
(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)						
*INDICATES SUBSTRATE ADDED						

FIG.61

HOST(PLASMID)	334(pRET22)	334(pRET22)	334(pRET22)	334(pRET22)	334(pRET22)	334(pRET22)
ADDED SUBSTRATE	50μM SA	50μM OA	50μM LA	50μM DLGA	25μM AA	50μM ADRENIC
	C18:0	C18:1n-9	C18:2n-6	C20:3n-6	C20:4n-6	C22:4n-6
FATTY ACID	% TOTAL LIPID					
C16:0	12.9	12.54	15.23	9.1	10.2	3.42
C16:1	37.71	23.83	24.87	16.61	18.375	7.66
C18:0	11.44	4.7	4.49	2.7	2.9	1.23
C18:1n-9	14.03	*16.87	9.54	6.74	6.39	2.99
C18:2n-6			16.87		0.15	0.28
C18:3n-6						
C20:2n-6						0.05
C20:3n-6				*44.34		
C20:4n-6				0.34	*25.78	0.26
C22:4n-6						*75.72
TOTAL LIPID (μg)	63	103	71	110	97	277

FIG.62A

HOST(PLASMID)	334(pRET22)	334(pRET22)	334(pRET22)	334(pRET22)
ADDED SUBSTRATE	50 μ M ALA	50 μ M PA	50 μ M EPA	50 μ M STA
	C18:3n-3	C18:0	C20:5n-3	C18:4n-3
FATTY ACID	% TOTAL LIPID			
C16:0	13.91	15.06	16.92	20.08
C16:1	14.74	31.77	23.57	20.17
C18:0	4.06	*4.85	4.94	6.02
C18:1n-9	6.65	13.59	10.46	9.29
C18:3n-3	*38.66			
C18:4n-3				*20.45
C20:4n-3				(12.57%) 2.94
C20:5n-3				
C22:5n-3			*15.48	
TOTAL LIPID (μ g)	80	84	81	60

*INDICATES SUBSTRATE ADDED
(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.62B

HOST(PLASMID)	334(pRET-22+pCGR-4)	334(pYX242+pYES2)
ADDED SUBSTRATE	50 μ M GLA	50 μ M GLA
FATTY ACID	% TOTAL LIPID	
C16:0	15.92	15.07
C16:1n-7	24.97	19.48
C18:0	8.52	6.48
C18:1n-7	3.9	1.61
C18:1n-9	18.48	12.71
C18:3n-6	*7.0	*10.54
C20:0	0	0
C20:3n-6	(27.81%) 4.36	(1.58%) 0.17
C20:4n-6	(27.55%) 4.32	0
TOTAL LIPID (μ g)	508	168

FIG.63A

HOST(PLASMID)	334(pRET-22+pCGR-4)	334(pYX242+pYES2)
ADDED SUBSTRATE	50 μ M STA	50 μ M STA
FATTY ACID	% TOTAL LIPID	
C16:0	18.74	16.21
C16:1n-7	21.35	26.09
C18:0	6.78	7.57
C18:1n-7	1.97	1.7
C18:1n-9	20.73	22.41
C18:4n-3	*6.05	*13.43
C20:0	0	0.45
C20:4n-3	(15.88%) 1.68	(4.73%) 0.69
C20:5n-3	(26.93%) 2.85	(3.22%) 0.47
TOTAL LIPID (μ g)	335	161

*INDICATES SUBSTRATE ADDED

(% CONVERSION)=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.63B

HSEL01	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
--------	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

FIG. 64A

HSEL01	208	LQFVLTIIQTSCGV IWP - C - - - - TFPLGWL YFQI - GYMISL I A	244
MELO4	211	VQFVLTITHTLSAVVKP - C - - - - GFPGCLIFQS - SYMMTLLVI	247
GLELO	248	TQFCMMSVQSSWDMYAMKVLGRPGYPF - FITALLWFWMTMLG	289
CEELO	218	VQFIIISCAVLAHLGYLMHFTNANCDFEPSVFKLAVFMDTTYLA	260
HSEL01	245	LFTNFYIQTYNKKGASRRKDLKDHQNGSM AAVNGHTNSFSPL	287
MELO4	248	LFLENFYIQTYRKKPV - - - KKELE - - - K - EVKNGF PKAHLIV	282
GLELO	290	LFYNF - - - YRKNAKLAKQAKADAK EKARKLQ	318
CEELO	261	LFVNFQLSYVLRGGKDKYKAVPKKKN	288
HSEL01	288	ENN VKPRKL R K D	299
MELO4	283	ANGMTDKKAQ	292

FIG.64B

1 ACAGGCGACT TTCTGTTGAA TTTGTTGCAG GCACAGCAAG GAAGGATGGC
51 AAACAGCAGC GTGTGGGATG ATGTGGTGGG CCGCGTGGAG ACCGGCGTGG
101 ACCAGTGGAT GGATGGCGCC AAGCCGTACG CACTCACCGA TGGGCTCCCG
151 ATGATGGACG TGTCCACCAT GCTGGCATTG GAGGTGGGAT ACATGGCCAT
201 GCTGCTCTTC GGCATCCCGA TCATGAAGCA GATGGAGAAG CCTTTTGAGC
251 TCAAGACCAT CAAGCTCTTG CACAACCTGT TTCTCTTCGG ACTTTCCTTG
301 TACATGTGCG TGGAGACCAT CCGCCAGGCT ATCCTCGGAG GCTACAAAGT
351 GTTTGGAAAC GACATGGAGA AGGGCAACGA GTCTCATGCT CAGGGCATGT
401 CTCGCATCGT GTACGTGTTT TACGTGTCCA AGGCATACGA GTTCTTGAT
451 ACCGCCATCA TGATCCTTTG CAAGAAGTTC AACCAGGTTT CCTTCTTGCA
501 TGTGTACCAC CATGCCACCA TTTTGGCCAT CTGGTGGGCT ATCGCCAAGT
551 ACGCTCCAGG AGGTGATGCG TACTTTTCAG TGATCCTAAA CTCTTTCATC
601 CACGTTATCA TGTACTCTTA CTACCCTGAA

FIG.65

1 TGDFLNLLQ AQQGRMANSS VWDDVVGRVE TGVDQWMDGA KPYALTDGLP
51 MMDVSTMLAF EVGYMAMLLF GIPIMQMEK PFELKTIKLL HNLFLFGLSL
101 YMCVETIRQA ILGGYKVFGN DMEKGNESHA QGMSRIVYVF YVSKAYEFLD
151 TAIMILCKKF NQVSFLHVVH HATIFAIWWA IAKYAPGGDA YFSVILNSFI
201 HVIMYSYYPE

FIG.66

```
1  ATGGCAAACA GCAGCGTGTG GGATGATGTG GTGGGCCGCG TGGAGACCGG
51 CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 TCCCGATGAT GGACGTGTCC ACCATGCTGG CATTGAGAGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AAGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGACAAA CTTGTTTCTC TTCGGACTTT
251 CCTTGTACAT GTGCGTGGAG ACCATCCGCC AGGCTATCCT CGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG TGTCTGCGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTCCTTC
451 TTGCATGTGT ACCACCATGC CACCATTTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCT CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTTCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA
```

FIG.67

```
1  ATGGCAAACA GCAGCGTGTG GGATGATGTG GTGGGCCGCG TGGAGACCGG
51  CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 CCCCgatgat GGACGTGTCC ACCATGCTGG CATTGAGGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AAGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGCACAA CTTGTTTCTC TTCGGACTTT
251 CCTTGTACAT GTGCGTGGAG ACCATCCGCC AGGCTATCCT CGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG CGTTCACGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTTCCTTC
451 TTGCATGTGT ACCACCATGC CACCATTTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCC CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGCACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTTCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA
```

FIG.68

```
1  ATGGCAACAG CAGCGTGTGG GATGATGTGG TGGGCCGCGT GGAGACCAGC
51  GTGGACCAGT GGATGGATGG CGCCAAGCCG TACGCACTCA CCGATGGGCT
101 CCCGATGATG GACGTGTCCA CCATGCTGGC ATTCGAGGTG GGATACATGG
151 CCATGCTGCT CTTCGGCATC CCGATCATGA AGCAGATGGA GAAGCCTTTT
201 GAGCTCAAGA CCATCAAGCT CTTGCACAAC TTGTTTCTCT TCGGACTTTC
251 CTTGTACATG TCGTGGGAGA CCATCCGCCA GGCTATCCTC GGAGGCTACA
301 AAGTGTTTGG AAACGACATG GAGAAGGGCA ACGAGTCTCA TGCTCAGGGC
351 ATGTCTCGCA TCGTGTACGT GTTCTACGTG TCCAAGGCAT ACGAGTTCTT
401 GGATACCGCC ATCATGATCC TTTGCAAGAA GTTCAACCAG GTTTCCTTCT
451 TGCAAGTGTA CCACCATGCC ACCATTTTTG CCATCTGGTG GGCTATCGCC
501 AAGTACGCTC CAGGAGGTGA TCGTACTTT TCAGTGATCC TCAACTCTTT
551 CGTGCACACC GTCATGTACG CATACTACTT CTTCTCCTCC CAAGGGTTCG
601 GGTTCGTGAA GCCAATCAAG CCGTACATCA CCACCCTTCA GATGACCCAG
651 TTCATGGCAA TGCTTGTGCA GTCCTTGAC GACTACCTCT TCCCATGCGA
701 CTACCCACAG GCTCTTGTGC AGCTTCTTGG AGTGTACATG ATCACCTTGC
751 TTGCCCTCTT CGGCAACTTT TTTGTGCAGA GCTATCTTAA AAAGCCAAAA
801 AAGAGCAAGA CCAACTAA
```

FIG.69

```
1  ATGGCAAACA GCAGCGTGTG GGATGATGTG GTGGGCCGCG TGGAGACCGG
51 CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 TCCCGATGAT GGACGTGTCC ACCATGCTGG CATTGAGGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AAGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGCACAA CTTGTTTCTC TTCGGACTTT
251 CTTGTACAT GTGCGTGGAG ACCATCCGCC AGGCTATCCT CGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG TGTTCTACGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTTCCTTC
451 TTGCATGTGT ACCACCATGC CACCGTTTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCT CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGCACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTTCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA
```

FIG.70

1 ATGGCAAACA GCAGCGTGTG GGATGGTGTG GTGGGCCGCG TGGAGACCGG
51 CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 TCCCGATGAT GGACGTGTCC ACCATGCTGG CATTGAGGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AAGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGCACAA CTTGTTTCTC TTCGGACTTT
251 CCTTGACAT GTGCGTGGAG ACCATCCGCC AGGCTATCCT CGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG TGTTCTACGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTCCTTC
451 TTGCATGCGT ACCACCATGC CACCATTTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCT CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTTCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA

FIG.71

```
1  ATGGCAAACA GCAGCGTGTG GGATGATGTG GTGGGCCGCG TGGAGACCGG
51  CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 TCCCGATGAT GGACGTGTCC ACCATGCTGG CATTGAGGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AAGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGACAA CTTGTTTCTC TTCGGACTTT
251 CTTGTACAT GTGCGTGGAG ACCATCCGCC AGGCTATCCT CGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG TGTTCTACGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTTCCTTC
451 TTGCATGTGT ACCACCATGC CACCATTTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCT CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTCCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA
```

FIG.72

```
1  ATGGCAAACA GCAGCGTGTG GGATGATGTG GTGGGCCGCG TGGAGACCGG
51  CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 TCCCGATGAT GGACGTGTCC ACCATGCTGG CATTGAGGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AGGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGACAA CTTGTTTCTC TTCGGACTTT
251 CCTTGACAT GTGCGTGGTG ACCATCCGCC AGGCTATCCT TGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG TGTTCTACGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTCCTTC
451 TTGCATGTGT ACCACCATGC CACCATTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCT CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTTCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA
```

FIG.73

1 MANSSVWDDV VGRVETGVDQ WMDGAKPYAL TDGLPMDVS TMLAFEVGYM
51 AMLLFGIPIM KQMEKPFELK TIKLLHNLFL FGLSLYMCVE TIRQAILGGY
101 KVFGNDMEKG NESHAQGMSR IVYVFCVSKA YEFLDTAIMI LCKKFNQVSF
151 LHVYHHATIF AIWWAIAKYA PGGDYFYSVI LNSFVHTVMY AYYFFSSQGF
201 GFVKPIKPYYI TTLQMTQFMA MLVQSLYDYL FPCDYPQALV QLLGVYMITL
251 LALFGNFFVQ SYLKKPKKSK TN*

FIG.74

1 MANSSVWDDV VGRVETGVDQ WMDGAKPYAL TDGPPMMDVS TMLAFEVGYM
51 AMLLFGIPIM KQMEKPFELK TIKLLHNLFL FGLSLYMCVE TIRQAILGGY
101 KVFGNDMEKG NESHAQGMSR IVYAFYVSKA YEFLDTAIMI LCKKFNQVSF
151 LHVYHHATIF AIWWAIAKYA PGGDAYFSVI LNSFVHTVMY AYYFFSSQGF
201 GFVKPIKPYI TTLQMTQFMA MLVQSLYDYL FPCDYPQALV QLLGVYMITL
251 LALFGNFFVQ SYLKKPKKSK TN*

FIG.75

1 MATAACGMMW WAAWRPAWTS GWMAPSRTHS PMGSR*WTCP PCWHSRWDTW
51 PCCSSASRS* SRWRSLLSSR PSSSCTTCFS SDFPCTCAWR PSARLSSEAT
101 KCLETTWRRRA TSLMLRACLA SCTCSTCPRH TSSWIPPS*S FARSTRFPS
151 CKCTTMPPFL PSGGLSPSTL QEVMRTFQ*S STLCTPSCT HTTSSPPKGS
201 GS*SQSSRTS PPFR*PSSWQ CLCSPCTTTS SHATTHRLLC SFLECT*SPC
251 LPSSATFLCR AILKSQKRAR PT

FIG.76

1 MANSSVWDDV VGRVETGVDQ WMDGAKPYAL TDGLPMDVS TMLAFEVGYM
51 AMLLFGIPIM KQMEKPFELK TIKLLHNLFL FGSLYMCVE TIRQAILGGY
101 KVFGNDMEKG NESHAQGMSR IVYVFYVSKA YEFLDTAIMI LCKKFNQVSF
151 LHVYHHATVF AIWWAIAKYA PGGDAYFSVI LNSFVHTVMY AYYFFSSQGF
201 GFVKPIKPYI TTLQMTQFMA MLVQSLYDYL FPCDYPQALV QLLGVYMITL
251 LALFGNFFVQ SYLKKPKKSK TN*

FIG.77

1 MANSSVWDGV VGRVETGVDQ WMDGAKPYAL TDGLPMDVS TMLAFEVGYM
51 AMLLFGIPIM KQMEKPFELK TIKLLHNLFL FGLSLYMCVE TIRQAILGGY
101 KVFGNDMEKG NESHAQGMSR IVYVFYVSKA YEFLDTAIMI LCKKFNQVSF
151 LHAYHHATIF AIWWAIAKYA PGGDAYFSVI LNSFVHTVMY AYYFFSSQGF
201 GFVKPIKPYI TTLQMTQFMA MLVQSLYDYL FPCDYPQALV QLLGVYMITL
251 LALFGNFFVQ SYLKKPKKSK TN*

FIG.78

1 MANSSVWDDV VGRVETGVDQ WMDGAKPYAL TDGLPMDVS TMLAFEVGYM
51 AMLLFGIPIM KQMEKPFELK TIKLLHNLFL FGLSLYMCVE TIRQAILGGY
101 KVFGNDMEKG NESHAQGMSR IVYVFYVSKA YEFLDTAIMI LCKKFNQVSF
151 LHVYHHATIF AIWWAIAKYA PGGDAYFSVI LNSFVHTVMY AYYFFSSQGF
201 GFVKPIKPYI TTLQMTQFMA MLVQSLYDYL FPCDYPQALV QLLGVYMITL
251 LALFGNFFVQ SYLKKPKKSK TN*

FIG.79

1 MANSSVWDDV VGRVETGVDQ WMDGAKPYAL TDGLPMDVS TMLAFEVGYM
51 AMLLFGIPIM RQMEKPFELK TIKLLHNLFL FGLSLYMCVV TIRQAILGGY
101 KVFGNDMEKG NESHAQGMSR IVYVFYVSKA YEFLDTAIMI LCKKFNQVSF
151 LHVYHHATIF AIWWAIAKYA PGGDAYFSVI LNSFVHTVMY AYYFFSSQGF
201 GFVKPIKPYI TTLQMTQFMA MLVQSLYDYL FPCDYPQALV QLLGVYMITL
251 LALFGNFFVQ SYLKKPKKSK TN*

FIG.80

334(PLASMID)	pRAT-4-A1	pRAT-4-A2	pRAT-4-A3	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pYX242
ADDED SUBSTRATE	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA
	% TOTAL LIPID							
C16:1n-7	16.95	5.22	15.52	11.06	14.49	15.69	15.18	11.24
C18:0	4.95	6.73	5.94	4.17	4.57	4.25	4.79	5.64
C18:1n-9	20.26	44.21	22.54	17.22	16.89	16.51	18.92	28.72
C18:1n-7	2.87	1.64	2.97	7.51	1.85	3.90	3.50	1.70
C18:2n-6	1.03	13.79	1.44	0.67	1.32	0.28	0.36	5.53
C18:3n-6	3.05	1.09	5.28	2.38	1.89	2.45	2.79	2.24
C20:3n-6	10.09	0.43	0.21	11.04	5.76	10.77	9.28	
C22:0		0.57						
% CONVERSION*	76.8%	28.4%	3.8%	82.2%	75.3%	81.5%	76.9%	

*% CONVERSION=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.81A

334 (PLASMID)	pRAT-4-A1	pRAT-4-A2	pRAT-4-A3	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pYX242
ADDED SUBSTRATE	EPA	EPA	EPA	EPA	EPA	EPA	EPA	EPA
	% TOTAL LIPID							
C16:1n-7	18.35	17.65	7.96	11.80	14.06	11.53	9.68	20.66
C18:0	3.41	3.69	2.57	2.90	3.18	3.10	3.20	8.49
C18:1n-9	15.22	16.44	10.69	12.28	13.01	11.50	12.85	16.27
C18:1n-7	1.36	2.12	1.60	5.62	2.13	2.02	2.35	1.33
C18:2n-6	1.07	0.40			0.17		0.21	0.83
C20:5n-3	23.24	31.54	46.96	37.98	40.40	32.68	43.67	13.55
C22:1n-9		0.09	0.12	0.09				
C22:5n-3	0.40	0.16		3.13	1.37	4.61	1.27	
% CONVERSION*	1.7%	0.5%		7.6%	3.3%	12.4%	2.8%	

*% CONVERSION=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.81B

pRAT-4-A1	1	M	A	N	S	S	V	W	D	D	V	V	G	R	V	E	T	G	V	D	Q	W	M	D	G	A	K	P	Y	A	L	T	D	G	L	P	M	M	D	V	S	T	M	L	A	F	E	V	47
pRAT-4-A2	1	M	A	N	S	S	V	W	D	D	V	V	G	R	V	E	T	G	V	D	Q	W	M	D	G	A	K	P	Y	A	L	T	D	G	L	P	M	M	D	V	S	T	M	L	A	F	E	V	47
pRAT-4-A4	1	M	A	N	S	S	V	W	D	D	V	V	G	R	V	E	T	G	V	D	Q	W	M	D	G	A	K	P	Y	A	L	T	D	G	L	P	M	M	D	V	S	T	M	L	A	F	E	V	47
pRAT-4-A6	1	M	A	N	S	S	V	W	D	D	V	V	G	R	V	E	T	G	V	D	Q	W	M	D	G	A	K	P	Y	A	L	T	D	G	L	P	M	M	D	V	S	T	M	L	A	F	E	V	47
pRAT-4-A7	1	M	A	N	S	S	V	W	D	D	V	V	G	R	V	E	T	G	V	D	Q	W	M	D	G	A	K	P	Y	A	L	T	D	G	L	P	M	M	D	V	S	T	M	L	A	F	E	V	47
pRAT-4-D1	1	M	A	N	S	S	V	W	D	D	V	V	G	R	V	E	T	G	V	D	Q	W	M	D	G	A	K	P	Y	A	L	T	D	G	L	P	M	M	D	V	S	T	M	L	A	F	E	V	47
pRAT-4-A1	48	G	Y	M	A	M	L	L	F	G	I	P	I	M	K	Q	M	E	K	P	F	E	L	K	T	I	K	L	L	H	N	L	F	L	F	G	L	S	L	Y	M	C	V	E	T	I	R	Q	94
pRAT-4-A2	48	G	Y	M	A	M	L	L	F	G	I	P	I	M	K	Q	M	E	K	P	F	E	L	K	T	I	K	L	L	H	N	L	F	L	F	G	L	S	L	Y	M	C	V	E	T	I	R	Q	94
pRAT-4-A4	48	G	Y	M	A	M	L	L	F	G	I	P	I	M	K	Q	M	E	K	P	F	E	L	K	T	I	K	L	L	H	N	L	F	L	F	G	L	S	L	Y	M	C	V	E	T	I	R	Q	94
pRAT-4-A6	48	G	Y	M	A	M	L	L	F	G	I	P	I	M	K	Q	M	E	K	P	F	E	L	K	T	I	K	L	L	H	N	L	F	L	F	G	L	S	L	Y	M	C	V	E	T	I	R	Q	94
pRAT-4-A7	48	G	Y	M	A	M	L	L	F	G	I	P	I	M	K	Q	M	E	K	P	F	E	L	K	T	I	K	L	L	H	N	L	F	L	F	G	L	S	L	Y	M	C	V	E	T	I	R	Q	94
pRAT-4-D1	48	G	Y	M	A	M	L	L	F	G	I	P	I	M	R	Q	M	E	K	P	F	E	L	K	T	I	K	L	L	H	N	L	F	L	F	G	L	S	L	Y	M	C	V	Y	T	I	R	Q	94
pRAT-4-A1	95	A	I	L	G	G	Y	K	V	F	G	N	D	M	E	K	G	N	E	S	H	A	Q	G	M	S	R	I	V	Y	V	F	C	V	S	K	A	Y	E	F	L	D	T	A	I	M	I	L	141
pRAT-4-A2	95	A	I	L	G	G	Y	K	V	F	G	N	D	M	E	K	G	N	E	S	H	A	Q	G	M	S	R	I	V	Y	V	F	C	V	S	K	A	Y	E	F	L	D	T	A	I	M	I	L	141
pRAT-4-A4	95	A	I	L	G	G	Y	K	V	F	G	N	D	M	E	K	G	N	E	S	H	A	Q	G	M	S	R	I	V	Y	V	F	C	V	S	K	A	Y	E	F	L	D	T	A	I	M	I	L	141
pRAT-4-A6	95	A	I	L	G	G	Y	K	V	F	G	N	D	M	E	K	G	N	E	S	H	A	Q	G	M	S	R	I	V	Y	V	F	C	V	S	K	A	Y	E	F	L	D	T	A	I	M	I	L	141
pRAT-4-A7	95	A	I	L	G	G	Y	K	V	F	G	N	D	M	E	K	G	N	E	S	H	A	Q	G	M	S	R	I	V	Y	V	F	C	V	S	K	A	Y	E	F	L	D	T	A	I	M	I	L	141
pRAT-4-D1	95	A	I	L	G	G	Y	K	V	F	G	N	D	M	E	K	G	N	E	S	H	A	Q	G	M	S	R	I	V	Y	V	F	C	V	S	K	A	Y	E	F	L	D	T	A	I	M	I	L	141
pRAT-4-A1	142	C	K	F	N	Q	V	S	F	L	H	V	Y	H	H	A	T	I	F	A	I	W	A	I	A	K	Y	A	P	G	G	D	A	Y	F	S	V	I	L	N	S	F	V	H	T	V	188		
pRAT-4-A2	142	C	K	F	N	Q	V	S	F	L	H	V	Y	H	H	A	T	I	F	A	I	W	A	I	A	K	Y	A	P	G	G	D	A	Y	F	S	V	I	L	N	S	F	V	H	T	V	188		
pRAT-4-A4	142	C	K	F	N	Q	V	S	F	L	H	V	Y	H	H	A	T	I	F	A	I	W	A	I	A	K	Y	A	P	G	G	D	A	Y	F	S	V	I	L	N	S	F	V	H	T	V	188		
pRAT-4-A6	142	C	K	F	N	Q	V	S	F	L	H	V	Y	H	H	A	T	I	F	A	I	W	A	I	A	K	Y	A	P	G	G	D	A	Y	F	S	V	I	L	N	S	F	V	H	T	V	188		
pRAT-4-A7	142	C	K	F	N	Q	V	S	F	L	H	V	Y	H	H	A	T	I	F	A	I	W	A	I	A	K	Y	A	P	G	G	D	A	Y	F	S	V	I	L	N	S	F	V	H	T	V	188		
pRAT-4-D1	142	C	K	F	N	Q	V	S	F	L	H	V	Y	H	H	A	T	I	F	A	I	W	A	I	A	K	Y	A	P	G	G	D	A	Y	F	S	V	I	L	N	S	F	V	H	T	V	188		

FIG.82A

pRAT-4-A1	189	M	Y	A	Y	F	F	S	S	Q	G	F	G	F	V	K	P	I	K	P	Y	I	T	T	L	Q	M	T	Q	F	M	A	M	L	V	Q	S	L	Y	D	Y	L	F	P	C	D	Y	235
pRAT-4-A2	189	M	Y	A	Y	F	F	S	S	Q	G	F	G	F	V	K	P	I	K	P	Y	I	T	T	L	Q	M	T	Q	F	M	A	M	L	V	Q	S	L	Y	D	Y	L	F	P	C	D	Y	235
pRAT-4-A4	189	M	Y	A	Y	F	F	S	S	Q	G	F	G	F	V	K	P	I	K	P	Y	I	T	T	L	Q	M	T	Q	F	M	A	M	L	V	Q	S	L	Y	D	Y	L	F	P	C	D	Y	235
pRAT-4-A6	189	M	Y	A	Y	F	F	S	S	Q	G	F	G	F	V	K	P	I	K	P	Y	I	T	T	L	Q	M	T	Q	F	M	A	M	L	V	Q	S	L	Y	D	Y	L	F	P	C	D	Y	235
pRAT-4-A7	189	M	Y	A	Y	F	F	S	S	Q	G	F	G	F	V	K	P	I	K	P	Y	I	T	T	L	Q	M	T	Q	F	M	A	M	L	V	Q	S	L	Y	D	Y	L	F	P	C	D	Y	235
pRAT-4-D1	189	M	Y	A	Y	F	F	S	S	Q	G	F	G	F	V	K	P	I	K	P	Y	I	T	T	L	Q	M	T	Q	F	M	A	M	L	V	Q	S	L	Y	D	Y	L	F	P	C	D	Y	235
pRAT-4-A1	236	P	Q	A	L	V	Q	L	L	G	V	Y	M	I	T	L	L	A	L	F	G	N	F	F	V	Q	S	Y	L	K	K	P	K	K	S	K	T	N								272		
pRAT-4-A2	236	P	Q	A	L	V	Q	L	L	G	V	Y	M	I	T	L	L	A	L	F	G	N	F	F	V	Q	S	Y	L	K	K	P	K	K	S	K	T	N								272		
pRAT-4-A4	236	P	Q	A	L	V	Q	L	L	G	V	Y	M	I	T	L	L	A	L	F	G	N	F	F	V	Q	S	Y	L	K	K	P	K	K	S	K	T	N								272		
pRAT-4-A6	236	P	Q	A	L	V	Q	L	L	G	V	Y	M	I	T	L	L	A	L	F	G	N	F	F	V	Q	S	Y	L	K	K	P	K	K	S	K	T	N								272		
pRAT-4-A7	236	P	Q	A	L	V	Q	L	L	G	V	Y	M	I	T	L	L	A	L	F	G	N	F	F	V	Q	S	Y	L	K	K	P	K	K	S	K	T	N								272		
pRAT-4-D1	236	P	Q	A	L	V	Q	L	L	G	V	Y	M	I	T	L	L	A	L	F	G	N	F	F	V	Q	S	Y	L	K	K	P	K	K	S	K	T	N								272		

FIG.82B

SCORES Init1: 259 Initn: 259 Opt: 509 z-score: 86.5 E(): 0.28
>>EN:HSEL01 (300 aa)
initn: 259 init1: 259 opt: 509 Z-score: 86.5 expect(): 0.28
Smith-Waterman score: 511; 34.0% identity in 265 aa overlap
(4-265:9-257)

	10	20	30	40	50	
TEL01	MANSSVWDDVVGVRVETGVDQWMDGAKPYALTDGLPMDVSTMLAFEVGYMAMLLF					
	: : :: : : : : : :: : :: :					
HSEL01	MEHFDASLSTYFKALLGPRDTRVKGW-----FLLDNYIPTFICSVI-----YLLIVWL					
	10	20	30	40		
	60	70	80	90	100	110
TEL01	GIPIMQMEKPFELKTIKLLHNLFLFGLSLYMCVETIRQAILGGYKVFVGNDEKGNESHA					
	: :: :: : :: : : : : ::					
HSEL01	GPKYMRN-KQPFSCRGILVVYNLGLTLLSLYMFCELVGVWEGKYNFFCQGTRTAGESDM					
	50	60	70	80	90	100
	120	130	140	150	160	170
TEL01	QGMSRIVYVVFYVSKAYEFLDTAIMILCKKFNQVSFLHVVYHHATIFAIWWAIKYAPGGDA					
	: : :: : :: :: : :::: :::: : :: :					
HSEL01	K-IIRVLWYYFYSKLIEFMDTFFFILRKNNHQITVLHVVYHASMNIWWFVMNWVPCGHS					
	110	120	130	140	150	160
	180	190	200	210	220	230
TEL01	YFSVILNSFVHTVMYAYFFSSQGFGFVKPI--KPYITTLMQTMAMLVQSLYDYLFP					
	:: : :: : : :: : : : : ::					
HSEL01	YFGATLNSFIHVLMSYYGLSS--VPSMRPYLWKKYITQGQLLQFVLTIIQTSCGVIWP					
	170	180	190	200	210	220
	240	250	260	270	280	290
TEL01	CDYPQALVQLLGVYMITLLALFGNFFVQSYLKKPKKSKTNXNCLHDMPLAGVRIDSESEL					
	: : : : : : :: :					
HSEL01	CTFPLGWLYFQIGYMISLIALFTNFIYQTYNKKGASRRKDHLKDHQNGSMAAVNGHTNSF					
	230	240	250	260	270	280

FIG.83

SCORES Init1: 278 Initn: 278 Opt: 515 z-score: 86.1 E(): 0.3
 >>EN:MEL04 (293 aa)
 initn: 278 init1: 278 opt: 515 Z-score: 86.1 expect(): 0.3
 Smith-Waterman score: 517; 34.1% identity in 267 aa overlap
 (8-270:16-265)

		10	20	30	40	50
TEL01		MANSSVWDDVVGRVETGVDQWMDGAKPYALTDGLPMDVSTMLAFEVGYMAM				
		::	::		: :	::: :
MEL04		MEQLKAFDNEVNAFLDNMFGPRDSRVRGW-----FLLDSYLP-----FILTITYLLS				
		10	20	30	40	
	60	70	80	90	100	110
TEL01	LLFGIPIMKQMEKP-FELKTIKLLHNLFLGSLYMCVETIRQAILGGYKVFGNDMEKGN					
	:	:	: :	:	:	::
MEL04	IWLGNKYMKN--RPALSLRGILTLYNLAITLLSAYMLVELILSSWEGGYNLQCNLDSAG					
	50	60	70	80	90	100
	120	130	140	150	160	170
TEL01	ESHAQMSRIVYVYVSKAYEFLDTAIMILCKKFNQVSFLHVVHHATIFAIWWAIKYP					
	:	::	:::::	:		
MEL04	EGDVR-VAKVLWWYYFSKLVFLDTIFFVLRKKTNQITFLHVVHHASMFNIWWCVLWIP					
	110	120	130	140	150	160
	180	190	200	210	220	
TEL01	GGDAYFSVILNSFVHTVMYAYYFFSSQGFGFVKPI--KPYITTLQMTQFMAMLVQSLYD					
	:::	:	:	: :	:	:
MEL04	CGQSFFGPTLNSFIHILMYSYYGLSV--FPSMHKYLWKKYLTQAQLVQFVLTITHTLA					
	170	180	190	200	210	220
	230	240	250	260	270	280
TEL01	YLFPCDYPQALVQLLGVYMITLLALFGNFFVQSYLKKPKKSKTNXNCLHDMPLAGVRIDS					
	:	:	::	::	::	::
MEL04	VVKPCGFPFGCLIFQSSYMMTLVILFLNFYIQTYRKKPVKKELQEKEVKNGFPAHLIVA					
	230	240	250	260	270	280

FIG.84

SCORES Initl: 412 Initn: 541 Opt: 628 z-score: 102.7 E(): 0.035
>>EN:GLELO (319 aa)
initn: 541 initl: 412 opt: 628 Z-score: 102.7 expect(): 0.035
Smith-Waterman score: 628; 43.4% identity in 244 aa overlap
(34-272:69-308)

	10	20	30	40	50	60
TEL01	SSVWDDVVG RVETGVDQWMDGAKPYALTDGLPMDVSTMLAFEVGYMAMLLFGIPIMKQM					
				: :	: : :: :: :: :	::
GLELO	LVAQAEKYIPTIVHHTRGFLVAVESPLARELPLMNP FHVLLIVLAYLVTVFVGMQIMKNF					
	40	50	60	70	80	90
	70	80	90	100	110	120
TEL01	EKPFEKTIKLLHNLFLFGLSLYMCVETIRQAILGGYKVF GNDMEKGNESHAQGM SRIVY					
	: : : : :	::	: : :: :	: : ::	::: ::	::: ::
GLELO	ER-FEVKTFSLLHNFLVSLISAYMCGGILYEAYQANYGLFENAAD--HTFKGLPMAKMIW					
	100	110	120	130	140	150
	130	140	150	160	170	180
TEL01	VFYVSKAYEFLDTAIMILCKKFNQVSFLHVYHHATIFAIWWAIAKYAPGGDAYFSVILNS					
	: : : :	: :: :: :	:: : : :			
GLELO	LFYFSKIMEFVDTMIMVLKKNRQISFLHVYHHSSIFTIWWLTVFVAPNGEAYFSAALNS					
	160	170	180	190	200	210
	190	200	210	220	230	
TEL01	FVHTVMYAYYFFSSQGFGFVKPIKPYITT LQMTQFMAMLVQSLYD-YLFPC----DYPQA					
	: :: : : : : :	:		:	:	
GLELO	FIHVIMYGYFLSALGFKQVSFIKFYITRSQMTQFCMMSVQSSWDHYAMKVLGRPGYPFF					
	220	230	240	250	260	270
	240	250	260	270	280	290
TEL01	LVQLLGVYMITLLALFGNFFVQSYLKKPKKSKTNXNCLHDMPLAGVRIDSESELRRHANS					
	:: : : : ::	:: ::				
GLELO	ITALLWFYMW TMLGLFYNFYRKNA-KLAKQAKADAAKEKARKLQ					
	280	290	300	310		

FIG.85

SCORES Init1: 94 Initn: 220 Opt: 302 z-score: 61.6 E(): 6.5
>>EN:CEELO (288 aa)
initn: 220 init1: 94 opt: 302 Z-score: 61.6 expect(): 6.5
Smith-Waterman score: 312; 34.3% identity in 239 aa overlap
(49-270:54-278)

	20	30	40	50	60	70	
TEL01	DQWMDGAKPYALTDGLPMDVSTMLAFEVGYMAMLLFGIPIMKQMEKPFELKTIKL-LHN						
					::	: : ::	:
CEELO	ATHGPKNFPDAEGRKFFADHFDVTIQASILYM-VVVFGTKWFMNRQPFQL-TIPLNIWN						
	30	40	50	60	70	80	
	80	90	100	110	120		
TEL01	LFLFGLSLYMCVE-----TI-RQAILGGY-KVFGNDMEKGNESHAQMSRIVVYFYVS						
	:: :: :	:	:: ::	: :::		:: ::	
CEELO	FILAAFSIAGAVKMTPEFFGTIANKGIVASYCKVF--DFTKGENGYW-----VWLFMAS						
	90	100	110	120	130		
	130	140	150	160	170	180	
TEL01	KAYEFLDTAIMILCKKFNQVSFLHVVYHHATIFAIWWAIKAYAPGGDAYFSVILNSFVHTV						
	: ::	:: :	:	:	: :	:: :	
CEELO	KLFELVDITIFLVLRRK--PLMFLHWYHHILTMIYAWYSHPLTPGFNRY-GIYLNFFVHAF						
	140	150	160	170	180	190	
	190	200	210	220	230	240	
TEL01	MYAYYFFSSQGFGFVKPIKPYITTLMQTMFM---AMLVQSLYDYLPF---CDYPQALVQL						
	: :	: :	: :	: ::	:	: :: :	
CEELO	MYSYYFLRSMKIRVPGFIAQAITSQIVQFIISCAVLAHLGYLMHFTNANCDFEPSVFKL						
	200	210	220	230	240	250	
	250	260	270	280	290	300	
TEL01	LGVYM-ITLLALFGNFFVQSYLKKPKKSKTNXNCLHDMPLAGVRIDSESELRRHANSIFF						
	: :		: :	:	:		
CEELO	-AVFMDTTYLALFVNFFLQSYVLRGGKDKYKAVPKKKNN						
	260	270	280				

FIG.86

334(PLASMID)	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pXY242	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pXY242	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pXY242
ADDED SUBSTRATE	GLA	GLA	GLA	GLA	GLA	AA	AA	AA	AA	AA	STA	STA	STA	STA	STA
	% TOTAL LIPID														
C16:1n-7	36.29	39.18	38.58	38.84	45.68	30.36	35.19	33.47	33.70	45.06	34.43	39.63	38.03	38.74	45.94
C18:0	2.78	3.13	2.73	3.05	1.76	2.51	2.78	2.56	2.72	1.65	4.01	3.13	3.06	3.76	1.82
C18:1n-9	15.89	16.55	15.92	16.34	13.23	14.60	15.14	15.39	15.26	13.01	16.05	16.51	17.37	19.07	14.10
C18:1w7	5.91	2.01	3.44	2.06	0.93	5.72	1.94	3.75	2.22	1.03	4.92	1.78	3.78	2.52	0.95
C18:3w6	2.10	1.96	1.60	2.19	5.29										
C18:4w3											0.83	0.88	0.53	1.09	3.42
C20:3w6	4.56	4.23	5.42	3.88					0.04						
C20:4w6						16.53	13.61	14.21	17.13	7.54					
C20:4w3											2.27	3.25	3.95	4.01	0.06
C22:4w6						0.57	0.20	0.75	0.18						
C22:4n-3											1.10	0.19	0.63	0.16	
% CONVERSION* OF C18	68.4%	68.4%	77.2%	64.0%							80.3%	79.6%	89.7%	79.3%	
% CONVERSION* OF C20						3.4%	1.5%	5.0%	1.0%		32.6%	5.5%	13.8%	3.8%	

FIG.87A

334(PLASMID)	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pYX242	pRAT-4-A4	pRAT-4-A6	pRAT-4-A7	pRAT-4-D1	pYX242
ADDED SUBSTRATE	EPA	EPA	EPA	EPA	EPA	NONE	NONE	NONE	NONE	NONE
	% TOTAL LIPID									
C16:1w7	31.51	33.92	34.31	34.74	47.12	39.54	41.51	41.44	43.02	46.45
C18:0	2.76	3.09	2.78	2.77	1.79	2.45	3.21	2.59	3.07	2.05
C18:1w9	15.69	15.79	16.49	15.22	13.95	18.56	19.08	17.98	18.80	14.03
C18:1w7	6.07	1.97	3.72	1.89	1.05	8.77	2.57	4.86	2.77	1.07
C20:1w7	0.22	0.08	0.13	0.07		0.61	0.07	0.14	0.07	
C20:5w3	11.88	15.20	11.32	13.91	6.41					
C22:5n-3	1.49	0.54	1.74	0.54						
% CONVERSION*	11.1%	3.4%	13.3%	3.7%						

*** CONVERSION=PRODUCT/(SUBSTRATE+PRODUCT)

FIG.87B

334(PLASMIID)	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7	pYX242	pRAT-4-A7
ADDED SUBSTRATE	LA	LA	ALA	ALA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA	GLA
% TOTAL LIPID																								
C16:1n-7	38.73	27.71	39.05	26.97	35.28	30.13	34.75	30.95	35.96	28.51	34.91	31.45	34.71	24.76	40.02	24.75								
C16:1n-5	0.37	0.17	0.32	0.20	0.25	0.21	0.27	0.18	0.33	0.22	0.31	0.21	0.32	0.20	0.40	0.21								
C18:0	2.21	2.80	2.14	3.27	2.12	3.14	2.44	3.28	2.15	2.86	2.55	3.52	2.42	2.71	2.75	3.26								
C18:1n-9	11.14	12.79	11.73	15.73	10.14	14.24	11.13	14.79	11.17	14.59	11.33	16.32	11.63	13.85	12.86	15.99								
C18:1n-7	0.85	3.19	0.73	3.84	0.64	3.31	0.78	3.01	0.96	3.71	0.95	3.89	1.06	3.49	1.11	3.87								
C18:2n-6	3.04	3.54																						
C18:3n-6					4.49	1.59																		
C18:3n-3			3.55	3.56																				
C18:4n-3							3.09	0.85																
C20:1n-7		0.10		0.09		0.24		0.24		0.09		0.10		0.22		0.27								
C20:2n-6		1.07								0.08														
C20:3n-6						5.84			9.08	14.13				0.06										
C20:4n-6													14.60	20.89										
C20:3n-3				2.71						0.10														
C20:4n-3								4.19			5.06	8.49												
C20:5n-3											0.16	0.25			6.47	15.32								
C22:4n-7				0.11	0.19	0.14	0.18	0.13	0.14	0.13	0.24	0.14	0.22		0.27									
C22:4n-6														0.76										
C22:4n-3								0.74				1.04												
C22:5n-3																1.99								
% CONVERSION* OF C18																								
% CONVERSION OF C20																								
% CONVERSION=PRODUCT/(SUBSTRATE+PRODUCT)																								
23.2%																								
43.3%																								
78.6%																								
85.3%																								
15.0%																								
10.9%																								
3.5%																								
11.5%																								

FIG.88

pRPL-6-1 DNA sequence

```
1  ATGATGTTGG CCGCAGGCTA TCTTCTAGTG CTCTCGGCCG CTCGCCAGAG
51  CTTCCAGCAG GACATTGACA ACCCCAACGG GGCCTACTCG ACCTCGTGGA
101 CTGGCCTGCC CATTGTGATG TCTGTGGTCT ATCTCAGCGG TGTGTTTGGG
151 CTCACAAAGT ACTTCGAGAA CCGGAAGCCC ATGACGGGGC TGAAGGACTA
201 CATGTTCACT TACAATCTCT ACCAGGTGAT CATCAACGTG TGGTGCGTGG
251 TGGCCTTTCT CCTGGAGGTG CGGCGTGCGG GCATGTCACT CATCGGCAAT
301 AAGGTGGACC TTGGGCCCAA CTCCTTCAGG CTCGGCTTCG TCACGTGGGT
351 GCACTACAAC AACAAGTACG TGGAGCTCCT CGACACCCTA TGGATGGTGC
401 TGCGCAAGAA GACGCAGCAG GTCTCCTTCC TCCACGTCTA TCATCACGTG
451 CTTCTGATGT GGGCCTGGTT CGTTGTCGTC AAGCTCGGCA ATGGTGGTGA
501 CGCATATTTT GCCGTCTCA TGAAGTCGAT CATCCACGTG ATGATGTATT
551 CCTACTACAC CATGGCGCTC CTGGGCTGGT CATGCCCTG GAAGCGCTAC
601 CTCACGCAGG CACAGCTCGT GCAGTTTTCG ATCTGCCTCG CCCACTCCAC
651 ATGGGCGGCA GTAACGGGTG CCTACCCGTG GCGAATTTGC TTGGTGGAGG
701 TGTGGGTGAT GGTGTCCATG CTGGTGCTCT TCACACGCTT CTACCGCCAG
751 GCCTATGCCA AGGAGGCGAA GGCCAAGGAG GCGAAAAAGC TCGCACAGGA
801 GGCATCACAG GCCAAGGCGG TCAAGGCGGA GTAA
```

FIG.89

pRPL-6-1 translated amino acid sequence

1 MMLAAGYLLV LSAARQSFQQ DIDNPNGAYS TSWTGLPIVM SVVYLSGVFG
51 LTKYFENRKP MTGLKDYMFT YNLYQVIINV WCVVAFLLLEV RRAGMSLIGN
101 KVDLGPNSFR LGFVTWVHYN NKYVELLDL WMVLRKKTQQ VSFLHVVHHV
151 LLMWAWFVVV KLGNGGDAYF AGLMNSIIHV MMYSYYTMAL LGWSCPWKRY
201 LTQAQLVQFC ICLAHSTWAA VTGAYPWRI LVEVWVMVSM LVLFTRFYRQ
251 AYAKEAKAKE AKKLAQEASQ AKAVKAE

FIG.90

pRPL-6-B2 DNA sequence

```

1  ATGATGTTGG CCGCAGGCTA TCTTCTAGTG CTCTCGGCCG CTCGCCAGAG
51  CTTCCAGCAG GACATTGACA ACCCCAACGG GGCCTACTCG ACCTCGTGGA
101 CTGGCCTGCC CATTGTGATG TCTGTGGTCT ATCTCAGCGG TGTGTTTGGG
151 CTCACAAAGT ACTTCGAGAA CCGGAAGCCC ATGACGGGGC TGAAGGACTA
201 CATGTTCACT TACAATCTCT ACCAGGTGAT CATCAACGTG TGGTGCGTGG
251 TGGCCTTTCT CCTGGAGGTG CGGCGTGCGG GCATGTCACT CATCGGCAAT
301 AAGGTGGACC TTGGGCCCAA CTCCTTCAGG CTCGGCTTCG TCACGTGGGT
351 GCACTACAAC AACAAGTACG TGGAGCTCCT CGACACCCTA TGGATGGTGC
401 TGCGCAAGAA GACGCAGCAG GTCTCCTTCC TCCACGTCTA TCATCACGTG
451 CTTCTGATGT GGGCCTGGTT CGTTGTGCGT AAGCTCGGCA ATGGTGGTGA
501 CGCATATTTT GGCGGTCTCA TGAACTCGAT CATCCACGTG ATGATGTATT
551 CCTACTACAC CATGGCGCTC CTGGGCTGGT CATGCCCTG GAAGCGCTAC
601 CTCACGCAGG CACAGCTCGT GCAGTTTTGC ATCTGCCTCG CCCACTCCAC
651 ATGGGCGGCA GTAACGGGTG CCTACCCGTG GCGAATTTGC TTGGTGGAGG
701 TGTGGGTGAT GGTGTCCATG CTGGTGCTCT TCACACGCTT CTACCGCCAG
751 GCCTATGCCA AGGAGGCGAA GGCCAAGGAG GCGAAAAAGC TCGCACAGGA
801 GGCATCACAG GCCAAGGCGG TCAAGGCGGA GTAA

```

FIG.91

pRPL-6-A3 DNA sequence

```
1  ATGATGTTGG CCGCAGGCTA TCTTCTAGTG CTCTCGGCCG CTCGCCAGAG
51  CTTCCAGCAG GACATTGACA ACCCCAACGG GGCCTACTCG ACCTCGTGGA
101 CTGGCCTGCC CATTGTGATG TCTGTGGTCT ATCTCAGCGG TGTGTTTGGG
151 CTCACAAAGT ACTTCGAGGA CCGGAAGCCC ATGACGGGGC TGAAGGACTA
201 CATGTTCACT TACAATCTCT ACCAGGTGAT CATCAACGTG TGGTGCGTGG
251 TGGCCTTTCT CCTGGAGGTG CGGCGTGCGG GCATGTCACT CATCGGCAAT
301 AAGGTGGACC TTGGGCCCAA CTCCTTCAGG CTCGGCTTCG TCACGTGGGT
351 GCACTACAAC AACAAGTACG TGGAGCTCCT CGACACCCTA TGGATGGTGC
401 TGCACAAGAA GACGCAGCAG GTCTCCTTCC TCCACGTCTA TCATCACGTG
451 CTTCTGATGT GGGCCTGGTT CGTTGTCGTC AAGCTCGGCA ATGGTGGTGA
501 CGCATATTTT GGCGGTCTCA TGAAGTCGAT CATCCACGTG ATGATGTATT
551 CCTACTACAC CATGGCGCTC CTGGGCTGGT CATGCCCTG GAAGCGCTAC
601 CTCACGCAGG CACAGCTCGT GCAGTTTTGC ATCTGCCTCG CCCACTCCAC
651 ATGGGCGGCA GTAACGGGTG CCTACCCGTG GCGAATTTGC TTGGTGGAGG
701 TGTGGGTGAT GGTGTCCATG CTGGTGCTCT TCACACGCTT CTAAGTCCAG
751 GCCTATGCCA AGGAGGCGAA GGCCAAGGAG GCGAAAAAGC TCGCACAGGA
801 GGCATCACAG GCCAAGGCGG TCAAGGCGGA GTAA
```

FIG.92

pRPL-6-B2 translated amino acid sequence

1 MMLAAGYLLV LSAARQSFQQ DIDNPNGAYS TSWTGLPIVM SVVYLSGVFG
51 LTKYFENRKP MTGLKDYMFT YNLYQVIINV WCVVAFLLLEV RRAGMSLIGN
101 KVDLGPNSFR LGFVTWVHYN NKYVELLDTL WMVLRKKTQQ VSFLHVVYHHV
151 LLMWAWFVVV KLGNGGDAYF GGLMNSIIHV MMYSYYTMAL LGWSCPWKRY
201 LTQAQLVQFC ICLAHSTWAA VTGAYPWRIC LVEVWVMVSM LVLFTRFYRQ
251 AYAKEAKAKE AKKLAQEASQ AKAVKAE

FIG.93

pRPL-6-A3 translated amino acid sequence

1 MMLAAGYLLV LSAARQSFQQ DIDNPNGAYS TSWTGLPIVM SVVYLSGVFG
51 LTKYFEDRKP MTGLKDYMFT YNLYQVIINV WCVVAFLELV RRAGMSLIGN
101 KVDLGPNFSR LGFVTWVHYN NKYVELLDTL WMVLRKKTQQ VSFLHVYHHV
151 LLMWAWFVVV KLGNGGDAYF GGLMNSIIHV MMYSYYTMAL LGWSCPWKRY
201 LTQAQLVQFC ICLAHSTWAA VTGAYPWRIE LVEVWVMVSM LVLFTRFYCQ
251 AYAKEAKAKE AKKLAQEASQ AKAVKAE

FIG.94

SCORES Init1: 282 Initn: 410 Opt: 512 z-score: 93.2 E(): 0.18
 >>DEL:MELO4 (293 aa)
 initn: 410 init1: 282 opt: 512 Z-score: 93.2 expect(): 0.18
 Smith-Waterman score: 512; 34.1% identity in 261 aa overlap
 (17-263:13-268)

	10	20	30	40	50
pRPL-6-B2.pe	MMLAAGYLLVLSAARQSFOQDIDNPNGAYSTSW----TGLP-IVMSVVYLSGVFGLTKYF				
MELO4	MEQLKAFDNEVNAFLDNMFGPRDSRVRGWFLLDSYLPFTILTITYLLSIWLGKMYM				
	10	20	30	40	50
	60	70	80	90	100
pRPL-6-B2.pe	ENRKPM TGLKDYMF TYNLYQVI INVCVVAFLLEVRRAGMSLIGNKVD-LGPNSFRLGFV				
MELO4	KNR-PALSLRGILTLYNLAITLLSAYMLVELILSSWEGGYNLQCQNLDSAGEGDVRVAKV				
	60	70	80	90	100
	120	130	140	150	160
pRPL-6-B2.pe	TWVHYNNKYVELLDTLWMVLRKKTQQVSFLHVYHHVLLMWAVFVVVKLGNGGDAYFGGLM				
MELO4	LWVYYFSKLVFELDTIFFVLRKKTNQITFLHVYHHASMFNIWCVLWIPCGQSFFGPTL				
	120	130	140	150	160
	180	190	200	210	220
pRPL-6-B2.pe	NSIIHVMYSYYTMALLG-----WSCPWKRYLTQAQLVQFCICLAHSTWAAVTG-AYPW				
MELO4	NSFIHILMYSYYGLSVFSPMHKYLW---WKKYLTQAQLVQFVLTITHLSAVVKPCGFPP				
	180	190	200	210	220
	230	240	250	260	270
pRPL-6-B2.pe	RICLV-EVWMVSMLVLFTRFYRQAYAKEAKEAKEAKKLAQEASQAKAVKAE				
MELO4	G-CLIFQSSYMMTLVILFLNFYIQTyrKKPVKKELQEKEVKNGFPKAHLIVANGMTDKKA				
	240	250	260	270	280
MELO4	Q				

FIG.95

SCORES Init1: 233 Initn: 319 Opt: 424 z-score: 84.7 E(): 0.52
 >>DEL:GLELO (319 aa)
 initn: 319 init1: 233 opt: 424 Z-score: 84.7 expect(): 0.52
 Smith-Waterman score: 433; 33.8% identity in 240 aa overlap
 (38-267:78-316)

	10	20	30	40	50	60
pRPL-6-B2.pe	LLVLSAARQSFQQDIDNPNGAYSTSWTGLPIVMSVVYLSGVFGLTKYFENRKPMTGLKDY					
GLELO	PTIVHHTRGFLVAVESPLARELPLMNPFHVLLIVLAYLVTVFVGMQIMKNFERFE-VKTF					
	50	60	70	80	90	100
	70	80	90	100	110	120
pRPL-6-B2.pe	MFTYNLYQVIINVWCVVAFLLEVRRAGMSLIGNKVDLGPNSFRLGFVTWVHYNNKYVELL					
GLELO	SLLHNFCLVSI SAYMCGGILYEAYQANYGLFENAADHTFKGLPMAKMIWLFYFSKIMEFV					
	110	120	130	140	150	160
	130	140	150	160	170	180
pRPL-6-B2.pe	DTLWMVLRRKKTQQVSFLHVYHHVLLMWAVFVVVKLGNGGDAYFGGLMNSIIHVMMYSYYT					
GLELO	DTMIMVLKKNRQISFLHVYHHSSIFTIWWLVTFVAPNGEAYFSAALNSFIHVIMYGYFF					
	170	180	190	200	210	220
	190	200	210	220	230	
pRPL-6-B2.pe	MALLGWS--CPWKRYLTQAQLVQFCICLAHSTW---AAVTG--AYPWICLVEVWVMVS					
GLELO	LSALGFKQVSFIKFYITRSQMTQFCMMSVQSSWDMYAMKVLGRPGYPFFITALLWFYMW					
	230	240	250	260	270	280
	240	250	260	270		
pRPL-6-B2.pe	MLVLFTRFYRQ--AYAKEAKAKEAKKLAQEASQAKAVKAE					
GLELO	MLGLFYNFYRKNAKLAKQAKADAAKEKARKLQ					
	290	300	310			

FIG.96

SCORES Init1: 211 Initn: 271 Opt: 436 z-score: 84.5 E(): 0.54
 >>DEL:HSEL01 (300 aa)
 initn: 271 initl1: 211 opt: 436 Z-score: 84.5 expect(): 0.54
 Smith-Waterman score: 436; 28.5% identity in 274 aa overlap
 (11-271:4-275)

pRPL-6-B2.pe MMLAAGYLLVLSAARQSFQQIDNPNGAYSTSWTGL- - - P-IVMSVVYLSGVFGLTKYF
 HSEL01 MEHFDASLSTYFKALLGPRDTRVKGWFLLDNYIPTFICSVIYLLIVWLGPKYM
 pRPL-6-B2.pe ENRKPM TGLKDYMF TYNLYQVI INVCVVAFLLEVRRAGMSLI- -GNKVDLGPNSFRLGF
 HSEL01 RNKQPF S-CRGILVVYNLGLTLLSLYMFCELV TGVWEGKY NFFCQGR TA-GE S DMKIIR
 pRPL-6-B2.pe VTWVHYNNKYVELLDTLWMVLRKKTQQVSFLHVVHVLMMWAWFVVVKLGNGGDAYFGGL
 HSEL01 VLWVYYFSKLIEFMDTFFFILRKNNHQITVLHVYH HASMLNIWVFMNVWPCGHSYFGAT
 pRPL-6-B2.pe MNSIIHVMMSYTYT MALLGWSCP- -WKRYLTQAQLVQFCICLAHSTWAAV-TGAYPWRI
 HSEL01 LNSFIHVLMSYTYGLSSVPSMRPYLWKKYITQGQLLQFVLTIIQTSCGVIWPCTFPLGW
 pRPL-6-B2.pe CLVEVWVMVSMVLVLFTRFYRQAYAKEAKAKEAKKLA- -QEASQAKAVKAE
 HSEL01 LYFQIGYMISLIALFTNFYIQTYNKKGASRRKDHLDKHQNGSMAAVNGHTNSFSPLENNV
 HSEL01 KPRKLRKD

FIG.97

SCORES Init1: 233 Initn: 284 Opt: 404 z-score: 77.9 E(): 1.2
 >>DEL:TEL01 (273 aa)
 initn: 284 init1: 233 opt: 404 Z-score: 77.9 expect(): 1.2
 Smith-Waterman score: 405; 32.5% identity in 246 aa overlap
 (26-257:23-267)

	10	20	30	40	50
pRPL-6-B2.pe	MMLAAGYLLVL	SAARQSFQQD	IDNPNGAYST	SWT-GLPIV-----	MSVVYLSGV-FGL
			: : ::	: :: : :	
TEL01	MANSSVWDDV	VGRVETGVDQ	WMDGAKPYAL	TDGLPMDVST	MLAFEVGYMAMLLFGI
	10	20	30	40	50
	60	70	80	90	100
pRPL-6-B2.pe	TKYFENRKPMT	GLKDYMTYNLY	QVIINVWCVVA	FLLEVRRAGMS	LIGNKVDLGPNSFRL
	: : : :	: : : :::	: : : ::::	: ::: : :	: :
TEL01	PIMKQMEKPF	E-LKTIKLLH	NLFLFGLSLY	MCVETIRQAIL	GGYKVFGNDMEKGNESHAQ
	60	70	80	90	100
	120	130	140	150	160
pRPL-6-B2.pe	GF--VTWVHY	NNKYVELLD	TLMVLRKKTQ	QVSFLHVYHH	VLLMWAFVVKLGNGGDAY
	: ::: :	: :	:	: : : :	: : : :
TEL01	GMSRIVYVFY	VSKAYEFLD	TAIMILCKF	NQVSFLHVYH	HATIFAIWWAI
	120	130	140	150	160
	170	180	190	200	210
pRPL-6-B2.pe	FGGLMNSIIH	VMMYSYYT	MAL--LGWSCP	WKRYLTQAQL	VQFCICLAHST
	: : : : :	: : : :	: : : :	: : : :	: : : :
TEL01	FSVILNSFV	HTVMYAYY	FFSSQGFGF	VKPIKPYITT	LQMTQFMAMLV
	180	190	200	210	220
	230	240	250	260	270
pRPL-6-B2.pe	WRICLVEVW	VMVSMVL	LFTRFYRQ	AYAKEAKAKE	AKKLAQEASQAKAVKAE
	: :	: : :	: : :		
TEL01	QALVQLLGV	YMITLLAL	FGNFFVQ	SYLKKPKK	SKTN
	240	250	260	270	

FIG.98

PERCENT CONVERSION OF SUBSTRATE EPA TO PRODUCT DPA

334(pYX242)	334(pRPL-6-1)	334(pRPL-6-B2)	334(pRPL-6-A3)
0.0%*	1.2%	5.1%	2.6%

*% CONVERSION=
$$\frac{[\% \text{ PRODUCT}]}{[\% \text{ PRODUCT} + \% \text{ SUBSTRATE}]} \times 100$$

FIG. 99

prPL-6-B2	1	M	M	L	A	A	G	Y	L	L	V	L	S	A	A	R	Q	S	F	Q	Q	D	I	D	N	P	N	G	A	Y	S	T	S	W	T	G	L	P	I	V	M	40	
prPL-6-A3	1	M	M	L	A	A	G	Y	L	L	V	L	S	A	A	R	Q	S	F	Q	Q	D	I	D	N	P	N	G	A	Y	S	T	S	W	T	G	L	P	I	V	M	40	
prPL-6-1	1	M	M	L	A	A	G	Y	L	L	V	L	S	A	A	R	Q	S	F	Q	Q	D	I	D	N	P	N	G	A	Y	S	T	S	W	T	G	L	P	I	V	M	40	
prPL-6-B2	41	S	V	V	Y	L	S	G	V	F	G	L	T	K	Y	F	F	E	N	R	K	P	M	T	G	L	K	D	Y	M	F	T	Y	N	L	Y	Q	V	I	I	N	V	80
prPL-6-A3	41	S	V	V	Y	L	S	G	V	F	G	L	T	K	Y	F	F	E	N	R	K	P	M	T	G	L	K	D	Y	M	F	T	Y	N	L	Y	Q	V	I	I	N	V	80
prPL-6-1	41	S	V	V	Y	L	S	G	V	F	G	L	T	K	Y	F	F	E	N	R	K	P	M	T	G	L	K	D	Y	M	F	T	Y	N	L	Y	Q	V	I	I	N	V	80
prPL-6-B2	81	W	C	V	V	A	F	L	L	E	V	R	R	A	G	M	S	L	I	G	N	K	V	D	L	G	P	N	S	F	R	L	G	F	V	T	W	V	H	Y	N	120	
prPL-6-A3	81	W	C	V	V	A	F	L	L	E	V	R	R	A	G	M	S	L	I	G	N	K	V	D	L	G	P	N	S	F	R	L	G	F	V	T	W	V	H	Y	N	120	
prPL-6-1	81	W	C	V	V	A	F	L	L	E	V	R	R	A	G	M	S	L	I	G	N	K	V	D	L	G	P	N	S	F	R	L	G	F	V	T	W	V	H	Y	N	120	
prPL-6-B2	121	N	K	Y	V	E	L	L	D	T	L	W	M	V	L	R	K	K	T	Q	Q	V	S	F	F	L	H	V	Y	H	H	V	L	L	M	W	A	W	F	V	V	V	160
prPL-6-A3	121	N	K	Y	V	E	L	L	D	T	L	W	M	V	L	R	K	K	T	Q	Q	V	S	F	F	L	H	V	Y	H	H	V	L	L	M	W	A	W	F	V	V	V	160
prPL-6-1	121	N	K	Y	V	E	L	L	D	T	L	W	M	V	L	R	K	K	T	Q	Q	V	S	F	F	L	H	V	Y	H	H	V	L	L	M	W	A	W	F	V	V	V	160
prPL-6-B2	161	K	L	G	N	G	G	D	A	Y	F	G	G	L	M	N	S	I	I	H	V	M	M	Y	S	Y	Y	T	M	A	L	L	G	W	S	C	P	W	K	R	Y	200	
prPL-6-A3	161	K	L	G	N	G	G	D	A	Y	F	G	G	L	M	N	S	I	I	H	V	M	M	Y	S	Y	Y	T	M	A	L	L	G	W	S	C	P	W	K	R	Y	200	
prPL-6-1	161	K	L	G	N	G	G	D	A	Y	F	A	G	L	M	N	S	I	I	H	V	M	M	Y	S	Y	Y	T	M	A	L	L	G	W	S	C	P	W	K	R	Y	200	
prPL-6-B2	201	L	T	Q	A	Q	L	V	Q	F	C	I	C	L	A	H	S	T	W	A	A	V	T	G	A	Y	P	W	R	I	C	L	V	E	V	W	V	M	V	S	M	240	
prPL-6-A3	201	L	T	Q	A	Q	L	V	Q	F	C	I	C	L	A	H	S	T	W	A	A	V	T	G	A	Y	P	W	R	I	C	L	V	E	V	W	V	M	V	S	M	240	
prPL-6-1	201	L	T	Q	A	Q	L	V	Q	F	C	I	C	L	A	H	S	T	W	A	A	V	T	G	A	Y	P	W	R	I	C	L	V	E	V	W	V	M	V	S	M	240	
prPL-6-B2	241	L	V	L	F	T	R	F	Y	R	Q	A	Y	A	K	E	A	K	E	A	K	K	L	A	Q	E	A	S	Q	A	K	A	V	K	A	E					277		
prPL-6-A3	241	L	V	L	F	T	R	F	Y	C	Q	A	Y	A	K	E	A	K	E	A	K	K	L	A	Q	E	A	S	Q	A	K	A	V	K	A	E					277		
prPL-6-1	241	L	V	L	F	T	R	F	Y	R	Q	A	Y	A	K	E	A	K	E	A	K	K	L	A	Q	E	A	S	Q	A	K	A	V	K	A	E					277		

FIG.100

334(plasmid)	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2
added substrate	LA	LA	GLA	GLA	DGLA	DGLA	AA	AA
	% total lipid							
C18:2n-6	4.68	1.85						
C20:2n-6								
C18:3n-6			7.33	5.94				
C20:3n-6					7.32	5.58		
C22:3n-6								
C20:4n-6							13.98	8.57
C22:4n-6								1.40
% CONVERSION*	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	14.0%

334(plasmid)	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2
added substrate	ALA	ALA	STA	STA	ETA	ETA	EPA	EPA
	% total lipid							
C18:3n-3	6.97	3.87						
C20:3n-3								
C18:4n-3			5.22	3.53				
C20:4n-3					4.17	3.85		
C22:4n-3								
C20:5n-3							11.36	15.43
C22:5n-3								2.54
% CONVERSION*	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	14.1%

334(plasmid)	pYX242	pRPL-6-B2	pYX242	pRPL-6-B2
added substrate	C20:0	C20:0	C20:1	C20:1
	% total lipid			
C20:0	0.85	1.12		
C22:0				
C20:1			3.20	4.07
C22:1				
% CONVERSION*	0.0%	0.0%	0.0%	0.0%

$$\% \text{CONVERSION} = [\text{PRODUCT} / (\text{PRODUCT} + \text{SUBSTRATE})] \times 100$$

FIG.101

ELONGASE GENES AND USES THEREOF

[0001] The subject application is a Continuation-In-Part of pending U.S. patent application Ser. No. 09/903,456 filed on Jul. 11, 2001, which is a Continuation-In-Part of pending U.S. patent application Ser. No. 09/624,670 filed on Jul. 24, 2000, which is a Continuation-In-Part of pending U.S. patent application Ser. No. 09/379,095 filed on Aug. 23, 1999, which is a Continuation-In-Part of pending U.S. patent application Ser. No. 09/145,828 filed on Sep. 2, 1998, all of which are herein incorporated in their entirety by reference.

BACKGROUND OF THE INVENTION

TECHNICAL FIELD

[0002] The subject invention relates to the identification of several genes involved in the elongation of long-chain polyunsaturated fatty acids (i.e., "elongases") and to uses thereof. In particular, the elongase enzyme is utilized in the conversion of one fatty acid to another. For example, elongase catalyzes the conversion of gamma linolenic acid (GLA) to dihomo- γ -linolenic acid (DGLA, 20:3n-6) and the conversion of stearidonic acid (STA, 18:4n-3) to (n-3)-eicosatetraenoic acid (20:4n-3). Elongase also catalyzes the conversion of arachidonic acid (AA, 20:4n-6) to adrenic acid (ADA, 22:4n-6), the conversion of eicosapentaenoic acid (EPA, 20:5n-3) to ω 3-docosapentaenoic acid (22:5n-3), and the conversion of α -linolenic acid (ALA, 18:3n-3) to 20:3n-3. DGLA, for example, may be utilized in the production of other polyunsaturated fatty acids (PUFAs), such as arachidonic acid (AA) which may be added to pharmaceutical compositions, nutritional compositions, animal feeds, as well as other products such as cosmetics.

BACKGROUND INFORMATION

[0003] The elongases which have been identified in the past differ in terms of the substrates upon which they act. Furthermore, they are present in both animals and plants. Those found in mammals have the ability to act on saturated, monounsaturated and polyunsaturated fatty acids. In contrast, those found in plants are specific for saturated or monounsaturated fatty acids. Thus, in order to generate polyunsaturated fatty acids in plants, there is a need for a PUFA-specific elongase.

[0004] In both plants and animals, the elongation process is believed to be the result of a four-step mechanism (Lassner et al., *The Plant Cell* 8:281-292 (1996)). CoA is the acyl carrier. Step one involves condensation of malonyl-CoA with a long-chain acyl-CoA to yield carbon dioxide and a β -ketoacyl-CoA in which the acyl moiety has been elongated by two carbon atoms. Subsequent reactions include reduction to β -hydroxyacyl-CoA, dehydration to an enoyl-CoA, and a second reduction to yield the elongated acyl-CoA. The initial condensation reaction is not only the substrate-specific step but also the rate-limiting step.

[0005] As noted previously, elongases, more specifically, those which utilize PUFAs as substrates, are critical in the production of long-chain polyunsaturated fatty acids which have many important functions. For example, PUFAs are important components of the plasma membrane of a cell where they are found in the form of phospholipids. They also serve as precursors to mammalian prostacyclins,

eicosanoids, leukotrienes and prostaglandins. Additionally, PUFAs are necessary for the proper development of the developing infant brain as well as for tissue formation and repair. In view of the biological significance of PUFAs, attempts are being made to produce them, as well as intermediates leading to their production, efficiently.

[0006] A number of enzymes are involved in PUFA biosynthesis including elongases (elo) (see FIG. 1). For example, linoleic acid (LA, 18:2- Δ 9,12 or 18:2n-6) is produced from oleic acid (OA, 18:1- Δ 9 or 18:1n-9) by a Δ 12 desaturase. GLA (18:3- Δ 6,9,12) is produced from linoleic acid by a Δ 6-desaturase. AA (20:4- Δ 5,8,11,14) is produced from dihomo- γ -linolenic acid (DCLA, 20:3- Δ 8,11,14) by a Δ 5-desaturase. As noted above, DGLA is produced from GLA by an elongase.

[0007] It must be noted that animals cannot desaturate beyond the Δ 9 position and therefore cannot convert oleic acid into linoleic acid. Likewise, α -linolenic acid (ALA, 18:3- Δ 9,12,15 or 18:3n-3) cannot be synthesized by mammals, since they lack Δ 15 desaturase activity. However, α -linolenic acid can be converted to stearidonic acid (STA, 18:4- Δ 6,9,12,15) by a Δ 6-desaturase (see PCT publication WO 96/13591; see also U.S. Pat. No. 5,552,306), followed by elongation to (n-3)-eicosatetraenoic acid (20:4- Δ 8,11,14,17 or 20:4n-3) in mammals and algae. This polyunsaturated fatty acid (i.e., 20:4- Δ 8,11,14,17) can then be converted to eicosapentaenoic acid (EPA, 20:5- Δ 5,8,11,14,17) by a Δ 5-desaturase. Other eukaryotes, including fungi and plants, have enzymes which desaturate at carbons 12 (see PCT publication WO 94/11516 and U.S. Pat. No. 5,443,974) and 15 (see PCT publication WO 93/11245). The major polyunsaturated fatty acids of animals therefore are either derived from diet and/or from desaturation and elongation of linoleic acid or α -linolenic acid. In view of the inability of mammals to produce these essential long chain fatty acids, it is of significant interest to isolate genes involved in PUFA biosynthesis from species that naturally produce these fatty acids and to express these genes in a microbial, plant or animal system which can be altered to provide production of commercial quantities of one or more PUFAs. Consequently, there is a definite need for the elongase enzyme, the gene encoding the enzyme, as well as recombinant methods of producing this enzyme. Additionally, a need exists for oils containing levels of PUFA beyond those naturally present as well as those enriched in novel PUFAs. Such oils can only be made by isolation and expression of the elongase gene.

[0008] One of the most important long chain PUFAs, noted above, is arachidonic acid (AA). AA is found in filamentous fungi and can also be purified from mammalian tissues including the liver and the adrenal glands. As noted above, AA production from DGLA is catalyzed by a Δ 5-desaturase, and DGLA production from γ -linolenic acid (GLA) is catalyzed by an elongase. However, until the present invention, no elongase had been identified which was active on substrate fatty acids in the pathways for the production of long chain PUFAs and, in particular, AA, eicosapentaenoic acid (EPA), adrenic acid, docosahexaenoic acid (DHA, 22:6n-3), ω 3-docosapentaenoic acid (22:5n-3) or ω 6-docosapentaenoic acid (22:5n-6).

[0009] Two genes appeared to be of interest in the present search for the elongase gene. In particular, the jojoba β -ketoacyl-coenzyme A synthase (KCS), or jojoba KCS (Cen-

Bank Accession #U37088), catalyzes the initial reaction of the fatty acyl-CoA elongation pathway (i.e., the condensation of malonyl-CoA with long-chain acyl-CoA (Lassner et al., *The Plant Cell* 8:281-292 (1996)). Jojoba KCS substrate preference is 18:0, 20:0, 20:1, 18:1, 22:1, 22:0 and 16:0. *Saccharomyces cerevisiae* elongase (ELO2) also catalyzes the conversion of long chain saturated and monounsaturated fatty acids, producing high levels of 22:0, 24:0, and also 18:0, 18:1, 20:0, 20:1, 22:0, 22:1, and 24:1 (Oh et al., *The Journal of Biological Chemistry* 272 (28):17376-17384 (1997); see also U.S. Pat. No. 5,484,724 for a nucleotide sequence which includes the sequence of ELO2; see PCT publication WO 88/07577 for a discussion of the sequence of a glycosylation inhibiting factor which is described in Example V). The search for a long chain PUFA-specific elongase in *Mortierella alpina* began based upon a review of the homologies shared between these two genes and by expression screening for PUFA-elongase activity.

SUMMARY OF THE INVENTION

[0010] The present invention encompasses an isolated nucleotide sequence comprising or complementary to at least about 50% of the nucleotide sequence shown in SEQ ID NO:1 (**FIG. 6**). This isolated sequence may be represented by SEQ ID NO:1. The sequence encodes a functionally active elongase which utilizes a polyunsaturated fatty acid or a monounsaturated fatty acid as a substrate. In particular, the sequence may be derived from a fungus of the genus *Mortierella* and may specifically be isolated from *Mortierella alpina*. Additionally, the present invention encompasses an isolated nucleotide sequence that encodes a polypeptide. The polypeptide comprises an amino acid sequence having at least 50% amino acid sequence identity to SEQ ID NO:13 and has elongase activity.

[0011] The present invention also includes a purified protein encoded by the above nucleotide sequence as well as a purified polypeptide which elongates polyunsaturated fatty acids or monounsaturated fatty acids and has at least about 50% amino acid similarity to the amino acid sequence of the purified protein (SEQ ID NO:13) encoded by the above nucleotide sequence.

[0012] Additionally, the present invention encompasses a method of producing an elongase enzyme comprising the steps of: a) isolating the nucleotide sequence comprising SEQ ID NO:1 (**FIG. 6**); b) constructing a vector comprising: i) the isolated nucleotide sequence operably linked to ii) a promoter; and c) introducing the vector into a host cell under time and conditions sufficient for expression of the elongase enzyme. The host cell may be a eukaryotic cell or a prokaryotic cell.

[0013] The prokaryotic cell may be, for example an *E. coli* cell, a cyanobacterial cell, or a *B. subtilis* cell. The eukaryotic cell may be, for example, a mammalian cell, an insect cell, a plant cell or a fungal cell. The fungal cell may be, for example, *Saccharomyces* spp., *Candida* spp., *Lipomyces* spp., *Yarrowia* spp., *Kluyveromyces* spp., *Hansenula* spp., *Aspergillus* spp., *Penicillium* spp., *Neurospora* spp., *Trichoderma* spp. or *Pichia* spp. In particular, the fungal cell may be a yeast cell such as *Saccharomyces* spp., in particular, *Saccharomyces cerevisiae*, *Candida* spp., *Hansenula* spp. or *Pichia* spp.

[0014] The invention also includes a vector comprising: a) a nucleotide sequence comprising SEQ ID NO:1 (**FIG. 6**)

operably linked to b) a promoter, as well as a host cell comprising this vector. The host may be a prokaryotic cell or a eukaryotic cell. Suitable examples of prokaryotic cells include *E. coli*, Cyanobacteria, and *B. subtilis* cells. Suitable examples of eukaryotic cells include a mammalian cell, an insect cell, a plant cell and a fungal cell. The fungal cell may be, for example, *Saccharomyces* spp., *Candida* spp., *Lipomyces* spp., *Yarrowia* spp., *Kluyveromyces* spp., *Hansenula* spp., *Aspergillus* spp., *Penicillium* spp., *Neurospora* spp., *Trichoderma* spp. and *Pichia* spp. In particular, the fungal cell may be, for example, a yeast cell such as, for example, *Saccharomyces* spp., in particular, *Saccharomyces cerevisiae*, *Candida* spp., *Hansenula* spp. and *Pichia* spp.

[0015] The present invention includes a plant cell, plant or plant tissue comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of at least one fatty acid selected from the group consisting of a monounsaturated fatty acid and a polyunsaturated fatty acid by the plant cell, plant or plant tissue. The polyunsaturated fatty acid may be, for example, dihomo- γ -linolenic acid (DGLA), 20:4n-3, and adrenic acid (ADA). The invention also includes one or more plant oils or fatty acids expressed by the plant cell, plant or plant tissue. Additionally, the present invention encompasses a transgenic plant comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0016] Furthermore, the present invention includes a transgenic, non-human mammal whose genome comprises a DNA sequence encoding an elongase operably linked to a promoter. The DNA sequence may be represented by SEQ ID NO:1 (**FIG. 6**). The present invention also includes a fluid (e.g., milk) produced by the transgenic, non-human wherein the fluid comprises a detectable level of at least one elongase or products thereof such as, for example, DGLA, ω 6-docosapentaenoic acid, ADA and/or 20:4n-3 (see **FIG. 1**).

[0017] Additionally, the present invention includes a method for producing a polyunsaturated fatty acid comprising the steps of: a) isolating said nucleotide sequence comprising SEQ ID NO:1 (**FIG. 6**); b) constructing a vector comprising the isolated nucleotide sequence; c) introducing the vector into a host cell under time and conditions sufficient for expression of elongase enzyme encoded by the isolated nucleotide sequence; and d) exposing the expressed elongase enzyme to a "substrate" polyunsaturated fatty acid in order to convert the substrate to a "product" polyunsaturated fatty acid. The substrate polyunsaturated fatty acid may be selected from the group consisting of, for example, γ -linolenic acid (GLA), stearidonic acid (STA) and arachidonic acid (AA), and the product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3, and ADA, respectively. The method may further comprise the step of exposing the product polyunsaturated fatty acid to at least one desaturase in order to convert the product polyunsaturated fatty acid to "another" polyunsaturated fatty acid. The product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3, and ADA. The another polyunsaturated fatty acid may be selected from the group consisting of, for example, AA, eicosapentaenoic acid (EPA), ω 6-docosapentaenoic acid, respectively, and the at least one

desaturase is $\Delta 5$ -desaturase, with respect to production of AA or EPA, and $\Delta 4$ -desaturase, with respect to production of $\omega 6$ -docosapentaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a "final" polyunsaturated fatty acid. The final polyunsaturated fatty acid may be, for example, docosahexaenoic acid (DHA), AA, $\omega 6$ -docosapentaenoic acid, or $\omega 3$ -docosapentaenoic acid.

[0018] Also, the present invention includes a nutritional composition comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method. The product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3 and ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA, or $\omega 6$ -docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, adrenic acid, $\omega 6$ -docosapentaenoic acid or $\omega 3$ -docosapentaenoic acid.

[0019] The nutritional composition may be, for example, an infant formula, a dietary supplement or a dietary substitute and may be administered to a human or an animal and may be administered enterally or parenterally. The nutritional composition may further comprise at least one macro-nutrient selected from the group consisting of coconut oil, soy oil, canola oil, monoglycerides, diglycerides, triglycerides, glucose, edible lactose, electrodialysed whey, electrodialysed skim milk, milk whey, soy protein, protein hydrolysates, sunflower oil, safflower oil, corn oil, and flax oil. It may also comprise at least one vitamin selected from the group consisting of Vitamins A, C, D, E, and B complex and at least one mineral selected from the group consisting of calcium magnesium, zinc, manganese, sodium, potassium, phosphorus, copper, chloride, iodine, selenium and iron.

[0020] Additionally, the present invention encompasses a pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method and 2) a pharmaceutically acceptable carrier. The composition may be administered to a human or an animal. It may also further comprise at least one element selected from the group consisting of a vitamin, a mineral, a salt, a carbohydrate, an amino acid, a free fatty acid, a preservative, an excipient, an anti-histamine, a growth factor, an antibiotic, a diluent, a phospholipid, an antioxidant, and a phenolic compound. It may be administered enterally, parenterally, topically, rectally, intramuscularly, subcutaneously, intradermally, or by any other appropriate means.

[0021] The present invention also includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated

fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, and ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA, or $\omega 6$ -docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, adrenic acid, $\omega 6$ -docosapentaenoic acid or $\omega 3$ -docosapentaenoic acid.

[0022] Moreover, the present invention also includes a cosmetic comprising a polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method.

[0023] Additionally, the present invention includes a method of preventing or treating a condition caused by insufficient intake or production of polyunsaturated fatty acids comprising administering to the patient the above nutritional composition in an amount sufficient to effect prevention or treatment.

[0024] The present invention also includes an isolated nucleotide sequence comprising or complementary to at least about 35% of the nucleotide sequence shown in SEQ ID NO:2 (**FIG. 22**). This sequence may be represented by SEQ ID NO:2. The sequence encodes a functionally active elongase which utilizes a polyunsaturated fatty acid as a substrate. This sequence may also be derived, for example, from a fungus of the genus *Mortierella*. In particular, it may be derived from *M. alpina*. Also, the present invention includes an isolated nucleotide sequence which encodes a polypeptide, wherein the polypeptide comprises an amino acid sequence having at least 30% amino acid sequence identity to SEQ ID NO:31 and which encodes an elongase.

[0025] Additionally, the present invention includes a purified protein encoded by the above nucleotide sequence (SEQ ID NO:31) as well as a purified polypeptide which elongates polyunsaturated fatty acids and has at least about 30% amino acid identity to the amino acid sequence of the purified protein (SEQ ID NO:31).

[0026] The present invention also includes a method of producing an elongase enzyme as described above. The sequence inserted in the vector is represented by SEQ ID NO:2 (**FIG. 22**). The host cell may be prokaryotic or eukaryotic. Suitable examples are described above.

[0027] The present invention also includes a vector comprising: a) a nucleotide sequence as represented by SEQ ID NO:2 (**FIG. 22**) operably linked to b) a promoter, as well as a host cell comprising this vector. Again, the host cell may be eukaryotic or prokaryotic. Suitable examples are described above.

[0028] The invention also includes a plant cell, plant or plant tissue comprising the above vector, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid by the plant cell, plant or plant tissue. The polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA. Additionally, the inven-

tion includes one or more plant oils or fatty acids expressed by the plant cell, plant or plant tissue.

[0029] Furthermore, the present invention also includes a transgenic plant comprising the above vector, wherein expression of the nucleotide sequence (SEQ ID NO:2) of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0030] The invention also includes a transgenic, non-human mammal whose genome comprises a DNA sequence (SEQ ID NO:2) encoding an elongase operably linked to a promoter. The invention also includes a fluid produced by this transgenic, non-human mammal wherein the fluid comprises a detectable level of at least one elongase or products thereof.

[0031] The present invention also includes a method for producing a polyunsaturated fatty acid comprising the steps of: a) isolating the nucleotide sequence represented by SEQ ID NO:2 (**FIG. 22**); b) constructing a vector comprising the isolated nucleotide sequence; c) introducing the vector into a host cell under time and conditions sufficient for expression of an elongase enzyme encoded by the isolated nucleotide sequence; and d) exposing the expressed elongase enzyme to a substrate polyunsaturated fatty acid in order to convert the substrate to a product polyunsaturated fatty acid. The substrate polyunsaturated fatty acid may be, for example, GLA, STA, or AA, the product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ω 6-docosapentaenoic acid, respectively.

[0032] The method may further comprise the step of exposing the expressed elongase enzyme to at least one desaturase in order to convert the product polyunsaturated fatty acid to another polyunsaturated fatty acid. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA, the another polyunsaturated fatty acid may be, for example, AA, EPA, or ω 6-docosapentaenoic acid, respectively, and the at least one desaturase is Δ 5-desaturase with respect to production of AA or EPA, and Δ 4-desaturase with respect to production of ω 6-docosapentaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a final polyunsaturated fatty acid. The final polyunsaturated fatty acid may be, for example, docosahexaenoic acid, AA, ω 6-docosapentaenoic acid, or ω 3-docosapentaenoic acid.

[0033] The invention also includes a nutritional composition comprising at least one polyunsaturated fatty acid selected from the product polyunsaturated fatty acid produced according to the method described with respect to SEQ ID NO:2, the another polyunsaturated fatty acid produced according to the method described with respect to SEQ ID NO:2, and the final polyunsaturated fatty acid produced according to the method described with respect to SEQ ID NO:2. The product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3 and ADA. The another polyunsaturated fatty acid may be selected from the group consisting of, for example, AA, EPA, and ω 6-docosapentaenoic acid. The final polyunsaturated fatty acid may be selected from the group consisting of, for example, DHA, AA, ω 6-docosapentaenoic acid, and ω 3-docosapentaenoic acid.

[0034] The other attributes of the composition are the same as those described above with respect to administration, characterization, components, etc.

[0035] The present invention also includes a pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method of noted above with respect to SEQ ID NO:2, the another polyunsaturated fatty acid produced according to the method described above with respect to SEQ ID NO:2, and the final polyunsaturated fatty acid produced according to the method described above with respect to SEQ ID NO:2, and 2) a pharmaceutically acceptable carrier. The characteristics of the above-described pharmaceutical composition (e.g., administration, components, etc.) also apply to this composition.

[0036] The present invention also includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of: the product polyunsaturated fatty acid produced according to the method described with respect to SEQ ID NO:2, the another polyunsaturated fatty acid produced according to the method described above with respect to SEQ ID NO:2, and the final polyunsaturated fatty acid produced according to the method described with respect to SEQ ID NO:2.

[0037] The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3 or ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA or ω 6-docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, adrenic acid, ω 6-docosapentaenoic acid or ω 3-docosapentaenoic acid.

[0038] The invention also includes a cosmetic comprising a polyunsaturated fatty acid selected from the group consisting of: the product polyunsaturated fatty acid produced according to the method described above with respect to SEQ ID NO:2, the another polyunsaturated fatty acid produced according to the method described above with respect to SEQ ID NO:2, and the final polyunsaturated fatty acid produced according to the method described above with respect to SEQ ID NO:2.

[0039] Additionally, the present invention includes a method of preventing or treating a condition caused by insufficient intake or production of polyunsaturated fatty acids comprising administering to the patient the nutritional composition described directly above in an amount sufficient to effect the prevention or treatment.

[0040] Furthermore, the present invention includes an isolated nucleotide sequence comprising or complementary to at least about 35% of the nucleotide sequence shown in SEQ ID NO:3 (**FIG. 43**). This sequence may be that represented by SEQ ID NO:3. This sequence encodes a functionally active elongase which utilizes a polyunsaturated fatty acid or a monounsaturated fatty acid as a substrate. The sequence is derived from a mammal such as, for example, a human. Also, the present invention includes an isolated nucleotide sequence which encodes a polypeptide, wherein the amino acid sequence of the polypeptide has at least about 30% amino acid sequence identity to SEQ ID NO:61 and has elongase activity.

[0041] The invention also includes a purified protein (SEQ ID NO:61) encoded by this nucleotide sequence. Also, the

invention includes a purified polypeptide which elongates polyunsaturated fatty acids or monounsaturated fatty acids and has at least about 30% amino acid similarity to the amino acid sequence of this purified protein (SEQ ID NO:61).

[0042] Additionally, the invention includes method of producing an elongase enzyme comprising the steps of: a) isolating the nucleotide sequence represented by SEQ ID NO:3 (FIG. 43); b) constructing a vector comprising: i) the isolated nucleotide sequence operably linked to ii) a promoter; and c) introducing said vector into a host cell under time and conditions sufficient for expression of the elongase enzyme. The host cell may be the same as that described above with respect to the corresponding methods utilizing SEQ ID NO:1 or 2.

[0043] The invention also includes a vector comprising: a) a nucleotide sequence as represented by SEQ ID NO:3 (FIG. 43) operably linked to b) a promoter, as well as a host cell comprising this vector. The host cell may be the same as that described above.

[0044] Moreover, the invention also includes a plant cell, plant or plant tissue comprising the above-described vector comprising SEQ ID NO:3, wherein expression of the nucleotide sequence of the vector results in production of at least one fatty acid selected from the group consisting of a monounsaturated fatty acid and a polyunsaturated fatty acid by said plant cell, plant or plant tissue. The polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3 or ADA. The invention also includes one or more plant oils or acids expressed by the plant cell, plant or plant tissue.

[0045] The invention also includes a transgenic plant comprising the vector comprising SEQ ID NO:3, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0046] Additionally, the present invention includes a transgenic, non-human mammal whose genome comprises a human DNA sequence encoding an elongase operably linked to a promoter. The DNA sequence is represented by SEQ ID NO:3 (FIG. 43). The invention also includes a fluid produced by said transgenic, non-human mammal wherein said fluid comprises a detectable level of at least one elongase or products thereof.

[0047] The invention also encompasses a method for producing a polyunsaturated fatty acid comprising the steps of: a) isolating the nucleotide sequence comprising SEQ ID NO:3 (FIG. 43); b) constructing a vector comprising said nucleotide sequence; c) introducing the vector into a host cell under time and conditions sufficient for expression of elongase enzyme encoded by the isolated nucleotide sequence; and d) exposing the expressed elongase enzyme to a substrate polyunsaturated fatty acid in order to convert the substrate to a product polyunsaturated fatty acid. The substrate polyunsaturated fatty acid may be, for example, GLA, STA or AA, and the product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA, respectively. The method may further comprise the step of exposing the product polyunsaturated fatty acid to at least one desaturase in order to convert the product polyunsaturated fatty acid to another polyunsaturated fatty acid. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3 and

ADA, the another polyunsaturated fatty acid may be, for example, AA, EPA, and ω 6-docosapentaenoic acid, respectively, and the at least one desaturase is Δ 5-desaturase with respect to production of AA or EPA and Δ 4-desaturase with respect to production of ω 6-docosapentaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a final polyunsaturated fatty acid. The final polyunsaturated fatty acid may be, for example, DHA, ADA, ω 6-docosapentaenoic acid, and ω 3-docosapentaenoic acid.

[0048] The nutritional composition comprising at least one polyunsaturated fatty acid which may be, for example, product polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:3, another polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:3, and the final polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:3. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA. The another polyunsaturated fatty acid may be selected from the group consisting of AA, EPA, or ω 6-docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, ADA, ω 6-docosapentaenoic acid, or ω 3-docosapentaenoic acid.

[0049] The other properties or characteristic of the nutritional composition (e.g., administration, components, etc.) as the same as those recited above with respect to the other nutritional compositions.

[0050] Moreover, the present invention also includes a pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of: the product polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:3, the another polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:3, and the final polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:3 and 2) a pharmaceutically acceptable carrier. The other properties of the composition (e.g., administration, additional components, etc.) are the same as those recited above with respect to the other pharmaceutical compositions.

[0051] The present invention also includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of: the product polyunsaturated fatty acid produced according to the method recited above with respect to SEQ ID NO:3, the another polyunsaturated fatty acid produced according to the method recited above with respect to SEQ ID NO:3, and the final polyunsaturated fatty acid produced according to the method recited above with respect to SEQ ID NO:3. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA. The polyunsaturated fatty acid may be, for example, AA, EPA, or ω 6-docosapentaenoic acid.

[0052] The final polyunsaturated fatty acid may be, for example, DHA, ADA, ω 6-docosapentaenoic acid or ω 3-docosapentaenoic acid.

[0053] Also, the present invention includes a cosmetic comprising a polyunsaturated fatty acid selected from the

group consisting of: the product polyunsaturated fatty acid produced according to the method recited above with respect to SEQ ID NO:3, said another polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:3, and the final polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:3.

[0054] A method of preventing or treating a condition caused by insufficient intake of polyunsaturated fatty acids comprising administering to the patient the nutritional composition recited above in connection with SEQ ID NO:3 in an amount sufficient to effect the prevention or treatment.

[0055] Additionally, the present invention includes an isolated nucleotide sequence comprising or complementary to at least about 35% of the nucleotide sequence shown in SEQ ID NO:4 (FIG. 46). The sequence may be represented by SEQ ID NO:4. It encodes a functionally active elongase which utilizes a polyunsaturated fatty acid as a substrate. The sequence may be derived or isolated from a nematode of the genus *Caenorhabditis* and, in particular, may be isolated from *C. elegans*. Further, the present invention includes an isolated nucleotide sequence which encodes a polypeptide, wherein the polypeptide comprises an amino acid sequence having at least about 30% amino acid identity to SEQ ID NO:62 and has elongase activity.

[0056] The present invention includes a purified protein (SEQ ID NO:62) encoded by the nucleotide sequence above. The invention also includes a purified polypeptide which elongates polyunsaturated fatty acids and has at least about 30% amino acid similarity to the amino acid sequence of the purified protein.

[0057] Additionally, the present invention includes a method of producing an elongase enzyme comprising the steps of: a) isolating the nucleotide sequence represented by SEQ ID NO:4 (FIG. 46); b) constructing a vector comprising: i) the isolated nucleotide sequence operably linked to ii) a promoter; and c) introducing the vector into a host cell under time and conditions sufficient for expression of the elongase enzyme. The properties of the host cell are the same as those described above in connection with SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:3.

[0058] The present invention also encompasses a vector comprising: a) a nucleotide sequence as represented by SEQ ID NO:4 (FIG. 46) operably linked to b) a promoter, as well as a host cell comprising this vector. The host cell has the same properties as those recited above in connection with the host cell recited above for SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3.

[0059] Moreover, the present invention includes a plant cell, plant or plant tissue comprising the above vector comprising SEQ ID NO:4, wherein expression of said nucleotide sequence of the vector results in production of a polyunsaturated fatty acid by the plant cell, plant or plant tissue. The polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA. The invention also includes one or more plant oils or fatty acids expressed by this plant cell, plant or plant tissue.

[0060] The invention also includes transgenic plant comprising the above vector including the nucleotide sequence corresponding to SEQ ID NO:4, wherein expression of the

nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0061] Additionally, the present invention includes a transgenic, non-human mammal whose genome comprises a *C. elegans* DNA sequence encoding an elongase operably linked to a promoter.

[0062] The DNA sequence may be represented by SEQ ID NO:4 (FIG. 46).

[0063] The invention also includes a fluid produced by the transgenic, non-human mammal, wherein the fluid comprises a detectable level of at least one elongase or products thereof.

[0064] The invention also includes a method for producing a polyunsaturated fatty acid comprising the steps of: a) isolating the nucleotide sequence represented by SEQ ID NO:4 (FIG. 46); b) constructing a vector comprising the isolated nucleotide sequence; c) introducing the vector into a host cell under time and conditions sufficient for expression of an elongase enzyme encoded by the isolated nucleotide sequence; and d) exposing the expressed elongase enzyme to a substrate polyunsaturated fatty acid in order to convert the substrate to a product polyunsaturated fatty acid. The substrate polyunsaturated fatty acid may be, for example, GLA, STA, or AA, and the product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA, respectively. The method may further comprise the step of exposing the expressed elongase enzyme to at least one desaturase in order to convert said product polyunsaturated fatty acid to another polyunsaturated fatty acid. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3 or ADA, the another polyunsaturated fatty acid may be, for example, AA, EPA or ω 6-docosapentaenoic acid, respectively, and the at least one desaturase is Δ 5-desaturase with respect to production of AA or EPA, and Δ 4-desaturase with respect to production of ω 6-docosapentaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a final polyunsaturated fatty acid. The final polyunsaturated fatty acid may be, for example, DHA, ADA, ω 6-docosapentaenoic acid, or ω 3-docosapentaenoic acid.

[0065] The invention also includes a nutritional composition comprising at least one polyunsaturated fatty acid selected from the group consisting of said the polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:4, the another polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:4, and the final polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, or ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA, or ω 6-docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, ADA, ω 6-docosapentaenoic acid, or ω 3-docosapentaenoic acid. The other characteristics of the composition are the same as those recited for the nutritional compositions present above.

[0066] Additionally, the present invention includes a pharmaceutical composition comprising 1) at least one polyun-

saturated fatty acid selected from the group consisting of: the product polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4, the another polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4, and the final polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4 and 2) a pharmaceutically acceptable carrier. The composition has the same properties (e.g., administration, added elements, etc.) as those described above with respect to the other pharmaceutical compositions.

[0067] The present invention also includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:4, the another polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4, and the final polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:4. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3 or ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA or ω 6-docosapentaenoic acid. The polyunsaturated fatty acid may be, for example, DHA, ADA, ω 6-docosapentaenoic acid or ω 3-docosapentaenoic acid.

[0068] Additionally, the present invention includes a cosmetic comprising a polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4, the another polyunsaturated fatty acid produced according to the method recited above in connection with SEQ ID NO:4 and the final polyunsaturated fatty acid produced according to the method described above in connection with SEQ ID NO:4.

[0069] Furthermore, the present invention encompasses a method of preventing or treating a condition caused by insufficient intake or production of polyunsaturated fatty acids comprising administering to the patient the nutritional composition recited with respect to SEQ ID NO:4 in an amount sufficient to effect the treatment or prevention.

[0070] The present invention also includes an isolated nucleotide sequence comprising or complementary to at least about 35% of the nucleotide sequence comprising SEQ ID NO:5 (**FIG. 54**). Thus, the sequence may be that represented by SEQ ID NO:5. The sequence may encode a functionally active elongase which utilizes a polyunsaturated fatty acid as a substrate. It may also be derived from a mammal such as, for example, a mouse.

[0071] The present invention also includes a purified protein (SEQ ID NO:65) encoded by the nucleotide sequence as well as a purified polypeptide which elongates polyunsaturated fatty acids and has at least about 30% amino acid sequence identity to the amino acid sequence of the protein. Also, the present invention includes an isolated nucleotide sequence encoding a polypeptide, wherein the polypeptide comprises an amino acid sequence having at least about 30% amino acid sequence identity to SEQ ID NO:65 and has elongase activity.

[0072] Additionally, the invention also includes a method of producing an elongase enzyme, as described above, in

which the nucleotide sequence isolated comprises either SEQ ID NO:5 or SEQ ID NO:6. The host cell utilized may be as described above.

[0073] The present invention also encompasses a vector comprising: a) a nucleotide sequence comprising SEQ ID NO:5 (**FIG. 54**) (or a nucleotide sequence comprising SEQ ID NO:6 (**FIG. 58**)) operably linked to b) a promoter, as well as a host cell comprising this vector. Again, the host cell may be as described above for the related methods using the other nucleotide sequences of the present invention.

[0074] Additionally, the invention includes a plant cell, plant or plant tissue comprising the vector comprising SEQ ID NO:5 or 6, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid by the plant cell, plant or plant tissue. When the nucleotide sequence of the vector comprises SEQ ID NO:5, the polyunsaturated fatty acid is selected from the group consisting of AA, ADA, GLA and STA. The invention also includes one or more plant oils or acids expressed by the plant cell, plant or plant tissue.

[0075] The present invention also includes a transgenic plant comprising the vector described above, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0076] Additionally, the present invention encompasses a transgenic, non-human mammal whose genome comprises a DNA sequence encoding an elongase, operably linked to a promoter, wherein the DNA sequence comprises SEQ ID NO:5 (**FIG. 54**) (or SEQ ID NO:6 (**FIG. 58**)). Also, the invention includes a fluid produced by this transgenic, non-human mammal, wherein the fluid comprises a detectable level of at least one elongase or products thereof.

[0077] The invention also includes a method for producing a polyunsaturated fatty acid, similar to the methods described above, except that the isolated nucleotide sequence comprises SEQ ID NO:5 (**FIG. 54**). The substrate polyunsaturated fatty acid may be selected from the group consisting of GLA, STA, AA, ADA and ALA, and the product polyunsaturated fatty acid may be selected from the group consisting of DGLA, 20:4n-3, ADA, ω 6-docosapentaenoic acid and STA, respectively. The method may further comprise the step of exposing the expressed elongase enzyme to at least one desaturase in order to convert the product polyunsaturated fatty acid to another polyunsaturated fatty acid. The product polyunsaturated fatty acid may be selected from the group consisting of DGLA, 20:4n-3, ADA and ω 6-docosapentaenoic acid, the another polyunsaturated fatty acid is selected from the group consisting of AA, EPA, ω 6-docosapentaenoic acid and docosahexaenoic acid respectively, and the at least one desaturase is Δ 5-desaturase with respect to production of AA or EPA, and Δ 4-desaturase with respect to production of ω 6-docosapentaenoic acid, and Δ 19-desaturase with respect to production of docosahexaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a final polyunsaturated fatty acid. The final polyunsaturated fatty acid may be selected from the group consisting of ADA, ω 3-docosapentaenoic acid and docosahexaenoic acid.

[0078] The present invention also includes a nutritional composition comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method above, the another polyunsaturated fatty acid is produced according to the method above, and the final polyunsaturated fatty acid produced according to the method above. The product polyunsaturated fatty acid may be selected from the group consisting of DGLA, 20:4n-3, ADA, and ω 6-docosapentaenoic acid and STA. The another polyunsaturated fatty acid is selected from the group consisting of AA, EPA, ω 6-docosapentaenoic acid and docosahexaenoic acid. The final polyunsaturated fatty acid is selected from the group consisting of ADA, ω 3-docosapentaenoic acid and docosahexaenoic acid. The nutritional composition may be selected from the group consisting of an infant formula, a dietary supplement and a dietary substitute.

[0079] The present invention also includes a pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method above, the another polyunsaturated fatty acid produced according to the method above, and the final polyunsaturated fatty acid produced according to the method above and 2) a pharmaceutically acceptable carrier.

[0080] Additionally, the present invention includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method above, the another polyunsaturated fatty acid produced according to the method above and the final polyunsaturated fatty acid produced according to the method above. The product polyunsaturated fatty acid may be selected from the group consisting of DGLA, 20:4n-3, ADA, ω 6-docosapentaenoic acid and STA. The another polyunsaturated fatty acid may be selected from the group consisting of AA, EPA, ω 6-docosapentaenoic acid and docosahexaenoic acid. The final polyunsaturated fatty acid may be selected from the group consisting of ADA, ω 3-docosapentaenoic acid and docosahexaenoic acid.

[0081] The invention includes a cosmetic comprising a polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method above, the another polyunsaturated fatty acid produced according to the method above and the final polyunsaturated fatty acid produced according to the method above.

[0082] Additionally, a method of preventing or treating a condition caused by insufficient intake of polyunsaturated fatty acids comprising administering to the patient the nutritional composition above in an amount sufficient to effect the prevention or treatment.

[0083] The present invention includes an isolated nucleotide sequence comprising or complementary to at least about 35% of the nucleotide sequence comprising SEQ ID NO:6 (FIG. 58). The isolated nucleotide sequence may comprise SEQ ID NO:6. The invention also includes a purified protein (SEQ ID NO:66) encoded by the nucleotide sequences. Furthermore, the present invention encompasses an isolated nucleotide sequence encoding a polypeptide, wherein the polypeptide comprises an amino acid sequence having at least about 35% amino acid sequence identity to SEQ ID NO:66 and has elongase activity.

[0084] Additionally, the present invention relates to an isolated nucleotide sequence comprising or complementary to at least about 35% of the nucleotide sequence shown in SEQ ID NO:7 (FIG. 72). This isolated sequence may be represented by SEQ ID NO:7. The sequence encodes a functionally active elongase which utilizes a polyunsaturated fatty acid or a monounsaturated fatty acid as a substrate. In particular, the sequence may be derived from a fungus of the genus *Thraustochytrium* and may specifically be isolated from *Thraustochytrium aureum*. The present invention also encompasses an isolated nucleotide sequence which encodes a polypeptide, wherein the polypeptide comprises an amino acid sequence having at least about 50% amino acid sequence identity to SEQ ID NO:80 and has elongase activity.

[0085] The present invention also includes a purified protein (SEQ ID NO:80) encoded by the above nucleotide sequence as well as a purified polypeptide which elongates polyunsaturated fatty acids or monounsaturated fatty acids and has at least about 50% amino acid identity to the amino acid sequence of the purified protein encoded by the above nucleotide sequence.

[0086] Additionally, the present invention encompasses a method of producing an elongase enzyme comprising the steps of: a) isolating the nucleotide sequence represented by SEQ ID NO:7 (FIG. 72); b) constructing a vector comprising: i) the isolated nucleotide sequence operably linked to ii) a promoter; and c) introducing the vector into a host cell under time and conditions sufficient for expression of the elongase enzyme. The host cell may be a eukaryotic cell or a prokaryotic cell.

[0087] The prokaryotic cell may be, for example an *E. coli* cell, a cyanobacterial cell, or a *B. subtilis* cell. The eukaryotic cell may be, for example, a mammalian cell, an insect cell, a plant cell or a fungal cell. The fungal cell may be, for example, *Saccharomyces* spp., *Candida* spp., *Lipomyces* spp., *Yarrowia* spp., *Kluyveromyces* spp., *Hansenula* spp., *Aspergillus* spp., *Penicillium* spp., *Neurospora* spp., *Trichoderma* spp. or *Pichia* spp. In particular, the fungal cell may be a yeast cell such as *Saccharomyces* spp., in particular, *Saccharomyces cerevisiae*, *Candida* spp., *Hansenula* spp. or *Pichia* spp.

[0088] The invention also includes a vector comprising: a) a nucleotide sequence as represented by SEQ ID NO:7 (FIG. 72) operably linked to b) a promoter, as well as a host cell comprising this vector. The host may be a prokaryotic cell or a eukaryotic cell. Suitable examples of prokaryotic cells include *E. coli*, *Cyanobacteria*, and *B. subtilis* cells. Suitable examples of eukaryotic cells include a mammalian cell, an insect cell, a plant cell and a fungal cell. The fungal cell may be, for example, *Saccharomyces* spp., *Candida* spp., *Lipomyces* spp., *Yarrowia* spp., *Kluyveromyces* spp., *Hansenula* spp., *Aspergillus* spp., *Penicillium* spp., *Neurospora* spp., *Trichoderma* spp. and *Pichia* spp. In particular, the fungal cell may be, for example, a yeast cell such as, for example, *Saccharomyces* spp., in particular, *Saccharomyces cerevisiae*, *Candida* spp., *Hansenula* spp. and *Pichia* spp.

[0089] The present invention includes a plant cell, plant or plant tissue comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of at least one fatty acid selected from the group consisting of a monounsaturated fatty acid and a polyun-

saturated fatty acid by the plant cell, plant or plant tissue. The polyunsaturated fatty acid may be, for example, dihomo- γ -linolenic acid (DGLA), eicosatetraenoic acid, adrenic acid and ω 3-docosapentaenoic acid. The invention also includes one or more plant oils or fatty acids expressed by the plant cell, plant or plant tissue. Additionally, the present invention encompasses a transgenic plant comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0090] Furthermore, the present invention includes a transgenic, non-human mammal whose genome comprises a DNA sequence encoding an elongase operably linked to a promoter. The DNA sequence may be represented by SEQ ID NO:7 (FIG. 72). The present invention also includes a fluid (e.g., milk) produced by the transgenic, non-human wherein the fluid comprises a detectable level of at least one elongase or products thereof such as, for example, DGLA, ω 6-docosapentaenoic acid, ADA and/or 20:4n-3 (see FIG. 1).

[0091] Additionally, the present invention includes a method for producing a polyunsaturated fatty acid comprising the steps of: a) isolating said nucleotide sequence represented by SEQ ID NO:7 (FIG. 72); b) constructing a vector comprising the isolated nucleotide sequence; c) introducing the vector into a host cell under time and conditions sufficient for expression of elongase enzyme encoded by the isolated nucleotide sequence; and d) exposing the expressed elongase enzyme to a "substrate" polyunsaturated fatty acid in order to convert the substrate to a "product" polyunsaturated fatty acid. The substrate polyunsaturated fatty acid may be selected from the group consisting of, for example, γ -linolenic acid (GLA), stearidonic acid (STA), arachidonic acid (AA) and EPA, and the product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3, ADA and ω 3-docosapentaenoic acid, respectively. The method may further comprise the step of exposing the product polyunsaturated fatty acid to at least one desaturase in order to convert the product polyunsaturated fatty acid to "another" polyunsaturated fatty acid. The product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3, and ADA. The another polyunsaturated fatty acid may be selected from the group consisting of, for example, AA, eicosapentaenoic acid (EPA), ω 6-docosapentaenoic acid, respectively, and the at least one desaturase is Δ 5-desaturase, with respect to production of AA or EPA, and Δ 4-desaturase, with respect to production of ω 6-docosapentaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a "final" polyunsaturated fatty acid. The final polyunsaturated fatty acid may be, for example, docosahexaenoic acid (DHA), ADA, ω 6-docosapentaenoic acid or ω 3-docosapentaenoic acid.

[0092] Also, the present invention includes a nutritional composition comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the "another" polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the

above-described method. The product polyunsaturated fatty acid may be selected from the group consisting of, for example, DGLA, 20:4n-3 and ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA, or ω 6-docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, adrenic acid, ω 6-docosapentaenoic acid or ω 3-docosapentaenoic acid.

[0093] The nutritional composition may be, for example, an infant formula, a dietary supplement or a dietary substitute and may be administered to a human or an animal and may be administered enterally or parenterally. The nutritional composition may further comprise at least one macro-nutrient selected from the group consisting of coconut oil, soy oil, canola oil, monoglycerides, diglycerides, triglycerides, glucose, edible lactose, electrodialysed whey, electrodialysed skim milk, milk whey, soy protein, protein hydrolysates, sunflower oil, safflower oil, corn oil, and flax oil. It may also comprise at least one vitamin selected from the group consisting of Vitamins A, C, D, E, and B complex and at least one mineral selected from the group consisting of calcium magnesium, zinc, manganese, sodium, potassium, phosphorus, copper, chloride, iodine, selenium and iron.

[0094] Additionally, the present invention encompasses a pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method and 2) a pharmaceutically acceptable carrier. The composition may be administered to a human or an animal. It may also further comprise at least one element selected from the group consisting of a vitamin, a mineral, a salt, a carbohydrate, an amino acid, a free fatty acid, a preservative, an excipient, an anti-histamine, a growth factor, an antibiotic, a diluent, a phospholipid, an antioxidant, and a phenolic compound. It may be administered enterally, parenterally, topically, rectally, intramuscularly, subcutaneously, intradermally, or by any other appropriate means.

[0095] The present invention also includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method. The product polyunsaturated fatty acid may be, for example, DGLA, 20:4n-3, and ADA. The another polyunsaturated fatty acid may be, for example, AA, EPA, or ω 6-docosapentaenoic acid. The final polyunsaturated fatty acid may be, for example, DHA, adrenic acid (ADA), ω 6-docosapentaenoic acid or ω 3-docosapentaenoic acid.

[0096] Moreover, the present invention also includes a cosmetic comprising a polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the above-described method, the another polyunsaturated fatty acid produced according to the above-described method, and the final polyunsaturated fatty acid produced according to the above-described method.

[0097] Additionally, the present invention includes a method of preventing or treating a condition caused by insufficient intake or production of polyunsaturated fatty acids comprising administering to the patient the above nutritional composition in an amount sufficient to effect prevention or treatment.

[0098] Also, the present invention encompasses an isolated nucleotide sequence or fragment thereof comprising or complementary to a nucleotide sequence encoding a polypeptide having elongase activity, wherein the amino acid sequence of the polypeptide has at least 50% identity to an amino acid sequence comprising SEQ ID NO:121. Additionally, the present invention includes an isolated nucleotide sequence or fragment thereof comprising or complementary to a nucleotide sequence having at least 50% nucleotide sequence identity to a nucleotide sequence comprising SEQ ID NO:119. The above nucleotide sequences may encode a functionally active elongase which utilizes a polyunsaturated fatty acid as a substrate. Furthermore, the sequences may be derived from the genus *Pavlova* sp.

[0099] The present invention also encompasses a purified protein encoded by the nucleotide sequences described above as well as a purified polypeptide which elongates polyunsaturated fatty acids and has at least about 30% amino acid identity to the amino acid sequence of the purified protein comprising SEQ ID NO:121.

[0100] Moreover, the present invention also includes a method of producing an elongase enzyme comprising the steps of: isolating a nucleotide sequence comprising SEQ ID NO:119; constructing a vector comprising: i) the isolated nucleotide sequence operably linked to ii) a promoter or other regulatory sequence; and introducing the vector into a host cell under time and conditions sufficient for expression of the elongase enzyme.

[0101] The host cell may be, for example, any one or more of the host cells referred to above in connection with the above cited sequences (e.g., SEQ ID NO:5, SEQ ID NO:7, etc.).

[0102] Furthermore, the present invention encompasses a vector comprising: a) a nucleotide sequence comprising SEQ ID NO:119 operably linked to b) a promoter or at least one regulatory sequence. The invention also includes a host cell comprising this vector. Once again, the host cell may be any one or more of the host cells described above in connection with the above-recited sequences (e.g., SEQ ID NO:5, SEQ ID NO:7, etc.).

[0103] Also, the present invention encompasses a plant cell, plant or plant tissue comprising, the vector above comprising SEQ ID NO:119, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid by the plant cell, plant or plant tissue. More specifically, the polyunsaturated fatty acid may be, for example, ADA or ω 3-docosapentaenoic acid. The invention also includes one or more plant oils or acids expressed by the plant cell, plant or plant tissue. The present invention also encompasses a transgenic plant comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

[0104] Additionally, the present invention includes a transgenic, non-human mammal whose genome comprises a

DNA sequence encoding an elongase, operably linked to a promoter or one or more other regulatory sequences, wherein the DNA sequence comprises SEQ ID NO:119. The invention also includes a fluid produced by the transgenic, non-human wherein the fluid comprises a detectable level of at least one elongase or products thereof.

[0105] Also, the present invention includes a method for producing a polyunsaturated fatty acid comprising the steps of: isolating a nucleotide sequence comprising SEQ ID NO:119; constructing a vector comprising the isolated nucleotide sequence; introducing the vector into a host cell under time and conditions sufficient for expression of an elongase enzyme encoded the isolated nucleotide sequence; and exposing the expressed elongase enzyme to a substrate polyunsaturated fatty acid in order to convert the substrate to a product polyunsaturated fatty acid. The substrate polyunsaturated fatty acid may be, for example, AA or EPA, and the product polyunsaturated fatty acid may be, for example, ADA or ω 3-docosapentaenoic acid (see FIG. 1). This method may further comprises the step of exposing the expressed elongase enzyme to at least one desaturase in order to convert said product polyunsaturated fatty acid to another polyunsaturated fatty acid. In this method, the product polyunsaturated fatty acid may be, for example, ADA or ω 3-docosapentaenoic acid, the another polyunsaturated fatty acid may be, for example, ω 6-docosapentaenoic acid, docohexaenoic acid or ω 3-docosapentaenoic acid. The at least one desaturase may be, for example, Δ 4-desaturase with respect to production of ω 6-docosapentaenoic acid or docohexaenoic acid, and Δ 19-desaturase with respect to production of ω 3-docosapentaenoic acid or docosahexaenoic acid. The method may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert the another polyunsaturated fatty acid to a final polyunsaturated fatty acid. The final polyunsaturated fatty acid may be, for example, docosahexaenoic acid.

[0106] It should also noted that the elongase of the present may be used in several of the pathways shown in FIG. 1, in which at least one desaturase and/or at least one "other" elongase has acted on a substrate or a product. For example, once DGLA has been converted to AA by a Δ 5-desaturase, the elongase of the present invention may then act on the AA in order to produce ADA. As another example, STA may be converted to ETA by an elongase and then to EPA by a Δ 5-desaturase. The EPA may then be converted to ω 3-docosapentaenoic acid by the elongase of the present invention (see FIG. 1). Thus, any pathway in which the elongase of the present invention is involved is encompassed by the present invention.

[0107] The present invention also includes a nutritional composition comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method or methods described above, the another polyunsaturated fatty acid produced according to the method described above, and the final polyunsaturated fatty acid produced according to the method described above. The product, "another" and final polyunsaturated fatty acids may be, for example, those described above in connection with SEQ ID

NO:121. This nutritional composition may be, for example, an infant formula, a dietary supplement and a dietary substitute.

[0108] The present invention also includes at least one pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the method or methods described above, the "another" polyunsaturated fatty acid produced according to the method described above, and the final polyunsaturated fatty acid produced according to the method described above and 2) a pharmaceutically acceptable carrier.

[0109] Also, the present invention includes an animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of the product polyunsaturated fatty acid produced according to the methods described above, the another polyunsaturated fatty acid produced according to the method described above and the final polyunsaturated fatty acid produced according to the method described above.

[0110] Additionally, the present invention includes a method of preventing or treating a condition caused by insufficient intake of polyunsaturated fatty acids comprising administering to the patient the nutritional composition above in an amount sufficient to effect the prevention or treatment.

[0111] All U.S. patents and publications referred to herein are hereby incorporated in their entirety by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0112] FIG. 1 represents various fatty acid biosynthesis pathways. The role of the elongase enzyme (elo) should be noted.

[0113] FIG. 2 represents the percent similarity and percent identity between the amino acid sequences of jojoba KCS (SEQ ID NO:8) and ELO2 (SEQ ID NO:9).

[0114] FIG. 3 represents the *S. cerevisiae* ELO2 sequence (SEQ ID NO:10) homologous to the jojoba KCS sequence (primer sequence underlined) of FIG. 2.

[0115] FIG. 4A shows the physical map of pRAE-2 containing the MAELO cDNA. FIG. 4B represents the physical map of the constitutive expression vector, pRAE-5, used for elongase enzyme production in yeast.

[0116] FIG. 5 represents a comparison of the nucleotide sequences of clones pRAE-5 (SEQ ID NO:11) and pRAE-6 (SEQ ID NO:12).

[0117] FIG. 6 illustrates the complete nucleotide sequence of *Mortierella alpina* elongase (MAELO) (SEQ ID NO:1).

[0118] FIG. 7 represents the amino acid sequence of the *Mortierella alpina* elongase translated from MAELO (see FIG. 6) (SEQ ID NO:13).

[0119] FIG. 8 represents an amino acid sequence alignment among 3 elongases: *S. cerevisiae* ELO2 (GNS1) (SEQ ID NO:14), *S. cerevisiae* ELO3 (SUR4) (SEQ ID NO:15) and the translated MAELO sequence as shown in FIG. 7.

[0120] FIG. 9 represents a comparison between the nucleotide sequence MAELO (SEQ ID NO:16) and the nucleotide sequence of ELO2 from *S. cerevisiae* (SEQ ID NO:17).

[0121] FIGS. 10A and 10B represents the PUFA elongase activity of MAELO expressed in baker's yeast.

[0122] FIG. 11 illustrates the PUFA elongase activity of MAELO when co-expressed with the $\Delta 5$ -desaturase cDNA from *M. alpina* to produce AA.

[0123] FIG. 12 compares the PUFA elongase activity of MAELO to the overexpression of ELO2 from *S. cerevisiae* in baker's yeast.

[0124] FIGS. 13 (SEQ ID NO:18 and SEQ ID NO:19), 14 (SEQ ID NO:20 and SEQ ID NO:21) and 15 (SEQ ID NO:22 and SEQ ID NO:23) represent three separate comparisons of amino acid sequences derived from *C. elegans* nucleotide sequences in the GenEMBL database with the translated MAELO.

[0125] FIG. 16 shows the comparison between amino acid translations of two different mammalian sequences in the GenEMBL database and the translated MAELO (SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26 and SEQ ID NO:27).

[0126] FIG. 17 shows the comparison of a translated DNA sequence (see published PCT application WO 88/07577) with the amino acid sequence derived from MAELO, which was detected during a database search (SEQ ID NO:28 and SEQ ID NO:29).

[0127] FIG. 18 shows the complete nucleotide sequence of the $\Delta 5$ -desaturase from *M. alpina* (SEQ ID NO:30).

[0128] FIG. 19 represents the initial GC-FAME analysis of MAD708 pool. The detection of a DGLA (C20:3n-6) peak should be noted.

[0129] FIG. 20 represents the PUFA elongase activity of the five MAD708 clones in yeast with GLA as substrate. All clones have apparent elongase activity.

[0130] FIG. 21 represents the DNA sequencing analysis of plasmid pRPB2. The analysis reveals an open reading frame of 957 bp in length.

[0131] FIG. 22 shows the complete nucleotide sequence of the *M. alpina* cDNA (SEQ ID NO:2), contained in the plasmid pRPB2, which is designated GLELO for its GLA elongase activity.

[0132] FIG. 23 represents the amino acid sequence of the *M. alpina* elongase (SEQ ID NO:31) translated from GLELO (see FIG. 22).

[0133] FIG. 24 illustrates the n-6 PUFA elongase activity in an induced culture of 334(pRPB2) when supplemented with GLA.

[0134] FIG. 25 represents the n-3 and n-6 PUFA elongase activity in an induced culture of 334(pRPB2) when supplemented with 25 μ M of other fatty acid substrates.

[0135] FIG. 26A illustrates the elongase activity of GLELO with GLA as a substrate when co-expressed with the *M. alpina* $\Delta 5$ -desaturase cDNA to produce AA. FIG. 26B illustrates the elongase activity of GLELO with STA as a substrate when co-expressed with the *M. alpina* $\Delta 5$ -desaturase cDNA to produce EPA.

[0136] FIG. 27 illustrates the comparison between the translated GLELO sequence (SEQ ID NO:32) (see FIG. 23) and the translated MAELO sequence (SEQ ID NO:33) (see FIG. 7).

- [0137] FIG. 28 represents a comparison of the amino acid sequence of 4 elongases: the translated amino acid sequence of GLELO (see FIG. 23), MAELO (see FIG. 7), *S. cerevisiae* ELO2 (GNS1), and *S. cerevisiae* ELO3 (SUR4). The histidine box is underlined.
- [0138] FIG. 29 represents an alignment between translated MAELO sequence (SEQ ID NO:34) and translated putative human homologue HS1 sequence (SEQ ID NO:35).
- [0139] FIG. 30 represents an alignment between the translated MAELO sequence (SEQ ID NO:36) and the translated putative human homologue HS2 sequence (SEQ ID NO:37).
- [0140] FIG. 31 shows an alignment between the translated MAELO sequence (SEQ ID NO:38) and the translated putative mouse homologue MM2 sequence (SEQ ID NO:39).
- [0141] FIG. 32 represents an alignment between the translated MAELO (SEQ ID NO:40) and the translated putative mouse homologue AI225632 sequence (SEQ ID NO:41).
- [0142] FIG. 33 illustrates an alignment between the translated GLELO sequence (SEQ ID NO:42) and the translated human homologue AI815960 sequence (SEQ ID NO:43).
- [0143] FIG. 34 shows an alignment between the translated GLELO sequence (SEQ ID NO:44) and the translated putative human homologue HS1 sequence (SEQ ID NO:45).
- [0144] FIG. 35 represents an alignment between the translated GLELO sequence (SEQ ID NO:46) and the translated putative human homologue sequence from AC004050 (SEQ ID NO:47).
- [0145] FIG. 36 illustrates an alignment between the translated GLELO sequence (SEQ ID NO:48) and the translated putative mouse homologue MM2 sequence (SEQ ID NO:49).
- [0146] FIG. 37 represents an alignment of the translated GLELO sequence (SEQ ID NO:50) and a translated putative mouse homologue AI225632 sequence (SEQ ID NO:51).
- [0147] FIG. 38 illustrates an alignment of the translated GLELO sequence (SEQ ID NO:52) and a translated putative mouse homologue U97107 (SEQ ID NO:53).
- [0148] FIG. 39 represents an alignment of the translated GLELO sequence (SEQ ID NO:54) and a translated putative *C. elegans* U68749 (F56H11.4) homologue sequence (SEQ ID NO:55).
- [0149] FIG. 40 shows an alignment between the translated MAELO sequence (SEQ ID NO:13) and a translated putative *C. elegans* U68749 (F56H11.4) homologue sequence (SEQ ID NO:56).
- [0150] FIG. 41 represents an alignment between the translated GLELO sequence (SEQ ID NO:57) and a translated putative *Drosophila melanogaster* homologue sequence, DM1 (SEQ ID NO:58).
- [0151] FIG. 42 illustrates an alignment between the translated MAELO sequence (SEQ ID NO:59) and a translated putative *Drosophila melanogaster* homologue sequence, DM1 (SEQ ID NO:60).
- [0152] FIG. 43 illustrates the complete nucleotide sequence of a human elongase HSELO1 (SEQ ID NO:3).
- [0153] FIG. 44 represents the deduced amino acid sequence of the human elongase HSELO1 (SEQ ID NO:61).
- [0154] FIG. 45 illustrates the elongase activity (PUFA and others) of an induced culture of 334(pRAE-58-A1) when supplemented with GLA or AA.
- [0155] FIG. 46 shows the complete nucleotide sequence of the *C. elegans* elongase CEELO (SEQ ID NO:4).
- [0156] FIG. 47 represents the deduced amino acid of *C. elegans* elongase CEELO (SEQ ID NO:62).
- [0157] FIG. 48 illustrates the PUFA elongase activity of an induced culture of 334(pRET-21) and 334(pRET-22) when supplemented with GLA and AA.
- [0158] FIG. 49 represents the complete nucleotide sequence of the putative human elongase gene HS3 (SEQ ID NO:63).
- [0159] FIG. 50 illustrates the deduced amino acid sequence of the putative human elongase enzyme HS3 (SEQ ID NO:64).
- [0160] FIG. 51 represents the elongase activity (PUFA and others) of HSELO1 expressed in baker's yeast when supplemented with GLA, AA, STA, EPA, OA, or ALA.
- [0161] FIG. 52 represents the elongase activity (PUFA and others) of HSELO1 expressed in baker's yeast when supplemented with 25 mM or 100 mM of GLA, AA, or EPA.
- [0162] FIGS. 53A, 53B, and 53C represent the elongase activity (PUFA and others) of HSELO1 expressed in baker's yeast when supplemented with PA, SA, ARA, BA, PTA, OA, EA, LA, GLA, DGLA, AA, ADA, ALA, STA, EPA, or DPA, or when no substrate is present.
- [0163] FIG. 54 represents the complete nucleotide sequence of mouse elongase MELO4 (SEQ ID NO:5).
- [0164] FIG. 55 represents the deduced amino acid sequence of the mouse elongase MELO4 (SEQ ID NO:65).
- [0165] FIG. 56 represents the PUFA elongase activity of MELO4 expressed in baker's yeast when supplemented with GLA, AA, ADA, STA, EPA, or DPA.
- [0166] FIGS. 57A, 57B, and 57C represent the PUFA elongase activity of MELO4 expressed in baker's yeast when supplemented with PA, SA, ARA, BA, PTA, OA, EA, LA, GLA, DGLA, AA, ADA, ALA, STA, EPA, or DPA, or when no substrate is present.
- [0167] FIG. 58 represents the complete nucleotide sequence of mouse elongase MELO7 (SEQ ID NO:6).
- [0168] FIG. 59 represents the deduced amino acid sequence of the mouse elongase MELO7 (SEQ ID NO:66).
- [0169] FIG. 60 represents the elongase activity (PUFA and others) of MELO7 expressed in baker's yeast when supplemented with GLA, AA, ADA, STA, EPA, or DPA.
- [0170] FIG. 61 shows the activity of the *C. elegans* elongase when expressed in yeast when no substrate is present and with addition of AA or GLA.
- [0171] FIG. 62 illustrates the PUFA elongase activity of an induced culture of 334(pRET22) when supplemented with 50 μ M of various substrates.

[0172] FIG. 63 represents the PUFA elongase activity with GLA (FIG. 63A) or STA (FIG. 63B) as a substrate when co-expressed with the *M. alpina* $\Delta 5$ -desaturase cDNA to produce AA or EPA, respectively.

[0173] FIG. 64 represents an amino acid sequence alignment among 4 elongases: HSELO1 (FIG. 44), MELO4 (FIG. 55), GLELO (FIG. 23), and CELO (FIG. 47).

[0174] FIG. 65 represents the consensus nucleotide sequence of the partial *Thraustochytrium aureum* (*T. aureum*) 7091 elongase gene (SEQ ID NO:67).

[0175] FIG. 66 represents the deduced amino acid sequence (SEQ ID NO:68) of the *T. aureum* 7091 elongase gene in FIG. 65.

[0176] FIG. 67 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:69), from plasmid pRAT-4-A1.

[0177] FIG. 68 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:70), from plasmid pRAT-4-A2.

[0178] FIG. 69 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:71), from plasmid pRAT-4-A3.

[0179] FIG. 70 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:72), from plasmid pRAT-4-A4.

[0180] FIG. 71 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:73), from plasmid pRAT-4-A6.

[0181] FIG. 72 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:7), from plasmid pRAT-4-A7.

[0182] FIG. 73 represents the complete nucleotide sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:74), from plasmid pRAT-4-D1.

[0183] FIG. 74 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:75), from plasmid pRAT-4-A1.

[0184] FIG. 75 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:76), from plasmid pRAT-4-A2.

[0185] FIG. 76 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:77), from plasmid pRAT-4-A3.

[0186] FIG. 77 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:78), from plasmid pRAT-4-A4.

[0187] FIG. 78 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:79), from plasmid pRAT-4-A6.

[0188] FIG. 79 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:80), from plasmid pRAT-4-A7.

[0189] FIG. 80 represents the deduced amino acid sequence of the *T. aureum* 7091 elongase TELO1 (SEQ ID NO:81), from plasmid pRAT-4-D1.

[0190] FIG. 81 represents the elongase activity of TELO1 expressed in baker's yeast when supplemented with GLA or EPA.

[0191] FIG. 82 represents an amino acid sequence alignment among the TELO1 elongases from the 7 plasmids: pRAT-4-A1 (FIG. 74), pRAT-4-A2 (FIG. 75), pRAT-4-A3 (FIG. 76), pRAT-4-A4 (FIG. 77), pRAT-4-A6 (FIG. 78), pRAT-4-A7 (FIG. 79), and pRAT-4-D1 (FIG. 80).

[0192] FIG. 83 represents an alignment between the translated TELO1 sequence (SEQ ID NO:82) and the translated HSELO1 sequence (SEQ ID NO:83).

[0193] FIG. 84 represents an alignment between the translated TELO1 sequence (SEQ ID NO:84) and the translated MELO4 sequence (SEQ ID NO:85).

[0194] FIG. 85 represents an alignment between the translated TELO1 sequence (SEQ ID NO:86) and the translated GLELO sequence (SEQ ID NO:87).

[0195] FIG. 86 represents an alignment between the translated TELO1 sequence (SEQ ID NO:88) and the translated CELO sequence (SEQ ID NO:89).

[0196] FIG. 87 represents the elongase activity of TELO1 expressed in baker's yeast when supplemented with GLA, AA, STA, EPA, or no substrate.

[0197] FIG. 88 represents the elongase activity of TELO1 expressed in baker's yeast when supplemented with LA, ALA, GLA, STA, DGLA, ETA, AA or EPA.

[0198] FIG. 89 represents the complete nucleotide sequence of the Pavlova sp. 459 elongase PELO1 (SEQ ID NO:117), from plasmid pRPL-6-1.

[0199] FIG. 90 represents the deduced amino acid sequence of the Pavlova sp. 459 elongase PELO1 (SEQ ID NO:118), from plasmid pRPL-6-1.

[0200] FIG. 91 represents the complete nucleotide sequence of the Pavlova sp. 459 elongase PELO1 (SEQ ID NO:119), from plasmid pRPL-6-B2.

[0201] FIG. 92 represents the complete nucleotide sequence of the Pavlova sp. 459 elongase PELO1 (SEQ ID NO:120), from plasmid pRPL-6-A3.

[0202] FIG. 93 represents the deduced amino acid sequence of the Pavlova sp. 459 elongase PELO1 (SEQ ID NO:121), from plasmid pRPL-6-B2.

[0203] FIG. 94 represents the deduced amino acid sequence of the Pavlova sp. 459 elongase PELO1 (SEQ ID NO:122), from plasmid pRPL-6-A3.

[0204] FIG. 95 represents an alignment between the translated PELO1 sequence (SEQ ID NO:118) and the translated MELO4 sequence (SEQ ID NO:65).

[0205] FIG. 96 represents an alignment between the translated PELO1 sequence (SEQ ID NO:119) and the translated GLELO sequence (SEQ ID NO:62).

[0206] FIG. 97 represents an alignment between the translated PELO1 sequence (SEQ ID NO:118) and the translated HSELO1 sequence (SEQ ID NO:61).

[0207] FIG. 98 represents an alignment between the translated PELO1 sequence (SEQ ID NO:118) and the translated TELO1 sequence (SEQ ID NO:65).

[0208] FIG. 99 represents the elongase activity of PELO1 expressed in baker's yeast when supplemented with GLA or EPA.

[0209] FIG. 100 represents an amino acid sequence alignment among the PELO1 elongases from the 3 plasmids: pRPL-6-B2, (FIG. 93), pRPL-6-A3 (FIG. 94), and pRPL-6-1 (FIG. 90) (see SEQ ID NO:121, SEQ ID NO:122 and SEQ ID NO:118, respectively).

[0210] FIG. 101 represents the elongase activity of PELO1 expressed in baker's yeast when supplemented with LA, GLA, DGLA, AA, ALA, STA, ETA, EPA, 20:0, or 20:1.

DETAILED DESCRIPTION OF THE INVENTION

[0211] The subject invention relates to nucleotide and encoded amino acid sequences of: elongase cDNAs derived from *Mortierella alpina*, an elongase cDNA derived from a human, an elongase cDNA derived from *C. elegans*, two elongase cDNAs derived from a mouse, an elongase cDNA derived from *Thraustochytrium aureum* and an elongase cDNA derived from Pavlova sp. 459. Furthermore, the subject invention also includes uses of the cDNAs and of the proteins encoded by the genes. For example, the genes and corresponding enzymes may be used in the production of polyunsaturated fatty acids and/or monounsaturated fatty acids such as, for example, DGLA, AA, ADA, EPA and/or DHA which may be added to pharmaceutical compositions, nutritional compositions, animal feeds, cosmetics, and to other valuable products.

The Elongase Genes and Enzymes Encoded Thereby

[0212] As noted above, an elongase enzyme encoded by an elongase cDNA is essential in the production of various polyunsaturated fatty acids, in particular, 20-24 carbon PUFAs. With respect to the present invention, the nucleotide sequence of the isolated *M. alpina* elongase cDNA (MAELO) is shown in FIG. 6, and the amino acid sequence of the corresponding purified protein or enzyme encoded by this nucleotide sequence is shown in FIG. 7. Additionally, the nucleotide sequence of the isolated GLA elongase cDNA (GLELO) is shown in FIG. 22, and the amino acid sequence of the corresponding purified protein or enzyme encoded by this nucleotide sequence is shown in FIG. 23. The nucleotide sequence of the isolated human sequence 1 (HSELO1) elongase is shown in FIG. 43, and the amino acid sequence of the corresponding purified protein or enzyme encoded by this sequence is shown in FIG. 44. Furthermore, the nucleotide sequence of the isolated *C. elegans* elongase cDNA (CEELO1) is shown in FIG. 46, and the amino acid sequence of the corresponding purified protein or enzyme encoded thereby is shown in FIG. 47. Additionally, the nucleotide sequence of the isolated mouse PUFA elongation enzyme (MELO4) is shown in FIG. 54, and the amino acid sequence of the corresponding purified protein or enzyme encoded thereby is shown in FIG. 55. Moreover, the nucleotide sequence of the second isolated mouse PUFA elongation enzyme (MELO7) is shown in FIG. 58, and the amino acid sequence of the corresponding purified protein or enzyme encoded thereby is shown in FIG. 59. Also, the nucleotide sequence of the isolated *T. aureum* elongase cDNA (TELO1) is shown in FIG. 72, and the amino acid

sequence of the corresponding purified protein or enzyme encoded thereby is shown in FIG. 79. Additionally, the nucleotide sequence of the isolated Pavlova sp. 459 elongase cDNA (PELO1) is shown in FIG. 91, and the amino acid sequence of the corresponding purified protein or enzyme encoded thereby is shown in FIG. 93.

[0213] As an example, several of the isolated elongases encoded by the cDNAs of the present invention elongate GLA to DGLA or elongate STA to 20:4n-3 or elongate AA to ADA. The production of arachidonic acid from DGLA, or EPA from 20:4n-3, is then catalyzed by, for example, a $\Delta 5$ -desaturase. Thus, neither AA (or EPA), nor DGLA (or 20:4n-3) nor ADA (or $\omega 3$ -docosapentaenoic acid), can be synthesized without at least one elongase cDNA and enzyme encoded thereby.

[0214] It should be noted that the present invention also encompasses nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 50%, preferably at least about 60%, and more preferably at least about 70% of the nucleotides in SEQ ID NO:1 (i.e., the nucleotide sequence of the MAELO cDNA described herein (see FIG. 6)). Furthermore, the present invention also includes nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 35%, preferably at least about 45%, and more preferably at least about 55% of the nucleotides in SEQ ID NO:2 (i.e., the nucleotide sequence of the GLELO cDNA described herein (see FIG. 22)). Additionally, the present invention also includes nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 35%, preferably at least about 45%, and more preferably at least about 55% of the nucleotides in SEQ ID NO:3 (i.e., the nucleotide sequence of the human sequence 1 (HSELO1) cDNA described herein (see FIG. 43)). In addition, the present invention also includes nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 35%, preferably at least about 45%, and more preferably at least about 55% of the nucleotides in SEQ ID NO:4 (i.e., the nucleotide sequence of the *C. elegans* cDNA, CEELO1, described herein (see FIG. 46)). Further, the present invention also includes nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 35%, preferably at least about 45%, and more preferably at least about 55% of the nucleotides in SEQ ID NO:5 or SEQ ID NO:6 (i.e., the nucleotide sequence of the mouse PUFA elongases MELO4 and MELO7, described herein (see FIGS. 54 and 58, respectively)). Also, the present invention encompasses nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 35%, preferably at least about 45%, and more preferably at least about 55% of the nucleotides in SEQ ID NO:7 (i.e., the nucleotide sequence of the *T. aureum* elongase gene TELO1, described herein (see FIG. 72)). Additionally, the present invention encompasses nucleotide sequences (and the corresponding encoded proteins) having sequences comprising or complementary to at least about 35%, preferably at least about 45%, and more preferably at least about 55% of the nucleotides in SEQ ID NO:119. It should be noted that the "most preferable" range, referred to in each instance, may be increased by increments of ten percent. For example, if "at least 55%" is the most preferable

range recited above, with respect to a particular sequence, such a range also naturally includes “at least 65% identity”, “at least 75% identity”, “at least 85% identity”, and “at least 95% identity”.

[0215] The sequence having the above-describe percent identity or complementary sequences may be derived from sources or sources other than from which the isolated, original sequences were derived (e.g., a eukaryote (e.g., *Thraustochytrium* spp. (e.g., *Thraustochytrium aureum* and *Thraustochytrium roseum*), *Schizochytrium* spp. (e.g., *Schizochytrium aggregatum*), *Conidiobolus* spp. (e.g., *Conidiobolus nanodes*), *Entomorphothora* spp. (e.g., *Entomorphothora exitalis*), *Saprolegnia* spp. (e.g., *Saprolegnia parasitica* and *Saprolegnia diclina*), *Leptomit* spp. (e.g., *Leptomit* *lacteus*), *Entomorphothora* spp., *Pythium* spp., *Porphyridium* spp. (e.g., *Porphyridium cruentum*), *Conidiobolus* spp., *Phytophthora* spp., *Penicillium* spp., *Coidosporium* spp., *Mucor* spp. (e.g., *Mucor circinelloides* and *Mucor javanicus*), *Fusarium* spp., *Aspergillus* spp., *Rhodotorula* spp., *Amphidinium carteri*, *Chaetoceros calcitrans*, *Cricosphaera carterae*, *Cryptocodinium cohnii*, *Cryptomonas ovata*, *Euglena gracilis*, *Gonyaulax polyedra*, *Gymnodinium* spp. (e.g. *Gymnodinium nelsoni*), *Gyrodinium cohnii*, *Isochrysis* spp. (e.g. *Isochrysis galbana*), Microalgae MK8805, *Nitzschia frustulum*, *Pavlova* spp. (e.g. *Pavlova lutheri*), *Phaeodactylum tricornutum*, *Prorocentrum cordatum*, *Rhodomonas lens*, and *Thalassiosira pseudonana*), a Psychrophilic bacteria (e.g., *Vibrio* spp. (e.g., *Vibrio marinus*)), a yeast (e.g., *Dipodascopsis uninucleata*), a non-mammalian organism such as a fly (e.g., *Drosophila melanogaster*) or *Caenorhabditis* spp. (e.g., *Caenorhabditis elegans*), or a mammal (e.g., a human or a mouse). Such sequences may also be derived from species within the genus *Mortierella*, other than the species *alpina*, for example, *Mortierella elongata*, *Mortierella exigua*, *Mortierella isabellina*, *Mortierella hygrophila*, and *Mortierella ramanniana*, va. *angulispora*.

[0216] Furthermore, the present invention also encompasses fragments and derivatives of the nucleotide sequences of the present invention (i.e., SEQ ID NO:1 (MAELO), SEQ ID NO:2 (GLELO), SEQ ID NO:3 (HSELO1), SEQ ID NO:4 (CEELO1)), SEQ ID NO:5 (MELO4), SEQ ID NO:6 (MELO7), SEQ ID NO:7 (TELO1), and SEQ ID NO:119 (PELO1)) as well as of the corresponding sequences derived from non-Mortierella or non-mammalian sources, etc., as described above, and having the above-described complementarity or correspondence/identity. Functional equivalents of the above-sequences (i.e., sequences having elongase activity) are also encompassed by the present invention.

[0217] For purposes of the present invention, “complementarity” is defined as the degree of relatedness between two DNA segments. It is determined by measuring the ability of the sense strand of one DNA segment to hybridize with the antisense strand of the other DNA segment, under appropriate conditions, to form a double helix. In the double helix, wherever adenine appears in one strand, thymine appears in the other strand. Similarly, wherever guanine is found in one strand, cytosine is found in the other. The greater the relatedness between the nucleotide sequences of two DNA segments, the greater the ability to form hybrid duplexes between the strands of two DNA segments.

[0218] The term “identity” refers to the relatedness of two sequences on a nucleotide-by-nucleotide basis over a particular comparison window or segment. Thus, identity is defined as the degree of sameness, correspondence or equivalence between the same strands (either sense or antisense) of two DNA segments (or two amino acid sequences). “Percentage of sequence identity” is calculated by comparing two optimally aligned sequences over a particular region, determining the number of positions at which the identical base or amino acid occurs in both sequences in order to yield the number of matched positions, dividing the number of such positions by the total number of positions in the segment being compared and multiplying the result by 100. Optimal alignment of sequences may be conducted by the algorithm of Smith & Waterman, Appl. Math. 2:482 (1981), by the algorithm of Needleman & Wunsch, J. Mol. Biol. 48:443 (1970), by the method of Pearson & Lipman, Proc. Natl. Acad. Sci. (USA) 85:2444 (1988) and by computer programs which implement the relevant algorithms (e.g., Clustal Macaw Pileup (<http://cmgm.stanford.edu/biochem218/11Multiple.pdf>); Higgins et al., CABIOS. 5L151-153 (1989)), FASTDB (Intelligenetics), BLAST (National Center for Biomedical Information; Altschul et al., Nucleic Acids Research 25:3389-3402 (1997)), PILEUP (Genetics Computer Group, Madison, Wis.) or GAP, BESTFIT, FASTA and TFASTA (Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, Madison, Wis.). (See U.S. Pat. No. 5,912,120.)

[0219] “Similarity” between two amino acid sequences is defined as the presence of a series of identical as well as conserved amino acid residues in both sequences. The higher the degree of similarity between two amino acid sequences, the higher the correspondence, sameness or equivalence of the two sequences. (“Identity between two amino acid sequences is defined as the presence of a series of exactly alike or invariant amino acid residues in both sequences.) The definitions of “complementarity”, “identity” and “similarity” are well known to those of ordinary skill in the art.

[0220] “Encoded by” refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 amino acids, more preferably at least 8 amino acids, and even more preferably at least 15 amino acids from a polypeptide encoded by the nucleic acid sequence.

[0221] The present invention also encompasses an isolated nucleotide sequence which encodes elongase activity and that is hybridizable, under moderately stringent conditions, to a nucleic acid having a nucleotide sequence comprising or complementary to the nucleotide sequences described above. A nucleic acid molecule is “hybridizable” to another nucleic acid molecule when a single-stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and ionic strength (see Sambrook et al., “Molecular Cloning: A Laboratory Manual, Second Edition (1989), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.)). The conditions of temperature and ionic strength determine the “stringency” of the hybridization. “Hybridization” requires that two nucleic acids contain complementary sequences. However, depending on the stringency of the hybridization, mismatches between bases may occur. The appropriate

stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation. Such variables are well known in the art. More specifically, the greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., supra). For hybridization with shorter nucleic acids, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook et al., supra).

[0222] As used herein, an “isolated nucleic acid fragment or sequence” is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA. (A “fragment” of a specified polynucleotide refers to a polynucleotide sequence which comprises a contiguous sequence of approximately at least about 6 nucleotides, preferably at least about 8 nucleotides, more preferably at least about 10 nucleotides, and even more preferably at least about 15 nucleotides, and most preferable at least about 25 nucleotides identical or complementary to a region of the specified nucleotide sequence.) Nucleotides (usually found in their 5'-monophosphate form) are referred to by their single letter designation as follows: “A” for adenylate or deoxyadenylate (for RNA or DNA, respectively), “C” for cytidylate or deoxycytidylate, “G” for guanylate or deoxyguanylate, “U” for uridylate, “T” for deoxythymidylate, “R” for purines (A or G), “Y” for pyrimidines (C or T), “K” for G or T, “H” for A or C or T, “I” for inosine, and “N” for any nucleotide.

[0223] The terms “fragment or subfragment that is functionally equivalent” and “functionally equivalent fragment or subfragment” are used interchangeably herein. These terms refer to a portion or subsequence of an isolated nucleic acid fragment in which the ability to alter gene expression or produce a certain phenotype is retained whether or not the fragment or subfragment encodes an active enzyme. For example, the fragment or subfragment can be used in the design of chimeric constructs to produce the desired phenotype in a transformed plant. Chimeric constructs can be designed for use in co-suppression or antisense by linking a nucleic acid fragment or subfragment thereof, whether or not it encodes an active enzyme, in the appropriate orientation relative to a plant promoter sequence.

[0224] The terms “homology”, “homologous”, “substantially similar” and “corresponding substantially” are used interchangeably herein. They refer to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate gene expression or produce a certain phenotype. These terms also refer to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially alter the functional properties of the resulting nucleic acid fragment relative to the initial, unmodified fragment. It is therefore understood, as those skilled in the art will appreciate, that the invention encompasses more than the specific exemplary sequences.

[0225] “Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence.

[0226] “Native gene” refers to a gene as found in nature with its own regulatory sequences. In contrast, “chimeric construct” refers to a combination of nucleic acid fragments that are not normally found together in nature. Accordingly, a chimeric construct may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that normally found in nature. (The term “isolated” means that the sequence is removed from its natural environment.)

[0227] A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric constructs. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

[0228] “Coding sequence” refers to a DNA sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include, but are not limited to, promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

[0229] “Promoter” refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Regulatory sequences (e.g., a promoter) can also be located within the transcribed portions of genes, and/or downstream of the transcribed sequences. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most host cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamoto and Goldberg, (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of some variation may have identical promoter activity.

[0230] An “intron” is an intervening sequence in a gene that does not encode a portion of the protein sequence. Thus, such sequences are transcribed into RNA but are then excised and are not translated. The term is also used for the

excised RNA sequences. An “exon” is a portion of the gene sequence that is transcribed and is found in the mature messenger RNA derived from the gene, but is not necessarily a part of the sequence that encodes the final gene product.

[0231] The “translation leader sequence” refers to a DNA sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner, R. and Foster, G. D. (1995) *Molecular Biotechnology* 3:225).

[0232] The “3' non-coding sequences” refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al., (1989) *Plant Cell* 1:671-680.

[0233] “RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. “Messenger RNA (mRNA)” refers to the RNA that is without introns and that can be translated into protein by the cell. “cDNA” refers to a DNA that is complementary to and synthesized from a mRNA template using the enzyme reverse transcriptase. The cDNA can be single-stranded or converted into the double-stranded form using the Klenow fragment of DNA polymerase I. “Sense” RNA refers to RNA transcript that includes the mRNA and can be translated into protein within a cell or in vitro. “Antisense RNA” refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (U.S. Pat. No. 5,107,065). The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. “Functional RNA” refers to antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes. The terms “complement” and “reverse complement” are used interchangeably herein with respect to mRNA transcripts, and are meant to define the antisense RNA of the message.

[0234] The term “endogenous RNA” refers to any RNA which is encoded by any nucleic acid sequence present in the genome of the host prior to transformation with the recombinant construct of the present invention, whether naturally-occurring or non-naturally occurring, i.e., introduced by recombinant means, mutagenesis, etc.

[0235] The term “non-naturally occurring” means artificial, not consistent with what is normally found in nature.

[0236] The term “operably linked” refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is regulated by the other.

For example, a promoter is operably linked with a coding sequence when it is capable of regulating the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in a sense or antisense orientation. In another example, the complementary RNA regions of the invention can be operably linked, either directly or indirectly, 5' to the target mRNA, or 3' to the target mRNA, or within the target mRNA, or a first complementary region is 5' and its complement is 3' to the target mRNA.

[0237] The term “expression”, as used herein, refers to the production of a functional end-product. Expression of a gene involves transcription of the gene and translation of the mRNA into a precursor or mature protein. “Antisense inhibition” refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. “Co-suppression” refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Pat. No. 5,231,020).

[0238] “Mature” protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or pro-peptides present in the primary translation product have been removed. “Precursor” protein refers to the primary product of translation of mRNA; i.e., with pre- and pro-peptides still present. Pre- and pro-peptides may be but are not limited to intracellular localization signals.

[0239] “Stable transformation” refers to the transfer of a nucleic acid fragment into a genome of a host organism, resulting in genetically stable inheritance. In contrast, “transient transformation” refers to the transfer of a nucleic acid fragment into the nucleus, or DNA-containing organelle, of a host organism resulting in gene expression without integration or stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. The preferred method of cell transformation of rice, corn and other monocots is the use of particle-accelerated or “gene gun” transformation technology (Klein et al., (1987) *Nature (London)* 327:70-73; U.S. Pat. No. 4,945,050), or an Agrobacterium-mediated method using an appropriate Ti plasmid containing the transgene (Ishida Y. et al., 1996, *Nature Biotech.* 14:745-750). The term “transformation” as used herein refers to both stable transformation and transient transformation.

[0240] Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter “Sambrook”).

[0241] The term “recombinant” refers to an artificial combination of two otherwise separated segments of sequence, e.g., by chemical synthesis or by the manipulation of isolated segments of nucleic acids by genetic engineering techniques.

[0242] “PCR” or “Polymerase Chain Reaction” is a technique for the synthesis of large quantities of specific DNA segments, consists of a series of repetitive cycles (Perkin Elmer Cetus Instruments, Norwalk, Conn.). Typically, the double stranded DNA is heat denatured, the two primers

complementary to the 3' boundaries of the target segment are annealed at low temperature and then extended at an intermediate temperature. One set of these three consecutive steps is referred to as a cycle.

[0243] Polymerase chain reaction ("PCR") is a powerful technique used to amplify DNA millions of fold, by repeated replication of a template, in a short period of time. (Mullis et al, *Cold Spring Harbor Symp. Quant. Biol.* 51:263-273 (1986); Erlich et al, European Patent Application 50,424; European Patent Application 84,796; European Patent Application 258,017, European Patent Application 237,362; Mullis, European Patent Application 201,184, Mullis et al U.S. Pat. No. 4,683,202; Erlich, U.S. Pat. No. 4,582,788; and Saiki et al, U.S. Pat. No. 4,683,194). The process utilizes sets of specific in vitro synthesized oligonucleotides to prime DNA synthesis. The design of the primers is dependent upon the sequences of DNA that are desired to be analyzed. The technique is carried out through many cycles (usually 20-50) of melting the template at high temperature, allowing the primers to anneal to complementary sequences within the template and then replicating the template with DNA polymerase.

[0244] The products of PCR reactions are analyzed by separation in agarose gels followed by ethidium bromide staining and visualization with UV transillumination. Alternatively, radioactive dNTPs can be added to the PCR in order to incorporate label into the products. In this case the products of PCR are visualized by exposure of the gel to x-ray film. The added advantage of radiolabeling PCR products is that the levels of individual amplification products can be quantitated.

[0245] The terms "recombinant construct", "expression construct" and "recombinant expression construct" are used interchangeably herein. These terms refer to a functional unit of genetic material that can be inserted into the genome of a cell using standard methodology well known to one skilled in the art. Such construct may be itself or may be used in conjunction with a vector. If a vector is used, then the choice of vector is dependent upon the method that will be used to transform host plants, as is well known to those skilled in the art. For example, a plasmid vector can be used. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells comprising any of the isolated nucleic acid fragments of the invention. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al., (1985) *EMBO J.* 4:2411-2418; De Almeida et al., (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

Production of the Elongase Enzyme

[0246] Once the gene encoding the elongase enzyme has been isolated, it may then be introduced into either a prokaryotic or eukaryotic host cell through the use of a vector or construct. The vector, for example, a bacteriophage, cosmid or plasmid, may comprise the nucleotide

sequence encoding the elongase enzyme, as well as any regulatory sequence (e.g., promoter) which is functional in the host cell and is able to elicit expression of the desaturase encoded by the nucleotide sequence. The regulatory sequence (e.g., promoter) is in operable association with or operably linked to the nucleotide sequence. (A regulatory sequence (e.g., promoter) is said to be "operably linked" with a coding sequence if the regulatory sequence affects transcription or expression of the coding sequence.) Suitable promoters include, for example, those from genes encoding alcohol dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, phosphoglucose isomerase, phosphoglycerate kinase, acid phosphatase, T7, TPI, lactase, metallothionein, cytomegalovirus immediate early, whey acidic protein, glucamylase, and promoters activated in the presence of galactose, for example, GAL1 and GAL10. Additionally, nucleotide sequences which encode other proteins, oligosaccharides, lipids, etc., may also be included within the vector as well as other non-promoter regulatory sequences such as a polyadenylation signal (e.g., the poly-A signal of SV-40T-antigen, ovalalbumin or bovine growth hormone). The choice of sequences present in the construct is dependent upon the desired expression products as well as the nature of the host cell.

[0247] As noted above, once the vector has been constructed, it may then be introduced into the host cell of choice by methods known to those of ordinary skill in the art including, for example, transfection, transformation and electroporation (see *Molecular Cloning: A Laboratory Manual*, 2nd ed., Vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press (1989)). The host cell is then cultured under suitable conditions permitting expression of the PUFA which is then recovered and purified.

[0248] It should also be noted that one may design a unique triglyceride or oil if one utilizes one construct or vector comprising the nucleotide sequences of two or more cDNAs (e.g., MAELO, GLELO, HSELO1, CELO1, TELO1, and/or PELO1). This vector may then be introduced into one host cell. Alternatively, each of the sequences may be introduced into a separate vector. These vectors may then be introduced into two host cells, respectively, or into one host cell.

[0249] Examples of suitable prokaryotic host cells include, for example, bacteria such as *Escherichia coli*, *Bacillus subtilis* as well as cyanobacteria such as *Spirulina* spp. (i.e., blue-green algae). Examples of suitable eukaryotic host cells include, for example, mammalian cells, plant cells, yeast cells such as *Saccharomyces* spp., *Lipomyces* spp., *Candida* spp. such as *Yarrowia* (*Candida*) spp., *Kluyveromyces* spp., *Pichia* spp., *Trichoderma* spp. or *Hansenula* spp., or fungal cells such as filamentous fungal cells, for example, *Aspergillus*, *Neurospora* and *Penicillium*. Preferably, *Saccharomyces cerevisiae* (baker's yeast) cells are utilized.

[0250] Expression in a host cell can be accomplished in a transient or stable fashion. Transient expression can occur from introduced constructs which contain expression signals functional in the host cell, but which constructs do not replicate and rarely integrate in the host cell, or where the host cell is not proliferating. Transient expression also can be accomplished by inducing the activity of a regulatable promoter operably linked to the gene of interest, although

such inducible systems frequently exhibit a low basal level of expression. Stable expression can be achieved by introduction of a construct that can integrate into the host genome or that autonomously replicates in the host cell. Stable expression of the gene of interest can be selected for through the use of a selectable marker located on or transfected with the expression construct, followed by selection for cells expressing the marker. When stable expression results from integration, the site of the construct's integration can occur randomly within the host genome or can be targeted through the use of constructs containing regions of homology with the host genome sufficient to target recombination with the host locus. Where constructs are targeted to an endogenous locus, all or some of the transcriptional and translational regulatory regions can be provided by the endogenous locus.

[0251] A transgenic mammal may also be used in order to express the enzyme of interest (i.e., the elongase) encoded by one or both of the above-described nucleotide sequences. More specifically, once the above-described construct is created, it may be inserted into the pronucleus of an embryo. The embryo may then be implanted into a recipient female. Alternatively, a nuclear transfer method could also be utilized (Schnieke et al., *Science* 278:2130-2133 (1997)). Gestation and birth are then permitted to occur (see, e.g., U.S. Pat. No. 5,750,176 and U.S. Pat. No. 5,700,671). Milk, tissue or other fluid samples from the offspring should then contain altered levels of PUFAs, as compared to the levels normally found in the non-transgenic animal. Subsequent generations may be monitored for production of the altered or enhanced levels of PUFAs and thus incorporation of the gene or genes encoding the elongase enzyme into their genomes. The mammal utilized as the host may be selected from the group consisting of, for example, a mouse, a rat, a rabbit, a pig, a goat, a sheep, a horse and a cow. However, any mammal may be used provided it has the ability to incorporate DNA encoding the enzyme of interest into its genome.

[0252] For expression of an elongase polypeptide, functional transcriptional and translational initiation and termination regions are operably linked to the DNA encoding the elongase polypeptide. Transcriptional and translational initiation and termination regions are derived from a variety of nonexclusive sources, including the DNA to be expressed, genes known or suspected to be capable of expression in the desired system, expression vectors, chemical synthesis, or from an endogenous locus in a host cell. Expression in a plant tissue and/or plant part presents certain efficiencies, particularly where the tissue or part is one which is harvested early, such as seed, leaves, fruits, flowers, roots, etc. Expression can be targeted to that location with the plant by utilizing specific regulatory sequence such as those of U.S. Pat. Nos. 5,463,174, 4,943,674, 5,106,739, 5,175,095, 5,420,034, 5,188,958, and 5,589,379. Alternatively, the expressed protein can be an enzyme which produces a product which may be incorporated, either directly or upon further modifications, into a fluid fraction from the host plant. Expression of an elongase gene or genes, or antisense elongase transcripts, can alter the levels of specific PUFAs, or derivatives thereof, found in plant parts and/or plant tissues. The elongase polypeptide coding region may be expressed either by itself or with other genes, in order to produce tissues and/or plant parts containing higher proportions of desired PUFAs or in which the PUFA composition more closely resembles that of human breast milk (Prieto et

al., PCT publication WO 95/24494). The termination region may be derived from the 3' region of the gene from which the initiation region was obtained or from a different gene. A large number of termination regions are known to and have been found to be satisfactory in a variety of hosts from the same and different genera and species. The termination region usually is selected as a matter of convenience rather than because of any particular property.

[0253] As noted above, a plant (e.g., *Glycine max* (soybean) or *Brassica napus* (canola)), plant cell, plant tissue, corn, potato, sunflower, safflower or flax may also be utilized as a host or host cell, respectively, for expression of the elongase enzyme(s) which may, in turn, be utilized in the production of polyunsaturated fatty acids. More specifically, desired PUFAs can be expressed in seed. Methods of isolating seed oils are known in the art. Thus, in addition to providing a source for PUFAs, seed oil components may be manipulated through the expression of the elongase genes, as well as perhaps desaturase genes, in order to provide seed oils that can be added to nutritional compositions, pharmaceutical compositions, animal feeds and cosmetics. Once again, a vector which comprises a DNA sequence encoding the elongase operably linked to a promoter, will be introduced into the plant tissue or plant for a time and under conditions sufficient for expression of the elongase gene. The vector may also comprise one or more genes which encode other enzymes, for example, $\Delta 4$ -desaturase, $\Delta 5$ -desaturase, $\Delta 6$ -desaturase, $\Delta 8$ -desaturase, $\Delta 9$ -desaturase, $\Delta 10$ -desaturase, $\Delta 12$ -desaturase, $\Delta 13$ -desaturase, $\Delta 15$ -desaturase, $\Delta 17$ -desaturase and/or $\Delta 19$ -desaturase. The plant tissue or plant may produce the relevant substrate (e.g., DGLA, GLA, STA, AA, ADA, EPA, 20:4n-3, etc.) upon which the enzymes act or a vector encoding enzymes which produce such substrates may be introduced into the plant tissue, plant cell, plant, or host cell of interest. In addition, substrate may be sprayed on plant tissues expressing the appropriate enzymes. Using these various techniques, one may produce PUFAs (e.g., n-6 unsaturated fatty acids such as DGLA, AA or ADA, or n-3 fatty acids such as EPA or DHA) by use of a plant cell, plant tissue, plant, or host cell of interest. It should also be noted that the invention also encompasses a transgenic plant comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in, for example, the seeds of the transgenic plant.

[0254] The substrates which may be produced by the host cell either naturally or transgenically, as well as the enzymes which may be encoded by DNA sequences present in the vector, which is subsequently introduced into the host cell, are shown in FIG. 1.

[0255] In view of the above, the present invention also encompasses a method of producing one of the elongase enzymes described above comprising the steps of: 1) isolating the desired nucleotide sequence of the elongase cDNA; 2) constructing a vector comprising said nucleotide sequence; and 3) introducing said vector into a host cell under time and conditions sufficient for the production of the elongase enzyme.

[0256] The present invention also encompasses a method of producing polyunsaturated fatty acids comprising exposing an acid to the elongase(s) produced as above such that the elongase converts the acid to a polyunsaturated fatty

acid. For example, when GLA is exposed to elongase, it is converted to DGLA. DGLA may then be exposed to $\Delta 5$ -desaturase which converts the DGLA to AA. The AA may then be converted to EPA by use of $\Delta 17$ -desaturase which may be, in turn, converted to DHA by use of elongase and a $\Delta 4$ -desaturase. Alternatively, elongase may be utilized to convert $18:4n-3$ to $20:4n-3$ which may be exposed to $\Delta 5$ -desaturase and converted to EPA. Elongase may also be used to convert $18:3n-3$ to $20:3n-3$, which may be, in turn, converted to $20:4n-3$ by a $\Delta 8$ -desaturase. Thus, elongase may be used in the production of polyunsaturated fatty acids which may be used, in turn, for particular beneficial purposes. (See FIG. 1 for an illustration of the many critical roles the elongase enzyme plays in several biosynthetic pathways.)

Uses of the Elongase Gene and Enzyme Encoded Thereby

[0257] As noted above, the isolated elongase cDNAs and the corresponding elongase enzymes (or purified polypeptides) encoded thereby have many uses. For example, each cDNA and corresponding enzyme may be used indirectly or directly in the production of polyunsaturated fatty acids, for example, DGLA, AA, ADA, $20:4n-3$ or EPA. ("Directly" is meant to encompass the situation where the enzyme directly converts the acid to another acid, the latter of which is utilized in a composition (e.g., the conversion of GLA to DGLA)). "Indirectly" is meant to encompass the situation where a fatty acid is converted to another fatty acid (i.e., a pathway intermediate) by elongase (e.g., GLA to DGLA) and then the latter fatty acid is converted to another fatty acid by use of a non-elongase enzyme (e.g., DGLA to AA by $\Delta 5$ -desaturase)). These polyunsaturated fatty acids (i.e., those produced either directly or indirectly by activity of the elongase enzyme) may be added to, for example, nutritional compositions, pharmaceutical compositions, cosmetics, and animal feeds, all of which are encompassed by the present invention. These uses are described, in detail, below.

Nutritional Compositions

[0258] The present invention includes nutritional compositions. Such compositions, for purposes of the present invention, include any food or preparation for human consumption including for enteral or parenteral consumption, which when taken into the body (a) serve to nourish or build up tissues or supply energy and/or (b) maintain, restore or support adequate nutritional status or metabolic function.

[0259] The nutritional composition of the present invention comprises at least one oil or acid produced by use of at least one elongase enzyme, produced using the respective elongase gene, and may either be in a solid or liquid form. Additionally, the composition may include edible macronutrients, vitamins and minerals in amounts desired for a particular use. The amount of such ingredients will vary depending on whether the composition is intended for use with normal, healthy infants, children or adults having specialized needs such as those which accompany certain metabolic conditions (e.g., metabolic disorders).

[0260] Examples of macronutrients which may be added to the composition include but are not limited to edible fats, carbohydrates and proteins. Examples of such edible fats include but are not limited to coconut oil, soy oil, and mono-

and diglycerides. Examples of such carbohydrates include but are not limited to glucose, edible lactose and hydrolyzed starch. Additionally, examples of proteins which may be utilized in the nutritional composition of the invention include but are not limited to soy proteins, electro dialysed whey, electro dialysed skim milk, milk whey, or the hydrolysates of these proteins.

[0261] With respect to vitamins and minerals, the following may be added to the nutritional compositions of the present invention: calcium, phosphorus, potassium, sodium, chloride, magnesium, manganese, iron, copper, zinc, selenium, iodine, and Vitamins A, E, D, C, and the B complex. Other such vitamins and minerals may also be added.

[0262] The components utilized in the nutritional compositions of the present invention will be of semi-purified or purified origin. By semi-purified or purified is meant a material which has been prepared by purification of a natural material or by synthesis.

[0263] Examples of nutritional compositions of the present invention include but are not limited to infant formulas, dietary supplements, dietary substitutes, and rehydration compositions. Nutritional compositions of particular interest include but are not limited to those utilized for enteral and parenteral supplementation for infants, specialist infant formulae, supplements for the elderly, and supplements for those with gastrointestinal difficulties and/or malabsorption.

[0264] The nutritional composition of the present invention may also be added to food even when supplementation of the diet is not required. For example, the composition may be added to food of any type including but not limited to margarines, modified butters, cheeses, milk, yogurt, chocolate, candy, snacks, salad oils, cooking oils, cooking fats, meats, fish and beverages.

[0265] In a preferred embodiment of the present invention, the nutritional composition is an enteral nutritional product, more preferably, an adult or pediatric enteral nutritional product. This composition may be administered to adults or children experiencing stress or having specialized needs due to chronic or acute disease states. The composition may comprise, in addition to polyunsaturated fatty acids produced in accordance with the present invention, macronutrients, vitamins and minerals as described above. The macronutrients may be present in amounts equivalent to those present in human milk or on an energy basis, i.e., on a per calorie basis.

[0266] Methods for formulating liquid or solid enteral and parenteral nutritional formulas are well known in the art. (See also the Examples below.)

[0267] The enteral formula, for example, may be sterilized and subsequently utilized on a ready-to-feed (RTF) basis or stored in a concentrated liquid or powder. The powder can be prepared by spray drying the formula prepared as indicated above, and reconstituting it by rehydrating the concentrate. Adult and pediatric nutritional formulas are well known in the art and are commercially available (e.g., Similac®, Ensure®, Jevity® and Alimentum® from Ross Products Division, Abbott Laboratories, Columbus, Ohio). An oil or fatty acid produced in accordance with the present invention may be added to any of these formulas.

[0268] The energy density of the nutritional compositions of the present invention, when in liquid form, may range from about 0.6 Kcal to about 3 Kcal per ml. When in solid or powdered form, the nutritional supplements may contain from about 1.2 to more than 9 Kcals per gram, preferably about 3 to 7 Kcals per gm. In general, the osmolality of a liquid product should be less than 700 mOsm and, more preferably, less than 660 mOsm.

[0269] The nutritional formula may include macronutrients, vitamins, and minerals, as noted above, in addition to the PUFAs produced in accordance with the present invention. The presence of these additional components helps the individual ingest the minimum daily requirements of these elements. In addition to the provision of PUFAs, it may also be desirable to add zinc, copper, folic acid and antioxidants to the composition. It is believed that these substance boost a stressed immune system and will therefore provide further benefits to the individual receiving the composition. A pharmaceutical composition may also be supplemented with these elements.

[0270] In a more preferred embodiment, the nutritional composition comprises, in addition to antioxidants and at least one PUFA, a source of carbohydrate wherein at least 5 weight % of the carbohydrate is indigestible oligosaccharide. In a more preferred embodiment, the nutritional composition additionally comprises protein, taurine, and carnitine.

[0271] As noted above, the PUFAs produced in accordance with the present invention, or derivatives thereof, may be added to a dietary substitute or supplement, particularly an infant formula, for patients undergoing intravenous feeding or for preventing or treating malnutrition or other conditions or disease states. As background, it should be noted that human breast milk has a fatty acid profile comprising from about 0.15% to about 0.36% as DHA, from about 0.03% to about 0.13% as EPA, from about 0.30% to about 0.88% as AA, from about 0.22% to about 0.67% as DGLA, and from about 0.27% to about 1.04% as GLA. Thus, fatty acids such as DGLA, AA, EPA and/or docosahexaenoic acid (DHA), produced in accordance with the present invention, can be used to alter, for example, the composition of infant formulas in order to better replicate the PUFA content of human breast milk or to alter the presence of PUFAs normally found in a non-human mammal's milk. In particular, a composition for use in a pharmacologic or food supplement, particularly a breast milk substitute or supplement, will preferably comprise one or more of AA, DGLA and GLA. More preferably, the oil blend will comprise from about 0.3 to 30% AA, from about 0.2 to 30% DGLA, and/or from about 0.2 to about 30% GLA.

[0272] Parenteral nutritional compositions comprising from about 2 to about 30 weight percent fatty acids calculated as triglycerides are encompassed by the present invention. The preferred composition has about 1 to about 25 weight percent of the total PUFA composition as GLA (U.S. Pat. No. 5,196,198). Other vitamins, particularly fat-soluble vitamins such as vitamin A, D, E and L-carnitine can optionally be included. When desired, a preservative such as alpha-tocopherol may be added in an amount of about 0.1% by weight.

[0273] In addition, the ratios of AA, DGLA and GLA can be adapted for a particular given end use. When formulated

as a breast milk supplement or substitute, a composition which comprises one or more of AA, DGLA and GLA will be provided in a ratio of about 1:19:30 to about 6:1:0.2, respectively. For example, the breast milk of animals can vary in ratios of AA:DGLA:GLA ranging from 1:19:30 to 6:1:0.2, which includes intermediate ratios which are preferably about 1:1:1, 1:2:1, 1:1:4. When produced together in a host cell, adjusting the rate and percent of conversion of a precursor substrate such as GLA and DGLA to AA can be used to precisely control the PUFA ratios. For example, a 5% to 10% conversion rate of DGLA to AA can be used to produce an AA to DGLA ratio of about 1:19, whereas a conversion rate of about 75% to 80% can be used to produce an AA to DGLA ratio of about 6:1. Therefore, whether in a cell culture system or in a host animal, regulating the timing, extent and specificity of elongase expression, as well as the expression of other desaturases, can be used to modulate PUFA levels and ratios. The PUFAs/acids produced in accordance with the present invention (e.g., AA and DGLA) may then be combined with other PUFAs/acids (e.g., GLA) in the desired concentrations and ratios.

[0274] Additionally, PUFA produced in accordance with the present invention or host cells containing them may also be used as animal food supplements to alter an animal's tissue or milk fatty acid composition to one more desirable for human or animal consumption.

Pharmaceutical Compositions

[0275] The present invention also encompasses a pharmaceutical composition comprising one or more of the fatty acids and/or resulting oils produced using at least one of the elongase cDNAs in accordance with the methods described herein. More specifically, such a pharmaceutical composition may comprise one or more of the acids and/or oils as well as a standard, well-known, non-toxic pharmaceutically acceptable carrier, adjuvant or vehicle such as, for example, phosphate buffered saline, water, ethanol, polyols, vegetable oils, a wetting agent or an emulsion such as a water/oil emulsion. The composition may be in either a liquid or solid form. For example, the composition may be in the form of a tablet, capsule, ingestible liquid or powder, injectible, or topical ointment or cream. Proper fluidity can be maintained, for example, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride and the like. Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening agents, flavoring agents and perfuming agents.

[0276] Suspensions, in addition to the active compounds, may comprise suspending agents such as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth or mixtures of these substances.

[0277] Solid dosage forms such as tablets and capsules can be prepared using techniques well known in the art. For example, PUFAs produced in accordance with the present invention can be tableted with conventional tablet bases such as lactose, sucrose, and cornstarch in combination with binders such as acacia, cornstarch or gelatin, disintegrating agents such as potato starch or alginic acid, and a lubricant

such as stearic acid or magnesium stearate. Capsules can be prepared by incorporating these excipients into a gelatin capsule along with antioxidants and the relevant PUFA(s). The antioxidant and PUFA components should fit within the guidelines presented above.

[0278] For intravenous administration, the PUFAs produced in accordance with the present invention or derivatives thereof may be incorporated into commercial formulations such as Intralipids™. The typical normal adult plasma fatty acid profile comprises 6.64 to 9.46% of AA, 1.45 to 3.11% of DGLA, and 0.02 to 0.08% of GLA. These PUFAs or their metabolic precursors can be administered alone or in combination with other PUFAs in order to achieve a normal fatty acid profile in a patient. Where desired, the individual components of the formulations may be provided individually, in kit form, for single or multiple use. A typical dosage of a particular fatty acid is from 0.1 mg to 20 g (up to 100 g) daily and is preferably from 10 mg to 1, 2, 5 or 10 g daily.

[0279] Possible routes of administration of the pharmaceutical compositions of the present invention include, for example, enteral (e.g., oral and rectal) and parenteral. For example, a liquid preparation may be administered, for example, orally or rectally. Additionally, a homogenous mixture can be completely dispersed in water, admixed under sterile conditions with physiologically acceptable diluents, preservatives, buffers or propellants in order to form a spray or inhalant.

[0280] The route of administration will, of course, depend upon the desired effect. For example, if the composition is being utilized to treat rough, dry, or aging skin, to treat injured or burned skin, or to treat skin or hair affected by a disease or condition, it may perhaps be applied topically.

[0281] The dosage of the composition to be administered to the patient may be determined by one of ordinary skill in the art and depends upon various factors such as weight of the patient, age of the patient, immune status of the patient, etc.

[0282] With respect to form, the composition may be, for example, a solution, a dispersion, a suspension, an emulsion or a sterile powder which is then reconstituted.

[0283] The present invention also includes the treatment of various disorders by use of the pharmaceutical and/or nutritional compositions described herein. In particular, the compositions of the present invention may be used to treat restenosis after angioplasty. Furthermore, symptoms of inflammation, rheumatoid arthritis, asthma and psoriasis may also be treated with the compositions of the invention. Evidence also indicates that PUFAs may be involved in calcium metabolism; thus, the compositions of the present invention may, perhaps, be utilized in the treatment or prevention of osteoporosis and of kidney or urinary tract stones.

[0284] Additionally, the compositions of the present invention may also be used in the treatment of cancer. Malignant cells have been shown to have altered fatty acid compositions. Addition of fatty acids has been shown to slow their growth, cause cell death and increase their susceptibility to chemotherapeutic agents. Moreover, the compositions of the present invention may also be useful for treating cachexia associated with cancer.

[0285] The compositions of the present invention may also be used to treat diabetes (see U.S. Pat. No. 4,826,877 and Horrobin et al., *Am. J. Clin. Nutr.* Vol. 57 (Suppl.) 732S-737S). Altered fatty acid metabolism and composition have been demonstrated in diabetic animals.

[0286] Furthermore, the compositions of the present invention, comprising PUFAs produced either directly or indirectly through the use of the elongase enzyme(s), may also be used in the treatment of eczema, in the reduction of blood pressure, and in the improvement of mathematics examination scores. Additionally, the compositions of the present invention may be used in inhibition of platelet aggregation, induction of vasodilation, reduction in cholesterol levels, inhibition of proliferation of vessel wall smooth muscle and fibrous tissue (Brenner et al., *Adv. Exp. Med. Biol.* Vol. 83, p.85-101, 1976), reduction or prevention of gastrointestinal bleeding and other side effects of non-steroidal anti-inflammatory drugs (see U.S. Pat. No. 4,666,701), prevention or treatment of endometriosis and premenstrual syndrome (see U.S. Pat. No. 4,758,592), and treatment of myalgic encephalomyelitis and chronic fatigue after viral infections (see U.S. Pat. No. 5,116,871).

[0287] Further uses of the compositions of the present invention include use in the treatment of AIDS, multiple sclerosis, and inflammatory skin disorders, as well as for maintenance of general health.

[0288] Additionally, the composition of the present invention may be utilized for cosmetic purposes. It may be added to pre-existing cosmetic compositions such that a mixture is formed or may be used as a sole composition.

Veterinary Applications

[0289] It should be noted that the above-described pharmaceutical and nutritional compositions may be utilized in connection with animals (i.e., domestic or non-domestic), as well as humans, as animals experience many of the same needs and conditions as humans. For example, the oil or acids of the present invention may be utilized in animal feed supplements, animal feed substitutes, animal vitamins or in animal topical ointments.

[0290] The present invention may be illustrated by the use of the following non-limiting examples:

EXAMPLE I

Determination of Codon Usage in *Mortierella alpina*

[0291] The 5' end of 1000 random cDNA clones were sequenced from *Mortierella alpina* cDNA library. The sequences were translated in six reading frames using GCG (Genetics Computer Group (Madison, Wis.)) with the FastA algorithm (Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444-2448 (1988)) to search for similarity between a query sequence and a group of sequences of the same type (nucleic acid or protein), specifically with the Swissprot database (GeneBio, Geneva, Switzerland). Many of the clones were identified as a putative housekeeping gene based on protein sequence homology to known genes. Twenty-one *M. alpina* cDNA sequences which matched with known, housekeeping genes in the database were selected (see Table 1 below). *M. alpina* codon bias table (see

Table 2) generated based on these 21 sequences as well as the full length *M. alpina* Δ5-(see FIG. 18), Δ6-, and Δ12-desaturase sequences. Since the FastA alignment between the putative protein coded by the *M. alpina* cDNA sequence and the known protein sequence was weak in some areas, only the codons from areas of strong homology were used.

TABLE 1

Clone #	Match	# of bp	# of a
193	Elongation factor 1-alpha	426	142
143	60S ribosomal protein L17	417	139
235	Actin I	360	120
299	40S ribosomal protein YS11	387	129
390	Ras-related protein rab-1a	342	114
65	40S ribosomal protein RP10	366	122
289	Ubiquitin-conjugating enzyme E2-16 KD	294	98
151	Ubiquinol-cytochrome C reductase	375	125
80	Initiation factor 5A-2	183	61
33	60S ribosomal protein L15	252	84
132	60S ribosomal protein L3-2	300	100
198	Histone H3	285	95
286	6-phosphogluconate dehydrogenase, decarboxylating	363	121
283	40S ribosomal protein S22	261	87
127	Elongation factor 2	231	77
197	Actin, gamma	252	84
496	40S ribosomal protein S16	270	90
336	Histone H4	219	73
262	Ubiquitin	228	76
188	Guanine nucleotide-binding protein beta subunit-like protein	213	71
81	Ubiquitin	228	76
21	TOTAL	6252	2084

[0292]

TABLE 2

Amino acid	Codon Bias	% used
Ala	GCC	63%
Arg	CGC	50%
Asn	AAC	97%
Asp	GAC	65%
Cys	TGC	87%
Gln	CAG	78%
Glu	GAG	85%
Gly	GGT	47%
His	CAC	91%
Ile	ATC	72%
Leu	CTC	49%
Lys	AAG	96%
Met	ATG	100%
Phe	TTC	78%
Pro	CCC	68%
Ser	TCC	46%
Thr	ACC	78%
Trp	TGG	100%
Tyr	TAC	95%
Val	GTC	72%
Stop	TAA	50%

EXAMPLE II

Cloning of a Full-length Elongase-like cDNA from *M. alpina*

[0293] The β-ketoacyl-coenzyme A synthase (KCS) from *jojoba* and the *Saccharomyces cerevisiae* elongase (ELO2) were aligned to determine an area of amino acid homology

(see FIG. 2). The codon bias was applied to the area of sequence corresponding to the homologous amino acids between the two elongases, and primers were designed based on this biased sequence (see FIG. 3). The cDNA was excised from the M11 *M. alpina* cDNA library (Knutzon et al., *J. Biol. Chem.* 273:29360-29366 (1998)), which contains approximately 6×10⁵ clones with an average insert size of 1.1 Kb. The excised cDNA was amplified with internal primer RO339 (5'-TTG GAG AGG AGG AAG CGA CCA CCG AAG ATG ATG-3') and a vector forward primer RO317 (5'-CAC ACAGGAAACAGC TAT GAC CAT GAT TAC G-3'). Polymerase Chain Reaction (PCR) was carried out in a 100 μl volume containing: 300 ng of excised *M. alpina* cDNA library, 50 pmole each primer, 10 μl of 10×buffer, 1 μl 10 mM PCR Nucleotide Mix (Boehringer Mannheim Corp., Indianapolis, Ind.) and 1.0 U of Taq Polymerase. Thermocycler conditions in Perkin Elmer 9600 (Norwalk, Conn.) were as follows: 94° C. for 2 mins., then 30 cycles of 94° C. for 1 min., 58° C. for 2 mins., and 72° C. for 3 mins. PCR was followed by an additional extension at 72° C. for 7 minutes.

[0294] The PCR amplified product was run on a gel, an amplified fragment of approximately 360 bp was gel purified, and the isolated fragment was directly sequenced using ABI 373A DNA Sequencer (Perkin Elmer, Foster City, Calif.). The sequence analysis package of GCG was used to compare the obtained sequence with known sequences. The sequence was translated in all six reading frames in the GCG Analysis Program using the FastA algorithm (Pearson and Lipman, supra). The Swissprot database (GeneBio, Geneva, Switzerland) of proteins was searched. This translated cDNA fragment was identified as a part of a putative elongase based on the homology of the putative protein sequence to the *S. cerevisiae* ELO2 (GNS1), having 41.3% identity in 63 amino acids.

[0295] New primers were designed based on the putative elongase sequence and the vector, pZL1 (Life Technologies, Inc., Gaithersburg, Md.) sequence used to construct *M. alpina* cDNA library. The *M. alpina* excised cDNA library was PCR amplified again using primers RO350 (5'-CAT CTC ATG GAT CCG CCA TGG CCG CCG CAA TCT TG-3'), which has an added BamHI restriction site (underlined), and the vector reverse primer RO352 (5'-ACG CGT ACG TAAAGC TTG-3') to isolate the full length *M. alpina* elongase cDNA, using previously described conditions. The termini of the approximately 1.5 Kb PCR amplified fragment was filled-in with T4 DNA polymerase (Boehringer Mannheim Corp., Indianapolis, Ind. to create blunt ends and cloned into the pCR-blunt vector (Invitrogen Corp., Carlsbad, Calif.). This resulted in two clones, pRAE-1 and pRAE-2 (see FIG. 4A). (Plasmid DNA pRAE-2 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, on Aug. 28, 1998, under the terms of the Budapest Treaty, and was accorded deposit number ATCC 203166.) The elongase cDNAs from these vectors were cut out as an EcoRI fragment and cloned into the EcoRI digested pYX242 (Novagen, Madison, Wis.) vector. The clones pRAE-5 and pRAE-6 (see FIG. 4B) have the elongase cDNAs from pRAE-1 and pRAE-2, respectively. (Plasmid DNA pRAE-5 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, on Aug. 28, 1998, under the terms of the Budapest Treaty, and was accorded deposit number ATCC 203167.) The sequenc-

ing of pRAE-5 and pRAE-6 revealed that 5' untranslated region of the elongase gene in pRAE-5 is 16 bp shorter than that in pRAE-6 (see FIG. 5). The complete *M. alpina* elongase cDNA sequence, designated MAELO was obtained from pRAE-2 (see FIG. 6). FIG. 7 is the amino acid sequence obtained from the translation of MAELO. The Swissprot database (GeneBio, Geneva, Switzerland) was searched again, as previously described, with the translated MAELO: MAELO has 44.3% identity in 317 amino acids with *S. cerevisiae* GNS1(ELO2) and 44.7% identity in 318 amino acids with *S. cerevisiae* SUR4(ELO3). The FastA alignment among the three elongases is shown in FIG. 8. At the nucleotide level (see FIG. 9), MAELO has 57.4% identity in 549 bp overlap with *S. cerevisiae* GNS1(ELO2) (GenBank Accession #S78624). However, the identity between the complete MAELO gene of 954 bp and *S. cerevisiae* GNS1 (ELO2) is 33.0%.

EXAMPLE III

Expression of *M. alpina* Elongase cDNA in Baker's Yeast

[0296] The constructs pRAE-5, and pRAE-6 were transformed into *S. cerevisiae* 334 (Hoveland et al., *Gene* 83:57-64 (1989)) and screened for elongase activity. The plasmid pCGN7875 (Calgene LLC, Davis, Calif.) containing jojoba KCS gene in pYES2 vector (Invitrogen Corp., Carlsbad, Calif.) was used as a positive control. The substrate used to detect elongase activity in *M. alpina* elongase (MAELO) was GLA and that in jojoba KCS was oleic acid (OA). The negative control strain was *S. cerevisiae* 334 containing pYX242 vector. The cultures were grown for 40-48 hours at 25° C., in selective media (Ausubel et al., *Short Protocols in Molecular Biology*, Ch. 13, p. 3-5 (1992)), in the presence of a particular substrate. The expression of the jojoba KCS gene cloned in pYES2 was under the control of GAL1 promoter, while the promoter in pYX242 is TP1, which is constitutive. Hence, the 334(pCGN7875) and 334(pYES2) cultures were induced with galactose. The GC-FAME analysis of the lipid fraction of each cell pellet was performed as previously described (Knutzon et al., supra).

[0297] The elongase activity results from different experiments are provided in FIGS. 10A and 10B. The jojoba KCS elongates long chain monounsaturated fatty acids 18:1n-9 to 20:1n-9. The amino acid homology between the *M. alpina* elongase (MAELO) and the *S. cerevisiae* ELO2 and ELO3 suggested that the proteins encoded by these genes may have similar substrate specificity. The activity of the *M. alpina* elongase, elongation (MAELO) of long chain monounsaturated and saturated fatty acids, is seen in the conversion of 18:1n-9 to 20:1n-9 and also in the synthesis of 24:0. The control strain, 334(pYX242) has very little or no detectable amount of 20:1 and 24:0 (see FIG. 10A). *M. alpina* elongase (MAELO) also acts on at least one PUFA, converting 18:3n-6(GLA) to 20:3n-6(DGLA). The percentage of the 20:3n-6 in total lipid is higher in the strain 334(pRAE-5) and 334(pRAE-6) with the *M. alpina* elongase (MAELO) cDNA when compared to that in the control 334(pYX242). The percentages of 20:3n-6 produced were 0.092% for 334(pYX242) vs. 0.324% for 334(pRAE-5) and 0.269% for 334(pRAE-6) (shown in parenthesis in FIGS. 10A and 10B). This difference in the fatty acid profile is also seen in the total amount of 20:3n-6 produced. Only 0.226 μ g of

20:3n-6 was produced by 334(pYX242) while 334(pRAE-5) and 334(pRAE-6) produced 2.504 μ g of 20:3n-6 and 1.006 μ g of 20:3n-6, respectively. Also, when no substrate is added, the level of 20:3n-6 is not detectable.

[0298] Once 20:3n-6 is generated by the *M. alpina* elongase (MAELO), the Δ 5-desaturase can convert it to AA in the desired expression system. To test this hypothesis, the constructs pRAE-5 and pCGR-4 (a Δ 5-desaturase containing plasmid) were co-transformed into *S. cerevisiae* 334 and screened for AA production. The substrate used was 25 OM GLA (18:3n-6). If the *M. alpina* elongase (MAELO) is active in yeast, then the substrate will be converted to DGLA(20:3n-6), which the Δ 5-desaturase will convert to AA(20:4n-6). The results in FIG. 11 confirm the production of AA and therefore, the activity of the *M. alpina* elongase (MAELO).

[0299] The expression of Δ 5-, Δ 6-, and Δ 12-desaturases, in yeast, along with the elongase, should result in the production of AA (see FIG. 1) without the need for an exogenous supply of fatty acids.

EXAMPLE IV

A Comparison of the Expression of *M. alpina* Elongase cDNA MAELO and *S. cerevisiae* Elongase ELO2 in Baker's Yeast

[0300] The ELO2 gene encoding for the yeast elongase was cloned from an *S. cerevisiae* genomic library (Origene, Rockville, Md.) using the primers RO514 (5'-GGC TAT GGA TCC ATG AAT TCA CTC GTT ACT CAA TAT G-3') and RO515 (5'-CCT GCC AAG CTT TTA CCT TTT TCT TCT GTG TTG AG-3') incorporating the restriction sites (underlined) BamHI and HindIII (respectively). The ELO2 gene was cloned into the vector pYX242 at the BamHI and HindIII sites, designated pRELO, transformed into the *S. cerevisiae* host 334 (Hoveland et al., supra) and screened for PUFA elongase activity. The vector plasmid was used as a negative control and 334(pRAE-5) was grown to compare the PUFA elongase activity. The cultures were grown as previously described with no galactose in the media and 25 μ M GLA added as a substrate. FIG. 12 shows that amount of 20:3n-6 or DGLA produced (elongated from 18:3n-6 or GLA) by 334(pRAE-5) was approximately 4 times the negative control containing the unaltered vector pYX242, while the two individual clones 334(pRELO-1) and 334(pRELO-2) were only twice the negative control. Additionally, when DGLA produced is expressed as a percent of the total lipids (shown in parenthesis, FIG. 12), the clones 334(pRELO-1) and 334(pRELO-2) produced 0.153% and 0.2% DGLA respectively, while 334(pYX242) produced 0.185% DGLA. Hence all these strains produced comparable percentages of DGLA. The strain 334(pRAE-5), however, produced 0.279% DGLA, an increase of 50.8% over 334(pYX242) (negative control). These data show that the *S. cerevisiae* elongase gene ELO2, even when overexpressed in yeast, does not elongate GLA to DGLA effectively. The *M. alpina* PUFA elongase activity is specific for this conversion as evidenced by the higher amount of DGLA produced compared to the control, 334(pYX242).

EXAMPLE V

Identification of Elongases from Other Sources
Using MAELO

[0301] The TFastA algorithm (Pearson and Lipman, supra) is used to search for similarity between a query peptide sequence and the database DNA sequence translated in each of the six reading frames. Translated MAELO was used as the query for a TFastA search in GCG with the GenEMBL database (6/98) from GCG to identify other potential elongase sequences based on their amino acid similarity comparisons to translated MAELO. For example, in FIGS. 13 and 14, two alignments are shown between translations of two different *C. elegans* sequences from chromosome III and MAELO. *C. elegans* DNA sequence (GenBank accession #Z68749) was annotated denoting similarity with GNS1 (ELO2), while the additional *C. elegans* DNA sequence (GenBank accession #U61954) was noted as similar to both GNS1 and SUR4 (ELO3). These are spliced DNA fragments in which the introns have been removed from the genomic sequence, and the exons assembled and translated. The amount of amino acid identity between the putative PUFA elongases from *C. elegans* and translated MAELO are around 30%. This would point towards a common function in the fatty acid metabolism, e.g., a PUFA elongase. FIG. 15 is another example of a translated *C. elegans* sequence (GenBank accession #AF003134) from chromosome III. The DNA sequence was identified that had DNA homology to the *S. cerevisiae* ELO2. Further inspection of this DNA sequence and its amino acid translation determined that there was homology to translated MAELO. *C. elegans*, therefore, may contain a PUFA elongase.

[0302] FIG. 16 shows the alignments of translated DNA sequences from mouse and human, respectively, with translated MAELO. The mouse sequence CIG30, GenBank accession #U97107, was isolated from brown adipose tissue and reported as being "similar to yeast SUR4 protein". As shown in FIG. 16, amino acids numbered 130 to 152 in the U97107 translation contain a high degree of similarity to the translated MAELO. The human sequence, GenBank accession #AC004050, from chromosome 4 was from an HTGS (High Throughput Genome Sequence). There were no annotations contained with this sequence. However, translated AC004050 had 28.7% identity in 150 amino acids with translated MAELO. This gene fragment could be a fragment of a human PUFA elongase based on its amino acid similarity to translated MAELO.

[0303] FIG. 17 shows the amino acid alignment of translated MAELO and a mammalian sequence (GenBank accession #I05465, International Publication #WO 88/07577) which claims that the protein derived from expression of this sequence is a glycosylation inhibition factor. The amino acid identities between the two proteins, signifying that there could be related function, such as PUFA elongase activity.

[0304] These examples of other translated DNA sequences and their homology to the translated MAELO illustrate that any of the above examples could potentially be a PUFA elongase. These examples are not inclusive of all the possible elongases. However, use of MAELO or its amino acid translation as a query for database searches can identify other genes which have PUFA elongase activities.

EXAMPLE VI

M. alpina cDNA Library Screening Using a Plaque
Hybridization Method

[0305] In an effort to isolate additional PUFA elongase genes from *M. alpina*, a conventional plaque hybridization method was used to screen an *M. alpina* cDNA library made in a lambda vector. The DNA probe was generated based on MAELO nucleotide sequence and was used to screen the M7+8 *M. alpina* cDNA library made in a Ziplox vector (Knutzon et al., *J. Biol. Chem.* 273:29360-29366 (1998)).

[0306] To make the DNA probe for screening the library, the MAELO cDNA was digested with NspI and PvuI restriction endonucleases. Three small DNA fragments, with an average size of approximately 300 bp, were produced and used as probes. The rationale for using a mixture of fragmented MAELO cDNA was based on the assumption that there might be a common region or domain in the amino acid sequence which is conserved among various PUFA elongases present in *M. alpina*. Using MAELO DNA probes, the cDNA library was screened by a plaque hybridization technique according to standard protocol (Sambrook et al., *Molecular Cloning*, 2nd Ed., Cold Spring Harbor, 1989).

[0307] Briefly, 50,000 primary clones were plated and transferred to nylon membranes. The membranes were denatured and hybridized with alpha-³²P-dCTP-labelled MAELO DNA probes overnight in the hybridization buffer which contained 20% formamide, 0.2% PVP, BSA, Ficoll, 0.1% SDS and 0.5 M NaCl. The filters were washed with 0.5×SSC at 37° C. and exposed to X-ray film for autoradiography. This procedure was repeated three times. Four clones (designated as F1, F2, F3, and F4) which hybridized repeatedly were picked and suspended in SM buffer (Sambrook et al., supra) containing 7% DMSO.

[0308] The largest open reading frame of each candidate was subcloned into yeast expression vector pYX242 (Novagen, Inc., Madison, Wis.). The cDNA clones F1 and F3 were subcloned into pYX242 at the EcoRI site while F2 and F4 were subcloned at NcoI/HindIII sites. The recombinant pYX242 containing each candidate was transformed into SC334 (Hoveland et al., supra) for expression in yeast. To determine the elongase activity, as well as substrate specificity, SC334 containing each cDNA clone was grown in minimal media lacking leucine in the presence of 25 μM of GLA substrate as described in Example III. The fatty acid analysis was performed as described in Knutzon et al. (*J. Biol. Chem.* 273:29360-29366 (1998)). The results indicated that none of these four cDNA clones showed any significant activity in converting GLA to DGLA. Thus, the hybridization approach appeared to be unsuccessful in identifying additional PUFA elongases.

EXAMPLE VII

Construction of Direct cDNA Expression Library of
M. alpina in Yeast

[0309] To identify PUFA elongase genes other than MAELO, a different approach was taken to screen the *M. alpina* cDNA library. In particular, since Baker's yeast is incapable of producing long chain PUFAs due to the absence of respective desaturases and elongases, an attempt was made to construct an expression cDNA library of *M. alpina*

in *Saccharomyces cerevisiae*. The vector pYES2 (Novagen, Inc., Madison, Wis.), containing the GAL1 promoter, was chosen for the expression of cDNA library in *S. cerevisiae*.

[0310] The conventional way by which a cDNA library is made (i.e., transformation of cDNA/vector ligated DNA mixture into host cells) is difficult in yeast because the transformation efficiency by direct electroporation of ligated DNA mix is very low compared to the efficiency of purified supercoiled plasmid DNA. However, the major advantage of this method is to avoid amplification of primary clones which happens when the library is made in *E. coli* as an intermediate. Due to the limitation in the number of colonies to be screened, it was decided to first optimize the efficiency of transformation in different *S. cerevisiae* strains using cDNA/vector ligated mix. The best results were obtained with a yield of $4\text{--}5 \times 10^5$ transformants per μg of ligated DNA in *S. cerevisiae* strain SC334 (Hoveland et al., supra).

[0311] To make a direct *M. alpina* cDNA expression library in yeast total RNA was isolated from the fungus. *M. alpina* fungus (ATCC #32221) was plated onto cornmeal agar (Difco Laboratories, Detroit, Mich.) and grown at room temperature for 3–4 days. Once fungus growth was visible, it was inoculated into 50 ml of potato dextrose broth and shaken at room temperature very slowly to formulate spores. Once spores were visible, the 50 ml culture was inoculated into a 1 liter culture of potato dextrose, and spores were grown for 72 hours. After filtering through sterile gauze, the cells were immediately frozen into liquid nitrogen for future RNA extraction. Total RNA was prepared from 36 g of cell pellet using the hot phenol/LiCl extraction method (Sambrook et al., supra). The cell pellets were homogenized in a 10 mM EDTA, 1% SDS and 200 mM sodium acetate, pH 4.8 solution. Phenol and chloroform were added to the homogenates, and the aqueous layer was extracted. The aqueous layer was back extracted one more time with phenol and chloroform. Then an equal volume of 4 M lithium chloride was added. The samples were ethanol precipitated on ice for 3 hours, and pellets were obtained by centrifugation. The RNA pellets were washed with 70% ethanol and resuspended in DEPC treated water. Total RNA was quantitated by spectrophotometry and visualized by agarose gel electrophoresis to confirm the presence of 28S and 18S ribosomal bands. Approximately, 15 mg of total RNA were obtained from 36 gram of cell pellet.

[0312] The library was constructed according to the standard protocol (Sambrook et al., *Molecular Cloning*, 2nd Ed., Cold Spring Harbor, 1989). Messenger RNA was prepared from the total RNA using oligo dT cellulose affinity purification. Messenger RNA was reverse transcribed with oligo dT primer containing a XhoI restriction site using AMV reverse transcriptase. Following first strand cDNA synthesis, the second strand of cDNA was synthesized by adding *E. coli* DNA polymerase, *E. coli* DNA ligase and RNase H.

[0313] The EcoRI adaptor was ligated into the blunt-ended cDNA by T4 DNA ligase. The cDNA sample was kinased using T4 polynucleotide kinase and digested with XhoI, diluted with column buffer and passed through a Sephacryl S-400 column. The DNA samples were eluted by high salt buffer. Samples containing DNA from 400–5,000 bps were pooled and used for ligation into a pYES2 vector (Invitrogen Corp., Carlsbad, Calif.). The cDNA was ligated into the EcoRI/XhoI digested pYES2 vector using T4 DNA ligase. A

large scale ligation reaction was carried out since a large amount of the ligated DNA (2–3 μg) is required in direct transformation of yeast.

[0314] To transform yeast cells directly with the cDNA/pYES2 ligated mixture, competent SC334 cells were prepared using the LiAc TRAFO method (Gietz et al., *Mol. Cell. Biol.*, 5: 255–269, 1995). Briefly, fresh culture of SC334 from the plate was inoculated into 50 ml YPD medium. The culture was grown at 30° C. with shaking until the OD at 600 had reached 1.0. Thirty ml of this starter was inoculated into 300 ml of YPD liquid medium and incubated with shaking until the cell number of the culture reached 35×10^6 cell/ml (approximately 3–4 h). The cells were harvested and washed with sterile water. The entire cell pellet was resuspended in 1.5 ml of freshly prepared 1×TE/LiAc (0.1M LiAc). These cells were used immediately for the transformations.

[0315] Seven hundred and fifty microliters of competent SC334 cells were aliquoted into 15 ml falcon tubes. Approximately 2 μg of cDNA/pYES2 ligated DNA were added to the cells along with carrier DNA and mixed gently. Three milliliters of sterile 40% PEG/LiAc was added to the cells and mixed gently but thoroughly. The cells were incubated at 30° C. for 30 min with shaking and subsequently given heat shock at 42° C. for 15 min. The cells were cooled, pelleted, and resuspended in 5 ml of 1×TE. A 100 μl aliquot of the above cells was plated onto fifty 150 mm selective agar plates lacking uracil (Ausubel et al., supra) and incubated at 30° C. for 3 days. A total of 8×10^5 primary clones were obtained. Five colonies were pooled in 1 ml minimal media lacking uracil (Ausubel et al., supra) and glycerol added to prepare stocks. A total of 5,000 pools were made for screening.

EXAMPLE VIII

MAD (*M. alpina* Direct) Screening in Yeast

[0316] The quality of the library was analyzed by determining the average size of the cDNAs in the library. Since the screening of the library was based on the expression of the cDNA, it was important to determine the average size of the cDNA present in the library. The expression library containing the longest cDNAs would be the best appropriate choice to isolate full-length cDNAs of interest. To this end, randomly selected pools were plated onto selective agar plates, as described in Example VII, to obtain individual colonies. Forty different yeast colonies were randomly picked, and each colony was inoculated into 5 ml of selective liquid medium lacking uracil (as described in Example VII) and grown, while shaking, for 24 hours at 30° c. Plasmid DNA was extracted from these colonies by the bead beating method (Hoffman et al., *Gene* 57:267 (1987)) adapted as follows:

[0317] Pellets from 5 ml of culture were lysed in 0.5 ml of a 100 mM NaCl, 10 mM Tris, pH 8.0, 1 mM EDTA and 0.1% SDS solution. Sterile 0.5 mm glass beads of equal volume were added and manually vortexed for 3 minutes. Two hundred microliters of the same buffer were added, and the mixture was vortexed for an additional minute. The samples were centrifuged on high for 2 minutes, and cytoplasmic extract was then transferred to a fresh tube. An equal volume of phenol/ CHCl_3 was added to the sample, vortexed and centrifuged again for 2 minutes. The aqueous layer was

re-extracted twice and precipitated with 0.3 M sodium acetate and approximately 2.5 volumes of ethanol for 30 minutes at -20°C . The precipitates were washed with 70% ethanol and resuspended in water. To eliminate RNA and any protein contamination, the plasmid DNAs isolated from 40 different samples were further purified using the QIAprep Spin Miniprep Kit according to the manufacturer's protocol (Qiagen Inc., Valencia, Calif.). The plasmid DNA samples were then restricted with EcoRI and XhoI restriction endonucleases to release the cDNA fragment, and the digest was analyzed on 1% agarose gel. The results indicated that the majority of the cDNAs of the direct library varied in length from 0.8 Kb to 1.5 Kb.

[0318] To screen the library, the glycerol stocks were thawed and approximately 0.5 ml was added to 5 ml of liquid selective media lacking uracil (Ausubel et al., supra) and grown at 30°C for 24 hours. The culture was then transferred into 50 ml of liquid selective medium lacking uracil with 2% galactose and 25 μM GLA (substrate for the elongase enzyme) for 24 hours at 25°C with shaking. The GC-FAME analysis of the lipid content in the cell pellet of each induced culture was performed as previously described (Knutzon et al., supra). The MAELO (pRAE-5 in pYX242 grown in selective media lacking leucine) was used as a positive control in each batch run. MAELO had consistently been able to convert 1.5% of GLA to DGLA (see Example III).

EXAMPLE IX

Identification of a cDNA Encoding a Potential PUFA Elongase

[0319] After screening and analyzing approximately 750 individual pools by GC-FAME analysis, as described in Example VIII, one pool of five colonies (i.e., MAD708) appeared to have significant enzymatic activity in converting GLA to DGLA. This activity was found to be approximately 5 fold higher than the *M. alpina* elongase activity (MAELO) in terms of DGLA/GLA ratio (**FIG. 19**). This pool was tested again under identical assay conditions to confirm the initial findings. The repeat experiment showed 9.5% conversion of GLA to DGLA and was again around 5 fold higher than *M. alpina* elongase activity (MAELO). These results strongly indicated that the MAD708 pool contained an elongase candidate which was specific for GLA as substrate. Since MAD708 was a pool of five different clones, it was necessary to isolate the individual cDNA clone which encoded for elongase activity from this pool. To do this, the original MAD708 glycerol stock was plated onto a selective media agar plate lacking uracil (Ausubel et al., supra). Thirty individual colonies were picked and grown in liquid selective medium, lacking uracil with 2% galactose, as previously described in Example VIII, in the presence of GLA. The cell pellet obtained from each culture was then subjected to fatty GC-FAME analysis (Knutzon et al., supra) along with a positive control of 334 (pRAE-5)(MAELO in pYX242). The fatty acid analysis from the 30 individual clones from the MAD708 expression pool in yeast revealed that 5 of the 30 clones showed elongase activity in converting GLA to DGLA. The fatty acid profiles of the active clones MAD708-2, MAD708-10, MAD708-18, MAD708-19 and MAD708-30 are shown in **FIG. 20**. As shown in this Figure, MAD708-2, 10, and 30 produced the most DGLA, approxi-

mately 25 fold more than MAELO (pRAE-5). These 3 converted in the range of 41% to 49% of GLA to DGLA. Other clones, MAD708-18 and MAD708-19, converted 8% and 21% of GLA to DGLA, respectively. All MAD708 clones converted a higher percentage of GLA to DGLA with respect to MAELO encoded elongase (3.4%).

EXAMPLE X

Characterization of cDNAs Encoding Elongase

[0320] Plasmid DNA was extracted from SC334 yeast clones (MAD708 pool) that showed significant GLA specific elongase activity by the bead beating method, as described in Example VIII. To determine the size of the cDNA insert, PCR was performed using each plasmid DNA obtained from positive elongase clones as a template. The forward primer RO541 (5'-GAC TAC TAG CAG CTG TAA TAC-3') and the reverse primer RO540 (5'-GTG AAT GTA AGC GTG ACA TAA-3') are in the multicloning site of the pYES2 vector and were used to amplify the cDNA insert within the EcoRI and XhoI sites. PCR reaction was performed in a 50 μl volume containing 4 μl of plasmid DNA, 50 pmole of each primer, 5 μl of 10 $\times$ buffer, 1 μl 10 μM PCR Nucleotide Mix (Boehringer Mannheim Corp., Indianapolis, Ind.) and 0.5 μl of High Five Taq polymerase (Boehringer Mannheim, Indianapolis, Ind.). The amplification was carried out as follows: 2 mins. denaturation at 94°C , then 94°C for 1 min, 55°C for 2 mins., and 72°C for 3 mins. for 30 cycles, and 7 mins. extension at 72°C at the end of the amplification. Analysis of PCR amplified products on a 1% agarose gel showed the sizes of the elongase cDNAs to be around 1.0-1.2 Kb. The plasmid DNAs, containing the potential elongase cDNAs, were designated as pRPB2, pRPB10, pRPB18, pRPB19, and pRPB30. Since the cDNA library was made in the pYES2 vector at the EcoRI and XhoI sites, the size of the cDNA present in each plasmid was further confirmed by digesting the above plasmids with EcoRI and XhoI.

[0321] The plasmid DNAs isolated from yeast were re-amplified in *E. coli* for long-term storage of the cDNA clones as well as for DNA sequencing. *E. coli* TOP10 (Invitrogen Corp., Carlsbad, Calif.) cells were transformed with the pRPB recombinant plasmids according to the manufacturer's protocol. The transformants obtained from each plasmid DNA were inoculated into LB containing ampicillin (50 $\mu\text{g}/\text{ml}$) and grown overnight at 37°C with shaking. Plasmid DNAs were isolated from these cultures by using QIAprep Spin Miniprep (Qiagen Inc., Valencia, Calif.) according to the manufacturer's protocol. The purified plasmid DNAs were then used for sequencing from both 5' and 3' ends. The DNA sequencing was performed by using a 373A Stretch ABI automated DNA sequencer (Perkin Elmer, Foster City, Calif.) according to the manufacturer's protocol. Primers used for sequencing were the forward primer RO541 (5'-GAT TAC TAG CAG CTG TAA TAC-3') and the reverse primer RO540 (5'-GTG AAT GTA AGC GTG ACA TAA-3') contained in the multicloning sites of the pYES2 vector. The obtained nucleotide sequences were transferred to Sequencher software program (Gene Codes Corporation, Ann Arbor, Mich.) for analysis. The DNA sequence analysis revealed that all five elongase cDNAs contained the identical nucleotide sequence with a common overlap of 301 nucleotides. Each DNA sequence contains a putative start site at the beginning of the 5' end and a stop codon with poly

A tail at the end of the 3' site. To further confirm the DNA sequence, internal forward primers R0728 (5'-GAG ACT TTG AGC GGT TCG-3') and R0730 (5'-TCT CTG CTG CGT TGA ACT CG-3'), along with reverse primers R0729 (5'-AAA GCT CTT GAC CTC GAA C-3') and R0731 (5'-AAC TTG ATG AAC GAC ACG TG-3') were designed within the cDNA, and used for sequencing of pRPB2, since this candidate possessed the highest elongase activity. The entire nucleotide sequence was analyzed by the Sequencer program (**FIG. 21**), and the longest open reading frame deduced from the entire cDNA sequence in pRPB2 appeared to be 957 bp in length (**FIG. 22**). The deduced open reading frame was then translated into the corresponding amino acid sequence, and the predicted sequence is shown in **FIG. 23**. The elongase encoded by the cDNA (pRPB2) identified from *M. alpina* appears to be a 318 amino acid long protein which is nearly identical in size with translated MAELO. This new elongase cDNA was designated as "GLELO" and its encoded protein has been named "GLA elongase".

[0322] Plasmid DNA pRPB2 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209 on Jul. 22, 1999 under the terms of the Budapest Treaty. It was accorded ATCC Deposit #PTA-402.

EXAMPLE XI

Biochemical Characterization of GLA Elongase (GLELO)

[0323] A. Confirmation of GLA Elongase Activity

[0324] To further confirm the activity of the GLA elongase encoded by the pRPB2 recombinant plasmid, elongase activity screening was repeated on the yeast clone SC334 containing pRPB2 plasmid. This experiment was also conducted to assure consistent lipid extraction and to detect the activity of GLA elongase by averaging four independent experiments. The *S. cerevisiae* 334 glycerol stock containing pRPB2 was plated onto minimal media agar plates lacking uracil. Individual colonies were randomly picked and grown in minimal medium lacking uracil, as described in Example VIII. The four independent cultures were combined, and a 5 ml aliquot was used as an inoculum for four separate 50 ml cultures. The cultures were then grown in the presence of GLA and were subjected to fatty acid analysis along with a negative control of *S. cerevisiae* 334 containing pYES2, as described in Example VIII. The average elongase activity from four independent cultures of 334(pRPB2) with 25 μ M GLA is shown in **FIG. 24**. The GLA elongase activity of each of the four independent samples of 334(pRPB2) appeared to be consistent with an average conversion of 62% GLA to DGLA.

[0325] B. Determination of GLELO Substrate Specificity for GLA Elongase

[0326] To analyze the substrate specificity of the GLA elongase, the culture of 334(pRPB2) was tested with different fatty acid substrates besides GLA (e.g., SA(18:0), OA(18:1), LA(18:2n-6), AA(20:4n-6), ADA(22:4n-6), ALA(18:3n-3), STA(18:4n-3), and EPA(20:5n-3)). Under identical assay conditions, the only other substrate utilized by the elongase enzyme was STA, a fatty acid from the n-3 pathway. GLA elongase was able to convert 73% of STA to 20:4n-3 (**FIG. 25**). From these experiments, it can be

concluded that the GLA elongase has substrate specificity for both GLA and STA, indicating that it possesses elongase activity along both the n-6 and n-3 pathways.

[0327] C. Co-expression of Fungal GLELO and Δ 5-Desaturase Gene in Yeast

[0328] Once DGLA (20:3n-6) is produced by the GLA elongase, the Δ 5-desaturase can convert it to AA (20:4n-6) in a desired co-expression system. This scheme, as depicted in **FIG. 1**, can be tested by co-transforming *S. cerevisiae* 334 with plasmids pRPB2 and pRPE31 (the recombinant plasmid pYX242 containing a Δ 5-desaturase cDNA (**FIG. 18**) cloned at the EcoRI site. The co-transformed yeast cultures were supplemented with 25 μ M GLA and analyzed for AA synthesis. If both elongase and Δ 5-desaturase enzymes are expressed, the GLA substrate will be converted to DGLA, which will then be converted to AA. The results in **FIG. 26A** indicate that the sequential action of GLA elongase and Δ 5-desaturase on GLA substrate resulted in an average conversion of 27% GLA to AA. Therefore, the GLA elongase has the ability to work with other enzymes in the n-6 PUFA synthetic pathway to produce desirable fatty acids.

[0329] To determine whether the above conversion is also true in n-3 pathways, the similar co-expression experiments were carried out in the presence of 25 μ M STA. Again, if both enzymes are expressed, the STA substrate will be converted to 20:4n-3 which will then be converted to EPA (20:5n-3) by the Δ 5-desaturase. **FIG. 26B** shows the results in which the production of EPA (approx. 40%) is observed. Once again, the GLA elongase demonstrates its ability to work with Δ 5-desaturase in the n-3 pathway to produce desirable fatty acids.

EXAMPLE XII

Sequence Comparison Between GLELO and Other Fungal Elongases

[0330] The sequence analysis package of GCG (see Example I) was used to compare the GLELO sequence with known protein sequences. The nucleotide sequence of GLELO open reading frame was first translated into amino acid sequence that was used as a query sequence to search Swissprot database (see Example I) using the FastA algorithm (see Example I). Based on amino acid sequence similarity, the best matches were found with *S. cerevisiae* YJT6 (an EST with unknown annotation) with 33.9% identity in 189 amino acid overlap, *S. cerevisiae* ELO2 (GNS1) with 25.8% identity in 295 amino acid overlap, and *S. cerevisiae* ELO3 (SUR4) with 25.2% identity in 313 amino acid overlap. The FastA alignment of GLELO with MAELO showed 30.9% identity in 275 amino acids (**FIG. 27**). GCG Pileup program creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments (see Example I), and was used with the elongases described above. The Pileup results indicate that there are many conserved regions among the elongases including a putative histidine box, which is underlined (Knutzon et. al., *J. Biol. Chem.* 273: 29360-29366, 1998) (**FIG. 28**). Thus, although GLELO has similarity with MAELO, the difference in their encoded elongases may presumably be due to their substrate preference. GLA elongase can convert a higher percentage of GLA to DGLA than *M. alpina* elongase. In addition, MAELO expression in *S. cerevisiae*

showed elongation of saturated and monounsaturated fatty acids in addition to GLA elongation to DGLA (see Example III).

EXAMPLE XIII

Identification of *M. alpina* MAELO Homologues in Mammals

[0331] The MAELO translated sequence was used to search the Unified Human Transcript Database of Abbott Laboratories, 100 Abbott Park Rd., Abbott Park, Ill. 60064. This database was searched using Basic Local Alignment Search Tool (BLAST) (Altschul et al., *Nuc. Acids Res.* 25:3389-3402 (1997)) which "is a set of similarity search programs designed to explore all of the available sequence databases regardless of whether the query is a protein or DNA." Specifically, the tblastn algorithm was used (i.e., a protein query search to a nucleotide database translated in six reading frames). The contig (CC) sequences in the Unified Human Transcript Database are consensus sequences representing groups of expressed sequence tags (EST) cDNAs derived from the public domain and from the Incyte LIFESEQ™ database of ESTs (Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, Calif. 94304) that are clustered together on the basis of defined sequence homology, and assembled on the basis of sequence overlap. Two sequences from this database, CC067284R1 and CC1484548T1 had 28% identity in 242 amino acid overlap and 28.6% identity in 266 amino acid overlap, respectively, with the translated MAELO sequence. The two derived and edited sequences were designated as hs1 and hs2, respectively, and copied into the sequence analysis software package of GCG (see Example I). The translated MAELO sequence was aligned with translated HS1 (28.5% identity in 242 amino acids) and HS2 (28.2% identity in 266 amino acids) cDNA sequences using the FastA algorithm, as shown in **FIGS. 29 and 30**, respectively. HS1 cDNA nucleotide sequence also had 86.9% identity in 844 bp with the I05465 nucleotide sequence (see Example V). The translated HS2 cDNA sequence had 100% identity with the amino acid sequence from GenBank with accession number W74824 (see published PCT application WO9839448).

[0332] The National Center for Biotechnology Information (NCBI at <http://www.ncbi.nlm.nih.gov/>) was used to conduct database searches using tblastn with the 28 amino acid sequence (DTIFIILRKQKLIFLHWYHHITTVLLYSW) translated from AC004050 (a human sequence identified in a TFASTA search, see Example V). This amino acid sequence contains a histidine box (underlined), which has a noted motif of desaturases (Knutzon et al., supra), and both PUFA elongases, MAELO and GLELO (see **FIG. 28**). A translated mouse sequence shown previously in Example V (GenBank Accession #U97107) and a translated *C. elegans* sequence (GenBank Accession #U41011) had the highest matches with this 28 amino acid query. The NCBI mouse EST database was searched again with tblastn, using translated U41011 as a query. An additional mouse sequence was identified (GenBank Accession #AF014033.1), annotated as "putative involvement in fatty acid elongation." Three longer sequences (GenBank Accession #'s AA591034, AA189549, and AA839346) were identified through a tblastn search of the mouse EST database with translated AF014033.1 and combined into one sequence designated as

mm2. The FastA alignment (see Example I) of translated mm2 and MAELO is shown in **FIG. 31**. Another related, but not identical mouse sequence (GenBank Accession #AI225632), was also identified in a tblastn search of the mouse EST database with AF014033.1. The FastA alignment with translated AI225632 to MAELO is shown in **FIG. 32**. The percent identity for both translated MM2 and AI225632 with translated MAELO is 30.4% in 191 and 115 amino acid overlap, respectively. The level of amino acid identity with translated MAELO with these two translated mouse sequences identifies them as putative homologues of PUFA elongases.

EXAMPLE XIV

Identification of *M. alpina* GLELO Homologues in Mammals

[0333] The TFASTA algorithm, which compares a protein sequence to the database DNA sequence translated in each of the six reading frames, was used with translated GLELO as the query. The GenEMBL database from GCG was used to identify other potential elongase sequences based on their amino acid similarity to translated GLELO. Three human sequences were found to have matches with the GLELO amino acid sequence. These sequences have GenBank accession numbers 1) AI815960, 2) AL034374, and 3) AC004050. AI815960, a *Homo sapiens* EST sequence, has 40.3% identity in 144 amino acid overlap with translated GLELO (see **FIG. 33**). A translated region of the human genomic sequence AL034374, derived from chromosome VI has 46.7% identity in a 60 amino acid overlap with translated GLELO. This homologous region in AL034374 appeared to be a part of the HS1 amino acid sequence which was shown to have homology with translated MAELO (see Example XIII). Therefore, HS1 sequence has similarity with both MAELO (see **FIG. 29**) as well as GLELO (see **FIG. 34**). A translated region of a human genomic sequence AC004050 from chromosome IV has 34.8% identity in 89 amino acid overlap with translated GLELO (see **FIG. 35**). The amino acid identities between GLELO and these human sequences indicate that the proteins derived from these human sequences could have related function, such as PUFA elongase activity.

[0334] To identify a mouse cDNA similar to GLELO, TFASTA searches were performed with the GenEMBL database using translated GLELO as a query. From the TFASTA searches, the three mouse sequences with the highest matches to translated GLELO were identified: (GenBank accession numbers 1) AF104033, 2) AI595258, and 3) U97107). AF104033 is annotated as "MUEL protein having putative fatty acid elongase with homology to yeast ELO3 (SUR4)" and is a part of the sequence of MM2. The MM2 sequence was initially derived from AF104033 mouse sequence, but the entire MM2 sequence was finally obtained through further mouse EST database searches and also shown to have homology with translated MAELO (see Example XIII and **FIG. 31**). When this MM2 amino acid sequence was aligned with translated GLELO sequence using FastA, a 34.6% identity in 211 amino acid overlap was found (see **FIG. 36**) indicating that MM2 also has homology with GLELO. AI595258 is a mouse cDNA clone having 5' similarity with yeast ELO3 elongase and is part of mouse EST cDNA AI225632. The AI225632 mouse sequence,

which is a longer sequence than AI595258, was shown to have similarity with translated MAELO (see FIG. 32). The AI225632 was also aligned with the translated GLELO, and the FastA alignment is shown in FIG. 37. A 35.3% identity in 199 amino acid overlap has been found. The third sequence, U97107, a mouse sequence, was annotated as "similar to yeast ELO3 (SUR4) gene." The FastA alignment of translated GLELO with U97107 is shown in FIG. 38 where a 23.7% identity in 279 amino acid overlap was found. Previously, a region of U97107 was also found to have a high degree of homology with MAELO based on a FastA alignment (see Example V and FIG. 16).

[0335] The above searches clearly indicate that the same human and mouse sequences were obtained by using either MAELO or GLELO as a query.

EXAMPLE XV

Identification of *M. alpina* GLELO and MAELO Homologues in Other PUFA Producing Organisms

[0336] A) *Caenorhabditis elegans*:

[0337] A putative amino acid sequence deduced from a chromosomal sequence of *C. elegans* (GenBank Accession #U41011) was able to identify a partial sequence contained in the mouse MM2 putative PUFA elongase which has amino acid similarity with both GLA elongase (GLELO) and *M. alpina* elongase (MAELO). It was therefore conceivable that *C. elegans* homologues of GLELO or MAELO might be present in the nematode database. The putative amino acid sequences derived from GLELO and MAELO sequences were used as queries independently to search the nematode databases. A BLAST search (see Example XIII) was performed on wormpep16 (blastp compares an amino acid query sequence against a nucleotide sequence database) and wormpep 16cDNAs (tblastn) databases which are predicted proteins and cDNAs obtained from the *C. elegans* genome sequencing project or EST's and their corresponding cDNA sequences, respectively. These sequence data were produced by the *C. elegans* Sequencing group, carried out jointly by the Sanger Centre and Genome Sequencing Center, and can be obtained from <ftp://ftp.sanger.ac.uk/pub/databases/wormpep/>. At least seven putative *C. elegans* translated sequences were identified by their amino acid sequence homology to the translated amino acid sequence of both GLELO and MAELO. The GenBank Accession #'s of those genomic sequences containing the deduced amino acids were identified as Z19154, U68749 (2 deduced proteins (F56H11.4 and F56H11.3 (wormpep Accession #'s)), U41011, U61954 (2 deduced proteins (F41H10.7 and F41H10.8 (wormpep Accession #'s)), and Z81058. Those underlined were identified in a previous search using translated MAELO as query (see Example V). As an example, the FastA amino acid alignments of translated U68749 (F56H11.4) with translated GLELO and MAELO are shown in FIGS. 39 and 40. Translated U68749 (F56H11.4) has 25-30% identity with both *M. alpina* elongase and GLA elongase in approximately a 200 amino acid overlap (see FIGS. 39 and 40). For all seven translated putative *C. elegans* cDNAs, the FastA alignments to translated GLELO was between 25-30% identity in a 200 amino acid overlap, while the identity was 26-34% in at least a 188 amino acid overlap for translated MAELO. The alignment similarities

indicate that either translated GLELO or MAELO can be used to identify potential genes from *C. elegans* with elongase activity.

[0338] B) *Drosophila melanogaster*:

[0339] The translated deduced cDNA from the genomic sequence U41011 (*C. elegans*) had its highest match with a *Drosophila melanogaster* EST, accession number AI134173 in a blastn search (compares a nucleotide query sequence against a nucleotide database) of the "other ESTs" database through NCBI (see Example XIII) and was assembled with an overlapping DNA EST fragment, accession number AI517255. The translated DNA fragment DM1, derived from the two overlapping sequences was aligned with translated GLELO as well as MAELO (see FIGS. 41 and 42) using FastA in GCG (see Example I). The alignments showed 27.2% identity with GLA elongase in a 206 amino acid overlap and 30% identity with *M. alpina* elongase in a 237 amino acid overlap. Thus, based on amino acid similarity, the DM1 could be a potential homologue to GLELO or MAELO having PUFA elongase-like activity. Moreover, using DNA sequences of GLELO and MAELO as queries for database searches, homologues with PUFA elongase activity from *Drosophila* can be identified.

EXAMPLE XVI

Cloning and Expression of a Human PUFA Elongase Homologue

[0340] Many potential PUFA elongase sequences were identified based on their amino acid similarities to translated GLELO and/or MAELO. To determine the potential elongase activities of these sequences, the cDNA encoding the full-length protein is then identified, cloned, and expressed, as demonstrated in the present example.

[0341] Primers RO719 (5'-GGT TCT CCC ATG GAA CAT TTT GAT GCA TC-3') and RO720 (5'-GGT TTC AAA GCT TTG ACT TCA ATC CTT CCG-3') were designed based on the putative HS1 sequence, and used to amplify the human liver Marathon-Ready cDNA (Clontech Laboratories, Inc., Palo Alto, Calif.). The polymerase chain reaction (PCR) was carried out in a 50 μ l volume containing: 5 μ l of human liver Marathon-Ready cDNA, 50 pmole each primer, 1 μ l 10 mM PCR Nucleotide Mix (Boehringer Mannheim Corp., Indianapolis, Ind.), 5 μ l 10xbuffer and 1.0 U of Advantage KlenTaq Polymerase Mix (Clontech Laboratories, Inc., Palo Alto, Calif.). Thermocycler conditions in Perkin Elmer 9600 (Norwalk, Conn.) were as follows: 94° C. for 2 mins, then 30 cycles of 94° C. for 1 min., 58° C. for 2 mins, and 72° C. for 3 mins. PCR was followed by an additional extension cycle at 72° C. for 7 minutes.

[0342] The PCR amplified product was run on a gel, an amplified fragment of approximately 960 bp was gel purified, the termini of the fragment filled-in with T4 DNA polymerase (Boehringer Mannheim, Corp., Indianapolis, Ind.), and cloned into pCR-Blunt Vector (Invitrogen Corp., Carlsbad, Calif.) following manufacturer's protocol. The new plasmid was designated as pRAE-52, and the putative PUFA elongase cDNA in this clone was sequenced using ABI 373A Stretch DNA Sequencer (Perkin Elmer, Foster City, Calif.). The putative PUFA elongase cDNA sequence in plasmid pRAE-52 is shown in FIG. 43, and the translated sequence is shown in FIG. 44.

[0343] The putative PUFA elongase cDNA from plasmid pRAE-52 was then digested with NcoI/HindIII, gel purified, and ligated into pYX242(NcoI/HindIII). The new plasmid was designated as pRAE-58-A1. (Plasmid 58-A1 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209 on Aug. 19, 1999, under the terms of the Budapest Treaty and was accorded deposit number PTA-566.) The construct pRAE-58-A1 was transformed into *S. cerevisiae* 334 (Hoveland et al., supra) and screened for elongase activity. The negative control strain was *S. cerevisiae* 334 containing pYX242 vector. The cultures were grown for 24 hours at 30° C., in selective media (Ausubel et al., supra), in the presence of 25 μ M of GLA or AA. In this study, DGLA or adrenic acid (ADA, 22:4n-6), respectively, was the predicted product of human elongase activity. When GLA was used as a substrate, the yeast cells containing the human elongase cDNA contained elevated levels of DGLA compared to control cells, 2.75% vs. 0.09% of total fatty acids, respectively (see FIG. 45). When AA was used as a substrate, the yeast cells containing the human elongase cDNA contained elevated levels of ADA compared to control cells, none detected vs. 1.21% of total fatty acids, respectively. Thus, the human elongase converts both 18 and 20 carbon chain long PUFAs to their respective elongated fatty acids.

[0344] The yeast cells containing the human elongase cDNA also had elevated levels of monounsaturated fatty acids including 18:1n-7, 20:1n-7, 20:1n-9, and 18:1n-5, compared to the control strain. Therefore, these results indicate that the identified human elongase is capable of utilizing PUFAs as well as monounsaturated fatty acids as substrates. Thus, this human sequence HSELO1, and its encoded protein (HSELO1p), possess elongase activity independent of substrate specificity.

[0345] To further confirm the substrate specificity of the human elongation enzyme, described above and referred to herein as HSELO1, the recombinant yeast strain 334(pRAE-58-A1) was grown in minimal media containing n-6 fatty acids GLA, AA, or n-3 fatty acids ALA, STA, or EPA. The lipid profiles of these yeast cultures, when examined by GC and GC-MS, indicated that there were accumulations of DGLA, ADA, ω 3-eicosatrienoic acid (ETra, C20:3n-3), ETA, and DPA, respectively (FIG. 51). The levels of these fatty acids were 7.29% (DGLA), 6.26% (ADA), 6.15% (ETra), 10.06% (ETA), and 6.66% (DPA), respectively, of the total fatty acids in the strain containing the pRAE-58-A1 sequence. These represented 78.3%, 42.7%, 30.4%, 79.2%, and 71.7% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms.

[0346] The yeast cells expressing the recombinant HSELO1 sequence, compared to the control cells, also contained significantly elevated levels of C18:1n-7 and C20:1n-7, and to a lesser extent, eicosenoic acid (EA, C20:1n-9) (FIG. 45). This finding suggested that the recombinant HSELO1 protein (HSELO1p) might also be involved in the elongation of monounsaturated fatty acids of 16 or 18 carbon lengths. To confirm this hypothesis, 25 μ M of exogenous OA was added as a substrate to the recombinant yeast strain 334(pRAE-58-A1). After incubation, the accumulation of EA at 2.25% of the total fatty acids demonstrated that the expressed HSELO1 enzyme could elongate monounsaturated fatty acids (FIG. 51). However, the conversion of OA to EA by recombinant HSELO1p was only 8.9%; this

conversion was significantly lower than the endogenous conversion of C16:1n-7 (to C18:1n-7) or C18:1n-7 (to C20:1n-7), which was 20.4% and 58.1%, respectively.

[0347] To determine whether the substrate concentration affects the conversion of 18 and 20 carbon fatty acids to the respective elongated products, two different concentrations of GLA, AA, and EPA were examined (FIG. 52). When 25 μ M of the substrates GLA, AA, and EPA were added exogenously, the levels of the fatty acids produced by two carbon elongation were 3.95% (DGLA), 2.91% (ADA), and 4.82% (DPA), respectively, of the total fatty acids in the lysates of 334(pRAE-58-A1). These represented 62.4%, 27.5%, and 70.3% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. When 100 μ M of the substrates GLA, AA, and EPA were added, the levels of the fatty acids produced by two carbon elongation were 9.56% (DGLA), 3.90% (ADA), and 11.50% (DPA), respectively, of the total fatty acids in the lysates of 334(pRAE-58-A1). These represented 39.8%, 15.7%, and 45.7% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. Although the addition of more substrates led to higher percentages of the two carbon elongated products, the overall conversion rate decreased by at least 35%.

[0348] To further confirm the substrate specificity of HSELO1p, the recombinant yeast strain 334(pRAE-58-A1) was grown in minimal media containing 25 μ M of saturated, monounsaturated, or PUFAs. The lipid profiles of these various substrates revealed that HSELO1p is not involved in the elongation of saturated fatty acids such as palmitic acid (PA, C16:0), stearic acid (SA, C18:0), arachidic acid (ARA, C20:0), behenic acid (BA, C22:0) (FIG. 53A). HSELO1p is also not involved in the elongation of monounsaturated fatty acids OA and EA. When PTA was added as a substrate, 12.76% of the total fatty acids was OA. However, this is not an increase in the level of OA compared to the samples where PTA was not added, as OA was 25-31% of the total fatty acids in all samples. HSELO1p is involved in the elongation of n-6 PUFAs LA, GLA, and AA, but not DGLA or ADA (FIG. 53B). The lipid profiles of these yeast cultures indicated that there were accumulations of C20:2n-6, DGLA, and ADA, respectively, but not C22:3n-6 or C24:4n-6. The levels of these fatty acids were 0.74% (C20:2n-6), 2.46% (DGLA), and 2.14% (ADA), respectively, of the total fatty acids in the lysates of 334(pRAE-58-A1). These represented 13.2%, 51.4%, and 27.1% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. HSELO1p is also involved in the elongation of n-3 PUFAs ALA, STA, and EPA, but not DPA (FIG. 53C). The lipid profiles of these yeast cultures indicated that there were accumulations of ETra, ETA, and DPA, respectively, but not C24:5n-3. The levels of these fatty acids were 1.03% ETra, 2.24% (ETA), and 3.19% (DPA), respectively, of the total fatty acids in the strain containing the pRAE-58-A1 sequence. These represented 22.2%, 61.9%, and 39.5% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. All results confirmed that the expression of HSELO1 from human liver in yeast resulted in the elongation of various long-chain PUFAs in n-6 and n-3 fatty acid pathways.

EXAMPLE XVII

Cloning, Expression and Characterization of a *C. elegans* PUFA Elongase

[0349] Several putative *C. elegans* elongases were identified with amino acid homology to both translated GLELO and MAELO. As with the human cDNA sequence, cloning of a cDNA and expression in yeast was used to determine if indeed it was a PUFA elongase. Primers RO738 (5'-AAT CAG GAA TTC ATG GCT CAG CAT CCG CTC GTT CAA C-3') and RO739 (5'-CCG CTTGTC GAC TTA GTT GTT CTT CTT CTT TGG CAC-3') with restriction sites EcoRI and SalI (underlined), respectively, were based on the putative cDNA sequence contained in the genomic sequence U68749 (wormpep cDNA accession #F56H11.4). A PCR amplification was performed in a 100 μ l volume containing: 250 ng excised *C. elegans* library cDNA (OriGene Technologies Inc., Rockville, Md.), 50 pmole each primer, 10 μ l 10 $\times$ reaction buffer (Boehringer Mannheim Corp., Indianapolis, Ind.), 1 μ l 10 mM PCR Nucleotide mix (Boehringer Mannheim Corp., Indianapolis, Ind.), and 2.5 U Taq polymerase (Boehringer Mannheim Corp., Indianapolis, Ind.). Thermocycler conditions in a Perkin Elmer 9600 (Norwalk, Conn.) were as follows: 95° C. for 5 mins, then 25 cycles of 94° C. for 30 secs, 55° C. for 2 mins, and 72° C. for 2 mins. PCR was followed by an additional cycle of 72° C. for 7 minutes.

[0350] The PCR amplified product was purified from an agarose gel, cut with EcoRI and SalI, ligated to pYX242 (Invitrogen Corp., Carlsbad, Calif.) (linearized with EcoRI and SalI) using the Rapid Ligation kit (Boehringer Mannheim Corp., Indianapolis, Ind.), according to the manufacturer's protocol and transformed into *E. coli* Top10 cells (Invitrogen Corp., Carlsbad, Calif.). The new plasmids, designated pRET-21 and pRET-22 (two individual clones from the ligation), were sequenced with the 373A Stretch DNA sequencer ABI (Perkin Elmer, Foster City, Calif.), and the cDNA sequences were identical. The 867 base cDNA nucleotide sequence of the plasmid pRET-22 containing the putative elongase is shown in FIG. 46 and the translated sequence of 288 amino acids is shown in FIG. 47. (Plasmid pRET-22 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209 on Aug. 19, 1999, under the terms of the Budapest Treaty and was accorded deposit number PTA-565.)

[0351] The plasmids pRET-21 and -22 were transformed into *S. cerevisiae* 334 as previously described (see Example III) and the resulting yeast cultures (334(pRET-21) and 334(pRET-22)) grown in 100 ml of selective media without leucine (Ausubel et al, supra) for 48 hours at 20° C. in the presence of 50 μ M GLA and AA. The cell pellets were collected and subjected to fatty acid analysis and the results shown in FIG. 48. DGLA, the predicted product from GLA elongation, was found to be an average of 1.79% of the total lipid in the two samples, versus 0.13% for the negative control (334 containing plasmid pYX242) indicating that the enzyme encoded by both pRET-21 and pRET-22 possessed GLA elongase activity. The percent conversion of GLA to DGLA by 334(pRET-21) and 334(pRET-22) was 11.1% and 19.4% respectively with an average of 15.25%. Interestingly, almost no elongation of AA or any endogenous fatty acid was observed (FIG. 48). These results indicate that the

elongase encoded by this newly identified *C. elegans* cDNA, CELO1, is able to specifically elongate GLA to DGLA, suggesting that it may be a *C. elegans* homologue of GLA elongase.

[0352] To further confirm the GLA elongation activity of CELO1, the experiment described in the paragraphs above was repeated with the exception that GLA and AA were added to cultures of 334(pRET-22) separately. Again, GLA was elongated to DGLA with a 38.2% conversion rate. No elongation activity of AA was detected as shown in FIG. 61. In this case, the percent conversion appears to be double that described in previous results (see FIG. 48) and may either be due to the absence of the additional substrate (AA) or the subculturing of the yeast. CELO1 has the additional activity of elongating endogenous 16:1n-7 to 18:1n-7 with a 9.12% conversion rate compared to 3.9% control culture 334(pYX242) under identical conditions. Thus, the *C. elegans* elongation enzyme possesses a major elongation activity for a C18 polyunsaturated fatty acid and a minor activity for a C16 monounsaturated fatty acid.

[0353] Additionally, to further determine the substrate specificity of CELO1, 50 μ M of each substrate besides GLA (e.g., SA (18:0), OA (18:1), LA (18:2n-6), DGLA (20:3n-6), AA (20:2n-6), ADA (22:4n-6), ALA (18:3n-3), PA (18:0), EPA (20:5n-3) and STA (18:4n-2)) was added individually to cultures of 334(pRET-22) and grown for 48 hours at 20° C., as described in Example XVII. STA was the only exogenously added substrate that was elongated. The CELO1 elongated 13% of STA incorporated to ETA (20:4n-3)(see FIG. 62).

[0354] Parallel to Examples III and XI, the *C. elegans* CELO1 gene in the plasmid pRET22 and *M. alpina* $\Delta 5$ desaturase (pCGR-4; see Example III) were co-expressed in yeast to determine if AA or EPA could be produced from exogenously added GLA or STA, respectively. When a yeast culture containing both pRET22 and pCGR4 plasmids was grown in the presence of 50 μ M GLA or STA in media lacking leucine and uracil, the percent conversion to the final products of AA and EPA, respectively, appeared identical (27% conversion) (see FIG. 63). Thus, simultaneous heterologous expression of CELO1 and a $\Delta 5$ desaturase results in the biosynthesis of AA and EPA from GLA and STA, respectively in yeast.

EXAMPLE XVIII

Isolation of a Putative Human Elongase cDNA
Based on AC004050 Sequence

[0355] To isolate the full length putative elongase cDNA based on the AC004050 sequence, primers RP735 (5'-CCT CCT GAA TTC CAQA CAC TAT TCA GCT TTC -3') and RO73 (5'-TAA TAC GAC TCA CTA TAG GG -3') were used to PCR amplify the human liver Marathon-Ready cDNA (Clontech Laboratories, Inc., Palo Alto, Calif.). The PCR was carried out using the Advantage™ cDNA PCR Kit (Clontech Laboratories, Inc., Palo Alto, Calif.) with 5 μ l of human liver Marathon-Ready cDNA and 50 pmole each primer following manufacturer's instructions. Thermocycler conditions in Perkin Elmer 9600 (Norwalk, Conn.) were as follows: 94° C. for 2 mins, then 30 cycles of 94° C. for 1 min., 58° C. for 2 mins., and 72° C. for 3 mins. PCR was followed by an additional extension at 72° C. for 7 mins.

[0356] The PCR amplified product was run on a gel, an amplified fragment of approximately 1 Kb was gel purified, the termini of the fragment were filled in with T4DNA polymerase (Boehringer Mannheim, Corp., Carlsbad, Calif.) following manufacturer's instructions. The new plasmid was designated as pRAE-59, and the putative PUFA elongase cDNA in this plasmid, designated as HS3, was sequenced using the ABI 373A Stretch Sequencer (Perkin Elmer, Foster City, Calif.). The putative PUFA elongase cDNA sequence HS3 is shown in FIG. 49, and the translated sequence is shown in FIG. 50.

EXAMPLE XIX

Cloning and Expression of a Mouse PUFA Elongation Enzyme

[0357] The National Center for Biotechnology Information (NCBI at <http://www.ncbi.nlm.nih.gov>) was used to conduct database searches using blastn with the mouse EST sequence AI225632 (see Example XIII). Three mouse EST sequences were identified (GenBank Accession #'s AI428130, AI595258, and AA061089), and assembled to generate a putative full-length elongation enzyme sequence, designated as MELO4. Primers RO819 (5'-ATG ATG CCA TGG AGC AGC TGA AGG CCT TTG-3') and RO820 (5'-CAG TCT CTG CTT TAA AAC AAG CTC GTC-3') were designed based on the putative full length mouse elongation enzyme sequence, and used to amplify the mouse brain Marathon-Ready cDNA (Clontech Laboratories, Inc., Palo Alto, Calif.). The Polymerase Chain Reaction (PCR) was carried out as previously described (Example XVI). The PCR amplified product was run on a gel, an amplified fragment of approximately 1,000 bp was gel purified, the termini of the fragment were digested with NcoI and DraI (Boehringer Mannheim, Corp., Indianapolis, Ind.), and the fragment was cloned into pYX242 (NcoI/HindIII). The new plasmid was designated as pRAE-84, and the putative PUFA elongation enzyme cDNA in this clone was sequenced using ABI 373A Stretch DNA Sequencer (Perkin Elmer, Foster City, Calif.). The putative PUFA elongation enzyme cDNA sequence in plasmid pRAE-84 is shown in FIG. 54, and the translated sequence is shown in FIG. 55.

[0358] The construct pRAE-84 was transformed into *S. cerevisiae* 334 (Hoveland et al., supra) and screened for elongase activity. The negative control strain was *S. cerevisiae* 334 containing pYX242 vector. The cultures were grown for 42-48 hours at 30° C., in selective media (Ausubel et al., supra), in the presence of 25 μ M of GLA, AA, ADA, STA, EPA, or DPA. The lipid profiles of these yeast cultures indicated that GLA was not elongated to the expected product of DGLA. However, there were accumulations of ADA, ω 6-tetracosatetraenoic acid (TTA, C24:4n-6), ETA, DPA, and ω 3-tetracosapentaenoic acid (TPA, C24:5n-3), respectively (FIG. 56). The n-6 fatty acid substrate AA was converted to ADA, which was subsequently converted to TTA, and the n-3 fatty acid EPA was converted to DPA, which was subsequently converted to TPA. The levels of these fatty acids were 0.64% (ADA), 1.07% (TTA), 1.47% (DPA), and 7.06% (TPA), respectively, of the total fatty acids in the strain containing the pRAE-84 sequence. These represented 10.4%, 62.6%, 32.7%, and 82.8% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. The C22 substrates ADA

and EPA were elongated to 2.4% (TTA) and 3.82% (TPA) of the total fatty acids. These represented 9.2% and 43.9% conversions of the substrate fatty acids, respectively. The expression of MELO4 in yeast results in the conversion of C20 and C22 fatty acids to the respective elongated products. The conversion rate of C22 to C24 fatty acids is much greater when the exogenously added substrate is C20 fatty acid.

[0359] To further confirm the substrate specificity of MELO4 protein (MELO4), the recombinant yeast strain 334(pRAE-84) was grown in minimal media containing 25 μ M of saturated, monounsaturated, or polyunsaturated fatty acids. The lipid profiles of these various substrates revealed that MELO4p is not involved in the elongation of saturated fatty acids such as PA, SA, APA, or BA (FIG. 57A). MELO4p is also not involved in the elongation of monounsaturated fatty acids PTA, OA, or EA. MELO4p is involved in the elongation of n-6 PUFAs AA and ADA, but not LA or DGLA (FIG. 57B). The lipid profiles of these yeast cultures indicated that there were accumulations of ADA and TTA, but not C20:2n-6 or C22:3n-6. When AA was added exogenously, the levels of product fatty acids were 0.5% (ADA) and 0.39% (TTA), and when ADA was added exogenously, the level of product fatty acid was 1.3% (TTA) of the total fatty acids in the strain containing the pRAE-84 sequence. These represented 8.7%, 43.8%, and 7.3% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. MELO4p is also involved in the elongation of GLA to DGLA. The lipid profile of the strain containing the pRAE-84 sequence, in presence of GLA, had 0.43% of DGLA, which represented 14.7% conversion of GLA to DGLA. MELO4p is also involved in the elongation of n-3 PUFAs EPA and DPA (FIG. 53C). The lipid profiles of these yeast cultures indicated that there were accumulations of DPA and TPA. When EPA was added, the levels of these fatty acids were 1.21% (DPA) and 3.38% (TPA), and when DPA was added, the level of the product fatty acid was 3.09% (TPA) of the total fatty acids in the strain containing the pRAE-84 sequence. These represented 24.0%, 73.6%, and 46.4% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. MELO4p is also involved in the elongation of STA to C22:4n-3. When STA was added, the levels of fatty acids produced by two-carbon elongation were 0.3% ETA and 0.23% C22:4n-3. These represented 11.1% and 43.4% conversions of substrate fatty acids to the products elongated by two carbon atoms. MELO4p also appeared to be involved in the elongation of ALA; however, the small amount of the fatty acid produced by two-carbon elongation (0.16% of ETrA) may not be significant. All results confirmed that the expression of MELO4 from mouse brain in yeast resulted in the elongation of C20 and C22 long-chain PUFAs in n-6 and n-3 fatty acid pathways.

EXAMPLE XX

Identification, Cloning, and Expression of HSELO1 Homologue from Mouse

[0360] The National Center for Biotechnology Information (NCBI at <http://www.ncbi.nlm.nih.gov>) was used to conduct database searches using blastn with the HSELO1 sequence. Two human EST sequences were identified (GenBank Accession #'s AI787925 and AI746838) and the

respective cDNA clones (I.M.A.G.E. Consortium Clone ID's 2076831 and 206182) were purchased through Research Genetics (Huntsville, Ala.). Primers RO833 (5'-GGT TTT ACC ATG GAA CAT TTC GAT GCG TCA C-3') and RO832 (5'-CGA CCT GCA GCT CGA GCA CA-3') were designed based on 5' sequence of the putative mouse elongation enzyme, and the cDNA clone vector, respectively. Primers RO833 and RO832 were used to amplify the mouse cDNA clone 2076182. The Polymerase Chain Reaction (PCR) was carried out as previously described (Example XVI). The termini of the PCR amplified product were filled-in with T4 DNA polymerase (Boehringer Mannheim, Corp., Indianapolis, Ind.) and the 5' region was digested with NcoI. The modified fragment was run on a gel, an amplified fragment of approximately 2.4 Kp was gel purified, and the fragment was cloned into pYX242 (NcoI/EcoRV). The new plasmid was designated as pRAE-87, and the putative PUFA elongation enzyme cDNA in this clone, MELO7, was sequenced using ABI 373A Stretch DNA Sequencer (Perkin Elmer, Foster City, Calif.). The putative PUFA elongation enzyme cDNA sequence in plasmid pRAE-87 (MELO7) is shown in **FIG. 58**, and the translated sequence is shown in **FIG. 59**.

[0361] The construct pRAE-87 was transformed into *S. cerevisiae* 334 (Hoveland et al., supra) and screened for elongase activity. The negative control strain was *S. cerevisiae* 334 containing pYX242 vector. The cultures were grown for 42-48 hours at 30° C., in selective media (Ausubel et al., supra), in the presence of 25 μ M of GLA, AA, STA, EPA, DPA, or ADA. The lipid profiles of the yeast cultures expressing MELO7 indicated that there were accumulations of DGLA, ADA, and ETA, respectively (**FIG. 60**). The levels of these fatty acids were 4.1% (DGLA), 6.33% (ADA), 3.4% (ETA), and 6.18% (DPA), respectively, of the total fatty acids in the strain containing the pRAE-87 sequence. These represented 78.7%, 36.0%, 81.0%, and 57.4% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. MELO7 protein (MELO7) was not involved in the elongation of ADA. MELO7p also appeared to be involved in further elongation of the fatty acid DPA produced by two-carbon elongation to TPA when EPA was the added substrate, and when DPA was added. However, the small amounts of the product fatty acids (0.27% and 0.25% of TPA) may not be significant. The yeast cells expressing the recombinant MELO7 sequence, compared to the control cells, also contained significantly elevated levels of C18:1n-7 and C20:1n-7. All results confirmed that the expression of MELO7 from mouse embryo in yeast resulted in the elongation of various long-chain PUFAs in n-6 and n-3 fatty acid pathways, and that MELO7 was a homologue of HSELO1.

EXAMPLE XXI

Cloning and Expression of a Fungal PUFA Elongation Enzyme

[0362] The translated amino acid sequences of the four elongases, HSELO1, MELO4, GLELO, and CELO, were aligned to identify areas of homology (**FIG. 64**). A primer was designed based on a block of sequences that showed similarities (underlined). Primer RO895 (5'-GTA GTA WGA GTA CAT GAT WAC GTG GAT RAA WGA GTT WAG-3') and vector primer RO898 (5'-CCC AGT CAC GAC GTT

GTA AAA CGA CGG CCA G-3') were used to amplify the cDNA from *Thraustochytrium aureum* 7091 (T7091). PCR was carried out in a 50 μ l volume containing: 1 μ l of T7091 cDNA, 0.2 μ M dNTP mix, 50 pM each primer, 5 μ l of 10 $\times$ buffer, 1.5 μ l of 50 mM MgSO₄, and 0.5 U of Taq DNA Polymerase. Thermocycler conditions in Perkin Elmer 9600 were as follows: 94° C. for 3 min, then 30 cycles of 95° C. for 45 sec., 55° C. for 30 sec., and 68° C. for 2 min. The PCR amplified mixture was run on a gel, an amplified fragment of approximately 750 bp was gel purified, and the isolated fragment was cloned into the pCR-blunt vector (Invitrogen, Co., Carlsbad, Calif.).

[0363] Twelve clones were prepared and sequenced. All twelve clones had the same sequence, and the consensus sequence was designated as cld6 (**FIG. 65**). The translated amino acid sequence of cld6 (**FIG. 66**) had 34.7% identity in 190 amino acids with HSELO1, 35.8% identity in 187 amino acids with MELO4, 45.6% identity in 160 amino acids with GLELO, and 32.9% identity in 155 amino acids with CELO. A new primer was designed based on the 5' sequence of cld6, at the first Met. RO1160 (5'-AAG GAA CCA TGG CAA ACA GCA GCG TGT GGG ATG-3'), which has an added NcoI site (underlined), and vector primer RO899 (5'-AGC GGA TAA CAA TTT CAC ACA GGAAAC AGC-3') were used to amplify the T7091 cDNA. The Polymerase Chain Reaction (PCR) was carried out as described above. The PCR amplified product was run on a gel, an amplified fragment of approximately 1.2 Kb was gel purified, the termini of the fragment were digested with NcoI and HindIII (Boehringer Mannheim, Corp., Indianapolis, Ind.), and the fragment was cloned into pYX242 (NcoI/HindIII). The new plasmids were designated as pRAT-4-A1, pRAT-4-A2, pRAT-4-A3, pRAT-4-A4, pRAT-4-A6, pRAT-4-A7, and pRAT-4-D1. (Plasmid DNA pRAT-4-A7 was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, on Apr. 19, 2001, under the terms of the Budapest Treaty, and was accorded accession number PTA-3301.) The putative PUFA elongation enzyme cDNAs in these clones were sequenced using ABI 373A Stretch DNA Sequencer (Perkin Elmer, Foster City, Calif.). The putative PUFA elongation enzyme cDNA sequences in plasmids pRAT-4-A1 through pRAT-4-D1 are shown in **FIGS. 67-73**, and the translated sequences are shown in **FIGS. 74-80**, respectively.

[0364] The constructs pPAT-4-A1 through pRAT-4-D1 were transformed into *S. cerevisiae* 334 (Hoveland et al., supra) and screened for elongase activity. *S. cerevisiae* 334 containing the unaltered pYX242 vector was used as a negative control. The cultures were grown for 48 hours at 24° C., in selective media (Ausubel et al., supra), in the presence of 25 μ M of GLA or EPA. In this study, DGLA or ω 3-docosapentaenoic acid (DPA, 22:5n-3), respectively, was the predicted product of the elongase activity. The lipid profiles of these yeast cultures indicated that GLA was elongated to DGLA in all the samples, except for 334(pRAT-4-A3) (**FIG. 81**). This was expected in that the cDNA sequence of pRAT-4-A3 had several stops at the 5' end of the sequence, which would result in a truncated version of the translated enzyme (**FIG. 76**). Although the DGLA level was very low for 334(pRAT-4-A2), the GLA level was the lowest in this sample; therefore, the conversion level of the fatty acid to its elongated form was significant at 28.4%. What was not expected was that there was definite substrate specificity among the expressed enzymes. The cultures of

334(pRAT-4-A1) and 334(pRAT-4-A2) had very low levels of DPA when EPA was added as a substrate, indicating that the expressed enzymes in these cultures preferred the elongation of the 18-carbon chain long PUFA, and not the 20-chain long PUFA, EPA. The cultures of 334(pRAT-4-A4), 334(pRAT-4-A6), 334(pRAT-4-A7), and 334(pRAT-D1) all had significant levels of DPA present, indicating that the expressed enzymes in these cultures were involved in the elongation of both 18- and 20-carbon chain long PUFAs to their respective elongated fatty acids.

[0365] The amino acid sequences of the 6 active clones were compared to determine if the substrate preferences were dictated by the translated sequences (**FIG. 82**). The cDNA sequences of pRAT-4-A1 and pRAT-4-A2 differed from other sequences in that they had a mutation at Y377C and V371A, respectively. This must be a critical region of the enzyme for 20-carbon chain elongation. The other sequences also had mutations, pRAT-4-A4 (I475V), pRAT-4-A6 (D26G and V458A), and pRAT-4-D1 (K182R and E269V); however, these mutations do not appear to interfere with 20-carbon chain elongation. The pRAT-4-A7 cDNA had a single, silent mutation at base 726 in comparison to the consensus sequence.

[0366] The translated amino acid sequence of the consensus TELO1 sequence had 34.0% identity in 265 amino acids with HSELO1 (**FIG. 83**), 34.1% identity in 267 amino acids with MELO4 (**FIG. 84**), 43.4% identity in 244 amino acids with GLELO (**FIG. 85**), and 34.3% identity in 239 amino acids with CELO (**FIG. 86**).

[0367] The yeast cells containing the fungal elongase cDNAs also had elevated levels of the monounsaturated fatty acid 18:1n-7, compared to the control strain (**FIG. 81**). Therefore, these results indicated that the identified fungal elongases are capable of utilizing PUFAs as well as monounsaturated fatty acids as substrates. Thus, these fungal sequences TELO1, and their encoded proteins (TELO1p), possess elongase activity independent of substrate specificity.

[0368] To further confirm the substrate specificity of the fungal elongation enzyme, described above and referred to herein as TELO1, the recombinant yeast strain 334(pRAT-4-A4), 334(pRAT-4-A6), 334(pRAT-4-A7), and 334(pRAT-4-D1) were grown in minimal media containing n-6 fatty acids GLA, AA, or n-3 fatty acids STA, or EPA, or no substrate. The lipid profiles of these yeast cultures, when examined by GC and GC-MS, indicated that there were accumulations of DGLA, ADA, ETA, ω 3-docosatetraenoic acid (DTA, C22:4n-3), and DPA, respectively (**FIG. 87**). The levels of these fatty acids were 3.88-5.42% (DGLA), 0.18-0.75% (ADA), 2.27-4.01% (ETA), 0.16-1.10% (DTA), and 0.54-1.74% (DPA), respectively, of the total fatty acids in the strains containing the TELO1 sequence. These represented 64.0-77.2%, 1.0-5.0%, 79.3-89.6%, 3.8-32.6%, and 3.4-13.3% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms.

[0369] The yeast cells expressing the recombinant TELO1 sequence, compared to the control cells, also contained significantly elevated levels of C18:1n-7 and decreased levels of C16:1n-7 (**FIG. 87**). This finding suggested that the recombinant TELO1 protein (TELOp) might also be involved in the elongation of monounsaturated fatty acids of 16-carbon length.

[0370] To further confirm the substrate specificity of TELOp, the recombinant yeast strain 334(pRAT-4-A7) was grown in minimal media containing 25 μ M of various PUFAs. The lipid profiles of these various substrates revealed that TELO1p is involved in the elongation of n-6 PUFAs LA, GLA, and AA, but not DGLA (**FIG. 88**). The lipid profiles of these yeast cultures indicated that there were accumulations of C20:2n-6, DGLA, and ADA, respectively, but not C22:3n-6. The levels of these fatty acids were 1.07% (C20:2n-6), 5.84% (DGLA), and 0.76% (ADA), respectively, of the total fatty acids in the lysates of 334(pRAT-4-A7). These represented 23.2%, 78.6%, and 3.5% conversions of the substrate fatty acids, respectively, to the products elongated by two-carbon atoms. TELO1p is also involved in the elongation of n-3 PUFAs ALA, STA, and EPA (**FIG. 88**). The lipid profiles of these yeast cultures indicated that there were accumulations of ETrA, ETA, and DPA, respectively. The levels of these fatty acids were 2.71% ETrA, 4.19% (ETA), and 1.99% (DPA), respectively, of the total fatty acids in the strain containing the TELO1 sequence. These represented 43.3%, 85.3%, and 11.5% conversions of the substrate fatty acids, respectively, to the products elongated by two-carbon atoms. When STA was added as a substrate, there was also an accumulation of DTA in the culture. The level of this fatty acid was 0.74%, which represented 15.0% conversion of ETA, the two-carbon chain elongated product from substrate STA. All results confirmed that the expression of TELO1 from T7091 in yeast resulted in the elongation of various long-chain PUFAs in n-6 and n-3 fatty acid pathways.

EXAMPLE XXII

Isolation of a Novel Elongase Gene from the Algae Pavlova sp. (CCMP459)

[0371] The fatty acid composition of the algae Pavlova sp. (CCMP 459) (Pav459) was investigated to determine the polyunsaturated fatty acids (PUFAs) produced by this organism. This algae showed a substantial amount of long chain PUFAs including eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3). Thus, this Pav459 was predicted to possess an elongase capable of converting EPA to ω 3-docosapentaenoic acid (DPA, 22:5n-3), which a Δ 4-desaturase can convert to DHA. The goal was therefore to isolate the predicted elongase gene from Pav459, and to verify the functionality of the enzyme by expression in an alternate host.

[0372] Frozen pellets of Pav459 were obtained from Provasoli-Guillard National Center for Culture of Marine Phytoplankton (CCMP, West Boothbay Harbor, Me.). These pellets were crushed in liquid nitrogen and total RNA was extracted from Pav459 by using the Qiagen RNeasy Maxi Kit (Qiagen, Valencia, Calif.), per manufacturers instructions. From this total RNA, mRNA was isolated using oligo dT cellulose resin, which was then used for the construction of a cDNA library using the pSport 1 vector (Invitrogen, Carlsbad, Calif.). The cDNA thus produced was directionally cloned (5'Sall/3'NotI) into pSport1 vector. The Pav459 library contained approximately 6.1×10^5 clones per ml, each with an average insert size of approximately 1200 bp. Two thousand five hundred primary clones from this library were sequenced from the 5' end using the T7 promoter primer (5'-TAA TAC GAC TCA CTA TTA GG-3'). Sequenc-

ing was carried out using the ABI BigDye sequencing kit (Applied Biosystems, Calif.) and the MegaBase Capillary DNA sequencer (Amersham biosciences, Piscataway, N.J.).

[0373] Two clones, designated 'pav06-C06' and pav07-G01,' which aligned to give a 500 bp sequence containing the 5' end of this novel elongase, were obtained from sequencing of the 2,500 library clones. This fragment shared 33.3% amino acid sequence identity with the mouse elongase MELO4 and 32.7% amino acid sequence identity with *T. aureum* elongase TELO1. To isolate the full-length gene, the EST clone pav06-C06 was used as a template for PCR reaction with 10 pmol of the 5' primer RO1327 (5'-TGC CCA TGA TGT TGG CCG CAG GCT ATC TTC TAG TG -3') and 10 pmol vector primer RO898 (5'-CCC AGT CAC GAC GTT GTA AAA CGA CGG CCA G-3'). PCR amplification was carried out using Platinum Taq DNA polymerase (Invitrogen, Carlsbad, Calif.) in a 50 μ l total volume containing: 1 μ l of the cDNA clone pav06-C06, PCR buffer containing 20 mM Tris-Cl, pH 8.4, 50 mM KCl (final concentration), 200 μ M each deoxyribonucleotide triphosphate, 10 pmole of each primer, 1.5 mM MgSO₄, and 0.5 μ l of Platinum Taq (HF) DNA polymerase. Amplification was carried out as follows using the Perkin Elmer 9700 machine: initial denaturation at 94° C. for 3 minute, followed by 35 cycles of the following: 94° C. for 45 sec, 55° C. for 30 sec, 68° C. for 2 min. The reaction was terminated at 4° C. The PCR amplified mixture was run on a gel, an amplified fragment of approximately 1.3 Kb was gel purified, and the isolated fragment was cloned into the pCR-blunt vector (Invitrogen, Carlsbad, Calif.). The recombinant plasmid was transformed into TOP10 supercompetent cells (Invitrogen, Carlsbad, Calif.), and prepared. The prepared recombinant plasmid was digested with EcoRI, run on a gel, and the digested fragment of approximately 1.2 Kb was gel purified, and cloned into pYX242 (EcoRI) vector (Novagen, Madison, Wis.). The new plasmid was designated as pRPL-6-1.

[0374] The plasmid pRPL-6-1 was prepared and sequenced using ABI 373A Stretch DNA Sequencer (Perkin Elmer, Foster City, Calif.). The putative elongation enzyme cDNA sequence in plasmid pRPL-6-1 is shown in FIG. 89, and the deduced amino acid sequence is shown in FIG. 90. The translated amino acid sequence of the cDNA in pRPL-6-1 had 33.7% identity in 261 amino acids with MELO4, 33.8% identity in 240 amino acids with GLELO, 28.1% identity in 274 amino acids with HSELO1, and 32.5% identity in 246 amino acids with TELO1.

[0375] The construct pRPL-6-1 was transformed into *S. cerevisiae* 334 (Hoveland et al., supra) and screened for elongase activity. *S. cerevisiae* 334 containing the unaltered pYX242 vector was used as a negative control. The cultures were grown for 44 hours at 24° C., in selective media (Ausubel et al., supra), in the presence of 25 μ M of GLA or EPA. In this study, DGLA or ω 3-docosapentaenoic acid (DPA, 22:5n-3), respectively, was the predicted product of the elongase activity. The lipid profiles of these yeast cultures indicated that while no conversion of GLA to DGLA was seen, EPA was elongated to DPA at a very low level (DPA was 0.34% of total fatty acids, while EPA was 32.28% of total fatty acids). This indicated that the expressed enzyme in this culture preferred the elongation of 20 carbon chain long PUFA, and not the 18 carbon chain long PUFA, GLA. It also indicated that a mutation might be

present in the DNA sequence, which is inhibiting the full activity of the expressed enzyme.

[0376] To isolate the full-length gene without mutations, RACE (rapid amplification of cDNA ends) ready cDNA was used as a target for the reaction. To prepare this material, approximately 5 μ g of total RNA was used according to the manufacturer's direction with the GeneRacer™ kit (Invitrogen, Carlsbad, Calif.) and Superscript II™ enzyme (Invitrogen, Carlsbad, Calif.) for reverse transcription to produce cDNA target. This cDNA was then used as a template for a PCR reaction with 50 pmols of the 5' primer RO1327 and 30 pmol GeneRacer™ 3' primer (5'-GCT GTC AAC GAT ACG CTA CGT AAC G-3'). PCR amplification was carried out using Platinum Taq DNA polymerase (Invitrogen, Carlsbad, Calif.) in a 50 μ l total volume containing: 2 μ l of the RACE ready cDNA, PCR buffer containing 20 mM Tris-Cl, pH 8.4, 50 mM KCl (final concentration), 200 μ M each deoxyribonucleotide triphosphate, 10 pmole of each primer, 1.5 mM MgSO₄, and 0.5 μ l of Platinum Taq (HF) DNA polymerase. Amplification was carried out as follows using the Perkin Elmer 9600 machine: initial denaturation at 94° C. for 3 minute, followed by 35 cycles of the following: 94° C. for 45 sec, 55° C. for 30 sec, 68° C. for 2 min. The reaction was terminated at 4° C.

[0377] The PCR amplified mixture was run on a gel, an amplified fragment of approximately 1.2 Kb was gel purified, and the isolated fragment was cloned into the PCR-blunt vector (Invitrogen, Carlsbad, Calif.). The recombinant plasmids were transformed into TOP10 supercompetent cells (Invitrogen, Carlsbad, Calif.), and prepared. The prepared recombinant plasmid was digested with EcoRI, run on a gel, and the digested fragment of approximately 1.2 Kb was gel purified, and cloned into pYX242 (EcoRI) vector (Novagen, Madison, Wis.). The new plasmids were designated as pRPL-6-B2 and pRPL-6-A3.

[0378] The plasmids pRPL-6-B2 and pRPL-6-A3 were prepared and sequenced using ABI 373A Stretch DNA Sequencer (Perkin Elmer, Foster City, Calif.). The putative elongation enzyme cDNA sequences in plasmids pRPL-6-B2 and pRPL-6-A3 are shown in FIGS. 91 and 92, and the deduced amino acid sequences are shown in FIGS. 93 and 94, respectively. The translated amino acid sequence of the cDNA in pRPL-6-B2 had 34.1% identity in 261 amino acids with MELO4 (FIG. 95), 33.8% identity in 240 amino acids with GLELO (FIG. 96), 28.5% identity in 274 amino acids with HSELO1 (FIG. 97), and 32.5% identity in 246 amino acids with TELO1 (FIG. 98). (Plasmid pRPL-6-B2 was deposited with the American Type Culture Collection, 10801 Manassas, Va. 20110-2209 under the terms of the Budapest Treaty and was accorded accession number PTA-4350.)

[0379] The constructs pRPL-6-B2 and pRPL-6-A3 were transformed into *S. cerevisiae* 334 (Hoveland et al., supra) and screened for elongase activity. *S. cerevisiae* 334 containing the unaltered pYX242 vector was used as a negative control. The cultures were grown for 44 hours at 24° C., in selective media (Ausubel et al., supra), in the presence of 25 μ M of GLA or EPA. In this study, DGLA or ω 3-docosapentaenoic acid (DPA, 22:5n-3), respectively, was the predicted product of the elongase activity. The lipid profiles of these yeast cultures indicated that GLA was not elongated to DGLA in any of the samples (data not shown). The cultures of 334(pRPL-6-B2) and 334(pRAT-6-A3) had significant

levels of conversion of the substrate EPA to DPA (FIG. 99), indicating that the expressed enzymes in these cultures preferred the elongation of 20-carbon chain long PUFA, and not the 18-chain long PUFA, GLA.

[0380] The amino acid sequences of the 3 clones were compared to determine if the substrate conversion levels were dictated by the translated sequences (FIG. 100). The cDNA sequence of pRPL-6-1 is different from pRPL-6-B2 at A512G. This single mutation dramatically reduced the conversion of the C20 substrate fatty acid to its elongated product. This must be a critical region of the enzyme for 20-carbon chain elongation. The cDNA sequence of pRPL-6-A3 is different from pRPL-6-B2 at D169N and C745R. These mutations reduced the conversion of the C20 substrate fatty acid to its elongated product, but the expressed enzyme was able to maintain some activity.

[0381] To further confirm the substrate specificity of the algal elongation enzyme, described above and referred to herein as PELOp, the recombinant yeast strain 334(pRPL-6-B2) was grown in minimal media containing n-6 fatty acids LA, GLA, DGLA, AA, or n-3 fatty acids ALA, STA, ETA, EPA, or 20:0, or 20:1. The lipid profiles of these yeast cultures, when examined by GC and GC-MS, indicated that there were accumulations of adrenic acid (ADA, 22:4-6) and EPA, respectively (FIG. 101). The levels of these fatty acids were 1.40% ADA and 2.54% EPA, respectively, of the total fatty acids in the strains containing the PELO1 sequence. These represented 14.0% and 14.1% conversions of the substrate fatty acids, respectively, to the products elongated by two carbon atoms. No elongation of the saturated fatty acid 20:0, or monounsaturated fatty acid 20:1 was seen. Also, no elongation of the C18 substrates LA, GLA, ALA, or STA was seen. These results indicated that the expressed enzyme activity in strain 334(pRPL-6-B2) was specific for the elongation of 20-carbon chain long PUFAs, and not the 18-chain long PUFA, or the 20-carbon chain long saturated or monounsaturated fatty acids.

Nutritional Compositions

[0382] The PUFAs described in the Detailed Description may be utilized in various nutritional supplements, infant formulations, nutritional substitutes and other nutritional solutions.

[0383] I. Infant Formulations

[0384] A. Isomil® Soy Formula with Iron:

[0385] Usage: As a beverage for infants, children and adults with an allergy or sensitivity to cows milk. A feeding for patients with disorders for which lactose should be avoided: lactase deficiency, lactose intolerance and galactosemia.

[0386] Features:

[0387] Soy protein isolate to avoid symptoms of cow's-milk-protein allergy or sensitivity.

[0388] Lactose-free formulation to avoid lactose-associated diarrhea.

[0389] Low osmolality (240 mOsm/kg water) to reduce risk of osmotic diarrhea.

[0390] Dual carbohydrates (corn syrup and sucrose) designed to enhance carbohydrate absorption and reduce the risk of exceeding the absorptive capacity of the damaged gut.

[0391] 1.8 mg of Iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.

[0392] Recommended levels of vitamins and minerals.

[0393] Vegetable oils to provide recommended levels of essential fatty acids.

[0394] Milk-white color, milk-like consistency and pleasant aroma.

[0395] Ingredients: (Pareve) 85% water, 4.9% corn syrup, 2.6% sugar (sucrose), 2.1% soy oil, 1.9% soy protein isolate, 1.4% coconut oil, 0.15% calcium citrate, 0.11% calcium phosphate tribasic, potassium citrate, potassium phosphate monobasic, potassium chloride, mono- and diglycerides, soy lecithin, carrageenan, ascorbic acid, L-methionine, magnesium chloride, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phyloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

[0396] B. Isomil® DF Soy Formula For Diarrhea:

[0397] Usage: As a short-term feeding for the dietary management of diarrhea in infants and toddlers.

[0398] Features:

[0399] First infant formula to contain added dietary fiber from soy fiber specifically for diarrhea management.

[0400] Clinically shown to reduce the duration of loose, watery stools during mild to severe diarrhea in infants.

[0401] Nutritionally complete to meet the nutritional needs of the infant.

[0402] Soy protein isolate with added L-methionine meets or exceeds an infant's requirement for all essential amino acids.

[0403] Lactose-free formulation to avoid lactose-associated diarrhea.

[0404] Low osmolality (240 mOsm/kg water) to reduce the risk of osmotic diarrhea.

[0405] Dual carbohydrates (corn syrup and sucrose) designed to enhance carbohydrate absorption and reduce the risk of exceeding the absorptive capacity of the damaged gut.

[0406] Meets or exceeds the vitamin and mineral levels recommended by the Committee on Nutrition of the American Academy of Pediatrics and required by the Infant Formula Act.

[0407] 1.8 mg of iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.

[0408] Vegetable oils to provide recommended levels of essential fatty acids.

[0409] Ingredients: (Pareve) 86% water, 4.8% corn syrup, 2.5% sugar (sucrose), 2.1% soy oil, 2.0% soy protein isolate,

1.4% coconut oil, 0.77% soy fiber, 0.12% calcium citrate, 0.11% calcium phosphate tribasic, 0.10% potassium citrate, potassium chloride, potassium phosphate monobasic, mono and diglycerides, soy lecithin, carrageenan, magnesium chloride, ascorbic acid, L-methionine, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phyloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

[0410] C. Isomil® SF Sucrose-Free Soy Formula With Iron:

[0411] Usage: As a beverage for infants, children and adults with an allergy or sensitivity to cow's-milk protein or an intolerance to sucrose. A feeding for patients with disorders for which lactose and sucrose should be avoided.

[0412] Features:

[0413] Soy protein isolate to avoid symptoms of cow's-milk-protein allergy or sensitivity.

[0414] Lactose-free formulation to avoid lactose-associated diarrhea (carbohydrate source is Polycose® Glucose Polymers).

[0415] Sucrose free for the patient who cannot tolerate sucrose.

[0416] Low osmolality (180 mOsm/kg water) to reduce risk of osmotic diarrhea.

[0417] 1.8 mg of iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.

[0418] Recommended levels of vitamins and minerals.

[0419] Vegetable oils to provide recommended levels of essential fatty acids.

[0420] Milk-white color, milk-like consistency and pleasant aroma.

[0421] Ingredients: (Pareve) 75% water, 11.8% hydrolyzed cornstarch, 4.1% soy oil, 4.1% soy protein isolate, 2.8% coconut oil, 1.0% modified cornstarch, 0.38% calcium phosphate tribasic, 0.17% potassium citrate, 0.13% potassium chloride, mono- and diglycerides, soy lecithin, magnesium chloride, ascorbic acid, L-methionine, calcium carbonate, sodium chloride, choline chloride, carrageenan, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phyloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

[0422] D. Isomil® 20 Soy Formula With Iron Ready To Feed, 20 Cal/fl oz.:

[0423] Usage: When a soy feeding is desired.

[0424] Ingredients: (Pareve) 85% water, 4.9% corn syrup, 2.6% sugar(sucrose), 2.1% soy oil, 1.9% soy protein isolate, 1.4% coconut oil, 0.15% calcium citrate, 0.11% calcium phosphate tribasic, potassium citrate, potassium phosphate monobasic, potassium chloride, mono- and diglycerides, soy

lecithin, carrageenan, ascorbic acid, L-methionine, magnesium chloride, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phyloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

[0425] E. Similac® Infant Formula:

[0426] Usage: When an infant formula is needed: if the decision is made to discontinue breastfeeding before age 1 year, if a supplement to breastfeeding is needed or as a routine feeding if breastfeeding is not adopted.

[0427] Features:

[0428] Protein of appropriate quality and quantity for good growth; heat-denatured, which reduces the risk of milk-associated enteric blood loss.

[0429] Fat from a blend of vegetable oils (doubly homogenized), providing essential linoleic acid that is easily absorbed.

[0430] Carbohydrate as lactose in proportion similar to that of human milk.

[0431] Low renal solute load to minimize stress on developing organs.

[0432] Powder, Concentrated Liquid and Ready To Feed forms.

[0433] Ingredients: (-D) Water, nonfat milk, lactose, soy oil, coconut oil, mono- and diglycerides, soy lecithin, ascorbic acid, carrageenan, choline chloride, taurine, m-inositol, alpha-tocopheryl acetate, zinc sulfate, niacinamide, ferrous sulfate, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, phyloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

[0434] F. Similac® NeoCare Premature Infant Formula With Iron:

[0435] Usage: For premature infants' special nutritional needs after hospital discharge. Similac NeoCare is a nutritionally complete formula developed to provide premature infants with extra calories, protein, vitamins and minerals needed to promote catch-up growth and support development.

[0436] Features:

[0437] Reduces the need for caloric and vitamin supplementation. More calories (22 Cal/fl oz) than standard term formulas (20 Cal/fl oz).

[0438] Highly absorbed fat blend, with medium-chain triglycerides (MCT oil) to help meet the special digestive needs of premature infants.

[0439] Higher levels of protein, vitamins and minerals per 100 calories to extend the nutritional support initiated in-hospital.

[0440] More calcium and phosphorus for improved bone mineralization.

[0441] Ingredients: -D Corn syrup solids, nonfat milk, lactose, whey protein concentrate, soy oil, high-oleic safflower oil, fractionated coconut oil (medium chain triglycerides), coconut oil, potassium citrate, calcium phosphate tribasic, calcium carbonate, ascorbic acid, magnesium chloride, potassium chloride, sodium chloride, taurine, ferrous sulfate, m-inositol, choline chloride, ascorbyl palmitate, L-carnitine, alpha-tocopheryl acetate, zinc sulfate, niacinamide, mixed tocopherols, sodium citrate, calcium pantothenate, cupric sulfate, thiamine chloride hydrochloride, vitamin A palmitate, beta carotene, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, phyloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

[0442] G. Similac Natural Care Low-Iron Human Milk Fortifier Ready To Use, 24 Cal/fl oz.:

[0443] Usage: Designed to be mixed with human milk or to be fed alternatively with human milk to low-birth-weight infants.

[0444] Ingredients: -D Water, nonfat milk, hydrolyzed cornstarch, lactose, fractionated coconut oil (medium-chain triglycerides), whey protein concentrate, soy oil, coconut oil, calcium phosphate tribasic, potassium citrate, magnesium chloride, sodium citrate, ascorbic acid, calcium carbonate, mono and diglycerides, soy lecithin, carrageenan, choline chloride, m-inositol, taurine, niacinamide, L-carnitine, alpha tocopheryl acetate, zinc sulfate, potassium chloride, calcium pantothenate, ferrous sulfate, cupric sulfate, riboflavin, vitamin A palmitate, thiamine chloride hydrochloride, pyridoxine hydrochloride, biotin, folic acid, manganese sulfate, phyloquinone, vitamin D3, sodium selenite and cyanocobalamin.

[0445] Various PUFAs of this invention can be substituted and/or added to the infant formulae described above and to other infant formulae known to those in the art.

[0446] II. Nutritional Formulations

[0447] A. Ensured®

[0448] Usage: ENSURE is a low-residue liquid food designed primarily as an oral nutritional supplement to be used with or between meals or, in appropriate amounts, as a meal replacement. ENSURE is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets. Although it is primarily an oral supplement, it can be fed by tube.

[0449] Patient Conditions:

- [0450] For patients on modified diets
- [0451] For elderly patients at nutrition risk
- [0452] For patients with involuntary weight loss
- [0453] For patients recovering from illness or surgery
- [0454] For patients who need a low-residue diet

[0455] Ingredients: -D Water, Sugar (Sucrose), Maltodextrin (Corn), Calcium and Sodium Caseinates, High-Oleic Safflower Oil, Soy Protein Isolate, Soy Oil, Canola Oil, Potassium Citrate, Calcium Phosphate Tribasic, Sodium Citrate, Magnesium Chloride, Magnesium Phosphate Dibasic, Artificial Flavor, Sodium Chloride, Soy Lecithin, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Gellan Gum,

Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Sodium Molybdate, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate.

[0456] B. Ensured® Bars:

[0457] Usage: ENSURE BARS are complete, balanced nutrition for supplemental use between or with meals. They provide a delicious, nutrient-rich alternative to other snacks. ENSURE BARS contain <1 g lactose/bar, and Chocolate Fudge Brownie flavor is gluten-free. (Honey Graham Crunch flavor contains gluten.)

[0458] Patient Conditions:

- [0459] For patients who need extra calories, protein, vitamins and minerals.
- [0460] Especially useful for people who do not take in enough calories and nutrients.
- [0461] For people who have the ability to chew and swallow
- [0462] Not to be used by anyone with a peanut allergy or any type of allergy to nuts.

[0463] Ingredients: Honey Graham Crunch—High-Fructose Corn Syrup, Soy Protein Isolate, Brown Sugar, Honey, Maltodextrin (Corn), Crisp Rice (Milled Rice, Sugar [Sucrose], Salt [Sodium Chloride] and Malt), Oat Bran, Partially Hydrogenated Cottonseed and Soy Oils, Soy Polysaccharide, Glycerine, Whey Protein Concentrate, Polydextrose, Fructose, Calcium Caseinate, Cocoa Powder, Artificial Flavors, Canola Oil, High-Oleic Safflower Oil, Nonfat Dry Milk, Whey Powder, Soy Lecithin and Corn Oil. Manufactured in a facility that processes nuts.

[0464] Vitamins and Minerals: Calcium Phosphate Tribasic, Potassium Phosphate Dibasic, Magnesium Oxide, Salt (Sodium Chloride), Potassium Chloride, Ascorbic Acid, Ferric Orthophosphate, Alpha-Tocopheryl Acetate, Niacinamide, Zinc Oxide, Calcium Pantothenate, Copper Gluconate, Manganese Sulfate, Riboflavin, Beta Carotene, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phyloquinone, Vitamin D3 and Cyanocobalamin.

[0465] Protein: Honey Graham Crunch—The protein source is a blend of soy protein isolate and milk proteins.

Soy protein isolate	74%
Milk proteins	26%

[0466] Fat: Honey Graham Crunch—The fat source is a blend of partially hydrogenated cottonseed and soybean, canola, high oleic safflower, oils, and soy lecithin.

Partially hydrogenated cottonseed and soybean oil	76%
Canola oil	8%
High-oleic safflower oil	8%
Corn oil	4%
Soy lecithin	4%

[0467] Carbohydrate: Honey Graham Crunch—The carbohydrate source is a combination of high-fructose corn syrup, brown sugar, maltodextrin, honey, crisp rice, glycerine, soy polysaccharide, and oat bran.

High-fructose corn syrup	24%
Brown sugar	21%
Maltodextrin	12%
Honey	11%
Crisp rice	9%
Glycerine	9%
Soy Polysaccharide	7%
Oat bran	7%

[0468] C. Ensure® High Protein:

[0469] Usage: Ensure HIGH PROTEIN is a concentrated, high-protein liquid food designed for people who require additional calories, protein, vitamins, and minerals in their diets. It can be used as an oral nutritional supplement with or between meals or, in appropriate amounts, as a meal replacement. ENSURE HIGH PROTEIN is lactose- and gluten-free, and is suitable for use by people recovering from general surgery or hip fractures and by patients at risk for pressure ulcers.

[0470] Patient Conditions:

[0471] For patients who require additional calories, protein, vitamins, and minerals, such as patients recovering from general surgery or hip fractures, patients at risk for pressure ulcers, and patients on low-cholesterol diets

[0472] Features:

[0473] Low in saturated fat

[0474] Contains 6 g of total fat and <5 mg of cholesterol per serving

[0475] Rich, creamy taste

[0476] Excellent source of protein, calcium, and other essential vitamins and minerals

[0477] For low-cholesterol diets

[0478] Lactose-free, easily digested

[0479] Ingredients: Vanilla Supreme: -D Water, Sugar (Sucrose), Maltodextrin (Corn), Calcium and Sodium Caseinates, High-Oleic Safflower Oil, Soy Protein Isolate, Soy Oil, Canola Oil, Potassium Citrate, Calcium Phosphate Tribasic, Sodium Citrate, Magnesium Chloride, Magnesium Phosphate Dibasic, Artificial Flavor, Sodium Chloride, Soy Lecithin, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Gellan Gum, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Sodium Molybdate, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

[0480] Protein:

[0481] The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	85%
Soy protein isolate	15%

[0482] Fat:

[0483] The fat source is a blend of three oils: high-oleic safflower, canola, and soy.

High-oleic safflower oil	40%
Canola oil	30%
Soy oil	30%

[0484] The level of fat in ENSURE HIGH PROTEIN meets American Heart Association (AHA) guidelines. The 6 grams of fat in ENSURE HIGH PROTEIN represent 24% of the total calories, with 2.6% of the fat being from saturated fatty acids and 7.9% from polyunsaturated fatty acids. These values are within the AHA guidelines of <30% of total calories from fat, <10% of the calories from saturated fatty acids, and <10% of total calories from polyunsaturated fatty acids.

[0485] Carbohydrate:

[0486] ENSURE HIGH PROTEIN contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla supreme, chocolate royal, wild berry, and banana), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla and other nonchocolate flavors:	
Sucrose	60%
Maltodextrin	40%
Chocolate:	
Sucrose	70%
Maltodextrin	30%

[0487] D. Ensure® Light

[0488] Usage: ENSURE LIGHT is a low-fat liquid food designed for use as an oral nutritional supplement with or between meals. ENSURE LIGHT is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

[0489] Patient Conditions:

[0490] For normal-weight or overweight patients who need extra nutrition in a supplement that contains 50% less fat and 20% fewer calories than Ensure.

[0491] For healthy adults who don't eat right and need extra nutrition.

[0492] Features:

[0493] Low in fat and saturated fat contains 3 g of total fat per serving and <5 mg cholesterol

- [0494] Rich, creamy taste
- [0495] Excellent source of calcium and other essential vitamins and minerals
- [0496] For low-cholesterol diets
- [0497] Lactose-free, easily digested

[0498] Ingredients:

[0499] French Vanilla: -D Water, Maltodextrin (Corn), Sugar (Sucrose), Calcium Caseinate, High-Oleic Safflower Oil, Canola Oil, Magnesium Chloride, Sodium Citrate, Potassium Citrate, Potassium Phosphate Dibasic, Magnesium Phosphate Dibasic, Natural and Artificial Flavor, Calcium Phosphate Tribasic, Cellulose Gel, Choline Chloride, Soy Lecithin, Carrageenan, Salt (Sodium Chloride), Ascorbic Acid, Cellulose Gum, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Pyridoxine Hydrochloride, Riboflavin, Chromium Chloride, Folic Acid, Sodium Molybdate, Biotin, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

[0500] Protein:

[0501] The protein source is calcium caseinate.

Calcium caseinate	100%
-------------------	------

[0502] Fat:

[0503] The fat source is a blend of two oils: high-oleic safflower and canola.

High-oleic safflower oil	70%
Canola oil	30%

[0504] The level of fat in ENSURE LIGHT meets American Heart Association (AHA) guidelines. The 3 grams of fat in ENSURE LIGHT represent 13.5% of the total calories, with 1.4% of the fat being from saturated fatty acids and 2.6% from polyunsaturated fatty acids. These values are within the AHA guidelines of <30% of total calories from fat, <10% of the, calories from saturated fatty acids, and <10% of total calories from polyunsaturated fatty acids.

[0505] Carbohydrate:

[0506] ENSURE LIGHT contains a combination of maltodextrin and sucrose. The chocolate flavor contains corn syrup as well. The mild sweetness and flavor variety (French vanilla, chocolate supreme, strawberry swirl), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

[0507] Vanilla and other nonchocolate flavors:

Sucrose	51%
Maltodextrin	49%

[0508] Chocolate:

Sucrose	47.0%
Corn Syrup	26.5%
Maltodextrin	26.5%

[0509] Vitamins and Minerals:

[0510] An 8-fl-oz serving of ENSURE LIGHT provides at least 25% of the RDIs for 24 key vitamins and minerals.

[0511] Caffeine:

[0512] Chocolate flavor contains 2.1 mg caffeine/8 fl oz.

[0513] E. Ensure Plus®

[0514] Usage: ENSURE PLUS is a high-calorie, low-residue liquid food for use when extra calories and nutrients, but a normal concentration of protein, are needed. It is designed primarily as an oral nutritional supplement to be used with or between meals or, in appropriate amounts, as a meal replacement. ENSURE PLUS is lactose- and gluten-free. Although it is primarily an oral nutritional supplement, it can be fed by tube.

[0515] Patient Conditions:

[0516] For patients who require extra calories and nutrients, but a normal concentration of protein, in a limited volume

[0517] For patients who need to gain or maintain healthy weight

[0518] Features:

[0519] Rich, creamy taste

[0520] Good source of essential vitamins and minerals

[0521] Ingredients:

[0522] Vanilla: -D Water, Corn Syrup, Maltodextrin (Corn), Corn Oil, Sodium and Calcium Caseinates, Sugar (Sucrose), Soy Protein Isolate, Magnesium Chloride, Potassium Citrate, Calcium Phosphate Tribasic, Soy Lecithin, Natural and Artificial Flavor, Sodium Citrate, Potassium Chloride, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenite, Phylloquinone, Cyanocobalamin and Vitamin D3.

[0523] Protein:

[0524] The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	84%
Soy protein isolate	16%

- [0525] Fat:
- [0526] The fat source is corn oil.

Corn oil	100%
----------	------

- [0527] Carbohydrate:
- [0528] ENSURE PLUS contains a combination of malto-dextrin and sucrose. The mild sweetness and flavor variety (vanilla, chocolate, strawberry, coffee, buffer pecan, and eggnog), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

- [0529] Vanilla, strawberry, butter pecan, and coffee fla-vors:

Corn Syrup	39%
Maltodextrin	38%
Sucrose	23%

- [0530] Chocolate and eggnog flavors:

Corn Syrup	36%
Maltodextrin	34%
Sucrose	30%

- [0531] Vitamins and Minerals:
- [0532] An 8-fl-oz serving of ENSURE PLUS provides at least 15% of the RDIs for 25 key Vitamins and minerals.
- [0533] Caffeine:
- [0534] Chocolate flavor contains 3.1 mg Caffeine/8 fl oz. Coffee flavor contains a trace amount of caffeine.

- [0535] F. Ensure Plus® HN

- [0536] Usage: ENSURE PLUS HN is a nutritionally com-plete high-calorie, high-nitrogen liquid food designed for people with higher calorie and protein needs or limited volume tolerance. It may be used for oral supplementation or for total nutritional support by tube. ENSURE PLUS HN is lactose- and gluten-free.

- [0537] Patient Conditions:
- [0538] For patients with increased calorie and protein needs, such as following surgery or injury.
- [0539] For patients with limited volume tolerance and early satiety.

- [0540] Features:
- [0541] For supplemental or total nutrition
- [0542] For oral or tube feeding
- [0543] 1.5 CaVmL,
- [0544] High nitrogen
- [0545] Calorically dense

- [0546] Ingredients:
- [0547] Vanilla: -D Water, Maltodextrin (Corn), Sodium and Calcium Caseinates, Corn Oil, Sugar (Sucrose), Soy Protein Isolate, Magnesium Chloride, Potassium Citrate, Calcium Phosphate Tribasic, Soy Lecithin, Natural and Artificial Flavor, Sodium Citrate, Choline Chloride, Ascor-bic Acid, Taurine, L-Carnitine, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Carrageenan, Cal-cium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochlo-ride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenite, Phylloquinone, Cyanocobalamin and Vita-min D3.

- [0548] G. Ensure® Powder:
- [0549] Usage: ENSURE POWDER (reconstituted with water) is a low-residue liquid food designed primarily as an oral nutritional supplement to be used with or between meals. Ensure POWDER is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

- [0550] Patient Conditions:
- [0551] For patients on modified diets
- [0552] For elderly patients at nutrition risk
- [0553] For patients recovering from illness/surgery
- [0554] For patients who need a low-residue diet

- [0555] Features:
- [0556] Convenient, easy to mix
- [0557] Low in saturated fat
- [0558] Contains 9 g of total fat and <5 mg of cho-lesterol per serving
- [0559] High in vitamins and minerals
- [0560] For low-cholesterol diets
- [0561] Lactose-free, easily digested

- [0562] Ingredients: -D Corn Syrup, Maltodextrin (Corn), Sugar (Sucrose), Corn Oil, Sodium and Calcium Caseinates, Soy Protein Isolate, Artificial Flavor, Potassium Citrate, Magnesium Chloride, Sodium Citrate, Calcium Phosphate Tribasic, Potassium Chloride, Soy Lecithin, Ascorbic Acid, Choline Chloride, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Chloride Hydrochloride, Cupric Sulfate, Pyridoxine Hydrochloride, Riboflavin, Vita-min A Palmitate, Folic Acid, Biotin, Sodium Molybdate, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

[0563] Protein:

[0564] The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	84%
Soy protein isolate	16%

[0565] Fat:

[0566] The fat source is corn oil.

Corn oil	100%
----------	------

[0567] Carbohydrate:

[0568] ENSURE POWDER contains a combination of corn syrup, maltodextrin, and sucrose. The mild sweetness of Ensure POWDER, plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, helps to prevent flavor fatigue and aid in patient compliance.

Vanilla:	
Corn Syrup	35%
Maltodextrin	35%
Sucrose	30%

[0569] H. Ensure® Pudding

[0570] Usage: ENSURE PUDDING is a nutrient-dense supplement providing balanced nutrition in a nonliquid form to be used with or between meals. It is appropriate for consistency-modified diets (e.g., soft, pureed, or full liquid) or for people with swallowing impairments. Ensure PUD-DING is gluten-free.

[0571] Patient Conditions:

[0572] For patients on consistency-modified diets (e.g., soft, pureed, or full liquid)

[0573] For patients with swallowing impairments

[0574] Features:

[0575] Rich and creamy, good taste

[0576] Good source of essential vitamins and minerals

[0577] Convenient-needs no refrigeration

[0578] Gluten-free

[0579] Nutrient Profile per 5 oz: Calories 250, Protein 10.9%, Total Fat 34.9%, Carbohydrate 54.2%

[0580] Ingredients:

[0581] Vanilla: -D Nonfat Milk, Water, Sugar (Sucrose), Partially Hydrogenated Soybean Oil, Modified Food Starch, Magnesium Sulfate, Sodium Stearoyl Lactylate, Sodium Phosphate Dibasic, Artificial Flavor, Ascorbic Acid, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Choline

Chloride, Niacinamide, Manganese Sulfate, Calcium Pan-tothenate, FD&C Yellow #5, Potassium Citrate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydro-chloride, Pyridoxine Hydrochloride, Riboflavin, FD&C Yel-low #6, Folic Acid, Biotin, Phylloquinone, Vitamin D3 and Cyanocobalamin.

[0582] Protein:

[0583] The protein source is nonfat milk. Nonfat milk 100%

[0584] Fat:

[0585] The fat source is hydrogenated soybean oil.

Hydrogenated soybean oil	100%
--------------------------	------

[0586] Carbohydrate:

[0587] ENSURE PUDDING contains a combination of sucrose and modified food starch. The mild sweetness and flavor variety (vanilla, chocolate, butterscotch, and tapioca) help prevent flavor fatigue. The product contains 9.2 grams of lactose per serving.

[0588] Vanilla and other nonchocolate flavors:

Sucrose	56%
Lactose	27%
Modified food starch	17%

[0589] Chocolate:

Sucrose	58%
Lactose	26%
Modified food starch	16%

[0590] I. Ensure® With Fiber:

[0591] Usage: ENSURE WITH FIBER is a fiber-contain-ing, nutritionally complete liquid food designed for people who can benefit from increased dietary fiber and nutrients. ENSURE WITH FIBER is suitable for people who do not require a low-residue diet. It can be fed orally or by tube, and can be used as a nutritional supplement to a regular diet or, in appropriate amounts, as a meal replacement. ENSURE WITH FIBER is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

[0592] Patient Conditions:

[0593] For patients who can benefit from increased dietary fiber and nutrients

[0594] Features:

[0595] New advanced formula-low in saturated fat, higher in vitamins and minerals

[0596] Contains 6 g of total fat and <5 mg of cho-lesterol per serving

[0597] Rich, creamy taste

[0598] Good source of fiber

- [0599] Excellent source of essential vitamins and minerals
- [0600] For low-cholesterol diets
- [0601] Lactose- and gluten-free
- [0602] Ingredients:
- [0603] Vanilla: -D Water; Maltodextrin (Corn), Sugar (Sucrose), Sodium and Calcium Caseinates, Oat Fiber, High-Oleic Safflower Oil, Canola Oil, Soy Protein Isolate, Corn Oil, Soy Fiber, Calcium Phosphate Tribasic, Magnesium Chloride, Potassium Citrate, Cellulose Gel, Soy Lecithin, Potassium Phosphate Dibasic, Sodium Citrate, Natural and Artificial Flavors, Choline Chloride, Magnesium Phosphate, Ascorbic Acid, Cellulose Gum, Potassium Chloride, Carrageenan, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Chromium Chloride, Biotin, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.
- [0604] Protein:
- [0605] The protein source is a blend of two high-biological-value proteins-casein and soy.
- [0606] Sodium and calcium caseinates 80%
- [0607] Soy protein isolate 20%
- [0608] Fat:
- [0609] The fat source is a blend of three oils: high-oleic safflower, canola, and corn.
- [0610] High-oleic safflower oil 40%
- [0611] Canola oil 40%
- [0612] Corn oil 20%
- [0613] The level of fat in ENSURE WITH FIBER meets American Heart Association (AHA) guidelines. The 6 grams of fat in ENSURE WITH FIBER represent 22% of the total calories, with 2.01% of the fat being from saturated fatty acids and 6.7% from polyunsaturated fatty acids. These values are within the AHA guidelines of $\leq 30\%$ of total calories from fat, $<10\%$ of the calories from saturated fatty acids, and $\leq 10\%$ of total calories from polyunsaturated fatty acids.
- [0614] Carbohydrate:
- [0615] ENSURE WITH FIBER contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla, chocolate, and butter pecan), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.
- [0616] Vanilla and other nonchocolate flavors:

Maltodextrin	66%
Sucrose	25%
Oat Fiber	7%
Soy Fiber	2%

[0617] Chocolate:

Maltodextrin	55%
Sucrose	36%
Oat Fiber	7%
Soy Fiber	2%

- [0618] Fiber:
- [0619] The fiber blend used in ENSURE WITH FIBER consists of oat fiber and soy polysaccharide. This blend results in approximately 4 grams of total dietary fiber per 8-fl. oz can. The ratio of insoluble to soluble fiber is 95:5.
- [0620] The various nutritional supplements described above and known to others of skill in the art can be substituted and/or supplemented with the PUFAs produced in accordance with the present invention.
- [0621] J. Oxepa™ Nutritional Product
- [0622] Oxepa is a low-carbohydrate, calorically dense, enteral nutritional product designed for the dietary management of patients with or at risk for ARDS. It has a unique combination of ingredients, including a patented oil blend containing eicosapentaenoic acid (EPA from fish oil), α -linolenic acid (GLA from borage oil), and elevated antioxidant levels.
- [0623] Caloric Distribution:
- [0624] Caloric density is high at 1.5 Cal/mL (355 Cal/8 fl oz), to minimize the volume required to meet energy needs.
- [0625] The distribution of Calories in Oxepa is shown in Table IV.

TABLE IV			
Caloric Distribution of Oxepa			
	per 8 fl oz.	per liter	% of Cal
Calories	355	1,500	—
Fat (g)	22.2	93.7	55.2
Carbohydrate (g)	25	105.5	28.1
Protein (g)	14.8	62.5	16.7
Water (g)	186	785	—

- [0626] Fat:
- [0627] Oxepa contains 22.2 g of fat per 8-fl oz serving (93.7 g/L).
- [0628] The fat source is an oil blend of 31.8% canola oil, 25% medium-chain triglycerides (MCTs), 20% borage oil, 20% fish oil, and 3.2% soy lecithin. The typical fatty acid profile of Oxepa is shown in Table V.
- [0629] Oxepa provides a balanced amount of polyunsaturated, monounsaturated, and saturated fatty acids, as shown in Table VI.
- [0630] Medium-chain triglycerides (MCTs)—25% of the fat blend—aid gastric emptying because they are absorbed by the intestinal tract without emulsification by bile acids.

[0631] The various fatty acid components of Oxepa™ nutritional product can be substituted and/or supplemented with the PUFAs produced in accordance with this invention.

TABLE V

Typical Fatty Acid Profile			
	% Total Fatty Acids	g/8 fl oz*	g/L*
Caproic (6:0)	0.2	0.04	0.18
Caprylic (8:0)	14.69	3.1	13.07
Capric (10:0)	11.06	2.33	9.87
Palmitic (16:0)	5.59	1.18	4.98
Palmitoleic	1.82	0.38	1.62
Stearic	1.94	0.39	1.64
Oleic	24.44	5.16	21.75
Linoleic	16.28	3.44	14.49
γ-Linolenic	3.47	0.73	3.09
α-Linolenic	4.82	1.02	4.29
Eicosapentaenoic	5.11	1.08	4.55
n-3-Docosapent- aenoic	0.55	0.12	0.49
Docosahexaenoic	2.27	0.48	2.02
Others	7.55	1.52	6.72

Fatty acids equal approximately 95% of total fat.

[0632]

TABLE VI

Fat Profile of Oxepa.	
% of total calories from fat	55.2
Polyunsaturated fatty acids	31.44 g/L
Monounsaturated fatty acids	25.53 g/L
Saturated fatty acids	32.38 g/L
n-6 to n-3 ratio	1.75:1
Cholesterol	9.49 mg/8 fl oz
	40.1 mg/L

[0633] Carbohydrate:

[0634] The carbohydrate content is 25.0 g per 8-fl-oz serving (105.5 g/L).

[0635] The carbohydrate sources are 45% maltodextrin (a complex carbohydrate) and 55% sucrose (a simple sugar), both of which are readily digested and absorbed.

[0636] The high-fat and low-carbohydrate content of Oxepa is designed to minimize carbon dioxide (CO2) production. High CO2 levels can complicate weaning in ventilator-dependent patients. The low level of carbohydrate also may be useful for those patients who have developed stress-induced hyperglycemia.

[0637] Oxepa is lactose-free.

[0638] Dietary carbohydrate, the amino acids from protein, and the glycerol moiety of fats can be converted to glucose within the body. Throughout this process, the carbohydrate requirements of glucose-dependent tissues (such as the central nervous system and red blood cells) are met. However, a diet free of carbohydrates can lead to ketosis, excessive catabolism of tissue protein, and loss of fluid and electrolytes. These effects can be prevented by daily ingestion of 50 to 100 g of digestible carbohydrate, if caloric

intake is adequate. The carbohydrate level in Oxepa is also sufficient to minimize gluconeogenesis, if energy needs are being met.

[0639] Protein:

[0640] Oxepa contains 14.8 g of protein per 8-fl-oz serving (62.5 g/L).

[0641] The total calorie/nitrogen ratio (150:1) meets the need of stressed patients.

[0642] Oxepa provides enough protein to promote anabolism and the maintenance of lean body mass without precipitating respiratory problems. High protein intakes are a concern in patients with respiratory insufficiency. Although protein has little effect on CO2 production, a high protein diet will increase ventilatory drive.

[0643] The protein sources of Oxepa are 86.8% sodium caseinate and 13.2% calcium caseinate.

[0644] The amino acid profile of the protein system in Oxepa meets or surpasses the standard for high quality protein set by the National Academy of Sciences.

*Oxepa is gluten-free.

Default settings for the analysis programs			
<u>GCG Programs</u>			
FastA Search			
<u>Default parameters:</u>			
Range of interest	Begin = 1	END = last protein or nucleic acid	
Search set	all of SwissProt (protein) or GenEMBL (nucleic acid)		
Word size	= (2) for protein = (6) for nucleic acid		
Expected scores	lists scores until E () value reaches 2.0		
TFastA search			
<u>Default parameters:</u>			
range of interest	Begin = 1	END = last nucleic acid	
Search set	all of GenEMBL		
Word size	wordsize = (2)		
Expected scores	lists scores until E() value reaches 2.0		
Pileup			
<u>Default parameters:</u>			
gap creation penalty	gap weight = 5		
gap extension penalty	gap length weight = 12		
plot figure	one page plot density = 2.7		
<u>Sequencher Program</u>			
<u>Default parameters:</u>			
Automatic Assembly	Dirty data algorithm = slower contig assembly but more rigorous comparisons between the sequences		
	minimum match = 85%		
	minimum overlap = 20		
<u>BLAST 2 (blastp, tblastn)</u>			
Default parameters:	V = 50	Lambda = .329	W = 3
	B = 50	K = 0.140	X = 22
	E = 10	H = 0.427	

-continued

Default settings for the analysis programs

blast n

Default parameters:

V = 100

Lambda = 1.37

W = 11

B = 250

K = 0.171

X1 = 22

E = 10

H = 1.31

X2 = 25

8/25 BLAST 2 Command Line Arguments

-v Hits

number of best scores to show

-b Alignments

number of best alignments to show

-e Expectation value (E)

[Real] default = 10.0

-m Alignment view options:

0 = pairwise,

1 = master-slave showing identities,

2 = master-slave, no identities,

3 = flat master-slave, show identities,

4 = flat master-slave, no identities,

5 = master-slave, no identities and blunt ends,

6 = flat master-slave, no identities and blunt ends [Integer]

default = 0

-F Filter query seq. (DUST with blastn, SEG with others)

[T/F]

default = T

-G Cost to open a gap (zero invokes default behavior)

[Integer]

default = 0

-E Cost to extend a gap (zero invokes default behavior)

[Integer]

default = 0

-X X dropoff value for gapped alignment (in bits) (zero invokes default behavior)

[Integer]

default = 0

-I Show GI's in defines

[T/F]

default = F

-q Penalty for a nucleotide mismatch (blastn only)

[Integer]

default = -3

-r Reward for a nucleotide match (blastn only)

[Integer]

default = 1

-f Threshold for extending hits default if zero

[Integer]

default = 0

-g Perform gapped alignment (not available with tblastx)

[T/F]

default = T

-q Query Genetic code to use

[Integer]

default = 1

-D DB Genetic code (for tblast [nx] only)

[Integer]

default = 1

-J Believe the query define

[T/F]

-continued

Default settings for the analysis programs												
default = F												
-M Matrix [String]												
default = BLOSUM62												
-W Word size default if zero												
[Integer]												
default = 0												
-z Effective length of the database (use zero for the real size) [Integer]												
default = 0												
-a Number of processors to use [Integer]												
default = site configurable												
(SeqServer.conf)												
Allowed and default values for gap open/gap extension cost (-G/-E) parameters:												
BLOSUM62												
-G	9	8	7	12	11	10						
-E	2	2	2	1	1	1						
BLOSUM50												
-G	12	11	10	9	15	14	13	12	18	17	16	15
-E	3	3	3	3	2	2	2	2	1	1	1	1
PAM250												
-G	13	12	11	10	15	14	13	12	19	18	17	16
-E	3	3	3	3	2	2	2	2	1	1	1	1
BLOSUM90												
-G	8	7	6	11	10	9						
-E	2	2	2	1	1	1						
PAM30												
-G	5	4	3	7	6	5	10	9	8			
-E	3	3	3	2	2	2	1	1	1			
PAM70												
-G	6	5	4	8	7	6	11	10	9			
-E	3	3	3	2	2	2	1	1	1			

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 122

<210> SEQ ID NO 1

<211> LENGTH: 954

<212> TYPE: DNA

<213> ORGANISM: Mortierella alpina

-continued

<400> SEQUENCE: 1

```
atggccgccc caatcttga caaggtcaac ttcggcattg atcagccctt cggaatcaag    60
ctcgacacct actttgtctc gccctatgaa ctcgtcaccg gaaagtccat cgactccttc    120
gtcttcacagg agggcgctac gcctctctcg acccagagag aggtcgccat gtggactatc    180
acttacttcg tcgtcatctt tgggtggtcg cagatcatga agagccagga cgccttcaag    240
ctcaagcccc tcttcacctt ccacaacttc ctctgacga tcgctgcgg atcgtgttg    300
ctctgttca tcgagaacct ggtcccatc ctgccagaa acggactttt ctacgccatc    360
tgcgacgacg gtgcctggac ccagcgctc gagctcctct actacctcaa ctacctggtc    420
aagtactggg agttggccga caccgtcttt ttggtcctca agaagaagcc tcttgagttc    480
ctgcactact tccaccactc gatgacctg gttctctgct ttgtccagct tggaggatac    540
atttcagtgt cctgggtccc tattaccctc aacttgactg tccacgtctt catgtactac    600
tactacatgc gtcctcgctc cgggtgttcg atctggtgga agcagtactt gaccactctc    660
cagatcgctc agttcgctct tgacctcgga ttcactact tctgcgcta cactacttc    720
gccttcacct acttcccctg ggctcccaac gtcggcaagt gcgccgttac cgaggggtgt    780
gctctctttg gctgcggact cctctccagc tatctcttgc tctttatcaa cttctaccgc    840
attacctaca atgccaaagg caaggcagcc aaggagcgtg gaagcaactt taccaccaag    900
actgtcaagt ccggcggatc gcccagaag ccctccaaga gcaagcacat ctaa    954
```

<210> SEQ ID NO 2

<211> LENGTH: 957

<212> TYPE: DNA

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 2

```
atggagtcca ttgcgcatt cctcccata aagatgccgc aagatctgtt tatggacctt    60
gccaccgcta tcggtgtccg ggccgcgccc tatgtcgatc ctctcgaggc cgcgtgttg    120
gcccaggccg agaagtacat ccccacgatt gtccatcaca cgcgtgggtt cctggtcgcg    180
gtggagtgcg ctttggtccc tgagctgccg ttgatgaacc cgttccactg gctgttgatc    240
gtgctcgctt atttggtcac ggtctttgtg ggcatgcaga tcatgaagaa ctttgagcgg    300
ttcgaggtea agacgttttc gtcctgcac aacttttgtc tggctctgat cagcgcttac    360
atgtgcggtg ggatcctgta cgaggcttat caggccaact atggactgtt tgagaacgct    420
gctgatcata ccttcaaggg tcttcctatg gccaaagatg tctggctctt ctacttctcc    480
aagatcatgg agtttgtcga caccatgatc atggtcctca agaagaacaa ccgccagatc    540
tccttcttgc acgtttacca ccacagctcc atcttcacca tctgggtggtt ggtcaccttt    600
gttgaccaca acggtgaagc ctacttctct gctgcgttga actcgttcat ccatgtgatc    660
atgtacggct actacttctt gtcggccttg ggcttcaagc aggtgtcgtt catcaagttc    720
tacatcacgc gtcgcgagat gacacagtto tgcagtatgt cgttcagtc ttctgtggac    780
atgtacgcca tgaaggctct tggccgcccc ggataccctt tcttcacac ggctctgctt    840
tggttctaca tgtggacctg gctcgtctc ttctacaact ttacagaaa gaacgccaag    900
ttggccaagc aggccaaggc cgacgtctcc aaggagaagg caaggaagtt gcagtaa    957
```

<210> SEQ ID NO 3

-continued

<211> LENGTH: 914

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```
atggaacatt ttgatgcac accatgtacc tatttcaagg cattgctagg ccctcgagat    60
actagagtaa aaggatgggt tcttctggac aattatatac ccacatttat ctgctctgtc    120
atatatttac taattgtatg gctgggacca aaatacatga ggaataaaca gccatttctt    180
tgccggggga ttttagtggt gtataacctt ggactcacac tgctgtctct gtatatgttc    240
tgtgagttag taacaggagt atgggaaggc aaatacaact tcttctgtca gggcacacgc    300
accgcaggag aatcagatat gaagattatc cgtgtcctct ggtggtacta cttctccaaa    360
ctcatagaa ttatggacac tttcttcttc atcctgcgca agaacaacca ccagatcacg    420
gtcctgcacg tctaccacca tgctcgtatg ctgaacatct ggtggtttgt gatgaactgg    480
gtccctcgcg gccactctta ttttggtgcc acacttaata gcttcatcca cgtcctcatg    540
tactcttact atggtttgtc gtcagtcctt tccatgcgtc catacctctg gtggaagaag    600
tacatcactc aggggcagct gcttcagttt gtgctgacaa tcaccagac cagctgcggg    660
gtcatctggc cgtgcacatt ccctcttggt tggttgtatt tccagattgg atacattatt    720
tccctgattg ctctcttcac aaacttctac attcagacct acaacaagaa aggggcctcc    780
cgaaggaaag accacctgaa ggaccaccag aatgggtccg tggctgctgt gaatggacac    840
accaacagct tttcacccct ggaaaacaat gtgaagccaa ggaagctgcg gaaggattga    900
agtcaaagaa ttga                                         914
```

<210> SEQ ID NO 4

<211> LENGTH: 867

<212> TYPE: DNA

<213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 4

```
atggctcagc atccgctcgt tcaacggctt ctcatgtca aattcgacac gaaacgattt    60
gtggctattg ctactcatgg gccaaagaat ttccctgacg cagaaggctg caagttcttt    120
gctgatcact ttgatgttac tattcaggct tcaatcctgt acatggctgt tgtgttcgga    180
acaaaatggg tcatgcgtaa tcgtcaacca ttccaattga ctattccact caacatctgg    240
aatttcatcc tcgcccgcatt ttccatcgca ggagctgtca aaatgacccc agagttcttt    300
ggaaccattg ccaacaaagg aattgtcgca tcctactgca aagtgtttga tttcacgaaa    360
ggagagaatg gatactgggt gtggctcttc atggcttcca aacttttcga acttgttgac    420
accatctctt tggttctccg taaacgtcca ctcatgttcc ttactggta tcaccatatt    480
ctcaccatga tctacgcctg gtactctcat ccattgacct caggattcaa cagatacggg    540
atztatctta actttgtcgt ccacgccttc atgtactctt actacttctt tcgctcgatg    600
aagattcgcg tgccaggatt catcgcccaa gctatcacat ctcttcaa at cgttcaattc    660
atcatctctt gcgcgttctt tgctcatctt gggtatctca tgcacttcac caatgcaaac    720
tgtgatttgc agccatcagt attcaagctc gcagttttca tggacacaa atacttggtc    780
cttttgcgta acttcttctt ccaatcatat gttctccgcg gaggaaaaga caagtacaag    840
gcagtgccaa agaagaagaa caactaa                                         867
```

-continued

<210> SEQ ID NO 5
<211> LENGTH: 879
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

atggagcagc tgaaggcctt tgataatgaa gtcaatgctt tcttgacaa catgtttgga	60
ccacgagatt ctcgagttcg cgggtggttc ctgctggact cttaccttc caccctcatc	120
ctcaccatca cgtacctgct ctcgatatgg ctgggtaaca agtacatgaa gaacaggcct	180
gtctgtgtctc tcaggggcat cctcaccttg tataacctcg caatcacact tctttctgcg	240
tatatgtctgg tggagctcat cctctccagc tgggaaggag gttacaactt gcagtgtcag	300
aatctcgaca gtgcaggaga aggtgatgtc cgggtagcca aggtcttctg gtggtactac	360
ttctccaaac tagtggagtt cctggacagc attttctttg ttctacgaaa aaagaccaat	420
cagatcacct tccttcatgt ctatcaccac gcgtccatgt tcaacatctg gtggtgtgtt	480
ttgaactgga taccttctgg tcaaagcttc tttggacca cctgaacag ctttatccac	540
attctcatgt actcctacta cggcctgtct gtgttccgt ccatgcacaa gtacctttgg	600
tggaagaagt acctcacaca ggctcagctg gtgcagttcg tactcaccat cagcacacg	660
ctgagtgcgc tggtagaagg ctgtggcttc cctttggct gtctcatctt ccagtcttcc	720
tatatgatga cgctggtcat cctgttctta aacttctata ttcagacata ccggaaaaag	780
ccagtgaaga aagagctgca agagaaagaa gtgaagaatg gtttccccaa agcccactta	840
attgtggcta atggcatgac ggacaagaag gotcaataa	879

<210> SEQ ID NO 6
<211> LENGTH: 900
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

atggaacatt tcgatgcgtc actcagtacc tatttcaagg ccttcctggg ccccgagat	60
acaagagtoa aaggatgggt cctcctggac aattacatcc ctacgtttgt ctgttctgtt	120
atttacttac tcattgtatg gctgggacca aaatacatga agaaccggca gccgttctct	180
tgccgaggca tcctgcagtt gtataacctt ggactcaccg tgctgtctct ctacatgttc	240
tatgagttgg tgacagggtg gtgggagggc aaatacaact ttttctgcca gggaacacgc	300
agcgcgggag aatccgatat gaagatcatc cgcgtcctct ggtggtacta cttctccaaa	360
ctcatcgaat tcatggacac ctttttcttc atccttcgca agaacaacca ccagatcacc	420
gtgtccatg tctaccacca cgctaccatg ctcaacatct ggtggtttgt gatgaactgg	480
gttccctgcg gccattcata ttttggtgcg aactcaaca gcttcatcca tgcctcatg	540
tactcgtact atggtctgtc ctccatcccg tccatgcgtc cctacctctg gtggaaaaag	600
tacatcactc aagggcagct ggtccagttt gtgctgacaa tcatccagac gacctgcggg	660
gtcttctggc catgctcctt ccctctcggt tggctgttct tccagattgg atacatgatt	720
tccctgattg ctctcttcac aaacttctac attcagactt acaacaagaa aggggcctct	780
cggaggaaag accacctgaa gggccaccag aacgggtctg tggccgccgt caacggacac	840
accaacagct tcccttcctt ggaaaacagc gtgaagccca ggaagcagcg aaaggattga	900

-continued

<210> SEQ ID NO 7
<211> LENGTH: 819
<212> TYPE: DNA
<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 7

```
atggcaaacac gcagcgtgtg ggatgatgtg gtgggccgcg tggagaccgg cgtggaccag      60
tggatggatg gcgccaagcc gtacgcactc accgatgggc tcccgatgat ggacgtgtcc      120
accatgctgg cattcgaggt gggtacatg gccatgtctc tcttcggcat cccgatcatg      180
aagcagatgg agaagccttt tgagctcaag accatcaagc tcttcacaaa cttgtttctc      240
ttcggacttt ccttgtacat gtgcgtggag accatccgcc aggctatcct cggaggctac      300
aaagtgtttg gaaacgacat ggagaaggcg aacgagtctc atgctcaggg catgtctcgc      360
atcgtgtacg tgttctactg gtccaaggca tacgagttct tggataccgc catcatgac      420
ctttgcaaga agttcaacca ggtttccttc ttgcatgtgt accacatgc caccattttt      480
gccatctggt gggctatcgc caagtacgct ccaggagggtg atgcgtactt ttcagtgatc      540
ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctctc ccaagggttc      600
gggttcgtga agccaatcaa gccgtacatc accacccttc agatgacca gttcatggca      660
atgcttgtgc agtccttgta cgactacctc ttcccatgcg actaccaca ggctcttggt      720
cagctccttg gagtgtacat gatcaccttg cttgccctct tcggcaactt ttttgtgcag      780
agctatctta aaaagccaaa aaagagcaag accaactaa      819
```

<210> SEQ ID NO 8
<211> LENGTH: 83
<212> TYPE: PRT
<213> ORGANISM: *Simmondsia chinensis*

<400> SEQUENCE: 8

```
Ala Thr Leu Pro Asn Phe Lys Ser Ser Ile Asn Leu His His Val Lys
 1             5             10             15
Leu Gly Tyr His Tyr Leu Ile Ser Asn Ala Leu Phe Leu Val Phe Ile
          20             25             30
Pro Leu Leu Gly Leu Ala Ser Ala His Leu Ser Ser Phe Ser Ala His
          35             40             45
Asp Leu Ser Leu Leu Phe Asp Leu Leu Arg Arg Asn Leu Leu Pro Val
          50             55             60
Val Val Cys Ser Phe Leu Phe Val Leu Leu Ala Thr Leu His Phe Leu
          65             70             75             80
Thr Arg Pro
```

<210> SEQ ID NO 9
<211> LENGTH: 80
<212> TYPE: PRT
<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 9

```
Ser Thr Leu Pro Pro Val Leu Tyr Ala Ile Thr Ala Tyr Tyr Val Ile
 1             5             10             15
Ile Phe Gly Gly Arg Phe Leu Leu Ser Lys Ser Lys Pro Phe Lys Leu
          20             25             30
Asn Gly Leu Phe Gln Leu His Asn Leu Val Leu Thr Ser Leu Ser Leu
```

-continued

35	40	45	
Thr Leu Leu Leu Leu Met Val Glu Gln Leu Val Pro Ile Ile Val Gln			
50	55	60	
His Gly Leu Tyr Phe Ala Ile Cys Asn Ile Gly Ala Trp Thr Gln Pro			
65	70	75	80
<210> SEQ ID NO 10			
<211> LENGTH: 240			
<212> TYPE: DNA			
<213> ORGANISM: <i>Saccharomyces cerevisiae</i>			
<400> SEQUENCE: 10			
tccaccctcc ccccgctcct ctacgccatc accgcctact acgtcatcat ctcggtggt			60
cgcttcctcc tctccaagtc caagcccttc aagctcaacg gtctcttcca gctccacaac			120
ctcgtcctca cctccctctc cctcaccctc ctctcctca tggtcgagca gctcgtcccc			180
atcatcgtcc agcacggtct ctacttcgcc atctgcaaca tcggtgcctg gaccagccc			240
<210> SEQ ID NO 11			
<211> LENGTH: 33			
<212> TYPE: DNA			
<213> ORGANISM: <i>Mortierella alpina</i>			
<400> SEQUENCE: 11			
gaattcaggc atggccgccc caatcttga caa			33
<210> SEQ ID NO 12			
<211> LENGTH: 49			
<212> TYPE: DNA			
<213> ORGANISM: <i>Mortierella alpina</i>			
<400> SEQUENCE: 12			
gaattcaggc atctcatgga tccgccatgg ccgcccgaat cttggacaa			49
<210> SEQ ID NO 13			
<211> LENGTH: 317			
<212> TYPE: PRT			
<213> ORGANISM: <i>Mortierella alpina</i>			
<400> SEQUENCE: 13			
Met Ala Ala Ala Ile Leu Asp Lys Val Asn Phe Gly Ile Asp Gln Pro			
1	5	10	15
Phe Gly Ile Lys Leu Asp Thr Tyr Phe Ala Gln Ala Tyr Glu Leu Val			
20	25	30	
Thr Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu Gly Val Thr Pro			
35	40	45	
Leu Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile Thr Tyr Phe Val			
50	55	60	
Val Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln Asp Ala Phe Lys			
65	70	75	80
Leu Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu Thr Ile Ala Ser			
85	90	95	
Gly Ser Leu Leu Leu Phe Ile Glu Asn Leu Val Pro Ile Leu Ala			
100	105	110	
Arg Asn Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly Ala Trp Thr Gln			
115	120	125	

-continued

Arg Leu Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val Lys Tyr Trp Glu
 130 135 140
 Leu Ala Asp Thr Val Phe Leu Val Leu Lys Lys Lys Pro Leu Glu Phe
 145 150 155 160
 Leu His Tyr Phe His His Ser Met Thr Met Val Leu Cys Phe Val Gln
 165 170 175
 Leu Gly Gly Tyr Thr Ser Val Ser Trp Val Pro Ile Thr Leu Asn Leu
 180 185 190
 Thr Val His Val Phe Met Tyr Tyr Tyr Tyr Met Arg Ser Ala Ala Gly
 195 200 205
 Val Arg Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu Gln Ile Val Gln
 210 215 220
 Phe Val Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala Tyr Thr Tyr Phe
 225 230 235 240
 Ala Phe Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly Lys Cys Ala Gly
 245 250 255
 Thr Glu Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu Ser Ser Tyr Leu
 260 265 270
 Leu Leu Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn Ala Lys Ala Lys
 275 280 285
 Ala Ala Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys Thr Val Lys Ser
 290 295 300
 Gly Gly Ser Pro Lys Lys Pro Ser Lys Ser Lys His Ile
 305 310 315

<210> SEQ ID NO 14

<211> LENGTH: 347

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 14

Met Asn Ser Leu Val Thr Gln Tyr Ala Ala Pro Leu Phe Glu Arg Tyr
 1 5 10 15
 Pro Gln Leu His Asp Tyr Leu Pro Thr Leu Glu Arg Pro Phe Phe Asn
 20 25 30
 Ile Ser Leu Trp Glu His Phe Asp Asp Val Val Thr Arg Val Thr Asn
 35 40 45
 Gly Arg Phe Val Pro Ser Glu Phe Gln Phe Ile Ala Gly Glu Leu Pro
 50 55 60
 Leu Ser Thr Leu Pro Pro Val Leu Tyr Ala Ile Thr Ala Tyr Tyr Val
 65 70 75 80
 Ile Ile Phe Gly Gly Arg Phe Leu Leu Ser Lys Ser Lys Pro Phe Lys
 85 90 95
 Leu Asn Gly Leu Phe Gln Leu His Asn Leu Val Leu Thr Ser Leu Ser
 100 105 110
 Leu Thr Leu Leu Leu Leu Met Val Glu Gln Leu Val Pro Ile Ile Val
 115 120 125
 Gln His Gly Leu Tyr Phe Ala Ile Cys Asn Ile Gly Ala Trp Thr Gln
 130 135 140
 Pro Leu Val Thr Leu Tyr Tyr Met Asn Tyr Ile Val Lys Phe Ile Glu
 145 150 155 160
 Phe Ile Asp Thr Phe Phe Leu Val Leu Lys His Lys Lys Leu Thr Phe
 165 170 175

-continued

Leu His Thr Tyr His His Gly Ala Thr Ala Leu Leu Cys Tyr Thr Gln
180 185 190
Leu Met Gly Thr Thr Ser Ile Ser Trp Val Pro Ile Ser Leu Asn Leu
195 200 205
Gly Val His Val Val Met Tyr Trp Tyr Tyr Phe Leu Ala Ala Arg Gly
210 215 220
Ile Arg Val Trp Trp Lys Glu Trp Val Thr Arg Phe Gln Ile Ile Gln
225 230 235 240
Phe Val Leu Asp Ile Gly Phe Ile Tyr Phe Ala Val Tyr Gln Lys Ala
245 250 255
Val His Leu Tyr Phe Pro Ile Leu Pro His Cys Gly Asp Cys Val Gly
260 265 270
Ser Thr Thr Ala Thr Phe Ala Gly Cys Ala Ile Ile Ser Ser Tyr Leu
275 280 285
Val Leu Phe Ile Ser Phe Tyr Ile Asn Val Tyr Lys Arg Lys Gly Thr
290 295 300
Lys Thr Ser Arg Val Val Lys Arg Ala His Gly Gly Val Ala Ala Lys
305 310 315 320
Val Asn Glu Tyr Val Asn Val Asp Leu Lys Asn Val Pro Thr Pro Ser
325 330 335
Pro Ser Pro Lys Pro Gln His Arg Arg Lys Arg
340 345

<210> SEQ ID NO 15

<211> LENGTH: 345

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 15

Met Asn Thr Thr Thr Ser Thr Val Ile Ala Ala Val Ala Asp Gln Phe
1 5 10 15
Gln Ser Leu Asn Ser Ser Ser Ser Cys Phe Leu Lys Val His Val Pro
20 25 30
Ser Ile Glu Asn Pro Phe Gly Ile Glu Leu Trp Pro Ile Phe Ser Lys
35 40 45
Val Phe Glu Tyr Phe Ser Gly Tyr Pro Ala Glu Gln Phe Glu Phe Ile
50 55 60
His Asn Lys Thr Phe Leu Ala Asn Gly Tyr His Ala Val Ser Ile Ile
65 70 75 80
Ile Val Tyr Tyr Ile Ile Ile Phe Gly Gly Gln Ala Ile Leu Arg Ala
85 90 95
Leu Asn Ala Ser Pro Leu Lys Phe Lys Leu Leu Phe Glu Ile His Asn
100 105 110
Leu Phe Leu Thr Ser Ile Ser Leu Val Leu Trp Leu Leu Met Leu Glu
115 120 125
Gln Leu Val Pro Met Val Tyr His Asn Gly Leu Phe Trp Ser Ile Cys
130 135 140
Ser Lys Glu Ala Phe Ala Pro Lys Leu Val Thr Leu Tyr Tyr Leu Asn
145 150 155 160
Tyr Leu Thr Lys Phe Val Glu Leu Ile Asp Thr Val Phe Leu Val Leu
165 170 175
Arg Arg Lys Lys Leu Leu Phe Leu His Thr Tyr His His Gly Ala Thr

-continued

180										185										190									
Ala	Leu	Leu	Cys	Tyr	Thr	Gln	Leu	Ile	Gly	Arg	Thr	Ser	Val	Glu	Trp														
	195						200					205																	
Val	Val	Ile	Leu	Leu	Asn	Leu	Gly	Val	His	Val	Ile	Met	Tyr	Trp	Tyr														
	210					215					220																		
Tyr	Phe	Leu	Ser	Ser	Cys	Gly	Ile	Arg	Val	Trp	Trp	Lys	Gln	Trp	Val														
	225				230					235					240														
Thr	Arg	Phe	Gln	Ile	Ile	Gln	Phe	Leu	Ile	Asp	Leu	Val	Phe	Val	Tyr														
			245						250					255															
Phe	Ala	Thr	Tyr	Thr	Phe	Tyr	Ala	His	Lys	Tyr	Leu	Asp	Gly	Ile	Leu														
			260				265						270																
Pro	Asn	Lys	Gly	Thr	Cys	Tyr	Gly	Thr	Gln	Ala	Ala	Ala	Ala	Tyr	Gly														
	275						280					285																	
Tyr	Leu	Ile	Leu	Thr	Ser	Tyr	Leu	Leu	Leu	Phe	Ile	Ser	Phe	Tyr	Ile														
	290					295					300																		
Gln	Ser	Tyr	Lys	Lys	Gly	Gly	Lys	Lys	Thr	Val	Lys	Lys	Glu	Ser	Glu														
	305				310					315					320														
Val	Ser	Gly	Ser	Val	Ala	Ser	Gly	Ser	Ser	Thr	Gly	Val	Lys	Thr	Ser														
				325					330					335															
Asn	Thr	Lys	Val	Ser	Ser	Arg	Lys	Ala																					
			340				345																						

<210> SEQ ID NO 16
<211> LENGTH: 587
<212> TYPE: DNA
<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 16

tctcgaccca gagagaggtc gccatgtgga ctatcactta cttcgtcgtc atctttggtg 60
gtcgcagat catgaagagc caggacgcct tcaagctcaa gccctcttc atcctccaca 120
acttctctct gacgatcgcg tccggatcgc tgttgctcct gttcatcgag aacctgggtcc 180
ccatcctcgc cagaaacgga cttttctacg ccatctgcga cgacgggtgc tggaccacgc 240
gcctcgagct cctctactac ctcaactacc tggtaagta ctgggagttg gccgacaccg 300
tcttttttgt cctcaagaag aagcctcttg agttcctgca ctacttccac cactcgatga 360
ccatggttct ctgctttgtc cagcttgagg gatacacttc agtgccttgg gtccctatta 420
ccctcaactt gactgtccac gtcttcatgt actactacta catgcgctcc gctgcgggtg 480
ttcgcatctg gtggaagcag tacttgacca ctctccagat cgtccagttc gttcttgacc 540
tcggattcat ctacttctgc gcctacacct acttcgcctt cacctac 587

<210> SEQ ID NO 17
<211> LENGTH: 590
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 17

cattaagcac tttgccccct gtgctatacg ccatcactgc ctattacgtt attatTTTTg 60
gtggcaggtt tttgttaagt aagtcgaaac catttaaatt aaatggcctt ttccaattgc 120
ataatttggt ttaacttca ctttcattga cgcttttatt gcttatgggt gaacaattag 180
tgccaattat tgttcagcac gggttatact tcgctatctg taatattggt gcttggtact 240

-continued

aaccgctcgt tacattatat tacatgaatt acattgtcaa gtttattgaa tttatagaca	300
cctttttctt ggtgctaaaa cataaaaaat tgacattttt gcatacttat caccatggcg	360
ctactgcctt attatgttac acccaattga tgggcaccac atctatttct tgggtcccta	420
tttcattgaa ccttggtgtt cacgtggtta tgtattggta ctatttcttg gctgccagag	480
gcatcagggt ctggtggaag gaatgggta ccagatttca aattatccaa tttgttttgg	540
atatcggttt catatatattt gctgtctacc aaaaagcagt tcacttgat	590

<210> SEQ ID NO 18
<211> LENGTH: 278
<212> TYPE: PRT
<213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 18

Arg Thr Phe Lys Met Met Asp Gln Ile Leu Gly Thr Asn Phe Thr Tyr	
1 5 10 15	
Glu Gly Ala Lys Glu Val Ala Arg Gly Leu Glu Gly Phe Ser Ala Lys	
20 25 30	
Leu Ala Val Gly Tyr Ile Ala Thr Ile Phe Gly Leu Lys Tyr Tyr Met	
35 40 45	
Lys Asp Arg Lys Ala Phe Asp Leu Ser Thr Pro Leu Asn Ile Trp Asn	
50 55 60	
Gly Ile Leu Ser Thr Phe Ser Leu Leu Gly Phe Leu Phe Thr Phe Pro	
65 70 75 80	
Thr Leu Leu Ser Val Ile Arg Lys Asp Gly Phe Ser His Thr Tyr Ser	
85 90 95	
His Val Ser Glu Leu Tyr Thr Asp Ser Thr Ser Gly Tyr Trp Ile Phe	
100 105 110	
Leu Trp Val Ile Ser Lys Ile Pro Glu Leu Leu Asp Thr Val Phe Ile	
115 120 125	
Val Leu Arg Lys Arg Pro Leu Ile Phe Met His Trp Tyr His His Ala	
130 135 140	
Leu Thr Gly Tyr Tyr Ala Leu Val Cys Tyr His Glu Asp Ala Val His	
145 150 155 160	
Met Val Trp Val Val Trp Met Asn Tyr Ile Ile His Ala Phe Met Tyr	
165 170 175	
Gly Tyr Tyr Leu Leu Lys Ser Leu Lys Val Pro Ile Pro Pro Ser Val	
180 185 190	
Ala Gln Ala Ile Thr Thr Ser Gln Met Val Gln Phe Ala Val Ala Ile	
195 200 205	
Phe Ala Gln Val His Val Ser Tyr Lys His Tyr Val Glu Gly Val Glu	
210 215 220	
Gly Leu Ala Tyr Ser Phe Arg Gly Thr Ala Ile Gly Phe Phe Met Leu	
225 230 235 240	
Thr Thr Tyr Phe Tyr Leu Trp Ile Gln Phe Tyr Lys Glu His Tyr Leu	
245 250 255	
Lys Asn Gly Gly Lys Lys Tyr Asn Leu Ala Lys Asp Gln Ala Lys Thr	
260 265 270	
Gln Thr Lys Lys Ala Asn	
275	

<210> SEQ ID NO 19

-continued

<211> LENGTH: 293
 <212> TYPE: PRT
 <213> ORGANISM: *Mortierella alpina*
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (293)...(293)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 293

<400> SEQUENCE: 19

Ala Gln Ala Tyr Glu Leu Val Thr Gly Lys Ser Ile Asp Ser Phe Val
 1 5 10 15
 Phe Gln Glu Gly Val Thr Pro Leu Ser Thr Gln Arg Glu Val Ala Met
 20 25 30
 Trp Thr Ile Thr Tyr Phe Val Val Ile Phe Gly Gly Arg Gln Ile Met
 35 40 45
 Lys Ser Gln Asp Ala Phe Lys Leu Lys Pro Leu Phe Ile Leu His Asn
 50 55 60
 Phe Leu Leu Thr Ile Ala Ser Gly Ser Leu Leu Leu Phe Ile Glu
 65 70 75 80
 Asn Leu Val Pro Ile Leu Ala Arg Asn Gly Leu Phe Tyr Ala Ile Cys
 85 90 95
 Asp Asp Gly Ala Trp Thr Gln Arg Leu Glu Leu Leu Tyr Tyr Leu Asn
 100 105 110
 Tyr Leu Val Lys Tyr Trp Glu Leu Ala Asp Thr Val Phe Leu Val Leu
 115 120 125
 Lys Lys Lys Pro Leu Glu Phe Leu His Tyr Phe His His Ser Met Thr
 130 135 140
 Met Val Leu Cys Phe Val Gln Leu Gly Gly Tyr Thr Ser Val Ser Trp
 145 150 155 160
 Val Pro Ile Thr Leu Asn Leu Thr Val His Val Phe Met Tyr Tyr Tyr
 165 170 175
 Tyr Met Arg Ser Ala Ala Gly Val Arg Ile Trp Trp Lys Gln Tyr Leu
 180 185 190
 Thr Thr Leu Gln Ile Val Gln Phe Val Leu Asp Leu Gly Phe Ile Tyr
 195 200 205
 Phe Cys Ala Tyr Thr Tyr Phe Ala Phe Thr Tyr Phe Pro Trp Ala Pro
 210 215 220
 Asn Val Gly Lys Cys Ala Gly Thr Glu Gly Ala Ala Leu Phe Gly Cys
 225 230 235 240
 Gly Leu Leu Ser Ser Tyr Leu Leu Leu Phe Ile Asn Phe Tyr Arg Ile
 245 250 255
 Thr Tyr Asn Ala Lys Ala Lys Ala Ala Lys Glu Arg Gly Ser Asn Phe
 260 265 270
 Thr Pro Lys Thr Val Lys Ser Gly Gly Ser Pro Lys Lys Pro Ser Lys
 275 280 285
 Ser Lys His Ile Xaa
 290

<210> SEQ ID NO 20
 <211> LENGTH: 238
 <212> TYPE: PRT
 <213> ORGANISM: *Caenorhabditis elegans*

<400> SEQUENCE: 20

Ser Leu Leu Thr Asn Gln Asp Glu Val Phe Pro His Ile Arg Ala Arg

-continued

1	5	10	15
Arg Phe Ile Gln Glu His Phe Gly Leu Phe Val Gln Met Ala Ile Ala	20	25	30
Tyr Val Ile Leu Val Phe Ser Ile Lys Arg Phe Met Arg Asp Arg Glu	35	40	45
Pro Phe Gln Leu Thr Thr Ala Leu Arg Leu Trp Asn Phe Phe Leu Ser	50	55	60
Val Phe Ser Ile Tyr Gly Ser Trp Thr Met Phe Pro Phe Met Val Gln	65	70	75
Gln Ile Arg Leu Tyr Gly Leu Tyr Gly Cys Gly Cys Glu Ala Leu Ser	85	90	95
Asn Leu Pro Ser Gln Ala Glu Tyr Trp Leu Phe Leu Thr Ile Leu Ser	100	105	110
Lys Ala Val Glu Phe Val Asp Thr Phe Phe Leu Val Leu Arg Lys Lys	115	120	125
Pro Leu Ile Phe Leu His Trp Tyr His His Met Ala Thr Phe Val Phe	130	135	140
Phe Cys Ser Asn Tyr Pro Thr Pro Ser Ser Gln Ser Arg Val Gly Val	145	150	155
Ile Val Asn Leu Phe Val His Ala Phe Met Tyr Pro Tyr Tyr Phe Thr	165	170	175
Arg Ser Met Asn Ile Lys Val Pro Ala Lys Ile Ser Met Ala Val Thr	180	185	190
Val Leu Gln Leu Thr Gln Phe Met Cys Phe Ile Tyr Gly Cys Thr Leu	195	200	205
Met Tyr Tyr Ser Leu Ala Thr Asn Gln Ala Arg Tyr Pro Ser Asn Thr	210	215	220
Pro Ala Thr Leu Gln Cys Leu Ser Tyr Thr Leu His Leu Leu	225	230	235

<210> SEQ ID NO 21

<211> LENGTH: 289

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (289)...(289)

<223> OTHER INFORMATION: Xaa = Unknown or Other at position 289

<400> SEQUENCE: 21

Glu Leu Val Thr Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu Gly	1	5	10	15
Val Thr Pro Leu Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile Thr	20	25	30	
Tyr Phe Val Val Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln Asp	35	40	45	
Ala Phe Lys Leu Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu Thr	50	55	60	
Ile Ala Ser Gly Ser Leu Leu Leu Leu Phe Ile Glu Asn Leu Val Pro	65	70	75	80
Ile Leu Ala Arg Asn Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly Ala	85	90	95	
Trp Thr Gln Arg Leu Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val Lys	100	105	110	

-continued

Tyr Trp Glu Leu Ala Asp Thr Val Phe Leu Val Leu Lys Lys Lys Pro
115 120 125
Leu Glu Phe Leu His Tyr Phe His His Ser Met Thr Met Val Leu Cys
130 135 140
Phe Val Gln Leu Gly Gly Tyr Thr Ser Val Ser Trp Val Pro Ile Thr
145 150 155 160
Leu Asn Leu Thr Val His Val Phe Met Tyr Tyr Tyr Tyr Met Arg Ser
165 170 175
Ala Ala Gly Val Arg Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu Gln
180 185 190
Ile Val Gln Phe Val Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala Tyr
195 200 205
Thr Tyr Phe Ala Phe Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly Lys
210 215 220
Cys Ala Gly Thr Glu Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu Ser
225 230 235 240
Ser Tyr Leu Leu Leu Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn Ala
245 250 255
Lys Ala Lys Ala Ala Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys Thr
260 265 270
Val Lys Ser Gly Gly Ser Pro Lys Lys Pro Ser Lys Ser Lys His Ile
275 280 285

Xaa

<210> SEQ ID NO 22
<211> LENGTH: 101
<212> TYPE: PRT
<213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 22

Met Leu Tyr Ser Ile Thr Arg Arg Cys Tyr Thr Phe Phe Val Thr Ser
1 5 10 15
Leu His Phe Tyr Gln Leu Tyr Val Thr Glu Cys Leu Glu Asn Val Ile
20 25 30
Phe Asn Val Leu Val Asn Gly Gln Ser Ile Asn Ser Arg Trp Lys Asp
35 40 45
Ala Glu Lys Thr Ile Thr Ser Phe Pro Phe His Phe Pro Gln Thr Phe
50 55 60
Phe Gln Gln Pro His Ile Leu Thr Leu His Phe Leu Phe Phe Val Phe
65 70 75 80
Val Ser Val Thr Leu Val Thr Val Phe Lys Lys Pro Lys Cys Glu Phe
85 90 95
Pro His Ser Leu Ala
100

<210> SEQ ID NO 23
<211> LENGTH: 115
<212> TYPE: PRT
<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 23

Met Ala Ala Ala Ile Leu Asp Lys Val Asn Phe Gly Ile Asp Gln Pro
1 5 10 15

-continued

Phe Gly Ile Lys Leu Asp Thr Tyr Phe Ala Gln Ala Tyr Glu Leu Val
 20 25 30
 Thr Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu Gly Val Thr Pro
 35 40 45
 Leu Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile Thr Tyr Phe Val
 50 55 60
 Val Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln Asp Ala Phe Lys
 65 70 75 80
 Leu Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu Thr Ile Ala Ser
 85 90 95
 Gly Ser Leu Leu Leu Leu Phe Ile Glu Asn Leu Val Pro Ile Leu Ala
 100 105 110
 Arg Asn Gly
 115

<210> SEQ ID NO 24
 <211> LENGTH: 272
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (272)...(272)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 272

<400> SEQUENCE: 24

Met Asp Thr Ser Met Asn Phe Ser Arg Gly Leu Lys Met Asp Leu Met
 1 5 10 15
 Gln Pro Tyr Asp Phe Glu Thr Phe Gln Asp Leu Arg Pro Phe Leu Glu
 20 25 30
 Glu Tyr Trp Val Ser Ser Phe Leu Ile Val Val Val Tyr Leu Leu Leu
 35 40 45
 Ile Val Val Gly Gln Thr Tyr Met Arg Thr Arg Lys Ser Phe Ser Leu
 50 55 60
 Gln Arg Pro Leu Ile Leu Trp Ser Phe Phe Leu Ala Ile Phe Ser Ile
 65 70 75 80
 Leu Gly Thr Leu Arg Met Trp Lys Phe Met Ala Thr Val Met Phe Thr
 85 90 95
 Val Gly Leu Lys Gln Thr Val Cys Phe Ala Ile Tyr Thr Asp Asp Ala
 100 105 110
 Val Val Arg Phe Trp Ser Phe Leu Phe Leu Leu Ser Lys Val Val Glu
 115 120 125
 Leu Gly Asp Thr Ala Phe Ile Ile Leu Arg Lys Arg Pro Leu Ile Phe
 130 135 140
 Val His Trp Tyr His His Ser Thr Val Leu Leu Phe Thr Ser Phe Gly
 145 150 155 160
 Tyr Lys Asn Lys Val Pro Ser Gly Gly Trp Phe Met Thr Met Asn Phe
 165 170 175
 Gly Val His Ser Val Met Tyr Thr Tyr Tyr Thr Met Lys Ala Ala Lys
 180 185 190
 Leu Lys His Pro Asn Leu Leu Pro Met Val Ile Thr Ser Leu Gln Ile
 195 200 205
 Leu Gln Met Val Leu Gly Thr Ile Phe Gly Ile Leu Asn Tyr Ile Trp
 210 215 220
 Arg Gln Glu Lys Gly Cys His Thr Thr Thr Glu His Phe Phe Trp Ser

-continued

225	230	235	240
Phe Met Leu Tyr Gly Thr Tyr Phe Ile Leu Phe Ala His Phe Phe His	245	250	255
Arg Ala Tyr Leu Arg Pro Lys Gly Lys Val Ala Ser Lys Ser Gln Xaa	260	265	270

<210> SEQ ID NO 25
 <211> LENGTH: 318
 <212> TYPE: PRT
 <213> ORGANISM: Mortierella alpina
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (318)...(318)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 318
 <400> SEQUENCE: 25

Met Ala Ala Ala Ile Leu Asp Lys Val Asn Phe Gly Ile Asp Gln Pro	1	5	10	15
Phe Gly Ile Lys Leu Asp Thr Tyr Phe Ala Gln Ala Tyr Glu Leu Val	20	25	30	
Thr Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu Gly Val Thr Pro	35	40	45	
Leu Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile Thr Tyr Phe Val	50	55	60	
Val Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln Asp Ala Phe Lys	65	70	75	80
Leu Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu Thr Ile Ala Ser	85	90	95	
Gly Ser Leu Leu Leu Leu Phe Ile Glu Asn Leu Val Pro Ile Leu Ala	100	105	110	
Arg Asn Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly Ala Trp Thr Gln	115	120	125	
Arg Leu Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val Lys Tyr Trp Glu	130	135	140	
Leu Ala Asp Thr Val Phe Leu Val Leu Lys Lys Lys Pro Leu Glu Phe	145	150	155	160
Leu His Tyr Phe His His Ser Met Thr Met Val Leu Cys Phe Val Gln	165	170	175	
Leu Gly Gly Tyr Thr Ser Val Ser Trp Val Pro Ile Thr Leu Asn Leu	180	185	190	
Thr Val His Val Phe Met Tyr Tyr Tyr Tyr Met Arg Ser Ala Ala Gly	195	200	205	
Val Arg Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu Gln Ile Val Gln	210	215	220	
Phe Val Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala Tyr Thr Tyr Phe	225	230	235	240
Ala Phe Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly Lys Cys Ala Gly	245	250	255	
Thr Glu Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu Ser Ser Tyr Leu	260	265	270	
Leu Leu Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn Ala Lys Ala Lys	275	280	285	
Ala Ala Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys Thr Val Lys Ser	290	295	300	

-continued

Gly Gly Ser Pro Lys Lys Pro Ser Lys Ser Lys His Ile Xaa
 305 310 315

<210> SEQ ID NO 26

<211> LENGTH: 178

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 26

Asn Leu Val Pro Ile Leu Ala Arg Asn Gly Leu Phe Tyr Ala Ile Cys
 1 5 10 15
 Asp Asp Gly Ala Trp Thr Gln Arg Leu Glu Leu Leu Tyr Tyr Leu Asn
 20 25 30
 Tyr Leu Val Lys Tyr Trp Glu Leu Ala Asp Thr Val Phe Leu Val Leu
 35 40 45
 Lys Lys Lys Pro Leu Glu Phe Leu His Tyr Phe His His Ser Met Thr
 50 55 60
 Met Val Leu Cys Phe Val Gln Leu Gly Gly Tyr Thr Ser Val Ser Trp
 65 70 75 80
 Val Pro Ile Thr Leu Asn Leu Thr Val His Val Phe Met Tyr Tyr Tyr
 85 90 95
 Tyr Met Arg Ser Ala Ala Gly Val Arg Ile Trp Trp Lys Gln Tyr Leu
 100 105 110
 Thr Thr Leu Gln Ile Val Gln Phe Val Leu Asp Leu Gly Phe Ile Tyr
 115 120 125
 Phe Cys Ala Tyr Thr Tyr Phe Ala Phe Thr Tyr Phe Pro Trp Ala Pro
 130 135 140
 Asn Val Gly Lys Cys Ala Gly Thr Glu Gly Ala Ala Leu Phe Gly Cys
 145 150 155 160
 Gly Leu Leu Ser Ser Tyr Leu Leu Leu Phe Ile Asn Phe Tyr Arg Ile
 165 170 175
 Thr Tyr

<210> SEQ ID NO 27

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Ser Leu Leu Val Val Lys Asp Leu Thr Tyr Leu Leu Pro Leu Cys Leu
 1 5 10 15
 Pro Gly Asp Thr Ile Phe Ile Ile Leu Arg Lys Gln Lys Leu Ile Phe
 20 25 30
 Leu His Trp Tyr His His Ile Thr Val Leu Leu Tyr Ser Trp Tyr Ser
 35 40 45
 Tyr Lys Asp Met Val Ala Gly Gly Gly Trp Phe Met Thr Met Asn Tyr
 50 55 60
 Gly Val His Ala Val Met Tyr Ser Tyr Tyr Ala Leu Arg Ala Ala Gly
 65 70 75 80
 Phe Arg Val Ser Arg Lys Phe Ala Met Phe Ile Thr Leu Ser Gln Ile
 85 90 95
 Thr Gln Met Leu Met Gly Cys Val Val Asn Tyr Leu Val Phe Cys Trp
 100 105 110

-continued

Met Gln His Asp Gln Cys His Ser His Phe Gln Asn Ile Phe Trp Ser
 115 120 125

Ser Leu Met Tyr Leu Ser Tyr Leu Val Leu Phe Cys His Phe Phe Phe
 130 135 140

Glu Ala Tyr
 145

<210> SEQ ID NO 28
 <211> LENGTH: 280
 <212> TYPE: PRT
 <213> ORGANISM: Mortierella alpina
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (280)...(280)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 280

<400> SEQUENCE: 28

Ser Phe Val Phe Gln Glu Gly Val Thr Pro Leu Ser Thr Gln Arg Glu
 1 5 10 15

Val Ala Met Trp Thr Ile Thr Tyr Phe Val Val Ile Phe Gly Gly Arg
 20 25 30

Gln Ile Met Lys Ser Gln Asp Ala Phe Lys Leu Lys Pro Leu Phe Ile
 35 40 45

Leu His Asn Phe Leu Leu Thr Ile Ala Ser Gly Ser Leu Leu Leu Leu
 50 55 60

Phe Ile Glu Asn Leu Val Pro Ile Leu Ala Arg Asn Gly Leu Phe Tyr
 65 70 75 80

Ala Ile Cys Asp Asp Gly Ala Trp Thr Gln Arg Leu Glu Leu Leu Tyr
 85 90 95

Tyr Leu Asn Tyr Leu Val Lys Tyr Trp Glu Leu Ala Asp Thr Val Phe
 100 105 110

Leu Val Leu Lys Lys Lys Pro Leu Glu Phe Leu His Tyr Phe His His
 115 120 125

Ser Met Thr Met Val Leu Cys Phe Val Gln Leu Gly Gly Tyr Thr Ser
 130 135 140

Val Ser Trp Val Pro Ile Thr Leu Asn Leu Thr Val His Val Phe Met
 145 150 155 160

Tyr Tyr Tyr Tyr Met Arg Ser Ala Ala Gly Val Arg Ile Trp Trp Lys
 165 170 175

Gln Tyr Leu Thr Thr Leu Gln Ile Val Gln Phe Val Leu Asp Leu Gly
 180 185 190

Phe Ile Tyr Phe Cys Ala Tyr Thr Tyr Phe Ala Phe Thr Tyr Phe Pro
 195 200 205

Trp Ala Pro Asn Val Gly Lys Cys Ala Gly Thr Glu Gly Ala Ala Leu
 210 215 220

Phe Gly Cys Gly Leu Leu Ser Ser Tyr Leu Leu Leu Phe Ile Asn Phe
 225 230 235 240

Tyr Arg Ile Thr Tyr Asn Ala Lys Ala Lys Ala Ala Lys Glu Arg Gly
 245 250 255

Ser Asn Phe Thr Pro Lys Thr Val Lys Ser Gly Gly Ser Pro Lys Lys
 260 265 270

Pro Ser Lys Ser Lys His Ile Xaa
 275 280

-continued

```

<210> SEQ ID NO 29
<211> LENGTH: 283
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Potential Mammalian Elongase
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (282)...(282)
<223> OTHER INFORMATION: Xaa = Unknown or Other at position 282

```

```

<400> SEQUENCE: 29

```

```

Pro Arg Tyr Lys Ser Gln Arg Met Val Pro Pro Gly Gln Leu His Pro
 1           5           10           15
Tyr Val Cys Leu Phe Cys Tyr Leu Leu Thr His Cys Met Ala Gly Thr
      20           25           30
Lys Ile His Glu Glu Pro Ala Ala Val Leu Leu Pro Ser Ile Leu Gln
      35           40           45
Leu Tyr Asn Leu Gly Leu Thr Leu Leu Ser Leu Tyr Met Phe Tyr Glu
      50           55           60
Leu Val Thr Gly Val Trp Glu Gly Lys Tyr Asn Phe Phe Cys Gln Gly
      65           70           75           80
Thr Arg Ser Ala Gly Glu Ser Asp Met Lys Ile Ile Arg Val Leu Trp
      85           90           95
Trp Tyr Tyr Phe Ser Lys Leu Ile Glu Phe Met Asp Thr Phe Phe Phe
      100          105          110
Ile Leu Arg Lys Asn Asn His Gln Ile Thr Val Leu His Val Tyr His
      115          120          125
His Ala Thr Met Leu Asn Ile Trp Trp Phe Val Met Asn Trp Val Pro
      130          135          140
Cys Gly His Ser Tyr Phe Gly Ala Thr Leu Asn Ser Phe Ile His Val
      145          150          155          160
Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Ser Ile Pro Ser Met Arg Pro
      165          170          175
Tyr Leu Trp Trp Lys Lys Tyr Ile Thr Gln Gly Gln Leu Val Gln Phe
      180          185          190
Val Leu Thr Ile Ile Gln Thr Thr Cys Gly Val Phe Trp Pro Cys Ser
      195          200          205
Phe Pro Leu Gly Trp Leu Phe Phe Gln Ile Gly Tyr Met Ile Ser Leu
      210          215          220
Ile Ala Leu Phe Thr Asn Phe Tyr Ile Gln Thr Tyr Asn Lys Lys Gly
      225          230          235          240
Ala Ser Arg Arg Lys Glu His Leu Lys Gly His Gln Asn Gly Ser Val
      245          250          255
Ala Ala Val Asn Gly His Thr Asn Ser Phe Pro Ser Leu Glu Asn Ser
      260          265          270
Val Lys Pro Arg Lys Gln Arg Lys Asp Xaa Gln
      275          280

```

```

<210> SEQ ID NO 30
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Mortierella alpina

```

```

<400> SEQUENCE: 30

```

```

Met Gly Thr Asp Gln Gly Lys Thr Phe Thr Trp Glu Glu Leu Ala Ala

```

-continued

1	5	10	15
His Asn Thr Lys Asp Asp Leu Leu Leu Ala Ile Arg Gly Arg Val Tyr	20	25	30
Asp Val Thr Lys Phe Leu Ser Arg His Pro Gly Gly Val Asp Thr Leu	35	40	45
Leu Leu Gly Ala Gly Arg Asp Val Thr Pro Val Phe Glu Met Tyr His	50	55	60
Ala Phe Gly Ala Ala Asp Ala Ile Met Lys Lys Tyr Tyr Val Gly Thr	65	70	80
Leu Val Ser Asn Glu Leu Pro Ile Phe Pro Glu Pro Thr Val Phe His	85	90	95
Lys Thr Ile Lys Thr Arg Val Glu Gly Tyr Phe Thr Asp Arg Asn Ile	100	105	110
Asp Pro Lys Asn Arg Pro Glu Ile Trp Gly Arg Tyr Ala Leu Ile Phe	115	120	125
Gly Ser Leu Ile Ala Ser Tyr Tyr Ala Gln Leu Phe Val Pro Phe Val	130	135	140
Val Glu Arg Thr Trp Leu Gln Val Val Phe Ala Ile Ile Met Gly Phe	145	150	155
Ala Cys Ala Gln Val Gly Leu Asn Pro Leu His Asp Ala Ser His Phe	165	170	175
Ser Val Thr His Asn Pro Thr Val Trp Lys Ile Leu Gly Ala Thr His	180	185	190
Asp Phe Phe Asn Gly Ala Ser Tyr Leu Val Trp Met Tyr Gln His Met	195	200	205
Leu Gly His His Pro Tyr Thr Asn Ile Ala Gly Ala Asp Pro Asp Val	210	215	220
Ser Thr Ser Glu Pro Asp Val Arg Arg Ile Lys Pro Asn Gln Lys Trp	225	230	235
Phe Val Asn His Ile Asn Gln His Met Phe Val Pro Phe Leu Tyr Gly	245	250	255
Leu Leu Ala Phe Lys Val Arg Ile Gln Asp Ile Asn Ile Leu Tyr Phe	260	265	270
Val Lys Thr Asn Asp Ala Ile Arg Val Asn Pro Ile Ser Thr Trp His	275	280	285
Thr Val Met Phe Trp Gly Gly Lys Ala Phe Phe Val Trp Tyr Arg Leu	290	295	300
Ile Val Pro Leu Gln Tyr Leu Pro Leu Gly Lys Val Leu Leu Leu Phe	305	310	315
Thr Val Ala Asp Met Val Ser Ser Tyr Trp Leu Ala Leu Thr Phe Gln	325	330	335
Ala Asn His Val Val Glu Glu Val Gln Trp Pro Leu Pro Asp Glu Asn	340	345	350
Gly Ile Ile Gln Lys Asp Trp Ala Ala Met Gln Val Glu Thr Thr Gln	355	360	365
Asp Tyr Ala His Asp Ser His Leu Trp Thr Ser Ile Thr Gly Ser Leu	370	375	380
Asn Tyr Gln Ala Val His His Leu Phe Pro Asn Val Ser Gln His His	385	390	395
Tyr Pro Asp Ile Leu Ala Ile Ile Lys Asn Thr Cys Ser Glu Tyr Lys	405	410	415

-continued

Val Pro Tyr Leu Val Lys Asp Thr Phe Trp Gln Ala Phe Ala Ser His
 420 425 430

Leu Glu His Leu Arg Val Leu Gly Leu Arg Pro Lys Glu Glu
 435 440 445

<210> SEQ ID NO 31

<211> LENGTH: 318

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 31

Met Glu Ser Ile Ala Pro Phe Leu Pro Ser Lys Met Pro Gln Asp Leu
 1 5 10 15

Phe Met Asp Leu Ala Thr Ala Ile Gly Val Arg Ala Ala Pro Tyr Val
 20 25 30

Asp Pro Leu Glu Ala Ala Leu Val Ala Gln Ala Glu Lys Tyr Ile Pro
 35 40 45

Thr Ile Val His His Thr Arg Gly Phe Leu Val Ala Val Glu Ser Pro
 50 55 60

Leu Ala Arg Glu Leu Pro Leu Met Asn Pro Phe His Val Leu Leu Ile
 65 70 75 80

Val Leu Ala Tyr Leu Val Thr Val Phe Val Gly Met Gln Ile Met Lys
 85 90 95

Asn Phe Glu Arg Phe Glu Val Lys Thr Phe Ser Leu Leu His Asn Phe
 100 105 110

Cys Leu Val Ser Ile Ser Ala Tyr Met Cys Gly Gly Ile Leu Tyr Glu
 115 120 125

Ala Tyr Gln Ala Asn Tyr Gly Leu Phe Glu Asn Ala Ala Asp His Thr
 130 135 140

Phe Lys Gly Leu Pro Met Ala Lys Met Ile Trp Leu Phe Tyr Phe Ser
 145 150 155 160

Lys Ile Met Glu Phe Val Asp Thr Met Ile Met Val Leu Lys Lys Asn
 165 170 175

Asn Arg Gln Ile Ser Phe Leu His Val Tyr His His Ser Ser Ile Phe
 180 185 190

Thr Ile Trp Trp Leu Val Thr Phe Val Ala Pro Asn Gly Glu Ala Tyr
 195 200 205

Phe Ser Ala Ala Leu Asn Ser Phe Ile His Val Ile Met Tyr Gly Tyr
 210 215 220

Tyr Phe Leu Ser Ala Leu Gly Phe Lys Gln Val Ser Phe Ile Lys Phe
 225 230 235 240

Tyr Ile Thr Arg Ser Gln Met Thr Gln Phe Cys Met Met Ser Val Gln
 245 250 255

Ser Ser Trp Asp Met Tyr Ala Met Lys Val Leu Gly Arg Pro Gly Tyr
 260 265 270

Pro Phe Phe Ile Thr Ala Leu Leu Trp Phe Tyr Met Trp Thr Met Leu
 275 280 285

Gly Leu Phe Tyr Asn Phe Tyr Arg Lys Asn Ala Lys Leu Ala Lys Gln
 290 295 300

Ala Lys Ala Asp Ala Ala Lys Glu Lys Ala Arg Lys Leu Gln
 305 310 315

-continued

<210> SEQ ID NO 32
<211> LENGTH: 279
<212> TYPE: PRT
<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 32

Val Ala Gln Ala Glu Lys Tyr Ile Pro Thr Ile Val His His Thr Arg
1 5 10 15
Gly Phe Leu Val Ala Val Glu Ser Pro Leu Ala Arg Glu Leu Pro Leu
20 25 30
Met Asn Pro Phe His Val Leu Leu Ile Val Leu Ala Tyr Leu Val Thr
35 40 45
Val Phe Val Gly Met Gln Ile Met Lys Asn Phe Glu Arg Phe Glu Val
50 55 60
Lys Thr Phe Ser Leu Leu His Asn Phe Cys Leu Val Ser Ile Ser Ala
65 70 75 80
Tyr Met Cys Gly Gly Ile Leu Tyr Glu Ala Tyr Gln Ala Asn Tyr Gly
85 90 95
Leu Phe Glu Asn Ala Ala Asp His Thr Phe Lys Gly Leu Pro Met Ala
100 105 110
Lys Met Ile Trp Leu Phe Tyr Phe Ser Lys Ile Met Glu Phe Val Asp
115 120 125
Thr Met Ile Met Val Leu Lys Lys Asn Asn Arg Gln Ile Ser Phe Leu
130 135 140
His Val Tyr His His Ser Ser Ile Phe Thr Ile Trp Trp Leu Val Thr
145 150 155 160
Phe Val Ala Pro Asn Gly Glu Ala Tyr Phe Ser Ala Ala Leu Asn Ser
165 170 175
Phe Ile His Val Ile Met Tyr Gly Tyr Tyr Phe Leu Ser Ala Leu Gly
180 185 190
Phe Lys Gln Val Ser Phe Ile Lys Phe Tyr Ile Thr Arg Ser Gln Met
195 200 205
Thr Gln Phe Cys Met Met Ser Val Gln Ser Ser Trp Asp Met Tyr Ala
210 215 220
Met Lys Val Leu Gly Arg Pro Gly Tyr Pro Phe Phe Ile Thr Ala Leu
225 230 235 240
Leu Trp Phe Tyr Met Trp Thr Met Leu Gly Leu Phe Tyr Asn Phe Tyr
245 250 255
Arg Lys Asn Ala Lys Leu Ala Lys Gln Ala Lys Ala Asp Ala Ala Lys
260 265 270
Glu Lys Ala Arg Lys Leu Gln
275

<210> SEQ ID NO 33
<211> LENGTH: 301
<212> TYPE: PRT
<213> ORGANISM: Mortierella alpina
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (301)...(301)
<223> OTHER INFORMATION: Xaa = Unknown or Other at position 301

<400> SEQUENCE: 33

Gly Ile Lys Leu Asp Thr Tyr Phe Ala Gln Ala Tyr Glu Leu Val Thr
1 5 10 15

-continued

Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu Gly Val Thr Pro Leu
 20 25 30
 Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile Thr Tyr Phe Val Val
 35 40 45
 Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln Asp Ala Phe Lys Leu
 50 55 60
 Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu Thr Ile Ala Ser Gly
 65 70 75 80
 Ser Leu Leu Leu Leu Phe Ile Glu Asn Leu Val Pro Ile Leu Ala Arg
 85 90 95
 Asn Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly Ala Trp Thr Gln Arg
 100 105 110
 Leu Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val Lys Tyr Trp Glu Leu
 115 120 125
 Ala Asp Thr Val Phe Leu Val Leu Lys Lys Lys Pro Leu Glu Phe Leu
 130 135 140
 His Tyr Phe His His Ser Met Thr Met Val Leu Cys Phe Val Gln Leu
 145 150 155 160
 Gly Gly Tyr Thr Ser Val Ser Trp Val Pro Ile Thr Leu Asn Leu Thr
 165 170 175
 Val His Val Phe Met Tyr Tyr Tyr Tyr Met Arg Ser Ala Ala Gly Val
 180 185 190
 Arg Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu Gln Ile Val Gln Phe
 195 200 205
 Val Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala Tyr Thr Tyr Phe Ala
 210 215 220
 Phe Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly Lys Cys Ala Gly Thr
 225 230 235 240
 Glu Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu Ser Ser Tyr Leu Leu
 245 250 255
 Leu Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn Ala Lys Ala Lys Ala
 260 265 270
 Ala Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys Thr Val Lys Ser Gly
 275 280 285
 Gly Ser Pro Lys Lys Pro Ser Lys Ser Lys His Ile Xaa
 290 295 300

<210> SEQ ID NO 34

<211> LENGTH: 289

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 34

Tyr Glu Leu Val Thr Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu
 1 5 10 15
 Gly Val Thr Pro Leu Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile
 20 25 30
 Thr Tyr Phe Val Val Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln
 35 40 45
 Asp Ala Phe Lys Leu Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu
 50 55 60
 Thr Ile Ala Ser Gly Ser Leu Leu Leu Leu Phe Ile Glu Asn Leu Val
 65 70 75 80

-continued

```

Pro Ile Leu Ala Arg Asn Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly
      85              90              95
Ala Trp Thr Gln Arg Leu Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val
      100              105              110
Lys Tyr Trp Glu Leu Ala Asp Thr Val Phe Leu Val Leu Lys Lys Lys
      115              120              125
Pro Leu Glu Phe Leu His Tyr Phe His His Ser Met Thr Met Val Leu
      130              135              140
Cys Phe Val Gln Leu Gly Gly Tyr Thr Ser Val Ser Trp Val Pro Ile
      145              150              155              160
Thr Leu Asn Leu Thr Val His Val Phe Met Tyr Tyr Tyr Tyr Met Arg
      165              170              175
Ser Ala Ala Gly Val Arg Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu
      180              185              190
Gln Ile Val Gln Phe Val Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala
      195              200              205
Tyr Thr Tyr Phe Ala Phe Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly
      210              215              220
Lys Cys Ala Gly Thr Glu Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu
      225              230              235              240
Ser Ser Tyr Leu Leu Leu Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn
      245              250              255
Ala Lys Ala Lys Ala Ala Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys
      260              265              270
Thr Val Lys Ser Gly Gly Ser Pro Lys Lys Pro Ser Lys Ser Lys His
      275              280              285
Ile

```

```

<210> SEQ ID NO 35
<211> LENGTH: 292
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (292)...(292)
<223> OTHER INFORMATION: Xaa = Unknown or Other at position 292

```

```

<400> SEQUENCE: 35

```

```

Ser Thr Tyr Phe Lys Ala Leu Leu Gly Pro Arg Asp Thr Arg Val Lys
  1          5              10              15
Gly Trp Phe Leu Leu Asp Asn Tyr Ile Pro Thr Phe Ile Cys Ser Val
      20              25              30
Ile Tyr Leu Leu Ile Val Trp Leu Gly Pro Lys Tyr Met Arg Asn Lys
      35              40              45
Gln Pro Phe Ser Cys Arg Gly Ile Leu Val Val Tyr Asn Leu Gly Leu
      50              55              60
Thr Leu Leu Ser Leu Tyr Met Phe Cys Glu Leu Val Thr Gly Val Trp
      65              70              75              80
Glu Gly Lys Tyr Asn Phe Phe Cys Gln Gly Thr Arg Thr Ala Gly Glu
      85              90              95
Ser Asp Met Lys Ile Ile Arg Val Leu Trp Trp Tyr Tyr Phe Ser Lys
      100              105              110
Leu Ile Glu Phe Met Asp Thr Phe Phe Phe Ile Leu Arg Lys Asn Asn

```

-continued

115					120					125					
His	Gln	Ile	Thr	Val	Leu	His	Val	Tyr	His	His	Ala	Ser	Met	Leu	Asn
130					135					140					
Ile	Trp	Trp	Phe	Val	Met	Asn	Trp	Val	Pro	Cys	Gly	His	Ser	Tyr	Phe
145					150					155					160
Gly	Ala	Thr	Leu	Asn	Ser	Phe	Ile	His	Val	Leu	Met	Tyr	Ser	Tyr	Tyr
				165					170					175	
Gly	Leu	Ser	Ser	Val	Pro	Ser	Met	Arg	Pro	Tyr	Leu	Trp	Trp	Lys	Lys
				180				185						190	
Tyr	Ile	Thr	Gln	Gly	Gln	Leu	Leu	Gln	Phe	Val	Leu	Thr	Ile	Ile	Gln
				195				200						205	
Thr	Ser	Cys	Gly	Val	Ile	Trp	Pro	Cys	Thr	Phe	Pro	Leu	Gly	Trp	Leu
				210				215						220	
Tyr	Phe	Gln	Ile	Gly	Tyr	Met	Ile	Ser	Leu	Ile	Ala	Leu	Phe	Thr	Asn
225					230					235					240
Phe	Tyr	Ile	Gln	Thr	Tyr	Asn	Lys	Lys	Gly	Ala	Ser	Arg	Arg	Lys	Asp
				245					250					255	
His	Leu	Lys	Asp	His	Gln	Asn	Gly	Ser	Met	Ala	Ala	Val	Asn	Gly	His
				260					265					270	
Thr	Asn	Ser	Phe	Ser	Pro	Leu	Glu	Asn	Asn	Val	Lys	Pro	Arg	Lys	Leu
				275				280						285	
Arg	Lys	Asp	Xaa												
				290											

<210> SEQ ID NO 36

<211> LENGTH: 291

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 36

Gln	Ala	Tyr	Glu	Leu	Val	Thr	Gly	Lys	Ser	Ile	Asp	Ser	Phe	Val	Phe
1				5					10					15	
Gln	Glu	Gly	Val	Thr	Pro	Leu	Ser	Thr	Gln	Arg	Glu	Val	Ala	Met	Trp
			20					25					30		
Thr	Ile	Thr	Tyr	Phe	Val	Val	Ile	Phe	Gly	Gly	Arg	Gln	Ile	Met	Lys
			35				40					45			
Ser	Gln	Asp	Ala	Phe	Lys	Leu	Lys	Pro	Leu	Phe	Ile	Leu	His	Asn	Phe
			50			55				60					
Leu	Leu	Thr	Ile	Ala	Ser	Gly	Ser	Leu	Leu	Leu	Leu	Phe	Ile	Glu	Asn
65				70					75					80	
Leu	Val	Pro	Ile	Leu	Ala	Arg	Asn	Gly	Leu	Phe	Tyr	Ala	Ile	Cys	Asp
			85					90						95	
Asp	Gly	Ala	Trp	Thr	Gln	Arg	Leu	Glu	Leu	Leu	Tyr	Tyr	Leu	Asn	Tyr
			100				105						110		
Leu	Val	Lys	Tyr	Trp	Glu	Leu	Ala	Asp	Thr	Val	Phe	Leu	Val	Leu	Lys
			115			120						125			
Lys	Lys	Pro	Leu	Glu	Phe	Leu	His	Tyr	Phe	His	His	Ser	Met	Thr	Met
			130			135						140			
Val	Leu	Cys	Phe	Val	Gln	Leu	Gly	Gly	Tyr	Thr	Ser	Val	Ser	Trp	Val
145				150					155					160	
Pro	Ile	Thr	Leu	Asn	Leu	Thr	Val	His	Val	Phe	Met	Tyr	Tyr	Tyr	Tyr
			165					170						175	

-continued

Met Arg Ser Ala Ala Gly Val Arg Ile Trp Trp Lys Gln Tyr Leu Thr
 180 185 190

Thr Leu Gln Ile Val Gln Phe Val Leu Asp Leu Gly Phe Ile Tyr Phe
 195 200 205

Cys Ala Tyr Thr Tyr Phe Ala Phe Thr Tyr Phe Pro Trp Ala Pro Asn
 210 215 220

Val Gly Lys Cys Ala Gly Thr Glu Gly Ala Ala Leu Phe Gly Cys Gly
 225 230 235 240

Leu Leu Ser Ser Tyr Leu Leu Leu Phe Ile Asn Phe Tyr Arg Ile Thr
 245 250 255

Tyr Asn Ala Lys Ala Lys Ala Ala Lys Glu Arg Gly Ser Asn Phe Thr
 260 265 270

Pro Lys Thr Val Lys Ser Gly Gly Ser Pro Lys Lys Pro Ser Lys Ser
 275 280 285

Lys His Ile
 290

<210> SEQ ID NO 37
 <211> LENGTH: 276
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (276)...(276)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 276

<400> SEQUENCE: 37

Val Asn Leu Tyr Gln Glu Val Met Lys His Ala Asp Pro Arg Ile Gln
 1 5 10 15

Gly Tyr Pro Leu Met Gly Ser Pro Leu Leu Met Thr Ser Ile Leu Leu
 20 25 30

Thr Tyr Val Tyr Phe Val Leu Ser Leu Gly Pro Arg Ile Met Ala Asn
 35 40 45

Arg Lys Pro Phe Gln Leu Arg Gly Phe Met Ile Val Tyr Asn Phe Ser
 50 55 60

Leu Val Ala Leu Ser Leu Tyr Ile Val Tyr Glu Phe Leu Met Ser Gly
 65 70 75 80

Trp Leu Ser Thr Tyr Thr Trp Arg Cys Asp Pro Val Asp Tyr Ser Asn
 85 90 95

Ser Pro Glu Ala Leu Arg Met Val Arg Val Ala Trp Leu Phe Leu Phe
 100 105 110

Ser Lys Phe Ile Glu Leu Met Asp Thr Val Ile Phe Ile Leu Arg Lys
 115 120 125

Lys Asp Gly Gln Val Thr Phe Leu His Val Phe His His Ser Val Leu
 130 135 140

Pro Trp Ser Trp Trp Trp Gly Val Lys Ile Ala Pro Gly Gly Met Gly
 145 150 155 160

Ser Phe His Ala Met Ile Asn Ser Ser Val His Val Ile Met Tyr Leu
 165 170 175

Tyr Tyr Gly Leu Ser Ala Phe Gly Pro Val Ala Gln Pro Tyr Leu Trp
 180 185 190

Trp Lys Lys His Met Thr Ala Ile Gln Leu Ile Gln Phe Val Leu Val
 195 200 205

Ser Leu His Ile Ser Gln Tyr Tyr Phe Met Ser Ser Cys Asn Tyr Gln

-continued

210	215	220
Tyr Pro Val Ile Ile His Leu Ile Trp Met Tyr Gly Thr Ile Phe Phe 225 230 235 240		
Met Leu Phe Ser Asn Phe Trp Tyr His Ser Tyr Thr Lys Gly Lys Arg 245 250 255		
Leu Pro Arg Ala Leu Gln Gln Asn Gly Ala Pro Gly Ile Ala Lys Val 260 265 270		
Lys Ala Asn Xaa 275		

<210> SEQ ID NO 38
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 38

Leu Leu Leu Leu Phe Ile Glu Asn Leu Val Pro Ile Leu Ala Arg Asn 1 5 10 15
Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly Ala Trp Thr Gln Arg Leu 20 25 30
Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val Lys Tyr Trp Glu Leu Ala 35 40 45
Asp Thr Val Phe Leu Val Leu Lys Lys Lys Pro Leu Glu Phe Leu His 50 55 60
Tyr Phe His His Ser Met Thr Met Val Leu Cys Phe Val Gln Leu Gly 65 70 75 80
Gly Tyr Thr Ser Val Ser Trp Val Pro Ile Thr Leu Asn Leu Thr Val 85 90 95
His Val Phe Met Tyr Tyr Tyr Tyr Met Arg Ser Ala Ala Gly Val Arg 100 105 110
Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu Gln Ile Val Gln Phe Val 115 120 125
Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala Tyr Thr Tyr Phe Ala Phe 130 135 140
Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly Lys Cys Ala Gly Thr Glu 145 150 155 160
Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu Ser Ser Tyr Leu Leu Leu 165 170 175
Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn Ala Lys Ala Lys Ala Ala 180 185 190
Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys Thr Val Lys Ser Gly Gly 195 200 205
Ser Pro Lys Lys Pro Ser Lys Ser Lys His Ile 210 215

<210> SEQ ID NO 39
 <211> LENGTH: 204
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 39

Ile Val Tyr Glu Phe Leu Met Ser Gly Trp Leu Ser Thr Tyr Thr Trp 1 5 10 15
Arg Cys Asp Pro Ile Asp Phe Ser Asn Ser Pro Glu Ala Leu Arg Met

-continued

20					25					30					
Val	Arg	Val	Ala	Trp	Leu	Phe	Met	Leu	Ser	Lys	Val	Ile	Glu	Leu	Met
		35					40					45			
Asp	Thr	Val	Ile	Phe	Ile	Leu	Arg	Lys	Lys	Asp	Gly	Gln	Val	Thr	Phe
	50					55					60				
Leu	His	Val	Phe	His	His	Ser	Val	Leu	Pro	Trp	Ser	Trp	Trp	Trp	Gly
65				70						75					80
Ile	Lys	Ile	Ala	Pro	Gly	Gly	Met	Gly	Ser	Phe	His	Ala	Met	Ile	Asn
			85						90					95	
Ser	Ser	Val	His	Val	Val	Met	Tyr	Leu	Tyr	Tyr	Gly	Leu	Ser	Ala	Leu
			100					105					110		
Gly	Pro	Val	Ala	Gln	Pro	Tyr	Leu	Trp	Trp	Lys	Lys	His	Met	Thr	Ala
		115					120					125			
Ile	Gln	Leu	Ile	Gln	Phe	Val	Leu	Val	Ser	Leu	His	Ile	Ser	Gln	Tyr
	130					135					140				
Tyr	Phe	Met	Pro	Ser	Cys	Asn	Tyr	Gln	Tyr	Pro	Val	Ile	Ile	His	Leu
145				150					155						160
Ile	Trp	Met	Tyr	Gly	Thr	Ile	Phe	Phe	Ile	Leu	Phe	Ser	Asn	Phe	Trp
			165						170					175	
Tyr	His	Ser	Tyr	Thr	Lys	Gly	Lys	Arg	Leu	Pro	Arg	Ala	Val	Gln	Gln
			180					185					190		
Asn	Gly	Ala	Pro	Ala	Thr	Thr	Lys	Val	Lys	Ala	Asn				
	195						200								

<210> SEQ ID NO 40

<211> LENGTH: 174

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 40

Tyr	Glu	Leu	Val	Thr	Gly	Lys	Ser	Ile	Asp	Ser	Phe	Val	Phe	Gln	Glu
1				5					10					15	
Gly	Val	Thr	Pro	Leu	Ser	Thr	Gln	Arg	Glu	Val	Ala	Met	Trp	Thr	Ile
			20				25						30		
Thr	Tyr	Phe	Val	Val	Ile	Phe	Gly	Gly	Arg	Gln	Ile	Met	Lys	Ser	Gln
	35					40					45				
Asp	Ala	Phe	Lys	Leu	Lys	Pro	Leu	Phe	Ile	Leu	His	Asn	Phe	Leu	Leu
	50					55					60				
Thr	Ile	Ala	Ser	Gly	Ser	Leu	Leu	Leu	Leu	Phe	Ile	Glu	Asn	Leu	Val
65				70					75					80	
Pro	Ile	Leu	Ala	Arg	Asn	Gly	Leu	Phe	Tyr	Ala	Ile	Cys	Asp	Asp	Gly
			85						90					95	
Ala	Trp	Thr	Gln	Arg	Leu	Glu	Leu	Leu	Tyr	Tyr	Leu	Asn	Tyr	Leu	Val
			100					105					110		
Lys	Tyr	Trp	Glu	Leu	Ala	Asp	Thr	Val	Phe	Leu	Val	Leu	Lys	Lys	Lys
	115					120						125			
Pro	Leu	Glu	Phe	Leu	His	Tyr	Phe	His	His	Ser	Met	Thr	Met	Val	Leu
	130					135						140			
Cys	Phe	Val	Gln	Leu	Gly	Gly	Tyr	Thr	Ser	Val	Ser	Trp	Val	Pro	Ile
145				150					155					160	
Thr	Leu	Asn	Leu	Thr	Val	His	Val	Phe	Met	Tyr	Tyr	Tyr	Tyr		
			165						170						

 -continued

<210> SEQ ID NO 41
 <211> LENGTH: 145
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 41

```

Asn Ala Phe Leu Asp Asn Met Phe Gly Pro Arg Asp Ser Arg Val Arg
 1             5             10             15
Gly Trp Phe Leu Leu Asp Ser Tyr Leu Pro Thr Phe Ile Leu Thr Ile
          20             25             30
Thr Tyr Leu Leu Ser Ile Trp Leu Gly Asn Lys Tyr Met Lys Asn Arg
          35             40             45
Pro Ala Leu Ser Leu Arg Gly Ile Leu Thr Leu Tyr Asn Leu Ala Ile
          50             55             60
Thr Leu Leu Ser Ala Tyr Met Leu Val Glu Leu Ile Leu Ser Ser Trp
65             70             75             80
Glu Gly Gly Tyr Asn Leu Gln Cys Gln Asn Leu Asp Ser Ala Gly Glu
          85             90             95
Gly Asp Val Arg Val Ala Lys Val Leu Val Trp Tyr Tyr Phe Ser Lys
          100            105            110
Leu Val Glu Phe Leu Asp Thr Ile Phe Phe Val Leu Arg Lys Lys Ala
          115            120            125
Asn Gln Ile Thr Phe Leu His Val Tyr His His Ala Ser Met Phe Asn
          130            135            140
Ile
145
  
```

<210> SEQ ID NO 42
 <211> LENGTH: 238
 <212> TYPE: PRT
 <213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 42

```

Leu Ile Val Leu Ala Tyr Leu Val Thr Val Phe Val Gly Met Gln Ile
 1             5             10             15
Met Lys Asn Phe Glu Arg Phe Glu Val Lys Thr Phe Ser Leu Leu His
          20             25             30
Asn Phe Cys Leu Val Ser Ile Ser Ala Tyr Met Cys Gly Gly Ile Leu
          35             40             45
Tyr Glu Ala Tyr Gln Ala Asn Tyr Gly Leu Phe Glu Asn Ala Ala Asp
          50             55             60
His Thr Phe Lys Gly Leu Pro Met Ala Lys Met Ile Trp Leu Phe Tyr
65             70             75             80
Phe Ser Lys Ile Met Glu Phe Val Asp Thr Met Ile Met Val Leu Lys
          85             90             95
Lys Asn Asn Arg Gln Ile Ser Phe Leu His Val Tyr His His Ser Ser
          100            105            110
Ile Phe Thr Ile Trp Trp Leu Val Thr Phe Val Ala Pro Asn Gly Glu
          115            120            125
Ala Tyr Phe Ser Ala Ala Leu Asn Ser Phe Ile His Val Ile Met Tyr
          130            135            140
Gly Tyr Tyr Phe Leu Ser Ala Leu Gly Phe Lys Gln Val Ser Phe Ile
145            150            155            160
  
```

Leu	Tyr	Asn	Leu	Gly	Ile	Thr	Leu	Leu	Ser	Ala	Tyr	Met	Leu	Ala	Glu
1				5					10					15	
Leu	Ile	Leu	Ser	Thr	Trp	Glu	Gly	Gly	Tyr	Asn	Leu	Gln	Cys	Gln	Asp
			20					25					30		
Leu	Thr	Ser	Ala	Gly	Glu	Ala	Asp	Ile	Arg	Val	Ala	Lys	Val	Leu	Trp
			35				40					45			
Trp	Tyr	Tyr	Phe	Ser	Lys	Ser	Val	Glu	Phe	Leu	Asp	Thr	Ile	Phe	Phe
	50					55						60			
Val	Leu	Arg	Lys	Lys	Thr	Ser	Gln	Ile	Thr	Phe	Leu	His	Val	Tyr	His
65					70					75					80
His	Ala	Ser	Met	Phe	Asn	Ile	Trp	Trp	Cys	Val	Leu	Asn	Trp	Ile	Pro
				85					90					95	
Cys	Gly	Gln	Ser	Phe	Phe	Gly	Pro	Thr	Leu	Asn	Ser	Phe	Ile	His	Ile
				100				105					110		
Leu	Met	Tyr	Ser	Tyr	Tyr	Gly	Leu	Ser	Val	Phe	Pro	Ser	Met	His	Lys
				115			120					125			
Tyr	Leu	Trp	Trp	Lys	Lys	Tyr	Leu	Thr	Gln	Ala	Gln	Leu	Val	Gln	Phe
	130					135					140				

Ala	Gln	Ala	Glu	Lys	Tyr	Ile	Pro	Thr	Ile	Val	His	His	Thr	Arg	Gly
1				5					10					15	
Phe	Leu	Val	Ala	Val	Glu	Ser	Pro	Leu	Ala	Arg	Glu	Leu	Pro	Leu	Met
			20					25					30		
Asn	Pro	Phe	His	Val	Leu	Leu	Ile	Val	Leu	Ala	Tyr	Leu	Val	Thr	Val
		35					40					45			
Phe	Val	Gly	Met	Gln	Ile	Met	Lys	Asn	Phe	Glu	Arg	Phe	Glu	Val	Lys
	50					55					60				
Thr	Phe	Ser	Leu	Leu	His	Asn	Phe	Cys	Leu	Val	Ser	Ile	Ser	Ala	Tyr
65					70					75				80	
Met	Cys	Gly	Gly	Ile	Leu	Tyr	Glu	Ala	Tyr	Gln	Ala	Asn	Tyr	Gly	Leu
				85					90					95	

-continued

Phe	Glu	Asn	Ala	Ala	Asp	His	Thr	Phe	Lys	Gly	Leu	Pro	Met	Ala	Lys
			100						105					110	
Met	Ile	Trp	Leu	Phe	Tyr	Phe	Ser	Lys	Ile	Met	Glu	Phe	Val	Asp	Thr
		115					120					125			
Met	Ile	Met	Val	Leu	Lys	Lys	Asn	Asn	Arg	Gln	Ile	Ser	Phe	Leu	His
		130				135					140				
Val	Tyr	His	His	Ser	Ser	Ile	Phe	Thr	Ile	Trp	Trp	Leu	Val	Thr	Phe
145					150					155					160
Val	Ala	Pro	Asn	Gly	Glu	Ala	Tyr	Phe	Ser	Ala	Ala	Leu	Asn	Ser	Phe
				165					170					175	
Ile	His	Val	Ile	Met	Tyr	Gly	Tyr	Tyr	Phe	Leu	Ser	Ala	Leu	Gly	Phe
			180					185						190	
Lys	Gln	Val	Ser	Phe	Ile	Lys	Phe	Tyr	Ile	Thr	Arg	Ser	Gln	Met	Thr
		195					200					205			
Gln	Phe	Cys	Met	Met	Ser	Val	Gln	Ser	Ser	Trp	Asp	Met	Tyr	Ala	Met
	210					215					220				
Lys	Val	Leu	Gly	Arg	Pro	Gly	Tyr	Pro	Phe	Phe	Ile	Thr	Ala	Leu	Leu
225					230					235					240
Trp	Phe	Tyr	Met	Trp	Thr	Met	Leu	Gly	Leu	Phe	Tyr	Asn	Phe	Tyr	Arg
				245					250					255	
Lys	Asn	Ala	Lys	Leu	Ala	Lys	Gln	Ala	Lys	Ala	Asp	Ala	Ala	Lys	Glu
			260				265							270	
Lys	Ala	Arg	Lys	Leu	Gln										
		275													

<210> SEQ ID NO 45

<211> LENGTH: 293

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

Met	Glu	His	Phe	Asp	Ala	Ser	Leu	Ser	Thr	Tyr	Phe	Lys	Ala	Leu	Leu
1				5					10					15	
Gly	Pro	Arg	Asp	Thr	Arg	Val	Lys	Gly	Trp	Phe	Leu	Leu	Asp	Asn	Tyr
			20					25					30		
Ile	Pro	Thr	Phe	Ile	Cys	Ser	Val	Ile	Tyr	Leu	Leu	Ile	Val	Trp	Leu
		35					40					45			
Gly	Pro	Lys	Tyr	Met	Arg	Asn	Lys	Gln	Pro	Phe	Ser	Cys	Arg	Gly	Ile
		50				55				60					
Leu	Val	Val	Tyr	Asn	Leu	Gly	Leu	Thr	Leu	Leu	Ser	Leu	Tyr	Met	Phe
65				70					75					80	
Cys	Glu	Leu	Val	Thr	Gly	Val	Trp	Glu	Gly	Lys	Tyr	Asn	Phe	Phe	Cys
			85						90					95	
Gln	Gly	Thr	Arg	Thr	Ala	Gly	Glu	Ser	Asp	Met	Lys	Ile	Ile	Arg	Val
			100					105					110		
Leu	Trp	Trp	Tyr	Tyr	Phe	Ser	Lys	Leu	Ile	Glu	Phe	Met	Asp	Thr	Phe
		115					120					125			
Phe	Phe	Ile	Leu	Arg	Lys	Asn	Asn	His	Gln	Ile	Thr	Val	Leu	His	Val
		130				135					140				
Tyr	His	His	Ala	Ser	Met	Leu	Asn	Ile	Trp	Trp	Phe	Val	Met	Asn	Trp
145				150						155				160	
Val	Pro	Cys	Gly	His	Ser	Tyr	Phe	Gly	Ala	Thr	Leu	Asn	Ser	Phe	Ile
				165					170					175	

-continued

His Val Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Ser Val Pro Ser Met
 180 185 190
 Arg Pro Tyr Leu Trp Trp Lys Lys Tyr Ile Thr Gln Gly Gln Leu Leu
 195 200 205
 Gln Phe Val Leu Thr Ile Ile Gln Thr Ser Cys Gly Val Ile Trp Pro
 210 215 220
 Cys Thr Phe Pro Leu Gly Trp Leu Tyr Phe Gln Ile Gly Tyr Met Ile
 225 230 235 240
 Ser Leu Ile Ala Leu Phe Thr Asn Phe Tyr Ile Gln Thr Tyr Asn Lys
 245 250 255
 Lys Gly Ala Ser Arg Arg Lys Asp His Leu Lys Asp His Gln Asn Gly
 260 265 270
 Ser Met Ala Ala Val Asn Gly His Thr Asn Ser Phe Ser Pro Leu Glu
 275 280 285
 Asn Asn Val Lys Pro
 290

<210> SEQ ID NO 46
 <211> LENGTH: 182
 <212> TYPE: PRT
 <213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 46

Phe Glu Asn Ala Ala Asp His Thr Phe Lys Gly Leu Pro Met Ala Lys
 1 5 10 15
 Met Ile Trp Leu Phe Tyr Phe Ser Lys Ile Met Glu Phe Val Asp Thr
 20 25 30
 Met Ile Met Val Leu Lys Lys Asn Asn Arg Gln Ile Ser Phe Leu His
 35 40 45
 Val Tyr His His Ser Ser Ile Phe Thr Ile Trp Trp Leu Val Thr Phe
 50 55 60
 Val Ala Pro Asn Gly Glu Ala Tyr Phe Ser Ala Ala Leu Asn Ser Phe
 65 70 75 80
 Ile His Val Ile Met Tyr Gly Tyr Tyr Phe Leu Ser Ala Leu Gly Phe
 85 90 95
 Lys Gln Val Ser Phe Ile Lys Phe Tyr Ile Thr Arg Ser Gln Met Thr
 100 105 110
 Gln Phe Cys Met Met Ser Val Gln Ser Ser Trp Asp Met Tyr Ala Met
 115 120 125
 Lys Val Leu Gly Arg Pro Gly Tyr Pro Phe Phe Ile Thr Ala Leu Leu
 130 135 140
 Trp Phe Tyr Met Trp Thr Met Leu Gly Leu Phe Tyr Asn Phe Tyr Arg
 145 150 155 160
 Lys Asn Ala Lys Leu Ala Lys Gln Ala Lys Ala Asp Ala Ala Lys Glu
 165 170 175
 Lys Ala Arg Lys Leu Gln
 180

<210> SEQ ID NO 47
 <211> LENGTH: 141
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT

-continued

<222> LOCATION: (141)...(141)

<223> OTHER INFORMATION: Xaa = Unknown or Other at position 141

<400> SEQUENCE: 47

```

Asp Thr Ile Phe Ile Ile Leu Arg Lys Gln Lys Leu Ile Phe Leu His
 1             5             10             15

Trp Tyr His His Ile Thr Val Leu Leu Tyr Ser Trp Tyr Ser Tyr Lys
             20             25             30

Asp Met Val Ala Gly Gly Gly Trp Phe Met Thr Met Asn Tyr Gly Val
             35             40             45

His Ala Val Met Tyr Ser Tyr Tyr Ala Leu Arg Ala Ala Gly Phe Arg
             50             55             60

Val Ser Arg Lys Phe Ala Met Phe Ile Thr Leu Ser Gln Ile Thr Gln
65             70             75             80

Met Leu Met Gly Cys Val Val Asn Tyr Leu Val Phe Cys Trp Met Gln
             85             90             95

His Asp Gln Cys His Ser His Phe Gln Asn Ile Phe Trp Ser Ser Leu
             100            105            110

Met Tyr Leu Ser Tyr Leu Val Leu Phe Cys His Phe Phe Phe Glu Ala
             115            120            125

Tyr Ile Gly Lys Met Arg Lys Thr Thr Lys Ala Glu Xaa
             130            135            140

```

<210> SEQ ID NO 48

<211> LENGTH: 241

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 48

```

Leu Leu Ile Val Leu Ala Tyr Leu Val Thr Val Phe Val Gly Met Gln
 1             5             10             15

Ile Met Lys Asn Phe Glu Arg Phe Glu Val Lys Thr Phe Ser Leu Leu
             20             25             30

His Asn Phe Cys Leu Val Ser Ile Ser Ala Tyr Met Cys Gly Gly Ile
             35             40             45

Leu Tyr Glu Ala Tyr Gln Ala Asn Tyr Gly Leu Phe Glu Asn Ala Ala
             50             55             60

Asp His Thr Phe Lys Gly Leu Pro Met Ala Lys Met Ile Trp Leu Phe
65             70             75             80

Tyr Phe Ser Lys Ile Met Glu Phe Val Asp Thr Met Ile Met Val Leu
             85             90             95

Lys Lys Asn Asn Arg Gln Ile Ser Phe Leu His Val Tyr His His Ser
             100            105            110

Ser Ile Phe Thr Ile Trp Trp Leu Val Thr Phe Val Ala Pro Asn Gly
             115            120            125

Glu Ala Tyr Phe Ser Ala Ala Leu Asn Ser Phe Ile His Val Ile Met
             130            135            140

Tyr Gly Tyr Tyr Phe Leu Ser Ala Leu Gly Phe Lys Gln Val Ser Phe
145            150            155            160

Ile Lys Phe Tyr Ile Thr Arg Ser Gln Met Thr Gln Phe Cys Met Met
             165            170            175

Ser Val Gln Ser Ser Trp Asp Met Tyr Ala Met Lys Val Leu Gly Arg
             180            185            190

```

-continued

Pro Gly Tyr Pro Phe Phe Ile Thr Ala Leu Leu Trp Phe Tyr Met Trp
 195 200 205
 Thr Met Leu Gly Leu Phe Tyr Asn Phe Tyr Arg Lys Asn Ala Lys Leu
 210 215 220
 Ala Lys Gln Ala Lys Ala Asp Ala Ala Lys Glu Lys Ala Arg Lys Leu
 225 230 235 240

Gln

<210> SEQ ID NO 49
 <211> LENGTH: 217
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 49

Ile Val Tyr Asn Phe Ser Leu Val Ile Leu Ser Leu Tyr Ile Val Tyr
 1 5 10 15
 Glu Phe Leu Met Ser Gly Trp Leu Ser Thr Tyr Thr Trp Arg Cys Asp
 20 25 30
 Pro Ile Asp Phe Ser Asn Ser Pro Glu Ala Leu Arg Met Val Arg Val
 35 40 45
 Ala Trp Leu Phe Met Leu Ser Lys Val Ile Glu Leu Met Asp Thr Val
 50 55 60
 Ile Phe Ile Leu Arg Lys Lys Asp Gly Gln Val Thr Phe Leu His Val
 65 70 75 80
 Phe His His Ser Val Leu Pro Trp Ser Trp Trp Trp Gly Ile Lys Ile
 85 90 95
 Ala Pro Gly Gly Met Gly Ser Phe His Ala Met Ile Asn Ser Ser Val
 100 105 110
 His Val Val Met Tyr Leu Tyr Tyr Gly Leu Ser Ala Leu Gly Pro Val
 115 120 125
 Ala Gln Pro Tyr Leu Trp Trp Lys Lys His Met Thr Ala Ile Gln Leu
 130 135 140
 Ile Gln Phe Val Leu Val Ser Leu His Ile Ser Gln Tyr Tyr Phe Met
 145 150 155 160
 Pro Ser Cys Asn Tyr Gln Tyr Pro Val Ile Ile His Leu Ile Trp Met
 165 170 175
 Tyr Gly Thr Ile Phe Phe Ile Leu Phe Ser Asn Phe Trp Tyr His Ser
 180 185 190
 Tyr Thr Lys Gly Lys Arg Leu Pro Arg Ala Val Gln Gln Asn Gly Ala
 195 200 205
 Pro Ala Thr Thr Lys Val Lys Ala Asn
 210 215

<210> SEQ ID NO 50
 <211> LENGTH: 178
 <212> TYPE: PRT
 <213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 50

Pro Thr Ile Val His His Thr Arg Gly Phe Leu Val Ala Val Glu Ser
 1 5 10 15
 Pro Leu Ala Arg Glu Leu Pro Leu Met Asn Pro Phe His Val Leu Leu
 20 25 30
 Ile Val Leu Ala Tyr Leu Val Thr Val Phe Val Gly Met Gln Ile Met

-continued

35	40	45
Lys Asn Phe Glu Arg Phe Glu Val Lys Thr Phe Ser Leu Leu His Asn 50 55 60		
Phe Cys Leu Val Ser Ile Ser Ala Tyr Met Cys Gly Gly Ile Leu Tyr 65 70 75 80		
Glu Ala Tyr Gln Ala Asn Tyr Gly Leu Phe Glu Asn Ala Ala Asp His 85 90 95		
Thr Phe Lys Gly Leu Pro Met Ala Lys Met Ile Trp Leu Phe Tyr Phe 100 105 110		
Ser Lys Ile Met Glu Phe Val Asp Thr Met Ile Met Val Leu Lys Lys 115 120 125		
Asn Asn Arg Gln Ile Ser Phe Leu His Val Tyr His His Ser Ser Ile 130 135 140		
Phe Thr Ile Trp Trp Leu Val Thr Phe Val Ala Pro Asn Gly Glu Ala 145 150 155 160		
Tyr Phe Ser Ala Ala Leu Asn Ser Phe Ile His Val Ile Met Tyr Gly 165 170 175		
Tyr Tyr		

<210> SEQ ID NO 51
<211> LENGTH: 148
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 51

Asn Glu Val Asn Ala Phe Leu Asp Asn Met Phe Gly Pro Arg Asp Ser 1 5 10 15	
Arg Val Arg Gly Trp Phe Leu Leu Asp Ser Tyr Leu Pro Thr Phe Ile 20 25 30	
Leu Thr Ile Thr Tyr Leu Leu Ser Ile Trp Leu Gly Asn Lys Tyr Met 35 40 45	
Lys Asn Arg Pro Ala Leu Ser Leu Arg Gly Ile Leu Thr Leu Tyr Asn 50 55 60	
Leu Ala Ile Thr Leu Leu Ser Ala Tyr Met Leu Val Glu Leu Ile Leu 65 70 75 80	
Ser Ser Trp Glu Gly Gly Tyr Asn Leu Gln Cys Gln Asn Leu Asp Ser 85 90 95	
Ala Gly Glu Gly Asp Val Arg Val Ala Lys Val Leu Val Trp Tyr Tyr 100 105 110	
Phe Ser Lys Leu Val Glu Phe Leu Asp Thr Ile Phe Phe Val Leu Arg 115 120 125	
Lys Lys Ala Asn Gln Ile Thr Phe Leu His Val Tyr His His Ala Ser 130 135 140	
Met Phe Asn Ile 145	

<210> SEQ ID NO 52
<211> LENGTH: 302
<212> TYPE: PRT
<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 52

Phe Met Asp Leu Ala Thr Ala Ile Gly Val Arg Ala Ala Pro Tyr Val 1 5 10 15	
--	--

-continued

```

Asp Pro Leu Glu Ala Ala Leu Val Ala Gln Ala Glu Lys Tyr Ile Pro
      20                      25                      30

Thr Ile Val His His Thr Arg Gly Phe Leu Val Ala Val Glu Ser Pro
      35                      40                      45

Leu Ala Arg Glu Leu Pro Leu Met Asn Pro Phe His Val Leu Leu Ile
      50                      55                      60

Val Leu Ala Tyr Leu Val Thr Val Phe Val Gly Met Gln Ile Met Lys
      65                      70                      75                      80

Asn Phe Glu Arg Phe Glu Val Lys Thr Phe Ser Leu Leu His Asn Phe
      85                      90                      95

Cys Leu Val Ser Ile Ser Ala Tyr Met Cys Gly Gly Ile Leu Tyr Glu
      100                     105                     110

Ala Tyr Gln Ala Asn Tyr Gly Leu Phe Glu Asn Ala Ala Asp His Thr
      115                     120                     125

Phe Lys Gly Leu Pro Met Ala Lys Met Ile Trp Leu Phe Tyr Phe Ser
      130                     135                     140

Lys Ile Met Glu Phe Val Asp Thr Met Ile Met Val Leu Lys Lys Asn
      145                     150                     155                     160

Asn Arg Gln Ile Ser Phe Leu His Val Tyr His His Ser Ser Ile Phe
      165                     170                     175

Thr Ile Trp Trp Leu Val Thr Phe Val Ala Pro Asn Gly Glu Ala Tyr
      180                     185                     190

Phe Ser Ala Ala Leu Asn Ser Phe Ile His Val Ile Met Tyr Gly Tyr
      195                     200                     205

Tyr Phe Leu Ser Ala Leu Gly Phe Lys Gln Val Ser Phe Ile Lys Phe
      210                     215                     220

Tyr Ile Thr Arg Ser Gln Met Thr Gln Phe Cys Met Met Ser Val Gln
      225                     230                     235                     240

Ser Ser Trp Asp Met Tyr Ala Met Lys Val Leu Gly Arg Pro Gly Tyr
      245                     250                     255

Pro Phe Phe Ile Thr Ala Leu Leu Trp Phe Tyr Met Trp Thr Met Leu
      260                     265                     270

Gly Leu Phe Tyr Asn Phe Tyr Arg Lys Asn Ala Lys Leu Ala Lys Gln
      275                     280                     285

Ala Lys Ala Asp Ala Ala Lys Glu Lys Ala Arg Lys Leu Gln
      290                     295                     300

```

<210> SEQ ID NO 53

<211> LENGTH: 271

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 53

```

Met Asp Thr Ser Met Asn Phe Ser Arg Gly Leu Lys Met Asp Leu Met
  1              5              10              15

Gln Pro Tyr Asp Phe Glu Thr Phe Gln Asp Leu Arg Pro Phe Leu Glu
      20              25              30

Glu Tyr Trp Val Ser Ser Phe Leu Ile Val Val Val Tyr Leu Leu Leu
      35              40              45

Ile Val Val Gly Gln Thr Tyr Met Arg Thr Arg Lys Ser Phe Ser Leu
      50              55              60

Gln Arg Pro Leu Ile Leu Trp Ser Phe Phe Leu Ala Ile Phe Ser Ile

```

-continued

65		70		75		80
Leu Gly Thr	Leu Arg	Met Trp Lys Phe	Met Ala Thr Val Met Phe Thr			
	85		90		95	
Val Gly Leu	Lys Gln Thr Val Cys Phe Ala Ile Tyr Thr Asp Asp Ala					
	100		105		110	
Val Val Arg	Phe Trp Ser Phe Leu Phe Leu Leu Ser Lys Val Val Glu					
	115		120		125	
Leu Gly Asp	Thr Ala Phe Ile Ile Leu Arg Lys Arg Pro Leu Ile Phe					
	130		135		140	
Val His Trp	Tyr His His Ser Thr Val Leu Leu Phe Thr Ser Phe Gly					
	145		150		155	
Tyr Lys Asn	Lys Val Pro Ser Gly Gly Trp Phe Met Thr Met Asn Phe					
	165		170		175	
Gly Val His	Ser Val Met Tyr Thr Tyr Tyr Thr Met Lys Ala Ala Lys					
	180		185		190	
Leu Lys His	Pro Asn Leu Leu Pro Met Val Ile Thr Ser Leu Gln Ile					
	195		200		205	
Leu Gln Met	Val Leu Gly Thr Ile Phe Gly Ile Leu Asn Tyr Ile Trp					
	210		215		220	
Arg Gln Glu	Lys Gly Cys His Thr Thr Thr Glu His Phe Phe Trp Ser					
	225		230		235	
Phe Met Leu	Tyr Gly Thr Tyr Phe Ile Leu Phe Ala His Phe Phe His					
	245		250		255	
Arg Ala Tyr	Leu Arg Pro Lys Gly Lys Val Ala Ser Lys Ser Gln					
	260		265		270	

<210> SEQ ID NO 54

<211> LENGTH: 265

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 54

Thr Arg Gly	Phe Leu Val Ala Val Glu Ser Pro Leu Ala Arg Glu Leu
1	5 10 15
Pro Leu Met	Asn Pro Phe His Val Leu Leu Ile Val Leu Ala Tyr Leu
	20 25 30
Val Thr Val	Phe Val Gly Met Gln Ile Met Lys Asn Phe Glu Arg Phe
	35 40 45
Glu Val Lys	Thr Phe Ser Leu Leu His Asn Phe Cys Leu Val Ser Ile
	50 55 60
Ser Ala Tyr	Met Cys Gly Gly Ile Leu Tyr Glu Ala Tyr Gln Ala Asn
65	70 75 80
Tyr Gly Leu	Phe Glu Asn Ala Ala Asp His Thr Phe Lys Gly Leu Pro
	85 90 95
Met Ala Lys	Met Ile Trp Leu Phe Tyr Phe Ser Lys Ile Met Glu Phe
	100 105 110
Val Asp Thr	Met Ile Met Val Leu Lys Lys Asn Asn Arg Gln Ile Ser
	115 120 125
Phe Leu His	Val Tyr His His Ser Ser Ile Phe Thr Ile Trp Trp Leu
	130 135 140
Val Thr Phe	Val Ala Pro Asn Gly Glu Ala Tyr Phe Ser Ala Ala Leu
145	150 155 160

```
<210> SEQ ID NO 55
<211> LENGTH: 265
<212> TYPE: PRT
<213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 55
```

Ala	Thr	His	Gly	Pro	Lys	Asn	Phe	Pro	Asp	Ala	Glu	Gly	Arg	Lys	Phe
1				5					10					15	
Phe	Ala	Asp	His	Phe	Asp	Val	Thr	Ile	Gln	Ala	Ser	Ile	Leu	Tyr	Met
			20					25					30		
Val	Val	Val	Phe	Gly	Thr	Lys	Trp	Phe	Met	Arg	Asn	Arg	Gln	Pro	Phe
		35					40					45			
Gln	Leu	Thr	Ile	Pro	Leu	Asn	Ile	Trp	Asn	Phe	Ile	Leu	Ala	Ala	Phe
	50					55					60				
Ser	Ile	Ala	Gly	Ala	Val	Lys	Met	Thr	Pro	Glu	Phe	Phe	Gly	Thr	Ile
65					70					75					80
Ala	Asn	Lys	Gly	Ile	Val	Ala	Ser	Tyr	Cys	Lys	Val	Phe	Asp	Phe	Thr
				85					90					95	
Lys	Gly	Glu	Asn	Gly	Tyr	Trp	Val	Trp	Leu	Phe	Met	Ala	Ser	Lys	Leu
			100					105					110		
Phe	Glu	Leu	Val	Asp	Thr	Ile	Phe	Leu	Val	Leu	Arg	Lys	Arg	Pro	Leu
		115					120					125			
Met	Phe	Leu	His	Trp	Tyr	His	His	Ile	Leu	Thr	Met	Ile	Tyr	Ala	Trp
	130					135					140				
Tyr	Ser	His	Pro	Leu	Thr	Pro	Gly	Phe	Asn	Arg	Tyr	Gly	Ile	Tyr	Leu
145					150					155					160
Asn	Phe	Val	Val	His	Ala	Phe	Met	Tyr	Ser	Tyr	Tyr	Phe	Leu	Arg	Ser
				165					170					175	
Met	Lys	Ile	Arg	Val	Pro	Gly	Phe	Ile	Ala	Gln	Ala	Ile	Thr	Ser	Leu
		180						185					190		
Gln	Ile	Val	Gln	Phe	Ile	Ile	Ser	Cys	Ala	Val	Leu	Ala	His	Leu	Gly
		195					200					205			
Tyr	Leu	Met	His	Phe	Thr	Asn	Ala	Asn	Cys	Asp	Phe	Glu	Pro	Ser	Val
	210					215					220				
Phe	Lys	Leu	Ala	Val	Phe	Met	Asp	Thr	Thr	Tyr	Leu	Ala	Leu	Phe	Val
225					230					235					240
Asn	Phe	Phe	Leu	Gln	Ser	Tyr	Val	Leu	Arg	Gly	Gly	Lys	Asp	Lys	Tyr
				245					250					255	

 -continued

Lys Ala Val Pro Lys Lys Lys Asn Asn
 260 265

<210> SEQ ID NO 56

<211> LENGTH: 288

<212> TYPE: PRT

<213> ORGANISM: *Caenorhabditis elegans*

<400> SEQUENCE: 56

Met Ala Gln His Pro Leu Val Gln Arg Leu Leu Asp Val Lys Phe Asp
 1 5 10 15
 Thr Lys Arg Phe Val Ala Ile Ala Thr His Gly Pro Lys Asn Phe Pro
 20 25 30
 Asp Ala Glu Gly Arg Lys Phe Phe Ala Asp His Phe Asp Val Thr Ile
 35 40 45
 Gln Ala Ser Ile Leu Tyr Met Val Val Val Phe Gly Thr Lys Trp Phe
 50 55 60
 Met Arg Asn Arg Gln Pro Phe Gln Leu Thr Ile Pro Leu Asn Ile Trp
 65 70 75 80
 Asn Phe Ile Leu Ala Ala Phe Ser Ile Ala Gly Ala Val Lys Met Thr
 85 90 95
 Pro Glu Phe Phe Gly Thr Ile Ala Asn Lys Gly Ile Val Ala Ser Tyr
 100 105 110
 Cys Lys Val Phe Asp Phe Thr Lys Gly Glu Asn Gly Tyr Trp Val Trp
 115 120 125
 Leu Phe Met Ala Ser Lys Leu Phe Glu Leu Val Asp Thr Ile Phe Leu
 130 135 140
 Val Leu Arg Lys Arg Pro Leu Met Phe Leu His Trp Tyr His His Ile
 145 150 155 160
 Leu Thr Met Ile Tyr Ala Trp Tyr Ser His Pro Leu Thr Pro Gly Phe
 165 170 175
 Asn Arg Tyr Gly Ile Tyr Leu Asn Phe Val Val His Ala Phe Met Tyr
 180 185 190
 Ser Tyr Tyr Phe Leu Arg Ser Met Lys Ile Arg Val Pro Gly Phe Ile
 195 200 205
 Ala Gln Ala Ile Thr Ser Leu Gln Ile Val Gln Phe Ile Ile Ser Cys
 210 215 220
 Ala Val Leu Ala His Leu Gly Tyr Leu Met His Phe Thr Asn Ala Asn
 225 230 235 240
 Cys Asp Phe Glu Pro Ser Val Phe Lys Leu Ala Val Phe Met Asp Thr
 245 250 255
 Thr Tyr Leu Ala Leu Phe Val Asn Phe Phe Leu Gln Ser Tyr Val Leu
 260 265 270
 Arg Gly Gly Lys Asp Lys Tyr Lys Ala Val Pro Lys Lys Lys Asn Asn
 275 280 285

<210> SEQ ID NO 57

<211> LENGTH: 282

<212> TYPE: PRT

<213> ORGANISM: *Mortierella alpina*

<400> SEQUENCE: 57

Ala Ala Leu Val Ala Gln Ala Glu Lys Tyr Ile Pro Thr Ile Val His
 1 5 10 15

-continued

His Thr Arg Gly Phe Leu Val Ala Val Glu Ser Pro Leu Ala Arg Glu
 20 25 30
 Leu Pro Leu Met Asn Pro Phe His Val Leu Leu Ile Val Leu Ala Tyr
 35 40 45
 Leu Val Thr Val Phe Val Gly Met Gln Ile Met Lys Asn Phe Glu Arg
 50 55 60
 Phe Glu Val Lys Thr Phe Ser Leu Leu His Asn Phe Cys Leu Val Ser
 65 70 75 80
 Ile Ser Ala Tyr Met Cys Gly Gly Ile Leu Tyr Glu Ala Tyr Gln Ala
 85 90 95
 Asn Tyr Gly Leu Phe Glu Asn Ala Ala Asp His Thr Phe Lys Gly Leu
 100 105 110
 Pro Met Ala Lys Met Ile Trp Leu Phe Tyr Phe Ser Lys Ile Met Glu
 115 120 125
 Phe Val Asp Thr Met Ile Met Val Leu Lys Lys Asn Asn Arg Gln Ile
 130 135 140
 Ser Phe Leu His Val Tyr His His Ser Ser Ile Phe Thr Ile Trp Trp
 145 150 155 160
 Leu Val Thr Phe Val Ala Pro Asn Gly Glu Ala Tyr Phe Ser Ala Ala
 165 170 175
 Leu Asn Ser Phe Ile His Val Ile Met Tyr Gly Tyr Tyr Phe Leu Ser
 180 185 190
 Ala Leu Gly Phe Lys Gln Val Ser Phe Ile Lys Phe Tyr Ile Thr Arg
 195 200 205
 Ser Gln Met Thr Gln Phe Cys Met Met Ser Val Gln Ser Ser Trp Asp
 210 215 220
 Met Tyr Ala Met Lys Val Leu Gly Arg Pro Gly Tyr Pro Phe Phe Ile
 225 230 235 240
 Thr Ala Leu Leu Trp Phe Tyr Met Trp Thr Met Leu Gly Leu Phe Tyr
 245 250 255
 Asn Phe Tyr Arg Lys Asn Ala Lys Leu Ala Lys Gln Ala Lys Ala Asp
 260 265 270
 Ala Ala Lys Glu Lys Ala Arg Lys Leu Gln
 275 280

<210> SEQ ID NO 58
 <211> LENGTH: 278
 <212> TYPE: PRT
 <213> ORGANISM: Drosophila melanogaster
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (235)...(235)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 235

<400> SEQUENCE: 58

Pro Thr Lys Met Ile Asn Met Asp Ile Ser Val Thr Pro Asn Tyr Ser
 1 5 10 15
 Tyr Ile Phe Asp Phe Glu Asn Asp Phe Ile His Gln Arg Thr Arg Lys
 20 25 30
 Trp Met Leu Glu Asn Trp Thr Trp Val Phe Tyr Tyr Cys Gly Ile Tyr
 35 40 45
 Met Leu Val Ile Phe Gly Gly Gln His Phe Met Gln Asn Arg Pro Arg
 50 55 60

-continued

Phe Gln Leu Arg Gly Pro Leu Ile Ile Trp Asn Thr Leu Leu Ala Met
 65 70 75 80
 Phe Ser Ile Met Gly Ala Ala Arg Thr Ala Pro Glu Leu Ile His Val
 85 90 95
 Leu Arg His Tyr Gly Leu Phe His Ser Val Cys Val Pro Ser Tyr Ile
 100 105 110
 Glu Gln Asp Arg Val Cys Gly Phe Trp Thr Trp Leu Phe Val Leu Ser
 115 120 125
 Lys Leu Pro Glu Leu Gly Asp Thr Ile Phe Ile Val Leu Arg Lys Gln
 130 135 140
 Pro Leu Ile Phe Leu His Trp Tyr His His Ile Thr Val Leu Ile Tyr
 145 150 155 160
 Ser Trp Phe Ser Tyr Thr Glu Tyr Thr Ser Ser Ala Arg Trp Phe Ile
 165 170 175
 Val Met Asn Tyr Cys Val His Ser Val Met Tyr Ser Tyr Tyr Ala Leu
 180 185 190
 Lys Ala Ala Arg Phe Asn Pro Pro Arg Phe Ile Ser Met Ile Ile Thr
 195 200 205
 Ser Leu Gln Leu Ala Gln Met Ile Ile Gly Cys Ala Ile Asn Val Trp
 210 215 220
 Ala Asn Gly Phe Leu Lys Thr His Gly Thr Xaa Ser Cys His Ile Ser
 225 230 235 240
 Gln Arg Asn Ile Asn Leu Ser Ile Ala Met Tyr Ser Ser Tyr Phe Val
 245 250 255
 Leu Phe Ala Arg Phe Phe Tyr Lys Ala Tyr Leu Ala Pro Gly Gly His
 260 265 270
 Lys Ser Arg Arg Met Ala
 275

<210> SEQ ID NO 59

<211> LENGTH: 286

<212> TYPE: PRT

<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 59

Val Thr Gly Lys Ser Ile Asp Ser Phe Val Phe Gln Glu Gly Val Thr
 1 5 10 15
 Pro Leu Ser Thr Gln Arg Glu Val Ala Met Trp Thr Ile Thr Tyr Phe
 20 25 30
 Val Val Ile Phe Gly Gly Arg Gln Ile Met Lys Ser Gln Asp Ala Phe
 35 40 45
 Lys Leu Lys Pro Leu Phe Ile Leu His Asn Phe Leu Leu Thr Ile Ala
 50 55 60
 Ser Gly Ser Leu Leu Leu Leu Phe Ile Glu Asn Leu Val Pro Ile Leu
 65 70 75 80
 Ala Arg Asn Gly Leu Phe Tyr Ala Ile Cys Asp Asp Gly Ala Trp Thr
 85 90 95
 Gln Arg Leu Glu Leu Leu Tyr Tyr Leu Asn Tyr Leu Val Lys Tyr Trp
 100 105 110
 Glu Leu Ala Asp Thr Val Phe Leu Val Leu Lys Lys Lys Pro Leu Glu
 115 120 125
 Phe Leu His Tyr Phe His His Ser Met Thr Met Val Leu Cys Phe Val
 130 135 140

-continued

Gln Leu Gly Gly Tyr Thr Ser Val Ser Trp Val Pro Ile Thr Leu Asn
 145 150 155 160
 Leu Thr Val His Val Phe Met Tyr Tyr Tyr Met Arg Ser Ala Ala
 165 170 175
 Gly Val Arg Ile Trp Trp Lys Gln Tyr Leu Thr Thr Leu Gln Ile Val
 180 185 190
 Gln Phe Val Leu Asp Leu Gly Phe Ile Tyr Phe Cys Ala Tyr Thr Tyr
 195 200 205
 Phe Ala Phe Thr Tyr Phe Pro Trp Ala Pro Asn Val Gly Lys Cys Ala
 210 215 220
 Gly Thr Glu Gly Ala Ala Leu Phe Gly Cys Gly Leu Leu Ser Ser Tyr
 225 230 235 240
 Leu Leu Leu Phe Ile Asn Phe Tyr Arg Ile Thr Tyr Asn Ala Lys Ala
 245 250 255
 Lys Ala Ala Lys Glu Arg Gly Ser Asn Phe Thr Pro Lys Thr Val Lys
 260 265 270
 Ser Gly Gly Ser Pro Lys Lys Pro Ser Lys Ser Lys His Ile
 275 280 285

<210> SEQ ID NO 60
 <211> LENGTH: 261
 <212> TYPE: PRT
 <213> ORGANISM: Drosophila melanogaster
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (218)...(218)
 <223> OTHER INFORMATION: Xaa = Unknown or Other at position 218
 <400> SEQUENCE: 60

Ile Phe Asp Phe Glu Asn Asp Phe Ile His Gln Arg Thr Arg Lys Trp
 1 5 10 15
 Met Leu Glu Asn Trp Thr Trp Val Phe Tyr Tyr Cys Gly Ile Tyr Met
 20 25 30
 Leu Val Ile Phe Gly Gly Gln His Phe Met Gln Asn Arg Pro Arg Phe
 35 40 45
 Gln Leu Arg Gly Pro Leu Ile Ile Trp Asn Thr Leu Leu Ala Met Phe
 50 55 60
 Ser Ile Met Gly Ala Ala Arg Thr Ala Pro Glu Leu Ile His Val Leu
 65 70 75 80
 Arg His Tyr Gly Leu Phe His Ser Val Cys Val Pro Ser Tyr Ile Glu
 85 90 95
 Gln Asp Arg Val Cys Gly Phe Trp Thr Trp Leu Phe Val Leu Ser Lys
 100 105 110
 Leu Pro Glu Leu Gly Asp Thr Ile Phe Ile Val Leu Arg Lys Gln Pro
 115 120 125
 Leu Ile Phe Leu His Trp Tyr His His Ile Thr Val Leu Ile Tyr Ser
 130 135 140
 Trp Phe Ser Tyr Thr Glu Tyr Thr Ser Ser Ala Arg Trp Phe Ile Val
 145 150 155 160
 Met Asn Tyr Cys Val His Ser Val Met Tyr Ser Tyr Tyr Ala Leu Lys
 165 170 175
 Ala Ala Arg Phe Asn Pro Pro Arg Phe Ile Ser Met Ile Ile Thr Ser
 180 185 190

-continued

Leu Gln Leu Ala Gln Met Ile Ile Gly Cys Ala Ile Asn Val Trp Ala
 195 200 205

Asn Gly Phe Leu Lys Thr His Gly Thr Xaa Ser Cys His Ile Ser Gln
 210 215 220

Arg Asn Ile Asn Leu Ser Ile Ala Met Tyr Ser Ser Tyr Phe Val Leu
 225 230 235 240

Phe Ala Arg Phe Phe Tyr Lys Ala Tyr Leu Ala Pro Gly Gly His Lys
 245 250 255

Ser Arg Arg Met Ala
 260

<210> SEQ ID NO 61
 <211> LENGTH: 299
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

Met Glu His Phe Asp Ala Ser Leu Ser Thr Tyr Phe Lys Ala Leu Leu
 1 5 10 15

Gly Pro Arg Asp Thr Arg Val Lys Gly Trp Phe Leu Leu Asp Asn Tyr
 20 25 30

Ile Pro Thr Phe Ile Cys Ser Val Ile Tyr Leu Leu Ile Val Trp Leu
 35 40 45

Gly Pro Lys Tyr Met Arg Asn Lys Gln Pro Phe Ser Cys Arg Gly Ile
 50 55 60

Leu Val Val Tyr Asn Leu Gly Leu Thr Leu Leu Ser Leu Tyr Met Phe
 65 70 75 80

Cys Glu Leu Val Thr Gly Val Trp Glu Gly Lys Tyr Asn Phe Phe Cys
 85 90 95

Gln Gly Thr Arg Thr Ala Gly Glu Ser Asp Met Lys Ile Ile Arg Val
 100 105 110

Leu Trp Trp Tyr Tyr Phe Ser Lys Leu Ile Glu Phe Met Asp Thr Phe
 115 120 125

Phe Phe Ile Leu Arg Lys Asn Asn His Gln Ile Thr Val Leu His Val
 130 135 140

Tyr His His Ala Ser Met Leu Asn Ile Trp Trp Phe Val Met Asn Trp
 145 150 155 160

Val Pro Cys Gly His Ser Tyr Phe Gly Ala Thr Leu Asn Ser Phe Ile
 165 170 175

His Val Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Ser Val Pro Ser Met
 180 185 190

Arg Pro Tyr Leu Trp Trp Lys Lys Tyr Ile Thr Gln Gly Gln Leu Leu
 195 200 205

Gln Phe Val Leu Thr Ile Ile Gln Thr Ser Cys Gly Val Ile Trp Pro
 210 215 220

Cys Thr Phe Pro Leu Gly Trp Leu Tyr Phe Gln Ile Gly Tyr Ile Ile
 225 230 235 240

Ser Leu Ile Ala Leu Phe Thr Asn Phe Tyr Ile Gln Thr Tyr Asn Lys
 245 250 255

Lys Gly Ala Ser Arg Arg Lys Asp His Leu Lys Asp His Gln Asn Gly
 260 265 270

Ser Val Ala Ala Val Asn Gly His Thr Asn Ser Phe Ser Pro Leu Glu
 275 280 285

-continued

Asn Asn Val Lys Pro Arg Lys Leu Arg Lys Asp
290 295

<210> SEQ ID NO 62

<211> LENGTH: 288

<212> TYPE: PRT

<213> ORGANISM: *Caenorhabditis elegans*

<400> SEQUENCE: 62

Met Ala Gln His Pro Leu Val Gln Arg Leu Leu Asp Val Lys Phe Asp
1 5 10 15
Thr Lys Arg Phe Val Ala Ile Ala Thr His Gly Pro Lys Asn Phe Pro
20 25 30
Asp Ala Glu Gly Arg Lys Phe Phe Ala Asp His Phe Asp Val Thr Ile
35 40 45
Gln Ala Ser Ile Leu Tyr Met Val Val Val Phe Gly Thr Lys Trp Phe
50 55 60
Met Arg Asn Arg Gln Pro Phe Gln Leu Thr Ile Pro Leu Asn Ile Trp
65 70 75 80
Asn Phe Ile Leu Ala Ala Phe Ser Ile Ala Gly Ala Val Lys Met Thr
85 90 95
Pro Glu Phe Phe Gly Thr Ile Ala Asn Lys Gly Ile Val Ala Ser Tyr
100 105 110
Cys Lys Val Phe Asp Phe Thr Lys Gly Glu Asn Gly Tyr Trp Val Trp
115 120 125
Leu Phe Met Ala Ser Lys Leu Phe Glu Leu Val Asp Thr Ile Phe Leu
130 135 140
Val Leu Arg Lys Arg Pro Leu Met Phe Leu His Trp Tyr His His Ile
145 150 155 160
Leu Thr Met Ile Tyr Ala Trp Tyr Ser His Pro Leu Thr Pro Gly Phe
165 170 175
Asn Arg Tyr Gly Ile Tyr Leu Asn Phe Val Val His Ala Phe Met Tyr
180 185 190
Ser Tyr Tyr Phe Leu Arg Ser Met Lys Ile Arg Val Pro Gly Phe Ile
195 200 205
Ala Gln Ala Ile Thr Ser Leu Gln Ile Val Gln Phe Ile Ile Ser Cys
210 215 220
Ala Val Leu Ala His Leu Gly Tyr Leu Met His Phe Thr Asn Ala Asn
225 230 235 240
Cys Asp Phe Glu Pro Ser Val Phe Lys Leu Ala Val Phe Met Asp Thr
245 250 255
Thr Tyr Leu Ala Leu Phe Val Asn Phe Phe Leu Gln Ser Tyr Val Leu
260 265 270
Arg Gly Gly Lys Asp Lys Tyr Lys Ala Val Pro Lys Lys Lys Asn Asn
275 280 285

<210> SEQ ID NO 63

<211> LENGTH: 798

<212> TYPE: DNA

<213> ORGANISM: *Homo sapiens*

<400> SEQUENCE: 63

atgaacatgt cagtgttgac ttacaagaa tatgaattcg aaaagcagtt caacgagaat 60

-continued

```

gaagccatcc aatggatgca ggaaaactgg aagaaatcctt tcctgttttc tgctctgtat    120
gctgccttta tattcggtagg tcggcaccta atgaataaac gagcaaagtt tgaactgagg    180
aagccattag tgctctggct tctgaccctt gcagtcctca gtatattcgg tgctcttcga    240
actggtgctt atatggtgta cattttgatg accaaaggcc tgaagcagtc agtttgtgac    300
cagggttttt acaatggacc tgtcagcaaa ttctgggctt atgcatttgt gctaagcaaa    360
gcacccgaac taggagatac aatattcatt attctgagga agcagaagct gatcttcctg    420
cactggtatc accacatcac tgtgctcctg tactcttggt actcctacaa agacatggtt    480
gccgggggag gttggttcat gactatgaac tatggcgtgc acgccgtgat gtactcttac    540
tatgccttgc gggcggcagg ttcccgagtc tcccgaagt ttgccatgtt catcaccttg    600
tcccatgatc ctcagatgct gatgggctgt gtggttaact acctggtctt ctgctggatg    660
cagcatgacc agtgtcactc tcactttcag aacatcttct ggtcctcact catgtacctc    720
agctaccttg tgctcttctg ccatttcttc tttaggcctt acatcggcaa aatgaggaaa    780
acaacgaaag ctgaatag                                     798

```

<210> SEQ ID NO 64

<211> LENGTH: 265

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

```

Met Asn Met Ser Val Leu Thr Leu Gln Glu Tyr Glu Phe Glu Lys Gln
 1             5             10            15
Phe Asn Glu Asn Glu Ala Ile Gln Trp Met Gln Glu Asn Trp Lys Lys
      20             25             30
Ser Phe Leu Phe Ser Ala Leu Tyr Ala Ala Phe Ile Phe Gly Gly Arg
      35             40             45
His Leu Met Asn Lys Arg Ala Lys Phe Glu Leu Arg Lys Pro Leu Val
      50             55             60
Leu Trp Ser Leu Thr Leu Ala Val Phe Ser Ile Phe Gly Ala Leu Arg
      65             70             75             80
Thr Gly Ala Tyr Met Val Tyr Ile Leu Met Thr Lys Gly Leu Lys Gln
      85             90             95
Ser Val Cys Asp Gln Gly Phe Tyr Asn Gly Pro Val Ser Lys Phe Trp
      100            105            110
Ala Tyr Ala Phe Val Leu Ser Lys Ala Pro Glu Leu Gly Asp Thr Ile
      115            120            125
Phe Ile Ile Leu Arg Lys Gln Lys Leu Ile Phe Leu His Trp Tyr His
      130            135            140
His Ile Thr Val Leu Leu Tyr Ser Trp Tyr Ser Tyr Lys Asp Met Val
      145            150            155            160
Ala Gly Gly Gly Trp Phe Met Thr Met Asn Tyr Gly Val His Ala Val
      165            170            175
Met Tyr Ser Tyr Tyr Ala Leu Arg Ala Ala Gly Phe Arg Val Ser Arg
      180            185            190
Lys Phe Ala Met Phe Ile Thr Leu Ser Gln Ile Thr Gln Met Leu Met
      195            200            205
Gly Cys Val Val Asn Tyr Leu Val Phe Cys Trp Met Gln His Asp Gln
      210            215            220

```

-continued

Cys His Ser His Phe Gln Asn Ile Phe Trp Ser Ser Leu Met Tyr Leu
 225 230 235 240

Ser Tyr Leu Val Leu Phe Cys His Phe Phe Phe Glu Ala Tyr Ile Gly
 245 250 255

Lys Met Arg Lys Thr Thr Lys Ala Glu
 260 265

<210> SEQ ID NO 65
 <211> LENGTH: 292
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 65

Met Glu Gln Leu Lys Ala Phe Asp Asn Glu Val Asn Ala Phe Leu Asp
 1 5 10 15

Asn Met Phe Gly Pro Arg Asp Ser Arg Val Arg Gly Trp Phe Leu Leu
 20 25 30

Asp Ser Tyr Leu Pro Thr Phe Ile Leu Thr Ile Thr Tyr Leu Leu Ser
 35 40 45

Ile Trp Leu Gly Asn Lys Tyr Met Lys Asn Arg Pro Ala Leu Ser Leu
 50 55 60

Arg Gly Ile Leu Thr Leu Tyr Asn Leu Ala Ile Thr Leu Leu Ser Ala
 65 70 75 80

Tyr Met Leu Val Glu Leu Ile Leu Ser Ser Trp Glu Gly Gly Tyr Asn
 85 90 95

Leu Gln Cys Gln Asn Leu Asp Ser Ala Gly Glu Gly Asp Val Arg Val
 100 105 110

Ala Lys Val Leu Trp Trp Tyr Tyr Phe Ser Lys Leu Val Glu Phe Leu
 115 120 125

Asp Thr Ile Phe Phe Val Leu Arg Lys Lys Thr Asn Gln Ile Thr Phe
 130 135 140

Leu His Val Tyr His His Ala Ser Met Phe Asn Ile Trp Trp Cys Val
 145 150 155 160

Leu Asn Trp Ile Pro Cys Gly Gln Ser Phe Phe Gly Pro Thr Leu Asn
 165 170 175

Ser Phe Ile His Ile Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Val Phe
 180 185 190

Pro Ser Met His Lys Tyr Leu Trp Trp Lys Lys Tyr Leu Thr Gln Ala
 195 200 205

Gln Leu Val Gln Phe Val Leu Thr Ile Thr His Thr Leu Ser Ala Val
 210 215 220

Val Lys Pro Cys Gly Phe Pro Phe Gly Cys Leu Ile Phe Gln Ser Ser
 225 230 235 240

Tyr Met Met Thr Leu Val Ile Leu Phe Leu Asn Phe Tyr Ile Gln Thr
 245 250 255

Tyr Arg Lys Lys Pro Val Lys Lys Glu Leu Gln Glu Lys Glu Val Lys
 260 265 270

Asn Gly Phe Pro Lys Ala His Leu Ile Val Ala Asn Gly Met Thr Asp
 275 280 285

Lys Lys Ala Gln
 290

<210> SEQ ID NO 66

-continued

<211> LENGTH: 299

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 66

Met Glu His Phe Asp Ala Ser Leu Ser Thr Tyr Phe Lys Ala Phe Leu
1 5 10 15
Gly Pro Arg Asp Thr Arg Val Lys Gly Trp Phe Leu Leu Asp Asn Tyr
20 25 30
Ile Pro Thr Phe Val Cys Ser Val Ile Tyr Leu Leu Ile Val Trp Leu
35 40 45
Gly Pro Lys Tyr Met Lys Asn Arg Gln Pro Phe Ser Cys Arg Gly Ile
50 55 60
Leu Gln Leu Tyr Asn Leu Gly Leu Thr Leu Leu Ser Leu Tyr Met Phe
65 70 75 80
Tyr Glu Leu Val Thr Gly Val Trp Glu Gly Lys Tyr Asn Phe Phe Cys
85 90 95
Gln Gly Thr Arg Ser Ala Gly Glu Ser Asp Met Lys Ile Ile Arg Val
100 105 110
Leu Trp Trp Tyr Tyr Phe Ser Lys Leu Ile Glu Phe Met Asp Thr Phe
115 120 125
Phe Phe Ile Leu Arg Lys Asn Asn His Gln Ile Thr Val Leu His Val
130 135 140
Tyr His His Ala Thr Met Leu Asn Ile Trp Trp Phe Val Met Asn Trp
145 150 155 160
Val Pro Cys Gly His Ser Tyr Phe Gly Ala Thr Leu Asn Ser Phe Ile
165 170 175
His Val Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Ser Ile Pro Ser Met
180 185 190
Arg Pro Tyr Leu Trp Trp Lys Lys Tyr Ile Thr Gln Gly Gln Leu Val
195 200 205
Gln Phe Val Leu Thr Ile Ile Gln Thr Thr Cys Gly Val Phe Trp Pro
210 215 220
Cys Ser Phe Pro Leu Gly Trp Leu Phe Phe Gln Ile Gly Tyr Met Ile
225 230 235 240
Ser Leu Ile Ala Leu Phe Thr Asn Phe Tyr Ile Gln Thr Tyr Asn Lys
245 250 255
Lys Gly Ala Ser Arg Arg Lys Asp His Leu Lys Gly His Gln Asn Gly
260 265 270
Ser Val Ala Ala Val Asn Gly His Thr Asn Ser Phe Pro Ser Leu Glu
275 280 285
Asn Ser Val Lys Pro Arg Lys Gln Arg Lys Asp
290 295

<210> SEQ ID NO 67

<211> LENGTH: 630

<212> TYPE: DNA

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 67

acaggcgact ttctgttgaa ttgtgtgcag gcacagcaag gaaggatggc aaacagcagc 60
gtgtgggatg atgtggtggg ccgcgtggag accggcgtgg accagtggat ggaaggcgcc 120
aagccgtacg cactcaccga tgggcctccg atgatggacg tgtocaccat gctggcattc 180

-continued

gagggtgggat acatggccat gctgctcttc ggcacccga tcatgaagca gatggagaag 240
ccttttgagc tcaagaccat caagctcttg cacaacttgt ttctcttcgg actttccttg 300
tacatgtgcg tggagaccat ccgccaggct atcctcggag gctacaaagt gtttggaac 360
gacatggaga agggcaacga gtctcatgct cagggcatgt ctgcacatgt gtacgtgttc 420
tacgtgtcca aggcatacga gttcttgat accgccatca tgatcctttg caagaagttc 480
aaccagggtt ccttcttgca tgtgtaccac catgccacca tttttgcat ctggtgggct 540
atcgccaagt acgtccagg aggtgatgcg tacttttcag tgatcctaaa ctctttcatc 600
cacgttatca tgtactctta ctaccctgaa 630

<210> SEQ ID NO 68
<211> LENGTH: 210
<212> TYPE: PRT
<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 68

Thr Gly Asp Phe Leu Leu Asn Leu Leu Gln Ala Gln Gln Gly Arg Met
1 5 10 15
Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr Gly
20 25 30
Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp Gly
35 40 45
Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly Tyr
50 55 60
Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu Lys
65 70 75 80
Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu Phe
85 90 95
Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile Leu
100 105 110
Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu Ser
115 120 125
His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser Lys
130 135 140
Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys Phe
145 150 155 160
Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe Ala
165 170 175
Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr Phe
180 185 190
Ser Val Ile Leu Asn Ser Phe Ile His Val Ile Met Tyr Ser Tyr Tyr
195 200 205
Pro Glu
210

<210> SEQ ID NO 69
<211> LENGTH: 819
<212> TYPE: DNA
<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 69

atggcaaaca gcagcgtgtg ggatgatgtg gtgggccgcg tggagaccgg cgtggaccag 60

-continued

tggatggatg gcgccaagcc gtacgcactc accgatgggc tcccgatgat ggacgtgtcc	120
accatgctgg cattcgaggt gggatacatg gccatgctgc tcttcggcat cccgatcatg	180
aagcagatgg agaagccttt tgagctcaag accatcaagc tcttgacaaa cttgtttctc	240
ttcggacttt ccttgtacat gtgcgtggag accatccgcc aggctatcct cggaggctac	300
aaagtgtttg gaaacgacat ggagaagggc aacgagtctc atgctcaggc catgtctcgc	360
atcgtgtacg tgttctgcgt gtccaaggca tacgagttct tggataccgc catcatgac	420
ctttgcaaga agttcaacca ggtttccttc ttgcatgtgt accacatgc caccattttt	480
gccatctggt gggctatcgc caagtacgct ccaggagggtg atgcgtactt ttcagtgtac	540
ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctcctc ccaagggttc	600
gggttcgtga agccaatcaa gccgtacatc accacccttc agatgaccca gttcatggca	660
atgcttgtgc agtccttgta cgactaccto ttcccatgcg actaccaca ggctcttgtg	720
cagcttcttg gagtgtacat gatcaccttg cttgccctct tcggcaactt ttttgtgcag	780
agctatctta aaaagccaaa aaagagcaag accaactaa	819

<210> SEQ ID NO 70

<211> LENGTH: 819

<212> TYPE: DNA

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 70

atggcaaacg gcagcgtgtg ggatgatgtg gtgggccgct tggagaccg cgtggaccag	60
tggatggatg gcgccaagcc gtacgcactc accgatgggc ccccgatgat ggacgtgtcc	120
accatgctgg cattcgaggt gggatacatg gccatgctgc tcttcggcat cccgatcatg	180
aagcagatgg agaagccttt tgagctcaag accatcaagc tcttgacaaa cttgtttctc	240
ttcggacttt ccttgtacat gtgcgtggag accatccgcc aggctatcct cggaggctac	300
aaagtgtttg gaaacgacat ggagaagggc aacgagtctc atgctcaggc catgtctcgc	360
atcgtgtacg cgttctacgt gtccaaggca tacgagttct tggataccgc catcatgac	420
ctttgcaaga agttcaacca ggtttccttc ttgcatgtgt accacatgc caccattttt	480
gccatctggt gggctatcgc caagtacgcc ccaggagggtg atgcgtactt ttcagtgtac	540
ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctcctc ccaagggttc	600
gggttcgtga agccaatcaa gccgtacatc accacccttc agatgaccca gttcatggca	660
atgcttgtgc agtccttgta cgactaccto ttcccatgcg actaccaca ggctcttgtg	720
cagcttcttg gagtgtacat gatcaccttg cttgccctct tcggcaactt ttttgtgcag	780
agctatctta aaaagccaaa aaagagcaag accaactaa	819

<210> SEQ ID NO 71

<211> LENGTH: 818

<212> TYPE: DNA

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 71

atggcaaacg gcagcgtgtg gatgatgtg tgggccgct tggagaccg cgtggaccag	60
ggatggatg gcccaagccg tacgcactca ccgatgggct cccgatgatg gacgtgtcca	120
ccatgctggc attcgaggtg ggatacatg ccgatgctgt cttcggcatc ccgatcatga	180

-continued

agcagatgga gaagcctttt gagctcaaga ccatcaagct cttgcacaac ttgtttctct	240
tcggactttc cttgtacatg tgcgtggaga ccatccgccg ggctatcctc ggaggtaca	300
aagtgtttgg aaacgacatg gagaagggca acgagtctca tgctcagggc atgtctcgca	360
tcgtgtacgt gttctacgtg tccaaggcat acgagttctt ggataccgcc atcatgatcc	420
tttgcaagaa gttcaaccag gtttccttct tgcaagtgtg ccaccatgcc accatttttg	480
ccatctggtg ggctatcgcc aagtacgctc caggaggtga tgcgtacttt tcagtgatcc	540
tcaactcttt cgtgcacacc gtcatgtacg catactactt cttctcctcc caagggttcg	600
ggttcgtgaa gccaatcaag ccgtacatca ccacccttca gatgaccagc ttcattggcaa	660
tgcttgtgca gtccttgtag gactacctct tcccatgcga ctaccacagc gctcttgtgc	720
agcttcttgg agtgtacatg atcaccttgc ttgccctctt cggcaacttt tttgtgcaga	780
gctatcttaa aaagccaaaa aagagcaaga ccaactaa	818

<210> SEQ ID NO 72

<211> LENGTH: 819

<212> TYPE: DNA

<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 72

atggcaaaca gcagcgtgtg ggatgatgtg gtgggccgcg tggagaccgg cgtggaccag	60
tggtatggatg gcgccaagcc gtacgcactc accgatgggc tcccgatgat ggacgtgtcc	120
accatgctgg cattcgaggt gggatacatg gccatgtgc tcttcggcat cccgatcatg	180
aagcagatgg agaagccttt tgagctcaag accatcaagc tcttgcaaaa cttgtttctc	240
ttcggacttt ccttgtacat gtgcgtggag accatccgcc aggotatcct cggaggctac	300
aaagtgtttg gaaacgacat ggagaagggc aacgagtctc atgctcaggg catgtctcgc	360
atcgtgtacg tgttctacgt gtccaaggca tacgagttct tggataccgc catcatgac	420
ctttgcaaga agttcaacca ggtttccttc ttgcatgtgt accaccatgc caccgttttt	480
gccatctggt gggctatcgc caagtacgct ccaggaggtg atgcgtactt ttcagtgatc	540
ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctcctc ccaagggttc	600
gggttcgtga agccaatcaa gccgtacatc accacccttc agatgacca gttcatggca	660
atgcttgtgc agtccttgta cgactacctc ttcccatgcg actaccacac ggctcttgtg	720
cagcttcttg gagtgtacat gatcaccttg cttgccctct tcggcaactt ttttgtgcag	780
agctatctta aaaagccaaa aaagagcaag accaactaa	819

<210> SEQ ID NO 73

<211> LENGTH: 819

<212> TYPE: DNA

<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 73

atggcaaaca gcagcgtgtg ggatggtgtg gtgggccgcg tggagaccgg cgtggaccag	60
tggtatggatg gcgccaagcc gtacgcactc accgatgggc tcccgatgat ggacgtgtcc	120
accatgctgg cattcgaggt gggatacatg gccatgtgc tcttcggcat cccgatcatg	180
aagcagatgg agaagccttt tgagctcaag accatcaagc tcttgcaaaa cttgtttctc	240
ttcggacttt ccttgtacat gtgcgtggag accatccgcc aggotatcct cggaggctac	300

-continued

```
aaagtgtttg gaaacgacat ggagaagggc aacgagtctc atgctcaggg catgtctcgc 360
atcgtgtacg tgttctacgt gtccaaggca tacgagttct tggataccgc catcatgac 420
ctttgcaaga agttcaacca ggtttccttc ttgcatgctg accaccatgc caccattttt 480
gccatctggt gggctatcgc caagtacgct ccaggagggtg atgcgtactt ttcagtgatc 540
ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctcctc ccaagggttc 600
gggttcgtga agccaatcaa gccgtacatc accacccttc agatgacca gttcatggca 660
atgcttgtgc agtccttgta cgactacctc ttcccatgcg actaccaca ggctcttggt 720
cagcttcttg gagtgtacat gatcaccttg cttgccctct tcggcaactt ttttgtgcag 780
agctatctta aaaagccaaa aaagagcaag accaactaa 819
```

<210> SEQ ID NO 74

<211> LENGTH: 819

<212> TYPE: DNA

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 74

```
atggcaaaca gcagcgtgtg ggatgatgtg gtgggccgcg tggagaccgg cgtggaccag 60
tggatggatg gcgccaagcc gtacgcactc accgatgggc tccgatgat ggacgtgtcc 120
accatgctgg cattcgaggt gggatacatg gccatgctgc tcttcggcat cccgatcatg 180
aggcagatgg agaagccttt tgagctcaag accatcaagc tcttgacaaa cttgtttctc 240
ttcggacttt ccttgtacat gtgcgtggtg accatccgcc aggctatcct tggaggctac 300
aaagtgtttg gaaacgacat ggagaagggc aacgagtctc atgctcaggg catgtctcgc 360
atcgtgtacg tgttctacgt gtccaaggca tacgagttct tggataccgc catcatgac 420
ctttgcaaga agttcaacca ggtttccttc ttgcatgtgt accaccatgc caccattttt 480
gccatctggt gggctatcgc caagtacgct ccaggagggtg atgcgtactt ttcagtgatc 540
ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctcctc ccaagggttc 600
gggttcgtga agccaatcaa gccgtacatc accacccttc agatgacca gttcatggca 660
atgcttgtgc agtccttgta cgactacctc ttcccatgcg actaccaca ggctcttggt 720
cagcttcttg gagtgtacat gatcaccttg cttgccctct tcggcaactt ttttgtgcag 780
agctatctta aaaagccaaa aaagagcaag accaactaa 819
```

<210> SEQ ID NO 75

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 75

```
Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr
 1          5          10          15
Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp
          20          25          30
Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly
          35          40          45
Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu
          50          55          60
Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu
```

-continued

65	70	75	80
Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile	85	90	95
Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu	100	105	110
Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Cys Val Ser	115	120	125
Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys	130	135	140
Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe	145	150	155
Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr	165	170	175
Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr	180	185	190
Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro	195	200	205
Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln	210	215	220
Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val	225	230	235
Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn	245	250	255
Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn	260	265	270

<210> SEQ ID NO 76

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 76

Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr	1	5	10	15
Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp	20	25	30	
Gly Pro Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly	35	40	45	
Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu	50	55	60	
Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu	65	70	75	80
Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile	85	90	95	
Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu	100	105	110	
Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Ala Phe Tyr Val Ser	115	120	125	
Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys	130	135	140	
Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe	145	150	155	160

-continued

Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr
165 170 175

Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr
180 185 190

Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro
195 200 205

Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln
210 215 220

Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val
225 230 235 240

Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn
245 250 255

Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn
260 265 270

<210> SEQ ID NO 77

<211> LENGTH: 265

<212> TYPE: PRT

<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 77

Met Ala Thr Ala Ala Cys Gly Met Met Trp Trp Ala Ala Trp Arg Pro
1 5 10 15

Ala Trp Thr Ser Gly Trp Met Ala Pro Ser Arg Thr His Ser Pro Met
20 25 30

Gly Ser Arg Trp Thr Cys Pro Pro Cys Trp His Ser Arg Trp Asp Thr
35 40 45

Trp Pro Cys Cys Ser Ser Ala Ser Arg Ser Ser Arg Trp Arg Ser Leu
50 55 60

Leu Ser Ser Arg Pro Ser Ser Ser Cys Thr Thr Cys Phe Ser Ser Asp
65 70 75 80

Phe Pro Cys Thr Cys Ala Trp Arg Pro Ser Ala Arg Leu Ser Ser Glu
85 90 95

Ala Thr Lys Cys Leu Glu Thr Thr Trp Arg Arg Ala Thr Ser Leu Met
100 105 110

Leu Arg Ala Cys Leu Ala Ser Cys Thr Cys Ser Thr Cys Pro Arg His
115 120 125

Thr Ser Ser Trp Ile Pro Pro Ser Ser Phe Ala Arg Ser Ser Thr Arg
130 135 140

Phe Pro Ser Cys Lys Cys Thr Thr Met Pro Pro Phe Leu Pro Ser Gly
145 150 155 160

Gly Leu Ser Pro Ser Thr Leu Gln Glu Val Met Arg Thr Phe Gln Ser
165 170 175

Ser Thr Leu Ser Cys Thr Pro Ser Cys Thr His Thr Thr Ser Ser Pro
180 185 190

Pro Lys Gly Ser Gly Ser Ser Gln Ser Ser Arg Thr Ser Pro Pro Phe
195 200 205

Arg Pro Ser Ser Trp Gln Cys Leu Cys Ser Pro Cys Thr Thr Thr Ser
210 215 220

Ser His Ala Thr Thr His Arg Leu Leu Cys Ser Phe Leu Glu Cys Thr
225 230 235 240

Ser Pro Cys Leu Pro Ser Ser Ala Thr Phe Leu Cys Arg Ala Ile Leu
245 250 255

 -continued

Lys Ser Gln Lys Arg Ala Arg Pro Thr
 260 265

<210> SEQ ID NO 78

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 78

Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr
 1 5 10 15

Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp
 20 25 30

Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly
 35 40 45

Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu
 50 55 60

Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu
 65 70 75 80

Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile
 85 90 95

Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu
 100 105 110

Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser
 115 120 125

Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys
 130 135 140

Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Val Phe
 145 150 155 160

Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr
 165 170 175

Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr
 180 185 190

Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro
 195 200 205

Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln
 210 215 220

Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val
 225 230 235 240

Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn
 245 250 255

Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn
 260 265 270

<210> SEQ ID NO 79

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: *Thraustochytrium aureum*

<400> SEQUENCE: 79

Met Ala Asn Ser Ser Val Trp Asp Gly Val Val Gly Arg Val Glu Thr
 1 5 10 15

Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp
 20 25 30

-continued

Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly
 35 40 45
 Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu
 50 55 60
 Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu
 65 70 75 80
 Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile
 85 90 95
 Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu
 100 105 110
 Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser
 115 120 125
 Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys
 130 135 140
 Phe Asn Gln Val Ser Phe Leu His Ala Tyr His His Ala Thr Ile Phe
 145 150 155 160
 Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr
 165 170 175
 Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr
 180 185 190
 Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro
 195 200 205
 Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln
 210 215 220
 Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val
 225 230 235 240
 Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn
 245 250 255
 Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn
 260 265 270

<210> SEQ ID NO 80

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 80

Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr
 1 5 10 15
 Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp
 20 25 30
 Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly
 35 40 45
 Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu
 50 55 60
 Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu
 65 70 75 80
 Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile
 85 90 95
 Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu
 100 105 110
 Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser

-continued

115	120	125
Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys 130 135 140		
Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe 145 150 155 160		
Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr 165 170 175		
Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr 180 185 190		
Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro 195 200 205		
Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln 210 215 220		
Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val 225 230 235 240		
Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn 245 250 255		
Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn 260 265 270		

<210> SEQ ID NO 81

<211> LENGTH: 272

<212> TYPE: PRT

<213> ORGANISM: Thraustochytrium aureum

<400> SEQUENCE: 81

Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr 1 5 10 15
Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp 20 25 30
Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly 35 40 45
Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Arg Gln Met Glu 50 55 60
Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu 65 70 75 80
Phe Gly Leu Ser Leu Tyr Met Cys Val Val Thr Ile Arg Gln Ala Ile 85 90 95
Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu 100 105 110
Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser 115 120 125
Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys 130 135 140
Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe 145 150 155 160
Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr 165 170 175
Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr 180 185 190
Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro 195 200 205

-continued

Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln
 210 215 220

Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val
 225 230 235 240

Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn
 245 250 255

Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn
 260 265 270

<210> SEQ ID NO 82
 <211> LENGTH: 292
 <212> TYPE: PRT
 <213> ORGANISM: Thraustochytrium aureum
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (273)...(273)
 <223> OTHER INFORMATION: Xaa = Unknown or other at position 273

<400> SEQUENCE: 82

Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr
 1 5 10 15

Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp
 20 25 30

Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly
 35 40 45

Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu
 50 55 60

Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu
 65 70 75 80

Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile
 85 90 95

Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu
 100 105 110

Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser
 115 120 125

Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys
 130 135 140

Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe
 145 150 155 160

Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr
 165 170 175

Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr
 180 185 190

Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro
 195 200 205

Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln
 210 215 220

Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val
 225 230 235 240

Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn
 245 250 255

Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn
 260 265 270

Xaa Asn Cys Leu His Asp Met Pro Leu Ala Gly Val Arg Ile Asp Ser

-continued

275	280	285
Glu Ser Glu Leu		
290		
 <210> SEQ ID NO 83		
<211> LENGTH: 284		
<212> TYPE: PRT		
<213> ORGANISM: Homo sapiens		
 <400> SEQUENCE: 83		
Met Glu His Phe Asp Ala Ser Leu Ser Thr Tyr Phe Lys Ala Leu Leu		
1 5 10 15		
Gly Pro Arg Asp Thr Arg Val Lys Gly Trp Phe Leu Leu Asp Asn Tyr		
20 25 30		
Ile Pro Thr Phe Ile Cys Ser Val Ile Tyr Leu Leu Ile Val Trp Leu		
35 40 45		
Gly Pro Lys Tyr Met Arg Asn Lys Gln Pro Phe Ser Cys Arg Gly Ile		
50 55 60		
Leu Val Val Tyr Asn Leu Gly Leu Thr Leu Leu Ser Leu Tyr Met Phe		
65 70 75 80		
Cys Glu Leu Val Thr Gly Val Trp Glu Gly Lys Tyr Asn Phe Phe Cys		
85 90 95		
Gln Gly Thr Arg Thr Ala Gly Glu Ser Asp Met Lys Ile Ile Arg Val		
100 105 110		
Leu Trp Trp Tyr Tyr Phe Ser Lys Leu Ile Glu Phe Met Asp Thr Phe		
115 120 125		
Phe Phe Ile Leu Arg Lys Asn Asn His Gln Ile Thr Val Leu His Val		
130 135 140		
Tyr His His Ala Ser Met Leu Asn Ile Trp Trp Phe Val Met Asn Trp		
145 150 155 160		
Val Pro Cys Gly His Ser Tyr Phe Gly Ala Thr Leu Asn Ser Phe Ile		
165 170 175		
His Val Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Ser Val Pro Ser Met		
180 185 190		
Arg Pro Tyr Leu Trp Trp Lys Lys Tyr Ile Thr Gln Gly Gln Leu Leu		
195 200 205		
Gln Phe Val Leu Thr Ile Ile Gln Thr Ser Cys Gly Val Ile Trp Pro		
210 215 220		
Cys Thr Phe Pro Leu Gly Trp Leu Tyr Phe Gln Ile Gly Tyr Met Ile		
225 230 235 240		
Ser Leu Ile Ala Leu Phe Thr Asn Phe Tyr Ile Gln Thr Tyr Asn Lys		
245 250 255		
Lys Gly Ala Ser Arg Arg Lys Asp His Leu Lys Asp His Gln Asn Gly		
260 265 270		
Ser Met Ala Ala Val Asn Gly His Thr Asn Ser Phe		
275 280		

<210> SEQ ID NO 84
 <211> LENGTH: 288
 <212> TYPE: PRT
 <213> ORGANISM: Thraustochytrium aureum
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: (273)...(273)
 <223> OTHER INFORMATION: Xaa = Unknown or other at position 273

-continued

<400> SEQUENCE: 84

Met Ala Asn Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr
1 5 10 15
Gly Val Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp
20 25 30
Gly Leu Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly
35 40 45
Tyr Met Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu
50 55 60
Lys Pro Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu
65 70 75 80
Phe Gly Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile
85 90 95
Leu Gly Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu
100 105 110
Ser His Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser
115 120 125
Lys Ala Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys
130 135 140
Phe Asn Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe
145 150 155 160
Ala Ile Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr
165 170 175
Phe Ser Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr
180 185 190
Tyr Phe Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro
195 200 205
Tyr Ile Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln
210 215 220
Ser Leu Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val
225 230 235 240
Gln Leu Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn
245 250 255
Phe Phe Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn
260 265 270
Xaa Asn Cys Leu His Asp Met Pro Leu Ala Gly Val Arg Ile Asp Ser
275 280 285

<210> SEQ ID NO 85

<211> LENGTH: 283

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 85

Met Glu Gln Leu Lys Ala Phe Asp Asn Glu Val Asn Ala Phe Leu Asp
1 5 10 15
Asn Met Phe Gly Pro Arg Asp Ser Arg Val Arg Gly Trp Phe Leu Leu
20 25 30
Asp Ser Tyr Leu Pro Thr Phe Ile Leu Thr Ile Thr Tyr Leu Leu Ser
35 40 45
Ile Trp Leu Gly Asn Lys Tyr Met Lys Asn Arg Pro Ala Leu Ser Leu
50 55 60

-continued

Arg Gly Ile Leu Thr Leu Tyr Asn Leu Ala Ile Thr Leu Leu Ser Ala
 65 70 75 80
 Tyr Met Leu Val Glu Leu Ile Leu Ser Ser Trp Glu Gly Gly Tyr Asn
 85 90 95
 Leu Gln Cys Gln Asn Leu Asp Ser Ala Gly Glu Gly Asp Val Arg Val
 100 105 110
 Ala Lys Val Leu Trp Trp Tyr Tyr Phe Ser Lys Leu Val Glu Phe Leu
 115 120 125
 Asp Thr Ile Phe Phe Val Leu Arg Lys Lys Thr Asn Gln Ile Thr Phe
 130 135 140
 Leu His Val Tyr His His Ala Ser Met Phe Asn Ile Trp Trp Cys Val
 145 150 155 160
 Leu Asn Trp Ile Pro Cys Gly Gln Ser Phe Phe Gly Pro Thr Leu Asn
 165 170 175
 Ser Phe Ile His Ile Leu Met Tyr Ser Tyr Tyr Gly Leu Ser Val Phe
 180 185 190
 Pro Ser Met His Lys Tyr Leu Trp Trp Lys Lys Tyr Leu Thr Gln Ala
 195 200 205
 Gln Leu Val Gln Phe Val Leu Thr Ile Thr His Thr Leu Ser Ala Val
 210 215 220
 Val Lys Pro Cys Gly Phe Pro Phe Gly Cys Leu Ile Phe Gln Ser Ser
 225 230 235 240
 Tyr Met Met Thr Leu Val Ile Leu Phe Leu Asn Phe Tyr Ile Gln Thr
 245 250 255
 Tyr Arg Lys Lys Pro Val Lys Lys Glu Leu Gln Glu Lys Glu Val Lys
 260 265 270
 Asn Gly Phe Pro Lys Ala His Leu Ile Val Ala
 275 280

<210> SEQ ID NO 86

<211> LENGTH: 295

<212> TYPE: PRT

<213> ORGANISM: Thraustochytrium aureum

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: (270)...(270)

<223> OTHER INFORMATION: Xaa = Unknown or other at position 270

<400> SEQUENCE: 86

Ser Ser Val Trp Asp Asp Val Val Gly Arg Val Glu Thr Gly Val Asp
 1 5 10 15
 Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp Gly Leu Pro
 20 25 30
 Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly Tyr Met Ala
 35 40 45
 Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu Lys Pro Phe
 50 55 60
 Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu Phe Gly Leu
 65 70 75 80
 Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile Leu Gly Gly
 85 90 95
 Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu Ser His Ala
 100 105 110

```
<210> SEQ ID NO 87
<211> LENGTH: 280
<212> TYPE: PRT
<213> ORGANISM: Mortierella alpina

<400> SEQUENCE: 87
```

Leu	Val	Ala	Gln	Ala	Glu	Lys	Tyr	Ile	Pro	Thr	Ile	Val	His	His	Thr
1				5					10					15	
Arg	Gly	Phe	Leu	Val	Ala	Val	Glu	Ser	Pro	Leu	Ala	Arg	Glu	Leu	Pro
			20					25					30		
Leu	Met	Asn	Pro	Phe	His	Val	Leu	Leu	Ile	Val	Leu	Ala	Tyr	Leu	Val
		35					40					45			
Thr	Val	Phe	Val	Gly	Met	Gln	Ile	Met	Lys	Asn	Phe	Glu	Arg	Phe	Glu
	50					55					60				
Val	Lys	Thr	Phe	Ser	Leu	Leu	His	Asn	Phe	Cys	Leu	Val	Ser	Ile	Ser
65					70				75						80
Ala	Tyr	Met	Cys	Gly	Gly	Ile	Leu	Tyr	Glu	Ala	Tyr	Gln	Ala	Asn	Tyr
			85						90					95	
Gly	Leu	Phe	Glu	Asn	Ala	Ala	Asp	His	Thr	Phe	Lys	Gly	Leu	Pro	Met
			100					105					110		
Ala	Lys	Met	Ile	Trp	Leu	Phe	Tyr	Phe	Ser	Lys	Ile	Met	Glu	Phe	Val
		115					120					125			
Asp	Thr	Met	Ile	Met	Val	Leu	Lys	Lys	Asn	Asn	Arg	Gln	Ile	Ser	Phe
	130					135					140				
Leu	His	Val	Tyr	His	His	Ser	Ser	Ile	Phe	Thr	Ile	Trp	Trp	Leu	Val
145					150					155				160	
Thr	Phe	Val	Ala	Pro	Asn	Gly	Glu	Ala	Tyr	Phe	Ser	Ala	Ala	Leu	Asn
			165						170					175	

-continued

Ser Phe Ile His Val Ile Met Tyr Gly Tyr Tyr Phe Leu Ser Ala Leu
180 185 190

Gly Phe Lys Gln Val Ser Phe Ile Lys Phe Tyr Ile Thr Arg Ser Gln
195 200 205

Met Thr Gln Phe Cys Met Met Ser Val Gln Ser Ser Trp Asp Met Tyr
210 215 220

Ala Met Lys Val Leu Gly Arg Pro Gly Tyr Pro Phe Phe Ile Thr Ala
225 230 235 240

Leu Leu Trp Phe Tyr Met Trp Thr Met Leu Gly Leu Phe Tyr Asn Phe
245 250 255

Tyr Arg Lys Asn Ala Lys Leu Ala Lys Gln Ala Lys Ala Asp Ala Ala
260 265 270

Lys Glu Lys Ala Arg Lys Leu Gln
275 280

<210> SEQ ID NO 88
<211> LENGTH: 283
<212> TYPE: PRT
<213> ORGANISM: Thraustochytrium aureum
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (255)...(255)
<223> OTHER INFORMATION: Xaa = Unknown or other at position 255

<400> SEQUENCE: 88

Asp Gln Trp Met Asp Gly Ala Lys Pro Tyr Ala Leu Thr Asp Gly Leu
1 5 10 15

Pro Met Met Asp Val Ser Thr Met Leu Ala Phe Glu Val Gly Tyr Met
20 25 30

Ala Met Leu Leu Phe Gly Ile Pro Ile Met Lys Gln Met Glu Lys Pro
35 40 45

Phe Glu Leu Lys Thr Ile Lys Leu Leu His Asn Leu Phe Leu Phe Gly
50 55 60

Leu Ser Leu Tyr Met Cys Val Glu Thr Ile Arg Gln Ala Ile Leu Gly
65 70 75 80

Gly Tyr Lys Val Phe Gly Asn Asp Met Glu Lys Gly Asn Glu Ser His
85 90 95

Ala Gln Gly Met Ser Arg Ile Val Tyr Val Phe Tyr Val Ser Lys Ala
100 105 110

Tyr Glu Phe Leu Asp Thr Ala Ile Met Ile Leu Cys Lys Lys Phe Asn
115 120 125

Gln Val Ser Phe Leu His Val Tyr His His Ala Thr Ile Phe Ala Ile
130 135 140

Trp Trp Ala Ile Ala Lys Tyr Ala Pro Gly Gly Asp Ala Tyr Phe Ser
145 150 155 160

Val Ile Leu Asn Ser Phe Val His Thr Val Met Tyr Ala Tyr Tyr Phe
165 170 175

Phe Ser Ser Gln Gly Phe Gly Phe Val Lys Pro Ile Lys Pro Tyr Ile
180 185 190

Thr Thr Leu Gln Met Thr Gln Phe Met Ala Met Leu Val Gln Ser Leu
195 200 205

Tyr Asp Tyr Leu Phe Pro Cys Asp Tyr Pro Gln Ala Leu Val Gln Leu
210 215 220

-continued

Leu Gly Val Tyr Met Ile Thr Leu Leu Ala Leu Phe Gly Asn Phe Phe
 225 230 235 240

Val Gln Ser Tyr Leu Lys Lys Pro Lys Lys Ser Lys Thr Asn Xaa Asn
 245 250 255

Cys Leu His Asp Met Pro Leu Ala Gly Val Arg Ile Asp Ser Glu Ser
 260 265 270

Glu Leu Arg Arg His Ala Asn Ser Ile Phe Phe
 275 280

<210> SEQ ID NO 89
 <211> LENGTH: 265
 <212> TYPE: PRT
 <213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 89

Ala Thr His Gly Pro Lys Asn Phe Pro Asp Ala Glu Gly Arg Lys Phe
 1 5 10 15

Phe Ala Asp His Phe Asp Val Thr Ile Gln Ala Ser Ile Leu Tyr Met
 20 25 30

Val Val Val Phe Gly Thr Lys Trp Phe Met Arg Asn Arg Gln Pro Phe
 35 40 45

Gln Leu Thr Ile Pro Leu Asn Ile Trp Asn Phe Ile Leu Ala Ala Phe
 50 55 60

Ser Ile Ala Gly Ala Val Lys Met Thr Pro Glu Phe Phe Gly Thr Ile
 65 70 75 80

Ala Asn Lys Gly Ile Val Ala Ser Tyr Cys Lys Val Phe Asp Phe Thr
 85 90 95

Lys Gly Glu Asn Gly Tyr Trp Val Trp Leu Phe Met Ala Ser Lys Leu
 100 105 110

Phe Glu Leu Val Asp Thr Ile Phe Leu Val Leu Arg Lys Arg Pro Leu
 115 120 125

Met Phe Leu His Trp Tyr His His Ile Leu Thr Met Ile Tyr Ala Trp
 130 135 140

Tyr Ser His Pro Leu Thr Pro Gly Phe Asn Arg Tyr Gly Ile Tyr Leu
 145 150 155 160

Asn Phe Val Val His Ala Phe Met Tyr Ser Tyr Tyr Phe Leu Arg Ser
 165 170 175

Met Lys Ile Arg Val Pro Gly Phe Ile Ala Gln Ala Ile Thr Ser Leu
 180 185 190

Gln Ile Val Gln Phe Ile Ile Ser Cys Ala Val Leu Ala His Leu Gly
 195 200 205

Tyr Leu Met His Phe Thr Asn Ala Asn Cys Asp Phe Glu Pro Ser Val
 210 215 220

Phe Lys Leu Ala Val Phe Met Asp Thr Thr Tyr Leu Ala Leu Phe Val
 225 230 235 240

Asn Phe Phe Leu Gln Ser Tyr Val Leu Arg Gly Gly Lys Asp Lys Tyr
 245 250 255

Lys Ala Val Pro Lys Lys Lys Asn Asn
 260 265

<210> SEQ ID NO 90
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Internal Primer RO339

<400> SEQUENCE: 90

ttggagagga ggaagcgacc accgaagatg atg 33

<210> SEQ ID NO 91
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward Primer RO317

<400> SEQUENCE: 91

cacacaggaa acagctatga ccatgattac g 31

<210> SEQ ID NO 92
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO350

<400> SEQUENCE: 92

catctcatgg atccgccatg gccgcgcgcaa tcttg 35

<210> SEQ ID NO 93
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer RO352

<400> SEQUENCE: 93

acgcgtacgt aaagcttg 18

<210> SEQ ID NO 94
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO514

<400> SEQUENCE: 94

ggctatggat ccatgaattc actcgttact caatatg 37

<210> SEQ ID NO 95
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO515

<400> SEQUENCE: 95

cctgccaaagc ttttaccttt ttcttctgtg ttgag 35

<210> SEQ ID NO 96
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward Primer RO541

<400> SEQUENCE: 96

-continued

gactactagc agctgtaata c 21

<210> SEQ ID NO 97
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer R0540

<400> SEQUENCE: 97

gtgaatgtaa gcgtgacata a 21

<210> SEQ ID NO 98
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Internal Forward Primer R0728

<400> SEQUENCE: 98

gagactttga gcggttcg 18

<210> SEQ ID NO 99
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Internal Forward Primer R0730

<400> SEQUENCE: 99

ttctctgtgc gttgaactcg 20

<210> SEQ ID NO 100
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer R0729

<400> SEQUENCE: 100

aaagctcttg acctcgaac 19

<210> SEQ ID NO 101
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer R0731

<400> SEQUENCE: 101

aacttgatga acgacacgtg 20

<210> SEQ ID NO 102
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer R0719

<400> SEQUENCE: 102

ggttctccca tggaacattt tgatgcac 29

-continued

<210> SEQ ID NO 103
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO720

<400> SEQUENCE: 103

ggtttcaaag ctttgacttc aatccttccg 30

<210> SEQ ID NO 104
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO738

<400> SEQUENCE: 104

aatcaggaat tcatggctca gcatccgctc gttcaac 37

<210> SEQ ID NO 105
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO739

<400> SEQUENCE: 105

ccgcttgtcg acttagttgt tcttcttctt tggcac 36

<210> SEQ ID NO 106
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RP735

<400> SEQUENCE: 106

cctcctgaat tccaacacta ttcagctttc 30

<210> SEQ ID NO 107
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO73

<400> SEQUENCE: 107

taatacgact cactataggg 20

<210> SEQ ID NO 108
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO819

<400> SEQUENCE: 108

atgatgccat ggagcagctg aaggcctttg 30

<210> SEQ ID NO 109
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Primer R0820

<400> SEQUENCE: 109

cagtctctgc tttaaaacaa gctcgtc 27

<210> SEQ ID NO 110
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer R0833

<400> SEQUENCE: 110

gggttttacca tggaacattt cgatgcgtca c 31

<210> SEQ ID NO 111
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer R0832

<400> SEQUENCE: 111

cgacctgcag ctcgagcaca 20

<210> SEQ ID NO 112
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer R0895
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (7)...(7)
<223> OTHER INFORMATION: w = a or t/u at position 7
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (19)...(19)
<223> OTHER INFORMATION: w = a or t/u at position 19
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (28)...(28)
<223> OTHER INFORMATION: r = g or a at position 28
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (31)...(31)
<223> OTHER INFORMATION: w = a or t/u at position 31
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (37)...(37)
<223> OTHER INFORMATION: w = a or t/u at position 37

<400> SEQUENCE: 112

gtagtawgag tacatgatwa cgtggatraa wgagttwag 39

<210> SEQ ID NO 113
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer R0898

<400> SEQUENCE: 113

cccagtcacg acgttgtaaa acgacggcca g 31

-continued

<210> SEQ ID NO 114
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO1160

<400> SEQUENCE: 114

aaggaacat ggcaaacagc agcgtgtggg atg 33

<210> SEQ ID NO 115
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer RO899

<400> SEQUENCE: 115

agcggataac aatttcacac aggaacagc 30

<210> SEQ ID NO 116
<211> LENGTH: 28
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 116

Asp Thr Ile Phe Ile Ile Leu Arg Lys Gln Lys Leu Ile Phe Leu His
1 5 10 15

Trp Tyr His His Ile Thr Val Leu Leu Tyr Ser Trp
20 25

<210> SEQ ID NO 117
<211> LENGTH: 834
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Pavlova sp. (specific species unknown)

<400> SEQUENCE: 117

atgatgttgg ccgcaggcta tcttctagtg ctctcggccg ctgccagag cttccagcag 60

gacattgaca accccaacgg ggccactctg acctcgtgga ctggcctgcc catttgtgatg 120

tctgtggtct atctcagcgg tgtgtttggg ctcaaaagt acttcgagaa ccggaagccc 180

atgacggggc tgaaggacta catgttcact tacaatctct accaggtgat catcaacgtg 240

tggtgcgtgg tggcctttct cctggagggt cggcgtgcgg gcatgtcact catcggcaat 300

aaggtggacc ttgggcccaa ctccctcagg ctcggtctcg tcacgtgggt gactacaac 360

aacaagtacg tggagctcct cgacacccta tggatggtgc tgcgcaagaa gacgcagcag 420

gtctccttcc tccacgtcta tcatcacgtg cttctgatgt gggcctggtt cgttgtctgtc 480

aagctcgcca atggtgtgta cgcataatct gccggtctca tgaactcgat catccacgtg 540

atgatgtatt cctactacac catggcgctc ctgggctggt catgcccctg gaagcgctac 600

ctcacgcagg cacagctcgt gcagttttgc atctgcctcg cccactccac atgggcggca 660

gtaacgggtg cctaccctgt gcgaatttgc ttggtggagg tgtgggtgat ggtgtocatg 720

ctggtgtctt tcacacgctt ctaccgccag gcctatgcca aggaggcgaa ggccaaggag 780

gcgaaaaagc tcgcacagga ggcacacag gccaaaggcg tcaaggcgga gtaa 834

-continued

<210> SEQ ID NO 118
<211> LENGTH: 277
<212> TYPE: PRT
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Pavlova sp. (specific species unknown)

<400> SEQUENCE: 118

Met Met Leu Ala Ala Gly Tyr Leu Leu Val Leu Ser Ala Ala Arg Gln
1 5 10 15

Ser Phe Gln Gln Asp Ile Asp Asn Pro Asn Gly Ala Tyr Ser Thr Ser
20 25 30

Trp Thr Gly Leu Pro Ile Val Met Ser Val Val Tyr Leu Ser Gly Val
35 40 45

Phe Gly Leu Thr Lys Tyr Phe Glu Asn Arg Lys Pro Met Thr Gly Leu
50 55 60

Lys Asp Tyr Met Phe Thr Tyr Asn Leu Tyr Gln Val Ile Ile Asn Val
65 70 75 80

Trp Cys Val Val Ala Phe Leu Leu Glu Val Arg Arg Ala Gly Met Ser
85 90 95

Leu Ile Gly Asn Lys Val Asp Leu Gly Pro Asn Ser Phe Arg Leu Gly
100 105 110

Phe Val Thr Trp Val His Tyr Asn Asn Lys Tyr Val Glu Leu Leu Asp
115 120 125

Thr Leu Trp Met Val Leu Arg Lys Lys Thr Gln Gln Val Ser Phe Leu
130 135 140

His Val Tyr His His Val Leu Leu Met Trp Ala Trp Phe Val Val Val
145 150 155 160

Lys Leu Gly Asn Gly Gly Asp Ala Tyr Phe Ala Gly Leu Met Asn Ser
165 170 175

Ile Ile His Val Met Met Tyr Ser Tyr Tyr Thr Met Ala Leu Leu Gly
180 185 190

Trp Ser Cys Pro Trp Lys Arg Tyr Leu Thr Gln Ala Gln Leu Val Gln
195 200 205

Phe Cys Ile Cys Leu Ala His Ser Thr Trp Ala Ala Val Thr Gly Ala
210 215 220

Tyr Pro Trp Arg Ile Cys Leu Val Glu Val Trp Val Met Val Ser Met
225 230 235 240

Leu Val Leu Phe Thr Arg Phe Tyr Arg Gln Ala Tyr Ala Lys Glu Ala
245 250 255

Lys Ala Lys Glu Ala Lys Lys Leu Ala Gln Glu Ala Ser Gln Ala Lys
260 265 270

Ala Val Lys Ala Glu
275

<210> SEQ ID NO 119
<211> LENGTH: 834
<212> TYPE: DNA
<213> ORGANISM: Unknown
<220> FEATURE:
<223> OTHER INFORMATION: Pavlova sp. (specific species unknown)

<400> SEQUENCE: 119

atgatgttgg ccgcaggcta tcttctagtg ctctcggccg ctgccagag cttccagcag 60
gacattgaca accccaacgg ggcctactcg acctcgtgga ctggcctgcc cattgtgatg 120

-continued

Ser Phe Gln Gln Asp Ile Asp Asn Pro Asn Gly Ala Tyr Ser Thr Ser
 20 25 30
 Trp Thr Gly Leu Pro Ile Val Met Ser Val Val Tyr Leu Ser Gly Val
 35 40 45
 Phe Gly Leu Thr Lys Tyr Phe Glu Asn Arg Lys Pro Met Thr Gly Leu
 50 55 60
 Lys Asp Tyr Met Phe Thr Tyr Asn Leu Tyr Gln Val Ile Ile Asn Val
 65 70 75 80
 Trp Cys Val Val Ala Phe Leu Leu Glu Val Arg Arg Ala Gly Met Ser
 85 90 95
 Leu Ile Gly Asn Lys Val Asp Leu Gly Pro Asn Ser Phe Arg Leu Gly
 100 105 110
 Phe Val Thr Trp Val His Tyr Asn Asn Lys Tyr Val Glu Leu Leu Asp
 115 120 125
 Thr Leu Trp Met Val Leu Arg Lys Lys Thr Gln Gln Val Ser Phe Leu
 130 135 140
 His Val Tyr His His Val Leu Leu Met Trp Ala Trp Phe Val Val Val
 145 150 155 160
 Lys Leu Gly Asn Gly Gly Asp Ala Tyr Phe Gly Gly Leu Met Asn Ser
 165 170 175
 Ile Ile His Val Met Met Tyr Ser Tyr Tyr Thr Met Ala Leu Leu Gly
 180 185 190
 Trp Ser Cys Pro Trp Lys Arg Tyr Leu Thr Gln Ala Gln Leu Val Gln
 195 200 205
 Phe Cys Ile Cys Leu Ala His Ser Thr Trp Ala Ala Val Thr Gly Ala
 210 215 220
 Tyr Pro Trp Arg Ile Cys Leu Val Glu Val Trp Val Met Val Ser Met
 225 230 235 240
 Leu Val Leu Phe Thr Arg Phe Tyr Arg Gln Ala Tyr Ala Lys Glu Ala
 245 250 255
 Lys Ala Lys Glu Ala Lys Lys Leu Ala Gln Glu Ala Ser Gln Ala Lys
 260 265 270
 Ala Val Lys Ala Glu
 275

<210> SEQ ID NO 122

<211> LENGTH: 277

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: Pavlova sp. (specific species unknown)

<400> SEQUENCE: 122

Met Met Leu Ala Ala Gly Tyr Leu Leu Val Leu Ser Ala Ala Arg Gln
 1 5 10 15
 Ser Phe Gln Gln Asp Ile Asp Asn Pro Asn Gly Ala Tyr Ser Thr Ser
 20 25 30
 Trp Thr Gly Leu Pro Ile Val Met Ser Val Val Tyr Leu Ser Gly Val
 35 40 45
 Phe Gly Leu Thr Lys Tyr Phe Glu Asp Arg Lys Pro Met Thr Gly Leu
 50 55 60
 Lys Asp Tyr Met Phe Thr Tyr Asn Leu Tyr Gln Val Ile Ile Asn Val
 65 70 75 80

-continued

Trp	Cys	Val	Val	Ala	Phe	Leu	Leu	Glu	Val	Arg	Arg	Ala	Gly	Met	Ser
				85					90					95	
Leu	Ile	Gly	Asn	Lys	Val	Asp	Leu	Gly	Pro	Asn	Ser	Phe	Arg	Leu	Gly
			100					105					110		
Phe	Val	Thr	Trp	Val	His	Tyr	Asn	Asn	Lys	Tyr	Val	Glu	Leu	Leu	Asp
		115					120					125			
Thr	Leu	Trp	Met	Val	Leu	Arg	Lys	Lys	Thr	Gln	Gln	Val	Ser	Phe	Leu
	130					135				140					
His	Val	Tyr	His	His	Val	Leu	Leu	Met	Trp	Ala	Trp	Phe	Val	Val	Val
	145				150					155					160
Lys	Leu	Gly	Asn	Gly	Gly	Asp	Ala	Tyr	Phe	Gly	Gly	Leu	Met	Asn	Ser
			165						170					175	
Ile	Ile	His	Val	Met	Met	Tyr	Ser	Tyr	Tyr	Thr	Met	Ala	Leu	Leu	Gly
		180						185					190		
Trp	Ser	Cys	Pro	Trp	Lys	Arg	Tyr	Leu	Thr	Gln	Ala	Gln	Leu	Val	Gln
		195					200					205			
Phe	Cys	Ile	Cys	Leu	Ala	His	Ser	Thr	Trp	Ala	Ala	Val	Thr	Gly	Ala
	210					215				220					
Tyr	Pro	Trp	Arg	Ile	Cys	Leu	Val	Glu	Val	Trp	Val	Met	Val	Ser	Met
	225				230					235					240
Leu	Val	Leu	Phe	Thr	Arg	Phe	Tyr	Cys	Gln	Ala	Tyr	Ala	Lys	Glu	Ala
			245						250					255	
Lys	Ala	Lys	Glu	Ala	Lys	Lys	Leu	Ala	Gln	Glu	Ala	Ser	Gln	Ala	Lys
			260				265						270		
Ala	Val	Lys	Ala	Glu											
		275													

1. An isolated nucleotide sequence or fragment thereof comprising or complementary to a nucleotide sequence encoding a polypeptide having elongase activity, wherein the amino acid sequence of said polypeptide has at least 50% identity to an amino acid sequence comprising SEQ ID NO:121.

2. An isolated nucleotide sequence or fragment thereof comprising or complementary to a nucleotide sequence having at least 50% nucleotide sequence identity to a nucleotide sequence comprising SEQ ID NO:119.

3. The isolated nucleotide sequence of claims 1 or 2 wherein said sequence encodes a functionally active elongase which utilizes a polyunsaturated fatty acid as a substrate.

4. The isolated nucleotide sequence of claim 1 wherein said sequence is derived from an algae.

5. The isolated nucleotide sequence of claim 4 wherein said algae is *Pavlova* sp.

6. A purified protein encoded by said nucleotide sequence of claims 1 or 2.

7. A purified polypeptide which elongates polyunsaturated fatty acids and has at least about 30% amino acid identity to the amino acid sequence of said purified protein of claim 6.

8. A method of producing an elongase enzyme comprising the steps of:

- a) isolating a nucleotide sequence comprising SEQ ID NO:119;
- b) constructing a vector comprising: i) said isolated nucleotide sequence operably linked to ii) a promoter;

c) introducing said vector into a host cell under time and conditions sufficient for expression of said elongase enzyme.

9. The method of claim 8 wherein said host cell is selected from the group consisting of a eukaryotic cell or a prokaryotic cell.

10. The method of claim 9 wherein said prokaryotic cell is selected from the group consisting of *E. coli*, *Cyanobacteria*, and *B. subtilis*.

11. The method of claim 9 wherein said eukaryotic cell is selected from the group consisting of a mammalian cell, an insect cell, a plant cell and a fungal cell.

12. The method of claim 11 wherein said fungal cell is selected from the group consisting of *Saccharomyces* spp., *Candida* spp., *Lipomyces starkey*, *Yarrowia* spp., *Kluyveromyces* spp., *Hansenula* spp., *Aspergillus* spp., *Penicillium* spp., *Neurospora* spp., *Trichoderma* spp. and *Pichia* spp.

13. The method of claim 12 wherein said fungal cell is a yeast cell selected from the group consisting of *Saccharomyces* spp., *Candida* spp., *Hansenula* spp. and *Pichia* spp.

14. The method of claim 13 wherein said yeast cell is *Saccharomyces cerevisiae*.

15. A vector comprising: a) a nucleotide sequence comprising SEQ ID NO:119 operably linked to b) a promoter.

16. A host cell comprising said vector of claim 15.

17. The host cell of claim 16 wherein said host cell is selected from the group consisting of a eukaryotic cell or a prokaryotic cell.

18. The host cell of claim 17 wherein said prokaryotic cell is selected from the group consisting of *E. coli*, Cyanobacteria, and *B. subtilis*.

19. The host cell of claim 17 wherein said eukaryotic cell is selected from the group consisting of a mammalian cell, an insect cell, a plant cell and a fungal cell.

20. The host cell of claim 19 wherein said fungal cell is selected from the group consisting of *Saccharomyces* spp., *Candida* spp., *Lipomyces starkey*, *Yarrowia* spp., *Kluyveromyces* spp., *Hansenula* spp., *Aspergillus* spp., *Penicillium* spp., *Neurospora* spp., *Trichoderma* spp. and *Pichia* spp.

21. The host cell of claim 20 wherein said fungal cell is a yeast cell selected from the group consisting of *Saccharomyces* spp., *Candida* spp., *Hansenula* spp. and *Pichia* spp.

22. The host cell of claim 21 wherein said yeast cell is *Saccharomyces cerevisiae*.

23. A plant cell, plant or plant tissue comprising said vector of claim 15, wherein expression of said nucleotide sequence of said vector results in production of a polyunsaturated fatty acid by said plant cell, plant or plant tissue.

24. The plant cell, plant or plant tissue of claim 23 wherein said polyunsaturated fatty acid is selected from the group consisting of ADA and ω 3-docosapentaenoic acid.

25. One or more plant oils or acids expressed by said plant cell, plant or plant tissue of claim 23.

26. A transgenic plant comprising said vector of claim 15, wherein expression of said nucleotide sequence of said vector results in production of a polyunsaturated fatty acid in seeds of said transgenic plant.

27. A transgenic, non-human mammal whose genome comprises a DNA sequence encoding an elongase, operably linked to a promoter, wherein said DNA sequence comprises SEQ ID NO:119.

28. A fluid produced by said transgenic, non-human mammal of claim 27 wherein said fluid comprises a detectable level of at least one elongase or products thereof.

29. A method for producing a polyunsaturated fatty acid comprising the steps of:

- a) isolating a nucleotide sequence comprising SEQ ID NO:119;
- b) constructing a vector comprising said isolated nucleotide sequence;
- c) introducing said vector into a host cell under time and conditions sufficient for expression of an elongase enzyme encoded by said isolated nucleotide sequence; and
- d) exposing said expressed elongase enzyme to a substrate polyunsaturated fatty acid in order to convert said substrate to a product polyunsaturated fatty acid.

30. The method according to claim 29, wherein said substrate polyunsaturated fatty acid is selected from the group consisting of AA and EPA, and said product polyunsaturated fatty acid is selected from the group consisting of ADA and ω 3-docosapentaenoic acid, respectively.

31. The method according to claim 29 further comprising the step of exposing said expressed elongase enzyme to at least one desaturase in order to convert said product polyunsaturated fatty acid to another polyunsaturated fatty acid.

32. The method according to claim 31 wherein said product polyunsaturated fatty acid is selected from the group consisting of ADA and ω 3-docosapentaenoic acid, said another polyunsaturated fatty acid is selected from the group consisting of ω 6-docosapentaenoic acid, ω 3-docosapentaenoic acid and docohexaenoic acid, and said at least one desaturase is Δ 4-desaturase with respect to production of ω 6-docosapentaenoic acid or docosahexaenoic acid, and Δ 19-desaturase with respect to production of ω 3-docosapentaenoic acid or docosahexaenoic acid.

33. The method of claim 32 further comprising the step of exposing said another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of at least one elongase and at least one additional desaturase in order to convert said another polyunsaturated fatty acid to a final polyunsaturated fatty acid.

34. The method of claim 33 wherein said final polyunsaturated fatty acid is docosahexaenoic acid.

35. A nutritional composition comprising at least one polyunsaturated fatty acid selected from the group consisting of said product polyunsaturated fatty acid produced according to the method of claim 29, said another polyunsaturated fatty acid produced according to the method of claim 31, and said final polyunsaturated fatty acid produced according to the method of claim 33.

36. The nutritional composition of claim 35 wherein said product polyunsaturated fatty acid is selected from the group consisting of ADA and ω 3-docosapentaenoic acid.

37. The nutritional composition of claim 35 wherein said another polyunsaturated fatty acid is selected from the group consisting of ω 6-docosapentaenoic acid, ω 3-docosapentaenoic acid and docosahexaenoic acid.

38. The nutritional composition of claim 35 wherein said final polyunsaturated fatty acid is docosahexaenoic acid.

39. The nutritional composition of claim 35 wherein said nutritional composition is selected from the group consisting of an infant formula, a dietary supplement and a dietary substitute.

40. A pharmaceutical composition comprising 1) at least one polyunsaturated fatty acid selected from the group consisting of said product polyunsaturated fatty acid produced according to the method of claim 29, said another polyunsaturated fatty acid produced according to the method of claim 31, and said final polyunsaturated fatty acid produced according to the method of claim 33 and 2) a pharmaceutically acceptable carrier.

41. An animal feed comprising at least one polyunsaturated fatty acid selected from the group consisting of said product polyunsaturated fatty acid produced according to the method of claim 29, said another polyunsaturated fatty acid produced according to the method of claim 31 and said final polyunsaturated fatty acid produced according to the method of claim 33.

42. A method of preventing or treating a condition caused by insufficient intake of polyunsaturated fatty acids comprising administering to said patient said nutritional composition of claim 35 in an amount sufficient to effect said prevention or treatment.