



PCT WELTORGANISATION FÜR GEISTIGES EIGENTUM
Internationales Büro
INTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)

<p>(51) Internationale Patentklassifikation ⁷ : C07K 7/64, 7/06, 5/08, 5/06</p>	A1	<p>(11) Internationale Veröffentlichungsnummer: WO 00/53627</p> <p>(43) Internationales Veröffentlichungsdatum: 14. September 2000 (14.09.00)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP00/01751</p> <p>(22) Internationales Anmeldedatum: 1. März 2000 (01.03.00)</p> <p>(30) Prioritätsdaten: 199 10 727.0 11. März 1999 (11.03.99) DE</p> <p>(71) Anmelder (für alle Bestimmungsstaaten ausser US): MERCK PATENT GMBH [DE/DE]; Frankfurter Strasse 250, D-64293 Darmstadt (DE).</p> <p>(72) Erfinder; und (75) Erfinder/Anmelder (nur für US): JONCZYK, Alfred [DE/DE]; Schepp Allee 57, D-64295 Darmstadt (DE). ARNOLD, Markus [DE/DE]; Wilhelm-Leuschner-Strasse 31, D-63263 Neu-Isenburg (DE).</p> <p>(74) Gemeinsamer Vertreter: MERCK PATENT GMBH; D-64271 Darmstadt (DE).</p>	<p>(81) Bestimmungsstaaten: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO Patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Veröffentlicht <i>Mit internationalem Recherchenbericht. Vor Ablauf der für Änderungen der Ansprüche zugelassenen Frist; Veröffentlichung wird wiederholt falls Änderungen eintreffen.</i></p>	
<p>(54) Title: METHOD FOR PRODUCING CYCLO-(Asp-DPhe-NMeVal-Arg-Gly)</p> <p>(54) Bezeichnung: VERFAHREN ZUR HERSTELLUNG VON CYCLO-(Asp-DPhe-NMeVal-Arg-Gly)</p> <p>(57) Abstract</p> <p>The invention relates to a method for producing the cyclic pentapeptide cyclo-(Arg-Gly-Asp-DPhe-NMeVal) by cyclising a linear pentapeptide chosen from the group H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH, H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg (Pbf)-OH, H-Asp(OBzl)-DPhe-NMeVal-Arg (Pbf)-Gly-OH, H-DPhe-NMeVal-Arg (Pbf)-Gly-Asp (OBzl)-OH or H-NMeVal-Arg (Pbf)-Gly-Asp (OBzl)-DPhe-OH, subsequently separating protective groups and optionally further converting into the physiologically unobjectionable salts thereof.</p> <p>(57) Zusammenfassung</p> <p>Verfahren zur Herstellung des cyclischen Pentapeptids cyclo-(Arg-Gly-Asp-DPhe-NMeVal) durch Cyclisierung eines linearen Pentapeptids, ausgewählt aus der Gruppe H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH, H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg (Pbf)-OH, H-Asp(OBzl)-DPhe-NMeVal-Arg (Pbf)-Gly-OH, H-DPhe-NMeVal-Arg (Pbf)-Gly-Asp (OBzl)-OH oder H-NMeVal-Arg (Pbf)-Gly-Asp (OBzl)-DPhe-OH, anschließender Schutzgruppenabspaltung und gegebenenfalls weitere Umwandlung in dessen physiologisch unbedenkliche Salze.</p>		

Process for the preparation of cyclo(Asp-DPhe-NMeVal-Arg-Gly)

The invention relates to a novel process for the preparation of the cyclic pentapeptide cyclo(Arg-Gly-Asp-DPhe-NMeVal).

Cyclic pentapeptides, amongst them also cyclo(Arg-Gly-Asp-DPhe-NMeVal) and its physiologically acceptable salts, are disclosed in EP 0 770 622. The present invention is to be regarded as a selection invention in relation to EP 0 770 622.

Generally, cyclic peptides are obtained by cyclization of a linear precursor molecule under the customary conditions of peptide synthesis.

In order that selective linkage of two amino acids or two segments, consisting of a number of amino acids, or alternatively a cyclization of a linear peptide, can be guaranteed, the corresponding functionalities of the amino acids which are not intended to participate in the reaction are to be blocked by appropriate protective groups. Various types of protective groups for amino, carboxyl, hydroxyl, thiol or carboxamide functions, and also for guanidine functions or for the imidazole nitrogen, were therefore developed which, in their combination, make possible a wide possibility of variation with respect to the optimization of the reactions mentioned beforehand. The synthesis of the linear precursor molecules, the linear peptides, can moreover be carried out by means of two methods, on the one hand by means of a solid-phase peptide synthesis, on the other hand in solution. In this case, stepwise couplings of the amino acids or fragment condensations of segments of amino acids are possible. The respective coupling steps can in turn be carried out using different condensation reagents, such as carbodiimides, carbodiimidazole, those of the uronium type such as TBTU, or according to mixed anhydride methods or active ester methods.

The invention was based on the object of developing a novel, improved process for the preparation of cyclo(Arg-Gly-Asp-DPhe-NMeVal), in comparison with the previously known processes.

5 It was surprisingly found that in the synthesis of the cyclopeptide cyclo(Arg-Gly-Asp-DPhe-NMeVal) by cyclization of a linear precursor molecule, the combination of the protective groups 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf) for the
10 guanidino group in the side chain of the arginine and benzyl (Bzl) for the carboxyl group in the side chain of the aspartic acid leads to an optimization with respect to the yield.

The invention therefore relates to a process
15 for the preparation of the cyclic pentapeptide
cyclo(Arg-Gly-Asp-DPhe-NMeVal)
by cyclization of a linear pentapeptide selected from the group consisting of

20 H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH,
H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-OH,
H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH,
H-DPhe-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-OH or
H-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-OH,
subsequent protective group removal and, if
25 appropriate, further conversion into its physiologically acceptable salts.

The invention furthermore relates to a process for the preparation of the cyclic pentapeptide
cyclo(Arg-Gly-Asp-DPhe-NMeVal)
30 as described, characterized in that the linear pentapeptide H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH is cyclized.

The reaction conditions of this cyclization of the linear peptides selected from the group consisting
35 of

H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH,
H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-OH,
H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH,

H-DPhe-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-OH or

H-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-OH

with respect to the choice of the dehydrating agent,
the inert solvent and the reaction temperature and the
5 further conversion into its physiologically acceptable
salts have already been disclosed in EP 0 770 622.

The removal of the benzyl protective group on
the side chain of the aspartic acid can be removed
under customary conditions (for this cf.: T.W. Greene,
10 P.G.M. Wuts, Protective Groups in Organic Chemistry, 2nd
Ed., Wiley, New York 1991 or P.J. Kocienski, Protecting
Groups, 1st Ed., George Thieme Verlag, Stuttgart - New
York, 1994, H. Kunz, H. Waldmann in Comprehensive
Organic Synthesis, Vol. 6 (Ed. B.M. Trost, I. Fleming,
15 E. Winterfeldt), Pergamon, Oxford. 1991, pp. 631-701),
e.g. by treating with hydrogen in the presence of a
catalyst (e.g. of a noble metal catalyst such as
palladium, expediently on a support such as carbon).
Suitable solvents are, for example, alcohols such as
20 methanol or ethanol or amides such as DMF or
alternatively mixtures with further inert solvents,
such as, for example, mixtures with water. As a rule,
the hydrogenolysis is carried out at temperatures
between approximately 0 and 100°C and pressures between
25 approximately 1 and 200 bar, preferably at 20-30°C and
1-10 bar.

The Pbf protective group, which was introduced
into peptide chemistry by L.A. Carpino et al., Tet.
Lett. 1993, 34, 7829-7832, is removed, for example, by
30 treating with 95% trifluoroacetic acid (TFA). The Pbf
protective group in this case shows a greater lability
with respect to TFA than the structurally similar
protective groups 4-methoxy-2,3,6-
trimethylphenylsulfonyl (Mtr) and 2,2,5,7,8-
35 pentamethylchroman-6-sulfonyl (Pmc), which are also
possible as side-chain protective groups with respect
to the synthesis of cyclo(Arg-Gly-Asp-DPhe-NMeVal).

TFA is preferably used in an excess without addition of a further solvent. TFA can also be employed as a mixture with an inert solvent, such as, for example, the combination TFA/dichloromethane in the ratio 6:4.
5 TFA can furthermore also be employed with an addition of 1-10%, preferably 2%, of water. The reaction temperature for the cleavage is expediently between approximately 0 and approximately 50°C, and the reaction is preferably carried out between 15 and 30°C
10 (room temperature).

The abbreviations of amino acids listed above and below stand for the radicals of the following amino acids:

Asp	Aspartic acid
15 Arg	Arginine
Gly	Glycine
Phe	Phenylalanine
Val	Valine

Furthermore, above and below the following have
20 the meanings:

Boc	tert-Butoxycarbonyl
Bzl	Benzyl
CHA	Cyclohexylamine
D	Characterization of a D-amino acid
25 DCCI	Dicyclohexylcarbodiimide
DMAP	Dimethylaminopyridine
DMF	Dimethylformamide
EDCI	N-Ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride
30 Et	Ethyl
Fmoc	9-Fluorenylmethoxycarbonyl
HOBt	1-Hydroxybenzotriazole
Me	Methyl
MTBE	Methyl tert-butyl ether
35 Mtr	4-Methoxy-2,3,6-trimethylphenylsulfonyl
NMe	N-methylated α -amino group
NMP	N-methylpyrrolidone
OtBu	tert-Butyl ester

	OMe	Methyl ester
	OEt	Ethyl ester
	Pbf	2,2,4,6,7-Pentamethyldihydrobenzofuran-5-sulfonyl
5	Pmc	2,2,5,7,8-pentamethylchroman-6-sulfonyl
	POA	Phenoxyacetyl
	Pr	Propyl
	Su	Succinimide
	TBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyl-
10		uronium tetrafluoroborate
	TFA	Trifluoroacetic acid
	Z	Benzyloxycarbonyl

It was furthermore surprisingly found that the specific selection of the side-chain protective groups Pbf on Arg and Bzl on Asp even in the synthesis of the linear peptides, which, as mentioned beforehand, are intermediates in the synthesis of cyclo(Arg-Gly-Asp-DPhe-NMeVal), leads to improved yields in the respective synthesis stages. As a result, the yield of cyclo(Arg-Gly-Asp-DPhe-NMeVal) is increased overall and as a result thereof the costs of the synthesis are lowered.

The improvement in yield is achieved in this case both in the solid-phase peptide synthesis and in the synthesis in solution of the linear peptides H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH, H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-OH, H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH, H-DPhe-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-OH or H-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-OH, in particular H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH.

The protected amino acids or amino acid fragments used in the two synthesis methods are customarily prepared by methods of amino acid and peptide synthesis, such as described in the standard works Principles of Peptide Synthesis, ed. M. Bodansky, Springer Verlag Berlin 1984; Houben Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry],

1.c., Volume 15/11, 1974, pages 1 to 806, Georg Thieme Verlag, Stuttgart; Calbiochem/Novabiochem Catalogue and Synthesis Handbook 1999; Synthesis Notes or Peptide Synthesis Protocols, eds. M.W. Pennington and B.M. Dunn
5 in Methods in Molecular Biology, Vol. 35, Humana Press Totowa N.J. 1994, namely under reaction conditions which are known and suitable for the reactions mentioned. Use can also be made in this case of variants which are known, but not mentioned here in
10 greater detail.

The underlying principles of solid-phase peptide synthesis have been laid down by B.F. Gysin and R.B. Merrifield (J. Am. Chem. Soc. 1972, 94, 3102 ff.). The solid-phase synthesis of the linear peptides
15 described beforehand, their removal and purification is carried out as described by A. Jonczyk and J. Meienhofer in Peptides, Proc. 8th Am. Pept. Symp., Eds. V. Hruby and D.H. Rich, Pierce Comp. III, p. 73-77, 1983 or analogously to the techniques described in
20 Angew. Chem. 1992, 104, 375-391.

Particularly preferably, the synthesis of one of the linear peptides, such as described beforehand, takes place convergently by fragment condensation.

The invention therefore furthermore relates to
25 a process for the preparation of the cyclic pentapeptide $\text{cyclo}(\text{Arg-Gly-Asp-DPhe-NMeVal})$, characterized in that the synthesis of the linear peptide $\text{H-Asp}(\text{OBzl})\text{-DPhe-NMeVal-Arg}(\text{Pbf})\text{-Gly-OH}$ is carried out convergently by fragment condensation of a
30 tripeptide $\text{R}^1\text{-Asp}(\text{OBzl})\text{-DPhe-NMeVal-OH}$, where R^1 is an amino protective group, with a dipeptide $\text{H-Arg}(\text{Pbf})\text{-Gly-R}^2$, where R^2 is a carboxyl protective group, and the protective groups R^1 and R^2 are then removed.

R^1 , as described beforehand, is an amino
35 protective group. The expression "amino protective group" is generally known and relates to groups which are suitable for protecting (or blocking) an amino group from chemical reactions. Typical groups of this

type are, in particular, unsubstituted or substituted acyl, aryl, aralkoxymethyl or aralkyl groups. As the amino protective groups are removed after the desired reaction (or reaction sequence), their nature and size is otherwise uncritical; however, those having 1-20 C atoms are preferred. The expression "acyl group" is to be interpreted in the widest sense in connection with the present process. It includes acyl groups derived from aliphatic, araliphatic, alicyclic, aromatic or heterocyclic carboxylic acids or sulfonic acids and also, in particular, alkoxy-carbonyl, alkenyloxy-carbonyl, aryloxy-carbonyl and especially aralkoxy-carbonyl groups. Examples of acyl groups of this type are formyl or alkanoyl such as acetyl, propionyl, butyryl; aralkanoyl such as phenylacetyl; aroyl such as benzoyl or toluyl; aryloxyalkanoyl such as phenoxyacetyl; alkoxy-carbonyl such as methoxy-carbonyl, ethoxy-carbonyl, 2,2,2-trichloroethoxy-carbonyl, Boc, 2-iodoethoxy-carbonyl; alkenyloxy-carbonyl such as aryloxy-carbonyl (Aloc), aralkyloxy-carbonyl such as CBZ (synonymous with Z), 4-methoxybenzyloxy-carbonyl (MOZ), 4-nitrobenzyloxy-carbonyl or 9-fluorenylmethoxy-carbonyl (Fmoc); 2-(phenylsulfonyl)ethoxy-carbonyl; trimethylsilyl-ethoxy-carbonyl (Teoc) or arylsulfonyl such as 4-methoxy-2,3,6-trimethylphenylsulfonyl (Mtr). Trityl (Trt) is furthermore also known as an amino protective group. Preferred amino protective groups are Boc, Fmoc and Aloc, furthermore Z, benzyl and acetyl. Boc is particularly preferred.

R^2 , as described beforehand, is a carboxyl protective group. The expression "carboxyl protective group" is also generally known and relates to groups which are suitable for protecting the hydroxyl group of a carboxylic acid from chemical reactions. Typical groups of this type are the abovementioned unsubstituted or substituted aryl, aralkyl, aroyl or acyl groups, furthermore also alkyl groups, alkyl, aryl

or aralkylsilyl groups. The nature and size of the carboxyl protective groups and synonymously therewith of the hydroxyl protective groups is uncritical, as they are removed again after the desired chemical reaction or reaction sequence; groups having 1-20, in particular 1-10, C atoms are preferred. Examples of hydroxyl protective groups are, inter alia, aralkyl groups such as benzyl, 4-methoxybenzyl or 2,4-dimethoxybenzyl, aroyl groups such as benzoyl or p-nitrobenzoyl, acyl groups such as acetyl or pivaloyl, p-toluenesulfonyl, alkyl groups such as methyl or tert-butyl, but also allyl, alkylsilyl groups such as trimethylsilyl (TMS), triisopropylsilyl (TIPS), tert-butyl dimethylsilyl (TBS) or triethylsilyl, trimethylsilylethyl or aralkylsilyl groups such as tert-butyl diphenylsilyl (TBDPS). Preferred hydroxyl protective groups are methyl, benzyl, acetyl, tert-butyl or TBS. Methyl and tert-butyl are particularly preferred.

In general, preferred protective groups for the carboxyl group of the aspartic acid side chain are linear or branched alkyl groups, such as methyl, ethyl or tert-butyl, or arylalkyl groups, such as benzyl; in the process according to the invention benzyl.

In general, preferred protective groups for the guanidino group of the arginine side chain are Z, Boc, Mtr or Pmc; in the process according to the invention Pbf.

The liberation of the protective group used in each case is known from the literature (e.g. T.W. Greene, P.G.M. Wuts, Protective Groups in Organic Chemistry, 2nd Ed., Wiley, New York 1991 or P.J. Kocienski, Protecting Groups, 1. Ed., Georg Thieme Verlag, Stuttgart - New York, 1994). Use can also be made in this case of variants which are known per se, but not mentioned here in greater detail.

The invention relates to a process, as described beforehand, characterized in that the

tripeptide R^1 -Asp(OBzl)-DPhe-NMeVal-OH, where R^1 is an amino protective group, is prepared by linear synthesis by reacting Z-DPhe-OH with H-NMeVal-OMe to give H-DPhe-NMeVal-OMe and then reacting this with an activated R^1 -
5 Asp(OBzl)-OH derivative and cleaving the methyl ester.

A preferred derivative of R^1 -Asp(OBzl)-OH is the succinimide R^1 -Asp(OBzl)-OSu. Further active esters which can be employed are known from the customary literature on peptide synthesis, as described
10 beforehand.

The invention furthermore relates to a process, such as described beforehand, characterized in that the dipeptide H-Arg(Pbf)-Gly- R^2 , where R^2 is a carboxyl protective group, is prepared by linear synthesis by
15 reacting Z-Arg(Pbf)-OH and the protective group Z is removed.

The invention likewise relates to the linear pentapeptides, selected from the group consisting of
H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH,
20 H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-OH,
H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH,
H-DPhe-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-OH or
H-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-OH,
as intermediates in the synthesis of cyclo(Arg-Gly-Asp-
25 DPhe-NmeVal).

The invention relates to a process for the preparation of the cyclic pentapeptide



by

- 30 a) reacting a dipeptide H-Arg(Pbf)-Gly- R^2 , prepared by linear synthesis of Z-Arg(Pbf)-OH with H-Gly- R^2 and subsequent removal of the protective group Z, with
- b) a tripeptide R^1 -Asp(OBzl)-DPhe-NMeVal-OH, prepared
35 by linear synthesis of Z-DPhe-OH with H-NMeVal-OMe and removal of the protective group Z to give H-DPhe-NMeVal-OMe and subsequent coupling of this

peptide to an active ester of R¹-Asp(OBzl)-OH and subsequent removal of the methyl ester, convergently to give the linear pentapeptide R¹-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-R²,

- 5 c) removing the protective groups R¹ and R²,
- d) cyclizing the liberated pentapeptide H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH to give cyclo(Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal),
- e) removing the benzyl protective group,
- 10 f) removing the Pbf protective group with TFA and

optionally converting the trifluoroacetate of cyclo(Arg-Gly-Asp-DPhe-NMeVal) produced by the steps a-f) into further physiologically acceptable salts.

- 15 Further physiologically acceptable salts are, for example, salts of inorganic acids, such as sulfuric acid, sulfurous acid, dithionic acid, nitric acid, hydrohalic acids such as hydrochloric acid or hydrobromic acid, phosphoric acids such as, for
- 20 example, orthophosphoric acid, sulfamic acid, furthermore of organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic mono- or polybasic carboxylic, sulfonic or sulfuric acids, e.g. formic acid, acetic acid, propionic acid, hexanoic
- 25 acid, octanoic acid, decanoic acid, hexadecanoic acid, octadecanoic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid,
- 30 nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, benzenesulfonic acid, trimethoxybenzoic acid, adamantanecarboxylic acid, p-toluenesulfonic acid, glycolic acid, embonic acid, chlorophenoxyacetic acid, aspartic acid, glutamic acid,
- 35 proline, glyoxylic acid, palmitic acid, parachlorophenoxyisobutyric acid, cyclohexanecarboxylic acid, glucose-1-phosphate, naphthalenemono- and disulfonic acids or laurylsulfuric acid.

Particularly preferred physiologically acceptable salts are the hydrochloride or the internal salts of cyclo(Arg-Gly-Asp-DPhe-NMeVal).

5 If R¹ is Boc and R² is tert-butyl, these terminal tert-butyl-like protective groups can be cleaved with formic acid without the side-chain protective groups Pbf and Bzl being attacked.

The following examples describe specific embodiments of the individual synthesis steps.

10 Above and below, all temperatures are indicated in °C.

Example 1:

Synthesis of Boc-Asp(OBzl)-DPhe-NMeVal-OH

15

1. 25.3 ml of trimethylchlorosilane are added dropwise to a solution of 26.5 g of Z-NMeVal-OH in 200 ml of methanol and the mixture is stirred overnight. The reaction mixture is freed from the solvent and the residue is taken up in methyl tert-butyl ether (MTBE) and washed with 5% Na₂CO₃ and water. After removal of the solvent, the residue is taken up in methanol and 1N HCl, mixed with water-moist Pd/C (10%) and hydrogenated in a gentle stream of H₂. After 20 completion of the reaction, the catalyst is filtered off, the filtrate is freed from the solvent and the residue is recrystallized from ethyl acetate. H-NMeVal-OMe hydrochloride is obtained in a yield of 75%.

2. A solution of 16.1 g of Z-DPhe-OH, 10 g of H-NMeVal-OMe hydrochloride and 10.1 ml of diisopropyl-ethylamine in 100 ml of dichloromethane is cooled to 0-5° and 11.35 g of EDCI are added. The mixture is first stirred for one hour at 0-5°, then overnight at room temperature. The solvent is removed and the 35 residue is taken up in MTBE and washed with Na₂CO₃ (5%), 1N HCl and water and dried. After removal of the solvent, Z-DPhe-NMeVal-OMe is obtained in a yield of 84.5%.

3. 12 g of Z-CPhe-NMeVal-OMe are dissolved in 80 ml of THF and 20 ml of water, mixed with 10 mg of thymolphthalein and treated dropwise with 10M NaOH until a blue colouration of the indicator is obtained.

5 On decolourization of the indicator, the mixture is repeatedly treated dropwise with 10M NaOH. When a decolourization of the indicator is no longer observed, it is adjusted to pH 2 with 10% aqueous KHSO₄ solution, methanol is removed and the product is extracted with
10 MTBE. After drying with Na₂SO₄, the CHA salt is precipitated from the filtrate by addition of 2.9 ml of CHA. Z-DPhe-NMeVal-OH x CHA is obtained in a yield of 90%.

4. 16.4 g of Z-DPhe-NMeVal-OH x CHA (cyclohexyl-
15 ammonium salts) are stirred in 250 ml of MTBE and 100 ml of H₃PO₄ (10%) until everything has dissolved. After removal of the aqueous phase, the organic phase is washed with water and saturated NaCl solution and dried. The solvent is removed and the residue is taken
20 up in 15.2 ml of 2N NaOH and 150 ml of THF and, after addition of the moist catalyst (1 g of Pd/C (10%)), hydrogenated in a gentle stream of H₂. The catalyst is filtered off and the clear solution is treated with 12.1 g of Boc-Asp(OBzl)-OSu and 4.5 ml of triethylamine
25 and stirred overnight at room temperature. After removal of the solvent, the residue is taken up in MTBE and washed with H₃PO₄ (10%), water and saturated NaCl solution. The organic phase is treated with 3.3 ml of CHA. The resulting salt Boc-Asp(OBzl)-DPhe-NMeVal-OH x
30 CHA is filtered off and dried in vacuo. The yield is 93%.

Example 2:

Synthesis of H-Arg(Pbf)-Gly-OtBu

35

1. 33.0 g of Z-Arg(Pbf)-OH x CHA are stirred in 300 ml of acetic acid and 300 ml of H₃PO₄ until everything has dissolved. After removal of the aqueous

phase, the organic phase is washed with water and saturated NaCl solution and dried. After removal of the solvent, the residue is dissolved in 250 ml of dichloromethane with 8.38 g of H-Gly-OtBu x HCl and cooled to 0°. 17.12 ml of diisopropylethylamine and 16.05 g of TBTU are then added and the mixture is stirred for 60 min at 0° and overnight at room temperature. The solvent is removed and replaced by 250 ml of ethyl acetate. After washing with Na₂CO₃ solution (5%), water and saturated NaCl solution, the solvent is removed. Z-Arg(Pbf)-Gly-OtBu is obtained in a yield of 86%.

2. A solution of 30 g of Z-Arg(Pbf)-Gly-OtBu in 350 ml of THF is mixed with 3 g of water-moist Pd/C (10%) and the Z group is removed by hydrogenation under a gentle stream of H₂. The catalyst is then filtered off and the solvent is removed. The residue is taken up in ethyl acetate and treated further as in Example 2.1. H-Arg(Pbf)-Gly-OtBu is obtained in a yield of 86%.

20

Example 3:

Synthesis of H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH

1. 1.23 g of Boc-Asp(OBzl)-DPhe-NMeVal-OH x CHA are converted into the free acid in the customary manner and this is dissolved in 12.5 ml of dichloromethane with 0.81 g of H-Arg(Pbf)-Gly-OtBu and 0.22 g of DMAP. The solution is cooled to 0-5° and treated with 0.345 g of EDCI. It is stirred for 2 hours at 0-5° and overnight at room temperature. The solvent is removed, the residue is taken up in MTBE and treated further as in Example 2.1. Boc-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OtBu is obtained in a yield of 82%.

2. 2.3 g of Boc-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OtBu are dissolved in 23 ml of 95% formic acid for the removal of the terminal protective groups and concentrated in vacuo after 30 min. The product is triturated with ether, filtered off and dried in vacuo.

35

H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH x HCOOH is obtained in a yield of 95%.

Example 4:

5 Synthesis of cyclo(Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal)

A solution of 11.9 g of H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH x HCOOH in 60 ml of NMP is added dropwise to a stirred solution of 7.25 g of TBTU and
10 7.45 ml of N-methylmorpholine in 180 ml of N-methylpyrrolidone. The reaction solution is stirred for 20 hours and then added dropwise to a solution of 47.5 g of NaHCO₃ in 1800 ml of water. The precipitate is filtered and dried in vacuo. Cyclo(Arg(Pbf)-Gly-
15 Asp(OBzl)-DPhe-NMeVal) is obtained in a yield of 73.4%.

Example 5:

Synthesis of cyclo(Arg-Gly-Asp-DPhe-NMeVal)

20 1. A solution of 2 g of cyclo(Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal) in 26 ml of THF is mixed with 0.5 g of Pd/C (10%). Hydrogen is passed through for 2 hours, the mixture is freed of the catalyst and the solvent is removed in vacuo. The product crystallizes
25 after addition of 32 ml of acetone, and is filtered off and dried. Cyclo(Arg(Pbf)-Gly-Asp-DPhe-NMeVal) is obtained in a yield of 83%.

2. 1.5 g of cyclo(Arg(Pbf)-Gly-Asp-DPhe-NMeVal) are dissolved in 15 ml of 95% TFA. After 1 hour, the
30 solution is added dropwise to 150 ml of isopropyl ether and the solid is filtered off and dried. The dried product is dissolved in 30 ml of isopropanol/water 1:2 and treated with ion exchanger III (acetate form; Merck KGaA). The filtered solution is concentrated and
35 freeze-dried. Cyclo(Arg-Gly-Asp-DPhe-NMeVal) is obtained as an internal salt in a yield of 96%.

Patent Claims

1. Process for the preparation of the cyclic
5 pentapeptide

cyclo(Arg-Gly-Asp-DPhe-NMeVal)

by cyclization of a linear pentapeptide selected from
the group consisting of

10 H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH,
H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-OH,
H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH,
H-DPhe-NMeVal-Arg(Pbf)-Gly-Asp(OBz)-OH or
H-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-OH,

subsequent protective group removal and, if
15 appropriate, further conversion into its
physiologically acceptable salts.

2. Process for the preparation of the cyclic
pentapeptide

cyclo(Arg-Gly-Asp-DPhe-NMeVal)

20 according to Claim 1, characterized in that the linear
pentapeptide H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH is
cyclized.

3. Process according to Claim 1 or 2,
characterized in that the synthesis of the linear
25 peptide H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH is
carried out convergently by fragment condensation of a
tripeptide R¹-Asp(OBzl)-DPhe-NMeVal-OH, where R¹ is an
amino protective group, with a dipeptide H-Arg(Pbf)-
Gly-R², where R² is a carboxyl protective group, and the
30 protective groups R¹ and R² are then removed.

4. Process according to Claims 1 to 3,
characterized in that the tripeptide R¹-Asp(OBzl)-DPhe-
NMeVal-OH, where R¹ is an amino protective group, is
prepared by linear synthesis by reacting Z-DPhe-OH with
35 H-NMeVal-OMe to give H-DPhe-NMeVal-OMe and this is then
reacted with an activated R¹-Asp(OBzl)-OH derivative
and the methyl ester is cleaved.

5. Process according to Claims 1 to 3, characterized in that the dipeptide H-Arg(Pbf)-Gly-R², where R² is a carboxyl protective group, is prepared by
5 linear synthesis by reacting Z-Arg(Pbf)-OH with H-Gly-R² and removing the protective group Z.

6. Linear pentapeptides, selected from the group consisting of

10 H-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-NMeVal-OH,
H-Gly-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-OH,
H-Asp(OBzl)-DPhe-NMeVal-Arg(Pbf)-Gly-OH,
H-DPhe-NMeVal-Arg(Pbf)-Gly-Asp(OBz)-OH or
H-NMeVal-Arg(Pbf)-Gly-Asp(OBzl)-DPhe-OH,
as intermediates in the synthesis of cyclo(Arg-Gly-Asp-
15 DPhe-NMeVal).