

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2222706 T4**

(12) **Oversættelse af ændret
europæisk patentskrift**

-
- (51) Int.Cl.: **C 07 K 16/28 (2006.01)** **A 61 K 39/395 (2006.01)** **A 61 P 29/00 (2006.01)**
A 61 P 37/06 (2006.01) **C 12 N 15/13 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-11-21**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om opretholdelse af patentet i ændret form: **2016-08-10**
- (86) Europæisk ansøgning nr.: **08861706.3**
- (86) Europæisk indleveringsdag: **2008-12-15**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-09-01**
- (86) International ansøgning nr.: **EP2008067499**
- (87) Internationalt publikationsnr.: **WO2009077483**
- (30) Prioritet: **2007-12-14 WO PCT/EP2007/063979** **2008-08-28 EP 08163163**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **Novo Nordisk A/S, Novo Allé, 2880 Bagsværd, Danmark**
- (72) Opfinder: **SVENSSON, Anders, Roskildevägen 17 B, 217 46 Malmö, Sverige**
URSÄ, Birgitte, Skt. Kjeldsgade 12, 3.th., 2100 København Ø, Danmark
WAGTMANN, Peter Andreas Nicolai Reumert, Vallerød Banevej 34, 2960 Rungsted Kyst, Danmark
PEDERSEN, Inger Lund, Vangsåvej 10, 2720 Vanløse, Danmark
- (74) Fuldmægtig i Danmark: **Budde Schou A/S, Hausergade 3, 1128 København K, Danmark**
- (54) Benævnelse: **Antistoffer der binder til NKG2D og brugen heraf**
- (56) Fremdragne publikationer:
WO-A-2007/005874
US-A1- 2005 158 307
US-A1- 2006 280 755
BAUER S ET AL: "Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA" SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, US, WASHINGTON, DC, vol. 285, no. 5428, 30 July 1999 (1999-07-30), pages 727-729, XP002177592 ISSN: 0036-8075 cited in the application
LI, P. ET AL: "Complex Structure of the activating immunoreceptor NKG2D and its MHC class I-like ligand MICA", NATURE IMMUNOLOGY, vol. 2, no. 5, May 2001 (2001-05), pages 443-451, XP009153395, WO-A2-02/068615
WO-A2-2005/097160
WO-A2-2007/042573
KWONG K.Y. ET AL: 'Generation, affinity maturation, and characterization of a human anti-human NKG2D monoclonal antibody with dual antagonistic and agonistic activity' J MOL BIOL. vol. 384, no. 5, 31 December 2008, pages 1143 - 1156
Experimental report concerning KYK-2.0.

Fortsættes ...

MINCHEVA-NILSSON L. ET AL: 'Placenta-derived soluble MHC class I chain-related molecules down-regulate NKG2D receptor on peripheral blood mononuclear cells during human pregnancy: a possible novel immune escape mechanism for fetal survival' THE JOURNAL OF IMMUNOLOGY vol. 176, 2006, pages 3585 - 3592

ANDRÉ P. ET AL: 'Comparative analysis of human NK cell activation induced by NKG2D and natural cytotoxicity' EUR. J. IMMUNOL. vol. 34, 2004, pages 961 - 971

GEFFROY-LUSEAU A. ET AL: 'Allogeneic T lymphocytes as a source of peptide-dependent T cells specific for myeloma cells' INTERNATIONAL IMMUNOLOGY vol. 17, no. 9, 18 July 2005, pages 1193 - 1200

TANAKA J. ET AL: 'Cytolytic activity and regulatory functions of inhibitory NK cell receptor-expressing T cells expanded from granulocyte colony-stimulating factor-mobilized peripheral blood mononuclear cells' BLOOD vol. 104, no. 3, 01 August 2004, pages 768 - 774

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates to antibodies against human NKG2D (hNKG2D) and their use in treating or preventing diseases and disorders in human patients, as further defined in the claims.

BACKGROUND OF THE INVENTION

[0002] The immunoreceptor NKG2D is normally expressed on human CD8⁺ T cells and NK cells. On pre-activated CD8⁺ cells, the human NKG2D (hNKG2D) homodimeric receptor functions as a co-stimulator of TCR and CD28+TCR signalling via its DAP10 association, whereas in NK cells it functions as a direct activator. Various ligands for hNKG2D have been identified and characterized, including the MHC Class I-related ligands MICA and MICB, the UL16-binding protein (ULBP) family, and the retinoic acid early transcript-1 (RAET1) family.

[0003] In chronic autoimmune diseases such as rheumatoid arthritis, hNKG2D is expressed on a sub-set of CD4⁺ CD28⁻ T cells and is involved in stimulation of their proliferation and IFN γ production, and MIC expression is upregulated (Groh et al., PNAS 2003;100:9452). It has also been shown that CD4⁺ hNKG2D-expressing T cells in Crohn's disease mediate inflammatory and cytotoxic responses through MICA interactions (Allez et al., Gastroenterology 2007;132:2346-2358). An initial suggestion that NKG2D is an essential driver in autoimmune inflammation came from the prevention and treatment of the inflammation leading to diabetes in a murine model of diabetes (NOD mice) by a monoclonal antibody (mAb) binding to and blocking murine NKG2D (CX5) (Ogasawara et al., Immunity 2004; 20:757-767), suggesting therapeutic applications for anti-NKG2D antibodies. Such applications have been described in, e.g., US20050158307, WO2005097160, WO2005115517, and WO2006024367.

[0004] Most murine mAbs against human NKG2D that have been described (see, e.g., Pende et al., Eur J Immunol 2001;31:1076-86, WO02068615; Bauer et al., Science 1999;285:727-9; Castriconi et al., PNAS 2003;100:4120-25; and André et al., Eur J Immunol 2004;34:1-11) or are commercially available (e.g., antibody 149810 from R&D Systems, MN, USA, and ON72 from Beckman Coulter Inc.), are immunogenic.

[0005] US2006280755 relates to NKG2D modulating agents for treatment of autoimmune diseases, inflammatory disorders and allograft rejection. The modulating agents are characterised as to reduce the interaction of NKG2D with DAP10, reduce the amount of NKG2D on the cell surface, reduce NKG2D-mediated activation and have a KD of less than 1nM. Said modulation agents include fully human monoclonal antibodies directed against NKG2D (and preparation thereof). However, no human monoclonal antibodies have been generated.

[0006] The only fully human anti-NKG2D mAb described in the literature reportedly had both agonistic and antagonistic effects on NKG2D-signalling (Kwong et al., J Mol Biol 2008;384:1143-1156), rendering it less suitable as a therapeutic agent for inflammatory and/or autoimmune disorders.

[0007] Accordingly, there is a need for anti-hNKG2D mAbs with optimal properties for therapeutic use in inflammatory and/or autoimmune diseases and disorders. The present invention addresses these and other needs and provides several additional benefits that will be described in the remainder of this document.

[0008] WO2007/005874 discloses a human monoclonal antibodies to programmed death ligand 1 (PD-L1) wherein antibody designated 5F8 comprises the identical VL region (i.e. SEQ. ID. NO:14) as the MS antibody of the current invention.

SUMMARY OF THE INVENTION

[0009] The present invention provides isolated anti-hNKG2D monoclonal antibodies useful for therapeutic applications in humans, as defined in the claim. The antibodies are fully human or humanized to minimize the risk for immune responses against the antibodies when administered to a patient. As described herein, other antigen-binding molecules such as, e.g., antigen-binding antibody fragments, antibody derivatives, and multi-specific molecules, can be designed or derived from such antibodies.

[0010] In one aspect, the antibodies are further characterized by one or more functional properties, or by a combination of functional properties. Exemplary properties include, *e.g.*, preventing hNKG2D-mediated activation of hNKG2D-expressing NK or T cell; competing with at least one natural hNKG2D ligand, or with several ligands, in binding to hNKG2D; reducing the amount of hNKG2D on the surface of a hNKG2D-expressing NK or T cell; binding also cynomolgous and/or rhesus NKG2D; binding only one antibody molecule per hNKG2D dimer; cross-linking no more than 2 hNKG2D dimers when added to hNKG2D-expressing NK and/or T cells; having insignificant agonist effect on hNKG2D signalling upon binding; and/or binding to hNKG2D with a dissociation constant (KD) of 1 nM or less. Certain anti-hNKG2D antibodies of the invention may also or alternatively compete with, bind to essentially the same epitope as, or bind with the same or higher affinity as, the human anti-hNKG2D antibody MS described herein. For example, in one embodiment, the antibodies are also or alternatively more capable of competing with or blocking hNKG2D-binding of MS than known murine anti-hNKG2D antibodies (*e.g.*, the ones described above). In one embodiment, the antibodies bind to the same hNKG2D epitope as MS. The skilled person will understand that antibodies provided by and/or used in embodiments of this invention may exhibit three, four, or more of the above-referenced features.

[0011] In another aspect, the antibodies also or alternatively comprise one or more paratopes and/or antigen-binding sequences that are identical or similar to MS paratopes and/or antigen-binding sequences described herein.

[0012] The present specification further describes nucleic acids encoding antibodies of the invention, expression vectors comprising such nucleic acids, host cells comprising such nucleic acids, host cells producing antibodies of the invention, and methods of producing anti-hNKG2D antibodies by culturing such host cells under appropriate conditions.

[0013] Antigen-binding fragments of such antibodies, as well as molecules comprising such antigen-binding fragments, including engineered antibody fragments, antibody derivatives, bispecific antibodies and other multispecific molecules, are also provided.

[0014] Pharmaceutical compositions and kits or other articles that comprise such antibodies or other molecules also are provided.

[0015] Further disclosed herein are methods of reducing or inhibiting hNKG2D activation, hNKG2D-signalling, or activation of hNKG2D-expressing NK or T cells by one or more functional properties, or by a combination of functional properties of the antibodies of the current invention.

[0016] Further provided for are antibodies and antigen-binding fragments as defined in the claims for use in reducing inflammation, and for use in treating or preventing autoimmune and/or inflammatory diseases or disorders, including, but not limited to rheumatoid arthritis, inflammatory bowel disease (IBD) including Crohn's disease and ulcerative colitis, systemic erythematosis lupus (SLE), psoriasis, psoriatic arthritis, multiple sclerosis, celiac disease, viral disease (such as, *e.g.*, viral hepatitis), and transplant rejection of various organs and tissues (including, but not limited to, heart and bone marrow).

DESCRIPTION OF THE DRAWINGS

[0017]

Figure 1 shows analyses of exemplary sera from hNKG2D-immunized mice from the KM mouse™ strain. Flow cytometry analysis on NKG2D-expressing BAF/3 cells or control cells (BAF/3 cells) at various time points revealed increasing titers of antibody with NKG2D-selective binding in 1 µl of serum (A). Pre-incubation with sera (1 µl) taken after the 6th immunization contained antibody capable of preventing MICA-Fc-binding to NKG2D (B).

Figure 2 shows an example of a human antibody in the form of a hybridoma supernatant bound specifically to NKG2D expressing cells (A) but not to the same cell-line not expressing NKG2D (B). Antibody was added to the cells in the form of hybridoma supernatant. The binding of a directly labelled positive control, murine anti-NKG2D antibody 149810, to NKG2D expressing cells (C) and non expressing cells (D), is also shown. The black outline represents background staining, and solid peaks represent specific staining.

Figure 3 demonstrates dose-response of NKG2D-binding to NKG2D-expressing cells of recombinantly expressed and purified fully human IgG4 antibodies (16F16, 16F31, MS, and 21 F2) as compared to commercial murine antibodies (ON72 and 149810).

Figure 4 shows the amino acid sequences for the heavy (H) and light (L) chains of human anti-hNKG2D antibodies 16F16, and 16F31 (A), and MS and 21F2 (B) of IgG4 isotype, highlighting variable regions (bold) and CDR regions (underlined). The corresponding sequence identifiers for the amino acid sequences and the various highlighted portions are provided in Table 1.

Figure 5 shows alignments of VH and VL sequences with the corresponding recombined germline sequences. CDR regions are indicated by bold Kabat numbers and somatic hypermutations are indicated by bold underlined text. (A) 16F16 IgG4 H chain; (B) 16F16 IgG4 L chain; (C) 16F31 IgG4 H chain; (D) 16F31 IgG4 L chain; (E) MS IgG4 H chain; (F) MS IgG4 L chain; (G) 21 F2 IgG4 H chain; (H) 21 F2 IgG4 L chain. SEQ ID NOS:27-30 correspond to recombined VH3_21/D3-9/JH4, VKI_L15/JK2, VH3_20/D3-10/JH6, and VKIII_A27/JK3, respectively, and SEQ ID NOS 60-63 correspond to recombined VH4_59/JH3, VKIII_A27/JK1, VH5_51/D3_10_R3/JH4, and VKIII_L6/JK1, respectively.

Figure 6 shows blockade of ligand- (MICA-) binding by an exemplary human anti-NKG2D antibody, demonstrated by blockade of ligand binding by preincubation with antibody in a hybridoma supernatant. The outline represents background, grey represents ligand binding without pre-incubation, and black with dotted line represents ligand binding with pre-incubation.

Figure 7 shows a dose-response curve obtained when analyzing various concentrations of recombinantly expressed and purified fully human anti-hNKG2D antibodies (16F16, 16F31, MS, and 21 F2; IgG4 isotype), giving the IC50 and dose needed for full blockade of 1µg MICA-mFc binding.

Figure 8 shows that NKG2D-binding of ON72 to NKG2D was completely prevented by pre-incubation with hybridoma supernatant containing 16F16. Outline represents background, gray represents ON72-binding without pre-incubation, and black dotted represents ON72-binding with pre-incubation.

Figure 9 shows the capability of fully human anti-hNKG2D antibodies to block the subsequent binding of murine anti-hNKG2D antibodies to NKG2D, or vice versa. (A) 16F16: Pre-incubation with recombinantly expressed and purified 16F16 (0.3 µg; IgG4 isotype) prevented ON72 (0.3 µg) from binding to NKG2D. Reversing the incubation order showed that pre-incubation with ON72 (0.3 µg) only prevented 8.5% of the binding of recombinantly expressed and purified 16F16 of the IgG4 isotype (0.3 µg), showing that some fraction of NKG2D remained available for binding by the fully human antibody, suggesting an at least partially different epitope to the one bound by ON72. Antibody 149810 only demonstrated approximately 50% cross-inhibition of recombinantly expressed and purified 16F16 (IgG4 isotype), when tested at 1:1 (0.3µg, 0.3µg) and at 3:1 (1µg, 0.3 µg) of antibody concentration, 149810 to 16F16, respectively, again likely showing differences in the binding epitope on NKG2D. The following incubation and detection combinations were used: detection of ON72 without (a) or with (b) 16F16 pre-incubation; detection of 16F16 without (c) or with (d) ON72 pre-incubation; detection of 149810 (0.3µg) without (e) or with (f) 16F16 pre-incubation (0.3µg); detection of 16F16 without (g) or with (h) 149810 pre-incubation (0.3µg); detection of 149810 (1µg) without (i) or with (j) 16F16 pre-incubation (0.3µg); detection of 16F16 (0.3µg) without (k) or with (l) 149810 pre-incubation (1µg): 16F16 detection (0.3µg). (B) MS: 0.1 µg murine anti-hNKG2D antibody was incubated with or without pre-incubation with 0.1 µg of MS antibody, followed by detection with anti-mouse antibody using flow cytometry. Incubation with 0,1 µg ON72, 1D11, or 149810 without pre-incubation with MS was normalized to 100%, and is shown in (a), (c), and (e), respectively. Incubation with 0.1 µg MS for 30 minutes followed by incubation and detection of ON72, 1D11, or 149810 is shown in (b), (d), and (f), respectively. Pre-incubation with MS inhibited 98%, 88%, and 96.5% of the NKG2D-binding of ON72, 1D11, and 149810, respectively, suggesting similar epitopes of at least some of the antibodies. (C) 21 F2: detection of ON72 binding with (a) or without (b) pre-incubation with 21 F2; detection of 1 D11 binding with (c) or without (d) pre-incubation with 21 F2; and detection of 149810 binding with (e) or without (f) 21 F2 (all antibodies at 0.1 µg).

Figure 10 shows staining of rhesus or cynomolgous (cyno) cells with ON72 and 16F16 antibody purified from original hybridoma. (A) cyno NK cells, (B) cyno CD8+ T cells, (C) rhesus NK cells, and (D) rhesus CD8+ T cells. The values presented are mean fluorescent intensity (MFI) of binding where the MFI of binding of secondary antibody alone has been subtracted. No binding to CD4+ T cells was observed in either species.

Figure 11 shows the binding of human antibody MS to human or cynomolgous CD8-positive cells in peripheral blood mononuclear cells (PBMCs) at different antibody concentrations, demonstrating that the affinity to human and cynomolgous NKG2D is similar.

Figure 12 shows that addition of ligand-blocking antibodies, (ON72 or recombinantly expressed and purified 16F16 (IgG4 isotype)), blocked NK-mediated killing of MICA-expressing target cells in a dose-dependent fashion in a ⁵¹Cr-release assay.

Figure 13 shows that recombinantly expressed and purified 16F16 and 16F31 (both IgG4 isotype) were capable of inhibiting killing of both MICA (A) and ULBP3 (B) bearing target cells (BaF/3) by NK-92 cells in a dose dependent manner, with near total blockade by 16F16 at 0.8 µg/ml for both ligands, and partial blockade by 16F31 at the highest tested dose of 20 µg/ml for both ligands.

Figure 14 shows that recombinantly expressed and purified MS, 21 F2, and 16F16 (all IgG4 isotype) were capable of inhibiting NK-mediated killing of ligand-expressing target cells. (A) inhibition of NK-92 cells killing of ULBP3-BaF/3 cells by MS or 21 F2. (B) inhibition of NKL cells killing BaF/3-MICA cells by MS or 16F16.

Figure 15 shows antibody-induced reduction of cell-surface NKG2D, using BaF/3 cells transfected with NKG2D and DAP10. The

figure shows the percentage of NKG2D receptor that remained on the surface of the cells after overnight incubation with ON72 (1 µg) or recombinantly expressed and purified human antibodies 16F16 (1 µg; IgG4 isotype) or 16F31 (3 µg; IgG4 isotype), as compared to control (cell surface NKG2D receptor after overnight incubation without anti-NKG2D antibody =100%).

Figure 16 shows MS antibody-induced reduction of cell-surface NKG2D, using BaF/3 cells transfected with NKG2D and DAP10 (performed as for figure 15) (A), or freshly prepared human NK cells from peripheral blood (B). In (B), the human NK cells were incubated overnight in the presence of human serum, to mimic a situation in blood with IgGs present, and varying concentrations of MS antibody. Maximum downmodulation was achieved even at the lowest concentration, corresponding to about 60% receptor saturation measured in binding assay under similar conditions on NKG2D+ NK cells.

Figure 17 shows the percentage reduction of cell-surface NKG2D on human NK cells after over-night incubation with indicated 21 F2 antibody concentrations.

Figure 18 shows the effect of ON72, MS, and 21 F2 on surface-presented NKG2D in different types of cells in human blood samples, at the indicated time points. The concentration of each antibody was 0.1 µg/ml. While not being limited to theory, the reduction of surface-presented NKG2D in the experiments likely represents NKG2D internalization. Figures (A) to (C) shows antibody-induced NKG2D internalization of NKG2D-expressing NK cells, CD8+ T cells, and δγT cells, respectively. MS and 21 F2 resulted in less reduction of surface-associated NKG2D than ON72.

Figure 19 shows the results of an assay testing for an agonistic effect of immobilized MS and ON72 on T cell proliferation, using 2 different sub-optimal doses of CD3 to allow for co-stimulation. (A) [CD3] = 0.1 ng/ml; (B) [CD3] = 0.3 ng/ml. T cell proliferation was assessed by CFSE dilution in a PBMC population stimulated with immobilized antibody as indicated for 3 days followed by IL-2 stimulation for four days. CD28 stimulation is included as a positive control of co-stimulation. No significant agonistic effect could be detected for MS, whereas ON72 had a low but significant effect on T cell proliferation.

Figure 20 depicts 3-dimensional superimposed representations of hNKG2D dimer complexed with Fab-fragment(s) of anti-NKG2D antibody (MS or hzON72) or with MICA ligand. The hNKG2D homodimer ('NKG2D') is shown in a surface representation with one of the monomers in a darker color than the other. The Fab fragments ('MS' and 'hzON72', respectively) are indicated in black tube style while the MICA ('MICA') is indicated in a light schematic secondary structure representation style. (A), (B) Superpositioning of the hNKG2D/MS Fab and hNKG2D/MICA crystal complex structures (Li et al, Nat Immunol 2001;2:443-451; PDB-code 1HYR, using the C-alpha-atoms of the common hNKG2D molecule as template). As both MICA and MS bind to the NKG2D dimer in an asymmetric manner, (A) and (B) show the two possible relative binding orientations of the two ligand molecules when bound to the NKG2D-dimer. There was a considerable overlap between the MICA and the MS Fab in superimposition calculations to hNKG2D for both orientations, demonstrating the ability of the MS Fab to block MICA from binding to the hNKG2D receptor. See Example 11. (C) Superpositioning of the hNKG2D/hzON72 Fab and the hNKG2D/MICA complex crystal structures. Each monomer of hNKG2D was bound by an hzON72 Fab. In a superimposition calculation to hNKG2D, also the hzON72 antibody made a considerable overlap to the MICA binding site, showing that hzON72 can block MICA binding to hNKG2D. See Example 12.

Figure 21 shows the epitope residues in the sequence (SEQ ID NO:2) of each NKG2D monomer unit of a hNKG2D dimer for MS Fab (A), hzON72 Fab (B) and a MICA molecule (C) in the sequences (SEQ ID NO:2) of the two hNKG2D monomer units. NKG2D residues within 4.0 Å distance from the crystal structure ligand atoms were considered to be part of the binding epitope and are underlined. Doubly underlined residues were involved in hydrogen-binding to the ligand. (A) Binding epitope for a single MS Fab on hNKG2D monomer units 1 and 2 in a hNKG2D dimer. Crystallographic monomers N and C were combined in the NKG2D monomer unit 1, and crystallographic monomers M and D were combined in the NKG2D monomer unit 2. In monomer unit 2, the Lys 150 side chain atom N ϵ was only involved in hydrogen-binding in one of the two crystallographically independent complexes. See also Tables 9-12. (B) Respective binding epitopes for 2 hzON72 Fabs simultaneously bound to hNKG2D monomer units 1 and 2. Trp 166 was involved in hydrogen-bonding in one of the crystallographically independent molecular complexes (one hzON72 Fab molecule in complex with one hNKG2D monomer) but the distance was too far for hydrogen-binding in the other. See Tables 14-15. (C) Binding epitope for a MICA molecule on hNKG2D monomer units 1 and 2. MICA showed an asymmetric binding to the hNKG2D dimer, and could therefore bind in two orientations relative to MS-Fab. The second orientation of MICA can be obtained simply by the interchange of the 2 monomer unit representations.

Figure 22 shows the hNKG2D molecules in surface representations with one of the monomers slightly darker than the other. The NKG2D atoms within 4.0 Å distance from their respective crystals structures MS/hzON72/MICA Fab atoms are colored in black and are shown for the MS Fab (A), 2 hzON72 Fabs (B) and, for MICA, (C) and (D). As both MICA and MS Fab bind to the NKG2D dimer in an asymmetric manner, the relative binding orientation to NKG2D can differ. This is indicated in the figure which shows the two possible relative binding orientations of MICA in (C) and (D) to MS. See also Tables 9-12 and 14-15.

DEFINITIONS

[0018] As used herein, "hNKG2D" and, unless otherwise stated or contradicted by context, the terms "NKG2D," also known as "NKG2-D," "CD314," "D12S2489E," "KLRK1," "killer cell lectin-like receptor subfamily K, member 1," and "KLRK1," refer to a human killer cell activating receptor gene, its mRNA (e.g., NCBI RefSeq NM_007360; SEQ ID NO:1), and its gene product (NCBI RefSeq NP_031386; SEQ ID NO:2), or naturally occurring variants thereof. In NK and T cells, the ligand-binding form of the hNKG2D receptor is a homodimer (Li et al, Nat Immunol 2001;2:443-451). The hNKG2D receptor is typically presented at the surface in complex with DAP10 (Wu et al, J Exp Med 2000;192:1059 et seq.; NCBI Accession No. AAG29425, AAD50293) and has been suggested to also form higher order complexes. Any activity attributed herein to hNKG2D, e.g., cell activation, antibody recognition, etc., can also be attributed to hNKG2D in the form of a complex or higher-order complexes with DAP10, and/or other components.

[0019] The term "antibody" herein is used in the broadest sense and specifically includes full-length monoclonal antibodies, polyclonal antibodies, and, unless otherwise stated or contradicted by context, antigen-binding fragments, antibody variants, and multispecific molecules thereof, so long as they exhibit the desired biological activity. Generally, a full-length antibody is a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen binding portion thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarily determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. General principles of antibody molecule structure and various techniques relevant to the production of antibodies are provided in, e.g., Harlow and Lane, ANTIBODIES: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1988).

[0020] An "antigen-binding fragment" of an antibody is a molecule that comprises a portion of a full-length antibody which is capable of detectably binding to the antigen, typically comprising one or more portions of at least the VH region. Antigen-binding fragments include multivalent molecules comprising one, two, three, or more antigen-binding portions of an antibody, and single-chain constructs wherein the VL and VH regions, or selected portions thereof, are joined by synthetic linkers or by recombinant methods to form a functional, antigen-binding molecule. While some antigen-binding fragments of an antibody can be obtained by actual fragmentation of a larger antibody molecule (e.g., enzymatic cleavage), most are typically produced by recombinant techniques.

[0021] The terms "antibody derivative" and "immunoconjugate" are used interchangeably herein to denote molecules comprising a full-length antibody or an antigen-binding fragment thereof, wherein one or more amino acids are chemically modified, e.g., by alkylation, PEGylation, acylation, ester formation or amide formation or the like, e.g., for linking the antibody to a second molecule. Exemplary modifications include PEGylation (e.g., cysteine-PEGylation), biotinylation, radiolabelling, and conjugation with a second agent (such as a cytotoxic agent),

[0022] A "multispecific molecule" comprises an antibody, or an antigen-binding fragment thereof, which is associated with or linked to at least one other functional molecule (e.g. another peptide or protein such as another antibody or ligand for a receptor) thereby forming a molecule that binds to at least two different binding sites or target molecules. Exemplary multispecific molecules include bi-specific antibodies and antibodies linked to soluble receptor fragments or ligands.

[0023] The term "human antibody", as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from (i.e., are identical or essentially identical to) human germline immunoglobulin sequences. Furthermore, if the antibody contains a constant region, the constant region also is "derived from" human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*). However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[0024] A "humanized" antibody is a human/non-human chimeric antibody that contains a minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which

residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit, or non-human primate having the desired specificity, affinity, and capacity. In some instances, FR residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FR residues are those of a human immunoglobulin sequence. The humanized antibody can optionally also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see, e.g., Jones et al., *Nature* 321:522-525 (1986); Riechmann et al., *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992), WO 92/02190, US Patent Application 20060073137, and US Patents 6750325, 6632927, 6639055, 6548640, 6407213, 6180370, 6054297, 5929212, 5895205, 5886152, 5877293, 5869619, 5821337, 5821123, 5770196, 5777085, 5766886, 5714350, 5693762, 5693761, 5530101, 5585089, and 5225539.

[0025] The term "hypervariable region" when used herein refers to the amino acid residues of an antibody that are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a "complementarity-determining region" or "CDR" (residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light-chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy-chain variable domain; (Kabat et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and/or those residues from a "hypervariable loop" (residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light-chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy-chain variable domain; Chothia and Lesk, *J. Mol. Biol.* 1987;196:901-917). Typically, the numbering of amino acid residues in this region is performed by the method described in Kabat *et al.*, *supra*. Phrases such as "Kabat position", "variable domain residue numbering as in Kabat" and "according to Kabat" herein refer to this numbering system for heavy chain variable domains or light chain variable domains. Using the Kabat numbering system, the actual linear amino acid sequence of a peptide may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or CDR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of CDR H2 and inserted residues (e.g. residues 82a, 82b, and 82c, etc. according to Kabat) after heavy chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a "standard" Kabat numbered sequence.

[0026] "Framework region" or "FR" residues are those VH or VL residues other than the CDRs as herein defined.

[0027] An "epitope" or "binding site" is an area or region on an antigen to which an antigen-binding peptide (such as an antibody) specifically binds. A protein epitope may comprise amino acid residues directly involved in the binding (also called the immunodominant component of the epitope) and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically antigen binding peptide (in other words, the amino acid residue is within the "solvent-excluded surface" and/or "footprint" of the specifically antigen binding peptide). The term epitope herein includes both types of amino acid binding sites in any particular region of a hNKG2D that specifically binds to an anti-hNKG2D antibody, or another hNKG2D-specific agent according to the invention, unless otherwise stated (e.g., in some contexts the invention relates to antibodies that bind directly to particular amino acid residues). NKG2Ds may comprise a number of different epitopes, which may include, without limitation, (1) linear peptide antigenic determinants, (2) conformational antigenic determinants which consist of one or more non-contiguous amino acids located near each other in a mature NKG2D conformation; and (3) post-translational antigenic determinants which consist, either in whole or part, of molecular structures covalently attached to a NKG2D, such as carbohydrate groups. Unless otherwise specified or contradicted by context, conformational antigenic determinants comprise NKG2D amino acid residues within about 4 Å distance from an atom of an antigen-binding peptide.

[0028] The "solvent excluded surface" is the area of a molecule which, in a computer calculation, cannot be reached by any water molecule, e.g., because of binding of the molecule to a ligand (Lee and Richards, *J Mol Biol* 1971;55:379-400).

[0029] The phrase "binds to essentially the same epitope or determinant as" an antibody of interest (e.g., MS or 21 F2) means that an antibody "competes" with the antibody of interest for NKG2D molecules to which the antibody of interest specifically binds.

[0030] A "paratope" is an area or region of an antigen-binding portion of an antibody that specifically binds an antigen. Unless otherwise stated or clearly contradicted by context, a paratope may comprise amino acid residues directly involved in epitope binding, several of which are typically in CDRs, and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the specifically bound antigen (in other words, the amino acid residue is within the "solvent-excluded surface" and/or "footprint" of the specifically bound antigen).

[0031] The ability of an anti-NKG2D antibody to "block" the binding of a NKG2D molecule to a natural NKG2D-ligand (*e.g.*, MICA), means that the antibody, in an assay using soluble or cell-surface associated NKG2D and ligand molecules, can detectably reduce the binding of a NKG2D-molecule to the ligand in a dose-dependent fashion, where the NKG2D molecule detectably binds to the ligand in the absence of the antibody. An exemplary assay for determining whether an anti-NKG2D antibody is capable of blocking MICA-binding is provided in Example 3. The same assay can be used for testing antibody-mediated blocking of other NKG2D ligands.

[0032] A "variant" of a polypeptide refers to a polypeptide having an amino acid sequence that is substantially identical to a reference polypeptide, typically a native or "parent" polypeptide. The polypeptide variant may possess one or more amino acid substitutions, deletions, and/or insertions at certain positions within the native amino acid sequence and/or additions at one or both termini.

[0033] The term "substantially identical" in the context of two amino acid sequences means that the sequences, when optimally aligned, such as by the programs GAP or BEST-FIT using default gap weights, share at least about 50 percent sequence identity. Typically sequences that are substantially identical will exhibit at least about 60, at least about 70, at least about 80, at least about 90, at least about 95, at least about 98, or at least about 99 percent sequence identity.

[0034] "Corresponding" amino acid positions in two substantially identical amino acid sequences are those aligned by any of the protein analysis software referred to herein.

[0035] A nucleic acid sequence (or element) is "operably linked" to another nucleic acid sequence (or element) when it is placed into a functional relationship with the other nucleic acid sequence. For example, DNA for a pre-sequence or secretory leader is operably linked to DNA for (*i.e.*, coding for expression of) a polypeptide if it is expressed as a pre-protein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome-binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, some elements, such as enhancers, do not have to be contiguous with a coding sequence in order to be operably linked. Linking typically is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers may be used in accordance with conventional practice.

[0036] An "isolated" molecule is a molecule that is the predominant species in the composition wherein it is found with respect to the class of molecules to which it belongs (*i.e.*, it makes up at least about 50% of the type of molecule in the composition and typically will make up at least about 70%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or more of the species of molecule, *e.g.*, peptide, in the composition). Commonly, a composition of an antibody molecule will exhibit 98%, 98%, or 99% homogeneity for antibody molecules in the context of all present peptide species in the composition or at least with respect to substantially active peptide species in the context of proposed use.

[0037] In the context of the present invention, "treatment" or "treating" refers to preventing, alleviating, managing, curing or reducing one or more symptoms or clinically relevant manifestations of a disease or disorder, unless contradicted by context. For example, "treatment" of a patient in whom no symptoms or clinically relevant manifestations of a disease or disorder have been identified is preventive or prophylactic therapy, whereas clinical, curative, or palliative "treatment" of a patient in whom symptoms or clinically relevant manifestations of a disease or disorder have been identified generally does not constitute preventive or prophylactic therapy. Each form of treatment may be considered a distinct aspect of the invention.

DESCRIPTION OF THE INVENTION

[0038] The present invention is based, in part, on anti-NKG2D antibodies with properties suitable for treating human patients suffering from NKG2D-related conditions, such as, *e.g.*, autoimmune and inflammatory diseases and disorders. Antibodies of the invention are fully human or humanized in order to minimize the risk for an immune response against the antibody by the patient's own immune system, and bind to hNKG2D in its active form, *i.e.*, a homodimer on the surface of a cell and associated with DAP10.

[0039] The antibodies of the invention are typically useful for treatment of conditions where NKG2D activity should be reduced. Such antibodies can reduce or inhibit activation of NKG2D-expressing NK and/or T cells by, *e.g.*, competing with or blocking one or more endogenous NKG2D-ligands for binding to NKG2D, down-modulating or otherwise reducing the amount of cell-surface NKG2D upon binding, and/or eliciting an ADCC or CDC response against the cells.

[0040] In one aspect, antibodies of the invention are antagonists and compete with one or more natural ligands such as MICA for binding to human NKG2D, thereby reducing ligand-induced NKG2D-activation. MICA molecules have been clearly implicated in inflammatory diseases, and, as shown in Example 3, several human antibodies were effective at blocking MICA-binding to cell-surface NKG2D, particularly MS and 21 F2 (reference), and epitope determination showed that MS Fab obstructed MICA from binding (Example 11, Figure 20). Both MS and 21F2 (reference) were also highly efficient in blocking NK-cell mediated cytotoxicity (Example 6). Thus, these results demonstrate that the invention provides antibodies having such properties.

[0041] In a more particular aspect, antibodies of the invention are efficient antagonists, but also have insignificant agonistic effect on hNKG2D signalling, thus not contributing to NKG2D-driven inflammation. For example, as shown in Figure 19, no co-stimulation of immobilized MS on CD3-triggered proliferation of PBMCs could be detected, whereas immobilized ON72 resulted in a small but significant co-stimulation. Without being limited to theory, this difference may at least in part be due to the differences in epitopes, shown in Figures 20-22. An antigen-binding portion of bivalent MS antibody binds strongly to one monomer in an hNKG2D dimer complex, but blocks binding of a second MS antibody (or a second antigen-binding portion of the same antibody) to the second monomer. By contrast, when an antigen-binding portion of a bivalent hzON72 antibody binds a first monomer in an hNKG2D dimer, it does not block the binding of a second hzON72 antibody (or a second antigen-binding portion of the same antibody) to the second monomer.

[0042] In one embodiment, the invention provides human or humanized anti-NKG2D anti-bodies as defined in the claims which, when added to NKG2D-expressing NK or T cells, cross-link not more than 2 hNKG2D dimers. Preferably, such antibodies are bivalent. A bivalent antibody (such as, e.g., MS) for which the binding of the antigen-binding portion to an NKG2D monomer unit blocks further binding to the second NKG2D monomer unit can at most crosslink 2 hNKG2D dimers only. By contrast, a bivalent antibody which can bind an NKG2D monomer unit in an hNKG2D dimer without blocking binding to the second NKG2D monomer unit in an hNKG2D dimer can result in cross-linking of any number of hNKG2D dimers. Clustering of surface receptors commonly occurs in receptor activation.

[0043] In one embodiment, the invention provides human or humanized anti-NKG2D anti-bodies as defined in the claims which, when added to NKG2D-expressing NK or T cells, bind strongly only to one monomer in an hNKG2D dimer complex. Without being limited to theory, strong binding to both monomers of the dimer can be a prerequisite for activation of the NKG2D-receptor. MICA and hzON72 bind strongly to both monomer units in an hNKG2D dimer. MS binding to hNKG2D dimer is, however, dominated by binding to one of the monomer units while binding to the second monomer unit is weak and unspecific, and with a smaller solvent-excluded surface area on the second hNKG2D monomer (Example 11). In separate and specific preferred embodiments, the ratio of the solvent-excluded surface areas from the first and second NKG2D monomer units by the binding of an antibody of the invention is more than about 1:1, at least about 2:1, or at least about 3:1.

[0044] In one embodiment, the invention provides human or humanized anti-NKG2D anti-bodies as defined in the claims which bind essentially the same epitope as MS. Without being limited to theory, interactions of a ligand with particular residues, or residue combinations, on the hNKG2D dimer could avoid or minimize agonist activity. In separate and specific embodiments, the epitope of an antibody of the invention as defined in the claims comprises at least one residue selected from, at least 3 residues selected from, at least 5 residues selected from, at least 8 residues selected from, at least 10 residues selected from, at least 12 residues selected from, or all of the residues selected from the group consisting of Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gin 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2).

[0045] In one aspect, the present invention provides a fully human antibody, or antigen-binding fragment thereof, as defined in the claims, that effectively prevents NKG2D-mediated cytotoxicity of a hNKG2D-expressing NK or T cell, competes with at least MICA in binding to hNKG2D; reduces the amount of cell-surface hNKG2D upon binding via, e.g., stimulating down-modulation of hNKG2D, internalization of hNKG2D and/or preventing reappearance of hNKG2D; has an affinity to hNKG2D of 10 nM or less, cross-reacts with cynomolgus and/or rhesus NKG2D; and is non-depleting, e.g., by having an IgG4 isotype. In a particular embodiment, the antibody is a non-depleting fully human antibody of the IgG4 isotype, with an affinity to hNKG2D of 1 nM or less, preferably 300 pM or less, which blocks at least 50%, at least 70%, or at least 90% of endogenous hNKG2D-ligand binding, and reduces the amount of cell-surface hNKG2D with at least 10%, at least 30%, or at least 50%. In another particular embodiment, the antibody is a bivalent non-depleting fully human antibody of the IgG4 isotype, with an affinity below 100 pM, which has an EC50 concentration below 0.01 ng/ml for blocking the binding of full saturation dose of MICA-Fc to cell-surface associated NKG2D, is capable of reducing the amount of cell-surface NKG2D with at least 75% upon binding, and, optionally, has an EC50 concentration for reducing a ligand-induced NK cell cytotoxicity that is lower than the EC50 concentration required for binding to cell-surface associated NKG2D. The antibody may further be capable of achieving, in an assay using NKG2D-expressing cells, its maximum level of hNKG2D down-modulation at a concentration lower than that required to obtain saturation of the hNKG2D receptors (*i.e.*, saturation dose).

[0046] The production, characterization, and use of antibodies specifically binding hNKG2D and having some or all of these properties are described in more detail in the following sections, including the Examples.

Anti-NKG2D antibodies

[0047] The antibodies of the invention are characterized by particular functional and/or structural features or properties. Assays to evaluate the functional activities of anti-hNKG2D antibodies are described in detail in the Examples, and structural properties such as, e.g., amino acid sequences, are described below.

Functional properties

[0048] The antibodies of the invention bind to hNKG2D. In one embodiment, an antibody of the invention binds to hNKG2D with high affinity, for example with a KD of 10^{-7} M or less, a KD of 10^{-8} M or less, a KD of 1 nM or less, a KD of 0.3 nM or less, a KD of 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, or 0.01 nM or less. In a particular embodiment, the antibody binds to hNKG2D with an affinity of 0.1 nM or less.

[0049] In one aspect, the invention provides antibodies, as defined in the claims, also binding to one or more NKG2D orthologs in monkey such as cynomolgous monkey (*Macaca fascicularis*, NCBI accession No. AJ426429) and rhesus monkey (*Macaca mulatta*, NCBI accession No. AJ554302), and/or to hNKG2D homodimer, correctly folded monomeric full-length hNKG2D, hNKG2D fragment comprising an extracellular portion of hNKG2D, denatured hNKG2D, or to any combination of the preceding NKG2D forms. For example, as demonstrated in Example 5, the binding of human antibodies 21F2 (reference) and MS to specific cynomolgous cell types were more than about 65% and about 75%, respectively, of their binding to the same human cell types, per the corresponding EC50 (*i.e.*, the half maximal effective concentration) values. Accordingly, in one embodiment, an antibody of the invention binds to cynomolgous and/or rhesus NKG2D with similar affinity or efficacy as it binds to hNKG2D. For example, an antibody can bind to NKG2D-expressing cynomolgous or rhesus NK or T cells with an EC50 of about 50% or more, about 65% or more, or about 75% or more, of the corresponding EC50 for a corresponding population of NKG2D-expressing human NK or T cells. Additionally or alternatively, an antibody can bind to cynomolgous or rhesus NKG2D with an affinity of about 30% or more, about 50% or more, about 65% or more, or about 75% or more, about 80% or more, about 85% or more, or about 90% or more, of the affinity for hNKG2D. Such antibodies have the advantage of allowing for toxicity testing in the most suitable animal model (or models) prior to use in humans.

[0050] In one particular aspect, antibodies of the invention also bind a form of NKG2D that known murine anti-hNKG2D antibodies such as ON72 do not bind. Specifically, as described in Example 3, pre-incubation with ON72 only blocked about 82% of subsequently added human 16F16 antibody (reference) from binding to hNKG2D, while pre-incubation with 16F16 blocked about 95% of subsequently added ON72 from binding to hNKG2D.

[0051] Furthermore, the antibodies of the invention can reduce or inhibit hNKG2D-mediated activation of NK or T cells, *i.e.*, antagonize the hNKG2D receptor. This may be tested in, e.g., one or more cytotoxicity assays described herein or known in the art. For example, an antibody inhibits hNKG2D-mediated activation of an NK or T cell if it inhibits the NK- or T cell-mediated killing of an NKG2D-ligand-expressing target cell by at least 10%, more preferably by at least 30%, even more preferably by at least 40%, at least 50%, at least 60%, at least 70%, at least 80% or at least 90%, as compared to target cell killing in the absence of any anti-hNKG2D antibody or in the presence of a non-specific, control antibody.

[0052] Antibodies of the invention that are hNKG2D antagonists can have no or low agonist activity. Such antibodies are human or humanized. Agonist activity may be tested in one of the assay described herein, or an assay known in the art. For example, one type of assay is a co-stimulation assay measuring proliferation of peripheral blood lymphocytes (PBMcs) stimulated with low levels of CD3 in the presence or absence of immobilized anti-NKG2D antibody (see Example 10). In such an assay, proliferation in the presence of an antibody of the invention is not more than 30%, not more than 20%, not more than 10%, not more than 5% or not significantly higher than in the absence of antibody. Preferably, proliferation in the presence of an antibody of the invention is not significantly higher than in the absence of antibody. In an additional or alternative embodiment, hNKG2D agonist activity of an antibody of the invention in an agonist assay is not more than 30%, not more than 20%, not more than 10%, not more than 5%, or not significantly higher than a control value. The control is preferably a negative control, such as, e.g., in the absence of antibody, in the absence of cell or another reagent, and/or in the presence of an irrelevant antibody. Preferably, agonist activity of an antibody of the invention is not significantly higher than a control value.

[0053] In another aspect, the invention provides antibodies as defined in the claims that have a lower, preferably substantially lower, EC50 concentration for blocking ligand-induced cytotoxicity than for binding to cell-surface NKG2D of an NK or T cell. For example, for ON72, the EC50 concentration for binding to cell-surface NKG2D expressed on BaF/3 cells (0.062 µg/ml) was similar to the EC50 concentration for blocking NK-cell mediated killing of ligand- (ULBP3-) expressing - target cells (0.065 µg/ml), whereas 21F2 (reference) had a lower, and MS a substantially lower, EC50 for blocking cytotoxicity (21 F2: 0.021 µg/ml; MS: 0.012 µg/ml) than for binding to cell-surface NKG2D (21 F2: 0.033 µg/ml; MS: 0.032 g/ml) (see Examples 6 and 9). Further, MS achieved maximum blocking of cytotoxicity at lower concentrations (a concentration corresponding only to about 80% saturation of cell-associated NKG2D-receptors, Figure 3) than 21F2 (reference) and 16F16 (reference) (which had concentrations corresponding to saturating concentrations or higher, Figure 3). Thus, in one embodiment, the invention provides human or humanized antibodies, as defined in the claims, that have a lower EC50 concentration for blocking ligand-induced cytotoxicity than for binding to cell-surface NKG2D of an NK or T cell. The EC50 for blocking cytotoxicity of NK or T cells of a cell line or other suitable preparation can be, e.g., about 95% or less, about 90% or less, about 85% or less, about 80% or less, about 70% or less, about 50% or less, or about 40% or less, of the EC50 for binding to cell-surface NKG2D of the same cell line or preparation. Exemplary cell lines for testing include NK-92 and NKL cells.

[0054] In another embodiment, the invention provides antibodies as defined in the claims, that achieve maximum blockage of NK cell cytotoxicity at a concentration lower than the concentration required to saturate the available hNKG2D-receptors. In a specific embodiment, the antibodies also compete with MS in binding to hNKG2D. In another specific embodiment, such antibodies bind to essentially the same hNKG2D epitope as MS.

[0055] The antibodies may reduce or inhibit NKG2D-mediated activation by, e.g., interfering with the hNKG2D-binding of one or more endogenous hNKG2D-ligands. For example, the antibodies may reduce or inhibit the hNKG2D-binding of MICA; MICB; ULBP1; ULBP2; ULBP4; and/or RAET1-family member; e.g., by reducing or inhibiting the hNKG2D-binding of MICA; or of MICA and MICB; or of MICA and ULBP3; or of MICA, MICB, and ULBP3; or of MICA, MICB, and all ULBP1, -2, -3, and 4; or of MICA, MICB, and one or more RAET1 family members. The ability of an antibody to inhibit hNKG2D-binding of endogenous NKG2D-ligands can be evaluated using binding or competition assays described herein. In one embodiment, antibodies of the invention are capable of inhibiting at least 30% of ligand binding, or at least 50% of ligand binding, or at least 70% of ligand binding, or at least 80%, or at least 90% of ligand binding. In another embodiment, the IC50 for an antibody of the invention to inhibit the hNKG2D-binding of 1 µg MICA-mFc is 1 nM or less, 0.5 nM or less, 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, or 0.02 nM or less, 0.01 nM or less, 0.005 or less, or 0.002 or less. In another embodiment, full blockage of 1 µg MICA-mFc binding is achieved at an antibody concentration of 5 nM or less, 1 nM or less, 0.7 nM or less, 0.5 nM or less, or 0.2 nM or less, 0.1 nM or less, 0.05 nM or less, or about 0.02 nM or less. In one embodiment, the invention provides antibodies, especially human antibodies, that are as efficient or more efficient in reducing or inhibiting ligand hNKG2D-binding, such as, e.g., MICA binding to hNKG2D, than any of ON72, BAT221, 5C6, 1D11, ECM217, and 149810.

[0056] Additionally or alternatively, an anti-hNKG2D antibody of the invention can be capable of reducing the amount of cell-surface hNKG2D upon (*i.e.*, following) binding. Reduction of cell-surface associated hNKG2D upon binding of an antibody can be an advantageous feature, since it reduces the number of hNKG2D receptors available for ligand binding and subsequent activation. Without being limited to theory, this reduction may be caused by NKG2D down-modulation, internalization, or other mechanism. As shown herein, anti-hNKG2D antibodies having a human Fc-region, such as human antibodies, are capable of effectively reducing the amount of cell-surface hNKG2D. For example, human anti-hNKG2D antibodies 16F16 (reference), MS, and 21F2 (reference) all reduced the amount of cell-surface hNKG2D with about 75% or more after overnight incubation in the absence of serum, with MS being the most effective, achieving 75-90% downmodulation at a low concentration (Figures 15-17). Also, in the presence of serum, an MS concentration corresponding to less than saturating concentration on hNKG2D-expressing BaF/3 cells achieved maximum downmodulation (Figure 16B). Accordingly, in one embodiment, the invention provides antibodies, as defined in the claims, binding to hNKG2D that are able to achieve maximum down-modulation of hNKG2D at less than saturating concentrations. In another embodiment, such antibodies also compete with MS in binding to hNKG2D. In another embodiment, such antibodies also bind to essentially the same hNKG2D epitope as MS. An antibody of the invention can be capable of reducing cell surface hNKG2D by at least 10%, at least 20%, at least 30%, at least 50%, at least 70%, or at least 90% as compared to cell-surface hNKG2D in the absence of anti-hNKG2D antibody or in the presence of a non-specific control antibody. Preferably, the antibodies achieve reduction of cell-surface NKG2D while causing no or minimal activation of NKG2D-receptor signalling, *i.e.*, with no or minimal agonist activity. Exemplary assays for evaluating cell surface hNKG2D and agonistic activity of anti-hNKG2D antibodies are described herein. In one embodiment, the invention provides antibodies, as defined in the claims, particularly human antibodies, which are capable of a higher degree of down-modulation than a control antibody selected from ON72, BAT221, 5C6, 1D11, ECM217, and 149810. In another embodiment, an anti-hNKG2D antibody of the invention can be capable of achieving maximum down-modulation of cell-surface NKG2D expressed by a cell or cell-line at a concentration lower

than a saturating concentration.

[0057] In another embodiment, the invention provides antibodies, as defined in the claims, compete with and/or bind to the same epitope on hNKG2D as MS. Such antibodies can be identified based on their ability to cross-compete with MS in standard hNKG2D binding assays as described herein. The ability of a test antibody to inhibit the binding of 16F16, 16F31, MS, or 21 F2 to hNKG2D demonstrates that the test antibody can compete with 16F16, 16F31, MS, or 21F2 for binding to hNKG2D and thus can bind to the same epitope on hNKG2D as 16F16, 16F31, MS, or 21F2. In a preferred embodiment, the antibody that binds to the same epitope on hNKG2D as MS is a human monoclonal antibody. Such human monoclonal antibodies can be prepared and isolated as described in the Examples.

[0058] In another preferred embodiment, the antibody binds to a different epitope than any of the mouse monoclonal antibodies ON72, BAT221, 5C6, 1D11, ECM217, and 149810, and cross-competes more with MS than with either of the listed mouse monoclonal antibodies.

[0059] In one embodiment, the epitope of an antibody of the invention comprises one or more residues selected from Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention comprises 5 or more residues selected from Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention comprises 8, 10, 12 or more residues selected from Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention comprises the residues Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention consists essentially of the residues Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention consists of one or more residues selected from Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention consists of the residues Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of hNKG2D (SEQ ID NO: 2).

[0060] In one embodiment, the epitope of an antibody of the invention comprises one or more residues involved in hydrogen-binding selected from Lys 150, Ser 151, Tyr 152, Ile 181, Met 184, Gln 185, Lys 197, Thr 205, and Asn 207 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention comprises 5 or more residues involved in hydrogen-binding selected from Lys 150, Ser 151, Tyr 152, Ile 181, Met 184, Gln 185, Lys 197, Thr 205, and Asn 207 of hNKG2D (SEQ ID NO: 2). In one embodiment, the epitope of an antibody of the invention comprises Lys 150, Ser 151, Tyr 152, Ile 181, Met 184, Gln 185, Lys 197, Thr 205, and Asn 207 of hNKG2D (SEQ ID NO: 2).

[0061] Preferred antibodies of the invention exhibit at least one, more preferably two, three, four, five or more, of the following properties: (a) prevents NKG2D-mediated activation of an NKG2D-expressing NK or T cell, optionally with an EC50 for reducing ligand-induced cytotoxicity lower than the EC50 for binding to the cell; (b) competes with at least one NKG2D ligand in binding to NKG2D, preferably with at least MICA and ULBP3; (c) reduces the amount of NKG2D on the surface of a NKG2D-expressing NK or T cell, preferably with at least 75%; (d) binds to cynomolgous and/or rhesus NKG2D, preferably with no less than 50% of the affinity by which it binds to hNKG2D; (e) binds to more than one form or conformation of NKG2D; (f) binds to NKG2D with a Kd of 1 nM or less, preferably 0.1 nM or less; (g) competes with MS in binding to hNKG2D, (h) competes more with MS than with any of ON72, BAT221, 5C6, 1D11, ECM217, and 149810 in binding to hNKG2D; (i) blocks more than 90% of 16F16, MS, or 21 F2 binding to cell-surface hNKG2D; (j) has insignificant agonist activity, and (k) binds to essentially the same epitope as MS. Any combination of the above-described functional features, and/or the functional features as described in the Examples, may be exhibited by an antibody of the invention.

Structural properties

[0062] A preferred antibody of the invention is the human monoclonal antibody MS produced, isolated, and structurally and functionally characterized as described in the Examples. Further human monoclonal antibodies disclosed herein are 16F16, 16F31 and 21F2. Full-length, variable, and CDR sequences of these antibodies are set forth in Table 1.

Table 1

Full-length, variable, and COR amino acid sequences for 16F164 (reference), 16F31 (reference), MS, and 21F2 (reference)			
Antibody portion	SEQ ID NO:	Antibody portion	SEQ ID NO:
16F16 IgG4 H chain	7	MS IgG4 H chain	40
16F16 L chain	8	MS L chain	41
16F31 IgG4 H chain	9	21F2 IgG4 H chain	42
16F16 L chain	10	21F2 L chain	43
16F16 VH region	11	MS VH region	44
16F16 VL region	12	MS VL region	45
16F31 VH region	13	21F2 VH region	46
16F31 VL region	14	21F2 VL region	47
16F16 VH CDR1	15	MS VH CDR1	48
16F16 VH CDR2	16	MS VH CDR2	49
16F16 VH CDR3	17	MS VH CDR3	50
16F16 VL CDR1	18	MS VL CDR1	51
16F16 VL CDR2	19	MS VL CDR2	52
16F16 VL CDR3	20	MS VL CDR3	53
16F31 VH CDR1	21	21F2 VH CDR1	54
16F31 VH CDR2	22	21 F2 VH CDR2	55
16F31 VH CDR3	23	21 F2 VH CDR3	56
16F31 VL CDR1	24	21F2 VL CDR1	57
16F31 VL CDR2	25	21 F2 VL CDR2	58
16F31 VL CDR3	26	21 F2 VL CDR3	59

[0063] Certain anti-NKG2D antibodies of the invention have the same or a similar paratope as MS. In one embodiment, the antibody has a paratope comprising residues corresponding to one or more of Tyr 33 and Trp 97 of the MS L chain (SEQ ID NO: 41), and/or to one or more of Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of the MS H chain (SEQ ID NO: 40). In one embodiment, the antibody has a paratope comprising residues corresponding to Tyr 33 and Trp 97 of the MS L chain (SEQ ID NO: 41), and/or to 3, 5, 7, 10 or more of Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of the MS H chain (SEQ ID NO: 40). In one embodiment, the antibody has a paratope comprising residues corresponding to Tyr 33 and Trp 97 of the MS L chain (SEQ ID NO: 41), and Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of the MS H chain (SEQ ID NO: 40). In one embodiment, the antibody has a paratope consisting essentially of residues corresponding to Tyr 33 and Trp 97 of the MS L chain (SEQ ID NO: 41), and Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of the MS H chain (SEQ ID NO: 40). In one embodiment, the antibody has a paratope consisting of residues corresponding to Tyr 33 and Trp 97 of the MS L chain (SEQ ID NO: 41), and Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of the MS H chain (SEQ ID NO: 40).

[0064] Given that all of 16F16, 16F31, 21F2, and MS can bind to hNKG2D, it may be possible to "mix and match" the respective VH and VL sequences of these antibodies to create other anti-hNKG2D binding molecules. The hNKG2D-binding of such "mixed and matched" antibodies can be tested using the binding assays described herein (e.g. flow cytometry, Biacore, ELISAs) and/or using a cytotoxicity assay as described herein. Preferably, when VH and VL chains are mixed and matched, a VH sequence from a particular VHNL pairing is replaced with a structurally similar VH sequence. Likewise, preferably a VL sequence from a particular VHNL pairing is replaced with a structurally similar VL sequence.

[0065] Accordingly, in one aspect, the present specification discloses an isolated monoclonal antibody, or antigen binding portion thereof, comprising: (a) a VH region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 11, 13, 44, and 46, and (b) a VL region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 12, 14, 45, and 47; wherein the antibody binds hNKG2D. Preferred heavy and light chain combinations include: (a) a VH

region comprising the amino acid sequence of SEQ ID NO: 11; and (b) a light chain variable region comprising the amino acid sequence of SEQ ID NO: 12; (a) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 13; and (b) a light chain variable region comprising the amino acid sequence of SEQ ID NO: 14; (a) a VH region comprising the amino acid sequence of SEQ ID NO: 44; and (b) a light chain variable region comprising the amino acid sequence of SEQ ID NO: 46; or (a) a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 45; and (b) a light chain variable region comprising the amino acid sequence of SEQ ID NO: 47.

[0066] In another aspect, the invention provides antibodies as defined in the claims that comprise the heavy chain and light chain CDR1s, CDR2s and CDR3s of MS. Also disclosed herein are other antibodies that comprise the heavy chain and light chain CDR1 s, CDR2s and/or CDR3s of 16F16, 16F31, MS, or 21 F2, or combinations thereof. The CDR regions are delineated using the Kabat system (Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). See, *e.g.*, Figures 4 and 5. Given that each of these antibodies can bind to hNKG2D and that antigen-binding specificity is provided primarily by the CDR 1, 2 and 3 regions, the VH CDR1, 2 and 3 sequences and VL CDR1, 2 and 3 sequences can be "mixed and matched" (*i.e.*, CDRs from different antibodies can be mixed and match, although each antibody can contain a VH CDR1, 2 and 3 and a VL CDR1, 2 and 3) to create other anti-hNKG2D binding molecules as disclosed herein. The hNKG2D-binding of such "mixed and matched" antibodies can be tested using the binding assays described above and in the Examples (*e.g.* flow cytometry, Biacore, or ELISAs). Preferably, when VH CDR sequences are mixed and matched, the CDR1, CDR2 and/or CDR3 sequence from a particular VH sequence is replaced with a structurally similar CDR sequence(s). Likewise, when VL CDR sequences are mixed and matched, the CDR1, CDR2 and/or CDR3 sequence from a particular VL sequence preferably is replaced with a structurally similar CDR sequence(s). For example, the VL CDR1s and CDR3s of 16F16, 16F31, MS, and 21F2 and the VL CDR2 sequences of MS and 21F2 share some structural similarity and therefore are amenable to mixing and matching. It will be readily apparent to the ordinarily skilled artisan that novel VH and VL sequences can be created by substituting one or more VH and/or VL CDR region sequences with structurally similar sequences from the CDR sequences disclosed herein for monoclonal antibodies 16F16, 16F31, MS, and 21F2.

[0067] Accordingly, in another aspect, described herein is an isolated monoclonal antibody, or antigen binding portion thereof comprising: (a) a VH CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 15, 21, 48, and 54; (b) a VH CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 16, 22, 49, and 55; (c) a VH CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 17, 23, 50, and 56; (d) a VL CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 24, 51, and 57; (e) a VL CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 19, 25, 52, and 57; and (f) a VL CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 20, 26, 53, and 59; wherein the antibody binds hNKG2D.

[0068] In one embodiment disclosed herein, the antibody comprises: (a) a VH CDR1 comprising SEQ ID NO:15; (b) a VH CDR2 comprising SEQ ID NO:16; (c) a VH CDR3 comprising SEQ ID NO:17; (d) a VL CDR1 comprising SEQ ID NO:18; (e) a VL CDR2 comprising SEQ ID NO: 19; and (f) a VL CDR3 comprising SEQ ID NO: 20.

[0069] In another embodiment disclosed herein, the antibody comprises: (a) a VH CDR1 comprising SEQ ID NO: 21; (b) a VH CDR2 comprising SEQ ID NO:22; (c) a VH CDR3 comprising SEQ ID NO:23; (d) a VL region CDR1 comprising SEQ ID NO:24; (e) a VL CDR2 comprising SEQ ID NO:25; and (f) a VL CDR3 comprising SEQ ID NO: 26.

[0070] As also defined in the claims, the antibody according to the invention comprises: (a) a VH CDR1 comprising SEQ ID NO: 48; (b) a VH CDR2 comprising SEQ ID NO:49; (c) a VH CDR3 comprising SEQ ID NO:50; (d) a VL region CDR1 comprising SEQ ID NO:51; (e) a VL CDR2 comprising SEQ ID NO:52; and (f) a VL CDR3 comprising SEQ ID NO: 53.

[0071] In another embodiment disclosed herein, the antibody comprises: (a) a VH CDR1 comprising SEQ ID NO: 54; (b) a VH CDR2 comprising SEQ ID NO:55; (c) a VH CDR3 comprising SEQ ID NO:56; (d) a VL region CDR1 comprising SEQ ID NO:57; (e) a VL CDR2 comprising SEQ ID NO:58; and (f) a VL CDR3 comprising SEQ ID NO: 59.

[0072] In another embodiment disclosed herein, the antibody comprises: (a) a VH CDR1 consisting of SEQ ID NO:15; (b) a VH CDR2 consisting of SEQ ID NO:16; (c) a VH CDR3 consisting of SEQ ID NO:17; (d) a VL CDR1 consisting of SEQ ID NO:18; (e) a VL CDR2 consisting of SEQ ID NO: 19; and (f) a VL CDR3 consisting of SEQ ID NO: 20.

[0073] In another embodiment disclosed herein, the antibody comprises: (a) a VH CDR1 consisting of SEQ ID NO: 21; (b) a VH CDR2 consisting of SEQ ID NO:22; (c) a VH CDR3 consisting of SEQ ID NO:23; (d) a VL region CDR1 consisting of SEQ ID NO:24; (e) a VL CDR2 consisting of SEQ ID NO:25; and (f) a VL CDR3 consisting of SEQ ID NO: 26.

[0074] In a preferred embodiment according to the invention, the antibody comprises: (a) a VH CDR1 consisting of SEQ ID NO: 48; (b) a VH CDR2 consisting of SEQ ID NO:49; (c) a VH CDR3 consisting of SEQ ID NO:50; (d) a VL region CDR1 consisting of SEQ ID NO:51; (e) a VL CDR2 consisting of SEQ ID NO:52; and (f) a VL CDR3 consisting of SEQ ID NO: 53.

[0075] In another embodiment according to the invention, the antibody comprises: (a) a VH CDR1 consisting of SEQ ID NO: 48; (b) a VH CDR2 consisting of SEQ ID NO:49; (c) a VH CDR3 consisting of SEQ ID NO:50; (d) a VL region CDR1 consisting of SEQ ID NO:51; (e) a VL CDR2 consisting of SEQ ID NO:52; and (f) a VL CDR3 consisting of SEQ ID NO: 53, and residues corresponding to one, two, or all of Gln 1, Asp 26, and Asp 27 in the MS H chain (SEQ ID NO: 40)

[0076] In another embodiment disclosed herein, the antibody comprises: (a) a VH CDR1 consisting of SEQ ID NO: 54; (b) a VH CDR2 consisting of SEQ ID NO:55; (c) a VH CDR3 consisting of SEQ ID NO:56; (d) a VL region CDR1 consisting of SEQ ID NO:57; (e) a VL CDR2 consisting of SEQ ID NO:58; and (f) a VL CDR3 consisting of SEQ ID NO: 59.

[0077] In certain embodiments, an antibody of the invention comprises a VH region from a particular germline H chain immunoglobulin gene, or a combination of particular germline H chain immunoglobulin genes; and/or a VL region from a particular germline L chain immunoglobulin gene, or a combination of particular germline L chain immunoglobulin genes. Such combinations can be obtained, *e.g.*, *in vivo* via somatic recombination in a B cell.

[0078] For example, in one embodiment, disclosed herein is an isolated anti-hNKG2D antibody, or an antigen-binding fragment thereof, wherein the antibody: (a) comprises a VH region from a human VH3_21, VH3_20, VH4_59, or VH5_51 gene recombined with a human D3-9, D3-10, or D3_10_R3 gene and a JH3, JH4 or JH6 gene, (b) comprises a VL region derived from a human VKI_L15 or VKIII_A27 or VKIII_L6 gene recombined with a human JK1, JK2 or JK3 gene, and (c) the antibody binds to hNKG2D.

[0079] In another embodiment, disclosed herein is an isolated anti-hNKG2D antibody, or an antigen-binding fragment thereof, comprising a VH region obtained by a recombination of human VH3_21, D3-9, and JH4 genes and a VL region obtained by a recombination of human VKI_L15 and JK2 genes.

[0080] In another embodiment, disclosed herein is an isolated anti-hNKG2D antibody, or an antigen-binding fragment thereof, comprising a VH region obtained by a recombination of human VH3_20, D3-10, and JH6 genes and a VL region obtained by a recombination of human VKIII_A27 and JK3 genes.

[0081] In another embodiment, the invention provides an isolated anti-hNKG2D antibody, or an antigen-binding fragment thereof, as defined in the claims, comprising a VH region obtained by a recombination of human VH4_59, a D gene, and JH3 genes and a VL region obtained by a recombination of human VKIII_A27 and JK1 genes.

[0082] In another embodiment, disclosed herein is an isolated anti-hNKG2D antibody, or an antigen-binding fragment thereof, comprising a VH region obtained by a recombination of human VH5_51, D3_10_R3, and JH4 genes and a VL region obtained by a recombination of human VKIII_L6 and JK1 genes.

[0083] In separate and specific embodiments, the invention provides isolated anti-NKG2D antibodies as defined in the claims, obtained by introducing one, two, three, four or more amino acid substitutions and/or somatic hypermutations in the VH and/or VL region of an anti-hNKG2D antibody described above.

[0084] As used herein, a human antibody comprises heavy or light chain variable regions "of" or "derived from" or that are "the product of" a particular germline sequence if the variable regions of the antibody are obtained from a system (as described below) that uses human germline immunoglobulin genes. Such "systems" include immunizing a transgenic mouse carrying human immunoglobulin genes with the antigen of interest or screening a human immunoglobulin gene library displayed on phage with the antigen of interest. A human antibody that is "of" or "derived from" or "the product of" a human germline immunoglobulin sequence can be identified as such by comparing the amino acid sequence of the human antibody to the amino acid sequences of human germline immunoglobulins and selecting the human germline immunoglobulin sequence that is closest in sequence (*i.e.*, greatest % identity) to the sequence of the human antibody. A human antibody that is "of" or "derived from" or "the product of" a particular human germline immunoglobulin sequence may contain amino acid differences as compared to the germline sequence, due to, for example, naturally-occurring somatic mutations or intentional introduction of site-directed mutation(s) (which may be selected substitutions).

[0085] However, a human antibody is typically at least 90% identical in amino acid sequence to an amino acid sequence encoded by a recombined germline immunoglobulin sequence and can usually be identified as human when compared to the

germline immunoglobulin amino acid sequences of other species (e.g., murine germline sequences). In certain cases, a human antibody may be at least 95%, or even at least 96%, 97%, 98%, or 99% identical in amino acid sequence to the amino acid sequence encoded by the recombined germline immunoglobulin gene.

[0086] Typically, a human antibody derived from a particular human germline sequence will display no more than 10 amino acid differences from the amino acid sequence encoded by the human germline immunoglobulin gene. In certain cases, the human antibody may display no more than 8, no more than 5, or even no more than 4, 3, 2, or 1 amino acid difference, or no amino acid difference, from the amino acid sequence encoded by the recombined germline immunoglobulin gene.

[0087] In yet another embodiment, an antibody of the invention, as defined in the claims, comprises heavy and light chain variable regions comprising amino acid sequences that are homologous to the amino acid sequences of the preferred antibodies described herein, and wherein the antibodies retain the desired functional properties of the anti-hNKG2D antibodies of the invention. For example, the invention provides an isolated antibody, or antigen binding portion thereof, as defined in the claims, comprising a heavy chain variable region and a light chain variable region, wherein: (a) the VH region comprises an amino acid sequence that is at least 80% identical to the amino acid sequence of SEQ ID NO: 44; (b) the VL region comprises an amino acid sequence that is at least 80% identical to the amino acid sequence of SEQ ID NO: 45; (c) the antibody binds to hNKG2D and exhibits at least one of the functional properties described herein, preferably several of the functional properties described herein.

[0088] In other embodiments, the VH and/or VL amino acid sequences may be 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth above. An antibody having VH and VL regions having high (*i.e.*, 80% or greater) identity to the VH and VL regions of the sequences set forth above, can be obtained by mutagenesis (e.g., site-directed or PCR-mediated mutagenesis) of nucleic acid molecules encoding SEQ ID NOs: 44-45, followed by testing of the encoded altered antibody for retained function (e.g., hNKG2D binding affinity, hNKG2D-ligand blocking, hNKG2D downmodulation, or reduction of NKG2D-mediated activation of an NK or T cell) using the functional assays described herein.

[0089] The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm in sequence-analysis software. Protein analysis software matches similar sequences using measures of similarity assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions.

[0090] The percent identity between two amino acid sequences can be determined, e.g., using the Needleman and Wunsch (J. Mol. Biol. 48:444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

[0091] Polypeptide sequences can also be compared using FASTA, applying default or recommended parameters. A program in GCG Version 6.1., FASTA (e.g., FASTA2 and FASTA3) provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, Methods Enzymol. 1990;183:63-98; Pearson, Methods Mol. Biol. 2000;132:185-219).

[0092] The percent identity between two amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (Comput. Appl. Biosci., 1988;11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0093] Another algorithm for comparing a sequence to a other sequences contained in a database is the computer program BLAST, especially blastp, using default parameters. See, e.g., Altschul et al., J. Mol. Biol. 1990;215:403-410; Altschul et al., Nucleic Acids Res. 1997;25:3389-402 (1997). The protein sequences of the present invention can there be used as a "query sequence" to perform a search against public databases to, for example, identify related sequences. Such searches can be performed using the XBLAST program (version 2.0) of Altschul, *et al.* 1990 (*supra*). BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the antibody molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, 1997 (*supra*).

When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Described herein is also an antibody that comprises a VH region comprising CDR1, CDR2 and CDR3 sequences and a VL region comprising CDR1, CDR2 and CDR3 sequences, wherein one or

more of these CDR sequences comprise specified amino acid sequences based on the preferred antibodies described herein; 16F16, 16F31, MS, or 21 F2, wherein one or more CDRs optionally contains one or more conservative amino acid modifications, and wherein the antibodies retain the desired functional properties of the anti-hNKG2D antibodies described herein. Accordingly, described is an isolated anti-body, or antigen-binding fragment thereof, comprising a heavy chain variable region comprising CDR1, CDR2, and CDR3 sequences and a light chain variable region comprising CDR1, CDR2, and CDR3 sequences, wherein: (a) the VH region CDR3 sequence comprises an amino acid sequence selected from the group consisting of amino acid sequences of SEQ ID NOs: 17, 23, 50, and 56; (b) the VL region CDR3 sequence comprises an amino acid sequence selected from the group consisting of amino acid sequences of SEQ ID NOs: 20, 26, 53 and 59; (c) one or more CDRs optionally contains one or more conservative amino acid modifications, and (d) the antibody binds to hNKG2D and exhibits at least one of the functional properties described herein, more preferably several of the functional properties described herein.

[0094] In a further embodiment disclosed herein, the VH region CDR2 sequence comprises an amino acid sequence selected from the group consisting of amino acid sequences of SEQ ID NOs: 16, 22, 49, and 55; and the VL region CDR2 sequence comprises an amino acid sequence selected from the group consisting of amino acid sequences of SEQ ID NOs: 19, 25, 52, and 58, wherein one or more CDRs optionally contains one or more conservative amino acid modifications.

[0095] In a still further embodiment disclosed herein, the VH region CDR1 sequence comprises an amino acid sequence selected from the group consisting of amino acid sequences of SEQ ID NOs: 15, 21, 48, and 54, and conservative modifications thereof; and the VL region CDR1 sequence comprises an amino acid sequence selected from the group consisting of amino acid sequences of SEQ ID NOs: 18, 24, 51, and 57, wherein one or more CDRs optionally contains one or more conservative amino acid modifications.

[0096] As used herein, the term "conservative amino acid modifications" is intended to refer to amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody containing the amino acid sequence. Such conservative modifications include amino acid substitutions, additions and deletions. Modifications can be introduced into an antibody of the invention by standard techniques known in the art, such as, e.g., site-directed mutagenesis and PCR-mediated mutagenesis. An antibody sequence comprising amino acid modifications as compared to a parent antibody is typically at least 90%, preferably at least 95%, 98%, or 99% identical to the corresponding amino acid sequence in the parent and/or comprises at most 10, preferably at most 5, 4, 3, 2 amino acid modifications as compared to the parent antibody sequence.

[0097] "Conservative" amino acid substitutions are typically those in which an amino acid residue is replaced with an amino acid residue having a side chain with similar physicochemical properties. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g. glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g. threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine).

[0098] Thus, one or more amino acid residues within the CDR regions of an antibody described herein can be replaced with other amino acid residues from the same side chain family and the altered antibody can be tested for retained function (*i.e.*, the functions set forth in (c), (d) and (e) above) using the functional assays described herein.

Antigen-binding fragments

[0099] The anti-hNKG2D antibodies of the invention may be prepared as full-length antibodies or antigen-binding fragments thereof. Examples of antigen-binding fragments include Fab, Fab', F(ab)2, F(ab')2, F(ab)3, Fv (typically the VL and VH domains of a single arm of an antibody), single-chain Fv (scFv; see *e.g.*, Bird et al., Science 1988;242:423-426; and Huston et al. PNAS 1988;85:5879-5883), dsFv, Fd (typically the VH and CH1 domain), and dAb (typically a VH domain) fragments; VH, VL, VhH, and V-NAR domains; monovalent molecules comprising a single VH and a single VL chain; minibodies, diabodies, triabodies, tetrabodies, and kappa bodies (see, *e.g.*, Ill et al., Protein Eng 1997;10:949-57); camel IgG; IgNAR; as well as one or more isolated CDRs or a functional paratope, where the isolated CDRs or antigen-binding residues or polypeptides can be associated or linked together so as to form a functional antibody fragment. Various types of antibody fragments have been described or reviewed in, *e.g.*, Holliger and Hudson, Nat Biotechnol 2005;23:1126-1136; WO2005040219, and published U.S. Patent Applications 20050238646 and 20020161201.

[0100] Antibody fragments can be obtained using conventional recombinant or protein engineering techniques, and the

fragments can be screened for antigen-binding or other function in the same manner as are intact antibodies.

[0101] Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of full-length antibodies (see, e.g., Morimoto et al., *Journal of Biochemical and Biophysical Methods*, 24:107-117 (1992); and Brennan et al., *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter et al., *Bio/Technology*, 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. In other embodiments, the antibody of choice is a single-chain Fv fragment (scFv). See WO 1993/16185; U.S. Pat. No. 5,571,894; and U.S. Pat. No. 5,587,458. The antibody fragment may also be a "linear antibody", e.g., as described in U.S. Pat. No. 5,641,870, for example. Such linear antibody fragments may be monospecific or bispecific.

Multispecific Molecules

[0102] In another aspect, the present invention features multispecific molecules (comprising an anti-hNKG2D antibody, or an antigen-fragment thereof, of the invention as defined in the claims. Such multispecific molecules include bispecific molecules comprising at least one first binding specificity for hNKG2D and a second binding specificity for a second target epitope.

[0103] One type of bispecific molecules are bispecific antibodies. Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Methods for making bispecific antibodies are known in the art, and traditional production of full-length bispecific antibodies is usually based on the coexpression of two immunoglobulin heavy-chain-light-chain pairs, where the two chains have different specificities (Millstein et al., *Nature*, 305: 537-539 (1983)). Bispecific antibodies can be prepared as full-length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies) or any other antigen-binding fragments described herein.

[0104] In the bispecific antibodies according to the present invention, at least one binding epitope is on the hNKG2D protein. The anti-NKG2D-binding moiety may be combined with second moiety that binds to a molecule on a pro-inflammatory leukocyte, e.g., a T-cell receptor molecule (e.g. CD2, CD3, CD4, or CD8), so as to focus cellular defense mechanisms to a pro-inflammatory hNKG2D-expressing cell. In this embodiment, the bispecific antibodies can, e.g., be used to direct cytotoxic agents to, or an ADCC/CDC attack on, pro-inflammatory cells that express NKG2D. The cytotoxic agent could be, e.g., saporin, an anti-interferon-alpha agent, a vinca alkaloid, the ricin A chain, methotrexate, or a radioactive isotope.

[0105] In another embodiment, the second moiety binds a cell-associated target that is presented on or expressed by cells associated with a disease state normally regulated by effector lymphocytes, such as cancer, viral infection, or the like. Thus, for example, a typical target may be a cell stress-associated molecule such as a MIC molecule (e.g., MIC-A or MIC-B) or a ULBP (e.g., Rae-1, H-60, ULBP2, ULBP3, HCMV UL18, or Rae-1 β) or a pathogen-associated molecule such as a viral hemagglutinin.

[0106] Other multispecific molecules include those produced from the fusion of a hNKG2D-binding antibody moiety to one or more other non-antibody proteins. Such multispecific proteins and how to construct them have been described in the art. See, e.g., Dreier et al. (*Bioconjug. Chem.* 9(4): 482-489 (1998)); U.S. Patent 6,046,310; U.S. Patent Publication No. 20030103984; European Patent Application 1 413 316; US Patent Publication No. 20040038339; von Strandmann et al., *Blood* (2006;107:1955-1962.), and WO 2004056873. According to the present invention, the non-antibody protein could be, for example, a suitable ligand for any of the antigens of "second moiety" described in the preceding section; e.g., a ligand for a T-cell or Fc receptor, or a cell-stress molecule such as MIC-A, MIC-B, ULBP, or a pathogen-associated molecule such as a viral hemagglutinin.

[0107] Multispecific molecules with more than two valencies are also contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.*, 147: 60 (1991).

[0108] The multispecific molecules of the present invention can be prepared by conjugating the constituent binding specificities using methods known in the art. For example, each binding specificity of the multispecific molecule can be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents can be used for covalent conjugation. Examples of cross-linking agents include protein A, carbodiimide, N-succinimidyl-S-acetyl-thioacetate (SATA), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), o-phenylenedimaleimide (oPDM), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), and sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (sulfo-SMCC) (see e.g., Karpovsky et al. (1984) *J. Exp. Med.* 160:1686; Liu, MA et al. (1985) *Proc. Natl. Acad. Sci. USA* 82:8648). Other methods include those described in Paulus (1985) *Behring Ins. Mitt.* No. 78, 118-132; Brennan et al. (1985) *Science* 229:81-83), and Glennie et al.

(1987) J. Immunol. 139: 2367-2375). Preferred conjugating agents are SATA and sulfo-SMCC, both available from Pierce Chemical Co. (Rockford, IL).

[0109] When the binding specificities are antibodies, they can be conjugated via sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In a particularly preferred embodiment, the hinge region is modified to contain an odd number of sulfhydryl residues, preferably one, prior to conjugation.

[0110] Alternatively, both binding specificities can be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful where the bispecific molecule is a mAb x mAb, mAb x Fab, Fab x F(ab')₂ or ligand x Fab fusion protein. A bispecific molecule of the invention can be a single chain molecule comprising one single chain antibody and a binding determinant, or a single chain bispecific molecule comprising two binding determinants. Bispecific molecules may comprise at least two single chain molecules. Methods for preparing bispecific molecules are described or reviewed in, for example in U.S. Patent Number 5,260,203; U.S. Patent Number 5,455,030; U.S. Patent Number 4,881,175; U.S. Patent Number 5,132,405; U.S. Patent Number 5,091, 513; U.S. Patent Number 5,476,786; U.S. Patent Number: 5,013,653; U.S. Patent Number 5,258,498; U.S. Patent Number 5,482,858; U.S. Patent application publication 20030078385, Kontermann et al., (2005) Acta Pharmacologica Sinica 26(1):1-9; Kostelny et al., (1992) J. Immunol. 148(5):1547-1553; Hollinger et al., (1993) PNAS (USA) 90:6444-6448; and Gruber et al. (1994) J. Immunol. 152: 5368.

Antibody variants

[0111] An antibody of the invention, as defined in the claims, further can be prepared using an antibody having one or more of the VH and/or VL sequences disclosed herein as starting material to engineer a modified antibody or antibody "variant", which modified antibody may have altered properties from the parent antibody. An antibody can be engineered by modifying one or more residues within one or both variable regions (*i.e.*, VH and/or VL), for example within one or more CDR regions and/or within one or more framework regions. Additionally or alternatively, an antibody can be engineered by modifying residues within the constant region(s), for example to alter the effector function(s) of the antibody. Additionally, from antigen-binding portions of an antibody, other constructs such as antigen-binding fragments, antibody derivatives, immunoconjugates, and multispecific molecules can be prepared.

[0112] Standard molecular biology techniques can be used to prepare and express the altered antibody sequence.

[0113] Though an antibody variant or derivative typically has at least one altered property as compared to the "parent" antibody, the antibody variant or derivative can retain one, some or most of the functional properties of the anti-hNKG2D antibodies described herein, which functional properties include, but are not limited to: (a) prevents NKG2D-mediated activation of an NKG2D-expressing NK or T cell, optionally with an EC₅₀ for reducing ligand-induced cytotoxicity lower than the EC₅₀ for binding to the cell; (b) competes with at least one NKG2D ligand in binding to NKG2D, preferably with at least MICA and ULBP3; (c) reduces the amount of NKG2D on the surface of a NKG2D-expressing NK or T cell, preferably with at least 75%; (d) binds to cynomolgous and/or rhesus NKG2D, preferably with substantially similar efficacy or affinity; (e) binds to more than one form or conformation of NKG2D; (f) binds to NKG2D with a K_d of 1 nM or less, preferably 0.1 nM or less; (g) competes with Ms (h) competes more with MS than with any of ON72, BAT221, 5C6, 1D11, ECM217, and 149810 in binding to hNKG2D; (i) blocks more than 90% of MS binding to cell-surface hNKG2D; (j) has less agonist activity on hNKG2D than any of ON72, BAT221, 5C6, 1D11, ECM217, and 149810. Any combination of the above-described functional features, and/or the functional features as described in the Examples, may be exhibited by an antibody of the invention.

[0114] The functional properties of the antibody variants and derivatives can be assessed using standard assays available in the art and/or described herein. For example, the ability of the antibody to bind hNKG2D can be determined using standard binding assays, such as those set forth in the Examples (e.g., Biacore, flow cytometry, or ELISAs).

Variable region modifications

[0115] One type of variable region engineering that can be performed is CDR grafting. Antibodies interact with target antigens predominantly through amino acid residues that are located in the six heavy and light chain complementarily determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to express recombinant antibodies that mimic the properties of specific naturally occurring antibodies by constructing expression vectors that

include CDR sequences from the specific naturally occurring antibody grafted onto framework sequences from a different antibody with different properties (see, e.g., Riechmann, L. et al. (1998) *Nature* 332:323-327; Jones, P. et al. (1986) *Nature* 321:522-525; Queen, C. et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:10029-10033; U.S. Patent No. 5,225,539 to Winter, and U.S. Patent Nos. 5, 530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen et al.) Accordingly, disclosed herein is also an isolated antibody, or antigen binding portion thereof, comprising: a VH region comprising CDR1, CDR2, and CDR3 sequences comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 15, 21, 48, and 54, SEQ ID NOs: 16, 22, 49, and 55, and SEQ ID NOs: 17, 23, 50, and 56, respectively, and a VL region comprising CDR1, CDR2, and CDR3 sequences comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 18, 24, 51, and 57, SEQ ID NOs: 19, 25, 52, and 58, and SEQ ID NOs: 20, 26, 53, and 59, respectively. Thus, such antibodies contain the VH and VL CDR sequences of antibodies 16F16, 16F31, MS, or 21F2, yet may contain framework sequences different from these antibodies.

[0116] Disclosed herein is also a chimeric or humanized version of a murine anti-hNKG2D monoclonal antibody, or antigen-binding fragment thereof, which binds hNKG2D, and the use of such antibodies (e.g., in the modulation of hNKG2D-mediated physiological processes in a mammalian host). In one embodiment, the murine antibody is one of ON72, BAT221, 5C6, 1D11, 149810, and ECM217. In another embodiment, the murine antibody is not one of ON72, BAT221, 5C6, 1D11, 149810, and ECM217. Thus, such antibodies contain the VH and VL CDR sequences of ON72, BAT221, 5C6, 1D11, 149810, or ECM217, or murine monoclonal antibody different from ON72, BAT221, 5C6, 1D11, 149810, ECM217, framework sequences different from these antibodies. In one embodiment, the humanized antibody is a humanized version of ON72, comprising e.g. the amino acid sequences of SEQ ID NOS:70 and 71 heavy- and light chain, respectively.

[0117] Such framework sequences can be obtained from public DNA databases or published references that include germline antibody gene sequences. For example, germline DNA sequences for human heavy and light chain variable region genes can be found in the "dBase" human germline sequence database (available on the Internet at www.mrc-cpe.cam.ac.uk/vbase), as well as in Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Tomlinson, I. M., et al. (1992) "The Repertoire of Human Germline VH Sequences Reveals about Fifty Groups of VH Segments with Different Hypervariable Loops" *J. Mol. Biol.* 227:776-798; and Cox, J. P. L. et al. (1994) "A Directory of Human Germ-line VH Segments Reveals a Strong Bias in their Usage" *Eur. J. Immunol.* 24:827-836.

[0118] Preferred framework sequences for use in the antibodies of the invention are those that are structurally similar to the framework sequences used by selected antibodies of the invention, e.g., similar to the VH3_21, D3-9, JH4, VKI_L15, and JK2, or VH3_20, D3-10, JH6, VKIII_A27, and JK3, or VH4_59, JH3, VKIII_A27, and JK1, or VH5_51, D3_10_R3 JH4, VKIII_L6, and JK1 framework sequences used by the 16F16 (reference), 16F31 (reference), MS, and 21F2 (reference) antibodies. The VH CDR1, 2 and 3 sequences of MS and the VL CDR1, 2 and 3 sequences of MS can be grafted onto framework regions that have the same sequence as that found in the germline immunoglobulin gene from which the framework sequence derive, or the CDR sequences can be grafted onto framework regions that contain one or more mutations as compared to the germline sequences. For example, it has been found that in certain instances it is beneficial to mutate residues within the framework regions to maintain or enhance the antigen binding ability of the antibody (see e.g., U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen et al.).

[0119] In another aspect of the invention, the structural features of anti-hNKG2D antibodies of the invention are used to create structurally related anti-hNKG2D antibodies that retain at least one functional property of the antibodies of the invention, including binding to hNKG2D. For example, CDR regions can be combined recombinantly with known framework regions to create additional, recombinantly-engineered, anti-hNKG2D antibodies of the invention. The starting material for the engineering method is one or more of the VH and/or VL sequences provided herein, or one or more CDR regions thereof. To create the engineered antibody, it is not necessary to actually prepare (*i.e.*, express as a protein) an antibody having one or more of the VH and/or VL sequences provided herein, or one or more CDR regions thereof. Rather, the information contained in the sequence(s) is used as the starting material to create a "second generation" sequence(s) derived from the original sequence(s) and then the "second generation" sequence(s) is prepared and expressed as a protein.

[0120] Accordingly, in another embodiment, the invention provides a method for preparing an anti-hNKG2D antibody, as defined in the claims, comprising: (a) providing: (i) a heavy chain variable region antibody sequence comprising a CDR1 sequence of SEQ ID NO: 48 a CDR2 sequence of SEQ ID NO: 49 and a CDR3 sequence of SEQ ID NO: 50; and (ii) a light chain variable region antibody sequence comprising a CDR1 sequence of SEQ ID NO: 51 a CDR2 sequence of SEQ ID NO: 52 and a CDR3 sequence of SEQ ID NO: 53; (b) altering at least one amino acid residue within the first antibody sequence and/or the second antibody sequence to create at least one altered antibody sequence; and (c) preparing the altered antibody sequence; and (d) expressing the altered antibody sequence as a protein.

[0121] Another type of variable region modification disclosed herein is to mutate amino acid residues within the VH and/or VL

CDR1, CDR2 and/or CDR3 regions to thereby improve one or more binding properties (e.g., affinity) of the antibody of interest. Site-directed mutagenesis or PCR-mediated mutagenesis can be performed to introduce the mutation(s) and the effect on antibody binding, or other functional property of interest, can be evaluated *in vitro* or *in vivo* assays as described herein and provided in the Examples. Preferably conservative modifications (as discussed above) are introduced. The mutations may be amino acid substitutions, additions or deletions. Moreover, typically no more than 8, more typically no more than 5 residues are altered within a single CDR region.

[0122] Accordingly, in another embodiment, disclosed herein are isolated anti-hNKG2D antibodies, comprising a heavy chain variable region comprising: (a) a VH CDR1 region comprising an amino acid sequence selected from SEQ ID NOs: 15, 21, 48, and 54, or an amino acid sequence having one, two, three, four, five, six, seven, or eight amino acid substitutions, deletions or additions as compared to an amino acid sequence selected from SEQ ID NOs:15, 21, 48, and 54; (b) a VH CDR2 region comprising an amino acid sequence selected from SEQ ID NOs:16, 22, 49, and 55, or an amino acid sequence having one, two, three, four, five, six, seven or eight amino acid substitutions, deletions or additions as compared to an amino acid sequence selected from SEQ ID NOs:16, 22, 49, and 55; (c) a VH CDR3 region comprising an amino acid sequence selected from SEQ ID NOs:17, 23, 50, and 56, or an amino acid sequence having one, two, three, four, five, six, seven or eight amino acid substitutions, deletions or additions as compared to an amino acid sequence selected from SEQ ID NOs:17, 23, 50, and 56; (d) a VL CDR1 region comprising an amino acid sequence selected from SEQ ID NO:18, 24, 51, and 57, or an amino acid sequence having one, two, three, four, five, six, seven, or eight amino acid substitutions, deletions or additions as compared to an amino acid sequence selected from SEQ ID NO:18, 24, 51, and 57; (e) a VL CDR2 region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs:19, 25, 52, and 58, or an amino acid sequence having one, two, three, four, five, six, seven, or eight amino acid substitutions, deletions or additions as compared to an amino acid sequence selected from the group consisting of SEQ ID NOs:19, 25, 52, and 58; and (f) a VL CDR3 region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 20, 26, 53, and 59, or an amino acid sequence having one, two, three, four, five, six, seven, or eight amino acid substitutions, deletions or additions as compared to an amino acid sequence selected from the group consisting of SEQ ID NOs:20, 26, 53 and 59.

[0123] Engineered antibodies of the invention include those in which modifications have been made to framework residues within VH and/or VL, e.g. to improve the properties of the antibody. Typically such framework modifications are made to decrease the immunogenicity of the antibody. For example, one approach is to "backmutate" one or more framework residues to the corresponding germline sequence. More specifically, an antibody that has undergone somatic mutation may contain framework residues that differ from the germline sequence from which the antibody is derived. Such residues can be identified by comparing the antibody framework sequences to the germline sequences from which the antibody is derived.

[0124] For example, for 16F16 (reference), amino acid residue R111 (Kabat residue 103; within FR4) of VH is an arginine whereas this residue in the corresponding germline sequence is a tryptophan (see Figure 5A). To return the framework region sequences to their germline configuration, some or all of the somatic mutations can be "backmutated" to the germline sequence by, for example, site-directed mutagenesis or PCR-mediated mutagenesis (e.g., residue 111 of the VH of 16F16 (reference) can be "backmutated" from threonine to alanine). As another example, for 16F31 (reference), amino acid residue Y95 (within FR3) of the VH region is a tyrosine whereas this residue in the corresponding germline sequence is a histidine (see Figure 5C). To return the framework region sequences to their germline configuration, the somatic mutation can be "backmutated" from tyrosine to histidine. As another example, for MS, Kabat residues 3, 6, and 7 of the VH region are histidine (H), aspartic acid (D), and D, respectively, whereas these residues in the corresponding germline sequences are glutamine (Q), glycine (G), and G, respectively (see Figure 5E). For 21F2 (reference) Kabat residues 13, 24, 76, and 93 are glutamic acid (E), asparagine (N), N, and G, respectively, whereas these residues in the corresponding germline sequences are lysine (K), G, serine (S), and alanine (A), respectively. To return the framework region sequences to their germline configuration, the somatic mutations can be similarly "backmutated". Such "backmutated" antibodies are also intended to be encompassed by the invention, as defined in the claims.

[0125] Another type of framework modification involves mutating one or more residues within the framework region, or even within one or more CDR regions, to remove T cell epitopes to thereby reduce the potential immunogenicity of the antibody. This approach is also referred to as "deimmunization" and is described in further detail in U.S. Patent Publication No. 20030153043 by Carr et al.

Fc modifications

[0126] In addition or as an alternative to modifications made within the framework or CDR regions, antibodies of the invention

may be engineered to include modifications within the Fc region, typically to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding, protein stability and/or antigen-dependent cellular cytotoxicity, or lack thereof. Furthermore, an antibody of the invention may be chemically modified (*e.g.*, one or more chemical moieties can be attached to the antibody) or be modified to alter its glycosylation, again to alter one or more functional properties of the antibody. Each of these embodiments is described in further detail below. The residues in the Fc region are numbered according to Kabat.

[0127] If desired, the class of an antibody may be "switched" by known techniques. Such techniques include, *e.g.*, the use of direct recombinant techniques (see *e.g.*, US Patent 4816397) and cell-cell fusion techniques (see *e.g.*, US Patent 5916771). For example, an antibody that was originally produced as an IgM molecule may be class switched to an IgG antibody. Class switching techniques also may be used to convert one IgG subclass to another, *e.g.*, from IgG1 to IgG2. Thus, the effector function of the antibodies of the invention may be changed by isotype switching to, *e.g.*, an IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM antibody for various therapeutic uses. Exemplary cDNA sequences for constant regions are available via, *e.g.*, GenBank (accessible via NCBI and other public websites), are as follows:

Human IgG1 constant heavy chain region: GenBank accession No.: J00228;

Human IgG2 constant heavy chain region: GenBank accession No.: J00230;

Human IgG3 constant heavy chain region: GenBank accession No.: X04646;

Human IgG4 constant heavy chain region: GenBank accession No.: K01316; and

Human kappa light chain constant region: GenBank accession No.: J00241.

[0128] In one embodiment, the hinge region of CH1 is modified such that the number of cysteine residues in the hinge region is altered, *e.g.*, increased or decreased. This approach is described further in U.S. Patent No. 5677425 by Bodmer *et al.* The number of cysteine residues in the hinge region of CH1 is altered to, for example, facilitate assembly of the light and heavy chains or to increase or decrease the stability of the antibody.

[0129] In another embodiment, the Fc hinge region of an antibody is mutated to decrease the biological half life of the antibody. More specifically, one or more amino acid mutations are introduced into the CH2-CH3 domain interface region of the Fc-hinge fragment such that the antibody has impaired Staphylococcal protein A (SpA) binding relative to native Fc-hinge domain SpA binding. This approach is described in further detail in U.S. Patent No. 6165745 by Ward *et al.* In another embodiment, the antibody is modified to increase its biological half life. Various approaches are possible. For example, one or more of the following mutations can be introduced: T252L, T254S, and T256F, as described in U.S. Patent No. 6277375 to Ward. Alternatively, to increase the biological half life, the antibody can be altered within the CH1 or CL region to contain a salvage receptor binding epitope taken from two loops of a CH2 domain of an Fc region of an IgG, as described in U.S. Patent Nos. 5869046 and 6121022 by Presta *et al.* In yet other embodiments, the Fc region is altered by replacing at least one amino acid residue with a different amino acid residue to alter the effector function(s) of the antibody. For example, one or more amino acids selected from amino acid residues 234, 235, 236, 237, 297, 318, 320 and 322 can be replaced with a different amino acid residue such that the antibody has an altered affinity for an effector ligand but retains the antigen-binding ability of the parent antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor or the C1 component of complement. This approach is described in further detail in U.S. Patent Nos. 5624821 and 5648260, both to Winter *et al.* In another example, one or more amino acids selected from amino acid residues 329, 331 and 322 can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Patent Nos. 6194551 by Idusogie *et al.* In another example, one or more amino acid residues within amino acid positions 231 and 239 are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in PCT Publication WO 94/29351 by Bodmer *et al.* In yet another example, the Fc region is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or to increase the affinity of the antibody for an Fcγ receptor by modifying one or more amino acids at the following positions: 238, 239, 248, 249, 252, 254, 255, 256, 258, 265, 267, 268, 269, 270, 272, 276, 278, 280, 283, 285, 286, 289, 290, 292, 293, 294, 295, 296, 298, 301, 303, 305, 307, 309, 312, 315, 320, 322, 324, 326, 327, 329, 330, 331, 333, 334, 335, 337, 338, 340, 360, 373, 376, 378, 382, 388, 389, 398, 414, 416, 419, 430, 434, 435, 437, 438 or 439. This approach is described further in PCT Publication WO 00/42072 by Presta. Moreover, the binding sites on human IgG1 for FcγRI, FcγRII, FcγRIII and FcRn have been mapped and variants with improved binding have been described (see Shields, R.L. *et al.* (2001) *J. Biol. Chem.* 276:6591-6604). Specific mutations at positions 256, 290, 298, 333, 334 and 339 were shown to improve binding to FcRn. Additionally, the following combination mutants were shown to improve FcγRIII binding: T256A/S298A, S298A/E333A, S298A/K224A and S298A/E333A/K334A.

[0130] The constant region may further be modified to stabilize the antibody, e.g., to reduce the risk of a bivalent antibody separating into two monovalent VH-VL fragments. For example, in an IgG4 constant region, residue S241 may be mutated to a proline (P) residue to allow complete disulphide bridge formation at the hinge (see, e.g., Angal et al., *Mol Immunol.* 1993;30:105-8).

Glycosylation modifications

[0131] In still another embodiment, the glycosylation of an antibody is modified. For example, an aglycosylated antibody can be made (*i.e.*, the antibody lacks glycosylation). Glycosylation can be altered to, for example, increase the affinity of the antibody for antigen. Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the antibody sequence. For example, one or more amino acid substitutions can be made that result in elimination of one or more variable region framework glycosylation sites to thereby eliminate glycosylation at that site. Such aglycosylation may increase the affinity of the antibody for antigen. Such an approach is described in further detail in U.S. Patent Nos. 5714350 and 6350861 by Co et al. Additionally or alternatively, an antibody can be made that has an altered type of glycosylation, such as a hypofucosylated antibody having reduced amounts of fucosyl residues or an antibody having increased bisecting GlcNac structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Such carbohydrate modifications can be accomplished by, for example, expressing the antibody in a host cell with altered glycosylation "machinery". Cells with such alterations have been described in the art and can be used as host cells in which to express recombinant antibodies of the invention to thereby produce an antibody with altered glycosylation. For example, EP1176195 by Hanai *et al.* describes a cell line with a functionally disrupted FUT8 gene, which encodes a fucosyl transferase, such that antibodies expressed in such a cell line exhibit hypofucosylation. PCT Publication WO 03/035835 by Presta describes a variant CHO cell line, Lec3 cells, with reduced ability to attach fucose to Asn(297)-linked carbohydrates, also resulting in hypofucosylation of antibodies expressed in that host cell (see also Shields, R.L. et al. (2002) *J. Biol. Chem.* 277:26733-26740). PCT Publication WO 99/54342 by Umana et al. describes cell lines engineered to express glycoprotein-modifying glycosyl transferases (e.g., beta(1,4)-N-acetylglucosaminyltransferase III (GnTIII)) such that antibodies expressed in the engineered cell lines exhibit increased bisecting GlcNac structures which results in increased ADCC activity of the antibodies (see also Umana et al. (1999) *Nat. Biotech.* 7:176 180).

[0132] In certain embodiments of the methods of engineering antibodies of the invention, mutations can be introduced randomly or selectively along all or part of an anti-hNKG2D antibody coding sequence (*e.g.*, MS coding sequence) and the resulting modified antibodies can be screened for binding activity and/or other functional properties as described herein. Mutational methods have been described in the art. For example, PCT Publication WO 02/092780 by Short describes methods for creating and screening antibody mutations using saturation mutagenesis, synthetic ligation assembly, or a combination thereof.

[0133] Alternatively, PCT Publication WO 03/074679 by Lazar et al. describes methods of using computational screening methods to optimize physiochemical properties of antibodies.

Antibody derivatives

[0134] Antibody derivatives (or immunoconjugates) within the scope of this invention include anti-hNKG2D antibodies conjugated or covalently bound to a second agent.

[0135] For example, in one aspect, the invention provides immunoconjugates comprising an antibody conjugated or covalently bonded to a cytotoxic agent. The term "cytotoxic agent" as used herein is a molecule that is capable of killing a cell bearing a hNKG2D receptor on its cell surface. Any type of moiety with a cytotoxic or cytotoxic effect can be conjugated to the present antibodies to form a cytotoxic conjugate of the present invention and to inhibit or kill specific NK receptor expressing cells, including therapeutic radioisotopes, toxic proteins, toxic small molecules, such as drugs, toxins, immunomodulators, hormones, hormone antagonists, enzymes, oligonucleotides, enzyme inhibitors, therapeutic radionuclides, angiogenesis inhibitors, chemotherapeutic drugs, vinca alkaloids, anthracyclines, epidophyllotoxins, taxanes, antimetabolites, alkylating agents, antibiotics, COX-2 inhibitors, SN-38, antimitotics, antiangiogenic and apoptotic agents, particularly doxorubicin, methotrexate, taxol, CPT-11, camptothecins, nitrogen mustards, gemcitabine, alkyl sulfonates, nitrosoureas, triazenes, folic acid analogs, pyrimidine analogs, purine analogs, platinum coordination complexes, *Pseudomonas* exotoxin, ricin, abrin, 5-fluorouridine, ribonuclease (RNase), DNase I, Staphylococcal enterotoxin-A, pokeweed antiviral protein, gelonin, diphtherin toxin, *Pseudomonas* exotoxin, and *Pseudomonas* endotoxin and others (see, e.g., Remington's Pharmaceutical Sciences, 19th Ed. (Mack Publishing

Co. 1995); Goodman and Gilman's The Pharmacological Basis of Therapeutics (McGraw Hill, 2001); Pastan et al. (1986) Cell 47:641; Goldenberg (1994) Cancer Journal for Clinicians 44:43; U.S. Pat. No. 6,077,499). It will be appreciated that a toxin can be of animal, plant, fungal, or microbial origin, or can be created de novo by chemical synthesis.

[0136] In another embodiment, the antibody is derivatized with a radioactive isotope, such as a therapeutic radionuclide or a radionuclide suitable for detection purposes. Any of a number of suitable radioactive isotopes can be used, including, but not limited to, I-131, Indium-111, Lutetium-171, Bismuth-212, Bismuth-213, Astatine-211, Copper-62, Copper-64, Copper-67, Yttrium-90, Iodine-125, Iodine-131, Phosphorus-32, Phosphorus-33, Scandium-47, Silver-111, Gallium-67, Praseodymium-142, Samarium-153, Terbium-161, Dysprosium-166, Holmium-166, Rhenium-186, Rhenium-188, Rhenium-189, Lead-212, Radium-223, Actinium-225, Iron-59, Selenium-75, Arsenic-77, Strontium-89, Molybdenum-99, Rhodium-105, Palladium-109, Praseodymium-143, Promethium-149, Erbium-169, Iridium-194, Gold-198, Gold-199, and Lead-211. In general, the radionuclide preferably has a decay energy in the range of 20 to 6,000 keV, preferably in the ranges 60 to 200 keV for an Auger emitter, 100-2,500 keV for a beta emitter, and 4,000-6,000 keV for an alpha emitter. Also preferred are radionuclides that substantially decay with generation of alpha-particles.

[0137] The antibody conjugates of the invention can be used to modify a given biological response, where the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, an enzymatically active toxin, or active fragment thereof, such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor or interferon- γ ; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

[0138] The second agent can be linked to the antibody directly or indirectly, using any of a large number of available methods. For example, an agent can be attached at the hinge region of the reduced antibody component via disulfide bond formation, using cross-linkers such as N-succinyl 3-(2-pyridyldithio)propionate (SPDP), or via a carbohydrate moiety in the Fc region of the antibody (see, e.g., Yu et al. (1994) Int. J. Cancer 56: 244; Wong, Chemistry of Protein Conjugation and Cross-linking (CRC Press 1991); Upešlacis et al., "Modification of Antibodies by Chemical Methods," in Monoclonal antibodies: principles and applications, Birch et al. (eds.), pages 187-230 (Wiley-Liss, Inc. 1995); Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in Monoclonal antibodies: Production, engineering and clinical application, Ritter et al. (eds.), pages 60-84 (Cambridge University Press 1995); Cattel et al. (1989) Chemistry today 7:51-58; Delprino et al. (1993) J. Pharm. Sci 82:699-704; Arpicco et al. (1997) Bioconjugate Chemistry 8:3; Reisfeld et al. (1989) Antibody, Immunicon. Radiopharm. 2:217). See, also, e.g. Amon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., 'The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev., 62:119-58 (1982).

[0139] For further discussion of types of cytotoxins, linkers and methods for conjugating therapeutic agents to antibodies, see also Saito, G. et al. (2003) Adv. Drug Deliv. Rev. 55:199-215; Trail, P.A. et al. (2003) Cancer Immunol. Immunother. 52:328-337; Payne, G. (2003) Cancer Cell 3:207-212; Allen, T.M. (2002) Nat. Rev. Cancer 2:750-763; Pastan, I. and Kreitman, R. J. (2002) Curr. Opin. Invest. Drugs 3:1089-1091; Senter, P.D. and Springer, C.J. (2001) Adv. Drug Deliv. Rev. 53:247-264.

[0140] In other embodiments, the second agent is a detectable moiety, which can be any molecule that can be quantitatively or qualitatively observed or measured. Examples of detectable markers useful in the conjugated antibodies of this invention are radioisotopes, fluorescent dyes, or a member of a complementary binding pair, such as a member of any one of: and antigen/antibody (other than an antibody to NKG2D), lectin/carbohydrate; avidin/biotin; receptor/ligand; or molecularly imprinted polymer/print molecule systems.

[0141] The second agent may also or alternatively be a polymer, intended to, e.g., increase the circulating half-life of the antibody. Exemplary polymers and methods to attach such polymers to peptides are illustrated in, e.g., U.S. Pat. Nos. 4766106; 4179337; 4495285; and 4609546. Additional illustrative polymers include polyoxyethylated polyols and polyethylene glycol (PEG) moieties. As used herein, the term "polyethylene glycol" is intended to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono (C1-C10) alkoxy- or aryloxy-polyethylene glycol or polyethylene glycol-maleimide. For example, a full-length antibody or antibody fragment can be conjugated to one or more PEG molecules with a molecular weight of

between about 1,000 and about 40,000, such as between about 2000 and about 20,000, e.g., about 3,000-12,000. To pegylate an antibody or fragment thereof, the antibody or fragment typically is reacted with polyethylene glycol (PEG), such as a reactive ester or aldehyde derivative of PEG, under conditions in which one or more PEG groups become attached to the antibody or antibody fragment. Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive PEG molecule (or an analogous reactive water-soluble polymer). In certain embodiments, the antibody to be pegylated is an aglycosylated antibody. Methods for pegylating proteins are known in the art and can be applied to the antibodies of the invention. See for example, EP154316 by Nishimura et al., International patent application PCT/US04/11494, and EP401384 by Ishikawa et al.

Nucleic acids

[0142] Another aspect of the invention pertains to nucleic acid molecules that encode the antibodies of the invention. The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is "isolated" or "rendered substantially pure" when purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art. See, F. Ausubel, et al., ed. (1987) *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York. A nucleic acid of the invention can be, for example, DNA or RNA and may or may not contain intronic sequences. In a preferred embodiment, the nucleic acid is a cDNA molecule. While the following paragraphs refer to DNA sequences or use thereof, the same methods or principles can generally be applied to mRNA sequences.

[0143] Nucleic acids of the invention can be obtained using standard molecular biology techniques. For antibodies expressed by hybridomas (e.g., hybridomas prepared from transgenic mice carrying human immunoglobulin genes as described further below), cDNAs encoding the light and heavy chains of the antibody made by the hybridoma can be obtained by standard PCR amplification or cDNA cloning techniques. For antibodies obtained from an immunoglobulin gene library (e.g., using phage display techniques), nucleic acids encoding the antibody can be recovered from the library.

[0144] Preferred nucleic acids molecules of the invention are those that encode (or comprise a nucleic acid sequence that encodes) the H and L chain sequences of the MS antibody of the IgG4 isotype. DNA sequences encoding the MS VH and VL sequences are those that encode for SEQ ID NOS:44 and 45, respectively.

[0145] Once DNA fragments encoding VH and VL segments are obtained, these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term "operatively linked", as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

[0146] The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (CH1, CH2 and CH3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region, but most preferably is an IgG4 constant region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH1 constant region.

[0147] The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U. S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region, but most preferably is a kappa constant region.

[0148] To create a scFv gene, the VH-and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly₄-Ser)₃, such that the VH and VL sequences can be expressed as a

contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see e.g., Bird et al. (1988) Science 242:423-426; Huston et al. (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883; McCafferty et al., (1990) Nature 348:552-554).

Antibody production

[0149] Monoclonal antibodies (mAbs) of the present invention can be produced by a variety of techniques, including conventional monoclonal antibody methodology e.g., the standard somatic cell hybridization technique of Kohler and Milstein (1975) Nature 256: 495. Although somatic cell hybridization procedures are preferred, in principle, other techniques for producing monoclonal antibody can be employed e.g., viral or oncogenic transformation of B lymphocytes.

[0150] One preferred animal system for preparing hybridomas is the murine system. Immunization protocols and techniques for isolation of immunized splenocytes for fusion are known in the art, as are fusion partners (e.g., murine myeloma cells) and fusion procedures. Humanized antibodies of the present invention can also be prepared based on the sequence of a murine monoclonal antibody using established techniques. For example, DNA encoding the heavy and light chain immunoglobulins can be obtained from the murine hybridoma of interest and engineered to contain non-murine (e.g., human) immunoglobulin sequences using standard molecular biology techniques. For example, to create a chimeric antibody (reference), the murine variable regions can be linked to human constant regions using methods known in the art (see e.g., U.S. Patent No. 4816567 to Cabilly et al.). To create a humanized antibody, the murine CDR regions can be inserted into a human framework using methods known in the art (see e.g., U.S. Patent No. 5225539 to Winter, and U.S. Patent Nos. 5530101; 5585089; 5693762 and 6180370 to Queen et al.).

[0151] In a preferred embodiment, the antibodies of the invention are human monoclonal antibodies. Such human monoclonal antibodies directed against hNKG2D can be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse system. These transgenic and transchromosomal mice include mice referred to herein as HuMAb mice and KM mice, respectively, and are collectively referred to herein as "human Ig mice." The HuMAb mouse (Medarex, Inc.) contains human immunoglobulin gene miniloci that encode unrearranged human heavy (p and y) and K light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous, u and K chain loci (see e.g., Lonberg, et al. (1994) Nature 368: 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or K, and, in response to immunization, the introduced human heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgGK monoclonal (Lonberg, N. *et al.* (1994), *supra*; reviewed in Lonberg, N. (1994) Handbook of Experimental Pharmacology 113:49-101; Lonberg, N. and Huszar, D. (1995) Intern. Rev. Immunol. 13:65-93, and Harding, F. and Lonberg, N. (1995) Ann. N. Y. Acad. Sci. 764:536-546). The preparation and use of HuMAb mice, and the genomic modifications carried by such mice, is further described in Taylor, L. et al. (1992) Nucleic Acids Research 20:6287-6295; Chen, J. et al. (1993) International Immunology 5: 647-656; Tuailon et al. (1993) Proc. Natl. Acad. Sci. USA 90:3720-3724; Choi et al. (1993) Nature Genetics 4: 117-123; Chen, J. et al. (1993) EMBO J. 12: 821-830; Tuailon et al. (1994) J. Immunol. 152:2912-2920; Taylor, L. et al. (1994) International immunology 6: 579-591; and Fishwild, D. et al. (1996) Nature Biotechnology 14: 845-851. See further, U. S. Patent Nos. 5545806; 5569825; 5625126; 5633425; 5789650; 5877397; 5661016; 5814318; 5874299; and 5770429; all to Lonberg and Kay; U.S. Patent No. 5545807 to Surani et al.; PCT Publication Nos. WO 92/03918, WO 93/12227, WO 94/25585, WO 97/13852, WO 98/24884 and WO 99/45962, all to Lonberg and Kay; and PCT Publication No. WO 01/14424 to Korman et al. In another embodiment, human antibodies of the invention can be raised using a mouse that carries human immunoglobulin sequences on transgenes and transchromosomes, such as a mouse that carries a human heavy chain transgene and a human light chain transchromosome. Such mice, referred to herein as "KM mice", are described in detail in PCT Publication WO 02/43478 to Ishida et al. Still further, alternative transgenic animal systems expressing human immunoglobulin genes are available in the art and can be used to raise anti-hNKG2D antibodies of the invention. For example, an alternative transgenic system referred to as the Xenomouse (Abgenix, Inc.) can be used; such mice are described in, for example, U.S. Patent Nos. 5939598; 6075181; 6114598; 6150584 and 6162963 to Kucherlapati et al. Moreover, alternative transchromosomal animal systems expressing human immunoglobulin genes are available in the art and can be used to raise anti-hNKG2D antibodies of the invention. For example, mice carrying both a human heavy chain transchromosome and a human light chain transchromosome, referred to as "TC mice" can be used; such mice are described in Tomizuka et al. (2000) Proc. Natl. Acad. Sci. USA 97:722-727. Furthermore, cows carrying human heavy and light chain transchromosomes have been described in the art (Kuroiwa et al. (2002) Nature Biotechnology 20:889-894) and can be used to raise anti-hNKG2D antibodies of the invention.

[0152] Human monoclonal antibodies of the invention can also be prepared using phage display methods for screening libraries of human immunoglobulin genes. Such phage display methods for isolating human antibodies are established in the art. See for example: U.S. Patent Nos. 5223409; 5403484; and 5571698 to Ladner et al.; U.S. Patent Nos. 5427908 and 5580717 to Dower et al.; U.S. Patent Nos. 5969108 and 6172197 to McCafferty et al.; and U.S. Patent Nos. 5885793; 6521404; 6544731; 6555313; 6582915 and 6593081 to Griffiths et al. Human monoclonal antibodies of the invention can also be prepared using SCID mice into

which human immune cells have been reconstituted such that a human antibody response can be generated upon immunization. Such mice are described in, for example, U.S. Patent Nos. 5476996 and 5698767 to Wilson et al.

[0153] When human Ig mice are used to raise human antibodies of the invention, such mice can be immunized with a purified or enriched preparation of hNKG2D antigen and/or cells expressing hNKG2D, as described by Lonberg, N. et al. (1994) *Nature* 368(6474): 856-859; Fishwild, D. et al. (1996) *Nature Biotechnology* 14: 845-851; and PCT Publication WO 98/24884 and WO 01/14424. Preferably, the mice will be 6-16 weeks of age upon the first infusion. For example, a purified or enriched preparation (5-50 µg) of hNKG2D antigen can be used to immunize the human Ig mice intraperitoneally. In the event that immunizations using a purified or enriched preparation of hNKG2D antigen do not result in antibodies, mice can also be immunized with cells expressing hNKG2D, e.g., a human NK or T-cell line, or a mammalian cell expressing recombinant hNKG2D with or without DAP10, to promote immune responses.

[0154] Detailed procedures to generate fully human monoclonal antibodies to hNKG2D are described in Example 1 below. The form and amount of antigen administered (e.g., hNKG2D polypeptide or cell expressing hNKG2D), as well as administration schedules and the possible use of adjuvants such as, e.g., complete Freund's adjuvant or incomplete Freund's adjuvant, are typically optimized for each antigen-mouse system according to established methods in the art.

[0155] The immune response can be monitored over the course of the immunization protocol with plasma samples being obtained by retroorbital bleeds, and the plasma or serum can be screened by ELISA (as described below), and mice with sufficient titers of anti-hNKG2D human immunoglobulin can be used for fusions. Mice can be boosted intravenously with antigen 3 days before sacrifice and removal of the spleen. It is expected that 2-3 fusions for each immunization may need to be performed.

[0156] To generate hybridomas producing human monoclonal antibodies of the invention, splenocytes and/or lymph node cells from immunized mice can be isolated and fused to an appropriate immortalized cell line, such as a mouse myeloma cell line. The resulting hybridomas can be screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic lymphocytes from immunized mice can be fused to one-sixth the number of P3X63-Ag8.653 nonsecreting mouse myeloma cells (ATCC, CRL 1580) with 50% PEG. Alternatively, the cells can be fused by electrofusion. Cells are plated at approximately 2×10^5 in a flat bottom microtiter plate, followed by a two week incubation in selective medium containing 20% fetal Clone Serum, 18% "653" conditioned media, 5% origen (IGEN), 4 mM L-glutamine, 1 mM sodium pyruvate, 5mM HEPES, 0.055 mM 2-mercaptoethanol, 50 units/ml penicillin, 50 mg/ml streptomycin, 50 mg/ml gentamycin and 1X HAT (Sigma; the HAT is added 24 hours after the fusion). After approximately two weeks, cells can be cultured in medium in which the HAT is replaced with HT. Individual wells can then be screened by ELISA for human monoclonal IgM and IgG antibodies. Once extensive hybridoma growth occurs, medium can be observed usually after 10-14 days. The antibody secreting hybridomas can be replated, screened again, and if still positive for human IgG, the monoclonal antibodies can be subcloned at least twice by limiting dilution. The stable subclones can then be cultured *in vitro* to generate small amounts of antibody in tissue culture medium for characterization. To purify human monoclonal antibodies, selected hybridomas can be grown in two-liter spinner-flasks for monoclonal antibody purification. Supernatants can be filtered and concentrated before affinity chromatography with protein A-sepharose (Pharmacia, Piscataway, N.J.). Eluted IgG can be checked by gel electrophoresis and high performance liquid chromatography to ensure purity. The buffer solution can be exchanged into PBS, and the concentration can be determined by spectroscopy. The monoclonal antibodies can be aliquoted and stored at -80°.

[0157] Antibodies of the invention can also be produced in a host cell transfectoma using, for example, a combination of recombinant DNA techniques and gene transfection methods as is well known in the art (e.g., Morrison, S. (1985) *Science* 229:1202).

[0158] For example, to express the antibodies, DNAs encoding partial or full-length light and heavy chains, can be obtained by standard molecular biology techniques (e.g. PCR amplification or cDNA cloning using a hybridoma that expresses the antibody of interest) and the DNAs can be inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences and may serve their intended function of regulating the transcription and translation of the antibody gene.

[0159] The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector or, more typically, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (e.g., ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present). The light and heavy chain variable regions of the antibodies described herein can be used to create full-length antibody genes of any antibody isotype by inserting them into expression vectors already encoding heavy chain

constant and light chain constant regions of the desired isotype such that the VH segment is operatively linked to the CH segment(s) within the vector and the VL segment is operatively linked to the CL segment within the vector. Additionally or alternatively, the recombinant expression vector can encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene can be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (*i.e.*, a signal peptide from a non-immunoglobulin protein).

[0160] In addition to the antibody chain genes, the recombinant expression vectors of the invention carry regulatory sequences that control the expression of the antibody chain genes in a host cell. The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.* polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel (Gene Expression Technology. Methods in Enzymology 185, Academic Press, San Diego, CA (1990)).

[0161] It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences, may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV), Simian Virus 40 (SV40), adenovirus, (*e.g.*, the adenovirus major late promoter (AdMLP) and polyoma. Alternatively, nonviral regulatory sequences may be used, such as the ubiquitin promoter or p-globin promoter. Still further, regulatory elements composed of sequences from different sources, such as the SRa promoter system, which contains sequences from the SV40 early promoter and the long terminal repeat of human T cell leukemia virus type 1 (Takebe, Y. et al. (1988) Mol. Cell. Biol. 8:466-472).

[0162] In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (*e.g.* origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see, *e.g.* U.S. Pat. Nos. 4399216, 4634665 and 5179017, all by Axel et al.). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr-host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

[0163] For expression of the light and heavy chains, the expression vector(s) encoding the heavy and light chains is transfected into a host cell by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, *e.g.*, electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like. Although it is theoretically possible to express the antibodies of the invention in either prokaryotic or eukaryotic host cells, expression of antibodies in eukaryotic cells, and most preferably mammalian host cells, is the most preferred because such eukaryotic cells, and in particular mammalian cells, are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active antibody. Prokaryotic expression of antibody genes has been reported to be ineffective for production of high yields of active antibody (Boss, M. A. and Wood, C. R. (1985) Immunology Today 6:12-13).

[0164] Preferred mammalian host cells for expressing the recombinant antibodies of the invention include Chinese Hamster Ovary (CHO cells) (including dhfr-CHO cells, described in Urlaub and Chasin, (1980) Proc. Natl. Acad. Sci. USA 77:4216-4220, used with a DHFR selectable marker, *e.g.*, as described in R. J. Kaufman and P. A. Sharp (1982) Mol. Biol. 159:601-621), NSO myeloma cells, COS cells and SP2 cells. In particular, for use with NSO myeloma cells, another preferred expression system is the GS gene expression system disclosed in WO 87/04462, WO 89/01036 and EP 338,841. When recombinant expression vectors encoding antibody genes are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods.

Antibody characterization

[0165] After production or purification, or as part of a screening or selection procedure, the functional characteristics of an anti-hNKG2D antibody of the invention can be investigated. Functional properties of interest include, *e.g.*, antibody binding specificity for hNKG2D, antibody competition with hNKG2D-ligands, antibody competition with reference antibodies (such as, *e.g.*, 16F16,

16F31, MS, and 21F2), the epitope to which the antibody binds, the affinity of the antibody-antigen interaction, and antagonistic/agonistic properties of the antibody.

[0166] The following are brief descriptions of exemplary assays for antibody characterization. Some are further described in subsequent sections and/or described in the Examples.

1. (1) Antibody specificity for hNKG2D can be evaluated by confirming that the monoclonal antibody (or, as part of animal screening procedures, serum containing polyclonal antibodies) binds NKG2D expressing cells but not NKG2D negative cells. Cell lines with or without NKG2D are incubated with antibody followed by incubation with secondary antibody directly labelled, and visualised by, *e.g.*, flow cytometry.
2. (2) Blockade of ligand binding can be evaluated by incubating cells expressing NKG2D with or without antibody or hybridoma supernatant, followed by incubation with a ligand-mFc protein and a secondary antibody specific for the ligand, and the level of ligand binding and blockade thereof determined by flow cytometry. Blockade can be calculated as the % ligand binding with pre-incubation compared to without pre-incubation, when lower binding is seen upon pre-incubation.
3. (3) Competition for binding site used by one or more reference anti-NKG2D antibodies can be evaluated in a similar manner, except that the pre-incubation can either performed with an antibody of the invention or the reference antibody (*e.g.*, ON72 or 149810), followed by incubation with and detection of the subsequently added antibody.
4. (4) Affinity parameters, including on- and off- rate, of antibodies can determined on a Biacore machine. For example, hNKG2D-Fc protein can be immobilized on a chip, the antibody passed over the chip, the on- and off-rates determined, and the KD calculated.
5. (5) Induction of NKG2D internalisation by antibodies can be measured by incubating hNKG2D-expressing cells with or without antibody overnight, followed by re-addition of the antibody and detection of the level of NKG2D (*i.e.* the level of antibody bound) in a flow cytometer.
6. (6) The ability of an antibody to block hNKG2D-ligand mediated killing can be assessed, using, *e.g.*, the NK cell lines NK92 or NKL as effector cells that kill ⁵¹Cr-loaded target cells expressing NKG2D ligand, either MICA, MICB, or ULBP1-4.
7. (7) Cross-reactivity of the human anti-NKG2D antibodies with monkey NK and CD8+ T cells but not CD4+ T cells (as in humans), can be demonstrated by flow cytometry after incubation of monkey and human PBMC's with hNKG2D antibody and secondary antibody, along with markers of the different cell types in PBMCs, and analysing NKG2D staining of the various subsets.
8. (8) Activation of NKG2D upon antibody binding can be measured as induction of cell-proliferation of CD8+ cells in a PBMC population upon stimulation via the T-cell receptor, CD28 and or NKG2D, with or without pre-stimulation (*e.g.*, via TCR, CD28 and IL-2 or IL-15).

Binding Assays

[0167] The present invention provides for antibodies, and antigen-binding fragments and immunoconjugates thereof, that bind hNKG2D. Any of a wide variety of assays can be used to assess binding of an antibody to hNKG2D. Protocols based upon ELISAs, radioimmunoassays, Western blotting, BIACORE, and other competition assays, *inter alia*, are suitable for use and are well known in the art. Further, several binding assays, including competition assays, are described in the Examples.

[0168] For example, simple binding assays can be used, in which a test antibody is incubated in the presence of a target protein or epitope (*e.g.*, NKG2D or a portion thereof), unbound antibodies are washed off, and the presence of bound antibodies is assessed using, *e.g.*, radiolabels, physical methods such as mass spectrometry, or direct or indirect fluorescent labels detected using, *e.g.*, cytofluorometric analysis (*e.g.* FACScan). Such methods are well known to those of skill in the art. Any amount of binding above the amount seen with a control, non-specific antibody indicates that the antibody binds specifically to the target.

[0169] In such assays, the ability of the test antibody to bind to the target cell or human NKG2D can be compared with the ability of a (negative) control protein, *e.g.* an antibody raised against a structurally unrelated antigen, or a non-Ig peptide or protein, to bind to the same target. Antibodies or fragments that bind to the target cells or NKG2D using any suitable assay with 25%, 50%, 100%, 200%, 1000%, or higher increased affinity relative to the control protein, are said to "specifically bind to" or "specifically interact with" the target, and are preferred for use in the therapeutic methods described below. The ability of a test antibody to affect the binding of a (positive) control antibody against NKG2D, *e.g.* 16F16, 16F31, MS, or 21 F2, may also be assessed.

[0170] In one aspect, the invention provides for anti-hNKG2D antibodies, as defined in the claims, sharing biological characteristics and/or substantial VH and/or VL sequence identity with MS. One exemplary biological characteristic is the binding

to the MS epitope, *i.e.*, the respective region in the extracellular domain of hNKG2D to which the MS antibody binds. To screen for antibodies that bind to the MS epitope, a routine cross-blocking assay, such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed.

[0171] In an exemplary cross-blocking or competition assay, MS (control) antibody and a test antibody are admixed (or pre-adsorbed) and applied to a sample containing NKG2D. In certain embodiments, one would pre-mix the control antibodies with varying amounts of the test antibody (e.g., 1:10 or 1:100) for a period of time prior to applying to the NKG2D-containing sample. In other embodiments, the control and varying amounts of test antibody can simply be admixed during exposure to the antigen/target sample. As long as one can distinguish bound from free antibodies (e.g., by using separation or washing techniques to eliminate unbound antibodies) and the control antibody from test antibody (e.g., by using species- or isotype-specific secondary antibodies, by specifically labeling the control antibody with a detectable label, or by using physical methods such as mass spectrometry to distinguish between different compounds) one will be able to determine if the test antibody reduces the binding of the control antibody to the antigen, indicating that the test antibody recognizes substantially the same epitope as the control. In this assay, the binding of the (labeled) control antibody in the presence of a completely irrelevant antibody is the control high value. The control low value is obtained by incubating the labeled (positive) control antibody with unlabeled control antibody, where competition would occur and reduce binding of the labeled antibody.

[0172] In a test assay, a significant reduction in labeled antibody reactivity in the presence of a test antibody is indicative of a test antibody that recognizes the same epitope, *i.e.*, one that "cross-reacts" with the labeled control antibody. Any test antibody or compound that reduces the binding of the labeled control to the antigen/target by at least 50% or more preferably 70%, at any ratio of control:test antibody or compound between about 1:10 and about 1:100 is considered to be an antibody or compound that binds to substantially the same epitope or determinant as the control. Preferably, such test antibody or compound will reduce the binding of the control to the antigen/target by at least 90%. Nevertheless, any compound or antibody that reduces the binding of a control antibody or compound to any measurable extent can be used in the present invention.

[0173] In one embodiment, competition can be assessed by a flow cytometry test. Cells bearing hNKG2D are incubated first with a control antibody that is known to specifically bind to the receptor (e.g., T or NK cells expressing hNKG2D or BaF/3 cell recombinantly expressing hNKG2D, and MS antibody), and then with the test antibody that may be labeled with, e.g., a fluorochrome or biotin. The test antibody is said to compete with the control if the binding obtained with preincubation with saturating amounts of control antibody is 80%, preferably, 50%, 40% or less of the binding (mean of fluorescence) obtained by the antibody without preincubation with the control. Alternatively, a test antibody is said to compete with the control if the binding obtained with a labeled control (by a fluorochrome or biotin) on cells preincubated with saturating amount of antibody to test is 80%, preferably 50%, 40%, or less of the binding obtained without preincubation with the antibody. See Example 5 for an exemplary antibody competition assay.

[0174] Similar cross-blocking assays can also be used to evaluate whether a test (humanized) antibody affects the binding of a natural ligand for human NKG2D, such as MICA, MICB, ULBP1, ULBP2, ULBP3, ULBP4, or a member of the RAET1 family, simply by exchanging 16F16, 16F31, MS, or 21F2 for a suitable form of the hNKG2D-ligand. One suitable form, described in the Examples, are fusion proteins of the ligand (e.g., MICA) with the Fc-portion of an antibody. Having the ligand conjugated to an Fc-region allows for detection of the fusion protein by antibodies specific for the animal species from which the Fc-region derives, using, e.g., goat-anti-mouse antibodies to detect a murine Fc-region.

[0175] In one embodiment, a cellular assay is used in which hNKG2D-expressing cells, e.g., CD4⁺CD28⁻ cells from rheumatoid arthritis patients (or the equivalent cells from another autoimmune or inflammatory disorder) are incubated with an NKG2D ligand such as MICA, MICB, or a ULBP protein, e.g., in the form of an Fc-fusion protein, or a cell expressing any of these ligands, and the ability of an anti-NKG2D antibody or other molecule to block the activation of the cell is assessed. In an alternative assay, a baseline level of activity for the NKG2D receptor is obtained in the absence of a ligand, and the ability of the antibody or compound to cause a decrease in the baseline activity level is detected. In one type of embodiment, a high-throughput screening approach is used to identify compounds capable of blocking the activation of the receptor, or otherwise downregulating it. See Example 3 for an exemplary ligand competition assay.

[0176] Preferably, monoclonal antibodies that recognize an NKG2D epitope will react with an epitope that is present on a substantial percentage of CD4⁺ T cells, particularly CD4⁺CD28⁻ T cells, in patients such as rheumatoid arthritis patients, but will not significantly react with other cells, *i.e.*, immune or non-immune cells that do not express NKG2D. Accordingly, once an antibody that specifically recognizes hNKG2D on NK or T cells, it can be tested for its ability to bind to T cells taken from patients with autoimmune or inflammatory disorders such as rheumatoid arthritis. It will be appreciated that the present invention can be used for the treatment of any disorder in which NKG2D activity is linked to the pathology of the disorder, regardless of the cell

type expressing the receptor (e.g., CD4+ T cells, CD8+ T cells, NK cells, etc.), and the antibodies can be tested for their ability to bind to the receptor on whichever cell type is relevant for the particular disorder. For example, if it is observed that a particular disorder is associated with excess activity or proliferation of NKG2D-expressing NK cells, then the antibodies can be developed and tested using NK cells expressing the same receptor.

[0177] In one embodiment, the antibodies are validated in an immunoassay to test its ability to bind to NKG2D-expressing cells, e.g. CD4+CD28- T cells taken from patients with rheumatoid arthritis. For example, peripheral blood lymphocytes (PBLs) are taken from a plurality of patients, and CD4+, preferably CD4+ CD28-, cells are enriched from the PBLs, e.g., by flow cytometry using relevant antibodies. The ability of a given antibody to bind to the cells is then assessed using standard methods well known to those in the art. Antibodies that are found to bind to a substantial proportion (e.g., 20%, 30%, 40%, 50%, 60%, 70%, 80% or more) of cells known to express NKG2D, e.g. NK cells, CD8 T cells, CD4 T cells from RA patients, etc., from a significant percentage of patients (e.g., 5%, 10%, 20%, 30%, 40%, 50% or more) can be deemed suitable for use in the present invention, both for diagnostic purposes to determine the expression of the NKG2D receptor in a patient's cells or for use in the herein-described therapeutic methods, e.g., for use as human-suitable blocking or, alternatively, cytotoxic antibodies. To assess the binding of the antibodies to the cells, the antibodies can either be directly or indirectly labeled. When indirectly labeled, a secondary, labeled antibody is typically added. The binding of the antibodies to the cells can then be detected using, e.g., cytofluorometric analysis (e.g. FACS). Such methods are well known in the art.

[0178] In some aspects of the invention, e.g., where it is not desired to kill NKG2D-expressing cells, the antibodies of the invention preferably do not demonstrate substantial specific binding to Fc receptors. Such antibodies may comprise constant regions of various heavy chains that are known not to bind Fc receptors. One such example is an IgG4 conconstant stant region. Alternatively, antibody fragments that do not comprise constant regions, such as Fab or F(ab')₂ fragments, can be used to avoid Fc receptor binding. Fc receptor binding can be assessed according to methods known in the art, including for example testing binding of an antibody to Fc receptor protein in a BIAcore assay. Also, any other antibody type can be used in which the Fc portion is modified to minimize or eliminate binding to Fc receptors (see, e.g., WO3101485). Assays such as, e.g., cell based assays, to assess Fc receptor binding are well known in the art, and are described in, e.g., WO3101485.

Functional assays

[0179] Any suitable physiological change that reflects NKG2D activity can be used to assess the utility of a test compound or antibody. For example, one can measure a variety of effects in, e.g., cell-based assays, such as changes in gene expression, cytokine production, signalling molecule phosphorylation, cell growth, cell proliferation, pH, intracellular second messengers, e.g., Ca²⁺, IP₃, cGMP, or cAMP, or activity such as cytotoxic activity or ability to activate other T cells. For example, the activity of the receptor can be assessed by detecting the expression of NKG2D-responsive genes, e.g., CD25, IFN-gamma, or TNF-alpha (see, e.g., Groh et al. (2003) PNAS 100: 9452-9457; André et al. (2004) Eur. J. Immunol 34: 1-11). Alternatively, NKG2D activity can be assessed by incubating CD4+CD28-NKG2D+ cells in the presence of a ligand or activating anti-NKG2D antibody, as well as an anti-CD3 antibody, to evaluate the ability of the compound or test antibody to inhibit the release of TNF-alpha or IFN-gamma by the T cells. Alternatively, CD4+CD28-NKG2D+ T cells can be incubated in the presence of ligand, e.g., MICA, MICB, ULBP-1, ULBP-2, ULBP-3, etc., or ligand-producing cells, e.g., autologous MIC+ RA synoviocytes, and the ability of the test antibody or compound to inhibit cytokine production (e.g., IFN-gamma or TNF-alpha), or T cell proliferation assessed.

[0180] *In vitro* assays can optionally use cells taken from patients with autoimmune or inflammatory disorders such as RA, e.g. CD4+CD28- cells expressing NKG2D taken from (or cell lines derived therefrom) patients with RA, but in general any NKG2D-expressing cells can be used. For example, non-RA immune cell lines, e.g. T cell lines, can be transfected with an NKG2D-encoding transgene and used in the present assays, so long that the expression of the receptor alters the activity of the cells in a detectable way, e.g., renders them activatable by NKG2D ligand. Cell lines can, for example, be established using CD4+CD28-NKG2D+ cells from RA patients, e.g. PBLs or T cells isolated from synovial tissue. Such cells can be cultured in the presence of IL-15 to ensure continued expression of NKG2D (see, e.g., Groh et al. (2003) PNAS 100: 9452-9457).

[0181] If an anti-hNKG2D antibody reduces or blocks NKG2D interactions with one or more of its ligands, or competes with an antibody known to block hNKG2D ligand interaction, it can be useful for reducing NKG2D-mediated activation of NK or T cells. This can be evaluated by a typical cytotoxicity assays. Example 6 describes an exemplary cytotoxicity assay, NKG2D-ligand mediated killing of target cells. Here, the ability of anti-hNKG2D antibodies to reduce or inhibit the NK cell-mediated killing of MICA-transfected BaF/3 is assessed by measuring target cell release of ⁵¹Cr.

[0182] In other aspects, it may desirable to ensure that antibodies of the invention lack substantial agonistic activity. Several

assays can be used for this purpose, including the following.

[0183] One assay can evaluate proliferation and cytokine production after activation with antibodies, either soluble or plate-bound, in combination with anti-CD3 and/or anti-CD28 antibodies, of PBMCs from healthy volunteers or IBD patients. In this method, PBMCs are purified by conventional methods from healthy subjects or inflammatory bowel disease (IBD) patients. The cells are stained with CFSE (from Molecular probes, cat #C34554). To 10^7 cells (in 0.5 ml PBS with 2% FCS) is added 1 μ l CFSE (0.5 mM) and the cells are incubated at 37°C for 10 min. Then, 2 ml FCS is added, and the mixture is left for 1 min at room temperature. The cells are then washed 3 times by centrifugation with RPMI-1640 medium (12 ml). After wash, the cells are resuspended in 1 ml media (e.g. RPMI-1640) with 2% FCS.

[0184] Ninety-six well plates are coated with 30 μ l anti-mouse Fc (Jackson - Immuno Research 115-006-008) for 2 hours at room temperature, and then washed with PBS. Antibodies (anti-CD3 Bioscience cat#14-0037-82, anti-CD28 cat#348046 Becton Dickinson) are added according to the scheme below and left in the well:

Cells alone

CD3 0.1 or 0.3 ng/ml

CD3 0.1 or 0.3 ng/ml +CD28 0.2 μ g/ml

CD3 0.1 or 0.3 ng/ml+CD28 0.2 μ g/ml+anti-NKG2D 0.2 μ g/ml

CD3 0.1 or 0.3 ng/ml+anti-NKG2D 0.2 μ g/ml

[0185] Next, 100.000 CFSE-labelled PBMCs are added and left for 3 days. Supernatant is then collected for analysis of cytokines, and the PBMCs are analysed by flow cytometry with regard to the type of lymphocyte with anti-CD56, anti-CD4, anti-CD8, and CFSE labelling for proliferation.

[0186] In another assay, the effect on the cytotoxic potential of CD8+ T cells towards a target cells lacking NKG2D ligands, is tested. If binding boosts the cytotoxic potential of the cells, agonistic activity is present. Briefly, IL-2 stimulated PBMC from healthy subjects are incubated with p815 cells expressing MICA, or with untransfected p815 cells and an anti-CD3 antibody, (which will lead to redirected killing by binding to the Fc receptors on p815 cells) and CD8 cytotoxic T cells. It is then analyzed whether an anti-NKG2D antibody that does not bind to p815 cells (e.g., an antibody of human IgG4 isotype) blocks MICA-NKG2D-directed binding and/or if the antibody boosts CD3-p815 redirected binding. In this manner, it can be shown that the activity of the CD8+ T cells is not enhanced by incubating p815 cells with an anti-CD3 antibody and an additional anti-NKG2D antibody, while the same anti-NKG2D antibody can shown to be functional by demonstrating that it blocks NK-MICA interaction on p815-induced killing in the same PBMC population.

[0187] In another assay, it can be explored whether NKG2D-signalling pathways and - molecules are activated by addition of one or more anti-NKG2D antibodies. NK cell lines (such as, e.g., NKL cells or NK-92 cells), or human NK or CD8+ T cells isolated from peripheral blood, can be used. For example, NKL cells can be incubated with a human anti-NKG2D antibody in solution or plate bound, with, e.g., Fc-MICA or irradiated MICA expressing cells as a control. After incubation for suitable time periods, (e.g., 5, 10, 30 min), the cells are lysed in the presence of protease and phosphatase inhibitors on ice, and analyzed for the levels of one or more phosphorylated signalling molecules that are known to be downstream of stimulation of NKG2D (e.g., Pi3K, Akt, and vav), by standard Western blotting techniques.

[0188] In animal-based assays, any physiological or pathological consequence of NKG2D activation in cells within the animal can be used to assess antibody or test compound activity.

[0189] For example, CD4+CD28-NKG2D+ cells can be introduced into the joints of an animal model, with or without co-administration of ligand producing cells such as MICA-producing synoviocytes, and inflammation or tissue damage is assessed. Test compounds or antibodies can then be introduced, and their ability to inhibit, slow, reverse, or in any way affect the inflammation or tissue damage is detected.

[0190] Experiments with rheumatoid arthritis (RA) synovial explants can also be performed to study the effects of blocking NKG2D on spontaneous release of pro-inflammatory cytokines (see, e.g. Brennan et al., Lancet 1989; 2 (8657);244-247). In such an assay, human or humanized anti-hNKG2D antibodies are tested on RA synovial membrane cultures and compared to, e.g.,

murine anti-hNKG2D antibodies at concentrations shown to be useful to block ligand binding and function of NKG2D. RA synovial cells are cultured for 48 hrs in the absence or presence of anti-NKG2D antibodies or an isotype control antibody. Known anti-inflammatory drugs can be used as positive controls. The effects of the anti-NKG2D antibodies are initially tested at concentrations up to 30 µg/ml on 6 RA synovial membranes. Viability of the cells is analysed in a assay staining living cells (e.g. a MTT assay) to determine if the added reagent has any cytotoxicity. ELISA is then used to detect cytokines such as, e.g., TNFα, IL-1β and IL-6 levels in culture supernatants.

[0191] Alternatively, antibodies of the invention can be tested in experimental models of, e.g., psoriasis or ulcerative colitis. Psoriasis-affected skin sample can be transplanted onto a SCID mouse together with the patients own PBMC's, and the effect of introduction of a test compound and their ability to inhibit, slow, reverse, or in any way affect the inflammation or tissue damage, can be detected. Kjellev et al. (Eur J Immunol 2008;37:1397-1406) and Ito et al. (Am J Physiol Gastrointest Liver Physiol 2008;294:G199-G207) describe experimental models for assessing treatment of ulcerative colitis using anti-murine NKG2D antibody.

Pharmaceutical Formulations

[0192] In one embodiment, the present invention provides a pharmaceutical composition or formulation comprising anti-hNKG2D antibodies as defined in the claims, together with one or more carriers.

[0193] Accordingly, one exemplary aspect of the invention is a pharmaceutical formulation comprising such an antibody which is present in a concentration from 1 mg/ml to 500 mg/ml, and wherein said formulation has a pH from 2.0 to 10.0. The formulation may further comprise a buffer system, preservative(s), tonicity agent(s), chelating agent(s), stabilizers, and/or surfactants. In one embodiment, the pharmaceutical formulation is an aqueous formulation, i.e., formulation comprising water. Such formulation is typically a solution or a suspension. In a further embodiment, the pharmaceutical formulation is an aqueous solution. The term "aqueous formulation" is defined as a formulation comprising at least 50 %w/w water. Likewise, the term "aqueous solution" is defined as a solution comprising at least 50 %w/w water, and the term "aqueous suspension" is defined as a suspension comprising at least 50 %w/w water.

[0194] In another embodiment, the pharmaceutical formulation is a freeze-dried formulation, whereto the physician or the patient may add solvents and/or diluents prior to administration.

[0195] In another embodiment, the pharmaceutical formulation is a dried formulation (e.g. freeze-dried or spray-dried) ready for use without any prior dissolution.

[0196] In a further aspect, the pharmaceutical formulation comprises an aqueous solution of such an antibody, and a buffer, wherein the antibody is present in a concentration from 1 mg/ml or above, and wherein said formulation has a pH from about 2.0 to about 10.0.

[0197] In a another embodiment, the pH of the formulation is in the range selected from the list consisting of from about 2.0 to about 10.0, about 3.0 to about 9.0, about 4.0 to about 8.5, about 5.0 to about 8.0, and about 5.5 to about 7.5.

[0198] In a further embodiment, the formulation includes a buffer that is selected from the group consisting of sodium acetate, sodium carbonate, citrate, glycylglycine, histidine, glycine, lysine, arginine, sodium dihydrogen phosphate, disodium hydrogen phosphate, sodium phosphate, and tris(hydroxymethyl)-aminomethan, bicine, tricine, malic acid, succinate, maleic acid, fumaric acid, tartaric acid, aspartic acid or mixtures thereof. Each one of these specific buffers constitutes an alternative embodiment of the invention.

[0199] In a further embodiment, the formulation also or alternatively comprises a pharmaceutically acceptable preservative. The preservative may be selected from, e.g., the group consisting of phenol, o-cresol, m-cresol, p-cresol, methyl p-hydroxybenzoate, propyl p-hydroxybenzoate, 2-phenoxyethanol, butyl p-hydroxybenzoate, 2-phenylethanol, benzyl alcohol, chlorobutanol, and thiomerosal, bronopol, benzoic acid, imidurea, chlorohexidine, sodium dehydroacetate, chlorocresol, ethyl p-hydroxybenzoate, benzethonium chloride, chlorphenesine (3p-chlorophenoxypropane-1,2-diol) or mixtures thereof. The preservative may, e.g., be present in a concentration from 0.1 mg/ml to 20 mg/ml, from 0.1 mg/ml to 5 mg/ml, from 5 mg/ml to 10 mg/ml, or from 10 mg/ml to 20 mg/ml. Each one of these specific preservatives constitutes an alternative embodiment of the invention. The use of a preservative in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0200] In a further embodiment, the formulation also or alternatively comprises an isotonic agent. The isotonic agent may be, *e.g.*, selected from the group consisting of a salt (*e.g.* sodium chloride), a sugar or sugar alcohol, an amino acid (*e.g.* L-glycine, L-histidine, arginine, lysine, isoleucine, aspartic acid, tryptophan, threonine), an alditol (*e.g.* glycerol (glycerine), 1,2-propanediol (propyleneglycol), 1,3-propanediol, 1,3-butanediol) polyethyleneglycol (*e.g.* PEG400) or mixtures thereof. Any sugar such as mono-, di-, or polysaccharides, or watersoluble glucans, including for example fructose, glucose, mannose, sorbose, xylose, maltose, lactose, sucrose, trehalose, dextran, pullulan, dextrin, cyclodextrin, soluble starch, hydroxyethyl starch and carboxymethylcellulose-Na may be used. In one embodiment, the sugar additive is sucrose. Sugar alcohol is defined as a C4-C8 hydrocarbon having at least one -OH group and includes, for example, mannitol, sorbitol, inositol, galactitol, dulcitol, xylitol, and arabitol. In one embodiment, the sugar alcohol additive is mannitol. The sugars or sugar alcohols mentioned above may be used individually or in combination. There is no fixed limit to the amount used, as long as the sugar or sugar alcohol is soluble in the liquid preparation and does not adversely effect the stabilizing effects achieved using the methods of the invention. The sugar or sugar alcohol concentration can, *e.g.*, be between about 1 mg/ml and about 150 mg/ml. The isotonic agent can be present in a concentration from, *e.g.*, 1 mg/ml to 50 mg/ml, from 1 mg/ml to 7 mg/ml, from 8 mg/ml to 24 mg/ml, or from 25 mg/ml to 50 mg/ml. Each one of these specific isotonic agents constitutes an alternative embodiment of the invention. The use of an isotonic agent in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0201] In a further embodiment, the formulation also or alternatively comprises a chelating agent. The chelating agent can, for example, be selected from salts of ethylenediaminetetraacetic acid (EDTA), citric acid, and aspartic acid, and mixtures thereof. The chelating agent may, for example, be present in a concentration from 0.1 mg/ml to 5 mg/ml, from 0.1 mg/ml to 2 mg/ml, or from 2 mg/ml to 5 mg/ml. Each one of these specific chelating agents constitutes an alternative embodiment of the invention. The use of a chelating agent in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0202] In a further embodiment of the invention the formulation also or alternatively comprises a stabilizer. The use of a stabilizer in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995. More particularly, compositions of the invention can be stabilized liquid pharmaceutical compositions whose therapeutically active components include a polypeptide that possibly exhibits aggregate formation during storage in liquid pharmaceutical formulations. By "aggregate formation" is intended a physical interaction between the polypeptide molecules that results in formation of oligomers, which may remain soluble, or large visible aggregates that precipitate from the solution. By "during storage" is intended a liquid pharmaceutical composition or formulation once prepared, is not immediately administered to a subject. Rather, following preparation, it is packaged for storage, either in a liquid form, in a frozen state, or in a dried form for later reconstitution into a liquid form or other form suitable for administration to a subject. By "dried form" is intended the liquid pharmaceutical composition or formulation is dried either by freeze drying (*i.e.*, lyophilization; see, for example, Williams and Polli (1984) J. Parenteral Sci. Technol. 38:48-59), spray drying (see Masters (1991) in Spray-Drying Handbook (5th ed; Longman Scientific and Technical, Essex, U.K.), pp. 491-676; Broadhead et al. (1992) Drug Devel. Ind. Pharm. 18:1169-1206; and Mumenthaler et al. (1994) Pharm. Res. 11:12-20), or air drying (Carpenter and Crowe (1988) Cryobiology 25:459-470; and Roser (1991) Biopharm. 4:47-53). Aggregate formation by a polypeptide during storage of a liquid pharmaceutical composition can adversely affect biological activity of that polypeptide, resulting in loss of therapeutic efficacy of the pharmaceutical composition. Furthermore, aggregate formation may cause other problems such as blockage of tubing, membranes, or pumps when the polypeptide-containing pharmaceutical composition is administered using an infusion system.

[0203] The pharmaceutical compositions of the invention may alternatively or further comprise an amount of an amino acid base sufficient to decrease aggregate formation by the polypeptide during storage of the composition. By "amino acid base" is intended an amino acid or a combination of amino acids, where any given amino acid is present either in its free base form or in its salt form. Where a combination of amino acids is used, all of the amino acids may be present in their free base forms, all may be present in their salt forms, or some may be present in their free base forms while others are present in their salt forms. In one embodiment, amino acids to use in preparing the compositions of the invention are those carrying a charged side chain, such as arginine, lysine, aspartic acid, and glutamic acid. Any stereoisomer (*i.e.*, L, D, or a mixture thereof) of a particular amino acid (*e.g.* methionine, histidine, imidazole, arginine, lysine, isoleucine, aspartic acid, tryptophan, threonine and mixtures thereof) or combinations of these stereoisomers, may be present in the pharmaceutical compositions of the invention so long as the particular amino acid is present either in its free base form or its salt form. In one embodiment the L-stereoisomer is used. Compositions of the invention may also be formulated with analogues of these amino acids. By "amino acid analogue" is intended a derivative of the naturally occurring amino acid that brings about the desired effect of decreasing aggregate formation by the polypeptide during storage of the liquid pharmaceutical compositions of the invention. Suitable arginine analogues include, for example, aminoguanidine, ornithine and N-monoethyl L-arginine, suitable methionine analogues include ethionine and buthionine

and suitable cysteine analogues include S-methyl-L cysteine. As with the other amino acids, the amino acid analogues are incorporated into the compositions in either their free base form or their salt form. In a further embodiment of the invention the amino acids or amino acid analogues are used in a concentration, which is sufficient to prevent or delay aggregation of the protein.

[0204] In a further embodiment of the invention methionine (or other sulphuric amino acids or amino acid analogous) may be added to inhibit oxidation of methionine residues to methionine sulfoxide when the polypeptide acting as the therapeutic agent is a polypeptide comprising at least one methionine residue susceptible to such oxidation. The term "inhibit" in this context is intended to mean minimal accumulation of methionine oxidized species over time. Inhibiting methionine oxidation results in greater retention of the polypeptide in its proper molecular form. Any stereoisomer of methionine (L or D) or combinations thereof can be used. The amount to be added should be an amount sufficient to inhibit oxidation of the methionine residues such that the amount of methionine sulfoxide is acceptable to regulatory agencies. Typically, this means that the composition contains no more than about 10% to about 30% methionine sulfoxide. Generally, this can be achieved by adding methionine such that the ratio of methionine added to methionine residues ranges from about 1:1 to about 1000:1, such as 10:1 to about 100:1.

[0205] In a further embodiment, the formulation further or alternatively comprises a stabilizer selected from the group of high molecular weight polymers or low molecular compounds. In a further embodiment of the invention the stabilizer is selected from polyethylene glycol (e.g. PEG 3350), polyvinyl alcohol (PVA), polyvinylpyrrolidone, carboxy-/hydroxycellulose or derivatives thereof (e.g. HPC, HPC-SL, HPC-L and HPMC), cyclodextrins, sulphur-containing substances as monothioglycerol, thioglycolic acid and 2-methylthioethanol, and different salts (e.g. sodium chloride). Each one of these specific stabilizers constitutes an alternative embodiment of the invention.

[0206] The pharmaceutical compositions may also or alternatively comprise additional stabilizing agents, which further enhance stability of a therapeutically active polypeptide therein. Stabilizing agents of particular interest to the present invention include, but are not limited to, methionine and EDTA, which protect the polypeptide against methionine oxidation, and a nonionic surfactant, which protects the polypeptide against aggregation associated with freeze-thawing or mechanical shearing.

[0207] In a further embodiment, the formulation further or alternatively comprises a surfactant. The surfactant may, for example, be selected from a detergent, ethoxylated castor oil, polyglycolized glycerides, acetylated monoglycerides, sorbitan fatty acid esters, polyoxypropylene-polyoxyethylene block polymers (eg. poloxamers such as Pluronic® F68, poloxamer 188 and 407, Triton X-100), polyoxyethylene sorbitan fatty acid esters, polyoxyethylene and polyethylene derivatives such as alkylated and alkoxyated derivatives (tweens, e.g. Tween-20, Tween-40, Tween-80 and Brij-35), monoglycerides or ethoxylated derivatives thereof, diglycerides or polyoxyethylene derivatives thereof, alcohols, glycerol, lectins and phospholipids (eg. phosphatidyl serine, phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl inositol, diphosphatidyl glycerol and sphingomyelin), derivatives of phospholipids (eg. dipalmitoyl phosphatidic acid) and lysophospholipids (eg. palmitoyl lysophosphatidyl-L-serine and 1-acyl-sn-glycero-3-phosphate esters of ethanolamine, choline, serine or threonine) and alkyl, alkoxy (alkyl ester), alkoxy (alkyl ether)-derivatives of lysophosphatidyl and phosphatidylcholines, e.g. lauroyl and myristoyl derivatives of lysophosphatidylcholine, dipalmitoylphosphatidylcholine, and modifications of the polar head group, that is cholines, ethanolamines, phosphatidic acid, serines, threonines, glycerol, inositol, and the positively charged DODAC, DOTMA, DCP, BISHOP, lysophosphatidylserine and lysophosphatidylthreonine, and glycerophospholipids (eg. cephalins), glyceroglycolipids (eg. galactopyransoide), sphingoglycolipids (eg. ceramides, gangliosides), dodecylphosphocholine, hen egg lysolecithin, fusidic acid derivatives- (e.g. sodium tauro-dihydrofusidate etc.), long-chain fatty acids and salts thereof C6-C12 (e.g., oleic acid and caprylic acid), acylcarnitines and derivatives, N^α-acylated derivatives of lysine, arginine or histidine, or side-chain acylated derivatives of lysine or arginine, N^α-acylated derivatives of dipeptides comprising any combination of lysine, arginine or histidine and a neutral or acidic amino acid, N^α-acylated derivative of a tripeptide comprising any combination of a neutral amino acid and two charged amino acids, DSS (docusate sodium, CAS registry no [577-11-7]), docusate calcium, CAS registry no [128-49-4]), docusate potassium, CAS registry no [7491-09-0]), SDS (sodium dodecyl sulphate or sodium lauryl sulphate), sodium caprylate, cholic acid or derivatives thereof, bile acids and salts thereof and glycine or taurine conjugates, ursodeoxycholic acid, sodium cholate, sodium deoxycholate, sodium taurocholate, sodium glycocholate, N-Hexadecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate, anionic (alkyl-aryl-sulphonates) monovalent surfactants, zwitterionic surfactants (e.g. N-alkyl-N,N-dimethylammonio-1-propanesulfonates, 3-cholamido-1-propyldimethylammonio-1-propanesulfonate, cationic surfactants (quaternary ammonium bases) (e.g. cetyl-trimethylammonium bromide, cetylpyridinium chloride), non-ionic surfactants (eg. Dodecyl β-D-glucopyranoside), poloxamines (eg. Tetronic's), which are tetrafunctional block copolymers derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine, or the surfactant may be selected from the group of imidazoline derivatives, or mixtures thereof. Each one of these specific surfactants constitutes an alternative embodiment of the invention.

[0208] The use of a surfactant in pharmaceutical compositions is well-known to the skilled person. For convenience reference is made to Remington: The Science and Practice of Pharmacy, 19th edition, 1995.

[0209] In a further embodiment, the formulation further or alternatively comprises protease inhibitors such as EDTA (ethylenediamine tetraacetic acid) and benzamidineHCl, but other commercially available protease inhibitors may also be used. The use of a protease inhibitor is particularly useful in pharmaceutical compositions comprising zymogens of proteases in order to inhibit autocatalysis.

[0210] It is possible that other ingredients may also or alternatively be present in the peptide pharmaceutical formulation of the present invention. Such additional ingredients may include wetting agents, emulsifiers, antioxidants, bulking agents, tonicity modifiers, chelating agents, metal ions, oleaginous vehicles, proteins (e.g., human serum albumin, gelatine or proteins) and a zwitterion (e.g., an amino acid such as betaine, taurine, arginine, glycine, lysine and histidine). Such additional ingredients, of course, should not adversely affect the overall stability of the pharmaceutical formulation of the present invention.

[0211] Pharmaceutical compositions containing an antibody according to the present invention may be administered to a patient in need of such treatment at several sites, for example, at topical sites, for example, skin and mucosal sites, at sites which bypass absorption, for example, administration in an artery, in a vein, in the heart, and at sites which involve absorption, for example, administration in the skin, under the skin, in a muscle or in the abdomen.

[0212] Administration of pharmaceutical compositions according to the invention may be through several routes of administration, for example, lingual, sublingual, buccal, in the mouth, oral, in the stomach and intestine, nasal, pulmonary, for example, through the bronchioles and alveoli or a combination thereof, epidermal, dermal, transdermal, vaginal, rectal, ocular, for example through the conjunctiva, uretal, and parenteral to patients in need of such a treatment.

[0213] Compositions of the current invention may be administered in several dosage forms, for example, as solutions, suspensions, emulsions, microemulsions, multiple emulsion, foams, salves, pastes, plasters, ointments, tablets, coated tablets, rinses, capsules, for example, hard gelatine capsules and soft gelatine capsules, suppositories, rectal capsules, drops, gels, sprays, powder, aerosols, inhalants, eye drops, ophthalmic ointments, ophthalmic rinses, vaginal pessaries, vaginal rings, vaginal ointments, injection solution, in situ transforming solutions, for example in situ gelling, in situ setting, in situ precipitating, in situ crystallization, infusion solution, and implants.

[0214] Compositions of the invention may further be compounded in, or attached to, for example through covalent, hydrophobic and electrostatic interactions, a drug carrier, drug delivery system and advanced drug delivery system in order to further enhance stability of the antibody, increase bioavailability, increase solubility, decrease adverse effects, achieve chronotherapy well known to those skilled in the art, and increase patient compliance or any combination thereof. Examples of carriers, drug delivery systems and advanced drug delivery systems include, but are not limited to, polymers, for example cellulose and derivatives, polysaccharides, for example dextran and derivatives, starch and derivatives, poly(vinyl alcohol), acrylate and methacrylate polymers, polylactic and polyglycolic acid and block copolymers thereof, polyethylene glycols, carrier proteins, for example albumin, gels, for example, thermogelling systems, for example block co-polymeric systems well known to those skilled in the art, micelles, liposomes, microspheres, nanoparticulates, liquid crystals and dispersions thereof, L2 phase and dispersions thereof, well known to those skilled in the art of phase behaviour in lipid-water systems, polymeric micelles, multiple emulsions, self-emulsifying, self-microemulsifying, cyclodextrins and derivatives thereof, and dendrimers.

[0215] Compositions of the current invention are useful in the formulation of solids, semisolids, powder and solutions for pulmonary administration of an antibody, using, for example a metered dose inhaler, dry powder inhaler and a nebulizer, all being devices well known to those skilled in the art.

[0216] Compositions of the current invention are specifically useful in the formulation of controlled, sustained, protracting, retarded, and slow release drug delivery systems. More specifically, but not limited to, compositions are useful in formulation of parenteral controlled release and sustained release systems (both systems leading to a many-fold reduction in number of administrations), well known to those skilled in the art. Even more preferably, are controlled release and sustained release systems administered subcutaneous. Without limiting the scope of the invention, examples of useful controlled release system and compositions are hydrogels, oleaginous gels, liquid crystals, polymeric micelles, microspheres, nanoparticles,

[0217] Methods to produce controlled release systems useful for compositions of the current invention include, but are not limited to, crystallization, condensation, co-crystallization, precipitation, co-precipitation, emulsification, dispersion, high pressure homogenisation, encapsulation, spray drying, microencapsulating, coacervation, phase separation, solvent evaporation to produce microspheres, extrusion and supercritical fluid processes. General reference is made to Handbook of Pharmaceutical

Controlled Release (Wise, D.L., ed. Marcel Dekker, New York, 2000) and Drug and the Pharmaceutical Sciences vol. 99: Protein Formulation and Delivery (MacNally E.J., ed. Marcel Dekker, New York, 2000).

[0218] Parenteral administration may be performed by subcutaneous, intramuscular, intraperitoneal or intravenous injection by means of a syringe, optionally a pen-like syringe. Alternatively, parenteral administration can be performed by means of an infusion pump. A further option is a composition which may be a solution or suspension for the administration of the antibody compound in the form of a nasal or pulmonic spray. As a still further option, the pharmaceutical compositions containing an antibody of the invention can also be adapted to transdermal administration, *e.g.* by needle-free injection or from a patch, optionally an iontophoretic patch, or transmucosal, *e.g.* buccal, administration.

[0219] The antibody can be administered via the pulmonary route in a vehicle, as a solution, suspension or dry powder using any of known types of devices suitable for pulmonary drug delivery. Examples of these comprise of, but are not limited to, the three general types of aerosol-generating for pulmonary drug delivery, and may include jet or ultrasonic nebulizers, metered-dose inhalers, or dry powder inhalers (Cf. Yu J, Chien YW. Pulmonary drug delivery: Physiologic and mechanistic aspects. Crit Rev Ther Drug Carr Sys 14(4) (1997) 395-453).

[0220] Based on standardised testing methodology, the aerodynamic diameter (d_a) of a particle is defined as the geometric equivalent diameter of a reference standard spherical particle of unit density (1 g/cm^3). In the simplest case, for spherical particles, d_a is related to a reference diameter (d) as a function of the square root of the density ratio as described by:

$$d_a = \sqrt{\frac{\rho}{\rho_o}} d$$

[0221] Modifications to this relationship occur for non-spherical particles (cf. Edwards DA, Ben-Jebria A, Langer R. Recent advances in pulmonary drug delivery using large, porous inhaled particles. J Appl Physiol 84(2) (1998) 379-385). The terms "MMAD" and "MMEAD" are well-described and known to the art (cf. Edwards DA, Ben-Jebria A, Langer R and represents a measure of the median value of an aerodynamic particle size distribution. Recent advances in pulmonary drug delivery using large, porous inhaled particles. J Appl Physiol 84(2) (1998) 379-385). Mass median aerodynamic diameter (MMAD) and mass median effective aerodynamic diameter (MMEAD) are used interchangeably, are statistical parameters, and empirically describe the size of aerosol particles in relation to their potential to deposit in the lungs, independent of actual shape, size, or density (cf. Edwards DA, Ben-Jebria A, Langer R. Recent advances in pulmonary drug delivery using large, porous inhaled particles. J Appl Physiol 84(2) (1998) 379-385). MMAD is normally calculated from the measurement made with impactors, an instrument that measures the particle inertial behaviour in air.

[0222] In a further embodiment, the formulation could be aerosolized by any known aerosolisation technology, such as nebulisation, to achieve a MMAD of aerosol particles less than $10 \mu\text{m}$, more preferably between $1.5 \mu\text{m}$, and most preferably between $1.3 \mu\text{m}$. The preferred particle size is based on the most effective size for delivery of drug to the deep lung, where protein is optimally absorbed (cf. Edwards DA, Ben-Jebria A, Langer A, Recent advances in pulmonary drug delivery using large, porous inhaled particles. J Appl Physiol 84(2) (1998) 379-385).

[0223] Deep lung deposition of the pulmonary formulations comprising the antibody may optionally be further optimized by using modifications of the inhalation techniques, for example, but not limited to: slow inhalation flow (*eg.* 30 L/min), breath holding and timing of actuation.

[0224] The term "stabilized formulation" refers to a formulation with increased physical stability, increased chemical stability or increased physical and chemical stability.

[0225] The term "physical stability" of the protein formulation as used herein refers to the tendency of the antibody to form biologically inactive and/or insoluble aggregates as a result of exposure of the antibody to thermo-mechanical stresses and/or interaction with interfaces and surfaces that are destabilizing, such as hydrophobic surfaces and interfaces. Physical stability of the aqueous antibody formulations is evaluated by means of visual inspection and/or turbidity measurements after exposing the formulation filled in suitable containers (*e.g.* cartridges or vials) to mechanical/physical stress (*e.g.* agitation) at different temperatures for various time periods. Visual inspection of the formulations is performed in a sharp focused light with a dark background. The turbidity of the formulation is characterized by a visual score ranking the degree of turbidity for instance on a scale from 0 to 3 (a formulation showing no turbidity corresponds to a visual score 0, and a formulation showing visual turbidity in daylight corresponds to visual score 3). A formulation is classified physical unstable with respect to antibody aggregation, when it

shows visual turbidity in daylight. Alternatively, the turbidity of the formulation can be evaluated by simple turbidity measurements well-known to the skilled person. Physical stability of the aqueous antibody formulations can also be evaluated by using a spectroscopic agent or probe of the conformational status of the antibody. The probe is preferably a small molecule that preferentially binds to a non-native conformer of the antibody. One example of a small molecular spectroscopic probe of protein structure is Thioflavin T. Thioflavin T is a fluorescent dye that has been widely used for the detection of amyloid fibrils. In the presence of fibrils, and perhaps other protein configurations as well, Thioflavin T gives rise to a new excitation maximum at about 450 nm and enhanced emission at about 482 nm when bound to a fibril protein form. Unbound Thioflavin T is essentially non-fluorescent at the wavelengths.

[0226] Other small molecules can be used as probes of the changes in protein structure from native to non-native states. For instance the "hydrophobic patch" probes that bind preferentially to exposed hydrophobic patches of a protein. The hydrophobic patches are generally buried within the tertiary structure of a protein in its native state, but become exposed as a protein begins to unfold or denature. Examples of these small molecular, spectroscopic probes are aromatic, hydrophobic dyes, such as anthracene, acridine, phenanthroline or the like. Other spectroscopic probes are metal-amino acid complexes, such as cobalt metal complexes of hydrophobic amino acids, such as phenylalanine, leucine, isoleucine, methionine, and valine, or the like.

[0227] The term "chemical stability" of the antibody formulation as used herein refers to chemical covalent changes in the antibody structure leading to formation of chemical degradation products with potential less biological potency and/or potential increased immunogenic properties compared to the native antibody structure. Various chemical degradation products can be formed depending on the type and nature of the native antibody and the environment to which the antibody is exposed. Elimination of chemical degradation can most probably not be completely avoided and increasing amounts of chemical degradation products is often seen during storage and use of the antibody formulation as well-known by the person skilled in the art. Most proteins are prone to deamidation, a process in which the side chain amide group in glutaminyl or asparaginyl residues is hydrolysed to form a free carboxylic acid. Other degradations pathways involves formation of high molecular weight transformation products where two or more protein molecules are covalently bound to each other through transamidation and/or disulfide interactions leading to formation of covalently bound dimer, oligomer and polymer degradation products (Stability of Protein Pharmaceuticals, Ahem. T.J. & Manning M.C., Plenum Press, New York 1992). Oxidation (of for instance methionine residues) can be mentioned as another variant of chemical degradation. The chemical stability of the antibody formulation can be evaluated by measuring the amount of the chemical degradation products at various time-points after exposure to different environmental conditions (the formation of degradation products can often be accelerated by for instance increasing temperature). The amount of each individual degradation product is often determined by separation of the degradation products depending on molecule size and/or charge using various chromatography techniques (e.g. SEC-HPLC and/or RP-HPLC).

[0228] Hence, as outlined above, a "stabilized formulation" refers to a formulation with increased physical stability, increased chemical stability or increased physical and chemical stability. In general, a formulation must be stable during use and storage (in compliance with recommended use and storage conditions) until the expiration date is reached.

[0229] In one embodiment of the invention the pharmaceutical formulation comprising the antibody is stable for more than 6 weeks of usage and for more than 3 years of storage.

[0230] In another embodiment of the invention the pharmaceutical formulation comprising the antibody is stable for more than 4 weeks of usage and for more than 3 years of storage.

[0231] In a further embodiment of the invention the pharmaceutical formulation comprising the antibody is stable for more than 4 weeks of usage and for more than two years of storage.

[0232] In an even further embodiment of the invention the pharmaceutical formulation comprising the antibody is stable for more than 2 weeks of usage and for more than two years of storage.

[0233] Suitable antibody formulations can also be determined by examining experiences with other already developed therapeutic monoclonal antibodies. Several monoclonal antibodies have been shown to be efficient in clinical situations, such as Rituxan (Rituximab), Herceptin (Trastuzumab) Xolair (Omalizumab), Bexar (Tositumomab), Campath (Alemtuzumab), Zevalin, Oncolym, Humira and similar formulations may be used with the antibodies of this invention. For example, a monoclonal antibody can be supplied at a concentration of 10 mg/mL in either 100 mg (10 mL) or 500 mg (50 mL) single-use vials, formulated for IV administration in 9.0 mg/mL sodium chloride, 7.35 mg/mL sodium citrate dihydrate, 0.7 mg/mL polysorbate 80, and sterile water for injection. The pH is adjusted to 6.5. Alternatively, the antibody can be formulated in a solution comprising histidin, sucrose, and Polysorbate 80.

Diagnostic applications

[0234] The hNKG2D-antibodies of the invention also have non-therapeutic applications. For example, anti-hNKG2D antibodies may also be useful in diagnostic assays for NKG2D protein, e.g. detecting its expression in specific cells, tissues, or serum. For example, anti-hNKG2D antibodies could be used in assays selecting patients for anti-hNKG2D treatment. For such purposes, the anti-hNKG2D antibodies could be used for analyzing for the presence of hNKG2D in serum or tissue specimens, testing for the presence of CD4+ T cells expressing NKG2D, or the presence of disease promoting cells expressing NKG2D (e.g., NK or CD4+ or CD8+ T cells). Such analyses could be combined with analyses testing, e.g., for the levels of soluble MICA in blood (see, e.g., WO2003089616 by Spies et al.).

[0235] For diagnostic applications, the antibody typically will be labeled with a detectable moiety. Numerous labels are available that can be generally grouped into the following categories:

1. (a) Radioisotopes, such as ^{35}S , ^{14}C , ^{125}I , ^3H , and ^{131}I . The antibody can be labeled with the radioisotope using the techniques described in Current Protocols in Immunology, Volumes 1 and 2, Coligen et al., Ed. Wiley-Interscience, New York, N.Y., Pubs. (1991), for example, and radioactivity can be measured using scintillation counting.
2. (b) Fluorescent labels such as rare-earth chelates (europium chelates) or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, Lissamine, phycoerythrin and Texas Red are available. The fluorescent labels can be conjugated to the antibody using the techniques disclosed in Current Protocols in Immunology, *supra*, for example. Fluorescence can be quantified using a fluorimeter.
3. (c) Various enzyme-substrate labels are available and U.S. Pat. No. 4,275,149 provides a review of some of these. The enzyme generally catalyzes a chemical alteration of the chromogenic substrate that can be measured using various techniques. For example, the enzyme may catalyze a color change in a substrate, which can be measured spectrophotometrically. Alternatively, the enzyme may alter the fluorescence or chemiluminescence of the substrate. Techniques for quantifying a change in fluorescence are described above. The chemiluminescent substrate becomes electronically excited by a chemical reaction and may then emit light that can be measured (using a chemiluminometer, for example) or donates energy to a fluorescent acceptor. Examples of enzymatic labels include luciferases (e.g., firefly luciferase and bacterial luciferase; U.S. Pat. No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, malate dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRPO), alkaline phosphatase, beta-galactosidase, glucoamylase, lysozyme, saccharide oxidases (e.g., glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like. Techniques for conjugating enzymes to antibodies are described in O'Sullivan et al, "Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme Immunoassay," in Methods in Enzym. (Ed., J. Langone & H. Van Vunakis), Academic Press, New York, 73:147-166 (1981).

[0236] Examples of enzyme-substrate combinations include, for example:

1. (i) Horseradish peroxidase (HRPO) with hydrogen peroxide as a substrate, wherein the hydrogen peroxide oxidizes a dye precursor (e.g., orthophenylene diamine (OPD) or 3,3',5,5'-tetramethyl benzidine hydrochloride (TMB));
2. (ii) alkaline phosphatase (AP) with para-nitrophenyl phosphate as chromogenic substrate; and
3. (iii) beta-D-galactosidase (beta-D-Gal) with a chromogenic substrate (e.g., p-nitrophenyl-beta-D-galactosidase) or fluorogenic substrate 4-methylumbelliferyl-p-beta-galactosidase.

[0237] Numerous other enzyme-substrate combinations are available to those skilled in the art. For a general review of these, see U.S. Pat. Nos. 4,275,149 and 4,318,980.

[0238] Sometimes, the label is indirectly conjugated with the antibody. The skilled artisan will be aware of various techniques for achieving this. For example, the antibody can be conjugated with biotin, and any of the three broad categories of labels mentioned above can be conjugated with avidin, or vice versa. Biotin binds selectively to avidin, and thus, the label can be conjugated with the antibody in this indirect manner. Alternatively, to achieve indirect conjugation of the label with the antibody, the antibody is conjugated with a small hapten (e.g., digoxin) and one of the different types of labels mentioned above is conjugated with an anti-hapten antibody (e.g., anti-digoxin antibody). Thus, indirect conjugation of the label with the antibody can be achieved.

[0239] In another embodiment of the invention, the anti-NKG2D antibody need not be labeled, and the presence thereof can be detected using a labeled secondary antibody that binds to the NKG2D antibody.

[0240] The antibodies of the present invention may be employed in any known assay method, such as competitive-binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp. 147-158 (CRC Press, Inc. 1987).

[0241] For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

[0242] The antibodies may also be used for *in vivo* diagnostic assays. Generally, the antibody is labeled with a radionuclide or a non-radioactive indicator detectable by, *e.g.*, nuclear magnetic resonance, or other means known in the art. Preferably, the label is a radiolabel, such as, *e.g.*, ^{125}I , ^{131}I , ^{67}Cu , $^{99\text{m}}\text{Tc}$, or ^{111}In . The labeled antibody is administered to a host, preferably via the bloodstream, and the presence and location of the labeled antibody in the host is assayed. This imaging technique is suitably used in the detection, staging and treatment of neoplasms. The radioisotope is conjugated to the protein by any means, including metal-chelating compounds or lactoperoxidase, or iodogen techniques for iodination.

[0243] As a matter of convenience, the antibodies of the present invention can be provided in a kit, *i.e.*, a packaged combination of reagents in predetermined amounts with instructions for performing the diagnostic assay. Where the antibody is labeled with an enzyme, the kit will include substrates and cofactors required by the enzyme (*e.g.*, a substrate precursor that provides the detectable chromophore or fluorophore). In addition, other additives may be included such as stabilizers, buffers (*e.g.*, a block buffer or lysis buffer) and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents that substantially optimize the sensitivity of the assay. Particularly, the reagents may be provided as dry powders, usually lyophilized, including excipients that on dissolution will provide a reagent solution having the appropriate concentration.

Therapeutic applications

[0244] The use of a human or humanized anti-hNKG2D antibody as defined the claims for treating a patient is also provided for by the present invention. In one embodiment, the invention provides for the use of a human or humanized antibody as defined the claims in the preparation of a pharmaceutical composition for administration to a human patient. Typically, the patient suffers from, or is at risk for, an autoimmune or inflammatory disease or disorder.

[0245] For example, in one aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for reducing or inhibiting hNKG2D-mediated activation of NK or T cells in a patient in need thereof, comprising the step of administering a human or humanized anti-NKG2D antibody to the patient, which antibody reduces or prevents ligand-mediated activation of the NKG2D receptor. In one embodiment, the use of a human or humanized anti-hNKG2D antibody as defined in the claims is directed at decreasing the activity of such lymphocytes in patients having a disease in which increased NK or T cell activity is detrimental, which involves, affects or is caused by cells susceptible to lysis by NK or T cells, or which is caused or characterized by increased NK and/or T cell activity, such as an autoimmune disease or disorder or an inflammatory condition. In one aspect, the invention provides an antibody as defined in the claims for use in reducing chronic inflammation in the patient.

[0246] Exemplary conditions or disorders to be treated with the antibodies of the invention, include, but are not limited to systemic lupus erythematosus, rheumatoid arthritis, juvenile chronic arthritis, psoriatic arthritis, osteoarthritis, spondyloarthropathies (ankylosing spondylitis), systemic sclerosis (scleroderma), idiopathic inflammatory myopathies (dermatomyositis, polymyositis), Sjogren's syndrome, vasculitis, systemic vasculitis, temporal arteritis, atherosclerosis, sarcoidosis, myasthenia gravis, autoimmune hemolytic anemia (immune pancytopenia, paroxysmal nocturnal hemoglobinuria), pernicious anemia, autoimmune thrombocytopenia (idiopathic thrombocytopenic purpura, immune-mediated thrombocytopenia), thyroiditis (Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, atrophic thyroiditis), diabetes mellitus, immune-mediated renal disease (glomerulonephritis, tubulointerstitial nephritis, autoimmune oophoritis), autoimmune orchitis, autoimmune uveitis, anti-phospholipid syndrome, demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or Guillain-Barre syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious hepatitis (hepatitis A, B, C, D, E and other non-hepatotropic viruses), autoimmune chronic active hepatitis, viral hepatitis, primary biliary cirrhosis, granulomatous hepatitis, Wegener's granulomatosis, Behcet's disease, and sclerosing cholangitis, inflammatory bowel diseases such as ulcerative colitis or Crohn's disease, celiac

disease, gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, dermatitis herpetiformis, psoriasis, pemphigus vulgaris, vitiligo (leukoderma), allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, chronic obstructive pulmonary disease, and transplantation associated diseases including graft rejection and graft-versus-host-disease.

[0247] For example, in one aspect, the anti-NKG2D antibody is used in combination with one or more other anti-inflammatory agents, including, but not limited to, analgesic agents, immunosuppressive agents (e.g., B- or T-cell antagonists such as B-cell depletion agents and T cell inhibiting agents; complement inhibiting agents), corticosteroids, and anti-TNFalpha agents or other anti-cytokine or anti-cytokine receptor agents, and anti-angiogenic agents. Specific examples include methotrexate, TSG-6, Rituxan® or other B-cell therapies, anti-IL12 (p40) antibodies, CTLA4-Fc fusion proteins, IL-1-receptor antagonists, IL-1 antibodies, IL-15 antibodies, IL-18 antibodies, and anti-IL6R antibodies. Further examples of combination therapies are provided below.

[0248] When one or more other agents or approaches are used in combination with the present therapy, there is no requirement for the combined results to be additive of the effects observed when each treatment is conducted separately. Although at least additive effects are generally desirable, any decrease in NKG2D activity or other beneficial effect above one of the single therapies would be of benefit. Also, there is no particular requirement for the combined treatment to exhibit synergistic effects, although this is certainly possible and advantageous. The NKG2D-based treatment may precede, or follow, the other treatment by, e.g., intervals ranging from minutes to weeks and months. It also is envisioned that more than one administration of either the anti-NKG2D composition or the other agent will be utilized. The agents may be administered interchangeably, on alternate days or weeks; or a cycle of anti-NKG2D treatment may be given, followed by a cycle of the other agent therapy. In any event, all that is required is to deliver both agents in a combined amount effective to exert a therapeutically beneficial effect, irrespective of the times for administration.

[0249] The following describes some selected inflammatory and/or autoimmune diseases or disorders for which anti-hNKG2D antibodies of the invention can be used as therapeutic agents. Preferably, the anti-hNKG2D antibody as defined in the claims is full-length bivalent MS, or an antigen-binding fragment, variant or derivative thereof.

Rheumatoid arthritis

[0250] Rheumatoid arthritis (RA) is a chronic systemic autoimmune inflammatory disease that mainly involves the synovial membrane of multiple joints with resultant injury to the articular cartilage. The pathogenesis is T lymphocyte dependent and is associated with the production of rheumatoid factors, auto antibodies directed against self IgG, with the resultant formation of immune complexes that attain high levels in joint fluid and blood. These complexes in the joint may induce the marked infiltrate of lymphocytes and monocytes into the synovium and subsequent marked synovial changes; the joint space is infiltrated by similar cells with the addition of numerous neutrophils. The pathological T cell produces cytokines and other soluble factors adding to the attraction and activation of other cells, and to the destruction of the tissue. Tissues affected are primarily the joints, often in symmetrical pattern. However, extra-articular disease also occurs in two major forms. One form is the development of extra-articular lesions with ongoing progressive joint disease and typical lesions of pulmonary fibrosis, vasculitis, and cutaneous ulcers. The second form of extra-articular disease is the so called Felty's syndrome which occurs late in the RA disease course, sometimes after joint disease has become quiescent, and involves the presence of neutropenia, thrombocytopenia and splenomegaly. This can be accompanied by vasculitis in multiple organs with formations of infarcts, skin ulcers and gangrene. Patients often also develop rheumatoid nodules in the subcutis tissue overlying affected joints; the nodules late stage have necrotic centers surrounded by a mixed inflammatory cell infiltrate. Other manifestations which can occur in RA include: pericarditis, pleuritis, coronary arteritis, interstitial pneumonitis with pulmonary fibrosis, keratoconjunctivitis sicca, and rheumatoid nodules.

[0251] Accordingly, in one aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for use in treating and/or preventing rheumatoid arthritis (RA). The use of a human or humanized anti-hNKG2D antibody as defined in the claims comprises delivering an effective amount of an anti-hNKG2D antibody to a patient having RA or being identified/diagnosed as being at substantial risk of developing RA, such that RA is treated or prevented. In one aspect, the anti-NKG2D antibody is demonstrated to be effective in ameliorating RA in an acceptable model of RA, such as is described in US Patent No. 6414218 and US Patent Publication No. 20030005469 (related principles and models are described in, e.g., Wooley, P. H., *Animal Models of Arthritis*, eds. J. H. Kippel and P. A. Dieppe, Mosby Publishers (London), 1998 ; Erning et al., *Arthritis Res*, 4 Suppl 3:S 133-40, 2002; Holmdahl et al., *Ageing Res Rev*, 1(1): 135-47, 2002; Anthony et al., *Clin Exp Rheumatol*, 17(2) :240-

4,1999; Durie et al., Clin Immunol Immunopathol, 73(1):11-8, 1994; and Muller-Ladner et al., Drugs Today (Bare), 35(4-5):379-88, 1999). In a further aspect, the antibody that is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expressing leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells (e.g., impairing the expansion and/or function of autoreactive CD8+ T cells) (in contrast to, e.g., at least some of the antibodies described in US Patent Publication No. 20040115198), without significantly depleting such cells (e.g., causing a reduction of about 10% or less of such cells as compared to a suitable control). In one aspect, the method results in a modulation of one or more biomarkers in a manner consistent with the treatment or prevention (as applicable) of RA (e.g., serum IL-6, TNF- α , IL-1, VEGF, TIFR, IL-2R, shed CD4, shed CD8, and/or C reactive protein). In another aspect, the practice of the method results in a detectable reduction of synovial inflammation in the peripheral joints of the patient/host. In one aspect, the method results in preventing radiographic deterioration and improving physical function in the patient or host as exhibited by, e.g., a reduction in radiographic progression in the patient or host, reduction in swollen and tender joints (as determined by acceptable analytical criteria), and/or significantly improved quality of life (e.g., as determined by a reduction in disability scores on the RA Health Assessment Questionnaire). The antibody can be used alone or in combination with one or more other anti-RA agent, such as a non-steroidal anti-inflammatory drug (NSAID), a COX-2 inhibitor, an analgesic, a corticosteroid (e.g., prednisone, hydrocortisone), gold, an immunosuppressant (e.g., methotrexate), a B-cell depletion agent (e.g., Rituxan®), a B-cell agonist (e.g., LymphoStat-B®) and an anti-TNF α agent (e.g., Embrel®, Humira® and Remicade®), an anti-IL1 receptor antagonist (e.g., Kineret®), an anti-IL-15 antibody, or a disease-modifying anti-rheumatic drug (DMARD).

Demyelinating diseases

[0252] Demyelinating diseases of the central and peripheral nervous systems, including Multiple Sclerosis (MS); idiopathic demyelinating polyneuropathy or Guillain-Barre syndrome; and Chronic Inflammatory Demyelinating Polyneuropathy, are believed to have an autoimmune basis and result in nerve demyelination as a result of damage caused to oligodendrocytes or to myelin directly. In MS there is evidence to suggest that disease induction and progression is dependent on T lymphocytes. MS is a demyelinating disease that is T lymphocyte-dependent and has either a relapsing-remitting course or a chronic progressive course. The etiology is unknown; however, viral infections, genetic predisposition, environment, and autoimmunity all contribute. Lesions contain infiltrates of predominantly T lymphocyte mediated, microglial cells and infiltrating macrophages; CD4+ T lymphocytes are the predominant cell type at lesions.

[0253] Thus, in another aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims treating and/or preventing MS. The use of a human or humanized anti-hNKG2D antibody as defined in the claims comprises delivering an effective amount of an anti-hNKG2D antibody to a human patient having MS or being identified/diagnosed as being at substantial risk of developing MS, such that MS is treated or prevented in the patient or host. In a particular aspect, the anti-NKG2D monoclonal antibody is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expression leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells, without significantly depleting such cells. The antibody can be used alone or in combination with other anti-MS agents such as Tyzabri®.

Inflammatory bowel disease

[0254] In CD8+ T cells in the intestine, NKG2D acts as a co-stimulator of CD28+ cells (Roberts et al., J Immunol 2001;167:5527-30). Furthermore, in inflammation of the intestine (celiac disease), NKG2D is upregulated, and intestinal epithelial lymphocytes are stimulated via NKG2D to kill and produce cytokines (Hue et al., Immunity 2004;21:367-77; Meresse et al., Immunity 2004;21:357-66). Additionally, IL-15, often found during intestinal inflammation, up-regulates NKG2D on intestinal epithelial lymphocytes (Roberts et al., J Immunology 2001;167:5527-30). Furthermore, during intestinal inflammation, MICA, a ligand for NKG2D, is upregulated (Meresse et al, *supra*, Hue et al., *supra*). NKG2D is also upregulated on pro-inflammatory lymphocytes of patients with Crohn's disease (Allez et al., presentation at 16th European Congress of Immunology (ECI2006), September 6-9, 2006; Paris, France).

[0255] Thus, in another aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for treating and/or preventing inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis. As shown by Kjellef et al. (Eur J Immunol 2008;37:1397-1406), inhibition of NKG2D-receptor function by early antibody therapy attenuated transfer-induced colitis in SCID mice, an animal model of colitis.

[0256] The use of a human or humanized anti-hNKG2D antibody as defined in the claims for treating an inflammatory bowel disease comprises delivering an effective amount of an anti-NKG2D antibody to a human patient having IBD or being

identified/diagnosed as being at substantial risk of developing IBD, such that IBD is treated or prevented in the patient. In a particular aspect, the inventive use of a human or humanized anti-hNKG2D antibody as defined in the claims for IBD treatment/prevention is practiced by use of an anti-NKG2D monoclonal antibody that is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expressing leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells, without significantly depleting such cells. The antibody can be used alone or in combination with other anti-IBD agents, such as drugs containing mesalamine (including sulfasalazine and other agents containing 5-aminosalicylic acid (5-ASA), such as olsalazine and balsalazide), non-steroidal anti-inflammatory drugs (NSAIDs), analgesics, corticosteroids (e.g., prednisone, hydrocortisone), TNF-inhibitors (including adalimumab (Humira®, etanercept (Enbrel® and infliximab (Remicade®), anti-IL12 antibodies, immunosuppressants (such as 6-mercaptopurine, azathioprine and cyclosporine A), and antibiotics.

Psoriasis

[0257] Psoriasis is a T lymphocyte-mediated inflammatory disease. Lesions contain infiltrates of T lymphocytes, macrophages and antigen processing cells, and some neutrophils.

[0258] Thus, in another aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for treating and/or preventing psoriasis. The use of a human or humanized anti-hNKG2D antibody as defined in the claims comprises delivering an effective amount of an anti-hNKG2D antibody to a human patient having psoriasis or being identified/diagnosed as being at substantial risk of developing psoriasis, such that psoriasis is treated or prevented in the patient. In a more particular aspect, the agent is an anti-NKG2D monoclonal antibody that is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expressing leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells, without significantly depleting such cells. The antibody can be used alone or in combination with one or more other anti-psoriasis treatments such as phototherapy, topical therapy (e.g., tar, topical glucocorticoids), or systemic therapy (e.g., methotrexate, a synthetic retinoid, cyclosporine), an anti-TNFalpha agent (e.g., Embrel®, Humira®, Remicade®), a T-cell inhibitor (e.g., Raptiva®), vitamin D analogs, p38 mitogen-activated protein kinase (MAPK) inhibitors, as well as a biologic agent such as Rituxan®.

Psoriatic arthritis

[0259] Psoriatic arthritis is a chronic inflammatory arthritic condition affecting the skin, the joints, the insertion sites of tendons, ligaments, and fascia, and is commonly associated with psoriasis. (approximately 7% of patients with psoriasis develop psoriatic arthritis). Much evidence suggests that a T-cell-mediated process drives the pathophysiology of psoriatic arthritis. Monocytes also play a role in psoriatic arthritis and are responsible for the production of matrix metalloproteinases, which may mediate the destructive changes in the joints of patients with psoriatic arthritis. Furthermore, NK cells are also found in affected joints, suggesting a role in the disease pathology.

[0260] Thus, in another aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for treating and/or preventing psoriatic arthritis. The use of a human or humanized anti-hNKG2D antibody as defined in the claims comprises delivering an effective amount of an anti-hNKG2D antibody to a human patient having psoriatic arthritis or being identified/diagnosed as being at substantial risk of developing psoriatic arthritis, such that the psoriatic arthritis is treated or prevented in the patient. In a more particular aspect, the agent is an anti-NKG2D monoclonal antibody that is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expressing leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells, without significantly depleting such cells. The antibody can be used alone or in combination with one or more other anti-psoriatic arthritis treatments such as nonsteroidal anti-inflammatory drugs (aspirin, ibuprofen), methotrexate, a synthetic retinoid, cyclosporine, a corticosteroid, an anti-TNFalpha agent (e.g., Embrel®, Humira®, Remicade®).

Systemic lupus erythematosus

[0261] In systemic lupus erythematosus (SLE), the central mediator of disease is the production of auto-reactive antibodies to self proteins/tissues and the subsequent generation of immune-mediated inflammation. Antibodies either directly or indirectly mediate tissue injury. Though T lymphocytes have not been shown to be directly involved in tissue damage, T lymphocytes are required for the development of auto-reactive antibodies. The genesis of the disease is thus T lymphocyte dependent. Multiple organs and systems are affected clinically including kidney, lung, musculoskeletal system, mucocutaneous, eye, central nervous system, cardiovascular system, gastrointestinal tract, bone marrow and blood.

[0262] Thus, in another aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for treating and/or preventing SLE. The use of a human or humanized anti-hNKG2D antibody as defined in the claims comprises delivering an effective amount of an anti-hNKG2D antibody to a human patient having SLE or being identified/diagnosed as being at substantial risk of developing SLE, such that the SLE is treated or prevented in the patient. In a more particular aspect, the agent is an anti-NKG2D monoclonal antibody that is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expressing leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells, without significantly depleting such cells. The antibody can be used alone or in combination with other anti-SLE agents, such as non-steroidal anti-inflammatory drugs (NSAIDs), analgesics, corticosteroids (*e.g.*, prednisone, hydrocortisone), immunosuppressants (such as cyclophosphamide, azathioprine, and methotrexate), antimalarials (such as hydroxychloroquine) and biologic drugs that inhibit the production of dsDNA antibodies (*e.g.* LIP 394).

Diabetes

[0263] Type I diabetes mellitus or insulin-dependent diabetes is the autoimmune destruction of pancreatic islet B cells; this destruction is mediated by auto-antibodies and auto-reactive T cells. Antibodies to insulin or the insulin receptor can also produce the phenotype of insulin-non-responsiveness.

[0264] Thus, in another aspect, an anti-NKG2D antibody is delivered to a patient suffering from or at substantial risk of developing type I diabetes mellitus in an amount and under conditions sufficient to treat or prevent the condition in the patient. The antibody can be used alone or in combination with other anti-diabetic agents, such as insulin, or beta cell growth or survival factors, or immunomodulatory antibodies such as anti-CD3 antibodies.

Transplantation

[0265] Transplantation associated diseases, including graft rejection and Graft-Versus-Host-Disease (GVHD) are T lymphocyte-dependent; inhibition of T lymphocyte function is ameliorative.

[0266] Thus, in another aspect, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for reducing the likelihood of transplant rejection (or reducing the severity or prolonging the time to onset of a transplant rejection-related condition, *i.e.*, to prolong allograft survival). The use of a human or humanized anti-hNKG2D antibody as defined in the claims comprising delivering an effective amount of an anti-hNKG2D antibody to a human patient that is about to be, is, or recently was the recipient of a tissue/organ transplant, such that the likelihood of rejection is detectably reduced (*e.g.*, as compared to a control). In a particular aspect, the anti-NKG2D antibody is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expression leukocytes and/or impairing expansion of NKG2D+ T cells or NK cells, without significantly depleting such cells. Examples of tissue transplants that can be treated include, but are not limited to, liver, lung, kidney, heart, small bowel, and pancreatic islet cells, as well as in bone marrow-transplantation and in the treatment of graft versus host disease (GVHD). The antibody can be used alone or in combination with other agents for inhibiting transplant rejection, such as immunosuppressive agents (*e.g.* cyclosporine, azathioprine, methylprednisolone, prednisolone, prednisone, mycophenolate mofetil, sirolimus, rapamycin, tacrolimus), anti-infective agents (*e.g.*, acyclovir, clotrimazole, ganciclovir, nystatin, trimethoprim-sulfamethoxazole), diuretics (*e.g.* bumetanide, furosemide, metolazone) and ulcer medications (*e.g.*, cimetidine, famotidine, lansoprazole, omeprazole, ranitidine, sucralfate). For hematopoietic transplantation, hematopoietic growth factor(s) (*e.g.*, erythropoietin, G-CSF, GM-CSF, IL-3, IL-11, thrombopoietin, etc.) or antimicrobial(s) (*e.g.*, antibiotic, antiviral, antifungal) may be administered as an adjunct therapy.

Other autoimmune or inflammatory diseases

[0267] In other separate aspects, the invention provides the use of a human or humanized anti-hNKG2D antibody as defined in the claims for treating and/or preventing other autoimmune or inflammatory diseases or disorders, comprising delivering an effective amount of an anti-hNKG2D antibody to a human patient having the disease or disorder or being identified/diagnosed as being at substantial risk of developing the disease or disorder, such that it is treated or prevented in the patient, where the disease or disorder is one described below. In a more particular aspect, the agent is an anti-NKG2D monoclonal antibody that is capable of detectably reducing ligand-induced NKG2D activation of NKG2D-expressing leukocytes and/or impairing expansion of

NKG2D+ T cells or NK cells, without significantly depleting such cells. The antibody can be used alone or in combination with one or more other therapeutic agents used for treating the disease or disorder.

[0268] Juvenile chronic arthritis is a chronic idiopathic inflammatory disease which begins often at less than 16 years of age. Its phenotype has some similarities to RA; some patients which are rheumatoid factor positive are classified as juvenile rheumatoid arthritis. The disease is sub-classified into three major categories: pauciarticular, polyarticular, and systemic. The arthritis can be severe and is typically destructive and leads to joint ankylosis and retarded growth. Other manifestations can include chronic anterior uveitis and systemic amyloidosis.

[0269] Spondyloarthropathies are a group of disorders with some common clinical features and the common association with the expression of HLA-B27 gene product. The disorders include: ankylosing spondylitis, Reiter's syndrome (reactive arthritis), arthritis associated with inflammatory bowel disease, spondylitis associated with psoriasis, juvenile onset spondyloarthropathy and undifferentiated spondyloarthropathy. Distinguishing features include sacroileitis with or without spondylitis; inflammatory asymmetric arthritis; association with HLA-B27 (a serologically defined allele of the HLA-B locus of class I MHC); ocular inflammation, and absence of autoantibodies associated with other rheumatoid disease. The cell most implicated as key to induction of the disease is the CD8+ T lymphocyte, a cell which targets antigen presented by class I MHC molecules. CD8+ T cells may react against the class I MHC allele HLA B27 as if it were a foreign peptide expressed by MHC class I molecules. It has been hypothesized that an epitope of HLA-B27 may mimic a bacterial or other microbial antigenic epitope and thus induce a CD8+ T cells response.

[0270] Systemic sclerosis (scleroderma) has an unknown etiology. A hallmark of the disease is induration of the skin; likely this is induced by an active inflammatory process. Scleroderma can be localized or systemic; vascular lesions are common and endothelial cell injury in the microvasculature is an early and important event in the development of systemic sclerosis; the vascular injury may be immune mediated. An immunologic basis is implied by the presence of mononuclear cell infiltrates in the cutaneous lesions and the presence of anti-nuclear antibodies in many patients. ICAM-1 is often unregulated on the cell surface of fibroblasts in skin lesions suggesting that T cell interaction with these cells may have a role in the pathogenesis of the disease. Other organs involved include: the gastrointestinal tract: smooth muscle atrophy and fibrosis resulting in abnormal peristalsis/motility; kidney: concentric subendothelial intimal proliferation affecting small arcuate and interlobular arteries with resultant reduced renal cortical blood flow, results in proteinuria, azotemia and hypertension; skeletal muscle: atrophy, interstitial fibrosis; inflammation; lung: interstitial pneumonitis and interstitial fibrosis; and heart: contraction band necrosis, scarring/fibrosis.

[0271] Idiopathic inflammatory myopathies including dermatomyositis, polymyositis and others are disorders of chronic muscle inflammation of unknown etiology resulting in muscle weakness. Muscle injury/inflammation is often symmetric and progressive. Autoantibodies are associated with most forms. These myositis-specific autoantibodies are directed against and inhibit the function of components, proteins and RNA's, involved in protein synthesis.

[0272] Sjogren's syndrome is due to immune-mediated inflammation and subsequent functional destruction of the tear glands and salivary glands. The disease can be associated with or accompanied by inflammatory connective tissue diseases. The disease is associated with autoantibody production against Ro and La antigens, both of which are small RNA-protein complexes. Lesions result in keratoconjunctivitis sicca, xerostomia, with other manifestations or associations including biliary cirrhosis, peripheral or sensory neuropathy, and palpable purpura.

[0273] Systemic vasculitis are diseases in which the primary lesion is inflammation and subsequent damage to blood vessels which results in ischemia/necrosis/degeneration to tissues supplied by the affected vessels and eventual end-organ dysfunction in some cases. Vasculitides can also occur as a secondary lesion or sequelae to other immune-inflammatory mediated diseases such as rheumatoid arthritis, systemic sclerosis, etc., particularly in diseases also associated with the formation of immune complexes. Diseases in the primary systemic vasculitis group include: systemic necrotizing vasculitis: polyarteritis nodosa, allergic angitis and granulomatosis, polyangiitis; Wegener's granulomatosis; lymphomatoid granulomatosis; and giant cell arteritis. Miscellaneous vasculitides include: mucocutaneous lymph node syndrome (MLNS or Kawasaki's disease), isolated CNS vasculitis, Behet's disease, thromboangiitis obliterans (Buerger's disease) and cutaneous necrotizing vasculitis. The pathogenic mechanism of most of the types of vasculitis listed is believed to be primarily due to the deposition of immunoglobulin complexes in the vessel wall and subsequent induction of an inflammatory response either via ADCC, complement activation, or both.

[0274] Sarcoidosis is a condition of unknown etiology which is characterized by the presence of epithelioid granulomas in nearly any tissue in the body; involvement of the lung is most common. The pathogenesis involves the persistence of activated macrophages and lymphoid cells at sites of the disease with subsequent chronic sequelae resultant from the release of locally and systemically active products released by these cell types.

[0275] Autoimmune hemolytic anemia including autoimmune hemolytic anemia, immune pancytopenia, and paroxysmal nocturnal hemoglobinuria is a result of production of antibodies that react with antigens expressed on the surface of red blood cells (and in some cases other blood cells including platelets as well) and is a reflection of the removal of those antibody coated cells via complement mediated lysis and/or ADCC/Fc-receptor-mediated mechanisms.

[0276] In autoimmune thrombocytopenia including thrombocytopenic purpura, and immune-mediated thrombocytopenia in other clinical settings, platelet destruction/removal occurs as a result of either antibody or complement attaching to platelets and subsequent removal by complement lysis, ADCC or Fc-receptor mediated mechanisms.

[0277] Thyroiditis including Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, and atrophic thyroiditis, are the result of an autoimmune response against thyroid antigens with production of antibodies that react with proteins present in and often specific for the thyroid gland. Experimental models exist including spontaneous models: rats (BUF and BB rats) and chickens (obese chicken strain); inducible models: immunization of animals with either thyroglobulin, thyroid microsomal antigen (thyroid peroxidase).

[0278] Immune mediated renal diseases, including glomerulonephritis and tubulointerstitial nephritis, are the result of antibody or T lymphocyte mediated injury to renal tissue either directly as a result of the production of autoreactive antibodies or T cells against renal antigens or indirectly as a result of the deposition of antibodies and/or immune complexes in the kidney that are reactive against other, non-renal antigens. Thus other immune-mediated diseases that result in the formation of immune-complexes can also induce immune mediated renal disease as an indirect sequelae. Both direct and indirect immune mechanisms result in inflammatory response that produces/induces lesion development in renal tissues with resultant organ function impairment and in some cases progression to renal failure. Both humoral and cellular immune mechanisms can be involved in the pathogenesis of lesions.

[0279] Inflammatory and Fibrotic Lung Disease, including Eosinophilic Pneumonias; Idiopathic Pulmonary Fibrosis, and Hypersensitivity Pneumonitis may involve a dysregulated immune-inflammatory response. Inhibition of that response would be of therapeutic benefit.

[0280] Autoimmune or Immune-mediated Skin Disease including Bullous Skin Diseases, Erythema Multiforme, and Contact Dermatitis are mediated by auto-antibodies, the genesis of which is T lymphocyte dependent.

[0281] Allergic diseases, including asthma; allergic rhinitis; atopic dermatitis; food hypersensitivity; and urticaria are T lymphocyte dependent. These diseases are predominantly mediated by T lymphocyte induced inflammation, IgE mediated-inflammation or a combination of both.

[0282] Another disease suitable for treatment with a human anti-NKG2D antibody is viral hepatitis, as shown in WO2007130642 and by Chen et al. (Hepatology, Vol. 46 (3) pp. 706-715 (2007)).

[0283] It will be understood that the effective amount of the NKG2D modulator, as well as the overall dosage regimen, may vary according to the disease and the patient's clinical status, which, in turn, may be reflected in one or more clinical parameters such as clinically accepted disease scores. For example, for rheumatoid arthritis, the severity of disease and/or outcome of treatment, may be evaluated by monitoring number of swollen joints; pain; mobility; and/or the official disease score ACR 20/50 or 70. For Type 1 diabetes, severity of disease and/or outcome of treatment may be evaluated by measuring blood glucose levels or variations thereof, Hb1c levels, the amount of insulin needed, and the like. For multiple sclerosis, brain inflammation can be assessed through scanning the brain. For hematopoietic transplant rejection, severity of the disease (failure to engraft) and/or outcome of treatment may be evaluated by evidence of prolonged neutropenia, thrombocytopenia, and red-cell transfusion dependence in patients that have undergone myeloablative conditioning, and by failure to observe chimerism in patients that have undergone non-myeloablative conditioning. In general, detectable effects on treatment outcome using the methods and compositions of the present invention include a decrease in the necessity for other treatments (including, e.g., a decrease in the amount and/or duration of other drugs or treatments), a decrease in number and/or duration of hospital stays, a decrease in lost work days due to illness, and the like. It will be further understood that the effective amount may be determined by those of ordinary skill in the art by routine experimentation, by constructing a matrix of values and testing different points in the matrix.

Dosages

[0284] For administration of the antibody, the dosage ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 5

mg/kg, of the host body weight. For example, dosages can be about 0.3 mg/kg body weight, about 1 mg/kg body weight, about 3 mg/kg body weight, about 5 mg/kg body weight or about 10 mg/kg body weight or within the range of 1-10 mg/kg. An exemplary treatment regime entails administration twice per week, once per week, once every two weeks, once every three weeks, once every four weeks, once a month, once every 3 months or once every three to 6 months. Preferred dosage regimens for an anti-hNKG2D antibody of the invention include about 1, 3 or 10 mg/kg body weight via intravenous administration or subcutaneous injection, with the antibody being given using one of the following dosing schedules: (i) loading doses every 1-3 weeks for 2-4 dosages, then every two; months (ii) every four weeks; (iii) every week, or any other optimal dosing. In some methods, two or more monoclonal antibodies with different binding specificities are administered simultaneously, in which case the dosage of each antibody administered falls within the ranges indicated. Antibody is usually administered on multiple occasions. Intervals between single dosages can be, for example, weekly, monthly, every three months or yearly. Intervals can also be irregular as indicated by measuring blood levels of antibody to the target antigen in the patient. In some methods, dosage is adjusted to achieve a plasma antibody concentration of about 1-1000 µg /ml and in some methods about 25-300 µg/ml. Alternatively, antibody can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, human antibodies show the longest half life, followed by humanized antibodies, chimeric antibodies, and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or non-prophylactic (e.g., palliative or curative). In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In palliative or curative applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

[0285] The appropriate doses of anti-inflammatory agents will approximate those already employed in clinical therapies wherein the anti-inflammatory agents are administered alone or in combination with other agents. Variation in dosage will likely occur depending on the condition being treated. The physician administering treatment will be able to determine the appropriate dose for the individual subject

Articles of manufacture

[0286] In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above is provided. For example, the article of manufacture can comprise a container containing a human or humanized anti-hNKG2D antibody as defined in the claims together with instructions directing a user to treat a disorder such as an autoimmune or inflammatory disease or disorder in a human with the antibody in an effective amount. The article of manufacture typically comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is the human or humanized anti-hNKG2D antibody as defined in the claims, or an antigen-binding fragment or antibody derivative (e.g., an immunoconjugate) comprising such an antibody. The label or package insert indicates that the composition is used for treating the condition of choice, such as, e.g., rheumatoid arthritis.

[0287] Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises the human or humanized antibody and (b) a second container with a composition contained therein, wherein the composition comprises a therapeutic agent other than the human or humanized antibody. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the first and second compositions can be used in combination to treat an autoimmune or inflammatory disease or disorder. Such therapeutic agents may be any of the adjunct therapies described in the preceding section. Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically acceptable buffer, such as bacteriostatic water for injection (BWFJ), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

EXAMPLES

[0288] Further details of the invention are illustrated by the following non-limiting Examples, in which the antibody MS according to the invention as well as various reference antibodies were used.

Example 1: Generation and initial screening of human monoclonal antibodies against hNKG2D**Materials and Methods**

[0289] Antigen. Soluble NKG2D-hFc fusion protein (R&D, cat: 1299-NK) or NKG2D expressed on the surface of cells (NK, BAF, or CHO) were used as antigens for immunization. The BAF cells were co-transfected with full-length NKG2D and DAP10. The CHO cells were transfected with an NKG2D point mutant that transports to the cell surface without DAP10 (Wu et al., Science 1999;385:730-2). The NK cells were primary NK cells naturally expressing NKG2D.

[0290] Mice. Fully human monoclonal antibodies against NKG2D were produced in the KM mouse™ strain of transgenic mice that express human antibody genes (PCT publication WO 02/43478 to Ishida et al.). In this mouse strain, the endogenous kappa light chain gene has been homozygously disrupted as described in Chen et al (1993) EMBO J. 12:811-820, and the endogenous mouse heavy chain has been homozygously disrupted as described in Example 1 of PCT Publication WO 01/09187 for Humab mice. The mouse strain carries a human kappa light chain transgene, KC05, as described in Fishwild et al (1996) Nature Biotechnology 14:845-851. The mouse strain also carries a human heavy chain transchromosome, SC20, as described in WO0243478.

[0291] Immunizations. In a first series of immunizations, animals were immunized intraperitoneally with alternating injections of NKG2D-transfected BAF cells and NKG2D-transfected CHO cells, or primary human NK cells with or without any adjuvant. Each mouse was immunized IP with 5×10^6 cells every or every other week (6 times in total). The mice were boosted with 5×10^6 NKG2D-transfected BAF cells intravenously 3 and 2 days before sacrifice and removal of the spleen. The animal experiments were performed according to Danish National Research Council guidelines.

[0292] In a second series of immunizations, animals were immunized intraperitoneally and in the foot path with NKG2D-hFc with different adjuvant. Each mouse was immunized 7 x 25ug NKG2F-hFc/Ribi/ip/sc, 1 x 25ug NKG2D-hFc/CFA/ip/sc, 1 x 25ug NKG2D-hFc/IFA/ip/sc, 1 x 30ug anti-CTLA4 + 40ug NKG2D-hFc/IFA/ip/sc, 1 x 25ug NKG2D-hFc/Ribi/ip/sc and boosted 2 x 30ug/PBS/ip/iv 3 and 2 days before sacrifice and removal of the spleen. The animal experiments were performed according to American National Research Council guidelines.

[0293] Screening of Mouse Sera. The sera from the immunized mice were screened by flow cytometry analysis for NKG2D-specificity and selected sera were also tested for their ability to neutralize binding of the MICA ligand, as described in Example 3. Mice that had generated high titers of antibodies that specifically bound NKG2D and neutralized MICA binding were selected for hybridoma production.

[0294] Generation of Hybridomas. The spleen from each selected immunized mouse was homogenised and a single cell suspension of splenocytes used for fusion to X61 Ag8653 myeloma cells (ATCC, CRL 1580). The fusions were performed using polyethyleneglycol. (PEG) 1500 as previously described (Harlow and Lane, ANTIBODIES: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1988)) and electrofusion using the The Cyto Pulse™ CEEF-50 Electrofusion System (Cyto Pulse Sciences, Inc.).

[0295] The fused cells were initially seeded in 96-well tissue culture plates in selective DMEM HAT medium, supplemented with 10% FBS and 5% origin (Hybridoma cloning Factor, BioVeris). The plates were incubated for 10-14 days with 1-2 medium changes, respectively, to DMEM HT medium supplemented with 5% FBS and 0.7% origin, before harvest and screening of the supernatants. Clones tested positive were expanded and subcloned by limiting dilution until stable clones had been generated. The selected clones were continuously screened for the presence of anti-NKG2D specific antibodies by FACS analysis as well as for their ability to neutralize MICA binding.

[0296] Screening of Hybridoma Supernatants. The primary screening of the hybridoma supernatants from the first series of immunizations was performed using direct ELISA or flow cytometry analysis (FACS) to test for the presence of anti-NKG2D specific antibodies. Briefly, the ELISA was performed by coating maxisorp plates with 50 µl 0.4 µg/ml mFc-NKG2D (comprising the extracellular portion of NKG2D fused to murine Fc and expressed in CHO cells) overnight in PBS at 4°C. followed by blocking with PBS, 0.05% Tween 20, for 15 min at room temperature. The plates were subsequently incubated with 50 µl hybridoma supernatant, and NKG2D-specific antibodies detected using Goat-Anti-human IgG-HRP Fcy Fragment specific (Jackson, 109-036-098). These incubations were performed for 1 hr at room temperature, and between each step the plates were washed with

PBS, 0.05% Tween 20. Bound antibodies were visualized using 100µl TMB substrate (Kem-En-Tec), and stopped with 4M H₃PO₄. The plates were read at 450 and 620 nm. For FACS, binding to NKG2D-expressing BaF/3 cells and control BaF/3 cells not expressing NKG2D was analyzed by incubation of 50000 cells in 10 µl with 90 µl hybridoma supernatant for 30 min at 4°C, followed by washing with PBS with 2% FCS, and subsequently incubated with secondary Goat-Anti-human IgG-HRP Fcγ Fragment specific (Jackson, 109-036-098). The cells were then analysed on a B&D FACSArray (BD Biosciences). Antibodies that only stained NKG2D-expressing BaF/3 cells and not control cells were deemed NKG2D-specific.

[0297] The primary screen for the second series of immunizations was a direct ELISA to test for the presence of anti-NKG2D specific antibodies. Briefly, the ELISA was performed by coating maxisorp plates with 1-2 mg/ml hFc-NKG2D (R&D Systems) overnight in PBS at 4°C, followed by blocking with PBS, 0.05% Tween 20, 5% chicken serum for 30-60 min at room temperature. The plates were subsequently incubated with 50 µl hybridoma supernatant and 50 µl blocking buffer, and NKG2D-specific antibodies detected using Anti-human IgG-HRP (Bethyl, A80-115P) in blocking buffer. These incubations were performed for 1 hr at room temperature, and between each step the plates were washed with PBS, 0.05% Tween 20. Bound antibodies were visualized using ABTS substrate (Moss Inc, product: ABTS-1000). The plates were read at 415 nm with Molecular Devices Software.

[0298] Hybridomas selected from an ELISA primary screen were subjected to a secondary screen using FACS, as described above.

[0299] Commercially available murine antibodies (149810 and ON72) were used as controls.

Results

[0300] Highly selective sera from immunized mice were identified by NKG2D-binding and ligand blocking ability (exemplary results shown in Figure 1A and 1B), and selected mice were used for fusion and hybridoma generation. About 2500 hybridomas were screened by ELISA and flow cytometry and NKG2D-specific clones identified. Figure 2 shows that human antibody in a hybridoma supernatant bound to NKG2D-expressing cells but not NKG2D-negative cells, comparing to a commercial antibody (149810). Antibodies from three hybridomas (16F16, 16F31 and 21 F2) from the first series of immunization, and several antibodies from the second series of immunizations (including MS), were selected for recombinant production and further testing.

Example 2: Recombinant production and sequencing

[0301] A second batch of several hundreds of hybridomas from fusions mice spleens expressing human antibodies were obtained from a separate round of immunization(s). These were screened for NKG2D-specificity using FACS in the same manner as described in Example 1. Antibodies from one hybridoma, MS, were selected for recombinant production and further testing.

[0302] The variable regions of the heavy and light chains of the antibodies were identified by PCR and subsequent sequencing of the isolated product, of mRNA from the hybridoma.

Materials and Methods

[0303] *RNA purification.* Total RNA was purified using RNeasy from Qiagen according to the manufactures instructions, except that β-mercaptoethanol was omitted from the procedure. The quality of the RNA was checked by light spectroscopy (260/280 nm, 1.8 < ratio < 2.0) and occasionally RNA degradation was evaluated using a bioanalyser.

[0304] *RT-PCR.* Full length cDNA was synthesised by SMART-RACE (kit from Clontech).

[0305] *PCR.* PCR was performed with the HFII polymerase from Clontech. The 5' primer (with EcoRI) annealed to a conserved sequence introduced during SMART-RACE. Two 3' primers were designed that anneal to conserved regions of the IgG (VH) and kappa chains (VL), respectively. Restriction sites were also present in the 3' primers (BsiWI (VL) and NheI (VH)). The PCR was performed in duplicate (to check for PCR introduced mutations) for all VH and VL amplifications. If the PCR reaction failed, the VL and VH were amplified using a degenerate 5' primer mix from Novagen.

[0306] *PCR product purification.* The PCR product (~ 550 bp) was separated on a 1 % agarose gel, excised, purified on GFX columns (from Amersham) and eluted in DNase free water.

[0307] *Ligation.* The PCR products and the expression vector (ampicillin resistance) were cut with appropriate restriction enzymes (VH, EcoRI + NheI and VL, EcoRI + BsiWI). The ligation of the variable domains into the isotype-dictating vector (IgG4 for NKG2D) was catalyzed by the T4-ligase (Roche). The plasmid used was pTT5 (Durocher et al., Nucleic Acids Res 2002;30(2):e9; Pham et al., Biotechnol Bioeng 2003;84(3):332-42).

[0308] *Check of insert in the expression vector (colony PCR).* Competent *E. coli* (Top10) were transformed with the ligation mix and ampicillin resistant clones were selected overnight. In total, 8 positive colonies for both VH and VL were picked. Via colony PCR and gel electrophoresis (1% agarose), all colonies were checked for inserts matching the expected size.

[0309] *Sequencing/miniprep.* An aliquot from all positive colony PCRs was prepared for sequencing (using ExoSAPit). In total, 32 PCR products were sequenced for each clone ((8*VH +8*VL)*2 (PCR in duplicate)). The sequences were analysed (using VectorNTI) and positive bacteria clones corresponding to the cloned VH and VL were up-scaled (mini/maxiprep), and the DNA purified for HEK293/6E transfection (GFX columns). If more than one VH and VL sequence was identified, then all possible VL and VH combinations were expressed in HEK293/6E cells.

[0310] *Recombinant production.* The identified variable regions of heavy and light chains were inserted into heavy and light chain human IgG4 framework respectively and expressed from two vectors in HEK293 cells at a high level. The antibodies were purified on a protein A column.

[0311] *Antibody expression in HEK293/6E cells.* HEK293 cells were passaged in Freestyle293 medium from Gibco. On the day of transfection, cells were diluted to a concentration of 1 million cells/ml. For a 30 ml transfection, 15 µg of heavy-chain vector and 15 µg of light-chain vector were mixed with 2 ml Opti-MEM and 40 µl 293fectin (then Freestyle293 medium to a total volume of 30 ml). After 6 days of incubation, cells were pelleted by centrifugation (1000 rpm, 10 min) and the supernatant was harvested for protein A purification.

[0312] *Purification.* The recombinantly expressed IgG4 variants of the human antibodies was purified on MabSelect™ SuRe protein-A columns. After column application of antibody, the column was washed with 10 column volumes of PBS buffer, and antibody eluted with 100 mM Glycine, 100 mM NaCl buffer, pH 3.0, followed by buffer exchange into PBS buffer using a HighTrap™ Desalting column. All operations were controlled by an Aktaexpress system from GE Healthcare Amersham Biosciences AB.

[0313] The typical concentration range of purified antibody was from 10 - 130 mg/l (0.3 - 3.3 mg/30 ml).

Results

[0314] cDNA sequences encoding 16F16 (IgG4) H chain, 16F16 L chain, 16F31 (IgG4) H chain, and 16F31 L chain are set forth in SEQ ID NOS:3-6, respectively, and respective sequence identifiers of full-length, variable, and CDR amino acid sequences of 16F16 (IgG4), 16F31 (IgG4), MS (IgG4) and 21 F2 (IgG4) are set forth in Table 1. Figure 4 shows the amino acid sequences for 16F16, 16F31, MS, and 21F2 of IgG4 isotype, high-lighting variable (bold) and CDR (bold underline) regions.

[0315] Using the JointMLc algorithm for IgH joint composition and the D-classification described in Ohm-Laursen et al., (Immunology 2006;119:265-77), the following germline sequences for the variable regions were identified for 16F16 and 16F31:

16F16 VH: VH3_21/D3-9/JH4 (SEQ ID NOS:31/32/33, respectively)

16F16 VL: VKI_L15/JK2 (SEQ ID NOS:34/35, respectively)

16F31 VH: VH3_20/D3-10/JH6 (SEQ ID NOS:36/(EL)/37, respectively)

16F31 VL: VKII_A27/JK3 (SEQ ID NOS:38/39, respectively)

MS VH: VH4_59/D3_27_R3/JH3 (SEQ ID NOS:64/(NWG)/65, respectively)

MS VL: VKIII_A27/JK1 (SEQ ID NOS:38/66, respectively)

21F2 VH: VH5_51/D3_10_R3/JH4 (SEQ ID NOS:67/68/33, respectively)

21F2 VL: VKIII_L6/JK1 (SEQ ID NOS:69/66, respectively)

[0316] Alignments of VH and VL sequences with the corresponding recombined germline sequences (SEQ ID NOS:27-30 correspond to recombined VH3_21/D3-9/JH4, VKI_L15/JK2, VH3_20/D3-10/JH6, and VKIII_A27/JK3, respectively, and SEQ ID NOS:60-63 correspond to recombined VH4_59/D7_27_3/JH3, VKIII_A27/JK1, VH5_51/D3_10_R3/JH4, and VKIII_L6/JK1, respectively) indicating somatic hypermutations, are shown in Figures 5A-5H.

Example 3: MICA blocking experiments

Materials and Methods

[0317] *Flow cytometry assays - MICA blockade.* For analysis of blockade of ligand binding, 50000 NKG2D/DAP10-expressing BaF/3 cells were incubated in 100 μ l total (PBS with 2%FBS at pH7.4) with varying amounts of hybridoma supernatant or purified antibody for 1h at 16°C, followed by incubation with mFc-MICA (for human antibodies) or hFc-MICA (for ON72) (1 μ g) for 30 min at 4°C. The cells were thereafter washed, and secondary Goat-Anti-mouse IgG-HRP Fcy Fragment specific, Jackson (109-036-151) was added for detection of MICA-mFc binding. The cells were then analysed on a B&D FACSAarray flow cytometer. The degree of reduction of MICA binding by preincubation was analysed as MFI (mean fluorescence intensity) of binding with pre-incubation in % of binding of MICA without preincubation.

[0318] A more detailed dose-response curve was also performed, analysing the concentrations of recombinantly expressed and purified antibody needed for 50% inhibition (IC50) and full blockade of 1 μ g MICA-mFc binding.

Results

[0319] The antibodies were analysed for their ability to block ligand binding. Figure 6 demonstrates that pre-incubation with a hybridoma supernatant virtually blocked all binding of the ligand, MICA. A dose-response curve was performed using recombinantly expressed antibodies, demonstrating IC50 and full blockade of a NKG2D saturating dose of MICA-mFc (1 μ g) binding at 0.017 and 0.2 nM 16F16 and at 0.16 and 0.7 nM 16F31 (Figure 7). The corresponding results for ON72 were 0.02 and 0.24 nM for IC50 and full blockade of 1 μ g MICA-Fc, respectively. Detailed results are shown in Table 2 below. The IC50 of MS and 21 F2 were the lowest, 0.0016 nM and 0.0048 nM, respectively.

Table 2

Antibody	IC50 (μ g/ml)	Full blockade of 1 μ g MICA-Fc (μ g/ml)
16F16	0.019	0.59
16F31	0.24	4.8
MS	0.0025	0.053
21F2	0.0063	0.16
ON72	0.019	0.54

Example 4: Competition with murine antibodies

Materials and Methods

[0320] *Flow cytometry assay - competition with murine antibodies.* For analysis of blockade of commercially available murine anti-hNKG2D antibodies, 50000 NKG2D-expressing cells were incubated in 100 μ l final (PBS with 2%FBS at pH7.4) with hybridoma supernatant or purified and recombinantly expressed antibody (at 0.3 μ g or as indicated) for 1 h at 16°C, followed by incubation with a murine anti-hNKG2D antibody (ON72, 149810, 1D11 or 5C6 (for 1D11 and 5C6, see, e.g., Bauer et al., Science

1999:285:727-9 and WO02068615); at 0.3µg or as indicated) for 30 min at 4°C. The cells were thereafter washed, and secondary Goat-Anti-mouse IgG-HRP Fcy Fragment specific, Jackson (109-036-151) was added for detection of binding of the murine antibody. The cells were analysed on a B&D FACSArray. This setup was also performed by pre-incubation with murine antibody followed by the hybridoma supernatant or purified human antibody, using secondary Goat-anti-human IgG-HRP Fcy Fragment specific (Jackson, 109-036-098) for detection. The degree of reduction of binding by pre-incubation was analysed as MFI of binding with pre-incubation in % of binding without pre-incubation.

Results

[0321] Pre-incubation of cells with a hybridoma supernatant followed by incubation with ON72 demonstrated that 95% of ON72-binding was blocked (Figure 8). Performing the same type of assay with recombinantly expressed 16F16 antibody demonstrated that 16F16 blocked 95% of ON72 binding, while pre-incubation with ON72 only blocked 82% of subsequent 16F16 binding (Figure 9A). Likewise, for 149810 and 16F16, only about 50% blockade was observed by either antibody, irrespective of the order of incubation or relative antibody concentration (Figure 9A). For the other available murine anti-hNKG2D antibodies, cross inhibition is presented in Table 2, demonstrating near full blockade of ON72 by 1 D11 and 5C6 and approximately 85 % by 149810. Performing the same type of assay with recombinantly expressed MS demonstrated that pre-incubation with MS inhibited 98% of ON72-binding, 88% of 1 D11-binding and 96.5% of 149810 binding (Figure 9B).

Table 3

"Pre-incubation" antibody	"Post-incubation" antibody	Inhibition (%)
149810	ON72	85
1D11	ON72	97
5C6	ON72	97

Example 5: Blood cell binding and cross-reactivity with monkey NKG2D

Materials and Methods

[0322] *Flow cytometry assay- man and monkey PBMC.* Peripheral blood mononuclear cells (PMBCs) were isolated from humans, or from Cynomolgous or Rhesus monkey. All animal work were performed according to Danish National Research Council guidelines. Each PBMC sample was labelled with a marker for the different cellular subsets (NK, CD8+, CD4+ and γδ T cells, as well as for NKG2D, with either ON72 or recombinantly expressed and purified 16F16. The cells were washed and analysed on a BD FACSDiva (BD Biosource) for staining of subsets of cells for NKG2D with the two antibodies. The MFI of the staining was calculated for the individual antibodies.

[0323] In a separate experiment, the binding of MS and 21F2 to both PBMC preparations and full blood from healthy volunteers or cynomolgous monkey was tested by adding a full dose range of MS or 21 F2 followed by detection with anti-human IgG4 antibody, and EC50 values were calculated. Briefly, incubation with antibody was performed at 4°C for 30 min followed by washing, then directly labelled secondary anti-hlgG4 antibody was added and incubated for 30 min at 4°C along with antibodies specific for the various cell populations of interest, CD8, CD4, NK and γδ-T cells, then the cells were washed twice in PBS with 2%FCS and the red blood cell lysed. The cells were then analyzed by flow cytometry and binding to the different cell populations assessed.

Results

[0324] Results for 16F16 and ON72 are shown in Figures 10A-10D. All NK and CD8+ T cells stained positive for NKG2D, whereas no CD4+ T cells stained positive in cynomolgous or rhesus PMBC's. The same results were obtained for human PMBC's that were run in parallel. This agrees with literature, i.e., that in man, NKG2D is normally expressed on NK cells and CD8+ T cells but not on CD4+ T cells.

[0325] Staining PMBC's from cynomolgous and rhesus monkey with ON72 or 16F16 demonstrated similar binding of the two antibodies to NK cells and CD8+ T cells, but no binding to CD4+ T cells. These results validated the two monkey strains as

suitable species for toxicity studies.

[0326] No cross-reactivity to mouse, rat, dog or pig NKG2D was observed with either commercially available antibody or any of 16F16 or 16F31.

[0327] Results for MS binding to human and cynomolgous CD8 T cells are shown in Figures 11A and 11B, respectively, and EC50 values for the binding of MS and 21 F2 to the PBMC preparations are shown in Table 4, along with the relative EC50 values for cynomolgous to human cells (%). This demonstrates that both MS and 21 F2 have very similar affinities to human and cynomolgous NKG2D.

Table 4

Cell Type	EC50 (µg/ml) for MS			EC50 (µg/ml) for 21 F2		
	Cyno	Human	%	Cyno	Human	%
Gd	0.0301	0.0355	85.5	0.0464	0.0572	81.1
CD8	0.0286	0.0357	80.1	0.0435	0.0605	71.9
NK	0.0309	0.0411	75.2	0.0475	0.0712	66.7

Example 6: Bioassay

[0328] To test that the fully human antibodies actually blocked activity of NKG2D, an NKG2D-ligand-driven cytotoxicity assay was developed. A 51Cr-release assay was used, where the target cells were loaded with the radioactive dye and its release measured as a consequence of NK killing of the cell.

Materials and Methods

[0329] *NKG2D-MICA interaction mediated killing assay.* Biological assays for measuring NKG2D-ligand mediated killing of target cells are suitable for testing anti-NKG2D antibodies. The NK cell lines NK92 or NKL (ATCC No. CRL-2407; Robertson et al., Exp Hematol 1996;24:406-15) both kill MICA-transfected BaF/3 cells in an NKG2D-dependent fashion, and can be used as effector cells, killing 51Cr-loaded target cells expressing an NKG2D-ligand (either MICA, MICB or ULBP1-4).

[0330] In a first assay, NKL cells were incubated for 4h with 51Cr-loaded, MICA-expressing BaF/3 cells in the ratio 10:1, in the presence or absence of 1 or 5 µg ON72 or recombinantly expressed 16F16 of IgG4 isotype. After incubation, the supernatant was transferred to microtiter plates, scintillant added, and the release of 51 Cr was measured, as a result of killing of the target cells, in a Topcounter (Wallach). The reduction in release of 51 Cr was a measure of inhibition of killing by the added antibody, and the percentage of cells that were killed was calculated.

[0331] In a second assay, NK-92 cells was incubated with either ⁵¹Cr-labelled MICA- or ULBP3 expressing BaF/3 cells, and the reduction in killing by addition of increasing concentration of recombinantly expressed and purified 16F16, 16F31, MS, or 21F2 of IgG4 isotype. The results are presented as % inhibition of killing.

Results

[0332] Addition of an ligand blocking antibody, using either ON72 or 16F16, blocked killing of MICA-bearing cells by NKL cells in a dose-dependent fashion, depicted as % inhibition of killing (See Figure 12). Control cells not expressing MICA were not killed by the NKL cells.

[0333] 16F16 also inhibited killing of both MICA- and ULBP-bearing BaF/3 target cells by NK-92 cells in a dose dependent manner (Figures 12A and 12B), depicted as % inhibition of the killing, with near total blockade at 0.8 µg/ml of both MICA- and ULBP-NKG2D induced killing. 16F31 (IgG4) blocked about 75% of killing at the highest dose tested (20 µg/ml; Figures 13A and 13B).

[0334] As shown in Figure 14A, MS and 21 F2 both inhibited killing of ULBP3-bearing cells by NK-92 cells in a 51Cr-release

assay. In Figure 14A, blocking of cytotoxicity is depicted as % inhibition, with 0 being the two cells incubated together without addition of antibody, with MS being more efficient. As shown in Figure 14B, maximum inhibition of killing of ligand-(MICA-) bearing cells by NKL cells in a 51Cr-release assay was obtained at a very low concentration of MS (0.01 µg/ml) while the highest tested concentration of 16F16 (0.1 µg/ml) only lead to about 40% inhibition. A summary of IC50 data are provided in Table 5.

Table 5

Antibody	EC50 (µg/ml)	
	ULPB3-expressing target cells	MICA-expressing target cells
16F16	0.35	0.14
16F31	14.8	14.9
MS	0.012	0.0016
21F2	0.021	
ON72	0.065	

Example 7: Antibody-induced NKG2D downmodulation

[0335] When hNKG2D expressing cells are incubated with antibody, down-modulation, e.g., via internalization, of NKG2D was shown to occur in a similar manner previously demonstrated for certain anti-mNKG2D antibodies in mouse models. This will lead to a different mode of action, and, possibly, a longer effect-time of the antibody. Here, down-modulation was analysed by measuring how much the NKG2D level decreased after overnight incubation with antibody.

Materials and Methods

[0336] *Flow cytometry assay- down-modulation.* Antibody-mediated down-modulation of NKG2D was analysed on different types of NKG2D-expressing cells by incubation overnight with ON72, 16F16, 16F31, MS, or 21F2. Without being limited to theory, differences in down-modulation can reflect differences in antigen-interaction, e.g., epitope. The human antibodies were recombinantly expressed as IgG4 isotype, which isotype binds the Fc-receptor with low affinity.

[0337] In a first experiment, 1 mL containing 1 µg of ON72 or 16F16; 3 µg 16F31; 0.1 µg MS, or 0.3 or 1 µg 21 F2, was added to NKG2D- and DAP10-expressing cells.

[0338] In a second experiment, freshly prepared NK cells were incubated with different amounts of MS (0; 0.003; 0.01; 0.03; 0.1; 0.3; 1 µg) or 21 F2 (0.1 µg) in the presence of 10% human serum to mimic the situation in full blood, with the presence of IgGs with higher affinity for the Fc receptors.

[0339] In a third experiment, 0.1 µg /ml MS, ON72, or 21 F2 was added to whole blood containing NK, CD8+, and γδ T cells. After incubation, staining with anti-CD8, anti-CD56 (NK cells), anti-γδ, and anti-human antibody identified binding to the various subtypes

[0340] As controls in the above experiments, cells were left untreated at 37°C overnight. The next day, both untreated and antibody-treated cells were incubated with 0.1 µg of the pre-treatment antibody, followed by either Goat-Anti-mouse IgG-HRP Fcy Fragment specific, Jackson (109-036-151) for detecting ON72, or Goat-Anti-human IgG-HRP Fcy Fragment specific, Jackson (109-036-098) for detecting human antibody. The cells were then washed and analyzed on a B&D FACSArray. The difference in staining between untreated and anti-body-treated staining levels was analyzed as a measure of NKG2D down-modulation, and the % remaining cell surface NKG2D was calculated as % staining after pre-treatment compared to staining of untreated cells.

Results

[0341] As shown in Figure 15, ON72, 16F16, and 16F31 all induced NKG2D down-modulation in NKG2D-expressing BAF/3 cells. For ON72 and 16F31, approximately 55% down-modulation was observed, as compared to about 75% down-modulation using 16F16. This suggests that 16F16 induces down-modulation more effectively, as it has a similar Kd value to that of ON72. Without being limited to theory, this might be due to different binding epitopes. After incubation with MS, there was about 95% reduction in

surface NKG2D (Figure 16A). In freshly isolated NK cells, an MS concentration corresponding only to about 60% of saturation (0.03 µg/ml) induced maximum internalization of cell-surface NKG2D in the presence of serum, with, in this case, only about 35% NKG2D available for binding after overnight incubation with antibody (Figure 16B). For 21 F2, in the absence of serum, a concentration of 1 µg/mL induced 78% (Figure 17) downmodulation of NKG2D.

[0342] In 3 different populations of NKG2D+ lymphocytes in whole blood, for MS and 21 F2 approximately 80% internalization was observed at 24 h, whereas ON72 induced internalization to nearly 100% (Figure 18). Additionally, ON72 induced internalization faster than MS and 21F2 (Figure 18), reaching 50-76% in 1h.

Example 8: Non-depleting IgG4 versions of human antibodies

[0343] Antibodies cross-linking cells in blood may induce depletion of the antibody-bound cells. However, the affinity of IgG4 antibodies to the activating Fc-receptors is so low compared to that of IgG1 that IgG4 antibodies do not lead to depletion.

Materials and Methods

[0344] *Whole blood cell depletion assay.* To demonstrate non-depletion of NKG2D-expressing cells, whole human blood is incubated with 1 µg of IgG4-versions of human antibodies for 4 hours, and the relative distribution of NKG2D-positive and NKG2D-negative cells analyzed and compared to whole human blood incubated in the absence of antibody. The analysis described in Example 4 can then be used for evaluation.

Example 9: Affinity determination

[0345] Surface plasmon resonance measurements were performed on a Biacore 1000 upgrade apparatus (Biacore GE Healthcare; Biacore Upgrade CA0396) at 25°C. In all Biacore experiments HBS-EP buffer (Biacore GE Healthcare; BR-1001-88) served as running buffer and sensorgrams were analyzed with Biaevaluation 4.1 software.

[0346] *Protein immobilisation.* Recombinant MICA-Fc proteins were obtained from R&D systems or were recombinantly produced. Recombinant ULBP-1, 2, 3, MICB and NKG2D-Fc were purchased from R&D systems. Recombinant NKG2D-Fc proteins were immobilized covalently to carboxyl groups in the dextran layer on a Sensor Chip CM5 (Biacore GE Healthcare; BR-1000-14). The sensor chip surface was activated with EDC/NHS (N-ethyl-N'-(3-dimethylaminopropyl) carbodiimidehydrochloride and N-hydroxysuccinimide (Biacore GE Healthcare; BR-1000-50)). Proteins were diluted to 10 µg/ml in coupling buffer (10 mM acetate, pH 5.2) and injected until the appropriate immobilization level was reached (i.e. 500 to 1000 RU). Deactivation of the remaining activated groups was performed using 100 mM ethanolamine pH 8 (Biacore GE Healthcare; BR-1000-50).

[0347] *Affinity measurements.* Human antibodies 16F16, 16F31, MS, and 21F2, both recombinantly expressed as IgG4 isotype, were compared to murine antibody ON72. For kinetic experiments, serial dilutions of soluble antibodies (from 0.3 to 30 nanoM) were injected for 2 min at a constant flow rate of 40 µl/min on dextran layers containing immobilized NKG2D-Fc proteins (500 to 1000 RU), and allowed to dissociate for 3 min before regeneration by a eight second injection of 500 mM NaCl and 10 mM NaOH buffer. The resulting sensorgrams were analysed by global fitting using the Langmuir model. The dissociation constant (KD) was calculated as $KD = k_d/k_a$.

[0348] *Affinity on cellular membrane-situated NKG2D.* Full dose-response curves for binding of antibody to cells expressing NKG2D were performed to analyze binding affinity to the naturally occurring receptor.

Results

[0349] The affinity determination results for 16F16 is shown in Table 6A, at two different NKG2D-densities on the chip in the Biacore, demonstrating high affinity (KD 1.72 E-10M) and slow off rate (k_d 3.7 E-5/s). ON72 was similar, with a $k_a=9.6E5/(M*s)$; $k_d=1.1$ E-4/s; and $KD=1.7E-10M$. Both MS and 21F2, however, had a higher affinity (KD 2.52 E-12M and 7.79 E-11M, respectively), and MS had a slower off-rate (k_d 1.45 E-05/s) (Table 6B).

[0350] Figure 3 demonstrates dose-dependent NKG2D binding of recombinantly produced and purified human antibodies 16F16, 16F31, MS and 21F2 to NKG2D- and DAP10-expressing BaF/3 cells, as compared to commercially available murine antibodies (ON72 and 149810), using flow cytometry. The EC50 values for binding were as follows:

16F16:	0.051 µg/ml (0.034 nM)
16F31:	0.31 µg/ml (0.21 nM)
MS:	0.032 µg/ml (0.021 nM)
21 F2:	0.033 µg/ml (0.023 nM)
ON72:	0.062 µg/ml (0.048 nM)
149810:	0.063 µg/ml (0.042 nM).

Table 6A

Biacore analysis of 16F16 NKG2D-binding						
NKG2D density (RU)	Ka (M(-1)s(-1))	Kd (s(-1))	Rmax	KD (M)	Chi	Fi
1200	2.08E+05	3.71E-05	1.61E+03	1.78E-10	0.56	v. good
600	4.19E+05	6.91E-05	657	1.65E-10	0.43	v. good
Mean ± SD				(1.72E-10±9.19E-12) M		

Table 6B

Biacore analysis of MS and 21 F2 NKG2D-binding						
NKG2D density (RU)	Ka (1/Ms)	Kd (1/s)	KD (M)	Rmax (RU)	Chi ² (RU ²)	fit
MS - 350	5.75+0E6	1.45E-05	2.52 E-12	89.7	2.9	v.good
21F2 - 400	1.85E+06	1.44E-04	7.79 E-11	112.3	1.42	v.good

Example 10 - Agonist activity of immobilized anti-NKG2D antibodies

[0351] To analyse antibody agonistic activity, proliferation of peripheral blood lymphocytes (PBMcs) stimulated with low levels of CD3 was assessed in the presence or absence of immobilized anti-NKG2D antibody, using CD28 as control. The stimulation was done under circumstances where NKG2D have been shown to act as a co-stimulatory molecule (Mashoo et al, Immunol. 2005;174:4480-4484), believed to reflect the triggering of NKG2D in the presence of pro-inflammatory cytokines as under chronic inflammatory conditions. In this assay, PBMcs were stimulated for 3 days with surface bound antibodies, followed by 4 days of IL-2 stimulation, and the proliferation assessed by CFSE dilution in either all lymphocytes, CD8+, or CD4+ T cells.

Materials and Methods

[0352] *PBMC proliferation assay.* PBMcs were purified by gradient centrifugation. Ninety-six-well Maxisorp plates were coated with anti-Fc antibody (Jackson - Immuno Research 115-006-008), then washed and followed by addition of anti-CD3 (0.1 or 0.3 ng/ml, Bioscience cat#14-0037-82), anti-NKG2D (MS or ON72, 0.2 µg/ml) and/or anti-CD28 antibody (0.2 µg/ml, Becton Dickison cat# 348040). Onehundred-and-fifty thousand PBMcs were added to each well, and the cells incubated at 37°C for 3 days. The cells were thereafter labelled with CFSE (molecular probes cat# C34554). Ten million cells were incubated in 0.5 ml 1 µM CFSE for 10 min at 37°C, followed by a wash, and 150,000 PBMcs per well in 60-well plates were incubated for 4 days with IL-2 (10 U/ml). Finally, the cells were stained with anti-CD8 and anti-CD4 antibody and the proliferation measured by CFSE dilution in either all lymphocytes (total presented in Figure 19) or CD8+ or CD4+ T cells (similar results obtained)

Results

[0353] As shown in Figure 19, MS did not significantly co-stimulate proliferation of lymphocytes at either 0.1 or 0.3 ng/ml CD3 stimulation, whereas ON72 resulted in a small but significant co-stimulation at both CD3 concentrations. See Table 7. In both cases the control, anti-CD28, gave strong co-stimulation, and anti-NKG2D did not add significantly to this. This shows that there is a difference in binding mode of the two antibodies, with immobilized ON72 having detectable agonistic activity, whereas MS is a more pure antagonist.

Table 7

Results from PBMC proliferation assay		
Stimulation	% proliferation	SEM
CD3 0.1	2,82	0,39
CD3 0.1+MS	3,60	0,65
CD3 0.1+ON72	7,12	1,35
CD3 0.1+CD28	24,53	4,71
CD3 0.1+CD28+MS	21,64	7,59
CD3 0.1+CD28+ON72	22,86	6,09
CD3 0.3	15,80	5,30
CD3 0.3+MS	17,32	4,41
CD3 0.3+ON72	28,38	7,00
CD3 0.3+CD28	39,90	6,91
CD3 0.3+CD28+MS	46,08	5,71
CD3 0.3+CD28+ON72	40,12	7,93

Example 11 - Crystal structure of soluble hNKG2D in complex with MS-Fab

[0354] The crystal structure of a soluble fragment of hNKG2D in complex with a Fab fragment of the human monoclonal antibody MS was solved and refined to 1.7 Å resolution with X-ray crystallography. The results confirmed that the antibody, when bound to hNKG2D, blocks the binding of a MICA molecule (Figs 20-22). It was also shown that each hNKG2D dimer bound only one MS Fab fragment. The MS Fab portion bound primarily to one of the two hNKG2D monomers ("NKG2D monomer unit 1"), but, although it only interacted weakly with the other monomer ("NKG2D monomer unit 2"), any further MS Fab was blocked from binding monomer unit 2.

[0355] A list of literature referred to in this Example is provided in Example 12.

Materials and Methods

[0356] Soluble hNKG2D (residues 89-216 of SEQ ID NO:2) and MS Fab (comprising a light chain corresponding to SEQ ID NO:41 and a heavy chain fragment corresponding to residues 1-213 of SEQ ID NO: 40) were mixed with a slight molar excess of hNKG2D and the complex was purified on a gel-filtration column. The complex was then concentrated to about 9.5 mg/ml. Crystals were grown with the hanging drop-technique in 17 % PEG3350, 200 mM sodium malonate and 100 mM bis-tris-propane buffer with a pH of 7.5. Crystals were transferred to a cryo-solution containing 75 % of the precipitant solution and 25 % of glycerole. The crystal was allowed to soak for about 15 seconds. The crystal was then flash frozen in liquid N₂ and kept at 100 K during data collection by a cryogenic N₂ gas stream. Crystallographic data were collected, originally to 2.4 Å resolution at a Rigaku 007HF rotating anode source and thereafter, using a new crystal, to 1.7 Å resolution at beam-line BL911-3(2) at MAX-lab, Lund, Sweden. Space group determination, integration and scaling of the data were made in the XDS software package(3). Cell parameters for the synchrotron data were determined to be 82.1, 54.2, 169.4 Å, 90°, 102.62° and 90°, respectively. Molecular replacement, using the MOLREP(4) and PHASER software program(5;6) of the CCP4 suite(7), a Fab molecule from the PDB-deposited(8) structure 1L71(9) and a hNKG2D molecule from the deposited 1 MPU structure(10), were used for structure determination. The Fab molecule was divided into two domains, the variable and the constant domains, and for the NKG2D a monomer was used as a search model in the molecular replacement calculations. Crystallographic refinements, using the REPMAC5 software program(11), was followed by computer graphics inspection of the electron density maps, model corrections and building using the Coot software program (12). The procedure was cycled until no further significant improvements in could be made to the model. The structure was originally interpreted in the C2 space group, using the rotating anode data. With the synchrotron data XDS(3) indicated a non-centric monoclinic space group though and data were integrated in space group P2₁, later changed to P2₁. A C-centered orthorhombic cell did also score highly. Also the point-LESS software(13) proposed the P2₁ as the correct space group when testing the synchrotron data. The PHASER software program was used for a new round of molecular

replacement, using the preliminary models prepared in the first round of molecular replacement in the C2 space group. Molecular replacement run successfully but as R- and R-free values (a comparisons of observed experimental data with, from the model, calculated data) during refinements did not decrease as expected (R- and R-Free of 0.35 and 0.43), despite reasonable electron density maps, further investigations of the data and refinements were started. Different space groups were tested, including P1, but did not improve the refinements. Instead data was inspected for twinning and data was transferred to the SHELXL refinement software program(14). Using a twin relation-ship of (h,k,l) -> (h,k,-h-l) the R- and R-free for all data dropped from 0.34 and 0.40 to 0.30 and 0.34; respectively, and with a refined twinning factor, BASF, of 0.25. Manual modifications to the model were made with the COOT graphics software program. Refinement was carried out in the SHELXL computer program. Final R- and R-free for all data, with no cut-offs, after 14 cycles of manual intervention and following refinements were 0.277 and 0.320, respectively, and the model showed a root-mean-square deviation (RMSD) from ideal bond lengths of 0.008 Å (Table 8). The refined twinning factor was calculated by SHELXL to be 0.26.

Results

[0357] As shown in Figures 20A, 20B, 21A, and 21C, MS-Fab effectively blocks MICA binding to both monomers of the hNKG2D dimer. However, while the MICA molecule bound to both monomers of the NKG2D dimer(1), the MS Fab bound primarily to one of the two monomers, herein denoted "NKG2D monomer unit 1" (Fig. 21A and 21C). The interactions between MS and NKG2D monomer unit 2 were found to be less specific (e.g., comprise no or fewer hydrogen-bonds), and less important in keeping the MS Fab/NKG2D complex together.

[0358] Calculation by the software program AREAIMOL of the CCP4 program suite(7) of the average areas excluded in pair-wise interactions gave for the two independent soluble hNKG2D/MS-Fab molecular complexes of the determined crystals structure totally 909 and 876 Å², respectively. The average areas excluded in pair-wise interaction between NKG2D monomer unit 1 and MS Fab were calculated to be, for the two independent complexes, 710 and 736 Å², respectively. The excluded areas for the other monomer ("NKG2D monomer unit 2") were substantially smaller, 227 and 158 Å², respectively.

[0359] The direct contacts between the hNKG2D and MS Fab were identified by running the CONTACTS software of the CCP4 program suite(7) using a cut-off distance of 4.0 Å between the MS-Fab and the hNKG2D molecules. The results for the two independent soluble hNKG2D/MS-Fab complex molecules of the crystal structure are shown in Tables 9-12. The resulting hNKG2D epitope for MS was found to comprise the following residues of hNKG2D (SEQ ID NO: 2): Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 (Fig. 21A). In NKG2D monomer unit 2, only 5 interactions were found, and only one residue (Tyr 152) was present in both of the crystallographically independent complexes. Further, the Lys 150 side chain atom Nε was only involved in hydrogen-binding in one of the complexes, and the remaining interactions were of weaker polar and hydrophobic type.

[0360] The MS hNKG2D epitope comprised residues located in the loop just before and the beginning of β-strand β3'(1), Lys 150-Tyr 152; in β5' and the loop after, Thr 180-Gln 185; in β5, Leu 191; in β6, Lys 197, Tyr 199 and Glu 201; and in the loop preceding and in the β7 strand, Thr 205-Thr 208. These contact areas agreed very well with what have been reported as the binding site for MICA on hNKG2D (1). MICA binds asymmetrically to the symmetric homodimer of NKG2D(10). This is also the case for the MS Fab binding to hNKG2D. Therefore, there will be two possible binding orientations to NKG2D for MS Fab relative to MICA. This is shown in Figure 20A, B and Figure 22, where it is clearly seen that the MS Fab blocks both possible MICA relative binding orientations.

[0361] The MS paratope for hNKG2D included residues Tyr 33 and Trp 97 of the MS light (L) chain (SEQ ID NO: 41, Tables 9-12), and residues Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of the heavy (H) chain (SEQ ID NO: 40, Tables 9-12). The hNKG2D epitope, and the residues involved in hydrogen-binding, are also indicated in the amino-acid sequence of hNKG2D in Figure 21A.

Example 12: Crystal structure of soluble hNKG2D in complex with hzON72-Fab

[0362] The crystal structure of soluble hNKG2D in complex with a humanized version of ON72 Fab fragment (hzON72) was solved and refined to 3.15 Å resolution with the use of X-ray crystallography. The results confirmed that the antibody, when bound to hNKG2D, will be able to block binding of MICA molecules to hNKG2D (Figures 20-22). It was also shown that each hNKG2D dimer can bind two hzON72 Fab portions simultaneously.

[0363] A list of literature referred to in this Example is provided at the end of the Example.

Materials and Methods

[0364] A soluble hNKG2D fragment (corresponding to residues 81-216 of SEQ ID NO:2) and hzON72 Fab (SEQ ID NO:70 and SEQ ID NO:71, heavy chain fragment and light chain, respectively) were mixed with a slight molar excess of hNKG2D and the complex was purified on a gel-filtration column. The complex was then concentrated to about 7.5 mg/ml. Crystals were grown with the hanging drop-technique in 1M LiSO₄ and 100 mM MES buffer pH 6.5. Crystals were transferred to a cryo-solution containing 75 % of the precipitant solution and 25 % of glycerole. The crystal was allowed to soak for about 15 seconds. The crystal was then flash frozen in liquid N₂ and kept at 100 K during data collection by a cryogenic N₂ gas stream. Crystallographic data to 3.15 Å resolution were collected using beam-line BL911-5(2) at MAX-lab, Lund, Sweden. Space group determination, integration and scaling of the data were made in the XDS software package(3). Cell parameters were determined to be 65.7, 93.3, 128.9 Å, 90°, 93.83° and 90°, respectively. Space group was determined to be P2₁ with space for one NKG2D dimer and two hzON72 Fab molecules, in the asymmetric unit. Molecular replacement, using the MOLREP(4) software program of the CCP4 suite(7) and the Fab molecule of the PDB-deposited(8) structure 1UJ3(15), and the hNKG2D dimer of the deposited 1 MPU structure(10), were used for structure determination. The Fab molecule was first tested in rotation function runs with different elbow angles from which the Fab with the highest scoring was picked. NKG2D was searched for as a dimer in the molecular replacement calculations. Crystallographic refinements, using non-crystallographic restraints between the two hNKG2D monomers and between the two Fab molecules, were made in the REFMAC5 software program(11) of the CCP4 program suite(7). The crystallographic refinement was followed by computer graphics inspection of the electron density maps, model corrections and building using the Coot software program (12). The procedure was cycled until no further significant improvements in could be made to the model. Final R- and R-free for all data were 0.216 and 0.268, respectively, and the model showed a root-mean-square deviation (RMSD) from ideal bond lengths of 0.012 Å (Table 13).

Results

[0365] A MICA molecule binds strongly to both monomers of NKG2D(1), but two hzON72-Fab molecules instead bound independently to each of the hNKG2D monomers, effectively blocking MICA binding to the NKG2D dimer (Figures 20C, 21B, C and 22). Calculation by the software program AREAIMOL of the CCP4 program suite(7) of the average areas excluded in pair-wise interactions gave for the two crystallographically independent soluble hNKG2D/hzON72-Fab molecular complexes (one hzON72 Fab molecule in complex with one hNKG2D monomer) in the determined crystals structure a total of 791 and 801 Å², respectively. The average areas excluded in pair-wise interaction between the soluble hNKG2D monomers and the heavy chains of hzON72-Fab were calculated to be, for the two crystallographically independent complexes, 642 and 631 Å², respectively, while for the light chains 208 and 242 Å², respectively.

[0366] The direct contacts between the hNKG2D to hzON72-Fab were identified by running the CONTACTS software of the CCP4 program suite using a cut-off distance of 4.0 Å between the hzON72-Fab and hNKG2D molecules. The results for the two independent soluble hNKG2D/hzON72-Fab molecules of the crystal structure are shown in Tables 14-15. The resulting hNKG2D epitope for hzON72 was found to comprise the following residues of hNKG2D (SEQ ID NO: 2): Ser 165, Trp 166, Leu 174, Ser 175, Pro 176, Asn 177, Leu 179, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Lys 186, Ala 193, Ser 194, Ser 195, Lys 197 and Tyr 199. The hzON72 paratope for hNKG2D included residues Tyr 1, Lys 92, Thr 93 and Leu 94 of the hzON72 light (L) chain (SEQ ID NO:71, Table 14-15), and residues Trp 33, Asp 52, Asp 55, Tyr 57, Asn 59, Tyr 60, Tyr 101, Asp 102, Gly 103, Tyr 104, Tyr 105 and Val 106 of the hzON72 heavy (H) chain (SEQ ID NO:70) The hNKG2D epitope for hzON72, and the residues involved in hydrogen-binding, are also indicated in the amino-acid sequence of hNKG2D in Figure 21B.

[0367] The hNKG2D epitope was comprised of residues located in the beginning of β-strand β4(1). Ser 165-Trp 166; in the loop before β5', Leu 174-Asn 177; in the β5' strand and the loop thereafter, Leu 179-Lys 186; the loop before β6, Ala 193-Ser 195; and in the β6 strand, Lys 197 and Tyr 199. These contact areas agreed very well with what have been reported as the binding site for MICA on hNKG2D(1) and it was clear that hzON72 antibody can block the MICA binding. This is shown in Figures 20C, 21B, C and 22.

Table 8. Results from the X-ray model refinement to the observed data of the NKG2D/MS-Fab complex by the software program SHELXL(14).

```

REMARK 2 RESOLUTION. 1.70 ANGSTROMS.
REMARK 3
REMARK 3 REFINEMENT.
REMARK 3 PROGRAM : SHELXL-97
REMARK 3 AUTHORS : G.M.SHELDRICK
REMARK 3
REMARK 3 DATA USED IN REFINEMENT.
REMARK 3 RESOLUTION RANGE HIGH (ANGSTROMS) : 1.70
REMARK 3 RESOLUTION RANGE LOW (ANGSTROMS) : 10.00
REMARK 3 DATA CUTOFF (SIGMA(F)) : 0.0
REMARK 3 COMPLETENESS FOR RANGE (%) : 92.1
REMARK 3 CROSS-VALIDATION METHOD : FREE R
REMARK 3 FREE R VALUE TEST SET SELECTION : RANDOM
REMARK 3
REMARK 3 FIT TO DATA USED IN REFINEMENT (NO CUTOFF).
REMARK 3 R VALUE (WORKING + TEST SET, NO CUTOFF) : 0.2766
REMARK 3 R VALUE (WORKING SET, NO CUTOFF) : 0.2772
REMARK 3 FREE R VALUE (NO CUTOFF) : 0.3203
REMARK 3 FREE R VALUE TEST SET SIZE (% NO CUTOFF) : 5.3
REMARK 3 FREE R VALUE TEST SET COUNT (NO CUTOFF) : 7735
REMARK 3 TOTAL NUMBER OF REFLECTIONS (NO CUTOFF) : 146970
REMARK 3
REMARK 3 FIT/AGREEMENT OF MODEL FOR DATA WITH F>4SIG(F).
REMARK 3 R VALUE (WORKING + TEST SET, F>4SIG(F)) : 0.2566
REMARK 3 R VALUE (WORKING SET, F>4SIG(F)) : 0.2573
REMARK 3 FREE R VALUE (F>4SIG(F)) : 0.3003
REMARK 3 FREE R VALUE TEST SET SIZE (% F>4SIG(F)) : 5.3
REMARK 3 FREE R VALUE TEST SET COUNT (F>4SIG(F)) : 6402
REMARK 3 TOTAL NUMBER OF REFLECTIONS (F>4SIG(F)) : 121240
REMARK 3
REMARK 3 NUMBER OF NON-HYDROGEN ATOMS USED IN REFINEMENT.
REMARK 3 PROTEIN ATOMS : 10194
REMARK 3 NUCLEIC ACID ATOMS : 0
REMARK 3 HETEROGEN ATOMS : 0
REMARK 3 SOLVENT ATOMS : 132
REMARK 3
REMARK 3 MODEL REFINEMENT.
REMARK 3 OCCUPANCY SUM OF NON-HYDROGEN ATOMS : 10326.00
REMARK 3 OCCUPANCY SUM OF HYDROGEN ATOMS : 0.00
REMARK 3 NUMBER OF DISCRETELY DISORDERED RESIDUES : 0
REMARK 3 NUMBER OF LEAST-SQUARES PARAMETERS : 41308
REMARK 3 NUMBER OF RESTRAINTS : 42586
REMARK 3
REMARK 3 RMS DEVIATIONS FROM RESTRAINT TARGET VALUES.
REMARK 3 BOND LENGTHS (A) : 0.008
REMARK 3 ANGLE DISTANCES (A) : 0.020
REMARK 3 SIMILAR DISTANCES (NO TARGET VALUES) (A) : 0.000
REMARK 3 DISTANCES FROM RESTRAINT PLANES (A) : 0.0247
REMARK 3 ZERO CHIRAL VOLUMES (A**3) : 0.027
REMARK 3 NON-ZERO CHIRAL VOLUMES (A**3) : 0.032
REMARK 3 ANTI-BUMPING DISTANCE RESTRAINTS (A) : 0.025
REMARK 3 RIGID-BOND ADP COMPONENTS (A**2) : 0.000
REMARK 3 SIMILAR ADP COMPONENTS (A**2) : 0.098
REMARK 3 APPROXIMATELY ISOTROPIC ADPS (A**2) : 0.000
REMARK 3
REMARK 3 BULK SOLVENT MODELING.
REMARK 3 METHOD USED : MOEWS & KRETSINGER, J.MOL.BIOL.91(1973)201-228
REMARK 3

```

```

REMARK 3 STEREOCHEMISTRY TARGET VALUES : ENGH AND HUBER
REMARK 3 SPECIAL CASE:
REMARK 3

```

```

REMARK 200 SOFTWARE USED: SHELX
REMARK 200 STARTING MODEL: NONE

```

```
REMARK 200 REMARK:
```

```
REMARK 280
```

```
REMARK 280 CRYSTAL
```

```
REMARK 280 SOLVENT CONTENT, VS (%) 42.2
```

```
REMARK 280 MATTHEWS COEFFICIENT, VM (ANGSTROMS**3/DA): 2.13
```

Table 9

hNKG2D monomer "N" (SEQ ID NO: 2) interactions with the "H", MS-Fab heavy chain (SEQ ID NO: 40) and "L", MS-Fab light chain (SEQ ID NO: 41). This is for the first of the crystallographically independent hNKG2D/MS-Fab complex molecules in the crystal. A cut-off of 4.0 Å was used. The contacts were identified by the CONTACT computer program of the CCP4 suite(7). In the last column "****" indicates a strong possibility for a hydrogen bond at this contact (distance < 3.3 Å) as calculated by CONTACT, "***" indicates a weak possibility (distance > 3.3 Å). Blank indicates that the program considered there to be no possibility of a hydrogen bond. Hydrogen-bonds are specific between a donor and an acceptor, are typically strong, and are easily identifiable.

hNKG2D			MS			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Lys	150 N	CG	Ser	31 H	CB	3.93	
			Ser	31 H	OG	2.84	
Lys	150 N	CD	Ser	31 H	OG	3.11	
Lys	150 N	CE	Ser	31 H	OG	3.78	

hNKG2D			MS			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Ser	151 N	CA	Ser	30 H	O	3.73	
			Ser	31 H	O	3.84	
			Ser	31 H	CA	3.79	
Ser	151 N	CB	Ser	30 H	O	3.26	
Ser	151 N	OG	Ser	30 H	O	2.30	***
			Ser	31 H	CA	3.99	
			Ser	30 H	C	3.26	
Tyr	152 N	CG	Tyr	33 H	CE2	3.91	
Tyr	152 N	CD1	Tyr	33 H	CE2	3.94	
Tyr	152 N	CE1	Tyr	33 H	CE2	3.77	
			Tyr	53 H	CE2	3.59	
			Tyr	53 H	CD2	3.42	
Tyr	152 N	cz	Tyr	33 H	CE2	3.58	
			Ser	30 H	O	3.88	
			Tyr	53 H	CD2	3.66	
			Tyr	32 H	C	3.84	
			Tyr	32 H	O	3.52	
			Tyr	33 H	CD2	3.67	
Tyr	152 N	OH	Tyr	53 H	CD2	3.28	
			Ser	52 H	CA	3.78	
			Ser	52 H	C	3.86	
			Tyr	53 H	N	3.06	***
			Tyr	32 H	C	3.38	
			Tyr	32 H	O	2.68	***
			Tyr	33 H	N	3.99	*
			Tyr	33 H	CD2	3.99	
			Tyr	53 H	CB	3.84	
Tyr	152 N	CE2	Tyr	33 H	CE2	3.56	
			Ser	30 H	O	3.93	
			Tyr	32 H	N	3.80	
			Tyr	32 H	CA	3.61	
			Tyr	32 H	C	3.37	
			Tyr	32 H	O	3.49	
			Tyr	33 H	N	3.81	
			Tyr	33 H	CD2	3.40	
Tyr	152 N	CD2	Tyr	33 H	CE2	3.73	
			Tyr	33 H	CD2	3.87	
Thr	180 N	CG2	Tyr	33 L	OH	3.43	
			Tyr	33 L	CE1	3.96	
Thr	180 N	C	Tyr	33 L	OH	3.94	
Ile	181 N	N	Tyr	33 L	OH	3.46	*
Ile	181 N	C	Tyr	33 L	OH	3.53	

hNKG2D			MS			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Ile	181 N	O	Tyr	33 L	OH	2.65	***
			Tyr	33 L	CZ	3.42	
			Tyr	33 L	CE2	3.33	
Ile	182 N	CD1	Tyr	33 L	OH	3.53	
			Tyr	33 L	CZ	3.51	
			Tyr	33 L	CE2	3.65	
Glu	183 N	O	Trp	97 L	CH2	3.67	
			Trp	97 L	CZ2	3.09	
Met	184 N	CE	Trp	98 H	CD1	3.75	
			Tyr	33 H	CB	3.85	
			Tyr	33 H	CG	3.71	
			Tyr	33 H	CD2	3.75	
Met	184 N	C	Asn	58 H	ND2	3.92	
Met	184 N	O	Asn	58 H	CB	3.95	
			Asn	58 H	CG	3.86	
			Asn	58 H	ND2	2.94	***
			Trp	97 L	CH2	3.62	
			His	50 H	CE1	3.55	
			His	50 H	NE2	3.45	*
Gln	185 N	CG	Asn	58 H	ND2	4.00	
Gln	185 N	CD	Ser	56 H	OG	3.04	
			Ser	56 H	O	3.92	
			Ala	57 H	O	3.52	
Gln	185 N	OE1	Ser	56 H	CB	3.77	
			Ser	56 H	OG	2.41	***
			Ala	57 H	O	3.95	*
Gln	185 N	NE2	Asn	58 H	CB	3.55	
			Ser	56 H	OG	2.99	***
			Ser	56 H	C	3.70	
			Ser	56 H	O	2.97	***
			Ala	57 H	O	2.64	***
			Asn	58 H	CA	3.96	
			Ala	57 H	C	3.27	
			Asn	58 H	N	3.78	*
Leu	191 N	CD1	Tyr	33 H	OH	3.05	
Lys	197 N	NZ	Asp	99 H	O	3.13	***
Tyr	199 N	OH	Trp	98 H	CD1	3.92	
Tyr	199 N	CD2	Tyr	33 H	CE2	3.67	
Glu	201 N	CG	Tyr	33 H	CZ	3.92	
			Tyr	33 H	OH	2.80	
Glu	201 N	CD	Tyr	33 H	OH	3.29	
			Ser	56 H	OG	3.60	

hNKG2D			MS			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Glu	201 N	OE1	Tyr	33 H	CZ	3.58	
			Tyr	33 H	OH	3.01	***
			Ser	56 H	CB	3.59	
			Ser	56 H	OG	3.14	***
			Ser	56 H	O	3.94	*
			Tyr	33 H	CE1	3.24	
Glu	201 N	OE2	Ser	56 H	CB	3.78	
			Ser	56 H	OG	3.26	***
Thr	205 N	OG1	Ser	56 H	CB	3.63	
			Ser	56 H	OG	3.95	*
Thr	205 N	CG2	Ser	56 H	CB	3.99	
			Ser	54 H	OG	3.18	
Pro	206 N	O	Ser	54 H	CB	3.03	
			Ser	54 H	OG	2.87	***
Asn	207 N	OD1	Ser	54 H	OG	3.67	*
Thr	208 N	N	Tyr	53 H	OH	3.43	*
Thr	208 N	CB	Tyr	53 H	OH	3.96	
Thr	208 N	OG1	Tyr	53 H	OH	3.57	*
Thr	208 N	CG2	Tyr	53 H	OH	3.37	

Table 10

hNKG2D monomer "C" (SEQ ID NO: 2) interactions with the "B", MS-Fab heavy chain (SEQ ID NO: 40) and "A", MS-Fab light chain (SEQ ID NO: 41). This is for the second of the crystallographically independent hNKG2D/MS-Fab complex molecules in the crystal. A cut-off of 4.0 Å was used. The contacts were identified by the CONTACT computer program of the CCP4 suite(7). In the last column "****" indicates a strong possibility for a hydrogen bond at this contact (distance < 3.3 Å) as calculated by CONTACT, "*" indicates a weak possibility (distance > 3.3 Å). Blank indicates that the program considered there to be no possibility of a hydrogen bond.

hNKG2D			MS			distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Lys	150 C	CG	Ser	31 B	OG	3.54	
Lys	150 C	CD	Asp	27 B	OD1	3.96	
			Ser	31 B	OG	3.56	
Lys	150 C	CE	Asp	27 B	CG	3.67	
			Asp	27 B	OD1	2.83	
			Asp	27 B	OD2	3.81	
			Ser	31 B	OG	3.30	
Lys	150 C	NZ	Asp	27 B	CG	3.30	
			Asp	27 B	OD1	2.76	****
			Asp	27 B	OD2	3.46	*
Ser	151 C	CA	Ser	30 B	O	3.46	
			Ser	31 B	O	3.80	
Ser	151 C	CB	Ser	30 B	C	3.99	
			Ser	30 B	O	3.00	
Ser	151 C	OG	Ser	30 B	CB	3.93	

hNKG2D			MS			distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
			Ser	30 B	C	3.58	
			Ser	30 B	O	2.47	***
Tyr	152 C	CD1	Tyr	33 B	CE2	3.93	
Tyr	152 C	CE1	Tyr	53 B	CE2	3.54	
			Tyr	53 B	CD2	3.44	
			Tyr	33 B	CE2	3.72	
Tyr	152 C	CZ	Tyr	32 B	O	3.62	
			Tyr	53 B	CD2	3.59	
			Tyr	33 B	CE2	3.61	
			Tyr	33 B	CD2	3.68	
Tyr	152 C	OH	Tyr	53 B	CB	3.88	
			Ser	52 B	CA	3.64	
			Ser	52 B	CB	3.73	
			Ser	52 B	C	3.75	
			Tyr	53 B	N	2.98	***
			Tyr	32 B	C	3.65	
			Tyr	32 B	O	2.92	***
			Tyr	53 B	CD2	3.25	
			Tyr	33 B	CD2	3.98	
Tyr	152 C	CE2	Tyr	32 B	N	3.82	
			Tyr	32 B	CA	3.74	
			Tyr	32 B	C	3.42	
			Tyr	32 B	O	3.43	
			Tyr	33 B	N	3.90	
			Tyr	33 B	CE2	3.72	
			Tyr	33 B	CD2	3.52	
			Ser	30 B	O	3.76	
Tyr	152 C	CD2	Tyr	33 B	CE2	3.93	
			Tyr	33 B	CD2	3.99	
Thr	180 C	CG2	Tyr	33 A	OH	3.46	
Thr	180 C	C	Tyr	33 A	OH	3.86	
Ile	181 C	N	Tyr	33 A	OH	3.37	*
Ile	181 C	CA	Tyr	33 A	OH	3.92	
Ile	181 C	C	Tyr	33 A	OH	3.38	
Ile	181 C	O	Tyr	33 A	CE2	3.11	
			Tyr	33 A	CZ	3.20	
			Tyr	33 A	OH	2.53	***
Ile	182 C	CG1	Trp	98 B	CZ2	3.52	
			Trp	98 B	CH2	3.37	
Ile	182 C	CD1	Tyr	33 A	CE2	3.55	
			Trp	98 B	CZ2	3.93	
			Tyr	33 A	CZ	3.48	

hNKG2D			MS			distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
			Tyr	33 A	OH	3.59	
Ile	182 C	CG2	Trp	98 B	CZ2	3.97	
Glu	183 C	O	Trp	97 A	CH2	3.27	
			Trp	97 A	CZ2	3.49	
Met	184 C	CA	His	50 B	NE2	3.99	
Met	184 C	CB	Tyr	33 B	CE1	3.74	
			Tyr	33 B	OH	3.98	
			Tyr	33 B	CZ	3.96	
Met	184 C	CG	Tyr	33 B	CD1	3.90	
			Tyr	33 B	CE1	3.52	
			Tyr	33 B	OH	3.95	
			Tyr	33 B	CE2	3.98	
			Tyr	33 B	CZ	3.57	
Met	184 C	CE	Tyr	33 B	CB	3.60	
			Tyr	33 B	CG	3.39	
			Tyr	33 B	CD1	3.68	
			Tyr	33 B	CD2	3.74	
Met	184 C	C	His	50 B	NE2	3.85	
Met	184 C	O	Asn	58 B	ND2	3.02	***
			His	50 B	CE1	3.14	
			His	50 B	NE2	2.99	***
Gln	185 C	CD	Ala	57 B	O	3.73	
			Asn	58 B	CB	3.48	
			Asn	58 B	CG	3.89	
			Asn	58 B	ND2	3.73	
Gln	185 C	OE1	Asn	58 B	CA	3.96	
			Asn	58 B	CB	3.03	
			Asn	58 B	CG	3.53	
			Asn	58 B	ND2	3.23	***
			His	50 B	ND1	3.91	*
			His	50 B	CE1	2.97	
			His	50 B	NE2	3.62	*
Gln	185 C	NE2	Ala	57 B	C	3.00	
			Asn	58 B	N	3.41	*
			Asn	58 B	CA	3.59	
			Ser	56 B	OG	3.64	*
			Ala	57 B	O	2.50	***
			Asn	58 B	CB	3.55	
Leu	191 C	CD1	Tyr	33 B	OH	3.20	
Lys	197 C	CD	Trp	98 B	CZ3	3.77	
			Trp	98 B	CH2	3.71	
Lys	197 C	CE	Trp	98 B	CZ3	3.54	

hNKG2D			MS			distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
			Asp	99 B	O	3.86	
Lys	197 C	NZ	Trp	98 B	CZ3	3.50	
			Asp	99 B	O	2.73	***
			Asp	99 B	OD1	3.56	*
			Asp	99 B	C	3.74	
Tyr	199 C	CE1	Trp	98 B	CE3	3.87	
			Trp	98 B	CZ3	3.29	
			Trp	98 B	CH2	3.84	
Tyr	199 C	CZ	Trp	98 B	CE3	3.64	
			Trp	98 B	CZ3	3.39	
Tyr	199 C	OH	Trp	98 B	CE3	3.01	
			Trp	98 B	CZ3	2.80	
Tyr	199 C	CD2	Tyr	33 B	CE2	3.57	
			Tyr	33 B	CD2	3.98	
Glu	201 C	CG	Tyr	33 B	OH	2.80	
Glu	201 C	CD	Ser	56 B	CB	3.72	
			Ser	56 B	OG	3.32	
			Tyr	33 B	OH	3.32	
Glu	201 C	OE1	Ser	56 B	CB	3.29	
			Ser	56 B	OG	3.04	***
			Tyr	33 B	CE1	3.58	
			Tyr	33 B	OH	3.08	***
			Tyr	33 B	CZ	3.80	
Glu	201 C	OE2	Ser	56 B	CB	3.49	
I			Ser	56 B	OG	2.87	***
Thr	205 C	OG1	Ser	56 B	CB	3.83	
I			Ser	56 B	OG	3.49	*
Thr	205 C	CG2	Ser	56 B	N	3.85	
			Ser	56 B	CA	3.99	
			Ser	56 B	CB	3.30	
			Ser	56 B	OG	3.84	
			Ser	54 B	OG	3.93	
Pro	206 C	CG	Ser	54 B	O	3.97	
Pro	206 C	C	Ser	54 B	OG	3.94	
Pro	206 C	O	Ser	54 B	CB	2.96	
			Ser	54 B	OG	2.71	***
Asn	207 C	OD1	Ser	54 B	OG	3.78	*
Thr	208 C	N	Tyr	53 B	OH	3.59	*
Thr	208 C	OG1	Tyr	53 B	OH	3.92	*
Thr	208 C	CG2	Tyr	53 B	OH	3.59	

Table 11

hNKG2D monomer "M" (SEQ ID NO: 2) interactions with the "H", MS-Fab heavy chain (SEQ ID NO: 40) and "L", MS-Fab light chain (SEQ ID NO: 41). This is for the first of the crystallographically independent hNKG2D/MS-Fab complex molecules in the crystal. A cut-off of 4.0 Å was used. The contacts were identified by the CONTACT computer program of the CCP4 suite(7). In the last column "****" indicates a strong possibility for a hydrogen bond at this contact (distance < 3.3 Å) as calculated by CONTACT, "***" indicates a weak possibility (distance > 3.3 Å). Blank indicates that the program considered there to be no possibility of a hydrogen bond.

hNKG2D			MG			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Tyr	152 M	OH	Asp	26 H	O	3.66	*
Met	184 M	CG	Gln	1 H	OE1	3.58	
Gln	185 M	NE2	Gln	1 H	OE1	3.92	*
Tyr	199 M	OH	Asp	26 H	OD2	3.87	*

Table 12

hNKG2D monomer "D" (SEQ ID NO: 2) interactions with the "B", MS-Fab heavy chain (SEQ ID NO: 40) and "A", MS-Fab light chain (SEQ ID NO: 41). This is for the second of the crystallographically independent hNKG2D/MS-Fab complex molecules in the crystal. A cut-off of 4.0 Å was used. The contacts were identified by the CONTACT computer program of the CCP4 suite(7). In the last column "****" indicates a strong possibility for a hydrogen bond at this contact (distance < 3.3 Å) as calculated by CONTACT, "***" indicates a weak possibility (distance > 3.3 Å). Blank indicates that the program considered there to be no possibility of a hydrogen bond.

hNKG2D			MS			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Lys	150 D	CE	Tyr	32 B	CE1	3.99	
			Tyr	32 B	CZ	3.78	
			Tyr	32 B	OH	2.80	
Lys	150 D	NZ	Tyr	32 B	CE1	3.00	
			Tyr	32 B	cz	3.05	
			Tyr	32 B	OH	2.60	***
Tyr	152 D	OH	Asp	26 B	O	3.99	*
			Gln	1 B	CA	3.72	

Table 13. Results from the X-ray model refinement to the observed data of the hNKG2D/hzON72-Fab complex by the REFMAC5 software program(11).

```

REMARK 3  REFINEMENT.
REMARK 3  PROGRAM      : REFMAC 5.2.0019
REMARK 3  AUTHORS      : MURSHUDOV,VAGIN,DODSON
REMARK 3
REMARK 3  REFINEMENT TARGET : MAXIMUM LIKELIHOOD
REMARK 3
REMARK 3  DATA USED IN REFINEMENT.
REMARK 3  RESOLUTION RANGE HIGH (ANGSTROMS) : 3.15
REMARK 3  RESOLUTION RANGE LOW  (ANGSTROMS) : 29.66
REMARK 3  DATA CUTOFF          (SIGMA(F)) : NONE
REMARK 3  COMPLETENESS FOR RANGE (%) : 100.00
REMARK 3  NUMBER OF REFLECTIONS : 25642
REMARK 3
REMARK 3  FIT TO DATA USED IN REFINEMENT.
REMARK 3  CROSS-VALIDATION METHOD : THROUGHOUT
REMARK 3  FREE R VALUE TEST SET SELECTION : RANDOM
REMARK 3  R VALUE (WORKING + TEST SET) : 0.21870

```

```

REMARK 3 R VALUE (WORKING SET) : 0.21608
REMARK 3 FREE R VALUE : 0.26854
REMARK 3 FREE R VALUE TEST SET SIZE (%) : 5.0
REMARK 3 FREE R VALUE TEST SET COUNT : 1350
REMARK 3
REMARK 3 FIT IN THE HIGHEST RESOLUTION BIN.
REMARK 3 TOTAL NUMBER OF BINS USED : 20
REMARK 3 BIN RESOLUTION RANGE HIGH : 3.150
REMARK 3 BIN RESOLUTION RANGE LOW : 3.231
REMARK 3 REFLECTION IN BIN (WORKING SET) : 1849
REMARK 3 BIN COMPLETENESS (WORKING+TEST) (%) : 100.00
REMARK 3 BIN R VALUE (WORKING SET) : 0.331
REMARK 3 BIN FREE R VALUE SET COUNT : 97
REMARK 3 BIN FREE R VALUE : 0.385
REMARK 3
REMARK 3 NUMBER OF NON-HYDROGEN ATOMS USED IN REFINEMENT.
REMARK 3 ALL ATOMS : 8737
REMARK 3
REMARK 3 B VALUES.
REMARK 3 FROM WILSON PLOT (A**2) : NULL
REMARK 3 MEAN B VALUE (OVERALL, A**2) : 49.049
REMARK 3 OVERALL ANISOTROPIC B VALUE.
REMARK 3 B11 (A**2) : -0.66
REMARK 3 B22 (A**2) : -3.44
REMARK 3 B33 (A**2) : 3.95
REMARK 3 B12 (A**2) : 0.00
REMARK 3 B13 (A**2) : -1.15
REMARK 3 B23 (A**2) : 0.00
REMARK 3
REMARK 3 ESTIMATED OVERALL COORDINATE ERROR.
REMARK 3 ESU BASED ON R VALUE (A) : NULL
REMARK 3 ESU BASED ON FREE R VALUE (A) : 0.502
REMARK 3 ESU BASED ON MAXIMUM LIKELIHOOD (A) : 0.407
REMARK 3 ESU FOR B VALUES BASED ON MAXIMUM LIKELIHOOD (A**2) : 54.265
REMARK 3
REMARK 3 CORRELATION COEFFICIENTS.
REMARK 3 CORRELATION COEFFICIENT FO-FC : 0.896
REMARK 3 CORRELATION COEFFICIENT FO-FC FREE : 0.844
REMARK 3
REMARK 3 RMS DEVIATIONS FROM IDEAL VALUES COUNT RMS WEIGHT
REMARK 3 BOND LENGTHS REFINED ATOMS (A) : 8971 ; 0.012 ; 0.022
REMARK 3 BOND ANGLES REFINED ATOMS (DEGREES) : 12204 ; 1.516 ; 1.948
REMARK 3 TORSION ANGLES, PERIOD 1 (DEGREES) : 1116 ; 7.673 ; 5.000
REMARK 3 TORSION ANGLES, PERIOD 2 (DEGREES) : 368 ; 37.729 ; 24.402
REMARK 3 TORSION ANGLES, PERIOD 3 (DEGREES) : 1464 ; 21.254 ; 15.000
REMARK 3 TORSION ANGLES, PERIOD 4 (DEGREES) : 32 ; 19.705 ; 15.000
REMARK 3 CHIRAL-CENTER RESTRAINTS (A**3) : 1338 ; 0.109 ; 0.200
REMARK 3 GENERAL PLANES REFINED ATOMS (A) : 6759 ; 0.004 ; 0.020
REMARK 3 NON-BONDED CONTACTS REFINED ATOMS (A) : 3866 ; 0.230 ; 0.200
REMARK 3 NON-BONDED TORSION REFINED ATOMS (A) : 5954 ; 0.312 ; 0.200
REMARK 3 H-BOND (X...Y) REFINED ATOMS (A) : 311 ; 0.169 ; 0.200
REMARK 3 SYMMETRY VDW REFINED ATOMS (A) : 56 ; 0.279 ; 0.200
REMARK 3 SYMMETRY H-BOND REFINED ATOMS (A) : 6 ; 0.260 ; 0.200
REMARK 3
REMARK 3 ISOTROPIC THERMAL FACTOR RESTRAINTS. COUNT RMS WEIGHT
REMARK 3 MAIN-CHAIN BOND REFINED ATOMS (A**2) : 5674 ; 0.279 ; 1.500
REMARK 3 MAIN-CHAIN ANGLE REFINED ATOMS (A**2) : 9064 ; 0.512 ; 2.000
REMARK 3 SIDE-CHAIN BOND REFINED ATOMS (A**2) : 3824 ; 0.749 ; 3.000

```

```

REMARK 3 SIDE-CHAIN ANGLE REFINED ATOMS (A**2): 3140 ; 1.231 ; 4.500
REMARK 3
REMARK 3 NCS RESTRAINTS STATISTICS
REMARK 3 NUMBER OF DIFFERENT NCS GROUPS : 3
REMARK 3
REMARK 3 NCS GROUP NUMBER : 1
REMARK 3 CHAIN NAMES : L A
REMARK 3 NUMBER OF COMPONENTS NCS GROUP : 1
REMARK 3 COMPONENT C SSSEQI TO C SSSEQI CODE
REMARK 3 1 L 1 L 214 1
REMARK 3 1 A 1 A 214 1
REMARK 3 GROUP CHAIN COUNT RMS WEIGHT
REMARK 3 TIGHT POSITIONAL 1 L (A): 1656 ; 0.05 ; 0.05
REMARK 3 TIGHT THERMAL 1 L (A**2): 1656 ; 0.06 ; 0.50
REMARK 3
REMARK 3 NCS GROUP NUMBER : 2
REMARK 3 CHAIN NAMES : H B
REMARK 3 NUMBER OF COMPONENTS NCS GROUP : 1
REMARK 3 COMPONENT C SSSEQI TO C SSSEQI CODE
REMARK 3 1 H 1 H 220 1
REMARK 3 1 B 1 B 220 1
REMARK 3 GROUP CHAIN COUNT RMS WEIGHT
REMARK 3 TIGHT POSITIONAL 2 H (A): 1668 ; 0.06 ; 0.05
REMARK 3 TIGHT THERMAL 2 H (A**2): 1668 ; 0.07 ; 0.50
REMARK 3
REMARK 3 NCS GROUP NUMBER : 3
REMARK 3 CHAIN NAMES : N C
REMARK 3 NUMBER OF COMPONENTS NCS GROUP : 1
REMARK 3 COMPONENT C SSSEQI TO C SSSEQI CODE
REMARK 3 1 N 88 N 215 1
REMARK 3 1 C 88 C 215 1
REMARK 3 GROUP CHAIN COUNT RMS WEIGHT
REMARK 3 TIGHT POSITIONAL 3 N (A): 1026 ; 0.03 ; 0.05
REMARK 3 TIGHT THERMAL 3 N (A**2): 1026 ; 0.04 ; 0.50
REMARK 3
REMARK 3
REMARK 3 TLS DETAILS
REMARK 3 NUMBER OF TLS GROUPS : 5
REMARK 3 ATOM RECORD CONTAINS RESIDUAL B FACTORS ONLY
REMARK 3
REMARK 3 TLS GROUP : 1
REMARK 3 NUMBER OF COMPONENTS GROUP : 2
REMARK 3 COMPONENTS C SSSEQI TO C SSSEQI
REMARK 3 RESIDUE RANGE : L 1 L 107
REMARK 3 RESIDUE RANGE : H 1 H 121
REMARK 3 ORIGIN FOR THE GROUP (A): 24.2040 90.5950 36.8960
REMARK 3 T TENSOR
REMARK 3 T11: -0.1079 T22: -0.1998
REMARK 3 T33: -0.2449 T12: 0.0010
REMARK 3 T13: -0.0388 T23: -0.0446
REMARK 3 L TENSOR
REMARK 3 L11: 1.1625 L22: 5.5207
REMARK 3 L33: 1.9806 L12: -0.2378
REMARK 3 L13: -0.5312 L23: -0.6045
REMARK 3 S TENSOR
REMARK 3 S11: 0.0803 S12: -0.0443 S13: 0.1447
REMARK 3 S21: 0.3782 S22: -0.2475 S23: 0.0045
REMARK 3 S31: -0.0073 S32: 0.0331 S33: 0.1671

```

```

REMARK 3
REMARK 3 TLS GROUP : 2
REMARK 3 NUMBER OF COMPONENTS GROUP : 2
REMARK 3 COMPONENTS C SSSEQI TO C SSSEQI
REMARK 3 RESIDUE RANGE : L 108 L 214
REMARK 3 RESIDUE RANGE : H 122 H 220
REMARK 3 ORIGIN FOR THE GROUP (A): 35.1200 96.1120 3.4800
REMARK 3 T TENSOR
REMARK 3 T11: -0.2849 T22: 0.0610
REMARK 3 T33: -0.1761 T12: -0.0495
REMARK 3 T13: -0.0038 T23: 0.0239
REMARK 3 L TENSOR
REMARK 3 L11: 2.5261 L22: 3.4556
REMARK 3 L33: 2.6465 L12: -0.8617
REMARK 3 L13: -0.0480 L23: -0.4377
REMARK 3 S TENSOR
REMARK 3 S11: -0.1130 S12: 0.4186 S13: 0.1914
REMARK 3 S21: -0.1933 S22: -0.0187 S23: -0.2357
REMARK 3 S31: -0.0662 S32: 0.2231 S33: 0.1317
REMARK 3
REMARK 3 TLS GROUP : 3
REMARK 3 NUMBER OF COMPONENTS GROUP : 2
REMARK 3 COMPONENTS C SSSEQI TO C SSSEQI
REMARK 3 RESIDUE RANGE : A 1 A 107
REMARK 3 RESIDUE RANGE : B 1 B 121
REMARK 3 ORIGIN FOR THE GROUP (A): 9.5860 42.5190 37.1590
REMARK 3 T TENSOR
REMARK 3 T11: -0.1183 T22: -0.2086
REMARK 3 T33: -0.1316 T12: -0.0149
REMARK 3 T13: -0.0254 T23: 0.0350
REMARK 3 L TENSOR
REMARK 3 L11: 1.2758 L22: 4.8903
REMARK 3 L33: 1.9328 L12: -0.4321
REMARK 3 L13: 0.5123 L23: 0.2320
REMARK 3 S TENSOR
REMARK 3 S11: 0.0143 S12: -0.0304 S13: -0.1607
REMARK 3 S21: 0.3252 S22: -0.0524 S23: -0.1532
REMARK 3 S31: 0.0565 S32: 0.0312 S33: 0.0380
REMARK 3
REMARK 3 TLS GROUP : 4
REMARK 3 NUMBER OF COMPONENTS GROUP : 2
REMARK 3 COMPONENTS C SSSEQI TO C SSSEQI
REMARK 3 RESIDUE RANGE : A 108 A 214
REMARK 3 RESIDUE RANGE : B 122 B 220
REMARK 3 ORIGIN FOR THE GROUP (A): -2.3250 37.1030 4.0740
REMARK 3 T TENSOR
REMARK 3 T11: -0.3236 T22: 0.0744
REMARK 3 T33: -0.1171 T12: -0.0180
REMARK 3 T13: 0.0587 T23: -0.0971
REMARK 3 L TENSOR
REMARK 3 L11: 2.6929 L22: 3.2562
REMARK 3 L33: 2.1386 L12: -0.1058
REMARK 3 L13: 0.1471 L23: -0.1863
REMARK 3 S TENSOR
REMARK 3 S11: -0.1360 S12: 0.4399 S13: -0.3031
REMARK 3 S21: -0.1914 S22: 0.0086 S23: 0.1138
REMARK 3 S31: -0.0130 S32: -0.2162 S33: 0.1274
REMARK 3
REMARK 3 TLS GROUP : 5
REMARK 3 NUMBER OF COMPONENTS GROUP : 2
REMARK 3 COMPONENTS C SSSEQI TO C SSSEQI
REMARK 3 RESIDUE RANGE : N 89 N 215
REMARK 3 RESIDUE RANGE : C 89 C 215
REMARK 3 ORIGIN FOR THE GROUP (A): 17.8890 66.4400 62.9460
REMARK 3 T TENSOR
REMARK 3 T11: 0.4354 T22: 0.1526
REMARK 3 T33: -0.1075 T12: -0.2178
REMARK 3 T13: -0.1984 T23: 0.0840
REMARK 3 L TENSOR
REMARK 3 L11: 1.4035 L22: 6.3187
REMARK 3 L33: 3.4618 L12: 1.1657
REMARK 3 L13: 0.1963 L23: 1.1874
REMARK 3 S TENSOR
REMARK 3 S11: 0.5552 S12: -0.7422 S13: -0.2312
REMARK 3 S21: 1.1999 S22: -0.1494 S23: -0.2659
REMARK 3 S31: 0.2701 S32: 0.0344 S33: -0.4059
REMARK 3
REMARK 3 BULK SOLVENT MODELLING.
REMARK 3 METHOD USED : MASK
REMARK 3 PARAMETERS FOR MASK CALCULATION
REMARK 3 VDW PROBE RADIUS : 1.20
REMARK 3 ION PROBE RADIUS : 0.80
REMARK 3 SHRINKAGE RADIUS : 0.80
REMARK 3
REMARK 3 OTHER REFINEMENT REMARKS: NULL

```

Table 14

hNKG2D monomer "N" (SEQ ID NO:2) interactions with the "H", hzON72-Fab heavy chain (SEQ ID NO:70) and "L", hzON72-Fab light chain (SEQ ID NO:71). This is for the first of the crystallographically independent hNKG2D/hzON72-Fab complex molecules in the crystal. A cut-off of 4.0 Å was used. The contacts were identified by the CONTACT computer program of the CCP4 suite(7). In the last column "****" indicates a strong possibility for a hydrogen bond at this contact (distance < 3.3 Å) as calculated by CONTACT, "***" indicates a weak possibility (distance > 3.3 Å). Blank indicates that the program considered there to be no possibility of a hydrogen bond.

hNKG2D			hzON72			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Ser	165 N	CB	Tyr	104 H	CE2	3.66	
Ser	165 N	OG	Tyr	104 H	CE2	3.53	
			Tyr	104 H	CD2	3.95	
Trp	166 N	NE1	Tyr	105 H	OH	3.38	*
Leu	174 N	CG	Tyr	104 H	CE1	3.94	
Leu	174 N	CD2	Tyr	104 H	CD1	3.78	
			Tyr	104 H	CE1	3.91	
Leu	174 N	C	Tyr	104 H	OH	3.97	
Leu	174 N	O	Tyr	104 H	CE1	3.61	
			Tyr	104 H	CZ	3.61	
			Tyr	104 H	OH	2.90	***
Ser	175 N	C	Tyr	104 H	CE1	3.81	
Ser	175 N	O	Tyr	104 H	CE1	3.79	
Pro	176 N	N	Tyr	104 H	CE1	3.84	
Pro	176 N	CA	Gly	103 H	O	3.99	
			Gly	103 H	CA	3.78	
Pro	176 N	CB	Gly	103 H	CA	3.81	
Asn	177 N	N	Gly	103 H	O	3.29	***
			Tyr	101 H	OH	3.79	*
Asn	177 N	CA	Tyr	101 H	OH	3.63	
Asn	177 N	OD1	Tyr	101 H	CE2	3.52	
Leu	179 N	C	Tyr	101 H	OH	3.54	
Leu	179 N	O	Gly	103 H	O	3.77	*
			Tyr	101 H	CE1	3.60	
			Tyr	101 H	CZ	3.35	
			Tyr	101 H	OH	2.33	***
Thr	180 N	CA	Tyr	104 H	O	3.40	
Thr	180 N	CG2	Trp	33 H	CZ2	3.76	
			Val	106 H	CG2	3.60	
			Trp	33 H	NE1	3.77	
			Tyr	101 H	CE1	3.73	
Thr	180 N	C	Tyr	104 H	O	3.55	
Ile	181 N	N	Tyr	104 H	O	2.78	***
Ile	181 N	CA	Tyr	104 H	O	3.78	
Ile	181 N	CB	Tyr	104 H	O	3.76	
Ile	181 N	CG2	Tyr	105 H	CE1	3.90	
			Tyr	105 H	CZ	3.63	
			Tyr	105 H	OH	3.76	
Ile	182 N	CD1	Asn	59 H	OD1	3.49	
Ile	182 N	CG2	Asn	59 H	CG	3.80	
			Asn	59 H	OD1	3.51	
			Asn	59 H	ND2	3.66	

hNKG2D			hzON72			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Glu	183 N	N	Leu	94 L	CD1	3.64	
Glu	183 N	CB	Lys	92 L	O	3.81	
Glu	183 N	CG	Tyr	105 H	OH	3.91	
Glu	183 N	CD	Tyr	105 H	CE1	3.87	
			Tyr	105 H	CZ	3.90	
			Tyr	105 H	OH	3.06	
Glu	183 N	OE1	Lys	92 L	CG	3.77	
			Lys	92 L	CD	3.95	
			Lys	92 L	CE	3.20	
			Thr	93 L	CG2	3.94	
			Tyr	105 H	OH	3.64	*
Glu	183 N	OE2	Tyr	105 H	CE1	3.95	
			Tyr	105 H	CZ	3.54	
			Tyr	105 H	OH	2.39	***
Glu	183 N	C	Leu	94 L	CD1	3.86	
Glu	183 N	O	Leu	94 L	N	3.16	***
			Thr	93 L	C	3.86	
			Leu	94 L	CD1	3.45	
			Thr	93 L	CA	3.51	
			Thr	93 L	CB	3.61	
Met	184 N	CG	Leu	94 L	CB	3.95	
Met	184 N	CE	Tyr	60 H	O	3.39	
			Asn	59 H	CB	3.92	
Met	184 N	O	Tyr	1 L	CD2	3.98	
			Leu	94 L	O	3.84	*
Lys	186 N	N	Tyr	1 L	OH	3.90	*
Lys	186 N	CB	Tyr	1 L	OH	3.78	
Ala	193 N	CB	Tyr	57 H	CE1	3.84	
Ser	194 N	O	Asp	55 H	CG	3.84	
			Asp	55 H	OD1	3.41	*
			Asp	55 H	OD2	3.47	*
Ser	195 N	CA	Asp	55 H	OD1	3.82	
Ser	195 N	CB	Asp	55 H	CG	3.46	
			Asp	55 H	OD1	2.61	
			Asp	55 H	OD2	3.81	
Ser	195 N	OG	Asp	55 H	CG	3.74	
			Asp	55 H	OD1	3.07	***
Ser	195 N	O	Asp	55 H	CG	3.71	
			Asp	55 H	OD1	3.81	*
			Asp	55 H	OD2	3.23	***
Lys	197 N	CG	Tyr	57 H	CD1	3.79	
			Tyr	57 H	CE1	3.95	

hNKG2D			hzON72			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
			Tyr	57 H	CD2	3.99	
			Tyr	57 H	CG	3.82	
Lys	197 N	CD	Tyr	57 H	CD1	3.84	
			Tyr	57 H	CG	3.74	
			Asp	55 H	OD2	3.17	
			Tyr	57 H	CB	3.89	
Lys	197 N	CE	Trp	33 H	CH2	3.75	
			Trp	33 H	CZ2	3.39	
			Asp	52 H	OD2	3.64	
			Asp	55 H	OD2	3.27	
			Tyr	57 H	CB	3.98	
Lys	197 N	NZ	Asp	55 H	CG	3.27	
			Trp	33 H	CZ2	3.68	
			Asp	52 H	CB	3.78	
			Asp	52 H	CG	3.38	
			Asp	52 H	OD2	2.53	***
			Asp	55 H	CB	3.57	
			Asp	55 H	OD2	2.36	***
			Tyr	57 H	CB	3.77	
Tyr	199 N	CD1	Tyr	57 H	OH	3.64	

Table 15

hNKG2D monomer "C" (SEQ ID NO:2) interactions with the "B", hzON72-Fab heavy chain (SEQ ID NO:70) and "A", hzON72-Fab light chain (SEQ ID NO:71). This is for the second of the crystallographically independent hNKG2D/hzON72-Fab complex molecules in the crystal. A cut-off of 4.0 Å was used. The contacts were identified by the CONTACT computer program of the CCP4 suite(7). In the last column "****" indicates a strong possibility for a hydrogen bond at this contact (distance < 3.3 Å) as calculated by CONTACT, "***" indicates a weak possibility (distance > 3.3 Å). Blank indicates that the program considered there to be no possibility of a hydrogen bond.

hNKG2D			hzON72			Distance [Å]	Possibly H-bond
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name		
Trp	166 C	NE1	Tyr	105 B	OH	3.26	***
Leu	174 C	C	Tyr	104 B	OH	3.78	
Leu	174 C	O	Tyr	104 B	CE1	3.62	
			Tyr	104 B	CZ	3.37	
			Tyr	104 B	OH	2.66	***
Ser	175 C	C	Tyr	104 B	CE1	3.27	
Ser	175 C	O	Tyr	104 B	CD1	3.74	
			Tyr	104 B	CE1	3.37	
Pro	176 C	N	Tyr	104 B	CD1	3.99	
			Tyr	104 B	CE1	3.13	
			Tyr	104 B	CZ	3.98	
			Tyr	104 B	OH	3.92	*
Pro	176 C	CA	Gly	103 B	CA	3.84	
			Tyr	104 B	CD1	3.70	

hNKG2D			hzON72				
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name	Distance [Å]	Possibly H-bond
			Tyr	104 B	CE1	3.24	
Pro	176 C	CB	Gly	103 B	CA	3.76	
			Tyr	104 B	CE1	3.94	
Pro	176 C	CG	Tyr	104 B	CE1	3.75	
			Tyr	104 B	OH	3.71	
Pro	176 C	CD	Tyr	104 B	CE1	3.77	
			Tyr	104 B	OH	3.68	
Asn	177 C	N	Gly	103 B	CA	3.94	
			Gly	103 B	C	3.79	
			Tyr	101 B	OH	3.92	*
			Gly	103 B	O	3.02	***
Asn	177 C	CA	Tyr	101 B	OH	3.49	
			Gly	103 B	O	3.66	
Asn	177 C	OD1	Asp	102 B	O	3.70	*
			Tyr	101 B	CE2	3.41	
Leu	179 C	C	Tyr	101 B	OH	3.99	
Leu	179 C	O	Tyr	101 B	CZ	3.78	
			Tyr	101 B	CE1	3.82	
			Tyr	104 B	O	3.71	*
			Tyr	101 B	OH	2.82	***
			Gly	103 B	O	3.90	*
Thr	180 C	CA	Tyr	104 B	O	3.36	
Thr	180 C	CG2	Trp	33 B	CZ2	3.99	
			Val	106 B	CG2	3.72	
			Tyr	101 B	CE1	3.84	
Thr	180 C	C	Tyr	104 B	O	3.76	
Ile	181 C	N	Tyr	104 B	O	3.15	***
Ile	181 C	CB	Tyr	105 B	CE2	3.99	
			Tyr	105 B	CZ	3.91	
Ile	181 C	CG2	Tyr	105 B	CE1	3.72	
			Tyr	105 B	CZ	3.49	
			Tyr	105 B	OH	3.43	
Ile	182 C	CD1	Asn	59 B	OD1	3.39	
Ile	182 C	CG2	Leu	94 A	CD2	3.97	
			Asn	59 B	CG	3.80	
			Asn	59 B	OD1	3.57	
			Asn	59 B	ND2	3.62	
			Leu	94 A	CD1	3.96	
Glu	183 C	N	Leu	94 A	CD1	3.44	
Glu	183 C	CD	Tyr	105 B	OH	3.53	
Glu	183 C	OE1	Lys	92 A	CE	3.17	
			Thr	93 A	CG2	3.35	

hNKG2D			hzON72				
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name	Distance [Å]	Possibly H-bond
Glu	183 C	OE2	Tyr	105 B	CZ	3.86	
			Tyr	105 B	OH	2.71	***
Glu	183 C	O	Thr	93 A	C	3.88	
			Leu	94 A	N	3.06	***
			Leu	94 A	CA	3.93	
			Leu	94 A	O	3.37	*
			Thr	93 A	CA	3.68	
			Thr	93 A	CB	3.54	
			Leu	94 A	CD1	3.94	
Met	184 C	CA	Leu	94 A	O	3.78	
Met	184 C	CE	Tyr	60 B	O	3.16	
Met	184 C	O	Leu	94 A	O	3.72	*
			Tyr	1 A	CG	3.98	
			Tyr	1 A	CD2	3.93	
Lys	186 C	N	Tyr	1 A	OH	3.81	*
Lys	186 C	CA	Tyr	1 A	OH	3.65	
Lys	186 C	CB	Tyr	1 A	OH	3.51	
Ala	193 C	CB	Tyr	57 B	CE1	3.90	
Ser	194 C	O	Asp	55 B	OD1	3.94	*
			Asp	55 B	OD2	3.84	*
Ser	195 C	CB	Asp	55 B	CG	3.63	
			Asp	55 B	OD1	2.94	
			Asp	55 B	OD2	3.78	
Ser	195 C	OG	Asp	55 B	CG	3.81	
			Asp	55 B	OD1	3.25	***
Ser	195 C	O	Asp	55 B	OD2	3.52	*
Lys	197 C	CG	Tyr	57 B	CZ	3.90	
			Tyr	57 B	CG	3.96	
			Tyr	57 B	CD1	3.80	
			Tyr	57 B	CE1	3.77	
Lys	197 C	CD	Asp	55 B	OD2	3.58	
			Tyr	57 B	CG	3.73	
			Tyr	57 B	CD1	3.67	
Lys	197 C	CE	Trp	33 B	CZ2	3.56	
			Asp	52 B	OD2	3.96	
			Asp	55 B	OD2	3.49	
			Trp	33 B	CH2	3.83	
Lys	197 C	NZ	Trp	33 B	CZ2	3.77	
			Asp	52 B	CG	3.77	
			Asp	52 B	OD2	2.83	***
			Asp	55 B	CB	3.88	
			Asp	55 B	CG	3.44	

hNKG2D			hzON72				
Res. Type	Res. # and Chain	Atom name	Res. Type	Res. # and Chain	Atom name	Distance [Å]	Possibly H-bond
			Asp	55 B	OD2	2.41	***
			Tyr	57 B	CB	3.78	

Reference List

[0368]

1. Li, P., Morris, D. L., Willcox, B. E., Steinle, A., Spies, T., and Strong, R. K. (2001) Nature Immunology 2, 443-451
2. Ursby, T., Mammen, C. B., Cerenius, Y., Svensson, C., Sommarin, B., Fodje, M. N., Kvik, A., Logan, D. T., Als-Nielsen, J., Thunnissen, M. M. G. M., Larsen, S., and Liljas, A. (2004) The New Macromolecular Crystallography Stations At MAX-lab: The MAD Station.
3. Kabsch, W. (1993) J. Appl. Crystallogr. 26, 795-800
4. Vagin, A. and Teplyakov, A. (1997) J. Appl. Crystallogr. 30, 1022-1025
5. McCoy, A. J., Grosse-Kunstleve, R. W., Storoni, L. C., and Read, R. J. (2005) Acta Crystallographica Section D Biological Crystallography 61, 458-464
6. McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., and Read, R. J. (2007) J. Appl. Crystallogr. 40, 658-674
7. Bailey, S. (1994) Acta Crystallogr. Sect. D-Biol. Crystallogr. 50, 760-763
8. Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N., and Bourne, P. E. (2000) Nucleic Acids Res. 28, 235-242
9. Vajdos, F. F., Adams, C. W., Breece, T. N., Presta, L. G., de Vos, A. M., and Sidhu, S. S. (2002) Journal of Molecular Biology 320, 415-428
10. McFarland, B. J., Kortemme, T., Yu, S. F., Baker, D., and Strong, R. K. (2003) Structure (Cambridge) 11, 411-422
11. Murshudov, G. N., Vagin, A. A., and Dodson, E. J. (1997) Acta Crystallogr. Sect. D-Biol. Crystallogr. 53, 240-255
12. Emsley, P. and Cowtan, K. (2004) Acta Crystallogr. Sect. D-Biol. Crystallogr. 60, 2126-2132
13. Evans, P. (1962) Acta Crystallogr D Biol Crystallogr. 2006. Jan. 72-82
14. Sheldrick, G. (2008) Act Cryst a 64, 112-122
15. Ohto, U., Mizutani, R., Nakamura, M., Adachi, H., and Satow, Y. (2004) J Synchrotron Radiat 11, 105-108

EXEMPLARY EMBODIMENTS

[0369] The following are exemplary embodiments of the invention as defined in the claims as well as reference embodiments:

1. An isolated human or humanized monoclonal antibody, or antigen-binding fragment thereof, which binds human NKG2D (hNKG2D).
2. The antibody or antigen-binding fragment of the preceding embodiment, which reduces hNKG2D-mediated activation of an hNKG2D-expressing NK or T cell.
3. The antibody or antigen-binding fragment of any preceding embodiment, which competes with at least one hNKG2D ligand in binding to hNKG2D.
4. The antibody or antigen-binding fragment of the preceding embodiment, wherein the ligand is MICA.
5. The antibody or antigen-binding fragment of any preceding embodiment, which reduces the amount of NKG2D on the surface of an NKG2D-expressing NK or T cell.
6. The antibody or antigen-binding fragment of any preceding embodiment, which, when immobilized, does not significantly co-stimulate CD3-triggered proliferation of peripheral blood mononuclear cells (PBMCs).
7. The antibody or antigen-binding fragment of any preceding embodiment, which, when immobilized, has no significant agonistic effect on hNKG2D-mediated activation of an hNKG2D-expressing NK or T cell.
8. The antibody or antigen-binding fragment of any preceding embodiment, which binds to cynomolgous and rhesus NKG2D.
9. The antibody of any preceding embodiment, which binds to hNKG2D with a KD of 1 nM or less.

10. 10. The antibody of any preceding embodiment, which binds to hNKG2D with a KD of 0.1 nM or less.
11. 11. The antibody or antigen-binding fragment of any preceding embodiment, which, when added to NKG2D-expressing NK or T cells, cross-links no more than two hNKG2D dimers.
12. 12. The antibody or antigen-binding fragment of any preceding embodiment, which binds strongly to only a first hNKG2D monomer in an hNKG2D dimer.
13. 13. The antibody or antigen-binding fragment of the preceding embodiment, which, when bound to the first hNKG2D monomer, blocks binding of the antibody or antigen-binding fragment to the second hNKG2D monomer.
14. 14. The antibody or antigen-binding fragment of any preceding embodiment, for which the ratio of the solvent-exclusion areas on the first and second hNKG2D monomers in an hNKG2D dimer is about 2:1 or more.
15. 15. The antibody or antigen-binding fragment of the preceding embodiment, for which the ratio is about 3:1 or more.
16. 16. The antibody or antigen-binding fragment of any preceding embodiment, which competes with a reference antibody in binding to NKG2D, wherein the reference antibody is selected from the group consisting of:
 1. (a) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:11 and a light-chain variable region comprising the sequence of SEQ ID NO:12;
 2. (b) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:13 and a light-chain variable region comprising the sequence of SEQ ID NO:14;
 3. (c) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:44 and a light-chain variable region comprising the sequence of SEQ ID NO:45; and
 4. (d) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:46 and a light-chain variable region comprising the sequence of SEQ ID NO:47.
17. 17. The antibody or antigen-binding fragment of any of the preceding embodiments, which binds to the same epitope of NKG2D as a reference antibody, wherein the reference antibody is selected from the group consisting of:
 1. (a) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:11 and a light-chain variable region comprising the sequence of SEQ ID NO:12;
 2. (b) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:13 and a light-chain variable region comprising the sequence of SEQ ID NO:14;
 3. (c) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:44 and a light-chain variable region comprising the sequence of SEQ ID NO:45; and
 4. (d) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:46 and a light-chain variable region comprising the sequence of SEQ ID NO:47.
18. 18. The antibody or antigen-binding fragment of any of the preceding embodiments, which binds to NKG2D with substantially the same KD as a reference antibody, wherein the reference antibody is selected from the group consisting of:
 1. (a) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:11 and a light-chain variable region comprising the sequence of SEQ ID NO:12;
 2. (b) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:13 and a light-chain variable region comprising the sequence of SEQ ID NO:14;
 3. (c) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:44 and a light-chain variable region comprising the sequence of SEQ ID NO:45; and
 4. (d) an antibody comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:46 and a light-chain variable region comprising the sequence of SEQ ID NO:47.
19. 19. The antibody or antigen-binding fragment of any of embodiments 16-18, wherein the reference antibody comprises a heavy-chain variable region comprising the sequence of SEQ ID NO:44 and a light-chain variable region comprising the sequence of SEQ ID NO:45.
20. 20. The antibody or antigen-binding fragment of the preceding embodiment, which binds to an epitope comprising Lys 150, Ser 151, Tyr 152, Thr 180, **Ile** 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and/or Thr 208 of SEQ ID NO:2.
21. 21. The antibody or antigen-binding fragment of the preceding embodiment, which binds to an epitope comprising 5 or more residues selected from Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of SEQ ID NO:2.
22. 22. The antibody or antigen-binding fragment of the preceding embodiment, which binds to an epitope comprising Lys 150, Ser 151, Tyr 152, Thr 180, Ile 181, Ile 182, Glu 183, Met 184, Gln 185, Leu 191, Lys 197, Tyr 199, Glu 201, Thr 205, Pro 206, Asn 207 and Thr 208 of SEQ ID NO:2.
23. 23. The antibody or antigen-binding fragment of any of embodiments 16-18, wherein the reference antibody comprises a heavy-chain variable region comprising the sequence of SEQ ID NO:46 and a light-chain variable region comprising the sequence of SEQ ID NO:47.
24. 24. The antibody or antigen-binding fragment of any preceding embodiment, comprising a heavy chain variable region that is the product of or derived from a set of human genes comprising:
 1. (a) VH3-21, D3-9, and JH4 genes;

2. (b) VH3_20, D3-10, and JH6 genes;
 3. (c) VH4_59, D7_27_R3, and JH3 genes; or
 4. (d) VH5_51, D3_10_R3 and JH4 genes.
25. 25. The antibody or antigen-binding fragment of any preceding embodiment, comprising a light-chain variable region that is the product of or derived from a set of human genes comprising
 1. (a) VKI_L15 and JK2 genes;
 2. (b) VKIII_A27 and JK3 genes;
 3. (c) VKIII_A27 and JK1 genes; or
 4. (d) VKIII_L6 and JK1 genes.
 26. 26. The antibody or antigen-binding fragment of any preceding embodiment, comprising a heavy chain variable region that is the product of or derived from a set of human genes comprising VH4_59, D7_27_R3, and JH3 genes, and a light-chain variable region that is the product of or derived from a set of human genes comprising VKIII_A27 and JK1 genes.
 27. 27. The antibody or antigen-binding fragment of any preceding embodiment, comprising a paratope comprising a residue corresponding to Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 or Asp 99 of SEQ ID NO:44 or Tyr 33 or Trp 97 of SEQ ID NO:45, or any combination thereof.
 28. 28. The antibody or antigen-binding fragment of the preceding embodiment, comprising a paratope comprising at least 5 residues selected from the residues corresponding to Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 or Asp 99 of SEQ ID NO:44 or Tyr 33 or Trp 97 of SEQ ID NO:45, or any combination thereof.
 29. 29. The antibody or antigen-binding fragment of the preceding embodiment, comprising a paratope comprising residues corresponding to Gln 1, Asp 26, Asp 27, Ser 30, Ser 31, Tyr 32, Tyr 33, His 50, Ser 52, Tyr 53, Ser 54, Ser 56, Ala 57, Asn 58, Trp 98 and Asp 99 of SEQ ID NO:44 and Tyr 33 and Trp 97 of SEQ ID NO:45.
 30. 30. The antibody or antigen-binding fragment of any preceding embodiment, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:48;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:49; and
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:50.
 31. 31. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:51;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:52; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:53.
 32. 32. The antibody or antigen-binding fragment of any preceding embodiment, comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:44, and a light-chain variable region comprising the sequence of SEQ ID NO:45.
 33. 33. The antibody or antigen-binding fragment of any of embodiments 1-25, comprising a heavy chain variable region that is the product of or derived from a set of human genes comprising VH5_51, D3_10_R3 and JH4 genes, and a light-chain variable region that is the product of or derived from a set of human genes comprising VKIII_L6 and JK1 genes.
 34. 34. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:54;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:55; and
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:56.
 35. 35. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:57;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:58; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:59.
 36. 36. The antibody or antigen-binding fragment of the preceding embodiment, comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:46, and a light-chain variable region comprising the sequence of SEQ ID NO:47.
 37. 37. The antibody or antigen-binding fragment of embodiment 1, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:15;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:16; and
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:17.
 38. 38. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:18;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:19; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:20.
 39. 39. The antibody or antigen-binding fragment of the preceding embodiment, comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:11, and a light-chain variable region comprising the sequence of SEQ ID NO:12.
 40. 40. The antibody or antigen-binding fragment of embodiment 1, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:21;

2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:22; and
3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:23.
41. 41. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:24;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:25; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:26.
42. 42. The antibody or antigen-binding fragment of the preceding embodiment, comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:13, and a light-chain variable region comprising the sequence of SEQ ID NO:14.
43. 43. The antibody or antigen-binding fragment of embodiment 1, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:48;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:49; and
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:50.
44. 44. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:51;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:52; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:53.
45. 45. The antibody or antigen-binding fragment of embodiment 1, comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:44, and a light-chain variable region comprising the sequence of SEQ ID NO:45.
46. 46. The antibody or antigen-binding fragment of embodiment 1, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:54;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:55; and
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:56.
47. 47. The antibody or antigen-binding fragment of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:57;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:58; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:59.
48. 48. The antibody or antigen-binding fragment of the preceding embodiment, comprising a heavy-chain variable region comprising the sequence of SEQ ID NO:46, and a light-chain variable region comprising the sequence of SEQ ID NO:47.
49. 49. The antibody or antigen-binding fragment of embodiment 1, comprising
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:15, 21, 48, or 54;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:16, 22, 49, or 55;
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:17, 23, 50, or 56;
 4. (d) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:18, 24, 51, or 57;
 5. (e) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:19, 25, 52, or 58; and
 6. (f) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:20, 26, 53, or 59;
 with no more than 8 conservative amino acid substitutions.
50. 50. The antibody or antigen-binding fragment of the preceding embodiment, comprising no more than 5 amino acid substitutions.
51. 51. The antibody of any preceding embodiment, which is human.
52. 52. The antibody or antigen-binding fragment of any preceding embodiment, which is bivalent.
53. 53. The antibody of the preceding embodiment, which is an IgG1, IgG2, IgG3, or IgG4 antibody.
54. 54. The antibody of the preceding embodiment, which is an IgG4 antibody.
55. 55. The antibody of any preceding embodiments, which comprises an S241P mutation in a VH chain.
56. 56. An isolated antibody comprising a heavy-chain sequence comprising the sequence of SEQ ID NO:7 and a light-chain sequence comprising the sequence of SEQ ID NO:8.
57. 57. An isolated antibody comprising a heavy-chain sequence comprising the sequence of SEQ ID NO:9 and a light-chain sequence comprising the sequence of SEQ ID NO:10.
58. 58. An isolated antibody comprising a heavy-chain sequence comprising the sequence of SEQ ID NO:40 and a light-chain sequence comprising the sequence of SEQ ID NO:41.
59. 59. An isolated antibody comprising a heavy-chain sequence comprising the sequence of SEQ ID NO:42 and a light-chain sequence comprising the sequence of SEQ ID NO:43.
60. 60. An isolated antibody that competes more with 16F16, 16F31, MS, or 21 F2 than with any of ON72, BAT221, 5C6, 1D11, ECM217, or 149810 in binding to hNKG2D, and prevents NKG2D-mediated activation of an NKG2D-expressing NK or T cell.
61. 61. The isolated antibody of the preceding embodiment, which competes more with MS than with ON72, BAT221, 5C6, 1D11, ECM217, or 149810.
62. 62. The isolated antibody of embodiment 60, which competes more with 21 F2 than with ON72, BAT221, 5C6, 1D11, ECM217, or 149810.

63. 63. An isolated antibody that binds cell surface-associated hNKG2D on an NK-cell preparation with a first half maximal effective concentration (EC50), and reduces killing of target cells expressing NKG2D-ligand mediated by the NK cell preparation with a second EC50, wherein the second EC50 is substantially lower than the first EC50.
64. 64. The antibody of the preceding embodiment, wherein the NK-cell preparation is NK-92 or NKL cells.
65. 65. The antibody of the preceding embodiment, wherein the ligand is ULBP-3 or MICA.
66. 66. The antibody of the preceding embodiment, wherein the second EC50 is no more than 80% of the first EC50.
67. 67. The antibody of the preceding embodiment, wherein the second EC50 is no more than 50% of the first EC50.
68. 68. An isolated antibody that binds hNKG2D, which antibody achieves its maximum reduction of NK-cell mediated killing of target cells expressing a ligand at a concentration lower than a concentration substantially saturating cell-surface associated NKG2D.
69. 69. An isolated antibody that binds hNKG2D, prevents NKG2D-mediated activation of an NKG2D-expressing NK or T cell, and is capable of downmodulating more than 70% of cell-surface-associated NKG2D.
70. 70. The antibody of the preceding embodiment, which is capable of downmodulating less than 90% of cell-surface-associated NKG2D of an NK cell.
71. 71. An isolated antibody that binds hNKG2D, and which binds to human and cynomolgous NKG2D-expressing cells with similar affinity.
72. 72. The antibody of the preceding embodiment, which has an EC50 for binding to cynomolgous CD8+ T cells in a PBMC preparation that is at least 50% of its EC50 for binding to human CD8+ T cells in a PBMC preparation.
73. 73. The antibody of the preceding embodiment, which has an EC50 for binding to cynomolgous CD8+ T cells in a PBMC preparation that is at least 65% of its EC50 for binding to human CD8+ T cells in a PBMC preparation.
74. 74. The isolated antibody of any preceding embodiments, which has an affinity for human NKG2D of no more than 0.1 nM.
75. 75. The antibody of any preceding embodiment, which has an affinity for human NKG2D of no more than 10 pM.
76. 76. The antibody of any of embodiments 60-75, comprising:
 1. (a) a heavy-chain variable region CDR1 comprising the sequence of SEQ ID NO:48 or 54;
 2. (b) a heavy-chain variable region CDR2 comprising the sequence of SEQ ID NO:49 or 55; and
 3. (c) a heavy-chain variable region CDR3 comprising the sequence of SEQ ID NO:50 or 56.
77. 77. The antibody of the preceding embodiment, comprising:
 1. (a) a light-chain variable region CDR1 comprising the sequence of SEQ ID NO:51 or 57;
 2. (b) a light-chain variable region CDR2 comprising the sequence of SEQ ID NO:52 or 58; and
 3. (c) a light-chain variable region CDR3 comprising the sequence of SEQ ID NO:53 or 59.
78. 78. The antibody of any of embodiments 60-77, which is human.
79. 79. An antigen-binding fragment of the antibody of any of embodiments 60-78.
80. 80. A nucleic acid encoding a variable-heavy or variable-light chain of the antibody or antigen-binding fragment of any preceding embodiment.
81. 81. An expression vector comprising the nucleic acid of the preceding embodiment.
82. 82. A host cell comprising the expression vector of the preceding embodiment.
83. 83. A host cell producing the antibody or antigen-binding fragment of any of embodiments 1-79.
84. 84. The host cell of any of embodiments 82 and 83, which is a CHO cell.
85. 85. A method of producing an anti-NKG2D antibody or antigen-binding fragment comprising culturing the host cell of any of embodiments 82 to 84 under suitable conditions and recovering said antibody or antigen-binding fragment thereof.
86. 86. A method of producing a variant anti-NKG2D antibody, or an antigen-binding fragment thereof, comprising
 1. (a) providing a heavy-chain variable region comprising a CDR1 sequence selected from SEQ ID NOS:15, 21, 48, and 54, a CDR2 sequence selected from SEQ ID NOS:16, 22, 49 and 55, and a CDR3 sequence selected from SEQ ID NOS:17, 23, 50 and 56;
 2. (b) providing a light-chain variable region comprising a CDR1 sequence selected from SEQ ID NOS:18, 24, 51 and 57, a CDR2 sequence selected from SEQ ID NOS:19, 25, 52 and 58, and a CDR3 sequence selected from SEQ ID NOS:20, 26, 53, and 59;
 3. (c) altering up to 8 amino acid residues in each of the heavy- and light chain variable regions to produce altered heavy- and light-chain variable regions; and
 4. (d) expressing a variant anti-NKG2D antibody comprising the altered heavy- and light-chain variable regions in a host cell.
87. 87. An immunoconjugate comprising the antibody or antigen-binding fragment of any of embodiments 1-79, linked to a therapeutic agent.
88. 88. A multispecific molecule comprising the antibody or antigen-binding fragment of any of embodiments 1-79, linked to a second moiety having a different binding specificity than the antibody.
89. 89. The multispecific molecule of the preceding embodiment, wherein the second moiety is a second antibody, or antigen-binding fragment thereof.

90. 90. A composition comprising the antibody or antigen-binding fragment of any of embodiments 1-79, and a pharmaceutically acceptable carrier.
91. 91. A method for preventing NKG2D-mediated activation of an NKG2D-expressing NK or T cell, comprising contacting the NK or T cell with the antibody or antigen-binding fragment of any of embodiments 1-79, wherein the antibody or antigen-binding fragment competes with at least one NKG2D ligand in binding to NKG2D.
92. 92. The method of the preceding embodiment, wherein the NKG2D ligand is MICA.
93. 93. The use of the antibody or antigen-binding fragment of any of embodiments 1-79 for reducing the amount of NKG2D on the surface of an NKG2D-expressing NK or T cell, comprising contacting the NK or T cell with the antibody or antigen-binding fragment of any of embodiments 1-79, wherein the antibody or antigen-binding fragment competes with at least one NKG2D ligand in binding to human NKG2D.
94. 94. The use of the antibody or antigen-binding fragment of any of embodiments 1-79 for treating an inflammatory or autoimmune disorder, comprising administering the composition of embodiment 90 to a human subject suffering from or at risk for an inflammatory or autoimmune disorder.
95. 95. The use of the antibody or antigen-binding fragment according to the preceding embodiment, wherein the patient is suffering from or at risk for an autoimmune disorder.
96. 96. The use of the antibody or antigen-binding fragment according to the preceding embodiment, wherein the autoimmune disorder is rheumatoid arthritis.
97. 97. The use of the antibody or antigen-binding fragment according to embodiment 95, wherein the disorder is multiple sclerosis.
98. 98. The use of the antibody or antigen-binding fragment according to embodiment 95, wherein the disorder is systemic lupus erythematosus (SLE).
99. 99. The use of the antibody or antigen-binding fragment according to embodiment 95, wherein the disorder is psoriasis.
100. 100. The use of the antibody or antigen-binding fragment according to embodiment 95, wherein the disorder is celiac disease.
101. 101. The use of the antibody or antigen-binding fragment according to embodiment 95, wherein the disorder is an inflammatory bowel disease.
102. 102. The use of the antibody or antigen-binding fragment according to the preceding embodiment, wherein the inflammatory bowel disease is ulcerative colitis.
103. 103. The use of the antibody or antigen-binding fragment according to embodiment 101, wherein the inflammatory bowel disease is Crohn's disease.
104. 104. The use of the antibody or antigen-binding fragment according to embodiment 94, wherein the human subject is suffering from or at risk for transplant rejection or graft-versus-host disease.
105. 105. The use of the antibody or antigen-binding fragment according to the preceding embodiment, wherein the transplant is a heart transplant or a bone marrow transplant.
106. 106. The use of the antibody or antigen-binding fragment according to any of embodiments 94 to 105, further comprising administering a second anti-inflammatory agent.
107. 107. The use of the antibody or antigen-binding fragment according to the preceding embodiment, wherein the second anti-inflammatory agent is selected from an immuno-suppressant, an analgesic, an anti-angiogenic agent, a corticosteroid, a B-cell depletion agent, a B-cell antagonist, a T-cell antagonist, a complement-inhibiting agent, an anti-cytokine agent, and an anti-cytokine receptor agent, and combinations thereof.
108. 108. The use of the antibody or antigen-binding fragment according to any of embodiments 106 and 107, wherein the second anti-inflammatory agent is an antagonist of TNFalpha activity.
109. 109. The use of the antibody or antigen-binding fragment for treating an inflammatory or autoimmune disorder, comprising administering the composition of embodiment 90 to a human subject at risk for an inflammatory or autoimmune disorder.
110. 110. The use of the antibody or antigen-binding fragment of the preceding embodiment, wherein the human subject is at risk for transplant rejection or graft-versus-host disease.
111. 111. The use of the antibody or antigen-binding fragment of any of embodiments 1-79, in the preparation of a medicament to treat an inflammatory or autoimmune disorder in a human subject.
112. 112. The antibody or antigen-binding fragment of any of embodiments 1-79, for use in treating an inflammatory or autoimmune disorder in a human subject.

[0370] All headings and sub-headings are used herein for convenience only and should not be construed as limiting the invention in any way,

[0371] Any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

[0372] The terms "a" and "an" and "the" and similar referents as used in the context of describing the invention are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context.

[0373] Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. Unless otherwise stated, all exact values provided herein are representative of corresponding approximate values (e.g., all exact exemplary values provided with respect to a particular factor or measurement can be considered to also provide a corresponding approximate measurement, modified by "about," where appropriate).

[0374] All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context.

[0375] The use of any and all examples, or exemplary language (e.g., "such as") provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise indicated. No language in the specification should be construed as indicating any element is essential to the practice of the invention unless as much is explicitly stated.

[0376] The citation of patent documents herein is done for convenience only and does not reflect any view of the validity, patentability and/or enforceability of such patent documents,

[0377] The description herein of any aspect or embodiment of the invention using terms such as "comprising", "having", "including" or "containing" with reference to an element or elements is intended to provide support for a similar aspect or embodiment of the invention that "consists of", "consists essentially of", or substantially comprises" that particular element or elements, unless otherwise stated or clearly contradicted by context (e.g., a composition described herein as comprising a particular element should be understood as also describing a composition consisting of that element, unless otherwise stated or clearly contradicted by context).

SEQUENCE LISTING

[0378]

<110> Novo Nordisk A/S

<120> ANTIBODIES AGAINST HUMAN NKG2D AND USES THEREOF

<130> 7923.504-WO

<160> 71

<170> PatentIn version 3.5

<210> 1

<211> 1770

<212> DNA

<213> Homo sapiens

<400> 1

```

gaggagtgga ttacatatc caacagttgt tattacattg gtaaggaaag aagaacttgg      60
gaagaaagag tttgctggcc tgtgcttoga agaactctga tctgctttct atagataatg    120
aggaagaaat ggtatgtgtg gggacttccc agttggctgt aagttgccat ttgaactaaa    180
cgaatatgat caggaactga ggacatatct aaattttcta gttttataga aggcctttat    240
ccacaagaat caagatcttc cctctctgag caggaatcct ttgtgcattg aagactttag    300
attcctctct gcggtagacg tgcacttata agtatattgat ggggtggatt cgtggtcgga    360
ggtctcgaca cagctgggag atgagtgaat ttcataatta taacttggat ctgaagaaga    420
gtgatttttc aacacgatgg caaaagcaaa gatgtccagt agtcaaaagc aaatgtagag    480
aaaatgcate tccatttttt ttctgctgct tcatcgctgt agccatggga atccgtttca    540
ttattatggt agcaatatgg agtgctgtat tccataaactc attattcaac caagaagttc    600
aaattccctt gaccgaaagt tactgtggcc catgtcctaa aaactggata tgttacaaaa    660
ataactgcta ccaatttttt gatgagagta aaaactggtg tgagagccag gcttcttgta    720
tgtctcaaaa tgccagcctt ctgaaagtat acagcaaaga ggaccaggat ttacttaaac    780
tggtgaagtc atatcattgg atgggactag tacacattcc aacaaatgga tcttggcagt    840
gggaagatgg ctccattctc tcaccaacc tactaacaat aattgaaatg cagaagggag    900
actgtgactc ctatgctctg agcttttaag gctatataga aaactgttca actccaaata    960
catacatctg catgcaaagg actgtgtaaa gatgatcaac catctcaata aaagccagga   1020
acagagaaga gattacacca gcggtaacac tgccaaccga gactaaagga aacaacaaaa   1080
aacaggacaa aatgacccaa gactgtcaga tttcttagac tccacaggac caaaccatag   1140
aacaatttca ctgcaaacat gcatgattct ccaagacaaa agaagagaga tcctaaaggc   1200

aattcagata tcccacaggc tgctctctcc accacaagcc cagagtggat gggctggggg   1260
aggggtgctg ttttaatttc taaaggtagg accaacaccc aggggatcag tgaagggaaga   1320
gaaggccagc agatcagtga gagtgcaacc ccaccctcca caggaaattg cctcatgggc   1380
agggccacag cagagagaca cagcatgggc agtgcccttc ctgctctgtg gggtcatgct   1440
gccactttta atgggtcctc caccacacgg ggtcaggggg gtggtgctgc ccagtgggc   1500
catgattatc ttaaaggcat tattotccag cottaagato ttaggaagctt tcctttgcta   1560
tgatttgtae ttgcttgagt cccatgactg tttctcttcc tctctttctt ccttttgtaa   1620
tagtaatatc catcctatgt ttgtccactc attgtatttt ggaagcacat aacttgtttg   1680
gtttcacagg ttcacagtta agaaggaatt ttgcctctga ataaatagaa tcttgagtct   1740
catgcaaaaa aaaaaaaaaa aaaaaaaaaa
catgcaaaaa aaaaaaaaaa aaaaaaaaaa      1770

```

<210> 2

<211> 216

<212> PRT

<213> Homo sapiens

<400> 2

```

Met Gly Trp Ile Arg Gly Arg Arg Ser Arg His Ser Trp Glu Met Ser
1      5      10      15

Glu Phe His Asn Tyr Asn Leu Asp Leu Lys Lys Ser Asp Phe Ser Thr
      20      25      30

Arg Trp Gln Lys Gln Arg Cys Pro Val Val Lys Ser Lys Cys Arg Glu
      35      40      45

Asn Ala Ser Pro Phe Phe Phe Cys Cys Phe Ile Ala Val Ala Met Gly
      50      55      60

Ile Arg Phe Ile Ile Met Val Ala Ile Trp Ser Ala Val Phe Leu Asn
65      70      75      80

Ser Leu Phe Asn Gln Glu Val Gln Ile Pro Leu Thr Glu Ser Tyr Cys
      85      90      95

Gly Pro Cys Pro Lys Asn Trp Ile Cys Tyr Lys Asn Asn Cys Tyr Gln
      100      105      110

Phe Phe Asp Glu Ser Lys Asn Trp Tyr Glu Ser Gln Ala Ser Cys Met
      115      120      125

Ser Gln Asn Ala Ser Leu Leu Lys Val Tyr Ser Lys Glu Asp Gln Asp
      130      135      140

Leu Leu Lys Leu Val Lys Ser Tyr His Trp Met Gly Leu Val His Ile
145      150      155      160

Pro Thr Asn Gly Ser Trp Gln Trp Glu Asp Gly Ser Ile Leu Ser Pro
      165      170      175

Asn Leu Leu Thr Ile Ile Glu Met Gln Lys Gly Asp Cys Ala Leu Tyr
      180      185      190

Ala Ser Ser Phe Lys Gly Tyr Ile Glu Asn Cys Ser Thr Pro Asn Thr
      195      200      205

Tyr Ile Cys Met Gln Arg Thr Val
      210      215

```

<210> 3

<211> 1344

<212> DNA

<213> Homo sapiens

<400> 3

```

gaggtgcagc tgggtggagtc tgggggaggc ctggtcaagc ctgggggggtc cctgagactc      60
tcctgtgcag cctctggatt cacccttcagt agctatagca tgaactgggt ccgccaggct      120
ccagggaagg ggotggagtg ggtctcatcc attactagta gtagtagtta cataactac      180
gcagactcag tgaaggggccg attcaccatc tccagagaca acgccaagaa ctactgtat      240
ctgcaaatga acagcctgag agccgaggac acggtctgtt attactgtgc gagagatagg      300
cgatattttg actggttccc tcttgactac cggggccagg gaaccctggt caccgtctcc      360
tcagctagca ccaaggggccc atccgtcttc cccctggcgc cctgctecag gagcacctcc      420
gagagcacag ccgccttggg ctgcctggtc aaggactact tccccgaacc ggtgacgggtg      480
tcgtggaact caggcgccct gaccagcggc gtgcacacct tccgggtgtt cctacagtcc      540
tcaggactct actccctcag cagcgtggtg accgtgccct ccagcagctt gggcargaaag      600
acctcacctt gcaacgtaga tcacaagccc agcaacacca aggtggacaa gagagttgag      660
tccaaatatg gtcccccatg cccaccatgc ccagcacctg agttcctggg gggaccatca      720

gtcttctctg tcccccaaaa acccaaggac actctcatga tctcccgac cctgaggtc      780
acgtgcgtgg tgggtggagtc gagccaggaa gaccccgagg tcagttcaa ctggtacgtg      840
gatggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagtt caacagcacg      900
taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaacgg caaggagtac      960
aagtgcaggg tctccaacaa aggcctcccc tcctccatcg agaaaacct ctccaaagcc      1020
aaagggcagc cccgagagcc acaggtgtac accctgcccc catcccagga ggagatgacc      1080
aagaaccagg tcagcctgac ctgcctggtc aaaggcttct accccagcga catcgccgtg      1140
gagtgggaga gcaatgggca gccggagaa aactacaaga ccacgcctcc cgtgctggac      1200
tccgacggct ccttcttctc ctacagcagg ctaaccgtgg acaagagcag gtggcaggag      1260
gggaatgtct tctcatgtct cgtgatgcat gaggctctgc acaaccacta cacacagaag      1320
agcctctccc tgtctctggg taaa                                     1344

```

<210> 4

<211> 642

<212> DNA

<213> Homo sapiens

<400> 4

```

gacatccaga tgaccagtc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc      60
atcacttgtc gggcgagtca ggggtattag agctggttag cctgggtatca gcagaaacca      120
gagaaagccc ctaagtcctt gatctatgct gcattccagt tgcaaaagtg ggtcccatca      180
aggttcagcg gcagtggatc tgggacagat ttcaactcca ccacagcag cctgcagcct      240
gaagattttg caacttatta ctgccaaacg tataatggtt acccgtacac ttttggccag      300
gggaccaagc tggagatcaa acgtacgggt gctgcaccat ctgtcttcat cttcccgcca      360
tctgatgagc agttgaaate tggaaactgcc tctgttgtgt gctgctgaa taacttctat      420
cccagagagg ccaaagtaca gtggaagggt gataacgccc tccaatcggg taactccag      480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg      540
ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc      600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gt                                     642

```

<210> 5

<211> 1347

<212> DNA

<213> Homo sapiens

<400> 5

```

gaggtgcagc tgggtggagtc tgggggaggt gtggtacggc ctgggggggtc cctgagactc      60
tcctgtgcag cctctggatt caectttgat gattatggca tgacctgggt ccgccaagct      120
ccagggaagg ggctggagtg ggtctctggt attaatlgga atggtggtag cacaggttat      180
gcagactctg tgaaggggccg attcaccatc tccagagaca acgccaagaa ctccctgtat      240
ctgcaaatga acagtctgag agccgaggac acggccttgt attactgtgc gagagagagg      300
gagttatact actactacta cggtttggac gtctggggcc aaggggaccac ggtcaccgtc      360
tcctcagcta gcaccaaggg cccatccgtc ttcccccctgg cgcctctgctc caggagcacc      420
tccgagagca cagccggcctt gggctgcctg gtcaaggact acttccccga accggtgacg      480
gtgtcgtgga actcaggcgc cctgaccagc ggcgtgcaca ccttcccggc tgtcctacag      540
tcctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcag cttgggcacg      600
aagacctaca cctgcaacgt agatcacaag cccagcaaca ccaaggtgga caagagagtt      660
gagtcctaat atggtccccc atgccacaca tgcccagcac ctgagttcct ggggggacca      720
tcagtcttcc tgttccccc aaacccaag gacctctca tgatctccc gaccctgag      780
gtcacgtgct tgggtggtgga cgtgagccag gaagacccc aggtccagtt caactggtac      840
gtggatggcg tggaggtgca taatgccaa acaaagccgc gggaggagca gttcaacagc      900
acgtaccgtg tggtcagcgt cctcacctc ctgcaccagg actggctgaa cggcaaggag      960
tacaagtgca aggtctccaa caaaggcctc ccgtctcca tcgagaaaa catctccaaa      1020
gccaaagggc agccccgaga gccacagggtg tacaccctgc ccccatccca ggaggagatg      1080
accaagaacc aggtcagcct gacctgcctg gtcaaaggct tctaccccag cgacatcgcc      1140
gtggagtggg agagcaatgg gcagccggag aacaactaca agaccacgcc tcccgtgctg      1200
gactccgacg gctccttctt cctctacagc aggctaaccg tggacaagag cagggtggcag      1260
gaggggaatg tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacacag      1320
aagagcctct cctgtctctt gggtaaa      1347

```

<210> 6

<211> 645

<212> DNA

<213> Homo sapiens

<400> 6

```

gaaattgtgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagccacc      60
ctctctgca gggccagtc gagtgtagc agcagctact tagcctggta ccagcagaaa      120
cctggccagg ctcccaggct cctcatctat ggtgcatcca gcagggccac tggcatccca      180
gacaggttca gtggcagtg gtctgggaca gacttcactc tcaccatcag cagactggag      240
cctgaagatt ttgcagtgt taactgtcag cagtatggta gctcaccatt cactttcggc      300
cctgggacca aagtggatat caaacgtacg gtggctgcac catctgtctt catcttcccg      360
ccatctgatg agcagttgaa atctggaact gcctctgttg tgtgctgct gaataacttc      420
tateccagag aggccaaagt acagtggaag gtggataacg cctccaatc gggtaacttc      480
caggagagtg tcacagagca ggacagcaag gacagcacct acagcctcag cagcaccctg      540
acgtgagaca aagcagacta cgagaaacac aaagtctacg cctgcgaagt ccccatcag      600
ggcctgagct cgcccgctac aaagagcttc aacaggggag agtgt      645

```

<210> 7

<211> 448

<212> PRT

<213> Homo sapiens

<400> 7

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ser Ile Thr Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asp Arg Arg Tyr Phe Asp Trp Phe Pro Leu Asp Tyr Arg Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser
 115 120 125
 Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala
 130 135 140
 Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val
 145 150 155 160
 Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala
 165 170 175
 Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val
 180 185 190
 Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His
 195 200 205
 Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly
 210 215 220
 Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser
 225 230 235 240
 Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 245 250 255
 Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro
 260 265 270
 Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
 275 280 285
 Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val
 290 295 300
 Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 305 310 315 320
 Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr
 325 330 335
 Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 340 345 350

Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
 355 360 365
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 370 375 380
 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 385 390 395 400
 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser
 405 410 415
 Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 420 425 430
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
 435 440 445
 <210> 8
 <211> 214
 <212> PRT
 <213> Homo sapiens
 <400> 8
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
 35 40 45
 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Gly Tyr Pro Tyr
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125
 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140
 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160
 Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205
 Phe Asn Arg Gly Glu Cys
 210

<210> 9
 <211> 449
 <212> PRT
 <213> Homo sapiens

<400> 9

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Arg Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Gly Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Gly Ile Asn Trp Asn Gly Gly Ser Thr Gly Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Arg Glu Leu Tyr Tyr Tyr Tyr Tyr Gly Leu Asp Val Trp
 100 105 110
 Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125
 Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr
 130 135 140
 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160
 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
 165 170 175
 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
 180 185 190
 Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp
 195 200 205
 His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr
 210 215 220
 Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro
 225 230 235 240
 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
 245 250 255
 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp
 260 265 270
 Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
 275 280 285
 Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val
 290 295 300

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
305 310 315 320

Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys
325 330 335

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
340 345 350

Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
355 360 365

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
370 375 380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
385 390 395 400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys
405 410 415

Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu
420 425 430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly
435 440 445

Lys

<210> 10

<211> 215

<212> PRT

<213> Homo sapiens

<400> 10

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
20 25 30

Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu

35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95
 Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg Thr Val Ala
 100 105 110
 Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
 115 120 125
 Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
 130 135 140
 Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
 145 150 155 160
 Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
 165 170 175
 Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
 180 185 190
 Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
 195 200 205
 Ser Phe Asn Arg Gly Glu Cys
 210 215

<210> 11

<211> 121

<212> PRT

<213> Homo sapiens

<400> 11

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ser Ile Thr Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asp Arg Arg Tyr Phe Asp Trp Phe Pro Leu Asp Tyr Arg Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 12

<211> 108

<212> PRT

<213> Homo sapiens

<400> 12

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
 35 40 45
 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Gly Tyr Pro Tyr
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
 100 105

<210> 13

<211> 122

<212> PRT

<213> Homo sapiens

<400> 13

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Arg Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Gly Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Gly Ile Asn Trp Asn Gly Gly Ser Thr Gly Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Arg Glu Leu Tyr Tyr Tyr Tyr Gly Leu Asp Val Trp
 100 105 110
 Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> 14

<211> 109

<212> PRT

<213> Homo sapiens

<400> 14

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95
 Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys Arg
 100 105

<210> 15

<211> 5

<212> PRT

<213> Homo sapiens

<400> 15

Ser Tyr Ser Met Asn
 1 5

<210> 16

<211> 17

<212> PRT

<213> Homo sapiens

<400> 16

Ser Ile Thr Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 17

<211> 12

<212> PRT

<213> Homo sapiens

<400> 17

Asp Arg Arg Tyr Phe Asp Trp Phe Pro Leu Asp Tyr
 1 5 10

<210> 18

<211> 11

<212> PRT

<213> Homo sapiens

<400> 18

Arg Ala Ser Gln Gly Ile Ser Ser Trp Leu Ala
 1 5 10

<210> 19

<211> 7

<212> PRT

<213> Homo sapiens

<400> 19

Ala Ala Ser Ser Leu Gln Ser
 1 5

<210> 20

<211> 9

<212> PRT

<213> Homo sapiens

<400> 20

Gln Gln Tyr Asn Gly Tyr Pro Tyr Thr
1 5

<210> 21

<211> 5

<212> PRT

<213> Homo sapiens

<400> 21

Asp Tyr Gly Met Thr
1 5

<210> 22

<211> 17

<212> PRT

<213> Homo sapiens

<400> 22

Gly Ile Asn Trp Asn Gly Gly Ser Thr Gly Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 23

<211> 13

<212> PRT

<213> Homo sapiens

<400> 23

Glu Arg Glu Leu Tyr Tyr Tyr Tyr Gly Leu Asp Val
1 5 10

<210> 24

<211> 12

<212> PRT

<213> Homo sapiens

<400> 24

Arg Ala Ser Gln Ser Val Ser Ser Ser Tyr Leu Ala
1 5 10

<210> 25

<211> 7

<212> PRT

<213> Homo sapiens

<400> 25

Gly Ala Ser Ser Arg Ala Thr
1 5

<210> 26

<211> 9

<212> PRT

<213> Homo sapiens

<400> 26

Gln Gln Tyr Gly Ser Ser Pro Phe Thr
1 5

<210> 27

<211> 111

<212> PRT

<213> Homo sapiens

<400> 27

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ser Ile Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
100 105 110

<210> 28

<211> 107

<212> PRT

<213> Homo sapiens

<400> 28

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro Tyr
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 29

<211> 118

<212> PRT

<213> Homo sapiens

<400> 29

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Arg Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30
 Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Gly Ile Asn Trp Asn Gly Gly Ser Thr Gly Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr His Cys
 85 90 95
 Ala Arg Tyr Tyr Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr
 100 105 110
 Thr Val Thr Val Ser Ser
 115

<210> 30

<211> 108

<212> PRT

<213> Homo sapiens

<400> 30

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95
 Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 100 105

<210> 31

<211> 98

<212> PRT

<213> Homo sapiens

<400> 31

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ser Ile Ser Ser Ser Ser Tyr Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg

<210> 32

<211> 5

<212> PRT

<213> Homo sapiens

<400> 32

Arg Tyr Phe Asp Trp
1 5

<210> 33

<211> 13

<212> PRT

<213> Homo sapiens

<400> 33

Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
1 5 10

<210> 34

<211> 95

<212> PRT

<213> Homo sapiens

<400> 34

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser Leu Ile
35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Tyr Pro
85 90 95

<210> 35

<211> 12

<212> PRT

<213> Homo sapiens

<400> 35

Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
1 5 10

<210> 36

<211> 98

<212> PRT

<213> Homo sapiens

<400> 36

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Arg Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30

Gly Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Gly Ile Asn Trp Asn Gly Gly Ser Thr Gly Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Leu Tyr His Cys
 85 90 95

Ala Arg

<210> 37

<211> 20

<212> PRT

<213> Homo sapiens

<400> 37

Tyr Tyr Tyr Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val
 1 5 10 15

Thr Val Ser Ser
 20

<210> 38

<211> 96

<212> PRT

<213> Homo sapiens

<400> 38

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30

Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45

Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80

Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95

<210> 39

<211> 12

<212> PRT

<213> Homo sapiens

<400> 39

Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 1 5 10

<210> 40

<211> 442

<212> PRT

<213> Homo sapiens

<400> 40

Gln Val His Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Asp Asp Ser Ile Ser Ser Tyr
 20 25 30
 Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 Gly His Ile Ser Tyr Ser Gly Ser Ala Asn Tyr Asn Pro Ser Leu Lys
 50 55 60
 Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80
 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95
 Asn Trp Asp Asp Ala Phe Asn Ile Trp Gly Gln Gly Thr Met Val Thr
 100 105 110
 Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro
 115 120 125
 Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val
 130 135 140
 Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala
 145 150 155 160
 Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly
 165 170 175
 Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly
 180 185 190
 Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys
 195 200 205
 Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys
 210 215 220
 Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 225 230 235 240
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 245 250 255

Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp
 260 265 270
 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 275 280 285
 Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 290 295 300
 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 305 310 315 320
 Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 325 330 335
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu
 340 345 350
 Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 355 360 365
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 370 375 380
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 385 390 395 400
 Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn
 405 410 415
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 420 425 430
 Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
 435 440
 <210> 41
 <211> 215
 <212> PRT
 <213> Homo sapiens
 <400> 41
 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95
 Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala
 100 105 110
 Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser
 115 120 125
 Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu
 130 135 140
 Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser
 145 150 155 160
 Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu
 165 170 175
 Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val
 180 185 190
 Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
 195 200 205
 Ser Phe Asn Arg Gly Glu Cys
 210 215

<210> 42

<211> 450

<212> PRT

<213> Homo sapiens

<400> 42

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Glu Pro Gly Glu
 1 5 10 15
 Ser Leu Lys Ile Ser Cys Lys Asn Ser Gly Tyr Ser Phe Thr Asn Tyr
 20 25 30
 Trp Val Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
 35 40 45
 Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe
 50 55 60
 Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Asn Thr Ala Tyr
 65 70 75 80
 Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
 85 90 95
 Gly Arg Leu Thr Met Phe Arg Gly Ile Ile Ile Gly Tyr Phe Asp Tyr
 100 105 110
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly
 115 120 125
 Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser
 130 135 140
 Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
 145 150 155 160
 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
 165 170 175
 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
 180 185 190
 Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val
 195 200 205
 Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys

210	215	220
Tyr Gly Pro Pro Cys	Pro Pro Cys Pro Ala	Pro Glu Phe Leu Gly Gly
225	230	235 240
Pro Ser Val Phe Leu	Phe Pro Pro Lys	Pro Lys Asp Thr Leu Met Ile
245	250	255
Ser Arg Thr	Pro Glu Val Thr Cys Val	Val Val Asp Val Ser Gln Glu
260	265	270
Asp Pro Glu Val Gln	Phe Asn Trp Tyr Val Asp	Gly Val Glu Val His
275	280	285
Asn Ala Lys Thr Lys	Pro Arg Glu Glu Gln Phe	Asn Ser Thr Tyr Arg
290	295	300
Val Val Ser Val Leu	Thr Val Leu His Gln Asp	Trp Leu Asn Gly Lys
305	310	315 320
Glu Tyr Lys Cys Lys	Val Ser Asn Lys	Gly Leu Pro Ser Ser Ile Glu
325	330	335
Lys Thr Ile Ser Lys	Ala Lys Gly Gln Pro Arg	Glu Pro Gln Val Tyr
340	345	350
Thr Leu Pro Pro Ser	Gln Glu Glu Met Thr Lys	Asn Gln Val Ser Leu
355	360	365
Thr Cys Leu Val Lys	Gly Phe Tyr Pro Ser Asp	Ile Ala Val Glu Trp
370	375	380
Glu Ser Asn Gly Gln	Pro Glu Asn Asn Tyr Lys	Thr Thr Pro Pro Val
385	390	395 400
Leu Asp Ser Asp Gly	Ser Phe Phe Leu Tyr	Ser Arg Leu Thr Val Asp
405	410	415
Lys Ser Arg Trp Gln	Glu Gly Asn Val Phe Ser	Cys Ser Val Met His
420	425	430
Glu Ala Leu His Asn	His Tyr Thr Gln Lys	Ser Leu Ser Leu Ser Leu
435	440	445
	Gly Lys	
	450	

<210> 43

<211> 214

<212> PRT

<213> Homo sapiens

<400> 43

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Trp
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110
 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125
 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140
 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160
 Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175
 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205
 Phe Asn Arg Gly Glu Cys
 210

<210> 44

<211> 115

<212> PRT

<213> Homo sapiens

<400> 44

Gln Val His Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Asp Asp Ser Ile Ser Ser Tyr
 20 25 30
 Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 Gly His Ile Ser Tyr Ser Gly Ser Ala Asn Tyr Asn Pro Ser Leu Lys
 50 55 60
 Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80
 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95
 Asn Trp Asp Asp Ala Phe Asn Ile Trp Gly Gln Gly Thr Met Val Thr
 100 105 110
 Val Ser Ser
 115

<210> 45

<211> 108

<212> PRT

<213> Homo sapiens

<400> 45

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30
 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45
 Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80
 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95
 Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 46

<211> 123

<212> PRT

<213> Homo sapiens

<400> 46

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Glu Pro Gly Glu
 1 5 10 15
 Ser Leu Lys Ile Ser Cys Lys Asn Ser Gly Tyr Ser Phe Thr Asn Tyr
 20 25 30
 Trp Val Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
 35 40 45
 Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe
 50 55 60
 Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Asn Thr Ala Tyr
 65 70 75 80
 Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
 85 90 95
 Gly Arg Leu Thr Met Phe Arg Gly Ile Ile Ile Gly Tyr Phe Asp Tyr
 100 105 110
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 47

<211> 107

<212> PRT

<213> Homo sapiens

<400> 47

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Trp
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 48

<211> 5

<212> PRT

<213> Homo sapiens

<400> 48

Ser Tyr Tyr Trp Ser
 1 5

<210> 49

<211> 16

<212> PRT

<213> Homo sapiens

<400> 49

His Ile Ser Tyr Ser Gly Ser Ala Asn Tyr Asn Pro Ser Leu Lys Ser
 1 5 10 15

<210> 50

<211> 7

<212> PRT

<213> Homo sapiens

<400> 50

Trp Asp Asp Ala Phe Asn Ile
 1 5

<210> 51

<211> 12

<212> PRT

<213> Homo sapiens

<400> 51

Arg Ala Ser Gln Ser Val Ser Ser Ser Tyr Leu Ala
 1 5 10

<210> 52

<211> 7

<212> PRT

<213> Homo sapiens

<400> 52

Gly Ala Ser Ser Arg Ala Thr
 1 5

<210> 53

<211> 9

<212> PRT

<213> Homo sapiens

<400> 53

Gln Gln Tyr Gly Ser Ser Pro Trp Thr
1 5

<210> 54

<211> 5

<212> PRT

<213> Homo sapiens

<400> 54

Asn Tyr Trp Val Gly
1 5

<210> 55

<211> 17

<212> PRT

<213> Homo sapiens

<400> 55

Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe Gln
1 5 10 15

Gly

<210> 56

<211> 14

<212> PRT

<213> Homo sapiens

<400> 56

Leu Thr Met Phe Arg Gly Ile Ile Ile Gly Tyr Phe Asp Tyr
1 5 10

<210> 57

<211> 11

<212> PRT

<213> Homo sapiens

<400> 57

Arg Ala Ser Gln Ser Val Ser Ser Tyr Leu Ala
1 5 10

<210> 58

<211> 7

<212> PRT

<213> Homo sapiens

<400> 58

Asp Ala Ser Asn Arg Ala Thr
1 5

<210> 59

<211> 9

<212> PRT

<213> Homo sapiens

<400> 59

Gln Gln Arg Ser Asn Trp Pro Trp Thr
1 5

<210> 60

<211> 116

<212> PRT

<213> Homo sapiens

<400> 60

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Tyr
 20 25 30

Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg Asn Trp Gly Asp Ala Phe Asp Ile Trp Gly Gln Gly Thr Met Val
 100 105 110

Thr Val Ser Ser
 115

<210> 61

<211> 108

<212> PRT

<213> Homo sapiens

<400> 61

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser
 20 25 30

Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu
 35 40 45

Ile Tyr Gly Ala Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu
 65 70 75 80

Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Gly Ser Ser Pro
 85 90 95

Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 62

<211> 122

<212> PRT

<213> Homo sapiens

<400> 62

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
 1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr
 20 25 30

Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
 35 40 45

Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe
 50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
 65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
 85 90 95

Ala Arg Ile Thr Met Val Arg Gly Val Ile Ile Tyr Phe Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 63

<211> 107

<212> PRT

<213> Homo sapiens

<400> 63

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro Trp
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 64

<211> 97

<212> PRT

<213> Homo sapiens

<400> 64

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Tyr
 20 25 30

Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg

<210> 65

<211> 16

<212> PRT

<213> Homo sapiens

<400> 65

Asp Ala Phe Asp Val Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser
 1 5 10 15

<210> 66

<211> 12

<212> PRT

<213> Homo sapiens

<400> 66

Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 1 5 10

<210> 67

<211> 98

<212> PRT

<213> Homo sapiens

<400> 67

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
 1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Tyr

20 25 30

Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
 35 40 45

Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser Pro Ser Phe
 50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
 65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
 85 90 95

Ala Arg

<210> 68

<211> 9

<212> PRT

<213> Homo sapiens

<400> 68

Ile Thr Met Val Arg Gly Val Ile Ile
 1 5

<210> 69

<211> 95

<212> PRT

<213> Homo sapiens

<400> 69

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser Asn Trp Pro
 85 90 95

<210> 70

<211> 220

<212> PRT

<213> Artificial

<220>

<223> Humanized sequence

<400> 70

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Glu Ile Asp Pro Ser Asp Ile Tyr Thr Asn Tyr Ala Gln Lys Phe
 50 55 60

Gln Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Gly Ile Tyr Asp Gly Tyr Tyr Val Tyr Gly Met Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125

Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr
 130 135 140

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160

Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
 165 170 175

Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
 180 185 190

Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp
 195 200 205

His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val
 210 215 220

<210> 71

<211> 214

<212> PRT

<213> Artificial

<220>

<223> Humanized sequence

<400> 71

```

Tyr Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15

Asp Arg Val Thr Ile Thr Cys Arg Ser Ser Gln Asp Ile Ser Asn Tyr
          20           25           30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
          35           40           45

Tyr Tyr Thr Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50           55           60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65           70           75           80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Gly Lys Thr Leu Pro Arg
          85           90           95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
          100          105          110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
          115          120          125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
          130          135          140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145          150          155          160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
          165          170          175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
          180          185          190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
          195          200          205

Phe Asn Arg Gly Glu Cys
          210

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US20050158307A \[0003\]](#)
- [WO2005097160A \[0003\]](#)
- [WO2005115517A \[0003\]](#)
- [WO2006024367A \[0003\]](#)
- [WO20060615A \[0004\] \[0320\]](#)
- [US2006280755A \[0005\]](#)
- [WO2007005874A \[0008\]](#)
- [WO9202190A \[0024\]](#)
- [US20060073137A \[0024\]](#)
- [US6750325B \[0024\]](#)
- [US6632927B \[0024\]](#)
- [US6639055B \[0024\]](#)

- [US6548640B \[0024\]](#)
- [US6407213B \[0024\]](#)
- [US6180370B \[0024\] \[0115\] \[0118\] \[0150\]](#)
- [US6054297A \[0024\]](#)
- [US5929212A \[0024\]](#)
- [US5895205A \[0024\]](#)
- [US5886152A \[0024\]](#)
- [US5877293A \[0024\]](#)
- [US5869619A \[0024\]](#)
- [US5821337A \[0024\]](#)
- [US5821123A \[0024\]](#)
- [US5770196A \[0024\]](#)
- [US5777085A \[0024\]](#)
- [US5766886A \[0024\]](#)
- [US5714350A \[0024\] \[0131\]](#)
- [US5693762A \[0024\] \[0115\] \[0118\] \[0150\]](#)
- [US5693761A \[0024\]](#)
- [US5530101A \[0024\] \[0115\] \[0118\] \[0150\]](#)
- [US5585089A \[0024\] \[0115\] \[0118\] \[0150\]](#)
- [US5225539A \[0024\] \[0115\] \[0150\]](#)
- [WO2005040219A \[0099\]](#)
- [US20050238646A \[0099\]](#)
- [US20020161201A \[0099\]](#)
- [WO199316185A \[0101\]](#)
- [US5571894A \[0101\]](#)
- [US5587458A \[0101\]](#)
- [US5641870A \[0101\]](#)
- [US6046310A \[0106\]](#)
- [US20030103984A \[0106\]](#)
- [EP1413316A \[0106\]](#)
- [US20040038339A \[0106\]](#)
- [WO2004056873A \[0106\]](#)
- [US5260203A \[0110\]](#)
- [US5455030A \[0110\]](#)
- [US4861175A \[0110\]](#)
- [US5132405A \[0110\]](#)
- [US5091513A \[0110\]](#)
- [US5476786A \[0110\]](#)
- [US5013653A \[0110\]](#)
- [US5258498A \[0110\]](#)
- [US5482858A \[0110\]](#)
- [US20030078385A \[0110\]](#)
- [US20030153043A \[0125\]](#)
- [US4816397A \[0127\]](#)
- [US5916771A \[0127\]](#)
- [US5677425A \[0128\]](#)
- [US6165745A \[0129\]](#)
- [US6277375B \[0129\]](#)
- [US5869046A \[0129\]](#)
- [US6121022A \[0129\]](#)
- [US5624821A \[0129\]](#)
- [US5648260A \[0129\]](#)
- [US6194551B \[0129\]](#)
- [WO9429351A \[0129\]](#)
- [WO0042072A \[0129\]](#)
- [US6350861B \[0131\]](#)
- [EP1176195A \[0131\]](#)
- [WO03035835A \[0131\]](#)

- [WO9954342A \[0131\]](#)
- [WO02092780A \[0132\]](#)
- [WO03074679A \[0133\]](#)
- [US6077499A \[0135\]](#)
- [US4766106A \[0141\]](#)
- [US4179337A \[0141\]](#)
- [US4495285A \[0141\]](#)
- [US4609546A \[0141\]](#)
- [EP154316A \[0141\]](#)
- [US0411494W \[0141\]](#)
- [EP401384A \[0141\]](#)
- [US4816567A \[0150\]](#)
- [US5545806A \[0151\]](#)
- [US5569825A \[0151\]](#)
- [US5625126A \[0151\]](#)
- [US5633425A \[0151\]](#)
- [US5789650A \[0151\]](#)
- [US5877397A \[0151\]](#)
- [US5661016A \[0151\]](#)
- [US5814318A \[0151\]](#)
- [US5874299A \[0151\]](#)
- [US5770429A \[0151\]](#)
- [US5545807A \[0151\]](#)
- [WO9203918A \[0151\]](#)
- [WO9312227A \[0151\]](#)
- [WO9425585A \[0151\]](#)
- [WO9713852A \[0151\]](#)
- [WO9824884A \[0151\] \[0153\]](#)
- [WO9945962A \[0151\]](#)
- [WO0114424PCT \[0151\]](#)
- [WO0243478A \[0151\] \[0290\] \[0290\]](#)
- [US5939598A \[0151\]](#)
- [US6075181A \[0151\]](#)
- [US6114598A \[0151\]](#)
- [US6150584A \[0151\]](#)
- [US6162963A \[0151\]](#)
- [US5223409A \[0152\]](#)
- [US5403484A \[0152\]](#)
- [US5571698A \[0152\]](#)
- [US5427908A \[0152\]](#)
- [US5580717A \[0152\]](#)
- [US5969108A \[0152\]](#)
- [US6172197B \[0152\]](#)
- [US5885793A \[0152\]](#)
- [US6521404B \[0152\]](#)
- [US6544731B \[0152\]](#)
- [US6555313B \[0152\]](#)
- [US6582915B \[0152\]](#)
- [US6593081B \[0152\]](#)
- [US5476996A \[0152\]](#)
- [US5698787A \[0152\]](#)
- [WO0114424A \[0153\]](#)
- [US4399216A \[0162\]](#)
- [US4634665A \[0162\]](#)
- [US5179017A \[0162\]](#)
- [WO8704462A \[0164\]](#)
- [WO8901036A \[0164\]](#)

- [EP338841A \[0164\]](#)
- [WO3101485A \[0178\]](#)
- [WO3101485A \[0179\]](#)
- [WO2003089616A \[0234\]](#)
- [US4275149A \[0235\] \[0237\]](#)
- [US4737456A \[0235\]](#)
- [US4318980A \[0237\]](#)
- [US6414218B \[0251\]](#)
- [US20030005469A \[0251\]](#)
- [US20040115198A \[0251\]](#)
- [WO2007130642A \[0282\]](#)
- [WO0109187A \[0290\]](#)

Non-patent literature cited in the description

- GROH et al.PNAS, 2003, vol. 100, 9452- [0003]
- ALLEZ et al.Gastroenterology, 2007, vol. 132, 2346-2358 [0003]
- OGASAWARA et al.Immunity, 2004, vol. 20, 757-767 [0003]
- PENDE et al.Eur J Immunol, 2001, vol. 31, 1076-86 [0004]
- BAUER et al.Science, 1999, vol. 285, 727-9 [0004] [0320]
- CASTRICONI et al.PNAS, 2003, vol. 100, 4120-25 [0004]
- ANDRÉ et al.Eur J Immunol, 2004, vol. 34, 1-11 [0004]
- KWONG et al.J Mol Biol, 2008, vol. 384, 1143-1156 [0006]
- LI et al.Nat Immunol, 2001, vol. 2, 443-451 [0017] [0018]
- WU et al.J Exp Med, 2000, vol. 192, 1059- [0018]
- HARLOWLANEANTIBODIES: A LABORATORY MANUALCold Spring Harbor Laboratory Press19880000 [0019] [0294]
- JONES et al.Nature, 1986, vol. 321, 522-525 [0024]
- RIECHMANN et al.Nature, 1988, vol. 332, 323-329 [0024]
- PRESTACurr. Op. Struct. Biol., 1992, vol. 2, 593-596 [0024]
- KABAT et al.Sequences of Proteins of Immunological InterestU.S. Department of Health and Human Services19910000 [0025]
- CHOTHIALESKJ. Mol. Biol, 1987, vol. 196, 901-917 [0025]
- LEERICHARDSJ Mol Biol, 1971, vol. 55, 379-400 [0028]
- NEEDLEMANWUNSCHJ. Mol. Biol., 1970, vol. 48, 444-453 [0090]
- PEARSONMethods Enzymol., 1990, vol. 183, 63-98 [0091]
- PEARSONMethods Mol. Biol., 2000, vol. 132, 185-219 [0091]
- E. MEYERSW. MILLERComput. Appl. Biosci., vol. 1988, 11-17 [0092]
- ALTSCHUL et al.J. Mol. Biol., 1990, vol. 215, 403-410 [0093]
- ALTSCHUL et al.Nucleic Acids Res., 1997, vol. 25, 3389-402 [0093]
- BIRD et al.Science, 1988, vol. 242, 423-426 [0099]
- HUSTON et al.PNAS, 1988, vol. 85, 5879-5883 [0099]
- III et al.Protein Eng, 1997, vol. 10, 949-57 [0099]
- HOLLIGERHUDSONNat Biotechnol, 2005, vol. 23, 1126-1136 [0099]
- MORIMOTO et al.Journal of Biochemical and Biophysical Methods, 1992, vol. 24, 107-117 [0101]
- BRENNAN et al.Science, 1985, vol. 229, 81- [0101]
- CARTER et al.Bio/Technology, 1992, vol. 10, 163-167 [0101]
- MILLSTEIN et al.Nature, 1983, vol. 305, 537-539 [0103]
- DREIERBioconj. Chem., 1998, vol. 9, 4482-489 [0106]
- STRANDMANN et al.Blood, 2006, vol. 107, 1955-1962 [0106]
- TUTT et al.J. Immunol, 1991, vol. 147, 60- [0107]
- KARPOVSKY et al.J. Exp. Med., 1984, vol. 160, 1686- [0108]
- LIU, MA et al.Proc. Natl. Acad. Sci. USA, 1985, vol. 82, 8648- [0108]
- PAULUSBehring Ins. Mitt., 1985, 78118-132 [0108]
- BRENNAN et al.Science, 1985, vol. 229, 81-83 [0108]

- **GLENNIE et al.**J. Immunol., 1987, vol. 139, 2367-2375 [0108]
- **KONTERMANN et al.**Acta Pharmacological Sinica, 2005, vol. 26, 11-9 [0110]
- **KOSTELNY et al.**J. Immunol., 1992, vol. 148, 51547-1553 [0110]
- **HOLLINGER et al.**PNAS, 1993, vol. 90, 6444-6448 [0110]
- **GRUBER**J. Immunol., 1994, vol. 152, 5368- [0110]
- **RIECHMANN, L.**Nature, 1998, vol. 332, 323-327 [0115]
- **JONES, P. et al.**Nature, 1986, vol. 321, 522-525 [0115]
- **QUEEN, C.**Proc. Natl. Acad. Sci. U.S.A., 1989, vol. 86, 10029-10033 [0115]
- **KABAT, E. A et al.**Sequences of Proteins of Immunological InterestU.S. Department of Health and Human Services19910000 [0117]
- **TOMLINSON, I. M. et al.**The Repertoire of Human Germline VH Sequences Reveals about Fifty Groups of VH Segments with Different Hypervariable LoopsJ. Mol. Biol., 1992, vol. 227, 776-798 [0117]
- **COX, J. P. L. et al.**A Directory of Human Germ-line VH Segments Reveals a Strong Bias in their UsageEur. J. Immunol., 1994, vol. 24, 827-836 [0117]
- **SHIELDS, R.L. et al.**J. Biol. Chem., 2001, vol. 276, 6591-6604 [0129]
- **ANGAL et al.**Mol Immunol., 1993, vol. 30, 105-8 [0130]
- **SHIELDS, R.L. et al.**J. Biol. Chem., 2002, vol. 277, 26733-26740 [0131]
- **UMANA et al.**Nat. Biotech., 1999, vol. 7, 176-180 [0131]
- Remington's Pharmaceutical SciencesMack Publishing Co. 19950000 [0135]
- Goodman and Gilman's The Pharmacological Basis of TherapeuticsMcGraw Hill20010000 [0135]
- **PASTAN et al.**Cell, 1986, vol. 47, 641- [0135]
- **GOLDENBERG**Cancer Journal for Clinicians, 1994, vol. 44, 43- [0135]
- **YU**Int. J. Cancer, 1994, vol. 56, 244- [0138]
- **WONG**Chemistry of Protein Conjugation and Cross-linkingCRC Press19910000 [0138]
- Modification of Antibodies by Chemical Methods**UPESLACIS et al.**Monoclonal antibodies: principles and applicationsWiley-Liss, Inc.19950000187-230 [0138]
- Production and Characterization of Synthetic Peptide-Derived Antibodies**PRICE**Monoclonal antibodies: Production, engineering and clinical applicationCambridge University Press1995000060-84 [0138]
- **CATTEL et al.**Chemistry today, 1989, vol. 7, 51-58 [0138]
- **DELPRINO et al.**J. Pharm. Sci, 1993, vol. 82, 699-704 [0138]
- **ARPICCO et al.**Bioconjugate Chemistry, 1997, vol. 8, 3- [0138]
- **REISFELD et al.**Antihody, Immunicon. Radiopharm, 1989, vol. 2, 217- [0138]
- Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy**AMON et al.**Monoclonal Antibodies And Cancer TherapyAlan R. Liss, Inc.19850000243-56 [0138]
- Antibodies For Drug Delivery**HELLSTROM et al.**Controlled Drug DeliveryMarcel Dekker, Inc.19870000623-53 [0138]
- Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review**THORPE et al.**Monoclonal Antibodies '84: Biological And Clinical Applications19850000475-506 [0138]
- Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer TherapyMonoclonal Antibodies For Cancer Detection And TherapyAcademic Press19850000303-16 [0138]
- The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates **THORPE**Immunol. Rev.19820000vol. 62, 119-58 [0138]
- **SAITO, G. et al.**Adv. Drug Deliv. Rev., 2003, vol. 55, 199-215 [0139]
- **TRAIL, P.A. et al.**Cancer Immunol. Immunother., 2003, vol. 52, 328-337 [0139]
- **PAYNE, G.**Cancer Cell, 2003, vol. 3, 207-212 [0139]
- **ALLEN, T.M.**Nat. Rev. Cancer, 2002, vol. 2, 750-763 [0139]
- **PASTAN, I.KREITMAN, R. J.**Curr. Opin. Investig. Drugs, 2002, vol. 3, 1089-1091 [0139]
- **SETER, P.D.**Adv. Drug Deliv. Rev.Springer, C.J.20010000vol. 53, 247-264 [0139]
- Current Protocols in Molecular BiologyGreene Publishing and Wiley Interscience19870000 [0142]
- **KABAT, E. A**Sequences of Proteins of Immunological InterestU.S. Department of Health and Human Services19910000 [0146]
- **KABAT, E. A et al.**Sequences of Proteins of Immunological InterestU. S. Department of Health and Human Services19910000 [0147]
- **BIRD et al.**Science19880000vol. 242, 423-426 [0148]
- **HUSTON et al.**Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 5879-5883 [0148]
- **MCCAFFERTY et al.**Nature, 1990, vol. 348, 552-554 [0148]
- **KOHLERMILSTEIN**Nature, 1975, vol. 256, 495- [0149]
- **LONBERG et al.**Nature, 1994, vol. 368, 856-859 [0151]
- **LONBERG, N.**Handbook of Experimental Pharmacology, 1994, vol. 113, 49-101 [0151]

- **LONBERG, N.HUSZAR, D.**Intern. Rev. Immunol., 1995, vol. 13, 65-93 [0151]
- **HARDING, F.LONBERG, N.**Ann. N. Y. Acad. Sci., 1995, vol. 764, 536-546 [0151]
- **TAYLOR, L. et al.**Nucleic Acids Research, 1992, vol. 20, 6287-6295 [0151]
- **CHEN, J. et al.**International Immunology, 1993, vol. 5, 647-656 [0151]
- **TUAILLON et al.**Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 3720-3724 [0151]
- **CHOI et al.**Nature Genetics, 1993, vol. 4, 117-123 [0151]
- **CHEN, J. et al.**EMBO J., 1993, vol. 12, 821-830 [0151]
- **TUAILLON et al.**J. Immunol., 1994, vol. 152, 2912-2920 [0151]
- **TAYLOR, L. et al.**International immunology, 1994, vol. 6, 579-591 [0151]
- **FISHWILD, D. et al.**Nature Biotechnology, 1996, vol. 14, 845-851 [0151]
- **TOMIZUKA et al.**Proc. Natl. Acad. Sci. USA, 2000, vol. 97, 722-727 [0151]
- **KUROIWA et al.**Nature Biotechnol, 2002, vol. 20, 889-894 [0151]
- **LONBERG, N. et al.**Nature, 1994, vol. 368, 6474856-859 [0153]
- **FISHWILD, D. et al.**NatureBiotechnolo,gy, 1996, vol. 14, 845-851 [0153]
- **MORRISON, S.**Science, 1985, vol. 229, 1202- [0157]
- **GOEDEL**Gene Expression Technology. Methods in EnzymologyAcademic Press19900000vol. 185, [0160]
- **TAKEBE, Y. et al.**Cell. Biol., 1988, vol. 8, 466-472 [0161]
- **BOSS, M. A.WOOD, C. R.**Immunology Today, 1985, vol. 6, 12-13 [0163]
- **URLAUBCHASIN**Proc. Nail. Acad. Sci., 1980, vol. 77, 4216-4220 [0164]
- **R. J. KAUFMANP. A. SHARP**Mol. Biol., 1982, vol. 159, 601-621 [0164]
- **A Laboratory Manual**Cold Spring Harbor Laboratory19880000 [0170]
- **GROH et al.**PNAS, 2003, vol. 100, 9452-9457 [0179]
- **ANDRÉ et al.**Eur. J. Immunol, 2004, vol. 34, 1-11 [0179]
- **GROH**PNAS, 2003, vol. 100, 9452-9457 [0180]
- **BRENNAN et al.**Lancet, 1989, vol. 2, 8657244-247 [0190]
- **KJELLEV et al.**Eur J Immunol, 2008, vol. 37, 1397-1406 [0191] [0255]
- **ITO et al.**Am J Physiol Gastrointest Liver Physiol, 2008, vol. 294, G199-G207 [0191]
- **REMINGTON**The Science and Practice of Pharmacy19950000 [0199] [0200] [0201] [0202]
- **WILLIAMSPOLLIJ.**Parenteral Sci. Technol., 1984, vol. 38, 48-59 [0202]
- **MASTERS**Spray-Drying HandbookLongman Scientific and Technical19910000491-676 [0202]
- **BROADHEAD et al.**Drug Devel. Ind. Pharm., 1992, vol. 18, 1169-1206 [0202]
- **MUMENTHALER et al.**Pharm. Res., 1994, vol. 11, 12-20 [0202]
- **CARPENTERCROWE**Cryobiology, 1988, vol. 25, 459-470 [0202]
- **ROSER**Biopharm., 1991, vol. 4, 47-53 [0202]
- **The Science and Practice of Pharmacy**19950000 [0208]
- **WISE, D.L.**Handbook of Pharmaceutical Controlled ReleaseMarcel Dekker20000000 [0217]
- **Drug and the Pharmaceutical Sciences**Protein Formulation and DeliveryMarcel Dekker20000000vol. 99, [0217]
- **YU JCHIEN YW**Pulmonary drug delivery: Physiologic and mechanistic aspectsCrit Rev Ther Drug Carr Sys, 1997, vol. 14, 4395-453 [0219]
- **EDWARDS DABEN-JEBRIA ALANGER R**Recent advances in pulmonary drug delivery using large, porous inhaled particlesJ Appl Physiol, 1998, vol. 84, 2379-385 [0221]
- **J Appl Physiol**, 1998, vol. 84, 2379-385 [0221] [0221] [0222]
- **AHEM. T.J.MANNING M.C.**Stability of Protein PharmaceuticalsPlenum Press19920000 [0227]
- **Current Protocols in Immunology**Wiley-Interscience19910000vol. 1 and 2, [0235]
- **Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme Immunoassay**O'SULLIVAN et al.Methods in Enzym.Academic Press19810000vol. 73, 147-166 [0235]
- **ZOLA**Monoclonal Antibodies: A Manual of TechniquesCRC Press, Inc.19870000147-158 [0240]
- **WOOLEY, P. H.**Animal Models of ArthritisMosby Publishers19980000 [0251]
- **ERNING et al.**Arthritis Res, 2002, vol. 4, 3133-40 [0251]
- **HOLMDAHL et al.**Ageing Res Rev, 2002, vol. 1, 1135-47 [0251]
- **ANTHONY et al.**Clin Exp Rheumatol, 1999, vol. 17, 2240-4 [0251]
- **DURIE et al.**Clin Immunol Immunopathol, 1994, vol. 73, 111-8 [0251]
- **MULLER-LADNER et al.**Drugs Today, 1999, vol. 35, 4-5379-88 [0251]
- **ROBERTS et al.**J Immunol, 2001, vol. 167, 5527-30 [0254]
- **HÜE et al.**Immunity, 2004, vol. 21, 367-77 [0254]
- **MERESSE et al.**Immunity, 2004, vol. 21, 357-66 [0254]
- **ROBERTS et al.**J Immunology, 2001, vol. 167, 5527-30 [0254]

- **ALLEZ et al.**European Congress of Immunology, 2006, [0254]
- **CHEN**Hepatology, 2007, vol. 46, 3706-715 [0282]
- **WU et al.**Science, 1999, vol. 385, 730-2 [0289]
- **CHEN et al.**EMBO J., 1993, vol. 12, 811-820 [0290]
- **FISHWILD et al.**Nature Biotechnology, 1996, vol. 14, 845-851 [0290]
- **DUROCHER et al.**Nucleic Acids Res, 2002, vol. 30, 2e9- [0307]
- **PHAM et al.**Biotechnol Bioeng, 2003, vol. 84, 3332-42 [0307]
- **OHM-LAURSEN et al.**Immunology, 2006, vol. 119, 265-77 [0315]
- **ROBERTSON et al.**Exp Hematol, 1996, vol. 24, 406-15 [0329]
- **MASHOO et al.**Immunol., 2005, vol. 174, 4480-4484 [0351]
- **LI, P.MORRIS, D. L.WILLCOX, B. E.STEINLE, A.SPIES, T.STRONG, R. K.**Nature Immunology, 2001, vol. 2, 443-451 [0368]
- **URSBY, T. MAMMEN, C. B.CERENIUS, Y.SVENSSON, C.SOMMARIN, B.FODJE, M. N.KVICK, A.LOGAN, D. T.LS-NIELSEN, J.THUNNISSEN, M. M. G. M.**The MAD StationThe New Macromolecular Crystallography Stations At MAX-lab, 2004, [0368]
- **KABSCH, W.J.**Appl. Crystallogr., 1993, vol. 26, 795-800 [0368]
- **VAGIN, A.TEPLYAKOV, A.J.**Appl. Crystallogr., 1997, vol. 30, 1022-1025 [0368]
- **MCCOY, A. J. GROSSE-KUNSTLEVE, R. W.STORONI, L. C.READ, R. J.**Acta Crystallographica Section D Biological Crystallography, 2005, vol. 61, 458-464 [0368]
- **MCCOY, A. J. GROSSE-KUNSTLEVE, R. W.ADAMS, P. D.WINN, M. D.STORONI, L. C.READ, R. J.**J. Appl. Crystallogr., 2007, vol. 40, 658-674 [0368]
- **BAILEY, S.**Acta Crystallogr. Sect. D-Biol. Crystellogr., 1994, vol. 50, 760-763 [0368]
- **BERMAN, H. M.WESTBROOK, J.FENG, Z.GILLILAND, G.BHAT, T. N. WEISSIG, H.SHINDYALOV, I. N.BOURNE, P. E.**Nucleic Acids Res., 2000, vol. 28, 235-242 [0368]
- **VAJDOS, F. F.ADAMS, C. W.BREECE, T. N.PRESTA, L. G.VOS, A. M.SIDHU, S. S.**Journal of Molecular Biology, 2002, vol. 320, 415-428 [0368]
- **MCFARLAND, B. J.KORTEMME, T.YU, S. F.BAKER, D.STRONG, R. K.**Structure (Cambridge), 2003, vol. 11, 411-422 [0368]
- **MURSHUDOV, G. N.VAGIN, A. ADODSON, E. J.**Acta Crystallogr. Sect. D-Biol. Crystallogr., 1997, vol. 53, 240-255 [0368]
- **EMSLEY, P.COWTAN, K.**Acta Crystallogr. Sect. D-Biol. Crystallogr., 2004, vol. 60, 2126-2132 [0368]
- **EVANS, P.**Acta Crystallogr D Biol Crystallogr. 2006. Jan, 1962, 72-82 [0368]
- **SHELDRIK, G.**Act Cryst a, 2008, vol. 64, 112-122 [0368]
- **OHTO, U.MIZUTANI, R.NAKAMURA, M.ADACHI, H.SATOW, Y.**J Synchrotron Radiat, 2004, vol. 11, 105-108 [0368]

PATENTKRAV

1. Isoleret humant eller humaniseret monoklonalt antistof eller antigenbindende fragment, der konkurrerer med et reference-antistof om binding til hNKG2D, hvor reference-antistoffet omfatter et tungkæde-variabelt område omfattende sekvensen ifølge SEQ ID NO:44 og et letkæde-variabelt område omfattende sekvensen ifølge SEQ ID NO:45, og hvor antistoffet eller det antigenbindende fragment yderligere omfatter:
- (a) et tungkæde-variabelt område CDR1 omfattende sekvensen ifølge SEQ ID NO:48;
 - (b) et tungkæde-variabelt område CDR2 omfattende sekvensen ifølge SEQ ID NO:49;
 - 10 og
 - (c) et tungkæde-variabelt område CDR3 omfattende sekvensen ifølge SEQ ID NO:50, og
 - (d) et letkæde-variabelt område CDR1 omfattende sekvensen ifølge SEQ ID NO:51;
 - (e) et letkæde-variabelt området CDR2 omfattende sekvensen ifølge SEQ ID NO:52,
 - 15 og
 - (f) et letkæde-variabelt område CDR3 omfattende sekvensen ifølge SEQ ID NO:53.
2. Isoleret humant eller humaniseret monoklonalt antistof eller antigenbindende fragment, der binder hNKG2D og omfatter et tungkæde-variabelt område omfattende sekvensen ifølge SEQ ID NO:44 og et letkæde-variabelt område omfattende sekvensen ifølge SEQ ID NO:45.
- 20
3. Antistof eller antigenbindende fragment ifølge krav 1 eller 2 til anvendelse til behandling af en inflammatorisk eller autoimmun forstyrrelse i et menneske.
- 25
4. Antistof eller antigenbindende fragment ifølge krav 1 eller 2 til anvendelse til behandling af en inflammatorisk eller autoimmun forstyrrelse i et menneske, hvor den inflammatoriske eller autoimmune forstyrrelse er rheumatoid arthritis, multipel sklerose, systemisk lupus erythomatosus (SLE), psoriasis, cøliaki, ulcerativ colitis, Crohns sygdom, transplantatafstødning eller graft-versus-vært-sygdom.
- 30
5. Antistof eller antigenbindende fragment ifølge krav 1 eller 2 til anvendelse til behandling af en inflammatorisk eller autoimmun forstyrrelse i et menneske, hvor den inflammatoriske eller autoimmune forstyrrelse er Crohns sygdom.

DRAWINGS




Fig. 1




Fig. 2




Fig. 3

16F16 heavy chain (IgG4):

EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSSITSSSSYIYYAD
SVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRRYFDWFFLDYRGQGTLVTVSSAST
 KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
 SSVVTVPSSSLGKTYTCNVDHKPSNTKVDKRVESKYGPCCPAPEFLGGPSVFLFPPKP
 KDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVL
 HQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKG
 FYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALH
 NHYTQKSLSLGK (SEQ ID NO:7)

16F16 light chain:

DIQMTQSPSSLSASVGRVITTCRASQGISSWLAWYQQKPEKAPKSLIYAASSLQSGVPSRF
SGSGSGTDFTLTITSSLPEDFATYYCQQYNGYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQ
 LKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADY
 EKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:8)

16F31 heavy chain (IgG4):

EVQLVESGGGVVRPVGSLRLSCAASGFTFDDYGMTWVRQAPGKGLEWVSGINWNGSGTGYAD
SVKGRFTISRDNAKNSLYLQMNSLRAEDTALYYCARERELYYYYYGLDVWGQGTTVTVSSAS
 TKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS
 LSSVTVPSSSLGKTYTCNVDHKPSNTKVDKRVESKYGPCCPAPEFLGGPSVFLFPPK
 PKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTV
 LHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVK
 GFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL
 HNHYTQKSLSLGK (SEQ ID NO:9)

16F31 light chain:

EIVLTQSPGTLSTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDR
FSGSGSGTDFTLTISRLEPEDFAVYYCQQYGSSPFTFGPGTKVDIKRTVAAPSVFIFPPSDE
 QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKAD
 YEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:10)

Fig. 4A

MS Heavy Chain (IgG4):

QVHLQESGPGLVKPSSETLSLTCTVSDDSISSYYWSWIRQPPGKGLEWIGHISYSGSANYNPS
LKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCANWDDAFNIWGQGTMTVTVSSASTKGPSVF
 PLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV
 PSSSLGKTKYTCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMI
 SRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLN
 GKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDI
 AVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQK
 SLSLSLGK (SEQ ID NO:40)

MS Light Chain:

EIVLTQSPGTLSTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDR
PSGSGSGTDFTLTISRLEPEDFAVYYCQQYGSSPWTFGQGTKVEIKRTVAAPSVFIFPPSDE
 QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSTLTLSKAD
 YEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:41)

21F2 Heavy chain (IgG4):

EVQLVQSGAEVKEPGESESLKISCKNSGYSFTNYWVGWVRQMPGKGLEWGMGIYIPGDS DTRYSP
SFQQQVTISADKSINTAYLQWSSSLKASDTAMYYCGRLTMFRGIIIGYFDYWGQGTTLVTVSSA
 STKGPSVFPPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLY
 SLSSVTVTPSSSLGKTKYTCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPP
 KPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLT
 VLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLV
 KGFYPSDI AVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL
 LHNHYTQKSLSLSLGK (SEQ ID NO:42)

21F2 Light chain:

EIVLTQSPATLSTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYDASNRATGIPARF
SGSGSGTDFTLTISISLEPEDFAVYYCQQRSNWFWTFGQGTKVEIKRTVAAPSVFIFPPSDEQ
 LKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSTLTLSKADY
 EKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:43)

Fig. 4B

Fig. 5A: 16F16 Heavy Chain

```

      1       2       3       4       5       6
123456789012345678901234567890123456789012345678901234567890
EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYSMN  WVRQAPGKGLWVSSITS  SSSYIYYA
EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYSMN  WVRQAPGKGLWVSSISS  SSSYIYYA
      7       8       9      10      11
12345678901234567890123456789012345678901234567890 <- Kabat
DSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDRRYFDWFFL  DYEGQGTLLTVSS  16F16  VH
DSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR/  /  DYEGQGTLLTVSS  VH3_21/D3-9/JH4

```

Fig. 5B: 16F16 Light Chain

```

      1       2       3       4       5       6
123456789012345678901234567890123456789012345678901234567890
DIQMTQSPSSLSASVGDRTITCRASQ  GISSWLAWYQQKPEKAPKSLIYAASSLQSGVPS
DIQMTQSPSSLSASVGDRTITCRASQ  GISSWLAWYQQKPEKAPKSLIYAASSLQSGVPS
      7       8       9      10
1234567890123456789012345678901234567890123456789 <- The Kabat
RFSGSGSGTDFTLTISRLEPEDFATYYCQYNGYP  YTFGQGTKLEIK  16F16  VL
RFSGSGSGTDFTLTISRLEPEDFATYYCQYNGYP//YTFGQGTKLEIK  VKI_L15/JK2

```

Fig. 5C: 16F31 Heavy Chain

```

      1       2       3       4       5       6
123456789012345678901234567890123456789012345678901234567890
EVQLVESGGGVVRPGGSLRLSCAASGFTFDDYGM  WVRQAPGKGLWVSGINW  NGGSTGYA
EVQLVESGGGVVRPGGSLRLSCAASGFTFDDYGM  WVRQAPGKGLWVSGINW  NGGSTGYA
      7       8       9      10      11
12345678901234567890123456789012345678901234567890 <- Kabat
DSVKGRFTISRDNAKNSLYLQMNSLRAEDTALYYCARERELYYYYYGL  DVMGQGTTVTVSS  16F31  VH
DSVKGRFTISRDNAKNSLYLQMNSLRAEDTALYYCAR/  /YYYYYGM  DVMGQGTTVTVSS  VH3_20/D3-
10/JH6

```

Fig 5D: 16F31 Light Chain

```

      1       2       3       4       5       6
123456789012345678901234567890123456789012345678901234567890
EIVLTQSPGTLSPGEPATLSCPASQS  VSSSYLAWYQQKPGQAPRLIYGASSRATGIPD
EIVLTQSPGTLSPGEPATLSCPASQS  VSSSYLAWYQQKPGQAPRLIYGASSRATGIPD
      7       8       9      10
1234567890123456789012345678901234567890123456789 <- Kabat
RFSGSGSGTDFTLTISRLEPEDFAVYYCQYGGSP  FTFGPGTKVDIK  16F31  VL
RFSGSGSGTDFTLTISRLEPEDFAVYYCQYGGSP//FTFGPGTKVDIK  VKIII_A27/JK3

```

Fig. 5E: MS Heavy Chain

MS Heavy Chain

1	2	3	4	5	6	
123456789012345678901234567890123456789012345678901234567890	Kabat					
QVHLQESGPGLVKPSSTLSLTCTVSDDSISSYYWWSWIRQPPGKGLEWIGHISYSGSANYN	MS VH					
QVQLQESGPGLVKPSSTLSLTCTVSGGSISSYYWWSWIRQPPGKGLEWIGYIYYSGSTNYN	VH4_59					
7	8	9	10	11		
12345678901234567890123456789012345678901234567890123456789012345678901234567890123456789012345678901234567890	Kabat					
PSLKSRTVISVDTSKNQFSLKLSSTVAADTAVYYCA_ NWQ DAFNINWGQGTMTVTSS	MS VH					
PSLKSRTVISVDTSKNQFSLKLSSTVAADTAVYYCAR/NWG/DAFDINWGQGTMTVTSS	VH4_59/D7_27_3/JH3					

Fig. 5F: MS Light Chain

1	2	3	4	5	6	
123456789012345678901234567890123456789012345678901234567890	Kabat					
EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPD	MS Light Chain					
EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPD	VKIII_A27					
7	8	9	10			
12345678901234567890123456789012345678901234567890123456789012345678901234567890123456789012345678901234567890	Kabat					
RFSGSGSGTDFTLTISRLEPEDFAVYYCQYGSSEF WTFGQGTKVEIK	MS VL					
RFSGSGSGTDFTLTISRLEPEDFAVYYCQYGSSEF/WTFGQGTKVEIK	VKIII_A27/JK1					

Fig. 5G: 21F2 Heavy Chain

1	2	3	4	5	6	
123456789012345678901234567890123456789012345678901234567890	Kabat					
EVQLVQSGAEVKEPGESLKISCKNSGYSTFTNYWGWVRQMPGKGLEWIMGIIYPGDSSTRYS	21F2 Heavy Chain					
EVQLVQSGAEVKEPGESLKISCKNSGYSTFTSIWGWVRQMPGKGLEWIMGIIYPGDSSTRYS	VH5_51					
7	8	9	10	11		
12345678901234567890123456789012345678901234567890123456789012345678901234567890123456789012345678901234567890	Kabat					
PSFQGVVISADKSIINTAYLQWSSSLKASDTAMYYCQR LTMFRGIIIIQYFDYWGQGLTVTVSS	21F2 Heavy Chain					
PSFQGVVISADKSIINTAYLQWSSSLKASDTAMYYCAR/ITMVRGVII/YFDYWGQGLTVTVSS	VH5_51/D3_10_R3/JH4					

Fig. 5H: 21F2 Light Chain

1	2	3	4	5	6	
123456789012345678901234567890123456789012345678901234567890	Kabat					
EIVLTQSPATLSLSPGSRATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYDASNRATGIPA	21F2 Light Chain					
EIVLTQSPATLSLSPGSRATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYDASNRATGIPA	VKIII_L6					
7	8	9	10			
12345678901234567890123456789012345678901234567890123456789012345678901234567890123456789012345678901234567890	Kabat					
RFSGSGSGTDFTLTISRLEPEDFAVYYCQQRSNWF WTFGQGTKVEIK	21F2 Light Chain					
RFSGSGSGTDFTLTISRLEPEDFAVYYCQQRSNWF/WTFGQGTKVEIK	VKIII_L6/JK1					




Fig. 6




Fig. 7






Fig. 8

A



B



C




Fig. 9




Fig. 10






Fig. 11

**Fig. 12**

A



B





Fig. 13

A



B






Fig. 14

**Fig. 15**

A



B







Fig. 16

**Fig. 17**

A



B



C





Fig. 18

A



B




Fig. 19




Fig. 20A




Fig. 20B




Fig. 20C

NKG2D monomer unit 1

```

1    MGWIRGRRSR HSWEMSEFHN YNLDLKKSDF STRWQKQRCF VVKSKCRENA
51   SPFFFCFIA VAMGIRFIIM VTIWSAVFLN SLFNQEVQIP LTESYCGPCF
101  KNWICYKNNC YQFFDESKNW YESQASCMSQ NASLLKVYSK EDQDLLKLVK
151  SYHWMGLVHI PTNGSWQWED GSILSPNLLT IIEMQKGDCA LYASSFKGYI
201  ENCSTPNTYI CMQRTV

```

NKG2D monomer unit 2

```

1    MGWIRGRRSR HSWEMSEFHN YNLDLKKSDF STRWQKQRCF VVKSKCRENA
51   SPFFFCFIA VAMGIRFIIM VTIWSAVFLN SLFNQEVQIP LTESYCGPCF
101  KNWICYKNNC YQFFDESKNW YESQASCMSQ NASLLKVYSK EDQDLLKLVK
151  SYHWMGLVHI PTNGSWQWED GSILSPNLLT IIEMQKGDCA LYASSFKGYI
201  ENCSTPNTYI CMQRTV

```

Fig. 21A

NKG2D monomer unit 1

```

1      MGWIRGRRSR HSWEMSEFHN YNLDLKKSD F STRWQKQRC P VVKS KCRENA
51     SPFFFCFFIA VAMGIRFIIM VTIWSAVFLN SLFNQEVQIP LTESYCGPCP
101    KNWICYKNNC YQFFDESKNW YESQASCMSQ NASLLKVYSK EDQDLLKLVK
151    SYHWMGLVHI PTNGSWQWED GSILSPNLIT IEMQKGDCA LYASSFKGYI
201    ENCSTPNTYI CMQRTV

```

NKG2D monomer unit 2

```

1      MGWIRGRRSR HSWEMSEFHN YNLDLKKSD F STRWQKQRC P VVKS KCRENA
51     SPFFFCFFIA VAMGIRFIIM VTIWSAVFLN SLFNQEVQIP LTESYCGPCP
101    KNWICYKNNC YQFFDESKNW YESQASCMSQ NASLLKVYSK EDQDLLKLVK
151    SYHWMGLVHI PTNGSWQWED GSILSPNLIT IEMQKGDCA LYASSFKGYI
201    ENCSTPNTYI CMQRTV

```

Fig. 21B

NKG2D monomer unit 1

```

1      MGWIRRRSR HSWEMSEFHN YNLDLKKSD F STRWQKQRC P VVKS KCRENA
51     SPFFFCFIA VAMGIRFIIM VTIWSAVFLN SLFNQEVQIP LTESYCGPCP
101    KNWICYKNNC YQFFDESKNW YESQASCMSQ NASLLKVYSK EDQDLLKLVK
151    SYHWMGLVHI PTNGSWQWED GSILSPNLLT I IEMQKGDCA LYASSFKGYI
201    ENCSTPNTYI CMQRTV

```

NKG2D monomer unit 2

```

1      MGWIRRRSR HSWEMSEFHN YNLDLKKSD F STRWQKQRC P VVKS KCRENA
51     SPFFFCFIA VAMGIRFIIM VTIWSAVFLN SLFNQEVQIP LTESYCGPCP
101    KNWICYKNNC YQFFDESKNW YESQASCMSQ NASLLKVYSK EDQDLLKLVK
151    SYHWMGLVHI PTNGSWQWED GSILSPNLLT I IEMQKGDCA LYASSFKGYI
201    ENCSTPNTYI CMQRTV

```

Fig. 21C




Fig. 22