



US00PP31976P2

(12) **United States Plant Patent**
Smith

(10) **Patent No.:** **US PP31,976 P2**

(45) **Date of Patent:** **Jul. 14, 2020**

(54) **CHRYSANTHEMUM PLANT NAMED**
'CIFZ0069'

(50) Latin Name: *Chrysanthemum x morifolium*
Varietal Denomination: **CIFZ0069**

(71) Applicant: **SYNGENTA CROP PROTECTION**
AG, Basel (CH)

(72) Inventor: **Mark A. Smith**, Alva, FL (US)

(73) Assignee: **Syngenta Crop Protection AG**, Basel
(CH)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/350,950**

(22) Filed: **Feb. 4, 2019**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/14 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./296**
CPC *A01H 6/1424* (2018.05)

(58) **Field of Classification Search**
USPC Plt./296
CPC *A01H 6/1424; A01H 5/02*
See application file for complete search history.

Primary Examiner — Keith O. Robinson

(74) *Attorney, Agent, or Firm* — Dale Skalla

(57) **ABSTRACT**

A new garden type *Chrysanthemum* plant named
'CIFZ0069' particularly distinguished by the medium size
plant with a round mound plant habit, a small sized (1.5
inch) flower, red-bronze decorative flower color and a natu-
ral season response in late September to mid-October

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Chrysanthemum x morifolium.
Varietal denomination: 'CIFZ0069'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new *Chrysanthemum*,
botanically known as *Chrysanthemum x morifolium*, and
hereinafter referred to by the variety name 'CIFZ0069'.

'CIFZ0069' is a product of a planned breeding program.
The new cultivar is a medium size plant with round mound
plant habit, a small sized (1.5 inch) flower, red-bronze
decorative flower color and a natural season response in late
September to mid-October.

'CIFZ0069' originated from 'CIFZ0031', a stock x-rayed
at 1750 RAD in December of 2015 in Gilroy, Calif.
'CIFZ0069' was selected from the resulting population on
Mar. 8, 2016 in Gilroy, Calif.

The female parent was the variety 'CIFZ0031', U.S. Pat.
No. 23,924.

When compared to the female parent, 'CIFZ0069' is/has:

a similar flower type with a bit smaller flower size but darker bronze/
almost red flower color versus orange of 'CIFZ0031'.
a couple days slower in blackcloth response and natural season response
than 'CIFZ0031'.
has a similar plant habit and similar or a bit smaller overall plant size than
'CIFZ0031'.

As 'CIFZ0069' is a mutant plant, there is no male parent.

The first act of asexual reproduction of 'CIFZ0069' was
accomplished when vegetative stem cuttings were propa-
gated from the initial selection in April 2016 in Gilroy, Calif.

BRIEF SUMMARY OF INVENTION

Horticultural examination of plants grown from cuttings
of the plant initiated in April 2016 and continuing thereafter,

2

has demonstrated that the combination of characteristics as
herein disclosed for 'CIFZ0069' are firmly fixed and are
retained through successive generations of asexual repro-
duction.

'CIFZ0069' has not been observed under all possible
environmental conditions. The phenotype may vary signifi-
cantly with variations in environment such as temperature,
light intensity and day length.

A Plant Breeder's Right for this cultivar has not yet been
applied for. 'CIFZ0069' has not been made publicly avail-
able prior to the effective filing date of this application,
notwithstanding any disclosure that may have been made
less than one year prior to the effective filing date of this
application by the inventor or another who obtained
'CIFZ0069' directly from the inventor.

The following traits have been repeatedly observed and
are determined to be basic characteristics of the new variety.
The combination of these characteristics distinguishes this
Chrysanthemum as a new and distinct variety.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographic drawings show typical
flower and foliage characteristics of 'CIFZ0069' with colors
being as true as possible with an illustration of this type.

The photographic drawings show in FIG. 1 a close view
of flower of the new variety and in FIG. 2 a flowering plant
from an outdoor trial.

The aforementioned photographs: FIG. 1, as well as FIG.
2, were taken in August 2018 both showing a plant from the
same blackcloth indoor trial in Gilroy, Calif.

These plants were about 10 weeks of age. One rooted
cutting per pot had been planted in a one gallon pot, not
pinched in week 24, and black clothed from week 25. Plants
started flowering in August.

DETAILED BOTANICAL DESCRIPTION

The measurements were taken in Gilroy, Calif. in August
2018 on the plants from the aforementioned trial.

Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.) 2015.

TABLE 1

DIFFERENCES BETWEEN THE NEW VARIETY 'CIFZ0069' AND TWO MOST SIMILAR VARIETIES:		
	'CIFZ0069'	'CIFZ0048', U.S. Plant Pat. No. 29,639
Flowering response (blackcloth):	Similar	Similar
Flowering response (natural):	Similar	Similar
Flower color:	Similar	Similar
Flower size:	Larger	Smaller
Flowering:	Not flowering uniform	Flowering very uniform
Plant habit:	Wider	More upright
	'CIFZ0069'	'Pavia Red', PP23,134
Flowering response (blackcloth):	2 weeks faster	2 weeks later
Flowering response (natural):	Similar	Similar
Flower size:	Similar	Similar
Plant habit:	Less round/more mound	Very round
Flower longevity	Long lasting	Shorter lasting
Plant size:	Smaller	Bigger

Plant:

Form, growth and habit.—Herbaceous garden-type, stems upright and outwardly spreading, freely branching, strong and moderately vigorous growth habit.

Plant height (above soil).—11-16 cm.

Plant height (inflorescence included).—17-29 cm.

Plant width.—33-35 cm.

Roots:

Number of days to initiate roots.—About 4 days at about 22° C.

Number of days to produce a rooted cutting.—14-16 days at 22° C.

Type.—Fine, fibrous, free branching.

Color.—RHS N155B.

Foliage:

Arrangement.—Alternate.

Immature leaf, color upper surface.—RHS 139A.

Immature leaf, color lower surface.—RHS 137C.

Mature leaf, color, upper surface.—RHS 137A.

Mature leaf, color lower surface.—RHS NN137C.

Length.—5.6 cm.

Width.—4.7-5.2 cm.

Shape.—Ovate, with distinct lobes.

Base shape.—Attenuate.

Apex shape.—Mucronate.

Margin.—5-lobed (palmate), edge dentate.

Texture, upper surface.—Bifid hairs.

Texture, lower surface.—Bifid hairs.

Color of veins, upper surface.—RHS 138B.

Color of veins, lower surface.—RHS N137A.

Pattern of veining.—Palmate.

Petiole color.—RHS 138A.

Petiole Length.—1.8-2.2 cm.

Diameter.—0.2-0.3 cm.

Texture.—Covered with bifid hairs.

Stem:

Quantity of main branches per plant.—5-8.

Color of stem.—RHS 138A.

Length of stem.—7-9 cm.

Diameter.—0.6 cm.

Length of internodes.—1.0-1.6 cm.

Texture.—Bifid hairs.

Color of peduncle.—RHS 138B.

Length of peduncle.—6.7-7.8 cm.

Peduncle diameter.—0.3 cm.

Texture.—Bifid hairs.

Inflorescence:

Type.—Compositae, solitary, decorative type inflorescences borne terminally above foliage, ligulate ray florets florets arranged on a capitulum.

Quantity of short days to flowering (response time).—Approximately 5.5 weeks.

Natural season flowering.—Late September to mid-October.

Quantity of inflorescences per plant.—130-140.

Lastingness of individual blooms on the plant.—More than six weeks from first color.

Fragrance.—Slightly spicy.

25 Bud (when showing color):

Color.—RHS 46A.

Length.—0.6-0.8 cm.

Width.—1.4 cm.

Shape.—Oblate.

30 Immature inflorescence:

Diameter.—3.4 cm.

Color of ray florets, upper surface.—RHS 46A.

Color of ray florets, lower surface.—RHS 46A overlain with RHS 7A.

35 Mature inflorescence:

Diameter.—4.3 cm.

Depth.—1.6 cm.

Total diameter of disc.—No disc.

Receptacle color.—RHS 141B.

Receptacle height.—0.5 cm.

Receptacle diameter.—0.3 cm.

Ray florets:

Average quantity of florets.—Between 230 and 240 in many whorls.

Color of florets, upper surface (abaxial).—RHS 46A.

Color inner surface (adaxial).—RHS 46A overlain with RHS 7A.

Length.—1.9-2.1 cm.

Width/diameter.—0.5 cm.

Shape.—Elliptical.

Apex shape.—Most obtuse and dentate.

Base shape.—Tube.

Margin.—Entire, small incisions may occur at the tip.

Texture, upper surface.—Papillate.

Lower surface.—Papillate.

55 Disc florets: None present.

Phyllaries:

Quantity.—20-30.

Color, upper surface.—RHS 137B.

Color, lower surface.—RHS 137C.

Length.—0.6 cm.

Width.—0.3 cm.

Shape.—Lanceolate.

Apex shape.—Acute.

Base.—Fused.

Margins.—Entire.

65

Texture, upper surface.—Glabrous.
Texture, lower surface.—Canescent.
 Reproductive organs:
Pistil.—One.
Length.—0.4-0.5 cm.
Style color.—RHS 150A.
Style length.—0.2-0.3 cm.
Stigma color.—RHS 7D.
Stigma shape.—Bi-parted.
Ovary color.—RHS 145C.
Ovary length.—0.15-0.2 cm.
Ovary width.—0.1 cm.

Androecium:

Stamens.—None (are normally present in disk florets, these are not developed).
Fertility/seed set.—Has not been determined to date.
 5 *Disease/pest resistance.*—Has not been determined to date.

What is claimed is:

1. A new and distinct variety of *Chrysanthemum* plant named 'CIFZ0069' substantially as illustrated and described herein.

* * * * *



FIG. 1




FIG. 2