



(86) Date de dépôt PCT/PCT Filing Date: 2015/12/22
(87) Date publication PCT/PCT Publication Date: 2016/06/30
(85) Entrée phase nationale/National Entry: 2017/06/23
(86) N° demande PCT/PCT Application No.: US 2015/067399
(87) N° publication PCT/PCT Publication No.: 2016/106326
(30) Priorité/Priority: 2014/12/23 (US62/095,850)

(51) Cl.Int./Int.Cl. *C07D 493/18* (2006.01),
A61K 31/352 (2006.01), *A61P 35/00* (2006.01),
A61P 35/02 (2006.01)
(71) Demandeur/Applicant:
SLOAN-KETTERING INSTITUTE FOR CANCER
RESEARCH, US
(72) Inventeur/Inventor:
KUNNARI, TERO, DE
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : POLYMORPHE DE LA GRANATICINE B
(54) Title: POLYMORPH OF GRANATICIN B





Figure 10



(57) **Abrégé/Abstract:**

The present invention provides a crystalline Form A of Compound 1, also referred to as Granaticin B, and pharmaceutically compositions thereof. The present invention also provides methods of treating a microbial infection, or a disease, disorder, or condition associated with abnormal cellular proliferation, using crystalline Form A of Compound 1 or pharmaceutical compositions thereof.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2016/106326 A1(43) International Publication Date
30 June 2016 (30.06.2016)

(51) International Patent Classification:

A61P 35/02 (2006.01) *A61K 31/122* (2006.01)

(21) International Application Number:

PCT/US2015/067399

(22) International Filing Date:

22 December 2015 (22.12.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/095,850 23 December 2014 (23.12.2014) US

(71) Applicant: **SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH** [US/US]; 1275 York Avenue, New York, NY 10065 (US).(72) Inventor: **KUNNARI, Tero**; Dinglerstrasse 16, 63739 Aschaffenburg (DE).(74) Agent: **BAKER, C., Hunter**; Wolf, Greenfield & Sacks, P.c., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: POLYMORPH OF GRANATICIN B




Figure 10

(57) Abstract: The present invention provides a crystalline Form A of Compound 1, also referred to as Granaticin B, and pharmaceutically compositions thereof. The present invention also provides methods of treating a microbial infection, or a disease, disorder, or condition associated with abnormal cellular proliferation, using crystalline Form A of Compound 1 or pharmaceutical compositions thereof.

POLYMORPH OF GRANATICIN B

RELATED APPLICATIONS

[0001] The present application claims priority under 35 U.S.C. § 119(e) to U.S. provisional patent application, U.S.S.N. 62/095,850, filed December 23, 2014, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] Granaticins are polyketide-derived antibiotics produced as secondary metabolites by *Streptomyces violaceoruber* (James *et al.*, *J Gen Microbiol.* 1989, 135(7), 1997-2003). Granaticins are benzoisochromanequinones (BIQs; Keller-Schierlein *et al. Helv. Chim. Acta*, 1968, 51, 1257-1268). Chemical modifications of granaticins have been carried out at C-10 via a C-C bond formation either by glycosylation or by dimerization (Hopwood, *Chem Rev.* 1997, 97, 2465-2497; Floss *et al.*, *J Nat Prod.* 1986, 49, 951; Toral-Barza *et al, Mol. Cancer Ther.* 2007, 6, 3028-3038; Salaski *et al.*, *J. Med. Chem.* 2009, 23, 2181-2184). Additional chemical or enzymatic modification can be made to 8,11-Ethanofuro[2,3-e]naphtho[2,3-c:6,7-c']dipyrans-2,6,13(9H)-trione, 3,3a,5,8,11,13b-hexahydro-7,8,12,15-tetrahydroxy-5,9-dimethyl-, (3aS,5S,8S,9R,11R,13bS,15R) of the 3- methyl-2-oxabicyclo[2.2.2]oct-5-ene-4,8-diol group of Granaticin A (*e.g.*, coupling a carbohydrate moiety) provides granaticin B (Compound **1** in Figure 10, also called MSK-777).

[0003] In addition to the anti-bacterial activity of granaticins, Frattini *et al.* has found that granaticins inhibit protein kinase pathways (Change *et al.*, *Antibiot.* 1975, 28, 156; PCT Application Publication WO 2011/112635). In particular, granaticins were found to inhibit Cdc kinase activity based upon hits identified from high-throughput screening (HTS) of over 300,000 compounds for their ability to inhibit a heterodimer of a kinase (Cdc7) and an activator (Dbf4) that phosphorylates serine and threonine residues (WO 2011/112635).

[0004] Granaticin B was prepared from isolating the culture filtrate of *Streptomyces lateritius* or *Streptomyces violaceoruber* (Elson *et al.*, *J. Antibiotic*, 1988, 41(4), 570-572; U.S. Patent No. 3836642; Barcza *et al.*, *Helv. Chim. Acta*, 1966, 49(6), 1736-1740). The current process for preparing granaticin B involves down-stream or chemical unit operation to purify the natural product from fermentation (Keller-Schierlein, *Helv. Chim. Acta.*, 1968, 51, 1257-1268; Gilpin, *J. Antibiot.*, 1988, 41(4), 570-572). This process involves multiple steps including extractions and chromatography. The yield is not desirable, and degradation of the desired product has been observed. Therefore, there is a need to develop

feasible a large-scale process for the production of granaticin B for clinical and research purposes.

Summary of the Invention

[0005] A polymorph of granaticin B (Compound **1**) has been discovered and named Form A. The present invention provides not only Form A but compositions thereof, which are useful in treating and studying diseases. In certain embodiments, Form A, and pharmaceutical compositions thereof, are useful in treating and/or preventing bacterial infections and/or proliferative diseases (*e.g.*, cancer). Form A is a stable crystalline form of Compound **1**. Form A has been characterized by various techniques as described herein including, but not limited to, x-ray powder diffraction, differential scanning calorimetry, and thermogravimetric Fourier-transform infrared thermogram.

[0006] Also provided herein are pharmaceutical compositions comprising Form A of Compound **1** and optionally a pharmaceutically acceptable excipient.

[0007] The present invention also provides methods of treating a proliferative disease (*e.g.*, cancer) or a bacterial infection using Form A or pharmaceutical compositions thereof.

[0008] Also provided herein are methods of preparing the crystalline Form A of Compound **1**. In certain embodiments, the methods involve evaporative crystallization from chloroform/methanol or methanol/acetone. The provided methods are useful in preparing crystalline Form A of Compound **1** on a large-scale (*e.g.* over 50 grams) because the traditional fermentation process would be too voluminous (*e.g.* over 100L) for a normal laboratory to handle.

DEFINITIONS

[0009] The term “polymorph” refers to a crystalline form of a compound (*e.g.*, Compound **1**), or a hydrate or solvate thereof, in a particular crystal packing arrangement. All polymorphs of a particular compound have the same elemental composition. The term “crystalline,” as used herein, refers to a solid state form which consists of orderly arrangement of structural units. Different crystalline forms of the same compound, or a hydrate, or solvate thereof, arise from different packing of the molecules in the solid state, which results in different crystal symmetries and/or unit cell parameter. Different crystalline forms usually have different X-ray diffraction patterns, infrared spectra, melting points, densities, hardness, crystal shapes, optical and electrical properties, stabilities, and/or solubility. Recrystallization solvent, rate of crystallization, storage temperature, and other

factors may cause one crystalline form to dominate in a particular preparing. Various polymorphs of a compound, or a hydrate or solvate thereof, can be prepared by crystallization under different conditions.

[0010] The term “solvate” refers to forms of a compound (*e.g.*, Compound **1**) that are associated with a solvent, usually by a solvolysis reaction. This physical association may include hydrogen bonding. Conventional solvents include water, methanol, ethanol, acetic acid, DMSO, THF, diethyl ether, and the like. In certain embodiments, solvates are formed using Class 3 solvents. Categories of solvents are defined in, for example, the International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH), “Impurities: Guidelines for Residual Solvents, Q3C(R3), (November 2005). A compound may be prepared, *e.g.*, in crystalline form, and may be solvated. Suitable solvates include pharmaceutically acceptable solvates and further include both stoichiometric solvates and non-stoichiometric solvates. In certain instances, the solvate is capable of isolation, for example, when one or more solvent molecules are incorporated in the crystal lattice of a crystalline solid.

[0011] The term “hydrate,” refers to a compound (*e.g.*, Compound **1**) which is associated with water. Typically, the number of the water molecules contained in a hydrate of a compound is in a definite ratio to the number of the compound molecules in the hydrate. Hydrates include both stoichiometric hydrates and non-stoichiometric hydrates. Therefore, a hydrate of a compound may be represented, for example, by the general formula $R \cdot xH_2O$, wherein R is the organic compound, and x is a number greater than 0. A given compound may form more than one type of hydrate, including, *e.g.*, monohydrates (stoichiometric, x is 1), lower hydrates (non-stoichiometric, x is a number greater than 0 and smaller than 1, *e.g.*, hemihydrates ($R \cdot 0.5H_2O$)), and polyhydrates (non-stoichiometric, x is a number greater than 1, *e.g.*, dihydrates ($R \cdot 2H_2O$) and hexahydrates ($R \cdot 6 H_2O$)).

[0012] It is also to be understood that compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed “isomers”.

[0013] As used herein, the term “impurity” refers to extraneous matter included in a compound or composition (*e.g.*, Form A of Compound **1**). Extraneous matter includes one or more substances that are different from the compound of interest. In certain embodiments, the extraneous matter is undesired extraneous matter. For example, when an anhydrous compound is desired, the solvent (*e.g.*, water) included with the compound is considered an

impurity. When a crystalline compound is desired, an amorphous form of the compound included with the compound is considered an impurity. When a certain polymorph of a compound is desired, a different polymorph of the compound included with the compound is considered an impurity. The term “substantially free of impurities” means that a compound (*e.g.*, Form A of Compound **1**), contains no significant amount of extraneous matter (*e.g.*, undesired extraneous matter). In certain embodiments, about 1 wt%, about 2 wt%, about 3 wt%, about 5 wt%, about 7 wt%, or about 10 wt% of extraneous matter in a composition is a significant amount of extraneous matter.

[0014] A “subject” to which administration is contemplated includes, but is not limited to, humans (*i.e.*, a male or female of any age group, *e.g.*, a pediatric subject (*e.g.*, infant, child, adolescent) or adult subject (*e.g.*, young adult, middle-aged adult, or senior adult)) and/or other non-human animals, for example, mammals (*e.g.*, primates (*e.g.*, cynomolgus monkeys, rhesus monkeys); commercially relevant mammals such as cattle, pigs, horses, sheep, goats, cats, and/or dogs) and birds (*e.g.*, commercially relevant birds such as chickens, ducks, geese, and/or turkeys). In certain embodiments, the animal is a mammal. In certain embodiments, the subject is a non-human animal. In certain embodiments, the animal is human.

[0015] The terms “administer,” “administering,” or “administration,” as used herein, refers to implanting, absorbing, ingesting, injecting, inhaling, or otherwise introducing a compound (*e.g.*, Form A of Compound **1**) or pharmaceutical composition thereof, in or on a subject. When a compound of the invention is provided in combination with one or more other active agents, “administration” and its variants are each understood to include concurrent and/or sequential introduction of the compound and the other active agents.

[0016] As used herein, the terms “condition,” “disease,” and “disorder” are used interchangeably.

[0017] As used herein, and unless otherwise specified, the terms “treat,” “treating” and “treatment” contemplate an action that occurs while a subject is suffering from the specified disease, disorder or condition, which reduces the severity of the disease, disorder or condition, or retards or slows the progression of the disease, disorder or condition (“therapeutic treatment”), and also contemplates an action that occurs before a subject begins to suffer from the specified disease, disorder or condition (“prophylactic treatment”).

[0018] For example, in certain embodiments, the terms “treatment,” “treat,” and “treating” refer to administering a medicament (*e.g.*, Form A of Compound **1** or a pharmaceutical composition thereof) in order to reverse, alleviate, delay the onset of, or inhibit the progress of a “pathological condition” (*e.g.*, a disease, disorder, or condition, or one or more signs or

symptoms thereof) in a subject. In some embodiments, treatment may be administered after one or more signs or symptoms have developed or have been observed. In other embodiments, treatment may be administered in the absence of signs or symptoms of the disease or condition. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms. Treatment may also be continued after symptoms have resolved, for example, to delay or prevent recurrence.

[0019] The terms “prevention,” “prevent,” and “preventing,” as used herein, refer to administering a medicament (*e.g.*, Form A of Compound **1** or a pharmaceutical composition thereof) beforehand to avert or forestall the appearance of one or more symptoms of a disease or disorder in a subject. The person of ordinary skill in the medical art recognizes that the terms “prevention,” “prevent,” and “preventing” are not absolute terms. In the medical art these terms are understood to refer to the prophylactic administration of a medicament to substantially diminish the likelihood or seriousness of a condition, or symptom of the condition, and this is the sense intended in this disclosure.

[0020] An “effective amount” of a compound described herein refers to an amount sufficient to elicit the desired biological response, *e.g.*, treating or preventing a condition. As will be appreciated by those of ordinary skill in this art, the effective amount of a compound described herein may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the condition being treated, the mode of administration, and the age and health of the subject. An effective amount encompasses therapeutic and prophylactic treatment. For example, in treating cancer or a bacterial infection, an effective amount of the compound may provide a therapeutic and/or prophylactic benefit in the treatment and/or prevention of the cancer or bacterial infection, or to delay or minimize one or more symptoms associated with the cancer or bacterial infection.

[0021] A “therapeutically effective amount” of a compound described herein is an amount sufficient to provide a therapeutic benefit in the treatment of a condition (*e.g.*, cancer or bacterial infection) or to delay or minimize one or more symptoms associated with the condition. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment of the condition. The term “therapeutically effective amount” can encompass an amount that improves overall therapy, reduces symptoms or causes of the condition, or enhances the therapeutic efficacy of another therapeutic agent.

[0022] A “prophylactically effective amount” of a compound described herein is an amount sufficient to prevent a condition (*e.g.*, cancer or bacterial infection), or one or more

symptoms associated with the condition or prevent its recurrence. A prophylactically effective amount of a compound means an amount of a therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the condition. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

[0023] The details of one or more embodiments of the invention are set forth herein. Other features, objects, and advantages of the invention will be apparent from the Detailed Description, the Figures, the Examples, and the Claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024] *Figure 1* depicts an X-Ray Powder Diffraction (XRPD) pattern of Form A of Compound **1**.

[0025] *Figure 2* depicts another XRPD pattern of Form A from another batch.

[0026] *Figure 3* depicts a Differential Scanning Calorimetry (DSC) thermogram of the Form A.

[0027] *Figure 4* depicts a DSC thermogram of Form A from another batch.

[0028] *Figure 5* depicts a Thermogravimetric (TG) thermogram of Form A.

[0029] *Figure 6* depicts a TG thermogram of Form A from another batch.

[0030] *Figure 7* depicts a Solid-state Cross-Polarization Magic Angle Spinning Carbon-13 Nuclear Magnetic Resonance (^{13}C CP/MAS - NMR) spectrum of Form A.

[0031] *Figure 8* depicts a ^{13}C CP/MAS - NMR spectrum of Form A from another batch.

[0032] *Figure 9* depicts an exemplary process for producing Form A of Compound **1** (*i.e.* MSK-777).

[0033] *Figure 10* depicts the structures (flat and stereochemical depictions) of Compound **1** (MSK-777, Granaticin B) and Compound **2** (Granaticin A).


[0034] *Figure 11* shows a typical HPLC-profile of the resin extract from fermentation of Compound **1** (see Example 1).

[0035] *Figures 12A-12B* show the raw XRD data of Form A prepared from the first batch.

[0036] *Figures 13A-13B* show the raw XRD data of Form A prepared from the second batch.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

[0037] Granaticin B (Compound **1**) is a secondary metabolite antibiotic produced by *Streptomyces violaceoruber*. Compound **1** has shown significant potential for the treatment of a variety of disorders including bacterial infections and proliferative diseases. (*see, e.g.*, WO 2011/112635).



Solid Form

[0038] A crystalline polymorph of Compound **1** that, as compared to amorphous Compound **1**, imparts improved physical characteristics such as stability and/or ease of formulation is desirable. Accordingly, provided herein is a crystalline form (denoted Form A) of Compound **1**.

[0039] The crystalline Form A of Compound **1** was found to be more stable than amorphous Compound **1**. Amorphous Compound **1** is less stable and prone to degradation during the traditional purification process of granaticin B. In some embodiments, the crystalline Form A of Compound **1** demonstrates improved solubility (*e.g.* higher dissolution rate) compared to the amorphous Compound **1**.

[0040] In some embodiments, Form A has a water content of about 0.01 wt% to about 5.0 wt%. In some embodiments, Form A has a water content of about 2.0 wt%. Form A is substantially non-hygroscopic.

[0041] In some embodiments, Form A is substantially free of impurities. As used herein, impurities include, but are not limited to, any extraneous matter such residual solvents, salts, or other forms of granaticin B. In some embodiments, Form A is 99% free of impurities. In some embodiments, Form A is 97% free of impurities. In some embodiments, Form A is 95% free of impurities. In some embodiments, Form A is 92% free of impurities. In some embodiments, Form A is 90% free of impurities. In certain embodiments, the impurities include extraneous matter, such as a salt forming acid, residual solvents, or any other

impurities that may result from the preparation and/or isolation of Compound **1**. In some embodiments, Form A is substantially free of amorphous Compound **1**. In some embodiments, Form A is substantially free of another crystalline form of Compound **1**. In some embodiments, Form A is substantially free of a salt of Compound **1**. In some embodiments, Form A is substantially free of a solvate of Compound **1**.

[0042] Different solid forms of a compound typically differ in their physical and chemical properties based on the arrangement of the molecules in the solid form (*e.g.*, the arrangement of the molecule in the crystal lattice). A given substance may give rise to a variety of solid forms, in particular a variety of crystalline forms, wherein each form has different and distinct physical and chemical properties, such as solubility profiles, thermodynamic and chemical stabilities, melting points, and/or x-ray diffraction peaks.

[0043] The crystalline Form A can be characterized by one or more of the characteristics described herein including, but not limited to, XRPD diffraction pattern and/or peaks, DSC thermogram, TG-FTIR thermogram, and/or NMR peaks, appearance, melting point, solubility, and stability. In certain embodiments, the crystalline Form A is characterized by XRPD diffraction pattern and/or peaks. In certain embodiments, the crystalline Form A is characterized by XRPD diffraction pattern and/or peaks, and at least one other technique as described herein (*e.g.*, DSC thermogram, DVS isotherm, or TG-FTIR thermogram).

[0044] In some embodiments, the crystalline Form A is characterized by an X-ray powder diffraction pattern substantially similar to the one depicted in *Figure 1*. In some embodiments, the crystalline Form A is characterized by an X-ray powder diffraction pattern substantially similar to the one depicted in *Figure 2*. In some embodiments, the crystalline Form A is characterized in that it has one or more peaks in its X-ray powder diffraction pattern selected from those in Table 1. In some embodiments, the crystalline Form A is characterized by at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, at least ten, at least eleven, at least twelve, at least thirteen, at least fourteen, at least fifteen, at least sixteen, or at least seventeen peaks in its X-ray powder diffraction pattern selected from those in Table 1. In some embodiments, the crystalline Form A of Compound **1** is characterized in that it has one or more peaks in its X-ray powder diffraction pattern selected from the strong and very strong peaks in Table 1. In some embodiments, the crystalline Form A of Compound **1** is characterized in that it has all the peaks in its X-ray powder diffraction pattern selected from the strong and very strong peaks in Table 1. In some embodiments, the crystalline Form A of Compound **1** is characterized in that it has one or more peaks in its X-ray powder diffraction pattern selected

from the very strong peaks in Table 1. In some embodiments, the crystalline Form A of Compound **1** is characterized in that it has all the peaks in its X-ray powder diffraction pattern selected from the very strong peaks in Table 1. In some embodiments, the crystalline Form A of Compound **1** is characterized in that it has both very strong peaks listed in Table 1 (*i.e.*, 19.02 and 23.16 angle theta-2).

Table 1. X-ray powder diffraction pattern.

Intensity %	Angle 2-Theta °	Intensity (relative)
15	4.8 ± 0.2	m
48	8.6 ± 0.2	w
100	9.0 ± 0.2	vs
33	9.8 ± 0.2	s
25	11.5 ± 0.2	s
10	12.5 ± 0.2	m
8	13.2 ± 0.2	w
15	15.3 ± 0.2	m
25	16.1 ± 0.2	s
10	16.9 ± 0.2	m
8	17.3 ± 0.2	w
20	18.2 ± 0.2	s
6	19.6 ± 0.2	w
5	20.2 ± 0.2	w
6	21.5 ± 0.2	w
8	21.9 ± 0.2	w
8	23.2 ± 0.2	w

[0045] The terms used in the tables herein have the following meanings: The term “vs” stands for “very strong.” The term “s” stands for “strong.” The term “m” stands for “medium.” The term “w” stands for “weak.”

[0046] In some embodiments, the crystalline Form A is characterized by one or more peaks in its X-ray powder diffraction pattern selected from those in Table 2. In some embodiments, the crystalline Form A is characterized by at least one, at least two, at least three, at least four, at least five, at least six, at least seven, at least eight, or at least nine peaks in its X-ray powder diffraction pattern selected from those in Table 2. In some embodiments, the characteristic peaks include the very strong peak indicated in Table 2. In some embodiments, the crystalline Form A is characterized by the very strong peak indicated in

Table 2, and at least one, at least two, at least three, or at least four of the other strong peaks in its X-ray powder diffraction pattern selected from those in Table 2.

Table 2. Select characteristic peaks from the X-ray powder diffraction pattern.

Intensity %	Angle 2-Theta °	Intensity (relative)
100	9.0 ± 0.2	vs
33	9.8 ± 0.2	s
25	11.5 ± 0.2	s
25	16.1 ± 0.2	s
20	18.2 ± 0.2	s

[0047] In some embodiments, the crystalline Form A has a DSC thermogram substantially similar to the one depicted in *Figure 3*. In some embodiments, the crystalline Form A has a DSC thermogram substantially similar to the one depicted in *Figure 4*. In some embodiments, the crystalline Form A is characterized in that it has a DSC thermogram with an endotherm having a peak temperature ($T_{\max}$) of about 190 ± 0.2 °C. In some embodiments, the crystalline Form A is characterized by a DSC endothermogram with phase transition in the range of about 140° C to about 250° C. In some embodiments, the crystalline Form A is characterized by a DSC endothermogram with phase transition in the range of about 155° C to about 225° C.

[0048] In some embodiments, the crystalline Form A has a TG-FTIR thermogram substantially similar to the one depicted in *Figure 5*. In some embodiments, the crystalline Form A has a TG-FTIR thermogram substantially similar to the one depicted in *Figure 6*. In some embodiments, the crystalline Form A is characterized in that it losses about 2.0 wt% of H₂O after the temperature of the crystalline Form A is increased from 0 °C to about 150 °C. This mass decrease is related to evaporation of water and residual solvent from the crystals. No indication of hydrate(s) is seen due to lack of stepwise reduction of the mass below the melting point.

[0049] In some embodiments, the crystalline Form A has an observed decomposition point of about 190 ± 0.2 °C.

[0050] In some embodiments, the crystalline Form A is stable for at least about 1 month, at least about 2 months, at least about 4 months, at least about 6 months, at least about 12 months, at least about 18 months, at least about 24 months, or at least about 3 years at about 25 °C and about 60% relative humidity. In some embodiments, the crystalline Form A has

substantially the same XRPD pattern post storage for at least about 1 month, at least about 2 months, at least about 4 months, at least about 6 months, at least about 12 months, at least about 18 months, at least about 24 months, or at least about 3 years at about 25 °C and about 60% relative humidity.

[0051] In some embodiments, the crystalline Form A is stable for at least about 1 month, at least about 2 months, at least about 4 months, at least about 6 months, at least about 8 months, at least about 10 months, at least about 12 months, at least about 18 months, or at least about 24 months at about 40 °C and about 75% relative humidity. In some embodiments, the crystalline Form A has substantially the same XRPD pattern post storage for at least about 1 month, at least about 2 months, at least about 4 months, at least about 6 months, at least about 8 months, at least about 10 months, at least about 12 months, at least about 18 months, or at least about 24 months at about 40 °C and about 75% relative humidity.

Pharmaceutical Compositions

[0052] In some embodiments, the present invention provides a composition comprising the crystalline Form A of Compound **1** as described herein and optionally a pharmaceutically acceptable excipient. In certain embodiments, the pharmaceutical compositions are useful for treating a disease, disorder, or condition described herein. In some embodiments, the disease is a bacterial infection. In some embodiments, the disease is a proliferative disease. In some embodiments, a provided composition is formulated for administration to a subject in need of such composition. In certain embodiments, a provided composition is formulated for oral administration to a subject. In certain embodiments, a provided composition is formulated into an oral dosage form. In certain embodiments, a provided composition is formulated into a tablet, powder, pill, capsule, or the like, for oral ingestion by a subject.

[0053] Suitable techniques, carriers, and excipients include those found within, for example, *Remington: The Science and Practice of Pharmacy*, 19th edition, Mack Publishing Company, Easton, PA 1995; Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton, PA 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, NY 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th edition, Lippincott Williams & Wilkins, 1999, all of which are incorporated herein by reference in their entireties.

[0054] In general, doses of provided pharmaceutical compositions employed for adult human treatment are typically in the range of about 0.01 mg to about 5000 mg per day. In

certain embodiments, doses employed for adult human treatment are from about 1 mg to about 1000 mg per day. In certain embodiments, a desired dose is conveniently presented in a single dose or in divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example, as two, three, four or more sub-doses per day.

[0055] It will be understood that a specific dosage and treatment regimen for any particular subject may depend on a variety of factors, including the activity of the specific compound employed, age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a provided compound in the composition may also depend upon the particular compound in the composition.

Methods of Preparing Form A

[0056] The present invention also provides methods of preparing the crystalline Form A of Compound **1** as described herein. In certain embodiments, the methods involve evaporative crystallization. In certain embodiments, the methods comprise mixing Compound **1** with chloroform and methanol to provide a mixture. In certain embodiments, the chloroform is water free. In certain embodiments, the solution of Compound **1**, chloroform, and methanol is substantially homogeneous. In certain embodiments, the solution of Compound **1**, chloroform, and methanol is substantially free of solid materials. In some embodiments, Compound **1** is first dissolved in chloroform (100%) resulting in a solution. Methanol is then added to the solution, and chloroform is removed gradually from the mixture by evaporation. During the evaporation process, methanol is gradually added to the solution. Eventually the crystallization solution contains almost 100% methanol.

[0057] In other embodiments, methods of preparing the crystalline Form A of Compound **1** as described herein comprise mixing Compound **1** with methanol and acetone to generate a mixture. In some embodiments, the methods further comprise adding aqueous hydrochloride to acidify the mixture. In some embodiments, the acidified mixture has a water ratio of about 65 vol%. In some embodiments, the concentration of Compound **1** is about 3-6 g/L. In some embodiments, the concentration of Compound **1** is about 4-5 g/L. In certain embodiments, the mixture of Compound **1**, methanol, and acetone is substantially homogeneous. In certain embodiments, the mixture of Compound **1**, methanol, and acetone is substantially free of solid materials. In some embodiments, Compound **1** is dissolved in acetone (100%) resulting in a solution. Methanol is then added to the solution, and acetone is gradually removed from the mixture by evaporation. During the evaporation process, methanol is gradually added to

the solution to reach the target concentration. Eventually the crystallization mixture contains almost 100% methanol.

[0058] In certain embodiments, the methods of preparing the crystalline Form A of Compound **1** further comprise lowering the temperature of the mixture to provide a solid. The steps of preparing the crystalline Form A from chloroform and methanol may be performed at any suitable temperature, *e.g.*, a suitable temperature of about -30 °C to about 65 °C. Other ranges are also possible. In certain embodiments, the suitable temperature is from about 0 °C to about 30 °C. In certain embodiments, the suitable temperature is from about 15 °C to about 25 °C. In certain embodiments, the suitable temperature is about 0 °C. In certain embodiments, the suitable temperature is about 23 °C (room temperature).

[0059] The steps of preparing the crystalline Form A from methanol and acetone may be performed at any suitable temperature, *e.g.*, a suitable temperature of about -30 °C to about 65 °C. Other ranges are also possible. In certain embodiments, the suitable temperature is from about 0 °C to about 30 °C. In certain embodiments, the suitable temperature is about 0 °C. In certain embodiments, the suitable temperature is about 4 °C. In certain embodiments, the suitable temperature is about 23 °C (room temperature). In certain embodiments, the steps of preparing the crystalline Form A from methanol and acetone may be performed at one temperature, followed by a lower temperature, followed by a further lower temperature. In certain embodiments, the methods of preparing the crystalline Form A from methanol and acetone comprise stirring the acidified mixture at about 23 °C (room temperature), followed by about 4 °C, followed by about 0 °C.

[0060] In certain embodiments, the methods of preparing the crystalline Form A of Compound **1** further comprise isolating the solid from the mixture.

[0061] Compound **1** useful in the preparation of the crystalline Form A may be substantially free of impurities. In certain embodiments, Compound **1** useful in the preparation of the crystalline Form A is about 90% free of impurities. In certain embodiments, Compound **1** useful in the preparation of the crystalline Form A is about 92% free of impurities. In certain embodiments, Compound **1** useful in the preparation of the crystalline Form A is about 95% free of impurities. In certain embodiments, Compound **1** useful in the preparation of the crystalline Form A is about 97% free of impurities. In certain embodiments, Compound **1** useful in the preparation of the crystalline Form A is about 99% free of impurities. In certain embodiments, Compound **1** useful in the preparation of the crystalline Form A is about 99.5% free of impurities. In certain embodiments, Compound **1**

useful in the preparation of the crystalline Form A includes a lactone ring opened product as an impurity.

[0062] Compound **1** may be present in the solution of Compound **1**, chloroform, and methanol at any suitable concentration (*e.g.*, about 0.003 kg/L, about 0.01 kg/L, about 0.02 kg/L, about 0.03 kg/L, about 0.04 kg/L, about 0.05 kg/L, about 0.06 kg/L, about 0.08 kg/L, about 0.1 kg/L, about 0.2 kg/L, about 0.5 kg/L, or about 1 kg/L), as the solubility of Compound **1** permits. In certain embodiments, the concentration of Compound **1** in the solution of Compound **1** and methanol is about 10-100 g/L.

[0063] When the mixture of the inventive methods comprises a solid, the solid may be isolated from the mixture by a process known in the art, such as by filtration and/or centrifuge. The solid isolated from the mixture may optionally be subject to a reduced pressure and/or a suitable temperature as described herein. In certain embodiments, the solid in the mixture comprises the crystalline Form A. In certain embodiments, the solid in the mixture comprises the crystalline Form A that is substantially free of impurities. In certain embodiments, the solid isolated from the mixture comprises the crystalline Form A. In certain embodiments, the solid isolated from the mixture comprises the crystalline Form A that is substantially free of impurities. In certain embodiments, the crystalline Form A isolated from the mixture comprises at least 99%, at least 99.2%, at least 99.4%, at least 99.5%, at least 99.6%, at least 99.7%, at least 99.8%, at least 99.9%, at least 99.95%, at least 99.99%, at least 99.995%, or at least 99.999% the crystalline Form A by weight.

[0064] In certain embodiments, the mixture is substantially free of solid materials before the temperature of the mixture is lowered. In certain embodiments, the mixture is heterogeneous after the temperature of it is lowered. In certain embodiments, the mixture comprises a solid after the temperature of the mixture is lowered. In certain embodiments, the mixture comprises a solid and a liquid after the temperature of the mixture is lowered.

[0001] A suitable condition may also include a suitable pressure under which one or more steps of the inventive methods are performed. In certain embodiments, the suitable pressure is about 1 atmosphere. A suitable pressure may also be higher or lower than 1 atmosphere (*i.e.*, a reduced pressure). A reduced pressure may be a pressure lower than about 10^{-1} atmosphere, lower than about 10^{-2} atmosphere, lower than about 10^{-3} atmosphere, lower than about 10^{-4} atmosphere, lower than about 10^{-5} atmosphere, lower than about 10^{-6} atmosphere, lower than about 10^{-7} atmosphere, lower than about 10^{-8} atmosphere, lower than about 10^{-9} atmosphere, lower than about 10^{-10} atmosphere, or lower than about 10^{-11} atmosphere.

[0002] A suitable condition may also include a suitable atmosphere under which one or more steps of the inventive methods are performed. In certain embodiments, the suitable atmosphere is air. In certain embodiments, the suitable atmosphere is an inert atmosphere. In certain embodiments, the suitable atmosphere is a nitrogen or argon atmosphere.

[0065] A suitable condition may also include a suitable time duration that one or more steps of the method lasts. In certain embodiments, the suitable time duration is in the order of minutes (*e.g.*, about 30 min), hours (*e.g.*, about 1 hour, about 2 hours, about 3 hours, about 6 hours, or about 12 hours), days (*e.g.*, about 1 day or about 2 days) or weeks (*e.g.*, about 1 week). For example, in the step of lowering the temperature of the mixture, the temperature of the mixture may be lowered over a suitable time duration described herein.

Treatment Methods

[0066] The present invention provides compounds and pharmaceutical compositions useful for inhibiting the growth of or kill rapidly dividing cells comprising administering an effective amount of the Crystalline Form A of Compound **1** as described herein to a subject in need of treatment. In some embodiments, the Crystalline Form A of Compound **1** also contemplates the treatment of a disease, disorder, or condition associated with abnormal cellular proliferation, such as cancer, autoimmune diseases, inflammatory diseases, and diabetic retinopathy.

[0067] In one aspect, provided is a method of treating a proliferative disease comprising administering an effective amount of the Crystalline Form A of Compound **1** of the present invention to a subject in need thereof.

[0068] Thus, in one aspect, provided is a method of treating cancer comprising administering an effective amount of the Crystalline Form A of Compound **1** of the present invention to a subject in need thereof.

[0069] In another aspect, provided is a method of treating an autoimmune disease comprising administering an effective amount of the Crystalline Form A of Compound **1** of the present invention to a subject in need thereof.

[0070] In yet another aspect, provided is a method of treating an inflammatory disease comprising administering the Crystalline Form A of Compound **1** of the present invention to a subject in need thereof.

[0071] A “proliferative disease” refers to a disease that occurs due to abnormal growth or extension by the multiplication of cells (Walker, *Cambridge Dictionary of Biology*; Cambridge University Press: Cambridge, UK, 1990). A proliferative disease may be

associated with: 1) the pathological proliferation of normally quiescent cells; 2) the pathological migration of cells from their normal location (*e.g.*, metastasis of neoplastic cells); 3) the pathological expression of proteolytic enzymes such as the matrix metalloproteinases (*e.g.*, collagenases, gelatinases, and elastases); or 4) the pathological angiogenesis as in proliferative retinopathy and tumor metastasis. Exemplary proliferative diseases include cancers (*i.e.*, “malignant neoplasms”), benign neoplasms, angiogenesis, inflammatory diseases, and autoimmune diseases.

[0072] The term “angiogenesis” refers to the physiological process through which new blood vessels form from pre-existing vessels. Angiogenesis is distinct from vasculogenesis, which is the *de novo* formation of endothelial cells from mesoderm cell precursors. The first vessels in a developing embryo form through vasculogenesis, after which angiogenesis is responsible for most blood vessel growth during normal or abnormal development. Angiogenesis is a vital process in growth and development, as well as in wound healing and in the formation of granulation tissue. However, angiogenesis is also a fundamental step in the transition of tumors from a benign state to a malignant one, leading to the use of angiogenesis inhibitors in the treatment of cancer. Angiogenesis may be chemically stimulated by angiogenic proteins, such as growth factors (*e.g.*, VEGF). “Pathological angiogenesis” refers to abnormal (*e.g.*, excessive or insufficient) angiogenesis that amounts to and/or is associated with a disease.

[0073] The terms “neoplasm” and “tumor” are used herein interchangeably and refer to an abnormal mass of tissue wherein the growth of the mass surpasses and is not coordinated with the growth of a normal tissue. A neoplasm or tumor may be “benign” or “malignant,” depending on the following characteristics: degree of cellular differentiation (including morphology and functionality), rate of growth, local invasion, and metastasis. A “benign neoplasm” is generally well differentiated, has characteristically slower growth than a malignant neoplasm, and remains localized to the site of origin. In addition, a benign neoplasm does not have the capacity to infiltrate, invade, or metastasize to distant sites. Exemplary benign neoplasms include, but are not limited to, lipoma, chondroma, adenomas, acrochordon, senile angiomas, seborrheic keratoses, lentigos, and sebaceous hyperplasias. In some cases, certain “benign” tumors may later give rise to malignant neoplasms, which may result from additional genetic changes in a subpopulation of the tumor’s neoplastic cells, and these tumors are referred to as “pre-malignant neoplasms.” An exemplary pre-malignant neoplasm is a teratoma. In contrast, a “malignant neoplasm” is generally poorly differentiated (anaplasia) and has characteristically rapid growth accompanied by progressive infiltration,

invasion, and destruction of the surrounding tissue. Furthermore, a malignant neoplasm generally has the capacity to metastasize to distant sites. The term “metastasis,” “metastatic,” or “metastasize” refers to the spread or migration of cancerous cells from a primary or original tumor to another organ or tissue and is typically identifiable by the presence of a “secondary tumor” or “secondary cell mass” of the tissue type of the primary or original tumor and not of that of the organ or tissue in which the secondary (metastatic) tumor is located. For example, a prostate cancer that has migrated to bone is said to be metastasized prostate cancer and includes cancerous prostate cancer cells growing in bone tissue.

[0074] The term “cancer” refers to a class of diseases characterized by the development of abnormal cells that proliferate uncontrollably and have the ability to infiltrate and destroy normal body tissues. See, e.g., *Stedman’s Medical Dictionary*, 25th ed.; Hensyl ed.; Williams & Wilkins: Philadelphia, 1990. Exemplary cancers include, but are not limited to, acoustic neuroma; adenocarcinoma; adrenal gland cancer; anal cancer; angiosarcoma (e.g., lymphangiosarcoma, lymphangioendotheliosarcoma, hemangiosarcoma); appendix cancer; benign monoclonal gammopathy; biliary cancer (e.g., cholangiocarcinoma); bladder cancer; breast cancer (e.g., adenocarcinoma of the breast, papillary carcinoma of the breast, mammary cancer, medullary carcinoma of the breast); brain cancer (e.g., meningioma, glioblastomas, glioma (e.g., astrocytoma, oligodendroglioma), medulloblastoma); bronchus cancer; carcinoid tumor; cervical cancer (e.g., cervical adenocarcinoma); choriocarcinoma; chordoma; craniopharyngioma; colorectal cancer (e.g., colon cancer, rectal cancer, colorectal adenocarcinoma); connective tissue cancer; epithelial carcinoma; ependymoma; endotheliosarcoma (e.g., Kaposi’s sarcoma, multiple idiopathic hemorrhagic sarcoma); endometrial cancer (e.g., uterine cancer, uterine sarcoma); esophageal cancer (e.g., adenocarcinoma of the esophagus, Barrett’s adenocarcinoma); Ewing’s sarcoma; ocular cancer (e.g., intraocular melanoma, retinoblastoma); familial hypereosinophilia; gall bladder cancer; gastric cancer (e.g., stomach adenocarcinoma); gastrointestinal stromal tumor (GIST); germ cell cancer; head and neck cancer (e.g., head and neck squamous cell carcinoma, oral cancer (e.g., oral squamous cell carcinoma), throat cancer (e.g., laryngeal cancer, pharyngeal cancer, nasopharyngeal cancer, oropharyngeal cancer)); hematopoietic cancers (e.g., leukemia such as acute lymphoblastic leukemia (ALL) (e.g., B-cell ALL, T-cell ALL), acute myeloid leukemia (AML) (e.g., B-cell AML, T-cell AML), biphenotypic acute leukemia, chronic myeloid leukemia (CML) (e.g., B-cell CML, T-cell CML), and chronic lymphocytic leukemia (CLL) (e.g., B-cell CLL, T-cell CLL)); lymphoma such as Hodgkin lymphoma (HL) (e.g., B-cell HL, T-cell HL) and non-Hodgkin lymphoma (NHL) (e.g., B-cell NHL such

as diffuse large cell lymphoma (DLCL) (*e.g.*, diffuse large B-cell lymphoma), follicular lymphoma, chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma (MCL), marginal zone B-cell lymphomas (*e.g.*, mucosa-associated lymphoid tissue (MALT) lymphomas, nodal marginal zone B-cell lymphoma, splenic marginal zone B-cell lymphoma), primary mediastinal B-cell lymphoma, Burkitt lymphoma, lymphoplasmacytic lymphoma (*i.e.*, Waldenström's macroglobulinemia), hairy cell leukemia (HCL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma and primary central nervous system (CNS) lymphoma; and T-cell NHL such as precursor T-lymphoblastic lymphoma/leukemia, peripheral T-cell lymphoma (PTCL) (*e.g.*, cutaneous T-cell lymphoma (CTCL) (*e.g.*, mycosis fungoides, Sezary syndrome), angioimmunoblastic T-cell lymphoma, extranodal natural killer T-cell lymphoma, enteropathy type T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, and anaplastic large cell lymphoma); a mixture of one or more leukemia/lymphoma as described above; and multiple myeloma (MM)), heavy chain disease (*e.g.*, alpha chain disease, gamma chain disease, mu chain disease); hemangioblastoma; hypopharynx cancer; inflammatory myofibroblastic tumors; immunocytic amyloidosis; kidney cancer (*e.g.*, nephroblastoma *a.k.a.* Wilms' tumor, renal cell carcinoma); liver cancer (*e.g.*, hepatocellular cancer (HCC), malignant hepatoma); lung cancer (*e.g.*, bronchogenic carcinoma, small cell lung cancer (SCLC), non-small cell lung cancer (NSCLC), adenocarcinoma of the lung); leiomyosarcoma (LMS); mastocytosis (*e.g.*, systemic mastocytosis); muscle cancer; myelodysplastic syndrome (MDS); mesothelioma; myeloproliferative disorder (MPD) (*e.g.*, polycythemia vera (PV), essential thrombocytosis (ET), agnogenic myeloid metaplasia (AMM) *a.k.a.* myelofibrosis (MF), chronic idiopathic myelofibrosis, chronic myelocytic leukemia (CML), chronic neutrophilic leukemia (CNL), hypereosinophilic syndrome (HES)); neuroblastoma; neurofibroma (*e.g.*, neurofibromatosis (NF) type 1 or type 2, schwannomatosis); neuroendocrine cancer (*e.g.*, gastroenteropancreatic neuroendocrine tumor (GEP-NET), carcinoid tumor); osteosarcoma (*e.g.*, bone cancer); ovarian cancer (*e.g.*, cystadenocarcinoma, ovarian embryonal carcinoma, ovarian adenocarcinoma); papillary adenocarcinoma; pancreatic cancer (*e.g.*, pancreatic adenocarcinoma, intraductal papillary mucinous neoplasm (IPMN), Islet cell tumors); penile cancer (*e.g.*, Paget's disease of the penis and scrotum); pinealoma; primitive neuroectodermal tumor (PNT); plasma cell neoplasia; paraneoplastic syndromes; intraepithelial neoplasms; prostate cancer (*e.g.*, prostate adenocarcinoma); rectal cancer; rhabdomyosarcoma; salivary gland cancer; skin cancer (*e.g.*, squamous cell carcinoma (SCC), keratoacanthoma (KA), melanoma, basal cell carcinoma (BCC)); small bowel cancer (*e.g.*, appendix cancer); soft

tissue sarcoma (*e.g.*, malignant fibrous histiocytoma (MFH), liposarcoma, malignant peripheral nerve sheath tumor (MPNST), chondrosarcoma, fibrosarcoma, myxosarcoma); sebaceous gland carcinoma; small intestine cancer; sweat gland carcinoma; synovioma; testicular cancer (*e.g.*, seminoma, testicular embryonal carcinoma); thyroid cancer (*e.g.*, papillary carcinoma of the thyroid, papillary thyroid carcinoma (PTC), medullary thyroid cancer); urethral cancer; vaginal cancer; and vulvar cancer (*e.g.*, Paget's disease of the vulva).

[0075] As used herein "inhibition," "inhibiting," and "inhibit", refer to the ability of a compound to reduce, slow, halt or prevent activity of a particular biological process in a cell relative to vehicle. In certain embodiments, the biological process is *in vitro* (*e.g.*, cellular assay). In certain embodiments, the biological process is *in vivo*.

[0076] In certain embodiments, the provided methods for cancer treatment is to treat leukemia, thyroid cancer, melanoma, ovarian cancer, lung cancer, prostate cancer, renal cell carcinoma, cervical cancer, or breast cancer.

[0077] In certain embodiments, the leukemia is acute myeloid leukemia, chronic lymphocytic leukemia, acute lymphoblastic leukemia, or biphenotypic acute leukemia.

[0078] In another aspect, provided herein is a method of inhibiting cell growth in a subject comprising administering to the subject with a therapeutically effective amount of the crystalline Form A as described herein or a pharmaceutical composition thereof.

[0079] In another aspect, provided herein is a method of inhibiting cell growth in a biological sample comprising contacting the biological sample with a therapeutically effective amount of the crystalline Form A as described herein or a pharmaceutical composition thereof.

[0080] In another aspect, provided herein is a method of inducing apoptosis of a cell in a subject comprising administering to the subject with a therapeutically effective amount of the crystalline Form A or a pharmaceutical composition thereof.

[0081] In another aspect, provided herein is a method of inducing apoptosis of a cell in a biological sample comprising contacting the biological sample with a therapeutically effective amount of the crystalline Form A as described herein or a pharmaceutical composition thereof.

[0082] In another aspect, the present invention provides compounds and pharmaceutical compositions useful for treating or preventing microbial infection comprising administering an effective amount of the Crystalline Form A of Compound **1** as described herein to a subject in need of treatment. In some embodiments, the microbial infection is a bacterial

infection. In one aspect, the present invention provides methods for inhibiting bacterial growth comprising administering an effective amount of the Crystalline Form A of Compound **1** as described herein to a subject in need of treatment.

[0083] In yet another aspect, provided is a method of treating or preventing microbial infection caused by pathogen that are resistant to other treatments, are multi-drug tolerant or resistant and/or that neither grow nor die in the presence of or as a result of other treatments. Such a method can be conducted *in vivo* (*i.e.*, by administration to a subject) or *in vitro* (*e.g.*, upon contact with bacteria in a cell culture). For example, in certain embodiments, provided is a method of treating and/or preventing a microbial infection resistant to other treatments comprising administering an effective amount of the Crystalline Form A of Compound **1** to a subject with the microbial infection. In certain embodiments, provided is a method of treating and/or preventing a microbial infection caused by bacteria that are multi-drug resistant, comprising administering an effective amount of the Crystalline Form A of Compound **1** to a subject with the infection. In certain embodiments, the infection-causing bacteria are resistant to other treatments. In some embodiments, the infection-causing bacteria are multi-drug tolerant. In some embodiments, the infection-causing bacteria are multi-drug resistant.

[0084] In another aspect, provided herein is a method for inhibiting bacterial cell growth comprising contacting bacteria with the crystalline Form A as described herein, or a pharmaceutical composition thereof. In another aspect, provided herein is a method for inducing bacterial hypersusceptibility comprising contacting bacteria with the crystalline Form A as described herein, or a pharmaceutical composition thereof.

[0085] In some embodiments of the provided methods, the bacteria are contacted with the crystalline Form A *in vitro*. In some embodiments of the provided methods, the bacteria are contacted with the crystalline Form A *in vivo*.

[0086] In some embodiments of the provided methods, the crystalline Form A of Compound **1** is administered with a second therapeutic agent as defined herein. In some embodiments, the second therapeutic agent is an antibiotic.

[0087] Exemplary bacterial infections include, but are not limited to, infections with a gram positive bacteria (*e.g.*, of the phylum *Actinobacteria*, phylum *Firmicutes*, or phylum *Tenericutes*); gram negative bacteria (*e.g.*, of the phylum *Aquificae*, phylum *Deinococcus-Thermus*, phylum *Fibrobacteres/Chlorobi/Bacteroidetes* (FCB), phylum *Fusobacteria*, phylum *Gemmatimonadest*, phylum *Nitrospirae*, phylum *Planctomycetes/Verrucomicrobia/Chlamydiae* (PVC), phylum *Proteobacteria*, phylum *Spirochaetes*, or phylum *Synergistetes*); or other bacteria (*e.g.*, of the phylum *Acidobacteria*,

phylum *Chloroflexi*, phylum *Chrysiogenetes*, phylum *Cyanobacteria*, phylum *Deferribacteres*, phylum *Dictyoglomi*, phylum *Thermodesulfobacteria*, or phylum *Thermotogae*).

[0088] In certain embodiments, the bacterial infection is an infection with a gram positive bacteria. In certain embodiments, the gram positive bacteria are bacteria of the phylum *Firmicutes*.

[0089] In certain embodiments, the bacteria are a member of the phylum *Firmicutes* and the genus *Enterococcus*, i.e., the bacterial infection is an *Enterococcus* infection. The term “*Enterococcus species*” refers to a genus of lactic acid bacteria of the phylum *Firmicutes*. They are Gram-positive cocci which often occur in pairs (diplococci for example *Diplococcus pneumoniae*). *Enterococci* are facultative anaerobic organisms. Exemplary *Enterococci* bacteria include, but are not limited to, *E. avium*, *E. durans*, *E. faecalis*, *E. faecium*, *E. gallinarum*, *E. solitarius*, *E. casseliflavus*, and *E. raffinosus*. In certain embodiments, the *Enterococcus* infection is an *E. faecalis* infection. In certain embodiments, the *Enterococcus* infection is an *E. faecium* infection.

[0090] In certain embodiments, the bacteria are a member of the phylum *Firmicutes* and the genus *Staphylococcus*, i.e., the bacterial infection is a *Staphylococcus* infection. The term “*Staphylococcus species*” refers to Gram-positive bacteria, which appear as grape-like clusters when viewed through a microscope and as large, round, golden-yellow colonies, often with .beta.-hemolysis, when grown on blood agar plates. Exemplary *Staphylococci* bacteria include, but are not limited to, *S. arlettae*, *S. aureus*, *S. auricularis*, *S. capitis*, *S. caprae*, *S. carnosus*, *S. chromogenes*, *S. cohi*, *S. condimenti*, *S. croceolyticus*, *S. delphini*, *S. devriesei*, *S. epidermis*, *S. equorum*, *S. felis*, *S. fluroettii*, *S. gallinarum*, *S. haemolyticus*, *S. hominis*, *S. hyicus*, *S. intermedius*, *S. kloosii*, *S. leei*, *S. lenus*, *S. lugdunensis*, *S. lutrae*, *S. lyticans*, *S. massiliensis*, *S. microti*, *S. muscae*, *S. nepalensis*, *S. pasteurii*, *S. penttenkoferi*, *S. piscifermentans*, *S. pseudointermedius*, *S. pseudolugdunensis*, *S. pulvereri*, *S. rostri*, *S. saccharolyticus*, *S. saprophyticus*, *S. schleiferi*, *S. sciuri*, *S. simiae*, *S. simulans*, *S. stepanovicii*, *S. succinus*, *S. vitulinus*, *S. warneri*, and *S. xylosus*. In certain embodiments, the *Staphylococcus* infection is an *S. aureus* infection. In certain embodiments, the *Staphylococcus* infection is an *S. epidermis* infection.

[0091] In certain embodiments, the bacteria are a member of the phylum *Firmicutes* and the genus *Streptococcus*, i.e., the bacterial infection is a *Streptococcus* infection. The term “*Streptococcus species*” refers to a genus of spherical, Gram-positive bacteria, and a member of the phylum *Firmicutes*. *Streptococci* are lactic acid bacteria. *Streptococcus species*

includes bacteria such as, for example, *S. hemolyticus*, *S. mitis*, *S. salivarius*, and *S. pneumoniae*. *Streptococcus* species are responsible for infectious diseases such as meningitis, bacterial pneumonia, endocarditis, erysipelas and necrotizing fasciitis (“flesh-eating” microbial infections).

[0092] In certain embodiments, the bacteria are a member of the phylum *Firmicutes* and the genus *Bacillus*, *i.e.*, the bacterial infection is a *Bacillus* infection. The term “*Bacillus species*” refers to a large number of diverse, rod-shaped Gram positive bacteria that are motile by peritrichous flagella and are aerobic such as *B. anthracis* or *B. subtilis*.

[0093] In certain embodiments, the bacteria are a member of the phylum *Actinobacteria* and the family *Mycobacterium*, *i.e.*, the bacterial infection is a *Mycobacterium* infection. The term “*Mycobacterium species*” refers to Gram-positive, non-motile, pleomorphic rods related to the actinomyces. Tuberculosis in humans is caused by *Mycobacterium tuberculosis*. MDR-TB (multi-drug resistant tuberculosis) describes strains of tuberculosis that are resistant to at least the two first-line TB drugs, isoniazid and rifampicin.

[0094] In certain embodiments, the bacterial infection is resistant to other antibiotic therapy. For example, in certain embodiments, the bacterial infection is vancomycin resistant (VR). In certain embodiments, the bacterial infection is a vancomycin-resistant *E. faecalis* infection. In certain embodiments, the bacterial infection is a vancomycin-resistant *E. faecium* infection. In certain embodiments, the bacterial infection is a methicillin-resistant (MR). In certain embodiments, the bacterial infection is a methicillin-resistant *S. aureus* (MRSA) infection.

[0095] In certain embodiments, the effective amount is a therapeutically effective amount. In certain embodiments, the effective amount is a prophylactically effective amount. In certain embodiments, the subject is suffering from at least one bacterial infection.

[0096] The Crystalline Form A of Compound **1** provided herein is typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject or organism will depend upon a variety of factors including the disease, disorder, or condition being treated and the severity of the disorder; the activity of the specific active ingredient employed; the specific composition employed; the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific active

ingredient employed; the duration of the treatment; drugs used in combination or coincidental with the specific active ingredient employed; and like factors well known in the medical arts.

[0097] The Crystalline Form A of Compound **1** and compositions provided herein can be administered by any route, including enteral (*e.g.*, oral), parenteral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, subcutaneous, intraventricular, transdermal, interdermal, rectal, intravaginal, intraperitoneal, topical (as by powders, ointments, creams, and/or drops), mucosal, nasal, bucal, sublingual; by intratracheal instillation, bronchial instillation, and/or inhalation; and/or as an oral spray, nasal spray, and/or aerosol. Specifically contemplated routes are oral administration, intravenous administration (*e.g.*, systemic intravenous injection), regional administration via blood and/or lymph supply, and/or direct administration to an affected site. In general the most appropriate route of administration will depend upon a variety of factors including the nature of the agent (*e.g.*, its stability in the environment of the gastrointestinal tract), the condition of the subject (*e.g.*, whether the subject is able to tolerate oral administration), *etc.*

[0098] The exact amount of Compound **1** required to achieve an effective amount will vary from subject to subject, depending, for example, on species, age, and general condition of a subject, severity of the side effects or disorder, identity of the mode of administration, and the like. The desired dosage can be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. In certain embodiments, the desired dosage can be delivered using multiple administrations (*e.g.*, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations).

[0099] In certain embodiments, an effective amount of Compound **1** for administration one or more times a day to a 70 kg adult human may comprise about 0.0001 mg to about 3000 mg, about 0.0001 mg to about 2000 mg, about 0.0001 mg to about 1000 mg, about 0.001 mg to about 1000 mg, about 0.01 mg to about 1000 mg, about 0.1 mg to about 1000 mg, about 1 mg to about 1000 mg, about 1 mg to about 100 mg, about 10 mg to about 1000 mg, or about 100 mg to about 1000 mg, of a compound per unit dosage form.

[00100] In certain embodiments, Compound **1** and the compositions thereof may be administered orally or parenterally at dosage levels sufficient to deliver from about 0.001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 50 mg/kg, preferably from about 0.1 mg/kg to about 40 mg/kg, preferably from about 0.5 mg/kg to about 30 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, and more

preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[00101] It will be appreciated that dose ranges as described herein provide guidance for the administration of provided pharmaceutical compositions to an adult. The amount to be administered to, for example, a child or an adolescent can be determined by a medical practitioner or person skilled in the art and can be lower or the same as that administered to an adult.

[00102] It will be also appreciated that the Crystalline Form A of Compound **1** or composition thereof, as described herein, can be administered in combination with a second therapeutic agent. The Crystalline Form A of Compound **1** or compositions thereof can be administered in combination with a second therapeutic agent to improve bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body. It will also be appreciated that the therapy employed may achieve a desired effect for the same disorder, and/or it may achieve different effects.

[00103] When “combination therapy” is employed, an effective amount can be achieved using a first amount of Compound **1**, or a pharmaceutical composition thereof, and a second amount of an additional suitable therapeutic agent.

[00104] In certain embodiments, Compound **1** or a pharmaceutical composition thereof as described herein, and the additional therapeutic agent are each administered in an effective amount (i.e., each in an amount which would be therapeutically effective if administered alone). In other embodiments, Compound **1** or a pharmaceutical composition thereof as described herein, and the additional therapeutic agent are each administered in an amount which alone does not provide a therapeutic effect (a sub-therapeutic dose). In yet other embodiments, Compound **1** or a pharmaceutical composition thereof as described herein can be administered in an effective amount, while the additional therapeutic agent is administered in a sub-therapeutic dose. In still other embodiments, Compound **1** or a pharmaceutical composition thereof as described herein, can be administered in a sub-therapeutic dose, while the additional therapeutic agent is administered in an effective amount.

[00105] As used herein, the terms “in combination” or “co-administration” can be used interchangeably to refer to the use of more than one therapy (*e.g.*, one or more prophylactic and/or therapeutic agents). The use of the terms does not restrict the order in which therapies (*e.g.*, prophylactic and/or therapeutic agents) are administered to a subject.

[00106] Co-administration encompasses administration of the first and second amounts of the compounds in an essentially simultaneous manner, such as in a single pharmaceutical

composition, for example, capsule or tablet having a fixed ratio of first and second amounts, or in multiple, separate capsules or tablets for each. In addition, such co- administration also encompasses use of each compound in a sequential manner in either order. When co-administration involves the separate administration of the first amount of Compound **1** or a pharmaceutical composition thereof as described herein, and a second amount of an additional therapeutic agent, the compounds are administered sufficiently close in time to have the desired therapeutic effect. For example, the period of time between each administration which can result in the desired therapeutic effect, can range from minutes to hours and can be determined taking into account the properties of each compound such as potency, solubility, bioavailability, plasma half-life, and kinetic profile. For example, Compound **1** or a pharmaceutical composition thereof as described herein, and the second therapeutic agent can be administered in any order within about 24 hours of each other, within about 16 hours of each other, within about 8 hours of each other, within about 4 hours of each other, within about 1 hour of each other or within about 30 minutes of each other.

[00107] More, specifically, a first therapy (*e.g.*, a prophylactic or therapeutic agent such as a compound described herein) can be administered prior to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy to a subject.

[00108] The Crystalline Form A of Compound **1** or compositions thereof can be administered concurrently with, prior to, or subsequent to, one or more additional therapeutically active agents. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. It will further be appreciated that the additional therapeutically active agent utilized in this combination can be administered together in a single composition or administered separately in different compositions. The particular combination to employ in a regimen will take into account compatibility of the Crystalline Form A of Compound **1** with the additional therapeutically active agent and/or the desired therapeutic effect to be achieved. In general, it is expected that additional therapeutically active agents utilized in combination do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

[00109] Exemplary second therapeutic agents include, but are not limited to, antibiotics, anti-viral agents, anesthetics, anti-coagulants, inhibitors of an enzyme, steroidal agents, steroidal or non-steroidal anti-inflammatory agents, antihistamine, immunosuppressant agents, antigens, vaccines, antibodies, decongestant, sedatives, opioids, pain-relieving agents, analgesics, anti-pyretics, hormones, and prostaglandins, *etc.* Therapeutically active agents include small organic molecules such as drug compounds (*e.g.*, compounds approved by the US Food and Drug Administration as provided in the Code of Federal Regulations (CFR)), peptides, proteins, carbohydrates, monosaccharides, oligosaccharides, polysaccharides, nucleoproteins, mucoproteins, lipoproteins, synthetic polypeptides or proteins, small molecules linked to proteins, glycoproteins, steroids, nucleic acids, DNAs, RNAs, nucleotides, nucleosides, oligonucleotides, antisense oligonucleotides, lipids, hormones, vitamins and cells.

[00110] In certain embodiments, the additional therapeutic agent is an antibiotic. Exemplary antibiotics include, but are not limited to, penicillins (*e.g.*, penicillin, amoxicillin), cephalosporins (*e.g.*, cephalexin), macrolides (*e.g.*, erythromycin, clarithromycin, azithromycin, troleandomycin), fluoroquinolones (*e.g.*, ciprofloxacin, levofloxacin, ofloxacin), sulfonamides (*e.g.*, co-trimoxazole, trimethoprim), tetracyclines (*e.g.*, tetracycline, chlortetracycline, oxytetracycline, demeclocycline, methacycline, sancycline, doxycycline, aureomycin, terramycin, minocycline, 6-deoxytetracycline, lymecycline, meclocycline, methacycline, rolitetracycline, and glycylcycline antibiotics (*e.g.*, tigecycline)), aminoglycosides (*e.g.*, gentamicin, tobramycin, paromomycin), aminocyclitol (*e.g.*, spectinomycin), chloramphenicol, sparsomycin, quinupristin/dalfoprisin (Syndercid™),

[00111] In some embodiments, the additional therapeutic agent is an anti-proliferative agent. In some embodiments, the anti-proliferative agent is an anti-cancer agent. As used herein, the anti-cancer agents encompass biotherapeutic anti-cancer agents as well as chemotherapeutic agents.

[00112] Exemplary biotherapeutic anti-cancer agents include, but are not limited to, interferons, cytokines (*e.g.*, tumor necrosis factor, interferon α , interferon γ), vaccines, hematopoietic growth factors, monoclonal serotherapy, immunostimulants and/or immunomodulatory agents (*e.g.*, IL-1, 2, 4, 6, or 12), immune cell growth factors (*e.g.*, GM-CSF) and antibodies (*e.g.* HERCEPTIN (trastuzumab), T-DM1, AVASTIN (bevacizumab), ERBITUX (cetuximab), VECTIBIX (panitumumab), RITUXAN (rituximab), BEXXAR (tositumomab)).

[00113] Exemplary chemotherapeutic agents include, but are not limited to, anti-estrogens (*e.g.* tamoxifen, raloxifene, and megestrol), LHRH agonists (*e.g.* goserelin and leuprolide), anti-androgens (*e.g.* flutamide and bicalutamide), photodynamic therapies (*e.g.* vertoporphin (BPD-MA), phthalocyanine, photosensitizer Pc4, and demethoxy-hypocrellin A (2BA-2-DMHA)), nitrogen mustards (*e.g.* cyclophosphamide, ifosfamide, trofosfamide, chlorambucil, estramustine, and melphalan), nitrosoureas (*e.g.* carmustine (BCNU) and lomustine (CCNU)), alkylsulphonates (*e.g.* busulfan and treosulfan), triazenes (*e.g.* dacarbazine, temozolomide), platinum containing compounds (*e.g.* cisplatin, carboplatin, oxaliplatin), vinca alkaloids (*e.g.* vincristine, vinblastine, vindesine, and vinorelbine), taxoids (*e.g.* paclitaxel or a paclitaxel equivalent such as nanoparticle albumin-bound paclitaxel (Abraxane), docosahexaenoic acid bound-paclitaxel (DHA-paclitaxel, Taxoprexin), polyglutamate bound-paclitaxel (PG-paclitaxel, paclitaxel poliglumex, CT-2103, XYOTAX), the tumor-activated prodrug (TAP) ANG1005 (Angiopep-2 bound to three molecules of paclitaxel), paclitaxel-EC-1 (paclitaxel bound to the erbB2-recognizing peptide EC-1), and glucose-conjugated paclitaxel, *e.g.*, 2'-paclitaxel methyl 2-glucopyranosyl succinate; docetaxel, taxol), epipodophyllins (*e.g.* etoposide, etoposide phosphate, teniposide, topotecan, 9-aminocamptothecin, camptothecin, irinotecan, crisnatol, mitomycin C), anti-metabolites, DHFR inhibitors (*e.g.* methotrexate, dichloromethotrexate, trimetrexate, edatrexate), IMP dehydrogenase inhibitors (*e.g.* mycophenolic acid, tiazofurin, ribavirin, and EICAR), ribonucleotide reductase inhibitors (*e.g.* hydroxyurea and deferoxamine), uracil analogs (*e.g.* 5-fluorouracil (5-FU), floxuridine, doxifluridine, irinotecan, tegafur-uracil, capecitabine), cytosine analogs (*e.g.* cytarabine (ara C), cytosine arabinoside, and fludarabine), purine analogs (*e.g.* mercaptopurine and Thioguanine), Vitamin D3 analogs (*e.g.* EB 1089, CB 1093, and KH 1060), isoprenylation inhibitors (*e.g.* lovastatin), dopaminergic neurotoxins (*e.g.* 1-methyl-4-phenylpyridinium ion), cell cycle inhibitors (*e.g.* staurosporine), actinomycin (*e.g.* actinomycin D, dactinomycin), bleomycin (*e.g.* bleomycin A2, bleomycin B2, peplomycin), anthracycline (*e.g.* daunorubicin, doxorubicin, pegylated liposomal doxorubicin, idarubicin, epirubicin, pirarubicin, zorubicin, mitoxantrone), MDR inhibitors (*e.g.* verapamil), Ca²⁺ ATPase inhibitors (*e.g.* thapsigargin), imatinib, thalidomide, lenalidomide, tyrosine kinase inhibitors (*e.g.*, axitinib (AG013736), bosutinib (SKI-606), cediranib (RECENTINTM, AZD2171), dasatinib (SPRYCEL®, BMS-354825), erlotinib (TARCEVA®), gefitinib (IRESSA®), imatinib (Gleevec®, CGP57148B, STI-571), lapatinib (TYKERB®, TYVERB®), lestaurtinib (CEP-701), neratinib (HKI-272), nilotinib (TASIGNA®), semaxanib (semaxinib, SU5416), sunitinib (SUTENT®, SU11248), toceranib

(PALLADIA®), vandetanib (ZACTIMA®, ZD6474), vatalanib (PTK787, PTK/ZK), trastuzumab (HERCEPTIN®), bevacizumab (AVASTIN®), rituximab (RITUXAN®), cetuximab (ERBITUX®), panitumumab (VECTIBIX®), ranibizumab (Lucentis®), nilotinib (TASIGNA®), sorafenib (NEXAVAR®), everolimus (AFINITOR®), alemtuzumab (CAMPATH®), gemtuzumab ozogamicin (MYLOTARG®), temsirolimus (TORISEL®), ENMD-2076, PCI-32765, AC220, dovitinib lactate (TKI258, CHIR-258), BIBW 2992 (TOVOK™), SGX523, PF-04217903, PF-02341066, PF-299804, BMS-777607, ABT-869, MP470, BIBF 1120 (VARGATEF®), AP24534, JNJ-26483327, MGCD265, DCC-2036, BMS-690154, CEP-11981, tivozanib (AV-951), OSI-930, MM-121, XL-184, XL-647, and/or XL228), proteasome inhibitors (*e.g.*, bortezomib (VELCADE)), mTOR inhibitors (*e.g.*, rapamycin, temsirolimus (CCI-779), everolimus (RAD-001), ridaforolimus, AP23573 (Ariad), AZD8055 (AstraZeneca), BEZ235 (Novartis), BGT226 (Novartis), XL765 (Sanofi Aventis), PF-4691502 (Pfizer), GDC0980 (Genetech), SF1126 (Semafoe) and OSI-027 (OSI)), oblimersen, gemcitabine, carminomycin, leucovorin, pemetrexed, cyclophosphamide, dacarbazine, procarbazine, prednisolone, dexamethasone, campathecin, plicamycin, asparaginase, aminopterin, methopterin, porfiromycin, melphalan, leurosidine, leurosine, chlorambucil, trabectedin, procarbazine, discodermolide, carminomycin,, aminopterin, and hexamethyl melamine.

EXAMPLES

[00114] In order that the invention described herein may be more fully understood, the following examples are set forth. These examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

Example 1. Production of MSK-777

[00115] MSK-777 production and purification process consists of following blocks: a) fermentation, b) resin wash and elution, c) concentration, d) RP-chromatography, e) extraction, f) concentration and g) crystallization.

[00116] MSK-777 is produced through fermentation of *Streptomyces* strain *Streptomyces violaceoruber*. Fermentation broth contains as product scavenger polymeric absorbent resin XAD7 (The current tradename is XAD7HP, was earlier XAD7). The XAD7 resin is a non-ionic aliphatic acrylic polymer, which traps the formed bacterial metabolites. This direct trapping increases the production rate (inhibiting the feed-back regulation) and prohibits the possible degradation of the already formed metabolites. After fermentation the resin is

separated from the fermentation broth with filtration. The resin is selectively washed with small amounts of organic solvent in water, followed by elution of the product with organic solvent or mixture of organic solvent and water.

[00117] As main metabolites of the fermentation Granaticin A (a biosynthetic precursor of the MSK777) and MSK-777 are detected. Additionally 5 - 10 minor granaticin group intermediates (side products, degradation products) and several other non-granaticin metabolites are detected. A typical fermentation profile analyzed from the scavenger resin extract consist of 40 - 60 % Granaticin A and 20 - 40 % MSK-777 (see Figure 11).

[00118] The polar impurities of MSK-777 on the scavenger resin are washed with (20 : 80) acetone - water mixture prior the elution of the desired product. The resin is extracted after the wash sequence with three aliquots of acetone -water (85 : 15) mixture. The combined eluate fractions are concentrated under vacuum to 20-30 % of the original volume. This is necessary to facilitate a fluent loading onto the following reverse preparation (RP) chromatographic system. The pH of the sample before chromatography is in the range of about 5.2 to 7.2. RP-chromatography is performed using Pharmaprep RP-18 (40 - 63) pm material (Merck). Before injecting the sample, the column is equilibrated with 35 % MeOH in water. The column is then loaded with (1 - 3) wt. % MSK-777 relative to stationary phase weight. Typical loading volumes are (2 - 4) times the column volume. After loading the column, chromatography is run with a 5-step gradient using increasing MeOH amounts (35/45/50 % MeOH) and descending pH (0/0.1 % AcH). Pure pooled MSK-777 fractions are combined and extracted with chloroform once using 1/5 of the volume. The chloroform phase is washed twice with 5 g/L NaHCO₃ and twice with 0.001 % HCl.

[00119] The chloroform phase is evaporated to dryness. Dry MeOH is added to solubilize the residue with an final concentration of MSK-777 of about 20- 50 g/L. The pH is adjusted with isopropanolic HCl to between 3.5 and 5.0. The formed suspension is filtered, washed quickly with n-pentane, and dried under vacuum for 24 h at room temperature.

Example 2. Preparation of Form A from Chloroform/Methanol (First Batch)

[00120] The MSK-777 produced from Example 1 was taken and further dried for 18 h at 45 °C under vacuum (<10 mbar). The batch had been crystallized from water-free chloroform/methanol. The obtained crystals appear as dark-red prisms under the microscope.

Example 3. Preparation of Form A from Methanol/Acetone (Second Batch)

[00121] A 5 g of MSK-777 produced from Example 1 was taken and dissolved in methanol - acetone (1:3; v:v). Dilute aqueous hydrochloric acid (pH = 4-5) was slowly added under vigorous mixing to the solution until water ratio of ca. 65 vol % was achieved. The final concentration of MSK-777 was about 4 -5 g/L. The mixture was stirred two hours at room temperature, followed by four hours at +4 °C and further overnight at 0 °C. Crystals were filtered and washed with 15 % methanol in water. Crystals were dried first in air, followed by vacuum at ambient temperature and finally at 45 °C and vacuum (<10 mbar). The obtained crystals appear as dark-red prisms.

*Example 4. Analysis and Characterization of MSK-777 before crystallization***4.1. UV-spectrum**

[00122] The UV spectrum of MSK-777 shows broad absorptions at 235, 280, 510 and 527 nm. These absorptions indicate a benzoisochromanequinones (BIQ) chromophore. The broad adsorption in the visible region (around 527 nm) is responsible for the intense red color of the substance.

[00123] Under basic conditions (pH > 8) a bathochromic shift of Vis-absorption (ca. 40 nm) is observed, indicating ionization of slightly acidic phenolic protons. This pH dependent indicatory behavior is characteristic to the BIQ chromophore and is visually observed as the solution turns from red to blue. The color change has an inflection point at ca. pH=8.5 corresponding to the approximate averaged pKa values of the acidic phenol protons.

4.2. Chemical degradation

[00124] MSK-777 shows a slow degradation under acidic conditions (pH < 4) at ambient temperature as observed by HPLC. The degradation is caused by hydrolysis of the glycosidic bond, forming Granaticin A, the biosynthetic precursor of MSK-777. Consistently, in a time dependent HPLC experiment a simultaneous increase of Granaticin A chromatographic areas and decrease of Granaticin B areas is observed. The hydrolysis experiment and the almost identical UV spectra indicate that MSK-777 (Granaticin B) and Granatioin A share the same core structure and chromophore. Therefore the structural difference between Granaticin A and MSK-777 arises from a hydrolysable part (sugar moiety) of the latter compound.

4.3. Low resolution mass spectroscopic analysis

[00125] Low resolution mass spectroscopic analysis was conducted using HPLC-MS in electrospray (API) modus. Both, negative and positive ionization modes were recorded simultaneously. Typical for phenolic compounds without a positive ionizable functional group, negative ionization provides a distinct mother ion peak ($[M-H]^-$) together with only a few weak degradation signals. The mother peak with mass of $M/z=558$ corresponds to the anticipated molecular form of MSK-777 ($C_{28}H_{30}O_{12}$). In contrast, positive ionization shows a complex degradation pattern with a molecular ion obtained by NH_4^+ adduct formation ($[M+NH_4]^+$) with a mass of 576. The cleavage of the sugar moiety (2-methyltetrahydropyran-3-ol, a loss of m/z 114) from MSK-777 (formation of Granaticin A) was detected in the positive ionization modus, giving a signal with a mass of 445. Similar, the Granaticin A mass spectrum shows the same molecular peak (mass 444, corresponding to the molecular formula of $C_{22}H_{20}O_{10}$) in line with a missing sugar moiety compared to MSK-777.

4.4. High resolution mass spectroscopic analysis

[00126] The high resolution mass spectroscopic analysis shows a molecular mass of 557.16736 ($[M-H]^-$). The difference to the theoretical calculated mass for $[M-H]^- = C_{28}H_{29}O_{12}$ is 1.63 ppm. Also the high resolution MS-MS degradation products (two major signals) give the correct anticipated masses. This result confirms theoretical elemental composition of $C_{28}H_{30}O_{12}$.

4.5. IR spectroscopic analysis

[00127] The FTIR spectrum of MSK-777 shows two strong signals at ca. 1800 and 1600 cm^{-1} due to carbonyl functions. Especially the signal at 1800 cm^{-1} is characteristic for quinone carbonyl system. The signal at 1600 cm^{-1} arises from the lactone carbonyl. The FTIR spectrum confirms the structure of MSK-777.

4.6. NMR

[00128] The data collected from a set of measurements (1D- ^1H , 1D-sel-TOCSY, 1D- ^{13}C , 2DCOSY, 2D-HMBC) confirms that the protons and carbons in MSK-777 were made matching the structure of Granaticin B.

*Example 5. Analysis and Characterization of Form A***5.1. Analytical data for two batches of Form A****Table 3.** Analytical Data for Crystalline MSK-777 Batches

Batch	Purity / %	Assay / %	Water / %	MeOH / ppm	DCM / ppm	Pentane / ppm
TK136_19_01	99,35	98,96	0,86	3484	<46	<54
TK136_19_03	99,41	99,97	0,97	671	<15	<54

5.2. XRD of Form A

[00129] In the X-ray powder diffraction analysis, both batches show sharp and well separated signals indicating crystallinity. Both batches show identical signal pattern, indicating that they share the same crystal structure. The peak positions and their intensities are tabulated in Table 4, below.

Table 4. XRPD Peak Positions and Intensities for Investigated MSK-777 Batches

Batch TK136_19_01		Batch TK136_19_03	
Intensity	2 θ	Intensity	2 θ
15	4,8	5	4,8
4,8	8,6	42	8,6
100	9,0	100	9,0
33	9,8	28	10,0
25	11,5	25	11,5
10	12,5	8	12,6
8	13,2	8	13,3
15	15,3	13	15,4
25	16,1	23	16,1
10	16,9	8	16,9
8	17,3	6	17,3
20	18,2	22	18,2
6	19,6	6	19,6
5	20,2	5	20,2
6	21,5	4	21,6
8	21,9	5	22,1
8	23,2	5	24,2

5.3. Thermal analysis of Form A

[00130] The DSC analysis shows for both samples a weak endothermic transition peak at 160 °C followed by a melting / decomposition point at ca. 190°C. The TG analysis shows an approx. 2 % decrease in mass in the temperature interval up to the temperature of 100 °C.

This mass decrease is related to evaporation of water and residual solvent from the crystals. No indication of hydrate(s) is seen (no stepwise reduction of the mass below the melting point). Beyond approx. 185 °C a rapid loss of the mass is detected indicating degradation of the product.

5.4. Solid Phase NMR analysis of Form A

[00131] Both batches show the same solid phase ^{13}C NMR spectra. The spectra appear to indicate an amorphous material instead of the expected crystalline phase; broad unresolved signals instead of well separated sharp signals. This result is explained by the observation that during the NMR sample pretreatment consisting of compacting the sample by pressure the initially crystalline material partially melted (i.e. became amorphous).

EQUIVALENTS AND SCOPE

[00132] In the claims articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[00133] Furthermore, the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims is introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, *e.g.*, in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements and/or features, certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth in haec verba herein. It is also noted that the terms “comprising” and “containing” are intended to be open and permits the inclusion of

additional elements or steps. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.


[00134] This application refers to various issued patents, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. If there is a conflict between any of the incorporated references and the instant specification, the specification shall control. In addition, any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Because such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the invention can be excluded from any claim, for any reason, whether or not related to the existence of prior art.

[00135] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments described herein. The scope of the present embodiments described herein is not intended to be limited to the above Description, but rather is as set forth in the appended claims. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present invention, as defined in the following claims.

CLAIMS

What is claimed is:

1. Crystalline Form A of Compound **1**:



2. The crystalline Form A of claim 1 characterized by an XRPD pattern substantially similar to the one depicted in *Figure 1*.
3. The crystalline Form A of claim 1 characterized by an XRPD pattern substantially similar to the one depicted in *Figure 2*.
4. The crystalline Form A of any one of preceding claims characterized by having four or more peaks in its XRPD pattern selected from those in the following table:

Angle (2-Theta °)
9.1 ± 0.2
10,0 ± 0.2
11.6 ± 0.2
16.1 ± 0.2

5. The crystalline Form A of any one of preceding claims characterized by having five or more peaks in its XRPD pattern selected from those in the following table:

Angle (2-Theta °)
9.1 ± 0.2
10,0 ± 0.2
11.6 ± 0.2
16.1 ± 0.2
18.3 ± 0.2

6. The crystalline Form A of any one of preceding claims characterized by having seven or more peaks in its XRPD pattern selected from those in the following table:

Angle (2-Theta °)
9.1 ± 0.2
10,0 ± 0.2
11.6 ± 0.2
16.1 ± 0.2
18.3 ± 0.2
4.5 ± 0.2
15.4 ± 0.2

7. The crystalline Form A of any one of preceding claims characterized by having nine or more peaks in its XRPD pattern selected from those in the following table:

Angle (2-Theta °)
9.1 ± 0.2
10,0 ± 0.2
11.6 ± 0.2
16.1 ± 0.2
18.3 ± 0.2
4.5 ± 0.2
15.4 ± 0.2
12,6 ± 0.2
17,0 ± 0.2

8. The crystalline Form A of any one of preceding claims characterized by having thirteen peaks in its XRPD pattern selected from those in the following table:

Angle (2-Theta °)
9.1 ± 0.2
10,0 ± 0.2
11.6 ± 0.2
16.1 ± 0.2
18.3 ± 0.2

Angle (2-Theta °)
4.5 ± 0.2
15.4 ± 0.2
12.6 ± 0.2
17,0 ± 0.2
13.3 ± 0.2
17.5 ± 0.2
21.6 ± 0.2
23.3 ± 0.2

9. The crystalline Form A of any one of preceding claims characterized by having seventeen peaks in its XRPD pattern selected from those in the following table:

Angle (2-Theta °)
9.1 ± 0.2
10,0 ± 0.2
11.6 ± 0.2
16.1 ± 0.2
18.3 ± 0.2
4.5 ± 0.2
15.4 ± 0.2
12.6 ± 0.2
17,0 ± 0.2
13.3 ± 0.2
17.5 ± 0.2
22,1 ± 0.2
23.3 ± 0.2
8.7 ± 0.2
19.7 ± 0.2
20.3 ± 0.2
21.6 ± 0.2

10. The crystalline Form A of any one of preceding claims characterized by a DSC endothermogram substantially similar to the one depicted in *Figure 3*.

11. The crystalline Form A of any one of claims 1-9 characterized by a DSC endothermogram substantially similar to the one depicted in *Figure 4*.

12. The crystalline Form A of any one of preceding claims characterized by a DSC endothermogram with a peak temperature ($T_{\max}$) of about 190 °C.

13. The crystalline Form A of any one of preceding claims characterized by a DSC endothermogram with phase transition in the range of about 140° C to about 250 °C.
14. The crystalline Form A of any one of claims 1-12 characterized by a DSC endothermogram with phase transition in the range of about 155 °C to about 225 °C.
15. The crystalline Form A of any one of preceding claims having an observed decomposition point of about 190 °C.
16. The crystalline Form A of any one of preceding claims characterized by a thermogravimetric spectrum substantially similar to the one depicted in *Figure 5*.
17. The crystalline Form A of any one of claims 1-15 characterized by a thermogravimetric spectrum substantially similar to the one depicted in *Figure 6*.
18. The crystalline Form A of any one of preceding claims characterized by an NMR spectrum substantially similar to the one depicted in *Figure 7*.
19. The crystalline Form A of any one of claims 1-17 characterized by an NMR spectrum substantially similar to the one depicted in *Figure 8*.
20. The crystalline Form A of any one of preceding claims, wherein the crystalline Form A is obtained from chloroform and methanol.
21. The crystalline Form A of claim 20, wherein the chloroform is free of water.
22. The crystalline Form A of any one of claims 1-19, wherein the crystalline Form A is obtained from methanol and acetone.
23. The crystalline Form A of any one of preceding claims, wherein the crystalline Form A is not a hydrate.

24. The crystalline Form A of any one of preceding claims, wherein the crystalline Form A is substantially free of amorphous Compound **1**.
25. The crystalline Form A of any one of preceding claims, wherein about 0.01 wt% to about 10 wt% of amorphous Compound **1** is present.
26. The crystalline Form A of any one of preceding claims, wherein about 0.01 wt% to about 5.0 wt% of water is present.
27. The crystalline Form A of any one of preceding claims, wherein the crystalline Form A is substantially free of impurities.
28. A method of preparing the crystalline Form A of claim 1 comprising crystalizing Compound **1** from chloroform and methanol.
29. The method of claim 28 wherein the chloroform is water-free.
30. A method of preparing the crystalline Form A of claim 1 comprising mixing Compound **1** with methanol and acetone to generate a mixture.
31. The method of claim 30 further comprising adding aqueous hydrochloride to acidify the mixture.
32. The method of claim 31, wherein about 65 vol% of water is present in the acidified mixture.
33. The method of any one of claims 30-32, wherein the concentration of Compound **1** is about 3-6 g/L.
34. The method of claim 33, wherein the concentration of Compound **1** is about 4-5 g/L.
35. The method of any one of claims 30-34 further comprising lowering the temperature of the mixture to provide a solid.

36. The method of claim 35 further comprising isolating the solid from the mixture.
37. A pharmaceutical composition comprising a crystalline Form A of any one of claims 1-27 and optionally a pharmaceutically acceptable excipient.
38. The pharmaceutical composition of claim 36, wherein the composition comprises a therapeutically effective amount of the crystalline Form A.
39. A method for treating a proliferative disease comprising administering to a subject in need thereof a therapeutically effective amount of crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38.
40. The method of claim 39, wherein the proliferative disease is cancer.
41. The method of claim 40, wherein the cancer is leukemia, thyroid cancer, melanoma, ovarian cancer, lung cancer, prostate cancer, renal cell carcinoma, cervical cancer, or breast cancer.
42. The method of claim 41, wherein the leukemia is acute myeloid leukemia, chronic lymphocytic leukemia, acute lymphoblastic leukemia, or biphenotypic acute leukemia.
43. A method of inhibiting cell growth in a subject, the method comprising administering to the subject with a therapeutically effective amount of crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38.
44. A method of inhibiting cell growth in a biological sample, the method comprising contacting the biological sample with a therapeutically effective amount of crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38.
45. A method of inducing apoptosis of a cell in a subject, the method comprising: administering to the subject with a therapeutically effective amount of crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38.
46. A method of inducing apoptosis of a cell in a biological sample, the method comprising contacting the biological sample with a therapeutically effective amount of

crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38.

47. The method of any one of claims 39-46, further comprising administering to the subject or biological sample a therapeutically effective amount of a second therapeutic agent, the pharmaceutically acceptable salt thereof, or the pharmaceutical composition thereof.

48. The method of claim 47, wherein the second therapeutic agent is an anti-proliferative agent.

49. A kit comprising the crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38; and instructions for administering the crystalline Form A or the pharmaceutical composition.

50. A method of treating a microbial infection in a subject comprising administering an effective amount of the crystalline Form A of any one of claims 1-27, or a pharmaceutical composition of claim 37 or 38 to the subject.

51. The method of claim 50, wherein the microbial infection is a bacterial infection.

52. The method of claim 51, wherein the bacterial infection is caused by a gram positive bacterium.

53. The method of claim 52, wherein the gram positive bacterium is of the genus *Staphylococcus*, *Streptococcus*, *Micrococcus*, *Peptococcus*, *Peptostreptococcus*, *Enterococcus*, *Bacillus*, *Clostridium*, *Lactobacillus*, *Listeria*, *Erysipelothrix*, *Propionibacterium*, *Eubacterium*, or *Corynebacterium*.

54. The method of any one of claims 49-53, wherein the subject is a human.




Figure 1




Figure 2

Datei: C:\...\DSC\TAI_DSC_Messaten\DSa5560.001
Bediener: gs
Versuchsdatum: 10-Apr-2012 12:04
Gerät: DSC Q2000 V24.4 Build 116

Probe: MSK-777 TK136_19_01
Größe: 1,7960 mg
DSC




Figure 3

Datei: C:\DSC\TAI_DSC_Messaten\DSa5561.001
Bediener: gs
Versuchsdatum: 10-Apr-2012 12:56
Gerät: DSC Q2000 V24.4 Build 116

Probe: MSK-777 TK136_19_03
Größe: 2,0290 mg
DSC






Figure 4



f43101	2012-04-11 08:15	Hauptansicht	Gerät: NETZSCH TG 209 Datei: TGa3885.dt2 Projekt: Kunnari Versuchs-ID: 07 Datum/Zeit: 10.04.2012 16:46:17 Labor: HERAEUS Operator:		Probe: MSK-777 TK136_19_01 Material: pharmabac.bt2 Korrektur-Datei: 091211.tt2 Tempkal. Datei: 23.0/10.0(K/min)/300.0 Bereich: TG 209 standard/K Probenr./TC: 12.634 mg Probenmasse:	Modus/Messart.: TG/Probe + Korrektur Segmente: 1/1 Tiegel: Al2O3 Atmosphäre: ----/---- / Ar/10 / Ar/---- Korr./Bereich: 020/500 mg Zyklen vor Messung: 0xVak	TGa3885.ngb
Dateipfad: D:\ngbwin\ta\data5\TGa3885.dt2			Bemerkung:				

Figure 5



f43101 2012-04-11 08:15 Hauptansicht			
Gerät:	NETZSCH TG 209	Probe:	MSK-777 TK136_19_03
Datei:	TGa3886.dt2	Material:	
Projekt:	Kunnari	Korrektur-Datei:	pharmabac.bt2
Versuchs-ID:	08	Tempkal. Datei:	091211.tt2
Datum/Zeit:	10.04.2012 17:33:42	Bereich:	23.0/10.0(K/min)/300.0
Labor:	HERAEUS	Probenr./TC:	TG 209 standard/K
Operator:		Probenmasse:	10.194 mg
Dateipfad: D:\ngbw\in\ta\data5\TGa3886.dt2		Bemerkung:	
Modus/Messart:		TG/Probe + Korrektur	
Segmente:		1/1	
Tiegel:		Al2O3	
Atmosphäre:		---/--- / Ar/10 / Ar/---	
Korr./Bereich:		020/500 mg	
Zyklen vor Messung:		0xVak	

Figure 6

7/17




Figure 7

8/17

```

NAME          4apr112.006
EXPNO          10
PROCNO         1
Date_          20120425
Time_          13.13
INSTRUM  AQura_GerNr6329
PROBHD    4 mm MAS BB/1H
PULPROG          cpramp
TD           4096
SOLVENT          CD2C12
NS           5067
DS            0
SWH          45045.047 Hz
FIDRES        10.997326 Hz
AQ           0.0455156 sec
RG           16384
DW           11.100 usec
DE           20.00 usec
TE           297.5 K
D1           15.00000000 sec

```

```

----- CHANNEL f1 -----
NUC1           13C
P15           2000.00 usec
PL1            4.50 dB
SF01          100.6228303 MHz

```

```

----- CHANNEL f2 -----
CPDPRG2        tppm15
NUC2            1H
P3             4.40 usec
P31            6.80 usec
PL2            -2.00 dB
PL12           -4.00 dB
SF02          400.1316005 MHz
SPNAM0         ramp.64
SPOAL0         0.500
SPOFFS0        0.00 Hz
SI             16384
SF            100.6129183 MHz
WDW            EM
SSB            0
LB             50.00 Hz
GB            0
PC             1.00
SR            149.28 Hz

```

Figure 7
(Continued)

9/17




Figure 8

10/17

```

NAME          4apr112.005
EXPNO          10
PROCNO         1
Date_          20120423
Time_          12.12
INSTRUM  AQura_GerNr6329
PROBHD    4 mm MAS BB/1H
PULPROG          cpramp
TD          4096
SOLVENT          CD2C12
NS          4780
DS           0
SWH          45045.047 Hz
FIDRES        10.997326 Hz
AQ          0.0455156 sec
RG          16384
DW          11.100 usec
DE          20.00 usec
TE          297.1 K
D1          15.00000000 sec

----- CHANNEL f1 -----
NUC1          13C
P15          2000.00 usec
PL1          4.50 dB
SFO1          100.6228303 MHz

----- CHANNEL f2 -----
CPDPRG2        tppm15
NUC2           1H
P3           4.40 usec
P31          6.80 usec
PL2          -2.00 dB
PL12         -4.00 dB
SF02          400.1316005 MHz
SPNAM0        ramp.64
SPOAL0         0.500
SPOFFS0        0.00 Hz
SI           16384
SF           100.6129183 MHz
WDW           EM
SSB           0
LB           50.00 Hz
GB           0
PC           1.00
SR           149.28 Hz

```

Figure 8
(continued)




Figure 9

12/17




Figure 10




Figure 11

14/17

Version : PKS_2.01

Title : MSK777 TK136_19_01 gemörsert

Diffractometer : Transmission
 Monochromator : Curved Germanium (111)
 Wavelength : 1.540598 Cu
 Detector : Image Plate
 Scan Mode : Transmission / Stationary PSD / Fixed omega
 Scan Type : 2Theta

! Raw data file used : E:\Verschiedene\V4560.rmb
 ! created : 04-Apr-12 11:56

! Peak search parameters : Expected halfwidth : 0.150
 ! Significance level : 10.0
 ! Peak height level : 500

Peaklist [Range 1 : 2Theta = 3.000 79.051 0.015 Imax = 37226]

D	2Theta	I(rel)	I(abs)	I(int)	FWHM	H	K	L
17.834480	4.9509	13.97	5124	0.00	0.1800			
10.203506	8.6592	51.74	18971	0.00	0.2100			
9.718068	9.0926	100.00	36665	0.00	0.2250			
8.878934	9.9540	33.26	12194	0.00	0.1500			
7.675597	11.5194	26.33	9654	0.00	0.2250			
7.035887	12.5708	13.00	4766	0.00	0.1800			
6.707521	13.1889	10.60	3887	0.00	0.1500			
5.804516	15.2521	15.85	5813	0.00	0.2250			
5.522740	16.0352	24.90	9131	0.00	0.1800			
5.259232	16.8444	13.89	5092	0.00	0.2100			
5.111773	17.3340	11.20	4107	0.00	0.2100			
4.884393	18.1476	17.20	6305	0.00	0.3300			
4.517023	19.6376	8.79	3223	0.00	0.2250			
4.384967	20.2351	4.30	1577	0.00	0.1800			
4.129109	21.5034	7.82	2869	0.00	0.2700			
4.055562	21.8981	12.54	4599	0.00	0.1800			
3.845650	23.1095	10.76	3946	0.00	0.2700			
3.747940	23.7206	4.63	1698	0.00	0.1500			
3.659094	24.3052	3.49	1281	0.00	0.1950			
3.534577	25.1752	1.53	561	0.00	0.1050			
3.345694	26.6219	3.79	1389	0.00	0.1200			
2.304619	39.0528	1.69	621	0.00	0.1650			

Figure 12A

15/17

Title : MSK777 TK136_19_01 Kristalle

Diffractometer : Transmission
 Monochromator : Curved Germanium (111)
 Wavelength : 1.540598 Cu
 Detector : Image Plate
 Scan Mode : Transmission / Stationary PSD / Fixed omega
 Scan Type : 2Theta

! Raw data file used : E:\Verschiedene\V4560_2.rmb
 ! created : 04-Apr-12 12:23

! Peak search parameters : Expected halfwidth : 0.150
 ! Significance level : 10.0
 ! Peak height level : 500

Peaklist [Range 1 : 2Theta = 3.000 79.051 0.015 Imax = 62198]

!	D	2Theta	I(rel)	I(abs)	I(int)	FWHM	H	K	L
	17.893715	4.9345	12.47	7705	0.00	0.1800			
	12.458025	7.0899	2.53	1561	0.00	0.3000			
	10.205769	8.6572	38.64	23866	0.00	0.2100			
	9.717583	9.0931	100.00	61769	0.00	0.2250			
	8.871635	9.9622	25.63	15832	0.00	0.1500			
	7.674469	11.5211	23.72	14654	0.00	0.2400			
	7.035125	12.5722	11.56	7141	0.00	0.1950			
	6.715550	13.1731	8.97	5543	0.00	0.2100			
	5.801803	15.2593	11.81	7297	0.00	0.2700			
	5.520470	16.0419	14.48	8945	0.00	0.1800			
	5.256653	16.8527	11.77	7269	0.00	0.2250			
	5.116567	17.3176	8.91	5506	0.00	0.1650			
	4.867994	18.2092	14.98	9251	0.00	0.2700			
	4.515440	19.6445	6.48	4004	0.00	0.2400			
	4.381221	20.2526	5.58	3445	0.00	0.2100			
	4.223195	21.0188	2.19	1355	0.00	0.1200			
	4.128156	21.5084	8.10	5004	0.00	0.1800			
	4.057230	21.8890	9.41	5812	0.00	0.2100			
	3.849586	23.0855	7.54	4657	0.00	0.3300			
	3.743406	23.7497	3.41	2107	0.00	0.1200			
	3.657998	24.3126	3.19	1971	0.00	0.1800			
	3.341307	26.6575	3.37	2080	0.00	0.1350			
	3.077495	28.9906	1.49	918	0.00	0.1200			

Figure 12B

16/17

Title : MSK777 TK136_19_03 gemörsert

Diffractometer : Transmission
 Monochromator : Curved Germanium (111)
 Wavelength : 1.540598 Cu
 Detector : Image Plate
 Scan Mode : Transmission / Stationary PSD / Fixed omega
 Scan Type : 2Theta

! Raw data file used : E:\Verschiedene\V4561.rmb
 ! created : 04-Apr-12 14:26

! Peak search parameters : Expected halfwidth : 0.150
 ! Significance level : 10.0
 ! Peak height level : 500

Peaklist [Range 1 : 2Theta = 3.000 79.051 0.015 Imax = 54845]

!	D	2Theta	I(rel)	I(abs)	I(int)	FWHM	H	K	L
	17.643456	5.0046	5.14	2790	0.00	0.1950			
	10.858358	8.1360	11.94	6488	0.00	0.3300			
	10.116099	8.7341	42.72	23211	0.00	0.1350			
	9.714836	9.0956	100.00	54338	0.00	0.1650			
	8.787724	10.0576	26.21	14244	0.00	0.1800			
	8.400547	10.5224	2.66	1443	0.00	0.0900			
	7.650923	11.5567	25.79	14013	0.00	0.1800			
	7.008681	12.6198	9.68	5260	0.00	0.1800			
	6.636741	13.3302	9.73	5288	0.00	0.1500			
	5.771197	15.3407	13.39	7278	0.00	0.1950			
	5.521335	16.0394	22.51	12229	0.00	0.1650			
	5.225974	16.9524	11.86	6446	0.00	0.1950			
	5.068121	17.4844	8.55	4644	0.00	0.1500			
	4.859704	18.2405	20.58	11181	0.00	0.2250			
	4.622404	19.1856	2.14	1160	0.00	0.2100			
	4.495121	19.7342	8.15	4426	0.00	0.1650			
	4.374690	20.2831	5.93	3221	0.00	0.1800			
	4.114008	21.5833	3.71	2017	0.00	0.1500			
	4.028483	22.0472	7.78	4226	0.00	0.1650			
	3.809758	23.3302	8.11	4409	0.00	0.1800			
	3.733845	23.8114	2.58	1400	0.00	0.1500			
	3.639885	24.4355	2.80	1520	0.00	0.1950			
	3.377888	26.3636	2.66	1444	0.00	0.2400			
	3.321714	26.8177	2.29	1245	0.00	0.1950			
	3.193658	27.9143	2.16	1173	0.00	0.1800			
	3.081099	28.9560	1.54	839	0.00	0.1350			
	2.882843	30.9956	1.98	1074	0.00	0.1350			
	2.306442	39.0207	2.23	1210	0.00	0.0750			

Figure 13A

17/17

Title : MSK777 TK136_19_03 Kristalle

Diffractometer : Transmission
 Monochromator : Curved Germanium (111)
 Wavelength : 1.540598 Cu
 Detector : Image Plate
 Scan Mode : Transmission / Stationary PSD / Fixed omega
 Scan Type : 2Theta



! Raw data file used : E:\Verschiedene\V4561_2.rmb
 ! created : 04-Apr-12 13:12

! Peak search parameters : Expected halfwidth : 0.150
 ! Significance level : 10.0
 ! Peak height level : 500



Peaklist [Range 1 : 2Theta = 3.000 79.051 0.015 Imax = 53112]

!	D	2Theta	I(rel)	I(abs)	I(int)	FWHM	H	K	L
	19.461641	4.5368	2.13	1125	0.00	0.1200			
	17.644384	5.0043	4.67	2470	0.00	0.2100			
	11.211878	7.8791	8.85	4682	0.00	0.3300			
	10.102257	8.7461	46.18	24423	0.00	0.1200			
	9.718712	9.0920	100.00	52891	0.00	0.1800			
	8.799019	10.0446	26.87	14214	0.00	0.2100			
	7.639626	11.5739	31.61	16717	0.00	0.1650			
	7.000416	12.6348	12.49	6607	0.00	0.1500			
	6.647510	13.3085	6.85	3625	0.00	0.2100			
	5.987799	14.7826	2.07	1095	0.00	0.0900			
	5.763396	15.3616	12.40	6558	0.00	0.1800			
	5.517025	16.0520	18.58	9830	0.00	0.1500			
	5.217324	16.9807	11.19	5919	0.00	0.2550			
	5.067416	17.4869	7.01	3707	0.00	0.1800			
	4.855331	18.2571	24.83	13134	0.00	0.2100			
	4.487427	19.7684	9.05	4784	0.00	0.1650			
	4.370926	20.3008	8.96	4741	0.00	0.2100			
	4.118873	21.5575	5.62	2971	0.00	0.1500			
	4.026230	22.0597	7.56	3999	0.00	0.1950			
	3.807685	23.3431	6.30	3332	0.00	0.1500			
	3.729910	23.8369	2.81	1486	0.00	0.1500			
	3.635557	24.4650	2.94	1553	0.00	0.1650			
	3.497540	25.4463	1.79	946	0.00	0.1200			
	3.357976	26.5228	2.72	1440	0.00	0.3300			
	3.149010	28.3183	1.67	886	0.00	0.1650			
	3.033001	29.4254	2.54	1343	0.00	0.1350			
	2.879860	31.0285	1.92	1016	0.00	0.2100			
	2.661836	33.6424	1.52	803	0.00	0.1950			

Figure 13B



(1) Granaticin B



(2) Granaticin A

Figure 10

