




(86) Date de dépôt PCT/PCT Filing Date: 1997/10/24
 (87) Date publication PCT/PCT Publication Date: 1998/05/07
 (45) Date de délivrance/Issue Date: 2010/05/11
 (85) Entrée phase nationale/National Entry: 1999/04/20
 (86) N° demande PCT/PCT Application No.: EP 1997/005900
 (87) N° publication PCT/PCT Publication No.: 1998/018940
 (30) Priorité/Priority: 1996/10/25 (DE19644478.0)

(51) Cl.Int./Int.Cl. *C12N 15/55* (2006.01),
A01H 5/00 (2006.01), *C12N 1/21* (2006.01),
C12N 15/82 (2006.01)
 (72) Inventeurs/Inventors:
SONNEWALD, UWE, DE;
EBNETH, MARCUS, DE;
SCHMIDT, RALF-MICHAEL, DE
 (73) Propriétaire/Owner:
BASF AKTIENGESELLSCHAFT, DE
 (74) Agent: ROBIC

(54) Titre : EXPRESSION DE GENES SPECIFIQUES AUX FEUILLES DANS DES PLANTES TRANSGENIQUES
 (54) Title: LEAF-SPECIFIC GENE EXPRESSION IN TRANSGENETIC PLANTS



(57) Abrégé/Abstract:

Promoters are disclosed which cause the permanent, leaf-specific expression of a coding nucleotide sequence controlled by said promoters, for example a sequence which confers a specific resistance or increases the photosynthesis capacity.




PCT
 WELTORGANISATION FÜR GEISTIGES EIGENTUM
 Internationales Büro
 INTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
 INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)

<p>(51) Internationale Patentklassifikation⁶ : C12N 15/55, 15/82, 1/21, A01H 5/00</p>	<p>A1</p>	<p>(11) Internationale Veröffentlichungsnummer: WO 98/18940 (43) Internationales Veröffentlichungsdatum: 7. Mai 1998 (07.05.98)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP97/05900 (22) Internationales Anmeldedatum: 24. Oktober 1997 (24.10.97) (30) Prioritätsdaten: 196 44 478.0 25. Oktober 1996 (25.10.96) DE (71) Anmelder (für alle Bestimmungsstaaten ausser US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE). (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): SONNEWALD, Uwe [DE/DE]; Martha-Brauttsch-Strasse 7a, D-06467 Hoym (DE). EBNETH, Marcus [DE/DE]; Bicklingerweg 16, D-06484 Quedlinburg (DE). SCHMIDT, Ralf-Michael [DE/DE]; Gräfensteinstrasse 14, D-67434 Neustadt (DE). (74) Anwälte: KINZEBACH, Werner usw.; Reitstötter, Kinzebach & Partner, Sternwartstrasse 4, D-81679 München (DE).</p>		<p>(81) Bestimmungsstaaten: AU, BR, CA, IL, JP, US, europäisches Patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Veröffentlicht <i>Mit internationalem Recherchenbericht. Vor Ablauf der für Änderungen der Ansprüche zugelassenen Frist. Veröffentlichung wird wiederholt falls Änderungen eintreffen.</i></p>

(54) Title: **LEAF-SPECIFIC GENE EXPRESSION IN TRANSGENETIC PLANTS**

(54) Bezeichnung: **BLATTSPEZIFISCHE EXPRESSION VON GENEN IN TRANSGENEN PFLANZEN**



(57) Abstract

Promoters are disclosed which cause the permanent, leaf-specific expression of a coding nucleotide sequence controlled by said promoters, for example a sequence which confers a specific resistance or increases the photosynthesis capacity.

LEAF-SPECIFIC GENE EXPRESSION IN TRANSGENETIC PLANTS

The present invention relates to nucleic acid sequences having promoter activity, which cause a leaf-specific expression, in plants, of encoding nucleotide sequences under the control of said promoters, to expression cassettes, recombinant vectors and microorganisms which embrace such regulatory sequences, to transgenic plants transformed with them, to a method of producing transgenic plants, and to a method of isolating the leaf-specific promoter.

10 It is known to transfer foreign genes into the genome of a plant in a targeted manner by means of genetic engineering methods. This process is termed transformation, and the resulting plants are termed transgenic. Transgenic plants are currently employed in various fields of biotechnology. The predominant aims are, on the one hand, crop protection and, on the other hand, an improved quality of the harvestable products. In order to express foreign genes efficiently in plants, regulation signals which allow ordered transcription are required. These include promoters and terminators. The terminators located on the 3' end of the encoding DNA serve to end transcription and, if appropriate, as a signal for polyadenylation of the mRNA formed. Promoters contain recognition sequences for RNA-polymerases and for transcriptional effectors. The promoters are responsible for the expression behavior of the foreign genes.

20

Herbicide-tolerant plants as they are disclosed in DE-A-3701623 are an example of genetic engineering measures in crop protection. Other examples are insect-resistant plants (Vaek et al. (1987) Plant Cell 5, 159-169), virus-resistant plants (Powell et al. (1986) Science 232, 738-743) and ozone-resistant plants (Van Camp et al. (1994) BioTech. 12, 165-168). Examples of quality improvements achieved by genetic engineering are: better keeping qualities in fruit (Oeller et al. (1991) Science 254, 437-439), increased starch production in potato tubers (Stark et al. (1992) Science 242, 419), altered starch (Visser et al. (1991) Mol. Gen. Genet. 225, 289-296) and lipid composition (Voelker et al. (1992) Science 257, 72-74) and production of plant-foreign polymers (Poirer et al. (1992) Science 256, 30 520-523).

A large number of promoters which control expression of foreign genes in plants is known. The most frequently used promoter is the 35S CaMV promoter (Franck et al. (1980) Cell 21, 285-294).

This promoter contains various recognition sequences for transcriptional effectors which, in their totality, lead to constitutive expression of the gene which has been introduced (Benfey et al. (1989) EMBO J. 8, 2195-2202). Frequently, 5 inducible or cell- or tissue-specific promoters are also employed.

Examples of inducible expression which have been described are, inter alia, the following: wound induction (DE-A-3843628, 10 DE-B-3837752), chemical induction (Ward et al. (1993) Plant Molec. Biol. 22, 361-366) and light induction (Fluhr et al. (1986) Science 232, 1106-1112).

DE-A-4207358 discloses a promoter which causes gene expression 15 specific to the stomatic cells, but no specific expression in mesophyll cells or in epidermal cells of leaves. An artificial change of the opening periods of the stomata allows the gas exchange of plants manipulated in this fashion to be regulated as 20 desired. Herbicide tolerance or herbicide resistance can not be mediated by such a promoter.

Other examples of cell- and tissue-specific expression are: seed-, tuber- and fruit-specific expression (compiled in Edwards 25 and Coruzzi (1990) Annu. Rev. Genet. 24, 275-303; DE-A-3843627), phloem-specific expression (Schmülling et al. (1989) Plant Cell 1, 665-670), root-nodule-specific expression (DE-A-3702497) and meristem-specific expression (Ito et al. (1994) Plant Molec. Biol. 24, 863-878). Examples of promoters in 30 chloroplast-containing cells are also known from Edwards and Coruzzi (1990), Annu. Rev. Genet. 24, 277-279. The promoters described in this publication cause expression either only in inducible form (for example the rbcS-3A promoter) or only in certain types of cells (for example the GS2 and GS3A promoters), 35 but the expression is not limited to certain parts of the plant.

The use of the above-described promoters is frequently problematic. For example, promoters which cause constitutive 40 expression of the genes under their control can be employed for generating herbicide-tolerant and pathogen-resistant plants, but have the disadvantage that the products of the genes under their control exist in all parts of the plant, including the parts of the plant which are harvested, which may be undesirable in some cases. Equally, inducible promoters are not without problems 45 since the conditions for induction in crop plants are typically difficult to control in the field.

In C3 plants, the promoter of phosphoenol pyruvate carboxylase, which originates from a C4 plant, leads to expression in the mesophyll of the leaf (Stockhaus et al., (1994), Mol. Gen. Genet. 245, 286-293). However, this promoter only mediates a low activity level in the mesophyll. Moreover, it also shows activity in roots. This poor organ specificity is undesired for many applications.

Promoters which cause leaf-specific, preferably permanent, expression of genes under their control have not been disclosed.

It would therefore be desirable to find routes to express genes in plants while avoiding the above disadvantages.

It is an object of the present invention to provide means which allow a targeted, organ-specific gene expression in plants. These means should, for example, be suitable for the expression of resistance genes and genes which modify the photosynthesis rate.

We have found that this object is achieved, surprisingly, by providing a new promoter which causes a preferentially permanent, leaf-specific expression, in plants, of an encoding nucleotide sequence under the control of said promoter independently of induction factors.

The invention is now illustrated in greater detail with reference to the figures which follow.

Figure 1 shows (A) a schematic representation of the BamHI fragment of potato clone FBP-1 which has been cloned into vector pUC19 and comprises approx. 7100 bp. The cytosolic FBPase(cy-FBPase) promoter region is shown in black; (B) the construction scheme of plasmid FBP:pBlue;

Figure 2 shows the nucleotide sequence of the potato cy-FBPase promoter. The region which is complementary to the 3' end of the 5' subfragment, of the cy-FBPase, used for the Southern hybridization is underlined; two palindromic sequence portions are underlined by a dotted line; the 5'-terminal sequence "GGATC" was added to the genomic DNA to produce a BamHI cleavage site;

Figure 3 shows (A) the construction scheme of plasmids FBP:GUS and FBP:GUS (DEL); (B) a schematic representation of plasmid FBP:GUS with the FBPase promoter comprising approx. 1700

bp, the GUS gene comprising approx. 1870 bp and the nopaline synthase terminator comprising approx. 260 bp, inserted into vector pBI 101;

5 Figure 4 shows a bar chart which illustrates the cy-FBPase-promoter-controlled leaf-specific GUS activity in transgenic potato plants; it shows the results obtained for two different transformation experiments with FBP:GUS (plant line "Me 1-22" and "Me 1-9"), comparing them with a control experiment ("control");

10 Figure 5 shows a bar chart which confirms the cy-FBPase-promoter-controlled, leaf-specific GUS activity in transformed tobacco plants. "TME-1/67" refers to the results obtained with vector FBP:GUS. "TME-11/13" refers to the results obtained with vector FBP:GUS (DEL). "WT" shows the results obtained for the wild type. The data shown are the amount of 4-methylumbelliferone formed, per milligram of protein per minute;

15 Figure 6 shows the histochemical detection of GUS activity in various tissues of the tobacco leaf in a transgenic plant. (A) Cross-section of the central vascular bundle of the source leaf, (B) epidermis, (C) cross-section of the petiole, (D) cross-section of the mesophyll of a source leaf. The sections were fixed for 20 minutes in 3% strength paraformaldehyde solution and subsequently incubated overnight in X-Gluc solution. Then, the chlorophyll was removed using 70% ethanol;

20 Figure 7 shows a Northern blot which illustrates the uniform leaf-specific GUS expression in transgenic tobacco plants, mediated by the potato cy-FBPase promoter;

25 Figure 8 shows a histochemical proof of the β -glucuronidase (GUS) activity in tobacco seedlings;

30 Figure 9 shows a schematic representation of plasmid pBin-FBP with the potato cy-FBPase promoter which comprises 1724 bp and the octopine synthase terminator which comprises 280 bp, inserted into vector pBin 19, and

35 Figure 10 shows a cDNA probe of the potato cy-FBPase gene (EMBL No.: X76946):

40

The term "gene" or "encoding (nucleotide) sequence" denotes for the purposes of the present invention a nucleotide sequence which encodes a specific, if appropriate hereditary, structure, for example at least one protein, at least one ribozyme or at least one antisense RNA; or function, for example resistance; or an altered profile of the plant constituents, such as oils, lipids, enzymes, proteins, biopolymers, so that, for example, the

45

nutritional value, the yield or the industrial usability of the plant is improved.

A "promoter" refers, in accordance with the invention, to a
5 nucleotide sequence region which governs transcription of a gene,
or the synthesis of the corresponding mRNA. The promoter
comprises a sequence which is positioned 5'-upstream of the
transcription start. It comprises, as essential sequence element,
at least the so-called "TATA" box. Other regulatory elements,
10 such as the "CAAT" box or a GC box, may also be present. In
addition, it may be necessary for the promoter sequence according
to the invention to have, in addition to the abovementioned
sequence section, a sequence positioned 3'-downstream of the
transcription start, for example a leader sequence or part
15 thereof, so as to show, or fully exert, the desired promoter
activity and/or promoter specificity.

"Resistance" for the purposes of the present invention denotes
20 the artificially induced resistance or tolerance of the
transgenic plants to herbicides and/or pathogens, for example
fungi, viruses or insects, to specific external conditions, such
as high concentrations of ozone, sulfur dioxide, nitric oxides or
other exogenous pollutants, and to high and low temperatures,
25 drought or UV light.

A "modification of the photosynthesis rate of a plant" embraces
the reduction in, and, in particular, increase of, the
photosynthetic activity of the transformed plants. This may be
30 effected for example by expressing genes which improve the light
utilization of the plant in a targeted manner, increase the
reaction rate of individual, rate-determining metabolic steps, or
have an effect on the exchange of substances with the
environment.

35 "Leaf specificity" for the purposes of the present invention
means that a foreign gene which is under the control of a
promoter according to the invention is expressed in the entire
leaf organ or in specific leaf tissues preferably in the
40 mesophyll (for example palisade parenchyma), but not in the shoot
or in other parts of the plant such as, in particular, the roots.
In particular, "leaf specificity" for the purposes of the present
invention also exists when the promoter according to the
invention favors the expression of a foreign gene in the leaf,
45 preferably in the mesophyll of leaves, in particular of source
leaves, in comparison with other plant organs, such as stem,
non-germinating tubers, fruits or seeds of the plant, and causes,

in leaves, a significantly higher expression, for example at least approximately 5 to 10 times, such as 10 to 100 times higher.

"Source leaves" of a plant are the old leaves of a plant which fix an excess of carbon by means of photosynthesis and thus export bound carbon into other plant organs, for example the younger "sink" leaves.

- 10 "Permanent" expression for the purposes of the present invention denotes an expression, of the gene under the control of the promoter according to the invention, which is essentially independent of exogenously applied chemical induction signals and which persists over one or more plant generations.

Firstly, the present invention relates to promoters which cause a preferentially permanent, leaf-specific expression, in plants, of encoding nucleotide sequences under the control of said promoters.

- 20 More particularly, the present invention relates to a promoter which causes a mesophyll-specific expression, in plants, of an encoding nucleotide sequence under the control of said promoter, characterized in that said promoter comprises a promoter sequence of a cytosolic fructose-1,6-bisphosphatase from plants of the genus *Solanum* which comprises the nucleotide sequence SEQ ID NO:1; SEQ ID NO:2 or SEQ ID NO:3.


- 30 The primary site of action of herbicides and of a large number of pathogens is the leaf tissue, so that a leaf-specific expression of the resistance genes in question would provide sufficient protection. Since photosynthesis equally proceeds in the leaf tissue, the modification and, in particular, improvement of the photosynthetic performance would require the leaf-specific expression of one or more genes which have an effect on the photosynthesis rate.

The promoters according to the invention now have the surprising advantage of being able to express resistance genes specifically at the actual site of action within the plant. On the other hand, it is possible, for the first time, to influence the photosynthesis rate in a targeted manner by using the promoters according to the invention. As surprisingly demonstrated by the test results, preferred promoters make possible, for the first

6a

time, specific localization and expression of a foreign gene in the mesophyll of leaves, in particular of source leaves, while no activity can be found in parenchymatic tissue and also in xylem, phloem and others.

A promoter according to the invention can be provided by isolating and characterizing promoters of leaf-specifically and preferentially permanently expressed genes. Preferred promoters are those which correspond essentially to the promoters of the cytosolic fructose-1,6-bisphosphatase genes (cy-FBPase genes)



from leaf-specific mesophyll cells of plants. Especially preferred according to the invention are cy-FBPase promoters which have been isolated from leaf-specific mesophyll cells of plants of the genus *Solanum* (potatoes), and functional
5 equivalents thereof. A preferred embodiment relates to a nucleotide sequence with the desired promoter activity which is isolated from *Solanum tuberosum* var. Desiree, or functional equivalents thereof. Particularly preferred is a promoter with a nucleotide sequence, chosen from SEQ ID NO: 1 to 5, or functional
10 equivalents of these sequences.

The transcription start point in the preferred nucleotide sequence in accordance with SEQ ID NO: 1 was determined with the aid of "primer extension" using an ALF (automatic laser
15 fluorescence DNA sequencer (Pharmacia)). To this end, a 5'-fluorescence-labeled oligonucleotide was constructed which is complementary to a 21 bp region in the promoter from +1577 to +1599 (SEQ ID NO: 1). With the aid of this primer, total RNA from source leaves was transcribed into single-stranded cDNA. The RNA
20 which was not recognized by the primer and the RNA portions of the cDNA/RNA hybrids were subsequently digested. The cDNA was then analyzed on the ALF simultaneously with the promoter-DNA sequenced with the same primer. It was possible to identify the transcription start point by comparing the signals in the
25 sequence gel. According to this, sequence SEQ ID NO: 1 comprises 1428 bp promoter region (SEQ ID NO: 4) and 292 bp 5'-untranslated region of the cyt FBPase. A TATA box sequence ("TTATAAA") was found 30 bp upstream of the start point, and a CAAT box ("ATCATCCAAACAT") 141 bp upstream of the start point. In
30 addition, several inverted and direct sequence repeats were found which show no homology to sequence repeats which have been found in other promoters.

35 The direct and inverted sequence repeats which have been with a length of at least 10 bp, are listed in the tables which follow.

40

45

Direct sequence repeats

	Fragment starting at base	Repeat starting at base	Size	Repeat sequence	
				5'	3'
5	225	1316	10	AAGGATATTT	
	269	1686	11	TCTTTTTTTTTT	
	825	1016	12	TCAAAAGTTATG	
	1039	1493	13	ATATGTGACGTGG	
10	1083	1535	11	ATAGAAACAAA	
	1085	1411	10	AGAAACAAA	
	1172	1608	12	GTGCCAACCACT	
	1203	1639	13	CTCTTCCACGTG	

15

Inverted sequence repeats

	Fragment starting at base	Repeat starting at base	Size	Repeat sequence	
				5'	3'
20	99	293	10	CAAACATTTT	
	275	275	10	TTTTTAAAAA	
	313	1132	10	AACTTCTGTT	
	535	535	10	TGCATATGCA	
	835	835	10	TGCAGCTGCA	
25	1269	1370	11	TGTATATCAAA	
	1293	1359	10	TCATCCAAAC	
	1401	1401	-10	TTTTATAAAA	
	1658	1658	12	TCTGACGTCAGA	

30

The position data are based in each case on the numbering of the nucleotide residues in accordance with SEQ ID NO:2.

The above list contains, in particular, two virtually identical
 35 palindromic sequence sections of 10 bp which contain in each case
 twice the motif TGCA. This motif exists in the complementary
 strand as ACGT, a box which has been identified by various
 workers as a regulatory sequence (for example Guliano et al.,
 (1988), Proc. Acad. Natl. Sci. USA, 85, 7089-7093) and as a
 40 binding site for DNA-binding leucine zipper proteins (for example
 Armstrong et al., (1992), Plant Cell, 4, 525-537). The binding of
 leucine zipper proteins is probably not affected by the
 orientation of the motif. These sequences are marked in Figure 2
 by underlining with a dotted line. The effect of the above
 45 part-sequences on the promoter activity and/or promoter
 specificity according to the invention can be tested by those

skilled in the art, for example with the aid of customary deletion experiments.

A further preferred embodiment of the invention relates to a
5 5'-shortened promoter sequence (SEQ ID NO: 3). It comprises a 817
bp promoter region (SEQ ID NO: 5) and a 292 bp 5'-untranslated
region. Surprisingly, the organ or tissue specificity of the
shortened promoter is identical to that of the above-described,
longer promoter, but the promoter activity of the shortened
10 promoter is different.

Functionally equivalent promoter sequences are, according to the
invention, those sequences which despite differing nucleotide
15 sequence still have the desired functions, ie. promoter activity
and tissue or organ specificity. A measure for the promoter
activity is, for example, the expression rate determined for a
certain marker gene which is under the regulatory control of the
promoter according to the invention. Examples of suitable marker
20 genes are the β -glucuronidase (GUS) gene from E. coli or the green
fluorescence protein (GFP) gene (Baulcombe et al., (1993), Plant
J., 7 (6), 1045-1053)). The organ or tissue specificity can be
determined readily by comparing the expression rates, for the
above marker genes, determined in individual tissues or organs of
25 the plant. Functional equivalents for the purposes of the present
invention embrace naturally occurring variants of the sequences
described herein as well as artificial nucleotide sequences, for
example those obtained by chemical synthesis and adapted to the
codon usage of a plant.

30 Functional equivalents are to be understood as meaning, in
particular, also natural or artificial mutations of an originally
isolated promoter sequence which furthermore show the desired
function. Mutations embrace substitutions, additions, deletions,
35 exchanges or insertions of one or more nucleotide radicals. Thus,
the present invention also embraces for example those nucleotide
sequences which are obtained by modifying the nucleotide sequence
in accordance with SEQ ID NO:1 to 5. The aim of such a
modification can be, for example, a further limitation of the
40 promoter sequence contained therein, or else, for example, the
insertion of further restriction enzyme cleavage sites.

Functional equivalents are also those promoter variants whose
promoter function is attenuated or more pronounced in comparison
45 with the wild type.

If appropriate, leaf-specifically expressed genes must first be identified by experiments prior to isolation of the promoter sequence, for example by subtractive hybridization (for example as described by R.A. Meyers, *Molecular Biology and Biotechnology* 5 (1995), VCH, pp. 698-699). Next, a genome library can be established from leaves of the donor organism using known processes, for example by isolating the total DNA, subsequent partial digestion, packaging of fragments of defined size in bacteriophages, infection of bacteria with the recombinant 10 bacteriophages and subsequent amplification of the genome library. The phages containing the genomic DNA can then be transferred for example onto nylon filters and hybridized with a radiolabeled cDNA of the previously identified leaf-specific gene. Hybridizing phage DNAs can be visualized by autoradiography 15 and then individualized. To isolate the phage DNA, it is possible to start from one individual plaque and to inoculate and incubate lytic agar plates and to obtain the DNA in a manner known per se, for example by phenol/chloroform extraction followed by precipitation with ethanol. The fragment lengths of the promoter 20 regions of the genomic clones which have been isolated can now be determined for example by Southern hybridizations using a 5'-cDNA probe of the leaf-specifically expressed gene following various restriction cleavages. Now, a promoter region can be cloned into a suitable vector, propagated, for example in *E. coli*, and the 25 complete nucleotide sequence of the promoter can be determined by sequencing. Suitable cloning vectors are, inter alia, pBR332, pUC series, M13mp series and pACYC184.

30 Expediently, the promoter may now be linked operatively with a suitable gene in an expression cassette so that the promoter is capable of controlling transcription of the gene fused to it. Operative linkage is to be understood as meaning the sequential arrangement of promoter, encoding sequence, terminator and, if 35 appropriate, other regulative elements, allowing each of the abovementioned elements to fulfil its function in gene expression as intended.

The present invention furthermore relates to such an expression 40 cassette. The invention furthermore relates to recombinant vectors, for example plasmids or viruses, which comprise at least one expression cassette according to the invention.

In principle, the nucleotide sequences downstream of the promoter 45 sequence of the expression cassette may contain all open reading frames which are possible for any peptide, and one or more introns. Examples which may be mentioned are: sequences for

enzymes; sequences which are complementary to a) a genome sequence, it being possible for the genome sequence to be an open reading frame; b) an intron; c) a non-encoding leader sequence; d) any sequence which - when integrated into the genome in a complementary fashion - inhibits transcription, mRNA-processing (for example splicing) or translation.

The nucleotide sequence inserted can be produced synthetically or obtained naturally or comprise a mixture of synthetic and natural DNA components. In general, synthetic nucleotide sequences are produced with codons preferred by plants. These codons preferred by plants can be determined from codons with the highest frequency of protein and which are expressed in the most interesting plant species. When preparing an expression cassette, various DNA fragments can be manipulated in order to obtain a nucleotide sequence which expediently reads in the correct direction and which is provided with a correct reading frame. To link the DNA fragments to each other, adaptors or linkers may be added to the fragments.

20

The promoter regions according to the invention and the terminator regions should expediently be provided in the direction of transcription with a linker or polylinker comprising one or more restriction sites for inserting this sequence. As a rule, the linker has 1 to 10, in most cases 1 to 8, preferably 2 to 6 restriction sites. In general, the linker has a size of less than 100 bp, frequently less than 60 bp, but at least 5 bp, inside the regulatory regions. The promoter according to the invention can be native or homologous to the host plant, but also foreign or heterologous thereto. The expression cassette according to the invention comprises, in the 5'-3' direction of transcription, the promoter according to the invention, any sequence and a region for transcriptional termination. Various termination regions can be exchanged for each other as desired.

35

It is furthermore possible to employ manipulations which provide suitable restriction cleavage sites or which remove excess DNA or restriction cleavage sites. Where insertions, deletions or substitutions, for example transitions and transversions, are suitable, it is possible to use in-vitro mutagenesis, "primer repair", restriction or ligation. In the case of suitable manipulations, for example restriction, "chewing back" or filling up overlapping sections for blunt ends, it is possible to provide complementary ends of the fragments for ligation.

45

Especially suitable encoding nucleotide sequences are tolerance- or resistance-inducing genes, genes which increase the photosynthetic performance of the plant, or marker genes such as the β -glucuronidase gene (GUS) from *Escherichia coli*. Examples of
5 suitable tolerance genes are those which increase the tolerance of a plant to temperature, drought, UV or environmental pollutants. Examples of suitable resistance genes are the bar gene from *Streptomyces hygrosopicus*, which mediates resistance to the total herbicide phosphinothricin, chitinase genes which
10 mediate tolerance to fungal infections, and ribozyme genes whose RNA transcripts are capable of recognizing and cleaving viral RNA with high specificity. These and other resistance genes are known from *Transgenic Plants and Crop Improvement*, in *Transgenic Plants*, Vol. 1, Engineering and Utilization, edited by S-d Kung
15 and R. Wu, Academic Press, 1993, Part III, pp. 243-372; and from R.H. Symons (1992), Small catalytic RNAs, *Ann. Rev. Biochem.* 61, 641-671. Examples of suitable genes for increasing the photosynthetic performance are the genes which encode sucrose phosphate synthase (SPS) or fructose-1,6-bisphosphatase (FBPase).

20

The fused construct can now be transferred into plant genomes by various known processes. Examples of suitable processes are protoplast transformation by polyethylene glycol-induced DNA
25 uptake, electroporation, sonication or microinjection, and the transformation of intact cells or tissue by micro- or macroinjection into tissue or embryos, tissue electroporation, incubation of dry embryos in DNA-containing solution, biolistic gene transfer and, especially preferably, *Agrobacterium* transformation. The abovementioned processes are described, for
30 example, in B. Jenes et al., *Techniques for Gene Transfer*; in *Transgenic Plants*, Vol. 1, Engineering and Utilization, edited by S-d Kung and R. Wu, Academic Press, 1993, pp. 128-143 and in Potrykus (1991) *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42, 205-225. The fused construct is preferably cloned into a vector
35 which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al. (1980) *Nucl. Acids Res.* 12, 8711). The present invention furthermore relates to such vectors and to microorganisms transformed with them, in particular
40 *Agrobacterium*.

40

Then, *agrobacteria* which have been transformed with a vector according to the invention can be used in a known manner for the transformation of plants, in particular crop plants such as
45 cereals, maize, soya, rice, cotton, sugarbeet, canola, sunflower, flax, potato, tobacco, tomato, oilseed rape, alfalfa, lettuce and the various tree, nut and grape vine species, for example by

bathing scarified leaves or leaf sections in an agrobacteria solution and subsequently culturing them in suitable media.

The present invention furthermore relates to the use of vectors
5 according to the invention for the transformation of plants. The transformation of plants by agrobacteria is known, inter alia, from F.F. White, Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization, edited by S-d Kung and R. Wu, Academic Press, 1993, pp. 15-38 and from
10 S.B. Gelvin, Molecular Genetics of T-DNA Transfer from Agrobacterium to Plants, also in Transgenic Plants, pp. 49-78. Transgenic plants which express the gene which is under the control of the promoter introduced and fused thereto, in a leaf-specific manner, can be regenerated from the transformed
15 cells of the scarified leaves or leaf sections in a known manner. The present invention furthermore relates to such transgenic plants, propagation material thereof and to plant cells, plant tissue or parts of plants.

20 The cloning steps carried out for the purposes of the present invention, for example restriction cleavage, agarose gel electrophoresis, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linking DNA
25 fragments, transformation of E. coli cells, growing bacteria, multiplication of phages and sequence analysis of recombinant DNA were carried out as described by Sambrook et al. (1989) Cold Spring Harbor Laboratory Press; ISBN 0-87969-309-6.

30 The invention is illustrated by the examples which now follow, but not limited thereto:

The bacterial strains used hereinbelow (E. coli, XL-1 Blue and P2 392) were obtained from Stratagene. The Agrobacterium tumefaciens
35 strain employed for the transformation of plants (C58C1 with the plasmid pGV 3850kan) was described by Debleare et al. (1985, Nucl. Acid Res. 13, 4777). The cloning vectors used were pUC19 (Yanish-Perron (1985) Gene 33, 103-119), pBlueScript SK (Stratagene), pBin19 (Bevan (1984) Nucl. Acids Res. 12,
40 8711-8720) and pBI101 (Jefferson et al. (1987) EMBO J. 6, 3901-3907).

Solanum tuberosum L variety Desirée was obtained from Vereinigte Saatzuchten eG Ebstorf.

45 Nicotiana tabacum L Samsun NN was obtained from Vereinigte Saatzuchten eG Ebstorf.

Example 1: Isolation of a leaf-specific promoter

1. Isolation of a leaf-specifically expressed gene

5 1.1. Probe used

A cDNA probe of the potato cy-FBPase gene, produced using reverse transcriptase (EMBL No.: X76946) was used for the experiments
10 described hereinbelow. The cDNA probe (Figure 10) embraced 1487 nucleotides. The region which encodes the structural gene (FBPase) embraces nucleotides 199 to 1218.

15 1.2. Establishing a genome library

To establish a genome library from potato (*Solanum tuberosum* var. Desiree), total DNA from potato leaves was isolated by the method described by Rogers et al. ((1985) *Plant Mol. Biol.* 5, pp. 69-76). 300 µg of the DNA were subsequently subjected to partial
20 digestion with the restriction enzyme *Sau3A*, and the fragments between 12 and 20 kb were isolated by means of sucrose gradient centrifugation, dialyzed and concentrated by shaking with butanol.

25 The DNA was ligated into *Bam*HI-digested EMBL3 arms obtained from Stratagene (11099 North Torrey Pines Road, La Jolla, CA 92037, USA) following the manufacturer's information and subsequently packaged in vitro (*Gigapack II Gold* packaging extracts, Stratagene, following the manufacturer's information). *E. coli*
30 bacteria of strain P2 392 (Stratagene) were infected with the recombinant lambda phages, the titer of the library was determined, and the library was subsequently amplified.

35 1.3. Screening of the genome library, and isolation of the cy-FBPase gene

To isolate a genome clone comprising the cy-FBPase gene, 3×10^5 phages were plated. After the phages had been transferred to
40 nylon filters (*Hybond*^N*, Amersham Buchler), the filters were baked for 2 hours at 80°C for fixing. They were subsequently prehybridized in *Hypo-Hybond*^N* buffer at 42°C.

45 1 l of *Hypo-Hybond*^N* buffer contains:

250 ml of 1M sodium phosphate buffer pH 7.2

50 ml of 5M NaCl

* trademarks

15

2 ml of 0.5M EDTA pH 8.0

2 ml of sonicated herring sperm DNA, 1 mg/l

400 ml of formamide

5 50 g of PEG 6000

70 g of SDS

200 ml of water

10 The potato cy-FBPase sample which had been radiolabeled with High-Prime (Boehringer Mannheim) was first denatured for 5 minutes at 95°C and then placed in the prehybridization solution. The filters were hybridized overnight at 42°C. After the radioactive hybridization solution had been stripped off, the
15 filters were washed for 20 minutes at 42°C in 2X SSC (an NaCl/sodium citrate buffer), 0.1% SDS. The filters were subsequently washed again for 20 minutes at the same temperature using 1X SSC, 0.1% SDS. Then, a film was placed on the filters and exposed overnight at -70°C.

20

4 hybridizing phage DNAs were visualized by autoradiography and individualized. Starting from in each case one individual plaque, in each case one lytic agar plate was inoculated, incubated overnight at 37°C, and the phages were rinsed off the next day
25 with 10 ml of phage buffer (SM). The phage supernatant was subsequently treated with chloroform and the bacteria were centrifuged off. The supernatant was treated with in each case one spatula-tip of DNase and RNase and the mixtures were incubated for 30 minutes at 37°C. After 100 µl of 0.5 M EDTA and
30 200 µl of 10% strength SDS solution had been added, the batch was incubated for a further 20 minutes at 65°C. Then, 4.5 ml of 3M potassium acetate solution pH 4.8 were added and the batch was mixed and centrifuged. The supernatant was subsequently extracted by shaking with phenol:chloroform:isoamyl alcohol (volumetric
35 ratio 25:24:1). After extraction, the DNA was precipitated from the supernatant by adding two volumes of ethanol, and the resulting sediment was dissolved in 600 µl of TE-RNase.

2. Localization of the promoter

40

The fragment lengths of the promoter regions of the 4 clones which had been isolated were determined by Southern hybridizations with a 5'-cDNA sample following various
45 restriction cleavages. Clone FBP-1 was selected for further analyses. An approximately 7100 b BamHI fragment of clone FBP-1 was cloned into the BamHI cleavage site of vector pUC19 for further characterization. Sequencing, Southern hybridization and

16

restriction analysis allowed the promoter region to be restricted to a 1724 base pair fragment (Figure 1A). The probes used for the Southern hybridization were the 5'-342 bp (*HincII/EcoRI*) and 3'-216 bp (*EcoRI/EcoRV*) subfragments of the cDNA of the cy-FBPase. It was known from the sequence of the cDNA of the cytosolic FBPase that the restriction enzyme *ScaI* cleaves in the non-encoding 5' region of the cDNA. A single *ScaI* cleavage site was found in the genomic clone FBP-1. Sequence information on the genomic clone made it possible to demonstrate that this was the cleavage site which corresponded to the cDNA. It was used to remove the promoter region from the encoding region. To this end, the promoter region *XbaI/ScaI* was excised from the clone FBP-1, the ends were filled up and ligated into the vector pBlueScript-SK (pBSK-), which had previously been *SpeI*-cleaved and filled up (termed FBP:pBlue). The preparation of FBP:pBlue is shown diagrammatically in Figure 1B. The complete DNA sequence was subsequently determined by sequencing (Figure 2).

Example 2: Preparation of a transformation vector

20

1. Preparation of plasmid FBP:GUS

The expression characteristics of the new promoter were analyzed by marker gene experiments. To this end, the cy-FBPase promoter was fused with the β -glucuronidase gene (GUS) from *E. coli*. The promoter was isolated from plasmid FBP:pBlue as a *BamHI* fragment and cloned into the *BamHI* cleavage site of expression vector pBI101 (Figure 3A) (Jefferson et al., (1987), *EMBO J.* 6, 3901-3907). The resulting plasmid FBP:GUS (Figure 3B) was subsequently employed for the transformation of *Agrobacterium tumefaciens*.

2. Preparation of the deletion construct FBP:GUS(DEL)

35

A deletion construct was prepared which comprises approximately 1.1 kb of the promoter sequence. To this end, a fragment comprising the GUS gene, the NOS terminator and 1100 bp of the promoter was excised from FBP:GUS by means of *EcoRI* digest. This fragment was isolated, purified and ligated into the *EcoRI* cleavage site of the vector pBin 19 (Figure 3A). The orientation of the fragment in pBin 19 was checked by cleavage with *BamHI*. *Agrobacterium tumefaciens* was also transformed with this vector.

45

Example 3: Transformation of *Agrobacterium tumefaciens*

The transformation of *Agrobacterium tumefaciens* was carried out following the method of Höfgen and Willmitzer (Nucl. Acid Res. 5 (1988) 16, 9877). The agrobacteria were grown in YEB medium (Vervliet et al., J. Gen. Virol. (1975) 26, 33).

Example 4: Transformation of the cy-FBPase promoter in tobacco and potato plants, and analysis of expression
10**1.1. Transformation of tobacco**

For the transformation of tobacco plants (*Nicotiana tabacum* L. 15 cv. Samsun NN), 10 ml of an overnight culture of *Agrobacterium tumefaciens* which had been transformed with FBP:GUS or FBP:GUS (DEL) and grown under selection were centrifuged, the supernatant was discarded, and the bacteria were resuspended in the same volume of an antibiotic-free medium. Leaf disks of sterile plants 20 (approximate diameter 1 cm) were bathed in this bacterial solution in a sterile Petri dish. The leaf disks were subsequently placed on MS medium (Murashige and Skoog, *Physiol. Plant.* (1962) 15,473) with 2% sucrose and 0.8% Bacto agar in Petri dishes. After 25 incubation in the dark for 2 days at 25°C, they were transferred to MS medium with 100 mg/l kanamycin, 500 mg/l claforan, 1 mg/l benzylaminopurine (BAP), 0.2 mg/l naphthylacetic acid (NAA), 1.6% glucose and 0.8% Bacto agar, and culturing was continued (16 hours light/8 hours dark). Growing shoots were transferred to 30 hormone-free MS medium with 2% sucrose, 250 mg/l claforan and 0.8% Bacto agar.

1.2. Transformation of potatoes

35 20 small leaves, scarified with a scalpel, of a sterile potato culture were placed into 10 ml of MS medium with 2% sucrose which contained 50 µl of an FBP:GUS or FBP:GUS(DEL) transformed *Agrobacterium tumefaciens* overnight culture which had been grown under selection. After gentle shaking for 5 minutes, the Petri 40 dishes were incubated at 25°C in the dark. After two days, the leaves were placed on MS medium with 1.6% glucose, 2 mg/l zeatin ribose, 0.02 mg/l gibberellic acid, 500 mg/l claforan, 50 mg/l kanamycin and 0.8% Bacto agar. After incubation at 25°C and 3000 45 Lux for one week, the claforan concentration in the medium was halved. Further culturing was carried out by known methods (Rocha-Sosa et al. (1989) *EMBO J.* 8, 23-29).

1.3. Expression analysis of the cy-FBPase promoter in transgenic tobacco and potato plants

In each case 60 transformed plants from amongst the transformed
5 tobacco and potato plants were regenerated, and the
 β -glucuronidase activity was determined. β -Glucuronidase was
detected as described by Martin et al. (1992) in: The GUS
Reporter System as a Tool to Study Plant Gene Expression in: GUS
Protocols: Using the GUS Gene as a Reporter of Gene Expression,
10 Academic Press, pp. 23-43. To analyze the expression in greater
detail, 40 tobacco plants and 21 potato plants were selected.
After the transformed plants had been transferred into the
greenhouse, the organ-specific β -glucuronidase expression was
determined.
15

Figure 4 shows a comparison of the enzyme activities in different
potato tissues. The data show that the promoter mediates
leaf-specific expression of the reporter gene.

20 Figure 5 shows a comparison of the GUS activity in various organs
of the wild type and of transgenic tobacco plants which carry the
GUS gene under the control of a variety of promoters according to
the invention. TME-1/67 refers to a plant which has been
25 transformed with FBP:GUS. TME-11/13 refers to a plant which has
been transformed with FBP:GUS (DEL).

The determination of the GUS activity in the organs sink leaf,
source leaf, stem and root of 9-week-old tobacco plants and of
30 seeds revealed that, in both transformation lines, the highest
activity is to be found in source leaves and a markedly lower
activity in sink leaves. The measurements in sink leaves always
differed greatly within one leaf. The activity in stem and root
was only marginally above the background activity measured in
35 wild-type tobacco. In these tissues, the promoter is not active.
Slightly elevated activity levels were measured in tobacco seeds
in comparison with wild-type seeds, even though no mRNA could be
detected in the Northern blot. GUS activity in seeds was also
found when seed homogenate was incubated in X-Gluc solution.
40 Wild-type seeds show no staining (not shown). It is possible that
the promoter is active in the course of seed development and not
in the mature seeds themselves. The GUS activity might be
explained by stored protein. Depending on the plant, the activity
found in the leaves was higher by a factor of approximately 10 to
45 50 in comparison with seeds.

19.

Figure 6 illustrates the cell specificity of the construct FBP:GUS according to the invention. Identical histological findings are obtained with the shortened promoter construct FBP:GUS(DEL). To investigate the cell specificity of the cyt
5 FBPase promoter in the leaf in greater detail, leaf cross-sections were prepared of fully unfolded tobacco leaves of 8-week-old plants which had been grown in the greenhouse. The sections were fixed in 3% paraformaldehyde for 20 minutes, stained overnight in X-Gluc (5-bromo-4-chloro-3-indolyl-
10 β -D-glucuronide) solution and the chlorophyll was subsequently removed using 70% ethanol. In addition, the epidermis was removed from the mesophyll and incubated separately to avoid contamination with dibromodichloro-indigo, which had been released into the staining solution by damaged mesophyll cells.

15

Figure 6 (A) shows a cross-section through the central vascular bundle of a source leaf. It can be seen clearly that, in the central vascular bundle, only the mesophyll cells were stained on the upper side. The parenchymatic tissue and also xylem, phloem
20 and other tissues localized here were not stained. Some of the epidermis cells seem to have been stained. This is probably not an activity in the cells, since the epidermis incubated in isolation shows no GUS activity in trichomes, stomata and epidermal cells (B). Staining of the epidermal cells (A) might be
25 attributed to contamination from the excised mesophyll cells, or to the fact that the sections contained several layers and mesophyll cells located behind the epidermal cells were seen through them. No blue staining was observed in the petiole (C). In the mesophyll cross-section, very strong expression was found
30 in the palisade parenchyma and a slightly less pronounced expression in the spongy parenchyma (D).

To investigate the expression in greater detail, the total RNA
35 was isolated from different organs of tobacco plants (transformed with FBP:GUS), and GUS-specific transcripts were detected by means of Northern analyses (Figure 7). The isolation was carried out as described by Logemann et al. (Anal. Biochem. (1987) 163,21). For the analysis, in each case 20 to 40 μ g of RNA were
40 separated in a 1.5% agarose gel comprising formaldehyde. After electrophoretic separation of the RNA molecules, the RNA was transferred to a nylon membrane by means of capillary transfer. The detection of specific transcripts was carried out as described by Amasino (Anal. Biochem. (1986) 152, 304). The cDNA
45 fragments employed as probe were radiolabeled with a Random Primed DNA Labeling Kit (Boehringer, Mannheim). GUS-specific transcripts were only detected in sink and source leaves, while no positive signal was identified in stalks, roots and seeds. A

20

comparison of the β -glucuronidase activity obtained in sink and source leaves demonstrates that the activity is much higher in source leaves. Histochemical analyses of seedlings revealed that uniform leaf-specific expression is guaranteed even in early 5 developmental stages of tobacco plants (Figure 8).

Example 5: Construction of the vector pBin-FBP

10 The promoter fragment was excised from pUC19 using BamHI/SacI; the ends were filled up with T4-DNA-polymerase. The fragment was subsequently separated from vector pUC19 by electrophoresis on a 1% TAE agarose gel and purified with glass milk (BIO 101.1070 Joshua Way, Vista, CA 92083, USA). It was then ligated into a
15 pBin19 derivative which had been cleaved with EcoRI and treated with Klenow enzyme. This derivative thus comprised an expression cassette embracing the cy-FBPase promoter and the terminator sequence of octopine synthase from *Agrobacterium tumefaciens* (Figure 9).

20

25

30

35

40

45

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: BASF Aktiengesellschaft
- (B) STREET: -
- (C) CITY: Ludwigshafen
- (E) COUNTRY: Deutschland
- (F) POSTAL CODE (ZIP): D-67056

(ii) TITLE OF INVENTION: Leaf-specific gene expression in transgenic plants

(iii) NUMBER OF SEQUENCES: 5

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1720 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Solanum tuberosum
- (B) STRAIN: Desiree
- (F) TISSUE TYPE: Leaf

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION:1429..1720
- (D) OTHER INFORMATION:/function= "part of the leader sequence of the cy-FBPase"

(ix) FEATURE:

- (A) NAME/KEY: promoter
- (B) LOCATION:1..1428

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION:1429
- (D) OTHER INFORMATION:/function= "start of transcription"

(ix) FEATURE:

- (A) NAME/KEY: TATA_signal
- (B) LOCATION:1399..1405

(ix) FEATURE:

- (A) NAME/KEY: CAAT_signal
- (B) LOCATION:1288..1300

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```

CCAGCTAATG CTGCTCTTGT CACTCAAAT GATGGTATCC CTCTCGTCAT CCAGTGTTTG      60
TCAAGTCCTG TTAGGAACAC AGTAAGGATA TAAACAAACA TTTTGTGGTC TTCTTGGTTA     120
TTGAGTGCTT GCTGTTCACT TGTTAAAATT GCACATATAC GTAGTGAGAA ACTCAACTGT     180
TGAGTACCAT TGATCCGTCA ATCTTGTCGA TAACTTTGAT AAGGATATTT CAGGCATCAG     240
ACATGTCACC TCTATAGAAC TTGGTCTTTT TTTTAAAAA TAAAAATAAA AATGTTTGGC     300
ATCATACGAA CTTCTGTTAC TTTAGGCTGT ATCCAGAATA AAATGTTGTT TCCTCATTCT     360
GGAATTAGTT GTTTTGCACA CGGAAGACTT TCGAAATTTA CTAATTGTGT TCGTCCGTCT     420
CAAACCTGGCT CACACTTTGG TGGTCAATTT TACTTCTCAA GGTAAGCAAT TACAGAATAT     480
GAATGTCGCT CTCCTCATAT TTATCCGAAC AATAAAAAAT GATATCTGTT TGCATATGCA     540
TGTAGATCAC ACAGGGGGGG GGGGGGGCGG GGATCTAAGG GAGCTAAATC TAATTTATAT     600
TAGTAGATGT TGAATCCGT TGGGCTTCAT TATGTGTTTT TACATATTCG TTTCTGAACC     660
ACCCCCACCC CGGTGAAAAA CATTGCTCTG CCACTGGCTC AATGTATTGA CACAAATGAA     720
CTTCAAACCTG GGCAGGTGAA TTATGCTCTA GGAGCATTGT ATTATCTATG CAATGCATCA     780
AACAAGGAAG AGATCTTAAA GCCAGAAGTA ATTGATGCAA TCAAAAGTTA TGCAGCTGCA     840
GGTGGAGTTA GTACAAGCTT CAGTAATTTG GCTCAGGCTT TCTTAGATCA ACATGTTCTT     900
CAGCTTAATT AAAATGGAGG AAACCAAAGA TTATGTTGTA AAATCATTIT CTATCCTAGA     960
TGGTCTATCG GAAACAATTT ATTTATTACT CCTATCCAAT TCATTATATT TTCAAAGTT    1020
ATGAAGTCCA CGAAATATGT GACGTGGGTA AAGAAGACCC ATGCCAAGCC AGTGGGATAT    1080
AGAAACAAAA CATGTAATAA AGAGAACAAA TAATGAGTTT CGAAAAGAAC AGAAGTTAGC    1140
ATAAGGACGA GAATCACATT ATCTTAGGTG CCAACCACTA ATCCTATGTA TCATTCTCCT    1200
CTTCCACGT GTCATCCTAC ACTTCCTTTG CCATCAGATT AGATAGCCCG GTTAGTACCT    1260
ACACTGTATA TCAAAAAATA CGTAACAATC ATCCAAACAT ATCATCGATC AAAGGATATT    1320
TATCTTGATG TGCTTTCGCC GTCCATTGTA ACGAGTTTGG ATGAATTTGA TATACACCCA    1380
CTCAGATATC AATATATTTT ATAAAAAGAA ACAAATTGA ATACTAGTAA TATCTATGTA    1440
GATATTTATT TTTTCAACAA TCCTGTAAGT TATAAGGATA ACTCACTTAT ATGTGACGTG    1500
GATAATGAAG AGCTAGGCAG GCAGTGAGAG ATAGAAACAA ATTAAGCAGA GACGAAAAAC    1560
AAATCAGTTA ACAGAATGAC GAATTGGATC ACGCTTTATC TTAGTGCCAA CCACTGATCC    1620
CATGCATCAC TCTGCTCTTT CCACGTGGCA TCCTCTGACG TCAGATCAGA TTCCTCTTCT    1680
TTCTTTTTTT TTTCTGTATA TATATGAGCA TTTTAGTAGT    1720

```

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1724 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Solanum tuberosum
- (B) STRAIN: Desiree
- (F) TISSUE TYPE: Leaf

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION:1433..1724
- (D) OTHER INFORMATION:/function= "Part of the leader sequence of the cy-FBPase"

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION:1..6
- (D) OTHER INFORMATION:/function= "BamHI Restriction Site"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

GGATCCAGCT AATGCTGCTC TTGTCACTCA AAATGATGGT ATCCCTCTCG TCATCCAGTG      60
TTTGTCAAGT CCTGTTAGGA ACACAGTAAG GATATAAACA AACATTTTGT GGTCTTCTTG      120
GTTATTGAGT GCTTGCTGTT CACTTGTTAA AATTGCACAT ATACGTAGTG AGAAACTCAA      180
CTGTTGAGTA CCATTGATCC GTCAATCTTG TCGATAACTT TGATAAGGAT ATTTCAGGCA      240
TCAGACATGT CACCTCTATA GAACTTGGTC TTTTTTTTTTA AAAATAAAAA TAAAAATGTT      300
TGGCATCATA CGAACTTCTG TTACTIONTAGG CTGTATCCAG AATAAAATGT TGTTTCCTCA      360
TTCTGGAATT AGTTGTTTTG CACACGGAAG ACTTTCGAAA TTTACTAATT GTGTTTCGTCC      420
GTCTCAAACCT GGCTCACACT TTGGTGGTCA ATTTTACTTC TCAAGGTAAG CAATTACAGA      480
ATATGAATGT CGCTCTCCTC ATATTTATCC GAACAATAAA AAATGATATC TGTTTGCATA      540
TGCATGTAGA TCACACAGGG GGGGGGGGGG GCGGGGATCT AAGGGAGCTA AATCTAATTT      600
ATATTAGTAG ATGTTGAATT CCGTTGGGCT TCATTATGTG TTTTACATA TTCGTTTCTG      660
AACCACCCCC ACCCCGGTGA AAAACATTGC TCTGCCACTG GCTCAATGTA TTGACACAAA      720
TGAACCTCAA ACTGGGCAGG TGAATTATGC TCTAGGAGCA TTGTATTATC TATGCAATGC      780
ATCAAACAAG GAAGAGATCT TAAAGCCAGA AGTAATTGAT GCAATCAAAA GTTATGCAGC      840
TGCAGGTGGA GTTAGTACAA GCTTCAGTAA TTTGGCTCAG GCTTCTTAG ATCAACATGT      900
TCCTCAGCTT AATTAAAATG GAGGAAACCA AAGATTATGT TGTAATAATCA TTTTCTATCC      960
TAGATGGTCT ATCGGAAACA ATTTATTTAT TACTCCTATC CAATTCATTA TATTTTCAA      1020
AGTTATGAAG TCCACGAAAT ATGTGACGTG GGTAAGAAG ACCCATGCCA AGCCAGTGGG      1080
ATATAGAAAC AAAACATGTA ATAAAGAGAA CAAATAATGA GTTTCGAAAA GAACAGAAGT      1140
TAGCATAAGG ACGAGAATCA CATTATCTTA GGTGCCAACC ACTAATCCTA TGTATCATTC      1200
TCCTCTTTCC ACGTGTCATC CTACACTTCC TTTGCCATCA GATTAGATAG CCCGGTTAGT      1260
ACCTACACTG TATATCAAAA AATACGTAAC AATCATCCAA ACATATCATC GATCAAAGGA      1320
TATTTATCTT GATGTGCTTT CGCCGTCCAT TGTAACGAGT TTGGATGAAT TTGATATACA      1380

```

CCCACTCAGA TATCAATATA TTTTATAAAA AGAAACAAAA TTGAATACTA GTAATATCTA 1440
 TGTAGATATT TATTTTTTCA ACAATCCTGT AAGTTATAAG GATAACTCAC TTATATGTGA 1500
 CGTGGATAAT GAAGAGCTAG GCAGGCAGTG AGAGATAGAA ACAAATTAAG CAGAGACGAA 1560
 AAACAAATCA GTTAACAGAA TGACGAATTG GATCACGCTT TATCTTAGTG CCAACCACTG 1620
 ATCCCATGCA TCACTCTGCT CTTTCCACGT GGCATCCTCT GACGTCAGAT CAGATTCCTC 1680
 TTCTTTCTTT TTTTTTCTG TATATATATG AGCATTTTAG TAGT 1724

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1109 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Solanum tuberosum
- (B) STRAIN: Desiree
- (F) TISSUE TYPE: Leaf

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION:1..6
- (D) OTHER INFORMATION:/function= "EcoRI Restriction Site"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GAATTCCGTT GGGCTTCATT ATGTGTTTTT ACATATTCGT TTCTGAACCA CCCCCACCCC 60
 GGTGAAAAC ATTGCTCTGC CACTGGCTCA ATGTATTGAC ACAAATGAAC TTCAAACTGG 120
 GCAGGTGAAT TATGCTCTAG GAGCATTGTA TTATCTATGC AATGCATCAA ACAAGGAAGA 180
 GATCTTAAAG CCAGAAGTAA TTGATGCAAT CAAAAGTTAT GCAGCTGCAG GTGGAGTTAG 240
 TACAAGCTTC AGTAATTTGG CTCAGGCTTT CTTAGATCAA CATGTTCCCTC AGCTTAATTA 300
 AAATGGAGGA AACCAAAGAT TATGTTGTAA AATCATTTTC TATCCTAGAT GGTCTATCGG 360
 AAACAATTTA TTTATTACTC CTATCCAATT CATTATATTT TCAAAAGTTA TGAAGTCCAC 420
 GAAATATGTG ACGTGGGTAA AGAAGACCCA TGCCAAGCCA GTGGGATATA GAAACAAAAC 480
 ATGTAATAAA GAGAACAAAT AATGAGTTTC GAAAAGAACA GAAGTTAGCA TAAGGACGAG 540
 AATCACATTA TCTTAGGTGC CAACCACTAA TCCTATGTAT CATTCTCCTC TTTCCACGTG 600
 TCATCCTACA CTTCTTTTGC CATCAGATTA GATAGCCCGG TTAGTACCTA CACTGTATAT 660
 CAAAAAATAC GTAACAATCA TCCAAACATA TCATCGATCA AAGGATATTT ATCTTGATGT 720
 GCTTTCGCCG TCCATTGTAA CGAGTTTGGG TGAATTTGAT ATACACCCAC TCAGATATCA 780
 ATATATTTTA TAAAAAGAAA CAAAATTGAA TACTAGTAAT ATCTATGTAG ATATTTATTT 840
 TTTCAACAAT CCTGTAAGTT ATAAGGATAA CTCACCTATA TGTGACGTGG ATAATGAAGA 900

GCTAGGCAGG CAGTGAGAGA TAGAAACAAA TTAAGCAGAG ACGAAAAACA AATCAGTTAA 960
 CAGAATGACG AATTGGATCA CGCTTTATCT TAGTGCCAAC CACTGATCCC ATGCATCACT 1020
 CTGCTCTTTC CACGTGGCAT CCTCTGACGT CAGATCAGAT TCCTCTTCTT TCTTTTTTTT 1080
 TTCTGTATAT ATATGAGCAT TTTAGTAGT 1109

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1428 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Solanum tuberosum
 (B) STRAIN: Desiree
 (F) TISSUE TYPE: Leaf

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CCAGCTAATG CTGCTCTTGT CACTCAAAT GATGGTATCC CTCTCGTCAT CCAGTGTTTG 60
 TCAAGTCCTG TTAGGAACAC AGTAAGGATA TAAACAAACA TTTTGTGGTC TTCTTGGTTA 120
 TTGAGTGCTT GCTGTTCACT TGTTAAAATT GCACATATAC GTAGTGAGAA ACTCAACTGT 180
 TGAGTACCAT TGATCCGTCA ATCTTGTCGA TAACTTTGAT AAGGATATTT CAGGCATCAG 240
 ACATGTCACC TCTATAGAAC TTGGTCTTTT TTTTAAAAA TAAAAATAAA AATGTTTGGC 300
 ATCATACGAA CTTCTGTTAC TTTAGGCTGT ATCCAGAATA AAATGTTGTT TCCTCATTCT 360
 GGAATTAGTT GTTTTGCACA CGGAAGACTT TCGAAATTTA CTAATTGTGT TCGTCCGTCT 420
 CAAACTGGCT CACTTTTGG TGGTCAATTT TACTTCTCAA GGTAAGCAAT TACAGAATAT 480
 GAATGTCGCT CTCCTCATAT TTATCCGAAC AATAAAAAAT GATATCTGTT TGCATATGCA 540
 TGTAGATCAC ACAGGGGGGG GGGGGGGCGG GGATCTAAGG GAGCTAAATC TAATTTATAT 600
 TAGTAGATGT TGAATTCCGT TGGGCTTCAT TATGTGTTTT TACATATTCG TTTCTGAACC 660
 ACCCCCACCC CGGTGAAAAA CATTGCTCTG CCACTGGCTC AATGTATTGA CACAAATGAA 720
 CTTCAAACCTG GGCAGGTGAA TTATGCTCTA GGAGCATTGT ATTATCTATG CAATGCATCA 780
 AACAAAGGAAG AGATCTTAAA GCCAGAAGTA ATTGATGCAA TCAAAGTTA TGCAGCTGCA 840
 GGTGGAGTTA GTACAAGCTT CAGTAATTTG GCTCAGGCTT TCTTAGATCA ACATGTTCCCT 900
 CAGCTTAATT AAAATGGAGG AAACCAAAGA TTATGTTGTA AAATCATTTT CTATCCTAGA 960
 TGGTCTATCG GAAACAATTT ATTTATTACT CCTATCCAAT TCATTATATT TTCAAAGTT 1020
 ATGAAGTCCA CGAAATATGT GACGTGGGTA AAGAAGACCC ATGCCAAGCC AGTGGGATAT 1080
 AGAAACAAAA CATGTAATAA AGAGAACAAA TAATGAGTTT CGAAAAGAAC AGAAGTTAGC 1140

ATAAGGACGA GAATCACATT ATCTTAGGTG CCAACCACTA ATCCTATGTA TCATTCTCCT 1200
 CTTTCCACGT GTCATCCTAC ACTTCCTTTG CCATCAGATT AGATAGCCCG GTTAGTACCT 1260
 ACACTGTATA TCAAAAATA CGTAACAATC ATCCAAACAT ATCATCGATC AAAGGATATT 1320
 TATCTTGATG TGCTTTCGCC GTCCATTGTA ACGAGTTTGG ATGAATTTGA TATACACCCA 1380
 CTCAGATATC AATATATTTT ATAAAAAGAA ACAAATTGA ATACTAGT 1428

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 817 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Solanum tuberosum
- (B) STRAIN: Desiree
- (F) TISSUE TYPE: Leaf

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION:1..6
- (D) OTHER INFORMATION:/function= "EcoRI Restriction Site"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GAATTCCGTT GGGCTTCATT ATGTGTTTTT ACATATTCGT TTCTGAACCA CCCCCACCCC 60
 GGTGAAAAC ATTGCTCTGC CACTGGCTCA ATGTATTGAC ACAAATGAAC TTCAAACCTGG 120
 GCAGGTGAAT TATGCTCTAG GAGCATTGTA TTATCTATGC AATGCATCAA ACAAGGAAGA 180
 GATCTTAAAG CCAGAAGTAA TTGATGCAAT CAAAAGTTAT GCAGCTGCAG GTGGAGTTAG 240
 TACAAGCTTC AGTAATTTGG CTCAGGCTTT CTTAGATCAA CATGTTCCCTC AGCTTAATTA 300
 AAATGGAGGA AACCAAAGAT TATGTTGTAA AATCATTTTC TATCCTAGAT GGTCTATCGG 360
 AAACAATTTA TTTATTACTC CTATCCAATT CATTATATTT TCAAAAGTTA TGAAGTCCAC 420
 GAAATATGTG ACGTGGGTAA AGAAGACCCA TGCCAAGCCA GTGGGATATA GAAACAAAAC 480
 ATGTAATAAA GAGAACAAT AATGAGTTTC GAAAAGAACA GAAGTTAGCA TAAGGACGAG 540
 AATCACATTA TCTTAGGTGC CAACCACTAA TCCTATGTAT CATTCTCCTC TTTCCACGTG 600
 TCATCCTACA CTTCTTTGTC CATCAGATTA GATAGCCCGG TTAGTACCTA CACTGTATAT 660
 CAAAAAATAC GTAACAATCA TCCAAACATA TCATCGATCA AAGGATATTT ATCTTGATGT 720
 GCTTTCGCCG TCCATTGTAA CGAGTTTGGG TGAATTTGAT ATACACCCAC TCAGATATCA 780
 ATATATTTTA TAAAAGAAA CAAAATTGAA TACTAGT 817

WHAT IS CLAIMED IS:

1. A promoter which causes a mesophyll-specific expression, in plants, of an encoding nucleotide sequence under the control of said promoter, characterized in that said promoter comprises a promoter sequence of a cytosolic fructose-1,6-bisphosphatase from plants of the genus *Solanum*, which comprises the nucleotide sequence SEQ ID NO:1; SEQ ID NO:2 or SEQ ID NO:3.
2. The promoter as claimed in claim 1, which comprises the promoter sequence of cytosolic fructose-1,6-bisphosphatase from leaf-specific mesophyll
10 cells of plants of the genus *Solanum*, which comprises the nucleotide sequence SEQ ID NO:1; SEQ ID NO:2 or SEQ ID NO:3.
3. The promoter as claimed in claim 1 or 2, which comprises a nucleotide sequence from nucleotide +1 to +1428 of SEQ ID NO:4 or +1 to +817 of SEQ ID NO:5.
4. An expression cassette, which comprises at least one promoter sequence defined in any one of claims 1 to 3.
5. The expression cassette as claimed in claim 4, wherein the expression of
20 the encoding nucleotide sequence under the control of the promoter according to any one of claims 1 to 3, imparts resistance to plants or an increase in the photosynthesis rate of the plant.
6. A recombinant vector which comprises the expression cassette defined in claim 4 or 5.
7. The vector as claimed in claim 6, selected from the group consisting of plasmids FBP:GUS and pBin-FBP.

8. A microorganism which contains the recombinant vector as claimed in either of claim 6 or 7.
9. The microorganism as claimed in claim 8, from the genus *Agrobacterium*.
10. The microorganism of claim 9, characterized in that the microorganism is of the species *Agrobacterium tumefaciens*.
11. Use of the vector as claimed in claim 6 or 7 or of the microorganism as claimed in any one of claims 8 to 10 for the transformation of plants, plant cells, plant tissues or parts of plants.
12. The use as claimed in claim 11, wherein the transformed plant is selected from the group consisting of cereal, maize, soya, rice, cotton, sugarbeet, canola, sunflower, flax, potato, tobacco, tomato, oilseed rape, alfalfa, lettuce and the various tree, nut and grape vine species.
13. The use of the promoter as claimed in any one of claims 1 to 3
 - a) for the generation of a plant bioreactor;
 - b) for altering the profile of plant constituents;
 - c) for manipulating the plant metabolism; and
 - d) for the transfer of resistance genes to plants.
14. A transgenic plant cell, transformed with the vector as claimed in either of claim 6 or 7 or with the microorganism as claimed in any one of claim 8 or 10.
15. The transgenic plant cell as claimed in claim 14, selected from cells of crop plants selected from the group consisting of cereal, maize, soya, rice, cotton, sugarbeet, canola, sunflower, flax, potato, tobacco, tomato, oilseed rape, alfalfa, lettuce and the various tree, nut and grape vine species.
16. A method of producing transgenic plants selected from the group of crop plants consisting of cereal, maize, soya, rice, cotton, sugarbeet, canola,

sunflower, flax, potato, tobacco, tomato, oilseed rape, alfalfa, lettuce and the various tree, nut and grape vine species, which comprises transforming plant cells, plant tissue, parts of plants or protoplasts with the vector as claimed in either of claim 6 or 7 or with the microorganism as claimed in either of claim 8 or 9, culturing the transformed cells, tissue, parts of plants or protoplasts in a growth medium and, if appropriate, regenerating plants from the culture.

10 17. A method of isolating the promoter causing mesophyll-specific expression, in plants, of an encoding nucleotide sequence under the control of said promoter, characterized in that said promoter comprises a promoter sequence of a cytosolic fructose-1,6-bisphosphatase from plants of the genus *Solanum*, which method comprises

- a) hybridizing a plant genome library with the complement of cytFBPase cDNA corresponding to EMBL No. X76946,
- b) isolating positive clones and
- c) testing the isolated clones for promoter activity.

18. A DNA molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, 2, 3, 4 and 5.

20 19. Functional equivalent of the promoter according to any one of claims 1 to 3, which causes a mesophyll-specific expression, in plants, of an encoding nucleotide sequence under the control of the promoter, characterized in that said functional equivalent is selected from the group consisting of

- a) artificial sequences, which are obtained by modifying the nucleotide sequence in accordance with SEQ ID NO:1 to 5 by adapting them to the codon usage of a plant; and
- b) mutants obtained from sequences according to any one of SEQ ID NO:1 to 5 by substitution, addition, deletion, reversal or insertion of one or more nucleotide according to SEQ ID NO:1 to SEQ ID NO:5, said mutants comprising at least one of the following repeat sequences with a minimum length of 10 bp:

32

10

5' AAGGATATTT 3'
5' TCTTTTTTTTTT 3'
5' TCAAAAGTTATG 3'
5' ATATGTGACGTGG 3'
5' ATAGAAACAAA 3'
5' AGAAACAAAA 3'
5' GTGCCAACCACT 3'
5' CTCCTTCCACGTG 3'
5' CAAACATTTT 3'
5' TTTTAAAAA 3'
5' AACTTCTGTT 3,
5' TGCATATGCA 3'
5' TGCAGCTGCA 3'
5' TGTATATCAAA 3,
5' TCATCCAAAC 3,
5, TTTTAAAAA 3'
5' TCTGACGTCAGA 3'.




Fig. 1A




Fig. 1B

3/12

GGATCCAGCTAATGCTGCTCTTGTCACCTCAAATGATGGTATCCCTCTCGTCATCCAGTGTTTGTC AAG
TCCTGTTAGGAACACAGTAAGGATATAAACAAACATTTTGTGGTCTTCTTGGTTATTGAGTGCTTGCTGTTCACT
TGTTAAAATTGCACATATACGTAGTGAGAACTCAACTGTTGAGTACCATMGATCCGTCAATCTTGTCGATAACT
TTGATAAGGATATTT CAGGCATCAGACATGTCACCTCTATAGA ACTTGGTCTTTTTTTTTTAAAAATAAAAAATAAA
AATGTTTGGCATCATACGA ACTTCTGTTACTTTAGGCTGTATCCAGAATAAAAATGTTGTTTCTCATTCTGGAAT
TAGTTGTTTTGCACACGGAAGACTTTCGAAATTTACTAATTGTGTTTCGTCCGTCTCAA ACTGGCTCACACTTTGG
TGGTCAATTTTACTTCTCAAGGTAAGCAATTACAGAATATGAATGTCGCTCTCCTCATATTTATCCGAACAATAA
AAAATGATATCTGTTTGCATATGCATGTAGATCACACACCCCCCCCCCCCCCGCCCTAGATTCCCTCGATTTAG
ATTAAATATAATCATCTACAAGAATTCCGTTGGGCTTCATTATGTGTTTTTACATATTCGTTTCTGAACCACCCC
CACCCCGGTGAAAACATTGCTCTGCCACTGGCTCAATGTATTGACACAAATGAACTTCAA ACTGGGCAGGTGAA
TTATGCTCTAGGAGCATTGTATTATCTATGCAATGCATCAAACAAGGAAGAGATCTTAAAGCCAGAAGTAATTGA
TGCAATCAAAGTTATG CAGCTGCAGGTGGAGTTAGTACAAGCTTCAGTAATTTGGCTCAGGCTTCTTAGATCA
ACATGTTCTCAGCTTAATTTAAAATGGAGGAAACCAAAGATTATGTTGTAAAATCATTTTCTATCCTAGATGGTC
TATCGGAAACAATTTATTTATTACTCCTATCCAATTCATTATATTTTCAAAGTTATGAAGTCCACGAAATATGT
GACGTGGGTAAAGAAGACCCATGCCAAGCCAGTGGGATATAGAAACAAAACATGTAATAAAGAGAACAATAATG
AGTTTCGAAAAGAACAGAAGTTAGCATAAGGACGAGAATCACATTATCTTAGGTGCCAACCACTAATCCTATGTA
TCATTCTCCTCTTTCCACGTGTCATCCTACACTTCCTTTGCCATCAGATTAGATAGCCCGGTTAGTACCTACACT
GTATATCAAAAAATACGTAACAATCATCAAACATATCATCGATCAAAGGATATTTATCTTGATGTGCTTTCCGCC
GTCCATTGTAACGAGTTTGGATGAATTTGATATACACCCACTCAGATATCAATATATTTTATAAAAAGAAACAAA
ATTGAATACTAGTAATATCTATGTAGATATTTATTTTTTCAACAATCCTGTAAGTTATAAGGATAACTCACTTAT
ATGTGACGTGGATAATGAAGAGCTAGGCAGGCAGTGAGAGATAGAAACAAATTAAGCAGAGACGAAAAACAATC
AGTTAACAGAATGACGAATTTGGATCAGCTTTATCTTAGTGCCAACCACTGATCCCATGCATCACTCTGCTCTTT
CCACGTGGCATCCTCTGACGTCAGATCAGATTCCTCTTCTTTCTTTTTTTTTTCTGTATATATGAGCATTMTA
GTAGT

Fig. 2






Fig. 3A

Fig. 3B




6/12

Fig. 4



7/12

Fig. 5



8/12

Fig. 6



A



C




B



D

9/12



1: Source leaf

2: Sink leaf

3: Stalk

4: Root

5: Seed

Fig. 7


10/12



Fig. 8

11/12


Fig. 9




12/12

aatcagtta acagaatgac gaattggatc acgctttatc ttagtgccaa ccactgatcc 60
 catgcatcac tctgctcttt ccacgtggca tectctgacg tcagatcaga ttcctcttct 120
 tctctttttt tttctgtata tataatgagca ttttagtagt actgctgccc caatctctta 180
 cataaaaatc gaagcacgat ggatcacgcy gggatcgcac accggacgga tttgatgaca 240
 ataacaaggt ttgtgttgaa tgagcagacy aagcaccctg aatcccgtgg agacttcagt 300
 attttgctca gtcacattgt tcttgctgc aagttcgtat gcactgctgt taacaaggca 360
 ggtttagcca aacttctagg acttgctggt gagactaatg tgcagggaga agatcaaaaag 420
 aaacttgatg tactctcaaa tgaagtgtt atcaaggctt tggtagcag taaccgaaca 480
 tgcattcttg tctctgaaga agatgaagaa gccacattg ttaggccagc taaccgtgga 540
 aaatactgty tagtttttga tccctggat ggatcacga acattgattg tgggttctct 600
 attggaacga tctttggaat ttacatgac aaagacggtc atgaaccaac actagatgat 660
 gtcttgcaac ctgggatgaa catgttagct gctggttact gcatgtatgg aagttcttgt 720
 acgctagttt tgagcactgg atctggagtt aatggtttta ccttgatcc ctctcttggc 780
 gagttcatcc taactcatcc tgacatcaag attcctaaga aaggaagat ttattcagtg 840
 aatgaaggaa atgccaaagaa ctgggacagt ccaacatcca aatatgtgca gagctgcaag 900
 tatcccgtg atggttcttc accaaaatct ttgagatata ttggaagtat ggttgctgat 960
 gttcatcgtc cattactcta tggaggcatc ttcttgacc cggagataa gaaaagcccc 1020
 aacgggaaac tgagggttct ctatgaagta tttcccattg catttctgat ggaacaagca 1080
 ggaggccaag catttactgg gaagcaacgg gcacttgact tagttccaga gaagatcac 1140
 gaacgctctc ctataattct tggtagttat gatgatgttg aggagatcaa aaagctctac 1200
 gctgctgaag agcaaaaactg atagatgtat ctataccatg taatcacttc actactcttg 1260
 ctggtgcaga tatcaaatct ctcaaatc agcaagtgt tactgtttat gttgcacaat 1320
 agctgctgty atgcgataat acgttcacat tactggttgt tctaactttt tgtcttgaag 1380
 tatctatttc tcatcaacaa taaatgttg aatagagaag ttctggctta ttattgttat 1440
 caaagtctt ttgtaatgct atccatttag aatcaagcta atttttt 1487

Fig. 10



 cy-FBPase Promoter