



(51) International Patent Classification:

C07D 235/30 (2006.01) C07D 235/14 (2006.01)
A61K 31/4184 (2006.01)

(21) International Application Number:

PCT/EP2013/051944

(22) International Filing Date:

31 January 2013 (31.01.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/593,459 1 February 2012 (01.02.2012) US
61/678,064 31 July 2012 (31.07.2012) US

(71) Applicant (for all designated States except CN): **EURO-CELTIQUE** [LU/LU]; 2, avenue Charles de Gaulle, L-1653 Luxembourg (LU).

(71) Applicant (for CN only): **CRYSTAL BIOPHARMA-CEUTICAL LLC** [US/US]; 341 Raven Circle, Wyoming, DE 19934 (US).

(72) Inventors: **CHEN, Yu**; 5078 Kingston Way, San Jose, CA 95130 (US). **CHEN, Yi**; 5662 Owens Drive #304, Pleasanton, CA 94588 (US).

(74) Agent: **MARKS & CLERK, LLP**; 62-68 Hills Road, Cambridge Cambridgeshire CB2 1LA (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— with international search report (Art. 21(3))

(54) Title: NOVEL THERAPEUTIC AGENTS



(57) Abstract: The present invention relates to a class of hydroxamic acid compounds of Formula (I), which act as alkylating agents and/or inhibitors of the HDAC pathway, having potential utility in the treatment of a neoplastic disease and immune diseases.

Novel Therapeutic Agents

The present invention relates to a class of hydroxamic acid compounds, which act as alkylating agents and/or inhibitors of the HDAC pathway, to uses thereof, to processes for the preparation thereof and compositions comprising said compounds. These compounds have potential utility in a variety of therapeutic areas including the treatment of a neoplastic disease and immune diseases.


Cancer is one of the most life threatening diseases in which cells in a part of the body experience out-of-control growth. According to latest data from American Cancer Society, it is estimated to have 1.6 million new cases of cancer in USA in 2011. Cancer is the second leading cause of death in the United States (second only to heart disease) and will claim more than 570,000 lives in 2011. In fact, it is estimated that 50% of all men and 33% of all women living in the United States will develop some type of cancer in their lifetime. Therefore cancer constitutes a major public health burden and represents a significant cost in the United States. For decades, surgery, chemotherapy, and radiation were the established treatments for various cancers. Patients usually receive a combination of these treatments depending upon the type and extent of their disease. But the chemotherapy is most important option for cancer patient when the surgery treatment is impossible.

Bendamustine, a well known chemotherapy first synthesized in 1963, consists of an alkylating nitrogen mustard moiety and a purine-like benzimidazole moiety with a suggested purine-analog effect (Barman Balfour JA, et al, *Drugs* **2001**; 61: 631–640). Bendamustine has been shown to have substantial activity against low-grade lymphomas (Herold M, et al., *Blood*, **1999**; 94, Suppl 1: 262a), multiple myelomas (Poenisch W, et al., *Blood* **2000**; 96, Suppl 1: 759a), and several solid tumors (Kollmannsberger C, et al., *Anticancer Drugs* **2000**; 11: 535–539). It was also reported that bendamustine effectively induces apoptosis in lymphoma cells (Chow KU, et al., *Haematologica*, **2001**; 86: 485–493). It has received FDA approval for the treatment of chronic lymphocytic leukemia (CLL) and for treatment of indolent B-cell non-Hodgkin's lymphoma (NHL) that has progressed during or within six months of treatment with rituximab or a rituximab-containing regimen.

In recent years, histone deacetylases (HDAC) has emerged as an important disease target for cancer treatment [Minucci, S. et al., *Nat Rev Cancer* **2006**, 6, 38-51]. The human HDAC enzymes have 18 isoforms grouped into Class I-IV according to their sequence homology. Class I, II and IV, commonly referred to as the classical HDACs, are comprised of 11 family

members. Class III HDACs consists of 7 enzymes and they are distinct from other HDAC family members, therefore are given a unique term sirtuins. The inhibition of HDAC enzyme leads to histone acetylation which is associated with the remodelling of chromatin and plays a key role in the epigenetic regulation of gene expression. In addition, HDAC inhibitors have been shown to evoke the acetylation of many important non-histone proteins such as HSP90, alpha-tubulin, Ku-70, Bcl-6, importin, cortactin, p53, STAT1, E2F1, GATA-1 and NF-kB, which can alter many important signaling networks related to cancer treatment. The underlying mechanism of action of HDAC inhibitors includes the differentiation, cell cycle arrest, inhibition of DNA repair, induction of apoptosis, upregulation of tumor suppressors, down regulation of growth factors, oxidative stress and autophagy. In the last decade, a large number of structurally diverse HDAC inhibitors have been identified and at least 12 HDAC inhibitors are currently in human clinical trials for cancer treatments, including short-chain fatty acid (valproic acid), hydroxamates (SAHA, LBH589, PXD101, JNJ-26481585, ITF2357, CUDC-101), cyclic tetrapeptides (FK-228), benzamide (MS-275), and several other compounds (CHR-3996, 4SC-201, SB939). Among them, SAHA and FK-228 has been approved by the US FDA for the treatment of advanced cutaneous T-cell lymphoma.

WO 2010/085377 refers to a class of hydroxamic acid derivatives, which inhibit the HDAC pathway and have potential utility in the treatment of a neoplastic disease or an autoimmune disease. Among the compounds disclosed is NL-101 having the structure shown below:




The biological assay showed that NL-101 potently inhibits HDAC enzyme (HDAC1 IC₅₀ of 9 nM). NL-101 was sent to NCI (NSC# 751447) for NCI-60 cell line panel screening. The data showed that NL-101 is about × 25-100 fold more potent than Bendamustine in the NCI-60 cell lines that are representative of a variety of human cancer type.

There is a continuing need for further pharmaceuticals useful for the treatment of cancer and auto-immune diseases, preferably having advantages over existing therapies, such as improved potency or selectivity, or reduced toxicity.

The present invention relates to a class of hydroxamic acid derivatives, which act as

alkylating agents and/or inhibitors of the HDAC pathway. The single dual-functional small molecules of the invention may attack the cancer cells synergistically from two distinct directions simultaneously (DNA damaging and the inhibitions of the HDAC pathway). Thus, the compounds of the present invention may be useful in treating a patient having a tumor, such as one treatable by Bendamustine and/or the inhibitors of HDAC pathway. The compounds of the invention may additionally be useful in the prevention and treatment of an immune disease.


Thus, in one aspect, this invention relates to a compound of Formula (I) or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof:



wherein

Z is $(\text{CR}_a\text{R}_b)_p\text{N}(\text{R}_a)(\text{CR}_a\text{R}_b)_q$;

X_1 and X_2 are each independently selected from halo and OSO_2R_c ;



Q is heteroaryl, which is optionally substituted with alkyl, alkenyl, alkynyl, cycloalkyl, halo, nitro, oxo, cyano or OR_e ;

R_a , R_b , R_d and R_e are each independently selected from H, alkyl, alkenyl and alkynyl;

R_c is selected from alkyl, alkenyl and alkynyl; and

p and q are each independently selected from 0, 1, 2, 3 and 4;

a tautomer thereof or a pharmaceutically acceptable salt, solvate or polymorph of said compound or tautomer.

Preferably, p and q are each independently selected from 1, 2, and 3. More preferably, p is 1 and q is 2; or p is 2 and q is 1; or p is 0 and q is 3; or p is 3 and q is 0; or p and q are both 2.

Preferably, Z is $(\text{CH}_2)_p\text{NH}(\text{CH}_2)_q$. Most preferably, Z is $(\text{CH}_2)_2\text{NH}(\text{CH}_2)$.


Preferably, X_1 and X_2 are each independently selected from halo. More preferably, X_1 and X_2 are each independently selected from chloro, bromo and iodo. Most preferably, X_1 and X_2 are both chloro.

Preferably, Q is an optionally substituted 9-10 membered heteroaryl. More preferably, Q is an optionally substituted benzimidazolyl. Yet more preferably, Q is benzimidazolyl



substituted by one or more alkyl groups. Even more preferably, Q is benzimidazolyl substituted by 1, 2, or 3 methyl groups. Most preferably, Q is benzimidazolyl substituted by a methyl group.

In a preferred embodiment, the compounds of the invention are represented by

5 Formula (II):



In more preferred embodiments, the compounds of the invention are represented by Formula (III) or Formula (III A):









10

In a more preferred embodiment, the compounds of the invention are represented by Formula (III) or Formula (III A) wherein X_1 and X_2 are each independently selected from halo, and Z is $(CH_2)_pNH(CH_2)_q$.

In a yet more preferred embodiment, the compounds of the invention are represented by

15 Formula (III) or Formula (III A) wherein X_1 and X_2 are both chloro, and Z is $(CH_2)_2NH(CH_2)_2$.


The following compounds are preferred:

 CY-102		
		

The alkene group in the compounds of Formula (I) may be in the form of either the (E) or (Z)-isomer, and are preferably the (E) isomer. In particular, the most preferred compound CY-102 is the (E)-isomer.

5 It should be recognized that the compounds of the present invention may be present and optionally administered in the form of salts or solvates. The invention encompasses any pharmaceutically acceptable salts and solvates of any one of the above-described compounds and modifications thereof.

The most preferred compound is the compound CY-102 or a pharmaceutically
10 acceptable salt, solvate or polymorph thereof:



Also within the scope of this invention is a pharmaceutical composition containing one
15 or more of the compounds, modifications, and/or salts or solvates thereof described above for use in treating a neoplastic disease, or an immune disorder, therapeutic uses thereof, and use of the compounds for the manufacture of a medicament for treating the disease / disorder.

This invention also relates to a method of treating a neoplastic disorder (*e.g.*, cancer, myelodysplastic syndrome, or myeloproliferative disease) by administering to a subject in need
20 thereof an effective amount of one or more of the compounds, modifications, and/or salts or solvates, and compositions thereof described above.

Furthermore, this invention relates to a method of treating an immune disease (*e.g.*, rheumatoid arthritis and multiple sclerosis) by administering to a subject in need thereof an effective amount of one or more of the compounds, modifications, and/or salts or solvates, and
25 compositions thereof described above.

The details of one or more embodiments of the invention are set forth in the description

below. Other features, objects, and advantages of the invention will be apparent from the description and from the claims. It should be understood that all embodiments / features of the invention (compounds, pharmaceutical compositions, methods of make / use, *etc*) described herein, including any specific features described in the examples and original claims, can
5 combine with one another unless not applicable or explicitly disclaimed.

Compounds of the invention may contain one or more asymmetric carbon atoms. Accordingly, the compounds may exist as diastereomers, enantiomers or mixtures thereof. The syntheses of the compounds may employ racemates, diastereomers or enantiomers as starting materials or as intermediates. Diastereomeric compounds may be separated by chromatographic
10 or crystallization methods. Similarly, enantiomeric mixtures may be separated using the same techniques or others known in the art. Each of the asymmetric carbon atoms may be in the R or S configuration and both of these configurations are within the scope of the invention.

A modified compound of any one of such compounds including a modification having an improved (*e.g.*, enhanced, greater) pharmaceutical solubility, stability, bioavailability and/or
15 therapeutic index as compared to the unmodified compound is also contemplated. The examples of modifications include but not limited to the prodrug derivatives, the deuterium-enriched compounds, and compound conjugates with polyethylene glycol, dextran, polyvinyl alcohol, carbohydrate polymer, antibody, small biomolecule such as Vitamin E or its derivatives, or mixtures thereof. For example:

- 20 • Prodrug derivatives: prodrugs, upon administration to a subject, are converted *in vivo* into active compounds of the present invention [*Nature Reviews of Drug Discovery*, **2008**, Volume 7, p255]. It is noted that in many instances, the prodrugs themselves also fall within the scope of the range of compounds according to the present invention. The prodrugs of the compounds of the present invention can be prepared by standard organic
25 reaction, for example, by reacting with a carbamylating agent (*e.g.*, 1,1-acyloxyalkylcarbonochloridate, para-nitrophenyl carbonate, or the like) or an acylating agent. Further examples of methods and strategies of making prodrugs are described in *Bioorganic and Medicinal Chemistry Letters*, **1994**, Vol. 4, p. 1985.
- 30 • Deuterium-enriched compounds: deuterium (D or ^2H) is a stable, non-radioactive isotope of hydrogen and has an atomic weight of 2.0144. Hydrogen naturally occurs as a mixture of the isotopes ^1H (hydrogen or protium), D (^2H or deuterium), and T (^3H or tritium). The natural abundance of deuterium is 0.015%. One of ordinary skill in the art recognizes that in all chemical compounds with a H atom, the H atom actually represents a mixture of H and D, with about 0.015% being D. Thus, compounds with a level of

deuterium that has been enriched to be greater than its natural abundance of 0.015%, should be considered unnatural and, as a result, novel over their nonenriched counterparts.

- Compound-polymer conjugates: Many anti-cancer agents exhibit excellent antitumor activity against *in vivo* animal xenografts. However, their water insolubility makes it difficult to administer these drugs. One approach to overcome the pharmaceutical and pharmacokinetic shortcomings of these poor soluble drugs is to covalently bind them to polymers such as polyethylene glycol, dextran, polyvinyl alcohol, and carbohydrate polymers. Using this approach, the water solubility of the anticancer agent can be improved such that the polymeric conjugate can be parenterally administered in aqueous medium.
- Compound-antibody conjugates: For many years it has been an aim of scientists in the field of specifically targeted drug therapy to use monoclonal antibodies (MAbs) for the specific delivery of toxic agents to human cancers. Conjugates of tumor-associated MAbs and suitable toxic agents have been developed. The toxic agent is most commonly a chemotherapy drug, although particle-emitting radionuclides, or bacterial or plant toxins have also been conjugated to MAbs, especially for the therapy of cancer (Sharkey and Goldenberg, CA Cancer J. Clin. **2006** July-August; 56(4):226-243). The advantages of using MAb-chemotherapy drug conjugates are that (a) the chemotherapy drug itself is structurally well defined; (b) the chemotherapy drug is linked to the MAb protein using very well defined conjugation chemistries, often at specific sites remote from the MAbs antigen binding regions; (c) MAb-chemotherapy drug conjugates can be made more reproducibly than chemical conjugates involving MAbs and bacterial or plant toxins, and as such are more amenable to commercial development and regulatory approval; and (d) the MAb-chemotherapy drug conjugates are orders of magnitude less toxic systemically than radionuclide MAb conjugates.

When the compounds of the present invention possess a free base form, the compounds can be prepared as a pharmaceutically acceptable acid addition salt by reacting the free base form of the compound with a pharmaceutically acceptable inorganic or organic acid, *e.g.*, hydrohalides such as hydrochloride, hydrobromide, hydroiodide; other mineral acids such as sulfate, nitrate, phosphate, *etc.*; and alkyl and monoarylsulfonates such as ethanesulfonate, toluenesulfonate and benzenesulfonate; and other organic acids and their corresponding salts such as acetate, tartrate, maleate, succinate, citrate, benzoate, salicylate and ascorbate. Further

acid addition salts of the present invention include, but are not limited to: adipate, alginate, arginate, aspartate, bisulfate, bisulfite, bromide, butyrate, camphorate, camphorsulfonate, caprylate, chloride, chlorobenzoate, cyclopentanepropionate, digluconate, dihydrogenphosphate, dinitrobenzoate, dodecylsulfate, fumarate, galactate (from mucic acid), galacturonate, glucoheptaoate, gluconate, glutamate, glycerophosphate, hemisuccinate, hemisulfate, heptanoate, hexanoate, hippurate, 2-hydroxyethanesulfonate, iodide, isethionate, iso-butyrate, lactate, lactobionate, malonate, mandelate, metaphosphate, methanesulfonate, methylbenzoate, monohydrogenphosphate, 2-naphthalenesulfonate, nicotinate, oxalate, oleate, pamoate, pectinate, persulfate, phenylacetate, 3-phenylpropionate, phosphonate and phthalate.

When the compounds of the present invention possess a free acid form, a pharmaceutically acceptable base addition salt can be prepared by reacting the free acid form of the compound with a pharmaceutically acceptable inorganic or organic base. Examples of such bases are alkali metal hydroxides including potassium, sodium and lithium hydroxides; alkaline earth metal hydroxides such as barium and calcium hydroxides; alkali metal alkoxides, *e.g.*, potassium ethanolate and sodium propanolate; and various organic bases such as ammonium hydroxide, piperidine, diethanolamine and N-methylglutamine. Also included are the aluminum salts of the compounds of the present invention. Further base salts of the present invention include, but are not limited to: copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium and zinc salts. Organic base salts include, but are not limited to, salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, *e.g.*, arginine, betaine, caffeine, chlorprocaine, choline, N,N'-dibenzylethylenediamine (benzathine), dicyclohexylamine, diethanolamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, iso-propylamine, lidocaine, lysine, meglumine, N-methyl-D-glucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethanolamine, triethylamine, trimethylamine, tripropylamine and tris-(hydroxymethyl)-methylamine (tromethamine). It should be recognized that the free acid forms will typically differ from their respective salt forms somewhat in physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid forms for the purposes of the present invention.

In one aspect, a pharmaceutically acceptable salt is a hydrochloride salt, hydrobromide salt, methanesulfonate, toluenesulfonate, acetate, fumarate, sulfate, bisulfate, succinate, citrate, phosphate, maleate, nitrate, tartrate, benzoate, biocarbonate, carbonate, sodium hydroxide salt,

calcium hydroxide salt, potassium hydroxide salt, tromethamine salt, or mixtures thereof.

The compound CY-102 is preferably formed and/or used as the hydrochloride salt.

Compounds of the present invention that comprise tertiary nitrogen-containing groups may be quaternized with such agents as (C₁₋₄) alkyl halides, *e.g.*, methyl, ethyl, iso-propyl and
5 tert-butyl chlorides, bromides and iodides; di-(C₁₋₄) alkyl sulfates, *e.g.*, dimethyl, diethyl and
diamyl sulfates; alkyl halides, *e.g.*, decyl, dodecyl, lauryl, myristyl and stearyl chlorides,
bromides and iodides; and aryl (C₁₋₄) alkyl halides, *e.g.*, benzyl chloride and phenethyl bromide.
Such salts permit the preparation of both water- and oil-soluble compounds of the invention.

Amine oxides, also known as amine-*N*-oxide and *N*-oxide, of anti-cancer agents with
10 tertiary nitrogen atoms have been developed as prodrugs [Mol Cancer Therapy. **2004** Mar;
3(3):233-44]. Compounds of the present invention that comprise tertiary nitrogen atoms may be
oxidized by such agents as hydrogen peroxide (H₂O₂), Caro's acid or peracids like *meta*-
Chloroperoxybenzoic acid (mCPBA) to form amine oxide.

The compound CY-102 may, for example, be used in the form of its *N*-oxide or a salt
15 thereof.

The invention encompasses pharmaceutical compositions comprising the compound of
the present invention and pharmaceutical excipients, as well as other conventional
pharmaceutically inactive agents. Any inert excipient that is commonly used as a carrier or
diluent may be used in compositions of the present invention, such as sugars, polyalcohols,
20 soluble polymers, salts and lipids. Sugars and polyalcohols which may be employed include,
without limitation, lactose, sucrose, mannitol, and sorbitol. Illustrative of the soluble polymers
which may be employed are polyoxyethylene, poloxamers, polyvinylpyrrolidone, and dextran.
Useful salts include, without limitation, sodium chloride, magnesium chloride, and calcium
chloride. Lipids which may be employed include, without limitation, fatty acids, glycerol fatty
25 acid esters, glycolipids, and phospholipids.

In addition, the pharmaceutical compositions may further comprise binders (*e.g.*, acacia,
cornstarch, gelatin, carbomer, ethyl cellulose, guar gum, hydroxypropyl cellulose,
hydroxypropyl methyl cellulose, povidone), disintegrating agents (*e.g.*, cornstarch, potato starch,
alginic acid, silicon dioxide, croscarmellose sodium, crospovidone, guar gum, sodium starch
30 glycolate, Primogel), buffers (*e.g.*, tris-HCL, acetate, phosphate) of various pH and ionic
strength, additives such as albumin or gelatin to prevent absorption to surfaces, detergents (*e.g.*,
Tween 20, Tween 80, Pluronic F68, bile acid salts), protease inhibitors, surfactants (*e.g.*, sodium
lauryl sulfate), permeation enhancers, solubilizing agents (*e.g.*, glycerol, polyethylene glycerol,
cyclodextrins), a glidant (*e.g.*, colloidal silicon dioxide), anti-oxidants (*e.g.*, ascorbic acid,

sodium metabisulfite, butylated hydroxyanisole), stabilizers (*e.g.*, hydroxypropyl cellulose, hydroxypropylmethyl cellulose), viscosity increasing agents (*e.g.*, carbomer, colloidal silicon dioxide, ethyl cellulose, guar gum), sweeteners (*e.g.*, sucrose, aspartame, citric acid), flavoring agents (*e.g.*, peppermint, methyl salicylate, or orange flavoring), preservatives (*e.g.*, Thimerosal, benzyl alcohol, parabens), lubricants (*e.g.*, stearic acid, magnesium stearate, polyethylene glycol, sodium lauryl sulfate), flow-aids (*e.g.*, colloidal silicon dioxide), plasticizers (*e.g.*, diethyl phthalate, triethyl citrate), emulsifiers (*e.g.*, carbomer, hydroxypropyl cellulose, sodium lauryl sulfate), polymer coatings (*e.g.*, poloxamers or poloxamines), coating and film forming agents (*e.g.*, ethyl cellulose, acrylates, polymethacrylates) and/or adjuvants.

In one embodiment, the pharmaceutical compositions are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

Additionally, the invention encompasses pharmaceutical compositions comprising any solid or liquid physical form of the compound of the invention. For example, the compounds can be in a crystalline form, in amorphous form, and have any particle size. The particles may be micronized, or may be agglomerated, particulate granules, powders, oils, oily suspensions or any other form of solid or liquid physical form.

DEFINITIONS:

The term “alkyl” refers to a straight or branched hydrocarbon containing 1-20 carbon atoms (*e.g.*, C₁-C₁₀). Examples of alkyl include, but are not limited to, methyl, methylene, ethyl, ethylene, n-propyl, i-propyl, n-butyl, i-butyl, and t-butyl. Preferably, the alkyl group has one to ten carbon atoms. More preferably, the alkyl group has one to four carbon atoms.

The term “alkenyl” refers to a straight or branched hydrocarbon containing 2-20 carbon atoms (*e.g.*, C₂-C₁₀) and one or more double bonds. Examples of alkenyl include, but are not limited to, ethenyl, propenyl, and allyl. Preferably, the alkylene group has two to ten carbon atoms. More preferably, the alkylene group has two to four carbon atoms.

The term “alkynyl” refers to a straight or branched hydrocarbon containing 2-20 carbon

atoms (*e.g.*, C₂-C₁₀) and one or more triple bonds. Examples of alkynyl include, but are not limited to, ethynyl, 1-propynyl, 1- and 2-butynyl, and 1-methyl-2-butynyl. Preferably, the alkynyl group has two to ten carbon atoms. More preferably, the alkynyl group has two to four carbon atoms.

5 The term “cycloalkyl” refers to a saturated hydrocarbon ring system having 3 to 30 carbon atoms (*e.g.*, C₃-C₁₂, C₃-C₈, C₃-C₆). Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

 The term “heteroaryl” refers to an aromatic 5-8 membered monocyclic, or an 8-12 membered bicyclic ring system having one or more heteroatoms selected from O, N, S, P and
10 Se). Examples of heteroaryl groups include pyridyl, furyl, imidazolyl, benzimidazolyl, pyrimidinyl, thienyl, quinolinyl, indolyl, and thiazolyl.

 “Halo” means fluoro, chloro, bromo or iodo.

 “Protected derivatives” means derivatives of compounds in which a reactive site are blocked with protecting groups. Protected derivatives are useful in the preparation of
15 pharmaceuticals or in themselves may be active as inhibitors. A comprehensive list of suitable protecting groups can be found in T.W.Greene, Protecting Groups in Organic Synthesis, 3rd edition, Wiley & Sons, 1999.

 “Pharmaceutically acceptable carrier” means a non-toxic solvent, dispersant, excipient, adjuvant, or other material which is mixed with the compounds of the present invention in order
20 to form a pharmaceutical composition, *i.e.*, a dose form capable of administration to the patient. Examples of pharmaceutically acceptable carrier includes suitable polyethylene glycol (*e.g.*, PEG400), surfactant (*e.g.*, Cremophor), or cyclopolysaccharide (*e.g.*, hydroxypropyl- β -cyclodextrin or sulfobutyl ether β -cyclodextrins), polymer, liposome, micelle, nanosphere, *etc.*

 “Therapeutically effective amount” of a composition described herein is meant an
25 amount of the composition which confers a therapeutic effect on the treated subject, at a reasonable benefit/risk ratio applicable to any medical treatment. The therapeutic effect may be objective (*i.e.*, measurable by some test or marker) or subjective (*i.e.*, subject gives an indication of or feels an effect). An effective amount of the composition described above may range from about 0.1 mg/kg to about 500 mg/kg, preferably from about 0.2 to about 50 mg/kg. Effective
30 doses will also vary depending on route of administration, as well as the possibility of co-usage with other agents. It will be understood, however, that the total daily usage of the compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the

disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or contemporaneously with the specific compound employed; and like factors well known in the medical arts.

As used herein, the term “treating” refers to administering a compound to a subject that has, for example, a neoplastic or immune disorder, or has a symptom of or a predisposition toward it, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disorder, the symptoms of or the predisposition toward the disorder. The term “an effective amount” refers to the amount of the active agent that is required to confer the intended therapeutic effect in the subject. Effective amounts may vary, as recognized by those skilled in the art, depending on route of administration, excipient usage, and the possibility of co-usage with other agents.

A “subject” refers to a human and a non-human animal. Examples of a non-human animal include all vertebrates, *e.g.*, mammals, such as non-human primates (particularly higher primates), dog, rodent (*e.g.*, mouse or rat), guinea pig, cat, and non-mammals, such as birds, amphibians, reptiles, *etc.* In a preferred embodiment, the subject is a human. In another embodiment, the subject is an experimental animal or animal suitable as a disease model.

When compounds according to the present invention exhibit insufficient solubility, methods for solubilizing the compounds may be used. Such methods are known to those of skill in this art, and include, but are not limited to, pH adjustment and salt formation, using co-solvents, such as ethanol, propylene glycol, polyethylene glycol (PEG) 300, PEG 400, DMA (10-30%), DMSO (10-20%), NMP (10-20%), using surfactants, such as polysorbate 80, polysorbate 20 (1-10%), cremophor EL, Cremophor RH40, Cremophor RH60 (5-10%), Pluronic F68/Poloxamer 188 (20-50%), Solutol HS15 (20-50%), Vitamin E TPGS, and d- α -tocopheryl PEG 1000 succinate (20-50%), using complexation such as HP β CD and SBE β CD (10-40%), and using advanced approaches such as micelle, addition of a polymer, nanoparticle suspensions, and liposome formation.

“Combination therapy” includes the administration of the subject compounds of the present invention in further combination with other biologically active ingredients (such as, but not limited to, a second and different antineoplastic agent) and non-drug therapies (such as, but not limited to, surgery or radiation treatment). For instance, the compounds of the invention can be used in combination with other pharmaceutically active compounds, or non-drug therapies,

preferably compounds that are able to enhance the effect of the compounds of the invention. The compounds of the invention can be administered simultaneously (as a single preparation or separate preparation) or sequentially to the other therapies. In general, a combination therapy envisions administration of two or more drugs/treatments during a single cycle or course of therapy.

In one embodiment, the compounds of the invention are administered in combination with one or more of traditional chemotherapeutic agents. The traditional chemotherapeutic agents encompass a wide range of therapeutic treatments in the field of oncology. These agents are administered at various stages of the disease for the purposes of shrinking tumors, destroying remaining cancer cells left over after surgery, inducing remission, maintaining remission and/or alleviating symptoms relating to the cancer or its treatment. Examples of such agents include, but are not limited to, alkylating agents such as Nitrosureas (*e.g.*, Carmustine, Lomustine and Streptozocin), ethylenimines (*e.g.*, thiotepa, hexamethylmelanine), Alkylsulfonates (*e.g.*, Busulfan), Hydrazines and Triazines (*e.g.*, Altretamine, Procarbazine, Dacarbazine and Temozolomide), and platinum based agents (*e.g.*, Carboplatin, Cisplatin, and Oxaliplatin); plant alkaloids such as Podophyllotoxins (*e.g.*, Etoposide and Teniposide), Taxanes (*e.g.*, Paclitaxel and Docetaxel), Vinca alkaloids (*e.g.*, Vincristine, Vinblastine and Vinorelbine); anti-tumor antibiotics such as Chromomycins (*e.g.*, Dactinomycin and Plicamycin), Anthracyclines (*e.g.*, Doxorubicin, Daunorubicin, Epirubicin, Mitoxantrone, and Idarubicin), and miscellaneous antibiotics such as Mitomycin and Bleomycin; anti-metabolites such as folic acid antagonists (*e.g.*, Methotrexate), pyrimidine antagonists (*e.g.*, 5-Fluorouracil, Fluorouridine, Cytarabine, Capecitabine, and Gemcitabine), purine antagonists (*e.g.*, 6-Mercaptopurine and 6-Thioguanine) and adenosine deaminase inhibitors (*e.g.*, Cladribine, Fludarabine, Nelarabine and Pentostatin); topoisomerase inhibitors such as topoisomerase I inhibitors (Topotecan, Irinotecan), topoisomerase II inhibitors (*e.g.*, Amsacrine, Etoposide, Etoposide phosphate, Teniposide), and miscellaneous anti-neoplastics such as ribonucleotide reductase inhibitors (Hydroxyurea), adrenocortical steroid inhibitor (Mitotane), anti-microtubule agents (Estramustine), and retinoids (Bexarotene, Isotretinoin, Tretinoin (ATRA)).

In one aspect of the invention, the compounds may be administered in combination with one or more targeted anti-cancer agents that modulate protein kinases involved in various disease states. Examples of such kinases may include, but are not limited ABL1, ABL2/ARG, ACK1, AKT1, AKT2, AKT3, ALK, ALK1/ACVRL1, ALK2/ACVR1, ALK4/ACVR1B, ALK5/TGFBR1, ALK6/BMPR1B, AMPK(A1/B1/G1), AMPK(A1/B1/G2), AMPK(A1/B1/G3), AMPK(A1/B2/G1), AMPK(A2/B1/G1), AMPK(A2/B2/G1), AMPK(A2/B2/G2), ARAF,

ARK5/NUAK1, ASK1/MAP3K5, ATM, Aurora A, Aurora B , Aurora C , AXL, BLK, BMPR2,
 BMX/ETK, BRAF, BRK, BRSK1, BRSK2, BTK, CAMK1a , CAMK1b, CAMK1d, CAMK1g ,
 CAMKIIa , CAMKIIb , CAMKIId , CAMKIIg , CAMK4, CAMKK1, CAMKK2, CDC7-DBF4,
 CDK1-cyclin A, CDK1-cyclin B, CDK1-cyclin E, CDK2-cyclin A, CDK2-cyclin A1, CDK2-
 5 cyclin E, CDK3-cyclin E, CDK4-cyclin D1, CDK4-cyclin D3, CDK5-p25, CDK5-p35, CDK6-
 cyclin D1, CDK6-cyclin D3, CDK7-cyclin H, CDK9-cyclin K, CDK9-cyclin T1, CHK1, CHK2,
 CK1a1 , CK1d , CK1epsilon , CK1g1 , CK1g2, CK1g3 , CK2a , CK2a2, c-KIT, CLK1 , CLK2,
 CLK3, CLK4, c-MER, c-MET, COT1/MAP3K8, CSK, c-SRC, CTK/MATK, DAPK1, DAPK2,
 DCAMKL1, DCAMKL2, DDR1, DDR2, DLK/MAP3K12, DMPK, DMPK2/CDC42BPG,
 10 DNA-PK, DRAK1/STK17A, DYRK1/DYRK1A, DYRK1B, DYRK2, DYRK3, DYRK4,
 EEF2K, EGFR, EIF2AK1, EIF2AK2, EIF2AK3, EIF2AK4/GCN2, EPHA1, EPHA2, EPHA3,
 EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHB1, EPHB2, EPHB3, EPHB4, ERBB2/HER2,
 ERBB4/HER4, ERK1/MAPK3, ERK2/MAPK1, ERK5/MAPK7, FAK/PTK2, FER, FES/FPS,
 FGFR1, FGFR2, FGFR3, FGFR4, FGR, FLT1/VEGFR1, FLT3, FLT4/VEGFR3, FMS,
 15 FRK/PTK5, FYN, GCK/MAP4K2, GRK1, GRK2, GRK3, GRK4, GRK5, GRK6, GRK7,
 GSK3a, GSK3b, Haspin, HCK, HGK/MAP4K4, HIPK1, HIPK2, HIPK3, HIPK4,
 HPK1/MAP4K1, IGF1R, IKKa/CHUK , IKKb/IKBKB, IKKe/IKBKE, IR, IRAK1, IRAK4,
 IRR/INSRR, ITK, JAK1, JAK2, JAK3, JNK1 , JNK2 , JNK3, KDR/VEGFR2, KHS/MAP4K5,
 LATS1, LATS2, LCK, LCK2/ICK, LKB1 , LIMK1, LOK/STK10, LRRK2, LYN, LYNB,
 20 MAPKAPK2, MAPKAPK3, MAPKAPK5/PRAK, MARK1, MARK2/PAR-1Ba, MARK3,
 MARK4, MEK1, MEK2, MEKK1, MEKK2, MEKK3, MELK, MINK/MINK1, MKK4, MKK6,
 MLCK/MYLK, MLCK2/MYLK2, MLK1/MAP3K9, MLK2/MAP3K10, MLK3/MAP3K11,
 MNK1, MNK2, MRCKa/, CDC42BPA, MRCKb/, CDC42BPB, MSK1/RPS6KA5,
 MSK2/RPS6KA4, MSSK1/STK23, MST1/STK4, MST2/STK3, MST3/STK24, MST4,
 25 mTOR/FRAP1, MUSK, MYLK3, MYO3b, NEK1, NEK2, NEK3, NEK4, NEK6, NEK7,
 NEK9, NEK11, NIK/MAP3K14, NLK, OSR1/OXSR1, P38a/MAPK14, P38b/MAPK11,
 P38d/MAPK13 , P38g/MAPK12 , P70S6K/RPS6KB1, p70S6Kb/, RPS6KB2, PAK1, PAK2,
 PAK3, PAK4, PAK5, PAK6, PASK, PBK/TOPK, PDGFRa, PDGFRb, PDK1/PDPK1,
 PDK1/PDHK1, PDK2/PDHK2 , PDK3/PDHK3, PDK4/PDHK4, PHKg1 , PHKg2 , PI3Ka,
 30 (p110a/p85a), PI3Kb, (p110b/p85a), PI3Kd, (p110d/p85a), PI3Kg(p120g), PIM1, PIM2, PIM3,
 PKA, PKAcb, PKAcb , PKCa , PKCb1 , PKCb2 , PKCd , PKCepsilon, PKCeta, PKCg ,
 PKCiota, PKCmu/PRKD1, PKCnu/PRKD3, PKCtheta, PKCzeta, PKD2/PRKD2, PKG1a ,
 PKG1b , PKG2/PRKG2, PKN1/PRK1, PKN2/PRK2, PKN3/PRK3, PLK1, PLK2, PLK3,
 PLK4/SAK, PRKX, PYK2, RAF1, RET, RIPK2, RIPK3, RIPK5, ROCK1, ROCK2,

RON/MST1R, ROS/ROS1, RSK1, RSK2, RSK3, RSK4, SGK1, SGK2, SGK3/SGKL, SIK1, SIK2, SLK/STK2, SNARK/NUAK2, SRMS, SSK/TSSK6, STK16, STK22D/TSSK1, STK25/YSK1, STK32b/YANK2, STK32c/YANK3, STK33, STK38/NDR1, STK38L/NDR2, STK39/STLK3, SRPK1, SRPK2, SYK, TAK1, TAOK1, TAOK2/TAO1, TAOK3/JIK, TBK1, 5 TEC, TESK1, TGFBR2, TIE2/TEK, TLK1, TLK2, TNIK, TNK1, TRKA, TRKB, TRKC, TRPM7/CHAK1, TSSK2, TSSK3/STK22C, TTBK1, TTBK2, TTK, TXK, TYK1/LTK, TYK2, TYRO3/SKY, ULK1, ULK2, ULK3, VRK1, VRK2, WEE1, WNK1, WNK2, WNK3, YES/YES1, ZAK/MLTK, ZAP70, ZIPK/DAPK3, KINASE, MUTANTS, ABL1(E255K), ABL1(F317I), ABL1(G250E), ABL1(H396P), ABL1(M351T), ABL1(Q252H), ABL1(T315I), 10 ABL1(Y253F), ALK (C1156Y), ALK(L1196M), ALK (F1174L), ALK (R1275Q), BRAF(V599E), BTK(E41K), CHK2(I157T), c-Kit(A829P), c-KIT(D816H), c-KIT(D816V), c-Kit(D820E), c-Kit(N822K), C-Kit (T670I), c-Kit(V559D), c-Kit(V559D/V654A), c-Kit(V559D/T670I), C-Kit (V560G), c-KIT(V654A), C-MET(D1228H), C-MET(D1228N), C-MET(F1200I), c-MET(M1250T), C-MET(Y1230A), C-MET(Y1230C), C-MET(Y1230D), C- 15 MET(Y1230H), c-Src(T341M), EGFR(G719C), EGFR(G719S), EGFR(L858R), EGFR(L861Q), EGFR(T790M), EGFR, (L858R,T790M) , EGFR(d746-750/T790M), EGFR(d746-750), EGFR(d747-749/A750P), EGFR(d747-752/P753S), EGFR(d752-759), FGFR1(V561M), FGFR2(N549H), FGFR3(G697C), FGFR3(K650E), FGFR3(K650M), FGFR4(N535K), FGFR4(V550E), FGFR4(V550L), FLT3(D835Y), FLT3(ITD), JAK2 20 (V617F), LRRK2 (G2019S), LRRK2 (I2020T), LRRK2 (R1441C), p38a(T106M), PDGFRa(D842V), PDGFRa(T674I), PDGFRa(V561D), RET(E762Q), RET(G691S), RET(M918T), RET(R749T), RET(R813Q), RET(V804L), RET(V804M), RET(Y791F), TIF2(R849W), TIF2(Y897S), and TIF2(Y1108F).

In another aspect of the invention, the subject compounds may be administered in 25 combination with one or more targeted anti-cancer agents that modulate non-kinase biological targets, pathway, or processes. Such targets pathways, or processes include but not limited to heat shock proteins (e.g. HSP90), poly-ADP (adenosine diphosphate)-ribose polymerase (PARP), hypoxia-inducible factors(HIF), proteasome, Wnt/Hedgehog/Notch signaling proteins, TNF-alpha, matrix metalloproteinase, farnesyl transferase, apoptosis pathway (e.g Bcl-xL, Bcl- 30 2, Bcl-w), histone deacetylases (HDAC), histone acetyltransferases (HAT), and methyltransferase (e.g histone lysine methyltransferases, histone arginine methyltransferase, DNA methyltransferase, etc).

In another aspect of the invention, the compounds of the invention are administered in combination with one or more of other anti-cancer agents that include, but are not limited to,

hormonal therapies (e.g. Tamoxifen, Fulvestrant, Clomifene, Anastrozole, Exemestane, Formestane, Letrozole, etc), vascular disrupting agent, gene therapy, RNAi cancer therapy, chemoprotective agents (e.g., amfostine, mesna, and dexrazoxane), antibody conjugate (e.g. brentuximab vedotin, ibritumomab tiuxetan), cancer immunotherapy such as Interleukin-2, cancer vaccines (e.g., sipuleucel-T) or monoclonal antibodies (e.g., Bevacizumab, Alemtuzumab, Rituximab, Trastuzumab, etc).

In another aspect of the invention, the subject compounds are administered in combination with radiation therapy or surgeries. Radiation is commonly delivered internally (implantation of radioactive material near cancer site) or externally from a machine that employs photon (x-ray or gamma-ray) or particle radiation. Where the combination therapy further comprises radiation treatment, the radiation treatment may be conducted at any suitable time so long as a beneficial effect from the co-action of the combination of the therapeutic agents and radiation treatment is achieved. For example, in appropriate cases, the beneficial effect is still achieved when the radiation treatment is temporally removed from the administration of the therapeutic agents, perhaps by days or even weeks.

In certain preferred embodiments, the compounds of the invention are administered in combination with one or more of radiation therapy, surgery, or anti-cancer agents that include, but are not limited to, DNA damaging agents, anti-metabolites, topoisomerase inhibitors, anti-microtubule agents, EGFR inhibitors, HER2 inhibitors, VEGFR2 inhibitors, BRAF inhibitors, Bcr-Abl inhibitors, PDGFR inhibitors, ALK inhibitors, PLK inhibitors, MET inhibitors, epigenetic agents, HSP90 inhibitors, PARP inhibitors, CHK inhibitors, aromatase inhibitor, estrogen receptor antagonist, and antibodies targeting VEGF, HER2, EGFR, CD50, CD20, CD30, CD33, etc.

In certain preferred embodiments, the compounds of the invention are administered in combination with one or more of abarelix, abiraterone acetate, aldesleukin, alemtuzumab, altretamine, anastrozole, asparaginase, bevacizumab, bexarotene, bicalutamide, bleomycin, bortezomib, brentuximab vedotin, busulfan, capecitabine, carboplatin, carmustine, cetuximab, chlorambucil, cisplatin, cladribine, clofarabine, clomifene, crizotinib, cyclophosphamide, dasatinib, daunorubicin liposomal, decitabine, degarelix, denileukin diftitox, denileukin diftitox, denosumab, docetaxel, doxorubicin, doxorubicin liposomal, epirubicin, eribulin mesylate, erlotinib, estramustine, etoposide phosphate, everolimus, exemestane, fludarabine, fluorouracil, fotemustine, fulvestrant, gefitinib, gemcitabine, gemtuzumab ozogamicin, goserelin acetate, histrelin acetate, hydroxyurea, ibritumomab tiuxetan, idarubicin, ifosfamide, imatinib mesylate, interferon alfa 2a, ipilimumab, ixabepilone, lapatinib ditosylate, lenalidomide, letrozole,

leucovorin, leuprolide acetate, levamisole, lomustine, mechlorethamine, melphalan, methotrexate, mitomycin C, mitoxantrone, nelarabine, nilotinib, oxaliplatin, paclitaxel, paclitaxel protein-bound particle, pamidronate, panitumumab, pegaspargase, peginterferon alfa-2b, pemetrexed disodium, pentostatin, raloxifene, rituximab, sorafenib, streptozocin, sunitinib maleate, tamoxifen, temsirolimus, teniposide, thalidomide, toremifene, tositumomab, trastuzumab, tretinoin, uramustine, vandetanib, vemurafenib, vinorelbine, zoledronate, radiation therapy, or surgery.

A wide variety of administration methods may be used in conjunction with the compounds of the present invention. Compounds of the present invention may be administered or coadministered orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery (for example by catheter or stent), subcutaneously, intraadiposally, intraarticularly, or intrathecally. The compounds according to the invention may also be administered or coadministered in slow release dosage forms.

Compounds may be in gaseous, liquid, semi-liquid or solid form, formulated in a manner suitable for the route of administration to be used. For oral administration, suitable solid oral formulations include tablets, capsules, pills, granules, pellets, sachets and effervescent, powders, and the like. Suitable liquid oral formulations include solutions, suspensions, dispersions, emulsions, oils and the like. For parenteral administration, reconstitution of a lyophilized powder is typically used. Tablets and iv infusion may be preferred.

The invention further provides methods for the prevention or treatment of a neoplastic disease or immune disease. In one embodiment, the invention relates to a method of treating a neoplastic disease or immune disease in a subject in need of treatment comprising administering to said subject a therapeutically effective amount of a compound of the invention. In one embodiment, the invention further provides for the use of a compound of the invention in the manufacture of a medicament for halting or decreasing a neoplastic disease or immune disease.

The neoplastic disease includes but not limited to lung cancer, head and neck cancer, central nervous system cancer, prostate cancer, testicular cancer, colorectal cancer, pancreatic cancer, liver cancer, stomach cancer, biliary tract cancer, esophageal cancer, gastrointestinal stromal tumor, breast cancer, cervical cancer, ovarian cancer, uterine cancer, leukemia, lymphomas, multiple myeloma, melanoma, basal cell carcinoma, squamous cell carcinoma, bladder cancer, renal cancer, sarcoma, mesothelioma, thymoma, myelodysplastic syndrome and myeloproliferative disease.

In certain embodiments, the neoplastic disease is a solid tumor. Representative treatable

solid tumors include melanoma, breast cancer, lung cancer (*e.g.*, small cell lung cancer (SCLC), or non-small cell lung cancer (NSCLC)), colon cancer, renal cancer, or sarcoma.kyl

In certain embodiments, the method may further include administering a second therapeutic agent known to be effective for treating the solid tumor.

5 For example, effective second therapeutic agent known to be effective for treating breast cancer includes: Methotrexate (Abitrexate, Folex, Folex PFS, Methotrexate LPF, Mexate-AQ); Paclitaxel (Taxol); Paclitaxel Albumin-stabilized Nanoparticle Formulation (Abraxane); Doxorubicin Hydrochloride (Adriamycin, Adriamycin PFS; Adriamycin RDF); Fluorouracil (Acrucil, Efudex, Fluoroplex); Everolimus (Afinitor); Anastrozole (Arimidex); Exemestane
10 (Aromasin); Capecitabine (Xeloda); Cyclophosphamide (Clafen, Cytosan, Neosar); Docetaxel (Taxotere); Epirubicin Hydrochloride (Ellence); Everolimus; Toremifene (Fareston); Fulvestrant (Faslodex); Letrozole (Femara); Gemcitabine Hydrochloride (Gemzar); Trastuzumab (Herceptin); Ixabepilone (Ixempra); Lapatinib Ditosylate; Tamoxifen Citrate (Nolvadex, Novaldex); Pertuzumab (Perjeta); Toremifene; Lapatinib Ditosylate (Tykerb); Doxorubicin
15 Hydrochloride & Cyclophosphamide; Doxorubicin Hydrochloride & Cyclophosphamide & Paclitaxel; Doxorubicin Hydrochloride & Cyclophosphamide & Fluorouracil; Cyclophosphamide & Methotrexate & Fluorouracil; Fluorouracil & Cyclophosphamide & Epirubicin Hydrochloride.

Effective second therapeutic agent known to be effective for treating small cell lung
20 cancer (SCLC) includes: Methotrexate (Abitrexate, Folex, Folex PFS, Methotrexate LPF, Mexate, Mexate-AQ); Etoposide (Toposar, VePesid); Etoposide Phosphate (Etopophos); Topotecan Hydrochloride (Hycamtin).

Effective second therapeutic agent known to be effective for treating non-small cell lung cancer (NSCLC) includes: Methotrexate (Abitrexate, Folex, Folex PFS, Methotrexate LPF,
25 Mexate, Mexate-AQ); Paclitaxel (Taxol); Paclitaxel Albumin-stabilized Nanoparticle Formulation (Abraxane); Pemetrexed Disodium (Alimta); Bevacizumab (Avastin); Carboplatin (Paraplat, Paraplatin); Cisplatin (Platinol, Platinol-AQ); Crizotinib (Xalkori); Erlotinib Hydrochloride; Gefitinib (Iressa); Gemcitabine Hydrochloride (Gemzar); Pemetrexed Disodium; Erlotinib Hydrochloride (Tarceva); Carboplatin & Paclitaxel; Gemcitabine Hydrochloride &
30 Cisplatin.

Other than the standard surgical treatment, effective second therapeutic agent known to be effective for treating melanoma includes: imiquimod (Zyclara, Aldara, Beselna, R-837); interferon (adjuvant therapy after surgery); Bacille Calmette-Guerin (BCG) vaccine; interleukin-2; Ipilimumab (Yervoy); Vemurafenib (Zelboraf); Dacarbazine (DTIC); Temozolomide

(Temodar); interferon & temozolomide; interferon, interleukin-2, and temozolomide; or isolated limb perfusion (ILF, infusing the limb with a heated solution of chemotherapy), depending on the specific stages of the melanoma at the time of diagnosis.

Effective second therapeutic agent known to be effective for treating colon cancer includes: Fluorouracil (Adrucil, Efudex, Fluoroplex); Bevacizumab (Avastin); Irinotecan Hydrochloride (Camptosar); Capecitabine (Xeloda); Cetuximab (Erbix); Oxaliplatin (Eloxatin); Leucovorin Calcium; Panitumumab (Vectibix); Regorafenib (Stivarga); Leucovorin Calcium (Wellcovorin); Ziv-Aflibercept (Zaltrap); Leucovorin Calcium & Fluorouracil & Irinotecan Hydrochloride; Leucovorin Calcium & Fluorouracil & Irinotecan Hydrochloride + Bevacizumab; Leucovorin Calcium (Folinic Acid) & Fluorouracil & Oxaliplatin; Capecitabine & Oxaliplatin.

Effective second therapeutic agent known to be effective for treating renal cancer includes: Fluorouracil (Adrucil, Efudex, Fluoroplex); Bevacizumab (Avastin); Irinotecan Hydrochloride (Camptosar); Cetuximab (Erbix); Panitumumab (Vectibix); Regorafenib (Stivarga); Ziv-Aflibercept (Zaltrap); Capecitabine & Oxaliplatin; Leucovorin Calcium (Folinic Acid) & Fluorouracil & Irinotecan Hydrochloride; Leucovorin Calcium & Fluorouracil & Irinotecan Hydrochloride + Bevacizumab; Leucovorin Calcium (Folinic Acid) & Fluorouracil & Oxaliplatin.

Preferred combinations of compounds of the invention, especially in combination with CY-102 or a pharmaceutically acceptable salt, solvate or polymorph thereof include combinations with:

Proteasome inhibitors (e.g. bortezomib, carfilzomib).

IMiDs (e.g. Thalidomide, lenalidomide, pomalidomide).

Platinum agents (e.g. cisplatin, carboplatin).

Folate antagonists (e.g. pemetrexed, pralatrexate).

CD30 antibodies and conjugates (e.g. brentuximab, vendotin).

Antibodies (also conjugated) to treat haematological malignancies like anti CD20 (e.g. ofatumumab, rituximab, GA101, etc).

B-cell receptor antagonists (e.g. ibrutinib).

PI3K antagonists (e.g. GS1101 or IPI145).

BTK inhibitors.

Taxanes (e.g. taxol, paclitaxel).

Antibodies (also conjugated) to treat ovarian cancer (e.g. alpha folate receptor mabs, CA125 antibodies).

Antibodies to treat multiple myeloma (e.g. elotuzumab, anti CD38 mabs).

Anthracyclines (e.g. doxorubicin, idarubicin).

Nucleoside analogues (purine antagonists) like cytarabine, fludarabine, gemcitabine.

PNP antagonists (e.g. forodesine).

5 Bcr-abl tyrosinekinase blockers (e.g. imatinib, dasatinib, ponatinib, nilotinib).

mTor antagonists (e.g. temsirolimus, everolimus).

Agents influencing the CD40 activation (e.g. CD40 antagonists, CD40 gene medicines).

Multi tyrosine kinase antagonists (e.g. sorafenib, axitinib).

10 Bifunctional antibodies (e.g. CD19/CD3, also conjugated, also recognising other CD epitopes).

Preferred combinations of compounds of the invention, especially in combination with CY-102 or a pharmaceutically acceptable salt, solvate or polymorph thereof include combinations with one or more, such as one, two or three, of the above-identified therapeutic agents.

15 Especially preferred combinations of compounds of the invention include combinations of CY-102 or a pharmaceutically acceptable salt, solvate or polymorph thereof and forodesine, optionally in combination with one or more, such as one, two or three, of the above-identified therapeutic agents.

20 The combinations of compounds of the invention include combinations of CY-102 or a pharmaceutically acceptable salt, solvate or polymorph thereof and forodesine, optionally in combination with one or more, such as one, two or three, of the above-identified therapeutic agents.

The treatment above may be in conjunction with other treatments, such as surgery, radiation therapy, laser therapy, stem cell transplant.

25 In a further aspect, the present invention is directed to combinations of one or more compounds of the invention with one or more additional therapeutic agents. In a further aspect, the present invention is directed to combinations of one or more compounds of the invention with one or more additional therapeutic agents for use as a medicament, and in particular, for use in the treatment of the diseases disclosed herein. In a further aspect, the present invention is
30 directed to the use of combinations of one or more compounds of the invention with one or more additional therapeutic agents in the treatment of the diseases disclosed herein. In preferred embodiments of all aspects of the invention, the disease to be treated is CLL.

In a further aspect, the present invention is directed to a kit comprising (a) a first pharmaceutical composition comprising one or more compounds of the present invention and (b)

a second pharmaceutical composition comprising one or more additional therapeutic agents as defined herein. In a further aspect, the present invention is directed to a kit comprising (a) a first pharmaceutical composition comprising one or more compounds of the present invention and (b) a second pharmaceutical composition comprising one or more additional therapeutic agents as defined herein for use as a medicament, and in particular, for use in the treatment of the diseases disclosed herein. In a further aspect, the present invention is directed to a kit comprising (a) a first pharmaceutical composition comprising one or more compounds of the present invention and (b) a second pharmaceutical composition comprising one or more additional therapeutic agents as defined herein in the treatment of the diseases disclosed herein. In preferred embodiments of all aspects of the invention, the disease to be treated is CLL.

In a further aspect, the present invention is directed to a product containing a compound of formula (I) as defined herein, or a tautomer thereof or a pharmaceutically acceptable salt, solvate or polymorph of said compound or tautomer, and one or more other therapeutic agents as defined herein, as a combined preparation for simultaneous, separate or sequential use in treating a neoplastic disease or an immune disease.

It is well known that immunosuppression is one of major side-effect of many conventional chemotherapeutics. For example, at low dose, cyclophosphamide can be used to treat immune diseases such as multiple sclerosis, rheumatoid arthritis and the suppression of transplant rejections (Emadi A, *et al*, *Nat Rev Clin Oncol*. **2009** Nov; 6(11):638-47; Perini P, *et al*. *Neurol Sci*. **2008** Sep; 29 Suppl 2:S233-4) and is also widely used in bone marrow transplantation “conditioning” and “mobilization” regimens, and for the treatment of refractory severe autoimmune conditions, such as systemic lupus erythematosus (SLE), minimal change disease, severe rheumatoid arthritis, Wegener’s granulomatosis (with trade name Cytoxan), scleroderma, and multiple sclerosis (with trade name Revimmune). In addition, HDAC has recently emerging as a promising target for treating immune disease [Szyf M. *Clin Rev Allergy Immunol*. **2010** Aug;39(1):62-77]. The compounds of present invention may therefore be used for treatment of an immune disease.

In a preferred embodiment, the immune disease is selected from the group consisting of the rejection of transplanted organs and tissues, a graft-versus-host disease, a non-autoimmune inflammatory disease, and an autoimmune disease, wherein said autoimmune disease is selected from the group consisting of acute disseminated encephalomyelitis, addison’s disease, ankylosing spondylitis, antiphospholipid antibody syndrome, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease, bullous pemphigoid, coeliac disease, chagas disease, chronic obstructive pulmonary disease, churg-strauss syndrome,


dermatomyositis, Crohn's disease, diabetes mellitus type 1, endometriosis, goodpasture's syndrome, graves' disease, guillain-barré syndrome, hashimoto's disease, hidradenitis suppurativa, idiopathic thrombocytopenic purpura, interstitial cystitis, lupus erythematosus, morphea, multiple sclerosis, myasthenia gravis, narcolepsy, neuromyotonia, pemphigus vulgaris, pernicious anaemia, polymyositis, primary biliary cirrhosis, psoriasis, psoriatic arthritis, rheumatoid arthritis, schizophrenia, scleroderma, temporal arteritis, vasculitis, vitiligo, and wegener's granulomatosis.


It should be understood that the invention is not limited to the particular embodiments shown and described herein, but that various changes and modifications may be made without departing from the spirit and scope of the invention as defined by the claims.

GENERAL SYNTHETIC METHODS

The compounds according to the present invention may be synthesized according to a variety of reaction schemes. Necessary starting materials may be obtained by standard procedures of organic chemistry. The compounds and processes of the present invention will be better understood in connection with the following representative synthetic schemes and examples, which are intended as an illustration only and not limiting of the scope of the invention. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art and such changes and modifications including, without limitation, those relating to the chemical structures, substituents, derivatives, and/or methods of the invention may be made without departing from the spirit of the invention and the scope of the appended claims.

A typical approach using $Z = (CH_2)_p$ as an example to illustrate the synthesis of the Formula (III) compounds is described in Scheme 1. X_1 and R_d in general Scheme 1 are the same as those described in the Summary section







Scheme 1

The commercially available starting material **1-1** (CAS#: 41939-61-1) can react with appropriate carboxylic acid to form the benzimidazole intermediate **1-2**, which can react with methyl acrylate by a Pd-catalyzed coupling to afford the cinnamate intermediate **1-3**. The intermediate (**1-3**) can be subsequently reduced, for example with H₂, Pd/C, to an amino-substituted intermediate (**1-4**), which can react with oxirane to easily afford intermediate (**1-5**). After that, intermediate **1-5** can be converted to intermediate (**1-6**) with high yield by reaction with a chlorinating reagent such as thionyl chloride or phosphorus pentachloride. Finally the hydroxylation of intermediate (**1-6**) in NH₂OH can afford the target compounds.

Alternatively, Formula (III) compounds can be synthesized according to the general Scheme 1A. X₁ and R_d in general Scheme 1A are the same as those described in the Summary section above.





Scheme 1A

1A-2 can be prepared by standard organic reactions. After that the commercially available starting material **1A-1** (CAS#: 41939-61-1) can react with **1A-2** to form the

benzimidazole intermediate **1A-3**, can be subsequently reduced, for example with H_2 , Pd/C, to an amino-substituted intermediate (**1A-4**), which can react with oxirane to easily afford intermediate (**1A-5**). After that, intermediate **1A-5** can be converted to intermediate (**1A-6**) with high yield by reaction with a chlorinating reagent such as thionyl chloride or phosphorus pentachloride. Finally the hydroxylation of intermediate (**1A-6**) in NH_2OH can afford the target compounds.

Similarly, a typical approach using Z is $(CH_2)_pNH(CH_2)_q$ as an example to illustrate the synthesis of the Formula (III) compounds is described in Scheme 2. X_1 and R_d in general Scheme 2 are the same as those described in the Summary section above.




Scheme 2

The starting material **2-1** can be converted to **2-2** by standard organic reactions. The secondary amine of intermediate (**2-2**) can be protected by a protecting group (-PG) such as Boc to yield intermediate (**2-3**), which undergo hydrolysis to afford the carboxylic acid intermediate **2-4**. After that, **2-4** can react with N^1 -methyl-4-nitrobenzene-1,2-diamine to form the benzimidazole intermediate **2-5**, which can react with methyl acrylate by a Pd-catalyzed coupling to afford the cinnamate intermediate **2-6**. The intermediate **2-6** can be subsequently reduced, for example Fe/NH_4Cl , Fe/HCl or $Zn/FeSO_4$, to an amino-substituted intermediate (**2-7**), which can react with oxirane to easily afford intermediate (**2-8**). After that **2-8** can be converted to intermediate (**2-9**) with high yield by reaction with a chlorinating reagent such as

thionyl chloride or phosphorus pentachloride. The de-protection of intermediate (**2-9**) affords the intermediate **2-10**. Finally the hydroxylamination of **2-10** in NH_2OH can afford the target compounds of Formula (III).

Alternatively, Formula (III) compounds can be synthesized according to the Scheme 2^a.


5 X_1 and R_d in general Scheme 2^a are the same as those described in the Summary section above.

Scheme 2^a

The starting material **2^a-1** with different p and q can be prepared by standard organic reactions. After that, **2^a-1** can be converted to carboxylic acid intermediate **2^a-2** with TFA. The secondary amine of intermediate **2^a-2** can be protected by a protecting group such as Boc to yield intermediate **2^a-3**, which can react with N¹-methyl-4-nitrobenzene-1,2-diamine to form the benzimidazole intermediate **2^a-4**. Next, the intermediate **2^a-4** can be subsequently reduced, for example Zn/AcOH, Fe/NH₄Cl, Fe/HCl or Zn/FeSO₄, to an amino-substituted intermediate (**2^a-5**), which can react with oxirane to easily afford alcohol intermediate (**2^a-6**). After that **2^a-6** can be converted to intermediate (**2^a-7**) with high yield by reaction with a chlorinating reagent such as thionyl chloride, MsCl/LiCl, or phosphorus pentachloride. The hydrolysis of ester **2^a-7**, e.g. in LiOH will afford carboxylic acid intermediate **2^a-8**, which can couple with NH₂OH to form the hydroxamic acid intermediate **2^a-9**. Finally, the de-protection of **2^a-9** afford the target

compounds of Formula (III).

As a further example, the several different approaches to synthesize CY-102 are described in the following scheme 2A:



Scheme 2A


As showed in Scheme 2A, CY-102-IV can be prepared by reacting CY-102-I with hydroxylamine, in the presence of a base such as for example potassium hydroxide. Said reaction is performed in an appropriate solvent, such as, for example, methanol. Finally, the de-Boc of CY-102-IV will lead to CY-102.

Another route showed in Scheme 2A for the preparation of CY-102 is follows: first, the hydrolysis of CY-102-I, e.g in LiOH or HCl to afford the carboxylic acid intermediate CY-102-II; next, CY-102-II can either couple with NH_2OH at the presence of appropriate reagents such as HATU/TEA/DCM to form CY-102-IV or can be converted CY-102-IV by such as the method reported in Tetrahedron Letters, 41, (2000), 6285-6288; finally, the de-Boc of CY-102-IV will lead to CY-102.


Alternative route to prepare CY-102 is first to hydrolyze CY-102-I e.g in LiOH or HCl to afford the carboxylic acid intermediate CY-102-II, which can coupled with O or N-protected hydroxylamine such as $\text{NH}_2\text{-O-THP}$, $\text{NH}_2\text{-O-Bn}$, N-t-Boc-O-THP , N-t-Boc-O-TBDMS , $\text{N,O-bis-(phenoxycarbonyl)-hydroxylamine}$, $\text{N,O-bis(tert-butoxycarbonyl)hydroxylamine}$, and $\text{N,N,O-tris-(trimethylsilyl)-hydroxylamine}$ to form intermediate CY-102-III. For example, CY-102-II can couple with $\text{NH}_2\text{-O-THP}$ in the presence of appropriate reagents such as $\text{N}^1\text{-(ethylcarbonimidoyl)-N,N-dimethyl-1,3-propanediamine, monohydrochloride (EDC)}$ and $\text{1-hydroxy-1H-benzotriazole (HOBT)}$ to form intermediate CY-102-III. This reaction may be performed in the presence of a base such as triethylamine, in a suitable solvent, such as, a mixture of dichloromethane and tetrahydrofuran. Finally CY-102 can be prepared by

deprotecting CY-102-III with an appropriate reagents, such as for example, trifluoro acetic acid. Said reaction is performed in an appropriate solvent, such as, for example, methanol or dichloromethane.

The approaches to synthesize the intermediate **CY-102-I**




are described in Scheme 2B-2C.



Scheme 2B


The commercially available starting material **2B-1** (CAS#: 41939-61-1) react with amine protected 3-aminopropanoic acid followed by a deprotection process to form the benzimidazole intermediate **2B-2**, which can react with (E)-methyl 3-(4-formylphenyl)acrylate to afford the cinnamate intermediate **2B-3**. The secondary amine of intermediate (**2B-3**) can be protected by a protecting group (-PG) such as Boc to yield intermediate (**2B-4**), which can be subsequently reduced, for example by Fe/NH₄Cl, Fe/HCl or Zn/FeSO₄, to an amino-substituted intermediate (**2B-5**). Intermediate **2B-5** can react with oxirane to easily afford intermediate (**2B-6**) which can be converted to intermediate (**CY-102-I**) with high yield by reaction with a chlorinating reagent such as thionyl chloride or phosphorus pentachloride.



Scheme 2C

The commercially available starting material **2C-1** (CAS#: 364-76-1) can react with oxirane to easily afford intermediate **2C-2**. The OH group of intermediate (**2C-2**) can be protected by a protecting group (-PG) to form the intermediate (**2C-3**). After that **2C-3** can react with NH_2CH_3 to afford intermediate **2C-4**, which can be reduced for example by $\text{Fe}/\text{NH}_4\text{Cl}$, Fe/HCl or Zn/FeSO_4 , to an amino-substituted intermediate (**2C-5**). At the same time, the commercially available starting material **2C-6** can be converted to the intermediate **2C-7** and then the Boc protected **2C-8** by standard organic reactions, which will react with **2C-5** to form the benzimidazole intermediate **2C-9**. Next, the OH group of **2C-9** will undergo the deprotection reaction to yield intermediate **2C-10**, which can be subsequently converted to **CY-102-I** with high yield by reaction with a chlorinating reagent such as thionyl chloride or phosphorus pentachloride.

The preferred method to prepare CY-102 as shown in Scheme 2D.



Scheme 2D

The commercially available starting material **2D-1** (4-bromobenzaldehyde) is converted to cinnamic intermediate **2D-2**. After that **2D-2** can react with tert-butyl 3-aminopropanoate to

form **2D-3**, which can be converted to carboxylic acid intermediate **2D-4** with an appropriate reagent, such as for example, trifluoro acetic acid. The Boc protection of amine of **2D-4** will lead to intermediate **2D-5**, which will react with N1-methyl-4-nitrobenzene-1,2-diamine(CAS#: 41939-61-1) to form intermediate **2D-6** followed by a cyclization reaction to form

5 benzimidazole intermediate **2D-7**. Intermediate **2D-7** can be reduced for example by Zn/AcOH, Fe/NH₄Cl, Fe/HCl or Zn/FeSO₄, to an amino-substituted intermediate (**2D-8**), which can react with oxirane to easily afford intermediate (**2D-9**). **2D-9** can be converted to intermediate **2D-10** with high yield by reaction with a chlorinating reagent such as thionyl chloride, MsCl/LiCl, or phosphorus pentachloride. The hydrolysis of **2D-10** e.g in LiOH will afford the carboxylic acid
10 intermediate **2D-11**, which can couple with NH₂OH at the presence of appropriate coupling reagents such as HATU/TEA/DCM to form intermediate **2D-12**. Finally, the de-Boc of **2D-12** will lead to the target molecule of **CY-102**.

EXAMPLES

15 The invention is illustrated by the following non-limiting examples.

Where NMR data are presented, ¹H spectra were obtained on either a Varian VXR-200 (200 MHz, ¹H), Varian Gemini-300 (300 MHz) or XL400 (400 MHz) and are reported as ppm down field from Me₄Si with number of protons, multiplicities, and coupling constants in Hertz indicated parenthetically. Where HPLC data are presented, analyses were performed using an
20 Agilent 1100 system. Where LC/MS data are presented, analyses were performed using an Agilent 6210 TOF LC/MS or an Applied Biosystems API-100 mass spectrometer and Shimadzu SCL-10A LC column: Altech platinum C18, 3 micron, 33 mmx7 mm ID; Samples were eluted using a linear gradient of 0-100% acetonitrile/pH4.50, 200 mM NH₄ acetate over 10 minutes with a flow rate of 3.0 mL/min. Chromatograms were generated over the range 240-400 nm
25 using a diode array detector.

In the following examples:

DCM = dichloromethane

Boc = tert-butyloxycarbonyl

HATU = *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate


30 TEA = triethanolamine

MsCl = methanesulfonyl chloride

DMF = dimethyl fluoride

THF = tetrahydrofuran

EA = ethyl acetate

Example 1: Preparation of CY-102

Scheme 2D

1.1: General procedure for Preparation of 2D-3: A mixture of **2D-1** (5.8 g, 31.8 mmol) and K_2CO_3 (13.2 g, 95.6 mmol) in 1,2-dichloroethane (150 mL) was stirred for 20 mins and filtered. To the filtrate was added **2D-2** (5 g, 24.51 mmol), and then $NaBH(OAc)_3$ (6.24 g, 29.4 mmol) was added in portions. The resulting mixture was stirred at r.t. overnight. The mixture was quenched with water and extracted with DCM. The organic phases were dried and concentrated. The residue was re-crystallized by DCM to give the product **2D-3** (4.0 g, yield 49.2%), as a white solid. HNMR-Analysis: 1H NMR ($CDCl_3$) δ : 7.67 (d, $J=16.04$ Hz, 1 H), 7.49 (d, $J=7.43$ Hz, 2 H), 7.35 (d, $J=7.43$ Hz, 2 H), 6.42 (d, $J=16.04$ Hz, 1 H), 4.27 (q, $J=6.91$ Hz, 2 H), 3.84 (s, 3 H), 2.87 (t, $J=5.87$ Hz, 3 H), 2.48 (t, $J=6.06$ Hz, 3 H), 1.44 (s, 11 H), 1.34 (t, $J=7.04$ Hz, 3 H).

1.2: General procedure for Preparation of 2D-4: To a suspension of **2D-3** (25.0 g, 75.1 mmol) in DCM (300 mL) was added TFA (30 mL) and the mixture was stirred at r.t. overnight. The mixture was concentrated, the residue was dissolved in DCM, adjusted to pH=7 with NaOH solution, the mixture was concentrated. The residue was dissolved in DCM and MeOH, then filtered and the filtrate was concentrated to give the crude product **2D-4** (20.0 g, yield 96.2%). HNMR-Analysis: 1H NMR ($DMSO-d_6$) δ : 1.23 (t, $J=7.04$ Hz, 3 H), 2.67 (t, $J=7.43$ Hz, 2 H), 3.01 - 3.12 (m, 2 H), 4.16 (d, $J=7.04$ Hz, 4 H), 6.67 (d, $J=16.04$ Hz, 1 H), 7.53 (d, $J=7.83$ Hz, 2 H), 7.63 (d, $J=16.04$ Hz, 1 H), 7.77 (d, $J=8.22$ Hz, 2 H), 9.13 (brs., 2 H).

1.3: General procedure for Preparation of 2D-5: A mixture of **2D-4** (20 g, 72.2 mmol) and Boc_2O (31.5 g, 144.4 mmol) in 1,4-dioxane (250 mL) was heated to reflux for 5 hrs. The mixture was concentrated and the residue was purified by column flash to give **2D-5** (22.1 g, yield 81.2%) as a white solid. HNMR-Analysis: 1H NMR ($CDCl_3$) δ : 1.33 (t, $J=7.24$ Hz, 3 H), 1.46 (brs., 9 H), 2.60 (brs., 2 H), 3.48 (brs., 2 H), 4.26 (q, $J=7.17$ Hz, 2 H), 4.47 (br. s., 2 H),

6.41 (d, $J=16.04$ Hz, 1 H), 7.23 (d, $J=6.26$ Hz, 2 H), 7.48 (d, $J=8.22$ Hz, 2 H), 7.66 (d, $J=16.04$ Hz, 1 H).

1.4: General procedure for Preparation of 2D-6: To a mixture of compound N¹-methyl-4-nitrobenzene-1,2-diamine (41 g, 0.11 mol) and TEA (20.4 g, 0.2 mol) in DCM (1000 mL) was added HATU (45.7 g, 0.12 mol) and **2D-5** (16.1 g, 0.11 mol) at 0°C and the reaction mixture was stirred at 20°C for 12 hrs. The reaction mixture was poured into water, washed with water for three times. The organic phase was dried over Na₂SO₄ and concentrated to give **2D-6** (50 g), as a red oil, which was used directly in the next step without further purification. ¹HNMR of **2D-6**: 1.44 (s, 9 H) 1.33(m, 3H) 2.67 (t, $J=6$ Hz, 2 H) 2.92 (s, 3 H) 3.18 (m, 2 H) 3.61 (t, $J=5.6$, 2H) 4.26 (q, $J=7.2$ Hz, 2H) 4.48 (s, 2 H) 6.41 (d, $J=16.$ Hz, 1 H) 6.57 (d, $J=9.2$ Hz, 1 H) 7.23(d, $J=7.6$, 2 H) 7.49 (d, $J=8$ Hz, 2 H) 7.65(d, $J=16.$ Hz, 1 H) 7.98-8.11(m, 2 H).

1.5: General procedure for Preparation of 2D-7: a mixture of compound **2D-6** (45 g, crude) in toluene and acetic acid (500 mL) was stirred at 100°C for 30 mins. The reaction mixture was concentrated to give **2D-7** (50 g), as a red oil, which was used directly in the next step without further purification. ¹HNMR-Analysis of **2D-7**: 1.27(t, 3H)1.33 (brs, 9 H) 3.05-3.18 (m, 4 H) 3.50 - 3.76 (m, 5 H) 4.20 (m, 2 H) 4.39 (s., 2 H) 6.31 (dd, $J=16.04$, 2.35 Hz, 1 H) 7.15 - 7.34 (m, 5 H) 7.48 - 7.60 (dd, $J=16,3.2$ Hz,1 H) 8.13 (d, $J=4.4$ Hz, 1 H) 8.52 (s, 1 H)

1. 6: General procedure for Preparation of 2D-8: To a mixture of compound **2D-7** (50 g, crude) and AcOH (20 mL) in DCM (1000 mL) was added Zn (15 g, 0.23 mol) at 0°C and the reaction mixture was stirred at 20°C for 1 h. The reaction mixture was filtered; the filtrate was concentrated to give the crude product (80 g) as red oil which was used to next step without further purification. ¹HNMR-Analysis of **2D-8**: 1.39 - 1.50 (m, 9 H) 3.11 (q, $J=7.30$ Hz, 3 H) 3.38 (br. s., 2 H) 3.67 (d, $J=11.74$ Hz, 3 H) 4.22 - 4.38 (m, 4 H) 6.36 (d, $J=16.04$ Hz, 1 H) 6.74 (d, $J=8.61$ Hz, 1 H) 6.99 - 7.20 (m, 3 H) 7.22 (s, 1 H) 7.33 (d, $J=6.65$ Hz, 2 H) 7.56 (d, $J=16.04$ Hz, 1 H).

1.7: General procedure for Preparation of 2D-9: a mixture of compound **2D-8** (80 g, crude) and ethylene oxide (80 mL) in water (1000 mL) and acetic acid (20 mL) was stirred at 23°C for 5 hrs. The reaction mixture was concentrated to give **2D-9** (63 g), as a red oil, which was used directly in the next step without further purification. ¹HNMR (MeOD 400MHz): 1.30 (m, 12 H) 3.22 (br. s., 2 H) 3.50 (d, $J=4.8$, 3 H) 3.563 (q , 1 H) 3.67 (m, 10 H) 4.23 (q, 2 H) 6.43 (d, 2 H) 6.38 (d, $J=16,1$ H) 6.91 (d, $J=8.4$, 2H) 7.22(t,2H) 7.29(d, 2H) 7.33 (d, $J=8$ Hz, 2 H) 7.44(q, 2H) 7.60 (t, 1 H).

1.8: General procedure for Preparation of 2D-10: to a mixture of compound **2D-9** (70 g, crude) and TEA (20.4 g, 0.2 mol) in DCM (1000 mL) was added MsCl (13.74 g, 0.12 mol) at

0°C and the reaction mixture was stirred at 20°C for 1 h. The reaction mixture was poured into water, washed with water three times. The organic phase was dried over Na₂SO₄ and concentrated to give the crude product (100 g). The crude product was dissolved in DMF (500 mL) and LiCl (16.8 g, 0.4 mol) and the resulting mixture was stirred at 100°C for 2 hrs. The mixture was concentrated and purified by silica gel chromatography to give **2D-10** (18 g).

¹HNMR (DMSO 400MHz): 1.25(m, 12 H) 3.03 (br. s., 2 H) 3.51 (m, 2 H) 3.58 - 3.69 (m, 10 H) 4.17 (q, J=7.6 Hz, 2 H) 4.45 (br. s., 2 H) 6.58 (d, J=16 Hz, 1 H) 6.8 (t, 1H) 6.9(br.s, 1H) 7.25 (d, J=8, 1H) 7.33(d, J=9.2, 1H) 7.60(d, J=16, 1H) 7.66 (d, J=7.2, 2 H).

1.9: General procedure for Preparation of 2D-11: A mixture of compound **2D-10** (36 g, 59.6 mmol) and LiOH.H₂O (3.78 g, 88 mmol) in a mixture of THF and water (600 mL) was stirred at 23°C for 5 hrs. The reaction mixture was acidified with HCl (1M) to pH=7 and the mixture was filtered. The solid was collected to give **2D-11** (20 g, yield: 59%), as a white solid, which was used directly in the next step without further purification.

1.10: General procedure for Preparation of 2D-12: To a mixture of **2D-11** (16.4 g, 28.52 mmol) and TEA (15.0 g, 0.147mol) in DCM (500 mL) was added HATU (16.8 g, 44 mmol) and NH₂OH-HCl (5.16 g, 73.7 mmol) in turn at 20°C. The reaction mixture was stirred at 20°C for 5 hrs. The mixture was poured into water, diluted with DCM, washed with water for three times. The organic phase was dried over Na₂SO₄ and concentrated to give the crude product. The crude product was purified with prep-HPLC to give **2D-12** (7 g, yield: 42%) as white solid.

1. 11: General procedure for Preparation of CY-102: a mixture of compound **2D-12** (7 g, 11.86 mmol) and HCl/EA (50 mL) in DCM (100 mL) was stirred at 23°C for 2 hrs. The reaction mixture was concentrated to give **CY-102** (5.875 g, yield: 95%) as a yellow powder. ¹HNMR(MeOD 400MHz): 3.73 (m, 8 H) 3.87 (m, 4 H) 4.04 (s, 3 H) 4.38 (s, 2 H) 6.50 (d, J=16Hz, 1 H) 6.88 (d, J=2Hz, 1H) 7.18 (dd, J=9.2, 2Hz, 1 H) 7.50 (d, J=16 Hz, 1 H) 7.68 (m, 5 H). m/z(MH⁺) is 490.

Example 2 Inhibition of Histone Deacetylase Enzymatic Activity



The following assay protocol is used to assess the inhibitory activity of the compounds of the invention against the HDAC enzymes (Hela Nuclear Extract assay):

- Buffer: 25 mM HEPES, pH 8.0, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl₂
- Substrate: Fluor-de-Lys substrate (Biomol, Cat. # KI-104) in a 50 mM stock solution in DMSO.
- Enzyme stock solution: 4 µg/mL enzyme in buffer.

To begin the assay, test compounds (2 µl in DMSO diluted to 13 µl in buffer for transfer

to the assay plate) are pre-incubated with enzyme (20 μ l of 4 μ g/mL stock solution) for 10 minutes at room temperature in 35 μ l pre-incubation volume. The reaction is started by bringing the temperature to 37°C and adding 15 μ l substrate. Total reaction volume is 50 μ l. The reaction is stopped after 20 minutes by adding 50 μ l developer, prepared as directed by Biomol (Fluor-de-Lys developer, Cat. # KI-105). Assay plate is incubated in the dark for 10 minutes at room temperature before reading (λ_{EX} = 360 nm, λ_{Em} = 470 nm, Cutoff filter at 435 nm). The HDAC inhibitors SAHA and TSA are used as reference compounds. Such assays, carried out with a range of doses of test compounds, allow the determination of an approximate IC₅₀ value.

As an example, the following table shows the results obtained for CY-102 and Bendamustine. In the HDAC (nucleare extract) assay, CY-102 is about 10-fold more potent than the FDA approved HDAC inhibitor SAHA.

Bendamustine		CY-102		
				
HDAC	CY-102	SAHA	Trichostatin A	Bendamustinie
IC ₅₀ (Nuclear Extract)	3.5 nM	26.4 nM	2.1 nM	N/A*
* No HDAC activity up to highest testing concentration of 10 μ M				

Example 3 Molecular Docketing Study

Computer modeling with the MOE program (Chemical Computer Group, Canada) was used to assess the interaction between CY-102 and HDAC8. The result (not shown) indicates that CY-102 tightly binds to HDAC8 at its catalytic center, which is consistent with the existing data showing that CY-102 is a strong HDAC inhibitor.

Example 4 Water Solubility

To measure water solubility, to approximately 10 mg of a sample in a tube-stoppered 10 mL graduated cylinder, increasing volumes of distilled water at room temperature were added according to the steps shown in the table below:

Water Solubility	step 1	step 2	step 3	step 4	step 5
Total volume of H ₂ O added (mL)	1	2	4	5	10
Approximate solubility (mg/mL)	10	5	2.5	2	1

After each addition of water to give the indicated total volume, the mixture was vortexed or sonicated for 1 min and was visually inspected for any undissolved parts of the sample. If, after a total of 10 mL of water had been added (step 5), the sample or parts of it remained undissolved, the contents of the measuring cylinder was transferred to a 100 mL measuring cylinder which was then filled up with water up to 100 mL (20 ml, 25 ml, 50 ml, 100 ml) and shaken. The approximate solubility was given in the table under that volume of added water in which complete dissolution of the sample occurred. If the substance was still apparently insoluble, further dilution was undertaken to ascertain whether the column elution or the flask solubility method should be used.

Using the method described above, water solubility of CY-102 was determined to be greater than about 20 mg/mL, which is at least about 200-fold more water soluble than NL-101.

Example 5 General *In vitro* Anti-proliferation Assay

Cell antiproliferation assay is performed by using the PerkinElmer ATPlite™ Luminescence Assay System. Briefly, the various test cancer cell lines are plated at a density of about 1×10^4 cells per well in Costar 96-well plates, and are incubated with different concentrations of compounds for about 72 hours in medium supplemented with 5% FBS. One lyophilized substrate solution vial is then reconstituted by adding 5 mL of substrate buffer solution, and is agitated gently until the solution is homogeneous. About 50 μ L of mammalian cell lysis solution is added to 100 μ L of cell suspension per well of a microplate, and the plate is shaken for about five minutes in an orbital shaker at ~700 rpm. This procedure is used to lyse the cells and to stabilize the ATP. Next, 50 μ L substrate solution is added to the wells and microplate is shaken for five minutes in an orbital shaker at ~700 rpm. Finally, the luminescence is measured by a PerkinElmer TopCount® Microplate Scintillation Counter. Such assays, carried out with a range of doses of test compounds, allow the determination of the cellular anti-antiproliferative IC₅₀ of the compounds of the present invention.

Example 6 *In vitro* Assay: NCI-60 DTP Human Tumor Cell Line Screen at 10 μ M

NL-101 and CY-102 was sent to U.S. National Cancer Institute (NCI) for NCI 60-cell

line screening using a single compound dose (10 μ M).

The human tumor cell lines of the cancer screening panel were grown in RPMI 1640 medium containing 5% fetal bovine serum (5% FBS) and 2 mM L-glutamine. For a typical screening experiment, cells were inoculated into 96-well microtiter plates in 100 μ L, at plating densities ranging from 5,000 to 40,000 cells/well depending on the doubling time of individual cell lines. After cell inoculation, the microtiter plates were incubated at 37°C, 5% CO₂, 95% air, and 100% relative humidity for 24 h prior to addition of experimental compounds. After 24 hr, two plates of each cell line were fixed *in situ* with TCA, to represent a measurement of the cell population for each cell line at the time of drug addition (Tz). Experimental drugs were solubilized in dimethyl sulfoxide at 400-fold the desired final maximum test concentration and stored frozen prior to use. At the time of drug addition, an aliquot of frozen concentrate was thawed and diluted to twice the desired final maximum test concentration with complete medium containing 50 μ g/ml gentamicin. Aliquots of 100 μ L of these different drug dilutions were added to the appropriate microtiter wells already containing 100 μ L of medium, resulting in the required final drug concentrations.

Following drug addition, the plates were incubated for an additional 48 hrs at 37°C, 5% CO₂, 95% air, and 100% relative humidity. For adherent cells, the assay was terminated by the addition of cold TCA. Cells were then fixed *in situ* by the gentle addition of 50 μ L of cold 50% (w/v) TCA (final concentration, 10% TCA) and incubated for 60 minutes at 4°C. The supernatant was discarded, and the plates were washed five times with tap water and air dried. Sulforhodamine B (SRB) solution (100 μ L) at 0.4% (w/v) in 1% acetic acid was added to each well, and plates were incubated for 10 minutes at room temperature. After staining, unbound dye was removed by washing five times with 1% acetic acid and the plates were air dried. Bound stain was subsequently solubilized with 10 mM trizma base, and the absorbance was read on an automated plate reader at a wavelength of 515 nm. For suspension cells, the methodology was the same except that the assay was terminated by fixing settled cells at the bottom of the wells by gently adding 50 μ L of 80% TCA (final concentration, 16 % TCA).

Using the seven absorbance measurements [time zero, (Tz), control growth, (C), and test growth in the presence of drug at the 10 μ M concentration levels (Ti)], percentage growth was calculated at each of the drug concentrations levels. Percentage growth inhibition was calculated as: $[(Ti-Tz)/(C-Tz)] \times 100$ for which $Ti \geq Tz$ or $[(Ti-Tz)/Tz] \times 100$ for which $Ti < Tz$.

The results of the assays for CY-102 and NL-101 are summarized in the table below.

Cell Panel	Cell Line	NL-101 Growth %	CY-102 Growth %
Leukemia	HL-60(TB)	18.98	-10.34
Leukemia	K-562	31.63	1.07
Leukemia	MOLT-4	18.01	2.17
Leukemia	CCRF-CEM	17.16	
Leukemia	RPMI-8226	49.23	2.28
Leukemia	SR	29.62	0.09
NSCLC	A549/ATCC	43.54	-32.69
NSCLC	EKVX	92.48	
NSCLC	HOP-62	13.56	-30.02
NSCLC	HOP-92	22.07	-35.89
NSCLC	NCI-H226	60.27	-20.77
NSCLC	NCI-H23	30.14	-6.76
NSCLC	NCI-H322M	75.30	-15.30
NSCLC	NCI-H460	25.56	2.61
NSCLC	NCI-H522	-5.90	
Colon Cancer	COLO 205	54.36	-81.34
Colon Cancer	HCC-2998	86.98	-80.55
Colon Cancer	HCT-116	23.73	-1.84
Colon Cancer	HCT-15	76.48	9.96
Colon Cancer	HT29	37.10	-50.67
Colon Cancer	KM12	65.79	-78.14
Colon Cancer	SW-620	30.40	2.33
CNS Cancer (Glioma)	SF-268	10.27	-31.46
CNS Cancer (Glioma)	SF-295	43.95	-53.87
CNS Cancer (Glioma)	SF-539	23.72	
CNS Cancer (Glioma)	SNB-19	55.62	-10.81
CNS Cancer (Glioma)	SNB-75	19.79	-48.45
CNS Cancer (Glioma)	U251	35.03	-39.66
Melanoma	LOX IMVI	-23.43	-26.41
Melanoma	MALME-3M	28.59	-64.17
Melanoma	M14	39.93	-76.75

Melanoma	MDA-MB-435	49.07	-73.24
Melanoma	SK-MEL-2	37.31	-21.36
Melanoma	SK-MEL-28	59.32	-21.36
Melanoma	SK-MEL-5	-6.55	-79.31
Melanoma	UACC-257	33.75	-36.39
Melanoma	UACC-62	12.99	-69.24
Ovarian Cancer	IGROV1	55.99	-14.43
Ovarian Cancer	OVCAR-3	50.40	-55.02
Ovarian Cancer	OVCAR-4	68.52	-19.00
Ovarian Cancer	OVCAR-5	66.28	-3.75
Ovarian Cancer	OVCAR-8	32.65	-26.88
Ovarian Cancer	NCI/ADR-RES	77.80	17.78
Ovarian Cancer	SK-OV-3	34.40	-41.53
Renal Cancer	786-0	19.82	-30.40
Renal Cancer	A498	49.70	-81.56
Renal Cancer	ACHN	14.30	-19.97
Renal Cancer	CAKI-1	27.64	-25.49
Renal Cancer	RXF 393	5.95	-35.55
Renal Cancer	SN12C	17.91	-1.53
Renal Cancer	TK-10	48.48	-28.81
Renal Cancer	UO-31	63.69	-9.19
Prostate Cancer	PC-3	51.40	-1.86
Prostate Cancer	DU-145	18.47	-16.06
Breast Cancer	MCF7	37.24	-15.28
Breast Cancer	MDA-MB-231	72.24	-35.33
Breast Cancer	HS 578T	24.01	8.17
Breast Cancer	BT-549	79.21	-33.26
Breast Cancer	T-47D	-16.57	-27.48
Breast Cancer	MDA-MB-468	-34.68	-21.52
	Median	36.35	-28.5

The results show that, when NL-101 and CY-102 were tested side-by-side at a single dose of about 10 μ M in 60 cancer cell lines of leukemia, multiple myeloma, non small cell lung cancer (NSCLC), breast cancer, melanoma, ovarian cancer, prostate cancer, colon cancer, CNS cancer, and renal cancer, the mean growth percent of NL-101 in the 60 cancer cell lines is 36%.

- 5 In contrast, the mean growth percent of CY-102 is -28%. Based on this data, the average cellular IC_{50} of CY-102 in the 60 cancer cell lines is expected to be at least 10-fold more potent than the IC_{50} of NL-101, which on average is about 2 μ M.

More impressively, CY-102 was found to be particularly potent in several solid tumor cell lines, such as breast cancer (*e.g.*, MCF7, MDA-MB-231, BT-549, T-47D, MDA-MB-468),
10 colon cancer (*e.g.*, COLO 205, HCC-2998, HT29, SW-620), renal (*e.g.*, A498), and particularly in melanoma (*e.g.*, MALME-3M, M14, MDA-MB-435, SK-MEL-5, UACC-62), suggesting that CY-102 may have wide applications in treating solid tumors. On the other hand, NL-101 appears to be more effective against hematological cancers such as leukemia, lymphoma, and multiple myeloma.

15 **Example 7 *In vitro* hERG Assay**

The hERG (Human Ether- α -go-go-Related-Gene) assay was used to assess cardiotoxic effects of drug candidates, CY-102. Results (not shown) demonstrated that CY-102 has much lower (about 5-10 fold less) cardiotoxicity compared to that of NL-101.

Example 8 *In vivo* Xenograft Studies


- 20 As compared to NL-101, CY-102 is much more potent in *in vitro* cellular antiproliferative assay (about 10-fold more potent, see above), shows much less *in vitro* cardiotoxicity in the hERG assay (about 5-10 fold less, see above), and is significantly more (>200 fold) soluble in water (see above). Thus, CY-102 is selected for *in vivo* studies in the xenograft models of Breast cancer (MBA-MD-231, MX-1), SCLC (H69, H526), Sarcoma (HT-1080, SJSA-1),
25 Melanoma(MDA-MB-435, SK-MEL-5), and NSCLC (H1975, HCC827, H3255, PC-9).

Athymic nude mice (CD-1 *nu/nu*) or SCID mice are obtained at age 6-8 weeks from vendors and acclimated for a minimum 7-day period. The cancer cells are then implanted into the nude mice. Depending on the specific tumor type, tumors are typically detectable about two weeks following implantation. When tumor sizes reach $\sim 100\text{-}200\text{ mm}^3$, the animals with
30 appreciable tumor size and shape are randomly assigned into groups of 8 mice each, including one vehicle control group and treatment groups. Dosing varies depending on the purpose and length of each study, which typically proceeds for about 3-4 weeks. Tumor sizes and body weight are typically measured three times per week. In addition to the determination of tumor

size changes, the last tumor measurement is used to generate the tumor size change ratio (T/C value), a standard metric developed by the National Cancer Institute for xenograft tumor evaluation. In most cases, %T/C values are calculated using the following formula: $\% T/C = 100 \times \Delta T/\Delta C$ if $\Delta T > 0$. When tumor regression occurred ($\Delta T < 0$), however, the following
5 formula is used: $\% T/T_0 = 100 \times \Delta T/T_0$. Values of $<42\%$ are considered significant.

Claims


1. A compound of Formula (I) or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof:



wherein

Z is $(\text{CR}_a\text{R}_b)_p\text{N}(\text{R}_a)(\text{CR}_a\text{R}_b)_q$;

X₁ and X₂ are each independently selected from halo and OSO_2R_c ;



Q is heteroaryl, which is optionally substituted with alkyl, alkenyl, alkynyl, cycloalkyl, halo, nitro, oxo, cyano or OR_e ;


R_a, R_b, R_d and R_e are each independently selected from H, alkyl, alkenyl and alkynyl;

R_c is selected from alkyl, alkenyl and alkynyl; and


p and q are each independently selected from 0, 1, 2, 3 and 4.

2. A compound according to claim 1 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein p is 1 and q is 2; or p is 2 and q is 1; or p is 0 and q is 3; or p is 3 and q is 0; or p and q are both 2.
3. A compound according to either claim 1 or claim 2 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein Z is $(\text{CH}_2)_p\text{NH}(\text{CH}_2)_q$.
4. A compound according to claim 3 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein Z is $(\text{CH}_2)_2\text{NH}(\text{CH}_2)$.
5. A compound according to any one of claims 1 to 4 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein X₁ and X₂ are each independently selected from chloro, bromo and iodo.


6. A compound according to claim 5 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein X_1 and X_2 are both chloro.
7. A compound according to any one of claims 1 to 6 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein Q is an optionally substituted benzimidazolyl.
8. A compound according to claim 7 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein Q is benzimidazolyl substituted by one or more alkyl groups.
9. A compound according to claim 8 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof, wherein the compound of Formula (I) is represented by Formula(II):



10. A compound according to claim 9 or an N-oxide thereof, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof wherein the compound of Formula (I) is represented by Formula (III):



11. A compound according to claim 10, which is



or a pharmaceutically acceptable salt, solvate, polymorph or tautomer thereof.

12. A compound according to claim 11, which is the hydrochloride salt or a solvate or polymorph thereof.

13. A pharmaceutical composition comprising a compound of formula (I) or an N-oxide thereof as defined in any one of claims 1 to 12, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or an N-oxide thereof, and a pharmaceutically acceptable diluent or carrier.
14. A combination comprising a compound of formula (I) or an N-oxide thereof as defined in any one of claims 1 to 12, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof, together one or more other therapeutic agents.
15. A combination according to claim 14 wherein the one or more other therapeutic agents is selected from:
 - proteasome inhibitors (e.g. bortezomib, carfilzomib),
 - IMiDs (e.g. Thalidomide, lenalidomide, pomalidomide),
 - platinum agents (e.g. cisplatin, carboplatin),
 - folate antagonists (e.g. pemetrexed, pralatrexate),
 - CD30 antibodies and conjugates (e.g. brentuximab, vendotin),
 - antibodies (also conjugated) to treat haematological malignancies like anti CD20 (e.g. ofatumumab, rituximab, GA101, etc),
 - B-cell receptor antagonists (e.g. ibrutinib),
 - PI3K antagonists (e.g. GS1101 or IPI145),
 - BTK inhibitors,
 - taxanes (e.g. taxol, paclitaxel),
 - antibodies (also conjugated) to treat ovarian cancer (e.g. alpha folate receptor mabs, CA125 antibodies),
 - antibodies to treat multiple myeloma (e.g. elotuzumab, anti CD38 mabs),
 - anthracyclines (e.g. doxorubicin, idarubicin),
 - nucleoside analogues (purine antagonists) like cytarabine, fludarabine, gemcitabine,
 - PNP antagonists (e.g. forodesine),
 - Bcr-abl tyrosinekinase blockers (e.g. imatinib, dasatinib, ponatinib, nilotinib),
 - mTor antagonists (e.g. temsirolimus, everolimus),

Agents influencing the CD40 activation (e.g. CD40 antagonists, CD40 gene medicines), multi tyrosine kinase antagonists (e.g. sorafenib, axitinib), and bifunctional antibodies (e.g. CD19/CD3, also conjugated, also recognising other CD epitopes).

16. A compound of formula (I) or an N-oxide thereof as defined in any one of claims 1 to 12, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof, or a combination according to claim either claim 14 or claim 15, for use as a medicament.
17. A compound of formula (I) or an N-oxide thereof as defined in any one of claims 1 to 12, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof, or a combination according to either claim 14 or claim 15, for use as a medicament for treating a neoplastic disease or an immune disease.
18. A method of treating a neoplastic disease or an immune disease, comprising administering to a subject in need thereof an effective amount of a compound of formula (I) or an N-oxide thereof as defined in any one of claims 1 to 12, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof, or a combination according to either claim 14 or claim 15.
19. A compound or combination for use, according to claim 17, or a method according to claim 18, wherein the neoplastic disease is a solid tumor.
20. A compound or combination for use, or a method according to claim 19, wherein the solid tumor is melanoma, breast cancer, lung cancer, colon cancer, renal cancer, or sarcoma.
21. A product containing a compound of formula (I) or an N-oxide thereof as defined in any one of claims 1 to 12, or a pharmaceutically acceptable salt, solvate, polymorph or tautomer of said compound of formula (I) or N-oxide thereof, and one or more other therapeutic agents as defined in claim 15, as a combined preparation for simultaneous, separate or sequential use in treating a neoplastic disease or an immune disease.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/051944

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D235/30 A61K31/4184 C07D235/14
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2008/050125 A1 (IMP INNOVATIONS LTD [GB]; ABOAGYE ERIC OFORI [GB]) 2 May 2008 (2008-05-02) claims 2, 17, 20 page 35; example 3; compounds C1-A	1
A	WO 2010/085377 A2 (CRYSTAL BIOPHARMACEUTICAL LLC [US]; CHEN YU [US]; CHEN YI [US]) 29 July 2010 (2010-07-29) cited in the application claims 1, 35, 40, 42	1-21



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

4 March 2013

Date of mailing of the international search report

11/03/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Guazzelli, Giuditta

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2013/051944

Patent document cited in search report		Publication date		Patent family member(s)		Publication date	
WO 2008050125	A1	02-05-2008	US	2010063144	A1	11-03-2010	
			WO	2008050125	A1	02-05-2008	

WO 2010085377	A2	29-07-2010	CA	2750413	A1	29-07-2010	
			CN	102186842	A	14-09-2011	
			EP	2389375	A2	30-11-2011	
			JP	2012515776	A	12-07-2012	
			US	2011269706	A1	03-11-2011	
			WO	2010085377	A2	29-07-2010	
