

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 February 2010 (25.02.2010)

PCT

(10) International Publication Number
WO 2010/021590 A1

(51) International Patent Classification:
A23L 1/303 (2006.01) *A23L 1/28* (2006.01)

(21) International Application Number:
PCT/SE2009/050942

(22) International Filing Date:
20 August 2009 (20.08.2009)

(25) Filing Language: Swedish

(26) Publication Language: English

(30) Priority Data:
0801820-2 21 August 2008 (21.08.2008) SE

(71) Applicant (for all designated States except US): VIA-SOLDE AB [SE/SE]; Box 292, S-791 27 Falun (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): EKLUND-JONSSON, Charlotte [SE/SE]; Rösbacka 575, S-449 90 Nol (SE). JOHANSSON, Sara [SE/SE]; Norra Gubberogatan 19B, S-416 63 Göteborg (SE). HJORTMO, Sofia [SE/SE]; Bokskogsbacken 67, S-422 56 Hisings Backa (SE). LINDQVIST, Helen [SE/SE]; Uggledals Byväg 39, S-427 40 Billdal (SE).

(74) Agent: ROMARE, Anette; Box 142, S-401 22 Göteborg (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: METHOD TO VITAMIN-D ENRICH FOOD USING YEAST AND UV LIGHT

(57) Abstract: Method for vitamin enrichment of foodstuffs containing ascomyceta such as yeast or the mould *Fusarium venenatum*, whereupon the foodstuff is illuminated with ultraviolet light with a wavelength of between 100-400 nm, in order for vitamin D₂ to form in the foodstuff.



WO 2010/021590 A1

METHOD TO VITAMIN-D ENRICH FOOD USING YEAST AND UV LIGHT

TECHNICAL FIELD

The invention relates to a method for enrichment of foodstuffs containing ascomycota such as yeasts, or the mould *Fusarium venenatum* with
5 vitamins.

BACKGROUND ART

Vitamin D is a fat-soluble vitamin that in humans amongst other things regulates the levels of calcium and phosphorus in the blood and regulates the mineralization of the skeleton. Together with sunlight (UVB) the vitamin
10 D₃, a form of vitamin D, can be formed in the skin of the human from the provitamin 7-Dehydrocholesterol.

The provitamin converts to the previtamin D₃ after exposure to ultraviolet light, the previtamin is then further transformed to the physiologically active vitamin D₃ with the aid of heat. Vitamin D₃ may also be received from various
15 food stuffs and especially animal food stuffs such as fish, cooking fat, eggs, dairy products and meat.

In fungi, for instance in yeast, ergosterol functions as a provitamin and can with the aid of ultraviolet light and heat be transformed into vitamin D₂. Vitamin D₂ is just as vitamin D₃ physiologically active in the human body.

20 As vitamin D is important for the intercalation of calcium in the skeleton it has been shown that a lack of vitamin D involves an increased risk of osteoporosis. In later years it has also been shown that an adequate intake of vitamin D may reduce the risk of developing other diseases, such as colon and breast cancer, cardiovascular disease, multiple sclerosis, psoriasis and
25 diabetes.

During periods when not enough sunlight is present to produce vitamin D in the skin, for instance during the winter season in northern latitudes, older

people and people with darker skin above all run a great risk of not receiving enough vitamin D. The diet is then of great importance as a source of vitamin D. Vegetarians and vegans that do not receive vitamin D from their diet have to rely solely on dietary supplements.

- 5 There is thus a need to supply foodstuffs able to give an addition of vitamin D especially to the individuals that have difficulties covering the need of vitamin D with an ordinary diet. One aim of the invention is therefore to provide such foodstuffs that are vegetarian and easily accessible. Another aim of the invention is to offer foodstuffs with an increased amount of vitamin D without
10 particular enrichment additives of vitamin since many consumers are sceptical towards additives in foodstuffs.

SUMMARY OF THE INVENTION

- According to the invention a method has now been developed for enriching foodstuffs containing ascomycota such as yeast or the mould *Fusarium*
15 *venenatum* with vitamins. The method according to the invention is mainly characterized in that ergosterol in foodstuffs is converted into vitamin D₂ through the foodstuff being placed under a source of ultraviolet light and the foodstuff being illuminated with ultraviolet light with a wavelength of between 100-400 nm.

- 20 It has been shown that good results are achieved with a light source that during the illumination step is placed at a distance of between 0-2 m from the foodstuff.

- One type of foodstuffs that are suitable for treatment with the method according to the invention is bread baked on rising dough, since such bread
25 contains yeast and thus also ergosterol that can be converted into vitamin D.

The bread is preferably portion-packed bread, as such bread offers a large exposed and illuminable surface relative to its volume. Especially suitable portion-packed bread are round loaves, for instance rye loaves.

It has been shown that the temperature at which the foodstuff is illuminated with ultraviolet light is of lesser importance. It is possible to obtain an increased level of vitamin D₂ in a foodstuff containing yeast at temperatures over 10 °C. However, products with a temperature below 10 °C can also
5 obtain an increased level of vitamin D₂ through the surface layer being heated during illumination.

To achieve suitable levels of vitamin D in the foodstuff it is sufficient to illuminate the foodstuff for a relatively short period of time. The illumination time can be 0.1-60 seconds, preferably 2-30 seconds and more preferably 3-
10 5 seconds. Short illumination times are especially suitable for continuous processes, for instance as bread or another foodstuff is brought forward on a conveyor belt and is allowed to pass under a light source.

The illumination time can advantageously be adjusted so that the foodstuff is treated to a predetermined amount of vitamin D₂, for instance to an amount
15 corresponding to 30 % of a recommended daily intake per serving of the foodstuff. The illumination time may preferably be adjusted to reflect any changes made to the recommended daily intake as a result of new research findings or recommendations from a public authority.


The method according to the invention is preferably applied to a continuous
20 process where the foodstuff is transported past the source of ultraviolet light at a speed adjusted to reach a predetermined level of illumination and thus a predetermined level of vitamin D in the foodstuff. The illumination time is determined in a continuous process partly by the speed of the conveyor belt, partly by the amount of light per surface unit that is delivered. It is
25 alternatively possible to illuminate foodstuffs in a non-continuous process for a specific period of time.

Some of the wavelengths within the mentioned interval 100-400 nm also have anti-microbial effects, for instance 254 nm. This leads to unwanted microorganisms being eliminated at the same time as vitamin D₂ is formed.

DESCRIPTION OF THE INVENTION

Conversion of ergosterol to vitamin D₂ takes place according to the following simplified chemical equation.

Chemical equation:



$h\nu$ = light of a specific wavelength

Δ = heat

10 Bread containing yeast, and thus also ergosterol, can be illuminated with ultraviolet light and thereby receive an increased level of vitamin D₂.

Bread containing ergosterol is illuminated according to the invention with ultraviolet light which leads to the formation of vitamin D₂. Vitamin D₂ is formed during illumination of ultraviolet light with wavelengths between 100-400 nm, at a distance of 0-2 m, at temperatures over 10 °C.

15 The advantages with a bread that contains vitamin D₂ is that a large number of consumers can receive their daily amount of vitamin D even during the winter season when the amount of sunlight in the Nordic countries is insufficient to form vitamin D in the skin. This is especially advantageous to vegetarians and vegans who do not eat fish, eggs or meat, foodstuffs that are
20 rich in vitamin D. Since vegans also do not eat dairy products, which in Sweden are enriched with vitamin D, this group derives particular advantage from a bread rich in vitamin D. Also older people and people with darker skin benefit from this invention since it is more difficult for them to produce vitamin D in the skin. Groups that also may take advantage of bread enriched with
25 vitamin D are people that are allergic to milk or that are lactose intolerant. Also pregnant and breastfeeding women need an increased addition of vitamin D since both their own and the foetus' or baby's need of vitamin D has to be met.

The invention thus admits that it is possible in a simple and natural way to increase the amount of vitamin D in foodstuffs containing ascomyceta such as yeast but also products containing the mould *Fusarium venenatum*. Initial studies have shown that that vegetarian products based on *Fusarium*
5 *venenatum* are suitable to enrich with vitamin D by the aid of ultraviolet light.

The level of vitamin D₂ obtained after exposure to ultraviolet light is dependent on the content of fungi and the shape of the product (i.e. the size of the exposed surface in relation to the volume) and will thus vary. The conversion of ergosterol to vitamin D₂ in products containing *Fusarium*
10 *venenatum* during illumination with ultraviolet light is very effective since the amount of ergosterol is high.

Due to the fact that ultraviolet light converts the ergosterol in the fungus to vitamin D, no additives are needed in the foodstuff. This is an advantage, both since consumers may be sceptical to additives in foodstuffs and since
15 enrichment of foodstuffs such as bread is difficult and affects recipes and the number of ingredients in the foodstuff. With the illumination with ultraviolet light the amount of vitamin D in the foodstuff can be increased without having to change the manufacturing process to an appreciable extent. The ultraviolet equipment can easily be incorporated into a continuous manufacturing line
20 producing for instance round loaves or other foodstuffs.

The recommended daily dosage of vitamin D₂ is at present 7.5 µg per day for adult, healthy individuals. A higher dosage of 10 µg is recommended for the elderly, children and individuals spending most of their time indoors.

In order to avoid overdoses it is therefore suitable that the dosage of vitamin
25 D₂ per serving of the foodstuff is below approximately 7.5 µg. An example of a suitable size of a serving of bread can be 1-3 pieces of bread. An advantage with the invention is that the illumination time can be adjusted to reflect changes in the recommended daily intake as a result of new research findings or recommendations from a public authority.

Today's stressful lifestyle also affects the development of the foodstuffs market and increases the proportion of fast food and easy cooked food. Many of these products are bread based, such as typical fast food accompaniments, for instance hot dog rolls or hamburger bread. The
5 demands on the nutritional content of the diet, and above all the nutritional density, increase more and more because we move less and due to this we cannot eat the same quantities of food as a person with high physical activity. Therefore, higher demands are also placed on fast food. The vitamin enrichment method according to the invention is of course especially suitable
10 also for bread that is a part of fast food, such as hot dog rolls, hamburger bread, pita bread or similar.

The ultraviolet rays only penetrate a short distance into the bread, which means that only a superficial layer gets elevated levels of vitamin D. It is therefore suitable to illuminate bread that has a large exposed surface in
15 relation to volume, for instance round loaves, such as rye loaves or other types of portion-packed bread. The above mentioned fast food breads of course fall into this category.

For round loaves with a large exposed surface it has been shown that a sufficient amount of vitamin D₂ is achieved already at relatively short
20 illumination times of a few seconds. Illumination times of 0.1-60 seconds, preferably 2-30 seconds and more preferably 3-5 seconds are generally enough for this type of foodstuffs.

Some of the wavelengths within the mentioned interval 100-400 nm also have anti-microbial effects, for instance 254 nm. This leads to unwanted
25 microorganisms being eliminated at the same time as vitamin D₂ is formed.

Since the ultraviolet fluorescent lamps lose effect over time it is suitable that they are exchanged after half of the manufacturer's recommended life length of the lamp. This is done in order to secure correct and even amounts of vitamin D in the treated foodstuff.

DESCRIPTION OF FIGURES

The invention will in the following section be more thoroughly described with reference to the figures in the attached drawings.

In this respect:

- 5 Figure 1 shows an illumination device on a laboratory scale for illumination of bread according to the invention.

Figure 2 a and b shows an upscaled prototype of the illumination device for illumination of bread according to the invention.

- 10 Figure 3 shows a diagram over the amount of vitamin D in relation to the illumination time and different wavelengths for illumination of different types of bread.

Figure 4 shows a diagram over the amount of vitamin D in relation to illumination time for illumination of different types of bread.

- 15 Figure 5 shows a staple diagram over the amount of vitamin D in rye loaves of different temperatures after illumination with ultraviolet light.

Figure 6 shows a staple diagram over the amount of vitamin D in rye loaves of different temperatures before illumination with ultraviolet light.

Figure 7 shows a diagram over the amount of vitamin D in rye loaves in relation to illumination time for extremely long illumination times.

- 20 Figure 8 shows a diagram over the amount of vitamin D in relation to illumination time for a foodstuff containing *Fusarium venenatum*.

DETAILED DESCRIPTION

Equipment

For the purpose of examining suitable illumination times among other things, stationary tests on a laboratory scale were conducted on foodstuffs containing ascomyceta such as yeast or the mould *Fusarium venenatum* with the aid of the illumination device shown in Figure 1. The illumination device 1 comprises a light-proof illumination box 2 in which a test platform 3 is arranged. The test platform 3 is accessible for placement of a sample 4, for instance a round loaf through a hatch 5 in the wall of the illumination box 2. A lamp hood 6 containing ultraviolet lamps is placed over the illumination box 2 for illumination of a sample 4 arranged on the test platform 3. The illumination box has an opening 8 over the test platform 3, which opening can be closed with the aid of a plate that functions as a shutter 9.

During the tests the ultraviolet lamps are lit and the sample 4 is placed on the test platform 3 through the hatch 5 in the wall of the illumination box 2. The shutter 9 is thereafter removed whereupon the sample 4 is exposed to the ultraviolet light. After the illumination time expires the shutter 9 is put back into place, the sample 4 is removed through the hatch 5 and the ultraviolet lamps are switched off.

For use of the invention in a continuous process, equipment similar to the equipment shown in Figure 2a and 2b can, for example, be used. The illumination device 11 shown in Figures 2a and 2b comprises a lamp holder 12 with ultraviolet fluorescent lamps protected by a quartz glass 10. The lamp holder 12 is mounted on a stand 14 that is equipped with wheels 15 so that it is easily movable, for instance for different placements or for cleaning. The lamp holder 12 is also equipped with a light-proof screen 17 mounted in such a way that it prevents illumination of ultraviolet light from the ultraviolet lamps to the sides.

In Figure 2b two illumination devices 11 are shown placed on each side of a conveyor belt 16 on which the foodstuff that is to be illuminated is transported. The lamp holders' 12 light-proof screens 17 are arranged parallel to the side edges 16 of the conveyor belt and form together with the
5 lamp holders 12 a screened light tunnel over a section of the conveyor belt 16, as shown in the figure.

Through a device such as the one shown in Figure 2, it is possible to easily illuminate foodstuffs containing yeast fungi or the mould *Fusarium venenatum* as a step in the manufacturing process. The illumination can be
10 performed without any major modifications to the existing manufacturing equipment, which is a considerable advantage of the invention. The treatment according to the invention does not affect the manufacturing process in any other aspect, nor does it have any negative effect on properties such as taste, consistency or appearance of the finished foodstuff.

15 The baked breads or a foodstuff containing *Fusarium venenatum* are transported into the light tunnel and are illuminated for 3-5 seconds whereupon they are transported on, possibly for packaging. In the suitably vertically adjustable lamp holders 12 the fluorescent lamps 13 are placed parallel to the running direction of the conveyor belt 16. The two illumination
20 parts are suitably locked together with a simple locking arrangement arranged between the lamp holders 12.

Example 1

Rye loaves (weight: approximately 34 g, thickness: approximately 1 cm, diameter: approximately 12 cm) were illuminated with ultraviolet light at a distance of 2,5 cm during four different time periods: 2 seconds, 4 seconds, 8 seconds and 12 seconds. Two ultraviolet lights with different wavelengths were tested: one shortwave (254 nm) and one midrange (302 nm).

The results of the measurements are shown in table 1 and are illustrated in the diagram in Figure 3.

Table 1. Amount of vitamin D₂ in rye loaf illuminated with 254nm and 302nm

| Wavelength | Time (s) | Amount of vitamin D ₂ (µg per rye loaf) |
|------------|----------|---|
| | 0 | 0,00 |
| 254nm | 2 | 0,33 |
| | 4 | 2,57 |
| | 8 | 3,97 |
| | 12 | 5,01 |
| 302nm | 2 | 0,24 |
| | 4 | 1,24 |
| | 8 | 3,03 |
| | 12 | 4,23 |

Example 2

A comparison of the amount of vitamin D₂ was made for four different types of bread: wheat loaves, rye loaves, coarse crispbread and regular crispbread.

The results are shown in table 2 and are illustrated in the diagram in Figure 4

- 5 *Table 2. Comparison of the amount of vitamin D₂ in different kinds of bread after illumination with ultraviolet light of 302 nm.*

| | Amount of vitamin D ₂ (µg per g of piece of bread) | | | |
|-------|---|------------|--------------------|---------------------|
| Time | Rye loaf | Wheat loaf | Coarse crisp bread | Regular crisp bread |
| 0s | 0,00 | 0,00 | 0,00 | 0,00 |
| 2s | 0,43 | 0,00 | 0,88 | 0,19 |
| 4s | 0,93 | 2,18 | 0,00 | 0,56 |
| 8s | 1,60 | 3,10 | 0,56 | 0,46 |
| 16s | 1,99 | 2,91 | 0,39 | 0,39 |
| 30s | 6,21 | 7,44 | 1,02 | 2,22 |
| 45s | 6,78 | 11,84 | 1,42 | 2,33 |
| 60s | 12,90 | 16,73 | 1,90 | 2,87 |
| 2min | 14,58 | 19,81 | 4,22 | 9,70 |
| 4min | 26,11 | 18,98 | 3,48 | 9,60 |
| 10min | 30,38 | 24,02 | 7,77 | 16,44 |
| 15min | 30,10 | 21,78 | 7,31 | 17,43 |

- 10 The results indicate that the amount of vitamin D₂ was greatest in rye cakes after illumination for 15 minutes. The lowest amount was obtained in coarse crispbread. However it seems that the amount of D₂ in illuminated wheat cakes increased the fastest with increased illumination times.

- 15 A comparison of the time required for the different kinds of bread to reach an amount of 7 µg of vitamin D₂ showed that the wheat cakes needed approximately 30 seconds, the rye cakes needed approximately 45 seconds, regular crispbread needed approximately 1.5 minutes and coarse crispbread needed approximately 10 minutes.

Example 3

In order to determine how different temperatures after illumination influence the formation of vitamin D₂, rye loaves were exposed to different temperatures after illumination with ultraviolet light of 302 nm. The illumination time was 2 minutes for each loaf. The results are shown in table 3 and in the diagram in Figure 5.

Table 3. Temperature of the rye loaves after illumination

| | Description | | T _{after illumination} | T _{after oven} | T _{after room temperature} |
|--------------------|---|---|---------------------------------|-------------------------|-------------------------------------|
| High temperature | 5 min oven 100°C | A | 18°C | 50,9°C | - |
| | | B | 17°C | 56°C | - |
| Medium temperature | 1 min oven 100°C, room temperature for 10 min | A | 17°C | 29,6°C | 26,2°C |
| | | B | 17°C | 32,4°C | 26,4°C |
| Low temperature | 20 min room temperature | A | 18°C | - | 26°C |
| | | B | 17°C | - | 22,3°C |
| Frozen directly | Frozen directly after illumination | A | 17°C | - | - |
| | | B | 17°C | - | - |

- 10 The results show that there is a slight reduction in the amount of vitamin D₂ in rye loaves if the bread is frozen directly after illumination. The highest amount was obtained when the breads first were placed in room temperature for 20 minutes and then frozen.

Example 4

In order to determine how different temperatures prior to illumination influence the formation of vitamin D₂, rye loaves were exposed to different temperatures prior to illumination of ultraviolet light, see table 4. The illumination time was 2 minutes at 302 nm for each loaf. The results are shown in table 5 and in the diagram in Figure 6.

Table 4. Table showing the different temperatures for the different rye loaves

| Designation | T_{initial} | T_{after illumination} | Other information |
|--------------------|----------------------------|---------------------------------------|--|
| Frozen A | -8,5°C | -6,0°C | |
| Frozen B | -9,8°C | -6,8°C | |
| Fridge cold A | 9,4°C | 13,4°C | |
| Fridge cold B | 9,8°C | 12,3°C | |
| Room temp A | 17,8°C | 18,7°C | |
| Room temp B | 20,7°C | 20,7°C | |
| Oven A | 33,0°C | 27,5°C | 1 minute at 100°C prior to illumination |
| Oven B | 24,0°C | 23,7°C | 1 minute at 100°C prior to illumination |
| Oven C | 52,5°C | 42,4°C | 1 minute at 200°C prior to illumination |
| Oven D | 43,9°C | 36,8°C | 1 minute at 200°C prior to illumination |
| Oven E | 70,3°C | 54,1°C | 2 minutes at 200°C prior to illumination |
| Oven F | 63,9°C | 50,1°C | 2 minute at 200°C prior to illumination |

Table 5. Average value of the amount of vitamin D₂ in rye loaf with different temperatures prior to illumination

| Temperature prior to illumination | Amount of vitamin D ₂ (µg per bread) |
|-----------------------------------|---|
| Frozen | 14.72 |
| Fridge cold | 19,58 |
| Room temperature | 18,65 |
| Oven (20-40°C) | 18,53 |
| Oven (20-40°C) | 19,59 |

- 5 The results show that there are no great variations in the amount of vitamin D₂ in the rye loaves after illumination regardless of the initial temperature.

On the other hand, the formation of vitamin D₂ did not seem to be favoured if the rye loaves were frozen during illumination with ultraviolet light. However, an increase in the amount of vitamin D₂ is also shown during treatment of frozen loaves. The reason for this is assumed to be that the surface layer of the bread becomes sufficiently hot in order for the vitamin transformation to occur.

10

Example 5

In order to verify that, regardless of the illumination time, toxic doses of vitamin D₂ are not formed, rye loaves were illuminated for extreme periods of time at 302 nm, see table 6 and Figure 7. The amount of vitamin D₂ formed
5 reached a maximum after 12 minutes and the amount was then 31 µg per rye loaf. The acute toxic dose for vitamin D₂ is 1250 µg per day.

Table 6. Amount of vitamin D₂ in rye loaf with different illumination times

| Illumination time (min) | Amount of vitamin D ₂ (µg per rye loaf) |
|-------------------------|--|
| 0,00 | 0,00 |
| 0,03 | 0,43 |
| 0,07 | 0,93 |
| 0,13 | 1,60 |
| 0,27 | 1,99 |
| 0,50 | 6,21 |
| 0,75 | 6,78 |
| 1,00 | 12,90 |
| 2,00 | 19,81 |
| 4,00 | 26,11 |
| 10,00 | 30,38 |
| 15,00 | 30,10 |
| 20,00 | 25,92 |

Example 6

Vegetarian mince containing *Fusarium venenatum* was illuminated with ultraviolet light of 302 nm for seven different periods of time: 2 seconds, 4 seconds, 8 seconds, 16 seconds, 30 seconds, 45 seconds and 60 seconds.

- 5 The distance between the lamp and the vegetarian mince was 2.5 cm. The results of the measurements are shown in table 7 and in the diagram in Figure 8.

Table 7. Amount of vitamin D₂ in vegetarian mince illuminated with 302 nm.

| Illumination time (s) | Amount of vitamin D ₂ (µg per g) |
|-----------------------|---|
| 0 | 0,00 |
| 2 | 1,19 |
| 4 | 1,18 |
| 8 | 1,71 |
| 16 | 7,79 |
| 30 | 8,49 |
| 45 | 15,50 |
| 60 | 18,06 |

CLAIMS

1. Method for vitamin enrichment of foodstuffs containing ascomyceta such as yeast or the mould *Fusarium venenatum*, characterized in that ergosterol in the foodstuff is transformed into vitamin D₂ through placement
5 under an ultraviolet lamp and the illumination of the foodstuff with ultraviolet light with wavelengths between 100-400 nm.
2. Method according to claim 1, characterized in that the source of ultraviolet light is placed at a distance of between 0-2 m from the foodstuff during illumination.
- 10 3. Method according to claim 1 or 2, characterized in that the foodstuff is a bread baked from rising dough.
4. Method according to claim 3, characterized in that the bread is portion packed bread.
5. Method according to claim 1 or 2, characterized in that the
15 foodstuff contains *Fusarium venenatum*.
6. Method according to any of the preceding claims, characterized in that the foodstuff is illuminated with ultraviolet light at a temperature from -70°C to 70°C, preferably at a temperature of 10°C to 70°C.
7. Method according to any of the preceding claims, characterized in
20 that the illumination time is 0,1-60 seconds, preferably 2-30 seconds and more preferably 3-5 seconds.
8. Method according to any of the preceding claims, characterized in that the illumination of the foodstuff takes place in a continuous process, whereupon the foodstuff is transported past the source of ultraviolet light at a
25 speed adapted to achieve a predetermined dose of illumination.




FIG. 1




FIG. 2a






FIG. 2b

214



3/4




FIG.5






FIG.6

4/4

FIG. 7FIG. 8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE2009/050942

A. CLASSIFICATION OF SUBJECT MATTER

IPC: see extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: A23L, A21D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, BIOSIS, CA, COMPDX, EMBASE, MEDLINE, NPL,
AGRICOLA, FSTA, FROSTI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X | US 20080138469 A1 (R. DEGRE ET AL), 12 June 2008 (12.06.2008), claim 1, page 1, column 2, Example 2 | 1-2,5-8 |
| A | page 6, column 2, Example 6, page 8, column 1, Example 7 -- | 1-8 |
| X | JP 8298981 A, ASAHI BREWERIES LTD, 1996-11-19: (abstract) Retrieved from: WPI database, WEEK 199705, AN 1997-045816 page 3, column 1, paragraph 0015, page 3, column 2, paragraph 0024 | 1-2,5-8 |
| A | page 4, column 2, paragraph 0031 -- | 1-8 |

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 November 2009

Date of mailing of the international search report

19-11-2009

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Johan Kjellgren / MRo
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE2009/050942

| C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|---|---|-----------------------|
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| X | Kawazoe T 'Efficient transformation of ergosterol to vitamin D-2 by ultraviolet light irradiation', Journal of the Japanese Society for Food Science and Technology, 1995, Vol. 42, No. 4, pages 262-267, (abstract) BIOSIS[online]: BIOSIS (R) Biological Abstracts, Inc USA [retrived on 30 januari 2009]. AN: PREV199598311933. Retrived from EPOQUE NPL database abstract | 1-2,5-8 |
| A | abstract -- | 1-8 |
| A | CN 101057658 A, WANJING XIAMEN FOODSTUFFS CO L, 2007-10-24; (abstract) Retrieved from: EPODOC database -- ----- | 8 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE2009/050942

International patent classification (IPC)

A23L 1/303 (2006.01)

A23L 1/28 (2006.01)

Download your patent documents at www.prv.se

The cited patent documents can be downloaded:

- From "Cited documents" found under our online services at www.prv.se (English version)
- From "Anförda dokument" found under "e-tjänster" at www.prv.se (Swedish version)

Use the application number as username. The password is **YOANJCHPZM**.

Paper copies can be ordered at a cost of 50 SEK per copy from PRV InterPat (telephone number 08-782 28 85).

Cited literature, if any, will be enclosed in paper form.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/SE2009/050942

US 20080138469 A1 12/06/2008 NONE
