

**(12) STANDARD PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2013226081 B2**

(54) Title  
**Compositions and kits for molecular counting**

(51) International Patent Classification(s)  
**C12Q 1/68** (2006.01)      **C12N 15/11** (2006.01)

(21) Application No: **2013226081**      (22) Date of Filing: **2013.02.27**

(87) WIPO No: **WO13/130674**

(30) Priority Data

(31) Number      (32) Date      (33) Country  
**61/745,385**      **2012.12.21**      **US**  
**61/603,921**      **2012.02.27**      **US**

(43) Publication Date: **2013.09.06**  
(44) Accepted Journal Date: **2018.06.14**

(71) Applicant(s)  
**Cellular Research, Inc.**

(72) Inventor(s)  
**Fu, Glenn K.;Fodor, Stephen P.A.;Wilhelmy, Julie**

(74) Agent / Attorney  
**Pizzeys Patent and Trade Mark Attorneys Pty Ltd, PO Box 291, WODEN, ACT, 2606, AU**

(56) Related Art  
**DE 102008025656 A1**





---

**Published:**

— *with international search report (Art. 21(3))*

## COMPOSITIONS AND KITS FOR MOLECULAR COUNTING

### CROSS-REFERENCE TO RELATED APPLICATIONS

**[0001]** This application claims the benefit of U.S. Provisional Application No. 61/603,921 filed February 27, 2012, and U.S. Provisional Application No. 61/745,385, filed December 21, 2012, both of which are incorporated by reference herein in their entirety.

### FIELD OF THE INVENTION

**[0002]** Methods and uses of molecular counting are disclosed. Molecules can be counted by sequencing and tracking the number of occurrences of a target molecule. Molecules can also be counted by hybridization of the molecule to a solid support and detection of the hybridized molecules. In some instances, the molecules to be counted are labeled. The molecules to be counted may also be amplified.

### BACKGROUND OF THE INVENTION

**[0003]** Accurate determination of the quantity of nucleic acids is necessary in a wide variety of clinical and research measurements. When dissolved in solution, the average concentration of nucleic acids (RNA or DNA) can be determined by UV light absorbance spectrophotometry or by fluorescent DNA-binding stains. However, the measurement required is often not just for the total amount of nucleic acids present, but specifically for one or more species of interest contained and mixed with all of the other nucleic acids within the sample. In these cases, the nucleic acid molecule of interest is usually distinguished from all of the other nucleic acids through a defined sequence of nucleotides that is unique to the species of interest. A short synthetic ribo- or deoxyribo- oligonucleotide with a complementary sequence to the nucleic acid of interest can be used for its detection and identification. For instance, the Polymerase Chain Reaction (PCR) uses a pair of these oligonucleotides to serve as annealing primers for repeated cycles of DNA polymerization mediated by DNA polymerase enzymes. DNA microarrays are another common detection method where oligonucleotides are immobilized on solid supports to hybridize to DNA molecules bearing complementary sequences. Although both PCR and microarray methods are capable of specific detection, accurate determination of the quantity of the detected molecules is difficult (especially when it is present in low abundance or when contained within a large background of other nucleic acids). In the case of PCR (also sometimes referred to as quantitative-PCR, qPCR, TaqMan, or real-time PCR), the amount of amplified DNA molecules represents an estimate of its concentration in the starting solution. In the case of microarrays, the amount of DNA hybridized is an estimate of its concentration in solution.

In both cases, only relative measurements of concentration can be made, and the absolute number of copies of nucleic acid in the sample cannot be precisely determined. However, when reference nucleic acids of pre-determined concentrations are included in the test, relative comparisons can be made to this standard reference to estimate the absolute number of copies of nucleic acids being detected.

**[0004]** Digital PCR is one method that can be used to determine the absolute number of DNA molecules of a particular nucleotide sequence (Sykes et al. *Biotechniques* 13: 444-449 (1992), Vogelstein et al. *Digital PCR. Proc Natl Acad Sci USA* 96: 9236-9241 (1999)). In this method, the nucleic acid solution is diluted and stochastically partitioned into individual containers so that there is on average less than one molecule in every two containers. PCR is then used to detect the presence of the nucleic acid molecule of interest in each container. If quantitative partitioning is assumed, the dynamic range is governed by the number of containers available for stochastic separation. Micro fabrication and picoliter-sized emulsion droplets can be used to increase the number of containers available thereby extending the measurement dynamic range (Fan et al. *Am J Obstet Gynecol* 200: 543 e541-547 (2009), Kalinina et al. *Nucleic Acids Res* 25: 1999-2004 (1997)). Due to the physical constraints of manufacturing large numbers of separate containers and in carrying out these larger numbers of reactions, in practice the digital PCR method is limited to investigations on only a small number of different DNA molecules at a time.

**[0005]** Recently, a new method to determine the absolute quantity of DNA molecules has been demonstrated where identical copies of individual DNA molecules can be counted after the stochastic attachment of a set of diverse nucleic acid labels (Fu et al. *Proc Natl Acad Sci USA* 108: 9026-9031 (2011)). Unlike digital PCR, this is a highly parallel method capable of counting many different DNA molecules simultaneously. In this method, each copy of a molecule randomly attaches to a short nucleic acid label by choosing from a large, non-depleting reservoir of diverse labels. The subsequent diversity of the labeled molecules is governed by the statistics of random choice, and depends on the number of copies of identical molecules in the collection compared to the number of kinds of labels. Once the molecules are labeled, they can be amplified so that simple present/absent threshold detection methods can be used for each. Counting the number of distinctly labeled targets reveals the original number of molecules of each species. Unlike digital PCR, which stochastically expands identical molecules into physical space, the method of stochastic labeling expands identical molecules into chemical space. An important distinction from digital PCR is that the stochastic labeling method does not require the challenging physical separation of identical

molecules into individual physical containers. The approach is practical, and after labeling, a simple detector device such as a microarray with complementary probe sequences to the labels can be used to identify and count the number of labels present. In addition, when stochastic labels are attached to DNA molecules that are prepared for DNA sequencing readouts, the labeling sequence can serve as discreet counting tags for absolute quantitation, or as unique identifiers to distinguish each originally tagged template from its amplified daughter molecules (Kinde et al. Proc Natl Acad Sci USA 108: 9530-9535 (2011)).

## SUMMARY OF THE INVENTION

**[0006]** In some embodiments is a digital reverse transcription method comprising: a) contacting a sample comprising a plurality of RNA molecules with a plurality of oligonucleotide tags to produce a labeled-RNA molecule, wherein: the plurality of RNA molecules comprise at least 2 mRNA molecules of different sequences; the plurality of oligonucleotide tags comprises at least 2 oligonucleotide tags of different sequences; and the plurality of oligonucleotide tags comprises an oligodT sequence; b) conducting a first strand synthesis reaction by contacting the labeled-RNA molecules with a reverse transcriptase enzyme to produce a labeled-cDNA molecule; and c) detecting the labeled-cDNA molecule by hybridizing the labeled-cDNA molecule to a solid support.

**[0007]** In some embodiments is a stochastic label-based hybridization chain reaction method comprising stochastically labeling one or more nucleic acid molecules with a plurality of hairpin oligonucleotide tags, wherein the hairpin oligonucleotide tag comprises an overhang; and the one or more nucleic acid molecules act as initiators for a hybridization chain reaction.

**[0008]** At least a portion of the hairpin oligonucleotide tag may hybridize to at least a portion of the one or more nucleic acid molecules. The hairpin oligonucleotide tag may comprise an oligodT sequence. The one or more nucleic acid molecules may comprise one or more adapters. At least a portion of the hairpin oligonucleotide tag may hybridize to at least a portion of the one or more adapters. At least one hairpin oligonucleotide tag of the plurality of hairpin oligonucleotide tags may comprise one or more labels. At least one hairpin oligonucleotide tag of the plurality of hairpin oligonucleotide tags may comprise two or more labels.

**[0009]** Each hairpin oligonucleotide tag of the plurality of hairpin oligonucleotide tags may comprise one or more labels. Each hairpin oligonucleotide tag of the plurality of hairpin oligonucleotide tags may comprise two or more labels. In some instances, the hairpin oligonucleotide tag does not comprise a label.

**[0010]** The plurality of hairpin oligonucleotide tags may comprise one or more hairpin oligonucleotide tags with a 5' overhang, hairpin oligonucleotide tags with a 3' overhang, or a combination thereof.

**[0011]** The stem portion of the hairpin oligonucleotide tag can be one or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be two or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be three or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be four or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be five or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be six or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be seven or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be eight or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be nine or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be ten or more nucleotides in length. The stem portion of the hairpin oligonucleotide tag can be 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100 or more nucleotides in length.

**[0012]** The loop portion of the hairpin oligonucleotide tag can be one or more nucleotides in length. The the loop portion of the hairpin oligonucleotide tag can be two or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be three or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be four or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be five or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be six or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be seven or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be eight or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be nine or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be ten or more nucleotides in length. The loop portion of the hairpin oligonucleotide tag can be 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100 or more nucleotides in length.

**[0013]** The hairpin oligonucleotide tag may comprise a unique identifier region. The unique identifier region can be in the loop portion of the hairpin oligonucleotide tag. The unique identifier region can be in the stem portion of the hairpin oligonucleotide tag. The unique identifier region can be in the overhang portion of the hairpin oligonucleotide tag.

**[0014]** The label may comprise a unique identifier region.

**[0015]** In some embodiments the oligonucleotide tag further comprises a unique identifier region. In some embodiments the unique identifier region is at least one nucleotide in length.

In some embodiments the oligonucleotide tag further comprises a universal primer binding site. In some embodiments the oligonucleotide tag is at least one nucleotide in length.

**[0016]** In some embodiments the solid support is an array. In some embodiments the solid support is an addressable array. In some embodiments the solid support is an Affymetrix 3K tag array, Arrayjet non-contact printed array, or Applied Microarrays Inc (AMI) array. In some embodiments the solid support is a bead.

**[0017]** Further disclosed herein is a cell analysis method comprising: a) contacting a sample comprising a plurality of molecules with a plurality of oligonucleotide tags to produce a labeled-molecule, wherein: the plurality of molecules comprise at least 2 molecules of different sequences; the plurality of oligonucleotide tags comprises at least 2 oligonucleotide tags of different sequences; and the sample is from at least one cell; and b) detecting the labeled-molecule by hybridizing the labeled-molecule to a solid support.

**[0018]** In some embodiments is a clonal amplification method comprising: a) stochastically labeling a plurality of molecules with a plurality of oligonucleotide tags to produce a labeled-molecule, wherein: the plurality of molecules comprise at least 2 molecules of different sequences; and the plurality of oligonucleotide tags comprises at least 2 oligonucleotide tags of different sequences; b) amplifying the labeled-molecules to produce a labeled-amplicon; and c) detecting the labeled-amplicon.

**[0019]** Further disclosed herein is a kit comprising: a) a plurality of oligonucleotide tags, wherein the oligonucleotide tag of the plurality of oligonucleotide tags comprises: a target specific region; and a unique identifier region; and b) an enzyme.

**[0020]** In some embodiments the enzyme is a reverse transcriptase enzyme. In some embodiments the enzyme is a ligase. In some embodiments the enzyme is a polymerase. In some embodiments the enzyme is an RNase. In some embodiments the enzyme is a DNase. In some embodiments the enzyme is an endonuclease.

**[0021]** In some embodiments the oligonucleotide tag is at least 25 nucleotides in length. In some embodiments the unique identifier region is at least 10 nucleotides in length. In some embodiments the target specific region is at least 10 nucleotides in length. In some embodiments the target specific region comprises an oligodT sequence. In some embodiments the oligonucleotide tag further comprises a universal primer binding site.

**[0022]** In some embodiments the kit further comprises a support. In some embodiments the support is a semi-solid support. In some embodiments the support is a solid support. In some embodiments the solid support is an array. In some embodiments the support is an addressable array. In some embodiments the support is an Affymetrix 3K tag array, Arrayjet

non-contact printed array, or Applied Microarrays Inc (AMI) array. In some embodiments the support is a bead.

**[0023]** In some embodiments the kit further comprises a primer. In some embodiments the primer is a universal primer. In some embodiments the primer binds to the oligonucleotide tag. In some embodiments the primer binds to the universal primer binding site of the oligonucleotide tag.

**[0024]** In some embodiments the kit further comprises a control oligo. In some embodiments the control oligo comprises at least 15 nucleotides. In some embodiments the control oligo is a bright hybridization control oligo. In some embodiments the control oligo is a spike-in template control. In some embodiments the oligonucleotide tag further comprises a label.

**[0025]** In some embodiments the primer further comprises a label. In some embodiments the control oligo further comprises a label. In some embodiments the label is a dye label. In some embodiments the label is a Cy3 dye. In some embodiments the label is a Tye563 dye.

**[0026]** In some embodiments the kit further comprises a buffer.

**[0027]** In some embodiments the kit further comprises a carrier.

**[0028]** In some embodiments the kit further comprises a detergent.

**[0029]** Further disclosed herein is a system for determining the absolute quantity of a plurality of nucleic acid molecules. The system may comprise a) a plurality of oligonucleotide tags; and b) a detector for detecting at least a portion of the oligonucleotide tags.

**[0030]** The detector may comprise an array detector, fluorescent reader, non-fluorescent detector, CR reader, or scanner. In some embodiments the method further comprises the fluorescent reader is a Sensovation or AG fluorescent reader. In some embodiments the method further comprises the scanner is a flatbed scanner.

**[0031]** The system may further comprise a thermal cycler. In some embodiments the system further comprises a sequencer. In some embodiments the system further comprises a hybridization chamber.

**[0032]** The system may further comprise a computer. In some embodiments the computer comprises a memory device. In some embodiments the memory device is capable of storing data. In some embodiments the system further comprises a software program. In some embodiments the system further comprises a computer-readable program.

**[0033]** In some embodiments the oligonucleotide tag further comprises a unique identifier region. In some embodiments the unique identifier region is at least 10 nucleotides in length.

In some embodiments the unique identifier region cannot hybridize to the molecule. In some embodiments the oligonucleotide tag further comprises a universal primer binding site. In some embodiments the oligonucleotide tag is at least 20 nucleotides in length. In some embodiments the oligonucleotide tag further comprises a target specific region. In some embodiments the target specific region comprises an oligodT sequence. In some embodiments the target specific region is at least 10 nucleotides in length. In some embodiments the method further comprises conducting a first strand synthesis reaction to produce a labeled-cDNA molecule.

**[0034]** In some embodiments the amplifying the labeled-molecule comprises conducting a polymerase chain reaction. Alternatively, amplifying the labeled-molecule may comprise conducting a non-PCR based amplification reaction. Amplifying the labeled-molecule may comprise exponential amplification of the labeled-molecule. Amplifying the labeled-molecule may comprise linear amplification of the labeled molecule. Amplifying the labeled-molecule may comprise hybridization chain reaction (HCR) based amplification method.

**[0035]** Amplifying the labeled-molecule may comprise amplifying at least the label portion of the labeled molecule, the molecule portion of the labeled molecule, or a combination thereof.

**[0036]** In some embodiments the method further comprises conducting a polymerase chain reaction on the labeled-molecule or any product thereof to produce a double-stranded labeled-molecule. In some embodiments conducting the polymerase chain reaction comprises annealing a first target specific primer to the labeled-molecule or any product thereof. In some embodiments conducting the polymerase chain reaction further comprises annealing a universal primer to the universal primer binding site of the oligonucleotide tag. In some embodiments the polymerase chain reaction comprises absolute PCR, HD-PCR, Next Gen PCR, digital RTA, or any combination thereof. In some embodiments the method comprises conducting a nested PCR reaction on the double-stranded labeled-cDNA molecule. In some embodiments conducting the nested PCR reaction comprises denaturing the labeled-molecule or any product thereof to produce a denatured single-stranded labeled-molecule or any product thereof. In some embodiments conducting the nested PCR reaction further comprises annealing a second target specific primer to the denatured single-stranded labeled-molecule or any product thereof. In some embodiments conducting the nested PCR reaction further comprises annealing a universal primer to the universal primer binding site of the oligonucleotide tag.

**[0037]** In some embodiments the method further comprises conducting a sequencing

reaction to determine the sequence of at least a portion of the oligonucleotide tag, at least a portion of the labeled-molecule, a product thereof, a complement thereof, a reverse complement thereof, or any combination thereof.

[0038] In some embodiments detecting the labeled-molecules or any products thereof comprises an array detector, fluorescent reader, non-fluorescent detector, CR reader, or scanner. In some embodiments the molecule is a nucleic acid molecule.

[0039] In some embodiments the nucleic acid molecule is a DNA molecule. In some embodiments the nucleic acid molecule is an RNA molecule. In some embodiments the molecule is a peptide. In some embodiments the peptide is a polypeptide.

[0040] In some embodiments the plurality of molecules is from a cell. In some embodiments the sample is from a single cell. In some embodiments the sample is from less than about 100 cells. In some embodiments the sample is from less than about 50 cells. In some embodiments the sample is from less than about 20 cells. In some embodiments the sample is from less than about 10 cells. In some embodiments the sample is from less than about 5 cells. In some embodiments the cell is a mammalian cell. In some embodiments the cell is a human cell. In some embodiments the cell is from a subject suffering from a disease or condition. In some embodiments the disease or condition is cancer. In some embodiments the disease or condition is a pathogenic infection. In some embodiments the disease or condition is a genetic disorder. In some embodiments the cell is from a healthy subject. In some embodiments the cell is a diseased cell. In some embodiments the diseased cell is a cancerous cell. In some embodiments the cell is a healthy cell. In some embodiments the cell is not a diseased or infected cell. In some embodiments the labeled-molecules are produced by stochastic labeling.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0041] The skilled artisan will understand that the drawings described below are for illustration purposes only. The drawings are not intended to limit the scope of the present teachings in any way.

[0042] FIG. 1 shows a schematic of labeling and detection of a target molecule

[0043] FIG. 2 shows signals for the detection of labels in hybridized molecules

[0044] FIG. 3 shows signals for the detection of labels in hybridized molecules

[0045] FIG. 4 shows signals for the detection of labels in hybridized molecules

[0046] FIG. 5 shows signals for the detection of labels in hybridized molecules

[0047] FIG. 6 shows signals for the detection of labels in hybridized molecules

[0048] FIG. 7 shows a schematic of detection of a labeled molecule by an array detector

[0049] FIG. 8 shows a schematic of stochastic labeling of a plurality of molecules

[0050] FIG. 9 Exemplary PCR primer consisting of a universal PCR sequence, a short label sequence and a target or gene-specific sequence.

[0051] FIG. 10 shows a schematic for the synthesis of of oligonucleotide tags

[0052] FIG. 11A shows a schematic for the synthesis of of oligonucleotide tags without target-specific sequence

[0053] FIG. 11B-D shows a schematic for the synthesis of of oligonucleotide tags

[0054] FIG. 12A-B depict degenerate oligonucleotide tags

[0055] FIG. 13 Additional Examples of Labeled Primers. A) Labeled Primer without generic primer sequence. B) Labeled Primer with universal target sequence

[0056] FIG. 14 Absolute PCR Protocol

[0057] FIG. 15 Formation of Primer Dimers

[0058] FIG. 16 Method to prevent the formation of primer artifacts

[0059] FIG. 17 Differences between a standard array and a digital array

[0060] FIG. 18 Digital microarray probes – detection using a combination of gene and label sequences

[0061] FIG. 19 Absolute quantitation of mRNA molecules by counting individual DNA molecules

[0062] FIG. 20 Digital microarray for RNA expression

[0063] FIG. 21 Digital microarray for DNA copy number

[0064] FIG. 22 Digital microarray for microRNAs

[0065] FIG. 23A Digital microarray for single cell pre-implantation genetic diagnosis (PGD) (a) cycle 0; (b) cycle 5; (c) cycle 10; (d) cycle 15

[0066] FIG. 23B shows a schematic of a method for single cell pre-implantation genetic diagnosis (PGD)

[0067] FIG. 24 Digital microarray for measuring fetal aneuploidy in maternal circulating nucleic acids –e.g., Trisomy 21

[0068] FIG. 25 Absolute quantitation of mRNA molecules by counting individual DNA molecules

[0069] FIG. 26 Labeling with an “inert” primer

[0070] FIG. 27 Emulsion PCR to prevent artifacts from out-competing cDNAs during amplification

[0071] FIG. 28 A method that does not rely on homopolymer tailing

[0072] FIG. 29 Linear amplification methods

[0073] FIG. 30 Labeling with strand switching

[0074] FIG. 31 Labeling by random priming

[0075] FIG. 32A-B show the results for the optimization of cDNA synthesis

[0076] FIG. 33 Schematic of stochastic labeling followed by HCR detection of nucleic acid molecules

[0077] FIG. 34 Schematic of stochastic labeling of hairping HCR oligonucleotides

[0078] FIG. 35 Schematic of the serial dilution scheme for the titration experiment with serial dilutions of kanamycin RNA

[0079] FIG. 36A-H Shows the scatter plots of results for the titration experiment with serial dilutions of kanamycin RNA

[0080] FIG. 37 Shows the Correlation graph for the titration experiment with serial dilutions of kanamycin RNA

[0081] FIG. 38 Schematic of the serial dilution scheme for the titration experiment with serial dilutions of human liver RNA to measure GAPDH expression

[0082] FIG. 39A-H Shows the scatter plots of results for the titration experiment with serial dilutions of human liver RNA to measure GAPDH expression

[0083] FIG. 40 Shows the correlation graph for the titration experiment with serial dilutions of human liver RNA to measure GAPDH expression

[0084] FIG. 41A-D Shows the scatter plots of results for the accurate measurements of control bacterial genes

[0085] FIG. 42 Shows the scatter plot for the validation of kanamycin counts by digital PCR experiment

[0086] FIG. 43 Schematic of the method for absolute quantitation of mRNA molecules directly from cell lysates

#### **DETAILED DESCRIPTION OF THE INVENTION**

[0087] Reference will now be made in detail to exemplary embodiments of the invention. While the invention will be described in conjunction with the exemplary embodiments, it will be understood that they are not intended to limit the invention to these embodiments. On the contrary, the invention is intended to cover alternatives, modifications and equivalents, which may be included within the spirit and scope of the invention.

[0088] The invention has many preferred embodiments and relies on many patents, applications and other references for details known to those of the art. Therefore, when a patent, application, or other reference, such as a printed publication, is cited or repeated below, it should be understood that it is incorporated by reference in its entirety for all

purposes and particularly for the proposition that is recited.

**[0089]** An individual is not limited to a human being, but may also be other organisms including, but not limited to, mammals, plants, bacteria, or cells derived from any of the above.

**[0090]** Throughout this disclosure, various aspects of this invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

**[0091]** Disclosed herein are methods, kits, and systems for detecting and/or quantifying molecules in a sample. In some instances, methods, kits, and systems for individually counting molecules in a sample are provided. Alternatively, methods, kits, and systems for determining the expression level of a gene or gene product are provided. In some instances, the methods comprise the attachment of an oligonucleotide tag to a molecule (e.g., RNA, DNA, protein) to form a labeled molecule. The oligonucleotide tag can comprise a target specific region, unique identifier region, universal primer binding region, detectable label region, or any combination thereof. In some instances, the attachment of the oligonucleotide tag to the molecule results in the formation of a unique junction comprising at least a portion of the oligonucleotide tag and at least a portion of the molecule. An expression level of a gene or gene product can be determined by detecting and/or quantifying at least a portion of the labeled molecule (e.g., unique junction, oligonucleotide tag, molecule). The absolute quantity of a target molecule can also be determined by detecting the number of unique oligonucleotide tags of the labeled molecules and/or the number of unique junctions in the labeled molecules.

**[0092]** Further disclosed herein are absolute PCR methods for amplifying and/or quantifying one or more molecules. A schematic of the absolute PCR protocol is depicted in FIG. 14. As shown in Step 1 of FIG. 14, an oligonucleotide tag (1404) comprising a universal primer binding site (1401), unique identifier region (1402) and a target specific region (1403) is hybridized to a target molecule (1405). As shown in Step 2 of FIG. 14, the oligonucleotide tag (1404) may act as a primer and a copy of the target molecule (1405) can

be synthesized by primer extension by a polymerase (e.g., DNA polymerase) to produce an amplicon (1406). The amplicon (1406) may comprise a universal primer binding site (1401), unique identifier region (1402) and a complement of target molecule (1411). As shown in Step 3 of FIG. 14, a reverse primer (1407) can anneal to the the amplicon (1406). As shown in Step 4 of FIG. 14, the amplicon (1406) can act as a template for synthesizing second amplicon (1408). The second amplicon (1408) can comprise a copy of the target molecule (1411') and a complement of the universal primer binding site (1401') and a complement of the unique identifier region (1402'). As shown in Step 5 of FIG. 14, the amplicons (1406, 1408) can act as templates for subsequent amplification with a forward primer (1409) comprising the universal primer binding site and a reverse primer (1410) comprising a target specific sequence. Each subsequent amplicon comprises the unique identifier region (1402). By incorporating the unique identifier region into each amplicon, the amplification efficiency and/or amplification bias can be determined. In addition, the quantity of the target molecules can be determined by counting the number of different unique identifier regions that are associated with each target molecule. The absolute PCR method can be used for subsequent analysis of the target molecules (Step 6 of FIG. 14). For example, the amplicons produced by the absolute PCR method can be used to detect and/or quantify one or more target molecules. Unincorporated oligonucleotide tags can be removed by purification of the amplicons.

**[0093] I. Labeling of molecules with oligonucleotide tags**

**[0094] A. Stochastic labeling of molecules**

**[0095]** The methods disclosed herein comprise the attachment of oligonucleotide tags to molecules in a sample. In some instances, attachment of the oligonucleotide tags to the molecules comprises stochastic labeling of the molecules. Methods for stochastically labeling molecules can be found, for example, in U.S. Serial Numbers 12/969,581 and 13/327,526. Generally, the stochastic labeling method comprises the random attachment of a plurality of oligonucleotide tags to one or more molecules. The plurality oligonucleotide tags are provided in excess of the one or more molecules to be labeled. In stochastic labeling, each individual molecule to be labeled has an individual probability of attaching to the plurality of oligonucleotide tags. The probability of each individual molecule to be labeled attaching to a particular tag can be about the same as any other individual molecule to be labeled. Accordingly, in some instances, the probability of any of the molecules in a sample finding any of the tags is assumed to be equal, an assumption that can be used in mathematical calculations to estimate the number of molecules in the sample. In some circumstances the

probability of attaching can be manipulated by, for example electing tags with different properties that would increase or decrease the binding efficiency of that tag with a individual molecule. The oligonucleotide tags can also be varied in numbers to alter the probability that a particular tag will find a binding partner during the stochastic labeling. For example one tag may be overrepresented in a pool of tags, thereby increasing the chances that the overrepresented tag finds at least one binding partner.

**[0096]** *B. Methods for attaching an oligonucleotide tag to a molecule*

**[0097]** Attachment of an oligonucleotide tag to a molecule can occur by a variety of methods, including, but not limited to, hybridization of the oligonucleotide tag to the molecule. In some instances, the oligonucleotide tag comprises a target specific region. The target specific region can comprise a sequence that is complementary to at least a portion of the molecule to be labeled. The target specific region can hybridize to the molecule, thereby producing a labeled molecule.

**[0098]** Attachment of the oligonucleotide tag to a molecule can occur by ligation. Ligation techniques comprise blunt-end ligation and sticky-end ligation. Ligation reactions can include DNA ligases such as DNA ligase I, DNA ligase III, DNA ligase IV, and T4 DNA ligase. Ligation reactions can include RNA ligases such as T4 RNA ligase I and T4 RNA ligase II.

**[0099]** Methods of ligation are described, for example in Sambrook et al. (2001) and the New England BioLabs catalog both of which are incorporated herein by reference for all purposes. Methods include using T4 DNA Ligase which catalyzes the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini in duplex DNA or RNA with blunt and sticky ends; Taq DNA Ligase which catalyzes the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini of two adjacent oligonucleotides which are hybridized to a complementary target DNA; E. coli DNA ligase which catalyzes the formation of a phosphodiester bond between juxtaposed 5'-phosphate and 3'-hydroxyl termini in duplex DNA containing cohesive ends; and T4 RNA ligase which catalyzes ligation of a 5' phosphoryl-terminated nucleic acid donor to a 3' hydroxyl-terminated nucleic acid acceptor through the formation of a 3'→5' phosphodiester bond, substrates include single-stranded RNA and DNA as well as dinucleoside pyrophosphates; or any other methods described in the art. Fragmented DNA may be treated with one or more enzymes, for example, an endonuclease, prior to ligation of adaptors to one or both ends to facilitate ligation by generating ends that are compatible with ligation.

**[00100]** In some instances, both ends of the oligonucleotide tag are attached to the

molecule. For example, both ends of the oligonucleotide tag can be hybridized and/or ligated to one or more ends of the molecule. In some instances, attachment of both ends of the oligonucleotide tag to both ends of the molecule results in the formation of a circularized labeled-molecule. Both ends of the oligonucleotide tag can also be attached to the same end of the molecule. For example, the 5' end of the oligonucleotide tag is ligated to the 3' end of the molecule and the 3' end of the oligonucleotide tag is hybridized to the 3' end of the molecule, resulting in a labeled-molecule with a hairpin structure at one end. In some instances the oligonucleotide tag is attached to the middle of the molecule.

**[00101]** In some instances, attachment of the oligonucleotide tag to the molecule comprises the use of one or more adaptors. Adaptors can comprise a target specific region on one end, which allows the attachment of the adaptor to the molecule, and an oligonucleotide tag specific region on the other end, which allows attachment of the oligonucleotide tag to the adaptor. Adaptors can be attached to the molecule and/or oligonucleotide by methods including, but not limited to, hybridization and/or ligation.

**[00102]** Methods for ligating adaptors to fragments of nucleic acid are well known. Adaptors may be double-stranded, single-stranded or partially single-stranded. In preferred aspects adaptors are formed from two oligonucleotides that have a region of complementarity, for example, about 10 to 30, or about 15 to 40 bases of perfect complementarity, so that when the two oligonucleotides are hybridized together they form a double stranded region. Optionally, either or both of the oligonucleotides may have a region that is not complementary to the other oligonucleotide and forms a single stranded overhang at one or both ends of the adaptor. Single-stranded overhangs may preferably be about 1 to about 8 bases, and most preferably about 2 to about 4. The overhang may be complementary to the overhang created by cleavage with a restriction enzyme to facilitate "sticky-end" ligation. Adaptors may include other features, such as primer binding sites and restriction sites. In some aspects the restriction site may be for a Type IIS restriction enzyme or another enzyme that cuts outside of its recognition sequence, such as EcoP151 (see, Mucke *et al. J Mol Biol* 2001, 312(4):687-698 and US 5,710,000 which is incorporated herein by reference in its entirety).

**[00103]** The oligonucleotide tag can be attached to any region of a molecule. For example, the oligonucleotide can be attached to the 5' or 3' end of a polynucleotide (e.g., DNA, RNA). For example, the target-specific region of the oligonucleotide tag comprises a sequence that is complementary to a sequence in the 5' region of the molecule. The target-specific region of the oligonucleotide tag can also comprise a sequence that is complementary to a sequence in

the 3' region of the molecule. In some instances, the oligonucleotide tag is attached a region within a gene or gene product. For example, genomic DNA is fragmented and an oligonucleotide tag is attached to the fragmented DNA. In other instances, an RNA molecule is alternatively spliced and the oligonucleotide tag is attached to the alternatively spliced variants. In another example, the polynucleotide is digested and the oligonucleotide tag is attached to the digested polynucleotide. In another example, the target-specific region of the oligonucleotide tag comprises a sequence that is complementary to a sequence within the molecule.

**[00104] II. Reverse Transcription**

**[00105]** In some instances, the methods disclosed herein comprise attachment of an oligonucleotide tag to an RNA molecule to produce a labeled-RNA molecule. The methods disclosed herein can further comprise reverse transcription of the labeled-RNA molecule to produce a labeled-cDNA molecule. In some instances, at least a portion of the oligonucleotide tag acts as a primer for the reverse transcription reaction. For example, as shown in FIG. 1, Steps 1A-B, an oligonucleotide tag comprising an oligodT sequence hybridizes to the polyA tail of an mRNA molecule. The oligodT portion of the oligonucleotide tag acts as a primer for first strand synthesis of the cDNA molecule.

**[00106]** In some instances the labeled cDNA molecule can be used as a molecule for a new stochastic labeling reaction. The labeled cDNA can have a first tag or set of tags from attachment to the RNA prior to reverse transcription and a second tag or set of tags attached to the cDNA molecule. These multiple labeling reactions can, for example, be used to determine the efficiency of events that occur between the attachment of the first and second tags, e.g. an optional amplification reaction or the reverse transcription reaction.

**[00107]** In another example, an oligonucleotide tag is attached to the 5' end of an RNA molecule to produce a labeled-RNA molecule. Reverse transcription of the labeled-RNA molecule can occur by the addition of a reverse transcription primer. In some instances, the reverse transcription primer is an oligodT primer, random hexanucleotide primer, or a target-specific oligonucleotide primer. Generally, oligo(dT) primers are 12–18 nucleotides in length and bind to the endogenous poly(A)+ tail at the 3' end of mammalian mRNA. Random hexanucleotide primers can bind to mRNA at a variety of complementary sites. Target-specific oligonucleotide primers typically selectively prime the mRNA of interest.

**[00108]** In some instances, the method comprises repeatedly reverse transcribing the labeled-RNA molecule to produce multiple labeled-cDNA molecules. The methods disclosed herein can comprise conducting at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15,

16, 17, 18, 19, or 20 reverse transcription reactions. The method can comprise conducting at least about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 reverse transcription reactions.

**[00109] III. Amplification of labeled molecules**

**[00110]** The methods disclosed herein may comprise amplification of the labeled molecules to produce labeled amplicons. Amplification of the labeled molecules can comprise PCR-based methods or non-PCR based methods. Amplification of the labeled molecules may comprise exponential amplification of the labeled molecules. Amplification of the labeled molecules may comprise linear amplification of the labeled molecules.

**[00111]** In some instances, amplification of the labeled molecules comprises non-PCR based methods. Examples of non-PCR based methods include, but are not limited to, multiple displacement amplification (MDA), transcription-mediated amplification (TMA), nucleic acid sequence-based amplification (NASBA), strand displacement amplification (SDA), real-time SDA, rolling circle amplification, or circle-to-circle amplification.

**[00112]** Amplification of the labeled molecules may comprise hybridization chain reaction (HCR) based methods (Dirks and Pierce, *PNAS*, 2004; Zhang et al., *Anal Chem*, 2012). HCR based methods may comprise DNA-based HCR. HCR based methods may comprise one or more labeled probes. The one or more labeled probes may comprise one or more oligonucleotide tags disclosed herein.

**[00113]** In some instances, the methods disclosed herein further comprise conducting a polymerase chain reaction on the labeled-molecule (e.g., labeled-RNA, labeled-DNA, labeled-cDNA) to produce a labeled-amplicon. The labeled-amplicon can be double-stranded molecule. The double-stranded molecule can comprise a double-stranded RNA molecule, a double-stranded DNA molecule, or a RNA molecule hybridized to a DNA molecule. One or both of the strands of the double-stranded molecule can comprise the oligonucleotide tag. Alternatively, the labeled-amplicon is a single-stranded molecule. The single-stranded molecule can comprise DNA, RNA, or a combination thereof. The nucleic acids of the present invention can comprise synthetic or altered nucleic acids.

**[00114]** The polymerase chain reaction can be performed by methods such as PCR, HD-PCR, Next Gen PCR, digital RTA, or any combination thereof. Additional PCR methods include, but are not limited to, allele-specific PCR, Alu PCR, assembly PCR, asymmetric PCR, droplet PCR, emulsion PCR, helicase dependent amplification HDA, hot start PCR, inverse PCR, linear-after-the-exponential (LATE)-PCR, long PCR, multiplex PCR, nested PCR, hemi-nested PCR, quantitative PCR, RT-PCR, real time PCR, single cell PCR, and

touchdown PCR.

**[00115]** In some instances, conducting a polymerase chain reaction comprises annealing a first target specific primer to the labeled-molecule. Alternatively or additionally, conducting a polymerase chain reaction further comprises annealing a universal primer to a universal primer binding site region of the oligonucleotide tag, wherein the oligonucleotide tag is on a labeled-molecule or labeled-amplicon. The methods disclosed herein can further comprise annealing a second target specific primer to the labeled-molecule and/or labeled-amplicon.

**[00116]** In some instances, the method comprises repeatedly amplifying the labeled-molecule to produce multiple labeled-amplicons. The methods disclosed herein can comprise conducting at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amplification reactions. Alternatively, the method comprises conducting at least about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amplification reactions.

**[00117]** Other suitable amplification methods include the ligase chain reaction (LCR) (for example, Wu and Wallace, *Genomics* 4, 560 (1989), Landegren et al., *Science* 241, 1077 (1988) and Barringer et al. *Gene* 89:117 (1990)), transcription amplification (Kwoh et al., *Proc. Natl. Acad. Sci. USA* 86, 1173 (1989) and W088/10315), self-sustained sequence replication (Guatelli et al., *Proc. Nat. Acad. Sci. USA*, 87, 1874 (1990) and W090/06995), selective amplification of target polynucleotide sequences (U.S. Patent No. 6,410,276), consensus sequence primed polymerase chain reaction (CP-PCR) (U.S. Patent No. 4,437,975), arbitrarily primed polymerase chain reaction (AP-PCR) (U.S. Patent Nos. 5,413,909, 5,861,245), rolling circle amplification (RCA) (for example, Fire and Xu, *PNAS* 92:4641 (1995) and Liu et al., *J. Am. Chem. Soc.* 118:1587 (1996)) and U.S. Pat. No. 5,648,245, strand displacement amplification (see Lasken and Egholm, *Trends Biotechnol.* 2003 21(12):531-5; Barker et al. *Genome Res.* 2004 May;14(5):901-7; Dean et al. *Proc Natl Acad Sci U S A.* 2002; 99(8):5261-6; Walker et al. 1992, *Nucleic Acids Res.* 20(7):1691-6, 1992 and Paez, et al. *Nucleic Acids Res.* 2004; 32(9):e71), Qbeta Replicase, described in PCT Patent Application No. PCT/US87/00880 and nucleic acid based sequence amplification (NABSA). (See, U.S. Patent Nos. 5,409,818, 5,554,517, and 6,063,603, each of which is incorporated herein by reference), Other amplification methods that may be used are described in, U.S. Patent Nos. 6,582,938, 5,242,794, 5,494,810, 4,988,617, and US Pub. No. 20030143599 each of which is incorporated herein by reference. DNA may also be amplified by multiplex locus-specific PCR or using adaptor-ligation and single primer PCR (See Kinzler and Vogelstein, *NAR* (1989) 17:3645-53. Other available methods of amplification, such as balanced PCR (Makrigiorgos, et al. (2002), *Nat Biotechnol*, Vol. 20, pp.936-9), may

also be used.

**[00118]** Molecular inversion probes ("MIPs") may also be used for amplification of selected targets. MIPs may be generated so that the ends of the pre-circle probe are complementary to regions that flank the region to be amplified. The gap can be closed by extension of the end of the probe so that the complement of the target is incorporated into the MIP prior to ligation of the ends to form a closed circle. The closed circle can be amplified and detected by sequencing or hybridization as previously disclosed in Hardenbol *et al.*, *Genome Res.* 15:269-275 (2005) and in U.S. Patent No. 6,858,412.

**[00119]** Amplification of the labeled molecule may comprise the use of one or more primers. FIG. 9 shows an exemplary forward and reverse primers. The forward primer (901) may comprise a universal PCR sequence (902), unique identifier sequence (903) and target sequence (904). The reverse primer (905) may comprise a target sequence.

**[00120]** Primers used in the method can be designed with the use of the Primer 3, a computer program which suggests primer sequences based on a user defined input sequence. Other primer designs may also be used, or primers may be selected by eye without the aid of computer programs. There are many options available with the program to tailor the primer design to most applications. Primer3 can consider many factors, including, but not limited to, oligo melting temperature, length, GC content, 3' stability, estimated secondary structure, the likelihood of annealing to or amplifying undesirable sequences (for example interspersed repeats) and the likelihood of primer-dimer formation between two copies of the same primer. In the design of primer pairs, Primer3 can consider product size and melting temperature, the likelihood of primer-dimer formation between the two primers in the pair, the difference between primer melting temperatures, and primer location relative to particular regions of interest to be avoided.

#### **[00121] IV. Sequencing**

**[00122]** In some aspects, the methods disclosed herein further comprise determining the sequence of the labeled-molecule or any product thereof (e.g., labeled-amplicons, labeled-cDNA molecules). Determining the sequence of the labeled-molecule or any product thereof can comprise conducting a sequencing reaction to determine the sequence of at least a portion of the oligonucleotide tag, at least a portion of the labeled-cDNA molecule, a complement thereof, a reverse complement thereof, or any combination thereof. In some instances only the tag or a portion of the tag is sequenced. Determining the sequence of the labeled-molecule or any product thereof can be performed by sequencing methods such as Helioscope<sup>TM</sup> single molecule sequencing, Nanopore DNA sequencing, Lynx Therapeutics'

Massively Parallel Signature Sequencing (MPSS), 454 pyrosequencing, Single Molecule real time (RNAP) sequencing, Illumina (Solexa) sequencing, SOLiD sequencing, Ion Torrent™, Ion semiconductor sequencing, Single Molecule SMRT(TM) sequencing, Polony sequencing, DNA nanoball sequencing, and VisiGen Biotechnologies approach. Alternatively, determining the sequence of the labeled-molecule or any product thereof can use sequencing platforms, including, but not limited to, Genome Analyzer IIx, HiSeq, and MiSeq offered by Illumina, Single Molecule Real Time (SMRT™) technology, such as the PacBio RS system offered by Pacific Biosciences (California) and the Solexa Sequencer, True Single Molecule Sequencing (tSMS™) technology such as the HeliScope™ Sequencer offered by Helicos Inc. (Cambridge, MA).

**[00123]** In some instances, determining the sequence of the labeled-molecule or any product thereof comprises paired-end sequencing, nanopore sequencing, high-throughput sequencing, shotgun sequencing, dye-terminator sequencing, multiple-primer DNA sequencing, primer walking, Sanger dideoxy sequencing, Maxim-Gilbert sequencing, pyrosequencing, true single molecule sequencing, or any combination thereof. Alternatively, the sequence of the labeled-molecule or any product thereof can be determined by electron microscopy or a chemical-sensitive field effect transistor (chemFET) array.

**[00124]** In another example, determining the sequence of labeled-molecules or any product thereof comprises RNA-Seq or microRNA sequencing. Alternatively, determining the sequence of labeled-molecules or any products thereof comprises protein sequencing techniques such as Edman degradation, peptide mass fingerprinting, mass spectrometry, or protease digestion.

**[00125]** The sequencing reaction can, in certain embodiments, occur on a solid or semi-solid support, in a gel, in an emulsion, on a surface, on a bead, in a drop, in a continuous follow, in a dilution, or in one or more physically separate volumes.

**[00126]** Sequencing may comprise sequencing at least about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more nucleotides or base pairs of the labeled molecule. In some instances, sequencing comprises sequencing at least about 200, 300, 400, 500, 600, 700, 800, 900, 1000 or more nucleotides or base pairs of the labeled molecule. In other instances, sequencing comprises sequencing at least about 1500; 2,000; 3,000; 4,000; 5,000; 6,000; 7,000; 8,000; 9,000; or 10,000 or more nucleotides or base pairs of the labeled molecule.

**[00127]** Sequencing may comprise at least about 200, 300, 400, 500, 600, 700, 800, 900, 1000 or more sequencing reads per run. In some instances, sequencing comprises sequencing at least about 1500; 2,000; 3,000; 4,000; 5,000; 6,000; 7,000; 8,000; 9,000; or 10,000 or more

sequencing reads per run.

**[00128] V. Detection Methods**

**[00129]** The methods disclosed herein can further comprise detection of the labeled-molecules and/or labeled-amplicons. Detection of the labeled-molecules and/or labeled-amplicons can comprise hybridization of the labeled-molecules to surface, e.g. a solid support. Alternatively, or additionally, detection of the labeled-molecules comprises contacting the labeled-molecules and/or labeled-amplicons with surface, e.g. a solid support. In some instances, the method further comprises contacting the labeled-molecules and/or labeled-amplicons with a detectable label to produce a detectable-label conjugated labeled-molecule. The methods disclosed herein can further comprise detecting the detectable-label conjugated labeled-molecule. Detection of the labeled-molecules or any products thereof (e.g., labeled-amplicons, detectable-label conjugated labeled-molecule) can comprise detection of at least a portion of the oligonucleotide tag, molecule, detectable label, a complement of the oligonucleotide tag, a complement of the molecule, or any combination thereof.

**[00130]** Detection of the labeled-molecules or any products thereof can comprise an emulsion. For example, the labeled-molecules or any products thereof can be in an emulsion. Alternatively, detection of the labeled-molecules or any products thereof comprises one or more solutions. In other instances, detection of the labeled-molecules comprises one or more containers.

**[00131]** Detection of the labeled-molecules or any products thereof (e.g., labeled-amplicons, detectable-label conjugated labeled-molecule) can comprise detecting each labeled-molecule or products thereof. For example, the methods disclosed herein comprise sequencing at least a portion of each labeled-molecule, thereby detecting each labeled-molecule.

**[00132]** In some instances, detection of the labeled-molecules and/or labeled-amplicons comprises electrophoresis, spectroscopy, microscopy, chemiluminescence, luminescence, fluorescence, immunofluorescence, colorimetry, or electrochemiluminescence methods. For example, the method comprises detection of a fluorescent dye. Detection of the labeled-molecule or any products thereof can comprise colorimetric methods. For example, the colorimetric method comprises the use of a colorimeter or a colorimetric reader. A non-limiting list of colorimeters and colorimetric readers include Sensovation's Colorimetric Array Imaging Reader (CLAIR), ESEQuant Lateral Flow Immunoassay Reader, SpectraMax 340PC 38, SpectraMax Plus 384, SpectraMax 190, VersaMax, VMax, and EMax.

**[00133]** Additional methods used alone or in combination with other methods to detect the labeled-molecules and/or amplicons can comprise the use of an array detector, fluorescence reader, non-fluorescent detector, CR reader, luminometer, or scanner. In some instances, detecting the labeled-molecules and/or labeled-amplicons comprises the use of an array detector. Examples of array detectors include, but are not limited to, diode-array detectors, photodiode array detectors, HPLC photodiode array detectors, pixel array detectors, Germanium array detectors, CMOS and CCD array detectors, Gated linear CCD array detectors, InGaAs photodiode array systems, and TE cooled CCD systems. The array detector can be a microarray detector. Non-limiting examples of microarray detectors include microelectrode array detectors, optical DNA microarray detection platforms, DNA microarray detectors, RNA microarray detectors, and protein microarray detectors.

**[00134]** In some instances, a fluorescence reader is used to detect the labeled-molecule and/or labeled-amplicons. The fluorescence reader can read 1, 2, 3, 4, 5, or more color fluorescence microarrays or other structures on biochips, on slides, or in microplates. In some instances, the fluorescence reader is a Sensovation Fluorescence Array imaging Reader (FLAIR). Alternatively, the fluorescence reader is a fluorescence microplate reader such as the Gemini XPS Fluorescence microplate reader, Gemini EM Fluorescence microplate reader, Finstruments® Fluoroskan filter based fluorescence microplate reader, PHERAstar microplate reader, FLUOstar microplate reader, POLARstar Omega microplate reader, FLUOstar OPTIMA multi-mode microplate reader and POLARstar OPTIMA multi-mode microplate reader. Additional examples of fluorescence readers include PharosFX<sup>TM</sup> and PharosFX Plus systems.

**[00135]** In some instances, detection of the labeled-molecule and/or labeled-amplicon comprises the use of a microplate reader. In some instances, the microplate reader is an xMark<sup>TM</sup> microplate absorbance spectrophotometer, iMark microplate absorbance reader, EnSpire® Multimode plate reader, EnVision Multilabel plate reader, VICTOR X Multilabel plate reader, FlexStation, SpectraMax Paradigm, SpectraMax M5e, SpectraMax M5, SpectraMax M4, SpectraMax M3, SpectraMax M2-M2e, FilterMax F series, Fluoroskan Ascent FL Microplate Fluorometer and Luminometer, Fluoroskan Ascent Microplate Fluorometer, Luminoskan Ascent Microplate Luminometer, Multiskan EX Microplate Photometer, Muliskan FC Microplate Photometer, and Muliskan GO Microplate Photometer. In some instances, the microplate reader detects absorbance, fluorescence, luminescence, time-resolved fluorescence, light scattering, or any combination thereof. In some embodiments, the microplate reader detects dynamic light scattering. The microplate reader,

can in some instances, detect static light scattering. In some instances, detection of the labeled-molecules and/or labeled-amplicons comprises the use of a microplate imager. In some instances, the microplate imager comprises ViewLux uHTS microplate imager and BioRad microplate imaging system.

**[00136]** Detection of labeled-molecules and/or products thereof can comprise the use of a luminometer. Examples of luminometers include, but are not limited to, SpectraMax L, GloMax®-96 microplate luminometer, GloMax®-20/20 single-tube luminometer, GloMax®-Multi<sup>+</sup> with Instinct<sup>TM</sup> software, GloMax®-Multi Jr single tube multimode reader, LUMIstar OPTIMA, LEADER HC<sup>+</sup> luminometer, LEADER 450i luminometer, and LEADER 50i luminometer.

**[00137]** In some instances, detection of the labeled-molecules and/or labeled-amplicons comprises the use of a scanner. Scanners include flatbed scanners such as those provided by Cannon, Epson, HP, Fujitsu, and Xerox. Additional examples of flatbed scanners include the FMBIO® fluorescence imaging scanners (e.g., FMBIO® II, III, and III *Plus* systems). Scanners can include microplate scanners such as the Arrayit ArrayPix<sup>TM</sup> microarray microplate scanner. In some instances, the scanner is a Personal Molecular Imager<sup>TM</sup> (PMI) system provided by Bio-rad.

**[00138]** Detection of the labeled-molecule can comprise the use of an analytical technique that measures the mass-to-charge ratio of charged particles, e.g. mass spectrometry. In some embodiments the mass-to-charge ratio of charged particles is measured in combination with chromatographic separation techniques. In some embodiments sequencing reactions are used in combination with mass-to-charge ratio of charged particle measurements. In some embodiments the tags comprise isotopes. In some embodiments the isotope type or ratio is controlled or manipulated in the tag library.

**[00139]** Detection of the labeled-molecule or any products thereof comprises the use of small particles and/or light scattering. For example, the amplified molecules (e.g., labeled-amplicons) are attached to haptens or directly to small particles and hybridized to the array. The small particles can be in the nanometer to micrometer range in size. The particles can be detected when light is scattered off of its surface.

**[00140]** A colorimetric assay can be used where the small particles are colored, or haptens can be stained with colorimetric detection systems. In some instances, a flatbed scanner can be used to detect the light scattered from particles, or the development of colored materials. The methods disclosed herein can further comprise the use of a light absorbing material. The light absorbing material can be used to block undesirable light scatter or reflection. The light

absorbing material can be a food coloring or other material. In some instances, detection of the labeled-molecule or any products thereof comprises contacting the labeled-molecule with an off-axis white light.

**[00141]** Detection of the labeled-molecule may comprise hybridization chain reaction (HCR). As depicted in FIG. 33, a sample comprising a plurality of nucleic acid molecules (3340) is stochastically labeled with a plurality of oligonucleotide tags (3330). The oligonucleotide tags (3330) comprise a unique identifier region (3310) and an adapter region (3320). Stochastically labeling the nucleic acid molecules can comprise attachment of one or more oligonucleotide tags (3330) to one or more ends of the nucleic acid molecule (3340) to produce one or more labeled-molecules (3345). The one or more labeled molecules can be contacted with a plurality of HCR probes (3350). The plurality of HCR probes (3350) may comprise a hairpin molecules with an overhang and one or more labels (3360, 3390). The plurality of HCR probes (3350) may comprise a mixture of hairpin molecules with 5' overhangs and hairpin molecules with 3' overhangs. The plurality of HCR probes may comprise a stem (3370, 3380). The sequence of the stem (3370, 3380) may be complementary to at least a portion of the oligonucleotide tag. The sequence of the stem (3370, 3380) may be complementary to the adapter region (3320) of the oligonucleotide tag. The adapter region (3320) of the oligonucleotide may act as an initiator for a hybridization chain reaction. As shown in FIG. 33, the stem (3370) of the HCR probe (3350) can hybridize to the adapter region (3320) of the labeled molecule (3345). Hybridization of the stem (3370) of the HCR probe (3350) to the adapter region (3320) of the labeled molecule (3345) can result in opening of the stem (e.g., 3370 and 3380 of the stem are no longer annealed) and linearization of the HCR probe (3350), which results in the formation of a labeled molecule hybridized to a HCR probe (3355). The linearized HCR probe can then act as an initiator for subsequent hybridization of another HCR probe. The stem of a second HCR probe can hybridize to the linearized HCR probe that has hybridized to the labeled molecule, resulting in linearization of the second HCR probe and the formation of a labeled-molecule containing two linearized HCR probes. The linearized second HCR probe can act as an initiator for another hybridization reaction. This process can be repeated multiple times to produce a labeled molecule with multiple linearized HCR probes (3375). The labels (3360, 3390) on the HCR probe can enable detection of the labeled molecule. The labels (3360, 3390) may be any type of label (e.g., fluorophore, chromophore, small molecule, nanoparticle, hapten, enzyme, antibody, magnet). The labels (3360 and 3390) may comprise fragments of a single label. The labels (3360, 3390) may generate a detectable signal when they are in close proximity. When

the HCR probe is a hairpin, the labels (3360 and 3390) may be too far away to produce a detectable signal. When the HCR probe is linearized and multiple linearized HCR probes are hybridized together, the labels (3360, 3390) may be in close enough proximity to generate a detectable signal. For example, a HCR probe (3350) may comprise two pyrene moieties as labels (3360, 3390). Alternatively, the labels may be nanoparticles. The HCR can enable attachment of multiple HCR probes to a labeled molecule, which can result in signal amplification. Stoachastic labeling followed by HCR may increase the sensitivity of detection, analysis and/or quantification of the nucleic acid molecules. Stochastic labeling followed by HCR may increase the accuracy of detection, analysis, and/or quantification of one or more nucleic acid molecules.

**[00142]** Additional methods and apparatus for signal detection and processing of intensity data are disclosed in, for example, U.S. Patents Nos. 5,143,854, 5,547,839, 5,578,832, 5,631,734, 5,800,992, 5,834,758, 5,856,092, 5,902,723, 5,936,324, 5,981,956, 6,025,601, 6,090,555, 6,141,096, 6,185,030, 6,201,639; 6,218,803; and 6,225,625, in U.S. Patent Pub. Nos. 20040012676 and 20050059062 and in PCT Application PCT/US99/06097 (published as WO99/47964), each of which also is hereby incorporated by reference in its entirety for all purposes.

**[00143]** Detection and/or quantification of the labeled molecules may comprise the use of computers or computer software. Computer software products may comprise a computer readable medium having computer-executable instructions for performing the logic steps of the method of the invention. Suitable computer readable medium include floppy disk, CD-ROM/DVD/DVD-ROM, hard-disk drive, flash memory, ROM/RAM, magnetic tapes, etc. The computer-executable instructions may be written in a suitable computer language or combination of several languages. Basic computational biology methods are described in, for example, Setubal and Meidanis et al., *Introduction to Computational Biology Methods* (PWS Publishing Company, Boston, 1997); Salzberg, Searles, Kasif, (Ed.), *Computational Methods in Molecular Biology*, (Elsevier, Amsterdam, 1998); Rashidi and Buehler, *Bioinformatics Basics: Application in Biological Science and Medicine* (CRC Press, London, 2000) and Ouellette and Bzevanis *Bioinformatics: A Practical Guide for Analysis of Gene and Proteins* (Wiley & Sons, Inc., 2<sup>nd</sup> ed., 2001). See also US 6,420,108.

**[00144]** Computer program products and software may be used for a variety of purposes, such as probe design, management of data, analysis, and instrument operation. See, U.S. Patent Nos. 5,593,839, 5,795,716, 5,733,729, 5,974,164, 6,066,454, 6,090,555, 6,185,561, 6,188,783, 6,223,127, 6,229,911 and 6,308,170. Computer methods related to genotyping

using high density microarray analysis may also be used in the present methods, see, for example, US Patent Pub. Nos. 20050250151, 20050244883, 20050108197, 20050079536 and 20050042654. Additionally, the present disclosure may have preferred embodiments that include methods for providing genetic information over networks such as the Internet as shown in U.S. Patent Pub. Nos. 20030097222, 20020183936, 20030100995, 20030120432, 20040002818, 20040126840, and 20040049354.

**[00145]** Detection and/or quantification of the labeled-molecules or any products thereof can comprise the use of one or more algorithms. Alternatively, or additionally, the methods, kits and compositions can further comprise a computer, software, printer, and/or electronic data or information.

**[00146]** The methods disclosed herein can further comprise the transmission of data/information. For example, data/information derived from the detection and/or quantification of the labeled-molecule or any products thereof are transmitted to another device and/or instrument. In some instances, the information obtained from an algorithm can also be transmitted to another device and/or instrument. Transmission of the data/information can comprise the transfer of data/information from a first source to a second source. The first and second sources can be in the same approximate location (e.g., within the same room, building, block, campus). Alternatively, first and second sources are in multiple locations (e.g., multiple cities, states, countries, continents, etc). In some embodiments a non-transitory computable readable media is used to store or analyze data generated using methods described herein.

**[00147]** Transmission of the data/information can comprise digital transmission or analog transmission. Digital transmission can comprise the physical transfer of data (a digital bit stream) over a point-to-point or point-to-multipoint communication channel. Examples of such channels are copper wires, optical fibres, wireless communication channels, and storage media. The data can be represented as an electromagnetic signal, such as an electrical voltage, radiowave, microwave, or infrared signal.

**[00148]** Analog transmission can comprise the transfer of a continuously varying analog signal. The messages can either be represented by a sequence of pulses by means of a line code (baseband transmission), or by a limited set of continuously varying wave forms (passband transmission), using a digital modulation method. The passband modulation and corresponding demodulation (also known as detection) can be carried out by modem equipment. According to the most common definition of digital signal, both baseband and passband signals representing bit-streams are considered as digital transmission, while an

alternative definition only considers the baseband signal as digital, and passband transmission of digital data as a form of digital-to-analog conversion.

**[00149]** The applications and uses of the systems and methods described herein can produce one or more result useful to diagnose a disease state of an individual, for example, a patient. In one embodiment, a method of diagnosing a disease comprises reviewing or analyzing data relating to the presence and/or the concentration level of a target in a sample. A conclusion based review or analysis of the data can be provided to a patient, a health care provider or a health care manager. In one embodiment the conclusion is based on the review or analysis of data regarding a disease diagnosis. It is envisioned that in another embodiment that providing a conclusion to a patient, a health care provider or a health care manager includes transmission of the data over a network.

**[00150]** Accordingly, business systems and methods using the systems and methods described herein are provided.

**[00151]** One aspect of the invention is a business method comprising screening patient test samples for the presence or absence of a biologically active analyte to produce data regarding the analyte, collecting the analyte data, providing the analyte data to a patient, a health care provider or a health care manager for making a conclusion based on review or analysis of the data regarding a disease diagnosis. In one embodiment the conclusion is provided to a patient, a health care provider or a health care manager includes transmission of the data over a network.

**[00152]** Accordingly FIG. 8 is a block diagram showing a representative example logic device through which reviewing or analyzing data relating to the present invention can be achieved. Such data can be in relation to a disease, disorder or condition in an individual. FIG. 8 shows a computer system (or digital device) 800 connected to an apparatus 820 for use with the scanning sensing system 824 to, for example, produce a result. The computer system 800 may be understood as a logical apparatus that can read instructions from media 811 and/or network port 805, which can optionally be connected to server 809 having fixed media 812. The system shown in FIG. 8 includes CPU 801, disk drives 803, optional input devices such as keyboard 815 and/or mouse 816 and optional monitor 807. Data communication can be achieved through the indicated communication medium to a server 809 at a local or a remote location. The communication medium can include any means of transmitting and/or receiving data. The communication medium can comprise a non-transitory computer readable media. For example, the communication medium can be a network connection, a wireless connection or an internet connection. Such a connection can

provide for communication over the World Wide Web. It is envisioned that data can be transmitted over such networks or connections for reception and/or review by a party 822. The receiving party 822 can be but is not limited to a patient, a health care provider or a health care manager.

**[00153]** In one embodiment, a computer-readable medium includes a medium suitable for transmission of a result of an analysis of an environmental or biological sample. The medium can include a result regarding a disease condition or state of a subject, wherein such a result is derived using the methods described herein. The computer readable media can be non-transitory.

**[00154]** Data Analysis: In some embodiments the scanner instrument produces the raw intensity values for each position on the array as well as the background intensity. Many methods can be used to calculate the number of molecules in the sample. For example, the values for the control positions on the array are removed from the dataset and a scatter plot is generated to provide an image of the data. This may occur with or without the background intensity subtracted from the raw data. A threshold intensity value can be established in order to classify the positive spots and the negative spots. All of the positive spots are summed to provide a total count of unique stochastic labels. This process can be automated in Microsoft excel or another computer software program.

**[00155]** An alternative to this strategy is the use of clustering algorithms such as k-means clustering. K-means clustering is a method of cluster analysis which aims to partition all of the observations into clusters in which each observation belongs to the cluster with the nearest mean. The data can be split into 2 or 3 clusters (or more, 3 clusters seems to produce the cleanest numbers so far) and the number of data points can be added up to determine the counts.

## **[00156] VI. Target molecules**

**[00157]** The methods, kits, and systems disclosed herein can be used in the stochastic labeling of molecules. Such molecules include, but are not limited to, polynucleotides and polypeptides. As used herein, the terms “polynucleotide” and “nucleic acid molecule” refers to a polymeric form of nucleotides of any length, either ribonucleotides, deoxyribonucleotides, locked nucleic acids (LNA) or peptide nucleic acids (PNAs), that comprise purine and pyrimidine bases, or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. A “polynucleotide” or “nucleic acid molecule” can consist of a single nucleotide or base pair. Alternatively, the “polynucleotide” or “nucleic acid molecule” comprises two or more nucleotides or base pairs. For example, the

“polynucleotide” or “nucleic acid molecule” comprises at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or base pairs. In another example, the polynucleotide comprises at least about 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, or 10000 nucleotides or base pairs. The backbone of the polynucleotide can comprise sugars and phosphate groups, as may typically be found in RNA or DNA, or modified or substituted sugar or phosphate groups. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components. Thus the terms nucleoside, nucleotide, deoxynucleoside and deoxynucleotide generally include analogs such as those described herein. These analogs are those molecules having some structural features in common with a naturally occurring nucleoside or nucleotide such that when incorporated into a nucleic acid or oligonucleoside sequence, they allow hybridization with a naturally occurring nucleic acid sequence in solution. Typically, these analogs are derived from naturally occurring nucleosides and nucleotides by replacing and/or modifying the base, the ribose or the phosphodiester moiety. The changes can be tailor made to stabilize or destabilize hybrid formation or enhance the specificity of hybridization with a complementary nucleic acid sequence as desired. In some instances, the molecules are DNA, RNA, or DNA-RNA hybrids. The molecules can be single-stranded or double-stranded. In some instances, the molecules are RNA molecules, such as mRNA, rRNA, tRNA, ncRNA, lncRNA, siRNA, or miRNA. The RNA molecules can be polyadenylated. Alternatively, the mRNA molecules are not polyadenylated. Alternatively, the molecules are DNA molecules. The DNA molecules can be genomic DNA. The DNA molecules can comprise exons, introns, untranslated regions, or any combination thereof.

**[00158]** In some instances, the molecules are polypeptides. As used herein, the term “polypeptide” refers to a molecule comprising at least one peptide. In some instances, the polypeptide consists of a single peptide. Alternatively, the polypeptide comprises two or more peptides. For example, the polypeptide comprises at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 peptides. Examples of polypeptides include, but are not limited to, amino acids, proteins, peptides, hormones, oligosaccharides, lipids, glycolipids, phospholipids, antibodies, enzymes, kinases, receptors, transcription factors, and ligands.

**[00159]** The methods, kits, and systems disclosed herein can be used to stochastically label individual occurrences of identical or nearly identical molecules and/or different molecules.

In some instances, the methods, kits, and systems disclosed herein can be used to stochastically label identical or nearly identical molecules (e.g., molecules comprise identical or nearly identical sequences). For example, the molecules to be labeled comprise at least about 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity. The nearly identical molecules may differ by less than about 100, 90, 80, 70, 60, 50, 40, 30, 25, 20, 25, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotides or base pairs. In some instances, the molecules to be labeled are variants of each other. For example, the molecules to be labeled may contain single nucleotide polymorphisms or other types of mutations. In another example, the molecules to be labeled are splice variants. In some instances, at least one molecule is stochastically labeled. In other instances, at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 identical or nearly identical molecules are stochastically labeled. Alternatively, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 identical or nearly identical molecules are stochastically labeled. In other instances, at least 1500; 2,000; 2500; 3,000; 3500; 4,000; 4500; 5,000; 6,000; 7,000; 8,000; 9,000; or 10000 identical or nearly identical molecules are stochastically labeled. In other instances; at least 15,000; 20,000; 25,000; 30,000; 35,000; 40,000; 45,000; 50,000; 60,000; 70,000; 80,000; 90,000; or 100,000 identical or nearly identical molecules are stochastically labeled.

**[00160]** In other instances, the methods, kits, and systems disclosed herein can be used to stochastically label different molecules. For example, the molecules to be labeled comprise less than 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% sequence identity. The different molecules may differ by at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100 or more nucleotides or base pairs. In some instances, at least one molecule is stochastically labeled. In other instances, at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 different molecules are stochastically labeled. Alternatively, at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 different molecules are stochastically labeled. In other instances, at least 1500; 2,000; 2500; 3,000; 3500; 4,000; 4500; 5,000; 6,000; 7,000; 8,000; 9,000; or 10000 different molecules are stochastically labeled. In other instances; at least 15,000; 20,000; 25,000; 30,000; 35,000; 40,000; 45,000; 50,000; 60,000; 70,000; 80,000; 90,000; or 100,000 different molecules are stochastically labeled.

**[00161]** The different molecules to be labeled can be present in the sample at different concentrations or amounts. For example, the concentration or amount of one molecule is greater than the concentration or amount of another molecule in the sample. In some instances, the concentration or amount of at least one molecule in the sample is at least about

1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times greater than the concentration or amount of at least one other molecule in the sample. In another example, the concentration or amount of one molecule is less than the concentration or amount of another molecule in the sample. The concentration or amount of at least one molecule in the sample can be at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times less than the concentration or amount of at least one other molecule in the sample.

**[00162]** In some instances, the molecules to be labeled are in one or more samples. The molecules to be labeled can be in two or more samples. The two or more samples can contain different amounts or concentrations of the molecules to be labeled. In some instances, the concentration or amount of one molecule in one sample can be greater than the concentration or amount of the same molecule in a different sample. For example, a blood sample might contain a higher amount of a particular molecule than a urine sample. Alternatively, a single sample is divided into two or more subsamples. The subsamples can contain different amounts or concentrations of the same molecule. The concentration or amount of at least one molecule in one sample can be at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times greater than the concentration or amount of the same molecule in another sample. Alternatively, the concentration or amount of one molecule in one sample can be less than the concentration or amount of the same molecule in a different sample. For example, a heart tissue sample might contain a higher amount of a particular molecule than a lung tissue sample. The concentration or amount of at least one molecule in one sample can be at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 or more times less than the concentration or amount of the same molecule in another sample. In some instances, the different concentrations or amounts of a molecule in two or more different samples is referred to as sample bias.

## **[00163] VII. Oligonucleotide tags**

**[00164]** In some embodiments, the methods, kits, and systems disclosed herein comprise a plurality of oligonucleotide tags. The oligonucleotide tags can comprise a target specific region, a unique identifier region, an adapter region, a universal primer binding site region, or any combination thereof. FIG. 10-13 shows exemplary oligonucleotide tags.

**[00165]** As shown in FIG. 10, the oligonucleotide tag (1004) may comprise a universal primer binding site (1001), unique identifier region (1002) and a target specific region (1003).

**[00166]** As shown in FIG. 11A, the oligonucleotide tag (1107) can comprise a universal primer binding site (1102), a unique identifier region (1103) and a target specific region (1105). The universal primer binding site (1102) may comprise a phosphorothioate linkage, as depicted by an “\*” in FIG. 11A. As shown in FIG. 11B, the oligonucleotide tag (1128) can comprise a universal primer binding site (1122), a unique identifier region (1123), bridge splint (1129), and a target specific region (1126). As shown in FIG. 11C, the oligonucleotide tag (1158) may comprise a universal primer binding site (1151), unique identifier region (1152), ligation sequence (1153), and a target specific sequence (1157). As shown in FIG. 11D, the oligonucleotide tag (1177) may comprise a universal primer binding site (1171), unique identifier region (1172), ligation sequence (1173), and a DNA target specific sequence (1178).

**[00167]** As shown in FIG. 12A, an oligonucleotide tag (1201) may comprise a universal primer binding site (1202), a unique identifier region comprising a degenerate sequence (1203) and a target specific region (1204). As shown in FIG. 12B, an oligonucleotide tag (1210) may comprise a universal primer binding site (1211), a unique identifier region (1215) comprising a degenerate sequence (1213) flanked by two flanking sequences (1212 and 1214) and a target specific region (1216).

**[00168]** The oligonucleotide tag may comprise one or more secondary structures. As shown in FIG. 13A, the oligonucleotide tag (1301) comprises a hairpin structure. The oligonucleotide tag (1301) can comprise a target specific region (1302), a cleavable stem (1303, 1304), and a unique identifier region (1305).

**[00169]** The oligonucleotide tag may comprise a target specific region that can hybridize to a plurality of different target molecules. For example, as shown in FIG. 13B, the oligonucleotide tag (1310) comprises a universal primer binding site (1311), unique identifier region (1312), and a universal target specific region (1313). The universal target specific region (1313) may comprise an oligodT sequence that enables hybridization to target molecules comprising a polyA or polyU sequence.

**[00170]** A method for synthesizing a plurality of oligonucleotide tags is depicted in FIG. 10. As shown in FIG. 10, oligonucleotide tags (1004) can be synthesized separately. The oligonucleotide tags (1004) can comprise a universal primer binding site (1001), a unique identifier region (1002), and a target specific region. The individual oligonucleotide tags can be pooled to produce a plurality of oligonucleotide tags (1005) comprising a plurality of different unique identifier regions.

**[00171]** A method for synthesizing a plurality of oligonucleotide tags is depicted in FIG.

11A. As shown in FIG. 11A, oligonucleotide fragments (1101) can be synthesized separately. The oligonucleotide fragments (1101) can comprise a universal primer binding site (1102) and a unique identifier region (1103). The universal primer binding site (1102) may comprise a phosphorothioate linkage, as depicted by an “\*” in FIG. 11A. As shown in Step 1 of FIG. 11A, the individual oligonucleotide fragments (1101) may be mixed to produce a plurality of oligonucleotide fragments (1104). The plurality of oligonucleotide fragments (1104) can be attached to a target specific region (1105). As shown in Step 2 of FIG. 11A, the target specific region can be ligated to the oligonucleotide tag to produce an oligonucleotide tag comprising a target specific region (1105). 5' and 3' exonucleases may be added to the reaction to remove non-ligated products (1105, 1101). The oligonucleotide tag (1106) comprising the universal primer binding site (1102), unique identifier region (1103) and target specific region (1105) may be resistant to 5' and 3' exonucleases. As shown in Step 3 of FIG. 11A, the 3' phosphate group from the ligated oligonucleotide tag (1106) can be removed to produce an oligonucleotide tag (1107) without a 3' phosphate group. The 3' phosphate group can be removed enzymatically. For example, a T4 polynucleotide kinase can be used to remove the 3' phosphate group.

**[00172]** Another method of synthesizing oligonucleotide tags is depicted in FIG. 11B. As shown in FIG. 11B, an oligonucleotide tag (1128) can be synthesized by ligating two oligonucleotide fragments (1121 and 1127). One oligonucleotide fragment (1121) may comprise a universal primer binding site (1122), unique identifier region (1123) and a left splint (1124). The other oligonucleotide fragment (1128) may comprise a right splint (1125) and a target specific region (1126). A ligase (e.g., T4 DNA ligase) can be used to join the two oligonucleotide fragments (1121 and 1127) to produce an oligonucleotide tag (1128). Double stranded ligation of the left splint (1124) and right splint (1125) can produce an oligonucleotide tag (1128) with a bridge splint (1129).

**[00173]** An alternative method of synthesizing an oligonucleotide tag by ligating two oligonucleotide fragments is depicted in FIG. 11C. As shown in FIG. 11C, an oligonucleotide tag (1158) is synthesized by ligating two oligonucleotide fragments (1150 and 1158). One oligonucleotide fragment (1150) may comprise a universal primer binding site (1151), unique identifier region (1152), and a ligation sequence (1153). The other oligonucleotide fragment (1158) may comprise a ligation sequence (1154) that is complementary to the ligation sequence (1153) of the first oligonucleotide fragment (1150), a complement of a target specific region (1155), and a label (1156). The oligonucleotide fragment (1159) may also comprise a 3' phosphate which prevents extension of the oligonucleotide fragment. As shown

in Step 1 of FIG. 11C, the ligation sequences (1153 and 1154) of the two oligonucleotide fragments may anneal and a polymerase can be used to extend the 3' end of the first oligonucleotide fragment (1150) to produce an oligonucleotide tag (1158). The oligonucleotide tag (1158) may comprise a universal primer binding site (1151), unique identifier region (1152), ligation sequence (1153), and a target specific sequence (1157). The target specific sequence (1157) of the oligonucleotide tag (1158) may be the complement of the complement of the target specific region (1155) of the second oligonucleotide fragment (1159). The oligonucleotide fragment comprising the label (1156) can be removed from the oligonucleotide tags (1158). For example, the label (1156) may comprise biotin and oligonucleotide fragments (1159) comprising the biotin label (1156) can be removed via streptavidin capture. In another example, the label (1156) may comprise a 5' phosphate and oligonucleotide fragments (1159) comprising the 5' phosphate (1156) can be removed via an exonuclease (e.g., Lambda exonuclease).

**[00174]** As depicted in FIG. 11D, a first oligonucleotide fragment (1170) comprising a universal primer binding site (1171), unique identifier region (1172), a first ligation sequence (1173) is annealed to a second oligonucleotide fragment (1176) comprising a second ligation sequence (1174) and an RNA complement of the target sequence (1175). Step 1 may comprise annealing the first and second ligation sequences (1173 and 1174) followed by reverse transcription of the RNA complement of the target sequence (1175) to produce an oligonucleotide tag (1177) comprising a universal primer binding site (1171), unique identifier region (1172), a first ligation sequence (1173), and a target specific region (1178). The oligonucleotide fragments comprising the RNA complement of the target sequence can be selectively degraded by RNase treatment.

**[00175]** The oligonucleotide tag can comprise at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or base pairs. In another example, the oligonucleotide tag comprises at least about 1500, 2,000; 2500, 3,000; 3500, 4,000; 4500, 5,000; 5500, 6,000; 6500, 7,000; 7500, 8,000; 8500, 9,000; 9500, or 10,000 nucleotides or base pairs.

**[00176]** The tags can be hexamers, e.g. random hexamers. The tags can be randomly generated from a set of mononucleotides. The tags can be assembled by randomly incorporating mononucleotides.

**[00177]** The tags can also be assembled without randomness, to generate a library of different tags which are not randomly generated but which includes sufficient numbers of different tags to practice the methods.

**[00178]** In some embodiments an oligonucleotide tag can comprise a cutback in a target molecule. The cutback can be, for example, a enzymatic digestion of one or both ends of a target molecule. The cutback can be used in conjunction with the addition of added oligonucleotide tags. The combination of the cutback and the added tags can contain information related to the particular starting molecule. By adding a random cutback to the tag a smaller diversity of the added tags may be necessary for counting the number of target molecules when detection allows a determination of both the random cutback and the added oligonucleotides.

**[00179]** The oligonucleotide tag can comprise a target specific region. The target specific region can comprise a sequence that is complementary to the molecule. In some instances, the molecule is an mRNA molecule and the target specific region comprises an oligodT sequence that is complementary to the polyA tail of the mRNA molecule. The target specific region can also act as a primer for DNA and/or RNA synthesis. For example, the oligodT sequence of the target specific region can act as a primer for first strand synthesis of a cDNA copy of the mRNA molecule. Alternatively, the target specific region comprises a sequence that is complementary to any portion of the molecule. In other instances, the target specific region comprises a random sequence that can be hybridized or ligated to the molecule. The target specific region can enable attachment of the oligonucleotide tag to the molecule. Attachment of the oligonucleotide tag can occur by any of the methods disclosed herein (e.g., hybridization, ligation). In some instances, the target specific region comprises a sequence that is recognized by one or more restriction enzymes. The target specific region can comprise at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or base pairs. In another example, the target specific region comprises at least about 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, or 10000 nucleotides or base pairs. Preferably, the target specific region comprises at least about 5-10, 10-15, 10-20, 10-30, 15-30, or 20-30 nucleotides or base pairs.

**[00180]** In some instances, the target specific region is specific for a particular gene or gene product. For example, the target specific region comprises a sequence complementary to a region of a p53 gene or gene product. Therefore, the oligonucleotide tags can only attach to molecules comprising the p53-specific sequence. Alternatively, the target specific region is specific for a plurality of different genes or gene products. For example, the target specific region comprises an oligodT sequence. Therefore, the oligonucleotide tags can attach to any molecule comprising a polyA sequence. In another example, the target specific region

comprises a random sequence that is complementary to a plurality of different genes or gene products. Thus, the oligonucleotide tag can attach to any molecule with a sequence that is complementary to the target specific region. In other instances, the target specific region comprises a restriction site overhang (e.g., EcoRI sticky-end overhang). The oligonucleotide tag can ligate to any molecule comprising a sequence complementary to the restriction site overhang.

**[00181]** The oligonucleotide tag disclosed herein often comprises a unique identifier region. The unique identifier region may be used to uniquely identify occurrences of target species thereby marking each species with an identifier that can be used to distinguish between two otherwise identical or nearly identical targets. The unique identifier region of the plurality of oligonucleotide tags can comprise a collection of different semiconductor nanocrystals, metal compounds, peptides, oligonucleotides, antibodies, small molecules, isotopes, particles or structures having different shapes, colors, barcodes or diffraction patterns associated therewith or embedded therein, strings of numbers, random fragments of proteins or nucleic acids, different isotopes, or any combination thereof. The unique identifier region can comprise a degenerative sequence. The unique identifier region can comprise at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or base pairs. In another example, the unique identifier region comprises at least about 1500; 2,000; 2500, 3,000; 3500, 4,000; 4500, 5,000; 5500, 6,000; 6500, 7,000; 7500, 8,000; 8500, 9,000; 9500, or 10,000 nucleotides or base pairs. Preferably, the unique identifier region comprises at least about 10-30, 15-40, or 20-50 nucleotides or base pairs.

**[00182]** In some instances, the oligonucleotide tag comprises a universal primer binding site. The universal primer binding site allows the attachment of a universal primer to the labeled-molecule and/or labeled-amplicon. Universal primers are well known in the art and include, but are not limited to, -47F (M13F), alfaMF, AOX3', AOX5', BGH\_r, CMV\_-30, CMV\_-50, CVM\_f, LACrmt, lamgda gt10F, lambda gt 10R, lambda gt11F, lambda gt11R, M13 rev, M13Forward(-20), M13Reverse, male, p10SEQP\_pQE, pA\_-120, pet\_4, pGAP Forward, pGL\_RVpr3, pGLpr2\_R, pKLAC1\_4, pQE\_FS, pQE\_RS, puc\_U1, puc\_U2, revers\_A, seq\_IRES\_tam, seq\_IRES\_zpet, seq\_ori, seq\_PCR, seq\_pIRES-, seq\_pIRES+, seq\_pSecTag, seq\_pSecTag+, seq\_retro+PSI, SP6, T3-prom, T7-prom, and T7-term\_Inv. Attachment of the universal primer to the universal primer binding site can be used for amplification, detection, and/or sequencing of the labeled-molecule and/or labeled-amplicon. The universal primer binding site can comprise at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11,

12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or base pairs. In another example, the universal primer binding site comprises at least about 1500; 2,000; 2500, 3,000; 3500, 4,000; 4500, 5,000; 5500, 6,000; 6500, 7,000; 7500, 8,000; 8500, 9,000; 9500, or 10,000 nucleotides or base pairs. Preferably, the universal primer binding site comprises 10-30 nucleotides or base pairs.

**[00183]** The oligonucleotide tag may comprise an adapter region. The adapter region may enable hybridization of one or more probes. The adapter region may enable hybridization of one or more HCR probes.

**[00184]** The oligonucleotide tag may comprise one or more labels.

**[00185]** The oligonucleotide tag may act as an initiator for a hybridization chain reaction (HCR). The adapter region of the oligonucleotide tag may act as an initiation for HCR. The universal primer binding site may act as an initiator for HCR.

**[00186]** In some instances, the oligonucleotide tag is single-stranded. In other instances, the oligonucleotide tag is double-stranded. The oligonucleotide tag can be linear. Alternatively, the oligonucleotide tag comprises a secondary structure. As used herein, “secondary structure” includes tertiary, quaternary, etc...structures. In some instances, the secondary structure is a hairpin, a stem-loop structure, an internal loop, a bulge loop, a branched structure or a pseudoknot, multiple stem loop structures, cloverleaf type structures or any three dimensional structure. In some instances, the secondary structure is a hairpin. The hairpin can comprise an overhang sequence. The overhang sequence of the hairpin can act as a primer for a polymerase chain reaction and/or reverse transcription reaction. The overhang sequence comprises a sequence that is complementary to the molecule to which the oligonucleotide tag is attached and the overhang sequence hybridizes to the molecule. The overhang sequence can be ligated to the molecule and acts as a template for a polymerase chain reaction and/or reverse transcription reaction. In some embodiments the tag comprises nucleic acids and/or synthetic nucleic acids and/or modified nucleic acids.

**[00187]** An oligonucleotide tag comprising a hairpin may act as a probe for a hybridization chain reaction. Further disclosed herein is a stochastic label-based hybridization chain reaction (HCR) method comprising stochastically labeling one or more nucleic acid molecules with an oligonucleotide tag, wherein the oligonucleotide tag is a hairpin and the one or more nucleic acid molecules act as initiators for a hybridization chain reaction. A schematic of a stochastic label-based hybridization reaction is depicted in FIG. 34. As shown in FIG. 34, one or more nucleic acid molecules (3480) are stochastically labeled with a plurality of hairpin oligonucleotide tags (3490) by initiating a hybridization chain reaction.

The hairpin oligonucleotide tags may comprise one or more labels (3410, 3470), an overhang (3420, 3420'), a stem (3430, 3460), and a loop (3450). The overhang region (3420) of the hairpin oligonucleotide tag (3490) may comprise a target specific region. The overhang region (3420) may comprise an oligodT sequence. The sample comprising the one or more nucleic acid molecules may be treated with one or more restriction nucleases prior to stochastic labeling. The overhang region (3420) may comprise a restriction enzyme recognition sequence. The sample comprising the one or more nucleic acid molecules may be contacted with one or more adapters prior to stochastic labeling to produce an adapter-nucleic acid molecule hybrid. The overhang region (3420) and the stem (3430) may be complementary to the one or more adapters. The loop (3450) of the oligonucleotide tag may comprise a unique identifier region. Hybridization of a first hairpin oligonucleotide tag (3490) to the nucleic acid molecules (3480) may result in the formation of a labeled molecule (3415), wherein the first hairpin oligonucleotide tag is linearized to produce a first linearized oligonucleotide tag. The first linearized oligonucleotide tag of labeled molecule (3415) can act as an initiator for hybridization of a second hairpin oligonucleotide tag to the labeled molecule (3415) to produce a labeled molecule with two linearized oligonucleotide tags (3425). The second linearized oligonucleotide tag can act as an initiator for another hybridization reaction. This process can be repeated multiple times to produce a labeled molecule with multiple linearized HCR probes (3435). The labels (3410, 3470) on the HCR probe can enable detection of the labeled molecule. The labels (3410, 3470) may be any type of label (e.g., fluorophore, chromophore, small molecule, nanoparticle, hapten, enzyme, antibody, magnet). The labels (3360 and 3390) may comprise fragments of a single label. The labels (3410, 3470) may generate a detectable signal when they are in close proximity. When the oligonucleotide tag is a hairpin, the labels (3360 and 3390) may be too far away to produce a detectable signal. When the hairpin oligonucleotide tag is linearized and multiple linearized hairpin oligonucleotide tags are hybridized together, the labels (3410, 3470) may be in close enough proximity to generate a detectable signal. For example, a hairpin oligonucleotide tag (3350) may comprise two pyrene moieties as labels (3410, 3470). Alternatively, the labels may be nanoparticles. The stochastic label-based HCR can enable attachment of multiple hairpin oligonucleotide tags to a labeled molecule, which can result in signal amplification. Stoachastic label-based HCR may increase the sensitivity of detection, analysis and/or quantification of the nucleic acid molecules. Stochastic label-based HCR may increase the accuracy of detection, analysis, and/or quantification of one or more nucleic acid molecules.

**[00188]** In some instances, the plurality of oligonucleotide tags comprises at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 different oligonucleotide tags. In other instances, the plurality of oligonucleotide tags comprises at least about 200; 300; 400; 500; 600; 700; 800; 900; 1,000; 2,000; 3,000; 4,000; 5,000; 6,000; 7,000; 8,000; 9,000; or 10000 different oligonucleotide tags. Alternatively; the plurality of oligonucleotide tags comprises at least about 20,000; 30,000; 40,000; 50,000; 60,000; 70,000; 80,000; 90,000; or 100,000 different oligonucleotide tags.

**[00189]** The number of oligonucleotide tags in the plurality of oligonucleotide tags is often in excess of the number of molecules to be labeled. In some instances, the number of oligonucleotide tags in the plurality of oligonucleotide tags is at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times greater than the number of molecules to be labeled.

**[00190]** The number of different oligonucleotide tags in the plurality of oligonucleotide tags is often in excess of the number of different molecules to be labeled. In some instances, the number of different oligonucleotide tags in the plurality of oligonucleotide tags is at least about 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times greater than the number of different molecules to be labeled.

**[00191]** In some instances, stochastic labeling of a molecule comprises a plurality of oligonucleotide tags, wherein the concentration of the different oligonucleotide tags in the plurality of oligonucleotide tags is the same. In such instances, the plurality of oligonucleotide tags comprises equal numbers of each different oligonucleotide tag. Additionally, the relative ratio of the different oligonucleotide tags in the plurality of oligonucleotide is 1:1:1...1.

**[00192]** In some instances, stochastic labeling of a molecule comprises a plurality of oligonucleotide tags, wherein the concentration of the different oligonucleotide tags in the plurality of oligonucleotide tags is different. In such instances, the plurality of oligonucleotide tags comprises different numbers of each different oligonucleotide tag. Additionally, the relative ratio of the different oligonucleotide tags in the plurality of oligonucleotide is not 1:1:1...1. In some instances, some oligonucleotide tags are present at higher concentrations than other oligonucleotide tags in the plurality of oligonucleotide tags. In some instances, stochastic labeling with different concentrations of oligonucleotide tags extends the sample measurement dynamic range without increasing the number of different labels used. For example, consider stochastically labeling 3 nucleic acid sample molecules with 10 different oligonucleotide tags all at equal concentration. We expect to observe 3

different labels. Now instead of 3 nucleic acid molecules, consider 30 nucleic acid molecules, and we expect to observe all 10 labels. In contrast, if we still used 10 different stochastic labels and alter the relative ratios of the labels to 1:2:3:4...10, then with 3 nucleic acid molecules, we would expect to observe between 1-3 labels, but with 30 molecules we would expect to observe only approximately 5 labels thus extending the range of measurement with the same number of stochastic labels.

**[00193]** The relative ratios of the different oligonucleotide tags in the plurality of oligonucleotide tags can be 1:X, where X is at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100. Alternatively, the relative ratios of “n” different oligonucleotide tags in the plurality of oligonucleotide tags is 1:A:B:C:...Z<sub>n</sub>, where A, B, C...Z<sub>n</sub> is at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100.

**[00194]** In some instances, the concentration of two or more different oligonucleotide tags in the plurality of oligonucleotide tags is the same. For “n” different oligonucleotide tags, the concentration of at least 2, 3, 4, ...n different oligonucleotide tags is the same. Alternatively, the concentration of two or more different oligonucleotide tags in the plurality of oligonucleotide tags is different. For “n” different oligonucleotide tags, the concentration of at least 2, 3, 4,...n different oligonucleotide tags is different. In some instances, for “n” different oligonucleotide tags, the difference in concentration for at least 2, 3, 4,...n different oligonucleotide tags is at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000-fold.

**[00195]** In some instances, at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 100% of the different oligonucleotide tags in the plurality of oligonucleotide tags have the same concentration. Alternatively, at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 100% of the different oligonucleotide tags in the plurality of oligonucleotide tags have a different concentration.

**[00196]** The sequences of the oligonucleotide tags may be optimized to minimize dimerization of oligonucleotide tags. FIG. 15 depicts the formation of oligonucleotide tag dimers when the oligonucleotide tag sequences are not optimized. As shown in FIG. 15, when oligonucleotide tag sequences are not optimized, a first oligonucleotide tag (1507) comprising a universal primer binding site (1501), a first unique identifier region (1502) and

a first target specific region (1503) can anneal to a second oligonucleotide tag (1508) comprising a universal primer binding site (1501), a second unique identifier region (1504) and a second target specific region (1505). The oligonucleotide tag dimer can be amplified and result in the formation of an amplicon (1506) comprising two universal primer binding sites on each end of the amplicon and a target specific region and a unique identifier region. Because the concentration of the oligonucleotide tags are far greater than the number of DNA templates, these oligonucleotide tag dimers can outcompete the labeled DNA molecules in an amplification reaction. Unamplified DNAs lead to false negatives, and amplified oligonucleotide tag dimers lead to high false positives. Thus, the oligonucleotide tags can be optimized to minimize oligonucleotide tag dimer formation. Alternatively, oligonucleotide tags that dimerize are discarded, thereby eliminating oligonucleotide tag dimer formation.

**[00197]** Alternatively, as depicted in FIG. 16, oligonucleotide tag dimer formation can be eliminated or reduced by incorporating one or more modifications into the oligonucleotide tag sequence. As shown in FIG. 16, an oligonucleotide tag (1610) comprising a universal primer binding site (1611), unique identifier region (1612), and target specific region (1613) comprising uracils and a 3' phosphate group is annealed to a target molecule (1616). The target molecule (1616) may be a restriction endonuclease digested fragment. The restriction endonuclease may recognize the recognition site depicted in FIG. 16. PCR amplification may comprise one or more forward primers (1618 and 1618) and one or more reverse primers (1614 and 1615). PCR amplification may comprise nested PCR with a forward primer (1618) specific for the universal primer binding site (1611) of the oligonucleotide tag and a forward primer (1617) specific for the target specific region (1613) of the oligonucleotide tag and reverse primers (1614 and 1615) that are specific for the target molecule. The target molecule can be amplified using a Pfu DNA polymerase, which cannot amplify template comprising one or more uracils. Thus, any dimerized oligonucleotide tags cannot be amplified by Pfu DNA polymerase.

**[00198] VIII. Detectable labels**

**[00199]** The methods, kits, and systems disclosed herein can further comprise a detectable label. The terms “detectable label” or “label” can be used interchangeably and refer to any chemical moiety attached to a nucleotide, nucleotide polymer, or nucleic acid binding factor, wherein the attachment may be covalent or non-covalent. Preferably, the label is detectable and renders the nucleotide or nucleotide polymer detectable to the practitioner of the invention. Detectable labels that may be used in combination with the methods disclosed herein include, for example, a fluorescent label, a chemiluminescent label, a quencher, a

radioactive label, biotin, pyrene moiety, gold, or combinations thereof. Non-limiting example of detectable labels include luminescent molecules, fluorochromes, fluorescent quenching agents, colored molecules, radioisotopes or scintillants.

**[00200]** In some instances, the methods disclosed herein further comprise attaching one or more detectable labels to the labeled-molecule or any product thereof (e.g., labeled-amplicon). The methods can comprise attaching two or more detectable labels to the labeled-molecule. Alternatively, the method comprises attaching at least about 3, 4, 5, 6, 7, 8, 9, or 10 detectable labels to a labeled-molecule. In some instances, the detectable label is a Cy<sup>TM</sup> label. The Cy<sup>TM</sup> label is a Cy3 label. Alternatively, or additionally, the detectable label is biotin. In some embodiments the detectable label is attached to a probe which binds the molecule or labeled molecule. This can occur, for example, after the molecule or labeled molecule has been hybridized to an array. In one example the molecule is bound to partners on an array. After the binding a probe with can bind the molecule is bound to the molecules on the array. This process can be repeated with multiple probes and labels to decrease the likelihood that a signal is the result of nonspecific binding of a label or nonspecific binding of the molecule to the array.

**[00201]** In some instances a donor acceptor pair can be used as the detectable labels. Either the donor or acceptor can be attached to a probe that binds a nucleic acid. The probe can be, for example, a nucleic acid probe that can bind to a the molecule or the labeled molecule. The corresponding donor or acceptor can be added to cause a signal.

**[00202]** In some instances, the detectable label is a Freedom dye, Alexa Fluor<sup>®</sup> dye, Cy<sup>TM</sup> dye, fluorescein dye, or LI-COR IRDyes<sup>®</sup>. In some instances, the Freedom dye is fluorescein (6-FAM<sup>TM</sup>, 6-carboxyfluorescein), MAX (NHS Ester), TYE<sup>TM</sup> 563, TEX 615, TYE<sup>TM</sup> 665, TYE 705. The detectable label can be an Alexa Fluor dye. Examples of Alexa Fluor<sup>®</sup> dyes include Alexa Fluor<sup>®</sup> 488 (NHS Ester), Alexa Fluor<sup>®</sup> 532 (NHS Ester), Alexa Fluor<sup>®</sup> 546 (NHS Ester), Alexa Fluor<sup>®</sup> 594 (NHS Ester), Alexa Fluor<sup>®</sup> 647 (NHS Ester), Alexa Fluor<sup>®</sup> 660 (NHS Ester), or Alexa Fluor<sup>®</sup> 750 (NHS Ester). Alternatively, the detectable label is a Cy<sup>TM</sup> dye. Examples of Cy<sup>TM</sup> dyes include, but are not limited to, Cy2, Cy3, Cy3B, Cy3.5, Cy5, Cy5.5, and Cy7. In some instances, the detectable label is a fluorescein dye. Non-limiting examples of fluorescein dyes include 6-FAM<sup>TM</sup> (Azide), 6-FAM<sup>TM</sup> (NHS Ester), Fluorescein dT, JOE (NHS Ester), TET<sup>TM</sup>, and HEX<sup>TM</sup>. In some instances, the detectable label is a LI-COR IRDyes<sup>®</sup>, such as 5' IRDye<sup>®</sup> 700, 5' IRDye<sup>®</sup> 800, or IRDye<sup>®</sup> 800CW (NHS Ester). In some instances, the detectable label is TYE<sup>TM</sup> 563. Alternatively, the detectable label is Cy3.

**[00203]** The detectable label can be Rhodamine dye. Examples of rhodamine dyes include, but are not limited to, Rhodamine Green<sup>TM</sup>-X (NHS Ester), TAMRA<sup>TM</sup>, TAMRA<sup>TM</sup> (NHS Ester), Rhodamine Red<sup>TM</sup>-X(NHS Ester), ROX<sup>TM</sup> (NHS Ester), and 5'TAMRA<sup>TM</sup> (Azide). In other instances, the detectable label is a WellRED Dye. WellRED Dyes include, but are not limited to, WellRED D4 dye, WellRED D3 dye, and WellRED D2 dye. In some instances, the detectable label is Texas Red<sup>®</sup>-X (NHS Ester), Lightcycler<sup>®</sup> 640 (NHS Ester), or Dy 750 (NHS Ester).

**[00204]** In some instances, detectable labels include a linker molecule. Examples of linker molecules include, but are not limited to, biotin, avidin, streptavidin, HRP, protein A, protein G, antibodies or fragments thereof, Grb2, polyhistidine, Ni<sup>2+</sup>, FLAG tags, myc tags. Alternatively, detectable labels include heavy metals, electron donors/acceptors, acridinium esters, dyes and calorimetric substrates. In other instances, detectable labels include enzymes such as alkaline phosphatase, peroxidase and luciferase.

**[00205]** A change in mass can be considered a detectable label, as is the case of surface plasmon resonance detection. The skilled artisan would readily recognize useful detectable labels that are not mentioned herein, which may be employed in the operation of the present invention.

**[00206]** In some instances, detectable labels are used with primers. For example, the universal primer is a labeled with the detectable label (e.g., Cy3 labeled universal primer, fluorophore labeled universal primer). Alternatively, the target specific primer is labeled with the detectable label (e.g., TYE 563-labeled target specific primer). In other instances, detectable labels are used with the oligonucleotide tags. For example, the oligonucleotide tag is labeled with a detectable label (e.g., biotin-labeled oligonucleotide tag). In other instances, detectable labels are used with the nucleic acid template molecule. Detectable labels can be used to detect the labeled-molecules or labeled-amplicons. Alternatively, detectable labels are used to detect the nucleic acid template molecule.

**[00207]** In some instances, the detectable label is attached to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, probe, HCR probe, and/or non-labeled molecule. Methods for attaching the detectable label to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled molecule include, but are not limited to, chemical labeling and enzymatic labeling. In some instances, the detectable label is attached by chemical labeling. In some embodiments, chemical labeling techniques comprise a chemically reactive group. Non-limiting examples of reactive groups include amine-reactive succinimidyl esters such as NHS-fluorescein or NHS-rhodamine, amine-reactive

isothiocyanate derivatives including FITC, and sulfhydryl-reactive maleimide-activated fluors such as fluorescein-5-maleimide. In some embodiments, reaction of any of these reactive dyes with another molecule results in a stable covalent bond formed between a fluorophore and the linker and/or agent. In some embodiments, the reactive group is isothiocyanates. In some embodiments, a label is attached to an agent through the primary amines of lysine side chains. In some embodiments, chemical labeling comprises a NHS-ester chemistry method.

**[00208]** Alternatively, the detectable label is attached by enzymatic labeling. Enzymatic labeling methods can include, but are not limited to, a biotin acceptor peptide/biotin ligase (AP/Bir A), acyl carrier protein/phosphopantetheine transferase (ACP/PPTase), human *O*<sup>6</sup>-alkylguanine transferase (hAGT), Q-tag/transglutaminase (TGase), aldehyde tag/formylglycine-generating enzyme, mutated prokaryotic dehalogenase (HaloTagTM), and farnesylation motif/protein farnesyltransferase (PFTase) methods. Affinity labeling can include, but is not limited to, noncovalent methods utilizing dihydrofolate reductase (DHFR) and Phe36Val mutant of FK506-binding protein 12 (FKBP12(F36V)) , and metal-chelation methods.

**[00209]** Crosslinking reagents can be used to attach a detectable label to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled molecule. In some instances, the crosslinking reagent is glutaraldehyde. Glutaraldehyde can react with amine groups to create crosslinks by several routes. For example, under reducing conditions, the aldehydes on both ends of glutaraldehyde couple with amines to form secondary amine linkages.

**[00210]** In some instances, attachment of the detectable label to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled molecule comprises periodate-activation followed by reductive amination. In some instances, Sulfo-SMCC or other heterobifunctional crosslinkers are used to conjugate the detectable to the primer, oligonucleotide tag, labeled-molecule, labeled-amplicon, and/or non-labeled molecule. For example, Sulfo-SMCC is used to conjugate an enzyme to a drug. In some embodiments, the enzyme is activated and purified in one step and then conjugated to the drug in a second step. In some embodiments, the directionality of crosslinking is limited to one specific orientation (e.g., amines on the enzyme to sulfhydryl groups on the antibody).

## **[00211] IX. Supports**

**[00212]** In some instances, the methods, kits, and systems disclosed herein comprise a support. The term “support” and “substrate” as used herein are used interchangeably and refer

to a material or group of materials having a rigid or semi-rigid surface or surfaces. The support or substrate can be a solid support. Alternatively, the support is a non-solid support. The support or substrate can comprise a membrane, paper, plastic, coated surface, flat surface, glass, slide, chip, or any combination thereof. In many embodiments, at least one surface of the solid support will be substantially flat, although in some embodiments it may be desirable to physically separate synthesis regions for different compounds with, for example, wells, raised regions, pins, etched trenches, or the like. According to other embodiments, the solid support(s) will take the form of beads, resins, gels, microspheres, or other geometric configurations. Alternatively, the solid support(s) comprises silica chips, microparticles, nanoparticles, plates, and arrays. Methods and techniques applicable to polymer (including protein) array synthesis have been described in U.S. Patent Pub. No. 20050074787, WO 00/58516, U.S. Patent Nos. 5,143,854, 5,242,974, 5,252,743, 5,324,633, 5,384,261, 5,405,783, 5,424,186, 5,451,683, 5,482,867, 5,491,074, 5,527,681, 5,550,215, 5,571,639, 5,578,832, 5,593,839, 5,599,695, 5,624,711, 5,631,734, 5,795,716, 5,831,070, 5,837,832, 5,856,101, 5,858,659, 5,936,324, 5,968,740, 5,974,164, 5,981,185, 5,981,956, 6,025,601, 6,033,860, 6,040,193, 6,090,555, 6,136,269, 6,269,846 and 6,428,752, in PCT Publication No. WO 99/36760 and WO 01/58593, which are all incorporated herein by reference in their entirety for all purposes. Patents that describe synthesis techniques in specific embodiments include U.S. Patent Nos. 5,412,087, 6,147,205, 6,262,216, 6,310,189, 5,889,165, and 5,959,098. Nucleic acid arrays are described in many of the above patents, but many of the same techniques may be applied to polypeptide arrays. Additional exemplary substrates are disclosed in U.S. Patent No. 5,744,305 and US Patent Pub. Nos. 20090149340 and 20080038559.

**[00213]** In some instances, the solid support is a bead. Examples of beads include, but are not limited to, streptavidin beads, agarose beads, magnetic beads, Dynabeads®, MACS® microbeads, antibody conjugated beads (e.g., anti-immunoglobulin microbead), protein A conjugated beads, protein G conjugated beads, protein A/G conjugated beads, protein L conjugated beads, oligo-dT conjugated beads, silica beads, silica-like beads, anti-biotin microbead, anti-fluorochrome microbead, and BcMag™ Carboxy-Terminated Magnetic Beads.

**[00214]** The solid support can be an array or microarray. The solid support can comprise discrete regions. The solid support can be an addressable array. In some instances, the array comprises a plurality of probes fixed onto a solid surface. The plurality of probes enables hybridization of the labeled-molecule and/or labeled-amplicon to the solid surface. The

plurality of probes comprises a sequence that is complementary to at least a portion of the labeled-molecule and/or labeled-amplicon. In some instances, the plurality of probes comprises a sequence that is complementary to the oligonucleotide tag portion of the labeled-molecule and/or labeled-amplicon. In other instances, the plurality of probes comprises a sequence that is complementary to the junction formed by the attachment of the oligonucleotide tag to the molecule.

**[00215]** The array can comprise one or more probes. The probes can be in a variety of formats as depicted in FIG. 18. As shown in FIG. 18A-18C, 18G and 18H, the array (1801, 1806, 1811, 1828, 1832) can comprise a probe (1804, 1809, 1814, 1836, 1835) comprising a sequence that is complementary to at least a portion of the target molecule (1802, 1807, 1813, 1829, 1833) and a sequence that is complementary to the unique identifier region of an oligonucleotide tag (1803, 1808, 1812, 1830, 1834). As shown FIG. 18A-18B, 18G and 18H, the sequence that is complementary to at least a portion of the target molecule (1802, 1807, 1829, 1833) can be attached to the array. As shown in FIG. 18C, the sequence that is complementary to the unique identifier region (1812) can be attached to the array. As shown in FIG. 18D-18F, the array (1816, 1820, 1824) can comprise a first probe (1817, 1821, 1825) comprising a sequence that is complementary to at least a portion of the target molecule and a second probe (1819, 1823, 1827) that is complementary to the unique identifier region. FIG. 18A-18H also depict the various ways in which a stochastically labeled molecule (1805, 1810, 1815, 1818, 1822, 1826, 1831, 1837) can hybridize to the arrays. For example, as shown in FIG. 18A and 18C, the junction of the unique identifier region and the target molecule of the stochastically labeled molecule (1805, 1815) can hybridize to the probe (1804, 1814) on the array. As shown in FIG. 18B, 18D-18H, there can be a gap in the regions of the stochastically labeled molecule (1810, 1818, 1822, 1826, 1831, 1837) that can hybridize to the probe on the array. As shown in FIG. 18D-18F and 18H, different regions of the stochastically labeled molecule (1818, 1822, 1826, 1837) can hybridize to two or more probes on the array. Thus, the array probes can be in many different formats. The array probes can comprise a sequence that is complementary to a unique identifier region, a sequence that is complementary to the target molecule, or a combination thereof. Hybridization of the stochastically labeled molecule to the array can occur by a variety of ways. For example, two or more nucleotides of the stochastically labeled molecule can hybridize to one or more probes on the array. The two or more nucleotides of the stochastically labeled molecule that hybridize to the probes may be consecutive nucleotides, non-consecutive nucleotides, or a combination thereof. The stochastically labeled molecule

that is hybridized to the probe can be detected by any method known in the art. For example, the stochastically labeled molecules can be directly detected. Directly detecting the stochastically labeled molecule may comprise detection of a fluorophore, hapten, or detectable label. The stochastically labeled molecules can be indirectly detected. Indirect detection of the stochastically labeled molecule may comprise ligation or other enzymatic or non-enzymatic methods.

**[00216]** The array can be in a variety of formats. For example, the array can be in a 16-, 32-, 48-, 64-, 80-, 96-, 112-, 128-, 144-, 160-, 176-, 192-, 208-, 224-, 240-, 256-, 272-, 288-, 304-, 320-, 336-, 352-, 368-, 384-, or 400-format. Alternatively, the array is in an 8x60K, 4x180K, 2x400K, 1x1M format. In other instances, the array is in an 8x15K, 4x44K, 2x105K, 1x244K format.

**[00217]** The array can comprise a single array. The single array can be on a single substrate. Alternatively, the array is on multiple substrates. The array can comprise multiple formats. The array can comprise a plurality of arrays. The plurality of arrays can comprise two or more arrays. For example, the plurality of arrays can comprise at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 arrays. In some instances, at least two arrays of the plurality of arrays are identical. Alternatively, at least two arrays of the plurality of arrays are different.

**[00218]** In some instances, the array comprises symmetrical chambered areas. For example, the array comprises 0.5 x 0.5 mm, 1 x 1 mm, 1.5 x 1.5 mm, 2 x 2 mm, 2.5 x 2.5 mm, 3 x 3 mm, 3.5 x 3.5 mm, 4 x 4 mm, 4.5 x 4.5 mm, 5 x 5 mm, 5.5 x 5.5 mm, 6 x 6 mm, 6.5 x 6.5 mm, 7 x 7 mm, 7.5 x 7.5 mm, 8 x 8 mm, 8.5 x 8.5 mm, 9 x 9 mm, 9.5 x 9.5 mm, 10 x 10 mm, 10.5 x 10.5 mm, 11 x 11 mm, 11.5 x 11.5 mm, 12 x 12 mm, 12.5 x 12.5 mm, 13 x 13 mm, 13.5 x 13.5 mm, 14 x 14 mm, 14.5 x 14.5 mm, 15 x 15 mm, 15.5 x 15.5 mm, 16 x 16 mm, 16.5 x 16.5 mm, 17 x 17 mm, 17.5 x 17.5 mm, 18 x 18 mm, 18.5 x 18.5 mm, 19 x 19 mm, 19.5 x 19.5 mm, or 20 x 20 mm chambered areas. In some instances, the array comprises 6.5 x 6.5 mm chambered areas. Alternatively, the array comprises asymmetrical chambered areas. For example, the array comprises 6.5 x 0.5 mm, 6.5 x 1 mm, 6.5 x 1.5 mm, 6.5 x 2 mm, 6.5 x 2.5 mm, 6.5 x 3 mm, 6.5 x 3.5 mm, 6.5 x 4 mm, 6.5 x 4.5 mm, 6.5 x 5 mm, 6.5 x 5.5 mm, 6.5 x 6 mm, 6.5 x 6.5 mm, 6.5 x 7 mm, 6.5 x 7.5 mm, 6.5 x 8 mm, 6.5 x 8.5 mm, 6.5 x 9 mm, 6.5 x 9.5 mm, 6.5 x 10 mm, 6.5 x 10.5 mm, 6.5 x 11 mm, 6.5 x 11.5 mm, 6.5 x 12 mm, 6.5 x 12.5 mm, 6.5 x 13 mm, 6.5 x 13.5 mm, 6.5 x 14 mm, 6.5 x 14.5 mm, 6.5 x 15 mm, 6.5 x 15.5 mm, 6.5 x 16 mm, 6.5 x 16.5 mm, 6.5 x 17 mm, 6.5 x 17.5 mm, 6.5 x 18 mm, 6.5 x 18.5 mm, 6.5 x 19 mm, 6.5 x 19.5 mm, or 6.5 x 20 mm chambered areas.

**[00219]** The array can comprise at least about 1  $\mu\text{m}$ , 2  $\mu\text{m}$ , 3  $\mu\text{m}$ , 4  $\mu\text{m}$ , 5  $\mu\text{m}$ , 6  $\mu\text{m}$ , 7  $\mu\text{m}$ , 8  $\mu\text{m}$ , 9  $\mu\text{m}$ , 10  $\mu\text{m}$ , 15  $\mu\text{m}$ , 20  $\mu\text{m}$ , 25  $\mu\text{m}$ , 30  $\mu\text{m}$ , 35  $\mu\text{m}$ , 40  $\mu\text{m}$ , 45  $\mu\text{m}$ , 50  $\mu\text{m}$ , 55  $\mu\text{m}$ , 60  $\mu\text{m}$ , 65  $\mu\text{m}$ , 70  $\mu\text{m}$ , 75  $\mu\text{m}$ , 80  $\mu\text{m}$ , 85  $\mu\text{m}$ , 90  $\mu\text{m}$ , 95  $\mu\text{m}$ , 100  $\mu\text{m}$ , 125  $\mu\text{m}$ , 150  $\mu\text{m}$ , 175  $\mu\text{m}$ , 200  $\mu\text{m}$ , 225  $\mu\text{m}$ , 250  $\mu\text{m}$ , 275  $\mu\text{m}$ , 300  $\mu\text{m}$ , 325  $\mu\text{m}$ , 350  $\mu\text{m}$ , 375  $\mu\text{m}$ , 400  $\mu\text{m}$ , 425  $\mu\text{m}$ , 450  $\mu\text{m}$ , 475  $\mu\text{m}$ , or 500  $\mu\text{m}$  spots. In some instances, the array comprises 70  $\mu\text{m}$  spots.

**[00220]** The array can comprise at least about 1  $\mu\text{m}$ , 2  $\mu\text{m}$ , 3  $\mu\text{m}$ , 4  $\mu\text{m}$ , 5  $\mu\text{m}$ , 6  $\mu\text{m}$ , 7  $\mu\text{m}$ , 8  $\mu\text{m}$ , 9  $\mu\text{m}$ , 10  $\mu\text{m}$ , 15  $\mu\text{m}$ , 20  $\mu\text{m}$ , 25  $\mu\text{m}$ , 30  $\mu\text{m}$ , 35  $\mu\text{m}$ , 40  $\mu\text{m}$ , 45  $\mu\text{m}$ , 50  $\mu\text{m}$ , 55  $\mu\text{m}$ , 60  $\mu\text{m}$ , 65  $\mu\text{m}$ , 70  $\mu\text{m}$ , 75  $\mu\text{m}$ , 80  $\mu\text{m}$ , 85  $\mu\text{m}$ , 90  $\mu\text{m}$ , 95  $\mu\text{m}$ , 100  $\mu\text{m}$ , 125  $\mu\text{m}$ , 150  $\mu\text{m}$ , 175  $\mu\text{m}$ , 200  $\mu\text{m}$ , 225  $\mu\text{m}$ , 250  $\mu\text{m}$ , 275  $\mu\text{m}$ , 300  $\mu\text{m}$ , 325  $\mu\text{m}$ , 350  $\mu\text{m}$ , 375  $\mu\text{m}$ , 400  $\mu\text{m}$ , 425  $\mu\text{m}$ , 450  $\mu\text{m}$ , 475  $\mu\text{m}$ , 500  $\mu\text{m}$ , 525  $\mu\text{m}$ , 550  $\mu\text{m}$ , 575  $\mu\text{m}$ , 600  $\mu\text{m}$ , 625  $\mu\text{m}$ , 650  $\mu\text{m}$ , 675  $\mu\text{m}$ , 700  $\mu\text{m}$ , 725  $\mu\text{m}$ , 750  $\mu\text{m}$ , 775  $\mu\text{m}$ , 800  $\mu\text{m}$ , 825  $\mu\text{m}$ , 850  $\mu\text{m}$ , 875  $\mu\text{m}$ , 900  $\mu\text{m}$ , 925  $\mu\text{m}$ , 950  $\mu\text{m}$ , 975  $\mu\text{m}$ , 1000  $\mu\text{m}$  feature pitch. In some instances, the array comprises 161  $\mu\text{m}$  feature pitch.

**[00221]** The array can comprise one or more probes. In some instances, the array comprises at least about 5, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, or 100 probes. Alternatively, the array comprises at least about 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, or 3000 probes. The array can comprise at least about 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, or 10000 probes. In some instances, the array comprises at least about 960 probes. Alternatively, the array comprises at least about 2780 probes. The probes can be specific for the plurality of oligonucleotide tags. The probes can be specific for at least a portion of the plurality of oligonucleotide tags. The probes can be specific for at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97% or 100% of the total number of the plurality of oligonucleotide tags. Alternatively, the probes are specific for at least about 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97% or 100% of the total number of different oligonucleotide tags of the plurality of oligonucleotide tags. In other instances, the probes are non-specific probes. For example, the probes can be specific for a detectable label that is attached to the labeled-molecule. The probe can be streptavidin.

**[00222]** The array can be a printed array. In some instances, the printed array comprises one or more oligonucleotides attached to a substrate. For example, the printed array comprises 5' amine modified oligonucleotides attached to an epoxy silane substrate.

**[00223]** Alternatively, the array comprises a slide with one or more wells. The slide can

comprise at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 wells. Alternatively, the slide comprises at least about 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 650, 700, 750, 800, 850, 900, 950, or 1000 wells. In some instances, the slide comprises 16 wells. Alternatively, the slide comprises 96 wells. In other instances, the slide comprises at least about 80, 160, 240, 320, 400, 480, 560, 640, 720, 800, 880, or 960 wells.

**[00224]** In some instances, the solid support is an Affymetrix 3K tag array, Arrayjet non-contact printed array, or Applied Microarrays Inc (AMI) array. Alternatively, the support comprises a contact printer, impact printer, dot printer, or pin printer.

**[00225]** The solid support can comprise the use of beads that self assemble in microwells. For example, the solid support comprises Illumina's BeadArray Technology. Alternatively, the solid support comprises Abbott Molecular's Bead Array technology, and Applied Microarray's FlexiPlex<sup>TM</sup> system.

**[00226]** In other instances, the solid support is a plate. Examples of plates include, but are not limited to, MSD multi-array plates, MSD Multi-Spot<sup>®</sup> plates, microplate, ProteOn microplate, AlphaPlate, DELFIA plate, IsoPlate, and LumaPlate.

## **[00227] X. Enzymes**

**[00228]** The methods, kits, and systems disclosed herein comprise one or more enzymes. Examples of enzymes include, but are not limited to, ligases, reverse transcriptases, polymerases, and restriction nucleases. In some instances, attachment of the oligonucleotide tag to the molecules comprises the use of one or more ligases. Examples of ligases include, but are not limited to, DNA ligases such as DNA ligase I, DNA ligase III, DNA ligase IV, and T4 DNA ligase, and RNA ligases such as T4 RNA ligase I and T4 RNA ligase II.

**[00229]** The methods, kits, and systems disclosed herein can further comprise the use of one or more reverse transcriptases. In some instances, the reverse transcriptase is a HIV-1 reverse transcriptase, M-MLV reverse transcriptase, AMV reverse transcriptase, and telomerase reverse transcriptase. In some instances, the reverse transcriptase is M-MLV reverse transcriptase.

**[00230]** In some instances, the methods, kits, and systems disclosed herein comprise the use of one or more polymerases. Examples of polymerases include, but are not limited to, DNA polymerases and RNA polymerases. In some instances, the DNA polymerase is a DNA polymerase I, DNA polymerase II, DNA polymerase III holoenzyme, and DNA polymerase IV. Commercially available DNA polymerases include, but are not limited to, Bst 2.0 DNA Polymerase, Bst 2.0 WarmStart<sup>TM</sup> DNA Polymerase, Bst DNA Polymerase, Sulfolobus DNA

Polymerase IV, Taq DNA Polymerase, 9°N™m DNA Polymerase, Deep VentR™ (exo-) DNA Polymerase, Deep VentR™ DNA Polymerase, Hemo KlenTaq™, LongAmp® Taq DNA Polymerase, OneTaq® DNA Polymerase, Phusion® DNA Polymerase, Q5™ High-Fidelity DNA Polymerase, Therminator™ γ DNA Polymerase, Therminator™ DNA Polymerase, Therminator™ II DNA Polymerase, Therminator™ III DNA Polymerase, VentR® DNA Polymerase, VentR® (exo-) DNA Polymerase, Bsu DNA Polymerase, phi29 DNA Polymerase, T4 DNA Polymerase, T7 DNA Polymerase, and Terminal Transferase. Alternatively, the polymerase is an RNA polymerases such as RNA polymerase I, RNA polymerase II, RNA polymerase III, *E. coli* Poly(A) polymerase, phi6 RNA polymerase (RdRP), Poly(U) polymerase, SP6 RNA polymerase, and T7 RNA polymerase.

**[00231]** In some instances, the methods, kits, and systems disclosed herein comprise one or more restriction enzymes. Restriction enzymes include type I, type II, type III, and type IV restriction enzymes. In some instances, Type I enzymes are complex, multisubunit, combination restriction-and-modification enzymes that cut DNA at random far from their recognition sequences. Generally, type II enzymes cut DNA at defined positions close to or within their recognition sequences. They can produce discrete restriction fragments and distinct gel banding patterns. Type III enzymes are also large combination restriction-and-modification enzymes. They often cleave outside of their recognition sequences and can require two such sequences in opposite orientations within the same DNA molecule to accomplish cleavage; they rarely give complete digests. In some instances, type IV enzymes recognize modified, typically methylated DNA and can be exemplified by the McrBC and Mrr systems of *E. coli*.

**[00232] XI. Miscellaneous Components**

**[00233]** The methods, kits, and systems disclosed herein can comprise the use of one or more reagents. Examples of reagents include, but are not limited to, PCR reagents, ligation reagents, reverse transcription reagents, enzyme reagents, hybridization reagents, sample preparation reagents, and reagents for nucleic acid purification and/or isolation.

**[00234]** The methods, kits, and systems disclosed herein can comprise the use of one or more buffers. Examples of buffers include, but are not limited to, wash buffers, ligation buffers, hybridization buffers, amplification buffers, and reverse transcription buffers. In some instances, the hybridization buffer is a commercially available buffer, such as TMAC Hyb solution, SSPE hybridization solution, and ECONO™ hybridization buffer. The buffers disclosed herein can comprise one or more detergents.

**[00235]** The methods, kits, and systems disclosed herein can comprise the use of one or

more carriers. Carriers can enhance or improve the efficiency of one or more reactions disclosed herein (e.g., ligation reaction, reverse transcription, amplification, hybridization). Carriers can decrease or prevent non-specific loss of the molecules or any products thereof (e.g., labeled-molecule, labeled-cDNA molecule, labeled-amplicon). For example, the carrier can decrease non-specific loss of a labeled-molecule through absorption to surfaces. The carrier can decrease the affinity of the molecule, labeled-molecule, or any product thereof to a surface or substrate (e.g., container, eppendorf tube, pipet tip). Alternatively, the carrier can increase the affinity of the molecule or any product thereof to a surface or substrate (e.g., bead, array, glass, slide, chip). Carriers can protect the molecule or any product thereof from degradation. For example, carriers can protect an RNA molecule or any product thereof from ribonucleases. Alternatively, carriers can protect a DNA molecule or any product thereof from a DNase. Examples of carriers include, but are not limited to, nucleic acid molecules such as DNA and/or RNA, or polypeptides. Examples of DNA carriers include plasmids, vectors, polyadenylated DNA, and DNA oligonucleotides. Examples of RNA carriers include polyadenylated RNA, phage RNA, phage MS2 RNA, *E.coli* RNA, yeast RNA, yeast tRNA, mammalian RNA, mammalian tRNA, short polyadenylated synthetic ribonucleotides and RNA oligonucleotides. The RNA carrier can be a polyadenylated RNA. Alternatively, the RNA carrier can be a non-polyadenylated RNA. In some instances, the carrier is from a bacteria, yeast, or virus. For example, the carrier can be a nucleic acid molecule or a polypeptide derived from a bacteria, yeast or virus. For example, the carrier is a protein from *Bacillus subtilis*. In another example, the carrier is a nucleic acid molecule from *Escherichia coli*. Alternatively, the carrier is a nucleic acid molecule or peptide from a mammal (e.g., human, mouse, goat, rat, cow, sheep, pig, dog, or rabbit), avian, amphibian, or reptile.

**[00236]** The methods, kits, and systems disclosed herein can comprise the use of one or more control agents. Control agents can include control oligos, inactive enzymes, non-specific competitors. Alternatively, the control agents comprise bright hybridization, bright probe controls, nucleic acid templates, spike-in templates, PCR amplification controls. The PCR amplification controls can be positive controls. In other instances, the PCR amplification controls are negative controls. The nucleic acid template controls can be of known concentrations. The control agents can comprise one or more labels.

**[00237]** Spike-in templates can be templates that are added to a reaction or sample. For example, a spike-in template can be added to an amplification reaction. The spike-in template can be added to the amplification reaction any time after the first amplification cycle. In some instances, the spike-in template is added to the amplification reaction after the 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup>,

6<sup>th</sup>, 7<sup>th</sup>, 8<sup>th</sup>, 9<sup>th</sup>, 10<sup>th</sup>, 11<sup>th</sup>, 12<sup>th</sup>, 13<sup>th</sup>, 14<sup>th</sup>, 15<sup>th</sup>, 20<sup>th</sup>, 25<sup>th</sup>, 30<sup>th</sup>, 35<sup>th</sup>, 40<sup>th</sup>, 45<sup>th</sup>, or 50<sup>th</sup> amplification cycle. The spike-in template can be added to the amplification reaction any time before the last amplification cycle. The spike-in template can comprise one or more nucleotides or nucleic acid base pairs. The spike-in template can comprise DNA, RNA, or any combination thereof. The spike-in template can comprise one or more labels.

**[00238]** The methods, kits, and systems disclosed herein can comprise the use of one or more pipet tips and/or containers (e.g., tubes, vials, multiwell plates). In some instances, the pipet tips are low binding pipet tips. Alternatively, or additionally, the containers can be low binding containers. Low binding pipet tips and low binding containers can have reduced leaching and/or subsequent sample degradation associated with silicone-based tips and non-low binding containers. Low binding pipet tips and low binding containers can have reduced sample binding as compared to non-low binding pipet tips and containers. Examples of low binding tips include, but are not limited to, Corning® DeckWorks™ low binding tips and Avant Premium low binding graduated tips. A non-limiting list of low-binding containers include Corning® Costar® low binding microcentrifuge tubes and Cosmobrand low binding PCR tubes and microcentrifuge tubes.

**[00239] XIII. Indications**

**[00240]** The methods disclosed herein may be used in gene expression monitoring, transcript profiling, library screening, genotyping, epigenetic analysis, methylation pattern analysis, tumor typing, pharmacogenomics, agrigenetics, pathogen profiling and detection and diagnostics. Gene expression monitoring and profiling methods have been shown in U.S. Patent Nos. 5,800,992, 6,013,449, 6,020,135, 6,033,860, 6,040,138, 6,177,248 and 6,309,822. Genotyping and uses therefore are shown in U.S. Patent Publication Nos. 20030036069 and 20070065816 and U.S. Patent Nos. 5,856,092, 6,300,063, 5,858,659, 6,284,460, 6,361,947, 6,368,799 and 6,333,179. Other uses are embodied in U.S. Patent Nos. 5,871,928, 5,902,723, 6,045,996, 5,541,061, and 6,197,506.

**[00241]** Disclosed herein are methods, kits and compositions for detection, monitoring, and/or prognosis of a disease or condition in a subject. Generally, the method comprises (a) stochastically labeling a molecule to produce a stochastically-labeled molecule; and (b) detecting and/or quantifying the stochastically-labeled molecule, thereby detecting, monitoring, and/or prognosing a disease or condition in a subject. Detecting a disease or condition can comprise diagnosing a disease or condition.

**[00242]** Monitoring a disease or condition in a subject can further comprise monitoring a therapeutic regimen. Monitoring a therapeutic regimen can comprise determining the efficacy

of a therapeutic regimen. In some instances, monitoring a therapeutic regimen comprises administrating, terminating, adding, or altering a therapeutic regimen. Altering a therapeutic regimen can comprise increasing or reducing the dosage, dosing frequency, or mode of administration of a therapeutic regimen. A therapeutic regimen can comprise one or more therapeutic drugs. The therapeutic drugs can be an anticancer drug, antiviral drug, antibacterial drug, antipathogenic drug, or any combination thereof.

**[00243] *A. Cancer***

**[00244]** In some instances, the disease or condition is a cancer. The molecules to be stochastically labeled can be from a cancerous cell or tissue. In some instances, the cancer is a sarcoma, carcinoma, lymphoma or leukemia. Sarcomas are cancers of the bone, cartilage, fat, muscle, blood vessels, or other connective or supportive tissue. Sarcomas include, but are not limited to, bone cancer, fibrosarcoma, chondrosarcoma, Ewing's sarcoma, malignant hemangioendothelioma, malignant schwannoma, bilateral vestibular schwannoma, osteosarcoma, soft tissue sarcomas (e.g. alveolar soft part sarcoma, angiosarcoma, cystosarcoma phylloides, dermatofibrosarcoma, desmoid tumor, epithelioid sarcoma, extraskeletal osteosarcoma, fibrosarcoma, hemangiopericytoma, hemangiosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, lymphosarcoma, malignant fibrous histiocytoma, neurofibrosarcoma, rhabdomyosarcoma, and synovial sarcoma).

**[00245]** Carcinomas are cancers that begin in the epithelial cells, which are cells that cover the surface of the body, produce hormones, and make up glands. By way of non-limiting example, carcinomas include breast cancer, pancreatic cancer, lung cancer, colon cancer, colorectal cancer, rectal cancer, kidney cancer, bladder cancer, stomach cancer, prostate cancer, liver cancer, ovarian cancer, brain cancer, vaginal cancer, vulvar cancer, uterine cancer, oral cancer, penile cancer, testicular cancer, esophageal cancer, skin cancer, cancer of the fallopian tubes, head and neck cancer, gastrointestinal stromal cancer, adenocarcinoma, cutaneous or intraocular melanoma, cancer of the anal region, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, cancer of the urethra, cancer of the renal pelvis, cancer of the ureter, cancer of the endometrium, cancer of the cervix, cancer of the pituitary gland, neoplasms of the central nervous system (CNS), primary CNS lymphoma, brain stem glioma, and spinal axis tumors. In some instances, the cancer is a skin cancer, such as a basal cell carcinoma, squamous, melanoma, nonmelanoma, or actinic (solar) keratosis.

**[00246]** In some instances, the cancer is a lung cancer. Lung cancer can start in the airways that branch off the trachea to supply the lungs (bronchi) or the small air sacs of the

lung (the alveoli). Lung cancers include non-small cell lung carcinoma (NSCLC), small cell lung carcinoma, and mesothelioma. Examples of NSCLC include squamous cell carcinoma, adenocarcinoma, and large cell carcinoma. The mesothelioma may be a cancerous tumor of the lining of the lung and chest cavity (pleura) or lining of the abdomen (peritoneum). The mesothelioma may be due to asbestos exposure. The cancer may be a brain cancer, such as a glioblastoma.

**[00247]** Alternatively, the cancer may be a central nervous system (CNS) tumor. CNS tumors may be classified as gliomas or nongliomas. The glioma may be malignant glioma, high grade glioma, diffuse intrinsic pontine glioma. Examples of gliomas include astrocytomas, oligodendrogiomas (or mixtures of oligodendrogioma and astocytoma elements), and ependymomas. Astrocytomas include, but are not limited to, low-grade astrocytomas, anaplastic astrocytomas, glioblastoma multiforme, pilocytic astrocytoma, pleomorphic xanthoastrocytoma, and subependymal giant cell astrocytoma. Oligodendrogiomas include low-grade oligodendrogiomas (or oligoastrocytomas) and anaplastic oligodendriogliomas. Nongliomas include meningiomas, pituitary adenomas, primary CNS lymphomas, and medulloblastomas. In some instances, the cancer is a meningioma.

**[00248]** The leukemia may be an acute lymphocytic leukemia, acute myelocytic leukemia, chronic lymphocytic leukemia, or chronic myelocytic leukemia. Additional types of leukemias include hairy cell leukemia, chronic myelomonocytic leukemia, and juvenile myelomonocytic leukemia.

**[00249]** Lymphomas are cancers of the lymphocytes and may develop from either B or T lymphocytes. The two major types of lymphoma are Hodgkin's lymphoma, previously known as Hodgkin's disease, and non-Hodgkin's lymphoma. Hodgkin's lymphoma is marked by the presence of the Reed-Sternberg cell. Non-Hodgkin's lymphomas are all lymphomas which are not Hodgkin's lymphoma. Non-Hodgkin lymphomas may be indolent lymphomas and aggressive lymphomas. Non-Hodgkin's lymphomas include, but are not limited to, diffuse large B cell lymphoma, follicular lymphoma, mucosa-associated lymphatic tissue lymphoma (MALT), small cell lymphocytic lymphoma, mantle cell lymphoma, Burkitt's lymphoma, mediastinal large B cell lymphoma, Waldenström macroglobulinemia, nodal marginal zone B cell lymphoma (NMZL), splenic marginal zone lymphoma (SMZL), extranodal marginal zone B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, and lymphomatoid granulomatosis.

**[00250]** *B. Pathogenic Infection*

**[00251]** In some instances, the disease or condition is a pathogenic infection. The molecules to be stochastically labeled can be from a pathogen. The pathogen can be a virus, bacterium, fungi, or protozoan. In some instances, the pathogen may be a protozoan, such as *Acanthamoeba* (e.g., *A. astronyxis*, *A. castellanii*, *A. culbertsoni*, *A. hatchetti*, *A. polyphaga*, *A. rhyosodes*, *A. healyi*, *A. divionensis*), *Brachiola* (e.g., *B. connori*, *B. vesicularum*), *Cryptosporidium* (e.g., *C. parvum*), *Cyclospora* (e.g., *C. cayetanensis*), *Encephalitozoon* (e.g., *E. cuniculi*, *E. hellem*, *E. intestinalis*), *Entamoeba* (e.g., *E. histolytica*), *Enterocytozoon* (e.g., *E. bieneusi*), *Giardia* (e.g., *G. lamblia*), *Isospora* (e.g., *I. belli*), *Microsporidium* (e.g., *M. africanum*, *M. ceylonensis*), *Naegleria* (e.g., *N. fowleri*), *Nosema* (e.g., *N. algerae*, *N. oocularum*), *Pleistophora*, *Trachipleistophora* (e.g., *T. anthropophthora*, *T. hominis*), and *Vittaforma* (e.g., *V. corneae*). The pathogen may be a fungus, such as, *Candida*, *Aspergillus*, *Cryptococcus*, *Histoplasma*, *Pneumocystis*, and *Stachybotrys*.

**[00252]** The pathogen can be a bacterium. Exemplary bacteria include, but are not limited to, *Bordetella*, *Borrelia*, *Brucella*, *Campylobacter*, *Chlamydia*, *Chlamydophila*, *Clostridium*, *Corynebacterium*, *Enterococcus*, *Escherichia*, *Francisella*, *Haemophilus*, *Helicobacter*, *Legionella*, *Leptospira*, *Listeria*, *Mycobacterium*, *Mycoplasma*, *Neisseria*, *Pseudomonas*, *Rickettsia*, *Salmonella*, *Shigella*, *Staphylococcus*, *Streptococcus*, *Treponema*, *Vibrio*, or *Yersinia*.

**[00253]** The virus can be a reverse transcribing virus. Examples of reverse transcribing viruses include, but are not limited to, single stranded RNA-RT (ssRNA-RT) virus and double-stranded DNA –RT (dsDNA-RT) virus. Non-limiting examples of ssRNA-RT viruses include retroviruses, alpharetrovirus, betaretrovirus, gammaretrovirus, deltaretrovirus, epsilonretrovirus, lentivirus, spuma virus, metavirus, and pseudoviruses. Non-limiting examples of dsDNA-RT viruses include hepadenovirus and caulimovirus. Alternatively, the virus is a DNA virus or RNA virus. The DNA virus can be a double-stranded DNA (dsDNA) virus. In some instances, the dsDNA virus is an adenovirus, herpes virus, or pox virus. Examples of adenoviruses include, but are not limited to, adenovirus and infectious canine hepatitis virus. Examples of herpes viruses include, but are not limited to, herpes simplex virus, varicella-zoster virus, cytomegalovirus, and Epstein-Barr virus. A non-limiting list of pox viruses includes smallpox virus, cow pox virus, sheep pox virus, monkey pox virus, and vaccinia virus. The DNA virus can be a single-stranded DNA (ssDNA) virus. The ssDNA virus can be a parvovirus. Examples of parvoviruses include, but are not limited to, parvovirus B19, canine parvovirus, mouse parvovirus, porcine parvovirus, feline panleukopenia, and Mink enteritis virus.

**[00254]** Alternatively, the virus is a RNA virus. The RNA virus can be a double-stranded RNA (dsRNA) virus, (+) sense single-stranded RNA virus ((+)ssRNA) virus, or (-) sense single-stranded ((-)ssRNA) virus. A non-limiting list of dsRNA viruses include reovirus, orthoreovirus, cypovirus, rotavirus, bluetongue virus, and phytoreovirus. Examples of (+) ssRNA viruses include, but are not limited to, picornavirus and togavirus. Examples of picornaviruses include, but are not limited to, enterovirus, rhinovirus, hepatovirus, cardiovirus, aphthovirus, poliovirus, parechovirus, erbovirus, kobuvirus, teschovirus, and coxsackie. In some instances, the togavirus is a rubella virus, Sindbis virus, Eastern equine encephalitis virus, Western equine encephalitis virus, Venezuelan equine encephalitis virus, Ross River virus, O'nyong'nyong virus, Chikungunya, or Semliki Forest virus. A non-limiting list of (-) ssRNA viruses include orthomyxovirus and rhabdovirus. Examples of orthomyxoviruses include, but are not limited to, influenza virus a, influenza virus B, influenza virus C, isavirus, and thogotovirus. Examples of rhabdoviruses include, but are not limited to, cytorhabdovirus, dichorhabdovirus, ephemeroirus, lyssavirus, novirhabdovirus, and vesiculovirus.

**[00255]** *C. Fetal disorders*

**[00256]** In some instances, the disease or condition is pregnancy. The methods disclosed herein can comprise diagnosing a fetal condition in a pregnant subject. The methods disclosed herein can comprise identifying fetal mutations or genetic abnormalities. The molecules to be stochastically labeled can be from a fetal cell or tissue. Alternatively, or additionally, the molecules to be labeled can be from the pregnant subject.

**[00257]** The methods, kits, and systems disclosed herein can be used in the diagnosis, prediction or monitoring of autosomal trisomies (e.g., Trisomy 13, 15, 16, 18, 21, or 22). In some cases the trisomy may be associated with an increased chance of miscarriage (e.g., Trisomy 15, 16, or 22). In other cases, the trisomy that is detected is a liveborn trisomy that may indicate that an infant will be born with birth defects (e.g., Trisomy 13 (Patau Syndrome), Trisomy 18 (Edwards Syndrome), and Trisomy 21 (Down Syndrome)). The abnormality may also be of a sex chromosome (e.g., XXY (Klinefelter's Syndrome), XYY (Jacobs Syndrome), or XXX (Trisomy X)). The molecule(s) to be labeled can be on one or more of the following chromosomes: 13, 18, 21, X, or Y. For example, the molecule is on chromosome 21 and/or on chromosome 18, and/or on chromosome 13.

**[00258]** Further fetal conditions that can be determined based on the methods, kits, and systems disclosed herein include monosomy of one or more chromosomes (X chromosome monosomy, also known as Turner's syndrome), trisomy of one or more chromosomes (13, 18,

21, and X), tetrasomy and pentasomy of one or more chromosomes (which in humans is most commonly observed in the sex chromosomes, e.g. XXXX, XYY, XXXY, XYY, XXXXX, XXXXY, XXXYY, XYYYY and XXYYY), monoploidy, triploidy (three of every chromosome, e.g. 69 chromosomes in humans), tetraploidy (four of every chromosome, e.g. 92 chromosomes in humans), pentaploidy and multiploidy.

### EXEMPLARY EMBODIMENTS

**[00259]** Disclosed herein, in some embodiments, are methods, kits, and systems for digital reverse transcription of an RNA molecule. In some instances, the method comprises (a) producing a labeled-RNA molecule by contacting a sample comprising a plurality of RNA molecules with a plurality of oligonucleotide tags, wherein (i) the plurality of RNA molecules comprises two or more RNA molecules comprising at least two different sequences; and (ii) the plurality of oligonucleotide tags comprises oligonucleotide tags comprising two or more different unique identifier sequences; (b) conducting a first strand synthesis reaction by contacting the labeled-RNA molecules with a reverse transcriptase enzyme to produce a labeled-cDNA molecule; and (c) detecting the labeled-cDNA molecule by hybridizing the labeled-cDNA molecule to a solid support.

**[00260]** Producing a labeled-RNA molecule can comprise attaching the oligonucleotide tag to the RNA molecule. In some instances, the oligonucleotide tag is attached to the RNA molecule by hybridization. In other instances, the oligonucleotide tag is attached to the RNA molecule by ligation. The attachment of the oligonucleotide tag can comprise the use of a ligase enzyme. The oligonucleotide tag can be attached to any portion of the RNA molecule. For example, the oligonucleotide tag can be attached to the 5' end of the RNA molecule. Alternatively, the oligonucleotide tag is attached to the 3' end of the RNA molecule. In other instances, the oligonucleotide tag is attached to an internal region of the RNA molecule. Attachment of the oligonucleotide tag to the RNA molecule can comprise the use of one or more adaptor molecules.

**[00261]** In some instances, the oligonucleotide tag comprises a target specific region. The target specific region can enable attachment of the plurality of oligonucleotide tags to at least one RNA molecule. The target specific region can enable attachment of the plurality of oligonucleotide tags to two or more different RNA molecules. In some instances, the target specific region enables attachment of the plurality of oligonucleotide tags to at least about 3, 4, or 5 different RNA molecules. Alternatively, the target specific region enables attachment of the plurality of oligonucleotide tags to at least about 6, 7, 8, 9, or 10 different RNA

molecules. In other instances, the target specific region enables attachment of the plurality of oligonucleotide tags to at least about 11, 12, 13, 14, or 15 different RNA molecules. The target specific region can comprise an oligodT sequence. Alternatively, the target specific region comprises a random sequence that can attach to any portion of the RNA molecule.

**[00262]** In some instances, the oligonucleotide tag further comprises a universal primer region. The unique identifier region can be placed between the universal primer region and the target specific region. The oligonucleotide tag can be at least one nucleotide in length. The unique identifier region can be at least one nucleotides in length. The target specific region can be at least one nucleotide in length. The universal primer region can be at least one nucleotide in length. The oligonucleotide tag can comprise one or more nucleotide moieties. Alternatively, or additionally, the nucleotide tag comprises one or more non-nucleotide moieties.

**[00263]** In some instances, producing the labeled-RNA molecule further comprises a dNTP mix, annealing buffer, ligase, ligation buffer, or any combination thereof. Conducting the first strand synthesis reaction can further comprise a first strand buffer, dithiothreitol (DTT), RNase inhibitor, DNA polymerase, or any combination thereof.

**[00264]** The first strand synthesis reaction can further comprise a thermal cycler. The first strand synthesis reaction can further comprise a thermal cycler program comprising 1 cycle of 50° C for 60 minutes, followed by 3 cycles of 94° C for 2 minutes, 58° C for 2 minutes, and 68° C for 2 minutes, followed by 1 cycle of 4° C for at least 2 minutes. The methods disclosed herein can further comprise contacting the labeled-cDNA molecule with a target specific primer. The target specific primer can be a uracil-containing DNA primer. The target specific primer can hybridize to the labeled-cDNA molecule and a polymerase chain reaction can be conducted to produce a double-stranded labeled-cDNA molecule.

**[00265]** The sample can be further treated with one or more enzymes to remove or degrade RNA molecules, labeled-RNA molecules, unbound oligonucleotide tags, and/or unbound target specific primers. For example, the sample can be treated with an RNase enzyme to remove the RNA molecules (labeled and/or unbound RNA molecules) from the sample. Alternatively, the sample can be treated with a uracil DNA glycosylase (UDG) to hydrolyze the uracil from the DNA.

**[00266]** The method can further comprise conducting a polymerase chain reaction (PCR) to produce labeled-amplicons. In some instances, the polymerase chain reaction is a nested PCR. The nested PCR can comprise conducting a first PCR comprising mixing the double-stranded labeled-cDNA molecule with a first PCR mixture comprising a first target specific

PCR primer, universal PCR primer, polymerase buffer, DNA polymerase, dNTP mix, or any combination thereof. The first PCR can be conducted in thermal cycler. The first PCR can comprise a thermal cycler program comprising 1 cycle of 94° C for 2 minutes, followed by 30 cycles of 94° C for 20 seconds, 58° C for 20 seconds, and 68° C for 20 seconds, followed by 1 cycle of 68° C for 4 minutes and 1 cycle of 4° C for at least 2 minutes. The nested PCR can comprise conducting a second PCR comprising mixing at least a portion of the amplicons produced in the first PCR reaction with a second PCR mixture comprising a second target specific PCR primer, labeled-universal PCR primer, polymerase buffer, DNA polymerase, dNTP mix, or any combination thereof. The second target specific primer can hybridize to a region in the labeled molecule that is downstream of the first target specific primer. The labeled-universal PCR primer is labeled with a detectable label. In some instances, the labeled-universal PCR primer is a Cy3-labeled universal PCR primer. Alternatively, the labeled-universal PCR primer is a TYE 563-labeled universal PCR primer. The second PCR can be conducted in thermal cycler. The second PCR can comprise a thermal cycler program comprising 1 cycle of 94° C for 2 minutes, followed by 30 cycles of 94° C for 20 seconds, 58° C for 20 seconds, and 68° C for 20 seconds, followed by 1 cycle of 68° C for 4 minutes and 1 cycle of 4° C for at least 2 minutes. The second PCR of the nested PCR can produce a labeled-amplicon comprising the cDNA molecule, oligonucleotide tag and the detectable label. In some instances, the labeled-cDNA molecule of step 1c is the labeled-amplicon produced by the second PCR of the nested PCR.

**[00267]** In some instances, detecting the labeled-cDNA molecule comprises hybridizing at least a portion of the sample comprising the labeled-amplicons comprising the cDNA molecule, oligonucleotide tag and the detectable label to a solid support. Hybridizing at least a portion of the sample comprising the labeled-amplicons can comprise a hybridization mixture comprising at least a portion of the sample comprising the labeled-amplicons produced in the second PCR of nested PCR, control oligo, hybridization buffer, or any combination thereof. The control oligo can comprise the detectable label conjugated to an oligonucleotide. The detectable label is the same as the detectable label in the labeled-amplicon. For example, the labeled-amplicon comprises a Cy3 label and the control oligo comprises a Cy3-labeled oligonucleotide. The labeled-amplicons in the hybridization mixture are denatured. In some instances, denaturing the labeled-amplicons comprises incubating the hybridization mixture at 95° C. In some instances, the hybridization mixture is incubated at 95° C for at least about 1, 2, 3, 4, or 5 minutes. After denaturation of the labeled-amplicons, the hybridization mixture is incubated at 4° C for at least 2 minutes. Hybridization

of the labeled-amplicon to the support can comprise adding at least a portion of the hybridization mixture to the solid support. In some instances, hybridization of the labeled-amplicon to the solid support comprises adding at least a portion of the hybridization mixture to a well of an AMI array slide. The labeled-amplicon can be hybridized to the support for at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, or 48 hours. The labeled-amplicon can be hybridized to the support for at least about 4 hours. The labeled-amplicon can be hybridized to the support overnight. Alternatively, the labeled-amplicon is hybridized to the support for about 12-14 hours. In other instances, the labeled-amplicon is hybridized to the support for about 3-5 hours, 4-6 hours, 6-8 hours, 8-10 hours, 9-11 hours, 13-15 hours, 14-16 hours, 17-19 hours, or 18-20 hours. Hybridization of the labeled-amplicon to the support can comprise contacting the support with the labeled-amplicon and incubating the labeled-amplicon and support at a hybridization temperature. In some instances, the hybridization temperature is at least about 20 °C, 21 °C, 22 °C, 23 °C, 24 °C, 25 °C, 26 °C, 27 °C, 28 °C, 29 °C, 30 °C, 32 °C, 34 °C, 36 °C, 38 °C, 40 °C, 45 °C, 50 °C, 55 °C, 60 °C, or 65 °C.

**[00268]** The solid support can comprise a plurality of probes. The plurality of probes can comprise a sequence that is complementary to at least a portion of the labeled-cDNA molecule or labeled-amplicon. The plurality of probes can be arranged on the solid support in discrete regions, wherein a discrete region on the solid support comprises probes of identical or near-identical sequences. In some instances, two or more discrete regions on the solid support comprise two different probes comprising sequences complementary to the sequence of two different unique identifier regions of the oligonucleotide tag.

**[00269]** The method further comprise covering the array slide with an adhesive to produce a sealed array slide. The sealed array slide can be incubated at 37° C. The sealed array slide can be incubated at 37° C overnight. In some instances, the sealed array is incubated at 37° C for at least about 12-14 hours. After incubating the sealed array at 37° C, the method can further comprise removing the sealed array from 37° C. The hybridization mixture can be removed from each well. The hybridization mixture can be stored at -20° C. Alternatively, the hybridization mixture is discarded.

**[00270]** The method can further comprise washing the wells with a first wash buffer. Washing the wells comprises adding a wash buffer to the well and then aspirating the wash buffer. Additionally, a second wash can be performed with the same or a second wash buffer. Once the wash buffers have been aspirated from the wells, the array slide can be scanned. In

some instances, the array slide is scanned dry (e.g., fluid is removed from the wells). Alternatively, the array slide is scanned wet (e.g., fluid is in the wells). The array slide can be scanned by a scanner.

**[00271]** The method can comprise fragmentation of the amplification products (e.g., labeled amplicons) to produce fragmented labeled-amplicons. The fragmented labeled-amplicons can be attached to the solid support. The methods disclosed herein can further comprise attaching a detectable label to the labeled-molecules, labeled-amplicons, or fragmented-labeled amplicons. The detectable label can be attached to the labeled-molecules, labeled-amplicons, or fragmented-labeled amplicons prior to attachment of the labeled-molecules, labeled-amplicons, or fragmented-labeled amplicons to the solid support. Alternatively, the detectable label is attached to the labeled-molecules, labeled-amplicons, or fragmented-labeled amplicons after attachment of the labeled-molecules, labeled-amplicons, or fragmented-labeled amplicons to the solid support. The methods disclosed herein can comprise attaching two or more detectable labels to the labeled-molecules, labeled-amplicons, or fragmented-labeled amplicons. In some instances, a detectable label is the labeled-cDNA molecule and the detectable label is incorporated into the labeled-amplicon. For example, a Cy3 universal PCR primer is annealed to the labeled-cDNA molecule. Amplification of the labeled-cDNA molecule with Cy3 universal PCR primer can produce a Cy3-labeled amplicons. The methods disclosed herein can further comprise attaching a second detectable label to the first-detectable labeled-molecule. For example, The methods disclosed herein can comprise attaching biotin to the Cy3-labeled amplicons to produce biotin/Cy3-labeled amplicons.

**[00272]** In some instances, detecting the labeled-cDNA molecule comprises a fluorescent reader. The fluorescent reader can be a Sensovation FLAIR instrument.

**[00273]** In some instances, the data from the scanner is stored on a computer. Alternatively, or additionally, the data from the scanner is exported. In some instances, the data from the scanner is transmitted electronically. Exportation and/or transmission of the data can comprise one or more computer networks.

**[00274]** Further disclosed herein are methods, kits, and systems for stochastically labeling a molecule. Generally, the method comprises contacting a sample comprising a plurality of molecules with a plurality of oligonucleotide tags and randomly attaching one or more oligonucleotide tags from the plurality of oligonucleotide tags to one or more molecules in the sample. The plurality of oligonucleotide tags comprises oligonucleotide tags comprising two or more different unique identifier regions.

**[00275]** In some instances, the methods, kits, and systems comprise concentrations of the different oligonucleotide tags in the plurality of oligonucleotide tags. For example, the different oligonucleotide tags are present in the plurality of oligonucleotide tags in the same concentration. Alternatively, the concentration of at least one oligonucleotide tag in the plurality of oligonucleotide tags is greater than the concentration of at least one other oligonucleotide tag in the plurality of oligonucleotide tags. The concentration of the at least one oligonucleotide tag in the plurality of oligonucleotide tags is at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 times greater than the concentration of the at least one other oligonucleotide tag in the plurality of oligonucleotide tags. In some instances, the concentration of at least one oligonucleotide tag in the plurality of oligonucleotide tags is less than the concentration of at least one other oligonucleotide tag in the plurality of oligonucleotide tags. The concentration of the at least one oligonucleotide tag in the plurality of oligonucleotide tags is at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 times less than the concentration of the at least one other oligonucleotide tag in the plurality of oligonucleotide tags. In some instances, at least about 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97%, or 100% of the different oligonucleotide tags in the plurality of oligonucleotide tags are present in the plurality of oligonucleotide tags in the same or similar concentration. Alternatively, at least about 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 97%, or 100% of the different oligonucleotide tags in the plurality of oligonucleotide tags are present in the plurality of oligonucleotide tags in different concentrations.

**[00276]** The oligonucleotide tags can further comprise a target specific region, universal primer binding site, or any combination thereof. In some instances, the unique identifier region is between the target specific region and the universal primer binding site. The oligonucleotide tags can be attached to the molecules by hybridization, ligation, or any combination thereof. In some instances, one or more oligonucleotide tags are attached to a molecule. The oligonucleotide tag can be attached to the 5' end of the molecule, 3' end of the molecule, an internal site within the molecule, or any combination thereof. One or both ends of the oligonucleotide tag can be attached to the molecule.

**[00277]** The molecule can be a polynucleotide. The polynucleotide can comprise RNA, DNA, or any combination thereof. The molecule can be an RNA molecule. The RNA molecule can be an mRNA. The molecule can be polyadenylated. Alternatively, the molecule

is not polyadenylated.

**[00278]** Further disclosed herein are digital pre-amplification methods for increasing the quantity of a nucleic acid molecule in a sample. Generally, the method comprises (a) stochastically labeling a nucleic acid molecule in a sample by any of the methods disclosed herein to produce a labeled-nucleic acid molecule, wherein the labeled-nucleic acid molecule comprises an oligonucleotide tag attached to the nucleic acid molecule; and (b) amplifying the labeled-nucleic acid molecule to produce a plurality of labeled-amplicons, wherein a labeled-amplicon in the plurality of labeled-amplicons is a copy of the labeled-nucleic acid molecule. The labeled-nucleic acid molecule of step (a) can be repeatedly amplified to increase the quantity of the nucleic acid molecule in the sample. The oligonucleotide tag comprises a unique identifier region that can be used to distinguish identical or nearly identical nucleic acid molecules.

**[00279]** Stochastic labeling of the nucleic acid molecule prior to amplification can enable the identification of clonally replicated molecules originating from the sample template parent molecule. Stochastic labeling of the nucleic acid molecule prior to amplification can allow for controlled amplification of the nucleic acid molecule, wherein the amplification of an individual nucleic acid molecule can be tracked and monitored by the oligonucleotide label. The digital pre-amplification method can account for the true abundance levels of nucleic acid molecules in a sample. This method can be particularly useful for samples comprising limited quantities of a nucleic acid molecule. For example, this method can be used to increase the quantity of a nucleic acid molecule from a single cell. Stochastic labeling of the nucleic acid molecule in the cell followed by amplification of the labeled-nucleic acid molecules can allow for more precise quantitative measurements of the nucleic acid molecules.

**[00280]** In some instances, the labeled-nucleic acid molecules in the sample are amplified at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 times. Alternatively, the labeled-nucleic acid molecules in the sample are amplified at least about 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, or 100 times.

**[00281]** Digital pre-amplification of the nucleic acid molecules can enable repeated sampling of the nucleic acid molecules in the sample without depletion of the original sample. Repeated sampling of the nucleic acid molecules in the sample can comprise conducting one or more measurements and/or experiments on the labeled-amplicons produced from the amplification or repeated amplification reactions conducted on the labeled-nucleic acid molecules. Repeated sampling of the nucleic acid molecules in the

sample can comprise measurements for detecting and/or quantifying a nucleic acid molecule. Repeated sampling of the nucleic acid molecule in the sample can comprise conducting additional experimentation on the nucleic acid molecules in the sample.

**[00282]** In some embodiments, methods, kits, and systems for gene-specific detection of labeled molecules are disclosed. The methods, kits, and systems can be used to increase the detection specificity for one or more genes of interest. A schematic of the method is depicted in FIG. 7. Generally, the method comprises: a) hybridizing at least one target molecule to a solid support; and b) hybridizing a labeled gene-specific oligo to the target molecule to produce a labeled-target molecule.

**[00283]** Further disclosed herein are methods, kits and systems for the absolute quantification of one or more molecules. FIG. 17 depicts a comparison of the quantification of two genes (gene A and gene B). Quantification of the two genes by a standard array readout can provide a relative quantification of genes A and B. In the standard array readout, the genes are amplified and the amplicons are hybridized to an array. The relative amounts of genes A and B can be detected by fluorescence and the intensity (e.g., brightness) of the signal can be used to determine that the quantity of gene B is greater than the quantity of gene A. The digital amplification method disclosed herein can be used to provide an absolute quantification of genes A and B. The absolute quantification method can comprise (a) stochastically labeling two or more genes with a plurality of oligonucleotide tags to produce a stochastically labeled molecule, wherein the plurality of oligonucleotide tags comprises two or more different unique identifier region; (b) amplifying the stochastically labeled molecule to produce one or more stochastically labeled amplicons; and (c) detecting the number of different unique identifier regions associated with each stochastically labeled amplicons, thereby determining the absolute quantity of two or more molecules. As shown in FIG. 17B, detecting the unique identifier regions comprises hybridizing the stochastically labeled amplicons to a solid support (e.g., array). The stochastically labeled amplicons can hybridize to discrete locations on the solid support and the number of different unique identifier regions can be determined by counting the number of discrete locations as detected by fluorescence.

**[00284]** FIG. 19 depicts a schematic of an absolute quantification method of one or more RNA molecules. As shown in Step 1 of FIG. 19, cDNA synthesis of one or more target RNA molecules comprises annealing the oligodT sequence (e.g., target specific region, 1920) of an oligonucleotide tag (1920) to the polyA tail of a mRNA molecule (1910). The oligonucleotide tag (1920) further comprises a unique identifier region (1940) and a universal primer binding site (1950). The unique identifier region (1940) may comprise a

predetermined sequence. Alternatively, the unique identifier region (1940) comprises a random sequence. The resulting cDNA molecule (1960) comprises a copy of the mRNA molecule, the unique identifier region (1940) and the universal primer binding site (1950). As shown in Step 2, the cDNA molecule (1960) can be amplified by nested PCR comprising a first forward primer (1980), a second forward primer (1990) and a reverse primer comprising universal primer (1970) to produce one or more labeled amplicons (e.g., amplicons comprising the unique identifier region). The forward primers (1980, 1990) may be gene-specific primers. The labeled amplicons can be detected by any method known in the art. Absolute quantitation of mRNA molecules can occur by the detection and counting of different unique identifier regions.

**[00285]** FIG. 20 depicts another method for quantifying one or more molecules. The method may comprise (a) reverse transcribing one or more RNA molecules using a plurality of oligonucleotide tags (2030) comprising two or more oligonucleotide tags (2020) comprising a target specific region (2050), a unique identifier region (2060) and a universal primer binding site (2070) to produce one or more stochastically labeled cDNA copies, wherein the stochastically labeled cDNA copies comprise the unique identifier region. The unique identifier region may comprise a random sequence. The method may further comprising amplifying the stochastically labeled cDNA copies to produce one or more stochastically labeled amplicons. Amplifying may comprise PCR and T7 amplification. The stochastically labeled amplicons may comprise the unique identifier region. The method may further comprise detecting the stochastically labeled cDNA copies or stochastically labeled amplicons. Detecting the stochastically labeled molecules can comprise hybridizing the stochastically labeled molecules to one or more digital arrays to determine the number of distinct labels for each gene of interest. Hybridization may require both the presence of the mRNA sequence, most likely a segment on the 3' exon of the gene, and the unique identifier region. The array may comprise 7 million features. The one or more molecules may be in a sample. The sample may comprise 20,000 different mRNA sequences. The method may comprise determining the number of copies of each mRNA present in the sample. The plurality of oligonucleotide tags may comprise 350 or more oligonucleotide tags. In some instances, a subset of the 350 oligonucleotide tags may be applied at a lower concentration to increase the effective dynamic range of measurement.

**[00286]** FIG. 25 depicts another method of absolute quantitation of mRNA molecules. As shown in FIG. 25, the method comprises (a) conducting a reverse transcription reaction with an oligonucleotide tag (2560) to produce a stochastically labeled cDNA molecule (2520),

wherein the stochastically labeled cDNA molecule comprises a cDNA copy of an mRNA molecule (2510), a unique identifier region (2540) and a universal primer binding site (2550); and (b) detecting the stochastically labeled cDNA molecule. The oligonucleotide tag (2560) can serve as a primer for the reverse transcription reaction. The oligonucleotide tag (2560) may comprise a target specific region (2530), unique identifier region (2540) and a universal primer binding site (2550). The method may further comprise absolutely quantifying the mRNA molecules based on the detection of the stochastically labeled cDNA molecules. Detection of the stochastically labeled cDNA molecules may comprise counting the number of different unique identifier regions that are associated with each type of cDNA molecule. The method may further comprise amplifying the stochastically labeled cDNA molecule prior to said detecting to produce one or more stochastically labeled amplicons.

**[00287]** Further disclosed herein are methods, kits and systems for determining the DNA copy number. A general schematic of the method is depicted in FIG. 21. As shown in step 1 of FIG. 21, a genomic DNA (2110) can be fragmented to produce a DNA fragment (2130). Fragmentation of the genomic DNA may occur by any method known in the art. For example, fragmentation may comprise mechanical shearing. Alternatively, fragmentation may comprise digestion of the genomic DNA with one or more restriction nuclease. As shown in Step 2 of FIG. 21, the DNA fragments (2120) can be stochastically labeled with a plurality of oligonucleotide tags (2140) to produce a stochastically labeled molecule (2170). The oligonucleotide tag (2140) may comprise an adapter sequence (2150) and a unique identifier region (2160). The adapter sequence (2150) may enable attachment of the oligonucleotide tag (2140) to the DNA fragments. The adapter sequence (2150) may comprise one or more nucleotides that can anneal to the DNA fragments. Each stochastically labeled molecule (2170) may comprise one or more oligonucleotide tags (2150). The method may further comprise amplifying the stochastically labeled molecules (2170) to produce one or more stochastically labeled amplicons. The method may further comprise removing one or more DNA fragments prior to amplification. Removing the one or more DNA fragments may comprise digesting the DNA fragments with one or more restriction enzymes prior to amplification to prevent the replication of certain fragments. The method may further comprise detecting the stochastically labeled molecules. Detection may comprise hybridization to digital arrays detects the number of distinct unique identifier regions ligated to each DNA fragment.

**[00288]** Further disclosed herein are methods, kits and systems for analyzing one or more RNA molecules. The RNA molecules may be a small RNA molecule. The small RNA

molecule may be a microRNA. FIG. 22 depicts a general method for analyzing a small RNA molecule. As shown in Step 1 of FIG. 22, one or more miRNA molecules (2210) are stochastically labeled with a first plurality of oligonucleotide tags (2230). The oligonucleotide tags (2230) may comprise an adapter sequence (2240) and a unique identifier region (2250). The adapter sequence (2240) may enable attachment of the oligonucleotide tag (2230) to the miRNA molecule (2220) to produce a 3'-stochastically labeled miRNA (2260). As shown in Step 2 of FIG. 22, the method may further comprise stochastically labeling the 3'-stochastically labeled microRNA (2260) with a second plurality of oligonucleotide tags (2270). The second plurality of oligonucleotide tags (2270) may comprise an adapter sequence (2290) and a unique identifier region (2280). The adapter sequence (2290) may enable attachment of the oligonucleotide tag (2270) to the 3'-stochastically labeled miRNA molecule (2260) to produce a 5' and 3'-stochastically labeled miRNA (2295). The method may further comprise reverse transcribing the stochastically labeled miRNA, amplifying the stochastically labeled miRNA, detecting the stochastically labeled miRNA, quantifying the miRNA by detecting the stochastically labeled miRNA, hybridizing the stochastically labeled miRNA to an array, or a combination thereof. The array may be a digital array. The miRNA molecule may comprise any of the miRNA sequences. For example, the miRNA molecule may comprise a sequence disclosed in miRBase 18 <http://www.mirbase.org/>, which was released Nov. 2011 and lists 1921 unique mature human miRNAs. An array of 2 million features can adequately detect 1000 labels ligated to the 1921 miRNAs.

**[00289]** The methods, kits and systems disclosed herein can be used for genetical diagnosis. For example, the methods, kits and systems disclosed herein can be used for single cell pre-implantation genetic diagnosis (PGD). Primary challenges with single-cell genomic DNA amplification assays can be from allele dropout and replication bias. As shown in the computation modeling analysis depicted in FIG. 23A where every molecule has a 0.8 probability of replication, molecules of 1:1 initial copy ratios can easily be distorted to 1:10 or greater just after a few replication cycles. However, when labels are first attached prior to amplification, counting labels to determine copy number is unaffected by replication bias, so long as replication occurs. Aneuploidy determination and large regions of deletion or amplification can be easily and accurately determined by the stochastic labeling method disclosed herein. FIG. 23B depicts a schematic of the general method. As shown in Step 1 of FIG. 23B, the method may comprise fragmenting a genomic DNA (gDNA, 2310) to produce one or more fragmented molecules (2320). Fragmentation of the gDNA (2310) may comprise any method known in the art. For example, fragmentation may comprise conducting a

restriction digest reaction. As shown in Step 2 of FIG. 23B, the fragmented DNA (2320) can be stochastically labeled with a plurality of oligonucleotide tags (2380) to produce one or more stochastically labeled molecules (2330). The stochastically labeled molecule (2330) may comprise one or more oligonucleotide tags (2380). The oligonucleotide tags (2380) may comprise unique identifier sequence (2350) and a universal primer binding site (2340). The stochastically labeled molecule (2380) may be amplified using one or more primers (2360, 2370) that can hybridize to the universal primer binding site (2340) to produce one or more stochastically labeled amplicons. As shown in Step 3 of FIG. 23B, the stochastically labeled molecules (2330) can be detected by a GeneChip detector (2395). The stochastically labeled molecule (2330) can hybridize to a probe (2390) on the GeneChip detector (2395).

**[00290]** The methods, kits, and systems disclosed herein can be used in fetal diagnostics. The method may comprise (a) fragmenting a nucleic acid molecule in a sample to produce one or more nucleic acid fragments; (b) stochastically labeling the one or more nucleic acid fragments with a plurality of oligonucleotide tags comprising a unique identifier region to produce one or more stochastically labeled molecules; and (c) detecting the stochastically labeled molecules by counting the number of unique identifier regions. The method may further comprise diagnosing a fetal genetic disorder based on the detection of the stochastically labeled molecules.

**[00291]** FIG. 24 depicts a general schematic for using the stochastic labeling method in fetal diagnostics. In 100 nanograms of circulating DNA there may be about 10,000 genome equivalents. In the first trimester of maternal plasma, the total concentration of the fetal DNA can be about 10% of the total DNA in the maternal plasma sample. The method, as depicted in FIG. 24, may comprise fragmenting the DNA molecules (2410). Fragmentation may comprise the use of a 4-base restriction enzyme cutter. The fragmented DNA molecules may be stochastically labeled with a plurality of oligonucleotide tags (2420). Stochastic labeling may comprise ligating one or more oligonucleotide tags to the fragmented DNA molecules to produce one or more stochastically labeled molecules. The stochastically labeled molecules may be amplified in a multiplex reaction (2430) to produce one or more stochastically labeled amplicons. The stochastically labeled amplicons may be detected on an array (2440). The array may comprise 5 million features. Diagnosis of a fetal genetic disorder (e.g., trisomy 21) can be based on the detection of the stochastically labeled amplicons (2450, 2460). The 100,000 oligonucleotide tags may be synthesized as described in: Methods for screening factorial chemical libraries, Stephen P. A. Fodor et al, US Patent number: 5541061, issued 7/30/1996.

**[00292]** FIG. 26 depicts a schematic for stochastic labeling of one or more molecules with an inert primer. The method may comprise (a) reverse transcribing an mRNA molecule (2610) with a primer (2620) comprising an oligodU sequence to produce a cDNA copy of the mRNA molecule (2630), wherein the cDNA copy comprises a 3' polyA tail and a 5' oligodT sequence; and (b) stochastically labeling the cDNA copy (2620) with an oligonucleotide tag (2640) comprising a universal primer binding site (2650), unique identifier region (2660) and an oligodU sequence (2670) to produce stochastically labeled cDNA molecule (2680). The method may further comprise a second stochastic labeling step to produce a stochastically labeled cDNA molecule, wherein both ends of the cDNA molecule are stochastically labeled with an oligonucleotide tag. The method may further comprise treating the sample with uracil DNA glycosylase (UDG) to remove the oligodU primer (2620) and the oligonucleotide tags comprising the oligodU sequence. The method may further comprise amplifying the stochastically labeled cDNA molecule to produce one or more stochastically labeled amplicons.

**[00293]** FIG. 27 depicts a schematic for analyzing one or more molecules. The method may comprise (a) reverse transcribing an mRNA molecule (2710) with an oligonucleotide tag (2720) comprising an oligodU sequence (2730), unique identifier region (2740), and a universal primer binding site (2750) to produce a cDNA copy (2760) of the mRNA molecule, wherein the cDNA copy (2760) comprises the unique identifier region (2740) and the universal primer binding site (2750); and (b) amplifying the cDNA copy with a first primer (2790) comprising an oligodU sequence and a second primer (2780) comprising the universal primer sequence to produce stochastically labeled amplicons. The method may comprise treating the molecules with one or more restriction enzymes. The method may further comprise conductig an emulsion PCR reaction on the stochastically labeled molecules.

**[00294]** The methods depicted in FIG. 26-27 may rely on homopolymer tailing. FIG. 28 depicts a method that does not rely on homopolymer tailing. As depicted in FIG. 28, the method may comprise reverse transcribing an mRNA molecule to produce a cDNA copy. Reverse transcription of the mRNA molecule may be carried out on a bead surface. The method may comprise RNase H digestion of the mRNA molecule. The method may comprise stochasticly labeling the cDNA copy with a plurality of oligonucleotide tags to produce one or more stochastically labeled cDNA molecules. The oligonucleotide tag may comprise a secondary structure. The secondary structure may be a hairpin. The oligonucleotide tag may comprise a universal primer binding site, unique identifier region, restriction enzyme recognition site, target specific region, or any combination thereof. The

loop portion of the hairpin oligonucleotide tag may comprise a universal primer binding sequence. The loop portion of the hairpin oligonucleotide tag may comprise a unique identifier region. The loop portion of the hairpin oligonucleotide tag may further comprise a restriction enzyme recognition site. The oligonucleotide tag may be single stranded. The oligonucleotide tag may be double stranded. The method may further comprise amplifying the stochastically labeled cDNA molecule to produce one or more stochastically labeled amplicons. The method may further comprise digesting the stochastically labeled amplicons with a restriction nuclease to produce a digested stochastically labeled amplicon. The method may further comprise ligating one or more primers to the digested stochastically labeled amplicon to produce a primer-stochastically labeled amplicon. The primer may be a sequencing primer. The method may further comprise sequencing the primer-stochastically labeled amplicon. This method may reduce or prevent un-intended incorporation of oligonucleotide tags during PCR amplification. This method may improve sequencing of the stochastically labeled molecules compared to the sequencing of the stochastically labeled molecules from a reaction based on homopolymer tails. This method may reduce or prevent sequencing errors. The oligonucleotide tag may comprise a 3' phosphate. The 3' phosphate can prevent extension of the 3' end during a PCR reaction, thereby reducing or preventing non-specific amplification.

**[00295]** FIG. 29 depicts a linear amplification method. The method may comprise reverse transcribing one or more mRNA molecules by stochastically labeling the one or more RNA molecules with a plurality of oligonucleotide tags to produce one or more cDNA copies of the mRNA molecules, wherein the cDNA copies comprise the oligonucleotide tag. The oligonucleotide tag may comprise a universal primer binding site, unique identifier region and an oligodT sequence. The method may further comprise synthesizing a DNA copy of the mRNA molecule by second strand synthesis. The method may comprise linear amplification of the stochastically labeled cDNA molecule. Linear amplification may comprise amplifying the stochastically labeled cDNA molecule by T7 RNA polymerase, nicking enzyme strand displacement synthesis or RiboSPLA (NuGEN). The method may further comprise attaching one or more sequencing primers to the stochastically labeled molecule. The method may further comprise amplifying the stochastically labeled molecule to produce one or more stochastically labeled amplicons. The method may further comprise sequencing the stochastically labeled amplicons. This method may comprise a low level of initial amplification followed by exponential PCR. This method may be independent of ligation. This method may reduce or prevent artifacts generated by PCR.

**[00296]** FIG. 30 depicts a method of stochastically labeling one or more molecules by strand switching. The method may comprise reverse transcribing a first strand synthesis in the presence of a strand-switch oligonucleotide to produce a stochastically labeled cDNA molecule. The method may further comprise amplifying the stochastically labeled cDNA molecule.

**[00297]** FIG. 31 depicts a method of stochastically labeling one or more molecules by random priming. The method may comprise reverse transcribing an mRNA molecule to produce a stochastically labeled cDNA copy. Reverse transcribing may comprise stochastically labeling one or more molecules with a plurality of oligonucleotide tags, wherein the oligonucleotide tag comprises an oligo dU sequence, a unique identifier sequence and a universal primer sequence. The oligonucleotide tag may further comprise a restriction enzyme recognition site. The method may further comprise removing the mRNA molecules with RNase H. The method may further comprise conducting a second strand synthesis reaction with a second set of oligonucleotide tags. The second set of oligonucleotide tags may comprise a universal primer binding site, a restriction enzyme recognition site, and a unique identifier region. The method may further comprise treating the sample with UDG to remove oligonucleotide tags comprising one or more uracils. The method may further comprise amplifying the stochastically labeled molecules. The method may further comprise attaching one or more adapters to the stochastically labeled molecules. The oligonucleotide tag may comprise any three nucleotides (e.g., C, G, T – no A; C, G, A – no T). The oligonucleotide tag may comprise any two nucleotides (e.g., G, T – no A, C; A, C – no G, T). As shown in FIG. 31, the method may comprise first strand cDNA synthesis with an oligo dT (or dU for subsequent removal with UDG) oligonucleotide tag bearing 12 variable label nucleotides (C/G/T – A was excluded to prevent spurious self-priming to the T/U string). However, instead of TdT tailing to generate the second PCR priming site, an oligonucleotide tag containing a quasi-random string and a PCR sequence is used.

**[00298]** FIG. 43 depicts a schematic of a method for absolute quantitation of one or more molecules directly from one or more cell lysates. As shown in FIG. 43, an intact cell (4310) comprising one or more DNA molecules (4320), RNA molecules (4330), proteins (4340), or a combination thereof is lysed to produce a lysed cell (4350). The one or more DNA molecules (4320), RNA molecules (4330) and/or proteins (4340) can be released from the cell. The quantity of one or more mRNA molecules (4330) can be determined by stochastically labeling the mRNA molecules with a plurality of oligonucleotide tags (4390). The oligonucleotide tag may comprise a target specific region (4360), unique identifier

region (4370) and a universal primer binding site (4380).

**[00299]** In some instances, the target molecule is a DNA molecule. Alternatively, the target molecule is an RNA molecule. In some instances, the methods disclosed herein further comprise reverse transcribing the RNA molecule. The labeled gene-specific oligo can comprise one or more nucleotides. The one or more nucleotides can be a deoxynucleotide. Alternatively, or additionally, the one or more nucleotides are a deoxyribonucleotide. The one or more nucleotides can be a synthetic nucleotide. The labeled gene-specific oligo can comprise at least about 5 nucleotides. Alternatively, the labeled gene-specific oligo comprises at least about 10 nucleotides. Alternatively, the labeled gene-specific oligo comprises at least about 12 nucleotides. The labeled gene-specific oligo can comprise at least about 15 nucleotides. The labeled gene-specific oligo can comprise at least about 17 nucleotides. The labeled gene-specific oligo can comprise at least about 20 nucleotides. In some instances, the labeled gene-specific oligo comprises at least about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 nucleotides.

**[00300]** The labeled gene-specific oligo can comprise a target specific region. The target specific region of the labeled gene-specific oligo can be at least partially complementary to at least a portion of the target molecule. In some instances, the target specific region comprises at least about 5 nucleotides that are complementary to at least a portion of the target molecule. Alternatively, the target specific region comprises at least about 10 nucleotides that are complementary to at least a portion of the target molecule. In other instances, the target specific region comprises at least about 12 nucleotides that are complementary to at least a portion of the target molecule. The target specific region can comprise at least about 15 nucleotides that are complementary to at least a portion of the target molecule. The target specific region can comprise at least about 17 nucleotides that are complementary to at least a portion of the target molecule. The target specific region can comprise at least about 20 nucleotides that are complementary to at least a portion of the target molecule. The target specific region can comprise at least about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 nucleotides that are complementary to at least a portion of the target molecule. The target specific region can comprise a sequence that is at least about 60% complementary to at least a portion of the target molecule. Alternatively, the target specific region comprises a sequence that is at least about 70% complementary to at least a portion of the target molecule. The target specific region can comprise a sequence that is at least about 80% complementary to at least a portion of the target molecule. The target specific region can comprise a sequence that is at least about 85% complementary to at least a portion of the

target molecule. The target specific region can comprise a sequence that is at least about 90% complementary to at least a portion of the target molecule. The target specific region can comprise a sequence that is at least about 95% complementary to at least a portion of the target molecule. The target specific region can comprise a sequence that is at least about 97% complementary to at least a portion of the target molecule. The target specific region can comprise a sequence that is at least about 98% complementary to at least a portion of the target molecule.

**[00301]** The labeled gene-specific oligo can comprise any label disclosed herein. In some instances, the label is a fluorophore. Alternatively, the label is a cyanine dye (e.g., Cy3, Cy5).

**[00302]** The solid support can be any solid support disclosed herein. In some instances, the solid support is a detector array. The detector array can comprise a plurality of probes. The target molecule can be hybridized to one or more probes of the plurality of probes on the detector array.

**[00303]** The method can further comprise amplifying the target molecule prior to hybridization to the solid support. The methods disclosed herein can further comprise sequencing the target molecules hybridized to the solid support. The methods disclosed herein can be used to prevent false-positive detection of PCR amplified DNAs that do not contain the gene of interest.

**[00304]** The method can further comprise detecting the labeled-target molecules. Methods to detect the labeled-target molecule can comprise any of the detection methods and instruments disclosed herein. In some instances, detecting the labeled-target molecule comprises detecting the label. Detecting the labeled-target molecule can comprise a fluorometer. Alternatively, detecting the labeled-target molecule comprises a luminometer. In other instances, detecting the labeled-target molecule comprises a plate reader.

**[00305]** Further disclosed herein are methods, kits, and systems for capturing and/or enriching a population of target molecules. FIG. 8 shows a schematic of the method. Generally, the method comprises: a) stochastically labeling one or more nucleic acid molecules in a sample to produce a stochastically labeled molecule; and b) capturing one or more stochastically labeled molecules to produce a captured molecule, wherein the captured molecule comprise a target molecule.

**[00306]** Capturing the stochastically labeled molecule can comprise the use of one or more gene-specific oligos. The gene-specific oligos can attach to a specific stochastically labeled molecule to produce an oligo linked molecule. In some instances, the methods disclosed herein further comprise isolating the oligo linked molecule from the sample. The gene-

specific oligo can comprise a label or tag. The label or tag can enable isolation of the oligo linked molecule.

**[00307]** Alternatively, capturing the stochastically labeled molecule can comprise contacting the sample comprising the stochastically labeled molecules with a solid support. In some instances, the stochastically labeled molecule comprising the target molecule hybridizes to the solid support, thereby capturing the stochastically labeled molecule. Alternatively, the stochastically labeled molecule hybridized to the solid support does not comprise the target molecule and capturing the stochastically labeled molecule comprises collecting any unbound stochastically labeled molecules (e.g., stochastically labeled molecules that are not hybridized to the solid support). The solid support can be any of the solid supports disclosed herein. In some instances, the solid support is an array. In other instances, the solid support is a bead. The bead can be a magnetic bead. In some instances, capturing the stochastically labeled molecule comprises the use of a magnet.

**[00308]** The method can further comprise amplification of the stochastically labeled molecule and/or captured molecule. Amplification of the stochastically labeled molecule and/or captured molecule can comprise any of the amplification methods disclosed herein. In some instances, amplification of the stochastically labeled molecule and/or captured molecule comprises PCR.

**[00309]** The methods disclosed herein can further comprise sequencing of the captured molecule. Sequencing can comprise any of the sequencing methods disclosed herein. In some instances, the captured molecules are directly sequenced on the solid support.

**[00310]** Further disclosed herein are methods, kits, and systems for digital detection and/or quantification of a nucleic acid molecule. Generally, the methods, kits, and systems comprise (a) stochastically labeling a nucleic acid molecule with a plurality of oligonucleotide tags to produce a stochastically labeled-nucleic acid molecule; and (b) detecting and/or quantifying the stochastically labeled-nucleic acid molecule. The nucleic acid molecule can be a DNA molecule. The nucleic acid molecule can be from a cell. Alternatively, the nucleic acid molecule is a cell-free molecule. The nucleic acid molecule can be derived from a subject. Alternatively, the nucleic acid molecule can be derived from a foreign subject. The foreign subject can be a pathogen (e.g., virus, bacteria, fungus).

**[00311]** The method can further comprise amplifying the stochastically labeled-nucleic acid molecules to produce stochastically-labeled nucleic acid molecule amplicons. The stochastically labeled-nucleic acid molecules or any products thereof (e.g., stochastically-labeled nucleic acid molecule amplicons) can be repeatedly amplified.

**[00312]** In some instances, the method further comprises attaching one or more detectable labels to the stochastically labeled-nucleic acid molecules or products thereof. In some instances, at least one detectable label is attached to the stochastically labeled-nucleic acid molecules or products thereof. Alternatively, at least two detectable labels are attached to the stochastically labeled-nucleic acid molecules or products thereof. The detectable label can be biotin. Alternatively, the detectable label is a fluorescent dye. The fluorescent dye can be a Cy<sup>TM</sup> dye or a TYE 563 dye. The Cy<sup>TM</sup> dye can be Cy3.

**[00313]** The method can further comprise hybridization of the stochastically labeled-nucleic acid molecules or any products thereof to a solid support. The solid support can be a bead. Alternatively, the solid support is an array.

**[00314]** The method can further comprise conducting a sequencing reaction to determine the sequence of at least a portion of the stochastically labeled-nucleic acid molecule or product thereof. In some instances, at least a portion of the oligonucleotide tag of the stochastically labeled-nucleic acid molecule or product thereof is sequences. For example, at least a portion of the unique identifier region of the oligonucleotide tag is sequenced. In another example, at least a portion of the target specific region of the oligonucleotide tag is sequenced. Alternatively, or additionally, at least a portion of the nucleic acid molecule of the stochastically labeled-nucleic acid molecule is sequenced.

**[00315]** Detection and/or quantification of the stochastically labeled-nucleic acid molecules can comprise detection and/or quantification of the stochastically-labeled cDNA copies and/or the stochastically-labeled nucleic acid molecule amplicons. Detection and/or quantification of the stochastically labeled-nucleic acid molecules can further comprise detection of one or more detectable labels attached to the stochastically labeled-nucleic acid molecules or products thereof. Detection and/or quantification of the stochastically labeled-nucleic acid molecules or products thereof can comprise any of the detection and/or quantification methods disclosed herein. For example, a fluorescence reader can be used to detect and/or quantify the stochastically labeled-nucleic acid molecules or products thereof. Alternatively, a microarray reader can be used to detect and/or quantify the stochastically labeled-nucleic acid molecules or products thereof.

**[00316]** Further disclosed herein are methods, kits, and systems for digital detection and/or digital quantification of viral molecules. Generally, the methods, kits, and systems comprise (a) stochastically labeling one or more viral molecules with a plurality of oligonucleotide tags to produce a stochastically labeled-viral molecule; and (b) detecting and/or quantifying the stochastically labeled-viral molecule. In some instances, the viral molecules are nucleic acid

molecules. The nucleic acid molecules can be DNA or RNA.

**[00317]** The method can further comprise conducting a reverse transcription reaction to produce a stochastically-labeled cDNA copy of the stochastically-labeled viral molecule (e.g., stochastically-labeled viral RNA molecule). The stochastically-labeled viral molecule can be repeatedly reverse transcribed to produce multiple stochastically-labeled cDNA copies of the stochastically-labeled viral molecule. The methods can further comprise amplifying the stochastically labeled-viral molecules or any products thereof (e.g., stochastically-labeled cDNA copy) to produce stochastically-labeled viral amplicons. The stochastically labeled-viral molecules can be repeatedly amplified. Alternatively, the products of the stochastically-labeled viral molecules can be repeatedly amplified. In some instances, the products of the stochastically-labeled viral molecules are the stochastically-labeled cDNA copies of the stochastically-labeled viral molecule. Alternatively, the products of the stochastically-labeled viral molecules are the stochastically-labeled viral amplicons.

**[00318]** In some instances, the method further comprises attaching one or more detectable labels to the stochastically labeled-viral molecules or products thereof. In some instances, at least one detectable label is attached to the stochastically labeled-viral molecules or products thereof. Alternatively, at least two detectable labels are attached to the stochastically labeled-viral molecules or products thereof. The detectable label can be biotin. Alternatively, the detectable label is a fluorescent dye. The fluorescent dye can be a Cy<sup>TM</sup> dye or a TYE 563 dye. The Cy<sup>TM</sup> dye can be Cy3.

**[00319]** The method can further comprise hybridization of the stochastically labeled-viral molecules or any products thereof to a solid support. The solid support can be a bead. Alternatively, the solid support is an array.

**[00320]** The method can further comprise conducting a sequencing reaction to determine the sequence of at least a portion of the stochastically labeled-viral molecule or product thereof. In some instances, at least a portion of the oligonucleotide tag of the stochastically labeled-viral molecule or product thereof is sequences. For example, at least a portion of the unique identifier region of the oligonucleotide tag is sequenced. In another example, at least a portion of the target specific region of the oligonucleotide tag is sequenced. Alternatively, or additionally, at least a portion of the viral molecule of the stochastically labeled-viral molecule is sequenced.

**[00321]** Detection and/or quantification of the stochastically labeled-viral molecules can comprise detection and/or quantification of the stochastically-labeled cDNA copies and/or the stochastically-labeled viral amplicons. Detection and/or quantification of the stochastically

labeled-viral molecules can further comprise detection of one or more detectable labels attached to the stochastically labeled-viral molecules or products thereof. Detection and/or quantification of the stochastically labeled-viral molecules or products thereof can comprise any of the detection and/or quantification methods disclosed herein. For example, a fluorescence reader can be used to detect and/or quantify the stochastically labeled-viral molecules or products thereof. Alternatively, a microarray reader can be used to detect and/or quantify the stochastically labeled-viral molecules or products thereof.

**[00322]** In some instances, digital detection and/or digital quantification of the viral molecules can be used to determine the viral load in a subject suffering from a viral infection. Alternatively, digital detection and/or digital quantification of the viral molecules can be used in the diagnosis and/or prognosis of a viral infection. In some instances, digital detection and/or digital quantification of the viral molecules can be used in monitoring an antiviral therapeutic regimen.

**[00323]** Further disclosed herein are methods, kits, and systems for digital detection and/or quantification of a biomarker. The methods, kits, and systems can be used to quantify a biomarker. Generally, the methods, kits, and systems comprise (a) stochastically labeling a biomarker with a plurality of oligonucleotide tags to produce a stochastically labeled-biomarker; and (b) detecting and/or quantifying the stochastically labeled-biomarker. The biomarker can be a cancer biomarker. The biomarker can be a nucleic acid molecule or a protein. The nucleic acid molecule can be a DNA molecule. Alternatively, the nucleic acid molecule can be a RNA molecule. The biomarker can be derived from a subject. Alternatively, the biomarker can be derived from a foreign subject. The foreign subject can be a pathogen (e.g., virus, bacteria, fungus).

**[00324]** The method can further comprise conducting a reverse transcription reaction to produce a stochastically-labeled cDNA copy of the stochastically-labeled biomarker (e.g., stochastically-labeled biomarker RNA molecule). The stochastically-labeled biomarker can be repeatedly reverse transcribed to produce multiple stochastically-labeled cDNA copies of the stochastically-labeled biomarker. The methods can further comprise amplifying the stochastically labeled-biomarkers or any products thereof (e.g., stochastically-labeled cDNA copy) to produce stochastically-labeled biomarker amplicons. The stochastically labeled-biomarkers can be repeatedly amplified. Alternatively, the products of the stochastically-labeled biomarkers can be repeatedly amplified. In some instances, the products of the stochastically-labeled biomarkers are the stochastically-labeled cDNA copies of the stochastically-labeled biomarker. Alternatively, the products of the stochastically-labeled

biomarkers are the stochastically-labeled biomarker amplicons.

**[00325]** In some instances, the method further comprises attaching one or more detectable labels to the stochastically labeled-biomarkers or products thereof. In some instances, at least one detectable label is attached to the stochastically labeled-biomarkers or products thereof. Alternatively, at least two detectable labels are attached to the stochastically labeled-biomarkers or products thereof. The detectable label can be biotin. Alternatively, the detectable label is a fluorescent dye. The fluorescent dye can be a Cy<sup>TM</sup> dye or a TYE 563 dye. The Cy<sup>TM</sup> dye can be Cy3.

**[00326]** The method can further comprise hybridization of the stochastically labeled-biomarkers or any products thereof to a solid support. The solid support can be a bead. Alternatively, the solid support is an array.

**[00327]** The method can further comprise conducting a sequencing reaction to determine the sequence of at least a portion of the stochastically labeled-biomarker or product thereof. In some instances, at least a portion of the oligonucleotide tag of the stochastically labeled-biomarker or product thereof is sequences. For example, at least a portion of the unique identifier region of the oligonucleotide tag is sequenced. In another example, at least a portion of the target specific region of the oligonucleotide tag is sequenced. Alternatively, or additionally, at least a portion of the biomarker of the stochastically labeled-biomarker is sequenced.

**[00328]** Detection and/or quantification of the stochastically labeled-biomarkers can comprise detection and/or quantification of the stochastically-labeled cDNA copies and/or the stochastically-labeled biomarker amplicons. Detection and/or quantification of the stochastically labeled-biomarkers can further comprise detection of one or more detectable labels attached to the stochastically labeled-biomarkers or products thereof. Detection and/or quantification of the stochastically labeled-biomarkers or products thereof can comprise any of the detection and/or quantification methods disclosed herein. For example, a fluorescence reader can be used to detect and/or quantify the stochastically labeled-biomarkers or products thereof. Alternatively, a microarray reader can be used to detect and/or quantify the stochastically labeled-biomarkers or products thereof.

**[00329]** In some instances, digital detection and/or digital quantification of the biomarkers can be used to diagnose or prognose a condition in a subject in need thereof. In some instances, digital detection and/or digital quantification of the biomarkers can be used to monitor a therapeutic regimen.

**[00330]** The condition can be a cancer. The cancer can be a sarcoma, carcinoma, leukemia,

or lymphoma.

**[00331]** Alternatively, the condition is a pathogenic infection. The pathogenic infection can be a bacterial or viral infection.

**[00332]** Further disclosed herein are methods, kits and systems for counting or determining a number of nucleic acid molecules in a sample. The method may comprise: (a) providing a plurality of oligonucleotide tags wherein a oligonucleotide tag comprises a unique identifier sequence, a target sequence, and an optional PCR primer sequence; (b) combining a sample comprising nucleic acid molecules with the plurality of labeled primers to form a labeled nucleic acid molecule, wherein each target nucleic acid molecule is capable of attaching to a oligonucleotide tag with a unique identifier sequence; and (c) detecting (i) the nucleic acid molecule, a complement of the nucleic acid molecule, a reverse complement of the nucleic acid molecule, or a portion thereof, and (ii) the oligonucleotide tag, a complement of the oligonucleotide tag, a reverse complement of the oligonucleotide tag, or a portion thereof to determine the count or number of different labeled nucleic acid molecules, thereby counting or determining a number of nucleic acid molecules in the sample. The method may comprise counting or determining a number of 10 or more different nucleic acid molecules. The method may comprise counting or determining a number of 20 or more different nucleic acid molecules. The different nucleic acid molecules may differ by 1 or more nucleotides or base pairs. The different nucleic acids may be counted simultaneously. Alternatively, the different nucleic acid molecules may be counted sequentially.

**[00333]** The method of counting or determining a number of nucleic acid molecules in a sample may comprise: (a) providing a plurality of oligonucleotide tags wherein a oligonucleotide tag comprises a unique identifier sequence, a target sequence, and an optional PCR primer sequence; (b) combining a sample comprising nucleic acid molecules with the plurality of labeled primers to form a labeled nucleic acid molecule, wherein the attachment of the nucleic acid molecule to the oligonucleotide tag forms a unique molecule-tag junction; and (c) detecting the unique molecule-tag junction, a complement of the unique molecule-tag junction, a reverse complement of the unique molecule-tag junction, or a portion thereof to determine the count or number of different labeled nucleic acid molecules, thereby counting or determining a number of nucleic acid molecules in the sample. The method may comprise counting or determining a number of 10 or more different nucleic acid molecules. The method may comprise counting or determining a number of 20 or more different nucleic acid molecules. The different nucleic acid molecules may differ by 1 or more nucleotides or base pairs. The different nucleic acids may be counted simultaneously. Alternatively, the different

nucleic acid molecules may be counted sequentially.

**[00334]** The method of counting or determining a number of nucleic acid molecules in a sample may comprise: (a) providing a plurality of oligonucleotide tags, wherein the oligonucleotide tag comprises a target-specific sequence, a unique identifier sequence comprising a ribonucleic acid, and an optional PCR primer sequence; (b) combining a sample comprising nucleic acid molecules with the plurality of oligonucleotide tags to form a labeled nucleic acid molecule, wherein a target nucleic acid molecule is capable of attaching to oligonucleotide tags with different unique identifier sequences; (c) synthesizing a copy of the labeled nucleic acid molecule, wherein the copy of the labeled nucleic acid molecule comprises a copy of the nucleic acid molecule and a copy of the oligonucleotide tag and the ribonucleic acid of the unique identifier sequence comprises replaced with a deoxyribonucleic acid; and (d) detecting the copy of the labeled nucleic acid molecule, a complement of the copy of the labeled nucleic acid molecule, a reverse complement of the copy of the labeled nucleic acid molecule, or a portion thereof to determine a count of the copy of the labeled nucleic acid molecule, thereby counting or determining a number of nucleic acid molecules in the sample. The method may comprise counting or determining a number of 10 or more different nucleic acid molecules. The method may comprise counting or determining a number of 20 or more different nucleic acid molecules. The different nucleic acid molecules may differ by 1 or more nucleotides or base pairs. The different nucleic acids may be counted simultaneously. Alternatively, the different nucleic acid molecules may be counted sequentially.

**[00335]** The method of counting or determining a number of RNA molecules in a sample may comprise: (a) combining a sample comprising RNA molecules with a plurality of oligonucleotide tags, wherein the oligonucleotide tag comprises an RNA-specific sequence, a unique identifier sequence, and an optional PCR primer sequence; (b) synthesizing a copy of an RNA molecule by attaching a oligonucleotide tag to the RNA molecule to form a labeled DNA molecule, wherein each RNA molecule is capable of attaching to oligonucleotide tags with different unique identifier sequences and each labeled DNA molecule comprises a copy of the RNA molecule and a copy of the oligonucleotide tag; and (c) detecting the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled DNA molecule, or a portion thereof to determine a count of the labeled DNA molecule, thereby counting or determining a number of RNA molecules in the sample.

**[00336]** The method of counting or determining a number of RNA molecules in a sample may comprise: (a) providing a plurality of oligonucleotide tags, wherein the oligonucleotide

tag comprises an RNA-specific sequence, a unique identifier sequence comprising a ribonucleic acid, and an optional PCR primer sequence; (b) combining a sample comprising RNA molecules with the plurality of oligonucleotide tags to form a labeled RNA molecule, wherein a target RNA molecule is capable of attaching to oligonucleotide tags with different unique identifier sequences; (c) synthesizing a copy of the labeled RNA molecule to form a labeled DNA molecule, wherein the labeled DNA molecule comprises a copy of the RNA molecule and a copy of the oligonucleotide tag and the ribonucleic acid of the unique identifier sequence comprises replaced with a deoxyribonucleic acid; and (d) detecting the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled DNA molecule, or a portion thereof to determine a count of the labeled DNA molecule, thereby counting or determining a number of RNA molecules in the sample.

**[00337]** The method of counting or determining a number of RNA molecules in a sample may comprise: (a) combining a sample comprising RNA molecules with a plurality of oligonucleotide tags to form a labeled RNA molecule, wherein each target RNA molecule is capable of attaching to a different label; (b) optionally attaching a second oligonucleotide tag to the labeled RNA molecule to form a dual-labeled RNA molecule; (c) synthesizing a copy of the labeled RNA molecule or dual-labeled RNA molecule to form a labeled DNA molecule or dual-labeled DNA molecule, wherein the labeled DNA molecule and the dual-labeled DNA molecule comprise a copy of the oligonucleotide tag and a copy of the RNA molecule; and (d) detecting the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled-DNA molecule, the dual-labeled DNA molecule, a complement of the dual-labeled DNA molecule, a reverse complement of the dual-labeled DNA molecule, or a portion thereof to count or determine the number of different labeled DNA molecules or different dual-labeled DNA molecules, thereby counting or determining a number of RNA molecules in the sample.

**[00338]** The method of counting or determining a number of RNA molecules in a sample may comprise: (a) combining a sample comprising RNA molecules with a plurality of labels to form a labeled RNA molecule, wherein each target RNA molecule is capable of attaching to a different label; (b) optionally attaching a second label to the labeled RNA molecule to form a dual-labeled RNA molecule; and (c) detecting the labeled RNA molecule, a complement of the labeled RNA molecule, a reverse complement of the labeled-RNA molecule, the dual-labeled RNA molecule, a complement of the dual-labeled RNA molecule, a reverse complement of the dual-labeled RNA molecule, or a portion thereof to count or determine the number of different labeled RNA molecules or different dual-labeled RNA

molecules, thereby counting or determining a number of RNA molecules in the sample.

**[00339]** The method of counting or determining a number of mRNA molecules in a sample may comprise: (a) providing a plurality of oligonucleotide tags, wherein the oligonucleotide tag comprises a target-specific sequence, a unique identifier sequence, and an optional PCR primer sequence; (b) combining a sample comprising mRNA molecules with the plurality of oligonucleotide tags to form a labeled mRNA molecule, wherein each target mRNA molecule is capable of attaching to a different oligonucleotide tag; (b) synthesizing a copy of the labeled mRNA molecule to form a labeled DNA molecule, wherein the labeled DNA molecule comprises a copy of the mRNA molecule and a oligonucleotide tag or a copy of the oligonucleotide tag; and (c) detecting the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled DNA molecule, or a portion thereof to determine a count of different labeled DNA molecules, thereby counting or determining a number of mRNA molecules in the sample.

**[00340]** In one aspect, polyadenylated RNA from a single cell is analyzed by the methods disclosed herein. After cell lysis the polyA RNA may be enriched by capture on a solid support, such as a bead, having oligo dT attached or the amplification can be performed on the lysate. A labeled-cDNA copy of the RNA is made by hybridizing a primer that has an oligo dT region and a label-tag region. The label-tag region being 5' of the oligo dT region. Preferably there is an amplification sequence that is 5' of the label-tag region so that the label-tag region, which is variable between primers, is between a 5' common amplification primer sequence and a 3' oligo dT region. Second strand cDNA is then synthesized using standard methods, for example use of RNaseH and DNA polymerase. The resulting dsDNA can then be linearly amplified depending on the amplification primer sequence. For example, if the amplification primer sequence is a T7 RNA polymerase promoter sequence, antisense RNA can be generated by IVT using T7 RNA pol. If the amplification prime sequence includes a site for a nicking enzyme (e.g. Nt. BspQI), nicking enzyme strand displacement can be used to generate DNA copies of the RNA targets. The copies can then be modified to include sequencing primers at one or both ends and the products can be sequenced. Sequence information is collected for the tag and enough of the adjacent sequence to provide an identification of the target.

**[00341]** In some instances, the oligonucleotide tag comprises a ribonucleic acid. The oligonucleotide tag may comprise a ribonucleic acid that is uracil. The oligonucleotide tag may comprise a ribonucleic acid that is cytosine. The oligonucleotide tag may comprise a ribonucleic acid that is adenine. The oligonucleotide tag may comprise a ribonucleic acid that

is guanosine.

**[00342]** The unique identifier sequence may comprise a predetermined sequence. The unique identifier sequence may comprise a random sequence.

**[00343]** The target-specific sequence of the oligonucleotide tag may be specific for a plurality of targets. In some aspects, the target-specific sequence of the oligonucleotide tag comprises an oligo dT sequence. In some aspects, the target-specific sequence of the oligonucleotide tag may comprise an oligo dU sequence. In some instances, the target-specific sequence does not comprise an oligo dT nor oligo dU sequence.

**[00344]** The copy of the labeled DNA molecule may be synthesized by a reverse transcriptase enzyme. The reverse transcriptase enzyme may be selected from a retroviral reverse transcriptase, a phage DNA polymerase, or a DNA polymerase.

**[00345]** The method may further comprise synthesizing a copy of the labeled nucleic acid molecule to replace a ribonucleic acid with a deoxyribonucleic acid.

**[00346]** In some aspects, the detecting step comprises detecting the copy of the labeled nucleic acid molecule, a complement of the copy of the labeled nucleic acid molecule, a reverse complement of the copy of the labeled nucleic acid molecule, or a portion thereof. In some aspects, the detection step may comprise hybridization of the nucleic acid molecule portion of the labeled nucleic acid molecule, a complement of the nucleic acid molecule portion of the labeled nucleic acid molecule, a reverse complement of the nucleic acid molecule portion of the labeled nucleic acid molecule, the oligonucleotide tag of the labeled nucleic acid molecule, a complement of the oligonucleotide tag of the labeled nucleic acid molecule, a reverse complement of the oligonucleotide tag of the labeled nucleic acid molecule, a portion thereof, or any combination thereof to a solid support. In some aspects, the detection step may comprise hybridization of the nucleic acid molecule portion of the copy of the labeled nucleic acid molecule, the oligonucleotide tag portion of the copy of the labeled nucleic acid molecule, a complement thereof, a reverse complement thereof, a portion thereof, or any combination thereof to a solid support.

**[00347]** In some aspects, the detecting step comprises detecting the copy of the oligonucleotide tag, a complement of the copy of the oligonucleotide tag, a reverse complement of the copy of the oligonucleotide tag, or a portion thereof.

**[00348]** The detection step may comprise hybridization of the unique molecule-tag junction, a complement of the unique molecule-tag junction, a reverse complement of the unique molecule-tag junction, or a portion thereof to a solid support. The detection step may comprise hybridization of a copy of the unique molecule-tag junction, a complement of the

copy of the unique molecule-tag junction, a reverse complement of the copy of the unique molecule-tag junction, or a portion thereof to a solid support.

**[00349]** In some aspects, the solid support comprises an array. The array may comprise probes attached to the surface. The array may further comprise a probe feature for each possible labeled nucleic acid molecule combination. In another aspect, the solid support may comprise a bead.

**[00350]** In some aspects, the detection step comprises sequencing of (i) the nucleic acid molecule portion of the labeled nucleic acid molecule, a complement thereof, a reverse complement thereof, or a portion thereof, and (ii) the oligonucleotide tag portion of the labeled nucleic acid molecule, a complement thereof, a reverse complement thereof, or a portion thereof. In some aspects, the detection step comprises sequencing of (i) the nucleic acid molecule portion of the copy of the labeled nucleic acid molecule, a complement thereof, a reverse complement thereof, or a portion thereof, and (ii) the oligonucleotide tag portion of the copy of the labeled nucleic acid molecule, a complement thereof, a reverse complement thereof, or a portion thereof.

**[00351]** In some aspects, the detection step may comprise sequencing the unique oligonucleotide tag-DNA junction, a complement of the unique oligonucleotide tag-DNA junction, a reverse complement of the unique oligonucleotide tag-DNA junction, or a portion thereof. In some aspects, the detection step may comprise sequencing the copy of the unique oligonucleotide tag-DNA junction, a complement of the copy of the unique oligonucleotide tag-DNA junction, a reverse complement of the copy of the unique oligonucleotide tag-DNA junction, or a portion thereof.

**[00352]** In another aspect, the labeled nucleic acid molecule is amplified. In another aspect, the copy of the labeled nucleic acid sequence is amplified. The amplification of the labeled nucleic acid molecule or the copy of the labeled nucleic acid molecule may comprise a PCR-based method. The PCR-based method may comprise qPCR. The PCR-based method may comprise RT-PCR. The PCR-based method may comprise emulsion PCR. The amplification of the nucleic acid molecule-labeled conjugate may comprise a non-PCR-based method. The non-PCR-based method may comprise multiple displacement amplification. The non-PCR-based method may comprise random priming by a strand displacement polymerase.

**[00353]** In another aspect, the sample is from at least one single cell. Alternatively, the sample is from a plurality of cells. The sample may be from less than about 100 cells.

**[00354]** In some aspects, the nucleic acid molecule is a DNA molecule. In another aspect, the nucleic acid molecule is an RNA molecule. The nucleic acid molecule may be an mRNA

molecule. The nucleic acid molecule may be a noncoding RNA molecule. The noncoding RNA molecule may be a small noncoding RNA molecule. The noncoding RNA molecule may be a long noncoding RNA molecule. The noncoding RNA molecule may be a microRNA molecule. In some aspects, the oligonucleotide tag is attached to the nucleic acid molecule by ligation. In another aspect, the oligonucleotide tag is attached to the nucleic acid molecule by hybridization.

**[00355]** In another aspect is a method of counting or determining a number of DNA molecules in a sample comprising: (a) providing a plurality of oligonucleotide tags wherein a oligonucleotide tag comprises a unique identifier sequence, a target sequence, and an optional PCR primer sequence; (b) combining a sample comprising DNA molecules with the plurality of labeled primers to form a labeled DNA molecule, wherein the labeled DNA molecule comprises a DNA molecule and a oligonucleotide tag and each target DNA molecule is capable of attaching to a different oligonucleotide tag; and (c) detecting (i) the DNA molecule, a complement of the DNA molecule, a reverse complement of the DNA molecule, or a portion thereof, and (ii) the oligonucleotide tag, a complement of the oligonucleotide tag, a reverse complement of the oligonucleotide tag, or a portion thereof to determine the count or number of different labeled DNA molecules, thereby counting or determining a number of DNA molecules in the sample.

**[00356]** In another aspect is a method of counting or determining a number of DNA molecules in a sample comprising: (a) providing a plurality of oligonucleotide tags wherein a oligonucleotide tag comprises a unique identifier sequence, a target sequence, and an optional PCR primer sequence; (b) combining a sample comprising DNA molecules with the plurality of labeled primers to form a labeled DNA molecule, wherein the attachment of the DNA molecule to the oligonucleotide tag forms a unique molecule-tag junction; and (c) detecting the unique molecule-tag junction, a complement of the unique molecule-tag junction, a reverse complement of the unique molecule-tag junction, or a portion thereof to determine the count or number of different labeled DNA molecules, thereby counting or determining a number of DNA molecules in the sample.

**[00357]** In another aspect is a method for determining a copy number of a target DNA in a sample comprising: (a) providing a plurality of adaptors, wherein the adaptors comprise a unique identifier sequence and each adaptor is capable of attaching to a plurality of different DNA molecules; (b) fragmenting a sample comprising genomic DNA to produce a sample comprising DNA fragments; (c) combining a plurality of adaptors with the sample comprising DNA fragments to form an adaptor-DNA fragment conjugate, wherein

substantially all of the DNA fragments are randomly attached to an adaptor with a unique identifier sequence; and (d) detecting the adaptor, a complement of the adaptor, a reverse complement of the adaptor or a portion thereof to determine the number of different adaptor-DNA fragment conjugates, thereby determining a copy number of a target DNA.

**[00358]** In another aspect is a method determining a copy number of a target DNA molecule in a sample comprising: (a) providing a plurality of adaptors, wherein the adaptors comprise a unique identifier sequence and the adaptors are capable of attaching to a plurality of different DNA molecules; (b) fragmenting a sample comprising genomic DNA to produce a sample comprising DNA fragments; (c) attaching adaptors to the DNA fragments, wherein substantially all of the DNA fragments capable of being randomly attached to an adaptor with a unique identifier sequence and the attachment of the adaptor to the DNA fragment forms a unique adaptor-DNA junction; and (d) detecting the unique adaptor-DNA junction, a complement of the unique adaptor-DNA junction, a reverse complement of the unique adaptor-DNA junction, or a portion thereof to determine the count or number of different unique adaptor-DNA junctions, thereby determining a copy number of a target DNA.

**[00359]** In some aspects, the adaptor comprises a ribonucleic acid. In some aspects, the ribonucleic acid is uracil. In some aspects, the ribonucleic acid is cytosine. In some aspects, the ribonucleic acid is adenine. In some aspects, the ribonucleic acid is guanine.

**[00360]** In some aspects, the method further comprises synthesizing a copy of the adaptor-DNA fragment conjugate to replace a ribonucleic acid sequence in the adaptor with a deoxyribonucleic acid sequence.

**[00361]** In some aspects, the detecting step comprises detecting the copy of the unique adaptor-DNA junction, a complement of the copy of the unique adaptor-DNA junction, a reverse complement of the copy of the unique adaptor-DNA junction, or a portion thereof. In some aspects, the detecting step comprises detecting the copy of the adaptor, a complement of the copy of the adaptor, a reverse complement of the copy of the adaptor, or a portion thereof.

**[00362]** In some aspects, the detection step comprises hybridization of the unique adaptor-DNA junction, a complement of the unique adaptor-DNA junction, a reverse complement of the unique adaptor-DNA junction, or a portion thereof to a solid support. In another aspect, the detection step comprises hybridization of the copy of the unique adaptor-DNA junction, a complement of the copy of the unique adaptor-DNA junction, a reverse complement of the copy of the unique adaptor-DNA junction, or a portion thereof to a solid support.

**[00363]** In some aspects, solid support comprises an array. In some aspects, the array

comprises probes attached to the surface. In some aspects, the array comprises a probe feature for each unique-adaptor DNA junction. In some aspects, the array comprises a probe feature for each copy of the unique-adaptor DNA junction. In another aspect, the solid support comprises a bead. In some aspects, the detection step comprises sequencing the unique adaptor-DNA junction, a complement of the unique adaptor-DNA junction, a reverse complement of the unique adaptor-DNA junction, or a portion thereof. In some aspects, the detection step comprises sequencing the copy of the unique adaptor-DNA junction, a complement of the copy of the unique adaptor-DNA junction, a reverse complement of the copy of the unique adaptor-DNA junction, or a portion thereof.

**[00364]** In some aspects, the detection step comprises sequencing the copy of the adaptor, a complement of the copy of the adaptor, a reverse complement of the copy of the adaptor, or a portion thereof. In some aspects, the adaptor-DNA fragment conjugate is amplified.

**[00365]** In some aspects is a method of determining the presence or absence of genetic abnormalities comprising: (a) providing a plurality of oligonucleotide tags wherein a oligonucleotide tag comprises a unique identifier sequence, a target sequence, and an optional PCR primer sequence; (b) combining a sample comprising genomic DNA with the plurality of labeled primers to form a genomic DNA-oligonucleotide tag conjugate, wherein each genomic DNA is capable of attaching to a oligonucleotide tag with a unique identifier sequence; and (c) detecting the genomic DNA-oligonucleotide tag conjugate, a complement of genomic DNA-oligonucleotide tag conjugate, a reverse complement of the genomic DNA-oligonucleotide tag conjugate, or a portion thereof to count or determine a number of different genomic DNA-oligonucleotide tag conjugates, thereby determining the presence or absence of genetic abnormalities.

**[00366]** In some aspects, the detecting step comprises detecting the genomic DNA, a complement of the genomic DNA, a reverse complement of the genomic DNA, or a portion thereof. In some aspects, the detecting step comprises detecting the oligonucleotide tag, a complement of the oligonucleotide tag, a reverse complement of the oligonucleotide tag, or a portion thereof.

**[00367]** In some aspects, the genetic abnormality comprises an aneuploidy. The aneuploidy may be monosomy. The monosomy may be monosomy of the sex chromosome. The aneuploidy may be trisomy. The trisomy may be trisomy 21. The trisomy may be trisomy 18. The trisomy may be trisomy 13. The aneuploidy may be tetrasomy. The aneuploidy may be pentasomy. In some aspects, the method further comprises diagnosing a genetic abnormality. In some aspects, the method may further comprise diagnosing Turner

syndrome. In some aspects, the method may further comprise diagnosing Down syndrome. In some aspects, the method may further comprise diagnosing Edwards syndrome. In some aspects, the method may further comprise diagnosing Patau syndrome. In some aspects, the genetic abnormality comprises a deletion in the genomic DNA. In some aspects, the genetic abnormality comprises a polymorphism. In some aspects, the genetic abnormality comprises a single gene disorders. In some aspects, the genetic abnormality comprises a chromosome translocation.

**[00368]** In some aspects, the sample is from an embryo. In some aspects, the sample comprises at least one cell from the embryo.

**[00369]** In some aspects, the method further comprises determining an implantation status of the embryo based on the detecting step. In some aspects, the genomic DNA is fragmented prior to attachment of the oligonucleotide tags.

**[00370]** In some aspects, the genomic DNA is fragmented by a restriction enzyme. In some aspects, the genomic DNA is fragmented by an allele-specific restriction enzyme.

**[00371]** In some aspects, the oligonucleotide tag comprises a ribonucleic acid. In some aspects, the ribonucleic acid is uracil. In some aspects, the ribonucleic acid is cytosine. In some aspects, the ribonucleic acid is adenine. In some aspects, the ribonucleic acid is guanine. In some aspects, the method further comprises synthesizing a copy of the genomic DNA-oligonucleotide tag conjugate to replace a ribonucleic acid sequence in the oligonucleotide tag with a deoxyribonucleic acid sequence.

**[00372]** In some aspects, the detecting step comprises detecting the copy of the genomic DNA-oligonucleotide tag conjugate, a complement of the copy of the genomic DNA-oligonucleotide tag conjugate, a reverse complement of the copy of the genomic DNA-oligonucleotide tag conjugate, or a portion thereof.

**[00373]** In some aspects, the copy of the genomic DNA-oligonucleotide tag conjugate is synthesized by a reverse transcriptase enzyme.

**[00374]** In some aspects, the detection step comprises hybridization of the genomic DNA-oligonucleotide tag conjugate, a complement of the genomic DNA-oligonucleotide tag conjugate, a reverse complement of the genomic DNA-oligonucleotide tag conjugate, or a portion thereof to a solid support. In some aspects, the detection step comprises hybridization of the genomic DNA, a complement of the genomic DNA, a reverse complement of the genomic DNA, or a portion thereof to a solid support. In some aspects, the detection step comprises hybridization of the oligonucleotide tag, a complement of the oligonucleotide tag, a reverse complement of the oligonucleotide tag, or a portion thereof to a solid support. In

some aspects, the detection step comprises hybridization of the copy of the genomic DNA-oligonucleotide tag conjugate, a complement of the copy of the genomic DNA-oligonucleotide tag conjugate, a reverse complement of the copy of the genomic DNA-oligonucleotide tag conjugate, or a portion thereof to a solid support. In some aspects, the detection step comprises hybridization of the copy of the genomic DNA, a complement of the copy of the genomic DNA, a reverse complement of the copy of the genomic DNA, or a portion thereof to a solid support. In some aspects, the detection step comprises hybridization of the copy of the oligonucleotide tag, a complement of the copy of the oligonucleotide tag, a reverse complement of the copy of the oligonucleotide tag, or a portion thereof to a solid support.

**[00375]** In some aspects, the detection step comprises sequencing of the genomic DNA-oligonucleotide tag conjugate, a complement of the genomic DNA-oligonucleotide tag conjugate, a reverse complement of the genomic DNA-oligonucleotide tag conjugate, or a portion thereof. In some aspects, the detection step comprises sequencing of the genomic DNA, a complement of the genomic DNA, a reverse complement of the genomic DNA, or a portion thereof. In some aspects, the detection step comprises sequencing of the oligonucleotide tag, a complement of the oligonucleotide tag, a reverse complement of the oligonucleotide tag, or a portion thereof. In some aspects, the detection step comprises sequencing of the copy of the genomic DNA-oligonucleotide tag conjugate, a complement of the copy of the genomic DNA-oligonucleotide tag conjugate, a reverse complement of the copy of the genomic DNA-oligonucleotide tag conjugate, or a portion thereof. In some aspects, the detection step comprises sequencing of the copy of the genomic DNA, a complement of the copy of the genomic DNA, a reverse complement of the copy of the genomic DNA, or a portion thereof. In some aspects, the detection step comprises sequencing of the copy of the oligonucleotide tag, a complement of the copy of the oligonucleotide tag, a reverse complement of the copy of the oligonucleotide tag, or a portion thereof.

**[00376]** In some aspects, the genomic DNA-oligonucleotide tag conjugate is amplified. In some aspects, the copy of the genomic DNA-oligonucleotide tag conjugate is amplified.

**[00377]** Further disclosed herein are kits and compositions for stochastically labeling a molecule (e.g., nucleic acids such as DNA and RNA molecules, or polypeptides such as proteins and enzymes). In some instances, the kits and compositions are used for stochastically labeling a polyadenylated molecule. The polyadenylated molecule can be a polyadenylated RNA molecule. Alternatively, the kits and compositions are used for stochastically labeling a DNA molecule.

**[00378]** In some instances, the kits comprise a stochastic label primer, universal PCR primer, dye-labeled primer, reverse transcriptase, UDG enzyme, polymerase, buffers, dNTP, array, gene specific primers, target specific primers, control oligo, or any combination thereof. Alternatively, the kits comprise a) a universal PCR primer; b) a Cy3 labeled universal PCR primer; c) a Cy3 TrueTag Grid; and d) an array. The array can be a 2x8 array. The kits disclosed herein can further comprise a stochastic label primer, carrier, control oligo, reverse transcriptase, UDG enzyme, polymerase, gene specific primers, target specific primers, dNTP, or any combination thereof.

**[00379]** The stochastic label primer can comprise a primer attached to an oligonucleotide tag, wherein the oligonucleotide tag comprises an oligo dT sequence, a unique identifier region, and a universal primer binding site, and wherein the universal primer binding site can enable annealing of the universal PCR primer of the kit to the stochastic label primer. In some instances, a stochastic label oligo dT primer is an oligonucleotide tag attached to an oligo dT primer.

**[00380]** The dye-labeled primer can comprise a primer labeled with a dye. The primer can be a universal PCR primer. Alternatively, the primer is a target-specific primer. The dye can be a fluorescent dye. In some instances, the dye is a Cy<sup>TM</sup> dye. In some instances, the Cy<sup>TM</sup> dye is a Cy3 dye.

**[00381]** The kits and compositions disclosed herein can further comprise a plurality of probes. In some instances, the plurality of probes is hybridized to the array. The plurality of probes can allow hybridization of the labeled-molecule to the array. The plurality of probes can comprise a sequence that is complementary to the stochastic label oligo dT. Alternatively, or additionally, the plurality of probes comprises a sequence that is complementary to the molecule.

**[00382]** The kits and compositions disclosed herein can further comprise one or more reagents to remove non-labeled molecules, excess primers, or excess oligonucleotide tags (or stochastic label primers) from the sample comprising labeled-molecules.

**[00383]** In some instances, the kits and compositions comprise a reverse transcriptase enzyme. The reverse transcriptase can be MMLV reverse transcriptase.

**[00384]** The kits and compositions can comprise a polymerase enzyme. The polymerase can be a Taq polymerase. For example, the Taq polymerase is a Titantium Taq polymerase.

**[00385]** In some instances, the kits and compositions comprise an enzyme. The enzyme can be an RNase enzyme. Alternatively, the enzyme is UDG. In other instances, the enzyme is a restriction enzyme. The enzyme can be a protease. In some instances, the enzyme is a

DNase enzyme. Alternatively, the enzyme is a ligase. The kits and compositions can comprise one or more reagents that can deactivate an enzyme disclosed herein.

**[00386]** In some instances, the kit further comprises a carrier substance. The carrier substance can increase the efficiency of a reaction (e.g., amplification, reverse transcription, ligation, hybridization). The carrier substance can be a nucleic acid molecule. The nucleic acid molecule can be an RNA molecule. The RNA molecule can be a polyadenylated RNA or phage RNA. The phage RNA can be RNA from a MS2 phage. Alternatively, the nucleic acid molecule is a plasmid.

**[00387]** The kit can further comprise a solid support. The solid support can be a bead. The bead can hybridize to the labeled-molecule. The bead can enable detection of the labeled molecule. The bead can be a streptavidin bead or biotin-labeled bead.

**[00388]** The kit can further comprise an algorithm for detecting and/or quantifying the labeled-molecule. Alternatively, or additionally, the kit comprises a software program for detecting/and or quantifying the labeled-molecule. In some instances, the kits further comprise a thermal cycler. The kits can further comprise one or more components for sequencing the labeled-molecule. The one or more components for sequencing can comprise a sequencer, one or more primers for sequencing, beads for sequencing, or any combination thereof. The kit can further comprise one or more components for detecting and/or quantifying the labeled-molecule. The one or more components for detecting and/or quantifying the labeled-molecule can comprise an array detector, array reader, bead detector, scanner, fluorometer, or any of the instruments or components disclosed herein.

## EXAMPLES

**[00389]** *Example 1. Absolute Counting Protocol*

**[00390]** Part 1. Reverse Transcription and Stochastic Labeling

**[00391]** In this step, the stochastic labels are annealed to the poly A RNA. To increase the overall efficiency of the subsequent reverse transcription reaction, a large amount of carrier RNA is also added to the sample.

**[00392]** In some instances, tips with low nucleic acid binding properties are used when pipetting extremely low concentrations of RNA. These special tips can be used for pipetting the RNA sample into the annealing master mix. If a dilution of the RNA is needed, low binding tubes can be used as well. Once the RNA has been added to the annealing master mix, regular tubes/tips can be used.

**[00393]** Make a master mix by combining the reagents listed below:

Water	7.8 $\mu$ l
-------	-------------

K562 Total RNA (1 $\mu$ g/ $\mu$ l)	1 $\mu$ l
10mM dNTP	1 $\mu$ l
Gene Specific dUTP Primer (10 $\mu$ M)	0.4 $\mu$ l
<u>Stochastic Labels (10<math>\mu</math>M)*</u>	<u>0.4<math>\mu</math>l</u>
Total	10.6 $\mu$ l

[00394] Add 2 $\mu$ l of the RNA sample to be analyzed.

[00395] Mix well by pipetting and spin briefly

[00396] Incubate at 65° for 5 minutes (Program 1), and then place the tubes on ice for at least 1 minute.

[00397] In this step, double stranded cDNA is created for the specific gene of interest.

Each cDNA molecule will now contain a primer site for the subsequent PCR step. Combine the following to make a master mix for reverse transcription:

5X First Strand Buffer	4 $\mu$ l
0.1M DTT	1 $\mu$ l
SuperRNaseIn (20U/ $\mu$ l)	1 $\mu$ l
MMLV RT	1 $\mu$ l
NEB Taq Polymerase	0.4 $\mu$ l

*The use MMLV RT and NEB Taq Polymerase instead of Superscript III and Titanium Taq can alternatively be used*

[00398] Add 7.4 $\mu$ l of master mix to each tube and mix by pipetting gently. Spin briefly.

[00399] Run the following program (Program 2) on the thermal cycler:

37° for 60 minutes

3 cycles of:

94° for 2 minutes

55° for 2 minutes

68° for 2 minutes

Then 4° forever

[00400] After the PCR reaction, it is necessary to digest the sample with Uracil DNA Glycosylase (UDG) to prevent the unincorporated primer from being amplified in the gene specific PCR.

[00401] To each reaction, add 0.5 $\mu$ l of UDG. Mix very well by pipetting. Transfer all liquid to a new PCR tube to ensure that there is no carryover of unmixed sample.

[00402] Incubate at 37° for 30 minutes, then 4°.

**[00403]** Part 2. Initial Gene Specific PCR

**[00404]** Combine the following reagents to make a master mix for PCR:

Nuclease-free water	10.9 $\mu$ l
10X NEB Taq Buffer	1.5 $\mu$ l
10mM dNTP	0.3 $\mu$ l
Gene Specific Primer (1 $\mu$ M)	1 $\mu$ l
Universal PCR primer (1 $\mu$ M)*	1 $\mu$ l
NEB Taq Polymerase	0.3 $\mu$ l

**[00405]** Final concentration of 0.05 $\mu$ M primer increases specificity of products

**[00406]** Add 5 $\mu$ l of labeled product from the previous step to a new PCR tube. Add 15 $\mu$ l PCR master mix to each sample.

**[00407]** Mix well by pipetting and spin briefly.

**[00408]** Run the following program (Program 4) on the thermal cycler:

94° for 2 minutes

30cycles of:

94° for 2 minutes

55° for 2 minutes

68° for 2 minutes

Then 68° for 4 minutes

4° forever

**[00409]** Part 3. Second, Nested PCR

**[00410]** Prepare the master mix for the second, nested, PCR in the pre-PCR area.

Nuclease-free water	39.5 $\mu$ l
10X NEB Taq Buffer	5 $\mu$ l
10mM dNTP	1 $\mu$ l
Gene Specific Nested Primer (10 $\mu$ M)	1 $\mu$ l
5Tye563 Labeled Universal PCR primer (10 $\mu$ M)*	1 $\mu$ l
NEB Taq Polymerase	0.5 $\mu$ l

**[00411]** Aliquot 48 $\mu$ l of master mix to a new PCR tube.

**[00412]** Add 2 $\mu$ l from the first PCR reaction to the tube in a separate room designated for post amplification processing to avoid contamination of the pre-PCR area. Perform all

subsequent steps in this area.

[00413] Mix well by pipetting and spin briefly.

[00414] Run the following program (Program 4) on the thermal cycler:

94° for 2 minutes

30cycles of:

94° for 2 minutes

55° for 2 minutes

68° for 2 minutes

Then 68° for 4 minutes

4° forever

[00415] Optional Step: Run 4ul of PCR product on a polyacrylamide 4-20% gradient TBE gel to assess size and purity

[00416] Part 4. Target Hybridization

[00417] Turn hyb oven on at 37°.

[00418] Prepare samples for hybridization to an Applied Microarray Inc. array slide. Add the following in a 0.2mL PCR tube:

Wash A (6X SSPE+0.01% Triton X-100) 55µl

Cy3 Control Oligo (760pM)\* 1µl

PCR product 20µl

---

Total 76µl

[00419] Mix by pipetting and spin briefly.

[00420] Incubate tubes at 95° to denature and then place on ice.

[00421] Remove adhesive seal from AMI array slide. Pipet each hybridization cocktail into a well of the AMI array slide. Make a note of the order in which the targets are added.

Cover slide with second strip of adhesive (included)

[00422] Place sealed array slide into humidity chamber and put into hybridization oven.

Incubate at 37° overnight.

[00423] Part 5. Array Wash and Scan

[00424] After the overnight hybridization, take the array slide out of the hybridization oven and remove adhesive cover. Pipet out remaining hybridization cocktail and save at -20° if desired.

[00425] Dispense 150µl Wash A to each used well. Aspirate liquid and dispense 150µl Wash B (0.6X SSPE+0.01% Triton X-100) to each well. Aspirate liquid and bring array slide

to scanner as the arrays will be scanned dry.

[00426] Turn on the Senovation FLAIR instrument. Wait 10 minutes for the machine to warm up.

[00427] Open the software and click, “Tray Open”. Place the array slide into the 4-slide holder. Be sure to seat the slide properly. In the software, click “Tray Close”.

[00428] Click the “Scan” Icon. A window appears with information about the scan to be performed. Modify the name of the scan if desired and select the appropriate wells to be scanned by clicking the “...” icon in the “scan positions” field. Click each well that is to be scanned. The software will circle each selected well in yellow. Click “ok”.

[00429] The Plate Overview window will appear showing the progress of the scan. Once a well is scanned, the color on the screen will turn from grey to green if the reference pattern has been detected and the grid has been positioned. If the reference pattern has not been detected, the well will be colored red. If any of the scans do not detect the reference, the grid may be manually aligned by clicking the “reanalyze” button at the top of the screen. This will display the grid, which can be positioned properly. Click the green “accept analysis” button at the top of the screen.

[00430] Once all of the grids have been aligned, the data can be exported. To obtain windows functionality, press the “windows” key on the keyboard and “D” simultaneously. Locate the scan results in the “my documents” folder under ArrayReader/senovation/arrayreader.scanresults. Open the appropriate scan folder and copy the TIFF images and the result .csv files to a flash drive or transfer through the network.

[00431] Proceed to data analysis either manually or with a computer software package.

[00432] *Example 2. Four experiments where 120 RNA molecules were added to a sample of background total RNA*

[00433] 240 copies of a polyadenylated nucleic acid fragment was added to a 10  $\mu$ L reaction containing 1X titanium Taq DNA polymerase buffer, 0.2  $\mu$ M dNTPs, 0.2  $\mu$ M of a pool of 960 oligo (dT) stochastic labels, 0.2  $\mu$ M of a second strand cDNA primer and 0.2  $\mu$ L of Taq DNA polymerase. In some reactions, an additional number of polyadenylated DNA fragments with sequences unrelated to the 240 copies of test nucleic acid fragment were also added. In reaction A,  $1 \times 10^{10}$  background polyadenylated DNA molecules were added to the reaction. In reaction B,  $1 \times 10^9$  background polyadenylated DNA molecules were added to the reaction. In reaction C,  $1 \times 10^6$  background polyadenylated DNA molecules were added to the reaction. And, in reaction D, no background polyadenylated DNA molecules were added to the reaction. 10 ng, 1ng or 1pg of randomly fragmented and polyadenylated human genomic

DNA was tested. After 3 cycles of incubation at 94 °C for 2 min, 45 °C for 2 min and 65 °C for 5 min, 1 unit of Uracil DNA glycosylase is added and the reaction is incubated for 30 min at 37 °C. Half of the reaction is then added into a 20 µL PCR reaction consisting of 1X Titanium buffer, 0.2 µM dNTP, 0.2 µM gene-specific forward primer, 0.2 µM universal reverse primer and 0.3 µL Titanium Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 68 °C for 20 sec. A final incubation at 68 °C for 4 min was performed. A nested PCR is performed following the same conditions as the first PCR, except that a nested Forward primer was used. 2 µL of a 1:25 dilution of the initial PCR was used as template for the nested PCR. PCR products were randomly fragmented with DNase, biotin-labeled with Terminal transferase enzyme and then hybridized to a detector array for 12 hours at 37 °C. Signals from hybridized DNAs were detected via staining with Streptavidin conjugated Phycoerytherin and imaging on a microarray scanner. FIG. 2A-D shows the signals from hybridized DNAs for reactions A-D, respectively. The number of labels present in the hybridized DNA is counted and used to determine the number of original copies of nucleic acid fragments.

Reaction	# of labels	# of original copies
A	122	130
B	116	124
C	109	114
D	115	122

**[00434] *Example 3. Comparison with digital PCR***

**[00435]** The concentration of an in vitro transcribed RNA was determined using an Agilent bioanalyzer instrument. 0.5 µg of the RNA was mixed with 2 µg of a K562 cell line total RNA which was used as a carrier. The RNA mixture in 3 µL was added to 1 µL of a 10 mM dNTP solution and 2 µL of a 10 µM pool of 960 oligo (dT) labels and 7 µL of water. This mixture was incubated at 65 °C for 5min and immediately chilled on ice. 4 µL of a first strand reaction buffer (250mM Tris-HCl (pH 8.3 at 25°C), 375mM KCl), 1 µL of 0.1 M DTT, 1 µL of RNase inhibitor (20 units) and 1 µL of superscript II reverse transcriptase (200 units) was added and the reaction was incubated at 50 °C for 60min and then at 70 °C for 15min. 1 µL of RNase H (2 units) was added and the reaction was incubated at 37 °C for 20min. Digital PCR was used to quantitate the number of copies of cDNA synthesized from the in vitro transcribed RNA. The sample was also test by stochastic labeling PCR. 90 copies

of the cDNA (as determined by digital PCR) was added to a 10  $\mu$ L reaction containing 1X titanium PCR buffer, 0.2  $\mu$ M dNTPs and 0.2  $\mu$ L of titanium taq polymerase. The reaction was incubated for 3 cycles at 94 °C for 2min, 55 °C for 2min and 68 °C for 2 min. 1 unit of uracil DNA glycosylase was added and the reaction was incubated at 37 °C for 30min. First and nested PCRs, fragmentation, biotin-labeling and array detection were performed as described in Example 2. FIG. 3 shows the signals from the labels in the hybridized DNA. The number of labels present in the hybridized DNA is counted and used to determine the number of original copies of nucleic acid fragments. 40 labels were present in the hybridized DNA and 41 copies were determined by stochastic labeling, as compared to 43 copies as determined by digital PCR. These results demonstrate that stochastic labeling is an effective method for determining the count of a molecule and its accuracy is comparable to digital PCR.

**[00436] *Example 4. RT yield increased with reaction carriers***

**[00437]** To test the effectiveness of carrier RNAs on improving the reverse transcription yield and as a means to reduce non-specific RNA or cDNA losses during reactions, copies of an in vitro transcribed polyadenylated RNA was tested with stochastic labeling following the protocol described in example 2. Additionally, total RNA isolated from mammalian cells, yeast, or *E. coli*, short polyadenylated synthetic ribonucleotide, yeast tRNA, or MS2 phage RNA were added to the reaction mixture. Each reaction used anywhere between 0.5  $\mu$ g to 2  $\mu$ g of carrier RNA. The number of RNA molecules reverse transcribed to cDNA was determined by the number of observed labels detected on the array, and in each case, the effectiveness of each carrier RNA tested could be easily determined. FIG. 4A-D shows the observed labels for reactions A-D, respectively.

Reaction	# of Input molecules	Carrier RNA	RNase H	# of labels
A	188	Total RNA from X	--	158
B	188	MS2 phage RNA	--	165
C	188	Yeast tRNA	--	1
D	188	--	RNase H	154

**[00438] *Example 5. Comparison of MMLV and RNase H minus MMLV reverse transcriptase***

**[00439]** The performance of the wild type MMLV reverse transcriptase was compared with the RNase H minus mutant version (Superscript III) of the enzyme. 375 copies of an in

vitro transcribed polyadenylated RNA was added to a carrier of 1  $\mu$ g of a K562 cell line total RNA. The RNAs were added into a 12.6  $\mu$ L reaction containing 1  $\mu$ L of a 10mM dNTP solution, 0.4  $\mu$ L of a 10  $\mu$ M second strand primer, 0.4  $\mu$ L of a 10 $\mu$ M pool of 960 oligo(dT) labels. The reaction was incubated at 65 °C for 5 min to denature the RNA, and then quickly chilled on ice. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units) and 0.4  $\mu$ L of Taq DNA polymerase (2 units) was added. Additionally, in reaction A, 1  $\mu$ L (200 units) of the RNase H minus mutant (Superscript III) was added. And, in reaction B, 1  $\mu$ L (200 units) of the wild type MMLV reverse transcriptase was added. The reactions were incubated at 42 °C for 60min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 68 °C for 2min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5  $\mu$ L of the reaction was then added to a 20  $\mu$ L PCR reaction consisting of 1X Titanium buffer, 0.2  $\mu$ M dNTP, 0.2  $\mu$ M gene-specific forward primer, 0.2  $\mu$ M universal reverse primer and 0.3  $\mu$ L Titanium Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 68 °C for 20 sec. A final incubation at 68 °C for 4 min was performed. A nested PCR is performed following the same conditions as the first PCR, except that a nested Forward primer was used. 2  $\mu$ L of a 1:25 dilution of the initial PCR was used as template for the nested PCR. PCR products were randomly fragmented with DNase, biotin-labeled with Terminal transferase ezyme and then hybridized to a detector array for 12 hours at 37 °C. Signals from hybridized DNAs were detected via staining with Streptavidin conjugated Phycoerytherin and imaging on a microarray scanner. FIG. 5A-B show the labels present in the hybridized DNA in reactions A and B, respectively. The number of labels present in the hybridized DNA is counted and used to determine the number of original copies of nucleic acid fragments.

Reaction	# Input RNA molecules	Reverse Transcriptase	# of labels
A	188	Superscript III	159
B	188	MMLV	124

[00440] *Example 6. Comparison of polymerases for second strand synthesis*

[00441] The performance of Taq polymerase was compared to Titanium Taq polymerase. 1875 copies of an in vitro transcribed polyadenylated RNA was added to reaction A. 188 copies of an in vitro transcribed polyadenylated RNA was added to reaction B. 1875 copies

of an in vitro transcribed polyadenylated RNA was added to reaction C. And, 188 copies of an in vitro transcribed polyadenylated RNA was added to reaction D. 1  $\mu$ g of carrier RNA from a K562 cell line was added to each of the reaction mixtures. The RNAs were added into a 12.6  $\mu$ L reaction containing 1  $\mu$ L of a 10mM dNTP solution, 0.4  $\mu$ L of a 10  $\mu$ M second strand primer, 0.4  $\mu$ L of a 10 $\mu$ M pool of 960 oligo(dT) labels. The reactions were incubated at 65 °C for 5 min to denature the RNA, and then quickly chilled on ice. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units), reverse transcriptase, and 0.4  $\mu$ L of Taq DNA polymerase (2 units) were added to each reaction. The reactions were incubated at 42 °C for 60min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 68 °C for 2min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5  $\mu$ L of reactions A and B were mixed with a 20  $\mu$ L PCR reaction consisting of 1X Taq buffer, 0.2  $\mu$ M dNTP, 0.2  $\mu$ M gene-specific forward primer, 0.2  $\mu$ M universal reverse primer and 0.3  $\mu$ L Taq polymerase. 5  $\mu$ L of reactions C and D were mixed with a 20  $\mu$ L PCR reaction consisting of 1X Titanium buffer, 0.2  $\mu$ M dNTP, 0.2  $\mu$ M gene-specific forward primer, 0.2  $\mu$ M universal reverse primer and 0.3  $\mu$ L Titanium Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 68 °C for 20 sec. A final incubation at 68 °C for 4 min was performed. A nested PCR is performed following the same conditions as the first PCR, except that a nested Forward primer was used. 2  $\mu$ L of a 1:25 dilution of the initial PCR was used as template for the nested PCR. PCR products were randomly fragmented with DNase, biotin-labeled with Terminal transferase enzyme and then hybridized to a detector array for 12 hours at 37 °C. Signals from hybridized DNAs were detected via staining with Streptavidin conjugated Phycoerytherin and imaging on a microarray scanner. FIG. 6A-D shows the labels present in the hybridized DNA in reactions A-D, respectively. The number of labels present in the hybridized DNA is counted and used to determine the number of original copies of nucleic acid fragments.

Reaction	# Input RNA molecules	Polymerase	# of labels
A	1875	Taq	--
B	188	Taq	157
C	1875	Titanium Taq	--
D	188	Titanium Taq	129

**[00442] *Example 7. Absolute quantitation of mRNA by counting individual DNA molecules***

[00443] mRNA molecules can be quantitated by the addition of labels prior to amplification of cDNA molecules (FIG. 19). Labeled cDNA molecules are formed by cDNA synthesis of an mRNA molecule by the addition of a deoxy-oligonucleotide primer with (1) an oligo dT sequence to anneal to the poly-A RNA tail; (2) a collection of predetermined or random sequence label tags; and (3) a common or universal PCR primer sequence. The labeled cDNA molecules are amplified using gene-specific primers and a common or universal PCR primer. After amplification, the number of labels of different sequence composition can be readily detected by hybridization, sequencing or other detection methods. The difficult task of counting the number of individual mRNA molecules in solution is transformed into the simple task of determining the number of types of different labels, each being present at high concentrations following amplification, provided that the initial label sequence diversity is sufficiently greater than the number of molecules present. Any other suitable method can also be used to incorporate labels into the RNA or cDNA molecules before or during amplification. Any other PCR or non-PCR based methods can also be used to amplify the RNA or cDNA molecules. Although helpful in these examples, amplification of the labeled molecules may not be required for detection.

**[00444] *Example 8. Digital microarray for RNA expression***

[00445] The mRNA is reverse transcribed using a pool of n oligo-dT label primers (random primers with labels may also be used) (FIG. 20). The cDNA can be optionally amplified with methods such as PCR and T7 amplification. The labels are amplified along with each cDNA molecule. cDNAs are hybridized to digital arrays to determine the number of distinct labels for each gene of interest. Hybridization requires both presence of the gene sequence, most likely a segment on the 3' exon of the gene, and one of the label sequences. An array with 7 million features is sufficient to detect a collection of 350 labels applied to a sample with 20,000 different mRNA sequences to determine the number of copies of each mRNA present in the sample. A subset of the 350 label primers may be applied at a lower concentration to increase the effective dynamic range of measurement. This method is particularly advantageous for sampling limiting amounts of starting material, such as in single cells.

**[00446] *Example 9. Digital microarray for DNA copy number***

[00447] Genomic DNA is digested into small fragments in one or more reactions using one or more restriction enzymes. Adaptors with label sequences are ligated to the DNA fragments (FIG. 21). The ligated fragments are optionally amplified. Ligated fragments may optionally

be digested with one or more restriction enzymes prior to amplification to prevent the replication of certain fragments, which is useful in the selective amplification of only fragments of interest. Hybridization to digital arrays detects the number of distinct labels ligated to each restricted fragment. Using 350 label sequences, an array of 7 million features can assay 20,000 fragments in the genome, which represents average intervals of 150kb in humans. Additionally, some allele specific fragments may be assayed by choosing restriction enzymes (e.g., 4 base cutters) specific for an allele of interest.

**[00448] *Example 10. Digital microarray for microRNAs***

**[00449]** Labels are attached to the 3' and 5' ends of microRNA by ligation or other means (FIG. 22). The label-microRNA complex is reverse transcribed to generate label-DNA products. The label-DNA products are optionally amplified. The label-DNA products are hybridized on digital array to detect the number of labels per microRNA. miRBase 18 (<http://www.mirbase.org/>) was released in November 2011 and lists 1921 unique mature human miRNAs. An array of 2 million features can adequately detect 1000 labels ligated to the 1921 miRNAs.

**[00450] *Example 11. Digital microarray for single cell pre-implantation genetic diagnosis (PGD)***

**[00451]** Primary challenge with single-cell genomic DNA amplification assays is from allele dropout and replication bias. As shown in the computation modeling analysis in FIG. 43 where every molecule has a 0.8 probability of replication, molecules of 1:1 initial copy ratios can easily be distorted to 1:10 or greater just after a few replication cycles.

**[00452]** However, when labels are first attached prior to amplification, counting labels to determine copy number is unaffected by replication bias, so long as replication occurs. Although, this does not solve the problem of allele dropouts, aneuploidy determination and large regions of deletion or amplification can be easily and accurately determined. This is particularly useful for PGD applications.

**[00453] *Example 12. Digital microarray for measuring fetal aneuploidy in maternal circulating nucleic acids***

**[00454]** Digital microarray can be used to measure fetal aneuploidy in maternal circulating nucleic acids. A sample comprising maternal circulating nucleic acids is provided. The DNA is fragmented using a 4 base cutter. Labels are attached to the fragmented DNA. Circulate and multiplex PCR to amplify 40 chromosome 21 markers and 10 control chromosome markers. Detect amplified label-DNA products on an array of 5 million features. The number

of copies of chromosome 21 can be used to determine the occurrence of fetal aneuploidy (FIG. 24).

**[00455] Example 13. Absolute quantitation of mRNA by counting individual DNA molecules**

**[00456]** mRNA molecules can be quantitated by the incorporation of labels during first-strand cDNA synthesis (FIG. 25). Labeled cDNA molecules are formed by cDNA synthesis of an mRNA molecule by the addition of a deoxy-oligonucleotide primer with (1) an oligo dT sequence to anneal to the poly-A RNA tail; (2) a collection of predetermined or random sequence label tags; and (3) a common or universal PCR primer sequence. After first-strand cDNA synthesis, the number of labels of different sequence composition can be readily detected by hybridization, sequencing or other detection methods. The difficult task of counting the number of individual mRNA molecules in solution is transformed into the simple task of determining the number of types of different labels, each being present at high concentrations following amplification, provided that the initial label sequence diversity is sufficiently greater than the number of molecules present. Any other suitable method can also be used to incorporate labels into the RNA or cDNA molecules before or during first-strand cDNA synthesis.

**[00457] Example 14: Titration experiment with serial dilutions of kanamycin RNA**

**[00458]** A titration curve was generated by performing serial dilutions of kanamycin RNA to illustrate the broad dynamic range of the absolute counting protocol. Each of 9 serial dilutions was normalized to a concentration of 0.25 fg/μl from 2.5 pg/μl, 1.25 pg/μl, 0.25 pg/μl, 0.125 pg/μl, 0.025 pg/μl, 12.5 fg/μl, 2.5 fg/μl, 1.25 fg/μl and 0.25 fg/μl. All of the dilutions were made using a dilution solution of 1ng/μl E. Coli total RNA in tubes pre-rinsed with a solution of 10ng/μl yeast RNA to hinder the sticking of the sample RNA to the walls of the tube. The samples were added to a 12.6 μl reaction containing 1 μg E. Coli total RNA, 1 μl of a 10 mM solution of dNTP's, 0.4 μl of a 10 uM dU primer specific for kanamycin and 0.4 μl of a 10 μM pool of 960 dT oligo labels. The reaction was incubated at 65 °C for 5 min to denature the RNA, and then quickly chilled on ice. 4 μL of a 5X first strand buffer, 1 μL of a 0.1M DTT, 1 μL of superase RNase inhibitor (20 units), 1 μL (200 units) of the wild type MMLV reverse transcriptase and 0.4 μL of Taq DNA polymerase (2 units) were added. The reactions were incubated at 37 °C for 60 min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 72 °C for 2 min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5 μL of the reaction was then added to a 20 μL PCR reaction consisting of 1X Taq Reaction buffer, 0.2 μM

dNTP, 0.05 uM gene-specific forward primer, 0.05  $\mu$ M universal reverse primer and 0.3  $\mu$ L Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 72 °C for 20 sec. A final incubation at 72 °C for 4 min was performed. A nested PCR was performed using a nested forward primer and the universal reverse primer with a Cy3 label attached. 0.5  $\mu$ l of the initial PCR was used as template for the nested PCR. PCR conditions were the same as for the first PCR except that the 58°C step was performed at 55°C. The samples were hybridized to a detector array at 37°C overnight and scanned the following day using a fluorescence reader to detect which positions on the array contained the Cy3 label. The number of positive spots was used to determine the initial concentration of sample. FIG. 35 shows the dilution scheme. FIG. 36A-H shows the scatter plots of results and Table 1 shows the results. FIG. 37 shows the correlation graph.

[00459] Table 1

FIG	Initial Concentration	Dilution Factor	Expected Count	Actual Count
36A	2.5 pg/ $\mu$ L	10000	130	199
36B	1.25 pg/ $\mu$ L	5000	130	178
36C	0.25 pg/ $\mu$ L	1000	130	170
36D	0.125 pg/ $\mu$ L	500	130	153
36E	1.025 pg/ $\mu$ L	50	130	154
36F	12.5 fg/ $\mu$ L	10	130	117
36G	2.5 fg/ $\mu$ L	5	130	95
36H	1.25 fg/ $\mu$ L	1	130	137

[00460] **Example 15: Titration experiment with serial dilutions of human liver RNA to measure GAPDH expression**

[00461] A titration curve was generated by performing serial dilutions of human liver total RNA to illustrate the ability of the stochastic labeling protocol to detect levels of gene expression. Each of 8 serial dilutions was normalized to a concentration of 1.25 pg/ $\mu$ L from 5000 pg/ $\mu$ L, 1250 pg/ $\mu$ L, 500 pg/ $\mu$ L, 125 pg/ $\mu$ L, 50 pg/ $\mu$ L, 12.5 pg/ $\mu$ L, 5 pg/ $\mu$ L and 1.25 pg/ $\mu$ L. All of the dilutions were made using a dilution solution of 1 ng/ $\mu$ l E. Coli total RNA in tubes pre-rinsed with a solution of 10 ng/ $\mu$ l yeast RNA to hinder the sticking of the sample RNA to the walls of the tube. The samples were added to a 12.6  $\mu$ l reaction containing 1  $\mu$ g E. Coli total RNA, 1  $\mu$ l of a 10 mM solution of dNTP's, 0.4  $\mu$ l of a 10 uM dU primer specific for GAPDH and 0.4  $\mu$ l of a 10  $\mu$ M pool of 960 dT oligo labels. The reaction was incubated at 65

°C for 5 min to denature the RNA, and then quickly chilled on ice. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1 M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units), 1  $\mu$ L (200 units) of the wild type MMLV reverse transcriptase and 0.4  $\mu$ L of Taq DNA polymerase (2 units) was added. The reactions were incubated at 37 °C for 60 min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 72 °C for 2 min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5  $\mu$ L of the reaction was then added to a 20  $\mu$ L PCR reaction consisting of 1X Taq Reaction buffer, 0.2  $\mu$ M dNTP, 0.05  $\mu$ M gene-specific forward primer, 0.05  $\mu$ M universal reverse primer and 0.3  $\mu$ L Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 72 °C for 20 sec. A final incubation at 72 °C for 4 min was performed. A nested PCR was performed using a nested forward primer and the universal reverse primer with a Cy3 label attached. 0.5 $\mu$ l of the initial PCR was used as template for the nested PCR. PCR conditions were the same as for the first PCR except that the 58°C step was performed at 55°C. The samples were hybridized to a detector array at 37°C overnight and scanned the following day using a fluorescence reader to detect which positions on the array contained the Cy3 label. The number of positive spots was used to determine the initial concentration of sample. FIG. 38 shows the dilution scheme. FIG. 39 shows the scatter plots of results and Table 2 shows the results. FIG. 40 shows and correlation graph.

[00462] Table 2.

FIG	Initial Concentration	Dilution Factor	Actual Count
39A	5000 pg/ $\mu$ L	4000	73
39B	1250 pg/ $\mu$ L	1000	63
39C	500 pg/ $\mu$ L	400	69
39D	125 pg/ $\mu$ L	100	46
39E	50 pg/ $\mu$ L	40	65
39F	12.5 pg/ $\mu$ L	10	38
39G	5 pg/ $\mu$ L	4	53
39H	1.25 pg/ $\mu$ L	1	37

[00463] **Example 16: Measurements of control bacterial genes**

[00464] The protocol was validated using Poly A bacterial control RNAs (Lys, Thr, Dap and Phe), as well as RNA from the Kanamycin resistance gene. 4 different dilutions of each control were used to validate the accuracy of the counts. The samples were added to a 12.6  $\mu$ l reaction containing 1 $\mu$ g E. Coli total RNA, 1  $\mu$ L of a 10 mM solution of dNTP's, 0.4  $\mu$ L of a

10  $\mu$ M gene specific dU primer and 0.4  $\mu$ l of a 10  $\mu$ M pool of 960 dT oligo labels. The reaction was incubated at 65 °C for 5 min to denature the RNA, and then quickly chilled on ice. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1 M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units), 1  $\mu$ L (200 units) of the wild type MMLV reverse transcriptase and 0.4  $\mu$ L of Taq DNA polymerase (2 units) was added. The reactions were incubated at 37 °C for 60min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 72 °C for 2 min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5  $\mu$ L of the reaction was then added to a 20  $\mu$ L PCR reaction consisting of 1X Taq Reaction buffer, 0.2  $\mu$ M dNTP, 0.05  $\mu$ M gene-specific forward primer, 0.05  $\mu$ M universal reverse primer and 0.3  $\mu$ L Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 72 °C for 20 sec. A final incubation at 72 °C for 4 min was performed. A nested PCR was performed using a nested forward primer and the universal reverse primer with a Cy3 label attached. 0.5 $\mu$ l of the initial PCR was used as template for the nested PCR. PCR conditions were the same as for the first PCR except that the 58 °C step was performed at 55 °C. The samples were hybridized to a detector array at 37 °C overnight and scanned the following day using a fluorescence reader to detect which positions on the array contained the Cy3 label. The number of positive spots was used to determine the initial concentration of sample. FIG. 41 shows scatter plots of the results from the lowest concentration dilutions and Table 3 displays a summary table of the results.

[00465] Table 3.

FIG.	Gene	Copies in reaction (manufacturer)	Copies measured (CR)	Copies measured (digital PCR)
41A	Lys ( <i>B. subtilis</i> )	190	195	
41B	Dap ( <i>B. subtilis</i> )	137	119	
41C	Phe ( <i>B. subtilis</i> )	162	116	
41D	Thr ( <i>B. subtilis</i> )	189	108	
42	Kanamycin resistance gene (plasmid)	750	608	520

**[00466] Example 17: Comparison of quantification of kanamycin RNA by stochastic labeling and digital PCR**

[00467] The counts of kanamycin RNA generated by stochastic labeling were compared to the counts obtained from digital PCR as another example of validation. 5  $\mu$ g of kanamycin RNA was added to a 13  $\mu$ l reaction containing 2  $\mu$ g of E. Coli total RNA, 1  $\mu$ l of a 10 mM solution of dNTP's and 2  $\mu$ l of a 10 uM solution of 960 dT oligo labels. The sample was heated to 65°C for 5 minutes, then chilled on ice. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units), 1  $\mu$ L (200 units) of Superscript III reverse transcriptase was added to the reaction. The sample was incubated at 50°C for 60 minutes, then heated to 70°C for 15 minutes, then cooled to 4°C. 2 units of RNase H were added and the sample was incubated at 37°C for 20 minutes. 29  $\mu$ l of TE was added after the final incubation. A 50 million-fold serial dilution was performed and 1  $\mu$ l was used in seventy-five 15  $\mu$ l digital PCR reactions. Each of these reactions contained 7.5  $\mu$ l of a 2X SYBR PCR master mix, 0.13  $\mu$ l of a 10 uM kanamycin forward primer and 0.13  $\mu$ l of a 10 uM kanamycin reverse primer. PCR conditions included an initial incubation at 95°C for 30 seconds followed by 45 cycles of 95°C for 15 seconds and 58°C for 60 seconds. A melting curve program followed the PCR for the purpose of validating the results. FIG. 42 shows the scatter plot of results and Table 3 shows the summary of the counts for kanamycin. FIG. 42 shows the dPCR results of 0.0002pg Kanamycin RNA using SYBR green qPCR reagents. As shown in FIG. 42, 50 positive wells were observed out of 75 reactions, n=104 molecules present in 0.0002pg (520 molecules present in 0.001pg).

**[00468] Example 18: Gene expression measurements in Liver RNA**

[00469] The gene expression values of targets of varying abundance were measured using stochastic labeling. Based on previous assumptions of transcript abundance, differing concentrations of human liver total RNA were used to test each of 9 genes; GAPDH, B2M, RPL19, SDHA, GUSB, TUBB, ABCF1, G6PD, and TBP. The RNA quantities used in each reaction were designed to target the ideal counting range of 1-300 molecules and they were 0.625 pg, 1.25 pg, 1.25 pg, 125 pg, 12.5 pg, 12.5 pg, 2500 pg, 650 pg and 650 pg, respectively. The samples were added to a 12.6  $\mu$ l reaction containing 1 $\mu$ g E. Coli total RNA, 1  $\mu$ l of a 10 mM solution of dNTP's, 0.4  $\mu$ l of a 10 uM gene specific dU primer and 0.4  $\mu$ l of a 10 uM pool of 960 dT oligo labels. The reaction was incubated at 65 °C for 5 min to denature the RNA, and then quickly chilled on ice. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1 M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units), 1  $\mu$ L (200 units) of the wild type

MMLV reverse transcriptase and 0.4  $\mu$ L of Taq DNA polymerase (2 units) was added. The reactions were incubated at 37 °C for 60 min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 72 °C for 2 min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5  $\mu$ L of the reaction was then added to a 20  $\mu$ L PCR reaction consisting of 1X Taq Reaction buffer, 0.2  $\mu$ M dNTP, 0.05  $\mu$ M gene-specific forward primer, 0.05  $\mu$ M universal reverse primer and 0.3  $\mu$ L Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 72 °C for 20 sec. A final incubation at 72 °C for 4 min was performed. A nested PCR was performed using a nested forward primer and the universal reverse primer with a Cy3 label attached. 0.5 $\mu$ l of the initial PCR was used as template for the nested PCR. PCR conditions were the same as for the first PCR except that the 58°C step was performed at 55°C. The samples were hybridized to a detector array at 37°C overnight and scanned the following day using a fluorescence reader to detect which positions on the array contained the Cy3 label. The number of positive spots was used to determine the initial concentration of sample. Table 4 shows a summary of the counts for all 9 genes.

[00470] Table 4.

Liver RNA	Gene	Copies measured by CR
10 picograms (~ 1 cell)	B2M	304
	RPL19	200
	GAPDH	376
10 picograms (~ 1 cell)	SDHA	82
	GUSB	19
	TUBB	34
100 picograms (~ 10 cells)	GP6D	30
	ABCF1	3
	TBP	15

[00471] **Example 19: Absolute quantitation of mRNA molecules directly from cell lysates**

[00472] This example describes a method to generate transcript counts directly from cell lysates. A range of 40-100 cells from the Ramos (RA1) cell line washed in PBS were placed in a PCR tube with the following reagents: 1 $\mu$ l Triton X-100 5%, 1 $\mu$ g E. Coli total RNA, 1 $\mu$ l of a 10mM solution of dNTP's, 0.4 $\mu$ l of a gene specific dU primer and 0.4 $\mu$ l of a 10uM pool of 960 dT Oligos. The samples were heated to 70°C for 10 minutes and chilled on ice to lyse

the cells and allow the primers to anneal. 4  $\mu$ L of a 5X first strand buffer, 1  $\mu$ L of a 0.1M DTT, 1  $\mu$ L of superase RNase inhibitor (20 units), 1  $\mu$ L (200 units) of the wild type MMLV reverse transcriptase and 0.4  $\mu$ L of Taq DNA polymerase (2 units) was added. Control samples were also performed for the same cell numbers without the reverse transcriptase. The reactions were incubated at 37 °C for 60 min, followed by 3 cycles of 94 °C for 2min, 55 °C for 2 min and 72 °C for 2 min. 1 unit of uracil DNA glycosylase was added and the reaction was mixed and moved to a new tube and incubated at 37 °C for 30min. 5  $\mu$ L of the reaction was then added to a 20  $\mu$ L PCR reaction consisting of 1X Taq Reaction buffer, 0.2  $\mu$ M dNTP, 0.05 uM gene-specific forward primer, 0.05  $\mu$ M universal reverse primer and 0.3  $\mu$ L Taq polymerase. PCR conditions were 94 °C for 2 min followed by 30 cycles of 94 °C for 20 sec, 58 °C for 20 sec and 72 °C for 20 sec. A final incubation at 72 °C for 4 min was performed. A nested PCR was performed using a nested forward primer and the universal reverse primer with a Cy3 label attached. 0.5 $\mu$ l of the initial PCR was used as template for the nested PCR. PCR conditions were the same as for the first PCR except that the 58°C step was performed at 55°C. The samples were hybridized to a detector array at 37C overnight and scanned the following day using a fluorescence reader to detect which positions on the array contained the Cy3 label. The number of positive spots was used to determine the initial concentration of the RPL19 transcript in the cells. FIG. 43 shows a diagram summarizing the adaptation of the stochastic labeling protocol directly to cells.

**[00473] Example 20. Optimization of cDNA synthesis**

**[00474]** Three cDNA synthesis reactions were conducted. The composition of the three reactions are described below.

**[00475]** Reaction 1: Std = control RNA + 10nM dT24 + Reverse Transcriptase

**[00476]** Reaction 2: Chum = control RNA +10ng poly A carrier RNA + 10nM dT24 + Reverse Transcriptase

**[00477]** Reaction 3: Bead = control RNA + 1 x 10<sup>6</sup> dT40 beads + Reverse Transcriptase

**[00478]** The reactions were incubated for 1 hour at 42 °C, then diluted to the indicated number of input RNA copies for 35 cycles of PCR. The PCR products for each reaction are shown in FIG. 32. As shown in FIG. 32A, the RNA conversion to cDNA is higher on beads than in-solution

**[00479]** Having now fully described the present invention in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious to one of ordinary skill in the art that the same can be performed by modifying or changing the

invention within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any specific embodiment thereof, and that such modifications or changes are intended to be encompassed within the scope of the appended claims.

**[00480]** All publications, patents and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains, and are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.

**We claim:**

1. A method of counting the number of occurrences of an mRNA target molecule in a sample comprising:
  - a). providing a plurality of oligonucleotide tags, wherein an oligonucleotide tag of the plurality of oligonucleotide tags comprises a target specific sequence, a unique identifier sequence, and an optional universal PCR primer sequence, and wherein the target specific sequence comprises an oligo dT sequence;
  - b). combining a sample comprising an mRNA target molecule comprising a polyA sequence with the plurality of oligonucleotide tags such that the oligo dT sequence of the oligonucleotide tag hybridizes with the polyA sequence of the mRNA target molecule to form a labeled mRNA molecule, wherein different occurrences of the mRNA target molecule are capable of hybridizing to different oligonucleotide tags, and wherein the unique identifier region uniquely identifies occurrences of the mRNA target molecule;
  - c). synthesizing a copy of the labeled mRNA target molecule by reverse transcription to form a labeled DNA molecule, wherein at least a portion of the oligonucleotide tag acts as a primer for the reverse transcription reaction, and wherein the labeled DNA molecule comprises a copy of the mRNA target molecule and an oligonucleotide tag;
  - d). detecting the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled DNA molecule, or a portion thereof; and
  - e). determining a count of different labeled DNA molecules, thereby counting the number of occurrences of the mRNA target molecule in the sample.
2. The method of claim 1, wherein the unique identifier sequence comprises a random sequence.
3. The method of any one of the preceding claims, wherein the unique identifier sequence comprises a predetermined sequence.

4. The method of any preceding claim, wherein the oligonucleotide tag is at least 20 nucleotides in length.
5. The method of any one of the preceding claims, further comprising amplifying the labeled DNA molecule by polymerase chain reaction (PCR).
6. The method of claim 5, wherein the polymerase chain reaction comprises nested PCR.
7. The method of claim 6, wherein the nested PCR comprises conducting the nested PCR on the labeled DNA molecule to produce nested PCR labeled amplicons.
8. The method of any one of claims 5-7, wherein the polymerase chain reaction comprises annealing a gene-specific primer to the labeled DNA molecule.
9. The method of any of claims 6-8, wherein the polymerase chain reaction comprises annealing a universal primer to the universal PCR primer sequence of the oligonucleotide tag.
10. The method of any one of the preceding claims, wherein the detection step d) comprises determining the sequence of the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled DNA molecule, an amplified product of the labeled DNA molecule, a portion thereof, or a combination thereof.
11. The method of any one of the preceding claims, wherein the detection step d) comprises hybridizing the labeled DNA molecule, a complement of the labeled DNA molecule, a reverse complement of the labeled DNA molecule, an amplified product of the labeled DNA molecule, a portion thereof, or a combination thereof to a solid support.
12. The method of claim 11, wherein the solid support comprises a bead.
13. The method of claim 11 or claim 12, wherein the solid support comprises an array.

14. The method of any one of the preceding claims, wherein the sample is from fewer than 50 cells.
15. The method of any one of the preceding claims, wherein the sample is from a single cell.
16. The method of any one of the preceding claims, wherein the sample comprises cell lysate.
17. The method of any one of the preceding claims, wherein the number of oligonucleotide tags having different identifier sequences in the plurality of oligonucleotide tags is at least 100.
18. The method of any one of the preceding claims, wherein the number of oligonucleotide tags having different identifier sequences in the plurality of oligonucleotide tags is at least 1,000.
19. The method of any one of the preceding claims, wherein the number of oligonucleotide tags having different identifier sequences in the plurality of oligonucleotide tags is at least 10,000.
20. The method of any one of the preceding claims, wherein the labeled mRNA molecule comprises a stochastically labeled mRNA molecule, wherein the number of oligonucleotide tags having different identifier sequences in the plurality of oligonucleotide tags is at least 5 times greater than the number of occurrences of the mRNA target molecule.

FIG. 1





FIG. 2.



It is a count of the number of labels detected (as bright spots) and  $n$  is the number of RNA molecules measured biased on  $k$ .

FIG. 3.




FIG. 4




FIG. 5.




FIG. 6.




FIG. 7.




FIG. 8.




FIG. 9




FIG. 10




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D




FIG. 12






FIG. 13



A.



B.

FIG. 14




FIG. 15





FIG. 16



## Recognition Site:

5'...GATC...3'  
3'...CTAG...5'

FIG. 17

Gene A = somewhat less than Gene B




FIG. 18





FIG. 19

Step 1.



Step 2: PCR amplification of the labeled cDNA molecules of interest




FIG. 20




FIG. 21




FIG. 22




FIG. 23A




FIG. 23B




FIG. 24





FIG. 25

Step 1.



Step 2. Detection of the labeled cDNA molecules

FIG. 26

FIG. 27

### Step 3. Preparing to Implement the Model in Practice

Step 3. Add sequencing strand primer (forward) and PCR primer to extend 2 cycles (reverse).

				2710
				2720
				2730
				2740
				2750
				2760
				2770
				2780

FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32



FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 36





FIG. 36





FIG. 36




FIG. 37




FIG. 38





FIG. 39

A.



B.




FIG. 39





FIG. 39





FIG. 39





FIG. 40

## GAPDH in Liver




FIG. 41




FIG. 41




FIG. 42




FIG. 43

