

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 3405490 T3**

-
- (51) Int.Cl.: **C 07 K 16/28 (2006.01)** **A 61 P 35/00 (2006.01)** **C 07 K 16/30 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-01-10**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2021-10-20**
- (86) Europæisk ansøgning nr.: **17700749.9**
- (86) Europæisk indleveringsdag: **2017-01-10**
- (87) Den europæiske ansøgnings publiceringsdag: **2018-11-28**
- (86) International ansøgning nr.: **IB2017050109**
- (87) Internationalt publikationsnr.: **WO2017125831**
- (30) Prioritet: **2016-01-21 US 201662281543 P** **2016-12-08 US 201662431766 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Pfizer Inc., 235 East 42nd Street, New York, NY 10017, USA**
- (72) Opfinder: **CHOU, Joyce Ching, 767 Silkoak Way, Sunnyvale, CA 94086, USA**
WONG, Oi Kwan, 15 Arroyo View Circle, Belmont, CA 94002, USA
SASU, Barbra Johnson, 2125 Bryant Street, No 316, San Francisco, California 94110, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **MONO- OG BISPECIFIKKE ANTISTOFFER MOD EPIDERMAL VÆKSTFAKTORRECEPTOR VARIANT III OG CD3 OG ANVENDELSER DERAFT**
- (56) Fremdragne publikationer:
WO-A1-2013/185010
WO-A1-2014/153002
WO-A1-2015/006482
WO-A1-2017/021370
US-A1- 2012 189 630
BRYAN D. CHOI ET AL: "A novel bispecific antibody recruits T cells to eradicate tumors in the "immunologically privileged" central nervous system", ONCOIMMUNOLOGY, vol. 2, no. 4, 27 April 2013 (2013-04-27), page e23639, XP055306563, ISSN: 2162-4011, DOI: 10.4161/onci.23639
B. D. CHOI ET AL: "Systemic administration of a bispecific antibody targeting EGFRvIII successfully treats intracerebral glioma", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 110, no. 1, 17 December 2012 (2012-12-17), pages 270-275, XP055156305, ISSN: 0027-8424, DOI: 10.1073/pnas.1219817110
CHAN-JUAN SHEN ET AL: "Chimeric antigen receptor containing ICOS signaling domain mediates specific and efficient antitumor effect of T cells against EGFRvIII expressing glioma", JOURNAL OF HEMATOLOGY & ONCOLOGY, BIOMED CENTRAL LTD, LONDON UK, vol. 6, no. 1, 9 May 2013 (2013-05-09), page 33, XP021151586, ISSN: 1756-8722, DOI: 10.1186/1756-8722-6-33

Fortsættes ...

IAN M ZITRON ET AL: "Targeting and killing of glioblastoma with activated T cells armed with bispecific antibodies", BMC CANCER, BIOMED CENTRAL, LONDON, GB, vol. 13, no. 1, 22 February 2013 (2013-02-22), page 83, XP021141258, ISSN: 1471-2407, DOI: 10.1186/1471-2407-13-83

DescriptionField

[0001] The present invention relates to antibodies, e.g., full length antibodies or antigen binding fragments thereof, that specifically bind to Epidermal Growth Factor Receptor Variant III (EGFRvIII). The invention further relates to heteromultimeric antibodies (e.g., bispecific antibodies), antibody conjugates (e.g., antibody-drug-conjugates), compositions comprising the EGFRvIII antibodies, and methods for producing and purifying such antibodies. Their uses in diagnostics and therapeutics are also provided.

Background

[0002] EGFR variant III (EGFRvIII), a tumor specific mutant of EGFR, is a product of genomic rearrangement which is often associated with wild-type EGFR gene amplification. EGFRvIII is formed by an in-frame deletion of exons 2-7, leading to deletion of 267 amino acids with a glycine substitution at the junction. The truncated receptor loses its ability to bind ligands but acquires constitutive kinase activity. Interestingly, EGFRvIII frequently co-expresses with full length wild-type EGFR in the same tumor cells. Moreover, EGFRvIII expressing cells exhibit increased proliferation, invasion, angiogenesis and resistance to apoptosis.

[0003] EGFRvIII is most often found in glioblastoma multiforme (GBM). It is estimated that 25-35% of GBM carries this truncated receptor. Moreover, its expression often reflects a more aggressive phenotype and poor prognosis. Besides GBM, expression of EGFRvIII has also been reported in other solid tumors such as non-small cell lung cancer, head and neck cancer, breast cancer, ovarian cancer and prostate cancer. In contrast, EGFRvIII is not expressed in healthy tissues. The lack of expression in normal tissues makes EGFRvIII an ideal target for developing tumor specific targeted therapy. To date, there has not been any FDA approved monoclonal antibody (e.g., monospecific or bispecific) against EGFRvIII identified with high affinity, high specificity, and high potency in treating cancers such as GBM. Accordingly, there remains a need for antibodies (e.g., monospecific or bispecific) treating cancers such as GBM with improved efficacy and safety profile, and suitable for use with human patients.

[0004] WO2013/185010 A1 discloses bispecific antibody engaging molecules which have one arm that specifically engages a tumor cell which expresses the human EGFRvIII mutant protein on its surface, and a second arm that specifically engages T cell activation ligand CD3.

Summary

[0005] In an aspect of the invention, there is provided an antibody which specifically binds to EGFRvIII according to claim 1. Claim 1 relates to antibodies comprising the CDRs of the antibody clones listed in Table 8A and Table 8B.

[0006] The invention disclosed herein is directed to antibodies (e.g., monospecific or bispecific antibodies) and antibody conjugates that specifically bind to Epidermal Growth Factor Receptor Variant III (EGFRvIII).

[0007] The present disclosure provides an isolated antibody which specifically binds to EGFRvIII, wherein the antibody comprises (a) a heavy chain variable (VH) region comprising (i) a VH complementarity determining region one (CDR1) comprising the sequence shown in SEQ ID NO: 62, 63, 64, 74, 75, 76, 80, 81, 82, 88, 89, 90, 93, 94, 95, 99, 100, 101, 109, 110, 111, 115, 116, 117, 121, 122, 123, 132, 133, 134, 137, 138, 139, 143, 144, or 145; (ii) a VH CDR2 comprising the sequence shown in SEQ ID NO: 65, 66, 68, 69, 70, 71, 77, 78, 83, 84, 86, 87, 91, 92, 96, 97, 98, 102, 103, 105, 106, 112, 113, 118, 119, 124, 125, 127, 128, 130, 131, 135, 136, 140, 141, 146, 147, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, or 237; and (iii) a VH CDR3 comprising the sequence shown in SEQ ID NO: 67, 72, 73, 79, 85, 104, 107, 108, 114, 120, 126, 129, 142, 148, 219, 220, 221, 222, 223, or 236; and/or a light chain variable (VL) region comprising (i) a VL CDR1 comprising the sequence shown in SEQ ID NO: 149, 154, 156, 159, 162, 165, 166, 168, 169, 170, 171, 173, 174, 176, 178, 181, 182, 185, 187, 190, 192, 195, 198, 238, or 239; (ii) a VL CDR2 comprising the sequence shown in SEQ ID NO: 150, 152, 155, 157, 160, 163, 172, 175, 179, 183, 186, 188, 191, 193, 196, or 199; and (iii) a VL CDR3 comprising the sequence shown in SEQ ID NO: 151, 153, 158, 161, 164, 167, 177, 180, 184, 189, 194, 197, or 200.

[0008] Also provided is an isolated antibody which specifically binds to EGFRvIII, wherein the antibody comprises: a VH region comprising a VH CDR1, VH CDR2, and VH CDR3 of the VH sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 30, 32, 34, 35, 37, 39, 41, 43, 44, 46, 48, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 202, 203, 204, 205, 206, 207, 208, 209, 210, 214, 216, 217, or 218; and/or a VL region comprising VL CDR1, VL CDR2, and VL CDR3 of the VL sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 36, 38, 40, 42, 45, 47, 49, 51, 211, 212, 213, or 215. In some embodiments, the VH region as described herein comprises a variant with one or several conservative amino acid substitutions in residues that are not within a CDR and/or the VL region as described herein comprises a variant with one or several amino acid substitutions in amino acids that are not

within a CDR. For example, in some embodiments, the VH or VL region can comprise an amino acid sequence described above or a variant thereof with no more than 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 conservative substitutions in residues that are not within a CDR.

[0009] Provided is an isolated antibody which specifically binds to EGFRvIII, wherein the antibody comprises: a VH region comprising the sequence shown in SEQ ID NO: 5, 9, 11, 15, 30, 37, or 41; and/or a VL region comprising the sequence shown in SEQ ID NO: 6, 10, 12, 16, 31, 38, or 42. In some embodiments, the VH region comprises the sequence shown in SEQ ID NO: 9 and the VL region comprises the sequence shown in SEQ ID NO: 10. In some embodiments, the VH region comprises the sequence shown in SEQ ID NO: 11 and the VL region comprises the sequence shown in SEQ ID NO: 12. In some embodiments, the VH region comprises the sequence shown in SEQ ID NO: 15 and the VL region comprises the sequence shown in SEQ ID NO: 16. In some embodiments, the VH region comprises the sequence shown in SEQ ID NO: 30 and the VL region comprises the sequence shown in SEQ ID NO: 31. In some embodiments, the VH region comprises the sequence shown in SEQ ID NO: 37 and the VL region comprises the sequence shown in SEQ ID NO: 38. In some embodiments, the VH region comprises the sequence shown in SEQ ID NO: 41 and the VL region comprises the sequence shown in SEQ ID NO: 42.

[0010] In another aspect, provided is a bispecific antibody wherein the bispecific antibody is a full-length human antibody, comprising a first antibody variable domain of the bispecific antibody specifically binding to a target antigen (e.g., EGFRvIII), and comprising a second antibody variable domain of the bispecific antibody capable of recruiting the activity of a human immune effector cell by specifically binding to an effector antigen (e.g., Cluster of differentiation 3 (CD3)) located on the human immune effector cell. In some examples, the first antibody variable domain comprises a heavy chain variable (VH) region comprising a VH CDR1, VH CDR2, and VH CDR3 of the VH sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 30, 32, 34, 35, 37, 39, 41, 43, 44, 46, 48, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 202, 203, 204, 205, 206, 207, 208, 209, 210, 214, 216, 217, or 218; and/or a light chain variable (VL) region comprising VL CDR1, VL CDR2, and VL CDR3 of the VL sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 36, 38, 40, 42, 45, 47, 49, 51, 211, 212, 213, or 215. In some examples, the first antibody variable domain comprises (a) a heavy chain variable (VH) region comprising (i) a VH complementarity determining region one (CDR1) comprising the sequence shown in SEQ ID NO: 62, 63, 64, 74, 75, 76, 80, 81, 82, 88, 89, 90, 93, 94, 95, 99, 100, 101, 109, 110, 111, 115, 116, 117, 121, 122, 123, 132, 133, 134, 137, 138, 139, 143, 144, or 145; (ii) a VH CDR2 comprising the sequence shown in SEQ ID NO: 65, 66, 68, 69, 70, 71, 77, 78, 83, 84, 86, 87, 91, 92, 96, 97, 98, 102, 103, 105, 106, 112, 113, 118, 119, 124, 125, 127, 128, 130, 131, 135, 136, 140, 141, 146, 147, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, or 237; and (iii) a VH CDR3 comprising the sequence shown in SEQ ID NO: 67, 72, 73, 79, 85, 104, 107, 108, 114, 120, 126, 129, 142, 148, 219, 220, 221, 222, 223, or 236; and/or (b) a light chain variable (VL) region comprising (i) a VL CDR1 comprising the sequence shown in SEQ ID NO: 149, 154, 156, 159, 162, 165, 166, 168, 169, 170, 171, 173, 174, 176, 178, 181, 182, 185, 187, 190, 192, 195, 198, 238, or 239; (ii) a VL CDR2 comprising the sequence shown in SEQ ID NO: 150, 152, 155, 157, 160, 163, 172, 175, 179, 183, 186, 188, 191, 193, 196, or 199; and (iii) a VL CDR3 comprising the sequence shown in SEQ ID NO: 151, 153, 158, 161, 164, 167, 177, 180, 184, 189, 194, 197, or 200.

[0011] In some embodiments, the second antibody variable domain comprises the VH and/or VL region specific against CD3. For example, the second antibody variable domain comprises a heavy chain variable (VH) region comprising a VH CDR1, VH CDR2, and VH CDR3 of the VH sequence shown in SEQ ID NO: 240; and/or a light chain variable (VL) region comprising a VL CDR1, VL CDR2, and VL CDR3 of the VL sequence shown in SEQ ID NO: 241. In some embodiments, the second antibody variable domain comprises (a) a VH region comprising (i) a VH CDR1 comprising the sequence shown in SEQ ID NO: 244, 110, or 245; (ii) a VH CDR2 comprising the sequence shown in SEQ ID NO: 246 or 247; and (iii) a VH CDR3 comprising the sequence shown in SEQ ID NO: 248; and/or a VL region comprising (i) a VL CDR1 comprising the sequence shown in SEQ ID NO: 249; (ii) a VL CDR2 comprising the sequence shown in SEQ ID NO: 250; and (iii) a VL CDR3 comprising the sequence shown in SEQ ID NO: 251.

[0012] In some embodiments, the antibodies described herein comprise a constant region. In some embodiments, the antibodies described herein are of the human IgG1, IgG2 or IgG2Δa, IgG3, or IgG4 subclass. In some embodiments, the antibodies described herein comprise a glycosylated constant region. In some embodiments, the antibodies described herein comprise a constant region having decreased binding affinity to one or more human Fc gamma receptor(s).

[0013] In some embodiments, both the first and the second antibody variable domains of the bispecific antibody comprise amino acid modifications at positions 223, 225, and 228 (e.g., (C223E or C223R), (E225R), and (P228E or P228R)) in the hinge region and at position 409 or 368 (e.g., K409R or L368E (EU numbering scheme)) in the CH3 region of human IgG2 (SEQ ID NO: 290).

[0014] In some embodiments, both the first and the second antibody variable domains of the bispecific antibody comprise amino acid modifications at position 265 (e.g., D265A) of the human IgG2.

[0015] In some embodiments, both the first and the second antibody variable domains of the bispecific antibody comprise amino acid modifications at one or more of positions 265 (e.g., D265A), 330 (e.g., A330S), and 331 (e.g., P331S) of the human IgG2. In some embodiments, both the first and the second antibody variable domains of the

bispecific antibody comprise amino acid modifications at each of positions 265 (e.g., D265A), 330 (e.g., A330S), and 331 (e.g., P331S) of the human IgG2.

[0016] In another aspect, the invention provides an isolated antibody comprising an acyl donor glutamine-containing tag engineered at a specific site of the EGFRvIII antibody of the present invention.

[0017] In one variation, the invention provides an isolated antibody comprising an acyl donor glutamine-containing tag and an amino acid modification at position 222, 340, or 370 of the EGFRvIII antibody of the present invention. In some embodiments, the amino acid modification is a substitution from lysine to arginine.

[0018] In some embodiments, the EGFRvIII antibody of the present invention further comprises a linker.

[0019] In another aspect, the invention provides a conjugate of the EGFRvIII antibody as described herein, wherein the antibody is conjugated to an agent, wherein the agent is selected from the group consisting of a cytotoxic agent, an immunomodulating agent, an imaging agent, a therapeutic protein, a biopolymer, and an oligonucleotide. In some embodiments, the agent is a cytotoxic agent including, but not limited to, an anthracycline, an auristatin, a camptothecin, a combretastatin, a dolastatin, a duocarmycin, an enediyne, a geldanamycin, an indolino-benzodiazepine dimer, a maytansine, a puromycin, a pyrrolobenzodiazepine dimer, a taxane, a vinca alkaloid, a tubulysin, a hemiasterlin, a spliceostatin, a pladienolide, and stereoisomers, isosteres, analogs, or derivatives thereof. For example, the cytotoxic agent is MMAD (Monomethyl Auristatin D), 0101 (2-methylalanyl-*N*-[(3*R*,4*S*,5*S*)-3-methoxy-1-*1*(2*S*)-2-[(1*R*,2*R*)-1-methoxy-2-methyl-3-oxo-3-[[[(1*S*)-2-phenyl-1-(1,3-thiazol-2-yl)ethyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-*N*-methyl-L-valinamide), 3377 (N,2-dimethylalanyl-*N*-{[(1*S*,2*R*)-4-[(2*S*)-2-[(1*R*,2*R*)-3-[(1*S*)-1-carboxyl-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-2-methoxy-1-[(1*S*)-1-methylpropyl]-4-oxobutyl]-*N*-methyl-L-valinamide), 0131 (2-methyl-L-prolyl-*N*-[(3*R*,4*S*,5*S*)-1-[(2*S*)-2-[(1*R*,2*R*)-3-[(1*S*)-1-carboxyl-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-*N*-methyl-L-valinamide), or 0121 (2-methyl-L-prolyl-*N*-[(3*R*,4*S*,5*S*)-1-[(2*S*)-2-[(1*R*,2*R*)-3-[(2*S*)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-*N*-methyl-L-valinamide).

[0020] In some embodiments, the present invention provides a conjugate comprising the formula: antibody-(acyl donor glutamine-containing tag)-(linker)-(cytotoxic agent).

[0021] In other embodiments, the invention provides pharmaceutical compositions comprising any of the antibodies or antibody conjugates of the invention.

[0022] The invention also provides cell lines that recombinantly produce any of the antibodies of the invention.

[0023] The invention also provides nucleic acids encoding any of the antibodies of the invention. The invention also provides nucleic acids encoding a heavy chain variable region and/or a light chain variable region of any of the antibodies of the invention.

[0024] The invention also provides kits comprising an effective amount of any of the antibodies or antibody conjugates of the invention.

[0025] The invention also provides the isolated antibodies, bispecific antibodies, or antibody conjugates of the invention for use in methods of treating subjects in need thereof, wherein the methods comprise administering said antibodies to said subject.

[0026] Also provided are the antibodies of the invention for use in methods of treating a condition associated with malignant cells expressing EGFRvIII in a subject comprising administering to a subject in need thereof an effective amount of a pharmaceutical composition comprising the antibodies or antibody conjugates as described herein. In some embodiments, the condition is a cancer. In some embodiments, the cancer is an EGFRvIII related cancer (e.g., any cancer with EGFRvIII expression) selected from the group consisting of glioblastoma multiform, anaplastic astrocytoma, giant cell glioblastoma, gliosarcoma, anaplastic oligodendroglioma, anaplastic ependymoma, anaplastic oligoastrocytoma, choroid plexus carcinoma, anaplastic ganglioglioma, pineoblastoma, pineocytoma, meningioma, medulloepithelioma, ependymoblastoma, medulloblastoma, supraentorial primitive neuroectodermal tumor, atypical teratoid/rhabdoid tumor, head and neck cancer, non-small cell lung cancer, breast cancer, ovarian cancer, prostate cancer, medulloblastoma, colorectal cancer, anal cancer, gastric cancer, thyroid cancer, mesothelioma, uterine cancer, and bladder cancer.

[0027] In another aspect, the invention provides the antibodies of the invention for use in a method of inhibiting tumor growth or progression in a subject who has malignant cells expressing EGFRvIII, comprising administering to the subject in need thereof an effective amount of a pharmaceutical composition comprising the isolated antibodies, bispecific antibodies, or antibody conjugates, as described herein.

[0028] In another aspect, the invention provides the antibodies of the invention for use in a method inhibiting metastasis of malignant cells expressing EGFRvIII in a subject, comprising administering to the subject in need thereof an effective amount of the pharmaceutical composition comprising the isolated antibodies, bispecific antibodies, or antibody conjugates, as described herein.

[0029] In another aspect, the invention provides the antibodies of the invention for use in a method inducing tumor regression in a subject who has malignant cells expressing EGFRvIII, comprising administering to the subject in need thereof an effective amount of the pharmaceutical composition of a pharmaceutical composition comprising the isolated antibodies, bispecific antibodies, or antibody conjugates, as described herein.

Brief Description of the Figures/Drawings**[0030]**

FIGs. 1A, 1B, and 1C show examples of FACS binding histograms of three EGFRvIII antibodies: mAb 42G9 (FIG. 1A), 32A10 (FIG. 1B) and 32G8 (FIG. 1C), to the three F98 cell lines: F98 (EGFR negative), F98-EGFRwt, and F98-EGFRvIII. The X-axis is fluorescence intensity; the Y-axis is percentage of maximum / normalized to mode.

FIGs. 2A, 2B and 2C depict histograms showing the expression of wild-type EGFR and EGFRvIII in GBM cell lines as measured by flow cytometry: LN229-EGFRvIII (FIG. 2A), LN18-EGFRvIII (FIG. 2B) and DKMG (FIG. 2C). EGFRvIII was detected with mAb 42G9 and EGFRwt was detected with an EGFR wild-type specific mAb. The X-axis is fluorescence intensity; the Y-axis is percentage of maximum / normalized to mode.

FIGs. 3A and 3B show graphs demonstrating the cytotoxicity of three EGFRvIII-CD3 bispecific antibodies in EGFRvIII transduced LN18-EGFRvIII (FIG. 3A) and parental LN18 (FIG. 3B) cells.

FIGs. 4A and 4B show graphs demonstrating the cytotoxicity of three EGFRvIII-CD3 bispecific antibodies in EGFRvIII transduced LN229-EGFRvIII (FIG. 4A) and parental LN229 (FIG. 4B) cells.

FIG. 5 shows a graph demonstrating the cytotoxicity of three EGFRvIII-CD3 bispecific antibodies in DKMG cells, which express endogenous EGFRvIII and EGFR wild-type proteins.

FIG. 6 shows a graph illustrating the *in vivo* anti-tumor activity of EGFRvIII-CD3 bispecific antibodies in a subcutaneous model of LN229-EGFRvIII GBM cell line.

Detailed Description

[0031] The invention disclosed herein provides antibodies (e.g., monospecific or bispecific) and antibody conjugates that specifically bind to EGFRvIII (e.g., human EGFRvIII). The invention also provides polynucleotides encoding these antibodies, compositions comprising these antibodies and antibody conjugates, and methods of making and using these antibodies and antibody conjugates. The present disclosure also provides methods for treating a condition associated with EGFRvIII-mediated pathologies in a subject, such as cancer.

General Techniques

[0032] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, immunology, virology, monoclonal antibody generation and engineering, which are within the skill of the art. Such techniques are explained fully in the literature, such as, Molecular Cloning: A Laboratory Manual, second edition (Sambrook et al., 1989) Cold Spring Harbor Press; Oligonucleotide Synthesis (M.J. Gait, ed., 1984); Methods in Molecular Biology, Humana Press; Cell Biology: A Laboratory Notebook (J.E. Cellis, ed., 1998) Academic Press; Animal Cell Culture (R.I. Freshney, ed., 1987); Introduction to Cell and Tissue Culture (J.P. Mather and P.E. Roberts, 1998) Plenum Press; Cell and Tissue Culture: Laboratory Procedures (A. Doyle, J.B. Griffiths, and D.G. Newell, eds., 1993-1998) J. Wiley and Sons; Methods in Enzymology (Academic Press, Inc.); Handbook of Experimental Immunology (D.M. Weir and C.C. Blackwell, eds.); Gene Transfer Vectors for Mammalian Cells (J.M. Miller and M.P. Calos, eds., 1987); Current Protocols in Molecular Biology (F.M. Ausubel et al., eds., 1987); PCR: The Polymerase Chain Reaction, (Mullis et al., eds., 1994); Current Protocols in Immunology (J.E. Coligan et al., eds., 1991); Short Protocols in Molecular Biology (Wiley and Sons, 1999); Immunobiology (C.A. Janeway and P. Travers, 1997); Antibodies (P. Finch, 1997); Antibodies: a practical approach (D. Catty, ed., IRL Press, 1988-1989); Monoclonal antibodies: a practical approach (P. Shepherd and C. Dean, eds., Oxford University Press, 2000); Using antibodies: a laboratory manual (E. Harlow and D. Lane (Cold Spring Harbor Laboratory Press, 1999); The Antibodies (M. Zanetti and J.D. Capra, eds., Harwood Academic Publishers, 1995).

Definitions

[0033] An "antibody" is an immunoglobulin molecule capable of specific binding to a target, such as a carbohydrate, polynucleotide, lipid, polypeptide, etc., through at least one antigen recognition site, located in the variable region of the immunoglobulin molecule. As used herein, the term encompasses not only intact polyclonal or monoclonal antibodies, but also antigen binding fragments thereof (such as Fab, Fab', F(ab')₂, Fv), single chain (ScFv) and domain antibodies (including, for example, shark and camelid antibodies), and fusion proteins comprising an antibody, and any other modified configuration of the immunoglobulin molecule that comprises an antigen recognition site. An antibody includes an antibody of any class, such as IgG, IgA, or IgM (or sub-class thereof), and the antibody need not be of any particular class. Depending on the antibody amino acid sequence of the constant region of its heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and

several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2. The heavy-chain constant regions that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

[0034] The term "antigen binding fragment" or "antigen binding portion" of an antibody, as used herein, refers to one or more fragments of an intact antibody that retain the ability to specifically bind to a given antigen (e.g., EGFRvIII). Antigen binding functions of an antibody can be performed by fragments of an intact antibody. Examples of binding fragments encompassed within the term "antigen binding fragment" of an antibody include Fab; Fab'; F(ab')₂; an Fd fragment consisting of the VH and CH1 domains; an Fv fragment consisting of the VL and VH domains of a single arm of an antibody; a single domain antibody (dAb) fragment (Ward et al., Nature 341:544-546, 1989), and an isolated complementarity determining region (CDR).

[0035] An antibody, an antibody conjugate, or a polypeptide that "preferentially binds" or "specifically binds" (used interchangeably herein) to a target (e.g., EGFRvIII protein) is a term well understood in the art, and methods to determine such specific or preferential binding are also well known in the art. A molecule is said to exhibit "specific binding" or "preferential binding" if it reacts or associates more frequently, more rapidly, with greater duration and/or with greater affinity with a particular cell or substance than it does with alternative cells or substances. An antibody "specifically binds" or "preferentially binds" to a target if it binds with greater affinity, avidity, more readily, and/or with greater duration than it binds to other substances. For example, an antibody that specifically or preferentially binds to an EGFRvIII epitope is an antibody that binds this epitope with greater affinity, avidity, more readily, and/or with greater duration than it binds to other EGFRvIII epitopes or non-EGFRvIII epitopes. It is also understood that by reading this definition, for example, an antibody (or moiety or epitope) that specifically or preferentially binds to a first target may or may not specifically or preferentially bind to a second target. As such, "specific binding" or "preferential binding" does not necessarily require (although it can include) exclusive binding. Generally, but not necessarily, reference to binding means preferential binding.

[0036] A "variable region" of an antibody refers to the variable region of the antibody light chain or the variable region of the antibody heavy chain, either alone or in combination. As known in the art, the variable regions of the heavy and light chain each consist of four framework regions (FR) connected by three complementarity determining regions (CDRs) also known as hypervariable regions. The CDRs in each chain are held together in close proximity by the FRs and, with the CDRs from the other chain, contribute to the formation of the antigen binding site of antibodies. There are at least two techniques for determining CDRs: (1) an approach based on cross-species sequence variability (i.e., Kabat et al. Sequences of Proteins of Immunological Interest, (5th ed., 1991, National Institutes of Health, Bethesda MD)); and (2) an approach based on crystallographic studies of antigen-antibody complexes (Al-lazikani et al., 1997, J. Molec. Biol. 273:927-948). As used herein, a CDR may refer to CDRs defined by either approach or by a combination of both approaches.

[0037] A "CDR" of a variable domain are amino acid residues within the variable region that are identified in accordance with the definitions of the Kabat, Chothia, the accumulation of both Kabat and Chothia, AbM, contact, and/or conformational definitions or any method of CDR determination well known in the art. Antibody CDRs may be identified as the hypervariable regions originally defined by Kabat et al. See, e.g., Kabat et al., 1992, Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, NIH, Washington D.C. The positions of the CDRs may also be identified as the structural loop structures originally described by Chothia and others. See, e.g., Chothia et al., Nature 342:877-883, 1989. Other approaches to CDR identification include the "AbM definition," which is a compromise between Kabat and Chothia and is derived using Oxford Molecular's AbM antibody modeling software (now Accelrys®), or the "contact definition" of CDRs based on observed antigen contacts, set forth in MacCallum et al., J. Mol. Biol., 262:732-745, 1996. In another approach, referred to herein as the "conformational definition" of CDRs, the positions of the CDRs may be identified as the residues that make enthalpic contributions to antigen binding. See, e.g., Makabe et al., Journal of Biological Chemistry, 283:1156-1166, 2008. Still other CDR boundary definitions may not strictly follow one of the above approaches, but will nonetheless overlap with at least a portion of the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. As used herein, a CDR may refer to CDRs defined by any approach known in the art, including combinations of approaches. The methods used herein may utilize CDRs defined according to any of these approaches. For any given embodiment containing more than one CDR, the CDRs may be defined in accordance with any of Kabat, Chothia, extended, AbM, contact, and/or conformational definitions.

[0038] As used herein, "monoclonal antibody" refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the

antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler and Milstein, Nature 256:495, 1975, or may be made by recombinant DNA methods such as described in U.S. Pat. No. 4,816,567. The monoclonal antibodies may also be isolated from phage libraries generated using the techniques described in McCafferty et al., Nature 348:552-554, 1990, for example.

[0039] As used herein, "humanized" antibody refers to forms of non-human (e.g. murine) antibodies that are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. Preferably, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences, but are included to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Preferred are antibodies having Fc regions modified as described in WO 99/58572. Other forms of humanized antibodies have one or more CDRs (CDR L1, CDR L2, CDR L3, CDR H1, CDR H2, or CDR H3) which are altered with respect to the original antibody, which are also termed one or more CDRs "derived from" one or more CDRs from the original antibody.

[0040] As used herein, "human antibody" means an antibody having an amino acid sequence corresponding to that of an antibody produced by a human and/or which has been made using any of the techniques for making human antibodies known to those skilled in the art or disclosed herein. This definition of a human antibody includes antibodies comprising at least one human heavy chain polypeptide or at least one human light chain polypeptide. One such example is an antibody comprising murine light chain and human heavy chain polypeptides. Human antibodies can be produced using various techniques known in the art. In one embodiment, the human antibody is selected from a phage library, where that phage library expresses human antibodies (Vaughan et al., Nature Biotechnology, 14:309-314, 1996; Sheets et al., Proc. Natl. Acad. Sci. (USA) 95:6157-6162, 1998; Hoogenboom and Winter, J. Mol. Biol., 227:381, 1991; Marks et al., J. Mol. Biol., 222:581, 1991). Human antibodies can also be made by immunization of animals into which human immunoglobulin loci have been transgenically introduced in place of the endogenous loci, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. This approach is described in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016. Alternatively, the human antibody may be prepared by immortalizing human B lymphocytes that produce an antibody directed against a target antigen (such B lymphocytes may be recovered from an individual or from single cell cloning of the cDNA, or may have been immunized *in vitro*). See, e.g., Cole et al. Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77, 1985; Boerner et al., J. Immunol., 147 (1):86-95, 1991; and U.S. Pat. No. 5,750,373.

[0041] The term "chimeric antibody" is intended to refer to antibodies in which the variable region sequences are derived from one species and the constant region sequences are derived from another species, such as an antibody in which the variable region sequences are derived from a mouse antibody and the constant region sequences are derived from a human antibody.

[0042] The terms "polypeptide", "oligopeptide", "peptide" and "protein" are used interchangeably herein to refer to chains of amino acids of any length. For example, the chain may be relatively short (e.g., 10-100 amino acids), or longer. The chain may be linear or branched, it may comprise modified amino acids, and/or may be interrupted by non-amino acids. The terms also encompass an amino acid chain that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art. It is understood that the polypeptides can occur as single chains or associated chains.

[0043] A "monovalent antibody" comprises one antigen binding site per molecule (e.g., IgG or Fab). In some instances, a monovalent antibody can have more than one antigen binding sites, but the binding sites are from different antigens.

[0044] A "monospecific antibody" comprises two identical antigen binding sites per molecule (e.g. IgG) such that the two binding sites bind identical epitope on the antigen. Thus, they compete with each other on binding to one antigen molecule. Most antibodies found in nature are monospecific. In some instances, a monospecific antibody can also be a monovalent antibody (e.g. Fab)

[0045] A "bivalent antibody" comprises two antigen binding sites per molecule (e.g., IgG). In some instances, the two binding sites have the same antigen specificities. However, bivalent antibodies may be bispecific.

[0046] A "bispecific" or "dual-specific" is a hybrid antibody having two different antigen binding sites. The two antigen

binding sites of a bispecific antibody bind to two different epitopes, which may reside on the same or different protein targets.

[0047] A "bifunctional" antibody is an antibody having identical antigen binding sites (i.e., identical amino acid sequences) in the two arms but each binding site can recognize two different antigens.

[0048] A "heteromultimer", "heteromultimeric complex", or "heteromultimeric polypeptide" is a molecule comprising at least a first polypeptide and a second polypeptide, wherein the second polypeptide differs in amino acid sequence from the first polypeptide by at least one amino acid residue. The heteromultimer can comprise a "heterodimer" formed by the first and second polypeptide or can form higher order tertiary structures where polypeptides in addition to the first and second polypeptide are present.

[0049] A "heterodimer", "heterodimeric protein", "heterodimeric complex", or "heteromultimeric polypeptide" is a molecule comprising a first polypeptide and a second polypeptide, wherein the second polypeptide differs in amino acid sequence from the first polypeptide by at least one amino acid residue.

[0050] The "hinge region", "hinge sequence", and variations thereof, as used herein, includes the meaning known in the art, which is illustrated in, for example, Janeway et al., *ImmunoBiology: the immune system in health and disease*, (Elsevier Science Ltd., NY) (4th ed., 1999); Bloom et al., *Protein Science* (1997), 6:407-415; Humphreys et al., *J. Immunol. Methods* (1997), 209:193-202.

[0051] The "immunoglobulin-like hinge region", "immunoglobulin-like hinge sequence", and variations thereof, as used herein, refer to the hinge region and hinge sequence of an immunoglobulin-like or an antibody-like molecule (e.g., immunoadhesins). In some embodiments, the immunoglobulin-like hinge region can be from or derived from any IgG1, IgG2, IgG3, or IgG4 subtype, or from IgA, IgE, IgD or IgM, including chimeric forms thereof, e.g., a chimeric IgG1/2 hinge region.

[0052] The term "immune effector cell" or "effector cell" as used herein refers to a cell within the natural repertoire of cells in the human immune system which can be activated to affect the viability of a target cell. The viability of a target cell can include cell survival, proliferation, and/or ability to interact with other cells.

[0053] Antibodies of the invention can be produced using techniques well known in the art, e.g., recombinant technologies, phage display technologies, synthetic technologies or combinations of such technologies or other technologies readily known in the art (see, for example, Jayasena, S.D., *Clin. Chem.*, 45: 1628-50, 1999 and Fellouse, F.A., et al, *J. Mol. Biol.*, 373(4):924-40, 2007).

[0054] As known in the art, "polynucleotide," or "nucleic acid," as used interchangeably herein, refer to chains of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a chain by DNA or RNA polymerase. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and their analogs. If present, modification to the nucleotide structure may be imparted before or after assembly of the chain. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. Other types of modifications include, for example, "caps", substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, for example, proteins (e.g., nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide(s). Further, any of the hydroxyl groups ordinarily present in the sugars may be replaced, for example, by phosphonate groups, phosphate groups, protected by standard protecting groups, or activated to prepare additional linkages to additional nucleotides, or may be conjugated to solid supports. The 5' and 3' terminal OH can be phosphorylated or substituted with amines or organic capping group moieties of from 1 to 20 carbon atoms. Other hydroxyls may also be derivatized to standard protecting groups. Polynucleotides can also contain analogous forms of ribose or deoxyribose sugars that are generally known in the art, including, for example, 2'-O-methyl-, 2'-O-allyl, 2'-fluoro- or 2'-azido-ribose, carbocyclic sugar analogs, alpha- or beta-anomeric sugars, epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, sedoheptuloses, acyclic analogs and abasic nucleoside analogs such as methyl riboside. One or more phosphodiester linkages may be replaced by alternative linking groups. These alternative linking groups include, but are not limited to, embodiments wherein phosphate is replaced by P(O)S ("thioate"), P(S)S ("dithioate"), (O)NR₂ ("amidate"), P(O)R, P(O)OR', CO or CH₂ ("formacetal"), in which each R or R' is independently H or substituted or unsubstituted alkyl (1-20 C) optionally containing an ether (-O-) linkage, aryl, alkenyl, cycloalkyl, cycloalkenyl or araldyl. Not all linkages in a polynucleotide need be identical. The preceding description applies to all polynucleotides referred to herein, including RNA and DNA.

[0055] As known in the art, a "constant region" of an antibody refers to the constant region of the antibody light chain or the constant region of the antibody heavy chain, either alone or in combination.

[0056] As used herein, "substantially pure" refers to material which is at least 50% pure (i.e., free from contaminants),

more preferably, at least 90% pure, more preferably, at least 95% pure, yet more preferably, at least 98% pure, and most preferably, at least 99% pure.

[0057] A "host cell" includes an individual cell or cell culture that can be or has been a recipient for vector(s) for incorporation of polynucleotide inserts. Host cells include progeny of a single host cell, and the progeny may not necessarily be completely identical (in morphology or in genomic DNA complement) to the original parent cell due to natural, accidental, or deliberate mutation. A host cell includes cells transfected in vivo with a polynucleotide(s) of this invention.

[0058] As known in the art, the term "Fc region" is used to define a C-terminal region of an immunoglobulin heavy chain. The "Fc region" may be a native sequence Fc region or a variant Fc region. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The numbering of the residues in the Fc region is that of the EU index as in Kabat. Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., 1991. The Fc region of an immunoglobulin generally comprises two constant regions, CH2 and CH3.

[0059] As used in the art, "Fc receptor" and "FcR" describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc γ RII receptors include Fc γ RIIA (an "activating receptor") and Fc γ RIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. FcRs are reviewed in Ravetch and Kinet, *Ann. Rev. Immunol.*, 9:457-92, 1991; Capel et al., *Immunomethods*, 4:25-34, 1994; and de Haas et al., *J. Lab. Clin. Med.*, 126:330-41, 1995. "FcR" also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., *J. Immunol.*, 117:587, 1976; and Kim et al., *J. Immunol.*, 24:249, 1994).

[0060] The term "compete", as used herein with regard to an antibody, means that a first antibody, or an antigen binding fragment (or portion) thereof, binds to an epitope in a manner sufficiently similar to the binding of a second antibody, or an antigen binding portion thereof, such that the result of binding of the first antibody with its cognate epitope is detectably decreased in the presence of the second antibody compared to the binding of the first antibody in the absence of the second antibody. The alternative, where the binding of the second antibody to its epitope is also detectably decreased in the presence of the first antibody, can, but need not be the case. That is, a first antibody can inhibit the binding of a second antibody to its epitope without that second antibody inhibiting the binding of the first antibody to its respective epitope. However, where each antibody detectably inhibits the binding of the other antibody with its cognate epitope or ligand, whether to the same, greater, or lesser extent, the antibodies are said to "cross-compete" with each other for binding of their respective epitope(s). Both competing and cross-competing antibodies are encompassed by the present disclosure. Regardless of the mechanism by which such competition or cross-competition occurs (e.g., steric hindrance, conformational change, or binding to a common epitope, or portion thereof), the skilled artisan would appreciate, based upon the teachings provided herein, that such competing and/or cross-competing antibodies are encompassed and can be useful for the methods disclosed herein.

[0061] A "functional Fc region" possesses at least one effector function of a native sequence Fc region. Exemplary "effector functions" include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity; phagocytosis; down-regulation of cell surface receptors (e.g. B cell receptor), etc. Such effector functions generally require the Fc region to be combined with a binding domain (e.g. an antibody variable domain) and can be assessed using various assays known in the art for evaluating such antibody effector functions.

[0062] A "native sequence Fc region" comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. A "variant Fc region" comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification, yet retains at least one effector function of the native sequence Fc region. In some embodiments, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, e.g. from about one to about ten amino acid substitutions, and preferably, from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% sequence identity with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably, at least about 90% sequence identity therewith, more preferably, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% sequence identity therewith.

[0063] The term "effector function" refers to the biological activities attributable to the Fc region of an antibody. Examples of antibody effector functions include, but are not limited to, antibody-dependent cell-mediated cytotoxicity (ADCC), Fc receptor binding, complement dependent cytotoxicity (CDC), phagocytosis, C1q binding, and down regulation of cell surface receptors (e.g., B cell receptor; BCR). See, e.g., U.S. Pat No. 6,737,056. Such effector functions generally require the Fc region to be combined with a binding domain (e.g., an antibody variable domain) and can be assessed using various assays known in the art for evaluating such antibody effector functions. An exemplary measurement of effector function is through Fc γ 3 and/or C1q binding.

[0064] As used herein "antibody-dependent cell-mediated cytotoxicity" or "ADCC" refers to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (e.g. natural killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. ADCC activity of a molecule of interest can be assessed using an *in vitro* ADCC assay, such as that described in U.S. Patent No. 5,500,362 or 5,821,337. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and NK cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in an animal model such as that disclosed in Clynes et al., 1998, PNAS (USA), 95:652-656.

[0065] "Complement dependent cytotoxicity" or "CDC" refers to the lysing of a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (e.g. an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, e.g. as described in Gazzano-Santoro et al., J. Immunol. Methods, 202: 163 (1996), may be performed.

[0066] As used herein, "treatment" is an approach for obtaining beneficial or desired clinical results. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, one or more of the following: reducing the proliferation of (or destroying) neoplastic or cancerous cells, inhibiting metastasis of neoplastic cells, shrinking or decreasing the size of EGFRvIII expressing tumor, remission of an EGFRvIII associated disease (e.g., cancer), decreasing symptoms resulting from an EGFRvIII associated disease (e.g., cancer), increasing the quality of life of those suffering from an EGFRvIII associated disease (e.g., cancer), decreasing the dose of other medications required to treat an EGFRvIII associated disease (e.g., cancer), delaying the progression of an EGFRvIII associated disease (e.g., cancer), curing an EGFRvIII associated disease (e.g., cancer), and/or prolong survival of patients having an EGFRvIII associated disease (e.g., cancer).

[0067] "Ameliorating" means a lessening or improvement of one or more symptoms as compared to not administering an EGFRvIII antibody (monospecific or bispecific). "Ameliorating" also includes shortening or reduction in duration of a symptom.

[0068] As used herein, an "effective dosage" or "effective amount" of drug, compound, or pharmaceutical composition is an amount sufficient to effect any one or more beneficial or desired results. For prophylactic use, beneficial or desired results include eliminating or reducing the risk, lessening the severity, or delaying the outset of the disease, including biochemical, histological and/or behavioral symptoms of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. For therapeutic use, beneficial or desired results include clinical results such as reducing incidence or amelioration of one or more symptoms of various EGFRvIII associated diseases or conditions (such as for example multiple myeloma), decreasing the dose of other medications required to treat the disease, enhancing the effect of another medication, and/or delaying the progression of the EGFRvIII associated disease of patients. An effective dosage can be administered in one or more administrations. For purposes of this invention, an effective dosage of drug, compound, or pharmaceutical composition is an amount sufficient to accomplish prophylactic or therapeutic treatment either directly or indirectly. As is understood in the clinical context, an effective dosage of a drug, compound, or pharmaceutical composition may or may not be achieved in conjunction with another drug, compound, or pharmaceutical composition. Thus, an "effective dosage" may be considered in the context of administering one or more therapeutic agents, and a single agent may be considered to be given in an effective amount if, in conjunction with one or more other agents, a desirable result may be or is achieved.

[0069] An "individual" or a "subject" is a mammal, more preferably, a human. Mammals also include, but are not limited to primates, horses, dogs, cats, mice and rats.

[0070] As used herein, "vector" means a construct, which is capable of delivering, and, preferably, expressing, one or more gene(s) or sequence(s) of interest in a host cell. Examples of vectors include, but are not limited to, viral vectors, naked DNA or RNA expression vectors, plasmid, cosmid or phage vectors, DNA or RNA expression vectors associated with cationic condensing agents, DNA or RNA expression vectors encapsulated in liposomes, and certain eukaryotic cells, such as producer cells.

[0071] As used herein, "expression control sequence" means a nucleic acid sequence that directs transcription of a nucleic acid. An expression control sequence can be a promoter, such as a constitutive or an inducible promoter, or an enhancer. The expression control sequence is operably linked to the nucleic acid sequence to be transcribed.

[0072] As used herein, "pharmaceutically acceptable carrier" or "pharmaceutical acceptable excipient" includes any material which, when combined with an active ingredient, allows the ingredient to retain biological activity and is non-reactive with the subject's immune system. Examples include, but are not limited to, any of the standard pharmaceutical carriers such as a phosphate buffered saline solution, water, emulsions such as oil/water emulsion, and various types of wetting agents. Preferred diluents for aerosol or parenteral administration are phosphate buffered saline (PBS) or normal (0.9%) saline. Compositions comprising such carriers are formulated by well known conventional methods (see, for example, Remington's Pharmaceutical Sciences, 18th edition, A. Gennaro, ed., Mack Publishing Co., Easton, PA, 1990; and Remington, The Science and Practice of Pharmacy 21st Ed. Mack Publishing, 2005).

[0073] The term "acyl donor glutamine-containing tag" or "glutamine tag" as used herein refers to a polypeptide or a protein containing one or more Gln residue(s) that acts as a transglutaminase amine acceptor. See, e.g., WO2012059882

and WO2015015448.

[0074] The term " k_{on} " or " k_a ", as used herein, refers to the rate constant for association of an antibody to an antigen. Specifically, the rate constants (k_{on}/k_a and k_{off}/k_d) and equilibrium dissociation constants are measured using whole antibody (i.e. bivalent) and monomeric EGFRvIII proteins (e.g., Histidine-tagged EGFRvIII fusion protein).

[0075] The term " k_{off} " or " k_d ", as used herein, refers to the rate constant for dissociation of an antibody from the antibody/antigen complex.

[0076] The term " K_D ", as used herein, refers to the equilibrium dissociation constant of an antibody-antigen interaction.

[0077] Reference to "about" a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter per se. For example, description referring to "about X" includes description of "X." Numeric ranges are inclusive of the numbers defining the range. Generally speaking, the term "about" refers to the indicated value of the variable and to all values of the variable that are within the experimental error of the indicated value (e.g. within the 95% confidence interval for the mean) or within 10 percent of the indicated value, whichever is greater. Where the term "about" is used within the context of a time period (years, months, weeks, days etc.), the term "about" means that period of time plus or minus one amount of the next subordinate time period (e.g. about 1 year means 11-13 months; about 6 months means 6 months plus or minus 1 week; about 1 week means 6-8 days; etc.), or within 10 per cent of the indicated value, whichever is greater.

[0078] It is understood that wherever embodiments are described herein with the language "comprising," otherwise analogous embodiments described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0079] Where aspects or embodiments of the invention are described in terms of a Markush group or other grouping of alternatives, the present invention encompasses not only the entire group listed as a whole, but each member of the group individually and all possible subgroups of the main group, but also the main group absent one or more of the group members. The present invention also envisages the explicit exclusion of one or more of any of the group members in the claimed invention.

[0080] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In case of conflict, the present specification, including definitions, will control. Throughout this specification and claims, the word "comprise," or variations such as "comprises" or "comprising" will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers. Unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular.

[0081] Exemplary methods and materials are described herein, although methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention. The materials, methods, and examples are illustrative only and not intended to be limiting.

EGFRvIII Antibodies and Methods of Making Thereof

[0082] The present invention provides an antibody that binds to EGFRvIII [e.g., human EGFRvIII (e.g., accession number: P00533 Feature Identifier VAR_066493, or GenBank Accession No. AJN69267; mrpsgtagaallallaalcp asraleekkgnvyvtdhgscvracgadsyemeedgvrkckkcegcprkvcngigefkdsisi natnikhfkncsisgdhlpvafgrdsfthtppldpqeldilk-tvkeitgflliqawpenrt dlhafenleirgrtkqhggfslavvslhnlsglrlsikeisdgdvsiisgnknlcyanntinwkk lfgtsgqktkiisngensckatgqvchalcspegcwgpeprdcvscmvsgrecvdkcnlle geprefvenseciqchpeclpqamnitctgrgpdnciqcahy-idgphvcvtcpagvmgenntlv wkyadaghvchlhpnctygtgpglegcptngkipiatgmvgalllllvalgiglmfrrr hivrktlrlrlqerelvepltpsgaeapnqallrilketefkkikvlvgsgafgtvykgwipeg ekvipvaikelreatspankeildeayvmasvndphvcrllgicltstvtql-itqlmpfgcll dyvrehkdngsqyllnwcvqjakgmnyledrrlvhrdlaamvlvktphvktdfglakllg aeekeyhaeggkvpikwmalesilhriythqsd-vwsygvvtwelmtfgskpydgipaseissil ekgerlpqppictidvymimvkcwmidadsrpkfrelliefskmardpqrylvigqdermhpls ptdsnfyralmdeedmddvdadeylipqqgffsspstsrtpllsslsatsnnstvacidmgl qscpikedsflqrysdpdgtaldtsiddtflpvpeyinqsvpkrp-agsvqnpvyhnpplnpap srdphyqdphstavgnppeylntvqptcvnstfdspahwaqkgshqisldnpyqqdfffpeakp ngifkgstaenaeyl-rvapqssefiga(SEQ ID NO: 201))] and characterized by any one or more of the following characteristics: (a) decrease or downregulate the protein expression of EGFRvIII; (b) treat, prevent, ameliorate one or more symptoms of a condition associated with malignant cells expressing EGFRvIII in a subject (e.g., cancer such as glioblastoma multiforme); (c) inhibit tumor growth or progression in a subject (who has a malignant tumor expressing EGFRvIII); (d) inhibit metastasis of cancer (malignant) cells expressing EGFRvIII in a subject (who has one or more malignant cells expressing EGFRvIII); (e) induce regression (e.g., long-term regression) of a tumor expressing EGFRvIII; (f) exert cytotoxic activity in malignant cells expressing EGFRvIII; (g) block EGFRvIII interaction with other yet to be identified factors; and/or (h) induce bystander effect that kill or inhibit growth of non-EGFRvIII expressing malignant cells in the vicinity.

[0083] In one aspect, provided is an isolated antibody which specifically binds to EGFRvIII, wherein the antibody comprises (a) a heavy chain variable (VH) region comprising (i) a VH complementarity determining region one (CDR1) comprising the sequence shown in SEQ ID NO: 62, 63, 64, 74, 75, 76, 80, 81, 82, 88, 89, 90, 93, 94, 95, 99, 100, 101, 109, 110, 111, 115, 116, 117, 121, 122, 123, 132, 133, 134, 137, 138, 139, 143, 144, or 145; (ii) a VH CDR2 comprising

the sequence shown in SEQ ID NO: 65, 66, 68, 69, 70, 71, 77, 78, 83, 84, 86, 87, 91, 92, 96, 97, 98, 102, 103, 105, 106, 112, 113, 118, 119, 124, 125, 127, 128, 130, 131, 135, 136, 140, 141, 146, 147, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, or 237; and iii) a VH CDR3 comprising the sequence shown in SEQ ID NO: 67, 72, 73, 79, 85, 104, 107, 108, 114, 120, 126, 129, 142, 148, 219, 220, 221, 222, 223, or 236; and/or (b) a light chain variable (VL) region comprising (i) a VL CDR1 comprising the sequence shown in SEQ ID NO: 149, 154, 156, 159, 162, 165, 166, 168, 169, 170, 171, 173, 174, 176, 178, 181, 182, 185, 187, 190, 192, 195, 198, 238, or 239; (ii) a VL CDR2 comprising the sequence shown in SEQ ID NO: 150, 152, 155, 157, 160, 163, 172, 175, 179, 183, 186, 188, 191, 193, 196, or 199; and (iii) a VL CDR3 comprising the sequence shown in SEQ ID NO: 151, 153, 158, 161, 164, 167, 177, 180, 184, 189, 194, 197, or 200.

[0084] In another aspect, provided is an isolated antibody which specifically binds to EGFRvIII, wherein the antibody comprises: a VH region comprising a VH CDR1, VH CDR2, and VH CDR3 of the VH sequence shown in SEQ ID NO:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 30, 32, 34, 35, 37, 39, 41, 43, 44, 46, 48, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 202, 203, 204, 205, 206, 207, 208, 209, 210, 214, 216, 217, or 218; and/or a VL region comprising VL CDR1, VL CDR2, and VL CDR3 of the VL sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 36, 38, 40, 42, 45, 47, 49, 51, 211, 212, 213, or 215.

[0085] Provided herein is an antibody having any one of partial light chain sequence as listed in Table 1 and/or any one of partial heavy chain sequence as listed in Table 1. In Table 1, the underlined sequences are CDR sequences according to Kabat and in bold according to Chothia.

Table 1

mAb	Light Chain	Heavy Chain
m62G7	DVVMTQTPLTSLVTIGQPASISCK <u>SSQSLLYSNGKTYL</u> NWLLQRP QSPKRLIYLVSKLDSGVPDRFTG SGSGTDFTLKISRVEAEDLGFYY <u>CVQDTHFPLT</u> FGAGTKLELK (SEQ ID NO: 2)	EVQLQQSGPELVKPGASVKISCKT <u>SGYTFD</u> YTLHWVKQSHVKSLEWI <u>GGIDPINGGTT</u> YNQKF KG KATLTV DKSSSTAYMELRSLTSEDSAVYYC <u>ARGEAMDS</u> WGQGTSTVTSS (SEQ ID NO: 1)
h62G7	DVVMTQSPLSLPVTLGQPASISC <u>KSSQSLLYSNGKTYL</u> NWFQQR GQSPRRLIYLVSKLDSGVPDRFS GSGSGTDFTLKISRVEAEDVGVY <u>YCVQDTHFPLT</u> FGGGTKVEIK (SEQ ID NO: 4)	QVQLVQSGAEVKKPGASVKVSCK <u>ASGYTFD</u> YTLHWVRQAPGQGLE <u>WMGGINPINGGTT</u> YNQKF KG GRVT MTRDTSTSTVYMESSLRSEDYAV <u>YYCARGEAMDS</u> WGQGTSLVTSS (SEQ ID NO: 3)
h62G7-EQ/L6	DVVMTQSPLSLPVTLGQPASISC <u>KSSQSLLYSNGKTYL</u> NWFQQR GQSPRRLIYQVSKLDSGVPDRFS GSGSGTDFTLKISRVEAEDVGVY <u>YCGQDTHFPLT</u> FGGGTKVEIK (SEQ ID NO: 6)	QVQLVQSGAEVKKPGASVKVSCK <u>ASGYTFD</u> YTLHWVRQAPGQGLE <u>WMGGIWPI</u> TGGTTYNQKF KG GRVT MTRDTSTSTVYMESSLRSEDYAV <u>YYCARGEAQGS</u> WGQGTSLVTSS (SEQ ID NO: 5)

(continued)

mAb	Light Chain	Heavy Chain
h62G7 H14/L1-DV	DVVMTQSP ¹ SLPVT ² LGQPAS ³ ISC <u>KSSQSLLYSNDKTYTNWFQ</u> QRP GQSPRRLIY <u>EVSKLDV</u> GV ⁴ PD ⁵ RF ⁶ S GSGSGDFTLKISRVEAEDVGVY YCGQDTHF ⁷ PLT ⁸ FGGGTKVEIK (SEQ ID NO: 8)	QVQLVQSGAEVKKPGASVKV ⁹ SCK ASGYTFTD ¹⁰ YTLHWVRQAPGQGLE WMGGIWPI ¹¹ TGGTTY ¹² NQKFKGRVT MTRDTSTSTVYMELSSLRSEDTAV YYCARGEAGEGS ¹³ WGQGT ¹⁴ LVTVSS (SEQ ID NO: 7)
42G9	EVVLTQSPATLSVSPGERATLSC <u>RASQSVRSNLAWYQKSGQ</u> AP RLLIYGSTIRAT ¹ GVPARFSGSGS GTEFTLTIS ² SLQSEDFAVYYC <u>QQ</u> <u>YSDWPFT</u> FGPGTKVDIK (SEQ ID NO: 10)	QVTLKESGPVLLKPTETLT ³ LCTVS <u>GFSLSNPRMGVSWIRQPPG</u> KALE WFAHIFSTDEKSL ⁴ KL ⁵ SL ⁶ RSRL ⁷ TL ⁸ SK DTSK ⁹ SVVLTMTNMAPVDSATYY CAR <u>DSSN</u> YEGYFD ¹⁰ FWGQGT ¹¹ LVTV SS (SEQ ID NO: 9)
32A10	EVVMTQSPATLSVSPGERVTLSC <u>RASQSVSSNFAWYQQRP</u> GQAP RLLLYGAT ¹ TRAT ² GLPGRFSGSGS GTENILTISS ³ LQSEDFAIYFC <u>QQY</u> <u>KDWPFT</u> FGPGSKVDIK (SEQ ID NO: 12)	QVTLKESGPVLVKPTETLT ⁴ LCTV <u>SGFSLSNARMGVSWIRQPPG</u> KAL EWLAHIFSTDEKSI ⁵ RRSL ⁶ RSRL ⁷ TL ⁸ S KDTSK ⁹ SVVLTMTNMDPVD ¹⁰ TATY FCAR <u>DSSN</u> YEGYFD ¹¹ YWGQGT ¹² LVTV VSS (SEQ ID NO: 11)

(continued)

mAb	Light Chain	Heavy Chain
20B9	EIVMTQSPATLSVSPGERATLSC <u>RVSQSIGANLAWYQQKFGQA</u> PR LLIY <u>GASTRAT</u> GIPVRFSGGGSG TEFTLTISLQSEDFAYSC <u>QQYIY</u> <u>WPFT</u> FGPGTTVDIK (SEQ ID NO: 14)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNARMGVSWIRQPPG</u> KAL EWLG <u>HIFSTDEKSYSTSLR</u> GRITIS KDTSRGLVVLTLTNMDPVDATYY CAR <u>DSSNYEGYFDF</u> WGPGLVTV SS (SEQ ID NO: 13)
14C11	EIVMTQSPATLSVSPGERATLSC <u>RASQSVSNLAWYQQKPGQA</u> P RLLIY <u>GASTRAT</u> GVPARFSGSDS GTEFSLTISLQSEDFAVYFC <u>QQ</u> <u>YKDWPF</u> TFGPGTKVEIK (SEQ ID NO: 16)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNARMGVSWIRQPPG</u> KAL EWF <u>AHIFSTDEKSFRTSLR</u> SLTL SKDTSKQVVLTMTNMDPVDAT YYCAR <u>DSSNYEGYF</u> DYWGQGILV TVSS (SEQ ID NO: 15)
21E11	DMVVTQSPATLSVSPGERATLSC <u>RASQSVGSDLAWYQQPPGQSP</u> RLLIY <u>GASTRAT</u> GVPARFSGSGS GTDFTLTITSLESEDFAVY <u>CQQY</u> <u>NDWPF</u> TFGPGTKVDIK (SEQ ID NO: 18)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNVRMGVSWIRQPPG</u> KAL EWF <u>AHIFSSDEKSIRRSLR</u> SLTL KDTSKQVVLTMTNMDPVDATY YCAR <u>DSSNYEGYFDF</u> WQGGLVT VSSN (SEQ ID NO: 17)

(continued)

mAb	Light Chain	Heavy Chain
49B11	EMEVTSQSPATLSVSPGERATLSC <u>RASQNI</u>GSDLAWYQQQSGQAP RLLI<u>SGASTRAT</u>GVPTRFSGSGS GTDFTLTITSLQSEDFAVYYC<u>QQ</u> <u>YNDWPFT</u>FGPGTKVDIK (SEQ ID NO: 20)	QVTLKESGPVLVKPTETLTLTCTV <u>SGFSL</u>SNVRMGVSWIRQPPGKAL EWFA<u>HIFSSDEKS</u>IRRSRLRLTSL KDTSKSQVVLMTNMDPVDATY YCARD<u>DSSN</u>YEGYFDYWGQGLVT VSS (SEQ ID NO: 19)
46E10	EVVMTQSPPNLSVSPGERATLSC <u>RASQSVTS</u>NFAWYQQRPQGSP RLLLY<u>GASTRAT</u>GVPGRFSGSG SGTENILTISSLQSEDFAVYFC<u>QQ</u> <u>YKDWPF</u>TFGPGSKVDIK (SEQ ID NO: 22)	QVTLKESGPVLVKPTETLTLTCTV <u>SGFSL</u>SNARMGVSWIRQPPGKAL EWLA<u>HIFSTDEKS</u>IRRSRLRLTSL KDTSKSQVVLMTNMDPVDATY CARD<u>DSSN</u>YEGYFDYWGQGLVT SS (SEQ ID NO: 21)
12H6	EVVMTQSPATLSVSPGERATLSC <u>RASQGVSS</u>NFAWYQQRPQGSP RLLLY<u>GASTRAT</u>GVPGRFSGSG SGTENILTISSLQSEDFAIYFC<u>QQ</u> <u>YKDWPF</u>TFGPGSKVDIK (SEQ ID NO: 24)	QVTLKESGPVLVKPTETLTLTCTV <u>SGFSL</u>SNARMGVSWIRQPPGKAL EWLA<u>HIFSTDEKS</u>IRRSRLRLTSL KDTSKSQVVLMTNMDPVDATY YCARD<u>DSSN</u>YEGYFDYWGQGLVT VSS (SEQ ID NO: 23)

(continued)

mAb	Light Chain	Heavy Chain
19A9	EVVMTQSPATLSVSPGERATLSC <u>RASQSVNRNLAWYQQKPGQAP</u> RLLIFG TS TRATGIPARFSGSGG TEFTLTIDSLQSEHSGLYYC QQY <u>NDWPF</u> TFGPGTKVDIK (SEQ ID NO: 26)	QVTLEESGPVLVKPTETLTCTV <u>SGFSLSNARMGVSWIRQPPGKAP</u> EWFA HIF STDEKSLRLSLRSRLTL SKDTSKSVVLTMTNMDPVDAT YYCARD SSN YEGYFDYWGGGTLV TVSS (SEQ ID NO: 25)
11B11	EVLMTQSPATLSVSPGERATLSC <u>RASQSVSTNFAWYQQRPGQAP</u> RLLLF GAS TRATGIPGRFSGSGS GTENILTISSLQSEDFAIYFC QQY <u>KDWPF</u> TFGPGSKVEIK (SEQ ID NO: 28)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNAKMGVSWIRQPPGKAL</u> EWLA HIF STDEKSIIRSLRSRLTM SKDTSKSVVLTMTNMDPVDAT YYCVRD SSN YEGYFDYWGGGTLV TVSS (SEQ ID NO: 27)
21E7	DVVLTQSPATLSVSPGERATLSC <u>RASQSVNSNLAWYQQNPGQAP</u> RLLIFG SS TRATGIPASFGSGSGG TEFTLTINSLQSEHSAYYC QQY <u>NDWPF</u> TFGPGTKVDIK (SEQ ID NO: 29)	QVTLEESGPVLVKPTETLTCTV <u>SGFSLSNARMGVSWIRQPPGKAP</u> EWFA HIF STDEKSLRLSLRSRLTL SKDTSKSVVLTMTNMDPVDAT YYCARD SSN YEGYFDYWGGGTLV TVSS (SEQ ID NO: 25)

(continued)

mAb	Light Chain	Heavy Chain
12B2	EVVMTQSPATLSVSPGERATLSC <u>RASQSVINNLA</u> WYQQKPGQAPR LLIY <u>GTSTRAT</u> DIPARFSGSGGT EFTLTISLQSEDFAVYYC <u>QDYN</u> <u>NWPFT</u> FGPGTKVDIK (SEQ ID NO: 31)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNPR</u> MGVSWIRQPPGKAL EWLGH <u>IFSSDEKSYRL</u> SLRSLRSLIS KDTSKSQVVLMTNMDPVDATY YCVR <u>DSSNYGGYFDY</u> WGQGLTV TVSS (SEQ ID NO: 30)
11F10	EIVMTQSPATLSVSPGERTTSLC <u>RASQSVGSNLA</u> WYQQKPGQAP RLLIY <u>GASTRASG</u> VPARFSGSGS GTEFTLTISLQSEDFAVYSC <u>QEY</u> <u>NNWPFT</u> FGQGTKVEIK (SEQ ID NO: 33)	QVTLKESGPVLVKPIETLTCTVC <u>GFSLNPR</u> MGVSWIRQPPGKALE WLGH <u>IFSSDEKSYRL</u> FLRSLRSLISK DTSKSQVVLMTNMDPVDATYY CAR <u>DSSDYEGYFDY</u> WGQGLVTV SS (SEQ ID NO: 32)
17G11	EVVMTQSPATLSVSPGERATLSC <u>RASQSVINNLA</u> WYQQKPGQAPR LLIY <u>GTSTRAT</u> DIPARFSGSGGT EFTLTISLQSEDFAVYYC <u>QDYN</u> <u>NWPFT</u> FGPGTKVDIK (SEQ ID NO: 31)	QVTLKESGPVLVKPTETLTCTVF <u>GFSLNPR</u> MGVSWIRQPPGKAPE WLGH <u>IFSSDEKSYRL</u> SLRSLRSLISK DTSKSQVVFXTNMDPDGPATYY CVR <u>DSSNYEEYFDY</u> WGQGLVTV SS (SEQ ID NO: 34)

(continued)

mAb	Light Chain	Heavy Chain
29D5	KIVMTQSPATLSVSPGERATLSC <u>RANQIVSSNLAWYQQKPGQAPR</u> LLVFGT<u>STRAT</u>GIPIRFGSGSGT EFTLTVSSSLQSEDFAVYVVC<u>QQYN</u> <u>DWPFT</u>FGPGTKVDIK (SEQ ID NO: 36)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNPR</u>MGVSWLRQPPGKAL EWFA<u>HIFSTDEKS</u>YSPSLRGRLTV SKDTSKSQVLTLTNMDPVDTATY YCARD<u>SSNVEGYFDY</u>WGQGTLLVT VSS (SEQ ID NO: 35)
30D8	DIVMTQSPLSLPVTPGEPASISCR <u>SSQSL</u>LHNKRNNYLDWFLQKPG QSPQLLIY<u>LASN</u>RASGVDPDRFSG GGSGTDFTLKISRVEAEDVGVYY <u>CMQAQQT</u>PITFGQGTRLEIK (SEQ ID NO: 38)	EVQLVESGGGLVKPGGSLRLSCE <u>ASGFTFSD</u>AWMSWVRQAPGKGL EWVVG<u>RIKSKTDGGTTD</u>YVVPLNG RFIISRDDSRNTLYLQLNNLKTEDT AVYYCTT<u>VPGSYGYW</u>GQGTLLTV SS (SEQ ID NO: 37)
20E12	DIVLTQSPLSLSVTPGEPASISCR <u>SSQSL</u>LYSNGKNYLDWFLHKPG QSPQLLIY<u>LGSN</u>RASGVDPDRFSG SGSGIDFILKISRVEAEDVGVYYC <u>MQAQQT</u>PITFGQGTRLEIK (SEQ ID NO: 40)	EVNLVESGGGLVKPGGSLRLSCE <u>ASGFTFSY</u>AWMSWVRQAPGKGL EWVVG<u>RIKSIADGGATDY</u>AAPVRN RFTISRDDSRNTLYLEMHSLKTED TAVYYCTT<u>IPGNDAFDM</u>WGQGTMT VTVSS (SEQ ID NO: 39)

(continued)

mAb	Light Chain	Heavy Chain
26B9	DIVLTQSPLSLVPTPGEPASISCR <u>SSQSLLLHRDGFNYLDWFLQKPG</u> QSPQLLIY <u>LASSRAS</u> GVPDRFSG SDSGTDFTLKISRVEAEDVGVYY <u>CMQALQTPIT</u> FGQGTRLEIK (SEQ ID NO: 42)	EVQLVESWGVLVKPGGSLRLSCA ASGFI <u>FNN</u> AWMSWVRQAPGKGLE WIGRIKS <u>SDGGTTDYAAPVKDRF</u> TISRDDSKDTLYLQMNGLKTEDTA VYFCTT <u>APGGPF</u> DYWGQGTLLTV SS (SEQ ID NO: 41)
32G8	DIVLTQSPLSLVPTPGEPASISCR <u>SSQSLLYSNGKNYLDWFLHKPG</u> QSPQLLIY <u>LGSNRAS</u> GVPDRFSG SGSGIDFILKISRVEAEDVGVYYC <u>MQAQQTPIT</u> FGQGTRLEIK (SEQ ID NO: 40)	EVNLVESGGGLVKPGGSLRLSCE ASGFTFSYAWMSWVRQAPGKGL EWVGRISITDGGVIDYAAPVRNR CTISRDDSRNTLYLEMHSLKTEDT AVYYCTT <u>IPGNDDFDM</u> WGQGRM VTVSS (SEQ ID NO: 43)
34E7	DIVLTQSPLSLVPTPGEPASISCR <u>STQSLLYSNGKNYLDWFLHKPG</u> QSPQLLIF <u>LGSIRAS</u> GVPDRFSG SGSGIDFILKISRVEAEDVGVYYC <u>MQAQQTPIT</u> FGQGTRLEIK (SEQ ID NO: 45)	EVNLVESGGGLVKPGGSLRLSCE ASGFTFSYAWMSWVRQAPGKGL EWVGRISINDGGATDYASPVNR RFTISRDDSRNMLYLEMHSLKTED TAVYYCTT <u>IPGNDAFDM</u> WGQGTLL VTVSS (SEQ ID NO: 44)

(continued)

mAb	Light Chain	Heavy Chain
20G5	DIVLTQSPLSLPVTGEPASIS <u>CR</u> <u>SSQSLLYSDRRNYLDWFLQKPG</u> QSPHLLIY <u>LGSYRAS</u> GVDPDRFSG SSGGTDFTLKISRVEAEDVGYY <u>CMQALQIPIT</u> FGQGTREIK (SEQ ID NO: 47)	EVQLVESGGDLVKPGGSLRLSCA AS <u>GFTFT</u> NAWMSWVRQAPGKGL EWV <u>GRIKSKIDGGTTDYAAPVKG</u> RFIISRDDSKNTLSLQMNSLKTEDT AMYYCTT <u>APGGPFDY</u> WGQGS TVSS (SEQ ID NO: 46)
C6	ELQSVLTQPPSASGTPGQRTIS <u>CSGSSSNIGSNYYWYQQLP</u> GT APKILIY <u>RNNQRPS</u> GVDPDRFSGS KSGTSASLAISGLRSEDEADYYC <u>AAWDDNLSGWVFGTG</u> TKLTVL (SEQ ID NO: 49)	QVQLVQSGAEVKKPGSSVKVSK AS <u>GDTFSS</u> NAISWVRQAPGQGLE WMG <u>VIIPIFGTADY</u> AQKFQGRVTIT ADESTSTAYMELSSLRSEDTAVYY CAR <u>HTYHEYAGGY</u> GG <u>AMDPWG</u> QGTLTVTVSS (SEQ ID NO: 48)
B5	DIQMTQSPSSLSASVGDRTITC <u>RASQISSYLNWYQQKPGKAPK</u> LLIY <u>AASSLQSG</u> VPSRFSGSGSG TDFTLTISLQPEDFATYYC <u>QQSY</u> <u>STPLTFGQGT</u> KVEIK (SEQ ID NO: 51)	EVQLLESGGGLVQPGGSLRLSCA AS <u>GFTFSN</u> YAMSWVRQAPGKGLE WV <u>DISGGGRTY</u> ADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAV YYCAR <u>AGLLYGGGVYPM</u> DIWGQ GTLTVTVSS (SEQ ID NO: 50)

(continued)

mAb	Light Chain	Heavy Chain
42G9-1	EVVLTQSPATLSVSPGERATLSC <u>RASQSVRSNLAWYQQKSGQAP</u> RLLIY <u>GSTIRAT</u> GVPARFSGSGS GTEFTLTISLQSEDFAVYYC <u>QQ</u> <u>YSDWPFT</u> FGPGTKVDIK (SEQ ID NO: 10)	QVTLKESGPVLLKPTETLTLTCTVS <u>GFSLSNPRMGVSWIRQPPGKALE</u> WFA <u>HIFSTDEKSLKLSLRSLTSLK</u> DTSKSQVVLTMTNMAPVDSATYY CAR <u>X₁X₂SNYEGYDF</u> WGQGTLV VSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 52)
32A10-1	EVVMTQSPATLSVSPGERVTLS <u>RASQSVSSNFAWYQQRPGQAP</u> RLLLY <u>GATTIRAT</u> GLPGRFSGSGS GTENILTISSLQSEDFAIYFC <u>QQY</u> <u>KDWPFT</u> FGPGSKVDIK (SEQ ID NO: 12)	QVTLKESGPVLVKPTETLTLTCTV <u>SGFSLSNARMGVSWIRQPPGKAL</u> EWLA <u>HIFSTDEKSIRRSRLTSL</u> KDTSKSQVVLTMTNMDPVDATY FCAR <u>X₁X₂SNYEGYDF</u> WGQGTLV TVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 53)

(continued)

mAb	Light Chain	Heavy Chain
20B9-1	EIVMTQSPATLSVSPGERATLSC <u>RVSQSIGANLAWYQQKFGQA</u>PR LLIY<u>GASTRAT</u>GIPVRFSGGGSG TEFTLTISLQSEDFAIYSC<u>QQYIY</u> <u>WPFT</u>FGPGTTVDIK (SEQ ID NO: 14)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNARMGVSWIRQPPG</u>KAL EWLG<u>HIFSTDEKSYSTSLR</u>GRITIS KDTSRGLVVLTLTNMDPVDATYY CAR<u>X₁X₂SNYEGYDF</u>FWGPGFLVT VSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 54)
14C11-1	EIVMTQSPATLSVSPGERATLSC <u>RASQSVSNNLAWYQQKPGQA</u>P RLLIY<u>GASTRAT</u>GVPARFSGSDS GTEFSLTISLQSEDFAVYFC<u>QQ</u> <u>YKDWPFT</u>FGPGTKVEIK (SEQ ID NO: 16)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNARMGVSWIRQPPG</u>KAL EWFA<u>HIFSTDEKSFRTSLR</u>SRLTL SKDTSKSQVLTMTNMDPVDAT YYCAR<u>X₁X₂SNYEGYDF</u>YWGGQIL VTVSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 55)

(continued)

mAb	Light Chain	Heavy Chain
21E11-1	DMVVTQSPATLSVSPGERATLSC <u>RASQSVGSDLAWYQQPPGQSP</u> RLLIY<u>GASTRAT</u>GVPARFSGSGS GTDFTLTITSLESEDFAVYYC<u>QQY</u> <u>NDWPFT</u>FGPGTKVDIK (SEQ ID NO: 18)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNVR</u>MGVSWIRQPPGKAL EWFA<u>HIFSSDEKS</u>IRRSRLRLTSL KDTSKSQVVLMTNMDPVDATY YCAR<u>X₁X₂SNYEGYFD</u>FWGQGTLV TVSSN, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 56)
49B11-1	EMEVTSQSPATLSVSPGERATLSC <u>RASQNI</u><u>GSDLAWYQQSQGQAP</u> RLLI<u>S</u><u>GASTRAT</u>GVPTRFSGSGS GTDFTLTITSLSQSEDFAVYYC<u>QQ</u> <u>YNDWPFT</u>FGPGTKVDIK (SEQ ID NO: 20)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLNVR</u>MGVSWIRQPPGKAL EWFA<u>HIFSSDEKS</u>IRRSRLRLTSL KDTSKSQVVLMTNMDPVDATY YCAR<u>X₁X₂SNYEGYFD</u>YWGQGTLV TVSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 57)

(continued)

mAb	Light Chain	Heavy Chain
46E10-1	EVVMTQSPPNLSVSPGERATLSC <u>RASQSVTSNFAWYQQRPGQSP</u> RLLLY <u>GASTRAT</u> GVPGRFSGSG SGTENILTISSLQSEDFAVYFC <u>QQ</u> <u>YKDWPF</u> TFGPGSKVDIK (SEQ ID NO: 22)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNARM</u> GVSWIRQPPGKAL EWL <u>AHIFSTDEKS</u> IRRSRLTSL KDTSKSQVVLMTNMDPVDATYY CAR <u>X₁X₂SNYEGYFDY</u> WGQGTLLV VSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 58)
12H6-1	EVVMTQSPATLSVSPGERATLSC <u>RASQGVSSNFAWYQQRPGQSP</u> RLLLY <u>GASTRAT</u> GVPGRFSGSG SGTENILTISSLQSEDFAIYFC <u>QQ</u> <u>YKDWPF</u> TFGPGSKVDIK (SEQ ID NO: 24)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNARM</u> GVSWIRQPPGKAL EWL <u>AHIFSTDEKS</u> IRRSRLTSL KDTSKSQVVLMTNMDPVDATY YCAR <u>X₁X₂SNYEGYFDY</u> WGQGTLLV TVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 59)

(continued)

mAb	Light Chain	Heavy Chain
19A9-1	EVVMTQSPATLSVSPGERATLSC <u>RASQSVNRNLAWYQQKPGQAP</u> RLLIFG <u>GTSTRAT</u> GIPARFSGSGG TEFTLTIDSLQSEHSGLYYC <u>QQY</u> <u>NDWPF</u> TFGPGTKVDIK (SEQ ID NO: 26)	QVTLEESGPVLVKPTETLTCTV <u>SGFSLSNARMGVSWIRQPPGKAP</u> EWFA <u>HIFSTDEKSLRSLRSLTL</u> SKDTSKSQVVLMTNMDPVDAT YYCAR <u>X₁X₂SNYEGYFDYWGQGT</u> L VTVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 60)
11B11-1	EVLMTQSPATLSVSPGERATLSC <u>RASQSVSTNFAWYQQRPQGAP</u> RLLLF <u>GASTRA</u> TGIPGRFSGSGS GTENILTISSLSQSEDFAIYFC <u>QQY</u> <u>KDWPF</u> TFGPGSKVEIK (SEQ ID NO: 28)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNAKMGVSWIRQPPGKAL</u> EWLA <u>HIFSTDEKSIRRSRLRSLTM</u> SKDTSKSQVVLMTNMDPVDAT YYCVR <u>X₁X₂SNYEGYFDYWGQGT</u> L VTVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 61)

(continued)

mAb	Light Chain	Heavy Chain
21E7-1	DVVLTQSPATLSVSPGERATLSC <u>RASQSVNSNLAWYQQNPGQAP</u> RLLIF <u>GSSTRAT</u> GIPASFSGSGG TEFTLTINSLQSEHSAVYYC <u>QQY</u> <u>NDWPFT</u> FGPGTKVDIK (SEQ ID NO: 29)	QVTLEESGPVLVKPTETLTCTV <u>SGFSLSNARMGVSWIRQPPGKAP</u> EWFA <u>HIFSTDEKSLRSLRSLTL</u> SKDTSKSQVVLMTNMDPVDAT YYCAR <u>X₁X₂SNYEGYFDYWGQGT</u> L VTVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 202)
12B2-1	EVVMTQSPATLSVSPGERATLSC <u>RASQSVINNLA</u> WYQQKPGQAPR LLIY <u>GTSTRAT</u> DIPARFSGSGGT EFTLTISLSQSEDFAVYYC <u>QDYN</u> <u>NWPFT</u> FGPGTKVDIK (SEQ ID NO: 31)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNPRMGVSWIRQPPGKAL</u> EWL <u>GHISSDEKSYRLSLRSLSIS</u> KDTSKSQVVLMTNMDPVDATY YCVR <u>X₁X₂SNYGGYFDYWGQGT</u> L VTVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 203)

(continued)

mAb	Light Chain	Heavy Chain
11F10-1	EIVMTQSPATLSVSPGERATLSC <u>RASQSVGSNLA</u>WYQQKPGQAP RLIIY<u>GASTRASG</u>VPARFSGSGS GTEFTLTISLQSEDFAVYSC<u>QKEY</u> <u>NNWPFT</u>FGQGTKVEIK (SEQ ID NO: 33)	QVTLKESGPVLVKPIETLTLTCTVC <u>GFSLSNPRMGVSWIRQPPGKALE</u> WLG<u>HIFSSDEKS</u>YRLFLRSRLSISK DTSKSQVVLTMTNMDPVDATYY CAR<u>X₁X₂SDYEGYFDY</u>WGQGLVT VSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 204)
17G11-1	EVVMTQSPATLSVSPGERATLSC <u>RASQSVINNLA</u>WYQQKPGQAPR LLIY<u>GTSTRATD</u>IPARFSGSGSGT EFTLTISLQSEDFAVYYC<u>QDYN</u> <u>NNWPFT</u>FGPGTKVDIK (SEQ ID NO: 31)	QVTLKESGPVLVKPTETLTLTCTVF <u>GFSLSNPRMGVSWIRQPPGKAPE</u> WLG<u>HIFSSDEKS</u>YRLSLRSRLSISK DTSKSQVVFXTNMDPGDPATYY CVR<u>X₁X₂SNYEEYFDY</u>WGQGLVT VSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 205)

(continued)

mAb	Light Chain	Heavy Chain
29D5-1	KIVMTQSPATLSVSPGERATLSC <u>RANQIVSSNLA</u> WYQQKPGQAPR LLVFGT <u>STRAT</u> GIPIRFGSGSGT EFTLTVSSSLQSEDFAVYVC <u>QQYN</u> <u>DWPFT</u> FGPGTKVDIK (SEQ ID NO: 36)	QVTLKESGPVLVKPTETLTCTV <u>SGFSLSNPR</u> MGVSWLRQPPGKAL EWFA <u>HIFSTDEKS</u> YSPSLRGRLTV SKDTSKSVVLTLTNMDPVDATY YCAR <u>X₁X₂SNYEGYFDY</u> WGQGTLV TVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 206)
30D8-1	DIVMTQSPSLPVTGPGEPAISCR <u>SSQSLLHNKRNNYL</u> DWFLQKPG QSPQLLIY <u>LASN</u> RASGVDPDRFSG GSGGTDFTLKISRVEAEDVGVYY <u>CMQAQQTPIT</u> FGQGTRLEIK (SEQ ID NO: 38)	EVQLVESGGGLVKPGGSLRLSCE <u>ASGFTFSD</u> AWMSWVRQAPGKGL EWVG <u>RIKSKTX₁X₂GTTD</u> YVVPLN <u>GRFI</u> SRDDSRNTLYQLNNLKTED TAVYYCTT <u>VPGSYG</u> WGQGTLLVT VSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 207)

(continued)

mAb	Light Chain	Heavy Chain
20E12-1	DIVLTQSPLSLVTPGEPASISCR <u>SSQSLLYSX₁X₂KNYLDWFLHKP</u> GQSPQLLIYL<u>G</u>SNRASGVDPDRFS GSGGIDFILKISRVEAEDVGYY <u>CMQAQQTPI</u>TFGQGTREIK, , wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 211)	EVNLVESGGGLVKPGGSLRLSCE ASGFTFSYAWMSWVRQAPGKGL EWVG<u>RIKSIX₁X₂GATDYAAPVRN</u> RFTISRDDSRNTLYLEMHSLKTED TAVYYCTT<u>IPGNDAFDM</u>WGQGT MTVSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 208)
26B9-1	DIVLTQSPLSLVTPGEPASISCR <u>SSQSLLHRX₁X₂FNYLDWFLQKP</u> GQSPQLLIYL<u>ASSRAS</u>GVDPDRFS GSDSGTDFTLKISRVEAEDVGYY YC<u>MQALQTPIT</u>FGQGTREIK, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 212)	EVQLVESWGVLVKPGGSLRLSCA ASGFIF<u>NNAWMSWVRQAPGKGLE</u> WIG<u>RIKSXSX₁X₂GTTDYAAPVKDR</u> FTISRDDSKDTLYLQMNGLKTEDT AVYFCTT<u>APGGPFDYWGQGT</u>LV TVSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 209)

(continued)

mAb	Light Chain	Heavy Chain
32G8-1	DIVLTQSPLSLSVTPGEPASISCR <u>SSQSLLYSX₁X₂KNYLDWFLHKP</u> GQSPQLLIYL<u>G</u>SNRASGVDPDRFS GSGGIDFILKISRVEAEDVGYY <u>CMQAQQTPIT</u>FGQGTRLEIK, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 213)	EVNLVESGGGLVKPGGSLRLSCE <u>ASGFTFSYAWMSWVRQAPGKGL</u> EWVG<u>RIKSITX₁X₂GVIDY</u>AAPVRN RCTISRDDSRNTLYLEMHSLKTED TAVYYCTT<u>IPGNDDFDM</u>WGQGRM VTVSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 210)
34E7-1	DIVLTQSPLSLSVTPGEPASISCR <u>STQSLLYSX₁X₂KNYLDWFLHKP</u> GQSPQLLI<u>FLG</u>SIRASGVDPDRFS GSGGIDFILKISRVEAEDVGYY <u>CMQAQQTPIT</u>FGQGTRLEIK, , wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 215)	EVNLVESGGGLVKPGGSLRLSCE <u>ASGFTFSYAWMSWVRQAPGKGL</u> EWVG<u>RIKSINX₁X₂GATDY</u>ASPVRN RFTISRDDSRNMLYLEMHSLKTED TAVYYCTT<u>IPGNDAFDM</u>WGQGTL VTVSS, wherein X₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 214)

(continued)

mAb	Light Chain	Heavy Chain
20G5-1	DIVLTQSPLSLPVT P GEPASIS C R <u>SSQSLLYSDRRNYLDWFLQKPG</u> QSPHLLIY <u>LGSYRASG</u> VPDRFSG SSGTDFTLKISRVEAEDVGYY <u>CMQALQIPIT</u> FGQGT R LEIK (SEQ ID NO: 47)	EVQLVESGGDLVKPGGSLRLSCA AS GFTFT NAWM S WVRQAPGKGL EWVGR IKSKIX₁X₂GT TDYAAPVKG RFIISRDSDSKNTLSLQMN S LKTEDT AMYYCTT APGGPFDY WGQGS L V TVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 216)
C6-1	ELQSVLTQPPSASGTPGQ R VTIS <u>CSGSSSNIGSNVYVY</u> QQLPGT APKILIY RNNQRPSG VPDRFSGS KSGTSASLAISGLRSEDEADYYC <u>AAWDDNLSGWVFGTGT</u> KLTVL (SEQ ID NO: 49)	QVQLVQSGAEVKKPGSSVKV S CK AS GDTFSS NAISWVRQAPGQ G LE WMG VIIPIFGTADY AQKFQGRVTIT ADE S TSTAYMELSSLRSEDTAVYY CAR HTYHEYAGGYGGAMX₁X₂W GGGLTVTVSS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 217)

(continued)

mAb	Light Chain	Heavy Chain
B5-1	DIQMTQSPSSLSASVGDRTITC <u>RASQSISSYLNWYQQKPGKAPK</u> LLIV <u>AASSLQSGVPSRFS</u> SGSGG TDFTLTISLQPEDFATYYC <u>QQSY</u> <u>STPLTF</u> GQGTKVEIK (SEQ ID NO: 51)	EVQLLESGGGLVQPGGSLRLSCA ASGFTFSN <u>YAMSWVRQAPGKGLE</u> WVSD <u>DISGGGRTYYAX₁X₂</u> VKGRF TISRDNSKNTLYLQMNSLRAEDTA VYYCAR <u>AGLLYGGGVYPM</u> DIWG QGTLVTVS, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 218)

[0086] Also provided herein are CDR portions of antigen binding domains of antibodies to EGFRvIII (including Chothia, Kabat CDRs, and CDR contact regions). Determination of CDR regions is well within the skill of the art. It is understood that in some embodiments, CDRs can be a combination of the Kabat and Chothia CDR (also termed "combined CRs" or "extended CDRs"). In some embodiments, the CDRs are the Kabat CDRs. In other embodiments, the CDRs are the Chothia CDRs. In other words, in embodiments with more than one CDR, the CDRs may be any of Kabat, Chothia, combination CDRs, or combinations thereof. Table 2 provides examples of CDR sequences provided herein.

Table 2

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
m62G7	TDYTLH (SEQ ID NO: 62) (Kabat); GYTFTD (SEQ ID NO: 63) (Chothia); GYTFTDYTLH (SEQ ID NO: 64) (extended)	GIDPINGGTTYNQKFK G (SEQ ID NO: 65) (Kabat) GIDPINGGTTY (SEQ ID NO: 66) (Chothia)	GEAMDS (SEQ ID NO: 67)
h62G7	TDYTLH (SEQ ID NO: 62) (Kabat); GYTFTD (SEQ ID NO: 63) (Chothia); GYTFTDYTLH (SEQ ID NO: 64) (extended)	GINPINGGTTYNQKFK G (SEQ ID NO: 68) (Kabat) GINPINGGTTY (SEQ ID NO: 69) (Chothia)	GEAMDS (SEQ ID NO: 67)
h62G7-H14	TDYTLH (SEQ ID NO: 62) (Kabat); GYTFTD (SEQ ID NO: 63) (Chothia); GYTFTDYTLH (SEQ ID NO: 64) (extended)	GIWPITGGTTYNQKFK G (SEQ ID NO: 70) (Kabat) GIWPITGGTTY (SEQ ID NO: 71) (Chothia)	GAEAGS (SEQ ID NO: 72)
h62G7-EQ	TDYTLH (SEQ ID NO: 62) (Kabat); GYTFTD (SEQ ID NO: 63) (Chothia); GYTFTDYTLH (SEQ ID NO: 64) (extended)	GIWPITGGTTYNQKFK G (SEQ ID NO: 70) (Kabat) GIWPITGGTTY (SEQ ID NO: 71) (Chothia)	GEAQGS (SEQ ID NO: 73)
42G9	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSTDEKSLKLSLRS (SEQ ID NO: 77) (Kabat) HIFSTDEKSL (SEQ ID NO: 78) (Chothia)	DSSNYEGYFDF (SEQ ID NO: 79)

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
32A10	SNARMGVS (SEQ ID NO: 80) (Kabat); GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSIRRLRS (SEQ ID NO: 83) (Kabat) HIFSTDEKSI (SEQ ID NO: 84) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)
20B9	SNARMGVS (SEQ ID NO: 80) (Kabat); GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSYSTSLRG (SEQ ID NO: 86) (Kabat) HIFSTDEKSY (SEQ ID NO: 87) (Chothia)	DSSNYEGYFDF (SEQ ID NO: 79)
14C11	NNARMGVS (SEQ ID NO: 88) (Kabat); GFSLNNAR (SEQ ID NO: 89) (Chothia); GFSLNNARMGVS (SEQ ID NO: 90) (extended)	HIFSTDEKSFRTSLRS (SEQ ID NO: 91) (Kabat) HIFSTDEKSF (SEQ ID NO: 92) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)
21E11	SNVRMGVS (SEQ ID NO: 93) (Kabat); GFSLSNVR (SEQ ID NO: 94) (Chothia); GFSLSNVRMGVS (SEQ ID NO: 95) (extended)	HIFSSDEKSIRRLRS (SEQ ID NO: 96) (Kabat) HIFSSDEKSI (SEQ ID NO: 97) (Chothia)	DSSNYEGYFDF (SEQ ID NO: 79)
49B11	SNVRMGVS (SEQ ID NO: 93) (Kabat); GFSLSNVR (SEQ ID NO: 94) (Chothia); GFSLSNVRMGVS (SEQ ID NO: 95) (extended)	HIFSSDEKSIRRLRS (SEQ ID NO: 96) (Kabat) HIFSSDEKSI (SEQ ID NO: 97) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)
46E10 12H6	SNARMGVS (SEQ ID NO: 80) (Kabat); GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSIRRLRS (SEQ ID NO: 83) (Kabat) HIFSTDEKSI (SEQ ID NO: 84) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)
19A9 21E7	SNARMGVS (SEQ ID NO: 80) (Kabat); GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSLRLSLRS (SEQ ID NO: 98) (Kabat) HIFSTDEKSL (SEQ ID NO: 78) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
11B11	SNAKMGVS (SEQ ID NO: 99) (Kabat); GFSLSNK (SEQ ID NO: 100) (Chothia); GFSLSNKMGVS (SEQ ID NO: 101) (extended)	HIFSTDEKSIRRLRS (SEQ ID NO: 83) (Kabat) HIFSTDEKSI (SEQ ID NO: 84) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)
12B2	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSSDEKSYRLSLRS (SEQ ID NO: 102) (Kabat) HIFSSDEKSY (SEQ ID NO: 103) (Chothia)	DSSNYGGYFDY (SEQ ID NO: 104)
11F10	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSSDEKSYRLFLRS (SEQ ID NO: 105) (Kabat) HIFSSDEKSY (SEQ ID NO: 103) (Chothia)	DSSDYEGYFDY (SEQ ID NO: 107)
17G11	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSSDEKSYRLSLRS (SEQ ID NO: 102) (Kabat) HIFSSDEKSY (SEQ ID NO: 103) (Chothia)	DSSNYEEYFDY (SEQ ID NO: 108)
29D5	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSTDEKSYSPSLRG (SEQ ID NO: 106) (Kabat) HIFSTDEKSY (SEQ ID NO: 87) (Chothia)	DSSNYEGYFDY (SEQ ID NO: 85)
30D8	SDAWMS (SEQ ID NO: 109) (Kabat); GFTFSD (SEQ ID NO: 110) (Chothia); GFTFSDAWMS (SEQ ID NO: 111) (extended)	RIKSKTDGGTTDYVPL NG (SEQ ID NO: 112) (Kabat) RIKSKTDGGTTDY (SEQ ID NO: 113) (Chothia)	VPGSYGY (SEQ ID NO: 114)
20E12	SYAWMS (SEQ ID NO: 115) (Kabat); GFTFSY (SEQ ID NO: 116) (Chothia); GFTFSYAWMS (SEQ ID NO: 117) (extended)	RIKSIADGGATDYAAP VRN (SEQ ID NO: 118) (Kabat) RIKSIADGGATDY (SEQ ID NO: 119) (Chothia)	IPGNDAFDM (SEQ ID NO: 120)

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
26B9	NNAWMS (SEQ ID NO: 121) (Kabat); GFIFNN (SEQ ID NO: 122) (Chothia); GFIFNNAWMS (SEQ ID NO: 123) (extended)	RIKSKSDGGTTDYAAP VKD (SEQ ID NO: 124) (Kabat) RIKSKSDGGTTDY (SEQ ID NO: 125) (Chothia)	APGGPFDY (SEQ ID NO: 126)
32G8	SYAWMS (SEQ ID NO: 115) (Kabat); GFTFSY (SEQ ID NO: 116) (Chothia); GFTFSYAWMS (SEQ ID NO: 117) (extended)	RIKSITDGGVIDYAAPV RN (SEQ ID NO: 127) (Kabat) RIKSITDGGVIDY (SEQ ID NO: 128) (Chothia)	IPGNDDFDM (SEQ ID NO: 129)
34E7	SYAWMS (SEQ ID NO: 115) (Kabat); GFTFSY (SEQ ID NO: 116) (Chothia); GFTFSYAWMS (SEQ ID NO: 117) (extended)	RIKSINDGGATDYASPV RN (SEQ ID NO: 130) (Kabat) RIKSINDGGATDY (SEQ ID NO: 131) (Chothia)	IPGNDAFDM (SEQ ID NO: 120)
20G5	TNAWMS (SEQ ID NO: 132) (Kabat); GFTFTN (SEQ ID NO: 133) (Chothia); GFTFTNAWMS (SEQ ID NO: 134) (extended)	RIKSKIDGGTTDYAAPV KG (SEQ ID NO: 135) (Kabat) RIKSKIDGGTTDY (SEQ ID NO: 136) (Chothia)	APGGPFDY (SEQ ID NO: 126)
C6	SSNAIS (SEQ ID NO: 137) (Kabat); GDTFSS (SEQ ID NO: 138) (Chothia); GDTFSSNAIS (SEQ ID NO: 139) (extended)	VIIPIFGTADYAQKFQG (SEQ ID NO: 140) (Kabat) VIIPIFGTADY (SEQ ID NO: 141) (Chothia)	HTYHEYAGGYGG AMDP (SEQ ID NO: 142)
B5	SNYAMS (SEQ ID NO: 143) (Kabat); GFTFSN (SEQ ID NO: 144) (Chothia); GFTFSNYAMS (SEQ ID NO: 145) (extended)	DISGGGGRTYYADSVK G (SEQ ID NO: 146) (Kabat) DISGGGGRTYY (SEQ ID NO: 147) (Chothia)	AGLLYGGGVYPM DI (SEQ ID NO: 148)

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
42G9-1	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSTDEKSLKLSLRS (SEQ ID NO: 77) (Kabat) HIFSTDEKSL (SEQ ID NO: 78) (Chothia)	X_1X_2 SNYEGYFDF, wherein X_1 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X_2 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 219)
32A10-1	SNARMGVS (SEQ ID NO: 80) (Kabat); GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSIRRLRS (SEQ ID NO: 83) (Kabat) HIFSTDEKSI (SEQ ID NO: 84) (Chothia)	X_1X_2 SNYEGYFDY, wherein X_1 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X_2 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
20B9-1	SNARMGVS (SEQ ID NO: 80) (Kabat); GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSYSTSLRG(S EQ ID NO: 86) (Kabat) HIFSTDEKSY (SEQ ID NO: 87) (Chothia)	X_1X_2 SNYEGYFDF, wherein X_1 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X_2 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 219)
14C11-1	NNARMGVS (SEQ ID NO: 88) (Kabat); GFSLNNAR (SEQ ID NO: 89) (Chothia); GFSLNNARMGVS (SEQ ID NO: 90) (extended)	HIFSTDEKSFRTSLRS (SEQ ID NO: 91) (Kabat) HIFSTDEKSF (SEQ ID NO: 92) (Chothia)	X_1X_2 SNYEGYFDY, wherein X_1 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X_2 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
21E11-1	SNVRMGVS (SEQ ID NO: 93) (Kabat); GFSLSNVR (SEQ ID NO: 94) (Chothia); GFSLSNVRMGVS (SEQ ID NO: 95) (extended)	HIFSSDEKSIRRLRS (SEQ ID NO: 96) (Kabat) HIFSSDEKSI (SEQ ID NO: 97) (Chothia)	X_1X_2 SNYEGYFDF, wherein X_1 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X_2 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 219)
49811-1	SNVRMGVS (SEQ ID NO: 93) (Kabat); GFSLSNVR (SEQ ID NO: 94) (Chothia); GFSLSNVRMGVS (SEQ ID NO: 95) (extended)	HIFSSDEKSIRRLRS(SE Q ID NO: 96) (Kabat) HIFSSDEKSI (SEQ ID NO: 97) (Chothia)	X_1X_2 SNYEGYFDY, wherein X_1 is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X_2 is R, H, K, D, E, S, T, N, Q, C,

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
			G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
46E10-1	SNARMGVS (SEQ ID NO: 80) (Kabat);	HIFSTDEKSIRRLSRS (SEQ ID NO: 83) (Kabat)	X ₁ X ₂ SNYEGYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
12H6-1	GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSI (SEQ ID NO: 84) (Chothia)	
19A9-1	SNARMGVS (SEQ ID NO: 80) (Kabat);	HIFSTDEKSLRLSRS (SEQ ID NO: 98) (Kabat)	
21E7-1	GFSLSNAR (SEQ ID NO: 81) (Chothia); GFSLSNARMGVS (SEQ ID NO: 82) (extended)	HIFSTDEKSL (SEQ ID NO: 78) (Chothia)	X ₁ X ₂ SNYEGYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
11B11-1	SNAKMGVS (SEQ ID NO: 99) (Kabat); GFSLSNAK (SEQ ID NO: 100) (Chothia); GFSLSNAKMGVS (SEQ ID NO: 101) (extended)	HIFSTDEKSIRRLSRS (SEQ ID NO: 83) (Kabat) HIFSTDEKSI (SEQ ID NO: 84) (Chothia)	X ₁ X ₂ SNYEGYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
12B2-1	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSSDEKSYRLSRS (SEQ ID NO: 102) (Kabat) HIFSSDEKSY (SEQ ID NO: 103) (Chothia)	X ₁ X ₂ SNYGGYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 221)
11F10-1	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSSDEKSYRLFLRS (SEQ ID NO: 105) (Kabat) HIFSSDEKSY (SEQ ID NO: 103) (Chothia)	X ₁ X ₂ SDYEGYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 222)

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
17G11-1	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSSDEKSYRLSLRS (SEQ ID NO: 102) (Kabat) HIFSSDEKSY (SEQ ID NO: 103) (Chothia)	X ₁ X ₂ SNYEEYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 223)
29D5-1	SNPRMGVS (SEQ ID NO: 74) (Kabat); GFSLSNPR (SEQ ID NO: 75) (Chothia); GFSLSNPRMGVS (SEQ ID NO: 76) (extended)	HIFSTDEKSYSPSLRG (SEQ ID NO: 106) (Kabat) HIFSTDEKSY (SEQ ID NO: 87) (Chothia)	X ₁ X ₂ SNYEGYFDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 220)
30D8-1	SDAWMS (SEQ ID NO: 109) (Kabat); GFTFS (SEQ ID NO: 110) (Chothia); GFTFSDAWMS (SEQ ID NO: 111) (extended)	RIKSKTX ₁ X ₂ GTTDYV PLNG, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 224) (Kabat) RIKSKTX ₁ X ₂ GTTDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 225) (Chothia)	VPGSYGY (SEQ ID NO: 114)
20E12-1	SYAWMS (SEQ ID NO: 115) (Kabat); GFTFSY (SEQ ID NO: 116) (Chothia); GFTFSYAWMS (SEQ ID NO: 117) (extended)	RIKSIAX ₁ X ₂ GATDYAAP VRN, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 226) (Kabat) RIKSIAX ₁ X ₂ GATDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 227) (Chothia)	IPGNDAFDM (SEQ ID NO: 120)
26B9-1	NNAWMS (SEQ ID NO: 121) (Kabat); GFIFNN (SEQ ID NO: 122) (Chothia); GFIFNNAWMS (SEQ ID NO: 123) (extended)	RIKSKSX ₁ X ₂ GTTDYAAP VKD, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 228) (Kabat) RIKSKSX ₁ X ₂ GTTDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 229) (Chothia)	APGGPFDY (SEQ ID NO: 126)

(continued)

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
32G8-1	SYAWMS (SEQ ID NO: 115) (Kabat); GFTFSY (SEQ ID NO: 116) (Chothia); GFTFSYAWMS (SEQ ID NO: 117) (extended)	RIKSITX ₁ X ₂ GVIDYAAP VRN, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 230) (Kabat) RIKSITX ₁ X ₂ GVIDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 231) (Chothia)	IPGNDDFDM (SEQ ID NO: 129)
34E7-1	SYAWMS (SEQ ID NO: 115) (Kabat); GFTFSY (SEQ ID NO: 116) (Chothia); GFTFSYAWMS (SEQ ID NO: 117) (extended)	RIKSINX ₁ X ₂ GATDYASP VRN, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 232) (Kabat) RIKSINX ₁ X ₂ GATDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 233) (Chothia)	IPGNDAFDM (SEQ ID NO: 120)
20G5-1	TNAWMS (SEQ ID NO: 132) (Kabat); GFTFTN (SEQ ID NO: 133) (Chothia); GFTFTNAWMS (SEQ ID NO: 134) (extended)	RIKSKIX ₁ X ₂ GTTDYAAP VKG, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 234) (Kabat) RIKSKIX ₁ X ₂ GTTDY, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 235) (Chothia)	APGGPFDDY (SEQ ID NO: 126)
C6-1	SSNAIS (SEQ ID NO: 137) (Kabat); GDTFSS (SEQ ID NO: 138) (Chothia); GDTFSSNAIS (SEQ ID NO: 139) (extended)	VIIPIFGTADYAQKFQG (SEQ ID NO: 140) (Kabat) VIIPIFGTADY (SEQ ID NO: 141) (Chothia)	HTYHEYAGGYGG AMX ₁ X ₂ , wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 236)
B5-1	SNYAMS (SEQ ID NO: 143) (Kabat); GFTFSN (SEQ ID NO: 144) (Chothia); GFTFSNYAMS (SEQ ID NO: 145) (extended)	DISGGGGRTYYAX ₁ X ₂ VKG, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 237) (Kabat) DISGGGGRTYY (SEQ ID NO: 147) (Chothia)	AGLLYGGGVYPM DI (SEQ ID NO: 148)
Light Chain			
mAb	CDRL1	CDRL2	CDRL3
m62G7 h62G7	KSSQSLLYSNGKTYLN (SEQ ID NO: 149)	LVSKLDS (SEQ ID NO: 150)	VQDTHFPLT (SEQ ID NO: 151)

(continued)

Light Chain			
mAb	CDRL1	CDRL2	CDRL3
5 h62G7-L6	KSSQSLLYSNGKTYLN (SEQ ID NO: 149)	QVSKLDS (SEQ ID NO: 152)	GQDTHFPLT (SEQ ID NO: 153)
10 h62G7-L1-DV	KSSQSLLYSNDKTYTN (SEQ ID NO: 154)	EVSKLDV (SEQ ID NO: 155)	GQDTHFPLT (SEQ ID NO: 153)
42G9	RASQSVRSNLA (SEQ ID NO: 156)	GSTIRAT (SEQ ID NO: 157)	QQYSDWPFT (SEQ ID NO: 158)
15 32A10	RASQSVSSNFA (SEQ ID NO: 159)	GATTRAT (SEQ ID NO: 160)	QQYKDWPFPT (SEQ ID NO: 161)
20B9	RVSQSIGANLA (SEQ ID NO: 162)	GASTRAT (SEQ ID NO: 163)	QQYIWPFT (SEQ ID NO: 164)
14C11	RASQSVSNLA (SEQ ID NO: 165)	GASTRAT (SEQ ID NO: 163)	QQYKDWPFPT (SEQ ID NO: 161)
20 21E11	RASQSVGSDLA (SEQ ID NO: 166)	GASTRAT (SEQ ID NO: 163)	QQYNDWPFT (SEQ ID NO: 167)
49B11	RASQNIQSDLA (SEQ ID NO: 168)	GASTRAT (SEQ ID NO: 163)	QQYNDWPFT (SEQ ID NO: 167)
25 46E10	RASQSVTSNFA (SEQ ID NO: 169)	GASTRAT (SEQ ID NO: 163)	QQYKDWPFPT (SEQ ID NO: 161)
12H6	RASQGVSSNFA (SEQ ID NO: 170)	GASTRAT (SEQ ID NO: 163)	QQYKDWPFPT (SEQ ID NO: 161)
30 19A9	RASQSVNRNLA (SEQ ID NO: 171)	GTSTRAT (SEQ ID NO: 172)	QQYNDWPFT (SEQ ID NO: 167)
11B11	RASQSVSTNFA (SEQ ID NO: 173)	GASTRAT (SEQ ID NO: 163)	QQYKDWPFPT (SEQ ID NO: 161)
35 21E7	RASQSVNSNLA (SEQ ID NO: 174)	GSSTRAT (SEQ ID NO: 175)	QQYNDWPFT (SEQ ID NO: 167)
12B2 17G11	RASQSVINNLA (SEQ ID NO: 176)	GTSTRAT (SEQ ID NO: 172)	QDYNNWPFT (SEQ ID NO: 177)
40 11F10	RASQSVGSNLA (SEQ ID NO: 178)	GASTRASG (SEQ ID NO: 179)	QEYNNWPFT (SEQ ID NO: 180)
29D5	RANQIVSSNLA (SEQ ID NO: 181)	GTSTRAT (SEQ ID NO: 172)	QQYNDWPFT (SEQ ID NO: 167)
45 30D8	RSSQSLLHNKRNNYLD (SEQ ID NO: 182)	LASNRA (SEQ ID NO: 183)	MQAQQTPI (SEQ ID NO: 184)
20E12 32G8	RSSQSLLYSNGKNYLD (SEQ ID NO: 185)	LGSNRA (SEQ ID NO: 186)	MQAQQTPI (SEQ ID NO: 184)
50 26B9	RSSQSLLHRDGFNYLD (SEQ ID NO: 187)	LASSRA (SEQ ID NO: 188)	MQALQTPIT (SEQ ID NO: 189)
34E7	RSTQSLLYSNGKNYLD (SEQ ID NO: 190)	LGSIRA (SEQ ID NO: 191)	MQAQQTPI (SEQ ID NO: 184)
55 20G5	RSSQSLLYSRRNYLD (SEQ ID NO: 192)	LGSYRA (SEQ ID NO: 193)	MQALQIPIT (SEQ ID NO: 194)

(continued)

Light Chain			
mAb	CDRL1	CDRL2	CDRL3
C6	SGSSSNIGSNYVY (SEQ ID NO: 195)	RNNQRPS (SEQ ID NO: 196)	AAWDDNLSGWV (SEQ ID NO: 197)
B5	RASQSISSYLN (SEQ ID NO: 198)	AASSLQS (SEQ ID NO: 199)	QQSYSTPLT (SEQ ID NO: 200)
20E12-1 32G8-1	RSSQSLLYSX ₁ X ₂ KNYLD, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 238)	LGSNRAS (SEQ ID NO: 186)	MQAQQTPIIT (SEQ ID NO: 184)
26B9-1	RSSQSLLHRX ₁ X ₂ FNYLD, wherein X ₁ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W, and X ₂ is R, H, K, D, E, S, T, N, Q, C, G, P, A, V, I, L, M, F, Y, or W (SEQ ID NO: 239)	LASSRAS (SEQ ID NO: 188)	MQALQTPIT (SEQ ID NO: 189)

[0087] The present disclosure provides an antibody that binds to and competes with the antibody as described herein, including m62G7, h62G7, h62G7-H14/L1-DV, h62G7-EQ/L6, 42G9, 32A10, 20B9, 14C11, 21E11, 49B11, 46E10, 12H6, 19A9, 21E7, 11B11, 12B2, 11F10, 17G11, 29D5, 30D8, 20E12, 26B9, 32G8, 34E7, 20G5, C6, B5, 42G9-1, 32A10-1, 20B9-1, 14C11-1, 21E11-1, 49B11-1, 46E10-1, 12H6-1, 19A9-1, 21E7-1, 11B11-1, 12B2-1, 11F10-1, 17G11-1, 29D5-1, 30D8-1, 20E12-1, 26B9-1, 32G8-1, 34E7-1, 20G5-1, C6-1, and B5-1.

[0088] In some embodiments, the invention also provides CDR portions of antibodies to EGFRvIII antibodies based on CDR contact regions. CDR contact regions are regions of an antibody that imbue specificity to the antibody for an antigen. In general, CDR contact regions include the residue positions in the CDRs and Vernier zones which are constrained in order to maintain proper loop structure for the antibody to bind a specific antigen. See, e.g., Makabe et al., J. Biol. Chem., 283:1156-1166, 2007. Determination of CDR contact regions is well within the skill of the art.

[0089] The binding affinity (K_D) of the EGFRvIII antibody as described herein to EGFRvIII (such as human EGFRvIII (e.g., (SEQ ID NO: 201))) can be about 0.001 to about 5000 nM. In some embodiments, the binding affinity is about any of 5000 nM, 4500 nM, 4000 nM, 3500 nM, 3000 nM, 2500 nM, 2000 nM, 1789 nM, 1583 nM, 1540 nM, 1500 nM, 1490 nM, 1064 nM, 1000 nM, 933 nM, 894 nM, 750 nM, 705 nM, 678 nM, 532 nM, 500 nM, 494 nM, 400 nM, 349 nM, 340 nM, 353 nM, 300 nM, 250 nM, 244 nM, 231 nM, 225 nM, 207 nM, 200 nM, 186 nM, 172 nM, 136 nM, 113 nM, 104 nM, 101 nM, 100 nM, 90 nM, 83 nM, 79 nM, 74 nM, 54 nM, 50 nM, 45 nM, 42 nM, 40 nM, 35 nM, 32 nM, 30 nM, 25 nM, 24 nM, 22 nM, 20 nM, 19 nM, 18 nM, 17 nM, 16 nM, 15 nM, 12 nM, 10 nM, 9 nM, 8 nM, 7.5 nM, 7 nM, 6.5 nM, 6 nM, 5.5 nM, 5 nM, 4 nM, 3 nM, 2 nM, 1 nM, 0.5 nM, 0.3 nM, 0.1 nM, 0.01 nM, or 0.001 nM. In some embodiments, the binding affinity is less than about any of 5000 nM, 4000 nM, 3000 nM, 2000 nM, 1000 nM, 900 nM, 800 nM, 250 nM, 200 nM, 100 nM, 50 nM, 30 nM, 20 nM, 10 nM, 7.5 nM, 7 nM, 6.5 nM, 6 nM, 5 nM, 4.5 nM, 4 nM, 3.5 nM, 3 nM, 2.5 nM, 2 nM, 1.5 nM, 1 nM, or 0.5 nM.

[0090] Bispecific antibodies, monoclonal antibodies that have binding specificities for at least two different antigens, can be prepared using the antibodies disclosed herein. Methods for making bispecific antibodies are known in the art (see, e.g., Suresh et al., Methods in Enzymology 121:210, 1986). Traditionally, the recombinant production of bispecific antibodies was based on the coexpression of two immunoglobulin heavy chain-light chain pairs, with the two heavy chains having different specificities (Millstein and Cuellar, Nature 305, 537-539, 1983). Accordingly, in one aspect, provided is a bispecific antibody wherein the bispecific antibody is a full-length human antibody, comprising a first antibody variable domain of the bispecific antibody specifically binding to a target antigen (e.g., EGFRvIII), and comprising a second antibody variable domain of the bispecific antibody capable of recruiting the activity of a human immune effector cell by specifically binding to an effector antigen located on the human immune effector cell.

[0091] The human immune effector cell can be any of a variety of immune effector cells known in the art. For example, the immune effector cell can be a member of the human lymphoid cell lineage, including, but not limited to, a T cell (e.g.,

a cytotoxic T cell), a B cell, and a natural killer (NK) cell. The immune effector cell can also be, for example without limitation, a member of the human myeloid lineage, including, but not limited to, a monocyte, a neutrophilic granulocyte, and a dendritic cell. Such immune effector cells may have either a cytotoxic or an apoptotic effect on a target cell or other desired effect upon activation by binding of an effector antigen.

[0092] The effector antigen is an antigen (e.g., a protein or a polypeptide) that is expressed on the human immune effector cell. Examples of effector antigens that can be bound by the heterodimeric protein (e.g., a heterodimeric antibody or a bispecific antibody) include, but are not limited to, human CD3 (or CD3 (Cluster of Differentiation) complex), CD16, NKG2D, NKp46, CD2, CD28, CD25, CD64, and CD89.

[0093] The target cell can be a cell that is native or foreign to humans. In a native target cell, the cell may have been transformed to be a malignant cell or pathologically modified (e.g., a native target cell infected with a virus, a plasmodium, or a bacterium). In a foreign target cell, the cell is an invading pathogen, such as a bacterium, a plasmodium, or a virus.

[0094] The target antigen is expressed on a target cell in a diseased condition (e.g., an inflammatory disease, a proliferative disease (e.g., cancer), an immunological disorder, a neurological disease, a neurodegenerative disease, an autoimmune disease, an infectious disease (e.g., a viral infection or a parasitic infection), an allergic reaction, a graft-versus-host disease or a host-versus-graft disease). A target antigen is not effector antigen. In embodiments of the invention, the target antigen is EGFRVIII.

[0095] In some examples, provided is a bispecific antibody wherein the bispecific antibody is a full-length human antibody, comprising a first antibody variable domain of the bispecific antibody specifically binding to a target antigen, and comprising a second antibody variable domain of the bispecific antibody capable of recruiting the activity of a human immune effector cell by specifically binding to an effector antigen located on the human immune effector cell, wherein the first antibody variable domain comprises a heavy chain variable (VH) region comprising a VH CDR1, VH CDR2, and VH CDR3 of the VH sequence shown in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 30, 32, 34, 35, 37, 39, 41, 43, 44, 46, 48, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 202, 203, 204, 205, 206, 207, 208, 209, 210, 214, 216, 217, or 218; and/or a light chain variable (VL) region comprising a VL CDR1, VL CDR2, and VL CDR3 of the VL sequence shown in SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 36, 38, 40, 42, 45, 47, 49, 51, 211, 212, 213, or 215.

[0096] In some examples, provided is a bispecific antibody wherein the bispecific antibody is a full-length human antibody, comprising a first antibody variable domain of the bispecific antibody specifically binding to a target antigen, and comprising a second antibody variable domain of the bispecific antibody capable of recruiting the activity of a human immune effector cell by specifically binding to an effector antigen located on the human immune effector cell, wherein the first antibody variable domain comprises (a) a heavy chain variable (VH) region comprising (i) a VH complementarity determining region one (CDR1) comprising the sequence shown in SEQ ID NO: 62, 63, 64, 74, 75, 76, 80, 81, 82, 88, 89, 90, 93, 94, 95, 99, 100, 101, 109, 110, 111, 115, 116, 117, 121, 122, 123, 132, 133, 134, 137, 138, 139, 143, 144, or 145; (ii) a VH CDR2 comprising the sequence shown in SEQ ID NO: 65, 66, 68, 69, 70, 71, 77, 78, 83, 84, 86, 87, 91, 92, 96, 97, 98, 102, 103, 105, 106, 112, 113, 118, 119, 124, 125, 127, 128, 130, 131, 135, 136, 140, 141, 146, 147, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, or 237; and (iii) a VH CDR3 comprising the sequence shown in SEQ ID NO: 67, 72, 73, 79, 85, 104, 107, 108, 114, 120, 126, 129, 142, 148, 219, 220, 221, 222, 223, or 236; and/or (b) a light chain variable (VL) region comprising (i) a VL CDR1 comprising the sequence shown in SEQ ID NO: 149, 154, 156, 159, 162, 165, 166, 168, 169, 170, 171, 173, 174, 176, 178, 181, 182, 185, 187, 190, 192, 195, 198, 238, or 239; (ii) a VL CDR2 comprising the sequence shown in SEQ ID NO: 150, 152, 155, 157, 160, 163, 172, 175, 179, 183, 186, 188, 191, 193, 196, or 199; and (iii) a VL CDR3 comprising the sequence shown in SEQ ID NO: 151, 153, 158, 161, 164, 167, 177, 180, 184, 189, 194, 197, or 200.

[0097] In some examples, the second antibody variable domain comprises a heavy chain variable (VH) region comprising a VH CDR1, VH CDR2, and VH CDR3 of the VH sequence shown in SEQ ID NO: 240; and/or a light chain variable (VL) region comprising VL CDR1, VL CDR2, and VL CDR3 of the VL sequence shown in SEQ ID NO: 241.

[0098] In some embodiments, the second antibody variable domain comprises (a) a heavy chain variable (VH) region comprising (i) a VH complementary determining region one (CDR1) comprising the sequence shown in SEQ ID NO: 244, 110, or 245; (ii) a VH CDR2 comprising the sequence shown in SEQ ID NO: 246 or 247; and (iii) a VH CDR3 comprising the sequence shown in SEQ ID NO: 248; and/or (b) a light chain variable (VL) region comprising (i) a VL CDR1 comprising the sequence shown in SEQ ID NO: 249; (ii) a VL CDR2 comprising the sequence shown in SEQ ID NO: 250; and (iii) a VL CDR3 comprising the sequence shown in SEQ ID NO: 251.

[0099] Table 3 shows the specific amino acid and nucleic acid sequences of the second antibody variable domain, which is specific to CD3. In Table 3, the underlined sequences are CDR sequences according to Kabat and in bold according to Chothia.

Table 3

mAb	Light Chain	Heavy Chain
h2B4_HNPS_VH 1d_T2 4K_VL	DIVMTQSPDLSAVSLGERATINC KSSQSLFNVRSRKNYLAWYQQK PGQPPKLLIS WASTRES GVPDF SGSGSGTDFTLTISSLQAEDVAV YYC KQSYDL FTFGSGTKLEIK (SEQ ID NO: 241)	EVQLVESGGGLVQPGGSLRLSCA ASGFTFSD YYMTWVRQAPGKGLE WVA FIRN ARGYTSDH NPSVKGR FTISRDNAKNSLYLQMNSLRAEDT AVYYCARD DRPSYYVL DYWGQGT VTVSS (SEQ ID NO: 240)
h2B4_HNPS_VH 1d_T2 4K_VL	GACATTGTGATGACTCAATCCC CCGACTCCCTGGCTGTGTCCCT CGGCGAAGCGGCAACTATCAAC TGTAAGAAGCAGCCAGTCCCTGT TCAACGTCCGTCGAGGAAGAA CTACCTGGCCTGGTATCAGCAG AAACCTGGCAGCCGCCGAAG CTTCTGATCTCATGGCCTCAA CTCGGAAAGCGGAGTGCCAG ATAGATTCTCCGGATCTGGCTC CGGAACCGACTTCACCCCTGACG ATTTCGAGCTTGCAAGCGGAGG ATGTGGCCGTGTACTACTGCAA GCAGTCCTACGACCTCTTCACC TTTGGTTCGGGCACCAAGCTGG AGATCAA (SEQ ID NO: 243)	GAAGTCCAACTTGTGGAATCGGG AGGAGGCCCTTGTGCAACCCGGT GGATCCCTGAGGCTGTCATGCG CGCCTCGGGCTTACCTTTTCC GATTACTACATGACCTGGTCAG ACAGGCCCTGGAAAGGGGTG GAATGGGTGGCATTATCCGGA ATAGAGCCCGCGGATACACTTCC GACCACAACCCAGCGTGAAGG GGCGGTTCAACATTAGCCGCGA CAACGCCAAGAACTCCCTCTACC TCCAAATGAACAGCCTGCGGGC GGAGGATACCGCTGTGTACTACT GCGCCCGCAGCCGGCCGTCCTA CTATGTGCTGGACTIONTGGGGC CAGGGTACTACGGTCAACCGTCT CCTCA (SEQ ID NO: 242)

[0100] Table 4 shows the examples of CDR sequences of the second antibody variable domain, which is specific to CD3.

Table 4

Heavy Chain			
mAb	CDRH1	CDRH2	CDRH3
h2B4_H NPS	SDYYMT (SEQ ID NO: 244) (Kabat); GFTFSD (SEQ ID NO: 110) (Chothia); GFTFSDYYMT (SEQ ID NO: 245) (Extended)	FIRNRARGYTSDH (SEQ ID NO: 246) (Kabat) FIRNRARGYTSDHNPSVKG (SEQ ID NO: 247) (Extended)	DRPSYYVLDY (SEQ ID NO: 248)
Light Chain			
mAb	CDRH1	CDRH2	CDRH3
h2B4-1d_ T24 K	KSSQSLFNVRSRKN YLA (SEQ ID NO: 249)	WASTRES (SEQ ID NO: 250)	KQSYDLFT (SEQ ID NO: 251)

[0101] In some embodiments, a bispecific antibody provided herein which contains a CD3-specific variable domain contains an anti-CD3 sequence as provided in U.S. Publication No. 20160297885.

[0102] According to one approach to making bispecific antibodies, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant region sequences. The fusion preferably is with an immunoglobulin heavy chain constant region, comprising at least part of the hinge, CH2 and CH3 regions. It is preferred to have the first heavy chain constant region (CH1), containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are cotransfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

[0103] In another approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. This asymmetric structure, with an immunoglobulin light chain in only one half of the bispecific molecule, facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations. This approach is described in PCT Publication No. WO 94/04690.

[0104] In another approach, the bispecific antibodies are composed of amino acid modification in the first hinge region in one arm, and the substituted/replaced amino acid in the first hinge region has an opposite charge to the corresponding amino acid in the second hinge region in another arm. This approach is described in International Patent Application No. PCT/US2011/036419 (WO2011/143545).

[0105] In another approach, the formation of a desired heteromultimeric or heterodimeric protein (e.g., bispecific antibody) is enhanced by altering or engineering an interface between a first and a second immunoglobulin-like Fc region (e.g., a hinge region and/or a CH3 region). In this approach, the bispecific antibodies may be composed of a CH3 region, wherein the CH3 region comprises a first CH3 polypeptide and a second CH3 polypeptide which interact together to form a CH3 interface, wherein one or more amino acids within the CH3 interface destabilize homodimer formation and are not electrostatically unfavorable to homodimer formation. This approach is described in International Patent Application No. PCT/US2011/036419 (WO2011/143545).

[0106] In another approach, the bispecific antibodies can be generated using a glutamine-containing peptide tag engineered to the antibody directed to an epitope (e.g., EGFRvIII) in one arm and another peptide tag (e.g., a Lys-containing peptide tag or a reactive endogenous Lys) engineered to a second antibody directed to a second epitope in another arm in the presence of transglutaminase. This approach is described in International Patent Application No. PCT/IB2011/054899 (WO2012/059882).

[0107] In some embodiments, the heterodimeric protein (e.g., bispecific antibody) as described herein comprises a

full-length human antibody, wherein a first antibody variable domain of the bispecific antibody specifically binding to a target antigen (e.g., EGFRvIII), and comprising a second antibody variable domain of the bispecific antibody capable of recruiting the activity of a human immune effector cell by specifically binding to an effector antigen (e.g., CD3) located on the human immune effector cell, wherein the first and second antibody variable domain of the heterodimeric protein comprise amino acid modifications at positions 223, 225, and 228 (e.g., (C223E or C223R), (E225R), and (P228E or P228R)) in the hinge region and at position 409 or 368 (e.g., K409R or L368E (EU numbering scheme)) in the CH3 region of human IgG2 (SEQ ID NO: 290).

[0108] In some embodiments, the first and second antibody variable domains of the heterodimeric protein comprise amino acid modifications at positions 221 and 228 (e.g., (D221R or D221E) and (P228R or P228E)) in the hinge region and at position 409 or 368 (e.g., K409R or L368E (EU numbering scheme)) in the CH3 region of human IgG1 (SEQ ID NO: 291).

[0109] In some embodiments, the first and second antibody variable domains of the heterodimeric protein comprise amino acid modifications at positions 228 (e.g., (P228E or P228R)) in the hinge region and at position 409 or 368 (e.g., R409 or L368E (EU numbering scheme)) in the CH3 region of human IgG4 (SEQ ID NO: 292).

[0110] The amino acid sequence of the wild type Fc regions of human IgG1, IgG2, and IgG4 are listed below:

IgG2 (SEQ ID NO: 290)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
SSGLYSLSSVVTVPSSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECPPCPAPPVA
GPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPRE
EQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTL
PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPMLDSDGSFFLYS
KLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

IgG1 (SEQ ID NO: 291)

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL
QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAP
ELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTK
PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQV
YTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFL
YSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

IgG4 (SEQ ID NO: 292)

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
SSGLYSLSSVVTVPSSSLGTQTYTCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEFLG
GPSVFLFPPKPKDTLMISRTPEVTCVVVDVSDQEDPEVQFNWYVDGVEVHNAKTKPRE
EQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTL
PPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYS
RLTVDKSRWQEGNVFCFSVMHEALHNHYTQKSLSLSPGK

[0111] The antibodies useful in the present invention can encompass monoclonal antibodies, polyclonal antibodies, antibody fragments (e.g., Fab, Fab', F(ab')₂, Fv, Fc, etc.), chimeric antibodies, bispecific antibodies, heteroconjugate antibodies, single chain (ScFv), mutants thereof, fusion proteins comprising an antibody portion (e.g., a domain antibody), humanized antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen recognition site of the required specificity, including glycosylation variants of antibodies, amino acid sequence variants of antibodies, and covalently modified antibodies. The antibodies may be murine, rat, human, or any other origin (including chimeric or humanized antibodies).

[0112] In some embodiments, the EGFRvIII antibody as described herein is a monoclonal antibody. For example, the

EGFRvIII antibody is a humanized monoclonal antibody or a chimeric monoclonal antibody.

[0113] In some embodiments, the antibody comprises a modified constant region, such as, for example without limitation, a constant region that has increased potential for provoking an immune response. For example, the constant region may be modified to have increased affinity to an Fc gamma receptor such as, e.g., FcγRI, FcγRIIA, or FcγRIII.

[0114] In some embodiments, the antibody comprises a modified constant region, such as a constant region that is immunologically inert, that is, having a reduced potential for provoking an immune response. In some embodiments, the constant region is modified as described in Eur. J. Immunol., 29:2613-2624, 1999; PCT Application No. PCT/GB99/01441; and/or UK Patent Application No. 98099518. The Fc can be human IgG1, human IgG2, human IgG3, or human IgG4. The Fc can be human IgG2 containing the mutation A330P331 to S330S331 (IgG2Δa), in which the amino acid residues are numbered with reference to the wild type IgG2 sequence. Eur. J. Immunol., 29:2613-2624, 1999. In some embodiments, the antibody comprises a constant region of IgG4 comprising the following mutations (Armour et al., Molecular Immunology 40 585-593, 2003): E233F234L235 to P233V234A235 (IgG4Δc), in which the numbering is with reference to wild type IgG4. In yet another embodiment, the Fc is human IgG4 E233F234L235 to P233V234A235 with deletion G236 (IgG4Δb). In another embodiment, the Fc is any human IgG4 Fc (IgG4, IgG4Δb or IgG4Δc) containing hinge stabilizing mutation S228 to P228 (Aalberse et al., Immunology 105, 9-19, 2002). In another embodiment, the Fc can be aglycosylated Fc.

[0115] In some embodiments, the constant region is aglycosylated by mutating the oligosaccharide attachment residue (such as Asn297) and/or flanking residues that are part of the glycosylation recognition sequence in the constant region. In some embodiments, the constant region is aglycosylated for N-linked glycosylation enzymatically. The constant region may be aglycosylated for N-linked glycosylation enzymatically or by expression in a glycosylation deficient host cell.

[0116] In some embodiments, the constant region has a modified constant region that removes or reduces Fc gamma receptor binding. For example, the Fc can be human IgG2 containing the mutation D265, in which the amino acid residues are numbered with reference to the wild type IgG2 sequence (SEQ ID NO: 290). Accordingly, in some embodiments, the constant region has a modified constant region having the sequence shown in SEQ ID NO: 252:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
SSGLYSLSVVTVPSNFGTQTYTCNVDHKPSNTKVDKTVRKCRVRCPRCPAPPVA
GPSVFLFPPKPKDTLMISRTPEVTCVAVVAVSHEDPEVQFNWYVDGVEVHNAKTKPRE
EQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPSSIEKTIKTKGQPREPQVYTL
PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPMLDSDGSFFLYS
RLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK.

[0117] The nucleic acid encoding the sequence shown in SEQ ID NO: 252 is shown in SEQ ID NO: 253:

GCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACC
TCCGAGAGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCG
GTGACGGTGTCTGTGGAACCTCAGGCGCTCTGACCAGCGGCGTGACACCTTCCCG
GCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTAGTGACCGTGCCCT
CCAGCAACTTCGGCACCCAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAA
CACCAAGGTGGACAAGACAGTTGAGCGCAAATGTCGTGTCAGGTGCCCAAGGTG
CCCAGCACCACTGTGGCAGGACCGTCAGTCTTCTTCCCCCAAAACCCAAG
GACACCCTCATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGCCGTG
AGCCACGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTG
CATAATGCCAAGACAAAGCCACGGGAGGAGCAGTTCAACAGCACGTTCCGTGTGG

TCAGCGTCCTCACCGTCGTGCACCAGGACTGGCTGAACGGCAAGGAGTACAAGT
 GCAAGGTCTCCAACAAAGGCCTCCCATCCTCCATCGAGAAAACCATCTCCAAAAC
 5 CAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGA
 GATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCCAGC
 GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACC
 10 ACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAGGCTCACCG
 TGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGA
 GGCTCTGCACAACCACTACACACAGAAGAGCCTCTCCCTGTCTCCGGGTAAA.

[0118] In some embodiments, the constant region has a modified constant region having the sequence shown in SEQ ID NO: 254:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
 SSGLYSLSSVVTVPSSNFGTQTYTCNVDPHKPSNTKVDKTVKCEVECEPCPAPPVA
 GPSVFLFPPKPKDTLMISRTPEVTCVAVSHEDPEVQFNWYVDGVEVHNAKTKPRE
 25 EQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPSSIEKTKGQPREPQVYTL
 PPSREEMTKNQVSLTCEVKGFYPSDIAVEWESNGQPENNYKTPPMLDSDGSFFLYS
 KLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK.

[0119] The nucleic acid encoding the sequence shown in SEQ ID NO: 254 is shown in SEQ ID NO: 255:

GCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAGCACC
 35 TCCGAGAGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCG
 GTGACGGTGTCTGTGGAACCTCAGGCGCTCTGACCAGCGGCGTGCACACCTTCCCG
 GCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTAGTGACCGTGCCCT
 40 CCAGCAACTTCGGCACCCAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAA
 CACCAAGGTGGACAAGACAGTTGAGCGCAAATGTGAGGTCGAGTGCCCAGAGTG
 CCCAGCACCACTGTGGCAGGACCGTCAGTCTTCTCTTCCCCCCTTCCCCAAACCAAG
 45 GACACCCTCATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGCCGTG
 AGCCACGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTG
 CATAATGCCAAGACAAAGCCACGGGAGGAGCAGTTCAACAGCACGTTCCGTGTGG
 50 TCAGCGTCCTCACCGTCGTGCACCAGGACTGGCTGAACGGCAAGGAGTACAAGT
 GCAAGGTCTCCAACAAAGGCCTCCCATCCTCCATCGAGAAAACCATCTCCAAAAC
 CAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGA
 55 GATGACCAAGAACCAGGTCAGCCTGACCTGCGAGGTCAAAGGCTTCTACCCCAGC

GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACC
 ACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCG
 5 TGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGA
 GGCTCTGCACAACCACTACACACAGAAGAGCCTCTCCCTGTCTCCGGGTAAA.

[0120] The amino acid of the human Kappa constant region is shown in SEQ ID NO: 256:

10 GTVAAPSVFIFPPSDEQLKSGTASVVCLLNFPREAKVQWKVDNALQSGNSQESVT
 EQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC. And the nucleic acid encoding the se-
 quence of SEQ ID NO: 256 is shown in SEQ ID NO: 257:

GGAAGTGTGGCTGCACCATCTGTCTTTCATCTTCCCGCCATCTGATGAGCAGTTGAA
 15 ATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCA
 AAGTACAGTGGAAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGT
 20 CACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGCT
 GAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAG
 GGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTAG.

25 **[0121]** One way of determining binding affinity of antibodies to EGFRvIII is by measuring binding affinity of the bivalent
 antibody to monomeric EGFRvIII protein. The affinity of an EGFRvIII antibody can be determined by surface plasmon
 resonance (Biacore™3000™ surface plasmon resonance (SPR) system, Biacore™, INC, Piscataway NJ) equipped with
 pre-immobilized anti-mouse Fc or anti-human Fc using HBS-EP running buffer (0.01M HEPES, pH 7.4, 0.15 NaCl, 3
 mM EDTA, 0.005% v/v Surfactant P20). Monomeric 8-histidine tagged human EGFRvIII extracellular domain can be
 30 diluted into HBS-EP buffer to a concentration of less than 0.5 µg/mL and injected across the individual chip channels
 using variable contact times, to achieve two ranges of antigen density, either 50-200 response units (RU) for detailed
 kinetic studies or 800-1,000 RU for screening assays. Regeneration studies have shown that 25 mM NaOH in 25% v/v
 ethanol effectively removes the bound EGFRvIII protein while keeping the activity of EGFRvIII antibodies on the chip for
 over 200 injections. Typically, serial dilutions (spanning concentrations of 0.1-10x estimated K_D) of purified 8-histidine
 35 tagged EGFRvIII samples are injected for 1 min at 100 µL/minute and dissociation times of up to 2 hours are allowed.
 The concentrations of the EGFRvIII proteins are determined by absorbance at 280nm based on sequence specific
 extinction coefficient of the 8-histidine tagged EGFRvIII protein. Kinetic association rates (k_{on} or k_a) and dissociation
 rates (k_{off} or k_d) are obtained simultaneously by fitting the data globally to a 1:1 Langmuir binding model (Karlsson, R.
 Roos, H. Fagerstam, L. Petersson, B. (1994). Methods Enzymology 6. 99-110) using the BIAevaluation program. Equi-
 40 librium dissociation constant (K_D) values are calculated as k_{off}/k_{on} . This protocol is suitable for use in determining binding
 affinity of an antibody to any monomeric EGFRvIII, including human EGFRvIII, EGFRvIII of another mammal (such as
 mouse EGFRvIII, rat EGFRvIII, or primate EGFRvIII), as well as different forms of EGFRvIII (e.g., glycosylated EGFRvIII).
 Binding affinity of an antibody is generally measured at 25°C, but can also be measured at 37°C.

[0122] The antibodies as described herein may be made by any method known in the art. For the production of
 45 hybridoma cell lines, the route and schedule of immunization of the host animal are generally in keeping with established
 and conventional techniques for antibody stimulation and production, as further described herein. General techniques
 for production of human and mouse antibodies are known in the art and/or are described herein.

[0123] It is contemplated that any mammalian subject including humans or antibody producing cells therefrom can be
 manipulated to serve as the basis for production of mammalian, including human and hybridoma cell lines. Typically,
 50 the host animal is inoculated intraperitoneally, intramuscularly, orally, subcutaneously, intraplantar, and/or intradermally
 with an amount of immunogen, including as described herein.

[0124] Hybridomas can be prepared from the lymphocytes and immortalized myeloma cells using the general somatic
 cell hybridization technique of Kohler, B. and Milstein, C., Nature 256:495-497, 1975 or as modified by Buck, D. W., et
 al., In Vitro, 18:377-381, 1982. Available myeloma lines, including but not limited to X63-Ag8.653 and those from the
 55 Salk Institute, Cell Distribution Center, San Diego, Calif., USA, may be used in the hybridization. Generally, the technique
 involves fusing myeloma cells and lymphoid cells using a fusogen such as polyethylene glycol, or by electrical means
 well known to those skilled in the art. After the fusion, the cells are separated from the fusion medium and grown in a
 selective growth medium, such as hypoxanthineaminopterin-thymidine (HAT) medium, to eliminate unhybridized parent

cells. Any of the media described herein, supplemented with or without serum, can be used for culturing hybridomas that secrete monoclonal antibodies. As another alternative to the cell fusion technique, EBV immortalized B cells may be used to produce the monoclonal antibodies of the subject invention. The hybridomas are expanded and subcloned, if desired, and supernatants are assayed for anti-immunogen activity by conventional immunoassay procedures (e.g.,

radioimmunoassay, enzyme immunoassay, or fluorescence immunoassay).

[0125] Hybridomas that may be used as source of antibodies encompass all derivatives, progeny cells of the parent hybridomas that produce monoclonal antibodies specific for EGFRvIII, or portions thereof.

[0126] Hybridomas that produce such antibodies may be grown *in vitro* or *in vivo* using known procedures. The monoclonal antibodies may be isolated from the culture media or body fluids, by conventional immunoglobulin purification procedures such as ammonium sulfate precipitation, gel electrophoresis, dialysis, chromatography, and ultrafiltration, if desired. Undesired activity, if present, can be removed, for example, by running the preparation over adsorbents made of the immunogen attached to a solid phase and eluting or releasing the desired antibodies off the immunogen. Immunization of a host animal with cells expressing human EGFRvIII, a human EGFRvIII protein, or a fragment containing the target amino acid sequence conjugated to a protein that is immunogenic in the species to be immunized, e.g., keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfo succinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups, can yield a population of antibodies (e.g., monoclonal antibodies).

[0127] If desired, the antibody (monoclonal or polyclonal) of interest may be sequenced and the polynucleotide sequence may then be cloned into a vector for expression or propagation. The sequence encoding the antibody of interest may be maintained in vector in a host cell and the host cell can then be expanded and frozen for future use. Production of recombinant monoclonal antibodies in cell culture can be carried out through cloning of antibody genes from B cells by means known in the art. See, e.g. Tiller et al., J. Immunol. Methods 329, 112, 2008; U.S. Pat. No. 7,314,622.

[0128] In an alternative, the polynucleotide sequence may be used for genetic manipulation to "humanize" the antibody or to improve the affinity, or other characteristics of the antibody. For example, the constant region may be engineered to more nearly resemble human constant regions to avoid immune response if the antibody is used in clinical trials and treatments in humans. It may be desirable to genetically manipulate the antibody sequence to obtain greater affinity to EGFRvIII and greater efficacy in inhibiting EGFRvIII.

[0129] There are four general steps to humanize a monoclonal antibody. These are: (1) determining the nucleotide and predicted amino acid sequence of the starting antibody light and heavy variable domains (2) designing the humanized antibody, i.e., deciding which antibody framework region to use during the humanizing process (3) the actual humanizing methodologies/techniques and (4) the transfection and expression of the humanized antibody. See, for example, U.S. Pat. Nos. 4,816,567; 5,807,715; 5,866,692; 6,331,415; 5,530,101; 5,693,761; 5,693,762; 5,585,089; and 6,180,370.

[0130] A number of "humanized" antibody molecules comprising an antigen binding site derived from a non-human immunoglobulin have been described, including chimeric antibodies having rodent or modified rodent V regions and their associated CDRs fused to human constant regions. See, for example, Winter et al. Nature 349:293-299, 1991, Lobuglio et al. Proc. Nat. Acad. Sci. USA 86:4220-4224, 1989, Shaw et al. J Immunol. 138:4534-4538, 1987, and Brown et al. Cancer Res. 47:3577-3583, 1987. Other references describe rodent CDRs grafted into a human supporting framework region (FR) prior to fusion with an appropriate human antibody constant region. See, for example, Riechmann et al. Nature 332:323-327, 1988, Verhoeyen et al. Science 239:1534-1536, 1988, and Jones et al. Nature 321:522-525, 1986. Another reference describes rodent CDRs supported by recombinantly engineered rodent framework regions. See, for example, European Patent Publication No. 0519596. These "humanized" molecules are designed to minimize unwanted immunological response toward rodent anti-human antibody molecules which limits the duration and effectiveness of therapeutic applications of those moieties in human recipients. For example, the antibody constant region can be engineered such that it is immunologically inert (e.g., does not trigger complement lysis). See, e.g. PCT Publication No. PCT/GB99/01441; UK Patent Application No. 9809951.8. Other methods of humanizing antibodies that may also be utilized are disclosed by Daugherty et al., Nucl. Acids Res. 19:2471-2476, 1991, and in U.S. Pat. Nos. 6,180,377; 6,054,297; 5,997,867; 5,866,692; 6,210,671; and 6,350,861; and in PCT Publication No. WO 01/27160.

[0131] The general principles related to humanized antibodies discussed above are also applicable to customizing antibodies for use, for example, in dogs, cats, primate, equines and bovines. Further, one or more aspects of humanizing an antibody described herein may be combined, e.g., CDR grafting, framework mutation and CDR mutation.

[0132] In one variation, fully human antibodies may be obtained by using commercially available mice that have been engineered to express specific human immunoglobulin proteins. Transgenic animals that are designed to produce a more desirable (e.g., fully human antibodies) or more robust immune response may also be used for generation of humanized or human antibodies. Examples of such technology are Xenomouse™ from Abgenix, Inc. (Fremont, CA) and HuMAb-Mouse® and TC Mouse™ from Medarex, Inc. (Princeton, NJ).

[0133] In an alternative, antibodies may be made recombinantly and expressed using any method known in the art. In another alternative, antibodies may be made recombinantly by phage display technology. See, for example, U.S. Pat.

Nos. 5,565,332; 5,580,717; 5,733,743; and 6,265,150; and Winter et al., *Annu. Rev. Immunol.* 12:433-455, 1994. Alternatively, the phage display technology (McCafferty et al., *Nature* 348:552-553, 1990) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B cell. Phage display can be performed in a variety of formats; for review see, e.g., Johnson, Kevin S. and Chiswell, David J., *Current Opinion in Structural Biology* 3:564-571, 1993. Several sources of V-gene segments can be used for phage display. Clackson et al., *Nature* 352:624-628, 1991, isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Mark et al., *J. Mol. Biol.* 222:581-597, 1991, or Griffith et al., *EMBO J.* 12:725-734, 1993. In a natural immune response, antibody genes accumulate mutations at a high rate (somatic hypermutation). Some of the changes introduced will confer higher affinity, and B cells displaying high-affinity surface immunoglobulin are preferentially replicated and differentiated during subsequent antigen challenge. This natural process can be mimicked by employing the technique known as "chain shuffling." (Marks et al., *Bio/Technol.* 10:779-783, 1992). In this method, the affinity of "primary" human antibodies obtained by phage display can be improved by sequentially replacing the heavy and light chain V region genes with repertoires of naturally occurring variants (repertoires) of V domain genes obtained from unimmunized donors. This technique allows the production of antibodies and antibody fragments with affinities in the pM-nM range. A strategy for making very large phage antibody repertoires (also known as "the mother-of-all libraries") has been described by Waterhouse et al., *Nucl. Acids Res.* 21:2265-2266, 1993. Gene shuffling can also be used to derive human antibodies from rodent antibodies, where the human antibody has similar affinities and specificities to the starting rodent antibody. According to this method, which is also referred to as "epitope imprinting", the heavy or light chain V domain gene of rodent antibodies obtained by phage display technique is replaced with a repertoire of human V domain genes, creating rodent-human chimeras. Selection on antigen results in isolation of human variable regions capable of restoring a functional antigen binding site, i.e., the epitope governs (imprints) the choice of partner. When the process is repeated in order to replace the remaining rodent V domain, a human antibody is obtained (see PCT Publication No. WO 93/06213). Unlike traditional humanization of rodent antibodies by CDR grafting, this technique provides completely human antibodies, which have no framework or CDR residues of rodent origin.

[0134] Antibodies may be made recombinantly by first isolating the antibodies and antibody producing cells from host animals, obtaining the gene sequence, and using the gene sequence to express the antibody recombinantly in host cells (e.g., CHO cells). Another method which may be employed is to express the antibody sequence in plants (e.g., tobacco) or transgenic milk. Methods for expressing antibodies recombinantly in plants or milk have been disclosed. See, for example, Peeters, et al. *Vaccine* 19:2756, 2001; Lonberg, N. and D. Huszar *Int. Rev. Immunol* 13:65, 1995; and Pollock, et al., *J Immunol Methods* 231:147, 1999. Methods for making derivatives of antibodies, e.g., humanized, single chain, etc. are known in the art.

[0135] Immunoassays and flow cytometry sorting techniques such as fluorescence activated cell sorting (FACS) can also be employed to isolate antibodies that are specific for EGFRvIII, or tumor antigens of interest.

[0136] The antibodies as described herein can be bound to many different carriers. Carriers can be active and/or inert. Examples of well-known carriers include polypropylene, polystyrene, polyethylene, dextran, nylon, amylases, glass, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation. In some embodiments, the carrier comprises a moiety that targets the myocardium.

[0137] DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors (such as expression vectors disclosed in PCT Publication No. WO 87/04462), which are then transfected into host cells such as E. coli cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. See, e.g., PCT Publication No. WO 87/04462. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant regions in place of the homologous murine sequences, Morrison et al., *Proc. Nat. Acad. Sci.* 81:6851, 1984, or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. In that manner, "chimeric" or "hybrid" antibodies are prepared that have the binding specificity of a monoclonal antibody herein.

[0138] The EGFRvIII antibodies as described herein can be identified or characterized using methods known in the

art, whereby reduction of EGFRvIII expression levels are detected and/or measured. In some embodiments, an EGFRvIII antibody is identified by incubating a candidate agent with EGFRvIII and monitoring binding and/or attendant reduction of EGFRvIII expression levels. The binding assay may be performed with purified EGFRvIII polypeptide(s), or with cells naturally expressing, or transfected to express, EGFRvIII polypeptide(s). In one embodiment, the binding assay is a competitive binding assay, where the ability of a candidate antibody to compete with a known EGFRvIII antibody for EGFRvIII binding is evaluated. The assay may be performed in various formats, including the ELISA format.

[0139] Following initial identification, the activity of a candidate EGFRvIII antibody can be further confirmed and refined by bioassays, known to test the targeted biological activities. Alternatively, bioassays can be used to screen candidates directly. Some of the methods for identifying and characterizing antibodies are described in detail in the Examples.

[0140] EGFRvIII antibodies may be characterized using methods well known in the art. For example, one method is to identify the epitope to which it binds, or "epitope mapping." There are many methods known in the art for mapping and characterizing the location of epitopes on proteins, including solving the crystal structure of an antibody-antigen complex, competition assays, gene fragment expression assays, and synthetic peptide-based assays, as described, for example, in Chapter 11 of Harlow and Lane, *Using Antibodies, a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1999. In an additional example, epitope mapping can be used to determine the sequence to which an antibody binds. Epitope mapping is commercially available from various sources, for example, Pepscan Systems (Edelhertweg 15, 8219 PH Lelystad, The Netherlands). The epitope can be a linear epitope, i.e., contained in a single stretch of amino acids, or a conformational epitope formed by a three-dimensional interaction of amino acids that may not necessarily be contained in a single stretch. Peptides of varying lengths (e.g., at least 4-6 amino acids long) can be isolated or synthesized (e.g., recombinantly) and used for binding assays with an EGFRvIII or other tumor antigen antibody. In another example, the epitope to which the EGFRvIII antibody binds can be determined in a systematic screening by using overlapping peptides derived from the EGFRvIII sequence and determining binding by the EGFRvIII antibody. According to the gene fragment expression assays, the open reading frame encoding EGFRvIII is fragmented either randomly or by specific genetic constructions and the reactivity of the expressed fragments of EGFRvIII with the antibody to be tested is determined. The gene fragments may, for example, be produced by PCR and then transcribed and translated into protein in vitro, in the presence of radioactive amino acids. The binding of the antibody to the radioactively labeled EGFRvIII is then determined by immunoprecipitation and gel electrophoresis. Certain epitopes can also be identified by using large libraries of random peptide sequences displayed on the surface of phage particles (phage libraries). Alternatively, a defined library of overlapping peptide fragments can be tested for binding to the test antibody in simple binding assays. In an additional example, mutagenesis of an antigen binding domain, domain swapping experiments and alanine scanning mutagenesis can be performed to identify residues required, sufficient, and/or necessary for epitope binding. For example, domain swapping experiments can be performed using a mutant EGFRvIII in which various fragments of the EGFRvIII protein have been replaced (swapped) with sequences from EGFRvIII from another species (e.g., mouse), or a closely related, but antigenically distinct protein (e.g., Trop-1). By assessing binding of the antibody to the mutant EGFRvIII, the importance of the particular EGFRvIII fragment to antibody binding can be assessed. In the case of EGFRvIII specific antibody (i.e. antibody that does not bind EGFRwt (wild type) or any other proteins), epitope can be deduced from the sequence alignment of EGFRvIII to EGFRwt.

[0141] Yet another method which can be used to characterize an EGFRvIII antibody is to use competition assays with other antibodies known to bind to the same antigen, i.e., various fragments on EGFRvIII, to determine if the EGFRvIII antibody binds to the same epitope as other antibodies. Competition assays are well known to those of skill in the art.

[0142] An expression vector can be used to direct expression of an EGFRvIII antibody. One skilled in the art is familiar with administration of expression vectors to obtain expression of an exogenous protein in vivo. See, e.g., U.S. Pat. Nos. 6,436,908; 6,413,942; and 6,376,471. Administration of expression vectors includes local or systemic administration, including injection, oral administration, particle gun or catheterized administration, and topical administration. In another embodiment, the expression vector is administered directly to the sympathetic trunk or ganglion, or into a coronary artery, atrium, ventricle, or pericardium.

[0143] Targeted delivery of therapeutic compositions containing an expression vector, or subgenomic polynucleotides can also be used. Receptor-mediated DNA delivery techniques are described in, for example, Findeis et al., *Trends Biotechnol.*, 1993, 11:202; Chiou et al., *Gene Therapeutics: Methods And Applications Of Direct Gene Transfer*, J.A. Wolff, ed., 1994; Wu et al., *J. Biol. Chem.*, 263:621, 1988; Wu et al., *J. Biol. Chem.*, 269:542, 1994; Zenke et al., *Proc. Natl. Acad. Sci. USA*, 87:3655, 1990; and Wu et al., *J. Biol. Chem.*, 266:338, 1991. Therapeutic compositions containing a polynucleotide are administered in a range of about 100 ng to about 200 mg of DNA for local administration in a gene therapy protocol. Concentration ranges of about 500 ng to about 50 mg, about 1 µg to about 2 mg, about 5 µg to about 500 µg, and about 20 µg to about 100 µg of DNA can also be used during a gene therapy protocol. The therapeutic polynucleotides and polypeptides can be delivered using gene delivery vehicles. The gene delivery vehicle can be of viral or non-viral origin (see generally, Jolly, *Cancer Gene Therapy*, 1:51, 1994; Kimura, *Human Gene Therapy*, 5:845, 1994; Connelly, *Human Gene Therapy*, 1995, 1:185; and Kaplitt, *Nature Genetics*, 6:148, 1994). Expression of such coding sequences can be induced using endogenous mammalian or heterologous promoters. Expression of the coding

sequence can be either constitutive or regulated.

[0144] Viral-based vectors for delivery of a desired polynucleotide and expression in a desired cell are well known in the art. Exemplary viral-based vehicles include, but are not limited to, recombinant retroviruses (see, e.g., PCT Publication Nos. WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; WO 93/11230; WO 93/10218; WO 91/02805; U.S. Pat. Nos. 5, 219,740 and 4,777,127; GB Pat. No. 2,200,651; and EP Pat. No. 0 345 242), alphavirus-based vectors (e.g., Sindbis virus vectors, Semliki forest virus (ATCC VR-67; ATCC VR-1247), Ross River virus (ATCC VR-373; ATCC VR-1246) and Venezuelan equine encephalitis virus (ATCC VR-923; ATCC VR-1250; ATCC VR 1249; ATCC VR-532)), and adeno-associated virus (AAV) vectors (see, e.g., PCT Publication Nos. WO 94/12649, WO 93/03769; WO 93/19191; WO 94/28938; WO 95/11984 and WO 95/00655). Administration of DNA linked to killed adenovirus as described in Curiel, Hum. Gene Ther., 1992, 3:147 can also be employed.

[0145] Non-viral delivery vehicles and methods can also be employed, including, but not limited to, polycationic condensed DNA linked or unlinked to killed adenovirus alone (see, e.g., Curiel, Hum. Gene Ther., 3:147, 1992); ligand-linked DNA (see, e.g., Wu, J. Biol. Chem., 264:16985, 1989); eukaryotic cell delivery vehicles (see, e.g., U.S. Pat. No. 5,814,482; PCT Publication Nos. WO 95/07994; WO 96/17072; WO 95/30763; and WO 97/42338) and nucleic charge neutralization or fusion with cell membranes. Naked DNA can also be employed. Exemplary naked DNA introduction methods are described in PCT Publication No. WO 90/11092 and U.S. Pat. No. 5,580,859. Liposomes that can act as gene delivery vehicles are described in U.S. Pat. No. 5,422,120; PCT Publication Nos. WO 95/13796; WO 94/23697; WO 91/14445; and EP 0524968. Additional approaches are described in Philip, Mol. Cell Biol., 14:2411, 1994 and in Woffendin, Proc. Natl. Acad. Sci., 91:1581, 1994.

[0146] In some embodiments, the invention encompasses compositions, including pharmaceutical compositions, comprising antibodies described herein or made by the methods and having the characteristics described herein. As used herein, compositions comprise one or more antibodies that bind to EGFRvIII, and/or one or more polynucleotides comprising sequences encoding one or more these antibodies. These compositions may further comprise suitable excipients, such as pharmaceutically acceptable excipients including buffers, which are well known in the art.

[0147] The invention also provides methods of making any of these antibodies. The antibodies of this invention can be made by procedures known in the art. The polypeptides can be produced by proteolytic or other degradation of the antibodies, by recombinant methods (i.e., single or fusion polypeptides) as described above or by chemical synthesis. Polypeptides of the antibodies, especially shorter polypeptides up to about 50 amino acids, are conveniently made by chemical synthesis. Methods of chemical synthesis are known in the art and are commercially available. For example, an antibody could be produced by an automated polypeptide synthesizer employing the solid phase method. See also, U.S. Pat. Nos. 5,807,715; 4,816,567; and 6,331,415.

[0148] In another alternative, the antibodies can be made recombinantly using procedures that are well known in the art. In one example, a polynucleotide comprises a sequence encoding the heavy chain and/or the light chain variable regions of antibody m62G7, h62G7, h62G7-H14/L1-DV, h62G7-EQ/L6, 42G9, 32A10, 20B9, 14C11, 21E11, 49B11, 46E10, 12H6, 19A9, 21E7, 11B11, 12B2, 11F10, 17G11, 29D5, 30D8, 20E12, 26B9, 32G8, 34E7, 20G5, C6, B5, 42G9-1, 32A10-1, 20B9-1, 14C11-1, 21E11-1, 49B11-1, 46E10-1, 12H6-1, 19A9-1, 21E7-1, 11B11-1, 12B2-1, 11F10-1, 17G11-1, 29D5-1, 30D8-1, 20E12-1, 26B9-1, 32G8-1, 34E7-1, 20G5-1, C6-1, or B5-1. The sequence encoding the antibody of interest may be maintained in a vector in a host cell and the host cell can then be expanded and frozen for future use. Vectors (including expression vectors) and host cells are further described herein.

[0149] Heteroconjugate antibodies, comprising two covalently joined antibodies, are also within the scope of the invention. Such antibodies have been used to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980), and for treatment of HIV infection (PCT Publication Nos. WO 91/00360 and WO 92/200373; EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents and techniques are well known in the art, and are described in U.S. Pat. No. 4,676,980.

[0150] Chimeric or hybrid antibodies also may be prepared *in vitro* using known methods of synthetic protein chemistry, including those involving cross-linking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

[0151] In the recombinant humanized antibodies, the Fc γ portion can be modified to avoid interaction with Fc γ receptor and the complement and immune systems. The techniques for preparation of such antibodies are described in WO 99/58572. For example, the constant region may be engineered to more resemble human constant regions to avoid immune response if the antibody is used in clinical trials and treatments in humans. See, for example, U.S. Pat. Nos. 5,997,867 and 5,866,692.

[0152] The invention encompasses modifications to the antibodies and polypeptides of the invention including variants shown in Table 5, including functionally equivalent antibodies which do not significantly affect their properties and variants which have enhanced or decreased activity and/or affinity. For example, the amino acid sequence may be mutated to obtain an antibody with the desired binding affinity to EGFRvIII. Modification of polypeptides is routine practice in the art and need not be described in detail herein. Examples of modified polypeptides include polypeptides with conservative

substitutions of amino acid residues, one or more deletions or additions of amino acids which do not significantly deleteriously change the functional activity, or which mature (enhance) the affinity of the polypeptide for its ligand, or use of chemical analogs.

[0153] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue or the antibody fused to an epitope tag. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody of an enzyme or a polypeptide which increases the half-life of the antibody in the blood circulation.

[0154] Substitution variants have at least one amino acid residue in the antibody molecule removed and a different residue inserted in its place. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 5 under the heading of "conservative substitutions." If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table 5, or as further described below in reference to amino acid classes, may be introduced and the products screened. In some embodiments, substitution variants of antibodies provided herein have no more than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 conservative substitution in the VH or VL region as compared to the reference parent antibody. In embodiments of the invention, the substitutions are not within a CDR of the VH or VL region.

Table 5: Amino Acid Substitutions

Original Residue (naturally occurring amino acid)	Conservative Substitutions	Exemplary Substitutions
Ala (A)	Val	Val; Leu; Ile
Arg (R)	Lys	Lys; Gln; Asn
Asn (N)	Gln	Gln; His; Asp, Lys; Arg
Asp (D)	Glu	Glu; Asn
Cys (C)	Ser	Ser; Ala
Gln (Q)	Asn	Asn; Glu
Glu (E)	Asp	Asp; Gln
Gly (G)	Ala	Ala
His (H)	Arg	Asn; Gln; Lys; Arg
Ile (I)	Leu	Leu; Val; Met; Ala; Phe; Norleucine
Leu (L)	Ile	Norleucine; Ile; Val; Met; Ala; Phe
Lys (K)	Arg	Arg; Gln; Asn
Met (M)	Leu	Leu; Phe; Ile
Phe (F)	Tyr	Leu; Val; Ile; Ala; Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr; Phe
Tyr (Y)	Phe	Trp; Phe; Thr; Ser
Val (V)	Leu	Ile; Leu; Met; Phe; Ala; Norleucine

[0155] Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring amino acid residues are divided into groups based on common side-chain properties:

- (1) Non-polar: Norleucine, Met, Ala, Val, Leu, Ile;
- (2) Polar without charge: Cys, Ser, Thr, Asn, Gln;
- (3) Acidic (negatively charged): Asp, Glu;
- (4) Basic (positively charged): Lys, Arg;
- (5) Residues that influence chain orientation: Gly, Pro; and
- (6) Aromatic: Trp, Tyr, Phe, His.

[0156] Non-conservative substitutions are made by exchanging a member of one of these classes for another class.

[0157] Any cysteine residue not involved in maintaining the proper conformation of the antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant cross-linking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability, particularly where the antibody is an antibody fragment such as an Fv fragment.

[0158] Amino acid modifications can range from changing or modifying one or more amino acids to complete redesign of a region, such as the variable region. Changes in the variable region can alter binding affinity and/or specificity. In some examples, no more than one to five conservative amino acid substitutions are made within a CDR domain. In other examples, no more than one to three conservative amino acid substitutions are made within a CDR domain. In still other examples, the CDR domain is CDR H3 and/or CDR L3.

[0159] Modifications also include glycosylated and nonglycosylated polypeptides, as well as polypeptides with other post-translational modifications, such as, for example, glycosylation with different sugars, acetylation, and phosphorylation. Antibodies are glycosylated at conserved positions in their constant regions (Jefferis and Lund, *Chem. Immunol.* 65:111-128, 1997; Wright and Morrison, *TibTECH* 15:26-32, 1997). The oligosaccharide side chains of the immunoglobulins affect the protein's function (Boyd et al., *Mol. Immunol.* 32:1311-1318, 1996; Wittwe and Howard, *Biochem.* 29:4175-4180, 1990) and the intramolecular interaction between portions of the glycoprotein, which can affect the conformation and presented three-dimensional surface of the glycoprotein (Jefferis and Lund, *supra*; Wyss and Wagner, *Current Opin. Biotech.* 7:409-416, 1996). Oligosaccharides may also serve to target a given glycoprotein to certain molecules based upon specific recognition structures. Glycosylation of antibodies has also been reported to affect antibody-dependent cellular cytotoxicity (ADCC). In particular, CHO cells with tetracycline-regulated expression of $\beta(1,4)$ -N-acetylglucosaminyltransferase III (GnTIII), a glycosyltransferase catalyzing formation of bisecting GlcNAc, was reported to have improved ADCC activity (Umana et al., *Mature Biotech.* 17:176-180, 1999).

[0160] Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine, asparagine-X-threonine, and asparagine-X-cysteine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0161] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[0162] The glycosylation pattern of antibodies may also be altered without altering the underlying nucleotide sequence. Glycosylation largely depends on the host cell used to express the antibody. Since the cell type used for expression of recombinant glycoproteins, e.g. antibodies, as potential therapeutics is rarely the native cell, variations in the glycosylation pattern of the antibodies can be expected (see, e.g. Hse et al., *J. Biol. Chem.* 272:9062-9070, 1997).

[0163] In addition to the choice of host cells, factors that affect glycosylation during recombinant production of antibodies include growth mode, media formulation, culture density, oxygenation, pH, purification schemes and the like. Various methods have been proposed to alter the glycosylation pattern achieved in a particular host organism including introducing or overexpressing certain enzymes involved in oligosaccharide production (U.S. Pat. Nos. 5,047,335; 5,510,261 and 5,278,299). Glycosylation, or certain types of glycosylation, can be enzymatically removed from the glycoprotein, for example, using endoglycosidase H (Endo H), N-glycosidase F, endoglycosidase F1, endoglycosidase F2, endoglycosidase F3. In addition, the recombinant host cell can be genetically engineered to be defective in processing certain types of polysaccharides. These and similar techniques are well known in the art.

[0164] Other methods of modification include using coupling techniques known in the art, including, but not limited to, enzymatic means, oxidative substitution and chelation. Modifications can be used, for example, for attachment of labels for immunoassay. Modified polypeptides are made using established procedures in the art and can be screened using standard assays known in the art, some of which are described below and in the Examples.

[0165] Other antibody modifications include antibodies that have been modified as described in PCT Publication No. WO 99/58572. These antibodies comprise, in addition to a binding domain directed at the target molecule, an effector

domain having an amino acid sequence substantially homologous to all or part of a constant region of a human immunoglobulin heavy chain. These antibodies are capable of binding the target molecule without triggering significant complement dependent lysis, or cell mediated destruction of the target. In some embodiments, the effector domain is capable of specifically binding FcRn and/or FcγRIIb. These are typically based on chimeric domains derived from two or more human immunoglobulin heavy chain C_H2 domains. Antibodies modified in this manner are particularly suitable for use in chronic antibody therapy, to avoid inflammatory and other adverse reactions to conventional antibody therapy.

[0166] The invention includes affinity matured embodiments. For example, affinity matured antibodies can be produced by procedures known in the art (Marks et al., *Bio/Technology*, 10:779-783, 1992; Barbas et al., *Proc Nat. Acad. Sci., USA* 91:3809-3813, 1994; Schier et al., *Gene*, 169:147-155, 1995; Yelton et al., *J. Immunol.*, 155:1994-2004, 1995; Jackson et al., *J. Immunol.*, 154(7):3310-9, 1995; Hawkins et al., *J. Mol. Biol.*, 226:889-896, 1992; and PCT Publication No. WO2004/058184).

[0167] The following methods may be used for adjusting the affinity of an antibody and for characterizing a CDR. One way of characterizing a CDR of an antibody and/or altering (such as improving) the binding affinity of a polypeptide, such as an antibody, termed "library scanning mutagenesis". Generally, library scanning mutagenesis works as follows. One or more amino acid positions in the CDR are replaced with two or more (such as 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) amino acids using art recognized methods. This generates small libraries of clones (in some embodiments, one for every amino acid position that is analyzed), each with a complexity of two or more members (if two or more amino acids are substituted at every position). Generally, the library also includes a clone comprising the native (unsubstituted) amino acid. A small number of clones, e.g., about 20-80 clones (depending on the complexity of the library), from each library are screened for binding affinity to the target polypeptide (or other binding target), and candidates with increased, the same, decreased, or no binding are identified. Methods for determining binding affinity are well-known in the art. Binding affinity may be determined using Biacore™ surface plasmon resonance analysis, which detects differences in binding affinity of about 2-fold or greater. Biacore™ is particularly useful when the starting antibody already binds with a relatively high affinity, for example a K_D of about 10 nM or lower. Screening using Biacore™ surface plasmon resonance is described in the Examples, herein.

[0168] Binding affinity may be determined using Kinexa Biocensor, scintillation proximity assays, ELISA, ORIGEN immunoassay (IGEN), fluorescence quenching, fluorescence transfer, and/or yeast display. Binding affinity may also be screened using a suitable bioassay.

[0169] In some examples, every amino acid position in a CDR is replaced (in some examples, one at a time) with all 20 natural amino acids using art recognized mutagenesis methods (some of which are described herein). This generates small libraries of clones (in some examples, one for every amino acid position that is analyzed), each with a complexity of 20 members (if all 20 amino acids are substituted at every position).

[0170] In some examples, the library to be screened comprises substitutions in two or more positions, which may be in the same CDR or in two or more CDRs. Thus, the library may comprise substitutions in two or more positions in one CDR. The library may comprise substitution in two or more positions in two or more CDRs. The library may comprise substitution in 3, 4, 5, or more positions, said positions found in two, three, four, five or six CDRs. The substitution may be prepared using low redundancy codons. See, e.g., Table 2 of Balint et al., *Gene* 137(1):109-18, 1993.

[0171] The CDR may be CDRH3 and/or CDRL3. The CDR may be one or more of CDRL1, CDRL2, CDRL3, CDRH1, CDRH2, and/or CDRH3. The CDR may be a Kabat CDR, a Chothia CDR, or an extended CDR.

[0172] Candidates with improved binding may be sequenced, thereby identifying a CDR substitution mutant which results in improved affinity (also termed an "improved" substitution). Candidates that bind may also be sequenced, thereby identifying a CDR substitution which retains binding.

[0173] Multiple rounds of screening may be conducted. For example, candidates (each comprising an amino acid substitution at one or more position of one or more CDR) with improved binding are also useful for the design of a second library containing at least the original and substituted amino acid at each improved CDR position (i.e., amino acid position in the CDR at which a substitution mutant showed improved binding). Preparation, and screening or selection of this library is discussed further below.

[0174] Library scanning mutagenesis also provides a means for characterizing a CDR, in so far as the frequency of clones with improved binding, the same binding, decreased binding or no binding also provide information relating to the importance of each amino acid position for the stability of the antibody-antigen complex. For example, if a position of the CDR retains binding when changed to all 20 amino acids, that position is identified as a position that is unlikely to be required for antigen binding. Conversely, if a position of CDR retains binding in only a small percentage of substitutions, that position is identified as a position that is important to CDR function. Thus, the library scanning mutagenesis methods generate information regarding positions in the CDRs that can be changed to many different amino acids (including all 20 amino acids), and positions in the CDRs which cannot be changed or which can only be changed to a few amino acids.

[0175] Candidates with improved affinity may be combined in a second library, which includes the improved amino acid, the original amino acid at that position, and may further include additional substitutions at that position, depending

on the complexity of the library that is desired, or permitted using the desired screening or selection method. In addition, if desired, adjacent amino acid position can be randomized to at least two or more amino acids. Randomization of adjacent amino acids may permit additional conformational flexibility in the mutant CDR, which may in turn, permit or facilitate the introduction of a larger number of improving mutations. The library may also comprise substitution at positions that did not show improved affinity in the first round of screening.

[0176] The second library is screened or selected for library members with improved and/or altered binding affinity using any method known in the art, including screening using Biacore™ surface plasmon resonance analysis, and selection using any method known in the art for selection, including phage display, yeast display, and ribosome display.

[0177] The invention also encompasses fusion proteins comprising one or more fragments or regions from the antibodies of this invention. In one example, a fusion polypeptide is provided that comprises at least 10 contiguous amino acids of the variable light chain region shown in SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 29, 31, 33, 36, 38, 40, 42, 45, 47, 49, 51, 211, 212, 213, or 215, and/or at least 10 amino acids of the variable heavy chain region shown in SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 30, 32, 34, 35, 37, 39, 41, 43, 44, 46, 48, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 202, 203, 204, 205, 206, 207, 208, 209, 210, 214, 216, 217, or 218. In other examples, a fusion polypeptide is provided that comprises at least about 10, at least about 15, at least about 20, at least about 25, or at least about 30 contiguous amino acids of the variable light chain region and/or at least about 10, at least about 15, at least about 20, at least about 25, or at least about 30 contiguous amino acids of the variable heavy chain region. In another example, the fusion polypeptide comprises one or more CDR(s). In still other examples, the fusion polypeptide comprises CDR H3 (VH CDR3) and/or CDR L3 (VL CDR3). For purposes of this invention, a fusion protein contains one or more antibodies and another amino acid sequence to which it is not attached in the native molecule, for example, a heterologous sequence or a homologous sequence from another region. Exemplary heterologous sequences include, but are not limited to a "tag" such as a FLAG tag or a 6His tag. Tags are well known in the art.

[0178] A fusion polypeptide can be created by methods known in the art, for example, synthetically or recombinantly. Typically, the fusion proteins of this invention are made by preparing an expressing a polynucleotide encoding them using recombinant methods described herein, although they may also be prepared by other means known in the art, including, for example, chemical synthesis.

[0179] This invention also provides compositions comprising antibodies conjugated (for example, linked) to an agent that facilitate coupling to a solid support (such as biotin or avidin). For simplicity, reference will be made generally to antibodies with the understanding that these methods apply to any of the EGFRvIII antibody embodiments described herein. Conjugation generally refers to linking these components as described herein. The linking (which is generally fixing these components in proximate association at least for administration) can be achieved in any number of ways. For example, a direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (e.g., a halide) on the other.

[0180] The invention also provides isolated polynucleotides encoding the antibodies of the invention, and vectors and host cells comprising the polynucleotide.

[0181] Accordingly, the present disclosure provides polynucleotides (or compositions, including pharmaceutical compositions), comprising polynucleotides encoding any of the following: m62G7, h62G7, h62G7-H14/L1-DV, h62G7-EQ/L6, 42G9, 32A10, 20B9, 14C11, 21E11, 49B11, 46E10, 12H6, 19A9, 21E7, 11B11, 12B2, 11F10, 17G11, 29D5, 30D8, 20E12, 26B9, 32G8, 34E7, 20G5, C6, B5, 42G9-1, 32A10-1, 20B9-1, 14C11-1, 21E11-1, 49B11-1, 46E10-1, 12H6-1, 19A9-1, 21E7-1, 11B11-1, 12B2-1, 11F10-1, 17G11-1, 29D5-1, 30D8-1, 20E12-1, 26B9-1, 32G8-1, 34E7-1, 20G5-1, C6-1, and B5-1, or any fragment or part thereof having the ability to bind EGFRvIII.

[0182] In another aspect, the invention provides polynucleotides encoding any of the antibodies (including antibody fragments) and polypeptides of the invention, such as antibodies and polypeptides having impaired effector function. Polynucleotides can be made and expressed by procedures known in the art.

[0183] In another aspect, the invention provides compositions (such as a pharmaceutical compositions) comprising any of the polynucleotides of the invention. In some embodiments, the composition comprises an expression vector comprising a polynucleotide encoding any of the antibodies of the invention.

[0184] Expression vectors, and administration of polynucleotide compositions are further described herein.

[0185] In another aspect, the present disclosure provides a method of making any of the polynucleotides described herein.

[0186] Polynucleotides complementary to any such sequences are also encompassed by the present invention. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

[0187] Polynucleotides may comprise a native sequence (i.e., an endogenous sequence that encodes an antibody or a portion thereof) or may comprise a variant of such a sequence. Polynucleotide variants contain one or more substitutions, additions, deletions and/or insertions such that the immunoreactivity of the encoded polypeptide is not diminished, relative to a native immunoreactive molecule. The effect on the immunoreactivity of the encoded polypeptide may generally be assessed as described herein. Variants preferably exhibit at least about 70% identity, more preferably, at least about 80% identity, yet more preferably, at least about 90% identity, and most preferably, at least about 95% identity to a polynucleotide sequence that encodes a native antibody or a portion thereof.

[0188] Two polynucleotide or polypeptide sequences are said to be "identical" if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, or 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

[0189] Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O., 1978, A model of evolutionary change in proteins - Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J., 1990, Unified Approach to Alignment and Phylogenies pp. 626-645 Methods in Enzymology vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M., 1989, CABIOS 5:151-153; Myers, E.W. and Muller W., 1988, CABIOS 4:11-17; Robinson, E.D., 1971, Comb. Theor. 11:105; Santou, N., Nes, M., 1987, Mol. Biol. Evol. 4:406-425; Sneath, P.H.A. and Sokal, R.R., 1973, Numerical Taxonomy the Principles and Practice of Numerical Taxonomy, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J., 1983, Proc. Natl. Acad. Sci. USA 80:726-730.

[0190] Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (i.e. the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

[0191] Variants may also, or alternatively, be substantially homologous to a native gene, or a portion or complement thereof. Such polynucleotide variants are capable of hybridizing under moderately stringent conditions to a naturally occurring DNA sequence encoding a native antibody (or a complementary sequence).

[0192] Suitable "moderately stringent conditions" include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1 % SDS.

[0193] As used herein, "highly stringent conditions" or "high stringency conditions" are those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

[0194] It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

[0195] The polynucleotides of this invention can be obtained using chemical synthesis, recombinant methods, or PCR. Methods of chemical polynucleotide synthesis are well known in the art and need not be described in detail herein. One of skill in the art can use the sequences provided herein and a commercial DNA synthesizer to produce a desired DNA

sequence.

[0196] For preparing polynucleotides using recombinant methods, a polynucleotide comprising a desired sequence can be inserted into a suitable vector, and the vector in turn can be introduced into a suitable host cell for replication and amplification, as further discussed herein. Polynucleotides may be inserted into host cells by any means known in the art. Cells are transformed by introducing an exogenous polynucleotide by direct uptake, endocytosis, transfection, F-mating or electroporation. Once introduced, the exogenous polynucleotide can be maintained within the cell as a non-integrated vector (such as a plasmid) or integrated into the host cell genome. The polynucleotide so amplified can be isolated from the host cell by methods well known within the art. See, e.g., Sambrook et al., 1989.

[0197] Alternatively, PCR allows reproduction of DNA sequences. PCR technology is well known in the art and is described in U.S. Patent Nos. 4,683,195, 4,800,159, 4,754,065 and 4,683,202, as well as PCR: The Polymerase Chain Reaction, Mullis et al. eds., Birkhäuser Press, Boston, 1994.

[0198] RNA can be obtained by using the isolated DNA in an appropriate vector and inserting it into a suitable host cell. When the cell replicates and the DNA is transcribed into RNA, the RNA can then be isolated using methods well known to those of skill in the art, as set forth in Sambrook et al., 1989, supra, for example.

[0199] Suitable cloning vectors may be constructed according to standard techniques, or may be selected from a large number of cloning vectors available in the art. While the cloning vector selected may vary according to the host cell intended to be used, useful cloning vectors will generally have the ability to self-replicate, may possess a single target for a particular restriction endonuclease, and/or may carry genes for a marker that can be used in selecting clones containing the vector. Suitable examples include plasmids and bacterial viruses, e.g., pUC18, pUC19, Bluescript (e.g., pBS SK+) and its derivatives, mp18, mp19, pBR322, pMB9, ColE1, pCR1, RP4, phage DNAs, and shuttle vectors such as pSA3 and pAT28. These and many other cloning vectors are available from commercial vendors such as BioRad, Strategene, and Invitrogen.

[0200] Expression vectors generally are replicable polynucleotide constructs that contain a polynucleotide according to the invention. It is implied that an expression vector must be replicable in the host cells either as episomes or as an integral part of the chromosomal DNA. Suitable expression vectors include but are not limited to plasmids, viral vectors, including adenoviruses, adeno-associated viruses, retroviruses, cosmids, and expression vector(s) disclosed in PCT Publication No. WO 87/04462. Vector components may generally include, but are not limited to, one or more of the following: a signal sequence; an origin of replication; one or more marker genes; suitable transcriptional controlling elements (such as promoters, enhancers and terminator). For expression (i.e., translation), one or more translational controlling elements are also usually required, such as ribosome binding sites, translation initiation sites, and stop codons.

[0201] The vectors containing the polynucleotides of interest can be introduced into the host cell by any of a number of appropriate means, including electroporation, transfection employing calcium chloride, rubidium chloride, calcium phosphate, DEAE dextran, or other substances; microprojectile bombardment; lipofection; and infection (e.g., where the vector is an infectious agent such as vaccinia virus). The choice of introducing vectors or polynucleotides will often depend on features of the host cell.

[0202] The invention also provides host cells comprising any of the polynucleotides described herein. Any host cells capable of over-expressing heterologous DNAs can be used for the purpose of isolating the genes encoding the antibody, polypeptide or protein of interest. Non-limiting examples of mammalian host cells include but not limited to COS, HeLa, and CHO cells. See also PCT Publication No. WO 87/04462. Suitable non-mammalian host cells include prokaryotes (such as *E. coli* or *B. subtilis*) and yeast (such as *S. cerevisiae*, *S. pombe*; or *K. lactis*). Preferably, the host cells express the cDNAs at a level of about 5 fold higher, more preferably, 10 fold higher, even more preferably, 20 fold higher than that of the corresponding endogenous antibody or protein of interest, if present, in the host cells. Screening the host cells for a specific binding to EGFRvIII is effected by an immunoassay or FACS. A cell overexpressing the antibody or protein of interest can be identified.

EGFRvIII Antibody Conjugates

[0203] The present invention also provides a conjugate (or immunoconjugate) of the EGFRvIII antibody as described herein, wherein the antibody is conjugated to an agent (e.g., a cytotoxic agent) for targeted immunotherapy (e.g., antibody-drug conjugates) either directly or indirectly via a linker. For example, a cytotoxic agent can be linked or conjugated to the EGFRvIII antibody as described herein for targeted local delivery of the cytotoxic agent moiety to tumors (e.g., EGFRvIII expressing tumor).

[0204] Methods for conjugating cytotoxic agent or other therapeutic agents to antibodies have been described in various publications. For example, chemical modification can be made in the antibodies either through lysine side chain amines or through cysteine sulfhydryl groups activated by reducing interchain disulfide bonds for the conjugation reaction to occur. See, e.g., Tanaka et al., FEBS Letters 579:2092-2096, 2005, and Gentle et al., Bioconjugate Chem. 15:658-663, 2004. Reactive cysteine residues engineered at specific sites of antibodies for specific drug conjugation with defined stoichiometry have also been described. See, e.g., Junutula et al., Nature Biotechnology, 26:925-932, 2008. Conjugation

using an acyl donor glutamine-containing tag or an endogenous glutamine made reactive (i.e., the ability to form a covalent bond as an acyl donor) by polypeptide engineering in the presence of transglutaminase and an amine (e.g., a cytotoxic agent comprising or attached to a reactive amine) is also described in international applications WO2012/059882 and WO2015015448.

[0205] In some embodiments, the EGFRvIII antibody or the conjugate as described herein comprises an acyl donor glutamine-containing tag engineered at a specific site of the antibody (e.g., a carboxyl terminus, an amino terminus, or at another site in the EGFRvIII antibody). In some embodiments, the tag comprises an amino acid glutamine (Q) or an amino acid sequence LQG, LLQGG (SEQ ID NO: 258), LLQG (SEQ ID NO: 259), LSLSQG (SEQ ID NO: 260), GGGLLQGG (SEQ ID NO: 261), GLLQG (SEQ ID NO: 262), LLQ, GSPLAQSHGG (SEQ ID NO: 263), GLLQGGG (SEQ ID NO: 264), GLLQGG (SEQ ID NO: 265), GLLQ (SEQ ID NO: 266), LLQLLQGA (SEQ ID NO: 267), LLQGA (SEQ ID NO: 268), LLQYQGA (SEQ ID NO: 269), LLQGSG (SEQ ID NO: 270), LLQYQG (SEQ ID NO: 271), LLQLLQG (SEQ ID NO: 272), SLLQG (SEQ ID NO: 273), LLQLQ (SEQ ID NO: 274), LLQLLQ (SEQ ID NO: 275), LLQGR (SEQ ID NO: 276), LLQGPP (SEQ ID NO: 277), LLQGPA (SEQ ID NO: 278), GLLQGPP (SEQ ID NO: 279), GLLQGA (SEQ ID NO: 280), LLQGP (SEQ ID NO: 281), LLQGP (SEQ ID NO: 282), LLQGP (SEQ ID NO: 283), LLQP (SEQ ID NO: 284), LLQPGK (SEQ ID NO: 285), LLQAPGK (SEQ ID NO: 286), LLQGAPG (SEQ ID NO: 287), LLQGAP (SEQ ID NO: 288), and LLQLQG (SEQ ID NO: 289).


[0206] Also provided is an isolated antibody comprising an acyl donor glutamine-containing tag and an amino acid modification at position 222, 340, or 370 of the antibody (EU numbering scheme) wherein the modification is an amino acid deletion, insertion, substitution, mutation, or any combination thereof. In some embodiments, the amino acid modification is a substitution from lysine to arginine (e.g., K222R, K340R, or K370R).

[0207] The agents that can be conjugated to the EGFRvIII antibodies of the present invention include, but are not limited to, cytotoxic agents, immunomodulating agents, imaging agents, therapeutic proteins, biopolymers, or oligonucleotides.


[0208] Examples of a cytotoxic agent include, but are not limited to, anthracycline, an auristatin, a dolastatin, a combretastatin, a duocarmycin, a pyrrolobenzodiazepine dimer, an indolino-benzodiazepine dimer, an enediyne, a geldanamycin, a maytansine, a puromycin, a taxane, a vinca alkaloid, a camptothecin, a tubulysin, a hemiasterlin, a spliceostatin, a pladienolide, and stereoisomers, isosteres, analogs, or derivatives thereof.

[0209] The anthracyclines are derived from bacteria *Streptomyces* and have been used to treat a wide range of cancers, such as leukemias, lymphomas, breast, uterine, ovarian, and lung cancers. Exemplary anthracyclines include, but are not limited to, daunorubicin, doxorubicin (i.e., adriamycin), epirubicin, idarubicin, valrubicin, and mitoxantrone.


[0210] Dolastatins and their peptidic analogs and derivatives, auristatins, are highly potent antimitotic agents that have been shown to have anticancer and antifungal activity. See, e.g., U.S. Pat. No. 5,663,149 and Pettit et al., *Antimicrob. Agents Chemother.* 42:2961-2965, 1998. Exemplary dolastatins and auristatins include, but are not limited to, dolastatin 10, auristatin E, auristatin EB (AEB), auristatin EFP (AEFP), MMAD (Monomethyl Auristatin D or monomethyl dolastatin 10), MMAF (Monomethyl Auristatin F or N-methylvaline-valine-dolaisoleuine-dolaproine-phenylalanine), MMAE (Monomethyl Auristatin E or N-methylvaline-valine-dolaisoleuine-dolaproine-norephedrine), 5-benzoylvaleric acid-AE ester (AEVB), and other novel auristatins (such as the ones described in U.S. Publication No. 2013/0129753). In some embodiments, the auristatin is 0101 (2-methylalanyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(1,3-thiazol-2-yl)ethyl]amino]propyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide) having the following structure:




[0211] In some embodiments, the auristatin is 3377 (N,2-dimethylalanyl-N-[(1S,2R)-4-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxyl-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-2-methoxy-1-[(1S)-1-methylpropyl]-4-oxobutyl]-N-methyl-L-valinamide) having the following structure:




[0212] In some embodiments, the auristatin is 0131-OMe (N,2-dimethylalanyl-N-[(3R,4S,5S)-3-methoxy-1-((2S)-2-[(1R,2R)-1-methoxy-3-[(2S)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino}-2-methyl-3-oxopropyl]pyrrolidin-1-yl)-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide) having the following structure:



[0213] In other embodiments, the auristatin is 0131 (2-methyl-L-prolyl-N-[(3R,4S,5S)-1-((2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl)-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide) having the following structure:



[0214] In other embodiments, the auristatin is 0121 (2-methyl-L-prolyl-N-[(3R,4S,5S)-1-((2S)-2-[(1R,2R)-3-[(2S)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino}-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl)-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide) having the following structure:



[0215] Camptothecin is a cytotoxic quinoline alkaloid which inhibits the enzyme topoisomerase I. Examples of camptothecin and its derivatives include, but are not limited to, topotecan and irinotecan, and their metabolites, such as SN-38.


[0216] Combretastatins are natural phenols with vascular disruption properties in tumors. Exemplary combretastatins and their derivatives include, but are not limited to, combretastatin A-4 (CA-4) and ombrabulin.

[0217] Duocarmycin and CC-1065 are DNA alkylating agents with cytotoxic potency. See Boger and Johnson, PNAS 92:3642-3649 (1995). Exemplary duocarmycin and CC-1065 include, but are not limited to, (+)-duocarmycin A and (+)-duocarmycin SA, (+)-CC-1065, and the compounds as disclosed in the international application PCT/IB2015/050280 including, but not limited to, N~2~-acetyl-L-lysyl-L-valyl-N~5~-carbamoyl-N-[4-(((2-(((1S)-1-(chloromethyl)-3-[(5-(((1S)-1-(chloromethyl)-5-(phosphonoxy)-1,2-dihydro-3H-benzo[e]indol-3-yl)carbonyl]thiophen-2-yl)carbonyl]-2,3-dihydro-1H-benzo[e]indol-5-yl)oxy)carbonyl](methyl)amino)ethyl)(methyl)carbamoyl]oxy)methyl]phenyl]-L-ornithinamide having the structure:

5

10

15



20


N~2~-acetyl-L-lysyl-L-valyl-N~5~-carbamoyl-N-[4-((2-(((8S)-8-(chloromethyl)-6-((3-((1S)-1-(chloromethyl)-8-methyl-5-(phosphonooxy)-1,6-dihydropyrrolo[3,2-e]indol-3(2H)-yl)carbonyl}bicyclo[1.1.1]pent-1-yl)carbonyl]-1-methyl-3,6,7,8-tetrahydropyrrolo[3,2-e]indol-4-yl)oxy)carbonyl](methyl)amino)ethyl)(methyl)carbamoyl]oxy)methyl)phenyl]-L-ornithinamide having the structure:

25

30

35

40



45

N~2~-acetyl-L-lysyl-L-valyl-N~5~-carbamoyl-N-[4-((2-(((8S)-8-(chloromethyl)-6-((4-((1S)-1-(chloromethyl)-8-methyl-5-(phosphonooxy)-1,6-dihydropyrrolo[3,2-e]indol-3(2H)-yl)carbonyl}pentacyclo[4.2.0.0~2,5~.0-3,8~.0~4,7~]oct-1-yl)carbonyl]-1-methyl-3,6,7,8-tetrahydropyrrolo[3,2-e]indol-4-yl)oxy)carbonyl](methyl)amino)ethyl)(methyl)carbamoyl]oxy)methyl)phenyl]-L-ornithinamide having the structure:

50


55

5

10

15

20



[0218] Eneidyne are a class of anti-tumor bacterial products characterized by either nine- and ten-membered rings or the presence of a cyclic system of conjugated tripledouble-triple bonds. Exemplary eneidyne include, but are not limited to, calicheamicin, esperamicin, unciamycin, dynemicin, and their derivatives.

[0219] Geldanamycins are benzoquinone ansamycin antibiotic that bind to Hsp90 (Heat Shock Protein 90) and have been used antitumor drugs. Exemplary geldanamycins include, but are not limited to, 17-AAG (17-N-Allylamino-17-Demethoxygeldanamycin) and 17-DMAG (17-Dimethylaminoethylamino-17-demethoxygeldanamycin).


[0220] Hemiasterlin and its analogues (e.g., HTI-286) bind to the tubulin, disrupt normal microtubule dynamics, and, at stoichiometric amounts, depolymerize microtubules.

[0221] Maytansines or their derivatives maytansinoids inhibit cell proliferation by inhibiting the microtubules formation during mitosis through inhibition of polymerization of tubulin. See Remillard et al., Science 189:1002-1005, 1975. Exemplary maytansines and maytansinoids include, but are not limited to, mertansine (DM1) and its derivatives as well as ansamitocin.

[0222] Pyrrolobenzodiazepine dimers (PBDs) and indolino-benzodiazepine dimers (IGNs) are anti-tumor agents that contain one or more imine functional groups, or their equivalents, that bind to duplex DNA. PBD and IGN molecules are based on the natural product athramycin, and interact with DNA in a sequence-selective manner, with a preference for purine-guanine-purine sequences. Exemplary PBDs and their analogs include, but are not limited to, SJG-136.

[0223] Spliceostatins and pladienolides are anti-tumor compounds which inhibit splicing and interacts with spliceosome, SF3b. Examples of spliceostatins include, but are not limited to, spliceostatin A, FR901464, and (2S,3Z)-5-[[[(2R,3R,5S,6S)-6-[(2E,4E)-5-[(3R,4R,5R,7S)-7-(2-hydrazinyl-2-oxoethyl)-4-hydroxy-1,6-dioxaspiro[2.5]oct-5-yl]-3-methylpenta-2,4-dien-1-yl]-2,5-dimethyltetrahydro-2H-pyran-3-yl]amino]-5-oxopent-3-en-2-yl] acetate having the structure of

45



50

Examples of pladienolides include, but are not limited to, Pladienolide B, Pladienolide D, or E7107.

[0224] Taxanes are diterpenes that act as anti-tubulin agents or mitotic inhibitors. Exemplary taxanes include, but are not limited to, paclitaxel (e.g., TAXOL®) and docetaxel (TAXOTERE®).

[0225] Tubulysins are natural products isolated from a strain of myxobacteria that has been shown to depolymerize microtubules and induce mitotic arrest. Exemplary tubulysins include, but are not limited to, tubulysin A, tubulysin B, and tubulysin D.

[0226] Vinca alkylolids are also anti-tubulin agents. Exemplary vinca alkylolids include, but are not limited to, vincristine, vinblastine, vindesine, and vinorelbine.

[0227] Accordingly, in some embodiments, the cytotoxic agent is selected from the group consisting of MMAD (Monomethyl Auristatin D), 0101 (2-methylalanyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-2-methyl-3-oxo-3-[(1S)-2-phenyl-1-(1,3-thiazol-2-yl)ethyl]aminopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide), 3377 (N,2-dimethylalanyl-N-[(1S,2R)-4-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-2-methoxy-1-[(1S)-1-methylpropyl]-4-oxobutyl]-N-methyl-L-valinamide), 0131 (2-methyl-L-prolyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(1S)-1-carboxy-2-phenylethyl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide), 0131-OMe (N,2-dimethylalanyl-N-[(3R,4S,5S)-3-methoxy-1-[(2S)-2-[(1R,2R)-1-methoxy-3-[(2S)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino]-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide), 0121 (2-methyl-L-prolyl-N-[(3R,4S,5S)-1-[(2S)-2-[(1R,2R)-3-[(2S)-1-methoxy-1-oxo-3-phenylpropan-2-yl]amino]-1-methoxy-2-methyl-3-oxopropyl]pyrrolidin-1-yl]-3-methoxy-5-methyl-1-oxoheptan-4-yl]-N-methyl-L-valinamide), and (2S,3Z)-5-[(2R,3R,5S,6S)-6-[(2E,4E)-5-[(3R,4R,5R,7S)-7-(2-hydrazinyl-2-oxoethyl)-4-hydroxy-1,6-dioxaspiro[2.5]oct-5-yl]-3-methylpenta-2,4-dien-1-yl]-2,5-dimethyltetrahydro-2H-pyran-3-yl]amino]-5-oxopent-3-en-2-yl acetate.

[0228] In some embodiments, the agent is an immunomodulating agent. Examples of an immunomodulating agent include, but are not limited to, gancyclovir, etanercept, tacrolimus, sirolimus, voclosporin, cyclosporine, rapamycin, cyclophosphamide, azathioprine, mycophenolate mofetil, methotrexate, glucocorticoid and its analogs, cytokines, stem cell growth factors, lymphotoxins, tumor necrosis factor (TNF), hematopoietic factors, interleukins (e.g., interleukin-1 (IL-1), IL-2, IL-3, IL-6, IL-10, IL-12, IL-18, and IL-21), colony stimulating factors (e.g., granulocyte-colony stimulating factor (G-CSF) and granulocyte macrophage-colony stimulating factor (GM-CSF)), interferons (e.g., interferons- α , β and γ), the stem cell growth factor designated "S 1 factor," erythropoietin and thrombopoietin, or a combination thereof.

[0229] In some embodiments, the agent moiety is an imaging agent (e.g., a fluorophore or a chelator), such as fluorescein, rhodamine, lanthanide phosphors, and their derivatives thereof, or a radioisotope bound to a chelator. Examples of fluorophores include, but are not limited to, fluorescein isothiocyanate (FITC) (e.g., 5-FITC), fluorescein amidite (FAM) (e.g., 5-FAM), eosin, carboxyfluorescein, erythrosine, Alexa Fluor® (e.g., Alexa 350, 405, 430, 488, 500, 514, 532, 546, 555, 568, 594, 610, 633, 647, 660, 680, 700, or 750), carboxytetramethylrhodamine (TAMRA) (e.g., 5-TAMRA), tetramethylrhodamine (TMR), and sulforhodamine (SR) (e.g., SR101). Examples of chelators include, but are not limited to, 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA), 1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA), 1,4,7-triazacyclononane-1-glutaric acid-4,7-acetic acid (deferoxamine), diethylenetriaminepentaacetic acid (DTPA), and 1,2-bis(o-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA).

[0230] In some embodiments, therapeutic or diagnostic radioisotopes or other labels (e.g., PET or SPECT labels) can be incorporated in the agent for conjugation to the EGFRvIII antibodies as described herein. Examples of a radioisotope or other labels include, but are not limited to, ^3H , ^{11}C , ^{13}N , ^{14}C , ^{15}N , ^{15}O , ^{35}S , ^{18}F , ^{32}P , ^{33}P , ^{47}Sc , ^{51}Cr , ^{57}Co , ^{58}Co , ^{59}Fe , ^{62}Cu , ^{64}Cu , ^{67}Cu , ^{67}Ga , ^{68}Ga , ^{75}Se , ^{76}Br , ^{77}Br , ^{86}Y , ^{89}Zr , ^{90}Y , ^{94}Tc , ^{95}Ru , ^{97}Ru , ^{99}Tc , ^{103}Ru , ^{105}Rh , ^{105}Ru , ^{107}Hg , ^{109}Pd , ^{111}Ag , ^{111}In , ^{113}In , ^{121}Te , ^{122}Te , ^{123}I , ^{124}I , ^{125}I , ^{125}Te , ^{126}I , ^{131}I , ^{131}In , ^{133}I , ^{142}Pr , ^{143}Pr , ^{153}Pb , ^{153}Sm , ^{161}Tb , ^{165}Tm , ^{166}Dy , ^{166}H , ^{167}Tm , ^{168}Tm , ^{169}Yb , ^{177}Lu , ^{186}Re , ^{188}Re , ^{189}Re , ^{197}Pt , ^{198}Au , ^{199}Au , ^{201}Tl , ^{203}Hg , ^{211}At , ^{212}Bi , ^{212}Pb , ^{213}Bi , ^{223}Ra , ^{224}Ac , or ^{225}Ac .

[0231] In some embodiments, the agent is a therapeutic protein including, but is not limited to, a toxin, a hormone, an enzyme, and a growth factor.

[0232] Examples of a toxin protein (or polypeptide) include, but are not limited to, diphtheria (e.g., diphtheria A chain), *Pseudomonas* exotoxin and endotoxin, ricin (e.g., ricin A chain), abrin (e.g., abrin A chain), modeccin (e.g., modeccin A chain), alpha-sarcin, Aleurites fordii proteins, dianthin proteins, ribonuclease (RNase), DNase I, *Staphylococcal* enterotoxin-A, pokeweed antiviral protein, gelonin, diphtherin toxin, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, mitogellin, restrictocin, phenomycin, enomycin, tricothecenes, inhibitor cystine knot (ICK) peptides (e.g., ceratotoxins), and conotoxin (e.g., KIIIA or SmIIIA).

[0233] In some embodiments, the agent is a biocompatible polymer. The EGFRvIII antibodies as described herein can be conjugated to the biocompatible polymer to increase serum half-life and bioactivity, and/or to extend *in vivo* half-lives. Examples of biocompatible polymers include water-soluble polymer, such as polyethylene glycol (PEG) or its derivatives thereof and zwitterion-containing biocompatible polymers (e.g., a phosphorylcholine containing polymer).

[0234] In some embodiments, the agent is an oligonucleotide, such as anti-sense oligonucleotides.

[0235] In another aspect, the invention provides a conjugate of the antibody as described herein, wherein the conjugate comprises the formula: antibody-(acyl donor glutamine-containing tag)-(linker)-(cytotoxic agent).

[0236] Examples of a linker containing one or more reactive amines include, but are not limited to, Ac-Lys-Gly (acetyl-lysine-glycine), aminocaproic acid, Ac-Lys- β -Ala (acetyl-lysine- β -alanine), amino-PEG2 (polyethylene glycol)-C2, amino-PEG3-C2, amino-PEG6-C2 (or amino PEG6-propionyl), Ac-Lys-Val-Cit-PABC (acetyl-lysine-valine-citrulline-p-aminobenzyloxycarbonyl), amino-PEG6-C2-Val-Cit-PABC, aminocaproyl-Val-Cit-PABC, [(3R,5R)-1-{3-[2-(2-aminoethoxy)ethoxy]propanoyl}]piperidine-3,5-diyl]bis-Val-Cit-PABC, [(3S,5S)-1-{3-[2-(2-aminoethoxy)ethoxy]propanoyl}]piperidine-3,5-diyl]bis-Val-Cit-PABC, putrescine, or Ac-Lys-putrescine.

Methods of Using the EGFRvIII Antibodies and the Antibody Conjugates Thereof

[0237] The antibodies or the antibody conjugates of the present invention are useful in various applications including, but are not limited to, therapeutic treatment methods and diagnostic treatment methods.

[0238] The antibodies (e.g., monospecific and bispecific) and the antibody conjugates obtained by the methods described above can be used as a medicament. In some embodiments, such a medicament can be used for treating cancer, including solid tumors and liquid tumors. In some embodiments, the cancer is EGFRvIII related cancer including, but not limited to, glioblastoma (e.g., glioblastoma multiform), anaplastic astrocytoma, giant cell glioblastoma, gliosarcoma, anaplastic oligodendroglioma, anaplastic ependymoma, anaplastic oligoastrocytoma, choroid plexus carcinoma, anaplastic ganglioglioma, pineoblastoma, pineocytoma, meningioma, medulloepithelioma, ependymblastoma, medulloblastoma, supraentorial primitive neuroectodermal tumor, atypical teratoid/rhabdoid tumor, mixed glioma, head and neck cancer, non-small cell lung cancer, breast cancer, ovarian cancer, prostate cancer, medulloblastoma, colorectal cancer, anal cancer, cervical cancer, renal cancer, skin cancer, pancreatic cancer, liver cancer, bladder cancer, gastric cancer, thyroid cancer, mesothelioma, uterine cancer, lymphoma, or leukemia.

[0239] Provided is a method of inhibiting tumor growth or progression in a subject who has malignant cells expressing EGFRvIII, comprising administering to the subject in need thereof an effective amount of a composition comprising the EGFRvIII antibodies or the EGFRvIII antibody conjugates as described herein. Provided is a method of inhibiting metastasis of cells expressing EGFRvIII in a subject, comprising administering to the subject in need thereof an effective amount of a composition comprising the EGFRvIII antibodies or the EGFRvIII antibody conjugates as described herein.

Provided is a method of inducing tumor regression in malignant cells in a subject, comprising administering to the subject in need thereof an effective amount of a composition comprising the EGFRvIII antibodies or the EGFRvIII antibody conjugates as described herein.

[0240] The antibody (e.g., monospecific or bispecific) or the antibody conjugate according to the invention can be used in the manufacture of a medicament for treatment of a cancer in a patient in need thereof.

[0241] In some embodiments, the treatment can be in combination with one or more therapies against cancer selected from the group of antibodies therapy, chemotherapy, cytokines therapy, targeted therapy, vaccine therapy, dendritic cell therapy, gene therapy, hormone therapy, surgical resection, laser light therapy and radiation therapy. For example, the antibody (monospecific or bispecific) or the antibody conjugate of the invention can be administered to a patient in conjunction with (e.g., before, simultaneously, or following) 1) standard of care, including radiation, surgical resection, chemotherapy (e.g., temozolomide, procarbazine, carmustine, lomustine, vincristine etc.), antibody therapy such as bevacizumab, anti-angiogenic therapy, and/or tumor treating fields; 2) vaccine, including EGFRvIII vaccine; 3) targeted therapy, such as kinase inhibitors (e.g. everolimus); and 4) immunotherapies, including but not limited to anti-PD-1, anti-PD-L1, anti-PD-L2, anti-41BB, anti-TIM3, anti-LAG3, anti-TIGIT, anti-OX40, anti-HVEM, anti-BTLA, anti-CD40, anti-CD47, anti-CSF1R, anti-CSF1, anti-MARCO, anti-IL8, anti-CXCR4, and anti-CTLA4 antibodies.

[0242] The administration of the antibodies (e.g., monospecific or bispecific) or the antibody conjugates according to the invention may be carried out in any convenient manner, including by aerosol inhalation, injection, ingestion, transfusion, implantation or transplantation. The compositions described herein may be administered to a patient subcutaneously, intradermally, intratumorally, intracranially, intranodally, intramedullary, intramuscularly, by intravenous or intralymphatic injection, or intraperitoneally. In one embodiment, the heteromultimeric antibody compositions of the invention are preferably administered by intravenous injection.

[0243] In some embodiments, the administration of the antibodies (e.g., monospecific or bispecific) or the antibody conjugates can comprise administration of, for example, about 0.01 to about 20 mg per kg body weight including all integer values of mg per kg within those ranges. In some embodiments, the administration of the antibodies or the antibody conjugates can comprise administration of about 0.1 to 10 mg per kg body weight including all integer values of mg per kg within those ranges. The heteromultimeric antibody can be administered in one or more doses. In some embodiments, said effective amount of the antibody or the antibody conjugate can be administered as a single dose. In some embodiments, said effective amount of antibodies can be administered as more than one dose over a period time. Timing of administration is within the judgment of managing physician and depends on the clinical condition of the patient. While individual needs vary, determination of optimal ranges of effective amounts of a given antibody (e.g., monospecific or bispecific) or antibody conjugate for a particular disease or conditions within the skill of the art. An effective amount means an amount which provides a therapeutic or prophylactic benefit. The dosage administered will be dependent upon the age, health and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment and the nature of the effect desired. In some embodiments, an effective amount of heteromultimeric antibody or composition comprising those antibodies are administered parenterally. In some embodiments, administration can be an intravenous administration. In some embodiments, administration can be directly done by injection within a tumor.

[0244] In some embodiments, anti-EGFRvIII antibodies provided herein may be used for diagnostic purposes, such in assays to identify EGFRvIII protein in samples (e.g. in immunohistochemistry assays) or in patients.

Compositions

[0245] In one aspect, the present invention provides a pharmaceutical composition comprising an antibody (e.g., monospecific or bispecific) or an antibody conjugate, of the invention or portion thereof as described above in a pharmaceutically acceptable carrier. In certain embodiments, the polypeptides of the invention may be present in a neutral form (including zwitter ionic forms) or as a positively or negatively-charged species. In some embodiments, the polypeptides may be complexed with a counterion to form a "pharmaceutically acceptable salt," which refers to a complex comprising one or more polypeptides and one or more counterions, where the counterions are derived from pharmaceutically acceptable inorganic and organic acids and bases.

[0246] The antibody (e.g., monospecific or bispecific) or the antibody conjugate, or portions thereof, may be administered alone or in combination with one or more other polypeptides of the invention or in combination with one or more other drugs (or as any combination thereof). The pharmaceutical compositions, methods and uses of the invention thus also encompass embodiments of combinations (co-administration) with other active agents, as detailed below.

[0247] As used herein, the terms "co-administration," "co-administered" and "in combination with," referring to the antibodies of the invention and one or more other therapeutic agents, is intended to mean, and does refer to and include the following: (i) simultaneous administration of such combination of a heterodimer disclosed herein and therapeutic agent(s) to a patient in need of treatment, when such components are formulated together into a single dosage form which releases said components at substantially the same time to said patient; (ii) substantially simultaneous administration of such combination of a heterodimer disclosed herein and therapeutic agent(s) to a patient in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at substantially the same time by said patient, whereupon said components are released at substantially the same time to said patient; (iii) sequential administration of such combination of a heterodimer disclosed herein and therapeutic agent(s) to a patient in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at consecutive times by said patient with a significant time interval between each administration, whereupon said components are released at substantially different times to said patient; and (iv) sequential administration of such combination of a heterodimer disclosed herein and therapeutic agent(s) to a patient in need of treatment, when such components are formulated together into a single dosage form which releases said components in a controlled manner whereupon they are concurrently, consecutively, and/or overlappingly released at the same and/or different times to said patient, where each part may be administered by either the same or a different route.

[0248] Generally, the antibody (e.g., monospecific or bispecific) or the antibody conjugate disclosed herein or portions thereof are suitable to be administered as a formulation in association with one or more pharmaceutically acceptable excipient(s). The term 'excipient' is used herein to describe any ingredient other than the compound(s) of the invention. The choice of excipient(s) will to a large extent depend on factors such as the particular mode of administration, the effect of the excipient on solubility and stability, and the nature of the dosage form. As used herein, "pharmaceutically acceptable excipient" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Some examples of pharmaceutically acceptable excipients are water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Additional examples of pharmaceutically acceptable substances are wetting agents or minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody.

[0249] Pharmaceutical compositions of the present invention and methods for their preparation will be readily apparent to those skilled in the art. Such compositions and methods for their preparation may be found, for example, in Remington's Pharmaceutical Sciences, 19th Edition (Mack Publishing Company, 1995). Pharmaceutical compositions are preferably manufactured under GMP conditions.

[0250] A pharmaceutical composition of the invention may be prepared, packaged, or sold in bulk, as a single unit dose, or as a plurality of single unit doses. As used herein, a "unit dose" is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage. Any method for administering peptides, proteins or antibodies accepted in the art may suitably be employed for the heterodimeric proteins and portions thereof disclosed herein.

[0251] The pharmaceutical compositions of the invention are typically suitable for parenteral administration. As used herein, "parenteral administration" of a pharmaceutical composition includes any route of administration characterized by physical breaching of a tissue of a subject and administration of the pharmaceutical composition through the breach in the tissue, thus generally resulting in the direct administration into the blood stream, into muscle, or into an internal organ. Parenteral administration thus includes, but is not limited to, administration of a pharmaceutical composition by injection of the composition, by application of the composition through a surgical incision, by application of the composition

through a tissue-penetrating non-surgical wound, and the like. In particular, parenteral administration is contemplated to include, but is not limited to, subcutaneous, intraperitoneal, intramuscular, intrasternal, intravenous, intraarterial, intrathecal, intraventricular, intraurethral, intracranial, intrasynovial injection or infusions; and kidney dialytic infusion techniques. Preferred embodiments include the intravenous and the subcutaneous routes.

[0252] Formulations of a pharmaceutical composition suitable for parenteral administration typically generally comprise the active ingredient combined with a pharmaceutically acceptable carrier, such as sterile water or sterile isotonic saline. Such formulations may be prepared, packaged, or sold in a form suitable for bolus administration or for continuous administration. Injectable formulations may be prepared, packaged, or sold in unit dosage form, such as in ampoules or in multi dose containers containing a preservative. Formulations for parenteral administration include, but are not limited to, suspensions, solutions, emulsions in oily or aqueous vehicles, pastes, and the like. Such formulations may further comprise one or more additional ingredients including, but not limited to, suspending, stabilizing, or dispersing agents. In one embodiment of a formulation for parenteral administration, the active ingredient is provided in dry (i.e. powder or granular) form for reconstitution with a suitable vehicle (e.g. sterile pyrogen free water) prior to parenteral administration of the reconstituted composition. Parenteral formulations also include aqueous solutions which may contain excipients such as salts, carbohydrates and buffering agents (preferably to a pH of from 3 to 9), but, for some applications, they may be more suitably formulated as a sterile non-aqueous solution or as a dried form to be used in conjunction with a suitable vehicle such as sterile, pyrogen-free water. Exemplary parenteral administration forms include solutions or suspensions in sterile aqueous solutions, for example, aqueous propylene glycol or dextrose solutions. Such dosage forms can be suitably buffered, if desired. Other parentally-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form, or in a liposomal preparation. Formulations for parenteral administration may be formulated to be immediate and/or modified release. Modified release formulations include controlled, delayed, sustained, pulsed, targeted and programmed release formulations. For example, in one aspect, sterile injectable solutions can be prepared by incorporating the heterodimeric protein, e.g., bispecific antibody, in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

[0253] An exemplary, non-limiting pharmaceutical composition of the invention is a formulation as a sterile aqueous solution having a pH that ranges from about 5.0 to about 6.5 and comprising from about 1 mg/mL to about 200 mg/mL of an antibody (e.g., monospecific or bispecific) or an antibody conjugate disclosed herein, from about 1 millimolar to about 100 millimolar of histidine buffer, from about 0.01 mg/mL to about 10 mg/mL of polysorbate 80, from about 100 millimolar to about 400 millimolar of trehalose, and from about 0.01 millimolar to about 1.0 millimolar of disodium EDTA dihydrate.

[0254] Dosage regimens may be adjusted to provide the optimum desired response. For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form, as used herein, refers to physically discrete units suited as unitary dosages for the patients/subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are generally dictated by and directly dependent on (a) the unique characteristics of the chemotherapeutic agent and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

[0255] Thus, the skilled artisan would appreciate, based upon the disclosure provided herein, that the dose and dosing regimen is adjusted in accordance with methods well-known in the therapeutic arts. That is, the maximum tolerable dose can be readily established, and the effective amount providing a detectable therapeutic benefit to a patient may also be determined, as can the temporal requirements for administering each agent to provide a detectable therapeutic benefit to the patient. Accordingly, while certain dose and administration regimens are exemplified herein, these examples in no way limit the dose and administration regimen that may be provided to a patient in practicing the present invention.

[0256] It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated, and may include single or multiple doses. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and

are not intended to limit the scope or practice of the claimed composition. Further, the dosage regimen with the compositions of this invention may be based on a variety of factors, including the type of disease, the age, weight, sex, medical condition of the patient, the severity of the condition, the route of administration, and the particular antibody employed. Thus, the dosage regimen can vary widely, but can be determined routinely using standard methods. For example,

doses may be adjusted based on pharmacokinetic or pharmacodynamic parameters, which may include clinical effects such as toxic effects and/or laboratory values. Thus, the present invention encompasses intra-patient dose-escalation as determined by the skilled artisan. Determining appropriate dosages and regimens are well-known in the relevant art and would be understood to be encompassed by the skilled artisan once provided the teachings disclosed herein.

[0257] For administration to human subjects, the total monthly dose of an antibody (e.g., monospecific or bispecific) or an antibody conjugate disclosed herein is typically in the range of about 0.5 to about 1200 mg per patient, depending, of course, on the mode of administration, mechanism of action, and target biology. For example, an intravenous monthly dose may require about 1 to about 1000 mg/patient. The total monthly dose may be administered in single or divided doses and may, at the physician's discretion, fall outside of the typical range given herein.

[0258] An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of an antibody (e.g., monospecific or bispecific) or an antibody conjugate, disclosed herein is about 1 to about 1000 mg/patient/month. In certain embodiments, the heterodimeric protein may be administered at about 1 to about 200 or about 1 to about 150 mg/patient/month.

Kits

[0259] The invention also provides kits for use in the instant methods. Kits of the invention include one or more containers comprising the antibody (e.g., monospecific or bispecific) or the antibody conjugate of the invention and instructions for use in accordance with any of the methods described herein. Generally, these instructions comprise a description of administration of the heterodimeric protein for the above described therapeutic treatments.

[0260] The instructions relating to the use of the antibody (e.g., monospecific or bispecific) or the antibody conjugate as described herein generally include information as to dosage, dosing schedule, and route of administration for the intended treatment. The containers may be unit doses, bulk packages (e.g., multi-dose packages) or subunit doses. Instructions supplied in the kits of the invention are typically written instructions on a label or package insert (e.g., a paper sheet included in the kit), but machine-readable instructions (e.g., instructions carried on a magnetic or optical storage disk) are also acceptable.

[0261] The kits of this invention are in suitable packaging. Suitable packaging includes, but is not limited to, vials, bottles, jars, flexible packaging (e.g., sealed Mylar or plastic bags), and the like. Also contemplated are packages for use in combination with a specific device, such as an inhaler, nasal administration device (e.g., an atomizer) or an infusion device such as a minipump. A kit may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The container may also have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is a bispecific antibody. The container may further comprise a second pharmaceutically active agent.

[0262] Kits may optionally provide additional components such as buffers and interpretive information. Normally, the kit comprises a container and a label or package insert(s) on or associated with the container.

[0263] The following examples are offered for illustrative purposes only. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description.

Examples

Reference Example 1: Affinity Determination for Recombinant Anti-EGFRvIII Murine-Human Chimeric Antibody and Humanized Antibodies

[0264] This example determines the affinity of chimeric and humanized anti-EGFRvIII antibodies at 25°C and 37°C.

[0265] Anti-EGFRvIII mouse (m) antibody, m62G7, generated from hybridomas was sequenced and subcloned into suitable vectors for expression as murine-human chimeric antibodies. The CDRs of mouse antibody m62G7 were grafted onto human framework and expressed as human IgG1 recombinant antibody, h62G7. Affinity variants of h62G7 were made by introducing mutations in the CDRs of the heavy and light chains. The affinities of recombinant anti-EGFRvIII chimeric antibody m62G7 and humanized h62G7 antibodies were measured on a surface plasmon resonance Biacore™ T200 biosensor equipped with a research-grade anti-human Fc coupled CM4 sensor chip (GE Healthcare Inc., Piscataway, NJ). Anti-EGFRvIII antibodies were then captured by anti-human Fc. Monomeric 8-histidine tagged human EGFRvIII extracellular domain was then injected as the analyte at 10-fold dilution series with top concentration at 1000

nM. Affinity of anti-EGFRvIII antibodies towards human EGFRvIII was measured at both 25°C and 37°C (Table 6). None of these antibodies showed detectable binding to 1000 nM 8-histidine tagged recombinant wild-type protein EGFRwt under the same assay condition.

[0266] In Table 6, variants of h62G7 are described with reference to the heavy chain variation then the light chain variation. For example, antibody clone "h62G7-EQ/L6" refers to the h62G7 clone containing the "EQ" variation in the heavy chain (also referred to herein as "h62G7-EQ") and the "L6" variation in the light chain (also referred to herein as "h62G7-L6"). These heavy chain and light chain amino acid sequences are provided in Table 2. Also, in the present application, a h62G7 variant may be referred to with either the heavy chain or the light chain variant written first — so, for example, "h62G7-EQ/L6" and "h62G7-L6/EQ" both refer to an antibody which contains a h62G7-EQ heavy chain and a h62G7-L6 light chain.

Table 6

Antibody	25°C			37°C		
	k_a (1/Ms)	k_d (1/s)	K_D (nM)	k_a (1/Ms)	k_d (1/s)	K_D (nM)
m62G7	7.30E+05	6.40E-02	88.7	8.00E+05	1.70E-01	207.0
h62G7-EQ/L6	2.40E+05	1.00E-02	43.8	6.60E+05	7.40E-02	112.8
h62G7-EQ/L1-DV	2.00E+05	1.20E-05	59.9	3.70E+05	6.90E-02	185.8
h62G7-H14/L1-DV	1.80E+04	2.00E-02	1087.9	6.60E+04	1.00E-01	1539.6
h62G7-H14/L6	1.30E+04	1.30E-02	992.2	4.30E+04	6.80E-02	1583.3

Example 2: Affinity Determination for Human Anti-EGFRvIII Antibodies

[0267] This example determines the affinity of various human anti-EGFRvIII antibodies at 37°C.

[0268] To generate human antibodies against EGFRvIII, transgenic Alivamab mice (Ablexis LLC, San Francisco, CA) were immunized with alternating schedule of rat glioblastoma cell line expressing EGFRvIII, F98-npEGFRvIII (American Type Culture Collection, Manassas, VA) and peptides (SEQ ID NO: 227: CGSGSGLEEKKGNYVVDH) directed to the junction region in EGFRvIII. Hybridomas were generated using standard techniques. To determine the binding affinity and specificity of these hybridomas to EGFRvIII, antibodies in culture supernatants were captured by anti-mouse Fc using Biacore™ T200 biosensor equipped with anti-mouse Fc coupled CM4 sensor chips (Biacore™ AB, Uppsala, Sweden - now GE Healthcare). Monomeric 8-histidine tagged human EGFRvIII extracellular domain was then injected as the analyte at 10-fold dilution series starting with top concentration 1000 nM. Affinity of anti-EGFRvIII antibodies towards human EGFRvIII was measured at 37°C (Table 7). None of these hybridoma antibodies showed detectable binding to 1000 nM 8-histidine tagged recombinant wild-type protein EGFRwt under the same assay condition.

Table 7

Antibody	EGFRvIII binding at 37°C		
	k_a (1/Ms)	k_d (1/s)	K_D (nM)
42G9	6.88E+04	5.63E-04	8.2
32A10	6.54E+04	6.26E-04	9.6
21E11	6.66E+04	6.32E-04	9.5
49B11	7.64E+04	6.95E-04	9.1
46E10	5.97E+04	7.16E-04	12.0
12H6	5.93E+04	7.33E-04	12.4
19A9	5.58E+04	1.04E-03	18.6
11B11	5.21 E+04	1.13E-03	21.7
21E7	6.52E+04	1.30E-03	19.9
20B9	4.67E+04	1.50E-03	32.1
12B2	7.38E+04	1.79E-03	24.3

(continued)

Antibody	EGFRvIII binding at 37°C		
	k_a (1/Ms)	k_d (1/s)	K_D (nM)
11F10	6.63E+04	2.81 E-03	42.4
17G11	5.61E+04	3.00E-03	53.5
29D5	1.02E+05	4.24E-03	41.6
14C11	7.55E+04	5.93E-03	78.5
20E12	3.99E+04	1.41E-02	353.4
20G5	1.25E+05	2.89E-02	231.2
26B9	1.31E+05	3.20E-02	244.3
30D8	1.61E+05	2.77E-02	172.0
32G8	6.82E+03	1.22E-02	1788.9
34E7	3.77E+04	1.28E-02	339.5

Example 3: Binding Specificity of Anti-EGFRvIII Antibodies to EGFRvIII expressing Cell Lines by Flow Cytometry

[0269] This example demonstrates the cell binding specificity of anti-EGFRvIII antibodies to EGFRvIII expressing cells.

[0270] To assess the cell binding specificity of anti-EGFRvIII antibodies generated from the AlivaMab mice, three isogenic rat glioblastoma cell lines and a human cancer cell line were used: F98 (does not express any form of human EGFR), F98-EGFRwt (expresses wild-type EGFR), F98-npEGFRvIII (expresses EGFRvIII) and A431 (an epidermoid carcinoma cell line with wild-type EGFR over-expression), all obtained from American Type Culture Collection (Manassas, VA). For cell staining, 500,000 cells were incubated with 50 μ l hybridoma supernatants for 45 min at 4°C, washed with binding buffer (PBS (Phosphate Buffered Saline) + 0.5% BSA (Bovine Serum Albumin)), followed by incubation with FITC-conjugated goat anti-mouse Fc specific secondary antibody from Jackson ImmunoResearch Laboratories (West Grove, PA). Tables 8A and 8B show mean fluorescent intensities (MFI) of EGFRvIII antibodies (except clone 20G5) on EGFRvIII expressing cell line were at least 10-fold higher than on non-expressing cell lines. FIG. 1A, FIG. 1B, and FIG. 1C show examples of the FACS binding histograms of three EGFRvIII specific clones which had been cloned and expressed as recombinant human IgG1 antibodies, 42G9 (FIG. 1A), 32A10 (FIG. 1B) and 32G8 (FIG. 1C), to the three F98 cell lines.

Table 8A

Antibody	F98		F98-EGFRwt		F98-EGFRvIII		A431	
	MFI	% positive	MFI	% positive	MFI	% positive	MFI	% positive
2nd Ab only	170	0.6	202	1.7	258	2.3	592	0.4
anti-EGFR(wt and vIII)	163	0.5	9608	98.3	5329	99.4	55240	100.0
42G9	159	0.4	185	1.6	3247	98.5	538	0.3
32A10	159	0.5	185	1.4	3349	98.3	531	0.2
21E11	159	0.3	184	1.3	3105	98.5	555	0.5
49B11	156	0.6	185	1.3	2980	98.5	599	0.8
46E10	158	0.4	187	1.6	2986	98.7	560	0.5
12H6	157	0.5	188	1.9	3445	98.3	569	0.8
19A9	158	0.5	168	1.6	3100	98.1	578	1.0
11B11	161	0.6	187	1.7	3391	98.2	589	1.2
21E7	159	0.3	184	1.3	3105	98.5	603	1.1
20B9	157	0.3	189	1.8	3418	98.3	558	0.7

(continued)

	F98		F98-EGFRwt		F98-EGFRvIII		A431	
Antibody	MFI	% positive	MFI	% positive	MFI	% positive	MFI	% positive
12B2	156	0.4	185	1.5	2749	97.9	571	0.8
11F10	155	0.5	187	1.6	3283	98.0	582	1.1
17G11	157	0.6	184	1.5	3357	98.1	556	0.7
29D5	155	0.3	185	1.3	2829	97.9	531	0.4
14C11	157	0.4	185	1.3	3213	98.2	580	0.8

Table 8B

	F98		F98-EGFRwt		F98-EGFRvIII		A431	
Antibody	MFI	% positive	MFI	% positive	MFI	% positive	MFI	% positive
2nd Ab only	235	0.2	252	0.2	322	1.3	185	0.7
anti-EGFR(wt and vIII)	245	0.3	6857	97.2	5827	99.4	44493	100.0
20E12	381	6.0	348	3.4	3976	97.9	302	2.6
20G5	1248	16.8	1070	12.6	4639	98.5	391	2.0
26B9	310	4.1	298	2.3	5405	98.6	276	1.7
30D8	296	4.0	280	1.7	5165	98.6	269	1.3
32G8	329	4.9	301	1.6	3734	98.6	271	1.2
34E7	485	6.9	371	4.0	4128	98.5	294	1.1

Reference Example 4: Affinity Determination for Fully Human Anti-EGFRvIII Antibodies from phage library

[0271] This example determines the affinity of various human anti-EGFRvIII antibodies at 25°C.

[0272] Human anti-EGFRvIII antibodies obtained from phage library screen were sequenced and subcloned into suitable vectors for expression as recombinant human IgG1 antibodies. The affinities of antibodies were measured at 25°C (Table 9) on a surface plasmon resonance Biacore™ T200 biosensor equipped with an anti-human Fc coupled CM4 sensor chip (GE Healthcare Inc., Piscataway, NJ). Anti-EGFRvIII antibodies were captured by anti-human Fc. Monomeric 8-histidine tagged human EGFRvIII extracellular domain was then injected as the analyte at 10-fold dilution series starting at 1000 nM. Among the two antibodies, only C6 showed very weak but detectable binding to 1000 nM 8-histidine tagged recombinant wild-type protein EGFRwt at 25°C.

Table 9

Antibody	EGFRvIII binding at 25°C		
	k _a (1/Ms)	k _d (1/s)	K _D (nM)
B5	2.08E+04	1.41E-02	677.9
C6	1.68E+04	8.94E-03	532.1

Example 5: Generation and Characterization of GBM Cell Lines Expression EGFRvIII

[0273] This example demonstrates the expression of wild-type EGFR and EGFRvIII in GBM cell lines.

[0274] Five GFP (green fluorescent protein) and luciferase transduced human glioblastoma cell lines, DKMG, LN18, LN18-EGFRvIII, LN229 and LN229-EGFRvIII were used for functional characterization. DKMG, which expresses both endogenous wild-type EGFR and EGFRvIII, was obtained from DSMZ (Braunschweig, Germany). LN18 and LN229, which express only wild-type EGFR, were obtained from American Type Culture Collection (Manassas, VA). To generate GFP-luciferase labeled cell lines, DKMG, LN18 and LN229 were transduced with lentivirus particles (Amsbio, Cambridge,

MA) encoding both GFP (green fluorescent protein) and luciferase in a bicistronic system. LN18-EGFRvIII and LN229-EGFRvIII were then generated by transduction of the parental cell lines, with a lentivirus vector encoding the full length EGFRvIII gene (SEQ ID NO: 201). Wild-type EGFR and EGFRvIII expression in each cell line was then analysed using flow cytometer. For cell staining, 300,000 cells were incubated with 3 μ g EGFR wild-type specific or EGFRvIII specific antibody in 100 μ l binding buffer (PBS (Phosphate Buffered Saline) + 2% FBS) for 45 min at 4°C, washed with binding buffer, followed by incubation with Alexa Fluor 647-conjugated goat anti-human Fc specific secondary antibody from Jackson ImmunoResearch Laboratories (West Grove, PA). FIGs. 2A-C show the expression profiles of wild-type EGFR and EGFRvIII in LN229-EGFRvIII, LN18-EGFRvIII and DKMG, respectively.

Example 6: In vitro cytotoxicity assays with EGFRvIII-CD3 bispecific antibodies

[0275] This example demonstrates the cytotoxicity of EGFRvIII-CD3 bispecific antibodies towards EGFRvIII expressing GBM cell lines.

[0276] To generate EGFRvIII-CD3 bispecific antibodies, the heavy-chain variable domains of anti-EGFRvIII and anti-CD3 antibodies were subcloned into the appropriate human IgG2 based bispecific vectors and expressed with their corresponding light-chain in HEK293 cells. Purification of the EGFRvIII-CD3 bispecific antibodies was done according to published methods (J Mol Biol, 2012, 3, pp204-219; US patent publication 2013/0115208). In these assays, the EGFRvIII-CD3 bispecific antibodies contain the anti-EGFRvIII sequence of anti-EGFRvIII clones h62G7-EQ/L6, 30D8, or 42G9.

[0277] Target cells in this Example were: EGFRvIII transduced LN18-EGFRvIII cells (FIG. 3A) and parental LN18 (FIG. 3B) cells; EGFRvIII transduced LN229-EGFRvIII (FIG. 4A) and parental LN229 (FIG. 4B) cells; and DKMG cells (which express endogenous EGFRvIII and EGFR wild-type proteins) (FIG. 5).

[0278] For the cytotoxicity assays, luciferase transduced target cells were plated in white 96-well plates at 10,000 cells/well in PBMC media (RPMI, 10% FBS, 2 mM L-glutamine, 1% Pen/Strep, 20 μ M β -mercaptoethanol, 10 mM HEPES, 1% non-essential amino acids, 1 mM sodium pyruvate) and incubated at 37 °C. Twenty-four hours later, activated T cells at the desired T:E (target:effector) ratio (10,000 T cells for 1:1, for LN18 and LN229 cells; 50,000 T cells for 1:5, for DKMG cells) were added to target cells along with the EGFRvIII-CD3 bispecific antibodies, negative control human IgG, negative control CD3 monovalent antibody in bispecific Fc backbone, or negative control bivalent anti-EGFRvIII mAb 42G9 in wild-type human IgG. Cells were incubated for another 24 h at 37 °C. To detect the amount of viable target cells at the end of assay, the media was discarded and 100 μ l of 150 μ g/ml luciferin was added to each well. Luminescence signal was acquired on SpectraMax M5 Plate Reader (Molecular Devices, Sunnyvale, CA). Percentage of live target cells was determined by normalizing the luminescence reading for each sample to that of control well containing only target cells.

[0279] The results are summarized in FIGs. 3A, 3B, 4A, 4B, and 5. In the graphs, the EGFRvIII-CD3 bispecific antibody data are represented by open symbols, and the negative control antibody data are represented by solid symbols

[0280] Target cells that expressed EGFRvIII showed a dose-dependent response to treatment with EGFRvIII-CD3 bispecific antibodies h62G7-EQ/L6/CD3 biFc, 30D8/CD3 biFc, and 42G9/CD3 biFc. In contrast, target cells that expressed EGFR wild-type protein only were not killed, thus indicating the specificity of the EGFRvIII-CD3 bispecific antibodies for cells expressing EGFRvIII. In addition, target cells that expressed EGFRvIII did not show a response to treatment with negative control antibodies human IgG, CD3 monovalent biFc, or 42G1 hIgG1 (anti-EGFRvIII antibody).

[0281] For example, LN18-EGFRvIII target cells treated with 0.01 nM h62G7-EQ/L6/CD3 biFc were only about 20% viable at the end of the assay. In contrast, LN18-EGFRvIII target cells treated with 0.1 nM control IgG, CD3 mono biFc, or 42G9 hIgG1 were about 100% viable at the end of the assay (FIG. 3A). In addition, parental cell line LN18 target cells treated with 0.01 nM h62G7-EQ/L6/CD3 biFc were about 100% viable at the end of the assay (FIG 3B).

[0282] In another example, LN229-EGFRvIII target cells treated with 0.01 nM 42G9/CD3 biFc were only about 35% viable at the end of the assay. In contrast, LN229-EGFRvIII target cells treated with 0.1 nM control IgG, CD3 mono biFc, or 42G9 hIgG1 were about 90-100% viable at the end of the assay (FIG. 4A). In addition, parental cell line LN229 target cells treated with 0.01 nM 42G9/CD3 biFc were about 100% viable at the end of the assay (FIG 4B).

[0283] In another example, DKMG target cells treated with 1 nM h62G7-EQ/L6/CD3 biFc were only about 35% viable at the end of the assay (FIG. 5). In contrast, DKMG target cells treated with 1 nM control IgG, CD3 mono biFc, or 42G9 hIgG1 were about 100% viable at the end of the assay (FIG. 5).

[0284] These data demonstrate that EGFRvIII-CD3 bispecific antibodies effectively mediate killing by T cells of EGFRvIII expressing cells.

Example 7: In Vivo Study of Anti-EGFRvIII-CD3 Bispecific Antibodies in a GBM model LN229-EGFRvIII

[0285] This example determines the *in vivo* anti-tumor activity of anti-EGFRvIII bispecific antibodies in a subcutaneous LN229-EGFRvIII GBM cell line model.

[0286] Three million LN229-EGFRvIII cells were implanted subcutaneously into 5-6 weeks old NSG mice (Jackson Laboratory, Sacramento, CA). Tumor volume was measured once a week by a caliper device and calculated with the following formula: Tumor volume = (length × width²) /2. On day 18 post tumor implantation, animals were randomized by tumor sizes into five animals per group. A single dose of 20 million fresh pan T cells was administered intraperitoneally, followed by bolus tail vein injection of 0.5 mg/kg of EGFRvIII-CD3 bispecific antibodies (the antibodies contained the anti-EGFRvIII sequence of anti-EGFRvIII clones h62G7-EQ/L6, 30D8, or 42G9), or CD3 monovalent control in bispecific Fc backbone.

[0287] The results are summarized in FIG. 6. In the graph, the EGFRvIII-CD3 bispecific antibody data are represented by open symbols, and the negative control antibody data are represented by solid symbols.

[0288] The EGFRvIII-CD3 bispecific antibodies h62G7-EQ/L6/CD3 biFc, 30D8/CD3 biFc, and 42G9/CD3 biFc inhibited the *in vivo* growth of the EGFRvIII-expressing LN229-EGFRvIII GBM cells. In contrast, the negative control antibody CD3 monovalent biFc and a no treatment control (i.e. the mouse was not dosed with T cells or antibody) did not inhibit the *in vivo* growth of the LN229-EGFRvIII GBM cells. For example, at day 37 post tumor implantation, the mean tumor volume for mice treated with the EGFRvIII-CD3 bispecific antibodies 30D8/CD3 biFc and 42G9/CD3 biFc was less than 100 mm³, whereas the mean tumor volume for mice without treatment or treated with the CD3 monovalent biFc was greater than 1200 mm³.

[0289] These data demonstrate the *in vivo* anti-tumor activities of EGFRvIII-CD3 bispecific antibodies against EGFRvIII expressing tumor cells.

SEQUENCE LISTING

[0290]

<110> PFIZER INC.

Wong, Oi Kwan

Chou, Joyce Ching

<120> ANTIBODIES SPECIFIC FOR EPIDERMAL GROWTH FACTOR RECEPTOR VARIANT III AND THEIR USES

<130> PC72271Prov2

<160> 292

<170> PatentIn version 3.5

<210> 1

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 1

	Glu	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala
	1				5					10					15	
5	Ser	Val	Lys	Ile	Ser	Cys	Lys	Thr	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr
				20					25					30		
10	Thr	Leu	His	Trp	Val	Lys	Gln	Ser	His	Val	Lys	Ser	Leu	Glu	Trp	Ile
			35					40					45			
15	Gly	Gly	Ile	Asp	Pro	Ile	Asn	Gly	Gly	Thr	Thr	Tyr	Asn	Gln	Lys	Phe
		50					55					60				
20	Lys	Gly	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr
	65					70					75					80
25	Met	Glu	Leu	Arg	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
					85					90					95	
30	Ala	Arg	Gly	Glu	Ala	Met	Asp	Ser	Trp	Gly	Gln	Gly	Thr	Ser	Val	Thr
				100					105					110		
35	Val	Ser	Ser													
			115													
40	<210> 2															
	<211> 112															
	<212> PRT															
	<213> Artificial Sequence															
45	<220>															
	<223> Synthetic Construct															
50	<400> 2															
55																

	Asp	Val	Val	Met	Thr	Gln	Thr	Pro	Leu	Thr	Leu	Ser	Val	Thr	Ile	Gly
	1				5					10					15	
5	Gln	Pro	Ala	Ser	Ile	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser
				20					25					30		
10	Asn	Gly	Lys	Thr	Tyr	Leu	Asn	Trp	Leu	Leu	Gln	Arg	Pro	Gly	Gln	Ser
			35					40					45			
15	Pro	Lys	Arg	Leu	Ile	Tyr	Leu	Val	Ser	Lys	Leu	Asp	Ser	Gly	Val	Pro
		50					55					60				
20	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
	65					70					75				80	
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Phe	Tyr	Tyr	Cys	Val	Gln	Asp
					85					90					95	
30	Thr	His	Phe	Pro	Leu	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Glu	Leu	Lys
				100					105					110		
	<210> 3															
	<211> 115															
	<212> PRT															
	<213> Artificial Sequence															
	<220>															
	<223> Synthetic Construct															
35	<400> 3															
40	Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
	1				5					10					15	
45	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr
				20					25					30		
50	Thr	Leu	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
			35					40					45			
55	Gly	Gly	Ile	Asn	Pro	Ile	Asn	Gly	Gly	Thr	Thr	Tyr	Asn	Gln	Lys	Phe
		50					55					60				
60	Lys	Gly	Arg	Val	Thr	Met	Thr	Arg	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr
	65					70					75				80	
65	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys

	85	90	95	
5	Ala Arg Gly Glu Ala Met Asp Ser Trp Gly Gln Gly Thr Leu Val Thr	100	105	110
	Val Ser Ser			
10	115			
	<210> 4			
	<211> 112			
	<212> PRT			
	<213> Artificial Sequence			
15				
	<220>			
	<223> Synthetic Construct			
20				
	Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly	1	5	10
				15
25	Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser	20	25	30
30	Asn Gly Lys Thr Tyr Leu Asn Trp Phe Gln Gln Arg Pro Gly Gln Ser	35	40	45
	Pro Arg Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro	50	55	60
35				
	Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile	65	70	75
				80
40	Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Val Gln Asp	85	90	95
45	Thr His Phe Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys	100	105	110
	<210> 5			
	<211> 115			
	<212> PRT			
50	<213> Artificial Sequence			
	<220>			
	<223> Synthetic Construct			
55				
	<400> 5			
	Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala	1	5	10
				15

	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr	
				20					25					30			
5	Thr	Leu	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
			35					40					45				
10	Gly	Gly	Ile	Trp	Pro	Ile	Thr	Gly	Gly	Thr	Thr	Tyr	Asn	Gln	Lys	Phe	
		50					55					60					
15	Lys	Gly	Arg	Val	Thr	Met	Thr	Arg	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr	
	65					70					75					80	
20	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		
25	Ala	Arg	Gly	Glu	Ala	Gln	Gly	Ser	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	
				100					105					110			
30	Val	Ser	Ser														
			115														
	<210> 6																
	<211> 112																
	<212> PRT																
	<213> Artificial Sequence																
	<220>																
	<223> Synthetic Construct																
35	<400> 6																

	Asp	Val	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Thr	Leu	Gly	
	1				5					10					15		
5	Gln	Pro	Ala	Ser	Ile	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser	
				20					25					30			
10	Asn	Gly	Lys	Thr	Tyr	Leu	Asn	Trp	Phe	Gln	Gln	Arg	Pro	Gly	Gln	Ser	
			35					40					45				
15	Pro	Arg	Arg	Leu	Ile	Tyr	Gln	Val	Ser	Lys	Leu	Asp	Ser	Gly	Val	Pro	
		50					55					60					
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75				80		
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Gly	Gln	Asp	
					85					90					95		
30	Thr	His	Phe	Pro	Leu	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	
					100				105					110			

<210> 7
 <211> 115
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

 <400> 7

1 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 5 5 10 15
 5 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 10 Thr Leu His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Gly Ile Trp Pro Ile Thr Gly Gly Thr Thr Tyr Asn Gln Lys Phe
 50 55 60
 15 Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80
 20 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 25 Ala Arg Gly Glu Ala Glu Gly Ser Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ser
 115
 30 <210> 8
 <211> 112
 <212> PRT
 <213> Artificial Sequence
 35 <220>
 <223> Synthetic Construct
 40 <400> 8
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
 1 5 10 15
 45 Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 50
 55

	Asn	Asp	Lys	Thr	Tyr	Thr	Asn	Trp	Phe	Gln	Gln	Arg	Pro	Gly	Gln	Ser	
			35					40					45				
5	Pro	Arg	Arg	Leu	Ile	Tyr	Glu	Val	Ser	Lys	Leu	Asp	Val	Gly	Val	Pro	
			50				55					60					
10	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75					80	
15	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Gly	Gln	Asp	
					85					90					95		
20	Thr	His	Phe	Pro	Leu	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	
				100					105					110			
	<210> 9																
	<211> 121																
	<212> PRT																
	<213> Artificial Sequence																
25	<220>																
	<223> Synthetic Construct																
	<400> 9																
30	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Leu	Lys	Pro	Thr	Glu	
	1				5					10					15		
35	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Pro	
				20					25					30			
40	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu	
			35					40					45				
45	Trp	Phe	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Leu	Lys	Leu	Ser	
	50					55						60					
50	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val	
	65					70					75					80	
55	Val	Leu	Thr	Met	Thr	Asn	Met	Ala	Pro	Val	Asp	Ser	Ala	Thr	Tyr	Tyr	
					85					90					95		
	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Phe	Trp	Gly	
				100					105					110			
	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser								
				115				120									

<210> 10
 <211> 107
 <212> PRT
 <213> Artificial Sequence

5

<220>
 <223> Synthetic Construct

10

<400> 10

Glu Val Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
 1 5 10 15

15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Arg Ser Asn
 20 25 30

20

Leu Ala Trp Tyr Gln Gln Lys Ser Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Gly Ser Thr Ile Arg Ala Thr Gly Val Pro Ala Arg Phe Ser Gly
 50 55 60

25

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Ser
 65 70 75 80

30

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Ser Asp Trp Pro Phe
 85 90 95

35

Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 100 105

<210> 11
 <211> 121
 <212> PRT
 <213> Artificial Sequence

40

<220>
 <223> Synthetic Construct

45

<400> 11

50

55

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu		
	1				5					10					15			
5		Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Ala	
				20						25					30			
10	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu		
			35					40					45					
	Trp	Leu	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser		
		50					55					60						
15																		
	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val		
	65					70					75					80		
20																		
	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Phe		
					85					90					95			
25	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly		
				100					105					110				
	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser									
				115				120										
30																		
	<210>	12																
	<211>	107																
	<212>	PRT																
	<213>	Artificial Sequence																
35																		
	<220>																	
	<223>	Synthetic Construct																
40																		
	<400>	12																

	Glu	Val	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly		
	1				5					10					15			
5		Glu	Arg	Val	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ser	Ser	Asn	
				20					25					30				
10	Phe	Ala	Trp	Tyr	Gln	Gln	Arg	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Leu		
			35					40					45					
15	Tyr	Gly	Ala	Thr	Thr	Arg	Ala	Thr	Gly	Leu	Pro	Gly	Arg	Phe	Ser	Gly		
		50					55					60						
20	Ser	Gly	Ser	Gly	Thr	Glu	Asn	Ile	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser		
	65					70					75					80		
25	Glu	Asp	Phe	Ala	Ile	Tyr	Phe	Cys	Gln	Gln	Tyr	Lys	Asp	Trp	Pro	Phe		
					85					90					95			
30	Thr	Phe	Gly	Pro	Gly	Ser	Lys	Val	Asp	Ile	Lys							
				100					105									

<210> 13

<211> 121

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 13

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu
	1				5					10					15	
5	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Ala
				20					25					30		
10	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu
			35					40					45			
15	Trp	Leu	Gly	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Tyr	Ser	Thr	Ser
	50						55					60				
20	Leu	Arg	Gly	Arg	Ile	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Arg	Gly	Leu	Val
	65					70					75					80
25	Val	Leu	Thr	Leu	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr
						85				90					95	
30	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Phe	Trp	Gly
				100					105					110		
35	Pro	Gly	Phe	Leu	Val	Thr	Val	Ser	Ser							
			115					120								
40	<210> 14 <211> 107 <212> PRT <213> Artificial Sequence															
45	<220> <223> Synthetic Construct															
50	<400> 14															
55	Glu	Ile	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly
	1				5					10					15	
60	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Val	Ser	Gln	Ser	Ile	Gly	Ala	Asn
				20					25					30		
65	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Phe	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile
			35					40					45			
70	Tyr	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Ile	Pro	Val	Arg	Phe	Ser	Gly
	50						55					60				
75	Gly	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser
	65					70					75					80

	Glu	Asp	Phe	Ala	Ile	Tyr	Ser	Cys	Gln	Gln	Tyr	Ile	Tyr	Trp	Pro	Phe	
					85					90					95		
5	Thr	Phe	Gly	Pro	Gly	Thr	Thr	Val	Asp	Ile	Lys						
				100					105								
	<210> 15																
	<211> 121																
10	<212> PRT																
	<213> Artificial Sequence																
	<220>																
	<223> Synthetic Construct																
15	<400> 15																
	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu	
20	1				5					10					15		
	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Asn	Asn	Ala	
				20					25					30			
25	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu	
			35					40					45				
30	Trp	Phe	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Phe	Arg	Thr	Ser	
		50					55					60					
	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val	
35	65					70					75					80	
	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr	
					85					90					95		
40	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly	
				100					105					110			
45	Gln	Gly	Ile	Leu	Val	Thr	Val	Ser	Ser								
				115				120									
	<210> 16																
	<211> 107																
50	<212> PRT																
	<213> Artificial Sequence																
	<220>																
	<223> Synthetic Construct																
55	<400> 16																

	Glu	Ile	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly	
	1				5					10				15			
5	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ser	Asn	Asn	
				20					25					30			
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	
10			35					40					45				
	Tyr	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	
		50					55					60					
15	Ser	Asp	Ser	Gly	Thr	Glu	Phe	Ser	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser	
	65					70					75				80		
	Glu	Asp	Phe	Ala	Val	Tyr	Phe	Cys	Gln	Gln	Tyr	Lys	Asp	Trp	Pro	Phe	
20					85					90					95		
	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Glu	Ile	Lys						
25				100					105								
	<210> 17																
	<211> 122																
	<212> PRT																
	<213> Artificial Sequence																
30	<220>																
	<223> Synthetic Construct																
	<400> 17																
35																	
40																	
45																	
50																	
55																	

[illegible]

	Asp	Met	Val	Val	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly
	1				5					10					15	
5	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Gly	Ser	Asp
				20					25					30		
10	Leu	Ala	Trp	Tyr	Gln	Gln	Pro	Pro	Gly	Gln	Ser	Pro	Arg	Leu	Leu	Ile
			35					40					45			
15	Tyr	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly
		50					55					60				
20	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Thr	Ser	Leu	Glu	Ser
	65					70					75					80
25	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Asn	Asp	Trp	Pro	Phe
					85					90					95	
30	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile	Lys					
				100					105							
	<210> 19															
	<211> 121															
	<212> PRT															
	<213> Artificial Sequence															
	<220>															
	<223> Synthetic Construct															
35	<400> 19															
40	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu
	1				5					10					15	
45	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Val
				20					25					30		
50	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu
			35					40					45			

	Trp	Phe	Ala	His	Ile	Phe	Ser	Ser	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser	
	50						55					60					
5	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val	
	65					70					75				80		
10	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr	
					85					90					95		
15	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly	
				100					105					110			
20	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser								
			115					120									
	<210> 20																
	<211> 107																
	<212> PRT																
	<213> Artificial Sequence																
25	<220>																
	<223> Synthetic Construct																
	<400> 20																
30	Glu	Met	Glu	Val	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly	
	1				5					10					15		
35	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Asn	Ile	Gly	Ser	Asp	
				20					25					30			
40	Leu	Ala	Trp	Tyr	Gln	Gln	Gln	Ser	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	
			35					40					45				
45	Ser	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Val	Pro	Thr	Arg	Phe	Ser	Gly	
	50						55					60					
50	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Thr	Ser	Leu	Gln	Ser	
	65					70					75				80		
55	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Asn	Asp	Trp	Pro	Phe	
					85					90					95		
	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile	Lys						
				100					105								
	<210> 21																
	<211> 121																
	<212> PRT																
	<213> Artificial Sequence																

<220>

<223> Synthetic Construct

<400> 21

5

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
1 5 10 15

10

Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
20 25 30

15

Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

20

Trp Leu Ala His Ile Phe Ser Thr Asp Glu Lys Ser Ile Arg Arg Ser
50 55 60

Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
65 70 75 80

25

Val Leu Ile Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

30

Cys Ala Arg Asp Ser Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

35

<210> 22

<211> 107

<212> PRT

<213> Artificial Sequence

40

<220>

<223> Synthetic Construct

<400> 22

45

50

55

	Glu	Val	Val	Met	Thr	Gln	Ser	Pro	Pro	Asn	Leu	Ser	Val	Ser	Pro	Gly		
	1				5					10					15			
5		Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Thr	Ser	Asn	
				20					25					30				
10	Phe	Ala	Trp	Tyr	Gln	Gln	Arg	Pro	Gly	Gln	Ser	Pro	Arg	Leu	Leu	Leu		
			35					40					45					
15	Tyr	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Val	Pro	Gly	Arg	Phe	Ser	Gly		
		50					55					60						
20	Ser	Gly	Ser	Gly	Thr	Glu	Asn	Ile	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser		
	65					70					75					80		
25	Glu	Asp	Phe	Ala	Val	Tyr	Phe	Cys	Gln	Gln	Tyr	Lys	Asp	Trp	Pro	Phe		
					85					90					95			
30	Thr	Phe	Gly	Pro	Gly	Ser	Lys	Val	Asp	Ile	Lys							
				100					105									
35	<210> 23																	
	<211> 121																	
	<212> PRT																	
	<213> Artificial Sequence																	
40	<220>																	
	<223> Synthetic Construct																	
45	<400> 23																	
50																		
55																		

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu		
	1				5					10					15			
5		Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Ala	
				20						25					30			
10	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu		
			35					40					45					
15	Trp	Leu	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser		
		50					55					60						
20	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val		
	65					70					75					80		
25	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr		
					85					90					95			
30	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly		
				100					105					110				
35	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser									
			115					120										
40	<210> 24																	
	<211> 107																	
	<212> PRT																	
	<213> Artificial Sequence																	
45	<220>																	
	<223> Synthetic Construct																	
50	<400> 24																	

Glu Val Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
 1 5 10 15
 5 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Gly Val Ser Ser Asn
 20 25 30
 10 Phe Ala Trp Tyr Gln Gln Arg Pro Gly Gln Ser Pro Arg Leu Leu Leu
 35 40 45
 Tyr Gly Ala Ser Thr Arg Ala Thr Gly Val Pro Gly Arg Phe Ser Gly
 50 55 60
 15 Ser Gly Ser Gly Thr Glu Asn Ile Leu Thr Ile Ser Ser Leu Gln Ser
 65 70 75 80
 20 Glu Asp Phe Ala Ile Tyr Phe Cys Gln Gln Tyr Lys Asp Trp Pro Phe
 85 90 95
 Thr Phe Gly Pro Gly Ser Lys Val Asp Ile Lys
 100 105
 25
 <210> 25
 <211> 121
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 35 <400> 25
 Gln Val Thr Leu Glu Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 1 5 10 15
 40 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
 20 25 30
 45 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Pro Glu
 35 40 45
 Trp Phe Ala His Ile Phe Ser Thr Asp Glu Lys Ser Leu Arg Leu Ser
 50 55 60
 Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 65 70 75 80
 55 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 85 90 95

	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly	
				100					105					110			
5	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser								
			115					120									
	<210> 26																
	<211> 107																
10	<212> PRT																
	<213> Artificial Sequence																
	<220>																
	<223> Synthetic Construct																
15	<400> 26																
	Glu	Val	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly	
20	1				5					10					15		
	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Asn	Arg	Asn	
				20					25					30			
25																	
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	
			35					40					45				
30	Phe	Gly	Thr	Ser	Thr	Arg	Ala	Thr	Gly	Ile	Pro	Ala	Arg	Phe	Ser	Gly	
	50						55					60					
	Ser	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Asp	Ser	Leu	Gln	Ser	
35	65					70					75					80	
	Glu	His	Ser	Gly	Leu	Tyr	Tyr	Cys	Gln	Gln	Tyr	Asn	Asp	Trp	Pro	Phe	
					85					90					95		
40																	
	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile	Lys						
				100					105								
	<210> 27																
45	<211> 121																
	<212> PRT																
	<213> Artificial Sequence																
	<220>																
50	<223> Synthetic Construct																
	<400> 27																

55

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu	
	1				5					10					15		
5	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Ala	
				20					25					30			
	Lys	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu	
10			35					40					45				
	Trp	Leu	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser	
		50					55					60					
15	Leu	Arg	Ser	Arg	Leu	Thr	Met	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val	
	65					70					75					80	
	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr	
20					85					90					95		
	Cys	Val	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly	
25				100					105					110			
	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser								
			115					120									
30	<210> 28																
	<211> 107																
	<212> PRT																
	<213> Artificial Sequence																
35	<220>																
	<223> Synthetic Construct																
	<400> 28																
40																	
45																	
50																	
55																	

	Glu	Val	Leu	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly
	1				5					10					15	
5	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ser	Thr	Asn
				20					25					30		
10	Phe	Ala	Trp	Tyr	Gln	Gln	Arg	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Leu
			35					40					45			
15	Phe	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Ile	Pro	Gly	Arg	Phe	Ser	Gly
		50					55					60				
20	Ser	Gly	Ser	Gly	Thr	Glu	Asn	Ile	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser
	65					70					75					80
25	Glu	Asp	Phe	Ala	Ile	Tyr	Phe	Cys	Gln	Gln	Tyr	Lys	Asp	Trp	Pro	Phe
					85					90					95	
30	Thr	Phe	Gly	Pro	Gly	Ser	Lys	Val	Glu	Ile	Lys					
				100					105							

<210> 29

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 29

Asp Val Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
 1 5 10 15
 5
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Asn Ser Asn
 20 25 30
 10
 Leu Ala Trp Tyr Gln Gln Asn Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45
 Phe Gly Ser Ser Thr Arg Ala Thr Gly Ile Pro Ala Ser Phe Ser Gly
 50 55 60
 15
 Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Asn Ser Leu Gln Ser
 65 70 75 80
 20
 Glu His Ser Ala Val Tyr Tyr Cys Gln Gln Tyr Asn Asp Trp Pro Phe
 85 90 95
 25
 Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 100 105
 <210> 30
 <211> 121
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 35 <400> 30
 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 1 5 10 15
 40
 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Pro
 20 25 30
 45
 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 35 40 45
 50
 Trp Leu Gly His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Ser
 50 55 60
 55

	Leu	Arg	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val
	65					70					75				80	
5	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr
					85					90					95	
10	Cys	Val	Arg	Asp	Ser	Ser	Asn	Tyr	Gly	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly
				100					105					110		
15	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser							
			115					120								
	<210> 31															
	<211> 107															
	<212> PRT															
	<213> Artificial Sequence															
20	<220>															
	<223> Synthetic Construct															
25	<400> 31															
	Glu	Val	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly
	1				5					10					15	
30	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ile	Asn	Asn
				20					25					30		
35	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile
			35					40					45			
40	Tyr	Gly	Thr	Ser	Thr	Arg	Ala	Thr	Asp	Ile	Pro	Ala	Arg	Phe	Ser	Gly
	50						55					60				
45	Ser	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser
	65					70					75				80	
50	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Asp	Tyr	Asn	Asn	Trp	Pro	Phe
					85					90					95	
55	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile	Lys					
				100					105							
	<210> 32															
	<211> 121															
	<212> PRT															
	<213> Artificial Sequence															
	<220>															
	<223> Synthetic Construct															

<400> 32

5 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Ile Glu
 1 5 10 15

 Thr Leu Thr Leu Thr Cys Thr Val Cys Gly Phe Ser Leu Ser Asn Pro
 20 25 30
 10
 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 35 40 45

 Trp Leu Gly His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Phe
 50 55 60
 15

 Leu Arg Ser Arg Leu Ser Ile Ser Lys Asp Thr Ser Lys Ser Gln Val
 65 70 75 80
 20

 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 85 90 95
 25

 Cys Ala Arg Asp Ser Ser Asp Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
 100 105 110

 30 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 33

<211> 107

35 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

40

<400> 33

45

50

55

	Glu	Ile	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Ser	Pro	Gly	
	1				5					10				15			
5	Glu	Arg	Thr	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Gly	Ser	Asn	
				20					25					30			
10	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	
			35					40					45				
15	Tyr	Gly	Ala	Ser	Thr	Arg	Ala	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	
		50					55					60					
20	Ser	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Ser	
	65					70					75					80	
25	Glu	Asp	Phe	Ala	Val	Tyr	Ser	Cys	Gln	Glu	Tyr	Asn	Asn	Trp	Pro	Phe	
					85					90					95		
30	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys						
				100					105								
35	<210> 34																
	<211> 121																
	<212> PRT																
	<213> Artificial Sequence																
40	<220>																
	<223> Synthetic Construct																
45	<400> 34																

1 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Ile Glu
 5 Thr Leu Thr Leu Thr Cys Thr Val Cys Gly Phe Ser Leu Ser Asn Pro
 10 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 15 Trp Leu Gly His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Phe
 20 Leu Arg Ser Arg Leu Ser Ile Ser Lys Asp Thr Ser Lys Ser Gln Val
 25 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 30 Cys Ala Arg Asp Ser Ser Asp Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
 35 Gln Gly Thr Leu Val Thr Val Ser Ser
 40 <210> 35
 <211> 121
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Synthetic Construct
 50 <400> 35
 55 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu

	1		5		10		15									
5	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Pro
				20					25					30		
10	Arg	Met	Gly	Val	Ser	Trp	Leu	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu
			35					40					45			
15	Trp	Phe	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Tyr	Ser	Pro	Ser
		50					55					60				
20	Leu	Arg	Gly	Arg	Leu	Thr	Val	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val
	65					70					75				80	
25	Val	Leu	Thr	Leu	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr
					85					90					95	
30	Cys	Ala	Arg	Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly
				100					105					110		
35	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser							
			115					120								
40	<210> 36															
	<211> 107															
	<212> PRT															
	<213> Artificial Sequence															
45	<220>															
	<223> Synthetic Construct															
50	<400> 36															
55																

```

1      Lys Ile Val Met Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
      1          5          10          15
5      Glu Arg Ala Thr Leu Ser Cys Arg Ala Asn Gln Ile Val Ser Ser Asn
      20          25          30
10     Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Val
      35          40          45
15     Phe Gly Thr Ser Thr Arg Ala Thr Gly Ile Pro Ile Arg Phe Ser Gly
      50          55          60
20     Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Val Ser Ser Leu Gln Ser
      65          70          75          80
25     Glu Asp Phe Ala Val Tyr Val Cys Gln Gln Tyr Asn Asp Trp Pro Phe
      85          90          95
      Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
      100          105
30     <210> 37
      <211> 118
      <212> PRT
      <213> Artificial Sequence
      <220>
      <223> Synthetic Construct
      <400> 37

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 5 Ser Leu Arg Leu Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Asp Ala
 20 25 30
 10 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Gly Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Val Val
 50 55 60
 15 Pro Leu Asn Gly Arg Phe Ile Ile Ser Arg Asp Asp Ser Arg Asn Thr
 65 70 75 80
 20 Leu Tyr Leu Gln Leu Asn Asn Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95
 25 Tyr Cys Thr Thr Val Pro Gly Ser Tyr Gly Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser
 115
 30 <210> 38
 <211> 112
 <212> PRT
 <213> Artificial Sequence
 35 <220>
 <223> Synthetic Construct
 40 <400> 38
 Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 45 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Asn
 50
 55

	20	25	30
5	Lys Arg Asn Asn Tyr Leu Asp Trp Phe Leu Gln Lys Pro Gly Gln Ser 35	40	45
10	Pro Gln Leu Leu Ile Tyr Leu Ala Ser Asn Arg Ala Ser Gly Val Pro 50	55	60
15	Asp Arg Phe Ser Gly Gly Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile 65	70	75
20	Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala 85	90	95
25	Gln Gln Thr Pro Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys 100	105	110

<210> 39

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 39

	Glu	Val	Asn	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Glu	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Tyr	Ala	
				20					25					30			
10	Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
			35					40					45				
15	Gly	Arg	Ile	Lys	Ser	Ile	Ala	Asp	Gly	Gly	Ala	Thr	Asp	Tyr	Ala	Ala	
		50					55					60					
20	Pro	Val	Arg	Asn	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Arg	Asn	Thr	
	65					70					75				80		
25	Leu	Tyr	Leu	Glu	Met	His	Ser	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	
					85					90					95		
30	Tyr	Cys	Thr	Thr	Ile	Pro	Gly	Asn	Asp	Ala	Phe	Asp	Met	Trp	Gly	Gln	
				100					105					110			
35	Gly	Thr	Met	Val	Thr	Val	Ser	Ser									
			115					120									
40																	
	<210> 40																
	<211> 112																
	<212> PRT																
	<213> Artificial Sequence																
45																	
	<220>																
	<223> Synthetic Construct																
50																	
	<400> 40																
55																	

Asp Ile Val Leu Thr Gln Ser Pro Leu Ser Leu Ser Val Thr Pro Gly
 1 5 10 15
 5 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 10 Asn Gly Lys Asn Tyr Leu Asp Trp Phe Leu His Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
 50 55 60
 15 Asp Arg Phe Ser Gly Ser Gly Ser Gly Ile Asp Phe Ile Leu Lys Ile
 65 70 75 80
 20 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala
 85 90 95
 25 Gln Gln Thr Pro Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys
 100 105 110
 <210> 41
 <211> 119
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 35 <400> 41
 Glu Val Gln Leu Val Glu Ser Trp Gly Val Leu Val Lys Pro Gly Gly
 1 5 10 15
 40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ile Phe Asn Asn Ala
 20 25 30
 45 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 Gly Arg Ile Lys Ser Lys Ser Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50
 55

	50	55	60
5	Pro Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asp Thr 65 70 75 80		
10	Leu Tyr Leu Gln Met Asn Gly Leu Lys Thr Glu Asp Thr Ala Val Tyr 85 90 95		
15	Phe Cys Thr Thr Ala Pro Gly Gly Pro Phe Asp Tyr Trp Gly Gln Gly 100 105 110		
20	Thr Leu Val Thr Val Ser Ser 115		
	<210> 42		
	<211> 112		
	<212> PRT		
	<213> Artificial Sequence		
25	<220>		
	<223> Synthetic Construct		
	<400> 42		
30	Asp Ile Val Leu Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly 1 5 10 15		
35	Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Arg 20 25 30		
40	Asp Gly Phe Asn Tyr Leu Asp Trp Phe Leu Gln Lys Pro Gly Gln Ser 35 40 45		
45	Pro Gln Leu Leu Ile Tyr Leu Ala Ser Ser Arg Ala Ser Gly Val Pro 50 55 60		
50	Asp Arg Phe Ser Gly Ser Asp Ser Gly Thr Asp Phe Thr Leu Lys Ile 65 70 75 80		
	Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala 85 90 95		
55	Leu Gln Thr Pro Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys 100 105 110		
	<210> 43		
	<211> 120		
	<212> PRT		
	<213> Artificial Sequence		

<220>

<223> Synthetic Construct

<400> 43

5

Glu Val Asn Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

10

Ser Leu Arg Leu Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Tyr Ala
20 25 30

15

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

20

Gly Arg Ile Lys Ser Ile Thr Asp Gly Gly Val Ile Asp Tyr Ala Ala
50 55 60

Pro Val Arg Asn Arg Cys Thr Ile Ser Arg Asp Asp Ser Arg Asn Thr
65 70 75 80

25

Leu Tyr Leu Glu Met His Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
85 90 95

30

Tyr Cys Thr Thr Ile Pro Gly Asn Asp Asp Phe Asp Met Trp Gly Gln
100 105 110

Gly Arg Met Val Thr Val Ser Ser
115 120

35

<210> 44

<211> 120

<212> PRT

<213> Artificial Sequence

40

<220>

<223> Synthetic Construct

<400> 44

45

50

55

Glu Val Asn Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 5 Ser Leu Arg Leu Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Tyr Ala
 20 25 30
 10 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Gly Arg Ile Lys Ser Ile Asn Asp Gly Gly Ala Thr Asp Tyr Ala Ser
 50 55 60
 15 Pro Val Arg Asn Arg Phe Thr Ile Ser Arg Asp Asp Ser Arg Asn Met
 65 70 75 80
 Leu Tyr Leu Glu Met His Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95
 25 Tyr Cys Thr Thr Ile Pro Gly Asn Asp Ala Phe Asp Met Trp Gly Gln
 100 105 110
 30 Gly Thr Leu Val Thr Val Ser Ser
 115 120
 <210> 45
 <211> 112
 35 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 40 <400> 45
 45
 50
 55

	Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Ser	Val	Thr	Pro	Gly
	1				5					10					15	
5	Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Thr	Gln	Ser	Leu	Leu	Tyr	Ser
				20					25					30		
10	Asn	Gly	Lys	Asn	Tyr	Leu	Asp	Trp	Phe	Leu	His	Lys	Pro	Gly	Gln	Ser
			35					40					45			
15	Pro	Gln	Leu	Leu	Ile	Phe	Leu	Gly	Ser	Ile	Arg	Ala	Ser	Gly	Val	Pro
		50					55					60				
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Ile	Asp	Phe	Ile	Leu	Lys	Ile
	65					70					75				80	
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Met	Gln	Ala
					85					90					95	
30	Gln	Gln	Thr	Pro	Ile	Thr	Phe	Gly	Gln	Gly	Thr	Arg	Leu	Glu	Ile	Lys
				100					105					110		
35	<210> 46															
	<211> 119															
	<212> PRT															
	<213> Artificial Sequence															
40	<220>															
	<223> Synthetic Construct															
45	<400> 46															
50																
55																

	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Asp	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Thr	Asn	Ala	
				20					25					30			
10	Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
			35					40					45				
15	Gly	Arg	Ile	Lys	Ser	Lys	Ile	Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	
		50					55					60					
20	Pro	Val	Lys	Gly	Arg	Phe	Ile	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	
	65					70					75					80	
25	Leu	Ser	Leu	Gln	Met	Asn	Ser	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Met	Tyr	
					85					90					95		
30	Tyr	Cys	Thr	Thr	Ala	Pro	Gly	Gly	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	
				100					105					110			
35	Ser	Leu	Val	Thr	Val	Ser	Ser										
				115													
40	<210> 47																
	<211> 112																
	<212> PRT																
	<213> Artificial Sequence																
45	<220>																
	<223> Synthetic Construct																
50	<400> 47																

	Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Pro	Val	Thr	Pro	Gly
	1				5					10					15	
5	Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Leu	Leu	Tyr	Ser
				20					25					30		
10	Asp	Arg	Arg	Asn	Tyr	Leu	Asp	Trp	Phe	Leu	Gln	Lys	Pro	Gly	Gln	Ser
			35					40					45			
15	Pro	His	Leu	Leu	Ile	Tyr	Leu	Gly	Ser	Tyr	Arg	Ala	Ser	Gly	Val	Pro
		50					55					60				
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
	65					70					75					80
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Met	Gln	Ala
						85				90					95	
30	Leu	Gln	Ile	Pro	Ile	Thr	Phe	Gly	Gln	Gly	Thr	Arg	Leu	Glu	Ile	Lys
				100					105					110		

<210> 48
 <211> 126
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 <400> 48

	Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ser	
	1				5					10					15		
5	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Asp	Thr	Phe	Ser	Ser	Asn	
				20					25					30			
10	Ala	Ile	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
			35					40					45				
15	Gly	Val	Ile	Ile	Pro	Ile	Phe	Gly	Thr	Ala	Asp	Tyr	Ala	Gln	Lys	Phe	
		50					55					60					
20	Gln	Gly	Arg	Val	Thr	Ile	Thr	Ala	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	
	65					70					75					80	
25	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		
30	Ala	Arg	His	Thr	Tyr	His	Glu	Tyr	Ala	Gly	Gly	Tyr	Tyr	Gly	Gly	Ala	
				100					105					110			
35	Met	Asp	Pro	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
			115					120					125				
40	<210> 49																
	<211> 112																
	<212> PRT																
	<213> Artificial Sequence																
45	<220>																
	<223> Synthetic Construct																
50	<400> 49																
55	Glu	Leu	Gln	Ser	Val	Leu	Thr	Gln	Pro	Pro	Ser	Ala	Ser	Gly	Thr	Pro	
	1				5					10					15		

	Gly	Gln	Arg	Val	Thr	Ile	Ser	Cys	Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	
				20					25					30			
5	Ser	Asn	Tyr	Val	Tyr	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	
			35					40					45				
	Ile	Leu	Ile	Tyr	Arg	Asn	Asn	Gln	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	
10		50					55					60					
	Phe	Ser	Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Ser	Gly	
	65					70					75					80	
15																	
	Leu	Arg	Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Ala	Ala	Trp	Asp	Asp	
					85					90					95		
20	Asn	Leu	Ser	Gly	Trp	Val	Phe	Gly	Thr	Gly	Thr	Lys	Leu	Thr	Val	Leu	
				100					105						110		

<210> 50

<211> 123

25 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

30

<400> 50

35

40

45

50

55

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

5 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Ser Tyr
 20 25 30

10 Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

15 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

20 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr Pro Leu
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

25

<210> 52
 <211> 121
 <212> PRT
 30 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

35 <220>
 <221> misc_feature
 <222> (100)..(101)
 <223> Xaa can be any naturally occurring amino acid

40 <400> 52

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Leu Lys Pro Thr Glu
 1 5 10 15

45 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Pro
 20 25 30

50

55

Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 35 40 45

5 Trp Phe Ala His Ile Phe Ser Thr Asp Glu Lys Ser Leu Lys Leu Ser
 50 55 60

10 Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 65 70 75 80

Val Leu Thr Met Thr Asn Met Ala Pro Val Asp Ser Ala Thr Tyr Tyr
 85 90 95

15 Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Phe Trp Gly
 100 105 110

20 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> 53
 <211> 121
 25 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

30 <220>
 <221> misc_feature
 <222> (100)..(101)
 <223> Xaa can be any naturally occurring amino acid

35 <400> 53

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 1 5 10 15

40 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
 20 25 30

45 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 35 40 45

50 Trp Leu Ala His Ile Phe Ser Thr Asp Glu Lys Ser Ile Arg Arg Ser
 50 55 60

Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 65 70 75 80

55 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Phe

```

5          Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
          100                      105                      110

          Gln Gly Thr Leu Val Thr Val Ser Ser
          115                      120

10
<210> 54
<211> 121
<212> PRT
<213> Artificial Sequence

15
<220>
<223> Synthetic Construct

20
<220>
<221> misc_feature
<222> (100)..(101)
<223> Xaa can be any naturally occurring amino acid

25
          Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
          1          5          10          15

30
          Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
          20          25          30

35
          Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
          35          40          45

          Trp Leu Gly His Ile Phe Ser Thr Asp Glu Lys Ser Tyr Ser Thr Ser
          50          55          60

40
          Leu Arg Gly Arg Ile Thr Ile Ser Lys Asp Thr Ser Arg Gly Leu Val
          65          70          75          80

45
          Val Leu Thr Leu Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
          85          90          95

          Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Phe Trp Gly
          100                      105                      110

          Pro Gly Phe Leu Val Thr Val Ser Ser
          115                      120

55
<210> 55
<211> 121
<212> PRT

```

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (100)..(101)

<223> Xaa can be any naturally occurring amino acid

<400> 55

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Asn Asn Ala
20 25 30

Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Phe Ala His Ile Phe Ser Thr Asp Glu Lys Ser Phe Arg Thr Ser
50 55 60

Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
65 70 75 80

Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
100 105 110

Gln Gly Ile Leu Val Thr Val Ser Ser
115 120

<210> 56

<211> 122

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (100)..(101)

<223> Xaa can be any naturally occurring amino acid

<400> 56

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu	
	1				5					10					15		
5	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Val	
				20					25					30			
	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu	
10			35					40					45				
	Trp	Phe	Ala	His	Ile	Phe	Ser	Ser	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser	
		50					55					60					
15	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val	
	65					70					75					80	
	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr	
20					85					90					95		
	Cys	Ala	Arg	Xaa	Xaa	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Phe	Trp	Gly	
25				100					105					110			
	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Asn							
				115				120									
30	<210> 57																
	<211> 121																
	<212> PRT																
	<213> Artificial Sequence																
35	<220>																
	<223> Synthetic Construct																
	<220>																
	<221> misc_feature																
40	<222> (100)..(101)																
	<223> Xaa can be any naturally occurring amino acid																
	<400> 57																

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu	
	1				5					10					15		
5	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Val	
				20					25					30			
10	Arg	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu	
			35					40					45				
15	Trp	Phe	Ala	His	Ile	Phe	Ser	Ser	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser	
		50					55					60					
20	Leu	Arg	Ser	Arg	Leu	Thr	Leu	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val	
	65					70					75					80	
25	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr	
					85					90					95		
30	Cys	Ala	Arg	Xaa	Xaa	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly	
				100					105					110			
35	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser								
			115					120									
40	<210> 58																
	<211> 121																
	<212> PRT																
	<213> Artificial Sequence																
45	<220>																
	<223> Synthetic Construct																
50	<220>																
	<221> misc_feature																
	<222> (100)..(101)																
	<223> Xaa can be any naturally occurring amino acid																
55	<400> 58																

1 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 5 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
 10 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 15 Trp Leu Ala His Ile Phe Ser Thr Asp Glu Lys Ser Ile Arg Arg Ser
 20 Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 25 Val Leu Ile Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 30 Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
 35 Gln Gly Thr Leu Val Thr Val Ser Ser
 40 <210> 59
 <211> 121
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Synthetic Construct
 50 <220>
 <221> misc_feature
 <222> (100)..(101)
 <223> Xaa can be any naturally occurring amino acid
 55 <400> 59

1 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 5 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
 10 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 15 Trp Leu Ala His Ile Phe Ser Thr Asp Glu Lys Ser Ile Arg Arg Ser
 20 Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 25 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 30 Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
 35 Gln Gly Thr Leu Val Thr Val Ser Ser
 40 <210> 60
 <211> 121
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Synthetic Construct
 50 <220>
 <221> misc_feature
 <222> (100)..(101)
 <223> Xaa can be any naturally occurring amino acid
 55 <400> 60

1 Gln Val Thr Leu Glu Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 5 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
 10 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Pro Glu
 15 Trp Phe Ala His Ile Phe Ser Thr Asp Glu Lys Ser Leu Arg Leu Ser
 20 Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 25 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 30 Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
 35 Gln Gly Thr Leu Val Thr Val Ser Ser
 40 <210> 61
 <211> 121
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Synthetic Construct
 50 <220>
 <221> misc_feature
 <222> (100)..(101)
 <223> Xaa can be any naturally occurring amino acid
 55 <400> 61

	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Thr	Glu
	1				5					10					15	
5	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Ser	Asn	Ala
				20					25					30		
10	Lys	Met	Gly	Val	Ser	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu
			35					40					45			
15	Trp	Leu	Ala	His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser
		50					55					60				
20	Leu	Arg	Ser	Arg	Leu	Thr	Met	Ser	Lys	Asp	Thr	Ser	Lys	Ser	Gln	Val
	65					70					75					80
25	Val	Leu	Thr	Met	Thr	Asn	Met	Asp	Pro	Val	Asp	Thr	Ala	Thr	Tyr	Tyr
					85					90					95	
30	Cys	Val	Arg	Xaa	Xaa	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr	Trp	Gly
				100					105					110		
35	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser							
			115					120								
40	<210> 62 <211> 6 <212> PRT <213> Artificial Sequence															
45	<220> <223> Synthetic Construct															
50	<400> 62															
55								Thr	Asp	Tyr	Thr	Leu	His			
								1				5				
55	<210> 63 <211> 6 <212> PRT <213> Artificial Sequence															
	<220> <223> Synthetic Construct															
	<400> 63															
								Gly	Tyr	Thr	Phe	Thr	Asp			
								1				5				
	<210> 64 <211> 10															

<212> PRT
 <213> Artificial Sequence

 <220>
 5 <223> Synthetic Construct

 <400> 64

 10 Gly Tyr Thr Phe Thr Asp Tyr Thr Leu His
 1 5 10

 <210> 65
 <211> 17
 <212> PRT
 15 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

 20 <400> 65

 Gly Ile Asp Pro Ile Asn Gly Gly Thr Thr Tyr Asn Gln Lys Phe Lys
 1 5 10 15

 25 Gly

 <210> 66
 <211> 11
 30 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct
 35
 <400> 66

 Gly Ile Asp Pro Ile Asn Gly Gly Thr Thr Tyr
 1 5 10
 40
 <210> 67
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 45
 <220>
 <223> Synthetic Construct

 <400> 67
 50
 Gly Glu Ala Met Asp Ser
 1 5

 <210> 68
 55 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 68

5

Gly	Ile	Asn	Pro	Ile	Asn	Gly	Gly	Thr	Thr	Tyr	Asn	Gln	Lys	Phe	Lys
1				5					10					15	

10

Gly

<210> 69

<211> 11

<212> PRT

15

<213> Artificial Sequence

<220>

<223> Synthetic Construct

20

<400> 69

Gly	Ile	Asn	Pro	Ile	Asn	Gly	Gly	Thr	Thr	Tyr
1				5					10	

25

<210> 70

<211> 17

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Synthetic Construct

<400> 70

35

Gly	Ile	Trp	Pro	Ile	Thr	Gly	Gly	Thr	Thr	Tyr	Asn	Gln	Lys	Phe	Lys
1				5					10					15	

40

Gly

<210> 71

<211> 11

<212> PRT

<213> Artificial Sequence

45

<220>

<223> Synthetic Construct

<400> 71

50

Gly	Ile	Trp	Pro	Ile	Thr	Gly	Gly	Thr	Thr	Tyr
1				5					10	

55

<210> 72

<211> 11

<212> PRT

<213> Artificial Sequence

<220>
 <223> Synthetic Construct

 <400> 72
 5

 Gly Ile Trp Pro Ile Thr Gly Gly Thr Thr Tyr
 1 5 10

 <210> 73
 10 <211> 6
 <212> PRT
 <213> Artificial Sequence

 <220>
 15 <223> Synthetic Construct

 <400> 73

 Gly Glu Ala Gln Gly Ser
 20 1 5

 <210> 74
 <211> 8
 <212> PRT
 25 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

 30 <400> 74

 Ser Asn Pro Arg Met Gly Val Ser
 1 5

 35 <210> 75
 <211> 8
 <212> PRT
 <213> Artificial Sequence

 40 <220>
 <223> Synthetic Construct

 <400> 75

 45 Gly Phe Ser Leu Ser Asn Pro Arg
 1 5

 <210> 76
 <211> 12
 50 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct
 55
 <400> 76

		Gly	Phe	Ser	Leu	Ser	Asn	Pro	Arg	Met	Gly	Val	Ser
		1				5					10		

5 <210> 77
 <211> 16
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

 <400> 77

		His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Leu	Lys	Leu	Ser	Leu	Arg	Ser
		1				5					10					15	

15 <210> 78
 <211> 10
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> Synthetic Construct

25 <400> 78

		His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Leu
		1				5					10

30 <210> 79
 <211> 11
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Synthetic Construct

 <400> 79

		Asp	Ser	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Phe
		1				5					10	

40 <210> 80
 <211> 8
 <212> PRT
 <213> Artificial Sequence

45 <220>
 <223> Synthetic Construct

50 <400> 80

		Ser	Asn	Ala	Arg	Met	Gly	Val	Ser
		1				5			

55 <210> 81
 <211> 8
 <212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 81

 Gly Phe Ser Leu Ser Asn Ala Arg
 1 5
 10
 <210> 82
 <211> 12
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct

 <400> 82
 20
 Gly Phe Ser Leu Ser Asn Ala Arg Met Gly Val Ser
 1 5 10

 <210> 83
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 25
 <220>
 <223> Synthetic Construct
 30
 <400> 83

 His Ile Phe Ser Thr Asp Glu Lys Ser Ile Arg Arg Ser Leu Arg Ser
 1 5 10 15
 35
 <210> 84
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct

 <400> 84
 45
 His Ile Phe Ser Thr Asp Glu Lys Ser Ile
 1 5 10

 50
 <210> 85
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 55
 <220>
 <223> Synthetic Construct

 <400> 85

	Asp Ser Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr
	1 5 10
5	<210> 86
	<211> 16
	<212> PRT
	<213> Artificial Sequence
10	<220>
	<223> Synthetic Construct
	<400> 86
15	His Ile Phe Ser Thr Asp Glu Lys Ser Tyr Ser Thr Ser Leu Arg Gly
	1 5 10 15
20	<210> 87
	<211> 10
	<212> PRT
	<213> Artificial Sequence
	<220>
	<223> Synthetic Construct
25	<400> 87
30	His Ile Phe Ser Thr Asp Glu Lys Ser Tyr
	1 5 10
	<210> 88
	<211> 8
	<212> PRT
	<213> Artificial Sequence
35	<220>
	<223> Synthetic Construct
	<400> 88
40	Asn Asn Ala Arg Met Gly Val Ser
	1 5
45	<210> 89
	<211> 8
	<212> PRT
	<213> Artificial Sequence
	<220>
	<223> Synthetic Construct
50	<400> 89
55	Gly Phe Ser Leu Asn Asn Ala Arg
	1 5
	<210> 90
	<211> 12
	<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

5

<400> 90

Gly Phe Ser Leu Asn Asn Ala Arg Met Gly Val Ser
1 5 10

10

<210> 91

<211> 16

<212> PRT

<213> Artificial Sequence

15

<220>

<223> Synthetic Construct

<400> 91

20

His Ile Phe Ser Thr Asp Glu Lys Ser Phe Arg Thr Ser Leu Arg Ser
1 5 10 15

25

<210> 92

<211> 10

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Synthetic Construct

<400> 92

35

His Ile Phe Ser Thr Asp Glu Lys Ser Phe
1 5 10

40

<210> 93

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

45

<400> 93

Ser Asn Val Arg Met Gly Val Ser
1 5

50

<210> 94

<211> 8

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Synthetic Construct

<400> 94

		Gly	Phe	Ser	Leu	Ser	Asn	Val	Arg					
		1				5								

5 <210> 95
 <211> 12
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

 <400> 95

		Gly	Phe	Ser	Leu	Ser	Asn	Val	Arg	Met	Gly	Val	Ser
		1				5					10		

15 <210> 96
 <211> 16
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> Synthetic Construct

25 <400> 96

		His	Ile	Phe	Ser	Ser	Asp	Glu	Lys	Ser	Ile	Arg	Arg	Ser	Leu	Arg	Ser
		1				5					10					15	

30 <210> 97
 <211> 10
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Synthetic Construct

 <400> 97

		His	Ile	Phe	Ser	Ser	Asp	Glu	Lys	Ser	Ile
		1				5					10

40 <210> 98
 <211> 16
 <212> PRT
 <213> Artificial Sequence

45 <220>
 <223> Synthetic Construct

50 <400> 98

		His	Ile	Phe	Ser	Thr	Asp	Glu	Lys	Ser	Leu	Arg	Leu	Ser	Leu	Arg	Ser
		1				5					10					15	

55 <210> 99
 <211> 8
 <212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 99

 Ser Asn Ala Lys Met Gly Val Ser
 1 5
 10
 <210> 100
 <211> 8
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct

 <400> 100
 20
 Gly Phe Ser Leu Ser Asn Ala Lys
 1 5

 <210> 101
 25
 <211> 12
 <212> PRT
 <213> Artificial Sequence

 <220>
 30
 <223> Synthetic Construct

 <400> 101

 Gly Phe Ser Leu Ser Asn Ala Lys Met Gly Val Ser
 35 1 5 10

 <210> 102
 <211> 16
 <212> PRT
 40
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

 <400> 102
 45
 His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Ser Leu Arg Ser
 1 5 10 15

 <210> 103
 50
 <211> 10
 <212> PRT
 <213> Artificial Sequence

 <220>
 55
 <223> Synthetic Construct

 <400> 103

	His Ile Phe Ser Ser Asp Glu Lys Ser Tyr
	1 5 10
5	<210> 104
	<211> 11
	<212> PRT
	<213> Artificial Sequence
10	<220>
	<223> Synthetic Construct
	<400> 104
15	Asp Ser Ser Asn Tyr Gly Gly Tyr Phe Asp Tyr
	1 5 10
20	<210> 105
	<211> 16
	<212> PRT
	<213> Artificial Sequence
	<220>
	<223> Synthetic Construct
25	<400> 105
30	His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Phe Leu Arg Ser
	1 5 10 15
35	<210> 106
	<211> 16
	<212> PRT
	<213> Artificial Sequence
	<220>
	<223> Synthetic Construct
	<400> 106
40	His Ile Phe Ser Thr Asp Glu Lys Ser Tyr Ser Pro Ser Leu Arg Gly
	1 5 10 15
45	<210> 107
	<211> 11
	<212> PRT
	<213> Artificial Sequence
	<220>
	<223> Synthetic Construct
50	<400> 107
55	Asp Ser Ser Asp Tyr Glu Gly Tyr Phe Asp Tyr
	1 5 10
	<210> 108
	<211> 11
	<212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 108
 Asp Ser Ser Asn Tyr Glu Glu Tyr Phe Asp Tyr
 1 5 10
 10
 <210> 109
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 <400> 109
 20
 Ser Asp Ala Trp Met Ser
 1 5
 <210> 110
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 25
 <220>
 <223> Synthetic Construct
 30
 <400> 110
 Gly Phe Thr Phe Ser Asp
 1 5
 35
 <210> 111
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct
 45
 <400> 111
 Gly Phe Thr Phe Ser Asp Ala Trp Met Ser
 1 5 10
 50
 <210> 112
 <211> 19
 <212> PRT
 <213> Artificial Sequence
 55
 <220>
 <223> Synthetic Construct
 <400> 112

Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Val Val Pro
 1 5 10 15

5 Leu Asn Gly

<210> 113

<211> 13

<212> PRT

10 <213> Artificial Sequence

<220>

<223> Synthetic Construct

15 <400> 113

Arg Ile Lys Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr
 1 5 10

20 <210> 114

<211> 7

<212> PRT

<213> Artificial Sequence

25 <220>

<223> Synthetic Construct

<400> 114

30 Val Pro Gly Ser Tyr Gly Tyr
 1 5

<210> 115

<211> 6

35 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

40

<400> 115

Ser Tyr Ala Trp Met Ser
 1 5

45

<210> 116

<211> 6

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Synthetic Construct

<400> 116

55

Gly Phe Thr Phe Ser Tyr
 1 5

<210> 117
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Construct
 10
 <400> 117
 Gly Phe Thr Phe Ser Tyr Ala Trp Met Ser
 1 5 10
 <210> 118
 <211> 19
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 20
 <400> 118
 Arg Ile Lys Ser Ile Ala Asp Gly Gly Ala Thr Asp Tyr Ala Ala Pro
 1 5 10 15
 Val Arg Asn
 30
 <210> 119
 <211> 13
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <223> Synthetic Construct
 <400> 119
 40
 Arg Ile Lys Ser Ile Ala Asp Gly Gly Ala Thr Asp Tyr
 1 5 10
 <210> 120
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 45
 <220>
 <223> Synthetic Construct
 50
 <400> 120
 Ile Pro Gly Asn Asp Ala Phe Asp Met
 1 5
 55
 <210> 121
 <211> 6
 <212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 121
 Asn Asn Ala Trp Met Ser
 1 5
 10
 <210> 122
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 <400> 122
 20
 Gly Phe Ile Phe Asn Asn
 1 5
 <210> 123
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 25
 <220>
 <223> Synthetic Construct
 30
 <400> 123
 Gly Phe Ile Phe Asn Asn Ala Trp Met Ser
 1 5 10
 35
 <210> 124
 <211> 19
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct
 45
 <400> 124
 Arg Ile Lys Ser Lys Ser Asp Gly Gly Thr Thr Asp Tyr Ala Ala Pro
 1 5 10 15
 50
 Val Lys Asp
 <210> 125
 <211> 13
 <212> PRT
 <213> Artificial Sequence
 55
 <220>

<223> Synthetic Construct

<400> 125

5 Arg Ile Lys Ser Lys Ser Asp Gly Gly Thr Thr Asp Tyr
1 5 10

<210> 126

$\langle 211 \rangle$ 8

10 <212> PRT

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Synthetic Construct

15

<400> 126

Ala Pro Gly Gly Pro Phe Asp Tyr
1 5

<210> 127

<211> 19

<212> PRT

<213> Artificial Sequence

25

$\langle 220 \rangle$

<223> Synthetic Construct

<400> 127

30

Arg Ile Lys Ser Ile Thr Asp Gly Gly Val Ile Asp Tyr Ala Ala Pro
1 5 10 15

35 Val Arg Asn

<210> 128

<211> 13

<212> PRT

40 <213> Artificial Sequence

<220>

<223> Synthetic Construct

45 <400> 128

Arg Ile Lys Ser Ile Thr Asp Gly Gly Val Ile Asp Tyr
1 5 10

50 <210> 129

<211> 9

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Synthetic Construct

<400> 129

Ile Pro Gly Asn Asp Asp Phe Asp Met
1 5

5 <210> 130
<211> 19
<212> PRT
<213> Artificial Sequence

10 <220>
<223> Synthetic Construct

<400> 130

15 Arg Ile Lys Ser Ile Asn Asp Gly Gly Ala Thr Asp Tyr Ala Ser Pro

1 5 10 15

20 Val Arg Asn

<210> 131
<211> 13
<212> PRT
25 <213> Artificial Sequence

<220>
<223> Synthetic Construct

30 <400> 131

Arg Ile Lys Ser Ile Asn Asp Gly Gly Ala Thr Asp Tyr
1 5 10

35 <210> 132
<211> 6
<212> PRT
<213> Artificial Sequence

40 <220>
<223> Synthetic Construct

<400> 132

45 Thr Asn Ala Trp Met Ser
1 5

<210> 133
<211> 6
50 <212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

55 <400> 133

		Gly	Phe	Thr	Phe	Thr	Asn											
		1				5												

5 <210> 134
 <211> 10
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

 <400> 134

15		Gly	Phe	Thr	Phe	Thr	Asn	Ala	Trp	Met	Ser							
		1				5					10							

20 <210> 135
 <211> 19
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

25 <400> 135

	Arg	Ile	Lys	Ser	Lys	Ile	Asp	Gly	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	Pro		
	1				5					10					15			

30 Val Lys Gly

35 <210> 136
 <211> 13
 <212> PRT
 <213> Artificial Sequence

40 <220>
 <223> Synthetic Construct

 <400> 136

45		Arg	Ile	Lys	Ser	Lys	Ile	Asp	Gly	Gly	Thr	Thr	Asp	Tyr				
		1				5					10							

50 <210> 137
 <211> 6
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

55 <400> 137

	Ser	Ser	Asn	Ala	Ile	Ser												
	1				5													

<210> 138
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Consturct
 10
 <400> 138
 Gly Asp Thr Phe Ser Ser
 1 5
 15
 <210> 139
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <223> Synthetic Consturct
 <400> 139
 25
 Gly Asp Thr Phe Ser Ser Asn Ala Ile Ser
 1 5 10
 30
 <210> 140
 <211> 17
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic Consturct
 35
 <400> 140
 Val Ile Ile Pro Ile Phe Gly Thr Ala Asp Tyr Ala Gln Lys Phe Gln
 1 5 10 15
 40
 Gly
 45
 <210> 141
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic Consturct
 50
 <400> 141
 55
 Val Ile Ile Pro Ile Phe Gly Thr Ala Asp Tyr
 1 5 10
 <210> 142
 <211> 17
 <212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Consturct
 5
 <400> 142
 10 His Thr Tyr His Glu Tyr Ala Gly Gly Tyr Tyr Gly Gly Ala Met Asp
 1 5 10 15
 Pro
 15 <210> 143
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 20 <220>
 <223> Synthetic Consturct
 <400> 143
 25 Ser Asn Tyr Ala Met Ser
 1 5
 30 <210> 144
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 35 <400> 144
 Gly Phe Thr Phe Ser Asn
 1 5
 40 <210> 145
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Synthetic Construct
 <400> 145
 50 Gly Phe Thr Phe Ser Asn Tyr Ala Met Ser
 1 5 10
 55 <210> 146
 <211> 17
 <212> PRT
 <213> Artificial Sequence
 <220>

<223> Synthetic Construct

<400> 146

⁵

Asp	Ile	Ser	Gly	Gly	Gly	Gly	Arg	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys
1				5					10					15	

Gly

10

<210> 147

$\langle 211 \rangle$ 11

<212> PRT

<213> Artificial Sequence

15

<220>

<223> Synthetic Construct

<400> 147

20

Asp Ile Ser Gly Gly Gly Gly Arg Thr Tyr Tyr
1 5 10

<210> 148

25

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

30

<223> Synthetic Construct

<400> 148

Ala Gly Leu Leu Tyr Gly Gly Gly Val Tyr Pro Met Asp Ile
1 5 10

35

<210> 149

<211> 16

<212> PRT

40

<213> Artificial Sequence

<220>

<223> Synthetic Construct

45

<400> 149

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Asn Gly Lys Thr Tyr Leu Asn
1 5 10 15

50

<210> 150

$\langle 211 \rangle$ 7

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Synthetic Construct

<400> 150

		Leu Val Ser Lys Leu Asp Ser
		1 5
5	<210> 151	
	<211> 9	
	<212> PRT	
	<213> Artificial Sequence	
10	<220>	
	<223> Synthetic Construct	
	<400> 151	
15		Val Gln Asp Thr His Phe Pro Leu Thr
		1 5
20	<210> 152	
	<211> 7	
	<212> PRT	
	<213> Artificial Sequence	
	<220>	
	<223> Synthetic Construct	
25	<400> 152	
30		Gln Val Ser Lys Leu Asp Ser
		1 5
	<210> 153	
	<211> 9	
	<212> PRT	
	<213> Artificial Sequence	
35	<220>	
	<223> Synthetic Construct	
	<400> 153	
40		Gly Gln Asp Thr His Phe Pro Leu Thr
		1 5
45	<210> 154	
	<211> 16	
	<212> PRT	
	<213> Artificial Sequence	
	<220>	
	<223> Synthetic Construct	
50	<400> 154	
55		Lys Ser Ser Gln Ser Leu Leu Tyr Ser Asn Asp Lys Thr Tyr Thr Asn
		1 5 10 15
	<210> 155	
	<211> 7	
	<212> PRT	

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 155

 Glu Val Ser Lys Leu Asp Val
 1 5
 10
 <210> 156
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct

 <400> 156
 20
 Arg Ala Ser Gln Ser Val Arg Ser Asn Leu Ala
 1 5 10

 <210> 157
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 25
 <220>
 <223> Synthetic Construct
 30
 <400> 157

 Gly Ser Thr Ile Arg Ala Thr
 1 5
 35
 <210> 158
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct

 <400> 158
 45
 Gln Gln Tyr Ser Asp Trp Pro Phe Thr
 1 5
 50
 <210> 159
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 55
 <220>
 <223> Synthetic Construct

 <400> 159

		Arg	Ala	Ser	Gln	Ser	Val	Ser	Ser	Asn	Phe	Ala
		1				5					10	

5 <210> 160
 <211> 7
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

 <400> 160

		Gly	Ala	Thr	Thr	Arg	Ala	Thr
		1				5		

15 <210> 161
 <211> 9
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> Synthetic Construct

25 <400> 161

		Gln	Gln	Tyr	Lys	Asp	Trp	Pro	Phe	Thr
		1				5				

30 <210> 162
 <211> 11
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Synthetic Construct

 <400> 162

		Arg	Val	Ser	Gln	Ser	Ile	Gly	Ala	Asn	Leu	Ala
		1				5					10	

40 <210> 163
 <211> 7
 <212> PRT
 <213> Artificial Sequence

45 <220>
 <223> Synthetic Construct

50 <400> 163

		Gly	Ala	Ser	Thr	Arg	Ala	Thr
		1				5		

55 <210> 164
 <211> 9
 <212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 164

 Gln Gln Tyr Ile Tyr Trp Pro Phe Thr
 1 5
 10
 <210> 165
 <211> 11
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct

 <400> 165
 20
 Arg Ala Ser Gln Ser Val Ser Asn Asn Leu Ala
 1 5 10

 <210> 166
 25
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 <220>
 30
 <223> Synthetic Construct

 <400> 166

 Arg Ala Ser Gln Ser Val Gly Ser Asp Leu Ala
 35 1 5 10

 <210> 167
 <211> 9
 <212> PRT
 40
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

 <400> 167
 45

 Gln Gln Tyr Asn Asp Trp Pro Phe Thr
 1 5
 50
 <210> 168
 <211> 11
 <212> PRT
 <213> Artificial Sequence

 55
 <220>
 <223> Synthetic Construct

 <400> 168

		Arg	Ala	Ser	Gln	Asn	Ile	Gly	Ser	Asp	Leu	Ala
		1				5					10	
5		<210> 169 <211> 11 <212> PRT <213> Artificial Sequence										
10		<220> <223> Synthetic Construct										
		<400> 169										
15		Arg	Ala	Ser	Gln	Ser	Val	Thr	Ser	Asn	Phe	Ala
		1				5					10	
20		<210> 170 <211> 11 <212> PRT <213> Artificial Sequence										
		<220> <223> Synthetic Construct										
25		<400> 170										
30		Arg	Ala	Ser	Gln	Gly	Val	Ser	Ser	Asn	Phe	Ala
		1				5					10	
35		<210> 171 <211> 11 <212> PRT <213> Artificial Sequence										
		<220> <223> Synthetic Construct										
		<400> 171										
40		Arg	Ala	Ser	Gln	Ser	Val	Asn	Arg	Asn	Leu	Ala
		1				5					10	
45		<210> 172 <211> 7 <212> PRT <213> Artificial Sequence										
50		<220> <223> Synthetic Construct										
		<400> 172										
55		Gly	Thr	Ser	Thr	Arg	Ala	Thr				
		1				5						
		<210> 173 <211> 11 <212> PRT										

<213> Artificial Sequence

<220>

<223> Synthetic Construct

5

<400> 173

10

Arg Ala Ser Gln Ser Val Ser Thr Asn Phe Ala
1 5 10

<210> 174

<211> 11

<212> PRT

<213> Artificial Sequence

15

<220>

<223> Synthetic Construct

<400> 174

20

Arg Ala Ser Gln Ser Val Asn Ser Asn Leu Ala
1 5 10

<210> 175

25

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

30

<223> Synthetic Construct

<400> 175

35

Gly Ser Ser Thr Arg Ala Thr
1 5

<210> 176

<211> 11

<212> PRT

40

<213> Artificial Sequence

<220>

<223> Synthetic Construct

45

<400> 176

Arg Ala Ser Gln Ser Val Ile Asn Asn Leu Ala
1 5 10

50

<210> 177

<211> 9

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Synthetic Construct

<400> 177

		Gln Asp Tyr Asn Asn Trp Pro Phe Thr
		1 5
5	<210> 178	
	<211> 11	
	<212> PRT	
	<213> Artificial Sequence	
	<220>	
10	<223> Synthetic Construct	
	<400> 178	
15		Arg Ala Ser Gln Ser Val Gly Ser Asn Leu Ala
		1 5 10
	<210> 179	
	<211> 8	
	<212> PRT	
20	<213> Artificial Sequence	
	<220>	
	<223> Synthetic Construct	
25	<400> 179	
		Gly Ala Ser Thr Arg Ala Ser Gly
		1 5
30	<210> 180	
	<211> 9	
	<212> PRT	
	<213> Artificial Sequence	
35	<220>	
	<223> Synthetic Construct	
	<400> 180	
40		Gln Glu Tyr Asn Asn Trp Pro Phe Thr
		1 5
	<210> 181	
	<211> 11	
45	<212> PRT	
	<213> Artificial Sequence	
	<220>	
	<223> Synthetic Construct	
50	<400> 181	
		Arg Ala Asn Gln Ile Val Ser Ser Asn Leu Ala
55		1 5 10
	<210> 182	
	<211> 16	
	<212> PRT	

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 182
 Arg Ser Ser Gln Ser Leu Leu His Asn Lys Arg Asn Asn Tyr Leu Asp
 1 5 10 15
 <210> 183
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 <400> 183
 20
 Leu Ala Ser Asn Arg Ala Ser
 1 5
 <210> 184
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 25
 <220>
 <223> Synthetic Construct
 30
 <400> 184
 Met Gln Ala Gln Gln Thr Pro Ile Thr
 1 5
 <210> 185
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct
 <400> 185
 Arg Ser Ser Gln Ser Leu Leu Tyr Ser Asn Gly Lys Asn Tyr Leu Asp
 1 5 10 15
 50
 <210> 186
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 55
 <220>
 <223> Synthetic Construct
 <400> 186

Leu Gly Ser Asn Arg Ala Ser
 1 5

5 <210> 187
 <211> 16
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

<400> 187

15 Arg Ser Ser Gln Ser Leu Leu His Arg Asp Gly Phe Asn Tyr Leu Asp
 1 5 10 15

20 <210> 188
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

25 <400> 188

Leu Ala Ser Ser Arg Ala Ser
 1 5

30 <210> 189
 <211> 9
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Synthetic Construct

<400> 189

40 Met Gln Ala Leu Gln Thr Pro Ile Thr
 1 5

45 <210> 190
 <211> 16
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

50 <400> 190

55 Arg Ser Thr Gln Ser Leu Leu Tyr Ser Asn Gly Lys Asn Tyr Leu Asp
 1 5 10 15

<210> 191
 <211> 7
 <212> PRT

<213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 5
 <400> 191

 Leu Gly Ser Ile Arg Ala Ser
 1 5
 10
 <210> 192
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct

 <400> 192
 20
 Arg Ser Ser Gln Ser Leu Leu Tyr Ser Asp Arg Arg Asn Tyr Leu Asp
 1 5 10 15

 <210> 193
 25
 <211> 7
 <212> PRT
 <213> Artificial Sequence

 <220>
 30
 <223> Synthetic Construct

 <400> 193

 Leu Gly Ser Tyr Arg Ala Ser
 1 5
 35

 <210> 194
 <211> 9
 <212> PRT
 40
 <213> Artificial Sequence

 <220>
 <223> Synthetic Construct

 <400> 194
 45

 Met Gln Ala Leu Gln Ile Pro Ile Thr
 1 5
 50
 <210> 195
 <211> 13
 <212> PRT
 <213> Artificial Sequence
 55
 <220>
 <223> Synthetic Construct

 <400> 195

		Ser	Gly	Ser	Ser	Ser	Asn	Ile	Gly	Ser	Asn	Tyr	Val	Tyr
		1					5				10			

5 <210> 196
 <211> 7
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

 <400> 196

					Arg	Asn	Asn	Gln	Arg	Pro	Ser
					1				5		

15 <210> 197
 <211> 11
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> Synthetic Construct

25 <400> 197

		Ala	Ala	Trp	Asp	Asp	Asn	Leu	Ser	Gly	Trp	Val
		1					5				10	

30 <210> 198
 <211> 11
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Synthetic Construct

 <400> 198

					Arg	Ala	Ser	Gln	Ser	Ile	Ser	Ser	Tyr	Leu	Asn
					1					5				10	

40 <210> 199
 <211> 7
 <212> PRT
 <213> Artificial Sequence

45 <220>
 <223> Synthetic Construct

50 <400> 199

					Ala	Ala	Ser	Ser	Leu	Gln	Ser
					1				5		

55 <210> 200
 <211> 9
 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

5

<400> 200

Gln Gln Ser Tyr Ser Thr Pro Leu Thr
1 5

10

<210> 201

<211> 943

<212> PRT

<213> Homo sapiens

15

<400> 201

20

25

30

35

40

45

50

55

	Met	Arg	Pro	Ser	Gly	Thr	Ala	Gly	Ala	Ala	Leu	Leu	Ala	Leu	Leu	Ala	
	1				5					10					15		
5	Ala	Leu	Cys	Pro	Ala	Ser	Arg	Ala	Leu	Glu	Glu	Lys	Lys	Gly	Asn	Tyr	
				20					25					30			
	Val	Val	Thr	Asp	His	Gly	Ser	Cys	Val	Arg	Ala	Cys	Gly	Ala	Asp	Ser	
10			35					40					45				
	Tyr	Glu	Met	Glu	Glu	Asp	Gly	Val	Arg	Lys	Cys	Lys	Lys	Cys	Glu	Gly	
		50					55					60					
15	Pro	Cys	Arg	Lys	Val	Cys	Asn	Gly	Ile	Gly	Ile	Gly	Glu	Phe	Lys	Asp	
	65					70					75					80	
	Ser	Leu	Ser	Ile	Asn	Ala	Thr	Asn	Ile	Lys	His	Phe	Lys	Asn	Cys	Thr	
20					85					90					95		
	Ser	Ile	Ser	Gly	Asp	Leu	His	Ile	Leu	Pro	Val	Ala	Phe	Arg	Gly	Asp	
25				100					105					110			
	Ser	Phe	Thr	His	Thr	Pro	Pro	Leu	Asp	Pro	Gln	Glu	Leu	Asp	Ile	Leu	
			115					120					125				
30	Lys	Thr	Val	Lys	Glu	Ile	Thr	Gly	Phe	Leu	Leu	Ile	Gln	Ala	Trp	Pro	
		130					135					140					
	Glu	Asn	Arg	Thr	Asp	Leu	His	Ala	Phe	Glu	Asn	Leu	Glu	Ile	Ile	Arg	
35	145					150					155					160	
	Gly	Arg	Thr	Lys	Gln	His	Gly	Gln	Phe	Ser	Leu	Ala	Val	Val	Ser	Leu	
					165					170					175		
40	Asn	Ile	Thr	Ser	Leu	Gly	Leu	Arg	Ser	Leu	Lys	Glu	Ile	Ser	Asp	Gly	
				180					185					190			
	Asp	Val	Ile	Ile	Ser	Gly	Asn	Lys	Asn	Leu	Cys	Tyr	Ala	Asn	Thr	Ile	
45			195					200					205				
	Asn	Trp	Lys	Lys	Leu	Phe	Gly	Thr	Ser	Gly	Gln	Lys	Thr	Lys	Ile	Ile	
50		210					215					220					
	Ser	Asn	Arg	Gly	Glu	Asn	Ser	Cys	Lys	Ala	Thr	Gly	Gln	Val	Cys	His	
	225					230					235					240	
55	Ala	Leu	Cys	Ser	Pro	Glu	Gly	Cys	Trp	Gly	Pro	Glu	Pro	Arg	Asp	Cys	

	245	250	255
5	Val Ser Cys Arg Asn Val Ser Arg Gly Arg Glu Cys Val Asp Lys Cys 260 265 270		
10	Asn Leu Leu Glu Gly Glu Pro Arg Glu Phe Val Glu Asn Ser Glu Cys 275 280 285		
15	Ile Gln Cys His Pro Glu Cys Leu Pro Gln Ala Met Asn Ile Thr Cys 290 295 300		
20	Thr Gly Arg Gly Pro Asp Asn Cys Ile Gln Cys Ala His Tyr Ile Asp 305 310 315 320		
	Gly Pro His Cys Val Lys Thr Cys Pro Ala Gly Val Met Gly Glu Asn 325 330 335		
25	Asn Thr Leu Val Trp Lys Tyr Ala Asp Ala Gly His Val Cys His Leu 340 345 350		
30	Cys His Pro Asn Cys Thr Tyr Gly Cys Thr Gly Pro Gly Leu Glu Gly 355 360 365		
	Cys Pro Thr Asn Gly Pro Lys Ile Pro Ser Ile Ala Thr Gly Met Val 370 375 380		
35	Gly Ala Leu Leu Leu Leu Leu Val Val Ala Leu Gly Ile Gly Leu Phe 385 390 395 400		
40	Met Arg Arg Arg His Ile Val Arg Lys Arg Thr Leu Arg Arg Leu Leu 405 410 415		
	Gln Glu Arg Glu Leu Val Glu Pro Leu Thr Pro Ser Gly Glu Ala Pro 420 425 430		
45	Asn Gln Ala Leu Leu Arg Ile Leu Lys Glu Thr Glu Phe Lys Lys Ile 435 440 445		
50	Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys Gly Leu Trp 450 455 460		
	Ile Pro Glu Gly Glu Lys Val Lys Ile Pro Val Ala Ile Lys Glu Leu 465 470 475 480		
55	Arg Glu Ala Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu Asp Glu Ala 485 490 495		

	Tyr	Val	Met	Ala	Ser	Val	Asp	Asn	Pro	His	Val	Cys	Arg	Leu	Leu	Gly	
				500					505					510			
5	Ile	Cys	Leu	Thr	Ser	Thr	Val	Gln	Leu	Ile	Thr	Gln	Leu	Met	Pro	Phe	
			515					520					525				
10	Gly	Cys	Leu	Leu	Asp	Tyr	Val	Arg	Glu	His	Lys	Asp	Asn	Ile	Gly	Ser	
		530					535					540					
15	Gln	Tyr	Leu	Leu	Asn	Trp	Cys	Val	Gln	Ile	Ala	Lys	Gly	Met	Asn	Tyr	
	545					550					555					560	
20	Leu	Glu	Asp	Arg	Arg	Leu	Val	His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Val	
					565					570					575		
25	Leu	Val	Lys	Thr	Pro	Gln	His	Val	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Ala	
				580					585					590			
30	Lys	Leu	Leu	Gly	Ala	Glu	Glu	Lys	Glu	Tyr	His	Ala	Glu	Gly	Gly	Lys	
			595					600					605				
35	Val	Pro	Ile	Lys	Trp	Met	Ala	Leu	Glu	Ser	Ile	Leu	His	Arg	Ile	Tyr	
		610					615					620					
40	Thr	His	Gln	Ser	Asp	Val	Trp	Ser	Tyr	Gly	Val	Thr	Val	Trp	Glu	Leu	
	625					630					635					640	
45	Met	Thr	Phe	Gly	Ser	Lys	Pro	Tyr	Asp	Gly	Ile	Pro	Ala	Ser	Glu	Ile	
					645					650					655		
50	Ser	Ser	Ile	Leu	Glu	Lys	Gly	Glu	Arg	Leu	Pro	Gln	Pro	Pro	Ile	Cys	
				660					665					670			
55	Thr	Ile	Asp	Val	Tyr	Met	Ile	Met	Val	Lys	Cys	Trp	Met	Ile	Asp	Ala	
			675					680					685				
60	Asp	Ser	Arg	Pro	Lys	Phe	Arg	Glu	Leu	Ile	Ile	Glu	Phe	Ser	Lys	Met	
		690					695					700					
65	Ala	Arg	Asp	Pro	Gln	Arg	Tyr	Leu	Val	Ile	Gln	Gly	Asp	Glu	Arg	Met	
	705					710					715					720	
70	His	Leu	Pro	Ser	Pro	Thr	Asp	Ser	Asn	Phe	Tyr	Arg	Ala	Leu	Met	Asp	
					725					730					735		
75	Glu	Glu	Asp	Met	Asp	Asp	Val	Val	Asp	Ala	Asp	Glu	Tyr	Leu	Ile	Pro	
				740					745					750			

	Gln	Gln	Gly	Phe	Phe	Ser	Ser	Pro	Ser	Thr	Ser	Arg	Thr	Pro	Leu	Leu	
			755					760					765				
5	Ser	Ser	Leu	Ser	Ala	Thr	Ser	Asn	Asn	Ser	Thr	Val	Ala	Cys	Ile	Asp	
			770					775					780				
10	Arg	Asn	Gly	Leu	Gln	Ser	Cys	Pro	Ile	Lys	Glu	Asp	Ser	Phe	Leu	Gln	
			785				790					795				800	
15	Arg	Tyr	Ser	Ser	Asp	Pro	Thr	Gly	Ala	Leu	Thr	Glu	Asp	Ser	Ile	Asp	
					805					810					815		
20	Asp	Thr	Phe	Leu	Pro	Val	Pro	Glu	Tyr	Ile	Asn	Gln	Ser	Val	Pro	Lys	
				820					825					830			
25	Arg	Pro	Ala	Gly	Ser	Val	Gln	Asn	Pro	Val	Tyr	His	Asn	Gln	Pro	Leu	
			835					840					845				
30	Asn	Pro	Ala	Pro	Ser	Arg	Asp	Pro	His	Tyr	Gln	Asp	Pro	His	Ser	Thr	
			850				855					860					
35	Ala	Val	Gly	Asn	Pro	Glu	Tyr	Leu	Asn	Thr	Val	Gln	Pro	Thr	Cys	Val	
			865				870				875					880	
40	Asn	Ser	Thr	Phe	Asp	Ser	Pro	Ala	His	Trp	Ala	Gln	Lys	Gly	Ser	His	
				885						890					895		
45	Gln	Ile	Ser	Leu	Asp	Asn	Pro	Asp	Tyr	Gln	Gln	Asp	Phe	Phe	Pro	Lys	
				900					905					910			
50	Glu	Ala	Lys	Pro	Asn	Gly	Ile	Phe	Lys	Gly	Ser	Thr	Ala	Glu	Asn	Ala	
			915					920					925				
55	Glu	Tyr	Leu	Arg	Val	Ala	Pro	Gln	Ser	Ser	Glu	Phe	Ile	Gly	Ala		
			930				935					940					
	<210>	202															
	<211>	121															
	<212>	PRT															
	<213>	Artificial Sequence															
	<220>																
	<223>	Synthetic Construct															
	<220>																
	<221>	misc_feature															
	<222>	(100)..(101)															
	<223>	Xaa can be any naturally occurring amino acid															
	<400>	202															

1 Gln Val Thr Leu Glu Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 5 1 5 10 15
 5 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Ala
 20 25 30
 10 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Pro Glu
 35 40 45
 15 Trp Phe Ala His Ile Phe Ser Thr Asp Glu Lys Ser Leu Arg Leu Ser
 50 55 60
 20 Leu Arg Ser Arg Leu Thr Leu Ser Lys Asp Thr Ser Lys Ser Gln Val
 65 70 75 80
 25 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
 85 90 95
 30 Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
 100 105 110
 35 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120
 40 <210> 203
 <211> 121
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Synthetic Construct
 50 <220>
 <221> misc_feature
 <222> (100)..(101)
 <223> Xaa can be any naturally occurring amino acid
 55 <400> 203
 1 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
 5 10 15
 50 Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Pro
 20 25 30
 55 Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 35 40 45
 Trp Leu Gly His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Ser

	50	55	60
5	Leu Arg Ser Arg Leu Ser Ile Ser Lys Asp Thr Ser Lys Ser Gln Val 65 70 75 80		
10	Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr 85 90 95		
15	Cys Val Arg Xaa Xaa Ser Asn Tyr Gly Gly Tyr Phe Asp Tyr Trp Gly 100 105 110		
20	Gln Gly Thr Leu Val Thr Val Ser Ser 115 120		
25	<210> 204 <211> 121 <212> PRT <213> Artificial Sequence <220> <223> Synthetic Construct <220> <221> misc_feature <222> (100)..(101) <223> Xaa can be any naturally occurring amino acid		
30	<400> 204		
35	Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu 1 5 10 15		
40	Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Pro 20 25 30		
45	Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu 35 40 45		
50	Trp Leu Gly His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Ser 50 55 60		
55	Leu Arg Ser Arg Leu Ser Ile Ser Lys Asp Thr Ser Lys Ser Gln Val 65 70 75 80		
	Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr 85 90 95		
	Cys Val Arg Xaa Xaa Ser Asn Tyr Gly Gly Tyr Phe Asp Tyr Trp Gly 100 105 110		

Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

5 <210> 205
<211> 121
<212> PRT
<213> Artificial Sequence

10 <220>
<223> Synthetic Construct

<220>
<221> misc_feature
<222> (83)..(83)
15 <223> Xaa can be any naturally occurring amino acid

<220>
<221> misc_feature
<222> (100)..(101)
20 <223> Xaa can be any naturally occurring amino acid

<400> 205

25 Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
1 5 10 15

30 Thr Leu Thr Leu Thr Cys Thr Val Phe Gly Phe Ser Leu Ser Asn Pro
20 25 30

Arg Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Pro Glu
35 40 45

35 Trp Leu Gly His Ile Phe Ser Ser Asp Glu Lys Ser Tyr Arg Leu Ser
50 55 60

40 Leu Arg Ser Arg Leu Ser Ile Ser Lys Asp Thr Ser Lys Ser Gln Val
65 70 75 80

45 Val Phe Xaa Met Thr Asn Met Asp Pro Gly Asp Pro Ala Thr Tyr Tyr
85 90 95

Cys Val Arg Xaa Xaa Ser Asn Tyr Glu Glu Tyr Phe Asp Tyr Trp Gly
100 105 110

50 Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

55 <210> 206
<211> 121
<212> PRT
<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (100)..(101)

<223> Xaa can be any naturally occurring amino acid

<400> 206

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Asn Pro
20 25 30

Arg Met Gly Val Ser Trp Leu Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Phe Ala His Ile Phe Ser Thr Asp Glu Lys Ser Tyr Ser Pro Ser
50 55 60

Leu Arg Gly Arg Leu Thr Val Ser Lys Asp Thr Ser Lys Ser Gln Val
65 70 75 80

Val Leu Thr Leu Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 207

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (56)..(57)

<223> Xaa can be any naturally occurring amino acid

<400> 207

	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Glu	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asp	Ala	
					20					25					30		
10	Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
			35					40					45				
15	Gly	Arg	Ile	Lys	Ser	Lys	Thr	Xaa	Xaa	Gly	Thr	Thr	Asp	Tyr	Val	Val	
	50						55					60					
	Pro	Leu	Asn	Gly	Arg	Phe	Ile	Ile	Ser	Arg	Asp	Asp	Ser	Arg	Asn	Thr	
	65					70					75					80	
20																	
	Leu	Tyr	Leu	Gln	Leu	Asn	Asn	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	
					85					90					95		
25	Tyr	Cys	Thr	Thr	Val	Pro	Gly	Ser	Tyr	Gly	Tyr	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
	Leu	Val	Thr	Val	Ser	Ser											
30				115													
	<210>	208															
	<211>	120															
	<212>	PRT															
35	<213>	Artificial Sequence															
	<220>																
	<223>	Synthetic Construct															
40	<220>																
	<221>	misc_feature															
	<222>	(56)..(57)															
	<223>	Xaa can be any naturally occurring amino acid															
45	<400>	208															
50																	
55																	

	Glu	Val	Asn	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Glu	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Tyr	Ala	
				20					25					30			
	Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
10			35				40						45				
	Gly	Arg	Ile	Lys	Ser	Ile	Ala	Xaa	Xaa	Gly	Ala	Thr	Asp	Tyr	Ala	Ala	
		50					55					60					
15	Pro	Val	Arg	Asn	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Arg	Asn	Thr	
	65					70					75					80	
	Leu	Tyr	Leu	Glu	Met	His	Ser	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	
20					85					90					95		
	Tyr	Cys	Thr	Thr	Ile	Pro	Gly	Asn	Asp	Ala	Phe	Asp	Met	Trp	Gly	Gln	
				100					105					110			
25																	
	Gly	Thr	Met	Val	Thr	Val	Ser	Ser									
			115					120									
30	<210> 209																
	<211> 119																
	<212> PRT																
	<213> Artificial Sequence																
35	<220>																
	<223> Synthetic Construct																
	<220>																
	<221> misc_feature																
40	<222> (56)..(57)																
	<223> Xaa can be any naturally occurring amino acid																
	<400> 209																

	Glu	Val	Gln	Leu	Val	Glu	Ser	Trp	Gly	Val	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ile	Phe	Asn	Asn	Ala	
				20					25					30			
10	Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile	
			35					40					45				
15	Gly	Arg	Ile	Lys	Ser	Lys	Ser	Xaa	Xaa	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	
		50					55					60					
20	Pro	Val	Lys	Asp	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asp	Thr	
	65					70					75					80	
25	Leu	Tyr	Leu	Gln	Met	Asn	Gly	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Val	Tyr	
					85				90						95		
30	Phe	Cys	Thr	Thr	Ala	Pro	Gly	Gly	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	
				100					105					110			
35	Thr	Leu	Val	Thr	Val	Ser	Ser										
				115													
40	<210>	210															
	<211>	120															
	<212>	PRT															
	<213>	Artificial Sequence															
45	<220>																
	<223>	Synthetic Construct															
50	<220>																
	<221>	misc_feature															
	<222>	(56)..(57)															
	<223>	Xaa can be any naturally occurring amino acid															
55	<400>	210															

Glu Val Asn Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Tyr Ala
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Gly Arg Ile Lys Ser Ile Thr Xaa Xaa Gly Val Ile Asp Tyr Ala Ala
 50 55 60

Pro Val Arg Asn Arg Cys Thr Ile Ser Arg Asp Asp Ser Arg Asn Thr
 65 70 75 80

Leu Tyr Leu Glu Met His Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
 85 90 95

Tyr Cys Thr Thr Ile Pro Gly Asn Asp Asp Phe Asp Met Trp Gly Gln
 100 105 110

Gly Arg Met Val Thr Val Ser Ser
 115 120

<210> 211
 <211> 112
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic Construct

<220>
 <221> misc_feature
 <222> (33)..(34)
 <223> Xaa can be any naturally occurring amino acid

<400> 211

Asp Ile Val Leu Thr Gln Ser Pro Leu Ser Leu Ser Val Thr Pro Gly
 1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

5 Xaa Xaa Lys Asn Tyr Leu Asp Trp Phe Leu His Lys Pro Gly Gln Ser
 35 40 45

10 Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
 50 55 60

15 Asp Arg Phe Ser Gly Ser Gly Ser Gly Ile Asp Phe Ile Leu Lys Ile
 65 70 75 80

20 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala
 85 90 95

25 Gln Gln Thr Pro Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys
 100 105 110

<210> 212
 <211> 112
 <212> PRT
 <213> Artificial Sequence

30 <220>
 <223> Synthetic Construct

35 <220>
 <221> misc_feature
 <222> (33)..(34)
 <223> Xaa can be any naturally occurring amino acid

40 Asp Ile Val Leu Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15

45 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Arg
 20 25 30

50 Xaa Xaa Phe Asn Tyr Leu Asp Trp Phe Leu Gln Lys Pro Gly Gln Ser
 35 40 45

55 Pro Gln Leu Leu Ile Tyr Leu Ala Ser Ser Arg Ala Ser Gly Val Pro
 50 55 60

65 Asp Arg Phe Ser Gly Ser Asp Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala

85 90 95

5 Leu Gln Thr Pro Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys
100 105 110

<210> 213
<211> 112
<212> PRT
10 <213> Artificial Sequence

<220>
<223> Synthetic Construct

15 <220>
<221> misc_feature
<222> (33)..(34)
<223> Xaa can be any naturally occurring amino acid

20 <400> 213

Asp Ile Val Leu Thr Gln Ser Pro Leu Ser Leu Ser Val Thr Pro Gly
1 5 10 15

25 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

30 Xaa Xaa Lys Asn Tyr Leu Asp Trp Phe Leu His Lys Pro Gly Gln Ser
35 40 45

35 Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
50 55 60

40 Asp Arg Phe Ser Gly Ser Gly Ser Gly Ile Asp Phe Ile Leu Lys Ile
65 70 75 80

45 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala
85 90 95

50 Gln Gln Thr Pro Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys
100 105 110

<210> 214
<211> 120
50 <212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic Construct

55 <220>
<221> misc_feature
<222> (56)..(57)

<223> Xaa can be any naturally occurring amino acid

<400> 214

```

5      Glu Val Asn Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
      1          5          10          15

10     Ser Leu Arg Leu Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Tyr Ala
      20          25          30

15     Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
      35          40          45

20     Gly Arg Ile Lys Ser Ile Asn Xaa Xaa Gly Ala Thr Asp Tyr Ala Ser
      50          55          60

25     Pro Val Arg Asn Arg Phe Thr Ile Ser Arg Asp Asp Ser Arg Asn Met
      65          70          75          80

30     Leu Tyr Leu Glu Met His Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
      85          90          95

35     Tyr Cys Thr Thr Ile Pro Gly Asn Asp Ala Phe Asp Met Trp Gly Gln
      100         105         110

40     Gly Thr Leu Val Thr Val Ser Ser
      115         120

```

35 <210> 215
 <211> 112
 <212> PRT
 <213> Artificial Sequence

40 <220>
 <223> Synthetic Construct

45 <220>
 <221> misc_feature
 <222> (33)..(34)
 <223> Xaa can be any naturally occurring amino acid

<400> 215

50

55

	Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Ser	Val	Thr	Pro	Gly	
	1				5					10					15		
5	Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Thr	Gln	Ser	Leu	Leu	Tyr	Ser	
				20					25					30			
10	Xaa	Xaa	Lys	Asn	Tyr	Leu	Asp	Trp	Phe	Leu	His	Lys	Pro	Gly	Gln	Ser	
			35					40					45				
15	Pro	Gln	Leu	Leu	Ile	Phe	Leu	Gly	Ser	Ile	Arg	Ala	Ser	Gly	Val	Pro	
		50					55					60					
20	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Ile	Asp	Phe	Ile	Leu	Lys	Ile	
	65					70					75					80	
25	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr	Cys	Met	Gln	Ala	
					85					90					95		
30	Gln	Gln	Thr	Pro	Ile	Thr	Phe	Gly	Gln	Gly	Thr	Arg	Leu	Glu	Ile	Lys	
				100					105					110			
35	<210> 216																
	<211> 119																
	<212> PRT																
	<213> Artificial Sequence																
40	<220>																
	<223> Synthetic Construct																
45	<220>																
	<221> misc_feature																
	<222> (56)..(57)																
	<223> Xaa can be any naturally occurring amino acid																
50	<400> 216																
55																	

	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Asp	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Thr	Asn	Ala	
				20					25					30			
10	Trp	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	
			35					40					45				
15	Gly	Arg	Ile	Lys	Ser	Lys	Ile	Xaa	Xaa	Gly	Thr	Thr	Asp	Tyr	Ala	Ala	
		50					55					60					
20	Pro	Val	Lys	Gly	Arg	Phe	Ile	Ile	Ser	Arg	Asp	Asp	Ser	Lys	Asn	Thr	
	65					70				75						80	
25	Leu	Ser	Leu	Gln	Met	Asn	Ser	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Met	Tyr	
					85				90						95		
30	Tyr	Cys	Thr	Thr	Ala	Pro	Gly	Gly	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	
				100					105					110			
35	Ser	Leu	Val	Thr	Val	Ser	Ser										
				115													
40	<210> 217																
	<211> 126																
	<212> PRT																
	<213> Artificial Sequence																
45	<220>																
	<223> Synthetic Construct																
50	<220>																
	<221> misc_feature																
	<222> (114)..(115)																
	<223> Xaa can be any naturally occurring amino acid																
55	<400> 217																

	Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ser	
	1				5					10					15		
5	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Asp	Thr	Phe	Ser	Ser	Asn	
				20					25					30			
10	Ala	Ile	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
			35					40					45				
15	Gly	Val	Ile	Ile	Pro	Ile	Phe	Gly	Thr	Ala	Asp	Tyr	Ala	Gln	Lys	Phe	
		50					55					60					
20	Gln	Gly	Arg	Val	Thr	Ile	Thr	Ala	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	
	65					70					75					80	
25	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		
30	Ala	Arg	His	Thr	Tyr	His	Glu	Tyr	Ala	Gly	Gly	Tyr	Tyr	Gly	Gly	Ala	
				100					105					110			
35	Met	Xaa	Xaa	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
			115					120					125				
40	<210> 218																
	<211> 122																
	<212> PRT																
	<213> Artificial Sequence																
45	<220>																
	<223> Synthetic Construct																
50	<220>																
	<221> misc_feature																
	<222> (62)..(63)																
	<223> Xaa can be any naturally occurring amino acid																
55	<400> 218																

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

5 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

10 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Asp Ile Ser Gly Gly Gly Gly Arg Thr Tyr Tyr Ala Xaa Xaa Val
 50 55 60

15 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

20 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

25 Ala Arg Ala Gly Leu Leu Tyr Gly Gly Gly Val Tyr Pro Met Asp Ile
 100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 115 120

30 <210> 219
 <211> 11
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Synthetic Construct

40 <220>
 <221> misc_feature
 <222> (1)..(2)
 <223> Xaa can be any naturally occurring amino acid

45 <400> 219

Xaa Xaa Ser Asn Tyr Glu Gly Tyr Phe Asp Phe
 1 5 10

50 <210> 220
 <211> 11
 <212> PRT
 <213> Artificial Sequence

55 <220>
 <223> Synthetic Construct

<220>
 <221> misc_feature

<222> (1)..(2)

<223> Xaa can be any naturally occurring amino acid

<400> 220

5

Xaa	Xaa	Ser	Asn	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr
1				5					10	

<210> 221

10

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

15

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (1)..(2)

20

<223> Xaa can be any naturally occurring amino acid

<400> 221

25

Xaa	Xaa	Ser	Asn	Tyr	Gly	Gly	Tyr	Phe	Asp	Tyr
1				5					10	

<210> 222

<211> 11

<212> PRT

30

<213> Artificial Sequence

<220>

<223> Synthetic Construct

35

<220>

<221> misc_feature

<222> (1)..(2)

<223> Xaa can be any naturally occurring amino acid

40

<400> 222

Xaa	Xaa	Ser	Asp	Tyr	Glu	Gly	Tyr	Phe	Asp	Tyr
1				5					10	

45

<210> 223

<211> 11

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

55

<222> (1)..(2)

<223> Xaa can be any naturally occurring amino acid

<400> 223

		Xaa	Xaa	Ser	Asn	Tyr	Glu	Glu	Tyr	Phe	Asp	Tyr
		1				5					10	

5 <210> 224
 <211> 19
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Synthetic Construct

15 <220>
 <221> misc_feature
 <222> (7)..(8)
 <223> Xaa can be any naturally occurring amino acid

20 <400> 224

		Arg	Ile	Lys	Ser	Lys	Thr	Xaa	Xaa	Gly	Thr	Thr	Asp	Tyr	Val	Val	Pro
		1				5					10					15	

25 Leu Asn Gly

25 <210> 225
 <211> 13
 <212> PRT
 <213> Artificial Sequence

30 <220>
 <223> Synthetic Construct

35 <220>
 <221> misc_feature
 <222> (7)..(8)
 <223> Xaa can be any naturally occurring amino acid

40 <400> 225

		Arg	Ile	Lys	Ser	Lys	Thr	Xaa	Xaa	Gly	Thr	Thr	Asp	Tyr
		1				5					10			

45 <210> 226
 <211> 19
 <212> PRT
 <213> Artificial Sequence

50 <220>
 <223> Synthetic Construct

55 <220>
 <221> misc_feature
 <222> (7)..(8)
 <223> Xaa can be any naturally occurring amino acid

55 <400> 226

Arg Ile Lys Ser Ile Ala Xaa Xaa Gly Ala Thr Asp Tyr Ala Ala Pro
 1 5 10 15

5 Val Arg Asn

<210> 227

<211> 13

<212> PRT

10 <213> Artificial Sequence

<220>

<223> Synthetic Construct

15 <220>

<221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

20 <400> 227

Arg Ile Lys Ser Ile Ala Xaa Xaa Gly Ala Thr Asp Tyr
 1 5 10

25 <210> 228

<211> 19

<212> PRT

<213> Artificial Sequence

30 <220>

<223> Synthetic Construct

<220>

<221> misc_feature

35 <222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

<400> 228

40 Arg Ile Lys Ser Lys Ser Xaa Xaa Gly Thr Thr Asp Tyr Ala Ala Pro
 1 5 10 15

Val Lys Asp

45

<210> 229

<211> 13

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Synthetic Construct

<220>

55 <221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

<400> 229

Arg	Ile	Lys	Ser	Lys	Ser	Xaa	Xaa	Gly	Thr	Thr	Asp	Tyr
1				5					10			

5

<210> 230

<211> 19

<212> PRT

<213> Artificial Sequence

10

<220>

<223> Synthetic Construct

<220>

15

<221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

<400> 230

20

Arg	Ile	Lys	Ser	Ile	Thr	Xaa	Xaa	Gly	Val	Ile	Asp	Tyr	Ala	Ala	Pro
1				5					10					15	

25

Val Arg Asn

<210> 231

<211> 13

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Synthetic Construct

35

<220>

<221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

40

<400> 231

Arg	Ile	Lys	Ser	Ile	Thr	Xaa	Xaa	Gly	Val	Ile	Asp	Tyr
1				5					10			

45

<210> 232

<211> 19

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Synthetic Construct

<220>

55

<221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

<400> 232

Arg Ile Lys Ser Ile Asn Xaa Xaa Gly Ala Thr Asp Tyr Ala Ser Pro
 1 5 10 15

5 Val Arg Asn

<210> 233

<211> 13

<212> PRT

10 <213> Artificial Sequence

<220>

<223> Synthetic Construct

15 <220>

<221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

20 <400> 233

Arg Ile Lys Ser Ile Asn Xaa Xaa Gly Ala Thr Asp Tyr
 1 5 10

25 <210> 234

<211> 19

<212> PRT

<213> Artificial Sequence

30 <220>

<223> Synthetic Construct

<220>

<221> misc_feature

35 <222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

<400> 234

40 Arg Ile Lys Ser Lys Ile Xaa Xaa Gly Thr Thr Asp Tyr Ala Ala Pro
 1 5 10 15

Val Lys Gly

45

<210> 235

<211> 13

<212> PRT

<213> Artificial Sequence

50

<220>

<223> Synthetic Construct

<220>

55 <221> misc_feature

<222> (7)..(8)

<223> Xaa can be any naturally occurring amino acid

<400> 235

Arg Ile Lys Ser Lys Ile Xaa Xaa Gly Thr Thr Asp Tyr
 1 5 10

<210> 236

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (16)..(17)

<223> Xaa can be any naturally occurring amino acid

<400> 236

His Thr Tyr His Glu Tyr Ala Gly Gly Tyr Tyr Gly Gly Ala Met Xaa
 1 5 10 15

Xaa

<210> 237

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (13)..(14)

<223> Xaa can be any naturally occurring amino acid

<400> 237

Asp Ile Ser Gly Gly Gly Gly Arg Thr Tyr Tyr Ala Xaa Xaa Val Lys
 1 5 10 15

Gly

<210> 238

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> misc_feature

<222> (10)..(11)

<223> Xaa can be any naturally occurring amino acid

<400> 238

5 Arg Ser Ser Gln Ser Leu Leu Tyr Ser Xaa Xaa Lys Asn Tyr Leu Asp
 1 5 10 15

<210> 239

<211> 16

10 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

15

<220>

<221> misc_feature

<222> (10)..(11)

<223> Xaa can be any naturally occurring amino acid

20

<400> 239

 Arg Ser Ser Gln Ser Leu Leu His Arg Xaa Xaa Phe Asn Tyr Leu Asp
 1 5 10 15

25

<210> 240

<211> 121

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Synthetic Construct

<400> 240

35

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

40

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30

45

50

55

Tyr Met Thr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

5 Ala Phe Ile Arg Asn Arg Ala Arg Gly Tyr Thr Ser Asp His Asn Pro
 50 55 60

10 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser
 65 70 75 80

15 Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
 85 90 95

Tyr Cys Ala Arg Asp Arg Pro Ser Tyr Tyr Val Leu Asp Tyr Trp Gly
 100 105 110

20 Gln Gly Thr Thr Val Thr Val Ser Ser
 115 120

25 <210> 241
 <211> 112
 <212> PRT
 <213> Artificial Sequence

30 <220>
 <223> Synthetic Construct

35 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

40 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Phe Asn Val
 20 25 30

Arg Ser Arg Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

45 Pro Pro Lys Leu Leu Ile Ser Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60

50 Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Lys Gln
 85 90 95

55 Ser Tyr Asp Leu Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 242
 <211> 363
 <212> **DNA**
 <213> Artificial Sequence

5

<220>
 <223> Synthetic Construct

10

<400> 242

gaagtccaac ttgtcgaatc gggaggaggc cttgtgcaac ccggtggatc cctgaggctg 60
 tcatgcgcgg cctcgggctt caccttttcc gattactaca tgacctgggt cagacaggcc 120
 cctggaaaagg ggttggaatg ggtggcattc atccggaata gagcccgcgg atacacttcc 180
 gaccacaacc ccagcgtgaa ggggcggttc accattagcc gcgacaacgc caagaactcc 240
 ctctacctcc aaatgaacag cctgcgggcg gaggataccg ctgtgtacta ctgcgcccgc 300
 gaccggccgt cctactatgt gctggactac tggggccagg gtactacggt caccgtctcc 360
 tca 363

25

<210> 243
 <211> 336
 <212> **DNA**
 <213> Artificial Sequence

30

<220>
 <223> Synthetic Construct

<400> 243

35

gacattgtga tgactcaatc ccccgactcc ctggctgtgt ccctcggcga acgcgcaact 60
 atcaactgta aaagcagcca gtccctgttc aacgtccggt cgaggaagaa ctacctggcc 120
 tggatatcagc agaaacctgg gcagccgccg aagcttctga tctcatgggc ctcaactcgg 180
 gaaagcggag tgccagatag attctccgga tctggctccg gaaccgactt caccctgacg 240
 atttcgagct tgcaagcgga ggatgtggcc gtgtactact gcaagcagtc ctacgacctc 300
 ttcacctttg gttcgggcac caagctggag atcaaa 336

45

<210> 244
 <211> 6
 <212> **PRT**
 <213> Artificial Sequence

50

<220>
 <223> Synthetic Construct

55

<400> 244

Ser Asp Tyr Tyr Met Thr
 1 5

<210> 245
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Construct
 <400> 245
 10
 Gly Phe Thr Phe Ser Asp Tyr Tyr Met Thr
 1 5 10
 <210> 246
 15
 <211> 13
 <212> PRT
 <213> Artificial Sequence
 <220>
 20
 <223> Synthetic Construct
 <400> 246
 25
 Phe Ile Arg Asn Arg Ala Arg Gly Tyr Thr Ser Asp His
 1 5 10
 <210> 247
 <211> 19
 <212> PRT
 30
 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 35
 <400> 247
 Phe Ile Arg Asn Arg Ala Arg Gly Tyr Thr Ser Asp His Asn Pro Ser
 1 5 10 15
 40
 Val Lys Gly
 <210> 248
 <211> 10
 45
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 50
 <400> 248
 Asp Arg Pro Ser Tyr Tyr Val Leu Asp Tyr
 1 5 10
 55
 <210> 249
 <211> 17
 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

5

<400> 249

10

Lys	Ser	Ser	Gln	Ser	Leu	Phe	Asn	Val	Arg	Ser	Arg	Lys	Asn	Tyr	Leu
1				5					10					15	

Ala

15

<210> 250

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

20

<223> Synthetic Construct

<400> 250

25

Trp Ala Ser Thr Arg Glu Ser
1 5

<210> 251

$\langle 211 \rangle$ 8

<212> PRT

30

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Synthetic Construct

35

<400> 251

Lys Gln Ser Tyr Asp Leu Phe Thr
1 5

40

<210> 252

<211> 326

<212> PRT

<213> Artificial Sequence

45

$\langle 220 \rangle$

<223> Synthetic Construct

<400> 252

50

55

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15

5 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

10 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser

15

20

25

30

35

40

45

50

55

	50		55		60												
5	Leu 65	Ser	Ser	Val	Val	Thr 70	Val	Pro	Ser	Ser	Asn 75	Phe	Gly	Thr	Gln	Thr 80	
10	Tyr	Thr	Cys	Asn 85	Val	Asp	His	Lys	Pro	Ser	Asn 90	Thr	Lys	Val	Asp 95	Lys	
15	Thr	Val	Glu	Arg 100	Lys	Cys	Arg	Val	Arg 105	Cys	Pro	Arg	Cys	Pro 110	Ala	Pro	
20	Pro	Val	Ala 115	Gly	Pro	Ser	Val	Phe 120	Leu	Phe	Pro	Pro	Lys 125	Pro	Lys	Asp	
25	Thr	Leu 130	Met	Ile	Ser	Arg	Thr 135	Pro	Glu	Val	Thr	Cys 140	Val	Val	Val	Ala	
30	Val 145	Ser	His	Glu	Asp 150	Pro	Glu	Val	Gln	Phe	Asn 155	Trp	Tyr	Val	Asp	Gly 160	
35	Val	Glu	Val	His 165	Asn	Ala	Lys	Thr	Lys 170	Pro	Arg	Glu	Glu	Gln	Phe	Asn 175	
40	Ser	Thr	Phe	Arg 180	Val	Val	Ser	Val	Leu 185	Thr	Val	Val	His	Gln	Asp 190	Trp	
45	Leu	Asn 195	Gly	Lys	Glu	Tyr	Lys 200	Cys	Lys	Val	Ser	Asn 205	Lys	Gly	Leu	Pro	
50	Ser 210	Ser	Ile	Glu	Lys	Thr 215	Ile	Ser	Lys	Thr	Lys	Gly 220	Gln	Pro	Arg	Glu	
55	Pro 225	Gln	Val	Tyr	Thr 230	Leu	Pro	Pro	Ser	Arg	Glu 235	Glu	Met	Thr	Lys	Asn 240	
	Gln	Val	Ser	Leu 245	Thr	Cys	Leu	Val	Lys	Gly 250	Phe	Tyr	Pro	Ser	Asp 255	Ile	
	Ala	Val	Glu	Trp 260	Glu	Ser	Asn	Gly	Gln 265	Pro	Glu	Asn	Asn	Tyr 270	Lys	Thr	
	Thr	Pro	Pro	Met 275	Leu	Asp	Ser	Asp 280	Gly	Ser	Phe	Phe	Leu 285	Tyr	Ser	Arg	
	Leu 290	Thr	Val	Asp	Lys	Ser	Arg 295	Trp	Gln	Gln	Gly	Asn 300	Val	Phe	Ser	Cys	

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
305 310 315 320

5 Ser Leu Ser Pro Gly Lys
325

<210> 253

<211> 978

10 <212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

15

<400> 253

gcctccacca agggcccatc ggtcttcccc ctggcgccct gctccaggag cacctccgag 60
20 agcacagcgg ccctgggctg cctgggtcaag gactacttcc ccgaaccggt gacgggtgtcg 120
tggaactcag gcgctctgac cagcggcgtg cacaccttcc cggctgtcct acagtccctca 180
ggactctact ccctcagcag cgtagtgtacc gtgccctcca gcaacttcgg caccagacc 240
25 tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc 300
aaatgtcgtg tcagggtgccc aagggtgccc gcaccacctg tggcaggacc gtcagtcttc 360
30 ctcttccccc caaaacccaa ggacaccctc atgatctccc ggaccctga ggtcacgtgc 420
gtgggtgggtg ccgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc 480
gtggaggtgc ataattgcaa gacaaagcca cgggaggagc agttcaacag cacgttccgt 540
35 gtgggtcagcg tcctcaccgt cgtgcaccag gactgggtga acggcaagga gtacaagtgc 600
aagggtctcca acaaaggcct cccatcctcc atcgagaaaa ccatctccaa aaccaaaggg 660
cagccccgag aaccacaggt gtacaccctg ccccatccc gggaggagat gaccaagaac 720
40 caggtcagcc tgacctgcct ggtcaaaggc ttctacccca gcgacatcgc cgtggagtgg 780
gagagcaatg ggcagccgga gaacaactac aagaccacac ctcccatgct ggactccgac 840
45 ggctccttct tcctctacag caggctcacc gtggacaaga gcagggtggca gcaggggaac 900
gtcttctcat gtcctgtgat gcatgaggct ctgcacaacc actacacaca gaagagcctc 960
tccctgtctc cgggtaaa 978

50

<210> 254

<211> 326

<212> PRT

<213> Artificial Sequence

55

<220>

<223> Synthetic Construct

<400> 254

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg

5

10

15

20

25

30

35

40

45

50

55

	1			5					10				15			
5	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
				20					25					30		
10	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
			35					40					45			
15	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
		50					55					60				
20	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Asn	Phe	Gly	Thr	Gln	Thr
	65					70					75					80
25	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
				85						90					95	
30	Thr	Val	Glu	Arg	Lys	Cys	Glu	Val	Glu	Cys	Pro	Glu	Cys	Pro	Ala	Pro
				100					105					110		
35	Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp
			115					120					125			
40	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Ala
		130					135					140				
45	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly
	145					150					155					160
50	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn
				165						170					175	
55	Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Val	His	Gln	Asp	Trp
				180					185					190		
60	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro
			195					200					205			
65	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Gln	Pro	Arg	Glu
		210					215					220				
70	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn
	225					230					235					240
75	Gln	Val	Ser	Leu	Thr	Cys	Glu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile
				245						250					255	

	Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr	
	260	265 270
5	Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys	
	275	280 285
10	Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys	
	290	295 300
15	Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu	
	305	310 315 320
	Ser Leu Ser Pro Gly Lys	
	325	
20	<210> 255	
	<211> 978	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Synthetic Construct	
	<400> 255	
30		
35		
40		
45		
50		
55		

gcctccacca agggcccatc ggtcttcccc ctggcgccct gctccaggag cacctccgag 60
 agcacagcgg ccctgggctg cctggtcaag gactacttcc ccgaaccggt gacggtgtcg 120
 5 tggaaactcag gcgctctgac cagcggcgtg cacaccttcc cggctgtcct acagtcctca 180
 ggactctact ccctcagcag cgtagtacc gtgccctcca gcaacttcgg caccagacc 240
 10 tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc 300
 aaatgtgagg tcgagtgcc agagtgtcca gcaccacctg tggcaggacc gtcagtcttc 360
 ctcttcccc caaaacccaa ggacaccctc atgatctccc ggaccctga ggtcacgtgc 420
 15 gtggtggtgg ccgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc 480
 gtggagggtgc ataatgccaa gacaaagcca cgggaggagc agttcaacag cacgttccgt 540
 gtggtcagcg tcctcaccgt cgtgcaccag gactggctga acggcaagga gtacaagtgc 600
 20 aaggtctcca acaaaggcct cccatcctcc atcgagaaaa ccatctccaa aaccaaaggg 660
 cagccccgag aaccacaggt gtacaccctg ccccatccc gggaggagat gaccaagaac 720
 caggtcagcc tgacctgca ggtcaaaggc ttctaccca gcgacatcg cgtggagtgg 780
 25 gagagcaatg ggcagccgga gaacaactac aagaccacac ctcccatgct ggactccgac 840
 ggctccttct tcctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac 900
 30 gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacaca gaagagcctc 960
 tccctgtctc cgggtaaa 978

35 <210> 256
 <211> 107
 <212> PRT
 <213> Homo sapiens

40 <400> 256

45

50

55

Gly Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 1 5 10 15
 5 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 20 25 30
 10 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 35 40 45
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 50 55 60
 15 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 65 70 75 80
 20 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 85 90 95
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 25 100 105

<210> 257

<211> 324

<212> DNA

30 <213> Homo sapiens

<400> 257

35 ggaactgtgg ctgcaccatc tgtcttcac tccccgccat ctgatgagca gttgaaatct 60
 ggaactgcct ctgttgtgtg cctgctgaat aacttctatc ccagagaggc caaagtacag 120
 tggaagggtg ataacgccct ccaatcgggt aactcccagg agagtgtcac agagcaggac 180
 40 agcaaggaca gcacctacag cctcagcagc accctgacgc tgagcaaagc agactacgag 240
 aaacacaaaag tctacgcctg cgaagtcacc catcagggcc tgagctcgcc cgtcacaaag 300
 agcttcaaca ggggagagtg ttag 324

<210> 258

<211> 8

<212> PRT

50 <213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 258

55 Leu Gln Gly Leu Leu Gln Gly Gly
 1 5

<210> 259
 <211> 4
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Construct
 <400> 259
 10
 Leu Leu Gln Gly
 1
 <210> 260
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 20
 <400> 260
 Leu Ser Leu Ser Gln Gly
 1 5
 25
 <210> 261
 <211> 8
 <212> PRT
 <213> Artificial Sequence
 30
 <220>
 <223> Synthetic Construct
 35
 <400> 261
 Gly Gly Gly Leu Leu Gln Gly Gly
 1 5
 40
 <210> 262
 <211> 5
 <212> PRT
 <213> Artificial Sequence
 45
 <220>
 <223> Synthetic Construct
 <400> 262
 50
 Gly Leu Leu Gln Gly
 1 5
 55
 <210> 263
 <211> 13
 <212> PRT
 <213> Artificial Sequence
 <220>

<223> Synthetic Construct

<400> 263

5 Leu Leu Gln Gly Ser Pro Leu Ala Gln Ser His Gly Gly
 1 5 10

<210> 264

<211> 7

10 <212> PRT

<213> Artificial Sequence

<220>

15 <223> Synthetic Construct

<400> 264

20 Gly Leu Leu Gln Gly Gly Gly
 1 5

<210> 265

<211> 6

<212> PRT

25 <213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 265

30 Gly Leu Leu Gln Gly Gly
 1 5

<210> 266

35 <211> 4

<212> PRT

<213> Artificial Sequence

<220>

40 <223> Synthetic Construct

<400> 266

45 Gly Leu Leu Gln
 1

<210> 267

<211> 8

<212> PRT

50 <213> Artificial Sequence

<220>

<223> Synthetic Construct

55 <400> 267

Leu Leu Gln Leu Leu Gln Gly Ala
1 5

<210> 268
 <211> 5
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Construct
 10
 <400> 268
 Leu Leu Gln Gly Ala
 1 5
 <210> 269
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 20
 <400> 269
 Leu Leu Gln Tyr Gln Gly Ala
 1 5
 <210> 270
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 30
 <220>
 <223> Synthetic Construct
 35
 <400> 270
 Leu Leu Gln Gly Ser Gly
 1 5
 <210> 271
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct
 45
 <400> 271
 Leu Leu Gln Tyr Gln Gly
 1 5
 <210> 272
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 55
 <220>

<223> Synthetic Construct

<400> 272

5 Leu Leu Gln Leu Leu Gln Gly
 1 5

<210> 273

<211> 5

10 <212> PRT

<213> Artificial Sequence

<220>

15 <223> Synthetic Construct

<400> 273

 Ser Leu Leu Gln Gly
 1 5

20

<210> 274

<211> 5

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Synthetic Construct

<400> 274

30

 Leu Leu Gln Leu Gln
 1 5

<210> 275

35 <211> 6

<212> PRT

<213> Artificial Sequence

<220>

40 <223> Synthetic Construct

<400> 275

 Leu Leu Gln Leu Leu Gln
 1 5

45

<210> 276

<211> 5

<212> PRT

50 <213> Artificial Sequence

<220>

<223> Synthetic Construct

55

<400> 276

 Leu Leu Gln Gly Arg
 1 5

<210> 277
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Construct
 <400> 277
 10
 Leu Leu Gln Gly Pro Pro
 1 5
 <210> 278
 15
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 <220>
 20
 <223> Synthetic Construct
 <400> 278
 25
 Leu Leu Gln Gly Pro Ala
 1 5
 <210> 279
 <211> 8
 <212> PRT
 30
 <213> Artificial Sequence
 <220>
 <223> Synthetic Construct
 35
 <400> 279
 Gly Gly Leu Leu Gln Gly Pro Pro
 1 5
 40
 <210> 280
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 45
 <220>
 <223> Synthetic Construct
 <400> 280
 50
 Gly Gly Leu Leu Gln Gly Ala
 1 5
 <210> 281
 <211> 7
 55
 <212> PRT
 <213> Artificial Sequence
 <220>

<223> Synthetic Construct

<400> 281

5 Leu Leu Gln Gly Pro Gly Lys
 1 5

<210> 282

<211> 6

10 <212> PRT

<213> Artificial Sequence

<220>

15 <223> Synthetic Construct

<400> 282

 Leu Leu Gln Gly Pro Gly
 1 5

20

<210> 283

<211> 5

<212> PRT

<213> Artificial Sequence

25

<220>

<223> Synthetic Construct

<400> 283

30

 Leu Leu Gln Gly Pro
 1 5

<210> 284

35 <211> 4

<212> PRT

<213> Artificial Sequence

<220>

40 <223> Synthetic Construct

<400> 284

 Leu Leu Gln Pro
 1

45

<210> 285

<211> 6

<212> PRT

50 <213> Artificial Sequence

<220>

<223> Synthetic Construct

55 <400> 285

 Leu Leu Gln Pro Gly Lys
 1 5

<210> 286
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <223> Synthetic Construct
 <400> 286
 10
 Leu Leu Gln Ala Pro Gly Lys
 1 5
 <210> 287
 <211> 7
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Synthetic Construct
 20
 <400> 287
 Leu Leu Gln Gly Ala Pro Gly
 1 5
 <210> 288
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 30
 <220>
 <223> Synthetic Construct
 35
 <400> 288
 Leu Leu Gln Gly Ala Pro
 1 5
 <210> 289
 <211> 6
 <212> PRT
 <213> Artificial Sequence
 40
 <220>
 <223> Synthetic Construct
 45
 <400> 289
 Leu Leu Gln Leu Gln Gly
 1 5
 <210> 290
 <211> 326
 <212> PRT
 <213> Homo sapiens
 55
 <400> 290

	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg	
	1				5					10					15		
5	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	
				20					25					30			
10	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	
			35					40					45				
15	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	
	50						55					60					
20	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Asn	Phe	Gly	Thr	Gln	Thr	
	65					70					75					80	
25	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	
				85						90					95		
30	Thr	Val	Glu	Arg	Lys	Cys	Cys	Val	Glu	Cys	Pro	Pro	Cys	Pro	Ala	Pro	
				100					105					110			
35	Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	
			115					120					125				
40	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	
	130						135					140					
45	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	
	145					150					155					160	
50	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	
				165					170						175		
55	Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Val	His	Gln	Asp	Trp	
				180					185					190			
60	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	
			195					200					205				
65	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Gln	Pro	Arg	Glu	
	210						215					220					
70	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	

	225		230		235		240
5	Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile	245		250		255	
10	Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr	260		265		270	
15	Thr Pro Pro Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys	275		280		285	
20	Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys	290		295		300	
25	Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu	305		310		315	320
30	Ser Leu Ser Pro Gly Lys	325					
35	<210> 291						
40	<211> 330						
45	<212> PRT						
50	<213> Homo sapiens						
55	<400> 291						

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 5 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 10 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 15 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 20 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 25 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 30
 35
 40
 45
 50
 55

	115	120	125
5	Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys 130 135 140		
10	Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp 145 150 155 160		
15	Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu 165 170 175		
20	Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu 180 185 190		
25	His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn 195 200 205		
30	Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly 210 215 220		
35	Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu 225 230 235 240		
40	Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr 245 250 255		
45	Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn 260 265 270		
50	Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe 275 280 285		
55	Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn 290 295 300		
	Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr 305 310 315 320		
	Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 325 330		

<210> 292

<211> 327

<212> PRT

<213> Homo sapiens

<400> 292

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg

5

10

15

20

25

30

35

40

45

50

55

	1			5				10				15				
5	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
				20					25					30		
10	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
			35					40					45			
15	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
		50					55					60				
20	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Lys	Thr
	65					70					75					80
25	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
				85						90					95	
30	Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro
				100					105					110		
35	Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys
			115					120					125			
40	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val
		130					135					140				
45	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp
	145					150					155					160
50	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe
				165						170					175	
55	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp
			180					185						190		
60	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu
			195					200					205			
65	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg
		210					215					220				
70	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys
	225					230					235					240
75	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp
				245						250					255	

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270

5 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285

10 Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320

15 Leu Ser Leu Ser Leu Gly Lys
 325

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2013185010 A1 [0004]
- US 4816567 A [0038] [0129] [0147]
- WO 9958572 A [0039] [0151] [0165]
- US 5545807 A [0040]
- US 5545806 A [0040]
- US 5569825 A [0040]
- US 5625126 A [0040]
- US 5633425 A [0040]
- US 5661016 A [0040]
- US 5750373 A [0040]
- US 6737056 B [0063]
- US 5500362 A [0064]
- US 5821337 A [0064]
- WO 2012059882 A [0073] [0106] [0204]
- WO 2015015448 A [0073] [0204]
- US 20160297885 A [0101]
- WO 9404690 A [0103]
- US 2011036419 W [0104] [0105]
- WO 2011143545 A [0104] [0105]
- WO IB2011054899 A [0106]
- GB 9901441 W [0114] [0130]
- GB 98099518 A [0114]
- US 7314622 B [0127]
- US 5807715 A [0129] [0147]
- US 5866692 A [0129] [0130] [0151]
- US 6331415 B [0129] [0147]
- US 5530101 A [0129]
- US 5693761 A [0129]
- US 5693762 A [0129]
- US 5585089 A [0129]
- US 6180370 B [0129]
- EP 0519596 A [0130]
- GB 9809951 A [0130]
- US 6180377 B [0130]
- US 6054297 A [0130]
- US 5997867 A [0130] [0151]
- US 6210671 B [0130]
- US 6350861 B [0130]
- WO 0127160 A [0130]
- US 5565332 A [0133]
- US 5580717 A [0133]
- US 5733743 A [0133]
- US 6265150 B [0133]
- WO 9306213 A [0133]
- WO 8704462 A [0137] [0200] [0202]
- US 6436908 B [0142]
- US 6413942 B [0142]
- US 6376471 B [0142]
- WO 9007936 A [0144]
- WO 9403622 PCT [0144]
- WO 9325698 PCT [0144]
- WO 9325234 PCT [0144]
- WO 9311230 PCT [0144]
- WO 9310218 PCT [0144]
- WO 9102805 PCT [0144]
- US 5219740 A [0144]
- US 4777127 A [0144]
- GB 2200651 A [0144]
- EP 0345242 A [0144]
- WO 9412649 A [0144]
- WO 9303769 PCT [0144]
- WO 9319191 PCT [0144]
- WO 9428938 PCT [0144]
- WO 9511984 PCT [0144]
- WO 9500655 PCT [0144]
- US 5814482 A [0145]
- WO 9507994 A [0145]
- WO 9617072 PCT [0145]
- WO 9530763 PCT [0145]
- WO 9742338 PCT [0145]
- WO 9011092 A [0145]
- US 5580859 A [0145]
- US 5422120 A [0145]
- WO 9513796 A [0145]
- WO 9423697 PCT [0145]
- WO 9114445 PCT [0145]
- EP 0524968 A [0145]
- US 4676980 A [0149]
- WO 9100360 A [0149]
- WO 92200373 PCT [0149]
- EP 03089 A [0149]
- US 5047335 A [0163]
- US 5510261 A [0163]
- US 5278299 A [0163]
- WO 2004058184 A [0166]
- US 4683195 A [0197]
- US 4800159 A [0197]
- US 4754065 A [0197]
- US 4683202 A [0197]
- US 5663149 A [0210]
- US 20130129753 A [0210]
- WO IB2015050280 A [0217]
- US 20130115208 A [0276]

Non-patent literature cited in the description

- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press, 1989 [0032]
- Oligonucleotide Synthesis. 1984 [0032]
- Methods in Molecular Biology. Humana Press [0032]
- Cell Biology: A Laboratory Notebook. Academic Press, 1998 [0032]
- Animal Cell Culture. 1987 [0032]
- **J.P. MATHER ; P.E. ROBERTS.** Introduction to Cell and Tissue Culture. Plenum Press, 1998 [0032]
- Cell and Tissue Culture: Laboratory Procedures. J. Wiley and Sons, 1993 [0032]
- Methods in Enzymology. Academic Press, Inc, [0032]
- Handbook of Experimental Immunology [0032]
- Gene Transfer Vectors for Mammalian Cells. 1987 [0032]
- Current Protocols in Molecular Biology. 1987 [0032]
- PCR: The Polymerase Chain Reaction. 1994 [0032]
- Current Protocols in Immunology. 1991 [0032]
- Short Protocols in Molecular Biology. Wiley and Sons, 1999 [0032]
- **C.A. JANEWAY ; P. TRAVERS.** Immunobiology, 1997 [0032]
- **P. FINCH.** Antibodies, 1997 [0032]
- Antibodies: a practical approach. IRL Press, 1988 [0032]
- Monoclonal antibodies: a practical approach. Oxford University Press, 2000 [0032]
- **E. HARLOW ; D. LANE.** Using antibodies: a laboratory manual. Cold Spring Harbor Laboratory Press, 1999 [0032]
- The Antibodies. Harwood Academic Publishers, 1995 [0032]
- **WARD et al.** *Nature*, 1989, vol. 341, 544-546 [0034]
- **KABAT et al.** Sequences of Proteins of Immunological Interest. National Institutes of Health, 1991 [0036]
- **AI-IAZIKANI et al.** *J. Molec. Biol.*, 1997, vol. 273, 927-948 [0036]
- **KABAT et al.** Sequences of Proteins of Immunological Interest. Public Health Service, NIH, 1992 [0037]
- **CHOTHIA et al.** *Nature*, 1989, vol. 342, 877-883 [0037]
- **MACCALLUM et al.** *J. Mol. Biol.*, 1996, vol. 262, 732-745 [0037]
- **MAKABE et al.** *Journal of Biological Chemistry*, 2008, vol. 283, 1156-1166 [0037]
- **KOHLER ; MILSTEIN.** *Nature*, 1975, vol. 256, 495 [0038]
- **MCCAFFERTY et al.** *Nature*, 1990, vol. 348, 552-554 [0038]
- **VAUGHAN et al.** *Nature Biotechnology*, 1996, vol. 14, 309-314 [0040]
- **SHEETS et al.** *Proc. Natl. Acad. Sci. (USA)*, 1998, vol. 95, 6157-6162 [0040]
- **HOOGENBOOM ; WINTER.** *J. Mol. Biol.*, 1991, vol. 227, 381 [0040]
- **MARKS et al.** *J. Mol. Biol.*, 1991, vol. 222 (581 [0040]
- **COLE et al.** Monoclonal Antibodies and Cancer Therapy. Alan R. Liss, 1985, 77 [0040]
- **BOERNER et al.** *J. Immunol.*, 1991, vol. 147 (1), 86-95 [0040]
- **JANEWAY et al.** Immunobiology: the immune system in health and disease. Elsevier Science Ltd, 1999 [0050]
- **BLOOM et al.** Protein Science. 1997, vol. 6, 407-415 [0050]
- **HUMPHREYS et al.** *J. Immunol. Methods*, 1997, vol. 209, 193-202 [0050]
- **JAYASENA, S.D.** *Clin. Chem.*, 1999, vol. 45, 1628-50 [0053]
- **FELLOUSE, F.A. et al.** *J. Mol. Biol.*, 2007, vol. 373 (4), 924-40 [0053]
- **KABAT et al.** Sequences of Proteins of Immunological Interest. Public Health Service, National Institutes of Health, 1991 [0058]
- **RAVETCH ; KINET.** *Ann. Rev. Immunol.*, 1991, vol. 9, 457-92 [0059]
- **CAPEL et al.** *Immunomethods*, 1994, vol. 4, 25-34 [0059]
- **DE HAAS et al.** *J. Lab. Clin. Med.*, 1995, vol. 126, 330-41 [0059]
- **GUYER et al.** *J. Immunol.*, 1976, vol. 117, 587 [0059]
- **KIM et al.** *J. Immunol.*, 1994, vol. 24, 249 [0059]
- **CLYNES et al.** *PNAS (USA)*, 1998, vol. 95, 652-656 [0064]
- **GAZZANO-SANTORO et al.** *J. Immunol. Methods*, 1996, vol. 202, 163 [0065]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990 [0072]
- Remington, The Science and Practice of Pharmacy. Mack Publishing, 2005 [0072]
- **MAKABE et al.** *J. Biol. Chem.*, 2007, vol. 283, 1156-1166 [0088]
- **SURESH et al.** *Methods in Enzymology*, 1986, vol. 121, 210 [0090]
- **MILLSTEIN ; CUELLO.** *Nature*, 1983, vol. 305, 537-539 [0090]
- *Eur. J. Immunol.*, 1999, vol. 29, 2613-2624 [0114]
- **ARMOUR et al.** *Molecular Immunology*, 2003, vol. 40, 585-593 [0114]
- **AALBERSE et al.** *Immunology*, 2002, vol. 105, 9-19 [0114]
- **KARLSSON, R. ROOS ; H. FAGERSTAM ; L. PETERSSON, B.** *Methods Enzymology*, 1994, vol. 6, 99-110 [0121]
- **KOHLER, B. ; MILSTEIN, C.** *Nature*, 1975, vol. 256, 495-497 [0124]
- **BUCK, D. W. et al.** *Vitro*, 1982, vol. 18, 377-381 [0124]

- TILLER et al. *J. Immunol. Methods*, 2008, vol. 329, 112 [0127]
- WINTER et al. *Nature*, 1991, vol. 349, 293-299 [0130]
- LOBUGLIO et al. *Proc. Nat. Acad. Sci. USA*, 1989, vol. 86, 4220-4224 [0130]
- SHAW et al. *J Immunol.*, 1987, vol. 138, 4534-4538 [0130]
- BROWN et al. *Cancer Res.*, 1987, vol. 47, 3577-3583 [0130]
- RIECHMANN et al. *Nature*, 1988, vol. 332, 323-327 [0130]
- VERHOEYEN et al. *Science*, 1988, vol. 239, 1534-1536 [0130]
- JONES et al. *Nature*, 1986, vol. 321, 522-525 [0130]
- DAUGHERTY et al. *Nucl. Acids Res.*, 1991, vol. 19, 2471-2476 [0130]
- WINTER et al. *Annu. Rev. Immunol.*, 1994, vol. 12, 433-455 [0133]
- MCCAFFERTY et al. *Nature*, 1990, vol. 348, 552-553 [0133]
- JOHNSON, KEVIN S. ; CHISWELL, DAVID J. *Current Opinion in Structural Biology*, 1993, vol. 3, 564-571 [0133]
- CLACKSON et al. *Nature*, 1991, vol. 352, 624-628 [0133]
- MARK et al. *J. Mol. Biol.*, 1991, vol. 222, 581-597 [0133]
- GRIFFITH et al. *EMBO J.*, 1993, vol. 12, 725-734 [0133]
- MARKS et al. *Bio/Technol.*, 1992, vol. 10, 779-783 [0133]
- WATERHOUSE et al. *Nucl. Acids Res.*, 1993, vol. 21, 2265-2266 [0133]
- PEETERS et al. *Vaccine*, 2001, vol. 19, 2756 [0134]
- LONBERG, N. ; D. HUSZAR. *Int. Rev. Immunol.*, 1995, vol. 13, 65 [0134]
- POLLOCK et al. *J Immunol Methods*, 1999, vol. 231, 147 [0134]
- MORRISON et al. *Proc. Nat. Acad. Sci.*, 1984, vol. 81, 6851 [0137]
- HARLOW ; LANE. *Using Antibodies, a Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1999 [0140]
- FINDEIS et al. *Trends Biotechnol.*, 1993, vol. 11, 202 [0143]
- CHIOU et al. *Gene Therapeutics: Methods And Applications Of Direct Gene Transfer*. 1994 [0143]
- WU et al. *J. Biol. Chem.*, 1988, vol. 263, 621 [0143]
- WU et al. *J. Biol. Chem.*, 1994, vol. 269, 542 [0143]
- ZENKE et al. *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 3655 [0143]
- WU et al. *J. Biol. Chem.*, 1991, vol. 266, 338 [0143]
- JOLLY. *Cancer Gene Therapy*, 1994, vol. 1, 51 [0143]
- KIMURA. *Human Gene Therapy*, 1994, vol. 5, 845 [0143]
- CONNELLY. *Human Gene Therapy*, 1995, vol. 1, 185 [0143]
- KAPLITT. *Nature Genetics*, 1994, vol. 6, 148 [0143]
- CURIEL. *Hum. Gene Ther.*, 1992, vol. 3, 147 [0144]
- WU. *J. Biol. Chem.*, 1989, vol. 264, 16985 [0145]
- PHILIP. *Mol. Cell Biol.*, 1994, vol. 14, 2411 [0145]
- WOFFENDIN. *Proc. Natl. Acad. Sci.*, 1994, vol. 91, 1581 [0145]
- JEFFERIS ; LUND. *Chem. Immunol.*, 1997, vol. 65, 111-128 [0159]
- WRIGHT ; MORRISON. *TibTECH*, 1997, vol. 15, 26-32 [0159]
- BOYD et al. *Mol. Immunol.*, 1996, vol. 32, 1311-1318 [0159]
- WITTWE ; HOWARD. *Biochem.*, 1990, vol. 29, 4175-4180 [0159]
- WYSS ; WAGNER. *Current Opin. Biotech.*, 1996, vol. 7, 409-416 [0159]
- UMANA et al. *Mature Biotech.*, 1999, vol. 17, 176-180 [0159]
- HSE et al. *J. Biol. Chem.*, 1997, vol. 272, 9062-9070 [0162]
- MARKS et al. *Bio/Technology*, 1992, vol. 10, 779-783 [0166]
- BARBAS et al. *Proc Nat. Acad. Sci. USA*, 1994, vol. 91, 3809-3813 [0166]
- SCHIER et al. *Gene*, 1995, vol. 169, 147-155 [0166]
- YELTON et al. *J. Immunol.*, 1995, vol. 155, 1994-2004 [0166]
- JACKSON et al. *J. Immunol.*, 1995, vol. 154 (7), 3310-9 [0166]
- HAWKINS et al. *J. Mol. Biol.*, 1992, vol. 226, 889-896 [0166]
- BALINT et al. *Gene*, 1993, vol. 137 (1), 109-18 [0170]
- A model of evolutionary change in proteins - Matrices for detecting distant relationships. DAYHOFF, M.O. *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation. 1978, vol. 5, 345-358 [0189]
- HEIN J. *Unified Approach to Alignment and Phylogenies*, 1990, 626-645 [0189]
- Methods in Enzymology. Academic Press, Inc, vol. 183 [0189]
- HIGGINS, D.G. ; SHARP, P.M. *CABIOS*, 1989, vol. 5, 151-153 [0189]
- MYERS, E.W. ; MULLER W. *CABIOS*, 1988, vol. 4, 11-17 [0189]
- ROBINSON, E.D. *Comb. Theor.*, 1971, vol. 11, 105 [0189]
- SANTOU, N. ; NES, M. *Mol. Biol. Evol.*, 1987, vol. 4, 406-425 [0189]
- SNEATH, P.H.A. ; SOKAL, R.R. *Numerical Taxonomy the Principles and Practice of Numerical Taxonomy*. Freeman Press, 1973 [0189]
- WILBUR, W.J. ; LIPMAN, D.J. *Proc. Natl. Acad. Sci. USA*, 1983, vol. 80, 726-730 [0189]

- PCR: The Polymerase Chain Reaction. Birkauser Press, 1994 [0197]
- **TANAKA et al.** *FEBS Letters*, 2005, vol. 579, 2092-2096 [0204]
- **GENTLE et al.** *Bioconjugate Chem.*, 2004, vol. 15, 658-663 [0204]
- **JUNUTULA et al.** *Nature Biotechnology*, 2008, vol. 26, 925-932 [0204]
- **PETTIT et al.** *Antimicrob. Agents Chemother.*, 1998, vol. 42, 2961-2965 [0210]
- **BOGER ; JOHNSON.** *PNAS*, 1995, vol. 92, 3642-3649 [0217]
- **REMILLARD et al.** *Science*, 1975, vol. 189, 1002-1005 [0221]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1995 [0249]
- *J Mol Biol*, 2012, vol. 3, 204-219 [0276]

Patentkrav

1. Isoleret antistof, som binder specifikt til epidermal
vækstfaktorreceptor variant III (EGFRvIII), hvor antistoffet
5 omfatter:

(a) en tung kæde variabel (VH) region, der omfatter (i) en VH-
komplementaritetsbestemmende region ét (CDR1), der omfatter
sekvensen vist i SEQ ID NO: 74 75 eller 76; (ii) en VH-CDR2, der
omfatter sekvensen vist i SEQ ID NO: 77 eller 78; og (iii) en VH-
10 CDR3, der omfatter sekvensen vist i SEQ ID NO: 79; og

en let kæde variabel (VL) region, der omfatter (i) en VL-CDR1,
der omfatter sekvensen vist i SEQ ID NO: 156; (ii) en VL-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 157; og (iii) en VL-
CDR3, der omfatter sekvensen vist i SEQ ID NO: 158; eller

15 (b) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 80, 81 eller 82; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 83 eller 84; og (iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter
20 sekvensen vist i SEQ ID NO: 159; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 160; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 161; eller

(c) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 80, 81 eller 82; (ii) en VH-CDR2,
25 der omfatter sekvensen vist i SEQ ID NO: 86 eller 87; og (iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 79; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 162; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der
30 omfatter sekvensen vist i SEQ ID NO: 164; eller

(d) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 88, 89 eller 90; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 91 eller 92; og (iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og

35 en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 165; (ii) en VL-CDR2, der omfatter

sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 161; eller

(e) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 93, 94 eller 95; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 96 eller 97; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 79; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 166; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 167; eller

(f) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 93, 94 eller 95; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 96 eller 97; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 168; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 167; eller

(g) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 80, 81 eller 82; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 83 eller 84; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 169; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 161; eller

(h) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 80, 81 eller 82; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 83 eller 84; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 170; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 161; eller

(i) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 80, 81 eller 82; (ii) en VH-CDR2,

der omfatter sekvensen vist i SEQ ID NO: 98 eller 78; og iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 171; (ii) en VL-CDR2, der omfatter
5 sekvensen vist i SEQ ID NO: 172; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 167; eller
(j) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 80, 81 eller 82; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 98 eller 78; og iii)
10 en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 174; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 175; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 167; eller
15 (k) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 99, 100 eller 101; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 83 eller 84; og iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
20 sekvensen vist i SEQ ID NO: 173; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 163; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 161; eller
(l) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 74, 75 eller 76; (ii) en VH-CDR2,
25 der omfatter sekvensen vist i SEQ ID NO: 102 eller 103; og iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 104; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 176; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 172; og (iii) en VL-CDR3, der
30 omfatter sekvensen vist i SEQ ID NO: 177; eller
(m) en VH-region, der omfatter (i) en CDR1, der omfatter
sekvensen vist i SEQ ID NO: 74, 75 eller 76; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 105 eller 103; og iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 107; og
35 en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 178; (ii) en VL-CDR2, der omfatter

sekvensen vist i SEQ ID NO: 179; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 180; eller

(n) en VH-region, der omfatter (i) en CDR1, der omfatter sekvensen vist i SEQ ID NO: 74, 75 eller 76; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 102 eller 103; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 108; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 176; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 172; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 177; eller

(o) en VH-region, der omfatter (i) en CDR1, der omfatter sekvensen vist i SEQ ID NO: 74, 75 eller 76; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 106 eller 87; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 85; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 181; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 172; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 167; eller

(p) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 109, 110 eller 111; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 112 eller 113; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 114; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 182; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 183; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 184; eller

(q) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 115, 116 eller 117; (ii) en VH-CDR2, der omfatter sekvensen vist i SEQ ID NO: 118 eller 119; og (iii) en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 120; og

en VL-region, der omfatter (i) en VL-CDR1, der omfatter sekvensen vist i SEQ ID NO: 185; (ii) en VL-CDR2, der omfatter sekvensen vist i SEQ ID NO: 186; og (iii) en VL-CDR3, der omfatter sekvensen vist i SEQ ID NO: 184; eller

(r) en VH-region, der omfatter (i) en VH-CDR1, der omfatter sekvensen vist i SEQ ID NO: 121, 122 eller 123; (ii) en VH-CDR2,

der omfatter sekvensen vist i SEQ ID NO: 124 eller 125; og iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 126; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 187; (ii) en VL-CDR2, der omfatter
5 sekvensen vist i SEQ ID NO: 188; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 189; eller

(s) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 115, 116 eller 117; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 127 eller 128; og (iii)
10 en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 129; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 185; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 186; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 184; eller

(t) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 115, 116 eller 117; (ii) en VH-CDR2,
der omfatter sekvensen vist i SEQ ID NO: 130 eller 131; og (iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 120; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
20 sekvensen vist i SEQ ID NO: 190; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 191; og (iii) en VL-CDR3, der
omfatter sekvensen vist i SEQ ID NO: 184; eller

(u) en VH-region, der omfatter (i) en VH-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 132, 133 eller 134; (ii) en VH-CDR2,
25 der omfatter sekvensen vist i SEQ ID NO: 135 eller 136; og (iii)
en VH-CDR3, der omfatter sekvensen vist i SEQ ID NO: 126; og
en VL-region, der omfatter (i) en VL-CDR1, der omfatter
sekvensen vist i SEQ ID NO: 192; (ii) en VL-CDR2, der omfatter
sekvensen vist i SEQ ID NO: 193; og (iii) en VL-CDR3, der
30 omfatter sekvensen vist i SEQ ID NO: 194.

2. Isoleret antistof ifølge krav 1, hvor antistoffet
omfatter:

(a) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 9;
35 og

en VL-region, der omfatter sekvensen vist i SEQ ID NO: 10; eller
(b) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 11;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 12; eller
5 (c) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 13;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 14; eller
(d) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 15;
og
10 en VL-region, der omfatter sekvensen vist i SEQ ID NO: 16; eller
(e) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 17;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 18; eller
(f) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 19;
15 og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 20; eller
(g) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 21;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 22; eller
20 (h) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 23;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 24; eller
(i) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 25;
og
25 en VL-region, der omfatter sekvensen vist i SEQ ID NO: 26; eller
(j) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 25;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 29; eller
(k) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 27;
30 og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 28; eller

- (l) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 30;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 31; eller
(m) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 32;
5 og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 33; eller
(n) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 34;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 31; eller
10 (o) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 35;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 36; eller
(p) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 37;
og
15 en VL-region, der omfatter sekvensen vist i SEQ ID NO: 38; eller
(q) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 39;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 40; eller
(r) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 41;
20 og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 42; eller
(s) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 43;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 40; eller
25 (t) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 44;
og
en VL-region, der omfatter sekvensen vist i SEQ ID NO: 45; eller
(u) en VH-region, der omfatter sekvensen vist i SEQ ID NO: 46;
og
30 en VL-region, der omfatter sekvensen vist i SEQ ID NO: 47.

3. Isoleret antistof ifølge krav 1, hvor antistoffet er et
bispecifikt antistof, hvor det bispecifikke antistof er et

fuldlængde-antistof, der yderligere omfatter et andet antistof variabelt domæne, der er i stand til at rekruttere aktiviteten af en human immuneffektorcelle ved specifik binding til et effektorantigen placeret på den humane immuneffektorcelle.

4. Isoleret antistof ifølge krav 2, hvor antistoffet er et bispecifikt antistof, hvor det bispecifikke antistof er et fuldlængde-antistof, der yderligere omfatter et andet antistof variabelt domæne, der er i stand til at rekruttere aktiviteten af en human immuneffektorcelle ved specifik binding til et effektorantigen placeret på den humane immuneffektorcelle.

5. Isoleret antistof ifølge et hvilket som helst af kravene 3-4, hvor det bispecifikke antistof omfatter aminosyremodifikationer i positionerne 223, 225 og 228 i hængselregionen og i position 409 eller 368 (EU-nummeringssystem) i CH3-regionen af et humant IgG2 (SEQ ID NO: 290); og eventuelt yderligere omfatter en aminosyremodifikation i en eller flere af positionerne 265, 330 og 331 i det humane IgG2.

6. Antistof ifølge krav 1 eller krav 2, hvor antistoffet omfatter en acyldonor glutamin-indeholdende tag, der er indført på et specifikt sted; eller hvor antistoffet omfatter en linker.

7. Nukleinsyre, der koder for antistoffet ifølge et hvilket som helst af kravene 1-6.

8. Vektor, der omfatter nukleinsyren ifølge krav 7.

9. Værtscelle, der omfatter nukleinsyren ifølge krav 7 eller vektoren ifølge krav 8.

10. Konjugat af antistoffet ifølge et hvilket som helst af kravene 1-2 og 6, hvor antistoffet er konjugeret til et

middel, hvor midlet er valgt fra gruppen, der består af et cytotoxisk middel, et immunmodulerende middel, et billeddannelsesmiddel, et terapeutisk protein, en biopolymer og et oligonukleotid.

5

11. Farmaceutisk sammensætning, som omfatter antistoffet ifølge et hvilket som helst af kravene 1-6 eller konjugatet ifølge krav 10.

10 12. Antistof ifølge et hvilket som helst af kravene 1-6, konjugat ifølge krav 10 eller farmaceutisk sammensætning ifølge krav 11 til anvendelse som et medikament.

13. Antistof ifølge et hvilket som helst af kravene 1-6, konjugat ifølge krav 10 eller farmaceutisk sammensætning ifølge krav 11 til anvendelse til behandling af: cancer; en tilstand, der er forbundet med maligne celler, der udtrykker EGFRvIII; tumorvækst, -progression eller -metastase hos et individ, der har maligne celler, der udtrykker EGFRvIII; eller
20 en EGFRvIII-relateret cancer valgt fra gruppen, der består af glioblastoma multiforme, anaplastisk astrocytom, kæmpecelleglioblastom, gliosarkom, anaplastisk oligodendrogliom, anaplastisk ependymom, anaplastisk oligoastrocytom, choroid plexus-karcinom, anaplastisk
25 gangliogliom, pineoblastom, pineocytom, meningiom, medulloepiteliom, ependymoblastom, medulloblastom, supraentorial primitiv neuroektoderm tumor, atypisk teratoid/rhabdoid tumor, blandet gliom, hoved- og halscancer, ikke-småcellet lungecancer, brystcancer, ovariecancer, 30 prostatacancer, medulloblastom, kolorektal cancer, anal cancer, cervixcancer, nyrecancer, hudcancer, pankreascancer, levercancer, blærecancer, ventrikelcancer, thyreoideacancer, mesoteliom, uterus cancer, lymfom og leukæmi.

35 14. Fremgangsmåde til fremstilling af et antistof ifølge et hvilket som helst af kravene 1-6, som omfatter dyrkning af værtscellen ifølge krav 9 under betingelser, der resulterer i produktion af antistoffet eller det bispecifikke antistof,

og isolering af antistoffet eller det bispecifikke antistof
fra værtscellen eller kulturen.

DRAWINGS




FIG. 1A**FIG. 1B****FIG. 1C**





FIG. 2A**FIG. 2B****FIG. 2C**

FIG. 3A

LN18-EGFRvIII

**FIG. 3B**

LN18




FIG. 4A
LN229-EGFRvIII




FIG. 4B
LN229




FIG. 5




FIG. 6

LN229-EGFRvIII

