

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Begrænsning af europæisk patentskrift

(10) **DK/EP 2793579 T6**

-
- (51) Int.Cl.: **A 01 N 43/08 (2006.01)** **A 01 N 37/50 (2006.01)** **A 01 N 43/54 (2006.01)**
A 01 N 43/56 (2006.01) **A 01 N 43/653 (2006.01)** **A 01 N 43/713 (2006.01)**
A 01 N 43/76 (2006.01) **A 01 N 43/78 (2006.01)** **A 01 N 43/80 (2006.01)**
A 01 N 43/82 (2006.01) **A 01 N 43/88 (2006.01)** **A 01 N 43/90 (2006.01)**
A 01 N 47/24 (2006.01) **A 01 N 47/28 (2006.01)** **A 01 P 3/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-05-28**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om begrænsning af patentet: **2018-02-14**
- (86) Europæisk ansøgning nr.: **12795452.7**
- (86) Europæisk indleveringsdag: **2012-12-06**
- (87) Den europæiske ansøgnings publiceringsdag: **2014-10-29**
- (86) International ansøgning nr.: **EP2012074586**
- (87) Internationalt publikationsnr.: **WO2013092224**
- (30) Prioritet: **2011-12-21 EP 11195032** **2012-10-26 EP 12190109**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **BASF SE, Carl-Bosch-Strasse 38, 67056 Ludwigshafen, am Rhein, Tyskland**
- (72) Opfinder: **RHEINHEIMER, Joachim, Merziger Strasse 24, 67063 Ludwigshafen, Tyskland**
TERTERYAN, Violeta, Waldparkstrasse 34, 68163 Mannheim, Tyskland
REDLICH, Stefan, C4, 18, 68159 Mannheim, Tyskland
KREMZOV, Doris, Husarenstrasse 31a, 69121 Heidelberg, Tyskland
ROSENBAUM, Claudia, Riedroder Weg 8, 64683 Einhausen, Tyskland
ROHRER, Sebastian Georgios, Darsbergerstrasse 36, 69239 Neckarsteinach, Tyskland
GRAMMENOS, Wassilios, Alexander-Fleming-Strasse 13, 67071 Ludwigshafen, Tyskland
PILGER, Christian, Berthold-Schwarz-Strasse 41, 67063 Ludwigshafen, Tyskland
GEWEHR, Markus, Goethestrasse 21, 56288 Kastellaun, Tyskland
STAMMLER, Gerd, Korngasse 9, 69221 Dossenheim, Tyskland
MONTAG, Jurith, Brucknerstrasse 14, 67061 Ludwigshafen, Tyskland
SAUTER, Hubert, Alte Reichenbacher Strasse 113, 72270 Baiersbronn, Tyskland
RÖHL, Franz, Fußgönheimer Weg 11, 67161 Gönnsheim, Tyskland
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **ANVENDELSE AF STROBILURINTYPE FORBINDELSER TIL BEKÆMPELSE AF FYTOPATOGENE SVAMPE, DER ER RESISTENTE OVER FOR QO-HÆMMERE**
- (56) Fremdragne publikationer: **EP-A2- 0 398 692**

Fortsættes ...

WO-A1-96/36229

WO-A1-98/21174

WO-A1-99/46246

WO-A2-2007/048735

DE-A1- 19 900 571

US-A1- 2009 306 142

SCHOBERT R ET AL: "Reactions of chelated η^3 -pentadienyl iron complexes with nucleophiles", JOURNAL OF ORGANOMETALLIC CHEMISTRY, ELSEVIER-SEQUOIA S.A. LAUSANNE, CH, vol. 689, no. 3, 9 February 2004 (2004-02-09), pages 575-584, XP004487846, ISSN: 0022-328X, DOI: 10.1016/J.JORGANCHEM.2003.11.015 cited in the application

Description


The present invention relates to the use of strobilurine type compounds of formula I and the N-oxides and the salts thereof
5 for combating phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors, and to methods for combating such fungi. The invention also relates to novel compounds, processes for preparing these compounds, to compositions comprising at least
10 one such compound, to plant health applications, and to seeds coated with at least one such compound.

Qo inhibitor fungicides, often referred to as strobilurin-type fungicides (Sauter 2007: Chapter 13.2. Strobilurins and other
15 complex III inhibitors. In: Krämer, W.; Schirmer, U. (Ed.) - Modern Crop Protection Compounds. Volume 2. Wiley-VCH Verlag 457-495), are conventionally used to control a number of fungal pathogens in crops. Qo inhibitors typically work by inhibiting respiration by binding to a ubiquinol oxidation center of
20 a cytochrome bc₁ complex (electron transport complex III) in mitochondria. Said oxidation center is located on the outer side of the inner mitochondrial membrane. A prime example of the use of Qo inhibitors includes the use of, for example, strobilurins on wheat for the control of *Septoria tritici* (also known as
25 *Mycosphaerella graminicola*), which is the cause of wheat leaf blotch. Unfortunately, widespread use of such Qo inhibitors has resulted in the selection of mutant pathogens which are resistant to such Qo inhibitors (Gisi et al., Pest Manag Sci 56, 833-841, (2000). Resistance to Qo inhibitors has been detected
30 in several phytopathogenic fungi such as *Blumeria graminis*, *Mycosphaerella fijiensis*, *Pseudoperonospora cubensis* or *Venturia inaequalis*. Although several resistance mechanisms have been detected meanwhile (e.g. Jabs et al. Phytomedizin 31, 15-16 (2001); Olaya et al., Pestic Sci 54, 230-236 (1998), the major
35 part of resistance to Qo inhibitors in agricultural uses has been attributed to pathogens containing a single amino acid residue substitution G143A in the cytochrome b gene for their cytochrome bc₁ complex, the target protein of Qo inhibitors. See, for

example, Lucas, Pestic Outlook 14(6), 268-70 (2003); and Fraaije et al., Phy-topathol 95(8), 933-41 (2005). Thus, new methods and compositions are desirable for controlling pathogen induced diseases in crops comprising plants subjected to pathogens that are resistant to Qo inhibitors. Furthermore, in many cases, in particular at low application rates, the fungicidal activity of the known fungicidal strobilurin analogue compounds is unsatisfactory, especially in case that a high proportion of the fungal pathogens contain a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors. Based on this, it was also an object of the present invention to provide compounds having improved activity and/or a broader activity spectrum against phytopathogenic harmful fungi.

"Qo inhibitor," as used herein, includes any substance that is capable of diminishing and/or inhibiting respiration by binding to a ubihydroquinone oxidation center of a cytochrome bc₁ complex in mitochondria. The oxidation center is typically located on the outer side of the inner mitochondrional membrane.

From WO 2009/155095, the use of a Qi inhibitor UK2A of formula




is known for combating phytopathogenic fungi that are resistant to Qo inhibitors. Qi inhibitors typically work by inhibiting respiration by binding to a ubihydroquinone oxidation center of a cytochrome bcl complex in mitochondria, the said oxidation center being located on the inner side of the inner mitochondrional membrane.

The strobilurin-analogue compounds according to the present invention differ from those described in the abovemention publication by the specific formula I and by inhibiting

respiration by binding to a ubihydroquinone oxidation center of a cytochrome bcl complex in mitochondria which defines them as Qo inhibitors. Besides the strobilurin analogue-specific structural elements R⁴, these compounds contain two specific carbon atoms bound by a double bond wherein the groups R¹ and R² are cis-oriented or the R¹ and R² together with the abovementioned two carbon atoms linking them form a phenyl ring if R⁴ is 4-methyl-1,4-dihydro-tetrazol-5-on-1-yl.

Accordingly, the present invention relates to the use of compounds of formula I



wherein:

R¹, R²

independently of each other are hydrogen, halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl or C₃-C₆-cycloalkyl-C₁-C₄-alkyl, wherein the groups R¹ and R² are cis-oriented, or

R¹ and R² together with the two carbon atoms linking them form a phenyl ring provided that R⁴ is 4-methyl-1,4-dihydro-tetrazol-5-one-1-yl (formula R4-7), and

wherein the aliphatic moieties of R¹ and/or R² or the abovementioned phenyl ring may carry 1, 2, 3 or up to the maximum number of identical or different groups R^a which independently of one another are selected from:

R^a halogen, CN, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl and C₁-C₄-haloalkoxy;

Y

is a direct bond or a divalent group selected from $-\text{OCH}_2-$, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{C}(\text{Z})=\text{N}-\text{O}-\text{CH}_2-$, $-\text{CHZ}-\text{C}(\text{Z})=\text{N}-\text{O}-\text{CH}_2-$, $-\text{O}-\text{N}=\text{C}(\text{Z})-\text{C}(\text{Z})=\text{N}-\text{O}-\text{CH}_2-$, $-\text{C}(=\text{O})-\text{C}(\text{Z})=\text{N}-\text{O}-\text{CH}_2-$ and $-\text{C}(\text{N}=\text{O}-\text{Z})-\text{C}(\text{Z})=\text{N}-\text{O}-\text{CH}_2-$,

5 where the bond depicted on the left side of the divalent group Y is attached to R^3 , and the bond depicted on the right side is attached to the carbon atom being substituted by R^2 , and

10 Z, which may be the same or different to any other Z, is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl;

R^3

15 is phenyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl wherein the ring member atoms of the heterocyclyl include besides carbon atoms 1, 2, 3 or 4 heteroatoms selected from the group of N, O and S;

20 wherein the cyclic groups R^3 may carry 1, 2, 3, 4 or up to the maximum possible number of identical or different groups R^b which independently of one another are selected from:

25 R^b , which may be the same or different to any other R^b , is amino, halogen, hydroxyl, oxo, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyl-oxy, C_3 - C_6 -alkynyloxy, C_1 - C_6 -alkoxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkenyloxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkynyloxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylcarbonyloxy, phenyl, naphthyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of N, O and S as
35 ring members; and wherein the aforementioned phenyl and heterocyclyl groups R^b are attached via a direct bond, an oxygen or sulfur atom, and two radicals R^b that are bound to adjacent ring member atoms of the cyclic group R^3 may form together with

said ring member atoms a fused 5-, 6- or 7-membered saturated, partially unsaturated or aromatic cycle, which may be a carbocycle or heterocycle, wherein the ring member atoms of the heterocycle include besides carbon atoms 1, 2, 3 or 4 heteroatoms selected from the group of N, O and S, and

where the aliphatic or cyclic groups R^b for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^c :

R^c , which may be the same or different to any other R^c , is halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_6 -alkoxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkenyloxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkynyloxyimino- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxyimino-, C_2 - C_6 -alkenyloxyimino-, C_2 - C_6 -alkynyloxyimino-, C_2 - C_6 -haloalkenyloxyimino-, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, phenyl or a 5-membered saturated, partially unsaturated or aromatic heterocyclyl which, in addition to carbon atoms, contains one to three heteroatoms from the group consisting of N, O and S as ring members; wherein the aforementioned cyclic groups R^c are attached via a direct bond, an oxygen or sulfur atom, and where the aliphatic or cyclic groups R^c for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^d :

R^d , which may be the same or different to any other R^d , is halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl;

or

R^3

is $-CR^A=N-O-R^B$, wherein

R^A is amino, hydroxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_1 - C_4 -



alkoxyimino-C₁-C₄-alkyl, C₁-C₄-alkylamino, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylcarbonyloxy, phenyl, phenyl-C₁-C₄-alkyl, naphthyl or a 3-to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members; and wherein the aforementioned cyclic R^A are attached via a direct bond, an oxygen or sulfur atom;

R^B is C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₁-C₄-alkoxyimino-C₁-C₄-alkyl, C₁-C₄-alkoxycarbonyl, phenyl, phenyl-C₁-C₄-alkyl, naphthyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members; where the aliphatic or cyclic groups R^A and/or R^B for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^e:

Re, which may be the same or different to any other Re, is halogen, hydroxyl, nitro, CN, carboxyl, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₈-alkynyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy;

R⁴

is a monovalent group selected from formulae R4-1 to R4-7



wherein the jagged line defines the point of attachment, and

X is a direct bond or a divalent group CH₂, O or NH,

5 R⁵ is C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy or C₃-C₆-cycloalkyl,

R⁶ is C₁-C₄-alkyl or C₁-C₄-haloalkyl;

10 and the N-oxides and the agriculturally acceptable salts thereof, for combating phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors.

15 Furthermore, the present invention also relates to methods for combating phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors using the abovementioned compounds of formula I.


20 Certain strobilurin type compounds of formula I, wherein R⁴ is 1-methoxycarbonyl-2-methoxy-ethen-1-yl (defined as R⁴-2 herein, wherein X is O) and R¹ is CF₃, are mentioned in WO 1998/021174: (E)-2-[1-methoxy-meth-(E)-ylidene]-5-(4-phenoxy-phenoxy)-3-trifluoromethylpent-3-enoic acid methyl ester (CAS No. 207852-
25 99-1); (E)-2-[1-methoxy-meth-(E)-ylidene]-5-(3-phenoxy-phenoxy)-3-trifluoromethyl-pent-3-enoic acid methyl ester (207853-00-7); (E)-2-[1-methoxy-meth-(E)-ylidene]-4-methyl-5-(3-phenoxy-phenoxy)-3-trifluoromethyl-pent-3-enoic acid methyl ester; and (E)-5,5,5-trifluoro-2-[1-methoxy-meth-(E)-ylidene]-
30 3-methyl-4-(4-phenoxy-phenoxy-methyl)-pent-3-enoic acid methyl ester. However, it is not mentioned that the strobilurine type compounds inhibit fungal pathogens containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors.

35 Further, preparation of the compound (2E,3Z)-2-(ethylidene)-5-phenyl-3-pentenoic acid methyl ester (CAS-No. 681026-20-0) has been described in J Organomet Chem 689, 575-584 (2004).

Further, certain strobilurin type compounds, wherein R¹ and R² together with the two carbon atoms linking them form a phenyl ring and wherein R⁴ is 1-methyl-1,4-dihydro-tetrazole-5-one-4-yl (R⁴-7) are known inter alia from WO 1996/036229, WO 1999/046246 and DE 199 00 571 A1.

The compounds according to the present invention differ from those described in the abovementioned publications that R¹ and R² do not form with the two carbon atoms linking them a phenyl ring, and that (E)-2-[1-methoxy-meth-(E)-ylidene]-5-(4-phenoxy-phenoxy)-3-trifluoromethyl-pent-3-enoic acid methyl ester (207852-99-1); (E)-2-[1-methoxy-meth-(E)-ylidene]-5-(3-phenoxy-phenoxy)-3-trifluoromethyl-pent-3-enoic acid methyl ester (207853-00-7); (E)-2-[1-methoxy-meth-(E)-ylidene]-4-methyl-5-(3-phenoxy-phenoxy)-3-trifluoromethyl-pent-3-enoic acid methyl ester; (E)-5,5,5-trifluoro-2-[1-methoxy-meth-(E)-ylidene]-3-methyl-4-(4-phenoxy-phenoxy-methyl)-pent-3-enoic acid methyl ester; and (2E,3Z)-2-(ethylidene)-5-phenyl-3-pentenoic acid methyl ester (CAS-No. 681026-20-0) are excluded.

Therefore, according to a second aspect, the invention provides compounds of formula I which are represented by formula I



wherein:

R¹, R²

independently of each other are hydrogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl or C₃-C₆-cycloalkyl-C₁-C₄-alkyl, wherein the groups R¹ and R² are cis-oriented,

wherein the aliphatic moieties of R¹ and/or R² may carry 1, 2, 3 or up to the maximum number of identical or different groups R^a

which independently of one another are selected from:

R^a CN, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl and C₁-C₄-haloalkoxy;

Y

is a direct bond or a divalent group selected from -OCH₂-, -CH₂-,
 , -CH₂CH₂-, -C(Z)=N-O-CH₂-, -CHZ-C(Z)=N-O-CH₂-, -O-N=C(Z)-C(Z)=N-
 10 O-CH₂-, -C(=O)-C(Z)=N-O-CH₂ and -C(=N-O-Z)-C(Z)=N-O-CH₂-,

where the bond depicted on the left side of the divalent group
 Y is attached to R³, and the bond depicted on the right side is
 attached to the carbon atom being substituted by R², and

Z, which may be the same or different to any other Z, is C₁-C₄-
 alkyl or C₁-C₄-haloalkyl;

R³

is phenyl or a 3- to 10-membered saturated, partially
 unsaturated or aromatic mono- or bicyclic heterocyclyl wherein
 the ring member atoms of the heterocyclyl include besides carbon
 atoms 1, 2, 3 or 4 heteroatoms selected from the group of N, O
 25 and S;

wherein the cyclic groups R³ may carry 1, 2, 3, 4 or up to the
 maximum possible number of identical or different groups R^b which
 independently of one another are selected from:

R^b, which may be the same or different to any other R^b, is
 amino, halogen, hydroxyl, oxo, nitro, CN, carboxyl, C₁-C₄-alkyl,
 C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-
 C₄-haloalkoxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₂-C₆-
 35 alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₆-alkoxyimino-C₁-C₄-alkyl, C₂-
 C₆-alkenyloxyimino-C₁-C₄-alkyl, C₂-C₆-alkynyloxyimino-C₁-C₄-
 alkyl, C₁-C₄-alkylamino, C₁-C₄-alkoxy-carbonyl, C₁-C₄-
 alkylcarbonyloxy, phenyl, naphthyl or a 3- to 10-membered

saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of N, O and S as ring members; and wherein the aforementioned phenyl and heterocyclyl groups R^b are attached via a direct bond, an oxygen or sulfur atom; and

two radicals R^b that are bound to adjacent ring member atoms of the cyclic group R^3 may form together with said ring member atoms a fused 5-, 6- or 7-membered saturated, partially unsaturated or aromatic cycle, which may be a carbocycle or hetero-cycle, wherein the ring member atoms of the fused heterocycle include besides carbon atoms 1, 2, 3 or 4 heteroatoms selected from the group of N, O and S, and where the aliphatic or cyclic groups R^b for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^c :

R^c , which may be the same or different to any other R^c , is halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_6 -alkoxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkenyloxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkynyloxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxyimino-, C_2 - C_6 -alkenyloxyimino-, C_2 - C_6 -alkynyloxyimino-, C_2 - C_6 -haloalkenyloxyimino-, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, phenyl or a 5-membered saturated, partially unsaturated or aromatic heterocyclyl which, in addition to carbon atoms, contains one to three heteroatoms from the group consisting of N, O and S as ring members; wherein the aforementioned cyclic groups R^c are attached via a direct bond, an oxygen or sulfur atom and where the aliphatic or cyclic groups R^c for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^d :

R^d , which may be the same or different to any other R^d , is halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl; or

R^3

is $-CR^A=N-O-R^B$, wherein

5 R^A is amino, hydroxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_4 -alkyl, naphthyl or a 3-to 10-membered saturated, partially unsaturated or aromatic
10 mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members; and wherein the aforementioned cyclic R^A are attached via a direct bond, an oxygen or sulfur atom;


15 R^B is C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, CC_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxycarbonyl, phenyl, phenyl- C_1 - C_4 -alkyl, naphthyl or a 3- to 10-membered saturated, partially unsaturated or
20 aromatic mono- or bicyclic heterocyclyl or which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members;

where the aliphatic or cyclic groups R^A and/or R^B for their part
25 may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^e :

R^e , which may be the same or different to any other R^e , is
30 halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy;

R^4

35 is a monovalent group selected from formulae R4-1 to R4-7



wherein the jagged line defines the point of attachment, and
where

5

X is a divalent group O or NH,

R⁵ is C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy
or C₃-C₆-cycloalkyl,

10

R⁶ is C₁-C₄-alkyl or C₁-C₄-haloalkyl;

and the N-oxides and the agriculturally acceptable salts
thereof, except for (2E,3Z)-2-(ethylidene)-5-phenyl-3-pentenoic
acid methyl ester (CAS-No. 681026-20-0).

15

The abovementioned compound has been disclosed earlier, see on
page 5 herein.

20 The term "compounds I" refers to compounds of formula I.
Likewise, this terminology applies to all sub-formulae, e. g.
"compounds I.2" refers to compounds of formula 1.2 or "compounds
V" refers to compounds of formula V, etc..

25 The compounds I can be obtained by various routes in analogy to
prior art processes known (e.g. WO 1998/021174, J Organomet Chem
689, 575-584 (2004), WO 1996/036229) and, advantageously, by the
synthesis shown in the following schemes and in the experimental
part of this application.

30

A suitable method to prepare compounds I is illustrated in scheme


1. It starts with the reduction of an acetylene compound II with a reducing agent like lithium aluminium hydride preferably in the presence of a solvent. Suitable solvents are inert against the reducing agent used and preferably selected from cyclic or aliphatic ethers like diethyl ether, tetrahydrofuran (THF), 1,4-dioxane, and methyl-tert.-butyl ether (MTBE). The reaction temperature can be between -40°C and 100°C, preferably -20 to 60°C. After a reduced intermediate has been formed, a tin compound of formula III, wherein Alk defines a suitable alkyl residue like methyl, ethyl, n-propyl, or n-butyl and wherein L is a leaving group such as halogen, ethoxy and methoxy, in particular methoxy, is added. The resulting intermediate IV is a stable compound which can be isolated and purified with the usual methods (for example extraction and chromatography).

Compound IV is further reacted with compound V to yield the intermediate VI applying the usual methods for coupling aliphatic alcohols with hydroxyl compounds IV. The Mitsunobu reaction has proven especially useful.

Compound VI is then coupled with compound VII, wherein LG is a leaving group, preferably being halogen (except fluoro) or a sulfonyloxy group such as triflate, preferably in the presence of a suitable catalyst such as known transition metal catalysts, more preferably palladium catalysts, wherein the ligand may be trifuryl phosphine, triphenyl phosphine, tritolyl phosphine or bidentate phosphine ligands. The Copper compounds, such as CuI_2 can be added to improve the reaction. A wide variety of solvents is possible here, with THF, 1,4-dioxane and amides like dimethylformamide (DMF) being preferred. The reaction temperatures can be -20 to 150°C, preferably 20 to 120°C.


The resulting compounds I, wherein Y is $-\text{OCH}_2-$ and R^2 is H, can be further modified. For example, if R^4 contains an ester group, VIII can be transformed into a methyl amide by reaction with methyl amine.

Scheme 1:



Another general method to prepare compounds I is illustrated in scheme 2. The starting materials VIII are either known or can be prepared analogous to known compounds. The Wittig-Horner reaction of compounds VIII with compounds IX illustrated here (see also Tetrahedron Lett. 1988, 29, 3361) can be replaced by the Wittig reaction if this results in better yields. These reactions as well as the reaction conditions are well known. A specific problem is the E/Z-ratio in the newly formed double bond. The desired isomer is usually accompanied by some undesired isomer, which has to be removed by purification known in the art (e. g. chromatography, distillation, crystallization, etc.).

Scheme 2:




A route to compounds I, wherein R⁴ is of formula R⁴-4, is illustrated in scheme 3. The compound X can be obtained for example from compound VI by direct reaction with iodine. The iodine in compounds X may be replaced by other suitable leaving groups, for example by bromine, chlorine or triflate. The sodium atom in the salt VII can be replaced by other suitable metal atoms, for example potassium, lithium, magnesium, calcium, etc.. The coupling reaction of X and XI is performed preferably in the

presence of a transition metal catalyst being preferably copper in the presence of a nitrogen containing ligand system (see e.g.: Tetrahedron Lett 2008, 49 (26), 4196-4199; Org Lett. 2004, 6 (11), 1809-1812).

5

Scheme 3:



If individual compounds I cannot be obtained by the routes described above, they can be prepared by derivatization of other compounds I.

10

If the synthesis yields mixtures of isomers in the case of oximes, a separation is generally not necessarily required since in some cases the individual isomers can be interconverted during work-up for use or during application (e. g. under the action of light, acids or bases). Such conversions may also take place after use, e. g. in the treatment of plants in the treated plant, or in the harmful fungus to be controlled.

15

In the definitions of the variables given above, collective terms are used which are generally representative for the substituents in question. The term "C_n-C_m" indicates the number of carbon atoms possible in each case in the substituent or substituent moiety in question.

20

25

The term "halogen" refers to fluorine, chlorine, bromine and iodine.

The term "C₁-C₆-alkyl" refers to a straight-chained or branched saturated hydrocarbon group having 1 to 6 carbon atoms, e.g. methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl (isobutyl), 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-

30

35

dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl. Likewise, the term "C₁-C₄-alkyl" refers to a
 5 straight-chained or branched alkyl group having 1 to 4 carbon atoms.

The term "C₁-C₄-haloalkyl" refers to a straight-chained or branched alkyl group having 1 to 4 carbon atoms (as defined
 10 above), wherein some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as mentioned above, for example chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl,
 15 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and pentafluoroethyl, 2-fluoropropyl, 3-fluoropropyl, 2,2-difluoropropyl, 2,3-difluoropropyl, 2-
 20 chloropropyl, 3-chloropropyl, 2,3-dichloropropyl, 2-bromopropyl, 3-bromopropyl, 3,3,3-trifluoropropyl, 3,3,3-trichloropropyl, CH₂-C₂F₅, CF₂-C₂F₅, CF(CF₃)₂, 1-(fluoromethyl)-2-fluoroethyl, 1-(chloromethyl)-2-chloroethyl, 1-(bromomethyl)-2-bromoethyl, 4-fluorobutyl, 4-chlorobutyl, 4-bromobutyl or
 25 nonafluorobutyl. Likewise, the term "C₁-C₆-haloalkyl" refers to a straight-chained or branched alkyl group having 1 to 6 carbon atoms.

The term "C₁-C₆-alkoxy" refers to a straight-chain or branched
 30 alkyl group having 1 to 6 carbon atoms which is bonded via an oxygen, at any position in the alkyl group, e.g. OCH₃, OCH₂CH₃, O(CH₂)₂CH₃, 1-methylethoxy, O(CH₂)₃CH₃, 1-methyl-propoxy, 2-methylpropoxy or 1,1-dimethylethoxy, O(CH₂)₄C or O(CH₂)₅CH₃. Likewise, the term "C₁-C₄-alkoxy" refers to a straight-chain or
 35 branched alkyl group having 1 to 4 carbon atoms which is bonded via an oxygen, at any position in the alkyl group.

The term "C₁-C₄-haloalkoxy" refers to a C₁-C₄-alkoxy group as

defined above, wherein some or all of the hydrogen atoms may be replaced by halogen atoms as mentioned above, for example, OCH_2F , OCHF_2 , OCF_3 , OCH_2Cl , OCHCl_2 , OCCl_3 , chlorofluoromethoxy, dichlorofluoromethoxy, chlorodifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 2-iodoethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-2-fluoroethoxy, 2-chloro-2,2-difluoroethoxy, 2,2-dichloro-2-fluoroethoxy, 2,2,2-trichloroethoxy, OC_2F_5 , 2-fluoropropoxy, 3-fluoropropoxy, 2,2-difluoropropoxy, 2,3-difluoropropoxy, 2-chloropropoxy, 3-chloropropoxy, 2,3-dichloropropoxy, 2-bromopropoxy, 3-bromopropoxy, 3,3,3-trifluoropropoxy, 3,3,3-trichloropropoxy, $\text{OCH}_2\text{-C}_2\text{F}_5$, $\text{OCF}_2\text{-C}_2\text{F}_5$, 1-(CH_2F)-2-fluoroethoxy, 1-(CH_2Cl)-2-chloroethoxy, 1-(CH_2Br)-2-bromoethoxy, 4-fluorobutoxy, 4-chlorobutoxy, 4-bromobutoxy or nonafluorobutoxy. Likewise, the term "C₁-C₆-haloalkoxy" refers to a C₁-C₆-alkoxy group as defined above, wherein some or all of the hydrogen atoms may be replaced by halogen atoms as mentioned above.

The term "C₁-C₄-alkoxy-C₁-C₄-alkyl" refers to alkyl having 1 to 4 carbon atoms, wherein one hydrogen atom of the alkyl radical is replaced by a C₁-C₄-alkoxy group. Likewise, the term "C₁-C₆-alkoxy-C₁-C₆-alkyl" refers to alkyl having 1 to 6 carbon atoms, wherein one hydrogen atom of the alkyl radical is replaced by a C₁-C₆-alkoxy group.

The term "C₁-C₄-alkylamino" refers to an amino radical carrying one C₁-C₄-alkyl group as substituent, e.g. methylamino, ethylamino, propylamino, 1-methylethylamino, butylamino, 1-methylpropylamino, 2-methylpropylamino, 1,1-dimethylethylamino and the like. Likewise, the term "C₁-C₆-alkylamino" refers to an amino radical carrying one C₁-C₆-alkyl group as substituent.

The term "di(C₁-C₄-alkyl)amino" refers to an amino radical carrying two identical or different C₁-C₄-alkyl groups as substituents, e. g. dimethylamino, diethylamino, di-n-propylamino, diisopropylamino, N-ethyl-N-methylamino, N-(n-propyl)-N-methylamino, N-(isopropyl)-N-methylamino, N-(n-butyl)-N-methylamino, N-(2-butyl)-N-methylamino, N-(isobutyl)-

N-methylamino, and the like. Likewise, the term "di(C₁-C₆-alkyl)amino" refers to an amino radical carrying two identical or different C₁-C₆-alkyl groups as substituents.

5 The term "C₁-C₄-alkoxyimino" refers to a divalent imino radical (C₁-C₄-alkyl-O-N=) carrying one C₁-C₄-alkoxy group as substituent, e.g. methylimino, ethylimino, propylimino, 1-methylethylimino, butylimino, 1-methylpropylimino, 2-methylpropylimino, 1,1-dimeth-ylethylimino and the like.

10

The term "C₁-C₆-alkoxyimino-C₁-C₄-alkyl" refers to alkyl having 1 to 4 carbon atoms (as defined above), wherein two hydrogen atoms of one carbon atom of the alkyl radical are replaced by a divalent C₁-C₆-alkoxyimino radical (C₁-C₆-alkyl-O-N=) as defined
15 above.

The term "C₂-C₆-alkenyloxyimino-C₁-C₄-alkyl" refers to alkyl having 1 to 4 carbon atoms (as defined above), wherein two hydrogen atoms of one carbon atom of the alkyl radical are
20 replaced by a divalent C₂-C₆-alkenyloxyimino radical (C₂-C₆-alkenyl-O-N=).

The term "C₂-C₆-alkynyloxyimino-C₁-C₄-alkyl" refers to alkyl having 1 to 4 carbon atoms (as defined above), wherein two
25 hydrogen atoms of one carbon atom of the alkyl radical are replaced by a divalent C₂-C₆-alkynyloxyimino radical (C₂-C₆-alkynyl-O-N=).

The term "C₁-C₄-alkylcarbonyl" refers to a C₁-C₄-alkyl radical
30 which is attached via a carbonyl group. The term "(C₁-C₆-alkoxy)carbonyl" refers to a C₁-C₆-alkoxy radical which is attached via a carbonyl group.

The term "C₁-C₆-alkylaminocarbonyl" refers to a C₁-C₆-alkylamino
35 radical which is attached via a carbonyl group. Likewise, the term "di(C₁-C₆-alkyl)aminocarbonyl" refers to a di(C₁-C₆)alkylamino radical which is attached via a carbonyl group.

The term "C₂-C₄-alkenyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 4 carbon atoms and a double bond in any position, e.g. ethenyl, 1-propenyl, 2-propenyl (allyl), 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl. Likewise, the term "C₂-C₆-alkenyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 6 carbon atoms and a double bond in any position.

The term "C₂-C₄-alkynyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 4 carbon atoms and containing at least one triple bond, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-methyl-2-propynyl. Likewise, the term "C₂-C₆-alkynyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 6 carbon atoms and at least one triple bond.

The term "C₃-C₆-cycloalkyl" refers to monocyclic, bicyclic, saturated hydrocarbon radicals having 3 to 6 carbon ring members, such as cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

Likewise, the term "C₃-C₆-cycloalkenyl" refers to unsaturated hydrocarbon radicals having 3 to 6 carbon ring members and a double bond in any position, such as cyclopropenyl, cyclobutenyl, cyclopentenyl or cyclohexenyl.

The term "C₃-C₆-cycloalkyl-C₁-C₄-alkyl" refers to alkyl having 1 to 4 carbon atoms (as defined above), wherein one hydrogen atom of the alkyl radical is replaced by a cycloalkyl radical having 3 to 6 carbon atoms.

The term "phenyl-C₁-C₄-alkyl" refers to alkyl having 1 to 4 carbon atoms (as defined above), wherein one hydrogen atom of the alkyl radical is replaced by a phenyl radical.

Agriculturally acceptable salts of compounds I encompass

especially the salts of those cations or the acid addition salts of those acids whose cations and anions, respectively, have no adverse effect on the fungicidal action of the compounds I. Suitable cations are thus in particular the ions of the alkali metals, preferably sodium and potassium, of the alkaline earth metals, preferably calcium, magnesium and barium, of the transition metals, preferably manganese, copper, zinc and iron, and also the ammonium ion which, if desired, may carry one to four C₁-C₄-alkyl substituents and/or one phenyl or benzyl substituent, preferably diisopropylammonium, tetramethylammonium, tetrabutylammonium, trimethylbenzylammonium, furthermore phosphonium ions, sulfonium ions, preferably tri(C₁-C₄-alkyl)sulfonium, and sulfoxonium ions, preferably tri(C₁-C₄-alkyl)sulfoxonium. Anions of useful acid addition salts are primarily chloride, bromide, fluoride, hydrogensulfate, sulfate, dihydrogenphosphate, hydrogenphosphate, phosphate, nitrate, bicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate, and the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate. They can be formed by reacting a compound of formula I with an acid of the corresponding anion, preferably of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid or nitric acid.

The compounds of formula I can be present in atropisomers arising from restricted rotation about a single bond of asymmetric groups. They also form part of the subject matter of the present invention.

Depending on the substitution pattern, the compounds of formula I and their N-oxides may have one or more centers of chirality, in which case they are present as pure enantiomers or pure diastereomers or as enantiomer or diastereomer mixtures. Both, the pure enantiomers or diastereomers and their mixtures are subject matter of the present invention.

In respect of the the variables, the embodiments of the intermediates correspond to the embodiments of the compounds I.

Preference is given to those compounds I and where applicable also to compounds of all sub-formulae provided herein, e. g. formulae I.1 and 1.2, and to the intermediates such as compounds II, III, IV and V, wherein the substituents and variables (such as k, R¹, R², R³, R⁴, R⁵, R⁶, X, Y, Z, R^A, R^B, R^a, R^b, R^c, R^d and R^e) have independently of each other or more preferably in combination (any possible combination of 2 or more substituents as defined herein) the following meanings:

Preference is also given to the uses, methods, mixtures and compositions, wherein the definitions (such as phytopathogenic fungi, treatments, crops, compounds II, further active ingredients, solvents, solid carriers) have independently of each other or more preferably in combination the following meanings and even more preferably in combination (any possible combination of 2 or more definitions as provided herein) with the preferred meanings of compounds I herein:

According to one embodiment of the invention, the invention also relates to a method for combating phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors, comprising: treating the phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors or the materials, plants, the soil or seeds that are at risk of being diseased from phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors with an effective amount of at least one compound I, or a composition comprising it thereof.

The term "phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors" is to be understood that at least 10% of the fungal isolates to be controlled contain a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors, more preferably at least 30%, even more preferably at least 50%, and most preferably at least 75% of the fungi, in

particular between 90 and 100%.

It has been observed under field conditions that populations of phytopathogenic fungi apparently consisting of non-resistant strains can readily develop resistance. The compounds can be applied under such conditions, too, in order to prevent the formation of resistance and the spread of resistant strains altogether. In this regard it is useful that they have strong activity against non-resistant phytopathogenic fungi also.

According to another embodiment, the method for combating phytopathogenic fungi, comprises: a) identifying the phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors, or the materials, plants, the soil or seeds that are at risk of being diseased from phytopathogenic fungi as defined herein, and b) treating said fungi or the materials, plants, the soil or seeds with an effective amount of at least one compound I, or a composition comprising it thereof.

According to another embodiment of the invention, the invention also relates to a method for combating phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors, comprising: treating the phytopathogenic fungi whereof at least 10% contain a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors or the materials, plants, the soil or seeds that are at risk of being diseased from phytopathogenic fungi containing a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors with an effective amount of at least one compound I, or a composition comprising it thereof; more preferably at least 30%, even more preferably at least 50%, and most preferably at least 75% of the fungi contain a mutation in the mitochondrial cytochrome b gene conferring resistance to Qo inhibitors.

According to one embodiment of the use and the method for combating phytopathogenic fungi, wherein the mutation in the

mitochondrial cytochrome b gene of the phytopathogenic fungi is G143A.

According to another embodiment, the phytopathogenic fungi are
5 selected from the group consisting of basidiomycetes, ascomycetes, and oomycetes.

According to a further embodiment, the phytopathogenic fungi are
selected from the group consisting of *Alternaria alternata*,
10 *Blumeria graminis*, *Pyricularia oryzae* (also known as *Magnaporthe grisea*),
Septoria tritici (also known as *Mycosphaerella graminicola*),
Mycosphaerella fijiensis, *Venturia inaequalis*,
Pyrenophora teres, *Pyrenophora tritici-repentis* and *Plasmopara viticola*,
in particular *Septoria tritici*.

15 One embodiment of the invention relates to compounds I, wherein
 R^1 is halogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_2 - C_6 -alkenyl, C_2 - C_6 -
alkenyloxy, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl or C_3 - C_6 -cycloalkyl-
 C_1 - C_4 -alkyl, wherein the aliphatic moieties of R^1 may carry 1,
20 2, 3 or up to the maximum number of identical or different groups
 R^b which independently of one another are selected from halogen,
CN, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkyl and C_1 - C_4 -
haloalkoxy; more preferably R^1 is halogen, C_1 - C_4 -alkyl, C_1 - C_4 -
haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl or C_1 - C_4 -
25 haloalkoxy- C_1 - C_4 -alkyl; even more preferably R^1 is halogen, C_1 -
 C_4 -alkyl, C_1 - C_4 -chloroalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -
alkyl or C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl; in particular C_1 - C_4 -alkyl.


According to another embodiment, if R^4 is 1-methoxycarbonyl-2-
30 methoxy-ethen-1-yl (R^4 -2, wherein X is O), R^1 cannot be CF_3 .

According to a further embodiment, R^2 is hydrogen, halogen, C_1 -
 C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl
or C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl; more preferably R^2 is hydrogen,
35 halogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, or C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl;
even more preferably R^2 is hydrogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy;
in particular hydrogen.


According to a further embodiment, R^2 is halogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_2 - C_6 -alkenyl, C_2 - C_6 -alkenyloxy, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl or C_3 - C_6 -cycloalkyl- C_1 - C_4 -alkyl, wherein the aliphatic moieties of R^2 may carry 1, 2, 3 or up to the maximum number of identical or different groups R^a which independently of one another are selected from halogen, CN, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkyl and C_1 - C_4 -haloalkoxy.

According to a further embodiment, Z is hydrogen or methyl, in particular methyl.


According to a further embodiment, Y is a divalent group selected from $-OCH_2-$, $-CH_2-$, $-CH_2CH_2-$, $-C(CH_3)=N-O-CH_2$ $-O-N=C(CH_3)-C(CH_3)=N-O-CH_2-$ and $-C(=N-O-CH_3)-C(CH_3)=N-O-CH_2-$; preferably $-OCH_2-$, $-CH_2CH_2-$, $-C(CH_3)=N-O-CH_2$ $-O-N=C(CH_3)-C(CH_3)=N-O-CH_2-$ or $-C(=N-O-CH_3)-C(CH_3)=N-O-CH_2-$, where the bond depicted on the left side of the divalent group Y is attached to R^3 , and the bond depicted on the right side is attached to the carbon atom being substituted by R^2 ; in particular Y is $-OCH_2$, which compounds are of formula I.1:




According to a further embodiment, Y is $-CH_2-$, which compounds are of formula I.2:



According to a further embodiment, Y is $-CH_2CH_2-$, which compounds are of formula I.3:




According to a further embodiment, Y is $-\text{C}(\text{CH}_3)=\text{N}-\text{O}-\text{CH}_2-$ which compounds are of formula I.4:




5

According to a further embodiment, Y is $-\text{O}-\text{N}=\text{C}(\text{CH}_3)-\text{C}(\text{CH}_3)=\text{N}-\text{O}-\text{CH}_2-$, which compounds are of formula I.5:



10

According to a further embodiment, Y is $-\text{C}(=\text{N}-\text{O}-\text{CH}_3)-\text{C}(\text{CH}_3)=\text{N}-\text{O}-\text{CH}_2-$ which compounds are of formula I.6:



15

Particularly preferred embodiments of the invention relate to compounds I, wherein the group Y is one of the following radicals Y-1 to Y-10, where the bond depicted on the left side of the divalent group Y is attached to R^3 , and the bond depicted on the right side is attached to the carbon atom being substituted by R^2 :

20

No.	Y
Y-1	$-\text{OCH}_2-$
Y-2	$-\text{CH}_2-$
Y-3	$-\text{CH}_2\text{CH}_2-$
Y-4	$-\text{C}(\text{CH}_3)=\text{N}-\text{O}-\text{CH}_2-$
Y-5	$-\text{O}-\text{N}=\text{C}(\text{CH}_3)-\text{C}(\text{CH}_3)=\text{N}-\text{O}-\text{CH}_2-$

Y-6	$\text{-C(=N-O-CH}_3\text{)-C(CH}_3\text{)=N-O-CH}_2\text{-}$
Y-7	$\text{-CH}_2\text{-C(CH}_3\text{)=N-O-CH}_2\text{-}$
Y-8	$\text{-C(=O)-C(CH}_3\text{)=N-O-CH}_2\text{-}$
Y-9	$\text{-CH}_2\text{OCH}_2\text{-}$
Y-10	$\text{-CH=N-O-CH}_2\text{-}$

Particularly preferred embodiments of the invention relate to compounds I, wherein the group Y is $\text{-OCH}_2\text{-}$.

5 According to a further embodiment, R^3 is phenyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl wherein the ring member atoms of the heterocyclyl include besides carbon atoms 1, 2, 3 or 4 heteroatoms selected from the group of N, O and S, wherein R^3
10 may carry 1, 2, 3, 4 or up to the maximum possible number of identical or different groups R^b as defined herein; more preferably said 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl is a 5- to 6-membered heteroaryl wherein the ring member atoms of the
15 heteroaryl include besides carbon atoms 1, 2 or 3 heteroatoms selected from the group of N, O and S.

According to a further embodiment, R^3 is phenyl, wherein the phenyl may carry 1, 2, 3, 4 or up to the maximum possible number
20 of identical or different groups R^b as defined herein.

According to a further embodiment, R^3 is a 5-membered-heteroaryl, wherein the ring member atoms of the heterocyclyl include besides carbon atoms 1, 2 or 3 heteroatoms selected from the
25 group of N, O and S, wherein the heteroaryl may carry 1,2,3,4 or up to the maximum possible number of identical or different groups R^b as defined herein; more preferably said heteroaryl is pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, 1,2,4-triazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl or 1,2,4-
30 thiadiazolyl.

According to a further embodiment, R^3 is pyrazolyl, imidazolyl,

oxazolyl, isoxazolyl, thiazolyl, 1,2,4-triazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl or 1,2,4-thiadiazolyl, which is substituted by phenyl, wherein said phenyl may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^c , which may be the same or different to any other R^c , wherein R^c is halogen, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, phenyl or a 5-membered saturated, partially unsaturated or aromatic heterocyclyl which, in addition to carbon atoms, contains one to three heteroatoms from the group consisting of N, O and S as ring members; and wherein the aforementioned heterocyclyl groups R^c are attached via a direct bond, an oxygen or sulfur atom and for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^d as defined herein.

According to a further embodiment, R^3 is a 6-membered-heteroaryl, wherein the ring member atoms of the heterocyclyl include besides carbon atoms 1, 2 or 3 heteroatoms selected from the group of N, O and S, wherein the heteroaryl may carry 1, 2, 3, 4 or up to the maximum possible number of identical or different groups R^b as defined herein; more preferably said heteroaryl is pyridinyl or pyrimidinyl.

According to a further embodiment, R^3 carries 1, 2 or 3 identical or different groups R^b .

According to a further embodiment, R^b independently of one another are selected from carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylcarbonyloxy, phenyl, naphthyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members; and wherein the aforementioned phenyl and heterocyclyl groups R^b are attached via a direct bond, an oxygen or sulfur atom.

According to a further embodiment, the aliphatic or cyclic groups R^b for their part carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^c , which, may be the same or different to any other R^c , selected from halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy and C_1 - C_4 -haloalkoxy; more preferably R^b for their part carry 1, 2 or 3 identical or different groups R^c .

According to a further embodiment, R^c , which, may be the same or different to any other R^c , is halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, phenyl or a 5-membered saturated, partially unsaturated or aromatic heterocyclyl which, in addition to carbon atoms, contains one to three heteroatoms from the group consisting of N, O and S as ring members; wherein the aforementioned cyclic groups R^c are attached via a direct bond, an oxygen or sulfur atom and for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^d as defined herein.

According to a further embodiment, two radicals R^b that are bound to adjacent ring member atoms of the cyclic group R^3 form together with said ring member atoms a fused 5-, 6- or 7-membered saturated, partially unsaturated or aromatic cycle, which may be a carbocycle or heterocycle, wherein the ring member atoms of the fused heterocycle include besides carbon atoms 1, 2, 3 or 4 heteroatoms selected from the group of N, O and S.

According to a further embodiment, R^3 is $-CR^A=N-O-R^B$, wherein R^A is amino, hydroxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylcarbonyloxy, phenyl, naphthyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members; and wherein the aforementioned phenyl and

heterocyclyl groups R^A are attached via a direct bond, an oxygen or sulfur atom, where the aliphatic or cyclic groups R^A may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^e , which may be the same or different to any other R^e , is halogen, hydroxyl, nitro, CN, carboxyl, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₈-alkynyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.

According to a further embodiment, R^3 is $-CR^A=N-O-R^B$, wherein R^B is hydroxyl, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₁-C₄-alkoxyimino-C₁-C₄-alkyl, C₁-C₄-alkoxycarbonyl, phenyl, naphthyl or a 3- to 10-membered saturated, partially unsaturated or aromatic mono- or bicyclic heterocyclyl which, in addition to carbon atoms, contains one to four heteroatoms from the group consisting of O, N and S as ring members, where the aliphatic or cyclic groups R^B may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^e , which may be the same or different to any other R^e , is halogen, hydroxyl, nitro, CN, carboxyl, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₈-alkynyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.

More preferably R^A and R^B , independently of each other, are C₁-C₄-alkyl which may carry 1, 2, 3 or up to the maximum possible number of identical or different halogen; in particular R^A and R^B are methyl.

According to a further embodiment, the aliphatic or cyclic groups R^A and/or R^B for their part may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^e , more preferably they carry 0, 1 or 3 identical or different groups R^e .

According to a further embodiment, R^e , which may be the same or different to any other R^e , is halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy or C₁-C₄-haloalkoxy.

According to a further embodiment, R^4 is $-C(=NOCH_3)-CONHCH_3$, -

$C(=NOCH_3)-COOCH_3$, $-C(=CHOCH_3)-COOCH_3$, $-C(=CHOCH_3)-CONHCH_3$, $-N(OCH_3)-COOCH_3$, $-N(CH_3)-COOCH_3$ or $-N(CH_2CH_3)-COOCH_3$

According to a further embodiment, R^4 is R4-1 as defined herein,
 5 wherein X is O.

According to a further embodiment, R^4 is R4-1 as defined herein,
 wherein X is NH.

10 According to a further embodiment, R^4 is R4-2 as defined herein,
 wherein X is O.

According to a further embodiment, R^4 is R4-2 as defined herein,
 wherein X is NH.

15 According to a further embodiment, R^4 is R4-3 as defined herein,
 wherein X is O.

According to a further embodiment, R^4 is R4-3 as defined herein,
 20 wherein X is NH.

According to a further embodiment, R^4 is R4-4 as defined herein,
 wherein X is O.

25 According to a further embodiment, R^4 is R4-4 as defined herein,
 wherein X is NH.

According to a further embodiment, R^5 is C_1 - C_4 -alkyl, C_1 - C_4 -
 haloalkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -haloalkoxy, more preferably C_1 -
 30 C_2 -alkyl or C_1 - C_2 -alkoxy, even more preferably methyl, ethyl or
 methoxy.

According to a further embodiment, R^4 is R4-4 as defined herein,
 wherein X is O and wherein R^5 is methyl, ethyl or methoxy.

35 According to a further embodiment, R^4 is R4-5 as defined herein,
 wherein X is O.

According to a further embodiment, R^4 is R4-5 as defined herein, wherein X is NH.



According to a further embodiment, R^4 is R4-6 as defined herein.

5

According to a further embodiment, R^4 is R4-7 as defined herein.

Further embodiments of the invention relate to compounds I, wherein the group R^3 is one of the following radicals R3-A to R3-B, wherein # indicates the point of attachment to the linker moiety Y:

10





















Line	R^3
R3-A	
R3-B	

Particularly preferred embodiments of the invention relate to compounds I, wherein the group R^3 is one of the following radicals R3-1 to R3-195, wherein # indicates the point of attachment to the linker moiety Y:

15

Table A.

Line	R ³	Line	R ³
R3-1		R3-8	
R3-2		R3-9	
R3-3		R3-10	
R3-4		R3-11	
R3-5		R3-12	
R3-6		R3-13	
R3-7			

Line	R ³	Line	R ³
R3-14		R3-24	
R3-15		R3-25	
R3-16		R3-26	
R3-17		R3-27	
R3-18		R3-28	
R3-19		R3-29	
R3-20		R3-30	
R3-21		R3-31	
R3-22		R3-32	
R3-23		R3-33	

Line	R ³
R3-34	
R3-35	
R3-36	
R3-37	
R3-38	
R3-39	
R3-40	
R3-41	
R3-42	











Line	R ³
R3-43	
R3-44	
R3-45	
R3-46	
R3-47	
R3-48	
R3-49	
R3-50	









Line	R ³
R3-51	
R3-52	
R3-53	
R3-54	
R3-55	
R3-56	
R3-57	
R3-58	
R3-59	
R3-60	

Line	R ³
R3-61	
R3-62	
R3-63	
R3-64	
R3-65	
R3-66	
R3-67	
R3-68	
R3-69	
R3-70	
R3-71	

Line	R ³	Line	R ³
R3-72		R3-82	
R3-73		R3-83	
R3-74		R3-84	
R3-75		R3-85	
R3-76		R3-86	
R3-77		R3-87	
R3-78		R3-88	
R3-79		R3-89	
R3-80		R3-90	
R3-81			

Line	R ³	Line	R ³
R3-91		R3-101	
R3-92		R3-102	
R3-93		R3-103	
R3-94		R3-104	
R3-95		R3-105	
R3-96		R3-106	
R3-97		R3-107	
R3-98		R3-108	
R3-99		R3-109	
R3-100		R3-110	
		R3-111	

Line	R ³
R3-112	
R3-113	
R3-114	
R3-115	
R3-116	
R3-117	
R3-118	
R3-119	
R3-120	
R3-121	

Line	R ³
R3-122	
R3-123	
R3-124	
R3-125	
R3-126	
R3-127	
R3-128	
R3-129	

Line	R ³	Line	R ³
R3-130		R3-139	
R3-131		R3-140	
R3-132		R3-141	
R3-133		R3-142	
R3-134		R3-143	
R3-135		R3-144	
R3-136		R3-145	
R3-137		R3-146	
R3-138		R3-147	
		R3-148	

Line	R ³	Line	R ³
R3-149		R3-159	
R3-150		R3-160	
R3-151		R3-161	
R3-152		R3-162	
R3-153		R3-163	
R3-154		R3-164	
R3-155		R3-165	
R3-156		R3-166	
R3-157			
R3-158			

Line	R ³	Line	R ³
R3-167		R3-175	
R3-168		R3-176	
R3-169		R3-177	
R3-170		R3-178	
R3-171		R3-179	
R3-172		R3-180	
R3-173		R3-181	
R3-174		R3-182	
		R3-183	

Line	R ³	Line	R ³
R3-184		R3-194	
R3-185		R3-195	
R3-186			
R3-187			
R3-188			
R3-189			
R3-190			
R3-191			
R3-192			
R3-193			

wherein # indicates the point of attachment to the linker moiety Y.

5

Preferred embodiments of the invention relate to compounds I, wherein the group R³ is R3-A, in particular R3-1 (1-(4-chlorophenyl)-pyrazol-3-yl).

The compounds I and the compositions according to the invention, respectively, are suitable as fungicides. They are distinguished by an outstanding effectiveness against a broad spectrum of
5 phytopathogenic fungi, including soil-borne fungi, which derive especially from the classes of the Plasmodiophoromycetes, Peronosporomycetes (syn. Oomycetes), Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes (syn. Fungi imperfecti). Some are systemically effective and
10 they can be used in crop protection as foliar fungicides, fungicides for seed dressing and soil fungicides. Moreover, they are suitable for controlling harmful fungi, which inter alia occur in wood or roots of plants.

15 The compounds I and the compositions according to the invention are particularly important in the control of a multitude of phytopathogenic fungi on various cultivated plants, such as cereals, e. g. wheat, rye, barley, triticale, oats or rice; beet, e. g. sugar beet or fodder beet; fruits, such as pomes,
20 stone fruits or soft fruits, e. g. apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries, blackberries or gooseberries; leguminous plants, such as lentils, peas, alfalfa or soybeans; oil plants, such as rape, mustard, olives, sunflowers, coconut, cocoa beans, castor oil plants, oil palms,
25 ground nuts or soybeans; cucurbits, such as squashes, cucumber or melons; fiber plants, such as cotton, flax, hemp or jute; citrus fruit, such as oranges, lemons, grape-fruits or mandarins; vegetables, such as spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, cucurbits or
30 paprika; lauraceous plants, such as avocados, cinnamon or camphor; energy and raw material plants, such as corn, soybean, rape, sugar cane or oil palm; corn; tobacco; nuts; coffee; tea; bananas; vines (table grapes and grape juice grape vines); hop; turf; sweet leaf (also called Stevia); natural rubber plants or
35 ornamental and forestry plants, such as flowers, shrubs, broad-leaved trees or evergreens, e. g. conifers; and on the plant propagation material, such as seeds, and the crop material of these plants.

Preferably, compounds I and compositions thereof, respectively are used for controlling a multitude of fungi on field crops, such as potatoes sugar beets, tobacco, wheat, rye, barley, oats, rice, corn, cotton, soybeans, rape, legumes, sunflowers, coffee or sugar cane; fruits; vines; ornamentals; or vegetables, such as cucumbers, tomatoes, beans or squashes.

The term "plant propagation material" is to be understood to denote all the generative parts of the plant such as seeds and vegetative plant material such as cuttings and tubers (e. g. potatoes), which can be used for the multiplication of the plant. This includes seeds, roots, fruits, tubers, bulbs, rhizomes, shoots, sprouts and other parts of plants, including seedlings and young plants, which are to be transplanted after germination or after emergence from soil. These young plants may also be protected before transplantation by a total or partial treatment by immersion or pouring.

Preferably, treatment of plant propagation materials with compounds I and compositions thereof, respectively, is used for controlling a multitude of fungi on cereals, such as wheat, rye, barley and oats; rice, corn, cotton and soybeans.

The term "cultivated plants" is to be understood as including plants which have been modified by breeding, mutagenesis or genetic engineering including but not limiting to agricultural biotech products on the market or in development (cf. <http://cera-gmc.org/>, see GM crop database therein). Genetically modified plants are plants, which genetic material has been so modified by the use of recombinant DNA techniques that under natural circumstances cannot readily be obtained by cross breeding, mutations or natural recombination. Typically, one or more genes have been integrated into the genetic material of a genetically modified plant in order to improve certain properties of the plant. Such genetic modifications also include but are not limited to targeted post-translational modification of protein(s), oligo- or polypeptides e. g. by glycosylation or

polymer additions such as prenylated, acetylated or farnesylated moieties or PEG moieties.

The compounds I and compositions thereof, respectively, are particularly suitable for controlling the following plant diseases:

Albugo spp. (white rust) on ornamentals, vegetables (e. g. *A. candida*) and sunflowers (e. g. *A. tragopogonis*); *Alternaria* spp. (Alternaria leaf spot) on vegetables, rape (*A. brassicola* or *brassicae*), sugar beets (*A. tenuis*), fruits, rice, soybeans, potatoes (e. g. *A. solani* or *A. alternata*), tomatoes (e. g. *A. solani* or *A. alternata*) and wheat; *Aphanomyces* spp. on sugar beets and vegetables; *Ascochyta* spp. on cereals and vegetables, e. g. *A. tritici* (anthracnose) on wheat and *A. hordei* on barley; *Bipolaris* and *Drechslera* spp. (teleomorph: *Cochliobolus* spp.), e. g. Southern leaf blight (*D. maydis*) or Northern leaf blight (*B. zeicola*) on corn, e. g. spot blotch (*B. sorokiniana*) on cereals and e.g. *B. oryzae* on rice and turfs; *Blumeria* (formerly *Erysiphe*) *graminis* (powdery mildew) on cereals (e. g. on wheat or barley); *Botrytis cinerea* (teleomorph: *Botryotinia fuckeliana*: grey mold) on fruits and berries (e. g. strawberries), vegetables (e. g. lettuce, carrots, celery and cabbages), rape, flowers, vines, forestry plants and wheat; *Bremia lactucae* (downy mildew) on lettuce; *Ceratocystis* (syn. *Ophiostoma*) spp. (rot or wilt) on broad-leaved trees and evergreens, e. g. *C. ulmi* (Dutch elm disease) on elms; *Cercospora* spp. (*Cercospora* leaf spots) on corn (e.g. Gray leaf spot: *C. zeae-maydis*), rice, sugar beets (e. g. *C. beticola*), sugar cane, vegetables, coffee, soybeans (e. g. *C. sojina* or *C. kikuchii*) and rice; *Cladosporium* spp. on tomatoes (e. g. *C. fulvum*: leaf mold) and cereals, e. g. *C. herbarum* (black ear) on wheat; *Claviceps purpurea* (ergot) on cereals; *Cochliobolus* (anamorph: *Helminthosporium* of *Bipolaris*) spp. (leaf spots) on corn (*C. carbonum*), cereals (e. g. *C. sativus*, anamorph: *B. sorokiniana*) and rice (e. g. *C. miyabeanus*, anamorph: *H. oryzae*); *Colletotrichum* (teleomorph: *Glomerella*) spp. (anthracnose) on cotton (e. g. *C. gossypii*), corn (e. g. *C. graminicola*:

Anthracnose stalk rot), soft fruits, potatoes (e. g. *C. coccodes*: black dot), beans (e. g. *C. lindemuthianum*) and soybeans (e. g. *C. truncatum* or *C. gloeosporioides*); *Corticium* spp., e. g. *C. sasakii* (sheath blight) on rice; *Corynespora cassiicola* (leaf spots) on soybeans and ornamentals; *Cycloconium* spp., e. g. *C. oleaginum* on olive trees; *Cylindrocarpon* spp. (e. g. fruit tree canker or young vine decline, teleomorph: *Nectria* or *Neonectria* spp.) on fruit trees, vines (e. g. *C. liriodendri*, teleomorph: *Neonectria liriodendri*: Black Foot Disease) and ornamentals; *Dematophora* (teleomorph: *Rosellinia*) necatrix (root and stem rot) on soybeans; *Diaporthe* spp., e. g. *D. phaseolorum* (damping off) on soybeans; *Drechslera* (syn. *Helminthosporium*, teleomorph: *Pyrenophora*) spp. on corn, cereals, such as barley (e. g. *D. teres*, net blotch) and wheat (e. g. *D. tritici-repentis*: tan spot), rice and turf; Esca (dieback, apoplexy) on vines, caused by *Formitiporia* (syn. *Phellinus*) *punctata*, *F. mediterranea*, *Phaeomoniella chlamydospora* (earlier *Phaeoacremonium chlamydosporum*), *Phaeoacremonium aleophilum* and/or *Botryosphaeria obtusa*; *Elsinoe* spp. on pome fruits (*E. pyri*), soft fruits (*E. veneta*: anthracnose) and vines (*E. ampelina*: anthracnose); *Entyloma oryzae* (leaf smut) on rice; *Epicoccum* spp. (black mold) on wheat; *Erysiphe* spp. (powdery mildew) on sugar beets (*E. betae*), vegetables (e. g. *E. pisi*), such as cucurbits (e. g. *E. cichoracearum*), cabbages, rape (e. g. *E. cruciferarum*); *Eutypa lata* (*Eutypa* canker or dieback, anamorph: *Cytosporina lata*, syn. *Libertella blepharis*) on fruit trees, vines and ornamental woods; *Exserohilum* (syn. *Helminthosporium*) spp. on corn (e. g. *E. turcicum*); *Fusarium* (teleomorph: *Gibberella*) spp. (wilt, root or stem rot) on various plants, such as *F. graminearum* or *F. culmorum* (root rot, scab or head blight) on cereals (e. g. wheat or barley), *F. oxysporum* on tomatoes, *F. solani* on soybeans and *F. verticillioides* on corn; *Gaeumannomyces graminis* (take-all) on cereals (e. g. wheat or barley) and corn; *Gibberella* spp. on cereals (e. g. *G. zeae*) and rice (e. g. *G. fujikuroi*: Bakanae disease); *Glomerella cingulata* on vines, pome fruits and other plants and *G. gossypii* on cotton; Grain-staining complex on rice; *Guignardia bidwellii* (black rot) on vines; *Gymnosporangium*

spp. on rosaceous plants and junipers, e. g. *G. sabinae* (rust) on pears; *Helminthosporium* spp. (syn. *Drechslera*, teleomorph: *Cochliobolus*) on corn, cereals and rice; *Hemileia* spp., e. g. *H. vastatrix* (coffee leaf rust) on coffee; *Isariopsis clavispora* (syn. *Cladosporium vitis*) on vines; *Macrophomina phaseolina* (syn. *phaseoli*) (root and stem rot) on soybeans and cotton; *Microdochium* (syn. *Fusarium*) *nivale* (pink snow mold) on cereals (e. g. wheat or barley); *Microsphaera diffusa* (powdery mildew) on soybeans; *Monilinia* spp., e. g. *M. laxa*, *M. fructicola* and *M. fructigena* (bloom and twig blight, brown rot) on stone fruits and other rosaceous plants; *Mycosphaerella* spp. on cereals, bananas, soft fruits and ground nuts, such as e. g. *M. graminicola* (anamorph: *Septoria tritici*, *Septoria* blotch) on wheat or *M. fijiensis* (black Sigatoka disease) on bananas; *Peronospora* spp. (downy mildew) on cabbage (e. g. *P. brassicae*), rape (e. g. *P. parasitica*), onions (e. g. *P. destructor*), tobacco (*P. tabacina*) and soybeans (e. g. *P. manshurica*); *Phakopsora pachyrhizi* and *P. meibomiae* (soybean rust) on soybeans; *Phialophora* spp. e. g. on vines (e. g. *P. tracheiphila* and *P. tetraspora*) and soybeans (e. g. *P. gregata*: stem rot); *Phoma lingam* (root and stem rot) on rape and cabbage and *P. betae* (root rot, leaf spot and damping-off) on sugar beets; *Phomopsis* spp. on suflowers, vines (e. g. *P. viticola*: can and leaf spot) and soybeans (e. g. stem rot: *P. phaseoli*, teleomorph: *Diaporthe phaseolorum*); *Physoderma maydis* (brown spots) on corn; *Phytophthora* spp. (wilt, root, leaf, fruit and stem rot) on various plants, such as paprika and cucurbits (e. g. *P. capsici*), soybeans (e. g. *P. megasperma*, syn. *P. sojae*), potatoes and tomatoes (e. g. *P. infestans*: late blight) and broad-leaved trees (e. g. *P. ramorum*: sudden oak death); *Plasmodiophora brassicae* (club root) on cabbage, rape, radish and other plants; *Plasmopara* spp., e. g. *P. viticola* (grapevine downy mildew) on vines and *P. halstedii* on sunflowers; *Podosphaera* spp. (powdery mildew) on rosaceous plants, hop, pome and soft fruits, e. g. *P. leucotricha* on apples; *Polymyxa* spp., e. g. on cereals, such as barley and wheat (*P. graminis*) and sugar beets (*P. betae*) and thereby transmitted viral diseases; *Pseudocercospora herpotrichoides* (eyespot, teleomorph: *Tapesia yallundae*) on

cereals, e. g. wheat or barley; *Pseudoperonospora* (downy mildew) on various plants, e. g. *P. cubensis* on cucurbits or *P. humili* on hop; *Pseudopezicula tracheiphila* (red fire disease or ,rotbrenner', anamorph: *Phialophora*) on vines; *Puccinia* spp. (rusts) on various plants, e. g. *P. triticina* (brown or leaf rust), *P. striiformis* (stripe or yellow rust), *P. hordei* (dwarf rust), *P. graminis* (stem or black rust) or *P. recondita* (brown or leaf rust) on cereals, such as e. g. wheat, barley or rye, *P. kuehnii* (orange rust) on sugar cane and *P. asparagi* on asparagus; *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (tan spot) on wheat or *P. teres* (net blotch) on barley; *Pyricularia* spp., e. g. *P. oryzae* (teleomorph: *Magnaporthe grisea*, rice blast) on rice and *P. grisea* on turf and cereals; *Pythium* spp. (damping-off) on turf, rice, corn, wheat, cotton, rape, sunflowers, soybeans, sugar beets, vegetables and various other plants (e. g. *P. ultimum* or *P. aphanidermatum*); *Ramularia* spp., e. g. *R. collo-cygni* (*Ramularia* leaf spots, Physiological leaf spots) on barley and *R. beticola* on sugar beets; *Rhizoctonia* spp. on cotton, rice, potatoes, turf, corn, rape, potatoes, sugar beets, vegetables and various other plants, e. g. *R. solani* (root and stem rot) on soybeans, *R. solani* (sheath blight) on rice or *R. cerealis* (*Rhizoctonia* spring blight) on wheat or barley; *Rhizopus stolonifer* (black mold, soft rot) on strawberries, carrots, cabbage, vines and tomatoes; *Rhynchosporium secalis* (scald) on barley, rye and triticale; *Sarocladium oryzae* and *S. attenuatum* (sheath rot) on rice; *Sclerotinia* spp. (stem rot or white mold) on vegetables and field crops, such as rape, sunflowers (e. g. *S. sclerotiorum*) and soybeans (e. g. *S. rolfsii* or *S. sclerotiorum*); *Septoria* spp. on various plants, e. g. *S. glycines* (brown spot) on soybeans, *S. tritici* (*Septoria* blotch) on wheat and *S.* (syn. *Stagonospora*) *nodorum* (*Stagonospora* blotch) on cereals; *Uncinula* (syn. *Erysiphe*) *necator* (powdery mildew, anamorph: *Oidium tuckeri*) on vines; *Setosphaeria* spp. (leaf blight) on corn (e. g. *S. turcicum*, syn. *Helminthosporium turcicum*) and turf; *Sphacelotheca* spp. (smut) on corn, (e. g. *S. reiliana*: head smut), sorghum und sugar cane; *Sphaerotheca fuliginea* (powdery mildew) on cucurbits; *Spongospora subterranea* (powdery scab) on

potatoes and thereby transmitted viral diseases; *Stagonospora* spp. on cereals, e. g. *S. nodorum* (*Stagonospora* blotch, teleomorph: *Leptosphaeria* [syn. *Phaeosphaeria*] *nodorum*) on wheat; *Synchytrium endobioticum* on potatoes (potato wart disease); *Taphrina* spp., e. g. *T. deformans* (leaf curl disease) on peaches and *T. pruni* (plum pocket) on plums; *Thielaviopsis* spp. (black root rot) on tobacco, pome fruits, vegetables, soybeans and cotton, e. g. *T. basicola* (syn. *Chalara elegans*); *Tilletia* spp. (common bunt or stinking smut) on cereals, such as e. g. *T. tritici* (syn. *T. caries*, wheat bunt) and *T. controversa* (dwarf bunt) on wheat; *Typhula incarnata* (grey snow mold) on barley or wheat; *Urocystis* spp., e. g. *U. occulta* (stem smut) on rye; *Uromyces* spp. (rust) on vegetables, such as beans (e. g. *U. appendiculatus*, syn. *U. phaseoli*) and sugar beets (e. g. *U. betae*); *Ustilago* spp. (loose smut) on cereals (e. g. *U. nuda* and *U. avenae*), corn (e. g. *U. maydis*: corn smut) and sugar cane; *Venturia* spp. (scab) on apples (e. g. *V. inaequalis*) and pears; and *Verticillium* spp. (wilt) on various plants, such as fruits and ornamentals, vines, soft fruits, vegetables and field crops, e. g. *V. dahliae* on strawberries, rape, potatoes and tomatoes.

The compounds I and compositions thereof, respectively, may be used for improving the health of a plant. The invention also relates to a method for improving plant health by treating a plant, its propagation material and/or the locus where the plant is growing or is to grow with an effective amount of compounds I and compositions thereof, respectively.

The term "plant health" is to be understood to denote a condition of the plant and/or its products which is determined by several indicators alone or in combination with each other such as yield (e. g. increased biomass and/or increased content of valuable ingredients), plant vigor (e. g. improved plant growth and/or greener leaves ("greening effect")), quality (e. g. improved content or composition of certain ingredients) and tolerance to abiotic and/or biotic stress. The above identified indicators for the health condition of a plant may be interdependent or may

result from each other.

The compounds of formula I can be present in different crystal modifications whose biological activity may differ. They are
5 likewise subject matter of the present invention.

The compounds I are employed as such or in form of compositions by treating the fungi or the plants, plant propagation materials, such as seeds, soil, surfaces, materials or rooms to
10 be protected from fungal attack with a fungicidally effective amount of the active substances. The application can be carried out both before and after the infection of the plants, plant propagation materials, such as seeds, soil, surfaces, materials or rooms by the fungi.

15 Plant propagation materials may be treated with compounds I as such or a composition comprising at least one compound I prophylactically either at or before planting or transplanting.

20 The invention also relates to agrochemical compositions comprising an auxiliary and at least one compound I according to the invention.

An agrochemical composition comprises a fungicidally effective
25 amount of a compound I. The term "effective amount" denotes an amount of the composition or of the compounds I, which is sufficient for controlling harmful fungi on cultivated plants or in the protection of materials and which does not result in a substantial damage to the treated plants. Such an amount can
30 vary in a broad range and is dependent on various factors, such as the fungal species to be controlled, the treated cultivated plant or material, the climatic conditions and the specific compound I used.

35 The compounds I, their N-oxides and salts can be converted into customary types of agrochemical compositions, e. g. solutions, emulsions, suspensions, dusts, powders, pastes, granules, pressings, capsules, and mixtures thereof. Examples for

composition types are suspensions (e.g. SC, OD, FS), emulsifiable concentrates (e.g. EC), emulsions (e.g. EW, EO, ES, ME), capsules (e.g. CS, ZC), pastes, pastilles, wettable powders or dusts (e.g. WP, SP, WS, DP, DS), pressings (e.g. BR, TB, DT),
5 granules (e.g. WG, SG, GR, FG, GG, MG), insecticidal articles (e.g. LN), as well as gel formulations for the treatment of plant propagation materials such as seeds (e.g. GF). These and further compositions types are defined in the "Catalogue of pesticide formulation types and international coding system",
10 Technical Monograph No. 2, 6th Ed. May 2008, CropLife International.

The compositions are prepared in a known manner, such as described by Mollet and Grubemann, Formulation technology, Wiley
15 VCH, Weinheim, 2001; or Knowles, New developments in crop protection product formulation, Agrow Reports DS243, T&F Informa, London, 2005.

Suitable auxiliaries are solvents, liquid carriers, solid
20 carriers or fillers, surfactants, dispersants, emulsifiers, wetters, adjuvants, solubilizers, penetration enhancers, protective colloids, adhesion agents, thickeners, humectants, repellents, attractants, feeding stimulants, compatibilizers, bactericides, anti-freezing agents, anti-foaming agents,
25 colorants, tackifiers and binders.

Suitable solvents and liquid carriers are water and organic solvents, such as mineral oil fractions of medium to high boiling point, e.g. kerosene, diesel oil; oils of vegetable or animal
30 origin; aliphatic, cyclic and aromatic hydrocarbons, e. g. toluene, paraffin, tetrahydronaphthalene, alkylated naphthalenes; alcohols, e.g. ethanol, propanol, butanol, benzylalcohol, cyclohexanol; glycols; DMSO; ketones, e.g. cyclohexanone; esters, e.g. lactates, carbonates, fatty acid
35 esters, gamma-butyrolactone; fatty acids; phosphonates; amines; amides, e.g. N-methylpyrrolidone, fatty acid dimethylamides; and mixtures thereof.

Suitable solid carriers or fillers are mineral earths, e.g. silicates, silica gels, talc, kaolins, limestone, lime, chalk, clays, dolomite, diatomaceous earth, bentonite, calcium sulfate, magnesium sulfate, magnesium oxide; polysaccharides, e.g. cellulose, starch; fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas; products of vegetable origin, e.g. cereal meal, tree bark meal, wood meal, nutshell meal, and mixtures thereof.

- 10 Suitable surfactants are surface-active compounds, such as anionic, cationic, nonionic and amphoteric surfactants, block polymers, polyelectrolytes, and mixtures thereof. Such surfactants can be used as emulsifier, dispersant, solubilizer, wetter, penetration enhancer, protective colloid, or adjuvant.
- 15 Examples of surfactants are listed in McCutcheon's, Vol.1: Emulsifiers & Detergents, McCutcheon's Directories, Glen Rock, USA, 2008 (International Ed. or North American Ed.).

- Suitable anionic surfactants are alkali, alkaline earth or ammonium salts of sulfonates, sulfates, phosphates, carboxylates, and mixtures thereof. Examples of sulfonates are alkylarylsulfonates, diphenylsulfonates, alpha-olefin sulfonates, lignine sulfonates, sulfonates of fatty acids and oils, sulfonates of ethoxylated alkylphenols, sulfonates of alkoxyated arylphenols, sulfonates of condensed naphthalenes, sulfonates of dodecyl- and tridecylbenzenes, sulfonates of naphthalenes and alkylnaphthalenes, sulfosuccinates or sulfosuccinamates. Examples of sulfates are sulfates of fatty acids and oils, of ethoxylated alkylphenols, of alcohols, of ethoxylated alcohols, or of fatty acid esters. Examples of phosphates are phosphate esters. Examples of carboxylates are alkyl carboxylates, and carboxylated alcohol or alkylphenol ethoxylates.

- 35 Suitable nonionic surfactants are alkoxyates, N-substituted fatty acid amides, amine oxides, esters, sugar-based surfactants, polymeric surfactants, and mixtures thereof. Examples of alkoxyates are compounds such as alcohols,

alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxyated with 1 to 50 equivalents. Ethylene oxide and/or propylene oxide may be employed for the alkoxylation, preferably ethylene oxide. Examples of N-

5 substituted fatty acid amides are fatty acid glucamides or fatty acid alkanolamides. Examples of esters are fatty acid esters, glycerol esters or monoglycerides. Examples of sugar-based surfactants are sorbitans, ethoxylated sorbitans, sucrose and glucose esters or alkylpolyglucosides. Examples of polymeric

10 surfactants are home- or copolymers of vinylpyrrolidone, vinyl-alcohols, or vinylacetate.

Suitable cationic surfactants are quaternary surfactants, for example quaternary ammonium compounds with one or two

15 hydrophobic groups, or salts of long-chain primary amines. Suitable amphoteric surfactants are alkylbetains and imidazolines. Suitable block polymers are block polymers of the A-B or A-B-A type comprising blocks of polyethylene oxide and polypropylene oxide, or of the A-B-C type comprising alkanol,

20 polyethylene oxide and polypropylene oxide. Suitable polyelectrolytes are polyacids or polybases. Examples of polyacids are alkali salts of polyacrylic acid or polyacid comb polymers. Examples of polybases are polyvinylamines or polyethyleneamines.

25 Suitable adjuvants are compounds, which have a neglectable or even no pesticidal activity themselves, and which improve the biological performance of the compound I on the target. Examples are surfactants, mineral or vegetable oils, and other

30 auxiliaries. Further examples are listed by Knowles, Adjuvants and additives, Agrow Reports DS256, T&F Informa UK, 2006, chapter 5.

Suitable thickeners are polysaccharides (e.g. xanthan gum,

35 carboxymethylcellulose), anorganic clays (organically modified or unmodified), polycarboxylates, and silicates.

Suitable bactericides are bronopol and isothiazolinone

derivatives such as alkyliothiazolinones and benzisothiazolinones.

5 Suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

Suitable anti-foaming agents are silicones, long chain alcohols, and salts of fatty acids.

10 Suitable colorants (e.g. in red, blue, or green) are pigments of low water solubility and water-soluble dyes. Examples are inorganic colorants (e.g. iron oxide, titan oxide, iron hexacyanoferrate) and organic colorants (e.g. alizarin-, azo- and phthalocyanine colorants).

15 Suitable tackifiers or binders are polyvinylpyrrolidons, polyvinylacetates, polyvinyl alcohols, polyacrylates, biological or synthetic waxes, and cellulose ethers.

20 Examples for composition types and their preparation are:

i) Water-soluble concentrates (SL, LS)

25 10-60 wt% of a compound I and 5-15 wt% wetting agent (e.g. alcohol alkoxylates) are dissolved in water and/or in a water-soluble solvent (e.g. alcohols) ad 100 wt%. The active substance dissolves upon dilution with water.

ii) Dispersible concentrates (DC)

30 5-25 wt% of a compound I and 1-10 wt% dispersant (e. g. polyvinylpyrrolidone) are dissolved in organic solvent (e.g. cyclohexanone) ad 100 wt%. Dilution with water gives a dispersion.

35 iii) Emulsifiable concentrates (EC)

15-70 wt% of a compound I and 5-10 wt% emulsifiers (e.g. calcium

dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in water-insoluble organic solvent (e.g. aromatic hydrocarbon) ad 100 wt%. Dilution with water gives an emulsion.

5 iv) Emulsions (EW, EO, ES)

5-40 wt% of a compound I and 1-10 wt% emulsifiers (e.g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in 20-40 wt% water-insoluble organic solvent (e.g. aromatic
10 hydrocarbon). This mixture is introduced into water ad 100 wt% by means of an emulsifying machine and made into a homogeneous emulsion. Dilution with water gives an emulsion.

15 v) Suspensions (SC, OD, FS)

In an agitated ball mill, 20-60 wt% of a compound I are comminuted with addition of 2-10 wt% dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate), 0.1-2 wt% thickener (e.g. xanthan gum) and water ad 100 wt% to give
20 a fine active substance suspension. Dilution with water gives a stable suspension of the active substance. For FS type composition up to 40 wt% binder (e.g. polyvinylalcohol) is added.

25 vi) Water-dispersible granules and water-soluble granules (WG, SG)

50-80 wt% of a compound I are ground finely with addition of dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate) ad 100 wt% and prepared as water-dispersible
30 or water-soluble granules by means of technical appliances (e.g. extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active substance.

35 vii) Water-dispersible powders and water-soluble powders (WP, SP, WS)

50-80 wt% of a compound I are ground in a rotor-stator mill with

addition of 1-5 wt% dispersants (e.g. sodium lignosulfonate), 1-3 wt% wetting agents (e.g. alcohol ethoxylate) and solid carrier (e.g. silica gel) ad 100 wt%. Dilution with water gives a stable dispersion or solution of the active substance.

5

viii) Gel (GW, GF)

In an agitated ball mill, 5-25 wt% of a compound I are comminuted with addition of 3-10 wt% dispersants (e.g. sodium
10 lignosulfonate), 1-5 wt% thickener (e.g. carboxymethylcellulose) and water ad 100 wt% to give a fine suspension of the active substance. Dilution with water gives a stable suspension of the active substance.

15 iv) Microemulsion (ME)

5-20 wt% of a compound I are added to 5-30 wt% organic solvent blend (e.g. fatty acid dimethylamide and cyclohexanone), 10-25 wt% surfactant blend (e.g. alcohol ethoxylate and arylphenol
20 ethoxylate), and water ad 100 %. This mixture is stirred for 1 h to produce spontaneously a thermodynamically stable microemulsion.

iv) Microcapsules (CS)

25

An oil phase comprising 5-50 wt% of a compound I, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), 2-15 wt% acrylic monomers (e.g. methylmethacrylate, methacrylic acid and a di- or triacrylate) are dispersed into an aqueous solution of
30 a protective colloid (e.g. polyvinyl alcohol). Radical polymerization initiated by a radical initiator results in the formation of poly(meth)acrylate microcapsules. Alternatively, an oil phase comprising 5-50 wt% of a compound I according to the invention, 0-40 wt% water insoluble organic solvent (e.g.
35 aromatic hydrocarbon), and an isocyanate monomer (e.g. diphenylmethene-4,4'-diisocyanatae) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). The addition of a polyamine (e.g.

hexamethylenediamine) results in the formation of polyurea microcapsules. The monomers amount to 1-10 wt%. The wt% relate to the total CS composition.

5 ix) Dustable powders (DP, DS)

1-10 wt% of a compound I are ground finely and mixed intimately with solid carrier (e.g. finely divided kaolin) ad 100 wt%.

10 x) Granules (GR, FG)

0.5-30 wt% of a compound I is ground finely and associated with solid carrier (e.g. silicate) ad 100 wt%. Granulation is achieved by extrusion, spray-drying or fluidized bed.

15

xi) Ultra-low volume liquids (UL)

1-50 wt% of a compound I are dissolved in organic solvent (e.g. aromatic hydrocarbon) ad 100 wt%.

20

The compositions types i) to xi) may optionally comprise further auxiliaries, such as 0.1-1 wt% bactericides, 5-15 wt% anti-freezing agents, 0.1-1 wt% anti-foaming agents, and 0.1-1 wt% colorants.

25

The agrochemical compositions generally comprise between 0.01 and 95%, preferably between 0.1 and 90%, and in particular between 0.5 and 75%, by weight of active substance. The active substances are employed in a purity of from 90% to 100%, preferably from 95% to 100% (according to NMR spectrum).

30

Solutions for seed treatment (LS), Suspoemulsions (SE), flowable concentrates (FS), powders for dry treatment (DS), water-dispersible powders for slurry treatment (WS), water-soluble powders (SS), emulsions (ES), emulsifiable concentrates (EC) and gels (GF) are usually employed for the purposes of treatment of plant propagation materials, particularly seeds. The compositions in question give, after two-to-tenfold dilution,

35

active substance concentrations of from 0.01 to 60% by weight, preferably from 0.1 to 40%, in the ready-to-use preparations. Application can be carried out before or during sowing. Methods for applying compound I and compositions thereof, respectively, on to plant propagation material, especially seeds include dressing, coating, pelleting, dusting, soaking and in-furrow application methods of the propagation material. Preferably, compound I or the compositions thereof, respectively, are applied on to the plant propagation material by a method such that germination is not induced, e. g. by seed dressing, pelleting, coating and dusting.

When employed in plant protection, the amounts of active substances applied are, depending on the kind of effect desired, from 0.001 to 2 kg per ha, preferably from 0.005 to 2 kg per ha, more preferably from 0.05 to 0.9 kg per ha, and in particular from 0.1 to 0.75 kg per ha.

In treatment of plant propagation materials such as seeds, e. g. by dusting, coating or drenching seed, amounts of active substance of from 0.1 to 1000 g, preferably from 1 to 1000 g, more preferably from 1 to 100 g and most preferably from 5 to 100 g, per 100 kilogram of plant propagation material (preferably seeds) are generally required.

When used in the protection of materials or stored products, the amount of active substance applied depends on the kind of application area and on the desired effect. Amounts customarily applied in the protection of materials are 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active substance per cubic meter of treated material.

Various types of oils, wetters, adjuvants, fertilizer, or micronutrients, and further pesticides (e.g. herbicides, insecticides, fungicides, growth regulators, safeners) may be added to the active substances or the compositions comprising them as premix or, if appropriate not until immediately prior to use (tank mix). These agents can be admixed with the

compositions according to the invention in a weight ratio of 1:100 to 100:1, preferably 1:10 to 10:1.

5 The user applies the composition according to the invention usually from a predosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. Usually, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray liquor or the agrochemical composition
10 according to the invention is thus obtained. Usually, 20 to 2000 liters, preferably 50 to 400 liters, of the ready-to-use spray liquor are applied per hectare of agricultural useful area.

15 According to one embodiment, individual components of the composition according to the invention such as parts of a kit or parts of a binary or ternary mixture may be mixed by the user himself in a spray tank and further auxiliaries may be added, if appropriate.

20 Mixing the compounds I or the compositions comprising them in the use form as fungicides with other fungicides results in many cases in an expansion of the fungicidal spectrum of activity being obtained or in a prevention of fungicide resistance development. Furthermore, in many cases, synergistic effects are
25 obtained.

The following list of active substances, in conjunction with which the compounds I can be used, is intended to illustrate the possible combinations but does not limit them:

30

1. A) Respiration inhibitors

o Inhibitors of complex III at Q_o site (e.g. strobilurins): azoxystrobin, coumethoxystrobin, coumoxystrobin,
35 dimoxystrobin, enestroburin, fenaminstrobin, fenoxystrobin/flufenoxystrobin, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxystrobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, trifloxystrobin, 2-[2-(2,5-

dimethyl-phenoxyethyl)-phenyl]-3-methoxy-acrylic acid methyl ester and 2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneaminoxyethyl)-phenyl)-2-methoxyimino-N-methyl-acetamide, pyribencarb, triclopyricarb/chlorodincarb, famoxadone, fenamidone;

o inhibitors of complex III at Q_i site: cyazofamid, amisulbrom, [(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[[3-(acetoxymethoxy)-4-methoxy-pyridine-2-carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutoxycarbonyloxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[[3-(1,3-benzodioxol-5-ylmethoxy)-4-methoxy-pyridine-2-carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate; (3S,6S,7R,8R)-3-[[[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]amino]-6-methyl-4,9-dioxo-8-(phenylmethyl)-1,5-dioxonan-7-yl] 2-methylpropanoate

o inhibitors of complex II (e. g. carboxamides): benodanil, benzovindiflupyr, bixafen, boscalid, carboxin, fenfuram, fluopyram, flutolanil, fluxapyroxad, furametpyr, isopyrazam, mepronil, oxycarboxin, penflufen, penthiopyrad, sedaxane, tecloftalam, thifluzamide, N-(4'-trifluoromethylthiobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-(2-(1,3,3-trimethyl-butyl)-phenyl)-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide, 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-indan-4-yl)pyrazole-4-carboxamide, 3-(trifluoromethyl)-1-methyl-N-(1,1,3-trimethyl-indan-4-yl)pyrazole-4-carboxamide, 1,3-dimethyl-N-(1,1,3-trimethylindan-4-yl)pyrazole-4-carboxamide, 3-(trifluoromethyl)-1,5-dimethyl-N-(1,1,3-trimethylindan-4-yl)-pyrazole-4-carboxamide, 3-(difluoromethyl)-1,5-dimethyl-N-(1,1,3-trimethylindan-4-yl)pyrazole-4-carboxamide, 1,3,5-trimethyl-N-(1,1,3-trimethylindan-4-yl)pyrazole-4-carboxamide;

o other respiration inhibitors (e.g. complex I, uncouplers): diflumetorim, (5,8-difluoroquinazolin-4-yl)-{2-[2-fluoro-4-(4-trifloromethylpyridin-2-yloxy)-phenyl]-ethyl}-amine; nitrophenyl derivatives: binapacryl, dinobuton, dinocap, fluazinam; ferimzone; organometal compounds: fentin salts, such as fentin-acetate, fentin chloride or fentin hydroxide; ametoctradin; and silthiofam;

2. B) Sterol biosynthesis inhibitors (SBI fungicides)

o C14 demethylase inhibitors (DMI fungicides): triazoles: azaconazole, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, oxpoconazole, paclobutrazole, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, triticonazole, uniconazole, 1-[rel-(2S;3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)-oxiranylmethyl]-5-thio-cyanato-1H-[1,2,4]triazole, 2-[rel-(2S;3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)-oxiranylmethyl]-2H-[1,2,4]triazole-3-thiol; imidazoles: imazalil, pefurazoate, prochloraz, triflumizol; pyrimidines, pyridines and piperazines: fenarimol, nuarimol, pyrifenox, triforine;

o Delta14-reductase inhibitors: aldimorph, dodemorph, dodemorph-acetate, fenpropimorph, tridemorph, fenpropidin, piperalin, spiroxamine; - Inhibitors of 3-keto reductase: fenhexamid;

3. C) Nucleic acid synthesis inhibitors

o phenylamides or acyl amino acid fungicides: benalaxyl, benalaxyl-M, kiralaxyl, metalaxyl, metalaxyl-M (mefenoxam), ofurace, oxadixyl;

o others: hymexazole, octhilinone, oxolinic acid,

bupirimate, 5-fluorocytosine, 5-fluoro-2-(p-
 tolylmethoxy)pyrimidin-4-amine, 5-fluoro-2-(4-
 fluorophenylmethoxy)pyrimidin-4-amine;

5 4. D) Inhibitors of cell division and cytoskeleton

o tubulin inhibitors, such as benzimidazoles,
 thiophanates: benomyl, carbendazim, fuberidazole,
 thiabendazole, thiophanate-methyl; triazolopyrimidines: 5-
 10 chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-
 [1,2,4]triazolo[1,5-a]pyrimidine

o other cell division inhibitors: diethofencarb,
 ethaboxam, pencycuron, fluopicolide, zoxamide, metrafenone,
 15 pyriofenone;

5. E) Inhibitors of amino acid and protein synthesis

o methionine synthesis inhibitors (anilino-
 20 pyrimidines): cyprodinil, mepanipyrim, pyrimethanil;

o protein synthesis inhibitors: blasticidin-S,
 kasugamycin, kasugamycin hydrochloride-hydrate, mildiomicin,
 streptomycin, oxytetracyclin, polyoxine, validamycin A;

25

6. F) Signal transduction inhibitors

o MAP / histidine kinase inhibitors: fluoroimid,
 iprodione, procymidone, vinclozolin, fencpiclonil, fludioxonil;
 30 o G protein inhibitors: quinoxifen;

7. G) Lipid and membrane synthesis inhibitors

o Phospholipid biosynthesis inhibitors: edifenphos,
 35 iprobenfos, pyrazophos, isoprothiolane;

o lipid peroxidation: dicloran, quintozone, tecnazene,
 tolclorfen-methyl, biphenyl, chloroneb, etridiazole;

o phospholipid biosynthesis and cell wall deposition:
dimethomorph, flumorph, man-dipropamid, pyrimorph,
benthiavalicarb, iprovalicarb, valifenalate and N-(1-(1-(4-
5 cyano-phenyl)ethanesulfonyl)-but-2-yl) carbamic acid-(4-
fluorophenyl) ester;

o compounds affecting cell membrane permeability and
fatty acids: propamocarb, propamocarb-hydrochlorid

10 o fatty acid amide hydrolase inhibitors: 1-[4-[4-[5-
(2,6-difluorophenyl)-4,5-dihydro-3-isoxazolyl]-2-thiazolyl]-1-
piperidinyl]-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-
yl]ethanone

15 8. H) Inhibitors with Multi Site Action

o inorganic active substances: Bordeaux mixture, copper
acetate, copper hydroxide, copper oxychloride, basic copper
20 sulfate, sulfur;

o thio- and dithiocarbamates: ferbam, mancozeb, maneb,
metam, metiram, propineb, thiram, zineb, ziram;

25 o organochlorine compounds (e.g. phthalimides,
sulfamides, chloronitriles): anilazine, chlorothalonil,
captafol, captan, folpet, dichlofluanid, dichlorophen,
flusulfamide, hexachlorobenzene, pentachlorophenole and its
salts, phthalide, tolylfluanid, N-(4-chloro-2-nitro-phenyl)-N-
30 ethyl-4-methyl-benzenesulfonamide;

o guanidines and others: guanidine, dodine, dodine free
base, guazatine, guazatine-acetate, iminoctadine, iminoctadine-
triacetate, iminoctadine-tris(albesilate), dithianon, 2,6-
35 dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-
1,3,5,7(2H,6H)-tetraone;

9. I) Cell wall synthesis inhibitors

o inhibitors of glucan synthesis: validamycin, polyoxin B; melanin synthesis inhibitors: pyroquilon, tricyclazole, carpropamid, dicyclomet, fenoxanil;

5

10. J) Plant defence inducers

o acibenzolar-S-methyl, probenazole, isotianil, tiadinil, prohexadione-calcium; phosphonates: fosetyl, fosetyl-
10 aluminum, phosphorous acid and its salts;

11. K) Unknown mode of action

o bronopol, chinomethionat, cyflufenamid, cymoxanil,
15 dazomet, debacarb, diclomezine, difenzoquat, difenzoquat-methylsulfate, diphenylamin, fenpyrazamine, flumetover, flusulfamide, flutianil, methasulfocarb, nitrapyrin, nitrothal-isopropyl, oxathiapiprolin, oxin-copper, proquinazid, tebufloquin, tecloftalam, triazoxide, 2-butoxy-6-iodo-3-
20 propylchromen-4-one, N-(cyclopropylmethoxyimino-(6-difluoromethoxy-2,3-difluoro-phenyl)-methyl)-2-phenyl acetamide, N'-(4-(4-chloro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine, N'-(4-(4-fluoro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine, N'-
25 (2-methyl-5-trifluoromethyl-4-(3-trimethylsilanyl-propoxy)-phenyl)-N-ethyl-N-methyl formamidine, N'-(5-difluoromethyl-2-methyl-4-(3-trimethylsilanyl-propoxy)-phenyl)-N-ethyl-N-methyl formamidine, methoxy-acetic acid 6-tert-butyl-8-fluoro-2,3-dimethyl-quinolin-4-yl ester, 3-[5-(4-methylphenyl)-2,3-
30 dimethyl-isoxazolidin-3-yl]-pyridine, 3-[5-(4-chloro-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine (pyrisoxazole), N-(6-methoxy-pyridin-3-yl) cyclopropanecarboxylic acid amide, 5-chloro-1-(4,6-dimethoxy-pyrimidin-2-yl)-2-methyl-1H-benzoimidazole, 2-(4-chloro-phenyl)-N-[4-(3,4-dimethoxy-
35 phenyl)-isoxazol-5-yl]-2-prop-2-ynyloxy-acetamide;

12. L) Antifungal biocontrol agents, plant bioactivators: *Ampelomyces quisqualis* (e.g. AQ 10® from Intrachem Bio GmbH &

Co. KG, Germany), *Aspergillus flavus* (e.g. AFLAGUARD® from Syngenta, CH), *Aureobasidium pullulans* (e.g. BOTECTOR® from bio-ferm GmbH, Germany), *Bacillus pumilus* (e.g. NRRL Accession No. B-30087 in SONATA® and BALLAD® Plus from AgraQuest Inc., USA),

5 *Bacillus subtilis* (e.g. isolate NRRL-Nr. B-21661 in RHAPSODY®, SERENADE® MAX and SERENADE® ASO from AgraQuest Inc., USA), *Bacillus subtilis* var. *amylolique-faciens* FZB24 (e.g. TAEGRO® from Novozyme Biologicals, Inc., USA), *Candida oleophila* I-82 (e.g. ASPIRE® from Ecogen Inc., USA), *Candida saitoana* (e.g.

10 BIOCURE® (in mixture with lysozyme) and BIOCOAT® from Micro Flo Company, USA (BASF SE) and Arysta), Chitosan (e.g. ARMOUR-ZEN from BotriZen Ltd., NZ), *Clonostachys rosea* f. *catenulata*, also named *Gliocladium catenulatum* (e.g. isolate J1446: PRESTOP® from Verdera, Finland), *Coniothyrium minitans* (e.g. CONTANS® from

15 Propphyta, Germany), *Cryphonectria parasitica* (e.g. *Endothia parasitica* from CNICM, France), *Cryptococcus albidus* (e.g. YIELD PLUS® from Anchor Bio-Technologies, South Africa), *Fusarium oxysporum* (e.g. BIOFOX® from S.I.A.P.A., Italy, FUSACLEAN® from Natural Plant Protection, France), *Metschnikowia fructicola*

20 (e.g. SHEMER® from Agrogreen, Israel), *Microdochium dimerum* (e.g. ANTIBOT® from Agrauxine, France), *Phlebiopsis gigantea* (e.g. ROTSOP® from Verdera, Finland), *Pseudozyma flocculosa* (e.g. SPORODEX® from Plant Products Co. Ltd., Canada), *Pythium oligandrum* DV74 (e.g. POLYVERSUM® from Remeslo SSRO,

25 Biopreparaty, Czech Rep.), *Reynoutria sachlinensis* (e.g. REGALIA® from Marrone BioInnovations, USA), *Talaromyces flavus* V117b (e.g. PROTUS® from Propphyta, Germany), *Trichoderma asperellum* SKT-1 (e.g. ECO-HOPE" from Kumiai Chemical Industry Co., Ltd., Japan), *T. atroviride* LC52 (e.g. SENTINEL® from

30 Agrimm Technologies Ltd, NZ), *T. harzianum* T-22 (e.g. PLANTSHIELD® der Firma BioWorks Inc., USA), *T. harzianum* TH 35 (e.g. ROOT PRO® from Mycontrol Ltd., Israel), *T. harzianum* T-39 (e.g. TRICHODEX® and TRICHODERMA 2000® from Mycontrol Ltd., Israel and Makhteshim Ltd., Israel), *T. harzianum* and *T. viride*

35 (e.g. TRICHOPEL from Agrimm Technologies Ltd, NZ), *T. harzianum* ICC012 and *T. viride* ICC080 (e.g. REMEDIER® WP from Isagro Ricerca, Italy), *T. polysporum* and *T. harzianum* (e.g. BINAB® from BINAB Bio-Innovation AB, Sweden), *T. stromaticum* (e.g.

TRICOVAB® from C.E.P.L.A.C., Brazil), *T. virens* GL-21 (e.g. SOILGARD® from Certis LLC, USA), *T. viride* (e.g. TRIECO® from Ecosense Labs. (India) Pvt. Ltd., Indien, BIO-CURE® F from T. Stanes & Co. Ltd., Indien), *T. viride* TV1 (e.g. *T. viride* TV1 from Agribiotec srl, Italy), *Ulocladium oudemansii* HRU3 (e.g. BOTRY-ZEN® from Botry-Zen Ltd, NZ);

13. M) Growth regulators

10 14. abscisic acid, amidochlor, ancymidol, 6-benzylaminopurine, brassinolide, butralin, chlormequat (chlormequat chloride), choline chloride, cyclanilide, daminozide, dikegulac, dimethipin, 2,6-dimethylpuridine, ethephon, flumetralin, flurprimidol, fluthiacet, forchlorfenuron, gibberellic acid, 15 inabenfide, indole-3-acetic acid, maleic hydrazide, mefluidide, mepiquat (mepiquat chloride), naphthaleneacetic acid, N-6-benzyladenine, paclobutrazol, prohexadione (prohexadione-calcium), prohydrojasmon, thidiazuron, triapenthenol, tributyl phosphorotrithioate, 2,3,5-tri-iodobenzoic acid, trinexapac-ethyl and uniconazole;

15. N) Herbicides

o acetamides: acetochlor, alachlor, butachlor, 25 dimethachlor, dimethenamid, flufenacet, mefenacet, metolachlor, metazachlor, napropamide, naproanilide, pethoxamid, pretilachlor, propachlor, thenylchlor;

o amino acid derivatives: bilanafos, glyphosate, 30 glufosinate, sulfosate;

o aryloxyphenoxypropionates: clodinafop, cyhalofop-butyl, fenoxaprop, fluazifop, haloxyfop, metamifop, propaquizafop, quizalofop, quizalofop-P-tefuryl;

35

o Bipyridyls: diquat, paraquat;

o (thio)carbamates: asulam, butylate, carbetamide,

desmedipham, dimepiperate, eptam (EPTC), esprocarb, molinate, orbencarb, phenmedipham, prosulfocarb, pyributicarb, thiobencarb, triallate;

- 5 o cyclohexanediones: butroxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxym, tralkoxydim;
- o dinitroanilines: benfluralin, ethalfluralin, oryzalin, pendimethalin, prodiamine, trifluralin;
- 10 o diphenyl ethers: acifluorfen, aclonifen, bifenox, diclofop, ethoxyfen, fomesafen, lactofen, oxyfluorfen;
- o hydroxybenzonitriles: bomoxynil, dichlobenil, ioxynil;
- 15 o imidazolinones: imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr;
- 20 o phenoxy acetic acids: clomeprop, 2,4-dichlorophenoxyacetic acid (2,4-D), 2,4-DB, dichlorprop, MCPA, MCPA-thioethyl, MCPB, Mecoprop;
- o pyrazines: chloridazon, flufenpyr-ethyl, fluthiacet, norflurazon, pyridate;
- 25 o pyridines: aminopyralid, clopyralid, diflufenican, dithiopyr, fluridone, fluroxypyr, picloram, picolinafen, thiazopyr;
- 30 o sulfonyl ureas: amidosulfuron, azimsulfuron, bensulfuron, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethoxysulfuron, flazasulfuron, flucetosulfuron, flupyr-sulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metazosulfuron, metsulfuron-methyl, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron,
- 35

trifloxysulfuron, triflusulfuron, tritosulfuron, 1-((2-chloro-6-propyl-imidazo[1,2-b]pyridazin-3-yl)sulfonyl)-3-(4,6-dimethoxy-pyrimidin-2-yl)urea;

- 5 o triazines: ametryn, atrazine, cyanazine, dimethametryn, ethiozin, hexazinone, metamiltron, metribuzin, prometryn, simazine, terbuthylazine, terbutryn, triaziflam;
- o ureas: chlorotoluron, daimuron, diuron, fluometuron, isoproturon, linuron, methabenzthiazuron, tebuthiuron;
- 10 o other acetolactate synthase inhibitors: bispyribac-sodium, cloransulam-methyl, diclosulam, florasulam, flucarbazone, flumetsulam, metosulam, ortho-sulfamuron, penoxsulam, propoxycarbazone, pyribambenz-propyl, pyribenzoxim,
- 15 pyriftalid, pyriminobac-methyl, pyrimisulfan, pyriothiobac, pyroxasulfone, pyroxsulam;
- o others: amicarbazone, aminotriazole, anilofos, beflubutamid, benazolin, bencarbazone, benfluresate, benzofenap,
- 20 bentazone, benzobicyclon, bicyclopypyrone, bromacil, bromobutide, butafenacil, butamifos, cafenstrole, carfentrazone, cinidon-ethyl, chlorthal, cinmethylin, clomazone, cumyluron, cyprosulfamide, dicamba, difenzoquat, diflufenzopyr, *Drechslera monoceras*, endothal, ethofumesate, etobenzanid, fenoxasulfone,
- 25 fentrazamide, flumiclorac-pentyl, flumioxazin, flupoxam, flurochloridone, flurtamone, indanofan, isoxaben, isoxaflutole, lenacil, propanil, propyzamide, quinclorac, quinmerac, mesotrione, methyl arsonic acid, naptalam, oxadiargyl, oxadiazon, oxaziclomefone, pentoxazone, pinoxaden, pyraclonil,
- 30 pyraflufen-ethyl, pyrasulfotole, pyrazoxyfen, pyrazolynate, quinclamine, saflufenacil, sulcotrione, sulfentrazone, terbacil, tefuryltrione, tembotrione, thiencarbazone, topramezone, (3-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)-phenoxy]-
- 35 pyridin-2-yloxy)-acetic acid ethyl ester, 6-amino-5-chloro-2-cyclopropyl-pyrimidine-4-carboxylic acid methyl ester, 6-chloro-3-(2-cyclopropyl-6-methyl-phenoxy)-pyridazin-4-ol, 4-amino-3-chloro-6-(4-chlorophenyl)-5-fluoro-pyridine-2-

carboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)-pyridine-2-carboxylic acid methyl ester, and 4-amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluoro-phenyl)-pyridine-2-carboxylic acid methyl ester.

5

16. 0) Insecticides

o organo(thio)phosphates: acephate, azamethiphos, azinphos-methyl, chlorpyrifos, chlorpyrifos-methyl,
 10 chlorfenvinphos, diazinon, dichlorvos, dicrotophos, dimethoate, disulfoton, ethion, fenitrothion, fenthion, isoxathion, malathion, methamidophos, methidathion, methyl-parathion, mevinphos, monocrotophos, oxydemeton-methyl, paraoxon, parathion, phenthoate, phosalone, phosmet, phosphamidon,
 15 phorate, phoxim, pirimiphos-methyl, profenofos, prothiofos, sulprophos, tetrachlorvinphos, terbufos, triazophos, trichlorfon;

o carbamates: alanycarb, aldicarb, bendiocarb,
 20 benfuracarb, carbaryl, carbofuran, carbosulfan, fenoxycarb, furathiocarb, methiocarb, methomyl, oxamyl, pirimicarb, propoxur, thiodicarb, triazamate;

o pyrethroids: allethrin, bifenthrin, cyfluthrin,
 25 cyhalothrin, cyphenothrin, cypermethrin, alpha-cypermethrin, beta-cypermethrin, zeta-cypermethrin, deltamethrin, esfenvalerate, etofenprox, fenpropathrin, fenvalerate, imiprothrin, lambda-cyhalothrin, permethrin, prallethrin, pyrethrin I and II, resmethrin, silafluofen, tau-fluvalinate,
 30 tefluthrin, tetramethrin, tralomethrin, transfluthrin, profluthrin, dimefluthrin;

o insect growth regulators: a) chitin synthesis inhibitors: benzoylureas: chlorfluazuron, cyramazin,
 35 diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron; buprofezin, diofenolan, hexythiazox, etoxazole, clofentazine; b) ecdysone antagonists: halofenozide, methoxyfenozide, tebufenozide,

azadirachtin; c) juvenoids: pyriproxyfen, methoprene, fenoxycarb; d) lipid biosynthesis inhibitors: spirotetramat, spiromesifen, spirotetramat;

- 5 o nicotinic receptor agonists/antagonists compounds: clothianidin, dinotefuran, flupyradifurone, imidacloprid, thiamethoxam, nitenpyram, acetamiprid, thiacloprid, 1-2-chloro-thiazol-5-ylmethyl)-2-nitrimino-3,5-dimethyl-[1,3,5]triazinane;
- 10 o GABA antagonist compounds: endosulfan, ethiprole, fipronil, vaniliprole, pyrafluprole, pyriprole, 5-amino-1-(2,6-dichloro-4-methyl-phenyl)-4-sulfinamoyl-1H-pyrazole-3-carbothioic acid amide;
- 15 o macrocyclic lactone insecticides: abamectin, emamectin, milbemectin, lepimectin, spinosad, spinetoram;
- 20 o mitochondrial electron transport inhibitor (METI) I acaricides: fenazaquin, pyridaben, tebufenpyrad, tolfenpyrad, flufenerim;
- 25 o METI II and III compounds: acequinocyl, fluacyprim, hydramethylnon;
- 30 o Uncouplers: chlorfenapyr;
- o oxidative phosphorylation inhibitors: cyhexatin, diafenthiuron, fenbutatin oxide, propargite;
- 35 o moulting disruptor compounds: cryomazine;
- o mixed function oxidase inhibitors: piperonyl butoxide;
- o sodium channel blockers: indoxacarb, metaflumizone;
- o others: benclothiaz, bifenazate, cartap, flonicamid,

pyridalyl, pymetrozine, sulfur, thiocyclam, flubendiamide, chlorantraniliprole, cyazypyr (HGW86), cyenopyrafen, flupyrazofos, cyflumetofen, amidoflumet, imicyafos, bistrifluron, and pyrifluquinazon.

5

The present invention furthermore relates to agrochemical compositions comprising a mixture of at least one compound I (component 1) and at least one further active substance useful for plant protection, e. g. selected from the groups A) to O) (component 2), in particular one further fungicide, e. g. one
10 or more fungicide from the groups A) to L), as described above, and if desired one suitable solvent or solid carrier. Those mixtures are of particular interest, since many of them at the same application rate show higher efficiencies against harmful
15 fungi. Furthermore, combating harmful fungi with a mixture of compounds I and at least one fungicide from groups A) to L), as described above, is more efficient than combating those fungi with individual compounds I or individual fungicides from groups A) to L). By applying compounds I together with at least one
20 active substance from groups A) to O) a synergistic effect can be obtained, i.e. more than simple addition of the individual effects is obtained (synergistic mixtures).

This can be obtained by applying the compounds I and at least
25 one further active substance simultaneously, either jointly (e. g. as tank-mix) or separately, or in succession, wherein the time interval between the individual applications is selected to ensure that the active substance applied first still occurs at the site of action in a sufficient amount at the time of
30 application of the further active substance(s). The order of application is not essential for working of the present invention.

In binary mixtures, i.e. compositions according to the invention
35 comprising one compound I (component 1) and one further active substance (component 2), e. g. one active substance from groups A) to O), the weight ratio of component 1 and component 2 generally depends from the properties of the active substances

used, usually it is in the range of from 1:100 to 100:1, regularly in the range of from 1:50 to 50:1, preferably in the range of from 1:20 to 20:1, more preferably in the range of from 1:10 to 10:1 and in particular in the range of from 1:3 to 3:1.

5

In ternary mixtures, i.e. compositions according to the invention comprising one compound I (component 1) and a first further active substance (component 2) and a second further active substance (component 3), e. g. two active substances from groups A) to O), the weight ratio of component 1 and component 2 depends from the properties of the active substances used, preferably it is in the range of from 1:50 to 50:1 and particularly in the range of from 1:10 to 10:1, and the weight ratio of component 1 and component 3 preferably is in the range of from 1:50 to 50:1 and particularly in the range of from 1:10 to 10:1.

Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group A) (component 2) and particularly selected from azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-methyl, orysastrobin, picoxystrobin, pyraclostrobin, trifloxystrobin; famoxadone, fenamidone; benzovindiflupyr, bixafen, boscalid, fluopyram, fluxapyroxad, isopyrazam, penflufen, penthiopyrad, sedaxane; ametoctradin, cyazofamid, fluazinam, fentin salts, such as fentin acetate.

Preference is given to mixtures comprising a compound of formula I (component 1) and at least one active substance selected from group B) (component 2) and particularly selected from cyproconazole, difenoconazole, epoxiconazole, fluquinconazole, flusilazole, flutriafol, metconazole, myclobutanil, penconazole, propiconazole, prothioconazole, triadimefon, triadimenol, tebuconazole, tetraconazole, triticonazole, prochloraz, fenarimol, triforine; dodemorph, fenpropimorph, tridemorph, fenpropidin, spiroxamine; fenhexamid.

Preference is given to mixtures comprising a compound of formula

I (component 1) and at least one active substance selected from group C) (component 2) and particularly selected from metalaxyl, (metalaxyl-M) mefenoxam, ofurace.

5 Preference is given to mixtures comprising a compound of formula I (component 1) and at least one active substance selected from group D) (component 2) and particularly selected from benomyl, carbendazim, thiophanate-methyl, ethaboxam, fluopicolide, zoxamide, metrafenone, pyriofenone.

10 Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group E) (component 2) and particularly selected from cyprodinil, mepanipyrim, pyrimethanil.

15 Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group F) (component 2) and particularly selected from iprodione, fludioxonil, vinclozolin, quinoxyfen.

20 Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group G) (component 2) and particularly selected from dimethomorph, flumorph, iprovalicarb, benthiavalicarb, 25 mandipropamid, propamocarb.

Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group H) (component 2) and particularly selected from copper 30 acetate, copper hydroxide, copper oxychloride, copper sulfate, sulfur, mancozeb, metiram, propineb, thiram, captafol, folpet, chlorothalonil, dichlofluanid, dithianon.

35 Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group I) (component 2) and particularly selected from carpropamid and fenoxanil.

Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group J) (component 2) and particularly selected from acibenzolar-S-methyl, probenazole, tiadinil, fosetyl, fosetyl-aluminium, H₃PO₃ and salts thereof.

Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group K) (component 2) and particularly selected from cymoxanil, proquinazid and *N*-methyl-2-{1-[(5-methyl-3-trifluoromethyl-1H-pyrazol-1-yl)-acetyl]-piperidin-4-yl}-*N*-[(1*R*)-1,2,3,4-tetrahydronaphthalen-1-yl]-4-thiazolecarboxamide.

Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group L) (component 2) and particularly selected from *Bacillus subtilis* strain NRRL No. B-21661, *Bacillus pumilus* strain NRRL No. B-30087 and *Ulocladium oudemansii*.

Accordingly, the present invention furthermore relates to compositions comprising one compound I (component 1) and one further active substance (component 2), which further active substance is selected from the column "Component 2" of the lines B-1 to B-369 of Table B.

A further embodiment relates to the compositions B-1 to B-369 listed in Table B, where a row of Table B corresponds in each case to a fungicidal composition comprising one of the in the present specification individualized compounds of formula I (component 1) and the respective further active substance from groups A) to O) (component 2) stated in the row in question. Preferably, the compositions described comprise the active substances in synergistically effective amounts.

Table B: Composition comprising one individualized compound I and one further active substance from groups A) to O)

Mixture	Component 1	Component 2
---------	-------------	-------------

B-1	one individualized compound I	Azoxystrobin
B-2	one individualized compound I	Coumethoxystrobin
B-3	one individualized compound I	Coumoxystrobin
B-4	one individualized compound I	Dimoxystrobin
B-5	one individualized compound I	Enestroburin
B-6	one individualized compound I	Fenaminstrobin
B-7	one individualized compound I	Fenoxystrobin/Flufenoxystrobin
B-8	one individualized compound I	Fluoxastrobin
B-9	one individualized compound I	Kresoxim-methyl
B-10	one individualized compound I	Metominostrobin
B-11	one individualized compound I	Orysastrobin
B-12	one	Picoxystrobin

	individualized compound I	
B-13	one individualized compound I	Pyraclostrobin
B-14	one individualized compound I	Pyrametostrobin
B-15	one individualized compound I	Pyraoxystrobin
B-16	one individualized compound I	Pyribencarb
B-17	one individualized compound I	Trifloxystrobin
B-18	one individualized compound I	Triclopyricarb/Chlorodincarb
B-19	one individualized compound I	2-[2-(2,5-dimethyl-phenoxy)methyl)- phenyl]-3-methoxy-acrylic acid methyl ester
B-20	one individualized compound I	2-(2-(3-(2,6-dichlorophenyl)-1-methyl- allylideneaminooxymethyl)-phenyl)-2- methoxyimino-N-methyl-acetamide
B-21	one individualized compound I	Benalaxyl
B-22	one individualized compound I	Benalaxyl-M
B-23	one individualized	Benodanil

	compound I	
B-24	one individualized compound I	Benzovindiflupyr
B-25	one individualized compound I	Bixafen
B-26	one individualized compound I	Boscalid
B-27	one individualized compound I	Carboxin
B-28	one individualized compound I	Fenfuram
B-29	one individualized compound I	Fenhexamid
B-30	one individualized compound I	Flutolanil
B-31	one individualized compound I	Fluxapyroxad
B-32	one individualized compound I	Furametpyr
B-33	one individualized compound I	Isopyrazam
B-34	one individualized compound I	Isotianil

B-35	one individualized compound I	Kiralaxyl
B-36	one individualized compound I	Mepronil
B-37	one individualized compound I	Metalaxyl
B-38	one individualized compound I	Metalaxyl-M
B-39	one individualized compound I	Ofurace
B-40	one individualized compound I	Oxadixyl
B-41	one individualized compound I	Oxycarboxin
B-42	one individualized compound I	Penflufen
B-43	one individualized compound I	Penthiopyrad
B-44	one individualized compound I	Sedaxane
B-45	one individualized compound I	Tecloftalam
B-46	one	Thifluzamide

	individualized compound I	
B-47	one individualized compound I	Tiadinil
B-48	one individualized compound I	2-Amino-4-methyl-thiazole-5-carboxylic acid anilide
B-49	one individualized compound I	N-(4'-trifluoromethylthiobiphenyl-2- yl)-3-difluoromethyl-1-methyl-1H- pyrazole-4-carboxamide
B-50	one individualized compound I	N-(2-(1,3,3-trimethyl-butyl)-phenyl)- 1,3-dimethyl-5-fluoro-1H-pyrazole-4- carboxamide
B-51	one individualized compound I	3-(difluoromethyl)-1-methyl-N-(1,1,3- trimethylindan-4-yl)pyrazole-4- carboxamide
B-52	one individualized compound I	3-(trifluoromethyl)-1-methyl-N-(1,1,3- trimethylindan-4-yl)pyrazole-4- carboxamide
B-53	one individualized compound I	1,3-dimethyl-N-(1,1,3-trimethylindan- 4-yl)pyrazole-4-carboxamide
B-54	one individualized compound I	3-(trifluoromethyl)-1,5-dimethyl-N- (1,1,3-trimethylindan-4-yl)pyrazole-4- carboxamide
B-55	one individualized compound I	3-(difluoromethyl)-1,5-dimethyl-N- (1,1,3-trimethylindan-4-yl)pyrazole-4- carboxamide
B-56	one individualized compound I	1,3,5-trimethyl-N-(1,1,3- trimethylindan-4-yl)pyrazole-4- carboxamide
B-57	one individualized	Dimethomorph

	compound I	
B-58	one individualized compound I	Flumorph
B-59	one individualized compound I	Pyrimorph
B-60	one individualized compound I	Flumetover
B-61	one individualized compound I	Fluopicolide
B-62	one individualized compound I	Fluopyram
B-63	one individualized compound I	Zoxamide
B-64	one individualized compound I	Carpropamid
B-65	one individualized compound I	Diclocymet
B-66	one individualized compound I	Mandipropamid
B-67	one individualized compound I	Oxytetracyclin
B-68	one individualized compound I	Silthiofam

B-69	one individualized compound I	N- (6-methoxy-pyridin-3-yl) cyclopropanecarboxylic acid amide
B-70	one individualized compound I	Azaconazole
B-71	one individualized compound I	Bitertanol
B-72	one individualized compound I	Bromuconazole
B-73	one individualized compound I	Cyproconazole
B-74	one individualizes compound I	Difenoconazole
B-75	one individualized compound I	Diniconazole
B-76	one individualized compound I	Diniconazole-M
B-77	one individualized compound I	Epoxiconazole
B-78	one individualized compound I	Fenbuconazole
B-79	one individualized compound I	Fluquinconazole
B-80	one	Flusilazole

	individualized compound I	
B-81	one individualized compound I	Flutriafol
B-82	one individualized compound I	Hexaconazol
B-83	one individualized compound I	Imibenconazole
B-84	one individualized compound I	Ipconazole
B-85	one individualized compound I	Metconazole
B-86	one individualized compound I	Myclobutanil
B-87	one individualized compound I	Oxpoconazol
B-88	one individualized compound I	Paclobutrazol
B-89	one individualized compound I	Penconazole
B-90	one individualized compound I	Propiconazole
B-91	one individualized	Prothioconazole

	compound I	
B-92	one individualized compound I	Simeconazole
B-93	one individualized compound I	Tebuconazole
B-94	one individualized compound I	Tetraconazole
B-95	one individualized compound I	Triadimefon
B-96	one individualized compound I	Triadimenol
B-97	one individualized compound I	Triticonazole
B-98	one individualized compound I	Uniconazole
B-99	one individualized compound I	1-[<i>rel</i> -(2 <i>S</i> ;3 <i>R</i>)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)-oxiranylmethyl]-5-thiocyanato-1 <i>H</i> -[1,2,4]triazole,
B-100	one individualized compound I	2-[<i>rel</i> -(2 <i>S</i> ;3 <i>R</i>)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)-oxiranylmethyl]-2 <i>H</i> -[1,2,4]triazole-3-thiol
B-101	one individualized compound I	Cyazofamid
B-102	one individualized compound I	Amisulbrom

B-103	one individualized compound I	Imazalil
B-104	one individualized compound I	Imazalil-sulfate
B-105	one individualized compound I	Pefurazoate
B-106	one individualized compound I	Prochloraz
B-107	one individualized compound I	Triflumizole
B-108	one individualized compound I	Benomyl
B-109	one individualized compound I	Carbendazim
B-110	one individualized compound I	Fuberidazole
B-111	one individualized compound I	Thiabendazole
B-112	one individualized compound I	Ethaboxam
B-113	one individualized compound I	Etridiazole
B-114	one	Hymexazole

	individualized compound I	
B-115	one individualized compound I	2-(4-Chloro-phenyl)-N-[4-(3,4-dimethoxy-phenyl)-isoxazol-5-yl]-2-prop-2-ynyloxy-acetamide
B-116	one individualized compound I	Fluazinam
B-117	one individualized compound I	Pyrifenox
B-118	one individualized compound I	3-[5-(4-Chloro-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine (Pyrisoxazole)
B-119	one individualized compound I	3-[5-(4-Methyl-phenyl)-2,3-dimethylisoxazolidin-3-yl]-pyridine
B-120	one individualized compound I	Bupirimate
B-121	one individualized compound I	Cyprodinil
B-122	one individualized compound I	5-Fluorocytosine
B-123	one individualized compound I	5-Fluoro-2-(p-tolylmethoxy)pyrimidin-4-amine
B-124	one individualized compound I	5-Fluoro-2-(4-fluorophenylmethoxy)-pyrimidin-4-amine
B-125	one individualized	Diflumetorim

	compound I	
B-126	one individualized compound I	(5,8-Difluoroquinazolin-4-yl)-{2-[2-fluoro-4-(4-trifluoromethylpyridin-2-yloxy)-phenyl]-ethyl}-amine
B-127	one individualized compound I	Fenarimol
B-128	one individualized compound I	Ferimzone
B-129	one individualized compound I	Mepanipyrim
B-130	one individualized compound I	Nitrapyrin
B-131	one individualized compound I	Nuarimol
B-132	one individualized compound I	Pyrimethanil
B-133	one individualized compound I	Triforine
B-134	one individualized compound I	Fenpiclonil
B-135	one individualized compound I	Fludioxonil
B-136	one individualized compound I	Aldimorph

B-137	one individualized compound I	Dodemorph
B-138	one individualized compound I	Dodemorph-acetate
B-139	one individualized compound I	Fenpropimorph
B-140	one individualized compound I	Tridemorph
B-141	one individualized compound I	Fenpropidin
B-142	one individualized compound I	Fluoroimid
B-143	one individualized compound I	Iprodione
B-144	one individualized compound I	Procymidone
B-145	one individualized compound I	Vinclozolin
B-146	one individualized compound I	Famoxadone
B-147	one individualized compound I	Fenamidone
B-148	one	Flutianil

	individualized compound I	
B-149	one individualized compound I	Octhilinone
B-150	one individualized compound I	Probenazole
B-151	one individualized compound I	Fenpyrazamine
B-152	one individualized compound I	Acibenzolar-S-methyl
B-153	one individualized compound I	Ametoctradin
B-154	one individualized compound I	Amisulbrom
B-155	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutyryloxymethoxy-4-methoxypyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-[1,5]dioxonan-7-yl]2-methylpropanoate
B-156	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl]2-methylpropanoate
B-157	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[[3-(acetoxymethoxy)-4-methoxy-pyridine-2-carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl]2-methylpropanoate
B-158	one individualized	[(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutoxycarbonyloxy-4-methoxy-

	compound I	pyridine-2-carbonyl) amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate
B-159	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[[3-(1,3-benzodioxol-5-ylmethoxy)-4-methoxy-pyridine-2-carbonyl] amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate
B-160	one individualized compound I	(3S,6S,7R,8R)-3-[[(3-hydroxy-4-methoxy-2-pyridinyl) carbonyl] amino]-6-methyl-4,9-dioxo-8-(phenylmethyl)-1,5-dioxonan-7-yl 2-methylpropanoate
B-161	one individualized compound I	Anilazin
B-162	one individualized compound I	Blasticidin-S
B-163	one individualized compound I	Captafol
B-164	one individualized compound I	Captan
B-165	one individualized compound I	Chinomothionat
B-166	one individualized compound I	Dazomet
B-167	one individualized compound I	Debacarb
B-168	one individualized	Diclomezine

	compound I	
B-169	one individualized compound I	Difenzoquat,
B-170	one individualized compound I	Difenzoquat-methylsulfate
B-171	one individualized compound I	Fenoxanil
B-172	one individualized compound I	Folpet
B-173	one individualized compound I	Oxolinsäure
B-174	one individualized compound I	Piperalin
B-175	one individualized compound I	Proquinazid
B-176	one individualized compound I	Pyroquilon
B-177	one individualized compound I	Quinoxyfen
B-178	one individualized compound I	Triazoxid
B-179	one individualized compound I	Tricyclazole

B-180	one individualized compound I	2-Butoxy-6-iodo-3-propyl-chromen-4-one
B-181	one individualized compound I	5-Chloro-1-(4,6-dimethoxy-pyrimidin-2-yl)-2-methyl-1H-benzoimidazole
B-182	one individualized compound I	5-Chloro-7-(4-methyl-piperidin-1-yl)-6-(2,4,6-trifluoro-phenyl)-[1,2,4]triazolo[1,5-a]pyrimidine
B-183	one individualized compound I	Ferbam
B-184	one individualized compound I	Mancozeb
B-185	one individualized compound I	Maneb
B-186	one individualized compound I	Metam
B-187	one individualized compound I	Methasulphocarb
B-188	one individualized compound I	Metiram
B-189	one individualized compound I	Propineb
B-190	one individualized compound I	Thiram
B-191	one	Zineb

	individualized compound I	
B-192	one individualized compound I	Ziram
B-193	one individualized compound I	Diethofencarb
B-194	one individualized compound I	Benthiavalicarb
B-195	one individualized compound I	Iprovalicarb
B-196	one individualized compound I	Propamocarb
B-197	one individualized compound I	Propamocarb hydrochlorid
B-198	one individualized compound I	Valifenalate
B-199	one individualized compound I	N- (1- (1- (4- cyanophenyl) ethanesulfonyl) -but-2-yl) carbamic acid- (4-fluorophenyl) ester
B-200	one individualized compound i	Dodine
B-201	one individualized compound I	Dodine free base
B-202	one individualized	Guazatine

	compound I	
B-203	one individualized compound I	Guazatine-acetate
B-204	one individualized compound I	Iminoctadine
B-205	one individualized compound I	Iminoctadine-triacetate
B-206	one individualized compound I	Iminoctadine-tris (albesilate)
B-207	one individualized compound I	Kasugamycin
B-208	one individualized compound I	Kasugamycin-hydrochloride-hydrate
B-209	one individualized compound I	Polyoxine
B-210	one individualized compound I	Streptomycin
B-211	one individualized compound I	Validamycin A
B-212	one individualized compound I	Binapacryl
B-213	one individualized compound I	Dicloran

B-214	one individualized compound I	Dinobuton
B-215	one individualized compound I	Dinocap
B-216	one individualized compound I	Nitrothal-isopropyl
B-217	one individualized compound I	Tecnazen
B-218	one individualized compound I	Fentin salts
B-219	one individualized compound I	Dithianon
B-220	one individualized compound I	2,6-dimethyl-1H,5H-[1,4]dithiino [2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetraone
B-221	one individualized compound I	Isoprothiolane
B-222	one individualized compound I	Edifenphos
B-223	one individualized compound I	Fosetyl, Fosetyl-aluminium
B-224	one individualized compound I	Iprobenfos
B-225	one	Phosphorous acid (H_3PO_3) and

	individualized compound I	derivatives
B-226	one individualized compound I	Pyrazophos
B-227	one individualized compound I	Tolclofos-methyl
B-228	one individualized compound I	Chlorothalonil
B-229	one individualized compound I	Dichlofluanid
B-230	one individualized compound I	Dichlorophen
B-231	one individualized compound I	Flusulfamide
B-232	one individualized compound I	Hexachlorbenzene
B-233	one individualized compound I	Pencycuron
B-234	one individualized compound I	Pentachlorophenol and salts
B-235	one individualized compound I	Phthalide
B-236	one individualized	Quintozene

	compound I	
B-237	one individualized compound I	Thiophanate Methyl
B-238	one individualized compound I	Tolylfluanid
B-239	one individualized compound I	N- (4-chloro-2-nitro-phenyl) -N-ethyl-4-methyl-benzenesulfonamide
B-240	one individualized compound I	Bordeaux mixture
B-241	one individualized compound I	Copper acetate
B-242	one individualized compound I	Copper hydroxide
B-243	one individualized compound I	Copper oxychloride
B-244	one individualized compound I	basic Copper sulfate
B-245	one individualized compound I	Sulfur
B-246	one individualized compound I	Biphenyl
B-247	one individualized compound I	Bronopol

B-248	one individualized compound I	Cyflufenamid
B-249	one individualized compound I	Cymoxanil
B-250	one individualized compound I	Diphenylamin
B-251	one individualized compound I	Metrafenone
B-252	one individualized compound I	Pyrioifenone
B-253	one individualized compound I	Mildiomycin
B-254	one individualized compound I	Oxin-copper
B-255	one individualized compound I	Oxathiapiprolin
B-256	one individualized compound I	Prohexadione calcium
B-257	one individualized compound I	Spiroxamine
B-258	one individualized compound I	Tebufloquin
B-259	one	Tolyfluanid

	individualized compound I	
B-260	one individualized compound I	N- (Cyclopropylmethoxyimino- (6- difluoromethoxy-2,3-difluoro-phenyl) - methyl)-2-phenyl acetamide
B-261	one individualized compound I	N'-(4-(4-chloro-3-trifluoromethyl- phenoxy)-2,5-dimethyl-phenyl)-N-ethyl- N-methyl formamidine
B-262	one individualized compound I	N'-(4-(4-fluoro-3-trifluoromethyl- phenoxy)-2,5-dimethyl-phenyl)-N-ethyl- N-methyl formamidine
B-263	one individualized compound I	N'-(2-methyl-5-trifluoromethyl-4-(3- trimethylsilanyl-propoxy)-phenyl)-N- ethyl-N-methyl formamidine
B-264	one individualized compound I	N'-(5-difluoromethyl-2-methyl-4-(3- trimethylsilanyl-propoxy)-phenyl)-N- ethyl-N-methyl formamidine
B-265	one individualized compound I	Methoxy-acetic acid 6-tert-butyl-8- fluoro-2,3-dimethyl-quinolin-4-yl ester
B-266	one individualized compound I	<i>Bacillus subtilis</i> NRRL No. B-21661
B-267	one individualized compound I	<i>Bacillus pumilus</i> NRRL No. B-36087
B-268	one individualized compound I	<i>Ulocladium oudemansii</i>
B-269	one individualized compound I	Carbaryl
B-270	one individualized	Carbofuran

	compound I	
B-271	one individualized compound I	Carbosulfan
B-272	one individualized compound I	Methomylthiodicarb
B-273	one individualized compound I	Bifenthrin
B-274	one individualized compound I	Cyfluthrin
B-275	one individualized compound I	Cypermethrin
B-276	one individualized compound I	alpha-Cypermethrin
B-277	one individualized compound I	zeta-Cypermethrin
B-278	one individualized compound I	Deltamethrin
B-279	one individualized compound I	Esfenvalerate
B-280	one individualized compound I	Lambda-cyhalothrin
B-281	one individualized compound I	Permethrin

B-282	one individualized compound I	Tefluthrin
B-283	one individualized compound I	Diﬂubenzuron
B-284	one individualized compound I	Flufenoxuron
B-285	one individualized compound I	Lufenuron
B-286	one individualized compound I	Teflubenzuron
B-287	one individualized compound I	Spirotetramate
B-288	one individualized compound I	Clothianidin
B-289	one individualized compound I	Dinotefuran
B-290	one individualized compound I	Imidacloprid
B-291	one individualized compound I	Thiamethoxam
B-292	one individualized compound I	Flupyradifurone
B-293	one	Acetamiprid

	individualized compound I	
B-294	one individualized compound I	Thiacloprid
B-295	one individualized compound I	Endosulfan
B-296	one individualized compound I	Fipronil
B-297	one individualized compound I	Abamectin
B-298	one individualized compound I	Emamectin
B-299	one individualized compound I	Spinosad
B-300	one individualized compound I	Spinetoram
B-301	one individualized compound I	Hydramethylnon
B-302	one individualized compound I	Chlorfenapyr
B-303	one individualized compound I	Fenbutatin oxide
B-304	one individualized	Indoxacarb

	compound I	
B-305	one individualized compound I	Metaflumizone
B-306	one individualized compound I	Flonicamid
B-307	one individualized compound I	Lubendiamide
B-308	one individualized compound I	Chlorantraniliprole
B-309	one individualized compound I	Cyazypyr (HGW86)
B-310	one individualized compound I	Cyflumetofen
B-311	one individualized compound I	Acetochlor
B-312	one individualized compound I	Dimethenamid
B-313	one individualized compound I	metolachlor
B-314	one individualized compound I	Metazachlor
B-315	one individualized compound I	Glyphosate

B-316	one individualized compound I	Glufosinate
B-317	one individualized compound I	Sulfosate
B-318	one individualized compound I	Clodinafop
B-319	one individualized compound I	Fenoxaprop
B-320	one individualized compound I	Fluazifop
B-321	one individualized compound I	Haloxifyfop
B-322	one individualized compound I	Paraquat
B-323	one individualized compound I	Phenmedipham
B-324	one individualized compound I	Clethodim
B-325	one individualized compound I	Cycloxydim
B-326	one individualized compound I	Profoxydim
B-327	one	Sethoxydim

	individualized compound I	
B-328	one individualized compound I	Tepraloxydim
B-329	one individualized compound I	Pendimethalin
B-330	one individualized compound I	Prodiamine
B-331	one individualized compound I	Trifluralin
B-332	one individualized compound I	Acifluorfen
B-333	one individualized compound I	Bromoxynil
B-334	one individualized compound I	Imazamethabenz
B-335	one individualized compound I	Imazamox
B-336	one individualized compound I	Imazapic
B-337	one individualized compound I	Imazapyr
B-338	one individualized	Imazaquin

	compound I	
B-339	one individualized compound I	Imazethapyr
B-340	one individualized compound I	2,4-Dichlorophenoxyacetic acid (2,4-D)
B-341	one individualized compound I	Chloridazon
B-342	one individualized compound I	Clopyralid
B-343	one individualized compound I	Fluroxypyr
B-344	one individualized compound I	Picloram
B-345	one individualized compound I	Picolinafen
B-346	one individualized compound I	Bensulfuron
B-347	one individualized compound I	Chlorimuron-ethyl
B-348	one individualized compound I	Cyclosulfamuron
B-349	one individualized compound I	Iodosulfuron

B-350	one individualized compound I	Mesosulfuron
B-351	one individualized compound I	Metsulfuron-methyl
B-352	one individualized compound I	Nicosulfuron
B-353	one individualized compound I	Rimsulfuron
B-354	one individualized compound I	Triflusulfuron
B-355	one individualized compound I	Atrazine
B-356	one individualized compound I	Hexazinone
B-357	one individualized compound I	Diuron
B-358	one individualized compound I	Florasulam
B-359	one individualized compound I	Pyroxasulfone
B-360	one individualized compound I	Bentazone
B-361	one	Cinidon-ethyl

	individualized compound I	
B-362	one individualized compound I	Cinmethylin
B-363	one individualized compound I	Dicamba
B-364	one individualized compound I	Diflufenzopyr
B-365	one individualized compound I	Quinclorac
B-366	one individualized compound I	Quinmerac
B-367	one individualized compound I	Mesotrione
B-368	one individualized compound I	Saflufenacil
B-369	one individualized compound I	Topramezone

The active substances referred to as component 2, their preparation and their activity against harmful fungi is known (cf.: <http://www.alanwood.net/pesticides/>); these substances are commercially available. The compounds described by IUPAC nomenclature, their preparation and their fungicidal activity are also known (cf. Can. J. Plant Sci. 48(6), 587-94, 1968; EP-A 141 317; EP-A 152 031; EP-A 226 917; EP-A 243 970; EP-A 256 503; EP-A 428 941; EP-A 532 022; EP-A 1 028 125; EP-A 1 035 122;

EP-A 1 201 648; EP-A 1 122 244, JP 2002316902; DE 19650197; DE 10021412; DE 102005009458; US 3,296,272; US 3,325,503; WO 98/46608; WO 99/14187; WO 99/24413; WO 99/27783; WO 00/29404; WO 00/46148; WO 00/65913; WO 01/54501; WO 01/56358; WO 02/22583; WO 02/40431; WO 03/10149; WO 03/11853; WO 03/14103; WO 03/16286; WO 03/53145; WO 03/61388; WO 03/66609; WO 03/74491; WO 04/49804; WO 04/83193; WO 05/120234; WO 05/123689; WO 05/123690; WO 05/63721; WO 05/87772; WO 05/87773; WO 06/15866; WO 06/87325; WO 06/87343; WO 07/82098; WO 07/90624, WO 11/028657).

The mixtures of active substances can be prepared as compositions comprising besides the active ingredients at least one inert ingredient by usual means, e. g. by the means given for the compositions of compounds I.

Concerning usual ingredients of such compositions reference is made to the explanations given for the compositions containing compounds I.


The mixtures of active substances according to the present invention are suitable as fungicides, as are the compounds of formula I. They are distinguished by an outstanding effectiveness against a broad spectrum of phytopathogenic fungi, especially from the classes of the Ascomycetes, Basidiomycetes, Deuteromycetes and Perono-sporomycetes (syn. Oomycetes). In addition, it is referred to the explanations regarding the fungicidal activity of the compounds and the compositions containing compounds I, respectively.

I. Synthesis examples

With due modification of the starting compounds, the procedures shown in the synthesis examples below were used to obtain further compounds I. The resulting compounds, together with physical data, are listed in Table I below.

I.1 Preparation of strobilurin type compounds I

Example 1: Preparation of (Z)-5-[1-(4-Chloro-phenyl)-1H-pyrazol-3-yloxy]-2-[(E)-methoxyimino]-3-methyl-pent-3-enic acid methyl amide (I-6)



Ex. 1a: (Z)-3-Tributylstannanyl-but-2-en-1-ol (3)

To 156.9 ml (156.9 mmol) of a 1 molar solution of lithium-aluminum-hydride in THF 0.77 g (14.3 mmol) sodium methylate have been added and the temperature has been lowered to 0°C afterwards. Then a solution of 10.0 g (142.7 mmol) 2-buten-1-ol in 108 ml THF has been added dropwise with stirring at this temperature. Stirring was continued for 36 h at 4°C. Thereafter, at about 0°C, 28.6 ml (292.5 mmol) ethyl acetate was added slowly while stirring. A strongly exothermic reaction was observed. Stirring was continued for 10 min without cooling. After cooling to about 0°C, 45.8 g (142.7 mmol) tri(n-butyl)stannyl-methanolate have been added dropwise while stirring. Stirring was continued for 2 d at 4°C. After addition of 112.9 g methanol stirring was continued for 1 h at ambient temperature. The reaction mixture was poured into 250 ml of water, extracted thrice with 150 ml diethyl ether each, the combined extracts were washed twice with 80 ml of water each, then once with 40 ml saturated aqueous sodium chloride solution, dried with sodium

sulfate and concentrated in vacuo. Yield 48.6 g oil, which have been purified by chromatography on 300 g silica gel with hexane/MTBE (10:1). Final yield 34.6 g (67%) oil. δ = 0.90 (m); 1.30 (m); 1.50 (m); 1.97 (s); 4.02 (t); 6.27 (t).

5

Ex. 1b : 1-(4-Chloro-phenyl)-3-((Z)-3-tributylstannanyl-but-2-enyloxy)-1H-pyrazole (5)

To 8.09 g (30.8 mmol) triphenylphosphine in 200 ml THF have been added with stirring at -75°C 6.23 g (30.8 mmol) azodicarbonic acid diisopropyl ester. The mixture has been stirred at this temperature for 5 min. Then 10.39 g (28.8 mmol) (Z)-3-tributylstannanyl-but-2-en-1-ol have been added dropwise and stirred for 5 min at -75°C. After addition of 4.00 g (20.6 mmol) 1-(4-chlorophenyl)-3-hydroxypyrazole at -75°C a red suspension was formed. The mixture was allowed to warm up to ambient temperature and stirred for 3 d. After removal of the solvents in vacuo 29 g oil have been collected and purified by chromatography on 120 g silica with MTBE/heptane. Yield 8.3 g (75%) oil. δ = 0.85 (m); 0.95 (m); 1.30 (m); 1.50 (m); 2.00 (s); 4.65 (d); 5.90 (d); 6.43 (t); 7.35 (d); 7.52 (d); 7.68 (d).

15

20

Ex. 1c: (Z)-5-[1-(4-Chloro-phenyl)-1H-pyrazol-3-yloxy]-2-[(E)-methoxyimino]-3-methyl-pent-3-enic acid methyl ester (7)

25

3.00 g (5.58 mmol) 1-(4-Chloro-phenyl)-3-((Z)-3-tributylstannanyl-but-2-enyloxy)-1H-pyrazole, 1.15 g (5.86 mmol) hydroxamic acid bromide (6), 0.155 g (0.67 mmol) tri(2-furyl)phosphine and 96 mg (0.17 mmol) bis(dibenzylidenacetone)-palladium have been stirred in 10 ml 1,4-dioxane for 4 d at about 80°C. After removal of the solvents in vacuo 4.3 g oil have been collected and purified by chromatography on 70 g silica with MTBE/heptane/1% triethylamine. Yield 1.2 g (59%) oil. δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.58 (d); 5.85 (d); 5.93 (t); 7.35 (d); 7.52 (d); 7.65 (d).


30

35

Ex. 1d: (Z)-5-[1-(4-Chloro-phenyl)-1H-pyrazol-3-yloxy]-2-[(E)-methoxyimino]-3-methyl-pent-3-enic acid methyl amide (I-6)


120 mg (0.33 mmol) (Z)-5-[1-(4-chloro-phenyl)-1H-pyrazol-3-yloxy]-2-[(E)-methoxy-imino]-3-methyl-pent-3-enic acid methyl ester and 0.99 ml (1.98 mmol) of a 2-molar solution of methyl amine in THF have been stirred overnight at ambient temperature in 2.0 ml of THF and 0.5 ml of water. After removal of the solvent in vacuo the product (120 mg) was collected in quantitative yield and good purity. Melting point 129-130°C.

Example 2: (Z)-5-[1-(4-Chloro-phenyl)-1H-pyrazol-3-yloxy]-2-[1-methoxy-meth-(E)-yl-idene]-3-methyl-pent-3-enoic acid methyl ester (compound I-7)



1.50 g (2.79 mmol) 1-(4-Chloro-phenyl)-3-((Z)-3-tributylstannanyl-but-2-enyloxy)-1H-pyrazole (5), 0.81 g (3.35 mmol) of ester compound (8) [Chem Comm 4, 423-425, (2006)], 0.223 g (0.28 mmol) [(R)-(+)-2,2"-Bis(diphenylphosphino)-1,1"-bisnaphthyl]-palladium(II)chloride have been stirred in 14 ml 1,4-dioxane for 4.5 d at about 100°C. After removal of the solvents in vacuo the crude product has been collected, which has been purified by chromatography on 50 g silica with MTBE/hexane/2% triethylamine. Yield 0.12 g, melting point 123-125°C.

Example 3: Methyl N-[(Z)-3-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-1-methyl-prop-1-enyl]-N-methyl-carbamate (I-10)



Ex. 3a: 1-(4-chlorophenyl)-3-[(Z)-3-iodobut-2-en-1-yloxy]pyrazole (10)

To 10.0 g (18.6 mmol) 1-(4-chloro-phenyl)-3-((Z)-3-tributylstannanyl-but-2-en-1-yloxy)-1H-pyrazole in 100 ml methylene chloride have been added 4.7 g (18.6 mmol) iodine at ambient temperature with stirring which was continued for 3 h. After removal of the solvents in vacuo the crude product was dissolved in 200 ml MTBE. 100 ml 20 % aqueous potassium fluoride solution have been added and the mixture was stirred for 2 h at ambient temperature. The aqueous layer was separated and extracted with 20 ml methyl-tert.-butyl ether twice. The combined organic phases have been washed twice with 20 ml of water each, dried with sodium sulfate, and the solvents were removed in vacuo. The crude product (7.2 g) has been purified by chromatography on 50 g silica with methyl-tert.-butylether/hexane (1:20). Yield 5.9 g, melting point 75-77°C.


Ex. 3b: Methyl N-[(Z)-3-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-1-methyl-prop-1-enyl]-N-methyl-carbamate

1.50 g (4.00 mmol) 1-(4-chlorophenyl)-3-[(Z)-3-iodobut-2-en-1-yloxy]pyrazole, 0.43 g (4.81 mmol) N-methylcarbamic acid methyl ester (9), 76 mg (0.4 mmol) copperiodide, 1.27 g (6.00 mmol) potassium phosphate and 71 mg (0.80 mmol) N,N'-dimethylethylenediamine in 14 ml toluene have been stirred for 1.5 d at 100°C. After removal of the solvents in vacuo the crude product has been purified by chromatography on 50 g silica with MTBE/hexane (1:3). Yield 0.67 g. ¹H-NMR (CDCl₃): δ = 1.90 (s); 3.05 (s); 3.70

(s); 4.68 (m); 5.63 (m); 5.90 (d); 7.35 (m); 7.55 (m); 7.68 (d).

Example 4: (Z,2E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,4-dimethyl-pent-3-enamide (compound I-79)

5



Ex. 4a: 1-[1-(4-Chlorophenyl)pyrazol-3-yl]oxypropan-2-one (11)

10 5.50 g (28.3 mmol) 1-(4-chlorophenyl)pyrazol-3-ol, 3.91 g (28.3 mmol) potassium carbonate and 50 mg sodium iodide in 30 ml DMF have been stirred for 5 min at ambient temperature. Then 2.62 g (28.3 mmol) chloroacetone have been added dropwise while stirring which was continued at 60°C for 5 h. The mixture was
 15 poured into excess 10 % aqueous lithium chloride solution and extracted with ethyl acetate three times. The combined extracts have been washed with 10 % lithium chloride solution twice and dried with sodium sulfate. After removal of the solvent in vacuo the crude product has been purified by chromatography on silica.
 20 Yield 6.5 g. The product was used for the next step.

Ex. 4b: Methyl (Z,2E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-4-methyl-pent-3-enoate and E-isomer (13)


To 5.26 g (21.0 mmol) 1-[1-(4-chlorophenyl)pyrazol-3-yl]oxypropan-2-one and 7.20 g (26.9 mmol) methyl (2Z)-3-diethoxyphosphoryl-2-methoxyimino-propanoate (which can be prepared as described for the dimethoxy derivative [(Tetrahedron Let 29, 3361-3364 (1988)] in 100 ml THF have been added at ambient temperature with stirring 2.59 g (23.1 mmol) potassium tert.-butylate. Stirring was continued over night. After removal of the solvent in vacuo the mixture was purified by chromatography on silica with heptane/ethyl acetate. 1.07 g of a 80:20 E:Z-mixture have been collected. This has been used directly for the next step.

Ex. 4c: Z,2E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-N,4-dimethyl-pent-3-enamide

0.68 g (1.87 mmol) methyl (Z,2E)-5-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-methoxyimino-4-methyl-pent-3-enoate and E-isomer from the previous reaction have been dissolved in 3.0 ml THF. 2.0 ml 40% aqueous methylamine have been added at ambient temperature with stirring which was continued over night. After removal of the solvent in vacuo the mixture was purified by chromatography on silica with a heptane/ethyl acetate gradient. 90 mg of the desired Z-isomer have been collected. ¹H-NMR (CDCl₃): δ = 2.03 (s); 2.85 (d); 3.97 (s); 4.64 (s); 5.87 (s); 6.02 (d); 6.65 (br); 7.35 (d); 7.50 (d); 7.67 (d).

Example 5: Methyl N-[(Z)-3-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-1-methyl-prop-1-enyl]-N-methoxy-carbamate (I-1)

30



Ex. 5a: Cis-3-methyloxiran-2-yl]methanol (15)

5 To 10.0 g (138.7 mmol) Z-but-2-en-1-ol in 140 ml dichloromethane (DCM) 37.6 g (152.6 mmol) 3-chloroperbenzoic acid (70 % purity) have been added at 0-5°C in small portions with stirring. Stirring was continued at 0°C for 2 h. 35.0 g calcium hydroxide have been added at 0-5°C with stirring which was continued for about 2 h at 0°C. The precipitate was filtered off, washed with DCM, and the filtrates dried with sodium sulfate. The solvent was largely removed at 380 mbar/30°C. The crude product (15.0 g, purity 75%) was used without further purification.

Ex. 5b: Racemic 1-(4-chlorophenyl)-3-[(2R,3S)-3-methyloxiran-2-yl]methoxy]pyrazole (16)

To 28.3 g (107.9 mmol) triphenylphosphine and 15.0g (127.7 mmol) cis-3-methyloxiran-2-yl]methanol (15) from the preceding experiment in 400 ml THF have been added with stirring at -75°C 22.9 g (113.0 mmol) azodicarbonic acid diisopropyl ester. The mixture has been stirred at this temperature for 5 min. Then 20.0 g (102.8 mmol) 1-(4-chlorophenyl)-3-hydroxypyrazole have been added with stirring at -70°C. The mixture was allowed to warm up to ambient temperature and stirred for about 1 d. After removal of the solvents in vacuo the crude product was stirred

with 200 ml of diisopropylether from which 42 g of a solid substance have been collected and further purified by chromatography on 330 g silica with MTBE/heptane. Yield 22.5 g (82%).

5

Ex. 5c: 5-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]-3-methoxy-4-methyl-oxazolidin-2-one (17)

To 1.08 g (10.0 mmol) methyl N-methoxycarbamate in 17 ml DMSO have been added with stirring at ambient temperature 1.00 g (8.9 mmol) potassium tert.-butylate. Stirring has been continued for 5 min before 2.00 g (7.56 mmol) 1-(4-chlorophenyl)-3-[[1-(2R,3S)-3-methyloxiran-2-yl]methoxy]pyrazole have been added. The mixture was stirred at 90°C for 20 h. After cooling to ambient temperature the reaction mixture was poured into 150 ml of water, extracted thrice with 30 ml ethyl acetate each, the combined extracts have been dried with sodium sulfate and the solvent removed in vacuo. The crude product (2.5 g) was used without further purification.

20

Ex. 5d: Racemic (2S,3R)-1-[[1-(4-chlorophenyl)pyrazol-3-yl]oxy-3-(methoxyamino)butan-2-ol (18)

To 10.0 g (30.0 mmol) 5-[[1-(4-chlorophenyl)pyrazol-3-yl]oxymethyl]-3-methoxy-4-methyl-oxazolidin-2-one in 100 ml ethanol has been added at ambient temperature with stirring 14.6 g (40.0 mmol) 21% sodium ethylate solution (in ethanol) and stirred over night. The solvent was largely removed in vacuo and the remaining mixture poured into 250 ml aqueous sodium dihydrogenphosphate solution, extracted thrice with 150 ml ethyl acetate each, dried with sodium sulfate to yield 8.7 g of an oil after evaporation of the solvent in vacuo. Further purified by chromatography on 120 g silica with MTBE/heptane. Yield 4.0 g (43%). ¹H-NMR (CDCl₃): δ = 1.22 (d); 3.17 (m); 3.57 (s); 3.91 (m); 4.33 (m); 4.47 (m); 5.93 (s); 7.35 (d); 7.52 (d); 7.68 (s).

35

Ex. 5e: Racemic methyl N-[(1R,2S)-3-[[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-hydroxy-1-methyl-propyl]-N-methoxy-carbamate (19)

To 1.00 g (3.2 mmol) Racemic (2S,3R-1-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-3-(methoxyamino)butan-2-ol in 7 ml THF 0.28 g (3.5 mmol) sodium hydrogencarbonate has been added.
 5 Then 0.33 g (3.5 mmol) methyl chloroformate has been added dropwise with stirring at ambient temperature. Stirring has been continued over night. The reaction mixture was poured into 10 ml of water, extracted thrice with 10 ml MTBE each, the combined
 10 extracts dried with sodium sulfate, and the solvent removed in vacuo. The crude product (1.3 g) was used without further purification.

Ex. 5f: Methyl N-[(Z)-3-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-1-methyl-prop-1-enyl]-N-methoxy-carbamate (I-1)

15 To 3.50 g (9.46 mmol) racemic methyl N-[(1R,2S)-3-[1-(4-chlorophenyl)pyrazol-3-yl]oxy-2-hydroxy-1-methyl-propyl]-N-methoxy-carbamate in 13 ml THF 2.61 g (9.94 mmol) triphenylphoshin has been added with stirring at ambient
 20 temperature. The mixture was cooled to -15°C. Then 2.11 g (10.41 mmol) azodicarbonic acid diisopropyl ester has been added with stirring which was continued for 1 h at 0°C and for about 1 d at ambient temperature. After removal of the solvents in vacuo the crude product was further purified by chromatography on 25
 25 g silica with MTBE/heptane/1% triethylamine. ¹H-NMR (CDCl₃): δ = 1.93 (s); 3.72 (s); 3.81 (s); 4.78 (d); 5.73 (t); 5.90 (d); 7.36 (d); 7.55 (d); 7.70 (d).

30 Table I. Compounds of formula I with physical data (melting point [°C]; ¹H-NMR (CDCl₃) (δ); HPLC/MS retention time [min])

No.	R ¹	R ²	R ³	R ⁴	X	Y	m.p. [°C]; ¹ H-NMR (δ); R _t [min]
I-1	CH ₃	H	R ³ - 1	R ⁴ -4, R ⁵ = OCH ₃	O	Y-1	δ = 1.93 (s); 3.72 (s); 3.81 (s); 4.78 (d); 5.73 (t); 5.90 (d); 7.36 (d); 7.55 (d); 7.70 (d)

I-2	CH ₃	H	R3-1	R4-4, R ⁵ = OCH ₃	NH	Y-1	
I-3	CH ₃	H	R3-1	R4-3	O	Y-1	
I-4	CH ₃	H	R3-1	R4-3	NH	Y-1	
I-5	CH ₃	H	R3-1	R4-1	O	Y-1	88-89°C
I-6	CH ₃	H	R3-1	R4-1	NH	Y-1	130°C
I-7	CH ₃	H	R3-1	R4-2	O	Y-1	123-125°C
I-8	CH ₃	H	R3-1	R4-2	NH	Y-1	
I-9	CH ₃	H	R3-1	R4-7		Y-1	
I-10	CH ₃	H	R3-1	R4-4, R ⁵ = CH ₃	O	Y-1	δ = 1.90 (s); 3.05 (s); 3.70 (s); 4.68 (m); 5.63 (m); 5.90 (d); 7.35 (m); 7.55 (m); 7.68 (d)
I-11	CH ₃	H	R3-2	R4-1	NH	Y-1	δ = 1.95 (s); 2.85 (d); 3.95 (s); 4.55 (m); 5.87 (m); 5.95 (m); 6.70 (br); 6.95 (m); 7.73 (m); 7.80 (m)
I-12	CH ₃	H	R3-3	R4-1	NH	Y-1	δ = 1.95 (s); 2.27 (s); 2.80 (d); 3.95 (s); 4.52 (m); 5.83 (d); 5.93 (m); 6.75 (br); 6.95 (m); 7.30 (m)
I-13	CH ₃	H	R3-4	R4-1	NH	Y-1	δ = 1.93 (s); 2.38 (s); 2.80 (d); 3.95 (s); 4.55 (m); 5.88 (d);

							5.95 (m); 6.75 (br); 7.13 (m); 7.28 (m); 7.41 (d); 7.64 (d)
I- 14	CH ₃	H	R3- 5	R4-1	NH	Y-1	125-127°C
I- 15	CH ₃	H	R3- 6	R4-1	NH	Y-1	δ = 1.95 (s); 2.85 (d); 3.95 (s); 4.55 (m); 5.90 (d); 5.95 (m); 6.70 (br); 7.32 (m); 7.50 (s); 7.54 (d); 7.70 (d)
I- 16	CH ₃	H	R3- 7	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.97 (s); 4.56 (m); 5.88 (d); 5.94 (m); 6.70 (br); 7.45 (m); 7.64 (s); 7.72 (d)
I- 17	CH ₃	H	R3- 8	R4-1	NH	Y-1	
I- 18	CH ₃	H	R3- 9	R4-1	NH	Y-1	δ = 1.95 (s); 2.33 (s); 2.88 (d); 3.97 (s); 4.56 (m); 5.85 (m); 5.95 (m); 6.70 (br); 7.03 (t); 7.30 (m); 7.44 (m); 7.62 (m)
I- 19	CH ₃	H	R3- 10	R4-1	NH	Y-1	
I- 20	CH ₃	H	R3- 11	R4-1	NH	Y-1	δ = 1.95 (s); 2.25 (s); 2.30 (s); 2.87 (d); 3.97 (s); 4.56 (m); 5.85 (d); 5.95 (m); 6.73 (br); 7.15 (d); 7.27 (m); 7.35 (d); 7.63 (d)
I- 21	CH ₃	H	R3- 12	R4-1	NH	Y-1	

I- 22	CH ₃	H	R3- 13	R4-1	NH	Y-1	
I- 23	CH ₃	H	R3- 14	R4-1	NH	Y-1	
I- 24	CH ₃	H	R3- 15	R4-1	NH	Y-1	
I- 25	CH ₃	H	R3- 16	R4-1	NH	Y-1	
I- 26	CH ₃	H	R3- 17	R4-1	NH	Y-1	
I- 27	CH ₃	H	R3- 14	R4-1	NH	Y-1	
I- 28	CH ₃	H	R3- 18	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.97 (s); 4.57 (m); 5.87 (m); 5.95 (m); 6.70 (br); 7.07 (m); 7.33 (m); 7.55 (m); 7.64 (m)
I- 29	CH ₃	H	R3- 19	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.97 (s); 4.58 (m); 5.88 (d); 5.94 (m); 6.70 (br); 7.15 (m); 7.30 (m); 7.43 (m); 7.64 (s); 7.70 (d)
I- 30	CH ₃	H	R3- 20	R4-1	NH	Y-1	
I- 31	CH ₃	H	R3- 21	R4-1	NH	Y-1	
I- 32	CH ₃	H	R3- 22	R4-1	NH	Y-1	δ = 1.95 (s); 2.37 (s); 2.87 (d); 3.95 (s); 4.55 (m); 5.86 (d); 5.93 (m); 6.70 (br); 7.20 (m); 7.45 (m); 7.65 (d)

I- 33	CH ₃	H	R3- 23	R4-1	NH	Y-1	$\delta = 1.23$ (m); 1.95 (s); 2.64 (m); 2.87 (d); 3.95 (s); 4.58 (m); 5.87 (d); 5.95 (m); 6.70 (br); 7.25 (m); 7.48 (m); 7.67 (d)
I- 34	CH ₃	H	R3- 24	R4-1	NH	Y-1	$\delta = 1.95$ (s); 2.90 (d); 3.97 (s); 4.58 (m); 5.92 (m); 6.70 (br); 7.67 (m); 7.75 (d)
I- 35	CH ₃	H	R3- 25	R4-1	NH	Y-1	
I- 36	CH ₃	H	R3- 26	R4-1	NH	Y-1	
I- 37	CH ₃	H	R3- 58	R4-1	NH	Y-1	
I- 38	CH ₃	H	R3- 28	R4-1	NH	Y-1	
I- 39	CH ₃	H	R3- 29	R4-1	NH	Y-1	
I- 40	CH ₃	H	R3- 30	R4-1	NH	Y-1	
I- 41	CH ₃	H	R3- 31	R4-1	NH	Y-1	
I- 42	CH ₃	H	R3- 32	R4-1	NH	Y-1	
I- 43	CH ₃	H	R3- 33	R4-1	NH	Y-1	
I- 44	CH ₃	H	R3- 34	R4-1	NH	Y-1	
I- 45	CH ₃	H	R3- 35	R4-1	NH	Y-1	
I- 46	CH ₃	H	R3- 36	R4-1	NH	Y-1	

I- 47	CH ₃	H	R3- 37	R4-1	NH	Y-1	
I- 48	CH ₃	H	R3- 38	R4-1	NH	Y-1	
I- 49	CH ₃	H	R3- 39	R4-1	NH	Y-1	
I- 50	CH ₃	H	R3- 40	R4-1	NH	Y-1	
I- 51	CH ₃	H	R3- 41	R4-1	NH	Y-1	
I- 52	CH ₃	H	R3- 42	R4-1	NH	Y-1	
I- 53	CH ₃	H	R3- 43	R4-1	NH	Y-1	
I- 54	CH ₃	H	R3- 44	R4-1	NH	Y-1	
I- 55	CH ₃	H	R3- 45	R4-1	NH	Y-1	
I- 56	CH ₃	H	R3- 46	R4-1	NH	Y-1	
I- 57	CH ₃	H	R3- 47	R4-1	NH	Y-1	
I- 58	CH ₃	H	R3- 45	R4-1	NH	Y-1	
I- 59	CH ₃	H	R3- 49	R4-1	NH	Y-1	
I- 60	CH ₃	H	R3- 50	R4-1	NH	Y-1	
I- 61	CH ₃	H	R3- 51	R4-1	NH	Y-1	
I- 62	CH ₃	H	R3- 52	R4-1	NH	Y-1	
I-	CH ₃	H	R3-	R4-1	NH	Y-1	

63			53				
I- 64	CH ₃	H	R3- 54	R4-1	NH	Y-1	
I- 65	CH ₃	H	R3- 55	R4-1	NH	Y-1	
I- 66	CH ₃	H	R3- 56	R4-1	NH	Y-1	
I- 67	CH ₃	H	R3- 57	R4-1	NH	Y-1	
I- 68	CH ₃	H	R3- 58	R4-4, = OCH ₃	R ⁵ O	Y-1	
I- 69	CH ₃	H	R3- 58	R4-4, = OCH ₃	R ⁵ NH	Y-1	
I- 70	CH ₃	H	R3- 58	R4-3	O	Y-1	
I- 71	CH ₃	H	R3- 58	R4-3	NH	Y-1	
I- 72	CH ₃	H	R3- 27	R4-1	O	Y-1	
I- 73	CH ₃	H	R3- 58	R4-2	O	Y-1	
I- 74	CH ₃	H	R3- 58	R4-2	NH	Y-1	
I- 75	CH ₃	H	R3- 58	R4-7		Y-1	
I- 76	C ₂ H ₅	H	R3- 1	R4-1	NH	Y-1	
I- 77	C ₂ H ₅	H	R3- 2	R4-1	NH	Y-1	
I- 78	C ₂ H ₅	H	R3- 58	R4-1	NH	Y-1	
I- 79	H	CH ₃	R3- 1	R4-1	NH	Y-1	$\delta = 2.03$ (s); 2.85 (d); 3.97 (s); 4.64 (s);

							5.87 (s); 6.02 (d); 6.65 (br); 7.35 (d); 7.50 (d); 7.67 (d)
I- 80	H	CH ₃	R3- 1	R4-1	O	Y-1	
I- 81	CH ₃	H	R3- 1	R4-1	NH	Y-3	
I- 82	CH ₃	H	R3- 59	R4-1	NH	Y-1	δ = 1.95 (s); 2.16 (s); 2.18 (s); 2.30 (s); 2.90 (s); 3.97 (s); 4.33 (m); 4.65 (m); 5.30 (m); 5.87 (m); 6.07 (m); 6.55 (s); 6.65 (br); 6.96 (s)
I- 83	CH ₃	H	R3- 60	R4-1	NH	Y-1	δ = 1.95 (s); 2.22 (s); 2.90 (d); 3.95 (s); 4.35 (m); 4.67 (m); 5.30 (m); 5.88 (m); 6.07 (m); 6.65 (br); 6.73 (d); 7.35 (d); 7.43 (s)
I- 84	CH ₃	H	R3- 61	R4-1	NH	Y-1	δ = 1.95 (s); 2.15 (s); 2.18 (s); 2.28 (s); 2.90 (d); 3.95 (s); 4.33 (m); 4.67 (s); 5.39 (d); 5.43 (d); 5.88 (m); 6.57 (s); 6.65 (br); 6.95 (s)
I- 85	CH ₃	H	R3- 62	R4-1	NH	Y-1	δ = 1.95 (s); 2.21 (s); 2.25 (s); 2.89 (d); 3.95 (s); 4.35 (m); 4.70 (s); 5.39 (d); 5.44 (d); 5.90 (m); 6.65 (br); 6.73 (d); 7.40 (d); 7.45 (s)
I-	CH ₃	H	R3-	R4-1	O	Y-1	δ = 1.73 (m); 1.93 (s);

86			63				2.15 (s); 2.30 (s); 3.85 (s); 4.05 (s); 4.33 (m); 4.60 (m); 5.75 (m); 5.85 (m); 6.55 (s); 6.95 (s)
I- 87	CH ₃	H	R3- 64	R4-1	O	Y-1	δ = 1.95 (s); 2.12 (s); 2.14 (s); 2.30 (s); 3.85 (s); 4.07 (s); 4.35 (m); 4.63 (m); 5.85 (m); 6.15 (m); 6.27 (m); 6.55 (s); 6.97 (s)
I- 88	CH ₃	H	R3- 65	R4-1	O	Y-1	δ = 1.95 (s); 2.17 (s); 2.19 (s); 3.85 (s); 4.05 (s); 4.35 (m); 4.65 (m); 5.87 (m); 6.17 (m); 6.31 (m); 6.72 (d); 7.35 (d); 7.43 (s).
I- 89	CH ₃	H	R3- 66	R4-1	NH	Y-1	δ = 1.95 (s); 2.12 (s); 2.14 (s); 2.30 (s); 2.90 (d); 3.95 (s); 3.97 (s); 4.32 (m); 5.87 (m); 6.57 (s); 6.65 (br); 6.97 (s)
I- 90	CH ₃	H	R3- 63	R4-1	NH	Y-1	δ = 1.73 (m); 1.93 (s); 2.15 (s); 2.30 (s); 2.90 (d); 3.95 (s); 4.33 (m); 4.55 (m); 5.75 (m); 5.90 (m); 6.57 (s); 6.65 (br); 6.95 (s)
1- 91	CH ₃	H	R3- 67	R4-1	NH	Y-1	δ = 1.73 (m); 1.93 (s); 2.18 (s); 2.20 (s); 2.90 (d); 3.95 (s); 4.35 (m); 4.60 (m);

							5.75 (m); 5.90 (m); 6.55 (br); 6.73 (m); 7.35 (m); 7.43 (s)
I-92	CH ₃	H	R3-64	R4-1	NH	Y-1	δ = 1.95 (s); 2.15 (s); 2.30 (s); 2.90 (d); 3.95 (s); 4.33 (m); 4.60 (m); 5.87 (m); 6.15 (m); 6.27 (m); 6.57 (s); 6.65 (br); 6.95 (s)
I-93	CH ₃	H	R3-65	R4-1	NH	Y-1	δ = 1.95 (s); 2.19 (s); 2.21 (s); 2.90 (d); 3.95 (s); 4.35 (m); 4.65 (m); 5.90 (m); 6.17 (m); 6.31 (m); 6.65 (br); 6.75 (d); 7.35 (m); 7.43 (s)
I-94	CH ₃	H	R3-59	R4-1	O	Y-1	δ = 1.95 (s); 2.15 (s); 2.30 (s); 3.85 (s); 4.07 (s); 4.33 (m); 4.63 (m); 5.30 (m); 5.87 (m); 6.07 (m); 6.53 (s); 6.97 (s)
I-95	CH ₃	H	R3-60	R4-1	O	Y-1	δ = 1.93 (s); 2.22 (s); 3.85 (s); 4.07 (s); 4.37 (m); 4.67 (m); 5.30 (m); 5.87 (m); 6.07 (m); 6.70 (d); 7.37 (m); 7.45 (s)
I-96	CH ₃	H	R3-61	R4-1	O	Y-1	δ = 1.95 (s); 2.15 (s); 2.17 (s); 2.28 (s); 3.87 (s); 4.07 (s); 4.33 (m); 4.67 (s); 5.40 (d); 5.43 (d); 5.87 (m); 6.53 (s); 6.95 (s)

I- 97	CH ₃	H	R3- 62	R4-1	O	Y-1	δ = 1.95 (s); 2.21 (s); 2.23 (s); 3.85 (s); 4.05 (s); 4.37 (m); 4.70 (s); 5.39 (d); 5.42 (d); 5.87 (m); 6.70 (d); 7.38 (m); 7.45 (s)
I- 98	CH ₃	H	R3- 68	R4-1	O	Y-1	δ = 1.23 (t); 1.87 (s); 2.08 (s); 2.27 (s); 3.70 (s); 3.95 (s); 4.10 (q); 4.30 (m); 5.90 (m); 6.65 (s); 7.00 (s)
I- 99	CH ₃	H	R3- 69	R4-1	O	Y-1	
I- 100	CH ₃	H	R3- 69	R4-1	NH	Y-1	
I- 101	CH ₃	H	R3- 69	R4-4, R ⁵ = OCH ₃	O	Y-1	
I- 102	CH ₃	H	R3- 70	R4-1	O	Y-6	
I- 103	CH ₃	H	R3- 70	R4-1	NH	Y-6	
I- 104	CH ₃	H	R3- 70	R4-4, R ⁵ = OCH ₃	O	Y-6	
I- 105	CH ₃	H	R3- 71	R4-1	O	Y-4	
I- 106	CH ₃	H	R3- 71	R4-1	NH	Y-4	
I- 107	CH ₃	H	R3- 71	R4-4, R ⁵ = OCH ₃	O	Y-4	
I- 108	CH ₃	H	R3- 72	R4-1	NH	Y-1	
I-	CH ₃	H	R3-	R4-1	O	Y-1	

109			72				
I- 110	CH ₃	H	R3- 72	R4-4, = OCH ₃	R ⁵ O	Y-1	
I- 111	CH ₃	H	R3- 72	R4-1	NH	Y-2	
I- 112	CH ₃	H	R3- 72	R4-1	O	Y-2	
I- 113	CH ₃	H	R3- 72	R4-4, = OCH ₃	R ⁵ O	Y-2	
I- 114	CH ₃	H	R3- 1	R4-4, = OCH ₃	R ⁵ O	Y-2	
I- 115	CH ₃	H	R3- 1	R4-4, = OCH ₃	R ⁵ NH	Y-2	
I- 116	CH ₃	H	R3- 1	R4-3	O	Y-2	
I- 117	CH ₃	H	R3- 1	R4-3	NH	Y-2	
I- 118	CH ₃	H	R3- 1	R4-1	O	Y-2	
I- 119	CH ₃	H	R3- 1	R4-1	NH	Y-2	
I- 120	CH ₃	H	R3- 1	R4-2	O	Y-2	
I- 121	CH ₃	H	R3- 1	R4-2	NH	Y-2	
I- 122	CH ₃	H	R3- 1	R4-7		Y-2	
I- 123	CH ₃	H	R3- 1	R4-4, = CH ₃	R ⁵ O	Y-2	
I- 124	CH ₃	H	R3- 73	R4-1	NH	Y-1	
I- 125	CH ₃	H	R3- 74	R4-1	NH	Y-1	

I- 126	CH ₃	H	R3- 75	R4-1	NH Y-1	
I- 127	CH ₃	H	R3- 76	R4-1	NH Y-1	
I- 128	CH ₃	H	R3- 77	R4-1	NH Y-1	
I- 129	CH ₃	H	R3- 78	R4-1	NH Y-1	
I- 130	CH ₃	H	R3- 78	R4-1	NH Y-1	
I- 131	CH ₃	H	R3- 80	R4-1	NH Y-1	
I- 132	CH ₃	H	R3- 81	R4-1	NH Y-1	
I- 133	CH ₃	H	R3- 82	R4-1	NH Y-1	
I- 134	CH ₃	H	R3- 83	R4-1	NH Y-1	
I- 135	CH ₃	H	R3- 84	R4-1	NH Y-1	
I- 136	CH ₃	H	R3- 85	R4-1	NH Y-1	
I- 137	CH ₃	H	R3- 86	R4-1	NH Y-1	
I- 138	CH ₃	H	R3- 87	R4-1	NH Y-1	
I- 139	CH ₃	H	R3- 88	R4-1	NH Y-1	
I- 140	CH ₃	H	R3- 89	R4-1	NH Y-1	
I- 141	CH ₃	H	R3- 90	R4-1	NH Y-1	
I-	CH ₃	H	R3-	R4-1	NH Y-1	

142			91				
I- 143	CH ₃	H	R3- 92	R4-1	NH	Y-1	
I- 144	CH ₃	H	R3- 93	R4-1	NH	Y-1	
I- 145	CH ₃	H	R3- 94	R4-1	NH	Y-1	
I- 146	CH ₃	H	R3- 95	R4-1	NH	Y-1	
I- 147	CH ₃	H	R3- 96	R4-1	NH	Y-1	
I- 148	CH ₃	H	R3- 97	R4-1	NH	Y-1	
I- 149	CH ₃	H	R3- 98	R4-1	NH	Y-1	
I- 150	CH ₃	H	R3- 99	R4-1	NH	Y-1	
I- 151	CH ₃	H	R3- 100	R4-1	NH	Y-1	
I- 152	CH ₃	H	R3- 101	R4-1	NH	Y-1	
I- 153	CH ₃	H	R3- 102	R4-1	NH	Y-1	
I- 154	CH ₃	H	R3- 103	R4-1	NH	Y-1	
I- 155	CH ₃	H	R3- 104	R4-1	NH	Y-1	
I- 156	CH ₃	H	R3- 105	R4-1	NH	Y-1	
I- 157	CH ₃	H	R3- 106	R4-1	NH	Y-1	
I- 158	CH ₃	H	R3- 107	R4-1	NH	Y-1	

I- 159	CH ₃	H	R3- 108	R4-1	NH	Y-1	
I- 160	CH ₃	H	R3- 109	R4-1	NH	Y-1	
I- 161	CH ₃	H	R3- 110	R4-1	NH	Y-1	
I- 162	CH ₃	H	R3- 111	R4-1	NH	Y-1	
I- 163	CH ₃	H	R3- 112	R4-1	NH	Y-1	
I- 164	CH ₃	H	R3- 113	R4-1	NH	Y-1	
I- 165	CH ₃	H	R3- 114	R4-1	NH	Y-1	
I- 166	CH ₃	H	R3- 115	R4-1	NH	Y-1	
I- 167	CH ₃	H	R3- 116	R4-1	NH	Y-1	
I- 168	CH ₃	H	R3- 117	R4-1	NH	Y-1	
I- 169	CH ₃	H	R3- 118	R4-1	NH	Y-4	
I- 170	CH ₃	H	R3- 119	R4-1	NH	Y-4	
I- 171	CH ₃	H	R3- 120	R4-1	NH	Y-4	
I- 172	CH ₃	H	R3- 121	R4-1	NH	Y-4	
I- 173	CH ₃	H	R3- 122	R4-1	NH	Y-4	
I- 174	CH ₃	H	R3- 123	R4-1	NH	Y-4	
I-	CH ₃	H	R3-	R4-1	NH	Y-4	

175			124				
I- 176	CH ₃	H	R3- 125	R4-1	NH	Y-4	
I- 177	CH ₃	H	R3- 126	R4-1	NH	Y-1	
I- 178	CH ₃	OCH ₃	R3- 1	R4-4, = OCH ₃	R ⁵ O	Y-1	
I- 179	CH ₃	OCH ₃	R3- 1	R4-4, = OCH ₃	R ⁵ O	Y-1	
I- 180	CH ₃	OCH ₃	R3- 1	R4-4, = OCH ₃	R ⁵ O	Y-1	
I- 181	CH ₃	H	R3- 1	R4-4, = OCH ₃	R ⁵ O	Y-1	
I- 182	CH ₃	H	R3- 1	R4-1	NH	Y-1	
I- 183	CH ₃	H	R3- 1	R4-1	O	Y-1	
I- 184	CH ₃	H	R3- 1	R4-2	NH	Y-1	
I- 185	CH ₃	H	R3- 1	R4-2	O	Y-1	
I- 186	CH ₃	H	R3- 127	R4-1	O	Y-1	79-81°C
I- 187	CH ₃	H	R3- 127	R4-1	NH	Y-1	δ = 1.95 (s); 2.12 (s); 2.18 (s); 2.87 (d); 3.97 (s); 3.98 (s); 4.35 (m); 5.90 (m); 6.65 (br); 7.35 (d); 7.43 (s)
I- 188	C ₂ H ₅	H	R3- 1	R4-1	O	Y-1	δ = 1.05 (m); 2.30 (m); 3.85 (s); 4.05 (s); 4.62 (m); 5.90 (m); 7.35 (m); 7.52 (m); 7.70 (d)

I- 189	CH ₃	H	R3- 127	R4-4, = CH ₃	R ⁵ O	Y-1	δ = 1.90 (s); 2.22 (s); 2.24 (s); 3.02 (s); 3.70 (s); 3.98 (s); 4.48 (m); 5.57 (m); 6.75 (m); 7.40 (m); 7.45 (s)
I- 190	CH ₃	H	R3- 66	R4-1	O	Y-1	δ = 1.95 (s); 2.10 (s); 2.13 (s); 2.30 (s); 3.85 (s); 3.92 (s); 4.05 (s); 4.35 (m); 5.87 (m); 6.53 (s); 6.97 (s)
I- 191	CH ₃	H	R3- 128	R4-1	O	Y-1	δ = 1.30 (t); 1.95 (s); 2.20 (s); 2.22 (s); 3.85 (s); 4.06 (s); 4.20 (q); 4.37 (m); 5.87 (m); 6.71 (d); 7.37 (m); 7.45 (s)
I- 192	CH ₃	H	R3- 128	R4-1	NH	Y-1	δ = 1.30 (t); 1.95 (s); 2.20 (s); 2.21 (s); 2.92 (d); 3.95 (s); 4.22 (m); 4.37 (m); 5.90 (m); 6.65 (br); 6.73 (m); 7.37 (m); 7.45 (s)
I- 193	CH ₃	H	R3- 68	R4-1	NH	Y-1	δ = 1.32 (t); 1.95 (s); 2.16 (s); 2.17 (s); 2.32 (s); 2.92 (d); 3.97 (s); 4.20 (q); 4.33 (m); 5.90 (m); 6.57 (s); 6.65 (br); 7.97 (s)
I- 194	CH ₃	H	R3- 5	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.05 (s); 4.53 (m); 5.87 (d); 5.93 (m); 7.07 (m); 7.25 (m);

							7.55 (m); 7.63 (d)
I- 195	CH ₃	H	R3- 6	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.05 (s); 1.55 (m); 5.88 (d); 5.93 (m); 7.33 (m); 7.48 (s); 7.53 (d); 7.70 (d)
I- 196	CH ₃	H	R3- 2	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.55 (m); 5.87 (d); 5.92 (m); 6.95 (m); 7.73 (d); 7.80 (m)
I- 197	CH ₃	H	R3- 4	R4-1	O	Y-1	δ = 1.95 (s); 2.37 (s); 3.85 (s); 4.07 (s); 4.55 (m); 5.85 (d); 5.92 (m); 7.12 (m); 7.28 (m); 7.43 (d); 7.67 (d)
I- 198	CH ₃	H	R3- 22	R4-1	O	Y-1	δ = 1.95 (s); 2.35 (s); 3.85 (s); 4.05 (s); 4.57 (m); 5.83 (d); 5.93 (m); 7.20 (d); 7.47 (d); 7.66 (d)
I- 199	CH ₃	H	R3- 24	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.60 (m); 5.95 (m); 7.67 (m); 7.77 (d)
I- 200	CH ₃	H	R3- 22	R4-1	O	Y-1	δ = 1.25 (m); 1.95 (s); 2.65 (m); 3.85 (s); 4.05 (s); 4.58 (m); 5.83 (d); 5.93 (m); 7.25 (m); 7.48 (m); 7.67 (d)
I- 201	CH ₃	H	R3- 19	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.58 (m); 5.87 (d); 5.93 (m); 7.15 (m); 7.30 (m);

							7.45 (m); 7.63 (s); 7.70 (d)
I- 202	CH ₃	H	R3- 7	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.58 (m); 5.90 (d); 5.95 (m); 7.45 (m); 7.67 (d); 7.73 (d)
I- 203	CH ₃	H	R3- 129	R4-1	O	Y-1	δ = 1.35 (d); 1.95 (s); 3.85 (s); 4.07 (s); 4.55 (m); 5.84 (d); 5.93 (m); 6.92 (m); 7.45 (m); 7.60 (d)
I- 204	CH ₃	H	R3- 11	R4-1	O	Y-1	δ = 1.95 (s); 2.27 (s); 2.32 (s); 3.85 (s); 4.07 (s); 4.58 (m); 5.85 (d); 5.95 (m); 7.15 (d); 7.30 (m); 7.37 (s); 7.66 (d)
I- 205	CH ₃	H	R3- 3	R4-1	O	Y-1	δ = 1.95 (s); 2.25 (s); 3.83 (s); 4.05 (s); 4.54 (m); 5.82 (d); 5.92 (m); 6.95 (m); 7.25 (m); 7.32 (d)
I- 206	CH ₃	H	R3- 9	R4-1	O	Y-1	δ = 1.93 (s); 2.31 (s); 3.85 (s); 4.07 (s); 4.56 (m); 5.85 (m); 5.95 (m); 7.03 (m); 7.33 (m); 7.43 (m); 7.64 (m).
I- 207	CH ₃	H	R3- 18	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.58 (m); 5.85 (d); 5.95 (m); 7.05 (m); 7.33 (m); 7.55 (m); 7.65 (d)
I- 208	CH ₃	H	R3- 129	R4-1	NH	Y-1	δ = 1.33 (d); 1.95 (s); 2.87 (d); 3.97 (s);

							4.55 (m); 5.84 (m); 5.95 (m); 6.73 (br); 6.92 (d); 7.45 (d); 7.62 (m)
I- 209	CH ₃	H	R3- 58	R4-1	O	Y-1	δ = 1.95 (s); 3.83 (s); 4.05 (s); 4.65 (m); 5.95 (m); 7.43 (d); 7.55 (d); 8.23 (s)
I- 210	Cl	CH ₃	R3- 1	R4-1	NH	Y-1	δ = 2.10 (s); 2.65 (d); 4.01 (s); 4.95 (s); 5.97 (d); 6.85 (br); 7.37 (d); 7.50 (d); 7.67 (d)
I- 211	phenyl ring*		R3- 1	R4-7 R ⁶ =CH ₃		Y-1	90°C
I- 212	phenyl ring*		R3- 98	R4-7 R ⁶ =CH ₃		Y-1	151°C
I- 213	phenyl ring*		R3- 130	R4-7 R ⁶ =CH ₃		Y-4	
I- 214	phenyl ring*		R3- 131	R4-7 R ⁶ =CH ₃		Y-4	139-142°C
I- 215	phenyl ring*		R3- 132	R4-7 R ⁶ =CH ₃		Y-4	58-63°C
I- 216	phenyl ring*		R3- 133	R4-7 R ⁶ =CH ₃		Y-1	111-113°C
I- 217	phenyl ring*		R3- 65	R4-7 R ⁶ =CH ₃		Y-1	78-88°C
I- 218	phenyl ring*		R3- 60	R4-7 R ⁶ =CH ₃		Y-1	83-88°C
I- 219	phenyl ring*		R3- 67	R4-7 R ⁶ =CH ₃		Y-1	87-89°C
I- 220	phenyl ring*		R3- 62	R4-7 R ⁶ =CH ₃		Y-1	82-88°C
I-	phenyl ring*		R3-	R4-7		Y-1	92-98°C

221		134	R ⁶ =CH ₃			
I- 222	phenyl ring*	R3- 135	R4-7 R ⁶ =CH ₃	Y-1	141-145°C	
I- 223	phenyl ring*	R3- 64	R4-7 R ⁶ =CH ₃	Y-1	88-91 °C	
I- 224	phenyl ring*	R3- 59	R4-7 R ⁶ =CH ₃	Y-1	82-86°C	
I- 225	phenyl ring*	R3- 63	R4-7 R ⁶ =CH ₃	Y-1	64-70°C	
I- 226	phenyl ring*	R3- 136	R4-7 R ⁶ =CH ₃	Y-1	101-108°C	
I- 227	phenyl ring*	R3- 137	R4-7 R ⁶ =CH ₃	Y-1	98-102°C	
I- 228	phenyl ring*	R3- 138	R4-7 R ⁶ =CH ₃	Y-1	71-75°C	
I- 229	phenyl ring*	R3- 139	R4-7 R ⁶ =CH ₃	Y-7	72-74°C	
I- 230	phenyl ring*	R3- 140	R4-7 R ⁶ =CH ₃	Y-7	δ = 1.65 (s); 3.32 (s); 3.67 (s); 5.20 (s); 7.07 (m); 7.27 (m); 7.47 (m)	
I- 231	phenyl ring*	R3- 141	R4-7 R ⁶ =CH ₃	Y-1	136-138°C	
I- 232	phenyl ring*	R3- 142	R4-7 R ⁶ =CH ₃	Y-7	δ = 1.65 (s); 3.43 (s); 3.70 (s); 5.20 (s); 7.30-7.65 (m)	
I- 233	phenyl ring*	R3- 127	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.678 min	
I- 234	phenyl ring*	R3- 143	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.092 min	
I- 235	phenyl ring*	R3- 144	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.065 min	
I-	phenyl ring*	R3-	R4-7	Y-1	R _t = 3.947 min	

236		145	R ⁶ =CH ₃		
I- 237	phenyl ring*	R3- 146	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.875 min
I- 238	phenyl ring*	R3- 147	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.100 min
I- 239	phenyl ring*	R3- 68	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.933 min
I- 240	phenyl ring*	R3- 148	R4-7 R ⁶ =CH ₃	Y-4	R _t = 3.496 min
I- 241	phenyl ring*	R3- 149	R4-7 R ⁶ =CH ₃	Y-4	R _t = 4.075 min
I- 242	phenyl ring*	R3- 150	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.128 min
I- 243	phenyl ring*	R3- 151	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.894 min
I- 244	phenyl ring*	R3- 152	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.875 min
I- 245	phenyl ring*	R3- 453	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.995 min
I- 246	phenyl ring*	R3- 154	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.115 min
I- 247	phenyl ring*	R3- 20	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.715 min
I- 248	phenyl ring*	R3- 157	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.714 min
I- 249	phenyl ring*	R3- 128	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.861 min
I- 250	phenyl ring*	R3- 158	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.251 min
I- 251	phenyl ring*	R3- 159	R4-7 R ⁶ =CH ₃	Y-1	105-107°C
I- 252	phenyl ring*	R3- 151	R4-7 R ⁶ =Ethyl	Y-1	102-104°C

I- 253	phenyl ring*	R3- 1	R4-7 R ⁶ =CHF ₂	Y-1	96-97°C
I- 254	phenyl ring*	R3- 128	R4-7 R ⁶ =Ethyl	Y-1	R _t = 4.219 min
I- 255	phenyl ring*	R3- 68	R4-7 R ⁶ =Ethyl	Y-1	R _t = 4.239 min
I- 256	phenyl ring*	R3- 146	R4-7 R ⁶ =Ethyl	Y-1	R _t = 4.176 min
I- 257	phenyl ring*	R3- 66	R4-7 R ⁶ =Ethyl	Y-1	R _t = 4.013 min
I- 258	phenyl ring*	R3- 160	R4-7 R ⁶ =CH ₃	Y-1	δ = 2.15 (s); 3.67 (s); 3.95 (s); 5.28 (s); 6.85 (m); 7.45 (m); 7.50 (m); 7.70 (m)
I- 259	phenyl ring*	R3- 161	R4-7 R ⁶ =CH ₃	Y-1	δ = 1.33 (t); 2.17 (s); 3.68 (s); 4.22 (q); 5.28 (s); 6.87 (d); 7.45 (m); 7.52 (m); 7.73 (m)
I- 260	phenyl ring*	R3- 162	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.924 min
I- 261	phenyl ring*	R3- 163	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.131 min
I- 262	phenyl ring*	R3- 164	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.141 min
I- 263	phenyl ring*	R3- 165	R4-7 R ⁶ =CH ₃	Y-1	R _t = 3.916 min
I- 264	phenyl ring*	R3- 166	R4-7 R ⁶ - CH ₃	Y-1	R _t = 4.220 min
I- 265	phenyl ring*	R3- 167	R4-7 R ⁶ =CH ₃	Y-1	R _t = 4.050 min
I- 266	phenyl ring*	R3- 149	R4-7 R ⁶ =CH ₃	Y-8	δ = 2.03 (s); 3.71 (s); 7.45-7.62 (m)

I- 267	phenyl ring*	R3- 168	R4-7 R ⁶ =CH ₃	Y-1	δ = 2.20 (s); 2.40 (s); 3.69 (s); 5.20 (s); 6.70 (s); 7.12 (s); 7.42 (s); 7.53 (m); 7.72 (m); 7.91 (s)
I- 268	phenyl ring*	R3- 169	R4-7 R ⁶ =CH ₃	Y-1	δ = 2.11 (s); 2.18 (s); 3.62 (s); 3.72 (s); 5.21 (s); 6.77 (s); 6.96 (s); 7.25 (s); 7.53 (m); 7.72 (m); 8,91 (s)
I- 269	phenyl ring*	R3- 170	R4-7 R ⁶ =CH ₃	Y-1	δ = 2.00 (s); 2.05 (s); 2.16 (s); 2.21 (s); 3.70 (s); 5.16 (s); 6.72 (s); 6.82 (s); 7.50 (m); 7.74 (m)
I- 270	phenyl ring*	R3- 171	R4-7 R ⁶ =CH ₃	Y-1	δ = 2.16 (s); 2.39 (s); 2.53 (s); 3.70 (s); 5.19 (s); 6.70 (s); 7.53 (m); 7.72 (m)
I- 271	phenyl ring*	R3- 172	R4-7 R ⁶ =CH ₃	Y-1	δ = 2.18 (s); 2.29 (s); 3.70 (s); 5.18 (s); 6.71 (s); 7.06 (s); 7.50 (m); 7.71 (m); 8.38 (s); 8.44 (s)
I- 272	phenyl ring*	R3- 173	R4-7 R ⁶ =CH ₃	Y-1	δ = 1.28 (m); 2.12 (s); 3.00 (broad); 3.58 (m); 3.69 (s); 5.52 (s); 6.28 (s); 7.45 (m); 7.72 (m)
I- 273	phenyl ring*	R3- 174	R4-7 R ⁶ =CH ₃	Y-1	δ = 0.97 (m); 1.02 (m); 1.20 (m); 1.90 (m); 2.05 (s); 2.62 (m); 3.68 (s); 5.51 (s); 5.84 (s); 7.50 (m); 7.73 (m)

I- 274	phenyl ring*		R3- 175	R4-7 R ⁶ =CH ₃		Y-8	R _t = 3.715 min
I- 275	CH ₃	H	R3- 20	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.58 (m); 5.92 (m); 7.45 (m); 7.52 (m); 7.75 (m); 7.84 (d)
I- 276	CH ₃	H	R3- 12	R4-1	O	Y-1	72-74°C
I- 277	CH ₃	H	R3- 15	R4-1	O	Y-1	73-75°C
I- 278	CH ₃	H	R3- 176	R4-1	O	Y-1	δ = 1.97 (s); 3.85 (s); 4.07 (s); 4.58 (d); 5.90 (d); 5.95 (m); 7.17 (t); 7.43 (m); 7.65 (m)
I- 279	CH ₃	H	R3- 98	R4-1	O	Y-1	90-92°C
I- 280	CH ₃	H	R3- 20	R4-1	NH	Y-1	101-103°C
I- 281	CH ₃	H	R3- 12	R4-1	NH	Y-1	112-115°C
I- 282	CH ₃	H	R3- 15	R4-1	NH	Y-1	93-95°C
I- 283	CH ₃	H	R3- 176	R4-1	NH	Y-1	84-86°C
I- 284	CH ₃	H	R3- 98	R4-1	NH	Y-1	δ = 1.95 (s); 2.27 (s); 2.93 (d); 3.97 (s); 4.48 (m); 5.93 (m); 6.42 (m); 6.67 (broad); 6.80 (d); 7.37 (d); 7.45 (m); 7.68 (m); 7.83 (m)
I-	CH ₃	H	R3-	R4-4, R ⁵	O	Y-1	δ = 1.93 (s); 3.72 (s);

285			1	= OCH ₃			3.81 (s); 4.78 (d); 5.73 (t); 5.90 (d); 7.36 (d); 7.55 (d); 7.70 (d)
I- 286	CH ₃	H	R3- 51	R4-1	O	Y-1	δ = 1.93 (s); 2.45 (s); 3.82 (s); 4.05 (s); 4.60 (d); 5.92 (t); 7.25-7.50 (m)
I- 287	C ₂ H ₅	H	R3- 1	R4-1	NH	Y-1	δ = 1.05 (t); 2.30 (q); 2.90 (d); 3.97 (s); 4.58 (m); 5.93 (m); 6.70 (broad); 7.35 (m); 7.53 (m); 7.68 (s)
I- 288	CH ₃	H	R3- 58	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.97 (s); 4.63 (m); 5.95 (m); 6.75 (broad); 7.43 (m); 7.55 (m); 8.23 (s)
I- 289	CH ₃	H	R3- 51	R4-1	NH	Y-1	δ = 1.93 (s); 2.45 (s); 2.83 (d); 3.95 (s); 4.58 (m); 5.92 (m); 6.73 (broad); 7.30-7.50 (m)
I- 290	CH ₃	H	R3- 1	R4-4, R ⁵ = C ₂ H ₅	O	Y-1	δ = 1.18 (m); 1.95 (s); 3.45 (broad); 3.73 (s); 4.67 (m); 5.70 (m); 5.90 (m); 7.35 (m); 7.53 (m); 7.67 (m)
I- 291	CH ₃	H	R3- 39	R4-1	O	Y-1	72°C
I- 292	CH ₃	H	R3- 44	R4-1	O	Y-1	106-110°C
I- 293	CH ₃	H	R3- 44	R4-1	NH	Y-1	177°C
I-	CH ₃	H	R3-	R4-1	O	Y-1	δ = 1.93 (s); 3.20 (m);

294			177				3.28 (m); 3.83 (s); 3.90 (m); 4.05 (s); 4.42 (m); 4.50 (m); 5.88 (m); 5.94 (s); 6.83 (m); 7.25 (m); 7.50 (s)
I- 295	CH ₃	H	R3- 39	R4-1	NH	Y-1	142°C
I- 296	Isobutyl	H	R3- 1	R4-1	O	Y-1	δ = 0.93 (m); 1.60 (m); 2.15 (d); 3.84 (s); 4.07 (s); 4.62 (m); 5.89 (m); 7.35 (d); 7.53 (d); 7.67 (s)
I- 297	CH ₃	H	R3- 29	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.65 (m); 5.95 (m); 7.27 (m); 7.87 (m); 8.35 (s)
I- 298	CH ₃	H	R3- 36	R4-1	O	Y-1	δ = 1.95 (s); 3.85 (s); 4.07 (s); 4.65 (m); 5.95 (m); 7.63 (m); 7.67 (m); 7.93 (s); 8.37 (s)
I- 299	Isobutyl	H	R3- 1	R4-1	NH	Y-1	177°C
I- 300	C ₂ H ₅	H	R3- 58	R4-1	O	Y-1	87°C
I- 301	CH ₃	H	R3- 29	R4-1	NH	Y-1	140°C
I- 302	CH ₃	H	R3- 36	R4-1	NH	Y-1	125°C
I- 303	CH ₃	H	R3- 33	R4-1	O	Y-1	δ = 1.95 (s); 3.83 (s); 4.05 (s); 4.65 (m); 5.95 (m); 7.37 (m); 7.57 (m); 8.27 (s)
I-	CH ₃	H	R3-	R4-1	O	Y-1	δ = 1.25 (m); 1.95 (s);

304			54				2.68 (m); 3.83 (s); 4.05 (s); 4.65 (m); 5.95 (m); 7.30 (m); 7.51 (m); 8.20 (s)
I- 305	CH ₃	H	R3- 33	R4-1	NH	Y-1	126°C
I- 306	CH ₃	H	R3- 54	R4-1	NH	Y-1	128°C
I- 307	CH ₃	H	R3- 178	R4-1	O	Y-1	δ = 1.95 (s); 2.27 (s); 3.82 (s); 4.07 (s); 4.62 (m); 5.95 (m); 7.28 (m); 7.36 (m); 7.93 (s)
I- 308	CH ₃	H	R3- 130	R4-1	O	Y-4	δ = 1.95 (s); 2.25 (s); 3.80 (s); 4.05 (s); 4.55 (m); 5.90 (m); 7.25-7.60 (m); 7.77 (m); 7.85 (s); 8.05 (s); 8.27 (s)
I- 309	CH ₃	H	R3- 130	R4-1	NH	CH =N - O- CH ₂ -	δ = 1.95 (s); 2.88 (d); 3.96 (s); 4.50 (m); 5.92 (m); 6.70 (broad); 7.25-7.60 (m); 7.76 (m); 8.05 (s); 8.08 (s); 8.27 (s)
I- 310	CH ₃	H	R3- 42	R4-1	O	Y-1	116°C
I- 311	C ₂ H ₅	H	R3- 58	R4-1	NH	Y-1	154°C
I- 312	CH ₃	H	R3- 30	R4-1	NH	Y-1	167°C
I- 313	CH ₃	H	R3- 178	R4-1	NH	Y-1	143°C
I- 314	CH ₃	H	R3- 42	R4-1	NH	Y-1	147°C

I- 315	CH ₃	H	R3- 40	R4-1	NH	Y-1	153°C
I- 316	CH ₃	H	R3- 177	R4-1	NH	Y-1	δ = 1.93 (s); 2.88 (d); 3.20 (m); 3.28 (m); 3.90 (m); 3.95 (s); 4.42 (m); 4.50 (m); 5.90 (m); 5.95 (s); 6.67 (broad); 9.86 (m); 7.25 (m); 7.50 (s)
I- 317	CH ₃	H	R3- 179	R4-1	NH	Y-1	δ = 1.83 (s); 1.93 (s); 2.18 (s); 2.42 (m); 2.70 (m); 2.88 (d); 3.83 (s); 3.95 (s); 4.35 (m); 5.90 (m); 6.65 (m); 6.93 (m)
I- 318	CH ₃	H	R3- 38	R4-1	O	Y-1	87°C
I- 319	CH ₃	H	R3- 32	R4-1	O	Y-1	δ = 1.93 (s); 2.27 (s); 3.80 (s); 4.05 (s); 4.62 (m); 5.95 (m); 7.30 (m); 7.93 (s)
I- 320	CH ₃	H	R3- 34	R4-1	NH	Y-1	120°C
I- 321	CH ₃	H	R3- 38	R4-1	NH	Y-1	143°C
I- 322	CH ₃	H	R3- 37	R4-1	NH	Y-1	117°C
I- 323	CH ₃	H	R3- 40	R4-1	O	Y-1	118°C
I- 324	CH ₃	H	R3- 37	R4-1	O	Y-1	δ = 1.93 (s); 2.23 (s); 2.35 (s); 3.80 (s); 4.05 (s); 4.62 (m); 5.95 (m); 7.15 (m); 7.91 (s)
I- 325	CH ₃	H	R3- 40	R4-1	NH	Y-1	136°C

325			43				
I- 326	CH ₃	H	R3- 34	R4-1	O	Y-1	δ = 1.95 (s); 3.83 (s); 4.05 (s); 4.63 (m); 5.95 (m); 7.67 (m); 7.82 (m); 8.42 (s)
I- 327	CH ₃	H	R3- 32	R4-1	NH	Y-1	δ = 2.07 (s); 2.42 (s); 2.92 (d); 4.08 (s); 4.78 (m); 6.08 (m); 6.93 (broad); 7.45 (m); 8.08 (s)
I- 328	CH ₃	H	R3- 41	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.95 (s); 4.65 (m); 5.93 (m); 6.73 (broad); 7.30 (m); 7.55 (m); 8.23 (s)
I- 329	CH ₃	H	R3- 28	R4-1	NH	Y-1	94°C
I- 330	CH ₃	H	R3- 50	R4-1	NH	Y-1	130°C
I- 331	CH ₃	H	R3- 41	R4-1	O	Y-1	δ = 1.95 (s); 3.87 (s); 4.06 (s); 4.63 (m); 5.93 (m); 7.32 (m); 7.55 (m); 8.23 (s)
I- 332	C ₂ H ₅	H	R3- 2	R4-1	O	Y-1	δ = 1.05 (t); 2.30 (m); 3.82 (s); 4.05 (s); 4.62 (d); 5.90 (m); 6.93 (m); 7.73 (m); 7.80 (m)
I- 333	CH ₃	H	R3- 180	R4-1	O	Y-1	δ = 1.95 (s); 2.27 (s); 3.85 (s); 4.07 (s); 4.40 (m); 5.88 (m); 6.66 (d); 6.78 (m); 7.37 (m); 7.45 (m); 7.83 (d)
I-	CH ₃	H	R3-	R4-1	O	Y-1	123-124°C

334			77				
I- 335	CH ₃	H	R3- 119	R4-1	O	Y-4	δ = 1.90 (s); 2.12 (s); 3.83 (s); 3.92 (s); 4.03 (s); 4.33 (m); 5.74 (m); 7.18-7.45 (m)
I- 336	CH ₃	H	R3- 80	R4-1	O	Y-1	133-134°C
I- 337	CH ₃	H	R3- 180	R4-1	NH	Y-1	δ = 1.95 (s); 2.27 (s); 2.90 (d); 3.97 (s); 4.40 (m); 5.88 (m); 6.66 (broad); 6.82 (m); 7.37 (m); 7.45 (m); 7.81 (d)
I- 338	CH ₃	H	R3- 79	R4-1	O	Y-1	94-97°C
I- 339	CH ₃	H	R3- 119	R4-1	NH	Y-4	δ = 1.90 (s); 2.12 (s); 2.85 (d); 3.92 (s); 4.33 (m); 5.75 (m); 6.60 (broad); 7.18-7.45 (m)
I- 340	CH ₃	H	R3- 130	R4-1	O	CH =N - O- CH ₂ -	δ = 1.95 (s); 3.82 (s); 4.05 (s); 4.52 (m); 5.87 (m); 7.25-7.60 (m); 7.75 (m); 7.80 (s); 8.05 (s); 8.26 (s)
I- 341	CH ₃	H	R3- 130	R4-1	NH	Y-4	δ = 1.95 (s); 2.25 (s); 2.83 (d); 3.97 (s); 4.54 (m); 5.93 (m); 6.75 (broad); 7.25-7.60 (m); 7.77 (m); 7.82 (s); 8.02 (s); 8.29 (s)
I- 342	CH ₃	H	R3- 181	R4-1	NH	Y-1	δ = 1.95 (s); 2.27 (s); 2.88 (d); 3.97 (s); 4.39 (m); 5.93 (m); 6.67 (broad); 6.80 (m);

							7.08 (m); 7.27 (m); 7.45 (m)
I- 343	CH ₃	H	R3- 182	R4-1	NH	Y-1	δ = 1.95 (s); 2.27 (s); 2.91 (d); 3.98 (s); 4.41 (m); 5.93 (m); 6.70 (broad); 6.85 (m); 7.37 (m); 7.65 (m)
I- 344	CH ₃	H	R3- 183	R4-1	NH	Y-1	δ = 1.95 (s); 2.25 (s); 2.91 (d); 3.98 (s); 4.40 (m); 5.93 (m); 6.66 (broad); 6.83 (m); 7.25-7.46 (m)
I- 345	CH ₃	H	R3- 184	R4-1	NH	Y-1	δ = 1.95 (s); 2.25 (s); 2.90 (d); 3.85 (s); 3.98 (s); 4.38 (m); 5.93 (m); 6.66 (broad); 6.82 (m); 7.30 (m)
I- 346	CH ₃	H	R3- 185	R4-1	NH	Y-1	δ = 1.95 (s); 2.20 (s); 2.88 (d); 3.95 (s); 4.30 (m); 4.98 (s); 5.92 (m); 6.65 (m); 6.80 (m); 7.40 (m)
I- 347	CH ₃	H	R3- 69	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.95 (s); 4.65 (m); 5.92 (m); 6.67 (broad); 6.87 (m); 7.25 (m); 7.67 (m)
I- 348	CH ₃	H	R3- 186	R4-1	NH	Y-1	δ = 1.42 (m); 1.95 (s); 2.27 (s); 2.91 (d); 2.96 (m); 3.98 (s); 4.42 (m); 5.93 (m); 6.70 (broad); 6.83 (m); 7.87 (m)
I- 349	C ₂ H ₅	H	R3- 5	R4-1	NH	Y-1	98°C

I- 350	CH ₃	H	R3- 88	R4-1	O	Y-1	δ = 1.95 (s); 3.53 (s); 3.85 (s); 4.05 (s); 4.50 (m); 5.65 (s); 5.90 (m); 7.28 (m); 7.50 (s)
I- 351	CH ₃	H	R3- 88	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.50 (s); 3.97 (s); 4.48 (m); 5.65 (m); 5.93 (m); 6.70 (broad); 7.28 (m); 7.50 (s)
I- 352	CH ₃	H	R3- 76	R4-1	NH	Y-1	δ = 1.95 (s); 2.27 (s); 2.87 (d); 3.95 (s); 4.63 (m); 5.93 (m); 6.66 (broad); 7.35 (m); 7.77 (m)
I- 353	CH ₃	H	R3- 187	R4-1	NH	Y-1	δ = 1.95 (s); 2.87 (d); 3.97 (s); 4.63 (m); 5.90 (d); 5.95 (m); 6.70 (broad); 7.48 (m); 7.85 (m); 7.97 (d)
I- 354	CH ₃	H	R3- 188	R4-1	NH	Y-1	δ = 1.93 (s); 2.77 (d); 3.93 (s); 4.54 (m); 5.92 (m); 6.80 (broad); 7.33 (m); 7.45 (m)
I- 355	CH ₃	H	R3- 189	R4-1	NH	Y-1	δ = 1.95 (s); 2.85 (d); 3.95 (s); 4.55 (m); 5.92 (m); 6.70 (broad); 7.23 (m); 7.40 (m); 7.64 (s); 7.78 (d)
I- 356	CH ₃	H	R3- 193	R4-1	NH	Y-1	δ = 1.94 (s); 2.87 (d); 3.94 (s); 4.44 (m); 5.17 (s); 5.67 (d); 5.88 (m); 6.70 (broad); 6.86 (m); 7.20 (m); 7.37 (d)
I-	CH ₃	H	R3-	R4-1	NH	Y-1	δ = 1.95 (s); 2.80 (d);

357			194				3.95 (s); 4.55 (m); 5.87 (d); 5.93 (m); 6.75 (broad); 7.25 (m); 7.35 (m); 7.45 (m); 7.55 (m); 7.70 (d)
I- 358	CH ₃	H	R3- 83	R4-1	NH	Y-1	δ = 1.95 (s); 2.90 (d); 3.95 (s); 4.57 (m); 5.92 (m); 6.13 (s); 6.83 (broad); 7.45 (m); 7.35 (m); 7.70 (m)
I- 359	CH ₃	H	R3- 195	R4-1	NH	Y-1	δ = 1.95 (s); 2.50 (s); 2.88 (d); 3.95 (s); 4.57 (m); 5.85 (d); 5.95 (m); 6.70 (broad); 6.90 (m); 7.60 (m); 8.33 (d)

* "phenyl ring" denotes that R¹ and R² together with the two carbon atoms linking them form a phenyl ring. Isobutyl = 2-methyl-1-propyl. m.p. = melting point; Rt = HPLC Retention time.

5

HPLC-data: RP-18 column (Chromolith Speed ROD 50x4.6 mm from Merck KgaA, Germany), 1.8 ml/min, injection volume 2 μ l, column temperature 40°C.

10 Eluent: Acetonitrile + 0,1 % trifluoroacetic acid (TFA) / water + 0,1 % TFA (gradient %:95 to 95:5 within 5 min), 40°C. MS: Quadrupole electrospray ionisation, 80 V (pos. mode).

II. Examples of the action against harmful fungi

15

The fungicidal action of the compounds of the formula I was demonstrated by the following experiments:

II.1 Microtiter tests

20

The active substances were formulated separately as a stock

solution in dimethyl sulfoxide (DMSO) at a concentration of 10 000 ppm.

Use example 1: Activity against the Septoria blotch pathogen caused by *Septoria tritici* in the microtiter test

Fungal strains used:

1. a) *Septoria tritici* (Qo-inhibitor sensitive, wild-type)
2. b) *Septoria tritici* (Qo-inhibitor-resistant, G143A mutant)

100 ml 2 % Malt extract in water at pH6.8 were inoculated with microspores from 2 week old cultures grown on 2 % malt extract + 2 % agar in Petri dishes and incubated for 3 days on a rotary shaker at 24°C und 150 rpm. The culture was harvested, glycerol was added (15 % (v/v) and kept frozen at -20°C in aliquots of 1 ml.

1 ml stock suspension was thawed and suspended into 800 ml of 2 % malt extract in water at pH 6.8. Compounds were diluted from stock solution in (dimethylsulfoxide) DMSO in 10 steps. The compound solutions were diluted 1/5 with sterile deionized water before use. 5 µl of the compound solutions were transferred into empty microplates. The plates were then filled with 195 µl of the microspore suspension of each strain.

The antifungal activity was determined by measuring the turbidity of a culture in 96-well microplates in the presence of test compounds. Fungal growth was measured by recording the optical density at 620 nm every 15 h for 150 h. The relative antifungal activity was calculated by comparison of the effect of the test compounds with the effect of a DMSO control and a standard fungicide.

IC₅₀-values (concentration of test compound resulting in 50% inhibition of fungal growth) were calculated from the resulting dose-response for each compound and strain. The initial

concentration of the test compounds and the 10 steps of dilution (1:4 each) allowed IC₅₀-values from 0.001 to 100 µmol/l (µM) to be assessed.

5 Table II.

Compound	Resistant <i>Septoria tritici</i> isolate (G143A mutation) (R- IC ₅₀) [µM]	Sensitive <i>Septoria tritici</i> isolate (wild type) (S- IC ₅₀) [µM]	Resistance factor RF = R-IC ₅₀ /S-IC ₅₀
Azoxystrobin	> 100	3.5	n.d.
Dimoxystrobin	> 100	7.1	n.d.
Enestroburin	> 100	4.5	n.d.
Kresoximmethyl	> 100	0.76	n.d.
Metominostrobin	> 100	> 100	n.d.
Orysastrobin	> 100	27	n.d.
Picoxystrobin	> 100	2.3	n.d.
Pyrametostrobin	> 100	> 100	n.d.
Pyraoxystrobin	> 100	1.6	n.d.
Pyraclostrobin	3.4	0.0012	2882
Trifloxystrobin	> 100	0.52	n.d.
I-5	2.2	0.083	26
I-6	8.3	0.34	24
I-82	6.8	1.1	6
I-83	7.6	1.6	4.6
I-84	16.1	0.5	33
I-85	22	0.8	29
I-90	26	1.0	26
I-91	27	1.6	17

I-94	12	0.6	22
I-95	29	2.2	14
I-97	49	1.7	29
I-191	37	1.3	29
I-211	1.1	0.44	2.6
I-212	8.3	1.4	6
I-213	0.057	0.076	0.7
I-215	48	1.7	28
I-216	12.0	6.3	2.0
I-217	63	6.3	10
I-218	79	5.6	14
I-221	10	3.4	3
I-223	19	2.7	7
I-224	5.4	1.7	3.2
I-225	72	3.4	21
I-228	10	3.4	3.0
I-230	25	3.2	8
I-233	2.7	0.24	12
I-234	4	2.4	1.7
I-235	51	13	4
I-236	24	6	4
I-238	12	6	1.9
I-243	15	4.1	3.6
I-244	1.3	0.6	2.0
I-247	9	5	1.9
I-249	11	2.4	5
I-250	97	19	5
I-255	42	19	2.2
I-256	24	21	1.1

I-257	67	15	4.5
I-260	3.3	2.3	1.4
I-261	5	2.9	1.6
I-262	1.6	1	1.6
I-265	4	3.8	1.1
I-267	5	2.9	1.7
I-269	74	8	10
I-270	57	6	10
I-271	7	1.9	3.9
I-273	22	5	4
I-274	19	9	2.0
I-287	18	1.5	12
I-288	28	3.5	8.0
I-291	18	1.0	18.0
I-292	33	7.9	4.2
I-298	31	3.6	8.6
I-303	4.6	0.27	17.0
I-304	9.1	0.53	17.2
I-305	32	9.3	3.4
I-307	67	19.0	3.5
I-310	11	0.39	28.2
I-319	102	17.0	6.0
I-324	38	1.9	20.0
I-326	18	1.2	15.0
I-331	25	1.8	13.9
I-339	21	1.0	21.0

While commercial strobilurine type fungicides are do not show any activity against the resistant *Septoria* strain, containing a mutation in the mitochondrial cytochrome b gene conferring

resistance to Qo inhibitors (G143A) while being active against sensitive wild type strains, compounds I have been active against both, the resistant and the wild-type strains. In general, the resistance factor ratio (RF) calculated from the IC₅₀ values determined for both Septoria strains, was below 30 for the compounds I according to the invention. However, resistance factor ratios for commercial strobilurine type compounds are in most cases greater than 100 and usually greater than several hundreds (for details see e.g. FRAC, Mutations associated with Qo1-resistance, Dec. 2006; <http://frac.info/frac/work/Mutations%20associated%20with%20Qo1%20resistance.pdf> and citations cited therein).


II.2 Comparative Examples

A) Field trials

Compounds used:

Compound I-6 was used as 50g/l EC formulation. Pyraclostrobin was used as commercial product HEADLINE.

Compounds used:



Pyraclostrobin

compound I-6

Trial 1: Efficacy against Septoria tritici on winter wheat

The trial was conducted under field conditions in Böhl-Iggelheim, Rhineland-palatinate, Germany. Seeds of winter wheat (cv. Riband) were planted and grown under standard conditions with adequate supply of water and nutrients. At growing stage GS 32 (April 14th, 2011), a first compound treatment (200 g a.i. per ha) was made with with a water volume of 400 L/ha, which was

repeated 21 days later at growing stage GS 39. No further fungicide treatments were applied. Infection with fungal pathogens (e.g. *Septoria tritici*) occurred naturally. The evaluation of the disease incidences for *Septoria tritici* 20, 33 and 46 days after the last treatment (DAA) are shown in table III. In the last row, the evaluation of the percentage of Qo inhibitor-resistant *Septoria tritici* isolates with the G143A mutation after the treatments is given.

Table III.

Treatment	Concentration (g a.i./ha)	Disease (%)			Percentage of G143A mutation (%) in <i>Septoria tritici</i> isolates
		20 DAA	33 DAA	46 DAA	
Compound I-6	200	1	6	36	100
Pyraclostrobin	200	8	32	93	100
untreated	-	13	47	96	65

In this test, the fungal pathogen *Septoria tritici* has been completely selected towards Qo inhibitor-resistant isolates by each of the treatments with the strobilurin-analogue compounds Pyraclostrobin and compound I-6. Due to this high resistance level, Pyraclostrobin showed insufficient control level although it has been used at commercial dose levels, whereas compound I-6 was capable to control the Qo inhibitor-resistant isolates of *Septoria tritici* with the G143 mutation.

Trial 2: Efficacy against *Septoria tritici* on winter wheat

This trial was conducted under field conditions in Limburgerhof, Rhineland-Palatinate, Germany. Seeds of winter wheat (cv. Riband) were planted and grown under standard conditions with adequate supply of water and nutrients. At growing stage GS 33 (April 8th, 2011), a first first compound treatment (200 g a.i. per ha) was made with a water volume of 400 L/ha, which was repeated 26 days later at growing stage GS 39. No further

fungicide treatments were applied. Infection with fungal pathogens (e.g. *Septoria tritici*) occurred naturally. The evaluation of the disease incidences for *Septoria tritici* 19 and 34 days after the last treatment (DAA) are shown in Table IV.

5 In the last row, the evaluation of the percentage of Qo inhibitor-resistant *Septoria tritici* isolates with the G143A mutation after the treatments is given.

Table IV.

Treatment	Concentration (g a.i./ha)	Disease (%)		Percentage of G143A mutation (%) in <i>Septoria tritici</i> isolates
		19 DAA	34 DAA	
Compound I-6	200	3	12	99
Pyraclostrobin	200	5	28	100
untreated	-	10	36	94

In this test, about the entire population of the fungal pathogen *Septoria tritici* has been Qo inhibitor-resistant (as evaluated at the end of the trial). Due to this high resistance level, Pyraclostrobin has shown antifungal activity only slightly above the untreated control although it has been used at commercial dose levels. However compound I-6 was capable to reduce the infection by Qo inhibitor-resistant *Septoria tritici* with the G143 mutation significantly.

B) Glass house trials

The spray solutions were prepared in several steps:

25 The stock solution was prepared as follows: 1.26 ml of a 1:1 mixture of cyclohexanone and dimethylsulfoxide was added to 8.4 mg of active ingredient. Next, 40.74 ml of a mixture of water, acetone (10%), the emulsifier Wettol (0.1 %) and the wetting agent Silwet (0.05%) was added. This stock solution was then
30 further diluted with the described solvent-emulsifier-water

mixture to the desired concentrations.

Trial 3: Control of leaf blotch on wheat caused by caused by two *Septoria tritici* isolates containing the G143A in the cytochrome b gene for their cytochrome bc₁ complex

Wheat plants were grown in pots. These plants were sprayed to run-off with an aqueous suspension, containing the desired concentration of active ingredient. The next day, the treated plants were inoculated with an aqueous suspension of *Septoria tritici*. After inoculation, the trial plants were covered with a lid and immediately transferred to a chamber with a relative humidity of about 83 to 85% and 19.5 to 20°C. After 4 days the lid was removed. Altogether, the trial plants were cultivated for about 28 days in that greenhouse chamber. The extent of fungal attack on the leaves was then visually assessed as % diseased leaf area.

Table VII.

Treatment	Conc. (ppm)	Resistant <i>Septoria tritici</i> isolate 1 (G143A mutation) Disease level (%)	Resistant <i>Septoria tritici</i> isolate 2 (G143A mutation) Disease level (%)
I-7	200	3	0
I-7	100	3	1
I-7	50	20	5
I-7	25	80	15
I-7	12.5	100	90
I-211	200	1	0
I-211	100	3	1
I-211	50	10	5
I-211	25	60	10
I-211	12.5	80	30

I-14	200	3	1
I-14	100	15	5
I-14	50	70	15
I-14	25	80	70
I-14	12.5	100	90
I-6	200	1	0
I-6	100	3	3
I-6	50	3	3
I-6	25	10	15
I-6	12.5	50	20
I-11	200	1	1
I-11	100	3	5
I-11	50	30	30
I-11	25	90	50
I-11	12.5	100	70
Pyraclostrobin	200	30	15
Pyraclostrobin	100	90	25
Pyraclostrobin	50	100	40
Pyraclostrobin	25	100	60
Pyraclostrobin	12.5	100	60
Trifloxystrobin	200	90	60
Trifloxystrobin	100	90	70
Trifloxystrobin	50	90	70
Trifloxystrobin	25	100	70
Trifloxystrobin	12.5	100	80

III. Molecular modeling

III.1 Structural Models of Wild-Type and G143A mutant binding site

Structural models of the binding site of wild-type and G143A cytochrome bc₁ complex were generated based on the crystallographic structure of bovine cytochrome bc₁ complex with azoxystrobin bound to the Qo-site (PDB: 1SQB: Esser et al. J Mol Biol 341, 281-302 (2004)).

The structure was imported into Schrödinger Maestro (version 9.0, Schrödinger, LLC, New York, NY, 2009).

Cytochrome b was isolated from the structure of the complex and treated with the Schrödinger Protein Preparation Wizard (Schrödinger Suite 2009 Protein Preparation Wizard; Epik version 2.0, Schrödinger, LLC, New York, NY, 2009; Impact version 5.5, Schrödinger, LLC, New York, NY, 2009, Prime version 2.1, Schrödinger, LLC, New York, NY, 2009).

This structure was used as the model for the wild-type binding site without further changes.

The model of the G143A mutant was generated by changing a hydrogen in Glycine 143 of the wild-type model into a methyl group, thereby generating S-Alanine, using Schrödinger Maestro. Amino acids in a sphere of 5 Å around the co-crystallized molecule of azoxystrobin were energy-minimized using Schrödinger MacroModel (version 9.7, Schrödinger, LLC, New York, NY, 2009).

III.2 Molecular Docking

Structures of inhibitors were prepared for docking with Schrödinger LigPrep (version 2.3, Schrödinger, LLC, New York, NY, 2009) and docked into the structural models using Schrödinger Glide (version 5.5, Schrödinger, LLC, New York, NY, 2009).

III.3 Graphics of inhibitors bound to cytochrome bc₁ complex

Graphical representations of poses from the docking runs were

generated using Molecular Operating Environment (MOE; 2010.10; Chemical Computing Group Inc., 1010 Sherbooke St. West, Suite #910, Montreal, QC, Canada, H3A 2R7, 2010) and refined using the GNU Image Manipulation Program (GIMP, version 2.6.8, 2008).

5

As illustrated in Fig. 1 molecular modelling of an artificial Qo inhibitor-resistant cytochrome bc₁ complex with the mutation G143A (see below for details) has been carried out. Docking of the commercial strobilurine analogue compound pyraclostrobin shows the steric clash resulting in impaired binding of this active ingredient in the G143A mutant cytochrome bc₁ complex. The compounds of the present invention largely avoid this steric clash by either replacing the central phenyl ring of pyraclostrobin with a smaller and/or more flexible two carbon unit which may be suitably substituted or by replacing the well known pharmacophores with the smaller tetrazolinone moiety R4-7.

10

15

20

III.4 Number of Van-der-Waals clashes between inhibitors and Alanine G143A

25

30

The docked ligands were transferred to the G143A binding site model keeping the coordinates from docking into the wild-type model. Thereby, complexes between the G143A binding site and ligands were formed. Those were used as starting structures for the following energy minimization using Schrödinger MacroModel. During these simulations, only the ligand was allowed to move freely, the protein was considered "frozen". The poses generated by this procedure were considered as the relaxed state of the inhibitors after introduction of the G143A mutation.

35

Two atoms are considered to be sterically clashing, if the distance between their centers is shorter than 0.9 times the sum of their Van-der-Waals radii. For the atoms pairs relevant for strobilurin interaction with Alanine 143 the respective distances are described in Table V.

Table V.

Atom 1	vdW-Radius 1	Atom 2	vdW-Radius 2	vdW-Sum	Clash Distance
C	1,70	C	1,70	3,40	3,06
C	1,70	N	1,55	3,25	2,93
C	1,70	O	1,52	3,22	2,90
C	1,70	H	1,20	2,90	2,61
H	1,20	N	1,55	2,75	2,48
H	1,20	O	1,52	2,72	2,45

After minimization in the G143A binding site as described above, the number of steric Van-der-Waals clashes with Alanine 143 was counted for each inhibitor. The results are given in the Table VI.


Table VI.

Compound	No. of clashes
I-5	3
I-6	3
Pyrametostrobin	6
Metominostrobin	6
Azoxystrobin	7
Enestroburin	7
Pyraclostrobin	7
Dimoxystrobin	7
Orysastrobin	7
Pyraoxystrobin	8
Picoxystrobin	8
Trifloxystrobin	8
Kresoxim-Methyl	8

It was found that compounds I have a small number of steric Van-
der-Waals clashes and also have unexpectedly high activity
against Qo inhibitor-resistant fungal strains harboring said
G143A mutation in microtiter titer tests as well as in field
5 trials on sites which have a high portion of Qo inhibitor-
resistant fungal strains harboring said G143A mutation (see
below).

Patentkrav

1. Anvendelse af forbindelser med formel I



5 hvor:

R¹ og R² uafhængigt af hinanden er hydrogen, halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl eller C₃-C₆-cycloalkyl-C₁-C₄-alkyl, hvor grupperne R¹ og R² are cis-orienterede, eller

10 R¹ og R² sammen med de to carbonatomer, der kobler dem, danner en phenylring,

forudsat at R⁴ er 4-methyl-1,4-dihydropyridazin-5-on-1-yl (formel R4-7), og

15 hvor de alifatiske dele af R¹ og/eller R² eller ovennævnte phenylring kan bære 1, 2, 3 eller op til det maksimale antal af identiske eller forskellige grupper R^a, som uafhængigt af hinanden er udvalgt blandt:

R^a, halogen, CN, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl og C₁-C₄- haloalkoxy;

20 Y er en direkte binding eller en divalent gruppe, der er udvalgt blandt -OCH₂-, -CH₂-, -CH₂CH₂-, -C(Z)=N-O-CH₂-, -CHZ-C(Z)=N-O-CH₂-, -O-N=C(Z)-C(Z)=N-O-CH₂-, -C(=O)-C(Z)=N-O-CH₂- og -C(=N-O-Z)-C(Z)=N-O-CH₂-,

25 hvor bindingen på venstre side af den divalente gruppe Y er bundet til R³, og bindingen på højre side er bundet til carbonatomet, der substitueres med R², og

Z, der kan være ens eller forskellig i forhold til eventuelle andre Z, er hydrogen, C₁-C₄-alkyl eller C₁-C₄-haloalkyl;

30 R³ er phenyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, hvor ringelementatomerne i heterocyclyl'en ud over carbonatomer indbefatter 1, 2, 3 eller 4 heteroatomer, der er udvalgt fra gruppen, der består af N, O og S;

35 hvor de cykliske grupper R³ kan bære 1, 2, 3, 4 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^b, som uafhængigt af hinanden er udvalgt blandt:

R^b, der kan være ens eller forskellig i forhold til eventuelle

andre R^b og er amino, halogen, hydroxyl, oxo, nitro, CN, carboxyl, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₂-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₆-alkoxyimino-C₁-C₄-alkyl, C₂-C₆-alkenyloxyimino-C₁-C₄-alkyl, C₂-C₆-alkynyloxyimino-C₁-C₄-alkyl, C₁-C₄-alkylamino, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylcarbonyloxy, phenyl, naphthyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, som foruden carbonatomer indeholder et til fire heteroatomer fra gruppen, der består af N, O og S, som ringelementer; og hvor ovennævnte phenyl- og heterocyclylgrupper R^b er bundet via en direkte binding, et oxygen- eller svovlatom, og to radikaler R^b, der er bundet til tilgrænsende ringelementatomer i den cykliske gruppe R³, sammen med ringelementatomerne kan danne en fusioneret 5-, 6- eller 7-leddet, mættet, delvist umættet eller aromatisk cyklus, som kan være en carbocyklus eller heterocyklus, hvor ringelementatomerne i heterocyklussen ud over carbonatomer indbefatter 1, 2, 3 eller 4 heteroatomer, der er udvalgt fra gruppen, der består af N, O og S,

og

hvor de alifatiske eller cykliske grupper R^b for deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige grupper

R^c, der kan være ens eller forskellig i forhold til eventuelle andre R^c og er halogen, hydroxyl, nitro, CN, carboxyl, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₈-alkynyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₆-alkoxyimino-C₁-C₄-alkyl, C₂-C₆-alkenyloxyimino-C₁-C₄-alkyl, C₂-C₆-alkynyloxyimino-C₁-C₄-alkyl, C₁-C₆-alkoxyimino-, C₂-C₆-alkenyloxyimino-, C₂-C₆-alkynyloxyimino-, C₂-C₆-haloalkenyloxyimino-, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, phenyl eller en 5-leddet, mættet, delvist umættet eller aromatisk heterocyclyl, som foruden carbonatomer indeholder et til tre heteroatomer fra gruppen, der består af N, O og S, som ringelementer; hvor ovennævnte cykliske grupper R^c er bundet via en direkte binding, et oxygen- eller svovlatom, og hvor de alifatiske eller cykliske grupper R^c for deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af

identiske eller forskellige grupper R^d :

R^d , der kan være ens eller forskellig i forhold til eventuelle andre R^d og er halogen, C_1 - C_4 -alkyl eller C_1 - C_4 -haloalkyl; eller

5 R^3 er $-CR^A=N-O-R^B$, hvor

R^A er amino, hydroxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl,


10 C_1 - C_4 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_4 -alkyl, naphthyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, som foruden carbonatomer indeholder et til fire heteroatomer fra gruppen, der består af O, N og S, som ringelementer; og hvor ovennævnte
15 cykliske R^A er bundet via en direkte binding, et oxygen- eller svovlatom;

R^B er C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxycarbonyl, phenyl, phenyl- C_1 - C_4 -alkyl, naphthyl
20 eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, som foruden carbonatomer indeholder et til fire heteroatomer fra gruppen, der består af O, N og S, som ringelementer;

hvor de alifatiske eller cykliske grupper R^A og/eller R^B for
25 deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^e :

R^e , der kan være ens eller forskellig i forhold til eventuelle andre R^e , er halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy eller
30 C_1 - C_4 -haloalkoxy;

R^4 er en monovalent gruppe, der er udvalgt blandt formlerne R4-1 til R4-7



hvor den takkede linje definerer påhæftningspunktet, og
X er en direkte binding eller en divalent gruppe CH₂, O eller NH,

- 5 R⁵ er C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy eller C₃-C₆-cycloalkyl,
R⁶ er C₁-C₄-alkyl eller C₁-C₄-haloalkyl;
og N-oxiderne og de landbrugsmæssigt acceptable salte deraf til
bekæmpelse af fytopatogene svampe, der indeholder en mutation i
10 det mitokondriske cytochrom b-gen, der bibringer resistens over
for Qo-hæmmere, hvor mutationen er G143A.

2. Anvendelse ifølge krav 1, hvor de fytopatogene svampe er
udvalgt fra gruppen, der består af *Alternaria alternata*,
15 *Blumeria graminis*, *Pyricularia oryzae* (også kaldet *Magnaporthe grisea*),
Septoria tritici (også kaldet *Mycosphaerella graminicola*),
Mycosphaerella fijiensis, *Venturia inaequalis*,
Pyrenophora teres, *Pyrenophora tritici-repentis* og *Plasmopara viticola*.

- 20 3. Anvendelse ifølge et hvilket som helst af kravene 1 til 2,
hvor den fytopatogene svamp er *Septoria tritici*.

4. Fremgangsmåde til bekæmpelse af fytopatogene svampe ifølge
25 et hvilket som helst af kravene 1 til 3, der omfatter:
behandling af de fytopatogene svampe ifølge et hvilket som helst
af kravene 1 til 3 eller materialer, planter, jord eller frø,
der er i risiko for at være inficeret med fytopatogene svampe
ifølge et hvilket som helst af kravene 1 til 3, med en effektiv
30 mængde af mindst én forbindelse med formel I ifølge krav 1 eller
en sammensætning, der omfatter denne.


5. Fremgangsmåde ifølge krav 4, der omfatter:

a) identificering af de fytopatogene svampe ifølge et hvilket som helst af kravene 1 til 3 eller materialer, planter, jord eller frø, der er i risiko for at være inficeret med fotopatogene svampe ifølge et hvilket som helst af kravene 1 til 3,

og

b) behandling af svampene eller materialerne, planterne, jorden eller frøene med en effektiv mængde af mindst én forbindelse med formel I ifølge krav 1 eller en sammensætning, der omfatter denne.

6. Forbindelser med formel I



hvor:

R¹ og R² uafhængigt af hinanden er hydrogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkenyloxy, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl eller C₃-C₆-cycloalkyl-C₁-C₄-alkyl, hvor grupperne R¹ og R² er cis-orienterede,

hvor de alifatiske dele af R¹ og/eller R² kan bære 1, 2, 3 eller op til det maksimale antal af identiske eller forskellige grupper R^a, som uafhængigt af hinanden er udvalgt blandt:

R^a, CN, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl og C₁-C₄-haloalkoxy;

Y er en direkte binding eller en divalent gruppe, der er udvalgt blandt -OCH₂-, -CH₂-, -CH₂CH₂-, -C(Z)=N-O-CH₂-, -CHZ-C(Z)=N-O-CH₂-, -O-N=C(Z)-C(Z)=N-O-CH₂-, -C(=O)-C(Z)=N-O-CH₂- og -C(=N-O-Z)-C(Z)=N-O-CH₂-,

hvor bindingen på venstre side af den divalente gruppe Y er bundet til R³, og bindingen på højre side er bundet til carbonatomet, der substitueres med R², og

Z, der kan være ens eller forskellig i forhold til eventuelle andre Z, er C₁-C₄-alkyl eller C₁-C₄-haloalkyl;

R³ er phenyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, hvor ringelementatomerne i heterocyclyl'en ud over carbonatomer indbefatter 1, 2, 3 eller 4 heteroatomer, der er udvalgt fra gruppen, der består af N, O og S;

hvor de cykliske grupper R^3 kan bære 1, 2, 3, 4 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^b , som uafhængigt af hinanden er udvalgt blandt:

R^b , der kan være ens eller forskellig i forhold til eventuelle andre R^b og er amino, halogen, hydroxyl, oxo, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_1 - C_6 -alkoxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkenyloxyimino- C_1 - C_4 -alkyl, C_2 -
 10 C_6 -alkynyloxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylcarbonyloxy, phenyl, naphthyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, som foruden carbonatomer indeholder et til fire heteroatomer fra gruppen, der består af
 15 N, O og S, som ringelementer; og hvor ovennævnte phenyl- og heterocyclylgrupper R^b er bundet via en direkte binding, et oxygen- eller svovlatom;

og

to radikaler R^b , der er bundet til tilgrænsende ringelementatomer
 20 i den cykliske gruppe R^3 , sammen med ringelementatomerne kan danne en fusioneret 5-, 6- eller 7-leddet, mættet, delvist umættet eller aromatisk cyklus, som kan være en carbocyklus eller heterocyklus, hvor ringelementatomerne i den fusionerede heterocyklus ud over carbonatomer indbefatter 1, 2, 3 eller 4
 25 heteroatomer, der er udvalgt fra gruppen, der består af N, O og S, og

hvor de alifatiske eller cykliske grupper R^b for deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^c :

30 hvor R^c , der kan være ens eller forskellig i forhold til eventuelle andre R^c , er halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_6 -alkoxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkenyloxyimino- C_1 - C_4 -alkyl, C_2 - C_6 -alkynyloxyimino- C_1 - C_4 -alkyl,
 35 C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxyimino-, C_2 - C_6 -alkenyloxyimino-, C_2 - C_6 -alkynyloxyimino-, C_2 - C_6 -haloalkenyloxyimino-, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, phenyl eller en 5-leddet, mættet, delvist umættet eller

aromatisk heterocyclyl, som foruden carbonatomer indeholder et til tre heteroatomer fra gruppen, der består af N, O og S, som ringelementer; hvor ovennævnte cykliske grupper R^c er bundet via en direkte binding, et oxygen- eller svovlatom, og hvor de

5 alifatiske eller cykliske grupper R^c for deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^d :

hvor R^d , der kan være ens eller forskellig i forhold til eventuelle andre R^d , er halogen, C_1 - C_4 -alkyl eller C_1 - C_4 -haloalkyl;

10 eller

R^3 er $-CR^A=N-O-R^B$, hvor

R^A er amino, hydroxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, C_2 - C_6 -alkenyloxy, C_3 - C_6 -alkynyloxy, C_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkylamino, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylcarbonyloxy, phenyl, phenyl- C_1 - C_4 -alkyl, naphthyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, som foruden

20 carbonatomer indeholder et til fire heteroatomer fra gruppen, der består af O, N og S, som ringelementer; og hvor ovennævnte cykliske R^A er bundet via en direkte binding, et oxygen- eller svovlatom;

R^B er C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -haloalkyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -cycloalkenyl, CC_1 - C_4 -alkoxyimino- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxycarbonyl, phenyl, phenyl- C_1 - C_4 -alkyl, naphthyl eller en 3- til 10-leddet, mættet, delvist umættet eller aromatisk mono- eller bicyklisk heterocyclyl, som foruden carbonatomer indeholder et til fire heteroatomer fra gruppen,

30 der består af O, N og S, som ringelementer;


hvor de alifatiske eller cykliske grupper R^A og/eller R^B for deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^e :

hvor R^e , der kan være ens eller forskellig i forhold til eventuelle andre R^e , er halogen, hydroxyl, nitro, CN, carboxyl, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy eller C_1 - C_4 -haloalkoxy;

35

R^4 er en monovalent gruppe, der er udvalgt blandt formlerne R_4 -

1 til R4-7



hvor den takkede linje definerer påhæftningspunktet, og hvor
X er en divalent gruppe O eller NH,

5 R⁵ er C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy
eller C₃-C₆-cycloalkyl,

R⁶ er C₁-C₄-alkyl eller C₁-C₄-haloalkyl;

og N-oxiderne og de landbrugsmæssigt acceptable salte deraf,
bortset fra (2E,3Z)-2-(ethyliden)-5-phenyl-3-

10 pentensyremethylester (CAS-nr. 681026-20-0).

7. Forbindelser ifølge krav 6, hvor R¹ og R² uafhængigt af
hinanden er hydrogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₂-C₆-alkenyl,
C₂-C₆-alkenyloxy, C₂-C₆-alkynyl, C₃-C₆-cycloalkyl eller C₃-C₆-
15 cycloalkyl-C₁-C₄-alkyl.

8. Forbindelser ifølge et hvilket som helst af kravene 6 til
7, hvor R¹ er hydrogen, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkoxy-
C₁-C₄-alkyl eller C₁-C₄-haloalkoxy-C₁-C₄-alkyl.

20

9. Forbindelser ifølge krav 8, hvor R¹ er methyl eller ethyl.

10. Forbindelser ifølge et hvilket som helst af kravene 6 til
9, hvor R² er hydrogen.

25

11. Forbindelser ifølge et hvilket som helst af kravene 6 til
10, hvor Y er -OCH₂-, -CH₂CH₂-, -C(CH₃)=N-O-CH₂-, -O-N=C(CH₃)-
C(CH₃)=N-O-CH₂- eller -C(=N-O-CH₃)-C(CH₃)=N-O-CH₂-.

30 12. Forbindelser ifølge et hvilket som helst af kravene 6 til
11, hvor R³ er pyrazolyl, imidazolyl, oxazolyl, isoxazolyl,
thiazolyl, 1,2,4-triazolyl, 1,2,4-oxadiazolyl, 1,3,4-

oxadiazolyl eller 1,2,4-thiadiazolyl, der er substitueret med phenyl, hvor phenyl'en kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige grupper R^c:

- 5 hvor R^c, der kan være ens eller forskellige i forhold til eventuelle andre R^c, er halogen, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkoxyimino-C₁-C₄-alkyl, phenyl eller en 5-leddet, mættet, delvist umættet eller aromatisk heterocyclyl, som foruden carbonatomer
 10 indeholder et til tre heteroatomer fra gruppen, der består af N, O og S, som ringelementer; og hvor ovennævnte heterocyclylgrupper R^c er bundet via en direkte binding, et oxygen- eller svovlatom og for deres del kan bære 1, 2, 3 eller op til det maksimale mulige antal af identiske eller forskellige
 15 grupper R^d.

13. Forbindelser ifølge et hvilket som helst af kravene 6 til 12, hvor R⁴ er -C(=NOCH₃)-CONHCH₃, -C(=NOCH₃)-COOCH₃, -C(=CHOCH₃)-COOCH₃, -C(=CHOCH₃)-CONHCH₃, -N(OCH₃)-COOCH₃, -N(CH₃)-COOCH₃ eller
 20 -N(CH₂CH₃)-COOCH₃.

14. Agrokemiske sammensætninger, hvilke sammensætninger omfatter et hjælpstof og mindst én forbindelse med formel I ifølge et hvilket som helst af kravene 6 til 13, et N-oxid eller
 25 et landbrugsmæssigt acceptabelt salt deraf.

15. Sammensætninger ifølge krav 14, der omfatter mindst ét yderligere aktivt stof.

- 30 16. Sammensætninger ifølge krav 15, hvor det yderligere aktive stof er udvalgt blandt klasserne A) til O):

A) Respirationshæmmere

- Hæmmere af kompleks III i Q_o-bindingsstedet (f.eks. strobiluriner): azoxystrobin, coumethoxystrobin,
 35 coumoxystrobin, dimoxystrobin, enestroburin, fenaminstrobin, fenoxystrobin/flufenoxystrobin, fluoxastrobin, kresoximmethyl, metominostrobin, orysastrobin, picoxystrobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, trifloxystrobin, 2-[2-(2,5-

dimethylphenoxyethyl)-phenyl]-3-methoxyacrylsyremethylester
og

2-(2-(3-(2,6-dichlorophenyl)-1-methyl-
allylidenaminoxyethyl)-phenyl)-2-methoxyimino-N-

methylacetamid, pyribencarb, triclopyricarb/chlordin-carb,

5 famoxadon, fenamidon;

- hæmmere of kompleks III i Q₁-bindingsstedet: cyazofamid,
amisulbrom,

[(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxy-4-
methoxypyridin-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-

dioxonan-7-yl]-2-methylpropanoat, [(3S,6S,7R,8R)-8-benzyl-3-

10 [[3-(acetoxymethoxy)-4-methoxypyridin-2-carbonyl]amino]-6-

methyl-4,9-dioxo-1,5-dioxonan-7-yl]-2-methylpropanoat,

[(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutoxycarbonyloxy-4-

methoxypyridin-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-

dioxonan-7-yl]-2-methylpropanoat, [(3S,6S,7R,8R)-8-benzyl-3-

15 [[3-(1,3-benzodioxol-5-ylmethoxy)-4-methoxypyridin-2-

carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl]-2-

methylpropanoat; (3S,6S,7R,8R)-3-[[3-hydroxy-4-methoxy-2-

pyridinyl)carbonyl]amino]-6-methyl-4,9-dioxo-8-(phenylmethyl)-

1,5-dioxonan-7-yl]-2-methylpropanoat;

20 - hæmmere of kompleks II (f.eks. carboxamider): benodanil,

benzovindiflupyr, bixafen, boscalid, carboxin, fenfuram,

fluopyram, flutolanil, fluxapyroxad, furametpyr, isopyrazam,

mepronil, oxycarboxin, penflufen, penthiopyrad, sedaxan,

tecloftalam, thifluzamid, N-(4'-trifluormethylthiobiphenyl-2-

25 yl)-3-difluormethyl-1-methyl-1H-pyrazol-4-carboxamid, N-(2-

(1,3,3-trimethylbutyl)-phenyl)-1,3-dimethyl-5-fluor-1H-

pyrazol-4-carboxamid, 3-(difluormethyl)-1-methyl-N-(1,1,3-

trimethylindan-4-yl)pyrazol-4-carboxamid, 3-(trifluormethyl)-

1-methyl-N-(1,1,3-trimethylindan-4-yl)pyrazol-4-carboxamid,

30 1,3-dimethyl-N-(1,1,3-trimethylindan-4-yl)pyrazol-4-

carboxamid, 3-(trifluormethyl)-1,5-dimethyl-N-(1,1,3-

trimethylindan-4-yl)-pyrazol-4-carboxamid, 3-(difluormethyl)-

1,5-dimethyl-N-(1,1,3-trimethylindan-4-yl)pyrazol-4-

carboxamid, 1,3,5-trimethyl-N-(1,1,3-trimethylindan-4-

35 yl)pyrazol-4-carboxamid;

- andre respirationshæmmere (f.eks. kompleks I, afkoblere):

diflumetorim, (5,8-difluorquinazolin-4-yl)-{2-[2-fluor-4-(4-

trifluormethylpyridin-2-yloxy)-phenyl]-ethyl}-amin;

nitrophenylderivater: binapacryl, dinobuton, dinocap, fluazinam; ferimzon;

organometalforbindelser: fentinsalte, såsom fentinacetat, fentinchlorid eller fentinhydroxid; ametoctradin; og
5 silthiofam;

B) Sterolbiosyntesehæmmere (SBI-fungicider)

- Hæmmere af C14-demethylase (DMI-fungicider): triazoler:

azaconazol, bitertanol, bromuconazol, cyproconazol, difenoconazol, diniconazol, diniconazol-M, epoxiconazol,

10 fenbuconazol, fluquinconazol, flusilazol, flutriafol, hexaconazol, imibenconazol, ipconazol, metconazol,

myclobutanil, oxpoconazol, paclobutrazol, penconazol, propiconazol, prothioconazol, simeconazol, tebuconazol,

tetraconazol, triadimefon, triadimenol, triticonazol,

15 uniconazol, 1-[rel-(2S;3R)-3-(2-chlorphenyl)-2-(2,4-

difluorphenyl)-oxiranylmethyl]-5-thiocyanat-1H-[1,2,4]triazol,

2-[rel-(2S;3R)-3-(2-chlorphenyl)-2-(2,4-difluorphenyl)-

oxiranylmethyl]-2H-[1,2,4]triazol-3-thiol; imidazoler:

imazalil, pefurazoat, prochloraz, triflumizol; pyrimidiner,

20 pyridiner og piperaziner: fenarimol, nuarimol, pyrifenox,

triforin;

- Hæmmere af delta14-reduktase: aldimorph, dodemorph,

dodemorphacetat, fenpropimorph, tridemorph, fenpropidin,

piperalin, spiroxamin;

25 - Hæmmere af 3-ketoreduktase: fenhexamid;

C) Nukleinsyresyntesehæmmere

- phenylamider eller acylaminosyrefungicider: benalaxyl,

benalaxyl-M, kiralaxyl, metalaxyl, metalaxyl-M (mefenoxam),

ofurac, oxadixyl;

30 - andre: hymexazol, octhilinon, oxolinsyre, bupirimat, 5-

fluorcytosin, 5-fluor-2-(p-tolylmethoxy)pyrimidin-4-amin, 5-

fluor-2-(4-fluorphenylmethoxy)pyrimidin-4-amin;

D) Hæmmere af celledeling og cytoskelet

- tubulinhæmmere, såsom benzimidazoler, thiophanater: benomyl,

35 carbendazim, fuberidazol, thiabendazol, thiophanatmethyl;

triazolopyrimidiner: 5-chlor-7-(4-methylpiperidin-1-yl)-6-

(2,4,6-trifluorphenyl)-[1,2,4]triazol[1,5-a]pyrimidin;

- andre celledelingshæmmere: diethofencarb, ethaboxam,

pencycuron, fluopicolid, zoxamid, metrafenon, pyriofenon;

E) Hæmmere af aminosyre- og proteinsyntese

- methioninsyntesehæmmere (anilinpyrimidiner): cyprodinil, mepanipyrin, pyrimethanil;

5 - proteinsyntesehæmmere: blasticidin-S, kasugamycin, kasugamycin-hydrochloridhydrat, mildiomicin, streptomycin, oxytetracyclin, polyoxin, validamycin A;

F) Signaltransduktionshæmmere

10 - MAP/histidinkinasehæmmere: fluoroimid, iprodion, procymidon, vinclozolin, fenpiclonil, fludioxonil;

- G-proteinhæmmere: quinoxifen;

G) Lipid- og membransyntesehæmmere

- Phospholipidbiosyntesehæmmere: edifenphos, iprobenfos, pyrazophos, isoprothiolan;

15 - lipidperoxidation: dicloran, quintozen, tecnazen, tolclofosmethyl, biphenyl, chlorneb, etridiazol;

- phospholipidbiosyntese og cellevægsdeponering: dimethomorph, flumorph, mandipropamid, pyrimorph, benthiavalicarb, iprovalicarb, valifenalat og N-(1-(1-(4-

20 cyanophenyl)ethansulfonyl)-but-2-yl)-carbaminsyre-(4-fluorphenyl)-ester;

- forbindelser, der påvirker cellemembranpermeabilitet og fedtsyrer: propamocarb, propamocarbhydrochlorid;

25 - fedtsyreamidhydrolasehæmmere: 1-[4-[4-[5-(2,6-difluorphenyl)-4,5-dihydro-3-isoxazolyl]-2-thiazolyl]-1-piperidinyl]-2-[5-methyl-3-(trifluormethyl)-1H-pyrazol-1-yl]ethanon;

H) Hæmmere med flere aktivitetssteder

- uorganiske aktive stoffer: Bordeaux-blanding, kobberacetat, kobberhydroxid, kobberoxychlorid, basisk kobbersulfat, svovl;

30 - thio- og dithiocarbamater: ferbam, mancozeb, maneb, metam, metiram, propineb, thiram, zineb, ziram;

- organochlorforbindelser (f.eks. phthalimider, sulfamider, chlornitriler): anilazin, chlorthalonil, captafol, captan, folpet, dichlofluand, dichlorphen, flusulfamid,

35 hexachlorbenzen, pentachlorphenol og dens salte, phthalid, tolylfluand, N-(4-chlor-2-nitrophenyl)-N-ethyl-4-methylbenzensulfonamid;

- guanidiner og andre: guanidin, dodin, dodin fri base,

guazatin, guazatincetat, iminoctadin, iminoctadintriacetat, iminoctadintris(albesilat), dithianon, 2,6-dimethyl-1H,SH-[1,4]dithiino[2,3-c:5,6-c']dipyrrol-1,3,5,7(2H,6H)-tetraon;

I) Cellevægssyntesehæmmere

- 5 - hæmmere af glucansyntese: validamycin, polyoxin B; melaninsyntesehæmmere: pyroquilon, tricyclazol, carpropamid, dicyclomet, fenoxanil;

J) Planteforsvarsinduktorer

- acibenzolar-S-methyl, probenazol, isotianil, tiadinil, 10 prohexadioncalcium; fosfonater: fosetyl, fosetylaluminum, phosphorsyre og dens salte;

K) Ukendt virkningsmekanisme

- bronopol, chinomethionat, cyflufenamid, cymoxanil, dazomet, debacarb, diclomezin, difenzoquat, difenzoquatmethylsulfat, 15 diphenylamin, fenpyrazamin, flumetover, flusulfamid, flutianil, methasulfocarb, nitrapyrin, nitrothalisopropyl, oxathiapiprolin, oxinkobber, proquinazid, tebufloquin, tecloftalam, triazoxid, 2-butoxy-6-iod-3-propylchromen-4-on, N-(cyclopropylmethoxyimino-(6-difluormethoxy-2,3-difluorphenyl)-methyl)-2-phenylacetamid, N'-(4-(4-chlor-3-trifluormethylphenoxy)-2,5-dimethylphenyl)-N-ethyl-N-methylformamidin, N'-(4-(4-fluor-3-trifluormethylphenoxy)-2,5-dimethylphenyl)-N-ethyl-N-methylformamidin, N'-(2-methyl-5-trifluormethyl-4-(3-trimethylsilanylpropoxy)-phenyl)-N-ethyl-N-methylformamidin, N'-(5-difluormethyl-2-methyl-4-(3-trimethylsilanylpropoxy)-phenyl)-N-ethyl-N-methylformamidin, methoxyeddikesyre-6-tert-butyl-8-fluor-2,3-dimethylquinolin-4-ylester, 3-[5-(4-methylphenyl)-2,3-dimethylisoxazolidin-3-yl]-pyridin, 3-[5-(4-chlorphenyl)-2,3-dimethylisoxazolidin-3-yl]-pyridin-(pyrisoxazol), N-(6-methoxypyridin-3-yl)-cyclopropancarboxylsyreamid, 5-chlor-1-(4,6-dimethoxypyrimidin-2-yl)-2-methyl-1H-benzoimidazol, 2-(4-chlorphenyl)-N-[4-(3,4-dimethoxyphenyl)-isoxazol-5-yl]-2-prop-2-ynyloxyacetamid;

- 35 L) Antifungale biologiske bekæmpelsesmidler, plantebioaktivatorer
Ampelomyces quisqualis, Aspergillus flavus, Aureobasidium pullulans, Bacillus pumilus, Bacillus subtilis, Bacillus

subtilis var. amyloliquefaciens FZB24, Candida oleophila I-82, Candida saitoana, Chitosan, Clonostachys rosea f. catenulata også kaldet Gliocladium catenulatum, Coniothyrium minitans, Cryphonectria parasitica, Cryptococcus albidus, Fusarium oxysporum, Metschnikowia fructicola, Microdochium dimerum, Phlebiopsis gigantea, Pseudozyma flocculosa, Pythium oligandrum DV74, Reynoutria sachlinensis, Talaromyces flavus V117b, Trichoderma asperellum SKT-1, T. atroviride LC52, T. harzianum T-22, T. harzianum TH 35, T. harzianum T-39, T. harzianum og T. viride, T. harzianum ICC012 og T. viride ICC080, T. polysporum og T. harzianum, T. stromaticum, T. virens GL-21, T. viride, T. viride TV1, Ulocladium oudemansii HRU3;

M) Vækstregulerende midler

abscisinsyre, amidochlor, ancymidol, 6-benzylaminopurin, brassinolid, butralin, chlormequat (chlormequatchlorid), cholinchlorid, cyclanilid, daminozid, dikegulac, dimethipin, 2,6-dimethylpuridin, ethephon, flumetralin, flurprimidol, fluthiacet, forchlorfenuron, gibberellinsyre, inabenfid, indol-3-eddikesyre, maleinhydrazid, mefluidid, mepiquat (mepiquatchlorid), naphthaleneddikesyre, N-6-benzyladenin, paclobutrazol, prohexadion (prohexadioncalcium), prohydrojasmon, thidiazuron, triapenthenol, tributylphosphortrithioat, 2,3,5-triiodbenzoesyre, trinexapacethyl og uniconazol;

N) Herbicider

- acetamider: acetochlor, alachlor, butachlor, dimethachlor, dimethenamid, flufenacet, mefenacet, metolachlor, metazachlor, napropamid, naproanilid, pethoxamid, pretilachlor, propachlor, thenylchlor;

- aminosyrederivater: bilanafos, glyphosat, glufosinat, sulfosat;

- aryloxyphenoxypropionater: clodinafop, cyhalofopbutyl, fenoxaprop, fluazifop, haloxyfop, metamifop, propaquizafop, quizalofop, quizalofop-P-tefuryl;

- Bipyridyler: diquat, paraquat;

- (thio)carbamater: asulam, butylat, carbetamid, desmedipham, dimepiperat, eptam (EPTC), esprocarb, molinat, orbencarb, phenmedipham, prosulfocarb, pyributicarb, thiobencarb,

triallat;

- cyclohexandioner: butroxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxym, tralkoxydim;

- dinitroaniliner: benfluralin, ethalfluralin, oryzalin, pendimethalin, prodiamin, trifluralin;

- diphenylethere: acifluorfen, aclonifen, bifenox, diclofop, ethoxyfen, fomesafen, lactofen, oxyfluorfen;

- hydroxybenzonitriler: bomoxynil, dichlobenil, ioxynil;

- imidazolinoner: imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr;

- phenoxyeddikesyrer: clomeprop, 2,4-dichlorphenoxyeddikesyre (2,4-D), 2,4-DB, dichlorprop, MCPA, MCPA-thioethyl, MCPB, Mecoprop;

- pyraziner: chloridazon, flufenpyrethyl, fluthiacet, norflurazon, pyridat;

- pyridiner: aminopyralid, clopyralid, diflufenican, dithiopyr, fluridon, fluroxypyr, picloram, picolinafen, thiazopyr;

- sulfonylurinstoffer: amidosulfuron, azimsulfuron, bensulfuron, chlorimuronethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethoxysulfuron, flazasulfuron, flucetosulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metazosulfuron, metsulfuronmethyl, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, 1-((2-chlor-6-propylimidazo[1,2-b]pyridazin-3-yl)sulfonyl)-3-(4,6-dimethoxypyrimidin-2-yl)urinstof;

- triaziner: ametryn, atrazin, cyanazin, dimethametryn, ethiozin, hexazinon, metamitron, metribuzin, prometryn, simazin, terbuthylazin, terbutryn, triaziflam;

- urinstoffer: chlortoluron, daimuron, diuron, fluometuron, isoproturon, linuron, methabenzthiazuron, tebuthiuron;

- andre acetolactatsyntasehæmmere: bispyribacnatrium, cloransulamethyl, diclosulam, florasulam, flucarbazon, flumetsulam, metosulam, ortho-sulfamuron, penoxsulam, propoxycarbazon, pyribambenzpropyl, pyribenzoxim, pyriftalid, pyriminobacmethyl, pyrimisulfan, pyriothiobac, pyroxasulfon,

pyroxsulam;

- andre: amicarbazon, aminotriazol, anilofos, beflubutamid, benazolin, bencarbazon, benfluresat, benzofenap, bentazon, benzobicyclon, bicyclopiron, bromacil, bromobutid, butafenacil, butamifos, cafenstrol, carfentrazon, cinidonethyl, chlorthal, cinmethylin, clomazon, cumyluron, cyprosulfamid, dicamba, difenzoquat, diflufenzopyr, Drechslera monoceras, endothal, ethofumesat, etobenzanid, fenoxasulfon, fentrazamid, flumicloracpentyl, flumioxazin, flupoxam, flurochloridon, flurtamon, indanofan, isoxaben, isoxaflutol, lenacil, propanil, propyzamid, quinclorac, quinmerac, mesotrion, methylarsonsyre, naptalam, oxadiargyl, oxadiazon, oxaziclomefon, pentoxazon, pinoxaden, pyraclonil, pyraflufenethyl, pyrasulfotol, pyrazoxyfen, pyrazolynat, quinoclamín, saflufenacil, sulcotrion, sulfentrazon, terbacil, tefuryltrion, tembotrion, thiencarbazon, topramezon, (3-[2-chlor-4-fluor-5-(3-methyl-2,6-dioxo-4-trifluormethyl-3,6-dihydro-2H-pyrimidin-1-yl)-phenoxy]-pyridin-2-yloxy)-eddikesyreethylester, 6-amino-5-chlor-2-cyclopropylpyrimidin-4-carboxylsyremethylester, 6-chlor-3-(2-cyclopropyl-6-methylphenoxy)-pyridazin-4-ol, 4-amino-3-chlor-6-(4-chlorphenyl)-5-fluorpyridin-2-carboxylsyre, 4-amino-3-chlor-6-(4-chlor-2-fluor-3-methoxyphenyl)-pyridin-2-carboxylsyremethylester og 4-amino-3-chlor-6-(4-chlor-3-dimethylamino-2-fluorphenyl)-pyridin-2-carboxylsyremethylester;

0) Insecticider

- organo(thio)phosphater: acephat, azamethiphos, azinphosmethyl, chlorpyrifos, chlorpyrifosmethyl, chlorfenvinphos, diazinon, dichlorvos, dicrotophos, dimethoat, disulfoton, ethion, fenitrothion, fenthion, isoxathion, malathion, methamidophos, methidathion, methylparathion, mevinphos, monocrotophos, oxydemetonmethyl, paraoxon, parathion, phenthoat, phosalon, phosmet, phosphamidon, phorat, phoxim, pirimiphosmethyl, profenofos, prothiofos, sulprophos, tetrachlorvinphos, terbufos, triazophos, trichlorfon;

- carbamater: alanycarb, aldicarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, fenoxycarb, furathiocarb, methiocarb, methomyl, oxamyl, pirimicarb, propoxur, thiodicarb,

triazamat;

- pyrethroider: allethrin, bifenthrin, cyfluthrin, cyhalothrin, cyphenothrin, cypermethrin, alfa-cypermethrin, beta-cypermethrin, zeta-cypermethrin, deltamethrin, esfenvalerat, etofenprox, fenpropathrin, fenvalerat, imiprothrin, lambda-cyhalothrin, permethrin, prallethrin, pyrethrin I og II, resmethrin, silafluofen, tau-fluvalinat, tefluthrin, tetramethrin, tralomethrin, transfluthrin, profluthrin, dimefluthrin;

10 - insektvækstregulerende midler: a) chitinsyntesehæmmere: benzoylurinstoffer: chlorfluazuron, cyramazin, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron; buprofezin, diofenolan, hexythiazox, etoxazol, clofentazine; b) ecdysonantagonister: 15 halofenozid, methoxyfenozid, tebufenozid, azadirachtin; c) juvenoider: pyriproxyfen, methopren, fenoxycarb; d) lipidbiosyntesehæmmere: spiroadiclofen, spiromesifen, spirotetramat;

- nicotinreceptor-agonist/antagonistforbindelser: clothianidin, 20 dinotefuran, imidacloprid, thiamethoxam, nitenpyram, acetamiprid, thiacloprid, 1-(2-chlorthiazol-5-ylmethyl)-2-nitrimino-3,5-dimethyl-[1,3,5]triazinan;

- GABA-antagonistforbindelser: endosulfan, ethiprole, fipronil, vaniliprol, pyrafluprol, pyriprol, 5-amino-1-(2,6-dichlor-4-methylphenyl)-4-sulfinamoyl-1H-pyrazol-3-carbothiosyreamid; 25

- makrocycliske lactoninsekticider: abamectin, emamectin, milbemectin, lepimectin, spinosad, spinetoram;

- mitokondrisk elektrontransporthæmmer (METI) I-acaricider: fenazaquin, pyridaben, tebufenpyrad, tolfenpyrad, flufenerim;

30 - METI II- og III-forbindelser: acequinocyl, fluacyprim, hydramethylnon;

- Afkoblenende midler: chlorfenapyr;

- hæmmere af oxidativ phosphorylering: cyhexatin, diafenthiuron, fenbutatinoxid, propargit;

35 - fældningsforstyrrende forbindelser: cryomazin;

- oxidasehæmmere med blandet funktion: piperonylbutoxid;

- natriumkanalblokerende midler: indoxacarb, metaflumizon;

- andre: benclothiaz, bifenazat, cartap, flonicamid, pyridalyl,

pymetrozin, svovl, thiocyclam, flubendiamid,
chlorantraniliprol, cyazypyr (HGW86), cyenopyrafen,
flupyrazofos, cyflumetofen, amidoflumet, imicyafos,
bistrifluron og pyrifluquinazon.

DRAWINGS

Structural Models of G143A mutant binding site




Fig. 1: The commercially available strobilurin analogue compound Pyraclostrobin modelled into an artificial cytochrome bc_1 complex Qo-binding site displays steric clash of the phenyl ring with the 143A methyl group as indicated by arrow in the upper right part of the picture (left). Compound I-6 according to the present invention does not show a steric clash (right).