

(19) **DANMARK**

(10) **DK/EP 2854764 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 9/08 (2006.01)** **A 61 K 9/00 (2006.01)** **A 61 K 9/10 (2006.01)**
A 61 K 31/196 (2006.01) **A 61 K 31/473 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2019-04-08**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-12-12**
- (86) Europæisk ansøgning nr.: **13736976.5**
- (86) Europæisk indleveringsdag: **2013-06-05**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-04-08**
- (86) International ansøgning nr.: **IL2013050487**
- (87) Internationalt publikationsnr.: **WO2013183055**
- (30) Prioritet: **2012-06-05 US 201261655633 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Neuroderm Ltd, Weizmann Science Park , 3 Golda Meir Street, 74036 Ness Ziona, Israel**
- (72) Opfinder: **YACOBY-ZEEVI, Oron, 204 Hanarkissim Street, 60946 Bitsaron, Israel**
NEMAS, Mara, 17/4 Ha'atsmaut Street, 70700 Gedera, Israel
- (74) Fuldmægtig i Danmark: **Dennemeyer & Associates S.A, P.O. Box 700425, DE-81304 Munich, Tyskland**
- (54) Benævnelse: **SAMMENSÆTNINGER, DER OMFATTER APOMORFIN OG ORGANISKE SYRER, SAMT ANVENDELSER DERAFT**
- (56) Fremdragne publikationer:
EP-A2- 1 270 007
US-A1- 2007 269 482
MING-JUN TSAI ET AL: "Oral apomorphine delivery from solid lipid nanoparticles with different monostearate emulsifiers: Pharmacokinetic and behavioral evaluations", JOURNAL OF PHARMACEUTICAL SCIENCES, vol. 100, no. 2, 25 February 2011 (2011-02-25), pages 547-557, XP055078666, ISSN: 0022-3549, DOI: 10.1002/jps.22285

DESCRIPTION

TECHNICAL FIELD

[0001] The present invention is generally directed to stable liquid pharmaceutical compositions of apomorphine, and uses thereof in treating neurological diseases or disorders, including Parkinson's disease, or conditions associated therewith.

BACKGROUND ART

[0002] Parkinson's disease is a progressive degenerative disease of the central nervous system. Although the primary cause of Parkinson's disease is not known, it is characterized by the degeneration of dopaminergic neurons of the substantia nigra. The substantia nigra is located in the midbrain and is involved in controlling voluntary movements. The degeneration of neurons causes a shortage of dopamine in the brain, which is believed to cause the observable symptoms of the disease. These symptoms include paucity of movement and rigidity, resting tremor, bradykinesia, and poor balance.

[0003] There are a variety of therapeutic treatments available for Parkinson's disease. The best known is levodopa, a dopamine precursor; however, treatment with levodopa can cause serious side-effects, especially over a long term. One such complication of long-term treatment with levodopa is the development of rapid fluctuations in clinical state, where a patient switches suddenly between mobility and immobility for periods ranging from a few minutes to a few hours. This phenomenon is known as the "on-off effect", the "on" state characterized by the levodopa benefit of early normal motor functioning and the "off" state characterized by akinesia - abrupt loss of mobility, e.g., where a patient may suddenly stop while walking. Approximately half of patients on levodopa therapy will develop such on-off effects after several years of therapy.

[0004] While apomorphine hydrochloride has proved effective in treating "off" episodes in patients with Parkinson's disease, a common and serious side effect of administering apomorphine hydrochloride by subcutaneous injection is the development of subcutaneous nodules at the injection site, which can become infected, necessitating treatment or surgical involvement. A majority of people on infused apomorphine develop nodules, and a new nodule may form every time the infusion needle is re-sited, which may happen on a daily basis. Such nodules may be painful, limit available infusion sites and interfere with absorption. Further, unstable compositions, e.g., having precipitate of apomorphine or other agents, may be the cause, or exacerbate, such nodule side effects. Thus, there is a need for new, stable formulations of apomorphine which are safe and effective for administration to patients.

[0005] EP 1 270 007 A2 relates to pharmaceutical compositions in the form of microemulsions comprising apomorphine, apomorphine hydrochloride or other salt.

SUMMARY OF INVENTION

[0006] This disclosure relates at least in part to the discovery that certain organic acids, e.g., acidic amino acids, form stable salts of apomorphine that are suitable for compositions allowing for subcutaneous administration without unwanted side effects.

[0007] In one aspect, the present invention thus relates to a stable liquid pharmaceutical composition, particularly an aqueous pharmaceutical composition, comprising apomorphine, an organic acid, e.g., an acidic amino acid, and a pharmaceutically acceptable carrier, wherein the molar ratio of apomorphine to the organic acid is about 1 to about 0.5-2.5, or about 1 to about 0.5-1.5, e.g., 1:1.2.

[0008] In a further aspect, the pharmaceutical composition of the present invention can be used for the treatment of a

neurological or movement disease or disorder, or a condition associated therewith, in a patient in need thereof, by administering a pharmaceutical composition as disclosed herein. In one embodiment, the method of the invention is aimed at treating a neurological disease or disorder characterized by reduced levels of dopamine in a patient's brain, e.g., Parkinson's disease, by administering the disclosed composition.

[0009] In one particular such aspect, the pharmaceutical composition of the present invention can be used for the treatment of Parkinson's disease in a patient in need thereof by administering a composition as disclosed herein.

[0010] In another particular such aspect, the pharmaceutical composition of the present invention can be used for the treatment of akinesia in a patient in need thereof, comprising administering to the patient a composition as disclosed herein, wherein the patient has been treated with levodopa.

[0011] In still another aspect, the present invention relates to a liquid pharmaceutical composition as defined above, for use in a treatment of a neurological or movement disease or disorder, or a condition associated therewith.

[0012] In yet another aspect, the present invention relates to a liquid or semi-solid pharmaceutical composition as defined above, for use in a treatment of a neurological or movement disease or disorder, or a condition associated therewith.

BRIEF DESCRIPTION OF DRAWINGS

[0013]

Fig. 1 shows the system used for evaluating the effect of apomorphine formulation composition on pig skin staining. Infusion pump set at 0.05 ml/h (**A**); Formulation delivered using a 22G butterfly (**B**); Full thickness pig skin (cutis and subcutis) is placed on top of a bottle containing PBS. The skin is directly in contact with the PBS. The skin and bottle are covered with parafilm to protect the system from air as much as possible (**C**); A 100 ml bottle filled to the top with warm PBS (**D**); Formulation, 0.4 ml, was administered subcutaneously via a 22G needle, at 0.05 ml/h (total of 8 h) (**E**); and Incubator set at 37°C. Samples were incubated for a period of about 17 h (**F**).

Fig. 2 shows the effect of the formulation composition on apomorphine-dependent skin staining following continuous subcutaneous infusion into full thickness pig skin of some of the compositions exemplified in Example 3 hereinafter, wherein panel (a) shows 1% apomorphine-HCl (APO-GO®), pH 3.5; panel (b) shows 1% apomorphine-glutamate, pH 3.5; panel (c) shows 1% apomorphine-glutamate with ascorbic acid, pH 3.3; panel (d) shows 1% apomorphine-glutamate with ascorbic acid and meglumine, pH 4.2; panel (e) shows 1% apomorphine-glutamate with meglumine, pH 4.3; panel (f) shows 1% apomorphine-aspartate, pH 3.6; panel (g) shows 1% apomorphine-aspartate with meglumine pH 4.2; panel (h) shows 1% apomorphine-tartrate, pH 2.9; panel (i) shows 1% apomorphine-hemi tartrate, pH 3.2; and panel (j) shows 1% apomorphine-tartrate with meglumine, pH 3.6.

Fig. 3 shows histological grading of skin samples (Grading of Subcutis (SC) Changes) 1, 2 or 3 weeks after subcutaneous administration of apomorphine solution containing 0.5% or 2.0% Tween-80.

DETAILED DESCRIPTION OF THE INVENTION

Compositions

[0014] In one aspect, the present invention relates to a stable liquid pharmaceutical composition, particularly an aqueous pharmaceutical composition, comprising apomorphine and an organic acid. Exemplary organic acids include, without being limited to, amino acids, carboxylic acids, and dicarboxylic acids. For example, contemplated carboxylic acids and/or dicarboxylic acids for use in the claimed composition may contain at least two, three, or four carbon

atoms, e.g., tartaric acid. Contemplated dicarboxylic acids for use in the claimed compositions may be hydrophilic or substituted with hydrophilic groups, e.g., hydroxyl groups.

[0015] In the present invention, the contemplated compositions comprise apomorphine and an acidic amino acid, e.g., a natural or unnatural amino acid. Contemplated acidic amino acids for use in the claimed compositions may be acidic natural amino acids such as aspartic acid or glutamic acid, or acidic unnatural amino acids such as cysteic acid.

[0016] Contemplated herein, e.g., is a stable pharmaceutical composition comprising apomorphine and an acidic organic acid, where the molar ratio of apomorphine to the acidic organic acid is about 1 to about 0.5, or about 1.0 to about 2.5, or about 1 to about 0.5 to 1.5 (e.g., about 1:0.5 to about 1.0 to about 1.5). For example, in certain embodiments, the molar ratio of apomorphine to the organic acid is about 1:0.5, or about 1:0.6, or about 1:0.7, or about 1:0.8, or about 1:0.9, or about 1:1, or about 1:1.1, or about 1:1.2, or about 1:1.3, or about 1:1.4, or about 1:1.5, or about 1:1.6, or about 1:1.7, or about 1:1.8, or about 1:1.9, or about 1:2.0, or about 1:2.1, or about 1:2.2, or about 1:2.3, or about 1:2.4, or about 1:2.5.

[0017] In a particular embodiment, contemplated herein is a stable pharmaceutical composition comprising apomorphine and an acidic amino acid, where the molar ratio of apomorphine to the acidic amino acid is about 1 to about 0.5-2.5. For example, in certain embodiments, the molar ratio of apomorphine to the amino acid is about 1:0.5, or about 1:0.6, or about 1:0.7, or about 1:0.8, or about 1:0.9, or about 1:1, or about 1:1.1, or about 1:1.2, or about 1:1.3, or about 1:1.4, or about 1:1.5, or about 1:1.6, or about 1:1.7, or about 1:1.8, or about 1:1.9, or about 1:2.0, or about 1:2.1, or about 1:2.2, or about 1:2.3, or about 1:2.4, or about 1:2.5. In more particular embodiments, the amino acid is an acidic amino acid such as, without being limited to, aspartic acid, glutamic acid, or a combination thereof.

[0018] The pharmaceutical composition has a pH of greater than 3 at 25°C. In certain particular embodiments, the pharmaceutical composition has a pH between about 3.0 and about 5.5, more particularly between about 3.0 and about 5.0 at 25°C. In more particular embodiments, the composition has a pH between about 3.1 and about 4.9, about 3.2 and about 4.8, about 3.3 and about 4.7, about 3.4 and about 4.6, about 3.5 and about 4.5, about 3.6 and about 4.4, about 3.7 and about 4.3, about 3.8 and about 4.2, or about 3.9 and about 4.1 at 25°C. In other embodiments, the pH of the composition is about 4 at 25°C. In other particular embodiments, the pharmaceutical composition has a pH between about 4.0 and about 5.0, or between about 4.0 and about 7.5, at 25°C.

[0019] In certain embodiments, the pharmaceutical composition comprises at least about 1% by weight apomorphine. In some embodiments, the composition comprises about 1% to about 4% by weight apomorphine, e.g., about 1.25%, about 1.5%, about 1.75%, about 2%, about 2.25%, about 2.5%, about 2.75%, about 3%, about 3.25%, about 3.5%, about 3.75%, or about 4% by weight apomorphine. In other embodiments, a contemplated composition comprises about 1% to about 2.5% by weight apomorphine, or about 0.5% to about 3% by weight apomorphine, or about 1.1% to about 2.6% by weight or more apomorphine.

[0020] In certain embodiments, the pharmaceutical compositions of the present invention are stable for 24 hours, or for at least 24 hours, or more, at room temperature, e.g., at 20°C to 30°C, e.g., at 25°C. For example, in some embodiments, the disclosed compositions are stable for 24 hours, for at least 48 hours, or for 7 days or more, at 25°C. In particular embodiments, the disclosed pharmaceutical compositions are stable for at least 48 hours at 25°C, or are even stable for longer periods at room temperature, e.g., for 3 days, 1 week, 1 month, or longer.

[0021] In certain embodiments, the pharmaceutical composition has substantially no precipitation of solids for at least 24 hours, 48 hours, 7 days, or 2 months, e.g., at 20°C to 30°C, e.g., at 25°C. In a particular embodiment, the pharmaceutical composition has substantially no precipitation of solids for at least 48 hours at, e.g., 20°C to 30°C, e.g., at 25°C.

[0022] In certain embodiments, the pharmaceutical composition of the present invention further comprises an amino sugar, a basic amino acid, one or more antioxidants, or a combination thereof.

[0023] In certain particular embodiments, the pharmaceutical compositions of the invention further comprises an amino sugar such as, without being limited to, meglumine, and may also comprise a basic amino acid and one or more antioxidants. Particular such compositions comprise apomorphine, aspartic acid and/or glutamic acid, and

meglumine.

[0024] In certain particular embodiments, the pharmaceutical compositions of the invention further comprises a basic amino acid such as, without limiting, arginine, and may also comprise an amino sugar and one or more antioxidants. Particular such compositions comprise apomorphine, aspartic acid and/or glutamic acid, and arginine.

[0025] In certain particular embodiments, the pharmaceutical compositions of the present invention further comprises at least one, i.e., one, two, three or more, antioxidant, i.e., an agent that inhibits the formation of oxidation products. Such an agent may be, e.g., tyrosinase inhibitors and/or o-quinone scavengers and/or Cu^{++} chelators and/or antioxidants, and/or tetrahydroquinolines. For example, contemplated formulations may include o-quinone scavengers such as N-acetyl cysteine, glutathione, ascorbic acid, Na-ascorbate, and/or L-cysteine. For example, a contemplated formulation may include ascorbic acid and a cysteine, e.g., L-cysteine or N-acetyl cysteine. In a contemplated formulation, the ratio of ascorbic acid to another antioxidant, e.g., L-cysteine or N-acetyl cysteine, may exist at a particular weight-to-weight ratio such as about 1:1, about 2:1, about 3:1, about 4:1, about 5:1, about 6:1, about 7:1, about 8:1, about 9:1, or about 10:1. In some embodiments, formulations may include an agent chosen from one or more of tyrosinase inhibitors such as captopril, methimazole, quercetin, arbutin, aloesin, N-acetylglucosamine, retinoic acid, α -tocopheryl ferulate, MAP (Mg ascorbyl phosphate), substrate analogues (e.g., sodium benzoate, L-phenylalanine), Cu^{++} chelators (for example, $\text{Na}_2\text{-EDTA}$, $\text{Na}_2\text{-EDTA-Ca}$), DMSA (succimer), DPA (D-penicillamine), trientine-HCl, dimercaprol, clioquinol, sodium thiosulfate, TETA, TEPA, curcumin, neocuproine, tannin, and/or cuprizone. Other contemplated anti-oxidants that may form part of a disclosed formulation include sulfite salts, e.g., sodium hydrogen sulfite or sodium metabisulfite, di-tert-butyl methyl phenols, tert-butyl-methoxyphenols, polyphenols, tocopherols and/or ubiquinones, including caffeic acid.

[0026] Contemplated antioxidants that can be included in disclosed compositions may be selected from, e.g., thiols such as aurothioglucose, dihydrolipoic acid, propylthiouracil, thioredoxin, glutathione, cysteine, cystine, cystamine, and thiodipropionic acid; sulfoximines such as buthionine-sulfoximines, homo-cysteine-sulfoximine, buthionine-sulphones, and penta-, hexa- and heptathionine-sulfoximine; metal chelators such as α -hydroxy-fatty acids, palmitic acid, phytic acid, lactoferrin, citric acid, lactic acid, malic acid, humic acid, bile acid, bile extracts, bilirubin, biliverdin, EDTA, EGTA, and DTPA; sodium metabisulfite; vitamins such as vitamin E, vitamin C, ascorbyl palmitate, Mg ascorbyl phosphate, and ascorbyl acetate; phenols such as butylhydroxytoluene, butylhydroxyanisole, ubiquinol, nordihydroguaiaretic acid, and trihydroxybutyrophene; benzoates such as coniferyl benzoate; uric acid; mannose; propyl gallate; selenium such as selenium-methionine; stilbenes such as stilbene oxide and trans-stilbene oxide; and combinations thereof.

[0027] In particular embodiments, a contemplated pharmaceutical composition further comprises one or more antioxidants each independently selected from ascorbic acid, Na-ascorbate, L-cysteine, N-acetylcysteine (NAC), glutathione (GSH), $\text{Na}_2\text{-EDTA}$, $\text{Na}_2\text{-EDTA-Ca}$, or sodium bisulfite. In specific such embodiments, a contemplated pharmaceutical composition further comprises ascorbic acid and/or sodium bisulfite.

[0028] In certain embodiments, the pharmaceutical composition of the present invention further comprises a surfactant, i.e., a compound that lowers the surface tension of a liquid. Non-limiting examples of suitable surfactants include polysorbate surfactants such as Tween-80, Tween-60, Tween-40, Tween-20, Tween-65 and Tween-85, and sorbitane surfactants such as Span 20, Span 40, Span 60, Span 80 and Span 85. In a particular such embodiment, the pharmaceutical composition of the invention further comprises Tween-80.

[0029] In certain embodiments, the pharmaceutical composition of the present invention further comprises a local anesthetic, i.e., a drug which causes a reversible loss of sensation for a limited region of the body while maintaining consciousness, and/or an anti-inflammatory agent. Examples of local anesthetics include, without being limited to, amide based local anesthetics such as lidocaine, prilocaine, bupivacaine, levobupivacaine, ropivacaine, mepivacaine, dibucaine, and etidocine, and ester based local anesthetics such as procaine, amethocaine, cocaine, benzocaine, and tetracaine. Examples of anti-inflammatory agents include, without limiting, non-steroidal anti-inflammatory agents such as diclofenac, ketorolac, salicylates ibuprofen, piroxicam, and benzydamine, and steroidal anti-inflammatory agents such as prednisone, dexamethasone, betamethasone, prednisone hydrocortisone or salts thereof.

[0030] The pharmaceutical compositions of the present invention are liquid solutions, i.e., substantially homogeneous liquid mixtures at room temperature, e.g., at 25°C. Such liquid mixtures may comprise water and/or other pharmaceutically acceptable carriers or excipients. In a particular embodiment, the disclosed composition is substantially aqueous.

[0031] In certain embodiments, a disclosed liquid or semi-solid formulation will be stable for a period of 1 day, 2 days, 3 days, 1 week, 1 month, or more at room temperature. In certain embodiments, the pharmaceutical composition as defined herein further comprises a pharmaceutically acceptable excipient such as, without being limited to, Tween-80, Tween-60, Tween-40, Tween-20, N-methylpyrrolidone (NMP), or polyvinylpyrrolidone (PVP), EDTA (or salts thereof), cysteine, N-acetylcysteine, and/or sodium bisulfite.

[0032] Another stable liquid pharmaceutical composition, particularly an aqueous pharmaceutical composition, comprises (i) apomorphine or a salt thereof; and (ii) either an emulsifier or surfactant, or both emulsifier and surfactant. In one particular such aspect, the composition comprises apomorphine, and an emulsifier, surfactant, or both. In another particular such aspect, the composition comprises a salt of apomorphine, and an emulsifier, surfactant, or both. The apomorphine salt may be any salt formed by reacting apomorphine with an organic acid. Particular organic acids include, without limiting, amino acids, more particularly acidic amino acid such as aspartic acid, glutamic acid and cysteic acid, as well as carboxylic acids and dicarboxylic acids containing at least two, three, or four carbon atoms, e.g., tartaric acid. Particular such compositions have substantially no precipitation of solids, for at least 48 hours at 25°C within a pH ranging from about 3 to about 7.5, or at a pH greater than 4. These compositions may be liquid solutions, i.e., substantially homogeneous liquid mixtures at room temperature, e.g., at 25°C, or semi-solid solutions formulated, e.g., as a gel, gum, or candy, and may comprise water and/or other pharmaceutically acceptable carriers or excipients. In particular embodiments, the disclosed compositions are substantially aqueous.

[0033] Such other pharmaceutical composition may comprise at least 1% by weight apomorphine, or about 1% to about 4%, e.g., about 1.25%, about 1.5%, about 1.75%, about 2%, about 2.25%, about 2.5%, about 2.75%, about 3%, about 3.25%, about 3.5%, about 3.75%, or about 4%, by weight apomorphine.

[0034] Such other pharmaceutical composition may also comprise 0.01% to 4%, e.g., 0.01%, 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2.0%, 2.2%, 2.4%, 2.6%, 2.8%, 3.0%, 3.2%, 3.4%, 3.6%, 3.8%, or 4.0%, emulsifier or surfactant.

[0035] Such other pharmaceutical composition may further comprise one or more, i.e., one, two, three or more, antioxidants, as defined above. In particular embodiments, a contemplated pharmaceutical composition further comprises one or more antioxidants each independently selected from ascorbic acid, Na-ascorbate, L-cysteine, NAC, GSH, Na₂-EDTA, Na₂-EDTA-Ca, or sodium bisulfite.

[0036] The stable pharmaceutical composition of the present invention is suitable for subcutaneous administration. Generally, a stable pharmaceutical composition may also be suitable for intravenous, intranasal, transdermal, sublingual, intramuscular, buccal, ocular, or intratracheal administration. In some embodiments, the composition of the present invention is continuously infused, e.g., subcutaneously via a small pump. Generally, a stable pharmaceutical composition may also be inhaled or administered parenterally.

[0037] The disclosed compositions may be administered subcutaneously and/or, e.g., substantially continuously. When administered subcutaneously and/or dermally, such compositions having apomorphine or a pharmaceutically acceptable salt thereof, and an organic acid, e.g., an acidic amino acid, may result in minimal local tissue damage, e.g., as compared to subcutaneous or dermal administration of a composition that includes apomorphine-HCl. Further, such apomorphine/organic acid compositions, when further including a basic amino acid such as arginine, and/or a basic amino sugar such as meglumine, and/or a surfactant such as Tween-80, may have more stability, e.g., may not precipitate and/or form unwanted oxidation products over time as compared to a composition having apomorphine-HCl. Further, such apomorphine/organic acid compositions, when further including a surfactant such as Tween-80 may have more stability, e.g., may not precipitate and/or form unwanted oxidation products over time as compared to a composition having apomorphine without a surfactant. As contemplated herein, subcutaneous administration may comprise the use of one or more infusion pumps and/or transdermal and/or dermal patches. Alternatively, a disclosed composition may be administered intraduodenally, intranasally, sublingually, intramuscularly,

ocularly, buccally, intratracheally or intravenously.

[0038] As used herein in the specification, the term "organic acid" refers to an organic compound with acidic properties such as carboxylic acids, dicarboxylic acids, sulfonic acids, alcohols, hydroxy acids, thiols, and thio-acids. For example, the organic acids for use in the claimed composition may contain at least two, three or four carbon atoms, e.g., tartaric acid. Examples of organic acids include amino acids such as aspartic acid, glutamic acid, and arginine, and dicarboxylic acids such as fumaric acid, oxalic acid, malonic acid, succinic acid, glutaric acid and maleic acid. Further examples of organic acids include lactic acid, malic acid, aconitic acid, citric acid, glycolic acid, ascorbic acid, formic acid, acetic acid, tartaric acid, and glucuronic acid.

[0039] The term "natural amino acid" refers to any of the amino acids found in proteins. Examples of natural amino acids include alanine, arginine, aspartic acid, glutamic acid, histidine and lysine.

[0040] The term "non-natural amino acid" refers to non-proteinogenic amino acids that either occur naturally or are chemically synthesized. Examples of non-natural amino acids include ornithine, β -alanine, 2-aminoadipic acid, 3-aminoadipic acid, γ -carboxyglutamic acid, hydroxylysine, 4-guanidinobutyric acid, 3-guanidinopropionic acid, 4-azidobutanoic acid and 5-azidopentanoic acid. Both D- and L- amino acids are contemplated herein.

[0041] The term "amino sugar" refers to a monosaccharide in which a non-glycosidic hydroxyl group is replaced by an amino or substituted amino group, such as meglumine, D-glucosamine, sialic acid, N-acetylglucosamine and galactosamine.

[0042] The term "anti-oxidant" refers to an agent that inhibits the formation of oxidation products.

[0043] The term "emulsifier", also known as "emulgent", refers to a substance that stabilizes an emulsion by increasing its kinetic stability, e.g., a "surface active substance" or a surfactant. The term "surfactant" refers to a surface active substance, i.e., a compound that lowers the surface tension of a liquid, the interfacial tension between two liquids, or that between a liquid and a solid.

[0044] The term "pharmaceutical composition" as used herein refers to a composition comprising at least one active agent as disclosed herein, formulated together with one or more pharmaceutically acceptable carriers.

[0045] The term "pharmaceutically acceptable carrier" or "pharmaceutically acceptable excipient" as used herein refers to any and all solvents, dispersion media, preservatives, antioxidants, coatings, isotonic and absorption delaying agents and surfactants that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. The compositions may contain other active compounds providing supplemental, additional, or enhanced therapeutic functions.

[0046] The terms "pharmaceutically acceptable" and "pharmacologically acceptable" include molecular entities and compositions that do not produce an adverse, allergic, or other untoward reactions when administered to an animal, or human, as appropriate. For human administration, preparations should meet sterility, pyrogenicity, general safety, and purity standards as required by a government drug regulatory agency, e.g., the United States Food and Drug Administration (FDA) Office of Biologics standards.

[0047] The term "physiologically acceptable pH" is understood to mean a pH of a composition that facilitates administration of the composition to the patient without significant adverse effects, e.g., a pH of about 3 to about 9.8 at 25°C, e.g., a pH of about 3.5 ± 0.5 to about 9.5 ± 0.3 , or about 3.5 to about 9.5.

Methods

[0048] In a further aspect, the present invention provides a pharmaceutical composition as disclosed herein for use in a method of treating a neurological or movement disease or disorder, or a condition associated therewith, in a patient in need thereof, comprising administering to said patient the pharmaceutical composition.

[0049] The neurological or movement diseases or disorders treated by this method are characterized by reduced levels of dopamine in the brain, and include restless leg syndrome, Parkinson's disease, secondary parkinsonism, Alzheimer's disease, Huntington's disease, Shy-Drager syndrome, dystonia, akinesia, alcohol and morphine addiction, erectile dysfunction, and conditions resulting from brain injury, including carbon monoxide or manganese intoxication.

[0050] In one particular embodiment, the invention provides a pharmaceutical composition as disclosed herein for use in a method of treating Parkinson's disease in a patient in need thereof, comprising administering to the patient the pharmaceutical composition.

[0051] In another particular embodiment, the invention provides a pharmaceutical composition as disclosed herein for use in a method of treating akinesia in a patient in need thereof, comprising administering to the patient the pharmaceutical composition, wherein the patient has been treated with levodopa.

[0052] In a further embodiment, the invention provides a pharmaceutical composition as disclosed herein for use in a method of treating a condition associated with a neurological disease or disorder in a patient in need thereof, comprising administering to the patient the pharmaceutical composition. The neurological disease or disorder contemplated herein is Parkinson's disease, Alzheimer's disease, or akinesia, and the conditions associated with a neurological disease or disorder include, without limiting, alcoholism, opiate addiction, and erectile dysfunction.

[0053] In still another aspect, the present invention relates to a liquid pharmaceutical composition as defined above for use in a method for the treatment of a neurological or movement disease or disorder as defined herein, or a condition associated therewith.

[0054] In yet another aspect, the present invention relates to a liquid pharmaceutical composition as defined above, for use in a method for the treatment of a neurological or movement disease or disorder as defined herein, or a condition associated therewith.

[0055] As used herein in the specification, the term "neurological disease or disorder" refers to a disease or disorder of the body's nervous system.

[0056] The term "movement disorder" refers to a nervous system condition that causes abnormal voluntary or involuntary movements, or slow, reduced movements.

[0057] The term "treated with levodopa" includes past or present treatment of a patient with levodopa.

Kits

[0058] A contemplated kit may include an apomorphine salt or a liquid composition having apomorphine and an organic acid (and optionally an additional therapeutic agent, e.g., levodopa, carbidopa, or entacapone). Such a composition may be liquid or a lyophilized powder that can be reconstituted into a liquid formulation, or, for example, may form part of a transdermal patch, and may be designed for administration by any suitable route such as transdermally, intravenously, subcutaneously, intradermally, intramuscularly, or intraduodenally.

[0059] In some embodiments, disclosed liquid compositions, e.g., comprising apomorphine and an organic acid, may be provided in, e.g., one, two or more pre-filled cartridges suitable for use by a patient or physician. For example, provided herein is a kit comprising a prefilled cartridge wherein a disclosed liquid formulation is disposed within the cartridge (e.g., a pre-filled cartridge having a single dose or a dose suitable for a single administration to a patient of an apomorphine and organic acid formulation (and optionally an additional therapeutic agent, e.g., levodopa or carbidopa)), and optionally instructions for use.

[0060] The invention now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention in any way.

EXAMPLES

Example 1. Preparation of apomorphine salt solution/formulation for subcutaneous administration

[0061] Sodium bicarbonate 0.1 N solution was added to a 1% apomorphine-HCl solution. The apomorphine-HCl solution was derived from APO-GO®, a commercially available product comprising a 10mg/ml apomorphine-HCl solution. The precipitate, being the apomorphine base, was washed twice with water. The precipitated apomorphine base was dissolved with either a 0.59% glutamic acid solution, 0.53% aspartic acid solution, or 0.6% tartaric acid solution, to form a final 1% apomorphine solution at a 1:1 molar ratio with the glutamic, aspartic, or tartaric acid, or a 1:0.5 molar ratio with the tartaric acid. In some instances, ascorbic acid and/or meglumine solutions were added to the apomorphine salt solutions to obtain a final concentration of 0.5% or 0.6%, respectively. **Table 1** shows the composition of apomorphine formulations which were prepared as described above.

Table 1: Composition of apomorphine formulations

APO-GO® (10 mg/ml Apo.HCl) µl	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
NaHCO ₃ (0.1 N)		310	310	310	310	310	310	310	310	310
		prec.	prec.	prec.	prec.	prec.	prec.	prec.	prec.	prec.
Glutamic acid (5.9 mg/ml) µl		800	800	800	800					
Aspartic acid (5.3 mg/ml) µl						800	800			
Tartaric acid (6 mg/ml) µl								800	400	800
Meglumine (60 mg/ml) µl				100	50		50			50+50
Ascorbic acid 5% µl			100	100						
Water µl		200	100		150	200	150	200	600	150
APO/Acid ratio		1 to 1	1 to 1	1 to 1	1 to 1	1 to 1	1 to 1	1 to 1	1 to 0.5	1 to 1
Solutions appearance	trans.	trans.	trans.	trans.	trans.	trans.	trans.	trans.	trans.	trans.
pH	4.19	3.54	3.29	4.18	4.34	3.60	4.21	2.85	3.19	3.60
Total µl	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
prec. = precipitated; trans. = transparent										

Example 2. Preparation of apomorphine salt solution/formulation for subcutaneous administration

[0062] Apomorphine-HCl hemi-hydrate was dissolved in 0.1% sodium bisulfite solution to form 1% apomorphine solution. Apomorphine was precipitated by adding the appropriate amount of 0.1N sodium bicarbonate solution. The precipitate was washed twice with water. The precipitated apomorphine base was dissolved with 1.47% glutamic acid solution to form a final 2.5% apomorphine solution at a 1:1 molar ratio with the glutamic acid. In some instances, ascorbic acid and/or meglumine and/or arginine solutions were added to the apomorphine-glutamate solution to obtain a final concentration of 0.5%, 0.6% or 0.56%, respectively. **Table 2** shows the composition of apomorphine formulations which were prepared as described above.

Table 2: Composition of apomorphine formulations

Formulation No.	1	2
APO.HCl (10 mg/ml) µl	1250	1250
NaHCO ₃ 0.1 N	412	412
	precipitated	precipitated

Formulation No.	1	2
Glutamic acid (14.7 mg/ml) μ l	400	400
Meglumine (60 mg/ml) μ l		50
L-arginine (56 mg/ml)		
Ascorbic acid 5% μ l		50
Water μ l	100	
APO/Acid ratio	1to1	1to1
APO conc. (mg/ml)	25	25
Solutions appearance	transparent	transparent
pH	3.60	4.13
Total μ l	500	500

Example 3. Physical stability of apomorphine salt solutions *in vitro*

[0063] Various apomorphine salt solutions were examined for stability after sitting at room temperature for a period of time. The results of the study are shown in **Table 3**. Notably, apomorphine salt solutions containing aspartate, glutamate, tartrate, maleate and optionally meglumine were stable over the examined time range.

Table 3: Composition of apomorphine salt solutions for stability testing

API		Acid name	Other	Anti-oxidant	pH	Color	Precipitation
Name	Amount						
Apomorphine-HCl (Apo-Go®)	1%	HCl	Examined after 50 hrs at RT, 8 hrs at 40°C and 48 hrs at RT	Na-bisulfite	3.5	Slightly greenish	Yes
Apomorphine	1%	Ascorbate		Ascorbic	4.0	Greenish	No
Apomorphine	1%	Ascorbate		Ascorbic	3.3	Greenish	No
Apomorphine	1%	Glutamate		-	3.5	Colorless	No
Apomorphine	1%	Aspartate		-	3.1	Colorless	No
Apomorphine	2.5%	Tartrate	Examined after 48 hrs at RT	-	2.5	Colorless	No
Apomorphine	2.5%	Maleate		-	2.8	Colorless	No
Apomorphine	2.5%	Citrate		-	2.4	Grayish	No
Apomorphine	2.5%	Succinate		-	3.5	Colorless	Yes
Apomorphine	2.5%	Ascorbate		Ascorbic	3.6	Greenish	No
Apomorphine	2.5%	Ascorbate	Meglumine*	Ascorbic Cysteine	4.2	Greenish	No
Apomorphine-HCl (Apo-Go®)	1%	HCl	*	-	3.5	Slightly greenish	No
Apomorphine	1%	Glutamate	*	-	3.5	Colorless	No
Apomorphine	1%	Glutamate	*	Ascorbic	3.3	Greenish	No
Apomorphine	1%	Glutamate	Meglumine*	Ascorbic	4.2	Grayish	No
Apomorphine	1%	Glutamate	Meglumine*	-	4.3	Colorless	No
Apomorphine	1%	Aspartate	*	-	3.6	Colorless	No
Apomorphine	1%	Aspartate	Meglumine*	-	4.2	Colorless	No
Apomorphine	1%	Tartrate	*	-	2.9	Colorless	No
Apomorphine	1%	Tartrate	*	-	3.2	Colorless	No

API		Acid name	Other	Anti-oxidant	pH	Color	Precipitation
Name	Amount						
Apomorphine	1%	Tartrate	Meglumine*	-	3.6	Colorless	No
* Examined after 48 hrs at room temperature (RT)							

Example 4. Preparation of apomorphine salt solution/formulation for subcutaneous administration

[0064] Apomorphine-HCl hemi-hydrate was dissolved in 0.1% sodium bisulfite solution to form a 1% apomorphine solution. Apomorphine was precipitated by adding the appropriate amount of 0.1N sodium bicarbonate solution. The precipitate was washed twice with water. The precipitated apomorphine base was dissolved with 0.94% glutamic acid solution to form a final concentration of 1% or 1.5% apomorphine solution at 1:1 molar ratio with the glutamic acid. In some instances, ascorbic acid and/or meglumine and/or arginine solutions were added to the apomorphine-glutamate solution to obtain a final concentration of 0.5%, 0.6%, or 0.56%, respectively. **Table 4** shows the composition of apomorphine formulations which were prepared as described above.

Table 4: Composition of apomorphine formulations

Formulation No (%)	1	2	3	4	5	6	7	8
APO.HCl (10 mg/ml) μ l	1000	1500	1000	1500	1000	1500	1000	1500
NaHCO ₃ 0.1 N	330	495	330	495	330	495	330	495
	prec.	prec.	prec.	prec.	prec.	prec.	prec.	prec.
Glutamic acid (9.4 mg/ml) μ l	500	750	500	750	500	750	550	826
Meglumine (60 mg/ml) μ l			100	100			100	87
L-Arginine (56 mg/ml)					100	100		
Ascorbic acid 5% μ l			100	100	100	100	100	87
Water μ l	500	250	300	50	300	50	250	
APO/Acid ratio	1 to 1	1 to 1	1 to 1	1 to 1	1 to 1	1 to 1	1 to 1.1	1 to 1.1
APO conc. (mg/ml)	10	15	10	15	10	15	10	15
Solutions appearance	trans.	trans.	trans.	trans.	trans.	trans.	trans.	trans.
pH	3.83	3.76	4.40	4.27	4.46	4.32	4.41	4.20
Total μ l	1000	1000	1000	1000	1000	1000	1000	1000
prec. = precipitated; trans. = transparent								

Example 5. Effect of formulation composition on apomorphine-dependent SC staining following continuous subcutaneous infusion into full thickness pig skin, ex vivo

[0065] Various apomorphine-salt formulations were prepared and administered subcutaneously by continuous infusion into full thickness pig skin (cutis and subcutis) at a rate of 0.05 ml/h. **Fig. 1** shows the system used for the procedure. During infusion, the apomorphine formulations were incubated at 37°C for a period of 17 hours.

[0066] **Fig. 2** shows the effect of the formulation composition on apomorphine-dependent skin staining following continuous subcutaneous infusion into full thickness pig skin of some of the compositions listed in **Table 3**, wherein panel (a) shows 1% apomorphine-HCl (APO-GO®), pH 3.5; panel (b) shows 1% apomorphine-glutamate, pH 3.5; panel (c) shows 1% apomorphine-glutamate with ascorbic acid, pH 3.3; panel (d) shows 1% apomorphine-glutamate with ascorbic acid and meglumine, pH 4.2; panel (e) shows 1% apomorphine-glutamate with meglumine, pH 4.3; panel (f) shows 1% apomorphine-aspartate, pH 3.6; panel (g) shows 1% apomorphine-aspartate with meglumine pH 4.2; panel (h) shows 1% apomorphine-tartrate, pH 2.9; panel (i) shows 1% apomorphine-hemi tartrate, pH 3.2; and

panel (j) shows 1% apomorphine-tartrate with meglumine, pH 3.6. As shown in **Fig. 2**, the formulation containing 1% apomorphine-HCl (APO-GO®) caused the most severe staining of pig skin *ex vivo*. Notably, apomorphine salt formulations containing acidic amino acids (aspartate and glutamate) and optionally meglumine caused significantly less staining of pig skin than apomorphine-tartrate and HCl salts.

Example 6. Preparation of glutamate, aspartate and tartrate salts of apomorphine

[0067] Apomorphine base preparation was performed in the following manner. Apomorphine hydrochloride (Johnson Matthey, Macfarlan Smith) was dissolved in 0.1% Sodium metabisulfite solution (10mg/ml). Sodium bicarbonate solution at 0.4M was slowly added to the solution with stirring. The mixture was then stirred for 30 minutes at room temperature. The white precipitate that formed was filtered under nitrogen, washed twice with sterile water and dried in a vacuum oven under dry nitrogen flow for 4 hours. The yield of apomorphine dry base was 75.5%.

[0068] Apomorphine salts were prepared by dissolving the apomorphine dry base in the appropriate acid solution at molar ratios ranging between 1:0.5 and 1:1.5.

[0069] Apomorphine glutamate solution was prepared by dissolving apomorphine base in 0.1M glutamic acid solution. Water was added to obtain solutions with a molar ratio of 1:1 or 1:1.2, apomorphine:glutamic acid, at a final concentration of 2% apomorphine. The pH of each solution was 4.7 and 4.4, respectively.

[0070] Apomorphine aspartate solution was prepared by dissolving apomorphine base in 0.1M aspartic acid solution. Water was added to obtain a molar ratio of 1:1, apomorphine:aspartic acid, at a final concentration of 2% apomorphine. The pH of the solution was 4.9.

[0071] Apomorphine tartrate solution was prepared by dissolving apomorphine base in 0.1M tartaric acid solution. Water was added to obtain solutions with a molar ratio of 1:1 or 1:0.5, apomorphine:tartaric acid, at a concentration of 2% apomorphine. The pH of each solution was 3.45 and 4.0, respectively.

[0072] Apomorphine malate solution was prepared by dissolving apomorphine base in 0.1M Malic acid solution. Water was added to obtain solutions with a molar ratio of 1:1 apomorphine:malic acid, at a concentration of 2% apomorphine. The pH of each solution was 4.0. **Table 5** shows the composition of each apomorphine organic salt solution.

Example 7. Preparation of apomorphine-glutamate formulations with Tween-80

[0073] Apomorphine-glutamate solutions were prepared in the following manner. Antioxidants were dissolved in water to obtain a 1% antioxidants solution. Tween-80 was dissolved in the antioxidants solution to obtain a final concentration of 4% Tween-80. The 4% Tween/antioxidants solution was further diluted in antioxidants solution to obtain the desired final concentration of Tween-80. Apomorphine base and glutamic acid were then weighed to obtain the desired molar ratio (e.g., 1:1 to 1:1.5). The prepared Tween-80/antioxidants solution was added and the solution was stirred until dissolution occurred.

Table 5: Composition of organic acid salts of apomorphine

Formulation No. (%)	1	2 (1:1.2)	3 (1:1.2)	4	5	6 (1 eq)	7 (1/2 eq)	8	9	10
APO-GO (μl)										2500
APO base (mg)	50	50	50	25	50	50	50	50	25	
Glutamic acid 14.7 mg/ml (μl)	1873	2250	2250	938						
Aspartic acid 13.3 mg/ml (μl)					1875				938	
Tartaric acid 15 mg/ml (μl)						1875	938			

Formulation No. (%)	1	2 (1:1.2)	3 (1:1.2)	4	5	6 (1 eq)	7 (1/2 eq)	8	9	10
Malic acid 13.4 mg/ml (μl)								1875		
Meglumine 60 mg/ml (μl)			125			188				
Water (μl)	627	250	125	1560	625	439	1562	625	1560	
Total volume (μl)	2500	2500	2500	2500	2500	2500	2500	2500	2500	
APO (mg/ml) expected	20	20	20	10	20	20	20	20	10	10
pH	4.68	4.40	4.76	4.69	4.93	3.45	3.98	3.98	4.59	3.90

Example 8. Comparison of stability between apomorphine-glutamate solutions and commercially available apomorphine-HCl solutions

[0074] The stability of apomorphine-glutamate formulations containing 1% apomorphine, was compared to APO-GO®, a commercially available form of 10mg/ml (1%) apomorphine-HCl solution. Comparisons were performed under physiological conditions approximated by diluting the solutions at 1:10 with phosphate-buffered saline solution, pH 7.5 (PBS). **Table 6** shows the composition of the apomorphine-glutamate solutions - referred to as ND1 and ND2 - which were synthesized for comparison to commercially available apomorphine-hydrochloride solution. **Table 7** shows the stability of apomorphine in the apomorphine-hydrochloride and apomorphine-glutamate solutions as measured in milligrams of apomorphine per gram of solution at various time points following synthesis of the apomorphine-glutamate solutions. **Table 8** shows the production of degradation products of apomorphine in the formulated solutions at various time points. **Table 9** shows the stability of apomorphine in the apomorphine-hydrochloride and apomorphine-glutamate solutions as measured in milligrams of apomorphine per gram of solution at various time points following exposure to physiological conditions. **Table 10** shows the production of degradation products of apomorphine in the solutions at various time points following exposure to physiological conditions. Degradation products in Tables 8 and 10 are shown as a percent of apomorphine peak area as measured by high-performance liquid chromatography (HPLC).

Ingredient (%)	ND1	ND2
Apo base (FW 267.32)	1.0	1.0
Glutamic acid (FW 147.13)	0.8	0.8
Tween-80	0.5	2.0
Anti-oxidants	0.6	0.6

Table 7: Apomorphine stability (mg apomorphine/g solution) in spomorphine-glutamate and apomorphine-hydrochloride solutions at different time points

ND1	9.9	10.0	10.0
ND2	10.0	10.1	10.1
Apo-Go®	11.8	ND	ND
ND = Not determined			

Table 8: Degradation products of apomorphine produced in formulations as a percent of peak area (% of apomorphine peak area)

Sample ID	Unknown r.t.: 5.5 min.			Unknown r.t.: 6.9 min.		
	t=0 hours	t=24 hours	t=6 days	t=0 hours	t=24 hours	t=6 days
ND1	0.12	0.12	0.11	0.08	0.08	0.10
ND2	0.09	0.09	0.09	0.07	0.07	0.09
Apo-Go®	0	ND	ND	0.03	ND	ND
ND = Not determined						

Table 9: Apomorphine stability (mg apomorphine/g solution) following exposure to physiological conditions (1:10 dilution in PBS)

Sample ID	t=0 hours	t=24 hours	t=6 days
ND1	0.92	0.94	0.90
ND2	0.95	0.94	0.92
Apo-Go [®] in PBS + 2% Tween-80	1.10	1.04	0.27
Apo-Go [®] in PBS + 0.5% Tween-80	1.11	0.94	NA

ND = Not Applicable; Concentration of apomorphine in Apo-Go[®] solution 6 days after exposure to physiological conditions is shown in gray.

Table 10: Degradation products of apomorphine produced under physiological conditions (1:10 dilution in PBS) as a percent of peak area (% of apomorphine peak area)

Sample ID	Unknown r.t.: 5.5 min.			Unknown r.t.: 6.9 min.			r.t.: 13.1 min.	r.t.: 20.85 min.
	t=0 hours	t=24 hours	t=6 days	t=0 hours	t=24 hours	t=6 days	t=6 days	t=6 days
ND1	0.14	0.15	0.14	0.08	0	0.12	0	0
ND2	0.09	0.10	0.09	0.07	0	0	0	0

Apo-Go [®] in PBS+2% Tween-80	0.26	5.23	36.3	0.10	1.41	4.47	1.02	5.5
Apo-Go [®] in PBS+0.5% Tween-80	0.23	12.77	NA	0.16	2.55	NA	NA	NA

NA = Not Applicable; Concentration of apomorphine in Apo-Go[®] solution 1 and 6 days after exposure to physiological conditions are shown in gray.

[0075] The results clearly demonstrated that under physiological conditions the glutamic acid salt of apomorphine was stable for at least 6 days at room temperature, whereas the commercial product APO-GO[®] was extensively degraded. Tween-80 at a concentration of 0.5% was sufficient to maintain chemical stability of apomorphine-glutamate solution, while even 2% Tween-80 was not sufficient to maintain chemical stability of Apomorphine-HCl (APO-GO[®]) solution.

Example 9. Analysis of formulation components on apomorphine-glutamate solution stability

[0076] The effect of Tween-80, antioxidants, and the molar ratio of glutamic acid to apomorphine on the physical stability of apomorphine-glutamate salt formulations under physiological conditions was tested by diluting the formulations at 1:10 or 1:20 with PBS. The composition of each of the tested formulations (F1-5) is shown in **Table 11**. Each component is shown as a percent of the total solution.

Table 11: Apomorphine glutamate salt formulations

Ingredient (%)	F1	F2	F3	F4	F5
Apomorphine	2.0	2.0	2.0	2.0	2.0
Glutamic acid	1.3	1.3	1.6	1.6	1.6
Tween-80	2.0	2.0	0	2.0	5.0
Antioxidants	0	0.6	0.6	0.6	0.6
Apo:Glut (molar ratio)	1:1.18		1:1.45		

[0077] Samples of each formulation were diluted in freshly prepared PBS as shown in **Tables 12** and **13**. Samples of formulations F1-F3 were diluted in PBS at 1:20 (Table 11). Samples of formulations F4 and F5 were diluted in PBS at 1:10 and 1:20. The appearance of each solution was observed 20 minutes and 5 hours after dilution, and pH and Tween concentration were measured.

Table 12: Observations of formulations F1, F2 and F3 following 1:20 dilution

	Dilution 1:20		
	F1	F2	F3
20 minutes	Micellar	Transparent	Precipitated
5 hours	Blue/Precipitated	Precipitated	Precipitated
pH measured	7.45	7.34	7.31
Tween-80 concentration after dilution	0.1	0.1	0

Table 13: Observations of formulations F4 and F5 following 1:10 or 1:20 dilution

	Dilution 1:10		Dilution 1:20	
	F4	F5	F4	F5
2 hours	Micellar	Micellar	Transparent	Transparent
5 hours	Micellar	Micellar	Micellar	Micellar
pH measured	7.09	7.07	7.34	7.32
Tween-80 concentration after dilution	0.1	0.25	0.1	0.25

[0078] Following dilution of the formulations with PBS, the formulations were physically stable for at least 5 hours only if they contained Tween-80, antioxidants, and glutamic acid at a molar ratio of greater than 1:1.2, apomorphine:glutamic acid. In the absence of Tween-80 and at a molar ratio of 1:1.18, apomorphine:glutamic acid, the formulations were not physically stable at a pH of 7 to 7.5 (see Tables 11 and 12, formulation F3). In the absence of antioxidants, the solution turned blue (see Tables 11 and 12, formulation F1).

Example 10. Analysis of skin inflammation following subcutaneous administration of apomorphine formulations

[0079] Formulations containing 1% apomorphine and glutamic acid at a molar ratio of 1:1.5, were continuously administered subcutaneously to pigs, at a rate of 4 ml/24 h via a mini-pump. Commercially available 1% apomorphine-HCl (Apo-Go®) was also administered separately to allow for comparison of effects generated by administration of each agent. The formulation of each solution administered is shown in **Table 14**. Skin samples were collected 1, 2, or 3 weeks post-infusion and subjected to histopathological evaluation. Results are shown in **Table 15**. The sites that were infused with the apomorphine-glutamate formulations were characterized by minimal to mild chronic subcutaneous inflammation, whereas the commercial apomorphine-HCl preparation exhibited severe subcutaneous inflammation with necrosis (see the gray boxes in **Table 15**). There was no difference in the severity of lesions observed between the sites administered with formulations containing 0.5% or 2% Tween-80 as shown in **Fig. 3**.

Table 14: Formulations for Subcutaneous Administration

		Formulation		
		1	2	Apo-Go®
Ingredient (%)	Apo base	1.00	1.00	-
	Apo-HCl	-	-	1.00
	Glutamic acid	0.82	0.82	-
	Tween-80	0.50	2.00	-
	Antioxidants	0.6	0.6	0.1

Table 15: Histological evaluation of skin exposed to apomorphine formulations

Sample collection	3 weeks post-application		2 weeks post-application		1 week post-application		
Formulation	1	2	1	2	1	2	ApoGo®
	Subcutis						
Chronic Inflammation	2	1	1.5	2	1.75	1.5	4
Cavity with							

Amorphous Eosinophilic Material	0	0	0	0	0.25	0.25	0
Necrosis	0	0	0	0	0	0	2

Grading of histological changes: 0 = No change, 1 = Minimal change, 2 = Mild change, 3 = Moderate change, 4 = Marked change

[0080] As shown in **Tables 16, 17** and **18**, HPLC analysis demonstrated that the stability of the tested formulations containing 0.5% or 2% Tween-80 (identified as ND1 and ND2) was similar between 4 and 6 days after preparation. The average apomorphine concentrations (AVG mg/g) are shown in gray in **Tables 16** and **17**.

Table 16: Analysis of apomorphine solutions 4 days post-formulation preparation

Sample ID	Sample	Weight (mg)	HPLC (ppm)	Dilution	Apomorphine (mg/g)	AVG (mg/g)	STDEV	RSD (%)
ND1	ts1	104.80	106.827	10	10.193	10.17	0.04	0.4
	ts2	103.60	105.046	10	10.140			
ND1a	ts1	102.90	103.895	10	10.097	10.10	NA	NA

ND2	ts1	100.01	99.409	10	9.940	9.97	0.04	0.4
	ts2	97.85	97.852	10	10.000			

Table 17: Analysis of apomorphine solutions 5 days post-formulation preparation

Sample ID	Sample	Weight (mg)	HPLC (ppm)	Dilution	Apomorphine (mg/g)	AVG (mg/g)	STDEV	RSD (%)
ND1a	ts1	96.85	96.968	10	10.012	10.00	0.020	0.20
	ts2	100.46	100.305	10	9.985			
ND2	ts1	98.79	100.305	10	10.153	10.09	0.086	0.85
	ts2	97.40	97.708	10	10.032			

Table 18: HPLC analysis of apomorphine solutions at 0 and 6 days post-formulation preparation

Sample ID	Unknown r.t.: 5.5 min		Unknown r.t.: 6.9 min	
	t=0	t=6 days	t=0	t=6days
ND1	0.08	ND	0.08	ND
ND1a	0.09	0.095	0.08	0.105
ND2	0.09	0.095	0.075	0.105
ND-Not Determined				

Example 11. Effect of Tween-80 on apomorphine solution stability

[0081] The effect of Tween-80 on the physical stability of apomorphine-HCl formulations and apomorphine-glutamate formulations was analyzed. All tested formulations contained proprietary combinations of antioxidants. Observations regarding the color of undiluted solutions and evidence of precipitation in those solutions were recorded after either 4 days or 1 week at 25°C. Additionally, observations regarding evidence of precipitation in diluted solutions were recorded after 4 days at 25°C. A 1:20 dilution of each solution was achieved by placing 2.85 ml of 50 mM PBS at pH 7.5 in a vial and adding 150 µl of each undiluted solution dropwise. The effect of Tween-80 on the stability of undiluted apomorphine-HCl formulations is shown in **Table 19**. The effect of Tween-80 on the stability of apomorphine-HCl formulations diluted 1:20 in PBS is shown in **Table 20**. The effect of Tween-80 on stability of apomorphine-glutamate formulations is shown in **Table 21**. The effect of Tween-80 on the stability of apomorphine-glutamate formulations diluted 1:20 in PBS is shown in **Table 22**.

Table 19: Effect of Tween-80 on stability of undiluted apomorphine-HCl formulations

	1% Apomorphine-HCl, increasing Tween-80 from 0% to 0.5%			2.5% Apomorphine-HCl, increasing Tween-80 from 2% to 4%		
Ingredient	F-1	F-2	F-3	F-4	F-5	F-6
Tween-80 (%)	0	0.25	0.50	2.00	3.00	4.00
Stability after 4 days at 25°C	No color No precipitation	No color No precipitation	No color <i>Precipitated</i>	No color <i>Precipitated</i>	No color No precipitation	No color No precipitation
Stability after 1 week at 25°C	No color <i>Very slight precipitation</i>	No color <i>Precipitated</i>	No color <i>Precipitated</i>	No color <i>Precipitated</i>	No color <i>Slight precipitation</i>	No color <i>Very slight precipitation</i>

Table 20: Effect of Tween-80 on stability of diluted apomorphine-HCl formulations

	1% Apomorphine-HCl, Increasing Tween-80 from 0% to 0.5%			2.5% Apomorphine-HCl, Increasing Tween-80 from 2% to 4%		
Ingredient	F-1	F-2	F-3	F-4	F-5	F-6
Tween-80 (%)	0	0.25	0.50	2.00	3.00	4.00
Appearance after 4 days at 25°C	<i>Slight precipitation</i>	Micellar	Micellar	Not Micellar <i>Very turbid</i>	Not Micellar <i>Very turbid</i>	Not Micellar <i>Very turbid</i>

Table 21: Effect of Tween-80 on stability of undiluted apomorphine-glutamate formulations

	1% Apomorphine-Glutamate, Increasing Tween-80 from 0% to 0.5%			2.5% Apomorphine-Glutamate, Increasing Tween-80 from 0% to 4%			
Ingredient	F-7	F-8	F-9	F-10	F-11	F-12	F-13
Tween-80 (%)	0	0.25	0.50	0	2.00	3.00	4.00
Stability after 4 days at 25°C	No color No precipitation	No color No precipitation	No color No precipitation	Very slightly yellow No precipitation	Very slightly yellow No precipitation	Very slightly yellow No precipitation	Very slightly yellow No precipitation
Stability after 1 week at 25°C	No color No precipitation	No color No precipitation	No color No precipitation	No color No precipitation	No color No precipitation	No color No precipitation	No color No precipitation

Table 22: Effect of Tween-80 on stability of diluted apomorphine-glutamate formulations

	1% Apomorphine-Glutamate, Increasing Tween-80 from 0% to 0.5%			2.5% Apomorphine-Glutamate, Increasing Tween-80 from 0% to 4%			
Ingredient	F-7	F-8	F-9	F-10	F-11	F-12	F-13
Tween-80 (%)	0	0.25	0.50	0	2.00	3.00	4.00
Appearance after 4 days at 25°C	<i>Precipitated</i>	Almost transparent	Micellar	<i>Strongly precipitated</i>	Micellar Turbid	Micellar Turbid	Micellar

[0082] Precipitation occurred within 1 week at room temperature in all of the tested apomorphine-HCl formulations, whereas no precipitation occurred in any of the tested apomorphine-glutamate formulations. However, following dilution at 1:20 with PBS, both apomorphine salts precipitated in the absence of Tween (e.g., see formulations F-1, F-7, and F-10 in Tables 20 and 22). The 2.5% apomorphine-HCl solution could not be prepared without Tween as it did not dissolve.

Example 12. Comparison of apomorphine-glutamate and apomorphine-HCl formulation stability

[0083] UV-HPLC was used to analyze and compare the chemical stabilities of the apomorphine-glutamate and

apomorphine-HCl formulations. **Table 23** summarizes apomorphine recovery after 1 month at 25°C from apomorphine salt formulations that include varying amounts of Tween-80. The results documented in **Table 23** show that the apomorphine-HCl formulation is less stable than the apomorphine-glutamate formulations of the instant invention.

Table 23: Apomorphine recovery from solution after 1 month at 25°C

Apomorphine Salt	Tween-80 (%)	Apomorphine recovery (%)
1% Apomorphine+Glutamic acid	0	99.0
1% Apomorphine+Glutamic acid	0.50	101.7
2.5% Apomorphine+Glutamic acid	0	94.6
2.5% Apomorphine+Glutamic acid	2.00	98.0
2.5% Apomorphine+Glutamic acid	3.00	98.3
2.5% Apomorphine+Glutamic acid	4.00	95.0
1% Apomorphine-HCl	0	89.3
1% Apomorphine-HCl	0	89.5

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [EP1270007A2](#) [0005]

**SAMMENSÆTNINGER, DER OMFATTER APOMORFIN OG ORGANISKE SYRER,
SAMT ANVENDELSER DERAFT****Patentkrav**

1. Flydende farmaceutisk sammensætning, der omfatter apomorfin, en sur aminosyre og
5 en farmaceutisk acceptabel bærer, hvor molforholdet mellem apomorfin og den sure
aminosyre er 1:0,5 til 1:2,5, fortrinsvis 1:1 til 1:1,5, hvor sammensætningen
 - a) er en opløsning,
 - b) har en pH-værdi på mere end 3 ved 25 °C og
 - c) er egnet til subkutan administration.
- 10 2. Farmaceutisk sammensætning ifølge krav 1, hvor den sure aminosyre er
asparaginsyre, glutaminsyre eller en kombination deraf.
3. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor sammensætningen har en
pH-værdi på mellem 3,0 og 5,5 eller mellem 4,0 og 7,5 ved 25 °C.
4. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 3, der omfatter
15 mindst 1 eller 1 til 4 vægt-% apomorfin.
5. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 4, hvor
sammensætningen i det væsentlige ikke har nogen udfældning af faste stoffer i mindst
48 timer ved 25 °C, fortrinsvis hvor sammensætningen er vandig.
6. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 5, der
20 endvidere omfatter et aminosukker, fortrinsvis meglumin, en basisk aminosyre, fortrinsvis
arginin, én eller flere antioxidant, fortrinsvis hvor hver især uafhængigt af hinanden er
valgt blandt ascorbinsyre, Na-ascorbat, L-cystein, N-acetylcystein (NAC), glutathion
(GSH), Na₂-EDTA, Na₂-EDTA-Ca, natriumbisulfid eller en kombination deraf.
7. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 6, der
25 endvidere omfatter et overfladeaktivt middel, fortrinsvis Tween-80.
8. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 7, der
endvidere omfatter et lokalbedøvelsesmiddel og/eller et anti-inflammatorisk middel,
fortrinsvis hvor lokalbedøvelsesmidlet er et amidbaseret lokalbedøvelsesmiddel såsom
lidocain, prilocain, bupivacain, levobupivacain, ropivacain, mepivacain, dibucain og
30 etidocin eller et esterbaseret lokalbedøvelsesmiddel såsom procain, amethocain, kokain,
benzocain og tetracain; og det antiinflammatoriske middel er et ikke-steroidt anti-
inflammatorisk middel såsom diclofenac, ketorolac, salicylater, ibuprofen, piroxicam og
benzydamin eller et steroidt anti-inflammatorisk middel såsom prednison, dexamethason,
betamethason, prednison-hydrocortison eller et salt deraf.

9. Flydende eller halvfast farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 8 til anvendelse ved behandling af en neurologisk sygdom eller lidelse eller en bevægelsessygdom eller -lidelse såsom Parkinsons sygdom, Alzheimers sygdom og akinesi eller en tilstand, som associeres dermed, såsom alkoholisme, opiatafhængighed og erektil dysfunktion, fortrinsvis til behandling af Parkinsons sygdom.
- 5

DRAWINGS

Fig. 1





Fig. 2

Fig. 3

