



(86) **Date de dépôt PCT/PCT Filing Date:** 2015/06/19
(87) **Date publication PCT/PCT Publication Date:** 2015/12/23
(85) **Entrée phase nationale/National Entry:** 2016/12/13
(86) **N° demande PCT/PCT Application No.:** AU 2015/000360
(87) **N° publication PCT/PCT Publication No.:** 2015/192172
(30) **Priorité/Priority:** 2014/06/20 (AU2014902374)

(51) **Cl.Int./Int.Cl. A01N 63/00** (2006.01),
C12N 1/20 (2006.01)
(71) **Demandeur/Applicant:**
THE FLINDERS UNIVERSITY OF SOUTH AUSTRALIA,
AU
(72) **Inventeurs/Inventors:**
FRANCO, CHRISTOPHER MILTON MATHEW, AU;
LE, HOANG XUYEN, AU;
BALLARD, ROSS ALEXANDER, AU
(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : INOCULANTS ET PROCEDES POUR LES UTILISER**
(54) **Title: INOCULANTS AND METHODS FOR USE THEREOF**

(57) **Abrégé/Abstract:**

The present invention relates to methods for enhancing at least one growth parameter of a leguminous plant via co-inoculation of a leguminous plant with at least one rhizobial microorganism together with at least one actinobacterial microorganism. In further aspects, the present invention also relates to leguminous plants co-inoculated with at least one rhizobial microorganism together with at least one actinobacterial microorganism, as well as specific actinobacterial strains and inoculant compositions which are useful in accordance with the present invention.



- (51) International Patent Classification:

A01N 63/00 (2006.01) C12R 1/465 (2006.01)

C12N 1/20 (2006.01)

(21) International Application Number:

PCT/AU2015/000360

(22) International Filing Date:

19 June 2015 (19.06.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2014902374 20 June 2014 (20.06.2014) AU

(71) Applicant: THE FLINDERS UNIVERSITY OF SOUTH AUSTRALIA [AU/AU]; Sturt Road, Bedford Park, South Australia 5042 (AU).

(72) Inventors: FRANCO, Christopher, Milton, Mathew; 1 Ayrbank Avenue, Stonyfell, South Australia 5066 (AU). LE, Hoang, Xuyen; Unit 49/2 Ayliffes Road, St Marys, South Australia 5042 (AU). BALLARD, Ross, Alexander; 21 Linfield Avenue, Belair, South Australia 5052 (AU).

(74) Agent: PHILLIPS ORMONDE FITZPATRICK; Level 16, 333 Collins Street, Melbourne, Victoria 3000 (AU).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))
- (54) Title: INOCULANTS AND METHODS FOR USE THEREOF
- (57) Abstract: The present invention relates to methods for enhancing at least one growth parameter of a leguminous plant via co-inoculation of a leguminous plant with at least one rhizobial microorganism together with at least one actinobacterial microorganism. In further aspects, the present invention also relates to leguminous plants co-inoculated with at least one rhizobial microorganism together with at least one actinobacterial microorganism, as well as specific actinobacterial strains and inoculant compositions which are useful in accordance with the present invention.
- WO 2015/192172 A1

INOCULANTS AND METHODS FOR USE THEREOF

5

PRIORITY CLAIM

This application claims priority to Australian provisional patent application 2014902374, filed 20 June 2014, the content of which is hereby incorporated by
10 reference.

FIELD

The present invention relates to methods for enhancing at least one growth
15 parameter of a leguminous plant via co-inoculation of a leguminous plant with at least one rhizobial microorganism together with at least one actinobacterial microorganism. In further aspects, the present invention also relates to leguminous plants co-inoculated with at least one rhizobial microorganism together with at least one actinobacterial microorganism, as well as specific
20 actinobacterial strains and inoculant compositions which are useful in accordance with the present invention.

BACKGROUND

25 There are around 44 to 66 million tonnes of nitrogen fixed from atmospheric nitrogen by symbiosis of rhizobial microorganisms and legumes every year, which is nearly half of the nitrogen used in agriculture around the world.

The interaction of legume and rhizobia is highly specific and each rhizobial
30 species has a distinct range of leguminous plant hosts for forming nodules and fixing nitrogen. Different steps of the nodulation process requires the exchange of various signals between host legumes and rhizobia.

Actinobacteria are a large group that includes different genera of Gram-positive bacteria with a high G-C content in their DNA. Actinobacteria are widely distributed in terrestrial environments and some, like the nitrogen-fixing symbionts *Frankia*, are known to form associations with plants through symbiotic relationships. Recent studies have also found that endophytic actinobacteria produced plant growth-promoting compounds such as indole-3-acetic acid (IAA) and siderophores.

Effects of actinobacteria on rhizobia and symbiosis with legumes have been noticed but not many studies have investigated this complex combination, although several studies have suggested antagonism occurring between actinobacteria and rhizobia.

For example, Antoun *et al.* (*Canadian Journal of Microbiology* 24: 558-562, 1978) disclosed antagonism tests between actinobacteria isolated from different soils and effective strains of rhizobia. These data demonstrated that some actinobacteria inhibit the growth of rhizobia *in vitro* and *in planta*. Specifically, thirty one percent of the 481 actinobacteria investigated inhibited two efficient rhizobia strains, *Rhizobium meliloti* A2 and S14.

In a further study, Damirgi and Johnson (*Agronomy Journal* 58: 223-224, 1966) disclosed that the number of nodules on soybeans inoculated with *Rhizobium japonicum* strains 122 and 123 in autoclaved soil were reduced by up to 35% and 53%, respectively, by treatment with the actinobacterium E8. They also isolated about 60 actinobacteria from one soil sample where there had been poor nodulation of clovers. However, 20 of 24 actinobacteria isolated from an experimental soybean field did not inhibit eight sensitive *R. japonicum* strains in *in vitro* tests.

30

Antagonism was also examined between actinobacteria and 12 strains of rhizobia from five soil samples by Patel (*Plant and Soil* 41: 395-402, 1974). It

- 3 -

was reported that about 23-70% of the actinobacteria inhibited the rhizobia strains.

In light of the above, identification of actinobacterial inoculants that are compatible with rhizobial microorganisms and that can enhance the growth and development of leguminous plants would be desirable.

DESCRIPTION

Nucleotide and amino acid sequences are referred to herein by a sequence identifier number (SEQ ID NO:). A summary of the sequence identifiers is provided below:

Sequence Identifier	Description
SEQ ID NO: 1	27f primer nucleotide sequence
SEQ ID NO: 2	1465r primer nucleotide sequence
SEQ ID NO: 3	LuP3 16S rRNA gene nucleotide sequence
SEQ ID NO: 4	LuP12A 16S rRNA gene nucleotide sequence
SEQ ID NO: 5	LuP30 16S rRNA gene nucleotide sequence
SEQ ID NO: 6	LuP47B 16S rRNA gene nucleotide sequence
SEQ ID NO: 7	EN23 16S rRNA gene nucleotide sequence
SEQ ID NO: 8	EN27 16S rRNA gene nucleotide sequence
SEQ ID NO: 9	LuP8 16S rRNA gene nucleotide sequence
SEQ ID NO: 10	LuP44 16S rRNA gene nucleotide sequence

A sequence listing is also provided at the end of the specification.

In a first aspect, the present invention provides a method for enhancing at least one growth parameter of a leguminous plant, the method comprising co-inoculating the leguminous plant with:

- at least one rhizobial microorganism; and
- at least one actinobacterial microorganism;

wherein the co-inoculated leguminous plant has at least one enhanced growth parameter relative to a leguminous plant of the same taxon that has not been co-inoculated.

- 5 An “actinobacterial microorganism” as referred to herein should be understood to include any microorganism of the phylum Actinobacteria.

In some embodiments the actinobacterial microorganism is an endophytic actinobacterial microorganism. An “endophytic” actinobacterial organism should
10 be understood to include any actinobacterial organism that is able to live within a plant for at least a part of its lifecycle without causing apparent disease. In some embodiments, an endophyte may also be able to complete its lifecycle in the absence of a plant host, and thus only be an opportunistic endophyte. In some embodiments, an endophytic actinobacterial microorganism refers to an
15 actinobacterial microorganism which may be isolated from surface-sterilised healthy plant tissue. For reference, an example of isolation of actinobacterial endophytes from surface sterilised plant tissue is set out in Coombs & Franco (*Appl. Environ. Micro.* 69(9): 5603-5608, 2003).

- 20 In some embodiments, the actinobacterial microorganism is from a genus selected from *Streptomyces*, *Microbispora* or *Micromonospora*.

In some embodiments, the actinobacterial microorganism is from the genus *Streptomyces*.

25

In some embodiments, the actinobacterial microorganism is from a species selected from:

Streptomyces drozdowiczii;

Streptomyces ciscaucasicus;

30

Streptomyces canus;

Streptomyces rishiriensis;

Streptomyces badius; or

Streptomyces parvus

In some embodiments, the actinobacterial microorganism is from a species selected from *Streptomyces ciscaucasicus*, *Streptomyces canus* or
5 *Streptomyces rishiriensis*.

In some embodiments, the actinobacterial microorganism comprises a 16S rRNA gene nucleotide sequence which is at least 90% identical to one or more of: SEQ ID NO: 3; SEQ ID NO: 4; SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7,
10 SEQ ID NO: 8, SEQ ID NO: 9 and/or SEQ ID NO: 10.

In some embodiments, the actinobacterial microorganism comprises a 16S rRNA gene nucleotide sequence which is at least 90% identical to one or more of SEQ ID NO: 5 and/or SEQ ID NO: 6.

15

In some embodiments the actinobacterial microorganism comprises a 16S rRNA gene nucleotide sequence which is at least 90%, at least 91%, at least 91.5%, at least 92%, at least 92.5%, at least 93%, at least 93.5%, at least 94%, at least 94.5%, at least 95%, at least 95.5%, at least 96%, at least 96.5%, at
20 least 97%, at least 97.5%, at least 98%, at least 98.1%, at least 98.2% at least 98.3%, at least 98.4%, at least 98.5%, at least 98.6%, at least 98.7%, at least 98.8%, at least 98.9%, at least 99%, at least 99.1%, at least 99.2% at least 99.3%, at least 99.4%, at least 99.5%, at least 99.6%, at least 99.7%, at least 99.8% at least 99.9% or 100% sequence identity to a comparison window of
25 one or more of: SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9 and/or SEQ ID NO: 10.

When comparing nucleic acid sequences to calculate a percentage identity, the compared nucleic acid sequences should be compared over a comparison
30 window of, for example, at least 100 nucleotide residues, at least 300 nucleotide residues, at least 600 nucleotide residues, at least 1000 nucleotide residues, at least 1100 nucleotide residues, at least 1200 nucleotide residues,

at least 1300 nucleotide residues or at least 1400 nucleotide residues. In some embodiments, the comparison window may comprise the region in each of the compared nucleotide sequences between and including the binding sites of the 27f primer (SEQ ID NO: 1) and the 1465r primer (SEQ ID NO: 2) on the
5 compared nucleotide sequences.

The comparison window may comprise additions or deletions (ie. gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences.
10 Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms such as the BLAST family of programs as, for example, disclosed by Altschul *et al.* (*Nucl. Acids Res.* 25: 3389-3402, 1997). A detailed discussion of sequence analysis can be found in Unit 19. 3 of Ausubel *et al.* (*Current Protocols in Molecular Biology*,
15 John Wiley & Sons Inc, 1994-1998, Chapter 15,1998).

A number of particularly useful actinobacterial microorganisms of the present invention have been deposited in accordance with the provisions of the Budapest Treaty on the International Recognition of the Deposit of
20 Microorganisms for the Purposes of Patent Procedure.

The first deposited microorganism, referred to herein as *Streptomyces* sp. LuP30, has been deposited at the National Measurement Institute (NMI), Australia on 12 December 2013 under accession number V13/030101.
25

Accordingly, some embodiments, the actinobacterial microorganism is *Streptomyces* sp. LuP30 as deposited with the National Measurement Institute, Australia under accession number V13/030101; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth
30 parameter of a leguminous plant when the actinobacterial microorganism co-inoculated onto a leguminous plant with with a rhizobial microorganism.

- 7 -

The second deposited microorganism, referred to herein as *Streptomyces* sp. LuP47B, has been deposited at the National Measurement Institute (NMI), Australia on 12 December 2013 under accession number V13/030100.

5 Accordingly, in some embodiments, the actinobacterial microorganism is *Streptomyces* sp. LuP47B as deposited with the National Measurement Institute, Australia under accession number V13/030100; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial
10 microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

In addition, in accordance with the present invention, two previously known organisms were also surprisingly identified to be particularly useful in
15 accordance with the method of the present invention:

In some embodiments the actinobacterial microorganism is *Streptomyces* sp. EN23 as described in PCT publication WO/2005/003328 and deposited as AGAL Deposit No. NM03/35605; or a mutant or derivative of said
20 microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

In some embodiments the actinobacterial microorganism is *Streptomyces* sp. EN27 as described in PCT publication WO/2005/003328 and deposited as AGAL Deposit No. NM03/35606; or a mutant or derivative of said
25 microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

30

A “mutant or derivative” of the subject actinobacterial microorganisms should be understood to encompass, for example, any spontaneous or induced mutant,

conjugation progeny or genetically modified form of the deposited strains which retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

5

Mutagenisation techniques that may be used to generate derivatives or mutants include, for example, chemical mutagenesis (eg. EMS mutagenesis), ionising radiation-induced mutagenesis (eg. X-ray mutagenesis, γ -ray mutagenesis and UV mutagenesis), genetic insertion mutagenesis methods (eg. transposon mutagenesis) and the like.

10

As set out above, the present invention contemplates a method for enhancing at least one growth parameter of a leguminous plant.

15 A “growth parameter” of a leguminous plant, as referred to herein, can include any measurable characteristic of the leguminous plant.

In some embodiments, the growth parameter is a length and/or mass of a shoot of the leguminous plant.

20

In some embodiments, the growth parameter is a length and/or mass of a root of the leguminous plant.

In some embodiments, the growth parameter is a number and/or mass of nodules of the leguminous plant.

25

In some embodiments, the growth parameter is a number and/or mass of seed pods and/or seed produced by the leguminous plant.

30 In some embodiments, the growth parameter is a concentration and/or amount of a nutrient in the leguminous plant.

In some embodiments, the nutrient is selected from: Boron, Calcium, Copper, Magnesium, Manganese, Phosphorous, Sodium, Sulphur, Nitrogen and/or Zinc.

5 The concentration and/or amount of the nutrient may be measured using any method known in the art to be suitable for the relevant nutrient. Such methods may include, for example, the methods described by: Kirsten (*Organic Elemental Analysis – Ultramicro, Micro and Traces Methods*. Academic Press, New York, 1984); Horwath (*Instrumental Organic Analysis*. Academic Press, New York, 1977); Colombo and Giazzi (*American Laboratory* 38-45, 1982);
10 Fraisse and Schmidt (*J. Microchem.* 22: 109-130, 1977); Hegedus (*Microchim. Acta* 441-446, 1977); and Baur and Dirscherl (*Microchim. Acta* 1: 299-244, 1980).

In some embodiments, the nutrient is Nitrogen.

15

In some embodiments, the growth parameter is a germination rate of a leguminous plant seed. In some embodiments, “germination rate” may refer to the proportion of seeds of a leguminous plant that successfully germinate. In some embodiments “germination rate” may refer to a speed of germination of a
20 leguminous plant seed, and/or a proportion of seeds of a leguminous plant that successfully germinate per unit time, for example the proportion of seeds of a leguminous plant that successfully germinate per 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19 or 20 days.

25 The “germination rate” of a leguminous plant seed may be assessed using any suitable laboratory based or field based method, as would be readily ascertained by those skilled in the art.

30 As set out above, the present invention contemplates “enhancement” of the one or more growth parameters of the leguminous plant. “Enhancement” of a growth parameter should be understood to include any improvement in a growth parameter in a co-inoculated leguminous plant relative to a leguminous plant of

- 10 -

the same taxon that has not been co-inoculated in accordance with the method of the present invention.

In some embodiments, enhancement of a growth parameter will include an increase in the measured value of the growth parameter. For example, an increase in any of:

- a length and/or mass of a shoot;
- a length and/or mass of a root;
- a number and/or mass of nodules;
- 10 a number and/or mass of seed pods and/or seed;
- a concentration and/or amount of a nutrient; or
- a germination rate.

should be considered an enhancement of such growth parameters.

15 In some embodiments, enhancement of a growth parameter may comprise enhancement within a particular time period. For example, in some embodiments, enhancement of the growth parameter may comprise enhancement over a time period of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 20 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 60, 70, 80, 90 or 100 days.

In some embodiments, an “increase” in a growth parameter may include, for example, a 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 25 2-fold, 5-fold, 10-fold, 20 fold, 50-fold, 100-fold increase in the growth parameter in a co-inoculated leguminous plant relative to a leguminous plant of the same taxon that has not been co-inoculated.

In some embodiments, however, “enhancement” of the growth parameter may include a decrease in the measured value of the growth parameter. For example a decrease in the concentration and/or amount of a pathogen, disease symptom and/or toxin in the plant, and or a decrease in the time of germination 30

- 11 -

of a leguminous plant seed, may be considered “enhancement” of such growth parameters.

In some embodiments, an “decrease” in a growth parameter may include, for example, a 1%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% decrease in the growth parameter in a co-inoculated leguminous plant relative to a leguminous plant of the same taxon that has not been co-inoculated.

As set out above, the present invention contemplates co-inoculating the leguminous plant with at least one rhizobial microorganism and at least one actinobacterial microorganism.

A “rhizobial microorganism” as referred to herein may include any microorganism that is capable of fixing nitrogen after becoming established in a root nodule of a leguminous plant.

Rhizobial microorganisms are a paraphyletic group that generally fall into two classes of the proteobacteria, the alpha- and beta-proteobacteria. Most rhizobial microorganisms belong to the order Rhizobiales, but several rhizobia occur in distinct bacterial orders of the proteobacteria.

Examples of rhizobial microorganisms include:

Bradyrhizobium spp., such as *B. canariense*, *B. elkanii*, *B. japonicum*, *B. liaoningense* and *B. yuanmingense*;

Ochrobactrum spp., such as *O. cytisi* and *O. lupini*;

Azorhizobium spp., such as *A. caulinodans* and *A. doebereineriae*;

Devosia spp., such as *D. neptuniae*;

Methylobacterium spp., such as *M. nodulans*;

Mesorhizobium spp., such as *M. albiziae*, *M. amorphae*, *M. chacoense*, *M. cicero*, *M. huakuii*, *M. loti*, *M. mediterraneum*, *M. plurifarum*, *M. septentrionale*, *M. temperatum*, and *M. tianshanense*;

Phyllobacterium spp., such as *P. ifriqiyense*, *P. leguminum*, and *P. trifoli*;
Rhizobium spp., such as *R. cellulosilyticum*, *R. daejeonense*, *R. etli*, *R. galegae*, *R. gallicum*, *R. giardinii*, *R. hainanense*, *R. huautlense*, *R. indigoferae*,
R. leguminosarum, *R. loessense*, *R. lupini*, *R. lusitanum*, *R. mongolense*, *R.*
5 *miluonense*, *R. sullae*, *R. tropici*, *R. undicola* and *R. yanglingense*;

Sinorhizobium spp. such as *S. abri*, *S. adhaerens*, *S. americanum*, *S. arboris*, *S. fredii*, *S. indiaense*, *S. kostiense*, *S. kummerowiae*, *S. medicae*, *S. meliloti*, *S. mexicanus*, *S. morelense*, *S. saheli*, *S. terangae* and *S. xinjiangense*;

10 *Ensifer* spp.;

Burkholderia spp., such as *B. caribensis*, *B. dolosa*, *B. mimosarum*, *B. phymatum* and *B. tuberum*;

Cupriavidus spp., such as *C. taiwanensis*; and

Herbaspirillum spp., such as *H. lusitanum*.

15

In some embodiments, the rhizobial microorganism is a *Rhizobium* spp., a *Sinorhizobium* spp. or *Ensifer* spp., or a *Bradyrhizobium* spp.

In some embodiments, the rhizobial microorganism is of the species
20 *Sinorhizobium meliloti* or *Sinorhizobium medicae*. In some embodiments, the rhizobial microorganism is selected from the list of: *S. meliloti* strain RRI128 (referred to hereafter as 'RRI128'), *Sinorhizobium* strain SRDI736 (referred to hereafter as 'SRDI736') or *S. medicae* strain WSM1115G (referred to hereafter as 'WSM1115G').

25

In some embodiments, the rhizobial microorganism is of the species *Rhizobium leguminosarum*. In some embodiments, the rhizobial microorganism is selected from the list of: *R. leguminosarum* bv. *viciae* (referred to hereafter as 'WSM1455') or *R. leguminosarum* bv. *trifolii* (referred to hereafter as
30 'WSM1325').

In some embodiments, the rhizobial microorganism is of the species

Bradyrhizobium sp. *lupini* or *Bradyrhizobium japonicum*. In some embodiments, the rhizobial microorganism is selected from the list of: *Bradyrhizobium* sp. *lupini* strain WSM471 (referred to hereafter as 'WSM471') or *Bradyrhizobium japonicum* strain CB1809 (referred to hereafter as 'CB1809').

5

A range of rhizobial microorganisms are available from a range of commercial culture collections, as would be readily ascertained by those skilled in the art. In relation to a range of the rhizobial microorganisms described herein, these organisms can be accessed from the rhizobium culture collection of the South
 10 Australian Research & Development Institute (Plant Research Centre, Hartley Grove, Urrbrae SA 5064, Australia; www.sardi.sa.gov.au).

As set out above, the present invention contemplates a method for enhancing at least one growth parameter of a leguminous plant.

15

A "leguminous plant" as referred to herein should be understood as any member of the Fabaceae (or Leguminosae) that can form nodules when infected with a rhizobial microorganism.

20 Examples of leguminous plants include:

Medicago spp., such as *Medicago sativa* (also referred to as lucerne or alfalfa);

Pisum spp., such as *Pisum abyssinicum* (syn. *P. sativum* subsp. *abyssinicum*), *Pisum fulvum*, *Pisum sativum*, *Pisum sativum* subsp. *elatius*
 25 (syn. *P. elatius*, *P. syriacum*) and *Pisum sativum* subsp. *sativum*;

Glycine spp., such as *Glycine max*, *Glycine albicans*, *Glycine aphyonota*, *Glycine arenaria*, *Glycine argyrea*, *Glycine canescens*, *Glycine clandestine*, *Glycine curvata*, *Glycine cyrtoloba*, *Glycine falcate*, *Glycine gracei*, *Glycine hirticaulis*, *Glycine hirticaulis* subsp. *leptosa*, *Glycine lactovirens*, *Glycine*
 30 *latifolia*, *Glycine latrobeana*, *Glycine microphylla*, *Glycine montis-douglas*, *Glycine peratosa*, *Glycine pescadrensis*, *Glycine pindanica*, *Glycine pullenii*,

- 14 -

Glycine rubiginosa, *Glycine stenophita*, *Glycine syndetika*, *Glycine tabacina*,
Glycine tomentella and *Glycine soja*;

Cicer spp., such as *Cicer arietinum*;

Vicia spp., such as *V. faba*;

5 *Vigna* spp., such as *V. aconitifolia*, *V. angularis*, *V. mungo*, *V. radiata*, *V. subterranean*, *V. umbellatta* or *V. unguiculata*

Lathyrus spp., such as *Lathyrus sativus* or *Lathyrus tuberosus*;

Lens spp., such as *L. culinaris*

Lablab spp., such as *L. purpureus*;

10 *Phaseolus* spp., such as *P. acutifolius*, *P. coccineus*, *P. lunatus*, *P. vulgaris*, *P. polyanthus* or *P. Dumosus*;

Psophocarpus spp., such as *P. tetragonolobus*;

Cajanus spp., such as *C. cajan*;

Stizolobium spp.;

15 *Cyamopsis* spp., such as *C. tetragonoloba*;

Canavalia spp., such as *C. ensiformis* or *C. gladiata*;

Macrotyloma spp., such as *M. uniflorum*;

Lupinus spp., such as *L. mutabilis* or *L. albus*; or

Erythrina spp., such as *E. herbacea*.

20 *Trifolium* spp., such as *Trifolium subterraneum*

In some embodiments, the leguminous plant is a *Medicago* spp. plant. In some
embodiments, the leguminous plant is a *Medicago sativa*, lucerne or alfalfa
plant.

25

In some embodiments, the leguminous plant is a *Trifolium* sp. In some
embodiments, the leguminous plant is a *Trifolium subterraneum* plant.

In some embodiments, the leguminous plant is *Pisum* sp. In some
30 embodiments, the leguminous plant is a *Pisum sativum* plant.

- 15 -

In some embodiments, the leguminous plant is a *Glycine* sp. In some embodiments, the leguminous plant is a *Glycine max* plant.

As set out above, the present invention contemplates co-inoculating a leguminous plant with a rhizobial microorganism and an actinobacterial microorganism. As referred to herein, “co-inoculating” should be understood to include any method or process wherein a leguminous plant (including a leguminous plant seed) is brought into contact with a rhizobial microorganism and an actinobacterial microorganism. In some embodiments co-inoculation may comprise the rhizobial microorganism and/or actinobacterial microorganism being applied to a leguminous plant seed. Examples of leguminous plant seed inoculation are described by Hartley *et al.* (*Crop and Pasture Science* 63: 858-865, 2012). In some embodiments co-inoculation may comprise the rhizobial microorganism and/or actinobacterial microorganism being applied to soil in which a leguminous plant is growing. In some embodiments, co-inoculation may comprise the rhizobial microorganism and/or actinobacterial microorganism being applied to root and/or shoot tissue of a leguminous plant.

In some embodiments, “co-inoculating” may also comprise where the actinobacterial microorganism or rhizobial microorganism is pre-existing in the environment (eg. soil) into which a leguminous plant is grown. For example, co-inoculation may comprise application of an actinobacterial microorganism to a plant or soil and wherein a natural or pre-existing rhizobial microorganism in the soil co-inoculates the plant.

In some embodiments, the leguminous plant is exposed to a pathogen and, when exposed to the pathogen, the co-inoculated leguminous plant has at least one enhanced growth parameter relative to a leguminous plant of the same taxon that has not been co-inoculated.

- 16 -

In some embodiments, the pathogen is a root pathogen. A “root pathogen” as referred to herein should be understood to include any leguminous plant pathogen that infects and/or damages the roots of the leguminous plant. Examples of root pathogens of leguminous plants include fungal, oomycete, bacterial and/or nematode pathogens.

In some embodiments, the root pathogen is a nematode root pathogen such as a *Pratylenchus* spp., such as *P. neglectus* or *P. thornei*.

In some embodiments, the root pathogen is an oomycete root pathogen such as a *Phytophthora* spp., such as *P. sojae*.

In some embodiments, the pathogen is a fungal pathogen. Leguminous plants may be subject to attack by a range of fungal pathogens, including fungal root pathogens. Examples of such pathogens include *Rhizoctonia* spp., *Pythium* spp. or *Aphanomyces* spp.

In some embodiments, the pathogen is a *Rhizoctonia* sp. In some embodiments, the pathogen is *Rhizoctonia solani*. In some embodiments the pathogen is *Rhizoctonia solani* AG8.

In a second aspect, the present invention provides a leguminous plant, a leguminous plant part or leguminous plant reproductive material, co-inoculated with at least one rhizobial microorganism and at least one actinobacterial microorganism.

The leguminous plant contemplated in the second aspect of the invention may be any leguminous plant as hereinbefore described with reference to the first aspect of the invention.

30

Reference herein to a plant, plant part or plant reproductive material should be understood to encompass tissues, organs, whole organisms and parts thereof.

In some embodiments, the term plant, plant part, or plant reproductive material should also be understood to encompass multicellular aggregations of cultured cells such as colonies, plant calli, suspension cultures and the like.

- 5 In some embodiments, the leguminous plant part or plant reproductive material may include a leguminous plant seed. As referred to herein, a plant “seed” should be understood to refer to a mature or immature plant seed. As such, the term “seed” includes, for example, immature seed carried by a maternal plant or seed released from the maternal plant. In some embodiments, the term “seed”
10 may encompass any seed plant sporophyte between the developmental stages of fertilisation and germination.

In some embodiments, the actinobacterial microorganism contemplated in accordance with the second aspect of the invention is an actinobacterial
15 microorganism as hereinbefore described with reference to the first aspect of the invention.

In some embodiments, the rhizobial microorganism contemplated in accordance with the second aspect of the invention is a rhizobial
20 microorganism as hereinbefore described with reference to the first aspect of the invention.

In a third aspect, the present invention provides an isolated actinobacterial microorganism as deposited with the National Measurement Institute, Australia
25 under accession number V13/030101; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

30 In a fourth aspect, the present invention provides an isolated actinobacterial microorganism as deposited with the National Measurement Institute, Australia under accession number V13/030100; or a mutant or derivative of said

microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

- 5 In some embodiments, the present invention provides any of the hereinbefore described isolated actinobacterial microorganisms when used according to the method of the first aspect of the invention.

10 In a fifth aspect, the present invention also provides an inoculant composition comprising an actinobacterial microorganism.

In some embodiments, the actinobacterial microorganism comprises an actinobacterial microorganism as hereinbefore described with respect to the first aspect of the invention.

15

In some embodiments, the inoculant composition further comprises a rhizobial microorganism. In some embodiments, the rhizobial microorganism comprises a rhizobial microorganism as hereinbefore described with respect to the first aspect of the invention.

20

In some embodiments, the inoculant composition comprises a carrier or additive. The carrier or additives used will depend on the nature of the inoculant composition. For example, the inoculant composition may be in the form of a liquid composition, a solid composition (such as a powder, pellet or granular composition) a seed coating or the like.

25

The inoculant compositions of the present invention may be adapted to be applied to a leguminous plant in any suitable way. For example, the inoculant composition could be adapted to be applied as a seed coating, applied as a solid or liquid composition to the foliage or roots of a plant, or applied as a solid or liquid composition to soil before, during or after sowing of a leguminous plant.

30

A range of useful carriers or additives would be readily apparent to those of skill in the art and may include, for example: one or more gums (including xanthan gum), clay or peat based carriers, one or more nutrients including carbon or nitrogen sources, one or more antifungal or antibacterial agents, one or more seed coating agents, one or more wetting agents and the like.

In some embodiments, the present invention provides the hereinbefore described inoculant composition when used according to the method of the first aspect of the invention.

The present invention is further described with reference to the following non-limiting examples:

BRIEF DESCRIPTION OF THE FIGURES AND TABLES

FIGURE 1 shows the cultivable actinobacterial endoflora isolated from roots and nodules of different legumes – including pea, lucerne, clover and medics.

FIGURE 2 shows the beneficial effects of endophytic actinobacteria on germination of lucerne seeds on agar plates after 36 hours incubation. The plate on the left shows surface-sterilised seeds with 0.85% saline; the plant on the right shows surface-sterilised seeds coated with a spore suspension of LuP83 in 0.85% saline.

FIGURE 3 shows indole acetic acid production by selected endophytic actinobacteria.

FIGURE 4 shows the stimulation of the growth of *Rhizobium leguminosarum* bv. *trifolii* strain WSM 1325 and *Bradyrhizobium lupini* strain WSM 471 by two actinobacteria LuP30 and LuP47B after 5 days growth of the rhizobia on YMA medium at different concentrations of rhizobium. Panel A - From top to bottom

- 20 -

and left to right: WSM 1325 at 2 weeks old with 10^3 CFU per plate; LuP30 plugs on WSM 1325 with bigger colonies; WSM 471 at 7 days old with 10^5 CFU per plate (A) ISP2 control plugs, (B) LuP47B plugs and (C) LuP30 plugs; WSM 1325 at 7 days old with 10^5 CFU per plate. Panel B - From right to left images the increase of the growth of two rhizobia when closer to the plug of LuP30 or LuP47B shown under microscopy.

FIGURE 5 shows enhanced plant growth, nodule sizes and nitrogen fixation in lucerne plants co-inoculated with selected endophytic actinobacteria and *S. meliloti* RRI128. (Rhi. = Rhizobium RRI128). (A) and (B), plants and nodules in the seventh week after adding the rhizobia. (C) and (D), plants and nodules in 45 days after adding the rhizobia.

FIGURE 6 shows the total nitrogen fixed in the shoots of lucerne inoculated with endophytic actinobacteria EN23, LuP30 and LuP47B together with rhizobium RRI128.

FIGURE 7 shows the effect of endophytic actinobacteria on the symbiosis of rhizobia and lucerne under nutrient limited conditions. Two surface sterilised lucerne seeds were sown into 65 g of autoclaved washed sand in a 50 ml tube containing 10mls of McKnight's + N starter (300 mg for 20 L) was added at day 0 and MQ water as required later. One seedling was kept in each tube with 12h light and 12h dark and five replicates for each treatment. One ml (around 10^8 cfu/ml) of the *S. meliloti* RRI 128 was added into each seedling for treatment plants after 6 days, and the plants were grown up to 7 weeks. Left plants were inoculated with RRI 128 only while right plants were co-applied with LuP5B and RRI 128

FIGURE 8 shows the response of lucerne plants co-inoculated with LuP30 or LuP47B and different concentrations of *S. meliloti* RRI 128. Panel A is a graphical representation showing shoot, root and total plant dry weights after three weeks at three rhizobia concentrations: (A) 5×10^2 , (B) 5×10^4 , (C) 5×10^6

CFU.ml⁻¹. Error bars: Mean $\pm$ S.E. Panel B is a photograph of representative lucerne plants in tubes three weeks after inoculation with *S. meliloti* RRI 128 at 5×10^2 CFU.ml⁻¹. Left- *S. meliloti* RRI 128 alone; right- *S. meliloti* RRI 128 plus LuP30.

5

FIGURE 9 is a graphical representation showing lucerne shoot dry weight response by impact of LuP30 and LuP47B after 10, 21 and 35 days inoculation with *S. meliloti* RRI 128. Asterisks indicate significant differences at $P < 0.05$ (*) or $P < 0.01$ (**).

10

FIGURE 10 is a graphical representation showing accumulation of N (¹⁴N and ¹⁵N) in lucerne plants inoculated with rhizobia and actinobacteria.

FIGURE 11 is a graphical representation showing the effects of LuP30 and LuP47B on growth and nodulation of clover when co-inoculated with *Rhizobium* WSM 1325. Asterisks indicate significant differences at $p \leq 0.05$ (*) or $p \leq 0.01$ (**).

TABLE 1 shows indole acetic acid production and phosphate solubilising activity of selected endophytic actinobacteria. (+) positive production; (-) no production.

TABLE 2 shows *in-vitro* interaction assay results between selected endophytic actinobacteria and three different rhizobia at various concentrations. (++) positive effects on growth of the rhizobium; (+) slightly positive effects (0) neutral effect on growth of the rhizobium; (-) slightly negative effects on growth of the rhizobium; (--) negative effects on growth of the rhizobium.

TABLE 3 shows the effects of six endophytic actinobacteria isolated from healthy wheat roots on the symbiosis of RRI128 and lucerne plants after seven weeks from planting. Seeds were coated with six different actinobacteria in 0.3% xanthan gum one day before planting. Inoculation with the RRI128 strain

30

occurred five days after planting. (n = 5 pots, 4 plants per pot).

TABLE 4 shows the effects of endophytic actinobacteria on the symbiosis of rhizobia strain RRI128 and lucerne plants after 45 days from planting. Seeds
5 were coated with six different actinobacteria in 0.3% xanthan gum one day before planting. Inoculation with the RRI128 strain occurred five days after planting. (n = 10 pots, 4 plants per pot).

TABLE 5 shows the effects of endophytic actinobacteria, EN23, LuP30 and
10 LuP47B on symbiosis of the rhizobia strain RRI128 and lucerne in terms of nitrogen content and trace elements in lucerne shoots at 45 days old.

TABLE 6 shows the effects of endophytic actinobacteria on symbiosis of RRI128 and lucerne plants after seven weeks from planting in nutrient limited
15 conditions. Seeds were coated with six different actinobacteria in 0.3% xanthan gum one day before planting. Inoculation of the RRI128 strain occurred five days after planting (n = 5 tubes, 1 plant per tube).

TABLE 7 shows the effect of actinobacterial and rhizobial coinoculation on
20 *Rhizoctonia* root rot of lucerne plants and growth characteristics of shoot and root dry weights.

TABLE 8 shows 16S rRNA gene sequence identities of selected endophytic actinobacteria using BLASTN on GenBank.

25

TABLE 9 shows plant responses due to treatment with *Streptomyces* spp. EN23, LuP30 and LuP47B alone or co-inoculation with *S. meliloti* RRI 128 at 7 weeks after planting (n=4). Control^a: uninoculated plants; Control^b: plants inoculated only with *S. meliloti* RRI 128; (A)= 3 ppm N, (B)= 25 ppm N, (C)= 50
30 ppm N; Asterisks indicate significant differences at $P < 0.05$ (*) or $P < 0.01$ (**).

TABLE 10 shows the effect of actinobacteria and soil N on nodule number per lucerne plant at 4 and 7 weeks after inoculated (n=4). Different letters in the same column indicate means are significantly different ($P < 0.05$).

- 5 TABLE 11 shows the effects of LuP30 and LuP47B on the number of nodules per Lucerne plant after 3 weeks inoculation with different concentrations of *S.meliloti* RRI 128. (n=4) Means $\pm$ SE.

TABLE 12 shows lucerne shoot dry weight in response to co-inoculation with
10 LuP30 or LuP47B and *S. meliloti* RRI 128 after 10, 21 and 35 days at 25 ppm N ($^{15}\text{NH}_4^{15}\text{NO}_3$). Asterisks indicate significant differences at $p < 0.05$.

TABLE 13 shows the accumulation of N (^{14}N and ^{15}N) in the shoot and root of lucerne plants inoculated with rhizobia and actinobacteria (n=4). Different
15 letters in the same column indicate means are significantly different ($P < 0.05$).

TABLE 14 shows the growth and nodulation response of clover to LuP30 and LuP47B after 4 and 7 weeks co-inoculation with *Rhizobium* WSM 1325, (n=4). Different letters in the same column indicate means are significantly different (P
20 < 0.05).

TABLE 15 shows the effects of two actinobacteria LuP30 and LuP47B on the growth of two rhizobial strains on agar plates at three concentrations after 7 days. (++) positive effects on rhizobial growth visible as a zone of increased
25 growth around the actinobacterial plug; (+) slightly positive effects, a smaller zone of increased growth around the actinobacterial plug; (0) neutral effect.

TABLE 16 shows the growth and nodulation of soybean (*Glycine max* cv. Djackal) in response to co-inoculation with each of four strains of actinobacteria
30 and *Bradyrhizobium* strain CB 1809, 4 weeks after inoculation.

TABLE 17 shows the effect of actinobacteria on the elemental content of soy shoots (amount per plant).

TABLE 18 shows the effect of endophytic actinobacteria (*Streptomyces* spp. LuP8, LuP3, LuP44 and LuP47B in combination with *Bradyrhizobium* strain CB1809) and 25 mg NH₄NO₃ per kg soil on soybean growth, nodulation and seeds after 7 weeks inoculation with the rhizobia (n=4). Different letters in the same column indicate means are significantly different ($P < 0.05$). Rhi = *Bradyrhizobium* sp. CB1809.

10

TABLE 19 shows nodule number, nodule weight, pod number and total plant biomass in pea plants grown in field trials at three sites (Riverton SA, Hart SA and Pimpinio Vic).

15 EXAMPLE 1 – MATERIALS AND METHODS

Isolation and identification of endophytic actinobacteria

Four different legumes including lucerne, pea, clover and medics were collected from different places and picked randomly at various stages of growth around South Australia. Different media used for isolation of endophytic actinobacteria were Humic acid Vitamin B agar (HV; Masayuki and Hideo, *Journal of Fermentation Technology* 65(5): 501-509, 1987), yeast extract-casein hydrolysate agar (YECD), tryptic soy agar (TSA) (Oxoid Limited, UK), tap water yeast extract agar (TWYE), all at pH of 7.2± 0.2. Benomyl (DuPont Qualicon, Wilmington, Del. USA) was added to each agar medium at a final concentration of 50 µg.ml⁻¹ to control fungal growth.

The plants were washed under running tap water to remove dust and soil attached to the roots and nodules. The roots and nodules were separated from the plants and air-dried overnight at room temperature. The dry roots and nodules were surface sterilized following the method of Coombs and Franco

30

(*Applied and Environmental Microbiology*, Vol. 69: 5603-5608, 2003). The surface sterilization process started by washing with absolute ethanol for 1 minute, followed by 6 minutes in 4% NaOCl, 30 seconds in absolute ethanol and a final wash with autoclaved R.O. water.

5

Surface sterilized nodules were snipped out from the roots and crushed in 0.9% saline until forming a homogenous mixture. The nodule suspension was spread onto the surface of at least three different isolation media. The sterilized roots were air dried before being cut into approximately 1 cm fragments by a blade or
10 scissors, and placed onto the different media plates. Plates were incubated at 27°C and 37°C.

The plates were checked regularly at least once per week from the first week until new single colonies could not be found. When colonies appeared, they
15 were transferred to half strength potato dextrose agar (HPDA) for purification. The single colonies were transferred onto three different media such as HPDA, oatmeal agar (ISP3) and mannitol soybean agar (MS) to distinguish them based on their different morphologies, colour and pigments produced (media recipes all per Atlas, *Handbook of Microbiological Media*, 1993).

20

Actinobacteria, Sinorhizobium meliloti and lucerne seeds

Lucerne seeds named 'SARDI Ten' and *Sinorhizobium meliloti* RRI128 (referred to as RRI128), which is a commercial inoculant for lucerne, were
25 supplied by the South Australian Research and Development Institute (SARDI). Seeds chosen for planting were similar in size and weight. Five endophytic actinobacteria (EN2, EN16, EN23, EN27, EN46) which were isolated from healthy wheat root and demonstrated to benefit plant growth of some cereals (see Patent Cooperation Treaty publication WO2005/003328, incorporated
30 herein by reference), together with 148 endophytic actinobacteria isolated from different legumes, were tested both *in vitro* and *in planta*.

Effects of endophytic actinobacteria on germination of lucerne – on agar

Lucerne seeds were placed in Petri dishes (usually 2-3 times the required amount) and surface sterilized by the following method: 30 seconds in 70%
5 (v/v) ethanol, 2-3 minutes in 3% (v/v) hypochlorite solution, rinsed three times in autoclaved R.O. water, remaining in the third rinse for 10 minutes then left under the laminar flow to dry. Five sterilized seeds were put on a McKnight's solution 1% agar plate with one drop of an isolated actinobacteria suspension (200 – 2000 cells) applied to each seed while the control seeds received one
10 drop of 0.9% saline. The plates were left under a 14 hour light cycle per day at room temperature (20-30°C) for 2 weeks. The number of germinated seeds and the length of roots were recorded.

Effects of endophytic actinobacteria on germination of lucerne - in sandy loam

15

Lucerne seeds were also sterilized as described above and the sandy loam was autoclaved at 121°C for 15 minutes. Twelve percent moisture sandy loam was made by adding McKnight's solution before adding 300 g of sandy loam with 20 sterilized seeds to a small basket, 10 cm wide and 20 cm length. The
20 seeds were sown with actinobacterial suspension applied on the top before slightly covering with sandy loam. The baskets were kept under a 14 hour light / 10 hour dark cycle at room temperature (20-30°C) for 2 weeks. The number of seeds germinated was recorded, and when germinated, the length of the roots was measured. A total of 148 well-sporulating actinobacteria were tested.

25

IAA production

The ability of the endophytic actinobacteria to produce IAA was examined following the method of Khanma *et al.* (*World Journal of Microbiology and*
30 *Biotechnology* 25: 649-655, 2009). A 6 mm diameter plug of actinobacteria which was grown on ISP2 for 5-7 days was transferred into 5 ml of yeast malt extract (YME) containing 0.2% L-Tryptophan. The broth was shaken at 125 rpm

- 27 -

- for 7 days at 27°C before centrifuging 1 ml of broth at 11,000 rpm for 15 min. The mixture of 0.5 ml supernatant and 1 ml Salkowski reagent (12 g of FeCl₃ per litre of 7.9 M H₂SO₄) was mixed well and kept in the dark for 30 minutes. The IAA production activity was measured using optical density (OD) at 530nm.
- 5 YME broth without L-tryptophan was used as the base line and pure IAA (Sigma) with different concentrations were used to make a standard curve.

Phosphate solubilisation activity

- 10 The phosphate solubilisation ability of selected isolates was detected following the method described by Beneduzi *et al.* (*Applied Soil Ecology* 39: 311-320, 2008). The actinobacteria isolates were inoculated on glucose yeast (GY) medium that contained 10 g of glucose, 2 g of yeast extract and 1.5% agar in 1 L of distilled water. Two solutions were added to the medium, the first was 5 g
- 15 of K₂HPO₄ in 50 ml distilled water and the second solution was 10 g of CaCl₂ in 100 ml of distilled water. These two solutions were autoclaved separately and added into the GY medium before pouring into plates. These two solutions changed the colour of the GY medium to white opaque showing the presence of insoluble calcium phosphate. A positive reaction was demonstrated by the
- 20 presence of a clear zone in the area surrounding the isolates.

Antagonism tests

- Rhizobial strains were grown on yeast mannitol agar (YMA; Graham, *Applied*
- 25 *Microbiology* 17(5): 769-770, 1969) plates or slants for 3-5 days before harvesting. The rhizobial strains were harvested and serially diluted in 0.9% saline. The OD at 600nm of the rhizobial solutions was checked and the number of colony-forming units was counted following the method of Miles and Misra (*Journal Hygiene* 38: 732-749, 1938). At the same time 100 µl of these
- 30 serial dilutions at different OD values were spread onto YMA plates and allowed to dry. Two plugs about 25 mm² of each actinobacterial strain grown on International Streptomyces Project 2 (ISP2; Atlas, *Handbook of Microbiological*

Media, 1993) medium for 7 days were placed on the surface of the inoculated YMA plates. The plates were incubated for 5-7 days at 27°C and checked daily for antagonistic activity against the rhizobia. Rhizobia and the actinobacteria that were grown separately as pure cultures on YMA plates were used as negative controls. Streptomycin, vancomycin and were used as positive antibacterial controls. All treatments were replicated twice and incubated at 27°C.

Interaction between endophytic actinobacteria and rhizobium on lucerne

10

Lucerne seeds were surface-sterilized as described above and sown in autoclaved pots. Each pot contained about 1 kg of sand and vermiculite mixture, and had two separate parts to allow easy drainage. Five and 148 endophytic actinobacteria isolated from surface-sterilized healthy wheat roots and surface-sterilized healthy root and nodules of four different legumes such as lucerne, pea, clover and medics, respectively, were coated on the surface of the lucerne seeds in a 0.3% (w/v) sterile xanthan gum solution. 100 ml of MQ water was added to each pot before planting the ten coated seeds. The top of the pot was covered with a thin layer of granulated plastic beads. Then, 200 mL of 1/80 McKnight's solution containing starter nitrogen (266 mg NH_4NO_3 per 20 L McKnight's solution) was gently added to each pot before covering with freezer bags and placing in the glasshouse. The number of seedlings was thinned down to four plants before adding 1 ml of rhizobium RRI128 around 10^8 CFU/ml at five days from planting. Every week 50 ml of nitrogen solution (1.2 g.L^{-1} of NH_4NO_3) was applied to each pot for nitrogen-treated plants and MQ water was added as required.

All treatments were replicated ten times completely randomized in the glasshouse, with the position of the pots changed randomly every week. The plants were harvested after the sixth to seventh week from sowing. The parameters examined were height and dry weight of the shoot, length and dry weight of the root, the number and dry weight of nodules per plant. Nodules

30

were removed, counted and dried at 60°C. Dry weight of each nodule was calculated by dividing total nodule weight by total nodule number of two plants with five replicates.

5 *SPAD 502 readings, total nitrogen analysis and N₂ fixed in the shoots*

During the sixth week, leaves of lucerne plants were measured by a SPAD 502 meter (Chlorophyll meter SPAD-502, Konica Minolta) designed to indicate the amount of chlorophyll present in plant leaves. The three biggest leaves were
 10 checked to get SPAD readings. Moreover, dry leaves of control plants (only treated with *Sinorhizobium*), and plants treated with rhizobium and EN23 and EN27, LuP30 and LuP47B harvested in the seventh week were analysed for the content of nitrogen and other elements. The leaves were dried at 60°C for 48 h to obtain constant weight and were ground to about 1 mm in size for
 15 analysis.

N₂ fixation associated with the shoots was calculated by the following equation:

$$\text{N}_2 \text{ fixed in the shoots} = (\%N \cdot \text{SDW})_{\text{treatment}} - (\%N \cdot \text{SDW})_{\text{uninoculated}}$$

20 *Nutrient limitation experiment*

Lucerne seeds were surface sterilized and coated with actinobacteria as described above. Two coated seeds were sown in a 50 ml centrifuge tube containing 65 g of autoclaved washed sand and 10 ml of McKnight's starter N
 25 (0.133 mg) added five times less than compared with normal 0.665 mg per plant in pot assays. This was then covered by sand and plastic beads. The number of seedlings was thinned down to one seedling before inoculating with 10⁸ CFU.ml⁻¹ RRI128 suspension. The tubes were kept inside the growth chamber with a 12 h light and 12 h dark cycle. Water was supplied as required
 30 until 7 weeks.

DNA extraction

In a sterile 1.5 ml eppendorf tube 10 µl of lysozyme was added to 500 µl of Tris-EDTA (TE) pH 7.4, before re-suspending 2-3 loops of actinobacterial cells into the mixture, which was then vortexed and spun later. The eppendorf tube was incubated at 37°C for 60 minutes before adding 10 µl of proteinase K and 32.5 µl of 10% SDS and incubated at 55°C for 60 minutes. Next, 100 µl of 5M NaCl and 65 µl of CTAB/NaCl were added and the mixture was incubated at 55°C for 10 minutes. Six hundred microlitres of phenolchloroform was added and the tube was left at room temperature for 30 minutes with intermittent shaking every 10 minutes. After centrifuging at 12,000 rpm for 15 minutes, the supernatant was transferred to a new sterile 1.5 ml eppendorf tube. An additional 500 µl of chloroform was added to the tube and left at room temperature for 15 minutes with mixing by inversion every 7-8 minutes before centrifuging at 12,000 rpm for 15 minutes. The aqueous phase was transferred into a new sterile 1.5 ml eppendorf tube before adding 20 µl of 10 mg.ml⁻¹ RNase and incubating at 37°C for 60 minutes. Then 500 µl of chloroform was added and the tube left at room temperature for 15 minutes. After centrifuging at 12,000 rpm for 15 minutes, the supernatant was transferred to a new sterile 1.5 ml eppendorf tube. An additional 500 µl of chloroform was added and the tube left at room temperature for 15 minutes (with mixing by inversion every 7-8 minutes). After centrifuging at 12,000 rpm for 15 minutes, the supernatant was transferred to a new sterile 1.5 ml eppendorf tube (steps repeated). A 0.1x volume of 3M Na Acetate (50 µl) and 3x volume of 100% ethanol (1 ml) was added to the tube before leaving at -20°C overnight. The supernatant was removed carefully so as not to disrupt the pellet after centrifuging at the maximum speed 16,000 rpm for 15 minutes. The pellet was washed twice with 70% ethanol and dried by placing tubes in a heating block at 55°C with the lids open for approximately 10 minutes. Finally, the pellet was re-suspended in 50 µl of sterile water.

PCR of 16S rRNA gene

A master mixture was prepared as 1 µl of dNTPs (10 mM), 1 µl of DNA Taq polymerase (5U/µl), 5 µl of ThermoPol buffer, 2 µl of 27f primer (AGAGTTTGGATCCTGGCTCAG; SEQ ID NO: 1), 2 µl of 1465r primer (TACGGYTACCTTGTTACGACTT; SEQ ID NO: 2), 2 µl of DNA sample and 37 µl of injection water. PCR was performed by heating the PCR tubes at 94°C for 2 minutes, followed by 40 cycles of 94°C for one minute, 52°C for one minute and 72°C for two minutes, and 72°C for 10 minutes. A 1.2% agarose gel containing 3 µl of GelRed (Biotium) in 40 ml agarose was used to separate the molecular weight of the PCR products. One microlitre of loading dye was mixed well with 2 µl of each PCR product before loading the gel, which was run in a running buffer 0.5x TBE at 70 V and 400 mA for 60 minutes. The products of PCR were sequenced by Macrogen, Korea. The resultant 16S rRNA sequences were compared to the GenBank database by using the National Centre for Biotechnology Information database (NCBI) BLASTN program, including the results of the highest matches for each isolate and the corresponding bit score and percentage identity.

EXAMPLE 2 - ISOLATION AND IDENTIFICATION OF ENDOPHYTIC ACTINOBACTERIA

225 endophytic actinobacteria were isolated from roots and nodules of pea, lucerne, clover and medic. 73 were from nodules and 152 were from roots. Based on their morphology, 126 cultures (56%) belong to the genus *Streptomyces*, 54 (24%) belong to *Microbispora*, 20 (8.89%) belong to *Micromonospora* and 25 cultures are as yet unidentified. Humid acid vitamin B agar (HV), yeast extract casein dextrose (YECD) and tap water yeast extract (TWYE) media successfully allowed growth of almost all of the isolates mentioned. 125 cultures were isolated from HV medium, 72 cultures were from TWYE, 26 were from YECD and one of them was isolated from TSA. There was not much difference in the number of cultures isolated at 37°C and 27°C,

which were 125 and 112 cultures, respectively. Eighty five cultures were isolated from roots of lucerne, while 65 isolates were from pea (42 from roots and 23 from nodules), 37 cultures were from clover (16 from roots and 26 from nodules) and 35 were from medics (12 from roots and 23 from nodules). Thirty
 5 two out of the 73 isolates from nodules were *Streptomyces*, 17 were *Microbispora*, 5 were *Micromonospora* and 19 were unidentified.

10 EXAMPLE 3 - EFFECTS OF ENDOPHYTIC ACTINOBACTERIA ON GERMINATION OF LUCERNE, IAA PRODUCTION AND PHOSPHATE SOLUBILISATION

Fifty six of 148 cultures (38%) isolated from lucerne promoted germination of lucerne seeds and 27 (18%) isolates negatively affected germination of lucerne seeds in terms of number of seeds germinated and length of roots on agar
 15 plates. In addition, 39 of 148 cultures improved germination of lucerne seeds with the presence of rhizobium on sandy loam.

As shown in Table 1 and Figure 3, a range of endophytic actinobacteria showed the ability to produce IAA. Table 1 also identifies cultures that were found to
 20 have phosphate solubilisation ability. They were LuP5B, LuP44 and LuP8A.

EXAMPLE 4 – ANTAGONISM TESTING

Two concentrations of rhizobia were tested for antagonism with 14 endophytic
 25 actinobacteria; five non-legume-isolated and nine legume-isolated cultures. As shown in Table 2, most of the endophytic actinobacteria had neutral or positive effects on growth of three rhizobia RRI128, SRDI736, WSM1115G, except for LuP10, EN28 and EN46. LuP10 increased growth of the RRI128 but inhibited growth of SRDI736 and WSM1115G, whereas LuP3, LuP30 and LuP47B
 30 increased growth of the three rhizobia at the various concentrations.

As shown in Figure 4, LuP30 and LuP47B showed significant rhizobial growth

stimulation *Rhizobium leguminosarum* bv. *trifolii* strain WSM 1325 and *Bradyrhizobium lupini* strain WSM 471.

EXAMPLE 5 - EFFECTS OF SIX ENDOPHYTIC ACTINOBACTERIA 5 ISOLATED FROM NON-LEGUMES

As shown in Table 3, some endophytic actinobacteria isolated from wheat roots showed beneficial interactions with the RRI128-inoculated lucerne plants while some were neutral, with no significant impacts on different parameters. EN2
10 significantly increased the fresh and dry weight of the shoot as well as the length of the root, while EN23 increased not only height, fresh and dry weight of the shoot but also length and fresh weight of the root. In particular, the average height of the shoot plants receiving combined treatment of RRI128 and EN23 was 15.2 cm, whilst plants treated with RRI128 only was 12.55 cm. EN23
15 increased the shoot height of the plant by 21.1%. Moreover, EN23 significantly increased the dry weight of each nodule, the total dry weight of nodules, nitrogen content of the plant as well as total nitrogen per plant, though it did not significantly increase the number of nodules per plant. Total mass per plant treated with EN23 increased by 25.7% compared with plants treated with the
20 rhizobium only control.

Treatment with EN27 resulted in slight increases in height, fresh and dry weight of the shoot, and fresh and dry weight of the root. Although EN27 significantly reduced the number of nodules, the fresh weight of each nodule and the total
25 dry weight of nodules per plant were higher than that of RRI128 only plants. Furthermore, EN27 also significantly increased the SPAD readings, nitrogen content (%N₂) and total nitrogen per plant. EN23 and EN27 increased the amount of N₂ fixed in the shoots, by 0.85 and 0.80 mg per plant, respectively, compared with the RRI128 only plants (Table 3). In contrast, EN16 significantly
30 reduced the number of nodules and total dry weight of nodules per plant after seven weeks from planting. EN28 and EN46 had no significant effect on the growth of lucerne plants with the RRI128. Nitrogen content was 2.725% of

mass for the control, 3.225% of mass for EN23-treated and 3.65% for EN27-treated plants. There was 1.38 mg of total N per control plant, while EN23 and EN27-treated plants had 2.23 mg and 2.18 mg total N, respectively.

5 EXAMPLE 6 – EFFECTS OF ENDOPHYTIC ACTINOBACTERIA ISOLATED FROM LEGUMES

The interactions between rhizobium RRI128 and 148 cultures isolated from legumes were screened in terms of plant growth and nitrogen fixation. LuP47B
 10 and LuP30 showed beneficial effects on the symbiosis of rhizobium and lucerne, with increased height of shoot, mass of shoot and plant and nitrogen fixation per plant. As shown in Table 4, treatment with these cultures led to an increase of 35.33% and 24.87% of shoot dry weight and 29.91% and 25.87% of total mass per plant, respectively. LuP47B also increased the height of the
 15 shoot significantly, up to 26.25%. Although LuP30 did not significantly promote the height of the shoot, it developed a longer root compared with plants treated with RRI128 only. In contrast, LuP10 increased the root biomass instead of root length.

20 As shown in Table 5, the combination of rhizobium RRI128 and EN23 significantly increased copper, phosphorous, sodium and nitrogen content in the shoot compared with the RRI128 alone. LuP30 and LuP47B treatment resulted in a significant increase all of the trace elements tested, except iron with the presence of the RRI128. Although the shoot dry weight of plants
 25 inoculated with the RRI128 and LuP30 was less than that of plants treated with RRI128 and LuP47B they showed higher amounts of copper, iron and zinc compared with LuP47B. EN23, LuP30 and LuP47B also increased nitrogen content in the shoot, with 0.35, 0.61 and 0.83 mg per each shoot, respectively (see Figure 6).

EXAMPLE 7 - EFFECTS OF ENDOPHYTIC ACTINOBACTERIA ON SYMBIOSIS OF RHIZOBIA AND LUCERNE WITH LIMITATION OF NUTRITIENT SUPPLY

5 As shown in Table 6 and Figure 7, in limited nutrient conditions LuP5B significantly increased all growth parameters, such as height of shoot, length of root, shoot and root dry weight, except for number of nodules. LuP47B treatment resulted in increased length of root and shoot dry weight but LuP30 did not increase other parameters of plant growth. Although LuP12A increased
10 root dry weight, it did not promote root length. In contrast, LuP3 increased the length of root but it did not increase the root dry weight (Table 6).

EXAMPLE 8 – BIOCONTROL ACTIVITY IN PLANTA

15 Fifty milliliter centrifugal tubes were used to screen for the biocontrol ability of actinobacterial strains against the fungal root pathogen *Rhizoctonia*. Forty five grams of autoclaved sandy loam were used at a 12% moisture content added as McKnight's starter nitrogen (266mg of NH_4NO_3) solution. Two millet seeds infected with *R. solani* AG8 strain W19 were added at the top of the sandy
20 loam, and a further 10 g of soil containing 12% moisture was added to cover the millet seeds. Two tubes without adding the infested millet seeds with the pathogen and without endophytic actinobacteria were as used as controls. The tubes were placed in a rack covered with aluminium foil and placed in a chamber for two weeks at 15°C in the absence of light.

25

Lucerne seeds were surface-sterilized and pre-germinated with actinobacterial suspensions on autoclaved moist filter paper in petri dishes. When the roots were about 1-3mm length they were dipped in 5ml (to cover all the seeds) of the rhizobial suspension (approximately 10^8 cfu/ml) for 3 minutes. Two pre-
30 germinated and coated seeds were transferred into each 50 ml tube and covered with 5g of soil (12% moisture) and a layer of plastic beads. The tubes were kept at 15°C in a growth chamber for 3 weeks. There were two replicates

of each treatment, and MQ water was added as required. The number of seedlings that emerged, the length of root and root damage were recorded.

Results of the biocontrol assay are shown in Table 7.

5

EXAMPLE 9 – IDENTIFICATION OF ACTINOBACTERIA USING 16S RRNA GENE SEQUENCING

10 LuP3, LuP12A, LuP30, LuP47B, EN23, EN27, LuP8 and LuP44 were putatively identified as *Streptomyces sp.* by 16S rRNA gene sequencing.

The determined 16S rRNA gene sequences for each organism were as follows:

Isolate	16S rRNA gene sequence (5'-3')	Sequence Identifier
LuP3	GTGGATTAGTGGCGAACGGGTGAGTAACACGTGGGCAATCTGCC CTTCACTCTGGGACAAGCCCTGGAAACGGGGTCTAATACCGGAT AATACTTTCTCCCTCCTGGGAGAAGGTTGAAAGCTCCGGCGGTG AAGGATGAGCCCGCGGCCTATCAGCTAGTTGGTGGGGTAATGGC CTACCAAGGCGACGACGGGTAGCCGGCCTGAGAGGGCGACCGGC CACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGC AGTGGGGAATATTGCACAATGGGCGAAAGCCTGATGCAGCGACG CCGCGTGAGGGATGACGGCCTTCGGGTTGTAAACCTCTTTCAGC AGGGAAGAAGCGAAAGTGACGGTACCTGCAGAAGAAGCGCCGGC TAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGCGCAAGCGT TGTCCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTC CGTCGGTTGTGAAAGCCCGGGGCTTAACCCCGGGTCTGCAGTCG ATACGGGCAGGCTAGAGTGTGGTAGGGGAGATCGGAATTCCTGG TGTAGCGGTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCG AAGGCGGATCTCTGGGCCATTACTGACGCTGAGGAGCGAAAGCG TGGGGAGCGAACAGGATTAGATACCCTGGTAGTCCACGCCGTAA ACGTTGGGAACATAAGGTGTTGGCGACATTCCACGTCGTCGGTGC CGCAGCTAACGCATTAAGTTCCCGCCTGGGGGAGTACGGCCGCA AGGCTAAACTCAAAGGAATTGACGGGGGCCCCGCACAAGCAGCGG AGCATGTGGCTTAATTCGACGCAACGCGAAGAACCTTACCAAGG CTTGACATACACCGGAAAGCATCAGAGATGGTGCCCCCCTTGTG	SEQ ID NO: 3

	GTCGGTGTACAGGTGGTGCATGGCTGTCGTCAGCTCGTGTCTGTG AGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTTGTTCTGT GTTGCCAGCATGCCCTTCGGGGTGATGGGGACTCACAGGAGACT GCCGGGGTCAACTCGGAGGAAGGTGGGGACGACGTCAAGTCATC ATGCCCCCTTATGTCTTGGGCTGCACACGTGCTACAATGGCCGGT ACAATGAGCTGCGATGCCGCGAGGCGGAGCGAATCTCAAAAAGC CGGTCTCAGTTCGGATTGGGGTCTGCAACTCGACCCCATGAAGT CGGAGTTGCTAGTAATCGCAGATCAGCATTGCTGCGGTGAATAC GTTCCCGGGCCTTGTACACACCGCCCGTCACGTCACGAAAGTCG GTAACACCCGAAGCCG	
LuP12A	GATGAACCACTTCGGTGGGGATTAGTGGCGAACGGGTGAGTAAC ACGTGGGCAATCTGCCCTTCACTCTGGGACAAGCCCTGGAAACG GGGTCTAATACCGGATACCACTACCGCAGGCATCTGTGGTGGTT GAAAGCTCCGGCGGTGAAGGATGAGCCCGCGGCCTATCAAGGTT GTTGGTGAGGTAATGGCTCACCAAGGCGACGACGGGTAGCCGGC CTGAGAGGGCGACCGGCCACACTGGGACTGAGACACGGCCCAGA CTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGAA AGCCTGATGCAGCGACGCCGCGTGAGGGATGACGGCCTTCGGGT TGTAACCTCTTTCAGCAGGGAAGAAGCGAAAGTGACGGTACCT GCAGAAGAAGCGCCGGCTAACTACGTGCCAGCAGCCGCGGTAAT ACGTAGGGCGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGC TCGTAGGCGGCTTGTACGTCGGGTGTGAAAGCCCGGGGCTTAA CCCCGGGTCTGCATTCGATACGGGCTAGCTAGAGTGTGGTAGGG GAGATCGGAATTCCTGGTGTAGCGGTGAAATGCGCAGATATCAG GAGGAACACCGGTGGCGAAGGCGGATCTCTGGGCCATTACTGAC GCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCT GGTAGTCCACGCCGTAAACGGTGGGAACTAGGTGTTGGCGACAT TCCACGTCGTCGGTGCCGCGAGCTAACGCATTAAGTTCCCCGCCT GGGGAGTACGGCCGCAAGGCTAAACTCAAAGGAATTGACGGGG GCCCCGACAAGCAGCGGAGCATGTGGCTTAATTCGACGCAACGC GAAGAACCTTACCAAGGCTTGACATACGCCGGAAGCATCAGAG ACGGTGCCCCCCTTGTGGTTCGGTGTACAGGTGGTGCATGGCTGT CGTCAGCTCGTGTCTGAGATGTTGGGTAAAGTCCCGCAACGAG CGCAACCCTTGTCTGTGTTGCCAGCATGCCCTTCGGGGTGATG GGGACTCACAGGAGACCGCCGGGGTCAACTCGGAGGAAGGTGGG GACGACGTCAAGTCATCATGCCCCCTTATGTCTTGGGCTGCACAC GTGCTACAATGGCAGGTACAATGAGCTGCGATACCGTGAGGTGG AGCGAATCTCAAAAAGCCTGTCTCAGTTCGGATTGGGGTCTGCA	SEQ ID NO: 4

	ACTCGACCCCATGAAGTCGGAGTTGCTAGTAATCGCAGATCAGC ATTGCTGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCG TCACGTCACGAAAGTCGGTAACACCCGAAGCCGGTGGCCTCAAC CC	
LuP30	CAGTCGAACGATGAACACTTCGGTGGGGATTAGTGGCGAACGGG TGAGTAACACGTGGGCAATCTGCCCTTCACTCTGGGACAAGCCC TGGAAACGGGGTCTAATACCGGATAACACTTCCACTCGCATGGG TGGAGGTTAAAAGCTCCGGCGGTGAAGGATGAGCCCGCGGCCTA TCAGCTTGTTGGTGAGGTAATGGCTCACCAAGGCGACGACGGGT AGCCGGCCTGAGAGGGCGACCGGCCACACTGGGACTGAGACACG GCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGGGATGACGGCC TTCGGGTTGTAAACCTCTTTCAGCAGGGAAGAAGCGAAAGTGAC GGTACCTGCAGAAGAAGCGCCGGCTAACTACGTGCCAGCAGCCG CGGTAATACGTAGGGCGCAAGCGTTGTCCGGAATTATTGGGCGT AAAGAGCTCGTAGGCGGTCTGTGCGTCGGATGTGAAAGCCCGG GGCTTAACCCCGGGTCTGCATTTCGATACGGGCAGACTAGAGTGT GGTAGGGGAGATCGGAATTCCTGGTGTAGCGGTGAAATGCGCAG ATATCAGGAGGAACACCGGTGGCGAAGGCGGATCTCTGGGCCAT TACTGACGCTGAGGAGCGAAAGCGTGGGGAGCGAACAGGATTAG ATACCCTGGTAGTCCACGCCGTAAACGGTGGGAACTAGGTGTTG GCGACATTCCACGTCGTCGGTGCCGCAGCTAACGCATTAAGTTC CCCGCCTGGGGAGTACGGCCGCAAGGCTAAACTCAAAGGAATT GACGGGGGCCCCGCACAAGCAGCGGAGCATGTGGCTTAATTCGAC GCAACGCGAAGAACCTTACCAAGGCTTGACATACACCGGAAACG GCCAGAGATGGTCGCCCCCTTGTGGTTCGGTGTACAGGTGGTGCA TGGCTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGTAAAGTCCCG CAACGAGCGCAACCCTTGTTCTGTGTTGCCAGCATGCCCTTCGG GGTGATGGGGACTCACAGGAGACTGCCGGGGTCAACTCGGAGGA AGGTGGGGACGACGTCAAGTCATCATGCCCCCTTATGTCTTGGGC TGCACACGTGCTACAATGGCCGGTACAAAGAGCTGCGAAGCCGT GAGGTGGAGCGAATCTCAAAAAGCCGGTCTCAGTTCGGATTGGG GTCTGCAACTCGACCCCATGAAGTCGGAGTTGCTAGTAATCGCA GATCAGCATTGCTGCGGTGAATACGTTCCCGGGCCTTGTACACA CCGCCCGTCACGTCACGAAAGTCGGTAACACCCGAAGCCGGTGG CCCAACC	SEQ ID NO: 5

LuP47B	GTGAGGTAATGGCTCACCAAGGCGACGACGGGTAGCCGGCCTGA GAGGGCGACCGGCCACACTGGGACTGAGACACGGCCCAGACTCC TACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGAAAGCC TGATGCAGCGACGCCGCGTGAGGGATGACGGCCTTCGGGTGTA AACCTCTTTCAGCAGGGAAGAAGCGAAAGTGACGGTACCTGCAG AAGAAGCGCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGT AGGGCGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGT AGGCGGCTTGTCACGTCGGGTGTGAAAGCCCGGGGCTTAACCCC GGGTCTGCATTTCGATACGGGCTAGCTAGAGTGTGGTAAGGGAGA TCGGAATTCCTGGTGTAGCGGTGAAATGCGCAGATATCAGGAGG AACACCGGTGGCGAAGGCGGATCTCTGGGCCATTACTGACGCTG AGGAGCGAAAGCGTGGGGAGCGAACAGGATTAGATACCCTGGTA GTCCACGCCGTAAACGGTGGGAACTAAGGTGTTGGCGACATTCC ACGTCGTCGGTGCCGCAGCTAACGCATTAAGTTCCCGCCCGGGG GAGTACGGCCGCAAGGCTAAAACTCAAAGGAATTGACGGGGGCC CGCACAAGCAGCGGAGCATG	SEQ ID NO: 6
EN23	ACGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGATGA AGCCGCTTCGGTGGTGGATTAGTGGCGAACGGGTGAGTAACACG TGGGCAATCTGCCCTTCACTCTGGGACAAGCCCTGGAAACGGGG TCTAATACCGGATAACACTCTGTCCCGCATGGGACGGGGTTGAA AGCTCCGGCGGTGAAGGATGAGCCCGCGGCCTATCAGCTTGTTG GTGGGGTAATGGCCTACCAAGGCGACGACGGGTAGCCGGCCTGA GAGGGCGACCGGCCACACTGGGACTGAGACACGGCCCAGACTCC TACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGAAAGCC TGATGCAGCGACGCCGCGTGAGGGATGACGGCCTTCGGGTGTA AACCTCTTTCAGCAGGGAAGAAGCGAAAGTGACGGTACCTGCAG AAGAAGCGCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGT AGGGCGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGT AGGCGGCTTGTCACGTCGGATGTGAAAGCCCGGGGCTTAACCCC GGGTCTGCATTTCGATACGGGCTAgCTAGAGTGTGGTAGGGGAGA TCGGAATTCCTGGTGTAgCGGTGAAATGCGCAGATATCAGGAGG AACACCGGTGGCGAAGGCGGATCTCTGGGCCATTACTGACgTcT GAGGAGCGAAAGCGTGGGgAgCGAACAGGATTAGATACCCTGgT AGTCCACGCCGTAAACGTTGGgAACTAGgTGTTGGCGACATTCC ACGTCGTCGGTGCCGCAGCTAACGCATTAAGTTCCCGCCTGGG GAGTACGGCCGCAAGGCTAAAACTCAAAGGAATTGACGGGGGCC CGCACAAGCAGCGGAGCATGTGGCTTAATTCGACGCAACGCGAA GAACCTTACCAAGGCTTGACATATAACGGAAAGCATCAGAGATG	SEQ ID NO: 7

	GTGCCCCCCTTGTGGTCGGTATACAGGTGGTGCATGGCTGTCTGT CAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGC AACCCTTGTTCTGTGTTGCCAGCATGCCCTTCGGGGTGATGGGG ACTCACAGGAGACTGCCGGGGTCAACTCGGAGGAAGGTGGGGAC GACGTCAAGTCATCATGCCCCCTTATGTCTTGGGCTGCACACGTG CTACAATGGCCGGTACAATGAGCTGCGATGCCGCGAGGCGGAGC GAATCTCAAAAAGCCGGTCTCAGTTCGGATTGGGGTCTGCAACT CGACCCCATGAAGTCGGAGTTGCTAGTAATCGCAGATCAGCATT GCTGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCGTCA CGTCACGAAAGTCGGTAACACCCGAAGCCGGTGGCCCAACCCTT GTGGGAGGGAGCTGTCTGAAGGTGGGACTGGCGATTG	
EN27	TTAANACATGCAANTCGAACGATGAACCCNGTTTCGGTGGTGGA TTAGTGGCGAACGGTGAGTAANANGTGGGCAATTTGCCCTTCAT TTTGGACAAGCCCTGGAAACGGGTTTAATACCGGATAACATTTT NTCCCGCATGGGANGGGGTGAAAGNTCCGGCGGTGAAGGATGA GCCCCGCGCCTATNAGCTTGTTGGTGGGGTAATGGCCTACCCAA GGGAGACGGGTAGCCGGCCTGAGAGGGCGACCGGCCACACTGGG AATGAGANACGGCCCAGAATCCTACGGGAGGCAGCAGTGGGGAA TATTGCACAATGGGCGAAAGCCTGATGCAGCGANGCCGCGTGAG GGATGACGGCCTTNGGGTTGTAAACCTTTTTNAGCAGGGAAGAA GCGAAAGTGACGGTACCTGCAGAAGAAGCGCCGGCTAAATAAGT GCCAGCAGCCGCGGTAATAAGTAGGGCGCAAGCGTTGTCCGGAA TTATTGGGCGTAAAGAGCTTGTAGGCGGCTTGTCANGTNGGATG TGAAAGCCCCGGGGNTTAACCCCGGGTTTGCATTTGATACGGGCT AGNTAGAGTGTGGTAGGGGAGATNGGAATTCCTGGTGTAGCGGT GAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGGAT CTCTGGGCCATTACTGACGCTGAGGAGCGAAAGCGTGGGGAGCG AACAGGATTAGATAACCCTGGTAGTCCACGCCGTAAACGTTGGGA ACTAGGTGTTGGCGACATTCCACGTCGTCGGTGCCGCAGCTAAC GCATTAAGTTCCCCGNCTGGGGAGTACGGCCGCAAGGCTAANAC TCAAAGGAATTGACGGGGGCCCCGNACAAGCAGCGGANCATGTGG CTTAATTCGACGCANCGCGAAGAACCCTTACCAAGGCTTGACATA TACCGGAAAGCATCAGAGATGGTGCCCCCCTTGTGGTCGNTATA CANGTGGTGCATGNCTGTCTGTACCTCGTGTCTGTGAGATGTTGG GTTAAGTCCCGCAACGAGCGCNACCCTTGNTCTGTGTTGNCANC ATGCCCTTCGGGGNTGATGGGGACTCACAGGANACTGNCCGGGG TCAACTCCGGANGAAGGTGGGTGACGAAGTCAAGGTCATCATGN CCCCTTATGTCTTGGTGCTGCACACGTGC	SEQ ID NO: 8

LuP8	AATGGGCTAAGTTCGAAACGATTGAACCACTTTCGGTGGGGATT AGTGGCGAACGGGTGAGTAACACGTGGGCAATCTGCCCTTCACT CTGGGACAAGCCCTGGAAACGGGGTCTAATACCGGATAACCACTA CCGCAGGCATCTGTGGTGGTTGAAAGCTCCGGCGGTGAAGGATG AGCCCGCGGCCTATCAGCTTGTTGGTGAGGTAATGGCTCACCAA GGCGACGACGGGTAGCCGGCCTGAGAGGGCGACCGGCCACACTG GGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGG AATATTGCACAATGGGCGAAAGCCTGATGCAGCGACGCCGCGTG AGGGATGACGGCCTTCGGGTGTGTAACCTCTTTCAGCAGGGAAG AAGCGAAAGTGACGGTACCTGCAGAAGAAGCGCCGGCTAACTAC GTGCCAGCAGCCGCGGTAATACGTAGGGCGCAAGCGTTGTCCGG AATTATTGGGCGTAAAGAGCTCGTAGGCGGCTTGTCACGTCGGG TGTGAAAGCCCGGGGCTTAACCCCGGGTCTGCATTGATACGGG CTAGCTAGAGTGTGGTAGGGGAGATCGGAATTCCTGGTGTAGCG GTGAAATGCGCAGATATCAGGAGGAACACCGGTGGCGAAGGCGG ATCTCTGGGCCATTACTGACGCTGAGGAGCGAAAGCGGGGGGGG CAAAAAGGGAACCCGGCCGGGGGGGG	SEQ ID NO: 9
LuP44	TCGGTGGGGATTAGTGGCGAACGGGTGAGTAACACGTGGGCAAT CTGCCCTTCACTCTGGGACAAGCCCTGGAAACGGGGTCTAATAC CGGATACCACTACCGCAGGCATCTGTGGTGGTTGAAAGCTCCGG CGGTGAAGGATGAGCCCGCGGCCTATCAGCTTGTTGGTGAGGTA ATGGCTCACCAAGGCGACGACGGGTAGCCGGCCTGAGAGGGCGA CCGGCCACACTGGGACTGAGACACGGCCCAGACTCCTACGGGAG GCAGCAGTGGGGAATATTGCACAATGGGCGAAAGCCTGATGCAG CGACGCCGCGTGAGGGATGACGGCCTTCGGGTGTGTAACCTCTT TCAGCAGGGAAGAAGCGAAAGTGACGGTACCTGCAGAAGAAGCG CCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGCGCA AGCGTTGTCCGGAATTATTGGGCGTAAAGAGCTCGTAGGCGGCT TGTACGTCGGGTGTGAAAGCCCGGGGCTTAACCCCGGGTCTGC ATTCGATACGGGCTAGCTAGAGTGTGGTAGGGGAGATCGGAATT CCTGGTGTAGCGGTGAAATGCGCAGATATCAGGAGGAACACCGG TGGCGAAGGCGGATCTCTGGGCCATTACTGACGCTGAGGAGCGA AAGCGTGGGGAGCGAACAGGATTAGATACCCTGGTAGTCCACGC CGTAAACGGTGGGAACTAGGTGTTGGCGACATTCCACGTCGTCG GTGCCGCAGCTAACGCATTAAGTTCCCCGCCTGGGGAGTACGGC CGCAAGGCTAAAACTCAAAGGAATTGACGGGGGCCCCGCACAAGC AGCGGAGCATGTGGCTTAATTCGACGCAACGCGAAGAACCTTAC CAAGGCTTGACATACGCCGGAAGCATCGGAGACGGGGTCCCCC	SEQ ID NO: 10

	TTGTGGTCGGTGTACAGGTGGTGCATGGCTGTCGTCAGCTCGTG TCGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCTTGT CCTGTGTTGCCAGCATGCCCTTCGGGGTGATGGGGACTCACAGG AGACCGCCGGGGTCAACTCGGAGGAAGGTGGGGACGACGTCAAG TCATCATGCCCCCTTATGTCTTGGGCTGCACACGTGCTACAATGG CAGGTACAATGAGCTGCGATAACCGTGAGGTGGAGCGAATCTCAA AAAGCCTGTCTCAGTTCGGATTGGGGTCTGCAACTCGACCCCAT GAAGTCGGAGTTGCTAGTAATCGCAGATCAGCATTGCTGCGGTG AATACGTTCCCGGGCCTTGTACACACCGCCCGTCACGTCACGAA AGTCGGTAACACCCGAAGCCGGTGGCCCAACCC	
--	---	--

As shown in Table 8, the closest match for LuP3 was *Streptomyces drozdowiczii* with up to 99.4% 16S rRNA gene sequence identity. The closest match for LuP30 was *Streptomyces rishiriensis*, which showed up to 99.9% 16S rRNA gene identity. LuP12A, LuP47B, LuP8 and LuP44 are all very close to both *Streptomyces ciscaucasicus* and *Streptomyces canus*, with >99% 16S rRNA gene sequence identity.

EXAMPLE 10 – EFFECT OF ACTINOBACTERIA/RHIZOBIA CO-INOCULATION ON GROWTH PARAMETERS OF LUCERNE AT DIFFERING N CONCENTRATIONS

The effects of three actinobacteria on the growth and symbiosis of lucerne and rhizobia was studied at three levels of NH₄NO₃ 3 ppm, 25 ppm and 50 ppm.

The factorial experiment comprised (i) 3 strains of actinobacteria, (ii) ± inoculation with *Sinorhizobium meliloti* RRI 128 and (iii) 3 levels of soil NH₄NO₃. The pots were prepared and watered with a nitrogen deficient McKnight's solution supplemented with NH₄NO₃ to provide soil nitrogen of 3, 25 and 50 ppm. Plant seeds designated to rhizobia treatments were inoculated with a suspension (1ml per plant containing 10⁸ CFU) of rhizobia 6 days after sowing. Each treatment was replicated eight times. Pots were arranged in a completely randomised design in a greenhouse and plants four pots of each treatment were harvested at 4 and 7 weeks after inoculation with the *S. meliloti* RRI 128.

As shown in Table 9, co-inoculation of each of EN23, LuP30 and LuP47B with *S. meliloti* RRI 128 was able to statistically significantly improve at least one of root dry weight or shoot dry weight over the un-inoculated or Rhizobium only controls at at least one concentration of N.

As shown in Table 10, co-inoculation of each of EN23, LuP30 and LuP47B with *S. meliloti* RRI 128 was able to statistically significantly improve the number of nodules over the un-inoculated or Rhizobium only controls at at least one concentration of N at the 4 week and/or 7 week sampling time.

Regarding nutrient levels, in the absence of the *S. meliloti* RRI 128 EN23, LuP30 or LuP47B reduced significantly iron and copper in shoot plants after 7 weeks at both 25 mg and 50 mg/kg_{nitrogen} NH₄NO₃ supply while sodium and molybdenum was increased. Total nitrogen in shoot plants were increased significantly with seeds coated with LuP47B at 50 mg N while EN23 and LuP30 did increase total amount of nitrogen shoot plant but not significant. The actinobacteria showed the best impact on the nutrient in shoot of lucerne at 25 mg N after 4 weeks inoculation with *S. meliloti* RRI 128 EN23. Three actinobacteria treatment plants had higher the content of iron, manganese, boron, copper, molybdenum, zinc, and macro elements such as calcium, potassium, phosphate and nitrogen.

EXAMPLE 11 – EFFECT OF ACTINOBACTERIA/RHIZOBIA CO-INOCULATION ON GROWTH PARAMETERS OF LUCERNE AT DIFFERING DOSING OF RHIZOBIA INOCULATION

Increasing *S. meliloti* RRI 128 dose concentrations resulted in slight increases number of nodules and of the growth of the plant. The number of nodules per plant increased from 4.3 to 7.0 and 8.8 nodules when the concentration of rhizobia was increased from 5×10^2 to 5×10^4 and 5×10^6 respectively (see Table 11). The significant effects of LuP30 and LuP47B on plant growth and

nodulation of lucerne plants were on 5×10^2 CFU.ml⁻¹ of *S. meliloti* RRI 128 (See Figure 8). The shoot dry weight and total mass per plant were increased up to about 50% to 60% and was similar with plants treated with the rhizobia at 10^4 and 10^6 CFU.ml⁻¹ (see Figure 8A). In addition, co-inoculation with either LuP30 or LuP47B individually with *S. meliloti* RRI 128 at 5×10^2 CFU.ml⁻¹ increased the number of nodules up to 7 and 9 per plant, respectively while control plants had 4.3 nodules per plant.

EXAMPLE 12 – ¹⁵N EXPERIMENT

10

Streptomyces spp. LuP30 and LuP47B were added as spores to lucerne seed with a sterile 0.3% xanthan gum solution and air dried before sowing. Seeds were treated with *S. meliloti* RRI 128. The planting process was as described in EXAMPLE 10.

15

The nitrogen supplied was ¹⁵NH₄¹⁵NO₃ (98%) with initial N concentration in soil (25 mg/kg ¹⁵NH₄¹⁵NO₃). Plants were harvested after 10, 21 and 35 days after inoculation with the *S. meliloti* RRI 128. Nitrogen in shoot and root materials was analysed by mass spectrometry to determine the proportions of plant N derived from the atmosphere and soil.

20

The plants were harvested at three times - at 10, 21 and 35 days after inoculation with *S. meliloti* RRI 128. The effectiveness of LuP30 and LuP47B was re-confirmed by the increase of the shoot dry weight and the number of nodules after 21 and 35 days co-inoculation with *S. meliloti* RRI 128 (Figure 9) and (Table 12). The amount of ¹⁵N and ¹⁴N in plants was estimated by spectrometry and the total N in whole plant co-inoculation with LuP30 or LuP47B was increased up to 40 % and 60 % respectively compared with plants treated *S. meliloti* RRI 128 alone (Figure 10). This was mostly due to greater accumulation of ¹⁴N (derived from N₂-fixation) which was increased by LuP30 or LuP47B by 47% and 72%, respectively. The actinobacteria significantly

25

30

- 45 -

increased the amount of ^{14}N in their plants while LuP47B also increased the amount of ^{15}N in their shoot and root (Table 13).

EXAMPLE 13 – EFFECT OF ACTINOBACTERIA ON GROWTH AND
5 SYMBIOSIS OF CLOVER

Clover cultivar Campeda (*Trifolium subterraneum* L.), was chosen to examine the effects of the two actinobacteria LuP30 and LuP47B which have shown an increase in growth and nitrogen fixation of lucerne in previous experiments.
10 Rhizobial strain *Rhizobium* WSM 1325 was inoculated on seeds of clover.

The factorial experiment comprised (i) two strains of actinobacteria (LuP30 and LuP47B), (ii) inoculation with rhizobia strain WSM1325 for clover. Growth of rhizobia and actinobacteria, plant growth media and nutrition, sowing and water
15 supply were as described above. The concentration of NH_4NO_3 was supplied at 25 mg per kg of sand and vermiculite where the actinobacteria LuP30 and LuP47B showed increased plant growth and nitrogen fixation for lucerne plants. Eight replicates for each treatment with four pots each harvested at 4 and 7 weeks after inoculation with rhizobia.

20

Co-inoculation of LuP30 with WSM 1325 increased the number of nodules after 7 weeks and nodule mass after 4 and 7 weeks in clover (Table 14). Actinobacteria strain LuP47B co-inoculated with *Rhizobium* WSM 1325 significantly increased the dry weight of shoot, total mass and number of
25 nodules per plant after 4 and 7 weeks inoculation with the *Rhizobium* while the nodule mass per plant was only increased after 7 weeks (see Figure 11 and Table 14). There was a significant change in the dry wt. of root of plants between the two harvests; for example, LuP30 increased root dry weight after 4 weeks while LuP47B increased root dry weight after 7 weeks.

30

EXAMPLE 14 – IN VITRO INTERACTION OF RHIZOBIA AND ACTINOBACTERIA

The interaction between LuP30 or LuP47B on the growth of two rhizobia
5 *Rhizobium* WSM 1325 and *Bradyrhizobium* WSM 471 was studied at three concentrations 10^4 , 10^6 and 10^8 CFU.ml⁻¹ (or 10^3 , 10^5 and 10^7 cells on each agar plate) of the two rhizobial strains. The two actinobacteria LuP30 and LuP47B were grown on ISP2 for 7-10 days and agar plugs of the well grown cultures were placed onto the agar plates containing the three rates of rhizobia. The
10 growth of the rhizobia was examined 5 to 14 days after adding the actinobacteria plugs.

LuP30 and LuP47B showed positive and non-antagonistic effects on the growth of both rhizobia (*Rhizobium* WSM 1325 and *Bradyrhizobium* WSM 471). At low
15 concentrations of rhizobia, less than 10^7 CFU.ml⁻¹ or 10^5 CFU.ml⁻¹ on each agar plate, LuP30 and LuP47B promoted a visible increase in the growth of both rhizobia on YMA medium after 5 days incubation at 27°C (Table 15). When the concentration of the rhizobia was more than 10^7 CFU.ml⁻¹ the effects of two actinobacteria LuP30 and LuP47B were not obvious on the growth of the two
20 rhizobia as analysed by visual observation, as was observed with the low rhizobial concentrations. These results show that rhizobial strains obtain growth benefits and are not inhibited by the two actinobacteria LuP30 and LuP47B.

EXAMPLE 15 – EFFECT OF ACTINOBACTERIA ON GROWTH AND 25 SYMBIOSIS OF SOYBEAN PLANTS (*GLYCINE MAX*)

The overall aim of this experiment was to evaluate a range of endophytic actinobacterial strains on the growth of soybean plants to determine whether these strains have a broad leguminous plant host range. The results of the
30 study of plants harvested 4 weeks after the addition of the *Bradyrhizbium* inoculum to the actinobacterial-treated plants showed that 4 of the 18 strains

tested had significantly improved plant growth and/ or nitrogen content of the soybean plants.

Materials and Methods

5

Soybean seeds (*Glycine max* cv. Djackal) were surface-sterilised and coated with spores of actinobacteria suspended in 0.3% (v/w) xanthan gum. Coated seeds (1 per pot) were sown into a pasteurised potting mix ~1 kg (50:50 by volume of sand:vermiculite) contained in 1.25 L pots. 200 ml of nitrogen deficient McKnight's nutrient solution was applied at sowing and supplemented to provide 25 mg of NH_4NO_3 per kg of potting media. Pots were arranged in a randomised block design with 5 replicates of each treatment. Plants were harvested at 4 weeks and 7 weeks after inoculation with *Bradyrhizobium* strain CB 1809.

15

Treatments were nil control, rhizobia with no actinobacteria, and rhizobium plus each of the following *Streptomyces* strains: *Str.* EN23, *Str.* EN27, *Str.* LuP3, *Str.* LuP5, LuP8, *Str.* LuP10, *Str.* LuP12A, *Str.* LuP30, *Str.* LuP44, *Str.* LuP46B, *Str.* LuP47B, *Str.* LuP73B, *Str.* LuP75, *Str.* PG3, *Str.* PG4, *Str.* PP1, *Str.* PP9, 20 CM23.

Parameters measured were:

- Length, dry weight of shoot and root.
- Number and total mass of nodules per plant.
- 25 • Nitrogen, P and trace elements in the plant shoots.

Effectiveness of Bradyrhizobium CB 1809 on nodulation and plant growth of soybean

30 Inoculation with *Bradyrhizobium* CB 1809 in the absence of actinobacteria resulted in abundant nodulation (around 120 nodules per plant) and increased shoot and root dry weights confirming the effectiveness of the *Bradyrhizobium*

strain with the cultivar Djackal. There were no obvious constraints to nodulation, in the testing system.

5 *Effect of actinobacteria on plant growth and symbiosis of soy plants at 4 weeks post inoculation*

Data for four (isolated from lucerne root) of the 18 actinobacteria tested are presented, based on their positive effects. Thirteen of the strains did not affect any of the parameters measured. As shown in Table 16, plants treated with *Str.* LuP8 and *Bradyrhizobium* CB 1809 strain showed increases in plant growth compared to plants only inoculated with CB1809. *Str.* LuP47B increased dry weight of shoots and total plant weight (+15%) and LuP30 increased total plant weight (+12%). *Str.* LuP30, LuP44 and LuP47B also improved the nodule mass per plant by 20, 22 and 29 % respectively.

15 As shown in Table 17, plants treated with LuP47B also had increased levels of iron, magnesium, phosphorus and nitrogen (27 %) compared with plants inoculated with *Bradyrhizobium* CB1809 only. LuP8 increased total N (23 %) and also iron. LuP30 increased iron content.

20 *Effect of actinobacteria on plant growth and symbiosis of soy plants at 7 weeks post inoculation*

As shown in Table 18, statistically significant results were:

- 25 - LuP47B increased height of shoots by 38%;
- LuP8 increased total nodule mass by 54%;
- LuP8 increased fresh weight of pods by 24 % and number of seeds per plant by 35%; and
- LuP47B increased the dry weight of seeds by 43% while LuP8 increased
- 30 total dry weight of shoot and pods per plant by 24%.

EXAMPLE 16 – EFFECT OF ACTINOBACTERIA ON GROWTH AND SYMBIOSIS OF FIELD PEA (*PISUM* SP.)

Pea field trials were sown Hart (28 May) and Riverton (10 June) in South
 5 Australia (SA), and at Pimpinio (15 May) in Victoria. The trials were arranged in
 randomised complete block design with 3 replicates, each comprising an
 uninoculated control and 3 inoculation treatments. Treatments were applied to
 Kasper field pea which was sown to achieve a seedling density of 50 plants/m².
 The rhizobia treatment (*Rhizobium leguminosarum* bv. *viciae* strain WSM1455)
 10 was applied at approximately 100 fold the rate recommended commercially.
 The co-inoculation treatment comprised the rhizobia treatment co-inoculated
 with *Streptomyces* sp. strain Lup47B, which was applied as spores to the seed
 before sowing.

15 Six plants were sampled from each plot at approximately 8 weeks after sowing
 and nodule number and nodule dry weight per plant determined. An additional
 ten plant shoots were sampled from each plot in Oct./Nov. (late pod fill) and
 used to estimate shoot biomass, pod number per plant and to estimate the % N
 derived from fixation using the ¹⁵N natural abundance method. Plots were
 20 machine harvested to estimate grain yield and subsamples used for the
 determination of grain protein (Total N Leco, CSBP).

Table 19 shows nodule number, nodule weight, pod number and total plant
 biomass in pea plants grown in field trials at three sites (Riverton SA, Hart SA
 25 and Pimpinio Vic). A summary of the results shown in table 19 is:

- A significant effect on nodulation was seen at Pimpinio, where the
 actinobacteria (LuP47B) / rhizobium coinoculation significantly increased the
 number of nodules relative to the rhizobium only control; and
- Actinobacteria (LuP47B) / rhizobium coinoculation increased biomass at all
 30 sites relative to rhizobium only inoculation, with two sites and the mean of all
 sites achieving statistically significant increases in biomass.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, 5 compositions and compounds referred to, or indicated in this specification, individually or collectively, and any and all combinations of any two or more of the steps or features.

Also, it must be noted that, as used herein, the singular forms “a”, “an” and “the” 10 include plural aspects unless the context already dictates otherwise. Thus, for example, reference to “a microorganism” includes a single microorganism as well as two or more microorganisms; “a leguminous plant” includes a single plant as well as two or more plants; and so forth.

15 Reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common general knowledge in any country.

Throughout this specification, unless the context requires otherwise, the word 20 “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

25 Reference is made to standard textbooks of molecular biology that contain methods for carrying out basic techniques encompassed by the present invention, including DNA restriction and ligation for the generation of the various genetic constructs described herein. See, for example, Maniatis *et al*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, New 30 York, 1982) and Sambrook *et al*. (2000, *supra*).

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for enhancing at least one growth parameter of a leguminous plant, the method comprising co-inoculating the leguminous plant with:
 - 5 at least one rhizobial microorganism; and
 - at least one actinobacterial microorganism;wherein the co-inoculated leguminous plant has at least one enhanced growth parameter relative to a leguminous plant of the same taxon that has not been co-inoculated.
- 10 2. The method of claim 1 wherein the actinobacterial microorganism is an endophytic actinobacterial microorganism.
3. The method of claim 1 wherein the actinobacterial microorganism is from
15 a genus selected from *Streptomyces*, *Microbispora* or *Micromonospora*.
4. The method of claim 1 wherein the actinobacterial microorganism is from the genus *Streptomyces*.
- 20 5. The method of claim 4 wherein actinobacterial microorganism is from a species selected from: *Streptomyces drozdowiczii*, *Streptomyces ciscaucasicus*, *Streptomyces canus*, *Streptomyces rishiriensis*, *Streptomyces badius* or *Streptomyces parvus*.
- 25 6. The method of claim 1 wherein the actinobacterial microorganism comprises a 16S rRNA gene nucleotide sequence which is at least 90% identical to one or more of: SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9 or SEQ ID NO: 10.
- 30 7. The method of claim 1 wherein the actinobacterial microorganism is *Streptomyces* sp. LuP30 as deposited under NMI accession number V13/030101; or a mutant or derivative of said microorganism that retains the

ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

5 8. The method of claim 1 wherein the actinobacterial microorganism is *Streptomyces* sp. LuP47B as deposited under NMI accession number V13/030100; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with
10 a rhizobial microorganism.

9. The method of claim 1 wherein the actinobacterial microorganism is *Streptomyces* sp. EN23 as deposited under as AGAL Deposit No. NM03/35605; or a mutant or derivative of said microorganism that retains the
15 ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism co-inoculated onto a leguminous plant with a rhizobial microorganism.

10. The method of claim 1 wherein the actinobacterial microorganism is
20 *Streptomyces* sp. EN27 as deposited as AGAL Deposit No. NM03/35606; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

25

11. The method of claim 1 wherein the growth parameter is a length and/or mass of a shoot of the leguminous plant.

12. The method of claim 1 wherein the growth parameter is a length and/or
30 mass of a root of the leguminous plant.

13. The method of claim 1 wherein the growth parameter is a number and/or

mass of nodules of the leguminous plant.

14. The method of claim 1 wherein the growth parameter is a number and/or mass of seed pods and/or seed produced by the leguminous plant.

5

15. The method of claim 1 wherein the growth parameter is a concentration and/or amount of a nutrient in the leguminous plant.

16. The method of claim 15 wherein the nutrient is selected from: Boron,
10 Calcium, Copper, Magnesium, Manganese, Phosphorous, Sodium, Sulphur, Nitrogen and/or Zinc.

17. The method of claim 16 wherein the nutrient is Nitrogen.

15 18. The method of claim 1 wherein the growth parameter is a germination rate of a leguminous plant.

19. The method of claim 1 wherein the rhizobial microorganism is a *Sinorhizobium* sp.

20

20. The method of claim 1 wherein the rhizobial microorganism is a *Rhizobium* sp.

21. The method of claim 1 wherein the rhizobial microorganism is a
25 *Bradyrhizobium* sp.

22. The method of claim 1 wherein the leguminous plant is selected from the group of: a *Medicago* sp., a *Trifolium* sp., a *Pisum* sp., or a *Glycine* sp.

30 23. The method of claim 1 wherein the leguminous plant is exposed to a pathogen and, when exposed to the pathogen, the co-inoculated leguminous

plant has at least one enhanced growth parameter relative to a leguminous plant of the same taxon that has not been co-inoculated.

24. The method of claim 23 wherein the pathogen is a root pathogen.

5

25. The method of claim 23 wherein the pathogen is a fungal pathogen.

26. The method of claim 25 wherein the pathogen is a *Rhizoctonia* sp.

10 27. The method of claim 26 wherein the pathogen is *Rhizoctonia solani*.

28. A leguminous plant, leguminous plant part or leguminous plant reproductive material, co-inoculated with at least one rhizobial microorganism and at least one actinobacterial microorganism.

15

29. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism is an endophytic actinobacterial microorganism.

20 30. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism is from a genus selected from *Streptomyces*, *Microbispora* or *Micromonospora*.

25 31. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism is from the genus *Streptomyces*.

30 32. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 31 wherein actinobacterial microorganism is from a species selected from: *Streptomyces drozdowiczii*, *Streptomyces ciscaucasicus*, *Streptomyces canus*, *Streptomyces rishiriensis*, *Streptomyces badius* or *Streptomyces parvus*.

33. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism comprises a 16S rRNA gene nucleotide sequence which is at least 90% identical to one or more of: SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10.

34. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism is *Streptomyces* sp. LuP30 as deposited under NMI accession number V13/030101; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

15

35. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism is *Streptomyces* sp. LuP47B as deposited under NMI accession number V13/030100; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

20

36. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the actinobacterial microorganism is *Streptomyces* sp. EN23 as deposited under as AGAL Deposit No. NM03/35605; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism co-inoculated onto a leguminous plant with a rhizobial microorganism.

25

30

37. The leguminous plant, leguminous plant part or leguminous plant

reproductive material of claim 28 wherein the actinobacterial microorganism is *Streptomyces* sp. EN27 as deposited as AGAL Deposit No. NM03/35606; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

38. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the rhizobial microorganism is a *Sinorhizobium* sp.

39. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the rhizobial microorganism is a *Rhizobium* sp.

40. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the rhizobial microorganism is a *Bradyrhizobium* sp.

41. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 28 wherein the leguminous plant is selected from the group of: a *Medicago* sp., a *Trifolium* sp., a *Pisum* sp., or a *Glycine* sp.

42. The leguminous plant, leguminous plant part or leguminous plant reproductive material of claim 41 wherein the leguminous plant has at least one enhanced growth parameter relative to a leguminous plant of the same taxon that has not been co-inoculated.

43. An isolated actinobacterial microorganism as deposited under NMI accession number V13/030101; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a

leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

44. An isolated actinobacterial microorganism as deposited under NMI
5 accession number V13/030100; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

10 45. An inoculant composition comprising at least an actinobacterial microorganism.

46. The inoculant composition of claim 45 wherein the actinobacterial microorganism is an endophytic actinobacterial microorganism.

15

47. The inoculant composition of claim 45 wherein the actinobacterial microorganism is from a genus selected from *Streptomyces*, *Microbispora* or *Micromonospora*.

20 48. The inoculant composition of claim 45 wherein the actinobacterial microorganism is from the genus *Streptomyces*.

49. The inoculant composition of claim 48 wherein actinobacterial microorganism is from a species selected from: *Streptomyces drozdowiczii*,
25 *Streptomyces ciscaucasicus*, *Streptomyces canus*, *Streptomyces rishiriensis*, *Streptomyces badius* or *Streptomyces parvus*.

50. The inoculant composition of claim 45 wherein the actinobacterial microorganism comprises a 16S rRNA gene nucleotide sequence which is at
30 least 90% identical to one or more of: SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10.

51. The inoculant composition of claim 45 wherein the actinobacterial microorganism is *Streptomyces* sp. LuP30 as deposited accession number V13/030101; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

52. The inoculant composition of claim 45 wherein the actinobacterial microorganism is *Streptomyces* sp. LuP47B as deposited under accession number V13/030100; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

15

53. The inoculant composition of claim 45 wherein the actinobacterial microorganism is *Streptomyces* sp. EN23 as deposited under as AGAL Deposit No. NM03/35605; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism co-inoculated onto a leguminous plant with a rhizobial microorganism.

54. The inoculant composition of claim 45 wherein the actinobacterial microorganism is *Streptomyces* sp. EN27 as deposited as AGAL Deposit No. NM03/35606; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

55. The inoculant composition of claim 45 wherein the inoculant composition further comprises a rhizobial microorganism.

- 59 -

56. The inoculant composition of claim 55 wherein the rhizobial microorganism is a *Sinorhizobium* sp.

57. The inoculant composition of claim 55 wherein the rhizobial
5 microorganism is a *Rhizobium* sp.

58. The inoculant composition of claim 55 wherein the rhizobial microorganism is a *Bradyrhizobium* sp.

10 59. The inoculant composition of claim 45 further comprising a carrier or additive.

60. The inoculant composition of any one of claims 45 to 59 when used according to the method of claim 1.

15

61. An isolated actinobacterial microorganism as deposited under NMI accession number V13/030101 or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a leguminous plant when the actinobacterial microorganism is co-inoculated onto
20 a leguminous plant with a rhizobial microorganism.

62. An isolated actinobacterial microorganism as deposited under NMI accession number V13/030100; or a mutant or derivative of said microorganism that retains the ability to enhance at least one growth parameter of a
25 leguminous plant when the actinobacterial microorganism is co-inoculated onto a leguminous plant with a rhizobial microorganism.

FIGURE 1





FIGURE 2




FIGURE 3



3/22

FIGURE 4

A



B



FIGURE 5





FIGURE 6



5/22

FIGURE 7





FIGURE 8




B




7/22

FIGURE 9




8/22

FIGURE 10



9/22

FIGURE 11



10/22

TABLE 1

Cultures	IAA production	Phosphate solubilisation
LuP3	+	-
LuP5B	-	++
LuP8	+	+
LuP10	-	-
LuP12A	+	-
LuP30	++	-
LuP44	+	+
LuP46(2)	+	-
LuP47B	++	-
LuP73B	+	-
LuP83	+	-

TABLE 2

Cultures	OD _{600nm} <=0.06 < =6x10 ⁷ cfu/ml			OD _{600nm} <=0.1 <= 1x10 ⁸ cfu/ml		
	RRI128	SRDI736	WSM1115G	RRI128	SRDI736	WSM1115G
EN16	0	0	0	0	0	0
EN23	0	0	++	0	0	+
EN27	0	0	0	0	0	0
EN28	0	0	-	0	0	-
EN46	-	0	0	0	0	0
LuP3	++	+	+	++	+	+
LuP8	0	0	+	0	0	++
LuP10	++	--	-	+	--	-
LuP12 A	0	0	0	0	0	++
LuP30	++	++	++	++	+	++
LuP44	0	+	+	0	0	+
LuP46(2)	0	+	0	0	0	+
LuP47B	+	+	+	0	0	++
LuP83	0	0	++	0	0	+

TABLE 3

Lucerne treatment	Height of shoot (cm)	Shoot dry weight (mg)	Length of root (cm)	Root dry weight (mg)	
Untreated*	9.21 ^a	30.23 ^a	24.64 ^{bc}	13.55 ^{ab}	
RRI128 only	12.55 ^b	50.63 ^b	23.57 ^{ab}	17.49 ^{ab}	
RRI128 + EN2	13.94 ^{bcd}	64.17 ^{cd}	30.13 ^d	20.30 ^b	
RRI128 + EN16	12.75 ^b	52.55 ^b	18.44 ^a	12.44 ^a	
RRI128 + EN23	15.20 ^d	69.15 ^d	28.66 ^{cd}	16.48 ^{ab}	
RRI128 + EN27	14.82 ^{cd}	59.71 ^{bcd}	20.79 ^{ab}	20.87 ^b	
RRI128 + EN28	13.06 ^{bc}	55.45 ^{bc}	19.63 ^{ab}	17.38 ^{ab}	
RRI128+ EN46	13.48 ^{bcd}	58.30 ^{bc}	22.00 ^{ab}	16.00 ^{ab}	
	Nodule fresh weight (mg)	No. of nodules	SPAD 502 reading	N content (%)	N per plant (mg)
Untreated*	0 ^a	0 ^a	ND	ND	ND
RRI128 only	0.58 ^b	22.22 ^{cd}	28.31 ^a	2.725 ^{a**}	1.38
RRI128 + EN2	1.02 ^{cd}	23.55 ^{cd}	29.64 ^{ab}	ND	ND
RRI128 + EN16	0.86 ^{bc}	17.42 ^b	29.62 ^{ab}	ND	ND
RRI128 + EN23	1.27 ^d	25.74 ^d	31.46 ^{bc}	3.225 ^{b**}	2.23
RRI128 + EN27	0.78 ^{bc}	16.85 ^b	31.83 ^c	3.65 ^{c**}	2.18
RRI128 + EN28	0.74 ^{bc}	21.55 ^{bcd}	28.91 ^a	ND	ND
RRI128+ EN46	0.91 ^{bc}	19.70 ^{bc}	28.95 ^a	ND	ND

Untreated*: seeds were coated with 0.3% xanthan gum without rhizobium. The different letters in the same column indicate a significant difference ($p < 0.05$). Data was analyzed using one-way ANOVA and Duncan test.

12/22

TABLE 4

Treatment	Height of shoot (cm)	Length of root (cm)	Shoot dry weight (mg)	Root dry weight (mg)	Total Mass (mg)	No of nodules	DW of nodule (mg)	Total nodule mass (mg)
Untreated*	4.85 ^a	18.00 ^a	11.75 ^a	15.25 ^a	27 ^a	0 ^a	0 ^a	0 ^a
RRI128 only	12.19 ^b	18.77 ^{ab}	78.4 ^b	35.6 ^b	114 ^b	22.10 ^b	0.1081 ^b	2.280 ^b
RRI128 + EN23	13.33 ^{bc}	23.80 ^{ab}	89.73 ^{bc}	43.56 ^b	132.29 ^{bc}	21.90 ^b	0.1238 ^{bc}	2.711 ^{bc}
RRI128 + EN27	12.21 ^{bc}	27.56 ^{bc}	80.76 ^{bc}	47.78 ^{bc}	127.54 ^{bc}	19.89 ^b	0.1054 ^b	2.0964 ^b
RRI128 + LuP3	12.96 ^{bc}	20.47 ^{ab}	83.27 ^{bc}	44.2 ^b	127.47 ^{bc}	22.50 ^b	0.1061 ^b	2.390 ^b
RRI128 + LuP5B	11.17 ^{bc}	23.45 ^{ab}	74.59 ^b	42.78 ^b	117.37 ^b	16.70 ^b	0.1356 ^{bc}	2.264 ^b
RRI128 + LuP10	14.22 ^{bc}	21.30 ^{ab}	80.93 ^{bc}	58.89 ^c	139.82 ^{bc}	18.22 ^b	0.1261 ^{bc}	2.297 ^b
RRI128 + LuP12A	13.95 ^{bc}	22.33 ^{ab}	81.9 ^{bc}	40.3 ^b	122.20 ^{bc}	18.30 ^b	0.1072 ^b	1.962 ^b
RRI128 + LuP30	13.83 ^{bc}	32.55 ^c	97.9 ^{cd}	45.6 ^{bc}	143.50 ^c	20.80 ^b	0.0985 ^b	2.0610 ^b
RRI128 + LuP44	12.91 ^{bc}	21.57 ^{ab}	83.7 ^{bc}	49.2 ^{bc}	132.90 ^{bc}	20.50 ^b	0.1070 ^b	2.1935 ^b
RRI128 + LuP47B	15.39 ^c	25.92 ^{bc}	106.1 ^d	42.00 ^b	148.10 ^c	20.50 ^b	0.1615 ^c	3.380 ^c

Untreated*: seeds were coated with 0.3% xanthan gum without rhizobium. The different letters in the same column indicate a significant difference ($p < 0.05$). Data was analyzed using one-way ANOVA and Duncan test.

TABLE 5

Treatment	RRI128 only	RRI128 + EN23	RRI128 + LuP30	RRI128 + LuP47B
Boron (mg)	3.759a	3.932a	4.621b	4.631b
Calcium (µg)	0.839a	0.933ab	1.002b	1.181c
Copper (µg)	0.602a	0.745b	0.860c	0.738b
Iron (µg)	23.66a	21.13a	24.38a	21.90a
Magnesium (mg)	0.732a	0.784a	0.904b	0.983b
Manganese (µg)	12.54a	12.89a	14.82b	15.18b
Phosphorus (mg)	0.086a	0.111b	0.120c	0.121c
Sodium (mg)	0.102a	0.144bc	0.137b	0.173c
Sulfur (mg)	0.191a	0.212ab	0.228bc	0.244c
Nitrogen (mg)	2.598a	2.949b	3.205c	3.431d
Zinc (µg)	18.48a	18.96ab	22.75c	21.58bc

The different letters in the same column indicate a significant difference ($p < 0.05$). Data was analyzed using one-way ANOVA and Duncan test.

TABLE 6

Treatment	Height of shoot	Length of root	Shoot dry weight	Root dry weight	Total Mass	No of nodules
Untreated*	3 ^a	15.25 ^b	6.9 ^a	7.9 ^a	14.8 ^a	0 ^a
RRI128 only	3.6 ^{abc}	11.83 ^a	17.33 ^{bc}	12.67 ^{ab}	30.00 ^b	6.33 ^{bcd}
RRI128 + LuP3	3.96 ^{bc}	15.8 ^{bc}	19.4 ^{bc}	17.2 ^{bcd}	36.6 ^{bcd}	7 ^{bcd}
RRI128 + LuP5B	4.34 ^c	18.7 ^c	20.4 ^c	18.8 ^{cde}	39.2 ^{cd}	6.2 ^{bcd}
RRI128 + LuP12A	4.18 ^{bc}	13.62 ^{ab}	18.2 ^{bc}	22 ^e	40.2 ^d	6 ^{bcd}
RRI128 + LuP30	3.56 ^{ab}	13.32 ^{ab}	15.4 ^b	15.6 ^{bc}	31 ^{bc}	9.8 ^{de}
RRI128 + LuP47B	3.74 ^{abc}	16.16 ^{bc}	20 ^c	16.2 ^{bcd}	36.2 ^{bcd}	10 ^{de}

Untreated*: seeds were coated with 0.3% xanthan gum without rhizobium. The different letters in the same column indicate a significant difference ($p < 0.05$). Data was analyzed using one way ANOVA and Duncan test.

TABLE 7

	Height of shoot (cm)	Shoot dry weight (mg per plant)	Root dry weight (mg per plant)	Total biomass (mg per plant)
Control 1	3.74 a	10.01 a	11.19 ab	21.20 a
Control 2	2.2 d	2.67 d	4.27 e	6.74 e
Control 3	2.95 c	6.6 c	4.9 de	11.50 de
R + EN16	3.07 c	6.82 bc	7.31 cde	14.13 cd
R + LuP10	3.59 ab	8.09 bc	7.75 cde	15.84 c
R + LuP12A	3.12 bc	7.67 bc	6.45 cde	14.12 cd
R + LuP30	2.95 c	7.3 bc	8.52 bcd	15.82 c
R + LuP44	2.97 c	6.85 bc	9.34 bc	16.19 bc
R + LuP46(2)	3.37 abc	8.51 ab	7.78 cde	16.29 bc
R + LuP47B	3.36 abc	7.27 bc	8.11 bcde	15.38 c
R + LuP73B	3.08 bc	7.33 bc	13.1 a	20.43 ab
R + LuP83	3.12 bc	7.16 bc	7.69 cde	14.75 cd

Control 1, only rhizobium RRI128 and no *R. solani*; Control 2, without rhizobium RRI128 and *R. solani* applied; Control 3, rhizobium RRI128 and *R. solani* applied; R +, rhizobium RRI128 plus endophytic actinobacteria.

TABLE 8

	Closest match in GenBank using BLASTN			
	Organism	Accession number	Bits	% Identity
LuP3	<i>Streptomyces drozdowiczii</i> NRRL B-24297	EF654097	2620	99.40
	<i>Streptomyces drozdowiczii</i> NBRC 101007	NR041424		99.40
LuP12A	<i>Streptomyces ciscaucasicus</i> NBRC 12872	AB184208	2704	99.78
	<i>Streptomyces canus</i> NBRC 12752	AB184118		99.78
LuP30	NRRL B-3239	EF178682	2728	99.90
	<i>Streptomyces rishiriensis</i> NBRC 13407	AB184383		99.90
LuP47B	<i>Streptomyces ciscaucasicus</i> NBRC 12872	AB184208	2334	99.49
	<i>Streptomyces canus</i> NBRC 12752	AB184118		99.49
EN23	<i>Streptomyces badius</i> NRRL B-2567	NR_043350.1	2590	99%
EN27	<i>Streptomyces parvus</i> Strain sj38	JX013965.1	1815	94%
LuP8	<i>Streptomyces ciscaucasicus</i> NBRC 12872	NR_041085.1	1251	99.2%
	<i>Streptomyces canus</i> NBRC 12752	NR_112259.1	1251	99.2%
LuP44	<i>Streptomyces ciscaucasicus</i> NBRC 12872	NR_041085.1	2477	99.7%
	<i>Streptomyces canus</i> NBRC 12752	NR_112259.1	2477	99.7%

16/22

TABLE 9

Treatment	Shoot:root			Shoot weight DM (mg /plant)			Root weight DM (mg /plant)		
	A	B	C	A	B	C	A	B	C
Single inoculation with <i>Streptomyces</i>									
Control ^a	1.1	1.4	1.9	19.7	79.3	173	17.8	55.8	91
EN23	1.2	2.1 [*]	2.9 ^{**}	19.9	103.5 [*]	215.9 [*]	17.3	48.6	75.1 [*]
LuP30	1.0	2.2 [*]	2.6 [*]	20.3	94.1 [*]	215.9 [*]	19.6	43.8	83.6
LuP47B	1.2	2.6 ^{**}	2.7 [*]	19.7	103 [*]	212 [*]	17.1	39.6 [*]	78.5
Co-inoculation with <i>S.meliloti</i> RRI 128									
Control ^b	1.8	2.7	2.1	229	268	348	129	100	169
EN23	2.1	2.3	2.5	238	340 [*]	436 ^{**}	114	151 [*]	172
LuP30	3.7 ^{**}	2.8	2.7 [*]	283 [*]	392 ^{**}	379 [*]	76 ^{**}	141 [*]	142 [*]
LuP47B	2.8 [*]	3.5 [*]	2.0	284 [*]	365 ^{**}	331	101 [*]	109	163

TABLE 10

Treatment	4 weeks			7 weeks		
	3mg N	25mg N	50mg N	3mg N	25mg N	50mg N
RRI 128	22 ^a	24 ^a	40 ^a	27 ^a	47 ^a	69 ^b
RRI 128 + EN23	21 ^a	53 ^b	52 ^b	30 ^{ab}	51 ^a	56 ^a
RRI 128 + LuP30	31 ^b	51 ^b	48 ^b	41 ^c	57 ^a	66 ^{ab}
RRI 128 + LuP47B	24 ^a	42 ^b	36 ^a	37 ^{bc}	49 ^a	71 ^b

TABLE 11

Actinobacteria	<i>S.meliloti</i> RRI 128 concentrations (CFU.ml ⁻¹)		
	5 x 10 ²	5 x 10 ⁴	5 x 10 ⁶
Number of nodules per plant			
Nil	4.3 ± 0.9	7.0 ± 0.9	8.8 ± 1.5
LuP30	7.0 ± 1.1	7.3 ± 2.3	8.5 ± 1.9
LuP47B	9.0 ± 1.8	7.8 ± 1.1	9.8 ± 2.9

TABLE 12

Treatment	Shoot:root			Shoot weight DM			Root weight DM			Number of nodules (#/plant)		
				(mg /plant)			(mg /plant)					
	10d	21d	35d	10d	21d	35d	10d	21d	35d	10d	21d	35d
Single inoculation with <i>Streptomyces</i>												
Control ^a	2.4	1.9	1.8	5.8	31.0	64.9	2.4	16.2	36.5	0	0	0
LuP30	2.6	1.9	1.9	5.7	30.1	67.2	2.2	16.2	36.3	0	0	0
LuP47B	2.0	1.9	1.7	5.1	28.8	65.1	2.5	15.1	37.6	0	0	0
Co-inoculation with <i>S.meliloti</i> RRI 128												
Control ^b	2.5	1.9	1.9	5.8	25.9	87.6	2.3	13.7	45.9	0	18.6	30.0
LuP30	2.3	2.2	2.1	5.8	33.7*	110.2*	2.5	15.3	52.8	0	24.6*	48.6*
LuP47B	2.3	2.1	1.9	5.4	34.3*	110.7*	2.3	16.1	59.6*	0	22.8*	37.2

TABLE 13

Treatment	Shoot (µg)					
	21d			35d		
	¹⁵ N	¹⁴ N	Total N	¹⁵ N	¹⁴ N	Total N
RRI 128 only	256 ^a	489 ^a	745 ^a	543 ^a	1732 ^a	2275 ^a
RRI 128 + LuP30	312 ^{ab}	503 ^a	816 ^a	658 ^a	2564 ^b	3222 ^b
RRI 128 + LuP47B	343 ^b	586 ^a	929 ^a	648 ^a	2956 ^b	3605 ^b
Treatment	Root (µg)					
	21d			35d		
	¹⁵ N	¹⁴ N	Total N	¹⁵ N	¹⁴ N	Total N
RRI 128 only	102 ^a	175 ^a	276 ^a	279 ^a	735 ^a	1014 ^a
RRI 128 + LuP30	116 ^{ab}	184 ^a	300 ^a	313 ^{ab}	1062 ^{ab}	1375 ^{ab}
RRI 128 + LuP47B	125 ^b	197 ^a	322 ^a	364 ^b	1280 ^b	1644 ^b

TABLE 14

Treatment	Root length (cm)		Root weight (mg DM/ plant)		Nodule number (#/plant)		Total nodule mass (mg)	
	4 w	7 w	4 w	7 w	4 w	7 w	4 w	7 w
Untreated*	20 ^a	30 ^a	35.0 ^a	46.0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
WSM 1325 only	30 ^b	34 ^{ab}	63.5 ^b	69.8 ^b	114 ^b	119 ^b	7.5 ^b	8.3 ^b
WSM 1325 + LuP30	36 ^c	39 ^b	71.7 ^c	70.6 ^b	127 ^{bc}	168 ^c	9.3 ^c	11.2 ^c
WSM 1325 + LuP47B	29 ^b	31 ^a	59.6 ^b	79.3 ^c	138 ^c	175 ^c	7.9 ^b	11.7 ^c

TABLE 15

Cultures	≤ 10 ³ cfu/plate		≤ 10 ⁵ cfu/plate		≤ 10 ⁷ cfu/plate	
	WSM 1325	WSM 471	WSM 1325	WSM 471	WSM 1325	WSM 471
LuP30	++	++	++	++	+	+
LuP47B	++	++	++	++	+	+

TABLE 16

Treatment	DW of shoot (mg)	DW of root (mg)	Total DW (mg)	Number of nodules	Nodule mass (mg/plant)
Untreated	1320 ^a	511.8 ^a	1760 ^a	0 ^a	0 ^a
CB 1809	2120 ^b	515 ^a	2640 ^b	119 ^b	115 ^b
CB 1809 + LuP8	2520 ^c	717 ^b	3240 ^c	97 ^b	115 ^b
CB 1809 + LuP30	2330 ^{bc}	614.6 ^{ab}	2940 ^c	134 ^b	139 ^c
CB 1809 + LuP44	2170 ^b	606.8 ^{ab}	2780 ^b	109 ^b	141 ^c
CB 1809 + LuP47B	2500 ^c	528.2 ^a	3030 ^c	122 ^b	149 ^c

TABLE 17

Treatment	Copper (µg)	Iron (µg)	Magnesium (mg)	Phosphorous (mg)	Sodium (mg)	Nitrogen (mg)
Untreated	13.8 ^a	94 ^a	7.1 ^a	2.8 ^a	153 ^a	13.2 ^a
CB 1809	25.6 ^{bc}	162 ^b	9.5 ^b	2.9 ^a	499 ^{cd}	49.8 ^b
CB 1809 + LuP8	23.3 ^{bc}	203 ^c	11.1 ^{bc}	3.3 ^{ab}	404 ^c	61.6 ^c
CB 1809 + LuP30	23.9 ^{bc}	208 ^c	10.8 ^{bc}	3.2 ^{ab}	577 ^d	59.8 ^{bc}
CB 1809 + LuP44	24.5 ^{bc}	180 ^{bc}	10.5 ^{bc}	3.1 ^{ab}	496 ^{cd}	51.9 ^{bc}
CB 1809 + LuP47B	27.5 ^c	214 ^c	12.0 ^c	3.6 ^b	559 ^d	63.1 ^c

TABLE 18

Treatment	H of shoot (cm)	DW of shoot (g)	DW of root (g)	No of nodules (#/plant)	Total nodule mass (mg)
Untreated	25.7a	0.88a	0.41a	0a	0.0a
Rhi only	30.7a	1.10ab	0.39a	86b	91.5bc
Rhi + LuP8	34.8ab	1.38bc	0.39a	80b	140.7d
Rhi + LuP30	30.4a	1.12ab	0.37a	67b	85.5bc
Rhi + LuP44	34.0ab	1.09ab	0.37a	71b	89.3bc
Rhi + LuP47B	42.3b	1.26abc	0.34a	85b	109.8bc
Rhi + B	29.4a	1.06ab	0.43a	91b	115.9c
N	30.0a	1.58c	0.58b	0a	0.0a

Treatment	No of Pods (#/plant)	FW of pods (g)	DW of pod + seed (g)	No seeds (#/plant)	DW of seed (g)	Total mass of shoot and pods (g)
Untreated	2.5a	1.3a	0.34a	5.33a	0.07a	1.22a
Rhi only	5.0bc	3.7b	0.92bcd	9.13b	0.30b	2.02bc
Rhi + LuP8	6.1c	4.6c	1.12d	12.38c	0.36bc	2.50d
Rhi + LuP30	4.5bc	3.7b	0.83b	9.75bc	0.31b	1.95bc
Rhi + LuP44	5.3bc	3.7b	0.95bcd	10.75bc	0.31b	1.81b
Rhi + LuP47B	5.1bc	4.2bc	1.06cd	10.38bc	0.43c	2.33cd
Rhi + B	4.3b	3.8b	0.88bc	8.88b	0.36bc	1.94bc
N	6.0bc	4.7c	1.09d	10.5bc	0.44c	2.68d

TABLE 19

	Riverton SA				Hart SA			
	Nodule #	Nodule weight	Pod #	Peak Bioma ss	Nodule #	Nodule weight	Pod #	Peak Bioma ss
	#/ plant	mg/ plant	# per 10 plants	g per 10 plants	#/plant	mg/ plant	# per 10 plants	g per 10 plants
No inoculation	36	6.1	76	165	45	7.1	80	149
WSM1455 (Group F)	32	9.3	85	182	56	8.2	106	184
Actinobacteria (Lup47b)	26	4.7	109	229	47	7.2	104	193
L.S.D. (P = 0.05)	NS	NS	NS	38	NS	NS	19	43
	Pimpinio Vic				Mean of all sites			
	Nodule #	Nodule weight	Pod #	Peak Bioma ss	Nodule #	Nodule weight	Pod #	Peak Bioma ss
	#/plant	mg/ plant	# per 10 plants	g per 10 plants	#/plant	mg/ plant	# per 10 plants	g per 10 plants
No inoculation	29	12.5	8	75	36	9.0	50	124
WSM1455 (Group F)	28	14.8	7	75	37	11.2	60	140
Actinobacteria (Lup47b)	58	9.9	4	91	45	7.5	65	163
L.S.D. (P = 0.05)	12	NS	NS	NS	NS	NS	10	18