

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2008329065 B2**

(54) Title
Brassica plant comprising a mutant indehiscent allele

(51) International Patent Classification(s)
A01H 5/00 (2006.01) **C12N 15/82** (2006.01)

(21) Application No: **2008329065** (22) Date of Filing: **2008.11.25**

(87) WIPO No: **WO09/068313**

(30) Priority Data

(31) Number	(32) Date	(33) Country
07023052.9	2007.11.28	EP
61/004,660	2007.11.29	US

(43) Publication Date: **2009.06.04**

(44) Accepted Journal Date: **2014.05.08**

(71) Applicant(s)
Bayer CropScience NV

(72) Inventor(s)
Den Boer, Bart;Laga, Benjamin;Lambert, Bart

(74) Agent / Attorney
Davies Collison Cave, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000

(56) Related Art
WO 2006/009649
US 7135621

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 June 2009 (04.06.2009)

(10) International Publication Number
WO 2009/068313 A3

(51) **International Patent Classification:**
C12N 15/82 (2006.01) A01H 5/00 (2006.01)

(21) **International Application Number:**
PCT/EP2008/010147

(22) **International Filing Date:**
25 November 2008 (25.11.2008)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
07023052.9 28 November 2007 (28.11.2007) EP
61/004,660 29 November 2007 (29.11.2007) US

KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) **Applicant (for all designated States except US):** BAYER BIOSCIENCE N.V. [BE/BE]; Technologiepark 38, B-9052 Gent (BE).

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) **Inventors; and**
(75) **Inventors/Applicants (for US only):** LAGA, Benjamin [BE/BE]; Rakestraat 4, B-8750 Wingene (BE). DEN BOER, Bart [NL/BE]; Wilgenstraat 21, B-9820 Merelbeke (BE). LAMBERT, Bart [BE/BE]; Maarschalk Haiglaan 27, bus 4A, B-8900 Leper (BE).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

(74) **Common Representative:** BAYER BIOSCIENCE N.V.; Technologiepark 38, B-9052 Gent (BE).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))

[Continued on next page]

(54) **Title:** BRASSICA PLANT COMPRISING A MUTANT INDEHISCENT ALLELE




Figure 1

BnIND-A1 (1622 bps)

(57) **Abstract:** This invention relates to crop plants of which the fruit dehiscence properties are modulated. More specifically the invention relates to improved methods and means for reducing seed shattering, or delaying seed shattering until after harvest, in plants, while maintaining at the same time an agronomically relevant threshability of the pods.

WO 2009/068313 A3

WO 2009/068313 A3



— *with sequence listing part of description (Rule 5.2(a))*

(88) Date of publication of the international search report:
3 September 2009

***Brassica* plant comprising a mutant *INDEHISCENT* allele**

FIELD OF THE INVENTION

This invention relates to the field of agricultural products, especially crop plants, particularly of the *Brassicaceae* family, in particular *Brassica* species, of which the fruit dehiscence properties are modulated. More specifically the invention relates to improved methods and means for reducing seed shattering, or delaying seed shattering until after harvest, in plants such as *Brassicaceae* plants, particularly *Brassicaceae* plants grown for seed production, while maintaining at the same time an agronomically relevant treshability of the pods. Provided are both wild type and mutant nucleic acid molecules encoding *Brassica* *INDEHISCENT* proteins (*IND*) and the proteins as such. Also provided are *Brassica* plants comprising at least two *IND* genes, in particular *Brassica napus* plants, and cells, parts, seeds and progeny thereof, characterized in that they comprise three full knock-out mutant *ind* alleles in their genome, whereby the fruit dehiscence properties are significantly altered. In addition, methods for generating *Brassica* plants in which seed shattering is reduced, or in which seed shattering is delayed until after harvest, while an agronomically relevant treshability of the pods is preferably maintained, are provided herein, as are seed pods and seeds obtainable from such plants. Further provided are detection tools (kits) and methods for detecting the presence of one or more mutant *ind* and/or wild type *IND* alleles in biological samples, as well as methods for transferring one or more mutant *ind* and/or wild type *IND* alleles to other plants and methods for combining different *ind* and/or *IND* alleles in plants. In particular, methods for combining a suitable number of mutant *ind* alleles, which encode non-functional or no *IND* proteins and/or *IND* proteins having significantly reduced activity *in vivo*, in such a way as to significantly reduce seed shattering, or to delay seed shattering until after harvest, while maintaining at the same time an agronomically relevant treshability of the pods. In addition uses of the plants, or parts thereof, and/or progeny thereof, seeds and seed oils and the methods and/or kits of the invention are provided. Also provided are methods and means to increase the yield, particularly grain and seed yield. The yield increase phenotype may be separate from the reduced or delayed seed shatter phenotype.

30

BACKGROUND OF THE INVENTION

Siliques or pods from *Brassica* plants release their seeds through a process called fruit dehiscence. A silique consists of two carpels joined margin to margin. The suture between the margins forms a thick rib, called replum. As pod maturity approaches, the two valves separate progressively from the replum, along designated lines of weakness in the pod, eventually

35

resulting in the shattering of the seeds that were attached to the replum. The dehiscence zone defines the exact location of the valve dissociation.

5 Shedding of seed (also referred to as “seed shatter” or “pod shatter”) by mature pods before or during crop harvest is a universal phenomenon with crops that develop dry dehiscent fruits. Premature seed shatter results in a reduced seed recovery, which represents a problem in crops that are grown primarily for the seeds, such as oil-producing *Brassica* plants, particularly oilseed rape. Another problem related to premature seed shattering is an increase in volunteer growth in the subsequent crop year. In oilseed rape, pod shatter-related yield losses are on average 20%
10 (Child *et al.*, 1998, *J Exp Bot* 49: 829-838), but can reach up to 50%, depending on the weather conditions (MacLeod, 1981, *Harvesting in Oilseed Rape*, pp. 107-120 Cambridge Agricultural Publishing, Cambridge).

Current commercial oilseed rape varieties are extremely susceptible to shattering. There is little
15 variation for resistance to shattering within existing breeding programs of *B. napus* but resistant lines have been found within the diploid parents of *B. napus* (*B. oleracea* and *B. rapa*) as well as within other members of the *Brassica* genus, notably *B. juncea*, *B. carinata* and *B. nigra*. Kadkol *et al.* (1986, *Aust. J. Botany* 34 (5): 595-601) report increased resistance towards shattering in certain accessions of *B. campestris* that was associated with the absence of a separation layer in
20 the region of attachment of the siliqua valves to the replum. Prakash and Chopra (1988, *Plant breeding* 101: 167-168) describe the introgression of resistance to shattering in *Brassica napus* from *Brassica juncea* through non-homologous recombination. Spence *et al.* (1996, *J of Microscopy* 181: 195-203) describe that some lines of *Brassica juncea* show a reduced tendency to shatter as compared to *Brassica napus* lines. Morgan *et al.*, 1998 (*Fields Crop Research* 58,
25 153-165) describe genetic variation for pod shatter resistance among lines of oilseed rape developed from synthetic *B. napus* and conclude that lines which required much energy to open their pods appeared to have increased vascularisation in the dehiscence zone and to have reduced cell wall degradation within the dehiscence zone. They further found a significant negative correlation between the length of the pod beak and the force needed to cause pod shattering.
30 Child and Huttly (1999, *Proc 10th Int. Rapeseed Congress*) describe variation in pod maturation in an irradiation-induced mutant *B. napus* and a population of its parent cultivar, Jet Neuf, wherein the most resistant wild-type and mutant plants showed much lignification of groups of cells throughout the dehiscence zone and wherein vascular traces situated close to the inner edge of the dehiscence zone in the mutant were described to help to secure the valves. Child *et al.*
35 (2003, *J Exp Botany* 54 (389): 1919-1930) further describe the association between increased

pod shatter resistance and changes in the vascular structure in pods of a resynthesized *Brassica napus* line. However, the traditional methods for breeding have been unsuccessful in introducing shatter resistance into rape cultivars, without interference with other desirable traits such as early flowering, maturity and blackleg resistance (Prakash and Chopra, 1990, Genetical Research 56: 1-2).

Several genes, which promote or inhibit pod dehiscence, have been identified in *Arabidopsis thaliana* through mutant analysis: Combined mutants in both *SHATTERPROOF1* (*SHP1*; initially referred to as *AGL1*) and *SHATTERPROOF2* (*SHP2*; initially referred to as *AGL5*) result in indehiscent siliques (i.e. siliques which remain closed upon maturity in *Arabidopsis thaliana*) (Liljegren *et al.*, 2000, Nature 404, 766-770). Similarly, mutants in the *INDEHISCENT* gene (referred to as *IND1*) in *Arabidopsis thaliana* (Liljegren *et al.*, 2004, Cell 116: 843-853; PCT publication WO 01/79517), as well as in *ALCATRAZ* (referred to as *ALC*; Rajani *et al.* 2001, Current Biology 11, 1914-1922) interfered with pod dehiscence leading to pod shatter resistance. Constitutive expression of *FRUITFUL* (*FUL*), a repressor of *SHP* and *IND*, in *Arabidopsis thaliana* also resulted in indehiscent siliques (Ferrandiz *et al.*, 2000, Science, 289, 436-438). These transcription factors are believed to form a non-linear transcriptional network that controls valve margin identity and pod shatter. Liljegren *et al.* (2004, Cell 116: 843-853) further describe that *IND*, an atypical basic helix-loop-helix (bHLH) gene, directs the differentiation of the valve margin into the separation and lignified layers in *Arabidopsis thaliana*. The layer of lignified cells adjacent to the separation layer along with the endocarp *b* layer (a single lignified cell layer in each valve) produce a spring-like tension within the drying fruit that contributes to its opening. Lignification of the valve endocarp *b* layer requires the activities of *IND*, *SHP*, *ALC*, and *FUL*, a MADS-domain transcription factor that is expressed throughout the valves (Liljegren *et al.*, 2004, *supra*; Mandel and Yanofsky, 1995, Plant Cell 7, 1763-1771). *FUL* and *REPLUMLESS* (*RPL*), a homeodomain transcription factor that is expressed in the replum (Roeder *et al.*, 2003, Curr Biol 13, 1630-1635), have been found to set the boundaries of the genes that confer valve margin identity (Gu *et al.*, 1998, Development 125, 1509-1517; Ferrandiz *et al.*, 2000, Science, 289, 436-438; Roeder *et al.*, 2003, *supra*). Finally, *FILAMENTOUS FLOWER* (*FIL*) and *YABBY3* (*YAB3*), two *YABBY*-family transcription factors (Sawa *et al.*, 1999, Genes Dev 13, 1079-1088; Siegfried *et al.*, 1999, Development 126, 4117-4128), and *JAGGED* (*JAG*), a C2H2 zinc-finger transcription factor (Dinneny *et al.*, 2004, Development 131, 1101-1110; Ohno *et al.*, 2004, Development 131, 1111-1122), were identified to redundantly contribute to proper valve and valve margin development by promoting the expression of *FUL* and *SHP* in a region-specific manner (Dinneny *et al.*, 2005, Development

132, 4687-4696). Genes for a number of hydrolytic enzymes, such as endopolygalacturonases, which play a role, during pod dehiscence, in the programmed breakdown of the dehiscence zone in pods from *Brassica* plants have also been identified (see e.g. WO 97/13865; Petersen *et al.*, Plant. Mol. Biol., 1996, 31:517-527).

5

Liljegren *et al.* (2004, Cell 116: 843-853) describe five mutant alleles of *Arabidopsis* IND. The lignified cells in the dehiscence zone are either absent or present in plants comprising these mutant alleles depending on the severity of the mutations (severe *ind* mutants do not contain lignified cells in the region corresponding to the inner part of the valve margin in wild-type plants), but in all cases the silique is indehiscent. Wu *et al.* (2006), Planta 224, 971-979) describe a sixth mutant allele of *Arabidopsis* IND. Plants comprising this mutant allele show no lignified cells at the junctions of the valve margin and the replum, contain fewer cells in a region of seven layers of cells, which appeared to encompass the commonly known dehiscence zone and replum border in wild-type plants, and exhibit incomplete cytokinesis in this layer.

10

15

US 2005/0120417 and US 2007/0006336 describe the identification and isolation of two *IND1* orthologs from *Brassica napus*.

20

WO99/00503, WO01/79517 and WO0159122 describe downregulation of the expression of the *Arabidopsis* *ALC*, *IND*, *AGL1* and *AGL5* genes and orthologs thereof using gene-silencing techniques (such as antisense suppression or cosuppression) and mutagenesis.

25

Vancanneyt *et al.*, 2002 (XIII International Conference on *Arabidopsis* Research, Sevilla, Spain June 28-July 2; 2002) reported that the expression of *FUL* from *A. thaliana* under control of a CaMV 35S promoter in oilseed rape resulted in a number of pod shatter resistant transformants. Pods of such pod shatter resistant lines had no dehiscence zone, and opening of the pods could only be achieved by random fracture of the valves by applying considerable pressure.

30

Vancanneyt *et al.*, 2002 (XIII International Conference on *Arabidopsis* Research, Sevilla, Spain June 28-July 2; 2002) also reported that silencing of the *IND* gene in *Arabidopsis thaliana* using so-called dsRNA silencing techniques resulted in almost complete pod shatter resistance. Ninety-eight percent of the transgenic *Arabidopsis* lines developed siliques, which did not open along the valve suture, and could only be opened by applying considerable pressure to the valves.

It is important to realize that while seed shattering constitutes an important problem in oilseed rape culture, which may be solved by developing pod shatter resistant lines, ultimately, separation of the seeds from the pods is still required. In normal agricultural practice this is achieved by treshing of the pods by a combine harvester. Treshing of the pods by a combine harvester must be complete and must cause minimum damage to the seeds thus released. However, as pod strength increases, the more severe action required to tresh them causes an unacceptable level of damage to the seed. The pods of pod shatter resistant *Brassicaceae* plants should thus not be so strong that they cannot be treshed in a combine harvester (Bruce *et al.* 2001, J. Agric. Engng Res. 80, 343-350).

WO 2004/113542 describes that moderate dsRNA gene silencing of genes involved in the development of the dehiscence zone and valve margins of pods in *Brassicaceae* plants allows the isolation of transgenic lines with increased pod shatter resistance and reduced seed shattering, the pods of which however may still be opened along the dehiscence zone by applying limited physical forces.

Despite the fact that sequences of specific *IND* genes and mutant sequences thereof, particularly *Arabidopsis* and *Brassica napus* *IND* gene sequences and mutant *Arabidopsis* *IND* gene sequences, are available in the art, a need remains for further *IND* gene sequences, *e.g.* to enable a specifically desired modification of seed shattering in plants, such as *Brassica napus* plants. The isolation of mutant alleles corresponding to *ind* in economically important *Brassicaceae* plants, such as oilseed rape, is a laborious and time consuming task. Moreover, such isolation may be complicated by the amphidiploidy in oilseed rape and the consequent functional redundancy of the corresponding genes.

SUMMARY OF THE INVENTION

The inventors have found that the fruit dehiscence properties in *Brassica* plants can be controlled by controlling the number of *IND* genes/alleles that are “functionally expressed” in seed pods, *i.e.* that result in functional (biologically active) *IND* protein. By combining a number of full knock-out mutant *IND* alleles (“*ind* alleles”), while maintaining a minimal number of wild type *IND* alleles, resulting in a minimal level of functional *IND* protein, the dehiscence properties of

the seed pods can be modified, more specifically pod shatter resistance can be increased and seed shattering can be reduced, or seed shattering can be delayed until after harvest, while maintaining at the same time an agronomically relevant treshability of the pods, such that the pods may still be opened along the dehiscence zone by applying limited physical forces. It is
5 thought that a minimal number of wild type *IND* alleles is needed to still enable the separation of the seeds from the pods, in particular by treshing of the pods by a combine harvester, such that the treshing of the pods is complete and causes minimum damage to the seeds thus released.

Thus, in a first aspect, the present invention provides a *Brassica* plant comprising at least two
10 *IND* genes, in particular a *Brassica napus* plant (and parts thereof, such as seed pods and seeds), characterized in that it comprises three full knock-out mutant *IND* alleles in its genome, in particular of an *IND-A1* and/or an *IND-C1* gene, and wherein the pod shatter resistance of the plant is significantly increased compared to the pod shatter resistance of a plant not comprising mutant *IND* alleles, but wherein the plant preferably maintains an agronomically relevant
15 treshability of the pods.

In another aspect, the invention provides (isolated) nucleic acid sequences encoding wild type and/or mutant *IND* proteins, as well as fragments thereof, and methods of using these nucleic acid sequences to modify the fruit dehiscence properties of plants. Also provided are the proteins
20 themselves and their use.

The invention further relates to a plant, and cells, parts, seeds and progeny thereof, comprising at least one full knock-out mutant *IND* allele, and thus a reduced amount of functional *IND* protein compared to a plant, and cells, parts, seeds and progeny thereof, comprising an *IND* allele
25 encoding the corresponding functional *IND* protein. Such plants, and cells, parts, seeds and progeny thereof, can be used for obtaining plants with modified fruit dehiscence properties, in particular for obtaining *Brassica* plants with significantly reduced seed shattering that maintain an agronomically relevant treshability of the pods. As used herein, "plant part" includes anything derived from a plant of the invention, including plant parts such as cells, tissues, organs, seeds,
30 seed pods, seed meal, seed cake, seed fats or oils.

In a further aspect, the invention relates to seed pods with modified shatter resistance, which can be obtained from a plant according to the present invention, and the use of said seed pods, for example for planting and growing progeny from the plants.

In yet another aspect of the invention, methods are provided for generating and selecting plants, and cells, parts, seeds and progeny thereof, containing at least one full knock-out *ind* allele. In particular, methods are provided for generating and selecting *Brassica* plants comprising at least two *IND* genes, in particular *Brassica napus* plants, and cells, parts, seeds and progeny thereof, containing at least one full knock-out mutant *ind* allele present at at least one of the at least two different *IND* loci in the genome, for example at at least one of the two different loci of the *Brassica* IND-A1 and IND-C1 gene, and to distinguish between the presence of mutant *ind* alleles and wild type *IND* alleles in a plant or plant part. Thus methods are provided (such as mutagenesis and/or marker assisted selection) for generating and/or identifying *ind* alleles or plants or plant parts comprising such alleles and for combining a suitable number of *ind* alleles and/or different types of *ind* alleles in a single plant, whereby the fruit dehiscence properties of this plant are significantly modified.

In another embodiment of the invention, the mutant *IND* alleles of the invention are used to increase the yield of harvested seed or grain from *Brassica* plants. The increased yield may be a consequence of reducing or delaying seed shattering, but may also be independent from the reduced or delayed seed shatter. In particular, *Brassica* plants are provided comprising at least two *IND* genes, or a cell, part, seed or progeny thereof, characterized in that these plants comprise two mutant homozygous *IND* alleles as herein described in their genome.

20

FIGURE LEGENDS

Figure 1 – Schematical representation of the *IND-A1* gene, encoding a wild-type IND-A1 protein from *Brassica napus* (SEQ ID NO: 5).

Figure 2 – Schematical representation of the *IND-C1* gene, encoding a wild-type IND-C1 protein from *Brassica napus* (SEQ ID NO: 7).

In Figure 1 and 2 the position of the mutations described in the Examples (named “EMSxx” according to their respective “*ind-x1*-EMSxx” name as described in the Examples) is indicated with vertical lines; the length and position of the *IND* specific probes (named “ID xx” according to their respective SEQ ID NO: xx) are indicated by horizontal lines below the schematical representation of the *IND* genes; the position of the *IND* specific primers (named “ID xx” according to their respective SEQ ID NO: xx) are indicated by arrowheads.

GENERAL DEFINITIONS

“Increase of pod shatter resistance” and “reduction of seed shattering”, as used herein, refers to a decreased seed shatter tendency and/or a delay in the timing of seed shattering, in particular until

35

after harvest, of *Brassica* plants, the fruits of which normally do not mature synchronously, but sequentially, so that some pods burst open and shatter their seeds before or during harvest. The level of resistance to pod shattering is positively correlated with and can, for example, be measured by determining the force needed to break pods in the 'tensile separation test' (Davies and Bruce, 1997, *J Mat Sci* 32: 5895-5899; Morgan *et al.*, 1998, *Fields Crop Research* 58, 153-165), the number of intact pods remaining after e.g. 20 sec ('IP20'; Morgan *et al.*, 1998, *supra*), 9.7 or 17 sec (Bruce *et al.*, 2002, *Biosystems Eng* 81(2): 179-184) in a 'random impact test', the pod sample half-life ('LD50') in a random impact test, i.e. the treatment time needed to cause the opening of 50% of the pods in tested pod samples, and the 'field score for shattering' (Morgan *et al.*, 1998, *supra*). Random impact tests (RITs) and algorithms to define the pod sample half-lives in such RITs have been described in Bruce *et al.*, 2002 (*supra*), Morgan *et al.*, 1998 (*supra*) and the Examples below. Both publications are hereby incorporated by reference. Briefly, a sample of intact mature pods is placed in a closed drum together with steel balls and the drum is then vigorously agitated for increasing periods of times (e.g. 10 s, 20 s, 40 s, 80 s). After each period, the drum is opened and the number of broken and damaged pods is counted. The most accurate estimation of the level of shattering resistance for each line is calculated by fitting a linear x linear curve to all the available data and estimating the time taken for half of the pods within a sample to be broken ("pod sample half-life" or "LD50"). It is important however that pods open mainly along the dehiscence zone, and are not simply pulverized, as may occur with indehiscent pods.

An "agronomically relevant increase of pod shatter resistance", as used herein, refers to an increase of pod shatter resistance in a plant which results in pod shatter-related yield losses in the field (pre-harvest) below those normally observed for that plant in the field. For oilseed rape, pod shatter-related yield losses in the field are reported to be about 11% for a season with on average good growth conditions and about 25% for a season with on average bad growth conditions. A positive correlation has been found between these levels of seed loss and the level of seed loss at 9.7 s and 17 s treatment time, respectively, in the random impact test as described by Bruce *et al.*, 2002 (*Biosystems Eng* 81(2): 179-184). Alternatively, to determine whether the level of resistance to pod shattering in a plant is agronomically relevant, the pod sample half-life ('LD50', see above) of the plant can be compared with the pod sample half-life of a plant known to have an average level of pod shatter resistance, such as, for oilseed rape, all currently commercially available oilseed rape varieties.

As used herein, “pod or seed shattering” or “fruit or pod dehiscence” refers to a process that takes place in a fruit after seed maturation, whereby the valves detach from the central septum freeing the seeds. The region that breaks (i.e. the “dehiscence zone”) runs the entire length of the fruit between the valves and the replum (external septum). At maturity, the “dehiscence zone” is essentially a non-lignified layer of cells between a region of lignified cells in the valve and the replum. Shattering occurs due to the combination of cell wall loosening in the dehiscence zone and the tensions established by the differential mechanical properties of the drying cells in the silique.

- 10 A *Brassica* “fruit”, as used herein, refers to an organ of a *Brassica* plant that develops from a gynoeceium composed of fused carpels, which, upon fertilization, grows to become a “(seed) pod” or “silique” that contains the developing seeds. A *Brassica* “(seed) pod” or “silique” consists of a fruit wall (carpel) enclosing two locules separated by the septum. The “dehiscence zones” develop at the carpel margins adjacent to the septum and run the length of the silique.
- 15 The cells of the dehiscence zone eventually begin to degrade and this weakens the contact between the carpel walls or valves and the septum. The loss of cellular cohesion is confined to the cells of the dehiscence zone and results from middle lamella breakdown (Meakin and Roberts, 1990, J Exp Bot 41, 995-1011).
- 20 “Dehiscence zones”, as used herein, refers to layers of simple, parenchymatous cells, contained in the sutures situated on both sides of the bi-valved pod of plants, in particular *Brassica* plants. The dehiscence zones are situated between the pod valve edge and a central replum that contains the main vascular bundle to the stalk or pedicel. Dissociation of the cells in the dehiscence zone takes place during pod senescence and is complete by the time the pods reach full maturity
- 25 (Meakin and Roberts, 1990, *supra*). Valve separation can then take place. The dehiscence zone contains vascular traces, which pass from the pod wall to the pedicel (stalk) and the replum. The process of pod shatter takes place only after external force fractures the delicate vascular threads, allowing the valves to separate and the seeds to fall to the ground. This occurs during disturbance of the canopy, for example by contact with the combine during harvesting. The vascular tissue
- 30 contains thickened, lignified cells, which form the collenchymatous groups of cells found adjacent to the conductive cells (Meakin and Roberts, 1990, *supra*). This provides rigidity to the tissue and presumably, some resistance to fracturing.

As used herein, “an agronomically relevant treshability” refers to the resistance of a pod, particularly an oilseed rape pod, to opening along the dehiscence zone of the pod with concurrent

release of the seeds, upon application of physical forces that allow complete opening of the pods while preventing damage to the seeds, as they are used e.g. in a combine harvester. A positive correlation has been found between a pod sample half-life ('LD50') in a random impact test and their treshability. Oilseed rape pod sample half-lives, as determined in a RIT performed as described in the Examples, which correspond to agronomically relevant treshability should not exceed 80 seconds. Typical sample half-life values for control lines of commercially available oilseed rape varieties are about 10 seconds. Thus, lines with significantly increased pod shatter resistance with agronomically relevant treshability have a pod sample half-life in RIT between about 10 and about 80 seconds, between about 10 and about 60 seconds, between about 10 and about 50 seconds, between about 20 and about 60 seconds, between about 20 and about 50 seconds, between about 40 and about 60 seconds, of about 57 seconds.

"Crop plant" refers to plant species cultivated as a crop, such as *Brassica napus* (AACC, 2n=38), *Brassica juncea* (AABB, 2n=36), *Brassica carinata* (BBCC, 2n=34), *Brassica rapa* (syn. *B. campestris*) (AA, 2n=20), *Brassica oleracea* (CC, 2n=18) or *Brassica nigra* (BB, 2n=16). The definition does not encompass weeds, such as *Arabidopsis thaliana*.

The term "nucleic acid sequence" (or nucleic acid molecule) refers to a DNA or RNA molecule in single or double stranded form, particularly a DNA encoding a protein or protein fragment according to the invention. An "endogenous nucleic acid sequence" refers to a nucleic acid sequence within a plant cell, e.g. an endogenous allele of an *IND* gene present within the nuclear genome of a *Brassica* cell. An "isolated nucleic acid sequence" is used to refer to a nucleic acid sequence that is no longer in its natural environment, for example *in vitro* or in a recombinant bacterial or plant host cell.

The term "gene" means a DNA sequence comprising a region (transcribed region), which is transcribed into an RNA molecule (e.g. into a pre-mRNA, comprising intron sequences, which is then spliced into a mature mRNA, or directly into a mRNA without intron sequences) in a cell, operable linked to regulatory regions (e.g. a promoter). A gene may thus comprise several operably linked sequences, such as a promoter, a 5' leader sequence comprising e.g. sequences involved in translation initiation, a (protein) coding region (cDNA or genomic DNA) and a 3' non-translated sequence comprising e.g. transcription termination sites. "Endogenous gene" is used to differentiate from a "foreign gene", "transgene" or "chimeric gene", and refers to a gene from a plant of a certain plant genus, species or variety, which has not been introduced into that plant by transformation (i.e. it is not a "transgene"), but which is normally present in plants of

that genus, species or variety, or which is introduced in that plant from plants of another plant genus, species or variety, in which it is normally present, by normal breeding techniques or by somatic hybridization, e.g., by protoplast fusion. Similarly, an “endogenous allele” of a gene is not introduced into a plant or plant tissue by plant transformation, but is, for example, generated
5 by plant mutagenesis and/or selection or obtained by screening natural populations of plants.

“Expression of a gene” or “gene expression” refers to the process wherein a DNA region, which is operably linked to appropriate regulatory regions, particularly a promoter, is transcribed into an RNA molecule. The RNA molecule is then processed further (by post-transcriptional
10 processes) within the cell, e.g. by RNA splicing and translation initiation and translation into an amino acid chain (polypeptide), and translation termination by translation stop codons. The term “functionally expressed” is used herein to indicate that a functional protein is produced; the term “not functionally expressed” to indicate that a protein with significantly reduced or no functionality (biological activity) is produced or that no protein is produced (see further below).

15

The terms “protein” or “polypeptide” are used interchangeably and refer to molecules consisting of a chain of amino acids, without reference to a specific mode of action, size, 3-dimensional structure or origin. A “fragment” or “portion” of an IND protein may thus still be referred to as a “protein”. An “isolated protein” is used to refer to a protein that is no longer in its natural
20 environment, for example *in vitro* or in a recombinant bacterial or plant host cell. The term “transcription factor” is used to refer to a protein consisting of at least two discrete domains – a DNA binding domain and an activation or repression domain - that operate together to modulate the rate of transcriptional initiation from target gene promoters (Ptashne, 1988, Nature 335, 683-689). The term “basic helix-loop-helix (bHLH) domain transcription factor” is used to refer to a
25 transcription factor comprising, apart from the bHLH DNA binding domain (Heim *et al.*, 2003, Mol Biol Evol 20, 735-747; Toledo-Ortiz *et al.*, 2003, Plant Cell 15, 1749-1770), domains which are known to be important for the regulation of gene expression which may be conserved at the amino acid level in related proteins from different species (Quong *et al.*, 1993, Mol Cell Biol 13, 792-800). Transcriptional regulators comprising a bHLH domain bind DNA through residues in
30 the basic region while the helix-loop-helix domain promotes dimerization, allowing family members to form hetero- or homodimers (Murre *et al.*, 1989, Cell 56, 777-783).

The term “IND gene” refers herein to a nucleic acid sequence encoding an INDEHISCENT (IND) protein, which is a bHLH domain transcription factor required for seed dispersal
35 (Liljegren *et al.*, 2004, Cell 116: 843-853).

As used herein, the term “allele(s)” means any of one or more alternative forms of a gene at a particular locus. In a diploid (or amphidiploid) cell of an organism, alleles of a given gene are located at a specific location or locus (loci plural) on a chromosome. One allele is present on
5 each chromosome of the pair of homologous chromosomes.

As used herein, the term “homologous chromosomes” means chromosomes that contain information for the same biological features and contain the same genes at the same loci but possibly different alleles of those genes. Homologous chromosomes are chromosomes that pair
10 during meiosis. “Non-homologous chromosomes”, representing all the biological features of an organism, form a set, and the number of sets in a cell is called ploidy. Diploid organisms contain two sets of non-homologous chromosomes, wherein each homologous chromosome is inherited from a different parent. In amphidiploid species, essentially two sets of diploid genomes exist, whereby the chromosomes of the two genomes are referred to as “homeologous chromosomes”
15 (and similarly, the loci or genes of the two genomes are referred to as homeologous loci or genes). A diploid, or amphidiploid, plant species may comprise a large number of different alleles at a particular locus.

As used herein, the term “heterozygous” means a genetic condition existing when two different
20 alleles reside at a specific locus, but are positioned individually on corresponding pairs of homologous chromosomes in the cell. Conversely, as used herein, the term “homozygous” means a genetic condition existing when two identical alleles reside at a specific locus, but are positioned individually on corresponding pairs of homologous chromosomes in the cell.

As used herein, the term “locus” (loci plural) means a specific place or places or a site on a
25 chromosome where for example a gene or genetic marker is found. For example, the “*IND-A1* locus” refers to the position on a chromosome of the A genome where the *IND-A1* gene (and two *IND-A1* alleles) may be found, while the “*IND-C1* locus” refers to the position on a chromosome of the C genome where the *IND-C1* gene (and two *IND-C1* alleles) may be found.

30 Whenever reference to a “plant” or “plants” according to the invention is made, it is understood that also plant parts (cells, tissues or organs, seed pods, seeds, severed parts such as roots, leaves, flowers, pollen, etc.), progeny of the plants which retain the distinguishing characteristics of the parents (especially the fruit dehiscence properties), such as seed obtained by selfing or crossing,

e.g. hybrid seed (obtained by crossing two inbred parental lines), hybrid plants and plant parts derived there from are encompassed herein, unless otherwise indicated.

5 A “molecular assay” (or test) refers herein to an assay that indicates (directly or indirectly) the presence or absence of one or more particular *IND* alleles at one or both *IND* loci (e.g. at one or both of the *IND-A1* or *IND-C1* loci). In one embodiment it allows one to determine whether a particular (wild type or mutant) allele is homozygous or heterozygous at the locus in any individual plant.

10 “Wild type” (also written “wildtype” or “wild-type”), as used herein, refers to a typical form of a plant or a gene as it most commonly occurs in nature. A “wild type plant” refers to a plant with the most common phenotype of such plant in the natural population. A “wild type allele” refers to an allele of a gene required to produce the wild-type phenotype. By contrast, a “mutant plant” refers to a plant with a different rare phenotype of such plant in the natural population or
15 produced by human intervention, e.g. by mutagenesis, and a “mutant allele” refers to an allele of a gene required to produce the mutant phenotype.

As used herein, the term “wild type *IND*” (e.g. wild type *IND-A1* or *IND-C1*), means a naturally occurring *IND* allele found within plants, in particular *Brassicaceae* plants, especially *Brassica*
20 plants, which encodes a functional *IND* protein (e.g. a functional *IND-A1* or *IND-C1*, respectively). In contrast, the term “mutant *IND*” (e.g. mutant *IND-A1* or *IND-C1*), as used herein, refers to an *IND* allele, which does not encode a functional *IND* protein, i.e. an *IND* allele encoding a non-functional *IND* protein (e.g. a non-functional *IND-A1* or *IND-C1*, respectively), which, as used herein, refers to an *IND* protein having no biological activity or a significantly
25 reduced biological activity as compared to the corresponding wild-type functional *IND* protein, or encoding no *IND* protein at all. Such a “mutant *IND* allele” (also called “full knock-out” or “null” allele) is a wild-type *IND* allele, which comprises one or more mutations in its nucleic acid sequence, whereby the mutation(s) preferably result in a significantly reduced (absolute or relative) amount of functional *IND* protein in the cell *in vivo*. As used herein, a “full knock-out
30 *IND* allele” is a mutant *IND* allele the presence of which in homozygous state at each *IND* locus in the plant (e.g. a *Brassica napus* plant with two full knock-out *IND-A1* alleles and two full knock-out *IND-C1* alleles) results in an increase of pod shatter resistance in that plant which is too high to be still agronomically relevant. Mutant alleles of the *IND* protein-encoding nucleic acid sequences are designated as “*ind*” (e.g. *ind-a1* or *ind-c1*, respectively) herein. Mutant alleles
35 can be either “natural mutant” alleles, which are mutant alleles found in nature (e.g. produced

spontaneously without human application of mutagens) or “induced mutant” alleles, which are induced by human intervention, e.g. by mutagenesis.

A “significantly reduced amount of functional IND protein” (e.g. functional IND-A1 or IND-C1 protein) refers to a reduction in the amount of a functional IND protein produced by the cell comprising a mutant *IND* allele by at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 100% (i.e. no functional IND protein is produced by the cell) as compared to the amount of the functional IND protein produced by the cell not comprising the mutant *IND* allele. This definition encompasses the production of a “non-functional” IND protein (e.g. truncated IND protein) having no biological activity *in vivo*, the reduction in the absolute amount of the functional IND protein (e.g. no functional IND protein being made due to the mutation in the *IND* gene), and/or the production of an IND protein with significantly reduced biological activity compared to the activity of a functional wild type IND protein (such as an IND protein in which one or more amino acid residues that are crucial for the biological activity of the encoded IND protein, as exemplified below, are substituted for another amino acid residue). The term “mutant IND protein”, as used herein, refers to an IND protein encoded by a mutant *IND* nucleic acid sequence (“*ind* allele”) whereby the mutation results in a significantly reduced and/or no IND activity *in vivo*, compared to the activity of the IND protein encoded by a non-mutant, wild type *IND* sequence (“*IND* allele”).

“Mutagenesis”, as used herein, refers to the process in which plant cells (e.g., a plurality of *Brassica* seeds or other parts, such as pollen, etc.) are subjected to a technique which induces mutations in the DNA of the cells, such as contact with a mutagenic agent, such as a chemical substance (such as ethylmethylsulfonate (EMS), ethylnitrosourea (ENU), etc.) or ionizing radiation (neutrons (such as in fast neutron mutagenesis, etc.), alpha rays, gamma rays (such as that supplied by a Cobalt 60 source), X-rays, UV-radiation, etc.), or a combination of two or more of these. Thus, the desired mutagenesis of one or more *IND* alleles may be accomplished by use of chemical means such as by contact of one or more plant tissues with ethylmethylsulfonate (EMS), ethylnitrosourea, etc., by the use of physical means such as x-ray, etc, or by gamma radiation, such as that supplied by a Cobalt 60 source. While mutations created by irradiation are often large deletions or other gross lesions such as translocations or complex rearrangements, mutations created by chemical mutagens are often more discrete lesions such as point mutations. For example, EMS alkylates guanine bases, which results in base mispairing: an alkylated guanine will pair with a thymine base, resulting primarily in G/C to A/T transitions. Following mutagenesis, *Brassica* plants are regenerated from the treated cells using known

techniques. For instance, the resulting *Brassica* seeds may be planted in accordance with conventional growing procedures and following self-pollination seed is formed on the plants. Alternatively, doubled haploid plantlets may be extracted to immediately form homozygous plants, for example as described by Coventry *et al.* (1988, Manual for Microspore Culture Technique for *Brassica napus*. Dep. Crop Sci. Techn. Bull. OAC Publication 0489. Univ. of Guelph, Guelph, Ontario, Canada). Additional seed that is formed as a result of such self-pollination in the present or a subsequent generation may be harvested and screened for the presence of mutant *IND* alleles. Several techniques are known to screen for specific mutant alleles, e.g., DeleteageneTM (Delete-a-gene; Li *et al.*, 2001, Plant J 27: 235-242) uses polymerase chain reaction (PCR) assays to screen for deletion mutants generated by fast neutron mutagenesis, TILLING (targeted induced local lesions in genomes; McCallum *et al.*, 2000, Nat Biotechnol 18:455-457) identifies EMS-induced point mutations, etc. Additional techniques to screen for the presence of specific mutant *IND* alleles are described in the Examples below.

As used herein, the term "non-naturally occurring" when used in reference to a plant, means a plant with a genome that has been modified by man. A transgenic plant, for example, is a non-naturally occurring plant that contains an exogenous nucleic acid molecule, e.g., a chimeric gene comprising a transcribed region which when transcribed yields a biologically active RNA molecule capable of reducing the expression of an endogenous gene, such as an *IND* gene according to the invention, and, therefore, has been genetically modified by man. In addition, a plant that contains a mutation in an endogenous gene, for example, a mutation in an endogenous *IND* gene, (e.g. in a regulatory element or in the coding sequence) as a result of an exposure to a mutagenic agent is also considered a non-naturally plant, since it has been genetically modified by man. Furthermore, a plant of a particular species, such as *Brassica napus*, that contains a mutation in an endogenous gene, for example, in an endogenous *IND* gene, that in nature does not occur in that particular plant species, as a result of, for example, directed breeding processes, such as marker-assisted breeding and selection or introgression, with a plant of the same or another species, such as *Brassica juncea* or *rapa*, of that plant is also considered a non-naturally occurring plant. In contrast, a plant containing only spontaneous or naturally occurring mutations, i.e. a plant that has not been genetically modified by man, is not a "non-naturally occurring plant" as defined herein and, therefore, is not encompassed within the invention. One skilled in the art understands that, while a non-naturally occurring plant typically has a nucleotide sequence that is altered as compared to a naturally occurring plant, a non-naturally occurring plant also can be genetically modified by man without altering its nucleotide sequence, for example, by modifying its methylation pattern.

The term “ortholog” of a gene or protein refers herein to the homologous gene or protein found in another species, which has the same function as the gene or protein, but is (usually) diverged in sequence from the time point on when the species harboring the genes diverged (i.e. the genes evolved from a common ancestor by speciation). Orthologs of the *Brassica napus* *IND* genes may thus be identified in other plant species (e.g. *Brassica juncea*, etc.) based on both sequence comparisons (e.g. based on percentages sequence identity over the entire sequence or over specific domains) and/or functional analysis.

10 A “variety” is used herein in conformity with the UPOV convention and refers to a plant grouping within a single botanical taxon of the lowest known rank, which grouping can be defined by the expression of the characteristics resulting from a given genotype or combination of genotypes, can be distinguished from any other plant grouping by the expression of at least one of the said characteristics and is considered as a unit with regard to its suitability for being propagated unchanged (stable).

The term “comprising” is to be interpreted as specifying the presence of the stated parts, steps or components, but does not exclude the presence of one or more additional parts, steps or components. A plant comprising a certain trait may thus comprise additional traits.

20

It is understood that when referring to a word in the singular (e.g. plant or root), the plural is also included herein (e.g. a plurality of plants, a plurality of roots). Thus, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there be one and only one of the elements. The indefinite article "a" or "an" thus usually means "at least one".

25

For the purpose of this invention, the "sequence identity" of two related nucleotide or amino acid sequences, expressed as a percentage, refers to the number of positions in the two optimally aligned sequences which have identical residues (x100) divided by the number of positions compared. A gap, i.e., a position in an alignment where a residue is present in one sequence but not in the other, is regarded as a position with non-identical residues. The “optimal alignment” of two sequences is found by aligning the two sequences over the entire length according to the Needleman and Wunsch global alignment algorithm (Needleman and Wunsch, 1970, J Mol Biol 48(3):443-53) in The European Molecular Biology Open Software Suite (EMBOSS, Rice *et al.*, 2000, Trends in Genetics 16(6): 276—277; see e.g.

35

<http://www.ebi.ac.uk/emboss/align/index.html>) using default settings (gap opening penalty = 10 (for nucleotides) / 10 (for proteins) and gap extension penalty = 0.5 (for nucleotides) / 0.5 (for proteins)). For nucleotides the default scoring matrix used is EDNAFULL and for proteins the default scoring matrix is EBLOSUM62.

5

"Substantially identical" or "essentially similar", as used herein, refers to sequences, which, when optimally aligned as defined above, share at least a certain minimal percentage of sequence identity (as defined further below).

10 "Stringent hybridization conditions" can be used to identify nucleotide sequences, which are substantially identical to a given nucleotide sequence. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequences at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH)
15 at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically stringent conditions will be chosen in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least 60°C. Lowering the salt concentration and/or increasing the temperature increases stringency. Stringent conditions for RNA-DNA hybridizations (Northern blots using a probe of e.g. 100nt) are for example those which include at least one wash in 0.2X SSC at 63°C
20 for 20min, or equivalent conditions.

"High stringency conditions" can be provided, for example, by hybridization at 65°C in an aqueous solution containing 6x SSC (20x SSC contains 3.0 M NaCl, 0.3 M Na-citrate, pH 7.0), 5x Denhardt's (100X Denhardt's contains 2% Ficoll, 2% Polyvinyl pyrrolidone, 2% Bovine
25 Serum Albumin), 0.5% sodium dodecyl sulphate (SDS), and 20 µg/ml denaturated carrier DNA (single-stranded fish sperm DNA, with an average length of 120 - 3000 nucleotides) as non-specific competitor. Following hybridization, high stringency washing may be done in several steps, with a final wash (about 30 min) at the hybridization temperature in 0.2-0.1× SSC, 0.1% SDS.

30

"Moderate stringency conditions" refers to conditions equivalent to hybridization in the above described solution but at about 60-62°C. Moderate stringency washing may be done at the hybridization temperature in 1x SSC, 0.1% SDS.

“Low stringency” refers to conditions equivalent to hybridization in the above described solution at about 50-52°C. Low stringency washing may be done at the hybridization temperature in 2x SSC, 0.1% SDS. See also Sambrook *et al.* (1989) and Sambrook and Russell (2001).

5 “Increased harvested yield” or “increased seed or grain yield” refers to the larger amount of seed or grain harvested from a plurality of plants, each comprising mutant *IND* alleles according to the invention, when compared to the amount of seed or grain harvested from a similar number of isogenic plants without the mutant *IND* alleles. Yield is typically expressed in volume units of harvested seed per surface units, such as bushels/acre or Kg/ha. The yield increase is typically expressed in percentage, whereby the yield of the reference or control plant is referred to as 100% and the yield of the plants according to the inventions is expressed in % relative to the yield of the control plant. Observed yield increases in Brassica plants according to the invention ranged from at least 101% to at least 124% and it is expected that higher yield increases are feasible. Yield increase may also range from 104% to 108% or 105% to 110%.

15

DETAILED DESCRIPTION

Brassica napus (genome AACC, $2n=4x=38$), which is an allotetraploid (amphidiploid) species containing essentially two diploid genomes (the A and the C genome) due to its origin from diploid ancestors, comprises two *IND* genes in its genome. It was found by the inventors that one *IND* gene is located on the A genome (herein referred to as “*IND-A1*”) and one on the C genome (herein referred to as “*IND-C1*”). The *IND-A1* gene is said to be “homeologous” to the *IND-C1* gene, i.e. the “A gene” is found on the A genome and originates from the diploid ancestor *B. rapa* (AA), while the “C gene” is found on the C genome of *B. napus* and originates from the diploid ancestor *B. oleracea* (CC).

25

As in any diploid genome, two “alleles” can be present *in vivo* for each *IND* gene at each *IND* locus in the genome (one allele being the gene sequence found on one chromosome and the other on the homologous chromosome). The nucleotide sequence of these two alleles may be identical (homozygous plant) or different (heterozygous plant) in any given plant, although the number of different possible alleles existing for each *IND* gene may be much larger than two in the species population as a whole.

It was moreover found that *Brassica napus* plants, which are homozygous for a full knockout *ind* allele in only one of the two *IND* genes, i.e. in *IND-A1* or *IND-C1*, do not show a significant increase in pod shatter resistance compared to *Brassica napus* plants not comprising mutant *IND*

35

alleles, while in *Brassica napus* plants, which are homozygous for a full knockout *ind* allele in both *IND* genes, pod shatter resistance is significantly increased, but the level of pod shatter resistance is too high to maintain an agronomically relevant treshability. By contrast, pod shatter resistance is significantly increased in *Brassica napus* plants comprising three full knockout *ind* alleles of the two *Brassica napus* *IND* genes, to a level whereby the plants maintain an agronomically relevant treshability of the pods. It is thought that the presence of three full knockout *ind* alleles in a *Brassica* plant comprising at least two *IND* genes, in particular in a *Brassica napus* plant comprising an *IND-A1* and an *IND-C1* gene, may be required in order to obtain a plant, which shows an increased pod shatter resistance, while maintaining an agronomically relevant treshability of the pods.

Thus in one embodiment of the invention, a *Brassica* plant comprising at least two *IND* genes, in particular a *Brassica napus* plant comprising an *IND-A1* and an *IND-C1* gene, comprising 3 *ind* alleles is provided herein, whereby the *ind* alleles result in a significantly reduced amount of functional *IND* protein of the type encoded by the wild-type equivalent of these mutant alleles and thus an overall significantly reduced amount of the functional *IND* proteins produced in the plant cells, specifically in the developing seed pods, *in vivo*.

It is further thought that by combining sufficient copies of specific (mutant) *ind* alleles with sufficient copies of specific (wild type) *IND* alleles in one plant, in particular a *Brassica* plant, it is possible to fine tune the amount and/or type of functional *IND* proteins made, which in turn influences the fruit dehiscence properties of the plant. The absolute and relative amount of the *IND* proteins can thus be tuned in such a way as to provide plants that produce sufficient *IND* protein(s) to enable an agronomically relevant treshability of the seed pods, while reducing seed shattering before or during harvest.

Thus in one embodiment of the invention, a plant, in particular a *Brassica* plant, is provided comprising at least one functionally expressed *IND* allele, which encodes a fully functional *IND* protein, while the remaining alleles may be (mutant) *ind* alleles.

In one aspect of the invention a *Brassica* plant comprising at least two *IND* genes, in particular a *Brassica napus* plant, comprising n-tuple *ind* alleles of at least 2 different *IND* genes in that *Brassica* plant, in particular of the *IND-A1* and *IND-C1* genes, is provided, whereby $n \leq 3$ (e.g. $n = 1, 2, \text{ or } 3$), so that at least one allele produces a functional *IND* protein.

In a further aspect of the invention an homozygous *IND* single mutant- (n=2, i.e. homozygous for a mutant allele of one *IND* gene), and/or an homozygous *IND* double mutant- (n=4, i.e. homozygous for a mutant allele of two *IND* genes) plant of a *Brassica* species comprising at least two *IND* genes, in particular of *Brassica napus*, is provided, whereby the mutant alleles are mutant alleles of 2 different *IND* genes in that *Brassica* plant, in particular of the *IND-A1* and/or *IND-C1* genes. Such mutant plants may, according to this invention, be used for breeding purposes. Thus in one embodiment of the invention, an homozygous *IND* single mutant *Brassica napus* plant is provided herein, wherein the genotype of the plant can be described as *ind-a1/ind-a1*, *IND-C1/IND-C1*, or *IND-A1/IND-A1*, *ind-c1/ind-c1*. In another embodiment of the invention, an homozygous *IND* double mutant *Brassica napus* plant is provided herein, wherein the genotype of the plant can be described as *ind-a1/ind-a1*, *ind-c1/ind-c1*.

In a further aspect of the invention the homozygous *IND* single (n=2) mutant plant of the *Brassica* species comprising at least two *IND* genes, in particular of *Brassica napus*, comprises a further mutant *IND* allele, wherein the mutant plant is heterozygous for the additional mutant *IND* allele (i.e., n=3), and wherein the mutant allele is a mutant allele of the remaining wild-type *IND* gene in that *Brassica* plant, in particular of the *IND-A1* or *IND-C1* gene. Thus in a further embodiment of the invention, an homozygous *IND* single mutant *Brassica napus* plant comprising one further mutant *IND* allele is provided herein, wherein the genotype of the plant can be described as *ind-a1/ind-a1*, *IND-C1/ind-c1*, or *IND-A1/ind-a1*, *ind-c1/ind-c1*.

Further provided herein are nucleic acid sequences of wild type and mutant *IND* genes/alleles from *Brassica* species, as well as the wild type and mutant *IND* proteins. Also provided are methods of generating and combining mutant and wild type *IND* alleles in *Brassica* plants, as well as *Brassica* plants and plant parts comprising specific combinations of wild type and mutant *IND* alleles in their genome, whereby seed shattering is reduced in these plants. The use of these plants for transferring mutant *IND* alleles to other plants is also an embodiment of the invention, as are the plant products of any of the plants described. In addition kits and methods for marker assisted selection (MAS) for combining or detecting *IND* genes and/or alleles are provided. Each of the embodiments of the invention is described in detail herein below.

The *Brassica* plants described herein which exhibit reduced or delayed seed shattering have an increase in the yield of harvested seed. However, it was observed, unexpectedly, that the harvested seed yield from *Brassica* plants comprising only two mutant *IND* alleles in homozygous state, i.e. wherein the genotype of the plant can be described as *ind-a1/ind-a1*, *IND-*

C1/ IND-C1, or *IND-A1/ IND-A1*, *ind-c1/ind-c1* was also significantly increased, when compared to isogenic *Brassica* plants not comprising the mutant IND alleles, despite the absence of an observable reduced or delayed seed shatter phenotype in the *Brassica* plants comprising the mutant IND alleles. The invention thus also provides *Brassica* plants comprising at least two
5 *IND* genes, wherein at least two alleles produce a functional IND protein, which plants have a higher seed yield. It will be clear that the two mutant alleles at the IND-A locus or at the IND-C locus may be the same mutant allele or a different mutant allele.

Nucleic acid sequences according to the invention

10 Provided are both wild type *IND* nucleic acid sequences encoding functional IND proteins and mutant *ind* nucleic acid sequences (comprising one or more mutations, preferably mutations which result in no or a significantly reduced biological activity of the encoded IND protein or in no IND protein being produced) of *IND* genes from *Brassicaceae*, particularly from *Brassica* species, especially from *Brassica napus*, but also from other *Brassica* crop species. For example,
15 *Brassica* species comprising an A and/or a C genome may comprise different alleles of *IND-A* or *IND-C* genes, which can be identified and combined in a single plant according to the invention. In addition, mutagenesis methods can be used to generate mutations in wild type *IND* alleles, thereby generating mutant *ind* alleles for use according to the invention. Because specific *IND* alleles are preferably combined in a plant by crossing and selection, in one embodiment the *IND*
20 and/or *ind* nucleic acid sequences are provided within a plant (i.e. endogenously), e.g. a *Brassica* plant, preferably a *Brassica* plant which can be crossed with *Brassica napus* or which can be used to make a “synthetic” *Brassica napus* plant. Hybridization between different *Brassica* species is described in the art, e.g., as referred to in Snowdon (2007, Chromosome research 15: 85-95). Interspecific hybridization can, for example, be used to transfer genes from, e.g., the C
25 genome in *B. napus* (AACC) to the C genome in *B. carinata* (BBCC), or even from, e.g., the C genome in *B. napus* (AACC) to the B genome in *B. juncea* (AABB) (by the sporadic event of illegitimate recombination between their C and B genomes). “Resynthesized” or “synthetic” *Brassica napus* lines can be produced by crossing the original ancestors, *B. oleracea* (CC) and *B. rapa* (AA). Interspecific, and also intergeneric, incompatibility barriers can be successfully
30 overcome in crosses between *Brassica* crop species and their relatives, e.g., by embryo rescue techniques or protoplast fusion (see e.g. Snowdon, above).

However, isolated *IND* and *ind* nucleic acid sequences (e.g. isolated from the plant by cloning or made synthetically by DNA synthesis), as well as variants thereof and fragments of any of these
35 are also provided herein, as these can be used to determine which sequence is present

endogenously in a plant or plant part, whether the sequence encodes a functional, a non-functional or no protein (e.g. by expression in a recombinant host cell as described below) and for selection and transfer of specific alleles from one plant into another, in order to generate a plant having the desired combination of functional and mutant alleles.

5

Nucleic acid sequences of *IND-A1* and *IND-C1* have been isolated from *Brassica napus* as depicted in the sequence listing. The wild type *IND* sequences are depicted, while the mutant *ind* sequences of these sequences, and of sequences essentially similar to these, are described herein below and in the Examples, with reference to the wild type *IND* sequences. The genomic *IND* protein-encoding DNA from *Brassica napus* does not comprise any introns.

10

“IND-A1 nucleic acid sequences” or “IND-A1 variant nucleic acid sequences” according to the invention are nucleic acid sequences encoding an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 2 or nucleic acid sequences having at least 80%, at least 85%, at least 90%, at least 95%, 96%, 97%, 98%, 99% or 100% sequence identity with SEQ ID NO: 1 or SEQ ID NO: 5. These nucleic acid sequences may also be referred to as being “essentially similar” or “essentially identical” to the *IND* sequences provided in the sequence listing.

15

20

“IND-C1 nucleic acid sequences” or “IND-C1 variant nucleic acid sequences” according to the invention are nucleic acid sequences encoding an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 4 (*IND-C1-long*) or with SEQ ID NO: 4 from the amino acid at position 16 to the amino acid at position 210 (*IND-C1-short*) or nucleic acid sequences having at least 80%, at least 85%, at least 90%, at least 95%, 96%, 97%, 98%, 99% or 100% sequence identity with SEQ ID NO: 3 (*IND-C1-long*), with SEQ ID NO:3 from the nucleotide at position 46 to the nucleotide at position 633 (*IND-C1-short*) or with SEQ ID NO: 7. These nucleic acid sequences may also be referred to as being “essentially similar” or “essentially identical” to the *IND* sequences provided in the sequence listing.

25

30

Thus the invention provides both nucleic acid sequences encoding wild type, functional *IND-A1* and *IND-C1* proteins, including variants and fragments thereof (as defined further below), as well as mutant nucleic acid sequences of any of these, whereby the mutation in the nucleic acid sequence preferably results in one or more amino acids being inserted, deleted or substituted in comparison to the wild type *IND* protein. Preferably the mutation(s) in the nucleic acid sequence

35

result in one or more amino acid changes (i.e. in relation to the wild type amino acid sequence one or more amino acids are inserted, deleted and/or substituted) whereby the biological activity of the IND protein is significantly reduced or completely abolished. A significant reduction in or complete abolishment of the biological activity of the IND protein refers herein to a reduction in or abolishment of the DNA binding activity, the dimerization capacity and/or transcriptional regulating activity of the IND protein, such that the pod shatter resistance of a plant expressing the mutant IND protein is increased as compared to a plant expressing the corresponding wild type IND protein.

10 To determine the functionality of a specific *IND* allele/protein in plants, particularly in *Brassica* plants, the level of resistance to pod shattering in the plants can be determined by performing macroscopical, microscopical and histological assays on fruits and flowers of the plants comprising the specific *IND* allele/protein and of corresponding wild type plants analogous to the assays performed on *Arabidopsis* fruits and flowers as described by Liljegren *et al.* (2004, *supra*) or as described in the Examples below. Briefly, changes in pod shatter resistance can be evaluated and/or measured, e.g., by macroscopical tests, such as inspection of the seed pods with naked eye to evaluate, e.g., the presence or absence of the valve margins, the length of the beak of the pods, etc.; a Manual Impact Test (MIT) to compare the level of pod shatter resistance between different mutant *IND* lines and corresponding wild type lines by evaluating the ease of pod opening upon gently twisting the pods; a Random Impact Test (RIT) to compare the treshability of seed pods from plants from different mutant *IND* lines and corresponding wild type lines, respectively, by measuring the half-life of pod samples of these lines; and/or by microscopic tests to examine, e.g., whether and how cells at the valve margin and the dehiscence zone of seed pods are affected by mutations in *IND*. Once the dimerization partner of the IND protein (e.g., the IND protein itself in case its functioning depends on the formation of an homodimer or another protein in case its functioning depends on the formation of an heterodimer) and/or the gene(s) the transcription of which is regulated by the IND protein are identified and characterized, the functionality of a specific *IND* allele/protein can alternatively be evaluated by recombinant DNA techniques as known in the art, e.g., by co-expressing both partners of the dimer in a host cell (e.g. a bacterium, such as *E. coli*) and evaluating if dimers can still be formed, if the dimers can still bind to the bHLH binding site of the regulated gene(s), and/or if the transcription of these gene(s) is still regulated by this binding.

Both endogenous and isolated nucleic acid sequences are provided herein. Also provided are fragments of the *IND* sequences and *IND* variant nucleic acid sequences defined above, for use

as primers or probes and as components of kits according to another aspect of the invention (see further below). A “fragment” of a *IND* or *ind* nucleic acid sequence or variant thereof (as defined) may be of various lengths, such as at least 10, 12, 15, 18, 20, 50, 100, 200, 500, 600 contiguous nucleotides of the *IND* or *ind* sequence (or of the variant sequence).

5

Nucleic acid sequences encoding functional IND proteins

The nucleic acid sequences depicted in the sequence listing encode wild type, functional IND proteins from *Brassica napus*. Thus, these sequences are endogenous to the *Brassica napus* plants from which they were isolated. Other *Brassica* crop species, varieties, breeding lines or wild accessions may be screened for other *IND* alleles, encoding the same IND proteins or variants thereof. For example, nucleic acid hybridization techniques (e.g. Southern blot analysis, using for example stringent hybridization conditions) or PCR-based techniques may be used to identify *IND* alleles endogenous to other *Brassica* plants, such as various *Brassica napus* varieties, lines or accessions, but also *Brassica juncea* (especially *IND* alleles on the A-genome), *Brassica carinata* (especially *IND* alleles on the C-genome) and *Brassica rapa* (A-genome) and *Brassica oleracea* (C-genome) plants, organs and tissues can be screened for other wild type *IND* alleles. To screen such plants, plant organs or tissues for the presence of *IND* alleles, the *IND* nucleic acid sequences provided in the sequence listing, or variants or fragments of any of these, may be used. For example whole sequences or fragments may be used as probes or primers. For example specific or degenerate primers may be used to amplify nucleic acid sequences encoding IND proteins from the genomic DNA of the plant, plant organ or tissue. These *IND* nucleic acid sequences may be isolated and sequenced using standard molecular biology techniques. Bioinformatics analysis may then be used to characterize the allele(s), for example in order to determine which *IND* allele the sequence corresponds to and which IND protein or protein variant is encoded by the sequence.

Whether a nucleic acid sequence encodes a functional IND protein can be analyzed by recombinant DNA techniques as known in the art, e.g., by a genetic complementation test using, e.g., an *Arabidopsis* plant, which is homozygous for a full knock-out *ind* mutant allele or a *Brassica napus* plant, which is homozygous for a full knock-out *ind* mutant allele of both the IND-A1 and IND-C1 gene.

In addition, it is understood that *IND* nucleic acid sequences and variants thereof (or fragments of any of these) may be identified *in silico*, by screening nucleic acid databases for essentially similar sequences. Likewise, a nucleic acid sequence may be synthesized chemically. Fragments

of nucleic acid molecules according to the invention are also provided, which are described further below. Fragments include nucleic acid sequences encoding only the bHLH domain, or smaller fragments comprising part of the bHLH domain, such as the basic domain or the HLH domain, etc.

5

Nucleic acid sequences encoding mutant IND proteins

Nucleic acid sequences comprising one or more nucleotide deletions, insertions or substitutions relative to the wild type nucleic acid sequences are another embodiment of the invention, as are fragments of such mutant nucleic acid molecules. Such mutant nucleic acid sequences (referred to as *ind* sequences) can be generated and/or identified using various known methods, as described further below. Again, such nucleic acid molecules are provided both in endogenous form and in isolated form. In one embodiment, the mutation(s) result in one or more changes (deletions, insertions and/or substitutions) in the amino acid sequence of the encoded IND protein (i.e. it is not a “silent mutation”). In another embodiment, the mutation(s) in the nucleic acid sequence result in a significantly reduced or completely abolished biological activity of the encoded IND protein relative to the wild type protein.

10
15

The nucleic acid molecules may, thus, comprise one or more mutations, such as:

(a) a “missense mutation”, which is a change in the nucleic acid sequence that results in the substitution of an amino acid for another amino acid;

20

(b) a “nonsense mutation” or “STOP codon mutation”, which is a change in the nucleic acid sequence that results in the introduction of a premature STOP codon and thus the termination of translation (resulting in a truncated protein); plant genes contain the translation stop codons “TGA” (UGA in RNA), “TAA” (UAA in RNA) and “TAG” (UAG in RNA); thus any nucleotide substitution, insertion, deletion which results in one of these codons to be in the mature mRNA being translated (in the reading frame) will terminate translation.

25

(c) an “insertion mutation” of one or more amino acids, due to one or more codons having been added in the coding sequence of the nucleic acid;

(d) a “deletion mutation” of one or more amino acids, due to one or more codons having been deleted in the coding sequence of the nucleic acid;

30

(e) a “frameshift mutation”, resulting in the nucleic acid sequence being translated in a different frame downstream of the mutation. A frameshift mutation can have various causes, such as the insertion, deletion or duplication of one or more nucleotides.

As already mentioned, it is desired that the mutation(s) in the nucleic acid sequence preferably result in a mutant protein comprising significantly reduced or no biological activity *in vivo* or in the production of no protein. Basically, any mutation which results in a protein comprising at least one amino acid insertion, deletion and/or substitution relative to the wild type protein can lead to significantly reduced or no biological activity. It is, however, understood that mutations in certain parts of the protein are more likely to result in a reduced function of the mutant IND protein, such as mutations leading to truncated proteins, whereby significant portions of the functional domains, such as the DNA binding domain ('b'), the dimerization domain ('HLH') and/or transcription regulating domains, are lacking.

10

According to The *Arabidopsis* Information Resource (TAIR) database (<http://www.arabidopsis.org/>), the *Arabidopsis* INDEHISCENT protein (locus At4g00120.1; SEQ ID NO: 10) is 198 amino acids in length and comprises a "basic helix-loop-helix (bHLH) dimerisation" domain located between the amino acids at position 121 and 168 (pfam domain PF00010), between the amino acids at position 124 and 173 (smart domain SM00353), or between the amino acids at position 112 and 168 (prosite domain PS50888) and an "helix-loop-helix (HLH) DNA binding" domain between the amino acids at position 114 and 196 or 198 (superfam domain G3D.4.10.280.10 or SSF47459, respectively) of SEQ ID NO: 10.

15

20

The IND-A1 protein of *Brassica* described herein is about 185 amino acids in length (SEQ ID NO:2) and the IND-C1 protein about 195 (SEQ ID NO:4 from the amino acid at position 16 to 210) or 210 (SEQ ID NO:4) amino acids and they comprise the "basic bHLH dimerisation" domain located between the amino acids at position 120 and 167 in SEQ ID NO: 2 and position 133 and 180 in SEQ ID NO: 4 (pfam domain PF00010), between the amino acids at position 123 and 172 in SEQ ID NO: 2 and position 136 and 185 in SEQ ID NO: 4 (smart domain SM00353), or between the amino acids at position 111 and 167 in SEQ ID NO: 2 and position 124 and 180 in SEQ ID NO: 4 (prosite domain PS50888) and the "HLH DNA binding" domain between the amino acids at position 127 and 208 or 210 in SEQ ID NO: 4 (superfam domain G3D.4.10.280.10 or SSF47459, respectively), as determined by optimally aligning the *Brassica* and *Arabidopsis* IND proteins and based on the annotation information in the TAIR database.

25

30

As described by Heim *et al.* (2003, Mol Biol Evol 20, 735-747), the consensus bHLH domain sequence of 133 *Arabidopsis* bHLH transcription factor genes consists of approximately 56 amino acids (Heim *et al.*, Fig. 1; corresponding to position 119-174 in SEQ ID NO: 10). This bipartite domain comprises (1) the basic region, located at the N-terminal end of the domain,

35

which is involved in DNA binding and consists of about 13 amino acids with a high number of basic residues (“b”; corresponding to position 119-131 in SEQ ID NO: 10), and (2) the helix-loop-helix region, located at the C-terminal end, which functions as a dimerization domain and is constituted of about 43 mainly hydrophobic amino acid residues (corresponding to position 132-174 in SEQ ID NO: 10) that form two amphipathic alpha-helices of about 15 amino acids (“H1”; corresponding to position 132-146 in SEQ ID NO: 10) and 22 amino acids (“H2”; corresponding to position 153-174 in SEQ ID NO: 10), respectively, separated by a loop region of about 6 and up to about 14 amino acids (“L”; corresponding to position 147-152 in SEQ ID NO: 10), which is the most divergent region of the bHLH domain in terms of size and amino acid composition.

10 The two alpha-helices promote dimerization, allowing the formation of homo- and/or heterodimers between different family members (Toledo-Ortiz *et al.*, 2003, Plant Cell 15: 1749-1770). While the bHLH domain is evolutionarily conserved (Atchley and Fitch, 1997, PNAS 94: 5172-5176), there is little sequence similarity between different bHLH family members beyond this domain (Morgenstern and Atchley, 1999, Mol Biol Evol 16: 1654-1663).

15

Within those bHLH proteins with proven ability to bind DNA, the amino acids at position 5, 9, and 13 of the consensus bHLH domain sequence defined by Heim *et al.* (*supra*) are the most critical. For non-plant bHLH proteins, it was shown that a His (H) residue at position 5, a Glu (E) residue at position 9 and an Arg (R) residue at position 13 (all within the basic region) were critical for DNA binding (Brownlie *et al.*, 1997, Structure 5, 509-520; Atchley *et al.*, 1999, J Mol Evol 48, 501-516; Ledent and Vervoort, 2001, Genome Res 11, 754-770). However, some plant proteins have a variation of the H-E-R configuration. For example, according to Heim *et al.* (*supra*), the 5-9-13 motif of the bHLH domain encoded by *Arabidopsis* gene At4g00120 (corresponding to the *Arabidopsis* *IND* gene represented in SEQ ID NO: 9) consists of amino acid residues Gln (Q), Ala (A) and Arg (R), respectively (corresponding to positions 123, 127 and 131, respectively, in SEQ ID NO: 10) (Figure 4 of Heim *et al.* (*supra*)). Such plant proteins, which have a variation of the H-E-R configuration, may further contain helix-breaking prolines in the basic region, e.g. members of Group VIII and X, characteristics that may interfere with affinity for DNA. These variations may enable these proteins to act as negative regulators, retaining the ability to dimerize with other bHLH proteins but lacking the ability to bind DNA. While the 5-9-13 motif is important for DNA binding, the DNA backbone is contacted by the basic residues at positions 10 and 12 (both Arg (R) in the consensus bHLH domain sequence), which are also conserved in the majority of plant proteins (corresponding to positions 128 and 130 in SEQ ID NO: 10).

35

Furthermore, Heim *et al.* (*supra*) describe that the highly conserved hydrophobic residues at position 16, 20, 23, 27 in helix1 (corresponding to position 134, 138, 141, 145 in SEQ ID NO: 10) and at position 36, 39, 43, 49, 53, and 56 in helix2 (corresponding to position 154, 157, 161, 167, 171, 174 in SEQ ID NO: 10), for example, the leucine residue at position 23 within the helix1 domain (corresponding to position 141 in SEQ ID NO: 10) and the conserved hydrophobic residues in helix 2 that are located to one side of the helix, are necessary for dimerization or stabilization of dimer formation.

Finally, Heim *et al.* (*supra*; Fig. 4) indicate conserved amino acid sequences outside the DNA binding domain, some of which are thought to act as activation domain or be important for interaction with other modules of the transcription complex or to be targets of signal transduction chains.

Table 1 IND proteins - amino acids (AA) regions and positions

		AtIND1 (SEQ ID NO: 10)	AtIND1 (SEQ ID NO: 9)	BnIND-A1 (SEQ ID NO: 2/6)	BnIND-C1a/b (SEQ ID 4/8 from 16-210 / SEQ ID 4/8)
<u>Coding region</u>	<u>TAIR:</u>	1-198 (198 AA)	1-594	1-185 (185 AA)	16-210 / 1-210 (195 / 210 AA)
	PF00010	121-168	361-504	120-167	133-180
	SM00353	124-173	370-519	123-172	136-185
	PS50888	112-168	334-504	111-167	124-180
	G3D.4.10.280.10	114-196	340-588	-	127-208
	SSF47459	114-198	340-594	-	127-210
	<u>Liljegren et al.</u>	30-198 (169 AA)	88-594		
<u>bHLH:</u>	<u>Heim et al.</u>	119-174	355-523	118-173	131-186
	<u>Toledo-Ortiz et al.</u>	115-167	343-501	114-166	127-179
	<u>Liljegren et al.</u>	119-167	355-501	118-166	131-179
b	<u>Heim et al.</u>	119-131	355-393	118-132	131-145
	<u>Toledo-Ortiz et al.</u>	115-131	343-393	114-132	127-145
	<u>Liljegren et al.</u>	119-131	355-393	118-132	131-145
H1	<u>Heim et al.</u>	132-146	394-438	133-145	146-158

	Toledo-Ortiz <i>et al.</i>	132-146	394-438	133-145	146-158
	Liljegren <i>et al.</i>	132-145	394-435	133-144	146-157
L	Heim <i>et al.</i>	147-152	439-456	146-151	159-164
	Toledo-Ortiz <i>et al.</i>	147-152	439-456	146-151	159-164
	Liljegren <i>et al.</i>	146-152	436-456	145-151	158-164
H2	Heim <i>et al.</i>	153-174	457-523	152-173	165-186
	Toledo-Ortiz <i>et al.</i>	153-167	457-501	152-166	165-179
	Liljegren <i>et al.</i>	153-167	457-501	152-166	165-179
<u>Conserved</u>	N (1 ^T)	115	343-345	114	127
<u>AA</u>	V (2 ^T)	116	346-348	115	128
	Q (5 ^H)	123	367-379	122	135
	A (9 ^H - 13 ^T)	127	379-381	126	139
	R (10 ^H - 14 ^T)	128	382-384	127	140
	R (12 ^H - 16 ^T)	130	388-390	129	142
	R (13 ^H)	131	391-393	130	143
	I (16 ^H - 20 ^T)	134	400-403	133	146
	S (21 ^T)	135	404-406	134	147
	I (20 ^H - 24 ^T)	138	412-414	137	150
	L (23 ^H - 27 ^T)	141	421-423	140	153
	K (28 ^T)	142	424-426	141	154
	V (27 ^H)	145	433-435	144	157
	K (39 ^T)	150	448-450	149	162
	T (42 ^T)	153	460-463	152	165
	A (36 ^H)	154	460-462	153	166
	M (45 ^T)	156	466-468	155	168
	L (39 ^H -46 ^T)	157	469-471	156	169
	A (49 ^T)	160	478-480	159	172
	I (43 ^H - 50 ^T)	161	481-483	160	173
	Y (52 ^T)	163	487-489	162	175
	T (53 ^T)	164	490-492	163	176
	L (49 ^H -56 ^T)	167	499-501	166	179
	V (53 ^H)	171	511-513	170	183
	L (56 ^H)	174	580-582	173 (A)	186

<i>At ind</i>	<i>ind-5</i> (W13>STOP) ^L	42	124-126	25	41
	<i>ind-2</i> (A26>FS) ^L	55	163-165	-	-
	<i>ind-6</i> ^W	Insertion after 61	Insertion after 185	-	-
	<i>ind-4</i> (Q63>STOP) ^L	92	274-276	91	104
	<i>ind-3</i> (R99>H) ^L	128	382-384	127	140
	<i>ind-1</i> (L112>F) ^L	141	421-423	140	153

Heim et al.,^H: Heim *et al.*, 2003, Mol Biol Evol 20, 735-747; Toledo-Ortiz *et al.*,^L: Toledo-Ortiz *et al.*, 2003, Plant Cell 15: 1749-1770; Liljegren *et al.*,^L: Liljegren *et al.*, 2004, Cell, 116, 843-853; ^W: Wu *et al.*, 2006, Planta 224, 971-979.

5

Similarly, as described by Toledo-Ortiz *et al.* (2003, Plant Cell 15: 1749-1770; Figure 1), the bHLH domain of the *Arabidopsis* bHLH transcription factor family consists of approximately 56 amino acids (Toledo-Ortiz *et al.*; corresponding to position 115-167 in SEQ ID NO: 10). This bipartite domain comprises (1) the basic region, located at the N-terminal end of the domain, which is involved in DNA binding and consists of about 17 amino acids with a high number of basic residues (“b”; corresponding to position 115-131 in SEQ ID NO: 10), and (2) the HLH region, located at the C-terminal end, which functions as a dimerization domain and is constituted of about 39 mainly hydrophobic amino acid residues (corresponding to position 132-167 in SEQ ID NO: 10) that form two amphipathic alpha-helices of about 15 amino acids (“H1” corresponding to position 132-146 in SEQ ID NO: 10, and “H2” corresponding to position 152-167 in SEQ ID NO: 10) separated by a loop region of about 9 amino acids (“L”; corresponding to position 147-151 in SEQ ID NO: 10), which is the most divergent region of the bHLH domain in terms of size and amino acid composition.

Based on patterns of sequence conservation, a hypothetical consensus motif, representing the most conserved amino acids in the bHLH region, including 19 amino acids dispersed across the bHLH domain (18 from b, H1 and H2; 1 from L) was generated by Atchley *et al.* (1999). The identified conserved amino acids correspond to the amino acids at position 1, 2, 13, 14, 16 (in b); 20, 21, 24, 27, 28 (in H1); 39 (in L); 42, 45, 46, 49, 50, 52, 53, and 56 (in H2) of the *Arabidopsis* bHLH domain defined by Toledo-Ortiz *et al.* (2003, *supra*), which correspond to the amino acids at position 115, 116, 127, 128, 130 (in b); 134, 135, 138, 141, 142 (in H1); 150 (in L); 153, 156, 157, 160, 161, 163, 164, and 167 (in H2) of SEQ ID NO: 10.

According to Liljegren *et al.* (2004, Cell, 116, 843-853), the bHLH domain of the *Arabidopsis IND* gene comprises a basic region of 13 amino acids (SEQ ID NO: 10 from the amino acid at position 119 to 131) and two alpha-helices of 14 and 15 amino acids, respectively, (SEQ ID NO: 10 from the amino acid at position 132-145 and from the amino acid at position 153 to 167, respectively) separated by a variable loop region of 7 amino acids (SEQ ID NO: 10 from the amino acid at position 146 to 152).

Optimal alignment of the *Arabidopsis IND* nucleic acid (SEQ ID NO: 9) and amino acid (SEQ ID NO: 10) sequences with *IND* nucleic acid sequences, in particular the *Brassica IND* nucleic acid (SEQ ID NO: 1 and 3) and amino acid (SEQ ID NO: 2 and 4) sequences of the present invention, allows to determine the positions of the corresponding conserved domains and amino acids in these *Brassica* sequences (see Table 1 for the *Brassica IND* sequences of SEQ ID NO: 1 to 4).

Thus in one embodiment, nucleic acid sequences comprising one or more of any of the types of mutations described above are provided. In another embodiment, *ind* sequences comprising one or more stop codon (nonsense) mutations, one or more missense mutations and/or one or more frameshift mutations are provided. Any of the above mutant nucleic acid sequences are provided *per se* (in isolated form), as are plants and plant parts comprising such sequences endogenously. In the tables herein below the most preferred *ind* alleles are described and seed deposits of *Brassica napus* seeds comprising one or more *ind* alleles have been deposited as indicated.

A nonsense mutation in an *IND* allele, as used herein, is a mutation in an *IND* allele whereby one or more translation stop codons are introduced into the coding DNA and the corresponding mRNA sequence of the corresponding wild type *IND* allele. Translation stop codons are TGA (UGA in the mRNA), TAA (UAA) and TAG (UAG). Thus, any mutation (deletion, insertion or substitution) that leads to the generation of an in-frame stop codon in the coding sequence will result in termination of translation and truncation of the amino acid chain. In one embodiment, a mutant *IND* allele comprising a nonsense mutation is an *IND* allele wherein an in-frame stop codon is introduced in the *IND* codon sequence by a single nucleotide substitution, such as the mutation of CAG to TAG, TGG to TAG, TGG to TGA, or CAA to TAA. In another embodiment, a mutant *IND* allele comprising a nonsense mutation is an *IND* allele wherein an in-frame stop codon is introduced in the *IND* codon sequence by double nucleotide substitutions, such as the mutation of CAG to TAA, TGG to TAA, or CGG to TAG or TGA. In yet another embodiment, a mutant *IND* allele comprising a nonsense mutation is an *IND* allele wherein an

in-frame stop codon is introduced in the *IND* codon sequence by triple nucleotide substitutions, such as the mutation of CGG to TAA. The truncated protein lacks the amino acids encoded by the coding DNA downstream of the mutation (i.e. the C-terminal part of the IND protein) and maintains the amino acids encoded by the coding DNA upstream of the mutation (i.e. the N-terminal part of the IND protein). In one embodiment, a mutant *IND* allele comprising a nonsense mutation is an *IND* allele wherein the nonsense mutation is present anywhere in front of the conserved Leu residue of the H2 domain (at position 56 in the consensus bHLH domain sequence as described by Heim *et al.*, 2003, see above), so that at least the conserved Leu residue is lacking. The more truncated the mutant IND protein is in comparison to the wild type IND protein, the more the truncation may result in a significantly reduced or no activity of the IND protein. Thus in another embodiment, a mutant *IND* allele comprising a nonsense mutation which results in a truncated protein of less than about 170 amino acids (lacking the conserved Leu), less than about 150 amino acids (lacking the H2 domain), less than about 145 amino acids (lacking the L and H2 domains), less than about 130 amino acids (lacking the HLH domain), less than about 115 amino acids (lacking the bHLH domain), or even less amino acids in length, such as mutant *IND* alleles corresponding to the *Arabidopsis ind-4* or *ind-5* (Liljegren *et al.*, 2004, *supra*) alleles are provided (see Table 1).

The Tables herein below describe a range of possible nonsense mutations in the *Brassica napus IND* sequences provided herein:

Table 2a Potential STOP codon mutations in *IND-A1* (SEQ ID NO: 1)

Amino acid position	Nucleotide position	Wild type → mutant codon	Wild type → mutant amino acid
25	74	tgg → tag	TRP → STOP
	75	tgg → tga	TRP → STOP
	74+75	tgg → taa	TRP → STOP
57	169	cag → tag	GLN → STOP
	169+171	cag → taa	GLN → STOP
91	271	caa → taa	GLN → STOP
98	292	cag → tag	GLN → STOP
	292+294	cag → taa	GLN → STOP
122	364	cag → tag	GLN → STOP (1)
	364+366	cag → taa	GLN → STOP
128	382+383	cgg → tag	ARG → STOP
	382+384	cgg → tga	ARG → STOP
	382+383+384	cgg → taa	ARG → STOP
138	412+413	cgg → tag	ARG → STOP
	412+414	cgg → tga	ARG → STOP

	412+413+414	<u>cgg</u> → <u>taa</u>	ARG → STOP
168	502+503	<u>cgg</u> → <u>tag</u>	ARG → STOP
	502+504	<u>cgg</u> → <u>tga</u>	ARG → STOP
	502+503+504	<u>cgg</u> → <u>taa</u>	ARG → STOP
169	505	<u>cag</u> → <u>tag</u>	GLN → STOP
	505+507	<u>cag</u> → <u>taa</u>	GLN → STOP
181	542	<u>tgg</u> → <u>tag</u>	TRP → STOP
	543	<u>tgg</u> → <u>tga</u>	TRP → STOP
	542+543	<u>tgg</u> → <u>taa</u>	TRP → STOP

(1) seeds comprising a mutant IND-A1 allele comprising this non-sense mutation (called hereinafter *ind-a1*-EMS01) have been deposited at the American Type Culture Collection (ATCC, 10801 University Boulevard, Manassas, VA 20110-2209, US) on November 20, 2007, under accession number PTA-8796

5

Table 2b Potential STOP codon mutations in *IND-C1* (SEQ ID NO: 3)

Amino acid position	Nucleotide position	Wild type → mutant codon	Wild type → mutant amino acid
41	122	<u>tgg</u> → <u>tag</u>	TRP → STOP
	123	<u>tgg</u> → <u>tga</u>	TRP → STOP
	122+123	<u>tgg</u> → <u>taa</u>	TRP → STOP
50	148	<u>caa</u> → <u>taa</u>	GLN → STOP (2)
73	271	<u>cag</u> → <u>tag</u>	GLN → STOP
	271+272	<u>cag</u> → <u>taa</u>	GLN → STOP
104	310	<u>caa</u> → <u>taa</u>	GLN → STOP
111	331	<u>cag</u> → <u>tag</u>	GLN → STOP
	331+333	<u>cag</u> → <u>taa</u>	GLN → STOP
135	403	<u>cag</u> → <u>tag</u>	GLN → STOP (3)
	403+405	<u>cag</u> → <u>taa</u>	GLN → STOP
141	421+422	<u>cgg</u> → <u>tag</u>	ARG → STOP
	421+423	<u>cgg</u> → <u>tga</u>	ARG → STOP
	421+422+423	<u>cgg</u> → <u>taa</u>	ARG → STOP
151	451+452	<u>cgg</u> → <u>tag</u>	ARG → STOP
	451+453	<u>cgg</u> → <u>tga</u>	ARG → STOP
	451+452+453	<u>cgg</u> → <u>taa</u>	ARG → STOP
181	541+542	<u>cgg</u> → <u>tag</u>	ARG → STOP
	541+543	<u>cgg</u> → <u>tga</u>	ARG → STOP
	541+542+543	<u>cgg</u> → <u>taa</u>	ARG → STOP
182	544	<u>cag</u> → <u>tag</u>	GLN → STOP
	544+546	<u>cag</u> → <u>taa</u>	GLN → STOP
187	559	<u>cag</u> → <u>tag</u>	GLN → STOP
	559+561	<u>cag</u> → <u>taa</u>	GLN → STOP
191	571	<u>cag</u> → <u>tag</u>	GLN → STOP
	571+573	<u>cag</u> → <u>taa</u>	GLN → STOP

(2) seeds comprising a mutant IND-C1 allele comprising this non-sense mutation (called hereinafter *ind-cl*-EMS01) have been deposited at the ATCC on November 20, 2007, under accession number PTA-8796

(3) seeds comprising a mutant IND-C1 allele comprising this non-sense mutation (called hereinafter *ind-c1*-EMS03) have been deposited at the ATCC on November 20, 2007, under accession number PTA-8795

5 Obviously, mutations are not limited to the ones shown in the above tables and it is understood that analogous STOP mutations may be present in *ind* alleles other than those depicted in the sequence listing and referred to in the tables above.

A missense mutation in an *IND* allele, as used herein, is any mutation (deletion, insertion or
10 substitution) in an *IND* allele whereby one or more codons are changed into the coding DNA and the corresponding mRNA sequence of the corresponding wild type *IND* allele, resulting in the substitution of one or more amino acids in the wild type *IND* protein for one or more other amino acids in the mutant *IND* protein. In one embodiment, a mutant *IND* allele comprising a missense mutation is an *IND* allele wherein one or more of the conserved amino acids indicated
15 above or in Table 1 is/are substituted. As indicated above, some of the conserved amino acids are more critical for the biological activity of the *IND* protein than others. Thus, missense mutations which result in the substitution of, e.g., the amino acids at position 5, 9, and 13 or at positions 10 and 12 of the consensus bHLH domain sequence defined by Heim *et al.* (*supra*) are more likely to result in a significantly reduced or no activity, due to a reduced ability to bind to the target
20 DNA, of the *IND* protein. Similarly missense mutations which result in the substitution of, e.g., the amino acids at position 16, 20, 23, 27 in helix1 or at positions 36, 39, 43, 49, 53, and 56 in helix2 of the consensus bHLH domain sequence defined by Heim *et al.* (*supra*) are more likely to result in a significantly reduced or no activity, due to a reduced dimerization ability, of the *IND* protein. Seeds comprising a mutant *IND*-A1 allele comprising a missense mutation which
25 causes the substitution of the Arg residue at position 10 of the consensus bHLH domain sequence defined by Heim *et al.* (*supra*) for an His residue (called hereinafter *ind-a1*-EMS05) have been deposited at the ATCC on November 20, 2007, under accession number PTA-8795. In another embodiment, a mutant *IND* allele comprising a missense mutation is an *IND* allele comprising a missense mutation corresponding to the missense mutation in the *Arabidopsis ind-1* or *ind-3* (Liljegren *et al.*, 2004, *supra*) alleles (see Table 1).
30

A frameshift mutation in an *IND* allele, as used herein, is a mutation (deletion, insertion, duplication, and the like) in an *IND* allele that results in the nucleic acid sequence being translated in a different frame downstream of the mutation. In one embodiment, a mutant *IND*
35 allele comprising a frameshift mutation is an *IND* allele comprising a frameshift mutation corresponding to the frameshift mutation in the *Arabidopsis ind-2* (Liljegren *et al.*, 2004, *supra*)

allele, wherein a single nucleotide is deleted within codon 26, which results in a frameshift and production of a truncated protein of 35 amino acids (according to Liljegen *et al.*, 2004, *supra*). In another embodiment, a mutant *IND* allele comprising a frameshift mutation is an *IND* allele comprising a frameshift mutation corresponding to the frameshift mutation in the *Arabidopsis ind-6* (Wu *et al.*, 2006, *supra*) allele, wherein a Ds transposon is inserted after nucleotide 183 causing an 8 nucleotide duplication at the insertion site, or to the corresponding revertant *Arabidopsis ind* alleles (see Wu *et al.*, 2006, *supra*, Fig. 1a).

Amino acid sequences according to the invention

10 Provided are both wild type (functional) *IND* amino acid sequences and mutant *IND* amino acid sequences (comprising one or more mutations, preferably mutations which result in a significantly reduced or no biological activity of the *IND* protein) from *Brassicaceae*, particularly from *Brassica* species, especially from *Brassica napus*, but also from other *Brassica* crop species. For example, *Brassica* species comprising an A and/or a C genome may encode
15 different *IND-A* or *IND-C* amino acids. In addition, mutagenesis methods can be used to generate mutations in wild type *IND* alleles, thereby generating mutant alleles which can encode further mutant *IND* proteins. In one embodiment the wild type and/or mutant *IND* amino acid sequences are provided within a *Brassica* plant (i.e. endogenously). However, isolated *IND* amino acid sequences (e.g. isolated from the plant or made synthetically), as well as variants
20 thereof and fragments of any of these are also provided herein.

Amino acid sequences of *IND-A1* and *IND-C1* proteins have been isolated from *Brassica napus* as depicted in the sequence listing. The wild type *IND* sequences are depicted, while the mutant *IND* sequences of these sequences, and of sequences essentially similar to these, are described
25 herein below, with reference to the wild type *IND* sequences.

As described above, the *IND* proteins of *Brassica* described herein are about 185-210 amino acids in length and comprise a number of structural and functional domains.

30 “*IND-A1* amino acid sequences” or “*IND-A1* variant amino acid sequences” according to the invention are amino acid sequences having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 2. These amino acid sequences may also be referred to as being “essentially similar” or “essentially identical” to the *IND* sequences provided in the sequence listing.

“IND-C1 amino acid sequences” or “IND-C1 variant amino acid sequences” according to the invention are amino acid sequences having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, 96%, 97%, 98%, 99% or 100% sequence identity with SEQ ID NO: 4 (IND-C1-long) or with SEQ ID NO:4 from the amino acid at position 16 to the amino acid at position 210 (IND-C1-short). These amino acid sequences may also be referred to as being “essentially similar” or “essentially identical” the *IND* sequences provided in the sequence listing.

Thus, the invention provides both amino acid sequences of wild type, functional IND-A1 and IND-C1 proteins, including variants and fragments thereof (as defined further below), as well as mutant amino acid sequences of any of these, whereby the mutation in the amino acid sequence preferably results in a significant reduction in or a complete abolishment of the biological activity of the IND protein as compared to the biological activity of the corresponding wild type IND protein. A significant reduction in or complete abolishment of the biological activity of the IND protein refers herein to a reduction in or abolishment of the DNA binding activity, the dimerization capacity and/or transcriptional regulating activity of the IND protein, such that the pod shatter resistance of a plant expressing the mutant IND protein is increased as compared to a plant expressing the corresponding wild type IND protein compared to the pod shatter resistance of a corresponding wild type plant.

Both endogenous and isolated amino acid sequences are provided herein. Also provided are fragments of the IND amino acid sequences and IND variant amino acid sequences defined above. A “fragment” of a IND amino acid sequence or variant thereof (as defined) may be of various lengths, such as at least 10, 12, 15, 18, 20, 50, 100, 150, 175, 180 contiguous amino acids of the *IND* sequence (or of the variant sequence).

25

Amino acid sequences of functional IND proteins

The amino acid sequences depicted in the sequence listing are wild type, functional IND proteins from *Brassica napus*. Thus, these sequences are endogenous to the *Brassica napus* plants from which they were isolated. Other *Brassica* crop species, varieties, breeding lines or wild accessions may be screened for other functional IND proteins with the same amino acid sequences or variants thereof, as described above.

In addition, it is understood that IND amino acid sequences and variants thereof (or fragments of any of these) may be identified *in silico*, by screening amino acid databases for essentially similar sequences. Fragments of amino acid molecules according to the invention are also

35

provided. Fragments include amino acid sequences of the bHLH domain, or smaller fragments comprising part of the bHLH domain, such as the basic domain or the HLH domain, etc.

Amino acid sequences of mutant IND proteins

5 Amino acid sequences comprising one or more amino acid deletions, insertions or substitutions relative to the wild type amino acid sequences are another embodiment of the invention, as are fragments of such mutant amino acid molecules. Such mutant amino acid sequences can be generated and/or identified using various known methods, as described above. Again, such amino acid molecules are provided both in endogenous form and in isolated form.

10

In one embodiment, the mutation(s) in the amino acid sequence result in a significantly reduced or completely abolished biological activity of the IND protein relative to the wild type protein. As described above, basically, any mutation which results in a protein comprising at least one amino acid insertion, deletion and/or substitution relative to the wild type protein can lead to significantly reduced or no biological activity. It is, however, understood that mutations in certain parts of the protein are more likely to result in a reduced function of the mutant IND protein, such as mutations leading to truncated proteins, whereby significant portions of the functional domains, such as the DNA binding domain ('b'), the dimerization domain ('HLH') and/or amino acids which are important in the regulation of transcription (See Table 1), are lacking or are being substituted.

20

Thus in one embodiment, mutant IND proteins are provided comprising one or more deletion or insertion mutations, whereby the deletion(s) or insertion(s) result(s) in a mutant protein which has significantly reduced or no activity *in vivo*. Such mutant IND proteins are IND proteins wherein at least 1, at least 2, 3, 4, 5, 10, 20, 30, 50, 100, 100, 150, 175, 180 or more amino acids are deleted or inserted as compared to the wild type IND protein, whereby the deletion(s) or insertion(s) result(s) in a mutant protein which has significantly reduced or no activity *in vivo*.

25

In another embodiment, mutant IND proteins are provided which are truncated whereby the truncation results in a mutant protein that has significantly reduced or no activity *in vivo*. Such truncated IND proteins are IND proteins which lack functional domains in the C-terminal part of the corresponding wild type IND protein and which maintain the N-terminal part of the corresponding wild type IND protein. Thus in one embodiment, a truncated IND protein comprising the N-terminal part of the corresponding wild type IND protein up to but not including the conserved Leu residue of the H2 domain (at position 56 in the consensus bHLH

35

domain sequence as described by Heim *et al.*, 2003, see above) is provided. The more truncated the mutant protein is in comparison to the wild type protein, the more the truncation may result in a significantly reduced or no activity of the IND protein. Thus in another embodiment, a truncated IND protein comprising the N-terminal part of the corresponding wild type IND protein lacking part or all of the second H domain, and/or lacking part or all of the L domain, and/or lacking part or all of the first H domain, and/or lacking part or all of the basic domain (as described above), or even more amino acids are provided (see Table above).

In yet another embodiment, mutant IND proteins are provided comprising one or more substitution mutations, whereby the substitution(s) result(s) in a mutant protein that has significantly reduced or no activity *in vivo*. Such mutant IND proteins are IND proteins whereby conserved amino acid residues which have a specific function, such as a function in DNA binding, dimerization or transcription regulation, are substituted. Thus in one embodiment, a mutant IND protein comprising a substitution of a conserved amino acid residue which has a biological function, such as the conserved amino acids of the basic domain, or the H1, L or H2 domain as indicated in Table 1 above, is provided.

Methods according to the invention

Mutant *ind* alleles may be generated (for example induced by mutagenesis) and/or identified using a range of methods, which are conventional in the art, for example using PCR based methods to amplify part or all of the *ind* genomic or cDNA.

Following mutagenesis, plants are grown from the treated seeds, or regenerated from the treated cells using known techniques. For instance, mutagenized seeds may be planted in accordance with conventional growing procedures and following self-pollination seed is formed on the plants. Alternatively, doubled haploid plantlets may be extracted from treated microspore or pollen cells to immediately form homozygous plants, for example as described by Coventry *et al.* (1988, Manual for Microspore Culture Technique for *Brassica napus*. Dep. Crop Sci. Techn. Bull. OAC Publication 0489. Univ. of Guelph, Guelph, Ontario, Canada). Additional seed which is formed as a result of such self-pollination in the present or a subsequent generation may be harvested and screened for the presence of mutant *IND* alleles, using techniques which are conventional in the art, for example polymerase chain reaction (PCR) based techniques (amplification of the *ind* alleles) or hybridization based techniques, e.g. Southern blot analysis, BAC library screening, and the like, and/or direct sequencing of *ind* alleles. To screen for the presence of point mutations (so called Single Nucleotide Polymorphisms or SNPs) in mutant

IND alleles, SNP detection methods conventional in the art can be used, for example oligoligation-based techniques, single base extension-based techniques or techniques based on differences in restriction sites, such as TILLING.

5 As described above, mutagenization (spontaneous as well as induced) of a specific wild-type *IND* allele results in the presence of one or more deleted, inserted, or substituted nucleotides (hereinafter called "mutation region") in the resulting mutant *IND* allele. The mutant *IND* allele can thus be characterized by the location and the configuration of the one or more deleted, inserted, or substituted nucleotides in the wild type *IND* allele. The site in the wild type *IND*
10 allele where the one or more nucleotides have been inserted, deleted, or substituted, respectively, is herein also referred to as the "mutation region or sequence". A "5' or 3' flanking region or sequence" as used herein refers to a DNA region or sequence in the mutant (or the corresponding wild type) *IND* allele of at least 20 bp, preferably at least 50 bp, at least 750 bp, at least 1500 bp, and up to 5000 bp of DNA different from the DNA containing the one or more deleted, inserted,
15 or substituted nucleotides, preferably DNA from the mutant (or the corresponding wild type) *IND* allele which is located either immediately upstream of and contiguous with (5' flanking region or sequence") or immediately downstream of and contiguous with (3' flanking region or sequence") the mutation region in the mutant *IND* allele (or in the corresponding wild type *IND* allele). A "joining region" as used herein refers to a DNA region in the mutant (or the
20 corresponding wild type) *IND* allele where the mutation region and the 5' or 3' flanking region are linked to each other. A "sequence spanning the joining region between the mutation region and the 5' or 3' flanking region thus comprises a mutation sequence as well as the flanking sequence contiguous therewith.

25 The tools developed to identify a specific mutant *IND* allele or the plant or plant material comprising a specific mutant *IND* allele, or products which comprise plant material comprising a specific mutant *IND* allele are based on the specific genomic characteristics of the specific mutant *IND* allele as compared to the genomic characteristics of the corresponding wild type *IND* allele, such as, a specific restriction map of the genomic region comprising the mutation
30 region, molecular markers or the sequence of the flanking and/or mutation regions.

Once a specific mutant *IND* allele has been sequenced, primers and probes can be developed which specifically recognize a sequence within the 5' flanking, 3' flanking and/or mutation regions of the mutant *IND* allele in the nucleic acid (DNA or RNA) of a sample by way of a
35 molecular biological technique. For instance a PCR method can be developed to identify the

mutant *IND* allele in biological samples (such as samples of plants, plant material or products comprising plant material). Such a PCR is based on at least two specific “primers”: one recognizing a sequence within the 5’ or 3’ flanking region of the mutant *IND* allele and the other recognizing a sequence within the 3’ or 5’ flanking region of the mutant *IND* allele, respectively; or one recognizing a sequence within the 5’ or 3’ flanking region of the mutant *IND* allele and the other recognizing a sequence within the mutation region of the mutant *IND* allele; or one recognizing a sequence within the 5’ or 3’ flanking region of the mutant *IND* allele and the other recognizing a sequence spanning the joining region between the 3’ or 5’ flanking region and the mutation region of the specific mutant *IND* allele (as described further below), respectively.

10

The primers preferably have a sequence of between 15 and 35 nucleotides which under optimized PCR conditions “specifically recognize” a sequence within the 5’ or 3’ flanking region, a sequence within the mutation region, or a sequence spanning the joining region between the 3’ or 5’ flanking and mutation regions of the specific mutant *IND* allele, so that a specific fragment (“mutant *IND* specific fragment” or discriminating amplicon) is amplified from a nucleic acid sample comprising the specific mutant *IND* allele. This means that only the targeted mutant *IND* allele, and no other sequence in the plant genome, is amplified under optimized PCR conditions.

15

20 PCR primers suitable for the invention may be the following:

- oligonucleotides ranging in length from 17 nt to about 200 nt, comprising a nucleotide sequence of at least 17 consecutive nucleotides, preferably 20 consecutive nucleotides selected from the 5’ or 3’ flanking sequence of a specific mutant *IND* allele or the complement thereof (i.e., for example, the sequence 5’ or 3’ flanking the one or more nucleotides deleted, inserted or substituted in the mutant *IND* alleles of the invention, such as the sequence 5’ or 3’ flanking the non-sense, mis-sense or frameshift mutations described above or the sequence 5’ or 3’ flanking the STOP codon mutations indicated in the above Tables or the substitution mutations indicated above or the complement thereof) (primers recognizing 5’ flanking sequences); or

25

- oligonucleotides ranging in length from 17 nt to about 200 nt, comprising a nucleotide sequence of at least 17 consecutive nucleotides, preferably 20 nucleotides selected from the sequence of the mutation region of a specific mutant *IND* allele or the complement thereof (i.e., for example, the sequence of nucleotides inserted or substituted in the *IND* genes of the invention or the complement thereof) (primers recognizing mutation sequences) .

30

The primers may of course be longer than the mentioned 17 consecutive nucleotides, and may e.g. be 18, 19, 20, 21, 30, 35, 50, 75, 100, 150, 200 nt long or even longer. The primers may entirely consist of nucleotide sequence selected from the mentioned nucleotide sequences of flanking and mutation sequences. However, the nucleotide sequence of the primers at their 5' end (i.e. outside of the 3'-located 17 consecutive nucleotides) is less critical. Thus, the 5' sequence of the primers may consist of a nucleotide sequence selected from the flanking or mutation sequences, as appropriate, but may contain several (e.g. 1, 2, 5, 10) mismatches. The 5' sequence of the primers may even entirely consist of a nucleotide sequence unrelated to the flanking or mutation sequences, such as e.g. a nucleotide sequence representing restriction enzyme recognition sites. Such unrelated sequences or flanking DNA sequences with mismatches should preferably be not longer than 100, more preferably not longer than 50 or even 25 nucleotides.

Moreover, suitable primers may comprise or consist of a nucleotide sequence spanning the joining region between flanking and mutation sequences (i.e., for example, the joining region between a sequence 5' or 3' flanking one or more nucleotides deleted, inserted or substituted in the mutant *IND* alleles of the invention and the sequence of the one or more nucleotides inserted or substituted or the sequence 3' or 5', respectively, flanking the one or more nucleotides deleted, such as the joining region between a sequence 5' or 3' flanking non-sense, missense or frameshift mutations in the *IND* genes of the invention described above and the sequence of the non-sense, missense or frameshift mutations, or the joining region between a sequence 5' or 3' flanking a potential STOP codon mutation as indicated in the above Tables or the substitution mutations indicated above and the sequence of the potential STOP codon mutation or the substitution mutations, respectively), provided the nucleotide sequence is not derived exclusively from either the mutation region or flanking regions.

It will also be immediately clear to the skilled artisan that properly selected PCR primer pairs should also not comprise sequences complementary to each other.

30

For the purpose of the invention, the "complement of a nucleotide sequence represented in SEQ ID No: X" is the nucleotide sequence which can be derived from the represented nucleotide sequence by replacing the nucleotides through their complementary nucleotide according to Chargaff's rules ($A \leftrightarrow T$; $G \leftrightarrow C$) and reading the sequence in the 5' to 3' direction, i.e. in opposite direction of the represented nucleotide sequence.

35

Examples of primers suitable to identify specific mutant *IND* alleles are described in the Examples.

- 5 As used herein, “the nucleotide sequence of SEQ ID No. Z from position X to position Y” indicates the nucleotide sequence including both nucleotide endpoints.

Preferably, the amplified fragment has a length of between 50 and 1000 nucleotides, such as a length between 50 and 500 nucleotides, or a length between 100 and 350 nucleotides. The specific primers may have a sequence which is between 80 and 100% identical to a sequence within the 5' or 3' flanking region, to a sequence within the mutation region, or to a sequence spanning the joining region between the 3' or 5' flanking and mutation regions of the specific mutant *IND* allele, provided the mismatches still allow specific identification of the specific mutant *IND* allele with these primers under optimized PCR conditions. The range of allowable mismatches however, can easily be determined experimentally and are known to a person skilled in the art.

Detection and/or identification of a “mutant *IND* specific fragment” can occur in various ways, e.g., via size estimation after gel or capillary electrophoresis or via fluorescence-based detection methods. The mutant *IND* specific fragments may also be directly sequenced. Other sequence specific methods for detection of amplified DNA fragments are also known in the art.

Standard PCR protocols are described in the art, such as in "PCR Applications Manual" (Roche Molecular Biochemicals, 2nd Edition, 1999) and other references. The optimal conditions for the PCR, including the sequence of the specific primers, is specified in a “PCR identification protocol” for each specific mutant *IND* allele. It is however understood that a number of parameters in the PCR identification protocol may need to be adjusted to specific laboratory conditions, and may be modified slightly to obtain similar results. For instance, use of a different method for preparation of DNA may require adjustment of, for instance, the amount of primers, polymerase, MgCl₂ concentration or annealing conditions used. Similarly, the selection of other primers may dictate other optimal conditions for the PCR identification protocol. These adjustments will however be apparent to a person skilled in the art, and are furthermore detailed in current PCR application manuals such as the one cited above.

Examples of PCR identification protocols to identify specific mutant *IND* alleles are described in the Examples.

Alternatively, specific primers can be used to amplify a mutant *IND* specific fragment that can be used as a “specific probe” for identifying a specific mutant *IND* allele in biological samples. Contacting nucleic acid of a biological sample, with the probe, under conditions that allow hybridization of the probe with its corresponding fragment in the nucleic acid, results in the formation of a nucleic acid/probe hybrid. The formation of this hybrid can be detected (e.g. labeling of the nucleic acid or probe), whereby the formation of this hybrid indicates the presence of the specific mutant *IND* allele. Such identification methods based on hybridization with a specific probe (either on a solid phase carrier or in solution) have been described in the art. The specific probe is preferably a sequence that, under optimized conditions, hybridizes specifically to a region within the 5’ or 3’ flanking region and/or within the mutation region of the specific mutant *IND* allele (hereinafter referred to as “mutant *IND* specific region”). Preferably, the specific probe comprises a sequence of between 10 and 1000 bp, 50 and 600 bp, between 100 to 500 bp, between 150 to 350bp, which is at least 80%, preferably between 80 and 85%, more preferably between 85 and 90%, especially preferably between 90 and 95%, most preferably between 95% and 100% identical (or complementary) to the nucleotide sequence of a specific region. Preferably, the specific probe will comprise a sequence of about 13 to about 100 contiguous nucleotides identical (or complementary) to a specific region of the specific mutant *IND* allele.

Specific probes suitable for the invention may be the following:

- oligonucleotides ranging in length from 13 nt to about 1000 nt, comprising a nucleotide sequence of at least 13 consecutive nucleotides selected from the 5’ or 3’ flanking sequence of a specific mutant *IND* allele or the complement thereof (i.e., for example, the sequence 5’ or 3’ flanking the one or more nucleotides deleted, inserted or substituted in the mutant *IND* alleles of the invention, such as the sequence 5’ or 3’ flanking the non-sense, mis-sense or frameshift mutations described above or the sequence 5’ or 3’ flanking the potential STOP codon mutations indicated in the above Tables or the substitution mutations indicated above), or a sequence having at least 80% sequence identity therewith (probes recognizing 5’ flanking sequences); or
- oligonucleotides ranging in length from 13 nt to about 1000 nt, comprising a nucleotide sequence of at least 13 consecutive nucleotides selected from the mutation sequence of a specific mutant *IND* allele or the complement thereof (i.e., for example, the sequence of

nucleotides inserted or substituted in the *IND* genes of the invention, or the complement thereof), or a sequence having at least 80% sequence identity therewith (probes recognizing mutation sequences).

5 The probes may entirely consist of nucleotide sequence selected from the mentioned nucleotide sequences of flanking and mutation sequences. However, the nucleotide sequence of the probes at their 5' or 3' ends is less critical. Thus, the 5' or 3' sequences of the probes may consist of a nucleotide sequence selected from the flanking or mutation sequences, as appropriate, but may consist of a nucleotide sequence unrelated to the flanking or mutation sequences. Such unrelated
10 sequences should preferably be not longer than 50, more preferably not longer than 25 or even not longer than 20 or 15 nucleotides.

Moreover, suitable probes may comprise or consist of a nucleotide sequence spanning the joining region between flanking and mutation sequences (i.e., for example, the joining region
15 between a sequence 5' or 3' flanking one or more nucleotides deleted, inserted or substituted in the mutant *IND* alleles of the invention and the sequence of the one or more nucleotides inserted or substituted or the sequence 3' or 5', respectively, flanking the one or more nucleotides deleted, such as the joining region between a sequence 5' or 3' flanking non-sense, mis-sense or frameshift mutations in the *IND* genes of the invention described above and the sequence of the
20 non-sense, mis-sense or frameshift mutations, or the joining region between a sequence 5' or 3' flanking a potential STOP codon mutation as indicated in the above Tables or the substitution mutations indicated above and the sequence of the potential STOP codon or substitution mutation, respectively), provided the mentioned nucleotide sequence is not derived exclusively from either the mutation region or flanking regions.

25

Examples of specific probes suitable to identify specific mutant *IND* alleles are described in the Examples.

Detection and/or identification of a "mutant *IND* specific region" hybridizing to a specific probe
30 can occur in various ways, e.g., via size estimation after gel electrophoresis or via fluorescence-based detection methods. Other sequence specific methods for detection of a "mutant *IND* specific region" hybridizing to a specific probe are also known in the art.

Alternatively, plants or plant parts comprising one or more mutant *ind* alleles can be generated
35 and identified using other methods, such as the "Delete-a-geneTM" method which uses PCR to

screen for deletion mutants generated by fast neutron mutagenesis (reviewed by Li and Zhang, 2002, *Funct Integr Genomics* 2:254-258), by the TILLING (Targeting Induced Local Lesions IN Genomes) method which identifies EMS-induced point mutations using denaturing high-performance liquid chromatography (DHPLC) to detect base pair changes by heteroduplex analysis (McCallum *et al.*, 2000, *Nat Biotech* 18:455, and McCallum *et al.* 2000, *Plant Physiol.* 123, 439-442), etc. As mentioned, TILLING uses high-throughput screening for mutations (e.g. using Cel 1 cleavage of mutant-wildtype DNA heteroduplexes and detection using a sequencing gel system). Thus, the use of TILLING to identify plants or plant parts comprising one or more mutant *ind* alleles and methods for generating and identifying such plants, plant organs, tissues and seeds is encompassed herein. Thus in one embodiment, the method according to the invention comprises the steps of mutagenizing plant seeds (e.g. EMS mutagenesis), pooling of plant individuals or DNA, PCR amplification of a region of interest, heteroduplex formation and high-throughput detection, identification of the mutant plant, sequencing of the mutant PCR product. It is understood that other mutagenesis and selection methods may equally be used to generate such mutant plants.

Instead of inducing mutations in *IND* alleles, natural (spontaneous) mutant alleles may be identified by methods known in the art. For example, ECOTILLING may be used (Henikoff *et al.* 2004, *Plant Physiology* 135(2):630-6) to screen a plurality of plants or plant parts for the presence of natural mutant *ind* alleles. As for the mutagenesis techniques above, preferably *Brassica* species are screened which comprise an A and/or a C genome, so that the identified *ind* allele can subsequently be introduced into other *Brassica* species, such as *Brassica napus*, by crossing (inter- or intraspecific crosses) and selection. In ECOTILLING natural polymorphisms in breeding lines or related species are screened for by the TILLING methodology described above, in which individual or pools of plants are used for PCR amplification of the *ind* target, heteroduplex formation and high-throughput analysis. This can be followed by selecting individual plants having a required mutation that can be used subsequently in a breeding program to incorporate the desired mutant allele.

The identified mutant alleles can then be sequenced and the sequence can be compared to the wild type allele to identify the mutation(s). Optionally functionality can be tested as indicated above. Using this approach a plurality of mutant *ind* alleles (and *Brassica* plants comprising one or more of these) can be identified. The desired mutant alleles can then be combined with the desired wild type alleles by crossing and selection methods as described further below. Finally a

single plant comprising the desired number of mutant *ind* and the desired number of wild type *IND* alleles is generated.

5 Oligonucleotides suitable as PCR primers or specific probes for detection of a specific mutant *IND* allele can also be used to develop methods to determine the zygosity status of the specific mutant *IND* allele.

10 To determine the zygosity status of a specific mutant *IND* allele, a PCR-based assay can be developed to determine the presence of a mutant and/or corresponding wild type *IND* specific allele:

15 To determine the zygosity status of a specific mutant *IND* allele, two primers specifically recognizing the wild-type *IND* allele can be designed in such a way that they are directed towards each other and have the mutation region located in between the primers. These primers may be primers specifically recognizing the 5' and 3' flanking sequences, respectively. This set of primers allows simultaneous diagnostic PCR amplification of the mutant, as well as of the corresponding wild type *IND* allele.

20 Alternatively, to determine the zygosity status of a specific mutant *IND* allele, two primers specifically recognizing the wild-type *IND* allele can be designed in such a way that they are directed towards each other and that one of them specifically recognizes the mutation region. These primers may be primers specifically recognizing the sequence of the 5' or 3' flanking region and the mutation region of the wild type *IND* allele, respectively. This set of primers, together with a third primer which specifically recognizes the sequence of the mutation region in the mutant *IND* allele, allow simultaneous diagnostic PCR amplification of the mutant *IND* gene, as well as of the wild type *IND* gene.

30 Alternatively, to determine the zygosity status of a specific mutant *IND* allele, two primers specifically recognizing the wild-type *IND* allele can be designed in such a way that they are directed towards each other and that one of them specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region. These primers may be primers specifically recognizing the 5' or 3' flanking sequence and the joining region between the mutation region and the 3' or 5' flanking region of the wild type *IND* allele, respectively. This set of primers, together with a third primer which specifically recognizes the joining region between the mutation region and the 3' or 5' flanking region of the mutant *IND* allele,

35

respectively, allow simultaneous diagnostic PCR amplification of the mutant *IND* gene, as well as of the wild type *IND* gene.

Alternatively, the zygosity status of a specific mutant *IND* allele can be determined by using
5 alternative primer sets that specifically recognize mutant and wild type *IND* alleles.

If the plant is homozygous for the mutant *IND* gene or the corresponding wild type *IND* gene, the diagnostic PCR assays described above will give rise to a single PCR product typical, preferably typical in length, for either the mutant or wild type *IND* allele. If the plant is
10 heterozygous for the mutant *IND* allele, two specific PCR products will appear, reflecting both the amplification of the mutant and the wild type *IND* allele.

Identification of the wild type and mutant *IND* specific PCR products can occur e.g. by size estimation after gel or capillary electrophoresis (e.g. for mutant *IND* alleles comprising a number
15 of inserted or deleted nucleotides which results in a size difference between the fragments amplified from the wild type and the mutant *IND* allele, such that said fragments can be visibly separated on a gel); by evaluating the presence or absence of the two different fragments after gel or capillary electrophoresis, whereby the diagnostic PCR amplification of the mutant *IND* allele can, optionally, be performed separately from the diagnostic PCR amplification of the wild
20 type *IND* allele; by direct sequencing of the amplified fragments; or by fluorescence-based detection methods.

Examples of primers suitable to determine the zygosity of specific mutant *IND* alleles are described in the Examples.
25

Alternatively, to determine the zygosity status of a specific mutant *IND* allele, a hybridization-based assay can be developed to determine the presence of a mutant and/or corresponding wild type *IND* specific allele:

30 To determine the zygosity status of a specific mutant *IND* allele, two specific probes recognizing the wild-type *IND* allele can be designed in such a way that each probe specifically recognizes a sequence within the *IND* wild type allele and that the mutation region is located in between the sequences recognized by the probes. These probes may be probes specifically recognizing the 5' and 3' flanking sequences, respectively. The use of one or, preferably, both of these probes

allows simultaneous diagnostic hybridization of the mutant, as well as of the corresponding wild type *IND* allele.

Alternatively, to determine the zygosity status of a specific mutant *IND* allele, two specific probes recognizing the wild-type *IND* allele can be designed in such a way that one of them specifically recognizes a sequence within the *IND* wild type allele upstream or downstream of the mutation region, preferably upstream of the mutation region, and that one of them specifically recognizes the mutation region. These probes may be probes specifically recognizing the sequence of the 5' or 3' flanking region, preferably the 5' flanking region, and the mutation region of the wild type *IND* allele, respectively. The use of one or, preferably, both of these probes, optionally, together with a third probe which specifically recognizes the sequence of the mutation region in the mutant *IND* allele, allow diagnostic hybridization of the mutant and of the wild type *IND* gene.

Alternatively, to determine the zygosity status of a specific mutant *IND* allele, a specific probe recognizing the wild-type *IND* allele can be designed in such a way that the probe specifically recognizes the joining region between the 5' or 3' flanking region, preferably the 5' flanking region, and the mutation region of the wild type *IND* allele. This probe, optionally, together with a second probe that specifically recognizes the joining region between the 5' or 3' flanking region, preferably the 5' flanking region, and the mutation region of the mutant *IND* allele, allows diagnostic hybridization of the mutant and of the wild type *IND* gene.

Alternatively, the zygosity status of a specific mutant *IND* allele can be determined by using alternative sets of probes that specifically recognize mutant and wild type *IND* alleles.

25

If the plant is homozygous for the mutant *IND* gene or the corresponding wild type *IND* gene, the diagnostic hybridization assays described above will give rise to a single specific hybridization product, such as one or more hybridizing DNA (restriction) fragments, typical, preferably typical in length, for either the mutant or wild type *IND* allele. If the plant is heterozygous for the mutant *IND* allele, two specific hybridization products will appear, reflecting both the hybridization of the mutant and the wild type *IND* allele.

Identification of the wild type and mutant *IND* specific hybridization products can occur e.g. by size estimation after gel or capillary electrophoresis (e.g. for mutant *IND* alleles comprising a number of inserted or deleted nucleotides which results in a size difference between the

35

hybridizing DNA (restriction) fragments from the wild type and the mutant *IND* allele, such that said fragments can be visibly separated on a gel); by evaluating the presence or absence of the two different specific hybridization products after gel or capillary electrophoresis, whereby the diagnostic hybridization of the mutant *IND* allele can, optionally, be performed separately from the diagnostic hybridization of the wild type *IND* allele; by direct sequencing of the hybridizing DNA (restriction) fragments; or by fluorescence-based detection methods.

Examples of probes suitable to determine the zygosity of specific mutant *IND* alleles are described in the Examples.

Furthermore, detection methods specific for a specific mutant *IND* allele that differ from PCR- or hybridization-based amplification methods can also be developed using the specific mutant *IND* allele specific sequence information provided herein. Such alternative detection methods include linear signal amplification detection methods based on invasive cleavage of particular nucleic acid structures, also known as InvaderTM technology, (as described e.g. in US patent 5,985,557 "Invasive Cleavage of Nucleic Acids", 6,001,567 "Detection of Nucleic Acid sequences by Invader Directed Cleavage, incorporated herein by reference), RT-PCR-based detection methods, such as Taqman, or other detection methods, such as SNPlex. Briefly, in the InvaderTM technology, the target mutation sequence may e.g. be hybridized with a labeled first nucleic acid oligonucleotide comprising the nucleotide sequence of the mutation sequence or a sequence spanning the joining region between the 5' flanking region and the mutation region and with a second nucleic acid oligonucleotide comprising the 3' flanking sequence immediately downstream and adjacent to the mutation sequence, wherein the first and second oligonucleotide overlap by at least one nucleotide. The duplex or triplex structure that is produced by this hybridization allows selective probe cleavage with an enzyme (Cleavase®) leaving the target sequence intact. The cleaved labeled probe is subsequently detected, potentially via an intermediate step resulting in further signal amplification.

A "kit", as used herein, refers to a set of reagents for the purpose of performing the method of the invention, more particularly, the identification of a specific mutant *IND* allele in biological samples or the determination of the zygosity status of plant material comprising a specific mutant *IND* allele. More particularly, a preferred embodiment of the kit of the invention comprises at least two specific primers, as described above, for identification of a specific mutant *IND* allele, or at least two or three specific primers for the determination of the zygosity status. Optionally, the kit can further comprise any other reagent described herein in the PCR

identification protocol. Alternatively, according to another embodiment of this invention, the kit can comprise at least one specific probe, which specifically hybridizes with nucleic acid of biological samples to identify the presence of a specific mutant *IND* allele therein, as described above, for identification of a specific mutant *IND* allele, or at least two or three specific probes for the determination of the zygosity status. Optionally, the kit can further comprise any other reagent (such as but not limited to hybridizing buffer, label) for identification of a specific mutant *IND* allele in biological samples, using the specific probe.

The kit of the invention can be used, and its components can be specifically adjusted, for purposes of quality control (e.g., purity of seed lots), detection of the presence or absence of a specific mutant *IND* allele in plant material or material comprising or derived from plant material, such as but not limited to food or feed products.

The term “primer” as used herein encompasses any nucleic acid that is capable of priming the synthesis of a nascent nucleic acid in a template-dependent process, such as PCR. Typically, primers are oligonucleotides from 10 to 30 nucleotides, but longer sequences can be employed. Primers may be provided in double-stranded form, though the single-stranded form is preferred. Probes can be used as primers, but are designed to bind to the target DNA or RNA and need not be used in an amplification process.

The term “recognizing” as used herein when referring to specific primers, refers to the fact that the specific primers specifically hybridize to a nucleic acid sequence in a specific mutant *IND* allele under the conditions set forth in the method (such as the conditions of the PCR identification protocol), whereby the specificity is determined by the presence of positive and negative controls.

The term “hybridizing”, as used herein when referring to specific probes, refers to the fact that the probe binds to a specific region in the nucleic acid sequence of a specific mutant *IND* allele under standard stringency conditions. Standard stringency conditions as used herein refers to the conditions for hybridization described herein or to the conventional hybridizing conditions as described by Sambrook et al., 1989 (Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbour Laboratory Press, NY) which for instance can comprise the following steps: 1) immobilizing plant genomic DNA fragments or BAC library DNA on a filter, 2) prehybridizing the filter for 1 to 2 hours at 65°C in 6 X SSC, 5 X Denhardt’s reagent, 0.5% SDS and 20 µg/ml denaturated carrier DNA, 3) adding the hybridization probe which has been

labeled, 4) incubating for 16 to 24 hours, 5) washing the filter once for 30 min. at 68°C in 6X SSC, 0.1 %SDS, 6) washing the filter three times (two times for 30 min. in 30ml and once for 10 min in 500ml) at 68°C in 2 X SSC, 0.1 %SDS, and 7) exposing the filter for 4 to 48 hours to X-ray film at -70°C.

5

As used in herein, a "biological sample" is a sample of a plant, plant material or product comprising plant material. The term "plant" is intended to encompass plant tissues, at any stage of maturity, as well as any cells, tissues, or organs taken from or derived from any such plant, including without limitation, any seeds, leaves, stems, flowers, roots, single cells, gametes, cell cultures, tissue cultures or protoplasts. "Plant material", as used herein refers to material that is obtained or derived from a plant. Products comprising plant material relate to food, feed or other products that are produced using plant material or can be contaminated by plant material. It is understood that, in the context of the present invention, such biological samples are tested for the presence of nucleic acids specific for a specific mutant *IND* allele, implying the presence of nucleic acids in the samples. Thus the methods referred to herein for identifying a specific mutant *IND* allele in biological samples, relate to the identification in biological samples of nucleic acids that comprise the specific mutant *IND* allele.

10

15

20

The present invention also relates to the combination of specific *IND* alleles in one plant, to the transfer of one or more specific mutant *IND* allele(s) from one plant to another plant, to the plants comprising one or more specific mutant *IND* allele(s), the progeny obtained from these plants and to plant cells, plant parts, and plant seeds derived from these plants.

25

Thus, in one embodiment of the invention a method for combining two or more selected mutant *IND* alleles in one plant is provided comprising the steps of:

(a) generating and/or identifying two or more plants each comprising one or more selected mutant *IND* alleles, as described above,

(b) crossing a first plant comprising one or more selected mutant *IND* alleles with a second plant comprising one or more other selected mutant *IND* alleles, collecting F1 seeds from the cross, and, optionally, identifying an F1 plant comprising one or more selected mutant *IND* alleles from the first plant with one or more selected mutant *IND* alleles from the second plant, as described above,

30

(c) optionally, repeating step (b) until an F1 plant comprising all selected mutant *IND* alleles is obtained,

35

(d) optionally,

- identifying an F1 plant, which is homozygous or heterozygous for a selected mutant *IND* allele by determining the zygosity status of the mutant *IND* alleles, as described above, or
- generating plants which are homozygous for one or more of the selected mutant *IND* alleles by performing one of the following steps:
 - 5 - extracting doubled haploid plants from treated microspore or pollen cells of F1 plants comprising the one or more selected mutant *IND* alleles, as described above,
 - selfing the F1 plants comprising the one or more selected mutant *IND* allele(s) for one or more generations (y), collecting F1 Sy seeds from the selfings, and identifying F1 Sy plants, which are homozygous for the one or more mutant *IND* allele, as
10 described above.

In another embodiment of the invention a method for transferring one or more mutant *IND* alleles from one plant to another plant is provided comprising the steps of:

- (a) generating and/or identifying a first plant comprising one or more selected mutant *IND* alleles, as described above, or generating the first plant by combining the one or more selected mutant *IND* alleles in one plant, as described above (wherein the first plant is
15 homozygous or heterozygous for the one or more mutant *IND* alleles)
- (b) crossing the first plant comprising the one or more mutant *IND* alleles with a second plant not comprising the one or more mutant *IND* alleles, collecting F1 seeds from the cross
20 (wherein the seeds are heterozygous for a mutant *IND* allele if the first plant was homozygous for that mutant *IND* allele, and wherein half of the seeds are heterozygous and half of the seeds are azygous for, i.e. do not comprise, a mutant *IND* allele if the first plant was heterozygous for that mutant *IND* allele), and, optionally, identifying F1 plants comprising one or more selected mutant *IND* alleles, as described above,
- 25 (c) backcrossing F1 plants comprising one or more selected mutant *IND* alleles with the second plant not comprising the one or more selected mutant *IND* alleles for one or more generations (x), collecting BCx seeds from the crosses, and identifying in every generation BCx plants comprising the one or more selected mutant *IND* alleles, as described above,
- 30 (d) optionally, generating BCx plants which are homozygous for the one or more selected mutant *IND* alleles by performing one of the following steps:
 - extracting doubled haploid plants from treated microspore or pollen cells of BCx plants comprising the one or more desired mutant *IND* allele(s), as described above,
 - selfing the BCx plants comprising the one or more desired mutant *IND* allele(s) for one or more generations (y), collecting BCx Sy seeds from the selfings, and identifying BCx

Sy plants, which are homozygous for the one or more desired mutant *IND* allele, as described above.

In one aspect of the invention, the first and the second plant are *Brassicaceae* plants, particularly
5 *Brassica* plants, especially *Brassica napus* plants or plants from another *Brassica* crop species.
In another aspect of the invention, the first plant is a *Brassicaceae* plant, particularly a *Brassica*
plant, especially a *Brassica napus* plant or a plant from another *Brassica* crop species, and the
second plant is a plant from a *Brassicaceae* breeding line, particularly from a *Brassica* breeding
10 *Brassica* crop species. "Breeding line", as used herein, is a preferably homozygous plant line
distinguishable from other plant lines by a preferred genotype and/or phenotype that is used to
produce hybrid offspring.

In yet another embodiment of the invention, a method for making a plant, in particular a
15 *Brassica* crop plant, such as a *Brassica napus* plant, of which the pod shatter resistance is
increased but which preferably maintains an agronomically relevant treshability of the pods is
provided comprising combining and/or transferring mutant *IND* alleles according to the
invention in or to one *Brassica* plant, as described above.

20 In one aspect of the invention, the plant is a *Brassica* plant comprising at least two *IND* genes
wherein pod shatter resistance is increased while maintaining an agronomically relevant
treshability of the pods by combining and/or transferring three mutant *IND* alleles according to
the invention in or to the *Brassica* plant, as described above.

25 In still another embodiment of the invention, a method for making a hybrid *Brassica* crop seed or
plant comprising at least two *IND* genes, in particular a hybrid *Brassica napus* seed or plant, of
which the pod shatter resistance is increased but which maintains an agronomically relevant
treshability of the pods is provided, comprising the steps of:

- (a) generating and/or identifying a first plant comprising a first and a second selected mutant
30 *IND* allele in homozygous state and a second plant comprising a third selected mutant *IND*
allele in homozygous state, as described above,
- (b) crossing the first and the second plant and collecting F1 hybrid seeds from the cross.

In one aspect of the invention, the first or the second selected mutant *IND* allele is the same
35 mutant *IND* allele as the third selected mutant *IND* allele, such that the F1 hybrid seeds are

homozygous for one mutant *IND* allele and heterozygous for the other. In another aspect of the invention, the first plant is used as a male parent plant and the second plant is used as a female parent plant. In one embodiment of the invention, the first plant is completely pod shatter resistant. Such plants may be obtained by sowing complete indehiscent seed pods obtained by selfing the plants and harvesting complete seed pods in stead of thrashing the seed pods to harvest the seeds.

SEQUENCES

10 *IND* genes

SEQ ID NO: 1: Coding DNA of the *IND-A1* gene encoding a wild-type IND-A1 protein from *Brassica napus*.

SEQ ID NO: 2: wild type IND-A1 protein encoded by SEQ ID NO: 1.

SEQ ID NO: 3: Coding DNA of the *IND-C1* gene encoding a wild-type IND-C1 protein from *Brassica napus*.

SEQ ID NO: 4: wild type IND-C1 protein encoded by SEQ ID NO: 3.

SEQ ID NO: 5: Genomic DNA of the *IND-A1* gene encoding a wild-type IND-A1 protein from *Brassica napus*.

SEQ ID NO: 6: wild type IND-A1 protein encoded by SEQ ID NO: 5.

20 SEQ ID NO: 7: Genomic DNA of the *IND-C1* gene encoding a wild-type IND-C1 protein from *Brassica napus*.

SEQ ID NO: 8: wild type IND-C1 protein encoded by SEQ ID NO: 7.

SEQ ID NO: 9: Coding DNA of the *Arabidopsis IND1* gene.

SEQ ID NO: 10: *Arabidopsis* IND1 protein encoded by SEQ ID NO: 9.

25 SEQ ID NO: 11: nucleotide sequence of an *IND* homologue from *Brassica napus* (BN1-*IND* - SEQ ID NO: 2 of WO04/113542)

SEQ ID NO: 12: nucleotide sequence of a second *IND* homologue from *Brassica napus* (BN2-*IND* - SEQ ID NO: 3 of WO04/113542)

30 *Primers and probes*

SEQ ID NO 13: Forward oligonucleotide for detection of IND-A1-EMS01

SEQ ID NO 14: Forward oligonucleotide for detection of IND-A1-WT

SEQ ID NO 15: Reverse oligonucleotide for detection of IND-A1-EMS01 and -WT

SEQ ID NO 16: Forward oligonucleotide for detection of IND-A1-EMS05

35 SEQ ID NO 17: Forward oligonucleotide for detection of IND-A1-WT

- SEQ ID NO 18: Reverse oligonucleotide for detection of IND-A1-EMS05 and -WT
SEQ ID NO 19: Reverse oligonucleotide for detection of IND-C1-EMS01
SEQ ID NO 20: Reverse oligonucleotide for detection of IND-C1-WT
SEQ ID NO 21: Forward oligonucleotide for detection of IND-C1-EMS01 and -WT
5 SEQ ID NO 22: Reverse oligonucleotide for detection of IND-C1-EMS03
SEQ ID NO 23: Reverse oligonucleotide for detection of IND-C1-WT
SEQ ID NO 24: Forward oligonucleotide for detection of IND-C1-EMS03 and -WT
SEQ ID NO 25: Oligonucleotide for detection of IND-A1-EMS01 and -WT
SEQ ID NO 26: Oligonucleotide for detection of IND-A1-EMS01
10 SEQ ID NO 27: Oligonucleotide for detection of IND-A1-WT
SEQ ID NO 28: Oligonucleotide for detection of IND-A1-EMS05 and -WT
SEQ ID NO 29: Oligonucleotide for detection of IND-A1-EMS05
SEQ ID NO 30: Oligonucleotide for detection of IND-A1-WT
SEQ ID NO 31: Oligonucleotide for detection of IND-C1-EMS01 and -WT
15 SEQ ID NO 32: Oligonucleotide for detection of IND-C1-EMS01
SEQ ID NO 33: Oligonucleotide for detection of IND-C1-WT
SEQ ID NO 34: Oligonucleotide for detection of IND-C1-EMS03 and -WT
SEQ ID NO 35: Oligonucleotide for detection of IND-C1-EMS03
SEQ ID NO 36: Oligonucleotide for detection of IND-C1-WT
20 SEQ ID NO 37: Forward oligonucleotide for detection of IND-A1
SEQ ID NO 38: Reverse oligonucleotide for detection of IND-A1
SEQ ID NO 39: Forward oligonucleotide for detection of IND-C1
SEQ ID NO 40: Reverse oligonucleotide for detection of IND-C1

25 Unless stated otherwise in the Examples, all recombinant DNA techniques are carried out according to standard molecular biological techniques as described in Sambrook and Russell (2001) *Molecular Cloning: A Laboratory Manual*, Third Edition, Cold Spring Harbor Laboratory Press, NY, in Volumes 1 and 2 of Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*,
Current Protocols, USA and in Volumes I and II of Brown (1998) *Molecular Biology LabFax*,
30 Second Edition, Academic Press (UK). Standard materials and methods for plant molecular work are described in *Plant Molecular Biology Labfax* (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK. Standard materials and methods for polymerase chain reactions can be found in Dieffenbach and Dveksler (1995) *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, and in
35 McPherson *et al.* (2000) *PCR - Basics: From Background to Bench*, First Edition, Springer

Verlag, Germany. Standard procedures for AFLP analysis are described in Vos *et al.* (1995, NAR 23:4407-4414) and in published EP patent application EP 534858.

EXAMPLES

5

Example 1 - Isolation of the DNA sequences of the *IND* genes

To determine the sequences of the *IND* genes of an elite spring oilseed rape breeding line, a Bacterial Artificial Chromosome (BAC) library of the line was screened as follows:

10 1.1. Isolation of BAC clones comprising an *IND* sequence

To identify *Escherichia coli* colonies containing a BAC clone comprising an *IND* sequence of the elite spring oilseed rape breeding line, a BAC library of the line (average clone size of more than 120 kb) arrayed as individual duplicated clones on high density nylon filters were screened by standard Southern hybridization procedures:

- 15
- A mixture of two probes with the sequence of SEQ ID NO: 2 of WO04/113542 (“Bn1-IND”) and SEQ ID NO: 3 of WO04/113542 (“BN2-IND”) (SEQ ID NO: 11 and 12, respectively) and labeled according to standard procedures were used for hybridizing to the DNA on the nylon membrane.
 - Pre-hybridization was performed for 2 hour at 65°C in 30 ml of the following hybridization

20 buffer: 6X SSC (20X SSC contains 3.0 M NaCl, 0.3 M Na citrate, pH 7.0), 5X Denhardt's (100X Denhardt's contains 2% Ficoll, 2% Polyvinyl pyrrolidone, 2% Bovine Serum Albumin), 0.5% SDS and 20 µg/ml denaturated carrier DNA (single-stranded fish sperm DNA, with an average length of 120 - 3000 nucleotides)
 - Hybridization was performed under the following conditions:

25

 - The labeled probe (20 ng of each sequence) was denaturated by heating for 5 minutes at 95°C and chilling on ice for 5 minutes and added to 15 ml of hybridization buffer (same buffer as for the pre-hybridization)
 - The hybridization was performed overnight at 65°C.
 - The blots were washed three times for 30 minutes at 65°C in the hybridization tubes (once

30 with 30ml 6xSSC with 0.1% SDS and twice with 30ml 2xSSC with 0.1% SDS) and one time for 10 minutes at 65°C with 500ml 2xSSC with 0.1% SDS in a box.
 - Kodak X-OMAT AR films were exposed to the radioactive blots for 4 hours at -70°C.
 - Based on the positive signals, 14 *E. coli* colonies containing a BAC clone comprising an *IND* sequence were picked up by screening the BAC library from the elite spring oilseed rape

35 breeding line (total n° of positives: 65) (hereinafter called “positive colonies”).

1.2. Isolation of BAC clones comprising a full-length *IND* sequence

To identify positive colonies comprising a BAC clone with a full-length genomic DNA sequence of one of the *IND* genes, a Southern blot analysis was performed on BAC clone DNA isolated from the positive colonies and on genomic DNA isolated from *Brassica napus*:

- 5 - BAC clone DNA was isolated through alkaline lysis as described in the art from the positive colonies grown up in 25 ml Luria Broth medium containing 25µg/ml chloramphenicol.
- Genomic DNA was isolated from leaf tissue of *B. napus* according to the cetyltrimethylammoniumbromide (CTAB) method (Doyle and Doyle, 1987, Phytochemistry
10 Bulletin 19:11-15).
- The DNA concentration of each preparation was estimated by comparing the band intensity of 1 µl of each sample to the band intensity of 1, 2, 4, 8 and 20 µl of a solution containing 25 ng/µl Lambda DNA (Life Technologies®) on a 1% TBE (Invitrogen®) agarose gel (Roche®) containing ethidiumbromide (ICN Biochemicals®).
- 15 - 100-200 ng of BAC clone DNA and 1,7 µg genomic DNA were digested with restriction enzyme EcoRI in a final reaction volume of 20 µl, applying conditions proposed by the manufacturer (New England Biolabs). The time of digestion and/or amount of restriction enzyme were adjusted to ensure complete digestion of the genomic DNA samples without non-specific degradation.
- 20 - After digestion, 2 µl of loading dye containing RNase (12,5 ml 1% xylene cyanol FF; 12,5 ml 1% bromophenol blue water soluble indicator; 25 ml glycerol; 100 µl 0.5M EDTA pH8; 1µl RNase (10mg/ml)) was added to the digested DNA samples and the samples were incubated for 30 min at 37°C.
- The samples were loaded on a 1% TAE agarose gel.
- 25 - Phage Lambda DNA (Fermentas®) digested with PstI or 1kbp DNA Ladder (Life Technologies) was included as size standard.
- After electrophoresis, the DNA samples (digested BAC clone and genomic DNA) were transferred to a nylon membrane (Hybond-N+ Amersham Pharmacia Biotech®) by dry alkali capillary blotting.
- 30 - The nylon membranes with digested BAC clone and genomic DNA were screened by standard Southern hybridization procedures as described above for the BAC library screenings, except that for the genomic DNA the Kodak XOMAT AR films were exposed to the radioactive blots for 2 days at -70°C.
- Based on a comparison between the hybridization patterns obtained after digestion of BAC
35 clone DNA of the identified positive colonies and of genomic DNA isolated from *Brassica*

napus with restriction enzyme EcoRI and hybridization with the probes, the BAC clones were grouped in 2 groups and for each of the 2 groups a BAC clone was selected containing a full-length *IND* sequence (named *IND-A1* and *IND-C1*).

- 5 - The *IND* sequences comprised in the BAC clones of the selected positive colonies were determined by standard sequencing techniques (Agowa).

Table 3: Hybridization pattern of digested BAC clone and genomic DNA hybridized to the *Bn1*- and *Bn2-IND* probes

DNA sample:	Genomic DNA from <i>B. napus</i>	BAC clone DNA from <i>B. napus</i>	Corresponds to
restricted with:	Estimated length of the hybridizing DNA fragments:		
EcoRI	8 kb 2.2 kb	8 kb 2.2 kb	<i>IND-A1</i> <i>IND-C1</i>

Example 2 - Characterization of *IND* gene sequences from *Brassica napus*

10 After sequencing the genomic DNA fragments (SEQ ID NO: 5 and 7, respectively), the coding regions of the *IND* sequences were determined with FgeneSH (Softberry, Inc. Mount Kisco, NY, USA) and est2genome (Rice *et al.*, 2000, Trends in Genetics 16 (6): 276—277; Mott, 1997, Comput. Applic. 13:477-478) as depicted in the sequence listing.

15 Comparison of hybridizing bands generated in a Southern blot analysis on genomic DNA isolated from *B. rapa* (AA), *B. oleracea* (CC) and *B. napus* (AACC) and on BAC clone DNA isolated from the positive colonies identified in Example 1 (restricted with EcoRI and hybridized to probe as described in Example 1) indicated that the *IND-A1* sequence originated from the A genome and the *IND-C1* sequence from the C genome.

20

The protein encoding regions of the *IND* genes of the elite spring oilseed rape breeding line are represented in SEQ ID NO:1 (*IND-A1*), SEQ ID NO:3 from the nucleotide at position 46 to the nucleotide at position 633 (*IND-C1-short*) and SEQ ID NO:3 (*IND-C1-long*), respectively. The, by these nucleic acid sequence encoded, *IND-A1* and *IND-C1* protein sequences are depicted in
25 SEQ ID NO:2 (*IND-A1*), SEQ ID NO:4 from the amino acid at position 16 to the amino acid at position 210 (*IND-C1-short*) and SEQ ID NO:4 (*IND-C1-long*), respectively.

The percentage (nucleotide) sequence identity between the complete coding regions of *IND-A1* and *IND-C1-long* is 81% and between the complete coding regions of *IND-A1* and *IND-C1-short*
30 is 87%, while the percentage (nucleotide) sequence identity between the regions encoding the

bHLH domains of *IND-A1* and *IND-C1-long* and *-short* (as determined according to Toledo-Ortiz *et al.*, 2003, Plant Cell 15, 1749-1770) is 98%. These percentages indicate that the *IND* genes are more conserved in the region encoding the bHLH domain than in the remaining part of the coding region.

5

Similarly, the percentage (amino acid) sequence identity between the complete *IND-A1* and *IND-C1-long* proteins is 75% and between the complete *IND-A1* and *IND-C1-short* proteins is 80%, while the percentage (amino acid) sequence identity between the bHLH domains of *IND-A1* and *IND-C1-long* and *-short* (as determined according to Toledo-Ortiz *et al.*, 2003, Plant Cell 15, 1749-1770) is 98%. These percentages indicate that the *IND* proteins are more conserved in the bHLH domain than in the remaining part of the *IND* proteins.

10

Example 3 - Expression of *Brassica IND* genes

To analyze the expression of the different *IND* genes in different tissues, RT-PCR assays specific for each *IND* gene were performed on total RNA isolated from *Brassica napus* leaves, pod walls, dehiscence zone tissue and seeds using the following primers:

15

INDA1F1 5' AGGAGAGGAAGAGATGGATCC 3' (SEQ ID No. 37)

INDA1R1 5' TGAGTGTGAGGCTGAAGAAGC 3' (SEQ ID No. 38)

for the *IND-A1* gene, and

20

INDC1F1 5' CCTCATCATCTCCTTATGAAC 3' (SEQ ID No. 39)

INDC1R 5' CGTATTGCATCTCCTTCATCT 3'. (SEQ ID No. 40)

for the *IND-C1* gene.

The results indicated that both *IND* genes, i.e. *IND-A1* and *IND-C1*, were not expressed in leaf tissue and seeds, but were expressed in dehiscence zone tissue and that the *IND-A1* gene was expressed in pod walls, while the *IND-C1* gene was not expressed in pod walls.

25

Example 4 - Generation and isolation of mutant *IND* alleles (*ind*)

Mutations in the *IND* genes identified in Example 1 were generated and identified as follows:

- 30,000 seeds from an elite spring oilseed rape breeding line (M0 seeds) were preimbibed for two hours on wet filter paper in deionized or distilled water. Half of the seeds were exposed to 0.8% EMS and half to 1% EMS (Sigma: M0880) and incubated for 4 hours.
- The mutagenized seeds (M1 seeds) were rinsed 3 times and dried in a fume hood overnight. 30,000 M1 plants were grown in soil and selfed to generate M2 seeds. M2 seeds were harvested for each individual M1 plant.

30

- Two times 4800 M2 plants, derived from different M1 plants, were grown and DNA samples were prepared from leaf samples of each individual M2 plant according to the CTAB method (Doyle and Doyle, 1987, Phytochemistry Bulletin 19:11-15).
- The DNA samples were screened for the presence of point mutations in the *IND* genes causing the introduction of STOP codons in the protein-encoding regions of the *IND* genes or the substitution of amino acids in the *IND* proteins, particularly in the bHLH domain of the *IND* proteins, by direct sequencing by standard sequencing techniques (Agowa) and analyzing the sequences for the presence of the point mutations using the NovoSNP software (VIB Antwerp).
- 10 - The following mutant *IND* alleles (*ind*) were thus identified:

Table 4a: STOP codon and substitution mutations in *IND-A1*

Amino acid position	Nucleotide position		Wild type → mutant codon	Wild type → mutant amino acid	M2 Plant No.	Allele No.
	<i>SEQ ID: 2/6</i>	<i>SEQ ID: 1</i>				
122	364	924	<u>c</u> ag → <u>t</u> ag	GLN → STOP (in b)	POSH101, POSH102, POSH103, POSH104	<i>ind-al-EMS01</i> , <i>ind-al-EMS02</i> , <i>ind-al-EMS03</i> , <i>ind-al-EMS04</i>
103	307	867	gat → <u>a</u> at	ASP → ASN	POSH105	<i>ind-al-EMS05</i>
127	380	940	<u>c</u> gt → <u>c</u> at	ARG → HIS (in b)	POSH105	<i>ind-al-EMS05</i>

Table 4b: STOP codon mutations in *IND-C1*

Amino acid position	Nucleotide position		Wild type → mutant codon	Wild type → mutant amino acid	M2 Plant No.	Allele No.
	<i>SEQ ID: 4/8</i>	<i>SEQ ID: 3</i>				
50	148	644	<u>c</u> aa → <u>t</u> aa	GLN → STOP	POSH106	<i>ind-cl-EMS01</i>
135	403	899	<u>c</u> ag → <u>t</u> ag	GLN → STOP (in b)	POSH108	<i>ind-cl-EMS03</i>

- 15 Reference seeds of plants comprising alleles *ind-al-EMS01* and *ind-cl-EMS01* in homozygous state have been deposited at the American Type Culture Collection (ATCC, 10801 University Boulevard, Manassas, VA 20110-2209, US) on November 20, 2007, under accession number PTA-8796 (strain designation 07MBBN001171) and reference seeds of plants comprising alleles *ind-al-EMS05* and *ind-cl-EMS03* in homozygous state have been deposited at the ATCC on
- 20 November 20, 2007, under accession number PTA-8795 (strain designation 07MBBN000530).

In conclusion, the above examples show how mutant *IND* alleles can be generated and isolated. Also, plant material comprising such mutant alleles can be used to combine selected mutant and/or wild type alleles in a plant, as described in the following examples.

5 **Example 5 - Identification of a *Brassica* plant comprising a mutant *Brassica IND* allele**

Brassica plants comprising the mutations in the *IND* genes identified in Example 4 were identified as follows:

- For each mutant *IND* gene identified in the DNA sample of an M2 plant, at least 50 M2 plants derived from the same M1 plant as the M2 plant comprising the *IND* mutation were grown and DNA samples were prepared from leaf samples of each individual M2 plant.
- The DNA samples were screened for the presence of the identified point *IND* mutation as described above in Example 4.
- Heterozygous and homozygous (as determined based on the electropherograms) M2 plants comprising the same mutation were selfed and M3 seeds were harvested.

15 **Example 6 - Analysis of the fruit dehiscence properties of *Brassica* plants comprising a mutant *Brassica IND* gene**

To determine the correlation between the presence of mutant *IND* genes in *Brassica* plants and the fruit dehiscence properties of the *Brassica* plants, the fruit dehiscence properties of *Brassica* plants comprising a mutant *IND* gene were analyzed in the glass house and in the field as follows:

- To examine whether and how the fruit valve margins and the dehiscence properties of seed pods were affected by mutations in *IND*, *ind* fruit was compared to wild-type fruit using the following macroscopic tests:

- (a) Inspection of the seed pods and plants in general with naked eye to determine differences in the phenotype of the pods and plants caused by the presence of certain mutant *IND* alleles. Determination of the phenotype of the pods: When the pods were fully grown and filled, just prior to yellowing, the degree of sharpness of the zone that delineates the valve and beak at the zone where both valves are not touching anymore (at distal end of pod) of 5 random pods (from different plants if multiple plants per line are available) was assessed and attributed a score from 1 to 5: 1 for a clear indentation and fine sharp zone that separates valve and beak; 2 for some indentation and clear, though more fuzzy, zone that separates valve from beak; 3 for valves and beak that are still well observable as two different tissues but with a very smooth transition between them; 4 for valves and beak that are barely observable as different tissues; 5 for a completely smoothed transition

between valves and beak without any clear differentiation between both tissue types, i.e. the less indentation between the valve and the beak at the distal end of the pods the higher the score. A score of 1 (sharp indentation between the valve and the beak) corresponds to a wildtype phenotype of the pods, more specifically a pod shatter sensitive phenotype of the pods; a score of 2 to 4 (more gradual transition between the valve and the beak) corresponds to a pod shatter resistant phenotype of the pods, wherein seed shattering is significantly reduced or delayed while an agronomically relevant treshability of the pods is maintained, such that the pods may still be opened along the dehiscence zone by applying limited physical forces; and a score of 5 (no indentation between the valve and the beak) corresponds to a pod shatter resistant phenotype of the pods, wherein seed shattering is reduced or delayed to a degree which does not allow an agronomically relevant treshability of the pods anymore, such that the pods cannot be opened along the dehiscence zone by applying limited physical forces.

(b) Manual Impact Test (MIT) to determine the increase in pod shatter resistance caused by the presence of certain mutant *IND* alleles: The level of pod shatter resistance of *Brassica* lines comprising the mutant *IND* alleles and *Brassica* lines comprising the corresponding wild type *IND* alleles was compared in a semi-quantitative way by determining the physical forces needed to open closed mature pods by manually applying torsion on the pods. A distinction was made between pods which completely open along the dehiscence zone at the slightest torsion, pods which open only at the base of the dehiscence zone and need stronger torsion to open completely and pods which can only be crushed and do not open along the dehiscence zone. The pod shatter resistance of the pods was attributed a score from 1 to 5 based on this physical force: 1 for pods which completely open along the dehiscence zone at the slightest torsion, 2-4 for pods which open only at the base of the dehiscence zone and need stronger torsion to open completely and 5 for pods which can only be crushed and do not open along the dehiscence zone.

(c) Random Impact Test (RIT) to determine the increase in pod shatter resistance caused by the presence of certain mutant *IND* alleles: The level of pod shatter resistance of *Brassica* lines comprising the mutant *IND* alleles and *Brassica* lines comprising the corresponding wild type *IND* alleles was compared in a quantitative way by determining the half life of samples of pods from both lines according to Bruce *et al.* (2002, *supra*). More specifically, two replicate samples of 20 intact mature pods from each line were subjected to a RIT. 20 pods were placed together with six steel balls of 12.5 mm diameter in a cylindrical container of diameter 20 cm with its axis vertical. The container

was then subjected to simple harmonic motion of frequency 4.98 Hz and of stroke 51 mm in the horizontal plane. The pods, checked for soundness before the test, were shaken for cumulative times of 10, 20, 40, and, if more than 50% of pods remained intact, 80s. The drum was opened after each period and the number of closed pods counted. The pods were examined and classed as "closed" if the dehiscence zone of both valves was still closed. Thus the pods were classed as "opened" if one or both of the valves was detached, so that the seed had been released. If the majority of the pods was broken or damaged without opening of the dehiscence zone, the sample was marked "uncountable". To give each point equal weighing, the data were made evenly spaced in the independent variable, time, by adding 1 and taking \log_{10} . The percentage of pods opened p was transformed by the logit transformation, i.e. $\text{logit } p = \log_e(p/100-p)$. A linear model was then fitted to the transformed time and percentage data and used to estimate the half-life.

- (d) Field tests to determine the relationship between pod shatter resistance, treshability and yield and the presence of certain mutant *IND* alleles in plants: The level of pod shatter resistance, treshability and yield of *Brassica* lines comprising the mutant *IND* alleles and *Brassica* lines comprising the corresponding wild type *IND* alleles was compared in a semi-quantitative way by determining and comparing the level of seed shattering (SHAT), combiner harvest ability (CHA1) and treshing ability (CHA2) and in a quantitative way by determining and comparing seed yield per plot after combining (YLDP) and seed yield after treshing of straw (YLDS) in the field between plots with *ind* plants and plots with wild-type plants. The plots were attributed a score of 1-9 to indicate the level of seed shattering on the plot before harvest: a score of 1 to indicate that practically all plants on the plot were shattering before harvest to a score of 9 to indicate that practically no plants on the plot were shattering before harvest. The plots were attributed a score of 1-5 to indicate the level of combiner harvest ability on the plot: a score of 1, to 3 or to 5 to indicate that it was difficult, to feasible, or to easy, respectively, to harvest the plot with a combiner. The plots were attributed a score of 1-5 to indicate the level of treshing ability of the plot: a score of 1, to 3 or to 5 to indicate that it was difficult, to feasible, or to easy, respectively, to manually harvest the seed remaining in the straw after combiner harvest. The seed yield per plot after combining (YLDP; expressed in grams per plot) was determined by harvesting the seeds per plot with a combine harvester and weighing the seeds and the seed yield after treshing of straw (YLDS; expressed in weight% of the straw) was determined by manually harvesting the seeds remaining in the straw after seed harvest with the combine harvester.

- To examine more closely whether and how cells at the valve margin of seed pods are affected by mutations in *IND*, sections of *ind* fruit were compared to sections of wild-type fruit by microscopic evaluation of the seed pods:
 - Explants: Explants of about 3 mm taken from the proximal and distal ends of pods of similar developmental stage (about 35 days after anthesis (DAA), a stage of development which closely corresponds to the onset of visible pericarp yellowing) and size were harvested from plants grown in a plant growth room (two pods for each genotype) and/or in the field. Both dehiscence zones were dissected from the pods.
 - Fixation: Fixation was done in 100mM K-phosphate buffer pH7 with 10% formalin and 0.25% glutaraldehyde for a total of 4 hours. Vacuum infiltration was done after 1 and 2 hours for 15 minutes. The fixative was renewed after each vacuum infiltration.
 - Dehydration: The specimen was rinsed 2 times 30 minutes with 100mM K-phosphate buffer pH7. Dehydration was done with technical ethanol diluted with 0.85% NaCl in water: 60 minutes (‘) in 50% ethanol, 90’ in 70% ethanol, 90’ in 80% ethanol, 90’ in 90% ethanol, 90’ in 95% ethanol, 90’ in 100% ethanol at room temperature
 - Embedding: Embedding was done with The Leica 7022-31731 Histo-resin or the Kulzer Histo-Technik 7100 (Heraeus) embedding kits, which are three component resin (a basic resin, an activator and a hardener) kits. The three components were used in the proportions as advised by the manufacturer as follows: the specimen were incubated for 4 hours in 50% ethanol/50% basic resin, overnight in 30% ethanol/70% basic resin (optional: at 4°C), for 2 to 4 hours in 100% basic resin, for one day in 100% basic resin after renewing the basic resin and vacuum infiltration for 20’ (optionally at 4°C), for one day in basic resin + activator (1%) (“infiltration medium”) after vacuum infiltration in this medium for 20 minutes. The specimen was washed with basic resin + activator (1%) + hardener (1 ml in 15 ml) (“embedding medium”). The embedding was done in flat embedding moulds (AGAR flat embedding moulds G3531 with cavities of about 300µl: 14 mm long x 6 mm wide x 4 mm deep): 100-125 µl of embedding medium/cavity was added, the embedding medium was polymerized at 55°C for about one hour, the tissue was put on the polymerized embedding medium (1 explant/cavity), the cavities were filed with embedding medium, the embedding medium was polymerized for 3 to 5 hours at 55°C, the moulds were could down, the plastic blocks were removed from the moulds and stored at room temperature in a sealed container (e.g. eppendorf tube).
 - Sectioning: The plastic blocks were glued with the flat side on a 1cm³ perspex block and trimmed squarely around the specimen. 4 µm sections (3 to 4 explants per genotype, about 25 sections per explant) were cut with a ralph glass knife (made on -1 position of

the histoknifemaker of Reichert-Jung using 6 mm thick glass rods under a cutting angle of about 6°) on the microtome. The sections were attached on glass slides treated with Vectabond (Vector laboratories).

- Demonstration of lignin: unstained sections mounted in Eukitt were examined using a microscope equipped for fluorescence (with Zeiss filter set 02). Lignin fluoresces clear bluish
- Evaluation of histology: unstained sections were visualized by using DIC-Normaski or autofluorescence (with Zeiss filter set 18 -- Excitation BP390-420; Emission LP450).

6.1. Correlation between the presence of one or two mutant *Brassica IND* alleles in *Brassica* plants and the fruit dehiscence properties of those *Brassica* plants

To determine the correlation between the presence of one *ind* in heterozygous state (genotype: *IND-A1/ind-a1*, *IND-C1/IND-C1*; or *IND-A1/IND-A1*, *IND-C1/ind-c1*) or in homozygous state (genotype: *ind-a1/ind-a1*, *IND-C1/IND-C1* or *IND-A1/IND-A1*, *ind-c1/ind-c1*) in a *Brassica* plant and the fruit dehiscence properties of the *Brassica* plant, the fruit dehiscence properties of *Brassica* plants identified in Example 5 (in particular homozygous M2 plants No. POSH101, POSH103, POSH104, POSH105 and POSH106 and heterozygous M2 plants No. POSH105; see Table 4a and b for the corresponding *ind* alleles) were grown in the glass house and analyzed as described above. No significant difference in phenotype and fruit dehiscence properties was observed between wild type plants and these heterozygous and homozygous single mutant plants.

Field tests with homozygous single *ind* mutant (genotype: *ind-a1/ind-a1*, *IND-C1/IND-C1* or *IND-A1/IND-A1*, *ind-c1/ind-c1*) and wild type plants (genotype: *IND-A1/IND-A1*, *IND-C1/IND-C1*) from segregating backcross 3 (BC3) populations showed however an increase in seed yield for the homozygous single *ind* mutant plants (see Table below).

Genotype	SHAT (1-9)	CHA1 (1-5)	CHA2 (1-5)	YLDP (in g/plot)	YieldWTSeg%	YLDS (in wt% of straw)
<i>ind-a1-01/ind-a1-01</i> , <i>IND-C1/IND-C1</i>	8.0	4.9	5.0	2636.0	106	0.8
<i>IND-A1/IND-A1</i> , <i>IND-C1/IND-C1</i>	7.8	4.9	5.0	2490.0	100	0.7
<i>ind-a1-05/ind-a1-05</i> , <i>IND-C1/IND-C1</i>	8.1	4.8	5.0	2450.9	103	0.3
<i>IND-A1/IND-A1</i> ,	7.6	5.0	4.8	2387.6	100	0.4

<i>IND-C1/IND-C1</i>						
<i>IND-A1/IND-A1, ind-c1-01/ind-c1-01</i>	8.3	4.9	5.0	2856.0	113	0.6
<i>IND-A1/IND-A1, IND-C1/IND-C1</i>	8.3	4.8	5.0	2517.3	100	0.3
<i>IND-A1/IND-A1, ind-c1-03/ind-c1-03</i>	8.6	4.7	4.9	2833.6	113	0.5
<i>IND-A1/IND-A1, IND-C1/IND-C1</i>	8.1	4.6	5.0	2510.7	100	0.4

6.2. Correlation between the presence of at least three mutant *Brassica* *IND* alleles in *Brassica* plants and the fruit dehiscence properties of those *Brassica* plants

To determine the correlation between the presence of at least three mutant *IND* alleles in a *Brassica* plant and the fruit dehiscence properties of the *Brassica* plant, the *Brassica* plants identified in Example 5, and/or progeny thereof, comprising the mutant *IND* alleles, were crossed with each other and the fruit dehiscence properties of the progeny *Brassica* plants was analyzed as described above.

10 Plant material:

Progeny (i.e., homozygous double mutant plants with genotype *ind-a1/ind-a1, ind-c1/ind-c1*; homozygous single and heterozygous single –i.e., triple- mutant plants with genotype *ind-a1/ind-a1, IND-C1/ind-c1* and *IND-A1/ind-a1, ind-c1/ind-c1*; and wild type plants with genotype *IND-A1/IND-A1, IND-C1/IND-C1*) of line 51, line 45, line 176 and line 48, which themselves are heterozygous (genotype: *IND-A1/ind-a1, IND-C1/ ind-c1*) for alleles *IND-A1-EMS01* and *IND-C1-EMS01* (line 51), alleles *IND-A1-EMS01* and *IND-C1-EMS03* (line 45), alleles *IND-A1-EMS05* and *IND-C1-EMS01* (line 176), and alleles *IND-A1-EMS05* and *IND-C1-EMS03* (line 48), respectively.

20 Macroscopical evaluation:

a) Inspection of the seed pods and plants with naked eye.

- The pods from double homozygous mutant *IND* sibling plants (genotype: *ind-a1/ind-a1, ind-c1/ind-c1*) derived from lines 51, 45, 176 and 48 showed an altered pod morphology, already at immature stage, as compared to pods from wild-type *IND* sibling plants. More specifically, the pods of the double homozygous mutant *IND* sibling plants showed a lack of proper valve margin definition, particularly apparent at both the proximal and distal end of the fruit, as compared to the pods from wild-type *IND* sibling plants, which showed clearly defined margins. Furthermore, the sharp indentation between the valve and the beak at the distal end of the pods in the wild-type sibling plants was largely absent in the double

homozygous *ind* sibling plants, which showed a more gradual transition between valve and beak tissue. The flowers of the double homozygous mutant *IND* sibling plants of line 51 sometimes displayed deformed petals under greenhouse conditions. Furthermore, the pods from plants derived from line 45 were in general smaller than the pods from plants derived from the other lines. Since this size difference occurred in both wild-type and mutant *ind* sibling plants derived from line 45, it is probably caused by a background mutation in this line.

- The pods from plants comprising one *ind* allele in homozygous state and one *ind* allele in heterozygous state (genotype: *ind-a1/ind-a1*, *IND-C1/ind-c1* or *IND-A1/ind-a1*, *ind-c1/ind-c1*) showed an intermediate phenotype. More specifically, the valve margins of the pods of these mutant *IND* sibling plants were in general better defined than in the double homozygous mutant *IND* sibling plants, but the sharp indentation between the valve and the beak at the distal end of the pods in the wild-type sibling plants was still largely absent in these mutant plants.

- Table 5a shows the visual pod scores attributed to the phenotype of the pods from plants grown in the field as described above:

Table 5a

Genotype	Line n°	visual pod score (1-5)
<i>IND-A1/IND-A1, IND-C1/IND-C1</i>	51	1
	45	1
	176	1
	48	1
<i>ind-a1/ind-a1, IND-C1/ind-c1</i>	51	3
	45	3
	176	2
	48	3
<i>IND-A1/ind-a1, ind-c1/ind-c1</i>	51	3
	45	3
	176	2
	48	3
<i>ind-a1/ind-a1, ind-c1/ind-c1</i>	51	5
	45	5
	176	4
	48	5

b) Manual Impact Test (MIT):

- The pods from plants comprising two mutant *IND* alleles in homozygous state (genotype: *ind-a1/ind-a1, ind-c1/ind-c1*) derived from lines 51, 45, 176 and 48 were completely pod

shatter resistant (pods did not open along the dehiscence zone even after applying a strong torsion).

- The pod shatter resistance of pods from plants comprising one *ind* allele in homozygous state and one *ind* allele in heterozygous state (genotype: *ind-a1/ind-a1*, *IND-C1/ind-c1* or *IND-A1/ind-a1*, *ind-c1/ind-c1*) was increased as compared to the pod shatter resistance of pods from their wild-type sibling plants, but the pods could still be opened along the dehiscence zone after applying limited physical forces.
- Table 5b shows the scores attributed to the pods from plants grown in the field based on the physical force needed to open closed mature pods by manually applying torsion on the pods as described above:

Table 5b

Genotype	Line n°	Score based on physical force needed to open closed mature pods (1-5)
<i>IND-A1/IND-A1, IND-C1/IND-C1</i>	51	1
	45	1
	176	1
	48	1
<i>ind-a1/ind-a1, IND-C1/ind-c1</i>	51	3
	45	3
	176	1
	48	2
<i>IND-A1/ind-a1, ind-c1/ind-c1</i>	51	3
	45	2
	176	1
	48	3
<i>ind-a1/ind-a1, ind-c1/ind-c1</i>	51	ND
	45	ND
	176	ND
	48	ND

ND: not determined

15 c) Random Impact Test:

- As shown in Table 5c, the half life of pod samples ('LD50') was significantly higher for pods from homozygous double mutants (genotype *ind-a1/ind-a1, ind-c1/ind-c1*) derived from line 51 than for pods of homozygous double mutants derived from line 45, indicating that homozygous double mutant plants comprising the IND-C1-EMS01 allele (line 51) were more pod shatter resistant than homozygous double mutant plants comprising the IND-C1-EMS03 allele (line 45).

- 5 - Table 5c further shows that the LD50 value was in general higher for pods from plants comprising one *ind-c1* allele in homozygous state and one *ind-a1* allele in heterozygous state (genotype: *IND-A1/ind-a1, ind-c1/ind-c1*) than for pods from plants comprising one *ind-a1* allele in homozygous state and one *ind-c1* allele in heterozygous state (genotype: *ind-a1/ind-a1, IND-C1/ind-c1*) indicating that the mutations in the *IND-C1* allele could have a stronger effect on pod shatter resistance than the mutations in the *IND-A1* allele.

Table 5c

Genotype	Line n°	LD50-glasshouse			LD50-field1	LD50-field2
			Lower 95%	Upper 95%		
<i>IND-A1/IND-A1, IND-C1/IND-C1</i>	51	8.61	6.56	11.08	8.9	6.8
	45	8.07	6.08	10.45	7.8	5.7
	176	ND			5.3	5.3
	48	11.42	7.42	14.9	9	5.3
	48	8.86	*	*		
<i>ind-a1/ind-a1, IND-C1/IND-C1</i>	48	9.86	5.89	13.3	ND	ND
<i>IND-A1/IND-A1, ind-c1/ind-c1</i>	48	5.98	2.87	8.6	ND	ND
<i>ind-a1/ind-a1, IND-C1/ind-c1</i>	51	14.22	11.33	17.79	21.1	21.4
	51	22.78	18.68	27.8		
	45	14.97	11.95	18.74	22.9	24.6
	45	10.32	8.05	13.05		
	176	ND			7.3	8.6
	48	7.21	3.04	9.7	10.1	9.4
<i>IND-A1/ind-a1, ind-c1/ind-c1</i>	51	48.31	39.94	58.73	16.9	22.6
	51	46.46	38.44	56.41		
	45	26.89	22.03	32.95	20.6	14.6
	45	17.5	13.96	22.01		
	176	ND			10.9	8.0
	48	30.14	25.49	36.8	18.3	16.5
<i>ind-a1/ind-a1, ind-c1/ind-c1</i>	51	163.28	116.12	237.62	ND	ND
	45	73.57	53.85	103.54	ND	ND
	176	ND			ND	ND
	48	115.99	66.35	523.4	ND	ND

* Insufficient data available to estimate the upper and lower bounds to LD50

10

d) Field tests

Table 5d shows the level of seed shattering (SHAT), combiner harvest ability (CHA1), threshing ability (CHA2), seed yield per plot after combining (YLDP) and seed yield after threshing of straw (YLDS) determined as described above for field plots with *ind* plants and wild-type plants

as indicated. The YieldWTSeg% value represents the YLDP as a percentage of the wildtype segregant within one line, i. e. within line 51, 45, 176 or 48, respectively.

Table 5d

Genotype	Line n°	SHAT (1-9)	CHA1 (1-5)	CHA2 (1-5)	YLDP (in g/plot)	YieldWTSeg%	YLDS (in wt% of straw)
<i>IND-A1/IND-A1, IND-C1/IND-C1</i>	51	8.1	4.9	5.0	2154.7	100	0.6
	45	8.4	4.2	4.8	1868.7	100	0.8
	176	8.1	4.6	5.0	1710.2	100	0.3
	48	7.9	4.7	5.0	1844.2	100	0.5
<i>ind-a1/ind-a1, IND-C1/ind-c1</i>	51	8.9	2.9	3.8	2450.7	114	4.5
	45	8.8	2.3	3.3	2304.2	123	7.6
	176	8.7	3.9	4.9	2189.6	128	0.6
	48	8.8	4.1	4.9	2419.1	131	1.4
<i>IND-A1/ind-a1, ind-c1/ind-c1</i>	51	8.9	3.3	4.3	2739.6	127	1.9
	45	8.8	2.6	3.4	2441.6	131	3.4
	176	8.7	4.1	4.9	2071.6	121	0.7
	48	8.8	3.6	4.1	2379.8	129	2.4
<i>ind-a1/ind-a1, ind-c1/ind-c1</i>	51	9.1	1.2	2.0	515.3	24	27.4
	45	9.0	1.0	2.0	424.4	23	27.4
	176	9.0	1.1	2.6	702.4	41	21.0
	48	9.0	1.0	1.9	447.3	24	27.7

5

Microscopical evaluation:

- Pods from plants comprising two *ind* alleles in homozygous state derived from line 45 (genotype *ind-a1/ind-a1, ind-c1/ind-c1*) grown under greenhouse conditions showed a lignification throughout the complete dehiscence zone and a poor differentiation of cells belonging to the dehiscence zone from neighboring cell types, such as the vascular tissue cells and the lignified layer of cells normally found at the inner pod wall (i.e. the *enb* cells) ("strong morphological phenotype"). A similar pod phenotype was observed for these plants grown under field conditions. By contrast, the dehiscence zones were still well differentiated and mostly non-lignified in pods from plants comprising two *ind* alleles in homozygous state derived from line 51 (genotype *ind-a1/ind-a1, ind-c1/ind-c1*) grown under greenhouse conditions but the dehiscence zones did show extra lignification where the pod walls come together ("weaker morphological phenotype"). Pods from these plants grown under field conditions showed a lignification pattern similar to that of pods from plants with genotype *IND-A1/ind-a1, ind-c1/ind-c1* derived from line 45 described below. When combined with the data obtained from the RIT, these data could indicate that these plants combine a "weaker morphological phenotype" with a higher pod shatter resistance.

20

- Pods from plants comprising one *ind* allele in homozygous state and one *ind* allele in heterozygous state derived from lines 45 and 51 (genotype: *ind-a1/ind-a1*, *IND-C1/ind-c1* or *IND-A1/ind-a1*, *ind-c1/ind-c1*) grown under greenhouse conditions did not show an obvious phenotype. Under field conditions, pods from plants with genotype *IND-A1/ind-a1*, *ind-c1/ind-c1* derived from lines 45 displayed a more subtle morphological phenotype than pods from plants of their homozygous double mutant siblings (see above). More specifically, lignification did not occur throughout the complete dehiscence zone but the pods of these plants only displayed a few extra layers of lignified cells where the inner pod wall is attached to the septum, either symmetrically at both sides of the septum or only unilaterally. A similar pod phenotype was observed for pods from plants with genotype *IND-A1/ind-a1*, *ind-c1/ind-c1* derived from lines 51 grown under field conditions. Pods from plants with genotype *ind-a1/ind-a1*, *IND-C1/ind-c1* derived from lines 45 and 51 did not show an obvious phenotype under field conditions.

15 **Example 7 - Detection and/or transfer of mutant *IND* genes into (elite) *Brassica* lines**

The mutant *IND* genes are transferred into (elite) *Brassica* breeding lines by the following method: A plant containing a mutant *IND* gene (donor plant), is crossed with an (elite) *Brassica* line (elite parent / recurrent parent) or variety lacking the mutant *IND* gene. The following introgression scheme is used (the mutant *IND* gene is abbreviated to *ind* while the wild type is depicted as *IND*):

Initial cross: *ind / ind* (donor plant) X *IND / IND* (elite parent)

F1 plant: *IND / ind*

BC1 cross: *IND / ind* X *IND / IND* (recurrent parent)

BC1 plants: 50% *IND / ind* and 50% *IND / IND*

25 The 50% *IND / ind* are selected using molecular markers (e.g. AFLP, PCR, InvaderTM, and the like; see also below) for the mutant *IND* allele (*ind*).

BC2 cross: *IND / ind* (BC1 plant) X *IND / IND* (recurrent parent)

BC2 plants: 50% *IND / ind* and 50% *IND / IND*

The 50% *IND / ind* are selected using molecular markers for the mutant *IND* allele (*ind*).

30 **Backcrossing is repeated until BC3 to BC6**

BC3-6 plants: 50% *IND / ind* and 50% *IND / IND*

The 50% *IND / ind* are selected using molecular markers for the mutant *IND* allele (*ind*). To reduce the number of backcrossings (e.g. until BC3 in stead of BC6), molecular markers can be used specific for the genetic background of the elite parent.

BC3-6 S1 cross: *IND / ind* X *IND / ind*

5 BC3-6 S1 plants: 25% *IND / IND* and 50% *IND / ind* and 25% *ind / ind*

Plants containing *ind* are selected using molecular markers for the mutant *IND* allele (*ind*). Individual BC3-6 S1 plants that are homozygous for the mutant *IND* allele (*ind / ind*) are selected using molecular markers for the mutant and the wild-type *IND* alleles. These plants are then used for seed production.

10

To select for plants comprising a point mutation in an *IND* allele, direct sequencing by standard sequencing techniques known in the art, such as those described in Example 4, can be used. Alternatively, PCR assays can be developed to discriminate plants comprising a specific point mutation in an *IND* allele from plants not comprising that specific point mutation. The following discriminating PCR assays were thus developed to detect the presence or absence and the

15

- Template DNA:

- Genomic DNA isolated from leaf material of homozygous or heterozygous mutant *Brassica* plants (comprising a mutant *IND* allele, called hereinafter “IND-Xx-EMSXX”).

20

- Wild type DNA control: Genomic DNA isolated from leaf material of wild type *Brassica* plants (comprising the wild type equivalent of the mutant *IND* allele, called hereinafter “IND-Xx-WT”).

- Positive DNA control: Genomic DNA isolated from leaf material of homozygous mutant *Brassica* plants known to comprise IND-Xx-EMSXX.

25

- Primers and length of the fragment amplified from the mutant and corresponding wild-type target *IND* gene are indicated in Table 6. Generally, each primer set consists of one primer specific for the mutant and the wild type target gene (e.g. primer POSH101R2 is specific for IND-A1-EMS01 and IND-A1-WT) and one primer specific for the nucleotide difference (e.g. primer POSH101MF1 is specific for the IND-A1-EMS01 and primer POSH101WF1 is specific for IND-A1-WT). Usually, the last nucleotide of the latter primer matches with the nucleotide difference (underlined nucleotide in Table 6), but one (or more) additional target specific nucleotide(s) may be added to improve the annealing between the primer and its target sequence (see e.g. bold nucleotide in primer POSH 108MR1', which is specific for the

30

IND-C1-EMS03 allele, as compared to primer POSH 108WR1', which is specific for the IND-C1-WT allele).

- PCR mix: 2.5 μ l 10x PCR buffer (15mM MgCl₂), 0.25 μ l dNTP's (20 mM), 1 μ l forward primer (10 μ M), 1 μ l reverse primer(10 μ M), 0.25 μ l Taq-polymerase (5U/ μ l), 19.5 μ l Milli-Q H₂O, 0.5 μ l DNA (20-50 ng/ μ l) = Total volume of 25 μ l;
- Thermocycling profile: 4 min at 95°C; 30x [1min at 95°C (denaturation) and 1 min at annealing temperature specified in Table 6 and 2 min at 72°C (elongation)]; 5min at 72°C; cool down to 4°C. The optimal annealing temperature was determined by temperature gradient PCR wherein the annealing temperature was varied between 57°C to 70°C on a MJ Research thermocycler PTC-200 (Biozym). The optimal annealing temperature for the wild type *IND* specific primers is that temperature at which a clear PCR fragment of the expected size can be detected (as described below) for the DNA sample from the wild type *Brassica* plant and not for the DNA sample from the mutant *Brassica* plant. The optimal annealing temperature for the mutant *IND* specific primers is that temperature at which a clear PCR fragment of the expected size can be detected (as described below) for the DNA sample from the mutant *Brassica* plant and not for the DNA sample from the wild type *Brassica* plant.
- After amplification, 5 μ l loading dye (orange dye) was added to 15 μ l of the PCR samples and the samples were loaded on a 1.5% agarose gel.
- The banding patterns obtained after amplification of genomic DNA of mutant *Brassica* plants are evaluated as follows:
 - Data from DNA samples isolated from leaf material of the mutant *Brassica* plants within a single PCR run and a single PCR mix should not be accepted unless:
 - the wild-type DNA control shows the PCR fragment of the expected size for the *IND - Xx-WT* specific PCR assay and no PCR fragment of the expected size for the *IND-Xx-EMSXX* specific PCR assay
 - the positive DNA control shows the PCR fragment of the expected size for the *IND-Xx-EMSXX* specific PCR assay and no PCR fragment of the expected size for the *IND-Xx-WT* specific PCR assay
 - Lanes showing no PCR product of the expected size for the *IND-Xx-WT* specific PCR assay and the PCR fragment of the expected size for the *IND-Xx-EMSXX* specific PCR assay, indicate that the corresponding plant from which the genomic template DNA was prepared, is a homozygous mutant for *IND-Xx-EMSXX*.
 - Lanes showing the PCR fragment of the expected size for the *IND -Xx-WT* specific PCR assay and the *IND-Xx-EMSXX* specific PCR assay, indicate that the corresponding plant

from which the genomic template DNA was prepared, is a heterozygous mutant for *IND-Xx-EMSXX*.

- 5 - Lanes showing the PCR fragment of the expected size for the *IND -Xx-WT* specific PCR assay and no PCR product of the expected size for the *IND-Xx-EMSXX* specific PCR assay, indicate that the corresponding plant from which the genomic template DNA was prepared, is a wild type plant.

Table 6:

Allele No.	Primers	Annealing t° (°C)	Size fragment (bp)	PCR
IND-A1-EMS01	5' AAGGGTAAGCGACGACCCTT 3' (POSH101MF1 - SEQ ID NO: 13) 5' GAGTGTGAGGCTGAAGAAGC 3' (POSH101R2 - SEQ ID NO: 15)	67	191	
IND-A1-WT	5' AAGGGTAAGCGACGACCCTC 3' (POSH101WF1 - SEQ ID NO: 14) 5' GAGTGTGAGGCTGAAGAAGC 3' (POSH101R2 - SEQ ID NO: 15)	71.1	191	
IND-A1-EMS05	5' CCTCAGACGGTGGTGGCTCA 3' (POSH105MF1 - SEQ ID NO: 16) 5' AGGGTCAGACATAGGAGCTC 3' (POSH 101R1 - SEQ ID NO: 18)	70	201	
IND-A1-WT	5' CCTCAGACGGTGGTGGCTCG 3' (POSH105WF1 - SEQ ID NO: 17) 5' AGGGTCAGACATAGGAGCTC 3' (POSH 101R1 - SEQ ID NO: 18)	72	201	
IND-C1-EMS01	5' GTGGTTAAAAGAGTTTTCTTA 3' (POSH106MR1 - SEQ ID NO: 19) 5' ATTAGCATGTAAAACACTAG 3' (POSH106F1 - SEQ ID NO: 21)	60.6	436	
IND-C1-WT	5' GTGGTTAAAAGAGTTTTCTTG 3' (POSH106WR1 - SEQ ID NO: 20) 5' ATTAGCATGTAAAACACTAG 3' (POSH106F1 - SEQ ID NO: 21)	62.8	436	
IND-C1-EMS03	5' ACGAGCCACCACCGTCTAG 3' (POSH 108MR1' - SEQ ID NO: 22) 5' GTTCAAAGCAGATGCAGCAG 3' (POSH106F2 - SEQ ID NO: 24)	70	369	
IND-C1-WT	5' ACGAGCCACCACCGTCTG 3' (POSH 108WR1' - SEQ ID NO: 23) 5' GTTCAAAGCAGATGCAGCAG 3' (POSH106F2 - SEQ ID NO: 24)	68.9	369	

- 10 Alternatively, Invader™ technology (Third Wave Agbio) can be used to discriminate plants comprising a specific point mutation in an *IND* allele from plants not comprising that specific point mutation. The following discriminating Invader™ probes were thus developed to detect the

presence or absence and the zygosity status of the mutant alleles identified in Example 4 (see Table 7:

- Probes specific for the mutant or corresponding wild-type target *IND* gene (indicated as “5’ flap1-x” and “5’ flap2-x”, respectively) and “invading” probes which can be used in combination with them are indicated in Table 7. Generally, each probe set consists of one probe specific for the mutant or the wild type target gene of which the first nucleotide after the 5’ flap sequence matches with the nucleotide difference (underlined nucleotide in Table 7) (the so-called “primary probe”; e.g. the probe with SEQ ID NO: 26 is specific for IND-A1-EMS01 and the probe with SEQ ID NO: 27 is specific for IND-A1-WT) and one probe specific for the nucleotides upstream of the nucleotide difference (the so-called “invader® oligo”; e.g. the probe with SEQ ID NO: 25 is specific for the nucleotides upstream of the nucleotide difference between IND-A1-EMS01 and IND-A1-WT). The last nucleotide of the latter primer may match with the nucleotide difference in the mutant (as indicated by the bold nucleotides in Table 6), but other nucleotides may be used as well for this last nucleotide as long as the primary probe and the invader® oligo are still able to form a single base overlap when hybridized to the target DNA to generate the specific invasive structure recognized by the Cleavase® enzymes (Third Wave Agbio).
- The Invader™ assay procedure and interpretation of the data are performed as prescribed by the manufacturer (Third Wave Agbio). Briefly, the nucleotide sequences indicated as “flap1” and “flap2” in Table 7 represent the sequences of the 5’ “flaps” which are cleaved from the primary probes in the primary phase of the Invader™ assay and which are complementary to sequences in FRET™ cassette 1 and 2, respectively, and not complementary to the target mutant or wild type sequences. If the primary probes are cleaved in the primary phase and the flap1-probe and/or flap2-probe hybridise to FRET™ cassette 1 and 2, respectively, in the secondary phase, a signal is generated indicative of the presence in the sample of the mutant or corresponding wild-type target *IND* gene, respectively.

Table 7

Allele No.	Probes	
IND-A1-EMS01	5’ GCCGACGAGCCACCACCGTCTT 3’	(SEQ ID NO: 25)
	5’ flap1- <u>A</u> AGGGTCGTCGCTT 3’	(SEQ ID NO: 26)
IND-A1-WT	5’ GCCGACGAGCCACCACCGTCTT 3’	(SEQ ID NO: 25)
	5’ flap2- <u>G</u> AGGGTCGTCGCT 3’	(SEQ ID NO: 27)
IND-A1-EMS05	5’ CGGATCTTCTCGCTTATCCTTTCTCTACGCCGAA 3’	(SEQ ID NO: 28)
	5’ flap1- <u>I</u> GAGCCACCACCG 3’	(SEQ ID NO: 29)
IND-A1-WT	5’ CGGATCTTCTCGCTTATCCTTTCTCTACGCCGAA 3’	(SEQ ID NO: 28)
	5’ flap2- <u>C</u> GAGCCACCACCG 3’	(SEQ ID NO: 30)

IND-C1-EMS01	5' AGGTGGATCTACCATGAAATGAGGATTGTGGTT AAAAGAGTTTTCTTT 3'	(SEQ ID NO: 31)
	5' flap1- <u>A</u> TGTAATGAGATCAATAGGTTTG 3'	(SEQ ID NO: 32)
IND-C1-WT	5' AGGTGGATCTACCATGAAATGAGGATTGTGGTT AAAAGAGTTTTCTTT 3'	(SEQ ID NO: 31)
	5' flap2- <u>G</u> TGTAATGAGATCAATAGGTTTG 3'	(SEQ ID NO: 33)
IND-C1-EMS03	5' CCGTAACGTAAGGGTAAGCGAGGACCCCA 3'	(SEQ ID NO: 34)
	5' flap1- <u>T</u> AGACGGTGGTGGC 3'	(SEQ ID NO: 35)
IND-C1-WT	5' CCGTAACGTAAGGGTAAGCGAGGACCCCA 3'	(SEQ ID NO: 34)
	5' flap2- <u>C</u> AGACGGTGGTGGC 3'	(SEQ ID NO: 36)

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A non-naturally occurring *Brassica* plant comprising at least two *IND* genes, or a cell, part, seed or progeny thereof, wherein the plant comprises two or three full knock-out mutant *IND* alleles in its genome.
2. The plant according to claim 1, wherein the *IND* genes are *IND-A1* or *IND-C1* genes.
3. The plant according to claim 1 or 2, wherein the *IND* genes comprise a nucleic acid molecule selected from the group consisting of:
 - (a) a nucleic acid molecule which comprises at least 90% sequence identity to SEQ ID NO: 1, SEQ ID NO: 3 from the nucleotide at position 46 to the nucleotide at position 633, SEQ ID NO: 3, SEQ ID NO: 5, or SEQ ID NO: 7; and
 - (b) a nucleic acid molecule encoding an amino acid sequence comprising at least 90% sequence identity to SEQ ID NO: 2, SEQ ID NO: 4 from the amino acid at position 16 to the amino acid at position 21 or SEQ ID NO: 4.
4. The plant according to any one of the preceding claims, wherein the mutant *IND* alleles are selected from the group consisting of *ind-a1-EMS01*, *ind-a1-EMS05*, *ind-cl-EMS01* and *ind-cl-EMS03*.
5. The plant according to any one of the preceding claims, which is homozygous for one mutant *IND* allele.
6. The plant according to any one of the preceding claims, which produces a significantly reduced amount of functional *IND* protein compared to the amount of functional *IND* protein produced by a corresponding plant not comprising mutant *IND* alleles.
7. The plant according to any one of the preceding claims, wherein the seed yield of the plant is increased, preferably significantly increased compared to the seed yield of a corresponding plant not comprising mutant *IND* alleles.
8. The plant according to any one of the preceding claims, comprising three full knock-out *IND* alleles.

9. The plant according to claim 8, wherein the seed shattering of the plant is significantly reduced or delayed compared to the seed shattering of a corresponding plant not comprising mutant *IND* alleles.
10. The plant according to claim 9, which maintains an agronomically relevant treshability of the pods.
11. The plant according to any one of the preceding claims, which is a plant from a *Brassica* crop species, preferably *Brassica napus*, *Brassica juncea*, or *Brassica carinata*.
12. A seed pod obtainable from a plant according to any one of claims 1 to 11.
13. A method for identifying a *Brassica* plant according to any one of claims 1 to 11, or a cell, part, seed or progeny thereof, said method comprising identifying a first, a second and, optionally, a third selected full knock-out mutant *IND* allele in a biological sample comprising determining the presence of a mutant *IND* specific region of each selected full knock-out mutant *IND* allele in a nucleic acid present in the biological sample.
14. The method according to claim 13, which further comprises subjecting the biological sample to a polymerase chain reaction assay using a set of at least two primers for each selected full knock-out mutant *IND* allele, said set being selected from the group consisting of:
 - a set of primers, wherein one of said primers specifically recognizes the 5' flanking region of the mutant *IND* allele and the other of said primers specifically recognizes the 3' flanking region of the mutant *IND* allele, respectively,
 - a set of primers, wherein one of said primers specifically recognizes the 5' or 3' flanking region of the mutant *IND* allele and the other of said primers specifically recognizes the mutation region of the mutant *IND* allele, and
 - a set of primers, wherein one of said primers specifically recognizes the 5' or 3' flanking region of the mutant *IND* allele and the other of said primers specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the mutant *IND* allele, respectively.

15. The method according to claim 14, wherein said set of primers is selected from the group consisting of:
- a set of primers comprising one primer comprising the sequence of SEQ ID NO: 13 and/or one primer comprising the sequence of SEQ ID NO: 15,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 16 and/or one primer comprising the sequence of SEQ ID NO: 18,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 19 and/or one primer comprising the sequence of SEQ ID NO: 21, and
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 22 and/or one primer comprising the sequence of SEQ ID NO: 24.
16. The method according to claim 13, which further comprises subjecting the biological sample to a hybridization assay using a set of specific probes comprising at least one specific probe for each selected full knock-out mutant *IND* allele, said set being selected from the group consisting of :
- a set of specific probes, wherein one of said probes specifically recognizes the 5' flanking region of the mutant *IND* allele, and the other of said probes specifically recognizes the 3' flanking region of the mutant *IND* allele,
 - a set of specific probes, wherein one of said probes specifically recognizes the 5' or 3' flanking region of the mutant *IND* allele, and the other of said probes specifically recognizes the mutation region of the mutant *IND* allele,
 - a set of specific probes, wherein one of said probes specifically recognizes the 5' or 3' flanking region of the mutant *IND* allele and the other of said probes specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the mutant *IND* allele, respectively, and
 - a specific probe which specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the mutant *IND* allele.
17. The method according to claim 16, wherein said set of probes is selected from the group consisting of:
- a set of probes comprising one probe comprising the sequence of SEQ ID NO: 25 and/or one probe comprising the sequence of SEQ ID NO: 26,

- a set of probes comprising one probe comprising the sequence of SEQ ID NO: 28 and/or one probe comprising the sequence of SEQ ID NO: 29,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 31 and/or one probe comprising the sequence of SEQ ID NO: 32, and
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 34 and/or one probe comprising the sequence of SEQ ID NO: 35.
18. The method according to any one of claims 13 to 17, further comprising determining the zygosity status of a selected full knock-out mutant *IND* allele, comprising determining the presence of a mutant and/or a corresponding wild type *IND* specific region in the genomic DNA of said plant, or a cell, part, seed or progeny thereof.
19. The method according to claim 18, which further comprises subjecting the genomic DNA of said plant, or a cell, part, seed or progeny thereof, to a polymerase chain reaction assay using a set of at least two or at least three primers for each selected full knock-out mutant *IND* allele, wherein at least two of said primers specifically recognize the wild type *IND* allele, said at least two primers being selected from the group consisting of:
- a first primer which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a second primer which specifically recognizes the 3' or 5' flanking region of the mutant and the wild type *IND* allele, respectively,
 - a first primer which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a second primer which specifically recognizes the mutation region of the wild type *IND* allele, and
 - a first primer which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a second primer which specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the wild type *IND* allele, respectively, and
- wherein at least two of said primers specifically recognize the selected full knock-out mutant *IND* allele, said at least two primers being selected from the group consisting of:

- the first primer which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and the second primer which specifically recognizes the 3' or 5' flanking region of the mutant and the wild type *IND* allele, respectively,
 - the first primer which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a third primer which specifically recognizes the mutation region of the mutant *IND* allele, and
 - - the first primer which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a third primer which specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the mutant *IND* allele, respectively.
20. The method according to claim 19, wherein said set of at least three primers is selected from the group consisting of:
- a set of primers comprising one primer comprising the sequence of SEQ ID NO: 13, one primer comprising the sequence of SEQ ID NO: 14, and/or one primer comprising the sequence of SEQ ID NO: 15,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 16, one primer comprising the sequence of SEQ ID NO: 17, and/or one primer comprising the sequence of SEQ ID NO: 18,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 19, one primer comprising the sequence of SEQ ID NO: 20, and/or one primer comprising the sequence of SEQ ID NO: 21, and
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 22, one primer comprising the sequence of SEQ ID NO: 23, and/or one primer comprising the sequence of SEQ ID NO: 24.
21. The method according to claim 18, which further comprises subjecting the genomic DNA of said plant, or a cell, part, seed or progeny thereof, to a hybridization assay using a set of at least two specific probes for said selected full knock-out mutant *IND* allele, wherein at least one of said specific probes specifically recognizes the wild type *IND* allele, said at least one probe selected from the group consisting of:

- a first probe which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a second probe which specifically recognizes the 3' and 5' flanking region of the mutant and the wild type *IND* allele, respectively,
- a first probe which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a second probe which specifically recognizes the mutation region of the wild type *IND* allele,
- a first probe which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a second probe which specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the wild type *IND* allele, respectively, and
- a probe which specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the wild type *IND* allele, and

wherein at least one of said specific probes specifically recognize(s) the selected full knock-out mutant *IND* allele, said at least one probe selected from the group consisting of:

- the first probe which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and the second probe which specifically recognizes the 3' or 5' flanking region of the mutant and the wild type *IND* allele, respectively,
- the first probe which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a third probe which specifically recognizes the mutation region of the mutant *IND* allele,
- the first probe which specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, and a third probe which specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the mutant *IND* allele, and
- a probe which specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the mutant *IND* allele.

22. The method according to claim 21, wherein said set of at least three specific probes is selected from the group consisting of:
- a set of probes comprising one probe comprising the sequence of SEQ ID NO: 25, one probe comprising the sequence of SEQ ID NO: 26, and/or one probe comprising the sequence of SEQ ID NO: 27,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 28, one probe comprising the sequence of SEQ ID NO: 29, and/or one probe comprising the sequence of SEQ ID NO: 30,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 31, one probe comprising the sequence of SEQ ID NO: 32, and/or one probe comprising the sequence of SEQ ID NO: 33, and
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 34, one probe comprising the sequence of SEQ ID NO: 35, and/or one probe comprising the sequence of SEQ ID NO: 36.
23. A kit when used for identifying a *Brassica* plant according to any one of claims 1 to 11, or a cell, part, seed or progeny thereof, said kit used for identifying a first, a second and, optionally, a third selected full knock-out mutant *IND* allele in a biological sample, comprising a set of primers or probes for each selected full knock-out mutant *IND* allele, said set selected from the group consisting of:
- a set of primers or probes, wherein one of said primers or probes specifically recognizes the 5' flanking region of the mutant *IND* allele and the other of said primers or probes specifically recognizes the 3' flanking region of the mutant *IND* allele,
 - a set of primers or probes, wherein one of said primers or probes specifically recognizes the 5' or 3' flanking region of the mutant *IND* allele and the other of said primers or probes specifically recognizes the mutation region of the mutant *IND* allele,
 - a set of primers or probes, wherein one of said primers specifically recognizes the 5' or 3' flanking region of the mutant *IND* allele and the other of said primers or probes specifically recognizes the joining region between the 3' or

- 5' flanking region and the mutation region of the mutant *IND* allele, respectively, and
- a probe which specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the mutant *IND* allele.
24. The method according to claim 14 or 16, or the kit according to claim 23, wherein for the first, the second, or the third selected full knock-out mutant *IND* allele
- said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 923 or 925 to 1622 or of the complement thereof, respectively; said mutation region has the nucleotide sequence of nucleotide 924 of SEQ ID NO: 5 or of the complement thereof; and said joining region comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 924 or 924 to 1622 or of the complement thereof, respectively, or
 - said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 866 or 868 to 1622 or of the complement thereof, respectively, or of SEQ ID NO: 5 from nucleotide 1 to 939 or 941 to 1622 or of the complement thereof, respectively; said mutation region has the nucleotide sequence of nucleotide 867 of SEQ ID NO: 5 or of the complement thereof, or of nucleotide 940 of SEQ ID NO: 5 or of the complement thereof; and said joining region comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 867 or 867 to 1622 or from the complement thereof, or of SEQ ID NO: 5 from nucleotide 1 to 940 or 940 to 1622 or from the complement thereof, respectively, or
 - said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 643 or 645 to 1593 or of the complement thereof, respectively; said mutation region has the nucleotide sequence of nucleotide 644 of SEQ ID NO: 7 or of the complement thereof; and said joining region comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 644 or 644 to 1593 or of the complement thereof, respectively, or
 - said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 898 or 900 to 1593 or from the complement thereof, respectively; said mutation region has the nucleotide sequence of nucleotide

899 of SEQ ID NO: 7 or of the complement thereof; and said joining region comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 899 or 899 to 1593 or from the complement thereof, respectively.

25. The kit according to claim 23 or 24, wherein said set of at least two primers and/or said set of at least two specific probes for each selected full knock-out mutant *IND* allele is selected from the group consisting of:
- a set of primers comprising one primer comprising the sequence of SEQ ID NO: 13 and/or one primer comprising the sequence of SEQ ID NO: 15,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 16 and/or one primer comprising the sequence of SEQ ID NO: 18,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 19 and/or one primer comprising the sequence of SEQ ID NO: 21,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 22 and/or one primer comprising the sequence of SEQ ID NO: 24,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 25 and/or one probe comprising the sequence of SEQ ID NO: 26,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 28 and/or one probe comprising the sequence of SEQ ID NO: 29,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 31 and/or one probe comprising the sequence of SEQ ID NO: 32, and
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 34 and/or one probe comprising the sequence of SEQ ID NO: 35.
26. A kit when used for determining the zygosity status of a selected full knock-out mutant *IND* allele in a *Brassica* plant according to any one of claims 1 to 11, or a cell, part, seed or progeny thereof, , said kit comprising a set of primers or probes, wherein at least two of said primers or at least one of said probes specifically recognize the wild type *IND* allele and wherein at least two of said primers or at least one of said probes specifically recognize the mutant *IND* allele for each selected full knock-out mutant *IND* allele, selected from the group consisting of:

- a set of at least two primers or probes, wherein a first primer or probe specifically recognizes the 5' flanking region of the mutant and the wild type *IND* allele and a second primer or probe specifically recognizes the 3' flanking region of the mutant and the wild type *IND* allele,
 - a set of at least three primers or probes, wherein a first primer or probe specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, a second primer or probe specifically recognizes the mutation region of the mutant *IND* allele, and a third primer or probe specifically recognizes the mutation region of the wild type *IND* allele,
 - a set of at least three primers or probes, wherein a first primer or probe specifically recognizes the 5' or 3' flanking region of the mutant and the wild type *IND* allele, a second primer or probe specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the mutant *IND* allele, respectively, and a third primer or probe specifically recognizes the joining region between the 3' or 5' flanking region and the mutation region of the wild type *IND* allele, respectively, and
 - a set of at least two probes, wherein a first probe specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the mutant *IND* allele and a second probe specifically recognizes the joining region between the 5' or 3' flanking region and the mutation region of the wild type *IND* allele.
27. The method according to claim 19 or 21, or the kit according to claim 26, wherein for the first, the second, or the third selected full knock-out mutant *IND* allele
- said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 923 or 925 to 1622 or of the complement thereof, respectively; said mutation region of the wild type *IND* allele has the nucleotide sequence of nucleotide 924 of SEQ ID NO: 5 or of the complement thereof; said mutation region of the mutant *IND* allele has the sequence t or the complement thereof; said joining region of the wild type *IND* allele comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 924 or 924 to 1622 or of the complement thereof, respectively; and said joining region of the mutant *IND* allele comprises the nucleotide sequence of SEQ ID NO: 5 from

nucleotide 1 to 923 followed by t or t followed by the nucleotide sequence SEQ ID NO: 5 from nucleotide 925 to 1622 or of the complement thereof, respectively, or

- said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 866 or 868 to 1622 or of the complement thereof, respectively, or of SEQ ID NO: 5 from nucleotide 1 to 939 or 941 to 1622 or of the complement thereof, respectively; said mutation region of the wild type *IND* allele has the nucleotide sequence of nucleotide 867 of SEQ ID NO: 5 or of the complement thereof, or of nucleotide 940 of SEQ ID NO: 5 or of the complement thereof; said mutation region of the mutant *IND* allele has the sequence a or the complement thereof; said joining region of the wild type *IND* allele comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 867 or 867 to 1622 or from the complement thereof, or of SEQ ID NO: 5 from nucleotide 1 to 940 or 940 to 1622 or from the complement thereof, respectively; and said joining region of the mutant *IND* allele comprises the nucleotide sequence of SEQ ID NO: 5 from nucleotide 1 to 866 followed by a or a followed by the nucleotide sequence SEQ ID NO: 5 from nucleotide 868 to 1622 or from the complement thereof, or of SEQ ID NO: 5 from nucleotide 1 to 939 followed by a or a followed by the nucleotide sequence SEQ ID NO: 5 from nucleotide 941 to 1622 or from the complement thereof, respectively, or
- said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 643 or 645 to 1593 or of the complement thereof, respectively; said mutation region of the wild type *IND* allele has the nucleotide sequence of nucleotide 644 of SEQ ID NO: 7 or of the complement thereof; said mutation region of the mutant *IND* allele has the sequence t or the complement thereof; and said joining region of the wild type *IND* allele comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 644 or 644 to 1593 or of the complement thereof, respectively; and said joining region of the mutant *IND* allele comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 643 followed by t or t followed by the nucleotide sequence SEQ ID NO: 7 from nucleotide 645 to 1593 or of the complement thereof, respectively, or

- said 5' or 3' flanking region comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 898 or 900 to 1593 or from the complement thereof, respectively; said mutation region of the wild type *IND* allele has the nucleotide sequence of nucleotide 899 of SEQ ID NO: 7 or of the complement thereof; said mutation region of the mutant *IND* allele has the sequence t or the complement thereof; and said joining region of the wild type *IND* allele comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 899 or 899 to 1593 or from the complement thereof, respectively; and said joining region of the mutant *IND* allele comprises the nucleotide sequence of SEQ ID NO: 7 from nucleotide 1 to 898 followed by t or t followed by the nucleotide sequence SEQ ID NO: 7 from nucleotide 900 to 1593 or of the complement thereof, respectively.
28. The kit according to claim 26 or 27, wherein said set of at least three primers or probes for each selected full knock-out mutant *IND* allele is selected from the group consisting of:
- a set of primers comprising one primer comprising the sequence of SEQ ID NO: 13, one primer comprising the sequence of SEQ ID NO: 14, and/or one primer comprising the sequence of SEQ ID NO: 15,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 16, one primer comprising the sequence of SEQ ID NO: 17, and/or one primer comprising the sequence of SEQ ID NO: 18,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 19, one primer comprising the sequence of SEQ ID NO: 20, and/or one primer comprising the sequence of SEQ ID NO: 21,
 - a set of primers comprising one primer comprising the sequence of SEQ ID NO: 22, one primer comprising the sequence of SEQ ID NO: 23, and/or one primer comprising the sequence of SEQ ID NO: 24,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 25, one probe comprising the sequence of SEQ ID NO: 26, and/or one probe comprising the sequence of SEQ ID NO: 27,

- a set of probes comprising one probe comprising the sequence of SEQ ID NO: 28, one probe comprising the sequence of SEQ ID NO: 29, and/or one probe comprising the sequence of SEQ ID NO: 30,
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 31, one probe comprising the sequence of SEQ ID NO: 32, and/or one probe comprising the sequence of SEQ ID NO: 33, and
 - a set of probes comprising one probe comprising the sequence of SEQ ID NO: 34, one probe comprising the sequence of SEQ ID NO: 35 and/or one probe comprising the sequence of SEQ ID NO: 36.
29. A method for combining two or three selected full knock-out mutant *IND* alleles in one *Brassica* plant comprising at least two *IND* genes, said method comprising the steps of:
- identifying at least two plants each comprising at least one selected full knock-out mutant *IND* allele using the methods as described in any one of claims 13 to 17,
 - crossing the at least two plants and collecting F1 hybrid seeds from the at least one cross, and
 - optionally, identifying an F1 plant comprising two or three selected mutant *IND* alleles using the method as described in any one of claims 13 to 17.
30. The method according to claim 29, which further comprises the step of identifying an F1 plant, which is homozygous or heterozygous for a selected mutant *IND* allele by determining the zygosity status of the selected mutant *IND* allele, using the methods according to any one of claims 28 to 22.
31. A method for making a plant according to any one of claims 1 to 11 comprising combining and/or transferring full knock-out mutant *IND* alleles in or to one *Brassica* plant, according to claim 29 or 30.
32. A method for making a hybrid *Brassica* seed or plant according to any one of claims 1 to 11 comprising the steps of:
- (a) identifying a first plant comprising a first and a second selected full knock-out mutant *IND* allele in homozygous state and a second plant comprising a

third selected full knock-out mutant *IND* allele in homozygous state according to any one of claims 18 to 22, and

- (b) crossing the first and the second plant and collecting F1 hybrid seeds from the cross.
33. The method according to claim 32, wherein the first or the second selected full knock-out mutant *IND* allele is the same mutant *IND* allele as the third selected full knock-out mutant *IND* allele.
34. The method according to claim 32 or 33, wherein the first plant is used as a male parent plant and the second plant is used as a female parent plant.
35. *Brassica* seed comprising at least two *IND* genes, wherein the seed comprises two or three full knock-out mutant *IND* alleles comprising the *ind-a1*-EMS01 and *ind-cl*-EMS01 alleles having been deposited at the ATCC on November 20, 2007, under accession number PTA-8796 and *Brassica* seed comprising the *ind-a1*-EMS05 and *ind-cl*-EMS03 alleles having been deposited at the ATCC on November 20, 2007, under accession number PTA-8795 or derivatives therefrom.
36. A *Brassica* plant, or a cell, part, seed or progeny thereof, comprising at least two *IND* genes, wherein the plant comprises two or three full knock-out mutant *IND* alleles obtained from *Brassica* seed comprising the *ind-a1*-EMS01 and *ind-cl*-EMS01 alleles having been deposited at the ATCC on November 20, 2007, under accession number PTA-8796 and *Brassica* seed comprising the *ind-a1*-EMS05 and *ind-cl*-EMS03 alleles having been deposited at the ATCC on November 20, 2007, under accession number PTA-8795 or derivatives therefrom.
37. A *Brassica* plant, or a cell, part, seed or progeny thereof, comprising at least two *IND* genes, wherein the plant comprises two or three full knock-out mutant *IND* alleles, said full knock-out mutant *IND* alleles selected from one or more of the group consisting of the *ind-a1*-EMS01, *ind-a1*-EMS05, *ind-cl*-EMS01 and *ind-cl*-EMS03 allele in its genome, obtained by propagation of and/or breeding with a *Brassica* plant grown from the seed of claim 35.
38. A seed comprising at least two *IND* genes, wherein the seed comprises two or three full knock-out mutant *IND* alleles, said full knock-out mutant *IND* alleles selected from one or more of the group consisting of the *ind-a1*-EMS01, *ind-a1*-EMS05,

ind-cl-EMS01 and *ind-cl*-EMS03 allele, reference seed comprising said allele having been deposited at the ATCC on November 20, 2007, under accession number PTA-8796 and PTA-8795.

39. A plant, or a cell, part, seed or progeny thereof, comprising at least two *IND* genes, wherein the plant comprises two or three full knock-out mutant *IND* alleles, said full knock-out mutant *IND* alleles selected from one or more of the group consisting of the *ind-a1*-EMS01, *ind-a1*-EMS05, *ind-cl*-EMS01 or *ind-cl*-EMS03 allele, produced from the seed of claim 38.
40. A method to increase the yield of *Brassica* plant comprising at least two *IND* genes, comprising introducing two mutant homozygous *IND* alleles in its genome.
41. Use of two or three full knock-out mutant *IND* alleles to increase the seed yield in a *Brassica* plant comprising at least two *IND* genes.
42. Use of two or three full knock-out mutant *IND* alleles to increase the shatter resistance of pods in a *Brassica* plant comprising at least two *IND* genes.




Figure 1




Figure 2

SEQUENCE LISTING

<110> Bayer BioScience N.V.
 <120> Brassica plant comprising a mutant INDEHISCENT allele
 <130> BCS 07-2014
 <160> 40
 <170> PatentIn version 3.3
 <210> 1
 <211> 558
 <212> DNA
 <213> Brassica napus wild type IND-A1 coding sequence

<220>
 <221> CDS
 <222> (1)..(555)

<400> 1
 atg tct ggc tca aaa gca gat gca gcc ata gcc cca ata gtc atg atg 48
 Met Ser Gly Ser Lys Ala Asp Ala Ala Ile Ala Pro Ile Val Met Met
 1 5 10 15
 gag cat cat cat ctc ctt atg aat tgg aac aaa cct att gat ctc att 96
 Glu His His His Leu Leu Met Asn Trp Asn Lys Pro Ile Asp Leu Ile
 20 25 30
 aca gaa gaa aac tct ttt aac cac aat cct cat ttc ata gta gat cca 144
 Thr Glu Glu Asn Ser Phe Asn His Asn Pro His Phe Ile Val Asp Pro
 35 40 45
 cct tcc gaa acc cta agc cac ttc cag ccc ccg ccg aca atc ttc tcc 192
 Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro Thr Ile Phe Ser
 50 55 60
 gat cac gga gga gga gag gaa gca gaa gaa gaa gaa gaa gaa gaa gga 240
 Asp His Gly Gly Gly Glu Glu Ala Glu Glu Glu Glu Glu Glu Glu Gly
 65 70 75 80
 gag gaa gag atg gat ccg atg aag aag atg caa tac gcg att gct gcc 288
 Glu Glu Glu Met Asp Pro Met Lys Lys Met Gln Tyr Ala Ile Ala Ala
 85 90 95
 atg cag ccc gta gac ctc gat cca gcc acc gtt cct aag ccg aac cgc 336
 Met Gln Pro Val Asp Leu Asp Pro Ala Thr Val Pro Lys Pro Asn Arg
 100 105 110
 cgt aac gta agg gta agc gac gac cct cag acg gtg gtg gct cgt cgg 384
 Arg Asn Val Arg Val Ser Asp Asp Pro Gln Thr Val Val Ala Arg Arg
 115 120 125
 cgt aga gaa agg ata agc gag aag atc cgg ata ttg aag agg atg gtg 432
 Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met Val
 130 135 140
 cca ggc ggt gca aag atg gac act gcc tcc atg ctc gac gaa gcc atc 480
 Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala Ile
 145 150 155 160
 cgc tac acc aag ttc ttg aaa cgg cag gtg agg cta gct tct tca gcc 528
 Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg Leu Ala Ser Ser Ala
 165 170 175
 tca cac tca gct tgg agc tcc tat gtc tga 558
 Ser His Ser Ala Trp Ser Ser Tyr Val

180

185

<210> 2
<211> 185
<212> PRT
<213> Brassica napus wild type IND-A1 coding sequence

<400> 2

Met Ser Gly Ser Lys Ala Asp Ala Ala Ile Ala Pro Ile Val Met Met
1 5 10 15
Glu His His His Leu Leu Met Asn Trp Asn Lys Pro Ile Asp Leu Ile
20 25 30
Thr Glu Glu Asn Ser Phe Asn His Asn Pro His Phe Ile Val Asp Pro
35 40 45
Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro Thr Ile Phe Ser
50 55 60
Asp His Gly Gly Gly Glu Glu Ala Glu Glu Glu Glu Glu Glu Gly
65 70 75 80
Glu Glu Glu Met Asp Pro Met Lys Lys Met Gln Tyr Ala Ile Ala Ala
85 90 95
Met Gln Pro Val Asp Leu Asp Pro Ala Thr Val Pro Lys Pro Asn Arg
100 105 110
Arg Asn Val Arg Val Ser Asp Asp Pro Gln Thr Val Val Ala Arg Arg
115 120 125
Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met Val
130 135 140
Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala Ile
145 150 155 160
Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg Leu Ala Ser Ser Ala
165 170 175
Ser His Ser Ala Trp Ser Ser Tyr Val
180 185

<210> 3
<211> 633
<212> DNA
<213> Brassica napus wild type IND-C1 coding sequence

<220>
<221> CDS
<222> (1)..(630)

<400> 3

atg tat aaa aga aag gtc tat gcg tct cta gtc caa aaa ctc tat atg 48
 Met Tyr Lys Arg Lys Val Tyr Ala Ser Leu Val Gln Lys Leu Tyr Met
 1 5 10 15

tct ggt tca aaa gca gat gca gca gcc ata gcc cca ata gtc atg atg 96
 Ser Gly Ser Lys Ala Asp Ala Ala Ala Ile Ala Pro Ile Val Met Met
 20 25 30

gag cct cat cat ctc ctt atg aac tgg aac aaa cct att gat ctc att 144
 Glu Pro His His Leu Leu Met Asn Trp Asn Lys Pro Ile Asp Leu Ile
 35 40 45

aca caa gaa aac tct ttt aac cac aat cct cat ttc atg gta gat cca 192
 Thr Gln Glu Asn Ser Phe Asn His Asn Pro His Phe Met Val Asp Pro
 50 55 60

cct tcc gaa acc cta agc cac ttc cag ccc ccg ccg aca gtc ttc tcc 240
 Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro Thr Val Phe Ser
 65 70 75 80

gat ccc gga gga gga gag gaa gca gaa gac gaa gaa gga gag gaa gag 288
 Asp Pro Gly Gly Gly Glu Glu Ala Glu Asp Glu Glu Gly Glu Glu Glu
 85 90 95

ata gat gag atg aag gag atg caa tac gcg att gct gcc atg cag ccc 336
 Ile Asp Glu Met Lys Glu Met Gln Tyr Ala Ile Ala Ala Met Gln Pro
 100 105 110

gta gac atc gat cca gcc acc gtt cct aag ccg aac cgc cgt aac gta 384
 Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro Asn Arg Arg Asn Val
 115 120 125

agg gta agc gag gac ccc cag acg gtg gtg gct cgt cgg cgt aga gaa 432
 Arg Val Ser Glu Asp Pro Gln Thr Val Val Ala Arg Arg Arg Arg Glu
 130 135 140

agg ata agc gag aag atc cgg ata ttg aag agg atg gtg cca ggc ggt 480
 Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met Val Pro Gly Gly
 145 150 155 160

gca aag atg gac act gcc tcc atg ctt gac gaa gcc atc cgc tac acc 528
 Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala Ile Arg Tyr Thr
 165 170 175

aag ttc ttg aaa cgg cag gtg agg ctt ctt cag cct cac act cag ctt 576
 Lys Phe Leu Lys Arg Gln Val Arg Leu Leu Gln Pro His Thr Gln Leu
 180 185 190

ggg gct cct atg tct gac cct tct cgc ctt tgt tat tac cac aac tcg 624
 Gly Ala Pro Met Ser Asp Pro Ser Arg Leu Cys Tyr Tyr His Asn Ser
 195 200 205

gat acc taa 633
 Asp Thr
 210

<210> 4
 <211> 210
 <212> PRT
 <213> Brassica napus wild type IND-C1 coding sequence
 <400> 4

Met Tyr Lys Arg Lys Val Tyr Ala Ser Leu Val Gln Lys Leu Tyr Met
 1 5 10 15

Ser Gly Ser Lys Ala Asp Ala Ala Ala Ile Ala Pro Ile Val Met Met

20 25 30
 Glu Pro His His Leu Leu Met Asn Trp Asn Lys Pro Ile Asp Leu Ile
 35 40 45
 Thr Gln Glu Asn Ser Phe Asn His Asn Pro His Phe Met Val Asp Pro
 50 55 60
 Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro Thr Val Phe Ser
 65 70 75 80
 Asp Pro Gly Gly Gly Glu Glu Ala Glu Asp Glu Glu Gly Glu Glu Glu
 85 90 95
 Ile Asp Glu Met Lys Glu Met Gln Tyr Ala Ile Ala Ala Met Gln Pro
 100 105 110
 Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro Asn Arg Arg Asn Val
 115 120 125
 Arg Val Ser Glu Asp Pro Gln Thr Val Val Ala Arg Arg Arg Arg Glu
 130 135 140
 Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met Val Pro Gly Gly
 145 150 155 160
 Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala Ile Arg Tyr Thr
 165 170 175
 Lys Phe Leu Lys Arg Gln Val Arg Leu Leu Gln Pro His Thr Gln Leu
 180 185 190
 Gly Ala Pro Met Ser Asp Pro Ser Arg Leu Cys Tyr Tyr His Asn Ser
 195 200 205
 Asp Thr
 210

<210> 5
 <211> 1622
 <212> DNA
 <213> Brassica napus wild type IND-A1 genomic sequence

<220>
 <221> CDS
 <222> (561)..(1118)

<400> 5
 ttgacaatc tacatacata accaacaaaa agtagaatac cttgaaaatc taaaacccaa 60
 aatgatgtgtaaactcaag cttgggtccag agcataaaaa aattaaagcc atcgctttgg 120
 taccacatat ttaaactgca gttttttttt tttttttggg gggggggggg ggggtaatat 180
 aaaaatataa ttaacaaaaa aaaattatga aacaattagc atgtaaaaaca ctaatctttt 240

ggttgtgaca aaacgttttc acaaatgttc tataaataaa ttcaagtgca ttttatctgc 300
aaaatatata ctttactca taaaataaga gcgtttaaaa cattcataca cgcactacat 360
tgacatgaca aaagaaatcc gcaaatcac atgatgtatg tcgaaaaaaa caaaaaatac 420
acatgatgta tatatagaga ggatagtatc taggaaataa gactatatta tatatataaa 480
gaaaatagag aaaagataaa aatataaatt ggtatgtata aaagaaaggc ctatgctct 540
ctagtccaaa aactctatat atg tct ggc tca aaa gca gat gca gcc ata gcc 593
Met Ser Gly Ser Lys Ala Asp Ala Ala Ile Ala
1 5 10
cca ata gtc atg atg gag cat cat cat ctc ctt atg aat tgg aac aaa 641
Pro Ile Val Met Met Glu His His His Leu Leu Met Asn Trp Asn Lys
15 20 25
cct att gat ctc att aca gaa gaa aac tct ttt aac cac aat cct cat 689
Pro Ile Asp Leu Ile Thr Glu Glu Asn Ser Phe Asn His Asn Pro His
30 35 40
ttc ata gta gat cca cct tcc gaa acc cta agc cac ttc cag ccc ccg 737
Phe Ile Val Asp Pro Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro
45 50 55
ccg aca atc ttc tcc gat cac gga gga gga gag gaa gca gaa gaa gaa 785
Pro Thr Ile Phe Ser Asp His Gly Gly Gly Glu Glu Ala Glu Glu Glu
60 65 70 75
gaa gaa gaa gaa gga gag gaa gag atg gat ccg atg aag aag atg caa 833
Glu Glu Glu Glu Gly Glu Glu Glu Met Asp Pro Met Lys Lys Met Gln
80 85 90
tac gcg att gct gcc atg cag ccc gta gac ctc gat cca gcc acc gtt 881
Tyr Ala Ile Ala Ala Met Gln Pro Val Asp Leu Asp Pro Ala Thr Val
95 100 105
cct aag ccg aac cgc cgt aac gta agg gta agc gac gac cct cag acg 929
Pro Lys Pro Asn Arg Arg Asn Val Arg Val Ser Asp Asp Pro Gln Thr
110 115 120
gtg gtg gct cgt cgg cgt aga gaa agg ata agc gag aag atc cgg ata 977
Val Val Ala Arg Arg Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile
125 130 135
ttg aag agg atg gtg cca ggc ggt gca aag atg gac act gcc tcc atg 1025
Leu Lys Arg Met Val Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met
140 145 150 155
ctc gac gaa gcc atc cgc tac acc aag ttc ttg aaa cgg cag gtg agg 1073
Leu Asp Glu Ala Ile Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg
160 165 170
cta gct tct tca gcc tca cac tca gct tgg agc tcc tat gtc tga 1118
Leu Ala Ser Ser Ala Ser His Ser Ala Trp Ser Ser Tyr Val
175 180 185
cccttcttgc ctttgttatt accacaactc ggatacctaa ttataattct atcacgcgtt 1178
tcatgttgat atatatagat aaatggtcga ataaggattt cgatcgaaga ttgtatgtac 1238
aataaatgat gtgtgtatct caattaatgt atgatatata tatatatatg tatgcagtat 1298
gcatttatat tctattctct ataaggaggc aacattgccg gattagggct ttgatcttat 1358
gcaagttttc cgaccaaaaa tatgaaatac ttgtttggat ataacatatg aatcggataa 1418

gtgttactag ttatataact ggaaaacaaa tgtctggaat aagaattccc gggagaacca 1478
 agcctttctc taatccctaa gattatagct actgaaacaa tgaacaatg aagaatcagt 1538
 tgggcattag taaaaaaaaa agaatcagtt gggttgctta taaaattttg ttataaaatt 1598
 tatgtcgtat gtgtgtagc cgta 1622

<210> 6
 <211> 185
 <212> PRT
 <213> Brassica napus wild type IND-A1 genomic sequence
 <400> 6

Met Ser Gly Ser Lys Ala Asp Ala Ala Ile Ala Pro Ile Val Met Met
 1 5 10 15
 Glu His His His Leu Leu Met Asn Trp Asn Lys Pro Ile Asp Leu Ile
 20 25 30
 Thr Glu Glu Asn Ser Phe Asn His Asn Pro His Phe Ile Val Asp Pro
 35 40 45
 Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro Thr Ile Phe Ser
 50 55 60
 Asp His Gly Gly Gly Glu Glu Ala Glu Glu Glu Glu Glu Glu Gly
 65 70 75 80
 Glu Glu Glu Met Asp Pro Met Lys Lys Met Gln Tyr Ala Ile Ala Ala
 85 90 95
 Met Gln Pro Val Asp Leu Asp Pro Ala Thr Val Pro Lys Pro Asn Arg
 100 105 110
 Arg Asn Val Arg Val Ser Asp Asp Pro Gln Thr Val Val Ala Arg Arg
 115 120 125
 Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met Val
 130 135 140
 Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala Ile
 145 150 155 160
 Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg Leu Ala Ser Ser Ala
 165 170 175
 Ser His Ser Ala Trp Ser Ser Tyr Val
 180 185

<210> 7
 <211> 1593
 <212> DNA
 <213> Brassica napus wild type IND-C1 genomic sequence

<220>
 <221> CDS
 <222> (497)..(1126)

<400> 7
 tgccatacat aaccacggat catagtcgac acctcaacgt gaagcaaatt tgacaatcta 60
 catacataac caacaaaaag tagaataccg tgaaaacctt aaccctaaat atgatgtaaa 120
 actcaagctt ggtccagagc ataaaaaat taaagccatc gctttggat cacatattta 180
 aacgtcagtt ttttttggg gaagtaatat aaaaatataa ttaacaagaa aatttatgaa 240
 ataattagca tgtaaaacac tagtcttttg gttgtgacaa aacgttttca caaatgttct 300
 ataaataaat tcaagcacat tttatctgca aaatatatac tttcactcat aaaataagag 360
 cgtttaaaac attcatatac gcactacatt gacatgacaa aagaaatccg caaatacaaa 420
 catatttagt tcggatatat ctaggaaata agactatatt atatatataa agaaattaga 480
 aaaaaagaaa attggt atg tat aaa aga aag gtc tat gcg tct cta gtc caa 532
 Met Tyr Lys Arg Lys Val Tyr Ala Ser Leu Val Gln
 1 5 10
 aaa ctc tat atg tct ggt tca aaa gca gat gca gca gcc ata gcc cca 580
 Lys Leu Tyr Met Ser Gly Ser Lys Ala Asp Ala Ala Ala Ile Ala Pro
 15 20 25
 ata gtc atg atg gag cct cat cat ctc ctt atg aac tgg aac aaa cct 628
 Ile Val Met Met Glu Pro His His Leu Leu Met Asn Trp Asn Lys Pro
 30 35 40
 att gat ctc att aca caa gaa aac tct ttt aac cac aat cct cat ttc 676
 Ile Asp Leu Ile Thr Gln Glu Asn Ser Phe Asn His Asn Pro His Phe
 45 50 55 60
 atg gta gat cca cct tcc gaa acc cta agc cac ttc cag ccc ccg ccg 724
 Met Val Asp Pro Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro
 65 70 75
 aca gtc ttc tcc gat ccc gga gga gga gag gaa gca gaa gac gaa gaa 772
 Thr Val Phe Ser Asp Pro Gly Gly Gly Glu Glu Ala Glu Asp Glu Glu
 80 85 90
 gga gag gaa gag ata gat gag atg aag gag atg caa tac gcg att gct 820
 Gly Glu Glu Glu Ile Asp Glu Met Lys Glu Met Gln Tyr Ala Ile Ala
 95 100 105
 gcc atg cag ccc gta gac atc gat cca gcc acc gtt cct aag ccg aac 868
 Ala Met Gln Pro Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro Asn
 110 115 120
 cgc cgt aac gta agg gta agc gag gac ccc cag acg gtg gtg gct cgt 916
 Arg Arg Asn Val Arg Val Ser Glu Asp Pro Gln Thr Val Val Ala Arg
 125 130 135 140
 cgg cgt aga gaa agg ata agc gag aag atc cgg ata ttg aag agg atg 964
 Arg Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met
 145 150 155
 gtg cca ggc ggt gca aag atg gac act gcc tcc atg ctt gac gaa gcc 1012
 Val Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala
 160 165
 atc cgc tac acc aag ttc ttg aaa cgg cag gtg agg ctt ctt cag cct 1060
 Ile Arg Tyr Thr Lys Phe Leu Lys Arg Gln Val Arg Leu Leu Gln Pro
 175 180 185

cac act cag ctt ggg gct cct atg tct gac cct tct cgc ctt tgt tat 1108
His Thr Gln Leu Gly Ala Pro Met Ser Asp Pro Ser Arg Leu Cys Tyr
190 195 200
tac cac aac tcg gat acc taattataat tctatcacgc gtttcatggt 1156
Tyr His Asn Ser Asp Thr
205 210
gatatatata gataaatggt tgaataagga tttcgatcga agattgtatg gctattgatt 1216
acattatata ttgtacaata aatgatgtgt gtatttctat taatgtatat atgatatata 1276
tctgtttgca gtatgcattt atattctatt ctttataggg aggcaacatg ccggattagg 1336
gctttgatcg tatgcaagtt ttccgaccaa aaatatgaaa tacttgtttg gatataacat 1396
atgaatcgga taagtgttac tagttatata actggaaaaa attgtttggt ataagaattc 1456
ccgggagaac caagcctttc tctaaccct aagatcatag ctactgaaat aatgaaaaaa 1516
aacaaaaaaa aaacaatgaa gaatcagttg ggcattagtc caaaaaaaaa aaagaatcag 1576
ttggattgct tataaaa 1593

<210> 8
<211> 210
<212> PRT
<213> Brassica napus wild type IND-C1 genomic sequence
<400> 8

Met Tyr Lys Arg Lys Val Tyr Ala Ser Leu Val Gln Lys Leu Tyr Met
1 5 10 15
Ser Gly Ser Lys Ala Asp Ala Ala Ala Ile Ala Pro Ile Val Met Met
20 25 30
Glu Pro His His Leu Leu Met Asn Trp Asn Lys Pro Ile Asp Leu Ile
35 40 45
Thr Gln Glu Asn Ser Phe Asn His Asn Pro His Phe Met Val Asp Pro
50 55 60
Pro Ser Glu Thr Leu Ser His Phe Gln Pro Pro Pro Thr Val Phe Ser
65 70 75 80
Asp Pro Gly Gly Gly Glu Glu Ala Glu Asp Glu Glu Gly Glu Glu Glu
85 90 95
Ile Asp Glu Met Lys Glu Met Gln Tyr Ala Ile Ala Ala Met Gln Pro
100 105 110
Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro Asn Arg Arg Asn Val
115 120 125
Arg Val Ser Glu Asp Pro Gln Thr Val Val Ala Arg Arg Arg Arg Glu
130 135 140

Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Met Val Pro Gly Gly
145 150 155 160

Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala Ile Arg Tyr Thr
165 170 175

Lys Phe Leu Lys Arg Gln Val Arg Leu Leu Gln Pro His Thr Gln Leu
180 185 190

Gly Ala Pro Met Ser Asp Pro Ser Arg Leu Cys Tyr Tyr His Asn Ser
195 200 205

Asp Thr
210

<210> 9
<211> 597
<212> DNA
<213> Arabidopsis thaliana IND1

<220>
<221> CDS
<222> (1)..(594)

<400> 9
atg gaa aat ggt atg tat aaa aag aaa gga gtg tgc gac tct tgt gtc 48
Met Glu Asn Gly Met Tyr Lys Lys Lys Gly Val Cys Asp Ser Cys Val
1 5 10 15
tcg tcc aaa agc aga tcc aac cac agc ccc aaa aga agc atg atg gag 96
Ser Ser Lys Ser Arg Ser Asn His Ser Pro Lys Arg Ser Met Met Glu
20 25 30
cct cag cct cac cat ctc ctc atg gat tgg aac aaa gct aat gat ctt 144
Pro Gln Pro His His Leu Leu Met Asp Trp Asn Lys Ala Asn Asp Leu
35 40 45
ctc aca caa gaa cac gca gct ttt ctc aat gat cct cac cat ctc atg 192
Leu Thr Gln Glu His Ala Ala Phe Leu Asn Asp Pro His His Leu Met
50 55 60
tta gat cca cct ccc gaa acc cta att cac ttg gac gaa gac gaa gag 240
Leu Asp Pro Pro Pro Glu Thr Leu Ile His Leu Asp Glu Asp Glu Glu
65 70 75 80
tac gat gaa gac atg gat gcg atg aag gag atg cag tac atg atc gcc 288
Tyr Asp Glu Asp Met Asp Ala Met Lys Glu Met Gln Tyr Met Ile Ala
85 90 95
gtc atg cag ccc gta gac atc gac cct gcc acg gtc cct aag ccg aac 336
Val Met Gln Pro Val Asp Ile Asp Pro Ala Thr Val Pro Lys Pro Asn
100 105 110
cgc cgt aac gta agg ata agc gac gat cct cag acg gtg gtt gct cgt 384
Arg Arg Asn Val Arg Ile Ser Asp Asp Pro Gln Thr Val Val Ala Arg
115 120 125
cgg cgt cgg gaa agg atc agc gag aag atc cga att ctc aag agg atc 432
Arg Arg Arg Glu Arg Ile Ser Glu Lys Ile Arg Ile Leu Lys Arg Ile
130 135 140
gtg cct ggt ggt gcg aag atg gac aca gct tcc atg ctc gac gaa gcc 480
Val Pro Gly Gly Ala Lys Met Asp Thr Ala Ser Met Leu Asp Glu Ala

145					150					155					160	
ata	cgt	tac	acc	aag	ttc	ttg	aaa	cgg	cag	gtg	agg	att	ctt	cag	cct	528
Ile	Arg	Tyr	Thr	Lys	Phe	Leu	Lys	Arg	Gln	Val	Arg	Ile	Leu	Gln	Pro	
				165					170					175		
cac	tct	cag	att	gga	gct	cct	atg	gct	aac	ccc	tct	tac	ctt	tgt	tat	576
His	Ser	Gln	Ile	Gly	Ala	Pro	Met	Ala	Asn	Pro	Ser	Tyr	Leu	Cys	Tyr	
			180					185					190			
tac	cac	aac	tcc	caa	ccc	tga										597
Tyr	His	Asn	Ser	Gln	Pro											
		195														

<210> 10
 <211> 198
 <212> PRT
 <213> Arabidopsis thaliana IND1

<400> 10

Met	Glu	Asn	Gly	Met	Tyr	Lys	Lys	Lys	Gly	Val	Cys	Asp	Ser	Cys	Val
1				5					10					15	
Ser	Ser	Lys	Ser	Arg	Ser	Asn	His	Ser	Pro	Lys	Arg	Ser	Met	Met	Glu
			20					25					30		
Pro	Gln	Pro	His	His	Leu	Leu	Met	Asp	Trp	Asn	Lys	Ala	Asn	Asp	Leu
		35					40					45			
Leu	Thr	Gln	Glu	His	Ala	Ala	Phe	Leu	Asn	Asp	Pro	His	His	Leu	Met
	50					55					60				
Leu	Asp	Pro	Pro	Pro	Glu	Thr	Leu	Ile	His	Leu	Asp	Glu	Asp	Glu	Glu
65					70					75					80
Tyr	Asp	Glu	Asp	Met	Asp	Ala	Met	Lys	Glu	Met	Gln	Tyr	Met	Ile	Ala
				85					90					95	
Val	Met	Gln	Pro	Val	Asp	Ile	Asp	Pro	Ala	Thr	Val	Pro	Lys	Pro	Asn
			100					105					110		
Arg	Arg	Asn	Val	Arg	Ile	Ser	Asp	Asp	Pro	Gln	Thr	Val	Val	Ala	Arg
		115					120					125			
Arg	Arg	Arg	Glu	Arg	Ile	Ser	Glu	Lys	Ile	Arg	Ile	Leu	Lys	Arg	Ile
	130					135					140				
Val	Pro	Gly	Gly	Ala	Lys	Met	Asp	Thr	Ala	Ser	Met	Leu	Asp	Glu	Ala
145					150					155					160
Ile	Arg	Tyr	Thr	Lys	Phe	Leu	Lys	Arg	Gln	Val	Arg	Ile	Leu	Gln	Pro
				165					170					175	
His	Ser	Gln	Ile	Gly	Ala	Pro	Met	Ala	Asn	Pro	Ser	Tyr	Leu	Cys	Tyr
			180					185					190		

Tyr His Asn Ser Gln Pro
195

<210> 11
<211> 643
<212> DNA
<213> Nucleotide sequence of a INDEHISCENT homologue from Brassica napus
(Bn1-IND - SEQ ID NO: 2 of W004/113542)

<400> 11
gaattcgccc ttcgcatgta taaaaagaag ggtctatgcg tctctagtcc aaaaactcta 60
tatgtctggt tcaaaagcag atgcagcagc catagcccca atagtcatga tggagcctca 120
tcattctcctt atgaactgga acaaacctat tgatctcatt acacaagaaa actcttttaa 180
ccacaatcct catttcatgg tagatccacc ttccgaaacc ctaagccact tccagccccc 240
gccgacagtc ttctccgatc ccggaggagg agaggaagca gaagacgaag aaggagagga 300
agagatagat gagatgaagg agatgcaata cgcgattgct gccatgcagc ccgtagacat 360
cgatccagcc accgttccta agccgaaccg ccgtaacgta agggtaagcg aggaccccca 420
gacggtggtg gctcgtcggc gtagagaaag gataagcag aagatccgga tattgaagag 480
gatggtgcca ggcggtgcaa agatggacac tgcctccatg cttgacgaag ccatccgcta 540
caccaagttc ttgaaacggc aggtgaggct tcttcagcct cacactcagc ttggggctcc 600
tatgtctgac ctttctcgcc tttgttatta ccacaactct caa 643

<210> 12
<211> 660
<212> DNA
<213> Nucleotide sequence of a second INDEHISCENT homologue from Brassica napus
(Bn2-IND - SEQ ID NO: 3 of W004/113542)

<400> 12
gaattcgccc ttggcatgta caagaagaaa ggtctatgcg tctctagtcc aaaaactcta 60
tatatgtctg gctcaaaagc agatgcagcc atagcccca tagtcatgat ggagcatcat 120
catctcctta tgaattgga caaacctatt gatctcatta cagaagaaaa ctcttttaac 180
cacaatcctc atttcatagt agatccacct tccgaaacct taagccactt ccagcccccg 240
ccgacaatct tctccggtca cggaggagga gaggaagcag cagaagaaga agaagaagaa 300
ggagaggaag agatggatcc gatgaagaag atgcaatacg cgattgctgc catgcagccc 360
gtagacctcg atccagccac cgttcctaag ccgaaccgcc gtaacgtaag ggtaagcgac 420
gaccctcaga cggtggtggc tcgtcggcgt agagaaagga taagcgagaa gatccggata 480
ttgaggagga tgggtgccagg cggtgcaaag atggacactg cctccatgct cgacgaagcc 540
atccgctaca ccaagttctt gaaacggcag gtgaggctag cttcttcagc ctcaactca 600
gcttgagct cctatgtctg acccttcttg cctttgttat taccataact cgcagccctg 660

<210> 13
<211> 20
<212> DNA
<213> Artificial

<220>
 <223> forward oligonucleotide for detection of IND-A1-EMS01
 <400> 13
 aagggtaagc gacgaccctt 20

<210> 14
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> Forward oligonucleotide for detection of IND-A1-WT
 <400> 14
 aagggtaagc gacgaccctc 20

<210> 15
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> Reverse oligonucleotide for detection of IND-A1-EMS01 and -WT
 <400> 15
 gagtgtgagg ctgaagaagc 20

<210> 16
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> Forward oligonucleotide for detection of IND-A1-EMS05
 <400> 16
 cctcagacgg tggtggctca 20

<210> 17
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> Forward oligonucleotide for detection of IND-A1-WT
 <400> 17
 cctcagacgg tggtggctcg 20

<210> 18
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> Reverse oligonucleotide for detection of IND-A1-EMS05 and -WT
 <400> 18
 agggtcagac ataggagctc 20

<210> 19

<211> 21
 <212> DNA
 <213> Artificial

 <220>
 <223> Reverse oligonucleotide for detection of IND-C1-EMS01

 <400> 19
 gtggttaaaa gagttttctt a 21

 <210> 20
 <211> 21
 <212> DNA
 <213> Artificial

 <220>
 <223> Reverse oligonucleotide for detection of IND-C1-WT

 <400> 20
 gtggttaaaa gagttttctt g 21

 <210> 21
 <211> 20
 <212> DNA
 <213> Artificial

 <220>
 <223> Forward oligonucleotide for detection of IND-C1-EMS01 and -WT

 <400> 21
 attagcatgt aaaacactag 20

 <210> 22
 <211> 19
 <212> DNA
 <213> Artificial

 <220>
 <223> Reverse oligonucleotide for detection of IND-C1-EMS03

 <400> 22
 acgagccacc accgtctag 19

 <210> 23
 <211> 18
 <212> DNA
 <213> Artificial

 <220>
 <223> Reverse oligonucleotide for detection of IND-C1-WT

 <400> 23
 acgagccacc accgtctg 18

 <210> 24
 <211> 21
 <212> DNA
 <213> Artificial

 <220>
 <223> Forward oligonucleotide for detection of IND-C1-EMS03 and -WT

 <400> 24
 gttcaaaagc agatgcagca g 21

<210> 25
 <211> 22
 <212> DNA
 <213> Artificial

 <220>
 <223> oligonucleotide for detection of IND-A1-EMS01 and -WT

 <400> 25
 gccgacgagc caccaccgtc tt 22

 <210> 26
 <211> 14
 <212> DNA
 <213> Artificial

 <220>
 <223> oligonucleotide for detection of IND-A1-EMS01

 <400> 26
 aagggtcgtc gctt 14

 <210> 27
 <211> 13
 <212> DNA
 <213> Artificial

 <220>
 <223> oligonucleotide for detection of IND-A1-WT

 <400> 27
 gagggtcgtc gct 13

 <210> 28
 <211> 34
 <212> DNA
 <213> Artificial

 <220>
 <223> oligonucleotide for detection of IND-A1-EMS05 and -WT

 <400> 28
 cggatcttct cgcttatcct ttctctacgc cgaa 34

 <210> 29
 <211> 13
 <212> DNA
 <213> Artificial

 <220>
 <223> oligonucleotide for detection of IND-A1-EMS05

 <400> 29
 tgagccacca ccg 13

 <210> 30
 <211> 13
 <212> DNA
 <213> Artificial

 <220>
 <223> oligonucleotide for detection of IND-A1-WT

<400> 30
cgagccacca ccg 13

<210> 31
<211> 48
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide for detection of IND-C1-EMS01 and -WT

<400> 31
aggtgatct accatgaaat gaggattgtg gttaaaagag ttttcttt 48

<210> 32
<211> 23
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide for detection of IND-C1-EMS01

<400> 32
atgtaatgag atcaataggt ttg 23

<210> 33
<211> 23
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide for detection of IND-C1-WT

<400> 33
gtgtaatgag atcaataggt ttg 23

<210> 34
<211> 29
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide for detection of IND-C1-EMS03 and -WT

<400> 34
ccgtaacgta agggaagcg aggacccca 29

<210> 35
<211> 14
<212> DNA
<213> Artificial

<220>
<223> oligonucleotide for detection of IND-C1-EMS03

<400> 35
tagacggtgg tggc 14

<210> 36
<211> 14
<212> DNA
<213> Artificial

<220>
 <223> oligonucleotide for detection of IND-C1-WT

 <400> 36
 cagacggtgg tggc 14

<210> 37
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> Forward oligonucleotide for detection of IND-A1

 <400> 37
 aggagaggaa gagatggatc c 21

<210> 38
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> Reverse oligonucleotide for detection of IND-A1

 <400> 38
 tgagtgtgag gctgaagaag c 21

<210> 39
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> Forward oligonucleotide for detection of IND-C1

 <400> 39
 cctcatcatc tccttatgaa c 21

<210> 40
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> Reverse oligonucleotide for detection of IND-C1

 <400> 40
 cgtattgcat ctccttcatc t 21