



US 20090175908A1

(19) **United States**(12) **Patent Application Publication**
Yang et al.(10) **Pub. No.: US 2009/0175908 A1**(43) **Pub. Date: Jul. 9, 2009**(54) **INFLUENZA HEMAGGLUTININ AND
NEURAMINIDASE VARIANTS**(60) Provisional application No. 60/479,078, filed on Jun.
16, 2003.(75) Inventors: **Chin-Fen Yang**, San Jose, CA
(US); **George Kemble**, Fremont,
CA (US); **Chongguang Liu**,
Fremont, CA (US)Correspondence Address:
MEDIMMUNE, LLC
Jonathan Klein-Evans
ONE MEDIMMUNE WAY
GAITHERSBURG, MD 20878 (US)(73) Assignee: **Medimmune, LLC**, Gaithersburg,
MD (US)(21) Appl. No.: **12/399,077**(22) Filed: **Mar. 6, 2009****Related U.S. Application Data**(62) Division of application No. 10/870,690, filed on Jun.
16, 2004.**Publication Classification**(51) **Int. Cl.****A61K 39/145** (2006.01)
C07K 14/00 (2006.01)
C07H 21/04 (2006.01)
C12N 7/01 (2006.01)
C12N 15/63 (2006.01)
C12N 5/10 (2006.01)
C12N 7/02 (2006.01)
A61K 38/16 (2006.01)
A61K 31/7088 (2006.01)
A61P 31/16 (2006.01)(52) **U.S. Cl. 424/206.1; 530/350; 536/23.1;**
435/235.1; 435/320.1; 435/365; 435/239;
514/12; 514/44(57) **ABSTRACT**Polypeptides, polynucleotides, methods, compositions, and
vaccines comprising influenza hemagglutinin and neuramini-
dase variants are provided.

Figure 1A

SEQ ID NO:1

ca A/Shandong/9/93

Nucleotide Sequence of ca_A_Shandong_9_93_HA

Entire molecule length: 1745 bp

```
1  caggggataa  ttctattaac  catgaagact  atcattgctt  tgagctacat
51  tttatgtctg  gttttcgctc  aaaaacttcc  cggaaatgac  aacagcacag
101 caacgctgtg  cctgggacat  catgcagtgc  caaacggaac  gctagtgaaa
151 acaatcacga  atgatcaaat  tgaagtgact  aatgctactg  agttggttca
201 gagttcctca  acaggtagaa  tatgcggcag  tcctcaccga  atccttgatg
251 gaaaaaactg  cacactgata  gatgctctat  tgggagaccc  tcattgtgat
301 ggcttcctca  ataaggaatg  ggaccttttt  gttgaacgca  gcaaagctta
351 cagcaactgt  tacccttatg  atgtgccgga  ttatgcctcc  cttcaggctac
401 tagttgcctc  atcaggcacc  ctggagttta  tcaatgaaga  cttcaattgg
451 actggagtcg  ctcaggatgg  gggaagctat  gcttgcaaaa  gaggatctgt
501 taacagtttc  tttagtagat  tgaattgggt  gcacaaatta  gaatacaaat
551 atccagcgct  gaacgtgact  atgccaaaca  atggcaaatt  tgacaaattg
601 tacatttggg  gggttcacca  cccgagcacg  gacagtgacc  aaaccagcct
651 atatgttcga  gcatcaggga  gagtcacagt  ctctaccaa  agaagccaac
701 aaactgtaac  ccgaatatc  gggctctagc  cctgggtaag  gggtcagtcc
751 agtagaataa  gcatctattg  gacaatagta  aaaccgggag  acatactttt
801 gattaatagc  acagggaatc  taattgctcc  tcggggttac  ttcaaaatac
851 gaaatgggaa  aagctcaata  atgaggtcag  atgcacccat  tggcaactgc
901 agttctgaat  gcatcactcc  aaatggaagc  attcccaatg  acaaaccttt
951 tcaaaatgta  aacagaatca  catatggggc  ctgcccaga  tatgttaagc
1001 aaaacactct  gaaattggca  acagggatgc  ggaatgtacc  agagaaacaa
1051 actagaggca  tattcggcgc  aatcgcagg  ttcatagaaa  atggttggga
1101 gggaatggta  gacggttggt  acggtttcag  gcatcaaaat  tctgagggca
1151 caggacaagc  agcagatctt  aaaagcactc  aagcagcaat  cgaccaaatc
1201 aacgggaaac  tgaatagggt  aatcgagaaa  acgaacgaga  aattccatca
1251 aatcgaaaaa  gaattctcag  aagtagaagg  gagaattcag  gacctcgaga
1301 aatatgttga  agacactaaa  atagatctct  ggtcttaca  cgcgagcctt
1351 cttgttgccc  tggagaacca  acatacaatt  gatctaactg  actcagaaat
1401 gaacaaactg  tttgaaaaaa  caaggaaagc  actgagggaa  aatgctgagg
1451 acatgggcaa  tggttgcttc  aaaatatacc  acaaattgtg  caatgcctgc
1501 atagggtaa  tcagaaatgg  aacttatgac  catgatgtat  acagagacga
1551 agcattaaac  aaccggttcc  agatcaaagg  tgttgagctg  aagtcaggat
1601 acaaagattg  gatcctatgg  atttcctttg  ccataatcat  ctttttgctt
1651 tgtgttggtt  tgctgggggt  catcatgtgg  gcttgccaaa  aaggcaacat
1701 taggtgcaac  atttgcatth  gagggcatta  attaaaaaca  ccctg
```

SEQ ID NO:35

Amino Acid Sequence of ca_A_Shandong_9_93_HA Entire molecule length: 566 aa

```
1  mktiialsyi  lclvfaqklp  gndnstatlc  lghhavpngt  lvkttindqi
51  evtnatelvq  ssstgricgs  phrildgknc  tlidallgdp  hodgfgnkew
101 dlfferskay  sncypydvdp  yaslrslvas  sgtlefined  fnwtgvaqdg
151 gsyackrgsv  nsffsrlnwl  hkleykypal  nvtmpnngkf  dklyiwgvhh
201 pstdsdqtsl  yvrasgrvtv  stkrsgqvtv  pnigrspwvr  gqsrisiyw
251 tivkpgdill  instgnliap  rgyfkirngk  ssimrsdapi  gncssecitp
301 ngsipndkpf  qnvnrityga  cpryvkqntl  klatgmrnvp  ekqtrgifga
351 iagfiengwe  gmvdgwygfr  hqnseggtga  adlkstqaai  dqingklrnl
401 iektnekfhq  iekefseveg  riqdlekyve  dtkidlwsyn  aellvalenq
451 htidltdsem  nklfektrkq  lrenaedmgn  gcfskiyhkd  nacigsirng
501 tydhdvyrde  alnnrfqikg  velksygdw  ilwisfaisc  fllocvllgf
551 imwacqkgni  rcnici
```

Figure 1B**SEQ ID NO:2**

Nucleotide Sequence of ca_A_Shandong_9_93_NA Entire molecule length: 1429 bp

```
1 aaagataata acaattggct ctgtttctct cactattgcc acaatatgct
51 tccttatgca aattgccatc ctggaacta ctgtaacatt gcacttcaag
101 caatatgagt gcaactcccc cccaaacaac caagtaatgc tgtgtgaacc
151 aacaataata gaaagaaaca taacagagat agtgtatctg accaacacca
201 ccatagagaa agaaatatgc cccaaactag cagaatacag aaattgggtca
251 aagccgcaat gtaaaattac aggatttgca cctttttcta aggacaattc
301 aattcggctt tcagctggtg gagacatttg ggtgacaaga gaaccttatg
351 tgtcatgcga tcctggcaag tgttatcaat ttgcccttgg acagggaaca
401 aactaaaca acaggcactc aaatgacaca gtacatgata ggaccctta
451 tcgaacccta ttgatgaatg agttgggtgt tccatttcac ttgggaacca
501 agcaagtgtg catagcatgg tccagctcaa gttgtcacga tggaaaagca
551 tggctgcatg tttgtgtaac tgggcatgat gaaaatgcaa ctgctagctt
601 catttacgat gggaggcttg tagatagtat tggttcatgg tccaaaaata
651 tcctcaggac ccaggagtcg gaatgcgttt gtatcaatgg aacttgtaca
701 gtagtaatga ctgatggaag tgcttcagaa agagctgata ctaaaatact
751 attcattgaa gaggggaaaa tcgttcatat tagcccatgg tcaggaagtg
801 ctcagcatgt cgaggagtgc tcctgttatc ctcgatatcc tgggtgcaga
851 tgtgtctgca gagacaactg gaaaggctcc aataggcca tcgtagatat
901 aaatgtgaaa gattatagca ttgtttccag ttatgtgtgc tcaggacttg
951 ttggagacac acccagaaaa aacgacagct ccagcagtag ctattgccgg
1001 aatcctaaca atgagaaagg gagtcatgga gtgaaaggct gggcctttga
1051 tgatggaaat gacgtgtgga tgggaagaac gatcagcgag gagttacgct
1101 caggttatga aaccttcaaa gtcattggag gctggtccaa acctaaactc
1151 aaattgcaga taaataggca agtcatagtt gacagaggta ataggctccg
1201 ttattcttgt attttctctg ttgaaggcaa aagctgcac aatcgggtgct
1251 tttatgtgga gttgataagg ggaaggaaac aggaaactga agtctgggtg
1301 acctcaaaca gtattgttgt gttttgtggc acctcaggta catatggaac
1351 aggctcatgg ccctgatggg gcggacatca atctcatgcc tatataagct
1401 ttcgcaatth tagaaaaaaa ctccttggt
```

SEQ ID NO:36

Amino Acid Sequence of ca_A_Shandong_9_93_NA Entire molecule length: 436 aa

```
1 mqiaailvttv tlhfkqyecn sppnnqvmle eptiiernit eivylnttti
51 ekeicpklae yrnwskpqck itgfapfskd nsirlsaggd iwtrepvys
101 cdpqkcyqfa lgqgttlennr hndtvhdrt pyrtllmnel gvpfhlgtkq
151 vciaawssssc hdgkawlhvc vtghdenata sflydgrlvd sigswsknil
201 rtqesecvci ngtctvmtgd gsaseradtk ilfieegkiv hisplsgsaq
251 hveecscypv ypgvrcvcrd nwkgnsrpiv dinvkdisiv ssyvcsglv
301 dtprkndsss ssycrnnpne kgshgvgkwa fddgndvwmg rtiseelrsg
351 yetfkviggw skpnsklqin rqvivdrngr sgysgifsve gkscinrcfy
401 velirgrkqe tevwwtsnsi vvfctsgty gtgswp
```

Figure 1C

SEQ ID NO:3

ca A/Johannesburg/33/94-Like

Nucleotide Sequence of ca_A_Johannesburg_33_94_Like_HA Entire molecule length: 1755 bp

```
1 agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg
51 agctacattt tatgtctggt tttcgctcaa aaacttcccc gaaatgacaa
101 cagcacagca acgctgtgcc tgggacacca tgcagtgcc aacggaacgc
151 tagtgaaaac aatcacgaat gatcaaattg aagtgactaa tgctactgag
201 ctggttcaga gttccccaac aggtagaata tgcgacagcc ctcaccgaat
251 ccttgatgga aagaactgca cactgataga tgctctattg ggagaccctc
301 attgtgatgg cttccaaaat aaggaatggg acctttttgt tgaacgcagc
351 aaagcttaca gcaactgtta cccttatgat gtgccggatt atgcctccct
401 taggtcacta gttgcctcat caggcacccct ggagtttata aacgaaaact
451 tcaattggac tggagtgcgt caggatggga aaagctatgc ttgcaaaagg
501 ggatctgtta acagtttctt tagtagattg aattggttgc acaaattaga
551 atacaaatat ccagcgctga acgtgactat gccaaacaat ggcaaatttg
601 acaaatgtga catttggggg gttcaccacc cgagcacgga cagtgtccaa
651 accagcctat atgtccgagc atcagggaga gtcacagtct ctaccaaagg
701 aagccaacaa actgtaatcc cggatatcgg gtatagacca tgggtaaggg
751 gtcagtcag tagaataagc atctattgga caatagtaaa accgggagac
801 atacttttga ttaatagcac agggaatcta attgctcctc ggggttactt
851 caaaatacga aatgggaaaa gctcaataat gaggtcagat gcaccatttg
901 gcaactgcag ttctgaatgc atcactccaa atggaagcat tcccaatgac
951 aaaccttttc aaaatgtaaa caggatcaca tatggggcct gcccagata
1001 tgttaagcaa aacactctga aattggcaac agggatgcgg aatgtaccag
1051 agaacaacaa tagaggcata ttccggcgaa tcgcaggttt catagaaaat
1101 ggttgggagg gaatggtaga cggtttgtag ggtttcaggc atcaaaatcc
1151 tgagggcaca ggacaagctg cagatcttaa aagcactcaa gcagcaatcg
1201 accaaatcaa cgggaaactg aataggttag tcgagaaaac gaacgagaaa
1251 ttccatcaaa tcgaaaaaga attctcagaa gtagaaggga gaattcagga
1301 cctcgagaaa tatgttgaag aactaaaaat agatctctgg tcttacaatg
1351 cggaacttct tgttgctctg gagaaccaac atacaattga tctaactgac
1401 tcagaaatga acaaaactgt tgaaagaaca aggaagcaac tgagggaaaa
1451 tgctgaggac atgggcaatg gttgtttcaa aatataccac aaatgtgaca
1501 atgcctgcat aggttcaatc agaaatggaa cttatgacca tgatgtatac
1551 agagacgaag cattaacaaa ccggttccag atcaaagggtg ttgagctgaa
1601 gtcaggatac aaagattgga ttctatggat ttcccttgcc atatcgtgct
1651 ttttgctttg tgttgttttg cttgggttca tcatgtgggc ctgccaaaaa
1701 ggcaacatta ggtgcaacat ttgcatttga gtgcattaat taaaaacacc
1751 cttgt
```

SEQ ID NO:37

Amino Acid Sequence of ca_A_Johannesburg_33_94_Like_HA Entire molecule length: 566 aa

```
1 mktiialsyi lclvfaqklp gndnstatlc lghhavpngt lvkttitndqi
51 evtnatelvq ssptgricds phrildgknc tlidallgdp hcdgfnkew
101 dliverskay sncypydvdp yaslslvas sgtlefinen fnwtgvaqdg
151 ksyackrgsv nsffsrlnwl hkleykypal nvtmnpngkf dklyiwgvhh
201 pstdsvqtsl yvrasgrvtv stkrsgqgtvi pdigyrrpwr gqssrisiyw
251 tivkpgdill instgnliap rgyfkirngk ssimrsdapi gncssecitp
301 ngsipndkpf qnvnrityga cpryvkqntl klatgmrvnp ekqtrgifga
351 iagfiengwe gmvdgwygfr hqnseggtgqa adlkstqaai dqingklrnl
401 vektnekfhq iekefseveg riqdlekyve dtkidlwsyn aellvalenq
451 htidltdsem nklfertrkq lrenaedmgn gcfktyhkcd nacigsirng
501 tydhdvyrde alnnrfqikg velksykw dw ilwisfaisc fllecvtllgf
551 imwacqkgni rcnici
```

Figure 1D**SEQ ID NO:4**

Nucleotide Sequence of ca_A_Johannesburg_33_94_Like_NA

Entire molecule length: 1354 bp

```
1  gaaaatgaat  ccaaatacaa  agataataac  aattggctct  gtttctctca
51  ctattgccac  aatatgcttc  cttatgcaaa  ttgccatcct  ggtaactact
101  gtaacattgc  atttcaagca  atatgagtgc  aactcccccc  caaacaacca
151  agtaatgctg  tgtgaaccaa  caataataga  aagaaacata  acagagatag
201  tgtatctgac  caacaccacc  atagagaaaag  aaatatgccc  caaactagca
251  gaatacagaa  attggtcaaa  gccgcaatgt  aaaattacag  gatttgcacc
301  tttttctaa  gacaattcaa  ttcggctttc  cgctggtgga  gacatttggg
351  tgacaagaga  accttatgtg  tcatgcgatc  ctggcaagtg  ttatcaattt
401  gccctcggac  agggaacaac  actaaacaac  aggcattcaa  atgacacagt
451  acatgatagg  accccttata  gaaccctatt  gatgaatgag  ttgggtgttc
501  catttcattt  gggaaccaag  caagtgtgca  tagcatggtc  cagctcaagt
551  tgtcacgatg  gaaaagcatg  gctgcatgtt  tgtgtaactg  ggcattgatg
601  aatgcaact  gctagcttca  ttacgatgg  gaggcttgta  gatagtattg
651  gttcatggtc  caaaaatatc  ctacaggacc  aggagtcgga  atgcgtttgt
701  atcaatggaa  cttgtacagt  agtaatgact  gatggaagtg  cttcagaaag
751  agctgatact  aaaatactat  tcattgaaga  ggggaaaatc  gttcatatta
801  gccattgtc  aggaagtgtc  cagcatgtcg  aggagtgtc  ctgttatcct
851  cgatatcctg  gtgtcagatg  tgtctgcaga  gacaactgga  aaggctccaa
901  taggcccata  gtagatataa  atgtgaaaga  ttatagcatt  gtttccagtt
951  atgtgtgtc  aggacttggt  ggagacacac  ccagaaaaaa  cgacagctcc
1001  agcagtagct  attgctggaa  tcctaacaat  gagaaagggg  gtcattggagt
1051  gaaaggctgg  gcctttgatg  atggaaatga  cgtgtgggatg  ggaagaacga
1101  tcagcgagga  gttacgctca  gggttatgaa  ccttcaaagt  cattggaggc
1151  tgggtccaa  ctaactcaa  attgcagata  aataggcaag  tcatagttag
1201  cagaggtaat  aggtccggtt  attctggtat  tttctctgtt  gaaggcaaaa
1251  gctgcatcaa  tcggtgcttt  tatgtggagt  tgataagggg  aaggaaacag
1301  gaaactgaag  tctggtggac  ctcaaacagt  attgttgtgt  tttgtggcac
1351  ttca
```

SEQ ID NO:38

Amino Acid Sequence of ca_A_Johannesburg_33_94_Like_NA

Entire molecule length: 451 aa

```
1  kmnpnqkiit  igsvsltiat  icflmqiail  vttvtlhfkq  yecnsppnnq
51  vmlceptiie  rniteivylt  nttiekeicp  klaeyrnwsk  pqckitgfap
101  fskdnsirls  aggdiwvtre  pyvscdpgkc  yqfalgggtt  lnnrhnndtv
151  hdrtpyrll  mnelgvpfhl  gtkqvciaws  ssschdkgaw  lhvcvtghde
201  natasfiydg  rlvdsigsws  knilrtqese  cvcingtctv  vmdtgsaser
251  adtkilfiee  gkivhispls  gsaqhveecs  cyprypgvrc  vcrdnwkgss
301  rpivdinvk  ysivssyvcs  glvgdtprkn  dsssssywcn  pnnekggghv
351  kgwafddgnd  vwmgrtisee  lrsgyetfkv  iggwskpnsk  lqinrqvivid
401  rgnrsgysgi  fsvegkscin  rcfyvelirg  rkqetevvwt  snsivvcfgt
451  s
```

Figure 1E

SEQ ID NO:5

ca A/Wuhan/395/95

Nucleotide Sequence of ca A/Wuhan/395/95 H3

Entire molecule length: 1762 bp

```
1 agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg
51 agctacattt tatgtctggg ttctgctcaa aaacttcccg gaaatgacaa
101 cagcacggca acgctgtgcc tgggacacca tgcagtgcc aacggaacgc
151 tagtgaaaac aatcacgaat gaccaaattg aagtgactaa tgctactgag
201 ctgggttcaga gttcctcaac aggtagaata tgcgacagtc ctaccogaat
251 ccttgatgga aaaaactgca cactgataga tgctctattg ggagaccctc
301 attgtgatgg cttccaaaat aaggaatggg acctttttgt tgaacgcagc
351 aaagcttaca gcaactgtta cccttatgat gtgccggatt atgcttcctt
401 taggtcacta gttgcctcat ccggcaccct ggagtttacc aatgaaggct
451 tcaattggac tggagtcgct caggatggaa caagctatgc ttgcaaaagg
501 ggatctgtta aaagtctctt tagtagattg aattggttgc acaaattaga
551 atacaaatat ccagcactga acgtgactat gccaaaacaat gacaaatttg
601 acaaattgta catttggggg gtccaccacc cgagtacgga cagtgaccaa
651 accagcatat atgttcaagc atcagggaga gtcacagtct ctacaaaaag
701 aagccaacaa actgtaatcc cgaatatcgg gtctagaccc tgggtaaggg
751 ggatctccag cagaataagc atctattgga caatagtaaa accgggagac
801 atacttttga ttaacagcac agggaatcta attgctcctc ggggttactt
851 caaaatacga agtgggaaaa gctcaataat gaggtcagat gcaccattg
901 gcaactgcaa ttctgaatgc atcactocaa atggaagcat tcccaatgac
951 aaaccttttc aaaatgtaaa caggatcaca tatggggcct gtcccagata
1001 tgtaagcaa aacactctga aattggcaac agggatgcgg aatgtaccag
1051 agaaacaaac tagaggcata ttcggcgcaa tcgcaggttt catagaaaat
1101 ggttgggagg gaatggtaga cggttggtac ggtttcaggc atcaaaattc
1151 tgagggcaca ggacaagcag cagatcttaa aagcactcaa gcagcaatca
1201 accaaatcaa cgggaaactg aatagggttaa tcgagaaaac gaacgagaaa
1251 ttccatcaaa tcgaaaaaga attctcagaa gtagaaggga gaattcagga
1301 cctcgagaaa tatgttgaag aactaaaaat agatctctgg tcttacaacg
1351 cggagcttct tgttgccctg gagaaccaac atacaattga totaactgac
1401 tcagaaatga acaaactgtt tgaaagaaca aggaagcaac tgagggaaaa
1451 tgctgaggac atgggcaatg gttgcttcaa aatataccac aaatgtgaca
1501 atgcctgcat agggatcaatc agaaatggaa cttatgacca tgatgtatac
1551 agagacgaag cattaaacaa ccggttocag atcaaagggtg ttgagctgaa
1601 gtcaggatac aaagattgga tcctatggat ttcccttgcc atatcatgct
1651 ttttgctttg tgttgctctg ctgggggttca tcatgtgggc ctgccaaaaa
1701 ggcaacatta ggtgcaacat ttgcatttga gtgcattaat taaaaacacc
1751 cttgtttcta ct
```

SEQ ID NO:39

Amino Acid Sequence of ca A/Wuhan/395/95 H3

Entire molecule length: 566 aa

```
1 mktiiaalsyi lclvfaqlkp gndnstatlc lghhavpngt lvkttitndqi
51 evtnatelvq ssstgricds phrildgknc tlidallgdp hcdgfgqnkew
101 dlfferskay sncypydvdp yasrlslvas sgleftneg fnwtgvaqdg
151 tsyackrgsv ksffsrlnwl hkleykypal nvtmnpndkf dklyiwgvhh
201 pstdsdqtsi yvqasgrvtv stkrsgqvti pnigrpwwr gissrisiyw
251 tivkpgdill instgnliap rgyfkirsgk ssimrsdapi gncnsecitp
301 gnsipndkpf qnvnrityga cpryvkqntl klatgmrvnp ekqtrgifga
351 iagfiengwe gmvdgwygfr hqnsegtgga adlkstqaai nqingklrnl
401 iektnekfhq iekefseveg riqdlekyve dtkidlwsyn aellvalenq
451 htidltdsem nklfertrkq lrenaedmgn gcfskiyhkcd nacigsirng
501 tydhdvyrde alnnrfqikg velksygdw ilwisfaisc fllecvllgf
551 imwacqkgni rcnici
```

Figure 1F

SEQ ID NO:6

Nucleotide Sequence of ca A/Wuhan/395/95 N2

Entire molecule length: 1451 bp

```
1 agcaaaaagca ggagtgaata tgaatccaaa tcaaaaagata ataactattg
51 gctctgtttc tctcactatt gccacaatat gcttccttat gcaaattgcc
101 atcctggtaa ctactgtaac attacatttc aagcaatatg aatgcaactc
151 ccccccaaac aaccaagtaa tgctgtgtga accaacaata atagaaagaa
201 acataacaga gatagtgtat ctgaccaaca ccaccataga gaaggaaata
251 tgccccaac tagcagaata cagaaattgg tcaaagccgc aatgtaaaat
301 tacaggattt gcaccttttt ctaaggacaa ttcaattcgg ctttcgctg
351 gtggggacat ttgggtgaca agagaacctt atgtgtcatg cgatcctgac
401 aagtgttatc aatttgccct tggacagggg acaacactaa acaacaggca
451 ttcaaagac acagtacatg ataggacccc ttatcgaacc ctattgatga
501 atgagttggg tgttccattt catttgggaa ccaagcaagt gtgcatagca
551 tgggtccagct caagtgtgca cgatggaaaa gcatggctgc atgttttgtt
601 aactgggcat gatgaaaatg caactgctag cttcatttac gatggaggc
651 ttgtagatag tattggttca tgggtccaaa aaatcctcag gaccaggag
701 tcggaatgcg tttgtatcaa tggaaacttg acagtagtaa tgactgatgg
751 aagtgcctca ggaagagctg atactaaaat actattcatt gaagagggga
801 aaatcgctca tattagccca ttgtcaggaa gtgctcagca tgcgaggag
851 tgctcctgtt atcctcgata ttctggtgtc agatgtgtct gcagagacaa
901 ctggaaaggc tccaataggc ccacgtaga tataaatgtg aaagattata
951 gcattgtttc cagttatgtg tgctcaggac ttgttgaga cacaccaga
1001 aaaaacgaca gctccagcag tagccattgc ctgaatccta acaatgagga
1051 aggggggcat ggagtgaag gctgggcctt tgatgatgga aatgacgtgt
1101 ggatgggaag aacgatcagc gagaagttac gctcaggtta tgaaaccttc
1151 aaagtcattg gaggtgtgtc caaacctaac tcaaattgc agataaatag
1201 acaagtcata gttgacagag gtaataggtc cggttattct ggtattttct
1251 ctggtgaagg caaaagctgc atcaatcggg gcttttatgt ggagttgata
1301 aggggaagga aacaggaaac tgaagtctgg tggacctcaa acagtattgt
1351 tgtgttttgt ggcacctcag gtacatatgg aacaggctca tggcctgatg
1401 gggcgacat caatctcatg cctatataag ctttcgcaat ttagaaaaa
1451 a
```

SEQ ID NO:40

Amino Acid Sequence of ca A/Wuhan/395/95 N2

Entire molecule length: 469 aa

```
1 mnpnqkiiti gsvsltiati cflmqiaailv ttvtlhfkqy ecnsppnnqv
51 mlceptiier niteivyltn ttiekeicpk laeyrnwskp qckitgfapf
101 skdnsirlsa ggdiwvtrep yvscdpdkcy qfalgggttl nnrhsndtvh
151 drtpyrllm nelgvpfhlq tkqvciaawss sschdgkawl hvcvtghden
201 atasfiydgr lvdsigswsk kilrtqesec vcingtctvv mtdgsasgra
251 dtkilfieeg kivhisplsg saqhveecsc yprysgvrcv crdnwksnr
301 pivdinvkdy sivssyvcsg lvgdtprknd sssshclnp nneegghgvk
351 gwafddgndv wmgrtisekl rsgyetfkvi ggwskpnskl qinrqvivr
401 gnrsgysgif svegkscinr cfyvelirgr kqetevwwts nsivvfcgts
451 gtygtgswpd gadinlmpi
```

Figure 1G

SEQ ID NO:7

ca A/Sydney/05/97

Nucleotide Sequence of ca A/Sydney/05/97 H3 Entire molecule length: 1762 bp

```
1 agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg
51 agctacattt tatgtctggt ttctgctcaa aaaattcccc gaaatgacaa
101 cagcacggca acgctgtgcc tgggacacca tgcagtgcc aacggaacgc
151 tagtgaaaac aatcacgaat gaccaaattg aagtgactaa tgctactgag
201 ctggttcaga gttcctcaac aggtagaata tgcgacagtc ctcaccgaat
251 ccttgatgga gaaaactgca cactgataga tgctctattg ggagaccctc
301 attgtgatgg cttccaaaat aaggaatggg acctttttgt tgaacgcagc
351 aaagcctaca gcaactgtta cccttatgat gtgccggatt atgcctccct
401 taggtcacta gttgcctcat ccggcacccct ggagtttaac aatgaaagct
451 tcaattggac tggagtcgct cagaatggaa caagctatgc ttgcaaaagg
501 agttctatta aaagtttctt tagtagattg aattggttgc accaattaaa
551 atacaaatat ccagcactga acgtgactat gccaaacaat gacaaatttg
601 acaaattgta catttggggg gttcaccacc cgagtacgga cagtgaacaa
651 accagcatat atgctcaagc atcagggaga gtcacagtct ccacaaaag
701 aagccaacaa actgtaatcc cgaatatcgg atctagacct tgggtaaggg
751 gtatctccag cagaataagc atccattgga caatagtaaa accgggagac
801 atacttttga ttaacagcac agggaatcta attgctcctc ggggttactt
851 caaaatacga agtgggaaaa gctcaataat gaggtcagat gcacccattg
901 gcaaatgcaa ttctgaatgc atcactccaa atggaagcat tcccaatgac
951 aaaccatttc aaaatgtaaa caggatcaca tatggggcct gtcccagata
1001 tgtaaagcaa aacactctga aattggcaac agggatgcg aatgtaccag
1051 agaacaacac tagaggcata ttcggcgcaa tcgcaggtt catagaaaat
1101 ggttgggagg gaatggtaga cggttggtac ggtttcaggc atcaaaattc
1151 tgagggcaca ggacaagcag cagatcttaa aagcactcaa gcagcaatca
1201 accaaatcaa cgggaaactg aataggttaa tcgagaaaac gaacgagaaa
1251 ttccatcaaa ttgaaaaaga attctcagaa gtagaaggga gaattcagga
1301 cctcgagaaa tatgttgagg aactaaaaat agatctcttg tcgtacaacg
1351 cggagcttct tgttgccctg gagaaccaac atacaattga tctaactgac
1401 tcagaaatga acaaactgtt tgaaagaaca aggaagcaac tgagggaaaa
1451 tgctgaggat atgggcaatg gttgtttcaa aatataccac aaatgtgaca
1501 atgcctgcat aggggtcaatc agaaatggaa cttatgacca tgatgtatac
1551 agagacgaag cattaacaaa ccggttccag atcaaagggt ttgagctgaa
1601 gtcaggatac aaagattgga tcctatggat ttcttttgcc atatcatggt
1651 ttttgctttg tgttgttttg ctggggttca tcatgtgggc ctgccaaaaa
1701 ggcaacatta ggtgcaacat ttgcatttga gtgcattaat taaaaacacc
1751 cttgttttcta ct
```

SEQ ID NO:41

Amino Acid Sequence of ca A/Sydney/05/97 H3 Entire molecule length: 566 aa

```
1 mkttiialsyi lclvfaqkip gndnstatlc lghhavpngt lvkttitndqi
51 evtnatelvq sstgricds phrildgenc tlidallgdp hcdgfnkew
101 dlflverskay sncypydvdp yaslrlvas sgtlefnnes fnwtgvaqng
151 tsyackrssi ksffsrlnwl hqlkykypal nvtmpnndkf dklyiwgvhh
201 pstdsdqtsi yaqasgrvtv stkrsqqtvi pnigrpwwr gissrisihw
251 tivkpgdill instgnliap rgyfkirsgk ssimrsdapi gkcncsecitp
301 ngsipndkpf qnvnrityga cpryvkqntl klatgmrnvp ekqtrgfiga
351 iagfiengwe gmvdgwygfr hqnsegtgga adlkstqaai nqingklrnl
401 iektnekfhq iekefseveg riqdlekyve dtkidlwsyn aellvalenq
451 htidltdsem nklfertrkq lrenaedmgm gcfkiahkcd nacigsirng
501 tydhdvyrde alnnrfqikg velksgykdw ilwisfaisc fllecvllgf
551 imwacqkgni rcnici
```

Figure 1H

SEQ ID NO:8

Nucleotide Sequence of ca A/Sydney/05/97 N2

Entire molecule length: 1467 bp

```
1 agcaaaagca ggagtaaaga tgaatccaaa tcaaaagata ataacgattg
51 gctctgtttc tctcactatt gccacaatat gcttccttat gcaaattgcc
101 atcctggtaa ctactgtaac attgcatttc aagcaatatg aatgcagctc
151 tcccccaaac aaccaagtaa tgctgtgtga accaacaata atagaaagaa
201 acataacaga gatagtgtat ctgaccaaca ccaccataga gaaggaaata
251 tgccccaaac tagcagaata cagaaattgg tcaaagccac aatgtaaaat
301 tacaggattt gcaccttttt ctaaggacaa ttcaattcgg ctttccgctg
351 gtggggacat ttgggtgaca agggaaacct atgtgtcgtg cgatcctgac
401 aagtgttatc aatttgccct tggacaggga acaacactaa acaacaggca
451 ttcaaatgac acagtacatg ataggacccc ttatcgaacc ctattgatga
501 atgagttggg tgttccattt catttgggaa ccaagcaagt gtgcatagca
551 tgggtccagct caagttgtca cgatggaaaa gcatggctgc atgtttgtgt
601 aactgggcat gatgaaaatg caactgctag cttcatttac gatgggaggc
651 ttgtagatag tattggttca tggtcacaaa aaatcctcag gaccaggag
701 tcggaatgcg tttgtatcaa tggaaacttg acagtagtaa tgactgatgg
751 gagtgcctca ggaagagctg atactaaaat actattcatt gaggagggga
801 aaatcgttca tatcagccca ctgtcaggaa gtgctcagca tgtcaggag
851 tgctcctggt atcctcgata tcctggtgtc agatgtgtct gcagagacaa
901 ctggaaaggc tccaataggc ccacgtaga tataaatgta aaggattata
951 gcattgtttc cagttatgtg tgctcaggac ttgttgagga cacaccaga
1001 aaaaacgaca gctccagcag tagtcattgc ctgaatccta acaatgagga
1051 aggggggtcat ggagtgaag gctgggcctt tgatgatgga aatgacgtgt
1101 ggatgggaag aacgatcagc gagaagttcc gctcaggtta tgaaaccttc
1151 aaagtcattg aaggctggtc caaacctaac tccaaattgc agataaatag
1201 gcaagtcata gttgacagag gtaataggtc cggttattct ggtattttct
1251 ctggtgaagg caaaagctgc atcaatcggg gcttttatgt ggagttgata
1301 aggggaagga aacaggaaac tgaagtctgg tggacctcaa acagtattgt
1351 tgtgttttgt ggcacctcag gtacatatgg aacaggctca tggcctgatg
1401 gggcggacat caatctcatg cctatataag ctttcgcaat ttagaaaaa
1451 aactccttgt ttctact
```

SEQ ID NO:42

Amino Acid Sequence of ca A/Sydney/05/97 N2

Entire molecule length: 469 aa

```
1 mnpnqkiiti gsvsltiati cflmqiaailv ttvtlhfkqy ecsspnnqv
51 mlceptiier niteivyltn ttiekeicpk laeyrnwskp qckitgfapf
101 skdnsirlsa ggdiwvtrep yvscdpdkcy qfalgggttl nnrhsndtvh
151 drtpyrllm nelgvpfhlg tkqvciaawss sschdgkawl hvcvtghden
201 atasfiydgr lvdsigswsk kilrtqesec vcingtctvv mtdgsasgra
251 dtkilfieeg kivhisplsg saqhveecsc yprypgvrcv crdnwksnr
301 pivdinvkdy sivssyvcsq lvgdtpkrnd ssssshclnp nneegghgvk
351 gwafddgndv wmgrtisekf rsgyetfkvi egwskpnskl qinrqvivdr
401 gnrsgysgif svegkscinr cfyvelirgr kqetevwwts nsivvcgts
451 gtygtgswpd gadinlmpi
```

Figure 11

SEQ ID NO:9

ca A/Panama/2007/99

Nucleotide Sequence of ca A/Panama/2007/99 H3 Entire molecule length: 1762 bp

```
1 agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg
51 agctacattt tatgtctgtt tttcgctcaa aaacttcccc gaaatgacaa
101 cagcacggca acgctgtgcc tggggcacca tgcagtgtca aacggaacgc
151 tagtgaaaac aatcacgaat gaccaaattg aagtgactaa tgctactgag
201 ctggttcaga gttcctcaac aggtagaata tgcgacagtc ctcaccaa
251 ccttgatgga gaaaactgca cactaataga tgctctattg ggagaccctc
301 attgtgatgg cttccaaaat aaggaatggg acctttttgt tgaacgcagc
351 aaagcctaca gcaactgtta cccttatgat gtgcgggatt atgcctccct
401 taggtcacta gttgcctcat ccggcacact ggagtttaac aatgaaagct
451 tcaattggac tggagtctgt cagaatggaa caagctctgc ttgcaaaagg
501 ggatctaata aaagtttctt tagtagattg aattgggtgc accaattaaa
551 atacaaatat ccagcactga acgtgactat gccaaacaat gaaaaattt
601 acaaatgtta ctttggggg gttctccacc cgagtacgga cagtgaacca
651 atcagcctat atgctcaagc atcagggaga gtcacagtct ctacccaaag
701 aagccaacaa actgtaatcc cgaatatcgg atctagacct tgggtaagg
751 gtgtctccag cagaataagc atctattgga caatagtaaa accgggagac
801 atacttttga ttaacagcac agggaatcta attgctcctc ggggttactt
851 caaaatacga agtgggaaaa gctcaataat gaggtcagat gcaccattg
901 gcaaatgcaa ttctgaatgc atcactccaa atggaagcat tcccaatgac
951 aaaccatttc aaaatgtaaa caggatcaca tatggggcct gtcccagata
1001 tgtaagcaa aacactctga aattggcaac agggatgcgg aatgtaccag
1051 agaaacaaac tagaggcata ttcggcgcaa tcgcggttt catagaaaat
1101 ggttgggagg gaatggtgga cggttggtac ggtttcaggc atcaaaattc
1151 tgagggcaca ggacaagcag cagatcttaa aagcactcaa gcagcaatca
1201 accaaatcaa cgggaaactg aataggttaa tcgagaaaac gaacgagaaa
1251 ttccatcaaa ttgaaaaaga attctcagaa gtagaaggga gaattcagga
1301 cctcgagaaa tatgttgagg acactaaaat agatctctgg tcgtacaacg
1351 cggagcttct tgttgccctg gagaaccaac atacaattga tctaactgac
1401 tcagaaatga acaaaactgt tgaaagaaca aagaagcaac tgagggaaaa
1451 tgctgaggat atgggcaatg gttgtttcaa aatataccac aaatgtgaca
1501 atgcctgcat agggatcaatc agaaatggaa cttatgacca tgatgtatac
1551 agagacgaag cattaacaa cgggttcag atcaaagggt ttgagctgaa
1601 gtcaggatac aaagattgga tcctatggat ttcctttgcc atatcatgct
1651 ttttgctttg tggtgttttg ctgggggttc tcatgtgggc ctgccccaaa
1701 ggcaacatta ggtgcaacat ttgcatttga gtgcattaat taaaaacacc
1751 cttgtttcta ct
```

SEQ ID NO:43

Amino Acid Sequence of ca A/Panama/2007/99 H3 Entire molecule length: 566 aa

```
1 mktiialsyi lclvfaqklp gndnstatlc lghhavsngt lvkttitndqi
51 evtnatelvq ssstgricds phqildgenc tlidallgdp hcdgfgqnkew
101 dlfverskay sncypydvdp yaslrslvas sgtlefnnes fnwtgvaqng
151 tssackrgsn ksffsrlnwl hqlkykypal nvtmnpnekf dklyiwgvlh
201 pstdsdqisl yaqasgrvtv stkrsgqvti pnigsrpwwr gvssrisiyw
251 tivkpgdill instgnliap rgyfkirsgk ssimrsdapi gkcnsecitp
301 ngsipndkpf qnvnrityga cpryvkqntl klatgmrvnp ekqtrgifga
351 iagfiengwe gmvdgwygfr hqnsegtgga adlkstqaai nqingklrnl
401 iektnekfhq iekefseveg riqdlekyve dtkidlwsyn aellvalenq
451 htidltldsem nklfertkkq lrenaedmgn gcfkiyhkcd nacigsirng
501 tydhdvyrde alnnrfqikg velksygdw ilwisfaisc fllecvllgf
551 imwacqkgni ronicl
```

Figure 1J

SEQ ID NO:10

Nucleotide Sequence of ca A/Panama/2007/99 N2 Entire molecule length: 1466 bp

```
1 agcaaaagca ggagtaaaga tgaatccaaa tcaaaagata ataacgattg
51 gctctgtttc tctcactatt gccacaatat gcttccttat gcaaatagcc
101 atcctggtaa ctactgtaac attgcatttc aagcaatatg aatgcaactc
151 ccccccaaac aaccaagtaa tgctgtgtga accaacaata atagaaagaa
201 acataacaga gatagtgtat ctgaccaaca ccaccataga gaaggaaata
251 tgccccaac tagcagaata cagaaattgg tcaaagccgc aatgtaaaat
301 tacaggattt gcaccttttt ctaaggataa ttcaattcgg ctttcgctg
351 gtggggacat ttgggtgaca agagaacctt atgtgtcatg cgatcctgac
401 aagtgttata aatttgccct tggacaggga acaacactaa acaacaggca
451 ttcaaataac acagtacatg ataggacccc ttatcgaacc ctattgatga
501 atgagttggg tgttccattt catttgggaa ccaagcaagt gtgtatagca
551 tgggtccagc caagttgtca cgatggaaaa gcatggctgc atgtttgtgt
601 aactgggcat gatgaaaatg caactgctag cttcatttac gatgggagac
651 ttgtagatag tattggttca tgggtccaaa aaatcctcag gaccaggag
701 tcggaatgcg tttgtatcaa tggaacttgt acagtagtaa tgactgatgg
751 gagtgcctca ggaagagctg atactaaaat acttttcatt gaggagggga
801 aaatcgctca tactagcaaa ttgtcaggaa gtgctcagca tgtcaggag
851 tgctcctgtt atcctcgata tcctggtgtc agatgtgtct gcagagacaa
901 ctggaaaggc tccaataggc ccacgtaga tataaatgta aaggattata
951 gcattgtttc cagttatgtg tgctcaggac ttgttgga caacaccaga
1001 aaaaacgaca gctccagcag tagccattgc ctggatccta acaatgaaga
1051 aggggggtcat ggagtgaag gctgggcctt tgatgatgga aatgacgtgt
1101 ggatgggaag aacgatcagc gagaagtcac gctcaggtta tgaaaccttc
1151 aaggtcattg aaggctggtc caaacctaac tccaaattgc agataaata
1201 gcaagtcata gttgaaagag gtaatatgtc cggttattct ggtattttct
1251 ctgttgaagg caaaagctgc atcaatcggg gcttttatgt ggagttgata
1301 aggggaagga aacaggaaac tgaagtctgg tggacctcaa acagtattgt
1351 tgtgttttgt ggcacctcag gtacatatgg aacaggctca tggcctgatg
1401 gggcggacat caatctcatg cctatataag ctttcgcaat tttagaaaaa
1451 actccttgtt tctact
```

SEQ ID NO:44

Amino Acid Sequence of ca A/Panama/2007/99 N2 Entire molecule length: 469 aa

```
1 mnpnqkiiti gsvsltiati cflmqiaailv ttvtlhfkqy ecnsppnnqv
51 mlceptiier niteivyltn ttiekeicpk laeyrnwskp qckitgfapf
101 skdnsirlsa ggdiwvtrep yvscdpdkcy qfalgqgttl nnrhsndtvh
151 drtpyrllm nelgvpfhlg tkqvciawss sschdgkawl hvcvtghden
201 atasfiydgr lvdsigswsk kilrtqesec vcingtctvv mtdgsasgra
251 dtkilfieeg kivhtsklsg saqhveecsc yprypgvrcv crdnwksnr
301 pivdinvkdy sivssyvcsg lvgdtpknd sssshcldp nneegghgvk
351 gwafddgndv wmgrtiseks rsgyetfkvi egwskpnskl qinrqviver
401 gnmsgysgif svegkscinr cfyvelirgr kqetevwwts nsivvfcgts
451 gtygtgswpd gadinlmpi
```

Figure 1K

SEQ ID NO:11

ca A/Wyoming/03/2003

Nucleotide Sequence of ca A/Wyoming/03/2003 H3 Entire molecule length: 1762 bp

```
1 agcaaaagca ggggataatt ctattaacca tgaagactat cattgcttta
51 agctacattc tatgtctggt tttctctcaa aagcttcccg gaaatgacaa
101 cagcacggca acgctgtgcc ttgggcacca tgcagtacca aacggaacga
151 tagtgaaaac aatcacgaat gaccaaattg aagttactaa tgctactgag
201 ctggttcaga gttcctcaac aggtggaata tgcgacagtc ctcatacagat
251 ccttgatgga gaaaactgca cactaataga tgctctattg ggagaccctc
301 agtgtgatgg cttccaaaat aagaaatggg acctttttgt tgaacgcagc
351 aaagcctaca gcaactgtta cccttatgat gtgccggatt atgcctccct
401 taggtcacta gttgcctcat ccggcacact ggagtttaac aatgaaagct
451 tcaattgggc tggagtcact cagaatggaa caagctctgc ttgcaaaagg
501 agatctaata aaagtttctt tagtagattg aattggttga cccacttaaa
551 atacaaatac ccagcattga acgtgactat gccaaacaat gaaaaatttg
601 acaaattgta catttggggg gttcaccacc cggttacgga cagtgaccaa
651 atcagcctat atgctcaagc atcaggaaga atcacagtct ctaccaaag
701 aagccaacaa actgtaatcc cgaatatcgg atatagacc agggtaaggg
751 atatctccag cagaataagc atctattgga caatagtaaa accgggagac
801 atacttttga ttaacagcac aggaaatcta attgctcctc ggggttactt
851 caaaatcgga agtgggaaaa gctcaataat gagatcagat gcaccattg
901 gcaaatgcaa ttctgaatgc atcactccaa atggaagcat tcccaatgac
951 aaaccatttc aaaatgtaaa caggatcaca tatggggcct gtcccagata
1001 tgттаagcaa aacactctga aattggcaac agggatgcga aatgtaccag
1051 agaaacaaac tagaggcata tttggcgcaa tcgcggttt catagaaaat
1101 ggttgaggag gaatggtgga cggttgttac ggtttcaggc atcaaaattc
1151 tgagggcaca ggacaagcag cagatctcaa aagcactcaa gcagcaatca
1201 accaaatcaa tgggaaactg aataggttaa tcgggaaaac aaacgagaaa
1251 ttccatcaga ttgaaaaaga attctcagaa gtagaaggga gaattcagga
1301 cctcgagaaa tatgttgagg aactaaaaat agatctctgg tcatacaacg
1351 cggagcttct tgttgccctg gaaaaccaac atacaattga tctaactgac
1401 tcagaaatga acaaaactgtt tgaaagaaca aagaagcaac tgagggaaaa
1451 tgctgaggat atgggcaatg gttgtttcaa aatataccac aaatgtgaca
1501 atgcctgcat agagtcaatc agaaatggaa cttatgacca tgatgtatac
1551 agagatgaag cattaacaa cgggttccag atcaaagggt ttgagctgaa
1601 gtcaggatac aaagattgga tcctatggat tccttttgcc atatcatgtt
1651 ttttgctttg tgttgctttg ttggggttca tcatgtgggc ctgccaaaaa
1701 ggcaacatta ggtgcaacat ttgcatttga gtgcattaat taaaaacacc
1751 cttgtttcta ct
```

SEQ ID NO:45

Amino Acid Sequence of ca A/Wyoming/03/2003 H3 Entire molecule length: 566 aa

```
1 mktiialsyi lclvfsqklp gndnstatlc lghhavpngt ivkttitndqi
51 evtnatelvq ssstggicds phqildgenc tlidallgdp qcdgfnkkw
101 dlflverskay sncypydvpd yaslrslvas sgtlefnnes fnwagvtqng
151 tssackrrsn ksffsrlnwl thlkykypal nvtmpnnekf dklyiwgvhh
201 pvtddsqisl yaqasgritv stkrsgqtvi pnigyrprvr dissrisiyw
251 tivkpgdill instgnliap rgyfkirsgk ssimrsdapi gkcnsecitp
301 ngsipndkpf qnvnrityga cpryvkqntl klatgmrnvp ekqtrgifga
351 iagfiengwe gmvdgwygfr hqnsegtgga adlkstqaai nqingklrnl
401 igktnekfhq iekefseveg riqdlekyve dtkidlwsyn aellvalenq
451 htidltldsem nklferrtkkq lrenaedmgn gcfkiyhkcd naciesirng
501 tydhdvyrde alnnrfqikg velksykdw ilwisfaisc fllevallgf
551 imwacqkgni rcnici
```

Figure 1L**SEQ ID NO:12**

Nucleotide Sequence of ca A/Wyoming/03/2003 N2

Entire molecule length: 1467 bp

```
1  agcaaaagca  ggagtaaaga  tgaatccaaa  tcaaaagata  ataacgattg
51  gctctgtttc  cctcaccatt  tccacaatat  gcttcttcat  gcaaattgcc
101 atcctgataa  ctactgtaac  attgcatttc  aagcaatatg  aattcaactc
151 ccccccaaac  aaccaagtga  tgctgtgtga  accaacaata  atagaaagaa
201 acataacaga  gatagtgtat  ctgaccaaca  ccaccataga  gaaggaaata
251 tgccccaac  tagcagaata  cagaaattgg  tcaaagccgc  aatgtaacat
301 tacaggattt  gcaccttttt  ctaaggacaa  ttcgattcgg  ctttcogctg
351 gtggggacat  ctgggtgaca  agagaacctt  atgtgtcatg  cgatcctgac
401 aagtgttatc  aatttgccct  tggacaggga  acaacactaa  acaacgtgca
451 ttcaaataac  acagtacatg  ataggacccc  ttatcggacc  ctattgatga
501 atgagttggg  tgttccattt  catctgggga  ccaagcaagt  gtgcatagca
551 tgggtccagc  caagttgtca  cgatggaaaa  gcatggctgc  atgttttgtg
601 aacgggggat  gatgaaaatg  caactgctag  cttcatttac  aatgggaggc
651 ttgtagatag  tattgtttca  tgggtccaaa  aaatcctcag  gacccaggag
701 tcagaatgca  tttgtatcaa  tggaaactgt  acagtagtaa  tgactgatgg
751 gagtgcttca  ggaaaagctg  atactaaaat  actattcatt  gagggaggga
801 aaattgttca  tactagcaca  ttatcaggaa  gtgctcagca  tgtcgaggag
851 tgctcctggt  atcctcgata  tcctgggtgt  agatgtgtct  gcagagacaa
901 ctggaaaggc  tccaataggc  ccatcgtaga  tataaacata  aaggattata
951 gcattgtttc  cagttatgtg  tgctcaggac  ttgttggaga  cacaccacaga
1001 aaaaacgaca  gctccagcag  tagccattgc  ttggatccaa  acaatgagga
1051 aggtggatca  ggagtgaag  gctgggcatt  tgatgatgga  aatgacgtgt
1101 ggatgggaag  aacgatcagc  gagaagttac  gctcaggata  tgaaaccttc
1151 aaagtcattg  aaggctggtc  caaccctaac  tccaaattgc  agataaatag
1201 gcaagtcata  gttgacagag  gtaacaggtc  cggttattct  ggtattttct
1251 ctgttgaaag  caaaagctgc  atcaatcggg  gcttttatgt  ggagttgata
1301 aggggaagaa  aacaggaaac  tgaagtcttg  tggacctcaa  acagtattgt
1351 tgtgttttgt  ggcacctcag  gtacatatgg  aacaggctca  tggcctgatg
1401 gggcggacat  caatctcatg  cctatataag  ctttcgcaat  tttagaaaaa
1451 aactccttgt  ttctact
```

SEQ ID NO:46

Amino Acid Sequence of ca A/Wyoming/03/2003 N2 Entire molecule length: 469 aa

```
1  mnpnqkiiti  gsvsltisti  cffmqiaili  ttvtlhfkqy  efnsppnnqv
51  mlceptiier  niteivyltn  ttiekeicpk  laeyrnwskp  qcnitgfapf
101 skdnsirlsa  ggdiwvtrp  yvscdpdkcy  qfalgqgttl  nnvhsndtvh
151 drtpyrllm  nelgvpfhlg  tkqvciawss  sschdgkawl  hvcvtgdden
201 atasfiyng  lvdsivswsk  kilrtqesec  vcingtctvv  mtdgsasgka
251 dtkilfieeg  kivhtstlsg  saqhveecsc  yprypgvrcv  crdnwkgsnr
301 pivdinikdy  sivssyvcs  lvgdtprknd  ssssshcldp  nneeegghgvk
351 gwafddgndv  wmgrtisekl  rsgyetfkvi  egwsnpnsl  qinrqvivr
401 gnrsgysgif  svegkscinr  cfyvelirgr  kqetevlwts  nsivvfcgts
451 gtygtgswpd  gadinlmpi
```

Figure 1M

SEQ ID NO:13

ca A/Texas/36/91

Nucleotide Sequence of ca A/Texas/36/91 H1

Entire molecule length: 1778 bp

```
1 agcaaaagca ggggaaaata aaaacaacca aaatgaaagc aaaactacta
51 gtcctgttat gtgcattttac agctacatat gcagacacaa tatgtatagg
101 ctaccatgcg aacaactcaa ccgacactgt tgacacagta cttgagaaga
151 acgtgacagt gacacactct gtcaacctac ttgaggacag tcacaacgga
201 aaactatgtc gactaaaggg aatagcccca ctacaattgg gtaattgcag
251 cgttgccgga tggatcttag gaaacccaaa atgcgaatca ctgttttcta
301 aggaatcatg gtcctacatt gcagaaacac caaacctga gaatggaaca
351 tgttacccag ggtattttcg cgactatgag gaactgaggg agcaattgag
401 ttcagtatca tcattcgaga gattcgaaat attcccaaaa gaaagctcat
451 ggccaacca caccgtaacc aaaggagtaa cgacatcatg ctcccataat
501 gggaaaagca gtttttacag aaatttgcta tggctgacga agaagaatgg
551 cttgtacca aatgtgagca agtcctatgt aaacaacaaa gagaaagaag
601 tccttgtaact atgggggtgt catcacccgt ctaacatagg ggaccaaaagg
651 gccatctatc atacagaaaa tgcttatgtc tctgtagtgt cttcacatta
701 tagcagaaga ttcaccccag aaatagcaaa aagacccaaa gtaagagatc
751 aagaaggaag aattaactac tactggactc tgctggaacc cggggacaca
801 ataataattg aggcaaattg aaatctaata gcgccatggg atgctttcgc
851 actgagtaga ggctttgggt caggaatcat cacctcaaac gcatcaatgg
901 atgaatgtga cgcgaagtgt caaacacccc agggagctat aaacagtagt
951 cttcctttcc agaatgtaca cccagtcaca ataggagagt gtccaaagta
1001 tgtcaggagt acaaaattaa ggatggttac aggactaagg aacatoccat
1051 ccattcaatc cagaggtttg tttggagcca ttgccggttt cattgaaggg
1101 ggggtggactg gaatgataga tggatgggat gggttatcat atcagaatga
1151 acaaggatct ggctatgctg cggaccaaaa aagcacacaa aatgccatta
1201 acgggattac aaacaagggt aattctgtaa tcgagaaaat gaacactcaa
1251 ttcacagctg tgggcaaaaga attcaacaaa ttagaaagaa ggatggaaaa
1301 cttaaataaa aaagttgatg atggatttct ggacatttgg acatataatg
1351 cagaattgtt ggttctactg gaaaatggaa ggactttgga ttttcatgac
1401 tcaaatgtga agaatctgta tgagaaagta aaaagccaat tgaagaataa
1451 tgccaaagaa atagggaacg ggtgttttga attctatcac aagtgttaaca
1501 atgaatgcat ggaaagtgtg aaaaatggaa cttatgacta tccaaaatat
1551 tccgaagaat caaagttaaa caggggaaaa attgatggag tgaaatttga
1601 atcaatggga gtctatcaga ttctggcgat ctactcaact gtcgccagtt
1651 cactggtgct tttggtctcc ctgggggcaa tcagcttctg gatgtgttct
1701 aatgggtctt tgcagtgtag aatatgcato tgagaccaga atttcagaaa
1751 tataagaaaa aacacccttg tttctact
```

SEQ ID NO:47

Amino Acid Sequence of ca A/Texas/36/91 H1

Entire molecule length: 566 aa

```
1 mkakllvllc aftatyadti cigyhannst dtvdtvlekn vtvthsvnl1
51 edshngklcr lkgiaplqlg ncsvagwilg npkceslfsk eswsyiaetp
101 npengtcypg yfadyeelre qlssvssfer feifpkessw pnhtvtkgvt
151 tscshngkss fyrnllwltk knglypnvsk syvnnkekev lvlwgvhpps
201 nigdqraiyl tenayvsvvs shysrrftpe iakrpkvrdq egrinyywtl
251 lepgdtiife angnliapwy afalsrgfgs giitsnasmd ecdakcqtpp
301 gainsslpfq nvhpvtigec pkyvrstklr mvtglrnips iqsrgrlfgai
351 agfieggwtg midgwygyhh qneqgsyyaa dqkstqnain gitnkvnsvi
401 ekmntqftav gkefnklerr menlnkkvdd gfldiwtyna ellvllengr
451 tldfhdsnvk nlyekvksql knnakeigng cfefyhkcnn ecmesvkngt
501 ydypkysees klnrgkidgv klesmgvyqi laiystvass lvllvslgai
551 sfwmcnsgsl qcrici
```

Figure 1N**SEQ ID NO:14**

Nucleotide Sequence of ca A/Texas/36/91 N1

Entire molecule length: 1463 bp

```
1 agcaaaagca ggagtttaaa atgaatccaa atcaaaaaat aataatcata
51 ggatcaatca gtatggcaat cggaataatt agtctaatat tgcaaatagg
101 aaatattatt tcaatatggg ctagccactc aatccaaact ggaagtcaaa
151 accacactgg aatatgcaac caaagaatca ttacatatga aaatagcacc
201 tgggtgaatc aaacatatgt taatattaac aacactaatg ttgttgctgg
251 aaaggacaaa acttcagtga cattggccgg caattcatct ctttgcccta
301 tccgtgggtg ggctatatac acaaaagaca acagcataag aattggttcc
351 aaaggagatg tttttgtcat aagagagcct tttatatcat gttctcactt
401 ggaatgcaga accttttttc tgaccaaggg tgctctatta aatgacaagc
451 attcaaattg gaccgttaag gacagaagcc cttatagggc cttaatgagc
501 tgtcctctag gtgaagctcc gtctccatac aattcaagat ttgaatcagt
551 tgcttggtca gcaagcgcac gccatgatgg catgggctgg ctaacaatcg
601 gaatttctgg tccagataat ggagcagtgg ctgtactaaa atacaacggc
651 ataataactg aaaccataaa aagttggaag aagcgaatat taagaacaca
701 agagtctgaa tgtgtctgtg tgaacggttc atgttttacc ataatgaccg
751 atggcccgag taatggggcc gcctcgtaca gaatcttcaa aatcgagaag
801 ggggaaggta ctaaataaat agagttggat gcacccaatt atcattacga
851 ggaatgttcc tgttaccag acaccggcac agtgatgtgt gtgtgcaggg
901 acaattggca cggttcaa atcgacctggg tgtcttttaa tcaaacctg
951 gattatcaaa taggatacat ctgcagtggg gtgttcgggtg acaatccgcg
1001 tcccaaagat ggagaaggca gctgtaatcc agtgactggt gatggagcag
1051 acggagtaaa ggggttttca tacagatatg gtaatggtgt ttggatagga
1101 aggactaaaa gtaacagact cagaaaggga tttgagatga tttgggatcc
1151 taatggatgg acagataccg acagtgat ttctgtgaaa caggatgtcg
1201 tggcaatgac tgattgggtc gggtagacgg gaagtttcgt tcaacatcct
1251 gagctaacag gattggactg tatgagacct tgcttctggg ttgaattaat
1301 cagagggcga cctagagaaa atacaacaat ctggactagt gggagcagca
1351 tttctttttt tggcgtaaat agcgatactg caaactgggtc ttggccagac
1401 ggtgccgagt tgccattcac cattgacaag tagtccgttg aaaaaaact
1451 ccttgtttct act
```

SEQ ID NO:48

Amino Acid Sequence of ca A/Texas/36/91 N1

Entire molecule length: 470 aa

```
1 mnpnqkiiii gsismaigii slilqignii siwashsiqt gsqnhtgicn
51 qriityenst wvnqtyvnin ntnvvagkdk tsvtlagnss lcpirgwaiy
101 tkdnsirigs kgdvfvirep fiscshlecr tffltqgall ndkhsngtvk
151 drspyrals cplgeapsy nsrfesvaws asachdgmw ltigisgpdn
201 gavavlkyng iitetikswk krilrtqese cvcvngscft imtdgpsnga
251 asyrifkiek gkvtksiel apnyhyeecs cypdtgtvmc vcrdnwhgsn
301 rpwvsfnqnl dyqigyicsg vfgdnprpkd gegscnpvtv dgadgvkgfs
351 yryngvwig rtksnrlrk femiwdpngw ttdsdfsvk qdvvamtdws
401 gysgsfvqhp eltgldcmrp cfwvelirgr prenttiwts gssisfcgvn
451 sdtanwswpd gaelpftidk
```

Figure 1O

SEQ ID NO:15

ca A/Shenzhen/227/95

Nucleotide Sequence of ca A/Shenzhen/227/95 H1 Entire molecule length: 1689 bp

```
1  aaatgaaagc  aaaactacta  gtcctgttgt  gtgcattttac  agctacatat
51  gcagacacaa  tatgtatagg  ctaccatgcg  aacaactcaa  ccgacactgt
101  tgacacagta  cttgagaaga  acgtgacagt  gacacactct  gtcaacctac
151  ttgaggacag  tcacaacgga  aaactatgcc  gactaaaagg  aacagcccca
201  ctacaattgg  gtaattgcag  cgttgccgga  tggatcttag  gaaaccaga
251  atgcgaatca  ctgttttcta  aggaatcatg  gtcctacatt  gcagaaacac
301  caaacctga  gaatggaaca  tgttaccag  ggtatttcgc  cgactatgag
351  gaactgaggg  agcaattgag  ctcagtatca  tcattcgaga  gattcgaaat
401  attccccaag  gaaagctcat  ggcccaaaca  caccgtaacc  aaaggagtga
451  cggcatcatg  ctcccataat  gggaaaagca  gttttttaca  aaatttgcta
501  tggttgacgg  aaaagaatgg  cttgtaccca  aatctgagca  agtcctatgt
551  aaacaacaag  gagaaagaag  tccttgtaact  atgggggtgt  catcacccgt
601  ctaacatagg  ggaccaaagg  gccatctatc  atacagaaaa  tgcttatgtc
651  tctgtagtgt  cttcacatta  tagcagaaga  ttcacccag  aaatagcaaa
701  aagacccaaa  gtaagaggtc  aagaaggag  aattaactac  tactggactc
751  tgctggaacc  cggggacaca  ataataattg  aggcaaatgg  aaatctaata
801  gcgccatggt  acgctttcgc  actgagtaga  ggctttgggt  caggaatcat
851  cacctcaacc  gcatcaatgg  gtgaatgtga  cgctaagtgt  caaacacccc
901  aaggagctat  aaacagtagt  cttcctttcc  agaatgtaca  ccagtcaca
951  ataggagagt  gtcccaagta  tgtcaggagt  acaaaattaa  ggatggttac
1001  aggactaaga  aacatcccat  ccattcaatc  tagaggtttg  tttggagcca
1051  ttgccggttt  cattgaagg  ggtggactg  gaatgataga  tggatggtat
1101  ggttatcatc  atcagaatga  acaaggatct  ggctatgctg  cagacaaaaa
1151  aagcacacaa  aatgccattg  atgggattac  aaacaagggt  aattctgtaa
1201  tcgagaaaat  gaacactcaa  ttcacagctg  taggcaaaga  attcaacaaa
1251  ttagagagaa  ggatggaaaa  cttaaataag  aaagttgatg  atggatttct
1301  ggacattttg  acatataatg  cagagttggt  ggttctcctg  gaaaatggaa
1351  ggactttggg  ttttcattg  tcaaattgta  agaattctgt  tgagaaagta
1401  aaaaaccaat  tgaagaataa  tgccaaagaa  atcggaacg  ggtgttttga
1451  attctatcac  aagtgtaa  atgaatgcat  ggaaagtgtg  aaaaatggaa
1501  cttatgacta  tccaaaatat  tccgaagaat  caaagttaaa  cagggaaaaa
1551  attgatggag  tgaaattgga  atcaatggga  gtctatcaga  ttctggcgat
1601  ctactcaact  gtcgccagtt  cactggtgct  tttggtctcc  ctgggggcaa
1651  tcagtttctg  gatgtgttct  aatgggtcct  tgcagtgtg
```

SEQ ID NO:49

Amino Acid Sequence of ca A/Shenzhen/227/95 H1 Entire molecule length: 562 aa

```
1  mkakllvllc  aftatyadti  cigyhannst  dtvdtvlekn  vtvthsvnl1
51  edshngklcr  lkgtaplqlg  ncsvagwilg  npeceslfsk  eswsyiaetp
101  npengtcypg  yfadyeelre  qlssvssfer  feifpkessw  pkhtvtkgvt
151  ascshngkss  fyknllwlte  knglypnlsk  syvnnkekev  lvlwgvhhps
201  nigdqraiyl  tenayvsivs  shysrrftpe  iakrpkvrgq  egrinywtl
251  lepgdtiife  angnliapwy  afalsrgfgs  giitstasmg  ecdakcqtqp
301  gainsslpfq  nvhpvtigec  pkyvrstklr  mvtglrnips  iqsrgrlfgai
351  agfieggwtg  midgwygyhh  qneqsggyaa  dqkstqnaid  gitnkvnsvi
401  ekmntqftav  gkefnklerr  menlnkkvdd  gflldiwtyna  ellvllengr
451  tlgfhdsnvk  nlyekvknql  knnakeigng  cfefyhkcnn  ecmesvkngt
501  ydypkysees  klnrekidgv  klesmgvyqi  laiystvass  lvlvlslgai
551  sfwmcsngsl  qc
```

Figure 1P**SEQ ID NO:16**

Nucleotide Sequence of ca A/Shenzhen/227/95 N1 Entire molecule length: 1447 bp

```
1 agcaaaagca ggagttttaa atgaatccaa atcaaaaaat aataaccatt
51 ggatcaatca gtattgcaat tggaataatt agtctgatat tgcaaatagg
101 aaatattatt tcaatatggg ctagccactc aatccaaact ggaagtcaaa
151 accacactgg aatatgcaac caaagaatca ttacatatga aaatagcacc
201 tgggtaaatac aaacatatgt taatattaac aacactaatg ttgttgctgg
251 aaaggacaaa acctcaatga cattggccgg caattcatct ctttgcccta
301 tccgtggatg ggctatatac acaaaagaca acagcataag aattggttcc
351 aaaggagatg tttttgtcat aagagagcct tttatatcat gttctcactt
401 ggaatgcaga accttttttc tgaccaagg tgctctatta aatgacaagc
451 attcaaatgg gaccgttaag gacagaagcc cttatagggc cttaatgagc
501 tgtcctctag gtgaagctcc gtctccatac aattcaagat ttgaatcagt
551 tgcttggtca gcaagcgcat gccatgatgg cttgggctgg ctaacaatcg
601 gaatttctgg tccagataat ggggcagtgg ctgtactaaa atacaacggc
651 ataataactg aaaccattaa aagttggaag aagcgaatat taagaacaca
701 agagtctgaa tgtgtctgta tgaacggttc atgttttacc ataatgaccg
751 atggcccag taatggggcc gcatcgtaca gaatcttcaa aatcgagaag
801 gggagagtta ctaaatacat agagttggat gcacccaatt atcattacga
851 ggaatgttca tgttaccag acaccggcac agtgatgtgt gtgtgcaggg
901 acaattggca cggttcaaat cgaccttggg tgtcttttaa tcaaaacctg
951 gattatcaaa taggatacat ctgcagtggg gtgttcgggtg acaatccgcg
1001 tcccaaagat ggagaaggca gctgtaatcc agtgactgtt gatggagcag
1051 acggagtaaa ggggttttca tacagatatg gtaatggtgt ttggatagga
1101 aggactaaaa gtaacagact cagaaaggga tttgagatga tttgggatcc
1151 taatggatgg acagataccg acagtgattt ctcaatgaaa caggatatcg
1201 tggcaatgac tgattggtca gggtagagcg gaagttttgt tcaacatcct
1251 gagctaacag gattggactg tatgagacct tgcttttggg ttgaattagt
1301 cagagggcta cctagagaaa atacaacaat ctggactagt gggagcagca
1351 tttctttttg tggcgtaaat agcgatactg caaactggtc ttggccagac
1401 ggtgccgagt tgccattcac cattgacaag tagtccgttg aaaaaaa
```

SEQ ID NO:50

Amino Acid Sequence of ca A/Shenzhen/227/95 N1 Entire molecule length: 470 aa

```
1 mnpnqkiiti gsisiaigii slilqignii siwashsiqt gsqnhtgicn
51 griityenst wvnqtyvnin ntnvvagkdk tsmtlagnss lcpirgwaiy
101 tkdnsirigs kgdvfvirep fiscshlecr tffltqgall ndkhsngtvk
151 drspyrals cplgeapspy nsrfesvaws asachdglw ltigisgpdn
201 gavavlkyng iitetikswk krilrtqese cvcmngscft imtdgpsnga
251 asyrifkiek grvtsiield apnyhyeecs cypdtgtvmc vcrdnwhgsn
301 rpwvsfnqnl dyqigyicsg vfgdnprpkd gegscnpvtv dgadgvkqfs
351 yryngvwig rtksnrlrkq femiwdpngw tdt dsdfsmk qdivamtdws
401 gysgsfvqhp eltglcmlrp cfwvelvrgl prenttiwts gssisfcgvn
451 sdtanwswpd gaelpftidk
```

Figure 1Q

SEQ ID NO:17

ca A/Beijing/262/95

Nucleotide Sequence of ca A/Beijing/262/95 H1 Entire molecule length: 1775 bp

```
1 agcaaaagca ggggaaaata aaaacaacca aaatgaaagc aaaactacta
51 gtctgtttat gtacatttac agctacatat gcagacacaa tatgtatagg
101 ctaccatgcc aacaactcaa ccgacactgt tgacacagta cttgagaaga
151 atgtgacagt gacacactct gtcaacctac ttgaggacag tcacaattgga
201 aaactatgtc tactaaaagg aatagcccca ctacaattgg gtaattgcag
251 cgttgccgga tggatcttag gaaaccacga atgcgaatca ctgatttcta
301 aggaatcatg gtcctacatt gtagagacac caaacctga gaatggaaca
351 tgttaccag ggtatttcgc cgactatgag gaactgaggg agcaattgag
401 ttcagtatca tcatttgaga gattcgaaat attcccaaaa gaaagctcat
451 ggcccaaaca caccgtaaca ggagtaacgg catcatgctc ccataatggg
501 aaaagcagtt tttacagaaa tttgctatgg ctgacggaga agaatggcct
551 gtacccaaat ctgagcaatt cctatgtgaa caacaaagag aaagaagtcc
601 ttgtactatg ggggtgtcat caccatcta acatagggga ccaaagggcc
651 atctatcata cagaaaacgc ttatgtctct gtagtgtctt cacattatag
701 cagaagattc accccagaaa tagcaaaaag acccaaagta agaggtcagg
751 aaggagaat caactactac tggactctgc tggaaaccgg ggacacaata
801 atatttgagg caaatggaaa tctaatagcg ccatggtatg ctttcgcact
851 gagtagaggc tttgggtcag gaatcatcac ctcaaagca ccaatgaatg
901 aatgtgatgc gaagtgtcaa acacctcagg gagctataaa cagtagtctt
951 cctttccaga atgtacacc agtcacaata ggagagtgtc caaagtatgt
1001 caggagtaca aaattaagga tggttacagg actaaggaat atcccatcca
1051 ttcaatccag aggtttgttt ggagccattg ccggtttcat tgaagggggg
1101 tggactggaa tgatggatgg gtggtatggt tatcatcatc agaatgagca
1151 aggatctggc tatgtgcag atcaaaaaag cacacaaaat gccattaacg
1201 ggattacaaa taagtgtaat tctgtaattg agaaaatgaa cactcaattc
1251 acagctgtgg gcaaagaatt caacaaatta gaaagaagga tggaaaactt
1301 aaataaaaaa gttgatgatg gatttctaga catttggaata tataatgcag
1351 aattgttgg tctactggaa aatgaaagga ctttggaatt ccatgactca
1401 aatgtgaaga atctgtatga gaaagtgaag agccaattaa agaataatgc
1451 caaagaaata gggaacgggt gttttgaatt ctatcacaag tgaacaatg
1501 aatgcatgga aagtgtgaaa aatggaactt atgactatcc aaaatattcc
1551 gaagaatcaa agttaaacag ggagaaaatt gatggagtga aattggaatc
1601 aatgggagtc tatcagattc tggcgatcta ctcaactgtc gccagttcac
1651 tggttctttt ggtctccctg ggggcaatca gcttctggat gtgttccaat
1701 gggctcttgc agtgtagaat atgcactctga gaccagaatt tcagaaatat
1751 aagaaaaaac acccttgttt ctact
```

SEQ ID NO:51

Amino Acid Sequence of ca A/Beijing/262/95 H1 Entire molecule length: 565 aa

```
1 mkakllvllc tftatyadti cigyhannst dtvdtvlekn vtvthsvnl1
51 edshngklcl lkgiaplqlg ncsvagwilg npeceslisk eswsyivetp
101 npengtcypg yfadyeelre qlssvssfer feifpkessw pkhtvtgvta
151 scshngkssf yrnllwltek nglypnlsns yvnnkekevl vlgvvhpsn
201 igdqraiht enayvsvvss hysrrftpei akrpkvrgqe grinywtll
251 epgdtiifea ngnliapwya falsrgfgsg iitsnapmne cdakcqtppg
301 ainsslpfqv vhpvtigecp kyvrstklrm vtglrnipsi qsrqlfgaia
351 gfieggwtgm mdgwygyhhq neqsgyaad qkstqnaing itnkvnsvie
401 kmntqftavg kefnklerrm enlnkkvddg fldiwtynae llvllenert
451 ldfhdsnvkn lyekvksqk nnakeigngc fefyhkcne cmesvknqty
501 dypkyseesk lnrekidgvk lesmgvyqil aiystvassl vllvslgais
551 fwmcsngslq crici
```

Figure 1R**SEQ ID NO:18**

Nucleotide Sequence of ca A/Beijing/262/95 N1 Entire molecule length: 1463 bp

```
1 agcaaaagca ggagttttaa atgaatccaa atcaaaaaat aataaccatt
51 ggatcaatca gtatagtaat cgggataatt agtctaattg tgcaaatagg
101 aaatattatt tcaatatggg ctagtcactc aatccaaact ggaagtcaaa
151 accacactgg aatatgcaac caaagaatca tcacatatga aaatagcacc
201 tgggtgaatc acacatatgt taatattaac aacactaatg ttgttgctgg
251 aaaggacaaa acttcagtga cattggccgg caattcatca ctttgttcta
301 tcagtggatg ggctatatac acaaaagaca acagcataag aattggttcc
351 aaaggagatg tttttgtcat aagagagcct tttatatcat gttctcactt
401 ggaatgcaga accttttttc tgacccaagg tgctctatta aatgacaaac
451 attcaaattg gaccgttaag gacagaagtc cttatagggc cttaatgagc
501 tgtcctctag gcgaagctcc gtctccatat aattcaaagt ttgaatcagt
551 tgcttggcca gcaagcgcat gtcgatgatg catgggctgg ttaacaatcg
601 gaatttctgg tccagataat ggagcagtgg ctgtactaaa atacaacggc
651 ataataactg aaaccataaa aagttggaaa aagcgaatat taagaacaca
701 agagtctgaa tgtgtctgtg tgaacgggtc atgttttacc ataatgaccg
751 atggcccag taatggggcc gcctcgtaca aaatcttcaa gattgagaag
801 gggaaggtta ctaaataaat agagttgaat gcaccaatt ctcattatga
851 ggaatgttcc tgttaccag acactggcac agtgatgtgt gtatgcaggg
901 acaattggca cggttcaaat cgacctggg tgtcttttaa tcaaaacctg
951 gattatcaaa taggatacat ctgcagtggg gtgttcgggtg acaatccgcg
1001 tcccaaagat ggagagggca gctgtaatcc agtgactgtt gatggagcag
1051 acggagtaaa ggggttttca tacagatatg gtaatggtgt ttggatagga
1101 aggactaaaa gtaacagact cagaaaggga tttgagatga tttgggatcc
1151 taatggatgg acagataccg acagtgattt ctcagtgaag caggatgttg
1201 tggcaatgac tgattggcca gggtagacgc gaagtttcgt tcaacatcct
1251 gagctaacag gattggactg tataagacct tgcttctggg ttgaattagt
1301 cagaggacgg cctagagaaa atacaacaat ctggactagt gggagcagca
1351 tttctttttg tggcgtaaat agtgatactg caaactggtc ttggccagac
1401 ggtgctgagt tgccattcac cattgacaag tagtccgttg aaaaaaact
1451 ccttgtttct act
```

SEQ ID NO:52

Amino Acid Sequence of ca A/Beijing/262/95 N1 Entire molecule length: 470 aa

```
1 mnpnqkiiti gsisivigii slmlqignii siwashsiqt gsqnhtgicn
51 qriityenst wvnhtyvnin ntnvvagkdk tsvtlagnss lcsisgwaiy
101 tkdnsirigs kgdvvfirep fiscshlecr tffltqgall ndkhsngtvk
151 drspyrals cplgeapspy nskfesvaws asachdgmw ltigisgpdn
201 gavavlkyn g iitetikswk krilrtqese cvcvngscft imtdgpsnga
251 asykifkiek gkvtksieln apnshyeecs cypdtgtvmc vcrdnwhgsn
301 rpwvsfnqnl dyqigyicsg vfgdnprpkd gegscnptvt dgadgvkqfs
351 yryngvwig rtksnrlrkg femiwdpngw tdt dsdfsvk qdvvamtdws
401 gysgsfvqhp eltgldcirp cfwvelvrgr prenttiwts gssisfcgvn
451 sdtanwswpd gaelpftidk
```

Figure 1S

SEQ ID NO:19

ca A/New Caledonia/20/99

Nucleotide Sequence of ca A/New Caledonia/20/99 H1

Entire molecule length: 1775 bp

```
1 agcaaaagca ggggaaaata aaaacaacca aaatgaaagc aaaactactg
51 gtcctgttat gtacattttac agctacatat gcagacacaa tatgtatagg
101 ctaccatgcc aacaactcaa ccgacactgt tgacacagta cttgagaaga
151 atgtgacagt gacacactct gtcaacctac ttgaggacag tcacaatgga
201 aaactatgtc tactaaaagg aatagcccca ctacaattgg gtaattgcag
251 cgttgccgga tggatcttag gaaacccaga atgcgaatta ctgatttcca
301 aggaatcatg gtccctacatt gtagaaacac caaatcctga gaattggaaca
351 tgttaccag ggtatttcgc cgactatgag gaactgaggg agcaattgag
401 ttcagtatct tcatttgaga gattcgaaat attcccaaaa gaaagctcat
451 ggcccaaaca caccgtaacc ggagtatcag catcatgctc ccataatggg
501 aaaaacagtt ttacagaaa ttgctatgg ctgacgggga agaattggtt
551 gtacccaaac ctgagcaagt cctatgtaaa caacaaagag aaagaagtcc
601 ttgtactatg ggggtgttcat caccgccta acatagggga ccaaagggcc
651 ctctatcata cagaaaatgc ttatgtctct gtagtgtctt cacattatag
701 cagaagattc accccagaaa tagccaaaag acccaaagta agagatcagg
751 aaggaagaat caactactac tggactctgc tggaaacctg ggatacaata
801 atatttgagg caaatggaaa tctaatagcg ccatggtatg cttttgcaact
851 gagtagaggc ttgggatcag gaatcatcac ctcaaagtga ccaatggatg
901 aatgtgatgc gaagtgtcaa acacctcagg gagctataaa cagcagtcct
951 ctttccaga atgtacaccc agtcacaata ggagagtgtc caaagtatgt
1001 caggagtgc aaattgagga tggttacagg actaaggaac atcccatcca
1051 ttcaatccag aggtttgttt ggagccattg ccggtttcat tgaagggggg
1101 tggactgga tggtagatgg gtggtatggt tatcatcatc agaatgagca
1151 aggatctggc tatgctgcag atcaaaaaag tacacaaaat gccattaacg
1201 ggattacaaa caaggtgaat tctgtaattg agaaaatgaa cactcaattc
1251 acagctgtgg gcaaagaatt caacaaattg gaaagaagga tggaaaactt
1301 aaataaaaaa gttgatgatg ggtttctaga catttggaca tataatgcag
1351 aattgttgg tctactggaa aatgaaagga ctttggattt ccatgactcc
1401 aatgtgaaga atctgtatga gaaagtaaaa agccaattaa agaataatgc
1451 caaagaaata ggaaacgggt gttttgaatt ctatcacaag tgtaacaatg
1501 aatgcatgga gagtgtgaaa aatggaactt atgactatcc aaaatattcc
1551 gaagaatcaa agttaaacag ggagaaaatt gatggagtga aattggaatc
1601 aatgggagtc tatcagattc tggcgatcta ctcaactgtc gccagttccc
1651 tggttctttt ggtctccctg ggggcaatca gcttctggat gtgttccaat
1701 gggcttttgc agtgtagaat atgcatctga gaccagaatt tcagaagtat
1751 aagaaaaaac acccttgttt ctact
```

SEQ ID NO:53

Amino Acid Sequence of ca A/ New Caledonia /20/99 H1

Entire molecule length: 565 aa

```
1 mkakllvllc tftatyadti cigyhannst dtvdtvlekn vtvthsvnll
51 edshngklcl lkgiaplqlg ncsvagwilg npecellisk eswsyivetp
101 npengtcypg yfadyeelre qlssvssfer feifpkessw pkhtvtgvs
151 scshngknsf yrnllwltgk nglpnlsls yvnnkekevl vlgvvhppn
201 igdqrallyt enayvsvss hysrrftpei akrpkvrdqe grinyvwtll
251 epgdtiifea ngnliapwya falsrgfgsg iitsnapmde cdakcqtppg
301 ainsslpfqv vhpvtigecp kyvrsaklrm vtglrnipsi qsrqlfgaia
351 gfieggwtgm vdgwygyhhq neqsggyaad qkstqnaing itnkvnsvie
401 kmntqftavg kefnklerrm enlnkkvddg fldiwtynae llvllenert
451 ldfhdsnvkn lyekvksqk nnakeigngc fefyhkcne cmesvkngty
501 dypkyseesk lnrekidgvk lesmgvyqil aiystvassl vllvslgais
551 fwmcsngslq crici
```

Figure 1T

SEQ ID NO:20

Nucleotide Sequence of ca A/New Caledonia/20/99 N1

Entire molecule length: 1463 bp

```
1 agcaaaagca ggagttttaa atgaatccaa atcaaaaaat aataaccatt
51 ggatcaatca gtatagcaat cggaataatt agtctaattg tgcaaatagg
101 aaatattatt tcaatatggg ctagtcaact aatccaaact ggaagtcaaa
151 accacactgg agtatgcaac caaagaatca tcacatatga aaacagcacc
201 tgggtgaatc acacatatgt taatattaac aacactaatg ttgttgctgg
251 aaaggacaaa acttcagtga cattggccgg caattcatct ctttgttcta
301 tcagtggatg ggctatatac acaaaagaca acagcataag aattggctcc
351 aaaggagatg tttttgtcat aagagaacct ttcatatcat gttctcactt
401 ggaatgcaga accttttttc tgaccaagg tgctctatta aatgacaaac
451 attcaaattg gaccgttaag gacagaagtc cttatagggc cttaatgagc
501 tgtcctctag gtgaagctcc gtccccatac aattcaaagt ttgaatcagt
551 tgcattggta gcaagcgcat gccatgatgg catgggctgg ttaacaatcg
601 gaatttctgg tccagacaat ggagctgtgg ctgtactaaa atacaacggc
651 ataataactg aaaccataaa agtttgaaa aagcgaatat taagaacaca
701 agagtctgaa tgtgtctgtg tgaacgggtc atgtttcacc ataatgaccg
751 atggcccgag taatggggcc gcctcgtaca aaatcttcaa gatcgaaaag
801 ggaaggtta ctaaatcaat agagttgaat gcaccaatt ttcattatga
851 ggaatgttcc tgttaccag acactggcac agtgatgtgt gtatgcaggg
901 acaactggca tgggtcaaat cgaccttggg tgtcttttaa tcaaaacctg
951 gattatcaaa taggatacat ctgcagtggg gtgttcgggtg acaatccgcg
1001 tcccaaagat ggagagggca gctgtaatcc agtgactgtt gatggagcag
1051 acggagtaaa ggggttttca taaaaatag gtaatggtgt ttggatagga
1101 aggactaaaa gtaacagact tagaaagggg tttgagatga tttgggatcc
1151 taatggatgg acagataccg acagtgattt ctcaagtga aagaggtgtg
1201 tggcaataac tgattgggtc gggtagacgc gaagtttcgt tcaacatcct
1251 gagttaacag gattggactg tataagacct tgcttctggg ttgagttagt
1301 cagaggactg cctagagaaa atacaacaat ctggactagt gggagcagca
1351 tttctttttg tggcgtaaat agtgatactg caaactgggtc ttggccagac
1401 ggtgctgagt tgccgttcac cattgacaag tagttcgttg aaaaaaact
1451 ccttgtttct act
```

SEQ ID NO:54

Amino Acid Sequence of ca A/New Caledonia/20/99 N1

Entire molecule length: 470 aa

```
1 mnpnqiiti gsisiaigii slmlqignii siwashsiqt gsnhtgvcn
51 qriityenst wvnhtyvnn ntnvvagkdk tsvtlagnss lcsisgwaiy
101 tkdnsirigs kgdvfvirep fiscshlecr tffltqgall ndkhsngtvk
151 drspyrals cplgeapspy nskfesvaws asachdgmw ltigisgpdn
201 gavavlkyng iitetikswk krilrtqese cvcvngscft imtdgpsnga
251 asykifkiek gkvtksieln apnfhyeecs cypdtgtvmc vcrdnwhgsn
301 rpwvsfnqnl dyqigyicsg vfgdnprpkd gegscnptv dgadgvgkfs
351 ykygngvwig rtksnrlrkg femiwdpngw tdt dsdfsvk qdvvaitsws
401 gysgsfvqhp eltglcdirp cfwvelvrgl prenttiwts gssisfcgvn
451 sdtanwswpd gaelpftidk
```

Figure 1U

SEQ ID NO:21

ca B/Ann Arbor/1/94

Nucleotide Sequence of ca B/Ann Arbor/1/94 HA Entire molecule length: 1879 bp

```
1  agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt
51  actactcatg gtagtaacat ccaacgcaga tcgaatctgc actgggataa
101 catctttcaa ctcacctcat gtgggtcaaaa cagctactca aggggaagtc
151 aatgtgactg gtgtgatacc actgacaaca acaccaacaa aatctcattt
201 tgcaaatctc aaaggaacaa agaccagagg gaaactatgc ccaaactgtc
251 tcaactgcac agatctggat gtggccttgg gcagaccaat gtgtataggg
301 atcacacctt cggcaaaagc ttcaatactc cacgaagtca gacctgttac
351 atcggggtgc ttctctataa tgcacgacag aacaaaaatc agacagctac
401 ccaatcttct cagaggatat gaacatatca gattatcaac ccataacgtt
451 atcaacgcag aaagggcacc aggaggaccc tacagacttg gaacctcagg
501 atcttgccct aacgttacca gtagaagcgg attcttcgca acaatggctt
551 gggctgtccc aagggaacaa aaaacagcaa cgaaccactt aacagtagaa
601 gtaccataca tttgtacaaa aggagaagac caaattactg tttgggggtt
651 ccattctgat aacaaaatcc aaatgaaaaa cctctatgga gactcaaatc
701 ctcaaaagtt cacctcatct gccaatggaa taaccacaca ttatgtttct
751 cagattggtg gcttcccaaa tcaaacagaa gacggagggc taccacaaag
801 cggcagaatt gttgttgatt acatggtgca aaaacctggg aaaacaggaa
851 caattgtcta tcaaagaggt gttttgttgc ctcaaaaggt gtgggtgtgca
901 agtggcagga gcaaggtaat aaaagggtcc ttgcctttaa ttgggtgaagc
951 agattgcctt cacgaaaaat acggtggatt aaacaaaagc aagccttact
1001 acacaggaga acatgcaaaa gccataggaa attgccaat atgggtgaaa
1051 acacctttaa agcttgccaa tggaaacaaa tatagacctc ccgcaaaact
1101 attaaaggaa aagggtttct tcggagctat tgctggtttc ttagaaggag
1151 gatgggaagg aatgattgca ggttggcacg gatacacatc tcatggagca
1201 catgggggtg cagtggcagc agaccttaag agtacgcaag aagccataaa
1251 caagataaca aaaaatctca attctttgag tgagctagaa gtaaagaatc
1301 ttcaaagact aagtggtgcc atggatgaac tccacaacga aatactcgag
1351 ctggatgaga aagtggatga tctcagagct gacacaataa gctcgcaat
1401 agagcttgca gtcttgcttt ccaatgaagg aataataaac agtgaagatg
1451 agcatctatt ggcacttgag agaaaactaa agaaaatgct gggtcctct
1501 gctgtagaca tagggaatgg atgcttcgaa accaaacaca agtgcaacca
1551 gacctgctta gacaggatag ctgctggcac ctttaatgca ggagaatttt
1601 ctcttccac ttttgattca ctgaatatta ctgctgcac tttaaatgat
1651 gatggattgg ataatacatc tatactgctc tactactcaa ctgcggcttc
1701 tagtttggtt gtaacattga tgatagctat ttttattgtt tatatggtct
1751 ccagagacaa tgtttcttgc tccatctgtc tatagggaaa attgagccct
1801 gtattttcct ttattgtggt gcttgtttgc ttgttgccat tacagagaaa
1851 cgttattgaa aaatgctctt gttactact
```

SEQ ID NO:55

Amino Acid Sequence of ca B/Ann Arbor/1/94 HA Entire molecule length: 583 aa

```
1  mkaiivllmv vtsnadriect gitssnsphv vktatqgevn vtgviplttt
51  ptkshfanlk gtktrgklcp nclnctdldv algrpmcigi tpsakasilh
101 evrpvtsgcf pimhdrtkir qlpnllrgye hirlsthvni naerapggpy
151 rlgtsghsdn kiqmknlygd snpqkftssa ngitthyvsq iggfnpqted
201 gglpqsgriv vdymvqkpgk tgtivyqrgv llpqkvwcas grskvikgsl
301 pligeadclh ekygglnksk pyytgehaka igncpiwvkt plklangtky
351 rppakllkek gffgaiagfl eggwegmiag whgytshgah gvavaadlks
401 tqeainkitk nlslselev knlqrlsgam delhneilel dekvddlrad
451 tissqielav llsnegiins edehllaler klkmlgpsa vdigngcfet
501 khkenqtcl driaagtnag efslptfddl nitaaslndd gldnhtilly
551 ystaasslav tlmiaifivy mvsrdnvscs icl
```

Figure 1V

SEQ ID NO:22

Nucleotide Sequence of ca B/Ann Arbor/1/94 NA

Entire molecule length: 1554 bp

```
1 agcagaagca gagcatcttc tcaaaactga agtaaagagg ccaaaaatga
51 acaatgctac cttcaactat acaaacgtta accctatttc tcacatcagg
101 gggagtgtta ttatcactat atgtgtcagc cttactgtca tacttattgt
151 attcggatat attgctaaaa ttttcaccaa aaataattgc accaacaacg
201 tcgttggact gcgcgaacgc atcaaatgtt caggctgtga accattctgc
251 aacaaaagag atgaaattcc ttccccaga accggagtgg acataccccc
301 gtttatcttg ccagggttca accttcacga aagcactctt aattagccct
351 catagatttg gagaagccaa aggaaactca gctcccttga taataaggga
401 acctttttatt gcttgtggac caaaggagtg caaacacttt gctctaacc
451 attatgcagc tcaaccaggg ggatactaca atggaacaag agaggacaga
501 aacaagctga ggcactctgat ttcagtcaac ttaggcaaaa tccaactgt
551 agaaaactcc attttccata tggcagcttg gagtggatcc gcatgccatg
601 atggtagaga atggacatat atcggagtgt atggtcctga cagtaatgca
651 ttgatcaaaa taaaatatgg agaagcatac actgacacat accattccta
701 tgcaaacaac atcctaagaa cacaagaaag tgcctgcaat tgcatcgggg
751 gagattgtta tcttatgata actgatggct cagcttcagg aattagtaaa
801 tgcagattcc ttaagatccg agagggtcga ataataaaaag aaatatttcc
851 aacaggaagg gtagagcaca ctgaagaatg cacatgcgga tttgccagca
901 acaaaacat agaatgtgcc tgtagagata acagttacac agcaaaaaga
951 ccctttgtca aattaaatgt ggagactgat acagctgaaa taagattgat
1001 gtgcacagag acttatttgg acacccccag accagatgat ggaagcataa
1051 cagggccttg cgaatcta atgggacaaag ggagtggagg tgtcaaggga
1101 ggatttgttc atcaaagaat ggcattccaag attggaagat ggtactccc
1151 aacgatgtct aaaactaaaa gaatggggat ggaactgtat gtcaagtatg
1201 atggagaccc atggactgac agtgacgccc ttgctcctag tggagtaatg
1251 gtctcaatgg aagaacctgg ttggtactct ttcggcttcg aaataaaaga
1301 taagaaatgt gatgtcccct gtattgggat agagatggta catgatggtg
1351 gaaaaaggac ttggcactca gcagcaacag ccatttactg tttaatgggc
1401 tcaggacagt tgctatggga cactgtcaca ggtgttaata tggctctgta
1451 atggaggaat ggttgaatct gttctaaacc ctttgttcct attttatttg
1501 aacaattgtc cttactggac ttaattgttt ctgaaaaatg ctcttgttac
1551 tact
```

SEQ ID NO:56

Amino Acid Sequence of ca B/Ann Arbor/1/94 NA

Entire molecule length: 465 aa

```
1 mlpstiqtl lfltsggvll slyvsallsy llysdillkf spkiiaptts
51 ldcanasnvq avnhsatkem kflppepewt yprlscqgst fqkallisph
101 rfgeakgnsa pliirepfia cgpkeckhfa lthyaagpgg yyngtredrn
151 klrhlisvnl gkiptvensi fhmaawsgsa chdgrewtyi gvdgpdnsal
201 ikikygeayt dtyhsyanni lrtqesacnc iggcdylmit dgsasgiskc
251 rflkiregri ikeifptgrv ehteectcgf asnktiecac rdnsytakrp
301 fvklnvetdt aeirlmctet yldtprpddg sitgpcesng dkgsggvkkg
351 fvhqrmaski grwysrtmsk tkrmgmelyv kydgdpwtds dalapsgvmv
401 smeepgwysf gfeikdkkcd vpcigiemvh dggkrtwhsa ataiyclmgs
451 gqllwdvtg vnmal
```

Figure 1W

SEQ ID NO:23

ca B/Yamanashi/166/98

Nucleotide Sequence of ca B/Yamanashi/166/98 HA

Entire molecule length: 1881 bp

```
1  agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt
51  actactcatg gtagtaacat ccaatgcaga tcgaatctgc actgggataa
101 catcgtcaaa ctacacctcat gtggtcaaaa cagctactca aggggaggtc
151 aatgtgactg gtgtgatacc actgacaaca acaccaacaa aatctcattt
201 tgcaaattctc aaaggaacaa agaccagagg gaaactatgc ccaacctgtc
251 tcaactgcac agatctggat gtggccttag gcagaccaat gtgtgtgggg
301 gtcacacctt cggcaaaagc ttcaatactc cacgaagtca ggctgtttac
351 atccggatgc ttctctataa tgcacgacag acaaaaaatc agacagctac
401 ccaatcttct cagaggatat gaaaaaatca gattatcaac ccaaatcggt
451 atcaacgcag aaaaggcacc aggaggaccc tacagacttg gaacctcagg
501 atcttgccct aacgctacca gtagaagcgg atttttcgca acaatggcct
551 gggctgtccc aaaggacaac acaaaaacag caacgaatcc actaacagta
601 gaagtaccac acatctgtac aaaagaagaa gaccaaatta ctgtttgggg
651 gttccattct gatgacaaaa cccaaatgaa aaacctctat ggagactcaa
701 atcctcaaaa gttcacctca tctgctaata gagtaaccac acattatggt
751 tctcagattg gcggcttccc ggatcaaaca gaagacggag ggctaccaca
801 aagcggcaga attgttggtt attacatggt gcaaaaacct gggaaacag
851 gaacaattgt ctatcaaaga ggtattttgt tgctcaaaa ggtgtggtgc
901 gcgagtggca ggagcaaagt aataaaaggg tccttgccct taattggtga
951 agcagattgc cttcacgaaa aatacgggtg attaaacaaa agcaagcctt
1001 actacacagg agaacatgca aaagccatag gaaattgcc aatatgggtg
1051 aaaacacctt tgaagcttgc caatggaacc aaatatagac ctctgcacaa
1101 actattaag gaaaggggtt tcttcggagc tattgctggt ttcttagaag
1151 gaggatggga aggaatgatt gcaggttggc acggatacac atctcacgga
1201 gcacatggag tggcagtggc agcagacctt aagagtacgc aagaagccat
1251 aaacaagata acaaaaaatc tcaattcttt gagtgagcta gaagtaaaga
1301 atcttcaaag actaagtggg gccatggatg aactccacaa cgaaatactc
1351 gagctggatg agaaagtgga tgatctcaga gctgacacaa taagtcaca
1401 aatagaactt gcagtcttgc tttccaacga aggaataata aacagtgaag
1451 atgagcatct attggcactt gagagaaaac taaagaaaat gctgggtccc
1501 tctgctgtag acatagggaa tggatgcttc gaaaccaaac acaagtgcaa
1551 ccagacctgc ttagacagga tagctgctgg cacctttaat gcaggagaat
1601 tttctcttcc cacttttgat tcaactgaata ttactgctgc atctttaaat
1651 gatgatggat tggataacca tactatactg ctctactact caactgctgc
1701 ttctagtttg gctgtaacat tgatgatagc tatttttatt gtttatatga
1751 tctccagaga caatgtttct tgctccatct gtctataggg aaattaagcc
1801 ctgtattttc ctttattgta gtgcttggtt gcttgttatc attacaaaga
1851 aacgttattg aaaaatgctc ttgttactac t
```

SEQ ID NO:57

Amino Acid Sequence of ca B/Yamanashi/166/98 HA

Entire molecule length: 584 aa

```
1  mkaiivllmv vtsnadriect gitssnsphv vktatqgevn vtgviplttt
51  ptkshfanlk gktrgklcp tclnctdlv algrpmcvgv tpsakasilh
101 evrpvtsgcf pimhdrtkir qlpnllrgye kirlstqivi naekapggpy
151 rlgtsgscpn atsrsgffat mawavpkdnn ktatnpltve vphictkeed
201 qitvwgfhsd dktqmknlyg dsnpqkftss angvtthyvs qiggfpdqte
251 dgglpqsgri vdyvmvqkpg ktgtivyqrg illpqkvwca sgrskvikgs
301 lpligeadcl hekygglnks kpyytgehak aigncpiwvk tplklangtk
351 yrppakllke rgffgaiagf leggewegmia gwhgytshga hgvavaadlk
401 stqeainkit knlnslsele vknlqrlsga mdelhneile ldekvdllra
451 dtissqiela vllsnegiin sedehllale rklkklgpps avdigngcfe
501 tkhkcngtcl driaagtfnf gefslptfde lnitaaslnd dglndhtill
551 yystaassla vtlmiaifiv ymisrdnpsc sicl
```

Figure 1X

SEQ ID NO:24

Nucleotide Sequence of ca B/Yamanashi/166/98 NA Entire molecule length: 1557 bp

```
1  agcagaagca gagcatcttc tcaaaactga ggcaaatagg ccaaaaatga
51  acaatgctac cttcaactat acaaacgtta accctatttc tcacatcagg
101 gggagtgtta ttatcactat atgtgtcagc ttcactgtca tacttactat
151 attcgatat attgctaaaa ttttcaccaa cagaaataac tgcaccaaca
201 atgccattga attgtgcaaa cgcatacaat gttcaggctg tgaaccgttc
251 tgcaacaaaa ggggtgacac ttctctccc agaaccggag tggacatacc
301 ctcgtttatc ttgccgggac tcaacctttc agaaagcact cctaattagc
351 cctcatagat tcggagaaac caaaggaaac tcagctccct tgataataag
401 ggaacctttt attgcttggtg gaccaaagga atgcagacac tttgctctaa
451 ccattatgc agcccaacca gggggatact acaatggaac aagagaagac
501 agaaacaagc tgaggcatct aatttcagtc aaattgggca aaatcccaac
551 agtagaaaac tccattttcc acatggcagc ttggagcggg tccgcatgcc
601 atgatggtag agaatggaca tatatcggag ttgatggccc tgacagtaat
651 gcattgctca aaataaaaata tggagaagca tatactgaca cataccattc
701 ctatgcaaac aacatccctaa gaacacaaga aagtgcctgc aattgcatcg
751 ggggagattg ttatcttatg ataactgatg gctcagcttc agggattagt
801 gaatgcagat ttcttaagat tcgagagggc cgaataataa aagaaatatt
851 tccaacagga agagtagaac atactgaaga atgcacatgc ggatttgcca
901 gcaataaaac catagaatgt gcctgtagag ataacagtta cacagcaaaa
951 agaccctttg tcaaattaaa tgtggagact gatacagcag aaataagatt
1001 gatgtgcaca gagacttact tggacacccc cagaccagat gatggaagca
1051 taacagggcc ttgtgaatct aatggggata aagggagtgg aggcatacaag
1101 ggaggatttg ttcatcaaag aatggcatcc aagattggaa ggtggtactc
1151 tcgaacgatg tctaaaacta aaaggatggg gatgggactg tatgtcaagt
1201 atgatggaga cccatggatt gacagtgatg cccttactct tagcggagta
1251 atggtttcaa tggaagaacc tggttggtat tcctttggct tcgaaaataa
1301 agataagaaa tgtgatgtcc cctgtattgg gatagagatg gtacatgatg
1351 gtggaaagaa gacttggcac tcagcagcaa cagccattta ctgtttaatg
1401 ggctcaggac aactgctatg ggacactgtc acaggcgttg atatggctct
1451 gtaatggagg aatggttgag tctgttctaa accctttgtt cctattttgt
1501 ttgaacaatt gtccttactg aacttaattg tttctgaaaa atgctcttgt
1551 tactact
```

SEQ ID NO:58

Amino Acid Sequence of ca B/Yamanashi/166/98 NA

Entire molecule length: 466 aa

```
1  mlpstiqtlt lfltsggvll slyvsaslsy llysdillkf spteitaptm
51  plncanasnv qavnrsatkq vtlplpepew typrlscpgs tfqkallisp
101 hrfgetkgns apliirepfi acgpkecrhf althyaapqg gyyngtreldr
151 nklrhliisvk lgkiptvens ifhmaawsgs achdgrewty igvdgpdnsa
201 llkikygeay tdtyhsyann ilrtqesacn ciggdacylmi tdgsasgise
251 crflkiiregr iikeifptgr vehteectcg fasnktieca crdnsytakr
301 pfvklinvetd taeirlmcte tyldtprpdd gsitgpcesn gdkgsggikg
351 gfvhqrmask igrwysrtms ktkrmgmgly vkydgdpwid sdaltlsgvm
401 vsmeepgwys fgfeikdkkc dvpcigiemv hdggkktwhs aataiyclmg
451 sgqllwdtvt gvdmal
```

Figure 1Y

SEQ ID NO:25

ca B/Johannesburg/5/99

Nucleotide Sequence of ca B_Johannesburg_5_99_HA

Entire molecule length: 1882 bp

```
1 agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt
51 actactcatg gtagtaacat ccaatgcaga tcgaatctgc actgggataa
101 catcgtcaaa ctcacctcat gtggtcaaaa cagctactca aggggagggtc
151 aatgtgactg gtgcgatacc attgacaaca acaccaacaa aatctcattt
201 tgcaaatctc aaaggaacaa agaccagagg gaaactatgc ccaacctgtc
251 tcaactgcac agatctggat gtggccttgg gcagaccaat gtgtgtgggg
301 atcacacctt cggcaaaaagc ttcaatactc cacgaagtca gacctgttac
351 atccggatgc tttcctataa tgcacgcacg aacaaaaatc agacagctac
401 ccaatcttct cagaggatat gaaaaaatca gattatcaac ccaaacggtt
451 atcaacgcag aaaaggcacc aggaggaccc tacagacttg gaacttcagg
501 atcttgccct aacgctacca gtaaaagcgg atttttcgca acaatggctt
551 gggctgtccc aagggacaac aacaaaacag caacgaatcc actaacagta
601 gaagtaccac acatctgtac aaaagaagaa gaccaaatta ctgtttgggg
651 gttccattct gatgacaaaa cccaaatgaa aaacctctat ggagactcaa
701 atcctcaaaa gttcacctca tctgctaata gaataaccac acattatggt
751 tctcagattg gcggcttccc ggaccaaaca gaagacggag ggctaccaca
801 aagcggcaga attgttggtg attacatggt gcaaaaacct gggaaaacag
851 gaacaattgt ctatcaaaga gggatcttgt tgccctcaaaa ggtgtggtgc
901 gcgagtggca ggagcaaagt aataaaaggg tccttgccct taattggtga
951 agcagattgc cttcacgaaa aatacggtag attaaacaaa agcaagcctt
1001 actacacagg agaacatgca aaagccatag gaaattgccc aatatgggtg
1051 aaaacacctt tgaagcttgc caatggaacc aagtatagac ctctgcaaaa
1101 actattaaag gaaaggggtt tcttcggagc tattgctggt ttcttagaag
1151 gaggatggga aggaatgatt gcaggttggc acggatacac atctcacgga
1201 gcacacggag tggcagtggc agcagacctt aagagtacgc aagaagccat
1251 aaacaagata acaaaaaatc tcaattcttt gagtgagtta gaagtaaaga
1301 accttcaaag actaagtggg gccatggatg aactccataa cgaaatactc
1351 gagctggatg agaaagtgga tgatctcaga gctgacacaa taagctcaca
1401 aatagaactt gcagtcttgc tttccaacga aggaataata aacagtgaag
1451 atgagcatct attggcactt gagagaaaac taaagaagat gctgggtccc
1501 tctgctatag acatagggaa tggatgcttc gaaaccaaac acaagtgcaa
1551 ccagacctgc ttagacagga tagctgctgg cacctttaat gcaggagaat
1601 tttctcttcc cacttttgat tcaactgaaca ttactgctgc atctttaaat
1651 gatgatggat tggataacca tactatactg ctctactact caactgctgc
1701 ttctagtttg gctgtaacat tgatgatagc tatttttatt gtttatatga
1751 tctccagaga caatgtttct tgctccatct gtctataagg aaaattaagc
1801 cctgtatttt cctttattgt agtgcttgtt tgcttggtat cattacaaag
1851 aaacgttatt gaaaaatgct cttgttacta ct
```

SEQ ID NO:59

Amino Acid Sequence of ca B_Johannesburg_5_99_HA

Entire molecule length: 584 aa

```
1 mkaiivllmv vtsnadriect gitssnsphv vktatqgevn vtgaiplttt
51 ptkshfanlk gtktrgklcp tclnctdldv algrpmcvgi tpsakasilh
101 evrpvtsgcf pimhdrtkir qlpnllrgye kirlstqnvi naekapggpy
151 rlgtsgscpn atsksgffat mawavprdnk ktatnpltve vphictkeed
201 qitvwgfhsd dktqmknlyg dsnpqkftss angitthyvs qiggfpgdte
251 dgglpqsgri vvdymvqkpg ktgtivyqrg illpqkvwca sgrskvikgs
301 lpligeadcl hekygglnks kpyytgehak aigncpiwvk tplklangtk
351 yrppakllke rgffgaiagf leggwegmia gwhgytshga hgvavaadlk
401 stqeainkit knlnslsele vknlqrlsga mdelhneile ldekvdldla
451 dtissqiela vllsnegiin sedehllale rkllkmlgps aidigngcfe
501 tkhkcnqtc1 driaagtfna gefslptfds lnitaaslnl dglndhtill
551 yystaassla vtlmiaifiv ymisrdnvsc sicl
```

Figure 1Z**SEQ ID NO:26**

Nucleotide Sequence of ca B_Johannesburg_5_99_NA

Entire molecule length: 1557 bp

```
1  agcagaagca gagcatcttc tcaaaactga ggcaaatagg ccaaaaatga
51  acaatgctac cctcaactat acaaacgtta accctattcc tcacatcagg
101 gggagtgtta ttatcactat atgtgtcagc ttactgtca tacttactat
151 attcgatat attgctaaaa ttttcaccaa cagaaataac tgcaccagca
201 atgcccttgg attgtgcaaa cgcatacaat gttcaggctg tgaaccgttc
251 tgcaacaaaa ggggtgacac ttcttctccc agaaccggag tggacatacc
301 cgcggtttatc ttgcccgggc tcaacctttc agaaagcact cctaattagc
351 cctcatagat tcggagaaac caaaggaaac tcagctccct tgataataag
401 ggaacctttt attgcttgtg gaccaaagga atgcaaacac tttgctctaa
451 cccattatgc agcccaacca gggggatact acaatggaac aagagaagac
501 agaaacaagc taaggcatct aatttcagtc aaatttggta aaatcccaac
551 agtagaaaac tccattttcc acatggcagc atggagcggg tccgcatgcc
601 atgatggtaa agaatggaca tatatcggag ttgatggccc tgacagtaat
651 gcattgctca aaataaaata tggagaagca tatactgaca cataccattc
701 ctatgcaaac aacatcctaa gaacacaaga aagtgcctgc aattgcatcg
751 ggggaaattg ttatcttatg ataactgatg gctcagcttc aggtattagt
801 gagtgcagat ttcttaagat tcgagagggc cgaataataa aagaaatatt
851 tccaacagga agagtaaaac atactgaaga atgcacatgc ggatttgcca
901 gcaataaaac catagaatgt gcctgtagag ataacagtta cacagcaaaa
951 agaccctttg tcaaattaaa tgtggagact gatacagcag aaataagatt
1001 gatgtgcaca gagacttatt tggacacccc cagaccagat gatggaagca
1051 taacagggcc ttgtgaatct aatggggata aagggagtgg aggcatacaag
1101 ggaggatttg ttcatcaaag aatggcatcc aagattggaa ggtggtactc
1151 tcgaacaatg tctaaaacta aaaggatggg gatgggactg tatgtcaagt
1201 atgatggaga cccatggact gacagtgatg cccttgctct tagtggagta
1251 atggtttcaa tggaagaacc tggttggtac tcctttggct tcgaaataaa
1301 agataagaaa tgtgatgtcc cctgtattgg gatagagatg gtacatgatg
1351 gtggaaagga gacttggcac tcagcagcaa cagccattta ctgtttaatg
1401 ggctcaggac aactgctatg ggacactgtc acaggtgttg atatggctct
1451 gtaatggagg aatggttgag tctgttctaa accctttgtt cctattttgt
1501 ttgaacaatt gtccttactg aacttaattg tttctgaaaa atgctcttgt
1551 tactact
```

SEQ ID NO:60

Amino Acid Sequence of ca B_Johannesburg_5_99_NA

Entire molecule length: 466 aa

```
1  mlpstiqtlt lfltsggvll slyvsaslsy llysdillkf spteitapam
51  pldcanasnv qavnrsatkg vtlllpepew typrlscpgs tfqkallisp
101 hrfgetkgns apliirepfi acgpkeckhf althyaapqg gyyngtredr
151 nkrlrhliav fgkiptvens ifhmaawsgs achdgkewty igvdgpdsna
201 llkikygeay tdyhsyann ilrtqesacn ciggnacylmi tdgsasgise
251 crflkiregr iikeifptgr vkhteectcg fasnktieca crdnsytakr
301 pfvklinvetd taeirlmcte tyldtprpdd gsitgpcesn gdkgsggikg
351 gfvhqrmask igrwysrtms ktkrmgmgly vkydgdpwtd sdalalsgvm
401 vsmeepgwys fgfeikdkkc dvpcigiemv hdggketwhs aataiyclmg
451 sgqllwdtvt gvdmal
```

Figure 1AA

SEQ ID NO:27

ca B/Victoria/504/2000

Nucleotide Sequence of ca B/Victoria/504/2000 HA Entire molecule length: 1879 bp

```
1 agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt
51 actactcatg gtagtaacat ccaacgcaga tcgaatctgc actgggataa
101 catcttcaaa ctcacctcat gtggtcaaaa cagctactca aggggaagtc
151 aatgtgactg gtgtgatacc actgacaaca acaccaacaa aatctcattt
201 tgcaaatctc aaaggaacaa agaccagagg gaaactatgc ccaaactgtc
251 tcaactgcac agatctggat gtggccttgg gcagaccaat gtgtataggg
301 atcacacctt cggcaaaaagc ttcaatactc cacgaagtca gacctgttac
351 atccgggtgc tttcctataa tgcacgacag aacaaaaatc agacagctac
401 ccaatcttct cagaggatat gaacatatca gattatcaac ccataacgtt
451 atcaacgcag aaagggcacc aggaggacc tacagacttg gaacctcagg
501 atcttgccct aacgttacca gtagaagcgg attcttcgca acaatggctt
551 gggctgtccc aagggacaac aaaacagcaa cgaaccact aacagtagaa
601 gtaccataca tttgtacaaa aggagaagac caaattactg tttgggggtt
651 ccattctgat aacaaaatcc aaatgaaaaa cctctatgga gactcaaatc
701 ctcaaaagtt cacctcatct gccaatggaa taaccacaca ttatgtttct
751 cagattgggtg gtttcccaaa tcaaacagaa gacggagggc taccacaaag
801 cggcagaatt gttgttgatt acatggtgca aaaacctggg aaaacaggaa
851 caattgtcta tcaaagaggt gttttgttgc ctcaaaagggt gtgggtgtgca
901 agtggcagga gcaaggtaat aaaagggtcc ttgcctttaa ttgggtgaagc
951 agattgcctt cacgaaaaat acggtggatt aaacaaaagc aagccttact
1001 acacaggaga acatgcacaaa gccataggaa attgcccaat atgggtgaaa
1051 acacctttaa agcttgccaa tggaaacaaa tatagacctc ccgcaaaact
1101 attaaaggaa aagggtttct tcggagctat tgctggtttc ttagaaggag
1151 gatgggaagg aatgattgca ggttggcacg gatacacatc tcatggagca
1201 catgggggtg cagtggcagc agaccttaag agtacgcaag aagccataaa
1251 caagataaca aaaaatctca attctttgag tgagctagaa gtaaagaatc
1301 ttcaaagact aagtgggtgc atggatgaac tccacaacga aatactcgag
1351 ctggatgaga aagtggatga tctcagagct gacacaataa gctcgcgaat
1401 agagcttgca gtcttgcttt ccaatgaagg aataataaac agtgaagatg
1451 agcatctatt ggcacttgag agaaaaactaa agaaaatgct gggtcctctc
1501 gctgtagaca tagggaatgg atgcttcgaa accaaacaca agtgcaacca
1551 gacctgctta gacaggatag ctgctggcac ctttaatgca ggagaatttt
1601 ctcttccac ttttgattca ctgaatatta ctgctgcac tttaaatgat
1651 gatggattgg ataatacatc tatactgctc tactactcaa ctgctgcttc
1701 tagtttggtt gtaacattga tgatagctat ttttattggt tatatgggtc
1751 ccagagacaa tgtttcttgc tccatctgtc tatagggaaa attgagccct
1801 gtattttcct ttattgtggt gcttggttgc ttgttgccat tacagagaaa
1851 cgttattgaa aaatgctctt gttactact
```

SEQ ID NO:61

Amino Acid Sequence of ca B/Victoria/504/2000 HA

Entire molecule length: 583 aa

```
1 mkaiivllmv vtsnadriect gitssnsphv vktatqgevn vtgviplttt
51 ptkshfanlk gtktrgklcp nclnctdldv algrpmcigi tpsakasilh
101 evrpvtsgcf pimhdrtkir qlpnllrgye hirlsthnvi naerapggpy
151 rlgtsgscpn vtsrsgffat mawavprdnk tatnpltvev pyictkgedq
201 itvwgfhsdn kiqmknlygd snpqkftssa ngitthyvsq iggfpnqted
251 gglpqsgriv vdymvqkpgk tgtivyqrgv llpqkvwcas grskvikgsl
301 pligeadclh ekygglnksk ppytgehaka igncpiwvkt plklangtky
351 rppakllkek gffgaiagfl egwegmiag whgytshgah gvavaadlks
401 tqeainkitk nlslselev knlqrlsgam delhneilel dekvddlrad
451 tissqielav llsnegiins edehllaler klkklmgpsa vdigngcfet
501 khkcngtcld riaagtfnag efslptfddl nitaaslndd gldnhtilly
551 ystaasslav tlmiaifivy mvsrndvscs icl
```

Figure 1BB**SEQ ID NO:28**Nucleotide Sequence of ca B/Victoria/504/2000 NA
bp

Entire molecule length: 1554

```
1  agcagaagca gagcatcttc tcaaaactga agtaaagagg ccaaaaatga
51  acaatgctac cttcaactat acaaacgtta accctatttc tcacatcagg
101 gggagtgtta ttatcactat atgtgtcagc cttactgtca tacttattgt
151 attcgatat attgctaaaa ttttcaccaa aaataattgc accaacaacg
201 tcgttggact gcgcgaacgc atcaaagtgt caggctgtga accattctgc
251 aacaaaagag atgaaattcc tccccccaga accggagtgg acataccccc
301 gtttatcttg ccagggttca accttcacaga aagcactctt aattagccct
351 catagatttg gagaagccaa aggaaactca gctcccttga taataaggga
401 accttttatt gcttgtggac caaaggagtg caaacacttt gctctaacc
451 attatgcagc tcaaccaggg ggatactaca atggaacaag agaggacaga
501 aacaagctga ggcactctgat ttcagtcaac ttaggcaaaa tcccaactgt
551 agaaaactcc attttccata tggcagcttg gagtggatcc gcatgccatg
601 atggtagaga atggacatat atcggagttg atggtcctga cagtaatgca
651 ttgatcaaaa taaaatatgg agaagcatac actgacacat accattccta
701 tgcaaaacaac atcctaagaa cacaagaaaag tgccctgcaat tgcacgggg
751 gagattgtta tcttatgata actgatggct cagcttcagg aattagtaaa
801 tgcagattcc ttaagatccg agagggtcga ataataaaag aaatatattcc
851 aacaggaagg gtagagcaca ctgaagaatg cacatgcgga tttgccagca
901 acaaaacccat agaatgtgcc tgtagagata acagttacac agcaaaaaga
951 ccctttgtca aattaaatgt ggagactgat acagctgaaa taagattgat
1001 gtgcacagag acttatttgg acacccccag accagatgat ggaagcataa
1051 cagggccttg cgaatcta atgggacaaaag ggagtggagg tgtcaaggga
1101 ggatttgttc atcaaagaat ggcacccaag attggaagat ggtactcccg
1151 aacgatgtct aaaactaaaa gaatggggat ggaactgtat gtcaagtatg
1201 atggagaccc atggactgac agtgacgccc ttgctcctag tggagtaatg
1251 gtctcaatgg aagaacctgg ttggtactct ttcggcttcg aaataaaaga
1301 taagaaatgt gatgtcccct gtattgggat agagatggta catgatgggtg
1351 gaaaaaggac ttggcactca gcagcaacag ccatttactg tttaatgggc
1401 tcaggacagt tgctatggga cactgtcaca ggtgttaata tggctctgta
1451 atggaggaat ggttgaatct gttctaaacc ctttgttcct attttatttg
1501 acaattgtc cttactggac ttaattgttt ctgaaaaatg ctcttgttac
1551 tact
```

SEQ ID NO:62

Amino Acid Sequence of ca B/Victoria/504/2000 NA

Entire molecule length: 465 aa

```
1  mlpstiqtlt lfltsggvll slyvsallsy llysdillkf spkiiaptts
51  ldcanasnvq avnhsatkem kflppepewt yprlscqgst fqkallishp
101 rlfgeakgnsa pliirepfia cgpkeckhfa lthyaagpgg yyngtredrn
151 klrhlisvnl gkiptvensi fhmaawsgsa chdgrewtyi gvdgpdsnal
201 ikikygeayt dtyhsyanni lrtqesacnc iggdcyllmit dgsasgiskc
251 rflkiregri ikeifptgrv ehteectcgf asnktiecac rdnsytakrp
301 fvklnvetdt aeirlmctet yldtprpddg sitgpcesng dkgsggvkqg
351 fvhqrmaski grwysrtmsk tkrmgmelyv kydgdpwtds dalapsgvmv
401 smeepgwysf gfeikdkkcd vpcigiemvh dggkrtwhsa ataiyclmgs
451 gqllwdvtg vnmal
```

Figure 1CC

SEQ ID NO:29

ca B/Hong Kong/330/01

Nucleotide Sequence of ca B/Hong Kong/330/01 HA Entire molecule length: 1885 bp

```
1  agcagaagca  gagcattttc  taatatccac  aaaatgaagg  caataattgt
51  actactcatg  gtagtaacat  ccaatgcaga  tcgaatctgc  actggaataa
101 catcgtcaaa  ctacccccat  gtggtcaaaa  ctgctactca  aggggaagtc
151 aatgtgactg  gtgtgatacc  actgacaaca  acaccacca  aatctcattt
201 tgcaaattct  aaaggaacaa  aaaccagagg  gaaactatgc  ccaaaatgtc
251 tcaactgcac  agatctggac  gtggccttgg  gcagaccaa  atgcacgggg
301 aacatacctt  cggcaaaagt  ttcaatactc  catgaagtaa  gacctgttac
351 atctgggtgc  ttctctataa  tgacgcagag  aacaaaaatt  agacagctgc
401 ccaatcttct  cagaggatac  gaacgtatca  ggttatcaa  ccataacgtt
451 atcaatgcag  aaaaagcacc  aggaggacc  tacaaaattg  gaacctcagg
501 gtcttgccct  aacgttacca  atggaaacgg  attcttcgca  acaatggctt
551 gggctgtccc  aaaaaacgaa  aacaacaaa  cagcaacaaa  ttcattaaca
601 atagaagtac  catacatttg  tacagaagga  gaagaccaa  ttaccgtttg
651 ggggttccac  tctgatagcg  aaacccaaat  ggcaaaactc  tatggagact
701 caaagcctca  gaagtccact  tcatctgcta  acggagtgc  cacacattac
751 gtttcacaga  ttggtggctt  cccaaatcaa  acagaagacg  gaggactacc
801 acaaagtgg  agaattggtt  ttgattacat  ggtgcaaaa  tctggaaaaa
851 caggaacaat  tacctatcaa  agaggatatt  tattgcctca  aaaagtgtgg
901 tgcgcaagt  gcaggagcaa  ggtaataaaa  ggatccttgc  ctttaattgg
951 agaagcagat  tgcctccacg  aaaaatacgg  tggattaaac  aaaagcaagc
1001 cttactatac  aggggaacat  gcaaaagcca  taggaaattg  cccaatatgg
1051 gtgaaaacac  ccttgaagct  ggccaatgga  accaaatata  gacctcctgc
1101 aaaactatta  aaggaaaggg  gtttcttcgg  agctattgct  ggtttcttag
1151 aaggaggatg  ggaagggaat  attgcagggt  ggcacggata  cacatcccat
1201 ggagcacatg  gagtagcagt  ggacgcagac  cttaagagta  ctcaagaagc
1251 cataaacaag  atcacaaaaa  atctcaactc  tttgagtgc  ctggaagtaa
1301 agaattctca  aagactaagc  ggagccatgg  atgaactcca  caacgaaata
1351 ctagaactag  atgagaaagt  ggatgatctc  agagctgata  caataagctc
1401 gcaaatagaa  ctgcagctct  tgctttccaa  tgaaggaata  ataaacagtg
1451 aagatgagca  tctcttggcg  cttgaaagaa  aactgaagaa  aatgctgggc
1501 ccctctgctg  tagagatagg  gaatggatgc  ttcgaaacca  aacacaagtg
1551 caaccagacc  tgcctcgata  gaatagctgc  tggcaccttt  aatgcaggag
1601 aattttctct  cccaccttt  gattcactaa  atattactgc  tgcattctta
1651 aatgacgatg  gattggataa  tcatactata  ctgctttact  actcaactgc
1701 tgcttccagt  ttggctgtaa  cattgatgat  agctatcttt  gttgtttata
1751 tgggtctccag  agacaatgtt  tcttgttcca  tctgtctata  aggaaagtta
1801 agccccgtat  tttcctttat  tgtagtactt  gtttgcttgc  tatcattaca
1851 aaaaaacgtt  attgaaaaat  gctcttggtt  ctact
```

SEQ ID NO:63

Amino Acid Sequence of ca B/Hong Kong/330/01 HA

Entire molecule length: 585 aa

```
1  mkaiivllmv  vtsnadriect  gitssnsphv  vktatqgevn  vtgviplttt
51  ptkshfanlk  gtktrgklcp  kclnctdlldv  algrpkctgn  ipsakvsilh
101 evrpvtsgcf  pimhdrtkir  qlpnllrgye  rirlsnhnvi  naekapggpy
151 kigtsgscpn  vtngngffat  mawavpknen  nktatnslti  evpyictege
201 dqitvwgfhs  dsetqmakly  gdsqpqkfts  sangvtthyv  sqiggfnpqt
251 edgglpqsg  ivvdymvqks  gktgtityqr  gillpqkvwc  asgrskvikg
301 slpligeadc  lhekygglnk  skpyytgeha  kaigncpiwv  ktplklangt
351 kyrppakllk  ergffgaiag  fleggewegmi  agwhgytshg  ahgvavaadl
401 kstqeainki  tknlslslsel  evknlqrlsg  amdelhneil  eldekvdldl
451 adtissqiel  avllsnegii  nsedehllal  erklkkmlgp  saveigngcf
501 etkhkcnqtc  ldriaagtfn  agefslptfd  slnitaasln  ddgldnhtil
551 lyystaassl  avtlmiaifv  vymvsrdnvs  csicl
```

Figure 1DD**SEQ ID NO:30**

Nucleotide Sequence of ca B/Hong Kong/330/01 NA Entire molecule length: 1544 bp

```
1 agcagagcat cttctcaaaa ctgaagcaaa taggccaaaa tgaacaatgc
51 taccctcaac tatacaaaaca ttaaccctat ttctcacatc agggggagtg
101 ttattatcac tatatgtgtc agccttactg tcatacttac tgtattcgga
151 tatattgcta aaattttcac caacaaaaat aattgcacca acaacgtcgt
201 tggactccgc gaacgcatca aattttcagg ccgtgaacca ttctgcaaca
251 aaagagatga catttcttct cccagaaccg gagtggacat accctcgttt
301 atcttgccag ggttcaacct ttcaaaaagc actcctaatt agccctcata
351 gattcggaga agccaaagga aactcagctc ccttgataat aagggaacct
401 tttattgctt gtggaccaa ggagtgtaaa cactttgctc taaccatta
451 tgcagctcaa ccagggggat actacaatgg aacaagagag gacagaaaca
501 agctgaggca tctgatttca gtcaacttag gcaaaatacc aactgtagaa
551 aactccattt tccacatggc agcttggagt ggggccgcat gccatgatgg
601 tagagagtgg acttatatcg gagttgatgg ccctgacagt aatgcattga
651 tcaaaataaa atatggagaa gcatacactg acacatacca ttcctatgca
701 aacaacatcc taagaacaca agaaagtgcc tgcaactgca tcgggggaga
751 ttgttatctt atgataactg atggctcagc ttcaggaatt agtaaatgca
801 gattccttaa gattcgagag ggtcgaatag taaaagaaat atttccaaca
851 ggaagagtag agcatactga agaatgcaca tgcggatttg ccagcaataa
901 aaccatagaa tgtgcctgta gagataacag ttacacagca aaaagacctt
951 ttgtcaaatt aaatgtggaa actgatacag cagaaataag attgatgtgc
1001 acagagactt atttggacac ccccagacca gatgatggaa gcataacagg
1051 gccttgcgaa tctaatgggg acaaaggagg tggaggatc aaggagggat
1101 ttgtccatca aagaatggca tccaagattg gaagatggta ctctcgaacg
1151 atgtctaaaa ctaaaagaat ggggatggaa ctgtatgtca agtatgatgg
1201 agacccatgg actgacagtg atgcccttgc tcctagtgga gtaatggtct
1251 caatagaaga acctggttgg tattctttcg gcttcgaaat aaaagataag
1301 aaatgcgatg tcccctgtat tgggatagag atggtacacg atggtggaaa
1351 aacaacttgg cactcagcag caacagccat ttactgttta atgggctcag
1401 gacagttgct atgggacact atcacaggtg ttgatatggc tctgtaatgg
1451 aggaatggtt gaatctgttc taaacccttt gttcctattt tgtttgaaca
1501 attgtcctta ctggacttaa ttgtttctga aaaatgctct tgtt
```

SEQ ID NO:64

Amino Acid Sequence of ca B/Hong Kong/330/01 NA

Entire molecule length: 466 aa

```
1 mlpstiqtl t lfltsggvll slyvsallsy llysdillkf sptkiiaptt
51 sldsanasnf qavnhsatke mtfllepew typrlscqgs tfqkallisp
101 hrfgeakgns apliirepfi acgpkeckhf althyaagpg gyyngtreldr
151 nkrlrhlsnv lgkiptvens ifhmaawsgs achdgrewty igvdgpdnsa
201 likikygeay tdyhsyann iltqesacn ciggd cylmi tdgsasgisk
251 crflkiregr ivkeifptgr vehteectg fasnktieca crdnsytakr
301 pfvklrvetd taeirlmcte tyldtprpdd gsitgpcesn gdksggikg
351 gfvhqrmask igrwysrtms ktkrmgmely vkydgdptwd sdalapsgvm
401 vsieepgwys fgfeikdkkc dvpcigiemv hdggkttwhs aataiyclmg
451 sgqllwdtit gvdmal
```

Figure 1EE

SEQ ID NO:31

ca B/Brisbane/32/2002

Nucleotide Sequence of ca B_Brisbane_32_2002_HA

Entire molecule length: 1885 bp

```
1 agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt
51 actactcatg gtagtaacat ccaatgcaga tcgaatctgc actgggataa
101 catcgtcaaa ctcaccccat gtggtcaaaa ctgctactca aggggaggtc
151 aatgtgactg gtgtgatacc actgacaaca acaccaccca aatctcattt
201 tgcaaatctc aaaggaacaa aaaccagagg gaaactatgc ccaaaatgcc
251 tcaactgcac agatctggac gtggccttgg gcagaccaa atgcacgggg
301 aacataccct cggcaaaagt ttcaatactc catgaagtca gacctgttac
351 atctgggtgc tttcctataa tgcacgacag aacaaaaatt agacagctgc
401 ccaatcttct cagaggatac gaacatatca gggttatcaac tcataacggt
451 atcaatgcag aaaaggcacc aggaggacc tacaaaaatt gaacctcagg
501 gtcttgccct aacgtttacca atggaaacgg atttttcgca acaatggctt
551 gggccgtccc aaaaaacgac aacaacaaaa cagcaacaaa ttcattaaca
601 atagaagtac catacatttg tacagaagga gaagaccaa ttaccgtttg
651 ggggttccac tctgataacg aagcccaaat ggcaaaactc tatggggact
701 caaagcccca gaagttcacc tcctctgcca acggagtgc cacacattac
751 gtttcacaga ttggtggctt cccaaatcaa acagaagacg gaggactacc
801 acaaagtggg agaattggtt ttgattacat ggtgcaaaaa tctgggaaaa
851 caggaacaat tacctatcaa agaggatatt tattgcctca aaaagtgtgg
901 tgcgcaagtg gcaggagcaa ggtaataaaa ggatccttgc ctttaattgg
951 agaagcagat tgctccacg aaaaatacgg tggattaaac aaaagcaagc
1001 cttactacac aggggaacat gcaaaggcca taggaaattg cccaatatgg
1051 gtgaaaacac ccttgaagct ggccaatgga accaaatata gacctcctgc
1101 aaaactatta aaggaaagag gtttcttcgg agctattgct ggtttcttag
1151 aaggaggatg ggaaggatg attgcagggt ggacaggata cacatcccat
1201 ggggcacatg gagtagcagt ggcagcagac cttaagagta ctcaagaagc
1251 cataaacaag ataacaaaaa atctcaactc tttgagttag ctggaagtaa
1301 agaatcttca aagactaagc ggtgccatgg atgaactcca caacgaaata
1351 ctagaactag acgagaaagt ggatgatctc agagctgata caataagctc
1401 acaaatagaa ctgcagctct tgctttccaa tgaaggaata ataaacagtg
1451 aagatgagca tctcttggcg cttgaaagaa agctgaagaa aatgctgggc
1501 ccctctgctg tagagatagg gaatggatgc ttcgaaacca aacacaagtg
1551 caaccagacc tgtctcgaca gaatagctgc tggtaacctt gatgcaggag
1601 aattttctct ccccactttt gattcactga atattactgc tgcattctta
1651 aatgacgatg gattggataa tcatactata ctgctttact actcaactgc
1701 tgctccagt ttggctgtaa cattgatgat agctatcttt gttgtttata
1751 tggctccag agacaatggt tcttgctcca tctgtctata aggaaagtta
1801 agccctgtat tttcctttat tgtagtgcct gtttgcttgt taccattaca
1851 aaaaaacggt attgaaaaat gctcttgcta ctact
```

SEQ ID NO:65

Amino Acid Sequence of ca B_Brisbane_32_2002_HA

Entire molecule length: 585 aa

```
1 mkaiivllmv vtsnadriat gitssnsphv vktatqgevn vtgviplttt
51 ptkshfanlk gtktrgklcp kclnctdldv algrpkctgn ipsakvsilh
101 evrpvtsgcf pimhdrtkir qlpnllrgye hirlsthvni naekapggpy
151 kigtsgscpn vtngngffat mawavpkndn nktatnslti evpyictege
201 dqitvwgfhs dneaqmakly gdsqpkfts sangvtthyv sqiggfnpqt
251 edgglpqsgv ivvdyvmvqks gktgtityqr gillpqkvwc asgrskvikg
301 slpligeadc lhekyyglnk skpyytgeha kaigncpiwv ktplklangt
351 kyrppakllk ergffgaiag fleggwegmi agwhgytshg ahgvavaadl
401 kstqeainki tknlslslsel evknlqrlsg amdelhneil eldekvdldl
451 adtissqiel avllsnegii nsedehllal erkklkmlgp saveigngcf
501 etkhkcnqtc ldriaagtfd agefslptfd slnitaasl n ddgldnhtil
551 lyystaassl avtlmiaifv vymvsrdnvs csicl
```

Figure 1FF**SEQ ID NO:32**

Nucleotide Sequence of ca B_Brisbane_32_2002_NA

Entire molecule length: 1557 bp

```
1 agcagaagca gagcatcttc tcaaaactga ggcaaatagg ccaaaaatga
51 acaatgctac cttcaactat acaaacgtta accctatttc tcacatcagg
101 gggagtatta ttatcactat atgtgtcagc ttcattgtca tacttactat
151 attcggatat attgctaaaa ttctcaccaa cagaaataac tgcaccaaca
201 atgccattgg attgtgcaaa cgcatcaaat gttcaggctg tgaaccgttc
251 tgcaacaaaa ggggtgacac ttcttctccc agaaccagag tggacatacc
301 cgcgtttatc ttgcccgggc tcaacctttc agaaagcact cctaattagc
351 cctcatagat tcggagaaaac caaaggaaac tcagctccct tgataataag
401 ggaacctttt attgcttgtg gaccaaagga atgcaaacac tttgctctaa
451 ccattatgc agcccaacca gggggatact acaatggaac aagaggagac
501 agaaacaagc tgaggcatct aatttcagtc aaattgggca aaatcccaac
551 atagaaaaac tccattttcc acatggcagc atggagcggg tccgcatagc
601 atgatggtaa agaattggaca tatatcggag ttgatggccc tgacaataat
651 gcattgctca aaataaaaata tggagaagca tatactgaca cataccattc
701 ctatgcaaac aacatcctaa gaacacaaga aagtgcctgc aattgcatcg
751 ggggaaattg ttatcttatg ataactgatg gctcagcttc aggtattagt
801 gaatgcagat ttcttaaaat tcgagagggc cgaataataa aagaaatatt
851 tccaacagga agagtaaaac atactgaaga atgcacatgc ggatttgcca
901 gcaataagac catagaatgt gcctgtagag ataacagtta cacagcaaaa
951 agaccctttg tcaaattaaa cgtggagact gatacagcag aaataagatt
1001 gatgtgcaca gagacttatt tggacacccc cagaccagat gatggaagca
1051 taacagggcc ttgtgaatct aatggggaca aaggagtggt aggcattcaag
1101 ggaggatttg ttcatcaaag aatggcatcc aagattggaa ggtggtactc
1151 tcgaacgatg tctaaaacta aaaggatggg gatgggactg tatgtcaagt
1201 atgatggaga cccatgggct gacagtgatg cccttgctct tagtggagta
1251 atggtttcaa tggagaacc tggttggtac tcctttggct tcgaaataaa
1301 agataagaaa tgtgatgtcc cctgtattgg aatagagatg gtacatgatg
1351 gtggaaaaga gacttggcac tcagcagcaa cagccattta ctgtttaatg
1401 ggctcaggac agctgctgtg ggacactgtc acaggtgttg atatggctct
1451 gtaatggagg aatggttgag tctgttctaa accctttgtt cctattttgt
1501 ttgaacaatt gtccttactg aacttaattg tttctgaaaa atgctcttgt
1551 tactact
```

SEQ ID NO:66

Amino Acid Sequence of ca B_Brisbane_32_2002_NA

Entire molecule length: 466 aa

```
1 mlpstiqtlf lfltsggvll slyvsaslsy llysdillkf spteitaptm
51 pldcanasnq qavnrsatkg vtlllpepew typrlscpgs tfqkallisp
101 hrfgetkgns apliirepfi acgpkeckhf althyaapqg gyyngtrgdr
151 nkrlrhlsvk lgkiptvens ifhmaawsgs achdgkewty igvdgpdnna
201 llkikygeay tdyhsyann ilrtqesacn ciggnclmi tdgsasgise
251 crflkiregr iikeifptgr vkhteectcg fasnktieca crdnsytakr
301 pfvklrvetd taeirlmcte tyldtprpdd gsitgpcesn gdksggikg
351 gfvhqrmasq igrwysrtms ktkrmgmgly vkydgdwad sdalalsgvm
401 vsmeepgwys fgfeikdkkc dvpcigiemv hdgkktwhs aataiyclmg
451 sgqllwdtvt gvdmal
```

Figure 1GG

SEQ ID NO:33

ca B/Jilin/20/2003

Nucleotide Sequence of ca B/Jilin/20/03 HA Entire molecule length: 1853 bp

```
1 tctaatatcc acaaaatgaa ggcaataatt gtactactca tggtagtaac
51 atccaatgca gatcgaatct gcaactgggat aacatcttca aactcacctc
101 atgtggtcaa aacagctact caaggggagg tcaatgtgac tgggtgaata
151 ccactgacaa caacaccaac aaaatcttat ttgcaaatc tcaaaggaac
201 aaggaccaga gggaaactat gtccagactg tctcaactgt acagatctgg
251 atgtggcctt gggcagacca atgtgtgtgg ggaccacacc ttcgcaaaa
301 gcttcaatac tccacgaagt cagacctgtt acatccgggt gctttcctat
351 aatgcacgac agaacaaaaa tcagacaact acccaatctt ctcaaggat
401 atgaaaatat cagattatca acccaaaacg ttatcgatgc agaaaatgca
451 ccaggaggac cctacagact tggaaacctc ggatcttgcc ctaacgctac
501 cagtaaaagc ggatttttctg caacaatggc ttgggctgtc ccaaaggaca
551 acaacaaaaa tgcaacgaac ccactaacag tagaagtacc atacgtttgt
601 acagaagggg aagaccaa atactgtttg gggttccatt cagataacaa
651 aacccaatg aagaacctct atggagactc aaatcctcaa aagttcacct
701 catctgctaa tggagtaacc acacattatg tttctcagat tggcggtctc
751 ccagctcaaa cagaagacga aggactacca caaagcggca gaattgttgt
801 tgattacatg gtgcaaaaaa ctaggaaaac aggaacaatt gtctatcaaa
851 gaggtgtttt gttgcctcaa aagggtgtgtt gcgcgagtgg caggagcaaa
901 gtaataaaaag ggtccttgcc tttaattggt gaagcagatt gccttcatga
951 aaaatacggg ggattaaaca aaagcaagcc ttactacaca ggagaacatg
1001 caaaagccat aggaaattgc ccaatatggg tgaaaacacc tttgaagctt
1051 gccaatggaa ccaaatatag acctcctgca aaactattaa aggaaagggg
1101 tttcttcgga gctattgctg gtttcctaga aggaggatgg gaaggaatga
1151 ttgcaggttg gcacggatac acatctcacg gagcacatgg agtggcagtg
1201 gcggcgagacc ttaagagtac gcaagaagct ataaacaaga taacaaaaaa
1251 tctcaattct ttgagttagc tagaagtaaa gaatcttcaa agactaagtg
1301 gtgccatgga tgaactccac aacgaaatac tcgagctgga tgagaaagtg
1351 gatgatctca gagctgacac tataagctcg caaatagaac ttgcagctct
1401 gctttccaat gaaggaataa taaacagtga agatgagcat ctattggcac
1451 ttgagagaaa actaaagaaa atgtctgggtc cctctgctgt agacatagga
1501 aatggatgct tcgaaaccaa acacaagtgc aaccagacct gcttagacag
1551 gatagctgct ggcaccttta atgcaggaga attttctctc cccacttttg
1601 attcactgaa cattactgct gcactcttaa atgatgatgg attggataac
1651 catactatac tgctctatta ctcaactgct gcttctagtt tggctgtaac
1701 attgatgcta gctattttta ttgtttatat ggtctccaga gacaacgttt
1751 catgctccat ctgtctataa ggaagattaa gccttgtatt ttcttttatt
1801 gtatgtgctt tttgcttgct atcattacaa agaaacgtta ttgaaaaaatg
1851 ctc
```

SEQ ID NO:67

Amino Acid Sequence of ca B/Jilin/20/03 HA

Entire molecule length: 584 aa

```
1 mkaiivllmv vtsnadriect gitssnsphv vktatqgevn vtgviplttt
51 ptksyfanlk gtrtrgklcp dclnctdl dv algrpmcvgt tpsakasilh
101 evrpvtsgcf pimhdrtkir qlpnllrgye nirlstqnv daenapggpy
151 rlgtsghscpn atsksgffat mawavpkdnn knatnpl tve vpyvcteged
201 qitvwghsd nktpmknlyg dsnpqkftss angvtthyvs qiggfpaqte
251 deglpqsgri vvdymvqkpr ktgtivyqrg vllpqkwca sgrskvikgs
301 lpligeadcl hekyggl nks kpyytgehak aigncpiwvk tplklangtk
351 yrppakllke rgffgaiagf leggwegmia gwhgytshga hgvavaadlk
401 stqeainkit knlnslsele vknqlrlsga mdelhneile ldekvdldra
451 dtissqiela vllsnegiin sedehllale rklkklm lgs avdigngcfe
501 tkhkcnqtcl driaagtfna gefslptfds lnitaaslnd dgldnhtill
551 yystaassla vtlmlaifiv ymvsrdnvsc sicl
```

Figure 1HH**SEQ ID NO:34**

Nucleotide Sequence of ca B/Jilin/20/03 NA Entire molecule length: 1529 bp

```
1  tctcaaaaact  gaggc aaaata  ggccaaaaat  gaacaatgct  accctcaact
51  atacaaacgt  taaccctatt  cctcacatca  gggggagtg  tattatcact
101 atatgtgtca  gcttcaactg  catacttact  atattcggat  atattgctaa
151 aatttttcaac  aacagaaaata  actgcaccaa  caatgccatt  ggattgtgca
201 aacgcatcaa  atgttcaggg  tgtgaaccgt  tctgcaacaa  aaggggtgac
251 acttcttctc  ccagaaccgg  agtggacata  cccgcgttta  tcttgcccg
301 gctcaacctt  tcagaaagca  ctcctaatta  gccctcatag  attcggagaa
351 accaaaggaa  actcagctcc  cttgataata  agggaacctt  ttattgcttg
401 tggaccaaaag  gaatgcaaac  actttgctct  aaccattat  gcagcccaac
451 cagggggata  ctacaatgga  acaaaagaag  acagaaacaa  gctgaggcat
501 ctaatttcag  tcaaattggg  caaatccca  acagtagaaa  actccatttt
551 ccacatggca  gcatggagcg  ggtccgcatg  ccatgatgg  aaagaatgga
601 catatatcgg  agttgatggc  cctgacagta  atgcattgct  caaaataaaa
651 tatggagaag  catatactga  cacataccat  tcctatgcaa  acaacatcct
701 aagaacacaa  gaaagtgcct  gcaattgcat  cgggggaaat  tgttatctta
751 tgataactga  tggctcagct  tcaggtatta  gtgagtgcag  atttcttaag
801 attcgagagg  gccgaataat  aaaagaaata  tttccaacag  gaagagtaaa
851 acatactgaa  gaatgcacat  gcggtttgc  cagcaataaa  accatagaat
901 gtgcctgtag  agataacagt  tacacagcaa  aaagaccctt  tgtcaaatta
951 aatgtggaga  ctgatacagc  agaaataaga  ttgatgtgca  cagagactta
1001 tttggacacc  cccgaccag  atgatggaag  cataacagg  ccttgtgaat
1051 ctaatgggaa  taaaggaggt  ggaggcatca  agggaggatt  tgttcatcaa
1101 agaatggcat  ccaaaattgg  aagggtgtac  tctcgaacaa  tgtctaaaac
1151 caaaaggatg  ggaatgggac  tgtatgtcaa  gtatgatgga  gacccatgga
1201 ctgacagtga  tgcccttgct  cttagtggag  taatggtttc  aatggaagaa
1251 cctggttggt  actcatttgg  cttcgaaata  aaagataaga  aatgtgatgt
1301 cccctgtatt  gggatagaga  tggtacatga  tgggtgaaa  gagacttggc
1351 actcagcagc  aacagccatt  tactgtttaa  tgggctcagg  acaactgttg
1401 tgggacactg  tcacaggtgt  tgatatggct  ctgtaatggg  ggaatggttg
1451 agtctgttct  aaaccctttg  ttctatttt  gtttgaacaa  ttgtccttgc
1501 tgaacttaat  tgtttctgaa  aaatgctct
```

SEQ ID NO:68

Amino Acid Sequence of ca B/Jilin/20/03 NA

Entire molecule length: 466 aa

```
1  mlpstiqtl  lfltsggvll  slyvsaslsy  llysdillkf  stteitaptm
51  pldcanasv  qavnrsatk  vtlllpepew  typrlscpgs  tfqkallisp
101 hrfgetkgns  apliirepfi  acgpkeckhf  althyaap  ggyngtkedr
151 nkrlrhlsvk  lgkiptvens  ifhmaawsgs  achdgkewty  igvdgpdnsa
201 llkikygeay  tdyhsyann  ilrtqesacn  ciggnclmi  tdgsasgise
251 crflkiregr  iikeifptgr  vkhteectcg  fasnktieca  crdnsytakr
301 pfvklinvetd  taeirlmcte  tyltprpdd  gsitgpcesn  gnkgsggikg
351 gfvhqrmask  igrwysrtms  ktkrmgmgly  vkydgdptd  sdalalsgvm
401 vsmeepgwys  fgfeikdkkc  dvpcigiemv  hdggketwhs  aataiyclmg
451 sgqllwdtvt  gvdmal
```

Figure 2

SEQ ID NO	HA or NA	Strain Name
SEQ ID NO:1	HA	ca A/Shandong/9/93
SEQ ID NO:2	NA	ca A/Shandong/9/93
SEQ ID NO:3	HA	ca A/Johannesburg/33/94-Like
SEQ ID NO:4	NA	ca A/Johannesburg/33/94-Like
SEQ ID NO:5	HA (H3)	ca A/Wuhan/395/95
SEQ ID NO:6	NA (N2)	ca A/Wuhan/395/95
SEQ ID NO:7	HA (H3)	ca A/Sydney/05/97
SEQ ID NO:8	NA (N2)	ca A/Sydney/05/97
SEQ ID NO:9	HA (H3)	ca A/Panama/2007/99
SEQ ID NO:10	NA (N2)	ca A/Panama/2007/99
SEQ ID NO:11	HA (H3)	ca A/Wyoming/03/2003
SEQ ID NO:12	NA (N2)	ca A/Wyoming/03/2003
SEQ ID NO:13	HA (H1)	ca A/Texas/36/91
SEQ ID NO:14	NA (N1)	ca A/Texas/36/91
SEQ ID NO:15	HA (H1)	ca A/Shenzhen/227/95
SEQ ID NO:16	NA (N1)	ca A/Shenzhen/227/95
SEQ ID NO:17	HA (H1)	ca A/Beijing/262/95
SEQ ID NO:18	NA (N1)	ca A/Beijing/262/95
SEQ ID NO:19	HA (H1)	ca A/New Caledonia/20/99
SEQ ID NO:20	NA (N1)	ca A/New Caledonia/20/99
SEQ ID NO:21	HA	ca B/Ann Arbor/1/94
SEQ ID NO:22	NA	ca B/Ann Arbor/1/94
SEQ ID NO:23	HA	ca B/Yamanashi/166/98
SEQ ID NO:24	NA	ca B/Yamanashi/166/98
SEQ ID NO:25	HA	ca B/Johannesburg/5/99
SEQ ID NO:26	NA	ca B/Johannesburg/5/99
SEQ ID NO:27	HA	ca B/Victoria/504/2000
SEQ ID NO:28	NA	ca B/Victoria/504/2000
SEQ ID NO:29	HA	ca B/Hong Kong/330/01
SEQ ID NO:30	NA	ca B/Hong Kong/330/01
SEQ ID NO:31	HA	ca B/Brisbane/32/2002
SEQ ID NO:32	NA	ca B/Brisbane/32/2002
SEQ ID NO:33	HA	ca B/Jilin/20/2003
SEQ ID NO:34	NA	ca B/Jilin/20/2003

INFLUENZA HEMAGGLUTININ AND NEURAMINIDASE VARIANTS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a division and claims the benefit under 35 U.S.C. § 120 of U.S. patent application Ser. No. 10/870,690, filed Jun. 16, 2004, which claims the benefit under 35 U.S.C. § 119(e) of the U.S. Provisional Patent Application Ser. No. 60/479,078, filed on Jun. 16, 2003, the disclosures of each of which are incorporated herein in their entirety for all purposes.

BACKGROUND OF THE INVENTION

[0002] Vaccines against various and evolving strains of influenza are important from a community health standpoint, as well as commercially, since each year numerous individuals are infected with different strains and types of influenza virus. Infants, the elderly, those without adequate health care and immuno-compromised persons are at special risk of death from such infections. Compounding the problem of influenza infections is that novel influenza strains evolve readily and can spread between various species, thereby necessitating the continuous production of new vaccines.

[0003] Numerous vaccines capable of producing a protective immune response specific for different influenza viruses/virus strains have been produced for over 50 years and include whole virus vaccines, split virus vaccines, surface antigen vaccines and live attenuated virus vaccines. However, while appropriate formulations of any of these vaccine types are capable of producing a systemic immune response, live attenuated virus vaccines have the advantage of also being able to stimulate local mucosal immunity in the respiratory tract. Considerable work in the production of influenza viruses, and fragments thereof, for production of vaccines has been done by the present inventors and co-workers; see, e.g., U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus."

[0004] Because of the continual emergence (or re-emergence) or different influenza strains, new influenza vaccines are continually desired. Such vaccines typically are created using antigenic moieties of the newly emergent virus strains so, therefore, polypeptides and polynucleotides of novel, newly emergent, or newly re-emergent virus strains (especially sequences of antigenic genes) are highly desirable. Furthermore, such sequences within preferred vectors are also quite highly desired.

[0005] The present invention provides new and/or newly isolated influenza hemagglutinin and neuraminidase variants, optionally within preferred vectors, that are capable of use in production of numerous types of vaccines as well as in research, diagnostics, etc. Numerous other benefits will become apparent upon review of the following

SUMMARY OF THE INVENTION

[0006] In some aspects herein, the invention comprises an isolated or recombinant polypeptide that is selected from: the polypeptides encoded by any one of the sequences of the sequence listing, e.g., SEQ ID NO:1 through SEQ ID NO:34, any one of the polypeptides encoded by the sequence listing,

e.g., SEQ ID NO:35 through SEQ ID NO:68; any polypeptide that is encoded by a polynucleotide sequence which hybridizes under highly stringent conditions over substantially the entire length of a polynucleotide sequence of the sequence listing; and, a fragment of any of the above wherein the sequence comprises a hemagglutinin or neuraminidase polypeptide, or a fragment thereof. In various embodiments, the isolated or recombinant polypeptides of the invention are substantially identical to about 300 contiguous amino acid residues of any of the above polypeptides. In yet other embodiments, the invention comprises isolated or recombinant polypeptides (comprising hemagglutinin or fragments thereof), that comprise an amino acid sequence that is substantially identical over at least about 350 amino acids; over at least about 400 amino acids; over at least about 450 amino acids; over at least about 500 amino acids; over at least about 502 amino acids; over at least about 550 amino acids; over at least about 559 amino acids; over at least about 565 amino acids; or over at least about 566 amino acids contiguous of any of the polypeptides of claim of any of the above polypeptides. In yet other embodiments, the invention comprises isolated or recombinant polypeptides (e.g., comprising neuraminidase or fragments thereof), that comprise an amino acid sequence that is substantially identical over at least about 350 amino acids; over at least about 400 amino acids; over at least about 436 amino acids; over at least about 450 amino acids; over at least about 451 amino acids; over at least about 465 amino acids; over at least about 466 amino acids; over at least about 469 amino acids; or over at least about 470 amino acids contiguous of any of the polypeptides of any of the above polypeptides. Of course, in some embodiments, the polypeptide sequence (e.g., as listed in the sequence listing herein, e.g., SEQ ID NO:35 through SEQ ID NO:68) comprises less than 565, 559, etc. amino acids. In such embodiments, the shorter listed polypeptides optionally comprise less than 565, 559, etc. amino acids. In yet other embodiments, the polypeptides of the invention optionally comprise fusion proteins, proteins with a leader sequence, a precursor polypeptide, proteins with a secretion signal or a localization signal, or proteins with an epitope tag, an E-tag, or a H is epitope tag, etc. In still other embodiments, the invention comprises a polypeptide comprising a sequence having at least 95%, at least 96%, at least 97%, at least 98%, at least 98.5%, at least 99%, at least 99.2%, at least 99.4%, at least 99.6%, at least 99.8%, or at least 99.9% sequence identity to at least one polypeptide listed above. In some embodiments, such polypeptides are immunogenic.

[0007] In other aspects, the invention comprises a composition with one or more polypeptide listed above, or fragments thereof. The invention also includes polypeptides that are specifically bound by a polyclonal antisera raised against at least 1 antigen that comprises at least one amino acid sequence described above, or a fragment thereof. Such antibodies specific for the polypeptides described above are also features of the invention. The polypeptides of the invention are optionally immunogenic.

[0008] The invention also encompasses immunogenic compositions comprising an immunologically effective amount of one or more of any of the polypeptides described above as well as methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus by administering to the individual an immunologically effective amount of any of the above polypeptides in a physiologically acceptable carrier.

[0009] Additionally, the invention has reassortant influenza virus that encode one or more of the polypeptides above, in addition to immunogenic compositions comprising an immunologically effective amount of such recombinant influenza virus. Methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus, through administering an immunologically effective amount of such recombinant influenza virus in a physiologically acceptable carrier are also part of the invention. Such virus can optionally comprise a 6:2 reassortant virus with 6 genes encoding regions from one or more donor virus (e.g., A/AA/6/60, B/Ann Arbor/1/66, or A/Puerto Rico/8/34, which is more commonly known as PR8) and 2 gene encoding regions (typically and preferably encoding HA and NA or fragments thereof) selected from SEQ ID NO:1 through SEQ ID NO:34 or from similar strains, as defined herein, to those having SEQ ID NO:1-34, etc. Immunogenic compositions comprising such reassortant (recombinant) virus are also features of the invention.

[0010] In other aspects, the invention comprises an isolated or recombinant nucleic acid that is selected from: any one of the polynucleotide sequences of the sequence listing, e.g., SEQ ID NO:1 through SEQ ID NO:34 (or complementary sequences thereof), any one of the polynucleotide sequences encoding a polypeptide of the sequence listing, e.g., SEQ ID NO:35 through SEQ ID NO:68 (or complementary polynucleotide sequences thereof), a polynucleotide sequence which hybridizes under highly stringent conditions over substantially the entire length of any of the above polynucleotide sequences, and a polynucleotide sequence comprising all or a fragment of any of the above polynucleotide sequences wherein the sequence encodes a hemagglutinin or neuraminidase polypeptide or a fragment thereof. Such nucleic acids can be DNA, RNA, cRNA, DNA:RNA hybrids, single stranded nucleic acid, double stranded nucleic acid, etc. The invention also includes an isolated or recombinant nucleic acid (e.g., comprising hemagglutinin or fragments thereof), that encodes an amino acid sequence which is substantially identical over at least about 300 amino acids of any of the above nucleic acids, or over at least about 350 amino acids; over at least about 400 amino acids; over at least about 450 amino acids; over at least about 500 amino acids; over at least about 502 amino acids; over at least about 550 amino acids; over at least about 559 amino acids; over at least about 565 amino acids; or over at least about 566 amino acids of any of the above nucleic acids. In yet other embodiments, the invention comprises isolated or recombinant nucleic acids (e.g., comprising neuraminidase or fragments thereof), that encode an amino acid sequence that is substantially identical over at least about 350 amino acids; over at least about 400 amino acids; over at least about 436 amino acids; over at least about 450 amino acids; over at least about 451 amino acids; over at least about 465 amino acids; over at least about 466 amino acids; over at least about 469 amino acids; or over at least about 470 amino acids contiguous of any of the polypeptides above. Again, in situations wherein the amino acid is less than, e.g., 566, 565, 559, etc. in length (e.g., see, Sequence Listing in FIG. 1) then it should be understood that the length is optionally less than 566, 565, 559, etc. The invention also includes any of the above nucleic acids that comprise a hemagglutinin or neuraminidase polypeptide, or fragment thereof. Other aspects of the invention include isolated or recombinant nucleic acids that encode a polypeptide (optionally a hemagglutinin or neuraminidase polypeptide) whose

sequence has at least 95% identity, at least 96% identity, at least 97% identity, at least 98% identity, at least 98.5% identity, at least 99% identity, at least 99.2% identity, at least 99.4% identity, at least 99.6% identity, at least 99.8% identity, or at least 99.9% identity to at least one of the above described polynucleotide. The invention also includes isolated or recombinant nucleic acids encoding a polypeptide of hemagglutinin or neuraminidase produced by mutating or recombining one or more above described polynucleotide sequence. The polynucleotide sequences of the invention can optionally comprise one or more of, e.g., a leader sequence, a precursor sequence, or an epitope tag sequence or the like, and can optionally encode a fusion protein (e.g., with one or more additional nucleic acid sequences). Such nucleic acids of the invention can optionally encode immunogenic polypeptides.

[0011] In yet other embodiments, the invention comprises a composition of matter having two or more above described nucleic acids or fragments thereof (e.g., a library comprising at least about 2, 5, 10, 50 or more nucleic acids). Such compositions can optionally be produced by cleaving one or more above described nucleic acid (e.g., mechanically, chemically, enzymatically with a restriction endonuclease/RNase/DNAse, etc.). Other compositions of the invention include, e.g., compositions produced by incubating one or more above described nucleic acid in the presence of deoxyribonucleotide triphosphates and a thermostable nucleic acid polymerase. Immunogenic compositions having an immunologically effective amount of any of the above nucleic acids are also within the current invention.

[0012] Also within the invention are reassortant influenza virus comprising any of the above nucleic acids. Such reassortant viruses can (and preferably are) 6:2 reassortant viruses with 6 gene encoding regions from one or more donor virus (e.g., A/AA/6/60, B/AA/1/66 (also sometimes referred to herein as B/Ann Arbor/1/66, or A/Puerto Rico/8/34) and 2 gene encoding regions from two sequences above (e.g., from SEQ ID NO:1-34, from similar strains to those encoded in SEQ ID NO:1-34, etc.). Preferably, such two regions encode hemagglutinin and/or neuraminidase. Immunogenic compositions with immunologically effective amounts of such reassortant/recombinant influenza virus are also within purview of the current invention.

[0013] Vectors comprising one or more nucleic acid from SEQ ID NO:1-34 (again, also from similar strains to those of the sequence identification numbers) or fragments thereof are also within the current invention. Such vectors (e.g., expression vectors) can optionally be plasmids, cosmids, phage, viruses, virus fragments, etc. Especially preferred embodiments comprise plasmid vectors useful in plasmid rescue methods to produce virus (e.g., typically reassortant/recombinant virus for use in vaccines). Such plasmid systems are exemplified in, e.g., U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus"; Hoffmann, E., 2000, *PNAS*, 97(11): 6108-6113; U.S. Published Patent Application No. 20020164770 to Hoffmann; and U.S. Pat. No. 6,544,785 issued Apr. 8, 2003 to Palese, et al. Cells transduced, transformed, transfected, etc. with such vectors are also within the current invention.

[0014] The invention also encompasses cells comprising at least one above described nucleic acid, or a cleaved or amplified fragment or product thereof. Such cells can optionally

express a polypeptide encoded by such nucleic acid. Other embodiments of the invention include vectors (e.g., plasmids, cosmids, phage, viruses, virus fragments, etc.) comprising any of above described nucleic acid. Such vectors can optionally comprise an expression vector. Cells transduced by such vectors are also within the current invention.

[0015] In some embodiments, the invention encompasses a virus (e.g., an influenza virus) comprising one or more above described nucleic acid (e.g., from SEQ ID NO:1-34 or from similar strains to such and optionally encoding hemagglutinin and/or neuraminidase), or one or more fragments thereof. Typically, such viruses are reassortant/recombinant viruses. Immunogenic compositions comprising such virus are also part of the current invention. Such viruses can comprise a reassortant virus such as a 6:2 reassortment virus (which comprises 6 gene encoding regions from one or more donor virus (e.g., a master donor virus or a backbone virus such as A/AA/6/60, B/AA/1/66, A/Puerto Rico/8/34, etc.) and 2 gene encoding regions from one or more above described nucleotide sequence, or one or more fragment thereof which can optionally comprise hemagglutinin and/or neuraminidase). Other reassortant/recombinant viruses can comprise 7:1 reassortments. Reassortment viruses (optionally live viruses) of the invention can include donor viruses that are one or more of, e.g., temperature-sensitive (ts), cold-adapted (ca), or attenuated (att). For example, reassortment viruses can comprise, e.g., A/Ann Arbor/6/60, B/Ann Arbor/1/66, A/Puerto Rico/8/34, etc. In many embodiments, the produced viruses are live viruses (e.g., to be used in vaccines, etc.). Other embodiments include dead or inactivated viruses (e.g., also capable of use in vaccines, etc.). Cells comprising any of the above viruses are also products of the invention.

[0016] Methods of producing reassortant/recombinant influenza virus through culturing a host cell harboring an influenza virus in a suitable culture medium under conditions permitting expression of nucleic acid; and, isolating or recovering the recombinant influenza virus from one or more of the host cell or the medium are also part of the invention. Thus, introducing a plurality of vectors having an influenza virus genome into a population of host cells wherein the vectors comprise at least 6 internal genome segments of a first influenza strain (again, e.g., A/AA/6/60, B/AA/1/66, A/PR/8/34, etc.) and at least one (and preferably two) genome segments are selected from a second influenza strain (e.g., preferably one or more nucleic acid as described above, e.g., from SEQ ID NO:1-34 or from a similar strain to such or optionally comprising a hemagglutinin and/or neuraminidase, etc.) is a feature of the invention. Preferably, the first strain of virus is cold-adapted and/or temperature sensitive and/or attenuated. Also preferably, such viruses are suitable for administration as part of an intranasal vaccine formulation. Of course, other embodiments are suitable for administration as killed or inactivated vaccine formulations, live/attenuated nonnasal vaccine formulations, etc. The vectors in such methods can comprise influenza A viruses and/or influenza B viruses. Host cells for such methods can optionally comprise, e.g., Vero cells, PerC6 cells (ECACC deposit number 96022940), MDCK cells, 293T cells, COS cells, etc. Typical embodiments do not comprise helper viruses in the method and yet other typical embodiments comprise eight plasmid vectors to contain the influenza genome.

[0017] In other embodiments herein, the invention comprises immunogenic compositions having an immunologically effective amount of the above described recombinant

influenza virus (e.g., a live virus). Other embodiments include methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus by administering to the individual an immunologically effective amount of the recombinant influenza virus of described above (optionally in a physiologically effective carrier).

[0018] Other aspects of the invention include methods of producing an isolated or recombinant polypeptide by culturing any host cell above, in a suitable culture medium under conditions permitting expression of nucleic acid and, isolating the polypeptide from one or more of the host cell or the medium in which it is grown.

[0019] Immunogenic compositions are also features of the invention. For example, immunogenic compositions comprising one or more of the polypeptides and/or nucleic acids described above (e.g., a sequence from SEQ ID NO:1-68 or from similar strains to such, etc.) and, optionally, an excipient such as a pharmaceutically acceptable excipient or one or more pharmaceutically acceptable administration component. Immunogenic compositions of the invention can also comprise one or more above described virus as well (e.g., along with one or more pharmaceutically acceptable administration component).

[0020] Methods of producing an influenza virus vaccine are also included in the invention. For example, the invention includes introducing a plurality of vectors (e.g., plasmid vectors) comprising an influenza genome (e.g., influenza A or B) into a population of host cells that is capable of supporting replication of such virus, culturing the cells, recovering a plurality of influenza viruses and providing one or more pharmaceutically acceptable excipient with such virus to an individual (e.g., one in need of such treatment). Such viruses can optionally be cold-adapted and/or temperature sensitive and/or attenuated and preferably are suitable for administration in an intranasal vaccine formulation. Such methods can include wherein the vectors have at least 6 internal genome segments of a first influenza strain and at least one genome segment (and preferably 2 segments) from another influenza strain (e.g., with sequence selected from SEQ ID NO:1-34 or from similar strains to such, etc.) which segment optionally codes for an immunogenic influenza surface antigen of the second influenza strain.

[0021] Methods of producing immunogenic responses in a subject through administration of an effective amount of any of the above viruses to a subject are also within the current invention. Additionally, methods of prophylactic or therapeutic treatment of a viral infection (e.g., viral influenza) in a subject through administration of one or more above described virus in an amount effective to produce an immunogenic response against the viral infection are also part of the current invention. Subjects for such treatment can include mammals (e.g., humans). Such methods can also comprise in vivo administration to the subject as well as in vitro or ex vivo administration to one or more cells of the subject. Additionally, such methods can also comprise administration of a composition of the virus and a pharmaceutically acceptable excipient that is administered to the subject in an amount effect to prophylactically or therapeutically treat the viral infection.

[0022] The invention also comprises compositions of matter having one or more sequence selected from SEQ ID NO:1 through SEQ ID NO:34, and a selected master donor virus, typically wherein the selected sequence and the master donor

virus comprise a 6:2 reassortment, i.e., the HA and NA herein reassorted with the other six influenza genes from the donor virus. Such donor viruses are typically ca, att, ts influenza strains. For example, typically donor strains can include, e.g., A/Ann Arbor/6/60, B/Ann Arbor/1/66, or A/Puerto Rico/8/34 and variants thereof. Those of skill in the art will appreciate that typically donor strains can vary from reassortant to reassortant. Thus, those variations are also encompassed within the current invention. Another element of the invention comprises one or more live attenuated influenza vaccine comprising the such compositions, e.g., those having sequences herein reassorted in a 6:2 manner with a selected master donor virus.

[0023] Other aspects of the invention include, compositions of matter comprising a hemagglutinin polynucleotide and/or a neuraminidase polynucleotide reassorted with one or more master donor virus, again typically a ca, att, ts influenza virus, wherein the polynucleotide comprises a same virus strain as one or more virus strain of SEQ ID NO:1 through SEQ ID NO:34. Such hemagglutinin and/or neuraminidase polynucleotide is typically determined to be “within the same strain” when it produces a titer that is within a four-fold range of another virus (e.g., ones having the sequences listed herein) as measured by a hemagglutinin inhibition assay. As described below, however, other common assays can also be utilized to determine whether polynucleotides (i.e., viruses comprising such) are within the same strain.

[0024] These and other objects and features of the invention will become more fully apparent when the following detailed description is read in conjunction with the accompanying figures appendix.

BRIEF DESCRIPTION OF THE FIGURES

[0025] FIG. 1 displays the Sequence Listing of variant hemagglutinin and neuraminidase nucleic acids and polypeptides of the invention.

[0026] FIG. 2 displays an alternative organization of variant hemagglutinin and neuraminidase sequences as found in FIG. 1.

DETAILED DESCRIPTION

[0027] The present invention includes polypeptide and polynucleotide sequences of influenza hemagglutinin and neuraminidase as well as vectors, viruses, vaccines, compositions and the like comprising such sequences and methods of their use. Additional features of the invention are described in more detail herein.

DEFINITIONS

[0028] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. The following definitions supplement those in the art and are directed to the current application and are not necessarily to be imputed to any related or unrelated case, e.g., to any commonly owned patent or application. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred materials and methods are described herein. Accordingly, the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting. Additional terms are defined and described throughout.

[0029] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a virus” includes a plurality of viruses; reference to a “host cell” includes mixtures of host cells, and the like.

[0030] The terms “nucleic acid,” “polynucleotide,” “polynucleotide sequence,” and “nucleic acid sequence” refer to single-stranded or double-stranded deoxyribonucleotide or ribonucleotide polymers, chimeras or analogues thereof, or a character string representing such, depending on context. As used herein, the term optionally includes polymers of analogs of naturally occurring nucleotides having the essential nature of natural nucleotides in that they hybridize to single-stranded nucleic acids in a manner similar to naturally occurring nucleotides (e.g., peptide nucleic acids). Unless otherwise indicated, a particular nucleic acid sequence of this invention optionally encompasses complementary sequences in addition to the sequence explicitly indicated. From any specified polynucleotide sequence, either the given nucleic acid or the complementary polynucleotide sequence (e.g., the complementary nucleic acid) can be determined.

[0031] The term “nucleic acid” or “polynucleotide” also encompasses any physical string of monomer units that can be corresponded to a string of nucleotides, including a polymer of nucleotides (e.g., a typical DNA or RNA polymer), PNAs, modified oligonucleotides (e.g., oligonucleotides comprising bases that are not typical to biological RNA or DNA in solution, such as 2'-O-methylated oligonucleotides), and the like. A nucleic acid can be e.g., single-stranded or double-stranded.

[0032] A “subsequence” is any portion of an entire sequence, up to and including the complete sequence. Typically, a subsequence comprises less than the full-length sequence. A “unique subsequence” is a subsequence that is not found in any previously determined influenza polynucleotide or polypeptide sequence. The phrase “substantially identical,” in the context of two nucleic acids or polypeptides (e.g., DNAs encoding a HA or NA molecule, or the amino acid sequence of a HA or NA molecule) refers to two or more sequences or subsequences that have at least about 90%, preferably 91%, most preferably 92%, 93%, 94%, 95%, 96%, 97%, 98%, 98.5%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm or by visual inspection.

[0033] The term “variant” with respect to a polypeptide refers to an amino acid sequence that is altered by one or more amino acids with respect to a reference sequence. The variant can have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. Alternatively, a variant can have “nonconservative” changes, e.g., replacement of a glycine with a tryptophan. Analogous minor variation can also include amino acid deletion or insertion, or both. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without eliminating biological or immunological activity can be found using computer programs well known in the art, for example, DNASTAR software. Examples of conservative substitutions are also described herein.

[0034] The term “gene” is used broadly to refer to any nucleic acid associated with a biological function. Thus,

genes include coding sequences and/or the regulatory sequences required for their expression. The term “gene” applies to a specific genomic sequence, as well as to a cDNA or an mRNA encoded by that genomic sequence.

[0035] The “neuraminidase” polypeptides of the invention show immunological cross reactivity with one or more known neuraminidase molecule from an influenza virus. The literature is replete with examples of such known neuraminidases (e.g., in GenBank, in publications from the CDC, etc.). Similarly, the “hemagglutinin” polypeptides of the invention show immunological cross-reactivity with one or more known hemagglutinin molecule from an influenza virus. Again, the literature is replete with examples of such known hemagglutinin molecules.

[0036] Genes also include non-expressed nucleic acid segments that, for example, form recognition sequences for other proteins. Non-expressed regulatory sequences include “promoters” and “enhancers,” to which regulatory proteins such as transcription factors bind, resulting in transcription of adjacent or nearby sequences. A “tissue specific” promoter or enhancer is one that regulates transcription in a specific tissue type or cell type, or types.

[0037] “Expression of a gene” or “expression of a nucleic acid” typically means transcription of DNA into RNA (optionally including modification of the RNA, e.g., splicing) or transcription of RNA into mRNA, translation of RNA into a polypeptide (possibly including subsequent modification of the polypeptide, e.g., post-translational modification), or both transcription and translation, as indicated by the context.

[0038] An “open reading frame” or “ORF” is a possible translational reading frame of DNA or RNA (e.g., of a gene), which is capable of being translated into a polypeptide. That is, the reading frame is not interrupted by stop codons. However, it should be noted that the term ORF does not necessarily indicate that the polynucleotide is, in fact, translated into a polypeptide.

[0039] The term “vector” refers to the means by which a nucleic acid can be propagated and/or transferred between organisms, cells, or cellular components. Vectors include plasmids, viruses, bacteriophages, pro-viruses, phagemids, transposons, artificial chromosomes, and the like, that replicate autonomously or can integrate into a chromosome of a host cell. A vector can also be a naked RNA polynucleotide, a naked DNA polynucleotide, a polynucleotide composed of both DNA and RNA within the same strand, a poly-lysine-conjugated DNA or RNA, a peptide-conjugated DNA or RNA, a liposome-conjugated DNA, or the like, that is not autonomously replicating. In many, but not all, common embodiments, the vectors of the present invention are plasmids.

[0040] An “expression vector” is a vector, such as a plasmid that is capable of promoting expression, as well as replication of a nucleic acid incorporated therein. Typically, the nucleic acid to be expressed is “operably linked” to a promoter and/or enhancer, and is subject to transcription regulatory control by the promoter and/or enhancer.

[0041] A “bi-directional expression vector” is characterized by two alternative promoters oriented in the opposite direction relative to a nucleic acid situated between the two promoters, such that expression can be initiated in both orientations resulting in, e.g., transcription of both plus (+) or sense strand, and negative (−) or antisense strand RNAs.

[0042] An “amino acid sequence” is a polymer of amino acid residues (a protein, polypeptide, etc.) or a character string representing an amino acid polymer, depending on context.

[0043] A “polypeptide” is a polymer comprising two or more amino acid residues (e.g., a peptide or a protein). The polymer can optionally comprise modifications such as glycosylation or the like. The amino acid residues of the polypeptide can be natural or non-natural and can be unsubstituted, unmodified, substituted or modified.

[0044] In the context of the invention, the term “isolated” refers to a biological material, such as a virus, a nucleic acid or a protein, which is substantially free from components that normally accompany or interact with it in its naturally occurring environment. The isolated biological material optionally comprises additional material not found with the biological material in its natural environment, e.g., a cell or wild-type virus. For example, if the material is in its natural environment, such as a cell, the material can have been placed at a location in the cell (e.g., genome or genetic element) not native to such material found in that environment. For example, a naturally occurring nucleic acid (e.g., a coding sequence, a promoter, an enhancer, etc.) becomes isolated if it is introduced by non-naturally occurring means to a locus of the genome (e.g., a vector, such as a plasmid or virus vector, or amplicon) not native to that nucleic acid. Such nucleic acids are also referred to as “heterologous” nucleic acids. An isolated virus, for example, is in an environment (e.g., a cell culture system, or purified from cell culture) other than the native environment of wild-type virus (e.g., the nasopharynx of an infected individual).

[0045] The term “chimeric” or “chimera,” when referring to a virus, indicates that the virus includes genetic and/or polypeptide components derived from more than one parental viral strain or source. Similarly, the term “chimeric” or “chimera,” when referring to a viral protein, indicates that the protein includes polypeptide components (i.e., amino acid subsequences) derived from more than one parental viral strain or source. As will be apparent herein, such chimeric viruses are typically reassortant/recombinant viruses. Thus, in some embodiments, a chimera can optionally include, e.g., a sequence (e.g., of HA and/or NA) from an A influenza virus placed into a backbone comprised of, or constructed/derived from a B influenza virus (e.g., B/AA/1/66, etc.) or a B influenza virus sequence placed into an A influenza virus backbone (i.e., donor virus) such as, e.g., A/AA/6/60, etc.

[0046] The term “recombinant” indicates that the material (e.g., a nucleic acid or protein) has been artificially or synthetically (non-naturally) altered by human intervention. The alteration can be performed on the material within, or removed from, its natural environment or state. Specifically, e.g., an influenza virus is recombinant when it is produced by the expression of a recombinant nucleic acid. For example, a “recombinant nucleic acid” is one that is made by recombining nucleic acids, e.g., during cloning, DNA shuffling or other procedures, or by chemical or other mutagenesis; a “recombinant polypeptide” or “recombinant protein” is a polypeptide or protein which is produced by expression of a recombinant nucleic acid; and a “recombinant virus,” e.g., a recombinant influenza virus, is produced by the expression of a recombinant nucleic acid.

[0047] The term “reassortant,” when referring to a virus (typically herein, an influenza virus), indicates that the virus includes genetic and/or polypeptide components derived

from more than one parental viral strain or source. For example, a 7:1 reassortant includes 7 viral genomic segments (or gene segments) derived from a first parental virus, and a single complementary viral genomic segment, e.g., encoding a hemagglutinin or neuraminidase such as those listed in the SEQ ID Table herein. A 6:2 reassortant includes 6 genomic segments, most commonly the 6 internal genes from a first parental virus, and two complementary segments, e.g., hemagglutinin and neuraminidase, from one or more different parental virus. Reassortant viruses can also, depending upon context herein, be termed as “chimeric” and/or “recombinant.”

[0048] The term “introduced” when referring to a heterologous or isolated nucleic acid refers to the incorporation of a nucleic acid into a eukaryotic or prokaryotic cell where the nucleic acid can be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA). The term includes such methods as “infection,” “transfection,” “transformation,” and “transduction.” In the context of the invention a variety of methods can be employed to introduce nucleic acids into cells, including electroporation, calcium phosphate precipitation, lipid mediated transfection (lipofection), etc.

[0049] The term “host cell” means a cell that contains a heterologous nucleic acid, such as a vector or a virus, and supports the replication and/or expression of the nucleic acid. Host cells can be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, avian or mammalian cells, including human cells. Exemplary host cells can include, e.g., Vero (African green monkey kidney) cells, BHK (baby hamster kidney) cells, primary chick kidney (PCK) cells, Madin-Darby Canine Kidney (MDCK) cells, Madin-Darby Bovine Kidney (MDBK) cells, 293 cells (e.g., 293T cells), and COS cells (e.g., COS1, COS7 cells), etc. In other embodiments, host cells can optionally include eggs (e.g., hen eggs, embryonated hen eggs, etc.).

[0050] An “immunologically effective amount” of influenza virus is an amount sufficient to enhance an individual’s (e.g., a human’s) own immune response against a subsequent exposure to influenza virus. Levels of induced immunity can be monitored, e.g., by measuring amounts of neutralizing secretory and/or serum antibodies, e.g., by plaque neutralization, complement fixation, enzyme-linked immunosorbent, or microneutralization assay.

[0051] A “protective immune response” against influenza virus refers to an immune response exhibited by an individual (e.g., a human) that is protective against disease when the individual is subsequently exposed to and/or infected with wild-type influenza virus. In some instances, the wild-type (e.g., naturally circulating) influenza virus can still cause infection, but it cannot cause a serious infection. Typically, the protective immune response results in detectable levels of host engendered serum and secretory antibodies that are capable of neutralizing virus of the same strain and/or subgroup (and possibly also of a different, non-vaccine strain and/or subgroup) in vitro and in vivo.

[0052] As used herein, an “antibody” is a protein comprising one or more polypeptides substantially or partially encoded by immunoglobulin genes or fragments of immunoglobulin genes. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as myriad immunoglobulin variable region genes. Light chains are classified as either

kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. A typical immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kD) and one “heavy” chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these light and heavy chains respectively. Antibodies exist as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce F(ab)₂, a dimer of Fab which itself is a light chain joined to VH-CH1 by a disulfide bond. The F(ab)₂ may be reduced under mild conditions to break the disulfide linkage in the hinge region thereby converting the (Fab)₂ dimer into a Fab’ monomer. The Fab’ monomer is essentially a Fab with part of the hinge region (see Fundamental Immunology, W. E. Paul, ed., Raven Press, N.Y. (1999) for a more detailed description of other antibody fragments). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such Fab’ fragments may be synthesized de novo either chemically or by utilizing recombinant DNA methodology. Thus, the term antibody, as used herein, includes antibodies or fragments either produced by the modification of whole antibodies or synthesized de novo using recombinant DNA methodologies. Antibodies include, e.g., polyclonal antibodies, monoclonal antibodies, multiple or single chain antibodies, including single chain Fv (sFv or scFv) antibodies in which a variable heavy and a variable light chain are joined together (directly or through a peptide linker) to form a continuous polypeptide, and humanized or chimeric antibodies.

Influenza Virus

[0053] The polypeptides and polynucleotides of the invention are variants of influenza HA and NA sequences. See, e.g., the Sequence Listing in FIG. 1 below. In general, influenza viruses are made up of an internal ribonucleoprotein core containing a segmented single-stranded RNA genome and an outer lipoprotein envelope lined by a matrix protein. The genome of influenza viruses is composed of eight segments of linear (–) strand ribonucleic acid (RNA), encoding the immunogenic hemagglutinin (HA) and neuraminidase (NA) proteins, and six internal core polypeptides: the nucleocapsid nucleoprotein (NP); matrix proteins (M); non-structural proteins (NS); and 3 RNA polymerase (PA, PB1, PB2) proteins. During replication, the genomic viral RNA is transcribed into (+) strand messenger RNA and (–) strand genomic cRNA in the nucleus of the host cell. Each of the eight genomic segments is packaged into ribonucleoprotein complexes that contain, in addition to the RNA, NP and a polymerase complex (PB1, PB2, and PA). The hemagglutinin molecule consists of a surface glycoprotein and acts to bind to N-Acetyl-Neuraminic acid (NeuNAc), also known as sialic acid, on host cell surface receptors. In some embodiments herein, the polypeptides of the invention (and polypeptides encoded by the polynucleotides of the invention) can act to bind NeuNAc whether in vitro or in vivo. Such action can in some embodiments also be done by fragments of hemagglutinin which retain hemagglutinin activity. Hemagglutinin is made up of

two subunits, HA1 and HA2 and the entire structure is about 550 amino acids in length and about 220 kD. Neuraminidase molecules cleave terminal sialic acid residues from cell surface receptors of influenza virus, thereby releasing virions from infected cells. neuraminidase also removes sialic acid from newly made hemagglutinin and neuraminidase molecules. In some embodiments herein, the polypeptides of the invention (and polypeptides encoded by the polynucleotides of the invention) can act to cleave sialic acid residues whether in vitro or in vivo. This action can also be done in some embodiments by fragments of neuraminidase which retain neuraminidase activity. The neuraminidase polypeptides of the invention show immunological cross reactivity with one or more known neuraminidase molecule from an influenza virus. The literature is replete with examples of such known neuraminidases (e.g., in GenBank, in publications from the CDC, etc.). Similarly, the hemagglutinin polypeptides of the invention show immunological cross-reactivity with one or more known hemagglutinin molecule from an influenza virus. Again, the literature is replete with examples of such known hemagglutinin molecules.

[0054] Influenza is commonly grouped into influenza A and influenza B categories, as well as a typically less important C category. Influenza A and influenza B viruses each contain eight segments of single stranded RNA with negative polarity. The influenza A genome encodes eleven polypeptides. Segments 1-3 encode three polypeptides, making up a RNA-dependent RNA polymerase. Segment 1 encodes the polymerase complex protein PB2. The remaining polymerase proteins PB1 and PA are encoded by segment 2 and segment 3, respectively. In addition, segment 1 of some influenza strains encodes a small protein, PB1-F2, produced from an alternative reading frame within the PB1 coding region. Segment 4 encodes the hemagglutinin (HA) surface glycoprotein involved in cell attachment and entry during infection. Segment 5 encodes the nucleocapsid nucleoprotein (NP) polypeptide, the major structural component associated with viral RNA. Segment 6 encodes a neuraminidase (NA) envelope glycoprotein. Segment 7 encodes two matrix proteins, designated M1 and M2, which are translated from differentially spliced mRNAs. Segment 8 encodes NS1 and NS2, two nonstructural proteins, which are translated from alternatively spliced mRNA variants. The eight genome segments of influenza B encode II proteins. The three largest genes code for components of the RNA polymerase, PB1, PB2 and PA. Segment 4 encodes the HA protein. Segment 5 encodes NP. Segment 6 encodes the NA protein and the NB protein. Both proteins, NB and NA, are translated from overlapping reading frames of a bicistronic mRNA. Segment 7 of influenza B also encodes two proteins: M1 and BM2. The smallest segment encodes two products: NS1 is translated from the full length RNA, while NS2 is translated from a spliced mRNA variant.

[0055] Influenza types A and B are typically associated with influenza outbreaks in human populations. However, type A influenza also infects other creatures as well, e.g., birds, pigs, and other animals. The type A viruses are categorized into subtypes based upon differences within their hemagglutinin and neuraminidase surface glycoprotein antigens. Hemagglutinin in type A viruses has 14 known subtypes and neuraminidase has 9 known subtypes. In humans, currently only about 3 different hemagglutinin and 2 different neuraminidase subtypes are known, e.g., H1, H2, H3, N1, and N2. In particular, two major subtypes of influenza A have been active in humans, namely, H1N1 and H3N2. H1N2,

however has recently been of concern. Influenza B viruses are not divided into subtypes based upon their hemagglutinin and neuraminidase proteins. As will be appreciated, the sequences contained within the sequence listing in FIG. 1 comprise a number of different subtypes of influenza. Thus, for example in the sequence listing A-H3N2 strains are exemplified by ca A/Shandong/9/93, ca A/Johannesburg/33/94-like, ca A/Wuhan/395/95, ca A/Sydney/05/97, ca A/Panama/2007/99, ca A/Wyoming/03/2003. A-H1N1 strains are shown in ca A/Texas/36/91, ca A/Shenzhen/227/95, ca A/Beijing/262/95, and ca A/New Caledonia/20/99, while B-HANA strains include ca B/Ann Arbor/1/94, ca B/Yamanashi/166/98, ca B/Johannesburg/5/99, ca B/Victoria/504/2000, ca B/Hong Kong/330/2001, ca B/Brisbane/32/2002, and ca B/Jilin/20/2003.

[0056] Different strains of influenza can be categorized based upon, e.g., the ability of influenza to agglutinate red blood cells (RBCs or erythrocytes). Antibodies specific for particular influenza strains can bind to the virus and, thus, prevent such agglutination. Assays determining strain types based on such inhibition are typically known as hemagglutinin inhibition assays (HI assays or HAI assays) and are standard and well known methods in the art to characterize influenza strains. Of course, those of skill in the art will be familiar with other assays, e.g., ELISA, indirect fluorescent antibody assays, immunohistochemistry, Western blot assays, etc. with which to characterize influenza strains and the use of and discussion herein of HI assays should not be necessarily construed as limiting.

[0057] Briefly, in typical HI assays, sera to be used for typing or categorization, which is often produced in ferrets, is added to erythrocyte samples in various dilutions, e.g., 2-fold, etc. Optical determination is then made whether the erythrocytes are clumped together (i.e., agglutinated) or are suspended (i.e., non-agglutinated). If the cells are not clumped, then agglutination did not occur due to the inhibition from antibodies in the sera that are specific for that influenza. Thus, the types of influenza are defined as being within the same strain. In some cases, one strain is described as being "like" the other, e.g., strain x is a "y-like" strain, etc. For example, if two samples are within four-fold titer of one another as measured by an HI assay, then they can be described as belonging to the same strain (e.g., both belonging to the "New Caledonia" strain or both being "Moscow-like" strains, etc.). In other words, strains are typically categorized based upon their immunologic or antigenic profile. An HAI titer is typically defined as the highest dilution of a serum that completely inhibits hemagglutination. See, e.g., Schild, et al., *Bull. Wld Hlth Org.*, 1973, 48:269-278, etc. Again, those of skill in the art will be quite familiar with categorization and classification of influenza into strains and the methods to do so.

[0058] From the above it will be appreciated that the current invention not only comprises the specific sequences listed herein, but also such sequences within various vectors (e.g., ones used for plasmid reassortment and rescue, see below) as well as hemagglutinin and neuraminidase sequences within the same strains as the sequences listed herein. Also, such same strains that are within various vectors (e.g., typically ones used for plasmid reassortment and rescue such as A/Ann Arbor/6/60 or B/Ann Arbor/1/66, A/Puerto Rico/8/34, etc.) are also included.

[0059] As used herein, the term "similar strain" should be taken to indicate that a first influenza virus is of the same or related strain as a second influenza virus. In typical embodi-

ments such relation is commonly determined through use of an HAI assay. Influenza viruses that fall within a four-fold titer of one another in an HAI assay are, thus, of a "similar strain." Those of skill in the art, however, will be familiar with other assays, etc. to determine similar strains, e.g., FRID, neutralization assays, etc. The current invention also comprises such similar strains (i.e., strains similar to the ones present in the sequence listing herein) in the various plasmids, vectors, viruses, methods, etc. herein. Thus, unless the context clearly dictates otherwise, descriptions herein of particular sequences (e.g., those in the sequence listing) or fragments thereof also should be considered to include sequences from similar strains to those (i.e., similar strains to those strains having the sequences in those plasmids, vectors, viruses, etc. herein). Also, it will be appreciated that the NA and HA polypeptides within such similar strains are, thus, "similar polypeptides" when compared between "similar strains."

Influenza Virus Vaccines

[0060] The sequences, compositions and methods herein are primarily, but not solely, concerned with production of influenza viruses for vaccines. Historically, influenza virus vaccines have primarily been produced in embryonated hen eggs using strains of virus selected or based on empirical predictions of relevant strains. More recently, reassortant viruses have been produced that incorporate selected hemagglutinin and/or neuraminidase antigens in the context of an approved attenuated, temperature sensitive master strain. Following culture of the virus through multiple passages in hen eggs, influenza viruses are recovered and, optionally, inactivated, e.g., using formaldehyde and/or β -propiolactone (or alternatively used in live attenuated vaccines). Thus, it will be appreciated that HA and NA sequences (as in the current invention) are quite useful in constructing influenza vaccines.

[0061] Attempts at producing recombinant and reassortant vaccines in cell culture have been hampered by the inability of some of the strains approved for vaccine production to grow efficiently under standard cell culture conditions. However, prior work by the inventors and their coworkers provided a vector system, and methods for producing recombinant and reassortant viruses in culture, thus, making it possible to rapidly produce vaccines corresponding to one or many selected antigenic strains of virus, e.g., either A or B strains, various subtypes or substrains, etc., e.g., comprising the HA and NA sequences herein. See, U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus." Typically, the cultures are maintained in a system, such as a cell culture incubator, under controlled humidity and CO₂, at constant temperature using a temperature regulator, such as a thermostat to insure that the temperature does not exceed 35° C. Reassortant influenza viruses can be readily obtained by introducing a subset of vectors corresponding to genomic segments of a master influenza virus, in combination with complementary segments derived from strains of interest (e.g., HA and NA antigenic variants herein). Typically, the master strains are selected on the basis of desirable properties relevant to vaccine administration. For example, for vaccine production, e.g., for production of a live attenuated vaccine, the master donor virus strain may be selected for an attenuated phenotype, cold adaptation and/or temperature sensitivity. As explained elsewhere herein and, e.g., in U.S. patent applica-

tion Ser. No. 10/423,828, etc., various embodiments of the invention utilize A/Ann Arbor (AA)/6/60 or B/Ann Arbor/1/66 or A/Puerto Rico/8/34 influenza strain as a "backbone" upon which to add HA and/or NA genes (e.g., such as those sequences listed herein, etc.) to create desired reassortant viruses. Thus, for example, in a 6:2 reassortant, 2 genes (i.e., NA and HA) would be from the influenza strain(s) against which an immunogenic reaction is desired, while the other 6 genes would be from the Ann Arbor strain, or other backbone strain, etc. The Ann Arbor virus is useful for its cold adapted, attenuated, temperature sensitive attributes. Of course, it will be appreciated that the HA and NA sequences herein are capable of reassortment with a number of other virus genes or virus types (e.g., a number of different "backbones" such as A/Puerto Rico/8/34, etc., containing the other influenza genes present in a reassortant, namely, the non-HA and non-NA genes). Live, attenuated influenza A virus vaccines against human influenza viruses were recently licensed in the United States. See above. Such vaccines are reassortant H1N1 and H1N2 viruses in which the internal protein genes of A/Ann Arbor (AA)/6/60 (H2N2) cold adapted (ca) virus confer the cold adapted, attenuation and temperature sensitive phenotypes of the AA ca virus on the reassortant viruses (i.e., the ones having the hemagglutinin and neuraminidase genes from the non-Ann Arbor strain). In some embodiments herein, the reassortants can also comprise 7:1 reassortants. In other words, only the HA or the NA is not from the backbone or MDV strain. Previous work has been reported with suitable backbone donor virus strains that optionally are within various embodiments of the current invention. See, e.g., U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus"; Maassab et al., *J. of Inf. Dis.*, 1982, 146:780-790; Cox, et al., *Virology*, 1988, 167:554-567; Wareing et al., *Vaccine*, 2001, 19:3320-3330; Clements, et al., *J Infect Dis.* 1990, 161(5):869-77, etc.,

[0062] In some embodiments, the sequences herein can optionally have specific regions removed (both or either in the nucleic acid sequence or the amino acid sequence). For example, for those molecules having a polybasic cleavage site, such sites can optionally be removed. Those of skill in the art will be familiar with various methods of removing such specific regions. The resulting shortened sequences are also contained within the current invention. See, e.g., Li et al., *J. of Infectious Diseases*, 179:1132-8, 1999

[0063] The terms "temperature sensitive," "cold adapted" and "attenuated" as applied to viruses (typically used as vaccines or for vaccine production) which optionally encompass the current sequences, are well known in the art. For example, the term "temperature sensitive" (ts) indicates, e.g., that the virus exhibits a 100 fold or greater reduction in titer at 39° C. relative to 33° C. for influenza A strains, or that the virus exhibits a 100 fold or greater reduction in titer at 37° C. relative to 33° C. for influenza B strains. The term "cold adapted" (ca) indicates that the virus exhibits growth at 25° C. within 100 fold of its growth at 33° C., while the term "attenuated" (att) indicates that the virus replicates in the upper airways of ferrets but is not detectable in their lung tissues, and does not cause influenza-like illness in the animal. It will be understood that viruses with intermediate phenotypes, i.e., viruses exhibiting titer reductions less than 100 fold at 39° C. (for A strain viruses) or 37° C. (for B strain viruses), or

exhibiting growth at 25° C. that is more than 100 fold than its growth at 33° C. (e.g., within 200 fold, 500 fold, 1000 fold, 10,000 fold less), and/or exhibit reduced growth in the lungs relative to growth in the upper airways of ferrets (i.e., partially attenuated) and/or reduced influenza like illness in the animal, are also useful viruses and can be used in conjunction with the HA and NA sequences herein.

[0064] Thus, the present invention can utilize growth, e.g., in appropriate culture conditions, of virus strains (both A strain and B strain influenza viruses) with desirable properties relative to vaccine production (e.g., attenuated pathogenicity or phenotype, cold adaptation, temperature sensitivity, etc.) in vitro in cultured cells. Influenza viruses can be produced by introducing a plurality of vectors incorporating cloned viral genome segments into host cells, and culturing the cells at a temperature not exceeding 35° C. When vectors including an influenza virus genome are transfected, recombinant viruses suitable as vaccines can be recovered by standard purification procedures. Using the vector system and methods of the invention, reassortant viruses incorporating the six internal gene segments of a strain selected for its desirable properties with respect to vaccine production, and the immunogenic HA and NA segments from a selected, e.g., pathogenic strain such as those in the sequence listing herein, can be rapidly and efficiently produced in tissue culture. Thus, the system and methods described herein are useful for the rapid production in cell culture of recombinant and reassortant influenza A and B viruses, including viruses suitable for use as vaccines, including live attenuated vaccines, such as vaccines suitable for intranasal administration.

[0065] In such embodiments, typically, a single Master Donor Virus (MDV) strain is selected for each of the A and B subtypes. In the case of a live attenuated vaccine, the Master Donor Virus strain is typically chosen for its favorable properties, e.g., temperature sensitivity, cold adaptation and/or attenuation, relative to vaccine production. For example, exemplary Master Donor Strains include such temperature sensitive, attenuated and cold adapted strains of A/Ann Arbor/6/60 and B/Ann Arbor/1/66, respectively.

[0066] For example, a selected master donor type A virus (MDV-A), or master donor type B virus (MDV-B), is produced from a plurality of cloned viral cDNAs constituting the viral genome. Embodiments include those wherein recombinant viruses are produced from eight cloned viral cDNAs. Eight viral cDNAs representing either the selected MDV-A or MDV-B sequences of PB2, PB1, PA, NP, HA, NA, M and NS are optionally cloned into a bi-directional expression vector, such as a plasmid (e.g., pAD3000), such that the viral genomic RNA can be transcribed from an RNA polymerase I (pol I) promoter from one strand and the viral mRNAs can be synthesized from an RNA polymerase II (pol II) promoter from the other strand. Optionally, any gene segment can be modified, including the HA segment (e.g., to remove the multi-basic cleavage site).

[0067] Infectious recombinant MDV-A or MDV-B virus can be then recovered following transfection of plasmids bearing the eight viral cDNAs into appropriate host cells, e.g., Vero cells, co-cultured MDCK/293T or MDCK/COS7 cells. Using the plasmids and methods described herein and, e.g., in U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus"; Hoffmann, E., 2000, *PNAS*, 97(11):6108-6113; U.S.

Published Patent Application No. 20020164770 to Hoffmann; and U.S. Pat. No. 6,544,785 issued Apr. 8, 2003 to Palese, et al., the invention is useful, e.g., for generating 6:2 reassortant influenza vaccines by co-transfection of the 6 internal genes (PB1, PB2, PA, NP, M and NS) of the selected virus (e.g., MDV-A, MDV-B) together with the HA and NA derived from different corresponding type (A or B) influenza viruses e.g., as shown in the sequence listings herein. For example, the HA segment is favorably selected from a pathogenically relevant H1, H3 or B strain, as is routinely performed for vaccine production. Similarly, the NA segment can be selected from a strain with emerging relevance as a pathogenic strain such as those in the sequence listing herein. Reassortants incorporating seven genome segments of the MDV and either the HA or NA gene of a selected strain (7:1 reassortants) can also be produced. It will be appreciated, and as is detailed throughout, the molecules of the invention can optionally be combined in any desired combination. For example, the HA and/or NA sequences herein can be placed, e.g., into a reassortant backbone such as A/AA/6/60, B/AA/1/66, A/Puerto Rico/8/34 (i.e., PR8), etc., in 6:2 reassortants or 7:1 reassortants, etc. Thus, as explained more fully below, there would be 6 backbone gene regions from the donor virus (again, e.g., A/AA/6/60, etc.) and 2 genes regions from a second strain (e.g., a wild-type strain, not the backbone donor virus). Such 2 gene regions are preferably the HA and NA genes. A similar situation arises for 7:1 reassortants, in which however, there are 7 gene regions from the background donor virus and 1 gene (either HA or NA) from a different virus (typically wild-type or one to which an immune response is desired). Also, it will be appreciated that the sequences herein (e.g., those in the sequence listing of FIG. 1, etc.) can be combined in a number of means in different embodiments herein. Thus, any of the sequences herein can be present singularly in a 7:1 reassortant (i.e., the sequence of the invention present with 7 backbone donor virus gene regions) and/or can be present with another sequence of the invention in a 6:2 reassortant. Within such 6:2 reassortants, any of the sequences of the invention can optionally be present with any other sequence of the invention. Typical, and preferred, embodiments comprise HA and NA from the same original wild-type strains however. For example, typical embodiments can comprise a 6:2 reassortant having 6 gene regions from a backbone donor virus such as A/AA/6/60 and the HA and NA gene regions from the same wild-type strain such as ca A/Shandong/9/93 or both HA and NA from ca A/Wuhan/395/95 or both HA and NA from ca B/Ann Arbor/1/94 (which would typically, but not exclusively, be present within a B influenza backbone donor virus such as B/Ann Arbor/1/66, etc.), etc. Of course, it will again be appreciated that the invention also includes such reassortant viruses wherein the non-background gene regions (i.e., the HA and/or NA regions) are from similar strains (i.e., strains that are similar strains to influenza strains having the sequences found in SEQ ID NO:1-34. The above references are specifically incorporated herein in their entirety for all purposes, e.g., especially for their teachings regarding plasmids, plasmid rescue of virus (influenza virus), multi-plasmid systems for virus rescue/production, etc.

[0068] Again, the HA and NA sequences of the current invention are optionally utilized in such plasmid reassortment vaccines (and/or in other ts, cs, ca, and/or att viruses and vaccines). However, it should be noted that the HA and NA sequences, etc. of the invention are not limited to specific

vaccine compositions or production methods, and can, thus, be utilized in substantially any vaccine type or vaccine production method which utilizes strain specific HA and NA antigens (e.g., the sequences of the invention).

FluMist™

[0069] As mentioned previously, numerous examples and types of influenza vaccine exist. An exemplary influenza vaccine is FluMist™ (MedImmune Vaccines Inc., Mt. View, Calif.) which is a live, attenuated vaccine that protects children and adults from influenza illness (Belshe et al. (1998) *The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children* *N Engl J Med* 338: 1405-12; Nichol et al. (1999) *Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial* *JAMA* 282: 137-44). In typical, and preferred, embodiments, the methods and compositions of the current invention are preferably adapted to/used with production of FluMist™ vaccine. However, it will be appreciated by those skilled in the art that the sequences, methods, compositions, etc. herein are also adaptable to production of similar or even different viral vaccines.

[0070] FluMist™ vaccine strains contain, e.g., HA and NA gene segments derived from the wild-type strains to which the vaccine is addressed (or, in some instances, to related strains) along with six gene segments, PB1, PB2, PA, NP, M and NS, from a common master donor virus (MDV). The HA and NA sequences herein, thus, are optionally part of various FluMist™ formulations. The MDV for influenza A strains of FluMist™ (MDV-A), was created by serial passage of the wild-type A/Ann Arbor/6/60 (A/AA/6/60) strain in primary chicken kidney tissue culture at successively lower temperatures (Maassab (1967) *Adaptation and growth characteristics of influenza virus at 25 degrees C* *Nature* 213:612-4). MDV-A replicates efficiently at 25° C. (ca, cold adapted), but its growth is restricted at 38 and 39° C. (ts, temperature sensitive). Additionally, this virus does not replicate in the lungs of infected ferrets (att, attenuation). The ts phenotype is believed to contribute to the attenuation of the vaccine in humans by restricting its replication in all but the coolest regions of the respiratory tract. The stability of this property has been demonstrated in animal models and clinical studies. In contrast to the ts phenotype of influenza strains created by chemical mutagenesis, the ts property of MDV-A does not revert following passage through infected hamsters or in shed isolates from children (for a recent review, see Murphy & Coelingh (2002) *Principles underlying the development and use of live attenuated cold-adapted influenza A and B virus vaccines* *Viral Immunol* 15:295-323).

[0071] Clinical studies in over 20,000 adults and children involving 12 separate 6:2 reassortant strains have shown that these vaccines are attenuated, safe and efficacious (Belshe et al. (1998) *The efficacy of live attenuated, cold-adapted, trivalent, intranasal influenza virus vaccine in children* *N Engl J Med* 338:1405-12; Boyce et al. (2000) *Safety and immunogenicity of adjuvanted and unadjuvanted subunit influenza vaccines administered intranasally to healthy adults* *Vaccine* 19:217-26; Edwards et al. (1994) *A randomized controlled trial of cold adapted and inactivated vaccines for the prevention of influenza A disease* *J Infect Dis* 169:68-76; Nichol et al. (1999) *Effectiveness of live, attenuated intranasal influenza virus vaccine in healthy, working adults: a randomized controlled trial* *JAMA* 282: 137-44). Reassortants carrying the six internal genes of MDV-A and the two HA and NA gene

segments of a wild-type virus (i.e., a 6:2 reassortant) consistently maintain ca, ts and att phenotypes (Maassab et al. (1982) *Evaluation of a cold-recombinant influenza virus vaccine in ferrets* *J. Infect. Dis.* 146:780-900).

[0072] Production of such reassorted virus using B strains of influenza is more difficult, however, recent work (see, e.g., U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus") has shown an eight plasmid system for the generation of influenza B virus entirely from cloned cDNA. Methods for the production of attenuated live influenza A and B virus suitable for vaccine formulations, such as live virus vaccine formulations useful for intranasal administration were also shown.

[0073] The system and methods described previously are useful for the rapid production in cell culture of recombinant and reassortant influenza A and B viruses, including viruses suitable for use as vaccines, including live attenuated vaccines, such as vaccines suitable for intranasal administration. The sequences, methods, etc. of the current invention, are optionally used in conjunction with, or in combination with, such previous work involving, e.g., reassorted influenza viruses for vaccine production to produce viruses for vaccines.

Methods and Compositions for Prophylactic Administration of Vaccines

[0074] As stated above, alternatively, or in addition to, use in production of FluMist™ vaccine, the current invention can be used in other vaccine formulations. In general, recombinant and reassortant viruses of the invention (e.g., those comprising polynucleotides of SEQ ID NO:1-34 or polypeptides of SEQ ID NO:35-68, or similar strains of the virus sequences within SEQ ID NO:1-68, or fragments of any of the previous) can be administered prophylactically in an immunologically effective amount and in an appropriate carrier or excipient to stimulate an immune response specific for one or more strains of influenza virus as determined by the HA and/or NA sequence. Typically, the carrier or excipient is a pharmaceutically acceptable carrier or excipient, such as sterile water, aqueous saline solution, aqueous buffered saline solutions, aqueous dextrose solutions, aqueous glycerol solutions, ethanol, allantoic fluid from uninfected hen eggs (i.e., normal allantoic fluid or NAD), or combinations thereof. The preparation of such solutions insuring sterility, pH, isotonicity, and stability is effected according to protocols established in the art. Generally, a carrier or excipient is selected to minimize allergic and other undesirable effects, and to suit the particular route of administration, e.g., subcutaneous, intramuscular, intranasal, etc.

[0075] A related aspect of the invention provides methods for stimulating the immune system of an individual to produce a protective immune response against influenza virus. In the methods, an immunologically effective amount of a recombinant influenza virus (e.g., an HA and/or an NA molecule of the invention), an immunologically effective amount of a polypeptide of the invention, and/or an immunologically effective amount of a nucleic acid of the invention is administered to the individual in a physiologically acceptable carrier.

[0076] Generally, the influenza viruses of the invention are administered in a quantity sufficient to stimulate an immune

response specific for one or more strains of influenza virus (i.e., against the HA and/or NA strains of the invention). Preferably, administration of the influenza viruses elicits a protective immune response to such strains. Dosages and methods for eliciting a protective immune response against one or more influenza strains are known to those of skill in the art. See, e.g., U.S. Pat. No. 5,922,326; Wright et al., *Infect. Immun.* 37:397-400 (1982); Kim et al., *Pediatrics* 52:56-63 (1973); and Wright et al., *J. Pediatr.* 88:931-936 (1976). For example, influenza viruses are provided in the range of about 1-1000 HID_{50} (human infectious dose), i.e., about 10^5 - 10^8 pfu (plaque forming units) per dose administered. Typically, the dose will be adjusted within this range based on, e.g., age, physical condition, body weight, sex, diet, time of administration, and other clinical factors. The prophylactic vaccine formulation is systemically administered, e.g., by subcutaneous or intramuscular injection using a needle and syringe, or a needle-less injection device. Alternatively, the vaccine formulation is administered intranasally, either by drops, large particle aerosol (greater than about 10 microns), or spray into the upper respiratory tract. While any of the above routes of delivery results in a protective systemic immune response, intranasal administration confers the added benefit of eliciting mucosal immunity at the site of entry of the influenza virus. For intranasal administration, attenuated live virus vaccines are often preferred, e.g., an attenuated, cold adapted and/or temperature sensitive recombinant or reassortant influenza virus. See above. While stimulation of a protective immune response with a single dose is preferred, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect.

[0077] Typically, the attenuated recombinant influenza of this invention as used in a vaccine is sufficiently attenuated such that symptoms of infection, or at least symptoms of serious infection, will not occur in most individuals immunized (or otherwise infected) with the attenuated influenza virus. In some instances, the attenuated influenza virus can still be capable of producing symptoms of mild illness (e.g., mild upper respiratory illness) and/or of dissemination to unvaccinated individuals. However, its virulence is sufficiently abrogated such that severe lower respiratory tract infections do not occur in the vaccinated or incidental host.

[0078] Alternatively, an immune response can be stimulated by ex vivo or in vivo targeting of dendritic cells with influenza viruses comprising the sequences herein. For example, proliferating dendritic cells are exposed to viruses in a sufficient amount and for a sufficient period of time to permit capture of the influenza antigens by the dendritic cells. The cells are then transferred into a subject to be vaccinated by standard intravenous transplantation methods.

[0079] While stimulation of a protective immune response with a single dose is preferred, additional dosages can be administered, by the same or different route, to achieve the desired prophylactic effect. In neonates and infants, for example, multiple administrations may be required to elicit sufficient levels of immunity. Administration can continue at intervals throughout childhood, as necessary to maintain sufficient levels of protection against wild-type influenza infection. Similarly, adults who are particularly susceptible to repeated or serious influenza infection, such as, for example, health care workers, day care workers, family members of young children, the elderly, and individuals with compromised cardiopulmonary function may require multiple immunizations to establish and/or maintain protective

immune responses. Levels of induced immunity can be monitored, for example, by measuring amounts of neutralizing secretory and serum antibodies, and dosages adjusted or vaccinations repeated as necessary to elicit and maintain desired levels of protection.

[0080] Optionally, the formulation for prophylactic administration of the influenza viruses also contains one or more adjuvants for enhancing the immune response to the influenza antigens. Suitable adjuvants include: complete Freund's adjuvant, incomplete Freund's adjuvant, saponin, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil or hydrocarbon emulsions, bacille Calmette-Guerin (BCG), *Corynebacterium parvum*, and the synthetic adjuvants QS-21 and MF59.

[0081] If desired, prophylactic vaccine administration of influenza viruses can be performed in conjunction with administration of one or more immunostimulatory molecules. Immunostimulatory molecules include various cytokines, lymphokines and chemokines with immunostimulatory, immunopotentiating, and pro-inflammatory activities, such as interleukins (e.g., IL-1, IL-2, IL-3, IL-4, IL-12, IL-13); growth factors (e.g., granulocyte-macrophage (GM)-colony stimulating factor (CSF)); and other immunostimulatory molecules, such as macrophage inflammatory factor, Flt3 ligand, B7.1; B7.2, etc. The immunostimulatory molecules can be administered in the same formulation as the influenza viruses, or can be administered separately. Either the protein (e.g., an HA and/or NA polypeptide of the invention) or an expression vector encoding the protein can be administered to produce an immunostimulatory effect.

[0082] The above described methods are useful for therapeutically and/or prophylactically treating a disease or disorder, typically influenza, by introducing a vector of the invention comprising a heterologous polynucleotide encoding a therapeutically or prophylactically effective HA and/or NA polypeptide (or peptide) or HA and/or NA RNA (e.g., an antisense RNA or ribozyme) into a population of target cells in vitro, ex vivo or in vivo. Typically, the polynucleotide encoding the polypeptide (or peptide), or RNA, of interest is operably linked to appropriate regulatory sequences, e.g., as described herein. Optionally, more than one heterologous coding sequence is incorporated into a single vector or virus. For example, in addition to a polynucleotide encoding a therapeutically or prophylactically active HA and/or NA polypeptide or RNA, the vector can also include additional therapeutic or prophylactic polypeptides, e.g., antigens, co-stimulatory molecules, cytokines, antibodies, etc., and/or markers, and the like.

[0083] Although vaccination of an individual with an attenuated influenza virus of a particular strain of a particular subgroup can induce cross-protection against influenza virus of different strains and/or subgroups, cross-protection can be enhanced, if desired, by vaccinating the individual with attenuated influenza virus from at least two strains, e.g., each of which represents a different subgroup. Additionally, vaccine combinations can optionally include mixes of pandemic vaccines and non-pandemic strains. Vaccine mixtures (or multiple vaccinations) can comprise components from human strains and/or non-human influenza strains (e.g., avian and human, etc.). Similarly, the attenuated influenza virus

vaccines of this invention can optionally be combined with vaccines that induce protective immune responses against other infectious agents.

Polynucleotides of the Invention

[0084] Probes

[0085] The HA and NA polynucleotides of the invention, e.g., as shown in the sequences herein such as SEQ ID NO:1 through SEQ ID NO:34, and fragments thereof, are optionally used in a number of different capacities alternative to, or in addition to, the vaccines described above. Other exemplary uses are described herein for illustrative purpose and not as limitations on the actual range of uses, etc. Different methods of construction, purification, and characterization of the nucleotide sequences of the invention are also described herein.

[0086] In some embodiments, nucleic acids including one or more polynucleotide sequence of the invention are favorably used as probes for the detection of corresponding or related nucleic acids in a variety of contexts, such as in nucleic hybridization experiments, e.g., to find and/or characterize homologous influenza variants (e.g., homologues to sequences herein, etc.) infecting other species or in different influenza outbreaks, etc. The probes can be either DNA or RNA molecules, such as restriction fragments of genomic or cloned DNA, cDNAs, PCR amplification products, transcripts, and oligonucleotides, and can vary in length from oligonucleotides as short as about 10 nucleotides in length to full length sequences or cDNAs in excess of 1 kb or more. For example, in some embodiments, a probe of the invention includes a polynucleotide sequence or subsequence selected, e.g., from among SEQ ID NO:1-SEQ ID NO:34, or sequences complementary thereto. Alternatively, polynucleotide sequences that are variants of one of the above-designated sequences are used as probes. Most typically, such variants include one or a few conservative nucleotide variations. For example, pairs (or sets) of oligonucleotides can be selected, in which the two (or more) polynucleotide sequences are conservative variations of each other, wherein one polynucleotide sequence corresponds identically to a first variant or and the other(s) corresponds identically to additional variants. Such pairs of oligonucleotide probes are particularly useful, e.g., for specific hybridization experiments to detect polymorphic nucleotides or to, e.g., detect homologous influenza HA and NA variants, e.g., homologous to the current HA and NA sequences, infecting other species or present in different (e.g., either temporally and/or geographically different) influenza outbreaks. In other applications, probes are selected that are more divergent, that is probes that are at least about 91% (or about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 98.5%, about 98.7%, about 99%, about 99.1%, about 99.2%, about 99.3%, about 99.4%, about 99.5%, or about 99.6% or more about 99.7%, about 99.8%, about 99.9% or more) identical are selected.

[0087] The probes of the invention, e.g., as exemplified by sequences derived from the sequences herein, can also be used to identify additional useful polynucleotide sequences according to procedures routine in the art. In one set of embodiments, one or more probes, as described above, are utilized to screen libraries of expression products or chromosomal segments (e.g., expression libraries or genomic libraries) to identify clones that include sequences identical to, or with significant sequence similarity to, e.g., one or more

probe of, e.g., SEQ ID NO:1-SEQ ID NO:34, i.e., variants, homologues, etc. It will be understood that in addition to such physical methods as library screening, computer assisted bioinformatic approaches, e.g., BLAST and other sequence homology search algorithms, and the like, can also be used for identifying related polynucleotide sequences. Polynucleotide sequences identified in this manner are also a feature of the invention.

[0088] Oligonucleotide probes are optionally produced via a variety of methods well known to those skilled in the art. Most typically, they are produced by well known synthetic methods, such as the solid phase phosphoramidite triester method described by Beaucage and Caruthers (1981) *Tetrahedron Letts* 22(20):1859-1862, e.g., using an automated synthesizer, or as described in Needham-Van Devanter et al. (1984) *Nucl Acids Res*, 12:6159-6168. Oligonucleotides can also be custom made and ordered from a variety of commercial sources known to persons of skill. Purification of oligonucleotides, where necessary, is typically performed by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) *J Chrom* 255:137-149. The sequence of the synthetic oligonucleotides can be verified using the chemical degradation method of Maxam and Gilbert (1980) in Grossman and Moldave (eds.) Academic Press, New York, *Methods in Enzymology* 65:499-560. Custom oligos can also easily be ordered from a variety of commercial sources known to persons of skill.

[0089] In other circumstances, e.g., relating to attributes of cells or organisms expressing the polynucleotides and polypeptides of the invention (e.g., those harboring virus comprising the sequences of the invention), probes that are polypeptides, peptides or antibodies are favorably utilized. For example, isolated or recombinant polypeptides, polypeptide fragments and peptides derived from any of the amino acid sequences of the invention and/or encoded by polynucleotide sequences of the invention, e.g., selected from SEQ ID NO:1 through SEQ ID NO:34, are favorably used to identify and isolate antibodies, e.g., from phage display libraries, combinatorial libraries, polyclonal sera, and the like.

[0090] Antibodies specific for any a polypeptide sequence or subsequence, e.g., of SEQ ID NO:35 through SEQ ID NO:68, and/or encoded by polynucleotide sequences of the invention, e.g., selected from SEQ ID NO:1 through SEQ ID NO:34, are likewise valuable as probes for evaluating expression products, e.g., from cells or tissues. In addition, antibodies are particularly suitable for evaluating expression of proteins comprising amino acid subsequences, e.g., of those given herein, or encoded by polynucleotides sequences of the invention, e.g., selected from those shown herein, in situ, in a tissue array, in a cell, tissue or organism, e.g., an organism infected by an unidentified influenza virus or the like. Antibodies can be directly labeled with a detectable reagent, or detected indirectly by labeling of a secondary antibody specific for the heavy chain constant region (i.e., isotype) of the specific antibody. Antibodies against specific amino acids sequences herein (e.g., SEQ ID NOs: 35-68) are also useful in determining whether other influenza viruses are within the same strain as the current sequences (e.g., through an HI assay, etc.). Additional details regarding production of specific antibodies are provided below.

[0091] Diagnostic Assays

[0092] The nucleic acid sequences of the present invention can be used in diagnostic assays to detect influenza (and/or hemagglutinin and/or neuraminidase) in a sample, to detect

hemagglutinin-like and/or neuraminidase-like sequences, and to detect strain differences in clinical isolates of influenza using either chemically synthesized or recombinant polynucleotide fragments, e.g., selected from the sequences herein. For example, fragments of the hemagglutinin and/or neuraminidase sequences comprising at least between 10 and 20 nucleotides can be used as primers to amplify nucleic acids using polymerase chain reaction (PCR) methods well known in the art (e.g., reverse transcription-PCR) and as probes in nucleic acid hybridization assays to detect target genetic material such as influenza RNA in clinical specimens.

[0093] The probes of the invention, e.g., as exemplified by unique subsequences selected from, e.g., SEQ ID NO:1 through SEQ ID NO:34, can also be used to identify additional useful polynucleotide sequences (such as to characterize additional strains of influenza) according to procedures routine in the art. In one set of preferred embodiments, one or more probes, as described above, are utilized to screen libraries of expression products or cloned viral nucleic acids (i.e., expression libraries or genomic libraries) to identify clones that include sequences identical to, or with significant sequence identity to the sequences herein. In turn, each of these identified sequences can be used to make probes, including pairs or sets of variant probes as described above. It will be understood that in addition to such physical methods as library screening, computer assisted bioinformatic approaches, e.g., BLAST and other sequence homology search algorithms, and the like, can also be used for identifying related polynucleotide sequences.

[0094] The probes of the invention are particularly useful for detecting the presence and for determining the identity of influenza nucleic acids in cells, tissues or other biological samples (e.g., a nasal wash or bronchial lavage). For example, the probes of the invention are favorably utilized to determine whether a biological sample, such as a subject (e.g., a human subject) or model system (such as a cultured cell sample) has been exposed to, or become infected with influenza, or particular strain(s) of influenza. Detection of hybridization of the selected probe to nucleic acids originating in (e.g., isolated from) the biological sample or model system is indicative of exposure to or infection with the virus (or a related virus) from which the probe polynucleotide is selected.

[0095] It will be appreciated that probe design is influenced by the intended application. For example, where several allele-specific probe-target interactions are to be detected in a single assay, e.g., on a single DNA chip, it is desirable to have similar melting temperatures for all of the probes. Accordingly, the lengths of the probes are adjusted so that the melting temperatures for all of the probes on the array are closely similar (it will be appreciated that different lengths for different probes may be needed to achieve a particular T_m where different probes have different GC contents). Although melting temperature is a primary consideration in probe design, other factors are optionally used to further adjust probe construction, such as selecting against primer self-complementarity and the like.

[0096] Vectors, Promoters and Expression Systems

[0097] The present invention includes recombinant constructs incorporating one or more of the nucleic acid sequences described herein. Such constructs optionally include a vector, for example, a plasmid, a cosmid, a phage, a virus, a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), etc., into which one or more of the polynucleotide sequences of the invention, e.g., comprising

any of SEQ ID NO:1 through SEQ ID NO:34, or a subsequence thereof etc., has been inserted, in a forward or reverse orientation. For example, the inserted nucleic acid can include a viral chromosomal sequence or cDNA including all or part of at least one of the polynucleotide sequences of the invention. In one embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

[0098] The polynucleotides of the present invention can be included in any one of a variety of vectors suitable for generating sense or antisense RNA, and optionally, polypeptide (or peptide) expression products (e.g., a hemagglutinin and/or neuraminidase molecule of the invention, or fragments thereof). Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, pseudorabies, adenovirus, adeno-associated virus, retroviruses and many others (e.g., pCDL). Any vector that is capable of introducing genetic material into a cell, and, if replication is desired, which is replicable in the relevant host can be used.

[0099] In an expression vector, the HA and/or NA polynucleotide sequence of interest is physically arranged in proximity and orientation to an appropriate transcription control sequence (e.g., promoter, and optionally, one or more enhancers) to direct mRNA synthesis. That is, the polynucleotide sequence of interest is operably linked to an appropriate transcription control sequence. Examples of such promoters include: LTR or SV40 promoter, *E. coli* lac or trp promoter, phage lambda P_L promoter, and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.

[0100] A variety of promoters are suitable for use in expression vectors for regulating transcription of influenza virus genome segments. In certain embodiments, the cytomegalovirus (CMV) DNA dependent RNA Polymerase II (Pol II) promoter is utilized. If desired, e.g., for regulating conditional expression, other promoters can be substituted which induce RNA transcription under the specified conditions, or in the specified tissues or cells. Numerous viral and mammalian, e.g., human promoters are available, or can be isolated according to the specific application contemplated. For example, alternative promoters obtained from the genomes of animal and human viruses include such promoters as the adenovirus (such as Adenovirus 2), papilloma virus, hepatitis-B virus, polyoma virus, and Simian Virus 40 (SV40), and various retroviral promoters. Mammalian promoters include, among many others, the actin promoter, immunoglobulin promoters, heat-shock promoters, and the like.

[0101] Various embodiments of the current invention can comprise a number of different vector constructions. Such constructions are typically and preferably used in plasmid rescue systems to create viruses for use in vaccines (e.g., in live attenuated vaccines, in killed or inactivated vaccines, etc.). Thus, the invention includes recombinant DNA molecules having a transcription control element that binds a DNA-directed RNA polymerase that is operatively linked to a DNA sequence that encodes an RNA molecule, wherein the RNA molecule comprises a binding site specific for an RNA-directed RNA polymerase of a negative strand RNA virus,

operatively linked to an RNA sequence comprising the reverse complement of a mRNA coding sequence of a negative strand RNA virus. Also, the invention includes a recombinant DNA molecule that, upon transcription yields an RNA template that contains an RNA sequence comprising the reverse complement of an mRNA coding sequence of a negative strand RNA virus, and vRNA terminal sequences. The invention also includes a recombinant DNA molecule that upon transcription yields a replicable RNA template comprising the reverse complement of an mRNA coding sequence of a negative strand RNA virus. Such above recombinant DNA molecules typically involve wherein the negative strand RNA virus is influenza (e.g., influenza A or B, etc.). Also, the RNA molecule in such embodiments is typically an influenza genome segment and the RNA template is typically an influenza genome segment. The recombinant DNA molecules typically comprise wherein the RNA template is replicable, wherein the negative strand RNA virus is influenza, and wherein the RNA template is an influenza genome segment. Thus, the nucleic acids influenza segments typically comprise HA and/or NA genes (the corresponding nucleic acid of which is, e.g., in FIG. 1, or within similar strains of the strains having the nucleic acids in, e.g., FIG. 1).

[0102] The invention also includes methods of preparing an RNA molecule comprising transcribing a recombinant DNA molecule with a DNA-directed RNA polymerase, wherein the DNA molecule comprises a transcription control element that binds a DNA-directed RNA polymerase that is operatively linked to a DNA sequence that encodes an RNA molecule, wherein the RNA molecule comprises a binding site specific for an RNA-directed RNA polymerase of a negative strand RNA virus, operatively linked to an RNA sequence comprising the reverse complement of an mRNA coding sequence of a negative strand RNA virus. The invention also includes a method of preparing an RNA molecule comprising transcribing a recombinant DNA molecule with a DNA-directed RNA polymerase, wherein the recombinant DNA molecule yields upon transcription an RNA molecule that contains an RNA sequence comprising the reverse complement of an mRNA coding sequence of a negative strand RNA virus, and vRNA terminal sequences. Furthermore, the invention includes a method of preparing an RNA molecule comprising transcribing a recombinant DNA molecule with a DNA-directed RNA polymerase, wherein the recombinant DNA molecule yields upon transcription a replicable RNA molecule comprising the reverse complement of an mRNA coding sequence of a negative strand RNA virus. Such methods typically comprise wherein the negative strand RNA virus is influenza, and wherein the RNA molecule is an influenza genome segment. Such methods preferably include wherein the DNA-directed RNA polymerase is pol I, pol II, T7 polymerase, T3 polymerase, or Sp6 polymerase. Thus, again, the influenza nucleic acid segments typically comprise HA and/or NA genes as described throughout.

[0103] Other methods within the invention include methods of constructing a DNA molecule comprising a transcription control element that binds a DNA-directed RNA polymerase that is operatively linked to a DNA sequence that encodes an RNA molecule, wherein the RNA molecule comprises a binding site specific for an RNA-directed RNA polymerase of an influenza virus, operatively linked to an RNA sequence comprising the reverse complement of an mRNA coding sequence of an influenza virus, wherein the DNA sequence comprises a nucleic acid corresponding to one or

more of SEQ ID NO:1-34 or a fragment thereof or of one or more nucleic acid sequence of a similar strain (e.g., a strain similar to such strains having the sequences found in the sequences of FIG. 1, etc.). Also, the invention includes a method of constructing a DNA molecule comprising a DNA sequence that upon transcription yields an RNA template that contains an RNA sequence comprising the reverse complement of an mRNA coding sequence of an influenza virus, and vRNA terminal sequences, wherein the DNA sequence comprises a nucleic acid corresponding to one or more of SEQ ID NO:1-34 or a fragment thereof, or of one or more nucleic acid of a similar strain (e.g., a strain similar to such strains that have the sequences found in FIG. 1, etc.). Such methods also include wherein the RNA template is replicable. Other methods of the invention include those of constructing a DNA molecule comprising a DNA sequence that upon transcription yields a replicable RNA template comprising the reverse complement of an mRNA coding sequence of an influenza virus. These methods of the invention typically include wherein the RNA molecule is an influenza genome segment, wherein the DNA-directed RNA polymerase is pol I, pol II, T7 polymerase, T3 polymerase, or Sp6 polymerase.

[0104] Transcription is optionally increased by including an enhancer sequence. Enhancers are typically short, e.g., 10-500 bp, cis-acting DNA elements that act in concert with a promoter to increase transcription. Many enhancer sequences have been isolated from mammalian genes (hemoglobin, elastase, albumin, alpha-fetoprotein, and insulin), and eukaryotic cell viruses. The enhancer can be spliced into the vector at a position 5' or 3' to the heterologous coding sequence, but is typically inserted at a site 5' to the promoter. Typically, the promoter, and if desired, additional transcription enhancing sequences are chosen to optimize expression in the host cell type into which the heterologous DNA is to be introduced (Scharf et al. (1994) *Heat stress promoters and transcription factors Results Probl Cell Differ* 20:125-62; Kriegler et al. (1990) *Assembly of enhancers, promoters, and splice signals to control expression of transferred genes Methods in Enzymol* 185: 512-27). Optionally, the amplicon can also contain a ribosome binding site or an internal ribosome entry site (IRES) for translation initiation.

[0105] The vectors of the invention also favorably include sequences necessary for the termination of transcription and for stabilizing the mRNA, such as a polyadenylation site or a terminator sequence. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. In one embodiment, the SV40 polyadenylation signal sequences can provide a bi-directional polyadenylation site that insulates transcription of (+) strand mRNA molecules from the Poll promoter initiating replication of the (-) strand viral genome.

[0106] In addition, as described above, the expression vectors optionally include one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells, in addition to genes previously listed, markers such as dihydrofolate reductase or neomycin resistance are suitable for selection in eukaryotic cell culture.

[0107] The vector containing the appropriate nucleic acid sequence as described above, as well as an appropriate promoter or control sequence, can be employed to transform a host cell permitting expression of the protein. While the vectors of the invention can be replicated in bacterial cells, most frequently it will be desirable to introduce them into mam-

malian cells, e.g., Vero cells, BHK cells, MDCK cell, 293 cells, COS cells, or the like, for the purpose of expression.

[0108] As described elsewhere, the HA and NA sequences herein, in various embodiments, can be comprised within plasmids involved in plasmid-rescue reassortment. See, e.g., U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus"; Hoffmann, E., 2000, *PNAS*, 97(11):6108-6113; U.S. Published Patent Application No. 20020164770 to Hoffmann; and U.S. Pat. No. 6,544,785 issued Apr. 8, 2003 to Palese, et al. The reassortants produced can include the HA and NA genes arranged with the 6 other influenza genes from the A/Ann Arbor/6/60 donor strain, the B/Ann Arbor/1/66 donor strain (and/or derivatives and modifications thereof), the A/Puerto Rico/8/34 donor strain, etc.

[0109] Additional Expression Elements

[0110] Most commonly, the genome segment encoding the influenza virus HA and/or NA protein includes any additional sequences necessary for its expression, including translation into a functional viral protein. In other situations, a minigene, or other artificial construct encoding the viral proteins, e.g., an HA and/or NA protein, can be employed. Again, in such case, it is often desirable to include specific initiation signals that aid in the efficient translation of the heterologous coding sequence. These signals can include, e.g., the ATG initiation codon and adjacent sequences. To insure translation of the entire insert, the initiation codon is inserted in the correct reading frame relative to the viral protein. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

[0111] If desired, polynucleotide sequences encoding additional expressed elements, such as signal sequences, secretion or localization sequences, and the like can be incorporated into the vector, usually, in-frame with the polynucleotide sequence of interest, e.g., to target polypeptide expression to a desired cellular compartment, membrane, or organelle, or to direct polypeptide secretion to the periplasmic space or into the cell culture media. Such sequences are known to those of skill, and include secretion leader peptides, organelle targeting sequences (e.g., nuclear localization sequences, ER retention signals, mitochondrial transit sequences), membrane localization/anchor sequences (e.g., stop transfer sequences, GPI anchor sequences), and the like.

[0112] Where translation of a polypeptide encoded by a nucleic acid sequence of the invention is desired, additional translation specific initiation signals can improve the efficiency of translation. These signals can include, e.g., an ATG initiation codon and adjacent sequences, an IRES region, etc. In some cases, for example, full-length cDNA molecules or chromosomal segments including a coding sequence incorporating, e.g., a polynucleotide sequence of the invention (e.g., as in the sequences herein), a translation initiation codon and associated sequence elements are inserted into the appropriate expression vector simultaneously with the polynucleotide sequence of interest. In such cases, additional translational control signals frequently are not required. However, in cases where only a polypeptide coding sequence, or a portion thereof, is inserted, exogenous translational control signals, including, e.g., an ATG initiation codon is often provided for expression of the relevant sequence. The initia-

tion codon is put in the correct reading frame to ensure transcription of the polynucleotide sequence of interest. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use (see, e.g., Scharf D. et al. (1994) *Results Probl Cell Differ* 20:125-62; Bittner et al. (1987) *Methods in Enzymol* 153:516-544).

[0113] Production of Recombinant Virus

[0114] Negative strand RNA viruses can be genetically engineered and recovered using a recombinant reverse genetics approach (U.S. Pat. No. 5,166,057 to Palese et al.). Such method was originally applied to engineer influenza viral genomes (Luytjes et al. (1989) *Cell* 59:1107-1113; Enami et al. (1990) *Proc. Natl. Acad. Sci. USA* 92:11563-11567), and has been successfully applied to a wide variety of segmented and nonsegmented negative strand RNA viruses, e.g., rabies (Schnell et al. (1994) *EMBO J.* 13: 4195-4203); VSV (Lawson et al. (1995) *Proc. Natl. Acad. Sci. USA* 92: 4477-4481); measles virus (Radecke et al. (1995) *EMBO J.* 14:5773-5784); rinderpest virus (Baron & Barrett (1997) *J. Virol.* 71: 1265-1271); human parainfluenza virus (Hoffman & Banerjee (1997) *J. Virol.* 71: 3272-3277; Dubin et al. (1997) *Virology* 235:323-332); SV5 (He et al. (1997) *Virology* 237:249-260); canine distemper virus (Gassen et al. (2000) *J. Virol.* 74:10737-44); and Sendai virus (Park et al. (1991) *Proc. Natl. Acad. Sci. USA* 88: 5537-5541; Kato et al. (1996) *Genes to Cells* 1:569-579). Those of skill in the art will be familiar with these and similar techniques to produce influenza virus comprising the HA and NA sequences of the invention. Recombinant influenza viruses produced according to such methods are also a feature of the invention, as are recombinant influenza virus comprising one or more nucleic acids and/or polypeptides of the invention. Of course, as will be appreciated by those of skill in the art, influenza virus in general (and those of the invention as well) are RNA viruses. Thus, when the present invention describes influenza viruses as comprising, e.g., the sequences of FIG. 1, etc., it is to be understood to mean the corresponding RNA version of the sequence. The nucleotide sequences in FIG. 1 comprise DNA versions (e.g., coding plus sense, etc.) of the genes (along with some untranslated regions in the nucleotide sequences). Those of skill in the art can easily convert between RNA and DNA sequences, between complementary nucleotide sequences, a corresponding RNA sequence, etc. Thus, for example, those of skill in the art can easily convert from a nucleotide sequence (e.g., one given in FIG. 1 such as SEQ ID NO:1) to the corresponding amino acid sequence or a corresponding complementary sequence, etc. Also, as will be evident, when such HA and/or NA sequences are described within DNA vectors, e.g., plasmids, etc. then the corresponding DNA version of the sequences are to be used.

[0115] Cell Culture and Expression Hosts

[0116] The present invention also relates to host cells that are introduced (transduced, transformed or transfected) with vectors of the invention, and the production of polypeptides of the invention by recombinant techniques. Host cells are genetically engineered (i.e., transduced, transformed or transfected) with a vector, such as an expression vector, of this invention. As described above, the vector can be in the form of a plasmid, a viral particle, a phage, etc. Examples of appropriate expression hosts include: bacterial cells, such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells,

such as *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Neurospora crassa*; or insect cells such as *Drosophila* and *Spodoptera frugiperda*.

[0117] Most commonly, mammalian cells are used to culture the HA and NA molecules of the invention. Suitable host cells for the replication of influenza virus (e.g., with the HA and/or NA sequences herein) include, e.g., Vero cells, BHK cells, MDCK cells, 293 cells and COS cells, including 293T cells, COS7 cells or the like. Commonly, co-cultures including two of the above cell lines, e.g., MDCK cells and either 293T or COS cells are employed at a ratio, e.g., of 1:1, to improve replication efficiency. Typically, cells are cultured in a standard commercial culture medium, such as Dulbecco's modified Eagle's medium supplemented with serum (e.g., 10% fetal bovine serum), or in serum free medium, under controlled humidity and CO₂ concentration suitable for maintaining neutral buffered pH (e.g., at pH between 7.0 and 7.2). Optionally, the medium contains antibiotics to prevent bacterial growth, e.g., penicillin, streptomycin, etc., and/or additional nutrients, such as L-glutamine, sodium pyruvate, non-essential amino acids, additional supplements to promote favorable growth characteristics, e.g., trypsin, β -mercaptoethanol, and the like.

[0118] The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the inserted polynucleotide sequences, e.g., through production of viruses. The culture conditions, such as temperature, pH and the like, are typically those previously used with the particular host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, e.g., Freshney (1994) *Culture of Animal Cells, a Manual of Basic Technique*, 3rd edition, Wiley-Liss, New York and the references cited therein. Other helpful references include, e.g., Paul (1975) *Cell and Tissue Culture*, 5th ed., Livingston, Edinburgh; Adams (1980) *Laboratory Techniques in Biochemistry and Molecular Biology-Cell Culture for Biochemists*, Work and Burdon (eds.) Elsevier, Amsterdam. Additional details regarding tissue culture procedures of particular interest in the production of influenza virus in vitro include, e.g., Merten et al. (1996) *Production of influenza virus in cell cultures for vaccine preparation*. in Cohen and Shafferman (eds.) *Novel Strategies in Design and Production of Vaccines*, which is incorporated herein in its entirety for all purposes. Additionally, variations in such procedures adapted to the present invention are readily determined through routine experimentation and will be familiar to those skilled in the art.

[0119] Cells for production of influenza virus (e.g., having the HA and/or NA sequences of the invention) can be cultured in serum-containing or serum free medium. In some case, e.g., for the preparation of purified viruses, it is typically desirable to grow the host cells in serum free conditions. Cells can be cultured in small scale, e.g., less than 25 ml medium, culture tubes or flasks or in large flasks with agitation, in rotator bottles, or on microcarrier beads (e.g., DEAE-Dextran microcarrier beads, such as Dormacell, Pfeifer & Langen; Superbead, Flow Laboratories; styrene copolymer-tri-methylamine beads, such as Hillex, SoloHill, Ann Arbor) in flasks, bottles or reactor cultures. Microcarrier beads are small spheres (in the range of 100-200 microns in diameter) that provide a large surface area for adherent cell growth per volume of cell culture. For example a single liter of medium can include more than 20 million microcarrier beads provid-

ing greater than 8000 square centimeters of growth surface. For commercial production of viruses, e.g., for vaccine production, it is often desirable to culture the cells in a bioreactor or fermenter. Bioreactors are available in volumes from under 1 liter to in excess of 100 liters, e.g., Cyto3 Bioreactor (Osmonics, Minnetonka, Minn.); NBS bioreactors (New Brunswick Scientific, Edison, N.J.); laboratory and commercial scale bioreactors from B. Braun Biotech International (B. Braun Biotech, Melsungen, Germany).

[0120] Regardless of the culture volume, in many desired aspects of the current invention, it is important that the cultures be maintained at an appropriate temperature, to insure efficient recovery of recombinant and/or reassortant influenza virus using temperature dependent multi plasmid systems (see, e.g., U.S. Application No. 60/420,708, filed Oct. 23, 2002, U.S. application Ser. No. 10/423,828, filed Apr. 25, 2003, and U.S. Application No. 60/574,117, filed May 25, 2004, all entitled "Multi-Plasmid System for the Production of Influenza Virus"), heating of virus solutions for filtration, etc. Typically, a regulator, e.g., a thermostat, or other device for sensing and maintaining the temperature of the cell culture system and/or other solution, is employed to insure that the temperature is at the correct level during the appropriate period (e.g., virus replication, etc.).

[0121] In some embodiments herein (e.g., wherein reassorted viruses are to be produced from segments on vectors) vectors comprising influenza genome segments are introduced (e.g., transfected) into host cells according to methods well known in the art for introducing heterologous nucleic acids into eukaryotic cells, including, e.g., calcium phosphate co-precipitation, electroporation, microinjection, lipofection, and transfection employing polyamine transfection reagents. For example, vectors, e.g., plasmids, can be transfected into host cells, such as COS cells, 293T cells or combinations of COS or 293T cells and MDCK cells, using the polyamine transfection reagent TransIT-LT1 (Mirus) according to the manufacturer's instructions in order to produce reassorted viruses, etc. Thus, in one example, approximately 1 μ g of each vector is introduced into a population of host cells with approximately 2 μ l of TransIT-LT1 diluted in 160 μ l medium, preferably serum-free medium, in a total volume of 200 μ l. The DNA:transfection reagent mixtures are incubated at room temperature for 45 minutes followed by addition of 800 μ l of medium. The transfection mixture is added to the host cells, and the cells are cultured as described via other methods well known to those skilled in the art. Accordingly, for the production of recombinant or reassortant viruses in cell culture, vectors incorporating each of the 8 genome segments, (PB2, PB1, PA, NP, M, NS, HA and NA, e.g., of the invention) are mixed with approximately 20 μ l TransIT-LT1 and transfected into host cells. Optionally, serum-containing medium is replaced prior to transfection with serum-free medium, e.g., Opti-MEM I, and incubated for 4-6 hours.

[0122] Alternatively, electroporation can be employed to introduce such vectors incorporating influenza genome segments into host cells. For example, plasmid vectors incorporating an influenza A or influenza B virus are favorably introduced into Vero cells using electroporation according to the following procedure. In brief, approximately 5×10^6 Vero cells, e.g., grown in Modified Eagle's Medium (MEM) supplemented with 10% Fetal Bovine Serum (FBS) are resuspended in 0.4 ml OptiMEM and placed in an electroporation cuvette. Twenty micrograms of DNA in a volume of up to 25 μ l is added to the cells in the cuvette, which is then mixed

gently by tapping. Electroporation is performed according to the manufacturer's instructions (e.g., BioRad Gene Pulser II with Capacitance Extender Plus connected) at 300 volts, 950 microFarads with a time constant of between 28-33 msec. The cells are remixed by gently tapping and approximately 1-2 minutes following electroporation 0.7 ml MEM with 10% FBS is added directly to the cuvette. The cells are then transferred to two wells of a standard 6 well tissue culture dish containing 2 ml MEM, 10% FBS. The cuvette is washed to recover any remaining cells and the wash suspension is divided between the two wells. Final volume is approximately 3.5 mL. The cells are then incubated under conditions permissive for viral growth, e.g., at approximately 33° C. for cold adapted strains.

[0123] In mammalian host cells, a number of expression systems, such as viral-based systems, can be utilized. In cases where an adenovirus is used as an expression vector, a coding sequence is optionally ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome will result in a viable virus capable of expressing the polypeptides of interest in infected host cells (Logan and Shenk (1984) *Proc Natl Acad Sci* 81:3655-3659). In addition, transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

[0124] A host cell strain is optionally chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the protein include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation. Post-translational processing, which cleaves a precursor form into a mature form, of the protein is sometimes important for correct insertion, folding and/or function. Additionally proper location within a host cell (e.g., on the cell surface) is also important. Different host cells such as COS, CHO, BHK, MDCK, 293, 293T, COS7, etc. have specific cellular machinery and characteristic mechanisms for such post-translational activities and can be chosen to ensure the correct modification and processing of the current introduced, foreign protein.

[0125] For long-term, high-yield production of recombinant proteins encoded by, or having subsequences encoded by, the polynucleotides of the invention, stable expression systems are optionally used. For example, cell lines, stably expressing a polypeptide of the invention, are transfected using expression vectors that contain viral origins of replication or endogenous expression elements and a selectable marker gene. For example, following the introduction of the vector, cells are allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced sequences. Thus, resistant clumps of stably transformed cells, e.g., derived from single cell type, can be proliferated using tissue culture techniques appropriate to the cell type.

[0126] Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The cells expressing said protein can be sorted, isolated and/or purified. The protein or fragment thereof produced by a recombinant cell can be secreted, membrane-bound, or retained intracel-

lularly, depending on the sequence (e.g., depending upon fusion proteins encoding a membrane retention signal or the like) and/or the vector used.

[0127] Expression products corresponding to the nucleic acids of the invention can also be produced in non-animal cells such as plants, yeast, fungi, bacteria and the like. In addition to Sambrook, Berger and Ausubel, all infra, details regarding cell culture can be found in Payne et al. (1992) *Plant Cell and Tissue Culture in Liquid Systems* John Wiley & Sons, Inc. New York, N.Y.; Gamborg and Phillips (eds.) (1995) *Plant Cell Tissue and Organ Culture*; Fundamental Methods Springer Lab Manual, Springer-Verlag (Berlin Heidelberg N.Y.) and Atlas and Parks (eds.) *The Handbook of Microbiological Media* (1993) CRC Press, Boca Raton, Fla.

[0128] In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the expressed product. For example, when large quantities of a polypeptide or fragments thereof are needed for the production of antibodies, vectors that direct high-level expression of fusion proteins that are readily purified are favorably employed. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the coding sequence of interest, e.g., sequences comprising those found herein, etc., can be ligated into the vector in-frame with sequences for the amino-terminal translation initiating methionine and the subsequent 7 residues of beta-galactosidase producing a catalytically active beta galactosidase fusion protein; pIN vectors (Van Heeke & Schuster (1989) *J Biol Chem* 264:5503-5509); pET vectors (Novagen, Madison Wis.); and the like. Similarly, in the yeast *Saccharomyces cerevisiae* a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH can be used for production of the desired expression products. For reviews, see Ausubel, infra, and Grant et al., (1987); *Methods in Enzymology* 153:516-544.

[0129] Nucleic Acid Hybridization

[0130] Comparative hybridization can be used to identify nucleic acids of the invention, including conservative variations of nucleic acids of the invention. This comparative hybridization method is a preferred method of distinguishing nucleic acids of the invention. In addition, target nucleic acids which hybridize to the nucleic acids represented by, e.g., SEQ ID NO:1 through SEQ ID NO:34 under high, ultra-high and ultra-ultra-high stringency conditions are features of the invention. Examples of such nucleic acids include those with one or a few silent or conservative nucleic acid substitutions as compared to a given nucleic acid sequence.

[0131] A test target nucleic acid is said to specifically hybridize to a probe nucleic acid when it hybridizes at least one-half as well to the probe as to the perfectly matched complementary target, i.e., with a signal to noise ratio at least one-half as high as hybridization of the probe to the target under conditions in which the perfectly matched probe binds to the perfectly matched complementary target with a signal to noise ratio that is at least about 5x-10x as high as that observed for hybridization to any of the unmatched target nucleic acids.

[0132] Nucleic acids "hybridize" when they associate, typically in solution. Nucleic acids hybridize due to a variety of well-characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. Numerous protocols for nucleic acid hybridization are well known in the art. An extensive guide to the hybridization

of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes* part I chapter 2, “Overview of principles of hybridization and the strategy of nucleic acid probe assays,” (Elsevier, N.Y.), as well as in Ausubel, Sambrook, and Berger and Kimmel, all below. Hames and Higgins (1995) *Gene Probes* 1 IRL Press at Oxford University Press, Oxford, England, (Hames and Higgins 1) and Hames and Higgins (1995) *Gene Probes* 2 IRL Press at Oxford University Press, Oxford, England (Hames and Higgins 2) provide details on the synthesis, labeling, detection and quantification of DNA and RNA, including oligonucleotides.

[0133] An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is 50% formalin with 1 mg of heparin at 42° C., with the hybridization being carried out overnight. An example of stringent wash conditions comprises a 0.2×SSC wash at 65° C. for 15 minutes (see, Sambrook, *infra* for a description of SSC buffer). Often the high stringency wash is preceded by a low stringency wash to remove background probe signal. An example low stringency wash is 2×SSC at 40° C. for 15 minutes. In general, a signal to noise ratio of 5× (or higher) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization.

[0134] After hybridization, unhybridized nucleic acids can be removed by a series of washes, the stringency of which can be adjusted depending upon the desired results. Low stringency washing conditions (e.g., using higher salt and lower temperature) increase sensitivity, but can produce nonspecific hybridization signals and high background signals. Higher stringency conditions (e.g., using lower salt and higher temperature that is closer to the T_m) lower the background signal, typically with primarily the specific signal remaining. See, also, Rapley, R. and Walker, J. M. eds., *Molecular Biomethods Handbook* (Humana Press, Inc. 1998).

[0135] “Stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and northern hybridizations are sequence dependent, and are different under different environmental parameters. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993), *supra*, and in Hames and Higgins, 1 and 2. Stringent hybridization and wash conditions can easily be determined empirically for any test nucleic acid. For example, in determining highly stringent hybridization and wash conditions, the hybridization and wash conditions are gradually increased (e.g., by increasing temperature, decreasing salt concentration, increasing detergent concentration and/or increasing the concentration of organic solvents such as formalin in the hybridization or wash), until a selected set of criteria is met. For example, the hybridization and wash conditions are gradually increased until a probe binds to a perfectly matched complementary target with a signal to noise ratio that is at least 5× as high as that observed for hybridization of the probe to an unmatched target.

[0136] In general, a signal to noise ratio of at least 2× (or higher, e.g., at least 5×, 10×, 20×, 50×, 100×, or more) than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization. Detection of at least stringent hybridization between two sequences in the context of the present invention indicates

relatively strong structural similarity to, e.g., the nucleic acids of the present invention provided in the sequence listings herein.

[0137] “Very stringent” conditions are selected to be equal to the thermal melting point (T_m) for a particular probe. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the test sequence hybridizes to a perfectly matched probe. For the purposes of the present invention, generally, “highly stringent” hybridization and wash conditions are selected to be about 5° C. lower than the T_m for the specific sequence at a defined ionic strength and pH (as noted below, highly stringent conditions can also be referred to in comparative terms). Target sequences that are closely related or identical to the nucleotide sequence of interest (e.g., “probe”) can be identified under stringent or highly stringent conditions. Lower stringency conditions are appropriate for sequences that are less complementary.

[0138] “Ultra high-stringency” hybridization and wash conditions are those in which the stringency of hybridization and wash conditions are increased until the signal to noise ratio for binding of the probe to the perfectly matched complementary target nucleic acid is at least 10× as high as that observed for hybridization to any unmatched target nucleic acids. A target nucleic acid which hybridizes to a probe under such conditions, with a signal to noise ratio of at least one-half that of the perfectly matched complementary target nucleic acid is said to bind to the probe under ultra-high stringency conditions.

[0139] In determining stringent or highly stringent hybridization (or even more stringent hybridization) and wash conditions, the hybridization and wash conditions are gradually increased (e.g., by increasing temperature, decreasing salt concentration, increasing detergent concentration and/or increasing the concentration of organic solvents, such as formamide, in the hybridization or wash), until a selected set of criteria are met. For example, the hybridization and wash conditions are gradually increased until a probe comprising one or more polynucleotide sequences of the invention, e.g., sequences or unique subsequences selected from those given herein and/or complementary polynucleotide sequences, binds to a perfectly matched complementary target (again, a nucleic acid comprising one or more nucleic acid sequences or subsequences selected from those given herein and/or complementary polynucleotide sequences thereof), with a signal to noise ratio that is at least 2× (and optionally 5×, 10×, or 100× or more) as high as that observed for hybridization of the probe to an unmatched target (e.g., a polynucleotide sequence comprising one or more sequences or subsequences selected from known influenza sequences present in public databases such as GenBank at the time of filing, and/or complementary polynucleotide sequences thereof), as desired.

[0140] Using the polynucleotides of the invention, or subsequences thereof, novel target nucleic acids can be obtained; such target nucleic acids are also a feature of the invention. For example, such target nucleic acids include sequences that hybridize under stringent conditions to a unique oligonucleotide probe corresponding to any of the polynucleotides of the invention.

[0141] Similarly, even higher levels of stringency can be determined by gradually increasing the hybridization and/or wash conditions of the relevant hybridization assay. For example, those in which the stringency of hybridization and wash conditions are increased until the signal to noise ratio

for binding of the probe to the perfectly matched complementary target nucleic acid is at least 10x, 20x, 50x, 100x, or 500x or more as high as that observed for hybridization to any unmatched target nucleic acids. The particular signal will depend on the label used in the relevant assay, e.g., a fluorescent label, a calorimetric label, a radioactive label, or the like. A target nucleic acid which hybridizes to a probe under such conditions, with a signal to noise ratio of at least one-half that of the perfectly matched complementary target nucleic acid, is said to bind to the probe under ultra-ultra-high stringency conditions.

[0142] Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

[0143] Cloning, Mutagenesis and Expression of Biomolecules of Interest

[0144] General texts which describe molecular biological techniques, which are applicable to the present invention, such as cloning, mutation, cell culture and the like, include Berger and Kimmel, *Guide to Molecular Cloning Techniques Methods in Enzymology* volume 152 Academic Press, Inc., San Diego, Calif. (Berger); Sambrook et al., *Molecular Cloning—A Laboratory Manual* (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 2000 ("Sambrook") and *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2002) ("Ausubel"). These texts describe mutagenesis, the use of vectors, promoters and many other relevant topics related to, e.g., the generation of HA and/or NA molecules, etc.

[0145] Various types of mutagenesis are optionally used in the present invention, e.g., to produce and/or isolate, e.g., novel or newly isolated HA and/or NA molecules and/or to further modify/mutate the polypeptides (e.g., HA and NA molecules) of the invention. They include but are not limited to site-directed, random point mutagenesis, homologous recombination (DNA shuffling), mutagenesis using uracil containing templates, oligonucleotide-directed mutagenesis, phosphorothioate-modified DNA mutagenesis, mutagenesis using gapped duplex DNA or the like. Additional suitable methods include point mismatch repair, mutagenesis using repair-deficient host strains, restriction-selection and restriction-purification, deletion mutagenesis, mutagenesis by total gene synthesis, double-strand break repair, and the like. Mutagenesis, e.g., involving chimeric constructs, is also included in the present invention. In one embodiment, mutagenesis can be guided by known information of the naturally occurring molecule or altered or mutated naturally occurring molecule, e.g., sequence, sequence comparisons, physical properties, crystal structure or the like.

[0146] The above texts and examples found herein describe these procedures as well as the following publications (and references cited within): Sieber, et al., *Nature Biotechnology*, 19:456-460 (2001); Ling et al., *Approaches to DNA mutagenesis: an overview*, *Anal Biochem* 254(2): 157-178 (1997); Dale et al., *Oligonucleotide-directed random mutagenesis using the phosphorothioate method*, *Methods Mol Biol* 57:369-374 (1996); I. A. Lorimer, I. Pastan, *Nucleic Acids Res* 23, 3067-8 (1995); W. P. C. Stemmer, *Nature* 370, 389-91 (1994); Arnold, *Protein engineering for unusual environ-*

ments, *Current Opinion in Biotechnology* 4:450-455 (1993); Bass et al., *Mutant Trp repressors with new DNA-binding specificities*, *Science* 242:240-245 (1988); Fritz et al., *Oligonucleotide-directed construction of mutations: a gapped duplex DNA procedure without enzymatic reactions in vitro*, *Nucl Acids Res* 16: 6987-6999 (1988); Kramer et al., *Improved enzymatic in vitro reactions in the gapped duplex DNA approach to oligonucleotide-directed construction of mutations*, *Nucl Acids Res* 16: 7207 (1988); Sakamar and Khorana, *Total synthesis and expression of a gene for the α -subunit of bovine rod outer segment guanine nucleotide-binding protein (transducin)*, *Nucl Acids Res* 14: 6361-6372 (1988); Sayers et al., *Y-T Exonucleases in phosphorothioate-based oligonucleotide-directed mutagenesis*, *Nucl Acids Res* 16:791-802 (1988); Sayers et al., *Strand specific cleavage of phosphorothioate-containing DNA by reaction with restriction endonucleases in the presence of ethidium bromide*, (1988) *Nucl Acids Res* 16: 803-814; Carter, *Improved oligonucleotide-directed mutagenesis using M13 vectors*, *Methods in Enzymol* 154: 382-403 (1987); Kramer & Fritz *Oligonucleotide-directed construction of mutations via gapped duplex DNA*, *Methods in Enzymol* 154:350-367 (1987); Kunkel, *The efficiency of oligonucleotide directed mutagenesis*, in *Nucleic Acids & Molecular Biology* (Eckstein, F. and Lilley, D. M. J. eds., Springer Verlag, Berlin)) (1987); Kunkel et al., *Rapid and efficient site-specific mutagenesis without phenotypic selection*, *Methods in Enzymol* 154, 367-382 (1987); Zoller & Smith, *Oligonucleotide-directed mutagenesis: a simple method using two oligonucleotide primers and a single-stranded DNA template*, *Methods in Enzymol* 154: 329-350 (1987); Carter, *Site-directed mutagenesis*, *Biochem J* 237:1-7 (1986); Eghtedarzadeh & Henikoff, *Use of oligonucleotides to generate large deletions*, *Nucl Acids Res* 14: 5115 (1986); Mandecki, *Oligonucleotide-directed double-strand break repair in plasmids of Escherichia coli: a method for site-specific mutagenesis*, *Proc Natl Acad Sci USA*, 83:7177-7181 (1986); Nakamaye & Eckstein, *Inhibition of restriction endonuclease Nci I cleavage by phosphorothioate groups and its application to oligonucleotide-directed mutagenesis*, *Nucl Acids Res* 14: 9679-9698 (1986); Wells et al., *Importance of hydrogen-bond formation in stabilizing the transition state of subtilisin*, *Phil Trans R Soc Lond A* 317: 415-423 (1986); Botstein & Shortle, *Strategies and applications of in vitro mutagenesis*, *Science* 229:1193-1201 (1985); Carter et al., *Improved oligonucleotide site-directed mutagenesis using M13 vectors*, *Nucl Acids Res* 13: 4431-4443 (1985); Grundstrom et al., *Oligonucleotide-directed mutagenesis by microscale 'shot-gun' gene synthesis*, *Nucl Acids Res* 13:3305-3316 (1985); Kunkel, *Rapid and efficient site-specific mutagenesis without phenotypic selection*, *Proc Natl Acad Sci USA* 82:488-492 (1985); Smith, *In vitro mutagenesis*, *Ann Rev Genet* 19:423-462 (1985); Taylor et al., *The use of phosphorothioate-modified DNA in restriction enzyme reactions to prepare nicked DNA*, *Nucl Acids Res* 13: 8749-8764 (1985); Taylor et al., *The rapid generation of oligonucleotide-directed mutations at high frequency using phosphorothioate-modified DNA*, *Nucl Acids Res* 13: 8765-8787 (1985); Wells et al., *Cassette mutagenesis: an efficient method for generation of multiple mutations at defined sites*, *Gene* 34:315-323 (1985); Kramer et al., *The gapped duplex DNA approach to oligonucleotide-directed mutation construction*, *Nucl Acids Res* 12: 9441-9456 (1984); Kramer et al., *PointMismatch Repair*, *Cell* 38:879-887 (1984); Nambiar et al., *Total synthesis and cloning of a gene coding for the*

ribonuclease S protein, *Science* 223: 1299-1301 (1984); Zoller & Smith, *Oligonucleotide-directed mutagenesis of DNA fragments cloned into M13 vectors*, *Methods in Enzymol* 100:468-500 (1983); and Zoller & Smith, *Oligonucleotide-directed mutagenesis using M13-derived vectors: an efficient and general procedure for the production of point mutations in any DNA fragment*, *Nucl Acids Res* 10:6487-6500 (1982). Additional details on many of the above methods can be found in *Methods in Enzymol* Volume 154, which also describes useful controls for trouble-shooting problems with various mutagenesis, gene isolation, expression, and other methods.

[0147] Oligonucleotides, e.g., for use in mutagenesis of the present invention, e.g., mutating libraries of the HA and/or NA molecules of the invention, or altering such, are typically synthesized chemically according to the solid phase phosphoramidite triester method described by Beaucage and Caruthers, *Tetrahedron Letts* 22(20):1859-1862, (1981) e.g., using an automated synthesizer, as described in Needham-VanDevanter et al., *Nucleic Acids Res*, 12:6159-6168 (1984).

[0148] In addition, essentially any nucleic acid can be custom or standard ordered from any of a variety of commercial sources, such as The Midland Certified Reagent Company (mcrc@oligos.com), The Great American Gene Company (www.genco.com), ExpressGen Inc. (www.expressgen.com), Operon Technologies Inc. (Alameda, Calif.) and many others. Similarly, peptides and antibodies can be custom ordered from any of a variety of sources, such as PeptideGenic (available at pkim@ccnet.com), HTI Bio-products, Inc. (www.htibio.com), BMA Biomedicals Ltd. (U.K.), Bio-Synthesis, Inc., and many others.

[0149] The present invention also relates to host cells and organisms comprising a HA and/or NA molecule or other polypeptide and/or nucleic acid of the invention or such HA and/or NA or other sequences within various vectors such as 6:2 reassortant influenza viruses, plasmids in plasmid rescue systems, etc. Host cells are genetically engineered (e.g., transformed, transduced or transfected) with the vectors of this invention, which can be, for example, a cloning vector or an expression vector. The vector can be, for example, in the form of a plasmid, a bacterium, a virus, a naked polynucleotide, or a conjugated polynucleotide. The vectors are introduced into cells and/or microorganisms by standard methods including electroporation (see, From et al., *Proc Natl Acad Sci USA* 82, 5824 (1985), infection by viral vectors, high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface (Klein et al., *Nature* 327, 70-73 (1987)). Berger, Sambrook, and Ausubel provide a variety of appropriate transformation methods. See, above.

[0150] Several well-known methods of introducing target nucleic acids into bacterial cells are available, any of which can be used in the present invention. These include: fusion of the recipient cells with bacterial protoplasts containing the DNA, electroporation, projectile bombardment, and infection with viral vectors, etc. Bacterial cells can be used to amplify the number of plasmids containing DNA constructs of this invention. The bacteria are grown to log phase and the plasmids within the bacteria can be isolated by a variety of methods known in the art (see, for instance, Sambrook). In addition, a plethora of kits are commercially available for the purification of plasmids from bacteria, (see, e.g., EasyPrep™, FlexiPrep™, both from Pharmacia Biotech; StrataClean™, from Stratagene; and, QIAprep™ from Qiagen). The isolated

and purified plasmids are then further manipulated to produce other plasmids, used to transfect cells or incorporated into related vectors to infect organisms. Typical vectors contain transcription and translation terminators, transcription and translation initiation sequences, and promoters useful for regulation of the expression of the particular target nucleic acid. The vectors optionally comprise generic expression cassettes containing at least one independent terminator sequence, sequences permitting replication of the cassette in eukaryotes, or prokaryotes, or both, (e.g., shuttle vectors) and selection markers for both prokaryotic and eukaryotic systems. Vectors are suitable for replication and integration in prokaryotes, eukaryotes, or preferably both. See, Gilman & Smith, *Gene* 8:81 (1979); Roberts, et al., *Nature*, 328:731 (1987); Schneider, B., et al., *Protein Expr Purif* 6435:10 (1995); Ausubel, Sambrook, Berger (all supra). A catalogue of Bacteria and Bacteriophages useful for cloning is provided, e.g., by the ATCC, e.g., The ATCC Catalogue of Bacteria and Bacteriophage (1992) Gherna et al. (eds.) published by the ATCC. Additional basic procedures for sequencing, cloning and other aspects of molecular biology and underlying theoretical considerations are also found in Watson et al. (1992) *Recombinant DNA Second Edition* Scientific American Books, NY. See, above.

Polypeptide Production and Recovery

[0151] In some embodiments, following transduction of a suitable host cell line or strain and growth of the host cells to an appropriate cell density, a selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period. In some embodiments, a secreted polypeptide product, e.g., a HA and/or NA polypeptide as in a secreted fusion protein form, etc., is then recovered from the culture medium. In other embodiments, a virus particle containing a HA and/or a NA polypeptide of the invention is produced from the cell. Alternatively, cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Eukaryotic or microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, or other methods, which are well known to those skilled in the art. Additionally, cells expressing a HA and/or a NA polypeptide product of the invention can be utilized without separating the polypeptide from the cell. In such situations, the polypeptide of the invention is optionally expressed on the cell surface and is examined thus (e.g., by having HA and/or NA molecules, or fragments thereof, e.g., comprising fusion proteins or the like) on the cell surface bind antibodies, etc. Such cells are also features of the invention.

[0152] Expressed polypeptides can be recovered and purified from recombinant cell cultures by any of a number of methods well known in the art, including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography (e.g., using any of the tagging systems known to those skilled in the art), hydroxylapatite chromatography, and lectin chromatography. Protein refolding steps can be used, as desired, in completing configuration of the mature protein. Also, high performance liquid chromatography (HPLC) can be employed in the final purification steps. In addition to the references noted herein, a variety of purification methods are

well known in the art, including, e.g., those set forth in Sandana (1997) *Bioseparation of Proteins*, Academic Press, Inc.; and Bollag et al. (1996) *Protein Methods*, 2nd Edition Wiley-Liss, NY; Walker (1996) *The Protein Protocols Handbook Humana Press*, NJ; Harris and Angal (1990) *Protein Purification Applications: A Practical Approach* IRL Press at Oxford, Oxford, England; Harris and Angal *Protein Purification Methods: A Practical Approach* IRL Press at Oxford, Oxford, England; Scopes (1993) *Protein Purification: Principles and Practice 3rd Edition* Springer Verlag, NY; Janson and Ryden (1998) *Protein Purification: Principles High Resolution Methods and Applications, Second Edition* Wiley-VCH, NY; and Walker (1998) *Protein Protocols on CD-ROM* Humana Press, NJ.

[0153] When the expressed polypeptides of the invention are produced in viruses, the viruses are typically recovered from the culture medium, in which infected (transfected) cells have been grown. Typically, crude medium is clarified prior to concentration of influenza viruses. Common methods include ultrafiltration, adsorption on barium sulfate and elution, and centrifugation. For example, crude medium from infected cultures can first be clarified by centrifugation at, e.g., 1000-2000×g for a time sufficient to remove cell debris and other large particulate matter, e.g., between 10 and 30 minutes. Optionally, the clarified medium supernatant is then centrifuged to pellet the influenza viruses, e.g., at 15,000×g, for approximately 3-5 hours. Following resuspension of the virus pellet in an appropriate buffer, such as STE (0.01 M Tris-HCl; 0.15 M NaCl; 0.0001 M EDTA) or phosphate buffered saline (PBS) at pH 7.4, the virus is concentrated by density gradient centrifugation on sucrose (60%-12%) or potassium tartrate (50%-10%). Either continuous or step gradients, e.g., a sucrose gradient between 12% and 60% in four 12% steps, are suitable. The gradients are centrifuged at a speed, and for a time, sufficient for the viruses to concentrate into a visible band for recovery. Alternatively, and for most large-scale commercial applications, virus is elutriated from density gradients using a zonal-centrifuge rotor operating in continuous mode. Additional details sufficient to guide one of skill through the preparation of influenza viruses from tissue culture are provided, e.g., in Furminger. *Vaccine Production*, in Nicholson et al. (eds.) *Textbook of Influenza* pp. 324-332; Merten et al. (1996) *Production of influenza virus in cell cultures for vaccine preparation*, in Cohen & Shafferman (eds.) *Novel Strategies in Design and Production of Vaccines* pp. 141-151, and U.S. Pat. No. 5,690,937. If desired, the recovered viruses can be stored at -80° C. in the presence of sucrose-phosphate-glutamate (SPG) as a stabilizer.

[0154] Alternatively, cell-free transcription/translation systems can be employed to produce polypeptides comprising an amino acid sequence or subsequence of, e.g., SEQ ID NO:35 through SEQ ID NO:68, or encoded by the polynucleotide sequences of the invention. A number of suitable in vitro transcription and translation systems are commercially available. A general guide to in vitro transcription and translation protocols is found in Tymms (1995) *In vitro Transcription and Translation Protocols: Methods in Molecular Biology* Volume 37, Garland Publishing, NY.

[0155] In addition, the polypeptides, or subsequences thereof, e.g., subsequences comprising antigenic peptides, can be produced manually or by using an automated system, by direct peptide synthesis using solid-phase techniques (see, Stewart et al. (1969) *Solid-Phase Peptide Synthesis*, WH Freeman Co, San Francisco; Merrifield J (1963) *J Am Chem*

Soc 85:2149-2154). Exemplary automated systems include the Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City, Calif.). If desired, subsequences can be chemically synthesized separately, and combined using chemical methods to provide full-length polypeptides.

[0156] Modified Amino Acids

[0157] Expressed polypeptides of the invention can contain one or more modified amino acids. The presence of modified amino acids can be advantageous in, for example, (a) increasing polypeptide serum half-life, (b) reducing/increasing polypeptide antigenicity, (c) increasing polypeptide storage stability, etc. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production (e.g., N-linked glycosylation at N-X-S/T motifs during expression in mammalian cells) or modified by synthetic means (e.g., via PEGylation).

[0158] Non-limiting examples of a modified amino acid include a glycosylated amino acid, a sulfated amino acid, a prenylated (e.g., farnesylated, geranylgeranylated) amino acid, an acetylated amino acid, an acylated amino acid, a PEG-ylated amino acid, a biotinylated amino acid, a carboxylated amino acid, a phosphorylated amino acid, and the like, as well as amino acids modified by conjugation to, e.g., lipid moieties or other organic derivatizing agents. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature. Example protocols are found in Walker (1998) *Protein Protocols on CD-ROM* Human Press, Towata, N.J.

[0159] Fusion Proteins

[0160] The present invention also provides fusion proteins comprising fusions of the sequences of the invention (e.g., encoding HA and/or NA polypeptides) or fragments thereof with, e.g., immunoglobulins (or portions thereof), sequences encoding, e.g., GFP (green fluorescent protein), or other similar markers, etc. Nucleotide sequences encoding such fusion proteins are another aspect of the invention. Fusion proteins of the invention are optionally used for, e.g., similar applications (including, e.g., therapeutic, prophylactic, diagnostic, experimental, etc. applications as described herein) as the non-fusion proteins of the invention. In addition to fusion with immunoglobulin sequences and marker sequences, the proteins of the invention are also optionally fused with, e.g., sequences which allow sorting of the fusion proteins and/or targeting of the fusion proteins to specific cell types, regions, etc.

[0161] Antibodies

[0162] The polypeptides of the invention can be used to produce antibodies specific for the polypeptides given herein and/or polypeptides encoded by the polynucleotides of the invention, e.g., those shown herein, and conservative variants thereof. Antibodies specific for the above mentioned polypeptides are useful, e.g., for diagnostic and therapeutic purposes, e.g., related to the activity, distribution, and expression of target polypeptides. For example, such antibodies can optionally be utilized to define other viruses within the same strain(s) as the HA/NA sequences herein.

[0163] Antibodies specific for the polypeptides of the invention can be generated by methods well known in the art. Such antibodies can include, but are not limited to, polyclonal, monoclonal, chimeric, humanized, single chain, Fab fragments and fragments produced by an Fab expression library.

[0164] Polypeptides do not require biological activity for antibody production (e.g., full length functional hemaggluti-

nin or neuraminidase is not required). However, the polypeptide or oligopeptide must be antigenic. Peptides used to induce specific antibodies typically have an amino acid sequence of at least about 4 amino acids, and often at least 5 or 10 amino acids. Short stretches of a polypeptide can be fused with another protein, such as keyhole limpet hemocyanin, and antibody produced against the chimeric molecule.

[0165] Numerous methods for producing polyclonal and monoclonal antibodies are known to those of skill in the art, and can be adapted to produce antibodies specific for the polypeptides of the invention, and/or encoded by the polynucleotide sequences of the invention, etc. See, e.g., Coligan (1991) *Current Protocols in Immunology* Wiley/Greene, NY; Paul (ed.) (1998) *Fundamental Immunology, Fourth Edition*, Lippincott-Raven, Lippincott Williams & Wilkins; Harlow and Lane (1989) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, NY; Stites et al. (eds.) *Basic and Clinical Immunology* (4th ed.) Lange Medical Publications, Los Altos, Calif., and references cited therein; Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d ed.) Academic Press, New York, N.Y.; and Kohler and Milstein (1975) *Nature* 256: 495-497. Other suitable techniques for antibody preparation include selection of libraries of recombinant antibodies in phage or similar vectors. See, Huse et al. (1989) *Science* 246: 1275-1281; and Ward, et al. (1989) *Nature* 341: 544-546. Specific monoclonal and polyclonal antibodies and antisera will usually bind with a K_D of, e.g., at least about 0.1 μM , at least about 0.01 μM or better, and, typically and at least about 0.001 μM or better.

[0166] For certain therapeutic applications, humanized antibodies are desirable. Detailed methods for preparation of chimeric (humanized) antibodies can be found in U.S. Pat. No. 5,482,856. Additional details on humanization and other antibody production and engineering techniques can be found in Borrebaeck (ed.) (1995) *Antibody Engineering 2nd Edition* Freeman and Company, NY (Borrebaeck); McCafferty et al. (1996) *Antibody Engineering, A Practical Approach* IRL at Oxford Press, Oxford, England (McCafferty), and Paul (1995) *Antibody Engineering Protocols Humana Press*, Towata, N.J. (Paul). Additional details regarding specific procedures can be found, e.g., in Ostberg et al. (1983), *Hybridoma* 2: 361-367, Ostberg, U.S. Pat. No. 4,634,664, and Engelman et al., U.S. Pat. No. 4,634,666.

[0167] Defining Polypeptides by Immunoreactivity

[0168] Because the polypeptides of the invention provide a variety of new polypeptide sequences (e.g., comprising HA and NA molecules), the polypeptides also provide new structural features which can be recognized, e.g., in immunological assays. The generation of antisera which specifically bind the polypeptides of the invention, as well as the polypeptides which are bound by such antisera, are features of the invention.

[0169] For example, the invention includes polypeptides (e.g., HA and NA molecules) that specifically bind to or that are specifically immunoreactive with an antibody or antisera generated against an immunogen comprising an amino acid sequence selected from one or more of the sequences given herein, etc. To eliminate cross-reactivity with other homologues, the antibody or antisera is subtracted with the HA and/or NA molecules found in public databases at the time of filing, e.g., the "control" polypeptide(s). Where the other control sequences correspond to a nucleic acid, a polypeptide encoded by the nucleic acid is generated and used for antibody/antisera subtraction purposes.

[0170] In one typical format, the immunoassay uses a polyclonal antiserum which was raised against one or more polypeptide comprising one or more of the sequences corresponding to the sequences herein, etc. or a substantial subsequence thereof (i.e., at least about 30% of the full length sequence provided). The set of potential polypeptide immunogens derived from the present sequences are collectively referred to below as "the immunogenic polypeptides." The resulting antisera is optionally selected to have low cross-reactivity against the control hemagglutinin and/or neuraminidase homologues and any such cross-reactivity is removed, e.g., by immunoabsorption, with one or more of the control hemagglutinin and neuraminidase homologues, prior to use of the polyclonal antiserum in the immunoassay.

[0171] In order to produce antisera for use in an immunoassay, one or more of the immunogenic polypeptides is produced and purified as described herein. For example, recombinant protein can be produced in a recombinant cell. An inbred strain of mice (used in this assay because results are more reproducible due to the virtual genetic identity of the mice) is immunized with the immunogenic protein(s) in combination with a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see, e.g., Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York, for a standard description of antibody generation, immunoassay formats and conditions that can be used to determine specific immunoreactivity). Additional references and discussion of antibodies is also found herein and can be applied here to defining polypeptides by immunoreactivity. Alternatively, one or more synthetic or recombinant polypeptide derived from the sequences disclosed herein is conjugated to a carrier protein and used as an immunogen.

[0172] Polyclonal sera are collected and titrated against the immunogenic polypeptide in an immunoassay, for example, a solid phase immunoassay with one or more of the immunogenic proteins immobilized on a solid support. Polyclonal antisera with a titer of 10^6 or greater are selected, pooled and subtracted with the control hemagglutinin and/or neuraminidase polypeptide(s) to produce subtracted pooled titrated polyclonal antisera.

[0173] The subtracted pooled titrated polyclonal antisera are tested for cross reactivity against the control homologue (s) in a comparative immunoassay. In this comparative assay, discriminatory binding conditions are determined for the subtracted titrated polyclonal antisera which result in at least about a 5-10 fold higher signal to noise ratio for binding of the titrated polyclonal antisera to the immunogenic polypeptides as compared to binding to the control homologues. That is, the stringency of the binding reaction is adjusted by the addition of non-specific competitors such as albumin or non-fat dry milk, and/or by adjusting salt conditions, temperature, and/or the like. These binding conditions are used in subsequent assays for determining whether a test polypeptide (a polypeptide being compared to the immunogenic polypeptides and/or the control polypeptides) is specifically bound by the pooled subtracted polyclonal antisera. In particular, test polypeptides which show at least a 2-5 $\times$ higher signal to noise ratio than the control receptor homologues under discriminatory binding conditions, and at least about a $\frac{1}{2}$ signal to noise ratio as compared to the immunogenic polypeptide(s), shares substantial structural similarity with the immunogenic polypeptide as compared to the known receptor, etc., and is, therefore a polypeptide of the invention.

[0174] In another example, immunoassays in the competitive binding format are used for detection of a test polypeptide. For example, as noted, cross-reacting antibodies are removed from the pooled antisera mixture by immunoabsorption with the control polypeptides. The immunogenic polypeptide(s) are then immobilized to a solid support which is exposed to the subtracted pooled antisera. Test proteins are added to the assay to compete for binding to the pooled subtracted antisera. The ability of the test protein(s) to compete for binding to the pooled subtracted antisera as compared to the immobilized protein(s) is compared to the ability of the immunogenic polypeptide(s) added to the assay to compete for binding (the immunogenic polypeptides compete effectively with the immobilized immunogenic polypeptides for binding to the pooled antisera). The percent cross-reactivity for the test proteins is calculated, using standard calculations.

[0175] In a parallel assay, the ability of the control protein(s) to compete for binding to the pooled subtracted antisera is optionally determined as compared to the ability of the immunogenic polypeptide(s) to compete for binding to the antisera. Again, the percent cross-reactivity for the control polypeptide(s) is calculated, using standard calculations. Where the percent cross-reactivity is at least 5-10× as high for the test polypeptides as compared to the control polypeptide(s) and/or where the binding of the test polypeptides is approximately in the range of the binding of the immunogenic polypeptides, the test polypeptides are said to specifically bind the pooled subtracted antisera.

[0176] In general, the immunoabsorbed and pooled antisera can be used in a competitive binding immunoassay as described herein to compare any test polypeptide to the immunogenic and/or control polypeptide(s). In order to make this comparison, the immunogenic, test and control polypeptides are each assayed at a wide range of concentrations and the amount of each polypeptide required to inhibit 50% of the binding of the subtracted antisera to, e.g., an immobilized control, test or immunogenic protein is determined using standard techniques. If the amount of the test polypeptide required for binding in the competitive assay is less than twice the amount of the immunogenic polypeptide that is required, then the test polypeptide is said to specifically bind to an antibody generated to the immunogenic protein, provided the amount is at least about 5-10× as high as for the control polypeptide.

[0177] As an additional determination of specificity, the pooled antisera is optionally fully immunosorbed with the immunogenic polypeptide(s) (rather than the control polypeptide(s)) until little or no binding of the resulting immunogenic polypeptide subtracted pooled antisera to the immunogenic polypeptide(s) used in the immunosorption is detectable. This fully immunosorbed antisera is then tested for reactivity with the test polypeptide. If little or no reactivity is observed (i.e., no more than 2× the signal to noise ratio observed for binding of the fully immunosorbed antisera to the immunogenic polypeptide), then the test polypeptide is specifically bound by the antisera elicited by the immunogenic protein.

Nucleic Acid and Polypeptide Sequence Variants

[0178] As described herein, the invention provides for nucleic acid polynucleotide sequences and polypeptide amino acid sequences, e.g., hemagglutinin and neuraminidase sequences, and, e.g., compositions and methods comprising said sequences. Examples of said sequences are dis-

closed herein. However, one of skill in the art will appreciate that the invention is not necessarily limited to those sequences disclosed herein and that the present invention also provides many related and unrelated sequences with the functions described herein, e.g., encoding a HA and/or a NA molecule.

[0179] One of skill will also appreciate that many variants of the disclosed sequences are included in the invention. For example, conservative variations of the disclosed sequences that yield a functionally identical sequence are included in the invention. Variants of the nucleic acid polynucleotide sequences, wherein the variants hybridize to at least one disclosed sequence, are considered to be included in the invention. Unique subsequences of the sequences disclosed herein, as determined by, e.g., standard sequence comparison techniques, are also included in the invention.

[0180] Silent Variations

[0181] Due to the degeneracy of the genetic code, any of a variety of nucleic acid sequences encoding polypeptides and/or viruses of the invention are optionally produced, some which can bear lower levels of sequence identity to the HA and NA nucleic acid and polypeptide sequences herein. The following provides a typical codon table specifying the genetic code, found in many biology and biochemistry texts.

TABLE 1

Codon Table			
Amino acids		Codon	
Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	UGC UGU
Aspartic acid	Asp	D	GAC GAU
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	UUC UUU
Glycine	Gly	G	GGA GGC GGG GGU
Histidine	His	H	CAC CAU
Isoleucine	Ile	I	AUA AUC AUU
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	UUA UUG CUA CUC CUG CUU
Methionine	Met	M	AUG
Asparagine	Asn	N	AAC AAU
Proline	Pro	P	CCA CCC CCG CCU
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGU
Serine	Ser	S	AGC AGU UCA UCC UCG UCU
Threonine	Thr	T	ACA ACC ACG ACU
Valine	Val	V	GUA GUC GUG GUU
Tryptophan	Trp	W	UGG
Tyrosine	Tyr	Y	UAC UAU

[0182] The codon table shows that many amino acids are encoded by more than one codon. For example, the codons AGA, AGG, CGA, CGC, CGG, and CGU all encode the

amino acid arginine. Thus, at every position in the nucleic acids of the invention where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described above without altering the encoded polypeptide. It is understood that U in an RNA sequence corresponds to T in a DNA sequence.

[0183] Such “silent variations” are one species of “conservatively modified variations,” discussed below. One of skill will recognize that each codon in a nucleic acid (except ATG, which is ordinarily the only codon for methionine, and TTG, which is ordinarily the only codon for tryptophan) can be modified by standard techniques to encode a functionally identical polypeptide. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in any described sequence. The invention, therefore, explicitly provides each and every possible variation of a nucleic acid sequence encoding a polypeptide of the invention that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code (e.g., as set forth in Table 1, or as is commonly available in the art) as applied to the nucleic acid sequence encoding a hemagglutinin or a neuraminidase polypeptide of the invention. All such variations of every nucleic acid herein are specifically provided and described by consideration of the sequence in combination with the genetic code. One of skill is fully able to make these silent substitutions using the methods herein.

[0184] Conservative Variations

[0185] Owing to the degeneracy of the genetic code, “silent substitutions” (i.e., substitutions in a nucleic acid sequence which do not result in an alteration in an encoded polypeptide) are an implied feature of every nucleic acid sequence of the invention which encodes an amino acid. Similarly, “conservative amino acid substitutions,” in one or a few amino acids in an amino acid sequence are substituted with different amino acids with highly similar properties, are also readily identified as being highly similar to a disclosed construct such as those herein. Such conservative variations of each disclosed sequence are a feature of the present invention.

[0186] “Conservative variation” of a particular nucleic acid sequence refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or, where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences, see, Table 2 below. One of skill will recognize that individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 4%, 3%, 2% or 1%) in an encoded sequence are “conservatively modified variations” where the alterations result in the deletion of an amino acid, addition of an amino acid, or substitution of an amino acid with a chemically similar amino acid. Thus, “conservative variations” of a listed polypeptide sequence of the present invention include substitutions of a small percentage, typically less than 5%, more typically less than 4%, 3%, 2% or 1%, of the amino acids of the polypeptide sequence, with a conservatively selected amino acid of the same conservative substitution group. Finally, the addition of sequences which do not alter the encoded activity of a nucleic acid molecule, such as the addition of a non-functional sequence, is a conservative variation of the basic nucleic acid.

TABLE 2

Conservative Substitution Groups				
1	Alanine (A)	Serine (S)	Threonine (T)	
2	Aspartic acid (D)	Glutamic acid (E)		
3	Asparagine (N)	Glutamine (Q)		
4	Arginine (R)	Lysine (K)		
5	Isoleucine (I)	Leucine (L)	Methionine (M)	Valine (V)
6	Phenylalanine (F)	Tyrosine (Y)	Tryptophan (W)	

[0187] Unique Polypeptide and Polynucleotide Subsequences

[0188] In one aspect, the invention provides a nucleic acid which comprises a unique subsequence in a nucleic acid selected from the sequence of HA and NA molecules disclosed herein (e.g., SEQ ID NO:1-34). The unique subsequence is unique as compared to a nucleic acids corresponding to nucleic acids such as, e.g., those found in GenBank or other similar public databases at the time of filing (e.g., other known or characterized hemagglutinin and/or neuraminidase nucleic acid molecules). Alignment can be performed using, e.g., BLAST set to default parameters. Any unique subsequence is useful, e.g., as a probe to identify the nucleic acids of the invention. See, above.

[0189] Similarly, the invention includes a polypeptide (e.g., from SEQ ID NO:35 through 68) which comprises a unique subsequence in a polypeptide selected from the sequence of HA and NA molecules disclosed herein. Here, the unique subsequence is unique as compared to a polypeptide corresponding to, e.g., the amino acid corresponding to polynucleotide sequences found in, e.g., GenBank or other similar public databases at the time of filing.

[0190] The invention also provides for target nucleic acids which hybridize under stringent conditions to a unique coding oligonucleotide which encodes a unique subsequence in a polypeptide selected from the sequences of HA and NA molecules of the invention wherein the unique subsequence is unique as compared to a polypeptide corresponding to any of the control polypeptides (sequences of, e.g., the nucleic acids corresponding to those found in, e.g., GenBank or other similar public databases at the time of filing). Unique sequences are determined as noted above.

[0191] Sequence Comparison Identity and Homology

[0192] The terms “identical” or percent “identity,” in the context of two or more nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms described below (or other algorithms available to persons of skill) or by visual inspection.

[0193] The phrase “substantially identical,” in the context of two nucleic acids or polypeptides (e.g., DNAs encoding a HA or NA molecule, or the amino acid sequence of a HA or NA molecule) refers to two or more sequences or subsequences that have at least about 90%, preferably 91%, most preferably 92%, 93%, 94%, 95%, 96%, 97%, 98%, 98.5%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or more nucleotide or amino acid residue identity, when compared and aligned for maximum correspondence, as measured using a sequence comparison algorithm or by visual inspection. Such “substantially identical” sequences are typically considered to be “homologous,” with-

out reference to actual ancestry. Preferably, "substantial identity" exists over a region of the amino acid sequences that is at least about 200 residues in length, more preferably over a region of at least about 250 residues, and most preferably the sequences are substantially identical over at least about 300 residues, 350 residues, 400 residues, 425 residues, 450 residues, 475 residues, 480 residues, 490 residues, 495 residues, 499 residues, 500 residues, 502 residues, 559 residues, 565 residues, or 566 residues, or over the full length of the two sequences to be compared when the amino acids are hemagglutinin or hemagglutinin fragments or which is substantially identical over at least about 350 amino acids; over at least about 400 amino acids; over at least about over at least about 436 amino acids, over at least about 450 amino acids; over at least about 451 amino acids; over at least about 465 amino acids; over at least about 466 amino acids; over at least about 469 amino acids; over at least about 470 amino acids; or over at least about 566 amino acids contiguous when the amino acid is neuraminidase.

[0194] For sequence comparison and homology determination, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0195] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv Appl Math* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J Mol Biol* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc Natl Acad Sci USA* 85:2444 (1988), by computerized implementations of algorithms such as GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis., or by visual inspection (see generally, Ausubel et al., *supra*).

[0196] One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J Mol Biol* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (see, Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes

to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see, Henikoff & Henikoff (1989) *Proc Natl Acad Sci USA* 89:10915).

[0197] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc Natl Acad Sci USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0198] Another example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle (1987) *J. Mol. Evol.* 35:351-360. The method used is similar to the method described by Higgins & Sharp (1989) *CABIOS* 5:151-153. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster can then be aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences can be aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program can also be used to plot a dendrogram or tree representation of clustering relationships. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison.

[0199] An additional example of an algorithm that is suitable for multiple DNA, or amino acid, sequence alignments is the CLUSTALW program (Thompson, J. D. et al. (1994) *Nucl. Acids. Res.* 22: 4673-4680). CLUSTALW performs multiple pairwise comparisons between groups of sequences and assembles them into a multiple alignment based on homology. Gap open and Gap extension penalties can be, e.g., 10 and 0.05 respectively. For amino acid alignments, the BLOSUM algorithm can be used as a protein weight matrix. See, e.g., Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919.

Digital Systems

[0200] The present invention provides digital systems, e.g., computers, computer readable media and integrated systems comprising character strings corresponding to the sequence information herein for the nucleic acids and isolated or recombinant polypeptides herein, including, e.g., the

sequences shown herein, and the various silent substitutions and conservative substitutions thereof. Integrated systems can further include, e.g., gene synthesis equipment for making genes corresponding to the character strings.

[0201] Various methods known in the art can be used to detect homology or similarity between different character strings (see above), or can be used to perform other desirable functions such as to control output files, provide the basis for making presentations of information including the sequences and the like. Examples include BLAST, discussed supra. Computer systems of the invention can include such programs, e.g., in conjunction with one or more data file or data base comprising a sequence as noted herein.

[0202] Thus, different types of homology and similarity of various stringency and length between various HA or NA sequences or fragments, etc. can be detected and recognized in the integrated systems herein. For example, many homology determination methods have been designed for comparative analysis of sequences of biopolymers, for spell-checking in word processing, and for data retrieval from various databases. With an understanding of double-helix pair-wise complement interactions among four principal nucleobases in natural polynucleotides, models that simulate annealing of complementary homologous polynucleotide strings can also be used as a foundation of sequence alignment or other operations typically performed on the character strings corresponding to the sequences herein (e.g., word-processing manipulations, construction of figures comprising sequence or subsequence character strings, output tables, etc.).

[0203] Thus, standard desktop applications such as word processing software (e.g., Microsoft Word™ or Corel WordPerfect™) and database software (e.g., spreadsheet software such as Microsoft Excel™, Corel Quattro Pro™, or database programs such as Microsoft Access™, Paradox™, GeneWorks™, or MacVector™ or other similar programs) can be adapted to the present invention by inputting a character string corresponding to one or more polynucleotides and polypeptides of the invention (either nucleic acids or proteins, or both). For example, a system of the invention can include the foregoing software having the appropriate character string information, e.g., used in conjunction with a user interface (e.g., a GUI in a standard operating system such as a Windows, Macintosh or LINUX system) to manipulate strings of characters corresponding to the sequences herein. As noted, specialized alignment programs such as BLAST can also be incorporated into the systems of the invention for alignment of nucleic acids or proteins (or corresponding character strings).

[0204] Systems in the present invention typically include a digital computer with data sets entered into the software system comprising any of the sequences herein. The computer can be, e.g., a PC (Intel x86 or Pentium chip-compatible DOS™, OS2™, WINDOWS™, WINDOWSNT™, WINDOWS95™, WINDOWS2000™, WINDOWS98™, LINUX based machine, a MACINTOSH™, Power PC, or a UNIX based (e.g., SUN™ work station) machine) or other commercially available computer that is known to one of skill. Software for aligning or otherwise manipulating sequences is available, or can easily be constructed by one of skill using a standard programming language such as Visualbasic, PERL, Fortran, Basic, Java, or the like.

[0205] Any controller or computer optionally includes a monitor which is often a cathode ray tube ("CRT") display, a flat panel display (e.g., active matrix liquid crystal display,

liquid crystal display), or others. Computer circuitry is often placed in a box which includes numerous integrated circuit chips, such as a microprocessor, memory, interface circuits, and others. The box also optionally includes a hard disk drive, a floppy disk drive, a high capacity removable drive such as a writeable CD-ROM, and other common peripheral elements. Inputting devices such as a keyboard or mouse optionally provide for input from a user and for user selection of sequences to be compared or otherwise manipulated in the relevant computer system.

[0206] The computer typically includes appropriate software for receiving user instructions, either in the form of user input into a set parameter fields, e.g., in a GUI, or in the form of preprogrammed instructions, e.g., preprogrammed for a variety of different specific operations. The software then converts these instructions to appropriate language for instructing the operation, e.g., of appropriate mechanisms or transport controllers to carry out the desired operation. The software can also include output elements for controlling nucleic acid synthesis (e.g., based upon a sequence or an alignment of sequences herein), comparisons of samples for differential gene expression, or other operations.

Kits and Reagents

[0207] The present invention is optionally provided to a user as a kit. For example, a kit of the invention contains one or more nucleic acid, polypeptide, antibody, or cell line described herein (e.g., comprising, or with, a HA and/or NA molecule of the invention). The kit can contain a diagnostic nucleic acid or polypeptide, e.g., antibody, probe set, e.g., as a cDNA micro-array packaged in a suitable container, or other nucleic acid such as one or more expression vector. The kit typically further comprises, one or more additional reagents, e.g., substrates, labels, primers, for labeling expression products, tubes and/or other accessories, reagents for collecting samples, buffers, hybridization chambers, cover slips, etc. The kit optionally further comprises an instruction set or user manual detailing preferred methods of using the kit components for discovery or application of diagnostic sets, etc.

[0208] When used according to the instructions, the kit can be used, e.g., for evaluating a disease state or condition, for evaluating effects of a pharmaceutical agent or other treatment intervention on progression of a disease state or condition in a cell or organism, or for use as a vaccine, etc.

[0209] In an additional aspect, the present invention provides system kits embodying the methods, composition, systems and apparatus herein. System kits of the invention optionally comprise one or more of the following: (1) an apparatus, system, system component or apparatus component; (2) instructions for practicing methods described herein, and/or for operating the apparatus or apparatus components herein and/or for using the compositions herein. In a further aspect, the present invention provides for the use of any apparatus, apparatus component, composition or kit herein, for the practice of any method or assay herein, and/or for the use of any apparatus or kit to practice any assay or method herein.

[0210] Additionally, the kits can include one or more translation system as noted above (e.g., a cell) with appropriate packaging material, containers for holding the components of the kit, instructional materials for practicing the methods herein and/or the like. Similarly, products of the translation systems (e.g., proteins such as HA and/or NA molecules) can

be provided in kit form, e.g., with containers for holding the components of the kit, instructional materials for practicing the methods herein and/or the like. Furthermore, the kits can comprise various vaccines (e.g., produced through plasmid rescue protocols) such as live attenuated vaccine (e.g., Flu-Mist™) comprising the HA and/or NA sequences herein.

[0211] To facilitate use of the methods and compositions of the invention, any of the vaccine components and/or compositions, e.g., reasserted virus in allantoic fluid, etc., and additional components, such as, buffer, cells, culture medium, useful for packaging and infection of influenza viruses for experimental or therapeutic vaccine purposes, can be packaged in the form of a kit. Typically, the kit contains, in addition to the above components, additional materials which can

include, e.g., instructions for performing the methods of the invention, packaging material, and a container.

[0212] While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. For example, all the techniques and apparatus described above may be used in various combinations. All publications, patents, patent applications, or other documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, or other document were individually indicated to be incorporated by reference for all purposes.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 68

<210> SEQ ID NO 1

<211> LENGTH: 1745

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 1

caggggataa ttctattaac catgaagact atcattgctt tgagctacat tttatgtctg	60
gttttcgctc aaaaacttcc cggaaatgac aacagcacag caacgctgtg cctgggacat	120
catgcagtgc caaacggaac gctagtgaac acaatcacga atgatcaaat tgaagtgact	180
aatgctactg agttgggtca gagttcctca acaggtagaa tatgcggcag tcctcaccga	240
atccttgatg gaaaaaactg cacactgata gatgctctat tgggagaccc tcattgtgat	300
ggcttcocaa ataaggaatg ggaccttttt gttgaacgca gcaaagctta cagcaactgt	360
taccttatg atgtgccgga ttatgctccc cttaggtcac tagttgcctc atcaggcacc	420
ctggagttaa tcaatgaaga cttcaattgg actggagtcg ctcaggatgg gggaagctat	480
gcttgcaaaa gaggatctgt taacagtttc tttagtagat tgaattgggt gcacaaatta	540
gaatacaaat atccagcgt gaacgtgact atgccaaaca atggcaaatt tgacaaattg	600
tacatttggg ggggttccca ccgagcacg gacagtgcac aaaccagcct atatgttcga	660
gcatcaggga gagtcacagt ctctacaaa agaagccaac aaactgtaac cccgaatatc	720
gggtctagac cctgggtaag gggtcagtcc agtagaataa gcatctattg gacaatagta	780
aaaccgggag acatactttt gattaatagc acagggaatc taattgctcc tcgggggttac	840
ttcaaaatc gaaatgggaa aagctcaata atgaggtcag atgcacccat tggcaactgc	900
agttctgaat gcatcactcc aaatggaagc attcccaatg acaaaccttt tcaaaatgta	960
aacagaatca catatggggc ctgccccaga tatgttaagc aaaacactct gaaattggca	1020
acagggatgc ggaatgtacc agagaaacaa actagaggca tattcggcgc aatcgcaggt	1080
ttcatagaaa atggttgga gggaaatgga gacggttgt acggtttcag gcatcaaaat	1140
tctgagggca caggacaagc agcagatctt aaaagcactc aagcagcaat cgaccaaatc	1200
aacgggaaac tgaataggtt aatcgagaaa acgaacgaga aattccatca aatcgaaaaa	1260
gaattctcag aagtagaagg gagaattcag gacctcgaga aatatgttga agacactaaa	1320
atagatctct ggtcttacia cgcggagctt cttgttgccc tggagaacca acatacaatt	1380

-continued

```

gatctaactg actcagaaat gaacaaactg tttgaaaaaa caaggaagca actgagggaa 1440
aatgctgagg acatggggcaa tgggtgcttc aaaatatacc acaaatgtga caatgcctgc 1500
ataggggtcaa tcagaaatgg aacttatgac catgatgtat acagagacga agcattaaac 1560
aaccgggtcc agatcaaagg tgttgagctg aagtcaggat acaagattg gatcctatgg 1620
atttcctttg ccatatcatg ctttttgctt tgtgttggtt tgctgggggtt catcatgtgg 1680
gcctgccaaa aaggcaacat taggtgcaac atttgcatth gagtgcatta attaaaaaca 1740
ccctg 1745

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1429
<212> TYPE: DNA
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 2

```

```

aaagataata acaattggct ctgtttctct cactattgcc acaatatgct tccttatgca 60
aattgccatc ctggtaacta ctgtaacatt gcacttcaag caatatgagt gcaactcccc 120
cccaaacaac caagtaatgc tgtgtgaacc aacaataata gaaagaaaca taacagagat 180
agtgatatctg accaacacca ccatagagaa agaaatatgc cccaaactag cagaatacag 240
aaattgggtca aagccgcaat gtaaaattac aggatttgca cctttttcta aggacaattc 300
aattcggtct tcagctgggt gagacatttg ggtgacaaga gaaccttatg tgctatgcga 360
tcctggcaag tgttatcaat ttgcccttgg acagggaaca aactaaaca acaggcactc 420
aaatgacaca gtacatgata ggacccctta tcgaacctta ttgatgaatg agttgggtgt 480
tccatttcat ttgggaacca agcaagtgtg catagcatgg tccagctcaa gttgtcacga 540
tggaagagca tggctgcatg tttgtgtaac tgggcatgat gaaatgcaa ctgctagctt 600
catttacgat gggagcttg tagatagtat tggttcatgg tccaaaaata tcctcaggac 660
ccaggagtgc gaatgcgttt gtatcaatgg aacttgtaaa gtagtaatga ctgatggaag 720
tgcttcagaa agagctgata ctaaaatact attcattgaa gaggggaaaa tcgttcatat 780
tagccattg tcaggaagtg ctcagcatgt cgaggagtgc tcctgttata ctcgatatac 840
tggtgtcaga tgtgtctgca gagacaactg gaaaggctcc aataggccca tcgtagatat 900
aaatgtgaaa gattatagca ttgtttccag ttatgtgtgc tcaggacttg ttggagacac 960
accagaaaaa aacgacagct ccagcagtag ctattgccgg aatcctaaca atgagaaagg 1020
gagtcatgga gtgaaaggct gggcctttga tgatggaaat gacgtgtgga tgggaagaac 1080
gatcagcgag gagttacgct caggttatga aaccttcaa gtcattggag gctgggtcaa 1140
acctaactcc aaattgcaga taaataggca agtcatagtt gacagaggta ataggtccgg 1200
ttattctggt attttctctg ttgaaggcaa aagctgcac aatcggtgct tttatgtgga 1260
gttgataagg ggaaggaaac aggaactga agtctggtgg acctcaaaca gtattgttgt 1320
gtttgtggc acctcaggta catatggaac aggtcatggt ccctgatggg gcggacatca 1380
atctcatgcc tatataagct ttcgcaattt tagaaaaaaa ctctctgtt 1429

```

```

<210> SEQ ID NO 3
<211> LENGTH: 1755
<212> TYPE: DNA
<213> ORGANISM: Influenza A virus

```

-continued

<400> SEQUENCE: 3

```

agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg agctacattt    60
tatgtctggt ttctgctcaa aaacttcccg gaaatgacaa cagcacagca acgctgtgcc    120
tgggacacca tgcagtgcc aacggaacgc tagtgaaaac aatcacgaat gatcaaattg    180
aagtactaa tgctactgag ctggttcaga gttccccaac aggtagaata tgcgacagcc    240
ctcacccaat ccttgatgga aagaactgca cactgataga tgctctattg ggagaccctc    300
attgtgatgg ctccaaaat aaggaatggg acctttttgt tgaacgcagc aaagcttaca    360
gcaactgtta cccttatgat gtgccggatt atgcctccct taggtcacta gttgcctcat    420
caggcaccct ggagtttacc aacgaaaact tcaattggac tggagtcgct caggatggga    480
aaagctatgc ttgcaaaagg ggatctgtta acagtttctt tagtagattg aattggttgc    540
acaaattaga atacaaatat ccagcgctga acgtgactat gccaaacaat ggcaaatattg    600
acaaattgta catttggggg gttcaccacc cgagcacgga cagtgtccaa accagcctat    660
atgtccgagc atcaggggaga gtcacagtct ctacccaaag aagccaacaa actgtaatcc    720
cggatatcgg gtatagacca tgggtaaggg gtcagtccag tagaataagc atctattgga    780
caatagttaa accggggagac atacttttga ttaatagcac agggaatcta attgctctc    840
ggggttacct caaaatcaga aatgggaaaa gctcaataat gaggtcagat gcacccattg    900
gcaactgcag ttctgaatgc atcactccaa atggaagcat tcccaatgac aaaccttttc    960
aaaatgtaaa caggatcaca tatggggcct gcccagata tgtaagcaa aacactctga   1020
aattggcaac agggatgcgg aatgtaccag agaacaacac tagaggcata ttcggcgcaa   1080
tcgcaggttt catagaaat ggttgggagg gaatggtaga cggttggtac ggtttcaggc   1140
atcaaaattc tgagggcaca ggacaagctg cagatcttaa aagcactcaa gcagcaatcg   1200
accaaataca cgggaaactg aatagggttag tcgagaaaac gaacgagaaa ttccatcaaa   1260
tcgaaaaaga attctcagaa gtagaaggga gaattcagga cctcgagaaa tatgttgaag   1320
acactaaaat agatctctgg tcttacaatg cggaacttct tgttgctctg gagaaccaac   1380
atacaattga tctaactgac tcagaaatga acaaaactgtt tgaagaaca aggaagcaac   1440
tgagggaaaa tgctgaggac atgggcaatg gttgtttcaa aatataccac aaatgtgaca   1500
atgcctgcac agggtaacac agaaatggaa cttatgacca tgatgtatac agagacgaag   1560
cattaacaaa ccggttcacg atcaaaggtg ttgagctgaa gtcaggatac aaagattgga   1620
ttctatggat ttcttttgc atatcgtgct ttttgctttg tgttgttttg cttgggttca   1680
tcatgtgggc ctgcaaaaaa ggcaacatta ggtgcaacat ttgcatttga gtgcattaat   1740
taaaaacacc cttgt                                     1755

```

<210> SEQ ID NO 4

<211> LENGTH: 1354

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 4

```

gaaaaatgaat ccaaatacaa agataataac aattggctct gttctctca ctattgccac    60
aatatgcttc cttatgcaaa ttgccatcct ggtaactact gtaacattgc atttcaagca   120
atatgagtgc aactcccccc caaacacca agtaatgctg tgtgaaccaa caataataga   180

```

-continued

```

aagaaacata acagagatag tgtatctgac caacaccacc atagagaaag aaatatgccc 240
caaaactagca gaatacagaa attggtcaaa gccgcaatgt aaaattacag gatttgcacc 300
tttttctaag gacaattcaa ttcggttttc cgctggtgga gacatttggg tgacaagaga 360
accttatgtg tcatgcgac ctggcaagtg ttatcaattt gccctcggac agggaacaac 420
actaaacaac aggcattcaa atgacacagt acatgatagg accccttacc gaaccctatt 480
gatgaatgag ttgggtgttc catttcattt gggaaccaag caagtgtgca tagcatggtc 540
cagctcaagt tgtcacgatg gaaaagcatg gctgcatgtt tgtgtaactg ggcatgatga 600
aaatgcaact gctagcttca ttacgatgg gaggtttgta gatagtattg gttcatggtc 660
caaaaatacc ctacgagacc aggagtcgga atgcgtttgt atcaatggaa cttgtacagt 720
agtaatgact gatggaagtg cttcagaaag agctgatact aaaatactat tcattgaaga 780
ggggaaaatc gttcatatta gccattgtc aggaagtgtc cagcatgtcg aggagtgtc 840
ctgttatcct cgatatcctg gtgtcagatg tgtctgcaga gacaactgga aaggctccaa 900
taggccatc gtagatataa atgtgaaaga ttatagcatt gtttccagtt atgtgtgctc 960
aggacttgtt ggagacacac ccagaaaaaa cgacagctcc agcagtagct attgctggaa 1020
tcctaacaat gagaaagggg gtcattggagt gaaaggctgg gcctttgatg atggaaatga 1080
cgtgtggatg ggaagaacga tcagcgagga gttacgtcca ggttatgaaa ccttcaaagt 1140
cattggaggc tgggtccaaac ctaactccaa attgcagata aataggcaag tcatagttga 1200
cagaggtaat aggtccggtt attctggtat tttctctgtt gaaggcaaaa gctgcatcaa 1260
tcggtgcttt tatgtggagt tgataagggg aaggaaacag gaaactgaag tctggtggac 1320
ctcaaacagt attgttgtgt tttgtggcac ttca 1354

```

```

<210> SEQ ID NO 5
<211> LENGTH: 1762
<212> TYPE: DNA
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 5

```

```

agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg agctacattt 60
tatgtctggt ttctgctcaa aaacttccc gaaatgacaa cagcacggca acgtgtgccc 120
tgggacacca tgcagtgcc aacggaacgc tagtgaaaac aatcacgaat gaccaaattg 180
aagtgactaa tgctactgag ctgggttcaga gttcctcaac aggtagaata tgcgacagtc 240
ctcacggaat ccttgatgga aaaaactgca cactgataga tgctctattg ggagaccctc 300
attgtgatgg cttccaaaat aaggaatggg accttttgt tgaacgcagc aaagcttaca 360
gcaactgtta cccttatgat gtgccgatt atgcttcct taggtcacta gttgcctcat 420
ccggcaccct ggagtttacc aatgaaggct tcaattggac tggagtcgct caggatggaa 480
caagctatgc ttgcaaaagg ggatctgtta aaagtctct tagtagattg aattggttgc 540
acaaattaga atacaaatat ccagcactga acgtgactat gccaaacaat gacaaatttg 600
acaaattgta catttggggg gttcaccacc cgagtacgga cagtgaacaa accagcatat 660
atgttcaagc atcagggaga gtcacagtct ctacccaaag aagccaacaa actgtaatcc 720
cgaatatcgg gtctagaccc tgggtaaggg ggatctccag cagaataagc atctattgga 780
caatagtaaa accgggagac atacttttga ttaacagcac agggaatcta attgctcctc 840

```

-continued

```

ggggttactt caaaatcacga agtgggaaaa gctcaataat gaggtcagat gcacccattg    900
gcaactgcaa ttctgaatgc atcactccaa atggaagcat tcccaatgac aaaccttttc    960
aaaatgtaaa caggatcaca tatggggcct gtcccagata tgtaagcaa aacactctga    1020
aattggcaac agggatgcgg aatgtaccag agaaacaaac tagaggcata ttcggcgcaa    1080
tcgcaggttt catagaaaat ggttgggagg gaatggtaga cggttggtac ggtttcaggc    1140
atcaaaattc tgagggcaca ggacaagcag cagatcttaa aagcactcaa gcagcaatca    1200
accaaatcaa cgggaaactg aatagggttaa tcgagaaaac gaacgagaaa ttccatcaaa    1260
tcgaaaaaga attctcagaa gtagaaggga gaattcagga cctcgagaaa tatgttgaag    1320
acactaaaat agatctctgg tcttacaacg cggagcttct tgttgccctg gagaaccaac    1380
atacaattga tctaactgac tcagaaatga acaaaactgtt tgaagaaca aggaagcaac    1440
tgagggaaaa tgctgaggac atgggcaatg gttgcttcaa aatataccac aaatgtgaca    1500
atgcctgcat agggtaactc agaaatggaa cttatgacca tgatgtatac agagacgaag    1560
cattaacaaa ccggttccag atcaaaggty ttgagctgaa gtcaggatac aaagattgga    1620
tcctatggat ttcctttgcc atatcatgct ttttgctttg tgtgttctg ctggggttca    1680
tcatgtgggc ctgcaaaaaa ggcaacatta ggtgcaacat ttgcatttga gtgcattaat    1740
taaaaacacc cttgtttcta ct                                         1762

```

<210> SEQ ID NO 6

<211> LENGTH: 1451

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 6

```

agcaaaagca ggagtgaaaa tgaatccaaa tcaaaagata ataactattg gctctgtttc    60
tctcactatt gccacaatat gcttccttat gcaaatgcc atcctggtaa ctactgtaac    120
attacatttc aagcaataty aatgcaactc ccccccaaac aaccaagtaa tgctgtgtga    180
accaacaata atagaaagaa acataacaga gatagtgtat ctgaccaaca ccaccataga    240
gaaggaaata tgcccaaac tagcagaata cagaaattgg tcaagccgc aatgtaaaat    300
tacaggattt gcaccttttt ctaaggacaa ttcaattcgg ctttccgctg gtggggacat    360
ttgggtgaca agagaacctt atgtgtcatg cgatcctgac aagtgttatc aatttgccct    420
tggaaggga acaacactaa acaacaggca ttcaaagac acagtacatg ataggacccc    480
ttatcgaacc ctattgatga atgagttggg tgttccattt catttgggaa ccaagcaagt    540
gtgcatagca tgggtccagt caagttgtca cgatggaaaa gcatggctgc atgttttgtt    600
aactgggcat gatgaaaatg caactgctag cttcatttac gatgggaggc ttgtagatag    660
tattggttca tgggtcaaaa aaatcctcag gaccaggag tcggaatgcy tttgtatcaa    720
tggaacttgt acagtagtaa tgactgatgg aagtgcctca ggaagagctg atactaaaat    780
actattcatt gaagagggga aaatcgttca tattagccca ttgtcaggaa gtgctcagca    840
tgtcgaggag tgctcctgtt atcctcgata ttctggtgtc agatgtgtct gcagagacaa    900
ctggaaaggc tccaataggc ccatcgtaga tataaatgtg aaagattata gcattgtttc    960
cagttatgtg tgctcaggac ttgttgaga cacaccaga aaaaacgaca gctccagcag    1020
tagccattgc ctgaatccta acaatgagga aggggggtcat ggagtgaag gctgggcctt    1080

```

-continued

tgatgatgga aatgacgtgt ggatgggaag aacgatcagc gagaagttac gctcagggtta	1140
tgaaaccttc aaagtcattg gaggctggtc caaacctaac tccaaattgc agataaatag	1200
acaagtcata gttgacagag gtaataggtc cgggtattct ggtattttct ctgttgaagg	1260
caaaagctgc atcaatcggg gcttttatgt ggagttgata aggggaagga aacaggaaac	1320
tgaagctctg tggacctcaa acagtattgt tgtgttttgt ggcacctcag gtacatatgg	1380
aacaggctca tggcctgatg gggcggacat caatctcatg cctatataag ctttcgcaat	1440
tttagaaaaa a	1451

<210> SEQ ID NO 7

<211> LENGTH: 1762

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 7

agcaaaagca ggggataatt ctattaacca tgaagactat cattgctttg agctacattt	60
tatgtctggt ttctgctcaa aaaattcccg gaaatgacaa cagcacggca acgctgtgcc	120
tgggacacca tgcagtgcc aacggaacgc tagtgaaaac aatcacgaat gaccaaattg	180
aagtgactaa tgctactgag ctggttcaga gttcctcaac aggtagaata tgcgacagtc	240
ctcaccgaat ccttgatgga gaaaactgca cactgataga tgctctattg ggagaccctc	300
attgtgatgg cttccaaat aaggaatggg acctttttgt tgaacgcagc aaagcctaca	360
gcaactgtta cccttatgat gtgccggatt atgcctcct taggtcacta gttgcctcat	420
ccggcaccct ggagttaac aatgaaagct tcaattggac tggagtcgct cagaatggaa	480
caagctatgc ttgcaaaagg agttctatta aaagtttctt tagtagattg aattggttgc	540
accaattaaa atacaaatat ccagcactga acgtgactat gccaaacaat gacaaatttg	600
acaaattgta catttggggg gttcaccacc cgagtacgga cagtgaacaa accagcatat	660
atgetcaagc atcaggggaga gtcacagtct ccacaaaag aagccaacaa actgtaatcc	720
cgaatatcgg atctagaccc tgggtaaggg gtatctccag cagaataagc atccattgga	780
caatagtaaa accgggagac atacttttga ttaacagcac agggaatcta attgctctc	840
ggggttactt caaaatcga agtgggaaaa gctcaataat gaggtcagat gcaaccattg	900
gcaaattgcaa ttctgaatgc atcactccaa atggaagcat tcccaatgac aaaccatttc	960
aaaaatgtaa caggatcaca tatggggcct gtcccagata tgtaagcaa aacactctga	1020
aattggcaac agggatgcgg aatgtaccag agaaacaaac tagaggcata tcggcgcaa	1080
tcgcaggttt catagaaaat ggttgggagg gaatggtaga cggttggtac ggtttcaggc	1140
atcaaaatc tgagggcaca ggacaagcag cagatcttaa aagcactcaa gcagcaatca	1200
accaaatcaa cgggaaactg aataggttaa tcgagaaaac gaacgagaaa ttccatcaaa	1260
ttgaaaaaga attctcagaa gtagaaggga gaattcagga cctcgagaaa tatgttgagg	1320
acactaaaat agatctctgg tcgtacaacg cggagcttct tgttgccctg gagaaccaac	1380
atacaattga tctaactgac tcagaaatga acaaaactgtt tgaagaaca aggaagcaac	1440
tgagggaaaa tgctgaggat atgggcaatg gttgtttcaa aatataccac aaatgtgaca	1500
atgctgcat agggccaatc agaaatggaa cttatgacca tgatgtatac agagacgaag	1560
cattaacaaa ccggttcag atcaaagggt ttgagctgaa gtcaggatac aaagattgga	1620

-continued

tcctatggat ttcctttgcc atatcatgtt ttttgctttg tgttgttttg ctggggttca	1680
tcatgtgggc ctgccaaaaa ggcaacatta ggtgcaacat ttgcatttga gtgcattaat	1740
taaaaacacc ctgttttcta ct	1762

<210> SEQ ID NO 8
 <211> LENGTH: 1467
 <212> TYPE: DNA
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 8

agcaaaaagca ggagtaaaga tgaatccaaa tcaaaagata ataacgattg gctctgtttc	60
tctcactatt gccacaatat gcttcottat gcaaattgcc atcctggtaa ctactgtaac	120
attgcatttc aagcaatatg aatgcagctc tcccccaaac aaccaagtaa tgctgtgtga	180
accaacaata atagaaagaa acataacaga gatagtgtat ctgaccaaca ccaccataga	240
gaaggaaata tgcccaaac tagcagaata cagaaattgg tcaaagccac aatgtaaaat	300
tacaggattt gcaccttttt ctaaggacaa ttcaattcgg ctttcgctg gtggggacat	360
ttgggtgaca agggaacctt atgtgtcgtg cgatcctgac aagtgttate aatttgcct	420
tggacaggga acaacactaa acaacaggga ttcaaatgac acagtacatg ataggacccc	480
ttatcgaaac ctattgatga atgagttggg tgttcattt catttgggaa ccaagcaagt	540
gtgcatagca tgggtccagct caagtgtca cgatggaaaa gcattggctgc atgtttgtgt	600
aactgggcat gatgaaaatg caactgctag cttcatttac gatgggaggc ttgtagatag	660
tattggttca tgggtccaaa aaatcctcag gaccaggag tcggaatgcg tttgtatcaa	720
tggaaactgt acagtagtaa tgactgatgg gagtgttca ggaagagctg atactaaaat	780
actattcatt gaggagggga aaatcgttca taccagccca ctgtcaggaa gtgctcagca	840
tgctgaggag tgctcctgtt atcctcgata tccgtgtgtc agatgtgtct gcagagacaa	900
ctggaaaaggc tccaataggc ccacgtaga tataaatgta aaggattata gcattgtttc	960
cagttatgtg tgctcaggac ttgttgaga cacaccaga aaaaacgaca gctccagcag	1020
tagtcattgc ctgaatccta acaatgagga agggggtcat ggagtgaag gctgggcctt	1080
tgatgatgga aatgacgtgt ggatgggaag aacgatcagc gagaagttcc gctcaggtta	1140
tgaacacctc aaagtcatg aaggctgtgc caaacctaac tccaaattgc agataaatag	1200
gcaagtcata gttgacagag gtaataggtc cggttattct ggtattttct ctgttgaagg	1260
caaaagctgc atcaatcgtt gcttttatgt ggagttgata aggggaagga aacaggaaac	1320
tgaagtctgg tggacctcaa acagtattgt tgtgtttgt ggcacctcag gtacatatgg	1380
aacaggctca tggcctgatg gggcggacat caatctcatg cctatataag ctttcgcaat	1440
tttagaaaaa aactccttgt ttctact	1467

<210> SEQ ID NO 9
 <211> LENGTH: 1762
 <212> TYPE: DNA
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 9

agcaaaaagca ggggataatt ctattaacca tgaagactat cattgctttg agctacattt	60
tatgtctggt tttcgtcaa aaacttccc gaaatgacaa cagcacggca acgctgtgcc	120

-continued

tggggcacca	tgctactgtca	aacggaacgc	tagtgaaaac	aatcacgaat	gaccaaattg	180
aagtactata	tgctactgag	ctgggttcaga	gttcctcaac	aggtagaata	tgcgacagtc	240
ctcaccaaat	ccttgatgga	gaaaactgca	cactaataga	tgctctattg	ggagaccctc	300
attgtgatgg	cttccaaaat	aaggaatggg	acctttttgt	tgaacgcagc	aaagcctaca	360
gcaactgtta	cccttatgat	gtgccggatt	atgcctccct	taggtcacta	gttgccctcat	420
ccggcacact	ggagttaaac	aatgaaagct	tcaattggac	tggagtcgct	cagaatggaa	480
caagctctgc	ttgcaaaaag	ggatctaata	aaagtttctt	tagtagattg	aattggttgc	540
accaattaaa	atacaaatat	ccagcactga	acgtgactat	gccaaacaat	gaaaaatttg	600
acaaattgta	catttggggg	gttctccacc	cgagtacgga	cagtgaacca	atcagcctat	660
atgctcaagc	atcagggaga	gtcacagtct	ctacccaaag	aagccaacaa	actgtaatcc	720
cgaatatcgg	atctagaccc	tgggtaaggg	gtgtctccag	cagaataagc	atctattgga	780
caatagtaaa	accgggagac	atacttttga	ttaacagcac	agggaatcta	attgctcttc	840
ggggttactt	caaaatcaga	agtgggaaaa	gctcaataat	gaggtcagat	gcacccattg	900
gcaaatgcaa	ttctgaatgc	atcactccaa	atggaagcat	tcccaatgac	aaaccatttc	960
aaaatgtaaa	caggatcaca	tatggggcct	gtcccagata	tgtaagcaa	aacactctga	1020
aattggcaac	agggatgcgg	aatgtaccag	agaacaaaac	tagaggcata	ttcggcgcaa	1080
tcgcggtttt	catagaaaat	ggttgggagg	gaatggtgga	cggttggtac	ggtttcaggc	1140
atcaaaattc	tgagggcaca	ggacaagcag	cagatcttaa	aagcactcaa	gcagcaatca	1200
accaaataca	cgggaaactg	aataggttaa	tcgagaaaac	gaacgagaaa	ttccatcaaa	1260
ttgaaaaaga	attctcagaa	gtagaaggga	gaattcagga	cctcgagaaa	tatgttgagg	1320
acactaaaat	agatctctgg	tcgtacaacg	cggagcttct	tgttgccctg	gagaaccaac	1380
atacaattga	tctaactgac	tcagaaatga	acaaactggt	tgaagaaca	aagaagcaac	1440
tgagggaaaa	tgctgaggat	atgggcaatg	gttgtttcaa	aataaccac	aatgtgaca	1500
atgcctgcat	agggtcaatc	agaaatggaa	cttatgacca	tgatgtatac	agagacgaag	1560
cattaataca	ccggttcacg	atcaaagggt	ttgagctgaa	gtcaggatac	aaagattgga	1620
tcctatggat	ttcctttgcc	atatcatgct	ttttgctttg	tgtgtttttg	ctgggggttca	1680
tcattgtggc	ctgccaaaaa	ggcaacatta	ggtgcaacat	ttgcatttga	gtgcattaat	1740
taaaaacacc	cttgtttcta	ct				1762

<210> SEQ ID NO 10

<211> LENGTH: 1466

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 10

agcaaaagca	ggagtaaaga	tgaatccaaa	tcaaaagata	ataacgattg	gctctgtttc	60
tctcactatt	gccacaatat	gcttcottat	gcaaatagcc	atcctggtaa	ctactgtaac	120
attgcatttc	aagcaatatg	aatgcaactc	ccccccaaac	aaccaagtaa	tgctgtgtga	180
accaacaata	atagaagaa	acataacaga	gatagtgtat	ctgaccaaca	ccaccataga	240
gaaggaaata	tgccccaaac	tagcagaata	cagaaattgg	tcaaagccgc	aatgtaaaat	300
tacaggattt	gcaccttttt	ctaaggataa	ttcaattcgg	ctttccgctg	gtggggacat	360

-continued

ttgggtgaca agagaacctt atgtgtcatg cgatcctgac aagtgttata aatttgccct	420
tggacagggg acaacactaa acaacaggca ttcaaatgac acagtacatg ataggacccc	480
ttatcgaacc ctattgatga atgagttggg tgttccattt catttgggaa ccaagcaagt	540
gtgtatagca tgggtccagct caagttgtca cgatggaaaa gcatggctgc atgtttgtgt	600
aactgggcat gatgaaaatg caactgctag cttcatttac gatgggagac ttgtagatag	660
tattggttca tgggtccaaa aaatcctcag gaccaggag tcggaatgag ttgtatcaa	720
tggaaactgt acagtagtaa tgactgatgg gagtgcttca ggaagagctg atactaaaat	780
acttttcatt gaggagggga aaatcgttca tactagcaaa ttgtcaggaa gtgctcagca	840
tgctgaggag tgctcctgtt atcctcgata tcttggtgtc agatgtgtct gcagagacaa	900
ctggaaaggc tccaataggc ccatcgtaga tataaatgta aaggattata gcattgtttc	960
cagttatgtg tgctcaggac ttgttgagga cacaccaga aaaaacgaca gctccagcag	1020
tagccattgc ctggatccta acaatgaaga aggggggtcat ggagtgaag gctgggcctt	1080
tgatgatgga aatgacgtgt ggatgggaag aacgatcagc gagaagtcac gctcaggtta	1140
tgaaaccttc aaggtcattg aaggtcgttc caaacctaac tccaaattgc agataaatag	1200
gcaagtcata gttgaaagag gtaatatgtc cgggtattct ggtattttct ctgttgaagg	1260
caaaagctgc atcaatcggt gcttttatgt ggagttgata aggggaagga aacaggaaac	1320
tgaagctcgg tggacctcaa acagtattgt tgtgttttgt ggcacctcag gtacatatgg	1380
aacaggctca tggcctgatg gggcggacat caatctcatg cctatataag ctttcgcaat	1440
tttagaaaaa actccttgtt tctact	1466

<210> SEQ ID NO 11

<211> LENGTH: 1762

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 11

agcaaaagca ggggataatt ctattaacca tgaagactat cattgcttta agtacattc	60
tatgtctggt tttctctcaa aagcttcccg gaaatgacaa cagcacggca acgctgtgcc	120
ttgggcacca tgcagtacca aacggaacga tagtgaaaac aatcacgaat gaccaaattg	180
aagttactaa tgctactgag ctggttcaga gttcctcaac aggtggaata tgcgacagtc	240
ctcatcagat ccttgatgga gaaaactgca cactaataga tgctctattg ggagaccctc	300
agtgtgatgg cttccaaaat aagaaatggg acctttttgt tgaacgcagc aaagcctaca	360
gcaactgtta cccttatgat gtgccggatt atgcctccct taggtcacta gttgcctcat	420
ccggcacact ggagtttaac aatgaaagct tcaattgggc tggagtcaact cagaatggaa	480
caagctctgc ttgcaaaagg agatctaata aaagtttctt tagtagattg aattggttga	540
cccacttaaa atacaaatac ccagcattga acgtgactat gccaaacaat gaaaaatttg	600
acaaattgta catttggggg gttcaccacc cggttacgga cagtgaacaa atcagcctat	660
atgtcaagc atcaggaaga atcacagtct ctacaaaaag aagccaacaa actgtaatcc	720
cgaatatcgg atatagaccc agggtaaggg atatctccag cagaataagc atctattgga	780
caatagtaaa accgggagac atacttttga ttaacagcac aggaaatcta attgctctc	840
ggggttactt caaaatcga agtgggaaaa gctcaataat gagatcagat gcaccattg	900

-continued

```

gcaaatgcaa ttctgaatgc atcactccaa atggaagcat tccaatgac aaaccatttc 960
aaaaatgtaaa caggatcaca tatggggcct gtcccagata tgtaagcaa aacactctga 1020
aattggcaac agggatgcga aatgtaccag agaaacaaac tagaggcata ttggcgcaa 1080
tcgcggttt catagaaaat gggtgggagg gaatggtgga cggttgttac ggtttcaggc 1140
atcaaaatc tgagggcaca ggacaagcag cagatctcaa aagcactcaa gcagcaatca 1200
acaaatcaa tgggaaactg aataggttaa tcgggaaaac aaacgagaaa ttccatcaga 1260
ttgaaaaaga attctcagaa gtagaaggga gaattcagga cctcgagaaa tatgttgagg 1320
acactaaaat agatctctgg tcatacaacy cggagcttct tgttgccctg gaaaaccaac 1380
atacaattga tctaactgac tcagaaatga acaactggt tgaaagaaca aagaagcaac 1440
tgagggaaaa tgctgaggat atgggcaatg gttgtttcaa aatataccac aaatgtgaca 1500
atgctgcat agagtcaatc agaaatggaa cttatgacca tgatgtatac agagatgaag 1560
cattaacaa cgggttcacg atcaaggtg ttgagctgaa gtcaggatac aaagattgga 1620
tcctatggat ttcctttgcc atatcatgtt ttttgctttg tgttgctttg ttggggttca 1680
tcatgtgggc ctgcaaaaaa ggcaacatta ggtgcaacat ttgcatttga gtgcattaat 1740
taaaaacacc cttgtttcta ct 1762

```

<210> SEQ ID NO 12

<211> LENGTH: 1467

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 12

```

agcaaaagca ggagtaaaga tgaatccaaa tcaaaagata ataacgattg gctctgtttc 60
cctcaccatt tccacaatat gcttcttcat gcaaattgcc atcctgataa ctactgtaac 120
attgcatttc aagcaatatg aattcaactc ccccccaaac aaccaagtga tgctgtgtga 180
accaacaata atagaagaa acataacaga gatagtgtat ctgaccaaca ccaccataga 240
gaaggaaata tgcccaaac tagcagaata cagaaattgg tcaagccgc aatgtaacat 300
tacaggattt gcaccttttt ctaaggacaa ttcgattcgg ctttcgctg gtggggacat 360
ctgggtgaca agagaacctt atgtgtcatg cgatcctgac aagtgttate aatttgcct 420
tggaacagga acaacactaa acaacgtgca ttcaaatgac acagtacatg ataggacccc 480
ttatcgacc ctattgatga atgagttggg tgttcattt catctgggga ccaagcaagt 540
gtgcatagca tgggtccagct caagtgtgca cgatggaaaa gcatggctgc atgtttgtgt 600
aacgggggat gatgaaaatg caactgctag cttcatttac aatgggaggc ttgtagatag 660
tattgtttca tgggtccaaa aaatcctcag gaccagagg tcagaatgcg ttgtatcaa 720
tggaacttgt acagtagtaa tgactgatgg gagtgcttca ggaaaagctg atactaaaat 780
actattcatt gaggagggga aaattgttca tactagcaca ttatcaggaa gtgctcagca 840
tgtcaggagg tgctcctgtt atcctcgata tcttggtgtc agatgtgtct gcagagacaa 900
ctggaaaggc tccaataggc ccatcgtaga tataaacata aggattata gcattgtttc 960
cagttatgtg tgctcaggac ttgttgaga cacaccaga aaaaacgaca gctccagcag 1020
tagccattgc ttgatccaa acaatgagga aggtggtcat ggagtgaag gctgggcatt 1080
tgatgatgga aatgacgtgt ggatgggaag aacgatcagc gagaagttac gctcaggata 1140

```

-continued

tgaaaccttc aaagtcattg aaggctggtc caaccctaac tccaaattgc agataaatag	1200
gcaagtcata gttgacagag gtaacaggtc cggttattct ggtattttct ctgttgaagg	1260
caaaagctgc atcaatcggg gcttttatgt ggagttgata aggggaagaa aacaggaaac	1320
tgaagtcttg tggacctcaa acagtattgt tgtgttttgt ggcacctcag gtacatatgg	1380
aacaggctca tggcctgatg gggcggacat caatctcatg cctatataag ctttcgcaat	1440
tttagaaaa aactccttgt ttctact	1467

<210> SEQ ID NO 13

<211> LENGTH: 1778

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 13

agcaaaaagca ggggaaaata aaaacaacca aaatgaaagc aaaactacta gtcctgttat	60
gtgcatttac agctacatat gcagacacaa tatgtatagg ctaccatgcg aacaactcaa	120
ccgacactgt tgacacagta cttgagaaga acgtgacagt gacacactct gtcaacctac	180
ttgaggacag tcacaacgga aaactatgtc gactaaaggg aatagcccca ctacaattgg	240
gtaattgcag cgttgcccga tggatcttag gaaacccaaa atgcgaatca ctgttttcta	300
aggaatcatg gtcctacatt gcagaaacac caaacccctga gaatggaaca tgttaccag	360
ggtatttcgc cgactatgag gaactgaggg agcaattgag ttcagtatca tcattcgaga	420
gattcgaaat attcccaaaa gaaagctcat ggcccaacca caccgtaacc aaaggagtaa	480
cgacatcatg ctcccataat gggaaaagca gtttttacag aaatttgcta tggctgacga	540
agaagaatgg cttgtaccca aatgtgagca agtcctatgt aaacaacaaa gagaaagaag	600
tcctgttact atgggggtgt catcacccgt ctaacatagg ggaccaaagg gccatctatc	660
atacagaaaa tgcttatgtc tctgtagtgt cttcacatta tagcagaaga ttcacccag	720
aaatagcaaa aagacccaaa gtaagagatc aagaaggaag aattaactac tactggactc	780
tgctggaacc cggggacaca ataatatgtg aggcaaatgg aaatctaata gcgccatggt	840
atgctttcgc actgagtaga ggctttgggt caggaatcat cacctcaaac gcatcaatgg	900
atgaatgtga cgcgaagtgt caaacacccc agggagctat aaacagtagt ctccctttcc	960
agaatgtaca cccagtcaca ataggagagt gtccaaagta tgtcaggagt acaaaattaa	1020
ggatggttac aggactaagg aacatcccat ccattcaatc cagaggtttg tttggagcca	1080
ttgccggttt cattgaaggg ggggtggactg gaatgataga tggatggtat ggttatcatc	1140
atcagaatga acaaggatct ggctatgctg cggaccaaaa aagcacacaa aatgccatta	1200
acgggattac aaacaagggt aattctgtaa tcgagaaaat gaacactcaa ttcacagctg	1260
tgggcaaaga attcaacaaa ttagaaagaa ggatggaaaa cttaaataaa aaagttgatg	1320
atggatttct ggacatttgg acatataatg cagaattgtt ggttctactg gaaaatggaa	1380
ggactttgga ttttcatgac tcaaagtga agaactgtga tgagaaagta aaaagccaat	1440
tgaagaataa tgccaaagaa atagggaacg ggtgttttga attctatcac aagtgtaca	1500
atgaatgcat ggaaagtgtg aaaaatggaa cttatgacta tccaaaatat tccgaagaat	1560
caaaagttaa caggggaaaa attgatggag tgaattgga atcaatggga gtctatcaga	1620
ttctggcgat ctactcaact gtcgccagtt cactggtgct tttggtctcc ctgggggcaa	1680

-continued

 tcagcttctg gatgtgttct aatgggtctt tgcagtgtag aatatgcac tgagaccaga 1740

atttcagaaa tataagaaaa aacacccttg tttctact 1778

<210> SEQ ID NO 14

<211> LENGTH: 1463

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 14

agcaaaagca ggagtttaaa atgaatccaa atcaaaaaat aataatcata ggatcaatca 60

gtatggcaat cggaataatt agtctaatat tgcaaatagg aaatattatt tcaatatggg 120

ctagccactc aatccaaact ggaagtcaaa accacactgg aatatgcaac caaagaatca 180

ttacatatga aaatagcacc tgggtgaatc aaacatatgt taatattaac aacactaatg 240

ttgttgctgg aaaggacaaa acttcagtga cattggccgg caattcatct ctttgcccta 300

tccgtgggtg ggctatatac acaaaagaca acagcataag aattgggtcc aaaggagatg 360

tttttgtcat aagagagcct tttatatcat gttctcactt ggaatgcaga accttttttc 420

tgaccaaggg tgctctatta aatgacaagc attcaaatgg gaccgttaag gacagaagcc 480

cttatagggc cttaatgagc tgtcctctag gtgaagctcc gtctccatac aattcaagat 540

ttgaatcagt tgcttggtca gcaagcgcat gccatgatgg catgggctgg ctaacaatcg 600

gaatttctgg tccagataat ggagcagtgg ctgtactaaa atacaacggc ataataactg 660

aaaccataaa aagttggaag aagcgaatat taagaacaca agagtctgaa tgtgtctgtg 720

tgaaagggtc atgttttacc ataatgacgg atggcccgag taatggggcc gcctcgtaca 780

gaatcttcaa aatcgagaag gggaagggtta ctaaatcaat agagtgggat gcaccaatt 840

atcattacga ggaatgttcc tgttaccag acaccggcac agtgatgtgt gtgtgcaggg 900

acaattggca cgtttcaaat cgaccttggg tgtcttttaa tcaaaacctg gattatcaaa 960

taggatacat ctgcagtggg gtgttcgggtg acaatccggc tcccaaagat ggagaaggca 1020

gctgtaatcc agtgactgtt gatggagcag acggagtaaa ggggttttca tacagatatg 1080

gtaatgggtg ttgatagga aggactaaaa gtaacagact cagaaaggga tttgagatga 1140

tttgggatcc taatggatgg acagataccg acagtgattt ctctgtgaaa caggatgtcg 1200

tggaatgac tgattgtgca gggtagacgg gaagtttcgt tcaacatcct gagctaacag 1260

gattggactg tatgagacct tgcttctggg ttgaattaat cagagggcga cctagagaaa 1320

atacaacaat ctggactagt gggagcagca tttctttttg tggcgtaa at agcgatactg 1380

caaaactggc ttggccagac ggtgcccaggt tgccattcac cattgacaag tagtccgttg 1440

aaaaaaaact ccttgtttct act 1463

<210> SEQ ID NO 15

<211> LENGTH: 1689

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 15

aaatgaaagc aaaactacta gtcctgttgt gtgcatttac agctacatat gcagacacaa 60

tatgtatagg ctaccatgcg aacaactcaa ccgacactgt tgacacagta cttgagaaga 120

acgtgacagt gacacactct gtcaacctac ttgaggacag tcacaacgga aaactatgcc 180

-continued

gactaaaagg aacagcccca ctacaattgg gtaattgcag cgttgccgga tggatcttag	240
gaaaccaga atgcgaatca ctgttttcta aggaatcatg gtccacatt gcagaaacac	300
caaaccctga gaatggaaca tgtaaccag ggtatttcgc cgactatgag gaactgaggg	360
agcaattgag ctacagtatca tcattcgaga gattcgaaat attccccaag gaaagctcat	420
ggcccaaaa caccgtaacc aaaggagtga cggcatcatg ctcccataat gggaaaagca	480
gtttttacaa aaatttgcta tggctgacgg aaaagaatgg cttgtacca aatctgagca	540
agtccatgt aaacaacaag gagaaagaag tccttgtagt atggggtgt catcacccgt	600
ctaacatagg ggaccaaagg gccatctatc atacagaaaa tgcttatgtc tctgtagtgt	660
cttcacatta tagcagaaga ttcaccccag aaatagcaaa aagacccaaa gtaagaggtc	720
aagaaggagg aattaactac tactggactc tgctggaacc cggggacaca ataataattg	780
aggcaaatgg aaatctaata gcgccatgg acgctttcgc actgagtaga ggctttgggt	840
caggaatcat cacctcaacc gcatcaatgg gtgaatgtga cgctaagtgt caaacacccc	900
aaggagctat aaacagtagt cttcctttcc agaatgtaca ccagtcaca ataggagagt	960
gtcccaagta tgtaggagt acaaaattaa ggatggttac aggactaaga aacatcccat	1020
ccattcaatc tagaggtttg tttggagcca ttgccggttt cattgaaggg gggtaggactg	1080
gaatgataga tggatggtat ggttatcatc atcagaatga acaaggatct ggctatgctg	1140
cagacaaaa aagcacacaa aatgccattg atgggattac aaacaagggtg aattctgtaa	1200
tcgagaaaa gaacactcaa ttcacagctg taggcaaaga attcaacaaa ttagagagaa	1260
ggatgaaaa cttaataaag aaagttagt atggatttct ggacatttgg acatataatg	1320
cagagttggt ggttctcctg gaaaatggaa ggactttggg ttttcatgac tcaaatgtga	1380
agaatctgta tgagaaagta aaaaaccaat tgaagaataa tgccaaagaa atcgggaacg	1440
ggtgttttga attctatcac aagtgtaca atgaatgcat ggaaagtgtg aaaaatggaa	1500
cttatgacta tccaaaatat tccgaagaat caaagttaaa cagggaaaaa attgatggag	1560
tgaaattgga atcaatggga gtctatcaga ttctggcgat ctactcaact gtcgccagtt	1620
cactggtgct tttggtctcc ctgggggcaa tcagtttctg gatgtgttct aatgggtctt	1680
tcagtgta	1689

<210> SEQ ID NO 16

<211> LENGTH: 1447

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 16

agcaaaagca ggagttaaaa atgaatccaa atcaaaaaat aataaccatt ggatcaatca	60
gtattgcaat tggaataatt agtctgatat tgcaaatagg aaatattatt tcaatatggg	120
ctagccactc aatccaaact ggaagtcaaa accacactgg aatatgcaac caaagaatca	180
ttacatatga aaatagcacc tgggtaaatc aaacatatgt taatattaac aacactaatg	240
ttgttgctgg aaaggacaaa acctcaatga cattggccgg caattcatct ctttgccta	300
tccgtggatg ggctatatac acaaaagaca acagcataag aattgggttc aaaggagatg	360
tttttgtcat aagagagcct tttatatcat gttctcaett ggaatgcaga accttttttc	420
tgaccaaggg tgctctatta aatgacaagc attcaaatgg gaccgttaag gacagaagcc	480

-continued

```

cttatagggc cttaatgagc tgtcctctag gtgaagctcc gtctccatac aattcaagat 540
ttgaatcagt tgcttggtca gcaagcgcac gccatgatgg cttgggctgg ctaacaatcg 600
gaatttctgg tccagataat ggggcagtggt ctgtactaaa atacaacggc ataataactg 660
aaaccattaa aagttggaag aagcgaatat taagaacaca agagtctgaa tgtgtctgta 720
tgaacgggtc atgttttacc ataatgaccg atggcccag taatggggcc gcatcgtaca 780
gaatcttcaa aatcgagaag gggagagtta ctaaatcaat agagttggat gcaccaatt 840
atcattacga ggaatgttca tgttaccag acaccggcac agtgatgtgt gtgtgcaggg 900
acaattggca cggttcfaat cgaccttggg tgtcttttaa tcaaacctg gattatcaaa 960
taggatacat ctgcagtggtg gtgttcgggtg acaatccgag tcccaaagat ggagaaggca 1020
gctgtaatcc agtgactgtt gatggagcag acggagtaaa ggggttttca tacagatatg 1080
gtaatgggtg ttgatagga aggactaaaa gtaacagact cagaaaggga tttgagatga 1140
tttgggatcc taatggatgg acagataccg acagtgattt ctcaatgaaa caggatatcg 1200
tggcaatgac tgattggtca ggggtacagc gaagttttgt tcaacatcct gagctaacag 1260
gattggactg tatgagacct tgcttttggg ttgaattagt cagagggcta cctagagaaa 1320
atacaacaat ctggactagt gggagcagca tttctttttg tggcgtaaat agcgatactg 1380
caaatgggtc ttggccagac ggtgcccagt tgccattcac cattgacaag tagtccgttg 1440
aaaaaaa 1447

```

<210> SEQ ID NO 17

<211> LENGTH: 1775

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 17

```

agcaaaagca ggggaaata aaaacaacca aaatgaaagc aaaactacta gtcctgttat 60
gtacatttac agctacatat gcagacacaa tatgtatagg ctaccatgcc aacaactcaa 120
ccgacactgt tgacacagta cttgagaaga atgtgacagt gacacactct gtcaacctac 180
ttgaggacag tcacaatgga aaactatgtc tactaaaagg aatagcccca ctacaattgg 240
gtaattgcag cgttgcccga tggatcttag gaaaccaga atgcgaatca ctgatttcta 300
aggaatcatg gtcctacatt gtagagacac caaacctga gaatggaaca tgttaccag 360
ggtatttcgc cgactatgag gaactgaggg agcaattgag ttcagtatca tcatttgaga 420
gattcgaaat attcccaaaa gaaagctcat ggcccaaaca caccgtaaca ggagtaacgg 480
catcatgctc ccataatggg aaaagcagtt tttacagaaa tttgctatgg ctgacggaga 540
agaatggcct gtacccaaat ctgagcaatt cctatgtgaa caacaaagag aaagaagtcc 600
ttgtactatg ggggtgttcat caccatcta acatagggga ccaagggcc atctatcata 660
cagaaaaacg ttatgtctct gtagtgtctt cacattatag cagaagattc accccagaaa 720
tagcaaaaag acccaaaagta agaggtcagg aaggaagaat caactactac tggactctgc 780
tggaaccggg ggacacaata atatttgagg caaatggaaa tctaatagcg ccatggtatg 840
ctttcgact gagtagaggc tttgggtcag gaatcatcac ctcaaatgca ccaatgaatg 900
aatgtgatgc gaagtgtcaa acacctcagg gagctataaa cagtagtctt cctttccaga 960
atgtacaccc agtcacaata ggagagtgtc caaagtatgt caggagtaca aaattaagga 1020

```

-continued

tggttacagg actaaggaat atcccatcca ttcaatccag aggtttgttt ggagccattg	1080
ccggtttcat tgaagggggg tggactggaa tgatggatgg gtggtatggt tatcatcatc	1140
agaatgagca aggatctggc tatgtgcag atcaaaaaag cacacaaaat gccattaacg	1200
ggattacaaa taaggtgaat tctgtaattg agaaaatgaa cactcaattc acagctgtgg	1260
gcaagaatt caacaaatta gaaagaagga tggaaaactt aaataaaaaa gttgatgatg	1320
gatttctaga catttgaca tataatgcag aattgttggg tctactggaa aatgaaagga	1380
ctttggattt ccatgactca aatgtgaaga atctgtatga gaaagtgaaa agccaattaa	1440
agaataatgc caaagaaata gggaacgggt gttttgaatt ctatcacaag tgtaacaatg	1500
aatgcatgga aagtgtgaaa aatggaactt atgactatcc aaaatattcc gaagaatcaa	1560
agttaaacag ggagaaaatt gatggagtga aattggaatc aatgggagtc tatcagattc	1620
tggcgatcta ctcaactgtc gccagttcac tggttctttt ggtctccctg ggggcaatca	1680
gcttctggat gtgttccaat gggctctttg agtgtagaat atgcatctga gaccagaatt	1740
tcagaaatat aagaaaaaac acccttgttt ctact	1775

<210> SEQ ID NO 18

<211> LENGTH: 1463

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 18

agcaaaagca ggagttaaa atgaatccaa atcaaaaaat aataaccatt ggatcaatca	60
gtatagtaat cgggataatt agtctaattg tgcaaatagg aaatattatt tcaatatggg	120
ctagtcactc aatccaaact ggaagtcaaa accacactgg aatatgcaac caaagaatca	180
tcacatatga aaatagcacc tgggtgaatc acacatatgt taatattaac aacactaatg	240
ttgttgctgg aaaggacaaa acttcagtga cattggccgg caattcatca ctttgttcta	300
tcagtggatg ggctatatac acaaaagaca acagcataag aattgggtcc aaaggagatg	360
tttttgtcat aagagagcct tttatatcat gttctcactt ggaatgcaga accttttttc	420
tgaccaaggg tgctctatta aatgacaaac attcaaatgg gaccgttaag gacagaagtc	480
cttatagggc cttaatgagc tgtcctctag gcgaagctcc gtctccatat aattcaaagt	540
ttgaatcagt tgcttggtca gcaagcgcat gtcgatgatg catgggctgg ttaacaatcg	600
gaatttctgg tccagataat ggagcagtgg ctgtactaaa atacaacggc ataataactg	660
aaaccataaa aagttggaag aagcgaatat taagaacaca agagtctgaa tgtgtctgtg	720
tgaacgggtc atgttttacc ataatgaccg atggcccag taatggggcc gcctcgtaca	780
aaatcttcaa gattgagaag ggggaaggta ctaaatcaat agagttegaat gcaccaatt	840
ctcattatga ggaatgttcc tgtaaccag aactggcac agtgatgtgt gtatgcaggg	900
acaattggca cggttcaaat cgaccttggg tgtcttttaa tcaaaacctg gattatcaaa	960
taggatacat ctgcagtggg gtgttcgggt acaatccggg tcccaaagat ggagagggca	1020
gctgtaatcc agtgactgtt gatggagcag acggagtaaa ggggttttca tacagatatg	1080
gtaatgggtg ttggatagga aggactaaaa gtaacagact cagaaaggga tttgagatga	1140
tttgggatcc taatggatgg acagataccg acagtgattt ctacagtgaag caggatgttg	1200
tggcaatgac tgattggtca gggtaacagc gaagtttcgt tcaacatcct gagctaacag	1260

-continued

gattggactg tataagacct tgcttctggg ttgaattagt cagaggacgg cctagagaaa	1320
atacaacaat ctggactagt gggagcagca tttctttttg tggcgtaaat agtgatactg	1380
caaaactggtc ttggccagac ggtgctgagt tgccattcac cattgacaag tagtccgttg	1440
aaaaaaaaact ccttgtttct act	1463

<210> SEQ ID NO 19

<211> LENGTH: 1775

<212> TYPE: DNA

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 19

agcaaaagca ggggaaaata aaaacaacca aaatgaaagc aaaactactg gtcctgttat	60
gtacatttac agctacatat gcagacacaa tatgtatagg ctaccatgcc aacaactcaa	120
ccgacactgt tgacacagta cttgagaaga atgtgacagt gacacactct gtcaacctac	180
ttgaggacag tcacaatgga aaactatgtc tactaaaagg aatagcccca ctacaattgg	240
gtaattgcag cgttgccgga tggatccttag gaaaccaga atgcgaatta ctgatttcca	300
aggaatcatg gtcctacatt gtagaaacac caaatcctga gaatggaaca tgttaccag	360
ggtatttcgc cgactatgag gaactgaggg agcaattgag ttcagtatct tcatttgaga	420
gattcgaaat attcccaaaa gaaagctcat ggcccaaaaca caccgtaacc ggagtatcag	480
catcatgctc ccataatggg aaaaacagtt ttacagaaa ttgctatgg ctgacgggga	540
agaatggttt gtacccaaac ctgagcaagt cctatgtaaa caacaaagag aaagaagtcc	600
ttgtactatg ggggtgttcat caccgccta acatagggga ccaagggcc ctctatcata	660
cagaaaaatgc ttatgtctct gtagtgtctt cacattatag cagaagattc accccagaaa	720
tagccaaaag acccaaaagta agagatcagg aaggaagaat caactactac tggactctgc	780
tggaaacctg ggatacaata atatttgagg caaatggaaa tctaatagcg ccatgggatg	840
cttttgact gagtagaggc tttggatcag gaatcatcac ctcaaagca ccaatggatg	900
aatgtgatgc gaagtgtcaa acacctcagg gagctataaa cagcagtctt cctttccaga	960
atgtacaccc agtcacaata ggagagtgtc caaagtatgt caggagtgc aaattgagga	1020
tggttacagg actaaggaa atcccatcca tccaatccag aggtttgttt ggagccattg	1080
ccggtttcat tgaagggggg tggactggaa tggtagatgg gtggtatggt tatcatcatc	1140
agaatgagca aggatctggc tatgtgcag atcaaaaaag tacacaaaat gccattaacg	1200
ggattacaaa caaggtgaat tctgtaattg agaaaatgaa cactcaattc acagctgtgg	1260
gcaaagaatt caacaaattg gaaagaagga tggaaaactt aaataaaaaa gttgatgatg	1320
ggtttctaga catttggaca tataatgcag aattgttggg tctactggaa aatgaaagga	1380
ctttggattt ccatgactcc aatgtgaaga atctgtatga gaaagtaaaa agccaattaa	1440
agaataatgc caaagaata ggaaacgggt gttttgaatt ctatcacaag tgtaacaatg	1500
aatgcatgga gagtgtgaaa aatggaactt atgactatcc aaaatatcc gaagaatcaa	1560
agttaaacag ggagaaaatt gatggagtga aattggaatc aatgggagtc tatcagattc	1620
tggcgatcta ctcaactgtc gccagttccc tggttctttt ggtctccctg ggggcaatca	1680
gcttctggat gtgttccaat gggctcttgc agtgtagaat atgcactctga gaccagaatt	1740
tcagaagtat aagaaaaaac acccttgttt ctact	1775

-continued

<210> SEQ ID NO 20
<211> LENGTH: 1463
<212> TYPE: DNA
<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 20

```
agcaaaagca ggagttttaa atgaatccaa atcaaaaaat aataaccatt ggatcaatca    60
gtatagcaat cggaataatt agtctaattg tgcaaatagg aaatattatt tcaatatggg    120
ctagtcaact aatccaaact ggaagtcaaa accacactgg agtatgcaac caaagaatca    180
tcacatatga aaacagcacc tgggtgaatc acacatatgt taatattaac aacactaatg    240
ttgttgctgg aaaggacaaa acttcagtga cattggccgg caattcatct ctttgttcta    300
tcagtggatg ggctatatac aaaaagaca acagcataag aattggctcc aaaggagatg    360
tttttgctat aagagaacct ttcatatcat gttctcactt ggaatgcaga accttttttc    420
tgaccaaggg tgctctatta aatgacaaac attcaaatgg gaccgttaag gacagaagtc    480
cttatagggc cttaatgagc tgtcctctag gtgaagctcc gtcccatcac aattcaaagt    540
ttgaatcagt tgcattggtc gcaagcgcac gccatgatgg catgggctgg ttaacaatcg    600
gaattttctg tccagacaat ggagctgtgg ctgtactaaa atacaacggc ataataactg    660
aaaccataaa aagttggaaa aagcgaatat taagaacaca agagtctgaa tgtgtctgtg    720
tgaacgggtc atgtttcacc ataatgaccg atggcccag taatggggcc gcctcgtaca    780
aaatcttcaa gatcgaaaag gggagggtta ctaaatcaat agagttgaat gcaccaatt    840
ttcattatga ggaatgttcc tgttaccagg aactggcac agtgatgtgt gtatgcaggg    900
acaactggca tggttcaaat cgacctgggg tgtcttttaa tcaaacctg gattatcaaa    960
taggatacat ctgcagtggg gtgttcgggt acaatccgag tcccaaagat ggagagggca   1020
gctgtaatcc agtgactgtt gatggagcag acggagtaaa ggggttttca taaaaatatg   1080
gtaatggtgt ttggatagga aggactaaaa gtaacagact tagaaagggg ttgagatga    1140
tttgggatcc taatggatgg acagataccg acagtgattt ctcagtgaat caggatgttg   1200
tggaataaac tgattggtc gggtagacgg gaagtctcgt tcaacatcct gagttaacag   1260
gattggactg tataagacct tgcttctggg ttgagttagt cagaggactg cctagagaaa   1320
atacaacaat ctggactagt gggagcagca tttctttttg tggcgtaaat agtgatactg   1380
caaaactggc ttggccagac ggtgctgagt tgccgttcac cattgacaag tagttcgttg   1440
aaaaaaaact ccttgtttct act                                         1463
```

<210> SEQ ID NO 21
<211> LENGTH: 1879
<212> TYPE: DNA
<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 21

```
agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt actactcatg    60
gtagtaacat ccaacgcaga tcgaatctgc actgggataa catcttcaaa ctcacctcat   120
gtggtcaaaa cagctactca aggggaagtc aatgtgactg gtgtgatacc actgacaaca   180
acaccaacaa aatctcattt tgcaaatctc aaaggaacaa agaccagagg gaaactatgc   240
ccaaactgtc tcaactgcac agatctggat gtggccttgg gcagaccaat gtgtataggg   300
```

-continued

```

atcacacctt cggcaaaagc ttcaatactc cacgaagtca gacctgttac atccgggtgc 360
tttctataa tgcacgacag aacaaaaatc agacagctac ccaatcttct cagaggatat 420
gaacatatca gattatcaac ccataacgtt atcaacgcag aaagggcacc aggaggaccc 480
tacagacttg gaacctcagg atcttgccct aacgttacca gtagaagcgg attcttcgca 540
acaatggctt gggctgtccc aagggaacaac aaaacagcaa cgaacccact aacagtagaa 600
gtaccataca tttgtacaaa aggagaagac caaattactg tttgggggtt ccattctgat 660
aacaaaaatc aaatgaaaaa cctctatgga gactcaaatc ctcaaaagtt cacctcatct 720
gccaatggaa taaccacaca ttatgtttct cagattggtg gcttcccaaa tcaaacagaa 780
gacggagggc taccacaaag cggcagaatt gttgttgatt acatggtgca aaaacctggg 840
aaaaacaggaa caattgtcta tcaaagaggt gttttgttgc ctcaaaaggt gtggtgtgca 900
agtggcagga gcaaggtaat aaaagggctc ttgcctttaa ttggtgaagc agattgcctt 960
cacgaaaaat acggtggatt aaacaaaagc aagccttact acacaggaga acatgcaaaa 1020
gccataggaa attgcccaat atgggtgaaa acacctttaa agcttgccaa tggaaccaa 1080
tatagacctc ccgcaaaact attaaaggaa aagggtttct tcggagctat tgctggtttc 1140
ttagaaggag gatgggaagg aatgattgca ggttggcacg gatacacatc tcatggagca 1200
catgggggtg cagtggcagc agaccttaag agtacgcaag aagccataaa caagataaca 1260
aaaaatctca attctttgag tgagctagaa gtaaagaatc ttcaaagact aagtgtgtgc 1320
atggatgaac tccacaacga aatactcgag ctggatgaga aagtggatga tctcagagct 1380
gacacaataa gctcgcaaat agagcttgca gtcttgcttt ccaatgaagg aataataaac 1440
agtgaagatg agcatctatt ggcacttgag agaaaactaa agaaaatgct gggtcctct 1500
gctgtagaca tagggaatgg atgcttcgaa accaaacaca agtgcaacca gacctgetta 1560
gacaggatag ctgctggcac ctttaatgca ggagaatttt ctcttccac ttttgattca 1620
ctgaatatta ctgctgcac tttaaatgat gatggattgg ataatacaca tatactgctc 1680
tactactcaa ctgcggtctc tagtttggt gtaacattga tgatagctat ttttattgtt 1740
tatatggtct ccagagacaa tgtttcttgc tccatctgtc tatagggaaa attgagccct 1800
gtattttctt ttattgttgt gcttgtttgc ttgttgccat tacagagaaa cgttattgaa 1860
aaatgctctt gttactact 1879

```

<210> SEQ ID NO 22

<211> LENGTH: 1554

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 22

```

agcagaagca gagcatcttc tcaaaactga agtaaagagg ccaaaatga acaatgctac 60
cttcaactat acaaacgtta accctatttc tcacatcagg gggagtgtta ttatcactat 120
atgtgtcagc cttactgtca tacttattgt attcgatat attgctaaaa ttttcaccaa 180
aaataattgc accaacaacg tcgttggtgact ggcgaacgc atcaaagtt caggctgtga 240
accattctgc aacaaaagag atgaaattcc ttccccaga accggagtgg acatacccc 300
gtttatcttg ccagggttca accttcocaga aagcactett aattagccct catagatttg 360
gagaagccaa aggaaactca gctcccttga taataaggga accttttatt gcttgtggac 420

```

-continued

caaaggagtg caaacacttt gctctaacc	attatgcagc tcaaccaggg ggatactaca	480
atggaacaag agaggacaga aacaagctga	ggcatctgat ttcagtcaac ttaggcaaaa	540
tcccaactgt agaaaactcc attttccata	tggcagcttg gaggatgcc gcatgccatg	600
atggtagaga atggacatat atcggagttg	atggtcctga cagtaatgca ttgatcaaaa	660
taaaatatgg agaagcatac actgacacat	accattccta tgcaaacac atcctaagaa	720
cacaagaaag tgctgcaat tgcacgggg	gagattgtta tcttatgata actgatggct	780
cagcttcagg aattagtaaa tgcagattcc	ttaagatccg agagggtcga ataataaaag	840
aaatatttcc aacaggaagg gtagagcaca	ctgaagaatg cacatgcgga ttgccagca	900
acaaaaccat agaatgtgcc tgtagagata	acagttacac agcaaaaaga ccctttgtca	960
aattaaatgt ggagactgat acagctgaaa	taagattgat gtgcacagag acttatttgg	1020
acacccccag accagatgat ggaagcataa	cagggccttg cgaatctaag ggggacaaag	1080
ggagtggagg tgtcaaggga ggatttgttc	atcaaagaat ggcatccaag attggaagat	1140
ggtactcccc aacgatgtct aaaactaaaa	gaatggggat ggaactgtat gtcaagtatg	1200
atggagaccc atggactgac agtgacgccc	ttgctcctag tggagtaatg gtctcaatgg	1260
aagaacctgg ttgtactct ttcggcttcg	aaataaaaaga taagaaatgt gatgtccct	1320
gtattgggat agagatggta catgatggtg	gaaaaaggac ttggcactca gcagcaacag	1380
ccatttactg tttaatgggc tcaggacagt	tgctatggga cactgtcaca ggtgttaata	1440
tggctctgta atggaggaat ggttgaatct	gttctaaacc ctttgttctt attttatttg	1500
aacaattgtc cttactggac ttaattgttt	ctgaaaaatg ctcttggtac tact	1554

<210> SEQ ID NO 23

<211> LENGTH: 1881

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 23

agcagaagca gagcattttc taatatccac	aaaatgaagg caataattgt actactcatg	60
gtagtaacat ccaatgcaga tcgaatctgc	actgggataa categtcaaa ctcacctcat	120
gtggtcaaaa cagctactca aggggaggtc	aatgtgactg gtgtgatacc actgacaaca	180
acaccaacaa aatctcattt tgcaaatctc	aaaggaacaa agaccagagg gaaactatgc	240
ccaacctgtc tcaactgcac agatctggat	gtggccttag gcagaccaat gtgtgtgggg	300
gtcacacctt cggcaaaagc ttcaatactc	cacgaagtca ggctgttac atccggatgc	360
tttctataa tgcacgacag aacaaaaatc	agacagctac ccaatcttct cagaggatat	420
gaaaaaatca gattatcaac ccaaatcgtt	atcaacgcag aaaaggcacc aggaggaccc	480
tacagacttg gaacctcagg atcttgccct	aacgctacca gtagaagcgg atttttcgca	540
acaatggctt gggctgtccc aaaggacaac	aacaaaacag caacgaatcc actaacagta	600
gaagtaccac acatctgtac aaaagaagaa	gaccaaatta ctgtttgggg gttccattct	660
gatgacaaaa cccaaatgaa aaacctctat	ggagactcaa atcctcaaaa gttcacctca	720
tctgctaatt gagtaaccac acattatgtt	tctcagattg gcggcttccc ggatcaaaaa	780
gaagacggag ggctaccaca aagcggcaga	attgttgttg attacatggg gcaaaaacct	840
gggaaaaacag gaacaattgt ctatcaaaga	ggtattttgt tgcctcaaaa ggtgtggtgc	900

-continued

```

gcgagtggca ggagcaaagt aataaaaggg tccttgcctt taattggtga agcagattgc 960
cttcacgaaa aatacgggtg attaaacaaa agcaagcctt actacacagg agaacatgca 1020
aaagccatag gaaattgccc aatatgggtg aaaacacctt tgaagcttgc caatggaacc 1080
aaatatagac ctcttgcaaa actattaaag gaaaggggtt tcttcggagc tattgctggt 1140
ttcttagaag gaggatggga aggaatgatt gcaggttggc acggatacac atctcacgga 1200
gcacatggag tggcagtggc agcagacctt aagagtacgc aagaagccat aaacaagata 1260
acaaaaaatc tcaattcttt gagttagcta gaagtaaaga atcttcaaag actaagtggg 1320
gccatggatg aactccacaa cgaaatactc gagctggatg agaaagtgga tgatctcaga 1380
gctgacacaa taagctcaca aatagaactt gcagtcttgc tttccaaaga aggaataata 1440
aacagtgaag atgagcatct attggcactt gagagaaaac taaagaaaat gctgggtccc 1500
tctgctgtag acatagggaa tggatgcttc gaaaccaaac acaagtgcaa ccagacctgc 1560
ttagacagga tagctgctgg cacctttaat gcaggagaat tttctcttcc cacttttgat 1620
tcactgaata ttactgctgc atctttaaat gatgatggat tggataacca tactatactg 1680
ctctactact caactgctgc ttctagtgtg gctgtaacat tgatgatagc tatttttatt 1740
gtttatatga tctccagaga caatgtttct tgctccatct gtctataggg aaattaagcc 1800
ctgtattttc ctttattgta gtgcttgttt gcttgttatc attacaaaga aacgttattg 1860
aaaaatgctc ttgttactac t 1881

```

<210> SEQ ID NO 24

<211> LENGTH: 1557

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 24

```

agcagaagca gagcatcttc tcaaaactga ggcaaatagg ccaaaaatga acaatgctac 60
cttcaactat acaaacgtta accctatttc tcacatcagg gggagtgtta ttatcactat 120
atgtgtcagc ttcactgtca tacttactat attcggatat attgctaaaa ttttcaccaa 180
cagaaataac tgcaccaaca atgccattga attgtgcaaa cgcatacaat gttcaggctg 240
tgaaccgttc tgcaacaaaa ggggtgacac ttctctctcc agaaccggag tggacatacc 300
ctcgtttata ttgcccgggc tcaaccttcc agaaagcact cctaattagc cctcatagat 360
tcggagaaac caaaggaaac tcagctccct tgataataag ggaacctttt attgcttgtg 420
gaccaaagga atgcagacac tttgtcttaa ccattatgc agcccaacca gggggatact 480
acaatggaac aagagaagac agaaacaagc tgaggcatct aatttcagtc aaattgggca 540
aaatcccaac agtagaaaac tccattttcc acatggcagc ttggagcggg tccgcatgcc 600
atgatggtag agaatggaca tatatcggag ttgatggccc tgacagtaat gcattgctca 660
aaataaaata tggagaagca tatactgaca cataccattc ctatgcaaac aacatcctaa 720
gaacacaaga aagtgcctgc aattgcactg ggggagattg ttatcttatg ataactgatg 780
gctcagcttc agggattagt gaatgcagat ttcttaagat tcgagagggc cgaataataa 840
aagaaatatt tccaacagga agagtagaac atactgaaga atgcacatgc ggatttgcca 900
gcaataaaac catagaatgt gcctgtagag ataacagtta cacagcaaaa agaccctttg 960
tcaaattaaa tgtggagact gatacagcag aaataagatt gatgtgcaca gagacttact 1020

```

-continued

tggacacccc	cagaccagat	gatggaagca	taacagggcc	ttgtgaatct	aatggggata	1080
aaggagtg	agcatcaag	ggaggatttg	ttcatcaaag	aatggcatcc	aagattggaa	1140
ggtggtactc	tcgaacgatg	tctaaaacta	aaaggatggg	gatgggactg	tatgtcaagt	1200
atgatggaga	cccatggatt	gacagtgatg	cccttactct	tagcggagta	atggtttcaa	1260
tggaagaacc	tggttggtat	tcctttggct	tcgaaataaa	agataagaaa	tgtgatgtcc	1320
cctgtattgg	gatagagatg	gtacatgatg	gtggaaagaa	gacttggcac	tcagcagcaa	1380
cagccattta	ctgtttaatg	ggctcaggac	aactgctatg	ggacactgtc	acaggcggtg	1440
atatggctct	gtaatggagg	aatggttgag	tctgttctaa	accctttgtt	cctattttgt	1500
ttgaacaatt	gtccttactg	aacttaattg	tttctgaaaa	atgctcttgt	tactact	1557

<210> SEQ ID NO 25

<211> LENGTH: 1882

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 25

agcagaagca	gagcattttc	taatatccac	aaaatgaagg	caataattgt	actactcatg	60
gtagtaacat	ccaatgcaga	tcgaatctgc	actgggataa	catcgtcaaa	ctcacctcat	120
gtggtcaaaa	cagctactca	aggggaggtc	aatgtgactg	gtgcgatacc	attgacaaca	180
acaccaacaa	aatctctatt	tgcaaatctc	aaaggaacaa	agaccagagg	gaaactatgc	240
ccaacctgtc	tcaactgcac	agatctggat	gtggccttgg	gcagaccaat	gtgtgtgggg	300
atcacacctt	cggcaaaagc	ttcaatactc	cacgaagtca	gacctgttac	atccggatgc	360
tttctataa	tgacgacag	aacaaaaatc	agacagctac	ccaatcttct	cagaggatat	420
gaaaaaatca	gattatcaac	ccaaaacgtt	atcaacgcag	aaaaggcacc	aggaggaccc	480
tacagacttg	gaacttcagg	atcttgcctt	aacgtacca	gtaaaagcgg	atttttcgca	540
acaatggctt	gggtgtccc	aagggaacac	aacaaaacag	caacgaatcc	actaacagta	600
gaagtaccac	acatctgtac	aaaagaagaa	gaccaaatta	ctgtttgggg	gttccattct	660
gatgacaaaa	cccaaatgaa	aaacctctat	ggagactcaa	atcctcaaaa	gttcacctca	720
tctgctaagt	gaataaccac	acattatggt	tctcagattg	gcggcttccc	ggaccaaaaa	780
gaagacggag	ggctaccaca	aagcggcaga	attgttgttg	attacatggg	gcaaaaacct	840
gggaaaaacag	gaacaattgt	ctatcaaaga	gggatcttgt	tgctcaaaa	ggtgtggtgc	900
gcgagtggca	ggagcaaagt	aataaaagg	tccttgctt	taattggtga	agcagattgc	960
cttcacgaaa	aatacgggtg	attaacaaa	agcaagcctt	actacacagg	agaacatgca	1020
aaagccatag	gaaattgccc	aatatgggtg	aaaacacctt	tgaagcttgc	caatggaacc	1080
aagtatagac	ctcctgcaaa	actattaaag	gaaaggggtt	tcttcggagc	tattgtcggg	1140
ttcttagaag	gaggatggga	aggaatgatt	gcaggttggc	acggatacac	atctcacgga	1200
gcacacggag	tggcagtggc	agcagacctt	aagagtacgc	aagaagccat	aaacaagata	1260
acaaaaaatc	tcaattcttt	gagtgtggtt	gaagtaaaga	accttcaaa	actaagtggg	1320
gccatggatg	aactccataa	cgaaatactc	gagctggatg	agaaagtggg	tgatctcaga	1380
gctgacacaa	taagctcaca	aatagaactt	gcagtcctgc	tttccaacga	aggaataata	1440
aacagtgaag	atgagcatct	attggcactt	gagagaaaac	taaagaagat	gctgggtccc	1500

-continued

tctgctatag acatagggaa tggatgcttc gaaaccaaac acaagtgcaa ccagacctgc	1560
ttagacagga tagctgctgg cacctttaat gcaggagaat tttctcttcc cacttttgat	1620
tcactgaaca ttactgctgc atctttaaat gatgatggat tggataacca tactatactg	1680
ctctactact caactgctgc ttctagtgtg gctgtaacat tgatgatagc tatttttatt	1740
gtttatatga tctccagaga caatgtttct tgctccatct gtctataagg aaaattaagc	1800
cctgtatttt cctttattgt agtgcttggt tgcttggtat cattacaaag aaacgttatt	1860
gaaaaatgct cttgttacta ct	1882

<210> SEQ ID NO 26

<211> LENGTH: 1557

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 26

agcagaagca gagcatcttc tcaaaactga ggcaaatagg ccaaaaatga acaatgctac	60
cctcaactat acaaacgtta accctattcc tcacatcagg gggagtgtta ttatcactat	120
atgtgtcagc ttactgtca tacttactat attcggatat attgctaaaa ttttcaccaa	180
cagaaataac tgcaccagca atgcccttgg attgtgcaa cgcacaaat gtccaggctg	240
tgaaccgttc tgcaacaaaa ggggtgacac ttcttctccc agaaccggag tggacatacc	300
cgcgtttatc ttgccggggt tcaaccttcc agaaagcact cctaattagc cctcatagat	360
tcggagaaac caaaggaaac tcagctccct tgataataag ggaacctttt attgcttggtg	420
gaccaaagga atgcaaacac ttgtctctaa cccattatgc agcccaacca gggggatact	480
acaatggaac aagagaagac agaaacaagc taaggcatct aatttcagtc aaatttggtg	540
aaatcccaac agtagaaaac tccattttcc acatggcagc atggagcggg tccgcatgcc	600
atgatggtaa agaattggaca tatatcggag ttgatggccc tgacagtaat gcattgctca	660
aaataaaata tggagaagca tatactgaca cataccattc ctatgcaaac aacatcctaa	720
gaacacaaga aagtgcctgc aattgcatcg ggggaaattg ttatcttatg ataactgatg	780
gctcagcttc aggtattagt gagtgcagat ttcttaagat tcgagagggc cgaataataa	840
aagaaatatt tccaacagga agagtaaaac atactgaaga atgcacatgc ggatttgcca	900
gcaataaaac catagaatgt gcctgtagag ataacagtta cacagcaaaa agaccctttg	960
tcaaattaaa tgtggagact gatacagcag aaataagatt gatgtgcaca gagacttatt	1020
tggacacccc cagaccagat gatggaagca taacagggcc ttgtgaatct aatggggata	1080
aaggagtggt aggcataaag ggaggatttg ttcacaaag aatggcatcc aagattggaa	1140
ggtggtactc tcgaacaatg tctaaaacta aaaggatggg gatgggactg tatgtcaagt	1200
atgatggaga cccatggact gacagtgatg cccttgctct tagtgagta atggtttcaa	1260
tggaagaacc tggttggtac tcctttggtc tcgaaataaa agataagaaa tgtgatgtcc	1320
cctgtattgg gatagagatg gtacatgatg gtgaaaagga gacttggcac tcagcagcaa	1380
cagccattta ctgtttaatg ggctcaggac aactgctatg ggacactgtc acaggtgttg	1440
atatggctct gtaattggagg aatggttgag tctgttctaa accctttggt cctattttgt	1500
ttgaacaatt gtccctactg aacttaattg tttctgaaaa atgctcttgt tactact	1557

-continued

```

<210> SEQ ID NO 27
<211> LENGTH: 1879
<212> TYPE: DNA
<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 27

agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt actactcatg      60
gtagtaacat ccaacgcaga tcgaatctgc actgggataa catcttcaaa ctcacctcat      120
gtggtcaaaa cagctactca aggggaagtc aatgtgactg gtgtgatacc actgacaaca      180
acaccaacaa aatctcatct tgcaaatctc aaaggaacaa agaccagagg gaaactatgc      240
ccaaactgtc tcaactgcac agatctggat gtggccttgg gcagaccaat gtgtataggg      300
atcacacctt cggcaaaagc ttcaatactc cacgaagtca gacctgttac atccgggtgc      360
tttctctata tgacagacag aacaaaaatc agacagctac ccaatcttct cagaggatat      420
gaacatatca gattatcaac ccataacgtt atcaacgcag aaagggcacc aggaggaccc      480
tacagacttg gaacctcagg atcttgccct aacgttacca gtagaagcgg attcttcgca      540
acaatggctt gggctgtccc aagggaacac aaaacagcaa cgaaccactt aacagtagaa      600
gtaccataca ttgtacaaa aggagaagac caaattactg tttgggggtt ccattctgat      660
aacaaaaatc aatgaaaaa cctctatgga gactcaaatc ctcaaaagtt cacctcatct      720
gccaatggaa taaccacaca ttatgtttct cagattggtg gcttcccaaa tcaaacagaa      780
gacggagggg taccacaaa cggcagaatt gttgttgatt acatggtgca aaaacctggg      840
aaaacaggaa caattgtcta tcaaagaggt gttttgttgc ctcaaaaggt gtggtgtgca      900
agtggcagga gcaaggtaat aaaagggctc ttgcctttaa ttggtgaagc agattgcctt      960
cacgaaaaat acggtggatt aaacaaaagc aagccttact acacaggaga acatgcaaaa      1020
gccataggaa attgccaat atgggtgaaa acacctttaa agcttgccaa tggaacaaa      1080
tatagacctc ccgcaaaact attaaaggaa aagggtttct tcggagctat tgctggtttc      1140
ttagaaggag gatgggaagg aatgattgca ggttggcacg gatacacatc tcatggagca      1200
catgggggtg cagtggcagc agaccttaag agtacgcaag aagccataaa caagataaca      1260
aaaaatctca attctttgag tgagctagaa gtaaagaatc ttcaaagact aagtggtgcc      1320
atggatgaac tccacaacga aatactcgag ctggatgaga aagtggatga tctcagagct      1380
gacacaataa gctcgcaaat agagcttgca gtcttgcttt ccaatgaagg aataataaac      1440
agtgaagatg agcatctatt ggcacttgag agaaaactaa agaaaatgct gggtcctct      1500
gctgtagaca tagggaatgg atgcttcgaa accaaacaca agtgcaacca gacctgctta      1560
gacaggatag ctgctggcac cttaaatgca ggagaatttt ctctccacac ttttgattca      1620
ctgaatatta ctgctgcac tttaaatgat gatggattgg ataatacatc tatactgctc      1680
tactactcaa ctgcggtctc tagtttggt gtaacattga tgatagctat tttattgtt      1740
tatatggtct ccagagacaa tgtttctgac tocatctgac tatagggaaa attgagccct      1800
gtattttcct ttattgtggt gcttgtttgc ttgttgccat tacagagaaa cgttattgaa      1860
aaatgctctt gttactact                                     1879

```

```

<210> SEQ ID NO 28
<211> LENGTH: 1554
<212> TYPE: DNA
<213> ORGANISM: Influenza B virus

```

-continued

<400> SEQUENCE: 28

```

agcagaagca gagcatcttc tcaaaactga agtaaagagg ccaaaaatga acaatgctac    60
cttcaactat acaaacgtta accctatttc tcacatcagg gggagtgtta ttatcactat    120
atgtgtcagc cttactgtca tacttattgt attcggatat attgctaaaa ttttcaccaa    180
aaataattgc accaacaacg tcggttgact gcgcgaacgc atcaaatgtt caggctgtga    240
accattctgc aacaaaagag atgaaattcc ttccccaga accggagtgg acatacccc    300
gtttatcttg ccagggttca accttcacaga aagcactctt aattagccct catagatttg    360
gagaagccaa aggaaactca gctcccttga taataagga accttttatt gcttgtggac    420
caaaggagtg caaacacttt gctctaacc attatgcagc tcaaccaggg ggatactaca    480
atggaacaag agaggacaga aacaagctga ggcactctgat ttcagtcaac ttaggcaaaa    540
tccccactgt agaaaactcc attttccata tggcagcttg gagtggatcc gcatgccatg    600
atggtagaga atggacatat atcggagtgt atggctctga cagtaatgca ttgatcaaaa    660
taaaatatgg agaagcatac actgacacat accattccta tgcaacaac atcctaagaa    720
cacaagaaa tgcttgaat tgcatcgggg gagattgtta tcttatgata actgatggct    780
cagcttcagg aattagtaaa tgcagattcc ttaagatccg agagggtcga ataataaaaag    840
aaatatcttc aacaggaagg gtagagcaca ctgaagaatg cacatgcgga tttgccagca    900
acaaaacat agaattgtgc ttagagata acagttacac agcaaaaaga ccttttgtca    960
aattaaatgt ggagactgat acagctgaaa taagattgat gtgcacagag acttatttgg    1020
acacccccag accagatgat ggaagcataa cagggccttg cgaatcta at ggggacaaaag    1080
ggagtggagg tgtcaagga ggatttgttc atcaagaat ggcaccaag attggaagat    1140
ggtactcccg aacgatgtct aaaaactaaa gaatggggat ggaactgtat gtcaagtatg    1200
atggagaccc atggactgac agtgacgccc ttgctcctag tggagtatg gtctcaatgg    1260
aagaacctgg ttgtactct ttcggtctg aaataaaaaga taagaaatgt gatgtccct    1320
gtattgggat agagatggta catgatgggt gaaaaaggac ttggcactca gcagcaacag    1380
ccatttactg ttaattgggc tcaggacagt tgctatggga cactgtcaca ggtgttaata    1440
tggtctgtga atggaggaat ggttgaatct gttctaaacc ctttgttctt attttatttg    1500
aacaattgtc cttactggac ttaattgttt ctgaaaaatg ctcttgttac tact        1554

```

<210> SEQ ID NO 29

<211> LENGTH: 1885

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 29

```

agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt actactcatg    60
gtagtaacat ccaatgcaga tcgaatctgc actggaataa catcgtcaaa ctcaccccat    120
gtggtcaaaa ctgctactca aggggaagtc aatgtgactg gtgtgatacc actgacaaca    180
acaccacca aatctctatt tgcaaatctc aaaggaacaa aaaccagagg gaaactatgc    240
ccaaaatgtc tcaactgcac agatctggac gtggccttgg gcagacccaa atgcacgggg    300
aacatacctt cggcaaaagt ttcaatactc catgaagtaa gacctgttac atctgggtgc    360
tttctataa tgcacgacag aacaaaaatt agacagctgc ccaatcttct cagaggatac    420

```

-continued

```

gaacgtatca gggttatcaaa ccataacgtt atcaatgcag aaaaagcacc aggaggaccc 480
tacaaaaattg gaacctcagg gtcttgccct aacgttacca atggaaacgg attcttcgca 540
acaatggctt gggctgtccc aaaaaacgaa aacaacaaaa cagcaacaaa ttcattaaca 600
atagaagtac catacatttg tacagaagga gaagaccaa ttaccgtttg ggggttccac 660
tctgatagcg aaacccaaat ggcaaaactc tatggagact caaagcctca gaagttcact 720
tcatctgcta acggagtgcac cacacattac gtttcacaga ttggtggctt cccaaatcaa 780
acagaagacg gaggactacc acaaagtggt agaattgttg ttgattacat ggtgcaaaaa 840
tctggaaaaa caggaacaat tacctatcaa agaggatattt tattgcctca aaaagtgtgg 900
tgcgcaagtg gcaggagcaa ggtaataaaa ggatccttgc ctttaattgg agaagcagat 960
tgctccacg aaaaatacgg tggattaaac aaaagcaagc cttactatac aggggaacat 1020
gcaaaagcca taggaaattg cccaatatgg gtgaaaacac ccttgaagct ggccaatgga 1080
accaaata gacctcctgc aaaactatta aaggaaagg gtttcttcgg agctattgct 1140
ggtttcttag aaggaggatg ggaaggaatg attgcaggtt ggcacggata cacatcccat 1200
ggagcacatg gagtagcagt ggcagcagac ctaagagta ctcaagaagc cataaacaag 1260
atcacaaaa atctcaactc tttgagtgcg ctggaagtaa agaactctca aagactaagc 1320
ggagccatgg atgaactcca caacgaaata ctagaactag atgagaaagt ggatgatctc 1380
agagctgata caataagctc gcaaatagaa ctgcagctct tgctttccaa tgaaggaata 1440
ataaacagtg aagatgagca tctcttgcg cttgaaagaa aactgaagaa aatgctgggc 1500
ccctctgctg tagagatagg gaatggatgc ttcgaaacca aacacaagtg caaccagacc 1560
tgctcgata gaatagctgc tggcaccttt aatgcaggag aattttctct cccaccttt 1620
gattcactaa atattactgc tgcacttta aatgacgatg gattggataa tcatactata 1680
ctgctttact actcaactgc tgcttcacgt ttggtgtaa cattgatgat agctatcttt 1740
gttgtttata tggctccag agacaatgtt tctgttcca tctgtctata aggaaagtta 1800
agcccgctat tttcctttat tgtagtactt gtttgcttgt tatcattaca aaaaaacgtt 1860
attgaaaaat gctcttgta ctact 1885

```

<210> SEQ ID NO 30

<211> LENGTH: 1544

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 30

```

agcagagcat cttctcaaaa ctgaagcaaa taggcaaaa tgaacaatgc taccctcaac 60
tatacaaaca ttaacctat ttctcacatc agggggagtg ttattatcac tatatgtgc 120
agcctactg tcatacttac tgtattcgga tatattgcta aaattttcac caacaaaaat 180
aattgcacca acaacgtcgt tggactccgc gaacgcata aattttcagg cctgtaacca 240
ttctgcaaca aaagagatga cattttctct cccagaaccg gaggggacat accctcggtt 300
atcttgccag gggtcaacct ttcaaaaagc actcctaatt agccctcata gattcggaga 360
agccaaagga aactcagctc cttgataat aagggaacct tttattgctt gtggacaaa 420
ggagtgtaaa cactttgctc taaccatta tgcagctcaa ccagggggat actacaatgg 480
aacaagagag gacagaaaca agctgaggca tctgatttca gtcaacttag gcaaaatacc 540

```

-continued

```

aactgtagaa aactccattt tccacatggc agcttgaggt gggtcgcgat gccatgatgg 600
tagagagtgg acttatatcg gagttgatgg ccctgacagt aatgcattga tcaaaataaa 660
atatggagaa gcatacactg acacatacca ttcctatgca aacaacatcc taagaacaca 720
agaaagtgcc tgcaactgca tcgggggaga ttgttatctt atgataactg atgggtcagc 780
ttcaggaatt agtaaatgca gattccttaa gattcgagag ggtcgaatag taaaagaaat 840
atttccaaca ggaagagtag agcactactga agaatgcaca tgcggatttg ccagcaataa 900
aaccatagaa tgtgcttgta gagataacag ttacacagca aaaagaccct ttgtcaaatt 960
aaatgtggaa actgatacag cagaaataag attgatgtgc acagagactt atttggacac 1020
ccccagacca gatgatggaa gcataacagg gccttgcgaa tctaattggg acaaagggag 1080
tggaggtatc aagggaggat ttgtccatca aagaatggca tccaagattg gaagatggta 1140
ctctcgaacg atgtctaaaa ctaaaagaat ggggatggaa ctgtatgtca agtatgatgg 1200
agacctatgg actgacagtg atgcccttgc tctagtggga gtaatggtct caatagaaga 1260
acctggttgg tattctttcg gcttcgaaat aaaagataag aaatgcgatg tcccctgtat 1320
tgggatagag atggtacacg atggtggaaa aacaacttgg cactcagcag caacagccat 1380
ttactgttta atgggctcag gacagttgct atgggacact atcacaggtg ttgatatggc 1440
tctgtaatgg aggaatggtt gaatctgttc taaacccttt gttcctatct tgtttgaaca 1500
attgtcctta ctggacttaa ttgtttctga aaaatgctct tggt 1544

```

<210> SEQ ID NO 31

<211> LENGTH: 1885

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 31

```

agcagaagca gagcattttc taatatccac aaaatgaagg caataattgt actactcatg 60
gtagtaacat ccaatgcaga tcgaatctgc actgggataa catcgtcaaa ctcaccccat 120
gtggtcaaaa ctgctactca aggggaggtc aatgtgactg gtgtgatacc actgacaaca 180
acaccacca aatctcattt tgcaaatctc aaaggaacaa aaaccagagg gaaactatgc 240
ccaaatgcc tcaactgcac agatctggac gtggccttgg gcagaccaa atgcacgggg 300
aacataccct cggcaaaagt ttcaatactc catgaagtca gacctgttac atctgggtgc 360
tttctataa tgcacgacag aacaaaaatt agacagctgc ccaatcttct cagaggatac 420
gaacatatca ggttatcaac tcataacgtt atcaatgcag aaaaggcacc aggaggaccc 480
tacaaaattg gaacctcagg gtcttgccct aacgttacca atggaaacgg atttttcgca 540
acaatggctt gggcgcgtcc aaaaaacgac aacaacaaaa cagcaacaaa ttcattaaca 600
atagaagtac catactttg tacagaagga gaagaccaa ttaccgtttg ggggttcac 660
tctgataacg aagcccaaat ggcaaaactc tatggggact caaagcccca gaagttcac 720
tcatctgcc acggagtgc cacacattac gtttcacaga ttggtggctt cccaaatcaa 780
acagaagacg gaggactacc acaaagtggg agaattgttg ttgattacat ggtgcaaaaa 840
tctgggaaaa caggaacaat tacctatcaa agaggtatct tattgcctca aaaagtgttg 900
tgcgcaagtg gcaggagcaa ggtaataaaa ggatccttgc ctttaatttg agaagcagat 960
tgcctccacg aaaaatcacg tggattaaac aaaagcaagc cttactacac aggggaacat 1020

```

-continued

```

gcaaaggcca taggaaattg cccaatatgg gtgaaaacac ccttgaagct ggccaatgga 1080
accaaataata gacctcctgc aaaactatta aaggaaagag gtttcttcgg agctattgct 1140
ggtttcttag aaggaggatg ggaaggaatg attgcaggtt ggcacggata cacatcccat 1200
ggggcacatg gagtagcagt ggcagcagac cttaaagata ctcaagaagc cataaacaag 1260
ataacaaaaa atctcaactc tttgagttag ctggaagtaa agaactttca aagactaagc 1320
ggtgccatgg atgaactcca caacgaaata ctagaactag acgagaaagt ggatgatctc 1380
agagctgata caataagctc acaaatagaa ctgcagctct tgctttccaa tgaaggata 1440
ataaacagtg aagatgagca tctcttggcg cttgaaagaa agctgaagaa aatgctgggc 1500
ccctctgctg tagagatagg gaatggatgc ttcgaaacca aacacaagtg caaccagacc 1560
tgtctcgaca gaatagctgc tggtagcttt gatgcaggag aattttctct cccactttt 1620
gattcactga atattactgc tgcacttta aatgacgatg gattggataa tcatactata 1680
ctgctttact actcaactgc tgcctcagc ttggctgtaa cattgatgat agctatcttt 1740
gttgtttata tggctccag agacaatgtt tcttgctcca tctgtctata aggaaagtta 1800
agccctgtat tttcctttat tgtagtgctt gtttgcttgt taccattaca aaaaaacgtt 1860
attgaaaaat gctcttgcta ctact 1885

```

<210> SEQ ID NO 32

<211> LENGTH: 1557

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 32

```

agcagaagca gagcatcttc tcaaaactga ggcaaatagg ccaaaaatga acaatgctac 60
cttcaactat acaaacgtta accctatttc tcacatcagg gggagtatta ttatcactat 120
atgtgtcagc ttcattgtca tacttactat attcggatat attgctaaaa ttctcaccaa 180
cagaataaac tgcaccaaca atgccattgg attgtgcaaa cgcacaaat gttcaggctg 240
tgaaccgttc tgcaacaaaa ggggtgacac ttcttctccc agaaccagag tggacatacc 300
cgcgtttatc ttgccggggt tcaacctttc agaaagcact cctaattagc cctcatagat 360
tcggagaaac caagggaaac tcagctccct tgataataag ggaacctttt attgcttggtg 420
gaccaaagga atgcaaacac tttgctctaa ccattatgc agcccaacca gggggatact 480
acaatggaac aagaggagac agaaacaagc tgaggcatct aatttcagtc aaattgggca 540
aaatcccaac agtagaaaac tccattttcc acatggcagc atggagcggg tccgcattgc 600
atgatggtaa agaattgaca tatatcggag ttgatggccc tgacaataat gcattgctca 660
aaataaaata tggagaagca tatactgaca cataccattc ctatgcaaac aacatcctaa 720
gaacacaaga aagtgcctgc aattgcatcg ggggaaattg ttatcttatg ataactgatg 780
gctcagcttc aggtattagt gaatgcagat ttcttaaaat tcgagagggc cgaataataa 840
aagaaatatt tccaacagga agagtaaaac atactgaaga atgcacatgc ggatttgcca 900
gcaataagac catagaatgt gcctgtagag ataacagtta cacagcaaaa agaccctttg 960
tcaaattaaa cgtggagact gatacagcag aaataagatt gatgtgcaca gagacttatt 1020
tggacacccc cagaccagat gatggaagca taacagggcc ttgtgaatct aatggggaca 1080
aaggagtggt aggcacaaag ggaggatttg ttcataaaag aatggcatcc aagattggaa 1140

```

-continued

```

ggtggtactc tcgaacgatg tctaaaacta aaaggatggg gatgggactg tatgtcaagt 1200
atgatggaga cccatgggct gacagtgatg cccttgctct tagtgagta atgggttcaa 1260
tggaagaacc tggttggtac tcctttggct tcgaaataaa agataagaaa tgtgatgtcc 1320
cctgtattgg aatagagatg gtacatgatg gtggaaaaga gacttggcac tcagcagcaa 1380
cagccattta ctgtttaatg ggctcaggac agctgctgtg ggacactgtc acaggtgttg 1440
atatggctct gtaatggagg aatggttgag tctgttctaa accctttgtt cctattttgt 1500
ttgaacaatt gtccttactg aacttaattg tttctgaaaa atgctcttgt tactact 1557

```

<210> SEQ ID NO 33

<211> LENGTH: 1853

<212> TYPE: DNA

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 33

```

tctaatatcc aaaaaatgaa ggcaataatt gtactactca tggtagtaac atccaatgca 60
gatcgaatct gactgggat aacatcttca aactcacctc atgtggtcaa aacagctact 120
caaggggagg tcaatgtgac tgggtgaata ccactgacaa caacaccaac aaaatcttat 180
tttgcaaatc tcaaaggaa aaggaccaga gggaaactat gtccagactg tctcaactgt 240
acagatctgg atgtggcctt gggcagacca atgtgtgtgg ggaccacacc ttgggcaaaa 300
gcttcaatac tccacgaagt cagacctgtt acatccgggt gctttcctat aatgcacgac 360
agaacaaaaa tcagacaact acccaatctt ctcagaggat atgaaaatat cagattatca 420
acccaaaaacg ttatcgatgc agaaaatgca ccaggaggac cctacagact tggaacctca 480
ggatcttgcc ctaacgctac cagtaaaagc ggatttttcg caacaatggc ttgggctgtc 540
ccaaaggaca acaacaaaaa tgcaacgaac cactaacag tagaagtacc atacgtttgt 600
acagaagggg aagaccaaat tactgtttgg gggttccatt cagataacaa aaccccaatg 660
aagaacctct atggagactc aaatcctcaa aagttcacct catctgctaa tggagtaacc 720
acacattatg tttctcagat tggcggtctc ccagctcaa cagaagacga aggactacca 780
caaagcggca gaattgttgt tgattacatg gtgcaaaaac ctaggaaaac aggaacaatt 840
gtctatcaaa gaggtgtttt gttgcctcaa aaggtgtggt gcgcgagtgg caggagcaaa 900
gtaataaaaag ggtccttgcc tttaattggt gaagcagatt gccttcatga aaaatacggg 960
ggattaacaa aaagcaagcc ttactacaca ggagaacatg caaaagccat aggaaattgc 1020
ccaatatggg tgaaaacacc tttgaagctt gccaatggaa ccaaatatag acctcctgca 1080
aaactattaa aggaaagggg tttcttcgga gctattgctg gtttcctaga aggaggatgg 1140
gaaggaatga ttgcaggttg gcacggatgc acatctcacg gagcacatgg agtggcagtg 1200
gcgcgagacc ttaaggtac gcaagaagct ataaacaaga taacaaaaaa tctcaattct 1260
ttgagtgagc tagaagtaaa gaatcttcaa agactaagtg gtgccatgga tgaactccac 1320
aacgaaatac tcgagctgga tgagaaagtg gatgatctca gagctgacac tataagctcg 1380
caaatagaac ttgcagtctt gctttccaat gaaggaataa taaacagtga agatgagcat 1440
ctattggcac ttgagagaaa actaaagaaa atgctgggtc cctctgctgt agacatagga 1500
aatggatgct tcgaaaccaa acacaagtgc aaccagacct gcttagacag gatagctgct 1560
ggcaccttta atgcaggaga attttctctc ccacttttg attcactgaa cattactgct 1620

```

-continued

```

gcacatctttaa atgatgatgg attggataac catactatac tgctctatta ctcaactgct 1680
gcttctagtgt tggctgtaac attgatgcta gctatctttaa ttgtttatat ggtctccaga 1740
gacaacgttt catgctccat ctgtctataa ggaagattaa gccttgattt ttcctttatt 1800
gtagtgtcttg ttgtctgtgc atcattacaa agaaacgtta ttgaaaaatg ctc 1853

```

```

<210> SEQ ID NO 34
<211> LENGTH: 1529
<212> TYPE: DNA
<213> ORGANISM: Influenza B virus

```

```

<400> SEQUENCE: 34

```

```

tctcaaaact gaggcaaata ggccaaaat gaacaatgct accctcaact atacaaacgt 60
taacctatt cctcacatca gggggagtgt tattatcact atatgtgtca gcttcactgt 120
catacttact atattcggtat atattgctaa aattttcaac aacagaaata actgcaccaa 180
caatgccatt ggattgtgca aacgcaccaa atgttcaggc tgtgaaccgt tctgcaacaa 240
aaggggtgac acttcttctc ccagaaccgg agtggacata cccgcgttta tcttgccggg 300
gctcaacctt tcagaaagca ctccctaatta gccctcatag attcgagaa accaaaggaa 360
actcagctcc cttgataata agggaacctt ttattgcttg tggaccaaag gaatgcaaac 420
actttgctct aacctattat gcagcccaac cagggggata ctacaatgga acaaaagaag 480
acagaaacaa gctgaggcat ctaatttcag tcaaattggg caaaatccca acagtagaaa 540
actccatttt ccacatggca gcatggagcg ggtccgcatg ccatgatggt aaagaatgga 600
catatatcgg agttgatggc cctgacagta atgcattgct caaaataaaa tatggagaag 660
catatactga cacataccat tcctatgcaa acaacatcct aagaacacaa gaaagtgcct 720
gcaattgcat cgggggaaat tgttatctta tgataactga tggctcagct tcagggtatta 780
gtgagtgcag atttcttaag attcgagagg gccgaataat aaaagaaata tttccaacag 840
gaagagtaaa acatactgaa gaatgcacat gcggatttgc cagcaataaa accatagaat 900
gtgcctgtag agataacagt tacacagcaa aaagaccctt tgtcaaatta aatgtggaga 960
ctgatacagc agaaataaga ttgatgtgca cagagactta tttggacacc ccagaccag 1020
atgatggaag cataacaggg ccttgtgaat ctaatgggaa taaagggagt ggaggcatca 1080
agggaggatt tgttcatcaa agaatggcat ccaaattgg aaggtggtac tctcgaacaa 1140
tgtctaaaac caaaaggatg ggaatgggac tgtatgtcaa gtatgatgga gacctatgga 1200
ctgacagtga tgcccttgct cttagtggag taatggtttc aatggaagaa cctggttggg 1260
actcatttgg cttcgaaata aaagataaga aatgtgatgt ccctgtatt gggatagaga 1320
tgggtacatga tgggtgaaag gagacttggc actcagcagc aacagccatt tactgtttaa 1380
tgggctcagg acaactgttg tgggacactg tcacaggtgt tgatattggt ctgtaatggg 1440
ggaatggttg agtctgttct aaacccttgg ttcctatctt gtttgaacaa ttgtccttgc 1500
tgaacttaat tgtttctgaa aaatgctct 1529

```

```

<210> SEQ ID NO 35
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 35

```

-continued

Met	Lys	Thr	Ile	Ile	Ala	Leu	Ser	Tyr	Ile	Leu	Cys	Leu	Val	Phe	Ala	1	5	10	15
Gln	Lys	Leu	Pro	Gly	Asn	Asp	Asn	Ser	Thr	Ala	Thr	Leu	Cys	Leu	Gly	20	25	30	
His	His	Ala	Val	Pro	Asn	Gly	Thr	Leu	Val	Lys	Thr	Ile	Thr	Asn	Asp	35	40	45	
Gln	Ile	Glu	Val	Thr	Asn	Ala	Thr	Glu	Leu	Val	Gln	Ser	Ser	Ser	Thr	50	55	60	
Gly	Arg	Ile	Cys	Gly	Ser	Pro	His	Arg	Ile	Leu	Asp	Gly	Lys	Asn	Cys	65	70	75	80
Thr	Leu	Ile	Asp	Ala	Leu	Leu	Gly	Asp	Pro	His	Cys	Asp	Gly	Phe	Gln	85	90	95	
Asn	Lys	Glu	Trp	Asp	Leu	Phe	Val	Glu	Arg	Ser	Lys	Ala	Tyr	Ser	Asn	100	105	110	
Cys	Tyr	Pro	Tyr	Asp	Val	Pro	Asp	Tyr	Ala	Ser	Leu	Arg	Ser	Leu	Val	115	120	125	
Ala	Ser	Ser	Gly	Thr	Leu	Glu	Phe	Ile	Asn	Glu	Asp	Phe	Asn	Trp	Thr	130	135	140	
Gly	Val	Ala	Gln	Asp	Gly	Gly	Ser	Tyr	Ala	Cys	Lys	Arg	Gly	Ser	Val	145	150	155	160
Asn	Ser	Phe	Phe	Ser	Arg	Leu	Asn	Trp	Leu	His	Lys	Leu	Glu	Tyr	Lys	165	170	175	
Tyr	Pro	Ala	Leu	Asn	Val	Thr	Met	Pro	Asn	Asn	Gly	Lys	Phe	Asp	Lys	180	185	190	
Leu	Tyr	Ile	Trp	Gly	Val	His	His	Pro	Ser	Thr	Asp	Ser	Asp	Gln	Thr	195	200	205	
Ser	Leu	Tyr	Val	Arg	Ala	Ser	Gly	Arg	Val	Thr	Val	Ser	Thr	Lys	Arg	210	215	220	
Ser	Gln	Gln	Thr	Val	Thr	Pro	Asn	Ile	Gly	Ser	Arg	Pro	Trp	Val	Arg	225	230	235	240
Gly	Gln	Ser	Ser	Arg	Ile	Ser	Ile	Tyr	Trp	Thr	Ile	Val	Lys	Pro	Gly	245	250	255	
Asp	Ile	Leu	Leu	Ile	Asn	Ser	Thr	Gly	Asn	Leu	Ile	Ala	Pro	Arg	Gly	260	265	270	
Tyr	Phe	Lys	Ile	Arg	Asn	Gly	Lys	Ser	Ser	Ile	Met	Arg	Ser	Asp	Ala	275	280	285	
Pro	Ile	Gly	Asn	Cys	Ser	Ser	Glu	Cys	Ile	Thr	Pro	Asn	Gly	Ser	Ile	290	295	300	
Pro	Asn	Asp	Lys	Pro	Phe	Gln	Asn	Val	Asn	Arg	Ile	Thr	Tyr	Gly	Ala	305	310	315	320
Cys	Pro	Arg	Tyr	Val	Lys	Gln	Asn	Thr	Leu	Lys	Leu	Ala	Thr	Gly	Met	325	330	335	
Arg	Asn	Val	Pro	Glu	Lys	Gln	Thr	Arg	Gly	Ile	Phe	Gly	Ala	Ile	Ala	340	345	350	
Gly	Phe	Ile	Glu	Asn	Gly	Trp	Glu	Gly	Met	Val	Asp	Gly	Trp	Tyr	Gly	355	360	365	
Phe	Arg	His	Gln	Asn	Ser	Glu	Gly	Thr	Gly	Gln	Ala	Ala	Asp	Leu	Lys	370	375	380	
Ser	Thr	Gln	Ala	Ala	Ile	Asp	Gln	Ile	Asn	Gly	Lys	Leu	Asn	Arg	Leu	385	390	395	400

-continued

```

Ile Glu Lys Thr Asn Glu Lys Phe His Gln Ile Glu Lys Glu Phe Ser
405                      410                      415

Glu Val Glu Gly Arg Ile Gln Asp Leu Glu Lys Tyr Val Glu Asp Thr
420                      425                      430

Lys Ile Asp Leu Trp Ser Tyr Asn Ala Glu Leu Leu Val Ala Leu Glu
435                      440                      445

Asn Gln His Thr Ile Asp Leu Thr Asp Ser Glu Met Asn Lys Leu Phe
450                      455                      460

Glu Lys Thr Arg Lys Gln Leu Arg Glu Asn Ala Glu Asp Met Gly Asn
465                      470                      475                      480

Gly Cys Phe Lys Ile Tyr His Lys Cys Asp Asn Ala Cys Ile Gly Ser
485                      490                      495

Ile Arg Asn Gly Thr Tyr Asp His Asp Val Tyr Arg Asp Glu Ala Leu
500                      505                      510

Asn Asn Arg Phe Gln Ile Lys Gly Val Glu Leu Lys Ser Gly Tyr Lys
515                      520                      525

Asp Trp Ile Leu Trp Ile Ser Phe Ala Ile Ser Cys Phe Leu Leu Cys
530                      535                      540

Val Val Leu Leu Gly Phe Ile Met Trp Ala Cys Gln Lys Gly Asn Ile
545                      550                      555                      560

Arg Cys Asn Ile Cys Ile
565

```

<210> SEQ ID NO 36

<211> LENGTH: 436

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 36

```

Met Gln Ile Ala Ile Leu Val Thr Thr Val Thr Leu His Phe Lys Gln
1                      5                      10                      15

Tyr Glu Cys Asn Ser Pro Pro Asn Asn Gln Val Met Leu Cys Glu Pro
20                      25                      30

Thr Ile Ile Glu Arg Asn Ile Thr Glu Ile Val Tyr Leu Thr Asn Thr
35                      40                      45

Thr Ile Glu Lys Glu Ile Cys Pro Lys Leu Ala Glu Tyr Arg Asn Trp
50                      55                      60

Ser Lys Pro Gln Cys Lys Ile Thr Gly Phe Ala Pro Phe Ser Lys Asp
65                      70                      75                      80

Asn Ser Ile Arg Leu Ser Ala Gly Gly Asp Ile Trp Val Thr Arg Glu
85                      90                      95

Pro Tyr Val Ser Cys Asp Pro Gly Lys Cys Tyr Gln Phe Ala Leu Gly
100                     105                     110

Gln Gly Thr Thr Leu Asn Asn Arg His Ser Asn Asp Thr Val His Asp
115                     120                     125

Arg Thr Pro Tyr Arg Thr Leu Leu Met Asn Glu Leu Gly Val Pro Phe
130                     135                     140

His Leu Gly Thr Lys Gln Val Cys Ile Ala Trp Ser Ser Ser Ser Cys
145                     150                     155                     160

His Asp Gly Lys Ala Trp Leu His Val Cys Val Thr Gly His Asp Glu
165                     170                     175

Asn Ala Thr Ala Ser Phe Ile Tyr Asp Gly Arg Leu Val Asp Ser Ile
180                     185                     190

```

-continued

Gly Ser Trp Ser Lys Asn Ile Leu Arg Thr Gln Glu Ser Glu Cys Val
 195 200 205
 Cys Ile Asn Gly Thr Cys Thr Val Val Met Thr Asp Gly Ser Ala Ser
 210 215 220
 Glu Arg Ala Asp Thr Lys Ile Leu Phe Ile Glu Glu Gly Lys Ile Val
 225 230 235 240
 His Ile Ser Pro Leu Ser Gly Ser Ala Gln His Val Glu Glu Cys Ser
 245 250 255
 Cys Tyr Pro Arg Tyr Pro Gly Val Arg Cys Val Cys Arg Asp Asn Trp
 260 265 270
 Lys Gly Ser Asn Arg Pro Ile Val Asp Ile Asn Val Lys Asp Tyr Ser
 275 280 285
 Ile Val Ser Ser Tyr Val Cys Ser Gly Leu Val Gly Asp Thr Pro Arg
 290 295 300
 Lys Asn Asp Ser Ser Ser Ser Ser Tyr Cys Arg Asn Pro Asn Asn Glu
 305 310 315 320
 Lys Gly Ser His Gly Val Lys Gly Trp Ala Phe Asp Asp Gly Asn Asp
 325 330 335
 Val Trp Met Gly Arg Thr Ile Ser Glu Glu Leu Arg Ser Gly Tyr Glu
 340 345 350
 Thr Phe Lys Val Ile Gly Gly Trp Ser Lys Pro Asn Ser Lys Leu Gln
 355 360 365
 Ile Asn Arg Gln Val Ile Val Asp Arg Gly Asn Arg Ser Gly Tyr Ser
 370 375 380
 Gly Ile Phe Ser Val Glu Gly Lys Ser Cys Ile Asn Arg Cys Phe Tyr
 385 390 395 400
 Val Glu Leu Ile Arg Gly Arg Lys Gln Glu Thr Glu Val Trp Trp Thr
 405 410 415
 Ser Asn Ser Ile Val Val Phe Cys Gly Thr Ser Gly Thr Tyr Gly Thr
 420 425 430
 Gly Ser Trp Pro
 435

<210> SEQ ID NO 37
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 37

Met Lys Thr Ile Ile Ala Leu Ser Tyr Ile Leu Cys Leu Val Phe Ala
 1 5 10 15
 Gln Lys Leu Pro Gly Asn Asp Asn Ser Thr Ala Thr Leu Cys Leu Gly
 20 25 30
 His His Ala Val Pro Asn Gly Thr Leu Val Lys Thr Ile Thr Asn Asp
 35 40 45
 Gln Ile Glu Val Thr Asn Ala Thr Glu Leu Val Gln Ser Ser Pro Thr
 50 55 60
 Gly Arg Ile Cys Asp Ser Pro His Arg Ile Leu Asp Gly Lys Asn Cys
 65 70 75 80
 Thr Leu Ile Asp Ala Leu Leu Gly Asp Pro His Cys Asp Gly Phe Gln
 85 90 95
 Asn Lys Glu Trp Asp Leu Phe Val Glu Arg Ser Lys Ala Tyr Ser Asn

-continued

100					105					110					
Cys 115	Tyr	Pro	Tyr	Asp	Val 120	Pro	Asp	Tyr	Ala	Ser 125	Leu	Arg	Ser	Leu	Val
Ala 130	Ser	Ser	Gly	Thr	Leu 135	Glu	Phe	Ile	Asn	Glu 140	Asn	Phe	Asn	Trp	Thr
Gly 145	Val	Ala	Gln	Asp	Gly 150	Lys	Ser	Tyr	Ala	Cys 155	Lys	Arg	Gly	Ser	Val 160
Asn 165	Ser	Phe	Phe	Ser	Arg 170	Leu	Asn	Trp	Leu	His 175	Lys	Leu	Glu	Tyr	Lys
Tyr 180	Pro	Ala	Leu	Asn	Val 185	Thr	Met	Pro	Asn	Asn 190	Gly	Lys	Phe	Asp	Lys
Leu 195	Tyr	Ile	Trp	Gly	Val 200	His	His	Pro	Ser	Thr 205	Asp	Ser	Val	Gln	Thr
Ser 210	Leu	Tyr	Val	Arg	Ala 215	Ser	Gly	Arg	Val	Thr 220	Val	Ser	Thr	Lys	Arg
Ser 225	Gln	Gln	Thr	Val	Ile 230	Pro	Asp	Ile	Gly	Tyr 235	Arg	Pro	Trp	Val	Arg 240
Gly 245	Gln	Ser	Ser	Arg	Ile 250	Ser	Ile	Tyr	Trp	Thr 255	Ile	Val	Lys	Pro	Gly
Asp 260	Ile	Leu	Leu	Ile	Asn 265	Ser	Thr	Gly	Asn	Leu 270	Ile	Ala	Pro	Arg	Gly
Tyr 275	Phe	Lys	Ile	Arg	Asn 280	Gly	Lys	Ser	Ser	Ile 285	Met	Arg	Ser	Asp	Ala
Pro 290	Ile	Gly	Asn	Cys	Ser 295	Ser	Glu	Cys	Ile	Thr 300	Pro	Asn	Gly	Ser	Ile
Pro 305	Asn	Asp	Lys	Pro	Phe 310	Gln	Asn	Val	Asn	Arg 315	Ile	Thr	Tyr	Gly	Ala 320
Cys 325	Pro	Arg	Tyr	Val	Lys 330	Gln	Asn	Thr	Leu	Lys 335	Leu	Ala	Thr	Gly	Met
Arg 340	Asn	Val	Pro	Glu	Lys 345	Gln	Thr	Arg	Gly	Ile 350	Phe	Gly	Ala	Ile	Ala
Gly 355	Phe	Ile	Glu	Asn	Gly 360	Trp	Glu	Gly	Met	Val 365	Asp	Gly	Trp	Tyr	Gly
Phe 370	Arg	His	Gln	Asn	Ser 375	Glu	Gly	Thr	Gly	Gln 380	Ala	Ala	Asp	Leu	Lys
Ser 385	Thr	Gln	Ala	Ala	Ile 390	Asp	Gln	Ile	Asn	Gly 395	Lys	Leu	Asn	Arg	Leu 400
Val 405	Glu	Lys	Thr	Asn	Glu 410	Lys	Phe	His	Gln	Ile 415	Glu	Lys	Glu	Phe	Ser
Glu 420	Val	Glu	Gly	Arg	Ile 425	Gln	Asp	Leu	Glu	Lys 430	Tyr	Val	Glu	Asp	Thr
Lys 435	Ile	Asp	Leu	Trp	Ser 440	Tyr	Asn	Ala	Glu	Leu 445	Leu	Val	Ala	Leu	Glu
Asn 450	Gln	His	Thr	Ile	Asp 455	Leu	Thr	Asp	Ser	Glu 460	Met	Asn	Lys	Leu	Phe
Glu 465	Arg	Thr	Arg	Lys	Gln 470	Leu	Arg	Glu	Asn	Ala 475	Glu	Asp	Met	Gly	Asn 480
Gly 485	Cys	Phe	Lys	Ile	Tyr 490	His	Lys	Cys	Asp	Asn 495	Ala	Cys	Ile	Gly	Ser
Ile 500	Arg	Asn	Gly	Thr	Tyr 505	Asp	His	Asp	Val	Tyr 510	Arg	Asp	Glu	Ala	Leu

-continued

```

Asn Asn Arg Phe Gln Ile Lys Gly Val Glu Leu Lys Ser Gly Tyr Lys
515                520                525

Asp Trp Ile Leu Trp Ile Ser Phe Ala Ile Ser Cys Phe Leu Leu Cys
530                535                540

Val Val Leu Leu Gly Phe Ile Met Trp Ala Cys Gln Lys Gly Asn Ile
545                550                555                560

Arg Cys Asn Ile Cys Ile
565

```

```

<210> SEQ ID NO 38
<211> LENGTH: 451
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

```

```

<400> SEQUENCE: 38

```

```

Lys Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Val Ser Leu
1              5              10              15

Thr Ile Ala Thr Ile Cys Phe Leu Met Gln Ile Ala Ile Leu Val Thr
20            25            30

Thr Val Thr Leu His Phe Lys Gln Tyr Glu Cys Asn Ser Pro Pro Asn
35            40            45

Asn Gln Val Met Leu Cys Glu Pro Thr Ile Ile Glu Arg Asn Ile Thr
50            55            60

Glu Ile Val Tyr Leu Thr Asn Thr Thr Ile Glu Lys Glu Ile Cys Pro
65            70            75            80

Lys Leu Ala Glu Tyr Arg Asn Trp Ser Lys Pro Gln Cys Lys Ile Thr
85            90            95

Gly Phe Ala Pro Phe Ser Lys Asp Asn Ser Ile Arg Leu Ser Ala Gly
100           105           110

Gly Asp Ile Trp Val Thr Arg Glu Pro Tyr Val Ser Cys Asp Pro Gly
115           120           125

Lys Cys Tyr Gln Phe Ala Leu Gly Gln Gly Thr Thr Leu Asn Asn Arg
130           135           140

His Ser Asn Asp Thr Val His Asp Arg Thr Pro Tyr Arg Thr Leu Leu
145           150           155           160

Met Asn Glu Leu Gly Val Pro Phe His Leu Gly Thr Lys Gln Val Cys
165           170           175

Ile Ala Trp Ser Ser Ser Ser Cys His Asp Gly Lys Ala Trp Leu His
180           185           190

Val Cys Val Thr Gly His Asp Glu Asn Ala Thr Ala Ser Phe Ile Tyr
195           200           205

Asp Gly Arg Leu Val Asp Ser Ile Gly Ser Trp Ser Lys Asn Ile Leu
210           215           220

Arg Thr Gln Glu Ser Glu Cys Val Cys Ile Asn Gly Thr Cys Thr Val
225           230           235           240

Val Met Thr Asp Gly Ser Ala Ser Glu Arg Ala Asp Thr Lys Ile Leu
245           250           255

Phe Ile Glu Glu Gly Lys Ile Val His Ile Ser Pro Leu Ser Gly Ser
260           265           270

Ala Gln His Val Glu Glu Cys Ser Cys Tyr Pro Arg Tyr Pro Gly Val
275           280           285

Arg Cys Val Cys Arg Asp Asn Trp Lys Gly Ser Asn Arg Pro Ile Val

```

-continued

290	295	300
Asp Ile Asn Val Lys	Asp Tyr Ser Ile Val	Ser Ser Tyr Val Cys Ser
305	310	315 320
Gly Leu Val Gly Asp	Thr Pro Arg Lys Asn	Asp Ser Ser Ser Ser
325	330	335
Tyr Cys Trp Asn Pro	Asn Asn Glu Lys Gly	Gly His Gly Val Lys Gly
340	345	350
Trp Ala Phe Asp Asp	Gly Asn Asp Val Trp	Met Gly Arg Thr Ile Ser
355	360	365
Glu Glu Leu Arg Ser	Gly Tyr Glu Thr Phe	Lys Val Ile Gly Gly Trp
370	375	380
Ser Lys Pro Asn Ser	Lys Leu Gln Ile Asn	Arg Gln Val Ile Val Asp
385	390	395 400
Arg Gly Asn Arg Ser	Gly Tyr Ser Gly Ile	Phe Ser Val Glu Gly Lys
405	410	415
Ser Cys Ile Asn Arg	Cys Phe Tyr Val Glu	Leu Ile Arg Gly Arg Lys
420	425	430
Gln Glu Thr Glu Val	Trp Trp Thr Ser Asn	Ser Ile Val Val Phe Cys
435	440	445
Gly Thr Ser		
450		

<210> SEQ ID NO 39

<211> LENGTH: 566

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 39

Met Lys Thr Ile Ile	Ala Leu Ser Tyr Ile	Leu Cys Leu Val Phe Ala
1	5	10 15
Gln Lys Leu Pro Gly	Asn Asp Asn Ser Thr	Ala Thr Leu Cys Leu Gly
20	25	30
His His Ala Val Pro	Asn Gly Thr Leu Val	Lys Thr Ile Thr Asn Asp
35	40	45
Gln Ile Glu Val Thr	Asn Ala Thr Glu Leu	Val Gln Ser Ser Ser Thr
50	55	60
Gly Arg Ile Cys Asp	Ser Pro His Arg Ile	Leu Asp Gly Lys Asn Cys
65	70	75 80
Thr Leu Ile Asp Ala	Leu Leu Gly Asp Pro	His Cys Asp Gly Phe Gln
85	90	95
Asn Lys Glu Trp Asp	Leu Phe Val Glu Arg	Ser Lys Ala Tyr Ser Asn
100	105	110
Cys Tyr Pro Tyr Asp	Val Pro Asp Tyr Ala	Ser Leu Arg Ser Leu Val
115	120	125
Ala Ser Ser Gly Thr	Leu Glu Phe Thr Asn	Glu Gly Phe Asn Trp Thr
130	135	140
Gly Val Ala Gln Asp	Gly Thr Ser Tyr Ala	Cys Lys Arg Gly Ser Val
145	150	155 160
Lys Ser Phe Phe Ser	Arg Leu Asn Trp Leu	His Lys Leu Glu Tyr Lys
165	170	175
Tyr Pro Ala Leu Asn	Val Thr Met Pro Asn	Asn Asp Lys Phe Asp Lys
180	185	190

-continued

Leu	Tyr	Ile	Trp	Gly	Val	His	His	Pro	Ser	Thr	Asp	Ser	Asp	Gln	Thr	195	200	205
Ser	Ile	Tyr	Val	Gln	Ala	Ser	Gly	Arg	Val	Thr	Val	Ser	Thr	Lys	Arg	210	215	220
Ser	Gln	Gln	Thr	Val	Ile	Pro	Asn	Ile	Gly	Ser	Arg	Pro	Trp	Val	Arg	225	230	235
Gly	Ile	Ser	Ser	Arg	Ile	Ser	Ile	Tyr	Trp	Thr	Ile	Val	Lys	Pro	Gly	245	250	255
Asp	Ile	Leu	Leu	Ile	Asn	Ser	Thr	Gly	Asn	Leu	Ile	Ala	Pro	Arg	Gly	260	265	270
Tyr	Phe	Lys	Ile	Arg	Ser	Gly	Lys	Ser	Ser	Ile	Met	Arg	Ser	Asp	Ala	275	280	285
Pro	Ile	Gly	Asn	Cys	Asn	Ser	Glu	Cys	Ile	Thr	Pro	Asn	Gly	Ser	Ile	290	295	300
Pro	Asn	Asp	Lys	Pro	Phe	Gln	Asn	Val	Asn	Arg	Ile	Thr	Tyr	Gly	Ala	305	310	315
Cys	Pro	Arg	Tyr	Val	Lys	Gln	Asn	Thr	Leu	Lys	Leu	Ala	Thr	Gly	Met	325	330	335
Arg	Asn	Val	Pro	Glu	Lys	Gln	Thr	Arg	Gly	Ile	Phe	Gly	Ala	Ile	Ala	340	345	350
Gly	Phe	Ile	Glu	Asn	Gly	Trp	Glu	Gly	Met	Val	Asp	Gly	Trp	Tyr	Gly	355	360	365
Phe	Arg	His	Gln	Asn	Ser	Glu	Gly	Thr	Gly	Gln	Ala	Ala	Asp	Leu	Lys	370	375	380
Ser	Thr	Gln	Ala	Ala	Ile	Asn	Gln	Ile	Asn	Gly	Lys	Leu	Asn	Arg	Leu	385	390	395
Ile	Glu	Lys	Thr	Asn	Glu	Lys	Phe	His	Gln	Ile	Glu	Lys	Glu	Phe	Ser	405	410	415
Glu	Val	Glu	Gly	Arg	Ile	Gln	Asp	Leu	Glu	Lys	Tyr	Val	Glu	Asp	Thr	420	425	430
Lys	Ile	Asp	Leu	Trp	Ser	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Ala	Leu	Glu	435	440	445
Asn	Gln	His	Thr	Ile	Asp	Leu	Thr	Asp	Ser	Glu	Met	Asn	Lys	Leu	Phe	450	455	460
Glu	Arg	Thr	Arg	Lys	Gln	Leu	Arg	Glu	Asn	Ala	Glu	Asp	Met	Gly	Asn	465	470	475
Gly	Cys	Phe	Lys	Ile	Tyr	His	Lys	Cys	Asp	Asn	Ala	Cys	Ile	Gly	Ser	485	490	495
Ile	Arg	Asn	Gly	Thr	Tyr	Asp	His	Asp	Val	Tyr	Arg	Asp	Glu	Ala	Leu	500	505	510
Asn	Asn	Arg	Phe	Gln	Ile	Lys	Gly	Val	Glu	Leu	Lys	Ser	Gly	Tyr	Lys	515	520	525
Asp	Trp	Ile	Leu	Trp	Ile	Ser	Phe	Ala	Ile	Ser	Cys	Phe	Leu	Leu	Cys	530	535	540
Val	Val	Leu	Leu	Gly	Phe	Ile	Met	Trp	Ala	Cys	Gln	Lys	Gly	Asn	Ile	545	550	555
Arg	Cys	Asn	Ile	Cys	Ile											565		

<210> SEQ ID NO 40

<211> LENGTH: 469

<212> TYPE: PRT

-continued

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 40

```

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Val Ser Leu Thr
 1           5           10           15

Ile Ala Thr Ile Cys Phe Leu Met Gln Ile Ala Ile Leu Val Thr Thr
20           25           30

Val Thr Leu His Phe Lys Gln Tyr Glu Cys Asn Ser Pro Pro Asn Asn
35           40           45

Gln Val Met Leu Cys Glu Pro Thr Ile Ile Glu Arg Asn Ile Thr Glu
50           55           60

Ile Val Tyr Leu Thr Asn Thr Thr Ile Glu Lys Glu Ile Cys Pro Lys
65           70           75           80

Leu Ala Glu Tyr Arg Asn Trp Ser Lys Pro Gln Cys Lys Ile Thr Gly
85           90           95

Phe Ala Pro Phe Ser Lys Asp Asn Ser Ile Arg Leu Ser Ala Gly Gly
100          105          110

Asp Ile Trp Val Thr Arg Glu Pro Tyr Val Ser Cys Asp Pro Asp Lys
115          120          125

Cys Tyr Gln Phe Ala Leu Gly Gln Gly Thr Thr Leu Asn Asn Arg His
130          135          140

Ser Asn Asp Thr Val His Asp Arg Thr Pro Tyr Arg Thr Leu Leu Met
145          150          155          160

Asn Glu Leu Gly Val Pro Phe His Leu Gly Thr Lys Gln Val Cys Ile
165          170          175

Ala Trp Ser Ser Ser Ser Cys His Asp Gly Lys Ala Trp Leu His Val
180          185          190

Cys Val Thr Gly His Asp Glu Asn Ala Thr Ala Ser Phe Ile Tyr Asp
195          200          205

Gly Arg Leu Val Asp Ser Ile Gly Ser Trp Ser Lys Lys Ile Leu Arg
210          215          220

Thr Gln Glu Ser Glu Cys Val Cys Ile Asn Gly Thr Cys Thr Val Val
225          230          235          240

Met Thr Asp Gly Ser Ala Ser Gly Arg Ala Asp Thr Lys Ile Leu Phe
245          250          255

Ile Glu Glu Gly Lys Ile Val His Ile Ser Pro Leu Ser Gly Ser Ala
260          265          270

Gln His Val Glu Glu Cys Ser Cys Tyr Pro Arg Tyr Ser Gly Val Arg
275          280          285

Cys Val Cys Arg Asp Asn Trp Lys Gly Ser Asn Arg Pro Ile Val Asp
290          295          300

Ile Asn Val Lys Asp Tyr Ser Ile Val Ser Ser Tyr Val Cys Ser Gly
305          310          315          320

Leu Val Gly Asp Thr Pro Arg Lys Asn Asp Ser Ser Ser Ser Ser His
325          330          335

Cys Leu Asn Pro Asn Asn Glu Glu Gly Gly His Gly Val Lys Gly Trp
340          345          350

Ala Phe Asp Asp Gly Asn Asp Val Trp Met Gly Arg Thr Ile Ser Glu
355          360          365

Lys Leu Arg Ser Gly Tyr Glu Thr Phe Lys Val Ile Gly Gly Trp Ser
370          375          380

```

-continued

Lys Pro Asn Ser Lys Leu Gln Ile Asn Arg Gln Val Ile Val Asp Arg
 385 390 395 400
 Gly Asn Arg Ser Gly Tyr Ser Gly Ile Phe Ser Val Glu Gly Lys Ser
 405 410 415
 Cys Ile Asn Arg Cys Phe Tyr Val Glu Leu Ile Arg Gly Arg Lys Gln
 420 425 430
 Glu Thr Glu Val Trp Trp Thr Ser Asn Ser Ile Val Val Phe Cys Gly
 435 440 445
 Thr Ser Gly Thr Tyr Gly Thr Gly Ser Trp Pro Asp Gly Ala Asp Ile
 450 455 460
 Asn Leu Met Pro Ile
 465

<210> SEQ ID NO 41
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 41

Met Lys Thr Ile Ile Ala Leu Ser Tyr Ile Leu Cys Leu Val Phe Ala
 1 5 10 15
 Gln Lys Ile Pro Gly Asn Asp Asn Ser Thr Ala Thr Leu Cys Leu Gly
 20 25 30
 His His Ala Val Pro Asn Gly Thr Leu Val Lys Thr Ile Thr Asn Asp
 35 40 45
 Gln Ile Glu Val Thr Asn Ala Thr Glu Leu Val Gln Ser Ser Ser Thr
 50 55 60
 Gly Arg Ile Cys Asp Ser Pro His Arg Ile Leu Asp Gly Glu Asn Cys
 65 70 75 80
 Thr Leu Ile Asp Ala Leu Leu Gly Asp Pro His Cys Asp Gly Phe Gln
 85 90 95
 Asn Lys Glu Trp Asp Leu Phe Val Glu Arg Ser Lys Ala Tyr Ser Asn
 100 105 110
 Cys Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser Leu Arg Ser Leu Val
 115 120 125
 Ala Ser Ser Gly Thr Leu Glu Phe Asn Asn Glu Ser Phe Asn Trp Thr
 130 135 140
 Gly Val Ala Gln Asn Gly Thr Ser Tyr Ala Cys Lys Arg Ser Ser Ile
 145 150 155 160
 Lys Ser Phe Phe Ser Arg Leu Asn Trp Leu His Gln Leu Lys Tyr Lys
 165 170 175
 Tyr Pro Ala Leu Asn Val Thr Met Pro Asn Asn Asp Lys Phe Asp Lys
 180 185 190
 Leu Tyr Ile Trp Gly Val His His Pro Ser Thr Asp Ser Asp Gln Thr
 195 200 205
 Ser Ile Tyr Ala Gln Ala Ser Gly Arg Val Thr Val Ser Thr Lys Arg
 210 215 220
 Ser Gln Gln Thr Val Ile Pro Asn Ile Gly Ser Arg Pro Trp Val Arg
 225 230 235 240
 Gly Ile Ser Ser Arg Ile Ser Ile His Trp Thr Ile Val Lys Pro Gly
 245 250 255
 Asp Ile Leu Leu Ile Asn Ser Thr Gly Asn Leu Ile Ala Pro Arg Gly
 260 265 270

-continued

```

Tyr Phe Lys Ile Arg Ser Gly Lys Ser Ser Ile Met Arg Ser Asp Ala
275                280                285

Pro Ile Gly Lys Cys Asn Ser Glu Cys Ile Thr Pro Asn Gly Ser Ile
290                295                300

Pro Asn Asp Lys Pro Phe Gln Asn Val Asn Arg Ile Thr Tyr Gly Ala
305                310                315                320

Cys Pro Arg Tyr Val Lys Gln Asn Thr Leu Lys Leu Ala Thr Gly Met
325                330                335

Arg Asn Val Pro Glu Lys Gln Thr Arg Gly Ile Phe Gly Ala Ile Ala
340                345                350

Gly Phe Ile Glu Asn Gly Trp Glu Gly Met Val Asp Gly Trp Tyr Gly
355                360                365

Phe Arg His Gln Asn Ser Glu Gly Thr Gly Gln Ala Ala Asp Leu Lys
370                375                380

Ser Thr Gln Ala Ala Ile Asn Gln Ile Asn Gly Lys Leu Asn Arg Leu
385                390                395                400

Ile Glu Lys Thr Asn Glu Lys Phe His Gln Ile Glu Lys Glu Phe Ser
405                410                415

Glu Val Glu Gly Arg Ile Gln Asp Leu Glu Lys Tyr Val Glu Asp Thr
420                425                430

Lys Ile Asp Leu Trp Ser Tyr Asn Ala Glu Leu Leu Val Ala Leu Glu
435                440                445

Asn Gln His Thr Ile Asp Leu Thr Asp Ser Glu Met Asn Lys Leu Phe
450                455                460

Glu Arg Thr Arg Lys Gln Leu Arg Glu Asn Ala Glu Asp Met Gly Asn
465                470                475                480

Gly Cys Phe Lys Ile Tyr His Lys Cys Asp Asn Ala Cys Ile Gly Ser
485                490                495

Ile Arg Asn Gly Thr Tyr Asp His Asp Val Tyr Arg Asp Glu Ala Leu
500                505                510

Asn Asn Arg Phe Gln Ile Lys Gly Val Glu Leu Lys Ser Gly Tyr Lys
515                520                525

Asp Trp Ile Leu Trp Ile Ser Phe Ala Ile Ser Cys Phe Leu Leu Cys
530                535                540

Val Val Leu Leu Gly Phe Ile Met Trp Ala Cys Gln Lys Gly Asn Ile
545                550                555                560

Arg Cys Asn Ile Cys Ile
565

```

<210> SEQ ID NO 42

<211> LENGTH: 469

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 42

```

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Val Ser Leu Thr
1          5          10          15

Ile Ala Thr Ile Cys Phe Leu Met Gln Ile Ala Ile Leu Val Thr Thr
20          25          30

Val Thr Leu His Phe Lys Gln Tyr Glu Cys Ser Ser Pro Pro Asn Asn
35          40          45

Gln Val Met Leu Cys Glu Pro Thr Ile Ile Glu Arg Asn Ile Thr Glu

```

-continued

50	55	60
Ile Val Tyr Leu Thr	Asn Thr Thr Ile Glu	Lys Glu Ile Cys Pro Lys
65	70	75 80
Leu Ala Glu Tyr Arg	Asn Trp Ser Lys Pro	Gln Cys Lys Ile Thr Gly
85	90	95
Phe Ala Pro Phe Ser	Lys Asp Asn Ser Ile	Arg Leu Ser Ala Gly Gly
100	105	110
Asp Ile Trp Val Thr	Arg Glu Pro Tyr Val	Ser Cys Asp Pro Asp Lys
115	120	125
Cys Tyr Gln Phe Ala	Leu Gly Gln Gly Thr	Thr Leu Asn Asn Arg His
130	135	140
Ser Asn Asp Thr Val	His Asp Arg Thr Pro	Tyr Arg Thr Leu Leu Met
145	150	155 160
Asn Glu Leu Gly Val	Pro Phe His Leu Gly	Thr Lys Gln Val Cys Ile
165	170	175
Ala Trp Ser Ser Ser	Ser Cys His Asp Gly	Lys Ala Trp Leu His Val
180	185	190
Cys Val Thr Gly His	Asp Glu Asn Ala Thr	Ala Ser Phe Ile Tyr Asp
195	200	205
Gly Arg Leu Val Asp	Ser Ile Gly Ser Trp	Ser Lys Lys Ile Leu Arg
210	215	220
Thr Gln Glu Ser Glu	Cys Val Cys Ile Asn	Gly Thr Cys Thr Val Val
225	230	235 240
Met Thr Asp Gly Ser	Ala Ser Gly Arg Ala	Asp Thr Lys Ile Leu Phe
245	250	255
Ile Glu Glu Gly Lys	Ile Val His Ile Ser	Pro Leu Ser Gly Ser Ala
260	265	270
Gln His Val Glu Glu	Cys Ser Cys Tyr Pro	Arg Tyr Pro Gly Val Arg
275	280	285
Cys Val Cys Arg Asp	Asn Trp Lys Gly Ser	Asn Arg Pro Ile Val Asp
290	295	300
Ile Asn Val Lys Asp	Tyr Ser Ile Val Ser	Ser Tyr Val Cys Ser Gly
305	310	315 320
Leu Val Gly Asp Thr	Pro Arg Lys Asn Asp	Ser Ser Ser Ser Ser His
325	330	335
Cys Leu Asn Pro Asn	Asn Glu Glu Gly Gly	His Gly Val Lys Gly Trp
340	345	350
Ala Phe Asp Asp Gly	Asn Asp Val Trp Met	Gly Arg Thr Ile Ser Glu
355	360	365
Lys Phe Arg Ser Gly	Tyr Glu Thr Phe Lys	Val Ile Glu Gly Trp Ser
370	375	380
Lys Pro Asn Ser Lys	Leu Gln Ile Asn Arg	Gln Val Ile Val Asp Arg
385	390	395 400
Gly Asn Arg Ser Gly	Tyr Ser Gly Ile Phe	Ser Val Glu Gly Lys Ser
405	410	415
Cys Ile Asn Arg Cys	Phe Tyr Val Glu Leu	Ile Arg Gly Arg Lys Gln
420	425	430
Glu Thr Glu Val Trp	Trp Thr Ser Asn Ser	Ile Val Val Phe Cys Gly
435	440	445
Thr Ser Gly Thr Tyr	Gly Thr Gly Ser Trp	Pro Asp Gly Ala Asp Ile
450	455	460

-continued

Asn Leu Met Pro Ile
465

<210> SEQ ID NO 43
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 43

Met Lys Thr Ile Ile Ala Leu Ser Tyr Ile Leu Cys Leu Val Phe Ala
1 5 10 15
Gln Lys Leu Pro Gly Asn Asp Asn Ser Thr Ala Thr Leu Cys Leu Gly
20 25 30
His His Ala Val Ser Asn Gly Thr Leu Val Lys Thr Ile Thr Asn Asp
35 40 45
Gln Ile Glu Val Thr Asn Ala Thr Glu Leu Val Gln Ser Ser Ser Thr
50 55 60
Gly Arg Ile Cys Asp Ser Pro His Gln Ile Leu Asp Gly Glu Asn Cys
65 70 75 80
Thr Leu Ile Asp Ala Leu Leu Gly Asp Pro His Cys Asp Gly Phe Gln
85 90 95
Asn Lys Glu Trp Asp Leu Phe Val Glu Arg Ser Lys Ala Tyr Ser Asn
100 105 110
Cys Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser Leu Arg Ser Leu Val
115 120 125
Ala Ser Ser Gly Thr Leu Glu Phe Asn Asn Glu Ser Phe Asn Trp Thr
130 135 140
Gly Val Ala Gln Asn Gly Thr Ser Ser Ala Cys Lys Arg Gly Ser Asn
145 150 155 160
Lys Ser Phe Phe Ser Arg Leu Asn Trp Leu His Gln Leu Lys Tyr Lys
165 170 175
Tyr Pro Ala Leu Asn Val Thr Met Pro Asn Asn Glu Lys Phe Asp Lys
180 185 190
Leu Tyr Ile Trp Gly Val Leu His Pro Ser Thr Asp Ser Asp Gln Ile
195 200 205
Ser Leu Tyr Ala Gln Ala Ser Gly Arg Val Thr Val Ser Thr Lys Arg
210 215 220
Ser Gln Gln Thr Val Ile Pro Asn Ile Gly Ser Arg Pro Trp Val Arg
225 230 235 240
Gly Val Ser Ser Arg Ile Ser Ile Tyr Trp Thr Ile Val Lys Pro Gly
245 250 255
Asp Ile Leu Leu Ile Asn Ser Thr Gly Asn Leu Ile Ala Pro Arg Gly
260 265 270
Tyr Phe Lys Ile Arg Ser Gly Lys Ser Ser Ile Met Arg Ser Asp Ala
275 280 285
Pro Ile Gly Lys Cys Asn Ser Glu Cys Ile Thr Pro Asn Gly Ser Ile
290 295 300
Pro Asn Asp Lys Pro Phe Gln Asn Val Asn Arg Ile Thr Tyr Gly Ala
305 310 315 320
Cys Pro Arg Tyr Val Lys Gln Asn Thr Leu Lys Leu Ala Thr Gly Met
325 330 335
Arg Asn Val Pro Glu Lys Gln Thr Arg Gly Ile Phe Gly Ala Ile Ala

-continued

340	345	350
Gly Phe Ile Glu Asn	Gly Trp Glu Gly Met	Val Asp Gly Trp Tyr Gly
355	360	365
Phe Arg His Gln Asn	Ser Glu Gly Thr Gly	Gln Ala Ala Asp Leu Lys
370	375	380
Ser Thr Gln Ala Ala	Ile Asn Gln Ile Asn	Gly Lys Leu Asn Arg Leu
385	390	395
Ile Glu Lys Thr Asn	Glu Lys Phe His Gln	Ile Glu Lys Glu Phe Ser
405	410	415
Glu Val Glu Gly Arg	Ile Gln Asp Leu Glu	Lys Tyr Val Glu Asp Thr
420	425	430
Lys Ile Asp Leu Trp	Ser Tyr Asn Ala Glu	Leu Leu Val Ala Leu Glu
435	440	445
Asn Gln His Thr Ile	Asp Leu Thr Asp Ser	Glu Met Asn Lys Leu Phe
450	455	460
Glu Arg Thr Lys Lys	Gln Leu Arg Glu Asn	Ala Glu Asp Met Gly Asn
465	470	475
Gly Cys Phe Lys Ile	Tyr His Lys Cys Asp	Asn Ala Cys Ile Gly Ser
485	490	495
Ile Arg Asn Gly Thr	Tyr Asp His Asp Val	Tyr Arg Asp Glu Ala Leu
500	505	510
Asn Asn Arg Phe Gln	Ile Lys Gly Val Glu	Leu Lys Ser Gly Tyr Lys
515	520	525
Asp Trp Ile Leu Trp	Ile Ser Phe Ala Ile	Ser Cys Phe Leu Leu Cys
530	535	540
Val Val Leu Leu Gly	Phe Ile Met Trp Ala	Cys Gln Lys Gly Asn Ile
545	550	555
Arg Cys Asn Ile Cys Ile		
565		

<210> SEQ ID NO 44

<211> LENGTH: 469

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 44

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Val Ser Leu Thr
1 5 10 15
Ile Ala Thr Ile Cys Phe Leu Met Gln Ile Ala Ile Leu Val Thr Thr
20 25 30
Val Thr Leu His Phe Lys Gln Tyr Glu Cys Asn Ser Pro Pro Asn Asn
35 40 45
Gln Val Met Leu Cys Glu Pro Thr Ile Ile Glu Arg Asn Ile Thr Glu
50 55 60
Ile Val Tyr Leu Thr Asn Thr Thr Ile Glu Lys Glu Ile Cys Pro Lys
65 70 75 80
Leu Ala Glu Tyr Arg Asn Trp Ser Lys Pro Gln Cys Lys Ile Thr Gly
85 90 95
Phe Ala Pro Phe Ser Lys Asp Asn Ser Ile Arg Leu Ser Ala Gly Gly
100 105 110
Asp Ile Trp Val Thr Arg Glu Pro Tyr Val Ser Cys Asp Pro Asp Lys
115 120 125

-continued

Cys	Tyr	Gln	Phe	Ala	Leu	Gly	Gln	Gly	Thr	Thr	Leu	Asn	Asn	Arg	His
130					135					140					
Ser	Asn	Asp	Thr	Val	His	Asp	Arg	Thr	Pro	Tyr	Arg	Thr	Leu	Leu	Met
145					150					155					160
Asn	Glu	Leu	Gly	Val	Pro	Phe	His	Leu	Gly	Thr	Lys	Gln	Val	Cys	Ile
165					170					175					
Ala	Trp	Ser	Ser	Ser	Ser	Cys	His	Asp	Gly	Lys	Ala	Trp	Leu	His	Val
180					185					190					
Cys	Val	Thr	Gly	His	Asp	Glu	Asn	Ala	Thr	Ala	Ser	Phe	Ile	Tyr	Asp
195					200					205					
Gly	Arg	Leu	Val	Asp	Ser	Ile	Gly	Ser	Trp	Ser	Lys	Lys	Ile	Leu	Arg
210					215					220					
Thr	Gln	Glu	Ser	Glu	Cys	Val	Cys	Ile	Asn	Gly	Thr	Cys	Thr	Val	Val
225					230					235					240
Met	Thr	Asp	Gly	Ser	Ala	Ser	Gly	Arg	Ala	Asp	Thr	Lys	Ile	Leu	Phe
245					250					255					
Ile	Glu	Glu	Gly	Lys	Ile	Val	His	Thr	Ser	Lys	Leu	Ser	Gly	Ser	Ala
260					265					270					
Gln	His	Val	Glu	Glu	Cys	Ser	Cys	Tyr	Pro	Arg	Tyr	Pro	Gly	Val	Arg
275					280					285					
Cys	Val	Cys	Arg	Asp	Asn	Trp	Lys	Gly	Ser	Asn	Arg	Pro	Ile	Val	Asp
290					295					300					
Ile	Asn	Val	Lys	Asp	Tyr	Ser	Ile	Val	Ser	Ser	Tyr	Val	Cys	Ser	Gly
305					310					315					320
Leu	Val	Gly	Asp	Thr	Pro	Arg	Lys	Asn	Asp	Ser	Ser	Ser	Ser	Ser	His
325					330					335					
Cys	Leu	Asp	Pro	Asn	Asn	Glu	Glu	Gly	Gly	His	Gly	Val	Lys	Gly	Trp
340					345					350					
Ala	Phe	Asp	Asp	Gly	Asn	Asp	Val	Trp	Met	Gly	Arg	Thr	Ile	Ser	Glu
355					360					365					
Lys	Ser	Arg	Ser	Gly	Tyr	Glu	Thr	Phe	Lys	Val	Ile	Glu	Gly	Trp	Ser
370					375					380					
Lys	Pro	Asn	Ser	Lys	Leu	Gln	Ile	Asn	Arg	Gln	Val	Ile	Val	Glu	Arg
385					390					395					400
Gly	Asn	Met	Ser	Gly	Tyr	Ser	Gly	Ile	Phe	Ser	Val	Glu	Gly	Lys	Ser
405					410					415					
Cys	Ile	Asn	Arg	Cys	Phe	Tyr	Val	Glu	Leu	Ile	Arg	Gly	Arg	Lys	Gln
420					425					430					
Glu	Thr	Glu	Val	Trp	Trp	Thr	Ser	Asn	Ser	Ile	Val	Val	Phe	Cys	Gly
435					440					445					
Thr	Ser	Gly	Thr	Tyr	Gly	Thr	Gly	Ser	Trp	Pro	Asp	Gly	Ala	Asp	Ile
450					455					460					
Asn	Leu	Met	Pro	Ile											
465															

<210> SEQ ID NO 45

<211> LENGTH: 566

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 45

Met	Lys	Thr	Ile	Ile	Ala	Leu	Ser	Tyr	Ile	Leu	Cys	Leu	Val	Phe	Ser
1				5					10					15	

-continued

Gln Lys Leu Pro Gly	Asn Asp Asn Ser Thr	Ala Thr Leu Cys Leu Gly
20	25	30
His His Ala Val Pro	Asn Gly Thr Ile Val	Lys Thr Ile Thr Asn Asp
35	40	45
Gln Ile Glu Val Thr	Asn Ala Thr Glu Leu	Val Gln Ser Ser Ser Thr
50	55	60
Gly Gly Ile Cys Asp	Ser Pro His Gln Ile	Leu Asp Gly Glu Asn Cys
65	70	75 80
Thr Leu Ile Asp Ala	Leu Leu Gly Asp Pro	Gln Cys Asp Gly Phe Gln
85	90	95
Asn Lys Lys Trp Asp	Leu Phe Val Glu Arg	Ser Lys Ala Tyr Ser Asn
100	105	110
Cys Tyr Pro Tyr Asp	Val Pro Asp Tyr Ala	Ser Leu Arg Ser Leu Val
115	120	125
Ala Ser Ser Gly Thr	Leu Glu Phe Asn Asn	Glu Ser Phe Asn Trp Ala
130	135	140
Gly Val Thr Gln Asn	Gly Thr Ser Ser Ala	Cys Lys Arg Arg Ser Asn
145	150	155 160
Lys Ser Phe Phe Ser	Arg Leu Asn Trp Leu	Thr His Leu Lys Tyr Lys
165	170	175
Tyr Pro Ala Leu Asn	Val Thr Met Pro Asn	Asn Glu Lys Phe Asp Lys
180	185	190
Leu Tyr Ile Trp Gly	Val His His Pro Val	Thr Asp Ser Asp Gln Ile
195	200	205
Ser Leu Tyr Ala Gln	Ala Ser Gly Arg Ile	Thr Val Ser Thr Lys Arg
210	215	220
Ser Gln Gln Thr Val	Ile Pro Asn Ile Gly	Tyr Arg Pro Arg Val Arg
225	230	235 240
Asp Ile Ser Ser Arg	Ile Ser Ile Tyr Trp	Thr Ile Val Lys Pro Gly
245	250	255
Asp Ile Leu Leu Ile	Asn Ser Thr Gly Asn	Leu Ile Ala Pro Arg Gly
260	265	270
Tyr Phe Lys Ile Arg	Ser Gly Lys Ser Ser	Ile Met Arg Ser Asp Ala
275	280	285
Pro Ile Gly Lys Cys	Asn Ser Glu Cys Ile	Thr Pro Asn Gly Ser Ile
290	295	300
Pro Asn Asp Lys Pro	Phe Gln Asn Val Asn	Arg Ile Thr Tyr Gly Ala
305	310	315 320
Cys Pro Arg Tyr Val	Lys Gln Asn Thr Leu	Lys Leu Ala Thr Gly Met
325	330	335
Arg Asn Val Pro Glu	Lys Gln Thr Arg Gly	Ile Phe Gly Ala Ile Ala
340	345	350
Gly Phe Ile Glu Asn	Gly Trp Glu Gly Met	Val Asp Gly Trp Tyr Gly
355	360	365
Phe Arg His Gln Asn	Ser Glu Gly Thr Gly	Gln Ala Ala Asp Leu Lys
370	375	380
Ser Thr Gln Ala Ala	Ile Asn Gln Ile Asn	Gly Lys Leu Asn Arg Leu
385	390	395 400
Ile Gly Lys Thr Asn	Glu Lys Phe His Gln	Ile Glu Lys Glu Phe Ser
405	410	415

-continued

Glu Val Glu Gly Arg Ile Gln Asp Leu Glu Lys Tyr Val Glu Asp Thr
 420 425 430
 Lys Ile Asp Leu Trp Ser Tyr Asn Ala Glu Leu Leu Val Ala Leu Glu
 435 440 445
 Asn Gln His Thr Ile Asp Leu Thr Asp Ser Glu Met Asn Lys Leu Phe
 450 455 460
 Glu Arg Thr Lys Lys Gln Leu Arg Glu Asn Ala Glu Asp Met Gly Asn
 465 470 475 480
 Gly Cys Phe Lys Ile Tyr His Lys Cys Asp Asn Ala Cys Ile Glu Ser
 485 490 495
 Ile Arg Asn Gly Thr Tyr Asp His Asp Val Tyr Arg Asp Glu Ala Leu
 500 505 510
 Asn Asn Arg Phe Gln Ile Lys Gly Val Glu Leu Lys Ser Gly Tyr Lys
 515 520 525
 Asp Trp Ile Leu Trp Ile Ser Phe Ala Ile Ser Cys Phe Leu Leu Cys
 530 535 540
 Val Ala Leu Leu Gly Phe Ile Met Trp Ala Cys Gln Lys Gly Asn Ile
 545 550 555 560
 Arg Cys Asn Ile Cys Ile
 565

<210> SEQ ID NO 46

<211> LENGTH: 469

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 46

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Val Ser Leu Thr
 1 5 10 15
 Ile Ser Thr Ile Cys Phe Phe Met Gln Ile Ala Ile Leu Ile Thr Thr
 20 25 30
 Val Thr Leu His Phe Lys Gln Tyr Glu Phe Asn Ser Pro Pro Asn Asn
 35 40 45
 Gln Val Met Leu Cys Glu Pro Thr Ile Ile Glu Arg Asn Ile Thr Glu
 50 55 60
 Ile Val Tyr Leu Thr Asn Thr Thr Ile Glu Lys Glu Ile Cys Pro Lys
 65 70 75 80
 Leu Ala Glu Tyr Arg Asn Trp Ser Lys Pro Gln Cys Asn Ile Thr Gly
 85 90 95
 Phe Ala Pro Phe Ser Lys Asp Asn Ser Ile Arg Leu Ser Ala Gly Gly
 100 105 110
 Asp Ile Trp Val Thr Arg Glu Pro Tyr Val Ser Cys Asp Pro Asp Lys
 115 120 125
 Cys Tyr Gln Phe Ala Leu Gly Gln Gly Thr Thr Leu Asn Asn Val His
 130 135 140
 Ser Asn Asp Thr Val His Asp Arg Thr Pro Tyr Arg Thr Leu Leu Met
 145 150 155 160
 Asn Glu Leu Gly Val Pro Phe His Leu Gly Thr Lys Gln Val Cys Ile
 165 170 175
 Ala Trp Ser Ser Ser Ser Cys His Asp Gly Lys Ala Trp Leu His Val
 180 185 190
 Cys Val Thr Gly Asp Asp Glu Asn Ala Thr Ala Ser Phe Ile Tyr Asn
 195 200 205

-continued

Gly Arg Leu Val Asp Ser Ile Val Ser Trp Ser Lys Lys Ile Leu Arg
 210 215 220
 Thr Gln Glu Ser Glu Cys Val Cys Ile Asn Gly Thr Cys Thr Val Val
 225 230 235 240
 Met Thr Asp Gly Ser Ala Ser Gly Lys Ala Asp Thr Lys Ile Leu Phe
 245 250 255
 Ile Glu Glu Gly Lys Ile Val His Thr Ser Thr Leu Ser Gly Ser Ala
 260 265 270
 Gln His Val Glu Glu Cys Ser Cys Tyr Pro Arg Tyr Pro Gly Val Arg
 275 280 285
 Cys Val Cys Arg Asp Asn Trp Lys Gly Ser Asn Arg Pro Ile Val Asp
 290 295 300
 Ile Asn Ile Lys Asp Tyr Ser Ile Val Ser Ser Tyr Val Cys Ser Gly
 305 310 315 320
 Leu Val Gly Asp Thr Pro Arg Lys Asn Asp Ser Ser Ser Ser Ser His
 325 330 335
 Cys Leu Asp Pro Asn Asn Glu Glu Gly Gly His Gly Val Lys Gly Trp
 340 345 350
 Ala Phe Asp Asp Gly Asn Asp Val Trp Met Gly Arg Thr Ile Ser Glu
 355 360 365
 Lys Leu Arg Ser Gly Tyr Glu Thr Phe Lys Val Ile Glu Gly Trp Ser
 370 375 380
 Asn Pro Asn Ser Lys Leu Gln Ile Asn Arg Gln Val Ile Val Asp Arg
 385 390 395 400
 Gly Asn Arg Ser Gly Tyr Ser Gly Ile Phe Ser Val Glu Gly Lys Ser
 405 410 415
 Cys Ile Asn Arg Cys Phe Tyr Val Glu Leu Ile Arg Gly Arg Lys Gln
 420 425 430
 Glu Thr Glu Val Leu Trp Thr Ser Asn Ser Ile Val Val Phe Cys Gly
 435 440 445
 Thr Ser Gly Thr Tyr Gly Thr Gly Ser Trp Pro Asp Gly Ala Asp Ile
 450 455 460
 Asn Leu Met Pro Ile
 465

<210> SEQ ID NO 47

<211> LENGTH: 566

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 47

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Ala Phe Thr Ala Thr Tyr
 1 5 10 15
 Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20 25 30
 Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35 40 45
 Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
 50 55 60
 Ala Pro Leu Gln Leu Gly Asn Cys Ser Val Ala Gly Trp Ile Leu Gly
 65 70 75 80
 Asn Pro Lys Cys Glu Ser Leu Phe Ser Lys Glu Ser Trp Ser Tyr Ile

-continued

85	90	95
Ala Glu Thr Pro Asn	Pro Glu Asn Gly Thr	Cys Tyr Pro Gly Tyr Phe
100	105	110
Ala Asp Tyr Glu Glu	Leu Arg Glu Gln Leu	Ser Ser Val Ser Ser Phe
115	120	125
Glu Arg Phe Glu Ile	Phe Pro Lys Glu Ser	Ser Trp Pro Asn His Thr
130	135	140
Val Thr Lys Gly Val	Thr Thr Ser Cys Ser	His Asn Gly Lys Ser Ser
145	150	155 160
Phe Tyr Arg Asn Leu	Leu Trp Leu Thr Lys	Lys Asn Gly Leu Tyr Pro
165	170	175
Asn Val Ser Lys Ser	Tyr Val Asn Asn Lys	Glu Lys Glu Val Leu Val
180	185	190
Leu Trp Gly Val His	His Pro Ser Asn Ile	Gly Asp Gln Arg Ala Ile
195	200	205
Tyr His Thr Glu Asn	Ala Tyr Val Ser Val	Val Ser Ser His Tyr Ser
210	215	220
Arg Arg Phe Thr Pro	Glu Ile Ala Lys Arg	Pro Lys Val Arg Asp Gln
225	230	235 240
Glu Gly Arg Ile Asn	Tyr Tyr Trp Thr Leu	Leu Glu Pro Gly Asp Thr
245	250	255
Ile Ile Phe Glu Ala	Asn Gly Asn Leu Ile	Ala Pro Trp Tyr Ala Phe
260	265	270
Ala Leu Ser Arg Gly	Phe Gly Ser Gly Ile	Ile Thr Ser Asn Ala Ser
275	280	285
Met Asp Glu Cys Asp	Ala Lys Cys Gln Thr	Pro Gln Gly Ala Ile Asn
290	295	300
Ser Ser Leu Pro Phe	Gln Asn Val His Pro	Val Thr Ile Gly Glu Cys
305	310	315 320
Pro Lys Tyr Val Arg	Ser Thr Lys Leu Arg	Met Val Thr Gly Leu Arg
325	330	335
Asn Ile Pro Ser Ile	Gln Ser Arg Gly Leu	Phe Gly Ala Ile Ala Gly
340	345	350
Phe Ile Glu Gly Gly	Trp Thr Gly Met Ile	Asp Gly Trp Tyr Gly Tyr
355	360	365
His His Gln Asn Glu	Gln Gly Ser Gly Tyr	Ala Ala Asp Gln Lys Ser
370	375	380
Thr Gln Asn Ala Ile	Asn Gly Ile Thr Asn	Lys Val Asn Ser Val Ile
385	390	395 400
Glu Lys Met Asn Thr	Gln Phe Thr Ala Val	Gly Lys Glu Phe Asn Lys
405	410	415
Leu Glu Arg Arg Met	Glu Asn Leu Asn Lys	Lys Val Asp Asp Gly Phe
420	425	430
Leu Asp Ile Trp Thr	Tyr Asn Ala Glu Leu	Leu Val Leu Leu Glu Asn
435	440	445
Gly Arg Thr Leu Asp	Phe His Asp Ser Asn	Val Lys Asn Leu Tyr Glu
450	455	460
Lys Val Lys Ser Gln	Leu Lys Asn Asn Ala	Lys Glu Ile Gly Asn Gly
465	470	475 480
Cys Phe Glu Phe Tyr	His Lys Cys Asn Asn	Glu Cys Met Glu Ser Val
485	490	495

-continued

Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu
 500 505 510
 Asn Arg Gly Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr
 515 520 525
 Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu
 530 535 540
 Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu
 545 550 555 560
 Gln Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 48
 <211> LENGTH: 470
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 48

Met Asn Pro Asn Gln Lys Ile Ile Ile Ile Gly Ser Ile Ser Met Ala
 1 5 10 15
 Ile Gly Ile Ile Ser Leu Ile Leu Gln Ile Gly Asn Ile Ile Ser Ile
 20 25 30
 Trp Ala Ser His Ser Ile Gln Thr Gly Ser Gln Asn His Thr Gly Ile
 35 40 45
 Cys Asn Gln Arg Ile Ile Thr Tyr Glu Asn Ser Thr Trp Val Asn Gln
 50 55 60
 Thr Tyr Val Asn Ile Asn Asn Thr Asn Val Val Ala Gly Lys Asp Lys
 65 70 75 80
 Thr Ser Val Thr Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Arg Gly
 85 90 95
 Trp Ala Ile Tyr Thr Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly
 100 105 110
 Asp Val Phe Val Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu
 115 120 125
 Cys Arg Thr Phe Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His
 130 135 140
 Ser Asn Gly Thr Val Lys Asp Arg Ser Pro Tyr Arg Ala Leu Met Ser
 145 150 155 160
 Cys Pro Leu Gly Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser
 165 170 175
 Val Ala Trp Ser Ala Ser Ala Cys His Asp Gly Met Gly Trp Leu Thr
 180 185 190
 Ile Gly Ile Ser Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr
 195 200 205
 Asn Gly Ile Ile Thr Glu Thr Ile Lys Ser Trp Lys Lys Arg Ile Leu
 210 215 220
 Arg Thr Gln Glu Ser Glu Cys Val Cys Val Asn Gly Ser Cys Phe Thr
 225 230 235 240
 Ile Met Thr Asp Gly Pro Ser Asn Gly Ala Ala Ser Tyr Arg Ile Phe
 245 250 255
 Lys Ile Glu Lys Gly Lys Val Thr Lys Ser Ile Glu Leu Asp Ala Pro
 260 265 270
 Asn Tyr His Tyr Glu Glu Cys Ser Cys Tyr Pro Asp Thr Gly Thr Val

-continued

275	280	285
Met Cys Val Cys Arg	Asp Asn Trp His Gly	Ser Asn Arg Pro Trp Val
290	295	300
Ser Phe Asn Gln Asn	Leu Asp Tyr Gln Ile	Gly Tyr Ile Cys Ser Gly
305	310	315 320
Val Phe Gly Asp Asn	Pro Arg Pro Lys Asp	Gly Glu Gly Ser Cys Asn
325	330	335
Pro Val Thr Val Asp	Gly Ala Asp Gly Val	Lys Gly Phe Ser Tyr Arg
340	345	350
Tyr Gly Asn Gly Val	Trp Ile Gly Arg Thr	Lys Ser Asn Arg Leu Arg
355	360	365
Lys Gly Phe Glu Met	Ile Trp Asp Pro Asn	Gly Trp Thr Asp Thr Asp
370	375	380
Ser Asp Phe Ser Val	Lys Gln Asp Val Val	Ala Met Thr Asp Trp Ser
385	390	395 400
Gly Tyr Ser Gly Ser	Phe Val Gln His Pro	Glu Leu Thr Gly Leu Asp
405	410	415
Cys Met Arg Pro Cys	Phe Trp Val Glu Leu	Ile Arg Gly Arg Pro Arg
420	425	430
Glu Asn Thr Thr Ile	Trp Thr Ser Gly Ser	Ser Ile Ser Phe Cys Gly
435	440	445
Val Asn Ser Asp Thr	Ala Asn Trp Ser Trp	Pro Asp Gly Ala Glu Leu
450	455	460
Pro Phe Thr Ile Asp	Lys	
465	470	

<210> SEQ ID NO 49

<211> LENGTH: 562

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 49

Met Lys Ala Lys	Leu Leu Val Leu Leu Cys	Ala Phe Thr Ala Thr Tyr
1	5	10 15
Ala Asp Thr Ile Cys	Ile Gly Tyr His Ala	Asn Asn Ser Thr Asp Thr
20	25	30
Val Asp Thr Val Leu	Glu Lys Asn Val Thr	Val Thr His Ser Val Asn
35	40	45
Leu Leu Glu Asp Ser	His Asn Gly Lys Leu	Cys Arg Leu Lys Gly Thr
50	55	60
Ala Pro Leu Gln Leu	Gly Asn Cys Ser Val	Ala Gly Trp Ile Leu Gly
65	70	75 80
Asn Pro Glu Cys Glu	Ser Leu Phe Ser Lys	Glu Ser Trp Ser Tyr Ile
85	90	95
Ala Glu Thr Pro Asn	Pro Glu Asn Gly Thr	Cys Tyr Pro Gly Tyr Phe
100	105	110
Ala Asp Tyr Glu Glu	Leu Arg Glu Gln Leu	Ser Ser Val Ser Ser Phe
115	120	125
Glu Arg Phe Glu Ile	Phe Pro Lys Glu Ser	Ser Trp Pro Lys His Thr
130	135	140
Val Thr Lys Gly Val	Thr Ala Ser Cys Ser	His Asn Gly Lys Ser Ser
145	150	155 160

-continued

Phe	Tyr	Lys	Asn	Leu	Leu	Trp	Leu	Thr	Glu	Lys	Asn	Gly	Leu	Tyr	Pro
165					170					175					
Asn	Leu	Ser	Lys	Ser	Tyr	Val	Asn	Asn	Lys	Glu	Lys	Glu	Val	Leu	Val
180					185					190					
Leu	Trp	Gly	Val	His	His	Pro	Ser	Asn	Ile	Gly	Asp	Gln	Arg	Ala	Ile
195					200					205					
Tyr	His	Thr	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Ser	Ser	His	Tyr	Ser
210					215					220					
Arg	Arg	Phe	Thr	Pro	Glu	Ile	Ala	Lys	Arg	Pro	Lys	Val	Arg	Gly	Gln
225					230					235					240
Glu	Gly	Arg	Ile	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Glu	Pro	Gly	Asp	Thr
245					250					255					
Ile	Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Phe
260					265					270					
Ala	Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Thr	Ala	Ser
275					280					285					
Met	Gly	Glu	Cys	Asp	Ala	Lys	Cys	Gln	Thr	Pro	Gln	Gly	Ala	Ile	Asn
290					295					300					
Ser	Ser	Leu	Pro	Phe	Gln	Asn	Val	His	Pro	Val	Thr	Ile	Gly	Glu	Cys
305					310					315					320
Pro	Lys	Tyr	Val	Arg	Ser	Thr	Lys	Leu	Arg	Met	Val	Thr	Gly	Leu	Arg
325					330					335					
Asn	Ile	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly
340					345					350					
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr
355					360					365					
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser
370					375					380					
Thr	Gln	Asn	Ala	Ile	Asp	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile
385					390					395					400
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Lys
405					410					415					
Leu	Glu	Arg	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe
420					425					430					
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn
435					440					445					
Gly	Arg	Thr	Leu	Gly	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu
450					455					460					
Lys	Val	Lys	Asn	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly
465					470					475					480
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asn	Asn	Glu	Cys	Met	Glu	Ser	Val
485					490					495					
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu
500					505					510					
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Val	Tyr
515					520					525					
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu
530					535					540					
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu
545					550					555					560

Gln Cys

-continued

```

<210> SEQ ID NO 50
<211> LENGTH: 470
<212> TYPE: PRT
<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 50

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Ile Ser Ile Ala
 1             5             10             15

Ile Gly Ile Ile Ser Leu Ile Leu Gln Ile Gly Asn Ile Ile Ser Ile
20             25             30

Trp Ala Ser His Ser Ile Gln Thr Gly Ser Gln Asn His Thr Gly Ile
35             40             45

Cys Asn Gln Arg Ile Ile Thr Tyr Glu Asn Ser Thr Trp Val Asn Gln
50             55             60

Thr Tyr Val Asn Ile Asn Asn Thr Asn Val Val Ala Gly Lys Asp Lys
65             70             75             80

Thr Ser Met Thr Leu Ala Gly Asn Ser Ser Leu Cys Pro Ile Arg Gly
85             90             95

Trp Ala Ile Tyr Thr Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly
100            105            110

Asp Val Phe Val Ile Arg Glu Pro Phe Ile Ser Cys Ser His Leu Glu
115            120            125

Cys Arg Thr Phe Phe Leu Thr Gln Gly Ala Leu Leu Asn Asp Lys His
130            135            140

Ser Asn Gly Thr Val Lys Asp Arg Ser Pro Tyr Arg Ala Leu Met Ser
145            150            155            160

Cys Pro Leu Gly Glu Ala Pro Ser Pro Tyr Asn Ser Arg Phe Glu Ser
165            170            175

Val Ala Trp Ser Ala Ser Ala Cys His Asp Gly Leu Gly Trp Leu Thr
180            185            190

Ile Gly Ile Ser Gly Pro Asp Asn Gly Ala Val Ala Val Leu Lys Tyr
195            200            205

Asn Gly Ile Ile Thr Glu Thr Ile Lys Ser Trp Lys Lys Arg Ile Leu
210            215            220

Arg Thr Gln Glu Ser Glu Cys Val Cys Met Asn Gly Ser Cys Phe Thr
225            230            235            240

Ile Met Thr Asp Gly Pro Ser Asn Gly Ala Ala Ser Tyr Arg Ile Phe
245            250            255

Lys Ile Glu Lys Gly Arg Val Thr Lys Ser Ile Glu Leu Asp Ala Pro
260            265            270

Asn Tyr His Tyr Glu Glu Cys Ser Cys Tyr Pro Asp Thr Gly Thr Val
275            280            285

Met Cys Val Cys Arg Asp Asn Trp His Gly Ser Asn Arg Pro Trp Val
290            295            300

Ser Phe Asn Gln Asn Leu Asp Tyr Gln Ile Gly Tyr Ile Cys Ser Gly
305            310            315            320

Val Phe Gly Asp Asn Pro Arg Pro Lys Asp Gly Glu Gly Ser Cys Asn
325            330            335

Pro Val Thr Val Asp Gly Ala Asp Gly Val Lys Gly Phe Ser Tyr Arg
340            345            350

Tyr Gly Asn Gly Val Trp Ile Gly Arg Thr Lys Ser Asn Arg Leu Arg

```

-continued

355	360	365
Lys Gly Phe Glu Met	Ile Trp Asp Pro Asn	Gly Trp Thr Asp Thr Asp
370	375	380
Ser Asp Phe Ser Met	Lys Gln Asp Ile Val	Ala Met Thr Asp Trp Ser
385	390	395 400
Gly Tyr Ser Gly Ser	Phe Val Gln His Pro	Glu Leu Thr Gly Leu Asp
405	410	415
Cys Met Arg Pro Cys	Phe Trp Val Glu Leu	Val Arg Gly Leu Pro Arg
420	425	430
Glu Asn Thr Thr Ile	Trp Thr Ser Gly Ser	Ser Ile Ser Phe Cys Gly
435	440	445
Val Asn Ser Asp Thr	Ala Asn Trp Ser Trp	Pro Asp Gly Ala Glu Leu
450	455	460
Pro Phe Thr Ile Asp	Lys	
465	470	

<210> SEQ ID NO 51

<211> LENGTH: 565

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 51

Met Lys Ala Lys	Leu Leu Val Leu	Leu Cys Thr Phe Thr	Ala Thr Tyr
1	5	10	15
Ala Asp Thr Ile Cys	Ile Gly Tyr His	Ala Asn Asn Ser Thr	Asp Thr
20	25	30	
Val Asp Thr Val Leu	Glu Lys Asn Val Thr	Val Thr His Ser Val	Asn
35	40	45	
Leu Leu Glu Asp Ser	His Asn Gly Lys Leu	Cys Leu Leu Lys Gly	Ile
50	55	60	
Ala Pro Leu Gln Leu	Gly Asn Cys Ser Val	Ala Gly Trp Ile Leu	Gly
65	70	75	80
Asn Pro Glu Cys Glu	Ser Leu Ile Ser Lys	Glu Ser Trp Ser Tyr	Ile
85	90	95	
Val Glu Thr Pro Asn	Pro Glu Asn Gly Thr	Cys Tyr Pro Gly Tyr	Phe
100	105	110	
Ala Asp Tyr Glu Glu	Leu Arg Glu Gln Leu	Ser Ser Val Ser Ser	Phe
115	120	125	
Glu Arg Phe Glu Ile	Phe Pro Lys Glu Ser	Ser Trp Pro Lys His	Thr
130	135	140	
Val Thr Gly Val Thr	Ala Ser Cys Ser His	Asn Gly Lys Ser Ser	Phe
145	150	155	160
Tyr Arg Asn Leu Leu	Trp Leu Thr Glu Lys	Asn Gly Leu Tyr Pro	Asn
165	170	175	
Leu Ser Asn Ser Tyr	Val Asn Asn Lys Glu	Lys Glu Val Leu Val	Leu
180	185	190	
Trp Gly Val His His	Pro Ser Asn Ile Gly	Asp Gln Arg Ala Ile	Tyr
195	200	205	
His Thr Glu Asn Ala	Tyr Val Ser Val Val	Ser Ser His Tyr Ser	Arg
210	215	220	
Arg Phe Thr Pro Glu	Ile Ala Lys Arg Pro	Lys Val Arg Gly Gln	Glu
225	230	235	240

-continued

Gly Arg Ile Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr Ile
 245 250 255
 Ile Phe Glu Ala Asn Gly Asn Leu Ile Ala Pro Trp Tyr Ala Phe Ala
 260 265 270
 Leu Ser Arg Gly Phe Gly Ser Gly Ile Ile Thr Ser Asn Ala Pro Met
 275 280 285
 Asn Glu Cys Asp Ala Lys Cys Gln Thr Pro Gln Gly Ala Ile Asn Ser
 290 295 300
 Ser Leu Pro Phe Gln Asn Val His Pro Val Thr Ile Gly Glu Cys Pro
 305 310 315 320
 Lys Tyr Val Arg Ser Thr Lys Leu Arg Met Val Thr Gly Leu Arg Asn
 325 330 335
 Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly Phe
 340 345 350
 Ile Glu Gly Gly Trp Thr Gly Met Met Asp Gly Trp Tyr Gly Tyr His
 355 360 365
 His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser Thr
 370 375 380
 Gln Asn Ala Ile Asn Gly Ile Thr Asn Lys Val Asn Ser Val Ile Glu
 385 390 395 400
 Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn Lys Leu
 405 410 415
 Glu Arg Arg Met Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe Leu
 420 425 430
 Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn Glu
 435 440 445
 Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu Lys
 450 455 460
 Val Lys Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly Cys
 465 470 475 480
 Phe Glu Phe Tyr His Lys Cys Asn Asn Glu Cys Met Glu Ser Val Lys
 485 490 495
 Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu Asn
 500 505 510
 Arg Glu Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr Gln
 515 520 525
 Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu Val
 530 535 540
 Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu Gln
 545 550 555 560
 Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 52

<211> LENGTH: 470

<212> TYPE: PRT

<213> ORGANISM: Influenza A virus

<400> SEQUENCE: 52

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Ile Ser Ile Val
 1 5 10 15

Ile Gly Ile Ile Ser Leu Met Leu Gln Ile Gly Asn Ile Ile Ser Ile
 20 25 30

-continued

Trp	Ala	Ser	His	Ser	Ile	Gln	Thr	Gly	Ser	Gln	Asn	His	Thr	Gly	Ile
35					40					45					
Cys	Asn	Gln	Arg	Ile	Ile	Thr	Tyr	Glu	Asn	Ser	Thr	Trp	Val	Asn	His
50					55					60					
Thr	Tyr	Val	Asn	Ile	Asn	Asn	Thr	Asn	Val	Val	Ala	Gly	Lys	Asp	Lys
65					70					75					80
Thr	Ser	Val	Thr	Leu	Ala	Gly	Asn	Ser	Ser	Leu	Cys	Ser	Ile	Ser	Gly
85					90					95					
Trp	Ala	Ile	Tyr	Thr	Lys	Asp	Asn	Ser	Ile	Arg	Ile	Gly	Ser	Lys	Gly
100					105					110					
Asp	Val	Phe	Val	Ile	Arg	Glu	Pro	Phe	Ile	Ser	Cys	Ser	His	Leu	Glu
115					120					125					
Cys	Arg	Thr	Phe	Phe	Leu	Thr	Gln	Gly	Ala	Leu	Leu	Asn	Asp	Lys	His
130					135					140					
Ser	Asn	Gly	Thr	Val	Lys	Asp	Arg	Ser	Pro	Tyr	Arg	Ala	Leu	Met	Ser
145					150					155					160
Cys	Pro	Leu	Gly	Glu	Ala	Pro	Ser	Pro	Tyr	Asn	Ser	Lys	Phe	Glu	Ser
165					170					175					
Val	Ala	Trp	Ser	Ala	Ser	Ala	Cys	His	Asp	Gly	Met	Gly	Trp	Leu	Thr
180					185					190					
Ile	Gly	Ile	Ser	Gly	Pro	Asp	Asn	Gly	Ala	Val	Ala	Val	Leu	Lys	Tyr
195					200					205					
Asn	Gly	Ile	Ile	Thr	Glu	Thr	Ile	Lys	Ser	Trp	Lys	Lys	Arg	Ile	Leu
210					215					220					
Arg	Thr	Gln	Glu	Ser	Glu	Cys	Val	Cys	Val	Asn	Gly	Ser	Cys	Phe	Thr
225					230					235					240
Ile	Met	Thr	Asp	Gly	Pro	Ser	Asn	Gly	Ala	Ala	Ser	Tyr	Lys	Ile	Phe
245					250					255					
Lys	Ile	Glu	Lys	Gly	Lys	Val	Thr	Lys	Ser	Ile	Glu	Leu	Asn	Ala	Pro
260					265					270					
Asn	Ser	His	Tyr	Glu	Glu	Cys	Ser	Cys	Tyr	Pro	Asp	Thr	Gly	Thr	Val
275					280					285					
Met	Cys	Val	Cys	Arg	Asp	Asn	Trp	His	Gly	Ser	Asn	Arg	Pro	Trp	Val
290					295					300					
Ser	Phe	Asn	Gln	Asn	Leu	Asp	Tyr	Gln	Ile	Gly	Tyr	Ile	Cys	Ser	Gly
305					310					315					320
Val	Phe	Gly	Asp	Asn	Pro	Arg	Pro	Lys	Asp	Gly	Glu	Gly	Ser	Cys	Asn
325					330					335					
Pro	Val	Thr	Val	Asp	Gly	Ala	Asp	Gly	Val	Lys	Gly	Phe	Ser	Tyr	Arg
340					345					350					
Tyr	Gly	Asn	Gly	Val	Trp	Ile	Gly	Arg	Thr	Lys	Ser	Asn	Arg	Leu	Arg
355					360					365					
Lys	Gly	Phe	Glu	Met	Ile	Trp	Asp	Pro	Asn	Gly	Trp	Thr	Asp	Thr	Asp
370					375					380					
Ser	Asp	Phe	Ser	Val	Lys	Gln	Asp	Val	Val	Ala	Met	Thr	Asp	Trp	Ser
385					390					395					400
Gly	Tyr	Ser	Gly	Ser	Phe	Val	Gln	His	Pro	Glu	Leu	Thr	Gly	Leu	Asp
405					410					415					
Cys	Ile	Arg	Pro	Cys	Phe	Trp	Val	Glu	Leu	Val	Arg	Gly	Arg	Pro	Arg
420					425					430					

-continued

Glu	Asn	Thr	Thr	Ile	Trp	Thr	Ser	Gly	Ser	Ser	Ile	Ser	Phe	Cys	Gly
435					440					445					
Val	Asn	Ser	Asp	Thr	Ala	Asn	Trp	Ser	Trp	Pro	Asp	Gly	Ala	Glu	Leu
450					455					460					
Pro	Phe	Thr	Ile	Asp	Lys										
465					470										

<210> SEQ ID NO 53
 <211> LENGTH: 565
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 53

Met	Lys	Ala	Lys	Leu	Leu	Val	Leu	Leu	Cys	Thr	Phe	Thr	Ala	Thr	Tyr
1				5					10					15	
Ala	Asp	Thr	Ile	Cys	Ile	Gly	Tyr	His	Ala	Asn	Asn	Ser	Thr	Asp	Thr
20				25					30						
Val	Asp	Thr	Val	Leu	Glu	Lys	Asn	Val	Thr	Val	Thr	His	Ser	Val	Asn
35				40					45						
Leu	Leu	Glu	Asp	Ser	His	Asn	Gly	Lys	Leu	Cys	Leu	Leu	Lys	Gly	Ile
50				55					60						
Ala	Pro	Leu	Gln	Leu	Gly	Asn	Cys	Ser	Val	Ala	Gly	Trp	Ile	Leu	Gly
65				70					75					80	
Asn	Pro	Glu	Cys	Glu	Leu	Leu	Ile	Ser	Lys	Glu	Ser	Trp	Ser	Tyr	Ile
85				90					95						
Val	Glu	Thr	Pro	Asn	Pro	Glu	Asn	Gly	Thr	Cys	Tyr	Pro	Gly	Tyr	Phe
100				105					110						
Ala	Asp	Tyr	Glu	Glu	Leu	Arg	Glu	Gln	Leu	Ser	Ser	Val	Ser	Ser	Phe
115				120					125						
Glu	Arg	Phe	Glu	Ile	Phe	Pro	Lys	Glu	Ser	Ser	Trp	Pro	Lys	His	Thr
130				135					140						
Val	Thr	Gly	Val	Ser	Ala	Ser	Cys	Ser	His	Asn	Gly	Lys	Asn	Ser	Phe
145				150					155					160	
Tyr	Arg	Asn	Leu	Leu	Trp	Leu	Thr	Gly	Lys	Asn	Gly	Leu	Tyr	Pro	Asn
165				170					175						
Leu	Ser	Lys	Ser	Tyr	Val	Asn	Asn	Lys	Glu	Lys	Glu	Val	Leu	Val	Leu
180				185					190						
Trp	Gly	Val	His	His	Pro	Pro	Asn	Ile	Gly	Asp	Gln	Arg	Ala	Leu	Tyr
195				200					205						
His	Thr	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Ser	Ser	His	Tyr	Ser	Arg
210				215					220						
Arg	Phe	Thr	Pro	Glu	Ile	Ala	Lys	Arg	Pro	Lys	Val	Arg	Asp	Gln	Glu
225				230					235					240	
Gly	Arg	Ile	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Glu	Pro	Gly	Asp	Thr	Ile
245				250					255						
Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Phe	Ala
260				265					270						
Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Asn	Ala	Pro	Met
275				280					285						
Asp	Glu	Cys	Asp	Ala	Lys	Cys	Gln	Thr	Pro	Gln	Gly	Ala	Ile	Asn	Ser
290				295					300						
Ser	Leu	Pro	Phe	Gln	Asn	Val	His	Pro	Val	Thr	Ile	Gly	Glu	Cys	Pro
305				310					315					320	

-continued

Lys Tyr Val Arg Ser Ala Lys Leu Arg Met Val Thr Gly Leu Arg Asn
 325 330 335
 Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly Phe
 340 345 350
 Ile Glu Gly Gly Trp Thr Gly Met Val Asp Gly Trp Tyr Gly Tyr His
 355 360 365
 His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser Thr
 370 375 380
 Gln Asn Ala Ile Asn Gly Ile Thr Asn Lys Val Asn Ser Val Ile Glu
 385 390 395 400
 Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn Lys Leu
 405 410 415
 Glu Arg Arg Met Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe Leu
 420 425 430
 Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn Glu
 435 440 445
 Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu Lys
 450 455 460
 Val Lys Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly Cys
 465 470 475 480
 Phe Glu Phe Tyr His Lys Cys Asn Asn Glu Cys Met Glu Ser Val Lys
 485 490 495
 Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu Asn
 500 505 510
 Arg Glu Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr Gln
 515 520 525
 Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu Val
 530 535 540
 Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu Gln
 545 550 555 560
 Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 54
 <211> LENGTH: 470
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A virus

<400> SEQUENCE: 54

Met Asn Pro Asn Gln Lys Ile Ile Thr Ile Gly Ser Ile Ser Ile Ala
 1 5 10 15
 Ile Gly Ile Ile Ser Leu Met Leu Gln Ile Gly Asn Ile Ile Ser Ile
 20 25 30
 Trp Ala Ser His Ser Ile Gln Thr Gly Ser Gln Asn His Thr Gly Val
 35 40 45
 Cys Asn Gln Arg Ile Ile Thr Tyr Glu Asn Ser Thr Trp Val Asn His
 50 55 60
 Thr Tyr Val Asn Ile Asn Asn Thr Asn Val Val Ala Gly Lys Asp Lys
 65 70 75 80
 Thr Ser Val Thr Leu Ala Gly Asn Ser Ser Leu Cys Ser Ile Ser Gly
 85 90 95
 Trp Ala Ile Tyr Thr Lys Asp Asn Ser Ile Arg Ile Gly Ser Lys Gly

-continued

100	105	110
Asp Val Phe Val Ile	Arg Glu Pro Phe Ile	Ser Cys Ser His Leu Glu
115	120	125
Cys Arg Thr Phe Phe	Leu Thr Gln Gly Ala	Leu Leu Asn Asp Lys His
130	135	140
Ser Asn Gly Thr Val	Lys Asp Arg Ser Pro	Tyr Arg Ala Leu Met Ser
145	150	155 160
Cys Pro Leu Gly Glu	Ala Pro Ser Pro Tyr	Asn Ser Lys Phe Glu Ser
165	170	175
Val Ala Trp Ser Ala	Ser Ala Cys His Asp	Gly Met Gly Trp Leu Thr
180	185	190
Ile Gly Ile Ser Gly	Pro Asp Asn Gly Ala	Val Ala Val Leu Lys Tyr
195	200	205
Asn Gly Ile Ile Thr	Glu Thr Ile Lys Ser	Trp Lys Lys Arg Ile Leu
210	215	220
Arg Thr Gln Glu Ser	Glu Cys Val Cys Val	Asn Gly Ser Cys Phe Thr
225	230	235 240
Ile Met Thr Asp Gly	Pro Ser Asn Gly Ala	Ala Ser Tyr Lys Ile Phe
245	250	255
Lys Ile Glu Lys Gly	Lys Val Thr Lys Ser	Ile Glu Leu Asn Ala Pro
260	265	270
Asn Phe His Tyr Glu	Glu Cys Ser Cys Tyr	Pro Asp Thr Gly Thr Val
275	280	285
Met Cys Val Cys Arg	Asp Asn Trp His Gly	Ser Asn Arg Pro Trp Val
290	295	300
Ser Phe Asn Gln Asn	Leu Asp Tyr Gln Ile	Gly Tyr Ile Cys Ser Gly
305	310	315 320
Val Phe Gly Asp Asn	Pro Arg Pro Lys Asp	Gly Glu Gly Ser Cys Asn
325	330	335
Pro Val Thr Val Asp	Gly Ala Asp Gly Val	Lys Gly Phe Ser Tyr Lys
340	345	350
Tyr Gly Asn Gly Val	Trp Ile Gly Arg Thr	Lys Ser Asn Arg Leu Arg
355	360	365
Lys Gly Phe Glu Met	Ile Trp Asp Pro Asn	Gly Trp Thr Asp Thr Asp
370	375	380
Ser Asp Phe Ser Val	Lys Gln Asp Val Val	Ala Ile Thr Asp Trp Ser
385	390	395 400
Gly Tyr Ser Gly Ser	Phe Val Gln His Pro	Glu Leu Thr Gly Leu Asp
405	410	415
Cys Ile Arg Pro Cys	Phe Trp Val Glu Leu	Val Arg Gly Leu Pro Arg
420	425	430
Glu Asn Thr Thr Ile	Trp Thr Ser Gly Ser	Ser Ile Ser Phe Cys Gly
435	440	445
Val Asn Ser Asp Thr	Ala Asn Trp Ser Trp	Pro Asp Gly Ala Glu Leu
450	455	460
Pro Phe Thr Ile Asp	Lys	
465	470	

<210> SEQ ID NO 55

<211> LENGTH: 583

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

-continued

<400> SEQUENCE: 55

Met	Lys	Ala	Ile	Ile	Val	Leu	Leu	Met	Val	Val	Thr	Ser	Asn	Ala	Asp	1	5	10	15
Arg	Ile	Cys	Thr	Gly	Ile	Thr	Ser	Ser	Asn	Ser	Pro	His	Val	Val	Lys	20	25	30	
Thr	Ala	Thr	Gln	Gly	Glu	Val	Asn	Val	Thr	Gly	Val	Ile	Pro	Leu	Thr	35	40	45	
Thr	Thr	Pro	Thr	Lys	Ser	His	Phe	Ala	Asn	Leu	Lys	Gly	Thr	Lys	Thr	50	55	60	
Arg	Gly	Lys	Leu	Cys	Pro	Asn	Cys	Leu	Asn	Cys	Thr	Asp	Leu	Asp	Val	65	70	75	80
Ala	Leu	Gly	Arg	Pro	Met	Cys	Ile	Gly	Ile	Thr	Pro	Ser	Ala	Lys	Ala	85	90	95	
Ser	Ile	Leu	His	Glu	Val	Arg	Pro	Val	Thr	Ser	Gly	Cys	Phe	Pro	Ile	100	105	110	
Met	His	Asp	Arg	Thr	Lys	Ile	Arg	Gln	Leu	Pro	Asn	Leu	Leu	Arg	Gly	115	120	125	
Tyr	Glu	His	Ile	Arg	Leu	Ser	Thr	His	Asn	Val	Ile	Asn	Ala	Glu	Arg	130	135	140	
Ala	Pro	Gly	Gly	Pro	Tyr	Arg	Leu	Gly	Thr	Ser	Gly	Ser	Cys	Pro	Asn	145	150	155	160
Val	Thr	Ser	Arg	Ser	Gly	Phe	Phe	Ala	Thr	Met	Ala	Trp	Ala	Val	Pro	165	170	175	
Arg	Asp	Asn	Lys	Thr	Ala	Thr	Asn	Pro	Leu	Thr	Val	Glu	Val	Pro	Tyr	180	185	190	
Ile	Cys	Thr	Lys	Gly	Glu	Asp	Gln	Ile	Thr	Val	Trp	Gly	Phe	His	Ser	195	200	205	
Asp	Asn	Lys	Ile	Gln	Met	Lys	Asn	Leu	Tyr	Gly	Asp	Ser	Asn	Pro	Gln	210	215	220	
Lys	Phe	Thr	Ser	Ser	Ala	Asn	Gly	Ile	Thr	Thr	His	Tyr	Val	Ser	Gln	225	230	235	240
Ile	Gly	Gly	Phe	Pro	Asn	Gln	Thr	Glu	Asp	Gly	Gly	Leu	Pro	Gln	Ser	245	250	255	
Gly	Arg	Ile	Val	Val	Asp	Tyr	Met	Val	Gln	Lys	Pro	Gly	Lys	Thr	Gly	260	265	270	
Thr	Ile	Val	Tyr	Gln	Arg	Gly	Val	Leu	Leu	Pro	Gln	Lys	Val	Trp	Cys	275	280	285	
Ala	Ser	Gly	Arg	Ser	Lys	Val	Ile	Lys	Gly	Ser	Leu	Pro	Leu	Ile	Gly	290	295	300	
Glu	Ala	Asp	Cys	Leu	His	Glu	Lys	Tyr	Gly	Gly	Leu	Asn	Lys	Ser	Lys	305	310	315	320
Pro	Tyr	Tyr	Thr	Gly	Glu	His	Ala	Lys	Ala	Ile	Gly	Asn	Cys	Pro	Ile	325	330	335	
Trp	Val	Lys	Thr	Pro	Leu	Lys	Leu	Ala	Asn	Gly	Thr	Lys	Tyr	Arg	Pro	340	345	350	
Pro	Ala	Lys	Leu	Leu	Lys	Glu	Lys	Gly	Phe	Phe	Gly	Ala	Ile	Ala	Gly	355	360	365	
Phe	Leu	Glu	Gly	Gly	Trp	Glu	Gly	Met	Ile	Ala	Gly	Trp	His	Gly	Tyr	370	375	380	
Thr	Ser	His	Gly	Ala	His	Gly	Val	Ala	Val	Ala	Ala	Asp	Leu	Lys	Ser				

-continued

385	390	395	400
Thr Gln Glu Ala Ile	Asn Lys Ile Thr Lys	Asn Leu Asn Ser Leu Ser	
405	410	415	
Glu Leu Glu Val Lys	Asn Leu Gln Arg Leu	Ser Gly Ala Met Asp Glu	
420	425	430	
Leu His Asn Glu Ile	Leu Glu Leu Asp Glu	Lys Val Asp Asp Leu Arg	
435	440	445	
Ala Asp Thr Ile Ser	Ser Gln Ile Glu Leu	Ala Val Leu Leu Ser Asn	
450	455	460	
Glu Gly Ile Ile Asn	Ser Glu Asp Glu His	Leu Leu Ala Leu Glu Arg	
465	470	475	480
Lys Leu Lys Lys Met	Leu Gly Pro Ser Ala	Val Asp Ile Gly Asn Gly	
485	490	495	
Cys Phe Glu Thr Lys	His Lys Cys Asn Gln	Thr Cys Leu Asp Arg Ile	
500	505	510	
Ala Ala Gly Thr Phe	Asn Ala Gly Glu Phe	Ser Leu Pro Thr Phe Asp	
515	520	525	
Ser Leu Asn Ile Thr	Ala Ala Ser Leu Asn	Asp Asp Gly Leu Asp Asn	
530	535	540	
His Thr Ile Leu Leu	Tyr Tyr Ser Thr Ala	Ala Ser Ser Leu Ala Val	
545	550	555	560
Thr Leu Met Ile Ala	Ile Phe Ile Val Tyr	Met Val Ser Arg Asp Asn	
565	570	575	
Val Ser Cys Ser Ile	Cys Leu		
580			

<210> SEQ ID NO 56

<211> LENGTH: 465

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 56

Met Leu Pro Ser Thr	Ile Gln Thr Leu Thr	Leu Phe Leu Thr Ser Gly
1	5	10
Gly Val Leu Leu Ser	Leu Tyr Val Ser Ala	Leu Leu Ser Tyr Leu Leu
20	25	30
Tyr Ser Asp Ile Leu	Leu Lys Phe Ser Pro	Lys Ile Ile Ala Pro Thr
35	40	45
Thr Ser Leu Asp Cys	Ala Asn Ala Ser Asn	Val Gln Ala Val Asn His
50	55	60
Ser Ala Thr Lys Glu	Met Lys Phe Leu Pro	Pro Glu Pro Glu Trp Thr
65	70	75
Tyr Pro Arg Leu Ser	Cys Gln Gly Ser Thr	Phe Gln Lys Ala Leu Leu
85	90	95
Ile Ser Pro His Arg	Phe Gly Glu Ala Lys	Gly Asn Ser Ala Pro Leu
100	105	110
Ile Ile Arg Glu Pro	Phe Ile Ala Cys Gly	Pro Lys Glu Cys Lys His
115	120	125
Phe Ala Leu Thr His	Tyr Ala Ala Gln Pro	Gly Gly Tyr Tyr Asn Gly
130	135	140
Thr Arg Glu Asp Arg	Asn Lys Leu Arg His	Leu Ile Ser Val Asn Leu
145	150	155
		160

-continued

Gly Lys Ile Pro Thr Val Glu Asn Ser Ile Phe His Met Ala Ala Trp
 165 170 175
 Ser Gly Ser Ala Cys His Asp Gly Arg Glu Trp Thr Tyr Ile Gly Val
 180 185 190
 Asp Gly Pro Asp Ser Asn Ala Leu Ile Lys Ile Lys Tyr Gly Glu Ala
 195 200 205
 Tyr Thr Asp Thr Tyr His Ser Tyr Ala Asn Asn Ile Leu Arg Thr Gln
 210 215 220
 Glu Ser Ala Cys Asn Cys Ile Gly Gly Asp Cys Tyr Leu Met Ile Thr
 225 230 235 240
 Asp Gly Ser Ala Ser Gly Ile Ser Lys Cys Arg Phe Leu Lys Ile Arg
 245 250 255
 Glu Gly Arg Ile Ile Lys Glu Ile Phe Pro Thr Gly Arg Val Glu His
 260 265 270
 Thr Glu Glu Cys Thr Cys Gly Phe Ala Ser Asn Lys Thr Ile Glu Cys
 275 280 285
 Ala Cys Arg Asp Asn Ser Tyr Thr Ala Lys Arg Pro Phe Val Lys Leu
 290 295 300
 Asn Val Glu Thr Asp Thr Ala Glu Ile Arg Leu Met Cys Thr Glu Thr
 305 310 315 320
 Tyr Leu Asp Thr Pro Arg Pro Asp Asp Gly Ser Ile Thr Gly Pro Cys
 325 330 335
 Glu Ser Asn Gly Asp Lys Gly Ser Gly Gly Val Lys Gly Gly Phe Val
 340 345 350
 His Gln Arg Met Ala Ser Lys Ile Gly Arg Trp Tyr Ser Arg Thr Met
 355 360 365
 Ser Lys Thr Lys Arg Met Gly Met Glu Leu Tyr Val Lys Tyr Asp Gly
 370 375 380
 Asp Pro Trp Thr Asp Ser Asp Ala Leu Ala Pro Ser Gly Val Met Val
 385 390 395 400
 Ser Met Glu Glu Pro Gly Trp Tyr Ser Phe Gly Phe Glu Ile Lys Asp
 405 410 415
 Lys Lys Cys Asp Val Pro Cys Ile Gly Ile Glu Met Val His Asp Gly
 420 425 430
 Gly Lys Arg Thr Trp His Ser Ala Ala Thr Ala Ile Tyr Cys Leu Met
 435 440 445
 Gly Ser Gly Gln Leu Leu Trp Asp Thr Val Thr Gly Val Asn Met Ala
 450 455 460
 Leu
 465

<210> SEQ ID NO 57

<211> LENGTH: 584

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 57

Met Lys Ala Ile Ile Val Leu Leu Met Val Val Thr Ser Asn Ala Asp
 1 5 10 15
 Arg Ile Cys Thr Gly Ile Thr Ser Ser Asn Ser Pro His Val Val Lys
 20 25 30
 Thr Ala Thr Gln Gly Glu Val Asn Val Thr Gly Val Ile Pro Leu Thr
 35 40 45

-continued

Thr	Thr	Pro	Thr	Lys	Ser	His	Phe	Ala	Asn	Leu	Lys	Gly	Thr	Lys	Thr
50					55					60					
Arg	Gly	Lys	Leu	Cys	Pro	Thr	Cys	Leu	Asn	Cys	Thr	Asp	Leu	Asp	Val
65					70					75					80
Ala	Leu	Gly	Arg	Pro	Met	Cys	Val	Gly	Val	Thr	Pro	Ser	Ala	Lys	Ala
85					90					95					
Ser	Ile	Leu	His	Glu	Val	Arg	Pro	Val	Thr	Ser	Gly	Cys	Phe	Pro	Ile
100					105					110					
Met	His	Asp	Arg	Thr	Lys	Ile	Arg	Gln	Leu	Pro	Asn	Leu	Leu	Arg	Gly
115					120					125					
Tyr	Glu	Lys	Ile	Arg	Leu	Ser	Thr	Gln	Ile	Val	Ile	Asn	Ala	Glu	Lys
130					135					140					
Ala	Pro	Gly	Gly	Pro	Tyr	Arg	Leu	Gly	Thr	Ser	Gly	Ser	Cys	Pro	Asn
145					150					155					160
Ala	Thr	Ser	Arg	Ser	Gly	Phe	Phe	Ala	Thr	Met	Ala	Trp	Ala	Val	Pro
165					170					175					
Lys	Asp	Asn	Asn	Lys	Thr	Ala	Thr	Asn	Pro	Leu	Thr	Val	Glu	Val	Pro
180					185					190					
His	Ile	Cys	Thr	Lys	Glu	Glu	Asp	Gln	Ile	Thr	Val	Trp	Gly	Phe	His
195					200					205					
Ser	Asp	Asp	Lys	Thr	Gln	Met	Lys	Asn	Leu	Tyr	Gly	Asp	Ser	Asn	Pro
210					215					220					
Gln	Lys	Phe	Thr	Ser	Ser	Ala	Asn	Gly	Val	Thr	Thr	His	Tyr	Val	Ser
225					230					235					240
Gln	Ile	Gly	Gly	Phe	Pro	Asp	Gln	Thr	Glu	Asp	Gly	Gly	Leu	Pro	Gln
245					250					255					
Ser	Gly	Arg	Ile	Val	Val	Asp	Tyr	Met	Val	Gln	Lys	Pro	Gly	Lys	Thr
260					265					270					
Gly	Thr	Ile	Val	Tyr	Gln	Arg	Gly	Ile	Leu	Leu	Pro	Gln	Lys	Val	Trp
275					280					285					
Cys	Ala	Ser	Gly	Arg	Ser	Lys	Val	Ile	Lys	Gly	Ser	Leu	Pro	Leu	Ile
290					295					300					
Gly	Glu	Ala	Asp	Cys	Leu	His	Glu	Lys	Tyr	Gly	Gly	Leu	Asn	Lys	Ser
305					310					315					320
Lys	Pro	Tyr	Tyr	Thr	Gly	Glu	His	Ala	Lys	Ala	Ile	Gly	Asn	Cys	Pro
325					330					335					
Ile	Trp	Val	Lys	Thr	Pro	Leu	Lys	Leu	Ala	Asn	Gly	Thr	Lys	Tyr	Arg
340					345					350					
Pro	Pro	Ala	Lys	Leu	Leu	Lys	Glu	Arg	Gly	Phe	Phe	Gly	Ala	Ile	Ala
355					360					365					
Gly	Phe	Leu	Glu	Gly	Gly	Trp	Glu	Gly	Met	Ile	Ala	Gly	Trp	His	Gly
370					375					380					
Tyr	Thr	Ser	His	Gly	Ala	His	Gly	Val	Ala	Val	Ala	Ala	Asp	Leu	Lys
385					390					395					400
Ser	Thr	Gln	Glu	Ala	Ile	Asn	Lys	Ile	Thr	Lys	Asn	Leu	Asn	Ser	Leu
405					410					415					
Ser	Glu	Leu	Glu	Val	Lys	Asn	Leu	Gln	Arg	Leu	Ser	Gly	Ala	Met	Asp
420					425					430					
Glu	Leu	His	Asn	Glu	Ile	Leu	Glu	Leu	Asp	Glu	Lys	Val	Asp	Asp	Leu
435					440					445					

-continued

Arg	Ala	Asp	Thr	Ile	Ser	Ser	Gln	Ile	Glu	Leu	Ala	Val	Leu	Leu	Ser
450					455					460					
Asn	Glu	Gly	Ile	Ile	Asn	Ser	Glu	Asp	Glu	His	Leu	Leu	Ala	Leu	Glu
465					470					475					480
Arg	Lys	Leu	Lys	Lys	Met	Leu	Gly	Pro	Ser	Ala	Val	Asp	Ile	Gly	Asn
485					490					495					
Gly	Cys	Phe	Glu	Thr	Lys	His	Lys	Cys	Asn	Gln	Thr	Cys	Leu	Asp	Arg
500					505					510					
Ile	Ala	Ala	Gly	Thr	Phe	Asn	Ala	Gly	Glu	Phe	Ser	Leu	Pro	Thr	Phe
515					520					525					
Asp	Ser	Leu	Asn	Ile	Thr	Ala	Ala	Ser	Leu	Asn	Asp	Asp	Gly	Leu	Asp
530					535					540					
Asn	His	Thr	Ile	Leu	Leu	Tyr	Tyr	Ser	Thr	Ala	Ala	Ser	Ser	Leu	Ala
545					550					555					560
Val	Thr	Leu	Met	Ile	Ala	Ile	Phe	Ile	Val	Tyr	Met	Ile	Ser	Arg	Asp
565					570					575					
Asn	Val	Ser	Cys	Ser	Ile	Cys	Leu								
580															

<210> SEQ ID NO 58

<211> LENGTH: 466

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 58

Met	Leu	Pro	Ser	Thr	Ile	Gln	Thr	Leu	Thr	Leu	Phe	Leu	Thr	Ser	Gly
1				5					10					15	
Gly	Val	Leu	Leu	Ser	Leu	Tyr	Val	Ser	Ala	Ser	Leu	Ser	Tyr	Leu	Leu
20				25					30						
Tyr	Ser	Asp	Ile	Leu	Leu	Lys	Phe	Ser	Pro	Thr	Glu	Ile	Thr	Ala	Pro
35				40					45						
Thr	Met	Pro	Leu	Asn	Cys	Ala	Asn	Ala	Ser	Asn	Val	Gln	Ala	Val	Asn
50				55					60						
Arg	Ser	Ala	Thr	Lys	Gly	Val	Thr	Leu	Pro	Leu	Pro	Glu	Pro	Glu	Trp
65				70					75						80
Thr	Tyr	Pro	Arg	Leu	Ser	Cys	Pro	Gly	Ser	Thr	Phe	Gln	Lys	Ala	Leu
85				90					95						
Leu	Ile	Ser	Pro	His	Arg	Phe	Gly	Glu	Thr	Lys	Gly	Asn	Ser	Ala	Pro
100				105					110						
Leu	Ile	Ile	Arg	Glu	Pro	Phe	Ile	Ala	Cys	Gly	Pro	Lys	Glu	Cys	Arg
115				120					125						
His	Phe	Ala	Leu	Thr	His	Tyr	Ala	Ala	Gln	Pro	Gly	Gly	Tyr	Tyr	Asn
130				135					140						
Gly	Thr	Arg	Glu	Asp	Arg	Asn	Lys	Leu	Arg	His	Leu	Ile	Ser	Val	Lys
145				150					155						160
Leu	Gly	Lys	Ile	Pro	Thr	Val	Glu	Asn	Ser	Ile	Phe	His	Met	Ala	Ala
165				170					175						
Trp	Ser	Gly	Ser	Ala	Cys	His	Asp	Gly	Arg	Glu	Trp	Thr	Tyr	Ile	Gly
180				185					190						
Val	Asp	Gly	Pro	Asp	Ser	Asn	Ala	Leu	Leu	Lys	Ile	Lys	Tyr	Gly	Glu
195				200					205						
Ala	Tyr	Thr	Asp	Thr	Tyr	His	Ser	Tyr	Ala	Asn	Asn	Ile	Leu	Arg	Thr
210				215					220						

-continued

Gln Glu Ser Ala Cys Asn Cys Ile Gly Gly Asp Cys Tyr Leu Met Ile
 225 230 235 240
 Thr Asp Gly Ser Ala Ser Gly Ile Ser Glu Cys Arg Phe Leu Lys Ile
 245 250 255
 Arg Glu Gly Arg Ile Ile Lys Glu Ile Phe Pro Thr Gly Arg Val Glu
 260 265 270
 His Thr Glu Glu Cys Thr Cys Gly Phe Ala Ser Asn Lys Thr Ile Glu
 275 280 285
 Cys Ala Cys Arg Asp Asn Ser Tyr Thr Ala Lys Arg Pro Phe Val Lys
 290 295 300
 Leu Asn Val Glu Thr Asp Thr Ala Glu Ile Arg Leu Met Cys Thr Glu
 305 310 315 320
 Thr Tyr Leu Asp Thr Pro Arg Pro Asp Asp Gly Ser Ile Thr Gly Pro
 325 330 335
 Cys Glu Ser Asn Gly Asp Lys Gly Ser Gly Gly Ile Lys Gly Gly Phe
 340 345 350
 Val His Gln Arg Met Ala Ser Lys Ile Gly Arg Trp Tyr Ser Arg Thr
 355 360 365
 Met Ser Lys Thr Lys Arg Met Gly Met Gly Leu Tyr Val Lys Tyr Asp
 370 375 380
 Gly Asp Pro Trp Ile Asp Ser Asp Ala Leu Thr Leu Ser Gly Val Met
 385 390 395 400
 Val Ser Met Glu Glu Pro Gly Trp Tyr Ser Phe Gly Phe Glu Ile Lys
 405 410 415
 Asp Lys Lys Cys Asp Val Pro Cys Ile Gly Ile Glu Met Val His Asp
 420 425 430
 Gly Gly Lys Lys Thr Trp His Ser Ala Ala Thr Ala Ile Tyr Cys Leu
 435 440 445
 Met Gly Ser Gly Gln Leu Leu Trp Asp Thr Val Thr Gly Val Asp Met
 450 455 460
 Ala Leu
 465

<210> SEQ ID NO 59

<211> LENGTH: 584

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 59

Met Lys Ala Ile Ile Val Leu Leu Met Val Val Thr Ser Asn Ala Asp
 1 5 10 15
 Arg Ile Cys Thr Gly Ile Thr Ser Ser Asn Ser Pro His Val Val Lys
 20 25 30
 Thr Ala Thr Gln Gly Glu Val Asn Val Thr Gly Ala Ile Pro Leu Thr
 35 40 45
 Thr Thr Pro Thr Lys Ser His Phe Ala Asn Leu Lys Gly Thr Lys Thr
 50 55 60
 Arg Gly Lys Leu Cys Pro Thr Cys Leu Asn Cys Thr Asp Leu Asp Val
 65 70 75 80
 Ala Leu Gly Arg Pro Met Cys Val Gly Ile Thr Pro Ser Ala Lys Ala
 85 90 95
 Ser Ile Leu His Glu Val Arg Pro Val Thr Ser Gly Cys Phe Pro Ile

-continued

100					105					110				
Met	His	Asp	Arg	Thr	Lys	Ile	Arg	Gln	Leu	Pro	Asn	Leu	Leu	Arg Gly
115					120					125				
Tyr	Glu	Lys	Ile	Arg	Leu	Ser	Thr	Gln	Asn	Val	Ile	Asn	Ala	Glu Lys
130					135					140				
Ala	Pro	Gly	Gly	Pro	Tyr	Arg	Leu	Gly	Thr	Ser	Gly	Ser	Cys	Pro Asn
145					150					155				160
Ala	Thr	Ser	Lys	Ser	Gly	Phe	Phe	Ala	Thr	Met	Ala	Trp	Ala	Val Pro
165					170					175				
Arg	Asp	Asn	Asn	Lys	Thr	Ala	Thr	Asn	Pro	Leu	Thr	Val	Glu	Val Pro
180					185					190				
His	Ile	Cys	Thr	Lys	Glu	Glu	Asp	Gln	Ile	Thr	Val	Trp	Gly	Phe His
195					200					205				
Ser	Asp	Asp	Lys	Thr	Gln	Met	Lys	Asn	Leu	Tyr	Gly	Asp	Ser	Asn Pro
210					215					220				
Gln	Lys	Phe	Thr	Ser	Ser	Ala	Asn	Gly	Ile	Thr	Thr	His	Tyr	Val Ser
225					230					235				240
Gln	Ile	Gly	Gly	Phe	Pro	Asp	Gln	Thr	Glu	Asp	Gly	Gly	Leu	Pro Gln
245					250					255				
Ser	Gly	Arg	Ile	Val	Val	Asp	Tyr	Met	Val	Gln	Lys	Pro	Gly	Lys Thr
260					265					270				
Gly	Thr	Ile	Val	Tyr	Gln	Arg	Gly	Ile	Leu	Leu	Pro	Gln	Lys	Val Trp
275					280					285				
Cys	Ala	Ser	Gly	Arg	Ser	Lys	Val	Ile	Lys	Gly	Ser	Leu	Pro	Leu Ile
290					295					300				
Gly	Glu	Ala	Asp	Cys	Leu	His	Glu	Lys	Tyr	Gly	Gly	Leu	Asn	Lys Ser
305					310					315				320
Lys	Pro	Tyr	Tyr	Thr	Gly	Glu	His	Ala	Lys	Ala	Ile	Gly	Asn	Cys Pro
325					330					335				
Ile	Trp	Val	Lys	Thr	Pro	Leu	Lys	Leu	Ala	Asn	Gly	Thr	Lys	Tyr Arg
340					345					350				
Pro	Pro	Ala	Lys	Leu	Leu	Lys	Glu	Arg	Gly	Phe	Phe	Gly	Ala	Ile Ala
355					360					365				
Gly	Phe	Leu	Glu	Gly	Gly	Trp	Glu	Gly	Met	Ile	Ala	Gly	Trp	His Gly
370					375					380				
Tyr	Thr	Ser	His	Gly	Ala	His	Gly	Val	Ala	Val	Ala	Ala	Asp	Leu Lys
385					390					395				400
Ser	Thr	Gln	Glu	Ala	Ile	Asn	Lys	Ile	Thr	Lys	Asn	Leu	Asn	Ser Leu
405					410					415				
Ser	Glu	Leu	Glu	Val	Lys	Asn	Leu	Gln	Arg	Leu	Ser	Gly	Ala	Met Asp
420					425					430				
Glu	Leu	His	Asn	Glu	Ile	Leu	Glu	Leu	Asp	Glu	Lys	Val	Asp	Asp Leu
435					440					445				
Arg	Ala	Asp	Thr	Ile	Ser	Ser	Gln	Ile	Glu	Leu	Ala	Val	Leu	Leu Ser
450					455					460				
Asn	Glu	Gly	Ile	Ile	Asn	Ser	Glu	Asp	Glu	His	Leu	Leu	Ala	Leu Glu
465					470					475				480
Arg	Lys	Leu	Lys	Lys	Met	Leu	Gly	Pro	Ser	Ala	Ile	Asp	Ile	Gly Asn
485					490					495				
Gly	Cys	Phe	Glu	Thr	Lys	His	Lys	Cys	Asn	Gln	Thr	Cys	Leu	Asp Arg
500					505					510				

-continued

```

Ile Ala Ala Gly Thr Phe Asn Ala Gly Glu Phe Ser Leu Pro Thr Phe
515          520          525

Asp Ser Leu Asn Ile Thr Ala Ala Ser Leu Asn Asp Asp Gly Leu Asp
530          535          540

Asn His Thr Ile Leu Leu Tyr Tyr Ser Thr Ala Ala Ser Ser Leu Ala
545          550          555          560

Val Thr Leu Met Ile Ala Ile Phe Ile Val Tyr Met Ile Ser Arg Asp
565          570          575

Asn Val Ser Cys Ser Ile Cys Leu
580

```

```

<210> SEQ ID NO 60
<211> LENGTH: 466
<212> TYPE: PRT
<213> ORGANISM: Influenza B virus

```

```

<400> SEQUENCE: 60

```

```

Met Leu Pro Ser Thr Ile Gln Thr Leu Thr Leu Phe Leu Thr Ser Gly
1          5          10          15

Gly Val Leu Leu Ser Leu Tyr Val Ser Ala Ser Leu Ser Tyr Leu Leu
20          25          30

Tyr Ser Asp Ile Leu Leu Lys Phe Ser Pro Thr Glu Ile Thr Ala Pro
35          40          45

Ala Met Pro Leu Asp Cys Ala Asn Ala Ser Asn Val Gln Ala Val Asn
50          55          60

Arg Ser Ala Thr Lys Gly Val Thr Leu Leu Leu Pro Glu Pro Glu Trp
65          70          75          80

Thr Tyr Pro Arg Leu Ser Cys Pro Gly Ser Thr Phe Gln Lys Ala Leu
85          90          95

Leu Ile Ser Pro His Arg Phe Gly Glu Thr Lys Gly Asn Ser Ala Pro
100         105         110

Leu Ile Ile Arg Glu Pro Phe Ile Ala Cys Gly Pro Lys Glu Cys Lys
115         120         125

His Phe Ala Leu Thr His Tyr Ala Ala Gln Pro Gly Gly Tyr Tyr Asn
130         135         140

Gly Thr Arg Glu Asp Arg Asn Lys Leu Arg His Leu Ile Ser Val Lys
145         150         155         160

Phe Gly Lys Ile Pro Thr Val Glu Asn Ser Ile Phe His Met Ala Ala
165         170         175

Trp Ser Gly Ser Ala Cys His Asp Gly Lys Glu Trp Thr Tyr Ile Gly
180         185         190

Val Asp Gly Pro Asp Ser Asn Ala Leu Leu Lys Ile Lys Tyr Gly Glu
195         200         205

Ala Tyr Thr Asp Thr Tyr His Ser Tyr Ala Asn Asn Ile Leu Arg Thr
210         215         220

Gln Glu Ser Ala Cys Asn Cys Ile Gly Gly Asn Cys Tyr Leu Met Ile
225         230         235         240

Thr Asp Gly Ser Ala Ser Gly Ile Ser Glu Cys Arg Phe Leu Lys Ile
245         250         255

Arg Glu Gly Arg Ile Ile Lys Glu Ile Phe Pro Thr Gly Arg Val Lys
260         265         270

His Thr Glu Glu Cys Thr Cys Gly Phe Ala Ser Asn Lys Thr Ile Glu

```

-continued

275	280	285
Cys Ala Cys Arg Asp	Asn Ser Tyr Thr Ala	Lys Arg Pro Phe Val Lys
290	295	300
Leu Asn Val Glu Thr	Asp Thr Ala Glu Ile	Arg Leu Met Cys Thr Glu
305	310	315 320
Thr Tyr Leu Asp Thr	Pro Arg Pro Asp Asp	Gly Ser Ile Thr Gly Pro
325	330	335
Cys Glu Ser Asn Gly	Asp Lys Gly Ser Gly	Gly Ile Lys Gly Gly Phe
340	345	350
Val His Gln Arg Met	Ala Ser Lys Ile Gly	Arg Trp Tyr Ser Arg Thr
355	360	365
Met Ser Lys Thr Lys	Arg Met Gly Met Gly	Leu Tyr Val Lys Tyr Asp
370	375	380
Gly Asp Pro Trp Thr	Asp Ser Asp Ala Leu	Ala Leu Ser Gly Val Met
385	390	395 400
Val Ser Met Glu Glu	Pro Gly Trp Tyr Ser	Phe Gly Phe Glu Ile Lys
405	410	415
Asp Lys Lys Cys Asp	Val Pro Cys Ile Gly	Ile Glu Met Val His Asp
420	425	430
Gly Gly Lys Glu Thr	Trp His Ser Ala Ala	Thr Ala Ile Tyr Cys Leu
435	440	445
Met Gly Ser Gly Gln	Leu Leu Trp Asp Thr	Val Thr Gly Val Asp Met
450	455	460
Ala Leu		
465		

<210> SEQ ID NO 61
 <211> LENGTH: 583
 <212> TYPE: PRT
 <213> ORGANISM: Influenza B virus

<400> SEQUENCE: 61

Met Lys Ala Ile Ile	Val Leu Leu Met Val	Val Thr Ser Asn Ala Asp
1	5	10 15
Arg Ile Cys Thr Gly	Ile Thr Ser Ser Asn	Ser Pro His Val Val Lys
20	25	30
Thr Ala Thr Gln Gly	Glu Val Asn Val Thr	Gly Val Ile Pro Leu Thr
35	40	45
Thr Thr Pro Thr Lys	Ser His Phe Ala Asn	Leu Lys Gly Thr Lys Thr
50	55	60
Arg Gly Lys Leu Cys	Pro Asn Cys Leu Asn	Cys Thr Asp Leu Asp Val
65	70	75 80
Ala Leu Gly Arg Pro	Met Cys Ile Gly Ile	Thr Pro Ser Ala Lys Ala
85	90	95
Ser Ile Leu His Glu	Val Arg Pro Val Thr	Ser Gly Cys Phe Pro Ile
100	105	110
Met His Asp Arg Thr	Lys Ile Arg Gln Leu	Pro Asn Leu Leu Arg Gly
115	120	125
Tyr Glu His Ile Arg	Leu Ser Thr His Asn	Val Ile Asn Ala Glu Arg
130	135	140
Ala Pro Gly Gly Pro	Tyr Arg Leu Gly Thr	Ser Gly Ser Cys Pro Asn
145	150	155 160

-continued

Val Thr Ser Arg Ser Gly Phe Phe Ala Thr Met Ala Trp Ala Val Pro	
165	170 175
Arg Asp Asn Lys Thr Ala Thr Asn Pro Leu Thr Val Glu Val Pro Tyr	
180	185 190
Ile Cys Thr Lys Gly Glu Asp Gln Ile Thr Val Trp Gly Phe His Ser	
195	200 205
Asp Asn Lys Ile Gln Met Lys Asn Leu Tyr Gly Asp Ser Asn Pro Gln	
210	215 220
Lys Phe Thr Ser Ser Ala Asn Gly Ile Thr Thr His Tyr Val Ser Gln	
225	230 235 240
Ile Gly Gly Phe Pro Asn Gln Thr Glu Asp Gly Gly Leu Pro Gln Ser	
245	250 255
Gly Arg Ile Val Val Asp Tyr Met Val Gln Lys Pro Gly Lys Thr Gly	
260	265 270
Thr Ile Val Tyr Gln Arg Gly Val Leu Leu Pro Gln Lys Val Trp Cys	
275	280 285
Ala Ser Gly Arg Ser Lys Val Ile Lys Gly Ser Leu Pro Leu Ile Gly	
290	295 300
Glu Ala Asp Cys Leu His Glu Lys Tyr Gly Gly Leu Asn Lys Ser Lys	
305	310 315 320
Pro Tyr Tyr Thr Gly Glu His Ala Lys Ala Ile Gly Asn Cys Pro Ile	
325	330 335
Trp Val Lys Thr Pro Leu Lys Leu Ala Asn Gly Thr Lys Tyr Arg Pro	
340	345 350
Pro Ala Lys Leu Leu Lys Glu Lys Gly Phe Phe Gly Ala Ile Ala Gly	
355	360 365
Phe Leu Glu Gly Gly Trp Glu Gly Met Ile Ala Gly Trp His Gly Tyr	
370	375 380
Thr Ser His Gly Ala His Gly Val Ala Val Ala Ala Asp Leu Lys Ser	
385	390 395 400
Thr Gln Glu Ala Ile Asn Lys Ile Thr Lys Asn Leu Asn Ser Leu Ser	
405	410 415
Glu Leu Glu Val Lys Asn Leu Gln Arg Leu Ser Gly Ala Met Asp Glu	
420	425 430
Leu His Asn Glu Ile Leu Glu Leu Asp Glu Lys Val Asp Asp Leu Arg	
435	440 445
Ala Asp Thr Ile Ser Ser Gln Ile Glu Leu Ala Val Leu Leu Ser Asn	
450	455 460
Glu Gly Ile Ile Asn Ser Glu Asp Glu His Leu Leu Ala Leu Glu Arg	
465	470 475 480
Lys Leu Lys Lys Met Leu Gly Pro Ser Ala Val Asp Ile Gly Asn Gly	
485	490 495
Cys Phe Glu Thr Lys His Lys Cys Asn Gln Thr Cys Leu Asp Arg Ile	
500	505 510
Ala Ala Gly Thr Phe Asn Ala Gly Glu Phe Ser Leu Pro Thr Phe Asp	
515	520 525
Ser Leu Asn Ile Thr Ala Ala Ser Leu Asn Asp Asp Gly Leu Asp Asn	
530	535 540
His Thr Ile Leu Leu Tyr Tyr Ser Thr Ala Ala Ser Ser Leu Ala Val	
545	550 555 560
Thr Leu Met Ile Ala Ile Phe Ile Val Tyr Met Val Ser Arg Asp Asn	

-continued

565 570 575

Val Ser Cys Ser Ile Cys Leu
580

<210> SEQ ID NO 62
<211> LENGTH: 465
<212> TYPE: PRT
<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 62

Met Leu Pro Ser Thr Ile Gln Thr Leu Thr Leu Phe Leu Thr Ser Gly
1 5 10 15

Gly Val Leu Leu Ser Leu Tyr Val Ser Ala Leu Leu Ser Tyr Leu Leu
20 25 30

Tyr Ser Asp Ile Leu Leu Lys Phe Ser Pro Lys Ile Ile Ala Pro Thr
35 40 45

Thr Ser Leu Asp Cys Ala Asn Ala Ser Asn Val Gln Ala Val Asn His
50 55 60

Ser Ala Thr Lys Glu Met Lys Phe Leu Pro Pro Glu Pro Glu Trp Thr
65 70 75 80

Tyr Pro Arg Leu Ser Cys Gln Gly Ser Thr Phe Gln Lys Ala Leu Leu
85 90 95

Ile Ser Pro His Arg Phe Gly Glu Ala Lys Gly Asn Ser Ala Pro Leu
100 105 110

Ile Ile Arg Glu Pro Phe Ile Ala Cys Gly Pro Lys Glu Cys Lys His
115 120 125

Phe Ala Leu Thr His Tyr Ala Ala Gln Pro Gly Gly Tyr Tyr Asn Gly
130 135 140

Thr Arg Glu Asp Arg Asn Lys Leu Arg His Leu Ile Ser Val Asn Leu
145 150 155 160

Gly Lys Ile Pro Thr Val Glu Asn Ser Ile Phe His Met Ala Ala Trp
165 170 175

Ser Gly Ser Ala Cys His Asp Gly Arg Glu Trp Thr Tyr Ile Gly Val
180 185 190

Asp Gly Pro Asp Ser Asn Ala Leu Ile Lys Ile Lys Tyr Gly Glu Ala
195 200 205

Tyr Thr Asp Thr Tyr His Ser Tyr Ala Asn Asn Ile Leu Arg Thr Gln
210 215 220

Glu Ser Ala Cys Asn Cys Ile Gly Gly Asp Cys Tyr Leu Met Ile Thr
225 230 235 240

Asp Gly Ser Ala Ser Gly Ile Ser Lys Cys Arg Phe Leu Lys Ile Arg
245 250 255

Glu Gly Arg Ile Ile Lys Glu Ile Phe Pro Thr Gly Arg Val Glu His
260 265 270

Thr Glu Glu Cys Thr Cys Gly Phe Ala Ser Asn Lys Thr Ile Glu Cys
275 280 285

Ala Cys Arg Asp Asn Ser Tyr Thr Ala Lys Arg Pro Phe Val Lys Leu
290 295 300

Asn Val Glu Thr Asp Thr Ala Glu Ile Arg Leu Met Cys Thr Glu Thr
305 310 315 320

Tyr Leu Asp Thr Pro Arg Pro Asp Asp Gly Ser Ile Thr Gly Pro Cys
325 330 335

-continued

Glu Ser Asn Gly Asp Lys Gly Ser Gly Gly Val Lys Gly Gly Phe Val
 340 345 350
 His Gln Arg Met Ala Ser Lys Ile Gly Arg Trp Tyr Ser Arg Thr Met
 355 360 365
 Ser Lys Thr Lys Arg Met Gly Met Glu Leu Tyr Val Lys Tyr Asp Gly
 370 375 380
 Asp Pro Trp Thr Asp Ser Asp Ala Leu Ala Pro Ser Gly Val Met Val
 385 390 395 400
 Ser Met Glu Glu Pro Gly Trp Tyr Ser Phe Gly Phe Glu Ile Lys Asp
 405 410 415
 Lys Lys Cys Asp Val Pro Cys Ile Gly Ile Glu Met Val His Asp Gly
 420 425 430
 Gly Lys Arg Thr Trp His Ser Ala Ala Thr Ala Ile Tyr Cys Leu Met
 435 440 445
 Gly Ser Gly Gln Leu Leu Trp Asp Thr Val Thr Gly Val Asn Met Ala
 450 455 460
 Leu
 465

<210> SEQ ID NO 63
 <211> LENGTH: 585
 <212> TYPE: PRT
 <213> ORGANISM: Influenza B virus

<400> SEQUENCE: 63

Met Lys Ala Ile Ile Val Leu Leu Met Val Val Thr Ser Asn Ala Asp
 1 5 10 15
 Arg Ile Cys Thr Gly Ile Thr Ser Ser Asn Ser Pro His Val Val Lys
 20 25 30
 Thr Ala Thr Gln Gly Glu Val Asn Val Thr Gly Val Ile Pro Leu Thr
 35 40 45
 Thr Thr Pro Thr Lys Ser His Phe Ala Asn Leu Lys Gly Thr Lys Thr
 50 55 60
 Arg Gly Lys Leu Cys Pro Lys Cys Leu Asn Cys Thr Asp Leu Asp Val
 65 70 75 80
 Ala Leu Gly Arg Pro Lys Cys Thr Gly Asn Ile Pro Ser Ala Lys Val
 85 90 95
 Ser Ile Leu His Glu Val Arg Pro Val Thr Ser Gly Cys Phe Pro Ile
 100 105 110
 Met His Asp Arg Thr Lys Ile Arg Gln Leu Pro Asn Leu Leu Arg Gly
 115 120 125
 Tyr Glu Arg Ile Arg Leu Ser Asn His Asn Val Ile Asn Ala Glu Lys
 130 135 140
 Ala Pro Gly Gly Pro Tyr Lys Ile Gly Thr Ser Gly Ser Cys Pro Asn
 145 150 155 160
 Val Thr Asn Gly Asn Gly Phe Phe Ala Thr Met Ala Trp Ala Val Pro
 165 170 175
 Lys Asn Glu Asn Asn Lys Thr Ala Thr Asn Ser Leu Thr Ile Glu Val
 180 185 190
 Pro Tyr Ile Cys Thr Glu Gly Glu Asp Gln Ile Thr Val Trp Gly Phe
 195 200 205
 His Ser Asp Ser Glu Thr Gln Met Ala Lys Leu Tyr Gly Asp Ser Lys
 210 215 220

-continued

```

Pro Gln Lys Phe Thr Ser Ser Ala Asn Gly Val Thr Thr His Tyr Val
225                230                235                240

Ser Gln Ile Gly Gly Phe Pro Asn Gln Thr Glu Asp Gly Gly Leu Pro
245                250                255

Gln Ser Gly Arg Ile Val Val Asp Tyr Met Val Gln Lys Ser Gly Lys
260                265                270

Thr Gly Thr Ile Thr Tyr Gln Arg Gly Ile Leu Leu Pro Gln Lys Val
275                280                285

Trp Cys Ala Ser Gly Arg Ser Lys Val Ile Lys Gly Ser Leu Pro Leu
290                295                300

Ile Gly Glu Ala Asp Cys Leu His Glu Lys Tyr Gly Gly Leu Asn Lys
305                310                315                320

Ser Lys Pro Tyr Tyr Thr Gly Glu His Ala Lys Ala Ile Gly Asn Cys
325                330                335

Pro Ile Trp Val Lys Thr Pro Leu Lys Leu Ala Asn Gly Thr Lys Tyr
340                345                350

Arg Pro Pro Ala Lys Leu Leu Lys Glu Arg Gly Phe Phe Gly Ala Ile
355                360                365

Ala Gly Phe Leu Glu Gly Gly Trp Glu Gly Met Ile Ala Gly Trp His
370                375                380

Gly Tyr Thr Ser His Gly Ala His Gly Val Ala Val Ala Ala Asp Leu
385                390                395                400

Lys Ser Thr Gln Glu Ala Ile Asn Lys Ile Thr Lys Asn Leu Asn Ser
405                410                415

Leu Ser Glu Leu Glu Val Lys Asn Leu Gln Arg Leu Ser Gly Ala Met
420                425                430

Asp Glu Leu His Asn Glu Ile Leu Glu Leu Asp Glu Lys Val Asp Asp
435                440                445

Leu Arg Ala Asp Thr Ile Ser Ser Gln Ile Glu Leu Ala Val Leu Leu
450                455                460

Ser Asn Glu Gly Ile Ile Asn Ser Glu Asp Glu His Leu Leu Ala Leu
465                470                475                480

Glu Arg Lys Leu Lys Lys Met Leu Gly Pro Ser Ala Val Glu Ile Gly
485                490                495

Asn Gly Cys Phe Glu Thr Lys His Lys Cys Asn Gln Thr Cys Leu Asp
500                505                510

Arg Ile Ala Ala Gly Thr Phe Asn Ala Gly Glu Phe Ser Leu Pro Thr
515                520                525

Phe Asp Ser Leu Asn Ile Thr Ala Ala Ser Leu Asn Asp Asp Gly Leu
530                535                540

Asp Asn His Thr Ile Leu Leu Tyr Tyr Ser Thr Ala Ala Ser Ser Leu
545                550                555                560

Ala Val Thr Leu Met Ile Ala Ile Phe Val Val Tyr Met Val Ser Arg
565                570                575

Asp Asn Val Ser Cys Ser Ile Cys Leu
580                585

```

<210> SEQ ID NO 64

<211> LENGTH: 466

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

-continued

<400> SEQUENCE: 64

```

Met Leu Pro Ser Thr Ile Gln Thr Leu Thr Leu Phe Leu Thr Ser Gly
 1           5           10           15

Gly Val Leu Leu Ser Leu Tyr Val Ser Ala Leu Leu Ser Tyr Leu Leu
20           25           30

Tyr Ser Asp Ile Leu Leu Lys Phe Ser Pro Thr Lys Ile Ile Ala Pro
35           40           45

Thr Thr Ser Leu Asp Ser Ala Asn Ala Ser Asn Phe Gln Ala Val Asn
50           55           60

His Ser Ala Thr Lys Glu Met Thr Phe Leu Leu Pro Glu Pro Glu Trp
65           70           75           80

Thr Tyr Pro Arg Leu Ser Cys Gln Gly Ser Thr Phe Gln Lys Ala Leu
85           90           95

Leu Ile Ser Pro His Arg Phe Gly Glu Ala Lys Gly Asn Ser Ala Pro
100          105          110

Leu Ile Ile Arg Glu Pro Phe Ile Ala Cys Gly Pro Lys Glu Cys Lys
115          120          125

His Phe Ala Leu Thr His Tyr Ala Ala Gln Pro Gly Gly Tyr Tyr Asn
130          135          140

Gly Thr Arg Glu Asp Arg Asn Lys Leu Arg His Leu Ile Ser Val Asn
145          150          155          160

Leu Gly Lys Ile Pro Thr Val Glu Asn Ser Ile Phe His Met Ala Ala
165          170          175

Trp Ser Gly Ser Ala Cys His Asp Gly Arg Glu Trp Thr Tyr Ile Gly
180          185          190

Val Asp Gly Pro Asp Ser Asn Ala Leu Ile Lys Ile Lys Tyr Gly Glu
195          200          205

Ala Tyr Thr Asp Thr Tyr His Ser Tyr Ala Asn Asn Ile Leu Arg Thr
210          215          220

Gln Glu Ser Ala Cys Asn Cys Ile Gly Gly Asp Cys Tyr Leu Met Ile
225          230          235          240

Thr Asp Gly Ser Ala Ser Gly Ile Ser Lys Cys Arg Phe Leu Lys Ile
245          250          255

Arg Glu Gly Arg Ile Val Lys Glu Ile Phe Pro Thr Gly Arg Val Glu
260          265          270

His Thr Glu Glu Cys Thr Cys Gly Phe Ala Ser Asn Lys Thr Ile Glu
275          280          285

Cys Ala Cys Arg Asp Asn Ser Tyr Thr Ala Lys Arg Pro Phe Val Lys
290          295          300

Leu Asn Val Glu Thr Asp Thr Ala Glu Ile Arg Leu Met Cys Thr Glu
305          310          315          320

Thr Tyr Leu Asp Thr Pro Arg Pro Asp Asp Gly Ser Ile Thr Gly Pro
325          330          335

Cys Glu Ser Asn Gly Asp Lys Gly Ser Gly Gly Ile Lys Gly Gly Phe
340          345          350

Val His Gln Arg Met Ala Ser Lys Ile Gly Arg Trp Tyr Ser Arg Thr
355          360          365

Met Ser Lys Thr Lys Arg Met Gly Met Glu Leu Tyr Val Lys Tyr Asp
370          375          380

Gly Asp Pro Trp Thr Asp Ser Asp Ala Leu Ala Pro Ser Gly Val Met
385          390          395          400

```

-continued

Val Ser Ile Glu Glu Pro Gly Trp Tyr Ser Phe Gly Phe Glu Ile Lys
 405 410 415

Asp Lys Lys Cys Asp Val Pro Cys Ile Gly Ile Glu Met Val His Asp
 420 425 430

Gly Gly Lys Thr Thr Trp His Ser Ala Ala Thr Ala Ile Tyr Cys Leu
 435 440 445

Met Gly Ser Gly Gln Leu Leu Trp Asp Thr Ile Thr Gly Val Asp Met
 450 455 460

Ala Leu
 465

<210> SEQ ID NO 65
 <211> LENGTH: 585
 <212> TYPE: PRT
 <213> ORGANISM: Influenza B virus

<400> SEQUENCE: 65

Met Lys Ala Ile Ile Val Leu Leu Met Val Val Thr Ser Asn Ala Asp
 1 5 10 15

Arg Ile Cys Thr Gly Ile Thr Ser Ser Asn Ser Pro His Val Val Lys
 20 25 30

Thr Ala Thr Gln Gly Glu Val Asn Val Thr Gly Val Ile Pro Leu Thr
 35 40 45

Thr Thr Pro Thr Lys Ser His Phe Ala Asn Leu Lys Gly Thr Lys Thr
 50 55 60

Arg Gly Lys Leu Cys Pro Lys Cys Leu Asn Cys Thr Asp Leu Asp Val
 65 70 75 80

Ala Leu Gly Arg Pro Lys Cys Thr Gly Asn Ile Pro Ser Ala Lys Val
 85 90 95

Ser Ile Leu His Glu Val Arg Pro Val Thr Ser Gly Cys Phe Pro Ile
 100 105 110

Met His Asp Arg Thr Lys Ile Arg Gln Leu Pro Asn Leu Leu Arg Gly
 115 120 125

Tyr Glu His Ile Arg Leu Ser Thr His Asn Val Ile Asn Ala Glu Lys
 130 135 140

Ala Pro Gly Gly Pro Tyr Lys Ile Gly Thr Ser Gly Ser Cys Pro Asn
 145 150 155 160

Val Thr Asn Gly Asn Gly Phe Phe Ala Thr Met Ala Trp Ala Val Pro
 165 170 175

Lys Asn Asp Asn Asn Lys Thr Ala Thr Asn Ser Leu Thr Ile Glu Val
 180 185 190

Pro Tyr Ile Cys Thr Glu Gly Glu Asp Gln Ile Thr Val Trp Gly Phe
 195 200 205

His Ser Asp Asn Glu Ala Gln Met Ala Lys Leu Tyr Gly Asp Ser Lys
 210 215 220

Pro Gln Lys Phe Thr Ser Ser Ala Asn Gly Val Thr Thr His Tyr Val
 225 230 235 240

Ser Gln Ile Gly Gly Phe Pro Asn Gln Thr Glu Asp Gly Gly Leu Pro
 245 250 255

Gln Ser Gly Arg Ile Val Val Asp Tyr Met Val Gln Lys Ser Gly Lys
 260 265 270

Thr Gly Thr Ile Thr Tyr Gln Arg Gly Ile Leu Leu Pro Gln Lys Val

-continued

275	280	285
Trp Cys Ala Ser Gly	Arg Ser Lys Val Ile	Lys Gly Ser Leu Pro Leu
290	295	300
Ile Gly Glu Ala Asp	Cys Leu His Glu Lys	Tyr Gly Gly Leu Asn Lys
305	310	315 320
Ser Lys Pro Tyr Tyr	Thr Gly Glu His Ala	Lys Ala Ile Gly Asn Cys
325	330	335
Pro Ile Trp Val Lys	Thr Pro Leu Lys Leu	Ala Asn Gly Thr Lys Tyr
340	345	350
Arg Pro Pro Ala Lys	Leu Leu Lys Glu Arg	Gly Phe Phe Gly Ala Ile
355	360	365
Ala Gly Phe Leu Glu	Gly Gly Trp Glu Gly	Met Ile Ala Gly Trp His
370	375	380
Gly Tyr Thr Ser His	Gly Ala His Gly Val	Ala Val Ala Ala Asp Leu
385	390	395 400
Lys Ser Thr Gln Glu	Ala Ile Asn Lys Ile	Thr Lys Asn Leu Asn Ser
405	410	415
Leu Ser Glu Leu Glu	Val Lys Asn Leu Gln	Arg Leu Ser Gly Ala Met
420	425	430
Asp Glu Leu His Asn	Glu Ile Leu Glu Leu	Asp Glu Lys Val Asp Asp
435	440	445
Leu Arg Ala Asp Thr	Ile Ser Ser Gln Ile	Glu Leu Ala Val Leu Leu
450	455	460
Ser Asn Glu Gly Ile	Ile Asn Ser Glu Asp	Glu His Leu Leu Ala Leu
465	470	475 480
Glu Arg Lys Leu Lys	Lys Met Leu Gly Pro	Ser Ala Val Glu Ile Gly
485	490	495
Asn Gly Cys Phe Glu	Thr Lys His Lys Cys	Asn Gln Thr Cys Leu Asp
500	505	510
Arg Ile Ala Ala Gly	Thr Phe Asp Ala Gly	Glu Phe Ser Leu Pro Thr
515	520	525
Phe Asp Ser Leu Asn	Ile Thr Ala Ala Ser	Leu Asn Asp Asp Gly Leu
530	535	540
Asp Asn His Thr Ile	Leu Leu Tyr Tyr Ser	Thr Ala Ala Ser Ser Leu
545	550	555 560
Ala Val Thr Leu Met	Ile Ala Ile Phe Val	Val Tyr Met Val Ser Arg
565	570	575
Asp Asn Val Ser Cys	Ser Ile Cys Leu	
580	585	

<210> SEQ ID NO 66

<211> LENGTH: 466

<212> TYPE: PRT

<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 66

Met Leu Pro Ser Thr	Ile Gln Thr Leu Thr	Leu Phe Leu Thr Ser Gly
1	5	10 15
Gly Val Leu Leu Ser	Leu Tyr Val Ser Ala	Ser Leu Ser Tyr Leu Leu
20	25	30
Tyr Ser Asp Ile Leu	Leu Lys Phe Ser Pro	Thr Glu Ile Thr Ala Pro
35	40	45

-continued

Thr	Met	Pro	Leu	Asp	Cys	Ala	Asn	Ala	Ser	Asn	Val	Gln	Ala	Val	Asn
50					55					60					
Arg	Ser	Ala	Thr	Lys	Gly	Val	Thr	Leu	Leu	Leu	Pro	Glu	Pro	Glu	Trp
65					70					75					80
Thr	Tyr	Pro	Arg	Leu	Ser	Cys	Pro	Gly	Ser	Thr	Phe	Gln	Lys	Ala	Leu
85					90					95					
Leu	Ile	Ser	Pro	His	Arg	Phe	Gly	Glu	Thr	Lys	Gly	Asn	Ser	Ala	Pro
100					105					110					
Leu	Ile	Ile	Arg	Glu	Pro	Phe	Ile	Ala	Cys	Gly	Pro	Lys	Glu	Cys	Lys
115					120					125					
His	Phe	Ala	Leu	Thr	His	Tyr	Ala	Ala	Gln	Pro	Gly	Gly	Tyr	Tyr	Asn
130					135					140					
Gly	Thr	Arg	Gly	Asp	Arg	Asn	Lys	Leu	Arg	His	Leu	Ile	Ser	Val	Lys
145					150					155					160
Leu	Gly	Lys	Ile	Pro	Thr	Val	Glu	Asn	Ser	Ile	Phe	His	Met	Ala	Ala
165					170					175					
Trp	Ser	Gly	Ser	Ala	Cys	His	Asp	Gly	Lys	Glu	Trp	Thr	Tyr	Ile	Gly
180					185					190					
Val	Asp	Gly	Pro	Asp	Asn	Asn	Ala	Leu	Leu	Lys	Ile	Lys	Tyr	Gly	Glu
195					200					205					
Ala	Tyr	Thr	Asp	Thr	Tyr	His	Ser	Tyr	Ala	Asn	Asn	Ile	Leu	Arg	Thr
210					215					220					
Gln	Glu	Ser	Ala	Cys	Asn	Cys	Ile	Gly	Gly	Asn	Cys	Tyr	Leu	Met	Ile
225					230					235					240
Thr	Asp	Gly	Ser	Ala	Ser	Gly	Ile	Ser	Glu	Cys	Arg	Phe	Leu	Lys	Ile
245					250					255					
Arg	Glu	Gly	Arg	Ile	Ile	Lys	Glu	Ile	Phe	Pro	Thr	Gly	Arg	Val	Lys
260					265					270					
His	Thr	Glu	Glu	Cys	Thr	Cys	Gly	Phe	Ala	Ser	Asn	Lys	Thr	Ile	Glu
275					280					285					
Cys	Ala	Cys	Arg	Asp	Asn	Ser	Tyr	Thr	Ala	Lys	Arg	Pro	Phe	Val	Lys
290					295					300					
Leu	Asn	Val	Glu	Thr	Asp	Thr	Ala	Glu	Ile	Arg	Leu	Met	Cys	Thr	Glu
305					310					315					320
Thr	Tyr	Leu	Asp	Thr	Pro	Arg	Pro	Asp	Asp	Gly	Ser	Ile	Thr	Gly	Pro
325					330					335					
Cys	Glu	Ser	Asn	Gly	Asp	Lys	Gly	Ser	Gly	Gly	Ile	Lys	Gly	Gly	Phe
340					345					350					
Val	His	Gln	Arg	Met	Ala	Ser	Lys	Ile	Gly	Arg	Trp	Tyr	Ser	Arg	Thr
355					360					365					
Met	Ser	Lys	Thr	Lys	Arg	Met	Gly	Met	Gly	Leu	Tyr	Val	Lys	Tyr	Asp
370					375					380					
Gly	Asp	Pro	Trp	Ala	Asp	Ser	Asp	Ala	Leu	Ala	Leu	Ser	Gly	Val	Met
385					390					395					400
Val	Ser	Met	Glu	Glu	Pro	Gly	Trp	Tyr	Ser	Phe	Gly	Phe	Glu	Ile	Lys
405					410					415					
Asp	Lys	Lys	Cys	Asp	Val	Pro	Cys	Ile	Gly	Ile	Glu	Met	Val	His	Asp
420					425					430					
Gly	Gly	Lys	Glu	Thr	Trp	His	Ser	Ala	Ala	Thr	Ala	Ile	Tyr	Cys	Leu
435					440					445					
Met	Gly	Ser	Gly	Gln	Leu	Leu	Trp	Asp	Thr	Val	Thr	Gly	Val	Asp	Met

-continued

450 455 460

Ala Leu
465

<210> SEQ ID NO 67
<211> LENGTH: 584
<212> TYPE: PRT
<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 67

Met Lys Ala Ile Ile Val Leu Leu Met Val Val Thr Ser Asn Ala Asp
1 5 10 15

Arg Ile Cys Thr Gly Ile Thr Ser Ser Asn Ser Pro His Val Val Lys
20 25 30

Thr Ala Thr Gln Gly Glu Val Asn Val Thr Gly Val Ile Pro Leu Thr
35 40 45

Thr Thr Pro Thr Lys Ser Tyr Phe Ala Asn Leu Lys Gly Thr Arg Thr
50 55 60

Arg Gly Lys Leu Cys Pro Asp Cys Leu Asn Cys Thr Asp Leu Asp Val
65 70 75 80

Ala Leu Gly Arg Pro Met Cys Val Gly Thr Thr Pro Ser Ala Lys Ala
85 90 95

Ser Ile Leu His Glu Val Arg Pro Val Thr Ser Gly Cys Phe Pro Ile
100 105 110

Met His Asp Arg Thr Lys Ile Arg Gln Leu Pro Asn Leu Leu Arg Gly
115 120 125

Tyr Glu Asn Ile Arg Leu Ser Thr Gln Asn Val Ile Asp Ala Glu Asn
130 135 140

Ala Pro Gly Gly Pro Tyr Arg Leu Gly Thr Ser Gly Ser Cys Pro Asn
145 150 155 160

Ala Thr Ser Lys Ser Gly Phe Phe Ala Thr Met Ala Trp Ala Val Pro
165 170 175

Lys Asp Asn Asn Lys Asn Ala Thr Asn Pro Leu Thr Val Glu Val Pro
180 185 190

Tyr Val Cys Thr Glu Gly Glu Asp Gln Ile Thr Val Trp Gly Phe His
195 200 205

Ser Asp Asn Lys Thr Pro Met Lys Asn Leu Tyr Gly Asp Ser Asn Pro
210 215 220

Gln Lys Phe Thr Ser Ser Ala Asn Gly Val Thr Thr His Tyr Val Ser
225 230 235 240

Gln Ile Gly Gly Phe Pro Ala Gln Thr Glu Asp Glu Gly Leu Pro Gln
245 250 255

Ser Gly Arg Ile Val Val Asp Tyr Met Val Gln Lys Pro Arg Lys Thr
260 265 270

Gly Thr Ile Val Tyr Gln Arg Gly Val Leu Leu Pro Gln Lys Val Trp
275 280 285

Cys Ala Ser Gly Arg Ser Lys Val Ile Lys Gly Ser Leu Pro Leu Ile
290 295 300

Gly Glu Ala Asp Cys Leu His Glu Lys Tyr Gly Gly Leu Asn Lys Ser
305 310 315 320

Lys Pro Tyr Tyr Thr Gly Glu His Ala Lys Ala Ile Gly Asn Cys Pro
325 330 335

```

<210> SEQ ID NO 68
<211> LENGTH: 466
<212> TYPE: PRT
<213> ORGANISM: Influenza B virus

<400> SEQUENCE: 68

Met Leu Pro Ser Thr Ile Gln Thr Leu Thr Leu Phe Leu Thr Ser Gly
1          5              10          15

Gly Val Leu Leu Ser Leu Tyr Val Ser Ala Ser Leu Ser Tyr Leu Leu
20          25          30

Tyr Ser Asp Ile Leu Leu Lys Phe Ser Thr Thr Glu Ile Thr Ala Pro
35          40          45

Thr Met Pro Leu Asp Cys Ala Asn Ala Ser Asn Val Gln Ala Val Asn
50          55          60

Arg Ser Ala Thr Lys Gly Val Thr Leu Leu Leu Pro Glu Pro Glu Trp
65          70          75          80

Thr Tyr Pro Arg Leu Ser Cys Pro Gly Ser Thr Phe Gln Lys Ala Leu
85          90          95

Leu Ile Ser Pro His Arg Phe Gly Glu Thr Lys Gly Asn Ser Ala Pro
100         105         110

```

-continued

Leu Ile Ile Arg Glu	Pro Phe Ile Ala Cys	Gly	Pro Lys Glu Cys Lys
115	120	125	
His Phe Ala Leu Thr	His Tyr Ala Ala Gln	Pro Gly Gly Tyr Tyr Asn	
130	135	140	
Gly Thr Lys Glu Asp	Arg Asn Lys Leu Arg	His Leu Ile Ser Val Lys	
145	150	155	160
Leu Gly Lys Ile Pro	Thr Val Glu Asn Ser	Ile Phe His Met Ala Ala	
165	170	175	
Trp Ser Gly Ser Ala	Cys His Asp Gly Lys	Glu Trp Thr Tyr Ile Gly	
180	185	190	
Val Asp Gly Pro Asp	Ser Asn Ala Leu Leu	Lys Ile Lys Tyr Gly Glu	
195	200	205	
Ala Tyr Thr Asp Thr	Tyr His Ser Tyr Ala	Asn Asn Ile Leu Arg Thr	
210	215	220	
Gln Glu Ser Ala Cys	Asn Cys Ile Gly Gly	Asn Cys Tyr Leu Met Ile	
225	230	235	240
Thr Asp Gly Ser Ala	Ser Gly Ile Ser Glu	Cys Arg Phe Leu Lys Ile	
245	250	255	
Arg Glu Gly Arg Ile	Ile Lys Glu Ile Phe	Pro Thr Gly Arg Val Lys	
260	265	270	
His Thr Glu Glu Cys	Thr Cys Gly Phe Ala	Ser Asn Lys Thr Ile Glu	
275	280	285	
Cys Ala Cys Arg Asp	Asn Ser Tyr Thr Ala	Lys Arg Pro Phe Val Lys	
290	295	300	
Leu Asn Val Glu Thr	Asp Thr Ala Glu Ile	Arg Leu Met Cys Thr Glu	
305	310	315	320
Thr Tyr Leu Asp Thr	Pro Arg Pro Asp Asp	Gly Ser Ile Thr Gly Pro	
325	330	335	
Cys Glu Ser Asn Gly	Asn Lys Gly Ser Gly	Gly Ile Lys Gly Gly Phe	
340	345	350	
Val His Gln Arg Met	Ala Ser Lys Ile Gly	Arg Trp Tyr Ser Arg Thr	
355	360	365	
Met Ser Lys Thr Lys	Arg Met Gly Met Gly	Leu Tyr Val Lys Tyr Asp	
370	375	380	
Gly Asp Pro Trp Thr	Asp Ser Asp Ala Leu	Ala Leu Ser Gly Val Met	
385	390	395	400
Val Ser Met Glu Glu	Pro Gly Trp Tyr Ser	Phe Gly Phe Glu Ile Lys	
405	410	415	
Asp Lys Lys Cys Asp	Val Pro Cys Ile Gly	Ile Glu Met Val His Asp	
420	425	430	
Gly Gly Lys Glu Thr	Trp His Ser Ala Ala	Thr Ala Ile Tyr Cys Leu	
435	440	445	
Met Gly Ser Gly Gln	Leu Leu Trp Asp Thr	Val Thr Gly Val Asp Met	
450	455	460	
Ala Leu			
465			

What is claimed is:

1. An isolated polypeptide comprising any one of the amino acid sequences SEQ ID NO:35-52 and 54-68.

2. An immunogenic composition comprising an immunologically effective amount of the polypeptide of claim 1.

3. An isolated polynucleotide encoding the polypeptide of claim 1.

4. The polynucleotide of claim 3, wherein the polynucleotide is DNA.

5. The polynucleotide of claim 3, wherein the polynucleotide is RNA.

6. An immunogenic composition comprising an immunologically effective amount of the polynucleotide of claim 3.

7. A reassortant influenza virus comprising the polynucleotide of claim 3.

8. The virus of claim 7, wherein the virus comprises 6 internal genome segments from one or more donor viruses.

9. The virus of claim 8, wherein one donor virus is A/Ann Arbor/6/60, or A/Puerto Rico/8/34.

10. An immunogenic composition comprising an immunologically effective amount of the recombinant influenza virus of claim 8.

11. A vector comprising the polynucleotide of claim 3.

12. The vector of claim 11, wherein the vector is a plasmid, a cosmid, a phage, or a virus.

13. The vector of claim 11, wherein the vector is an expression vector.

14. An isolated cell comprising the vector of claim 11.

15. The virus of claim 8, wherein the 6 internal genome segments of the one or more donor viruses are selected for comprising one or more phenotypic attributes selected from the group consisting of: attenuated, cold adapted and temperature sensitive.

16. A method for producing the reassortant influenza virus of claim 7 in cell culture, the method comprising:

- i) introducing a plurality of vectors into a population of host cells capable of supporting replication of influenza viruses, which plurality of vectors comprises nucleotide sequences corresponding to at least 6 internal genome segments of a first influenza strain; and one genome segment encoding a polypeptide comprising any one of the amino acid sequences SEQ ID NO:35-52 and 54-68;
- ii) culturing the population of host cells; and,
- iii) recovering the influenza virus.

17. The method of claim 16, wherein the at least 6 internal genome segments of the first influenza virus strain are selected for comprising one or more phenotypic attributes selected from the group consisting of: attenuated, cold adapted and temperature sensitive.

18. The influenza virus produced by the method of claim 16, wherein the influenza virus is suitable for administration in an intranasal vaccine formulation.

19. The method of claim 16, wherein the first influenza strain is an influenza A strain.

20. The method of claim 16, wherein the first influenza strain is A/Ann Arbor/6/60, or A/Puerto Rico/8/34.

21. The method of claim 16, wherein the plurality of vectors is a plurality of plasmid vectors.

22. The method of claim 16, wherein the population of host cells comprises one or more of: Vero cells, PerC6 cells, MDCK cells, 293T cells, or COS cells.

23. The method of claim 16, wherein the method does not comprise use of a helper virus.

24. The method of claim 16, wherein the plurality of vectors consists of eight vectors.

25. An immunogenic composition comprising the polypeptide of claim 1.

26. The composition of claim 25, further comprising an excipient.

27. The composition of claim 26, wherein the excipient is a pharmaceutically acceptable excipient.

28. An immunogenic composition comprising the polynucleotide of claim 3.

29. The composition of claim 28, further comprising an excipient.

30. The composition of claim 29, wherein the excipient is a pharmaceutically acceptable excipient.

31. An immunogenic composition comprising the reassortant virus of claim 7.

32. The composition of claim 31 wherein the reassortant virus is a 6:2 reassortment virus comprising 6 internal genome segments from one or more donor viruses.

33. The composition of claim 32, wherein the 6 internal genome segments of the one donor virus are selected for comprising one or more phenotypic attributes selected from the group consisting of: attenuated, cold adapted and temperature sensitive.

34. The composition of claim 33, wherein one donor virus is A/Ann Arbor/6/60, or A/Puerto Rico/8/34.

35. A live attenuated influenza vaccine comprising the composition of claim 31.

36. The composition of claim 31, further comprising one or more pharmaceutically acceptable excipient.

37. A method of prophylactic or therapeutic treatment of a viral infection in a subject, the method comprising: administering to the subject the virus of claim 7 in an amount effective to produce an immunogenic response against the viral infection.

38. The method of claim 37, wherein the subject is a mammal.

39. The method of claim 38, wherein the mammal is a human.

40. The method of claim 37, wherein the virus is formulated using at least one pharmaceutically acceptable excipient.

* * * * *