




(86) Date de dépôt PCT/PCT Filing Date: 2019/03/22
(87) Date publication PCT/PCT Publication Date: 2019/09/26
(45) Date de délivrance/Issue Date: 2024/05/14
(85) Entrée phase nationale/National Entry: 2020/09/18
(86) N° demande PCT/PCT Application No.: CA 2019/050353
(87) N° publication PCT/PCT Publication No.: 2019/178699
(30) Priorité/Priority: 2018/03/23 (US62/647,151)

(51) Cl.Int./Int.Cl. *A61K 39/385* (2006.01),
A61K 41/00 (2020.01), *A61P 31/00* (2006.01),
A61P 35/00 (2006.01), *A61P 37/04* (2006.01),
C07C 323/41 (2006.01), *C07D 207/40* (2006.01),
C07D 211/88 (2006.01), *C07H 15/04* (2006.01),
C07H 15/10 (2006.01)
(72) Inventeurs/Inventors:
SHIAO, TZE CHIEH, CA;
ROY, RENE, CA
(73) Propriétaire/Owner:
KORANEX CAPITAL, CA
(74) Agent: ROBIC AGENCE PI S.E.C./ROBIC IP AGENCY
LP

(54) Titre : GLYCOCONJUGUES DE PRECISION EN TANT QU'OUTILS THERAPEUTIQUES
(54) Title: PRECISION GLYCOCONJUGATES AS THERAPEUTIC TOOLS

Photolytic thiol-ene "click chemistry"



R₁ = Cysteine residues on T-cell epitope (ex. Tetanus toxoid)

R₂ = Carbohydrate antigens serving as B-cell epitope

(57) Abrégé/Abstract:

The present description relates to glycoconjugates, glycoconjugate immunogens and glycoconjugate vaccines comprising carbohydrate antigens coupled to immunogenic carrier proteins, or materials used for detection and screening of resulting antibodies. Improved methods of more directly and precisely conjugating carbohydrate antigens to free thiol groups of immunogenic carrier proteins are described, including "click-chemistry" approaches based on photocatalytic thiol-ene reactions.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

26 September 2019 (26.09.2019)



(10) International Publication Number

WO 2019/178699 A1

(51) International Patent Classification:

A61K 39/385 (2006.01) C07C 323/41 (2006.01)
 A61K 41/00 (2006.01) C07D 207/40 (2006.01)
 A61P 31/00 (2006.01) C07D 211/88 (2006.01)
 A61P 35/00 (2006.01) C07H 15/04 (2006.01)
 A61P 37/04 (2006.01) C07H 15/10 (2006.01)

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
 TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
 KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

(21) International Application Number:

PCT/CA2019/050353

(22) International Filing Date:

22 March 2019 (22.03.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/647,151 23 March 2018 (23.03.2018) US

(71) Applicant: **KORANEX CAPITAL** [CA/CA]; 424, rue
 Guy, Bureau 202, Montreal, Québec H3J 1S6 (CA).

(72) Inventors: **SHIAO, Tze Chieh**; 4145 rue de Chambly,
 Montreal, Québec H1X 3L1 (CA). **ROY, René**; 325 Place
 Lussac, Terrebonne, Québec J6Y 1E6 (CA).

(74) Agent: **ROBIC LLP**; 630 René-Lévesque W Blvd, 20th
 Floor, Montréal, Québec H3B 1S6 (CA).


(81) Designated States (unless otherwise indicated, for every
 kind of national protection available): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
 CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
 DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
 HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
 KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
 OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
 SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
 kind of regional protection available): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
 UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
 TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: PRECISION GLYCOCONJUGATES AS THERAPEUTIC TOOLS

Fig. 2

Photolytic thiol-ene "click chemistry"

R₁ = Cysteine residues on T-cell epitope (ex. Tetanus toxoid)R₂ = Carbohydrate antigens serving as B-cell epitope

(57) Abstract: The present description relates to glycoconjugates, glycoconjugate immunogens and glycoconjugate vaccines comprising carbohydrate antigens coupled to immunogenic carrier proteins, or materials used for detection and screening of resulting antibodies. Improved methods of more directly and precisely conjugating carbohydrate antigens to free thiol groups of immunogenic carrier proteins are described, including "click-chemistry" approaches based on photocatalytic thiol-ene reactions.



WO 2019/178699 A1

PRECISION GLYCOCONJUGATES AS THERAPEUTIC TOOLS

The present description relates to glycoconjugate therapeutic tools. More specifically, the present description relates to carbohydrate antigens non-randomly coupled to free thiol groups of immunogenic and antigenic carrier peptides and proteins, and improved methods of producing same using, for example, photocatalytic thiol-ene “click chemistry” reactions. Applications as antigens, immunogens, vaccines, and in diagnostics are also described.

BACKGROUND

The ultimate objective of immunotherapy is to trigger the innate and adaptive responses of the immune system in a way similar to that produced during an infection or tumor progression. The principal interfaces between the innate and adaptive immune responses are the antigen-presenting cells (APCs), and particularly dendritic cells (DCs). APCs are able to recognize microorganisms through pattern recognition receptors (PRRs) such as Toll-like receptors (TLRs). On recognition of microbial surface determinants or aberrant and unnatural antigens, APCs undergo maturation and activation leading to a redistribution of MHC molecules from intracellular compartments to the cell surface, secretion of cytokines and chemokines. The microorganisms or tumors and their related antigenic markers can be engulfed by the APC through an endocytic pathway where it is typically degraded. The peptides and covalently linked antigens released by protein processing are then displayed on MHC class II molecules and are recognized by CD4⁺ T cells which in turn undergo functional maturation into different subsets, such as Th1 or Th2 cells, upon co-stimulatory signals received from the APC. Th1 cells lead to a predominantly pro-inflammatory response with the secretion of IFN- γ and TNF- α , whereas Th2 cells secrete typical cytokines. Albeit Th1 cells are mainly associated with a cell-mediated response, both types of Th cells support the production of antibodies by B cells, which in turn influences antibody isotype and function. For example, IL-12 and TNF- α are associated with the differentiation of Th1 cells and production of type 1 IgG subclasses, whereas IL-6 and other Th2 cytokines contribute to the type 2 IgG subclass (IgG1) production. It is thus desirable to be able to tailor vaccine-induced immunity to an appropriate response to deal with a pathogen or tumor antigen of interest.

Carbohydrates, as opposed to proteins and peptides, are T cell independent antigens not properly equipped to trigger the participation of Th cells and hence, cannot induce immune cell proliferation, antibody class switching, and affinity/specificity maturation. The major early advances initially encountered with carbohydrate-based vaccines have been supported by the discovery that, when properly conjugated to protein carriers, serving as T cell dependent epitopes, bacterial capsular polysaccharides

became capable of acquiring the requisite immunochemical ability to produce opsonophagocytic antibodies.

Traditionally, strategies for conjugating carbohydrate antigens to carrier proteins have relied on either reductive amination of aldehyde-derived sugars onto the ϵ -amino groups of the lysine residues, or simply amide coupling reactions. In both cases, partial and random carbohydrate antigen conjugation generally occurs. Furthermore, if all amide partners (amines from lysine or acid from glutamic/aspartic acids) are used for carbohydrate conjugation, far too many carbohydrate antigens become attached to the carrier proteins, thus resulting in masking potentially essential T cell peptide epitopes with the inherent diminution/elimination of immunogenicity. Thus, current strategies for preparing glycoconjugate vaccines are inadequate and face significant regulatory and/or commercial obstacles, since the preparations lack the necessary homogeneity in terms of their carbohydrate distribution and reproducibility (i.e., the attachment points of the sugars onto the proteins are randomly distributed and in various densities from batch to batch). Thus, glycoconjugate vaccines having greater carbohydrate antigen homogeneity, more precisely characterizable structures, and reproducibility from batch to batch would be highly desirable.

SUMMARY

The present description relates to glycoconjugate immunogens comprising carbohydrate antigens directly coupled to immunogenic carrier proteins at precise (non-random) positions. More particularly, the carrier proteins comprise one or more free thiol groups (e.g., corresponding to the side chains of cysteine residues) and the carbohydrate antigens are conjugated to the carrier proteins at one or more of these free thiol groups. The present description also relates to improved methods for synthesizing glycoconjugate immunogens/vaccines involving directly conjugating carbohydrate antigens to free thiol groups of carrier proteins, for example using "click-chemistry" approaches (e.g., photocatalytic thiol-ene reactions). The improved conjugation methods described herein may be performed under conditions sufficiently mild (e.g., use of only water-soluble reagents, the absence of organic solvents, or use concentrations of organic solvents sufficiently low (e.g., < 5%) to avoid carrier protein denaturation, and/or at relatively neutral pH) to avoid destroying the activity, antigenicity, and/or structure (e.g., cleavage of native disulfide bridges and/or denaturation) of a carrier protein, without affecting the specificity of the conjugation. Furthermore, photocatalytic thiol-ene reactions described herein may be performed under ultraviolet light in the presence of a catalyst (e.g., 355 nm or 365 nm), or under short-wave ultraviolet light (e.g., at 254 nm) in the absence of a catalyst, further simplifying the process.

In some aspects, the present description relates to a method for producing a glycoconjugate immunogen, the method comprising: (a) providing a carbohydrate antigen covalently linked to a terminal alkene (alkenyl carbohydrate antigen), the terminal alkene being directly conjugatable to a thiol group via

a thiol-ene reaction; (b) providing a carrier protein having one or more free thiol groups; and (c) performing a photocatalytic thiol-ene reaction to directly conjugate the carbohydrate antigen to the carrier protein at the one or more free thiol groups, thereby producing the glycoconjugate immunogen; wherein the carrier protein is immunogenic when administered to a subject, and wherein conjugation of the carbohydrate antigen to the carrier protein increases the immunogenicity of the carbohydrate antigen upon administration to the subject, as compared to administration of the unconjugated carbohydrate antigen.

In some embodiments, the photocatalytic thiol-ene reaction is performed under reaction conditions that avoid carrier protein denaturation. In embodiments, the alkenyl carbohydrate antigen is water-soluble and the photocatalytic thiol-ene reaction is performed under reaction conditions that retain the carrier protein's activity, antigenicity, and/or structure (e.g., in the absence of any organic solvent, or in the presence of an organic solvent at a concentration sufficiently low (e.g., < 5%) to avoid carrier protein denaturation). The photocatalytic thiol-ene reaction may be performed in the presence of a catalyst (e.g., a water-soluble or water-insoluble catalyst) under irradiation under short-wave (e.g., 254 nm) and/or long-wave (e.g., 355 or 365 nm) ultraviolet light.

In some embodiments, the photocatalytic thiol-ene reaction comprises reacting between 1 to 200 molar equivalents of the alkenyl carbohydrate antigen per free thiol group of the carrier protein; and/or the photocatalytic thiol-ene reaction is performed at a pH between about 3 and 10, 3.5 and 9.5, 4 and 9, 4.5 and 8.5, 5 and 8, 5.5 and 8, 6 and 8, or 6.5 and 7.5.

In some embodiments, the carbohydrate antigen, following conjugation to the carrier protein, is not cleavable from the carrier protein by an endogenous enzyme of the subject. In some embodiments, the alkenyl carbohydrate antigen is covalently linked to the terminal alkene, and/or the carbohydrate antigen is conjugated to the carrier protein, via a glycosidic bond, such as is an O-glycosidic bond, an S-glycosidic bond, an N-glycosidic bond, or a C-glycosidic bond, or a bond obtained by reductive amination, such as between an allyl amine and a reducing sugar (including bacterial CPS). In some embodiments, the carbohydrate antigen comprises a B cell epitope, and/or induces a humoral immune response in the subject. In some embodiments, the carbohydrate antigen comprises a T cell epitope, and/or induces a cell-mediated immune response in the subject. In some embodiments, the carbohydrate antigen comprises both a B cell epitope and a T cell epitope, and/or induces both a humoral and a cell-mediated immune response in the subject. In some embodiments, the carbohydrate antigen is or comprises a tumor associated carbohydrate antigen (TACA), such as Tn, S-Tn, *Thomsen-Friedenreich* (TF), (2,3)-S-TF, (2,6)-S-TF, Globo H, GD2, GD3, GM2, GM3, N-glycolyl-GM3, Lea, sLea, Lex, sLex, or any combination thereof. In some embodiments, the photocatalytic thiol-ene reaction conjugates at least two of the same carbohydrate antigen or more than one type of carbohydrate antigen to the carrier protein, thereby producing a multi-valent glycoconjugate immunogen (e.g., comprising 2, 3, 4, 5, 6, 7, 8, 9, 10,

11, 12, 13, 14, 15, or more of the same or different types of carbohydrate antigens conjugated to the carrier protein). In some embodiments, the carbohydrate antigen is or comprises a viral polysaccharide antigen, or a bacterial capsular polysaccharide (CPS) (e.g., a Pneumococcal and/or Streptococcal polysaccharide serotype, meningococcal CPS; influenza (such as influenza type a or b) CPS).

5 In some embodiments, the alkenyl carbohydrate antigen is linked to the terminal alkene via a linker (e.g., a linker as described herein).

In some embodiments, the carrier protein comprises one or more cysteine residues having the one or more free thiol groups. In some embodiments, the carrier protein comprises a T cell epitope, and/or induces a cell-mediated immune response in the subject.

10 In some embodiments, the carrier protein comprises a B cell epitope, and/or induces a humoral immune response in the subject. In some embodiments, the carrier protein comprises both a B cell epitope and a T cell epitope, and/or induces both a humoral and a cell-mediated immune response in the subject. In some embodiments, the carrier protein is, is from, or comprises: Tetanus Toxoid (TT), Diphtheria Toxoid (DT), cross-reacting material 197 (CRM197), Meningococcal Outer Membrane Protein Complex (OMPC), H. Influenzae Protein D (HiD), a virus-like particle (VLP), a cytokine, an immunogenic peptide
15 such as Tetanus Toxin 831-844 (SEQ ID NO: 1 or 2), albumin (such as bovine serum albumin or human serum albumin), or an immunogenic fragment thereof. In some embodiments, the carrier protein is a protein having one or more disulfide bridges, and wherein: (i) the one or more disulfide bridges remain unaffected following said photocatalytic thiol-ene reaction; or (ii) the carrier protein is pre-treated with a
20 reducing agent to expose one or more additional free thiol groups for conjugation to the carbohydrate antigen.

In some embodiments, the total number of carbohydrate antigens comprised in the glycoconjugate immunogen is equal to the number of free thiol groups available on the carrier protein prior to conjugation. In some embodiments, the glycoconjugate immunogen induces a cell-mediated
25 immune response to the carbohydrate antigen upon administration to the subject.

In some embodiments, the methods described herein further comprise: (d) purifying the glycoconjugate immunogen.


In some aspects, the present description relates to a method for producing a glycoconjugate vaccine, the method comprising formulating the glycoconjugate immunogen prepared by a method
30 described herein with a pharmaceutically acceptable excipient, and/or an adjuvant (e.g., an inorganic compound, a mineral oil, a microbial derivative, a plant derivative, a cytokine, squalene, alum, aluminum hydroxide, aluminum phosphate, calcium phosphate hydroxide, a toll-like receptor agonist, an immunostimulatory polynucleotide (e.g., CPG), an immunostimulatory lipid, Freund's adjuvant, RIBI's adjuvant, QS-21, muramyl dipeptide, or any combination thereof).

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more carbohydrate antigens and an immunogenic carrier protein having one or more solvent-accessible cysteine residues, wherein the one or more carbohydrate antigens are linked to the immunogenic carrier protein at the one or more solvent-accessible cysteine residues, and wherein
5 conjugation of the one or more carbohydrate antigens to the immunogenic carrier protein increases the immunogenicity of the one or more carbohydrate antigens upon administration to a subject, as compared to administration of the unconjugated carbohydrate antigen. In some embodiments, the one or more carbohydrate antigens are linked to the one or more solvent-accessible cysteine residues via a linker as described herein and the one or more carbohydrate antigens is/are as described herein. In some
10 embodiments, the synthetic glycoconjugate immunogen is a multi-valent glycoconjugate immunogen as described herein, and uses a carrier protein as described herein. In some embodiments, the total number of carbohydrate antigens comprised in the glycoconjugate immunogen is equal to the number of solvent-accessible cysteine residues on the carrier protein. In some embodiments, the synthetic glycoconjugate immunogen induces a cell-mediated immune response to the carbohydrate antigen upon administration to
15 the subject. In some embodiments, the synthetic glycoconjugate immunogen is prepared by a method described herein.

In some aspects, the present description relates to a glycoconjugate vaccine produced by a method as described herein, and/or comprising the synthetic glycoconjugate immunogen as described herein, and a pharmaceutically acceptable excipient, and/or an adjuvant. In some embodiments, the
20 glycoconjugate vaccine is a prophylactic vaccine or a therapeutic vaccine.


In some aspects, the present description relates to a method of immunizing, vaccinating, or treating a subject comprising administering to the subject the glycoconjugate immunogen produced by a method as described herein, the glycoconjugate vaccine produced by a method as described herein, a synthetic glycoconjugate immunogen as described herein, or a glycoconjugate vaccine as described
25 herein. In some embodiments, the present description relates to a synthetic glycoconjugate immunogen or a glycoconjugate vaccine as described herein, for use in immunizing, vaccinating, or treating a subject having a disease, or for detecting the presence of an antibody that specifically binds to the glycoconjugate, or for detecting said immunization, vaccination, or treatment (e.g., in a biological sample from a subject).

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein the synthetic glycoconjugate immunogen has the structure:



wherein

- y is at least 1;
- **SA** is selected from the group consisting of sugar antigens or immunogenic portions thereof, and when y is more than 1, **SA** are identical or different;
- $[\text{S}]_z\text{PC}$ is a carrier protein having one or more sulfur atoms originating from the one or more free thiol groups;
- z is at least 1 and is at least equal to y ; and
- **L** is a linker selected from the group consisting of linkers having the structure:




wherein:

- **X** is O, S, NR_1 , or CH_2 ;
- R_1 is $-\text{H}$, $-\text{COH}$, $-\text{COCH}_3$, or $-\text{COEt}$;
- n is 1, 2, 3, 4, or 5; and
- R_2 is H or Me; and

when y is more than 1, **L** are identical or different;

or a stereoisomer thereof.


In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein the synthetic glycoconjugate immunogen has the structure:



wherein

- y is at least 1;
- **SA** is selected from the group consisting of sugar antigens or immunogenic portions thereof, and when y is more than 1, **SA** are identical or different;

- **[S]_z-PC** is a carrier protein having one or more sulfur atoms originating from the one or more free thiol groups;
- **z** is at least 1 and is at least equal to **y**; and
- **L** is a linker selected from the group consisting of linkers having the structure:




wherein:

- **X** is S, NR₁, CH₂ or O;
- **R₁** is -H, -COH, -COMe, or -COEt;
- **n** is 1, 2, 3, 4, or 5;
- **R₂** is H or Me;
- **q** is 1, 2, 3, 4, or 5;
- **R₃** and **R₄** are each a hydrogen atom and **m** is 1, 2, 3, 4 or 5, or **R₃** and **R₄** form together a radical -CO-CH₂- or a radical -CO-CH₂-CH₂- with the carbonyl linked to the nitrogen atom, and **m** is 1; and
- when **y** is more than 1, **L** are identical or different;


or a stereoisomer thereof.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein the synthetic glycoconjugate immunogen has the structure:




wherein

- **y** is at least 1;
- **SA** is selected from the group consisting of sugar antigens or immunogenic portions thereof, and when **y** is more than 1, **SA** are identical or different;
- **[S]_z-PC** is a carrier protein having one or more sulfur atoms originating from the one or more free thiol groups;
- **z** is at least 1 and is at least equal to **y**; and
- **L** is a linker selected from the group consisting of linkers having the structure:



wherein

- **X** is S, NR₁, CH₂ or O;
- **R₁** is -H, -COH, -COMe, or -COEt;
- **n** is 1, 2, 3, 4, or 5;
- **R₂** is H or Me;
- **q** is 1, 2, 3, 4, or 5;
- **r** is 1, 2, 3, 4 or 5;
- **R₅** is S-PC, a covalent bond, or a radical of structure:




wherein **R₃** and **R₄** are each a hydrogen atom and **m** is 1, 2, 3, 4 or 5, or **R₃** and **R₄** form together a radical -CO-CH₂- or a radical -CO-CH₂-CH₂- with the carbonyl linked to the nitrogen atom, and **m** is 1; and

when **y** is more than 1, **L** are identical or different;

or a stereoisomer thereof.

In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as defined above, wherein the linker has the structure:




wherein

- **X** is S, NR₁, CH₂ or O;
- **R₁** is -H, -COH, -COMe, or -COEt;
- **n** is 1, 2, 3, 4, or 5;
- **R₂** is H or Me;
- **q** is 1, 2, 3, 4, or 5; and

- *r* is 1, 2, 3, 4 or 5.

In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as defined above, wherein the linker has the structure:



5 wherein

- *X* is S, NR₁, CH₂ or O;

- *R*₁ is -H, -COH, -COMe, or -COEt;


- *n* is 1, 2, 3, 4, or 5;

- *R*₂ is H or Me;

10 - *q* is 1, 2, 3, 4, or 5; and

- *r* is 1 or 2.

In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as defined above, wherein the linker has the structure:



15 wherein

- *X* is S, NR₁, CH₂ or O;

- *R*₁ is -H, -COH, -COMe, or -COEt;

- *n* is 1, 2, 3, 4, or 5;

- *R*₂ is H or Me;


20 - *q* is 1, 2, 3, 4, or 5; and

- *r* is 1 or 2.

In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as designed above, wherein *y* is an integer varying from 1 to 50, 1 to 40, 1 to 30, 1 to 20, 1 to 10, 1 to 5, or 1 to 3.


25

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein each linker has the structure:



- 5 wherein: **SA** is a sugar antigen or a portion thereof; **S-PC** is a carrier protein; **X** is O, S, NR₁, or CH₂; **R₁** is -H, -COH, -COCH₃, or -COEt; **n** is 1, 2, 3, 4, or 5; and **R₂** is H or Me; or a stereoisomer thereof.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein each linker has the structure:



- 10 wherein: **SA** is a sugar antigen or a portion thereof; **S-PC** is a carrier protein; **X** is S, NR₁, CH₂ or O; **R₁** is -H, -COH, -COMe, or -COEt; **n** is 1, 2, 3, 4, or 5; **R₂** is H or Me; **q** is 1, 2, 3, 4, or 5; **R₃** and **R₄** are each a hydrogen atom and **m** is 1, 2, 3, 4 or 5, or **R₃** and **R₄** form together a radical -CO-CH₂- or a radical -CO-CH₂-CH₂- with the carbonyl linked to the nitrogen atom, and **m** is 1; or a stereoisomer thereof.

- 15 In some embodiments, the sugar antigen is a carbohydrate antigen as described herein, the carrier protein is as described herein, the synthetic glycoconjugate immunogen is a multi-valent glycoconjugate immunogen as described herein.

- 20 In some aspects, the present description relates to a vaccine comprising the synthetic glycoconjugate immunogen as described herein, and a pharmaceutically acceptable excipient and/or an adjuvant as described herein.

In some aspects, the present description relates to the use of the glycoconjugate immunogen prepared by a method as described herein, or a synthetic glycoconjugate immunogen as described herein, for the manufacture of a vaccine.

- 25 In some aspects, the present description relates to the use of the glycoconjugate immunogen prepared by a method as described herein, a glycoconjugate vaccine produced by a method as described herein, a synthetic glycoconjugate immunogen as described herein, or a vaccine as described herein, for the treatment of a subject having a disease associated with increased expression of said one or more carbohydrate or sugar antigens.

General Definitions

Headings, and other identifiers, e.g., (a), (b), (i), (ii), etc., are presented merely for ease of reading the specification and claims. The use of headings or other identifiers in the specification or claims does not necessarily require the steps or elements be performed in alphabetical or numerical order or the order in which they are presented.

The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one” but it is also consistent with the meaning of “one or more”, “at least one”, and “one or more than one”.

The term “about” is used to indicate that a value includes the standard deviation of error for the device or method being employed in order to determine the value. In general, the terminology “about” is meant to designate a possible variation of up to 10%. Therefore, a variation of 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10% of a value is included in the term “about”. Unless indicated otherwise, use of the term “about” before a range applies to both ends of the range.

As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

As used herein, the term “protein” (e.g., in the expression “carrier protein”) means any peptide-linked chain of amino acids, which may or may not comprise any type of modification (e.g., chemical or post-translational modifications such as acetylation, phosphorylation, glycosylation, sulfation, sumoylation, prenylation, ubiquitination, etc.), so long as the modifications do not destroy the immunogenicity of the glycoconjugate immunogens and glycoconjugate vaccines described herein. For further clarity, the terms “protein” and “carrier protein” as used herein encompass both peptides and polypeptides.

As used herein, the term “administration” may comprise administration routes such as parenteral (e.g., subcutaneously, intradermally, intramuscularly, or intravenously), oral, transdermal, intranasal, etc., so long as the route of administration results in the generation of an immune response in the subject.

As used herein, “subject” generally refers to a mammal, including primates, and particularly to a human.

Other objects, advantages and features of the present description will become more apparent upon reading of the following non-restrictive description of specific embodiments thereof, given by way of example only with reference to the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

In the appended drawings:

Fig. 1 shows examples of typical chemical structures of key tumor-associated carbohydrate antigens (TACAs) from human glycoproteins (MUCs).

5 **Fig. 2** shows a proposed “click chemistry” approach for conjugating a carbohydrate antigen (B cell epitope) directly to an available thiol group of an immunogenic carrier protein (e.g., T cell epitope such as Tetanus Toxoid).

10 **Fig. 3** shows an example of a “precision glycoconjugate” prepared by a photocatalytic thiol-ene reaction to conjugate Tn or TF antigens to six native “free” cysteine residues on the carrier protein Tetanus Toxoid.

Fig. 4 shows an example of a “precision glycoconjugate” prepared by using Tn or TF antigens linked to thiol-specific iodoacetamido or maleimido groups, which are then conjugated to “free” cysteine residues on a carrier protein via a reaction a pH below 8 (iodoacetamido or maleimido groups) or a thiol-ene reaction (maleimido group).

15 **Fig. 5** shows reaction schemes for the syntheses of allyl Tn antigen and allyl TF antigen ready for conjugation to carrier proteins, as described in Examples 2-6.

Fig. 6A-6D shows the ¹H-NMR (**Fig. 6A**) and ¹³C-NMR (**Fig. 6B**) spectra, as well as mass spectrometry results (**Fig. 6C and 6D**) for allyl 2-acetamido-3,6-di-O-pivaloyl-2-deoxy- α -D-glucopyranoside (Compound 2).

20 **Fig. 7A-7D** shows the ¹H-NMR (**Fig. 7A**) and ¹³C-NMR (**Fig. 7B**) spectra, as well as mass spectrometry results (**Fig. 7C and 7D**) for allyl 2-acetamido-2-deoxy- α -D-galactopyranoside (allyl Tn).

Fig. 8A-8C shows the ¹H-NMR (**Fig. 8A**) spectra, as well as mass spectrometry results (**Fig. 8B & 8C**) for allyl 2-acetamido-4,6-O-benzylidene-2-deoxy- α -D-galactopyranoside (Compound 4).

25 **Fig. 9A-9D** shows the ¹H-NMR (**Fig. 9A**) and ¹³C-NMR (**Fig. 9B**) spectra, as well as mass spectrometry results (**Fig. 9C and 9D**) for allyl (2,3,4,6-tetra-O-benzoyl- β -D-galactopyranosyl)-(1 $\rightarrow$ 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- α -D-galactopyranoside (Compound 6)

Fig. 10A-10D shows the ¹H-NMR (**Fig. 10A**) and ¹³C-NMR (**Fig. 10B**) spectra, as well as mass spectrometry results (**Fig. 10C and 10D**) for Compound 7.

30 **Fig. 11A-11D** shows the ¹H-NMR (**Fig. 11A**) and ¹³C-NMR (**Fig. 11B**) spectra, as well as mass spectrometry results (**Fig. 11C and 11D**) for allyl (β -D-galactopyranosyl)-(1 $\rightarrow$ 3)-2-acetamido-2-deoxy- α -D-galactopyranoside (allyl TF).

Fig. 12 shows the results of Dynamic Light Scattering (DLS) analyses of the TF-TT glycoconjugate alone (upper left panel), peanut lectin alone (upper right panel), or both combined (lower panel).

Fig. 13 shows the thiol concentration of the peptide dTT831-840 as a function of time in the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn and O-Allyl-TF.

Fig. 14 shows the fold reduction of thiols of the peptide dTT831-840 in the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn as a function of the peptide's cysteine position and the presence of AAPH or DMPA.

Fig. 15 shows the reactivity of lectins to peptide dTT831-840 conjugated to Tn by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn as a function of the position of the peptide's cysteine and the presence of AAPH or DMPA.

Fig. 16 shows the kinetics of thiol concentration of the peptide dTT831-840 in the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn at different wave lengths in the presence of AAPH or DMPA.

Fig. 17 shows the reactivity of the lectin *Vicia Villosa* to the peptide dTT831-840 conjugated to Tn by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn at different wave lengths in the presence of AAPH or DMPA.

Fig. 18 shows the kinetics of thiol concentration of the peptide dTT831-840 in the photocatalytic thiol-ene conjugation reaction with C-Allyl and O-Allyl sugars.

Fig. 19 shows the reactivity of lectins and anti-TF antibody to peptide dTT831-840 conjugated by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn.

Fig. 20 shows the reactivity of anti-TF antibody to reduced dTT or reduced dTT conjugated by the photocatalytic thiol-ene conjugation reaction with O-Allyl-TF.

Fig. 21 shows the molar ratio of thiols of dTT at various steps of the photocatalytic thiol-ene conjugation reaction and following reduction of the conjugated dTT.

Fig. 22 shows the reactivity to lectins and anti-TF mAb of dTT conjugated by the photocatalytic thiol-ene conjugation reaction to O-Allyl-Tn.

Fig. 23 shows the molar ratio of thiol of chemically thiolated dTT conjugated or not by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn in the presence of low or high amount of AAPH.

Fig. 24 shows the reactivity of the lectin *Vicia Villosa* to chemically thiolated dTT conjugated by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn in the presence of low or high amount of AAPH at two different times.

Fig. 25 shows the reactivity of lectins and anti-TF antibody to reduced BSA conjugated by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn.

Fig. 26 shows the relationship between the reducing conditions and the molar ratio of thiols of reduced BSA.

Fig. 27 shows the reactivity of the Tn-specific *Vicia Villosa* lectin and anti-TF antibody to BSA with various free thiol molar ratio and conjugated by the photocatalytic thiol-ene conjugation reaction with various amount of O-Allyl-Tn.

Fig. 28 shows the relationship between the reducing conditions, the molar ratio of thiols of reduced BSA and the TF reactivity of reduced BSA conjugated by the photocatalytic thiol-ene conjugation reaction with O-Allyl-TF.

Fig. 29 shows the molar ratio of thiols adducts in BSA in function of time by chemical thiolation alone or thiolation with simultaneous photocatalytic thiol-ene conjugation reaction in the presence or absence of AAPH and O-Allyl-Tn.

Fig. 30 shows the reactivity of the Tn-specific *Vicia Villosa* lectin with thiolated BSA or BSA simultaneously thiolated and conjugated by photocatalytic thiol-ene conjugation reaction in the presence or absence of AAPH and O-Allyl-Tn.

Fig. 31 shows the relationship between the thiol molar ratio and the reactivity of the Tn-specific *Vicia Villosa* lectin of BSA conjugated by the photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn in a function of time.

Fig. 32 shows the reactivity of the Tn-specific *Vicia Villosa* lectin to the native or thiolated BSA conjugated by photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn in the presence of low or high amount of AAPH and at two wave-lengths.

Fig. 33 shows the relationship between the galactose molar ratio and the reactivity of the anti-TF mAb to BSA at different steps of thiolation and photocatalytic thiol-ene conjugation reaction with O-Allyl-Tn at low or high amount of AAPH.

Fig. 34 shows the chemical structures of the dTT831-844-Cys- β Ala and the Cys-dTT831-844- β Ala with a cysteine residue at the C- and N-terminal, respectively together with their tabulated LC-MS data.

Fig. 35 shows the detailed LC-MS data of the peptide dTT831-844-Cys- β Ala.

Fig. 36 shows the detailed LC-MS data of the peptide Cys-dTT831-844- β Ala.

Fig. 37 shows the photolytic AAPH-catalyzed thiol-ene reaction of the O-Allyl Tn on the C-terminal peptide dTT831-844-Cys- β Ala.

Fig. 38 shows the LC-MS profile of the O-Allyl Tn on the C-terminal peptide

Fig. 39 shows the photolytic AAPH-catalyzed thiol-ene reaction of the O-Allyl Tn on the N-terminal peptide Cys-dTT831-844- β Ala.

Fig. 40 shows the LC-MS profile of the O-Allyl Tn on the N-terminal peptide.

SEQUENCE LISTING

This application contains a Sequence Listing in computer readable form created March 19, 2019 having a size of about 4 KB.

DETAILED DESCRIPTION

The present description relates to glycoconjugate immunogens (e.g., for use as vaccines, in diagnostics, or for generating diagnostic or therapeutic tools such as specific anti-glycoconjugate antibodies). The glycoconjugate immunogens described herein generally comprise carbohydrate antigens coupled to immunogenic carrier proteins. More particularly, the carrier proteins comprise one or more free thiol groups (e.g., corresponding to the side chains of cysteine residues) and the carbohydrate antigens are conjugated to the carrier proteins at one or more of these free thiol groups. The present description also relates to improved methods for synthesizing glycoconjugate immunogens/vaccines involving directly conjugating carbohydrate antigens to free thiol groups of carrier proteins, for example using “click-chemistry” approaches (e.g., photocatalytic thiol-ene reactions). The improved conjugation methods described herein may be performed under conditions sufficiently mild (e.g., use of only water-soluble reagents, the absence of organic solvents, use concentrations of organic solvents sufficiently low (e.g., < 5%) to avoid carrier protein denaturation, and/or at relatively neutral pH) to avoid destroying the activity, antigenicity, and/or structure (e.g., cleavage of native disulfide bridges and/or denaturation) of a carrier protein, without affecting the specificity of the conjugation. Furthermore, photocatalytic thiol-ene reactions described herein may be performed under ultraviolet light in the presence of a catalyst (e.g., 355 nm or 365 nm), or under short-wave ultraviolet light (e.g., at 254 nm) in the absence of a catalyst, further simplifying the process.

Because of the greater precision offered by the conjugation methods described herein, the glycoconjugate immunogens/vaccines described herein may have greater homogeneity (in terms of their carbohydrate distributions), reproducibility, and may be easier to characterize as compared to glycoconjugates that rely on random coupling to other amino acid side chains (e.g., amines from lysine, or acid from glutamate/aspartate residues). Such characteristics may facilitate regulatory approval and/or commercialization of glycoconjugate vaccines, both of which have historically proven difficult based on traditional approaches.

In some aspects, the present description relates to a method for producing a glycoconjugate immunogen (e.g., for administration to a subject). As used here, the term “subject” generally refers to a living being (e.g., animal or human) that is able to mount an immune response to a glycoconjugate immunogen as described herein, preferably leading to the production of antibodies that specifically bind to the glycoconjugate immunogen. In some embodiments, a subject described herein may be a patient to

be treated therapeutically (e.g., via vaccination with a glycoconjugate immunogen described herein) or may be employed as a means for generating tools (e.g., antibodies) for research, diagnostic, and/or therapeutic purposes. The method for producing a glycoconjugate immunogen generally comprises providing a carbohydrate antigen having (e.g., chemically modified to comprise) a thiol-specific functional group; providing a carrier protein having one or more free thiol groups (e.g., one or more solvent-accessible cysteine residues, and/or cysteines not involved in disulfide bridges); and reacting the carbohydrate antigen with the carrier protein, thereby coupling of the carbohydrate antigen to the carrier protein at one or more predictable (non-random) attachment points corresponding to the positions of the free thiol groups of the carrier protein. In some embodiments, the method may further comprise purifying or isolating the glycoconjugate immunogen, which may then be formulated as a vaccine (e.g., comprising an adjuvant).

As used herein, the term “**glycoconjugate**” refers to a carbohydrate antigen (e.g., an antigenic monosaccharide, di-saccharide, oligo-saccharide, or polysaccharide) coupled to a carrier protein in order to enhance the immunogenicity of carbohydrate antigen in a subject of interest. The expressions “carbohydrate antigen” and “sugar antigen” carry the same meaning as used herein. The term “**immunogen**” refers to an agent that is capable of being specifically bound by components of the immune system (e.g., by an antibody and/or lymphocytes), and generating a humoral and/or cell-mediated immune response in a subject of interest. As used herein, the term “immunogen” in an expression such as “glycoconjugate immunogen” refers to the ability (i.e., physical characteristic or property) of the glycoconjugate without limiting the glycoconjugate itself to a particular use (e.g., as an immunogen for generating an immune response in a subject). For example, in some embodiments, a glycoconjugate immunogen described herein may be employed in diagnostic assays or methods (e.g., in vitro methods) to detect the presence or absence of an antibody that binds to the glycoconjugate immunogen in a biological sample (e.g., from a subject). In some embodiments, the glycoconjugate immunogens described herein may be used for screening, identifying, or evaluating antibodies that bind specifically to the glycoconjugate immunogen (e.g., monoclonal antibodies that are diagnostically or therapeutically applicable).

In some aspects, the present description relates to the use of a “click chemistry-type” approach for directly conjugating carbohydrate antigens to free thiol groups of carrier proteins. More specifically, as shown in **Fig. 2**, photocatalytic thiol-ene reactions are preferred in which carbohydrate antigens (R_2) are modified to contain a terminal alkene functionality (alkenyl carbohydrate antigen) suitable for the direct covalent attachment to the cysteine thiol groups (-SH) of carrier proteins (R_1) using a photocatalytic thiol reaction. Referring to **Fig. 3**, a carbohydrate antigen comprising a B-cell epitope (such as the tumor-associated carbohydrate antigens Tn and TF) may be photochemically conjugated via a thiol-ene reaction

to an immunogenic carrier protein such as tetanus toxoid (TT), which is known to have six free thiol groups owing to its six available cysteine residues (excluding four cysteine residues involved in disulfide bridges). Conjugation methods described herein, in the case of the tetanus toxoid carrier protein shown in Fig. 3, preferably result in precisely six carbohydrate antigens being conjugated at predictable (non-random) attachment points. The methods described herein are applicable to any mono-, oligo-, and polysaccharides, natural or synthetic, that can be made to end with an alkene group (terminal alkene).

In some embodiments, the carbohydrate antigens described herein may be chemically modified to be linked (directly or indirectly via a linker or spacer) to a terminal alkene (e.g., via a glycosidic bond or a bond obtained by reductive amination, such as between an allyl amine and a reducing sugar, preferably using NaBH_4 and/or NaBH_3CN), wherein the terminal alkene group of the alkenyl carbohydrate antigen is conjugatable to a free thiol of a carrier protein (e.g., one or more solvent-accessible cysteine residues, and/or cysteines not involved in disulfide bridges) via a thiol-ene reaction (e.g., a photocatalytic thiol-ene reaction). The terminal alkene group of the alkenyl carbohydrate antigen may be a monosubstituted alkene, a vinyl group, or an allyl group. In preferred embodiments, the alkenyl carbohydrate antigen is water-soluble, enabling its use in aqueous-based thiol-ene reactions described herein.

As used herein, the “**glycosidic bond**” may comprise one or more of an S-glycosidic bond, an N-glycosidic bond, an O-glycosidic bond, or a C-glycosidic bond, or a bond obtained by reductive amination of a reducing sugar (e.g., using NaBH_4 or preferably NaBH_3CN). In some embodiments, the glycosidic bond may be one that is not cleavable by an endogenous enzyme (e.g., a glycohydrolase) of the subject to be administered. Such an uncleavable glycosidic bond may result in a glycoconjugate immunogen having a longer half-life following administration to the subject, which may in turn generate a more favorable immune response for therapeutic and/or antibody-generation purposes. In some embodiments, the glycosidic bond may be an S-glycosidic bond, an N-glycosidic bond, or a C-glycosidic bond, or a bond obtained by reductive amination, such as between an allyl amine and a reducing sugar.

In some embodiments, the use of a “**linker**” or “**spacer**” is preferred and such terms are used herein to refer to a chemical linkage that provides sufficient physical separation of the carbohydrate antigen from the carrier protein to which it is conjugated to allow the carbohydrate antigen to be recognized by the immune system of a subject (e.g., as opposed to being masked by the carrier protein). In some embodiments, the linker may comprise a chain of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 contiguous residues selected from C, S, N, and O.

As used herein, the term “**conjugatable**” refers to the ability or capability of at least two molecules (e.g., a carbohydrate antigen and a carrier protein) to be covalently bonded to one another via a chemical reaction, regardless of whether the molecules are actually covalently bonded to one another. In

contrast, the term “**conjugated**” refers to at least two molecules (e.g., a carbohydrate antigen and a carrier protein) which are covalently bonded to one another.

In some embodiments, different types of carbohydrate antigens may be made to end with a terminal alkene group that is directly conjugatable to a thiol group via a thiol-ene reaction. For example, carbohydrate antigens having a terminal alkene group may be synthesized by adapting the approaches described herein in **Examples 2-7** for the synthesis of the allyl Tn and allyl TF reactants.

In some embodiments, the carbohydrate antigens having a terminal alkene group may be conjugated to free thiol groups of carrier proteins using photocatalytic thiol-ene reactions, for example, by adapting the approaches described herein in **Examples 8 and 9** for the synthesis of the Tn-TT and the TF-TT conjugates. For example, one or more alkenyl carbohydrate antigens dissolved in an aqueous solvent (e.g., a buffer such as PBS), is mixed with a carrier protein also dissolved in an aqueous solvent (e.g., a buffer such as PBS) in a vessel (e.g., quartz cell) suitable for ultraviolet light irradiation. The mixture is then irradiated under short-wave, medium-wave or long-wave ultraviolet light (e.g., having a peak wavelength at about 254 nm, at about 355 nm, or at about 365 nm), in the presence or absence of a catalyst (e.g., a photoinitiator or activator). As used herein in the context of the thiol-ene reactions, the terms “**catalyst**,” “**photoinitiator**,” and “**activator**” may be used interchangeably to refer to substances that accelerate conjugation of a carbohydrate antigen to a carrier protein at the one or more free thiol groups via a photocatalytic thiol-ene reaction.

In some embodiments, the catalyst may be a water-soluble photoinitiator such as a water-soluble free radical-generating azo compound; 2,2'-azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride (Vazo 44 or VA-044); 2,2'-azobis(2-amidinopropane) dihydrochloride (AAPH); metals or metal ions having photoinitiator activity; or any derivative thereof having photoinitiator activity in a photocatalytic thiol-ene reaction described herein. In some embodiments, the catalyst may be a water-soluble photoinitiator such as a water-soluble peroxide such as *tert*-butyl hydroperoxide or benzoylperoxide, or ammonium persulfate, or other suitable catalysts.

In some embodiments, the catalyst may be a water-insoluble photoinitiator such as a water-insoluble free radical-generating azo compound; 2,2-dimethoxy-2-phenylacetophenone (DMPA), azobisisobutyronitrile (AIBN), 2,2'-azobis(2-methylpropionitrile), 4,4'-azobis(4-cyanopentanoic acid) (ACVA), 1,1'-Azobis(cyanocyclohexane) (ACHN), diazenedicarboxylic acid bis(*N,N*-dimethylamide) (TMAD); azodicarboxylic acid dipiperidide (ADD), or any derivative thereof having photoinitiator activity in a photocatalytic thiol-ene reaction described herein.

In some embodiments, the photocatalytic thiol-ene reactions described herein may comprise controlling the ratio of molar equivalents of the alkenyl carbohydrate antigen (i.e., carbohydrate antigen comprising a terminal alkene) per free thiol group of the carrier protein, for example, to reduce, minimize,

or avoid sugar polymerization. In some embodiments, the photocatalytic thiol-ene reactions described herein may comprise reacting between 1 to 300, 1 to 250, 1 to 200, 1 to 100, or 1 to 10, 1 to 5, or 1 to 2 molar equivalents of the carbohydrate antigen per free thiol group of the carrier protein.

In some embodiments, the photocatalytic thiol-ene reactions described herein are performed for sufficient time to achieve at least a 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50-fold reduction in total free thiol concentration in the carrier protein (e.g., as determined by Ellman test). In some embodiments, the photocatalytic thiol-ene reactions described herein are performed for 10 to 300, 10 to 270, 10 to 240, 10 to 210, 10 to 180, 10 to 150, 10 to 120, 10 to 90, 10 to 60, or 10 to 30 minutes.

In some embodiments, the photocatalytic thiol-ene reactions described herein may be performed at a pH between about 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, or 4.0, and about 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, or 10. In some embodiments, the photocatalytic thiol-ene reactions described herein may be performed at a pH that avoids carrier protein denaturation. In some embodiments, the photocatalytic thiol-ene reactions described herein may be performed at a pH of between 4 and 5, preferably at pH 4.4 (e.g., in an acetate buffer).

In some embodiments, the use of a water-soluble photoinitiator (over a photoinitiator requiring an organic solvent) may be advantageous, as the thiol-ene conjugation reactions described herein may be performed only using aqueous reagents and aqueous solvents, since organic solvents may contribute to denaturation of the carrier protein, as well as undesired/unpredictable conjugation to non-free cysteine residues (e.g., cysteines involved in intramolecular and/or intermolecular disulfide bonds), as described in Dondoni et al., 2009 and Dondoni et al., 2012.

In alternative embodiments, a water-insoluble photoinitiator may be employed together with an organic solvent, as needed, for dissolution thereof. Preferably, the presence or concentration of the organic solvent shall not contribute to denaturation of the carrier protein, as well as undesired/unpredictable conjugation to non-free cysteine residues (e.g., cysteines involved in intramolecular and/or intermolecular disulfide bonds). For example, such a photoinitiator may be one such as 2,2-dimethoxy-2-phenylacetophenone (DMPA).

In alternate embodiments to the thiol-ene approach described herein, the carbohydrate antigens described herein may be modified to have one or more thiol-specific functional groups such as iodoacetamides, maleimides, benzylic halides, or bromomethylketones, which react by S-alkylation of thiols to generate stable thioether products (e.g., carried out at relatively low pHs such as below 9, 8, 7.5, or 7, and above 5, 5.5, or 6, which may avoid unwanted, random carbohydrate antigen conjugations to lysine residues). Examples of carbohydrate antigens that have been modified to comprise thiol-specific functional groups (e.g., iodoacetamido and maleimido groups) are shown in Fig. 4. In some

embodiments, the maleimido groups qualify as terminal alkenes that are conjugatable to thiol groups of carrier proteins via a thiol-ene reaction.

In some embodiments, the carbohydrate antigens described herein may comprise one or more B cell epitopes, and/or may induce a humoral immune response, and/or may comprise a T cell epitope, and/or induce a cell-mediated immune response in a subject upon administration. In some embodiments, glycoconjugate immunogens described herein may induce at least a cell-mediated immune response (e.g., in addition to a humoral response) to the carbohydrate antigen upon administration to a subject.

In some embodiments, the carbohydrate antigens described herein may be or comprise, for example, a tumor associated carbohydrate antigen (TACA). Glycoproteins and glycolipids of the outer cell membranes of cancer cells overexpress particular O-glycans. These glycoproteins constitute a family of proteins collectively known as mucins (MUCs) with MUC1 representing the most widely investigated. MUCs on normal cells are heavily O-glycosylated due to active glycosyltransferases which give rise to complex glycosylation patterns. In cancer cells, down-regulation of key glycosyltransferases trigger the accumulation of shorter glycans, giving rise to much more limited O-glycosylation on mucins. The consequence of these altered glycosylation patterns between normal and cancerous cells is the over accumulation of TACAs, which are otherwise cryptic (masked) by the complex glycosylation on normal tissues. The most common TACAs include the Tn antigen (a monosaccharide having the structure N-acetylgalactosamine (GalNAc)), the TF antigen (*Thomsen-Friedenreich* antigen, a disaccharide having the structure Gal β 1-3GalNAc α 1), and their cognate sialylated analogs, respectively. Overexpression of both the Tn and TF antigens on the cell surface of several types of tumor cells contributes to cancer cell adhesion and severe metastasis to sites containing receptor lectins (lungs, liver, lymph nodes, etc.). Accordingly, in some embodiments, the TACA described herein is, is from, or comprises: Tn antigen, STn antigen, *Thomsen-Friedenreich* (TF) antigen, (2,3)-S-TF, (2,6)-S-TF, Globo H, GD2, GD3, GM2, GM3, N-glycolyl-GM3, Lea, sLea, Lex, sLex, or any combination thereof. The structures of some common TACAs are shown in Fig. 1. As used herein, the expression “is from”, when used in the context of carbohydrate antigens, refers to carbohydrate variants derived from a known carbohydrate antigen, wherein the variant retains at least the antigenicity of the known carbohydrate antigen. For persons skilled in art, the synthesis of alkenyl-ending TACAs can be performed by both chemical as well as by chemoenzymatic processes, as described in Danishefsky et al., 2015.

In some embodiments, the carbohydrate antigens described herein may be or comprise carbohydrate antigens associated with infectious agents, such as, but not limited to, bacteria and/or viruses, or associated with infections, such as, but not limited to bacterial infections and/or viral infections. In some embodiments, the carbohydrate antigens described herein may be or comprise, for example, a viral polysaccharide antigen, or a bacterial capsular polysaccharide (CPS). In some

embodiments, the bacterial CPS is, is from, or comprises a Pneumococcal and/or Streptococcal polysaccharide serotype, meningococcal CPS, or influenza (such as influenza type a and b) CPS.

In some embodiments, the conjugation methods described herein may conjugate the same or more than one type of carbohydrate antigen to the carrier protein, thereby producing a multi-valent glycoconjugate immunogen. In some embodiments, the multi-valent glycoconjugate immunogens described herein may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more of the same or different types of carbohydrate antigens conjugated to a carrier protein. In some embodiments, the multi-valent glycoconjugate immunogens described herein may comprise more than one carbohydrate antigen that is conjugated to a single free thiol group on the carrier protein (e.g., via branched linker). In some embodiments, the multi-valent glycoconjugate immunogens described herein may comprise a plurality (e.g., at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more) of carbohydrate antigens that are conjugated to a single free thiol group on the carrier protein as a dendrimer (e.g., via linkers having extensive branching). In some embodiments, the carrier proteins described herein comprise one or more free thiol groups. As used herein, “free thiol” or “free thiol group” refers to carrier proteins having one or more sulfhydryl groups that are available for chemical modification and/or conjugation (e.g., to a carbohydrate antigen as described herein). In some embodiments, the free thiol concentration of a given carrier protein may be measured for example using the Ellman test using a cysteine standard curve, as described in the present Examples. The free thiol concentration of a given carrier protein may be expressed as fold reduction in free thiol concentration. In some embodiments, the glycoconjugates described herein may have at least a 50, 45, 40, 35, 30, 25, 20, 15, 10, 5, 4, 3, or 2-fold decrease in free thiol concentration following a conjugation method described herein (e.g., as measured by Ellman test).

In some embodiments, the carrier proteins described herein may be engineered to add one or more further cysteine residues, for example at the amino terminus, the carboxy terminus, or any solvent-accessible position of the carrier protein therebetween (as opposed to at positions that are not solvent accessible or “buried” within the three-dimensional structure of the protein carrier). In some embodiments, free thiol groups of protein carriers described herein may be defined as cysteines that are readily conjugatable to an alkenyl carbohydrate antigen by a photocatalytic thiol-ene reaction, when the reaction is performed under conditions sufficiently mild to not destroy the immunogenicity, structure, or activity of the carrier protein. In some embodiments, the term “activity” as used herein in relation to a carrier protein refers to the ability of the carrier protein to preserve the immunogenicity of the carbohydrate antigen (e.g., to an antibody that is known to specifically bind to the carbohydrate antigen).

In some embodiments, the free thiol group may refer to one or more solvent-accessible cysteine residues of the carrier protein, and/or cysteines that are not involved in disulfide bridges, which in some cases may be important to maintain the structure of the carrier protein. In some embodiments, the carrier

protein may be a protein having one or more disulfide bridges, and wherein the one or more disulfide bridges remain unaffected (i.e., intact) following conjugation to the carbohydrate antigen. In some embodiments, the carrier proteins described herein do not comprise free thiol-groups at their N and/or C-termini. Alternatively, in some embodiments, the carrier proteins described herein may comprise, or be engineered to comprise, a free thiol-group at their N and/or C-termini. In some embodiments, the carrier protein may be a protein having one or more disulfide bridges, and wherein the carrier protein may be pre-treated with a reducing agent (e.g., dithiothreitol (DTT), 2-mercaptoethanol, tris(2-carboxyethyl)phosphine (TCEP), or 2-mercaptoethylamine-HCl) to expose one or more additional free thiol groups for conjugation to the carbohydrate antigen. Such pre-treatments may be useful for example when the denatured (reduced) carrier protein has higher immunogenicity in a subject than the native (unreduced) carrier protein.

In some embodiments, the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, may be pre-treated with a reducing agent (e.g., as described herein). This pre-treatment may expose additional free thiol groups available for conjugation. In some embodiments, the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, may be pre-treated with a thiolating agent (e.g., 2-imminothiolane, N-hydroxysuccinimide dithiopropionate (DPS)). This pre-treatment may be employed to increase the number free thiol groups available for conjugation. In some embodiments, the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, may be pre-treated with a thiolating agent, and subsequently pre-treated with a reducing agent.

In some embodiments, it may be advantageous to avoid having too many multiple carbohydrate antigens conjugated to adjacent positions on the carrier proteins. In some embodiments, the carrier protein may preferably lack a cysteine-rich domain (e.g., a segment of at least 4, 5, 6, 7, 8, 9, or 10 consecutive amino acids comprising at least 50% of cysteine residues).

In some embodiments, the carrier protein may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 total cysteine residues. In some embodiments, the carrier protein may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 total free cysteine residues.

In some embodiments, the total number of carbohydrate antigens comprised in the glycoconjugate immunogens described herein is equal to the number of free thiol groups available on the carrier protein prior to conjugation. In some embodiments, the glycoconjugate immunogens described herein may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 carbohydrate antigens per carrier protein. In some embodiments, the present description relates to a composition comprising glycoconjugate immunogens having about or at least 70%, 75%, 80%, 85%, 90%, or 95% homogeneity in terms of carbohydrate conjugation species (e.g., at least 90% of

glycoconjugate immunogen species/molecules in the composition have the same number of carbohydrate antigens conjugated to the carrier protein).

In some embodiments, the carrier proteins described herein are proteins (e.g., peptides or polypeptides) that are preferably immunogenic to a subject that is to be administered to the glycoconjugate immunogen. Preferably, conjugation of the carbohydrate antigen to the carrier protein increases the immunogenicity of the carbohydrate antigen upon administration to the subject, as compared to administration of the unconjugated carbohydrate antigen (i.e., the carbohydrate antigen administered to the subject alone).

In some embodiments, the carrier proteins may comprise a T cell epitope, and/or induce a cell-mediated immune response in the subject upon administration.

In some embodiments, the carrier protein is a protein that is exogenous to the subject to be administered, which preferably has no (close) ortholog in the subject. In the context of human vaccine production, a carrier protein described herein refers to a “**carrier protein suitable for human use**” or simply “**suitable carrier protein**”, which means a carrier protein that is antigenically distinct from human proteins such that the carrier protein would not be considered as a “self-antigen” in humans. The use of carrier proteins that are too antigenically similar to corresponding human proteins may result in the carrier protein being considered as a “self-antigen”, which may not be ideal in human vaccines. For example, glycoconjugate immunogens consisting of TF antigen *randomly* conjugated to the ϵ -amino groups of lysine residues of bovine serum albumin (BSA) have been previously described and characterized (e.g., Demian et al., 2014; Rittenhouse-Diakun et al., 1998; Heimburg et al., 2006; Tati et al., 2017). However, not only was the level of carbohydrate on the 59 lysine residues of BSA random and inefficient (no more than 4 to 6 TF antigens were conjugated per BSA molecule), BSA would not be suitable as a carrier protein in human vaccines because it is too antigenically similar to human albumin. In some embodiments, the carrier protein is not albumin (e.g., bovine serum albumin).

Preferably, the carrier protein described herein may be a protein that has already received regulatory (e.g., FDA) approval for administration to human subjects (e.g., in approved vaccines). In some embodiments, the carrier protein is, is from, or comprises Tetanus Toxoid (TT), Diphtheria Toxoid (DT), cross-reacting material 197 (CRM197), Meningococcal Outer Membrane Protein Complex (OMPC), *H. Influenzae* Protein D (HiD), a cytokine, or an immunogenic fragment (or variant) thereof.

In some embodiments, the carrier protein may be TT which contains 10 cysteine residues, 4 of which being engaged in disulfide bridges, giving rise to a glycoconjugate immunogen having 6 conjugated carbohydrate antigens. In some embodiments, the carrier protein may be CRM197 which contains 4 cysteine residues, giving rise to a glycoconjugate immunogen having 4 conjugated carbohydrate antigens.

In some embodiments, the carrier proteins may be engineered to introduce additional cysteine residues in order to introduce additional free thiol groups for conjugation to carbohydrate antigens. In some embodiments, the cysteine residues may be engineered at solvent exposed portions of the carrier protein.

5 In some aspects, the present description relates to a method for producing a glycoconjugate vaccine or an immune response-triggering composition. The method may comprise formulating the glycoconjugate immunogen described herein with a pharmaceutically acceptable excipient, and/or an adjuvant. In some embodiments, the adjuvant is or comprises: an inorganic compound, a mineral oil, a microbial derivative, a plant derivative, a cytokine, squalene, alum, aluminum hydroxide, aluminum
10 phosphate, calcium phosphate hydroxide, a toll-like receptor agonist, an immunostimulatory polynucleotide (e.g., CPG), an immunostimulatory lipid, Freund's adjuvant, RIBI's adjuvant, QS-21, muramyl dipeptide, TiterMax™, Steviune™, Stimune™, or any combination thereof.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more carbohydrate antigens and an immunogenic carrier protein having one or more
15 solvent-accessible cysteine residues, wherein the one or more carbohydrate antigens are linked to the immunogenic carrier protein at the one or more solvent-accessible cysteine residues. The conjugation of the one or more carbohydrate antigens to the immunogenic carrier protein increases the immunogenicity of the one or more carbohydrate antigens upon administration to a subject, as compared to administration of the unconjugated carbohydrate antigen. As used herein, the term “synthetic” refers to a compound that
20 is not a product of nature, which is produced by human intervention.

In some embodiments, glycoconjugate immunogens described herein may comprise more than one species of carbohydrate antigen (e.g., more than one type of TACA) conjugated to the same carrier protein. For example, glycoconjugate immunogens described herein may comprise at least 2, 3, 4, 5, 6, 7,
8, 9, 10, 11, 12, 13, 14, 15 or more types or species of carbohydrate antigen (e.g., TACA). In some
25 embodiments, glycoconjugate immunogens described herein may comprise any combination of TACAs selected from Tn, S-Tn, *Thomsen-Friedenreich* (TF), (2,3)-S-TF, (2,6)-S-TF, Globo H, GD2, GD3, GM2, GM3, N-glycolyl-GM3, Lea, sLea, Lex, and sLex. In some embodiments, ratio in the combination of each TACAs may vary with the targeted tumor and may comprise between 1 to 20 molar ratios. In this way, the glycoconjugate immunogens described herein may be tailored, for example, to specific forms of
30 cancer that are associated with increased expression of particular combinations of multiple TACAs.

In some aspects, the present description relates to a glycoconjugate vaccine which is produced by a method described herein, and/or which comprises a synthetic glycoconjugate immunogen described herein, and further comprises a pharmaceutically acceptable excipient, and/or an adjuvant. As used herein, the term “vaccine” refers to a composition comprising a glycoconjugate immunogen described


herein that is administered a subject to provide a therapeutic benefit to the subject. In some embodiments, the glycoconjugate vaccines described herein may be a prophylactic vaccine or a therapeutic vaccine.

Vaccine compositions can be administered in dosages and by techniques well known to those skilled in the medical or veterinary arts, taking into consideration such factors as the age, sex, weight, species and condition of the recipient animal, and the route of administration. The route of administration can be percutaneous, via mucosal administration (e.g., oral, nasal, ocular) or via a parenteral route (e.g., intradermal, intramuscular, subcutaneous). Vaccine compositions can be administered alone, or can be co-administered or sequentially administered with other treatments or therapies. Forms of administration may include suspensions and preparations for parenteral, subcutaneous, intradermal or intramuscular administration (e.g., injectable administration) such as sterile suspensions or emulsions. Vaccines may be administered as a spray or mixed in food and/or water or delivered in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose, or the like. The compositions can contain auxiliary substances such as wetting or emulsifying agents, pH buffering agents, adjuvants, gelling or viscosity enhancing additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. Standard pharmaceutical texts, such as "Remington's Pharmaceutical Sciences," 1990 may be consulted to prepare suitable preparations, without undue experimentation.

In some aspects, the present description relates to a method of immunizing, vaccinating, or treating a subject comprising administering to the subject a glycoconjugate immunogen or a glycoconjugate vaccine described herein.


In some aspects, the present description relates to a carbohydrate antigen chemically modified to be linked to a sulfhydryl-specific (thiol specific) functional group. In some embodiments, the carbohydrate antigen is chemically modified to be linked to a terminal alkene (e.g., via a glycosidic bond), wherein the terminal alkene group is conjugatable to a free thiol of a carrier protein via a thiol-ene reaction. In some embodiments, the carbohydrate antigen and the terminal alkene group are linked via a linker as described herein. In some embodiments, the terminal alkene group may be a vinyl group or an allyl group.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein each linker has the structure:




wherein: **SA** is a sugar antigen or a portion thereof; **S-PC** is a carrier protein; **X** is O, S, NR₁, or CH₂; **R₁** is -H, -COH, -COCH₃, or -COEt; **n** is 1, 2, 3, 4, or 5; and **R₂** is H or Me; or a stereoisomer (e.g., diastereomer) thereof.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein each linker has the structure:




wherein: **SA** is a sugar antigen or a portion thereof; **S-PC** is a carrier protein; **X** is S, NR₁, CH₂ or O; **R₁** is -H, -COH, -COMe, or -COEt; **n** is 1, 2, 3, 4, or 5; **R₂** is H or Me; **q** is 1, 2, 3, 4, or 5; **R₃** and **R₄** are each a hydrogen atom and **m** is 1, 2, 3, 4 or 5, or **R₃** and **R₄** form together a radical -CO-CH₂- or a radical -CO-CH₂-CH₂- with the carbonyl linked to the nitrogen atom, and **m** is 1; or a stereoisomer (e.g., diastereomer) thereof. In some embodiments, when **X** is O, the sugar antigen does not comprise Tn or STn.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein the synthetic glycoconjugate immunogen has the structure:



wherein

- **y** is at least 1;
- **SA** is selected from the group consisting of sugar antigens or immunogenic portions thereof, and when **y** is more than 1, **SA** are identical or different;
- [**S**]_{**z**}-**PC** is a carrier protein having one or more sulfur atoms originating from the one or more free thiol groups;
- **z** is at least 1 and is at least equal to **y**; and
- **L** is a linker selected from the group consisting of linkers having the structure:




wherein:

- **X** is O, S, NR₁, or CH₂;
- **R₁** is -H, -COH, -COCH₃, or -COEt;
- **n** is 1, 2, 3, 4, or 5; and
- **R₂** is H or Me; and

when **y** is more than 1, **L** are identical or different;


or a stereoisomer thereof.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein the synthetic glycoconjugate immunogen has the structure:



wherein

- **y** is at least 1;
- **SA** is selected from the group consisting of sugar antigens or immunogenic portions thereof, and when **y** is more than 1, **SA** are identical or different;
- [**S**]_{**z**}-**PC** is a carrier protein having one or more sulfur atoms originating from the one or more free thiol groups;
- **z** is at least 1 and is at least equal to **y**; and
- **L** is a linker selected from the group consisting of linkers having the structure:



wherein:


- **X** is S, NR₁, CH₂ or O;
- **R₁** is -H, -COH, -COMe, or -COEt;
- **n** is 1, 2, 3, 4, or 5;
- **R₂** is H or Me;
- **q** is 1, 2, 3, 4, or 5;

- R_3 and R_4 are each a hydrogen atom and m is 1, 2, 3, 4 or 5, or R_3 and R_4 form together a radical $-CO-CH_2-$ or a radical $-CO-CH_2-CH_2-$ with the carbonyl linked to the nitrogen atom, and m is 1; and

when y is more than 1, L are identical or different;


or a stereoisomer thereof.

In some aspects, the present description relates to a synthetic glycoconjugate immunogen comprising one or more sugar antigens covalently linked to one or more free thiol groups of a carrier protein via a linker, wherein the synthetic glycoconjugate immunogen has the structure:




wherein

- y is at least 1;
- SA is selected from the group consisting of sugar antigens or immunogenic portions thereof, and when y is more than 1, SA are identical or different;
- $[S]_z-PC$ is a carrier protein having one or more sulfur atoms originating from the one or more free thiol groups;
- z is at least 1 and is at least equal to y ; and
- L is a linker selected from the group consisting of linkers having the structure:



wherein


- X is S, NR_1 , CH_2 or O;
- R_1 is -H, -COH, -COMe, or -COEt;
- n is 1, 2, 3, 4, or 5;
- R_2 is H or Me;
- q is 1, 2, 3, 4, or 5;
- r is 1, 2, 3, 4 or 5;
- R_5 is S-PC, a covalent bond or a radical of structure:



wherein R_3 and R_4 are each a hydrogen atom and m is 1, 2, 3, 4 or 5, or R_3 and R_4 form together a radical $-CO-CH_2-$ or a radical $-CO-CH_2-CH_2-$ with the carbonyl linked to the nitrogen atom, and m is 1; and

when y is more than 1, L are identical or different;
or a stereoisomer thereof.


In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as defined above, wherein the linker has the structure:



wherein

- X is S, NR_1 , CH_2 or O;
- R_1 is -H, -COH, -COMe, or -COEt;
- n is 1, 2, 3, 4, or 5;
- R_2 is H or Me;
- q is 1, 2, 3, 4, or 5; and
- r is 1, 2, 3, 4 or 5.

In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as defined above, wherein the linker has the structure:




wherein

- X is S, NR_1 , CH_2 or O;
- R_1 is -H, -COH, -COMe, or -COEt;
- n is 1, 2, 3, 4, or 5;

- R_2 is H or Me;
- q is 1, 2, 3, 4, or 5; and
- r is 1 or 2.

5 In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as defined above, wherein the linker has the structure:



wherein

- X is S, NR_1 , CH_2 or O;
- R_1 is -H, -COH, -COMe, or -COEt;
- n is 1, 2, 3, 4, or 5;
- R_2 is H or Me;
- q is 1, 2, 3, 4, or 5; and
- r is 1 or 2.

15 In some embodiments, the present description relates to the synthetic glycoconjugate immunogen as described above, wherein y is an integer varying from 1 to 50, 1 to 40, 1 to 30, 1 to 20, or 1 to 10, 1 to 5, or 1 to 2.

20 In some embodiments, the sugar antigen may be a carbohydrate antigen as described herein, and/or the carrier protein may be a carrier protein as described herein. In some embodiments, the synthetic glycoconjugate immunogen may be a multi-valent glycoconjugate immunogen, for example, comprising 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more of the same or different types of carbohydrate antigens conjugated to a carrier protein. In some embodiments, the glycoconjugate immunogen may induce a cell-mediated immune response to the carbohydrate antigen upon
25 administration to a subject.

In some aspects, the present description relates to a vaccine comprising a synthetic glycoconjugate immunogen as described herein, and a pharmaceutically acceptable excipient and/or an adjuvant (e.g., an adjuvant as described herein).

30 In some aspects, the present description relates to the use of the glycoconjugate immunogen prepared by a method as described herein, or a synthetic glycoconjugate immunogen as described herein, for the manufacture of a vaccine.

In some aspects, the present description relates to the use of the glycoconjugate immunogen prepared by a method as described herein, the glycoconjugate vaccine produced by a method as described herein, a synthetic glycoconjugate immunogen as described herein, or a vaccine as described herein, for the treatment of a subject having a disease associated with increased expression of said one or more carbohydrate or sugar antigens.

In some aspects, the present description relates to the use of the glycoconjugate immunogen prepared by a method as described herein, the glycoconjugate vaccine produced by a method as described herein, a synthetic glycoconjugate immunogen as described herein, or a vaccine as described herein, for producing an antibody that specifically binds to the glycoconjugate immunogen.

In some aspects, the present description relates to the use of the glycoconjugate immunogen prepared by a method as described herein, the glycoconjugate vaccine produced by a method as described herein, a synthetic glycoconjugate immunogen as described herein, or a vaccine as described herein, for detecting an antibody that specifically binds to the glycoconjugate immunogen.

In some aspects, the present description relates to a method for producing a glycoconjugate, the method comprising: (a) providing a carbohydrate antigen covalently linked to a terminal alkene (alkenyl carbohydrate antigen), the terminal alkene being directly conjugatable to a thiol group via a thiol-ene reaction; (b) providing a conjugate material suitable for conjugation to the carbohydrate antigen via a thiol-ene reaction as described herein, said conjugate material having one or more free thiol groups; and (c) performing a photocatalytic thiol-ene reaction to directly conjugate the carbohydrate antigen to the conjugate material at the one or more free thiol groups, thereby producing the glycoconjugate. In some embodiments, the conjugate material is or comprises a polymer, a polypeptide, a carrier protein as defined herein, a solid support, a particle, or any other material having a free thiol group suitable for conjugation to the carbohydrate antigen via a thiol-ene reaction as described herein.

In some aspects, the present description relates to the use of a glycoconjugate produced by a method described herein, for detecting or screening for the presence of an antibody that specifically binds to a carbohydrate antigen or a tumor-circulating cell comprising a carbohydrate antigen, or for detecting the presence of antibodies resulting from an immunization or vaccination with a carbohydrate antigen. In some embodiments, the detection or screening may be performed via any suitable detection method, such as an immunosorbent assay, ELISA, microarray, or immunoblot analysis.

In some aspects, the present description relates a method of treating a subject comprising administering a glycoconjugate or glycoconjugate immunogen as defined herein or produced by a method as described herein, to generate an immune response in said subject to a carbohydrate antigen, and screening a biological sample from said subject for the presence of antibodies that specifically binds to the carbohydrate antigen.

Other objects, advantages and features of the present description will become more apparent upon reading of the following non-restrictive description of specific embodiments thereof, given by way of example only with reference to the accompanying drawings.

ITEMS

1. A method for producing a glycoconjugate immunogen, the method comprising:

- (a) providing a water-soluble carbohydrate antigen covalently linked to a terminal alkene, the terminal alkene being directly conjugatable to a thiol group via a thiol-ene reaction and wherein the carbohydrate antigen is unprotected and water-soluble;
- (b) providing a carrier protein having one or more free thiol groups; and
- (c) performing a photocatalytic thiol-ene reaction to directly conjugate the carbohydrate antigen to the carrier protein at the one or more free thiol groups, thereby producing the glycoconjugate immunogen, wherein the photocatalytic thiol-ene reaction is performed:
 - (i) at a pH between 6 and 8;
 - (ii) under reaction conditions that avoid carrier protein denaturation, and/or that retain the carrier protein's activity, antigenicity, and/or structure; and
 - (iii) in the absence of any organic solvent, or in the presence of an organic solvent at a concentration sufficiently low to avoid carrier protein denaturation;

wherein the carrier protein is immunogenic when administered to a subject, and wherein conjugation of the carbohydrate antigen to the carrier protein increases the immunogenicity of the carbohydrate antigen upon administration to the subject, as compared to administration of the unconjugated carbohydrate antigen, and wherein the carbohydrate antigen comprises a B cell epitope and induces a humoral immune response in the subject.

2. The method of item 1, wherein the photocatalytic thiol-ene reaction is performed at a pH between 6.5 and 7.5.

3. The method of item 1 or 2, wherein said photocatalytic thiol-ene reaction is performed in the presence of a water-soluble catalyst.

4. The method of item 3, wherein the water-soluble catalyst is: a free radical-generating azo compound; 2,2'-azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride; 2,2'-azobis(2-amidinopropane) dihydrochloride (AAPH); metals or metal ions having photoinitiator activity; a peroxide; *tert*-butyl hydroperoxide; benzoylperoxide; ammonium persulfate; or any derivative thereof having photoinitiator activity.

5. The method of item 1 or 2, wherein said photocatalytic thiol-ene reaction is performed in the presence of a water-insoluble catalyst.
6. The method of item 5, wherein the water-insoluble catalyst is: a free radical-generating azo compound; 2,2-dimethoxy-2-phenylacetophenone (DMPA); azobisisobutyronitrile (AIBN); 2,2'-azobis(2-methylpropionitrile); 4,4'-Azobis(4-cyanopentanoic acid) (ACVA); 1,1'-azobis(cyanocyclohexane) (ACHN); diazenedicarboxylic acid bis(*N,N*-dimethylamide) (TMAD); azodicarboxylic acid dipiperidide (ADD); or any derivative thereof having photoinitiator activity.
7. The method of any one of items 1 to 6, wherein said photocatalytic thiol-ene reaction comprises irradiation under short-wave ultraviolet light.
8. The method of item 7, wherein said photocatalytic thiol-ene reaction comprises irradiation at about 254 nm.
9. The method of any one of items 1 to 7, wherein said photocatalytic thiol-ene reaction comprises irradiation under long-wave ultraviolet light.
10. The method of item 9, wherein said photocatalytic thiol-ene reaction comprises irradiation at about 355 nm or 365 nm.
11. The method of any one of items 1 to 10, wherein said photocatalytic thiol-ene reaction comprises reacting between 1 to 200 molar equivalents of the carbohydrate antigen per free thiol group of the carrier protein.
12. The method of any one of items 1 to 11, wherein said carbohydrate antigen, following conjugation to the carrier protein, is not cleavable from the carrier protein by an endogenous enzyme of the subject.
13. The method of any one of items 1 to 12, wherein said carbohydrate antigen is covalently linked to the terminal alkene.
14. The method of any one of items 1 to 12, wherein said carbohydrate antigen is conjugated to the carrier protein, via an O-glycosidic bond, an S-glycosidic bond, an N-glycosidic bond, or a C-glycosidic bond, or a bond obtained by reductive amination between an allyl amine and a reducing sugar.
15. The method of any one of items 1 to 14, wherein the carbohydrate antigen is or comprises a tumor associated carbohydrate antigen (TACA), a viral polysaccharide antigen, or a bacterial capsular polysaccharide (CPS).
16. The method of item 15, wherein the TACA is or is from: Tn, S-Tn, *Thomsen-Friedenreich* (TF), (2,3)-S-TF, (2,6)-S-TF, Globo H, GD2, GD3, GM2, GM3, N-glycolyl-GM3, Lea, sLea, Lex, sLex, or any combination thereof.

17. The method of any one of items 1 to 16, wherein said photocatalytic thiol-ene reaction conjugates more than one type of carbohydrate antigen to the carrier protein.

18. The method of item 15, wherein the bacterial CPS is or is from a Pneumococcal and/or Streptococcal polysaccharide serotype, meningococcal CPS, or influenza CPS.

19. The method of any one of items 1 to 18, wherein the carbohydrate antigen in (a) is linked to the terminal alkene via a linker.

20. The method of any one of items 1 to 19, wherein the carrier protein comprises one or more cysteine residues having the one or more free thiol groups, or optionally is engineered to add one or more further cysteine residues at the amino terminus, the carboxy terminus, or at a solvent-accessible position of the carrier protein.

21. The method of any one of items 1 to 20, wherein the carrier protein is or is from: Tetanus Toxoid (TT), Diphtheria Toxoid (DT), cross-reacting material 197 (CRM197), Meningococcal Outer Membrane Protein Complex (OMPC), H. Influenzae Protein D (HiD), a cytokine, an immunogenic peptide, Tetanus Toxin 831-844, albumin, or an immunogenic fragment thereof.

22. The method of any one of items 1 to 21, wherein:

(i) the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, is pre-treated with a reducing agent;

(ii) the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, is pre-treated with a thiolating agent;

(iii) the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, is pre-treated with a thiolating agent, and subsequently pre-treated with a reducing agent;

(iv) the carrier protein is a protein having one or more disulfide bridges, and wherein the one or more disulfide bridges remain unaffected following said photocatalytic thiol-ene reaction;

(v) the carrier protein is a protein having one or more disulfide bridges, and wherein the carrier protein is pre-treated with a reducing agent to expose one or more additional free thiol groups for conjugation to the carbohydrate antigen; or

(iv) the carrier protein is a protein having one or more disulfide bridges, and wherein the carrier protein is pre-treated with a thiolating agent to expose one or more additional free thiol groups for conjugation to the carbohydrate antigen, and subsequently pre-treated with a reducing agent.

23. The method of any one of items 1 to 22, wherein the total number of carbohydrate antigens comprised in the glycoconjugate immunogen is equal to the number of free thiol groups available on the carrier protein prior to conjugation.

24. The method of any one of items 1 to 23, wherein the glycoconjugate immunogen induces a cell-mediated immune response to the carbohydrate antigen upon administration to the subject.

25. The method of any one of items 1 to 24, further comprising: (d) purifying the glycoconjugate immunogen.

26. A method for producing a glycoconjugate vaccine or an adaptive immune response-triggering composition, the method comprising formulating the glycoconjugate immunogen prepared by the method of any one of items 1 to 25 with a pharmaceutically acceptable excipient, and/or an adjuvant.

27. The method of item 26, wherein the adjuvant is or comprises: an inorganic compound, a mineral oil, a microbial derivative, a plant derivative, a cytokine, squalene, alum, aluminum hydroxide, aluminum phosphate, calcium phosphate hydroxide, a toll-like receptor agonist, an immunostimulatory polynucleotide, an immunostimulatory lipid, Freund's adjuvant, RIBI's adjuvant, QS-21, muramyl dipeptide, or any combination thereof.

EXAMPLES

Example 1: General Methods

Reactions were carried out under argon atmosphere using commercially available HPLC grade. Commercially available reagents (Sigma Aldrich) were used without further purification. *N*-Acetyl-D-galactosamine and *N*-acetylneuraminic acid were provided from Rose Scientific Ltd. Alberta, Canada. The Fmoc- β -Ala-Wang resin and Fmoc amino acid were available commercially from Peptide Technologies Ltd, Pierrefonds, Qc, Canada. Progress of reactions was monitored by thin-layer chromatography using silica gel 60 F₂₅₄ coated plates (E. Merck). The conjugation by the click thiol-ene photoreaction was done in a quartz cuvette (10x10 mm path length, Fisher Scientific Canada, Cat. No. 14-958-130) placed between two hand held UV 365 nm lamps (UV-AC Hand Lamp, Dual 254/365nm UV; 115V-60Hz, 0.16 amps, VWR Canada, Cat. No. 89131-492). Flash chromatography was performed using ZEOprep™ silica gel 60 (40-63 μ m) from Canadian Life Science. Detection was carried out under UV light or by spraying with 20% ethanolic sulfuric acid or molybdate or KMnO₄ solution followed by heating. NMR spectra were recorded on Bruker ULTRASHIELD™ 300 MHz and Bruker Avance™III HD 600 MHz spectrometers. Proton and carbon chemical shifts (δ) are reported in ppm relative to the chemical shift of residual CHCl₃, which was set at 7.26 ppm (¹H) and 77.16 ppm (¹³C). Coupling constants (*J*) are reported in Hertz (Hz), and the following abbreviations are used for peak multiplicities: singlet (s), doublet (d), doublet of doublets (dd), doublet of doublet with equal coupling constants (t_{ap}), triplet (t), multiplet (m). Analysis and assignments were made using COSY (Correlated Spectroscopy)

and HSQC (Heteronuclear Single Quantum Coherence) experiments. High-resolution mass spectra (HRMS) were measured with a LC-MS-TOF (Liquid Chromatography Mass Spectrometry Time Of Flight) instrument from Agilent technologies in positive and/or negative electrospray mode by the analytical platform of UQAM. Either protonated ions (M+H)⁺ or sodium adducts (M+Na)⁺ were used for empirical formula confirmation. The native TT and TT-conjugate were dialyzed using 2000 KDa benzoylated dialysis tubing (Sigma-Aldrich (Ontario, Canada). The thiol contents of both native and conjugated TT were determined by the Ellman test at 412 nm (Ellman, G. L. *Arch. Biochem. Biophys.* 1959, 82, 70-77). The total sugar content of the TT-conjugate was determined by the colorimetric DuBois test measured at 492 nm (Dubois, M.; Gilles, K. A.; Hamilton, J. K.; Rebers, P. A.; Smith, F. Colorimetric Method for Determination of Sugars and Related Substances. *Anal. Chem.*, 1956, 28, 350-356) using a UV/VIS Ultrospec 100 *prot* spectrophotometer (Biochrom, USA). Dynamic Light Scattering (DLS), particle size distributions were measured in PBS using a Zetasizer Nano S90 from Malvern. The mouse monoclonal IgG3 antibody JAA-F11 was produced as previously described in Rittenhouse-Diakun et al., 1998.


General Solid Phase Peptide Synthesis (SPPS) procedure

The procedure of Solid-Phase Peptide Synthese (SPPS) was followed under litterature procedure (Papadopoulos et al., 2012) and started with Fmoc-β-Ala-Wang resin (650 mg, 0.34 mmol, 1.0 equiv.; 100-200 mesh, loading = 0.52 mmol/g). The reactions were conducted by rotation agitation in Econo-Pac disposable columns 1.5 x 14 cm (20 mL) (Bio-Rad Laboratories, ON, Canada). The resin was swollen in CH₂Cl₂ during 1 h, then filtered and reconditioned in DMF during 1 h. The Fmoc- protecting group of the commercial resin or of amino acids were removed with a solution of 20% piperidine in DMF (5 mL, 2 x 5 min then 1 x 10 min). The solvents and reagents were removed by filtration, and the resin was washed with DMF, CH₂Cl₂ and MeOH (3X with each solvent). The presence of free amino groups was verified by a Kaiser test or TNBS test. The free amines on the resin were treated with a solution de preactivated Fmoc amino acid : 3 equiv of amino acid, 3 equiv of HBTU (*N,N,N',N'*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uronium hexafluorophosphate) and a catalytic amount of HOBT (1-hydroxybenzotriazole) in DMF at 4 °C (10 min). DIPEA (Diisopropylamine, 9 equiv) was then added into the mixture and stirred at room temperature for 1 h 30 min. Completion of the coupling was determined using Kaiser or a TNBS colorimetric test. After filtration, the resin was washed and the Fmoc removal procedure was again repeated. At the end of the synthetic sequences, the last free amine was capping by acetylation (Ac₂O/DIPEA/DMF 1:1:8, 1 h). After filtration, the solutions were drained off, the resin was dried under vacuum and the cleavage was carried out using trifluoroacetic acid/water/ethanedithiol/triisopropylsilane (94.0/2.5/2.5/1.0) for 3 h. The resulting peptide was precipitated with methyl *tert*-butyl ether and isolated

from the resin bead by centrifugation (20 min, 2000 rpm, 3X). The precipitates were dried carefully with a stream of air jet. The crude peptide was solubilised in H₂O to separate from resin. The solution was then lyophilized to afford the desired peptide.

- 5 **dTT831-844-Cys-βAla.** From Fmoc-β-Ala-Wang resin (650 mg, 0.34 mmol, 1.0 equiv.; 100-200 mesh, loading = 0.52 mmol/g), the desired peptide **dTT831-844-Cys-βAla** was isolated as white powder (62 mg, 0.034 mmol, 10%). ESI⁺-LC-MS: [M+2H]²⁺ calcd for C₈₃H₁₃₅O₂₄N₁₉S, 906.9819; found, 906.9849; CAN/H₂O 5 to 95% 5.42 min.
- 10 **Cys-dTT831-844-βAla.** From Fmoc-β-Ala-Wang resin (650 mg, 0.34 mmol, 1.0 equiv.; 100-200 mesh, loading = 0.52 mmol/g), the desired peptide **Cys-dTT831-844-βAla** was isolated as white powder (62 mg, 0.034 mmol, 10%). ESI⁺-LC-MS: [M+2H]²⁺ calcd for C₈₃H₁₃₅O₂₄N₁₉S, 906.9819; found, 906.9844; CAN/H₂O 5 to 95% 5.32 min

15 **Example 2: Allyl 2-acetamido-3,6-di-O-pivaloyl-2-deoxy-α-D-glucopyranoside (Compound 2)**




Compound 2

- 20 Referring to Fig. 5, Acetyl chloride (2.76 mL, 38.80 mmol, 3.43 equiv.) was added dropwise to allylic alcohol (20.8 mL) at 0°C under argon atmosphere. At room temperature, *N*-acetyl-D-glucosamine (Compound 1) (2.50 g, 11.3 mmol, 1.00 eq.) was added. The reaction mixture was stirred at 70°C for 3 hours, then quenched by adding solid NaHCO₃ until pH 7. The suspension was filtered through a pad of Celite, washing several times with MeOH. The solvent was removed under reduced pressure, and the
- 25 crude allyl 2-acetamido-2-deoxy-D-glucosamine was precipitated by trituration with Et₂O/Ethanol. The solvent was then removed under reduced pressure several times after trituration. To a suspension of crude allyl 2-acetamido-2-deoxy-α-D-glucopyranoside intermediate, in the mixture of dry dichloromethane-pyridine (45 mL, v/v, 1:2) under nitrogen atmosphere at -15°C, pivaloyl chloride (3.90 mL, 31.64 mmol, 2.80 equiv.) was then added dropwise. The reaction mixture was stirred for 2 hours warmed to room
- 30 temperature to give the desired α-anomer (Compound 2) (R_f = 0.32) together with some β-anomer (R_f = 0.18); hexanes/EtOAc 1:1). The mixture was diluted then with CH₂Cl₂ and the organic phase was

successively washed with HCl (1M) several times, saturated aqueous solution of KHSO₄, saturated solution of NaHCO₃, and brine. The organic phase was dried over Na₂SO₄, filtered and evaporated under reduced pressure. The yellowish oil was purified by flash chromatography on silica gel (Hexane-EtOAc 6:4 to 1:1) to afford the desired compound allyl 2-acetamido-3,6-di-*O*-pivaloyl-2-deoxy- α -D-glucopyranoside (Compound 2) as white solid (4.85 g, 6.78 mmol, 60%). *R*_f = 0.32; hexanes/EtOAc 1:1; **Figs. 6A & 6B**: ¹H NMR (CDCl₃, 600 MHz): δ 5.87 (dddd, 1H, *J*_{H,H} = 16.8, 10.5, 6.2, 5.3 Hz, OCH₂CH=CH₂), 5.77 (d, 1H, *J*_{NH,H2} = 9.7 Hz, NH), 5.31-5.25 (m, 1H, OCH₂CH=CH₂), 5.22 (dd, 1H, *J*_{H,H} = 10.4, 1.3 Hz, OCH₂CH=CH₂), 5.09 (dd, 1H, *J*_{3,4} = 10.7, *J*_{2,3} = 9.3 Hz, H-3), 4.83 (d, 1H, *J*_{1,2} = 3.7 Hz, H-1), 4.39 (m, 1H, H-6a), 4.35-4.25 (m, 2H, H-6b and H-2), 4.19 (m, 1H, OCH₂), 4.02-3.93 (m, 1H, OCH₂), 3.85 (m, 1H, H-5), 3.59-3.48 (m, 1H, H-4), 3.03 (d, 1H, *J*_{4,OH} = 5.1 Hz, OH-4), 1.93 (s, 3H, NHCOCH₃), 1.23 (s, 9H, *tert*-Butyl) and 1.19 ppm (s, 9H, *tert*-Butyl); ¹³C NMR (CDCl₃, 150 MHz): δ 179.8, 179.1 (*tert*-BuCO), 169.7 (NHCO), 133.2 (OCH₂CH=CH₂), 118.1 (OCH₂CH=CH₂), 96.4 (C-1), 73.4 (C-3), 70.5 (C-5), 69.1 (C-4), 68.2 (OCH₂), 63.1 (C-6), 51.4 (C-2), 39.0, 38.9 (2 x C(CH₃)₃), 27.2, 27.0 (2 x C(CH₃)₃) and 23.2 ppm (CH₃). **Fig. 6C & 6D**: ESI⁺-HRMS: [M+H]⁺ calcd for C₂₁H₃₆O₈N, 430.2435; found, 430.2445. The β -anomer was isolated as white solid (971 mg, 2.26 mmol, 20%). *R*_f = 0.18; hexanes/EtOAc 1:1; ¹H NMR (CDCl₃, 300 MHz): δ 6.00 (d, 1H, *J*_{NH,H2} = 9.3 Hz, NH), 5.95-5.75 (m, 1H, OCH₂CH=CH₂), 5.35-5.03 (m, 3H, OCH₂CH=CH₂ and H-3), 4.55 (d, 1H, *J*_{1,2} = 8.4 Hz, H-1), 4.47-4.25 (m, 3H, H-6a, 6b and OCH₂), 4.14-3.90 (m, 2H, OCH₂ and H-2), 3.65-3.43 (m, 2H, H-5 and H-4), 3.23 (sb, 1H, OH-4), 1.92 (s, 3H, NHCOCH₃), 1.23 (s, 9H, *tert*-Butyl) and 1.20 ppm (s, 9H, *tert*-Butyl).

Example 3: Allyl 2-acetamido-2-deoxy- α -D-galactopyranoside (Allyl Tn)



Allyl Tn

Referring to **Fig. 5**, di-*O*-pivaloyl compound (Compound 2) (5.50 g, 12.80 mmol, 1.0 equiv.) in a mixture of dry dichloromethane-pyridine (126 mL, v/v 20:1) was cooled to -35°C under argon atmosphere. Trifluoromethanesulfonic anhydride (2.58 mL, 15.36 mmol, 1.2 equiv.) was then added and the mixture was stirred at this temperature. The temperature was warmed to room temperature for 2 hours. Water (12 mL) was then added into the solution. The mixture was heated, and stirred at reflux ($\approx$ 50°C)

overnight (12 hours). After reaching room temperature, the reaction mixture was diluted with dichloromethane and washed with 1M aqueous HCl several times. The organic layer was washed with H₂O, saturated NaHCO₃, and brine. The organic layer was dried over Na₂SO₄, filtered and evaporated under reduced pressure. The crude product was treated under Zemplén condition (1M sodium methoxide solution in methanol, 40 mL, pH 9). The solution was stirred at 50°C overnight. After cooling to room temperature, the solution was neutralized by addition on ion-exchange resin (Amberlite® IR 120, H⁺), filtered, washed with MeOH, and the solvent was removed under reduced pressure. Allyl T_N was isolated by precipitation in MeOH/EtOAc/Hexanes as white solid after lyophilisation (2.36 g, 9.10 mmol, 71%). R_f = 0.32; EtOAc/MeOH 4:1; **Fig. 7A & 7B**: ¹H NMR (CD₃OD, 600 MHz): δ 5.99-5.88 (m, 1H, OCH₂CH=CH₂), 5.31 (dd, 1H, *J*_{trans} = 17.3, *J*_{gem} = 1.3 Hz, OCH₂CH=CH₂), 5.17 (dd, 1H, *J*_{cis} = 10.5 Hz, OCH₂CH=CH₂), 4.86 (d, 1H, *J*_{1,2} = 3.8 Hz, *H*-1), 4.27 (dd, 1H, *J*_{2,3} = 11.0 Hz, *H*-2), 4.20 (m, 1H, OCH₂), 4.00 (m, 1H, OCH₂), 3.89 (dd, *J*_{3,4} = *J*_{4,5} = 2.6 Hz, *H*-4), 3.85-3.77 (m, 2H, *H*-3 and *H*-5), 3.72 (m, 2H, *H*-6a and *H*-6b) and 1.99 ppm (s, 3H, CH₃); ¹³C NMR (CD₃OD, 150 MHz): δ 172.5 (NHCO), 134.2 (OCH₂CH=CH₂), 116.1 (OCH₂CH=CH₂), 96.6 (C-1), 71.2 (C-3), 69.0 (C-4), 68.3 (C-5), 67.8 (OCH₂), 61.4 (C-6), 50.2 (C-2) and 21.2 ppm (CH₃). **Fig. 7C & 7D**: ESI⁺-HRMS: [M+H]⁺ calcd for C₁₁H₂₀O₆N, 262.1285; found, 262.1294.


Procedure B: Allyl 2-acetamido-2-deoxy-α-D-galactopyranoside can also be directly prepared from *N*-acetylgalactosamine (GalNAc) according to literature procedure (Feng et al., 2004: To a solution of *N*-acetylgalactosamine (442 mg, 2 mmol, 1.0 equiv.) in allyl alcohol (8 mL) at room temperature was added BF₃·Et₂O (250 μL, 2 mmol, 1.0 equiv.), and the mixture was stirred at 70 °C for 2 hours. The solution was cooled to room temperature and the solvent was removed under reduced pressure. The dry crude product was dissolved in minimum EtOH (5 mL). The desire allyl T_N product was precipitated in diisopropyl ether and isolated as white solid (417 mg, 1.60 mmol, 80%).

The **C-Allyl GalNAc** analog (**Fig. 5**) [1-(2'-Acetamido-2'-deoxy-α-D-galactopyranosyl)-2-propene] has been prepared according to literature procedure (Cipolla, et al., 2000 : 3-(2-Acetamido-3,4,6-tri-*O*-acetyl-2-deoxy-α-D-galactopyranosyl)-1-propene (Cui et al., 1998) (371 mg, 1.00 mmol, 1.0 equiv.) was treated under Zemplén condition (1M sodium methoxide solution in methanol, 5 mL, pH 8-9). The solution was stirred at room temperature for 1 h. The reaction mixture was neutralized by addition on ion-exchange resin (Amberlite® IR 120, H⁺), filtered, washed with MeOH, and the solvent was removed under reduced pressure. C-Allyl T_N was purified by chromatography on silica gel (EtOAc/MeOH 9:1 to 4:1) followed by crystallisation in EtOH as white solid (213 mg, 0.87 mmol, 87%). R_f = 0.28; EtOH 4:1; mp 230 °C (Litt. 215-217 °C, EtOAc/EtOH); According to literature NMR data : ¹H NMR (CD₃OD, 600 MHz): δ 5.81 (m,

1H, $^1\text{CH}_2\text{CH}=\text{CH}_2$), 5.08 (dd, 1H, $J_{\text{trans}} = 17.2$, $J_{\text{gem}} = 1.7$ Hz, $^1\text{CH}_2\text{CH}=\text{CH}_2$), 5.02 (dd, 1H, $J_{\text{cis}} = 10.2$ Hz, $^1\text{CH}_2\text{CH}=\text{CH}_2$), 4.22 (dd, 1H, $J = 9.3$, 5.0 Hz, $H-2'$), 4.14 (dt, 1H, $J = 10.0$, 5.0 Hz, $H-1'$), 3.91 (dd, 1H, $J = 3.0$ Hz, $H-4'$), 3.82-3.64 (m, 4H, $H-3'$, $H-5'$ and $H-6'$ ab), 2.45 (m, 1H, $H-1a$), 3.17 (m, 1H, $H-1b$) and 1.97 ppm (s, CH_3); ^{13}C NMR (CD_3OD , 150 MHz): δ 173.6 (NHCO), 136.2 ($^1\text{CH}_2\text{CH}=\text{CH}_2$), 117.1 ($^1\text{CH}_2\text{CH}=\text{CH}_2$), 72.9 ($C-1'$), 69.7 ($C-4'$), 69.5 ($C-3'$ and $C-5'$), 61.8 ($C-6'$), 52.0 ($C-2'$), 32.4 ($^1\text{CH}_2$) and 22.5 ppm (CH_3). ESI⁺-LCMS: $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{11}\text{H}_{20}\text{O}_5\text{N}$, 246.1336; found, 246.1332; CAN/ H_2O 5 to 95% 1.4 min.

The **S-Allyl GalNAc** analog (**Fig. 5**) was prepared according to literature : Knapp et al., 2002.


Example 4: Allyl 2-acetamido-4,6-O-benzylidene-2-deoxy- α -D-galactopyranoside (Compound 4)



Compound 4

Referring to **Fig. 5**, to a solution of allyl GalNAc (**Tn**) (2.35 g, 9.0 mmol, 1.0 equiv.) and benzaldehyde dimethylacetal (6.75 mL, 45.0 mmol, 5.0 equiv.) in dry DMF (20 mL) was added a catalytic amount of *p*-toluenesulfonic acid monohydrate. The mixture was stirred at room temperature. After 5 hours, the mixture was diluted with CHCl_3 and washed with a saturated aqueous solution of NaHCO_3 . The organic layer was separated and washed with water, dried over Na_2SO_4 , and concentrated to afford white solid. The benzylidene acetal (**4**) was isolated by precipitation in EtOAc/Hexanes as white solid (2.64 g, 7.56, 84%). $R_f = 0.21$; DCM/MeOH 9.0:0.5; **Fig. 8A**: ^1H NMR (CDCl_3 , 300 MHz): δ 7.59-7.46 (m, 2H, $H\text{-ar}$), 7.43-7.31 (m, 3H, $H\text{-ar}$), 5.91 (m, 1H, $\text{OCH}_2\text{CH}=\text{CH}_2$), 5.75 (d, 1H, $J_{\text{NH,H2}} = 9.0$ Hz, NH), 5.58 (s, 1H, PhCH), 5.34-5.17 (m, 2H, $\text{OCH}_2\text{CH}=\text{CH}_2$), 5.01 (d, 1H, $J_{1,2} = 3.5$ Hz, $H-1$), 4.56-4.42 (ddd, 1H, $J_{2,3} = 10.9$ Hz, $J_{2,\text{OH}} = 9.1$ Hz, $H-2$), 4.34 (dd, 1H, $J_{5,6a} = 1.5$ Hz, $J_{6a,6b} = 12.5$ Hz, $H-6a$), 4.19 (m, 2H, $H-4$ and OCH_2), 4.04 (m, 1H, dd, 1H, $J_{5,6b} = 1.6$ Hz, $J_{6a,6b} = 12.5$ Hz, $H-6b$), 4.01 (m, OCH_2), 3.86 (dd, 1H, $J_{3,4} = 10.9$ Hz, $H-3$), 3.71 (sb, 1H, $H-5$), 2.80 (d, 1H, $J_{3,\text{OH}} = 10.7$ Hz, OH-3) and 2.05 ppm (s, 3H, CH_3); **Fig. 8B & 8C**: ESI⁺-HRMS: $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{24}\text{O}_6\text{N}$, 350.1598; found, 350.1608.


Example 5: Allyl (2,3,4,6-tetra-O-benzoyl- β -D-galactopyranosyl)-(1 $\rightarrow$ 3)-2-acetamido-4,6-O-benzylidene-2-deoxy- α -D-galactopyranoside (Compound 6)



Compound 6

Referring to Fig. 5, Compound 4 (2.0 g, 5.72 mmol, 1.0 equiv.) and mercuric cyanide (2.17 g, 8.60 mmol, 1.5 equiv.) were dissolved in the mixture of anhydrous nitromethane-toluene (100 mL, 3:2, v/v) containing 4 Å molecular sieves under argon atmosphere. The mixture was stirred at room temperature for 30 min. 2,3,4,6-Tetra-O-benzoyl- α -D-galactopyranosyl bromide (Compound 5) (5.66 g, 8.58 mmol, 1.5 equiv.) was added into the mixture. The solution was stirred at 70°C for five hours, then kept stirring at room temperature overnight (8 hours). After total consumption of the starting material (Compound 4) as indicated by TLC (DCM/MeOH 9.0:0.5), the solvent was removed under reduced pressure. The residue was dissolved in EtOAc followed by filtration through a celite pad. The filtrate was successively washed with 10% aqueous potassium iodide solution, saturated sodium hydrogen carbonate solution and water, then dried over Na₂SO₄. The solvent was evaporated under reduced pressure to afford a white foam. The crude product was purified by chromatography on silica gel using a gradient of 100% hexanes to hexanes/EtOAc 1:2 to afford the desired disaccharide (Compound 6) as white solid (4.98, 5.38 mmol, 94%). mp : 110-111°C, R_f = 0.20; hexanes/EtOAc 1:2; Fig. 9A & 9B: ¹H NMR (CDCl₃, 600 MHz): δ 8.06-7.19 (m, 5H, *H*-ar), 5.98 (dd, 1H, $J_{3,4'} = 3.3$ Hz, $J_{4,5'} = 1.0$ Hz, H-4^{II}), 5.85-5.78 (m, 2H, OCH₂CH=CH₂ and H-2^{II}), 5.60 (dd, 1H, $J_{2,3'} = 10.2$ Hz, $J_{3,4'} = 3.4$ Hz, H-3^{II}), 5.48 (sb, 1H, NH), 5.23 (m, 3H, OCH₂CH=CH₂ and H-1), 4.68 (dd, 1H, $J_{5',6a'} = 6.9$ Hz, $J_{6a',6b'} = 11.4$ Hz, H-6a^I), 4.63-4.58 (m, 1H, H-2), 4.46-4.36 (m, 3H, H-4, H-5 and H-6b^{II}), 4.14-4.07 (m, 3H, H-6a, OCH₂ and H-3), 3.96 (m, 1H, OCH₂), 3.75 (m, 1H, H-6b), 3.51 (m, 1H, H-5) and 1.40 ppm (s, 3H, CH₃); ¹³C NMR (CDCl₃, 150 MHz): δ 170.0 (NHCO), 166.0, 165.5, 165.4, 165.2 (CO), 137.6-126.2 (multi, 30 C-arom), 133.2 (OCH₂CH=CH₂), 117.8 (OCH₂CH=CH₂), 102.0 (C-1^{II}), 100.9 (CPhCH), 97.3 (C-1^I), 76.1 (C-3), 75.4 (C-4), 71.7 (C-3^{II} and C-5^{II}), 70.2 (C-2^{II}), 69.1 (C-6), 68.6 (OCH₂), 68.1 (C-4^{II}), 62.9 (C-5), 62.6 (C-6^I), 48.2 (C-2) and 22.5 ppm (CH₃). Fig. 9C & 9D: ESI⁺-HRMS: [M+H]⁺ calcd for C₅₂H₅₀O₁₅N, 928.3175; found, 928.3133.

Example 6: Allyl (β -D-galactopyranosyl)-(1 $\rightarrow$ 3)-2-acetamido-2-deoxy- α -D-galactopyranoside
(Allyl TF)



Allyl TF

Referring to **Fig. 5**, a solution of compound **6** (1.12 g, 1.20 mmol, 1.0 equiv.) in 1M sodium methoxide in methanol (12 mL, pH 8-9) was stirred at room temperature until consumption of starting material. After 1 h 30 min, the solution was neutralized by the addition of ion-exchange resin (Amberlite IR 120, H⁺),
 10 filtered, washed with MeOH, and the solution was suspended with silica gel, filtered, and the solvent removed under reduced pressure. The silica gel was washed with 100% EtOAc several times followed by washing with second solution (EtOAc/MeOH/H₂O 1:1:0.1). The combined filtrate was evaporated under reduced pressure to afford the intermediate (Compound **7**) as white solid. *R_f* = 0.20; CHCl₃/MeOH/H₂O 11:6:1; **Fig. 10A & 10B**: ¹H NMR (D₂O, 600 MHz): δ 7.47-7.39 (m, 2H, *H*-ar), 7.37-7.26 (m, 3H, *H*-ar),
 15 5.82 (m, 1H, OCH₂CH=CH₂), 5.61 (s, 1H, PhCH), 5.20-5.09 (m, 2H, OCH₂CH=CH₂), 4.88 (d, 1H, *J*_{1,2} = 3.4 Hz, *H*-1), 4.47 (dd, *H*-4), 4.37-4.25 (m, 2H, *H*-2 and *H*-1^{II}), 4.13-3.98 (m, 4H, *H*-3, *H*-6a, *H*-6b, OCH₂), 3.96-3.88 (m, 1H, *H*-5), 4.56-4.42 (ddd, 1H, *J*_{2,3} = 10.9 Hz, *J*_{2,OH} = 9.1 Hz, *H*-2), 4.34 (dd, 1H, *J*_{5,6a} = 1.5 Hz, *J*_{6a,6b} = 12.5 Hz, *H*-6a), 4.19 (m, 2H, *H*-4 and OCH₂), 4.04 (m, 1H, dd, 1H, *J*_{5,6b} = 1.6 Hz, *J*_{6a,6b} = 12.5 Hz, *H*-6b), 4.01 (m, OCH₂), 3.86 (dd, 1H, *J*_{3,4} = 10.9 Hz, *H*-3), 3.71 (sb, 1H, *H*-5), 2.80 (d,
 20 1H, *J*_{3,OH} = 10.7 Hz, OH-3) and 2.05 ppm (s, 3H, CH₃); 3.83 (s, 1H, OCH₂), 3.72 (d, 1H, *J*_{3',4'} = *J*_{4',5'} = 3.2 Hz, *H*-4^{II}), 3.65-3.55 (m, 2H, *H*-6a,b), 3.49, (m, 1H, *H*-5^{II}), 3.43 (dd, 1H, *J*_{2',3'} = 10.0 Hz, *J*_{3',4'} = 3.3 Hz, *H*-3^{II}), 3.33-3.24 (m, 1H, *H*-2^{II}) and 1.86 ppm (s, 3H, CH₃); ¹³C NMR (D₂O, 150 MHz): δ 174.6 (NHCO), 136.8 (*C*-arom), 133.6 (OCH₂CH=CH₂), 129.9, 128.7, 126.5 (*C*-arom), 118.0 (OCH₂CH=CH₂), 104.9 (*C*-1^{II}), 101.3 (CHPh), 97.0 (*C*-1), 76.0 (*C*-4), 75.0 (*C*-3), 75.0 (*C*-5^{II}), 72.4 (*C*-3^{II}), 70.4 (*C*-2^{II}), 69.0 (*C*-6),
 25 68.7 (OCH₂CH=CH₂), 68.6 (*C*-4^{II}), 63.0 (*C*-5), 61.0 (*C*-6^{II}), 48.6 (*C*-2) and 22.0 ppm (CH₃). *R_f* = 0.38; EtOAc/MeOH/H₂O 7:3:0.1; *R_f* = 0.46; ACN/MeOH/H₂O 7:2:1. **Fig. 10C & 10D**: ESI⁺-HRMS: [M+H]⁺ calcd for C₂₄H₃₄O₁₁N, 512.2126; found, 512.2119.


The white solid intermediate was then dissolved in 10 mL of 60% aqueous acetic acid and the resulting
 30 solution was stirred at 60°C for 1.5 hours. The solvent was removed under reduced pressure, and the residue was lyophilized to afford the final allyl TF as white solid (427 mg, 1.0 mmol, 84%). mp = 230-

232 °C; $R_f = 0.53$; $\text{CHCl}_3/\text{MeOH}/\text{H}_2\text{O}$ 11:6:1; **Fig. 11A & 11B:** ^1H NMR (D_2O , 600 MHz): δ 5.80 (m, 1H, $\text{OCH}_2\text{CH}=\text{CH}_2$), 5.19 (dd, 1H, $J_{\text{trans}} = 17.3$ Hz, $\text{OCH}_2\text{CH}=\text{CH}_2$), 5.09 (dd, 1H, $J_{\text{cis}} = 10.4$ Hz, $\text{OCH}_2\text{CH}=\text{CH}_2$), 4.77 (d, 1H, $J_{1,2} = 3.7$ Hz, $H-1$), 4.29 (d, 1H, $J_{1,2} = 3.7$ Hz, $H-1$), 4.29 (d, 1H, $J_{1,2} = 7.8$ Hz, $H-1^{\text{II}}$), 4.16 (dd, 1H, $J_{2,3} = 11.2$ Hz, $J_{1,2} = 3.7$ Hz, $H-2$), 4.08-4.01 (m, 2H, $H-4$ and OCH_2), 3.92-3.82 (m, 3H, $H-3$, $H-5$ and OCH_2), 3.73 (dd, 1H, $H-4^{\text{II}}$), 3.63-3.52 (m, 4H, $H-6\text{a,b}$ and $H-6'\text{a,b}$), 3.47 (m, 2H, $H-3^{\text{II}}$ and $H-5^{\text{II}}$), 3.39 (dd, 1H, $J_{2',3'} = 10.0$ Hz, $J_{1',2'} = 7.7$ Hz, $H-2^{\text{II}}$) and 1.85 ppm (s, 3H, CH_3); ^{13}C NMR (150 MHz, CDCl_3): δ 174.6 (NHCO), 133.7 ($\text{OCH}_2\text{CH}=\text{CH}_2$), 117.9 ($\text{OCH}_2\text{CH}=\text{CH}_2$), 104.7 ($\text{C}-1^{\text{II}}$), 96.4 ($\text{C}-1$), 77.2 ($\text{C}-3$), 75.0 ($\text{C}-5^{\text{II}}$), 72.5 ($\text{C}-3^{\text{II}}$), 70.7 ($\text{C}-5$), 70.6 ($\text{C}-2^{\text{II}}$), 68.8 ($\text{C}-4$), 68.6 ($\text{C}-4^{\text{II}}$), 68.4 (OCH_2), 61.2 ($\text{C}-6^{\text{II}}$), 61.0 ($\text{C}-6$), 48.6 ($\text{C}-2$) and 22.0 ppm (CH_3). **Fig. 11C & 11D:** ESI $^+$ -HRMS: $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{29}\text{O}_{11}\text{NNa}$, 446.1633; found, 446.1613.

Example 7: Purification of Tetanus Toxoid monomer

Tetanus toxoid (TT) monomer was obtained by gel filtration chromatography before conjugation. One milliliter of a liquid preparation containing 4.5 mg/ml protein (as determined by the modified Lowry protein assay) was loaded onto a XK16-100 column filled with Superdex®200 Prep Grade (GE Healthcare Life Sciences, Uppsala, Sweden) equilibrated in PBS (20 mM NaHPO_4 [pH 7.2], 150 mM NaCl) and eluted with the same buffer. The protein eluted from the column in two peaks: the earlier-eluting peak contained oligomerized toxoid, and the later-eluting peak, corresponding to a Mr of 150,000, contained TT monomer. Fractions corresponding to the later (monomer) peak were pooled, desalted against deionized water, concentrated using a Centricon® Plus-70 centrifugal filter device (30K Ultracel PL membrane; Millipore, Billerica, MA), and then lyophilized.

Example 8: Tn-Tetanus Toxoid conjugate (Tn-TT conjugate)




Tn-TT conjugate

Procedure A: To a solution of allyl Tn (0.15 mg, 0.56 μmol , 18.0 equiv.) in 500 μL PBS was added 1.5 mL of PBS containing 4.7 mg (0.03 μmol) of tetanus toxoid in a quartz cell. The solution was irradiated at 254 nm for 7 hours after degazing under argon atmosphere. The solution was dialyzed for 24

hours against several washes of doubly distilled water, and lyophilized to give the Tn-TT-A conjugate as white solid (2.0 mg, 43%). **Procedure B:** Under the same thiol-ene click reaction, a catalytic amount of DMPA in acetonitrile (100 μ L) was added into the solution of allyl Tn (3 mg, 10.8 μ mol, 180.0 equiv.) containing Tetanus toxoid in PBS (2 mL, 4.5 mg/mL, 0.06 μ mol). The solution was irradiated at 365 nm for 15 min. The solution was then dialyzed for 24 hours, and lyophilized to give Tn-TT-B conjugate as white solid (5.4 mg, 61%).

Example 9: TF-Tetanus Toxoid conjugate (TF-TT conjugate)



TF-TT conjugate

Procedure A: To a solution of allyl TF (0.24 mg, 0.56 μ mol, 18.0 equiv.) in 500 μ L PBS was added 1.5 mL of PBS containing 4.7 mg (0.03 μ mol) of tetanus toxoid in a quartz cell. The solution was irradiated at 254 nm for 7 hours after degazing under argon atmosphere. The solution was then dialyzed for 24 hours, and lyophilized to give TF-TT-A conjugate as white solid (2.1 mg, 45%). **Procedure B:** Under the same thiol-ene click reaction, a catalytic amount of DMPA in acetonitrile (100 μ L) was added into the solution of allyl TF (5 mg, 10.8 μ mol, 180.0 equiv.) with Tetanus toxoid in PBS (2 mL, 4.5 mg/mL, 0.06 μ mol). The solution was irradiated at 365 nm for 15 min. The solution was then dialyzed for 24 hours, and lyophilized to give TF-TT-B conjugate as white solid (5.3 mg, 60%).

Example 10: Characterization of Tn-TT and TF-TT conjugates

The tetanus toxoid (TT) alone and the vaccine conjugates (Tn-TT-A; Tn-TT-B; TF-TT-A; and TF-TT-B) were analyzed by SDS gel electrophoresis, colorimetric analyses for the presence of sugar content (Dubois test), thiol content by the Ellmann test, Dynamic Light Scattering (DLS) (Fig. 12), and reactivity with the known mouse monoclonal antibody JAA-F11 using double radial immunodiffusion. The SDS gel electrophoresis results clearly indicated the monomeric form of the conjugate that also stained positive for the presence of the carbohydrate antigens (Tn and TF). The colorimetric analyses (Dubois test) confirmed the presence of the carbohydrate antigens (10% by weight) and that there were no residual free thiol groups on the carrier protein following conjugation. The double radial immunodiffusion analyses revealed

a precipitation band, clearly indicating the cross-reactivity of the new TF-TT conjugate with the anti-TF monoclonal antibody JAA-F11, thus confirming the presence of the immunogenic carbohydrate antigen (TF) on the TF-TT conjugate and that no precipitation band was observed with the carrier protein (tetanus toxoid) alone.

HPLC analysis of the conjugates. HPLC analysis of the glycoconjugate preparations was done by size exclusion chromatography. The chromatographic separation was performed with three 8- by 300-mm Shodex OHpak gel filtration columns connected in series (two SB-804 and one SB-803) preceded by a SB-807G guard column (Showa Denko). The glycoconjugate immunogens were eluted with 0.1 M NaNO₃ at a flow rate of 0.4 mL/min using a Knauer Smartline system equipped with a differential refractometer (RI) detector model 2300 and a UV detector model 2600 at wavelength of 280 nm. The conjugate preparation (8-mg/mL solution in the mobile phase) was injected using a 50-μL injection loop. In selected experiments, the fractions eluting at the void volume, which correspond to the conjugate fractions, were pooled, dialyzed against water Spectra/Por; Molecular weight cut-off (MWCO), 12,000 to 14,000 [Spectrum Laboratories]), and lyophilized. This corresponds to the 2:1 fractionated conjugate.

Example 11: Effect of the conjugation of O-Allyl-sugars by the thiol-ene photoreaction on the free thiol of peptide dTT831-844-Cys-βAla

The peptide dTT831-844-Cys-βAla (Tetanus Toxin (831-844); MW: 1813, SEQ ID NO: 2) was solubilized in water at a concentration of 3.7e-3 M and (500 uL) of peptide was stirred at room temperature with (6 uL, 0.1M, 3.0 equiv.) O-Allyl-Tn or O-Allyl-TF and water-soluble catalyst AAPH (2,2'-Azobis(2-methylpropionitrile, 23 uL, 0.025M, 3.0 equiv. in a quartz cuvette (10x10 mm path length, Fisher) placed between two hand held UV365 nm lamp (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH. The solution was sampled at 0, 45, 90 and 135 minutes and free thiol concentration was measured by the Ellman test using a cysteine standard curve. Fig. 13 shows that the free thiol concentration is reduced in a time dependent manner as a result of the photo thiol-ene reaction.

Example 12: Effect of cysteine positioning/accessibility in dTT831-844-Cys-βAla, and the type of activator employed, on conjugation of O-Allyl-Tn and immunoreactivity of the thiol-ene photoreaction product

The peptide dTT831-844 was synthesized with an N-terminal or C-terminal cysteine and solubilized in water at a concentration of 2 mM (1 mL) that was stirred at room temperature with 200 μ L, 0.1 M, 10 equiv.) of O-Allyl-Tn and a catalytic amount (0.2 equiv.) of activator AAPH in a quartz cuvette (10x10 mm path length, Fisher) placed between two hand held UV365nm lamp (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of activator and allowed to react for 60 min. The free thiol concentration was measured by the Ellman test using a cysteine standard curve at time 0 and 60 min. The fold reduction in free thiol correspond to the thiol concentration at $t=0$ / $t=60$.

The results are shown in Fig. 14, which presents the fold reduction in free thiol concentration for dTT831-844 peptides having an N-terminal (“SH-peptide”) or C-terminal (“Peptide-SH”) cysteine, and using either a water-soluble (AAPH) or water-insoluble activator (DMPA), in the thiol-ene photoreaction conjugation.

The immunoreactivity of the thiol-ene photoreaction product was measured by an Enzyme-linked lectin assay (ELLA). 1 μ g of peptide in 100 μ L PBS pH 7.4 was allowed to adsorb to 96-well plates (Maxisorp™, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with PBS-T+1% BSA blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 μ L of a 1/100 dilution of the lectin *Vicia Villosa* (anti-Tn VVA) or *Peanut agglutinin* (anti-TF PNA) coupled to the horseradish peroxidase (VVA-hrp, PNA-hrp, EY LABs). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells were washed 4 times with PBS-T and 100 μ L of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 μ L of 0.5N sulfuric acid and the plate was read in a plate reader at OD 450nm (Biotek EL808).

The results are shown in Fig. 15, which presents lectin reactivity to the different Tn-peptides (i.e., unconjugated: “pep-Cys” and “Cys-pep”; Tn-conjugated: “Tn-pep” and “pep-Tn”; “unc”: uncoated plate) using either a water-soluble (AAPH) or water-insoluble activator (DMPA). Interestingly, TT peptides with a C-terminal cysteine were specifically recognized by the anti-Tn lectin VVA, and the conjugation reaction was most effective with the water-soluble activator AAPH. None of the Tn glycoconjugates were recognized by the anti-TF lectin (peanut lectin, PNA) used as a negative control.

Example 13: Effect of wavelength on thiol-ene photoreaction conjugation of peptide dTT831-844-Cys-βAla to O-Allyl-Tn

The peptide dTT831-844-Cys-βAla (MW: 1813, SEQ ID NO: 2) was stirred at room temperature with O-Allyl-Tn 0.55 mM in 1 mL water and AAPH (1.44 mg, 5.5 nmol, 10.0 equiv.) or (0.3 mg, 1.1 nmol, 2.0 equiv.) DMPA in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at either 365nm or 355nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH and turned off to stop the reaction after (60 min). The peptide was also conjugated to O-Allyl-Tn in the absence of activator and at UV254nm. The free thiol concentration was measured by the Ellman test using a cysteine standard curve. Fig. 16 shows the decrease in free thiol over time (minutes) for three different conjugation conditions (“355nm+AAPH”; “365nm+DMPA”; and “254nm”).

The immunoreactivity of the thiol-ene photoreaction product was measured by Enzyme-linked lectin assay (ELLA). The indicated quantity of peptide in 100 μL PBS pH 7.4 was allowed to adsorb to 96-well plates (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and the wells filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 μL of a 1/100 dilution of the lectin *Vicia Villosa* coupled to the horseradish peroxidase (VVA-hrp, EY LABs). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells were washed 4 times with PBS-T and 100 μL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 μL of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

The ELLA results are presented in Fig. 17, which shows VVA lectin reactivity to various quantities of Tn-peptides conjugated by thiol-ene photoreaction at 365nm in the presence of AAPH or DMPAP, or at 254nm in absence of activator. The conjugation products were detected by the lectin, while the unconjugated peptide (“peptide”) or O-Allyl-Tn alone (“Tn”) were unreactive. Interestingly, Fig. 17 shows that conjugation at 254nm in the absence of activator generated a glycoconjugate product that was more immunoreactive than the ones generated by 365nm or 355nm in the presence of activator.

Example 14: Conjugation and reactivity of C-Allyl and O-Allyl saccharides to peptide dTT831-844-Cys-βAla by the thiol-ene photoreaction

The dTT831-844-Cys-βAla peptide (MW: 1813) was solubilized in water at a concentration of 0.55 mM and 1 mL of it was stirred at room temperature with O-Allyl-Tn, O-Allyl-TF, or C-Allyl-Tn (100 μL, 11 mM, 2.0 equiv) and AAPH (1.44 mg, 5.5 nmol, 10.0 equiv.) in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH. The solution was sampled at the indicated time and free thiol concentration was measured by the Ellman test using a cysteine standard curve. **Fig. 18**

shows the decrease in free thiol over time (minutes) for the different allyl saccharides tested: O-Allyl-Tn ("Tn"), O-Allyl-TF ("TF"), or C-Allyl-Tn ("cTn").

The immunoreactivity of the thiol-ene photoreaction product was measured by enzyme-linked lectin assay and enzyme-linked immunosorbent assay. The indicated quantity of peptide conjugate in 100 μL PBS pH 7.4 was allowed to adsorb to 96-well plates (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with PBS-T+1% BSA blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 μL of a 1/100 dilution of the lectin *Vicia Villosa*, or *Peanut agglutinin* coupled to the horseradish peroxidase (VVA-hrp, PNA-hrp, EY LABs), or 0.1 μg/ml of TF-specific purified murine monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells containing JAAF11 were washed with PBS-T then further incubated for 60 minutes with secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). Wells were then washed 4 times with PBS-T and 100 μL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 μL of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 19 shows that only the peptide conjugate generated with the O-Allyl-Tn (but not with the C-Allyl-Tn) was recognized by the anti-Tn lectin VVA. These results suggest that, while being conjugated to the peptide, the C-Allyl-Tn does not have a conformation that allows binding to the lectin.

Example 15: Conjugation of O-Allyl-TF to chemically reduced dTT by thiol-ene photoreaction

Detoxified tetanus toxoid (dTT) was dialysed in PBS pH 7.4 (MWCO 2,000) then incubated with 1000 eq DTT. After a 2 hours incubation at RT and rotation, the reduced protein solution was washed in PBS pH 7.4 by centrifugal filtration (MWCO 10,000, Amicon). The reduced protein (1 mL, 3.51 mg/mL, 0.02 nmol) was then stirred with O-Allyl-TF (30.0 equiv.) and AAPH (2.0 equiv.) in a final volume of 1.0 mL

PBS in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction then stopped after 2h.

- 5 The free thiol concentration of the reduced dTT generated by the DTT treatment compared to native dTT was measured on buffer exchanged dTT sample by an Ellman test using a cysteine standard curve and the protein concentration was measured by a Bradford assay using a BSA standard curve.

To measure the immunoreactivity of the dTT samples, 1 µg of protein sample was diluted into 100 µL of
10 PBS pH 7.4 and allowed to adsorb to the well of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of 0.1 µg/mL of TF-specific purified murine monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the plate was washed with PBS-T then further
15 incubated for 60 minutes with 100 µL of 1/1000 dilution of the secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). Wells were then washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 µL of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

20

Fig. 20 shows the immunoreactivity dTT-TF to JAAF11 (“dTT(DTT)-TF”), while the reduced dTT (“dTT(DTT)”) or native dTT (data not shown) were unreactive. This indicates that the O-Allyl-TF was indeed conjugated to dTT by the thiol-ene photo reaction in an immunoreactive conformation.

25 **Example 16: Conjugation of O-Allyl-Tn to chemically reduced dTT by thiol-ene photoreaction**

dTT was dialysed in PBS pH 7.4 (MWCO 30,000) then 0.5 mL (5.8 mg/mL) was incubated with 50 or 500 eq DTT at RT under rotation. After 2h, the reduced protein solution was washed in PBS pH 7.4 by centrifugal filtration (MWCO 10,000, Amicon). The protein reduced with 500 eq of DTT (70 uL, 0.5
30 mmol/mL) was then stirred with 0.1M O-Allyl-TF (60.0 equiv.) and 0.1M AAPH (4.0 equiv.) in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction then stopped after 45 minutes. To verify that the free thiols of the reduced dTT have formed a reduction-resistant thio-ether bond as a result of conjugation, or re-

oxidized into di-sulfide bond, the conjugated product was treated with 100 Eq of DTT for 1h then washed by centrifugal filtration, and then the thiol content was measured before and after the reducing treatment. The free thiol dosage was measured by an Ellman test using a cysteine standard curve. The protein concentration was measured by a Bradford assay using a BSA standard curve.

Fig. 21 shows the molar ratio of free thiols in dTT after the different conjugation conditions. The results indicate that the dTT protein was reduced in a dose-dependent manner and reached 9 free thiols upon treatment with 500 Eq DTT (out of a maximal theoretical number of 10 cysteines in dTT). Conjugation of O-Allyl-Tn or O-Allyl-TF by the thiol-ene photo-reaction lowered the free thiol to at or below the level of the non-reduced dTT. The resistance of the dTT-Tn conjugate to reduction confirmed the presence of a reduction-resistant thio-ether bond that formed by the thiol-ene photo reaction.

To measure the immunoreactivity of the dTT samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin *Vicia Villosa* or *Peanut Agglutinin* coupled to the horseradish peroxidase (VVA-hrp, PNA-hrp EY LABs), or 0.1 µg/mL of TF-specific purified murine monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells with JAAF11 were washed with PBS-T then further incubated for 60 minutes with 100 µL of 1/1000 dilution of the secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the plate washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 µL of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 22 shows the high reactivity of the dTT-Tn conjugate to the lectin VVA compared to the unconjugated dTT.

The native dTT and dTT-Tn conjugate were analyzed by SDS-PAGE electrophoresis on a 10% polyacrylamide gel, then stained with Coomassie blue. The gel electrophoresis results showed that the dTT-Tn conjugate migrated at higher molecular weight than the native non-conjugated dTT (data not shown).

Example 17: Effect of AAPH concentration on the conjugation of O-Allyl-Tn to chemically thiolated dTT by thiol-ene photoreactivity

dTT was dialysed in PBS pH 8 (MWCO 30,000) then 100 μ L (0.8 mg) was stirred with 200 Eq/protein of a solution of 0.1 M 2-imminothiolane hydrochloride in water (Sigma), 200 Eq/protein of 0.1 M O-Allyl-TF in water, and 0.8 Eq/protein or approximately 450 Eq/protein AAPH in water in a final volume of 124 μ L in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. Samples were retrieved during the photoreaction at 15 minutes and at the completion of the reaction at 45 minutes. The conjugation product was buffer exchanged to PBS pH 7.4 by size exclusion chromatography using SephadexTM G25-filled mini-spin column. The free thiol concentration was measured by Ellman assay using a cysteine standard curve. The protein concentration was measured by a Bradford assay using BSA as standard.

Fig. 23 shows that the ratio of free thiol in the dTT samples subsequent to chemical thiolation with 2-imminothiolane and thiol-ene photoreaction in the absence of O-Allyl-Tn reaches 25-30, while the presence of O-Allyl-Tn reduces the thiol ratio of dTT to approximately 20 and 5 in a dose-dependent manner respective to the AAPH concentration, suggesting that a thio-ether bond was formed by the thiol-ene photo-reaction.

To measure the immunoreactivity of the dTT samples, 1 μ g of protein sample was diluted into 100 μ L of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-TweenTM 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 μ L of a 1/100 dilution of the lectin *Vicia Villosa* coupled to the horseradish peroxidase (VVA-hrp, EY LABs). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the plate washed 4 times with PBS-T and 100 μ L of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 μ L of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 24 shows the reactivity of dTT samples to the lectin VVA-hrp. Higher reactivity of Tn conjugates compared to native unconjugated dTT was observed. Further, the reactivity correlated with the duration of the photo-reaction and the amount of AAPH activator used.

The dTT conjugates (5 µg) were analyzed by SDS-PAGE electrophoresis on a 10% polyacrylamide gel either stained with Coomassie blue or transferred to a membrane (Immobilon-P, Merck Millipore) for western blot analysis. The membrane was blocked in PBS-T + 1% BSA for 1h, washed with PBS-T, then incubated with a 1/100 dilution of the lectin *Vicia Villosa* coupled to the horseradish peroxidase (VVA-hrp, EY LABs) in PBS-T for 1h at RT or overnight at 4°C. The membrane was then extensively washed with PBS-T and the bound VVA-hrp was detected with the horseradish peroxidase chromogenic substrate 4-Chloro 1-Naphtol (0.3 mg/mL in PBS containing 0.03% hydrogen peroxide, Sigma). The results from both the Coomassie-stained SDS-PAGE and corresponding Western blot stained with VVA-hrp, revealed that the dTT-Tn migrated at a higher molecular weight than the native dTT, and was specifically reactive to VVA, indicating that the product is indeed conjugated to Tn (data not shown).

Example 18: Conjugation of O-Allyl-Tn and O-Allyl-TF to chemically reduced BSA by thiol-ene photoreaction

BSA (7 mg in 1 mL PBS, pH 7.4) was incubated with 600 eq DTT for 2 h then washed in PBS pH 7.4 by centrifugal filtration (MWCO 10,000, Amicon). The reduced protein was then conjugated to O-Allyl-Tn or to O-Allyl-TF by the thiol-ene photoreaction. Reduced BSA (600 µL, 5.3 mg/mL, 0.047 nmol) was stirred with O-Allyl-sugar (60 eq) and AAPH (4.0 equiv.) and PBS pH 7.4 (720 µL) in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. After 45 min., the product was washed with PBS pH 7.4 by centrifugal filtration (MWCO 10,000, Amicon) to remove any un-reacted O-Allyl and reagents. The free thiol concentration of the reduced BSA was measured by Ellman assay using a cysteine standard curve, and the protein concentration by a Bradford assay using BSA as standard.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin *Vicia Villosa*, or *Peanut agglutinin* coupled to the horseradish peroxidase (VVA-hrp, PNA-hrp, EY LABs) or 0.1 µg/mL of TF-specific purified murine monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells containing JAAF11 were washed with PBS-T then further incubated for 60 minutes with secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). Wells were then washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well.

The reaction was stopped by the addition of 100 μ L of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 25 shows that the BSA-TF and BSA-Tn generated under these conditions were highly reactive and specific to anti-TF JAAF11 monoclonal antibody and for the anti-Tn lectin VVA, respectively.

The BSA-Tn and BSA-TF conjugates (5 μ g) were analyzed by SDS-PAGE electrophoresis on a 10% polyacrylamide gel stained with Coomassie blue. The reduction of BSA under the conditions used generated a molar ratio of 19.29 thiols as measured by the Ellman test. Following conjugation with either O-Allyl-sugars, the Ellman's test was negative, suggesting that the free thiols were chemically coupled to the O-Allyl sugars by the thiol-ene photoreaction (data not shown). The Coomassie-stained SDS-PAGE gel showed both BSA-conjugates migrated at the expected molecular weight of BSA, and to higher molecular weight species (data not shown).

Example 19: Conjugation of O-Allyl-Tn to chemically reduced BSA by thiol-ene photoreaction

BSA (500 μ L, 7 mg/mL) was incubated with 400, 200 or 1000 Eq of DTT (0.5 M) for 1h at RT and rotation. The reduced BSA solution was then dialysed overnight in PBS pH 7.4. The free thiol concentration of the reduced BSA was then measured by Ellman assay using a cysteine standard curve, and the protein concentration by a Bradford assay using BSA as standard, from which the molar ratio of free thiol/BSA was calculated.

Fig. 26 shows a correlation between the amount of free thiol generated and the amount of the reducing agent DTT, which reaches about 10 thiol/BSA when the reduction was conducted with 1000 Eq DTT under the conditions employed.

The reduced protein was then conjugated to various ratios of O-Allyl-Tn. Reduced BSA (5 nmols) was stirred with 100 Eq of O-Allyl-Tn/reduced BSA or 100 Eq of O-Allyl-Tn/BSA-thiol with AAPH and PBS pH 7.4 in a final volume of 500 μ L in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. After 45 minutes, the product was washed with PBS pH 7.4 by centrifugal filtration (MWCO 10,000, Amicon) to remove unreacted O-Allyl and reagents.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin *Vicia Villosa*, coupled to the horseradish peroxidase (VVA-hrp, EY LABs) or 0.1 µg/mL of TF-specific purified murine monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells containing JAAF11 were washed with PBS-T then further incubated for 60 minutes with secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the plate was washed 4 times with PBS-T and 100ul of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 µL of 0.5N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 27 shows the reactivity of the various BSA-Tn conjugates to the lectin VVA and to the anti-TF monoclonal antibody JAAF11 (+goat anti-mouse IgG-hrp conjugate). Two of the BSA-conjugates, conjugated at the highest ratio of O-Allyl, were found to be highly reactive to VVA specifically, as a result of the thiol-ene photoreaction.

The BSA-Tn conjugates (5 µg) were analyzed by SDS-PAGE electrophoresis on a 10% polyacrylamide gel either stained with Coomassie blue or for glycoproteins according to the manufacturer's instructions (Pierce). The results showed that the BSA conjugates that migrated at a high molecular weight were positively stained for glycan, while the native unconjugated BSA was not (data not shown).

Example 19: Conjugation of O-Allyl-TF to chemically thiolated BSA by thiol-ene photoreaction

BSA (1 mL, 0.088 µmol, 5.89 mg/mL, PBS pH 8.0) was incubated with 5, 10, or 20 Eq of 2-iminothiolane (5, 10, 20 µL, 0.1 mmol/mL) for 1 h. Thiolated BSA was then washed by centrifugal filtration (MWCO 10,000, Amicon) to remove unreacted reagents before conjugating to O-Allyl-TF by thiol-ene photoconjugation. Thiolated BSA (440; 480; 575 µL, 6.8; 6.4; 5.2 mg/mL) was stirred with 3 eq/thiol of O-Allyl-Tn and AAPH (0.2 Eq/thiol) and PBS pH 7.4 in a final volume of 1 mL in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. After 1 h, the product was washed with PBS pH 7.4 by centrifugal filtration (MWCO 10,000, Amicon) to remove un-reacted O-Allyl and other reagents. The free

thiol concentrations of the BSA samples were measured by Ellman assay using a cysteine standard curve, and the protein concentration by a Bradford assay using BSA as standard.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin peanut agglutinin coupled to the horseradish peroxidase (PNA-hrp, EY Labs), or 0.1 µg/mL of TF-specific purified murine monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells containing JAAF11 were washed with PBS-T then further incubated for 60 minutes with secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). Wells were then washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each wells. The reaction was stopped by the addition of 100 µL of 0.5 N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

In Fig. 28, the dual axis graph shows the reactivity of the BSA-TF conjugates to the anti-TF lectin PNA or mAb JAAF11 in relation to the thiol ratio of the respective conjugates measured before TF conjugation. These results show that JAAF11 reactivity is detected with the BSA that was the most thiolated.

Example 20: Conjugation of O-Allyl-Tn to chemically thiolated BSA by thiol-ene photoreaction

BSA (250 µL, 4.3 mg/mL) was incubated with 200 eq of 0.1 M 2-imminothiolane (Sigma) in the presence or the absence of 200 Eq of 0.1 M O-Allyl-Tn and approximately 450 Eq AAPH and PBS pH 8 in a final volume of 316 µL in a quartz cuvette (10x10mm path length, Fisher) placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. Samples were taken at the indicated time during the reaction. The reaction was stopped after 45 minutes after which the product was buffer exchanged to PBS pH 7.4 by gel filtration chromatography using Sephadex™ G25 mini spin column. The free thiol concentration of the BSA samples were measured by Ellman assay using a cysteine standard curve, and the protein concentration by a Bradford assay using BSA as standard.

Fig. 29 shows a time-dependent increase of free thiol function upon reaction of the BSA with 2-imminothiolane alone or in the presence of 365nm light, while the addition of AAPH in the presence or the absence of O-Allyl-Tn in the photoreaction prevent the formation of detectable free thiols.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin *Vicia Villosa* coupled to the horseradish peroxidase (VVA-hrp, EY Labs). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells were then washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each wells. The reaction was stopped by the addition of 100 µL of 0.5 N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

In **Fig. 30**, the reactivity of the BSA conjugates to the lectin VVA indicates that only the BSA generated by the thiol-ene photoreaction in the presence of O-Allyl-Tn is detected by the VVA, while the product generated in the absence of O-Allyl-Tn is negative in the ELLA. These results suggest that the AAPH and UV365nm light catalyse the re-oxidation of the free disulfide into di-sulfide in the absence of O-Allyl-sugar, which competes with the formation of thio-ether bond in the presence of O-Allyl-Tn.

Example 21: Conjugation of O-Allyl-Tn to chemically thiolated BSA by thiol-ene photoreaction

BSA (250 µL, 4.3 mg/mL) was incubated with 200 Eq of 0.1 M 2-imminothiolane (Sigma), 200 Eq of 0.1 M O-Allyl-Tn, and either 0.8 Eq of AAPH or approximately 450 Eq AAPH and PBS pH 8 in a final volume of 316 µL in a quartz cuvette (10x10mm path length, Fisher). The effect of the anti-oxidant Vitamin C (approx. 5 mM) on conjugation was tested in the condition containing a high amount of AAPH. The cuvettes were placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. Samples were taken at the indicated time during the reaction. The reaction was stopped after 45 minutes after which the product was buffer exchanged to PBS pH 7.4 by gel filtration chromatography using Sephadex™ G25 mini spin column. The free thiol concentration of the conjugated BSA samples was measured by Ellman assay using a cysteine standard curve, and the protein concentration by a Bradford assay using BSA as standard.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin *Vicia Villosa* coupled to the horseradish peroxidase (VVA-hrp, EY Labs). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells were then washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 µL of 0.5 N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 31 having a dual Y-axis shows an increase in reactivity of the conjugation product to the lectin VVA as a function of time, and a corresponding decrease in free-thiol as a result of the thiol-ene photoconjugation of the O-Allyl-Tn to BSA. These results suggest that the reaction was more effective at higher concentrations of AAPH. Product generated in the presence of vitamin C was not reactive to the VVA lectin (not shown).

The BSA-Tn conjugates (5 µg) were analyzed by SDS-PAGE electrophoresis on a 10% polyacrylamide gel stained with Coomassie blue. The results showed a higher amount of high molecular weight species in the reaction at high AAPH than at lower AAPH. Vitamin C prevented the formation of high molecular weight species (data not shown).

Example 22: Conjugation of O-Allyl-Tn to BSA by thiol-ene photoreaction in function of protein thiolation

BSA (93 µL, 11.7 mg/mL) was incubated with 200 Eq of 0.1 M 2-imminothiolane (Sigma), 200 Eq of 0.1 M O-Allyl-Tn and either 0.08 Eq, 0.8 Eq or approximately 450 Eq AAPH and PBS pH 8 in a final volume of 316 µL in a quartz cuvette (10x10mm path length, Fisher). In some samples, the 2-imminothiolane and/or the AAPH was omitted and replaced by PBS pH 8. The cuvette was placed between two hand held UV lamps at 365nm or at 254nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. The reaction was stopped after 45 minutes after which the product was buffer exchanged to PBS pH 7.4 by gel filtration chromatography using Sephadex™ G25 mini spin column. The protein concentration was measured by a Bradford assay using BSA as standard.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 1/100 dilution of the lectin *Vicia Villosa* coupled to the horseradish peroxidase (VVA-hrp, EY Labs). After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells were then washed 4 times with PBS-T and 100 µL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each well. The reaction was stopped by the addition of 100 µL of 0.5 N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 32 shows the minimal reactivity of product to VVA when the conjugation is performed in absence of protein thiolation or when the thiolated protein is conjugated in the absence of activator AAPH. However, a VVA reactive product is generated by UV254nm in the absence of protein thiolation and AAPH.

Example 23: Absence of BSA-TF reactivity at high conjugation ratio

BSA (93 µL, 11.7 mg/mL) was incubated in the presence or absence of 200 Eq of 0.1 M 2-iminothiolane (Sigma), 200 Eq of 0.1 M O-Allyl-TF and 0.8 Eq or approximately 450 Eq AAPH and PBS pH 8 in a final volume of 316 µL in a quartz cuvette (10x10mm path length, Fisher) on the bench at RT or placed between two hand held UV lamps at 365nm (0.16 amps, VWR) at a distance of 2-5 cm from the cuvette. The UV was turned on immediately following the addition of AAPH to initiate the reaction. The reaction was stopped after 45 minutes after which the product was buffer exchanged to PBS pH 7.4 by gel filtration chromatography using Sephadex™ G25 mini spin column. The protein concentration was measured by a Bradford assay using BSA as a standard. The conjugated galactose of the TF epitope was measured by the method of Dubois using a galactose standard.

To measure the immunoreactivity of the BSA samples, 1 µg of protein sample was diluted into 100 µL of PBS pH 7.4 and allowed to adsorb to the wells of a 96-well plate (Maxisorp, Nunc) for a minimum of 1h at RT or overnight at 4°C. The wells were then washed twice with PBS-Tween™ 0.05% (PBS-T) and then filled with blocking solution for 30 minutes. The wells were then washed 3 times with PBS-T and filled with 100 µL of a 0.1 µg/mL dilution of the anti-TF monoclonal antibody JAAF11. After an incubation of 1h at RT with gentle shaking or 12h at 4°C, the wells were then washed 4 times with PBS-T and further incubated for 60 minutes with 100 µL of 1/1000 dilution of secondary antibody goat anti-mouse IgG (H+L)-hrp (Jackson ImmunoResearch). Wells were then washed 4 times with PBS-T and 100

μL of hrp substrate was added (Ultra TMB-ELISA, Thermo Scientific) to each wells. The reaction was stopped by the addition of 100 μL of 0.5 N sulfuric acid and the plate was read in a plate reader at 450nm (Biotek EL808).

Fig. 33 having a dual Y-axis shows the lower immunoreactivity of the BSA-TF with a high galactose ratio compared to the lower galactose conjugate. The detection of galactose in the two conjugates confirms that TF is conjugated to BSA but conjugated TF immunoreactivity is compromised at high ratio.

Example 24: Conjugation of O-Allyl-Tn to peptide dTT831-844-Cys-(Tn)-βAla BSA by thiol-ene photoreaction

Fig. 34 shows the chemical structures of the dTT831-844-Cys-βAla and the Cys-dTT831-844-βAla with a cysteine residue at the C- and N-terminal, respectively together with their tabulated LC-MS data.

Fig. 35 shows the detailed LC-MS data of the peptide dTT831-844-Cys-βAla (C-terminal).

Fig. 36 shows the detailed LC-MS data of the peptide Cys-dTT831-844-βAla (N-terminal).

Procedure A: To a solution of peptide in water (1 mL, 2mM) was added a solution of O-allyl-Tn in H₂O (200 μL, 0.1M, 10 equiv) and a catalytic amount of AAPH in UV cuvette. The mixture was stirred at room temperature under 365 nm irradiation for 60 min. Ellman's test showed the reaction was finished in 45 min. The solution was then lyophilized to afford the white power and LC-MS showed the presence of desired Tn-conjugated peptide; LC-MS: [M] calcd for C₉₄H₁₅₂O₃₁N₂₀S, 2089.0653; found, 2089.0780; CAN/H₂O 5 to 95% 5.08 min.

Procedure B in organic solvent: **dTT831-844-Cys-(Tn)-βAla (Fig. 34)** To a solution of peptide in MeOH (1 mL, 2 mM) was added a solution of Tn in H₂O (200 μL, 0.1M, 10 equiv) and a catalytic amount of DMPAP in UV cuvette. The mixture was stirred at room temperature under 365 nm irradiation for 60 min. Ellman's test showed the reaction was finished in 45 min. The solution was then lyophilized to afford the white power and LC-MS constated the presence of desired peptide-TN. LC-MS: [M] calcd for C₉₄H₁₅₂O₃₁N₂₀S, 2089.0653; found, 2089.0675; CAN/H₂O 5 to 95% 5.07 min.

Fig. 37 shows the photolytic AAPH-catalyzed thiol-ene reaction of the O-Allyl Tn on the C-terminal peptide dTT831-844-Cys-βAla.

Fig. 38 shows the LC-MS profile of the O-Allyl Tn on the C-terminal peptide

Example 25: Conjugation of O-Allyl-Tn to peptide Cys-(Tn)-dT831-844- βAla by thiol-ene photoreaction

Procedure A: To a solution of peptide Cys-(Tn)-dT831-844- βAla (Fig. 33) in water (2 mL, 0.482 mM) was added a solution of Tn in H₂O (100 μL, 0.1M, 10 equiv) and a catalytic amount of AAPH in UV cuvette. The mixture was stirred at room temperature under 365 nm irradiation for 60 min. Ellman's test showed the reaction to be finished in 45 min. The solution was then lyophilized to afford the white power and LC-MS showed the presence of desired peptide-Tn conjugate LC-MS: [M] calcd for C₉₄H₁₅₂O₃₁N₂₀S, 2089.0653; found, 2089.0719; CAN/H₂O 5 to 95% 5.11 min.

Procedure B: Cys-(Tn)-dT831-844- βAla (Fig. 33) To a solution of peptide in methanol (2 mL, 0.482 mM) was added a solution of Tn in H₂O (100 μL, 0.1M, 10 equiv) and a catalytic amount of DMPA in UV cuvette. The mixture was stirred at room temperature under 365 nm irradiation for 60 min. Ellman's test showed the reaction to be finished in 45 min. The solution was then lyophilized to afford the white power and LC-MS showed the presence of desired peptide-Tn. LC-MS: [M] calcd for C₉₄H₁₅₂O₃₁N₂₀S, 2089.0653; found, 2089.0817; CAN/H₂O 5 to 95% 5.10 min.

Fig. 39 shows the photolytic AAPH-catalyzed thiol-ene reaction of the O-Allyl Tn on the N-terminal peptide Cys-dT831-844-βAla.

Fig. 40 shows the detailed LC-MS profile of the the O-Allyl Tn on the N-terminal peptide.

REFERENCES:

Cipolla et al., "Stereoselective synthesis of α- C-glycosides of N-acetylgalactosamine", *Tetrahedron Asymm.* (2000), 11 : 295-303.

Cui et al., "Stereocontrolled allylation of 2-amino-2-deoxy sugar derivatives by a free-radical procedure", *Carbohydr. Res.*, (1998), 309: 319-330.

Danishefsky et al., "Development of Globo-H Cancer Vaccine", *Acc. Chem Res.*, (2015), 48(3): 643-652.

Demian et al., "Direct targeted glycation of the free sulfhydryl group of cysteine residue (Cys-34) of BSA. Mapping of the glycation sites of the anti-tumor Thomsen-Friedenreich neoglycoconjugate vaccine prepared by Michael addition reaction," *J. Mass Spectrom.*, (2014), 49: 1223-1233.

Dondoni et al., "A new ligation strategy for peptide and protein glycosylation: Photoinduced thiol-ene coupling", *Chem. Eur. J.*, (2009), 15: 11444-11449.

Dondoni et al., "Recent applications in thiol-ene coupling as a click process for glycoconjugation", *Chem. Soc. Rev.*, (2012), 41: 573-586.

5 Feng et al., "Chemo-enzymatic synthesis of fluorinated 2-N-acetamidoglycosaminic acid nucleotides using UDP-GlcNAc 2-epimerase", *Org. Biomol. Chem.* (2004), 2: 1617-1623.

Heimburg et al., "Inhibition of spontaneous breast cancer metastasis by anti-Thomsen-Friedenreich antigen monoclonal antibody JAA-F11.", *Neoplasia*, (2006) 8(11): 939-48.

Knapp et al., "Synthesis of α -GalNAc Thioconjugates from an α -GalNAc Mercaptan", *J. Org. Chem.*
10 (2002), 67: 2995-2999.

Papadopoulos, "Diazo transfer and click chemistry in the solid phase syntheses of lysine-based glycodendrimers as antagonists against Escherichia coli FimH." *Molecular Pharmaceutics* (2012), 9(3): 394-403.

Rittenhouse-Diakun et al., "Development and characterization of monoclonal antibody to T-antigen: (gal
15 beta1-3GalNAc-alpha-O)", *Hybridoma*, (1998), 17: 165-173.

Tati et al., "Humanization of JAA-F11, a Highly Specific Anti-Thomsen-Friedenreich Pancarcinoma Antibody and InVitro Efficacy Analysis.", *Neoplasia*, (2017), 19(9): 716-733.

CLAIMS:

1. A method for producing a glycoconjugate immunogen, the method comprising:

- (a) providing a water-soluble carbohydrate antigen covalently linked to a terminal alkene, the terminal alkene being directly conjugatable to a thiol group via a thiol-ene reaction and wherein the carbohydrate antigen is unprotected and water-soluble;
- (b) providing a carrier protein having one or more free thiol groups; and
- (c) performing a photocatalytic thiol-ene reaction to directly conjugate the carbohydrate antigen to the carrier protein at the one or more free thiol groups, thereby producing the glycoconjugate immunogen, wherein the photocatalytic thiol-ene reaction is performed:
 - (i) at a pH between 6 and 8;
 - (ii) under reaction conditions that avoid carrier protein denaturation, and/or that retain the carrier protein's activity, antigenicity, and/or structure; and
 - (iii) in the absence of any organic solvent, or in the presence of an organic solvent at a concentration sufficiently low to avoid carrier protein denaturation;

wherein the carrier protein is immunogenic when administered to a subject, and wherein conjugation of the carbohydrate antigen to the carrier protein increases the immunogenicity of the carbohydrate antigen upon administration to the subject, as compared to administration of the unconjugated carbohydrate antigen, and wherein the carbohydrate antigen comprises a B cell epitope and induces a humoral immune response in the subject.

2. The method of claim 1, wherein the photocatalytic thiol-ene reaction is performed at a pH between 6.5 and 7.5.

3. The method of claim 1 or 2, wherein said photocatalytic thiol-ene reaction is performed in the presence of a water-soluble catalyst.

4. The method of claim 3, wherein the water-soluble catalyst is: a free radical-generating azo compound; 2,2'-azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride; 2,2'-azobis(2-amidinopropane) dihydrochloride (AAPH); metals or metal ions having photoinitiator activity; a peroxide; *tert*-butyl hydroperoxide; benzoylperoxide; ammonium persulfate; or any derivative thereof having photoinitiator activity.

5. The method of claim 1 or 2, wherein said photocatalytic thiol-ene reaction is performed in the presence of a water-insoluble catalyst.

6. The method of claim 5, wherein the water-insoluble catalyst is: a free radical-generating azo compound; 2,2-dimethoxy-2-phenylacetophenone (DMPA); azobisisobutyronitrile (AIBN); 2,2'-azobis(2-methylpropionitrile); 4,4'-Azobis(4-cyanopentanoic acid) (ACVA); 1,1'-azobis(cyanocyclohexane) (ACHN); diazenedicarboxylic acid bis(*N,N*-dimethylamide) (TMAD); azodicarboxylic acid dipiperidide (ADD); or any derivative thereof having photoinitiator activity.

7. The method of any one of claims 1 to 6, wherein said photocatalytic thiol-ene reaction comprises irradiation under short-wave ultraviolet light.

8. The method of claim 7, wherein said photocatalytic thiol-ene reaction comprises irradiation at about 254 nm.

9. The method of any one of claims 1 to 7, wherein said photocatalytic thiol-ene reaction comprises irradiation under long-wave ultraviolet light.

10. The method of claim 9, wherein said photocatalytic thiol-ene reaction comprises irradiation at about 355 nm or 365 nm.

11. The method of any one of claims 1 to 10, wherein said photocatalytic thiol-ene reaction comprises reacting between 1 to 200 molar equivalents of the carbohydrate antigen per free thiol group of the carrier protein.

12. The method of any one of claims 1 to 11, wherein said carbohydrate antigen, following conjugation to the carrier protein, is not cleavable from the carrier protein by an endogenous enzyme of the subject.

13. The method of any one of claims 1 to 12, wherein said carbohydrate antigen is covalently linked to the terminal alkene.

14. The method of any one of claims 1 to 12, wherein said carbohydrate antigen is conjugated to the carrier protein, via an O-glycosidic bond, an S-glycosidic bond, an N-glycosidic bond, or a C-glycosidic bond, or a bond obtained by reductive amination between an allyl amine and a reducing sugar.

15. The method of any one of claims 1 to 14, wherein the carbohydrate antigen is or comprises a tumor associated carbohydrate antigen (TACA), a viral polysaccharide antigen, or a bacterial capsular polysaccharide (CPS).

5

16. The method of claim 15, wherein the TACA is or is from: Tn, S-Tn, *Thomsen-Friedenreich* (TF), (2,3)-S-TF, (2,6)-S-TF, Globo H, GD2, GD3, GM2, GM3, N-glycolyl-GM3, Lea, sLea, Lex, sLex, or any combination thereof.

10 17. The method of any one of claims 1 to 16, wherein said photocatalytic thiol-ene reaction conjugates more than one type of carbohydrate antigen to the carrier protein.

18. The method of claim 15, wherein the bacterial CPS is or is from a Pneumococcal and/or Streptococcal polysaccharide serotype, meningococcal CPS, or influenza CPS.

15

19. The method of any one of claims 1 to 18, wherein the carbohydrate antigen in (a) is linked to the terminal alkene via a linker.

20. The method of any one of claims 1 to 19, wherein the carrier protein comprises one or more cysteine residues having the one or more free thiol groups, or optionally is engineered to add one or more further cysteine residues at the amino terminus, the carboxy terminus, or at a solvent-accessible position of the carrier protein.

21. The method of any one of claims 1 to 20, wherein the carrier protein is or is from: Tetanus Toxoid (TT), Diphtheria Toxoid (DT), cross-reacting material 197 (CRM197), Meningococcal Outer Membrane Protein Complex (OMPC), H. Influenzae Protein D (HiD), a cytokine, an immunogenic peptide, Tetanus Toxin 831-844, albumin, or an immunogenic fragment thereof.

22. The method of any one of claims 1 to 21, wherein:
30 (i) the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, is pre-treated with a reducing agent;
(ii) the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, is pre-treated with a thiolating agent;

- (iii) the carrier protein, prior to or together with performing the photocatalytic thiol-ene reaction, is pre-treated with a thiolating agent, and subsequently pre-treated with a reducing agent;
- (iv) the carrier protein is a protein having one or more disulfide bridges, and wherein the one or more disulfide bridges remain unaffected following said photocatalytic thiol-ene reaction;
- 5 (v) the carrier protein is a protein having one or more disulfide bridges, and wherein the carrier protein is pre-treated with a reducing agent to expose one or more additional free thiol groups for conjugation to the carbohydrate antigen; or
- (iv) the carrier protein is a protein having one or more disulfide bridges, and wherein the carrier protein is pre-treated with a thiolating agent to expose one or more additional free thiol groups
- 10 for conjugation to the carbohydrate antigen, and subsequently pre-treated with a reducing agent.

23. The method of any one of claims 1 to 22, wherein the total number of carbohydrate antigens comprised in the glycoconjugate immunogen is equal to the number of free thiol groups available on the carrier protein prior to conjugation.

15

24. The method of any one of claims 1 to 23, wherein the glycoconjugate immunogen induces a cell-mediated immune response to the carbohydrate antigen upon administration to the subject.

20 25. The method of any one of claims 1 to 24, further comprising: (d) purifying the glycoconjugate immunogen.

26. A method for producing a glycoconjugate vaccine or an adaptive immune response-triggering composition, the method comprising formulating the glycoconjugate immunogen prepared by the method of any one of claims 1 to 25 with a pharmaceutically acceptable excipient, and/or an adjuvant.

25

27. The method of claim 26, wherein the adjuvant is or comprises: an inorganic compound, a mineral oil, a microbial derivative, a plant derivative, a cytokine, squalene, alum, aluminum hydroxide, aluminum phosphate, calcium phosphate hydroxide, a toll-like receptor agonist, an immunostimulatory polynucleotide, an immunostimulatory lipid, Freund's adjuvant, RIBI's adjuvant, QS-21, muramyl dipeptide, or any combination thereof.

30

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5






Fig. 6C

Compound Table					
Compound Label	RT	Mass	Abund	Formula	Tgt Mass
Cpd 1: C21 H35 N O8	0.1	429.2372	654630	C21 H35 N O8	429.2363
					Diff (ppm)
					2.16

Compound Label	RT	Algorithm	Mass
Cpd 1: C21 H35 N O8	0.1	Find By Formula	429.2372




Fig. 6D




Fig. 7A




Fig. 7B





Fig. 7C





Fig. 7D



MS Spectrum Peak List

m/z	Calc m/z	Diff(ppm)	z	Abund	Formula	Ion
262.1294	262.1285	3.29	1	4575	C11 H20 N O6	(M+H)+
281.1036				3013		
282.2803				2279		
284.1125				499971		
284.3188				35672		
284.5274				2930		
285.1149				56535		
286.1165				8170		
288.1456				40848		
300.084	300.0844	-1.48	1	1776	C11 H19 K N O6	(M+K)+

Fig. 8A




Fig. 8B





Fig. 8C



MS Spectrum Peak List

m/z	Calc m/z	Diff(ppm)	z	Abund	Formula	Ion
292.1189				268697		
292.3287				12192		
293.1218				39788		
294.1243				5059		
350.1608	350.1598	2.94	1	146470	C18 H24 N O6	(M+H)+
351.1639	351.1631	2.17	1	26665	C18 H24 N O6	(M+H)+
352.1666	352.1655	3.25	1	3895	C18 H24 N O6	(M+H)+
372.1425	372.1418	2.07	1	83058	C18 H23 N Na O6	(M+Na)+
373.1457	373.145	1.83	1	15140	C18 H23 N Na O6	(M+Na)+
388.1167	388.1157	2.72	1	5672	C18 H23 K N O6	(M+K)+

Fig. 9A




Fig. 9B




Fig. 9C




Fig. 9D




Fig. 10A




Fig. 10B




Fig. 10C





Fig. 10D



MS Spectrum Peak List

m/z	Calc m/z	Diff(ppm)	z	Abund	Formula	Ion
350.16				495714		
351.1626				91586		
512.2119	512.2126	-1.39	1	126315	C ₂₄ H ₃₄ N O ₁₁	(M+H) ⁺
529.2383	529.2392	-1.64	1	20391	C ₂₄ H ₃₇ N ₂ O ₁₁	(M+NH ₄) ⁺
534.1943	534.1946	-0.47	1	103350	C ₂₄ H ₃₃ N Na O ₁₁	(M+Na) ⁺
550.1667	550.1685	-3.25	1	6050	C ₂₄ H ₃₃ K N O ₁₁	(M+K) ⁺
1023.4185	1023.418	0.53	1	295	C ₄₈ H ₆₇ N ₂ O ₂₂	(2M+H) ⁺
1040.4393	1040.4445	-5.01	1	1591	C ₄₈ H ₇₀ N ₃ O ₂₂	(2M+NH ₄) ⁺
1045.3997	1045.3999	-0.25	1	171974	C ₄₈ H ₆₆ N ₂ Na O ₂₂	(2M+Na) ⁺
1061.3716	1061.3739	-2.14	1	1026	C ₄₈ H ₆₆ K N ₂ O ₂₂	(2M+K) ⁺

Fig. 11A




Fig. 11B





Fig. 11C





Fig. 11D



Size (d.nm):	% Number:	St Dev (d.nm):	Size (d.nm):	% Number:	St Dev (d.nm):
Peak 1: 73.36	100.0	2.037	Peak 1: 6.128	100.0	1.442
Z-Average (d.nm):			Z-Average (d.nm):		
PdI: 0.315	0.0	0.000	PdI: 0.518	0.0	0.000
Intercept: 0.864	0.0	0.000	Intercept: 0.847	0.0	0.000
Result quality : Refer to quality report			Result quality : Refer to quality report		




Figure 1




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30




Fig. 31




Fig. 32







Fig. 33



dT831-844-Cys-βAla



Cys-dTT831-844-βAla




Compound Table

Compound Label	Data File	RT	Mass	Abund	Formula	Tgt Mass	Diff (ppm)
Cpd 1: C83 H133 N19 O24 S	Roy_Chichi_181019_STC-GV-III-023_LC_5-95-10_UV_10ul.d	5.32	1811.952	211331	C83 H133 N19 O24 S	1811.9492	1.59
Cpd 1: C83 H133 N19 O24 S	Roy_Chichi_181019_STC-GV-III-024_LC_5-95-10_UV_10ul.d	5.42	1811.9526	331559	C83 H133 N19 O24 S	1811.9492	1.88


dTT831-844-Cys- β Ala

Fig. 35



Cys-dTT831-844- β Ala

Fig. 36



dT831-844-Cys-βAla-O-Tn




Fig. 38

dTT831-844-Cys-βAla-O-Tn




Fig. 39

O-Tn-Cys-dTT831-844-βAla






Fig. 40

O-Tn-Cys-dTT831-844-βAla



Photolytic thiol-ene "click chemistry"



R₁ = Cysteine residues on T-cell epitope (ex. Tetanus toxoid)

R₂ = Carbohydrate antigens serving as B-cell epitope