



(86) Date de dépôt PCT/PCT Filing Date: 2000/10/02

(87) Date publication PCT/PCT Publication Date: 2001/04/19

(45) Date de délivrance/Issue Date: 2012/07/03

(85) Entrée phase nationale/National Entry: 2002/04/12

(86) N° demande PCT/PCT Application No.: US 2000/027187

(87) N° publication PCT/PCT Publication No.: 2001/027128

(30) Priorités/Priorities: 1999/10/12 (US60/158,773);
2000/04/05 (US60/194,615)

(51) Cl.Int./Int.Cl. *C07H 15/203* (2006.01),
A61K 31/70 (2006.01), *A61K 31/7034* (2006.01),
A61K 45/06 (2006.01), *A61P 3/10* (2006.01)


(72) Inventeurs/Inventors:
ELLSWORTH, BRUCE, US;
WASHBURN, WILLIAM N., US;
SHER, PHILIP M., US;
WU, GANG, US;
MENG, WEI, US

(73) Propriétaire/Owner:
BRISTOL-MYERS SQUIBB COMPANY, US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : INHIBITEURS DE SGLT2 A BASE DE GLUCOSIDE C-ARYLE

(54) Title: C-ARYL GLUCOSIDE SGLT2 INHIBITORS AND METHOD



(57) Abrégé/Abstract:

SGLT2 inhibiting compounds are provided having formula (I) where R^1 , R^2 , and R^{2a} are independently hydrogen, OH, OR^5 , lower alkyl, CF_3 , $OCHF_2$, OCF_3 , SR^{5i} or halogen, or two of R^1 , R^2 and R^{2a} together with the carbons to which they are attached can form an annelated five, six or seven membered carbocycle or heterocycle; R^3 and R^4 are independently hydrogen, OH, OR^{5a} , $OAryl$, OCH_2Aryl , lower alkyl, cycloalkyl, CF_3 , $-OCHF_2$, $-OCF_3$, halogen, $-CN$, $-CO_2R^{5b}$, $-CO_2H$, $-COR^{6b}$, $-CH(OH)R^{6c}$, $-CH(OR^{5h})R^{6d}$, $-CONR^{6a}R^{6a}$, $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2Aryl$, $Aryl$, $-SR^{5e}$, $-SOR^{5f}$, SO_2R^{5g} , SO_2Aryl , or a five, six or seven membered heterocycle, or R^3 and R^4 together with the carbons to which they are attached form an annelated five, six or seven membered carbocycle or heterocycle; R^5 , R^{5a} , R^{5b} , R^{5c} , R^{5d} , R^{5e} , R^{5f} , R^{5g} , R^{5h} , and R^{5i} are independently lower alkyl; R^6 , R^{6a} , R^{6b} , R^{6c} and R^{6d} are independently hydrogen, alkyl, aryl, alkylaryl or cycloalkyl, or R^6 and R^{6a} together with the nitrogen to which they are attached form an annelated five, six or seven membered heterocycle; A is O, S, NH, or $(CH_2)_n$ where n is 0 - 3. A method is also provided for treating diabetes and related diseases employing an SGLT2 inhibiting amount of the above compound alone or in combination with another antidiabetic agent or other therapeutic agent.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2001 (19.04.2001)

PCT

(10) International Publication Number
WO 01/27128 A1

(51) International Patent Classification⁷: **C07H 15/203**,
A61K 31/70, A61P 3/10

08540 (US). **MENG, Wei** [CN/US]; 404 Bollen Court,
Pennington, NJ 08534 (US).

(21) International Application Number: PCT/US00/27187

(74) Agents: **KILCOYNE, John, M.** et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(22) International Filing Date: 2 October 2000 (02.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/158,773 12 October 1999 (12.10.1999) US
60/194,615 5 April 2000 (05.04.2000) US

(71) Applicant (for all designated States except US): **BRISTOL-MYERS SQUIBB COMPANY** [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **ELLSWORTH, Bruce** [US/US]; 7 Glenview Drive, Princeton, NJ 08540 (US). **WASHBURN, William, N.** [US/US]; 120 Pleasant Valley Road, Titusville, NJ 08560 (US). **SHER, Philip, M.** [US/US]; 18 Mifflin Court, Plainsboro, NJ 08536 (US). **WU, Gang** [US/US]; 390 Rosedale Road, Princeton, NJ

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: C-ARYL GLUCOSIDE SGLT2 INHIBITORS



(57) Abstract: SGLT2 inhibiting compounds are provided having formula (I) where R¹, R², and R^{2a} are independently hydrogen, OH, OR⁵, lower alkyl, CF₃, OCHF₂, OCF₃, SR⁵ⁱ or halogen, or two of R¹, R² and R^{2a} together with the carbons to which they are attached can form an annelated five, six or seven membered carbocycle or heterocycle; R³ and R⁴ are independently hydrogen, OH, OR^{5a}, OAryl, OCH₂Aryl, lower alkyl, cycloalkyl, CF₃, -OCHF₂, -OCF₃, halogen, -CN, -CO₂R^{5b}, -CO₂H, -COR^{6b}, -CH(OH)R^{6c}, -CH(OR^{5h})R^{6d}, -CONR^{6a}, -NHCOR^{5c}, -NHSO₂R^{5d}, -NHSO₂Aryl, Aryl, -SR^{5e}, -SOR^{5f}, SO₂R^{5g}, SO₂Aryl, or a five, six or seven membered heterocycle, or R³ and R⁴ together with the carbons to which they are attached form an annelated five, six or seven membered carbocycle or heterocycle; R⁵, R^{5a}, R^{5b}, R^{5c}, R^{5d}, R^{5e}, R^{5f}, R^{5g}, R^{5h}, and R⁵ⁱ are independently lower alkyl; R⁶, R^{6a}, R^{6b}, R^{6c} and R^{6d} are independently hydrogen, alkyl, aryl, alkylaryl or cycloalkyl, or R⁶ and R^{6a} together with the nitrogen to which they are attached form an annelated five, six or seven membered heterocycle; A is O, S, NH, or (CH₂)_n where n is 0 - 3. A method is also provided for treating diabetes and related diseases employing an SGLT2 inhibiting amount of the above compound alone or in combination with another antidiabetic agent or other therapeutic agent.



WO 01/27128 A1

C-ARYL GLUCOSIDE SGLT2 INHIBITORS AND METHODField of the Invention

5 The present invention relates to C-aryl glucosides which are inhibitors of sodium dependent glucose transporters found in the intestine and kidney (SGLT2) and to a method for treating diabetes, especially type II diabetes, as well as hyperglycemia, hyperinsulinemia, 10 obesity, hypertriglyceridemia, Syndrome X, diabetic complications, atherosclerosis and related diseases, employing such C-aryl glucosides alone or in combination with one, two or more other type antidiabetic agent and/or one, two or more other type therapeutic agents 15 such as hypolipidemic agents.

Background of the Invention

Approximately 100 million people worldwide suffer 20 from type II diabetes (NIDDM), which is characterized by hyperglycemia due to excessive hepatic glucose production and peripheral insulin resistance, the root causes for which are as yet unknown. Hyperglycemia is considered to be the major risk factor for the development of diabetic 25 complications, and is likely to contribute directly to the impairment of insulin secretion seen in advanced NIDDM. Normalization of plasma glucose in NIDDM patients would be predicted to improve insulin action, and to offset the development of diabetic complications. An 30 inhibitor of the sodium-dependent glucose transporter SGLT2 in the kidney would be expected to aid in the normalization of plasma glucose levels, and perhaps body weight, by enhancing glucose excretion.

The development of novel, safe, and orally active 35 antidiabetic agents is also desired in order to

complement existing therapies, including the sulfonylureas, thiazolidinediones, metformin, and insulin, and to avoid the potential side effects associated with the use of these other agents.

5 Hyperglycemia is a hallmark of type II diabetes (NIDDM); consistent control of plasma glucose levels in diabetes can offset the development of diabetic complications and beta cell failure seen in advanced disease. Plasma glucose is normally filtered in the
10 kidney in the glomerulus and actively reabsorbed in the proximal tubule. SGLT2 appears to be the major transporter responsible for the reuptake of glucose at this site. The SGLT specific inhibitor phlorizin or closely related analogs inhibit this reuptake process in
15 diabetic rodents and dogs resulting in normalization of plasma glucose levels by promoting glucose excretion without hypoglycemic side effects. Long term (6 month) treatment of Zucker diabetic rats with an SGLT2 inhibitor has been reported to improve insulin response to
20 glycemia, improve insulin sensitivity, and delay the onset of nephropathy and neuropathy in these animals, with no detectable pathology in the kidney and no electrolyte imbalance in plasma. Selective inhibition of SGLT2 in diabetic patients would be expected to normalize
25 plasma glucose by enhancing the excretion of glucose in the urine, thereby improving insulin sensitivity, and delaying the development of diabetic complications.

Ninety percent of glucose reuptake in the kidney occurs in the epithelial cells of the early S1 segment of
30 the renal cortical proximal tubule, and SGLT2 is likely to be the major transporter responsible for this reuptake. SGLT2 is a 672 amino acid protein containing 14 membrane-spanning segments that is predominantly expressed in the early S1 segment of the renal proximal
35 tubules. The substrate specificity, sodium dependence,

and localization of SGLT2 are consistent with the properties of the high capacity, low affinity, sodium-dependent glucose transporter previously characterized in human cortical kidney proximal tubules. In addition,
5 hybrid depletion studies implicate SGLT2 as the predominant Na^+ /glucose cotransporter in the S1 segment of the proximal tubule, since virtually all Na-dependent glucose transport activity encoded in mRNA from rat kidney cortex is inhibited by an antisense
10 oligonucleotide specific to rat SGLT2. SGLT2 is a candidate gene for some forms of familial glucosuria, a genetic abnormality in which renal glucose reabsorption is impaired to varying degrees. None of these syndromes investigated to date map to the SGLT2 locus on chromosome
15 16. However, the studies of highly homologous rodent SGLTs strongly implicate SGLT2 as the major renal sodium-dependent transporter of glucose and suggest that the glucosuria locus that has been mapped encodes an SGLT2 regulator. Inhibition of SGLT2 would be predicted to
20 reduce plasma glucose levels via enhanced glucose excretion in diabetic patients.

SGLT1, another Na-dependent glucose cotransporter that is 60% identical to SGLT2 at the amino acid level, is expressed in the small intestine and in the more
25 distal S3 segment of the renal proximal tubule. Despite their sequence similarities, human SGLT1 and SGLT2 are biochemically distinguishable. For SGLT1, the molar ratio of Na^+ to glucose transported is 2:1, whereas for SGLT2, the ratio is 1:1. The K_m for Na^+ is 32 and 250-
30 300 mM for SGLT1 and SGLT2, respectively. K_m values for uptake of glucose and the nonmetabolizable glucose analog α -methyl-D-glucopyranoside (AMG) are similar for SGLT1 and SGLT2, i.e. 0.8 and 1.6 mM (glucose) and 0.4 and 1.6 mM (AMG) for SGLT1 and SGLT2 transporters, respectively.

However, the two transporters do vary in their substrate specificities for sugars such as galactose, which is a substrate for SGLT1 only.

Administration of phlorizin, a specific inhibitor of SGLT activity, provided proof of concept *in vivo* by promoting glucose excretion, lowering fasting and fed plasma glucose, and promoting glucose utilization without hypoglycemic side effects in several diabetic rodent models and in one canine diabetes model. No adverse effects on plasma ion balance, renal function or renal morphology have been observed as a consequence of phlorizin treatment for as long as two weeks. In addition, no hypoglycemic or other adverse effects have been observed when phlorizin is administered to normal animals, despite the presence of glycosuria. Administration of an inhibitor of renal SGLTs for a 6-month period (Tanabe Seiyaku) was reported to improve fasting and fed plasma glucose, improve insulin secretion and utilization in obese NIDDM rat models, and offset the development of nephropathy and neuropathy in the absence of hypoglycemic or renal side effects.

Phlorizin itself is unattractive as an oral drug since it is a nonspecific SGLT1/SGLT2 inhibitor that is hydrolyzed in the gut to its aglycone phloretin, which is a potent inhibitor of facilitated glucose transport. Concurrent inhibition of facilitative glucose transporters (GLUTs) is undesirable since such inhibitors would be predicted to exacerbate peripheral insulin resistance as well as promote hypoglycemia in the CNS. Inhibition of SGLT1 could also have serious adverse consequences as is illustrated by the hereditary syndrome glucose/galactose malabsorption (GGM), in which mutations in the SGLT1 cotransporter result in impaired glucose uptake in the intestine, and life-threatening diarrhea and dehydration. The biochemical differences between

SGLT2 and SGLT1, as well as the degree of sequence divergence between them, allow for identification of selective SGLT2 inhibitors.


The familial glycosuria syndromes are conditions in which intestinal glucose transport, and renal transport of other ions and amino acids, are normal. Familial glycosuria patients appear to develop normally, have normal plasma glucose levels, and appear to suffer no major health deficits as a consequence of their disorder, despite sometimes quite high (110-114 g/daily) levels of glucose excreted. The major symptoms evident in these patients include polyphagia, polyuria and polydipsia, and the kidneys appear to be normal in structure and function. Thus, from the evidence available thus far, defects in renal reuptake of glucose appear to have minimal long term negative consequences in otherwise normal individuals.

The following references disclose O-aryl glucosides SGLT2 inhibitors for treating diabetes.

20


EP 598359A1

(Tanabe


Seiyaku)discloses compounds of the following structure AA $R^1 = \text{H, acyl,}$ $R^2 = \text{H, Me}$ $R^3, R^4 \text{ can be a variety of substituents}$

25


EP 0850948A1 discloses structures of the following genus B

B

JP 09188625A expands upon structure **B** to include
 5 examples of **B** where R^3 is H and where the 5 membered ring
 is saturated as well as the counterparts of
 benzothiophenes ($\text{O} = \text{S}$) and indenenes ($\text{O} = \text{CH}_2$).




10 JP 09124685A expands upon structure **B** for $R^3 = \text{H}$ to
 include derivatives of mono acylated C6 hydroxyl where
 the acyl group is a substituted benzoic or pyridyl
 carboxylic acid or a urethane generated from the
 corresponding phenol.




15

JP 09124684 discloses derivatives of structure **B**



EP 773226-A1 discloses derivatives of structure **B**



5

JP 08027006-A discloses derivatives of structure **A** where various combinations of the glucose hydroxyl are acylated and appears to be similar to EP 598359A1

10 EP 684254-A1 appears to encompass derivatives of structure **B** disclosed in JP 09188625A.

Other disclosures and publications which disclose SGLT2 inhibitors include the following:

15 K. Tsujihara et al, *Chem. Pharm. Bull.* **44**, 1174-1180 (1996)


M. Hongu et al, *Chem. Pharm. Bull.* **46**, 22-33 (1998)

M. Hongu et al, *Chem. Pharm. Bull.* **46**, 1545-1555 (1998)


A.Oku et al, *Diabetes*, **48**, 1794-1800 (1999)

20 JP 10245391 (Dainippon) discloses 500 structures as hypoglycemic agents for treatment of diabetes. These are O-glucosides of hydroxylated coumarins.

WO 98/31697 discloses compounds of the structure



Where Ar includes, among others, phenyl, biphenyl, diphenylmethane, diphenylethane, and diphenylether, and
 5 R^1 is a glycoside, R^2 is H, OH, amino, halogen, carboxy, alkyl, cycloalkyl, or carboxamido, and R^3 is hydrogen, alkyl, or acyl, and k, m, and n are independently 1 - 4.
 A subset of compounds disclosed in WO 98/31697 contains compounds of the following structures




A is O or $(CH_2)_x$ where $x = 0-3$
 R^3 is hydrogen, alkyl or acyl group where n is 1-4
 R^2 is hydrogen, alkyl, OH, NH_2 , halogen, CO_2H or carboximide where k is 1-4

10 which are disclosed for use in the treatment or prevention of inflammatory diseases, autoimmune diseases, infections, cancer, and cancer metastasis, reperfusion disorders, thrombosis, ulcer, wounds, osteoporosis,
 15 diabetes mellitus and atherosclerosis, among others.

Description of the Invention

In accordance with the present invention, C-aryl glucoside compounds are provided which have the structure
 20 I



wherein

R^1 , R^2 and R^{2a} are independently hydrogen, OH, OR^5 , alkyl, CF_3 , $OCHF_2$, OCF_3 , SR^{5i} or halogen, or two of R^1 , R^2 and R^{2a} together with the carbons to which they are
 5 attached can form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, and/or SO_2 ;

R^3 and R^4 are independently hydrogen, OH, OR^{5a} , OAr yl, OCH_2Ar yl, alkyl, cycloalkyl, CF_3 , $-OCHF_2$, $-OCF_3$, halogen, $-CN$, $-CO_2R^{5b}$, $-CO_2H$, $-COR^{6b}$, $-CH(OH)R^{6c}$, $-CH(OR^{5h})R^{6d}$, $-CONR^{6a}$, $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2Ar$ yl, Ar yl, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$, $-SO_2Ar$ yl, or a five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the
 10 ring which are N, O, S, SO, and/or SO_2 , or R^3 and R^4 together with the carbons to which they are attached form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the
 15 ring which are N, O, S, SO, and/or SO_2 ;

R^5 , R^{5a} , R^{5b} , R^{5c} , R^{5d} , R^{5e} , R^{5f} , R^{5g} , R^{5h} and R^{5i} are
 20 independently alkyl;

R^6 , R^{6a} , R^{6b} , R^{6c} and R^{6d} are independently hydrogen, alkyl, aryl, alkylaryl or cycloalkyl, or R^6 and R^{6a} together with the nitrogen to which they are attached
 25 form an annelated five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, and/or SO_2 ;

A is O, S, NH, or $(CH_2)_n$ where n is 0 - 3, and a pharmaceutically acceptable salt thereof, all
 30 stereoisomers thereof, and all prodrug esters thereof.

The compounds of formula I of the invention as defined above also include the proviso that where A is $(CH_2)_n$ where n is 0, 1, 2, or 3 or A is O, and at least one of R^1 , R^2 , and R^{2a} is OH or OR^5 , then at least one of
 35 R^1 , R^2 , and R^{2a} is CF_3 , OCF_3 , or $OCHF_2$ and/or at least one

of R^3 and R^4 is CF_3 , $-OCHF_2$, $-OCF_3$, $CH(OR^{5h})R^{6d}$, $CH(OH)R^{6c}$, COR^{6b} , $-CN$, $-CO_2R^{5b}$, $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2Aryl$, $Aryl$, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$ or $-SO_2Aryl$.

Preferred compounds of formula I as defined above
5 include the proviso that where A is $(CH_2)_n$ where n is 0, 1, 2, or 3 or A is O, and at least one of R^1 , R^2 , R^{2a} , R^3 and R^4 is OH or OR^5 , then at least one of R^1 , R^2 , and R^{2a} is CF_3 , OCF_3 , or $OCHF_2$ and/or at least one of R^3 and R^4 is CF_3 , $-OCHF_2$, $-OCF_3$, $-CN$, $-CO_2R^{5b}$, $CH(OR^{5h})R^{6d}$, $-NHCOR^{5c}$,
10 $-NHSO_2R^{5d}$, $-NHSO_2Aryl$, $Aryl$, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$, $-SO_2Aryl$ or halogen.

The compounds of formula I possess activity as inhibitors of the sodium dependent glucose transporters found in the intestine and kidney of mammals and are
15 useful in the treatment of diabetes and the micro- and macrovascular complications of diabetes such as retinopathy, neuropathy, nephropathy, and wound healing.

The present invention provides for compounds of formula I, pharmaceutical compositions employing such
20 compounds and for methods of using such compounds.

In addition, in accordance with the present invention, a method is provided for treating or delaying the progression or onset of diabetes, especially type I and type II diabetes, including complications of
25 diabetes, including retinopathy, neuropathy, nephropathy and delayed wound healing, and related diseases such as insulin resistance (impaired glucose homeostasis), hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, obesity, hyperlipidemia
30 including hypertriglyceridemia, Syndrome X, atherosclerosis and hypertension, and for increasing high density lipoprotein levels, wherein a therapeutically effective amount of a compound of structure I is administered to a human patient in need of treatment.

In addition, in accordance with the present invention, a method is provided for treating diabetes and related diseases as defined above and hereinafter, wherein a therapeutically effective amount of a
5 combination of a compound of structure I and another type of antidiabetic agent and/or another type of therapeutic agent such as a hypolipidemic agent is administered to a human patient in need of treatment.


The conditions, diseases, and maladies collectively
10 referred to as "Syndrome X" (also known as Metabolic Syndrome) are detailed in Johannsson *J. Clin. Endocrinol. Metab.*, 82, 727-34 (1997).

The term "other type of therapeutic agents" as employed herein refers to one or more antidiabetic agents
15 (other than SGLT2 inhibitors of formula I), one or more anti-obesity agents, anti-hypertensive agents, anti-platelet agents, anti-atherosclerotic agents and/or one or more lipid-lowering agents (including anti-atherosclerosis agents).

20 In the above method of the invention, the compound of structure I of the invention will be employed in a weight ratio to the one, two or more antidiabetic agent and/or one, two or more other type therapeutic agent (depending upon its mode of operation) within the range
25 from about 0.01:1 to about 300:1, preferably from about 0.1:1 to about 10:1.

Preferred are compounds of formula IA

IA



wherein A is CH₂ or O or S and is linked meta to the glucoside;

R¹, R² and R^{2a} are independently selected from H, lower alkyl, halogen, OR⁵, or OCHF₂ or two of R¹, R² and R^{2a} are H and the other is lower alkyl, halogen, OR⁵ or OCHF₂;

R³ and R⁴ are independently selected from lower alkyl, OR^{5a}, -OCHF₂, -SR^{5e}, OH, -CO₂R^{5b}, -3,4-(OCH₂O)-, -COR^{6b}, -CH(OH)R^{6c}, -CH(OR^{5h})R^{6d}, CF₃, R^{5c}-C(=O)-NH-, -SOR^{5f}, -SO₂R^{5g}, aryl, -NHSO₂Aryl, -NHSO₂R^{5d}, COOH, thiadiazole, tetrazole, -OCH₂Aryl, -OCF₃, OAryl, or H.

More preferred are compounds of formula I where A is CH₂;

R¹ is hydrogen, halogen or lower alkyl;


R² and R^{2a} are each H;

R³ is H;

R⁴ is lower alkyl, -COR^{6b}, -CH(OH)R^{6c}, -CH(OR^{5h})R^{6d}, R^{5a}O, -OCHF₂, -OCF₃ or -SR^{5e}.

Most preferred are compounds of formula I of the structure IB

IB




where R¹ is hydrogen, halogen or lower alkyl and R⁴ is lower alkyl, R^{5a}O, -OCHF₂, or -SR^{5e}. It is preferred that R¹ be linked *para* to the glucoside bond and the R⁴ substituent be linked at the *para* position.

Detailed Description of the Invention

The compounds of formula I of the invention can be prepared as shown in the following reaction schemes and description thereof wherein temperatures are expressed in
 5 degrees Centigrade.

Compounds of formula I can be prepared as shown in Scheme 1 by treatment of compounds of formula II
 II




(where Bn = benzyl)


10 with H₂ in the presence of a catalyst such as 1) Pd/C employing a solvent such as MeOH or EtOH or 2) preferably Pd(OH)₂ using a solvent such as EtOAc. Alternatively, compounds of formula I can be prepared by treatment of compounds of formula II with a Lewis acid such BBr₃, BCl₃,
 15 or BCl₃·Me₂S in a solvent such as CH₂Cl₂ at -78°. Compounds of formula I can also be prepared by treatment of compounds of formula II in a solvent such as EtSH containing BF₃·Et₂O, at 20°.

20 Compounds of formula II (which are novel intermediates) can be prepared by treatment of compounds of formula III with silanes such as Et₃SiH or preferably (iPr)₃SiH in a solvent such as MeCN or mixtures of MeCN/CH₂Cl₂ containing a Lewis acid such as BF₃·Et₂O at -30°.

III




Compounds of formula III (which are novel intermediates) can be prepared by coupling of a compound
 5 of formula IV
 IV




with compound V.

10 V




Compounds of formula IV are activated for coupling by treatment with *n*-BuLi or *t*-BuLi at -78° in a solvent such as THF prior to addition of lactone V. Preparation
 15 of lactone V is described in. R. Benhaddou, S Czernecki, et al., *Carbohydr. Res.*, **260** (1994), 243-250.

Scheme 1




Compounds of formula IV where A is $(\text{CH}_2)_n$ where $n = 1-3$ can be prepared as shown in Scheme 2 by treatment of compounds of formula VI

5 VI




with silanes such as Et_3SiH in a solvent such as MeCN or CH_2Cl_2 containing a Lewis acid such as $\text{BF}_3 \cdot \text{Et}_2\text{O}$ or TFA at
10 -30° to $+60^\circ$.

Compounds of formula VI can be prepared by coupling commercially available bromobenzaldehydes of formula VII
VII



with either the lithium or magnesium organometallic derivative of compounds of formula VIII
VIII




5

in a solvent such as Et₂O or THF using conditions familiar to those skilled in the art.

Compounds of formula VIII are either commercially available or readily prepared by standard methods known to those skilled in the art.

10

Scheme 2



Compounds of formula I where R⁴ is CH(OR^{5h})R^{6d} can be prepared by treatment of compounds of formula I where R⁴ is COR^{6b} sequentially with 1) an acetylating agent such as Ac₂O in a solvent such as pyridine alone or CH₂Cl₂ containing 1.5 equivalents of a base such as Et₃N, 2) a reducing agent such as NaBH₄ in a solvent such as EtOH, 3) an alkylating agent such as R^{5h}Br or R^{5h}I in the presence of a base such as NAH in a solvent such as DMF, and 4) alkaline ester hydrolysis conditions such as LiOH in a 2:3:1 mixture of THF/MeOH/H₂O.


20

Compounds of formula I where R^4 is $CH(OH)R^{6c}$ can be prepared by treatment of compounds of formula I where R^4 is COR^{6b} with a reducing agent such as $NaBH_4$ in a solvent such as EtOH.

- 5 Compounds of formula I where R^4 is COR^{6b} can be prepared by treatment of compounds of formula II where R^4 is COR^{6b} with a Lewis acid such as BCl_3 or BBr_3 at -78° in a solvent such as CH_2Cl_2 .


- 10 Compounds of formula II where A is CH_2 and R^4 is $-COR^{6b}$ can be prepared as shown in Scheme 3 by coupling commercially available or readily accessible compounds of formula IX

IX



15


where Z is Br or Cl with compounds of formula X



- 20 by heating the two components in a solvent such as PhMe in the presence of a catalyst such as $Pd(PPh_3)_4$.


Compounds of formula X (which are novel intermediates) can be prepared from compounds of formula XI

XI




by treatment with $(\text{Bu}_3\text{Sn})_2$ and a catalyst such as $\text{Pd}(\text{Ph}_3\text{P})_4$ in a solvent such as toluene.

- 5 Compounds of formula XI (which are novel intermediates) can be prepared from compounds of formula XII
XII
XII



- 10 by treatment with silanes such as $i\text{Pr}_3\text{SiH}$ or Et_3SiH in a solvent such as MeCN containing a Lewis acid such as $\text{BF}_3 \cdot \text{Et}_2\text{O}$ at -30° .


- Compounds of formula XII (which are novel intermediates) can be prepared by coupling compound V
15 with the organolithium obtained upon treatment of compounds of formula XIII
XIII




with $n\text{-BuLi}$ or $t\text{-BuLi}$ at -78° in THF.

20

Scheme 3




An alternative synthesis (Scheme 4) of compounds of
 formula IV where A is CH₂ entails reduction of compounds
 5 of formula XIV
 XIV



with a reducing agent such as Et₃SiH in a solvent such as
 10 MeCN or CH₂Cl₂ or mixtures thereof containing a catalyst
 such as BF₃·Et₂O.


Compounds of formula XIV can be readily prepared by
 Friedel-Craft acylation of commercially available
 hydrocarbons of formula XV

XV




with readily available acid chlorides of formula XVI

5 XVI




in a solvent such as CS₂ containing two equivalents of a Lewis Acid such as AlCl₃ or AlBr₃.

10

Scheme 4


Compounds of formula II where A is a bond can be prepared as shown in Scheme 5 by coupling compounds of formula XI with compounds of formula XVII

15 XVII




or the corresponding boronic acid XVIII.

XVIII




20 Coupling entails heating in the presence of a catalyst such as Pd(PPh₃)₄ employing a solvent such as 3:1

PhMe/EtOH containing Na_2CO_3 . Compounds of formula XVIII are either commercially available or can be prepared upon treatment of compounds of formula XVII with BCl_3 in a solvent such as CH_2Cl_2 . Compounds of formula XVII can be prepared by heating compounds of formula XIX




in a solvent such as DMSO containing a catalyst such as $\text{PdCl}_2 \cdot \text{dppf}$ and a base such as KOAc with compound XX.

XX




Scheme 5



Compounds of formula II, where $\text{A} = \text{CH}_2$ and $\text{R}^2 = \text{OH}$, can be prepared as shown in Scheme 6 upon sequential treatment of compounds of formula XXI


XXI



with a base such as NaH followed by heating with compounds of formula IX in a solvent such as PhMe.


5 Compounds of formula XXI can be prepared from compounds of formula XXII

XXII



10 by treatment with silanes such as Et₃SiH or i-Pr₃SiH in a solvent such as MeCN containing a Lewis acid such as BF₃·Et₂O at -30°.


Compounds of formula XXII can be prepared by coupling the compound of formula V with activated metallated derivatives of compounds of formula XXIII



15 which are prepared by sequential treatment of XXIII with a base such as NaH, KH, or KOtBu followed by an alkyllithium such as nBuLi or tBuLi in a solvent such as dry THF.


20

Scheme 6




Compounds of formula I, where A = O or NH, can be prepared as shown in Scheme 7 by coupling compounds of formula XXIV

5 XXIV



with commercially available compounds of formula XXV where X = O or NH


10 XXV



by heating in a solvent such as pyridine containing a base such as Et₃N, a catalyst such as Cu(OAc)₂ and molecular sieves.

Compounds of formula XXIV (which are novel intermediates) can be prepared by treating compounds of formula XXVI with BCl_3 in a solvent such as CH_2Cl_2 at -78° .


5 XXVI



Compounds of formula XXVI (which are novel intermediates) can be prepared by heating compounds of formula XI with compounds of formula XX in a solvent such as DMSO containing a catalyst such as $\text{PdCl}_2 \cdot \text{dppf}$ and a base such as KOAc.

10


Scheme 7




15

Compounds of formula IV where A is O or NH can be prepared as shown in Scheme 8 by coupling compounds of formula XVIII

XVIII




with compounds of formula XXVII where X = O or NH
XXVII



by heating in a solvent such as pyridine containing a
base such as Et₃N, a catalyst such as Cu(OAc)₂ and
molecular sieves.


10

Scheme 8

Compounds of formula IV where A is S can be prepared
as shown in Scheme 9 by coupling aryl disulfides of
formula XXVIII

15


XXVIII



with the organolithium obtained upon metalation of
compounds of formula XIII with *n*-BuLi or *t*-BuLi at -78°
in THF.

20

Scheme 9



Listed below are definitions of various terms used in the description of the instant invention. These definitions apply to the terms as they are used throughout the specification (unless they are otherwise limited in specific instances) either individually or as part of a larger group.

The following abbreviations are employed herein:

Ph = phenyl

Bn = benzyl

t-Bu = tertiary butyl

Me = methyl

Et = ethyl

TMS = trimethylsilyl

TMSN₃ = trimethylsilyl azide

TBS = tert-butyldimethylsilyl

THF = tetrahydrofuran

Et₂O = diethyl ether

EtOAc = ethyl acetate

DMF = dimethyl formamide

MeOH = methanol

EtOH = ethanol

i-PrOH = isopropanol

HOAc or AcOH = acetic acid

TFA = trifluoroacetic acid

i-Pr₂NEt = diisopropylethylamine

Et₃N = triethylamine

- DMAP = 4-dimethylaminopyridine
NaBH₄ = sodium borohydride
LiAlH₄ = lithium aluminum hydride
n-BuLi = n-butyllithium
- 5 Pd/C = palladium on carbon
KOH = potassium hydroxide
NaOH = sodium hydroxide
LiOH = lithium hydroxide
K₂CO₃ = potassium carbonate
- 10 NaHCO₃ = sodium bicarbonate
EDC (or EDC.HCl) or EDCI (or EDCI.HCl) or EDAC = 3-ethyl-3'-(dimethylamino)propyl- carbodiimide hydrochloride (or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride)
- 15 HOBT or HOBT.H₂O = 1-hydroxybenzotriazole hydrate
HOAT = 1-Hydroxy-7-azabenzotriazole
Ph₃P = triphenylphosphine
Pd(OAc)₂ = Palladium acetate
(Ph₃P)₄Pd⁰ = tetrakis triphenylphosphine palladium
- 20 Ar = argon
N₂ = nitrogen
min = minute(s)
h or hr = hour(s)
L = liter
- 25 mL = milliliter
μL = microliter
g = gram(s)
mg = milligram(s)
mol = moles
- 30 mmol = millimole(s)
meq = milliequivalent
RT = room temperature
sat or sat'd = saturated
aq. = aqueous
- 35 TLC = thin layer chromatography

HPLC = high performance liquid chromatography

LC/MS = high performance liquid chromatography/mass spectrometry

MS or Mass Spec = mass spectrometry

5 NMR = nuclear magnetic resonance


mp = melting point

dppf = diphenylphosphinoferrocene

Unless otherwise indicated, the term "lower alkyl" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons containing 1 to 8 carbons, and the terms "alkyl" and "alk" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons containing 1 to 20 carbons, preferably 1 to 10 carbons, more preferably 1 to 8 carbons, in the normal chain, such as methyl, ethyl, propyl, isopropyl, butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl, dodecyl, the various branched chain isomers thereof, and the like as well as such groups including 1 to 4 substituents such as halo, for example F, Br, Cl or I or CF₃, alkyl, alkoxy, aryl, aryloxy, aryl(aryl) or diaryl, arylalkyl, arylalkyloxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkylalkyloxy, optionally substituted amino, hydroxy, hydroxyalkyl, acyl, alkanoyl, heteroaryl, heteroaryloxy, cycloheteroalkyl, arylheteroaryl, arylalkoxycarbonyl, heteroarylalkyl, heteroarylalkoxy, aryloxyalkyl, aryloxyaryl, alkylamido, alkanoylamino, arylcarbonylamino, nitro, cyano, thiol, haloalkyl, trihaloalkyl and/or alkylthio.

Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1

or 2 double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclicalkyl, bicyclicalkyl and tricyclicalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 3 to 10 carbons, forming
 5 the ring and which may be fused to 1 or 2 aromatic rings as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl,



any of which groups may be optionally substituted with 1 to 4 substituents such as halogen, alkyl, alkoxy, hydroxy, aryl, aryloxy, arylalkyl, cycloalkyl,
 15 alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the alkyl substituents.

The term "cycloalkenyl" as employed herein alone or as part of another group refers to cyclic hydrocarbons
 20 containing 3 to 12 carbons, preferably 5 to 10 carbons and 1 or 2 double bonds. Exemplary cycloalkenyl groups include cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclohexadienyl, and cycloheptadienyl, which may be optionally substituted as defined for
 25 cycloalkyl.

The term "alkanoyl" as used herein alone or as part of another group refers to alkyl linked to a carbonyl group.

Unless otherwise indicated, the term "lower alkenyl" as used herein by itself or as part of another group
 30 refers to straight or branched chain radicals of 2 to 8 carbons, and the term "alkenyl" as used herein by itself

or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons, and more preferably 2 to 8 carbons in the normal chain, which include one to six double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, hydroxy, heteroaryl, cycloheteroalkyl, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, alkylthio and/or any of the alkyl substituents set out herein.

Unless otherwise indicated, the term "lower alkynyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 8 carbons, and the term "alkynyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons in the normal chain, which include one triple bond in the normal chain, such as 2-propynyl, 3-butynyl, 2-butynyl, 4-pentynyl, 3-pentynyl, 2-hexynyl, 3-hexynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 3-octynyl, 3-nonynyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like, and which may be optionally substituted with 1 to 4 substituents, namely, halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, amino, heteroaryl, cycloheteroalkyl, hydroxy, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, and/or alkylthio, and/or any of the alkyl substituents set out herein.

The terms "arylalkyl", "arylalkenyl" and "arylalkynyl" as used alone or as part of another group





refer to alkyl, alkenyl and alkynyl groups as described above having an aryl substituent.


Where alkyl groups as defined above have single bonds for attachment to other groups at two different carbon atoms, they are termed "alkylene" groups and may optionally be substituted as defined above for "alkyl".

Where alkenyl groups as defined above and alkynyl groups as defined above, respectively, have single bonds for attachment at two different carbon atoms, they are termed "alkenylene groups" and "alkynylene groups", respectively, and may optionally be substituted as defined above for "alkenyl" and "alkynyl".

Suitable alkylene, alkenylene or alkynylene groups $(\text{CH}_2)_m$ or $(\text{CH}_2)_p$ (where p is 1 to 8, preferably 1 to 5, and m is 1 to 5, preferably 1 to 3, which includes alkylene, alkenylene or alkynylene groups) as defined herein, may optionally include 1, 2, or 3 substituents which include alkyl, alkenyl, halogen, cyano, hydroxy, alkoxy, amino, thioalkyl, keto, C_3 - C_6 cycloalkyl, alkylcarbonylamino or alkylcarbonyloxy.

Examples of $(\text{CH}_2)_m$ or $(\text{CH}_2)_p$, alkylene, alkenylene and alkynylene include $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$,






10

The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine, with chlorine or fluorine being preferred.

15

The term "metal ion" refers to alkali metal ions such as sodium, potassium or lithium and alkaline earth metal ions such as magnesium and calcium, as well as zinc and aluminum.

Unless otherwise indicated, the term "aryl" or "Aryl" as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including 1-naphthyl and 2-naphthyl) and may optionally include one to three additional rings fused to a carbocyclic ring or a heterocyclic ring (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings for example



and may be optionally substituted through available carbon atoms with 1, 2, or 3 groups selected from hydrogen, halo, haloalkyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkyl-alkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, alkoxycarbonyl, arylcarbonyl, arylalkenyl, aminocarbonylaryl, arylthio, arylsulfinyl, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino and arylsulfonaminocarbonyl and/or any of the alkyl substituents set out herein.

Unless otherwise indicated, the term "lower alkoxy", "alkoxy", "aryloxy" or "aralkoxy" as employed herein
30 alone or as part of another group includes any of the

above alkyl, aralkyl or aryl groups linked to an oxygen atom.

Unless otherwise indicated, the term "substituted amino" as employed herein alone or as part of another group refers to amino substituted with one or two substituents, which may be the same or different, such as alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl and thioalkyl. These substituents may be further substituted with a carboxylic acid and/or any of the alkyl substituents as set out above. In addition, the amino substituents may be taken together with the nitrogen atom to which they are attached to form 1-pyrrolidinyl, 1-piperidinyl, 1-azepinyl, 4-morpholinyl, 4-thiamorpholinyl, 1-piperazinyl, 4-alkyl-1-piperazinyl, 4-arylalkyl-1-piperazinyl, 4-diarylalkyl-1-piperazinyl, 1-pyrrolidinyl, 1-piperidinyl, or 1-azepinyl, optionally substituted with alkyl, alkoxy, alkylthio, halo, trifluoromethyl or hydroxy.

Unless otherwise indicated, the term "lower alkylthio", "alkylthio", "arylthio" or "aralkylthio" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to a sulfur atom.


Unless otherwise indicated, the term "lower alkylamino", "alkylamino", "arylamino", or "arylalkylamino" as employed herein alone or as part of another group includes any of the above alkyl, aryl or arylalkyl groups linked to a nitrogen atom.

Unless otherwise indicated, the term "acyl" as employed herein by itself or as part of another group, as defined herein, refers to an organic radical linked to a

carbonyl $\left(\begin{array}{c} \text{O} \\ \parallel \\ \text{C} \end{array} \right)$ group; examples of acyl groups include any

of the alkyl substituents attached to a carbonyl, such as alkanoyl, alkenoyl, aroyl, aralkanoyl, heteroaroyl, cycloalkanoyl, cycloheteroalkanoyl and the like.



Unless otherwise indicated, the term
 5 "cycloheteroalkyl" as used herein alone or as part of another group refers to a 5-, 6- or 7-membered saturated or partially unsaturated ring which includes 1 to 2 hetero atoms such as nitrogen, oxygen and/or sulfur, linked through a carbon atom or a heteroatom, where
 10 possible, optionally via the linker $(CH_2)_p$ (where p is 1, 2 or 3), such as





and the like. The above groups may include 1 to 4 substituents such as alkyl, halo, oxo and/or any of the alkyl substituents set out herein. In addition, any of the cycloheteroalkyl rings can be fused to a cycloalkyl, aryl, heteroaryl or cycloheteroalkyl ring.
 25

Unless otherwise indicated, the term "heteroaryl" as used herein alone or as part of another group refers to a 5- or 6- membered aromatic ring which includes 1, 2, 3 or 4 hetero atoms such as nitrogen, oxygen or sulfur, and

such rings fused to an aryl, cycloalkyl, heteroaryl or cycloheteroalkyl ring (e.g., benzothiophenyl or indolyl), and includes possible N-oxides. The heteroaryl group may optionally include 1 to 4 substituents such as any of the
 5 the alkyl substituents set out above. Examples of heteroaryl groups include the following:



10



15 and the like.

The term "cycloheteroalkylalkyl" as used herein alone or as part of another group refers to cycloheteroalkyl groups as defined above linked through a C atom or heteroatom to a $(CH_2)_p$ chain.

20 The term "heteroarylalkyl" or "heteroarylalkenyl" as used herein alone or as part of another group refers to a heteroaryl group as defined above linked through a C atom or heteroatom to a $-(CH_2)_p-$ chain, alkylene or alkenylene as defined above.

25 The term "five, six or seven membered carbocycle or heterocycle" as employed herein refers to cycloalkyl or cycloalkenyl groups as defined above or heteroaryl groups


or cycloheteroaryl groups as defined above, such as thiadiazole, tetrazole, imidazole, or oxazole.

The term "polyhaloalkyl" as used herein refers to an "alkyl" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as CF₃CH₂, CF₃ or CF₃CF₂CH₂.

The term "polyhaloalkyloxy" as used herein refers to an "alkoxy" or "alkyloxy" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as CF₃CH₂O, CF₃O or CF₃CF₂CH₂O.

The term "prodrug esters" as employed herein includes esters and carbonates formed by reacting one or more hydroxyls of compounds of formula I with alkyl, alkoxy, or aryl substituted acylating agents employing procedures known to those skilled in the art to generate acetates, pivalates, methylcarbonates, benzoates and the like. In addition, prodrug esters which are known in the art for carboxylic and phosphorus acid esters such as methyl, ethyl, benzyl and the like.

Examples of such prodrug esters include



Where the compounds of structure I are in acid form they may form a pharmaceutically acceptable salt such as alkali metal salts such as lithium, sodium or potassium, alkaline earth metal salts such as calcium or magnesium as well as zinc or aluminum and other cations such as ammonium, choline, diethanolamine, lysine (D or L), ethylenediamine, t-butylamine, t-octylamine, tris-(hydroxymethyl)aminomethane (TRIS), N-methyl glucosamine (NMG), triethanolamine and dehydroabietylamine.

All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of the carbon atoms including any one of the R substituents. Consequently, compounds of formula I can exist in enantiomeric or diastereomeric forms or in mixtures thereof. The processes for preparation can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric or enantiomeric products are prepared, they can be separated by conventional methods for example, chromatographic or fractional crystallization.

Where desired, the compounds of structure I may be used in combination with one or more other types of antidiabetic agents and/or one or more other types of therapeutic agents which may be administered orally in the same dosage form, in a separate oral dosage form or by injection.

The other type of antidiabetic agent which may be optionally employed in combination with the SGLT2 inhibitor of formula I may be 1,2,3 or more antidiabetic agents or antihyperglycemic agents including insulin secretagogues or insulin sensitizers, or other antidiabetic agents preferably having a mechanism of action different from SGLT2 inhibition and may include biguanides, sulfonyl ureas, glucosidase inhibitors, PPAR γ agonists such as thiazolidinediones, α P2 inhibitors, PPAR α/γ dual agonists, dipeptidyl peptidase IV (DP4) inhibitors, and/or meglitinides, as well as insulin, glucagon-like peptide-1 (GLP-1), PTP1B inhibitors, glycogen phosphorylase inhibitors and/or glucos-6-phosphatase inhibitors.

The other types of therapeutic agents which may be optionally employed in combination with the SGLT2 inhibitors of formula I include anti-obesity agents, antihypertensive agents, antiplatelet agents,
5 antiatherosclerotic agents and/or lipid lowering agents.

The SGLT2 inhibitors of formula I may also be optionally employed in combination with agents for treating complications of diabetes. These agents include PKC inhibitors and/or AGE inhibitors.

10 It is believed that the use of the compounds of structure I in combination with 1, 2, 3 or more other antidiabetic agents produces antihyperglycemic results greater than that possible from each of these medicaments alone and greater than the combined additive anti-
15 hyperglycemic effects produced by these medicaments.

The other antidiabetic agent may be an oral antihyperglycemic agent preferably a biguanide such as metformin or phenformin or salts thereof, preferably metformin HCl.

20 Where the other antidiabetic agent is a biguanide, the compounds of structure I will be employed in a weight ratio to biguanide within the range from about 0.01:1 to about 100:1, preferably from about 0.1:1 to about 5:1.

The other antidiabetic agent may also preferably be
25 a sulfonyl urea such as glyburide (also known as glibenclamide), glimepiride (disclosed in U.S. Patent No. 4,379,785), glipizide, gliclazide or chlorpropamide, other known sulfonylureas or other antihyperglycemic agents which act on the ATP-dependent channel of the β -
30 cells, with glyburide and glipizide being preferred, which may be administered in the same or in separate oral dosage forms.

The compounds of structure I will be employed in a weight ratio to the sulfonyl urea in the range from about

WO 01/27128

PCT/US00/27187

0.01:1 to about 100:1, preferably from about 0.2:1 to about 10:1.

The oral antidiabetic agent may also be a glucosidase inhibitor such as acarbose (disclosed in U.S. Patent No. 4,904,769) or miglitol (disclosed in U.S. Patent No. 4,639,436), which may be administered in the same or in a separate oral dosage forms.

The compounds of structure I will be employed in a weight ratio to the glucosidase inhibitor within the range from about 0.01:1 to about 100:1, preferably from about 0.5:1 to about 50:1.

The compounds of structure I may be employed in combination with a PPAR γ agonist such as a thiazolidinedione oral anti-diabetic agent or other insulin sensitizers (which has an insulin sensitivity effect in NIDDM patients) such as troglitazone (Warner-Lambert's Rezulin[®], disclosed in U.S. Patent No. 4,572,912), rosiglitazone (SKB), pioglitazone (Takeda), Mitsubishi's MCC-555 (disclosed in U.S. Patent No. 5,594,016), Glaxo-Welcome's GL-262570, englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer, isaglitazone (MIT/J&J), JTT-501 (JPNT/P&U), L-805645 (Merck), R-119702 (Sankyo/WL), NN-2344 (Dr. Reddy/NN), or YM-440 (Yamanouchi), preferably rosiglitazone and pioglitazone.

The compounds of structure I will be employed in a weight ratio to the thiazolidinedione in an amount within the range from about 0.01:1 to about 100:1, preferably from about 0.2:1 to about 10:1.

The sulfonyl urea and thiazolidinedione in amounts of less than about 150 mg oral antidiabetic agent may be incorporated in a single tablet with the compounds of structure I.

The compounds of structure I may also be employed in combination with an antihyperglycemic agent such as

insulin or with glucagon-like peptide-1 (GLP-1) such as GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as disclosed in U.S. Patent No. 5,614,492 to Habener, the disclosure of which is incorporated herein by reference),
5 as well as AC2993 (Amylen) and LY-315902 (Lilly), which may be administered via injection, intranasal, or by transdermal or buccal devices.

Where present, metformin, the sulfonyl ureas, such as glyburide, glimepiride, glipyrider, glipizide,
10 chlorpropamide and gliclazide and the glucosidase inhibitors acarbose or miglitol or insulin (injectable, pulmonary, buccal, or oral) may be employed in formulations as described above and in amounts and dosing as indicated in the Physician's Desk Reference (PDR).

15 Where present, metformin or salt thereof may be employed in amounts within the range from about 500 to about 2000 mg per day which may be administered in single or divided doses one to four times daily.

Where present, the thiazolidinedione anti-diabetic
20 agent may be employed in amounts within the range from about 0.01 to about 2000 mg/day which may be administered in single or divided doses one to four times per day.

Where present insulin may be employed in formulations, amounts and dosing as indicated by the
25 Physician's Desk Reference.

Where present GLP-1 peptides may be administered in oral buccal formulations, by nasal administration or parenterally as described in U.S. Patent Nos. 5,346,701 (TheraTech), 5,614,492 and 5,631,224 which are
30 incorporated herein by reference.

The other antidiabetic agent may also be a PPAR α/γ dual agonist such as AR-HO39242 (Astra/Zeneca), GW-409544 (Glaxo-Wellcome), KRP297 (Kyorin Merck) as well as those disclosed by Murakami et al, "A Novel Insulin Sensitizer
35 Acts As a Coligand for Peroxisome Proliferation -

Activated Receptor Alpha (PPAR alpha) and PPAR gamma.
Effect on PPAR alpha Activation on Abnormal Lipid
Metabolism in Liver of Zucker Fatty Rats", Diabetes 47,
1841-1847 (1998), and in WO 2001/021602.

5

employing dosages as set out therein, which
compounds designated as preferred are preferred for use
herein.

10 The other antidiabetic agent may be an α P2 inhibitor
such as disclosed in WO 2000/059506

15 , employing dosages as set out
herein. Preferred are the compounds designated as
preferred in the above application.

The other antidiabetic agent may be a DP4 inhibitor
such as disclosed in WO99/38501, WO99/46272, WO99/67279
(PROBIODRUG), WO99/67278 (PROBIODRUG), WO99/61431
20 (PROBIODRUG), NVP-DPP728A (1-[[[2-[(5-cyanopyridin-2-
yl)amino]ethyl]amino]acetyl]-2-cyano-(S)-pyrrolidine)
(Novartis) (preferred) as disclosed by Hughes et al,
Biochemistry, 38(36), 11597-11603, 1999, TSL-225
(tryptophyl-1,2,3,4-tetrahydroisoquinoline-3-carboxylic
25 acid (disclosed by Yamada et al, Bioorg. & Med. Chem.
Lett. 8 (1998) 1537-1540, 2-cyanopyrrolidides and 4-
cyanopyrrolidides as disclosed by Ashworth et al, Bioorg.
& Med. Chem. Lett., Vol. 6, No. 22, pp 1163-1166 and
2745-2748 (1996) employing dosages as set out in the
30 above references.

The meglitinide which may optionally be employed in
combination with the compound of formula I of the
invention may be repaglinide, nateglinide (Novartis) or
KAD1229 (PF/Kissei), with repaglinide being preferred.

The SGLT2 inhibitor of formula I will be employed in a weight ratio to the meglitinide, PPAR γ agonist, PPAR α/γ dual agonist, $\alpha P2$ inhibitor or DP4 inhibitor within the range from about 0.01:1 to about 100:1, preferably
5 from about 0.2:1 to about 10:1.

The hypolipidemic agent or lipid-lowering agent which may be optionally employed in combination with the compounds of formula I of the invention may include 1,2,3 or more MTP inhibitors, HMG CoA reductase inhibitors,
10 squalene synthetase inhibitors, fibric acid derivatives, ACAT inhibitors, lipoxxygenase inhibitors, cholesterol absorption inhibitors, ileal Na^+ /bile acid cotransporter inhibitors, upregulators of LDL receptor activity, bile acid sequestrants, and/or nicotinic acid and derivatives
15 thereof.

MTP inhibitors employed herein include MTP inhibitors disclosed in U.S. Patent No. 5,595,872, U.S. Patent No. 5,739,135, U.S. Patent No. 5,712,279, U.S. Patent No. 5,760,246, U.S. Patent No. 5,827,875, U.S.
20 Patent No. 5,885,983 and U.S. Application Serial No. 09/175,180 filed October 20, 1998, now U.S. Patent No. 5,962,440. Preferred are each of the preferred MTP inhibitors disclosed in each of the above patents and applications.

25

The hypolipidemic agent may be an HMG CoA reductase inhibitor which includes, but is not limited to, mevastatin and related compounds as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related
30 compounds as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds such as disclosed in U.S. Patent No. 4,346,227, simvastatin and related compounds as disclosed in U.S. Patent Nos. 4,448,784 and 4,450,171. The hypolipidemic agent may also be the

compounds disclosed in WO 2001/096311 and WO 2001/096347.

Other HMG CoA reductase inhibitors which may be employed herein include, but are not limited to, fluvastatin, disclosed in U.S. Patent No. 5,354,772, cerivastatin disclosed in U.S. Patent Nos. 5,006,530 and 5,177,080, atorvastatin disclosed in U.S. Patent Nos. 4,681,893, 5,273,995, 5,385,929 and 5,686,104, atavastatin (Nissan/Sankyo's nisvastatin (NK-104)) disclosed in U.S. Patent No. 5,011,930, Shionogi-Astra/Zeneca visastatin (ZD-4522) disclosed in U.S. Patent No. 5,260,440, and related statin compounds disclosed in U.S. Patent No. 5,753,675, pyrazole analogs of mevalonolactone derivatives as disclosed in U.S. Patent No. 4,613,610, indene analogs of mevalonolactone derivatives as disclosed in PCT application WO 86/03488, 6-[2-(substituted-pyrrol-1-yl)-alkyl]pyran-2-ones and derivatives thereof as disclosed in U.S. Patent No. 4,647,576, Searle's SC-45355 (a 3-substituted pentanedioic acid derivative) dichloroacetate, imidazole analogs of mevalonolactone as disclosed in PCT application WO 86/07054, 3-carboxy-2-hydroxy-propane-phosphonic acid derivatives as disclosed in French Patent No. 2,596,393, 2,3-disubstituted pyrrole, furan and thiophene derivatives as disclosed in European Patent Application No. 0221025, naphthyl analogs of mevalonolactone as disclosed in U.S. Patent No. 4,686,237, octahydronaphthalenes such as disclosed in U.S. Patent No. 4,499,289, keto analogs of mevinolin (lovastatin) as disclosed in European Patent Application No. 0,142,146 A2, and quinoline and pyridine derivatives disclosed in U.S. Patent No. 5,506,219 and 5,691,322.

In addition, phosphinic acid compounds useful in inhibiting HMG CoA reductase suitable for use herein are disclosed in GB 2205837.

The squalene synthetase inhibitors suitable for use herein include, but are not limited to, α -phosphonosulfonates disclosed in U.S. Patent No. 5,712,396, those disclosed by Biller et al, J. Med. Chem., 1988, Vol. 31, No. 10, pp 1869-1871, including isoprenoid (phosphinylmethyl)phosphonates as well as other known squalene synthetase inhibitors, for example, as disclosed in U.S. Patent No. 4,871,721 and 4,924,024 and in Biller, S.A., Neuenschwander, K., Ponpipom, M.M., and Poulter, C.D., Current Pharmaceutical Design, 2, 1-40 (1996).

In addition, other squalene synthetase inhibitors suitable for use herein include the terpenoid pyrophosphates disclosed by P. Ortiz de Montellano et al, J. Med. Chem., 1977, 20, 243-249, the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-PP) analogs as disclosed by Corey and Volante, J. Am. Chem. Soc., 1976, 98, 1291-1293, phosphinylphosphonates reported by McClard, R.W. et al, J.A.C.S., 1987, 109, 5544 and cyclopropanes reported by Capson, T.L., PhD dissertation, June, 1987, Dept. Med. Chem. U of Utah, Abstract, Table of Contents, pp 16, 17, 40-43, 48-51, Summary.

Other hypolipidemic agents suitable for use herein include, but are not limited to, fibric acid derivatives, such as fenofibrate, gemfibrozil, clofibrate, bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds as disclosed in U.S. Patent No. 3,674,836, probucol and gemfibrozil being preferred, bile acid sequestrants such as cholestyramine, colestipol and DEAE-Sephadex (Secholex®, Policexide®), as well as lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphosphorylcholine (SPC, Roche), aminocyclodextrin (Tanabe Seiyoku), Ajinomoto AJ-814 (azulene derivative),

melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid, acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin,

5 poly(diallylmethylamine) derivatives such as disclosed in U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents.

10 The other hypolipidemic agent may be an ACAT inhibitor such as disclosed in, Drugs of the Future 24, 9-15 (1999), (Avasimibe); "The ACAT inhibitor, Cl-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters", Nicolosi et al,

15 Atherosclerosis (Shannon, Irel). (1998), 137(1), 77-85; "The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by selective suppression of the hepatic secretion of ApoB100-containing lipoprotein", Ghiselli, Giancarlo,

20 Cardiovasc. Drug Rev. (1998), 16(1), 16-30; "RP 73163: a bioavailable alkylsulfinyl-diphenylimidazole ACAT inhibitor", Smith, C., et al, Bioorg. Med. Chem. Lett. (1996), 6(1), 47-50; "ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic

25 activities in experimental animals", Krause et al, Editor(s): Ruffolo, Robert R., Jr.; Hollinger, Manfred A., Inflammation: Mediators Pathways (1995), 173-98, Publisher: CRC, Boca Raton, Fla.; "ACAT inhibitors: potential anti-atherosclerotic agents", Sliskovic et al,

30 Curr. Med. Chem. (1994), 1(3), 204-25; "Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. 6. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). 7.

35 Development of a series of substituted N-phenyl-N'-[(1-

phenylcyclopentyl)methyl]ureas with enhanced hypocholesterolemic activity", Stout et al, Chemtracts: Org. Chem. (1995), 8(6), 359-62, or TS-962 (Taisho Pharmaceutical Co. Ltd).

5 The hypolipidemic agent may be an upregulator of LD2 receptor activity such as MD-700 (Taisho Pharmaceutical Co. Ltd) and LY295427 (Eli Lilly).

 The hypolipidemic agent may be a cholesterol absorption inhibitor preferably Schering-Plough's
10 SCH48461 as well as those disclosed in Atherosclerosis 115, 45-63 (1995) and J. Med. Chem. 41, 973 (1998).

 The hypolipidemic agent may be an ileal Na⁺/bile acid cotransporter inhibitor such as disclosed in Drugs of the Future, 24, 425-430 (1999).

15 Preferred hypolipidemic agents are pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, cerivastatin, atavastatin and rosuvastatin.

 The amounts and dosages employed
20 will be as indicated in the Physician's Desk Reference and/or in the patents set out above.

 The compounds of formula I of the invention will be employed in a weight ratio to the hypolipidemic agent (where present), within the range from about 500:1 to
25 about 1:500, preferably from about 100:1 to about 1:100.

 The dose administered must be carefully adjusted according to age, weight and condition of the patient, as well as the route of administration, dosage form and regimen and the desired result.

30 The dosages and formulations for the hypolipidemic agent will be as disclosed in the various patents and applications discussed above.

 The dosages and formulations for the other hypolipidemic agent to be employed, where applicable,

will be as set out in the latest edition of the Physicians' Desk Reference.

For oral administration, a satisfactory result may be obtained employing the MTP inhibitor in an amount within the range of from about 0.01 mg/kg to about 500 mg and preferably from about 0.1 mg to about 100 mg, one to four times daily.

A preferred oral dosage form, such as tablets or capsules, will contain the MTP inhibitor in an amount of from about 1 to about 500 mg, preferably from about 2 to about 400 mg, and more preferably from about 5 to about 250 mg, one to four times daily.

For oral administration, a satisfactory result may be obtained employing an HMG CoA reductase inhibitor, for example, pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin in dosages employed as indicated in the Physician's Desk Reference, such as in an amount within the range of from about 1 to 2000 mg, and preferably from about 4 to about 200 mg.

The squalene synthetase inhibitor may be employed in dosages in an amount within the range of from about 10 mg to about 2000 mg and preferably from about 25 mg to about 200 mg.

A preferred oral dosage form, such as tablets or capsules, will contain the HMG CoA reductase inhibitor in an amount from about 0.1 to about 100 mg, preferably from about 5 to about 80 mg, and more preferably from about 10 to about 40 mg.

A preferred oral dosage form, such as tablets or capsules will contain the squalene synthetase inhibitor in an amount of from about 10 to about 500 mg, preferably from about 25 to about 200 mg.

The other hypolipidemic agent may also be a lipooxygenase inhibitor including a 15-lipooxygenase (15-LO) inhibitor such as benzimidazole derivatives as

disclosed in WO 97/12615, 15-LO inhibitors as disclosed
in WO 97/12613, isothiazolones as disclosed in
WO 96/38144, and 15-LO inhibitors as disclosed by
Sendobry et al "Attenuation of diet-induced
5 atherosclerosis in rabbits with a highly selective 15-
lipoxygenase inhibitor lacking significant antioxidant
properties, Brit. J. Pharmacology (1997) 120, 1199-1206,
and Cornicelli et al, "15-Lipoxygenase and its
Inhibition: A Novel Therapeutic Target for Vascular
10 Disease", Current Pharmaceutical Design, 1999, 5, 11-20.

The compounds of formula I and the hypolipidemic
agent may be employed together in the same oral dosage
form or in separate oral dosage forms taken at the same
time.

15 The compositions described above may be administered
in the dosage forms as described above in single or
divided doses of one to four times daily. It may be
advisable to start a patient on a low dose combination
and work up gradually to a high dose combination.

20 The preferred hypolipidemic agents are pravastatin,
simvastatin, lovastatin, atorvastatin, fluvastatin,
cerivastatin, atavastatin and rosuvastatin.

When the other type of therapeutic agent which may
be optionally employed with the SGLT2 inhibitor of
25 formula I is 1, 2, 3 or more of an anti-obesity agent, it
may include a beta 3 adrenergic agonist, a lipase
inhibitor, a serotonin (and dopamine) reuptake inhibitor,
a thyroid receptor beta drug, an anorectic agent, an NPY
antagonist, a Leptin analog and/or an MC4 agonist.

30 The beta 3 adrenergic agonist which may be
optionally employed in combination with a compound of
formula I may be AJ9677 (Takeda/Dainippon), L750355
(Merck), or CP331648 (Pfizer) or other known beta 3
agonists as disclosed in U.S. Patent Nos. 5,541,204,

5,770,615, 5,491,134, 5,776,983 and 5,488,064, with
AJ9677, L750,355 and CP331648 being preferred.

The lipase inhibitor which may be optionally
employed in combination with a compound of formula I may
5 be orlistat or ATL-962 (Alizyme), with orlistat being
preferred.

The serotonin (and dopamine) reuptake inhibitor
which may be optionally employed in combination with a
compound of formula I may be sibutramine, topiramate
10 (Johnson & Johnson) or axokine (Regeneron), with
sibutramine and topiramate being preferred.

The thyroid receptor beta compound which may be
optionally employed in combination with a compound of
formula I may be a thyroid receptor ligand as disclosed
15 in WO97/21993 (U. Cal SF), WO99/00353 (KaroBio) and
GB98/284425 (KaroBio), with compounds of the KaroBio
applications being preferred.

The anorectic agent which may be optionally employed
in combination with a compound of formula I may be
20 dexamphetamine, phentermine, phenylpropanolamine or
mazindol, with dexamphetamine being preferred.

The various anti-obesity agents described above may
be employed in the same dosage form with the compound of
formula I or in different dosage forms, in dosages and
25 regimens as generally known in the art or in the PDR.

Examples of the anti-platelet agent(s) which may be
optionally employed in combinations of this invention
include abciximab, ticlopidine, eptifibatide,
dipyridamole, aspirin, anagrelide, tirofiban and/or
30 clopidogrel.

Examples of the anti-hypertensive agent(s) which may
be optionally employed in combinations of this invention
include ACE inhibitors, calcium antagonists, alpha-
blockers, diuretics, centrally acting agents,

angiotensin-II antagonists, beta-blockers and vasopeptidase inhibitors.

Examples of ACE inhibitors include lisinopril, enalapril, quinapril, benazepril, fosinopril, ramipril, 5 captopril, enalaprilat, moexipril, trandolapril and perindopril; examples of calcium antagonists include amlodipine, diltiazem, nifedipine, verapamil, felodipine, nisoldipine, isradipine and nicardipine; examples of alpha-blockers include terazosin, doxazosin and prazosin; 10 examples of diuretics include hydrochlorothiazide, torasemide, furosemide, spironolactone and indapamide; examples of centrally acting agents include clonidine and guanfacine; examples of angiotensin-II antagonists include losartan, valsartan, irbesartan, candesartan and 15 telmisartan; examples of beta-blockers include metoprolol, propranolol, atenolol, carvedilol and sotalol; and examples of vasopeptidase inhibitors include omapatrilat and gemopatrilat.

In carrying out the method of the invention, a 20 pharmaceutical composition will be employed containing the compounds of structure I, with or without another antidiabetic agent and/or antihyperlipidemic agent, or other type therapeutic agent, in association with a pharmaceutical vehicle or diluent. The pharmaceutical 25 composition can be formulated employing conventional solid or liquid vehicles or diluents and pharmaceutical additives of a type appropriate to the mode of desired administration. The compounds can be administered to mammalian species including humans, monkeys, dogs, etc. 30 by an oral route, for example, in the form of tablets, capsules, granules or powders, or they can be administered by a parenteral route in the form of injectable preparations, or they can be administered intranasally or in transdermal patches. The dose for 35 adults is preferably between 10 and 2,000 mg per day,

which can be administered in a single dose or in the form of individual doses from 1-4 times per day.

A typical injectable preparation is produced by aseptically placing 250 mg of compounds of structure I
5 into a vial, aseptically freeze-drying and sealing. For use, the contents of the vial are mixed with 2 mL of physiological saline, to produce an injectable preparation.

SGLT2 inhibitor activity of the compounds of the
10 invention may be determined by use of an assay system as set out below.

Assay for SGLT2 Activity


15 The mRNA sequence for human SGLT2 (GenBank #M95549) was cloned by reverse-transcription and amplification from human kidney mRNA, using standard molecular biology techniques. The cDNA sequence was stably transfected into CHO cells, and clones were assayed for SGLT2
20 activity essentially as described in Ryan et al. (1994). Evaluation of inhibition of SGLT2 activity in a clonally selected cell line was performed essentially as described in Ryan et al., with the following modifications. Cells were grown in 96-well plates for 2-4 days to 75,000 or
25 30,000 cells per well in F-12 nutrient mixture (Ham's F-12), 10% fetal bovine serum, 300 ug/ml Geneticin and penicillin-streptomycin. At confluence, cells were washed twice with 10 mM Hepes/Tris, pH 7.4, 137 mM N-methyl-D-glucamine, 5.4 mM KCl, 2.8 mM CaCl₂, 1.2 mM MgSO₄. Cells
30 then were incubated with 10 μ M [¹⁴C]AMG, and 10 μ M inhibitor (final DMSO =0.5%) in 10 mM Hepes/Tris, pH 7.4, 137 mM NaCl, 5.4 mM KCl, 2.8 mM CaCl₂, 1.2 mM MgSO₄ at 37°C for 1.5 hr. Uptake assays were quenched with ice cold 1X PBS containing 0.5 mM phlorizin, and cells were
35 then lysed with 0.1% NaOH. After addition of MicroScint

scintillation fluid, the cells were allowed to shake for 1 hour, and then [^{14}C]AMG was quantitated on a TopCount scintillation counter. Controls were performed with and without NaCl. For determination of EC_{50} values, 10
5 inhibitor concentrations were used over 2 log intervals in the appropriate response range, and triplicate plates were averaged across plates.

Ryan MJ, Johnson G, Kirk J, Fuerstenberg SM, Zager RA and
10 Torok-Storb B. **1994**. HK-2: an immortalized proximal tubule epithelial cell line from normal adult human kidney. *Kidney International* **45**: 48-57.

The following Working Examples represent preferred
15 embodiments of the present invention. All temperatures are expressed in degrees Centigrade unless otherwise indicated.

Example 1



A. 3-Bromo-4'-ethylbenzylhydroxide


Dry Mg turnings (4.4g, 0.178 mol) under Ar were
25 stirred overnight whereupon 100 mL of dry Et_2O was added followed by addition over 1 hr of *p*-bromoethylbenzene (22g, 0.119 mol) in 20 mL of Et_2O . (In the event the reaction did not start, 0.5 ml of 1,2-dibromoethane was added). After stirring overnight, *m*-bromobenzaldehyde
30 (11g, 0.06 mol) in 20 mL of Et_2O was slowly added. The resulting light solution was monitored by HPLC over 4-6

hr to determine when complete. The reaction, after quenching with saturated aq. NH_4Cl , was extracted 3x with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated using a rotary
 5 evaporator. The resulting yellow oil was chromatographed on silica gel using 5% EtOAc/hexane to elute nonpolar impurities and 7-9% EtOAc/hexane to elute 12.4 g (71%) of 3-bromo-4'-ethylbenzhydrol as a light yellow oil.

10 B. 3-Bromo-4'-ethyldiphenylmethane

To a stirred -30° solution of Part A 3-bromo-4'-ethylbenzhydrol (12.4g, 0.0426 mol) in 120 mL of MeCN was added $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (6.04g, 0.0426 mol) followed by Et_3SiH (9.9g, 0.852 mol). The dark reaction after stirring 1 hr
 15 at -30° was warmed slowly to -5° . When complete by tlc, the reaction was quenched by addition of saturated aq. K_2CO_3 . After addition of 100 mL of H_2O , the mixture was extracted 3x with Et_2O . The combined organic layers were washed with brine, dried over Na_2SO_4 . After concentration
 20 using a rotary evaporator, 3-bromo-4'-ethyldiphenylmethane (11.17g, 95%) was obtained as a light yellow oil that was used without further purification.

C.




25

To a stirred -78° solution of Part B 3-bromo-4'-ethyldiphenylmethane (10.9g, 0.04 mol) in 100 mL of dry THF under Ar was added 25.7 mL of 1.7 M $t\text{-BuLi}$ in hexane
 30 over 20 min. After 1hr 2,3,4,6-tetra-O-benzyl- $\beta\text{-D-}$

glucolactone (23.5 g, 0.0437 mol) in 30 mL of THF was added over 15 min. The solution was stirred for 1 hr at -78° prior to quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 2 fold with EtOAc prior to washing with H_2O followed by brine. After drying over Na_2SO_4 and concentration using a rotary evaporator, 29.2 g of the desired title lactol was obtained as a colorless syrup that was carried forward without further purification.

10

D.




To a stirred -30° solution of Part C lactol (29.1g, 0.04 mol) in 100 mL of MeCN was added $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (5.62g, 0.04 mol) followed by Et_3SiH (9.21g, 0.08 mol). After 2 hr, when tlc showed the reaction to be complete, saturated aq. K_2CO_3 was added and the suspension stirred 1 hr at 20° prior to diluting with H_2O and Et_2O . The combined organic layers from 3 Et_2O extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator to yield 28.3 g of a light yellow syrup. Chromatography on silica gel with 5% EtOAc/hexane eluted nonpolar impurities followed slowly by the desired beta anomer and then the alpha anomer. Fractions enriched in the beta anomer could be further purified by either triterating with hexane or by recrystallization from EtOH to yield 6 g of the desired title beta tetra-O-benzyl C-glucoside. (Note when Et_3SiH is the reducing agent, a 5:1 beta/alpha anomer mixture is

25

30

obtained whereas when $i\text{Pr}_3\text{SiH}$ is substituted a 30:1 mixture is obtained.)

E.



5

A solution of Part D tetra-O-benzyl C-glucoside (2.4g, 3.35 mmol) in EtOAc (100 mL) containing 10% Pd(OH)₂/C (0.35 g) was stirred overnight under 1 atmos. H₂. After HPLC showed the reaction to be complete, the catalyst was filtered and the solvent removed using a rotary evaporator to obtain 1.1 g of the desired beta C-glucoside (92%) as a white crystalline solid.

15

HPLC retention time: 7.04 min, 100% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 8 min gradient 0-100% B hold 5 min at 100% B. Solvent A: 10% MeOH/H₂O + 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.

20

¹H NMR (500 MHz, CD₃OD) δ 7.27 (s, 1H), 7.23 (d, 2H, J=4.95 Hz), 7.1-7.0 (m, 5H), 4.08 (d, 1H, J=9.3Hz), 3.91 (s, 2H), 3.9 (dd, 1H, J=2.2, 11 Hz), 3.68 (dd, 1H, J=5.5, 11.5 Hz), 3.5-3.35 (m, 4H), 2.57 (q, 2H, J=7.2 Hz), 1.18 (t, 3H, J=7.2 Hz)


25

¹³C NMR (125 MHz, CD₃OD) δ 143, 142.8, 141, 140, 129.9, 129.6, 129.5, 129.1, 128.8, 126.7, 83.8, 82.3, 79.9, 76.4, 72.0, 63.2, 42.5, 29.4, 16.2.

30

Anal Calcd for $C_{21}H_{26}O_5$ LC-MS $[M+NH_4]$ 376; found 376.

Example 2



A. 3-Bromo-4'-methoxybenzhydrol


To a stirred -78° solution of *m*-dibromobenzene (70.9g, 0.3 mol) in 200 mL of dry THF under Ar was added 117 mL of 2.56 M *n*-BuLi (0.3 mol) in hexane over 10 min. After 30 min, *p*-methoxybenzaldehyde (27.2 g, 0.02 mol) in 50 mL of THF was added over 20 min. The solution was stirred for 1 hr at -78° (complete by tlc) prior to quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 2 fold with EtOAc prior to washing with H_2O followed by brine. After drying over Na_2SO_4 and concentration using a rotary evaporator, 103 g of 3-bromo-4'-methoxybenzhydrol was obtained as a yellow oil that was carried forward without further purification.

B. 3-Bromo-4'-methoxydiphenylmethane

To a stirred -40° solution of crude Part A 3-bromo-4'-methoxybenzhydrol (103g, 0.2 mol) in 300 mL of MeCN was added Et_3SiH (64 mL, 0.4 mol) followed by $BF_3 \cdot Et_2O$ (27.7g, 0.2 mol). When complete by tlc, the reaction was quenched by addition of saturated aq. K_2CO_3 (25 mL). After addition of 100 mL of H_2O , the mixture was extracted 3x with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 . After concentration using a rotary evaporator, the crude title 3-bromo-4'-


methoxydiphenylmethane (92g) was chromatographed on silica gel using 9% EtOAc/hexane to eluted 17 g of clean product followed by less pure fractions.

5 c.




To a stirred -78° solution of Part B 3-bromo-4'-
10 methoxydiphenylmethane (9.6g, 0.035 mol) in 50 mL of dry THF under Ar was added 14 mL of 2.5 M *n*-BuLi in hexane over 5 min. After stirring 30 min, 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (12.5 g, 0.023 mol) in 20 mL of THF was added over 10 min. The solution was stirred for
15 1 hr at -78° whereupon tlc analysis indicated the reaction was complete. After quenching with saturated aq. NH_4Cl (25 mL) and warming to 20° , the reaction was diluted with EtOAc (200 mL). The organic layer was washed with H_2O followed by brine. After drying over Na_2SO_4 and
20 concentration using a rotary evaporator, the desired title lactol was chromatographed on silica gel using 12.5% EtOAc/hexane to elute 8.1 g of >90% lactol followed by 9.7g of >80% purity.

D.



- 5 To a stirred -40° solution of Part C lactol (7.8g, 0.019 mol) in 100 mL of MeCN was added Et_3SiH (3.42 mL, 0.04 mol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (1.37 mL, 0.02 mol). After 1 hr, when tlc showed the reaction to be complete, saturated aq. K_2CO_3 (10mL) was added and the suspension
- 10 stirred 1 hr at 20° prior to extracting 3x with EtOAc. The combined organic layers were washed with H_2O , brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator to yield 8 g of crude product. Chromatography on silica gel with 5% EtOAc/hexane eluted nonpolar
- 15 impurities followed by 0.92 g of pure title β -tetra-O-benzyl C-glucoside followed by a 6.5g containing both anomers.

E.



20

- The above two fractions of Part D compound were hydrogenated separately over 10% $\text{Pd}(\text{OH})_2$ (2% by weight) for overnight at 1 atmosphere H_2 in EtOAc (12.5 mL/g of Part D compound). After filtration and solvent removal, the hydrogenolysis product of the mixed fractions was
- 25


purified by prep HPLC using a YMC S10 reverse phase column. The combined material yielded 1.85 g of pure β anomer as a white solid.

- 5 HPLC retention time: 6.04 min, Zorbax C-18 4.6x75mm column, 2.5 mL/min, detection at 220nm; 8 min gradient 0-100% B hold 3 min at 100% B. Solvent A: 10% MeOH/H₂O + 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.
- 10 ¹H NMR (400 MHz, CD₃OD) δ 7.28 (s, 1H), 7.24 (d, 2H, J=3Hz), 7.09 (m, 3H), 6.79 (d, 2H, J=7Hz), 4.08 (d, 1H, J= 8.8 Hz), 3.88 (s, 2H), 3.75 (d, 1H, J=12 Hz), 3.73 (s, 3H), 3.65 (dd, 1H, J=12, 3 Hz), 3.4 (m, 4H).
- 15 ¹³C NMR (100MHz, CD₃OD) δ 158.6, 142.1, 140.2, 133.8, 130.0, 128.7, 128.6, 128.3, 125.8, 82.9, 81.3, 79.0, 75.5, 71.1, 62.5, 55.1, 41.1.


Anal Calcd for C₂₀H₂₄O₆ LC-MS (M-H) 359; found 359.

20

Example 3




A.



25


To a stirred -78° solution of *m*-dibromobenzene (12.6g, 53 mmol) in 50 mL of dry THF under Ar was added 20 mL of 2.56 M *n*-BuLi (51 mmol) in hexane over 10 min. After 40 min, 2,3,4,6-tetra-*O*-benzyl- β -D-glucolactone (12 g, 22 mmol) in 30 mL of THF was added over 15 min. The solution was stirred for 1 hr at -78° (complete by tlc) prior to quenching with saturated aq. NH_4Cl (40 mL). After warming to 20° , the reaction was diluted 2 fold with EtOAc prior to washing with H_2O followed by brine. After drying over Na_2SO_4 and concentration using a rotary evaporator, 20 g of crude title lactol was obtained as an oil that was carried forward without further purification.

15 B.




To a stirred -45° solution of crude Part A lactol (20g, 0.2 mol) in 60 mL of MeCN was added Et_3SiH (7.8 mL, 45 mmol) followed by slow addition over 20 min of $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (4.2 mL, 22 mmol). When complete by tlc after an hour, the reaction was quenched by addition of saturated aq. K_2CO_3 (25 mL) and the mixture was extracted 3x with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated using a rotary evaporator. The resulting oil was triterated with 50 mL hexane whereupon solid precipitated after standing for 1 hr. This material was collected by filtration, washed with cold hexane twice and air dried to yield 8.9 g of the desired title β -*m*-bromophenyl-C-glucoside.

C.




5 A solution of Part B β -*m*-bromophenyl C-glucoside (1.36 g, 2 mmol), $\text{Pd}(\text{PPh}_3)_4$ (70 mg, 0.06 mmol), and hexabutylstannane (2.724 g, 6 mmol) in dry toluene (10 mL) was heated with stirring under Ar at 80° for 15 hr. After removal of toluene using a rotary evaporator, the residue was chromatographed on silica gel using 12:1 EtOAc/hexane to elute the desired title aryl stannane (761 mg), plus mixed fractions, which after a second column yielded an additional 92 mg of clean title stannane for a total yield of 48%, followed by 230 mg of recovered starting Part B β -*m*-bromophenyl-C-glucoside.

D.



20 A mixture of Part E stannane (2.66 g, 3 mmol), *p*-trifluoromethoxybenzyl chloride (1.04 g, 6 mmol), and $\text{Pd}(\text{PPh}_3)_4$ (100 mg, 0.09 mmol) was refluxed under Ar in THF (1 mL) for 15 hr. After removal of THF with a rotary evaporator, the residue was chromatographed on silica gel using 10:1 hexane/EtOAc to elute 1.3 g of the desired title tetrabenzyl ether.

E.



5

Conversion to the final free glucoside was achieved by stirring 295 mg of Part D tetrabenzyl ether with Pd(OH)₂ (15 mg) in EtOAc (3 mL) under 1 atmos of H₂ for 15 hr. The title product (104 mg) was isolated after
 10 filtration, Prep HPLC, and removal of solvent.


HPLC retention time: 7.21 min, Zorbax C-18 4.6x75mm column, 2.5 mL/min, detection at 220nm; 8 min gradient 0-100% B hold 3 min at 100% B. Solvent A: 10% MeOH/H₂O +
 15 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.

¹H NMR (400 MHz, CD₃OD) δ 7.3 (m, 5H), 7.15 (m, 3H), 4.10 (d, 1H, J= 8.8 Hz), 3.99 (s, 2H), 3.9 (d, 1H, J=12 Hz), 3.7 (dd, 1H, J=12, 3 Hz), 3.4 (m, 4H).

20


Anal Calcd for C₂₀H₂₁F₃O₆ LC-MS (M-H) 413; found 413

Example 4



25


A.



5 A mixture of Example 3 Part B β -*m*-bromophenyl-C-glucoside (3.0 g, 4.41 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (153 mg, 0.13 mmol), and hexabutyl-distannane (6.0 g, 13.2 mmol) in dry toluene (5 mL) was heated with stirring under Ar at 88° for 3 hr whereupon tlc analysis indicated the reaction
 10 was 90% complete. The reaction was terminated after a total of 5 hr. After removal of toluene using a rotary evaporator, the residue was chromatographed on silica gel using 1:8 EtOAc/hexane to elute the 2.95 g of desired aryl stannane.


15

B.



20 A mixture of Part A stannane (2.66 g, 3 mmol), *p*-methylthiobenzyl chloride (1.04 mg, 6.0 mmol), and tetrakis(triphenylphosphine)palladium (100 mg, 0.09 mmol) was refluxed under Ar in THF (5 mL) for 15 hr. After removal of THF with a rotary evaporator, the residue was
 25 chromatographed on silica gel using 6:1 hexane/EtOAc to elute 1.2 g of the desired title tetra-O-benzyl ether followed by 600 mg of title tetra-O-benzylether containing Ph_3P .

C.




5 1 M $\text{BCl}_3/\text{CH}_2\text{Cl}_2$ (6 mL, 8 mmol) was added over 5 minutes to a stirred -78° solution of Part B tetrabenzyl ether (295 mg, 0.4 mmol) under Ar in CH_2Cl_2 (0.25 mL). After 30 min, when tlc analysis indicated the reaction was complete, 30 mL of 2:1 $\text{CH}_2\text{Cl}_2/\text{PhMe}$ followed by 2 mL of
 10 MeOH were added. The volume was reduced by half using a rotary evaporator and 10 mL of MeOH added. After repeating this process 3x, all the volatiles were removed under vacuum. The residue was chromatographed on silica gel using 5% MeOH/ CH_2Cl_2 to eluted 143 mg of the desired
 15 glucoside in 90% purity. This material was further purified by reverse phase preparative HPLC to yield 104 mg of the final desired glucoside.

HPLC retention time: 6.69 min, Zorbax C-18 4.6x75mm
 20 column, 2.5 mL/min, detection at 220nm; 8 min gradient 0-100% B hold 3 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .


^1H NMR (400 MHz, CD_3OD) δ 7.27 (s, 1H), 7.25 (d, 2H, $J=2\text{Hz}$), 7.15 (m, 5H), 4.09 (d, 1H, $J=8.8\text{ Hz}$), 3.92 (s, 2H), 3.86 (d, 1H, $J=12\text{ Hz}$), 3.68 (dd, 1H, $J=12, 3\text{ Hz}$), 3.4 (m, 4H), 2.43 (s, 3H).

Anal Calcd for $\text{C}_{20}\text{H}_{24}\text{O}_6\text{S}$ LC-MS (M-H) 375; found 375

30

Example 5


A.



5

To a stirred suspension of 60% NaH (180 mg, 4.5 mmol) in THF (7 mL) under Ar was added 2-bromophenol (350 μ L, 3 mmol). After stirring for 15 min, the reaction was cooled to -78° and 1.4 M *t*-BuLi/hexane (2.36 mL, 3.3 mmol) was added dropwise. After 10 min, the solution was transferred via cannula to a stirred -78° solution of 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (1.62 g, 3.0 mmol) in THF (5 mL). The reaction was quenched after 15 min by slow addition of sat. $\text{NH}_4\text{Cl}/\text{H}_2\text{O}$ and then allowed to warm to 20° whereupon 200 mL of EtOAc was added. The organic layer was washed successively with H_2O and brine, dried over MgSO_4 , and concentrated. Chromatography on silica gel with 3:1 hexane/EtOAc yielded 390 mg of the desired title lactol.

B.




25

To a stirred 3:1 mixture of MeCN/CH₂Cl₂ (4 mL) containing Part A lactol (390 mg, 0.62 mmol) at -30° was added Et₃SiH (197 µL, 1.23 mmol) and BF₃·Et₂O (78 µL, 0.62 mmol). After 1 hr the reaction was quenched by addition
5 of 1 mL of sat. K₂CO₃, warmed to 20° and diluted with 100 mL EtOAc. The organic layer was washed successively with H₂O and brine, dried over MgSO₄, and concentrated. Chromatography on silica gel with 3:1 hexane/EtOAc yielded 269 mg of desired title phenolic C-glucoside.


10

C.



15 To a PhMe solution (1.1 mL) of Part B phenol (139 mg, 0.22 mmol) under Ar was added 60% NaH (11 mg, 0.27 mmol). After 10 min, 4-methylbenzyl bromide (46 mg, 0.25 mmol) was added as a solid to the blue solution which was then heated at 80° for 3.5 hr until complete by tlc
20 analysis. After cooling followed by addition of aqueous NH₄Cl, the reaction was diluted with EtOAc. The organic layer was washed successively with H₂O and brine, dried over MgSO₄, and concentrated. Chromatography on silica gel with 5:1 hexane/EtOAc yielded 71 mg of the desired
25 title tetra-O-benzylglucoside.

D.




5 Subsequent hydrogenolysis of Part C tetra-O-benzyl
glucoside over Pd/C in MeOH under 1 atmos H₂ yielded the
final title product which was purified by preparative
HPLC using a C18 reverse phase column a 45-90% MeOH/H₂O
gradient over 10 min to elute the desired β -C-glucoside
10 (2 mg).

HPLC retention time: 6.754 min, 100% pure, YMC S3 ODS
4.6x50mm, 2.5 mL/min, detection at 220nm; 8 min gradient
0-100% B hold 5 min at 100% B. Solvent A: 10% MeOH/H₂O +
15 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.

¹H NMR (500 MHz, CD₃OD) δ 7.15 (dd, 1H, J = 1.1, 7.7 Hz),
7.07 (d, 2H, J= 8.3 Hz), 7.02 (d, 2H, J=8.3 Hz), 6.96
(dd, 1H, J=1.2, 7.7 Hz), 6.77 (t, 1H, J= 7.7 Hz), 4.44
20 (d, 1H, J= 8.8 Hz), 3.89 (s, 2H), 3.87 (d, 1H, J=2.2 Hz),
3.75 (dd, 1H, J=4.9, 12.1), 3.49-3.41 (m, 4H), 2.26 (s,
3H).


Anal Calcd for C₂₀H₂₄O₆ LC-MS [M+H] 361; found 361.

25

Example 65 A. p-Chloromethylacetophenone

To a stirred solution of *p*-chloromethylbenzoyl chloride (390 mg, 2.06 mmol) in 8 mL THF at -20° under Ar was added tributylphosphine (406 mg, 2.29 mMol). After stirring the resulting yellow solution for 20 min at -20°
 10 -15° , 0.7 mL of 3M methyl magnesium bromide in ether (2.1 mmol) was added in one portion to generate a red solution which subsequently became orange over a 10 min period. The reaction was quenched by addition of 1N aq. HCl. After dilution with H₂O, the mixture was extracted
 15 3x with EtOAc, washed with H₂O prior to drying over Na₂SO₄. The residue obtained after removal of volatiles was chromatographed on silica gel using 5% EtOAc/hexane to elute 171 mg (50%) of *p*-chloromethylacetophenone.


20 B.



A mixture of the stannane described in Example 3
 25 Part C (300 mg, 0.33 mmol), *p*-chloromethylacetophenone (114 mg, 0.66 mmol), and Pd(PPh₃)₄ (20 mg, 0.09 mmol),

triphenylphosphine oxide (180 mg, 0.65 mmol), K_2CO_3 (75 mg, 0.55 mmol) was heated at 70° under Ar in THF (0.3 ml) for 16 hr. After removal of THF with a rotary evaporator, the residue was chromatographed on silica gel using 20:1 to 10:1 hexane/EtOAc to elute the desired tetrabenzyl ether (170 mg, 70%).

C.



10

A solution of Part B tetrabenzyl ether (60 mg, 0.08 mmol) in CH_2Cl_2 (5 mL) under Ar was cooled to -78° prior to the addition of 0.8 mL of 1 M BCl_3 in CH_2Cl_2 . After stirring for 1 hr at -78° , a second 0.8 mL portion of 1 M BCl_3 was added to the stirred reaction. After a second hour, 0.5 mL of PhMe was added followed by dropwise addition of 0.5 mL of MeOH. The volatiles were removed using a rotary evaporator; the process repeated after addition of 3 mL of a 2:1 mixture of CH_2Cl_2 /MeOH. Chromatography of the resulting residue on silica gel eluting with 5% MeOH/EtOAc yielded 20 mg of tetraol final product in 67% yield.

HPLC retention time: 2.35 min, 100% pure, YMC S3 ODS 4.6x50mm, 2.5 mL/min, detection at 220nm; 4 min gradient 0-100% B hold 4 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .


^1H -NMR (500 MHz, CD_3OD): δ 7.88 (d, 2H), 7.27-7.34 (m, 5H), 7.13 (d, 1H), 4.09 (d, 1H), 4.03 (s, 2H), 3.85 (d, 1H), 3.68 (dd, 1H), 3.35-3.48 (m, 4H), 2.55 (s, 3H)

5 ^{13}C -NMR (500 MHz, CD_3OD): δ 200.3, 148.8, 141.4, 141.2, 136.3, 130.2, 129.7, 129.6, 129.3, 127.0, 83.6, 82.2, 79.8, 76.4, 71.9, 63.1, 42.7, 26.6

Anal Calcd for $\text{C}_{21}\text{H}_{24}\text{O}_6$ LC-MS ($\text{M}+\text{NH}_4^+$): 390.2; found: 390.2

10

Example 7



15

A stirred solution of the final product of Example 6 (15 mg, 0.04 mmol) in 5 mL of EtOH was cooled to -20° whereupon NaBH_4 (5 mg, 0.13 mmol) was added. After 20 min being complete by tlc analysis, the reaction was quenched with a few drops of saturated aq. NH_4Cl . After removal of the volatiles, the residue was chromatographed on silica gel. Elution with 5% MeOH/EtOAc yielded 10 mg (67%) of the desired product.

20

25

HPLC retention time: 5.2 min, 100% pure, YMC S3 ODS 4.6x50mm, 2.5 mL/min, detection at 220nm; 8 min gradient 0-100% B hold 5 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .


^1H -NMR (500 MHz, CD_3OD): δ 7.21–7.32 (m, 5H), 7.16 (d, 2H), 7.10–7.11 (m, 1H), 4.77 (q, 1H), 4.08 (d, 1H), 3.94 (s, 2H), 3.86 (dd, 1H), 3.68 (dd, 1H), 3.34–3.48 (m, 4H), 1.40 (d, 3H)

5 ^{13}C -NMR (500 MHz, CD_3OD): δ 145.2, 142.5, 141.5, 140.9, 129.8, 129.6, 129.5, 129.2, 126.7, 126.6, 83.7, 82.2, 79.8, 76.4, 72.0, 63.2, 42.5, 25.5

Anal Calcd for $\text{C}_{21}\text{H}_{26}\text{O}_6$ LC-MS ($\text{M}+\text{NH}_4^+$): 392.2; found: 392.1

10

Example 8



A. 5-Bromo-2-methylbenzoic Acid

A mixture of *o*-toluic acid (28g, 206mmol), iron powder (0.74g, 13mmol), and Br_2 (42g, 260 mmol) were stirred at 0° for 2 hr. At this point the reaction, which had proceeded ~40%, was diluted with 25 mL of CH_2Cl_2 to facilitate stirring. The reaction was then heated at 45° for 16 hr to drive to completion. Upon cooling, the reaction was diluted with CH_2Cl_2 , washed 2x with 10% NaHSO_3 , 1x with brine prior to drying over Na_2SO_4 . After removal of the volatiles, the residue comprising a 2:1 mixture of 5-bromo to 3-bromotoluic acid was recrystallized from 95% EtOH to yield 14.4g of 5-bromo-2-methylbenzoic acid.

B. 5-Bromo-2-methyl-4'-methoxybenzophenone

To a stirred suspension of 5-bromo-2-methylbenzoic acid (1.29 g, 6 mmol) in 12 mL of CH_2Cl_2 containing oxalyl chloride (8 mmol) was added 2 drops of DMF. Once the


30

vigorous evolution of gas ceased, the reaction was stirred 6 hr prior to removal of the volatiles using a rotary evaporator. After dissolving the crude 5-bromo-2-methylbenzoyl chloride in 15 ml of CS₂, the stirred
5 mixture was cooled to 4° prior to adding anisole (0.7 g, 6.6 mmol) followed by AlCl₃ (1.7 g, 12 mmol). The reaction, after warming to 20° over 1 hr, was stirred for 15 hr prior to quenching with 1N HCl. Subsequently, the suspension was diluted with 50 ml H₂O and stirred until
10 all solids were in solution. The mixture was extracted 3x with EtOAc. The combined organic extracts were washed 1x with 1N HCl, H₂O, aq NaHCO₃, and brine prior to drying over Na₂SO₄. After removal of the volatiles, the resulting tan solid was recrystallized from 95% EtOH to
15 yield 1.6g of 5-bromo-2-methyl-4'-methoxybenzophenone.

C. 5-Bromo-2-methyl-4'-methoxydiphenylmethane


A solution of Et₃SiH (2.5 mL, 15.5 mmol), BF₃·Et₂O (1.3 mL, 10 mmol), and 5-bromo-2-methyl-4'-
20 methoxybenzophenone (1.6g, 5.25 mmol) in 11 mL of a 1:4 mixture CH₂Cl₂/MeCN was stirred overnight at 20°. Since by HPLC 5% of starting ketone remained, the solution was heated to 40° for 1 hr prior to quenching with 10% NaOH. After dilution with H₂O, the reaction was extracted 3x
25 with EtOAc. The combined organic layers were washed 2x with H₂O and once with brine before drying over Na₂SO₄. After removal of the volatiles, the residue was chromatographed on silica gel using hexane to elute 5-bromo-2-methyl-4'-methoxydiphenylmethane as a colorless
30 oil (1.4g, 95%)

D.



5 To a stirred -78° solution of Part C 5-bromo-2-methyl-4'-methoxydiphenylmethane (0.43g, 1.5 mmol) in 7 mL of dry THF under Ar was added dropwise 0.9 mL of 1.8 M *n*-BuLi in hexane. After 2 hr, 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (0.88g, 1.6 mmol) in 3 mL of THF was added
 10 over 1 min. The solution was stirred for 2 hr at -78° prior to quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 2 fold with H_2O prior to 3 extractions with EtOAc. The combined EtOAc fractions were washed with brine and dried over Na_2SO_4 .
 15 After concentration using a rotary evaporator, 1.1 g of the desired title lactol was obtained as a colorless syrup that was carried forward without further purification.


20 E.



To a stirred -30° solution of Part D lactol (1.1g, 1.47 mmol) in 10 mL of MeCN was added $i\text{Pr}_3\text{SiH}$ (0.7g, 4.5 mmol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.38g, 2.6 mmol). After 3 hr at -40° - -30° , the reaction was complete by tlc showed.

Saturated aq. K_2CO_3 was added and the suspension stirred 1 hr at 20° prior to diluting with H_2O and EtOAc. The combined organic layers from 3 EtOAc extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator to yield 1.2 g of a light yellow syrup. Chromatography on silica gel with 10% EtOAc/hexane eluted nonpolar impurities followed by the desired beta C-arylglucoside (0.54g).

10 F.



A solution of Part E tetra-O-benzyl C-glucoside (515 mg, 0.7 mmol) in EtOAc (10 mL) containing 10% $Pd(OH)_2/C$ (80 mg) was stirred overnight under 1 atmos. H_2 . After HPLC showed the reaction to be complete, the catalyst was filtered and the solvent removed using a rotary evaporator to obtain a white glassy solid that was further purified by preparative HPLC using a C_{18} reverse phase column to obtain 220 mg of the desired beta C-glucoside as a colorless syrup.

HPLC retention time: 6.43 min, 100% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 8 min gradient 0-100% B hold 5 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .

^1H NMR (500 MHz, CD_3OD) δ 7.20 (s, 1H), 7.18 (d, 1H, $J=7Hz$), 7.11 (d, 1H, $J=7Hz$), 6.89 (ABq, 4H), 4.07 (d, 1H,


J=9Hz), 3.90 (s, 2H), 3.87 (m, 1H), 3.70 (s, 3H), 3.68 (dd, 1H), 3.48-3.30 (m, 4H), 2.16 (s, 3H).

¹³C NMR (125 MHz, CD₃OD) δ 159.3, 140.3, 138.3, 137.4,
 5 133.7, 131.0, 130.8, 130.6, 126.9, 114.7, 83.5, 82.1,
 79.8, 76.3, 71.9, 63.1, 55.6, 59.6, 19.5.

Anal Calcd for C₂₁H₂₆O₆ LC-MS [M-H] 373; found 373.

10

Example 9



A. 5-Bromo-2-methyl-4'-hydroxydiphenylmethane


To a stirred -78° 10 mL CH₂Cl₂ solution of 5-bromo-2-methyl-4'-methoxydiphenylmethane (1.0g, 3.4 mmol) (See
 15 Example 8, Part C for preparation) was added 4.12 mL of a 1M BBr₃/ CH₂Cl₂. After 2 hr, the reaction was maintained at -40° for 20 hr whereupon HPLC indicated no starting ether remained. The reaction was quenched with aq. NaOH, extracted 3x with CH₂Cl₂, washed with brine prior to
 20 drying over Na₂SO₄. After removal of the volatiles, 0.84g of 5-bromo-2-methyl-4'-hydroxydiphenylmethane was obtained as a syrup which was used without further purification.

25 B. 5-Bromo-2-methyl-4'-benzyloxydiphenylmethane

A 10 mL DMF solution containing Part A 5-bromo-2-methyl-4'-hydroxydiphenylmethane (735 mg, 2.65 mmol), benzyl bromide (548 mg, 3.2 mmol), and K₂CO₃ (732 mg, 5.3 mmol) was stirred overnight. The reaction was then
 30 heated at 60° for 6 hr to drive the conversion from 80%

to 100%. After dilution with H₂O, the reaction was extracted 3x with EtOAc. The combined EtOAc layers were washed with H₂O and brine prior to drying over Na₂SO₄. The residue, after solvent removal under vacuum was chromatographed on silica gel using 3% EtOAc/hexane to elute 785 mg of 5-bromo-2-methyl-4'-benzyloxydiphenylmethane as a colorless syrup.

C.




10

To a stirred -78° solution of Part B 5-bromo-2-methyl-4'-benzyloxydiphenylmethane (0.43g, 1.2 mmol) in 7 mL of dry THF under Ar was added 0.68 mL of 1.9 M *n*-BuLi in hexane dropwise. After 30 min, 2,3,4,6-tetra-O-benzyl-β-D-glucolactone (0.7g, 1.3 mmol) in 3 mL of THF was added over 1 min. The solution was stirred for 0.75 hr at -78° prior to quenching with saturated aq. NH₄Cl. After warming to 20°, the reaction was diluted 2 fold with H₂O prior to 3 extractions with EtOAc. The combined EtOAc fractions were washed with brine and dried over Na₂SO₄. After concentration using a rotary evaporator, 0.96 g of the desired title lactol was obtained as a colorless syrup that was carried forward without further purification.

20


25

D.



- 5 To a stirred -30° solution of Part C lactol (0.96g, 1.16 mmol) in 10 mL of MeCN was added $i\text{Pr}_3\text{SiH}$ (0.37g, 2.3 mmol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.2g, 1.4 mmol). After 3 hr at -40° - -30° , saturated aq. K_2CO_3 was added and the suspension stirred 1 hr at 20° prior to diluting with H_2O
- 10 and EtOAc. The combined organic layers from 3 EtOAc extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator to yield 1.2 g of a light yellow syrup. Chromatography on silica gel with 9% EtOAc/hexane eluted nonpolar impurities; 10%
- 15 EtOAc/hexane eluted the desired beta C-arylglucoside (0.26g).

E.




20

- A solution of Part D penta-O-benzyl C-glucoside (255 mg, 0.31 mmol) in EtOAc (10 mL) containing 10% $\text{Pd}(\text{OH})_2/\text{C}$ (65 mg) was stirred 24 hr under 1 atmos. H_2 . After HPLC
- 25 showed the reaction to be complete, the catalyst was filtered and the solvent removed using a rotary

evaporator to obtain 115mg of a white glassy solid that was used without further purification.

F.



5 A threaded tube containing a magnetic stirrer, 4 mL of iPrOH and Part E phenolic C-glucoside (80 mg, 0.16 mmol) was cooled to -78° whereupon 1.5 g of CHClF_2 was added by condensing the gas. After adding 3 mL of 25%
10 aq. NaOH, the tube was sealed with a Teflon stopper and heated to 70° for 2 hr. By HPLC reaction contained a 2:3 mixture of starting phenol to desired ether. (Efforts to drive the conversion by prolonged reaction times were not successful.) After cooling, sufficient 1N HCl was added
15 to bring the pH to 2 whereupon most volatiles were removed using a rotary evaporator. The residue, after dissolution in 2:1 MeOH/ H_2O was purified by preparative HPLC equipped with a YMC S5 C_{18} reverse phase column (20x100 mm) employing a 10 min linear gradient with 45%-
20 90% aq MeOH at 20 mL/min to yield 40 mg of the desired phenolic ether.

HPLC retention time: 6.6 min, 95% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 8 min
25 gradient 0-100% B hold 5 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .

^1H NMR (400 MHz, CD_3OD) δ 7.22 (s, 1H), 7.20 (m, 1H), 7.12
30 (m, 1H), 7.06 (ABq, 4H), 6.73 (t, 1H, $J=27\text{Hz}$), 4.09 (d,


1H, J=9Hz), 3.98 (s, 2H), 3.89 (d, 1H), 3.68 (dd, 1H),
3.47-3.30 (m, 4H), 2.17 (s, 3H).

¹³C NMR (100 MHz, CD₃OD) δ 138.7, 138.2, 137.7, 136.6,
5 130.3, 130.2, 130.1, 126.4, 119.3, 117.0, 82.7, 81.4,
79.0, 75.6, 71.1, 62.3, 49.0, 38.8, 18.6.

Anal Calcd for C₂₁H₂₄F₂O₆ LC-MS [M+NH₄] 428; found 428.

10

Example 10



A. 5-Bromo-2-methyl-4'-thiomethylbenzophenone


AlCl₃ (535 mg, 4 mmol) was added to a 4° stirred 5 mL CS₂ solution of crude 5-bromo-2-methylbenzoyl chloride
15 (466mg, 2 mmol) (for preparation see Example 8, part B) and thioanisole (270mg, 2.3 mmol). The reaction, after warming to 20° over 1 hr, was stirred for 2 hr prior to quenching with 1N HCl. Subsequently, the suspension was diluted with 50 ml H₂O and stirred until all solids were
20 in solution. The mixture was extracted 3x with EtOAc. The combined organic extracts were washed 1x with 1N HCl, H₂O, aq NaHCO₃, and brine prior to drying over Na₂SO₄. After removal of the volatiles, the residue was chromatographed on silica gel using 15% EtOAc/hexane to
25 elute 450mg of 5-bromo-2-methyl-4'-thiomethylbenzophenone as a white solid.

B. 5-Bromo-2-methyl-4'-thiomethyldiphenylmethane

A solution of Et₃SiH (0.45 mL, 2.85 mmol), BF₃·Et₂O
30 (0.3 mL, 2.4 mmol), and Part A 5-bromo-2-methyl-4'-


thiomethylbenzophenone (450mg, 1.4 mmol) in 3 mL of a 1:9 mixture $\text{CH}_2\text{Cl}_2/\text{MeCN}$ was stirred overnight at 20° . After quenching with 10% NaOH and dilution with H_2O , the reaction was extracted 3x with EtOAc. The combined organic layers were washed 2x with H_2O and once with brine before drying over Na_2SO_4 . After removal of the volatiles, the residue was chromatographed on silica gel using 5% EtOAc/hexane to elute 416mg of 5-bromo-2-methyl-4'-thiomethyldiphenylmethane as a colorless oil.

C.




To a stirred -78° solution of Part B 5-bromo-2-methyl-4'-thiomethyldiphenylmethane (200mg, 0.65 mmol) in 10 mL of dry THF under Ar was added dropwise 0.42 mL of 1.8 M *n*-BuLi in hexane. After 2 hr, this solution was transferred by cannula to a stirred -78° solution of 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (0.88 g, 1.6 mmol) in 5 mL of THF. The solution was stirred for 2 hr at -78° before quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 2 fold with H_2O prior to 3 extractions with EtOAc. The combined EtOAc fractions were washed with brine and dried over Na_2SO_4 . After concentration using a rotary evaporator, 550mg of the desired title lactol was obtained as a colorless syrup that was carried forward without further purification.

D.



- 5 To a stirred -40° solution of Part C lactol (550mg, 0.72 mmol) in 6 mL of MeCN was added $i\text{Pr}_3\text{SiH}$ (0.22 mL, 1.0 mmol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.11 mL, 0.8 mmol). After 1.5 hr at -40° - -30° , when tlc showed the reaction to be complete, saturated aq. K_2CO_3 was added and the suspension
- 10 stirred 1 hr at 20° prior to diluting with H_2O and EtOAc. The combined organic layers from 3 EtOAc extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator. Chromatography of the residue on silica gel using 9% EtOAc/hexane as eluant eluted
- 15 240mg of the desired beta C-arylglucoside.

E.



20

- A solution of Part D tetra-O-benzyl C-glucoside (70mg, 0.1 mmol) in EtSH (1.5 mL) containing $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.24 mL, 2 mmol) was stirred at 20° for 2 hr. After 1 more hr following addition of an additional 0.12 mL of
- 25 $\text{BF}_3 \cdot \text{Et}_2\text{O}$, the reaction was complete. The reaction was quenched by slow addition of 0.4 mL of pyridine prior to dilution with aq. NH_4Cl . The combined organic layers from

3 EtOAc extractions were washed with brine, dried over Na₂SO₄, and concentrated using a rotary evaporator. The residue was purified by preparative HPLC using a C₁₈ reverse phase column to obtain 20mg of the desired beta C-glucoside as a white lyophilate after lyophilization.


HPLC retention time: 3.8 min, 95% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 4 min gradient 0-100% B hold 4 min at 100% B. Solvent A: 10% MeOH/H₂O + 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.

¹H NMR (500 MHz, CD₃OD) δ 7.21-7.11 (m, 5H), 7.05 (d, 2H, J=8.0 Hz), 4.08 (d, 1H, J=9.1 Hz), 3.98 (s, 2H), 3.87 (d, 1H, J=12.6 Hz), 3.68 (dd, 1H, J=5.2, 12.1 Hz), 3.49-3.30 (m, 4H), 2.41 (s, 3H).

¹³C NMR (125 MHz, CD₃OD) δ 139.8, 138.9, 138.4, 137.5, 137.1, 131.1, 130.9, 129.1, 130.3, 127.8, 127.1, 83.6, 82.2, 79.8, 76.4, 72.0, 63.2, 39.9, 19.5, 16.1.

Anal Calcd for C₂₁H₂₆O₅S LC-MS [M+NH₄] 408; found 408.

Example 11



A. 5-Bromo-2-chloro-4'-thiomethylbenzophenone


To a stirred suspension of commercial 5-bromo-2-chlorobenzoic acid (506mg, 2.12 mmol) in 10 mL of CH₂Cl₂ containing oxalyl chloride (2.4 mmol) was added 2 drops of DMF. Once the vigorous evolution of gas ceased, the

reaction was stirred 1.5 hr before removal of the volatiles using a rotary evaporator. After dissolving the crude 5-bromo-2-chlorobenzoyl chloride in 8 ml of CS₂, the stirred mixture was cooled to 4° prior to adding
5 thioanisole (260mg, 2.12 mmol) followed by AlCl₃ (566mg, 4.25 mmol). The reaction, after warming to 20° over 1 hr, was stirred for 20 hr prior to quenching with 1N HCl. Subsequently, the suspension was diluted with 50 ml H₂O and stirred until all solids were in solution. The
10 mixture was extracted 3x with EtOAc. The combined organic extracts were washed 1x with 1N HCl, H₂O, aq NaHCO₃, and brine prior to drying over Na₂SO₄. After removal of the volatiles, the 710mg of crude of 5-bromo-2-chloro-4'-thiomethylbenzophenone was not further
15 purified.

B. 5-Bromo-2-chloro-4'-thiomethyldiphenylmethane

A solution of Et₃SiH (1.4 mL, 8.8 mmol), BF₃·Et₂O (0.83 mL, 6.6 mmol), and Part A 5-bromo-2-chloro-4'-
20 thiomethylbenzophenone (710mg, 2.1 mmol) in 10 mL of a 1:4 mixture CH₂Cl₂/MeCN was stirred 2 hr at 20°. After quenching with 10% NaHCO₃ and dilution with H₂O, the reaction was extracted 3x with EtOAc. The combined organic layers were washed 2x with H₂O and once with
25 brine before drying over Na₂SO₄. After removal of the volatiles, the residue was chromatographed on silica gel using 5% EtOAc/hexane to elute 630mg of 5-bromo-2-chloro-4'-thiomethyldiphenylmethane as a colorless oil.

C.




5 To a stirred -78° solution of Part B 5-bromo-2-chloro-4'-thiomethyldiphenylmethane (200mg, 0.61 mmol) in 6 mL of dry THF under Ar was added 0.48 mL of 1.5 M *n*-BuLi in hexane dropwise. After 35 minutes, this solution was transferred by cannula to a stirred -78° solution of

10 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (361mg, 0.67 mmol) in 5 mL of THF. The solution was stirred for 1.5 hr at -78° prior to quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 2 fold with H_2O prior to 3 extractions with EtOAc. The combined

15 EtOAc fractions were washed with brine and dried over Na_2SO_4 . After concentration using a rotary evaporator, the residue was chromatographed on silica gel using 20% EtOAc/hexane to elute 250mg of the desired title lactol.


20 D.



To a stirred -30° solution of Part C lactol (250mg, 0.32 mmol) in 5 mL of MeCN was added $i\text{Pr}_3\text{SiH}$ (0.10 mL, 0.56 mmol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.048 mL, 0.38 mmol). After 0.5 hr at -30° , when tlc showed the reaction to be

complete, saturated aq. NaHCO_3 was added and the suspension stirred 1 hr at 20° prior to diluting with H_2O and EtOAc. The combined organic layers from 3 EtOAc extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator. Chromatography of the residue on silica gel using 9% EtOAc/hexane as eluant eluted 200mg of the desired beta C-arylglucoside.

E.



10

A solution of Part D tetra-O-benzyl C-glucoside (60mg, 0.1 mmol) in EtSH (2 mL) containing $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.24 mL, 2 mmol) was stirred at 20° for 3 hr. The reaction was quenched by slow addition of 0.4 mL of pyridine prior to dilution with aq. NH_4Cl . The combined organic layers from 3 EtOAc extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator. The residue was purified by preparative HPLC using a C_{18} reverse phase column to obtain 21.5mg of the desired beta C-glucoside as a white lyophilate after lyophilization.

HPLC retention time: 3.96 min, 95% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 4 min gradient 0-100% B hold 4 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .

^1H NMR (400 MHz, CD_3OD) δ 7.36-7.27 (m, 3H), 7.15 (d, 2H, $J=8.3$ Hz), 7.11 (d, 2H, $J=8.3$ Hz), 4.10-4.04 (m, 3H),

30


3.87 (d, 1H, J=12 Hz), 3.70 (dd, 1H, J=7.1, 11.8 Hz),
3.47-3.26 (m, 4H), 2.42 (s, 3H).

¹³C NMR (100 MHz, CD₃OD) δ 140.1, 139.3, 138.0, 137.5,
5 134.5, 132.0, 130.4, 130.2, 128.4, 128.0, 82.9, 82.8,
82.2, 79.7, 76.5, 71.8, 63.1, 39.5, 16.1.

Anal Calcd for C₂₀H₂₃ClO₅S LC-MS [M-H] 409; found 409.

10

Example 12



A. 5-Bromo-2-chloro-4'-methoxybenzophenone


To a stirred suspension of commercial 5-bromo-2-chlorobenzoic acid (506mg, 2.12 mmol) in 10 mL of CH₂Cl₂
15 containing oxalyl chloride (2.4 mmol) was added 2 drops of DMF. Once the vigorous evolution of gas ceased, the reaction was stirred 1.5 hr prior to removal of the volatiles using a rotary evaporator. After dissolving the crude 5-bromo-2-chlorobenzoyl chloride in 8 ml of
20 CS₂, the stirred mixture was cooled to 4° prior to adding anisole (240mg, 2.12 mmol) followed by AlCl₃ (566mg, 4.25 mmol). The reaction, after warming to 20° over 1 hr, was stirred for 20 hr prior to quenching with 1N HCl. Subsequently, the suspension was diluted with 50 ml H₂O
25 and stirred until all solids were in solution. The mixture was extracted 3x with EtOAc. The combined organic extracts were washed 1x with 1N HCl, H₂O, aq NaHCO₃, and brine prior to drying over Na₂SO₄. After removal of the volatiles, the residue was chromatographed

on silica gel using 15% EtOAc/hexane to elute 450mg of 5-bromo-2-chloro-4'-methoxybenzophenone.

B. 5-Bromo-2-chloro-4'-methoxydiphenylmethane

5 A solution of Et_3SiH (0.45 mL, 2.85 mmol), $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.3 mL, 2.4 mmol), and 5-bromo-2-chloro-4'-methoxybenzophenone (450mg, 1.4 mmol) in 3 mL of a 1:9 mixture $\text{CH}_2\text{Cl}_2/\text{MeCN}$ was stirred overnight at 20° . After quenching with 10% NaOH and dilution with H_2O , the
10 reaction was extracted 3x with EtOAc. The combined organic layers were washed 2x with H_2O and once with brine before drying over Na_2SO_4 . After removal of the volatiles, the residue was chromatographed on silica gel using 2% EtOAc/hexane to elute 416mg of 5-bromo-2-chloro-
15 4'-methoxydiphenylmethane as a colorless oil.

C.




20

To a stirred -78° solution of Part B 5-bromo-2-chloro-4'-methoxydiphenylmethane (212mg, 0.68 mmol) in 8 mL of dry THF under Ar was added 0.36 mL of 1.9 M *n*-BuLi in hexane dropwise. After 30 minutes, this solution was
25 transferred by cannula to a stirred -78° solution of 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (0.39g, 0.71 mmol) in 5 mL of THF. The solution was stirred for 2 hr at -78° prior to quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 2 fold
30 with H_2O prior to 3 extractions with EtOAc. The combined

EtOAc fractions were washed with brine and dried over Na_2SO_4 . After concentration using a rotary evaporator, the residue was chromatographed on silica gel using 20% EtOAc/hexane to elute 142mg of the desired title lactol.


5

D.



10 To a stirred -40° solution of Part C lactol (142mg, 0.18 mmol) in 1.5 mL of MeCN was added $i\text{Pr}_3\text{SiH}$ (0.041 mL, 0.2 mmol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.026 mL, 0.2 mmol). After 2 hr at -40° , when tlc showed the reaction to be complete, saturated aq. NaHCO_3 was added and the diluted
 15 with H_2O and CH_2Cl_2 . The combined organic layers from 3 CH_2Cl_2 extractions were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator. Chromatography of the residue on silica gel using 25% EtOAc/hexane as eluant eluted 139mg of the desired beta
 20 C-arylglucoside.

E.



25

A solution of Part D tetra-O-benzyl C-glucoside (136 mg, 0.18 mmol) in EtSH (1.0 mL) containing $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.46

mL, 3.6 mmol) was stirred at 20° for 4 hr. The reaction was diluted with CH₂Cl₂ and then concentrated using a rotary evaporator. The residue, after being dissolved in CH₂Cl₂, was washed with aq. NH₄Cl, H₂O, brine, dried over Na₂SO₄, and concentrated using a rotary evaporator. The crude product was purified by preparative HPLC using a C₁₈ reverse phase column to obtain 26mg of the desired beta C-glucoside as a white lyophilate after lyophilization.

10 HPLC retention time: 3.07 min, 95% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 4 min gradient 0-100% B hold 4 min at 100% B. Solvent A: 10% MeOH/H₂O + 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.

15


¹H NMR (500 MHz, CD₃OD) δ 7.35-7.28 (m, 3H), 7.1 (d, 2H, J=8.8 Hz), 6.8 (d, 2H, J=8.3 Hz), 4.05-3.90 (m, 3H), 3.80 (d, 1H, J=12.3 Hz), 3.67 (s, 3H), 3.61 (dd, 1H, J=4.8, 11.9 Hz), 3.42-3.25 (m, 4H) Hz).

20

¹³C NMR (125 MHz, CD₃OD) δ 159.6, 140.0, 139.9, 134.5, 133.0, 131.9, 130.8, 130.1, 114.8, 82.9, 82.2, 79.8, 76.5, 71.9, 63.1, 55.6, 39.2.

25 Anal Calcd for C₂₀H₂₃ClO₆ LC-MS [M+NH₄] 412; found 412.

Example 13



30


A. 5-Bromo-2-methoxy-4'-ethylbenzhydrol

To a stirred -78° solution of *p*-bromoethylbenzene (2.03g, 11 mmol) in 10 mL of dry THF under Ar was added 5 mL of 2.5 M *n*-BuLi (12 mmol) in hexane over 10 min. The
5 temperature was allowed to rise to -10° over 2 hr whereupon the reaction was cooled to -78° before adding solid 5-bromo-2-methoxybenzaldehyde (2.15 g, 10 mmol). After stirring overnight at 20° , the reaction was quenched with saturated aq. NH_4Cl and diluted 5 fold with
10 H_2O prior to 3 extractions with EtOAc. The combined EtOAc fractions were washed with brine and dried over Na_2SO_4 . After concentration using a rotary evaporator, the residue was chromatographed on silica gel using 10% EtOAc/hexane to elute 1.44g of 5-bromo-2-methoxy-4'-
15 ethylbenzhydrol.

B. 5-Bromo-2-methoxy-4'-ethyldiphenylmethane


A 9 mL solution of 1:8 $\text{CH}_2\text{Cl}_2/\text{MeCN}$ containing crude Part A 5-bromo-2-methoxy-4'-ethylbenzhydrol (1.44g, 4.5
20 mmol), Et_3SiH (0.75 mL, 5 mmol, and $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.6 mL, 6.4 mmol) was stirred overnight at 20° . After quenching with saturated aq. NaOH , the mixture was extracted 3x with EtOAc. The combined EtOAc fractions were washed with brine and dried over Na_2SO_4 . After concentration
25 using a rotary evaporator, the residue was chromatographed on silica gel using 2% EtOAc/hexane to elute 1.28g of 5-bromo-2-methoxy-4'-ethyldiphenylmethane.

C.



- 5 To a stirred -78° solution of Part B 5-bromo-2-methoxy-4'-ethyldiphenylmethane (0.25g, 0.82 mmol) in 7 mL of dry THF under Ar was added dropwise 0.5 mL of 1.8 M *n*-BuLi in hexane. After 2 hr, 2,3,4,6-tetra-O-benzyl- β -D-glucolactone (0.48g, 0.9 mmol) in 3 mL of THF was added
- 10 over 1 min. The solution was stirred for 2 hr at -78° prior to quenching with saturated aq. NH_4Cl . After warming to 20° , the reaction was diluted 5 fold with H_2O prior to 3 extractions with EtOAc. The combined EtOAc fractions were washed with brine and dried over Na_2SO_4 .
- 15 After concentration using a rotary evaporator, 0.67g of the desired title lactol was obtained as a light yellow syrup that was carried forward without further purification.


20 D.



- To a stirred -30° solution of Part C lactol (450mg, 0.59 mmol) in 10 mL of MeCN was added $i\text{Pr}_3\text{SiH}$ (0.2 mL, 0.9 mmol) followed by $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (0.1 mL, 0.7 mmol). After 1.5 hr at -40° , the reaction being complete by tlc was
- 25

quenched by addition of aq. NaHCO_3 and subsequently extracted 3x with EtOAc. The combined organic layers were washed with brine, dried over Na_2SO_4 , and concentrated using a rotary evaporator. Chromatography of the residue on silica gel with 10% EtOAc/hexane eluted 320mg of the desired beta C-arylglucoside.

E.



10


A solution of Part D tetra-O-benzyl C-glucoside (320 mg, 0.7 mmol) in EtOAc (15 mL) containing 10% $\text{Pd}(\text{OH})_2/\text{C}$ (30 mg) was stirred overnight under 1 atm. H_2 . After HPLC showed the reaction to be complete, the catalyst was filtered and the solvent removed using a rotary evaporator. The crude product was further purified by preparative HPLC using a C_{18} reverse phase column to obtain 24 mg of the desired beta C-glucoside as a white solid after lyophilization.

HPLC retention time: 3.84 min, 95% pure, YMC S5 C-18 4.6x50mm column, 2.5 mL/min, detection at 220nm; 4 min gradient 0-100% B hold 4 min at 100% B. Solvent A: 10% MeOH/ H_2O + 0.2 % H_3PO_4 . Solvent B: 90% MeOH/ H_2O + 0.2 % H_3PO_4 .

^1H NMR (500 MHz, CD_3OD) δ 7.23 (d, 1H, $J=7$ Hz), 7.17 (s, 1H), 7.05 (ABq, 4H), 6.89 (d, 1H, $J=7$ Hz), 4.02 (d, 1H $J=9\text{Hz}$), 3.92-3.83 (m, 3H), 3.76 (s, 3H), 3.66 (dd, 1H), 3.45-3.29 (m, 4H), 2.55 (q, 2H), 1.16 (t, 3H).


Anal Calcd for $C_{22}H_{28}O_6$ LC-MS $[M+NH_4]$ 406; found 406.

5

Example 1410 A. N-Ethyl-N-4-methoxybenzyl-2,6-dihydroxybenzamide


To a stirred solution of N-ethyl-4-methoxybenzyl
amine (1.07g, 6.49 mmol) in DMF (10 mL) was added 2,6-
dihydroxybenzoic acid (1.0g, 6.49 mmol) followed by HOAt
(0.97g, 7.14 mmol) and EDC (1.31g, 6.81 mmol). After
15 stirring overnight, the reaction was diluted with EtOAc
prior to washing 3x with H_2O . The combined aqueous
layers were extracted once with EtOAc. The organic
fractions were combined, washed once with brine, and
dried over Na_2SO_4 prior to concentrating using a rotary
20 evaporator. The residue was chromatographed on silica
gel using 75% EtOAc/hexane as the eluent. The resulting
promising impure fractions were further purified by
silica gel chromatography. A total of 631mg of the
desired N-ethyl-N-4-methoxybenzyl 2,6-dihydroxybenzamide
25 was obtained.

B.



5 A stirred suspension of the Part A amide (630mg, 2.09 mmol), CdCO_3 (939mg, 5.44 mmol) in toluene (30 mL) was refluxed for 1.5 hr using a Dean Stark trap prior to the addition of 2,3,4,6-tetra-O-acetyl- α -D-glucosopyranosyl bromide (1.12g, (2.72 mmol). After 15 hr
10 of reflux, no starting amide remained by tlc analysis. The hot suspension was filtered through celite which was washed with hot PhMe and then 3x with hot CHCl_3 . After removal of the volatiles using a rotary evaporator, the residue was chromatographed on silica gel. A mixture of
15 O-glucosides was eluted with 1:1 EtOAc/hexane prior to the tetraacetate of the desired title C-glucoside; 172mg of severely contaminated title C-glucoside was obtained.

C.



20

Impure Part B ester was stirred in 6:1 EtOH/ H_2O (1.4 mL) containing KOH (140mg, 2.5 mmol) for 16 hr. The

resulting solution was cooled to 4°, acidified to pH 5, and then extracted 2x with EtOAc. The combined EtOAc layers were washed with brine, and dried over Na₂SO₄ prior to concentrating using a rotary evaporator. The residue
5 was purified by prep HPLC with a C₁₈ YMC reverse phase column using a 45-90% MeOH/H₂O gradient over 30 min to elute the desired title C-glucoside (7.8 mg).

HPLC: 99.1%; Shimadzu LC-6A, YMC S3 ODS (6.0 X 150 mm);
10 flow rate of 1.5 mL/min; detection at 220nm; gradient elution 0-100% B over 30 minutes (A = 90% H₂O, 10% MeOH, 0.2% H₃PO₄, and B = 90% MeOH, 10% H₂O, 0.2% H₃PO₄); retention time = 23.4 minutes.


15 ¹H NMR (400 MHz, CD₃OD): δ 1.22 (3H, t, J = 7.2 Hz), 3.4-3.5 (6H, m), 3.73 (3H, s), 3.74 (1H, m), 3.77 (1H, m), 3.8-3.9 (2H, m), 4.36 (1H, d, J = 9.3 Hz), 6.77 (2H, d, J = 8.6 Hz), 7.11 (2H, d, J = 8.6 Hz), 7.18 (1H, s)

20 ¹³C NMR (125 MHz, CD₃OD): δ 14.9, 35.1, 35.1, 55.7, 62.5, 71.2, 75.8, 79.6, 80.3, 82.3, 104.8, 114.7, 117.1, 122.7, 130.7, 134.5, 134.6, 151, 159.3, 161, 171.9


Anal Calcd for C₂₃H₂₉NO₉ LC-MS [M-H] 462; found 462.

25

Example 15




A.



5 A mixture of Example 3 Part B β -*m*-bromophenyl-C-glucoside (100mg, 0.14 mmol), *p*-methylphenylboronic acid (59mg, 0.43 mmol), Na_2CO_3 (46mg, 0.43 mmol), and $\text{Pd}(\text{PPh}_3)_4$ (153mg, 0.13 mmol) in 3:1 PhMe/EtOH were stirred under Ar at 80° for 15 hr. After removal of the volatiles using a rotary evaporator, the residue was chromatographed on silica gel. 10:1 hexane/EtOAc eluted the desired title biphenyl C-glucoside (90mg) as a clear oil.

B.




15 To a -78° stirred CH_2Cl_2 solution (0.4 mL) of Part A tetra-O-benzyl ether (65mg, 0.09 mmol) under Ar was added 0.37 mL of a 1M BCl_3 in CH_2Cl_2 . After 1 hr, the reaction was quenched with 2 mL of MeOH and allowed to warm to 20°. After adjusting the pH to ~7 with aqueous NaHCO_3 , the suspension was extracted 2x with CH_2Cl_2 . The combined organic layers were dried over MgSO_4 and concentrated. 25 The resulting residue, after purification by preparative HPLC using a C_{18} reverse phase column, yielded 6.6mg of final title product. (Note the product is partially destroyed by the strongly acidic medium generated after the MeOH quench of the BCl_3 .)

HPLC retention time: 6.353 min, 100% pure, Zorbax C-18
4.6x50mm, 2.5 mL/min, detection at 220nm; 8 min gradient
0-100% B hold 5 min at 100% B. Solvent A: 10% MeOH/H₂O +
5 0.2 % H₃PO₄. Solvent B: 90% MeOH/H₂O + 0.2 % H₃PO₄.

¹H NMR (400 MHz, CD₃OD) δ 7.65 (s, 1H), 7.53-7.50 (m,
3H), 7.39-3.37 (m, 2H), 7.23 (d, 2H, J= 7.9 Hz), 4.20 (d,
1H, J= 9.3 Hz), 3.89 (dd, 1H, J= 2.2, 11.9 Hz), 3.71 (dd,
10 1H, J= 5.7, 11.9 Hz), 3.50-3.40 (m, 4H), 2.36 (s, 3H)
Anal Calcd for C₁₉H₂₂O₅ Low Res MS [M-H] 329; found 329

Examples 16 to 80


The compounds of Examples 16 to 80 set out in the
15 following Tables 1 and 2 were prepared employing
procedures of Examples 1 to 15 and reaction Schemes 1 to
9 above. It will be appreciated that compounds wherein A,
which may linked at the ortho, meta, or para position of
the aryl ring attached to the glucoside, may be any one
20 of (CH₂)_n, O, NH or S while R¹, R², R^{2a}, R³ and R⁴ may be
any of the substituents as defined above, may be prepared
employing the procedures of Examples 1 to 15 and reaction
Schemes 1 to 9.

Table 1

Example	A	R ³	Method of Example #	LC/MS or MS (M + H) ⁺
16	CH ₂	4-Me	1	345
17	CH ₂	4-OH	1	347
18	CH ₂	3-Me	2	345
19	CH ₂	H	3	331
20	CH ₂	3-OMe	3	361
21	CH ₂	4-CO ₂ Me	3	389
22	CH ₂	3, 4- (OCH ₂ O)	3	375
23	CH ₂	4-CF ₃	3	399
24	CH ₂	4-NHAc	3	388
25	CH ₂	4-SO ₂ Me	3	409
26	CH ₂	4-Ph	3	407
27	CH ₂	4-NHSO ₂ Ph-4'-Me	3	500
28	CH ₂	4-NHSO ₂ Me	3	424
29	CH ₂	4-CO ₂ H	3	375
30	CH ₂	4-Thiadiazole	3	415
31	CH ₂	4-Tetrazole	3	399
32	CH ₂	4-OCH ₂ Ph-4'-CN	1	462
33	CH ₂	4-OCHF ₂	1	397
34	CH ₂	4- <i>i</i> Pr	3	373

35	CH ₂	2- <i>i</i> Pr	3	373
36	CH ₂	4-O- <i>n</i> Pr	1	389
37	CH ₂	4-Tetrazole-2'-Me	3	413
38	CH ₂	4-Tetrazole-1'-Me	3	413
39	CH ₂	4-OPh	1	423
40	CH ₂	4- <i>n</i> Pr	1	373
41	CH ₂	4- <i>n</i> Bu	1	387
42	CH ₂	4-SO ₂ Et	1	423
43	CH ₂	4-SO ₂ - <i>n</i> Pr	1	437
44	CH ₂	4-SO ₂ Ph	3	471
45	CH ₂	4-SOMe	4	393
46	Bond	H	15	317
47	Bond	3-Me	15	331
48	Bond	4-MeO	15	347
49	(CH ₂) ₂	H	1	343 (M-H)
50	(CH ₂) ₂	4-Me	1	357 (M-H)
51	(CH ₂) ₃	H	1	376 (M+NH ₄)
52	(CH ₂) ₃	4-Me	1	390 (M+NH ₄)
53	(CH ₂) ₃	3-Me	1	390 (M+NH ₄)
54	Bond (para link)	H	15	317
55	CH ₂ (ortho link)	H	1	331
56	CH ₂ (ortho link)	4-Et	1	376 (M+NH ₄)

57	O	4-Me	Scheme 8	364 (M+NH ₄)
58	S	4-Me	Scheme 9	380 (M+NH ₄)


Table 2

Example	A	R ¹	R ²	R ³	Method of Example #	LC/MS or MS (M + H) ⁺
59	CH ₂	2-Me	H	4-Et	1	371 (M-H)
60	CH ₂	4-Me	H	4-Et	8	371 (M-H)
61	CH ₂	4-Me	H	4-SO ₂ Me	8	445 (M+Na)
62	CH ₂	4-Me	H	4-OH	9	359 (M-H)
63	CH ₂	4-Me	H	4-S(O)Me	10	407 (M+H)
64	CH ₂	4-Me	H	4-F	8	385 (M+NH ₄)
65	CH ₂	4-Me	H	4-Cl	8	377 (M-H)
66	CH ₂	4-Me	H	4-Me	8	357 (M-H)
67	CH ₂	4-Me	H	H	8	343 (M-H)
68	CH ₂	4-Me	6-Me	4-OMe	1	406 (M+NH ₄)
69	CH ₂	4-F	H	4-OMe	1	396 (M+NH ₄)
70	CH ₂	4-Cl	H	4-SOMe	11	427 (M+H)
71	CH ₂	4-Cl	H	4-SO ₂ Me	11	441 (M-H)
72	CH ₂	4-Cl	H	4-OCHF ₂	9	448 (M+NH ₄)

73	CH ₂	4-Et	H	4-OMe	8	406 (M+NH ₄)
74	CH ₂	4-iPr	H	4-OMe	8	420 (M+NH ₄)
75	CH ₂	4-iPr	H	4-SMe	10	417 (M-H)
76	CH ₂	4-iPr	H	4-SO ₂ Me	10	439 (M-H)
77	CH ₂	4,5-OCH ₂ O	H	4-Et	1	403 (M +H)
78	CH ₂	5-Me	H	4-Et	1	390 (M+NH ₄)
79	CH ₂	5-Me	6-Me	4-OMe	1	406 (M+NH ₄)
80	CH ₂	6-Me	H	4-Et	8	395 (M+Na)

We claim:

1. A compound having the structure



wherein

R^1 , R^2 and R^{2a} are independently hydrogen, OH, OR^5 , (C_1-C_{20}) alkyl, CF_3 , $OCHF_2$, OCF_3 , SR^{5i} or halogen, or two of R^1 , R^2 and R^{2a} together with the carbons to which they are attached can form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 ;

R^3 and R^4 are independently hydrogen, OH, OR^{5a} , $O-(C_6-C_{10})$ aryl, $OCH_2-(C_6-C_{10})$ aryl, (C_1-C_{20}) alkyl, (C_3-C_{20}) cycloalkyl, CF_3 , $-OCHF_2$, $-OCF_3$, halogen, $-CN$, $-CO_2R^{5b}$, $-CO_2H$, COR^{6b} , $-CH(OH)R^{6c}$, $-CH(OR^{5h})R^{6d}$, $-CONR^6R^{6a}$, $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2-(C_6-C_{10})$ aryl, (C_6-C_{10}) aryl, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$, or $-SO_2-(C_6-C_{10})$ aryl, or a five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 , or R^3 and R^4 together with the carbons to which they are attached form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 ;

R^5 , R^{5a} , R^{5b} , R^{5c} , R^{5d} , R^{5e} , R^{5f} , R^{5g} , R^{5h} and R^{5i} are independently (C_1-C_{20}) alkyl;


R^6 , R^{6a} , R^{6b} , R^{6c} , and R^{6d} are independently hydrogen, (C_1-C_{20}) alkyl, (C_6-C_{10}) aryl, (C_1-C_{20}) alkyl (C_6-C_{10}) aryl or

(C₃-C₂₀)cycloalkyl, or R⁶ and R^{6a} together with the nitrogen to which they are attached form an annelated five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO₂; A is O, S, NH, or (CH₂)_n where n is 0 - 3, or a pharmaceutically acceptable salt thereof;

with the proviso that where A is (CH₂)_n where n is 0, 1, 2, or 3 or A is O, and at least one of R¹, R², and R^{2a} is OH or OR⁵, then at least one of R¹, R², and R^{2a} is CF₃, OCF₃, or OCHF₂ or at least one of R³ and R⁴ is CF₃, -OCHF₂, -OCF₃, -CN, -CO₂R^{5b}, CH(OR^{5h})R^{6d}, CH(OH)R^{6c}, COR^{6b}, -NHCOR^{5c}, -NHSO₂R^{5d}, -NHSO₂-(C₆-C₁₀)aryl, (C₆-C₁₀)aryl, -SR^{5e}, -SOR^{5f}, -SO₂R^{5g} or -SO₂-(C₆-C₁₀)aryl, or at least one of R¹, R², and R^{2a} is CF₃, OCF₃, or OCHF₂ and at least one of R³ and R⁴ is CF₃, -OCHF₂, -OCF₃, -CN, -CO₂R^{5b}, CH(OR^{5h})R^{6d}, CH(OH)R^{6c}, COR^{6b}, -NHCOR^{5c}, -NHSO₂R^{5d}, -NHSO₂-(C₆-C₁₀)aryl, (C₆-C₁₀)aryl, -SR^{5e}, -SOR^{5f}, -SO₂R^{5g} or -SO₂-(C₆-C₁₀)aryl.

2. The compound as defined in Claim 1 with the proviso that where A is (CH₂)_n where n is 0, 1, 2, or 3 or A is O, and at least one of R¹, R², R^{2a}, R³ and R⁴ is OH or OR⁵, then at least one of R¹, R², and R^{2a} is CF₃, OCF₃, or OCHF₂ or at least one of R³ and R⁴ is CF₃, -OCHF₂, -OCF₃, -CN, -CO₂R^{5b}, CH(OR^{5h})R^{6d}, -NHCOR^{5c}, -NHSO₂R^{5d}, -NHSO₂-(C₆-C₁₀)aryl, (C₆-C₁₀)aryl, -SR^{5e}, -SOR^{5f}, -SO₂R^{5g}, -SO₂-(C₆-C₁₀)aryl or halogen, or at least one of R¹, R², and R^{2a} is CF₃, OCF₃, or OCHF₂ and at least one of R³ and R⁴ is CF₃, -OCHF₂, -OCF₃, -CN, -CO₂R^{5b}, CH(OR^{5h})R^{6d}, -NHCOR^{5c}, -NHSO₂R^{5d}, -NHSO₂-(C₆-C₁₀), (C₆-C₁₀)aryl, -SR^{5e}, -SOR^{5f}, -SO₂R^{5g}, or -SO₂-(C₆-C₁₀)aryl.

3. The compound as defined in Claim 1 having the structure



4. The compound as defined in Claim 1 wherein A is $(\text{CH}_2)_n$.

5. The compound as defined in Claim 3 wherein A is CH_2 or O or S.

6. The compound as defined in Claim 1 wherein A is CH_2 , O or S;

R^1 , R^2 and R^{2a} are independently selected from H, $(\text{C}_1\text{-C}_8)$ alkyl, halogen, OR^5 , or OCHF_2 , or two of R^1 , R^2 and R^{2a} are H and the other is $(\text{C}_1\text{-C}_8)$ alkyl, halogen, OR^5 , or OCHF_2 ;

R^3 and R^4 are independently selected from $(\text{C}_1\text{-C}_8)$ alkyl, OR^{5a} , $-\text{OCHF}_2$, $-\text{SR}^{5e}$, OH, CO_2R^{5b} , $-\text{COR}^{6b}$, $-\text{CH}(\text{OH})\text{R}^{6c}$,


$-\text{CH}(\text{OR}^{5h})\text{R}^{6d}$, CF_3 , $\text{R}^{5c}-\overset{\text{O}}{\parallel}\text{C}-\text{NH}-$, $-\text{SOR}^{5f}$, $-\text{SO}_2\text{R}^{5g}$, $(\text{C}_6\text{-C}_{10})$ aryl, $-\text{NHSO}_2-(\text{C}_6\text{-C}_{10})$ aryl, $-\text{NHSO}_2\text{R}^{5d}$, CO_2H , thiadiazole, tetrazole, $\text{OCH}_2-(\text{C}_6\text{-C}_{10})$ aryl, $-\text{OCF}_3$, $\text{O}-(\text{C}_6\text{-C}_{10})$ aryl, and H, or together form $-3,4-(\text{O}-\text{CH}_2-\text{O})-$.


7. The compound as defined in Claim 6 wherein A is CH_2 ; R^1 is hydrogen, halogen or $(\text{C}_1\text{-C}_8)$ alkyl; R^2 and R^{2a} are each H; R^3 is H; R^4 is $(\text{C}_1\text{-C}_8)$ alkyl, $-\text{COR}^{6b}$, $-\text{CH}(\text{OH})\text{R}^{6c}$, $-\text{CH}(\text{OR}^{5h})\text{R}^{6d}$, OR^{5a} , $-\text{OCHF}_2$, $-\text{OCF}_3$ or $-\text{SR}^{5e}$.

8. The compound as defined in Claim 7 where A is CH₂; R¹ is hydrogen, halogen or (C₁-C₈) alkyl; and R⁴ is (C₁-C₈) alkyl, R^{5a}O, -OCHF₂, or -SR^{5e}.


9. The compound as defined in Claim 7 wherein R⁴ is 4-C₂H₅.

10. The compound as defined in Claim 3 having the structure






11. The compound as defined in Claim 1 having the structure set out as follows:



where A is CH₂ and meta to the glucoside, R¹, R² and R^{2a} are each H, and R³ is as follows:

4-Me, 4-OH, 3-Me, H, 3-OMe, 4-CO₂Me, 3,4-(OCH₂O), 4-CF₃, 4-NHAc, 4-SO₂Me, 4-Ph, 4-NHSO₂Ph-4'-Me, 4-NHSO₂Me, 4-CO₂H, 4-thiadiazole, 4-tetrazole, 4-OCH₂Ph-4'-CN, 4-OCHF₂, 4-isopropyl, 2-isopropyl, 4-O-n-propyl, 4-tetrazole-2'-Me, 4-tetrazole-1'-Me, 4-OPh, 4-n-propyl, 4-n-butyl, 4-SO₂Et, 4-SO₂-n-propyl, 4-SO₂Ph or 4-SOMe.


12. The compound as defined in Claim 1 having the structure:



where	A:	R ³ :
	Bond (meta link)	H
	Bond (meta link)	3-Me
	Bond (meta link)	4-MeO
	(CH ₂) ₂ (meta link)	H
	(CH ₂) ₂ (meta link)	4-Me

(CH ₂) ₃ (meta link)	H
(CH ₂) ₃ (meta link)	4-Me
(CH ₂) ₃ (meta link)	3-Me
Bond (para link)	H
CH ₂ (ortho link)	H
CH ₂ (ortho link)	4-Et
O (meta link)	4-Me
S (meta link)	4-Me.

13. The compound as defined in Claim 1 having the structure:




where

A:	R¹:	R²:	R³:
CH ₂	2-Me	H	4-Et
CH ₂	4-Me	H	4-Et
CH ₂	4-Me	H	4-SO ₂ Me
CH ₂	4-Me	H	4-OH
CH ₂	4-Me	H	4-S(O)Me
CH ₂	4-Me	H	4-F
CH ₂	4-Me	H	4-Cl

CH ₂	4-Me	H	4-Me
CH ₂	4-Me	H	H
CH ₂	4-Me	6-Me	4-OMe
CH ₂	4-F	H	4-OMe
CH ₂	4-Cl	H	4-SOMe
CH ₂	4-Cl	H	4-SO ₂ Me
CH ₂	4-Cl	H	4-OCHF ₂
CH ₂	4-Et	H	4-OMe
CH ₂	4-iPr	H	4-OMe
CH ₂	4-iPr	H	4-SMe
CH ₂	4-iPr	H	4-SO ₂ Me
CH ₂	4,5-OCH ₂ O	H	4-Et
CH ₂	5-Me	H	4-Et
CH ₂	5-Me	6-Me	4-OMe
CH ₂	6-Me	H	4-Et-

14. The compound as defined in Claim 1 having the structure



15. A pharmaceutical composition comprising a compound as defined in any one of claims 1-14 and a pharmaceutically acceptable carrier thereof.

16. A pharmaceutical combination comprising the compound as defined in any one of claims 1-14 and an antidiabetic agent other than an SGLT2 inhibitor, an agent for treating the complications of diabetes, an anti-obesity agent, an antihypertensive agent, an antiplatelet agent, an antiatherosclerotic agent, or a lipid-lowering agent.

17. The pharmaceutical combination as defined in Claim 16 comprising said compound and an antidiabetic agent.

18. The combination as defined in Claim 17 wherein the antidiabetic agent is one or more of a biguanide, a sulfonyl urea, a glucosidase inhibitor, a PPAR γ agonist, a PPAR α/γ dual agonist, an α P2 inhibitor, a DP4 inhibitor, an insulin sensitizer, a glucagon-like peptide-1 (GLP-1), insulin, a meglitinide, a PTP1B inhibitor, a glycogen phosphorylase inhibitor, or a glucos-6-phosphatase inhibitor.

19. The combination as defined in Claim 18 wherein the antidiabetic agent is one or more of metformin, glyburide, glimepiride, glipyrizide, glipizide, chlorpropamide, gliclazide, acarbose, miglitol, pioglitazone, troglitazone, rosiglitazone, insulin, (S)-2-(2-benzoylphenylamino)-3-(4-(2-(5-methyl-2-phenyloxazol-4-yl)ethoxy)phenyl)propanoic acid, isaglitazone, 4-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]benzyl]-3,5-isoxazolidinedione, balaglitazone, 2-(2-(2-(4-phenoxy-2-propylphenoxy)ethyl)-1H-indol-5-yl)acetic acid, (Z)-1,4-bis{4-[(3,5-dioxo-1,2,4-oxadiazolidin-2-yl)methyl]phenoxy}but-2-ene, (+/-)-5-[4-(5-

Methoxy-1H-benzimidazol-2-ylmethoxy)benzyl]thiazolin-2,4-dione hydrochloride, ((3-((2R)-((2R)-3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yloxy)acetic acid, repaglinide, nateglinide, mitiglinide, tesaglitazar, (2S)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]-2-[(4-oxo-4-phenyl-butan-2-yl)amino]propanoic acid, 5-((2,4-dioxo-5-thiazolidinyl)methyl)-2-methoxy-N-((4-(trifluoromethyl)phenyl)methyl)benzamide, Exendin-4, Des-HAEGTFTSDVSSYLEGQAA-Arg26-EFIAWLV-Lys-(octanoyl)-GRG, or, 1-((2-((5-cyano-2-pyridinyl)amino)ethyl)amino)acetyl-2-pyrrolidinecarbonitrile.

20. The combination as defined in Claim 17 wherein said compound is present in a weight ratio to the antidiabetic agent within the range from 0.01 to 300:1.

21. The combination as defined in Claim 16 wherein the anti-obesity agent is a beta 3 adrenergic agonist, a lipase inhibitor, a serotonin and dopamine reuptake inhibitor, a thyroid receptor beta compound, or an anorectic agent.

22. The combination as defined in Claim 21 wherein the anti-obesity agent is orlistat, celistat, ((3-((2R)-((2R)-3-chlorophenyl)-2-hydroxyethyl)amino)propyl)-1H-indol-7-yloxy)acetic acid, N-[4-[2-[3-(6-Aminopyridin-3-yloxy)-2(S)-hydroxypropylamino]ethyl]phenyl]-4-isopropylbenzenesulfonamide, 4-(2-(2-(6-aminopyridin-3-yl)-2(R)-hydroxyethylamino)ethoxy)phenyl)acetic acid, sibutramine, topiramate, axokine, dexamphetamine, phentermine, phenylpropanolamine, or mazindol.

wherein

R^1 , R^2 and R^{2a} are independently hydrogen, OH, OR^5 , (C_1-C_8) alkyl, CF_3 , $OCHF_2$, OCF_3 , SR^{5i} or halogen, or two of R^1 , R^2 and R^{2a} together with the carbons to which they are attached can form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 ;

R^3 and R^4 are independently hydrogen, OH, OR^{5a} , $O-(C_6-C_{10})$ aryl, $OCH_2-(C_6-C_{10})$ aryl, (C_1-C_8) alkyl, (C_3-C_{20}) cycloalkyl, CF_3 , $-OCHF_2$, $-OCF_3$, halogen, $-CN$, $-CO_2R^{5b}$, $-CO_2H$, $-CONR^6R^{6a}$, $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2-(C_6-C_{10})$ aryl, (C_6-C_{10}) aryl, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$, $-SO_2-(C_6-C_{10})$ aryl, or a five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 , or R^3 and R^4 together with the carbons to which they are attached form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 ;


R^5 , R^{5a} , R^{5b} , R^{5c} , R^{5d} , R^{5e} , R^{5f} , R^{5g} and R^{5i} are independently (C_1-C_8) alkyl;

R^6 and R^{6a} are independently hydrogen, (C_1-C_{20}) alkyl, (C_6-C_{10}) aryl, (C_1-C_{20}) alkyl (C_6-C_{10}) aryl, (C_3-C_{20}) cycloalkyl, or R^6 and R^{6a} together with the nitrogen to which they are attached form an annelated five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 ;

A is O, S, NH, or $(CH_2)_n$ where n is 0 - 3, or a pharmaceutically acceptable salt thereof, with the proviso that where A is $(CH_2)_n$ where n is 0,1,2, or 3 or A is O, and at least one of R^1 , R^2 , R^{2a} , R^3 and R^4 is OH or OR^5 , then at least one of R^1 , R^2 , and R^{2a} is CF_3 , OCF_3 , or $OCHF_2$ or at

least one of R^3 and R^4 is CF_3 , $-OCHF_2$, $-OCF_3$, $-CN$, $-CO_2R^{5b}$, $CH(OR^{5h})R^{6d}$, $CH(OH)R^{6c}$, COR^{6b} , $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2-(C_6-C_{10})aryl$, $(C_6-C_{10})aryl$, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$, $-SO_2-(C_6-C_{10})aryl$ or halogen, or at least one of R^1 , R^2 , and R^{2a} is CF_3 , OCF_3 , or $OCHF_2$ and at least one of R^3 and R^4 is CF_3 , $-OCHF_2$, $-OCF_3$, $-CN$, $-CO_2R^{5b}$, $CH(OR^{5h})R^{6d}$, $CH(OH)R^{6c}$, COR^{6b} , $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2-(C_6-C_{10})aryl$, $(C_6-C_{10})aryl$, $-SR^{5e}$, $-SOR^{5f}$, $-SO_2R^{5g}$, $-SO_2-(C_6-C_{10})aryl$ or halogen;

or a compound of the structure



wherein

R^1 , R^2 and R^{2a} are independently hydrogen, OH, OR^5 , (C_1-C_8) alkyl, CF_3 , $OCHF_2$, OCF_3 , SR^{5i} or halogen, or two of R^1 , R^2 and R^{2a} together with the carbons to which they are attached can form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO_2 ;

R^3 and R^4 are independently hydrogen, OH, OR^{5a} , $O-(C_6-C_{10})aryl$, $OCH_2-(C_6-C_{10})aryl$, (C_1-C_8) alkyl, $(C_3-C_{20})cycloalkyl$, CF_3 , $-OCHF_2$, $-OCF_3$, halogen, $-CN$, $-CO_2R^{5b}$, $-CO_2H$, $-CONR^6R^{6a}$, $-NHCOR^{5c}$, $-NHSO_2R^{5d}$, $-NHSO_2-(C_6-C_{10})aryl$,

(C₆-C₁₀)aryl, -SR^{5e}, -SOR^{5f}, -SO₂R^{5g}, -SO₂-(C₆-C₁₀)aryl, or a five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO₂, or R³ and R⁴ together with the carbons to which they are attached form an annelated five, six or seven membered carbocycle or heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO₂;

R⁵, R^{5a}, R^{5b}, R^{5c}, R^{5d}, R^{5e}, R^{5f}, R^{5g} and R⁵ⁱ are independently (C₁-C₈) alkyl;

R⁶ and R^{6a} are independently hydrogen, (C₁-C₂₀)alkyl, (C₆-C₁₀)aryl, (C₁-C₂₀)alkyl(C₆-C₁₀)aryl, (C₃-C₂₀)cycloalkyl, or R⁶ and R^{6a} together with the nitrogen to which they are attached form an annelated five, six or seven membered heterocycle which may contain 1 to 4 heteroatoms in the ring which are N, O, S, SO, or SO₂;


A is O, S, NH, or (CH₂)_n where n is 0 - 3, or a pharmaceutically acceptable salt thereof.

27. Use of a therapeutically effective amount of a compound as defined in any one of claims 1-14 for treating or delaying the progression or onset of diabetes, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, delayed wound healing, insulin resistance, hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, hyperlipidemia, obesity, hypertriglyceridemia, Syndrome X, diabetic complications, atherosclerosis or hypertension, or for increasing high density lipoprotein levels in a mammalian species.

28. Use of a therapeutically effective amount of a compound as defined in any one of claims 1-14 in the manufacture of a medicament for treating or delaying the

progression or onset of diabetes, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, delayed wound healing, insulin resistance, hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, hyperlipidemia, obesity, hypertriglyceridemia, Syndrome X, diabetic complications, atherosclerosis or hypertension, or for increasing high density lipoprotein levels in a mammalian species.

29. A use as defined in claim 27 or 28 where the compound has the structure



30. Use of a therapeutically effective amount of a compound as defined in any one of claims 1-14 alone or in combination with another antidiabetic agent, an agent for treating the complications of diabetes, an anti-obesity agent, an antihypertensive agent, an antiplatelet agent, an antiatherosclerotic agent, or a lipid-lowering agent for treating type II diabetes in a mammalian species.

31. Use of a therapeutically effective amount of a compound as defined in any one of claims 1-14 alone or in combination with another antidiabetic agent, in the manufacture of a medicament for treating the complications of diabetes, an anti-obesity agent, an antihypertensive agent, an antiplatelet agent, an antiatherosclerotic agent,

or a lipid-lowering agent for treating type II diabetes in a mammalian species.

