



US 20050229267A1

(19) **United States**(12) **Patent Application Publication**
Rommens et al.(10) **Pub. No.: US 2005/0229267 A1**(43) **Pub. Date: Oct. 13, 2005**(54) **PRECISE BREEDING****Publication Classification**(75) Inventors: **Caius Rommens**, Boise, ID (US);
Jingson Ye, Boise, ID (US); **Hua Yan**,
Boise, ID (US); **Kathy M. M. Swords**,
Boise, ID (US); **Jaime**
Menendez-Humara, Boise, ID (US);
W. Leigh Brinkerhoff, Meridian, ID
(US); **Craig Richael**, Meridian, ID
(US)(51) **Int. Cl.⁷** **C07H 21/04**; A01H 1/00;
C12N 15/82
(52) **U.S. Cl.** **800/278**; 800/294; 435/468;
536/23.6(57) **ABSTRACT**Correspondence Address:
FOLEY AND LARDNER
SUITE 500
3000 K STREET NW
WASHINGTON, DC 20007 (US)(73) Assignee: **J.R. Simplot Company**(21) Appl. No.: **10/505,079**(22) PCT Filed: **Feb. 20, 2003**(86) PCT No.: **PCT/US03/04947****Related U.S. Application Data**(60) Provisional application No. 60/357,661, filed on Feb.
20, 2002. Provisional application No. 60/377,602,
filed on May 6, 2002.

The present invention relates to a new plant breeding process. The process improves the agronomic performance of crop plants by using genetic material that is also used in classical breeding. Instead of sexually recombining entire genomes at random, as is done in classical breeding, specific genetic elements are rearranged in vitro and inserted back into individual plant cells. Plants obtained through this new plant breeding process do not contain foreign nucleic acid but only contain nucleic acid from the plant species selected for transformation or plants that are sexually compatible with the selected plant species. Plants developed through this new plant breeding process are provided. In particular, potato plants displaying improved tuber storage and health characteristics are provided.

Figure 1. Diagrams for some P-DNA vectors




Figure 2. Alignment of potato and tobacco invertase inhibitor proteins**A.**

```

St-inh1      MRNLFPIMLITNLALNNDNNNNNNNNNNYLIHATCRETPYYSLCLTTLQSGPRSNEVE 60
Nt-inhh      MRNLFPIFMLITNLAFN-DNNNSNN-----IINTTCRATTNYPLCLTTLHSDPRTSEAE 53
              *****:*****.* ****.*          :*:*** *. *.*****:*.**:*.*
St-inh1      GGDAITTLGLIMVDAVKSKSIEIMEKIKELEKSNPEWRAPLSQCYVAYNAVLRADVTAV 120
Nt-inhh      GAD-LTTLGLVMVDAVKLKSIEIMKSIKKLEKSNPELRPLSQCYIVYYAVLHADVTAV 112
              *. * :*****:***** *****:*.**:****** * *****:.* **:******
St-inh1      EALKKGAPKFAEDGMDDVVAEAQTCEYSFNYYNKLDFFISNLSREIIELSKVAKSIIRML 180
Nt-inhh      EALKRGVPKFAENGMDVVAEAEETCEFSFK-YNGLVSPVSDMNKEIIELSSVAKSIIRML 171
              *****:*.*****:*** **..**:****:***: ** * *:***:..*****.*****
St-inh1      L 181
Nt-inhh      L 172
              *

```

B.

```

St-inh1      MRNLFPIMLITNLALNNDNNNNNNNNNNYLIHATCRETPYYSLCLTTLQSGPRSNEVE 60
Nt-inh1      MKNLIFLTMFLTILLQTNANN-----LVETTCKNTPNYQLCLKTLLSDKRS--AT 48
              *:***: :*:*.* * . * **          *:..***:*** *.*****.*. ** .
St-inh1      GGDAITTLGLIMVDAVKSKSIEIMEKIKELEKSNP--EWRAPLSQCYVAYNAVLRADVTV 118
Nt-inh1      G--DITTLALIMVDAIKAKANQAATISKLRHSNPPAAWKGPLKNCAFSYKVILTASLPE 106
              * ****.*****:*.**:* : .*:*.**.* *:..**:* .*:..* *..
St-inh1      AVEALKKGAPKFAEDGMDDVVAEAQTCEYSFNYYNKLDFFISNLSREIIELSKVAKSIIR 178
Nt-inh1      AIEALTKGDPKFAEDGMVGSSGDAQECE--EYFKGSKSPFSALNIAVHELSDVGRAIVR 163
              *:***.* ***** . .:** ** *:.. .*:.*. : ***.*:~*:
St-inh1      MLL 181
Nt-inh1      NLL 166
              **

```

Figure 3. Gene-free expression cassettes


<i>Promoter</i>	<i>SOI*</i>	<i>Spacer</i>	<i>Total size</i>
P:GBSS-small	Leader a/w* <i>R1</i>	Ubi intron	1729-bp
P:GBSS-small	Leader a/w <i>R1</i>	GBSS spacer	1397-bp
P:GBSS-large	Leader a/w <i>R1</i>	Ubi intron	2005-bp
P:GBSS-large	Leader a/w <i>R1</i>	GBSS spacer	1397-bp
P:GBSS-small	Trailer a/w <i>R1</i>	GBSS spacer	2042-bp
P:GBSS-small	Trailer a/w <i>R1</i>	Ubi intron	1705-bp
P:GBSS-large	Trailer a/w <i>R1</i>	GBSS spacer	2313-bp
P:GBSS-large	Trailer a/w <i>R1</i>	Ubi intron	1981-bp
P:GBSS-small	Leader a/w <i>Phosph.</i>	GBSS spacer	-
P:GBSS-small	Leader a/w <i>Phosph.</i>	Ubi intron	-
P:GBSS-large	Leader a/w <i>Phosph.</i>	GBSS spacer	1852-bp
P:GBSS-large	Leader a/w <i>Phosph.</i>	Ubi intron	2184-bp
P:GBSS-large	Leader a/w <i>PPO</i>	Ubi intron	1958-bp
P:GBSS-large	Leader a/w <i>PPO</i>	GBSS spacer	1626-bp
P:GBSS-small	Leader a/w <i>PPO</i>	Ubi intron	-
P:GBSS-small	Leader a/w <i>PPO</i>	GBSS spacer	-

*: sequence-of-interest; **: "associated with"

Figure 4. Alignment of the 3'-end of tuber-expressed PPO genes and trailers associated with these genes. Stop codons ('TAA') are underlined. The trailer sequence used for genetic modification of potato plants was isolated from 'P-PPO3', and is downstream from the stop codon (TTAGTC...ACAATT).

P-PPO3	CTGGCGATAACGGAAGTGTGGAGGATATTGGTTTGGGAAGATGAAGATACTATTGCGGTG	60
PPOM-41	CTGGCGATAACGGAAGTGTGGAGGATATTGGATTGGAAGATGAAGATACTATTGCGGTG	60
PPOM-44	CTGGCGATAACGGAAGTGTGGAGGATAATGGATTGGAAGATGAAGGTACTATNGCGGTG	60
P-PPO3	ACTCTGGTGCCAAAGAGAGGGTGGTGAAGGTATCTCCATTGAAAGTGCGACGATCAGTCTT	120
PPOM-41	ACTTTGGTTCCAAAAGTAGGTGGTGAAGGTGTATCCATTGAAAGTGTGGAGATCAAGCTT	120
PPOM-44	ACTTTGGTTCCAAAAGTTGGTGGTGAAGGTGTATCCATTGAAAGTGCGGAGATCAAGCTT	120
P-PPO3	GCAGATTGTTAATTAGTCTCTA-TTGA-ATCTGCTG---AGATTACAC-TTTGATGGAT	173
PPOM-41	GAGGATTGTTAAGTCCTCATGAGTTGGTGGCTACGGTACCAAATTTATGTTTAATTAGT	180
PPOM-44	GAGGATTGTTAAGTCCTCATGAGTTGGTGGCTATGGTACCAAATTNTATGTTTAATTAGT	180
P-PPO3	GATGCTCTGTT--TTTGTTCCTTCTGTTTTCCTC-TGTTGAAATCAGCTTTGTT	230
PPOM-41	ATTAATGTGTGTATGTGTTTGATTATGTTTCGGTTAAATGTATCAGCTGGATAGCTGAT	240
PPOM-44	ATTAATGTGTG---TGTTTGATTATGTTTCGGTTAAATGTATCANCTGGATAGCTGAT	236
P-PPO3	-GCTTGATTTC---ATTGAAGTTGTTATTCAAGAA-TAAATCAGTTA-CAATT-----	277
PPOM-41	TACTAGCCTTGCCAGTTGTTAATGCTATGTATGAAATAAATAAATAAATGGTTGCTCTTCT	300
PPOM-44	TACTAGCCTTCCAGTTGTTAATGCTATGTATGAAATACATAAATAAATGGTTGCTCTCC	296

Figure 5. Diagrams for some LifeSupport-vectors.



PRECISE BREEDING

FIELD OF THE INVENTION

[0001] This application claims priority to U.S. provisional application Ser. numbers 60/357,661 and 60/377,602, which are incorporated herein by reference. The present invention relates to methods for improving the nutritional, health, and agronomic characteristics of a plant by modifying specific, well-characterized DNA in the plant's genome. As opposed to classical plant breeding, the inventive process does not introduce unknown or potentially toxic genes into the plant genetic make-up. Furthermore, the inventive method, unlike conventional genetic engineering strategies, does not incorporate nucleic acids from foreign species, i.e., species that are not inter-fertile with the plant to be modified by genetic engineering, into the plant genome. Plants developed through the inventive plant breeding process display improved agronomic characteristics. Particularly preferred plants of the present invention include potatoes that exhibit improved health and tuber storage characteristics, and turf-grasses that exhibit improved disease and drought tolerance.

BACKGROUND

[0002] The agronomic performance of plants has typically been improved by either classical plant breeding or genetic engineering. Classical breeding typically results in the transfer of unknown nucleic acids from one plant to another. Genetic engineering techniques introduce foreign nucleic acids into the plant genome, i.e., DNA that is not from a plant or that is not from a plant that is naturally interfertile with the plant to be modified by genetic engineering. For example, genetic engineering introduces non-plant nucleic acids into a plant genome. Both classical breeding and genetic engineering strategies create plant genomes that contain undesirable and unwanted genetic material, and the resultant cross-bred or transgenic plants can exhibit unfavorable traits. The inadequacies of both strategies can prove harmful to the transgenic plants, as well as to the animals and humans who consume such products.

[0003] Conventional Breeding Relies on the Transfer of Unknown DNA

[0004] Plant breeding typically relies on the random recombination of plant chromosomes to create varieties that have new and improved characteristics. Thus, by screening large populations of progeny that result from plant crosses, breeders can identify those plants that display a desired trait, such as an increase in yield, improved vigor, enhanced resistance to diseases and insects, or greater ability to survive under drought conditions. However, classical breeding methods are laborious and time-consuming, and new varieties typically display only relatively modest improvements.

[0005] Furthermore, classical plant breeding typically results in the transfer of hundreds of unknown genes into a plant genome. It is likely that some of those transferred genes encode potentially harmful allergens, such as patatin, lectins, chitinases, proteases, thaumatin-like proteins, lipid transfer proteins, amylases, trypsin inhibitors, and seed storage proteins (Breiteneder et al., *J Allergy Clin Immunol* 106: 27-36).

[0006] Similarly, introgressed genes can be involved in the biosynthesis of toxins including lathrogens, hydrazines,

glucosinolates and goitrogens, coumarins, saponins, alkaloids, glycoalkaloids, biogenic amines, enzyme inhibitors, such as lectins (haemagglutinins), trypsin inhibitors, chelating substances such as phytates and oxalates, ribotoxins, antimicrobial peptides, amino acids such as beta-N-oxalyl-L-alanine, atractyloside, oleandrine, taxol, and isoquinoline (Pokorny, *Cas Lek Cesk* 136: 267-20 70, 1997). The risk of inadvertently introducing such poisons into human and animal food supplies is further increased through efforts to "untap" the genetic diversity of wild crop relatives that have not been used before for food consumption (Hoisington et al., *Proc Natl Acad Sci USA* 96: 593743, 1999).

[0007] Although classical plant breeding can easily introduce genes involved in undesirable anti-nutritional compounds into food crops and plants, it cannot easily remove them. For instance, it took about 15 years to reduce harmful phytate levels in corn and rice by inactivating *Lpa* genes (Raboy, *J Nutr* 132: 503S-505S, 2002). The long timeframe for realizing positive results is not practical, especially since there is an urgent need for methods that more effectively and efficiently improve the quality of food crops. One example of a gene that only recently was found to be associated with the synthesis of anti-nutritional compounds is the polyphenol oxidase (PPO) gene, which oxidizes certain phenolic compounds to produce mutagenic, carcinogenic and cytotoxic agents like phenoxyl radicals and quinoid derivatives (Kagan et al., *Biochemistry* 33: 9651-60, 1994). The presence of multiple copies of this gene in the genome of plants such as potato makes it particularly difficult to reduce PPO activity through breeding.

[0008] Even more time is needed for the removal of anti-nutritional compounds if little or nothing is known about their genetic basis. For instance, no genes have been linked to the accumulation of high concentrations of acrylamide, a potent neurotoxin and mutagen, in some potatoes that are heated to 160° C. or higher (Tareke et al., *J Agric Food Chem.* 50: 4998-5006, 2002). It is therefore very difficult to efficiently develop new potato varieties that produce less acrylamide during processing using conventional breeding. Thus, there is a need to grow potatoes and other carbohydrate-rich foods, such as wheat, with reduced levels of such dangerous compounds, but without the use of unknown or foreign nucleic acids.

[0009] Other anti-nutritional compounds that can accumulate during processing and are difficult to minimize or eliminate through breeding include the Maillard-reaction products N-Nitroso-N-(3-keto-1,2-butanediol)-3'-nitrotyramine (Wang et al., *Arch Toxicol* 70: 10-5, 1995), and 5-hydroxymethyl-2-furfural (Janzowski et al., *Food Chem Toxicol* 38: 801-9, 2000). Additional Maillard reaction products that have not been well characterized are also known to display mutagenic properties (Shibamoto, *Prog Clin Biol Res* 304: 359-76, 1989).

[0010] It can be equally difficult to rapidly increase levels of positive nutritional compounds in food crops due to the inherent imprecision of conventional plant breeding. For instance, it would be desirable to increase levels of "resistant starch" (Topping et al., *Physiol Rev* 81: 1031-64, 2001) in a variety of crops. Such starch is ultimately responsible for promoting immune responses, suppressing potential pathogens, and reducing the incidence of diseases including colorectal cancer (Bird et al., *Curr Issues Intest Microbiol* 1:

25-37, 2000). However, the only available plants with increased levels of resistant starch are low-yielding varieties like maize mutants “amylose extender”, “dull”, and “sugary-2.” Creation of new high resistant starch sources, such as potato, would enable broader dietary incorporation of this health-promoting component.

[0011] The inability to safely manipulate the genotypes of plants often leads to the use of external chemicals to induce a desired phenotype. Despite numerous breeding programs to delay tuber sprouting, for example, no potato varieties are available commercially that can be stored for months without treatment with sprout inhibitors. The latter, such as isopropyl-N-chlorophenyl-carbamate (CIPC), is linked to acute toxicity and tumor development, and can be present in processed potato foods at concentrations between 1 mg/kg and 5 mg/kg.

[0012] Genetic Engineering Relies on the Transfer of Foreign DNA

[0013] Genetic engineering can be used to modify, produce, or remove certain traits from plants. While there has been limited progress in improving the nutritional value and health characteristics of plants, most improvements target plant traits that promote ease of crop cultivation. Thus, certain plants are resistant to the glyphosate herbicide because they contain the bacterial gene 5-enolpyruvylshikimate-3-phosphate synthase (Padgett et al., *Arch Biochem Biophys.* 258: 564-73, 1987). Similarly, genetic engineering has produced insect-, viral-, and fungal-resistant plant varieties (Shah et al., *Trends in Biotechnology* 13: 362-368, 1995; Gao et al., *Nat Biotechnol.* 18: 1307-10, 2000; Osusky et al., *Nat Biotechnol.* 18: 1162-6, 2000), but few with enhanced nutrition or health benefits.

[0014] According to standard, well-known techniques, genetic “expression cassettes,” comprising genes and regulatory elements, are inserted within the borders of *Agrobacterium*-isolated transfer DNAs (“T-DNAs”) and integrated into plant genomes. Thus, *Agrobacterium*-mediated transfer of T-DNA material typically comprises the following standard procedures: (1) in vitro recombination of genetic elements, at least one of which is of foreign origin, to produce an expression cassette for selection of transformation, (2) insertion of this expression cassette, often together with at least one other expression cassette containing foreign DNA, into a T-DNA region of a binary vector, which usually consists of several hundreds of basepairs of *Agrobacterium* DNA flanked by T-DNA border sequences, (3) transfer of the sequences located between the T-DNA borders, often accompanied with some or all of the additional binary vector sequences from *Agrobacterium* to the plant cell, and (4) selection of stably transformed plant cells. See, e.g., U.S. Pat. Nos. 4,658,082, 6,051,757, 6,258,999, 5,453,367, 5,767,368, 6,403,865, 5,629,183, 5,464,763, 6,201,169, 5,990,387, 4,693,976, 5,886,244, 5,221,623, 5,736,369, 4,940,838, 6,153,812, 6,100,447, 6,140,553, 6,051,757, 5,731,179, 5,149,645 and EP 0 120,516, EP 0 257,472, EP 0 561,082, 1,009,842A1, 0 853,675A1, 0 486,233B1, 0 554,273A1, 0 270,822A1, 0 174,166A1, and WO 01/25459.

[0015] Thus, genetic engineering methods rely on the introduction of foreign nucleic acids into the food supply. Those techniques transfer complex fusions of a few to more than 20 genetic elements isolated from viruses, bacteria, and plants, that are not indigenous to the transformed plant

species. Such foreign elements include regulatory elements such as promoters and terminators, and genes that are involved in the expression of a new trait or function as markers to identify or select for transformation events. Despite the testing of foods containing foreign DNA for safety prior to regulatory approval, many consumers are concerned about the long-term effects of eating foods that express foreign proteins, which are produced by genes obtained from other, non-plant species.

[0016] One commonly used regulatory element is the 35S “super” promoter of cauliflower mosaic virus (CaMV), which is typically used in plant engineering to induce high levels of expression of transgenes to which it is directly linked. However, the 35S promoter also can enhance the expression of native genes in its vicinity (Weigel et al., *Plant Physiol.*, 122: 1003-13, 2000). Such promoters may thus induce unpredictable alterations in the expression of endogenous genes, possibly resulting in undesirable effects such as increased alkaloid production. Preferred “strong” promoters are generally those isolated from viruses, such as rice tungro bacilliform virus, maize streak virus, cassava vein virus, mirabilis virus, peanut chlorotic streak caulimovirus, figwort mosaic virus and chlorella virus. Other frequently used promoters are cloned from bacterial species and include the promoters of the nopaline synthase and octopine synthase gene.

[0017] To obtain appropriate termination of gene translation, terminator sequences are fused to the 3' end of transgenes and include genetic elements from the nopaline synthase and octopine synthase genes from *Agrobacterium*. Other genetic elements may be used to further enhance gene expression or target the expressed protein to certain cell compartments. These elements include introns to boost transgene expression and signal peptide sequences to target the foreign gene to certain cellular compartments, often derived from foreign plant species.

[0018] Certain genes involved in expression of a new trait are most frequently derived from foreign sources. If native genes are used, they are often inverted to silence the expression of that gene in transgenic plants and co-transformed with foreign DNA such as a selectable marker. The main disadvantage of this “antisense” technology is that the inverted DNA usually contains new and uncharacterized open reading frames inserted between a promoter and terminator. Thus, potato plants that were genetically modified with antisense constructs derived from the starch related gene R1 (Kossmann et al., U.S. Pat. No. 6,207,880), the L- and H-type glucan phosphorylase genes (Kawchuk et al., U.S. Pat. No. 5,998,701, 1999), the polyphenol oxidase gene (Steffens, U.S. Pat. No. 6,160,204, 2000), and genes for starch branching enzymes I and II (Schwall et al., *Nature Biotechnology* 18: 551-554, 2000) all potentially express new peptides consisting of at least 50 amino acids (Table 1). These new peptides may interfere with plant development and/or reduce the nutritional value of potato, and are therefore undesirable.

[0019] Conventional marker genes are incorporated into genetic constructs and used to select for transformation events. They confer either antibiotic or herbicide resistance (U.S. Pat. No. 6,174,724), a metabolic advantage (U.S. Pat. No. 5,767,378), or a morphologically abnormal phenotype (U.S. Pat. No. 5,965,791) to the transformed plant. Such

markers are typically derived from bacterial sources. Furthermore, because of the infidelity of T-DNA transfer, about 75% of transformation events in plants such as tomato, tobacco, and potato contain plasmid “backbone” sequences in addition to the T-DNA (Kononov et al., *Plant J.* 11: 945-57, 1997). The presence of such backbone sequences is undesirable because they are foreign and typically contain origins of replication and antibiotic resistance gene markers.

[0020] There do exist various methods for removing elements like foreign marker genes, but few are easily applicable to plant genetic engineering. According to one such method, the marker gene and desired gene or nucleotide sequence are placed on different vectors. The infection of plants with either a single *Agrobacterium* strain carrying both vectors (U.S. Pat. No. 6,265,638) or two *Agrobacterium* strains each of which carries one of the vectors can occasionally result in unlinked integration events, which may be separated genetically through outbreeding. The main disadvantage of this method is that the genetic separation of loci can be very laborious and time-consuming, especially if T-DNA integration events are linked. Furthermore, this method is not widely applicable in apomictic plants, which reproduce asexually, such as Kentucky bluegrass, or vegetatively propagated crops such as potato, which cannot be readily bred due to inbreeding depression, high levels of heterozygosity, and low fertility levels.

[0021] Another method for removing foreign genetic elements relies on inserting the foreign gene, like the selectable marker, into a transposable element. The modified transposable element may then be spliced out from the genome at low frequencies. Traditional crosses with untransformed plants must then be performed to separate the transposed element from the host (U.S. Pat. No. 5,482,852). As described for the previous method, this alternative method cannot be used for vegetatively propagated or apomictic plant systems.

[0022] A third method of removing a marker gene uses the Cre/lox site-specific recombination system of bacteriophage P1 (Dale & Ow, *Proc. Natl. Acad. Sci. USA*, 88: 10558-62, 1991). Insertion of a marker gene together with the Cre recombinase gene and a chimeric gene involved in induction of Cre (both with their own promoters and terminators) between two lox sites leads to excision of the region delineated by the lox sites during the regeneration process (Zuo et al., *Nat. Biotechnol.*, 19: 157-61, 2001). This complicated process is inefficient and not reliable, and may cause genome instability.

[0023] Recent studies report that some plant genes themselves may be used as transformation markers. Examples of such plant markers include Pga22 (Zuo et al., *Curr Opin Biotechnol.* 13: 173-80, 2002), Cki1 (Kakimoto, *Science* 274: 982-985, 1996) and Esr1 (Banno et al., *Plant Cell* 13: 2609-18, 2001). All of the genes, however, trigger cytokinin responses, which confer an undesirable phenotype to the transformed plant. Furthermore, such plant markers would still need to be removed upon transformation by any of the methods described above.

[0024] Alternative methods to transform plants are also based on the in vitro recombination of foreign genetic elements, and rely on bacterial plasmid sequences for maintenance in *E. coli*, parts of which are co-integrated during the transformation process. Examples of such methods to trans-

form plants with foreign DNA are described in U.S. Pat. Nos. 5,591,616, 6,051,757, 4,945,050, 6,143,949, 4,743,548, 5,302,523, and 5,284,253.

[0025] Marker-free transgenic plants may also be obtained by omitting any selection procedures prior to regeneration. A disadvantage of this method is that most events generated through this method will represent untransformed or chimeric plants because they will usually not be derived from single transformed plant cells. It is extremely difficult and laborious to use a marker-free procedure for the identification of transgenic plants that contain the same DNA insertion(s) in all their cells.

[0026] Thus, there is a very important need to improve plants beyond that which can be accomplished through the classical breeding crosses and conventional genetic engineering techniques, and which does not rely on the insertion of unknown or foreign nucleic acid into a plant genome. Accordingly, the present invention provides methods and compositions for precisely modifying a plant's own genetic material. Thus, the inventive “precise breeding” strategy does not induce undesirable phenotypes and does not introduce unknown or foreign nucleic acid into a plant genome.

SUMMARY

[0027] The present invention provides methods of genetically enhancing the nutritional value and agronomic performance of a plant without the permanent or stable incorporation of either unknown or foreign DNA into the genome of that plant. According to the methods of the present invention, specific, well-characterized nucleic acids, gene elements, and genes are isolated from a desired plant species or from a plant species that is sexually compatible with the desired plant, modified, and then reinserted back into the genome of the desired plant species. The modification may entail mutating the isolated nucleic acid sequence, deleting parts of the isolated nucleic acid, or simply joining the isolated nucleic acid to another polynucleotide, such as subcloning the isolated nucleic acid into a plasmid vector.

[0028] Accordingly, transgenic plants produced by the inventive methodology do not possess genomes that comprise any foreign species' nucleic acids. Thus, the methods of the present invention produces a transgenic plant whose genome does not comprise a non-plant species promoter, does not comprise a non-plant species terminator, does not comprise a non-plant species 5'-untranslated region, does not comprise a non-plant species 3'-untranslated region, does not comprise a non-plant species marker gene, does not comprise a non-plant species regulatory element, does not comprise a non-plant species gene, and does not comprise any other polynucleotide that is obtained from a non-plant species genome.

[0029] Thus, the present invention provides a method for producing a stable transgenic plant that exhibits a modified phenotype that is not exhibited by the non-transformed plant, comprising (a) transforming plant cells with a desired polynucleotide; (b) growing plants from the transformed cells; and (c) selecting a plant stably transformed with said desired polynucleotide which exhibits a new phenotype that is not exhibited by plants grown from the corresponding non-transformed plant cells. Preferably, the desired polynucleotide consists essentially of (i) nucleic acid sequences that are isolated from and/or native to the genome of the

plant cells, or to other plants of the same species, or are isolated from and/or native to the genome of a plant species that is sexually compatible with the plant from which the plant cells were isolated; and (ii) at least one DNA sequence that is a border-like sequence that has a sequence that is native to the genome of said plant cells or is native to the genome of plant cells of the same species, or is native to a plant that is sexually compatible with the plant from which the plant cells were isolated, and wherein the border-like sequence is capable of stably integrating the desired polynucleotide into the genome of said plant cells.

[0030] A preferred method of the present invention entails producing a transgenic plant that exhibits a modified phenotype that is not exhibited by the non-transformed plant, comprising (a) infecting explants with *Agrobacterium* carrying (i) a “P-DNA” vector, which contains a desired polynucleotide that is native to the transgenic plant, and (ii) a “LifeSupport” vector that contains an expression cassette containing a selectable marker gene; (b) selecting for transient expression of the selectable marker gene, preferably for 1-10 days, for 3-7 days, or for 4-5 days; (c) transferring explants to regeneration media to allow shoot formation; (d) screening populations of shoots to determine which comprise at least one copy of the desired polynucleotide in their genomes and, of those, which shoots do not contain any foreign nucleic acids, such as the selectable marker gene, in their genomes; and (e) allowing shoots which contain the desired polynucleotide in their genomes but not any marker gene DNA, to develop into whole plants, wherein the resultant whole plants exhibit a modified phenotype that is not exhibited by plants grown from non-transformed plant cells of the same species.

[0031] According to such a method, the desired polynucleotide (i) consists essentially of only elements that are isolated from and/or native to the genome of the plant cell species or sexually compatible species thereof; (ii) comprises at least one border element that has a sequence that is isolated from, or native to, the genome of the plant cell species or sexually compatible species thereof, and is capable of stably integrating the desired polynucleotide into the genome of a plant cell exposed to the vector; and (iii) is stably integrated into the genome of the transformed plant; wherein the method does not integrate non-plant species or foreign DNA into the genome of the transformed plant.

[0032] Furthermore, any selectable marker gene may be used as an indicator of successful transformation. For instance, a “neomycin phosphotransferase” marker gene, or an “hpt” marker gene may be used to confer resistance to the aminoglycoside antibiotics, kanamycin and hygromycin respectively. Other marker genes include the “bar” marker gene, which confers resistance to herbicide phosphinothricin; the “DHFR” marker gene, which confers resistance to methotrexate; and the “ESPS” marker gene, which confers resistance to Round-up herbicide. It is well known in the art how to follow expression of such marker genes to determine whether or not the marker gene has been stably expressed into the genome of a transformed plant cell. Accordingly, the skilled artisan knows how to follow expression of the marker gene to determine that the marker gene is only transiently expressed in the transformed plant cell.

[0033] In another aspect of the invention, there is provided a method of making a stably transformed plant comprising

the steps of: (1) identifying a target gene; (2) isolating a leader or trailer DNA sequence associated with said target gene; (3) optionally modifying said isolated leader or trailer DNA; (4) operably linking said leader or trailer DNA to native regulatory elements to form an expression cassette; (5) inserting said expression cassette into a P-DNA that is located on a binary vector, wherein the binary vector also carries an operable cytokinin gene such that the inadvertent insertion of additional binary vector sequences, which are of foreign origin, are detected by expression of the cytokinin gene; (6) introducing the modified binary vector into *Agrobacterium*; (7) stably integrating the rearranged native DNA into the genomes of plant cells using LifeSupport-mediated transformation; (8) regenerating plant cells that contain the rearranged native DNA; (9) discarding plants that display a cytokinin-overproducing phenotype and do not fully regenerate; and (10) maintaining for further analysis the desirable plants that are indistinguishable from untransformed plants.

[0034] In another aspect of the instant invention, a method of modifying the expression of a trait in a selected plant species is provided. In one embodiment, the method comprises (1) identifying the trait to be modified; (2) constructing a recombinant DNA molecule consisting essentially of genetic elements isolated from, or native to, the selected plant species, wherein the recombinant DNA molecule, when integrated into the genome of the selected plant species, modifies the expression of the trait in the transformed plant species; (3) stably integrating the recombinant DNA molecule into cells of the selected plant species using LifeSupport-mediated transformation; and (4) identifying transformed plants exhibiting modified expression of the trait.

[0035] In a preferred embodiment, polynucleotide that is native to a desired plant is inserted into the desired plant's genome by infecting explants with two different *Agrobacterium* strains. A first *Agrobacterium* strain is capable of transferring the native DNA from P-DNA vectors to plant cells; a second strain can transfer a T-DNA carrying an expression cassette for a selectable marker gene to plant cells. Examples of the latter vector include the so-called, “LifeSupport” vectors described herein. By preferably selecting plants that transiently express the marker gene for 1-10 days, for 3-7 days, or for 4-5 days, and subsequently transferring explants to regeneration media, a population of events is obtained, part of which represents plants that contain at least one copy of the polynucleotide, but which lack any copies of the T-DNA or marker gene.

[0036] In another embodiment, a single *Agrobacterium* strain is used that carries both a P-DNA vector, which houses the desired, native gene of interest or polynucleotide between P-DNA border-like sequences, and a LifeSupport vector, which contains a marker gene. The marker gene may, or may not, be inserted between P-DNA border-like sequences, T-DNA border sequences, or other T-DNA-like border sequences.

[0037] Thus, in another preferred embodiment, the P-DNA vector contains at least two expression cassettes, one of which comprises a native screenable or selectable marker gene driven by a native promoter and followed by a native terminator.

[0038] By preferably selecting for at least 2 days and more preferably for at least 5 days for native marker gene expres-

sion and subsequently transferring explants to regeneration media, a population of events is obtained that represent plants containing at least one copy of the introduced DNA stably integrated into their genomes. In preferred embodiments, the plant-derived marker gene encodes a mutant 5-enolpyruvul-3-phosphoshikimic acid synthase or tryptophan decarboxylase. In a more preferred embodiment, the selectable marker encodes for salt tolerance. In a most preferred embodiment, the salt tolerance gene has the nucleotide sequence shown in SEQ ID 35 and is used to select for transformation events in potato.

[0039] In yet another embodiment, the modified expression of the trait is characterized by an increase in expression, a decrease in expression, or in undetectable expression.

[0040] In another aspect of the instant invention, a plant made by the method of (1) identifying the trait to be modified; (2) constructing a recombinant DNA molecule consisting essentially of genetic elements isolated from the selected plant species, wherein the recombinant DNA molecule when integrated into the genome of the selected plant species modifies the expression of the trait in the transformed plant species; (3) stably integrating the recombinant DNA molecule into cells of the selected plant species through LifeSupport-mediated transformation; and (4) identifying transformed plants exhibiting modified expression of the trait, is provided.

[0041] In a further aspect, a method of modifying expression of a trait in a selected plant species is provided. This method comprises (1) identifying the trait to be modified; (2) constructing a recombinant DNA molecule consisting essentially of (a) genetic elements isolated from the selected plant species, wherein the genetic elements when integrated into the genome of the selected plant species modifies the expression of the trait in the transformed plant species; and (b) a selectable marker gene that is isolated from the same plant species; (3) stably integrating the recombinant DNA molecule into cells of the selected plant species through LifeSupport-mediated transformation; (4) detecting the selectable marker gene; and (5) identifying transformed plants exhibiting modified expression of the trait.

[0042] In yet one other aspect, a plant exhibiting a modified expression of a trait is provided. In one embodiment, the plant has stably integrated into its genome a recombinant DNA molecule consisting essentially of genetic elements isolated from a plant of the same species, or from a plant that is sexually compatible with that species, wherein the recombinant DNA molecule modifies the expression of the trait.

[0043] In another aspect of the present invention, an isolated nucleotide sequence referred to as "plant-DNA" ("P-DNA") is provided. In a preferred embodiment, the P-DNA itself lacks any genes or parts thereof and is delineated by terminal, T-DNA "border-like" sequences that share at least 50%, at least 75%, at least 90% or at least 95% sequence identity with the nucleotide sequence of the T-DNA borders of any virulent *Agrobacterium* strain, and which support an efficient transfer of the entire P-DNA from *Agrobacterium* to plant cells.

[0044] In a preferred embodiment a "border-like" sequence promotes and facilitates the integration of a polynucleotide to which it is linked. In another preferred embodiment, each terminal sequence of the modified P-DNA is

between 5-100 bp in length, 10-80 bp in length, 15-75 bp in length, 15-60 bp in length, 15-50 bp in length, 15-40 bp in length, 15-30 bp in length, 16-30 bp in length, 20-30 bp in length, 21-30 bp in length, 22-30 bp in length, 23-30 bp in length, 24-30 bp in length, 25-30 bp in length, or 26-30 bp in length. More preferably, the border-like sequence is between 20 and 28 nucleotides in length.

[0045] In a preferred embodiment, the P-DNA left and right border sequences of the present invention are isolated from and/or are native to the genome of a plant that is to be modified and are not identical in nucleotide sequence to any known *Agrobacterium*-derived T-DNA border sequence. Thus, in one embodiment, a P-DNA border sequence may possess 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more nucleotides that are different from a T-DNA border sequence from an *Agrobacterium* species, such as *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. Alternatively, in another embodiment, a P-DNA border, or a border-like sequence of the present invention has at least 95%, at least 90%, at least 80%, at least 75%, at least 70%, at least 60% or at least 50% sequence identity with a T-DNA border sequence from an *Agrobacterium* species, such as *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. More preferably, a native plant P-DNA border sequence that shares greater than or equal to 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, or 60% nucleotide sequence identity with an *Agrobacterium* T-DNA border sequence.

[0046] In another preferred embodiment, a border-like sequence can be isolated from a plant genome and then modified or mutated to change the efficiency by which they are capable of integrating a nucleotide sequence into another nucleotide sequence. In another embodiment, other polynucleotide sequences may be added to or incorporated within a border-like sequence of the present invention. Thus, in yet another embodiment, a P-DNA left border or a P-DNA right border may be modified so as to possess 5'- and 3'-multiple cloning sites, or additional restriction sites. In a further embodiment, a P-DNA border sequence may be modified to increase the likelihood that backbone DNA from the accompanying vector is not integrated into the plant genome.

[0047] In an even more preferred embodiment, the P-DNAs are isolated from any plant by using degenerate primers in a polymerase chain reaction. In one preferred embodiment, the P-DNA is derived from potato, is delineated by 25-bp termini with 80 and 88% identity to conventional T-DNA borders, respectively, and has the nucleotide sequence shown in SEQ ID NO. 1. In another most preferred embodiment, the P-DNA is derived from wheat, is delineated by 25-bp termini with 72% and 92% identity with conventional T-DNA borders, respectively, and contains the nucleotide sequence shown in SEQ ID NO. 34.

[0048] Such a P-DNA may be modified so as to comprise other polynucleotides positioned between the border-like sequences. In a preferred embodiment, the modified P-DNA consists essentially of, in the 5'- to 3'-direction, a first border-like sequence that promotes DNA transfer, a promoter, a desired polynucleotide that is operably linked to the

promoter, a terminator and a second border-like sequence that also promotes DNA transfer. In one other embodiment, the desired polynucleotide represents one or several copies of a leader, a trailer or a gene in sense and/or antisense orientations. In a more preferred embodiment, the modified P-DNA contains expression cassettes for both a mutant PPO gene and an invertase inhibitor gene.

[0049] Thus, in a preferred embodiment, the desired polynucleotide comprises a sense and antisense sequence of a leader sequence. In a more preferred embodiment, the leader sequence is associated with a gene that is endogenous to a cell of the selected plant species. In yet a more preferred embodiment, the leader is associated with a gene that is selected from the group consisting of a PPO gene, an R1 gene, a type L or H alpha glucan phosphorylase gene, an UDP glucose glucosyltransferase gene, a HOS1 gene, a S-adenosylhomocysteine hydrolase gene, a class II cinnamate 4-hydroxylase gene, a cinnamoyl-coenzyme A reductase gene, a cinnamoyl alcohol dehydrogenase gene, a caffeoyl coenzyme A O-methyltransferase gene, an actin depolymerizing factor gene, a Nin88 gene, a Lol p 5 gene, an allergen gene, a P450 hydroxylase gene, an ADP-glucose pyrophosphorylase gene, a proline dehydrogenase gene, an endo-1,4-beta-glucanase gene, a zeaxanthin epoxidase gene, and a 1-aminocyclopropane-1-carboxylate synthase gene.

[0050] In yet another preferred embodiment, the desired polynucleotide sequence comprises a sense and antisense sequence of a trailer sequence. In a preferred embodiment, the trailer sequence is associated with a gene selected from the group consisting of a PPO gene, an R1 gene, a type L or H alpha glucan phosphorylase gene, an UDP glucose glucosyltransferase gene, a HOS1 gene, a S-adenosylhomocysteine hydrolase gene, a class II cinnamate 4-hydroxylase gene, a cinnamoyl-coenzyme A reductase gene, a cinnamoyl alcohol dehydrogenase gene, a caffeoyl coenzyme A O-methyltransferase gene, an actin depolymerizing factor gene, a Nin88 gene, a Lol p 5 gene, an allergen gene, a P450 hydroxylase gene, an ADP-glucose pyrophosphorylase gene, a proline dehydrogenase gene, an endo-1,4-beta-glucanase gene, a zeaxanthin epoxidase gene, and a 1-aminocyclopropane-1-carboxylate synthase gene.

[0051] In a preferred embodiment, the desired polynucleotide, such as a gene, is isolated from, and/or is native to the plant that is to be transformed. In another preferred embodiment, the desired polynucleotide is modified or mutated. In one embodiment, a mutation to the isolated polynucleotide may render the desired nucleotide greater than or equal to 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, or 60% dissimilar to its unmutated form.

[0052] In a preferred embodiment of the present invention, the promoter of an expression cassette located within a P-DNA is a constitutive promoter. In a more preferred embodiment the constitutive promoter is the promoter of the Ubiquitin-3 gene of potato. In an even more preferred embodiment the constitutive promoter is the promoter of the Ubiquitin-7 gene of potato.

[0053] In another embodiment, the promoter of an expression cassette located within a P-DNA is a regulatable promoter. In a more preferred embodiment, the regulat-

able promoter is sensitive to temperature. In an even more preferred embodiment, the regulatable promoter is a ci21A promoter or a C17 promoter, each isolated from potato (Schneider et al., *Plant Physiol.* 113: 335-45, 1997; Kirch et al., *Plant Mol Biol* 33: 897-909, 1997).

[0054] In another embodiment, the promoter of an expression cassette located within a P-DNA can be regulated in a temporal fashion. In a preferred embodiment, the promoter is an rbcS promoter (Ueda et al., *Plant Cell* 1: 217-27, 1989).

[0055] In yet another embodiment, the promoter of an expression cassette located within a P-DNA is regulated by any one of abscisic acid, wounding, methyl jasmonate or gibberellic acid. In a further embodiment, this promoter is a promoter selected from either a Rab 16A gene promoter, an α -amylase gene promoter or a pin2 gene promoter.

[0056] In another embodiment, the promoter of an expression cassette located within a P-DNA is a tissue-specific promoter. In a particularly preferred embodiment, this promoter is a GBSS promoter isolated from *S. tuberosum*.

[0057] In one embodiment, the present invention provides a P-DNA vector that is capable of replication in both *E.coli* and *Agrobacterium*, and contains either a P-DNA or a modified P-DNA. In a preferred embodiment, this vector also contains an expression cassette for a cytokinin gene in its backbone to enable the selection against backbone integration events.

[0058] In another preferred embodiment, the desired nucleotide sequence further comprises a spacer element. In a more preferred embodiment, the spacer element is a Ubi intron sequence or a GBSS spacer sequence.

[0059] In another preferred embodiment, the desired nucleotide sequence comprises a mutated native gene encoding a functionally inactive protein, which reduces the overall activity of that protein if expressed in transgenic plants. In yet a more preferred embodiment, this mutated gene encodes a functionally inactive polyphenol oxidase lacking a copper binding domain.

[0060] In another preferred embodiment, the desired nucleotide sequence comprises a native gene encoding a functionally active protein. In yet a more preferred embodiment, this gene encodes for a protein with homology to the tobacco vacuolar invertase inhibitor.

[0061] In another embodiment, the terminator of an expression cassette located within a P-DNA is a Ubi3 terminator sequence or a 3'-untranslated region of a gene of a selected plant species.

[0062] In another aspect of the instant invention, a method for modifying a target plant cell is provided. In one embodiment, the method comprises: (1) inserting a modified P-DNA into the genome of at least one cell in the target plant cell using LifeSupport-mediated transformation; and (2) observing if there is a phenotypic change in the target plant cell; wherein the promoter in the modified P-DNA transcribes the sense and/or antisense untranslated sequences associated with a native gene to reduce expression of that native gene, thereby modifying the target plant cell. In another preferred embodiment, the promoter in the modified P-DNA transcribes a gene to overexpress that gene in the target plant cell.

[0063] In yet another aspect, there is provided a method of making a transgenic plant cell of a selected plant species that contains a modified P-DNA. The method comprises - co-transfecting a plant cell of the selected plant species with a P-DNA vector and a LifeSupport vector that comprises a marker gene flanked by a T-DNA left border and a T-DNA right border and a mutant *virD2* gene inserted into the vector backbone, and selecting for a plant cell that transiently expresses the marker gene, and isolating a plant cell that contains the modified P-DNA integrated into its genome but does not contain any nucleotides from the LifeSupport vector. In a preferred embodiment, the marker gene confers resistance to kanamycin. In a most preferred embodiment the yeast ADH terminator follows the kanamycin resistance gene.

[0064] In a preferred embodiment, the plant cell of the selected plant species targeted for transformation is in culture. In another preferred embodiment, the plant cell of the selected plant species targeted for transformation is within a plant.

[0065] The present invention also provides a plant of the selected species that comprises at least one cell with a genome that contains a modified P-DNA. In a preferred embodiment, the modified P-DNA consists essentially of, in the 5'- to 3'-direction, a first terminus that functions like a T-DNA border followed by P-DNA sequences, a promoter, a desired nucleotide sequence operably linked to both a promoter, a terminator and additional P-DNA sequences delineated by a second terminus. In another embodiment, the desired polynucleotide represents one or several copies of a leader, a trailer and a gene in the sense and/or antisense orientation.

[0066] In another embodiment, a plant that comprises at least one cell with a genome that contains a modified P-DNA is envisioned.

[0067] In another aspect of the invention, a method for reducing the expression of a gene in a selected plant species is provided. The method comprises the LifeSupport-mediated transformation of a plant cell from a selected plant species with a P-DNA vector, wherein the modified P-DNA of this vector is stably integrated into the genome of the plant cell. In another aspect of the invention, the modified P-DNA comprises a desired polynucleotide that reduces expression of an endogenous gene from the selected plant species.

[0068] In another aspect of the instant invention, a gene native to the selected plant species may be mutated and reintroduced into the plant using the inventive methods. Preferably, the mutated gene, for instance a mutated PPO gene, is integrated into the plant cell genome using a P-DNA vector.

[0069] The present invention also provides a method for reducing the undesirable expression of the polyphenol oxidase gene in a selected plant species. In a preferred embodiment, the method comprises integrating into a genome of a selected plant species a modified P-DNA comprised only of nucleotide sequences isolated from the selected plant species or from a plant that is sexually compatible with the selected plant species, consisting essentially of, in the 5'- to 3'-direction, a first P-DNA terminus that functions like a T-DNA border followed by flanking P-DNA sequences; a promoter;

a desired nucleotide which is a sense-oriented trailer nucleotide sequence associated with a specific PPO gene; an antisense-oriented sequence of the trailer nucleotide sequence from the specific PPO gene; a termination sequence, and additional P-DNA sequences delineated by a second terminus that functions like a T-DNA border, wherein the promoter produces a double-stranded RNA molecule that reduces the expression of the specific PPO gene, thereby reducing black spot bruising in specific tissues of the plant. In another embodiment, the sense- and antisense-oriented nucleotide sequences from the leader nucleotide sequences are obtained from the 5'-untranslated region preceding the specific PPO gene. In a further embodiment, the sense- and antisense-oriented leader or trailer sequence associated with the PPO gene may be separated by another polynucleotide sequence, referred to herein, as either an intron or a "spacer." In a preferred embodiment, the leader or trailer sequence is associated with a potato PPO gene. In a more preferred embodiment, the leader or trailer sequence is associated with a potato PPO gene that is expressed in potato tubers. In a most preferred embodiment, the leader or trailer sequence is associated with a potato PPO gene that is expressed in all parts of the potato tuber except for the epidermis.

[0070] The present invention also provides a method for reducing acrylamide production, sprout-induction during storage, phosphate accumulation, and/or cold-induced sweetening in tubers of a selected plant species.

[0071] In a preferred embodiment, the method comprises the LifeSupport-mediated transformation of a selected plant species with a modified P-DNA comprised only of nucleotide sequences isolated from the selected plant species, or from plants that are sexually compatible with the selected plant species, consisting essentially of, in the 5'- to 3'-direction, a first P-DNA with a left border-like sequence, a promoter, a desired nucleotide sequence, which is a sense-oriented nucleotide sequence from the leader sequence associated with the R1 gene, an antisense-oriented sequence from this leader sequence, a termination sequence, and a right border-like sequence. Upon expression, a leader-RNA duplex is produced that reduces expression of the R1 gene, thereby reducing cold-induced sweetening in the plant. In another embodiment, the desired sense- and antisense-oriented nucleotide sequences represent the trailer associated with the R1 gene. In a further embodiment, the sense- and antisense-oriented leader or trailer associated with R1 may be separated by another polynucleotide sequence, referred to herein, as either an intron or a "spacer."

[0072] In another preferred embodiment, the method comprises the LifeSupport-mediated transformation of a selected plant species with a modified P-DNA that is similar to the one described above but contains a leader- or trailer sequence associated with an alpha glucan phosphorylase gene.

[0073] In yet another preferred embodiment, the method comprises the LifeSupport-mediated transformation of a selected plant species with a modified P-DNA that contains an invertase inhibitor gene.

[0074] In another preferred embodiment, the modified P-DNA described in the preceding paragraphs are used to reduce the accumulation of additional undesirable products of the Maillard reaction, which occurs during the heating of

carbohydrate-rich foods such as potato tubers. These undesirable products include advanced glycation end products (AGEs) that have been associated with various pathologies.

[0075] The present invention also provides a method for increasing resistant starch levels in the storage organs of plants and food crops.

[0076] In a preferred embodiment, the method comprises the LifeSupport-mediated transformation of a selected plant species with a modified P-DNA that contains an expression cassette for a fusion of the trailer sequences associated with the starch branching enzyme I and II genes.

[0077] The present invention also provides isolated nucleotide sequences comprising the promoters of the potato GBSS gene and the potato proteinase inhibitor gene, which are predominantly expressed in tubers. The isolated promoters have the nucleotide sequence shown in SEQ ID NO.: 6 and SEQ ID NO.:40, respectively.

[0078] In one aspect, the present invention provides a method of modifying a trait of a selected plant comprising:

[0079] a. stably transforming cells from the selected plant with a desired polynucleotide, wherein the desired polynucleotide consists essentially of a nucleic acid sequence that is native to the selected plant, native to a plant from the same species, or is native to a plant that is sexually interfertile with the selected plant,

[0080] b. obtaining a stably transformed plant from the transformed plant cells wherein the transformed plant contains the desired polynucleotide stably integrated into the genome and wherein the desired polynucleotide modifies the trait.

[0081] In a preferred embodiment, the method further comprises co-transfecting the plant cells with a selectable marker gene that is transiently expressed in the plant cells, and identifying transformed plant cells, and transformed plants obtained from the transformed plant cells, wherein the selectable marker gene is not stably integrated and the desired polynucleotide is stably integrated into the genome.

[0082] In a preferred embodiment, the desired polynucleotide comprises a P-DNA, GBSS promoter, Ubi7 promoter, Ubi3 promoter, PIP promoter, modified PPO gene, invertase inhibitor gene, salt tolerance gene, R1-associated leader, phosphorylase-associated leader, R1-associated trailer, SBE-associated trailers, Ubi-intron, GBSS spacer, UbiT.

[0083] In another preferred embodiment, a "plant" of the present invention is a monocotyledenous plant, selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.

[0084] In yet another embodiment, a "plant" of the present invention is a dicotyledenous plant, selected from the group consisting of avocado, potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsettia, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

[0085] In yet another embodiment, plants and plant cells of the present inventive methods are transformed via *Agrobacterium*-mediated transformation. Preferably, the *Agrobacterium*-mediated transformation relies on the use of at

least one binary vector. In yet another embodiment, the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector. In a preferred embodiment the first binary vector contains the desired polynucleotide and the second binary vector contains a selectable marker gene, wherein the selectable marker gene is operably linked to a promoter and a terminator.

[0086] According to the present methods, the trait that is modified is selected from the group consisting of enhanced health and nutritional characteristics, improved storage, enhanced yield, enhanced salt tolerance, enhanced heavy metal tolerance, increased drought tolerance, increased disease tolerance, increased insect tolerance, increased water-stress tolerance, enhanced cold and frost tolerance, enhanced color, enhanced sweetness, improved vigor, improved taste, improved texture, decreased phosphate content, increased germination, increased micronutrient uptake, improved starch composition, improved flower longevity.

[0087] The present invention also encompasses a plant made by the present methods.

[0088] In another aspect, the present invention provides a method of modifying a trait in a selected plant comprising:

[0089] (a) identifying the trait to be modified;

[0090] (b) constructing a first polynucleotide consisting essentially of native genetic elements isolated from the selected plant, a plant from the same species, or a plant that is sexually interfertile with the selected plant, wherein the native genetic elements are capable of modifying the expression of a gene that controls the trait

[0091] (c) constructing a second polynucleotide comprising a selectable marker gene that is operably linked to a promoter and a terminator;

[0092] (d) co-transfecting plant cells from the selected plant with the first and second polynucleotides;

[0093] (e) selecting for the transient expression of the selectable marker gene;

[0094] (f) screening for plant cells stably transformed with the first polynucleotide but do not contain the second DNA molecule integrated into the genome; and

[0095] (g) obtaining a stably transformed plant from the transformed plant cells that exhibit a modified expression of the trait.

[0096] In one embodiment, the genetic elements comprise at least one of a promoter, sequence of interest, terminator enhancer, intron, spacer, or regulatory elements. In another embodiment, method of claim 4, wherein the plant cells are transfected with the first polynucleotide before the second polynucleotide or vice versa.

[0097] In one embodiment, the sequence of interest is a gene. In another embodiment, the gene is a mutated or wild-type polyphenol oxidase gene or a mutated or wild-type R1 gene. In one other embodiment, the sequence of interest is a leader or trailer sequence, wherein the leader or trailer sequence represents a sequence upstream or downstream of a gene that is native to the plant cell. In yet another

embodiment, the sequence of interest comprises a sense-oriented leader sequence operably linked to an antisense leader sequence. In another embodiment, the sequence of interest comprises a sense-oriented trailer sequence operably linked to an antisense trailer sequence. In another embodiment, the promoter is an inducible promoter. In another embodiment, the terminator is a yeast ADH terminator sequence.

[0098] According to the present invention, a leader construct comprises in 5'-to 3'-direction, a promoter, a sense-oriented leader sequence, the antisense sequence of the leader, and a terminator, wherein expression of the leader construct produces a double-stranded RNA molecule that facilitates the down-regulation of expression of the gene to which it is associated. In one other embodiment, the leader sequence is associated with, and located upstream of, the coding region of the PPO gene, the R1 gene, an L-type phosphorylase gene, or an alpha glucan phosphorylase gene.

[0099] In another embodiment, the trailer construct comprises in 5'-to 3'-direction, a promoter, a sense-oriented trailer sequence, the antisense sequence of the trailer, and a terminator, wherein expression of the trailer construct produces a double-stranded RNA molecule that facilitates the down-regulation of expression of the gene to which it is associated. In a preferred embodiment, the trailer sequence is associated with, and located downstream of, the coding region of the PPO gene, the R1 gene, an L-type phosphorylase gene, or an alpha glucan phosphorylase gene.

[0100] The method further comprises exposing the plant cell to a second vector that comprises a marker element, wherein the marker is transiently expressed in the transformed plant and is not stably integrated into the genome of the transformed plant. In one embodiment, the marker is a herbicide resistance gene, an antibiotic resistance gene, or NPTII.

[0101] Preferably, the plant cells are transformed via *Agrobacterium*-mediated transformation. In one embodiment, the *Agrobacterium*-mediated transformation relies on the use of at least one binary vector. In yet another embodiment, the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector. In one other embodiment, the first binary vector carries the first polynucleotide and the second binary vector carries the second polynucleotide.

[0102] The present invention provides another method of modifying the expression of a gene in a selected plant comprising:

[0103] (a) identifying the functional gene;

[0104] (b) constructing a first polynucleotide consisting essentially of native genetic elements isolated from the selected plant, a plant of the same species as the selected plant, or a plant that is sexually interfertile with the selected plant, wherein the native genetic elements are capable of modifying the expression of the gene;

[0105] (c) constructing a second polynucleotide comprising a functional selectable marker gene;

[0106] (d) co-transfecting plant cells from the selected plant with the first and second polynucleotides;

[0107] (e) selecting for the transient expression of the selectable marker gene;

[0108] (f) screening for plant cells stably transformed with the first polynucleotide but do not contain the second polynucleotide integrated into the genome; and

[0109] (g) obtaining a transformed plant from the transformed plant cells that exhibit modified expression of the gene.

[0110] Preferably, the plant cells are transformed via *Agrobacterium*-mediated transformation. In one embodiment, the *Agrobacterium*-mediated transformation relies on the use of at least one binary vector. In yet another embodiment, the

[0111] *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector. In one other embodiment, the first binary vector carries the first polynucleotide and the second binary vector carries the second polynucleotide.

[0112] In another embodiment, the first polynucleotide comprises at least one of a P-DNA, GBSS promoter, Ubi7 promoter, Ubi3 promoter, PIP promoter, modified PPO gene, invertase inhibitor gene, salt tolerance gene, R1-associated leader, phosphorylase-associated leader, R1-associated trailer, SBE-associated trailers, Ubi-intron, GBSS spacer, UbiT.

[0113] In another embodiment, the second polynucleotide comprises at least one of a selectable marker gene, an omega-mutated virD2 polynucleotide, a codA polynucleotide, and a codA::upp fusion polynucleotide.

[0114] The present invention also encompasses a plant made by such method.

[0115] In one other embodiment, a transgenic plant is provided which exhibits a modified expression of a trait compared to the non-transgenic plant from which it was derived, wherein the transgenic plant is stably transformed with a desired polynucleotide consisting essentially of native genetic elements isolated from the plant, a plant in the same species, or a plant that is sexually interfertile with the plant, and wherein the polynucleotide modifies the expression of the trait.

[0116] In another preferred embodiment, the "plant" of the present invention is a monocotyledonous plant, selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.

[0117] In yet another embodiment, the "plant" of the present invention is a dicotyledonous plant, selected from the group consisting of avocado, potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsettia, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

[0118] In another embodiment, the trait is selected from the group consisting of enhanced health and nutritional characteristics, improved storage, enhanced yield, enhanced salt tolerance, enhanced heavy metal tolerance, increased drought tolerance, increased disease tolerance, increased insect tolerance, increased water-stress tolerance, enhanced cold and frost tolerance, enhanced color, enhanced sweet-

ness, improved vigor, improved taste, improved texture, decreased phosphate content, increased germination, increased micronutrient uptake, improved starch composition, improved flower longevity.

[0119] In another embodiment, the desired polynucleotide comprises at least one of a P-DNA, GBSS promoter, Ubi7 promoter, Ubi3 promoter, PIP promoter, modified PPO gene, invertase inhibitor gene, salt tolerance gene, R1-associated leader, phosphorylase-associated leader, R1-associated trailer, SBE-associated trailers, Ubi-intron, GBSS spacer, UbiT.

[0120] The present invention also encompasses an isolated, border-like nucleotide sequence ranging in size from 20 to 100 bp that shares between 52% and 96% sequence identity with a T-DNA border sequence from *Agrobacterium tumefaciens*. In a preferred embodiment, the isolated nucleotide sequence is isolated from a monocotyledonous plant, selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm. In another embodiment, the nucleotide sequence is isolated from a dicotyledonous plant selected from the group consisting of potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsettia, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

[0121] In yet another embodiment, the isolated nucleotide sequence is isolated from potato, and has a nucleotide sequence shown in either SEQ ID NO. 94 or 95. In a preferred embodiment, the isolated nucleotide sequence shares 52% sequence identity with a T-DNA border sequence from *Agrobacterium tumefaciens*. The present invention encompasses a vector that comprises such nucleotide sequences.

[0122] The present invention also provides method of making a plant stably transformed with a desired polynucleotide comprising:

[0123] (a) isolating a P-DNA that is flanked by border-like sequences from the plant wherein the border-like sequences share between 52% and 96% sequence identity with an *Agrobacterium tumefaciens* T-DNA border sequence;

[0124] (b) inserting the desired polynucleotide between the P-DNA border-like sequences to form a P-DNA construct ; and

[0125] (c) transforming a plant cell from the plant with the P-DNA construct; and

[0126] (d) recovering a plant from the transformed plant cell stably transformed with the P-DNA construct.

[0127] In one embodiment, the P-DNA construct is carried on a vector comprised of a backbone integration marker gene and transformed plant cells are selected that do not contain the backbone integration marker gene. In another embodiment, the backbone integration marker gene is a cytokinin gene. In another embodiment, plant shoots are not selected that exhibit a cytokinin-overproducing phenotype. In yet another embodiment, the backbone integration marker gene is the IPT gene, and plant shoots are not selected that exhibit an abnormal phenotype or cannot develop roots.

[0128] In one other embodiment, the plant cells are from a monocotyledonous plant selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.

[0129] In another embodiment, the plant cells are from a dicotyledonous plant selected from the group consisting of potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsettia, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

[0130] Preferably, the plant cells are transformed via *Agrobacterium*-mediated transformation. In one embodiment, the *Agrobacterium*-mediated transformation relies on the use of at least one binary vector. In yet another embodiment, the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector. In one other embodiment, the first binary vector carries the first polynucleotide and the second binary vector carries the second polynucleotide. In one further embodiment, the second binary vector comprises at least one of a negative selectable marker gene and an omega-mutated virD2 gene, wherein the negative selectable marker gene is positioned within the right T-DNA border and the left T-DNA border, and wherein the omega-mutated virD2 gene is positioned within the backbone of the second binary vector. In a preferred embodiment, the second binary vector comprises both a negative selectable marker gene positioned within the right T-DNA border and the left T-DNA border, and an omega-mutated virD2 gene positioned within the backbone of the second binary vector.

[0131] The present invention also provides a P-DNA consisting essentially of, in the 5'- to 3'-direction, a first T-DNA border-like sequence, a promoter, a desired polynucleotide sequence operably linked to the promoter, a terminator, and a second T-DNA border-like sequence, wherein the border-like sequences have less than 100% sequence identity with T-DNA border sequences

[0132] In a preferred embodiment, the T-DNA border-like sequences, the promoter, the desired polynucleotide, and the terminator, are all isolated from the same plant, the same plant species, or plants that are sexually interfertile.

[0133] In another embodiment, the P-DNA further consists essentially of a selectable marker gene.

[0134] In yet another embodiment, the T-DNA border-like sequences, the promoter, the desired polynucleotide, the terminator and the selectable marker gene, are all isolated from the same plant, the same plant species, or plants that are sexually interfertile.

[0135] In yet another embodiment, the desired polynucleotide sequence in the P-DNA is a sequence upstream or downstream of the coding region of a gene, wherein the upstream sequence is a leader sequence, and wherein the downstream sequence is a trailer sequence. In this embodiment, the T-DNA border-like sequences, the promoter, the leader sequence, the trailer sequence, the terminator and the selectable marker gene are all isolated from the same plant, the same plant species, or plants that are sexually interfertile.

[0136] In another embodiment, vectors comprising such P-DNA constructs are provided by the present invention.

[0137] In another embodiment, the promoter is a regulatable promoter. In yet another embodiment, the regulatable promoter is sensitive to temperature. In a preferred embodiment, the regulatable promoter is a wheat wcs120 promoter. In another embodiment, the promoter is under temporal regulation. In yet another embodiment, the promoter is a carboxylase promoter. In a further embodiment, the carboxylase promoter is a maize carboxylase promoter.

[0138] The promoter may be regulated by any one of abscisic acid, wounding, methyl jasmonate or gibberellic acid. In another embodiment, the promoter is a promoter selected from either a Rab 16A gene promoter, an α -amylase gene promoter or a pin2 gene promoter. In yet another embodiment, the promoter is a tissue-specific promoter.

[0139] In one other embodiment, the leader sequence is a part of a 5'-untranslated region of a gene that is endogenous to a cell of the selected plant species. In another embodiment, the 5'-untranslated region is upstream of a start codon of a gene that is selected from the group consisting of a PPO gene, an R1 gene, a HOS 1 gene, a S-adenosylhomocysteine hydrolase gene, a class II cinnamate 4-hydroxylase gene, a cinnamoyl-coenzyme A reductase gene, a cinnamoyl alcohol dehydrogenase gene, a caffeoyl coenzyme A O-methyltransferase gene, an actin depolymerizing factor gene, a Nin88 gene, a Lol p 5 gene, an allergen gene, a P450 hydroxylase gene, an ADP-glucose pyrophosphorylase gene, a pr6 line dehydrogenase gene, an endo-1,4-beta-glucanase gene, a zeaxanthin epoxidase gene, and a 1-aminocyclopropane-1-carboxylate synthase gene.

[0140] In another embodiment, the trailer sequence is a part of the 3'-untranslated region of a gene that is downstream of a termination codon of a gene selected from the group consisting of a PPO gene, an R1 gene, a HOS1 gene, a S-adenosylhomocysteine hydrolase gene, a class II cinnamate 4-hydroxylase gene, a cinnamoyl-coenzyme A reductase gene, a cinnamoyl alcohol dehydrogenase gene, a caffeoyl coenzyme A O-methyltransferase gene, an actin depolymerizing factor gene, a Nin88 gene, a Lol p 5 gene, an allergen gene, a P450 hydroxylase gene, an ADP-glucose pyrophosphorylase gene, a proline dehydrogenase gene, an endo-1,4-beta-glucanase gene, a zeaxanthin epoxidase gene, and a 1-aminocyclopropane-1-carboxylate synthase gene.

[0141] The present vector may further comprise a spacer element that is either an Ubi intron sequence or a GBSS spacer sequence. In another embodiment, the vector comprises a terminator that is a Ubi3 terminator sequence or a 3'-untranslated region of an endogenous plant gene.

[0142] In another embodiment, the vector comprises a selectable marker gene operably linked to a constitutive promoter and a Cre gene operably linked to an inducible promoter, wherein the selectable marker gene and the Cre gene are flanked by a first recombinase recognition site and a second recombinase recognition site. In another embodiment, the first recombinase recognition site and the second recombinase recognition site are lox sites.

[0143] In another embodiment, the inducible promoter is a temperature-sensitive promoter, a chemically-induced promoter, or a temporal promoter. In yet another embodiment, the inducible promoter is a Ha hsp17.7 G4 promoter, a wheat wcs120 promoter, a Rab 16A gene promoter, an α -amylase gene promoter, a pin2 gene promoter, a carboxylase pro-

motor. In yet another preferred embodiment, further comprises a plant-derived marker gene. In another preferred embodiment, the plant-derived marker gene is an enolpyruvul-3-phosphoshikimic acid synthase gene.

[0144] In another aspect of the present invention, a method for modifying a plant cell is provided, comprising integrating a P-DNA sequence into the genome of a plant cell, wherein the P-DNA consists essentially of, in the 5'- to 3'-direction, a first T-DNA border-like sequence, a promoter, a desired polynucleotide sequence operably linked to the promoter, a terminator, and a second T-DNA border-like sequence, wherein the border-like sequences have less than 100% sequence identity with T-DNA border sequences, and wherein the T-DNA border-like sequences, the promoter, the desired polynucleotide, and terminator, are all isolated from or native to the genome of the plant cell, wherein the desired polynucleotide comprises sense and antisense sequences of a leader sequence or trailer sequence that are associated with the upstream or downstream non-coding regions of a gene in the plant, and wherein expression of the desired polynucleotide produces a double-stranded RNA transcript that targets the gene associated with the desired polynucleotide, thereby modifying the plant cell.

[0145] The present invention also encompasses a method for modifying a plant, comprising:

[0146] (i) transfecting at least one cell in the plant with the vector of the present invention;

[0147] (ii) selecting a cell expressing the functional selectable marker;

[0148] (iii) isolating the cell expressing the functional selectable marker;

[0149] (iii) inducing the expression of the functional Cre gene in the isolated cell;

[0150] (iv) culturing the isolated cell; and

[0151] (ii) observing the phenotype of cultured cells;

[0152] wherein a phenotype that is different to an untransfected plant cell indicates that the target plant cell has been modified.

[0153] In a preferred embodiment the selecting step of this and other methods of the present invention is performed by identifying which cells are resistant to an antibiotic.

[0154] In another aspect, a method for identifying a target plant cell whose genome contains a P-DNA, comprises co-transfecting a plant target cell with the vector of the present invention and a second *Agrobacterium*-derived vector that comprises a marker gene flanked by a T-DNA left border and a T-DNA right border and a omega-mutated virD2 gene, wherein the P-DNA is integrated into the genome of the plant target cell, and wherein no part of the second *Agrobacterium*-derived vector is integrated into the genome of the plant target cell. In a preferred embodiment, the marker in the second *Agrobacterium*-derived vector is a neomycin phosphotransferase gene.

[0155] In another aspect, the method for identifying a target plant cell whose genome contains at least a part of an integration cassette is provided, further comprises selecting cells that survive temporary growth on a kanamycin-containing media, wherein the genomes of the selected cells

contain only the integration cassette. In one embodiment, the target plant cell is within a plant. A plant comprising at least one cell whose genome comprises such a P-DNA is also encompassed by the present invention.

[0156] The present invention also encompasses a plant comprising at least one cell whose genome is artificially manipulated to contain only plant-derived nucleic acids, wherein no cells of the plant contain foreign nucleic acids integrated into the cell genome.

[0157] The present invention also encompasses a polynucleotide comprising the polynucleotide sequence of SEQ ID NO. 93, wherein the polynucleotide is between 20 and 80 nucleotides in length. In one embodiment, the polynucleotide is between 21 and 70 nucleotides in length, between 22 and 50 nucleotides in length, between 23 and 40 nucleotides in length, or between 24 and 30 nucleotides in length.

[0158] In another aspect, the invention encompasses a tuber-specific promoter as shown in SEQ ID NO. 40.

[0159] The present invention also encompasses an *Agrobacterium*-based method of making transgenic plant cells that do not contain a selectable marker gene stably integrated in nuclear DNA comprising:

[0160] a. constructing a first binary vector comprised of a polynucleotide consisting essentially of a desired functional gene operably linked to T-DNA borders or T-DNA border-like sequences at the 5' and 3' ends of the desired functional gene;

[0161] b. constructing a second binary vector comprised of a functional selectable marker gene operably linked to T-DNA borders or T-DNA border-like sequences at the 5' and 3' ends of the functional selectable marker gene;

[0162] c. incubating plants cells with:

[0163] i. an *Agrobacterium* strain carrying the first and the second binary vectors; or

[0164] ii. a first *Agrobacterium* strain carrying the first binary vector and a second *Agrobacterium* strain carrying the second binary vector;

[0165] d. selecting plant cells wherein the desired functional gene is integrated into plant nuclear DNA without integration of the selectable marker gene into plant nuclear DNA following incubation for an appropriate time period on a medium containing an appropriate selection agent.

[0166] In a preferred embodiment, the selectable marker gene is a herbicide resistance gene or an antibiotic resistance gene. In another preferred embodiment, the antibiotic resistance gene is the nNPTII gene. In another embodiment, the antibiotic resistance gene is the npt II structural gene operably linked to the promoter from the Ubiquitin-7 gene and the terminator from yeast alcohol dehydrogenase 1 (ADH1) gene. According to this method, the plant cells are first incubated with the first *Agrobacterium* strain and then subsequently incubated with the second *Agrobacterium* strain or -vice versa.

[0167] In a preferred embodiment, the first binary vector further comprises a binary integration marker gene that can be used to detect plant cells stably transformed with binary

vector backbone sequences. In another embodiment, the binary vector integration marker gene is selected from the group consisting of herbicide resistance gene, antibiotic resistance gene, or NPTII. In yet another embodiment, the second binary vector further comprises a gene fusion between the bacterial cytosine deaminase (coda) and uracil phosphoribosyltransferase (upp) genes, which is inserted between the T-DNA or T-DNA border-like sequences, and plant cells are exposed to 5-fluorocytosine following incubation with the first and second *Agrobacterium* strains in order to select against those plant cells transformed with the second binary vector.

[0168] In yet another embodiment, the secondary binary vector further comprises a gene that reduces the probability of backbone integration. In one embodiment, such a gene is the omega-mutated virD2 gene, wherein the omega-mutated virD2 gene reduces the frequency of integration of the selectable marker gene into the plant nuclear DNA.

[0169] The present invention also encompasses an isolated nucleotide sequence comprising the GBSS promoter isolated from *S. tuberosum*. In a preferred embodiment, this isolated nucleotide sequence has the nucleotide sequence that is SEQ ID. NO. 6 or 13.

BRIEF DESCRIPTION OF THE DRAWINGS

[0170] FIG. 1. Schematic illustrations of some P-DNA vectors used in the present invention. P-DNA region is indicated as grey box. "ipt"=expression cassette for the ipt gene; "npt"=expression cassette for the nptII gene; "mPPO"=expression cassette for a modified PPO gene; "INH"=expression cassette for an invertase inhibitor gene; "GUS"=expression cassette for the GUS gene; "LPPO"=expression cassette for a sense and antisense copy of the leader associated with a PPO gene; "LPH"=expression cassette for a sense and antisense copy of the leader associated with a phosphorylase gene; "Alf"=expression cassette for a potato Alfin homolog. See text for details.

[0171] FIG. 2. Gene-free expression cassettes

[0172] FIG. 3. Alignment of potato and tobacco invertase inhibitor proteins. "St"=*Solanum tuberosum* (potato); "Nt"=*Nicotiana tabacum* (tobacco)

[0173] FIG. 4. Alignment of trailers associated with various PPO genes.

[0174] FIG. 5. Schematic illustrations of some LifeSupport vectors used in the present invention. "codA" is an expression cassette for the codA gene; "codA::upp" is an expression cassette for the codA gene fused to upp; "QvirD2" is an expression cassette for the QvirD2 gene.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0175] The "precise breeding" strategy of the present invention improves the agronomic performance, nutritional value, and health characteristics of plants and crops without introducing unknown nucleic acid, or nucleic acid from a foreign species into a plant species genome, and without producing undesirable phenotypes or harmful side-effects.

[0176] Thus, the present invention provides a transgenic plant, and methods for making such a plant that do not integrate nucleic acid from non-plant species into that

plant's genome. Nucleic acids, promoters, regulatory elements, other non-coding gene sequences, markers, polynucleotides, and genes that are integrated into the selected plant genome are all preferably isolated from the plant that is to be transformed, plants of the same species to be transformed, or plants that are sexually interfertile with the plant to be transformed. Such "native" nucleic acids can be mutated, modified or cojoined with other native nucleic acids in an expression cassette and reintegrated into the selected plant genome, according to the methods described herein. Accordingly, the genotype and phenotype of the transgenic plant is altered using only that selected plant's own nucleic acid, or using nucleic acid from a plant that is sexually compatible with the selected plant.

[0177] To facilitate the production of such transgenic plants, the present invention makes use of the fact that not all T-DNA vectors used in *Agrobacterium*-mediated transformation are actually integrated into the plant genome; i.e., while a vector may be taken up by the plant cell, an actual integration event may not occur. According to the present invention, one may use such a vector to carry a selectable marker gene into a plant cell. Plant cells can then be screened to determine whether the marker has been stably integrated into the plant genome by determining for how long the marker gene is expressed. Accordingly, plant cells that only transiently express the selectable marker gene are desired because they represent cells that took up, but did not integrate into their genomes, the selectable marker gene.

[0178] Thus, by co-transforming a plant with such a "marker vector" and also with another vector that contains the desired native gene or polynucleotide, one can select plant cells that took up both vectors and, from those, determine which cells possess genomes that contain only the desired gene or polynucleotide. The "marker vector" can be modified to further reduce the possibility that the marker will be integrated into the plant genome. The present invention provides such "marker vectors" in the form of "Life-Support" vectors.

[0179] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Generally, the nomenclature used herein, and the laboratory procedures in cell culture, molecular genetics, and nucleic acid chemistry and hybridization described herein, are those well known and commonly employed in the art. Standard techniques are used for recombinant nucleic acid methods, polynucleotide synthesis, microbial culture, cell culture, tissue culture, transformation, transfection, transduction, analytical chemistry, organic synthetic chemistry, chemical syntheses, chemical analysis, and pharmaceutical formulation and delivery. Generally, enzymatic reactions and purification and/or isolation steps are performed according to the manufacturers' specifications. The techniques and procedures are generally performed according to conventional methodology disclosed, for example, in *Molecular cloning a laboratory manual*, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), and *Current protocols in molecular biology*, John Wiley & So Baltimore, Md. (1989).

[0180] Amino acid sequence: as used herein, includes an oligopeptide, peptide, polypeptide, or protein and fragments thereof, that are isolated from, native to, or naturally occurring in a plant, or are synthetically made but comprise the nucleic acid sequence of the endogenous counterpart.

[0181] Artificially manipulated: as used herein, "artificially manipulated" means to move, arrange, operate or control by the hands or by mechanical means or recombinant means, such as by genetic engineering techniques, a plant or plant cell, so as to produce a plant or plant cell that has a different biological, biochemical, morphological, or physiological phenotype and/or genotype in comparison to unmanipulated, naturally-occurring counterpart.

[0182] Asexual propagation: producing progeny by generating an entire plant from leaf cuttings, stem cuttings, root cuttings, tuber eyes, stolons, single plant cells protoplasts, callus and the like, that does not involve fusion of gametes.

[0183] Backbone: nucleic acid sequence of a binary vector that excludes the T-DNA or P-DNA sequence intended for transfer.

[0184] Border and Border-like sequences: "border sequences" are specific *Agrobacterium*-derived sequences. Typically, a left border sequence and a right border sequence flank a T-DNA and they both function as recognition sites for virD2-catalyzed nicking reactions. Such activity releases nucleic acid that is positioned between such borders. See Table 2 below for examples of border sequences. The released nucleic acid, complexed with virD2 and virE2, is targeted to plant cell nuclei where the nucleic acid is often integrated into the genome of the plant cell. Usually, two border sequences, a left-border and a right-border, are used to integrate a nucleotide sequence that is located between them into another nucleotide sequence. It is also possible to use only one border, or more than two borders, to accomplish integration of a desired nucleic acid in such fashion.

[0185] According to the present invention, a "border-like" sequence is isolated from the selected plant species that is to be modified, or from a plant that is sexually-compatible with the plant species to be modified, and functions like the border sequences of *Agrobacterium*. That is, a border-like sequence of the present invention promotes and facilitates the integration of a polynucleotide to which it is linked. A plant-DNA, i.e., P-DNA, of the present invention preferably contains border-like sequences.

[0186] A border-like sequence of a P-DNA is between 5-100 bp in length, 10-80 bp in length, 15-75 bp in length, 15-60 bp in length, 15-50 bp in length, 15-40 bp in length, 15-30 bp in length, 16-30 bp in length, 20-30 bp in length, 21-30 bp in length, 22-30 bp in length, 23-30 bp in length, 24-30 bp in length, 25-30 bp in length, or 26-30 bp in length.

[0187] The border-like sequences of the present invention can be isolated from any plant, such as from potato and wheat. See SEQ ID NO. 1 and SEQ ID NO. 34, for sequences which contain, at either end, the border-like sequences isolated from potato and wheat respectively. Thus, a P-DNA left and right border sequences of use for the present invention are isolated from and/or native to the genome of a plant that is to be modified. A P-DNA border-like sequence is not identical in nucleotide sequence to any known *Agrobacterium*-derived T-DNA border sequence.. Thus, a P-DNA border-like sequence may possess 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more nucleotides that are different from a T-DNA border sequence from an *Agrobacterium* species, such as *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. That is, a P-DNA border, or a border-like sequence of the present invention has at least 95%, at least 90%, at least 80%, at least 75%, at least 70%, at least 60% or at least 50% sequence identity

with a T-DNA border sequence from an *Agrobacterium* species, such as *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*, but not 100% sequence identity. As used herein, the descriptive terms "P-DNA border" and "P-DNA border-like" are exchangeable.

[0188] A native P-DNA border sequence is greater than or equal to 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, 60%, 59%, 58%, 57%, 56%, 55%, 54%, 53%, 52%, 51% or 50% similar in nucleotide sequence to a *Agrobacterium* T-DNA border sequence. A border-like sequence can, therefore, be isolated from a plant genome and be modified or mutated to change the efficiency by which they are capable of integrating a nucleotide sequence into another nucleotide sequence. Other polynucleotide sequences may be added to or incorporated within a border-like sequence of the present invention. Thus, a P-DNA left border or a P-DNA right border may be modified so as to possess 5'- and 3'-multiple cloning sites, or additional restriction sites. A P-DNA border sequence may

be modified to increase the likelihood that backbone DNA from the accompanying vector is not integrated into the plant genome.

[0189] Table 2 below depicts the sequences of known T-DNA border sequences and sequences identified herein as border-like sequences. None of the sequences identified as "border-like" in Table 2 have been identified previously as having a T-DNA border-like structure. The potato border-like sequences were isolated by the present inventive methods using degenerate primers in polymerase chain reactions from potato genomic DNA. The present invention encompasses the use of any P-DNA border-like sequence for transferring a cojoined polynucleotide into the genome of a plant cell.

[0190] Indeed, the present invention encompasses any border-like sequence that has the nucleic acid sequence structure of SEQ ID NO. 93: ANGATNTATN6GT (SEQ ID NO. 93), where "N" is any nucleotide, such as those represented by "A," "G," "C," or "T." This sequence represents the consensus sequence of border-like nucleic acids identified by the present invention.

TABLE 2

"Border" and "Border-Like" sequences		
<u>Agrobacterium T-DNA borders</u>		
TGACAGGATATATTGGCGGGTAAAC (SEQ ID NO.41)	Agrobacterium nopaline strains (RB)	
TGGCAGGATATATTGTGGTGTAAC (SEQ ID NO.42)	Agrobacterium nopaline strains (LB)	
TGGCAGGATATATACCGTTGTAATT (SEQ ID NO.43)	Agrobacterium octopine strains (RB)	
CGGCAGGATATATTCAATTGTAATT (SEQ ID NO.44)	Agrobacterium octopine strains (LB)	
TGGTAGGATATATACCGTTGTAATT (SEQ ID NO.45)	LB mutant	
TGGCAGGATATATGGTACTGTAATT (SEQ ID NO.46)	LB mutant	
YGRYAGGATATATWSNVBKGTAAWY (SEQ ID NO.47)	Border motif	
<u>Border-like sequences</u>		
CGGCAGGATATATCCTGATGTAAAT (SEQ ID NO.48)	R. leguminosarum	
TGGCAGGAGTTATTCGAGGGTAAAC (SEQ ID NO.49)	T. tengcongensis	
TGACAGGATATATCGTGATGTCAAC (SEQ ID NO.50)	Arabidopsis thaliana	
GGGAAGTACATATTGGCGGGTAAAC (SEQ ID NO.51)	A. thaliana CHR1v07142002	
TTACAGGATATATTAATATGTATGA (SEQ ID NO.52)	Oryza sativa AC078894	
TAACATGATATATTCCTTGTAAAT (SEQ ID NO.53)	Homo sapiens clone HQ0089	
TGACAGGATATATGGTAATGTAAAC (SEQ ID NO.54)	potato (left border sequence)*	
TGGCAGGATATATACCGATGTAAAC (SEQ ID NO.55)	potato (right border sequence)*	

Y = C or T; R = A or G; K = G or T; M = A or C; W = A or T; S = C or G; V = A, C, or G; B = C, G, or T.

The accession numbers for the border-like sequences are: *Oryza sativa* chromosome 10 BAC OSJNBa0096G08 genomic sequence (AC078894.11); *Arabidopsis thaliana* chromosome 3 (NM_114337.1); *Arabidopsis thaliana* chromosome 1 (NM_105664.1); *T. tengcongensis* strain MB4T, section 118 of 244 of the complete genome (AE013091.1); Homo sapiens clone HQ0089 (AF090888.1); *Rhizobium* Clone: rhiz98e12.glk.

*potato left and right border sequences were obtained and isolated according to the presently-described inventive methods.

[0191] Carrier DNA: a “carrier DNA” is a DNA segment that is used to carry certain genetic elements and deliver them into a plant cell. In conventional foreign DNA transfer, this carrier DNA is often the T-DNA of *Agrobacterium*, delineated by border sequences. The carrier DNA described here is obtained from the selected plant species to be modified and contains ends that may be structurally and functionally different from T-DNA borders but shares with such T-DNAs the ability to support both DNA transfer from *Agrobacterium* to the nuclei of plant cells or certain other eukaryotes and the subsequent integration of this DNA into the genomes of such eukaryotes.

[0192] Consisting essentially of: a composition “consisting essentially of” certain elements is limited to the inclusion of those elements, as well as to those elements that do not materially affect the basic and novel characteristics of the inventive composition. Thus, so long as the composition does not affect the basic and novel characteristics of the instant invention, that is, does not contain foreign DNA that is not from the selected plant species or a plant that is sexually compatible with the selected plant species, then that composition may be considered a component of an inventive composition that is characterized by “consisting essentially of” language.

[0193] Degenerate primer: a “degenerate primer” is an oligonucleotide that contains sufficient nucleotide variations that it can accommodate base mismatches when hybridized to sequences of similar, but not exact, homology.

[0194] Dicotyledon (dicot): a flowering plant whose embryos have two seed leaves or cotyledons. Examples of dicots include, but are not limited to, tobacco, tomato, potato, sweet potato, cassava, legumes including alfalfa and soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

[0195] Regulatory sequences: refers to those sequences which are standard and known to those in the art, that may be included in the expression vectors to increase and/or maximize transcription of a gene of interest or translation of the resulting RNA in a plant system. These include, but are not limited to, promoters, peptide export signal sequences, introns, polyadenylation, and transcription termination sites. Methods of modifying nucleic acid constructs to increase expression levels in plants are also generally known in the art (see, e.g. Rogers et al., 260 *J. Biol. Chem.* 3731-38, 1985; Cornejo et al., 23 *Plant Mol. Biol.* 567: 81, 1993). In engineering a plant system to affect the rate of transcription of a protein, various factors known in the art, including regulatory sequences such as positively or negatively acting sequences, enhancers and silencers, as well as chromatin structure may have an impact. The present invention provides that at least one of these factors may be utilized in engineering plants to express a protein of interest. The regulatory sequences of the present invention are native genetic elements, i.e., are isolated from the selected plant species to be modified.

[0196] Foreign: “foreign,” with respect to a nucleic acid, means that that nucleic acid is derived from non-plant organisms, or derived from a plant that is not the same species as the plant to be transformed or is not derived from a plant that is not interfertile with the plant to be transformed, does not belong to the species of the target plant.

[0197] According to the present invention, foreign DNA or RNA represents nucleic acids that are naturally occurring

in the genetic makeup of fungi, bacteria, viruses, mammals, fish or birds, but are not naturally occurring in the plant that is to be transformed. Thus, a foreign nucleic acid is one that encodes, for instance, a polypeptide that is not naturally produced by the transformed plant. A foreign nucleic acid does not have to encode a protein product. According to the present invention, a desired transgenic plant is one that does not contain any foreign nucleic acids integrated into its genome.

[0198] Native genetic elements, on the other hand, can be incorporated and integrated into a selected plant species genome according to the present invention. Native genetic elements are isolated from plants that belong to the selected plant species or from plants that are sexually compatible with the selected plant species. For instance, native DNA incorporated into cultivated potato (*Solanum tuberosum*) can be derived from any genotype of *S. tuberosum* or any genotype of a wild potato species that is sexually compatible with *S. tuberosum* (e.g., *S. demissum*).

[0199] Gene: “gene” refers to the coding region and does not include nucleotide sequences that are 5'- or 3'- to that region. A functional gene is the coding region operably linked to a promoter or terminator.

[0200] Genetic rearrangement: refers to the reassociation of genetic elements that can occur spontaneously in vivo as well as in vitro which introduce a new organization of genetic material. For instance, the splicing together of polynucleotides at different chromosomal loci, can occur spontaneously in vivo during both plant development and sexual recombination. Accordingly, recombination of genetic elements by non-natural genetic modification techniques in vitro is akin to recombination events that also can occur through sexual recombination in vivo.

[0201] In frame: nucleotide triplets (codons) are translated into a nascent amino acid sequence of the desired recombinant protein in a plant cell. Specifically, the present invention contemplates a first nucleic acid linked in reading frame to a second nucleic acid, wherein the first nucleotide sequence is a gene and the second nucleotide is a promoter or similar regulatory element.

[0202] Integrate: refers to the insertion of a nucleic acid sequence from a selected plant species, or from a plant that is from the same species as the selected plant, or from a plant that is sexually compatible with the selected plant species, into the genome of a cell of a selected plant species. “Integration” refers to the incorporation of only native genetic elements into a plant cell genome. In order to integrate a native genetic element, such as by homologous recombination, the present invention may “use” non-native DNA as a step in such a process. Thus, the present invention distinguishes between the “use of” a particular DNA molecule and the “integration” of a particular DNA molecule into a plant cell genome.

[0203] Introduction: as used herein, refers to the insertion of a nucleic acid sequence into a cell, by methods including infection, transfection, transformation or transduction.

[0204] Isolated: “isolated” refers to any nucleic acid or compound that is physically separated from its normal, native environment. The isolated material may be maintained in a suitable solution containing, for instance, a solvent, a buffer, an ion, or other component, and may be in purified, or unpurified, form.

[0205] Leader: Transcribed but not translated sequence preceding (or 5' to) a gene.

[0206] LifeSupport Vector: a LifeSupport vector is a construct that contains an expressible selectable marker gene, such as a neomycin phosphotransferase marker, that is positioned between T-DNA or T-DNA-like borders. The LifeSupport vector may be modified to limit integration of such a marker, as well as other polynucleotides, that are situated between the border or border-like sequences, into a plant genome. For instance, a LifeSupport vector may comprise a mutated *virD2*, *codA::upp* fusion, or any combination of such genetic elements. Thus, a modified *virD2* protein will still support T-DNA transfer to plant nuclei but will limit the efficiency of a subsequent genomic integration of T-DNAs (Shurvinton et al., *Proc Natl Acad Sci USA*, 89: 11837-11841, 1992; Mysore et al., *Mol Plant Microbe Interact*, 11: 668-683, 1998). Alternatively, *codA::upp* gene fusion can be used as negative selectable marker prior to regeneration. In one preferred construct, the LifeSupport vector comprises the *npt* marker operably linked to the yeast ADH terminator element.

[0207] Monocotyledon (monocot): a flowering plant whose embryos have one cotyledon or seed leaf. Examples of monocots include, but are not limited to turf grass, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, and palm.

[0208] Native: a "native" genetic element refers to a nucleic acid that naturally exists in, originates from, or belongs to the genome of a plant that is to be transformed. Thus, any nucleic acid, gene, polynucleotide, DNA, RNA, mRNA, or cDNA molecule that is isolated either from the genome of a plant or plant species that is to be transformed or is isolated from a plant or species that is sexually compatible or interfertile with the plant species that is to be transformed, is "native" to, i.e., indigenous to, the plant species. In other words, a native genetic element represents all genetic material that is accessible to plant breeders for the improvement of plants through classical plant breeding. Any variants of a native nucleic acid also are considered "native" in accordance with the present invention. In this respect, a "native" nucleic acid may also be isolated from a plant or sexually compatible species thereof and modified or mutated so that the resultant variant is greater than or equal to 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, or 60% similar in nucleotide sequence to the unmodified, native nucleic acid isolated from a plant. A native nucleic acid variant may also be less than about 60%, less than about 55%, or less than about 50% similar in nucleotide sequence.

[0209] A "native" nucleic acid isolated from a plant may also encode a variant of the naturally occurring protein product transcribed and translated from that nucleic acid. Thus, a native nucleic acid may encode a protein that is greater than or equal to 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, or 60% similar in amino acid sequence to the unmodified, native protein expressed in the plant from which the nucleic acid was isolated.

[0210] Naturally occurring nucleic acid: this phrase means that the nucleic acid is found within the genome of a selected plant species and may be a DNA molecule or an RNA molecule. The sequence of a restriction site that is normally present in the genome of a plant species can be engineered into an exogenous DNA molecule, such as a vector or oligonucleotide, even though that restriction site was not physically isolated from that genome. Thus, the present invention permits the synthetic creation of a nucleotide sequence, such as a restriction enzyme recognition sequence, so long as that sequence is naturally occurring in the genome of the selected plant species or in a plant that is sexually compatible with the selected plant species that is to be transformed.

[0211] Operably linked: combining two or more molecules in such a fashion that in combination they function properly in a plant cell. For instance, a promoter is operably linked to a structural gene when the promoter controls transcription of the structural gene.

[0212] P-DNA: according to the present invention, P-DNA ("plant-DNA") is isolated from a plant genome and comprises at each end, or at only one end, a T-DNA border-like sequence. The border-like sequence preferably shares at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 90% or at least 95%, but less than 100% sequence identity, with a T-DNA border sequence from an *Agrobacterium* species, such as *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. Thus, P-DNAs can be used instead of T-DNAs to transfer a nucleotide sequence from *Agrobacterium* to another polynucleotide sequence. The P-DNA may be modified to facilitate cloning and should preferably not naturally encode proteins or parts of proteins. The P-DNA is characterized in that it contains, at each end, at least one border sequence, referred to as either a "P-DNA border sequence" or "P-DNA border-like sequence," which are interexchangeable terms. See the definition of a "border sequence" and "border-like" above. A P-DNA may also be regarded as a "T-DNA-like" sequence, see definition below.

[0213] Plant: includes angiosperms and gymnosperms such as potato, tomato, tobacco, alfalfa, lettuce, carrot, strawberry, sugarbeet, cassava, sweet potato, soybean, maize, turf grass, wheat, rice, barley, sorghum, oat, oak, eucalyptus, walnut, and palm. Thus, a plant may be a monocot or a dicot. The word "plant," as used herein, also encompasses plant cells, seed, plant progeny, propagule whether generated sexually or asexually, and descendants of any of these, such as cuttings or seed. Plant cells include suspension cultures, callus, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, seeds and microspores. Plants may be at various stages of maturity and may be grown in liquid or solid culture, or in soil or suitable media in pots, greenhouses or fields. Expression of an introduced leader, trailer or gene sequences in plants may be transient or permanent. A "selected plant species" may be, but is not limited to, a species of any one of these "plants."

[0214] Precise breeding: refers to the improvement of plants by stable introduction of nucleic acids, such as native genes and regulatory elements isolated from the selected plant species, or from another plant in the same species as the selected plant, or from species that are sexually compatible with the selected plant species, into individual plant

cells, and subsequent regeneration of these genetically modified plant cells into whole plants. Since no unknown or foreign nucleic acid is permanently incorporated into the plant genome, the inventive technology makes use of the same genetic material that is also accessible through conventional plant breeding.

[0215] Plant species: the group of plants belonging to various officially named plant species that display at least some sexual compatibility.

[0216] Plant transformation and cell culture: broadly refers to the process by which plant cells are genetically modified and transferred to an appropriate plant culture medium for maintenance, further growth, and/or further development.

[0217] Recombinant: Was used herein, broadly describes various technologies whereby genes can be cloned, DNA can be sequenced, and protein products can be produced. As used herein, the term also describes proteins that have been produced following the transfer of genes into the cells of plant host systems.

[0218] Selectable marker: a “selectable marker” is typically a gene that codes for a protein that confers some kind of resistance to an antibiotic, herbicide or toxic compound, and is used to identify transformation events. Examples of selectable markers include the streptomycin phosphotransferase (spt) gene encoding streptomycin resistance, the phosphomannose isomerase (pmi) gene that converts mannose-6-phosphate into fructose-6 phosphate; the neomycin phosphotransferase (nptII) gene encoding kanamycin and geneticin resistance, the hygromycin phosphotransferase (hpt or aphiv) gene encoding resistance to hygromycin, acetolactate synthase (als) genes encoding resistance to sulfonylurea-type herbicides, genes coding for resistance to herbicides which act to inhibit the action of glutamine synthase such as phosphinothricin or basta (e.g., the bar gene), or other similar genes known in the art.

[0219] Sense suppression: reduction in expression of an endogenous gene by expression of one or more an additional copies of all or part of that gene in transgenic plants.

[0220] T-DNA-Like: a “T-DNA-like” sequence is a nucleic acid that is isolated from a selected plant species, or from a plant that is sexually compatible with the selected plant species, and which shares at least 75%, 80%, 85%, 90%, or 95%, but not 100%, sequence identity with *Agrobacterium* species T-DNA. The T-DNA-like sequence may contain one or more border or border-like sequences that are each capable of integrating a nucleotide sequence into another polynucleotide. A “P-DNA,” as used herein, is an example of a T-DNA-like sequence.

[0221] Trailer: Transcribed but not translated sequence following (or 3'to) a gene.

[0222] Transcribed DNA: DNA comprising both a gene and the untranslated leader and trailer sequence that are associated with that gene, which is transcribed as a single mRNA by the action of the preceding promoter.

[0223] Transcription and translation terminators: the expression vectors of the present invention typically have a transcription termination region at the opposite end from the transcription initiation regulatory region. The transcription termination region may be selected, for stability of the

mRNA to enhance expression and/or for the addition of polyadenylation tails added to the gene transcription product (Alber & Kawasaki, *Mol. & Appl. Genetics* 4: 19-34, 1982). Illustrative transcription termination regions include the E9 sequence of the pea RBCS gene (Mogen et al., *Mol. Cell Biol.*, 12: 5406-14, 1992) and the termination signals of various ubiquitin genes.

[0224] Transformation of plant cells: a process by which DNA is stably integrated into the genome of a plant cell. “Stably” refers to the permanent, or non-transient retention and/or expression of a polynucleotide in and by a cell genome. Thus, a stably integrated polynucleotide is one that is a fixture within a transformed cell genome and can be replicated and propagated through successive progeny of the cell or resultant transformed plant. Transformation may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of nucleic acid sequences into a prokaryotic or eukaryotic host cell, including *Agrobacterium*-mediated transformation protocols, viral infection, whiskers, electroporation, heat shock, lipofection, polyethylene glycol treatment, micro-injection, and particle bombardment.

[0225] Transgene: a gene that will be inserted into a host genome, comprising a protein coding region. In the context of the instant invention, the elements comprising the transgene are isolated from the host genome.

[0226] Transgenic plant: a genetically modified plant which contains at least one transgene.

[0227] Using/Use of: The present invention envisions the use of nucleic acid from species other than that of the selected plant species to be transformed to facilitate the integration of native genetic elements into a selected plant genome, so long as such foreign nucleic acid is not stably integrated into the same host plant genome. For instance, the plasmid, vector or cloning construct into which native genetic elements are cloned, positioned or manipulated may be derived from a species different to that from which the native genetic elements were derived.

[0228] Variant: a “variant,” as used herein, is understood to mean a nucleotide or amino acid sequence that deviates from the standard, or given, nucleotide or amino acid sequence of a particular gene or protein. The terms, “isoform,” “isotype,” and “analog” also refer to “variant” forms of a nucleotide or an amino acid sequence. An amino acid sequence that is altered by the addition, removal or substitution of one or more amino acids, or a change in nucleotide sequence, may be considered a “variant” sequence. The variant may have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. A variant may have “nonconservative” changes, e.g., replacement of a glycine with a tryptophan. Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted may be found using computer programs well known in the art such as Vector NTI Suite (InforMax, MD) software.

[0229] It is understood that the present invention is not limited to the particular methodology, protocols, vectors, and reagents, etc., described herein, as these may vary. It is

also to be understood that the terminology used herein is used for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention. It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to “a gene” is a reference to one or more genes and includes equivalents thereof known to those skilled in the art and so forth. Indeed, one skilled in the art can use the methods described herein to express any native gene (known presently or subsequently) in plant host systems.

[0230] P-DNA Vectors

[0231] *Agrobacterium*-mediated transformation methods are the preferred means of incorporating recombinant DNA into plant cells. According to the present invention, a binary vector was developed to produce genetically modified potato plants that contain only native potato nucleic acids. Such a vector is different from conventional, *Agrobacterium*-mediated, transformation vectors in three ways: (1) instead of an *Agrobacterium*-derived T-DNA sequence delineated by T-DNA borders, the present vector contains a native plant DNA (P-DNA) fragment that is flanked by border-like sequences, which support P-DNA transfer from *Agrobacterium* to plant cells although they are structurally and functionally different from T-DNA borders, (2) the backbone of the present vector may contain a marker that, if integrated into the plant cell's genome, prevents these cells from developing into mature plants, and (3) the present vector does not contain a foreign selectable marker gene between P-DNA termini.

[0232] The present invention demonstrates, surprisingly, that P-DNA fragments flanked by border-like sequences support DNA transfer from *Agrobacterium* into plant cells. P-DNA can be isolated from the genome of any plant by using primers that are designed on the basis of homology between the termini of a potato P-DNA and conventional T-DNA borders. Such fragments can then be tested and, if efficacious, used to transform that plant with native DNA exclusively. It is also possible to search plant genomic databases for DNA fragments with regions that show homology with T-DNA borders by using programs such as ‘blastn’ (Altschul et al., *J Mol Biol* 215: 403-10, 1990). The identified P-DNAs may then be modified to increase their utility. For instance, internal fragments of the isolated P-DNAs may be deleted and restriction sites may be added to facilitate cloning. It may also be efficacious to introduce point mutations at the terminal sequences to render the P-DNA more effective in transferring DNA.

[0233] Any gene expression cassette can be inserted between P-DNA border-like sequences. For potato transformations, such an expression cassette could consist of a potato promoter, operably linked to a potato gene and/or a leader or trailer sequence associated with that gene, and followed by a potato terminator. The expression cassette may contain additional potato genetic elements such as a signal peptide sequence fused in frame to the 5'-end of the gene, and a potato intron that could, for instance, be placed between promoter and gene-of-interest to enhance expression. For transformation of wheat with a modified P-DNA, all genetic elements that are inserted on the wheat P-DNA, including the P-DNA itself would be derived from wheat or plant species that are sexually compatible with wheat.

[0234] Another way to isolate P-DNAs is by generating a library of *Agrobacterium* strains that contain random plant DNA fragments instead of a T-DNA flanking a selectable marker gene. Explants infected with this library can be placed on proliferation medium that contains an appropriate selectable agent to identify P-DNAs that support the transfer of the marker gene from the vector in *Agrobacterium* to the plant cell.

[0235] It is possible that not just the native modified P-DNA, but also additional plasmid sequences are co-transferred from *Agrobacterium* to the plant cell during the transformation process. For the purposes of the present invention, this is an undesirable process because such plasmid “backbone” sequences represent non-plant, foreign DNA, such as bacterial DNA. The present invention prevents transformed plant cells that contain backbone sequences from developing into mature plants. Thus, the present invention makes it possible to distinguish backbone-containing and backbone-free transformation events during the regenerated shoot phase.

[0236] The method to select or screen against backbone integration events relies on the presence of an expression cassette for a marker, such as the isopentenyl phosphotransferase (IPT) gene, in the vector backbone, outside of the P-DNA. Upon backbone integration, the accumulation of IPT-induced cytokinin will alter the shape of transformed shoots, and prevent these shoots to develop roots. Instead of the IPT gene, any other gene that alters the shape, texture or color of the transformed plant's leaves, roots, stem, height or some other morphological feature can be used to screen and/or select against backbone integration events. Such a gene is referred to herein as a “backbone integration marker.” Thus, the transformed plant that exhibits an altered morphological feature attributable to the expression of the backbone integration marker gene is known to, contain in its genome foreign DNA in addition to the desired P-DNA. Accordingly, plants that exhibit a phenotype associated with the backbone integration marker are not desired.

[0237] The present invention is not limited to the use of only an IPT gene as a backbone integration marker; other genes can be used in such fashion. For example, a backbone integration marker may be an *Agrobacterium* transzeatine synthase (TZS) gene (Krall et al., *FEBS Lett* 527: 315-8, 2002) or a recessive *Arabidopsis* gene *hod* (Catterou et al., *Plant J* 30: 273-87, 2002). This method can be more easily applied for use in the present invention than some methods that insert toxic genes in vector backbone sequences. See, for instance, EP 1 009,842.

[0238] By positioning a backbone integration marker gene, such as a functional cytokinin gene upstream or downstream of the P-DNA, it is straightforward to distinguish between transformation events. Transformed plants that exhibit an altered morphological feature are discarded because they contain non-native DNA sequences integrated into the genome.

[0239] Another strategy for identifying plants that are stably transformed with only native DNA, is to employ the polymerase chain reaction. By using primers that are specifically designed to detect backbone sequences, plants can be identified and discarded that contain foreign backbone sequences in addition to the P-DNA. Other primer sets can subsequently be used to confirm the intact transfer of the

P-DNA. Thus, by either using the expression of a gene to change a morphological feature of a plant, or by screening for stably integrated foreign DNA in a transformed plant, plants stably transformed with only native DNA sequences can be identified and selected.

[0240] Genetic elements from a particular host plant can be inserted into the P-DNA sequence of a binary vector capable of replication in both *E. coli* and *Agrobacterium*. Introduction of the resulting vectors into disarmed *Agrobacterium* strains such as LBA4404 can be accomplished through electroporation, triparental mating or heat-shock treatment of chemically competent cells. The new strains can then be used to transform individual plant cells through infection of whole plants or explants.

[0241] Genetic elements from a particular host plant can be inserted into the P-DNA sequence of a binary vector capable of replication in both *E. coli* and *Agrobacterium*. Introduction of the resulting vectors into *Agrobacterium* strains such as LBA4404 can be accomplished through electroporation, triparental mating or heat-shock treatment of chemically competent cells. The new strains can then be used to transform individual plant-cells through infection of whole plants or explants. LBA4404 contains the disarmed Ti-plasmid pAL4404, which carries the virulence functions and a streptomycin resistance gene.

[0242] LifeSupport Vectors

[0243] Although the stable integration of bacterial marker genes into the genomes of plant cells facilitates the identification of transformation events, such modifications of plant genomes are undesirable because marker genes represent foreign DNA. Use of a foreign marker gene can be avoided by developing new *Agrobacterium*-based transformation methods.

[0244] One preferred embodiment is a novel method that relies on the use of two *Agrobacterium* strains: one strain containing a binary vector with a selectable marker gene intended for transient expression in plant nuclei, and another strain carrying the P-DNA with the actual sequences of interest intended for stable integration in plant genome (see Example 7).

[0245] Upon co-infection with the *Agrobacterium* strains, some plant cells will receive both a T-DNA with the marker gene and a P-DNA with the sequences of interest. Instead of subsequently selecting for stable integration of the marker gene by subjecting the infected explants for a long period of time to the appropriate antibiotic, explants are only briefly exposed to the antibiotic. In this way, all plant cells that transiently express the marker gene will survive. Because T-DNAs will in most cases degrade due to endogenous nuclease activities rather than stably integrate into their host's genome, the majority of plant cells that survived the transient selection are shown here to develop into shoots lacking a marker gene. The present invention, furthermore, demonstrates that a significant proportion of these marker-free shoots contain stably integrated P-DNAs.

[0246] There are various tools to enhance the efficiency of marker-free transformation. First, the present invention demonstrates that this frequency can be increased by sequentially infecting-explants with two *Agrobacterium* strains carrying the T-DNA/marker and P-DNA/sequences-

of-interest, respectively. Explants are first infected with the P-DNA strain, and after about 4 to 6 hours with the T-DNA strain.

[0247] Second, the T-DNA strain can be modified to express an omega-mutated virD2 gene. The modified virD2 protein will still support T-DNA transfer to plant nuclei but limit the efficiency of a subsequent genomic integration of T-DNAs (Shurvinton et al., *Proc Natl Acad Sci USA*, 89: 11837-11841, 1992; Mysore et al., *Mol Plant Microbe Interact*, 11: 668-683, 1998). The most preferred method of expressing a modified virD2 gene is by inserting an omega-mutated virD2 gene driven by the virD promoter in the backbone of the T-DNA vector.

[0248] Third, stable T-DNA integration can be further impaired by inserting telomere sequences close to the left- and right-border sequences of the T-DNA (Chiurazzi & Signer, *Plant Mol. Biol.*, 26: 923-934, 1994).

[0249] Fourth, the size of the T-DNA region carrying the marker gene can be increased to enhance the frequency of T-DNAs and P-DNAs moving together into the plant cell nucleus, and to reduce the frequency of genomic integration of the T-DNA.

[0250] Fifth, the frequency of T-DNAs and P-DNAs moving together into the plant cell nucleus can also be enhanced by using a single *Agrobacterium* strain carrying two compatible binary vectors with the T-DNA and P-DNA, respectively. An example of is two compatible binary vectors are a pSIM 1301-derived vector and a pBI121-derived vector.

[0251] Because the transiently expressed marker gene will usually not integrate into the plant genome, it is not necessary that both this gene and its regulatory sequences represent native DNA. In fact, it may be advantageous to use foreign regulatory sequences to promote high levels of transient gene expression in infected plant cells. A surprising discovery of the present invention is that an expression cassette containing the GUS gene followed by the terminator of the yeast alcohol dehydrogenase 1 (ADH1) was transiently expressed at high levels in potato cells. A similar construct with the yeast CYC1 terminator, however, did not function adequately. It may also be possible to enhance transient expression levels by operably linking a marker gene to a non-native promoter. Examples of such promoters are, e.g., synthetic promoters such as glucocorticoid-inducible promoters (Mori et al., *Plant J.*, 27: 79-86, 2001; Bohner et al., *Mol. Gen. Genet.*, 264: 860-70 2001), and non-native promoters such as the 35S promoters of cauliflower mosaic virus and figwort mosaic virus, and fungal promoters.

[0252] As an alternative to the two-strain *Agrobacterium*-mediated transformation approach described above, plants may also be transformed with a single strain that contains a P-DNA with both a native marker gene and the actual sequences of interest. The present invention demonstrates that it is possible to use salt tolerance genes as native markers for transformation. Such salt tolerance genes include crop homologs of the *Arabidopsis* genes SOS1 (Shi et al., *Nat Biotechnol.* 2002), AtNHX1 (Apse et al., *Science*. 285: 1256-8, 1999), Avp1 (Gaxiola et al., *Proc Natl Acad Sci USA*. 98: 11444-9, 2001), and CBF3 Kasuga et al., *Nat Biotechnol.* 17: 287-91, 1999).

[0253] The rearrangements of genetic elements accomplished through the inventive Precise Breeding methodology

could also occur spontaneously through the process of genetic recombination. For instance, all plants contain elements that can transpose from one to another chromosomal location. By inserting into promoters or genes, such transposable elements can enhance, alter, and/or reduce gene expression. For instance, the AMu4 insertion of the maize Mutator element in the promoter of the transcriptional regulator gene P-wr causes stripy red pericarps. Insertion of the same element in the promoter of the leaf-specific MADS-box gene ZMM19 resulted in expression of this gene in the inflorescences of maize, causing a foliaceous elongation of the glumes and other changes in male and female inflorescences, resulting in the famous phenotype of pod corn. Because of its bizarre tassels and ears, pod corn was of religious significance for certain native American tribes. Many genes are also rearranged through other transposon-induced modifications such as inversions, deletions, additions, and ectopic recombinations (Bennetzen, *Plant Mol Biol* 42: 251-69, 2000). Furthermore, plant DNA rearrangements frequently occur through the process of intragenic recombination. For instance, by recombining genes involved in resistance against specific pathogens, plants are able to develop resistance genes with new specificities and, thus, co-evolve with their pathogens (Ellis et al., *Trends Plant Sci* 5: 373-9, 2000). Another example of intragenic recombination relates to how plants reproduce: plants transition from cross-fertilizing to self-fertilizing by recombining genes involved in self-incompatibility (Kusaba et al., *Plant Cell* 13: 627-43, 2001). Other processes that promote genome evolution include, for instance, chromosome breakage and interchromosomal recombination.

[0254] Enhancing Tie Nutritional Value of Plants and Food Crops

[0255] To modify negative traits such as acrylamide accumulation during processing, glycoalkaloid accumulation, accumulation of undesirable advanced glycation products, CIPC accumulation, low levels of resistant starch, bruise susceptibility, cold-induced sweetening, disease susceptibility, low yield and low quality in crop plants through precise breeding, at least one specific expression cassette is incorporated into a host genome. Three different methods are used to eliminate negative traits: (1) overexpression of genes that prevent the occurrence of negative traits, (2) overexpression of mutated versions of genes associated with negative traits in order to titrate out the wild-type gene products with non-functional proteins, and (3) silencing specific genes that are associated with a negative trait by expressing at least one copy of a leader or trailer fragment associated with that gene in the sense and/or antisense orientation.

[0256] One example of an endogenous gene that is associated with a negative trait in potato and can be modified in vitro so that it encodes a non-functional protein is the polyphenol oxidase (PPO) gene. Upon impact injury, the PPO gene product is released from the plastid into the cytoplasm (Koussevitzky et al., *J. Biol. Chem.*, 273: 27064-9, 1998), where it will mediate the oxidation of phenols to create a variety of phenoxyl radicals and quinoid derivatives, which are toxic and/or ultimately form undesirable polymers that leave dark discolorations, or "black spots" in the crop.

[0257] Overexpressing a mutant PPO gene that contains a non-functional copper-binding domain can lower the activity of all PPO genes that are mainly expressed in tubers and

associated organs such as sprouts. The mutations render the polyphenol oxidase protein inactive because it is unable to bind copper. The skilled artisan would know where to make point mutations that would, in this case, compromise the function of a gene product. The applicants identified the copper binding domain in potato PPO by aligning the potato PPO protein sequence with a sweet potato PPO protein sequence (Klabunde et al., *Nat Struct. Biol.*, 5:1084-90, 1998). Areas of conservation, particularly those containing conserved histidine residues in copper-binding sites, were targets for inactivating the transgene product. Because the almost complete absence of PPO activity in such organs may negatively impact the plant's ability to resist pathogens, the present invention also describes an improved method of only lowering a specific PPO gene that is predominantly expressed in all parts of the mature tuber except for the epidermis. Silencing of this specific PPO gene by using a trailer sequence associated with that gene does not reduce PPO expression in the tuber epidermis, the part of the tuber that is most directly exposed to pathogens attempting to infect.

[0258] Enzymatic browning induced by the PPO gene not only reduces the quality of potato tubers; it also negatively affects crop foods such as wheat, avocado, banana, lettuce, apple, and pears.

[0259] Other genes that are associated with negative traits and can be silenced by using the leader or trailer sequences associated with those genes include the potato R1 gene and L-type phosphorylase genes. Both genes are involved in the degradation of starch to reducing sugars, such as glucose and fructose, which upon heating participate in the Maillard reaction to produce toxic products such as acrylamide. The present invention demonstrates that a reduction of cold-induced sweetening by lowering R1 or phosphorylase activity leads to a reduction of both non-enzymatic browning and acrylamide accumulation during the frying process of potatoes.

[0260] The invention also demonstrates the utility of overexpressing certain native genes in genetically modified crops. Levels of Maillard-reaction products such as acrylamide were reduced significantly by lowering the conversion of sucrose to reducing sugars through overexpression of a newly isolated vacuolar invertase inhibitor gene in potato.

[0261] The present invention also predicts that potato tubers displaying either an increased level of invertase inhibitor expression or a reduced level of R1 or phosphorylase expression will not require the intensive treatment with chemical sprout inhibitors such as CIPC prior to storage because their lowered levels of reducing sugars will (1) delay sprouting, and (2) allow storage at lower temperatures, thus further delaying sprouting. The highly reduced CIPC-residue levels, or the absence thereof, further enhances the nutritional value of processed foods derived from plants containing certain modified P-DNAs described here.

[0262] Thus, French fries or chips derived from tubers that contain the modified P-DNA will contain strongly reduced CIPC residue levels, further boosting their nutritional value.

[0263] The effect of simultaneously downregulating the expression of the PPO and either R1 or phosphorylase genes in potato tubers is synergistic because reducing sugars are

not only required for non-enzymatic browning through the Maillard reaction but also for browning mediated by the PPO enzyme. Decreased levels of reducing sugars in transgenic potato tubers will, therefore, also limit PPO activity and black spot bruise susceptibility. Thus, PPO, R1, and phosphorylase genes, and/or the leader or trailer sequences that are associated with these genes, represent DNA segments of interest that can be isolated, modified and reintroduced back into the plant to down-regulate the expression of these genes.

[0264] Apart from developing bruise resistance and reduced cold-induced sweetening, there are many other traits that can be introduced through Precise Breeding without using foreign DNA. For instance, disease resistance genes can be isolated from wild potato species and inserted into the genomes of disease susceptible varieties.

[0265] The Environmental Benefits of Modified Plants and Crops

[0266] As described above, reduced levels of either R1 or phosphorylase result in a reduced phosphorylation of starch. This reduction in starch phosphorylation results in a 90% decrease in phosphate content of potato tubers (Vikso-Nielsen, *Biomacromolecules*, 2: 83643, 2001). This will result in a reduction in phosphate levels in wastewaters from potato processing plants, which are currently about 25-40 mg/L. Thus, the use of low-phosphate tubers will reduce the release of phosphates into the environment and help to protect important ecosystems. Furthermore, low-phosphate potatoes may require less phosphate fertilization for optimal growth and yield, which would support a more sustainable agriculture by delaying the depletion of available phosphate resources.

[0267] Enhancing the Agricultural Performance of Plants and Food Crops

[0268] Apart from reduced bruise susceptibility and reduced cold-sweetening, which are two important processing traits, the present invention also provides salt tolerance, an increasingly important input trait. Some of the modified P-DNA constructs described in the present invention contain a salt tolerance gene as native marker for transformation. Importantly, the utility of this gene is not limited to a screening step in the transformation procedure. Overexpression of the salt tolerance gene in potato plants reduces stress symptoms induced by high salinity soil levels, and will make it possible to grow new varieties containing a modified P-DNA on a growing percentage of agricultural lands that contain salinity levels exceeding the maximum 2 millimhos/cm electrical conductivity levels that are optimal for growing conventional varieties.

[0269] Using Regulatory Elements Isolated from a Selected Plant Species or from a Species Sexually Compatible with the Selected Plant Species

[0270] Once the leader, gene or trailer has been isolated from the plant species of interest, and optionally modified, it can be operably linked to a plant promoter or similar regulatory element for appropriate expression in plants. Regulatory elements such as these serve to express untranslated sequences associated with a gene of interest in specific tissues or at certain levels or at particular times.

[0271] Dependent on the strategy involved in modifying the trait, it may be necessary to limit silencing to a particular

region of the plant. The promoter normally driving the expression of the endogenous gene may not be suitable for tissue-specific expression. As described in the section above, stable integration of bacterial or viral regulatory components, such as the cauliflower mosaic virus 35S "super" promoter, can result in unpredictable and undesirable events. Thus, one aspect of the present invention uses promoters that are isolated from the selected host plant species.

[0272] In a preferred embodiment of the instant invention, for use in *S. tuberosum*, the leader or trailer sequences associated with R1, phosphorylase, and PPO genes are operably linked to the granule-bound starch synthase gene promoter (Rohde et al., *J Gen & Breed*, 44, 311-315, 1990). This promoter has been used frequently by others to drive gene expression and is particularly active in potato tubers (van der Steege et al., *Plant Mol Biol*, 20: 19-30, 1992; Beaujean et al., *Biotechnol. Bioeng*, 70: 9-16, 2000; Oxenboll et al., *Proc Natl Acad Sci USA*, 9: 7639-44, 2000). This promoter may also be used, in a preferred embodiment, for expression of the modified leader or trailer sequences of R1, phosphorylase, and PPO genes.

[0273] Alternatively, other potato promoters can be operably linked to sequences of interest from potato. Such promoters include the patatin gene promoter (Bevan et al., *Nucleic Acids Res*, 14: 4625-38, 1986), or a fragment thereof, that promotes expression in potato tubers, the potato UDP-glucose pyrophosphorylase gene promoter (U.S. Pat. No. 5,932,783) and the promoter of the ubiquitin gene (Garbarino et al., *Plant Physiol*, 109: 1371-8, 1995).

[0274] The transcription of leaders and/or trailers can also be regulated by using inducible promoters and regulatory regions that are operably linked in a construct to a polynucleotide of interest. Examples of inducible promoters include those that are sensitive to temperature, such as heat or cold shock promoters. For instance, the potato ci21A-, and C17-promoters are cold-inducible (Kirch et al., *Plant Mol. Biol*, 33: 897-909, 1997; Schneider et al., *Plant Physiol*, 113: 335-45, 1997).

[0275] Other inducible promoters may be used that are responsive to certain substrates like antibiotics, other chemical substances, or pH. For instance, abscisic acid and gibberellic acid are known to affect the intracellular pH of plant cells and in so doing, regulate the Rab 16A gene and the alpha-amylase 1/6-4 promoter (Heimovaara-Dijkstra et al., *Plant Mol Biol*, 4 815-20, 1995). Abscisic acid, wounding and methyl jasmonate are also known to induce the potato pin2 promoter (Lorberth et al., *Plant J*, 2: 477-86, 1992).

[0276] In another example, some nucleotide sequences are under temporal regulation and are activated to express a downstream sequence only during a certain developmental stage of the plant or during certain hours of the day. For instance, the potato promoter of the small subunit of ribulose-1,5-bisphosphate carboxylase (rbcS) gene can direct cell-specific, light-regulated expression (Fritz et al., *Proc Natl Acad Sci USA*, 88: 4458-62, 1991). The skilled artisan is well versed in these exemplary forms of inducible promoters and regulatory sequences.

[0277] The use of certain polyadenylation signals may also be useful in regulating expression, by varying the stability of the mRNA transcript. In particular, some polyadenylation signals when operably linked to the 3' end of a polynucleotide cause the mRNA transcript to become accessible to degradation.

[0278] Thus, it is possible to regulate expression of a gene by operably linking it with one or more of such promoters, regulatory sequences, 3' polyadenylation signals, 3' untranslated regions, signal peptides and the like. According to the instant invention, DNA sequences and regulatory elements such as those described herein, and which will ultimately be integrated into a plant genome, are obtained from DNA of the selected plant species to be modified by the Precise Breeding process of the present invention. That is, DNA sequences and regulatory elements that are derived, isolated and cloned from other species, such as from bacteria, viruses, microorganisms, mammals, birds, reptiles and sexually incompatible plant species are not integrated into the genome of the transformed plant. DNA foreign to the selected plant species genome may be used in the present invention to create a transformation construct, so long as that foreign DNA is not integrated into a plant genome.

[0279] Not only does the present invention provide a method for transforming a plant species by integrating DNA obtained from the selected plant species, or from a plant that is sexually-compatible with the selected plant species, it also provides a means by which the expression of that DNA can be regulated. Accordingly, it is possible to optimize the expression of a certain sequence, either by tissue-specific or some other strategy, as previously described.

[0280] Using 3' Terminator Sequences Isolated from a Selected Plant Species

[0281] In addition to regulatory elements that initiate transcription, the native expression cassette also requires elements that terminate transcription at the 3'-end from the transcription initiation regulatory region. The transcription termination region and the transcription initiation region may be obtained from the same gene or from different genes. The transcription termination region may be selected, particularly for stability of the mRNA to enhance expression.

[0282] This particular element, the so-called "3'-untranslated region" is important in transporting, stabilizing, localizing and terminating the gene transcript. In this respect, it is well known to those in the art, that the 3'-untranslated region can form certain hairpin loop. Accordingly, the present invention envisions the possibility of operably linking a 3' untranslated region to the 3' end of a cloned polynucleotide such that the resultant mRNA transcript may be exposed to factors which act upon sequences and structures conferred by the 3' untranslated region.

[0283] A 3' sequence of the ubiquitin gene can be subcloned from the plant species from which the promoter and transgene were isolated and inserted downstream from a transgene to ensure appropriate termination of transcription. Both exemplary transgenes can be fused to the terminator sequence of the potato Ubiquitin gene (Ubi3) regardless of which promoter is used to drive their expression.

EXAMPLES

Example 1

Cloning of P-DNAs

[0284] This example demonstrates that T-DNA borders are specific to *Agrobacterium*. It also shows that plants contain T-DNA border-like sequences, and it provides the sequence

of DNA fragments isolated from potato and wheat that are delineated by such border-like sequences.

[0285] Conventional transformation systems use *Agrobacterium*-derived T-DNAs as vehicles for the transfer of foreign DNA from *Agrobacterium* to plant cells (Schilperoort et al., U.S. Pat. No. 4,940,838, 1990). Although T-DNAs usually comprise several hundreds of basepairs, delineated by a left-border (LB) and right-border (RB) repeat, they can also merely consist of such borders. The T-DNA borders play an essential role in the DNA transfer process because they function as specific recognition sites for virD2-catalyzed nicking reaction. The released single stranded DNA, complexed with *Agrobacterium* virD2 and virE2, is transferred to plant cell nuclei where it often integrates successfully into the plant genome. All T-DNA borders that have been used for foreign DNA transfer are derived from nopaline and octopine strains of *Agrobacterium tumefaciens* and *A. rhizogenes* (Table 2). These borders and often some flanking *Agrobacterium* DNA are present in thousands of binary vectors including, for example, pPAM (AY027531), pJawoh1 (AF408413), pYL156 (AF406991), pINDEX (AF294982), pC1300 (AF294978), pBI121 (AF485783), pLH9000 (AF458478), pAC161 (AJ315956), BinHytOP (Z37515), pHELLSGATE (AJ311874), pBAR-35S (AJ251014), pGreen (AJ007829), pBIN19 (X77672), pCAMBIA (AF354046), pX6-GFP (AF330636), pER8 (AF309825), pBI101 (U12639), pSKI074 (AF218466), pAJ1 (AC138659), pAC161 (AJ315956), pSLJ8313 (Y18556), and pGV4939 (AY147202). Recently, two homologs of T-DNA borders were identified in the chrysopine-type Ti plasmid pTiChry5 (Palanichelvam et al., *Mol Plant Microbe Interact* 13: 1081-91, 2000). The left border homolog is identical to an inactive border homolog located in the middle of the T-DNA of pTi15955. The right border homolog is unusually divergent from the sequence of functional T-DNA borders. It is therefore unlikely that these homologs are functionally active in supporting DNA transfer from pTiChry5 to plant cells.

[0286] Development of a new method that makes it possible to transform plants with only native DNA requires, in the first place, a replacement of the T-DNA including LB and RB. Unfortunately, advanced BLAST searches of public databases including those maintained by The National Center For Biotechnology Information, The Institute for Genomic Research, and SANGER failed to identify any border sequences in plants. It was therefore necessary to consider plant DNA sequences that are similar but not identical to T-DNA borders, designated here as "border-like" (border-like). Examples of plant border-like sequences that were identified in public databases are shown in Table 2. The challenge in trying to replace T-DNA borders with border-like sequences is that border sequences are highly conserved (see Table 2). A large part of these sequences is also highly conserved in the nick regions of other bacterial DNA transfer systems such as that of IncP, PC194, and ϕ X174, indicating that these sequences are essential for conjugative-like DNA transfer (Waters et al., *Proc Natl Acad Sci* 88:1456-60, 1991). Because there are no reliable data on border sequence requirements, the entire border seems therefore important in the nicking process. A single study that attempted to address this issue by testing the efficacy of border mutants in supporting DNA transfer is unreliable because negative controls did not appear to function appropriately (van Haaren et al., *Plant Mol Biol* 13: 523-531,

1989). Furthermore, none of the results of this study were confirmed molecularly. Despite these concerns, two possibly effective border mutants are shown in Table 2 as well.

[0287] Based on the homology among border sequences, a T-DNA border motif was identified (Table 2). Although this motif comprises 13,824 variants, many of which may not function—or may be inadequate—in transferring DNA, it represents the broadest possible definition of what a T-DNA border sequence is or may be. This border motif was then used to search publicly available DNA databases for homologs using the “Motif Alignment and Search Tool” (Bailey and Gribskov, *Bioinformatics* 14: 48-54, 1998) and “advanced BLASTN” (“penalty for nucleotide mismatch”=-1; “expect”=10⁵; Altschul et al., *Nucleic Acids Res* 25: 3389-3402, 1997). Again, these searches did not identify any identical matches in organisms other than *Agrobacterium*.

[0288] To try and increase the chance of isolating a potato DNA fragment containing border-like sequences that correspond to the border motif, DNA was isolated from 100 genetically diverse accessions (the so-called “core collection,” provided by the US Potato Genebank, WI). This DNA was pooled and used as template for polymerase chain reactions using a variety of oligonucleotides designed to anneal to borders or border-like sequences. Amplified fragments were sequence analyzed, and the sequence was then confirmed using inverse PCR with nested primers. One of the potato DNA fragments that was of particular interest contains a novel sequence without any major open reading frames that is delineated by border-like sequences (Table 2). One of the border-like sequences of this fragment contains at least 5 mismatches with T-DNA borders; the other border-like sequence contains at least 2 mismatches. Although both sequences contain one mismatch with the border motif, they were tested for their ability to support DNA transfer. For that purpose, the fragment was first reduced in size to 0.4-kilo basepairs by carrying out an internal deletion (SEQ ID NO.: 1). The resulting fragment was designated “P-DNA” (plant DNA) to distinguish it from the *Agrobacterium*-derived T-DNA. A similar fragment was isolated from the genome of the potato variety Russet Ranger, but has not been used for any further experiments.

[0289] Based on the divergence between P-DNA and T-DNA borders, the elongase amplification system (Life Technologies) was used with the following degenerate primers to isolate a P-DNA from wheat: 5'-GTTTACANHNB-NATATATCCTGYCA -3' (Bor-F) (SEQ ID NO. 56), and 5'-TGRCAGGATATATNVNDNTGTAAAC -3' (Bor-R) (SEQ ID NO. 57). The resulting 825-bp fragment is shown in SEQ ID NO.: 2, and was used to replace the T-DNA of a conventional binary vector. The efficacy of this construct can be tested by inserting an expression cassette for the GUS gene between P-DNA termini, and infecting wheat with an *Agrobacterium* strain carrying the resulting vector.

Example 2

Tobacco Transformation with P-DNA Vectors

[0290] This Example demonstrates that, despite structural (sequence divergence) and functional (transformation frequencies) differences between P-DNA termini and T-DNA borders, a P-DNA can be used in a similar way as a T-DNA to transfer DNA from *Agrobacterium* to tobacco cells.

[0291] A T-DNA-free vector that can be maintained in both *E. coli* and *A. tumefaciens* was obtained by removing the entire T-DNA region of the conventional binary vector pCAMBIA1301 (Cambia, AU). This was accomplished by simultaneously ligating a 5.9 kb SacII—SphI fragment of pSIM1301 with 2 fragments amplified from pCAMBIA1301 using the oligonucleotides pairs: 5'-CCGCGGTGATCA-CAGGCAGCAAC-3' (SEQ ID NO. 58) and 5'-AAGCTTC-CAGCCAGCCAACAGCTCCCCGAC-3' (SEQ ID NO. 59), and 5'-AAGCTTGGCTACTAGTGC-GAGATCTCTAAGAGAAAAAGAGCGTTTA-3' (SEQ ID NO. 60), and 5'-GCATGCTCGAGATAGGTGACCA-CATACAAATGGACGAACGG-3' (SEQ ID NO. 61), respectively.

[0292] To make it possible to screen against backbone integration events, an expression cassette comprising the *Agrobacterium* isopentenyl transferase (IPT) gene driven by the Ubi3 promoter and followed by the Ubi3 terminator (SEQ ID NO.: 3) was inserted as 2.6 kbp SacII fragment into the backbone of the T-DNA-free vector described above, yielding pSIM100-OD-IPT. Transformed plant cells expressing the IPT gene are expected to accumulate cytokinins and grow into abnormal shoots that cannot develop roots.

[0293] The 0.4 kb P-DNA fragment described in Example 1 was inserted into pSIM100-OD-IPT to generate pSIM 111 (FIG. 1; SEQ ID NO.: 4).

[0294] To test whether pSIM111 can be used to obtain transformed plants carrying P-DNAs (including any sequences located between P-DNA termini) without the additional vector backbone, a neomycin phosphotransferase (NPTII) gene expression cassette was inserted into the P-DNA of pSIM111 to create pSIM108 (FIG. 1).

[0295] The efficacy of P-DNA termini in supporting DNA transfer was tested by comparing transformation frequencies between pSIM108 and a control vector that contained a modified P-DNA with conventional T-DNA borders. This control vector, designated pSIM109, was generated by amplification of the entire P-DNA containing the NPTII gene expression cassette with the oligonucleotide pairs: 5'-ACTAGTGTTTACCCGCCAATATATCCTGTCAGAG-3' (SEQ ID NO. 62), and 5'-AAGCTTTGGCAGGATATAT-TGTGGTGTAAACGAAG-3' (SEQ ID NO. 63). A second control vector that was used for these experiments is the conventional binary vector pBI121 (Genbank accession number AF485783), which contains the same NPTII expression cassette inserted on a regular T-DNA. The binary vectors were introduced into *Agrobacterium tumefaciens* LBA4404 cells as follows. Competent LB4404 cells (50 uL) were incubated for 5 minutes at 37° C. in the presence of 1 µg of vector DNA, frozen for about 15 seconds in liquid nitrogen (about -196° C.), and incubated again at 37° C. for 5 minutes. After adding 1 mL of liquid broth (LB), the treated cells were grown for 3 hours at 28° C. and plated on LB/agar containing streptomycin (100 mg/L) and kanamycin (100 mg/L). The vector DNAs were then isolated from overnight cultures of individual LBA4404 colonies and examined by restriction analysis to confirm the presence of intact plasmid DNA.

[0296] Test transformations of the model plant tobacco were carried out by growing a 10-fold dilution of overnight-grown LBA4404::pSIM108 cells for 5-6 hours, precipitating

the cells for 15 minutes at 2,800 RPM, washing them with MS liquid medium (Phytotechnology) supplemented with sucrose (3%, pH 5.7) and resuspending the cells in the same medium to an OD_{600nm} of 0.2. The suspension was then used to infect leaf explants of 4-week-old in vitro grown *Nicotiana tabacum* plants. Infected tobacco explants were incubated for 2 days on co-culture medium (1/10 MS salts, 3% sucrose, pH 5.7) containing 6 g/L agar at 25° C. in a Percival growth chamber (16 hrs light) and subsequently transferred to M401/agar medium containing timentine (150 mg/L) and kanamycin (100 mg/L). The number of calli per explant that developed within the next 4 weeks is shown in Table 3. Our data demonstrate that P-DNAs delineated by either native termini or conventional T-DNA borders are about 50% more effective in transforming tobacco than T-DNAs. The increased efficiency of P-DNA transfer may be due to either its different CG content or other unknown structural features of the P-DNA.

Example 3

Potato Transformation with P-DNA Vectors

[0297] This Example demonstrates that a P-DNA can be used in a similar way as a T-DNA to transfer DNA from *Agrobacterium* to potato cells.

[0298] Potato transformations were carried out by infecting stem explants of 4-week-old in vitro grown Russet Ranger plantlets with *Agrobacterium* strains according to the following procedure. Ten-fold dilutions of overnight-grown cultures were grown for 5-6 hours, precipitated for 15 minutes at 2,800 RPM, washed with MS liquid medium (Phytotechnology) supplemented with sucrose (3%, pH 5.7), and resuspended in the same medium to an OD_{600nm} of 0.2. The resuspended cells were then used to infect 0.4-0.6 mm internodal potato segments. Infected stems were incubated for 2 days on co-culture medium (1/10 MS salts, 3% sucrose, pH 5.7) containing 6 g/L agar at 22° C. in a Percival growth chamber (16 hrs light) and subsequently transferred to callus induction medium (CIM, MS medium supplemented with 3% sucrose 3, 2.5 mg/L of zeatin riboside, 0.1 mg/L of naphthalene acetic acid, and 6 g/L of agar) containing timentine (150 mg/L) and kanamycin (100 mg/L). After 1 month of culture on CIM, explants were transferred to shoot induction medium (SIM, MS medium supplemented with 3% sucrose, 2.5 mg/L of zeatin riboside, 0.3 mg/L of gibberelic acid GA3, and 6 g/L of agar) containing timentine and kanamycin (150 and 100 mg/L respectively). After 3-4 weeks, the number of explants developing transgenic calli and/or shooting was counted. As shown in tobacco, the number of stem explants infected with pSIM108 that showed calli was higher than those in control experiments with the conventional binary vector pBI121 (Table 3). Shoots that subsequently arose from these calli could be grouped into two different classes. The first class of shoots was phenotypically indistinguishable from control shoots transformed with LBA::pBI121. The second class of shoots displayed an EPT phenotype. Shoots of the latter class were stunted in growth, contained only very small leaves, displayed a light-green to yellow color, and were unable to root upon transfer to hormone-free media. To confirm that shoots with an IPT phenotype contained the IPT gene stably integrated in their genomes, all shoots were transferred to Magenta boxes containing MS medium supplemented with

3% sucrose and timentine 150 mg/L, allowed to grow for 3 to 4 additional weeks, and used to isolate DNA. This plant DNA served as template in PCR reactions with an oligonucleotide pair designed to anneal to the IPT gene: 5'-GTC CAA CTT GCA CAG GAA AGA C-3', and 5'-CAT GGA TGA AAT ACT CCT GAG C-3'. As shown in Table 4, the PCR experiment confirmed a strict correlation between IPT phenotype and presence of the IPT gene. The presence of backbone DNA was also examined in plants obtained from a transformation with pBI121. This was done by performing PCR reactions on DNA isolated from the transformation events with the 'pBI121 backbone primers': 5'-CGGTG-TAAGTGAAGTGCAGTTGCCATG-3' (SEQ ID NO. 64), and 5'-CATCGGCCTCACTCATGAGCAGATTG-3' (SEQ ID NO. 65). Amplification of a 0.7 kbp band is indicative for backbone integration. By comparing the data presented in Table 4, it can be concluded that backbone integration frequencies are similar for P-DNA vectors and T-DNA vectors.

[0299] A second PCR experiment was carried out to test whether IPT-free plants did not contain any other backbone sequences. Because the IPT expression cassette is positioned close to the left border-like sequences, the oligonucleotide pair for this experiment was designed to anneal to backbone sequences close to the right border-like sequence: 5'-CACGCTAAGTGCCGGCCGTCCGAG-3' (SEQ ID NO. 66), and 5'-TCCTAATCGACGGCGCACCGGCTG-3' (SEQ ID NO. 67). Data from this experiment confirm that plants that are positive for the IPT gene are also positive for this other part of the backbone.

[0300] Similar experiments were carried out with the potato variety Russet Burbank. Based on an assessment of IPT phenotypes, the backbone integration frequencies for pSIM108 and pSIM109 were shown to be comparable to those in Russet Ranger (see Tables 4 and 5).

Example 4

Potato Invertase Inhibitor Gene

[0301] Using conventional transformation methods, this Example demonstrates that overexpressing a novel potato invertase inhibitor gene enhances the processing and health characteristics of potato tubers.

[0302] The following primers were designed to amplify a new potato homolog of the tobacco vacuolar invertase inhibitor Nt-inhh1 (Greiner et al., Nature Biotechnology, 17, 708-711, 1999): 5'-AAAGTTGAATTCAAATGAGAAATT-TATTC-3' (SEQ ID NO. 68), and 5'-TTTAAAGCTTTTCAT-AATAACATTCTAAT-3' (SEQ ID NO. 69). The amplification reaction was performed by mixing the following components: 4 µl plant DNA, 2 µl forward primer (10 pM/ml), 2 µl reverse primer, 25 µl Hot Start Master Mix (Qiagen Catalog Nr. 203443), and 17 µl water. This reaction mix was subjected to the following polymerase chain reaction (PCR) conditions using a PTC-100 thermocycler (MJ Research): (1) 5 minutes at 95° C. (1 cycle), (2) 1 minute at 94° C., 1 minute at 45° C. and 4 minutes at 72° C. (35 cycles), and (3) 10 minutes at 72OC (1 cycle). The total product was loaded on a 0.8% agarose gel, and a 540 base pair band was purified from gel using QIAquick Gel Extraction Kit (Qiagen, Calif.). This purified fragment was then ligated into pGEM-T Easy (Promega, Wis.) and transformed

into *E. coli* DH5-alpha using Max Efficiency Competent Cells (GibcoBRL, MD). Sequence analysis of recombinant plasmid DNA isolated from transformed DH5-alpha revealed the presence of a single open reading frame consisting of 543 base pairs that encodes for a putative 181-amino acid protein (SEQ ID NO.: 5); clustal-alignment revealed 70% homology to Nt-inhh (**FIG. 2**). This high level of homology extends to the 15-amino acid N-terminal domain, indicating that the potato homolog is targeted to the vacuole. Interestingly, the potato invertase inhibitor homolog, designated St-inh1, shares only 43% homology with the patented tobacco cell wall invertase inhibitor designated Nt-inh1 (Patent WO98/04722; **FIG. 2**).

[0303] Although the St-inh1 gene is present in unmodified potato tubers, its expression level is inadequate for full inhibition of invertase and reduced cold-induced sweetening. To increase the storage characteristics of potato, the St-inh1 gene was fused to a new tuber-enhanced promoter of the granule-bound starch synthase (GBSS) gene, which is known to promote high levels of gene expression in tubers. The GBSS promoter was isolated from the potato cultivar Russet Ranger by carrying out a PCR reaction using the forward primer 5'-GAACCATGCATCTCAATC-3' (SEQ ID NO. 70) and the reverse primer 5'-GTCAGGATCCCTACCAAGCTACAGATGAAC-3' (SEQ ID NO. 71). Sequence analysis of the amplified product cloned in pGEM-T demonstrated that this new promoter contains 658 basepairs (SEQ ID NO.: 6). The resulting promoter/gene fusion was then ligated to the 3' regulatory sequence of the potato ubiquitin gene (UbiT; SEQ ID NO.: 7), thus ensuring appropriate termination of transcription of the invertase inhibitor gene.

[0304] This expression cassette was inserted between T-DNA borders of a binary vector, and the resulting vector pSIM320 was used to transform Russet Ranger as described above. Three cuttings of nine independent transgenic lines were planted in soil and grown for four weeks in a growth chamber (11 hrs light; 20° C.). At least 3 minitubers were then harvested from each line and transferred to a refrigerator set at 4° C. to induce cold-sweetening. After 4 weeks, the glucose levels in these cold-stored minitubers were determined by using either an Accu-Chek meter and test strips (Roche Diagnostics, IN) or a glucose oxidase/peroxidase reagent (Megazyme, Ireland). These levels were compared with the average glucose levels in both 6 untransformed lines and 6 "vector control" lines transformed with a pSIM110-derived vector lacking the invertase inhibitor gene. As shown in Table 6, three transgenic lines accumulated less than 40% of the glucose in "vector control" lines demonstrating that the potato invertase inhibitor homolog is functionally active.

[0305] The following experiment showed that the amount of reducing sugars present in tubers correlates with acrylamide production during tuber processing. Russet Ranger potato tubers were freshly harvested from the field and stored at 4° C. to induce cold-sweetening; control tubers were stored at 18° C. After 4 weeks, glucose levels were determined in both groups of tubers. Subsequently, tubers were washed, blanched for either 8 minutes or 12 minutes at 165° F., cut into 0.290x0.290 shoestring strips, dipped in a 1% sodium acid pyrophosphate solution at 160° F., dried at 160° F. until 14±2% dryer weight loss is achieved, fried at 390° F. for 40 seconds to attain 64±2% first fry moisture, and

frozen for 20 minutes at -15° F., shaking the tray 2-3 times in the first 6 minutes. The resulting French fries were then analyzed for acrylamide levels by Covance laboratory (WI). As shown in Table 7, the glucose levels in tubers stored at 18° C. were below the detection level of 0.1 mg/g whereas cold-stored tubers contained on average 3.4 mg/g glucose. This table also shows that fries produced from the latter potatoes contain about 10-fold higher levels of acrylamide than fries produced from potatoes stored at 18° C. Even by using a shorter blanch time for 18° C.-stored potatoes than for 4° C.-stored potatoes to produce fries with a similar color (color ids of 78 and 71, respectively), a 5-fold difference in acrylamide accumulation was obtained (Table 7). Thus, there appears to be a straight correlation between the amount of reducing sugars such as glucose in tubers and the accumulation of acrylamide in fries derived from these tubers.

[0306] To determine whether the reduced glucose levels in pSIM320 lines would limit the processing-induced accumulation of acrylamide, cold-stored pSIM320 minitubers were processed by cutting into wedges, blanching for 8 minutes, dipping in 0.5% SAPP for 30 seconds, drying for 4.5 minutes at 160° F., frying for 40 seconds at 380° F., freezing for 15 minutes at -15° F., and finally drying for 3 minutes and 10 seconds at 160° F. The processed material was then shipped to Covance laboratory for acrylamide determinations. As shown in Table 6, French fries obtained from minitubers with the lowest amounts of glucose accumulated the lowest levels of acrylamide. A 40% reduction in glucose levels in lines "320-2" and "320-4" is associated with a 5-fold reduction in acrylamide levels.

Example 5

Leader and Trailer Sequences Associated with the Potato R1 Gene

[0307] Using conventional transformation methods, this Example demonstrates that a novel leader sequence associated with the potato R1 gene can be used effectively to enhance the processing and health characteristics of potato tubers. It also predicts that a novel trailer associated with that same gene can be exploited in the same way.

[0308] As an alternative to overexpressing the invertase inhibitor gene, methods were developed to limit acrylamide production without using any actual gene sequences. One such method is based on silencing the tuber-expressed R1 gene. Previously, it was shown that this starch-related gene can be silenced through antisense expression of a 1.9-kb gene fragment derived from that gene (Kossmann et al., U.S. Pat. No. 6,207,880). However, the antisense expression of large DNA fragments is undesirable because such fragments contain new open reading frames (Table 1). As a safer approach to the one described above, a small leader sequence associated with the R1 gene was isolated from potato. This leader was obtained by performing a rapid amplification of cDNA ends with the 5' RACE kit supplied by GIBCO BRL on total RNA from the tubers of Russet Ranger potato plants. Sequence analysis demonstrated that the R1-associated leader consists of 179 basepairs (SEQ ID NO.: 8). Both a sense and antisense copy of this leader sequence, separated by the potato Ubiquitin intron (SEQ ID NO.: 9), were placed between the GBSS promoter and UbiT. The resulting expression cassette for the leader sequence associated with R1 is shown in **FIG. 3** (SEQ ID NO.: 10).

A similar cassette containing a spacer derived from the GBSS promoter (SEQ ID NO.: 11)-instead of the Ubi intron-separating the sense and antisense copies of the R1 trailer is shown in (FIG. 3; SEQ ID NOs.: 12). Additional variants with a longer version of the GBSS promoter (SEQ ID NO.: 13) are shown in FIG. 3 (SEQ ID NOs.: 14-15).

[0309] To test the efficacy of the R1-associated leader in limiting acrylamide production, the expression cassette shown in FIG. 3 was inserted as KpnI—XbaI fragment between T-DNA borders of a binary vector. An *Agrobacterium* LBA4404 strain carrying the resulting vector pSIM332 was used to transform Russet Ranger potato. To induce tuber formation, 25 shoots representing independent transformation events were transferred to soil and placed in a growth chamber (11 hours light, 25° C.). After three weeks, at least 3 minitubers/line were stored for 4 weeks at 4° C. to induce starch mobilization. The glucose levels in these cold-stored minitubers were subsequently determined as described in Example 4, and compared with the average glucose levels in untransformed plants and vector controls. As shown in Table 8, minitubers derived from all 25 lines displayed reduced levels of glucose after cold-storage. An approximate 2-fold reduction in acrylamide levels in expected in French fries derived from minitubers displaying reduced R1 expression levels compared to controls. Much stronger effects of down-regulating R1 gene expression are anticipated in mature tubers.

[0310] As an alternative to the leader-based approach, expression cassettes that contained both a sense and antisense copy of the trailer sequence associated with R1 were generated. This trailer was obtained by performing a reverse transcription polymerase chain reaction (RT-PCR) on total RNA isolated from microtubers of the potato cultivar Russet Ranger. Complementary DNA was generated using the Omniscript RT Kit (Qiagen, Calif.) and then used as a template for a PCR reaction with Hot start DNA polymerase (Qiagen, Calif.) with the gene-specific reverse primer R1-1 (5'-GTTTCAGACAAGACCACAGATGTGA-3'). Sequence analysis of the amplified DNA fragment, cloned in pGEM-T demonstrated that the trailer associated with R1 consists of 333 basepairs (SEQ ID NO.: 16). The sense and antisense copies of the trailer were separated by either the Ubi intron or the GBSS spacer- and sandwiched between GBSS promoter and Ubi3 terminator (FIG. 3; SEQ ID NOs.: 17-18). Similar versions with the larger GBSS promoter are shown in FIG. 3 (SEQ ID NOs.: 19-20).

[0311] Glucose and acrylamide levels can be determined as described above. Tubers displaying about 50% or greater reductions in glucose concentrations are expected to also accumulate about 50% less acrylamide during the frying process. The improved health and storage characteristics of modified, plants can be confirmed in mature field-grown tubers.

[0312] Phosphate levels in potato tubers can be determined by using AOAC Method 995.11 Phosphorus (Total) in Foods (45.1.33 Official Methods of Analysis of AOAC International, 17th Edition). Samples are prepared by dry ashing in a muffle furnace followed with an acid digestion. The dissolved samples are then neutralized and treated with a molybdate-ascorbic acid solution and compared to a series of phosphorus standards (treated similarly). A dual beam spectrophotometer would be used for the calorimetric analy-

sis at 823 nanometers. A significant decrease in phosphate content, which is beneficial for the environment, is expected.

Example 6

Leader Sequence Associated with the L-Alpha Glucan Phosphorylase Gene

[0313] Using conventional transformation methods, this Example demonstrates that a novel leader sequence associated with the potato L-alpha glucan phosphorylase gene can be used to effectively enhance the processing and health characteristics of potato tubers.

[0314] Previously, it was shown that cold-induced sweetening can be reduced through antisense expression of 0.9-kb fragments derived from alpha glucan phosphorylase genes (Kawchuk et al., U.S. Pat. No. 5,998,701, 1999). However, the antisense expression of these relatively large DNA fragments is undesirable because they contain new and uncharacterized open reading frames that may impact the nutritional quality of foods if expressed in transgenic plants (Table 1).

[0315] As a safer approach to the one described above, small leader and trailer sequences that are associated with a L-type glucan phosphorylase gene were isolated from RNA of mature tubers. The primer pair used for this purpose is: 5'-GGATCCGAGTGTGGGTAAGTAATTAAG-3' (SEQ ID NO. 72), and 5'-GAATTCTGTGCTCTCTATGCAAATCTAGC-3' (SEQ ID NO. 73). The resultant leader sequence of 273 bp was amplified and is shown in SEQ ID NO.: 21. Similarly, the "direct" primer, 5'-GGAACATGTAAGCTGTGG-3' (SEQ ID NO. 74), was used -with an oligo-dT primer to amplify a 158 bp "trailer sequence" that is associated with the L-type phosphorylase gene (SEQ ID NO.: 22).

[0316] Expression cassettes were then designed using these trailer or leader sequences to modify the expression of L-type phosphorylase gene and, in so doing, lowering acrylamide levels in fried products by limiting starch mobilization. These cassettes were constructed in a similar way as described in Example 5, and are depicted in FIG. 3 (SEQ ID Nos.: 23-26). An *Agrobacterium* strain containing a binary vector with this expression cassette, designated pSIM216, was used to infect potato stems, and generate 25 transgenic plants. Minitubers derived from these plants were stored for 4 weeks at 4° C. to induce cold-sweetening. The cold-stored minitubers were then analyzed for glucose levels. As shown in Table 9, minitubers from all transgenic lines displayed reduced glucose levels.

[0317] Four lines that displayed at least 50% reduced glucose concentrations (lines 216-2, 216-5, 216-10, and 216-21) were used to assess processing-induced acrylamide levels. Although acrylamide levels in fried tubers derived from the first three lines were similar to those of controls, French fries that were derived from line 216-21 accumulated only 45% of the wild-type acrylamide levels (136 vs. 305 parts per billion). These results confirm the experiments described in Example 4 for tubers overexpressing the potato invertase inhibitor gene, in that relatively large reductions in glucose (and fructose) concentrations are needed to limit the heating-induced acrylamide accumulation in cold-stored minitubers. Because silencing of the phosphorylase gene is expected to be more effective in mature "216" tubers,

reductions in acrylamide levels are also anticipated to be more pronounced in the French fries produced from such tubers. The improved health and storage characteristics of modified plants can be confirmed in mature tubers.

Example 7

Modified Polyphenol Oxidase Gene

[0318] Using conventional transformation methods, this Example demonstrates that a modified polyphenol oxidase gene lacking a functional copper-binding site can be used effectively to reduce bruise susceptibility in tubers.

[0319] Previously, it was shown that black spot bruise susceptibility can be reduced through antisense expression of the 1.8-kb PPO gene (Steffens, U.S. Pat. No. 6,160,204, 2000). However, expression of the reverse complement of this large gene is undesirable because it contains new and uncharacterized open reading frames encoding peptides consisting of more than 100 amino acids, which may potentially impact the nutritional quality of foods (Table 1). As a safer approach to the one described above, the PPO gene was modified to encode a non-functional protein.

[0320] The wild-type potato PPO gene was isolated from Russet Ranger by using a polymerase chain reaction (PCR) method. First, genomic DNA was isolated from sprouts of Russet Ranger. The potato PPO gene was then amplified from the potato genomic DNA using DNA polymerase and oligonucleotide primers: 5': CGAATTCATGGCAAGCT-TGTGCAATAG-3' (PPO-F) (SEQ ID NO. 75), and 5'-CGAATTCTTAACAATCTGCAAGACTGATCG-3' (PPO-R) (SEQ ID NO. 76). These were designed to complement the 5'- and 3'-ends of the potato PPO gene. The amplified 1.6 kb fragment was cloned into a pGEM-T EASY vector (Promega) and confirmed to represent a functional PPO gene by sequence analysis (SEQ ID NO.: 27).

[0321] The copper binding domain in potato PPO was identified by aligning this protein with a sweet potato PPO protein that was shown to contain conserved Cysteine (Cys) residue at position 92, Glutamine residue (Glu) at position 236, and Histidine (His) residues at positions 88, 109, 118, 240, 244 and 274 coordinating the two active site coppers (Klabunde et al., Nature Structural Biol., 5: 1084-1090, 1998). These Cys, Glu, and His residues are also present in potato PPO.

[0322] The inactive PPO gene was created by using a PCR mutation replacement approach. Three fragments were amplified by Proof Start Taq DNA Polymerase (Qiagen) using 3 pairs of primers and wild-type Russet Ranger PPO as a template. The sequences of the first pair, designated P1-F and P2-R, respectively, are: 5'-GAGAGATCT-TGATAAGACACAACC-3' (SEQ ID NO. 77), and 5'-CATTACC¹ATAAGCC²CAC³TGTATATTAG CTTGT-TGC-3' (SEQ ID NO. 78) (1: "A" to "C" mutation, resulting in Cysteine to Glycine substitution at position 186; 2: "A" to "C" mutation, resulting in Cysteine to Tryptophan substitution at position 183; 3: "A" to "C" mutation, resulting in Histidine to Glutamine substitution at position 182). The sequences of the second pair, designated P3-F and P4-R, respectively, are 5'-GTGCTTATAGAATTGGTGGC-3' (SEQ ID NO. 79), and 5'-TAGTTCCTCCGGGAGTTCAGTG-3' (SEQ ID NO. 80). The sequences of the third pair, designated P5-F and P6-R, respectively, are

5'-CTCCCGGGAACTATAGG⁴AAACATTC CTCT⁵CGGTCCTGTCCACATCTGGTC-3' (SEQ ID NO. 81) and 5'-GTGTGATATCTGTTCTTTTCC-3' (SEQ ID NO. 82) (4: "A" to "G" mutation, resulting in Glutamine to Glycine substitution at position 326; 5: "A" to "T" mutation, resulting in Histidine to Leucine substitution at position 330).

[0323] An 80 bp fragment was amplified using primer P1-F and P2-R and digested with BglIII. This fragment contains one sticky end (BglIII) and one blunt end, and carries three mutations in copper binding site I. A 0.4 kb fragment amplified using primer P3-F and P4-R and digested with XmaI contains one blunt end and one sticky end (XmaI). A 0.2 Kb fragment was amplified using primer P5-F and P6-R and digested with XmaI and EcoRV. This third fragment with a sticky end (XmaI) and a blunt end (EcoRV) has two mutations in copper binding site II. The BglIII and EcoRV fragment from cloned wild-type potato PPO was then replaced with the above three ligated PCR amplified fragments. The presence of a total of 5 point mutations in the modified PPO gene was confirmed by sequence analysis (SEQ ID NO.: 28). To create an expression cassette for modified PPO (mPPO), the following four fragments were simultaneously ligated together: (1) a BamHI-HindIII fragment containing the GBSS promoter, (2) a HindIII-SacI fragment containing mutant PPO, (3) a SacI-KpnI fragment containing the Ubi-3 terminator, and (4) plasmid pBluescript, digested with KpnI and BamHI. This expression cassette was then inserted between borders of a binary vector to create pSIM314.

[0324] The efficacy of the mPPO gene expression cassette was assessed by transforming Russet Ranger stem explants with pSIM314. Nodal cuttings of transgenic plants containing this expression cassette were placed on MS medium supplemented with 7% sucrose. After a 5-week incubation period in the dark at 18° C., microtubers were isolated and assayed for PPO activity. For this purpose, 1 g of potato tubers was pulverized in liquid nitrogen. This powder was then added to 5 ml of 50 mM MOPS (3-(N-morpholino) propane-sulfonic acid) buffer (pH 6.5) containing 50 mM catechol, and incubated at room temperature with rotation for about 1 hour. The solid fraction was then precipitated, and the supernatant transferred to another tube to determine PPO activity by measuring the change of OD-410 over time. As shown in Table 10, microtubers isolated from some of the transgenic lines displayed a significantly reduced polyphenol oxidase activity compared to either untransformed controls or controls transformed with a construct not containing the mutant PPO gene. The strongest reduction in PPO activity was observed in lines "314-9", "314-17", and "314-29". To test whether expression of the mutant PPO gene also reduced PPO activity in minitubers, rooted plantlets of transgenic lines were planted in soil and incubated in a growth chamber for 4 weeks. A PPO assay on isolated minitubers demonstrated that reduced PPO activity in microtubers correlated in most cases with reduced activities in minitubers (Table 10). Transgenic lines displaying a reduced PPO activity can be propagated and tested both in the greenhouse and the field to confirm the "low bruise" phenotype in mature tubers. Because micro- and minitubers express a variety of polyphenol oxidases, some of which share only limited sequence homology with the targeted polyphenol oxidase that is predominantly expressed in mature tubers, an even more profound reduction of PPO activity may be anticipated in the mature tubers of lines such

as “314-9” and “314-17”. The data indicate that overexpression of a functionally inactive PPO gene can result in reduced bruise susceptibility. The improved health and storage characteristics of modified plants can also be confirmed in mature field-grown tubers.

Example 8

Trailer Sequence of a Polyphenol Oxidase Gene that is Specific for the Non-Epidermal Tissues of Potato Tubers

[0325] Using conventional transformation methods, this Example demonstrates that a novel trailer sequence associated with the potato PPO gene can be used effectively to reduce bruise susceptibility in tubers.

[0326] Reverse transcription PCR was used to also isolate the trailer sequence associated with the PPO gene expressed in potato tubers. The primers for the first PCR reaction were PPO-1 (5'-GAATGAGCTTGACAAGGCGGAG-3', (SEQ ID NO. 83)) and oligo-dT; primers for a second nested PCR reaction were PPO-2 (5'-CTGGCGATAACGGAAGTGTG-3', (SEQ ID NO. 84)) and oligo-dT. Sequence analysis of the amplified DNA fragments cloned into pGEM-T revealed the presence of a 154-bp trailer (SEQ ID NO.: 29). A sense and antisense copy of this trailer, separated by the Ubi intron, was then fused to the GBSS promoter and Ubi3 terminator as described above to generate an expression cassette shown in FIG. 3 (SEQ ID NO.: 30). An alternative construct containing the trailer segments separated by a GBSS spacer is shown in FIG. 3 (SEQ ID NO.: 31). Similar versions with the larger GBSS promoter are shown in FIG. 3 (SEQ ID NOs.: 32-33). Interestingly, the trailer of the PPO gene that is predominantly expressed in mature tubers (indicated with P-PPO3 in FIG. 4) is different from the trailer of PPO genes that are predominantly expressed in other tissues including microtubers (indicated with PPOM-41 and PPOM-44 in FIG. 4). Because of the low homology between trailers associated with different PPO genes, the use of the P-PPO3 trailer will result in a silencing of the mature tuber-specific PPO gene only. This very specific gene silencing would be difficult to accomplish with sequences derived from the PPO gene itself, thus demonstrating the advantage of using non-coding sequences for gene silencing. To visualize the extend of PPO activity, 0.5 mL of 50 mM catechol was pipetted on the cut surfaces of sliced genetically modified minitubers. Compared to controls, visual browning of the tuber regions was about 5 to 10-fold reduced. Interestingly, though, no reduced browning was observed in the potato skin. It appears that the trailer sequence used specifically silenced the PPO gene that is predominantly expressed in cortex and pith but not in the epidermal skin. This unexpected finding may be beneficial for tubers to protect themselves against some pathogens attempting to infect through the skin because the PPO gene may play some role in certain defense responses. To quantitatively determine PPO activity, an assay was performed as described in Example 7. Table 11 shows up to 80% reduction of PPO activity in transformed minitubers compared to untransformed controls. The level of reduction is expected to be even greater in mature tubers because these tubers express the targeted PPO gene more predominantly than mini- and microtubers. The improved characteristics of lines such as “217-7” and “217-26” can be confirmed in mature tubers.

Example 9

An Expression Cassette to Increase Levels of Resistant Starch

[0327] Increasing the amylose/amylopectin ratios in tubers can further enhance the nutritional value of potato products. One method that makes it possible to increase amylose content is based on the antisense expression of genes encoding for the starch branching enzyme (SBE) I and II (Schwall et al., *Nature Biotechnology* 18: 551-554, 2000). The disadvantages of this method are that (1) the efficiency of simultaneously silencing two different genes through exploitation of antisense technologies is very low, (2) the antisense expression of the relatively large SBE-I and SBE-II gene sequences results in the undesirable expression of open reading frames (Table 1) (3) corresponding constructs that harbor the two antisense expression cassettes are unnecessarily large and complex, thus, increasing chances of recombination and lowering transformation frequencies.

[0328] Our approach to increase amylose content in potato is based on the expression of the trailer sequences that are associated with-both genes. These trailers (SEQ ID No.:34 and 35) were isolated with the primer pairs 5'-GTCCATGATGTCTTCAGGGTGGA-3' (SEQ ID NO. 85), and 5'-CTAATATTTGATATATGTGATTGT-3' (SEQ ID NO. 86), and 5'-ACGAACCTGTGATCGCCITTGAAAG-3' (SEQ ID NO. 87), and 5'-ACTAAGCAAAACCTGCTGAAGCCC-3' (SEQ ID NO. 88). A single promoter drives expression of a sense and antisense fusion of both trailers, separated by the Ubiquitin-7 intron, and followed by the Ubiquitin-3 terminator. The size of the entire expression cassette is only 2.5-kb.

Example 10

Development of Marker-Free Transformation Methods

[0329] This Example demonstrates that plants can be transformed effectively without to need for stable integration of selectable marker genes.

[0330] This method is the first to take advantage of the phenomenon that DNAs targeted to the nuclei of plant cells often fails to subsequently integrate into the plant cell's genome. The inventors made the surprising discovery that it is possible to select for cells that temporarily express a non-integrating T-DNA containing a selectable marker gene by placing infected explants for 5 days on a plant medium with the appropriate selective agent. A second phenomenon that was applied to develop the current method is that T-DNAs from different binary vectors often target the same plant cell nucleus. By using two different binary vectors, one containing the selectable marker on a T-DNA, and the other one carrying a T-DNA or P-DNA with the actual sequences of interest, it was possible to apply a transient selection system and obtain populations of calli, shoots or plants, a significant portion of which represents marker-free transformation events.

[0331] A conventional binary vector designated pSIM011 was used to represent the vector with the “sequence of interest”, which is, in this test case, an expression cassette for the beta glucuronidase (GUS) gene located on a conventional T-DNA. The second binary vector that was used

for these experiments contains an expression cassette comprising the neomycin phosphotransferase (NPTII) gene driven by the strong promoter of the Ubiquitin-7 gene and followed by the terminator sequences of the nopaline synthase (nos) gene between the borders of the T-DNA of a pSIM011-derivative.

[0332] Surprisingly, a strong level of transient NPTII gene expression levels could also be obtained by replacing the nos terminator with the terminator of the yeast alcohol dehydrogenase 1 (ADH1) gene (Genbank accession number V01292, SEQ ID NO. 56). This finding is interesting because the yeast ADH1 terminator does not share homology with any plant terminator. Importantly, it should be noted here that many yeast terminators do not function adequately in plants. For instance, almost no GUS gene expression was observed in a similar experiment as described above with the GUS gene followed by the yeast iso-1-cytochrome c (CYC1) terminator (Genbank accession number SCCYT1). An improved vector carrying the selectable marker gene NPTII was generated by replacing the nos terminator with the yeast ADH1 terminator. The binary vector containing a selectable marker gene for transient transformation is designated "LifeSupport" (FIG. 5).

[0333] Potato stem explants were simultaneously infected with two *A. tumefaciens* LBA4404 strains containing pSIM011 and LifeSupport, respectively. A 1/10 dilution of overnight-grown cultures of each strain were grown for 5-6 hours before they were precipitated, washed and resuspended an OD_{600nm} of 0.4 as described in Example 3. The resuspended cells were then used to infect 0.4-0.6 cm internodal potato segments at a final density of each bacteria of 0.2 (OD_{600nm}). Infected stems were treated as in Example 3 with a main difference: the selection with kanamycin was limited to the first 5 days of culture on callus induction medium. Then, explants were allowed to further develop in fresh CIM and SIM containing only timentine 150 mg/L but no selective antibiotic. Within about 3 months from the infection day leaves from shoots derived from calli developed in 40-60% of the infected stems were both tested for GUS expression and PCR analyzed to identify events that contained the sequences of interest but no marker gene. As shown in Table 12, 11% of shoots represented marker-free transformation events.

[0334] The two-strain approach described above was also used to transform tobacco. Shoots that developed within about 2 months were GUS assayed and PCR analyzed. The high frequency of marker-free transformation events identified (18%) implies that the developed method is applicable to plant species other than potato (Table 12).

[0335] Importantly, sequential rather than simultaneous infection with the two different *Agrobacterium* strains resulted in an increase in the efficiency of marker-free transformation. The surprising effect of sequential infections was discovered by infecting potato stem explants with the *Agrobacterium* strain containing pSIM011, placing the infected explants on co-cultivation plates for 4 hours, and then re-infecting them with the LifeSupport vector. The doubly infected explants were treated as previously described in this example. As shown in Table 13, the lag time of 4 hours between the two different infections resulted in a 2-fold increased frequency of marker-free transformation events in potato.

Example 11

Precise Breeding with pSIM340

[0336] This Example demonstrates the efficacy of precise breeding. The health and agronomic characteristics of potato plants are enhanced by inserting potato genetic elements (see Examples 1, 4, and 7) into potato, using marker-free transformation (Example 10).

[0337] A binary vector containing two expression cassettes for the invertase inhibitor and mutant polyphenol oxidase genes inserted between P-DNA termini, designated pSIM340 (FIG. 1), was created by inserting both expression cassettes of mutant PPO and invertase inhibitor into a binary vector pSIM112'. Potato stem explants were infected simultaneously with pSIM340 and a further improved LifeSupport vector. The infected explants were then co-cultivated, subjected to transient selection, and induced to proliferate and develop shoots as discussed earlier. After 3 months, small shoots were transferred to new media and allowed to grow for 3 additional weeks. Shoots were then phenotypically analyzed, and leaf material was collected for molecular analyses to determine the presence of backbone, marker gene and P-DNA with the sequences of interest, as described in Examples 2 and 3. As shown in Table 14, 1.2% of events represented a plant that contained the modified P-DNA of pSIM340 without LifeSupport. This frequency of marker-free transformation is lower than found for a T-DNA, again revealing a functional difference between P-DNA and T-DNA.

Example 12

Selecting Against Stable Integration of LifeSupport T-DNAs

[0338] This Example demonstrates that the efficiency of precise breeding methods can be increased by selecting against stable integration of LifeSupport T-DNAs using the bacterial cytosine deaminase gene.

[0339] The previous example demonstrates that the efficiency of marker-free transformation is several-fold lower with a modified P-DNA than with a conventional T-DNA. To improve the efficiency of generating shoots only containing a modified P-DNA, an expression cassette for a suicide gene fusion comprising the bacterial cytosine deaminase (codA) and uracil phosphoribosyltransferase (upp) genes (InvivoGen, CA) was inserted between T-DNA borders of the LifeSupport vector, generating pSIM346 (FIG. 5). Potato stem explants were infected with one strain carrying pSIM340 and the other carrying pSIM346, and subsequently placed on the following media: (1) co-cultivation media for 2 days, (2) CIMTK media to select for transient marker gene expression for 5 days, (3) CIMT media to allow proliferation of plant cells that transiently expressed the marker gene for 30 days, (4) SIMT media with 500 mg/L of non-toxic 5-fluorocytosine (5-FC), which will be converted by plant cells expressing codA::upp into the toxic 5-fluorouracil (5-FU), to select against stable integration of the LifeSupport TDNA. Callus gave rise to shoots on SIMT within 4

weeks. These shoots were transferred to MS media with timentin and allowed to grow until sufficient tissue was available for PCR analysis. DNA was then extracted from 100 shoots and used to determine the presence of P-DNA, LifeSupport and backbone. As shown in Table 15, none of the shoots analyzed contained a LifeSupport T-DNA, indicating, for the first time, that the *codA::upp* gene fusion can be used as negative selectable marker prior to regeneration. More importantly, our results demonstrate that a negative selection against LifeSupport T-DNA integration increases the frequency of shoots that only contain a modified P-DNA. By coupling a positive selection for transient marker gene expression with a negative selection against stable integration of the *codA::upp* gene fusion, the frequency of shoots only containing a modified P-DNA is about 5-fold higher than by only employing the positive selection for transient marker gene expression (Table 15).

[0340] An even greater increase in the efficiency of marker-free transformation was obtained by using the LifeSupport vector pSIM350 (FIG. 5), which is similar to pSIM346 but contains the *codA* gene instead of the *codA::upp* gene fusion. Potato stem explants simultaneously infected with pSIM340 and pSIM350 were treated as described above, and 51 resulting shoots were molecularly tested for the occurrence of events only containing the T-DNA region from pSIM340. Interestingly, this PCR analysis revealed that some shoots contained the *codA* gene (Table 15). This finding demonstrates that *codA* is not as tight a negative selectable marker as *codA::upp* in plants. More importantly, a large number of shoots (29%) were shown to represent marker-free transformation events.

[0341] Efficiencies can be further increased by not infecting explants simultaneously with pSIM340 and pSIM350 but sequentially. By infecting the explants with pSIM340 and re-infecting them with pSIM350 after 4 hours, marker-free transformation frequencies are expected to be approximately 30-40%.

Example 13

Impairing Integration of LifeSupport T-DNAs

[0342] This Example demonstrates that the efficiency of precise breeding methods can be increased by impairing integration of the LifeSupport T-DNA into the plant genome using an omega-mutated *virD2* gene.

[0343] It has been shown that the omega domain of the *Agrobacterium* protein *virD2* is important for the ability of that protein to support T-DNA integration into plant genomes (Mysore et al., *Mol Plant Microbe Interact* 11: 668-83, 1998). Based on this observation, modified LifeSupport vectors were created that contain an expression cassette for an omega-mutated *virD2* protein inserted into the *SacI* site in their backbone sequences. The expression cassette was obtained by amplifying a 2.2-kb DNA fragment from plasmid pCS45 (courtesy of Dr. Walt Ream—Oregon State University, OR, USA-, SEQ ID NO.: 36). A LifeSupport-derivative carrying this expression cassette, designated pSIM401Ω (FIG. 5), was used to support the transformation

of potato plants with the modified P-DNA of pSIM340. After transient selection and shoot induction, 100 shoots were molecularly tested for the presence of transgenes. As shown in Table 15, 4.4% of shoots only contained the modified P-DNA, indicating that the use of omega-*virD2* increases the efficiency of marker-free transformation about 4-fold (Table 15).

[0344] Efficiencies are further improved by increasing the size of the LifeSupport T-DNA from 3.7 kb (in pSIM401Ω) to 8.1 kb (in the pSIM401Ω-derivative designated pSIM341Ω; FIG. 5). By regenerating shoots from potato stem explants simultaneously infected with pSIM340 and pSIM341Ω, 7 of 81 analyzed events (7%) were shown to represent marker-free transformation events (Table 15).

[0345] A further improvement can be obtained by infecting explants sequentially rather than simultaneously with pSIM340 and LifeSupport. In a similar way as described in Example 10, the frequency of plants that only contain a modified P-DNA can be about doubled by infecting the explants with pSIM340 and re-infecting them with LifeSupport after 4 hours.

Example 14

Development of a 1-Strain Approach

[0346] This Example demonstrates that high frequencies of marker-free transformation can also be obtained by using a single *Agrobacterium* strain that contains both the P-DNA vector and Life Support

[0347] Two compatible binary vectors were created that can be maintained simultaneously in *Agrobacterium*. Instead of using this system to stably integrate two T-DNAs carrying the DNA-of-interest and a marker gene, respectively (Komari et al. U.S. Pat. No. 5,731,179, 1998), it is intended for integration of only the modified P-DNA.

[0348] The first vector, designated pSIM356, contains an expression cassette comprising the GUS gene driven by the Ubi7 promoter and followed by UbiT inserted between P-DNA termini. The backbone portion of this vector contains bacterial origins of replication from pVS1 and pBR322, a spectinomycin resistance gene for bacterial selection, and an expression cassette for the IPT gene to enable selection against backbone integration in plants (FIG. 1). The second vector, designated pSIM363, contains an expression cassette comprising the NPTII gene driven by the Ubi7 promoter and followed by the yeast ADH1 terminator inserted between conventional T-DNA borders (FIG. 5). The backbone portion of this vector contains bacterial origins of replication from ColE1 (Genbank number V00268) and ori V (Genbank number M20134), and a kanamycin resistance gene for bacterial selection.

[0349] The concept of increasing marker-free transformation frequencies using pSIM356 and pSIM363 was tested in 100 tobacco shoots. As shown in Table 16, about 19% of regenerated shoots contained the DNA of interest without marker gene. An increase in marker-free transformation efficiency was also found by applying this 1-strain approach to potato. Nine of 60 independent shoots tested (15%) contained the pSIM340 T-DNA and lacked the LifeSupport T-DNA (Table 16).

[0350] The 1-strain approach can be combined with the method described in Example 12 to couple a positive selection for transient marker gene expression with a negative selection against stable integration of the coda gene. For this purpose, the LifeSupport vector pSIM365 was developed (FIG. 5). An *Agrobacterium* strain carrying this vector together with a P-DNA vector can be used to efficiently develop plants that only contain an expression cassette-of-interest located within a P-DNA stably integrated in their genomes.

Example 15

Precise Breeding Method Relying on a Native Marker

[0351] Apart from transforming crop plants with P-DNAs that only contain the desirable sequences to introduce beneficial traits, the present invention also provides a method of transforming such plants with P-DNAs that contain an additional native marker gene. Our novel and native marker genes of choice are potato homologs of the *Arabidopsis* vacuolar Na⁺/H⁺ antiporter gene and alfalfa alfin-1 gene. Expression of these genes do not only allow the identification of transformation events, but also provides salt tolerance to transformed plants. High salinity levels in an increasing acreage of agricultural land will therefore less affect potato plants containing the salt tolerance marker.

[0352] Two versions of a vacuolar Na⁺/H⁺-antiporter homolog, designated Pst (Potato salt tolerance) were amplified from cDNA of a late blight resistant variety obtained from the US Potato Genbank (WI), designated "LBR4", using the oligonucleotide pair 5'-CCCGGGATGGCTTCTGTGCTGGCT-3' (SEQ ID NO. 89) and 5'-GGTACCTCATGGACCCTGTTCCGT-3' (SEQ ID NO. 90). Their sequences are shown in SEQ ID NO.:37 and 38. A third gene (SEQ ID NO.:39) with homology to alfin-1 was amplified from LBR4 potato DNA using the primers 5'-CCCGGGTATGGAAAATTCGGTACCCAGGACTG-3' (SEQ ID NO. 91) and 5'-ACTAGTTAACTCTAGCTCTCTTGC -3' (SEQ ID NO. 92). The efficacy of the Pst genes to function as transformation marker was assessed by inserting a fusion with the Ubi7 promoter between conventional T-DNA borders of a modified pSIM341 vector. After a transient selection period, kanamycin-resistant cells are allowed to proliferate and develop shoots. These shoots are then transferred to media that contain 100-150 mM sodium chloride. Salt-tolerant shoots represent transformation events that contain the T-DNA of the modified pSIM341.

Example 16

Tuber-Specific Promoter

[0353] A newly isolated tuber-specific promoter can replace the GBSS promoter used to develop the expression cassettes described in previous examples. This promoter was isolated from the genome of Russet Burbank potato plants by using the inverse polymerase chain reaction with primers specific for a potato proteinase inhibitor gene (Genbank

Accession D17332) (SEQ ID NO. 39). The efficacy of the PIP promoter was tested by creating a binary vector that contains the GUS gene driven by this promoter and an expression cassette for the NPTII marker gene. A similar construct with the PIP promoter replaced by the GBSS promoter was used as control. Transformed shoots were obtained by infecting stem explants with *Agrobacterium* strains carrying the binary vectors, co-cultivation for 2 days, and selection on CIMTK medium for 2 months. These shoots were transferred to new media to induce root formation, and then planted into soil. Tubers can be assayed for GUS expression after a 3-month growth period in the green house.

Example 17

Preferred Constructs and Transformation Methods for Precise Breeding

[0354] Apart from pSIM340, many other vectors can be used to improve potato plants by transforming them with modified P-DNAs. Two of such vectors contain an expression cassette for a sense and antisense copy of the trailer associated with a PPO gene that is expressed in all tuber tissues except for the epidermis (see Example 8). Vector pSIM370 contains an additional expression cassette for a sense and antisense copy of the leader associated with phosphorylase gene (see Example 6). Vector pSIM371 contains a third expression cassette for the potato alfin-1 homolog (FIG. 1).

[0355] A third alternative vector, designated pSIM372, contains both an expression cassette for the potato alfin-1 homolog, and an expression cassette for a sense and antisense copy of a fusion of the PPO-associated trailer, R1-associated leader, and phosphorylase-associated leader.

[0356] The preferred LifeSupport vector for a 1-strain approach is pSIM365. For a 2-strain approach, the preferred vector is pSIM367, which contains expression cassettes for both NPTII and coda between T-DNA borders, and an additional expression cassette for omega virD2 in the plasmid backbone (FIG. 5).

[0357] Potato stem explants are infected with 1 strain carrying both pSIM365 and any of the vectors pSIM370, 371, and 372, or sequentially with 2 strains carrying pSIM366 and any of the preferred vectors-of-interest, respectively. After a 2-day co-cultivation and a 5-day transient selection period, the explants are transferred to media for proliferation/regeneration and elimination of *Agrobacterium*. Thirty days later, explants are transferred again to the same media but now also containing 5-FU to eliminate events containing LifeSupport T-DNAs. Shoots that subsequently arise on calli are transferred to regeneration media that may contain 100-200 mM salt to screen for salt tolerant events. The IPT-negative shoots are allowed to root and develop into mature plants. A large proportion of these plants (10%-100%) are predicted to represent marker-free and backbone-free plants containing a P-DNA with nucleotide sequences of interest stably integrated into their genomes.

TABLE 1

<u>Potentially expressed uncharacterized peptides in antisense potato lines</u>						
Gene (size of fragment used)	Predicted peptides encoded by ORFs in reverse-complemented DNA					
R1 (1.9-kb)	MSSTSNVGQD	CLAEVTISYQ	WVGRVINYNF	FLLIHWYTVV	EASTGITFQI	FPIGIRSEDD RSFYEKADRF
	AWVT					
	MSSESTFSKT	PNGRATDVGI	PTEEGTFPFR	YAILRDLAPT	ISLVNSSADI	A
	MSEGVGFKSK	ILPSFAWRS	NILGSKHVAK	QTFPFLARTE	TCERTSGMSG	VIRATAPSGI SSSPLTDFAT
	KIVGFS					
GLTP (1-kb)	VCSPALKADK	SKSADGTCVD	HSRRLIVVLV	LYPGMGTSYA	TAFISSPPIQ	YLFPSDPVET FP
	MLGSLVLPKS	PENRKQAVPN	PHFQEQHLVP	EKPFLDCGQ	GFSKLPQMHQ	
	MVNFLTQGI	DMETAFGSPK	MGFGKEQFG	ACVSRSEMDE	SGIGAVMVEQ	VCSICSRHFV LSMQI
GHTP (0.9-kb)	MLEGSMWPWN	QESMKRAFLN	HHFLMLHLFP	AQRPPQAADP	VCLKHQHMC	GCLSFQLHLS KLAPGDTPLI
	SSMFALD					
	MKLCSSIILS	IIKQKQVEIL	RACFGFPETK	TISVFSSVSW	NWHIICKSL	
	MTKKPDRKDN	IMPYNFPGTK	FLQPIFRNFF	LPSLCDKLLK	KSISVPQAIT	PCWKVQCGHG IKKA
PPO (1.8-kb)	TILKLDLHTF	NGHFFTASFW	NQSHRNSIFI	FQSNILQQFS	YRQLESNTGN	MISITSMNM RQASITPCKL
	RLIKLICIH	LVHVQRHIEP	YIVPIIRYF	IECQYLLLI	FLLCCP	
	MKGKEKPREM	NLQFFTTFNV	STVAISTMNI	SLLFKAKRVK	GVFIKFPHST	RSQILILGYVL LIRRMRSRGAD
	AEFSHRREL	VRNTIDLIGY	RRATTVYYIN	TFFYMGSTTR	LEIRRWYRCS	SR
	MEWALARNRI	PFFYCPNSLR	TSHGKGYDFH	RRKRIQSSTN	LYLLNPFFSR	QLISIHSTSC PHWHGGSKKS
	DLNRVSRNYP	CLHRFFDEVC	HRSRCEPEYE	GCFQ		
SBE A (1.2-kb)	MNNITHSPIL	IPFLEQLNPF	ISNCHMQPIV	KANTPILNGN	TKCRHSANIF	TNGNCIWEKP MNKIVDQHQI
	HNSIHISCES	KVFLVVPSES	HR			
	MKFRYPSPFN	PIVTSILIIC	NAIPRSINDV	DGLSRAIKSY	ISLSISQNAI	VLSPTRA
SBE B (2.6-kb)	MVNIMTSSSM	ATKFPSITVQ	CNSVLPWQVT	SNFIPFVCVL	WVEVEYKYQV	TTFKHNNLII IIAAAYLF
	MAKLVTHEIE	VPLSSQGHCE	KMDHLVKRNS	SINNRRSICQ	ARHARIHLFV	H
	MFETKLNSGV	VWNDWLTVNI	RNSNTPNTKL	VLLHHVVRTV	PSIEIANNFV	FLSSRSPTFI DYATIFPVES
	KF					
	MLYTSLYISY	LSNSMLLPSW	TNLHHSYSLN	NLSTYLGPL	PGGNQNQFLP	QKQAGQGPAY QKHLRQ

[0358]

TABLE 3

Transformation efficiency		
Binary vector	Calli/tobacco leaf explant $\pm$ SE	Calli/potato stem explant $\pm$ SE
pBI121	7.8 $\pm$ 0.6	0.31 $\pm$ 0.10
pSIM108	10.2 $\pm$ 0.6	0.59 $\pm$ 0.07
pSIM109	12.8 $\pm$ 0.6	0.47 $\pm$ 0.05

[0359]

TABLE 4

Backbone integration resulting from Russet Ranger transformation				
Binary vector	Total Nr.	IPT phenotype	PCR ⁺ for IPT	PCR ⁺ for 0.6 kb backbone fragment
pBI121	98	NA	NA	54 (55%)
pSIM108	193	138 (71%)	137 (71%)	NA
pSIM109	133	82 (62%)	80 (60%)	NA

NA: not applicable

[0360]

TABLE 5

Backbone integration resulting from Russet Burbank transformation		
Binary vector	Total Nr.	IPT phenotype
PSIM108	79	49 (60%)
PSIM109	72	60 (84%)

[0361]

TABLE 6

Acrylamide levels in French fries derived from cold-stored pSIM320 minitubers		
Line	glucose mg/g (%-reduced)	acrylamide (PPB)
Untransformed	10.2	469
Vector control	10.2	NA
320-2	5.4 (47%)	95
320-4	5.8 (43%)	107
320-7	8.7 (14%)	353
320-9	7.4 (27%)	137
320-17	6.0 (41%)	506
320-21	8.5 (16%)	428
320-33	6.6 (35%)	516

NA: not available

[0362]

TABLE 7

Acrylamide levels in French fries derived from untransformed mature tubers		
	Stored at 18° C. (color id.*)	Stored at 4° C. (color id.*)
Glucose levels	<0.1 mg/g	3.4 mg/g
8-minute blanch	53 PPB (78)	603 PPB (56)
12-minute blanch	28 PPB (84)	244 PPB (71)

*a higher value indicates a lighter color of the finished Fry product

[0363]

TABLE 8

Glucose levels in cold-stored pSIM332 minitubers	
Line	glucose mg/g (%-reduced)
Untransformed control	11.6 ± 0.5
Vector control	11.5 ± 0.5
332-1	5.4 (53%)
332-2	4.8 (58%)
332-4	7.0 (39%)
332-5	5.8 (50%)
332-6	6.9 (40%)
332-7	6.0 (48%)
332-8	6.8 (41%)
332-9	6.6 (43%)
332-10	5.4 (53%)
332-11	6.1 (47%)
332-12	6.4 (44%)
332-13	6.4 (44%)
332-15	7.7 (33%)
332-16	6.5 (43%)
332-17	5.3 (54%)
332-18	7.1 (38%)
332-21	6.3 (46%)
332-22	5.4 (53%)
332-23	4.2 (63%)
332-31	6.0 (48%)
332-34	6.2 (48%)
332-35	6.4 (44%)
332-39	6.7 (41%)
332-40	7.5 (35%)
332-41	5.7 (50%)

[0364]

TABLE 9

Glucose levels in cold-stored pSIM216 minitubers	
Line	glucose mg/g (%-reduced)
Untransformed control	11.6 ± 0.5
Vector control	11.5 ± 0.5
216-2	5.5 (52%)
216-3	8.8 (23%)
216-4	7.4 (36%)
216-5	5.8 (50%)
216-8	8.4 (27%)
216-10	5.1 (56%)
216-11	10.1 (19%)
216-12	9.3 (19%)
216-13	6.4 (44%)
216-15	8.8 (23%)
216-16	9.7 (16%)
216-17	6.4 (44%)
216-19	8.7 (24%)

TABLE 9-continued

Glucose levels in cold-stored pSIM216 minitubers	
Line	glucose mg/g (%-reduced)
216-21	3.2 (72%)
216-24	9.4 (18%)
216-26	9.3 (19%)
216-29	7.1 (38%)
216-30	8.2 (29%)
216-32	9.3 (19%)
216-34	7.1 (38%)
216-35	7.8 (32%)
216-38	7.1 (38%)
216-42	8.1 (30%)
216-44	9.4 (18%)
216-45	10.2 (11%)

[0365]

TABLE 10

PPO activity in potato lines expressing a modified PPO gene		
Line	OD-410/gram	
	micro tubers (%-reduced)	mini-tubers (%-reduced)
Untransformed controls	24.59 ± 2.22	20.07 ± 1.21
Vector controls	22.59 ± 3.36	19.55 ± 1.43
314-1	2.36 (90%)	17.8 (11%)
314-2	41.52 (-76%)	21.3 (-7%)
314-4	18.40 (22%)	5.4 (73%)
314-5	8.49 (64%)	19.1 (4%)
314-7	16.04 (32%)	16 (20%)
314-8	14.86 (37%)	17 (15%)
314-9	5.43 (77%)	4.3 (78%)
314-12	19.35 (18%)	19.6 (2%)
314-13	18.17 (23%)	15.4 (23%)
314-14	18.64 (21%)	17.32 (13%)
314-16	13.92 (41%)	18.2 (9%)
314-17	5.19 (78%)	2.4 (88%)
314-20	26.66 (-13%)	13.2 (34%)
314-21	11.32 (52%)	17.6 (12%)
314-22	13.45 (43%)	18.8 (6%)
314-23	5.19 (78%)	20.4 (-2%)
314-24	15.10 (36%)	19.6 (2%)
314-25	23.12 (2%)	19 (5%)
314-26	13.45 (43%)	17.8 (11%)
314-27	26.42 (-12%)	19.4 (3%)
314-28	31.85 (-35%)	19.4 (3%)
314-29	3.77 (84%)	14.8 (26%)
314-31	23.83 (-1%)	21.2 (-6%)
314-32	28.78 (-22%)	20 (0%)

[0366]

TABLE 11

PPO activity in potato minitubers expressing a modified trailer sequence associated with the PPO gene	
Line	OD-410/gram (%-reduced)
Untransformed controls	20.6 ± 1.3
Vector controls	17.9 ± 2.1
217-1	12.5 (39.4%)
217-4	12.6 (38.6%)
217-5	11.3 (45.0%)
217-6	6.1 (70.4%)

TABLE 11-continued

PPO activity in potato minitubers expressing a modified trailer sequence associated with the PPO gene	
Line	OD-410/gram (%-reduced)
217-7	5.7 (72.5%)
217-9	10.4 (49.6%)
217-10	15.2 (26.3%)
217-11	15.2 (26.3%)
217-12	6.6 (67.9%)
217-14	15.4 (25.4%)
217-15	13.5 (34.6%)
217-16	6.0 (71.0%)
217-17	9.7 (53.0%)
217-19	8.6 (58.4%)
217-21	14.2 (31.1%)
217-22	9.7 (53.0%)
217-23	15.2 (26.3%)
217-24	8.2 (60.1%)
217-25	11.9 (42.2%)
217-26	3.1 (84.8%)
217-27	6.2 (69.9%)
217-29	7.2 (65.1%)

[0367]

TABLE 12

Marker-free transformation with the LifeSupport vector + pSIM011				
Plant	Co-transformed	Marker only	Gene-of-interest only	Untransformed
Potato	0%	33%	11%	56%
Tobacco	20%	26%	18%	36%

Co-transformed: PCR-positive for both GUS and NPT
 Gene-of-interest only: PCR-positive for GUS
 Untransformed: Plants are PCR-negative for both GUS and NPT

[0368]

TABLE 13

Sequential potato transformation with the LifeSupport vector and pSIM011				
Time window	Co-transformed	Marker only	Gene-of-interest only	Untransformed
0 hrs	9%	36%	9%	46%
4 hrs	20%	30%	20%	30%

Untransformed: Plants are PCR-negative for marker and gene-of-interest

[0369]

TABLE 14

Marker-free transformation with the P-DNA vector pSIM340 + LifeSupport				
Plant	Co-transformed	Marker only	Gene-of-interest only	Untransformed
Potato	17%	52.8%	1.2%	29%

Co-transformed: PCR-positive for both the PPO gene of pSIM340 and the NPT gene from LifeSupport

Untransformed: Plants are PCR-negative for PPO and NPTII

[0370]

TABLE 15

Marker-free potato transformation with pSIM340 + improved LifeSupport vectors				
LifeSupport vector	Co-transformed	Marker only	Gene-of-interest only	Untransformed
PSIM346	0%	0%	4%	96%
PSIM350	10%	10%	29%	51%
PSIM401Ω	6%	34%	5%	55%
pSIM341Ω	16%	23%	7%	54%

Co-transformed: PCR-positive for both the PPO gene of pSIM340 and the NPT gene from LifeSupport

Untransformed: Plants are PCR-negative for PPO and NPTII

[0371]

TABLE 16

Marker-free potato transformation with a single <i>Agrobacterium</i> strain carrying both pSIM356 and pSIM363				
Plant	Co-transformed	Marker only	Gene-of-interest only	Untransformed
Tobacco	50%	15%	19%	16%
Potato	22%	5%	15%	58%

Co-transformed: PCR-positive for both the GUS gene of pSIM356 and the NPT gene from LifeSupport

Untransformed: Plants are PCR-negative for PPO and NPTII

[0372]

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 124

<210> SEQ ID NO 1

<211> LENGTH: 416

<212> TYPE: DNA

<213> ORGANISM: *Solanum tuberosum*

<400> SEQUENCE: 1

-continued

gtttacagta ccatatatcc tgtcagaggt atagaggcat gactggcatg atcactaaat	60
tgatgccac agaggagact tataacctac aggggcacgt agttctagga cttgaaagt	120
actgaccgta gtccaactcg gtataaagcc tactcccaac taaatatatg aaatttatag	180
cataactgca gatgagctcg attctagagt aggtaccgag ctggaattcc ttactcctcc	240
acaaagccgt aactgaagcg acttctattt ttctcaacct tcggacctga cgatcaagaa	300
tctcaatagg tagttcttca taagtgaagc tatccttcat agctacactt tctaaaggta	360
cgatagattt tggatcaacc acacacactt cgtttacacc ggtatatatc ctgcca	416

<210> SEQ ID NO 2

<211> LENGTH: 824

<212> TYPE: DNA

<213> ORGANISM: Triticum sp.

<400> SEQUENCE: 2

tggcaggata tatgagtgtg taaacaacca taatcaggct gtaattatca agagaactaa	60
tgacaagaag cagagcttat caagtgttcc gtccagctgt aacatgggca caaaagcttg	120
cttgatgcat gtcctggttt tcaaagagca atgtattctc aggtacctgc acgtttgatc	180
ccctaccacg tacaagacga gcagaaagga catgtctgca gaaacttaga cacatccatt	240
gcagactcgt tccgaagcat caggagagta gtcagcaatg gtcactctgct gatgtaaatt	300
aattgattgt tggtaataca attttaacag caatatatat aatatatcaa tagtatattg	360
aactatgaaa gactgtaatc atatataaca gcatacaaat tgcctgtgga acaagaggag	420
ctcatcaagt gtttagttca gaaatagcta accaagaatg caatataata ggggtactga	480
gtcccttca aaattactaa cttcagaaat agctaacca gaatgcaatg gcattgcata	540
atttaacaa ctgtcagcac caatctctga ctgaaggcag tttaccatt cagaagagca	600
cacattttct gaacgacaac tctgagcggg gattgttgac agcagcaatt aatctggcct	660
caagatgggt tccaacaaca tagatcagat acagcactca agcaccat aatcagccag	720
tactgatctg gttaccactg caattgatta acagatgaac tgtgaaatta agatttaact	780
gacagtaata tataccagtt ggcaggatat atccctctgt aaac	824

<210> SEQ ID NO 3

<211> LENGTH: 2595

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Expression cassette for the IPT gene

<400> SEQUENCE: 3

ctgcagccaa agcacatact tatcgattta aatttcacg aagagattaa tatcgaataa	60
tcatatacat actttaata catacaaat tttaaatata tatatctggt atataattaa	120
ttttttaaag tcatgaagta tgtatcaaat acacatatgg aaaaaattaa ctattcataa	180
tttaaaaaat agaaaagata catctagtga aattaggtgc atgtatcaaa tacattagga	240
aaagggcata tatcttgatc tagataatta acgattttga tttatgtata atttccaaat	300
gaaggtttat atctacttca gaaataacaa tatactttta tcagaacatt caacaaagta	360
acaaccaact agagtgaata atacacattg ttctctaaac atacaaaatt gagaaaagaa	420

-continued

tctcaaaatt tagagaaaca aatctgaatt tctagaagaa aaaaaataatt atgcactttg	480
ctattgctcg aaaaaataat gaaagaaatt agactttttt aaaagatgtt agactagata	540
tactcaaaag ctatcaaagg agtaatatc ttcttacatt aagtatttta gttacagtcc	600
tgtaattaaa gacacatttt agattgtatc taaacttaaa tgtatctaga atacatatat	660
ttgaatgcat catatacatg tatccgacac accaattctc ataaaaagcg taatatccta	720
aactaattta tccttcaagt caacttaagc ccaatatata ttttcattctc taaaggccca	780
agtggcacaa aatgtcaggc ccaattacga agaaaagggc ttgtaaaacc ctaataaagt	840
ggcactggca gagcttacac tctcattcca tcaacaaaga aaccctaaaa gccgcagcgc	900
cactgatttc tctctccag gcgaagatgc agatcttcgt gaagacccta acggggaaga	960
cgatcaccct agaggttgag tcttccgaca ccatcgacaa tgtcaaagcc aagatccagg	1020
acaaggaagg gattccccc gaccagcagc gtttgatttt cgccggaaag cagcttgagg	1080
atggtcgtac tcttgccgac tacaacatcc agaaggagtc aactctccat ctctgtctcc	1140
gtctccgtgg tgggtggtac atggacctgc atctaatttt cggtccaact tgcacaggaa	1200
agacgacgac cgcgatagct cttgcccagc agacagggct tccagtcctt tcgcttgatc	1260
gggtccaatg ctgtctctaa ctatcaaccg gaagcggacg accaacagtg gaagaactga	1320
aaggaacgac gcgtctctac cttgatgacg ggcctctggt ggagggtatc atcgacagca	1380
agcaagctca tcataggctg atcgaggagg tgtataatca tgaggccaac ggcgggctta	1440
ttcttgaggg aggatccacc tcggtgtctc actgcatggc gcgaaacagc tattggagtg	1500
cagattttcg ttggcatatt attcgcacaa agttaccga ccaagagacc ttcataaaag	1560
cgcccaaggc cagagttaag cagatgttgc accccgctgc aggccattct attattcaag	1620
agttggttta tctttggaat gaacctcggc tgaggcccat tctgaaagag atcgatggat	1680
atcgatatgc catgttgttt gctagccaga accagatcac ggcagatatg ctattgcagc	1740
ttgacgcaaa tatggaaggt aagttgatta atgggatcgc tcaggagtat ttcattccatg	1800
cgcgccaaca ggaacagaaa ttcccccaag ttaacgcagc cgctttcgac ggattcgaag	1860
gtcatccggt cggaatgtat taggttacgc cagccctcgc tcgcacctgt cttcatctgg	1920
ataagatggt cgtaattggt tttggctttg tcctgtttgt gcagggcggc aaatacttcc	1980
gacaatccat cgtgtcttca aactttatgc tgggtgaacaa gtcttagttt ccacgaaagt	2040
attatgttaa attttaaaat ttcgatgtat aatgtggcta taattgtaaa aataaactat	2100
cgtaagtgtg cgtgttatgt ataattgtc taaatgttta atatatatca tagaacgcaa	2160
taaatattaa atatagcgct tttatgaaat ataaatacat cattacaagt gttttatatt	2220
tcgggtggac tagtttttaa tgtttagcaa atgtcctatc agttttctct tttgtcgaa	2280
cggtaattta gagttttttt tgctatatgg attttcgttt ttgatgtatg tgacaaccct	2340
cgggattgtt gatatttttc aaaactaaga gtttttgctt attgttctcg tctatttttg	2400
atatcaatct tagttttata tcttttctag ttctctacgt gttaaatgtt caacacacta	2460
gcaatttggc tgcagcgtat ggattatgga actatcaagt ctgtgggatc gataaatatg	2520
cttctcagga atttgagatt ttacagtcct tatgtctatt gggttgagta taatatagta	2580
aaaaaatagg aattc	2595

-continued

<211> LENGTH: 9323
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: pSIM11
nucleotide sequence

<400> SEQUENCE: 4

agctttggca ggatatatac cgggtgtaaac gaagtgtgtg tggttgatcc aaaatctatc 60
gtacctttag aaagtgtagc tatgaaggat agtctcactt atgaagaact acctattgag 120
attcttgatc gtcaggctcg aaggttgaga aaaatagaag tcgcttcagt tacggctttg 180
tggaggagta agggtagccta ctctagaatc gagctcatcg ttatgctata aatttcatat 240
atttagttgg gagtaggctt tataccgagt tggactacgg tcagtcactt tcaagtccta 300
gaactacgtg cccctgtagg ttataagtct cctctgtggg catcaattta gtgatcatgc 360
cagtcatgcc tctatacctc tgacaggata tatggtactg taaacactag ttgtgaataa 420
gtcgcgtgtg atgtttgttt gagatctcta agagaaaaga gcgtttatta gaataacgga 480
tatttaaaag ggcgtgaaaa ggtttatccg ttcgtccatt tgtatgtggt cacctatctc 540
gagcatgcc accacagggt tcccctcggg atcaaagtac tttgatccaa cccctccgct 600
gctatagtgc agtcggcttc tgacgttcag tgcagccgtc ttctgaaaac gacatgtcgc 660
acaagtccta agttacgcga cagggtgccc cctgcccctt ttctggcgt tttctgtgcg 720
cgtgttttag tcgcataaag tagaatactt gcgactagaa ccggagacat tacgccatga 780
acaagagcgc cgcgcgtggc ctgctgggct atgcccgcgt cagcaccgac gaccaggact 840
tgaccaacca acggggccga ctgcacgcgg ccggctgcac caagctgttt tccgagaaga 900
tcaccggcac caggcgcgac cgcccgagc tggccaggat gcttgaccac ctacgccctg 960
gcgacgttgt gacagtgaac aggcctagacc gcctggcccg cagcaccgcg gacctactgg 1020
acattgccga gcgcataccg gaggcggcgg ccggcctgcg tagcctggca gagccgtggg 1080
ccgacaccac cagcggcgcc ggccgcatgg tgttgaccgt gttcgccggc attgccgagt 1140
tcgagcgttc cctaatacgc gaccgcaccc ggagcgggcg cgaggccgccc aaggcccgag 1200
gcgtgaagtt tggcccccgc cctaccccca ccccggcaca gatcgcgcac gcccgcgagc 1260
tgatcgacca ggaaggccgc accgtgaaag aggcggctgc actgcttggc gtgcacgct 1320
cgacctgta ccgcgcactt gagcgacgag aggaagtga cccaccgag gccaggcggc 1380
gcggtgcctt ccgtgaggac gcattgacg aggcgcagc cctggcggcc gccgagaatg 1440
aacgccaaga ggaacaagca tgaaccgca ccaggacggc caggacgaac cgtttttcat 1500
taccgaagag atcgaggcgg agatgatcgc ggccgggtac gtgttcgagc cgcccgcgca 1560
cgtctcaacc gtgcggctgc atgaaatcct ggccgggttg tctgatgcca agctggcggc 1620
ctggccggcc agcttgccg ctgaagaaac cgagcgccgc cgtctaaaaa ggtgatgtgt 1680
atttgagtaa aacagcttgc gtcattgcgt cgctgcgtat atgatgcgat gagtaaataa 1740
acaaatacgc aagggaagc catgaaggt atcgctgtac ttaaccagaa aggcgggtca 1800
ggcaagacga ccatcgcaac ccatctagcc cgcccccctg aactcgccgg gcccgatgtt 1860
ctgttagtcg attccgatcc ccagggcagt gcccgcgatt gggcgccgct gcgggaagat 1920
caaccgctaa ccgttgtcgg catcgaccgc ccgacgattg acccgacgt gaaggccatc 1980
ggccggcgcg acttcgtagt gatcgacgga gcgccccagg cggcggaact ggctgtgtcc 2040

-continued

gcgatcaagg	cagccgactt	cgtgctgatt	ccggtgcagc	caagccctta	cgacatatgg	2100
gccaccgccc	acctgggtgga	gctgggttaag	cagcgcattg	aggtcacgga	tggaaggcta	2160
caagcggcct	ttgtcgtgtc	gcgggcgac	aaaggcacgc	gcacgcgcgg	tgagggtgcc	2220
gaggcgctgg	ccgggtacga	gctgcccatt	cttgagtccc	gtatcacgca	gcgcgtgagc	2280
taccagggca	ctgccgccgc	cggcacaaac	gttcttgaat	cagaacccga	gggcgacgct	2340
gcccgcgagg	tccaggcgct	ggccgctgaa	attaaatcaa	aactcatttg	agttaatgag	2400
gtaagagaa	aatgagcaaa	agcacaacaa	cgctaagtgc	cggccgtccg	agcgcacgca	2460
gcagcaaggc	tgcaacgttg	gccagcctgg	cagacacgcc	agccatgaag	cgggtcaact	2520
ttcagtgcc	ggcggaggat	cacaccaagc	tgaagatgta	cgcggtacgc	caaggcaaga	2580
ccattaccga	gctgctatct	gaatacatcg	cgcagctacc	agagtaaatg	agcaaatgaa	2640
taaatgagta	gatgaatttt	agcggctaaa	ggaggcgcca	tggaatatca	agaacaacca	2700
ggcaccgacg	ccgtggaatg	ccccatgtgt	ggaggaacgg	gcggttgccc	aggcgtaagc	2760
ggctgggttg	tctgccggcc	ctgcaatggc	actggaaccc	ccaagcccga	ggaatcgccg	2820
tgacggtcgc	aaaccatccg	gcccgggtaca	aatcggcgcg	gcgctgggtg	atgacctggt	2880
ggagaagttg	aaggccgcgc	aggccgcccc	gcggcaacgc	atcgaggcag	aagcacgccc	2940
cgggtgaatcg	tggcaagcgg	ccgctgatcg	aatccgcaaa	gaatcccggc	aaccgcccgg	3000
agccggtgcg	ccgtcgatta	ggaagccgcc	caagggcgac	gagcaaccag	attttttcgt	3060
tccgatgctc	tatgacgtgg	gcaccgcgca	tagtcgcagc	atcatggacg	tgcccgtttt	3120
ccgtctgtcg	aagcgtgacc	gacgagctgg	cgaggtgac	cgctacgagc	ttccagacgg	3180
gcacgtagag	gtttccgcag	ggccggccgg	catggccagt	gtgtgggatt	acgacctggt	3240
actgatggcg	gtttcccatc	taaccgaatc	catgaaccga	taccgggaag	ggaagggaga	3300
caagcccggc	cgcgtgtttc	gtccacacgt	tgcggacgta	ctcaagtctc	gccggcgagc	3360
cgatggcgga	aagcagaaag	acgacctggt	agaaacctgc	attcgggtta	acaccacgca	3420
cgttgccatg	cagcgtacga	agaaggccaa	gaacggccgc	ctggtgacgg	tatccgaggg	3480
tgaagccttg	attagccgct	acaagatcgt	aaagagcgaa	accgggcggc	cggagtacat	3540
cgagatcgag	ctagctgatt	ggatgtaccg	cgagatcaca	gaaggcaaga	acccgacgct	3600
gctgacgggt	caccccgatt	actttttgat	cgatcccggc	atcggccggt	ttctctaccg	3660
cctggcacgc	cgcgcgcgag	gcaaggcaga	agccagatgg	ttgttcaaga	cgatctacga	3720
acgcagtggc	agcgcgggag	agttcaagaa	gttctgtttc	accgtgcgca	agctgatcgg	3780
gtcaaatgac	ctgccggagt	acgatttgaa	ggaggaggcg	gggcaggctg	gcccgatcct	3840
agtcatgcgc	taccgcaacc	tgatcgaggg	cgaagcatcc	gccggttcct	aatgtacgga	3900
gcagatgcta	gggcaaatg	ccctagcagg	ggaaaaaggt	cgaaaaggtc	tctttcctgt	3960
ggatagcacg	tacattggga	acccaaagcc	gtacattggg	aaccggaacc	cgtacattgg	4020
gaacccaaag	ccgtacattg	ggaaccggtc	acacatgtaa	gtgactgata	taaaagagaa	4080
aaaaggcgat	ttttccgcct	aaaactcttt	aaaacttatt	aaaactctta	aaaccgcct	4140
ggcctgtgca	taactgtctg	gccagcgcac	agccgaagag	ctgcaaaaag	cgcctaccct	4200
tcggtcgctg	cgtcccttac	gccccgccgc	ttcgcgtcgg	cctatcgccg	ccgctggccg	4260
ctcaaaaatg	gctggcctac	ggccaggcaa	tctaccaggg	cgcggacaag	ccgcgcgctc	4320

-continued

gccactcgac	cgccggcgcc	cacatcaagg	caccctgcct	cgcgcggttc	ggtgatgacg	4380
gtgaaaacct	ctgacacatg	cagctcccgg	agacggtcac	agcttgtctg	taagcggatg	4440
ccgggagcag	acaagcccgt	cagggcgcg	cagcgggtgt	tggcgggtgt	cggggcgcag	4500
ccatgaccca	gtcacgtagc	gatacgggag	tgtatactgg	cttaactatg	cggcatcaga	4560
gcagattgta	ctgagagtgc	accatatgcy	gtgtgaaata	ccgcacagat	gcgtaaggag	4620
aaaataccgc	atcaggcgct	cttccgcttc	ctcgtcact	gactcgctgc	gctcggctcg	4680
tcggctgcgg	cgagcggat	cagctcactc	aaaggcggt	atacggttat	ccacagaatc	4740
aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	4800
aaaggccgcy	ttgctggcgt	ttttccatag	gctccgcccc	cctgacgagc	atcacaaaaa	4860
tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	taaagatacc	aggcgtttcc	4920
ccctggaagc	tccctcgtgc	gctctcctgt	tccgaccctg	ccgcttaccg	gatacctgtc	4980
cgcccttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	ggtatctcag	5040
ttcggtgtag	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccga	5100
ccgctgcgcc	ttatccggta	actatcgtct	tgagtccaac	ccggtaaagc	acgacttatt	5160
gccactggca	gcagccactg	gtaacaggat	tagcagagcg	aggatatgtg	gcggtgctac	5220
agagttcttg	aagtgggtgc	ctaactacgg	ctacactaga	aggacagtat	ttggtatctg	5280
cgctctgctg	aagccagtta	ccttcggaaa	aagagttggt	agctcttgat	ccggcaaaaca	5340
aaccaccgct	ggtagcgggt	gtttttttgt	ttgcaagcag	cagattacgc	gcagaaaaaa	5400
aggatctcaa	gaagatcctt	tgatcttttc	tacggggtct	gacgctcagt	ggaacgaaaa	5460
ctcacgttaa	gggatttttg	tcatgcattc	taggtactaa	aacaattcat	ccagtaaaat	5520
ataatatatt	ttttctccc	aatcaggctt	gatccccagt	aagtcaaaaa	atagctcgac	5580
atactgttct	tccccgatat	cctccctgat	cgaccggacg	cagaaggcaa	tgtcatacca	5640
ctgtccgcc	ctgccgcttc	tcccgaagtc	aataaagcca	cttactttgc	catctttcac	5700
aaagatgttg	ctgtctccca	ggtcgccgtg	ggaaaagaca	agttcctctt	cgggcctttc	5760
cgtctttaa	aaatcataca	gctcgcgcgg	atctttaaat	ggagtgtctt	cttcccagtt	5820
ttcgcaatcc	acatcgccca	gatcgttatt	cagtaagtaa	tccaattcgg	ctaagcggct	5880
gtctaagcta	ttcgtatagg	gacaatccga	tatgtcgatg	gagtgaagaa	gcctgatgca	5940
ctccgcatac	agctcgataa	tcttttcagg	gctttgttca	tcttcatact	cttccgagca	6000
aaggacgcca	tcggcctcac	tcatgagcag	attgctccag	ccatcatgcc	gttcaaagtg	6060
caggaccttt	ggaacaggca	gctttccttc	cagccatagc	atcatgtcct	tttcccgctc	6120
cacatcatag	gtgggtccct	tataccggct	gtccgtcatt	tttaaatata	ggttttcatt	6180
ttctcccacc	agcttatata	ccttagcagg	agacattcct	tccgtatctt	ttacgcagcg	6240
gtatttttcg	atcagttttt	tcaattccgg	tgatattctc	attttagcca	tttattattt	6300
ccttcctctt	ttctacagta	tttaaagata	ccccagaag	ctaattataa	caagacgaac	6360
tccaattcac	tgctccttgc	attctaaaac	cttaaatacc	agaaaacagc	tttttcaaag	6420
ttgttttcaa	agttggcgta	taacatagta	tcgacggagc	cgattttgaa	accgcggatc	6480
ctgcagccaa	agcacatact	tatcgattta	aatttcacgc	aagagattaa	tatcgataaa	6540
tcataatacat	actttaaaata	cataacaaat	tttaaataca	tatatctggt	atataattaa	6600

-continued

ttttttaaag	tcatgaagta	tgtatcaaat	acacatatgg	aaaaaattaa	ctattcataa	6660
tttaaaaaat	agaaaagata	catctagtga	aattagggtgc	atgtatcaaa	tacattagga	6720
aaagggcata	tatcttgatc	tagataatta	acgattttga	tttatgtata	atttccaaat	6780
gaaggtttat	atctacttca	gaaataacaa	tatactttta	tcagaacatt	caacaaagta	6840
acaaccaact	agagtgaaaa	atacacattg	ttctctaacc	atacaaaatt	gagaaaagaa	6900
tctcaaaatt	tagagaaaca	aatctgaatt	tctagaagaa	aaaaataatt	atgcactttg	6960
ctattgctcg	aaaaataaat	gaaagaaatt	agactttttt	aaaagatgtt	agactagata	7020
tactcaaaag	ctatcaaagg	agtaatatcc	ttcttacatt	aagtatttta	gttacagtcc	7080
tgtaattaaa	gacacatttt	agattgtatc	taaacttaaa	tgtatctaga	atacatatat	7140
ttgaatgcat	catatacatg	tatccgacac	accaattctc	ataaaaagcg	taatatcccta	7200
aactaattta	tccttcaagt	caacttaagc	ccaatataca	ttttcatctc	taaaggccca	7260
agtggcacia	aatgtcaggc	ccaattacga	agaaaagggc	ttgtaaacc	ctaataaagt	7320
ggcactggca	gagcttacac	tctcattcca	tcaacaaaga	aaccctaaaa	gccgcagcgc	7380
cactgatttc	tctcctccag	gcgaagatgc	agatcttcgt	gaagacccta	acggggaaga	7440
cgatcaccct	agaggttgag	tcttccgaca	ccatcgacaa	tgtcaaagcc	aagatccagg	7500
acaaggaagg	gattccccca	gaccagcagc	gtttgatttt	cgccggaaag	cagcttgagg	7560
atggtcgtac	tcttgccgac	tacaacatcc	agaaggagtc	aactctccat	ctcgtgctcc	7620
gtctccgtgg	tggtggatcc	atggacctgc	atctaatttt	cggccaact	tgacacaggaa	7680
agacgcagac	cgcgatagct	cttgcccagc	agacagggct	tccagtcctt	tcgcttgatc	7740
gggtccaatg	ctgtcctcaa	ctatcaaccg	gaagcggacg	accaacagtg	gaagaactga	7800
aaggaacgac	gcgtctctac	cttgatgatc	ggcctctggt	ggagggtatc	atcgcagcca	7860
agcaagctca	tcataggtcg	atcgaggagg	tgtataatca	tgaggccaac	ggcgggctta	7920
ttcttgaggg	aggatccacc	tcggtgctca	actgcatggc	gcgaacagc	tattggagtg	7980
cagattttcg	ttggcatatt	attcgccaca	agttaccgca	ccaagagacc	ttcatgaaag	8040
cggccaaggc	cagagttaag	cagatgttgc	accccgctgc	aggccattct	attattcaag	8100
agttggttta	tctttggaat	gaacctcggc	tgaggcccat	tctgaaagag	atcgatggat	8160
atcgatatgc	catgttggtt	gctagccaga	accagatcac	ggcagatatg	ctattgcagc	8220
ttgacgcaaa	tatggaaggt	aagttgatta	atgggatcgc	tcaggagtat	ttcatccatg	8280
cgcgccaaca	ggaacagaaa	ttccccaaag	ttaacgcagc	cgctttcgac	ggattcgaag	8340
gtcatccggt	cggaatgtat	taggttacgc	cagccctgcg	tcgcacctgt	cttcatctgg	8400
ataagatggt	cgtaattggt	tttggttttg	tcctgttggtg	gcagggcggc	aaatacttcc	8460
gacaatccat	cgtgtcttca	aactttatgc	tggtgaacaa	gtcttagttt	ccaagaaagt	8520
attatgttaa	atttttaaat	ttcgatgtat	aatgtggcta	taattgtaaa	aataaactat	8580
cgtaagtgtg	cgtgttatgt	ataattttgc	taaatgttta	atatatatca	tagaacgcaa	8640
taaatattaa	atatagcgct	tttatgaaat	ataaatacat	cattacaagt	tgtttatatt	8700
tcgggtggac	tagttttttta	tgtttagcaa	atgtcctatc	agttttctct	ttttgtcgaa	8760
cggtaattta	gagttttttt	tgctatatgg	attttcgttt	ttgatgtatg	tgacaaccct	8820
cgggattggt	gattttatttc	aaaactaaga	gtttttgctt	attgttctcg	tctatttttg	8880

-continued

```

atatcaatct tagttttata tcttttctag ttctctacgt gttaaagtgt caacacacta 8940
gcaatttggc tgcagcgtat ggattatgga actatcaagt ctgtggggtc gataaatatg 9000
cttctcagga atttgagatt ttacagtctt tatgctcatt gggttgagta taatatagta 9060
aaaaaatagg aattctatcc gcggtgatca caggcagcaa cgctctgtca tcgttacaat 9120
caacatgcta ccctccgcga gatcatccgt gtttcaaacc cggcagctta gttgccgttc 9180
ttccgaatag catcggtaac atgagcaaag tctgccgcct tacaacggct ctcccgtga 9240
cgccgtcccc gactgatggg ctgcctgtat cgagtgggtga ttttgtgccg agctgccggg 9300
cggggagctg ttggctggct gga 9323

```

```

<210> SEQ ID NO 5
<211> LENGTH: 546
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

```

```

<400> SEQUENCE: 5

```

```

atgagaaatt tattccccat attgatgcta atcaccaatt tggcactcaa caacgataac 60
aacaacaaca acaacaacaa caataattat aatctcatac acgcaacgtg tagggagacc 120
ccatattact ccctatgtct caccacccta caatccggtc cacgtagtaa cgagggtgag 180
ggtggtgatg ccatcaccac cctaggcctc atcatggtgg acgcggtgaa atcaaaagtcc 240
atagaaataa tggaaaaaat aaaagagcta gagaaatcga accctgagtg gcgggccccca 300
cttagccagt gttacgtggc gtataatgcc gtcctacgag ccgatgtaac ggtagccgtt 360
gaagccttaa agaagggtgc ccccaaatTT gctgaagatg gtatggatga tgttgttgct 420
gaagcacaaa cttgtgagta tagttttaat tattataata aattggattt tccaatttct 480
aatttgagta gggaaataat tgaactatca aaagttgcta aatccataat tagaatgtta 540
ttatga 546

```

```

<210> SEQ ID NO 6
<211> LENGTH: 658
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

```

```

<400> SEQUENCE: 6

```

```

gaaccatgca tctcaatctt aatactaaaa aatgcaacaa aattctagtg gagggaccag 60
taccagtaca ttagatatta tcttttatta ctataataat attttaatta acacgagaca 120
taggaatgtc aagtggtagc ggtaggaggg agttggttca gttttttaga tactaggaga 180
cagaaccgga gggggccatt gcaaggccca agttgaagtc cagccgtgaa tcaacaaaga 240
gagggcccat aatactgtcg atgagcattt ccctataata cagtgtccac agttgccttc 300
cgctaaggga tagccacccg ctattctctt gacacgtgtc actgaaacct gctacaata 360
aggcaggcac ctctctatct tcacactcac tcaactcac agctcaacaa gtggttaactt 420
ttactcatct cctccaatta tttctgattt catgcatgtt tccctacatt ctattatgaa 480
tcgtgttatg gtgtataaac gttgtttcat atctcatctc atctattctg attttgattc 540
tcttgctac tgaatttgac cctactgtaa tcggtgataa atgtgaatgc ttcctcttct 600
tcttcttctt ctacagaaatc aattttctgt ttgtttttgt tcactctgtg cttggtag 658

```

-continued

```

<210> SEQ ID NO 7
<211> LENGTH: 355
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 7

ttttaatggt tagcaaatgt cctatcagtt ttctcttttt gtcgaacggg aatttagagt    60
tttttttgct atatggattt tcggttttga tgtatgtgac aacctcggg attgttgatt    120
tatttcaaaa ctaagagttt ttgcttattg ttctcgtcta ttttgatat caatcttagt    180
tttatatcct ttctagtctt ctacgtgtta aatgttcaac acactagcaa ttgggctgca    240
gcgtatggat tatggaacta tcaagtctgt gggatcgata aatatgcttc tcaggaattt    300
gagattttac agtctttatg ctcatgggt tgagtataat atagtaaaaa aatag        355

```

```

<210> SEQ ID NO 8
<211> LENGTH: 179
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 8

accttatctt actaccactt tccactctcc aatccccata ctctctgctc caatcttcat    60
tttgcttcgt gaattcatct tcatcgaatt tctcgacgct tcttcgctaa tttcctcgtt    120
acttcactaa aaatcgacgt ttctagctga acttgagtga attaagccag tgggaggat    179

```

```

<210> SEQ ID NO 9
<211> LENGTH: 569
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 9

gttagaaaac ttctctatct ttggtttttg totgtttaga ttctcgaatt agctaatacag    60
gtgctgttat agcccttaat tttagatttt ttttcggttg ttttgatgga aaaggcctaa    120
aatttgagtt tttttacggt ggtttgatgg aaaaggccta caattggagt tttccccgtt    180
gttttgatga aaaagccctt agtttgagat tttttttctg tcgattcgat tctaaagggt    240
taaaattaga gtttttcatat ttgtttgatg aaaaaggcct taaatttgag tttttccggt    300
tgatttgatg aaaaagccct agaatttggt ttttttcgtc ggtttgattc tgaaggccta    360
aaatttgagt ttctccggct gttttgatga aaaagcccta aatttgagtt tctccggctg    420
ttttgatgaa aaagccctaa atttgagttt tttccccgtg ttttagattg ttgggtttta    480
attctcgaat cagctaatac gggagtgtga aaagccctaa aatttgagtt tttttcgttg    540
ttctgattgt tgtttttatg aatttgacg          569

```

```

<210> SEQ ID NO 10
<211> LENGTH: 1738
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Expression
cassette for a sense and antisense copy of the
leader associated with the R1 gene

```

```

<400> SEQUENCE: 10

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg    60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac    120

```

-continued

```

gagacatagg aatgtcaagt ggtacggta ggaggaggtt ggttcagttt tttagatact 180
aggagacaga accggagggg cccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa 240
caaagagagg gcccataata ctgtcgtatg gcatttcctt ataatacagt gtccacagtt 300
gccttccgct aagggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta 360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat 420
ccaccttatt tcactaccac tttccactct ccaatcccca tactctctgc tccaatcttc 480
atthtgcttc gtgaattcat cttcatcgaa tttctcgacg cttcttcgct aatttcctcg 540
ttacttcact agaaatcgac gtttctagct gaacttgagt gaattaagcc agtgggagga 600
tgaattcaag gttagaaatc ttctctatth ttggtttttg tctgtttaga ttctcgaatt 660
agctaatacg gtgctgttat agcccttaat tttgagtttt ttttcggttg ttttgatgga 720
aaaggcctaa aatttgagtt tttttacgth ggthtgatgg aaaaggccta caattggagt 780
tttccccgth gttthgatga aaaagccctt agthtgagat tttttttctg tctgattcgat 840
tctaaaggth taaaattaga gtttttacct ttgthtgatg aaaaaggcct taaatttgag 900
tttttccgth tgatttgatg aaaaagccctt agaatttgth ttttttctgth gttthgattc 960
tgaaggccta aaatttgagt ttctccgctt gttthgatga aaaagccta aatttgagth 1020
tctccgctg ttttgatgaa aaagccctaa atthgagtht tttccccgth ttttagattg 1080
tttgthttta attctcgaat cagctaatac gggagtgthg aaagccctaa aatttgagth 1140
tttttctgth ttctgattgt tgtttttatg aatttgacga tggatatcat cctccactg 1200
gcttaattca ctcaagttca gctagaaacg tctgatttcta gtgaagtaac gaggaaatta 1260
gcgaagaagc gtcgagaaat tctgatgaag tgaattcacg aagcaaatg aagattggag 1320
cagagagtat ggggattgga gagtggaaag tggtagtgaa ataaggtaag cttttgattt 1380
taatgtthg caaatgtctt atcagthttt tctttttgth gaacggtaat ttagagthtt 1440
ttthgctata tggatthttc tttthgatgt atgtgacaac cctcgggatt gthgatttat 1500
ttcaaaacta agagthtttg cttattgttc tctctattht tggatatcaa tcttagthtt 1560
atatctthtt tagthtctta cgtgttaaat gttcaacaca ctacgaattt ggtgacgth 1620
tatggattat ggaactatca agtctgtggg atcgataaat atgcttctca ggaatttgag 1680
atthttacag ttttatgctc attgggttga gtataatata gtaaaaaaat agtctaga 1738

```

<210> SEQ ID NO 11

<211> LENGTH: 237

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
spacer sequence

<400> SEQUENCE: 11

```

gtaactttta ctcatctcct ccaattatth ctgatttcat gcatgtttcc ctacattcta 60
ttatgaatcg tgttatggth tataaacgth gttcatatc tcactctatc tattctgatt 120
ttgattctct tgccactgga atthgacctt actgtaatcg gtgataaatg tgaatgcttc 180
ctcttcttct tcttcttctc agaaatcaat ttctgttttg tttthgttca tctgtag 237

```

<210> SEQ ID NO 12

-continued

```

<211> LENGTH: 1406
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative
        expression cassette for a sense and antisense
        coopy of the leader associated with the R1 gene

<400> SEQUENCE: 12

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg      60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac      120
gagacatagg aatgtcaagt ggtagcggta ggaggagggt ggttcagttt tttagatact      180
aggagacaga accggagggg cccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa      240
caaagagagg gcccataata ctgtcgtatg gcatttccct ataatacagt gtccacagtt      300
gccttccgct aagggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta      360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat      420
ccaccttatt tcactaccac ttccactctt ccaatcccca tactctctgc tccaatcttc      480
atthtgcttc gtgaattcat cttcatcgaa tttctcgacg cttcttcgct aatttcctcg      540
ttacttcact agaaatcgac gtttctagct gaacttgagt gaattaagcc agtgggagga      600
tgaattcgtg gtaactttta ctcatctcct ccaattattt ctgatttcat gcatgtttcc      660
ctacattcta ttatgaatcg tggtatggtg tataaacggt gtttcatatc tcattctcatc      720
tattctgatt ttgattctct tgcctactga atttgaccct actgtaatcg gtgataaatg      780
tgaatgcttc ctcttcttct tcttcttctc agaaatcaat ttctgttttg tttttgttca      840
tctgtagctt gatatcatcc tcccactggc ttaattcact caagttcagc tagaaacgtc      900
gatttttagt gaagtaacga ggaaattagc gaagaagcgt cgagaaattc gatgaagatg      960
aattcacgaa gcaaaatgaa gattggagca gagagtatgg ggattggaga gtggaaagtg     1020
gtagtgaaat aaggtaagct tttgatttta atgttttagca aatgtcctat cagttttctc     1080
tttttgctga acggtaattt agagtttttt ttgctatatg gattttcgtt tttgatgtat     1140
gtgacaaccc tcgggattgt tgatttattt caaaactaag agtttttgct tattgtttctc     1200
gtctatthtg gatatcaatc ttagttttat atcttttcta gttctctacg tgttaaatgt     1260
tcaacacact agcaatttgg ctgcagcgta tggattatgg aactatcaag tctgtgggat     1320
cgataaatat gcttctcagg aatttgagat ttacagctct ttatgctcat tgggttgagt     1380
ataatatagt aaaaaaatag tctaga                                           1406

<210> SEQ ID NO 13
<211> LENGTH: 686
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 13

gaaccatgca tctcaatctt aataactaaa aatgcaacaa aattctagtg gagggaccag      60
taccagtaca ttagatatta tcttttatta ctataataat attttaatta acacgagaca     120
taggaatgtc aagtggtagc ggtaggaggg agttgggttca gttttttaga tactaggaga     180
cagaaccgga ggggccattt gcaaggccca agttgaagtc cagccgtgaa tcaacaaaga     240
gagggcccat aatactgtcg atgagcattt ccctataata cagtgtccac agttgccttc     300

```

-continued

cgctaaggga tagccaccg ctattctctt gacacgtgtc actgaaacct gctacaaata	360
aggcaggcac ctctcatctc tcactcacac tcaactcacac agctcaacaa gtggttaactt	420
ttactcatct cctccaatta tttctgattt catgcatggt tccctacatt ctattatgaa	480
tcgtgttatg gtgtataaac gttgtttcat atctcatctc atctattctg attttgattc	540
tcttgccctac tgaatttgac cctactgtaa tcggtgataa atgtgaatgc ttcctcttct	600
tcttcttctt ctacagaaatc aatttctgtt ttgtttttgt tcactgttag ctgggtagat	660
tccccctttt gtagaccaca catcac	686

<210> SEQ ID NO 14

<211> LENGTH: 2046

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative expression cassette for a sense and antisense copy of the leader associated with the R1 gene

<400> SEQUENCE: 14

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg	60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac	120
gagacatagg aatgtcaagt ggtagcggta ggaggaggtt ggttcagttt tttagatact	180
aggagacaga accggagggg cccattgcaa ggccaagtt gaagtccagc cgtgaatcaa	240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt	300
gccttcgcgt aaggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta	360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat	420
cctcatattc tagttgtatg ttgttcagag aagaccacag atgtgatcat attctcattg	480
tatcagatct gtgaccatt acctgatacc tcccatgaag ttacctgtat gattatacgt	540
gatccaaagc catcacatca tgttcacctt cagctatttg aggagaagtg agaagtagga	600
attgcaatat gaggaataat aagaaaaact ttgtaaaagc taaattagct gggtagata	660
tagggagaaa tgtgtaaaca ttgtactata tatagtatat acacacgcat tatgtattgc	720
attatgcact gaataatacc gcagcatcaa agaaggaatt caaggtaga aatctctct	780
atttttggtt tttgtctgtt tagattctcg aattagctaa tcagggtctg ttatagccct	840
taattttgag ttttttttcg gttgttttga tggaaaaggc ctaaaatttg agttttttta	900
cgttggtttg atggaaaagg cctacaattg gagttttccc cgttggtttg atgaaaagc	960
ccctagtttg agattttttt tctgtogatt cgattctaaa ggtttaaaat tagagttttt	1020
acatttgttt gatgaaaag gccttaaatt tgagtttttc cgggtgattt gatgaaaag	1080
ccctagaatt tgtgtttttt cgtcggtttg attctgaagg cctaaaattt gagtttctcc	1140
ggctgttttg atgaaaagc cctaaatttg agtttctccg gctgttttga tgaaaagcc	1200
ctaaatttga gttttttccc cgtgttttag attgtttggt ttttaattctc gaatcagcta	1260
atcagggagt gtgaaaagcc ctaaaatttg agtttttttc gttgttctga ttgtgtttt	1320
tatgaatttg cagatggata tccttctttg atgctgcggt attattcagt gcataatgca	1380
atacataatg cgtgtgtata tactatatat agtacaatgt ttacacattt ctccctatat	1440
cataccagc taatttagct ttacaaaagt ttttcttatt attcctcata ttgcaattcc	1500

-continued

tacttctcac ttctcctcca atagctgaag gtgaacatga tgtgatggct ttggatcacg	1560
tataatcata caggtaactt catgggaggt atcaggtaag tggtcacaga tctgatacaa	1620
tgagaatatg atcacatctg tggctcttctc tgaacaacat acaactagaa tatgaaagct	1680
tttgatttta atgtttagca aatgtcctat cagttttctc tttttgtcga acggtaat	1740
agagtttttt ttgctatatg gattttcgtt tttgatgtat gtgacaaccc tcgggattgt	1800
tgatttattt caaaactaag agtttttctt tattgttctc gtctattttg gatatacaatc	1860
ttagttttat atcttttcta gttctctacg tgttaaatgt tcaacacact agcaatttgg	1920
ctgcagcgta tggattatgg aactatcaag tctgtgggat cgataaata gcttctcagg	1980
aatttgagat ttacagctct ttatgctcat tgggttgagt ataatatagt aaaaaaatag	2040
tctaga	2046

<210> SEQ ID NO 15

<211> LENGTH: 1714

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative expression cassette for a sense and antisense copy of the leader associated with the R1 gene

<400> SEQUENCE: 15

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg	60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac	120
gagacatagg aatgtcaagt ggtagcggta ggaggaggtt ggttcagttt ttagataact	180
aggagacaga accggagggg cccattgcaa ggccaagt gaagtccagc cgtgaatcaa	240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt	300
gccttcgctt aaggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta	360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat	420
cctcatattc tagttgatg ttgttcagag aagaccacag atgtgatcat attctcattg	480
tatcagatct gtgaccatt acctgatacc tccatgaag ttacctgtat gattatacgt	540
gatccaaagc catcacatca tgttcacctt cagctattgg aggagaagtg agaagtagga	600
attgcaatat gaggaataat aagaaaaact ttgtaaaagc taaattagct gggatatgata	660
tagggagaaa tgtgtaaaca ttgtactata tatagtatat acacacgcat tatgtattgc	720
attatgcact gaataatacc gcagcatcaa agaaggaatt cgtggtaact ttactcatc	780
tcctccaatt atttctgatt tcatgcatgt ttccctacat tctattatga atcgtgttat	840
ggtgtataaa cgttgtttca tatctcatct catctattct gattttgatt ctcttgcccta	900
ctgaatttga ccctactgta atcggtgata aatgtgaatg ctccctcttc ttcttcttct	960
tctcagaaat caatttctgt tttgtttttg ttcattctgta gcttgatata ctctttgat	1020
gctgcgggat tattcagtgc ataagcaat acataatgog tgtgtatata ctatatatag	1080
tacaatgttt acacatttct ccctatatca taccagcta atttagcttt taaaaagttt	1140
ttcttattat tcctcatatt gcaattccta cttctcactt ctctccaat agctgaaggt	1200
gaacatgatg tgatggcttt ggatcacgta taatcataca ggtaacttca tgggaggtat	1260
caggtaagtg gtcacagatc tgatacaatg agaatatgat cacatctgtg gtcttctctg	1320

-continued

```

aacaacatac aactagaata tgaaagcttt tgattttaat gtttagcaaa tgcctatca 1380
gttttctctt ttgtcgaac ggtaatttag agtttttttt gctatatgga ttttcgtttt 1440
tgatgtatgt gacaaccctc gggattgttg atttatttca aaactaagag tttttgctta 1500
ttgttctcgt ctattttgga tatcaatctt agttttatat cttttctagt tctctacgtg 1560
ttaaatgttc aacacactag caatttggct gcagcgtatg gattatggaa ctatcaagtc 1620
tgtgggatcg ataaatatgc ttctcaggaa ttgagattt tacagtcttt atgctcattg 1680
ggttgagtat aatatagtaa aaaaatagtc taga 1714

```

```

<210> SEQ ID NO 16
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

```

```

<400> SEQUENCE: 16

```

```

tcatattcta gttgtatgt gttcagagaa gaccacagat gtgatcatat tctcattgta 60
tcagatctgt gaccacttac ctgatacctc ccatgaagtt acctgtatga ttatacgtga 120
tccaaagcca tcacatcatg ttcaccttca gctattggag gagaagtga aagtaggaat 180
tgcaatatga ggaataataa gaaaaacttt gtaaaagcta aattagctgg gtatgatata 240
gggagaaatg tgtaaacatt gtactatata tagtatatac acacgcatta tgtattgcat 300
tatgcactga ataataccgc agcatcaaag aag 333

```

```

<210> SEQ ID NO 17
<211> LENGTH: 2046
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative
expression cassette for a sense and antisense copy
of the trailer associated with the R1 gene

```

```

<400> SEQUENCE: 17

```

```

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg 60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac 120
gagacatagg aatgtcaagt ggtagcggta ggaggagggt ggttcagttt tttagatact 180
aggagacaga accggagggg ccatttgcaa ggcccaagtt gaagtccagc cgtgaatcaa 240
caaagagagg gcccataata ctgtcgtatg gcatttccct ataatacagt gtccacagtt 300
gccttccgct aagggatagc cccccgctat tctcttgaca cgtgtcactg aaacctgcta 360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat 420
cctcatattc tagttgtatg ttgttcagag aagaccacag atgtgatcat attctcattg 480
tatcagatct gtgaccactt acctgatacc tccatgaag ttacctgtat gattatacgt 540
gatocaaagc catcacatca tgttcacctt cagctattgg aggagaagtg agaagtagga 600
attgcaatat gaggaataat aagaaaaact ttgtaaaagc taaattagct gggtatgata 660
tagggagaaa tgtgtaaaca ttgtactata tatagtatat acacacgcat tatgtattgc 720
attatgcact gaataatacc gcagcatcaa agaaggaatt caagggttaga aatcttctct 780
atttttgggt ttgtctgtt tagattctcg aattagctaa tcagggtctg ttatagccct 840
taattttgag ttttttttcg gttgttttga tggaaaaggc ctaaaatttg agttttttta 900

```

-continued

cgttggtttg atggaagc cctacaattg gagttttccc cgttggtttg atgaaaaagc	960
ccctagtttg agattttttt tctgtcgatt cgattctaaa ggtttaaaat tagagttttt	1020
acatttggtt gatgaaaaag gccttaaaatt tgagtttttc cggttgattt gatgaaaaag	1080
ccctagaatt tgtgtttttt cgtcgggttg attctgaagg cctaaaattt gagtttctcc	1140
ggctggtttg atgaaaaagc cctaaatttg agtttctccg gctgttttga tgaaaaagcc	1200
ctaaatttga gttttttccc cgtgttttag attgtttggt ttttaattctc gaatcagcta	1260
atcagggagt gtgaaaaagc ctaaaatttg agtttttttc gttgttctga ttgttggttt	1320
tatgaatttg cagatggata tccttctttg atgctgcggg attattcagt gcataatgca	1380
atacataatg cgtgtgtata tactatatat agtacaatgt ttacacattt ctccctatat	1440
catacccgag taatttagct ttacaaaagt ttttcttatt attcctcata ttgcaattcc	1500
tacttctcac ttctcctcca atagctgaag gtgaacatga tgtgatggct ttggatcacg	1560
tataatcata caggtaactt catgggaggt atcaggtaag tggtcacaga tctgatacaa	1620
tgagaatatg atcacatctg tggctctctc tgaacaacat acaactagaa tatgaaagct	1680
tttgatttta atgttttagc aatgtcctat cagttttctc tttttgtcga acggtaattt	1740
agagtttttt ttgctatatg gattttcgtt tttgatgtat gtgacaaccc tcgggattgt	1800
tgatttattt caaaactaag agtttttgct tattgttctc gtctattttg gatatcaatc	1860
ttagttttat atcttttcta gttctctacg tgttaaatgt tcaacacact agcaatttgg	1920
ctgcagcgta tggattatgg aactatcaag tctgtgggat cgataaatat gcttctcagg	1980
aatttgagat ttacacgtct ttatgctcat tgggttgagt ataatatagt aaaaaaatag	2040
tctaga	2046

<210> SEQ ID NO 18

<211> LENGTH: 1714

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative expression cassette for a sense and antisense copy of the trailer associated with the R1 gene

<400> SEQUENCE: 18

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caaaaaatt ctagtggagg	60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac	120
gagacatagg aatgtcaagt ggtagcggta ggaggaggtt gggtcagttt ttagataact	180
aggagacaga accggagggg cccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa	240
caaagagagg gcccataata ctgtcgtatg gcatttccct ataatacagt gtccacagtt	300
gccttccgct aagggatagc cccccgctat tctcttgaca cgtgtcactg aaacctgcta	360
caataaagc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat	420
cctcatattc tagttgtatg ttgttcagag aagaccacag atgtgatcat attctcattg	480
tatcagatct gtgaccactt acctgatacc tccatgaag ttacctgtat gattatacgt	540
gatccaaagc catcacatca tgttcacctt cagctatttg aggagaagtg agaagtagga	600
attgcaatat gaggaataat aagaaaaact ttgtaaaagc taaattagct gggatatgata	660
tagggagaaa tgtgtaaaca ttgtactata tatagtatat acacacgcat tatgtattgc	720

-continued

```

attatgcact gaataatacc gcagcatcaa agaaggaatt cgtggtaact ttactcactc 780
tcctccaatt atttctgatt tcatgcatgt ttccctacat tctattatga atcgtgttat 840
ggtgtataaa cgttgtttca tatctcatct catctattct gatatttgatt ctcttgccca 900
ctgaatttga ccctactgta atcggtgata aatgtgaatg ctccctcttc ttcttcttct 960
tctcagaaat caatttctgt ttgtttttg ttcatctgta gcttgatata ctctttgat 1020
gctgcggtat tattcagtgc ataatgcaat acataatgcg tgtgtatata ctatatatag 1080
tacaatgttt acacatttct ccctatatca taccagcta atttagcttt taaaaagttt 1140
ttcttattat tcctcatatt gcaattccta ctctcactt ctccccaat agctgaaggt 1200
gaacatgatg tgatggcttt ggatcacgta taatcataca ggtaacttca tgggaggtat 1260
caggtaagtg gtcacagatc tgatacaatg agaatatgat cacatctgtg gtcttctctg 1320
aacaacatac aactagaata tgaaagcttt tgattttaat gtttagcaa tgcctatca 1380
gttttctctt ttgtgcgaac ggtaatttag agtttttttt gctatatgga ttttcgtttt 1440
tgatgtatgt gacaaccctc gggattgttg atttatttca aaactaagag tttttgctta 1500
ttgttctcgt ctattttgga tatcaatctt agttttatat cttttctagt tctctacgtg 1560
ttaaatgttc aacacactag caatttggtc gcagcgtatg gattatggaa ctatcaagtc 1620
tgtgggatcg ataaatatgc ttctcaggaa ttgagattt tacagtcttt atgctcattg 1680
ggttgagtat aatatagtaa aaaaatagtc taga 1714

```

<210> SEQ ID NO 19

<211> LENGTH: 2322

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative expression cassette for a sense and antisense copy of the trailer associated with the R1 gene

<400> SEQUENCE: 19

```

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caaaaaaatt ctagtggagg 60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac 120
gagacatagg aatgtcaagt ggtagcggta ggaggaggtt ggttcagttt tttagatact 180
aggagacaga accggagggg ccatttgcaa ggcccaagtt gaagtccagc cgtgaatcaa 240
caaagagagg gcccataata ctgtcgtatg gcatttccct ataatacagt gtccacagtt 300
gccttccgct aagggatagc ccccgcgtat tctcttgaca cgtgtcactg aaacctgcta 360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caacaagtgg 420
taacttttac tcatctcctc caattatttc tgatttcatg catgtttccc tacattctat 480
tatgaatcgt gttatgggtg ataaacgttg tttcatatct catctcatct attctgattt 540
tgattctctt gcctactgaa ttgacccta ctgtaatcgg tgataaatgt gaatgcttcc 600
tcttcttctt ctcttcttca gaaatcaatt tctgttttgt tttgttcat ctgtagcttg 660
gtagattccc cttttttag accacacatc acggatcctc atattctagt tgtatgttgt 720
tcagagaaga ccacagatgt gatcatattc tcattgtatc agatctgtga ccacttacct 780
gatacctccc atgaagttag ctgtatgatt atacgtgatc caaagccatc acatcatgtt 840
caccttcagc tattggagga gaagtgaaga gtaggaattg caatatgagg aataataaga 900

```

-continued

```

aaaactttgt aaaagctaaa ttagctgggt atgatatagg gagaaatgtg taaacattgt    960
actatatata gtatatacac acgcattatg tattgcatta tgcactgaat aataccgcag    1020
catcaaagaa ggaattcaag gttagaaatc ttctctatct ttggtttttg tctgtttaga    1080
ttctcgaatt agctaatacag gtgctgttat agcccttaat tttagatttt ttttcgggtg    1140
ttttgatgga aaaggcctaa aatttgagtt tttttacggt ggtttgatgg aaaaggccta    1200
caattggagt tttccccgtt gttttgatga aaaagccctt agtttgagat tttttttctg    1260
tcgattcgat tctaaagggt taaaattaga gtttttacat ttgtttgatg aaaaaggcct    1320
taaatttgag tttttccggt tgatttgatg aaaaagccct agaatttggt ttttttcgtc    1380
ggtttgattc tgaaggccta aaatttgagt ttctccggct gttttgatga aaaagcccta    1440
aatttgagtt tctccggctg ttttgatgaa aaagccctaa atttgagttt tttccccgtg    1500
tttttagatt ttgtgtttta attctcgaat cagctaatac gggagtgatg aaagccctaa    1560
aatttgagtt tttttcgttg ttctgattgt tgtttttatg aatttgcaga tggatatcct    1620
tctttgatgc tgcgggtatta ttcagtgcac aatgcaatac ataatgcgtg tgtatatact    1680
atatatagta caatgtttac acatttctcc ctatatcata cccagctaata ttagctttta    1740
caaagttttt ctattatttc ctcatattgc aattcctact tctcacttct cctccaatag    1800
ctgaaggatg acatgatgtg atggcttttg atcacgtata atcatacagg taacttcacg    1860
ggaggtatca ggtaagtggg cacagatctg atacaatgag aatatgatca catctgtggt    1920
cttctctgaa caacatacaa ctagaatatg aaagcttttg attttaatgt ttagcaaatg    1980
tcctatcagt tttctctttt tgtcgaacgg taatttagag ttttttttgc tatatggatt    2040
ttcgtttttg atgtatgtga caaccctcgg gattgttgat ttatttcaa actaagagtt    2100
tttgcttatt gttctcgtct attttggata tcaatcctag ttttatatct tttctagttc    2160
tctacgtgtt aaatgttcaa cacactagca atttggtctg agcgtatgga ttaatggaact    2220
atcaagtctg tgggatcgat aaatatgctt ctcaggaatt tgagatttta cagtctttat    2280
gctcattggg ttgagtataa tatagtaaaa aaatagtcta ga                        2322

```

<210> SEQ ID NO 20

<211> LENGTH: 1714

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative expression cassette for a sense and antisense copy of the trailer associated with the R1 gene

<400> SEQUENCE: 20

```

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg    60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac    120
gagacatagg aatgtcaagt ggtacgggta ggaggggagt gggtcagttt ttagataact    180
aggagacaga accggagggg cccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa    240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt    300
gccttcogct aagggatagc caccgctat tctcttgaca cgtgtcactg aaacctgcta    360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat    420
cctcatattc tagttgtatg ttgttcagag aagaccacag atgtgatcat attctcattg    480

```

-continued

tatcagatct gtgaccactt acctgatacc tcccatgaag ttacctgtat gattatacgt	540
gatccaaagc catcacatca tgttcacett cagctattgg aggagaagtg agaagtagga	600
attgcaatat gaggaataat aagaaaaact ttgtaaaagc taaattagct gggtagata	660
tagggagaaa tgtgtaaaca ttgtactata tatagtatat acacacgcat tatgtattgc	720
attatgcact gaataatacc gcagcatcaa agaaggaatt cgtggtaact tttactcatc	780
tcctccaatt atttctgatt tcatgcatgt ttccctacat tctattatga atcgtgttat	840
gggtataaaa cgttgtttca tatctcatct catctattct gattttgatt ctcttgccca	900
ctgaatttga ccctactgta atcggtgata aatgtgaatg cttcctcttc ttcttcttct	960
tctcagaaat caatttctgt tttgtttttg ttcactctgta gcttgatata cttctttgat	1020
gctgcgggat tattcagtgc ataatgcaat acataatgcg tgtgtatata ctatatatag	1080
tacaatgttt acacatttct ccctatatca taccagcta atttagcttt taaaaagttt	1140
ttcttattat tcctcatatt gcaattccta cttctcactt ctctccaat agctgaagg	1200
gaacatgatg tgatggcttt ggatcacgta taatcataca ggtaacttca tgggaggat	1260
caggtaagtg gtcacagatc tgatacaatg agaatatgat cacatctgtg gtcttctctg	1320
aacaacatac aactagaata tgaaagcttt tgattttaat gtttagcaaa tgcctatca	1380
gttttctctt tttgtcgaac ggtaatttag agtttttttt gctatatgga ttttcgtttt	1440
tgatgtatgt gacaaccctc gggattgttg atttatttca aaactaagag tttttgctta	1500
ttgttctcgt ctattttgga tatcaatctt agttttatat cttttctagt tctctacgtg	1560
ttaaagtgtc aacacactag caatttggtc gcagcgtatg gattatggaa ctatcaagtc	1620
tgtaggcatc ataaatatgc ttctcaggaa tttgagattt tacagtcttt atgctcattg	1680
ggttgagtat aatatagtaa aaaaatagtc taga	1714

<210> SEQ ID NO 21
 <211> LENGTH: 273
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 21

ttagagtgtg ggtaagtaat taagttaggg atttgtggga aatggacaaa tataagagag	60
tgagggggag tagtgaggga gattttcgtg cttttattga taaataaaaa aagggtgaca	120
tttaatttcc acaaggaggac gcaacacaac acacttaatt cctgtgtgtg aatcaataat	180
tgacttctcc aatcttcacg aataaaataa ttcacaatcc tcactctctt atcactctca	240
ttcgaaaagc tagatttgca tagagagcac aaa	273

<210> SEQ ID NO 22
 <211> LENGTH: 158
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 22

gagggggaag tgaatgaaaa ataacaaagg cacagtaagt agtttctctt tttatcatgt	60
gatgaaggta tataatgtat gtgtaaggag atgatgttat taccacataa taagagatga	120
agagtctcat tttctgctta aaaaaacaat tcactggc	158

<210> SEQ ID NO 23

-continued

<211> LENGTH: 1917
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Expression cassette for a sense and antisense copy of the leader associated with the L glucan phosphorylase gene

<400> SEQUENCE: 23

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg	60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac	120
gagacatagg aatgtcaagt ggtagcggta ggaggagggt ggttcagttt tttagatact	180
aggagacaga accggagggg ccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa	240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt	300
gccttcgctt aagggatagc caccgctat tctcttgaca cgtgtcactg aaacctgcta	360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat	420
ccgagtgtgg gtaagtaatt aagttaggga tttgtgggaa atggacaaat ataagagagt	480
gcaggggagt agtgcaggag attttcgtgc ttttattgat aaataaaaaa aggggtgacat	540
ttaatttcca caagaggacg caacacaaca cacttaattc ctgtgtgtga atcaataatt	600
gacttctcca atcttcatca ataaaaaat tcacaatcct cactctctta tcactctcat	660
tcgaaaagct agatttgcac agagagcaca gaattcaagg ttgaaatct tctctatttt	720
tggtttttgt ctgtttgatg tctcgaatta gctaatacagg tgctgttata gcccttaatt	780
ttgagttttt tttcggttgt tttgatggaa aaggcctaaa atttgagttt ttttacgttg	840
gtttgatgga aaaggcctac aattggagtt ttccccgttg ttttgatgaa aaagcccta	900
gtttgagatt tttttctgtg cgattcgatt ctaaaggttt aaaattagag tttttacatt	960
tgtttgatga aaaaggcctt aaatttgagt ttttcggtt gatttgatga aaaagcccta	1020
gaatttggtg tttttcgtcg gtttgattct gaaggcctaa aatttgagtt tctccggctg	1080
ttttgatgaa aaagccctaa atttgagttt ctccggctgt tttgatgaaa aagccctaaa	1140
tttgagtttt tttcccggtg ttttagattg ttggttttaa ttctcgaatc agctaatacag	1200
ggagtgatga aagccctaaa atttgagttt ttttcggtgt tctgattggt gtttttatga	1260
atttgacagat ggatatctgt gctctctatg caaatctagc ttttcgaatg agagtataa	1320
gagagtgagg atttggaatt attttattga tgaagattgg agaagtcaat tattgattca	1380
cacacaggaa ttaagtggtg tgtgttgctg cctcttgagg aaattaaatg tcaccctttt	1440
tttatttatc aataaaagca cgaaaatctc ctgcactact ccctgcact ctcttatatt	1500
tgccattttc ccacaaatcc ctaacttaat tacttaccac cactctaagc ttttgatttt	1560
aatgttttagc aaatgtccta tcagttttct ctttttgctg aacggtaatt tagagttttt	1620
tttgctatat ggattttcgt ttttgatgta tgtgacaacc ctgggattg ttgattttatt	1680
tcaaaactaa gagtttttgc ttattgttct cgtctatttt ggatatcaat cttagtttta	1740
tatcttttct agttctctac gtgttaaatg ttcaacacac tagcaatttg gctgcagcgt	1800
atggattatg gaactatcaa gtctgtggga tcgataaata tgcttctcag gaatttgaga	1860
ttttacagtc tttatgctca ttgggttgag tataatatag taaaaaata gtctaga	1917

<210> SEQ ID NO 24

-continued

```

<211> LENGTH: 1585
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative
        expression cassette for a sense and antisense copy of the leader
        associated with the L glucan phosphorylase gene

<400> SEQUENCE: 24

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg      60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac      120
gagacatagg aatgtcaagt ggtagcggta ggaggagggt gggttcagttt tttagatact      180
aggagacaga accggagggg ccatttgcaa ggcccaagtt gaagtccagc cgtgaatcaa      240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt      300
gccttcgctt aagggatagc caccgctat tctcttgaca cgtgtcactg aaacctgcta      360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat      420
ccgagtgtgg gtaagtaatt aagttaggga tttgtgggaa atggacaaat ataagagagt      480
gcaggggagt agtcaggag attttcgtgc ttttattgat aaataaaaaa agggtgacat      540
ttaatttcca caaggaggac caacacaaca cacttaattc ctgtgtgtga atcaataatt      600
gacttctcca atcttcatca ataaaaaat tcacaatcct cactctctta tcactctcat      660
tcgaaaagct agatttgcac agagagcaca gaattcgtgg taacttttac tcactctctc      720
caattatttc tgatttcacg catgtttccc tacattctat tatgaatcgt gttatggtgt      780
ataaacgttg tttcatatct catctcatct attctgattt tgattctctt gcctactgaa      840
tttgacccta ctgtaatcgg tgataaatgt gaatgccttc tcttcttctt cttcttctca      900
gaaatcaatt tctgttttgt tttgttcat ctgtagcttg atatctgtgc tctctatgca      960
aatctagctt ttcgaatgag agtgataaga gaggtaggat tgtgaattat tttattgatg      1020
aagattggag aagtcaatta ttgattcaca cacaggaatt aagtgtgttg tgttgcgtcc      1080
tcttgtggaa attaaatgac accctttttt tatttatcaa taaaagcacg aaaatctcct      1140
gcactactcc cctgcactct cttatatttg tccatttccc acaaatccct aacttaatta      1200
cttaccacac ctctaagctt ttgattttaa tgttttagcaa atgtcctatc agttttctct      1260
ttttgtcgaa cggtaattha gagttttttt tgctatatgg attttcgttt ttgatgtatg      1320
tgacaaccct cgggatttgt gatttatttc aaactaaga gtttttgctt attgttctcg      1380
tctatttttg atatcaatct tagttttata tcttttctag ttctctacgt gttaaatgtt      1440
caacacacta gcaatttggc tgcagcgtat ggattatgga actatcaagt ctgtgggatc      1500
gataaatatg cttctcagga atttgagatt ttacagtctt tatgctcatt gggttgagta      1560
taatatagta aaaaaatagt ctaga                                          1585

<210> SEQ ID NO 25
<211> LENGTH: 2193
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative
        expression cassette for a sense and antisense copy of the leader
        associated with the L glucan phosphorylase gene

<400> SEQUENCE: 25

```

-continued

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg	60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac	120
gagacatagg aatgtcaagt ggtagcggta ggaggaggtt ggttcagttt tttagatact	180
aggagacaga accggagggg cccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa	240
caaagagagg gcccataata ctgtcgtatg gcatttcctt ataatacagt gtccacagtt	300
gccttccgct aagggatagc cccccgtat tctcttgaca cgtgtcactg aaacctgcta	360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caacaagtgg	420
taacttttac tcatctcctc caattatttc tgatttcatt catgtttccc tacattctat	480
tatgaatcgt gttatgggtg ataaacgttg ttccatatct catctcatct attctgattt	540
tgattctctt gcctactgaa ttgacccta ctgtaatcgg tgataaatgt gaatgcttcc	600
tcttctctct cttcttctca gaaatcaatt tctgttttgt tttgttcat ctgtagcttg	660
gtagattccc cttttttag accacacatc acggatccga gtgtgggtaa gtaattaagt	720
tagggatttg tgggaaatgg acaaatataa gagagtgcag gggagtagtg caggagattt	780
tcgtgctttt attgataaat aaaaaagg tgacatttaa tttccacaag aggacgcaac	840
acaacacact taattcctgt gtgtgaatca ataattgact tctccaatct tcatcaataa	900
aataattcac aatcctcact ctcttatcac tctcattcga aaagctagat ttgcatagag	960
agcacagaat tcaaggttag aaatcttctc tatttttggt tttgtctgt ttagattctc	1020
gaattagcta atcaggtgct gttatagccc ttaattttga gtttttttc ggttgtttt	1080
atggaaaagg cctaaaaatt gagttttttt acgttggttt gatggaaaag gcctacaatt	1140
ggagttttcc ccgttgtttt gatgaaaag cccctagttt gagatttttt ttctgtcgat	1200
tcgattctaa aggtttaaaa ttagagtttt tacatttggt tgatgaaaaa ggccttaaat	1260
ttgagttttt ccggttgatt tgatgaaaaa gccctagaat ttgtgttttt tcgtcggttt	1320
gattctgaag gcctaaaaatt tgagtttctc cggctgtttt gatgaaaaag ccctaaattt	1380
gagtttctcc ggctgttttg atgaaaaagc cctaaatttg agtttttcc ccgtgtttta	1440
gattgttttg ttttaattct cgaatcagct aatcaggag tgatgaaaag cctaaaaatt	1500
gagttttttt cgttgttctg attgttgtt ttatgaattt gcagatggat atctgtgctc	1560
tctatgcaa tctagctttt cgaatgagag tgataagaga gtgaggattg tgaattattt	1620
tattgatgaa gattggagaa gtcaattatt gattcacaca caggaattaa gtgtgttggtg	1680
ttgcgtcctc ttgtggaaat taaatgtcac ccttttttta tttatcaata aaagcacgaa	1740
aatctcctgc actactcccc tgcactctct tataattgtc catttcccac aaatccctaa	1800
cttaattact taccacact ctaagctttt gattttaatg tttagcaaat gtcctatcag	1860
ttttctcttt ttgtcgaacg gtaatttaga gtttttttg ctatatggat tttcgttttt	1920
gatgtatgtg acaacctcgg ggattgttga tttatttcaa aactaagagt ttttgcttat	1980
tgttctcgct tattttggat atcaatctta gttttataac ttttctagtt ctctacgtgt	2040
taaatgttca acacactagc aatttggtcg cagcgtatgg attatggaac tatcaagtct	2100
gtgggatcga taaatatgct tctcaggaat ttgagatttt acagtcttta tgctcattgg	2160
gttgagtata atatagtaaa aaaatagtct aga	2193

-continued

```

<211> LENGTH: 1861
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Alternative
        expression cassette for a sense and antisense copy of the leader
        associated with the L glucan phosphorylase gene

<400> SEQUENCE: 26

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg      60
gaccagtacc agtacattag atattatctt ttattactat aataaatatt taattaacac      120
gagacatagg aatgtcaagt ggtagcggta ggaggagggt ggttcagttt tttagatact      180
aggagacaga accggagggg ccattgcaa ggcccaagtt gaagtccagc cgtgaatcaa      240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt      300
gccttcgctt aagggatagc caccgctat tctcttgaca cgtgtcactg aaacctgcta      360
caataaaggc aggcacctcc tcattctcac actcactcac tcacacagct caacaagtgg      420
taacttttac tcattctctc caattatttc tgatttcacg catgtttccc tacattctat      480
tatgaatcgt gttatgggtg ataaacgttg tttcatatct catctcatct attctgattt      540
tgattctctt gcctactgaa ttgacccta ctgtaatcgg tgataaatgt gaatgcttcc      600
tcttctctct cttctcttca gaaatcaatt tctgttttgt tttgttcat ctgtagcttg      660
gtagattccc cttttttagt accacacatc acggatccga gtgtgggtaa gtaattaagt      720
tagggatttg tgggaaatgg acaaatataa gagagtgcag gggagtagtg caggagattt      780
tcgtgctttt attgataaat aaaaaagggt tgacatttaa tttccacaag aggacgcaac      840
acaacacact taattcctgt gtgtgaatca ataattgact tctccaatct tcatcaataa      900
aataattcac aatcctcact ctcttatcac tctcattcga aaagctagat ttgcatagag      960
agcacagaat tcgtggtaac ttttactcat ctctccaat tatttctgat ttcatgcacg      1020
tttccctaca ttctattatg aatcgtgtta tgggtgataa acgttgtttc atatctcatc      1080
tcattctatt tgattttgat tctcttgctt actgaatttg accctactgt aatcggtgat      1140
aaatgtgaat gcttcctctt cttctcttcc ttctcagaaa tcaatttctg ttttggtttt      1200
gttcatctgt agcttgatat ctgtgctctc tatgcaaata tagcttttcg aatgagagtg      1260
ataagagagt gaggattgtg aattatttta ttgatgaaga ttggagaagt caattattga      1320
ttcacacaca ggaattaagt gtgttggtgt ggcctctctt ttggaaatta aatgtcacc      1380
tttttttatt tatcaataaa agcacgaaaa tctcctgcac tactccctg cactctctta      1440
tatttgacca tttccacaa atccctaact taattactta cccacactct aagcttttga      1500
ttttaatggt tagcaaatgt cctatcagtt ttctcttttt gtcgaacggt aatttagagt      1560
tttttttgct atatggattt tcgtttttga tgtatgtgac aacctcggg attgttgatt      1620
tatttcaaaa ctaagagttt ttgcttattg ttctcgtcta ttttgatat caatcttagt      1680
tttatatctt ttctagtctt ctacgtgtta aatgttcaac aactagcaa tttggctgca      1740
gcgtatggat tatggaacta tcaagtctgt gggatcgata aatatgcttc tcaggaattt      1800
gagattttac agtctttatg ctcatgggtt tgagtataat atagtaaaaa aatagtctag      1860
a                                                                                   1861

```

<210> SEQ ID NO 27

-continued

<211> LENGTH: 1788

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 27

```
atggcaagct tgtgcaatag tagtagtaca tctctcaaaa ctccttttac ttcttcctcc    60
acttctttat cttccactcc taagccctct caacttttca tccatggaaa acgtaaccaa    120
atgttcaaaag tttcatgcaa ggttatcaat aataacggtg accaaaacgt tgaacgaat    180
tctgttgatc gaagaaatgt tcttcttggc ttaggtggtc tttatggtgt tgctaattgt    240
ataccattag ctgcatccgc tgctccaact ccacctcctg atctctcgtc ttgtagtata    300
gccaggatta acgaaaaatca ggtggtgccg tacagttggt gcgcgcctaa gcctgatgat    360
atggagaaaag ttccgtatta caagttccct tctatgacta agctccgtgt ccgtcagcct    420
gctcatgaag ctaatgagga gtatatggcc aagtacaatc tggcgattag tcgaatgaga    480
gatcttgata agacacaacc tttaaaccct attggtttta agcaacaagc taatatacat    540
tgtgcttatt gtaatggtgc ttatagaatt ggtggcaaag agttacaagt tcataattct    600
tggtctttct tcccgttcca tagatggtac ttgtacttcc acgagagaat cgtgggaaaa    660
ttcattgatg atccaacttt cgctttgcc aattggaatt gggaccatcc aaagggatg    720
cgttttcctg ccatgtatga tcgtgaaggg acttcccttt tcgatgtaac acgtgaccaa    780
agtcaccgaa atggagcagt aatcgatctt ggttttttgc gcaatgaagt cgaacaact    840
caactccagt tgatgagcaa taatttaaca ctaatgtacc gtcaaatggt aactaatgct    900
ccatgtcctc ggatgttctt tgggtggcct tatgatctcg ggattaacac tgaactcccg    960
ggaactatag aaaacattcc tcacggctct gtccacatct ggtctggtac agtgagaggt   1020
tcaactttgc ccaatggtgc aatatcaaac ggtgagaata tgggtcattt ttactcagct   1080
gctttggacc cggttttctt ttgccatcac agcaatgtgg atcggatgtg gagegaatgg   1140
aaagcgacag gagggaaaag aacagatata acacataaag gttggttgaa ctccgagttc   1200
tttttctatg atgaaaaatga aaacccttac cgtgtgaaag tccgagactg tttggacacg   1260
aagaagatgg ggtatgatta tgcaccaatg gccaccccggt ggcgtaactt caagccaata   1320
acaaaaacta cagctgggaa agtgaataca gcttctcttc cgccagctag caatgtattc   1380
ccagtggcta aactcgacaa agcaatttcg ttttccatca ataggccgac ttcgtcaagg   1440
actcaacaag agaaaaatgc acaagaggag atgttgacat tcagtagcat aagatatgat   1500
aacagagggg acataagggt cgatgtgttc ctgaacgtgg acaataatgt gaatgcgaat   1560
gagcttgaca agcgggagtt tgcggggagt tatactagtt tgccacatgt tcatagagct   1620
ggtgagacta atcatatcgc gactgttgat ttccagctgg cgataacgga actgttgagg   1680
gatattgggt tggaagatga agatactatt gcggtgactc tggtgccaaa gagaggtggt   1740
gaaggtatct ccattgaaag tgcgacgata agtcttgacg attgttaa                   1788
```

<210> SEQ ID NO 28

<211> LENGTH: 1788

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 28

```
atggcaagct tgtgcaatag tagtagtaca tctctcaaaa ctccttttac ttcttcctcc    60
```

-continued

acttctttat cttccactcc taagccctct caacttttca tccatggaaa acgtaaccaa	120
atgttcaaag ttcatgcaa ggttatcaat aataacggtg accaaaacgt tgaaacgaat	180
tctgttgatc gaagaaatgt tcttcttggc ttaggtggtc tttatggtgt tgctaagtct	240
ataccattag ctgcattccg tgcctcaact ccacctctcg atctctcgtc ttgtagtata	300
gccaggatta acgaaaatca ggtggtgccg tacagttggt gcgcgctaa gcctgatgat	360
atggagaaag ttccgtatta caagttccct tctatgacta agctccgtgt ccgtcagcct	420
gctcatgaag ctaatgagga gtatatgcc aagtacaatc tggcgattag tcgaatgaga	480
gatcttgata agacacaacc tttaaacctt attggtttta agcaacaagc taatatacag	540
tgggcttatg gtaatggtgc ttatagaatt ggtggcaaag agttacaagt tcataattct	600
tggcttttct tcccggtcca tagatggtac ttgtacttcc acgagagaat cgtgggaaaa	660
ttcattgatg atccaacttt cgctttgcc aattggaatt gggaccatcc aaagggtatg	720
cgttttcctg ccatgtatga tcgtgaaggg acttcccttt tcgatgtaac acgtgaccaa	780
agtcaccgaa atggagcagt aatcgatctt ggttttttcg gcaatgaagt cgaacaact	840
caactccagt tgatgagcaa taatttaaca ctaatgtacc gtcaaagggt aactaatgct	900
ccatgtcctc ggatgttctt tgggtggcct tatgatctcg ggattaacac tgaactcccg	960
ggaactatag gaaacattcc tctcggctct gtccacatct ggtctggtac agtgagaggt	1020
tcaactttgc ccaatggtgc aatatcaaac ggtgagaata tgggtcattt ttactcagct	1080
gctttggacc cggttttctt ttgccatcac agcaatgtgg atcggatgtg gagcgaatgg	1140
aaagcgacag gagggaaaag aacagatatc acacataaag gttggttgaa ctccgagttc	1200
tttttctatg atgaaaatga aaacccttac cgtgtgaaag tccgagactg tttggacacg	1260
aagaagatgg ggtatgatta tgcaccaatg gccaccccg ggcgtaactt caagccaata	1320
acaaaaacta cagctgggaa agtgaatata gcttctcttc cgccagctag caatgtattc	1380
ccagtggcta aactcgacaa agcaatttcg tttccatca ataggccgac ttcgtcaagg	1440
actcaacaag agaaaaatgc acaagaggag atgttgacat tcagtagcat aagatatgat	1500
aacagagggg acataagggt cgatgtgttc ctgaacgtgg acaataatgt gaatgcgaat	1560
gagcttgaca aggcggagtt tgcggggagt tatactagtt tgccacatgt tcatagagct	1620
ggtgagacta atcatatcgc gactgttgat ttccagctgg cgataacgga actgttgag	1680
gatattgggt tggaaagatga agatactatt gcggtgactc tggtgccaaa gagaggtggt	1740
gaaggatatc ccattgaaag tgcgacgatc agtcttgacg attgttaa	1788

<210> SEQ ID NO 29

<211> LENGTH: 154

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 29

ttagtctcta ttgaatctgc tgagattaca ctttgatgga tgatgctctg tttttgtttt	60
cttggtctgt tttttcctct gttgaaatca gctttgttgc ttgatttcat tgaagttggt	120
attcaagaat aaatcagtta caattatggt tggg	154

<210> SEQ ID NO 30

<211> LENGTH: 1691

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Expression cassette for a sense and antisense copy of the trailer associated with a PPO gene

<400> SEQUENCE: 30

```

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg      60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac      120
gagacatagg aatgtcaagt ggtagcggta ggaggggagt ggttcagttt ttagatact      180
aggagacaga accggagggg cccattgcaa ggccaagt gaagtccagc cgtgaatcaa      240
caaagagagg gcccataata ctgctgatga gcatttcctt ataatacagt gtccacagtt      300
gccttcgcgt aagggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta      360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat      420
ccttagtctc tattgaatct gctgagatta cactttgatg gatgatgctc tgtttttgtt      480
ttcttgttct gttttttcct ctggtgaaat cagctttggt gcttgatttc attgaagttg      540
ttattcaaga ataatcagtg tacaattatg gaattcaagg ttagaaatct tctctatttt      600
tggtttttgt ctgtttgatg tctcgaatta gctaatacagg tgctgttata gcccttaatt      660
ttgagttttt tttcggttgt ttgatggaa aaggcctaaa atttgagttt ttttacgttg      720
gtttgatgga aaaggcctac aattggagtt ttccccgttg ttttgatgaa aaagcccta      780
gtttgagatt ttttttctgt cgattcgatt ctaaaggttt aaaattagag tttttacatt      840
tgtttgatga aaaaggcctt aaatttgagt ttttcggtt gatttgatga aaaagcccta      900
gaatttggtt tttttcgtcg gtttgattct gaaggcctaa aatttgagtt tctccggctg      960
ttttgatgaa aaagccctaa atttgagttt ctccggctgt tttgatgaaa aagccctaaa     1020
tttgagtttt ttccccgtgt ttagattgtt ttggttttaa ttctcgaatc agctaatacag     1080
ggagtgatgaa aagccctaaa atttgagttt tttcgttgtt tctgattggt gtttttatga     1140
atttgcatg gcatatcctt ctttgatgct gatccataat tgtaactgat ttattcttga     1200
ataacaactt caatgaaatc aagcaacaaa gctgatttca acagaggaaa aaacagaaca     1260
agaaaacaaa aacagagcat catccatcaa agtgtaatct cagcagattc aatagagact     1320
aagcttttga ttttaagtgt tagcaaatgt cctatcagtt ttctcttttt gtcgaacggg     1380
aatttagagt tttttttgct atatggattt tcgtttttga tgtatgtgac aacctcgagg     1440
attgttgatt tatttcaaaa ctaagagttt ttgcttattg ttctcgtcta ttttgatat     1500
caatcttagt tttatatctt ttctagttct ctacgtgtta aatgttcaac aactagcaa     1560
tttggtgca gcgtatggat tatggaacta tcaagtctgt gggatcgata aatagcttc     1620
tcaggaattt gagattttac agtctttatg ctcattgggt tgagtataat atagtaaaaa     1680
aatagtctag a                                     1691

```

<210> SEQ ID NO 31

<211> LENGTH: 1359

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Expression cassette for a sense and antisense copy of the trailer associated with a PPO gene

-continued

<400> SEQUENCE: 31

```

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg      60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac      120
gagacatagg aatgtcaagt ggtagcggta ggagggagtt ggttcagttt ttagatact      180
aggagacaga accggagggg cccattgcaa ggccaagtt gaagtccagc cgtgaatcaa      240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt      300
gccttccgct aagggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta      360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caagaaggat      420
ccttagtctc tattgaatct gctgagatta cactttgatg gatgatgctc tgtttttgtt      480
ttcttgttct gttttttcct ctggtgaaat cagctttggt gcttgatttc attgaagttg      540
ttattcaaga ataaatcagt tacaattatg gaattcgtgg taacttttac tcatctctc      600
caattatttc tgatttcatt catgtttccc tacattctat tatgaatcgt gttatggtgt      660
ataaacgttg ttcatatctc catctcatct attctgattt tgattctctt gcctactgaa      720
tttgacccta ctgtaatcgg tgataaatgt gaatgcttcc tcttcttctt cttcttctca      780
gaaatcaatt tctgttttgt tttgttcat ctgtagcttg atatccttct ttgatgctga      840
tccataattg taactgattt attcttgaat aacaacttca atgaaatcaa gcaacaaagc      900
tgatttcaac agaggaaaaa acagaacaag aaaacaaaaa cagagcatca tccatcaaag      960
tgtaatctca gcagattcaa tagagactaa gcttttgatt ttaatgttta gcaaatgtcc     1020
tatcagtttt ctctttttgt cgaacggtaa ttttagagtt ttttgctat atggattttc     1080
gtttttgatg tatgtgacaa ccctcgggat tgttgattta tttcaaaact aagagttttt     1140
gcttattgtt ctgcgtatatt ttggatatca atcttagttt tatactcttt ctagttctct     1200
acgtgttaaa tgttcaacac actagcaatt tggctgcagc gtatggatta tggaactatc     1260
aagtctgtgg gatcgataaa tatgcttctc aggaatttga gattttacag tctttatgct     1320
cattgggttg agtataatat agtaaaaaaa tagtctaga                               1359

```

<210> SEQ ID NO 32

<211> LENGTH: 1967

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Expression cassette for a sense and antisense copy of the trailer associated with a PPO gene

<400> SEQUENCE: 32

```

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg      60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac      120
gagacatagg aatgtcaagt ggtagcggta ggagggagtt ggttcagttt ttagatact      180
aggagacaga accggagggg cccattgcaa ggccaagtt gaagtccagc cgtgaatcaa      240
caaagagagg gcccataata ctgtcgatga gcatttcctt ataatacagt gtccacagtt      300
gccttccgct aagggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta      360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caacaagtgg      420
taacttttac tcatctctc caattatttc tgatttcatt catgtttccc tacattctat      480

```

-continued

tatgaatcgt gttatggtgt ataaacgttg tttcatatct catctcatct attctgattt	540
tgattctctt gcctactgaa ttgacccta ctgtaatcgg tgataaatgt gaatgcttcc	600
tcttctctct cttctcttca gaaatcaatt tctgttttgt tttgttcat ctgtagcttg	660
gtagattccc cttttttag accacacatc acggatcctt agtctctatt gaatctgtg	720
agattacact ttgatggatg atgctctgtt tttgttttct tgttctgttt ttctctctgt	780
tgaaatcagc tttgttgctt gatttcattg aagttgttat tcaagaataa atcagttaca	840
attatggaat tcaaggttag aaatcttctc tatttttggg tttgtctgtt ttagattctc	900
gaattagcta atcaggtgct gttatagccc ttaattttga gtttttttcc ggtgtttttg	960
atggaaaagg cctaaaaatt gagttttttt acgttggttt gatggaaaag gcctacaatt	1020
ggagttttcc ccgtgtttt gatgaaaaag ccctagttt gagatttttt ttctgtcgat	1080
tcgattctaa aggttttaaa ttagagtttt tacatttggt tgatgaaaaa ggccttaaat	1140
ttgagttttt ccggttgatt tgatgaaaaa gccctagaat ttgtgttttt tcgtcggttt	1200
gattctgaag gcctaaaatt tgagtttctc cggctgtttt gatgaaaaag ccctaaaatt	1260
gagtttctcc ggctgttttg atgaaaaagc cctaaatttg agttttttcc ccgtgtttta	1320
gattgttttg ttttaattct cgaatcagct aatcaggag tgtgaaaaag cctaaaattt	1380
gagttttttt cgttgttctg attgtgtttt ttatgaattt gcagatggat atccttcttt	1440
gatgctgac cataattgta actgatttat tottgaataa caacttcaat gaaatcaagc	1500
aacaaagctg atttcaacag aggaaaaac agaacaagaa aacaaaaaca gagcatcatc	1560
catcaaagtg taatctcagc agattcaata gagactaagc ttttgatttt aatgtttagc	1620
aaatgtccta tcagttttct ctttttgcg aacggtaatt tagagttttt ttgctatat	1680
ggattttcgt ttttgatgta tgtgacaacc ctcgggattg ttgatttatt tcaaaactaa	1740
gagtttttgc ttattgttct cgtctatttt ggatatcaat cttagtttta tatcttttct	1800
agttctctac gtgttaaatg ttcaacacac tagcaatttg gctgcagcgt atggattatg	1860
gaactatcaa gtctgtggga tcgataaata tgcttctcag gaatttgaga ttttacagtc	1920
tttatgctca ttgggttgag tataatatag taaaaaata gtctaga	1967

<210> SEQ ID NO 33

<211> LENGTH: 1635

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Expression cassette for a sense and antisense copy of the trailer associated with a PPO gene

<400> SEQUENCE: 33

ggtaccgaac catgcatctc aatcttaata ctaaaaaatg caacaaaatt ctagtggagg	60
gaccagtacc agtacattag atattatctt ttattactat aataatattt taattaacac	120
gagacatagg aatgtcaagt ggtagcggta ggagggagtt ggttcagttt ttagatact	180
aggagacaga accggagggg cccattgcaa ggccaagtt gaagtccagc cgtgaatcaa	240
caaagagagg gcccataata ctgtcgtatg gcatttcctt ataatacagt gtccacagtt	300
gccttccgct aagggatagc caccgcgtat tctcttgaca cgtgtcactg aaacctgcta	360
caaataaggc aggcacctcc tcattctcac actcactcac tcacacagct caacaagtg	420

-continued

taactttttac tcatctcctc caattatttc tgatttcatg catgtttccc tacattctat	480
tatgaatcgt gttatgggtg ataaacgttg tttcatatct catctcatct attctgattt	540
tgattctcct gcctactgaa ttgacccta ctgtaatcgg tgataaatgt gaatgcttcc	600
tcttcttctt cttcttctca gaaatcaatt tctgttttgt tttgttcat ctgtagcttg	660
gtagattccc cttttttag accacacatc acggatcctt agtctctatt gaatctgctg	720
agattacact ttgatggatg atgctctgtt tttgttttct tgttctgttt tttcctctgt	780
tgaaatcagc tttgttgctt gatttcattg aagttgttat tcaagaataa atcagttaca	840
attatggaat tcgtggtaac ttttactcat ctccccaat tatttctgat ttcatgcatg	900
ttccctaca ttctattatg aatcgtgtta tgggtataaa acgttgtttc atatctcatc	960
tcatctattc tgattttgat tctcttgctt actgaatttg accctactgt aatcggtgat	1020
aaatgtgaat gcttcctctt cttcttcttc ttctcagaaa tcaatttctg tttgtttttt	1080
gttcatctgt agcttgatat ccttctttga tgctgatcca taattgtaac tgatttattc	1140
ttgaataaca acttcaatga aatcaagcaa caaagctgat ttcaacagag gaaaaaacag	1200
aacaagaaaa caaaaacaga gcatcatcca tcaaagtgt atctcagcag attcaataga	1260
gactaagcct ttgattttta tgttttagca atgtcctatc agttttctct tttgtcgaa	1320
cggtaattta gagttttttt tgctatatgg attttcgttt ttgatgtatg tgacaaccct	1380
cgggattggt gattttattc aaaactaaga gtttttgctt attgttctcg tctatttttg	1440
atatcaatct tagttttata tcttttctag ttctctacgt gttaaatgtt caacacacta	1500
gcaatttggc tgcagcgtat ggattatgga actatcaagt ctgtgggacg gataaatatg	1560
cttctcagga atttgagatt ttacagctct tatgctcatt gggttgagta taatatagta	1620
aaaaaatagt ctaga	1635

<210> SEQ ID NO 34
 <211> LENGTH: 240
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 34

gtccatgatg tcttcagggt ggtagcattg actgatggca tcatagtttt ttttttaaaa	60
gtatttcctc tatgcatatt attagtatcc aataaattta ctggttggtg tacatagaaa	120
aagtgcattt gcatgtatgt gtttctctga aattttcccc agtttttggt gctttgcctt	180
tggagccaag tctctatatg tataagaaaa ctaagaacaa tcacatatat caaatattag	240

<210> SEQ ID NO 35
 <211> LENGTH: 228
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 35

acgaacttgt gatcgcggtg aaagatttga acgctacata gagcttcttg acgtatcttg	60
caatattgca tcagtcttg cggaatttca tgtgacaaca aggtttgcaa tttttccac	120
tattagtagt gcaacgatat acgcagagat gaagtgtga acaaacatat gtaaaatcga	180
tgaatttatg tcgaatgctg ggacgggctt cagcaggttt tgcttagt	228

<210> SEQ ID NO 36

-continued

<211> LENGTH: 2204
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Expression
cassette for an omega-mutated virD2 gene

<400> SEQUENCE: 36

ccgcgggtttt ctctccatcg cgtcagagggc cggtttttcgt cggcatcgaa gagggccact 60
cgtttaccgt catttgccaa agcagcgcaa agggccatga gtgcggtggt tttgccagca 120
ccccctttga aagagcaaaa cgtcaaaagt tgcataattct gatcccgctt gtcctgtgaa 180
acggagtgc tttgtatttt tgttcgtata aatgtttttg tgattatcga tgagtaaaag 240
cgttggtaca ctatttttta tttcaaattc gttataatta aattgcaatt gtagcaatta 300
tattcggttt ttcctgtaaa tatactgttg atttcataatc gagtagggct agactttaat 360
ctgtctaccc gggcacattt cgtgctggag tattcagacc ttccgctttt tttggaggaa 420
gctatgtcaa aacacaccag agtcacgtcg agtgagactg ccatcaacca gcatcgatcc 480
ctgaacgttg aaggggttaa ggtcgtgagt gcccgctctgc gatcggccga gtatgaaacc 540
ttttcctatc aagcgcgcct gctgggactt tcggatagta tggcaattcg cgttgcggtg 600
cgtcgcacgc ggggctttct cgaaatagat gcacacaccc gagaaaagat ggaagccata 660
cttcagtcga tcggaatact ctcaagtaat gtatccatgc ttctatctgc ctacgccgaa 720
gacctcgtat cggatctgga ggctgtgcga gatgaacgta ttgcttttgg tgaggctttc 780
gccgccctcg atggcctact ccgctccatt ttgtccgtat cccggcgacg gatcgacggt 840
tgctcgtat tgaagggtgc cttgtagcac ttgaccacgc acctgacggg agaaaattgg 900
atgcccgatc gcgctcaagt aatcattcgc attgtgccag gaggtggaac caagaccctt 960
cagcagataa tcaatcagtt ggagtacctg tcccgtaagg gaaagctgga actgcagcgt 1020
tcagcccggc atctcgatat tcccgttccg ccggatcaaa tccgtgagct tgcccaaagc 1080
tgggttacgg agggccggat ttatgacgaa agtcagtcag acgatgatag gcaacaagac 1140
ttaacaacac acattattgt aagcttcccc gcaggtagcg accaaaccgc agcttatgaa 1200
gccagccggg aatgggcagc cgagatgttt gggtcaggat acgggggttg ccgctataac 1260
tatctgacag cctaccacgt cgaccgcgat catccacatt tacatgtcgt ggtcaatcgt 1320
cgggaacttc tggggcacgg gtggctgaaa atatccaggc gccatcccca gctgaattat 1380
gacggcttac ggaaaaagat ggcagagatt tcacttcgtc acggcatagt cctggatgag 1440
acttcgcgag cagaaaaggg aatagcagag cgaccaatca catatgctga acatcgccgc 1500
cttgagcgga tgcaggctca aaagattcaa ttcgaagata cagattttga tgagacctcg 1560
cctgaggaag atcgtcggga cctcagtcga tcgttcgacg catttcgacg ggacccatct 1620
accggcgaac cggaccgtgc aaccgcacat gacaaacaac cgcttgaaca gacgcccgt 1680
ttccaggagt ccgccggctc cagcatcaaa gccgacgcac ggatccgctg atcattggag 1740
agcgagcgga gtgcccaccc atccgcgtcc aaaatccctg taattgggca tttcgggatt 1800
gagacttcct atgtcgtgta agccagcgtg cgaaaacgaa gcggcatttt cgttacttct 1860
cgcccggtga ctgacgttgc catgcacaca gtcaagcgcc agcagcgatc aaaacgacgt 1920
aatgacgagg aggcagggtc gagcgagcga aaccgtaaag gattgaaggc tgcgcaagtt 1980
gattccgagg caaatgtcgg tgagcaagac actcgcgatg acagcaacaa ggcggctgat 2040

-continued

```

ccggtgtctg cttccatcgg taccgagcaa ccggaagctt ctccaaagcg tccgcgtgac 2100
cgtcacgatg gagaattggg tggacgcaaa cgtgcaagag gtaatcgtcg ctcgagctcg 2160
agcgggggga cctagagaca ggaaggaccg aataatggcc gcgg 2204

```

```

<210> SEQ ID NO 37
<211> LENGTH: 1621
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

```

```

<400> SEQUENCE: 37

```

```

atggcttctg tgctggcttc tctgtttcca aaactgggct ctttgggtac ttcagatcat 60
gcttctgttg tatccatcaa cctctttgtg gcaactcctt gtgcttgcat catcattggt 120
catctcttgg aggagaaccg ctgggttaat gagtccatta ctgccctcat aattggtttg 180
tgtacaggag tggttatctt gctcgtaagt ggtggaaaga gtcacacct tctggttttc 240
agtgaagatc tctttttcat atatgtactt cctccaatca tatttaatgc agggtttcag 300
gtaaaaaaga agcaattttt cgtaaaactt attactataa tgatgttcgg agccattggt 360
accctgggtc catgtgccat tatatcatta ggtgccattc aaactttcaa gaagttggac 420
attgaatttc tagatatgtg ggattatctt gcaattggag caatatttgc tgccacagat 480
tccgtctgca cattgcaggc cctacatcag gatgagacac ccctccttta cagtcttgta 540
tttgagaag gaggttgtaa tgatgctaca tcggtggtgc ttttcaatgc tattcaaac 600
ttcgacctta cgagcatgaa tcccagtata gccctcagtt tccttgcaa cttcttctat 660
ctgttccttg ctagcacttt actgggagca ggaactggtc ttcttagtgc ttacattatc 720
aagaagctat attttggcag gcaactccaca gatcgtgagg ttgcccttat gatgctcatg 780
gcttacttat catacttgct ggccgaatta ttctatttga gtgggattct caccgtcttt 840
ttctgtggtg ttgtaatgtc tcaactacact tggcacaatg tgaccgagag ttcaagagtc 900
actacaaggc acacttttgc aactttgtca tttcttgtag agactttcct ctctctctat 960
gtcggcatgg atgcttttga tatcgagaag tggaaatttg ttggtgacag gcctggatta 1020
tcaatttcgg tgagtccaat actgatggga ctaatcttgc ttgggagagc tgcctttggt 1080
tttcatttat cattcttctc caacttaatg aagaaatcct cggagcaaaa aattaccttt 1140
aggcagcaag tgataatatg gtgggcaggc ttgatgagag gcgcagtgtc catggcactg 1200
gcataataata agttcactcg tgggggacac actcaactgc aggacaatgc aataatgatt 1260
accagcacga taaccattgt tctattcagc acaatggtat tcggtttaat gacaaaaccc 1320
cttataagtc tcctgctgcc accacagagg caattgagta cagtgtcatc aggcgcaaatt 1380
actocaaagt ctctaacagc cccactccta ggcagtcgag aggactctga agttgattta 1440
aatgttccag atcttctcctc cccaccaagt ttgaggatgc tacttacgc accaagtcac 1500
aaagtgcacg ggtactggcg caagtttgac gatgcattca tgcgccctat gtttgggtgt 1560
cggggatttg ctctctctgc ccctggttct ccaacggaac aggttccatg aggtaccaat 1620
c 1621

```

```

<210> SEQ ID NO 38
<211> LENGTH: 1620
<212> TYPE: DNA
<213> ORGANISM: Solanum tuberosum

```

-continued

<400> SEQUENCE: 38

```

atggcttctg tgcgtgcttc tctgtttcca aaactgggct ctttgggtac ttcagatcat    60
gcttctgttg tatccatcaa cctctttgtg gcaactcctt gtgcttgcat catcattggg    120
catctcttgg aggagaaccg ctgggttaat gagtccatta ctgccctcat aattggtttg    180
tgtacaggag tggttatctt gctcgtaagt ggtggaaaga actcacacct tctggttttc    240
agtgaagatc tctttttcat atatgtactt cctccaatca tatttaatgc agggtttcag    300
gtaaaaaaga agcaattttt cgtgaacttc attactataa tgatgttcgg agccattggg    360
accttggtct catgtgccat tatatcatta ggtgcaattc aaactttcaa gaagttggac    420
attgaatttc tagatatagg ggattatctt gcaattggag caatatttgc tgccacagat    480
tccgtctgca cattgcaggc cctacatcag gatgagacac ccctccttta cagtcttgta    540
tttgagaag gagttgtaaa tgatgctaca tcggtggtgc ttttcaatgc tattcaaaac    600
tttgacctta cgagcgtgaa tcccagtata gccctcagtt tccttgcaa cttcttctat    660
ctgttccttg ctagcacttt actgggagca ggaactggtc ttcttagtgc ttacattatc    720
aagaagctgt attttggcag gcaactccaca gatcgtgagg ttgcccttat gatgctcatg    780
gcttacttat catacatgct ggctgaacta ttctatttga gtgggattct cactgtattt    840
ttctgtggta ttgtaatgtc tcattacact tggcacaatg tgaccgagag ttcaagagtc    900
actacaaggc acgcttttgc aactttgtca tttcttgca agactttcct ctctctctat    960
gtcggcatgg atgcttttga tatcgagaag tggaaatttg ttggtgacag gcctggatta    1020
tcaatttccg tgagtccaat actgatggga ttaatcttgc tggggagagc tgcctttgtt    1080
tttcatttat cattcttctc caacttaatg aagaaatcct cggagcaaaa aattaccttt    1140
aggcagcaag tgataatatg gtgggcaggc ttgatgagag gcgcagtgtc catggcactg    1200
gcataataa agttcactcg tgggggacac actcaactgc aggacaatgc aataatgatt    1260
accagcacga taaccattgt tctattcagc acaatggtat tcggtttaat gacaaaaccc    1320
cttataagtc tcctgtctgc accacagagg caattgagta cagtgtcatc aggtgcaa    1380
actocaaagt ctctaacagc cccactccta ggcagtcgag aggactctga agttgattta    1440
aatgttccag atcttctcct cccaccaagt ttgaggatgc tacttacgc accaagtc    1500
aaagtgcacg ggtactggcg caagtttgac gatgcattca tgcgccctat gtttggtggg    1560
cggggatttg ctctctctgc ccctggttct ccaacggaac aggttccatg aggtacaatc    1620

```

<210> SEQ ID NO 39

<211> LENGTH: 747

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 39

```

atggaaaatt cggtaaccag gactgtagaa gaagtattca acgatttcaa aggtcgtaga    60
gtcgtgttaa tcaaagcact aactacagat gtcgagaagt tttatcaatc gtgtgatcct    120
gaaaaggaga acttgtgtct ctatgggctt cctaatgaaa catgggaagt aaacctccct    180
gtagaggagg tgccctccaga acttcggag ccagcattgg gcataaactt cgcacgtgat    240
ggaatgcaag agaaagactg gttatcactt gttgctgttc acagtgatc atggctgctt    300
tctgttgcat tttacttttg tgcaagggtt gggttcggca agagtgaag gaagaggctt    360

```

-continued

ttccaaatga taaatgatct cccaacagtg tttgaagttg ttaccggagc tgctaaacag	420
acacgtgatc cccctcacia caatagcaac aaaagcaaat caagtggaaa gcctcgacag	480
ccagagtccc aactcaaggc agtaaagggtg tctccaccta aaatggagaa cgacagtggg	540
gaggaggaag aagaagaaga ggatgaacaa ggagcaactc tctgtggagc ttgtggtgat	600
aattatgcc a ctgatgaatt ctggatttgc tgtgatattt gtgagagatg gttccatggc	660
aaatgtgtga agattacccc agcaaaagct gagcatatca agcagtacaa gtgtcctagt	720
tgcatgtgca agagagctag agtttaa	747

<210> SEQ ID NO 40
 <211> LENGTH: 741
 <212> TYPE: DNA
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 40

tgacatctgc caataaagcc aagaataatt ggcattaaca tgaccaaaaa aatggtttgg	60
cagcattaag tcaataaaaa aagctacttt aatataaaat aatattaaaa tgcttaataa	120
ccaacagttt ataagaaggt taatgttaac atggatgagg aatgaccaa aggggaatta	180
tatattaacc tttaaatcaa tctaattctc tctttttggt tctagctata ttactcgat	240
agataaactc tcttacttga cgaatTTTT gatacaagaa gacataattc atcatgattt	300
taattcgtcg tgtcaaatTT attaaatagt ttaattttta tcgtaaattt agatatgaaa	360
tttaaaaaaa aataaatata tacatatttg aagaatacat aaaaagtaca tataaatcac	420
aaatatttaa taattcaaga tattaataca catagaaaa taattactta caaagaaatt	480
cttatttgaa tcctctaata tcgagaagtg caacacaaac tgagacgaag aaaatgaata	540
atatttgata agaaatttat tataattgaa tgaccattta agtaattacg ggtaataaca	600
acacaataag gaactgtagt catTTTTaat acatggcaag gaatatgaga gtgtgatgag	660
tctataaata gaaggcttca ttagtgtaga ggagtcacaa acaagcaata cacaataaaa	720
attagtagct taaacaagat g	741

<210> SEQ ID NO 41
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Agrobacterium sp.

<400> SEQUENCE: 41

tgacaggata tattggcggg taaac	25
-----------------------------	----

<210> SEQ ID NO 42
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Agrobacterium sp.

<400> SEQUENCE: 42

tgacaggata tattgtggtg taaac	25
-----------------------------	----

<210> SEQ ID NO 43
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Agrobacterium sp.

<400> SEQUENCE: 43

-continued

tggcaggata tataccgttg taatt 25

<210> SEQ ID NO 44
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Agrobacterium sp.

<400> SEQUENCE: 44

cggcaggata tattcaattg taatt 25

<210> SEQ ID NO 45
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Agrobacterium sp.

<400> SEQUENCE: 45

tggtaggata tataccgttg taatt 25

<210> SEQ ID NO 46
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Agrobacterium sp.

<400> SEQUENCE: 46

tggcaggata tatggtactg taatt 25

<210> SEQ ID NO 47
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Consensus
sequence
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (16)
<223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 47

ygryaggata tatwsnvbkg taawy 25

<210> SEQ ID NO 48
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Rhizobium leguminosarum

<400> SEQUENCE: 48

cggcaggata tatcctgatg taaat 25

<210> SEQ ID NO 49
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Thermoanaerobacter tengcongensis

<400> SEQUENCE: 49

tggcaggagt tattcgaggg taaac 25

<210> SEQ ID NO 50
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Arabidopsis thaliana

-continued

<400> SEQUENCE: 50

tgacaggata tatcgtgatg tcaac 25

<210> SEQ ID NO 51

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 51

gggaagtaca tattggcggg taaac 25

<210> SEQ ID NO 52

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Oryza sativa

<400> SEQUENCE: 52

ttacaggata tattaatatg tatga 25

<210> SEQ ID NO 53

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

taacatgata tattcccttg taaat 25

<210> SEQ ID NO 54

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 54

tgacaggata tatggtaatg taaac 25

<210> SEQ ID NO 55

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 55

tggcaggata tataccgatg taaac 25

<210> SEQ ID NO 56

<211> LENGTH: 292

<212> TYPE: DNA

<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 56

ttcttcgcca gaggttttgt caagtctcca atcaaggttg tcggcttgtc taccttgcca 60

gaaatttacg aaaagatgga aaagggtcaa atcgttggtg gatacgttgt tgacacttct 120

aaataagcga atttcttatg atttatgatt tttattatta aataagttat aaaaaaata 180

agtgtataca aattttaaag tgactcttag gttttaaaac gaaaattctt attcttgagt 240

aactctttcc tgtaggtcag gttgctttct caggtatagc atgaggtcgc tc 292

<210> SEQ ID NO 57

<211> LENGTH: 25

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (14)
<223> OTHER INFORMATION: a, t, c or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (16)
<223> OTHER INFORMATION: a, t, c or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (18)
<223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 57

tgrcaggata tatnvndntg taaac 25

<210> SEQ ID NO 58
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 58

ccgcggtgat cacaggcagc aac 23

<210> SEQ ID NO 59
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 59

aagcttccag ccagccaaca gctccccgac 30

<210> SEQ ID NO 60
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 60

aagcttggct actagtgcga gatctctaag agaaaagagc gtta 45

<210> SEQ ID NO 61
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 61

gcatgctcga gataggtgac cacatacaaa tggacgaacg g 41

<210> SEQ ID NO 62
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

-continued

<400> SEQUENCE: 62

actagtgttt acccgccaat atatcctgtc agag

34

<210> SEQ ID NO 63

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 63

aagctttggc aggatatt gtggtgtaaa cgaag

35

<210> SEQ ID NO 64

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 64

cgggtgaagt gaactgcagt tgccatg

27

<210> SEQ ID NO 65

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 65

catcggcctc actcatgagc agattg

26

<210> SEQ ID NO 66

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 66

cacgctaagt gccggccgtc cgag

24

<210> SEQ ID NO 67

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 67

tcctaatacga cggcgaccg gctg

24

<210> SEQ ID NO 68

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 68

-continued

aaagttgaat tcaaatgaga aatttattc 29

<210> SEQ ID NO 69
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 69

ttttaagctt tcataataac attctaatt 28

<210> SEQ ID NO 70
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 70

gaaccatgca tctcaatc 18

<210> SEQ ID NO 71
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 71

gtcaggatcc ctaccaagct acagatgaac 30

<210> SEQ ID NO 72
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 72

ggatccgagt gtgggtaagt aattaag 27

<210> SEQ ID NO 73
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 73

gaattctgtg ctctctatgc aaatctagc 29

<210> SEQ ID NO 74
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 74

ggaacattga agctgtgg 18

<210> SEQ ID NO 75

-continued

<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 75

cgaattcatg gcaagcttgt gcaatag 27

<210> SEQ ID NO 76
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 76

cgaattctta acaatctgca agactgatcg 30

<210> SEQ ID NO 77
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 77

gagagatctt gataagacac aacc 24

<210> SEQ ID NO 78
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (7)
<223> OTHER INFORMATION: "a" to "c" mutation
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (14)
<223> OTHER INFORMATION: "a" to "c" mutation
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (17)
<223> OTHER INFORMATION: "a" to "c" mutation

<400> SEQUENCE: 78

cattaccata agcccactgt atattagctt gttgc 35

<210> SEQ ID NO 79
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 79

gtgcttatag aattggtggc 20

<210> SEQ ID NO 80
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 80

tagttcccgg gagttcagtg 20

<210> SEQ ID NO 81

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (17)

<223> OTHER INFORMATION: "a" to "g" mutation

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (29)

<223> OTHER INFORMATION: "a" to "t" mutation

<400> SEQUENCE: 81

ctcccgaggaa ctataggaaa cattcctctc ggtcctgtcc acatctggtc 50

<210> SEQ ID NO 82

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 82

gtgtgatatc tgttcttttc c 21

<210> SEQ ID NO 83

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 83

gaatgagctt gacaaggcgg ag 22

<210> SEQ ID NO 84

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 84

ctggcgataa cggaactgtt g 21

<210> SEQ ID NO 85

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 85

gtccatgatg tcttcagggt ggta 24

<210> SEQ ID NO 86

-continued

<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 86

ctaataatttg atatattgga ttgt 24

<210> SEQ ID NO 87
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 87

acgaacttgt gatcgcttg aaag 24

<210> SEQ ID NO 88
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 88

actaagcaaa acctgctgaa gcc 24

<210> SEQ ID NO 89
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 89

cccggtatg cttctgtgct ggct 24

<210> SEQ ID NO 90
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 90

ggtacctcat ggacctgtt ccgt 24

<210> SEQ ID NO 91
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 91

cccggtatg gaaaattcgg taccaggac tg 32

<210> SEQ ID NO 92
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 92

actagttaaa ctctagctct ctg

25

<210> SEQ ID NO 93

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (2)

<223> OTHER INFORMATION: a, t, c or g

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (6)

<223> OTHER INFORMATION: a, t, c or g

<220> FEATURE:

<221> NAME/KEY: modified_base

<222> LOCATION: (10)..(15)

<223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 93

angatntatn nnnntgt

18

<210> SEQ ID NO 94

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Triticum sp.

<400> SEQUENCE: 94

tggcaggata tatgagtgtg taaac

25

<210> SEQ ID NO 95

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Triticum sp.

<400> SEQUENCE: 95

ttggcaggat atatccctct gtaaac

26

<210> SEQ ID NO 96

<211> LENGTH: 74

<212> TYPE: PRT

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 96

Met Ser Ser Thr Ser Asn Val Gly Gln Asp Cys Leu Ala Glu Val Thr
1 5 10 15

Ile Ser Tyr Gln Trp Val Gly Arg Val Ile Asn Tyr Asn Phe Phe Leu
20 25 30

Leu Ile His Trp Tyr Thr Val Val Glu Ala Ser Thr Gly Ile Thr Phe
35 40 45

Gln Ile Phe Pro Ile Gly Ile Arg Ser Glu Asp Asp Arg Ser Phe Tyr
50 55 60

Glu Lys Ala Asp Arg Phe Ala Trp Val Thr
65 70

<210> SEQ ID NO 97

<211> LENGTH: 51

-continued

<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum
<400> SEQUENCE: 97
Met Ser Ser Glu Ser Thr Phe Ser Lys Thr Pro Asn Gly Arg Ala Thr
1 5 10 15
Asp Val Gly Ile Pro Thr Glu Glu Gly Thr Phe Pro Phe Arg Tyr Ala
20 25 30
Ile Leu Arg Asp Leu Ala Pro Thr Ile Ser Leu Val Asn Ser Ser Ala
35 40 45
Asp Ile Ala
50

<210> SEQ ID NO 98
<211> LENGTH: 76
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum
<400> SEQUENCE: 98
Met Ser Glu Gly Val Gly Phe Lys Ser Lys Ile Leu Pro Ser Phe Ala
1 5 10 15
Trp Arg Ser Ala Asn Ile Leu Gly Ser Lys His Val Ala Lys Gln Thr
20 25 30
Phe Pro Phe Leu Ala Arg Thr Glu Thr Cys Glu Arg Thr Ser Gly Met
35 40 45
Ser Gly Val Ile Arg Ala Thr Ala Pro Ser Gly Ile Ser Ser Ser Pro
50 55 60
Leu Thr Asp Phe Ala Thr Lys Ile Val Gly Phe Ser
65 70 75

<210> SEQ ID NO 99
<211> LENGTH: 62
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum
<400> SEQUENCE: 99
Val Cys Ser Pro Ala Leu Lys Ala Asp Lys Ser Lys Ser Ala Asp Gly
1 5 10 15
Thr Cys Val Asp His Ser Arg Arg Leu Ile Val Val Leu Val Leu Tyr
20 25 30
Pro Gly Met Gly Thr Ser Tyr Ala Thr Ala Phe Ile Ser Ser Pro Pro
35 40 45
Ile Gln Tyr Leu Phe Pro Ser Asp Pro Val Glu Thr Phe Pro
50 55 60

<210> SEQ ID NO 100
<211> LENGTH: 50
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum
<400> SEQUENCE: 100
Met Leu Gly Ser Leu Val Leu Pro Lys Ser Pro Glu Asn Arg Lys Gln
1 5 10 15
Ala Val Pro Asn Pro His Phe Gln Glu Gln His Leu Val Pro Glu Lys
20 25 30
Pro His Phe Leu Asp Cys Gly Gln Gly Phe Ser Lys Leu Pro Gln Met
35 40 45

-continued

His Gln
50

<210> SEQ ID NO 101
<211> LENGTH: 65
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 101

Met Val Asn Phe Leu Thr Gln Gly Ile Val Asp Met Glu Thr Ala Phe
1 5 10 15
Gly Ser Pro Lys Met Gly Gly Phe Gly Lys Glu Gln Phe Gly Ala Cys
20 25 30
Val Ser Arg Ser Glu Met Asp Glu Ser Gly Ile Gly Ala Val Met Val
35 40 45
Glu Gln Val Cys Ser Ile Cys Ser Arg His Phe Val Leu Ser Met Gln
50 55 60
Ile
65

<210> SEQ ID NO 102
<211> LENGTH: 77
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 102

Met Leu Glu Gly Ser Met Trp Pro Trp Asn Gln Glu Ser Met Lys Arg
1 5 10 15
Ala Phe Leu Asn His His Phe Leu Met Leu His Leu Phe Pro Ala Gln
20 25 30
Arg Pro Pro Gln Ala Ala Asp Pro Val Cys Leu Lys His Gln His Met
35 40 45
His Cys Gly Cys Leu Ser Phe Gln Leu His Leu Ser Lys Leu Ala Pro
50 55 60
Gly Asp Thr Pro Leu Ile Ser Ser Met Phe Ala Leu Asp
65 70 75

<210> SEQ ID NO 103
<211> LENGTH: 49
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 103

Met Lys Leu Cys Ser Ser Ile Ile Leu Ser Ile Ile Lys Gln Lys Gln
1 5 10 15
Val Glu Ile Leu Arg Ala Cys Phe Gly Phe Pro Glu Thr Lys Thr Ile
20 25 30
Ser Val Phe Ser Ser Val Ser Trp Asn Trp His Ile Ile Cys Lys Ser
35 40 45
Leu

<210> SEQ ID NO 104
<211> LENGTH: 64
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 104

-continued

```

Met Thr Lys Lys Pro Asp Arg Lys Asp Asn Ile Met Pro Tyr Asn Phe
 1           5           10           15
Pro Gly Thr Lys Phe Leu Gln Pro Ile Phe Arg Asn Phe Phe Leu Pro
          20           25           30
Ser Leu Cys Asp Lys Leu Leu Lys Lys Ser Ile Ser Val Pro Gln Ala
          35           40           45
Ile Thr Pro Cys Trp Lys Val Gln Cys Gly His Gly Ile Lys Lys Ala
 50           55           60

```

```

<210> SEQ ID NO 105
<211> LENGTH: 115
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

```

```

<400> SEQUENCE: 105

```

```

Thr Ile Leu Lys Leu Asp Leu His Thr Phe Asn Gly His Phe Phe Thr
 1           5           10           15
Ala Ser Phe Trp Asn Gln Ser His Arg Asn Ser Ile Phe Ile Phe Gln
          20           25           30
Ser Asn Ile Leu Gln Gln Phe Ser Tyr Arg Gln Leu Glu Ser Asn Thr
          35           40           45
Gly Asn Met Ile Ser Ile Thr Ser Met Asn Met Arg Gln Ala Ser Ile
          50           55           60
Thr Pro Cys Lys Leu Arg Leu Ile Lys Leu Ile Cys Ile His Ser Leu
          65           70           75           80
Val His Val Gln Lys His Ile Glu Pro Tyr Ile Val Pro Ile Ile Ile
          85           90           95
Arg Tyr Phe Ile Glu Cys Gln Tyr Leu Leu Leu Leu Ile Phe Leu Leu
          100          105          110
Cys Cys Pro
          115

```

```

<210> SEQ ID NO 106
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

```

```

<400> SEQUENCE: 106

```

```

Met Lys Gly Lys Glu Lys Pro Arg Glu Met Asn Leu Gln Phe Phe Thr
 1           5           10           15
Thr Asn Phe Val Ser Thr Val Ala Ile Ser Thr Met Asn Ile Ser Leu
          20           25           30
Leu Phe Lys Ala Lys Arg Val Lys Gly Val Phe Ile Lys Phe Pro His
          35           40           45
Ser Thr Arg Ser Gln Leu Ile Leu Gly Tyr Val Leu Leu Ile Arg Arg
          50           55           60
Met Ser Arg Gly Ala Asp Ala Glu Phe Ser His Arg Arg Glu Leu Val
          65           70           75           80
Val Arg Asn Thr Ile Asp Leu Ile Gly Tyr Arg Arg Ala Thr Thr Val
          85           90           95
Tyr Tyr Ile Asn Thr Phe Phe Tyr Met Gly Ser Thr Thr Arg Leu Glu
          100          105          110
Ile Arg Arg Trp Tyr Arg Cys Ser Ser Arg
          115          120

```

-continued

<210> SEQ ID NO 107

<211> LENGTH: 104

<212> TYPE: PRT

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 107

Met Glu Trp Ala Leu Ala Arg Asn Arg Ile Pro Phe Phe Tyr Cys Pro
1 5 10 15
Asn Ser Leu Arg Thr Ser His Gly Lys Gly Tyr Asp Phe His Arg Arg
20 25 30
Lys Arg Ile Gln Ser Ser Thr Asn Leu Tyr Leu Leu Asn Pro Phe Phe
35 40 45
Ser Arg Gln Leu Ile Ser Ile His Ser Thr Ser Cys Pro His Trp His
50 55 60
Gly Gly Ser Lys Lys Ser Asp Leu Asn Arg Val Ser Arg Asn Tyr Pro
65 70 75 80
Cys Leu His Arg Phe Phe Asp Glu Val Cys His Arg Ser Arg Cys Glu
85 90 95
Pro Glu Tyr Glu Gly Cys Phe Gln
100

<210> SEQ ID NO 108

<211> LENGTH: 92

<212> TYPE: PRT

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 108

Met Asn Asn Ile Thr His Ser Pro Ile Leu Ile Pro Phe Leu Glu Gln
1 5 10 15
Leu Asn Pro Phe Ile Ser Asn Cys His Met Gln Pro Ile Val Lys Ala
20 25 30
Asn Thr Pro Ile Leu Asn Gly Asn Thr Lys Cys Arg His Ser Ala Asn
35 40 45
Ile Phe Thr Asn Gly Asn Cys Ile Trp Glu Lys Pro Met Asn Lys Ile
50 55 60
Val Asp Gln His Gln Ile His Asn Ser Ile His Ile Ser Cys Glu Ser
65 70 75 80
Lys Val Phe Leu Val Val Pro Ser Glu Ser His Arg
85 90

<210> SEQ ID NO 109

<211> LENGTH: 57

<212> TYPE: PRT

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 109

Met Lys Phe Arg Tyr Pro Ser Pro Pro Asn Pro Ile Val Thr Ser Leu
1 5 10 15
Ile Ile Leu Cys Asn Ala Ile Pro Arg Ser Ile Asn Asp Val Asp Gly
20 25 30
Leu Ser Arg Ala Ile Lys Ser Tyr Ile Ser Leu Ser Ile Ser Gln Asn
35 40 45
Ala Ile Val Leu Ser Pro Thr Arg Ala
50 55

-continued

<210> SEQ ID NO 110
<211> LENGTH: 70
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 110
Met Val Asn Ile Met Thr Ser Ser Ser Met Ala Thr Lys Phe Pro Ser
1 5 10 15
Ile Thr Val Gln Cys Asn Ser Val Leu Pro Trp Gln Val Thr Ser Asn
20 25 30
Phe Ile Pro Phe Val Cys Val Leu Trp Val Glu Val Glu Tyr Lys Tyr
35 40 45
Gln Val Thr Thr Phe Lys His Asn Asn Leu Ile Ile Ile Ile His Ala
50 55 60
Ala Tyr Tyr Leu Phe Ser
65 70

<210> SEQ ID NO 111
<211> LENGTH: 51
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 111
Met Ala Lys Leu Val Thr His Glu Ile Glu Val Pro Leu Ser Ser Gln
1 5 10 15
Gly His Cys Glu Lys Met Asp His Leu Val Lys Arg Asn Ser Ser Ile
20 25 30
Asn Asn Arg Arg Ser Ile Cys Gln Ala Arg His Ala Arg Ile His Leu
35 40 45
Phe Val His
50

<210> SEQ ID NO 112
<211> LENGTH: 72
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 112
Met Phe Glu Thr Lys Leu Asn Ser Gly Val Val Trp Asn Asp Trp Leu
1 5 10 15
Thr Val Asn Ile Arg Asn Ser Asn Thr Pro Asn Thr Lys Leu Val Leu
20 25 30
Leu His His Val Val Arg Thr Val Pro Ser Ile Glu Ile Ala Asn Asn
35 40 45
Phe Val Phe Leu Ser Ser Arg Ser Pro Phe Thr Ile Asp Tyr Ala Thr
50 55 60
Ile Phe Pro Val Glu Ser Lys Phe
65 70

<210> SEQ ID NO 113
<211> LENGTH: 66
<212> TYPE: PRT
<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 113
Met Leu Tyr Thr Ser Leu Tyr Ile Ser Tyr Leu Ser Asn Ser Met Leu
1 5 10 15

-continued

Leu Pro Ser Trp Thr Asn Leu His His Ser Tyr Ser Leu Asn Asn Leu
 20 25 30

Ser Thr Tyr Leu Gly Leu Pro Leu Pro Gly Gly Asn Gln Asn Gln Phe
 35 40 45

Leu Pro Gln Lys Gln Ala Gly Gln Gly Pro Ala Tyr Gln Lys His Leu
 50 55 60

Arg Gln
 65

<210> SEQ ID NO 114
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 114

gttcagacaa gaccacagat gtga

24

<210> SEQ ID NO 115
 <211> LENGTH: 181
 <212> TYPE: PRT
 <213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 115

Met Arg Asn Leu Phe Pro Ile Leu Met Leu Ile Thr Asn Leu Ala Leu
 1 5 10 15

Asn Asn Asp Asn Asn Asn Asn Asn Asn Asn Asn Asn Tyr Asn Leu
 20 25 30

Ile His Ala Thr Cys Arg Glu Thr Pro Tyr Tyr Ser Leu Cys Leu Thr
 35 40 45

Thr Leu Gln Ser Gly Pro Arg Ser Asn Glu Val Glu Gly Gly Asp Ala
 50 55 60

Ile Thr Thr Leu Gly Leu Ile Met Val Asp Ala Val Lys Ser Lys Ser
 65 70 75 80

Ile Glu Ile Met Glu Lys Ile Lys Glu Leu Glu Lys Ser Asn Pro Glu
 85 90 95

Trp Arg Ala Pro Leu Ser Gln Cys Tyr Val Ala Tyr Asn Ala Val Leu
 100 105 110

Arg Ala Asp Val Thr Val Ala Val Glu Ala Leu Lys Lys Gly Ala Pro
 115 120 125

Lys Phe Ala Glu Asp Gly Met Asp Asp Val Val Ala Glu Ala Gln Thr
 130 135 140

Cys Glu Tyr Ser Phe Asn Tyr Tyr Asn Lys Leu Asp Phe Pro Ile Ser
 145 150 155 160

Asn Leu Ser Arg Glu Ile Ile Glu Leu Ser Lys Val Ala Lys Ser Ile
 165 170 175

Ile Arg Met Leu Leu
 180

<210> SEQ ID NO 116
 <211> LENGTH: 172
 <212> TYPE: PRT
 <213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 116

-continued

```

Met Arg Asn Leu Phe Pro Ile Phe Met Leu Ile Thr Asn Leu Ala Phe
 1           5           10           15
Asn Asp Asn Asn Asn Ser Asn Asn Ile Ile Asn Thr Thr Cys Arg Ala
          20           25           30
Thr Thr Asn Tyr Pro Leu Cys Leu Thr Thr Leu His Ser Asp Pro Arg
      35           40           45
Thr Ser Glu Ala Glu Gly Ala Asp Leu Thr Thr Leu Gly Leu Val Met
      50           55           60
Val Asp Ala Val Lys Leu Lys Ser Ile Glu Ile Met Lys Ser Ile Lys
      65           70           75           80
Lys Leu Glu Lys Ser Asn Pro Glu Leu Arg Leu Pro Leu Ser Gln Cys
          85           90           95
Tyr Ile Val Tyr Tyr Ala Val Leu His Ala Asp Val Thr Val Ala Val
      100           105           110
Glu Ala Leu Lys Arg Gly Val Pro Lys Phe Ala Glu Asn Gly Met Val
      115           120           125
Asp Val Ala Val Glu Ala Glu Thr Cys Glu Phe Ser Phe Lys Tyr Asn
      130           135           140
Gly Leu Val Ser Pro Val Ser Asp Met Asn Lys Glu Ile Ile Glu Leu
      145           150           155           160
Ser Ser Val Ala Lys Ser Ile Ile Arg Met Leu Leu
          165           170

```

<210> SEQ ID NO 117

<211> LENGTH: 181

<212> TYPE: PRP

<213> ORGANISM: Solanum tuberosum

<400> SEQUENCE: 117

```

Met Arg Asn Leu Phe Pro Ile Leu Met Leu Ile Thr Asn Leu Ala Leu
 1           5           10           15
Asn Asn Asp Asn Asn Asn Asn Asn Asn Asn Asn Asn Tyr Asn Leu
          20           25           30
Ile His Ala Thr Cys Arg Glu Thr Pro Tyr Tyr Ser Leu Cys Leu Thr
      35           40           45
Thr Leu Gln Ser Gly Pro Arg Ser Asn Glu Val Glu Gly Gly Asp Ala
      50           55           60
Ile Thr Thr Leu Gly Leu Ile Met Val Asp Ala Val Lys Ser Lys Ser
      65           70           75           80
Ile Glu Ile Met Glu Lys Ile Lys Glu Leu Glu Lys Ser Asn Pro Glu
          85           90           95
Trp Arg Ala Pro Leu Ser Gln Cys Tyr Val Ala Tyr Asn Ala Val Leu
      100           105           110
Arg Ala Asp Val Thr Val Ala Val Glu Ala Leu Lys Lys Gly Ala Pro
      115           120           125
Lys Phe Ala Glu Asp Gly Met Asp Asp Val Val Ala Glu Ala Gln Thr
      130           135           140
Cys Glu Tyr Ser Phe Asn Tyr Tyr Asn Lys Leu Asp Phe Pro Ile Ser
      145           150           155           160
Asn Leu Ser Arg Glu Ile Ile Glu Leu Ser Lys Val Ala Lys Ser Ile
          165           170           175
Ile Arg Met Leu Leu

```

-continued

180

<210> SEQ ID NO 118

<211> LENGTH: 166

<212> TYPE: PRT

<213> ORGANISM: *Nicotiana tabacum*

<400> SEQUENCE: 118

```

Met Lys Asn Leu Ile Phe Leu Thr Met Phe Leu Thr Ile Leu Leu Gln
 1             5             10             15
Thr Asn Ala Asn Asn Leu Val Glu Thr Thr Cys Lys Asn Thr Pro Asn
 20             25             30
Tyr Gln Leu Cys Leu Lys Thr Leu Leu Ser Asp Lys Arg Ser Ala Thr
 35             40             45
Gly Asp Ile Thr Thr Leu Ala Leu Ile Met Val Asp Ala Ile Lys Ala
 50             55             60
Lys Ala Asn Gln Ala Ala Val Thr Ile Ser Lys Leu Arg His Ser Asn
 65             70             75             80
Pro Pro Ala Ala Trp Lys Gly Pro Leu Lys Asn Cys Ala Phe Ser Tyr
 85             90             95
Lys Val Ile Leu Thr Ala Ser Leu Pro Glu Ala Ile Glu Ala Leu Thr
100            105            110
Lys Gly Asp Pro Lys Phe Ala Glu Asp Gly Met Val Gly Ser Ser Gly
115            120            125
Asp Ala Gln Glu Cys Glu Glu Tyr Phe Lys Gly Ser Lys Ser Pro Phe
130            135            140
Ser Ala Leu Asn Ile Ala Val His Glu Leu Ser Asp Val Gly Arg Ala
145            150            155            160
Ile Val Arg Asn Leu Leu
165

```

<210> SEQ ID NO 119

<211> LENGTH: 277

<212> TYPE: DNA

<213> ORGANISM: Unknown Organism

<220> FEATURE:

<223> OTHER INFORMATION: Description of Unknown Organism: P-PPO3
nucleotide sequence

<400> SEQUENCE: 119

```

ctggcgataa cggaactgtt ggaggatatt ggtttgaag atgaagatac tattgcggtg      60
actctggtgc caaagagagg tggatgaagt atctccattg aaagtgcgac gatcagtcct      120
gcagattgtt aattagtctc tattgaatct gctgagatta cactttgatg gatgatgctc      180
tgtttttgtt ttcttgttct gttttttcct ctgttgaaat cagctttgtt gcttgatttc      240
attgaagttg ttattcaaga ataaatcagt tacaatt                                277

```

<210> SEQ ID NO 120

<211> LENGTH: 300

<212> TYPE: DNA

<213> ORGANISM: Unknown Organism

<220> FEATURE:

<223> OTHER INFORMATION: Description of Unknown Organism: PPOM-41
nucleotide sequence

<400> SEQUENCE: 120

```

ctggcgataa cggaactgtt ggaggatatt ggattgaag atgaagatac tattgcggtg      60

```

-continued

acttttggttc caaaagtagg tgggtgaaggt gtatccattg aaagtgtgga gatcaagctt 120
gaggattggtt aagtcctcat gagttggttg ctacggtacc aaattttatg ttttaattagt 180
attaatgtgt gtatgtgttt gattatgttt cggttaaaat gtatcagctg gatagctgat 240
tactagcctt gccagttggtt aatgctatgt atgaaataaa taaataaatg gttgtcttct 300

<210> SEQ ID NO 121
<211> LENGTH: 296
<212> TYPE: DNA
<213> ORGANISM: Unknown Organism
<220> FEATURE:
<223> OTHER INFORMATION: Description of Unknown Organism: PPOM-44
nucleotide sequence
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (54)
<223> OTHER INFORMATION: a, t, c or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (166)
<223> OTHER INFORMATION: a, t, c or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (223)
<223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 121

ctggcgataa cggaactggt ggaggataat ggattggaag atgaaggtac tatngcggta 60
acttttggttc caaaagtttg tgggtgaaggt gtatccattg aaagtgcgga gatcaagctt 120
gaggattggtt aagtcctcat gagttggttg ctatggtacc aaattntatg ttttaattagt 180
attaatgtgt gtgtttgatt atgtttcggg taaaatgtat canctggata gctgattact 240
agccttccca gttgttaatg ctatgtatga aatacataaa taaatggttg tcttcc 296

<210> SEQ ID NO 122
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 122

gtccaacttg cacaggaaag ac 22

<210> SEQ ID NO 123
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 123

catggatgaa atactcctga gc 22

<210> SEQ ID NO 124
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer
<220> FEATURE:
<221> NAME/KEY: modified_base

-continued

```

<222> LOCATION: (8)
<223> OTHER INFORMATION: a, t, c or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (10)
<223> OTHER INFORMATION: a, t, c or g
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (12)
<223> OTHER INFORMATION: a, t, c or g

```

```

<400> SEQUENCE: 124

```

```

gtttacanhn bnatatatcc tgyca

```

25

What is claimed is:

1. A method of modifying a trait of a selected plant comprising:

a. stably transforming cells from said selected plant with a desired polynucleotide, wherein said desired polynucleotide consists essentially of a nucleic acid sequence that is native to said selected plant, native to a plant from the same species, or is native to a plant that is sexually interfertile with said selected plant,

b. obtaining a stably transformed plant from said transformed plant cells wherein said transformed plant contains said desired polynucleotide stably integrated into the genome and wherein said desired polynucleotide modifies said trait.

2. The method according to claim 1, further comprising co-transfecting said plant cells with a selectable marker gene that is transiently expressed in said plant cells, and identifying transformed plant cells, and transformed plants obtained from said transformed plant cells, wherein said selectable marker gene is not stably integrated and said desired polynucleotide is stably integrated into the genome.

3. The method of claim 1, wherein the plant is a monocotyledonous plant.

4. The method of claim 3, wherein said monocotyledonous plant is selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.

5. The method of claim 1, where in the plant is a dicotyledonous plant.

6. The method of claim 5, wherein said dicotyledonous plant is selected from the group consisting of potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsetta, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

7. The method of claim 1, wherein said trait is selected from the group consisting

of enhanced health and nutritional characteristics, improved storage, enhanced yield, enhanced salt tolerance, enhanced heavy metal tolerance, increased drought tolerance, increased disease tolerance, increased insect tolerance, increased water-stress tolerance, enhanced cold and frost tolerance, enhanced color, enhanced sweetness, improved vigor, improved taste, improved texture, decreased phosphate content,

increased germination, increased micronutrient uptake, improved starch composition, improved flower longevity.

8. The method of claim 1, wherein said desired polynucleotide comprises a P- DNA, GBSS promoter, Ubi7 promoter, Ubi3 promoter, PIP promoter, modified PPO gene, invertase inhibitor gene, salt tolerance gene, R1-associated leader, phosphorylase-associated leader, R1-associated trailer, SBE-associated trailers, Ubi-intron, GBSS spacer, UbiT.

9. The method of claim 1, wherein said plant cells are transformed via *Agrobacterium*-mediated transformation.

10. The method of claim 9, wherein *Agrobacterium*-mediated transformation relies on the use of at least one binary vector.

11. The method of claim 10, wherein the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector.

12. The method of claim 11, wherein the first binary vector contains said desired polynucleotide and the second binary vector contains a functional selectable marker gene, wherein said functional selectable marker gene is operably linked to a promoter and a terminator.

13. A plant made by the method of claim 1.

14. A method of modifying a trait in a selected plant comprising:

(a) identifying the trait to be modified;

(b) constructing a first polynucleotide consisting essentially of native genetic elements isolated from said selected plant, a plant from the same species, or a plant that is sexually interfertile with said selected plant, wherein said native genetic elements are capable of modifying the expression of a functional gene that controls said trait

(c) constructing a second polynucleotide comprising a functional selectable marker gene;

(d) co-transfecting plant cells from said selected plant with said first and second polynucleotides;

(e) selecting for the transient expression of said functional selectable marker gene;

(f) screening for plant cells stably transformed with said first polynucleotide but do not contain said second polynucleotide integrated into the genome; and

- (g) obtaining a stably transformed plant from said transformed plant cells that exhibit modified expression of said trait.
15. The method of claim 14, wherein the plant is a monocotyledenous plant.
16. The method of claim 15, wherein said monocotyledenous plant is selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.
17. The method of claim 14, where in the plant is a dicotyledenous plant.
18. The method of claim 17, wherein said dicotyledenous plant is selected from the group consisting of avocado, potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsetta, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.
19. The method of claim 14, wherein said trait is selected from the group consisting of enhanced health and nutritional characteristics, improved storage, enhanced yield, enhanced salt tolerance, enhanced heavy metal tolerance, increased drought tolerance, increased disease tolerance, increased insect tolerance, increased water-stress tolerance, enhanced cold and frost tolerance, enhanced color, enhanced sweetness, improved vigor, improved taste, improved texture, decreased phosphate content, increased germination, increased micronutrient uptake, improved starch composition, improved flower longevity.
20. The method of claim 14, wherein said genetic elements comprise at least one of a promoter, sequence of interest, terminator, enhancer, intron, spacer, or regulatory elements.
21. The method of claim 14, wherein said plant cells are transformed via *Agrobacterium*-mediated transformation.
22. The method of claim 21, wherein *Agrobacterium*-mediated transformation relies on the use of at least one binary vector.
23. The method of claim 22, wherein the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector.
24. The method of claim 23, wherein the first binary vector carries said first polynucleotide and the second binary vector carries said second polynucleotide.
25. A method of modifying the expression of a functional gene in a selected plant comprising:
- constructing a first polynucleotide consisting essentially of native genetic elements isolated from said selected plant, a plant of the same species as said selected plant, or a plant that is sexually interfertile with said selected plant, wherein said-native genetic elements are capable of modifying the expression of said functional gene;
 - constructing a second polynucleotide comprising a selectable marker gene operably linked to a promoter and terminator;
 - co-transfecting plant cells from said selected plant with said first and second polynucleotides;
 - selecting for the transient expression of said selectable marker gene;
 - screening for plant cells stably transformed with said first polynucleotide but do not contain said second polynucleotide integrated into the genome; and
 - obtaining a transformed plant from said transformed plant cells that exhibit modified expression of said gene.
26. The method of claim 25, wherein said plant is a monocotyledenous plant.
27. The method of claim 26, wherein said monocotyledenous plant is selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.
28. The method of claim 25, where in the plant is a dicotyledenous plant.
29. The method of claim 28, wherein said dicotyledenous plant is selected from the group consisting of potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsetta, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.
30. The method of claim 25, wherein said plant cells are transfected via *Agrobacterium*-mediated transformation.
31. The method of claim 30, wherein *Agrobacterium*-mediated transformation relies on the use of at least one binary vector.
32. The method of claim 31, wherein the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector.
33. The method of claim 32, wherein the first binary vector carries said first polynucleotide and the second binary vector carries said second polynucleotide.
34. The method of claim 25, wherein said first polynucleotide comprises at least one of a P-DNA, GBSS promoter, Ubi7 promoter, Ubi3 promoter, PIP promoter, modified PPO gene, PPO-associated trailer, invertase inhibitor gene, salt tolerance gene, R1-associated leader, phosphorylase-associated leader, R1-associated trailer, SBE-associated trailers, Ubi-intron, GBSS spacer, UbiT.
35. The method of claim 25, wherein said second polynucleotide comprises at least one of a selectable marker gene, an omega-mutated virD2 polynucleotide, a codA polynucleotide, and a codA::upp fusion polynucleotide.
36. A plant made by the method of claim 25.
37. A transgenic plant exhibiting modified expression of a trait compared to the non-transgenic plant from which it was derived, wherein said transgenic plant is stably transformed with a desired polynucleotide consisting essentially of native genetic elements isolated from said plant, a plant in the same species, or a plant that is sexually interfertile with said plant, and wherein said polynucleotide modifies the expression of said trait.
38. The plant according to claim 37, wherein said plant is a monocotyledenous plant.
39. The plant according to claim 38, wherein said monocotyledenous plant is selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.
40. The plant according to claim 37, wherein said plant is a dicotyledenous plant.
41. The plant according to claim 40, wherein said dicotyledenous plant is selected from the group consisting of potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet

potato, pepper, cotton, poinsetta, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

42. The trait according to claim 37, wherein said trait is selected from the group consisting of enhanced health and nutritional characteristics, improved storage, enhanced yield, enhanced salt tolerance, enhanced heavy metal tolerance, increased drought tolerance, increased disease tolerance, increased insect tolerance, increased water-stress tolerance, enhanced cold and frost tolerance, enhanced color, enhanced sweetness, improved vigor, improved taste, improved texture, decreased phosphate content, increased germination, increased micronutrient uptake, improved starch composition, improved flower longevity.

43. The desired polynucleotide according to claim 37, wherein said polynucleotide comprises at least one of a P-DNA, GBSS promoter, Ubi7 promoter, Ubi3 promoter, PIP promoter, modified PPO gene, PPO-associated trailer, invertase inhibitor gene, salt tolerance gene, R1-associated leader, phosphorylase-associated leader, R1-associated trailer, SBE-associated trailers, Ubi-intron, GBSS spacer, UbiT.

44. An isolated, border-like nucleotide sequence ranging in size from 20 to 100 bp that shares between 52% and 96% sequence identity with a T-DNA border sequence from *Agrobacterium tumefaciens*.

45. The isolated nucleotide sequence of claim 44 wherein said nucleotide sequence is isolated from a monocotyledonous plant.

46. The isolated nucleotide of claim 45, wherein said monocotyledonous plant is selected from the group consisting of wheat, turf, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.

47. The isolated nucleotide sequence of claim 44, wherein said nucleotide sequence is isolated from a dicotyledonous plant.

48. The isolated nucleotide sequence of claim 47, wherein said dicotyledonous plant is selected from the group consisting of potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsetta, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

49. The nucleotide sequence of claim 44, isolated from potato, which has a nucleotide sequence shown in either SEQ ID NO. 54 or 55.

50. The nucleotide sequence of claim 44, isolated from wheat, which has a nucleotide sequence shown in either SEQ ID NO. 94 or 95.

51. A method of making a plant stably transformed with a desired polynucleotide comprising:

- (a) isolating a P-DNA that is flanked by border-like sequences from said plant wherein said border-like sequences share between 52% and 96% sequence identity with an *Agrobacterium tumefaciens* T-DNA border sequence;
- (b) inserting said desired polynucleotide between said P-DNA border-like sequences to form a P-DNA construct; and
- (c) transforming a plant cell from said plant with said P-DNA construct; and

(d) recovering a plant from said transformed plant cell stably transformed with said P-DNA construct.

52. The method according to claim 51, wherein said P-DNA construct is carried on a vector comprised of a backbone integration marker gene and transformed plant cells are selected that do not contain said backbone integration marker gene.

53. The method according to claim 52, wherein said backbone integration marker gene is selected from the group consisting of is a cytokinin gene, and wherein plant shoots are not selected that exhibit a cytokinin-overproducing phenotype.

54. The method according to claim 53, wherein said backbone integration marker gene is the IPT gene, and plant shoots are not selected that exhibit an abnormal phenotype or cannot develop roots.

55. The method of claim 52, wherein said plant cells are from a monocotyledonous plant.

56. The method of claim 55, wherein said monocotyledonous plant is selected from the group consisting of wheat, turf grass, cereal, maize, rice, oat, wheat, barley, sorghum, orchid, iris, lily, onion, banana, sugarcane, sorghum, and palm.

57. The method of claim 52, wherein the plant cells are from a dicotyledonous plant.

58. The method of claim 57, wherein said dicotyledonous plant is selected from the group consisting of avocado, potato, tobacco, tomato, sugarbeet, broccoli, cassava, sweet potato, pepper, cotton, poinsetta, legumes, alfalfa, soybean, carrot, strawberry, lettuce, oak, maple, walnut, rose, mint, squash, daisy, and cactus.

59. The method of claim 51, wherein said plant cells are transfected via *Agrobacterium*-mediated transformation.

60. The method of claim 59, wherein *Agrobacterium*-mediated transformation relies on the use of at least one binary vector.

61. The method of claim 60, wherein the *Agrobacterium*-mediated transformation method uses a first binary vector and a second binary vector.

62. The method of claim 61, wherein the first binary vector carries said P-DNA construct.

63. The method of claim 61, wherein the second binary vector comprises at least one of a negative selectable marker gene and an omega-mutated virD2 gene, wherein the negative selectable marker gene is positioned within the right T-DNA border and the left T-DNA border, and wherein the omega-mutated virD2 gene is positioned within the backbone of the second binary vector.

64. A vector comprising the nucleotide sequence of claim 44.

65. A P-DNA consisting essentially of, in the 5' to 3'-direction, a first T-DNA border-like sequence, a promoter, a desired polynucleotide sequence operably linked to the promoter, a terminator, and a second T-DNA border-like sequence, wherein the border-like sequences have less than 100% sequence identity with T-DNA border sequences.

66. The P-DNA of claim 65, wherein said T-DNA border-like sequences, said promoter, said desired polynucleotide, and said terminator, are all isolated from the same plant, the same plant species, or plants that are sexually interfertile.

67. The P-DNA of claim 65, further consisting essentially of a selectable marker gene.

68. The P-DNA of claim 67, wherein said T-DNA border-like sequences, said promoter, said desired polynucleotide,

said terminator and said selectable marker gene, are all isolated from the same plant, the same plant species, or plants that are sexually infertile.

69. The P-DNA of claim 65, wherein the desired polynucleotide sequence is a sequence upstream or downstream of the coding region of a gene, wherein the upstream sequence is a leader sequence, and wherein the downstream sequence is a trailer sequence.

70. The P-DNA of claim 69, wherein said T-DNA border-like sequences, said promoter, said leader sequence, said trailer sequence, said terminator and said selectable marker gene are all isolated from the same plant, the same plant species, or plants that are sexually infertile.

71. An isolated nucleotide sequence comprising the GBSS promoter isolated from *S. tuberosum*.

72. The isolated nucleotide sequence of claim 71, which has the nucleotide sequence that is SEQ ID. NO. 6 or 13.

73. A vector comprising the P-DNA of claim 65.

74. A vector according to claim 74, wherein the promoter is a regulatable promoter.

75. A vector according to claim 74, wherein the regulatable promoter is sensitive to temperature.

76. A vector according to claim 75, wherein the regulatable promoter is a wheat wcs120 promoter.

77. A vector according to claim 74, wherein the promoter is under temporal regulation.

78. A vector according to claim 77, wherein the promoter is a carboxylase promoter.

79. A vector according to claim 78, wherein the carboxylase promoter is a maize carboxylase promoter.

80. A vector according to claim 74, wherein the promoter is regulated by any one of abscisic acid, wounding, methyl jasmonate or gibberellic acid.

81. A vector according to claim 80, wherein the promoter is a promoter selected from either a Rab 16A gene promoter, an α -amylase gene promoter or a pin2 gene promoter.

82. A vector according to claim 73, wherein the promoter is a tissue-specific promoter.

83. A vector according to claim 69, wherein the leader sequence is a part of a 5'-untranslated region associated with a gene that is endogenous to a cell of the selected plant species.

84. A vector according to claim 69, wherein the 5'-untranslated region is upstream of a start codon of a gene that is selected from the group consisting of a PPO gene, an R1 gene, a HOS1 gene, a S-adenosylhomocysteine hydrolase gene, a class II cinnamate 4-hydroxylase gene, a cinnamoyl-coenzyme A reductase gene, a cinnamoyl alcohol dehydrogenase gene, a caffeoyl coenzyme A O-methyltransferase gene, an actin depolymerizing factor gene, a Nin88 gene, a Lol p 5 gene, an allergen gene, a P450 hydroxylase gene, an ADP-glucose pyrophosphorylase gene, a proline dehydrogenase gene, an endo-1,4-beta-glucanase gene, a zeaxanthin epoxidase gene, and a 1-aminocyclopropane-1-carboxylate synthase gene.

85. A vector according to claim 69, wherein the trailer sequence is a part of the 3'-untranslated region associated with a gene that is downstream of a termination codon of a gene selected from the group consisting of a PPO gene, an R1 gene, a HOS1 gene, a S-adenosylhomocysteine hydrolase gene, a class II cinnamate 4-hydroxylase gene, a cinnamoyl-coenzyme A reductase gene, a cinnamoyl alcohol dehydrogenase gene, a caffeoyl coenzyme A O-methyltransferase gene, an actin depolymerizing factor gene, a Nin88

gene, a Lol p 5 gene, an allergen gene, a P450 hydroxylase gene, an ADP-glucose pyrophosphorylase gene, a proline dehydrogenase gene, an endo-1,4-beta-glucanase gene, a zeaxanthin epoxidase gene, and a 1-aminocyclopropane-1-carboxylate synthase gene.

86. A vector according to claim 65, further comprising a spacer element that is either an Ubi intron sequence or a GBSS spacer sequence.

87. A vector according to claim 65, wherein the terminator is a Ubi3 terminator sequence or a 3'-untranslated region of an endogenous plant gene.

88. A vector according to claim 65, further comprising a selectable marker gene operably linked to a constitutive promoter and a Cre gene operably linked to an inducible promoter, wherein the selectable marker gene and the Cre gene are flanked by a first recombinase recognition site and a second recombinase recognition site.

89. A vector according to claim 88, wherein the first recombinase recognition site and the second recombinase recognition site are lox sites.

90. A vector according to claim 88, wherein the inducible promoter is a temperature-sensitive promoter, a chemically-induced promoter, or a temporal promoter.

91. A vector according to claim 88, wherein the inducible promoter is a Ha hsp17.7 G4 promoter, a wheat wcs120 promoter, a Rab 16A gene promoter, an α -amylase gene promoter, a pin2 gene promoter, a carboxylase promoter.

92. A vector according to claim 65, further comprising a plant-derived marker gene.

93. A vector according to claim 92, wherein the plant-derived marker gene is an enolpyruvul-3-phosphoshikimic acid synthase gene, a salt-tolerance gene, or the PST1 gene, the PST2 gene, or the PST3 gene.

94. A method for modifying a plant cell, comprising integrating a P-DNA sequence into the genome of a plant cell, wherein the P-DNA consists essentially of, in the 5'- to 3'-direction, a first T-DNA border-like sequence, a promoter, a desired polynucleotide sequence operably linked to the promoter, a terminator, and a second T-DNA border-like sequence, wherein the border-like sequences have less than 100% sequence identity with T-DNA border sequences, and wherein the T-DNA border-like sequences, the promoter, the desired polynucleotide, and terminator, are all isolated from or native to the genome of the plant cell, wherein the desired polynucleotide comprises sense and antisense sequences of a leader sequence or trailer sequence that are associated with the upstream or downstream non-coding regions of a gene in the plant, and wherein expression of the desired polynucleotide produces a double-stranded RNA transcript that targets the gene associated with the desired polynucleotide, thereby modifying the plant cell.

95. A method for modifying a plant, comprising:

- (i) transfecting at least one cell in the plant with the vector of claim 88;
- (ii) selecting a cell expressing the selectable marker;
- (iii) isolating the cell expressing the selectable marker;
- (iii) inducing the expression of the Cre gene in the isolated cell;
- (iv) culturing the isolated cell; and
- (ii) observing the phenotype of cultured cells;

wherein a phenotype that is different to an untransfected plant cell indicates that the target plant cell has been modified.

96. A method according to claim 14 or 95, wherein the selecting step is performed by identifying which cells are resistant to an antibiotic.

97. A method for identifying a target plant cell whose genome contains a P-DNA, comprising co-transfecting a plant target cell with the vector of claim 65 and a second *Agrobacterium*-derived vector that comprises a marker gene flanked by a T-DNA borders or T-DNA border-like sequences and a omega-mutated virD2 gene, wherein the P-DNA of the vector of claim 65 is integrated into the genome of the plant target cell, and wherein no part of the second *Agrobacterium*-derived vector is integrated into the genome of the plant target cell, and wherein the omega-mutated virD2 gene is in the backbone.

98. A method according to claim 97, wherein the marker in the second *Agrobacterium*-derived vector is a neomycin phosphotransferase gene.

99. A method for identifying a target plant cell whose genome contains at least a part of an integration cassette according to claim 98, further comprising selecting cells that survive temporary growth on a kanamycin-containing media, wherein the genomes of the selected cells contain only the integration cassette.

100. A method according to any one of claims 94, 95, or 97, wherein the target plant cell is within a plant.

101. A plant comprising at least one cell whose genome comprises a P-DNA according to claim 65.

102. A plant comprising at least one cell whose genome is artificially manipulated to contain only plant-derived nucleic acids, wherein no cells of the plant contain foreign nucleic acids integrated into the cell genome.

103. The plant of claim 102, wherein the cell is capable of expressing at least one of the plant-derived nucleic acids, which expression modifies a trait associated with the plant.

104. A method for reducing black spot bruising in a selected plant species, comprising (i) integrating into a genome of a selected plant species, a P-DNA, delineated by border-like sequences, comprised only of polynucleotides native to, or isolated from the selected plant species, or comprised of polynucleotides native to, or isolated from a plant species that is sexually compatible with the selected plant species, wherein the P-DNA consists essentially of, in the 5'- to 3'-direction, a promoter; a sense-orientated leader nucleotide sequence from a PPO gene; an antisense-orientated sequence of the leader nucleotide sequence; and a termination sequence, wherein the promoter produces a double-stranded RNA molecule, and wherein the double-stranded RNA molecule brings about a reduction in the expression of the endogenous PPO gene, thereby reducing black spot bruising in the plant.

105. A method for reducing black spot bruising in a selected plant species, comprising integrating into a genome of a selected plant species a P-DNA, delineated by border-like sequences, comprised only of nucleotide sequences native to, or isolated from the selected plant species or comprised of polynucleotides native to, or isolated from a plant species that is sexually compatible with the selected plant species, wherein the P-DNA consists essentially of, in the 5'- to 3'-direction, a promoter; a sense-orientated trailer nucleotide sequence from a PPO gene; an antisense-orientated sequence of the trailer nucleotide sequence; and a termina-

tion sequence, wherein the promoter produces a double-stranded RNA molecule, and wherein the double-stranded RNA molecule brings about a reduction in the expression of the PPO gene, thereby reducing black spot bruising in the plant.

106. A method for reducing cold-induced sweetening in a selected plant species, comprising integrating into a genome of a selected plant species a P-DNA, delineated by border-like sequences, comprised only of nucleotide sequences isolated from the selected plant species or or comprised of polynucleotides native to, or isolated from a plant species that is sexually compatible with the selected plant species, wherein the P-DNA consists essentially of, in the 5'- to 3'-direction, a promoter; a sense-orientated leader nucleotide sequence associated with an R1 gene; an antisense-orientated sequence of the leader nucleotide sequence; and a termination sequence, wherein the promoter produces a double-stranded RNA molecule, and wherein the double-stranded RNA molecule reduces the expression of the R1 gene, thereby reducing cold-induced sweetening in the plant.

107. A method for reducing cold-induced sweetening in a selected plant species, comprising integrating into a genome of a selected plant species an P-DNA, delineated by border-like sequences, comprised only of nucleotide sequences isolated from the selected plant species or comprised of polynucleotides native to, or isolated from a plant species that is sexually compatible with the selected plant species, wherein the P-DNA consists essentially of, in the 5'- to 3'-direction, a promoter; a sense-orientated trailer nucleotide sequence associated with an R1 gene; an antisense-orientated sequence of the trailer nucleotide sequence; and a termination sequence, wherein the promoter produces a double-stranded RNA molecule, and wherein the double-stranded RNA molecule reduces the expression of the R1 gene, thereby reducing cold-induced sweetening in the plant.

108. The method of claim 8, wherein the sequence of interest is a gene.

109. The method of claim 108, wherein the gene is a modified polyphenol oxidase, polyphenol oxidase gene, or a modified R1 gene, or an R1 gene.

110. The method of claim 8, wherein the promoter is an inducible promoter.

111. The method of claim 8, wherein the terminator is a yeast ADH terminator sequence.

112. The method of claim 8, wherein the sequence of interest is a leader or trailer sequence, wherein the leader or trailer sequence represents a sequence upstream or downstream of a gene that is native to the plant cell.

113. The method of claim 112, wherein the sequence of interest comprises a sense-orientated leader sequence operably linked to an antisense leader sequence.

114. The method of claim 112, wherein the sequence of interest comprises a sense-orientated trailer sequence operably linked to an antisense trailer sequence.

115. The method of claim 113, wherein the leader construct comprises in 5'- to 3'-direction, a promoter, a sense-orientated leader sequence, the antisense sequence of the leader, and a terminator, wherein expression of the leader construct produces a double-stranded RNA molecule that facilitates the down-regulation of expression of the gene to which it is associated.

116. The method of claim 115, wherein the leader sequence is associated with, and located upstream of, the

coding region of the PPO gene, the R1 gene, an L-type phosphorylase gene, or an alpha glucan phosphorylase gene.

117. The method of claim 113, wherein the trailer construct comprises in 5'- to 3'-direction, a promoter, a sense-oriented trailer sequence, the antisense sequence of the trailer, and a terminator, wherein expression of the trailer construct produces a double-stranded RNA molecule that facilitates the down-regulation of expression of the gene to which it is associated.

118. The method of claim 117, wherein the trailer sequence is associated with, and located downstream of, the coding region of the PPO gene, the R1 gene, an L-type phosphorylase gene, or an alpha glucan phosphorylase gene.

119. The method of claim 9, further comprising exposing the plant cell to a second vector that comprises a marker element, wherein the marker is transiently expressed in the transformed plant and is not stably integrated into the genome of the transformed plant.

120. The method of claim 119, wherein the marker is a herbicide resistance gene.

121. The method of claim 120, wherein the cytokinin gene is an antibiotic resistance gene.

122. The method of claim 119, wherein the marker is a NPTII.

123. A polynucleotide comprising the polynucleotide sequence of SEQ ID NO. 93, wherein the polynucleotide is between 20 and 80 nucleotides in length.

124. The polynucleotide of claim 123, wherein the polynucleotide is between 21 and 70 nucleotides in length.

125. The polynucleotide of claim 123, wherein the polynucleotide is between 22 and 50 nucleotides in length.

126. The polynucleotide of claim 123, wherein the polynucleotide is between 23 and 40 nucleotides in length.

127. The polynucleotide of claim 123, wherein the polynucleotide is between 24 and 30 nucleotides in length.

128. The method of claim 4, wherein the plant cells are transfected with the first polynucleotide before the second polynucleotide.

129. A tuber-specific promoter as shown in SEQ ID NO. 40.

130. An *Agrobacterium*-based method of making transgenic plant cells that do not contain a selectable marker gene stably integrated in nuclear DNA comprising:

- a. constructing a first binary vector comprised of a polynucleotide consisting essentially of a desired gene operably linked to T-DNA borders or T-DNA border-like sequences at the 5' and 3' ends of said desired gene;
- b. constructing a second binary vector comprised of a selectable marker gene operably linked to T-DNA borders or T-DNA border-like sequences at the 5' and 3' ends of said selectable marker gene;
- c. incubating plants cells with:
 - i. an *Agrobacterium* strain carrying said first and said second binary vectors; or
 - ii. a first *Agrobacterium* strain carrying said first binary vector and a second *Agrobacterium* strain carrying said second binary vector;

d. selecting plant cells wherein said desired gene is integrated into plant nuclear DNA without integration of said selectable marker gene into plant nuclear DNA following incubation for an appropriate time period on a medium containing an appropriate selection agent.

131. The method according to claim 130, wherein said selectable marker gene is a herbicide resistance gene or an antibiotic resistance gene.

132. The method according to claim 131, wherein said antibiotic resistance gene is the nptII gene.

133. The method according to claim 132, wherein said antibiotic resistance gene is the npt II structural gene operably linked to the promoter from the Ubiquitin-7 gene and the terminator from yeast alcohol dehydrogenase 1 (ADH1) gene.

134. The method according to claim 130, wherein said plant cells are first incubated with said first *Agrobacterium* strain and then subsequently incubated with said second *Agrobacterium* strain or vice versa.

135. The method according to claim 130, wherein said first binary vector further comprises a binary integration marker gene that can be used to detect plant cells stably transformed with binary vector backbone sequences.

136. The method according to claim 135, wherein said binary vector integration marker gene is selected from the group consisting of a herbicide resistance gene, antibiotic resistance gene, or NPTII.

137. The method according to claim 130, wherein said second binary vector further comprises a gene fusion between the bacterial cytosine deaminase (codA) and uracil phosphoribosyltransferase (upp) genes, which is inserted between the T-DNA or T-DNA border-like sequences, and plant cells are exposed to 5-fluorocytosine following incubation with said first and second *Agrobacterium* strains in order to select against those plant cells transformed with said second binary vector.

138. The method according to claim 130, wherein said secondary binary vector further comprises the omega-mutated virD2 gene, wherein said omega-mutated virD2 gene reduces the frequency of integration of said selectable marker gene into said plant nuclear DNA.

139. The isolated, border-like nucleotide sequence of claim 44, wherein the sequence is 25 nucleotides in length.

140. The method of claim 61, wherein the second binary vector comprises at least one of a negative selectable marker gene and a gene that impairs integration, wherein the negative selectable marker gene is positioned within the right T-DNA border and the left T-DNA border, and wherein the gene that impairs integration is positioned within the backbone of the second binary vector.

141. The method of claim 99, wherein the temporary growth is from 1 to 5 days.

* * * * *