



(51) International Patent Classification:

C07D 495/14 (2006.01) *A61P 35/00* (2006.01)
A61K 31/437 (2006.01) *A61P 37/00* (2006.01)
A61P 25/00 (2006.01)

(21) International Application Number:

PCT/US20 15/028224

(22) International Filing Date:

29 April 2015 (29.04.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/986,738 30 April 2014 (30.04.2014) US
 61/986,789 30 April 2014 (30.04.2014) US

(71) Applicant: **INCYTE CORPORATION** [US/US]; 1801
 Augustine Cut-Off, Wilmington, Delaware 19803 (US).

(72) Inventors: **ZHOU, Jiacheng**; 46 Bay Boulevard, Newark,
 Delaware 19702 (US). **LIU, Pingli**; 205 Pilot Court, Ne-
 wark, Delaware 19702 (US). **CHEN, Shili**; 810 Colorado
 Drive, Newark, Delaware 19713 (US). **WU, Yongzhong**; 2
 Robert Court, Chadds Ford, Pennsylvania 19317 (US).
WANG, Dengjin; 2205 F Prior Road, Wilmington,
 Delaware 19809 (US). **JIA, Zhongjiang**; 516 Silver Fox
 Road, Kennett Square, Pennsylvania 19348 (US). **QIAO,
 Lei**; 1376 Stonegate Drive, Downingtown, Pennsylvania
 19335 (US). **FRIETZE, William**; 900 Merrybell Lane,
 Kennett Square, Pennsylvania 19348 (US). **XIA, Michael**;
 304 Clubhouse Lane, Wilmington, Delaware 19810 (US).

DAI, Yingrui; 108 Sassafrass Court, Wilmington,
 Delaware 19808 (US).

(74) Agents: **GOODSON, Susanne H.** et al; Fish & Richard-
 son P.C., P.O. Box 1022, Minneapolis, Minnesota 55440-
 1022 (US).


(81) Designated States (*unless otherwise indicated, for every
 kind of national protection available*): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
 BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
 DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
 HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
 KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
 MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
 PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
 SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
 kind of regional protection available*): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
 TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
 TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
 DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
 LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
 SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PROCESSES OF PREPARING A JAK1 INHIBITOR AND NEW FORMS THERETO



1a

(57) Abstract: This invention relates to processes for preparing a JAK1 inhibitor having Formula 1a: as well as new forms of the in-
 hibitor.

PROCESSES OF PREPARING A JAK1 INHIBITOR AND NEW FORMS THERETO

This application claims the benefit of priority of Ser. No. 61/986,738, filed April 30, 2014, and Ser. No. 61/986,789, filed April 30, 2014, each of which is
5 incorporated herein by reference in its entirety.

TECHNICAL FIELD

This invention relates to processes for preparing a JAK1 inhibitor, as well as new forms of the inhibitor.

BACKGROUND

10 Protein kinases (PKs) regulate diverse biological processes including cell growth, survival, differentiation, organ formation, morphogenesis, neovascularization, tissue repair, and regeneration, among others. Protein kinases also play specialized roles in a host of human diseases including cancer. Cytokines, low-molecular weight polypeptides or glycoproteins, regulate many pathways involved in the host
15 inflammatory response to sepsis. Cytokines influence cell differentiation, proliferation and activation, and can modulate both pro-inflammatory and anti-inflammatory responses to allow the host to react appropriately to pathogens. Signaling of a wide range of cytokines involves the Janus kinase family (JAKs) of protein tyrosine kinases and Signal Transducers and Activators of Transcription (STATs). There are four known mammalian JAKs: JAK1 (Janus kinase-1), JAK2,
20 JAK3 (also known as Janus kinase, leukocyte; JAKL; and L-JAK), and TYK2 (protein-tyrosine kinase 2).

Cytokine-stimulated immune and inflammatory responses contribute to pathogenesis of diseases: pathologies such as severe combined immunodeficiency
25 (SCID) arise from suppression of the immune system, while a hyperactive or inappropriate immune/inflammatory response contributes to the pathology of autoimmune diseases (*e.g.*, asthma, systemic lupus erythematosus, thyroiditis, myocarditis), and illnesses such as scleroderma and osteoarthritis (Ortmann, R. A., T. Cheng, *et al.* (2000) *Arthritis Res* 2(1): 16-32).

Deficiencies in expression of JAKs are associated with many disease states. For example, Jak1^{-/-} mice are runted at birth, fail to nurse, and die perinatally (Rodig, S. J., M. A. Meraz, *et al.* (1998) *Cell* 93(3): 373-83). Jak2^{-/-} mouse embryos are anemic and die around day 12.5 postcoitum due to the absence of definitive erythropoiesis.

The JAK/STAT pathway, and in particular all four JAKs, are believed to play a role in the pathogenesis of asthmatic response, chronic obstructive pulmonary disease, bronchitis, and other related inflammatory diseases of the lower respiratory tract. Multiple cytokines that signal through JAKs have been linked to inflammatory diseases/conditions of the upper respiratory tract, such as those affecting the nose and sinuses (*e.g.*, rhinitis and sinusitis) whether classically allergic reactions or not. The JAK/STAT pathway has also been implicated in inflammatory diseases/conditions of the eye and chronic allergic responses.

Activation of JAK/STAT in cancers may occur by cytokine stimulation (*e.g.* IL-6 or GM-CSF) or by a reduction in the endogenous suppressors of JAK signaling such as SOCS (suppressor of cytokine signaling) or PIAS (protein inhibitor of activated STAT) (Boudny, V., and Kovarik, J., *Neoplasia* 49:349-355, 2002). Activation of STAT signaling, as well as other pathways downstream of JAKs (*e.g.*, Akt), has been correlated with poor prognosis in many cancer types (Bowman, T., *et al. Oncogene* 19:2474-2488, 2000). Elevated levels of circulating cytokines that signal through JAK/STAT play a causal role in cachexia and/or chronic fatigue. As such, JAK inhibition may be beneficial to cancer patients for reasons that extend beyond potential anti-tumor activity.

JAK2 tyrosine kinase can be beneficial for patients with myeloproliferative disorders, *e.g.*, polycythemia vera (PV), essential thrombocythemia (ET), myeloid metaplasia with myelofibrosis (MMM) (Levin, *et al. Cancer Cell*, vol. 7, 2005: 387-397). Inhibition of the JAK2V617F kinase decreases proliferation of hematopoietic cells, suggesting JAK2 as a potential target for pharmacologic inhibition in patients with PV, ET, and MMM.

Inhibition of the JAKs may benefit patients suffering from skin immune disorders such as psoriasis, and skin sensitization. The maintenance of psoriasis is believed to depend on a number of inflammatory cytokines in addition to various


chemokines and growth factors (JCI, 113:1664-1675), many of which signal through JAKs (*Adv Pharmacol.* 2000;47:113-74).

Thus, new or improved agents which inhibit kinases such as JAKs are continually needed for developing new and more effective pharmaceuticals that are aimed at augmentation or suppression of the immune and inflammatory pathways (such as immunosuppressive agents for organ transplants), as well as agents for the prevention and treatment of autoimmune diseases, diseases involving a hyperactive inflammatory response (*e.g.*, eczema), allergies, cancer (*e.g.*, prostate, leukemia, multiple myeloma), and some immune reactions (*e.g.*, skin rash or contact dermatitis or diarrhea) caused by other therapeutics. The compounds of the invention, as well as its compositions and methods described herein are directed toward these needs and other ends.

SUMMARY


The present invention provides process of preparing a compound of Formula

Ia:



Ia

comprising reacting a compound of Formula Ib:



Ib

with a strong base in the presence of a first solvent component.

The present application provides an anhydrous form of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile.

The present application also provides a monohydrate form of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile.

5 The present application further provides various intermediates useful in the synthesis of the compound of Formula Ia.

DESCRIPTION OF DRAWINGS

Figure 1 shows a DSC thermogram characteristic of the compound of Example 9.

10 Figure 2 shows a TGA thermogram characteristic of the compound of Example 9.

Figure 3 shows an XRPD pattern characteristic of the compound of Example 9.

Figure 4 shows a DSC thermogram characteristic of the compound of Example 10.

15 Figure 5 shows a TGA thermogram characteristic of the compound of Example 10.

Figure 6 shows an XRPD pattern characteristic of the compound of Example 10.

20 Figure 7 shows a Vapor Sorption/Desorption Isotherm (1st cycle) characteristic of the compound of Example 9.

Figure 8 shows a Vapor Sorption/Desorption Isotherm (4th cycle) characteristic of the compound of Example 9.

Figure 9 shows an XRPD pattern overlay of the compound of Example 9 before and after a Dynamic Vapor Sorption (DVS) study.

25 Figure 10 shows an XRPD pattern characteristic of the compound of Example 9 after 5 days in a humidity chamber set at 75% relative humidity and 40 °C.

Figure 11 shows a TGA thermogram characteristic of the compound of Example 9 after 5 days in a humidity chamber set at 75% relative humidity and 40 °C.

30 Figure 12 shows a DSC thermogram of the compound of Example 9 after 5 days in a humidity chamber set at 75% relative humidity and 40 °C.


Figure 13 shows an XRPD pattern characteristic of the compound of Example 9 after 7 days in a humidity chamber set at 90% relative humidity and 30 °C.

Figure 14 shows a TGA thermogram of the compound of Example 9 after 7 days in a humidity chamber set at 90% relative humidity and 30 °C.

Figure 15 shows a DSC thermogram of the compound of Example 9 after 7 days in a humidity chamber set at 90% relative humidity and 30 °C.


DETAILED DESCRIPTION

The present invention provides, *inter alia*, processes and intermediates for making 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile, which is a compound Formula Ia:



Ia

as well as new forms of the compound of Formula Ia. The compound of Formula Ia includes anhydrous and hydrated forms of the compound. For example, the present application provides a new anhydrous form and a new monohydrate form of the compound of Formula Ia. The monohydrate form has Formula II:



II.

The present application provides an anhydrous form of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile.

In some embodiments, the anhydrous form is substantially isolated. In some embodiments, the anhydrous form is crystalline.

In some embodiments, the anhydrous form is characterized by a differential scanning cal thermogram having an endothermic peak having an onset at about 185


°C. In some embodiments, the anhydrous form is characterized by a differential scanning calorimetry (DSC) thermogram having an endothermic peak at about 190 °C. In some embodiments, the anhydrous form has a DSC thermogram substantially as shown in Figure 1.

5 In some embodiments, the anhydrous form is characterized by a thermogravimetric analysis (TGA) thermogram having a weight loss of about 1.25% up to 200 °C. In some embodiments, the anhydrous form has a TGA thermogram substantially as shown in Figure 2.

10 In some embodiments, the anhydrous form has at least one XRPD peak, in terms of 2-theta, selected from about 8.8°, about 10.5°, about 16.3°, about 17.5°, about 19.1°, about 21.2°, about 24.0°, and about 26.4°. In some embodiments, the anhydrous form has at least two XRPD peaks, in terms of 2-theta, selected from about 8.8°, about 10.5°, about 16.3°, about 17.5°, about 19.1°, about 21.2°, about 24.0°, and about 26.4°. In some embodiments, the anhydrous form has at least three XRPD
15 peaks, in terms of 2-theta, selected from about 8.8°, about 10.5°, about 16.3°, about 17.5°, about 19.1°, about 21.2°, about 24.0°, and about 26.4°. In some embodiments, the anhydrous form has at least four XRPD peaks, in terms of 2-theta, selected from about 8.8°, about 10.5°, about 16.3°, about 17.5°, about 19.1°, about 21.2°, about 24.0°, and about 26.4°. In some embodiments, the anhydrous form has at least four
20 XRPD peaks, in terms of 2-theta, selected from about 8.8°, about 16.3°, about 19.1°, about 24.0°, and about 26.4°.

 In some embodiments, the anhydrous form has an XRPD pattern substantially as shown in Figure 3. In some embodiments, the anhydrous form has a Vapor Sorption/Desorption Isotherm substantially as shown in Figure 7. In some
25 embodiments, the anhydrous form has a Vapor Sorption/Desorption Isotherm substantially as shown in Figure 8.

 The present application also provides a monohydrate form of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile monohydrate, which is a compound of Formula II:



II.

In some embodiments, the monohydrate form has a water content of about 5%.

In some embodiments, the monohydrate form is substantially isolated. In some
5 embodiments, the monohydrate form is crystalline.

In some embodiments, the monohydrate form is characterized by a DSC
thermogram having an endothermic peak having an onset at about 69 °C. In some
embodiments, the monohydrate form is characterized by a DSC thermogram having
an endothermic peak at about 106 °C.

10 In some embodiments, the monohydrate form is characterized by a DSC
thermogram having an endothermic peak having an onset at about 131 °C. In some
embodiments, the monohydrate form is characterized by a DSC thermogram having
an endothermic peak at about 140 °C. In some embodiments, the monohydrate form
has a DSC thermogram substantially as shown in Figure 4.

15 In some embodiments, the monohydrate form is characterized by a TGA
thermogram having a weight loss of about 5% up to about 150 °C. In some
embodiments, the monohydrate form has a TGA thermogram substantially as shown
in Figure 5.

In some embodiments, the monohydrate form has at least one XRPD peak, in
20 terms of 2-theta, selected from about 8.5°, about 8.7°, about 14.6°, about 15.1°, about
19.4°, about 25.1°, and about 26.5°. In some embodiments, the monohydrate form
has at least two XRPD peaks, in terms of 2-theta, selected from about 8.5°, about 8.7°,
about 14.6°, about 15.1°, about 19.4°, about 25.1°, and about 26.5°. In some
embodiments, the monohydrate form has at least three XRPD peaks, in terms of 2-
25 theta, selected from about 8.5°, about 8.7°, about 14.6°, about 15.1°, about 19.4°,
about 25.1°, and about 26.5°. In some embodiments, the monohydrate form has at
least four XRPD peaks, in terms of 2-theta, selected from about 8.5°, about 8.7°,
about 14.6°, about 15.1°, about 19.4°, about 25.1°, and about 26.5°. In some


embodiments, the monohydrate form has an XRPD pattern substantially as shown in Figure 6.

An XRPD pattern of reflections (peaks) is typically considered a fingerprint of a particular crystalline form. It is well known that the relative intensities of the XRPD peaks can widely vary depending on, *inter alia*, the sample preparation technique, crystal size distribution, various filters used, the sample mounting procedure, and the particular instrument employed. In some instances, new peaks may be observed or existing peaks may disappear, depending on the type of the instrument or the settings. As used herein, the term "peak" refers to a reflection having a relative height/intensity of at least about 4% of the maximum peak height/intensity. Moreover, instrument variation and other factors can affect the 2-theta values. Thus, peak assignments, such as those reported herein, can vary by plus or minus about 0.2° (2-theta), and the term "substantially" and "about" as used in the context of XRPD herein is meant to encompass the above-mentioned variations.

In the same way, temperature readings in connection with DSC, TGA, or other thermal experiments can vary about ± 3 °C depending on the instrument, particular settings, sample preparation, etc. Accordingly, a crystalline form reported herein having a DSC thermogram "substantially" as shown in any of the Figures or the term "about" is understood to accommodate such variation.

The present application further provides a process of forming the anhydrous form of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile, comprising:

- (i) adding n-heptane to a first mixture comprising a compound of Formula II:



II

and a first solvent component, wherein the first solvent component comprises ethyl acetate.

In some embodiments, the first solvent component further comprises methanol.


In some embodiments, the process further comprises before (i):

- (ii) dissolving the compound of Formula II in methanol to form a second mixture comprising the compound of Formula Ia and a second solvent component comprising methanol;
- (iii) polish filtering the second mixture; and
- (iv) adding ethyl acetate to the second mixture to form the first mixture.

In some embodiments, the process further comprises heating the first mixture to a temperature from about 60 °C to about 75 °C.


In some embodiments, the process further comprises removing at least a portion of the ethyl acetate and methanol to form the first mixture via distillation.

The present application further provides a process of preparing a compound of Formula Ia:



Ia

comprising reacting a compound of Formula Ib:



Ib

with a strong base in the presence of a first solvent component.

In some embodiments, the strong base is an alkoxide base.

In some embodiments, the strong base is an alkali metal alkoxide.

In some embodiments, the strong base is potassium tert-butoxide.

In some embodiments, the first solvent component comprises isopropyl alcohol.

In some embodiments, the first solvent component comprises isopropyl alcohol and tetrahydrofuran.


5 In some embodiments, the reacting of the composition with the strong base is conducted at a temperature of about room temperature.

In some embodiments, wherein about 0.05 to about 0.15 equivalents of strong base is used based on 1 equivalent of the compound of Formula Ib.

10 In some embodiments, the compound of Formula Ib is present in a composition comprising the compound of Formula Ib and the compound of Formula Ia prior to said reacting of said compound of Formula Ib with said strong base.


In some embodiments, the strong base that is reacted with a compound of Formula Ib is referred to as a first strong base.

15 In some embodiments, the compound of Formula Ia is prepared as a compound of Formula II:




II.

20 In some embodiments, the composition comprising a compound of Formula Ia and a compound of Formula Ib is prepared by a process comprising reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb:




IIa



IIIb

with a compound of Formula IV:



in the presence of a **Ci-6** alkyloxonium reagent and a second solvent component.

In some embodiments, before reacting, the compound of Formula IV is
 5 reacted with said **Ci-6** alkyloxonium reagent in the second solvent component.

In some embodiments, the reacting step is referred to as the reacting of a composition comprising a compound of Formula IIa and a compound of Formula IIIb.

In some embodiments, the **Ci-6** alkyloxonium reagent is triethyloxonium
 10 tetrafluoroborate.

In some embodiments, the second solvent component comprises ethanol.

In some embodiments, the second solvent component comprises tetrahydrofuran.


In some embodiments, the second solvent component comprises ethanol and
 15 tetrahydrofuran.

In some embodiments, the reacting of the composition of the compound of Formula IIa and the compound of Formula IIIb with the compound of Formula IV is conducted at a temperature at reflux.


In some embodiments, about 2 to about 4 equivalents of the compound of
 20 Formula IV is used based on 1 equivalent of the combined amount of the compound of Formula IIa and the compound of Formula IIIb.

In some embodiments, about 2 to about 4 equivalents of the alkyloxonium reagent is used based on 1 equivalent of the combined amount of the compound of Formula IIa and the compound of Formula IIIb.

In some embodiments, the composition comprising a compound of Formula IIa and a compound of Formula IIIb is prepared by a process comprising reacting a composition comprising a compound of Formula Va and a compound of Formula Vb:
 25



Va



Vb

with hydrogen gas in the presence of a hydrogenation catalyst and a third solvent component.

5 In some embodiments, the hydrogenation catalyst is palladium on carbon.


In some embodiments, the third solvent component comprises 2,2,2-trifluoroethanol.

In some embodiments, the reacting is run at a temperature from about 20 °C to about 35 °C.

10 In some embodiments, the reacting step is referred to as the reacting of a composition comprising a compound of Formula Va and a compound of Formula Vb.


In some embodiments, about 5 to about 15 weight % of the hydrogenation catalyst is used based on total mass of the compound of Formula Va and the compound of Formula Vb.

15 In some embodiments, the composition comprising a compound of Formula Va and a compound of Formula Vb is prepared by a process comprising reacting a compound of Formula VI:



VI

20 or a salt thereof, with a compound of Formula VII:




VII

in the presence of an tertiary amine base and a fourth solvent component, wherein:

X is a halo group.

In some embodiments, the compound of Formula VI, or salt thereof, is a salt of Formula Via:



Via.

5 In some embodiments, X is chloro.

In some embodiments, the tertiary amine base is N-ethyl-N-isopropylpropan-2-amine.


In some embodiments, the tertiary amine base that is present in the reaction of the compound of Formula VI is referred to as a first tertiary amine base.

10 In some embodiments, the fourth solvent component comprises N,N-dimethylformamide.

In some embodiments, the reacting is run at a temperature from about 65 °C to about 75 °C.

15 In some embodiments, the reacting step is referred to as the reacting of a compound of Formula VI.


In some embodiments, the compound of Formula VI, or salt thereof, is a salt of Formula Via:



Via;

20 the fourth solvent component comprises N,N-dimethylformamide;
the tertiary amine base is N-ethyl-N-isopropylpropan-2-amine; and
X is chloro.

The present application further provides a process of preparing a compound of Formula VI, comprising reacting a compound of Formula VIII:



under deprotection conditions to form the compound of Formula VI, or a salt thereof;

wherein P^1 is an amine protecting group.

5 In some embodiments, P^1 is t-butoxycarbonyl.

In some embodiments, the deprotection conditions involve reacting the compound of Formula VIII with a strong acid.


In some embodiments, the strong acid is HCl.

10 In some embodiments, the strong acid that is reacted with a compound of Formula VIII is referred to as a first strong acid.

In some embodiments, the reacting is run at a temperature from about 0 °C to about 40 °C.

In some embodiments, the reacting step is referred to as the reacting of a compound of Formula VIII.

15 In some embodiments, the compound of Formula VIII is prepared by a process comprising reacting a compound of Formula IX:



20 in the presence of a strong base and a fifth solvent component, wherein indicates a *cis*- or *trans*- bond; and P^1 is an amine protecting group. In some embodiments, the P^1 is t-butoxycarbonyl.

In some embodiments, the strong base is a C₆-alkoxide base.

In some embodiments, the alkoxide base is an alkali metal C₆-alkoxide.

In some embodiments, the alkoxide base is sodium methoxide.

25 In some embodiments, the strong base is present in a catalytic amount.

In some embodiments, the catalytic amount comprises about 0.010 to about 0.020 equivalents of the strong base based on 1 equivalent of the compound of Formula IX.

In some embodiments, the strong base that is present in the reaction of the compound of Formula IX is referred to as a second strong base.

In some embodiments, the fifth solvent component comprises tetrahydrofuran.

In some embodiments, the reacting is run at a temperature from about 0 °C to 5 °C.

In some embodiments, the reacting step is referred to as the reacting of a compound of Formula IX.

In some embodiments, the compound of Formula IX is prepared by a process comprising reacting a compound of Formula X:



in the presence of a strong acid and a sixth solvent component, wherein indicates a *cis*- or *trans*- bond; and P¹ is an amine protecting group. In some embodiments, P¹ is t-butoxycarbonyl.

15 In some embodiments, the strong acid is p-toluenesulfonic acid.

In some embodiments, the strong acid is present in a catalytic amount.

In some embodiments, the catalytic amount comprises about 0.005 to about 0.015 equivalents of the strong acid based on 1 equivalent of the compound of Formula X.


20 In some embodiments, the strong acid that is present in the reaction of the compound of Formula X is referred to as a second strong acid.

In some embodiments, the sixth solvent component comprises methanol.


In some embodiments, the reacting of the compound of Formula X in the presence of the strong acid is run at a temperature of about room temperature.

25 In some embodiments, the compound of Formula X is prepared by a process comprising:

(i) reacting a compound of Formula XII:



with a Wittig reagent in the presence of a seventh solvent component, wherein said Wittig reagent is prepared by a process comprising reacting a compound of Formula XXI:



XXI

in the presence of a strong base, wherein R^a and R^b are each independently Ci-6 alkyl; and

P¹ is an amine protecting group. In some embodiments, P¹ is t-butoxycarbonyl.

In some embodiments, before (i), the compound of Formula XXI is reacted with the strong base in said second solvent component.

In some embodiments, R^a and R^b are each ethyl.

In some embodiments, the strong base is potassium tert-butoxide.

In some embodiments, the strong base that is present in the reaction of the compound of Formula XXI is referred to as a base.


In some embodiments, the seventh solvent component comprises tetrahydrofuran.

In some embodiments, the reacting is run at a temperature at about 0 °C to about 5 °C.

In some embodiments, the reacting step is referred to as the reacting of a compound of Formula XII.

In some embodiments, about 1.0 to about 2.0 equivalents of the compound of Formula XXI is used based on 1 equivalent of the compound of Formula XII.

In some embodiments, the compound of Formula XII is prepared by a process comprising oxidizing a compound of Formula XIII:



XIII

wherein P¹ is an amine protecting group. In some embodiments, P¹ is t-butoxycarbonyl.

In some embodiments, the oxidizing comprises adding a first oxidizing agent to the compound of Formula XIII to form a first mixture.

In some embodiments, the first oxidizing agent is 2,2,6,6-tetramethyl-1-piperidinyloxy free radical (TEMPO).

5 In some embodiments, the adding of said first oxidizing agent is conducted in the presence of tetra-N-butylammonium chloride.


In some embodiments, the adding of said first oxidizing agent is run at a temperature of about room temperature.

10 In some embodiments, the oxidizing further comprises adding a base and a second oxidizing agent to said first mixture.

In some embodiments, the second oxidizing agent is an N-halosuccinimide compound.


In some embodiments, the second oxidizing agent is N-chlorosuccinimide.

15 In some embodiments, the compound of Formula XIII is prepared by a process comprising reacting a compound of Formula XV:



XV

with a compound of Formula XIV:



XIV


20

in the presence of boron trifluoride diethyl etherate and an eighth solvent component, wherein P¹ is an amine protecting group. In some embodiments, P¹ is t-butoxycarbonyl.

In some embodiments, the eighth solvent component comprises acetone.

25 In some embodiments, the reacting is run at a temperature of about room temperature.

In some embodiments, the compound of Formula XV is prepared by a process comprising reacting a compound of Formula XVI:



XVI

30

in the presence of a reducing agent and a ninth solvent component, wherein P¹ is an amine protecting group. In some embodiments, P¹ is t-butoxycarbonyl.

In some embodiments, the reducing agent is lithium borohydride.


In some embodiments, the reducing agent that is present in the reaction of the compound of Formula XVI is referred to as a first reducing agent.

In some embodiments, the ninth solvent component comprises tetrahydrofuran.

In some embodiments, the reacting is run at a temperature from about 0 °C to about 40 °C.

In some embodiments, the reacting step is referred to as the reacting of a compound of Formula XVI.

In some embodiments, the compound of Formula VIII is prepared by a process comprising reacting a compound of Formula XVII:



XVII

with hydrogen gas in the presence of a hydrogenation catalyst and a tenth solvent component;

wherein P¹ is an amine protecting group.

The \curvearrowright group in Formula XVII indicates that the CN group can be E or Z relative to the oxygen of the pyran ring. In some embodiments, P¹ is t-butoxycarbonyl.


In some embodiments, the hydrogenation catalyst is palladium on carbon.

In some embodiments, about 5 to about 15 weight % of the hydrogenation catalyst is used based on total mass of the compound of Formula XVII.

In some embodiments, the tenth solvent component comprises methanol.


In some embodiments, the compound of Formula XVII is prepared by

(i) reacting a compound of Formula XVIII:



XVIII

with a Wittig reagent in the presence of an eleventh solvent component, wherein said Wittig reagent is prepared by a process comprising reacting a compound of Formula XXII:



XXII

with a strong base, wherein:

R^{a1} , R^{b1} , and R^{c1} are each independently C₁-6 alkyl;

Y^- is a halide ion; and

P^1 is an amine protecting group.

In some embodiments, P^1 is t-butoxycarbonyl.

In some embodiments, Y^- is Cl^- .

In some embodiments, R^{a1} , R^{b1} , and R^{c1} are each methyl.

In some embodiments, R^{a1} , R^{b1} , and R^{c1} are each n-butyl.

In some embodiments, the strong base is potassium tert-butoxide.

In some embodiments, the strong base is lithium hexamethyldisilazide.


In some embodiments, the strong base that is reacted with a compound of Formula XXII is referred to as a fourth strong base.

In some embodiments, the eleventh solvent component comprises tetrahydrofuran.

In some embodiments, the eleventh solvent component comprises N,N-dimethylacetamide.

In some embodiments, the eleventh solvent component comprises tetrahydrofuran and N,N-dimethylacetamide.

In some embodiments, the compound of Formula XVIII is formed by a process comprising reacting a compound of Formula XIX:



XIX

in the presence of a weak organic acid and a twelfth solvent component, wherein P¹ is an amine protecting group. In some embodiments, P¹ is t-butoxycarbonyl.

5 In some embodiments, the weak organic acid is acetic acid.


In some embodiments, the twelfth solvent component comprises toluene.

In some embodiments, the reacting is run at a temperature at reflux.

In some embodiments, the reacting step is referred to as the reacting of a compound of Formula XIX.


10 In some embodiments, the compound of Formula XIX is formed by a process comprising:

(i) reacting a compound of Formula XX:




XX

15 with a compound of Formula XXIII:



XXIII

in the presence of an amine base and an thirteenth solvent component to form a compound of Formula XXIV:



XXIV

wherein

R² is C1-4 alkoxy;

X¹ is halo; and

25 P¹ is an amine protecting group.

In some embodiments, P¹ is t-butoxycarbonyl.

In some embodiments, the compound of Formula XXIII is ethyl chloroformate.

In some embodiments, the amine base is N-methylmorpholine.

5 In some embodiments, the amine base that is present in the reaction of the compound of Formula XX is referred to as a second tertiary amine base.

In some embodiments, the thirteenth solvent component comprises tetrahydrofuran.

In some embodiments, the reacting is run at a temperature at about 0 °C.

10 In some embodiments, the reacting step is referred to as the reacting of a compound of Formula XX.

In some embodiments, the process further comprises reducing said compound of Formula XXIV with a reducing agent.

In some embodiments, the reducing agent is sodium borohydride.

15 In some embodiments, the reducing agent is dissolved in a fourteenth solvent component.



In some embodiments, the reducing agent that is present in the reducing of the compound of Formula XXIV is referred to as a second reducing agent.

In some embodiments, the fourteenth solvent component comprises diglyme.


20 In some embodiments, the reducing is run at a temperature from about 70 °C to about 80 °C.

In some embodiments, the reducing step is referred to as the reducing of a compound of Formula XXIV.

In some embodiments, the composition comprising a compound of Formula Ia and a compound of Formula Ib is prepared by a process comprising reacting a composition comprising a compound of Formula IIa and a compound of Formula IIb:



with (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole:




or a salt thereof.

In some embodiments, the reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb is performed at a temperature from about 60 °C to about 70 °C.

In some embodiments, the reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb is performed in the presence of a fifteenth solvent component.

In some embodiments, the fifteenth solvent component comprises methanol.

In some embodiments, the (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole is prepared by reacting (*R*)-2,2,5-trimethyloxazolidin-4-one:



with trimethyloxonium tetrafluoroborate.

In some embodiments, about 1 equivalent of trimethyloxonium tetrafluoroborate is used based on 1 equivalent of (*R*)-2,2,5-trimethyloxazolidin-4-one.

In some embodiments, the reacting of (*R*)-2,2,5-trimethyloxazolidin-4-one is performed at about room temperature.


In some embodiments, the reacting of (*R*)-2,2,5-trimethyloxazolidin-4-one is performed in the presence of a sixteenth solvent component.

In some embodiments, the sixteenth solvent component comprises anhydrous dichloromethane.


In some embodiments, the steps of reacting (*R*)-2,2,5-trimethyloxazolidin-4-one with trimethyloxonium tetrafluoroborate to form (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole and reacting (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole with a composition comprising a compound of Formula IIa and a compound of

Formula IIIb are conducted in the same pot without isolation of (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole.

In some embodiments, the composition comprising a compound of Formula Ia and a compound of Formula Ib is prepared by a process comprising reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb:



with (*R*)-ethyl 2-hydroxypropanimidate:



In some embodiments, the reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb is performed at a temperature from about 75 °C to about 85 °C.

In some embodiments, the composition comprising a compound of Formula IIa and a compound of Formula IIIb further comprises a seventeenth solvent component.


In some embodiments, the seventeenth solvent component comprises ethanol.

In some embodiments, prior to the reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb, the (*R*)-ethyl 2-hydroxypropanimidate is dissolved in an eighteenth solvent component.

In some embodiments, the eighteenth solvent component comprises tetrahydrofuran.

In some embodiments, prior to the reacting the composition comprising a compound of Formula IIa and a compound of Formula IIIb, the composition comprising a compound of Formula IIa, a compound of Formula IIIb, and the seventeenth solvent component is heated at a temperature from about 75 °C to about 85 °C.

In some embodiments, the (*R*)-ethyl 2-hydroxypropanimidate is prepared by reacting (*R*)-1-cyanoethyl acetate:



with ethanol in the presence of a strong acid.

In some embodiments, the strong acid used in the reaction of (*R*)-1-cyanoethyl acetate with ethanol is hydrogen chloride.

In some embodiments, about 2.9 to about 3.1 equivalents of ethanol is used based on 1 equivalent of (*R*)-ethyl 2-hydroxypropanimidate.


In some embodiments, the (*R*)-1-cyanoethyl acetate and ethanol are combined to form an ethanol mixture.

In some embodiments, the strong acid is added to the ethanol mixture to form a further mixture.

In some embodiments, the strong acid is added at a temperature from about -45 °C to about -35 °C.

In some embodiments, adding the strong acid to the ethanol mixture to form a further mixture further comprises heating the further mixture to room temperature.

In some embodiments, the (*R*)-1-cyanoethyl acetate is formed by reacting (*R*)-1-amino-1-oxopropan-2-yl acetate:



with cyanuric chloride.

In some embodiments, about 0.4 to about 0.5 equivalents of cyanuric chloride is used based on 1 equivalent of (*R*)-1-amino-1-oxopropan-2-yl acetate.

In some embodiments, prior to the reacting of (*R*)-1-amino-1-oxopropan-2-yl acetate, the (*R*)-1-amino-1-oxopropan-2-yl acetate is mixed with a nineteenth solvent component.


In some embodiments, the nineteenth solvent component comprises *N,N*-dimethylformamide.

In some embodiments, prior to the reacting of (*R*)-1-amino-1-oxopropan-2-yl acetate, the cyanuric chloride is dissolved in a twentieth solvent component.

In some embodiments, the twentieth solvent component comprises 2-methoxy-2-methylpropane.

In some embodiments, the reacting of (*R*)-1-amino-1-oxopropan-2-yl acetate is performed at about room temperature.

In some embodiments, the (*R*)-1-amino-1-oxopropan-2-yl acetate is prepared by protecting (*R*)-2-hydroxypropanamide:



with acetyl chloride in the presence of an third tertiary amine base.

In some embodiments, the third tertiary amine base is 4-methylmorpholine.


In some embodiments, about 1 to about 1.1 equivalents of acetyl chloride is used based on 1 equivalent of (*R*)-2-hydroxypropanamide.

In some embodiments, the protecting is performed in the presence of a twenty-first solvent component.

In some embodiments, the twenty-first solvent component comprises tetrahydrofuran.

The present application further provides intermediates useful in the preparation of the compound of Formula Ia.

Accordingly, the present application provides a compound of Formula VIII:




5

wherein P¹ is an amine protecting group.


In some embodiments, P¹ is t-butoxycarbonyl.

The present application also provides a salt of Formula Via:



Via.

The present application further provides a compound of Formula Va:




Va

5

or salt thereof.

The present application further provides a compound of Formula Vb:




Vb

10

or salt thereof.

The present application further provides a compound of Formula IIa:




IIa

or salt thereof.

15

The present application further provides a compound of Formula IIIb:



IIIb

26

or salt thereof.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment (while the embodiments are intended to be
5 combined as if written in multiply dependent form). Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

At various places in the present specification, substituents of compounds of the invention are disclosed in groups or in ranges. It is specifically intended that the
10 invention include each and every individual subcombination of the members of such groups and ranges. For example, the term "Ci-6 alkyl" is specifically intended to individually disclose methyl, ethyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, and C₆ alkyl.

As used here, the term "Wittig reagent" refers to a ylide formed by the reaction of a phosphonate compound of formula $P(=O)(\text{Ci-6 alkoxy})_2(\text{Ci-6 alkyl-CN})$,
15 wherein the alkoxy and alkyl groups each have 1 to 6 carbons, in the presence of a strong base (e.g., potassium tert-butoxide). In some embodiments, the term "Wittig reagent" refers a ylide formed by the reaction of a phosphonium salt (e.g., cyanomethyl (trimethyl)phosphonium chloride or tributyl(cyanomethyl)phosphonium chloride) in the presence of a strong base (e.g., potassium tert-butoxide).

As used herein, the term "C_{n-m} alkyl", employed alone or in combination with
20 other terms, refers to a saturated hydrocarbon group that may be straight-chain or branched, having n to m carbon atoms. In some embodiments, the alkyl group contains 1 to 6, 1 to 4 or 1 to 3 carbon atoms. Examples of alkyl moieties include, but are not limited to, chemical groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl,
25 isobutyl, sec-butyl, tert-butyl, w-pentyl, 2-methyl-1-butyl, 3-pentyl, w-hexyl, 1,2,2-trimethylpropyl, and the like.

As used herein, the term "Ci-6 alkoxy", employed alone or in combination with other terms, refers to a group of formula -O-alkyl, wherein the alkyl group has 1
30 to 6 carbons. Example alkoxy groups include methoxy, ethoxy, propoxy (e.g., n-propoxy and isopropoxy), butoxy (e.g., n-butoxy, tert-butoxy), and the like.

As used herein, the term "Ci-6 alkyloxonium reagent" refers to a reagent having a cation of group of formula $(\text{Ci-6 alkyl})_3\text{O}^+$, wherein the Ci-6 alkyl group

refers to a linear or branched alkyl group having 1 to 6 carbons. Example **Ci-6** alkyloxonium reagents include trimethyloxonium tetrafluoroborate, triethyloxonium tetrafluoroborate (Meerwein's reagent), triethyloxonium hexafluorophosphate, triethyloxonium hexachloroantimonate, and the like. Counter ions include, but are not
5 limited to tetrafluoroborate, hexafluorophosphate, and hexachloroantimonate.

As used herein, the term "amino" refers to a group of formula -NH_2 .

As used herein, the term "amine base" refers to a mono-substituted amine group (i.e., primary amine base), di-substituted amine group (i.e., secondary amine base), or a tri-substituted amine group (i.e., tertiary amine base). Example mono-
10 substituted amine bases include methyl amine, ethyl amine, propyl amine, butyl amine, and the like. Example di-substituted amine bases include dimethylamine, diethylamine, dipropylamine, dibutylamine, pyrrolidine, piperidine, azepane, morpholine, and the like. In some embodiments, the tertiary amine has the formula N(R')_3 , wherein each R' is independently **Ci-6** alkyl, 3-10 member cycloalkyl, 4-10
15 membered heterocycloalkyl, 1-10 membered heteroaryl, and 5-10 membered aryl, wherein said 3-10 member cycloalkyl, 4-10 membered heterocycloalkyl, 1-10 membered heteroaryl, and 5-10 membered aryl are optionally substituted by 1, 2, 3, 4, 5, or 6 **Ci-6** alkyl groups. Example tertiary amine bases include trimethylamine, triethylamine, tripropylamine, triisopropylamine, tributylamine, tri-tert-butylamine,
20 N,N-dimethylethanamine, N-ethyl-N-methylpropan-2-amine, N-ethyl-N-isopropylpropan-2-amine, N-methylmorpholine, and the like. In some embodiments, the term "tertiary amine base" refers to a group of formula N(R)_3 , wherein each R is independently a linear or branched **Ci-6** alkyl group.

As used herein, the term "cycloalkyl", employed alone or in combination with
25 other terms, refers to a non-aromatic cyclic hydrocarbon moiety, which may optionally contain one or more alkenylene groups as part of the ring structure. In some embodiments, cycloalkyl is a 3-10 membered cycloalkyl, which is monocyclic or bicyclic. In some embodiments, cycloalkyl is a 3-6 or 3-7 monocyclic cycloalkyl. Exemplary cycloalkyl groups include 1,2,3,4-tetrahydro-naphthalene, cyclopropyl,
30 cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, adamantyl, and

the like. In some embodiments, the cycloalkyl group is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, the term "aryl", employed alone or in combination with other terms, refers to a monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings) aromatic hydrocarbon, such as, but not limited to, phenyl, 1-naphthyl, 2-naphthyl, anthracenyl, phenanthrenyl, and the like. In some embodiments, aryl is C₆₋₁₀ aryl. In some embodiments, the aryl group is a naphthalene ring or phenyl ring. In some embodiments, the aryl group is phenyl.

As used herein, the term "heteroaryl", employed alone or in combination with other terms, refers to a monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings) aromatic hydrocarbon moiety, having one or more heteroatom ring members selected from nitrogen, sulfur and oxygen. In some embodiments, heteroaryl is a 5-10 membered heteroaryl, which is monocyclic or bicyclic, comprising 1 to 9 carbon atoms and 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur and oxygen. Example heteroaryl groups include, but are not limited to, pyridine, pyrimidine, pyrazine, pyridazine, pyrrole, pyrazole, azolyl, oxazole, thiazole, imidazole, furan, thiophene, quinoline, isoquinoline, indole, benzothiophene, benzofuran, benzisoxazole, imidazo[1,2-b]thiazole, purine, or the like.

As used herein, the term "heterocycloalkyl", employed alone or in combination with other terms, refers to non-aromatic ring system, which may optionally contain one or more alkenylene or alkynylene groups as part of the ring structure, and which has at least one heteroatom ring member independently selected from nitrogen, sulfur and oxygen. When the heterocycloalkyl groups contains more than one heteroatom, the heteroatoms may be the same or different. Heterocycloalkyl groups can include mono- or polycyclic (e.g., having 2, 3 or 4 fused, spirocyclic, or bridged rings) ring systems. In some embodiments, heterocycloalkyl is 5-10 membered heterocycloalkyl, which is monocyclic or bicyclic, comprising 2 to 9 carbon atoms and 1, 2, 3, or 4 heteroatom ring members independently selected from nitrogen, sulfur, and oxygen. Examples of heterocycloalkyl groups include 1,2,3,4-tetrahydro-quinoline, azetidene, azepane, pyrrolidine, piperidine, piperazine, morpholine, thiomorpholine, pyran, and a 2-oxo-1,3-oxazolidine ring.

As used herein, the term "alkali metal" includes lithium, sodium, and potassium.

As used herein, "halo" or "halogen", employed alone or in combination with other terms, includes fluoro, chloro, bromo, and iodo.

5 Example "N-halosuccinimide compound" include, but are not limited to, N-chlorosuccinimide, N-bromosuccinimide, and N-iodosuccinimide.

As used herein, the term "alkoxide base" refers to a base having a group of formula (Ci-6 alkyl)O⁻, wherein the Ci-6 alkyl refers to a linear or branched alkyl group having 1 to 6 carbons. Example alkoxide bases include, but are not limited to,
10 methoxide, ethoxide, propoxide, isopropoxide, n-butoxide, iso-butoxide, tert-butoxide, and the like.

As used herein, the term "alkali metal alkoxide" refers to a group of formula M(O-Ci-6 alkyl), wherein M refers to an alkali metal (e.g., lithium, sodium, or potassium) and Ci-6 alkyl refers to a linear or branched alkyl group having 1 to 6
15 carbons. Example alkali metal alkoxides include, but are not limited to, sodium methoxide, sodium ethoxide, sodium isopropoxide, sodium tert-butoxide, lithium methoxide, lithium ethoxide, potassium methoxide, potassium ethoxide, potassium tert-butoxide, and the like.

As used herein the term, "hydrogenation catalyst" refers to a metal (e.g.,
20 palladium, nickel, or rhodium) catalyst suitable to catalyze a hydrogenation reaction (i.e., reaction of a compound with hydrogen gas). Example hydrogenation catalysts include, but are not limited to, palladium on carbon, Lindlar's catalyst (palladium deposited on calcium carbonate or barium sulfate), Wilkinson's catalyst, $\text{HRuCl(PPh}_3)_3$, $\text{RhCl(PPh}_3)_3$, $[\text{Rh(COD)Cl}]_2$, $[\text{Ir(COD)(PMePh}_2)_2]^+$, $[\text{Rh(1,5-cyclooctadiene)(PPh}_3)_2]^+$, PtO_2 (Adam's catalyst), palladium on carbon, palladium
25 black, Lindlar's catalyst (palladium deposited on calcium carbonate or barium sulfate and treated with lead), and the like. In some embodiments, the hydrogenation catalyst is one found in Nishimura, Heterogeneous Catalytic Hydrogenation for Organic Synthesis, Edition 1, Wiley (April 17, 2001) or Chalonier, Homogeneous
30 Hydrogenation, Edition 1, Springer Netherlands (December 6, 2010), each of which is incorporated herein by reference in its entirety.

Appropriate P¹ protecting groups include, but are not limited to the protecting groups for amines delineated in Wuts and Greene, *Protective Groups in Organic Synthesis*, 4th ed., John Wiley & Sons: New Jersey, pages 696-887 (and, in particular, pages 872-887) (2007), which is incorporated herein by reference in its entirety.

5 Example amine protecting groups include, but are not limited to, benzyloxycarbonyl (Cbz), 2,2,2-trichloroethoxycarbonyl (Troc), 2-(trimethylsilyl)ethoxycarbonyl (Teoc), 2-(4-trifluoromethylphenylsulfonyl)ethoxycarbonyl (Tsc), t-butoxycarbonyl (BOC), 1-adamantyloxycarbonyl (Adoc), 2-adamantylcarbonyl (2-Adoc), 2,4-dimethylpent-3-yloxycarbonyl (Doc), cyclohexyloxycarbonyl (Hoc), 1,1-dimethyl-2,2,2-
10 trichloroethoxycarbonyl (TcBOC), vinyl, 2-chloroethyl, 2-phenylsulfonyl ethyl, allyl, benzyl, 2-nitrobenzyl, 4-nitrobenzyl, diphenyl-4-pyridylmethyl, N',N'-dimethylhydrazinyl, methoxymethyl, t-butoxymethyl (Bum), benzyloxymethyl (BOM), or 2-tetrahydropyranyl (THP), tri(Ci-4 alkyl)silyl (e.g., tri(isopropyl)silyl), 1,1-diethoxymethyl, or N-pivaloyloxymethyl (POM).

15 Example oxidizing agents include Dess-Martin periodinane, 2,2,6,6-tetramethyl-1-piperidinyloxy free radical (TEMPO), N-halosuccinimide (e.g., N-bromosuccinimide, N-chlorosuccinimide, N-iodosuccinimide), and the like.

Example reducing agents include alkali metal borohydrides (e.g., lithium borohydride, sodium borohydride, potassium borohydride, etc.), alkali metal
20 aluminum hydrides (e.g., lithium aluminum hydride, sodium aluminum hydride), hydrogen gas (e.g., H₂/Pd on carbon), and the like.

Example acids can be inorganic or organic acids and include, but are not limited to, strong and weak acids. Some example acids include hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, p-toluenesulfonic acid, 4-
25 nitrobenzoic acid, methanesulfonic acid, benzenesulfonic acid, trifluoroacetic acid, and nitric acid. Some weak acids include, but are not limited to acetic acid, propionic acid, butanoic acid, benzoic acid, tartaric acid, pentanoic acid, hexanoic acid, heptanoic acid, octanoic acid, nonanoic acid, and decanoic acid.

Some example strong bases include, but are not limited to, hydroxide,
30 alkoxides, metal amides, metal hydrides, metal dialkylamides and arylamines, wherein; alkoxides include lithium, sodium and potassium salts of methyl, ethyl and t-butyl oxides; metal amides include sodium amide, potassium amide and lithium

amide; metal hydrides include sodium hydride, potassium hydride and lithium hydride; and metal dialkylamides include lithium, sodium, and potassium salts of methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, trimethylsilyl and cyclohexyl substituted amides.

5 All compounds, and pharmaceutically acceptable salts thereof, can be found together with other substances such as water and solvents (e.g., hydrates and solvates) or can be isolated.

In some embodiments, the compounds of the invention, or salts thereof, are substantially isolated. By "substantially isolated" is meant that the compound is at
10 least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compounds of the invention. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99%
15 by weight of the compounds of the invention, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

As used herein, the term "reacting" is used as known in the art and generally refers to the bringing together of chemical reagents in such a manner so as to allow their interaction at the molecular level to achieve a chemical or physical
20 transformation. In some embodiments, the reacting involves two reagents, wherein one or more equivalents of second reagent are used with respect to the first reagent. The reacting steps of the processes described herein can be conducted for a time and under conditions suitable for preparing the identified product.

The reactions of the processes described herein can be carried out in suitable
25 solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially nonreactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, e.g., temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out
30 in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected. In some

embodiments, reactions can be carried out in the absence of solvent, such as when at least one of the reagents is a liquid or gas.

A "solvent component" may be one solvent or a mixture of two or more solvents.

5 As used herein, "second", "third", "fourth", etc. as a prefix to the phrase "solvent component" is used to differentiate the solvent component from other solvent components used in earlier or later steps of the process and does not indicate that multiple solvents must be present.

Suitable solvents can include halogenated solvents such as carbon
10 tetrachloride, bromodichloromethane, dibromochloromethane, bromoform, chloroform, bromochloromethane, dibromomethane, butyl chloride, dichloromethane, tetrachloroethylene, trichloroethylene, 1,1,1-trichloroethane, 1,1,2-trichloroethane, 1,1-dichloroethane, 2-chloropropane, α,α,α -trifluorotoluene, 1,2-dichloroethane, 1,2-dibromoethane, hexafluorobenzene, 1,2,4-trichlorobenzene, 1,2-dichlorobenzene,
15 chlorobenzene, fluorobenzene, mixtures thereof and the like.

Suitable ether solvents include: dimethoxymethane, tetrahydrofuran, 1,3-dioxane, 1,4-dioxane, furan, diethyl ether, ethylene glycol dimethyl ether, ethylene glycol diethyl ether, diethylene glycol dimethyl ether (diglyme), diethylene glycol diethyl ether, triethylene glycol dimethyl ether, anisole, t-butyl methyl ether, mixtures
20 thereof and the like.

Suitable protic solvents can include, by way of example and without limitation, water, methanol, ethanol, 2-nitroethanol, 2-fluoroethanol, 2,2,2-trifluoroethanol, ethylene glycol, 1-propanol, 2-propanol, 2-methoxy ethanol, 1-butanol, 2-butanol, i-butyl alcohol, t-butyl alcohol, 2-ethoxyethanol, diethylene
25 glycol, 1-, 2-, or 3- pentanol, neo-pentyl alcohol, t-pentyl alcohol, diethylene glycol monomethyl ether, diethylene glycol monoethyl ether, cyclohexanol, benzyl alcohol, phenol, or glycerol.

Suitable aprotic solvents can include, by way of example and without limitation, tetrahydrofuran (THF), N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMA), 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU), 1,3-dimethyl-2-imidazolidinone (DMI), N-methylpyrrolidinone (NMP),
30 formamide, N-methylacetamide, N-methylformamide, acetonitrile, dimethyl

sulfoxide, propionitrile, ethyl formate, methyl acetate, hexachloroacetone, acetone, ethyl methyl ketone, ethyl acetate, sulfolane, N,N-dimethylpropionamide, tetramethylurea, nitromethane, nitrobenzene, or hexamethylphosphoramide.

Suitable hydrocarbon solvents include benzene, cyclohexane, pentane, hexane, 5 toluene, cycloheptane, methylcyclohexane, heptane, ethylbenzene, m-, o-, or p-xylene, octane, indane, nonane, or naphthalene.

The reactions of the processes described herein can be carried out in air or under an inert atmosphere. Typically, reactions containing reagents or products that are substantially reactive with air can be carried out using air-sensitive synthetic 10 techniques that are well known to the skilled artisan.

In some embodiments, preparation of compounds can involve the addition of acids or bases to affect, for example, catalysis of a desired reaction or formation of salt forms such as acid addition salts.

Example acids can be inorganic or organic acids. Inorganic acids include 15 hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, and nitric acid. Organic acids include formic acid, acetic acid, propionic acid, butanoic acid, benzoic acid, 4-nitrobenzoic acid, methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, tartaric acid, trifluoroacetic acid, propiolic acid, butyric acid, 2-butynoic acid, vinyl acetic acid, pentanoic acid, hexanoic acid, heptanoic acid, 20 octanoic acid, nonanoic acid and decanoic acid.

Example bases include lithium hydroxide, sodium hydroxide, potassium hydroxide, lithium carbonate, sodium carbonate, potassium carbonate, and sodium bicarbonate. Some example strong bases include, but are not limited to, hydroxide, alkoxides, metal amides, metal hydrides, metal dialkylamides and arylamines, 25 wherein; alkoxides include lithium, sodium and potassium salts of methyl, ethyl and t-butyl oxides; metal amides include sodium amide, potassium amide and lithium amide; metal hydrides include sodium hydride, potassium hydride and lithium hydride; and metal dialkylamides include lithium, sodium, and potassium salts of methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, trimethylsilyl and cyclohexyl 30 substituted amides.

Upon carrying out preparation of compounds according to the processes described herein, the usual isolation and purification operations such as concentration,

filtration, extraction, solid-phase extraction, recrystallization, chromatography, and the like may be used, to isolate the desired products.

The expression, "room temperature," as used herein, is understood in the art, and refer generally to a temperature (*e.g.* a reaction temperature) that is about the
5 temperature of the room in which the reaction is carried out, for example, a temperature from about 20 °C to about 30 °C.

The reactions of the processes described herein can be carried out at appropriate temperatures which can be readily determined by the skilled artisan. Reaction temperatures will depend on, for example, the melting and boiling points of
10 the reagents and solvent, if present; the thermodynamics of the reaction (*e.g.*, vigorously exothermic reactions may need to be carried out at reduced temperatures); and the kinetics of the reaction (*e.g.*, a high activation energy barrier may need elevated temperatures).

Preparation of the compounds described herein can involve the protection and
15 deprotection of various chemical groups (*e.g.*, protection and deprotection of amine groups). The need for protection and deprotection, and the selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Wuts and Greene, Protective Groups in Organic Synthesis, 4th ed., John Wiley & Sons: New Jersey, (2007), which
20 is incorporated herein by reference in its entirety. Adjustments to the protecting groups and formation and cleavage methods described herein may be adjusted as necessary in light of the various substituents.

As used herein, the term "deprotection conditions" refers to conditions suitable to cleave an amine protecting group. In some embodiments, deprotection
25 conditions may include cleavage of a protecting group in the presence of a strong acid, in the presence of a strong base, in the presence of a reducing agent, or in the presence of an oxidizing agent. Deprotection of an amine protecting group can be accomplished by methods known in the art for the removal of particular protecting groups for amines, such as those in Wuts and Greene, Protective Groups in Organic
30 Synthesis, 4th ed., John Wiley & Sons: New Jersey, pages 696-887 (and, in particular, pages 872-887) (2007), which is incorporated herein by reference in its entirety. In some embodiments, the treating comprises reacting the protected compound under

acidic conditions (e.g., hydrochloric acid or trifluoroacetic acid). In some embodiments, the temperature is about room temperature, at a temperature from about 15 °C to about 40 °C, or at a temperature from about 15 °C to about 30 °C.

Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ^1H or ^{13}C), infrared spectroscopy, spectrophotometry (e.g., UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC) or thin layer chromatography (TLC).

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The present invention also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the present invention include the non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, alcohols (e.g., methanol, ethanol, iso-propanol, or butanol) or acetonitrile (ACN) are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and *Journal of Pharmaceutical Science*, 66, 2 (1977), each of which is incorporated herein

by reference in its entirety. In some embodiments, the compounds described herein include the N-oxide forms.

Methods

5 The compound of Formula 1a, 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo [4,5-d]thieno [3,2-b]pyridin- 1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile, is a selective inhibitor of JAK1. The compound of Formula 1a inhibits JAK1 preferentially over JAK2 (e.g., have a JAK1/JAK2 IC₅₀ ratio >10).

 JAK1 plays a central role in a number of cytokine and growth factor signaling
10 pathways that, when dysregulated, can result in or contribute to disease states. For example, IL-6 levels are elevated in rheumatoid arthritis, a disease in which it has been suggested to have detrimental effects (Fonesca, J.E. et al, Autoimmunity Reviews, 8:538-42, 2009). Because IL-6 signals, at least in part, through JAK1, antagonizing IL-6 directly or indirectly through JAK1 inhibition is expected to
15 provide clinical benefit (Guschin, D., N., et al Embo J 14:1421, 1995; Smolen, J. S., et al. Lancet 371:987, 2008). Moreover, in some cancers JAK1 is mutated resulting in constitutive undesirable tumor cell growth and survival (Mullighan CG, Proc Natl Acad Sci U S A.106:9414-8, 2009; Flex E., et al.J Exp Med. 205:751-8, 2008). In other autoimmune diseases and cancers elevated systemic levels of inflammatory
20 cytokines that activate JAK1 may also contribute to the disease and/or associated symptoms. Therefore, patients with such diseases may benefit from JAK1 inhibition. Selective inhibitors of JAK1 may be efficacious while avoiding unnecessary and potentially undesirable effects of inhibiting other JAK kinases.

 Selective inhibitors of JAK1, relative to other JAK kinases, may have multiple
25 therapeutic advantages over less selective inhibitors. With respect to selectivity against JAK2, a number of important cytokines and growth factors signal through JAK2 including, for example, erythropoietin (Epo) and thrombopoietin (Tpo) (Parganas E, et al. Cell. 93:385-95, 1998). Epo is a key growth factor for red blood cells production; hence a paucity of Epo-dependent signaling can result in reduced
30 numbers of red blood cells and anemia (Kaushansky K, NEJM 354:2034-45, 2006). Tpo, another example of a JAK2-dependent growth factor, plays a central role in controlling the proliferation and maturation of megakaryocytes - the cells from which

platelets are produced (Kaushansky K, NEJM 354:2034-45, 2006). As such, reduced Tpo signaling would decrease megakaryocyte numbers (megakaryocytopenia) and lower circulating platelet counts (thrombocytopenia). This can result in undesirable and/or uncontrollable bleeding. Reduced inhibition of other JAKs, such as JAK3 and Tyk2, may also be desirable as humans lacking functional version of these kinases have been shown to suffer from numerous maladies such as severe-combined immunodeficiency or hyperimmunoglobulin E syndrome (Minegishi, Y, et al. Immunity 25:745-55, 2006; Macchi P, et al. Nature. 377:65-8, 1995). Therefore a JAK1 inhibitor with reduced affinity for other JAKs would have significant advantages over a less-selective inhibitor with respect to reduced side effects involving immune suppression, anemia and thrombocytopenia.

Another aspect of the present invention pertains to methods of treating a JAK-associated disease or disorder in an individual (*e.g.*, patient) by administering to the individual in need of such treatment a therapeutically effective amount or dose of a compound of the present invention or a pharmaceutical composition thereof. A JAK-associated disease can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of the JAK, including overexpression and/or abnormal activity levels. A JAK-associated disease can also include any disease, disorder or condition that can be prevented, ameliorated, or cured by modulating JAK activity.

JAK inhibitors are useful in treating various JAK-associated diseases or disorders. Examples of JAK-associated diseases include diseases involving the immune system including, for example, organ transplant rejection (*e.g.*, allograft rejection and graft versus host disease). Further examples of JAK-associated diseases include autoimmune diseases such as multiple sclerosis, rheumatoid arthritis, juvenile arthritis, psoriatic arthritis, type I diabetes, lupus, psoriasis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, myasthenia gravis, immunoglobulin nephropathies, myocarditis, autoimmune thyroid disorders, chronic obstructive pulmonary disease (COPD), and the like. In some embodiments, the autoimmune disease is an autoimmune bullous skin disorder such as pemphigus vulgaris (PV) or bullous pemphigoid (BP).

Further examples of JAK-associated diseases include allergic conditions such as asthma, food allergies, eszematous dermatitis, contact dermatitis, atopic dermatitis (atropic eczema), and rhinitis. Further examples of JAK-associated diseases include viral diseases such as Epstein Barr Virus (EBV), Hepatitis B, Hepatitis C, HIV,
5 HTLV 1, Varicella-Zoster Virus (VZV) and Human Papilloma Virus (HPV).

Further examples of JAK-associated diseases include diseases associated with cartilage turnover, for example, gouty arthritis, septic or infectious arthritis, reactive arthritis, reflex sympathetic dystrophy, algodystrophy, Tietze syndrome, costal athropathy, osteoarthritis deformans endemica, Mseleni disease, Handigodu disease,
10 degeneration resulting from fibromyalgia, systemic lupus erythematosus, scleroderma, or ankylosing spondylitis.

Further examples of JAK-associated disease include congenital cartilage malformations, including hereditary chondrolysis, chondrodysplasias, and pseudo-chondrodysplasias (e.g., microtia, enotia, and metaphyseal
15 chondrodysplasia).

Further examples of JAK-associated diseases or conditions include skin disorders such as psoriasis (for example, psoriasis vulgaris), atopic dermatitis, skin rash, skin irritation, skin sensitization (*e.g.*, contact dermatitis or allergic contact dermatitis). For example, certain substances including some pharmaceuticals when topically
20 applied can cause skin sensitization. In some embodiments, co-administration or sequential administration of at least one JAK inhibitor of the invention together with the agent causing unwanted sensitization can be helpful in treating such unwanted sensitization or dermatitis. In some embodiments, the skin disorder is treated by topical administration of at least one JAK inhibitor of the invention.

Further examples of JAK-associated diseases or conditions include those characterized by solid tumors (*e.g.*, prostate cancer, renal cancer, hepatic cancer, pancreatic cancer, gastric cancer, breast cancer, lung cancer, cancers of the head and neck, thyroid cancer, glioblastoma, Kaposi's sarcoma, Castleman's disease, uterine leiomyosarcoma, melanoma etc.), hematological cancers (*e.g.*, lymphoma, leukemia
30 such as acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML) or multiple myeloma), and skin cancer such as cutaneous T-cell lymphoma (CTCL) and cutaneous B-cell lymphoma. Example CTCLs include Sezary syndrome and mycosis

fungoides. Other examples of JAK-associated diseases or conditions include pulmonary arterial hypertension.

Other examples of JAK-associated diseases or conditions include inflammation-associated cancers. In some embodiments, the cancer is associated with inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is ulcerative colitis. In some embodiments, the inflammatory bowel disease is Crohn's disease. In some embodiments, the inflammation-associated cancer is colitis-associated cancer. In some embodiments, the inflammation-associated cancer is colon cancer or colorectal cancer. In some embodiments, the cancer is gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumor (GIST), adenocarcinoma, small intestine cancer, or rectal cancer.

JAK-associated diseases can further include those characterized by expression of: JAK2 mutants such as those having at least one mutation in the pseudo-kinase domain (*e.g.*, JAK2V617F); JAK2 mutants having at least one mutation outside of the pseudo-kinase domain; JAK1 mutants; JAK3 mutants; erythropoietin receptor (EPOR) mutants; or deregulated expression of CRLF2.

JAK-associated diseases can further include myeloproliferative disorders (MPDs) such as polycythemia vera (PV), essential thrombocythemia (ET), myelofibrosis with myeloid metaplasia (MMM), primary myelofibrosis (PMF), chronic myelogenous leukemia (CML), chronic myelomonocytic leukemia (CMML), hypereosinophilic syndrome (HES), systemic mast cell disease (SMCD), and the like. In some embodiments, the myeloproliferative disorder is myelofibrosis (*e.g.*, primary myelofibrosis (PMF) or post polycythemia vera/essential thrombocythemia myelofibrosis (Post-PV/Post-ET MF)). In some embodiments, the myeloproliferative disorder is post-essential thrombocythemia myelofibrosis (Post-ET MF). In some embodiments, the myeloproliferative disorder is post polycythemia vera myelofibrosis (Post-PV MF).

JAK-associated diseases further include myelodysplastic syndrome (MDS).

As used herein, myelodysplastic syndromes are intended to encompass heterogeneous and clonal hematopoietic disorders that are characterized by ineffective hematopoiesis on one or more of the major myeloid cell lineages. Myelodysplastic syndromes are associated with bone marrow failure, peripheral blood cytopenias, and

a propensity to progress to acute myeloid leukemia (AML). Moreover, clonal cytogenetic abnormalities can be detected in about 50% of cases with MDS. In 1997, The World Health Organization (WHO) in conjunction with the Society for Hematopathology (SH) and the European Association of Hematopathology (EAHP) proposed new classifications for hematopoietic neoplasms (Harris, et al, *J Clin Oncol* 1999;17:3835-3849; Vardiman, et al., *Blood* 2002;100:2292-2302). For MDS, the WHO utilized not only the morphologic criteria from the French-American-British (FAB) classification but also incorporated available genetic, biologic, and clinical characteristics to define subsets of MDS (Bennett, et al, *Br J Haematol* 1982;51:189-199). In 2008, the WHO classification of MDS (Table 1) was further refined to allow precise and prognostically relevant subclassification of unilineage dysplasia by incorporating new clinical and scientific information (Vardiman, et al, *Blood* 2009;114:937-951; Swerdlow, et al, WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, 4th Edition. Lyon France: IARC Press; 2008:88-103; Bunning and Germing, "Myelodysplasia; syndromes/neoplasms" in Chapter 5, Swerdlow, et al, eds. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, (ed. 4th edition): Lyon, France: IARC Press;2008:88-103).

Table 1. 2008 WHO Classification for De Novo Myelodysplastic Syndrome

Subtype	Blood	Bone Marrow
Refractory cytopenia with unilineage dysplasia (RCUD)	Single or Bicytopenia	Dysplasia in $\geq 10\%$ of 1 cell line, $< 5\%$ blasts
Refractory anemia with ring sideroblasts (RARS)	Anemia, no blasts	$\geq 15\%$ of erythroid precursors w/ring sideroblasts, erythroid dysplasia only, $< 5\%$ blasts
Refractory cytopenia with multilineage dysplasia	Cytopenia(s), $< 1 \times 10^9/L$ monocytes	Dysplasia in $\geq 10\%$ of cells in ≥ 2 hematopoietic lineages, $\pm 15\%$ ring sideroblasts, $< 5\%$ blasts
Refractory anemia with excess blasts-1 (RAEB-1)	Cytopenia(s), $\leq 2\%$ to 4% blasts, $< 1 \times 10^9/L$ monocytes	Unilineage or multilineage dysplasia, No Auer rods, 5% to 9% blasts
Refractory anemia with excess blasts-2 (RAEB-2)	Cytopenia(s), $\leq 5\%$ to 19% blasts, $< 1 \times 10^9/L$ monocytes	Unilineage or multilineage dysplasia, \pm Auer rods, 10% to 19% blasts
Myelodysplastic syndrome, unclassified (MDS-U)	Cytopenias	Unilineage or no dysplasia but characteristic MDS cytogenetics, $< 5\%$ blasts
MDS associated with isolated del(5q)	Anemia, platelets normal or increased	Unilineage erythroid. Isolated del(5q), $< 5\%$ blasts

5

In some embodiments, the present application provides a method of treating a myelodysplastic syndrome (MDS) in patient in need thereof, comprising administering to said patient a compound of Formula Ia, or a pharmaceutically acceptable salt thereof, wherein said MDS is selected from refractory cytopenia with unilineage dysplasia (RCUD), refractory anemia with ring sideroblasts (RARS), refractory cytopenia with multilineage dysplasia, refractory anemia with excess blasts-1 (RAEB-1), refractory anemia with excess blasts-2 (RAEB-2), myelodysplastic syndrome, unclassified (MDS-U), and MDS associated with isolated del(5q).

15

In some embodiments, the myelodysplastic syndrome is refractory cytopenia with unilineage dysplasia (RCUD).

In some embodiments, the myelodysplasia syndrome is refractory anemia with ring sideroblasts (RARS).

In some embodiments, the myelodysplasia syndrome is refractory cytopenia with multilineage dysplasia.

5 In some embodiments, the myelodysplastic syndrome is refractory anemia with excess blasts- 1 (RAEB-1).

In some embodiments, the myelodysplastic syndrome is refractory anemia with excess blasts-2 (RAEB-2).

10 In some embodiments, the myelodysplastic syndrome is myelodysplastic syndrome, unclassified (MDS-U).

In some embodiments, the myelodysplastic syndrome is myelodysplastic syndrome associated with isolated del(5q).

In some embodiments, the myelodysplastic syndrome is refractory to erythropoiesis-stimulating agents.

15 Other examples of JAK-associated diseases or conditions include ameliorating the dermatological side effects of other pharmaceuticals by administration of the compound of the invention. For example, numerous pharmaceutical agents result in unwanted allergic reactions which can manifest as acneiform rash or related dermatitis. Example pharmaceutical agents that have such undesirable side effects include anti-cancer drugs such as gefitinib, cetuximab, erlotinib, and the like. The
20 compounds of the invention can be administered systemically or topically (e.g., localized to the vicinity of the dermatitis) in combination with (e.g., simultaneously or sequentially) the pharmaceutical agent having the undesirable dermatological side effect. In some embodiments, the compound of the invention can be administered
25 topically together with one or more other pharmaceuticals, where the other pharmaceuticals when topically applied in the absence of a compound of the invention cause contact dermatitis, allergic contact sensitization, or similar skin disorder. Accordingly, compositions of the invention include topical formulations containing the compound of the invention and a further pharmaceutical agent which can cause
30 dermatitis, skin disorders, or related side effects.

Further JAK-associated diseases include inflammation and inflammatory diseases. Example inflammatory diseases include sarcoidosis, inflammatory diseases

of the eye (*e.g.*, iritis, uveitis, scleritis, conjunctivitis, or related disease), inflammatory diseases of the respiratory tract (*e.g.*, the upper respiratory tract including the nose and sinuses such as rhinitis or sinusitis or the lower respiratory tract including bronchitis, chronic obstructive pulmonary disease, and the like), inflammatory myopathy such as myocarditis, and other inflammatory diseases. In some embodiments, the inflammation disease of the eye is blepharitis.

Further JAK-associated diseases include ischemia reperfusion injuries or a disease or condition related to an inflammatory ischemic event such as stroke or cardiac arrest, endotoxin-driven disease state (*e.g.*, complications after bypass surgery or chronic endotoxin states contributing to chronic cardiac failure), anorexia, cachexia, fatigue such as that resulting from or associated with cancer, restenosis, sclerodermitis, fibrosis, conditions associated with hypoxia or astrogliosis such as, for example, diabetic retinopathy, cancer, or neurodegeneration, and other inflammatory diseases such as systemic inflammatory response syndrome (SIRS) and septic shock.

Other JAK-associated diseases include gout and increased prostate size due to, *e.g.*, benign prostatic hypertrophy or benign prostatic hyperplasia, as well as bone resorption diseases such as osteoporosis or osteoarthritis, bone resorption diseases associated with: hormonal imbalance and/or hormonal therapy, autoimmune disease (*e.g.* osseous sarcoidosis), or cancer (*e.g.* myeloma).

Further JAK-associated diseases include a dry eye disorder. As used herein, "dry eye disorder" is intended to encompass the disease states summarized in a recent official report of the Dry Eye Workshop (DEWS), which defined dry eye as "a multifactorial disease of the tears and ocular surface that results in symptoms of discomfort, visual disturbance, and tear film instability with potential damage to the ocular surface. It is accompanied by increased osmolarity of the tear film and inflammation of the ocular surface." Lemp, "The Definition and Classification of Dry Eye Disease: Report of the Definition and Classification Subcommittee of the International Dry Eye Workshop", *The Ocular Surface*, 5(2), 75-92 April 2007, which is incorporated herein by reference in its entirety. In some embodiments, the dry eye disorder is selected from aqueous tear-deficient dry eye (ADDE) or evaporative dry eye disorder, or appropriate combinations thereof. In some embodiments, the dry eye

disorder is Sjogren syndrome dry eye (SSDE). In some embodiments, the dry eye disorder is non-Sjogren syndrome dry eye (NSSDE).

Further JAK-associated diseases include conjunctivitis, uveitis (including chronic uveitis), chorioiditis, retinitis, cyclitis, scleritis, episcleritis, or iritis. Other
5 JAK-associated diseases include respiratory dysfunction or failure associated with viral infection, such as influenza and SARS.

Combination Therapies

One or more additional pharmaceutical agents such as, for example,
10 chemotherapeutics, anti-inflammatory agents, steroids, immunosuppressants, as well as PBK δ , mTor, Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors such as, for example, those described in WO 2006/056399, which is incorporated herein by reference in its entirety, or other agents can be used in combination with the compounds described herein for treatment of JAK-associated diseases, disorders or
15 conditions. The one or more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

Example chemotherapeutics include proteasome inhibitors (*e.g.*, bortezomib), thalidomide, revlimid, and DNA-damaging agents such as melphalan, doxorubicin, cyclophosphamide, vincristine, etoposide, carmustine, and the like.

20 Example steroids include corticosteroids such as dexamethasone or prednisone.

Example Bcr-Abl inhibitors include the compounds, and pharmaceutically acceptable salts thereof, of the genera and species disclosed in U.S. Pat. No. 5,521,184, WO 04/005281, and U.S. Ser. No. 60/578,491, all of which are
25 incorporated herein by reference in their entirety.

Example suitable Flt-3 inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 03/037347, WO 03/099771, and WO 04/046120, all of which are incorporated herein by reference in their entirety.

Example suitable RAF inhibitors include compounds, and their
30 pharmaceutically acceptable salts, as disclosed in WO 00/09495 and WO 05/028444, both of which are incorporated herein by reference in their entirety.

Example suitable FAK inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 04/080980, WO 04/056786, WO 03/024967, WO 01/064655, WO 00/053595, and WO 01/014402, all of which are incorporated herein by reference in their entirety.

5 In some embodiments, the additional pharmaceutical agent is selected from IMiDs, an anti-IL-6 agent, an anti-TNF- α agent, a hypomethylating agent, and a biologic response modifier (BRM).

Generally, a BRM is a substance made from living organisms to treat disease, which may occur naturally in the body or may be made in the laboratory. Examples
10 of BRMs include IL-2, interferon, various types of colony-stimulating factors (CSF, GM-CSF, G-CSF), monoclonal antibodies such as abciximab, etanercept, infliximab, rituximab, trastuzumab, and high dose ascorbate.

In some embodiments, the anti-TNF- α agent is infliximab, and etanercept.

In some embodiments, the hypomethylating agent is a DNA methyltransferase
15 inhibitor. In some embodiments, the DNA methyltransferase inhibitor is selected from 5 azacytidine and decitabine.

Generally, IMiDs are as immunomodulatory agents. In some embodiments, the IMiD is selected from thalidomide, lenalidomide, pomalidomide, CC-1 1006, and CC-10015.

20 In some embodiments, the additional pharmaceutical agent is selected from anti-thymocyte globulin, recombinant human granulocyte colony-stimulating factor (G-CSF), granulocyte-monocyte CSF (GM-CSF), a erythropoiesis-stimulating agent (ESA), and cyclosporine.

In some embodiments, the additional pharmaceutical agent is an additional
25 JAK inhibitor. In some embodiments, the additional JAK inhibitor is tofacitinib or ruxolitinib.

In some embodiments, one or more of the compounds of the invention can be used in combination with one or more other kinase inhibitors including imatinib, particularly for treating patients resistant to imatinib or other kinase inhibitors.

30 In some embodiments, a suitable chemotherapeutical agent can be selected from antimetabolite agents, topoisomerase 1 inhibitors, platinum analogs, taxanes, anthracyclines, and EGFR inhibitors, and combinations thereof.

In some embodiments, antimetabolite agents include capecitabine, gemcitabine, and fluorouracil (5-FU).

In some embodiments, taxanes include paclitaxel, Abraxane® (paclitaxel protein-bound particles for injectable suspension), and Taxotere® (docetaxel).

5 In some embodiments, platinum analogs include oxaliplatin, cisplatin, and carboplatin.

In some embodiments, topoisomerase 1 inhibitors include irinotecan and topotecan.

10 In some embodiments, anthracyclines include doxorubicin or liposomal formulations of doxorubicin.

In some embodiments, the chemotherapeutic is FOLFIRINOX (5-FU, lecovorin, irinotecan and oxaliplatin). In some embodiments, the chemotherapeutic agent is gemcitabine and Abraxane® (paclitaxel protein-bound particles for injectable suspension).

15 In some embodiments, one or more JAK inhibitors of the invention can be used in combination with a chemotherapeutic in the treatment of cancer, such as multiple myeloma, and may improve the treatment response as compared to the response to the chemotherapeutic agent alone, without exacerbation of its toxic effects. Examples of additional pharmaceutical agents used in the treatment of
20 multiple myeloma, for example, can include, without limitation, melphalan, melphalan plus prednisone [MP], doxorubicin, dexamethasone, and Velcade (bortezomib). Further additional agents used in the treatment of multiple myeloma include Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors. Additive or synergistic effects are desirable outcomes of combining a JAK inhibitor of the present invention
25 with an additional agent. Furthermore, resistance of multiple myeloma cells to agents such as dexamethasone may be reversible upon treatment with a JAK inhibitor of the present invention. The agents can be combined with the present compounds in a single or continuous dosage form, or the agents can be administered simultaneously or sequentially as separate dosage forms.

30 In some embodiments, a corticosteroid such as dexamethasone is administered to a patient in combination with at least one JAK inhibitor where the dexamethasone is administered intermittently as opposed to continuously.

In some further embodiments, combinations of one or more JAK inhibitors of the invention with other therapeutic agents can be administered to a patient prior to, during, and/or after a bone marrow transplant or stem cell transplant.

In some embodiments, the additional therapeutic agent is fluocinolone acetonide (Retisert®), or rimexolone (AL-2178, Vexol, Alcon).

In some embodiments, the additional therapeutic agent is cyclosporine (Restasis®).

In some embodiments, the additional therapeutic agent is a corticosteroid. In some embodiments, the corticosteroid is triamcinolone, dexamethasone, fluocinolone, cortisone, prednisolone, or flumetholone.

In some embodiments, the additional therapeutic agent is selected from Dehydrex™ (Holies Labs), Civamide (Opko), sodium hyaluronate (Vismed, Lantibio/TRB Chemedica), cyclosporine (ST-603, Sirion Therapeutics), ARG101(T) (testosterone, Argentis), AGR1012(P) (Argentis), ecabet sodium (Senju-Ista), gefarnate (Santen), 15-(s)-hydroxyeicosatetraenoic acid (15(S)-HETE), cevilemine, doxycycline (ALTY-0501, Alacrity), minocycline, iDestrin™ (NP50301, Nascent Pharmaceuticals), cyclosporine A (Nova22007, Novagali), oxytetracycline (Duramycin, MOLI1901, Lantibio), CF101 (2S,3S,4R,5R)-3,4-dihydroxy-5-[6-[(3-iodophenyl)methylamino]purin-9-yl]-N-methyl-oxolane-2-carbamyl, Can-Fite Biopharma), voclosporin (LX212 or LX214, Lux Biosciences), ARG103 (Agentis), RX-10045 (synthetic resolvins analog, Resolvix), DYN15 (Dyanmis Therapeutics), rivoglitazone (DE011, Daiichi Sankyo), TB4 (RegeneRx), OPH-01 (Ophtalmis Monaco), PCS101 (Pericor Science), REV1-31 (Evolutec), Lacritin (Senju), rebamipide (Otsuka-Novartis), OT-551 (Othera), PAI-2 (University of Pennsylvania and Temple University), pilocarpine, tacrolimus, pimecrolimus (AMS981, Novartis), loteprednol etabonate, rituximab, diquafosol tetrasodium (INS365, Inspire), KLS-0611 (Kissei Pharmaceuticals), dehydroepiandrosterone, anakinra, efalizumab, mycophenolate sodium, etanercept (Embrel®), hydroxychloroquine, NGX267 (TorreyPines Therapeutics), actemra, gemcitabine, oxaliplatin, L-asparaginase, or thalidomide.

In some embodiments, the additional therapeutic agent is an anti-angiogenic agent, cholinergic agonist, TRP-1 receptor modulator, a calcium channel blocker, a

mucin secretagogue, MUC1 stimulant, a calcineurin inhibitor, a corticosteroid, a P2Y2 receptor agonist, a muscarinic receptor agonist, an mTOR inhibitor, another JAK inhibitor, Bcr-Abl kinase inhibitor, Flt-3 kinase inhibitor, RAF kinase inhibitor, and FAK kinase inhibitor such as, for example, those described in WO 2006/056399, which is incorporated herein by reference in its entirety. In some embodiments, the additional therapeutic agent is a tetracycline derivative (e.g., minocycline or doxycycline). In some embodiments, the additional therapeutic agent binds to FKBP12.

In some embodiments, the additional therapeutic agent is an alkylating agent or DNA cross-linking agent; an anti-metabolite/demethylating agent (e.g., 5-fluorouracil, capecitabine or azacitidine); an anti-hormone therapy (e.g., hormone receptor antagonists, SERMs, or aromatase inhibitor); a mitotic inhibitor (e.g., vincristine or paclitaxel); an topoisomerase (I or II) inhibitor (e.g. mitoxantrone and irinotecan); an apoptotic inducers (e.g. ABT-737); a nucleic acid therapy (e.g. antisense or RNAi); nuclear receptor ligands (e.g., agonists and/or antagonists: all-trans retinoic acid or bexarotene); epigenetic targeting agents such as histone deacetylase inhibitors (e.g. vorinostat), hypomethylating agents (e.g. decitabine); regulators of protein stability such as Hsp90 inhibitors, ubiquitin and/or ubiquitin like conjugating or deconjugating molecules; or an EGFR inhibitor (erlotinib).

In some embodiments, the additional therapeutic agent(s) are demulcent eye drops (also known as "artificial tears"), which include, but are not limited to, compositions containing polyvinylalcohol, hydroxypropyl methylcellulose, glycerin, polyethylene glycol (e.g. PEG400), or carboxymethyl cellulose. Artificial tears can help in the treatment of dry eye by compensating for reduced moistening and lubricating capacity of the tear film. In some embodiments, the additional therapeutic agent is a mucolytic drug, such as N-acetyl-cysteine, which can interact with the mucoproteins and, therefore, to decrease the viscosity of the tear film.

In some embodiments, the additional therapeutic agent includes an antibiotic, antiviral, antifungal, anesthetic, anti-inflammatory agents including steroidal and non-steroidal anti-inflammatories, and anti-allergic agents. Examples of suitable medicaments include aminoglycosides such as amikacin, gentamycin, tobramycin, streptomycin, netilmycin, and kanamycin; fluoroquinolones such as ciprofloxacin, norfloxacin, ofloxacin, trovafloxacin, lomefloxacin, levofloxacin, and enoxacin;

naphthylidene; sulfonamides; polymyxin; chloramphenicol; neomycin; paramomycin; colistimethate; bacitracin; vancomycin; tetracyclines; rifampin and its derivatives ("rifampins"); cycloserine; beta-lactams; cephalosporins; amphotericins; fluconazole; flucytosine; natamycin; miconazole; ketoconazole; corticosteroids; diclofenac; 5 flurbiprofen; ketorolac; suprofen; cromolyn; lodoxamide; levocabastin; naphazoline; antazoline; pheniramine; or azalide antibiotic.

Pharmaceutical Formulations and Dosage Forms

When employed as pharmaceuticals, the compounds of the invention can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, epidermal, ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (*e.g.*, by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral or parenteral. 15 Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, *e.g.*, intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus dose, or may be, for example, by a continuous perfusion pump. 20 Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

This invention also includes pharmaceutical compositions which contain, as the active ingredient, the compound of the invention or a pharmaceutically acceptable salt thereof, in combination with one or more pharmaceutically acceptable carriers (excipients). In some embodiments, the composition is suitable for topical administration. In making the compositions of the invention, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When 30 the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which

acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft
5 and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it can be milled to a particle size of less than 200
10 mesh. If the active compound is substantially water soluble, the particle size can be adjusted by milling to provide a substantially uniform distribution in the formulation, *e.g.* about 40 mesh.

The compounds of the invention may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet
15 formation and for other formulation types. Finely divided (nanoparticulate) preparations of the compounds of the invention can be prepared by processes known in the art, *e.g.*, see International App. No. WO 2002/000196.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth,
20 gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions
25 of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

In some embodiments, the pharmaceutical composition comprises silicified microcrystalline cellulose (SMCC) and at least one compound described herein, or a
30 pharmaceutically acceptable salt thereof. In some embodiments, the silicified microcrystalline cellulose comprises about 98% microcrystalline cellulose and about 2% silicon dioxide w/w.

In some embodiments, the composition is a sustained release composition comprising at least one compound described herein, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier. In some
5 embodiments, the composition comprises at least one compound described herein, or a pharmaceutically acceptable salt thereof, and at least one component selected from microcrystalline cellulose, lactose monohydrate, hydroxypropyl methylcellulose, and polyethylene oxide. In some embodiments, the composition comprises at least one
10 compound described herein, or a pharmaceutically acceptable salt thereof, and microcrystalline cellulose, lactose monohydrate, and hydroxypropyl methylcellulose. In some embodiments, the composition comprises at least one compound described herein, or a pharmaceutically acceptable salt thereof, and microcrystalline cellulose, lactose monohydrate, and polyethylene oxide. In some
15 embodiments, the composition further comprises magnesium stearate or silicon dioxide. In some embodiments, the microcrystalline cellulose is Avicel PHI 02™. In some embodiments, the lactose monohydrate is Fast-flo 3 16™. In some embodiments, the hydroxypropyl methylcellulose is hydroxypropyl methylcellulose 2208 K4M (e.g., Methocel K4 M Premier™) and/or hydroxypropyl methylcellulose 2208 K100LV (e.g., Methocel K00LV™). In some embodiments, the polyethylene oxide is polyethylene oxide WSR 1105 (e.g., Polyox WSR 1105™).

20 In some embodiments, a wet granulation process is used to produce the composition. In some embodiments, a dry granulation process is used to produce the composition.

The compositions can be formulated in a unit dosage form, each dosage containing from about 5 to about 1,000 mg (1 g), more usually about 100 mg to about
25 500 mg, of the active ingredient. In some embodiments, each dosage contains about 10 mg of the active ingredient. In some embodiments, each dosage contains about 50 mg of the active ingredient. In some embodiments, each dosage contains about 25 mg of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit
30 containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

In some embodiments, the compositions of the invention contain from about 5 mg to about 50 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 5 mg to about 10 mg, about 10 mg to about 15 mg, about 15 mg to about 20 mg, about 20 mg to about 25 mg, about 25 mg to about 30 mg, about 30 mg to about 35 mg, about 35 mg to about 40 mg, about 40 mg to about 45 mg, or about 45 mg to about 50 mg of the active ingredient.

In some embodiments, the compositions of the invention contain from about 50 mg to about 500 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 50 mg to about 100 mg, about 100 mg to about 150 mg, about 150 mg to about 200 mg, about 200 mg to about 250 mg, about 250 mg to about 300 mg, about 350 mg to about 400 mg, or about 450 mg to about 500 mg of the active ingredient.

In some embodiments, the compositions of the invention contain from about 500 mg to about 1,000 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 500 mg to about 550 mg, about 550 mg to about 600 mg, about 600 mg to about 650 mg, about 650 mg to about 700 mg, about 700 mg to about 750 mg, about 750 mg to about 800 mg, about 800 mg to about 850 mg, about 850 mg to about 900 mg, about 900 mg to about 950 mg, or about 950 mg to about 1,000 mg of the active ingredient.

The active compound may be effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the

composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, about 0.1 to about 1000 mg of the active ingredient of the present invention.

5 The tablets or pills of the present invention can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration
10 in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

 The liquid forms in which the compounds and compositions of the present
15 invention can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

 Compositions for inhalation or insufflation include solutions and suspensions
20 in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in can be nebulized by use of inert gases. Nebulized solutions may be
25 breathed directly from the nebulizing device or the nebulizing device can be attached to a face masks tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

 Topical formulations can contain one or more conventional carriers. In some
30 embodiments, ointments can contain water and one or more hydrophobic carriers selected from, for example, liquid paraffin, polyoxyethylene alkyl ether, propylene glycol, white Vaseline, and the like. Carrier compositions of creams can be based on

water in combination with glycerol and one or more other components, e.g. glycerinemonostearate, PEG-glycerinemonostearate and cetylstearyl alcohol. Gels can be formulated using isopropyl alcohol and water, suitably in combination with other components such as, for example, glycerol, hydroxyethyl cellulose, and the like. In
5 some embodiments, topical formulations contain at least about 0.1, at least about 0.25, at least about 0.5, at least about 1, at least about 2, or at least about 5 wt % of the compound of the invention. The topical formulations can be suitably packaged in tubes of, for example, 100 g which are optionally associated with instructions for the treatment of the select indication, e.g., psoriasis or other skin condition.

10 The amount of compound or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the
15 symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

The compositions administered to a patient can be in the form of
20 pharmaceutical compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to
25 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of a compound of the present invention can vary according to, for example, the particular use for which the treatment is made, the
30 manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a compound of the invention in a pharmaceutical composition can vary depending upon

a number of factors including dosage, chemical characteristics (*e.g.*, hydrophobicity), and the route of administration. For example, the compounds of the invention can be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration. Some typical dose ranges are from about 1 µg/kg to about 1 g/kg of body weight per day. In some embodiments, the dose range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day. The dosage is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, formulation of the excipient, and its route of administration. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

The compositions of the invention can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed hereinabove.

In some embodiments, the compound, or pharmaceutically acceptable salt thereof, is administered as an ophthalmic composition. Accordingly, in some embodiments, the methods comprise administration of the compound, or pharmaceutically acceptable salt thereof, and an ophthalmically acceptable carrier. In some embodiments, the ophthalmic composition is a liquid composition, semi-solid composition, insert, film, microparticles or nanoparticles.

In some embodiments, the ophthalmic composition is a liquid composition. In some embodiments, the ophthalmic composition is a semi-solid composition. In some embodiments, the ophthalmic composition is a topical composition. The topical compositions include, but are not limited to liquid and semi-solid compositions. In some embodiments, the ophthalmic composition is a topical composition. In some embodiments, the topical composition comprises aqueous solution, an aqueous suspension, an ointment or a gel. In some embodiments, the ophthalmic composition is topically applied to the front of the eye, under the upper eyelid, on the lower eyelid and in the cul-de-sac. In some embodiments, the ophthalmic composition is sterilized. The sterilization can be accomplished by known techniques like sterilizing filtration of the solution or by heating of the solution in the ampoule ready for use. The ophthalmic compositions of the invention can further contain pharmaceutical

excipients suitable for the preparation of ophthalmic formulations. Examples of such excipients are preserving agents, buffering agents, chelating agents, antioxidant agents and salts for regulating the osmotic pressure.

As used herein, the term "ophthalmically acceptable carrier" refers to any material that can contain and release the compound, or pharmaceutically acceptable salt thereof, and that is compatible with the eye. In some embodiments, the ophthalmically acceptable carrier is water or an aqueous solution or suspension, but also includes oils such as those used to make ointments and polymer matrices such as used in ocular inserts. In some embodiments, the composition may be an aqueous suspension comprising the compound, or pharmaceutically acceptable salt thereof. Liquid ophthalmic compositions, including both ointments and suspensions, may have a viscosity that is suited for the selected route of administration. In some embodiments, the ophthalmic composition has a viscosity in the range of from about 1,000 to about 30,000 centipoise.

In some embodiments, the ophthalmic compositions may further comprise one or more of surfactants, adjuvants, buffers, antioxidants, tonicity adjusters, preservatives (e.g., EDTA, BAK (benzalkonium chloride), sodium chlorite, sodium perborate, polyquaterium-1), thickeners or viscosity modifiers (e.g., carboxymethyl cellulose, hydroxymethyl cellulose, polyvinyl alcohol, polyethylene glycol, glycol 400, propylene glycol hydroxymethyl cellulose, hydroxypropyl-guar, hyaluronic acid, and hydroxypropyl cellulose) and the like. Additives in the formulation may include, but are not limited to, sodium chloride, sodium bicarbonate, sorbic acid, methyl paraben, propyl paraben, chlorhexidine, castor oil, and sodium perborate.

Aqueous ophthalmic compositions (solutions or suspensions) generally do not contain physiologically or ophthalmically harmful constituents. In some embodiments, purified or deionized water is used in the composition. The pH may be adjusted by adding any physiologically and ophthalmically acceptable pH adjusting acids, bases or buffers to within the range of about 5.0 to 8.5. Ophthalmically acceptable examples of acids include acetic, boric, citric, lactic, phosphoric, hydrochloric, and the like, and examples of bases include sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate, tromethamine, trishydroxymethylamino-methane, and the like. Salts and buffers

include citrate/dextrose, sodium bicarbonate, ammonium chloride and mixtures of the aforementioned acids and bases.

In some embodiments, the methods involve forming or supplying a depot of the therapeutic agent in contact with the external surface of the eye. A depot refers to a source of therapeutic agent that is not rapidly removed by tears or other eye clearance mechanisms. This allows for continued, sustained high concentrations of therapeutic agent to be present in the fluid on the external surface of the eye by a single application. Without wishing to be bound by any theory, it is believed that absorption and penetration may be dependent on both the dissolved drug concentration and the contact duration of the external tissue with the drug containing fluid. As the drug is removed by clearance of the ocular fluid and/or absorption into the eye tissue, more drug is provided, e.g. dissolved, into the replenished ocular fluid from the depot. Accordingly, the use of a depot may more easily facilitate loading of the ocular tissue for more insoluble therapeutic agents. In some embodiments, the depot can remain for up to eight hours or more. In some embodiments, the ophthalmic depot forms includes, but is not limited to, aqueous polymeric suspensions, ointments, and solid inserts.

In some embodiments, the ophthalmic composition is an ointment or gel. In some embodiment, the ophthalmic composition is an oil-based delivery vehicle. In some embodiments, the composition comprises a petroleum or lanolin base to which is added the active ingredient, usually as 0.1 to 2%, and excipients. Common bases may include, but are not limited to, mineral oil, petrolatum and combinations thereof. In some embodiments, the ointment is applied as a ribbon onto the lower eyelid.

In some embodiment, the ophthalmic composition is an ophthalmic insert. In some embodiments, the ophthalmic insert is biologically inert, soft, bio-erodible, viscoelastic, stable to sterilization after exposure to therapeutic agents, resistant to infections from air borne bacteria, bio- erodible, biocompatible, and/or viscoelastic. In some embodiments, the insert comprises an ophthalmically acceptable matrix, e.g., a polymer matrix. The matrix is typically a polymer and the therapeutic agent is generally dispersed therein or bonded to the polymer matrix. In some embodiments, the therapeutic agent may be slowly released from the matrix through dissolution or hydrolysis of the covalent bond. In some embodiments, the polymer is bioerodible

(soluble) and the dissolution rate thereof can control the release rate of the therapeutic agent dispersed therein. In another form, the polymer matrix is a biodegradable polymer that breaks down such as by hydrolysis to thereby release the therapeutic agent bonded thereto or dispersed therein. In further embodiments, the matrix and therapeutic agent can be surrounded with an additional polymeric coating to further control release. In some embodiments, the insert comprises a biodegradable polymer such as polycaprolactone (PCL), an ethylene/vinyl acetate copolymer (EVA), polyalkyl cyanoacrylate, polyurethane, a nylon, or poly (dl-lactide-co-glycolide) (PLGA), or a copolymer of any of these. In some embodiments, the therapeutic agent is dispersed into the matrix material or dispersed amongst the monomer composition used to make the matrix material prior to polymerization. In some embodiments, the amount of therapeutic agent is from about 0.1 to about 50%, or from about 2 to about 20%. In further embodiments, the biodegradable or bioerodible polymer matrix is used so that the spent insert does not have to be removed. As the biodegradable or bioerodible polymer is degraded or dissolved, the therapeutic agent is released.

In further embodiments, the ophthalmic insert comprises a polymer, including, but are not limited to, those described in Wagh, et al, "Polymers used in ocular dosage form and drug delivery systems", *Asian J. Pharm.*, pages 12-17 (Jan. 2008), which is incorporated herein by reference in its entirety. In some embodiments, the insert comprises a polymer selected from polyvinylpyrrolidone (PVP), an acrylate or methacrylate polymer or copolymer (e.g., Eudragit® family of polymers from Rohm or Degussa), hydroxymethyl cellulose, polyacrylic acid, poly(amidoamine) dendrimers, poly(dimethyl siloxane), polyethylene oxide, poly(lactide-co-glycolide), poly(2-hydroxyethylmethacrylate), poly(vinyl alcohol), or poly(propylene fumarate). In some embodiments, the insert comprises Gelfoam® R. In some embodiments, the insert is a polyacrylic acid of 450 kDa-cysteine conjugate.

In some embodiments, the ophthalmic composition is a ophthalmic film. Polymers suitable for such films include, but are not limited to, those described in Wagh, et al. (*ibid*), In some embodiments, the film is a soft-contact lens, such as ones made from copolymers of N,N-diethylacrylamide and methacrylic acid crosslinked with ethyleneglycol dimethacrylate.

In some embodiments, the ophthalmic composition comprises microspheres or nanoparticles. In some embodiment, the microspheres comprise gelatin. In some embodiments, the microspheres are injected to the posterior segment of the eye, in the choroidal space, in the sclera, intravitreally or sub-retinally. In some embodiments, the microspheres or nanoparticles comprises a polymer including, but not limited to, those described in Wagh, et al. (*ibid*), which is incorporated herein by reference in its entirety. In some embodiments, the polymer is chitosan, a polycarboxylic acid such as polyacrylic acid, albumin particles, hyaluronic acid esters, polyitaconic acid, poly(butyl)cyanoacrylate, polycaprolactone, poly(isobutyl)caprolactone, poly(lactic acid-co-glycolic acid), or poly(lactic acid). In some embodiments, the microspheres or nanoparticles comprise solid lipid particles.

In some embodiments, the ophthalmic composition comprises an ion-exchange resin. In some embodiments, the ion-exchange resin is an inorganic zeolite or synthetic organic resin. In some embodiments, the ion-exchange resin includes, but is not limited to, those described in Wagh, et al. (*ibid*), which is incorporated herein by reference in its entirety. In some embodiments, the ion-exchange resin is a partially neutralized polyacrylic acid.

In some embodiments, the ophthalmic composition is an aqueous polymeric suspension. In some embodiments, the therapeutic agent or a polymeric suspending agent is suspended in an aqueous medium. In some embodiments, the aqueous polymeric suspensions may be formulated so that they retain the same or substantially the same viscosity in the eye that they had prior to administration to the eye. In some embodiments, they may be formulated so that there is increased gelation upon contact with tear fluid.

Labeled Compounds and Assay Methods

Another aspect of the present invention relates to labeled compounds of the invention (radio-labeled, fluorescent-labeled, etc.) that would be useful not only in imaging techniques but also in assays, both *in vitro* and *in vivo*, for localizing and quantitating JAK in tissue samples, including human, and for identifying JAK ligands by inhibition binding of a labeled compound. Accordingly, the present invention includes JAK assays that contain such labeled compounds.

The present invention further includes isotopically-labeled compounds of the invention. An "isotopically" or "radio-labeled" compound is a compound of the invention where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (*i.e.*, naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I and ^{131}I . The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for *in vitro* JAK labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I , ^{35}S or will generally be most useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br will generally be most useful.

It is to be understood that a "radio-labeled " or "labeled compound" is a compound that has incorporated at least one radionuclide. In some embodiments the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br . In some embodiments, the compound incorporates 1, 2, or 3 deuterium atoms.

The present invention can further include synthetic methods for incorporating radio-isotopes into compounds of the invention. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and an ordinary skill in the art will readily recognize the methods applicable for the compounds of invention.

A labeled compound of the invention can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (*i.e.*, test compound) which is labeled can be evaluated for its ability to bind a JAK by monitoring its concentration variation when contacting with the JAK, through tracking of the labeling. For example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to a JAK (*i.e.*, standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the JAK directly correlates to its binding affinity. Conversely, in some other screening assays, the standard compound is labeled and test compounds are unlabeled. Accordingly, the

concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.

5 *Kits*


The present invention also includes pharmaceutical kits useful, for example, in the treatment or prevention of JAK-associated diseases or disorders, such as cancer, which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention. Such
10 kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for
15 administration, and/or guidelines for mixing the components, can also be included in the kit.

Synthesis

The compound of Formula 1a can be synthesized according to the
20 embodiments above and as further illustrated by Schemes I-III below. In Scheme I, the nucleophilic coupling of (i) and (ii) in the presence of an amine base (*e.g.*, *N,N*-diisopropylethylamine) affords the isomeric mixture of compounds (iii-a) and (iii-b). Subsequent reduction of the nitro group under hydrogenation conditions (*e.g.*, in the presence of hydrogen gas and a hydrogen catalyst (*e.g.* palladium on carbon)) affords
25 the isomeric mixture of compounds (iv-a) and (iv-b). Coupling of (iv-a) and (iv-b) with (R)-(+)-lactamide in the presence of an alkyloxonium reagent (*e.g.*, triethyloxonium tetrafluoroborate) and subsequent cyclization affords the a mixture of the fused-tricyclic compounds of Formula 1a and Formula 1b. Subsequent reaction under reverse racemization conditions (*e.g.* in the presence of a strong base (*e.g.*
30 potassium *tert*-butoxide)) yields a single isomer of the compound of Formula 1a, 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-


yl)tetrahydro-2H-pyran-2-yl)acetonitrile, which may be optionally hydrated to form the monohydrate compound of Formula II.

Scheme I




- 5 In Scheme II, compounds of Formula VI can be produced by reaction of **(i)** in the presence of a reducing agent (*e.g.*, lithium borohydride) affords the amine-protected diol **(ii)**, which is asymmetrically protected in the presence of 2,2-dimethoxypropane and boron trifluoride diethyl etherate to form oxazolidine **(iii)**. Reaction of the free alcohol **(iii)** under oxidizing conditions affords the aldehyde **(iv)**, which is then reacted under modified Wittig-Horner conditions (*e.g.* in the presence of diethyl (cyanomethyl)phosphonate and a potassium tert-butoxide) to form a mixture of the *cis*- and *trans*- isomers **(v)**. Deprotection of **(v)** in the presence of a strong acid (*e.g.*, p-TsOH) affords compound **(vi)**, which is cyclized in the presence of a strong base (*e.g.* sodium methoxide) to form the amine-protected pyran of compound **(vii)**. Deprotection under standard conditions (*e.g.*, in the presence of a strong acid) affords the amine salt **(viii)**.
- 10
- 15

Scheme II



In Scheme III, compounds of Formula VI can be produced. Accordingly, asymmetric carbonyl reduction of carboxylic acid (**i**) affords the hydroxy-ester compound (**ii**), which is subsequently cyclized in the presence of a weak organic acid (*e.g.* acetic acid) to form amino-lactone (**iii**). Reaction of (**iii**) under modified Wittig conditions (*e.g.*, in the presence of cyanomethyl (trimethyl)phosphonium chloride and potassium tert-butoxide, or alternatively in the presence of tributyl(cyanomethyl)phosphonium chloride and potassium tert-butoxide) afford the mixture of *cis*- and *trans*- isomers of (**iv**). Reduction of the alkene (**iv**) in the presence of a hydrogen gas and a hydrogenation catalyst (*e.g.*, palladium on carbon) affords the mixture of *R*- and *S*- isomers of (**v**). Deprotection of the amine under standard conditions (*e.g.*, in the presence of a strong acid) affords amine salt (**vi**).


Scheme III



Intermediate (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole can be prepared, for example, according to the process shown in Scheme IV. For example,


reaction of (*R*)-2-hydroxypropanamide (i) with 2,2-dimethoxypropane (ii) in the presence of a boron-trihalogen etherate (e.g., BF₃OEt₂) affords (*R*)-2,2,5-trimethyloxazolidin-4-one (iii). Reaction of compound (iii) with a methylating agent (e.g., Me₃O⁺BF₄⁻) and subsequent reaction in the presence of a base (e.g., sodium carbonate) affords (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole (v).

Scheme IV



A mixture of the compounds of Formulae 1a and 1b can also be prepared, for example, according to the process shown in Scheme V. (*R*)-2,2,5-trimethyloxazolidin-4-one (i) is first reacted with a methylating agent (e.g., Me₃O⁺BF₄⁻) to form (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole (ii), which is subsequently reacted with a mixture of 2-((2*R*,5*S*)-5-(6-aminothieno[3,2-*Z'*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothieno[3,2-*Z'*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile (iii) to afford a mixture of the compounds of Formulae 1a and 1b (iv).

Scheme V




A mixture of the compounds of Formulae 1a and 1b can also be prepared, for example, according to the process shown in Scheme VI. (*R*)-2-hydroxypropanamide (i) is first protected under standard alcohol protection conditions (e.g., reaction with acetyl chloride) to form (*R*)-1-amino-1-oxopropan-2-yl acetate (ii), which is then reacted with cyanuric chloride to form (*R*)-1-cyanoethyl acetate (iii). Subsequent reaction of (iii) with ethanol in the presence of an acid catalyst (e.g., hydrogen chloride) affords (*R*)-ethyl 2-hydroxypropanimidate (iv) which is then reacted with a

mixture of 2-((2*R*,5*S*)-5-(6-aminothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile (**v**) to form a mixture of the compounds of Formulae 1a and 1b (**vi**).

5

Scheme VI



EXAMPLES

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially the same results.

Open access prep. LC-MS purification of some of the compounds prepared was performed on Waters mass directed fractionation systems. The basic equipment setup, protocols, and control software for the operation of these systems have been described in detail in literature. *See e.g.* "Two-Pump At Column Dilution Configuration for Preparative LC-MS", K. Blom, *J. Combi. Chem.*, 4, 295 (2002); "Optimizing Preparative LC-MS Configurations and Methods for Parallel Synthesis Purification", K. Blom, R. Sparks, J. Doughty, G. Everlof, T. Haque, A. Combs, *J. Combi. Chem.*, 5, 670 (2003); and "Preparative LC-MS Purification: Improved Compound Specific Method Optimization", K. Blom, B. Glass, R. Sparks, A. Combs, *J. Combi. Chem.*, 6, 874-883 (2004). The compounds separated were typically subjected to analytical liquid chromatography mass spectrometry (LCMS) for purity under the following conditions: Instrument; Agilent 1100 series, LC/MSD, Column: Waters Sunfire™ Cis 5 μm, 2.1 x 5.0 mm, Buffers: mobile phase A: 0.025% TFA in

water and mobile phase B: 0.025% TFA in acetonitrile; gradient 2% to 80% of B in 3 minutes with flow rate 1.5 mL/minute.

Some of the compounds prepared were analyzed using Differential Scanning Calorimetry (DSC). Typical DSC conditions are as follows:

5 TA Instruments Differential Scanning Calorimetry, Model Q200 with autosampler. Temperature range: 30 - 350 °C; Temperature ramp rate: 10 °C/min; Tzero aluminum sample pan and lid; nitrogen gas flow at 50 mL/min.

Some of the compounds prepared were analyzed using Thermogravimetric Analysis (TGA). Typical TGA conditions are as follows:

10 TA Instrument Thermogravimetric Analyzer, Model Q500. Temperature range: 20 °C to 600 °C; Temperature ramp rate: 20 °C/min; nitrogen purge flow: 40 mL/min; Balance purge flow: 60 mL/min; Sample purge flow; platinum sample pan.

PerkinElmer Thermogravimetric Analyzer, Model Pyris 1. Temperature range: 25°C to 300 °C; Temperature ramp rate: 10 °C/min; Nitrogen purge gas flow: 60 mL/min; TGA ceramic crucible sample holder.

Some of the compounds prepared were analyzed using X-Ray Powder Diffraction (XRPD). Typical XRPD conditions are as follows:

20 Bruker D2 PHASER X-Ray Powder Diffractometer instrument; X-ray radiation wavelength: 1.05406 Å CuK α 1; X-ray power: 30 KV, 10 mA; Sample powder: dispersed on a zero-background sample holder; General measurement conditions: Start Angle - 5 degree; Stop Angle - 60 degree; Sampling - 0.015 degree; Scan speed - 2 degree/min.

25 Rigaku MiniFlex X-ray Powder Diffractometer; X-ray radiation is from Copper Cu at 1.054056 Å with K β filter; X-ray power: 30 KV, 15 mA; Sample powder: dispersed on a zero-background sample holder. General measurement condition: Start Angle - 3 degree; Stop Angle - 45 degree; Sampling - 0.02 degree; Scan speed - 2 degree/min.

Some of the compounds prepared were analyzed using Dynamic Vapor Sorption (DVS). Typical DVS conditions are as follows:

30 SGA-100 Symmetric Vapor Sorption Analyzer from VTI Corporation. The moisture uptake profile was completed in four cycles in 10% relative humidity (RH) increments with the first adsorption from 25% to 95% RH, followed by desorption in


10% increments from 95% to 5% RH. The equilibration criteria were 0.0050 wt% in 5 minutes with a maximum equilibration time of 180 minutes; Data logging interval: 2.00 min or 0.0100 wt % . All adsorption and desorption were performed at room temperature (25 °C). No pre-drying step was applied.

5 Some of the compounds prepared were analyzed after treatment in a humidity chamber. Typical humidity chamber conditions are as follows:


Temperature: 40 °C; Relative humidity: 75%; Time: 5 days.

Temperature: 30 °C; Relative humidity: 90%; Time: 7 days.

10 **Example 1. 2-((2R,5S)-5-Aminotetrahydro-2H-pyran-2-yl)acetonitrile hydrochloride and 2-((2S,5S)-5-aminotetrahydro-2H-pyran-2-yl)acetonitrile hydrochloride (Procedure 1)**



Step 1. (S)-tert-Butyl 1,5-dihydroxypentan-2-ylcarbamate



15


In a 5-necked, 5-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, a reflux condenser, an addition funnel, and a nitrogen inlet was placed lithium borohydride (29.9 g, 1.37 mol, 2.0 eq.) at room temperature. Tetrahydrofuran (THF, 1.12 L) was charged to the flask while maintaining the internal temperature at below 40 °C. The resulting solution was then cooled to 0 - 5 °C before a solution of (5)-dimethyl 2-(tert-butoxycarbonylamino)pentanedioate (189 g, 0.687 mol) in THF (0.60 L) was added over a period greater than 50 minutes via the addition funnel. The resulting reaction mixture was gradually warmed to room temperature and stirred at room temperature for 17 h and then cooled to 15 °C.

20

25 Methanol (MeOH, 415 mL) was then added dropwise to the reaction mixture over 3 h to quench residual lithium borohydride. During addition of methanol, vigorous gas

evolution was observed. The quenched reaction mixture was then concentrated under reduced pressure to remove most of the solvents and the residue was treated with ethyl acetate (EtOAc, 350 mL) and water (300 mL). The resulting suspension was stirred at room temperature for 15 minutes before being filtered through a sintered funnel. The two phases of the filtrate were separated and the aqueous phase was extracted with EtOAc (100 mL). The combined organic phases were washed with brine (150 mL), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to afford crude (5)-*tert*-butyl 1,5-dihydroxypentan-2-ylcarbamate (135 g, 89.7%) as a white solid, which was used in the subsequent reaction without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 6.42 (d, *J* = 8.5 Hz, 1H), 4.52 (t, *J* = 5.5 Hz, 1H), 4.33 (t, *J* = 5.1 Hz, 1H), 3.42 - 3.21 (m, 4H), 3.18 (dt, *J* = 10.5, 6.1 Hz, 1H), 1.54 - 1.44 (m, 1H), 1.41 - 1.28 (m, 11H), 1.22 - 1.16 (m, 1H) ppm.


Step 2. (S)-tert-Butyl 4-(3-hydroxypropyl)-2,2-dimethyloxazolidine-3-carboxylate



In a 5-necked, 5-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, an addition funnel, and a nitrogen inlet was placed (*S*)-*tert*-butyl 1,5-dihydroxypentan-2-ylcarbamate (105 g, 0.479 mol) and acetone (4.0 L) at room temperature. The solution was then treated with 2,2-dimethoxypropane (64.8 g, 0.622 mol, 1.3 eq.) and boron trifluoride diethyl etherate (BF₃-Et₂O, 6.1 mL, 0.048 mol, 0.10 eq.) sequentially at room temperature. The resulting reaction mixture was stirred at room temperature for 18 h before triethylamine (TEA, 40.0 mL, 0.287 mol, 0.60 eq.) was added. The resulting mixture was then concentrated under reduced pressure to remove most of the solvents and the residue was treated with EtOAc (1000 mL) and water (200 mL). The two phases were separated and the organic phase was washed with brine (200 mL), dried over sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography (S102, gradient elution with 0 to 50% of ethyl acetate in hexanes) to afford (5)-*tert*-butyl 4-(3-hydroxypropyl)-2,2-dimethyloxazolidine-3-carboxylate (95.0 g, 76.5%) as a colorless oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.39 (t, *J* = 5.1 Hz, 1H), 3.85 (dd, *J* =


8.7, 5.7 Hz, 1H), 3.72 (s, 1H), 3.66 (dd, $J = 8.8, 1.3$ Hz, 1H), 3.42 - 3.32 (m, 2H), 1.61-1.54 (m, 1H), 1.51 - 1.24 (m, 18H) ppm.

Step 3. (S)-tert-Butyl 2,2-dimethyl-4-(3-oxopropyl)oxazolidine-3-carboxylate




5 In a 5-necked 5-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, an addition funnel, and a nitrogen inlet was placed (*S*)-*tert*-butyl 4-(3-hydroxypropyl)-2,2-dimethyloxazolidine-3-carboxylate (110.0 g, 0.424 mol) and dichloromethane (DCM, 1200 mL) at room temperature. The resulting solution was then treated with 2,2,6,6-tetramethyl-1-piperidinyloxy free radical
10 (TEMPO, 6.6 g, 0.042 mol, 0.10 eq.) and *tetra-n*-butylammonium chloride (11.8 g, 0.042 mol, 0.10 eq.) at room temperature before an aqueous solution of sodium bicarbonate (174 g, 2.08 mol, 5.0 eq) and an aqueous solution of potassium carbonate (28.7 g, 0.208 mol, 0.50 eq) were sequentially added to the reaction mixture at room temperature. While the resulting mixture was stirred vigorously, *N*-chlorosuccinimide
15 (69.1 g, 0.518 mol, 1.22 eq.) was added at room temperature. The resulting reaction mixture was then stirred at room temperature for an additional 3 h before being transferred to a separation funnel. The two phases were separated and the aqueous phase was extracted with DCM (250 mL). The combined organic layers were washed with brine (2 x 300 mL), dried over anhydrous sodium sulfate, concentrated under
20 reduced pressure, and dried under vacuum to afford crude (*S*)-*tert*-butyl 2,2-dimethyl-4-(3-oxopropyl)oxazolidine-3-carboxylate (109.1 g, 100%) as a pale orange oil, which was used in the subsequent reaction without further purification. ^1H NMR (300 MHz, DMSO- d_6) δ 9.64 (s, 1H), 4.01 - 3.60 (m, 3H), 2.39 (dd, $J = 10.9, 4.0$ Hz, 2H), 1.84 - 1.64 (m, 2H), 1.47-1.38 (m, 15H) ppm.

25 *Step 4. (S)-tert-Butyl 4-(4-cyanobut-3-enyl)-2,2-dimethyloxazolidine-3-carboxylate*



In a 3-necked, 3-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, and an addition funnel under nitrogen atmosphere was placed potassium *tert*-butoxide (1 M solution in THF, 1000 mL, 1.00 mol, 1.11 eq.) and tetrahydrofuran (THF, 700 mL) at room temperature. The solution was cooled to 0 - 5 °C and diethyl cyanomethylphosphonate (177.2 g, 1.00 mol, 1.11 eq.) was added dropwise via the addition funnel while maintaining the internal temperature at below 10 °C. The resulting reaction mixture was then stirred at room temperature for 2 h, and labeled as Solution A for the subsequent reaction. In a separate 5-necked, 12-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, an addition funnel, and a nitrogen inlet was placed (5)-*tert*-butyl 2,2-dimethyl-4-(3-oxopropyl)oxazolidine-3-carboxylate (232.0 g, 0.901 mol) and THF (3500 mL) at room temperature. The resulting solution was cooled to 0 - 5 °C, and was labeled as Solution B. Solution A was then added to Solution B dropwise while maintaining the internal temperature at below 5 °C. The resulting reaction mixture was gradually warmed to room temperature and stirred at room temperature for 1 h. Water (500 mL) was charged to the reaction mixture and two phases were separated. The aqueous phase was then extracted with EtOAc (2 x 100 mL). The combined organic phases were washed with brine (2 x 100 mL), dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, gradient elution with 0 to 30% of ethyl acetate in hexanes) to provide (5)-*tert*-butyl 4-(4-cyanobut-3-enyl)-2,2-dimethyloxazolidine-3-carboxylate (179.0 g, 70.8%) as a colorless oil. ¹H NMR (300 MHz, DMSO-*d*₆) δ 6.84 (dt, *J* = 16.4, 6.8 Hz, 0.5H), 6.66 (dt, *J* = 10.9, 7.7 Hz, 0.5H), 5.83 - 5.51 (m, 1H), 3.92 - 3.49 (m, 3H), 2.35 - 2.05 (m, 2H), 1.73 - 1.52 (m, 2H), 1.45 - 1.38 (m, 15H).


Step 5. (*S*)-*tert*-Butyl 6-cyano-1-hydroxyhex-5-en-2-ylcarbamate



In a 5-necked, 5-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, an addition funnel, and a nitrogen inlet was placed (*S*)-*tert*-butyl 4-(4-cyanobut-3-enyl)-2,2-dimethyloxazolidine-3-carboxylate (179.0 g, 0.638

mol) and methanol (2000 mL) at room temperature. The resulting colorless solution was cooled to 0 - 5 °C before being treated with catalytic amount of p-toluenesulfonic acid (PTSA, 12.1 g, 0.064 mol, 0.10 eq.). The resulting reaction mixture was gradually warmed to room temperature and stirred at room temperature for an additional 17 h. An aqueous solution of sodium bicarbonate (100 mL) was then added to the reaction mixture at room temperature and the resulting mixture was concentrated under reduced pressure to about 500 mL. The residue was then extracted with ethyl acetate (2 x 500 mL). The combined organic phases were washed with brine (2 x 100 mL), dried over anhydrous magnesium sulfate, concentrated under reduced pressure, and dried under vacuum to afford crude (5)-tert-butyl 6-cyano-1-hydroxyhex-5-en-2-ylcarbamate (144.2 g, 94%) as a light pink oil, which was used in the subsequent reaction without further purification. ¹H NMR (300 MHz, CDCl₃) δ 6.72 (dt, *J* = 16.3, 6.9 Hz, 0.5H), 6.53 (dt, *J* = 10.9, 7.6 Hz, 0.5H), 5.51 - 5.08 (m, 1H), 4.73 (s, 1H), 3.83 - 3.34 (m, 3H), 2.49 (dt, *J* = 11.3, 5.9 Hz, 1H), 2.31 (q, *J* = 7.0 Hz, 1H), 1.74-1.57 (m, 3H), 1.44 (s, 9H).

Step 6. (S)-tert-Butyl 6-(cyanomethyl)tetrahydro-2H-pyran-3-ylcarbamate (mixture of cis- and trans- isomers)



In a 5-necked, 5-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, an addition funnel, and a nitrogen inlet was placed (S)-tert-butyl 6-cyano-1-hydroxyhex-5-en-2-ylcarbamate (173.0 g, 0.720 mol) and tetrahydrofuran (1700 mL) at room temperature. The solution was cooled to 0 - 5 °C before a solution of sodium methoxide (0.5 M solution in methanol, 21.30 mL, 0.011 mol, 0.015 eq) in THF (90 mL) was added dropwise via the addition funnel while maintaining the internal temperature at below 5 °C. The resulting reaction mixture was stirred at below 5 °C for an additional 1 h before brine (500 mL) and tert-butyl methyl ether (TBME, 400 mL) were added to the reaction mixture sequentially. The

two phases were separated and the aqueous phase was extracted with TBME (250 mL). The combined organic phases were washed with brine (250 mL), dried over anhydrous magnesium sulfate, concentrated under reduced pressure, and dried under vacuum to afford crude (5)-*tert*-butyl 6-(cyanomethyl)tetrahydro-2 *H*-pyran-3-


5 ylcarbamate (173.0 g, 100%) as a viscous solid. The crude product, obtained as a mixture of *cis*- and *trans*-isomers, was used in the subsequent reaction without further purification. ¹H NMR (400 MHz, CDCl₃) δ 5.13 (s, 0.5H), 4.25 (s, 0.5H), 4.10 (ddd, *J* = 10.9, 4.7, 2.2 Hz, 0.5H), 3.89 (d, *J* = 12.1 Hz, 0.5H), 3.72 (s, 0.5H), 3.66 - 3.57 (m, 1.5H), 3.56 - 3.49 (m, 0.5H), 3.03 (t, *J* = 10.8 Hz, 0.5H), 2.54 (dd, *J* = 6.0, 2.5
10 Hz, 2H), 2.19 - 2.08 (m, 0.5H), 2.00 (d, *J* = 13.2 Hz, 0.5H), 1.87 (d, *J* = 13.1 Hz, 0.5H), 1.65 - 1.62 (m, 2H), 1.44 (d, *J* = 6.5 Hz, 9H), 1.38 - 1.24 (m, 0.5H).

Step 7. 2-((2R,5S)-5-Aminotetrahydro-2H-pyran-2-yl)acetonitrile hydrochloride and 2-((2S,5S)-5-aminotetrahydro-2H-pyran-2-yl)acetonitrile hydrochloride (mixture of trans- and cis- isomers)


15 In a 5-necked, 5-L round bottom flask equipped with an overhead mechanical stirrer, a thermocouple, an addition funnel, and a nitrogen inlet was placed (*S*)-*tert*-butyl 6-(cyanomethyl)tetrahydro-2 *H*-pyran-3-ylcarbamate (a mixture of *cis*- and *trans*- isomers, 173.0 g, 0.720 mol) and dichloromethane (1730 mL) at room temperature. The resulting solution was then treated with a solution of 4.0 M
20 hydrogen chloride (HCl) in 1,4-dioxane (720 mL, 2.880 mol, 4.0 eq.) via the addition funnel. The resulting reaction mixture was stirred at room temperature for 2 h before *tert*-butyl methyl ether (TBME, 1000 mL) was added. The resulting suspension was stirred at room temperature for 30 minutes. The white solids formed were collected by filtration, washed with TBME (280 mL), and dried in a vacuum oven with nitrogen
25 sweeping at 50 °C to afford 2-((2*R*,5*S*)-5-aminotetrahydro-2 *H*-pyran-2-yl)acetonitrile hydrochloride and 2-((2*S*,5*S*)-5-aminotetrahydro-2 *H*-pyran-2-yl)acetonitrile hydrochloride (114.0 g, 89.6%) as a mixture of *trans*- and *cis*- isomers. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.31 (s, 3H), 4.03 (ddd, *J* = 10.8, 4.5, 2.2 Hz, 0.5H), 3.92 (d, *J* = 12.6 Hz, 0.5H), 3.71 - 3.58 (m, 1H), 3.51 (ddp, *J* = 8.8, 4.4, 2.1 Hz, 0.5H), 3.36
30 (s, 0.5H), 3.25 (s, 0.5H), 3.04 (s, 0.5H), 2.87 - 2.56 (m, 2H), 2.08 (d, *J* = 12.7 Hz, 0.5H), 1.98 - 1.72 (m, 1.5H), 1.70 - 1.46 (m, 1.5H), 1.36 (qd, *J* = 13.2, 3.9 Hz, 0.5H)

ppm; ^{13}C NMR (101 MHz, DMSO- d_6) δ 118.48 (118.33), 72.52 (72.07), 67.75 (67.56), 45.53 (44.36), 28.59 (24.33), 26.78 (24.06), 23.42 (23.14) ppm; LCMS calculated for $\text{C}_7\text{H}_{12}\text{N}_2\text{O}$: 140.2 (free amine), Found: 141 ($\text{M}^+ + \text{H}$); Salt ratio (HCl to the free amine): 1.00.

5 **Example 2. 2-((2R,5S)-5-Aminotetrahydro-2H-pyran-2-yl)acetonitrile hydrochloride and 2-((2S,5S)-5-aminotetrahydro-2H-pyran-2-yl)acetonitrile hydrochloride (Procedure 2)**



Step 1. methyl (4S)-4-[(tert-butoxycarbonyl)amino]-5-hydroxypentanoate



10

A solution of (2S)-2-[(tert-butoxycarbonyl)amino]-5-methoxy-5-oxopentanoic acid (40.00 g, 153.1 mmol) in tetrahydrofuran (600 mL) was treated with 4-methylmorpholine (17 g, 168.1 mmol, 1.1 eq.) at room temperature. The resulting mixture was cooled to 0 °C before being treated with ethyl chloroformate (18 g, 165.9 mmol, 1.08 eq.) dropwise. The resulting reaction mixture was stirred at 0 °C for an additional 20 minutes before being filtered through Celite. The Celite bed was washed with THF (50 mL) and the clear filtrate solution was cooled to 0 °C. The cooled solution was then treated with solid NaBH_4 (3.00 g, 79.3 mmol, 0.52 eq.) followed by a solution of NaBH_4 (2.9 g, 76.1 mmol, 0.50 eq.) in diglyme (40 mL). The resulting reaction mixture was stirred at room temperature for 3 h before being treated with an aqueous 1.0 M HCl solution (100 mL). The mixture was then stirred at room temperature for 10 minutes. The inorganic solids were filtered off and rinsed with EtOAc (100 mL). Ethyl acetate (1100 mL) was then added into the filtrate and the combined filtrate solution was concentrated under reduced pressure to remove the solvents (1200 mL). The remaining mixture was then washed with saturated aqueous sodium bicarbonate (50 mL) and water (2 X 50 mL). The organic phase was then concentrated under reduced pressure to afford crude methyl (4S)-4-[(tert-


15

20

25

butoxycarbonyl)amino]-5-hydroxypentanoate (37.9 g), which was used in the subsequent reaction without further purification.

Step 2. (S)-tert-butyl (6-oxotetrahydro-2H-pyran-3-yl)carbamate




5

A solution of crude methyl (4S)-4-[(tert-butoxycarbonyl)amino]-5-hydroxypentanoate (37.9 g, 153.1 mmol) in toluene (600 mL) and acetic acid (200 mL) was heated to reflux for 60 minutes. The solvents were then removed by the azeotropic distillation at atmospheric pressure. During the azeotropic distillation, an additional amount of toluene (600 mL) was introduced and a total of 1200 mL of the mixed solvents was removed. The residual mixture was then cooled to room temperature over 3 h and stirred at room temperature for 2 h. The solids were collected by filtration, washed with n-heptane (100 mL), and dried under reduced pressure at below 40 °C to afford the desired tert-butyl [(3S)-6-oxotetrahydro-2H-pyran-3-yl]carbamate (25 g, 75% overall yield for two steps). ¹H NMR (400 MHz, CDCl₃) δ 4.39 (dd, *J* = 11.5, 5.4 Hz, 1H), 4.18 (dd, *J* = 11.4, 4 Hz, 1H), 4.00 (s, 1H), 2.69 - 2.54 (m, 2H), 2.22 (dq, *J* = 13.2, 6.6 Hz, 1H), 1.86 (dq, *J* = 14.6, 7.3 Hz, 1H), 1.43 (s, 9H); GCMS Calculated for C₁₀H₁₇NO₄: 215.25; Found: 215.1.

15

Step 3. tert-Butyl [(3S,6Z)-6-(cyanomethylene)tetrahydro-2H-pyran-3-yl]carbamate and tert-butyl [(3S,6E)-6-(cyanomethylene)tetrahydro-2H-pyran-3-yl]carbamate

20



Method A.

A solution of cyanomethyl (trimethyl)phosphonium chloride (8.8 g, 58.06 mmol, 1.25 eq.) in *N,N*-dimethylacetamide (DMAC, 50 ml) was degassed and cooled to 0 °C. The chilled solution was then treated with a solution of 1.0 M lithium hexamethyldisilazide in THF (LHMDS, 53.4 mL, 53.4 mmol, 1.15 eq.) at 0 °C. The

25

resulting mixture was stirred at 0 °C for 60 minutes before being treated with a solution of *tert*-butyl [(3*S*)-6-oxotetrahydro-2 *H*-pyran-3-yl]carbamate (10 g, 46.46 mmol) in THF (40 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 2 h followed by stirring overnight at room temperature. The reaction mixture was then quenched with water (100 mL) and the two phases were separated. The aqueous phase was extracted with *tert*-butyl methyl ether (TBME, 2 X 150 mL). The combined organic phase was washed with water (2 X 150 mL) before being concentrated under reduced pressure to afford the crude desired product (11 g, 99%) as a mixture of *tert*-butyl [(3*S*,6*Z*)-6-(cyanomethylene)tetrahydro-2 *H*-pyran-3-yl]carbamate and *tert*-butyl [(3*S*',6*E*)-6-(cyanomethylene)tetrahydro-2 *H*-pyran-3-yl]carbamate, which was used in the subsequent reaction without further purification and separation. Analytical samples of the individual isomers were obtained by silica gel column chromatography (SiO₂, gradient elution with 0 - 30% of *n*-Butyl methyl ether (MTBE) in heptane).

For (3*S*,6*E*)-isomer: ¹H NMR (400 MHz, CDCl₃) δ 4.82 (s, 1H), 4.67 (s, 1H), 4.12 (dd, 1H), 3.93 (m, 1H), 3.81 (dd, 1H), 2.79 (dt, 1H), 2.58 (m, 1H), 2.12 (dt, 1H), 1.69 (dt, 1H), 1.40 (s, 9H); GCMS Calculated for C₁₂H₁₈N₂O₃: 238.28; Found: 238.1.

For (3*S*,6*Z*)-isomer: ¹H NMR (400 MHz, CDCl₃) δ 4.81 (s, 1H), 4.48 (s, 1H), 4.19 (dt, 1H), 3.95 (m, 2H), 2.48 (m, 2H), 2.09 (m, 1H), 1.58 (m, 1H), 1.40 (s, 9H); GCMS Calculated for C₁₂H₁₈N₂O₃: 238.28; Found: 238.1.

Method B.


A mixture of tributyl(cyanomethyl)phosphonium chloride (593 g, 2.14 mol) in THF (5.9 L) was treated with solid potassium *tert*-butoxide (220 g, 1.97 mol) in portions at 0 °C over 5 minutes. The resulting mixture was gradually warmed to room temperature over 4 h. The resulting ylide solution was then treated with (5)-*tert*-butyl (6-oxotetrahydro-2 *H*-pyran-3-yl)carbamate (184 g, 0.855 mol) as a solid at room temperature and the resulting reaction mixture was heated at 70 °C and stirred at 70 °C for 16 h. When NMR analysis indicated that the starting material was consumed, the reaction mixture was gradually cooled to room temperature. The resulting slurry was poured into a 50-L separation funnel containing saturated aqueous sodium bicarbonate (5 L) and ice (1000 g). Ethyl acetate (6 L) was added and the mixture was stirred for 10 minutes. The two layers were separated and the aqueous layer was extracted with

EtOAc (3 L). The combined organic layers were washed with saturated aqueous sodium chloride (3 X 4 L), dried over sodium sulfate, and concentrated under reduced pressure. The resulting dark oil was dissolved in DCM (500 mL) and purified by column chromatography (S102, gradient elution with 20 - 40% ethyl acetate in heptanes) to afford the desired product (177 g, 87% yield) as a mixture of *tert*-butyl [(3*S*,6*Z*)-6-(cyanomethylene)tetrahydro-2 *H*-pyran-3-yl]carbamate and *tert*-butyl [(3*S'*,6*E*)-6-(cyanomethylene)tetrahydro-2 *H*-pyran-3-yl]carbamate, which was used in the subsequent reaction without further purification and separation. Analytical samples of the two isomers were obtained by silica gel column chromatography (S102, gradient elution with 0 - 30 % of *t*-butyl methyl ether (TBME) in heptane).

For (3*S*,6*E*)-isomer: ^1H NMR (400 MHz, CDCl₃) δ 4.82(s, 1H), 4.67(s, 1H), 4.12(dd, 1H), 3.93(m, 1H), 3.81(dd,1H), 2.79 (dt, 1H), 2.58 (m, 1H), 2.12(dt, 1H), 1.69(dt, 1H), 1.40 (s, 9H); GCMS Calculated for C₁₂H₁₈N₂O₃: 238.28; Found: 238.1.

For (3*S*,6*Z*)-isomer: ^1H NMR (400 MHz, CDCl₃) δ 4.81 (s, 1H), 4.48 (s,1H), 4.19 (dt, 1H), 3.95 (m, 2H), 2.48 (m, 2H), 2.09 (m, 1H), 1.58 (m, 1H), 1.40 (s, 9H); GCMS Calculated for C₁₂H₁₈N₂O₃: 238.28; Found: 238.1.

*Step 4. tert-Butyl [(3*S*,6*S*)-6-(cyanomethyl)tetrahydro-2*H*-pyran-3-yl]carbamate and tert-butyl [(3*S*, 6*R*)-6-(cyanomethyl)tetrahydro-2*H*-pyran-3-yl]carbamate*



A mixture of *tert*-butyl [(3*S*,6*Z*)-6-(cyanomethylene)tetrahydro-2 *H*-pyran-3-yl]carbamate and *tert*-butyl [(3*S'*,6*E*)-6-(cyanomethylene)tetrahydro-2 *H*-pyran-3-yl]carbamate (110 g, 0.462 mol) and palladium on carbon (10 wt % Pd-C, 50% wet, 14 g) in methanol (1.1 L) was hydrogenated at 20 psi until hydrogen uptake ceased (4 to 12 h). The reaction mixture was filtered through Celite (100 g) and the Celite bed was washed with methanol (500 mL). The combined filtrate was concentrated under reduced pressure to afford the crude reduction products as an off-white solid. The crude products were further dried by azeotropic distillation with toluene (2 x 500 mL) and *n*-heptane (2 x 500 mL) under reduced pressure to remove residual methanol and

water to afford *tert*-butyl [(3*S*,6*S*)-6-(cyanomethyl)tetrahydro-2 *H*-pyran-3-yl]carbamate and *tert*-butyl [(3*S*,6*R*)-6-(cyanomethyl)tetrahydro-2 *H*-pyran-3-yl]carbamate (111 g, nearly quantitative yield) as a mixture of diastereomers, which was used directly in the subsequent reaction without further purification and

5 separation. ¹H NMR (300 MHz, CDCl₃, as a mixture of two diastereomers) δ 5.15 (br s, 1 H), 4.29 (br s, 1 H), 4.14 - 4.07 (m, 1 H), 3.94 - 3.87 (m, 1 H), 3.77 - 3.44 (m, 5 H), 3.09 - 3.01 (m, 1 H), 2.58 - 2.49 (m, 4 H), 2.18 - 2.09 (m, 1 H), 2.05 - 1.95 (m, 1 H), 1.92 - 1.81 (m, 1 H), 1.76 - 1.51 (m, 5 H), 1.46 (s, 9 H), 1.44 (s, 9 H); GCMS calculated for C₁₂H₂₀N₂O₃: 240.30; Found: 240.1.

10 *Step 5. 2-((2*S*,5*S*)-5-Aminotetrahydro-2*H*-pyran-2-yl)acetonitrile hydrochloride and 2-((2*R*,5*S*)-5-aminotetrahydro-2*H*-pyran-2-yl)acetonitrile hydrochloride*

A solution of *tert*-butyl [(3*S*,6*S*)-6-(cyanomethyl)tetrahydro-2 *H*-pyran-3-yl]carbamate and *tert*-butyl [(3*S*,6*R*)-6-(cyanomethyl)tetrahydro-2 *H*-pyran-3-yl]carbamate (240 g, 0.999 mol) in dichloromethane (2.44 L) was treated with a

15 solution of 4 M HCl in 1,4-dioxane (1 L, 3.99 mol, 4.0 eq.) dropwise over 30 minutes while keeping the internal temperature at below 25 °C. The resulting reaction mixture was stirred at room temperature for 3 h. When ¹H NMR analysis of the filtered sample indicated that the reaction was complete, *tert*-butyl methyl ether (TBME, 675 mL) was added to the reaction mixture and the resulting mixture was stirred at room


20 temperature for 30 minutes. The white solids formed were collected by filtration under a blanket of nitrogen, washed with TBME (600 mL), and further dried in a vacuum oven under a blanket of nitrogen for 16 h to afford 2-((2*S*,5*S*)-5-aminotetrahydro-2 *H*-pyran-2-yl)acetonitrile hydrochloride and 2-((2*R*,5*S*)-5-aminotetrahydro-2 *H*-pyran-2-yl)acetonitrile hydrochloride (151.7 g, 86% yield) as a

25 mixture of *cis*- and *trans*-isomers. ³Q NMR (400 MHz, DMSO-*d*₆) (a mixture of *cis*- and *trans*- isomers) δ 8.31 (s, 3H), 4.03 (ddd, *J* = 10.8, 4.5, 2.2 Hz, 0.5H), 3.92 (d, *J* = 12.6 Hz, 0.5H), 3.71 - 3.58 (m, 1H), 3.51 (ddp, *J* = 8.8, 4.4, 2.1 Hz, 0.5H), 3.36 (s, 0.5H), 3.25 (s, 0.5H), 3.04 (s, 0.5H), 2.87 - 2.56 (m, 2H), 2.08 (d, *J* = 12.7 Hz, 0.5H), 1.98 - 1.72 (m, 1.5H), 1.70 - 1.46 (m, 1.5H), 1.36 (qd, *J* = 13.2, 3.9 Hz, 0.5H) ppm;

30 ¹³C NMR (101 MHz, DMSO-*d*₆) δ 118.48 (118.33), 72.52 (72.07), 67.75 (67.56), 45.53 (44.36), 28.59 (24.33), 26.78 (24.06), 23.42 (23.14) ppm; LCMS Calculated for

C₇H₁₂N₂O: 140.2 (free amine), Found: 141 (M⁺ + H); salt ratio (HCl to the free amine): 1.00.

Example 3. 2-((2*S*,5*S*)-5-(6-Nitrothieno [3,2-6]pyridin-7-ylamino)tetrahydro- 2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-nitrothieno [3,2-*Z*]pyridin-7-ylamino)tetrahydro-2 *H*-pyran-2-yl)acetonitrile




A mixture of 2-((2*S*,5*S*)-5-aminotetrahydro-2 *H*-pyran-2-yl)acetonitrile hydrochloride and 2-((2*R*,5*S*)-5-aminotetrahydro-2 *H*-pyran-2-yl)acetonitrile hydrochloride (2464 g, 13.95 mol, 1.16 eq.), and 7-chloro-6-nitrothieno[3,2-*b*]pyridine (2581 g, 12.02 mol) in *N,N*-dimethylformamide (DMF, 12.3 L) was treated with diisopropylethylamine (DIPEA, 4897 g, 46.13 mol, 3.84 eq.) at room temperature. The resulting reaction mixture was heated to 65 - 75 °C for 2 - 6 h until HPLC showed the reaction was complete. Water (36.9 L) was added to the reaction mixture at above 55 °C to precipitate the crude desired product and the resulting mixture was gradually cooled to 15 - 30 °C and stirred at 15 - 30 °C for 1 - 2 h. The yellow solids were collected by filtration, washed with water (36.9 L) and *n*-heptane (30.8 L), and dried in a vacuum oven at 45 - 50 °C to constant weight to afford the crude desired product (3550 g, 92.8%) as a mixture of 2-((2*R*,5*S*)-5-(6-nitrothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-nitrothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2 *H*-pyran-2-yl)acetonitrile, which was used in the subsequent reaction without further purification and separation. Analytical samples of the *cis*-isomer and *trans*-isomers were obtained by column chromatography separation (SiO₂, gradient elution with 0-50% of ethyl acetate in heptane).

For **(2*R*,5*S*)-isomer**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.09 (s, 1H), 8.87 (d, *J* = 9.1 Hz, 1H), 8.35 (d, *J* = 5.5 Hz, 1H), 7.53 (d, *J* = 5.5 Hz, 1H), 4.32 (ddq, *J* = 14.4, 10.1, 4.3 Hz, 1H), 4.17 (ddd, *J* = 10.9, 4.5, 2.2 Hz, 1H), 3.66 (ddd, *J* = 9.2, 6.8, 4.5

Hz, 1H), 3.52 (t, $J = 10.6$ Hz, 1H), 2.85 (dd, $J = 17.0, 4.4$ Hz, 1H), 2.76 (dd, $J = 17.0, 6.8$ Hz, 1H), 2.28-2.23 (m, 1H), 191-1.81 (m, 2H), 1.61 - 1.43 (m, 1H) ppm; ^{13}C NMR (101 MHz, DMSO- d_6) δ 159.30, 147.11, 145.78, 136.17, 125.59, 125.47, 118.47, 116.54, 71.79, 70.18, 50.25, 29.95, 29.18, 23.14 ppm; $\text{C}_{14}\text{H}_{14}\text{N}_4\text{O}_3\text{S}$ (MW 318.3), LCMS (EI) m/z 319 ($\text{M}^+ + \text{H}$).

For **(2S,5S)-isomer**: ^1H NMR (400 MHz, DMSO- d_6) δ 9.69 (d, $J = 8.5$ Hz, 1H), 9.14 (s, 1H), 8.37 (d, $J = 5.5$ Hz, 1H), 7.54 (d, $J = 5.5$ Hz, 1H), 4.49 (d, $J = 8.6$ Hz, 1H), 4.03 (d, $J = 11.9$ Hz, 1H), 3.89 (d, $J = 10.4$ Hz, 1H), 3.75 (dd, $J = 7.1, 3.6$ Hz, 1H), 2.84 (dd, $J = 17.1, 4.3$ Hz, 1H), 2.71 (dd, $J = 17.1, 7.0$ Hz, 1H), 2.07-2.04 (m, 2H), 1.75 - 1.40 (m, 2H) ppm; ^{13}C NMR (101 MHz, DMSO- d_6) δ 159.15, 147.26, 145.49, 136.22, 125.56, 125.28, 118.16, 116.77, 72.53, 70.31, 47.99, 27.84, 25.08, 23.65 ppm; $\text{C}_{14}\text{H}_{14}\text{N}_4\text{O}_3\text{S}$ (MW 318.3), LCMS (EI) m/z 319 ($\text{M}^+ + \text{H}$).

Example 4. 2-((2*R*,5*S*)-5-(6-Aminothiено [3,2-6]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothiено [3,2-6]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile



A solution of 2-((2*R*,5*S*)-5-(6-nitrothiено [3,2-6]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-nitrothiено [3,2-6]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile (2000 g, 6.28 mol) in 2,2,2-trifluoroethanol (8.1 L) was treated with palladium on carbon (10 wt % Pd-C, 50% wet, 200 g) at room temperature. The resulting reaction mixture was degassed and refilled with nitrogen gas three times followed by hydrogen gas, three times. The hydrogenation reaction was run at 20 - 35 °C with 50 psi of hydrogen pressure.


When HPLC showed the reaction was complete, the pressure was reduced to atmospheric before being degassed and refilled with nitrogen three times. The mixture was then filtered through Celite (667 g) and the Celite bed was rinsed with methanol (8 L). The filtrate was then concentrated under reduced pressure at below

50 °C to afford the desired crude product (2242 g) as a mixture of 2-((2*R*,5*S*)-5-(6-aminothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2 *H*-pyran-2-yl)acetonitrile. This mixture, obtained as foam which contained residual solvents was inseparable on a silica gel column, and was used in the subsequent reaction without further purification.

For **(2*R*,5*S*)-isomer**: ^1H NMR (400 MHz, DMSO- d_6) δ 7.93 (s, 1H), 7.55 (d, J = 5.6 Hz, 1H), 7.20 (d, J = 5.6 Hz, 1H), 4.91 (d, J = 9.7 Hz, 1H), 4.82 (s, 2H), 4.06 - 3.77 (m, 2H), 3.55 (dtt, J = 8.7, 4.5, 2.1 Hz, 1H), 3.24 (t, J = 10.3 Hz, 1H), 2.92 - 2.62 (m, 2H), 2.20 - 1.97 (m, 1H), 1.88 - 1.70 (m, 1H), 1.66 - 1.51 (m, 1H), 1.50 - 1.37 (m, 1H) ppm; ^{13}C NMR (101 MHz, DMSO- d_6) δ 150.28, 136.88, 134.86, 128.10, 124.86, 124.82, 118.55, 117.95, 72.14, 71.70, 49.30, 30.79, 29.74, 23.32 ppm; $\text{C}_{14}\text{H}_{16}\text{N}_4\text{O}_2$ (MW 288.3), LCMS (EI) *m/z* 289 (M^+ + H).

For **(2*S*,5*S*)-isomer**: ^1H NMR (400 MHz, DMSO- d_6) δ 8.01 (s, 1H), 7.59 (d, J = 5.5 Hz, 1H), 7.22 (d, J = 5.5 Hz, 1H), 5.09 (d, J = 9.6 Hz, 1H), 4.76 (s, 2H), 3.99 (d, J = 9.6 Hz, 1H), 3.87 (d, J = 11.7 Hz, 1H), 3.75 (dd, J = 11.7, 2.2 Hz, 1H), 3.66 (dddd, J = 10.3, 7.2, 4.4, 2.6 Hz, 1H), 2.85 - 2.62 (m, 2H), 1.86 - 1.69 (m, 2H), 1.63 (qd, J = 12.7, 12.2, 4.4 Hz, 1H), 1.52 - 1.44 (m, 1H) ppm; ^{13}C NMR (101 MHz, DMSO- d_6) δ 150.81, 137.91, 135.84, 128.17, 125.36, 124.83, 118.71, 118.61, 72.37, 70.23, 46.81, 27.08, 24.94, 23.44 ppm; $\text{C}_{14}\text{H}_{16}\text{N}_4\text{O}_2$ (MW 288.3), LCMS (EI) *m/z* 289 (M^+ + H).

Example 5. 2-((2*tf*,5*S*)-5-(2-((*tf*)-1-Hydroxyethyl)-1 *H*-imidazo[4,5-*i*]thieno[3,2-6]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-*c*]thieno[3,2-6]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile



In reactor A, a suspension of (*R*)-(+)-lactamide (2045 g, 22.96 mol, 3.0 eq.) in THF (10.1 L) was treated with Et₃O-BF₄ (4405 g, 22.96 mol, 3.0 eq.) at -10 to 0 °C. The resulting mixture in reactor A was then stirred at below 10 °C until a clear solution was formed. In reactor B, a mixture of 2-((2*R*,5*S*)-5-(6-aminothieno[3,2-
5 &]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothieno[3,2-*S*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile (2207 g, 7.65 mol) in ethanol (66.2 L) was heated to reflux to provide a clear solution. The solution in reactor A was then gradually added to the mixture in reactor B while maintaining the mixture in reactor B at reflux. The resulting reaction mixture in
10 reactor B was maintained at reflux until HPLC showed the reaction was complete. The reaction mixture in reactor B was then cooled to 10 - 30 °C before an aqueous solution of sodium bicarbonate (664 g, 7.4 L) was gradually added. During addition of aqueous sodium bicarbonate to adjust the pH to 8 - 9, the internal temperature was controlled at below 35 °C. The resulting mixture was then stirred at room temperature
15 for an additional 30 minutes before being filtered through Celite (1 kg). After the filtration, the Celite bed was rinsed with ethanol (5.0 L). The combined filtrate was concentrated under reduced pressure at below 60 °C and dichloromethane (17.7 L) was added to dissolve the residue. The resulting DCM solution was then treated with a 1N aqueous HCl solution (19.0 L) to adjust the pH of the aqueous phase to 1. The
20 mixture was then stirred for 30 minutes at room temperature. The two phases were separated and the DCM phase was discarded. The acidic aqueous phase was then washed with DCM (8.6 L). The two phases were separated and the DCM phase was discarded. The acidic aqueous phase was then treated with ethyl acetate (30.6 L) followed by solid sodium carbonate (540 g). The resulting mixture was stirred at
25 room temperature until the aqueous phase reached a pH of 9 - 10. w-Heptane (60.0 L) was then added to the mixture, and the resulting mixture was stirred at room temperature for at least 1 h. The solids were collected by filtration and washed with water (8.6 L) and w-heptane (8.6 L) before being dissolved in a mixture of methanol and DCM. The resulting solution of MeOH and DCM was then passed through a
30 silica gel bed and the silica gel bed was washed with a mixture of MeOH and DCM (1 to 9 by volume). The combined solution was concentrated under reduced pressure to afford a first crop of the desired product (1450 g), a mixture 2-((2*R*,5*S*)-5-(2-((*R*)-1-

hydroxyethyl)-1 *H*-imidazo[4,5-*i*]thieno[3,2-*/?*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*',5*S*')-5-(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-


i]thieno[3,2-*/?*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile, as white foam.

The two phases of the filtrate were then separated and the aqueous phase was
 5 extracted with DCM (2 X 25.0 L). The combined organic extracts were concentrated under reduced pressure and the residue was purified by silica gel column chromatography (SiO₂, gradient elution with 0.5% to 7.5% MeOH in DCM) to afford a second crop of the desired product (515 g; total 1965 g, 75%), also as a mixture 2-((2*R*,5*S*')-5-(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-*i*]thieno[3,2-*/?*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*',5*S*')-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*i*]thieno[3,2-*/?*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile, as
 10 white foam. The mixture of two diastereomers was inseparable by silica gel column chromatography, and was used in the subsequent reaction without further separation.

For **(2*R*,5*S*')-isomer**: ³H NMR (500 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s, 1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s, 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* = 12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* = 12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ 157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87, 62.95, 51.67, 30.39, 28.44, 24.09, 22.36
 20 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI) *m/e* 343 (M⁺ + H).


For **(2*S*,5*S*')-isomer**: ³H NMR (400 MHz, CD₃CN) δ 8.93 (s, 1H), 7.80 (d, *J* = 5.5 Hz, 1H), 7.63 (d, *J* = 5.5 Hz, 1H), 5.38 (s, 1H), 5.00 (dq, *J* = 9.4, 4.7 Hz, 1H), 4.43 (dd, *J* = 11.8, 9.3 Hz, 1H), 4.26 (dd, *J* = 8.6, 3.7 Hz, 1H), 4.16 (s, 1H), 4.03 - 3.89 (m, 1H), 3.17 (dd, *J* = 17.1, 8.6 Hz, 1H), 2.87 (dd, *J* = 17.2, 5.4 Hz, 1H), 2.57 (dtd, *J* = 14.8, 10.8, 4.2 Hz, 1H), 2.20 - 1.99 (m, 2H), 1.88-1.82 (m, 1H), 1.73 (d, *J* = 6.4 Hz, 3H) ppm; ¹³C NMR (101 MHz, CD₃CN) δ 158.04, 152.88, 142.12, 136.81, 136.46, 127.44, 127.14, 119.04, 116.97, 70.00, 64.83, 64.03, 54.55, 28.06, 25.73, 22.46, 21.94 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI) *m/e* 343 (M⁺ + H).

Example 6. 2-((2*R*,5*S*')-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*d*]thieno[3,2-*Z*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile monohydrate (Crude product)




A mixture of 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-
i]/thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-
(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-
5 pyran-2-yl)acetonitrile (800 g, 2.34 mol) in 2-propanol (IPA, 16 L) was treated with a
solution of 1.0 M potassium tert-butoxide in THF (234 mL, 0.234 mol, 0.10 eq.) at
room temperature. The resulting reaction mixture was stirred at room temperature for
20 h. When HPLC showed the base-promoted racemization reaction was complete,
the reaction mixture was treated with a 0.1 M aqueous HCl solution (approximately
10 2.4 L) to adjust the pH to 6 - 7. The mixture was then concentrated under reduced
pressure at below 50 °C to partially remove IPA (approximately 9.6 L) and the residue
was treated with water (10.5 L) at room temperature. The resulting mixture was then
stirred at room temperature for 2 h. The solids were collected by filtration, washed
with a mixture of IPA and water (2.4 L, 1 to 2 by volume), and dried under reduced
15 pressure at below 50 °C to constant weight to provide crude 2-((2*R*,5*S*)-5-(2-((*R*)-1-
hydroxyethyl)-1 *H*-imidazo[4,5-*i*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-
yl)acetonitrile monohydrate (657 g, 78%) as a white solid. ¹H NMR (500 MHz,
DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s,
1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s,
20 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* =
12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* =
12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ
157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87,
62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI)
25 *m/e* 343 (M⁺ + H); water content by KF: 5.15%.

**Example 7. 2-((2*tf*,5*S*)-5-(2-((*tf*)-1-Hydroxyethyl)-1 *H*-imidazo[4,5-*d*]thieno[3,2-
Z<]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile (Anhydrous Crystalline
Free Base)**




A solution of 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-
i]/thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile monohydrate (650
g, 1.80 mol) in methanol (7 L) was polish filtered before being concentrated at
5 atmospheric pressure to remove methanol (5 L). The resulting concentrated solution
was then treated with ethyl acetate (8.5 L) and the atmospheric distillation was
continued to remove solvents (5.3 L). *w*-Heptane (7.8 L) was then gradually added to
the residual mixture at 60 - 75 °C and the resulting mixture was gradually cooled
down to room temperature and stirred at room temperature for at least 1.5 h. The
10 solids were collected by filtration, washed with *w*-heptane (2.0 L), and dried in
vacuum at below 50 °C to constant weight to afford anhydrous crystalline 2-((2*R*,5*S*)-
5-(2-((*R*)-1-hydroxy ethyl)-1*H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-
2*H*-pyran-2-yl)acetonitrile (586 g, 95%) as a white solid. ¹H NMR (500 MHz,
DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s,
1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s,
15 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* =
12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* =
12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ
157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87,
20 62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI)
m/z 343 (M⁺ + H).

**Example 8. 2-((2*tf*,5*S*)-5-(2-((*tf*)-1-Hydroxyethyl)-1*H*-imidazo [4,5-*i*]thieno [3,2-
Z>]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile Monohydrate**



A suspension of anhydrous crystalline 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*i*]thieno[3,2-*h*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile (585 g, 1.71 mol) in water (5.9 L) was stirred at room temperature for 2 h. When XRPD confirmed that the conversion of the anhydrous crystalline form to the monohydrate crystalline form was complete, the solids were collected by filtration and washed with water (2.9 L). The wet cake was dried at 20 - 50 °C either in a vacuum oven or by pulling a vacuum from the filter until the water content analyzed by KF was $5 \pm 0.3\%$ to afford crystalline 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*i*]thieno[3,2-*h*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile monohydrate (602 g, 98%) as a white solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s, 1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s, 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* = 12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* = 12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ 157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87, 62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm; LCMS Calculated for: C₁₇H₁₈N₄O₂S: 342.4, Found: 343 (M⁺ + H); Water content: 5.15%.

Example 9. 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile (Anhydrous Crystalline Free Base)



A solution of amorphous ((2*R*,5*S*)-5-{2-[(1*R*)-1-hydroxyethyl]-1*H*-imidazo[4,5-*i*]thieno[3,2-*h*]pyridin-1-yl}tetrahydro-2*H*-pyran-2-yl)acetonitrile (84.5 g, 247 mmol) in methanol (MeOH, 800 mL) was polish filtered and transferred into a 5-neck 5 L round bottom flask equipped with a reflux condenser. Additional 200 mL of methanol was used to wash the filter funnel. To the resulted methanol solution, ethyl acetate (2000 mL) was added and the resulting clear solution was heated to 70


°C over 30 minutes. The mixture of methanol and ethyl acetate was then distilled at atmospheric pressure. An additional 2000 mL of ethyl acetate was added when the collected solvent mixture reached 1700 mL and the atmospheric distillation was continued. When the total collected solvent mixture reached 4000 mL (in about 8 h),
5 the remaining solution in the flask became cloudy and solids gradually formed. The atmospheric distillation was continued until the total collected solvent mixture reached 4250 mL, and the remaining solvent mixture, mainly ethyl acetate, in the flask was about 750 mL. w-Heptane (1150 mL) was then added slowly to the flask at approximately 70 °C (EtOAc to w-Heptane approximately at 1 to 1.5), and the
10 resulting mixture was gradually cooled to room temperature. The mixture was agitated at room temperature for an additional 2 - 3 h to initiate crystallization. The solids were collected by filtration, washed with w-heptane (2 x 300 mL), and dried at 50 °C in vacuum oven with N₂ sweeping to afford ((2*R*,5*S*)-5-{2-[(1*R*)-1-hydroxyethyl]-1*H*-imidazo[4,5-*i*]thieno[3,2-*h*]pyridin-1-yl}tetrahydro-2-*H*-pyran-2-yl)acetonitrile (78.1 g, 92.4%) as white anhydrous crystalline powder.
15

C₁₇H₁₈N₄O₂S (MW 342.42), LCMS (EI) *m/e* 343 (*M*⁺ + *H*); ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.97 (s, 1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.30 (t, *J* = 11.0 Hz, 1H), 4.17 (s, 1H), 4.02 (s, 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.86 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.76 - 2.56 (m, 1H), 2.19 (d, *J* = 10.6 Hz, 1H), 2.06 (d, *J* = 13.2 Hz, 1H), 1.72 (dd, *J* = 12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (100 MHz, DMSO-*d*₆) δ 157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87, 62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm.; water content by KF: 1.15% by weight. The DSC thermogram shown in Figure 1 revealed one endothermic event with an onset at
25 185.05 °C, which relates to the compound melting, and the peak at 189.71 °C. The endothermic event at about 260 °C associated with the decomposition of the compound. The TGA thermogram shown in Figure 2 revealed a weight loss of 1.25% up to 200 °C. Significant weight loss was observed at temperatures above 200 °C due to decomposition of the compound. The XRPD pattern was determined for the title
30 compound and is shown in Figure 3. A list of 2-theta peaks is provided in Table 2 below.

Table 2

2-Theta	d (Å)	Net Intensity	Gross Intensity	Relative Intensity (%)
6.608	13.3654	608	931	2.7
8.802	10.0379	8330	8970	37.6
9.699	9.1114	1560	2280	7.0
10.016	8.8244	2260	3010	10.2
10.574	8.3595	6020	6800	27.2
13.207	6.6983	2970	3870	13.4
14.509	6.1000	1490	2560	6.7
15.092	5.8659	3980	5180	18.0
16.329	5.4241	21900	23300	98.8
16.889	5.2454	1350	2870	6.1
17.554	5.0482	5510	7120	24.9
18.017	4.9194	4010	5680	18.1
19.081	4.6474	11000	12800	49.6
19.726	4.4971	3490	5320	15.8
21.174	4.1927	7880	9770	35.6
21.648	4.1018	2740	4640	12.4
22.466	3.9544	1080	2980	4.9
23.387	3.8007	3290	5160	14.9
24.037	3.6993	12100	13900	54.7
25.096	3.5456	626	2370	2.8
26.204	3.3980	5000	6720	22.6
26.420	3.3708	6510	8220	29.4
26.858	3.3169	1190	2880	5.4
27.681	3.2201	1320	2970	6.0
27.933	3.1916	1200	2830	5.4
29.318	3.0439	1060	2600	4.8
31.169	2.8672	1510	2870	6.8
31.300	2.8555	1560	2910	7.1
33.466	2.6755	607	1730	2.7
35.550	2.5233	826	1820	3.7
40.831	2.2083	501	1450	2.3
41.961	2.1514	966	1920	4.4
44.759	2.0232	618	1520	2.8
51.694	1.7669	229	1020	1.0

Example 10. 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile monohydrate



5

In a 2-L round bottom flask equipped with a reflux condenser, a thermocouple and a magnetic stirring bar, was charged amorphous 2-((2R,5S)-5-(2-((R)-1-

hydroxyethyl)-1 *H*-imidazo[4,5-*i*]thieno[3,2-*h*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile (93.0 g, 272 mmol) and isopropyl alcohol (IPA, 1116 mL) at room temperature. The resulting suspension was heated to 50 - 55 °C to obtain a clear solution. The solution was polish filtered through a filter funnel at 50 °C. An additional amount of isopropyl alcohol (IPA, 47 mL) was used to wash the filter funnel. To the combined solution was added water (930 mL) over 30 minutes while maintaining the internal temperature at 45 - 50 °C. White precipitate gradually appeared upon the addition of the water. The IPA was removed by rotary evaporation under vacuum with a water bath at 50 °C. During the distillation, an additional amount of water (1209 mL) was added to keep the total volume of the mixture at about 1500 mL. After the vacuum distillation, the resulting mixture was gradually cooled to room temperature and agitated at room temperature for 1 - 2 h. The solids were collected by filtration, washed with water (186 mL), and dried on a filter funnel with a housing vacuum at room temperature for 16 h to afford 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-*i*]thieno[3,2-*h*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile monohydrate (91.6 g, 94%) as white crystalline powder. ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s, 1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s, 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* = 12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* = 12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ 157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87, 62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm; C₁₇H₁₈N₄O₂S (MW 342.42), LCMS (EI) *m/z* 343 (M⁺ + H). Water content by KF: 5.15% by weight. The melting/decomposition range was determined by DSC and is shown in Figure 4. The DSC thermogram revealed one dehydration endothermic event with an onset at 69.37 °C and the peak at 106.41 °C, followed by the melting with an onset at 131.01 °C and a peak at 139.99 °C. The TGA thermogram (Figure 5) showed a weight loss of 4.9% up to 150 °C, which corresponds to the water content of the mono-hydrate. After 200 °C the compound starts to decompose. The XRPD pattern was determined for the title compound and is shown in Figure 6. A list of 2-theta peaks is provided in Table 3 below.

Table 3

2-Theta	d (Å)	Net Intensity	Gross Intensity	Relative Intensity (%)
8.477	10.4224	28200	28700	91.9
8.678	10.1819	14500	15000	47.3
11.368	7.7776	2600	3320	8.5
13.150	6.7274	564	1320	1.8
13.568	6.5211	162	922	0.5
14.592	6.0656	23400	24300	76.1
15.064	5.8765	21200	22100	69.0
15.789	5.6082	9180	10100	29.9
16.917	5.2369	6130	7140	19.9
17.427	5.0846	3000	4020	9.8
18.599	4.7668	7670	8720	25.0
19.388	4.5747	20300	21400	66.2
20.160	4.4011	4640	5700	15.1
21.091	4.2089	6690	7710	21.8
21.463	4.1368	839	1840	2.7
22.460	3.9554	3070	4150	10.0
22.871	3.8853	1370	2510	4.5
23.366	3.8040	6710	7900	21.8
23.503	3.7822	8550	9750	27.8
24.518	3.6279	706	2000	2.3
25.065	3.5499	29900	31200	97.3
25.674	3.4670	3830	5200	12.4
25.878	3.4401	11100	12500	36.2
26.306	3.3857	12500	13900	40.7
26.547	3.3549	18100	19500	58.9
27.574	3.2323	6810	8210	22.2
28.299	3.1511	717	2100	2.3
28.625	3.1160	3130	4490	10.2
29.200	3.0559	1670	3000	5.4
29.828	2.9930	1100	2370	3.6
30.451	2.9332	3810	5010	12.4
30.715	2.9085	5430	6600	17.7
31.001	2.8823	1830	2960	6.0
31.335	2.8524	1820	2910	5.9
32.496	2.7531	420	1420	1.4
33.388	2.6815	4050	5030	13.2
34.026	2.6327	1220	2180	4.0
34.291	2.6130	861	1800	2.8
35.285	2.5416	724	1630	2.4
35.729	2.5111	1990	2900	6.5
36.639	2.4507	684	1590	2.2
36.931	2.4320	744	1640	2.4
37.470	2.3983	2760	3630	9.0
37.686	2.3850	1160	2020	3.8

39.059	2.3043	567	1430	1.8
39.713	2.2678	767	1680	2.5
40.791	2.2103	908	1870	3.0
41.862	2.1563	2100	3070	6.8
42.406	2.1298	511	1470	1.7
43.684	2.0704	531	1430	1.7
44.647	2.0280	1760	2680	5.7
45.207	2.0042	462	1430	1.5
45.857	1.9773	428	1430	1.4
47.419	1.9157	1510	2540	4.9
47.976	1.8947	1750	2780	5.7
48.744	1.8667	896	1890	2.9
49.480	1.8406	470	1420	1.5
51.479	1.7737	811	1630	2.6
52.307	1.7476	452	1270	1.5
53.685	1.7060	321	1130	1.0

Example 11. Dynamic Vapor Sorption


A Dynamic Vapor Sorption (DVS) study was conducted on the compound of Example 9 to monitor the hygroscopicity, hydrate formation and dehydration, as well as solid form transformation. The moisture uptake profile was completed in four cycles in 10% relative humidity (RH) increments with the first adsorption from 25% to 95% RH, followed by desorption in 10% increments from 95% to 5% RH. The equilibration criteria were as follows: 0.0050 wt% in 5 minutes with a maximum equilibration time of 180 minutes. All adsorption and desorption were performed at room temperature (25 °C). No pre-drying step was applied for the sample. Water absorption of the anhydrous form was slow below 85% RH and fast above 85% RH. Changes in the XRPD patterns before and after DVS suggest solid form change. After DVS cycle 4, the anhydrous form was fully converted to the hydrate form. Under ambient storage conditions, the anhydrous form will slowly absorb water and convert to hydrate form. The dehydration process from the hydrate to the anhydrous form is very slow, even at 5% RH. The hydrate form is physically stable in a wide range of humidity conditions. Vapor sorption/desorption isotherms are shown in Figure 7 and Figure 8. An XRPD pattern overlay before and after DVS is shown in Figure 9.

Example 12. Humidity Chamber Study A


The compound of Example 9 was placed in a humidity chamber set at 40 °C and 75% relative humidity for 5 days. The resulting sample was analyzed by XRPD, TGA, and DSC as shown in Figures 10 to 12. Based on XRPD data, the anhydrous crystalline free base was converted to the crystalline hydrate form. The water content by KF was 4.6% and the weight loss observed by TGA was 5.01%.

Example 13. Humidity Chamber Study B

The compound of Example 9 was placed in a humidity chamber set at 30 °C and 90% relative humidity (RH) for 7 days. The resulting sample was analyzed by XRPD, TGA, and DSC as shown in Figures 13 to 15. Based on XRPD, the anhydrous crystalline free base was converted to the crystalline hydrate form. The XRPD patterns of the 30 °C / 90% RH sample corresponded to the hydrate form obtained from an aqueous suspension of the anhydrous crystalline free base drug substance. The water content by KF was 4.72% and the weight loss observed by TGA was 5.05%.

Example 14. (R)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole

Step 1. (R)-2,2,5-TrimethyloxazoUdin-4-one




To a solution of (2R)-2-hydroxypropanamide (30 g, 337 mmol) in acetone was added 2,2-dimethoxypropane (51.8 mL, 421 mmol, 1.25 equiv) and boron trifluoride diethyl etherate (BF₃-Et₂O, 92.13 mL, 16.8 mmol, 0.05 equiv) at room temperature. The resulting reaction mixture was then stirred at room temperature overnight. When TLC indicated the reaction was complete, the solvent was removed under reduced pressure at below 35 °C and the residue was added to methyl *n*-butyl ether (MTBE, 350 mL). The solution was treated with saturated aqueous sodium bicarbonate

(NaHCCCh) solution (15 mL), solid sodium bicarbonate (NaHCO₃, 4 g), and solid sodium sulfate (Na₂SO₄, 30 g) and the resulting mixture was stirred at room temperature for 30 minutes. The solids were filtered off and washed with methyl *t*-butyl ether (MTBE, 50 mL). The two phases of the combined filtrates were separated
 5 and the organic phase was washed with water (30 mL) and dried with sodium sulfate (Na₂SO₄, 20 g). After removal of the drying agent, the filtrate was concentrated under reduced pressure and the solvent was switched into *n*-hexane to provide desired product. The solids were collected by filtration, washed with *n*-hexane and dried to give the desired product, (*R*)-2,2,5-trimethyloxazolidin-4-one, (42 g, 97%) as white
 10 crystalline solids. ¹H NMR (400 MHz, CD₂Cl₂) δ 7.66 (s, 1H), 4.37 (q, *J* = 6.7 Hz, 1H), 1.47 (s, 3H), 1.44 (s, 3H), 1.36 (d, *J* = 6.7 Hz, 3H) ppm; ¹³C NMR (101 MHz, CD₂Cl₂) δ 175.58 (s), 90.70 (s), 73.76 (s), 29.97 (s), 28.52 (s), 18.60 (s) ppm.

Step 2. (R)-4-Methoxy-2,2,5-trimethyl-2,5-dihydrooxazole

15 To a solution of (*R*)-2,2,5-trimethyloxazolidin-4-one (20 g, 155 mmol) in dichloromethane (DCM, 200 mL) was added trimethoxonium tetrafluoride (Me₃O⁺BF₄⁻, 24.1 g, 155 mmol, 1.0 equiv) at room temperature. The resulting reaction mixture was stirred at room temperature for 6 hours. When ³¹P-NMR indicated the reaction was complete, the reaction mixture was added to
 20 dichloromethane (DCM, 100 mL) and cooled to 0 - 5 °C. The resulting solution was then treated with a 2M aqueous sodium carbonate solution (Na₂CO₃, 230 mL) while maintaining the internal temperature at 0 - 5 °C. The two phases were separated and the organic phase was washed with water (2 x 100 mL), dried over sodium sulfate (Na₂SO₄, 50 g), and concentrated under reduced pressure to give the crude desired
 25 product, (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole, (19 g, 87%) as yellow oil, which can be directly used for the subsequent reaction without further purification. The pure desired product (12 g; bp 50 °C at 50 mba) was obtained by vacuum distillation as colorless oil. ¹H NMR (400 MHz, CD₂Cl₂) δ 4.58 (q, *J* = 6.6 Hz, 1H), 3.80 (s, 3H), 1.41 (s, 3H), 1.35 (s, 3H), 1.29 (d, *J* = 6.6 Hz, 3H) ppm; ¹³C
 30 NMR (101 MHz, CD₂Cl₂) δ 169.80 (s), 103.90 (s), 75.08 (s), 56.46 (s), 30.50 (s), 28.69 (s), 20.07 (s) ppm.

Example 15. Alternative preparation of 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*c*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(2-((*S*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile



5

To a flask was added (*R*)-2,2,5-trimethyloxazolidin-4-one (6.7 g, 52 mmol), trimethyloxonium tetrafluoroborate ($\text{Me}_3\text{O}^+\text{BF}_4^-$, 7.7 g, 52 mmol), and anhydrous dichloromethane (DCM, 50 mL) at room temperature. The resulting mixture was stirred at room temperature for 2 hours. When ^1H NMR revealed that the formation of (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole was complete, the reaction mixture was charged with a mixture of 2-((2*R*,5*S*)-5-(6-aminothieno[3,2-*Z'*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothieno[3,2-*Z'*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile (Example 4, 5.0 g, 17.3 mmol) at room temperature. The solvent was then switched into methanol (80 mL) by addition of methanol to the reaction mixture and distillation to remove dichloromethane. The resulting reaction mixture was then heated to 65°C for 1.5 hours. When HPLC showed the reaction was complete, the solvent was removed under reduced pressure and the residue was treated with dichloromethane (80 mL). The resulting mixture was then treated with saturated aqueous sodium bicarbonate solution (NaHCO_3 , 10 mL) followed by solid NaHCO_3 (11.6 g, 139 mmol) before being stirred at room temperature for 30 min. The mixture was dried over Na_2SO_4 and the liquid was filtered. The solids were removed by filtration and the two phases of the filtrate were separated. The organic phase was washed with water (2 x 25 mL), dried over Na_2SO_4 , and concentrated. The residue was then dissolved in 0.1 M aqueous hydrochloric acid (HCl) and the resulting acidic aqueous solution was washed with dichloromethane (2 x 20 mL). The acidic aqueous solution was then neutralized with NaHCO_3 to precipitate the crude desired product. The solids were collected by filtration, washed with water, and dried to give the desired product (5.0 g, 84% yield) as a mixture of 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-

20

25


i)]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile.

The mixture of two diastereomers is inseparable by silica gel column chromatography and was used directly in the subsequent reaction (see, *e.g.*, Example 6) without further separation.


For **(2*R*,5*S*)-isomer**: ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s, 1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s, 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* = 12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* = 12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ 157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87, 62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI) *m/e* 343 (M⁺ + H).

For **(2*S*,5*S*)-isomer**: ¹H NMR (400 MHz, CD₃CN) δ 8.93 (s, 1H), 7.80 (d, *J* = 5.5 Hz, 1H), 7.63 (d, *J* = 5.5 Hz, 1H), 5.38 (s, 1H), 5.00 (dq, *J* = 9.4, 4.7 Hz, 1H), 4.43 (dd, *J* = 11.8, 9.3 Hz, 1H), 4.26 (dd, *J* = 8.6, 3.7 Hz, 1H), 4.16 (s, 1H), 4.03 - 3.89 (m, 1H), 3.17 (dd, *J* = 17.1, 8.6 Hz, 1H), 2.87 (dd, *J* = 17.2, 5.4 Hz, 1H), 2.57 (dtd, *J* = 14.8, 10.8, 4.2 Hz, 1H), 2.20 - 1.99 (m, 2H), 1.88-1.82 (m, 1H), 1.73 (d, *J* = 6.4 Hz, 3H) ppm; ¹³C NMR (101 MHz, CD₃CN) δ 158.04, 152.88, 142.12, 136.81, 136.46, 127.44, 127.14, 119.04, 116.97, 70.00, 64.83, 64.03, 54.55, 28.06, 25.73, 22.46, 21.94 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI) *m/e* 343 (M⁺ + H).

Example 16. Alternative preparation of 2-((2*R*,5*S*)-5-(2-((*R*)-1-Hydroxyethyl)-1 *H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(2-((*S*)-1-hydroxyethyl)-1 *H*-imidazo[4,5-*d*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2 *H*-pyran-2-yl)acetonitrile




Step 1. (R)-l-Amino-l-oxopropan-2-yl acetate



To a 1-L 3-necked round bottom flask equipped with a reflux condenser, a nitrogen inlet, a thermal couple and a magnetic stirring bar under nitrogen was charged (*R*)-2-hydroxypropanamide (49.95 g, 543.8 mmol) and tetrahydrofuran (THF, 300 mL) at room temperature. The resulting yellow solution was cooled to 0 - 5 °C by an ice-water bath before acetyl chloride (40.60 mL, 571.0 mmol, 1.05 equiv) was charged while maintaining the internal temperature at below 20 °C. After the internal temperature was dropped back to below 5 °C, 4-methylmorpholine (59.2 mL, 538.0 mmol, 0.99 equiv) was charged while maintaining the internal temperature to below 10 °C. The resulting reaction mixture was then stirred at room temperature for 4 hours before the mixture was filtered to remove the solids. The solids were washed with THF (60 mL) and the filtrate was concentrated under reduced pressure to provide the crude desired product, (*R*)-1-amino-1-oxopropan-2-yl acetate, (75.0 g, 105.2%) as a white to off-white solid, which was used directly in next reaction without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.42 (s, 1 H), 7.14 (s, 1 H), 4.84 (q, *J* = 7.6 Hz, 1 H), 2.03 (s, 3 H), 1.29 (d, *J* = 7.6 Hz, 3 H) ppm.


Step 2. (R)-1-Cyanoethyl acetate



To a 3-L 3-necked round bottom flask equipped with a reflux condenser, a nitrogen inlet, a thermal couple and a magnetic stirring bar under nitrogen was charged crude (*R*)-1-amino-1-oxopropan-2-yl acetate (70.0 g, 533.8 mmol) and *N,N*-dimethylformamide (DMF, 280 mL) at room temperature. The resulting light yellow solution was then treated with cyanuric chloride (44.0 g, 240.0 mmol) in 2-methoxy-2-methylpropane (TBME, 600 mL) at room temperature. The reaction mixture was stirred at room temperature for 16 hours. The white solids were filtered and washed with TBME (800 mL). The combined organic filtrates were washed sequentially with

saturated aqueous sodium bicarbonate solution (NaHCO₃, 200 mL), water (4 × 200 mL), and brine (100 mL). The resultant organic phase was then dried with anhydrous MgSO₄ (15 g) and concentrated under reduced pressure to provide the crude desired product, (*R*)-1-cyanoethyl acetate, (42.03 g, 69.6%) as a colorless to light yellow oil, which was used directly in the subsequent reaction without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 5.46 (q, *J* = 7.6 Hz, 1 H), 2.09 (s, 3 H), 1.53 (d, *J* = 7.6 Hz, 3 H) ppm.

Step 3. (R)-ethyl 2-hydroxypropanimidate



To a 100-mL round bottom flask equipped with a magnetic stirring bar was charged crude (*R*)-1-cyanoethyl acetate (1.021 g, 8.665 mmol) and ethanol (EtOH, 1.518 mL, 26.0 mmol) at room temperature. The resulting solution was then cooled to -40 °C before being bubbled in hydrogen chloride (HCl) gas for 45 seconds. The reaction mixture was then gradually warmed to room temperature. After 16 hours, the reaction mixture was cooled to 0 °C before tetrahydrofuran (THF, 2.0 mL, 25 mmol) and 4-methylmorpholine (0.865 mL, 7.87 mmol) were sequentially charged. The solids were removed by filtration and the crude desired product, (*R*)-ethyl 2-hydroxypropanimidate, in the filtrate was used directly in the subsequent reaction without further purification. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.14 (s, 1 H), 4.52 - 4.40 (m, 3 H), 1.36 - 1.31 (m, 6 H) ppm.

Step 4. 2-((2R, 5S)-5-(2-((R)-1-Hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile and 2-((2S,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile

To a 3-necked 100-mL round bottom flask equipped with a nitrogen inlet, a thermocouple and a magnetic stirring bar under nitrogen was charged a mixture of 2-((2*R*,5*S*)-5-(6-aminothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(6-aminothieno[3,2-*b*]pyridin-7-ylamino)tetrahydro-2*H*-pyran-2-yl)acetonitrile (Example 4, 0.30 g, 1.04 mmol) in ethanol (EtOH, 4.0 mL)

at room temperature. The mixture was then heated to 80 °C to provide a clear solution before a solution of the crude (*R*)-ethyl 2-hydroxypropanimidate in THF generated in previous step was added to the ethanol solution at 80 °C. When the reaction was completed after 1 hour as indicated by HPLC, the reaction mixture was cooled to room temperature. The cooled reaction mixture was filtered and the filtrate was concentrated under reduced pressure to afford the crude desired product (0.3 g, 84% yield) as a mixture of 2-((2*R*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*i*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile and 2-((2*S*,5*S*)-5-(2-((*R*)-1-hydroxyethyl)-1*H*-imidazo[4,5-*i*]thieno[3,2-*b*]pyridin-1-yl)tetrahydro-2*H*-pyran-2-yl)acetonitrile, which is inseparable by silica gel column chromatography and was used directly in the subsequent reaction without further separation (see, *e.g.*, Example 6).

For (**2*R*,5*S***)**Isomer**: ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.00 (s, 1H), 8.03 (d, *J* = 5.5 Hz, 1H), 7.69 (d, *J* = 5.5 Hz, 1H), 5.96 (s, 1H), 5.22 (s, 1H), 5.11 (s, 1H), 4.31 (t, *J* = 11.0 Hz, 1H), 4.22 - 4.08 (m, 1H), 4.02 (s, 1H), 2.96 (dd, *J* = 17.0, 4.2 Hz, 1H), 2.84 (dd, *J* = 17.0, 6.6 Hz, 1H), 2.65 (qd, *J* = 12.7, 4.3 Hz, 1H), 2.19 (d, *J* = 10.0 Hz, 1H), 2.06 (d, *J* = 13.4 Hz, 1H), 1.72 (dd, *J* = 12.6, 3.7 Hz, 1H), 1.66 (d, *J* = 6.5 Hz, 3H) ppm; ¹³C NMR (101 MHz, DMSO-*d*₆) δ 157.95, 152.11, 141.57, 136.18, 135.24, 127.98, 126.79, 119.08, 116.12, 72.42, 68.87, 62.95, 51.67, 30.39, 28.44, 24.09, 22.36 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI) *m/e* 343 (M⁺ + H).

For (**2*S*,5*S***)-**isomer**: ¹H NMR (400 MHz, CD₃CN) δ 8.93 (s, 1H), 7.80 (d, *J* = 5.5 Hz, 1H), 7.63 (d, *J* = 5.5 Hz, 1H), 5.38 (s, 1H), 5.00 (dq, *J* = 9.4, 4.7 Hz, 1H), 4.43 (dd, *J* = 11.8, 9.3 Hz, 1H), 4.26 (dd, *J* = 8.6, 3.7 Hz, 1H), 4.16 (s, 1H), 4.03 - 3.89 (m, 1H), 3.17 (dd, *J* = 17.1, 8.6 Hz, 1H), 2.87 (dd, *J* = 17.2, 5.4 Hz, 1H), 2.57 (dtd, *J* = 14.8, 10.8, 4.2 Hz, 1H), 2.20 - 1.99 (m, 2H), 1.88-1.82 (m, 1H), 1.73 (d, *J* = 6.4 Hz, 3H) ppm; ¹³C NMR (101 MHz, CD₃CN) δ 158.04, 152.88, 142.12, 136.81, 136.46, 127.44, 127.14, 119.04, 116.97, 70.00, 64.83, 64.03, 54.55, 28.06, 25.73, 22.46, 21.94 ppm; C₁₇H₁₈N₄O₂S (MW 342.4), LCMS (EI) *m/e* 343 (M⁺ + H).

Example A: *In vitro* JAK Kinase Assay

Compounds herein were tested for inhibitory activity of JAK targets according to the following *in vitro* assay described in Park *et al*, *Analytical Biochemistry* **1999**,

269, 94-104. The catalytic domains of human JAK1 (a.a. 837-1142), JAK2 (a.a. 828-1132) and JAK3 (a.a. 781-1124) were expressed using baculovirus in insect cells and purified. The catalytic activity of JAK1, JAK2 or JAK3 was assayed by measuring the phosphorylation of a biotinylated peptide. The phosphorylated peptide was detected
5 by homogenous time resolved fluorescence (HTRF). IC₅₀s of compounds were measured for each kinase in the 40 μ L reactions that contain the enzyme, ATP and 500 nM peptide in 50 mM Tris (pH 7.8) buffer with 100 mM NaCl, 5 mM DTT, and 0.1 mg/mL (0.01%) BSA. For the 1 mM IC₅₀ measurements, ATP concentration in the reactions was 1 mM. Reactions were carried out at room temperature for 1 hour
10 and then stopped with 20 μ L 45 mM EDTA, 300 nM SA-APC, 6 nM Eu-Py20 in assay buffer (Perkin Elmer, Boston, MA). Binding to the Europium labeled antibody took place for 40 minutes and HTRF signal was measured on a PHERA star plate reader (BMG, Cary, NC). The compound of Formula 1a was tested in the Example A assay at 1 mM ATP and found to have an IC₅₀ at JAK1 of \leq 100 nM and an IC₅₀ at
15 JAK2 of \leq 1000 nM with a JAK2/JAK1 ratio $>$ 10.

Example B: Cellular Assays

Cancer cell lines dependent on cytokines and hence JAK/STAT signal transduction, for growth, can be plated at 6000 cells per well (96 well plate format) in
20 RPMI 1640, 10% FBS, and 1 nG/nL of appropriate cytokine. Compounds can be added to the cells in DMSO/media (final concentration 0.2% DMSO) and incubated for 72 hours at 37 $^{\circ}$ C, 5% CO₂. The effect of compound on cell viability is assessed using the CellTiter-Glo Luminescent Cell Viability Assay (Promega) followed by TopCount (Perkin Elmer, Boston, MA) quantitation. Potential off-target effects of
25 compounds are measured in parallel using a non-JAK driven cell line with the same assay readout. All experiments are typically performed in duplicate.

The above cell lines can also be used to examine the effects of compounds on phosphorylation of JAK kinases or potential downstream substrates such as STAT proteins, Akt, Shp2, or Erk. These experiments can be performed following an
30 overnight cytokine starvation, followed by a brief preincubation with compound (2 hours or less) and cytokine stimulation of approximately 1 hour or less. Proteins are then extracted from cells and analyzed by techniques familiar to those schooled in the

art including Western blotting or ELISAs using antibodies that can differentiate between phosphorylated and total protein. These experiments can utilize normal or cancer cells to investigate the activity of compounds on tumor cell survival biology or on mediators of inflammatory disease. For example, with regards to the latter, cytokines such as IL-6, IL-12, IL-23, or IFN can be used to stimulate JAK activation resulting in phosphorylation of STAT protein(s) and potentially in transcriptional profiles (assessed by array or qPCR technology) or production and/or secretion of proteins, such as IL-17. The ability of compounds to inhibit these cytokine mediated effects can be measured using techniques common to those schooled in the art. To assess compound effects on JAK2, primary cells or cell lines can be stimulated with JAK2-dependent growth factors such as GM-CSF or Tpo, proteins extracted from cells and analyzed by techniques familiar to those schooled in the art including Western blotting or ELISAs using antibodies that can differentiate between phosphorylated and total protein.

Compounds herein can also be tested in cellular models designed to evaluate their potency and activity against mutant JAKs, for example, the JAK2V617F mutation found in myeloid proliferative disorders. These experiments often utilize cytokine dependent cells of hematological lineage (*e.g.* BaF/3) into which the wild-type or mutant JAK kinases are ectopically expressed (James, C, *et al. Nature* 434:1 144-1 148; Staerk, J., *et al. JBC* 280:41893-41899). Endpoints include the effects of compounds on cell survival, proliferation, and phosphorylated JAK, STAT, Akt, or Erk proteins.

Certain compounds herein can be evaluated for their activity inhibiting T-cell proliferation. Such as assay can be considered a second cytokine (*i.e.* JAK) driven proliferation assay and also a simplistic assay of immune suppression or inhibition of immune activation. The following is a brief outline of how such experiments can be performed. Peripheral blood mononuclear cells (PBMCs) are prepared from human whole blood samples using Ficoll Hypaque separation method and T-cells (fraction 2000) can be obtained from PBMCs by elutriation. Freshly isolated human T-cells can be maintained in culture medium (RPMI 1640 supplemented with 10% fetal bovine serum, 100 U/ml penicillin, 100 µg/ml streptomycin) at a density of 2×10^6 cells/ml at 37 °C for up to 2 days. For IL-2 stimulated cell proliferation analysis, T-cells are first

treated with Phytohemagglutinin (PHA) at a final concentration of 10 µg/mL for 72 hours. After washing once with PBS, 6000 cells/well are plated in 96-well plates and treated with compounds at different concentrations in the culture medium in the presence of 100 U/mL human IL-2 (ProSpec-Tany TechnoGene; Rehovot, Israel).

5 The plates are incubated at 37 °C for 72h and the proliferation index is assessed using CellTiter-Glo Luminescent reagents following the manufactory suggested protocol (Promega; Madison, WI).

Example C: *In vivo* anti-tumor efficacy

10 Compounds herein can be evaluated in human tumor xenograft models in immune compromised mice. For example, a tumorigenic variant of the TNA-6 plasmacytoma cell line can be used to inoculate SCID mice subcutaneously (Burger, R., *et al. Hematol J.* 2:42-53, 2001). Tumor bearing animals can then be randomized into drug or vehicle treatment groups and different doses of compounds can be
15 administered by any number of the usual routes including oral, i.p., or continuous infusion using implantable pumps. Tumor growth is followed over time using calipers. Further, tumor samples can be harvested at any time after the initiation of treatment for analysis as described above (Example B) to evaluate compound effects on JAK activity and downstream signaling pathways. In addition, selectivity of the
20 compound(s) can be assessed using xenograft tumor models that are driven by other known kinases (*e.g.* Bcr-Abl) such as the K562 tumor model.

Example D: Murine Skin Contact Delayed Hypersensitivity Response Test

Compounds herein can also be tested for their efficacies (of inhibiting JAK
25 targets) in the T-cell driven murine delayed hypersensitivity test model. The murine skin contact delayed-type hypersensitivity (DTH) response is considered to be a valid model of clinical contact dermatitis, and other T-lymphocyte mediated immune disorders of the skin, such as psoriasis (*Immunol Today.* 1998 Jan;19(1):37-44). Murine DTH shares multiple characteristics with psoriasis, including the immune
30 infiltrate, the accompanying increase in inflammatory cytokines, and keratinocyte hyperproliferation. Furthermore, many classes of agents that are efficacious in

treating psoriasis in the clinic are also effective inhibitors of the DTH response in mice (Agents Actions. 1993 Jan;38(1-2):1 16-21).

On Day 0 and 1, Balb/c mice are sensitized with a topical application, to their shaved abdomen with the antigen 2,4,dinitro-fluorobenzene (DNFB). On day 5, ears are measured for thickness using an engineer's micrometer. This measurement is recorded and used as a baseline. Both of the animals' ears are then challenged by a topical application of DNFB in a total of 20 μ L (10 μ L on the internal pinna and 10 μ L on the external pinna) at a concentration of 0.2%. Twenty-four to seventy-two hours after the challenge, ears are measured again. Treatment with the test compounds is given throughout the sensitization and challenge phases (day -1 to day 7) or prior to and throughout the challenge phase (usually afternoon of day 4 to day 7). Treatment of the test compounds (in different concentration) is administered either systemically or topically (topical application of the treatment to the ears). Efficacies of the test compounds are indicated by a reduction in ear swelling comparing to the situation without the treatment. Compounds causing a reduction of 20% or more were considered efficacious. In some experiments, the mice are challenged but not sensitized (negative control).

The inhibitive effect (inhibiting activation of the JAK-STAT pathways) of the test compounds can be confirmed by immunohistochemical analysis. Activation of the JAK-STAT pathway(s) results in the formation and translocation of functional transcription factors. Further, the influx of immune cells and the increased proliferation of keratinocytes should also provide unique expression profile changes in the ear that can be investigated and quantified. Formalin fixed and paraffin embedded ear sections (harvested after the challenge phase in the DTH model) are subjected to immunohistochemical analysis using an antibody that specifically interacts with phosphorylated STAT3 (clone 58E12, Cell Signaling Technologies). The mouse ears are treated with test compounds, vehicle, or dexamethasone (a clinically efficacious treatment for psoriasis), or without any treatment, in the DTH model for comparisons. Test compounds and the dexamethasone can produce similar transcriptional changes both qualitatively and quantitatively, and both the test compounds and dexamethasone can reduce the number of infiltrating cells. Both systemically and topical administration of the test compounds can produce inhibitive

effects, *i.e.*, reduction in the number of infiltrating cells and inhibition of the transcriptional changes.

Example E: *In vivo* anti-inflammatory activity

5 Compounds herein can be evaluated in rodent or non-rodent models designed to replicate a single or complex inflammation response. For instance, rodent models of arthritis can be used to evaluate the therapeutic potential of compounds dosed preventatively or therapeutically. These models include but are not limited to mouse or rat collagen-induced arthritis, rat adjuvant-induced arthritis, and collagen antibody-
10 induced arthritis. Autoimmune diseases including, but not limited to, multiple sclerosis, type I-diabetes mellitus, uveoretinitis, thyroiditis, myasthenia gravis, immunoglobulin nephropathies, myocarditis, airway sensitization (asthma), lupus, or colitis may also be used to evaluate the therapeutic potential of compounds herein. These models are well established in the research community and are familiar to those
15 schooled in the art (Current Protocols in Immunology, Vol 3., Coligan, J.E. *et al*, Wiley Press.; *Methods in Molecular Biology*: Vol. 225, Inflammation Protocols., Winyard, P.G. and Willoughby, D.A., Humana Press, 2003.).

**Example F: Animal Models for the Treatment of Dry Eye, Uveitis, and
20 Conjunctivitis**

 Agents may be evaluated in one or more preclinical models of dry eye known to those schooled in the art including, but not limited to, the rabbit concanavalin A (ConA) lacrimal gland model, the scopolamine mouse model (subcutaneous or
25 transdermal), the Botulinum mouse lacrimal gland model, or any of a number of spontaneous rodent auto-immune models that result in ocular gland dysfunction (e.g. NOD-SCID, MRL/lpr, or NZB/NZW) (Barabino et al, Experimental Eye Research 2004, 79, 613-621 and Schrader et al, Developmental Ophthalmology, Karger 2008, 41, 298-312, each of which is incorporated herein by reference in its entirety). Endpoints in these models may include histopathology of the ocular glands and eye
30 (cornea, etc.) and possibly the classic Schirmer test or modified versions thereof (Barabino et al.) which measure tear production. Activity may be assessed by dosing

via multiple routes of administration (e.g. systemic or topical) which may begin prior to or after measurable disease exists.

Agents may be evaluated in one or more preclinical models of uveitis known to those schooled in the art. These include, but are not limited to, models of experimental autoimmune uveitis (EAU) and endotoxin induced uveitis (EIU). EAU experiments may be performed in the rabbit, rat, or mouse and may involve passive or activate immunization. For instance, any of a number of retinal antigens may be used to sensitize animals to a relevant immunogen after which animals may be challenged ocularly with the same antigen. The EIU model is more acute and involves local or systemic administration of lipopolysaccharide at sublethal doses. Endpoints for both the EIU and EAU models may include fundoscopic exam, histopathology amongst others. These models are reviewed by Smith et al. (Immunology and Cell Biology 1998, 76, 497-512, which is incorporated herein by reference in its entirety). Activity is assessed by dosing via multiple routes of administration (e.g. systemic or topical) which may begin prior to or after measurable disease exists. Some models listed above may also develop scleritis/episcleritis, choroiditis, cyclitis, or iritis and are therefore useful in investigating the potential activity of compounds for the therapeutic treatment of these diseases.

Agents may also be evaluated in one or more preclinical models of conjunctivitis known those schooled in the art. These include, but are not limited to, rodent models utilizing guinea-pig, rat, or mouse. The guinea-pig models include those utilizing active or passive immunization and/or immune challenge protocols with antigens such as ovalbumin or ragweed (reviewed in Groneberg, D.A., et al, Allergy 2003, 58, 1101-1113, which is incorporated herein by reference in its entirety). Rat and mouse models are similar in general design to those in the guinea-pig (also reviewed by Groneberg). Activity may be assessed by dosing via multiple routes of administration (e.g. systemic or topical) which may begin prior to or after measurable disease exists. Endpoints for such studies may include, for example, histological, immunological, biochemical, or molecular analysis of ocular tissues such as the conjunctiva.

Example G: *In vivo* protection of bone

Compounds may be evaluated in various preclinical models of osteopenia, osteoporosis, or bone resorption known to those schooled in the art. For example, ovariectomized rodents may be used to evaluate the ability of compounds to affect signs and markers of bone remodeling and/or density (W.S.S. Jee and W. Yao, J Musculoskel. Nueron. Interact., 2001, 1(3), 193-207, which is incorporated herein by reference in its entirety). Alternatively, bone density and architecture may be evaluated in control or compound treated rodents in models of therapy (e.g. glucocorticoid) induced osteopenia (Yao, et al. Arthritis and Rheumatism, 2008, 58(6), 3485-3497; and *id.* 58(11), 1674-1686, both of which are incorporated herein by reference in its entirety). In addition, the effects of compounds on bone resorption and density may be evaluable in the rodent models of arthritis discussed above (Example E). Endpoints for all these models may vary but often include histological and radiological assessments as well as immunohisotology and appropriate biochemical markers of bone remodeling.

Example H: S100A9 Transgenic Mouse Model


It was previously shown that *S100A9* transgenic mice display bone marrow accumulation of MDSC accompanied by development of progressive multilineage cytopenias and cytological dysplasia similar to MDS. Further, early forced maturation of MDSC by either *a/-trans-retinoic acid* treatment or active immunoreceptor tyrosine-based activation motif-bearing (ITAM-bearing) adapter protein (DAP12) interruption of CD33 signaling rescued the hematologic phenotype and mitigated the disease. This system can be useful to test the effects on JAK1 inhibition on MDS-like disease in a preclinical model. *J. Clin. Invest.*, 123(11):4595-4611 (2013). Accordingly, a JAK1 selective inhibitor is dosed by oral gavage. The compound's ability to reduce the cytopenias and cytological dysplasia observed in the *S100A9* transgenic mice is monitored.

Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. U.S. Ser. No. 14/068,796, filed October 31, 2013, is incorporated herein by reference in its


entirety. Each reference, including all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety.

WHAT IS CLAIMED IS:

1. A process of preparing a compound of Formula 1a:


**1a**

comprising reacting a compound of Formula 1b:

**1b**


with a first strong base in the presence of a first solvent component.

2. The process of claim 1, wherein the first strong base is an alkoxide base.
3. The process of claim 1, wherein the first strong base is an alkali metal C₁-6 alkoxide.
4. The process of claim 1, wherein the first strong base is potassium *tert*-butoxide.
5. The process of any one of claims 1 to 4, wherein the compound of Formula 1b is present in a composition comprising the compound of Formula 1b and the compound of Formula 1a prior to said reacting of said compound of Formula 1b with said strong base.
6. The process of any one of claims 1 to 5, wherein the compound of Formula 1a is prepared as a compound of Formula II:




II.

7. The process of claim 5, wherein the composition comprising a compound of Formula Ia and a compound of Formula Ib is prepared by a process comprising reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb:




IIa



IIIb

with a compound of Formula IV:



IV


in the presence of a **Ci-6** alkyloxonium reagent and a second solvent component.

8. The process of claim 7, wherein before reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb, the compound of Formula IV is reacted with said **Ci-6** alkyloxonium reagent in the second solvent component.

9. The process of any one of claims 7 to 8, wherein the **Ci-6** alkyloxonium reagent is triethyloxonium tetrafluoroborate.

10. The process of any one of claims 7 to 9, wherein the composition comprising a compound of Formula IIa and a compound of Formula IIIb is prepared by a process


comprising reacting a composition comprising a compound of Formula Va and a compound of Formula Vb:




with hydrogen gas in the presence of a hydrogenation catalyst and a third solvent component.

11. The process of claim 10, wherein the hydrogenation catalyst is palladium on carbon.

12. The process of any one of claims 10 to 11, wherein the composition comprising a compound of Formula Va and a compound of Formula Vb is prepared by a process comprising reacting a compound of Formula VI:




or a salt thereof, with a compound of Formula VII:



in the presence of a first tertiary amine base and a fourth solvent component, wherein:


X is a halo group.

13. The process of claim 12, wherein the compound of Formula VI, or salt thereof, is a salt of Formula Via:



Via.


14. The process of any one of claims 12 to 13, wherein X is chloro.
15. The process of any one of claims 12 to 14, wherein the first tertiary amine base is N-ethyl-N-isopropylpropan-2-amine.
16. The process of any one of claims 12 to 15, wherein the compound of Formula VI, or salt thereof, is a salt of Formula VIa:



Via;


the fourth solvent component comprises N,N-dimethylformamide;
 the first tertiary amine base is N-ethyl-N-isopropylpropan-2-amine; and
 X is chloro.

17. The process of any one of claims 12 to 16, wherein the compound of Formula VI, or salt thereof, is produced by a process comprising reacting a compound of Formula VIII:



under deprotection conditions to form the compound of Formula VI, or salt thereof;
 wherein P¹ is an amine protecting group.


18. The process of claim 17, wherein the deprotection conditions involve reacting the compound of Formula VIII with a first strong acid.
19. The process of any one of claims 17 to 18, wherein the first strong acid is HCl.
20. The process of any one of claims 17 to 19, wherein P¹ is t-butoxycarbonyl.
21. The process of any one of claims 17 to 20, wherein the compound of Formula VIII is prepared by a process comprising reacting a compound of Formula IX:



IX

in the presence of a second strong base and a fifth solvent component, wherein $\Lambda\Lambda$ indicates a *cis*- or *trans*- bond.

22. The process of claim 21, wherein the second strong base is an **Ci-6** alkoxide base.
23. The process of claim 22, wherein the alkoxide base is sodium methoxide.
24. The process of any one of claims 21 to 23, wherein said compound of Formula IX is prepared by a process comprising reacting a compound of Formula X:




X

in the presence of a second strong acid and a sixth solvent component, wherein $\sim\sim\sim$ indicates a *cis*- or *trans*- bond.

25. The process of claim 24, wherein the second strong acid is p-toluenesulfonic acid.


26. The process of any one of claims 24 to 25, wherein said compound of Formula X is prepared by a process comprising:

(i) reacting a compound of Formula XII:



XII

with a Wittig reagent in the presence of a seventh solvent component, wherein said Wittig reagent is prepared by a process comprising reacting a compound of Formula XXI:



XXI


in the presence of a third strong base, wherein R^a and R^b are each independently C_{1-6} alkyl.

27. The process of claim 26, wherein before (i), the compound of Formula XXI is reacted with the third strong base in said seventh solvent component.

28. The process of any one of claims 26 to 27, wherein R^a and R^b are each ethyl.


29. The process of any one of claims 27 to 28, wherein the third strong base is potassium *tert*-butoxide.

30. The process of any one of claims 26 to 29, wherein said compound of Formula XII is prepared by a process comprising oxidizing a compound of Formula XIII:




XIII.

31. The process of claim 30, wherein said oxidizing comprises adding a first oxidizing agent to the compound of Formula XIII to form a first mixture.
32. The process of claim 31, wherein the first oxidizing agent is 2,2,6,6-tetramethyl-1-piperidinyloxy free radical (TEMPO).
33. The process of any one of claims 31 to 32, wherein said adding of said first oxidizing agent is conducted in the presence of tetra-N-butylammonium chloride.
34. The process of any one of claims 30 to 33, wherein the oxidizing further comprises adding a base and a second oxidizing agent to said first mixture.
35. The process of claim 34, wherein the second oxidizing agent is an N-halosuccinimide compound.
36. The process of claim 34, wherein the second oxidizing agent is N-chlorosuccinimide.
37. The process of any one of claims 30 to 36, wherein said compound of Formula XIII is prepared by a process comprising reacting a compound of Formula XV:




with a compound of Formula XIV:



in the presence of boron trifluoride diethyl etherate and an eighth solvent component.

38. The process of claim 37, wherein said compound of Formula XV is prepared by a process comprising reacting a compound of Formula XVI:




XVI

in the presence of a first reducing agent and a ninth solvent component.

39. The process of claim 38, wherein the first reducing agent is lithium borohydride.

40. The process of any one of claims 17 to 20, wherein the compound of Formula VIII is prepared by a process comprising reacting a compound of Formula XVII:




XVII

with hydrogen gas in the presence of a hydrogenation catalyst and a tenth solvent component.

41. The process of claim 40, wherein the hydrogenation catalyst is palladium on carbon.


42. The process of any one of claims 40 to 41, wherein said compound of Formula XVII is prepared by

(i) reacting a compound of Formula XVIII:



XVIII

with a Wittig reagent in the presence of an eleventh solvent component, wherein said Wittig reagent is prepared by a process comprising reacting a compound of Formula XXII:




XXII

with a fourth strong base, wherein:

R^{a1} , R^{b1} , and R^{c1} are each independently C₁₋₆ alkyl;

Y^- is a halide ion.

43. The process of claim 42, wherein Y^- is Cl^- .
44. The process of any one of claims 42 to 43, wherein R^{a1} , R^{b1} , and R^{c1} are each methyl.
45. The process of any one of claims 42 to 43, wherein R^{a1} , R^{b1} , and R^{c1} are each n-butyl.
46. The process of any one of claims 42 to 45, wherein the fourth strong base is potassium *tert*-butoxide.
47. The process of any one of claims 42 to 45, wherein the fourth strong base is lithium hexamethyldisilazide.
48. The process of any one of claims 42 to 47, wherein said compound of Formula XVIII is formed by a process comprising reacting a compound of Formula XIX:




XIX

in the presence of a weak organic acid and a twelfth solvent component.


49. The process of claim 48, wherein the weak organic acid is acetic acid.

50. The process of any one of claims 48 to 49, wherein said compound of Formula XIX is formed by a process comprising:


(1) reacting a compound of Formula XX:



with a compound of Formula XXIII:



in the presence of a second tertiary amine base and an thirteenth solvent component to form a compound of Formula XXIV:



wherein

R^{a2} is C1-4 alkoxy;

X¹ is halo.


51. The process of claim 50, wherein the compound of Formula XXIII is ethyl chloroformate.

52. The process of any one of claims 50 to 51, wherein the second tertiary amine base is N-methylmorpholine.

53. The process of any one of claims 50 to 52, further comprising reducing said compound of Formula XXIV with a second reducing agent.


54. The process of claim 53, wherein said second reducing agent is sodium borohydride.

55. An anhydrous form of a compound of Formula Ia:



Ia.

56. The form of claim 55, wherein the anhydrous form is crystalline.
57. A process of forming the anhydrous form of a compound of Formula Ia of claim 55, comprising:
- (i) adding n-heptane to a first mixture comprising a compound of Formula II:



II


and a first solvent component, wherein the first solvent component comprises ethyl acetate.

58. The process of claim 57, wherein the first solvent component further comprises methanol.
59. The process of claim 58, further comprising before (i):
- (ii) dissolving the compound of Formula II in methanol to form a second mixture comprising the compound of Formula Ia and a second solvent component comprising methanol;
- (iii) polish filtering the second mixture; and
- (iv) adding ethyl acetate to the second mixture to form the first mixture.

60. The process of any one of claims 58 to 59, further comprising heating the first mixture to a temperature from about 60 °C to about 75 °C.

61. The process of any one of claims 58 to 60, comprising removing at least a portion of the ethyl acetate and methanol to form the first mixture via distillation.


62. A compound of Formula VIII:




wherein P¹ is an amine protecting group.

63. The compound of claim 62, wherein P¹ is t-butoxycarbonyl.

64. A salt of Formula Via:




65. A compound of Formula Va:




or salt thereof.

66. A compound of Formula Vb:




or salt thereof.

67. A compound of Formula IIIa:




or salt thereof.

68. A compound of Formula IIIb:



or salt thereof.

69. A process for preparing a compound of Formula II:




comprising suspending an anhydrous form of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile in a solvent component comprising water.

70. The process of claim 69, wherein said suspending is conducted at about room temperature.

71. The process of claim 69, wherein the solvent component is water.

72. A process of preparing a compound of Formula II:



comprising isolating the compound of Formula II from a first mixture comprising of 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile and a solvent component comprising isopropanol and water.


73. The process of claim 72, further comprising, prior to said isolating, the steps of:

- (a) dissolving 2-((2R,5S)-5-(2-((R)-1-hydroxyethyl)-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl)tetrahydro-2H-pyran-2-yl)acetonitrile in a solvent component comprising isopropanol to form a second mixture; and
- (b) adding water to said second mixture to form said first mixture, wherein the solvent component comprises isopropanol and water.


74. The process of claim 73, further comprising removing at least a portion of said solvent component from said first mixture prior to said isolating.

75. The process of claim 73, where (a) and (b) are conducted at a temperature from about 40 °C to about 60 °C.

76. The process of claim 5, wherein the composition comprising a compound of Formula Ia and a compound of Formula Ib is prepared by a process comprising reacting a composition comprising a compound of Formula IIa and a compound of Formula IIb:




with (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole:



or a salt thereof.


77. The process of claim 76, wherein said (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole is prepared by reacting (*R*)-2,2,5-trimethyloxazolidin-4-one:




with trimethyloxonium tetrafluoroborate.

78. The process of claim 77, wherein the steps of reacting (*R*)-2,2,5-trimethyloxazolidin-4-one with trimethyloxonium tetrafluoroborate to form (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole and reacting (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole with a composition comprising a compound of Formula IIa and a compound of Formula IIb are conducted in the same pot without isolation of (*R*)-4-methoxy-2,2,5-trimethyl-2,5-dihydrooxazole.


79. The process of claim 5, wherein the composition comprising a compound of Formula Ia and a compound of Formula Ib is prepared by a process comprising reacting a composition comprising a compound of Formula IIa and a compound of Formula IIIb:



with (*R*)-ethyl 2-hydroxypropanimidate:




80. The process of claim 79, wherein said (*R*)-ethyl 2-hydroxypropanimidate is prepared by reacting (*R*)-1-cyanoethyl acetate:



with ethanol in the presence of a strong acid.


81. The process of claim 80, wherein said strong acid used in the reaction of (*R*)-1-cyanoethyl acetate with ethanol is hydrogen chloride.

82. The process of any one of claims 80 to 81, wherein said (*R*)-1-cyanoethyl acetate is formed by reacting (*R*)-1-amino-1-oxopropan-2-yl acetate:




with cyanuric chloride.

83. The process of claim 82, wherein said (*R*)-1-amino-1-oxopropan-2-yl acetate is prepared by protecting (*R*)-2-hydroxypropanamide:



with acetyl chloride in the presence of an third tertiary amine base.

84. The process of claim 83, wherein said third tertiary amine base is 4-methylmorpholine.







FIG. 2

**FIG. 3**




FIG. 4




FIG. 5





FIG. 6

7/15

**FIG. 7**

8/15




FIG. 8




FIG. 9





FIG. 10



Universal V4.5A TA Instruments

FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2015/028224

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D495/14 A61K31/437 A61P25/00 A61P35/00 A61P37/00
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal , CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	wo 2014/071031 AI (INCYTE CORP [US]) 8 May 2014 (2014-05-08) see example 20, step 2, example 73, step 3 including the cyano intermediate, steps 4 and 5, starting material in step 2 of example 74 and reactant of example 91 -----	55,63-70
A	wo 2007/044050 A2 (SQUIBB BRISTOL MYERS CO [US] ; PITTS WILLIAM J [US] ; DAS JAGABANDHU [US] 19 April 2007 (2007-04-19) see the compounds of claim 1 and the processes for their production -----	1-84



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 July 2015

Date of mailing of the international search report

21/07/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Traegl er-Goel del , M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2015/028224

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014071031 A1	08-05-2014	AU 2013337824 A1	14-05-2015
		CA 2888816 A1	08-05-2014
		TW 201422624 A	16-06-2014
		US 2014121198 A1	01-05-2014
		WO 2014071031 A1	08-05-2014

WO 2007044050 A2	19-04-2007	US 2006178393 A1	10-08-2006
		WO 2007044050 A2	19-04-2007
