



US008222010B2

(12) **United States Patent**  
**Franklin et al.**

(10) **Patent No.:** **US 8,222,010 B2**  
(45) **Date of Patent:** **\*Jul. 17, 2012**

(54) **RENEWABLE CHEMICAL PRODUCTION FROM NOVEL FATTY ACID FEEDSTOCKS**

(75) Inventors: **Scott Franklin**, La Jolla, CA (US); **Aravind Somanchi**, Redwood City, CA (US); **Karen Espina**, San Francisco, CA (US); **George Rudenko**, Mountain View, CA (US); **Penelope Chua**, San Francisco, CA (US)

(73) Assignee: **Solazyme, Inc.**, South San Francisco, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/981,409**

(22) Filed: **Dec. 29, 2010**

(65) **Prior Publication Data**

US 2011/0165634 A1 Jul. 7, 2011

**Related U.S. Application Data**

(63) Continuation of application No. 12/628,149, filed on Nov. 30, 2009, now Pat. No. 7,883,882.

(60) Provisional application No. 61/118,590, filed on Nov. 28, 2008, provisional application No. 61/118,994, filed on Dec. 1, 2008, provisional application No. 61/174,357, filed on Apr. 30, 2009, provisional application No. 61/219,525, filed on Jun. 23, 2009.

(51) **Int. Cl.**

- C12P 7/64** (2006.01)
- C12P 21/02** (2006.01)
- C12N 9/16** (2006.01)
- C12N 9/90** (2006.01)
- C12N 9/10** (2006.01)
- C12N 9/02** (2006.01)

(52) **U.S. Cl.** ..... **435/134; 435/196; 435/233; 435/193; 435/189; 435/69.1**

(58) **Field of Classification Search** ..... **435/134, 435/196, 233, 193, 189, 69.1**  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,235,056	A	3/1941	Walmesley
3,475,274	A	10/1969	Harned
3,957,578	A	5/1976	Narita et al.
3,983,008	A	9/1976	Shinozaki et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN 1940021 A 4/2007

(Continued)

OTHER PUBLICATIONS

St. Onge et al., "Physiological effects of medium-chain triglycerides: potential agents in the prevention of obesity," J Nutrition 132:329-332, 2002.\*

(Continued)





*Primary Examiner* — Rosanne Kosson

(74) *Attorney, Agent, or Firm* — Alston & Bird LLP

(57) **ABSTRACT**

Disclosed herein are methods of manufacturing renewable chemicals through the manufacture of novel triglyceride oils followed by chemical modification of the oils. Methods such as transesterification, hydrogenation, hydrocracking, deoxygenation, isomerization, interesterification, hydroxylation, hydrolysis and saponification are disclosed. Novel oils containing fatty acid chain lengths of C8, C10, C12 or C14 are also disclosed and are useful as feedstocks in the methods of the invention.

**19 Claims, 12 Drawing Sheets**



U.S. PATENT DOCUMENTS			FOREIGN PATENT DOCUMENTS		
4,341,038 A	7/1982	Bloch et al.	2009/0099260 A1	4/2009	Namal Senanayake et al.
4,373,434 A	2/1983	Alexander et al.	2009/0145392 A1	6/2009	Clark et al.
4,519,845 A	5/1985	Ou	2009/0148918 A1	6/2009	Trimbur et al.
4,673,490 A	6/1987	Subramanian et al.	2009/0197312 A1	8/2009	Pastinen et al.
4,755,467 A	7/1988	Scopes et al.	2009/0211150 A1	8/2009	Wu et al.
4,901,635 A	2/1990	Williams	2009/0298159 A1	12/2009	Wu et al.
4,992,605 A	2/1991	Craig et al.	2009/0305942 A1	12/2009	Day et al.
5,091,116 A	2/1992	Krishnamurthy et al.	2010/0021912 A1	1/2010	Farese et al.
5,130,242 A	7/1992	Barclay	2010/0058651 A1	3/2010	Knuth et al.
5,252,198 A	10/1993	Harrison et al.	2010/0120643 A1	5/2010	Brown et al.
5,270,177 A	12/1993	Ramos Lazcano et al.	2010/0151112 A1	6/2010	Franklin et al.
5,304,481 A	4/1994	Davies et al.	2010/0151538 A1	6/2010	Franklin et al.
5,338,673 A	8/1994	Thepenier et al.	2010/0151539 A1	6/2010	Franklin et al.
5,354,878 A	10/1994	Connemann et al.	2010/0151567 A1	6/2010	Franklin et al.
5,391,724 A	2/1995	Kindl et al.	2010/0170144 A1	7/2010	Day et al.
5,395,455 A	3/1995	Scott et al.	2010/0323413 A1	12/2010	Trimbur et al.
5,455,167 A *	10/1995	Voelker et al. .... 800/281	2010/0323414 A1	12/2010	Trimbur et al.
5,492,938 A *	2/1996	Kyle et al. .... 514/786	2011/0014665 A1	1/2011	Trimbur et al.
5,597,400 A	1/1997	Nonomura et al.	2011/0015417 A1	1/2011	Trimbur et al.
5,680,812 A	10/1997	Linsgeseder	2011/0047863 A1	3/2011	Trimbur et al.
5,685,218 A	11/1997	Kemper	2011/0190522 A1	8/2011	Trimbur et al.
5,693,507 A	12/1997	Daniell et al.	2011/0250658 A1	10/2011	Franklin et al.
5,826,500 A	10/1998	Kemper	2011/0252696 A1	10/2011	Franklin et al.
5,888,947 A	3/1999	Lambert et al.	2012/0034662 A1	2/2012	Hu et al.
5,900,370 A	5/1999	Running			
6,166,231 A	12/2000	Hoeksema	CN	101037639 A	9/2007
6,255,505 B1	7/2001	Bijl et al.	CN	101130513 A	2/2008
6,410,281 B1 *	6/2002	Barclay ..... 435/134	DE	2756977 A1	6/1978
6,441,208 B2	8/2002	Bijl et al.	EP	1178118 A1	2/2002
6,589,923 B2	7/2003	Lenuck et al.	EP	1681337 A1	7/2006
6,680,426 B2	1/2004	Daniell et al.	EP	1741767 A1	1/2007
6,750,048 B2	6/2004	Ruecker et al.	JP	60006799 A	1/1985
7,053,267 B2	5/2006	Knauf et al.	WO	WO 92/11373 A1	7/1992
7,081,567 B2	7/2006	Xue et al.	WO	WO 94/10288 A2	5/1994
7,135,620 B2	11/2006	Daniell et al.	WO	WO 95/13390 A2	5/1995
7,268,276 B2	9/2007	Ruezinsky et al.	WO	WO 99/64618 A1	11/1999
7,309,602 B2	12/2007	David	WO	WO 00/61740 A1	10/2000
7,351,558 B2	4/2008	Ruecker et al.	WO	WO 2004/101753 A2	11/2004
7,662,598 B2	2/2010	Ruecker et al.	WO	WO 2005/035693 A2	4/2005
7,678,931 B2	3/2010	Fichtali et al.	WO	WO 2007/027669 A1	3/2007
7,781,193 B2	8/2010	Ruecker et al.	WO	WO 2007/038566 A2	4/2007
7,851,199 B2	12/2010	Bailey et al.	WO	WO 2007/134294 A2	11/2007
7,883,882 B2 *	2/2011	Franklin et al. .... 435/196	WO	WO 2008/002643 A2	1/2008
7,935,515 B2 *	5/2011	Franklin et al. .... 435/257.1	WO	WO 2008/011811 A1	1/2008
8,029,579 B2	10/2011	Knuth et al.	WO	WO 2008/083352 A1	7/2008
2002/0012979 A1	1/2002	Berry	WO	WO 2008/134836 A2	11/2008
2002/0144455 A1	10/2002	Bertrand et al.	WO	WO 2008/151149 A2	12/2008
2002/0178467 A1	11/2002	Dehesh	WO	WO 2009/126843 A2	10/2009
2003/0082595 A1	5/2003	Jiang et al.	WO	WO 2010/045368 A2	4/2010
2003/0097686 A1	5/2003	Knauf et al.	WO	WO 2010/063031 A2	6/2010
2003/0211594 A1	11/2003	Rosebrook	WO	WO 2010/063032 A2	6/2010
2004/0230085 A1	11/2004	Jakkula et al.	WO	WO 2011/130573 A1	10/2011
2004/0235123 A1	11/2004	Liao et al.			
2005/0005333 A1	1/2005	Ruezinsky et al.			
2005/0084941 A1	4/2005	Abe et al.			
2005/0102716 A1	5/2005	Venkatramesh et al.			
2005/0112735 A1	5/2005	Zappi et al.			
2006/0107346 A1	5/2006	Schneeberger et al.			
2006/0122410 A1	6/2006	Fichtali et al.			
2006/0153826 A1	7/2006	Arnould et al.			
2006/0162006 A9	7/2006	Sherman et al.			
2006/0199984 A1	9/2006	Kuechler et al.			
2006/0225341 A1	10/2006	Rohr et al.			
2007/0004016 A1	1/2007	Picataggio et al.			
2007/0048848 A1	3/2007	Sears			
2007/0118916 A1	5/2007	Puzio et al.			
2007/0275438 A1	11/2007	David et al.			
2007/0286908 A1	12/2007	Clampitt			
2008/0038804 A1	2/2008	Du et al.			
2008/0160593 A1	7/2008	Oyler			
2008/0206379 A1	8/2008	Fabritius et al.			
2008/0256666 A1	10/2008	Zhu et al.			
2009/0004715 A1	1/2009	Trimbur et al.			
2009/0011480 A1	1/2009	Trimbur et al.			
2009/0018300 A1	1/2009	Bloom et al.			
2009/0035842 A1	2/2009	Trimbur et al.			
2009/0047721 A1	2/2009	Trimbur et al.			
2009/0061493 A1	3/2009	Trimbur et al.			

## OTHER PUBLICATIONS

PCT Search Report for application PCT/US2011/038464 mailed Nov. 3, 2011.

PCT Written Opinion of the International Searching Authority for application PCT/US2011/038464 mailed Nov. 3, 2011.

Abate et al., "Production of ethanol by a flocculent *Saccharomyces* sp. in a continuous upflow reactor using sucrose, sugar-cane juice, and molasses as the carbon source," *MIRCEN Journal*, 3:401-409, (1987).

Abbadì et al., "Knockout of the regulatory site of 3-ketoacyl-ACP synthase III enhances short- and medium-chain acyl-ACP synthesis," *The Plant Journal*, 24(1):1-9, (2000).

Alvarez et al., "Triacylglycerols in prokaryotic microorganisms," *Appl Microbiol Biotechnol*, 60:367-376, (2002).

Angerbauer et al., "Conversion of sewage sludge into lipids by *Lipomyces starkeyi* for biodiesel production," *Bioresource Technology*, 99:3051-3056, (2008).

Broun et al., "A bifunctional oleate 12-hydroxylase: desaturase from *Lesquerella fendleri*," *The Plant Journal*, 13(2):201-210, (1998).

Canakci et al., "Biodiesel production from oils and fats with high free fatty acids," *Transactions of the ASAE*, 44(6):1429-1436, (2001).

Cartens et al., "Eicosapentaenoic Acid (20:5n-3) from the Marine Microalga *Phaeodactylum tricoratum*," *Journal of the American Oil Chemists' Society*, 73(8):1025-1031, (1996).

- Chen et al., "High cell density culture of microalgae in heterotrophic growth," *Trends in Biotechnology*, 14:421-426, (1996).
- Chisti et al., "Biodiesel from microalgae," *Biotechnology Advances*, 25:294-306, (2007).
- Covello et al., "Functional Expression of the Extraplasmidial *Arabidopsis thaliana* Oleate Desaturase Gene (FAD2) in *Saccharomyces cerevisiae*," *Plant Physiol.*, 111:223-226, (1996).
- Dai et al., "Biodiesel generation from oleaginous yeast *Rhodotorula glutinis* with xylose assimilating capacity," *African Journal of Biotechnology*, 6(18):2130-2134, (2007).
- Dehesh et al., "Production of high levels of 8:0 and 10:0 fatty acids in transgenic canola by overexpression of Ch FatB2, a thioesterase cDNA from *Cuphea hookeriana*," *The Plant Journal*, 9(2):167-172, (1996).
- Demirbas, "Fuel Conversional Aspects of Palm Oil and Sunflower Oil," *Energy Sources*, 25:457-466, (2003).
- Dunahay et al., "Manipulation of Microalgal Lipid Production Using Genetic Engineering," *Applied Biochemistry and Biotechnology*, 57/58:223-231, (1996).
- El-Fadaly et al., "Single Cell Oil Production by an Oleaginous Yeast Strain in a Low Cost Cultivation Medium," *Research Journal of Microbiology*, 4(8):301-313, (2009).
- European Search Report and European Search Opinion for application EP08769988 mailed Jul. 1, 2011.
- European Search Report and European Search Opinion for application EP11158642 mailed Jul. 1, 2011.
- Evans et al., "A comparison of the oleaginous yeast, *Candida curvata*, grown on different carbon sources in continuous and batch culture," *Lipids*, 18(09):623-629, (1983).
- Facciotti et al., "Improved stearate phenotype in transgenic canola expressing a modified acyl-acyl carrier protein thioesterase," *Nat Biotechnol.*, 17(6):593-597, (1999).
- Gill et al., "Lipid Accumulation in an Oleaginous Yeast (*Candida 107*) Growing on Glucose in Single-Stage Continuous Culture," *Applied and Environmental Microbiology*, 33(02):231-239, (1977).
- Hall et al., "Lipid Accumulation in an Oleaginous Yeast (*Candida 107*) Growing on Glucose Under Various Conditions in a One- and Two-Stage Continuous Culture," *Applied and Environmental Microbiology*, 33(3):577-584, (2009).
- Heise et al., "Factors Controlling Medium-Chain Fatty Acid Synthesis in Plastids From *Cuphea* Embryos," *Prog. Lipid Res.*, 33(1/2):87-95, (1994).
- Helmensteine, "Gasoline and Octane Ratings," About.com, Chemistry, 1 page, (2011). [Retrieved from the Internet Oct. 3, 2011: <URL: <http://chemistry.about.com/cs/howthingswork/a/aa070401a.htm>>].
- Heredia et al., "Simultaneous utilization of glucose and xylose by *Candida curvata* D in continuous culture," *Biotechnology Letters*, 10(01):25-30, (1988).
- Hossain et al., "The effect of the sugar source on citric acid production by *Aspergillus niger*," *Appl Microbiol Biotechnol*, 19:393-397, (1984).
- Huber et al., "Synthesis of Transportation Fuels from Biomass: Chemistry, Catalysts, and Engineering," *Chem. Rev.*, 106: 4044-4098, (2006).
- Jaworski et al., "Industrial oils from transgenic plants," *Current Opinion in Plant Biology*, 6:178-184, (2003).
- Jha et al., "Cloning and functional expression of an acyl-ACP thioesterase FatB type from *Diploknema (Madhuca) butyracea* seeds in *Escherichia coli*," *Plant Physiology and Biochemistry*, 44:645-655, (2006).
- Kong et al., "Microbial production of lipids by cofermentation of glucose and xylose with *Lipomyces starkeyi* 2#," *Chinese Journal of Bioprocess Engineering*, 05(02):36, (2007). Abstract.
- Li et al., "Large-scale biodiesel production from microalga *Chlorella protothecoides* through heterotrophic cultivation in bioreactors," *Biotechnology and Bioengineering*, 98(04):764-771, (2007).
- Li et al., "Enzymatic transesterification of yeast oil for biodiesel fuel production," *The Chinese Journal of Process Engineering*, 07(01):137-140 (2007), and machine translation.
- Li et al., "High-density cultivation of oleaginous yeast *Rhodospiridium toruloides* Y4 in fed-batch culture," *Enzyme and Microbial Technology*, 41:312-317, (2007).
- Li et al., "Screening of oleaginous yeasts for broad-spectrum carbohydrates assimilating capacity," *China Biotechnology*, 25(12):39-44 (2005), and machine translation.
- Liu et al., "Biodiesel production by direct methanolysis of oleaginous microbial biomass," *Journal of Chemical Technology and Biotechnology*, 82:775-780, (2007).
- Mayer et al., "A Structural Model of the Plant Acyl-Acyl Carrier Protein Thioesterase FatB Comprises Two Helix/4-Stranded Sheet Domains, the N-terminal Domain Containing Residues That Affect Specificity and the C-terminal Domain Containing Catalytic Residues," *The Journal of Biological Chemistry*, 280(5):3621-3627, (2005).
- Meesters et al., "High-cell-density cultivation of the lipid accumulating yeast *Cryptococcus curvatus* using glycerol as a carbon source," *Applied Microbiology and Biotechnology*, 45:575-579, (1996).
- Mekhedov et al., "Toward a Functional Catalog of the Plant Genome. A Survey of Genes for Lipid Biosynthesis," *Plant Physiology*, 122:389-401, (2000).
- Murakami et al., "Lipid Composition of Commercial Bakers' Yeasts Having Different Freeze-tolerance in Frozen Dough," *Biosci. Biotechnol. Biochem.*, 60(11):1874-1876, (1996).
- Nes et al., "Sterol phylogenesis and algal evolution," *Proc Natl Acad Sci U S A*, 87(19):7565-7569, (1990).
- Papanikolaou et al., "Lipid production by *Yarrowia lipolytica* growing on industrial glycerol in a single-stage continuous culture," *Bioresource Technology*, 82:43-49, (2002).
- PCT International Preliminary Report on Patentability (Chapter I) of May 31, 2011 for application PCT/US09/066142.
- PCT International Preliminary Report on Patentability (Chapter I) of Oct. 12, 2010 for application PCT/US2009/040123.
- PCT Search Report for application PCT/US2009/040123 mailed Oct. 5, 2009.
- PCT Search Report for application PCT/US2011/0032582 mailed Aug. 9, 2011.
- PCT Written Opinion of the International Searching Authority for application PCT/US2011/0032582 mailed Aug. 9, 2011.
- Popov et al., "Separation of the total lipid extract of microalgae," *Doklady Akademii Sel'skokhozyaistvennykh Nauk v Bolgarii*, 3(1):45-54, (1970). Abstract only.
- Radakovits et al., "Genetic Engineering of Algae for Enhanced Biofuel Production," *Eukaryotic Cell*, 9(04): 486-501, (2010).
- Rehm et al., "Heterologous expression of the acyl-acyl carrier protein thioesterase gene from the plant *Umbellularia californica* mediates polyhydroxyalkanoate biosynthesis in recombinant *Escherichia coli*," *Appl Microbiol Biotechnol*, 55:205-209, (2001).
- Roy et al., "Production of Intracellular Fat by the Yeast *Lipomyces starkeyi*," *Indian Journal of Experimental Biology*, 16(4):511-512, (1978).
- Running et al., "The pathway of L-ascorbic acid biosynthesis in the colourless microalga *Prototheca moriformis*," *Journal of Experimental Botany*, 54(389): 1841-1849, (2003).
- Salou et al., "Growth and Energetics of *Leuconostoc oenos* during Cometabolism of Glucose with Citrate or Fructose," *Applied and Environmental Microbiology*, 60(5): 1459-1466, (1994).
- Schütt et al., "The role of acyl carrier protein isoforms from *Cuphea lanceolata* seeds in the de-novo biosynthesis of medium-chain fatty acids," *Publication, Planta*, 205:263-268, (1998).
- Schütt et al., "β-Ketoacyl-acyl carrier protein synthase IV: a key enzyme for regulation of medium-chain fatty acid synthesis in *Cuphea lanceolata* seeds," *Planta*, 215:847-854, (2002).
- Suh et al., "What limits production of unusual monoenoic fatty acids in transgenic plants?," *Planta*, 215:584-595, (2002).
- Tan et al., "Saponified Palm Kernel Oil and Its Major Free Fatty-Acids as Carbon Substrates for the Production of Polyhydroxyalkanoates in *Pseudomonas-Putida* Pgal," *Applied Microbiology and Biotechnology*, 47(3):207-211, (1997).
- Thavarungkul et al., "Batch injection analysis for the determination of sucrose in sugar cane juice using immobilized invertase and thermometric detection," *Biosensors & Bioelectronics*, 14:19-25, (1999).
- U.S. Appl. No. 12/131,766, Advisory Action mailed Oct. 13, 2011.

- U.S. Appl. No. 12/131,766, Non-Final Office Action mailed Aug. 1, 2011.
- U.S. Appl. No. 12/131,773, Requirement for Restriction/Election mailed Aug. 6, 2009.
- U.S. Appl. No. 12/131,783, Non-Final Office Action mailed Jun. 6, 2011.
- U.S. Appl. No. 12/131,783, Requirement for Restriction/Election mailed Apr. 19, 2011.
- U.S. Appl. No. 12/499,033, Final Office Action mailed Oct. 19, 2011.
- U.S. Appl. No. 12/499,033, Non-Final Office Action mailed Feb. 18, 2011.
- U.S. Appl. No. 12/499,033, Requirement for Restriction/Election mailed Oct. 15, 2010.
- U.S. Appl. No. 12/628,144, Non-Final Office Action mailed Jun. 7, 2011.
- U.S. Appl. No. 12/628,147, Examiner Interview Summary Record mailed Mar. 3, 2011.
- U.S. Appl. No. 12/628,147, Non-Final Office Action mailed Oct. 25, 2011.
- U.S. Appl. No. 12/772,163, Requirement for Restriction/Election mailed Jun. 24, 2011.
- U.S. Appl. No. 12/772,164, Non-Final Office Action mailed Oct. 12, 2011.
- U.S. Appl. No. 12/772,164, Requirement for Restriction/Election mailed Jul. 20, 2011.
- U.S. Appl. No. 12/772,170, Non-Final Office Action mailed Sep. 13, 2011.
- U.S. Appl. No. 12/772,170, Requirement for Restriction/Election mailed Jul. 13, 2011.
- U.S. Appl. No. 12/772,173, Requirement for Restriction/Election mailed Oct. 26, 2011.
- U.S. Appl. No. 12/772,174, Requirement for Restriction/Election mailed Aug. 10, 2011.
- U.S. Appl. No. 13/073,757, Non-Final Office Action mailed Aug. 15, 2011.
- Van De Loo et al., "An oleate 12-hydroxylase from *Ricinus communis* L. is a fatty acyl desaturase homolog," *Proc. Natl. Acad. Sci. USA*, 92:6743-6747, (1995).
- Voelker et al., "Alteration of the Specificity and Regulation of Fatty Acid Synthesis of *Escherichia coli* by Expression of a Plant Medium Chain Acyl-Acyl Carrier Protein Thioesterase," *Journal of Bacteriology*, 176(23):7320-7327, (1994).
- Voelker et al., "Broad-Range and Binary-Range Acyl-Acyl-Carrier-Protein Thioesterases Suggest an Alternative Mechanism for Medium-Chain Production in Seeds," *Plant Physiol.*, 114:669-677, (1997).
- Voetz et al., "Three Different cDNAs Encoding Acyl Carrier Proteins from *Cuphea lanceolata*," *Plant Physiol.*, 106:785-786, (1994).
- Wang et al., "Quantitative estimate of the effect of cellulase components during degradation of cotton fibers," *Carbohydrate Research*, 339:819-824, (2004).
- Wiberg et al., "The distribution of caprylate, caprate and laurate in lipids from developing and mature seeds of transgenic *Brassica napus* L.," *Planta*, 212:33-40, (2000).
- Wu et al., "A Comparative Study of Gases Generated from Simviant Thermal Degradation of Autotrophic and Heterotrophic *Chlorella*," *Progress in Natural Science*, 2(4):311-318, (1992).
- Wu et al., "New Discoveries in Study on Hydrocarbons From Thermal Degradation of Heterotrophically Yellowing Algae," *Science in China*, 37(3):326-35, (1994).
- Xu et al., "High quality biodiesel production from a microalga *Chlorella protothecoides* by heterotrophic growth in fermenters," *Journal of Biotechnology*, 126:499-507, (2006).
- Zarowska et al., "Production of Citric Acid on Sugar Beet Molasses by Single and Mixed Cultures of *Yarrowia Lipolytica*," *Electronic Journal of Polish Agricultural Universities*, 4(2):1-7, (2001). [Retrieved from the Internet Oct. 3, 2011: <URL: <http://www.ejpau.media.pl/volume4/issue2/biotechnology/art-01.html>>].
- Zhao et al., "Toward cheaper microbial oil for biodiesel oil," *China Biotechnology*, 25(02):8-11 (2005).
- U.S. Appl. No. 60/941,581, filed Jun. 1, 2007, Trimbur et al.
- U.S. Appl. No. 60/959,174, filed Jul. 10, 2007, Trimbur et al.
- U.S. Appl. No. 60/968,291, filed Aug. 27, 2007, Trimbur et al.
- U.S. Appl. No. 61/024,069, filed Jan. 28, 2008, Trimbur et al.
- U.S. Appl. No. 61/118,590, filed Jan. 28, 2008, Franklin et al.
- U.S. Appl. No. 61/118,994, filed Dec. 1, 2008, Franklin et al.
- U.S. Appl. No. 61/174,357, filed Apr. 30, 2009, Franklin et al.
- U.S. Appl. No. 61/219,525, filed Jun. 23, 2009, Franklin et al.
- A Look Back at the U.S. Department of Energy's Aquatic Species Program: Biodiesel from Algae, NREL/TP-580-24190, John Sheehan, Terri Dunahay, John Benemann and Paul Roesler (1998).
- Bonaventure et al., "Disruption of the FATB Gene in *Arabidopsis* Dethonstrates an Essential Role of Saturated Fatty Acids in Plant Growth," *The Plant Cell* 15:1020-1033 (2003).
- Borza et al., "Multiple Metabolic Roles for the Nonphotosynthetic Plastid of the Green Alga *Prototheca wickerhamii*," *Eukaryotic Cell* 4(2):253-261 (2005).
- Bouchard et al., "Characterization of Depolymerized Cellulosic Residues," *Wood Sci. Technol.*, 23:343-355 (1989).
- Chasan, "Engineering Fatty Acids—The Long and Short of It," *Plant Cell*, 7:235-237 (1995).
- Chen et al., "Highly Effective Expression of Rabbit Neutrophil Peptide-1 Gene in *Chlorella ellipsoidea* Cells," *Current Genetics*, 39:365-370 (2001).
- Chow et al., "Electrotransformation of *Chlorella vulgaris*," *Plant Cell Reports*, 18:778-780 (1999).
- Cobley et al., Construction of Shuttle Plasmids Which Can Be Efficiently Mobilized From *Escherichia coli* Into the Chromatically Adapting Cyanobacterium, *Plasmid*, 30:90-105 (1993).
- Courchesne et al., "Enhancement of Lipid Production Using Biochemical, Genetic and Transcription Factor Engineering Approaches," *J Biotechnol. Epub*, 141(1-2):31-41 (2009).
- Dawson et al., "Stable Transformation of *Chlorella*: Rescue of Nitrate Reductase-Deficient Mutants with the Nitrate Reductase Gene," *Current Microbiology*, 35:356-362 (1997).
- Deng et al., "Ionic Liquid as a Green Catalytic Reaction Medium for Esterifications," *J. Mol. Catalysis A: Chemical*, 165:33-36 (2001).
- Eccleston et al., "Medium-chain Fatty Acid Biosynthesis and Utilization in *Brassica mapus* Plants Expressing Lauroyl-Acyl Carrier Protein Thioesterase," *Planta* 198:46-53 (1996).
- El-Sheekh, MM., Stable Transformation of the Intact Cells of *Chlorella kessleri* With High Velocity Microprojectiles, *Biologia Plantarum* 42(2): 209-216 (1999).
- Falciatore et al., "Transformation of Nonselectable Reporter Genes in Marine Diatoms," *Marine Biotechnology*; 1:239-251 (1999).
- Fukuda et al., "Biodiesel Fuel Production by Transesterification of Oils," *J. Biosci. Bioeng.*, 92(5):405-416 (2001).
- GenBank: "Codon Usage Database file for *Chlorella vulgaris*," Jun. 2007, [Retrieved from the Internet Aug. 26, 2010: <URL: <http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=3077>>].
- GenBank: Accession No. L42851.1, "*Prototheca wickerhamii* large subunit ribosomal RNA (rrnL) gene, partial sequence; chloroplast gene for chloroplast product," Nov. 21, 2001.
- Grinna et al., "Size Distribution and General Structural Features of N-Linked Oligosaccharides from the Methylotrophic Yeast, *Pichia pastoris*," *Yeast*, 5:107-115 (1989).
- Guo-Zhong et al., "The Actin Gene Promoter-driven Bar as a Dominant Selectable Marker for Nuclear Transformation of *Dunaliella salina*," *Acta Genetica Sinica*, 32(4): 424-433 (2005).
- Gusakov et al., "Design of Highly Efficient Cellulase Mixtures for Enzymatic Hydrolysis of Cellulose," *Biotechnol. and Bioengineering*, 97(5):1028-1038 (2007).
- Hallman et al., "Reporter Genes and Highly Regulated Promoters as Tools for Transformation Experiments in *Volvox carteri*," *Proc Natl Acad Sci U S A.*, 91(24):11562-6 (1994).
- Hawkins et al., "Expression of Human Growth Hormone by the Eukaryotic Alga, *Chlorella*," *Current Microbiology*, 38:335-341 (1999).
- Heifetz, "Genetic Engineering of the Chloroplast," *Biochimie*, 82:655-666 (2000).
- Hsieh et al., "A PII-like Protein in *Arabidopsis*: Putative Role in Nitrogen Sensing," *Proc Natl Acad Sci USA*, 95(23):13965-13970 (1998).
- Hu et al., "Microalgal Triacylglycerols as Feedstocks for Biofuel Production: Perspectives and Advances," *The Plant Journal* 54:621-639 (2008).

- Huang et al., "Expression of Mercuric Reductase From *Bacillus megaterium* MB1 in Eukaryotic Microalga *Chlorella* sp. DT: An Approach for Mercury Phytoremediation," *Appl. Microbiol. Biotechnol.*, 72:197-205 (2006).
- Huber et al., "Production of Liquid Alkanes by Aqueous-Phase Processing of Biomass-Derived Carbohydrates," *Science*, 308: 1446-1450 (2005).
- Jarvis et al. "Transient Expression of Firefly Luciferase in Protoplasts of the Green Alga *Chlorella ellipsoidea*," *Current Genet.*, 19: 317-322 (1991).
- Jeoh et al., "Cellulose Digestibility of Pretreated Biomass Is Limited by Cellulose Accessibility," *Biotechnol Bioeng.*, 98(1):112-122 (2007).
- Kalscheuer et al., "Establishment of a Gene Transfer System for *Rhodococcus opacus* PD630 Based on Electroporation and its Application for Recombinant Biosynthesis of Poly(3-hydroxyalkanoic acids)," *Applied Microbiology and Biotechnology*, 52(4):508-515 (1999).
- Kang et al., "The regulation activity of *Chlorella* virus gene 5' upstream sequence in *Escherichia coli* and eucaryotic alage," *Institute of Microbiology, Chinese Academy of Sciences, Beijing*, 16(4):443-6 (2000). Abstract only.
- Katayama et al., "Alpha-Linolenate and Photosynthetic Activity in *Chlorella protothecoides*," *Plant Physiol.*, 42:308-313 (1967).
- Kim et al. "Stable Integra and Functional Expression of Flounder Growth Hormone Gene in Transformed Microalga, *Chlorella ellipsoidea*," (*Mar. Biotechnol.* 4:63-73 (2002).
- Koksharova, "Genetic Tools for Cyanobacteria," *Appl Microbiol Biotechnol*, 58(2):123-37 (2002).
- Lawford et al., "Performance Testing of *Zymomonas mobilis* Metabolically Engineered for Conformation of Glucose, Xylose, and Arabinose," *Appl Biochem Biotechnol.*, 98-100:429-48 (2002).
- Li et al., "Articles: Biocatalysts and Bioreactor Design, Biofuels From Microalgae," *Biotechnol. Prog.* 24:815-820 (2008).
- Lumbreras et al., "Efficient Foreign Gene Expression in *Chlamydomonas reinhardtii* Mediated by an Endogenous Intron," *Plant Journal*, 14(4):441-447 (1998).
- Maruyama et al., "Introduction of Foreign DNA Into *Chlorella saccharophila* by Electroporation," *Biotechnology Techniques*, 8:821-826 (2004).
- Matsuka et al., *Plant and Cell Physiol.*, 7: 149-162 (1966).
- Miao et al., "Biodiesel Production From Heterotrophic Microalgal Oil," *Biosource Technology*, 97:841-846 (2006).
- Miao et al., "High Yield Bio-Oil Production from Fast Pyrolysis by Metabolic Controlling of *Chlorella* Protothecoides," *J. Biotech.*, 110:85-93 (2004).
- Muller et al., "Nitrogen and hormonal responsiveness of the 22 kDa alpha-zein and b-32 genes in maize endosperm is displayed in the absence of the transcriptional regulator Opaque-2," *Plant J.*, 12(2):281-291, Abstract (1997).
- Onai et al., "Natural Transformation of the Thermophilic Cyanobacterium *Thermosynechococcus elongatus* BP-1: A Simple and Efficient Method for Gene Transfer," *Mol Genet Genomics*, 271(1):50-9 (2004).
- Papanikolaou et al., *J. Chem. Technol. Biotechnol.*, 79: 1189-1196 (2004).
- Park et al., "Isolation and Characterization of *Chlorella* Virus From Fresh Water in Korea and Application in *Chlorella* Transformation System," *Plant Pathol. J.*, 21(1): 13-20 (2005).
- PCT International Preliminary Report on Patentability (Chapter I) of Dec. 7, 2009 for application PCT/US08/65563.
- PCT Search Report of Aug. 20, 2010 for application PCT/US2009/066142.
- PCT Search Report of Nov. 5, 2010 for application PCT/US2009/066141.
- PCT Search Report of Nov. 6, 2008 for application PCT/US08/65563.
- PCT Written Opinion of the International Search Authority of Aug. 20, 2010 for application PCT/US2009/066142.
- PCT Written Opinion of the International Searching Authority of Nov. 6, 2008 for application PCT/US08/65563.
- PCT Written Opinion of the International Searching Authority of Nov. 5, 2010 for application PCT/US2009/066141.
- Rastogi et al., "Footprinting of the spinach nitrite reductase gene promoter reveals the preservation of nitrate regulatory elements between fungi and higher plants," *Plant Mol Biol.*, 34(3):465-476, Abstract (1997).
- Rosenberg et al., "A Green Light for Engineered Algae: Redirecting Metabolism to Fuel a Biotechnology Revolution," *Current Opinion in Biotechnology. Tissue, Cell and Pathway Engineering*, E-Pub 19:430-436 (2008).
- Running et al., "Extracellular production of L-ascorbic acid by *Chlorella* protothecoides, *Prototheca* species, and mutants of *P. moriformis* during aerobic culturing at low pH," *Journal of Industrial Microbiology & Biotechnology*, 29:93-98 (2002).
- Son et al., "Induction and Cultures of Mountain Genseng Adventitious Roots and AFLP Analysis for Identifying Mountain Ginseng," *Biotechnol. Bioprocess Eng.*, 4:119-123 (1999).
- Takashima et al., "Further Notes on the Growth and Chlorophyll Formation of *Chlorella* Protothecoides," *Plant & Cell Physiol.*, 5:321-332 (1964).
- Tan et al., "Establishment of a Micro-Particle Bombardment Transformation System for *Dunaliella salina*," *J Microbiol.*;43(4):361-5 (2005).
- Thiry et al., "Optimizing scale-up fermentation processes," *Trends in Biotechnology*, 20: 103-105 (2002).
- Trimble et al., "Structure of Oligosaccharides on *Saccharomyces SUC2* Invertase Secreted by the Methylophilic Yeast *Pichia pastoris*," *J. Biol. Chem.*, 266(34):22807-22817 (1991).
- U.S. Appl. No. 12/131,766, Non-Final Office Action mailed Dec. 10, 2009.
- U.S. Appl. No. 12/131,766, Non-Final Office Action mailed Nov. 23, 2010.
- U.S. Appl. No. 12/131,766, Requirement for Restriction/Election mailed Aug. 17, 2010.
- U.S. Appl. No. 12/131,766, Requirement for Restriction/Election mailed Aug. 5, 2009.
- U.S. Appl. No. 12/131,773, Final Office Action mailed Mar. 21, 2011.
- U.S. Appl. No. 12/131,773, Non-Final Office Action mailed Jun. 25, 2010.
- U.S. Appl. No. 12/131,773, Non-Final Office Action mailed Dec. 15, 2009.
- U.S. Appl. No. 12/131,773, Requirement for Restriction/Election mailed Jun. 25, 2010.
- U.S. Appl. No. 12/131,793, Final Office Action mailed Mar. 30, 2010.
- U.S. Appl. No. 12/131,793, Non-Final Office Action mailed Sep. 16, 2009.
- U.S. Appl. No. 12/131,793, Requirement for Restriction/Election mailed Aug. 6, 2009.
- U.S. Appl. No. 12/131,804, Final Office Action mailed Feb. 2, 2011.
- U.S. Appl. No. 12/131,804, Non-Final Office Action mailed Mar. 3, 2010.
- U.S. Appl. No. 12/131,804, Requirement for Restriction/Election mailed Nov. 18, 2009.
- U.S. Appl. No. 12/131,804, Requirement for Restriction/Election mailed Sep. 17, 2009.
- U.S. Appl. No. 12/194,389, Final Office Action mailed Jan. 5, 2011.
- U.S. Appl. No. 12/194,389, Non-Final Office Action mailed Feb. 4, 2010.
- U.S. Appl. No. 12/194,389, Requirement for Restriction/Election mailed Oct. 5, 2010.
- U.S. Appl. No. 12/194,389, Requirement for Restriction/Election mailed Nov. 2, 2009.
- U.S. Appl. No. 12/628,144, Final Office Action mailed Nov. 16, 2010.
- U.S. Appl. No. 12/628,144, Non-Final Office Action mailed Jul. 8, 2010.
- U.S. Appl. No. 12/628,147, Final Office Action mailed Oct. 1, 2010.
- U.S. Appl. No. 12/628,147, Non-Final Office Action mailed May 25, 2010.
- U.S. Appl. No. 12/628,149, Non-Final Office Action mailed Jun. 25, 2010.
- U.S. Appl. No. 12/628,149, Non-Final Office Action mailed Sep. 16, 2010.
- U.S. Appl. No. 12/628,149, Notice of Allowance mailed Dec. 15, 2010.

- U.S. Appl. No. 12/628,150, Non-Final Office Action mailed Apr. 29, 2010.
- U.S. Appl. No. 12/628,150, Non-Final Office Action mailed Oct. 13, 2010.
- U.S. Appl. No. 12/628,150, Notice of Allowance mailed Mar. 21, 2011.
- UNIPROT Accession No. P41758. Phosphoglycerate kinase, chloroplastic. Nov. 25, 2008. [Retrieved from the Internet Aug. 12, 2010: <<http://www.uniprot.org/uniprot/P41758.txt?version=46>>].
- UNIPROT Accession No. P41758. Phosphoglycerate kinase, chloroplastic. Jul. 22, 2008. [Retrieved from the Internet Aug. 25, 2010: <<http://http://www.uniprot.org/uniprot/P41758.txt?version=44>>]; amino acids 1-60.
- Van Gerpen, "Commercial Biodiesel Production," Oilseed and Biodiesel Workshop, Billings, Montana, Jan. 9, 2008, downloaded from [http://www.deq.state.mt.us/Energy/bioenergy/Biodiesel\\_Production\\_Educ\\_Presentations/10Montana\\_Production\\_Jan\\_2008\\_JVP.pdf](http://www.deq.state.mt.us/Energy/bioenergy/Biodiesel_Production_Educ_Presentations/10Montana_Production_Jan_2008_JVP.pdf) on Mar. 10, 2010.
- Van Gerpen, *Fuel Processing Technology*, 86: 1097-1107 (2005).
- Westphal, et al., "Vipp1 Deletion of Mutant Synechocystis: A Connection Between Bacterial Phage Shock and Thylakoid Biogenesis," *Proc Natl Acad Sci U S A.*, 98(7):4243-8 (2007).
- Wirth et al., "Transforamtion of Various Species of Gram-Negative Bacteria Belonging to 11 Difference Genera by Electroporation," *Mol Gen Genet.*; 216(1):175-177 (1989).
- Wolk et al., "Construction of Shuttle Vectors Capable of Conjugative Transfer From *Escherichia coli* to Nitrogen-Fixing Filamentous Cyanobacteria," *Proc Natl Acad Sci U S A.*, 81(5):1561-1565 (1984).
- Wyman et al., "Comparative Sugar Recovery Data From Laboratory Scale Application of Leading Pretreatment Technologies to Corn Stover," *Bioresour Technol.*, 96(18):2026-2032 (2005).
- Da Silva et al., "Effect of *n*-dodecane on *Cryptocodinium cohnii* fermentations and DHA production," *J Ind Microbiol Biotechnol*, DOI 10.1007/s10295-006-0081-8, 9 pages, (2006).
- Dormann et al., "Cloning and Expression in *Escherichia coli* of a Novel Thioesterase from *Arabidopsis thaliana* Specific for Long-Chain Acyl-Acyl Carrier Proteins," *Archives of Biochemistry and Biophysics*, 316(1):612-618, 1995.
- El-Sheekh et al., "Variation of Some Nutritional Constituents and Fatty Acid Profiles of *Chlorella vulgaris* Beijerinck Grown under Auto and Heterotrophic Conditions," *International Journal of Botany*, 5(2):153-159, (2009).
- Neish et al., "Carbohydrate Nutrition of *Chlorella Vulgaris*" *Canadian Journal of Botany*, 29:68-78, (1951).
- Papanikolaou et al., "Yarrowia lipolytica as a potential producer of citric acid from raw glycerol," *Journal of Applied Microbiology*, 92:737-744, (2002).
- PCT Search Report for application PCT/US2011/038463 mailed Jan. 18, 2012.
- PCT Written Opinion of the International Searching Authority for application PCT/US2011/038463 mailed Jan. 18, 2012.
- Sud et al., "Lipid Composition and Sensitivity of *Prototheca wickerhamii* to Membrane-Active Antimicrobial Agents," *Antimicrobial Agents and Chemotherapy*, 16:486-490, (1979).
- U.S. Appl. No. 12/628,144, Final Office Action mailed Dec. 5, 2011.
- U.S. Appl. No. 12/772,170, Final Office Action mailed Feb. 21, 2012.
- U.S. Appl. No. 12/772,173, Non-Final Office Action mailed Dec. 16, 2011.
- U.S. Appl. No. 12/772,174, Non-Final Office Action mailed Nov. 29, 2011.
- U.S. Appl. No. 13/029,061, Requirement for Restriction/Election mailed Nov. 29, 2011.
- U.S. Appl. No. 13/045,500, Non-Final Office Action mailed Mar. 9, 2012.
- U.S. Appl. No. 13/073,757, Non-Final Office Action mailed Dec. 29, 2011.
- U.S. Appl. No. 13/073,757, Notice of Allowance mailed Apr. 17, 2012.

\* cited by examiner

Figure 1




Figure 2






Figure 3







Figure 6




Figure 7b




Figure 7a




Figure 7d





Figure 7c






Figure 8




Figure 9

Figure 10




Figure 11






Figure 12





Figure 13



## RENEWABLE CHEMICAL PRODUCTION FROM NOVEL FATTY ACID FEEDSTOCKS

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 12/628,149, filed Nov. 30, 2009, now U.S. Pat. No. 7,883,882, issued on Feb. 8, 2011, which claims the benefit under 35 U.S.C. 119(e) of U.S. Provisional Patent Application No. 61/118,590, filed Nov. 28, 2008, U.S. Provisional Patent Application No. 61/118,994, filed Dec. 1, 2008, U.S. Provisional Patent Application No. 61/174,357, filed Apr. 30, 2009, and U.S. Provisional Patent Application No. 61/219,525, filed Jun. 23, 2009. Each of these applications is incorporated herein by reference in its entirety for all purposes.

### REFERENCE TO A SEQUENCE LISTING

This application includes a sequence listing as shown in pages 1-163, appended hereto.

### FIELD OF THE INVENTION

The present invention relates to the production of oils, fuels, and oleochemicals made from microorganisms. In particular, the disclosure relates to oil-bearing microalgae, methods of cultivating them for the production of useful compounds, including lipids, fatty acid esters, fatty acids, aldehydes, alcohols, and alkanes, and methods and reagents for genetically altering them to improve production efficiency and alter the type and composition of the oils produced by them.

### BACKGROUND OF THE INVENTION

Fossil fuel is a general term for buried combustible geologic deposits of organic materials, formed from decayed plants and animals that have been converted to crude oil, coal, natural gas, or heavy oils by exposure to heat and pressure in the earth's crust over hundreds of millions of years. Fossil fuels are a finite, non-renewable resource.

Increased demand for energy by the global economy has also placed increasing pressure on the cost of hydrocarbons. Aside from energy, many industries, including plastics and chemical manufacturers, rely heavily on the availability of hydrocarbons as a feedstock for their manufacturing processes. Cost-effective alternatives to current sources of supply could help mitigate the upward pressure on energy and these raw material costs.

PCT Pub. No. 2008/151149 describes methods and materials for cultivating microalgae for the production of oil and particularly exemplifies the production of diesel fuel from oil produced by the microalgae *Chlorella protothecoides*. There remains a need for improved methods for producing oil in microalgae, particularly for methods that produce oils with shorter chain length and a higher degree of saturation and without pigments, with greater yield and efficiency. The present invention meets this need.

### SUMMARY OF THE INVENTION

The invention provides cells of the genus *Prototheca* comprising an exogenous gene, and in some embodiments the cell is a strain of the species *Prototheca moriformis*, *Prototheca krugani*, *Prototheca stagnora* or *Prototheca zopfii* and in other embodiment the cell has a 23S rRNA sequence with at

least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19. In some cells the exogenous gene is coding sequence and is in operable linkage with a promoter, and in some embodiments the promoter is from a gene endogenous to a species of the genus *Prototheca*. In further embodiments the coding sequence encodes a protein selected from the group consisting of a sucrose invertase, a fatty acyl-ACP thioesterase, a fatty acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty aldehyde decarbonylase, an acyl carrier protein and a protein that imparts resistance to an antibiotic. Some embodiments of a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14, including acyl-ACP thioesterases with at least 50, 60, 70, 80, or 90% amino acid identity with one or more sequences selected from the group consisting of SEQ ID NOs: 59, 61, 63 and 138-140. In further embodiments the coding sequence comprises a plastid targeting sequence from microalgae, and in some embodiments the microalgae is a species of the genus *Prototheca* or *Chlorella* as well as other genera from the family Chlorellaceae. In some embodiments the plastid targeting sequence has at least 20, 25, 35, 45, or 55% amino acid sequence identity to one or more of SEQ ID NOs: 127-133 and is capable of targeting a protein encoded by an exogenous gene not located in the plastid genome to the plastid. In other embodiments the promoter is upregulated in response to reduction or elimination of nitrogen in the culture media of the cell, such as at least a 3-fold upregulation as determined by transcript abundance in a cell of the genus *Prototheca* when the extracellular environment changes from containing at least 10 mM or 5 mM nitrogen to containing no nitrogen. In further embodiments the promoter comprises a segment of 50 or more nucleotides of one of SEQ ID NOs: 91-102. In other embodiments the cell has a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19. In other embodiments the exogenous gene is integrated into a chromosome of the cell.

In additional embodiments of cells of the invention, the cell is of the genus *Prototheca* and comprises an exogenous fatty acyl-ACP thioesterase gene and a lipid profile of at least 4% C8-C14 of total lipids of the cell, an amount of C8 that is at least 0.3% of total lipids of the cell, an amount of C10 that is at least 2% of total lipids of the cell, an amount of C12 that is at least 2% of total lipids of the cell, an amount of C14 that is at least 4% of total lipids of the cell, and an amount of C8-C14 that is 10-30%, 20-30%, or at least 10, 20, or 30% of total lipids of the cell. In some embodiments the cell further comprises an exogenous sucrose invertase gene. In some embodiments the cell is a strain of the species *Prototheca moriformis*, *Prototheca krugani*, *Prototheca stagnora* or *Prototheca zopfii*, and in other embodiment the cell has a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19. In other embodiments the exogenous fatty acyl-ACP thioesterase gene is integrated into a chromosome of the cell. Other embodiments of the invention comprise methods of making triglyceride compositions of a lipid profile of at least 4% C8-C14 w/w or area percent of the triglyceride composition, an amount of C8 that is at least 0.3% w/w or area percent, an amount of C10 that is at least 2% w/w or area percent, an amount of C12 that is at least 2% w/w or area percent, an amount of C14 that is at least 4% w/w or area percent, and an amount of C8-C14 that is 10-30%, 20-30%, or at least 10, 20, or 30% w/w or area percent. The invention also comprises methods of making triglyceride compositions comprising cultivating the foregoing cells, wherein the cells also com-

prise an exogenous gene encoding a sucrose invertase and sucrose is provided as a carbon source. In some embodiments the sucrose invertase has at least 50, 60, 70, 80, or 90% amino acid identity to one or more of SEQ ID NOs: 3, 20-29 and 90.

Embodiments of the invention include triglyceride oil compositions as well as cells containing triglyceride oil compositions comprising a lipid profile of at least 4% C8-C14 and one or more of the following attributes: 0.1-0.4 micrograms/ml total carotenoids, less than 0.4 micrograms/ml total carotenoids, less than 0.001 micrograms/ml lycopene; less than 0.02 micrograms/ml beta carotene, less than 0.02 milligrams of chlorophyll per kilogram of oil; 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil; 0.2-0.5 milligrams of total tocotrienols per gram of oil, less than 0.4 milligrams of total tocotrienols per gram of oil, 4-8 mg per 100 grams of oil of campesterol, and 40-60 mg per 100 grams of oil of stigmasterol. In some embodiments of the invention the triglyceride oil compositions have a lipid profile of at least 4% C8-C14 w/w or area percent of the triglyceride composition, an amount of C8 that is at least 0.3% w/w or area percent, an amount of C10 that is at least 2% w/w or area percent, an amount of C12 that is at least 2% w/w or area percent, an amount of C14 that is at least 4% w/w or area percent, and an amount of C8-C14 that is 10-30%, 20-30%, or at least 10, 20, or 30% w/w or area percent. In other embodiments the triglyceride oil composition is blended with at least one other composition selected from the group consisting of soy, rapeseed, canola, palm, palm kernel, coconut, corn, waste vegetable, Chinese tallow, olive, sunflower, cotton seed, chicken fat, beef tallow, porcine tallow, microalgae, macroalgae, *Cuphea*, flax, peanut, choice white grease, lard, *Camelina sativa*, mustard seed, cashew nut, oats, lupine, kenaf, *calendula*, hemp, coffee, linseed (flax), hazelnut, *euphorbia*, pumpkin seed, coriander, *camellia*, sesame, safflower, rice, tung tree, cocoa, copra, pium poppy, castor beans, pecan, jojoba, *jatropha*, *macadamia*, Brazil nuts, avocado, petroleum, or a distillate fraction of any of the preceding oils.

Methods of the invention also include processing the aforementioned oils of by performing one or more chemical reactions from the list consisting of transesterification, hydrogenation, hydrocracking, deoxygenation, isomerization, interesterification, hydroxylation, hydrolysis to yield free fatty acids, and saponification. The invention also includes hydrocarbon fuels made from hydrogenation and isomerization of the aforementioned oils and fatty acid alkyl esters made from transesterification of the aforementioned oils. In some embodiments the hydrocarbon fuel is made from triglyceride isolated from cells of the genus *Prototheca* wherein the ASTM D86 T10-T90 distillation range is at least 25° C. In other embodiments the fatty acid alkyl ester fuel is made from triglyceride isolated from cells of the genus *Prototheca*, wherein the composition has an ASTM D6751 A1 cold soak time of less than 120 seconds.

The invention also includes composition comprising (a) polysaccharide comprising one or more monosaccharides from the list consisting of 20-30 mole percent galactose; 55-65 mole percent glucose; and 5-15 mole percent mannose; (b) protein; and (c) DNA comprising a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19; and (d) an exogenous gene. In some embodiments the exogenous gene is selected from a sucrose invertase and a fatty acyl-ACP thioesterase, and in further embodiments the composition further comprises lipid with a lipid profile of at least 4% C8-C14. In other embodiments the composition is formulated for consumption as an animal feed.

The invention includes recombinant nucleic acids encoding promoters that are upregulated in response to reduction or elimination of nitrogen in the culture media of a cell of the genus *Prototheca*, such as at least a 3-fold upregulation as determined by transcript abundance when the extracellular environment changes from containing at least 10 mM or 5 mM nitrogen to containing no nitrogen. In some embodiments the recombinant nucleic acid comprises a segment of 50 or more nucleotides of one of SEQ ID NOs: 91-102. The invention also includes nucleic acid vectors comprising an expression cassette comprising (a) a promoter that is active in a cell of the genus *Prototheca*; and (b) a coding sequence in operable linkage with the promoter wherein the coding sequence contains the most or second most preferred codons of Table 1 for at least 20, 30, 40, 50, 60, or 80% of the codons of the coding sequence. In some vectors the coding sequence comprises a plastid targeting sequence in-frame with a fatty acyl-ACP thioesterase, including thioesterase that have hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14. Some vectors include plastid targeting sequences that encode peptides that are capable of targeting a protein to the plastid of a cell of the genus *Prototheca*, including those from microalgae and those wherein the plastid targeting sequence has at least 20, 25, 35, 45, or 55% amino acid sequence identity to one or more of SEQ ID NOs. 127-133 and is capable of targeting a protein to the plastid of a cell of the genus *Prototheca*. Additional vectors of the invention comprise nucleic acid sequences endogenous to the nuclear genome of a cell of the genus *Prototheca*, wherein the sequence is at least 200 nucleotides long, and some vectors comprise first and second nucleic acid sequences endogenous to the nuclear genome of a cell of the genus *Prototheca*, wherein the first and second sequences (a) are each at least 200 nucleotides long; (b) flank the expression cassette; and (c) are located on the same *Prototheca* chromosome no more than 5, 10, 15, 20, and 50 kB apart.

The invention also includes a recombinant nucleic acid with at least 80, 90, 95 or 98% nucleotide identity to one or both of SEQ ID NOs: 134-135 and a recombinant nucleic acid encoding a protein with at least 80, 90, 95 or 98% amino acid identity to one or both of SEQ ID NOs: 136-137.

The invention also comprises methods of producing triglyceride compositions, comprising (a) culturing a population of cells of the genus *Prototheca* in the presence of a fixed carbon source, wherein: (i) the cells contain an exogenous gene; (ii) the cells accumulate at least 10, 20, 30, 40, 60, or 70% of their dry cell weight as lipid; and (iii) the fixed carbon source is selected from the group consisting of sorghum and depolymerized cellulosic material; and (b) isolating lipid components from the cultured microorganisms. In some embodiments the fixed carbon source is depolymerized cellulosic material selected from the group consisting of corn stover, *Miscanthus*, forage sorghum, sugar beet pulp and sugar cane bagasse, optionally that has been subjected to washing with water prior to the culturing step. In some methods the fixed carbon source is depolymerized cellulosic material and the glucose level of the depolymerized cellulosic material is concentrated to a level of at least 300 g/liter, at least 400 g/liter, at least 500 g/liter, or at least 600 g/liter of prior to the culturing step and is fed to the culture over time as the cells grow and accumulate lipid. In some methods the exogenous gene encodes a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14, and in some methods the triglyceride has a lipid profile of at least 4% C8-C14 and one or more of the following attributes: 0.1-0.4 micrograms/ml total carotenoids; less than 0.02 milligrams of

chlorophyll per kilogram of oil; 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil; 0.2-0.5 milligrams of total tocotrienols per gram of oil, 4-8 mg per 100 grams of oil of campesterol, and 40-60 mg per 100 grams of oil of stigmasterol.

Further methods of the invention include producing a triglyceride composition, comprising: (a) culturing a population of microorganisms in the presence of depolymerized cellulosic material, wherein: (i) the depolymerized cellulosic material is subjected to washing with water prior to the culturing step; (ii) the cells accumulate at least 10, 20, 30, 40, 60, or 70% of their dry cell weight as lipid; and (iii) the depolymerized cellulosic material is concentrated to at least 300, 400, 500, or 600 g/liter of glucose prior to the cultivation step; (iv) the microorganisms are cultured in a fed-batch reaction in which depolymerized cellulosic material of at least 300, 400, 500, or 600 g/liter of glucose is fed to the microorganisms; and (b) isolating lipid components from the cultured microorganisms. In some embodiments the fixed carbon source is depolymerized cellulosic material selected from the group consisting of corn stover, *Miscanthus*, forage sorghum, sugar beet pulp and sugar cane bagasse. In further embodiments the microorganisms are a species of the genus *Prototheca* and contain an exogenous gene, including a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14. A further method of the invention comprises manufacturing triglyceride oil comprising cultivating a cell that has a 23S rRNA sequence with at least 90 or 96% nucleotide identity to SEQ ID NO: 30 in the presence of sucrose as a carbon source.

The invention also includes methods of manufacturing a chemical comprising performing one or more chemical reactions from the list consisting of transesterification, hydrogenation, hydrocracking, deoxygenation, isomerization, interesterification, hydroxylation, hydrolysis, and saponification on a triglyceride oil, wherein the oil has a lipid profile of at least 4% C8-C14 and one or more of the following attributes: 0.1-0.4 micrograms/ml total carotenoids; less than 0.02 milligrams of chlorophyll per kilogram of oil; 0.10-0.60 milligrams of gamma tocopherol per 100 grams of oil; 0.1-0.5 milligrams of total tocotrienols per gram of oil, 1-8 mg per 100 grams of oil of campesterol, and 10-60 mg per 100 grams of oil of stigmasterol. Some methods are performed by manufacturing the oil by cultivating a cell of the genus *Prototheca* that comprises an exogenous fatty acyl-ACP thioesterase gene that encodes a fatty acyl-ACP thioesterase having hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14. In some methods the hydrolysis reaction is selected from the group consisting of saponification, acid hydrolysis, alkaline hydrolysis, enzymatic hydrolysis, catalytic hydrolysis, and hot-compressed water hydrolysis, including a catalytic hydrolysis reaction wherein the oil is split into glycerol and fatty acids. In further methods the fatty acids undergo an amination reaction to produce fatty nitrogen compounds or an ozonolysis reaction to produce mono- and dibasic-acids. In some embodiments the oil undergoes a triglyceride splitting method selected from the group consisting of enzymatic splitting and pressure splitting. In some methods a condensation reaction follows the hydrolysis reaction. Other methods include performing a hydroprocessing reaction on the oil, optionally wherein the product of the hydroprocessing reaction undergoes a deoxygenation reaction or a condensation reaction prior to or simultaneous with the hydroprocessing reaction. Some methods additionally include a gas removal reaction. Additional methods include processing the aforementioned oils by perform-

ing a deoxygenation reaction selected from the group consisting of: a hydrogenolysis reaction, hydrogenation, a consecutive hydrogenation-hydrogenolysis reaction, a consecutive hydrogenolysis-hydrogenation reaction, and a combined hydrogenation-hydrogenolysis reaction. In some methods a condensation reaction follows the deoxygenation reaction. Other methods include performing an esterification reaction on the aforementioned oils, optionally an interesterification reaction or a transesterification reaction. Other methods include performing a hydroxylation reaction on the aforementioned oils, optionally wherein a condensation reaction follows the hydroxylation reaction.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1 and 2 illustrate the growth curves of *Prototheca* species and *Chlorella luteoviridis* strain SAG 2214 grown on sorghum as the carbon source.

FIG. 3 shows time course growth of SAG 2214 on glucose and sucrose.

FIG. 4 shows maps of the cassettes used in *Prototheca* transformations, as described in Example 3.

FIG. 5 shows the results of Southern blot analysis on three transformants of UTEX strain 1435, as described in Example 3.

FIG. 6 shows a schematic of the codon optimized and non-codon optimized *suc2* (yeast sucrose invertase (*yInv*)) transgene construct. The relevant restriction cloning sites are indicated and arrows indicate the direction of transcription.

FIG. 7a shows the results of *Prototheca moriformis* grown on cellulosic-derived sugars (corn stover, beet pulp, sorghum cane, *Miscanthus* and glucose control). Growth is expressed in optical density measurements ( $A_{750}$  readings).

FIG. 7b shows the results of growth experiments using *Prototheca moriformis* using different levels of corn stover-derived cellulosic sugar as compared to glucose/xylose control.

FIG. 7c shows the impact that xylose has on the lipid production in *Prototheca* cultures.

FIG. 7d shows the impact of salt concentration ( $\text{Na}_2\text{SO}_4$ ) and antifoam on the growth (in dry cell weight (DCW)) of *Prototheca*.

FIG. 8 shows the impact of hydrothermal treatment of various cellulosic materials (sugar cane bagasse, sorghum cane, *Miscanthus* and beet pulp) and the resulting sugar stream on the growth of *Prototheca*.

FIG. 9 shows decreasing levels of hydroxymethyl furfurals (HMF) and furfurals in cellulosic biomass (sugar cane bagasse, sorghum cane, *Miscanthus* and beet pulp) after repeated cycles of hydrothermal treatment.

FIG. 10 shows a schematic of a saccharification process of cellulosic materials to generate sugar streams suitable for use in heterotrophic oil production in a fermentor.

FIG. 11 shows decreasing levels of HMF and furfurals in exploded sugar cane bagasse after repeated cycles of hydrothermal treatment.

FIG. 12 shows a schematic of thioesterase constructs used in *Prototheca* transformations. The heterologous beta-tubulin (driving  $\text{Neo}^R$ ) and glutamate dehydrogenase promoters are derived from *Chlamydomonas reinhardtii* and *Chlorella sorokiniana*, respectively. The nitrate reductase 3'UTR was derived from *Chlorella vulgaris*. The relevant restriction cloning sites are indicated and arrows indicate the direction of transcription.

FIG. 13 shows a chromatogram of renewable diesel produced from *Prototheca* triglyceride oil.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention arises from the discovery that *Prototheca* and certain related microorganisms have unexpectedly advantageous properties for the production of oils, fuels, and other hydrocarbon or lipid compositions economically and in large quantities, as well as from the discovery of methods and reagents for genetically altering these microorganisms to improve these properties. The oils produced by these microorganisms can be used in the transportation fuel, petrochemical, and/or food and cosmetic industries, among other applications. Transesterification of lipids yields long-chain fatty acid esters useful as biodiesel. Other enzymatic and chemical processes can be tailored to yield fatty acids, aldehydes, alcohols, alkanes, and alkenes. In some applications, renewable diesel, jet fuel, or other hydrocarbon compounds are produced. The present invention also provides methods of cultivating microalgae for increased productivity and increased lipid yield, and/or for more cost-effective production of the compositions described herein.

This detailed description of the invention is divided into sections for the convenience of the reader. Section 1 provides definitions of terms used herein. Section 2 provides a description of culture conditions useful in the methods of the invention. Section 3 provides a description of genetic engineering methods and materials. Section 4 provides a description of genetic engineering of *Prototheca* to enable sucrose utilization. Section 5 provides a description of genetic engineering of *Prototheca* to modify lipid biosynthesis. Section 6 describes methods for making fuels and chemicals. Section 7 discloses examples and embodiments of the invention. The detailed description of the invention is followed by examples that illustrate the various aspects and embodiments of the invention.

#### I. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton et al., *Dictionary of Microbiology and Molecular Biology* (2nd ed. 1994); *The Cambridge Dictionary of Science and Technology* (Walker ed., 1988); *The Glossary of Genetics*, 5th Ed., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, *The Harper Collins Dictionary of Biology* (1991). As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

“Active in microalgae” refers to a nucleic acid that is functional in microalgae. For example, a promoter that has been used to drive an antibiotic resistance gene to impart antibiotic resistance to a transgenic microalgae is active in microalgae.

“Acyl carrier protein” or “ACP” is a protein that binds a growing acyl chain during fatty acid synthesis as a thiol ester at the distal thiol of the 4'-phosphopantetheine moiety and comprises a component of the fatty acid synthase complex.

“Acyl-CoA molecule” or “acyl-CoA” is a molecule comprising an acyl moiety covalently attached to coenzyme A through a thiol ester linkage at the distal thiol of the 4'-phosphopantetheine moiety of coenzyme A.

“Area Percent” refers to the area of peaks observed using FAME GC/FID detection methods in which every fatty acid in the sample is converted into a fatty acid methyl ester

(FAME) prior to detection. For example, a separate peak is observed for a fatty acid of 14 carbon atoms with no unsaturation (C14:0) compared to any other fatty acid such as C14:1. The peak area for each class of FAME is directly proportional to its percent composition in the mixture and is calculated based on the sum of all peaks present in the sample (i.e. [area under specific peak/total area of all measured peaks]×100). When referring to lipid profiles of oils and cells of the invention, “at least 4% C8-C14” means that at least 4% of the total fatty acids in the cell or in the extracted glycerolipid composition have a chain length that includes 8, 10, 12 or 14 carbon atoms.

“Axenic” is a culture of an organism free from contamination by other living organisms.

“Biodiesel” is a biologically produced fatty acid alkyl ester suitable for use as a fuel in a diesel engine.

“Biomass” is material produced by growth and/or propagation of cells. Biomass may contain cells and/or intracellular contents as well as extracellular material, includes, but is not limited to, compounds secreted by a cell.

“Bioreactor” is an enclosure or partial enclosure in which cells are cultured, optionally in suspension.

“Catalyst” is an agent, such as a molecule or macromolecular complex, capable of facilitating or promoting a chemical reaction of a reactant to a product without becoming a part of the product. A catalyst increases the rate of a reaction, after which, the catalyst may act on another reactant to form the product. A catalyst generally lowers the overall activation energy required for the reaction such that it proceeds more quickly or at a lower temperature. Thus, a reaction equilibrium may be more quickly attained. Examples of catalysts include enzymes, which are biological catalysts; heat, which is a non-biological catalyst; and metals used in fossil oil refining processes.

“Cellulosic material” is the product of digestion of cellulose, including glucose and xylose, and optionally additional compounds such as disaccharides, oligosaccharides, lignin, furfurals and other compounds. Nonlimiting examples of sources of cellulosic material include sugar cane bagasses, sugar beet pulp, corn stover, wood chips, sawdust and switchgrass.

“Co-culture”, and variants thereof such as “co-cultivate” and “co-ferment”, refer to the presence of two or more types of cells in the same bioreactor. The two or more types of cells may both be microorganisms, such as microalgae, or may be a microalgal cell cultured with a different cell type. The culture conditions may be those that foster growth and/or propagation of the two or more cell types or those that facilitate growth and/or proliferation of one, or a subset, of the two or more cells while maintaining cellular growth for the remainder.

“Cofactor” is any molecule, other than the substrate, required for an enzyme to carry out its enzymatic activity.

“Complementary DNA” or “cDNA” is a DNA copy of mRNA, usually obtained by reverse transcription of messenger RNA (mRNA) or amplification (e.g., via polymerase chain reaction (“PCR”)).

“Cultivated”, and variants thereof such as “cultured” and “fermented”, refer to the intentional fostering of growth (increases in cell size, cellular contents, and/or cellular activity) and/or propagation (increases in cell numbers via mitosis) of one or more cells by use of selected and/or controlled conditions. The combination of both growth and propagation may be termed proliferation. Examples of selected and/or controlled conditions include the use of a defined medium (with known characteristics such as pH, ionic strength, and carbon source), specified temperature, oxygen tension, carbon diox-

ide levels, and growth in a bioreactor. Cultivate does not refer to the growth or propagation of microorganisms in nature or otherwise without human intervention; for example, natural growth of an organism that ultimately becomes fossilized to produce geological crude oil is not cultivation.

“Cytolysis” is the lysis of cells in a hypotonic environment. Cytolysis is caused by excessive osmosis, or movement of water, towards the inside of a cell (hyperhydration). The cell cannot withstand the osmotic pressure of the water inside, and so it explodes.

“Delipidated meal” and “delipidated microbial biomass” is microbial biomass after oil (including lipids) has been extracted or isolated from it, either through the use of mechanical (i.e., exerted by an expeller press) or solvent extraction or both. Delipidated meal has a reduced amount of oil/lipids as compared to before the extraction or isolation of oil/lipids from the microbial biomass but does contain some residual oil/lipid.

“Expression vector” or “expression construct” or “plasmid” or “recombinant DNA construct” refer to a nucleic acid that has been generated via human intervention, including by recombinant means or direct chemical synthesis, with a series of specified nucleic acid elements that permit transcription and/or translation of a particular nucleic acid in a host cell. The expression vector can be part of a plasmid, virus, or nucleic acid fragment. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

“Exogenous gene” is a nucleic acid that codes for the expression of an RNA and/or protein that has been introduced (“transformed”) into a cell. A transformed cell may be referred to as a recombinant cell, into which additional exogenous gene(s) may be introduced. The exogenous gene may be from a different species (and so heterologous), or from the same species (and so homologous), relative to the cell being transformed. Thus, an exogenous gene can include a homologous gene that occupies a different location in the genome of the cell or is under different control, relative to the endogenous copy of the gene. An exogenous gene may be present in more than one copy in the cell. An exogenous gene may be maintained in a cell as an insertion into the genome or as an episomal molecule.

“Exogenously provided” refers to a molecule provided to the culture media of a cell culture.

“Expeller pressing” is a mechanical method for extracting oil from raw materials such as soybeans and rapeseed. An expeller press is a screw type machine, which presses material through a caged barrel-like cavity. Raw materials enter one side of the press and spent cake exits the other side while oil seeps out between the bars in the cage and is collected. The machine uses friction and continuous pressure from the screw drives to move and compress the raw material. The oil seeps through small openings that do not allow solids to pass through. As the raw material is pressed, friction typically causes it to heat up.

“Fatty acyl-ACP thioesterase” is an enzyme that catalyzes the cleavage of a fatty acid from an acyl carrier protein (ACP) during lipid synthesis.

“Fatty acyl-CoA/aldehyde reductase” is an enzyme that catalyzes the reduction of an acyl-CoA molecule to a primary alcohol.

“Fatty acyl-CoA reductase” is an enzyme that catalyzes the reduction of an acyl-CoA molecule to an aldehyde.

“Fatty aldehyde decarboxylase” is an enzyme that catalyzes the conversion of a fatty aldehyde to an alkane.

“Fatty aldehyde reductase” is an enzyme that catalyzes the reduction of an aldehyde to a primary alcohol.

“Fixed carbon source” is a molecule(s) containing carbon, typically an organic molecule, that is present at ambient temperature and pressure in solid or liquid form in a culture media that can be utilized by a microorganism cultured therein.

“Homogenate” is biomass that has been physically disrupted.

“Hydrocarbon” is (a) a molecule containing only hydrogen and carbon atoms wherein the carbon atoms are covalently linked to form a linear, branched, cyclic, or partially cyclic backbone to which the hydrogen atoms are attached. The molecular structure of hydrocarbon compounds varies from the simplest, in the form of methane (CH<sub>4</sub>), which is a constituent of natural gas, to the very heavy and very complex, such as some molecules such as asphaltenes found in crude oil, petroleum, and bitumens. Hydrocarbons may be in gaseous, liquid, or solid form, or any combination of these forms, and may have one or more double or triple bonds between adjacent carbon atoms in the backbone. Accordingly, the term includes linear, branched, cyclic, or partially cyclic alkanes, alkenes, lipids, and paraffin. Examples include propane, butane, pentane, hexane, octane, and squalene.

“Hydrogen:carbon ratio” is the ratio of hydrogen atoms to carbon atoms in a molecule on an atom-to-atom basis. The ratio may be used to refer to the number of carbon and hydrogen atoms in a hydrocarbon molecule. For example, the hydrocarbon with the highest ratio is methane CH<sub>4</sub> (4:1).

“Hydrophobic fraction” is the portion, or fraction, of a material that is more soluble in a hydrophobic phase in comparison to an aqueous phase. A hydrophobic fraction is substantially insoluble in water and usually non-polar.

“Increase lipid yield” refers to an increase in the productivity of a microbial culture by, for example, increasing dry weight of cells per liter of culture, increasing the percentage of cells that constitute lipid, or increasing the overall amount of lipid per liter of culture volume per unit time.

“Inducible promoter” is a promoter that mediates transcription of an operably linked gene in response to a particular stimulus.

“In operable linkage” is a functional linkage between two nucleic acid sequences, such a control sequence (typically a promoter) and the linked sequence (typically a sequence that encodes a protein, also called a coding sequence). A promoter is in operable linkage with an exogenous gene if it can mediate transcription of the gene.

“In situ” means “in place” or “in its original position”.

“Limiting concentration of a nutrient” is a concentration of a compound in a culture that limits the propagation of a cultured organism. A “non-limiting concentration of a nutrient” is a concentration that supports maximal propagation during a given culture period. Thus, the number of cells produced during a given culture period is lower in the presence of a limiting concentration of a nutrient than when the nutrient is non-limiting. A nutrient is said to be “in excess” in a culture, when the nutrient is present at a concentration greater than that which supports maximal propagation.

“Lipase” is a water-soluble enzyme that catalyzes the hydrolysis of ester bonds in water-insoluble, lipid substrates. Lipases catalyze the hydrolysis of lipids into glycerols and fatty acids.

“Lipid modification enzyme” refers to an enzyme that alters the covalent structure of a lipid. Examples of lipid modification enzymes include a lipase, a fatty acyl-ACP thioesterase, a fatty acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, and a fatty aldehyde decarboxylase.

“Lipid pathway enzyme” is any enzyme that plays a role in lipid metabolism, i.e., either lipid synthesis, modification, or degradation, and any proteins that chemically modify lipids, as well as carrier proteins.

“Lipids” are a class of molecules that are soluble in non-polar solvents (such as ether and chloroform) and are relatively or completely insoluble in water. Lipid molecules have these properties, because they consist largely of long hydrocarbon tails which are hydrophobic in nature. Examples of lipids include fatty acids (saturated and unsaturated); glycerides or glycerolipids (such as monoglycerides, diglycerides, triglycerides or neutral fats, and phosphoglycerides or glycerophospholipids); nonglycerides (sphingolipids, sterol lipids including cholesterol and steroid hormones, prenol lipids including terpenoids, fatty alcohols, waxes, and polyketides); and complex lipid derivatives (sugar-linked lipids, or glycolipids, and protein-linked lipids). “Fats” are a subgroup of lipids called “triacylglycerides.”

“Lysate” is a solution containing the contents of lysed cells.

“Lysis” is the breakage of the plasma membrane and optionally the cell wall of a biological organism sufficient to release at least some intracellular content, often by mechanical, viral or osmotic mechanisms that compromise its integrity.

“Lysing” is disrupting the cellular membrane and optionally the cell wall of a biological organism or cell sufficient to release at least some intracellular content.

“Microalgae” is a eukaryotic microbial organism that contains a chloroplast or plastid, and optionally that is capable of performing photosynthesis, or a prokaryotic microbial organism capable of performing photosynthesis. Microalgae include obligate photoautotrophs, which cannot metabolize a fixed carbon source as energy, as well as heterotrophs, which can live solely off of a fixed carbon source. Microalgae include unicellular organisms that separate from sister cells shortly after cell division, such as *Chlamydomonas*, as well as microbes such as, for example, *Volvox*, which is a simple multicellular photosynthetic microbe of two distinct cell types. Microalgae include cells such as *Chlorella*, *Dunaliella*, and *Prototheca*. Microalgae also include other microbial photosynthetic organisms that exhibit cell-cell adhesion, such as *Agmenellum*, *Anabaena*, and *Pyrobotrys*. Microalgae also include obligate heterotrophic microorganisms that have lost the ability to perform photosynthesis, such as certain dinoflagellate algae species and species of the genus *Prototheca*.

“Microorganism” and “microbe” are microscopic unicellular organisms.

“Naturally co-expressed” with reference to two proteins or genes means that the proteins or their genes are co-expressed naturally in a tissue or organism from which they are derived, e.g., because the genes encoding the two proteins are under the control of a common regulatory sequence or because they are expressed in response to the same stimulus.

“Osmotic shock” is the rupture of cells in a solution following a sudden reduction in osmotic pressure. Osmotic shock is sometimes induced to release cellular components of such cells into a solution.

“Polysaccharide-degrading enzyme” is any enzyme capable of catalyzing the hydrolysis, or saccharification, of any polysaccharide. For example, cellulases catalyze the hydrolysis of cellulose.

“Polysaccharides” or “glycans” are carbohydrates made up of monosaccharides joined together by glycosidic linkages. Cellulose is a polysaccharide that makes up certain plant cell walls. Cellulose can be depolymerized by enzymes to yield

monosaccharides such as xylose and glucose, as well as larger disaccharides and oligosaccharides.

“Promoter” is a nucleic acid control sequence that directs transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription.

“Recombinant” is a cell, nucleic acid, protein or vector, that has been modified due to the introduction of an exogenous nucleic acid or the alteration of a native nucleic acid. Thus, e.g., recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes differently than those genes are expressed by a non-recombinant cell. A “recombinant nucleic acid” is a nucleic acid originally formed in vitro, in general, by the manipulation of nucleic acid, e.g., using polymerases and endonucleases, or otherwise is in a form not normally found in nature. Recombinant nucleic acids may be produced, for example, to place two or more nucleic acids in operable linkage. Thus, an isolated nucleic acid or an expression vector formed in vitro by ligating DNA molecules that are not normally joined in nature, are both considered recombinant for the purposes of this invention. Once a recombinant nucleic acid is made and introduced into a host cell or organism, it may replicate using the in vivo cellular machinery of the host cell; however, such nucleic acids, once produced recombinantly, although subsequently replicated intracellularly, are still considered recombinant for purposes of this invention. Similarly, a “recombinant protein” is a protein made using recombinant techniques, i.e., through the expression of a recombinant nucleic acid.

“Renewable diesel” is a mixture of alkanes (such as C10:0, C12:0, C14:0, C16:0 and C18:0) produced through hydrogenation and deoxygenation of lipids.

“Saccharification” is a process of converting biomass, usually cellulosic or lignocellulosic biomass, into monomeric sugars, such as glucose and xylose. “Saccharified” or “depolymerized” cellulosic material or biomass refers to cellulosic material or biomass that has been converted into monomeric sugars through saccharification.

“Sonication” is a process of disrupting biological materials, such as a cell, by use of sound wave energy.

“Species of furfural” is 2-furancarboxaldehyde or a derivative that retains the same basic structural characteristics.

“Stover” is the dried stalks and leaves of a crop remaining after a grain has been harvested.

“Sucrose utilization gene” is a gene that, when expressed, aids the ability of a cell to utilize sucrose as an energy source. Proteins encoded by a sucrose utilization gene are referred to herein as “sucrose utilization enzymes” and include sucrose transporters, sucrose invertases, and hexokinases such as glucokinases and fructokinases.

## II. CULTIVATION

The present invention generally relates to cultivation of *Prototheca* strains, particularly recombinant *Prototheca* strains, for the production of lipid. For the convenience of the reader, this section is subdivided into subsections. Subsection 1 describes *Prototheca* species and strains and how to identify new *Prototheca* species and strains and related microalgae by genomic DNA comparison. Subsection 2 describes bioreactors useful for cultivation. Subsection 3 describes media for



cultivation. Subsection 4 describes oil production in accordance with illustrative cultivation methods of the invention.

#### 1. *Prototheca* Species and Strains

*Prototheca* is a remarkable microorganism for use in the production of lipid, because it can produce high levels of lipid, particularly lipid suitable for fuel production. The lipid produced by *Prototheca* has hydrocarbon chains of shorter chain length and a higher degree of saturation than that produced by other microalgae. Moreover, *Prototheca* lipid is generally free of pigment (low to undetectable levels of chlorophyll and certain carotenoids) and in any event contains much less pigment than lipid from other microalgae. Moreover, recombinant *Prototheca* cells provided by the invention can be used to produce lipid in greater yield and efficiency, and with reduced cost, relative to the production of lipid from other microorganisms. Illustrative *Prototheca* strains for use in the methods of the invention include In addition, this microalgae grows heterotrophically and can be genetically engineered as *Prototheca wickerhamii*, *Prototheca stagnora* (including UTEX 327), *Prototheca portoricensis*, *Prototheca moriformis* (including UTEX strains 1441, 1435), and *Prototheca zopfii*. Species of the genus *Prototheca* are obligate heterotrophs.

Species of *Prototheca* for use in the invention can be identified by amplification of certain target regions of the genome. For example, identification of a specific *Prototheca* species or strain can be achieved through amplification and sequencing of nuclear and/or chloroplast DNA using primers and methodology using any region of the genome, for example using the methods described in Wu et al., *Bot. Bull. Acad. Sin.* (2001) 42:115-121 Identification of *Chlorella* spp. isolates using ribosomal DNA sequences. Well established methods of phylogenetic analysis, such as amplification and sequencing of ribosomal internal transcribed spacer (ITS1 and ITS2 rDNA), 23S rRNA, 18S rRNA, and other conserved genomic regions can be used by those skilled in the art to identify species of not only *Prototheca*, but other hydrocarbon and lipid producing organisms with similar lipid profiles and production capability. For examples of methods of identification and classification of algae also see for example *Genetics*, 2005 August; 170(4):1601-10 and *RNA*, 2005 April; 11(4):361-4.

Thus, genomic DNA comparison can be used to identify suitable species of microalgae to be used in the present invention. Regions of conserved genomic DNA, such as but not limited to DNA encoding for 23S rRNA, can be amplified from microalgal species and compared to consensus sequences in order to screen for microalgal species that are taxonomically related to the preferred microalgae used in the present invention. Examples of such DNA sequence comparison for species within the *Prototheca* genus are shown below. Genomic DNA comparison can also be useful to identify microalgal species that have been misidentified in a strain collection. Often a strain collection will identify species of microalgae based on phenotypic and morphological characteristics. The use of these characteristics may lead to miscategorization of the species or the genus of a microalgae. The use of genomic DNA comparison can be a better method of categorizing microalgae species based on their phylogenetic relationship.

Microalgae for use in the present invention typically have genomic DNA sequences encoding for 23S rRNA that have at least 99%, least 95%, at least 90%, or at least 85% nucleotide identity to at least one of the sequences listed in SEQ ID NOS: 11-19.

For sequence comparison to determine percent nucleotide or amino acid identity, typically one sequence acts as a ref-

erence sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally Ausubel et al., supra).

Another example algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (at the web address www.ncbi.nlm.nih.gov). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., supra.). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. For identifying whether a nucleic acid or polypeptide is within the scope of the invention, the default parameters of the BLAST programs are suitable. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a word length (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix. The TBLASTN program (using protein sequence for nucleotide sequence) uses as defaults a word length (W) of 3, an expectation (E) of 10, and a BLOSUM 62 scoring matrix. (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)).

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a refer-

ence sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

Other considerations affecting the selection of microorganisms for use in the invention include, in addition to production of suitable lipids or hydrocarbons for production of oils, fuels, and oleochemicals: (1) high lipid content as a percentage of cell weight; (2) ease of growth; (3) ease of genetic engineering; and (4) ease of biomass processing. In particular embodiments, the wild-type or genetically engineered microorganism yields cells that are at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, or at least 70% or more lipid. Preferred organisms grow heterotrophically (on sugars in the absence of light).

## 2. Bioreactor

Micromicroorganisms are cultured both for purposes of conducting genetic manipulations and for production of hydrocarbons (e.g., lipids, fatty acids, aldehydes, alcohols, and alkanes). The former type of culture is conducted on a small scale and initially, at least, under conditions in which the starting microorganism can grow. Culture for purposes of hydrocarbon production is usually conducted on a large scale (e.g., 10,000 L, 40,000 L, 100,000 L or larger bioreactors) in a bioreactor. *Prototheca* are typically cultured in the methods of the invention in liquid media within a bioreactor. Typically, the bioreactor does not allow light to enter.

The bioreactor or fermentor is used to culture microalgal cells through the various phases of their physiological cycle. Bioreactors offer many advantages for use in heterotrophic growth and propagation methods. To produce biomass for use in food, microalgae are preferably fermented in large quantities in liquid, such as in suspension cultures as an example. Bioreactors such as steel fermentors can accommodate very large culture volumes (40,000 liter and greater capacity bioreactors are used in various embodiments of the invention). Bioreactors also typically allow for the control of culture conditions such as temperature, pH, oxygen tension, and carbon dioxide levels. For example, bioreactors are typically configurable, for example, using ports attached to tubing, to allow gaseous components, like oxygen or nitrogen, to be bubbled through a liquid culture. Other culture parameters, such as the pH of the culture media, the identity and concentration of trace elements, and other media constituents can also be more readily manipulated using a bioreactor.

Bioreactors can be configured to flow culture media through the bioreactor throughout the time period during which the microalgae reproduce and increase in number. In some embodiments, for example, media can be infused into the bioreactor after inoculation but before the cells reach a desired density. In other instances, a bioreactor is filled with culture media at the beginning of a culture, and no more culture media is infused after the culture is inoculated. In other words, the microalgal biomass is cultured in an aqueous medium for a period of time during which the microalgae reproduce and increase in number; however, quantities of aqueous culture medium are not flowed through the bioreactor throughout the time period. Thus in some embodiments, aqueous culture medium is not flowed through the bioreactor after inoculation.

Bioreactors equipped with devices such as spinning blades and impellers, rocking mechanisms, stir bars, means for pressurized gas infusion can be used to subject microalgal cultures to mixing. Mixing may be continuous or intermittent. For example, in some embodiments, a turbulent flow regime of gas entry and media entry is not maintained for reproduc-

tion of microalgae until a desired increase in number of said microalgae has been achieved.

Bioreactor ports can be used to introduce, or extract, gases, solids, semisolids, and liquids, into the bioreactor chamber containing the microalgae. While many bioreactors have more than one port (for example, one for media entry, and another for sampling), it is not necessary that only one substance enter or leave a port. For example, a port can be used to flow culture media into the bioreactor and later used for sampling, gas entry, gas exit, or other purposes. Preferably, a sampling port can be used repeatedly without altering compromising the axenic nature of the culture. A sampling port can be configured with a valve or other device that allows the flow of sample to be stopped and started or to provide a means of continuous sampling. Bioreactors typically have at least one port that allows inoculation of a culture, and such a port can also be used for other purposes such as media or gas entry.

Bioreactors ports allow the gas content of the culture of microalgae to be manipulated. To illustrate, part of the volume of a bioreactor can be gas rather than liquid, and the gas inlets of the bioreactor to allow pumping of gases into the bioreactor. Gases that can be beneficially pumped into a bioreactor include air, air/CO<sub>2</sub> mixtures, noble gases, such as argon, and other gases. Bioreactors are typically equipped to enable the user to control the rate of entry of a gas into the bioreactor. As noted above, increasing gas flow into a bioreactor can be used to increase mixing of the culture.

Increased gas flow affects the turbidity of the culture as well. Turbulence can be achieved by placing a gas entry port below the level of the aqueous culture media so that gas entering the bioreactor bubbles to the surface of the culture. One or more gas exit ports allow gas to escape, thereby preventing pressure buildup in the bioreactor. Preferably a gas exit port leads to a "one-way" valve that prevents contaminating microorganisms from entering the bioreactor.

## 3. Media

Microalgal culture media typically contains components such as a fixed nitrogen source, a fixed carbon source, trace elements, optionally a buffer for pH maintenance, and phosphate (typically provided as a phosphate salt). Other components can include salts such as sodium chloride, particularly for seawater microalgae. Nitrogen sources include organic and inorganic nitrogen sources, including, for example, without limitation, molecular nitrogen, nitrate, nitrate salts, ammonia (pure or in salt form, such as, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and NH<sub>4</sub>OH), protein, soybean meal, cornsteep liquor, and yeast extract. Examples of trace elements include zinc, boron, cobalt, copper, manganese, and molybdenum in, for example, the respective forms of ZnCl<sub>2</sub>, H<sub>3</sub>BO<sub>3</sub>, CoCl<sub>2</sub>·6H<sub>2</sub>O, CuCl<sub>2</sub>·2H<sub>2</sub>O, MnCl<sub>2</sub>·4H<sub>2</sub>O and (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>·4H<sub>2</sub>O.

Micromicroorganisms useful in accordance with the methods of the present invention are found in various locations and environments throughout the world. As a consequence of their isolation from other species and their resulting evolutionary divergence, the particular growth medium for optimal growth and generation of lipid and/or hydrocarbon constituents can be difficult to predict. In some cases, certain strains of microorganisms may be unable to grow on a particular growth medium because of the presence of some inhibitory component or the absence of some essential nutritional requirement required by the particular strain of microorganism.

Solid and liquid growth media are generally available from a wide variety of sources, and instructions for the preparation of particular media that is suitable for a wide variety of strains of microorganisms can be found, for example, online at <http://www.utex.org/>, a site maintained by the University of Texas at Austin, 1 University Station A6700, Austin, Tex., 78712-

0183, for its culture collection of algae (UTEX). For example, various fresh water and salt water media include those described in PCT Pub. No. 2008/151149, incorporated herein by reference.

In a particular example, Proteose Medium is suitable for axenic cultures, and a 1 L volume of the medium (pH ~6.8) can be prepared by addition of 1 g of proteose peptone to 1 liter of Bristol Medium. Bristol medium comprises 2.94 mM NaNO<sub>3</sub>, 0.17 mM CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.3 mM MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.43 mM, 1.29 mM KH<sub>2</sub>PO<sub>4</sub>, and 1.43 mM NaCl in an aqueous solution. For 1.5% agar medium, 15 g of agar can be added to 1 L of the solution. The solution is covered and autoclaved, and then stored at a refrigerated temperature prior to use. Another example is the *Prototheca* isolation medium (PIM), which comprises 10 g/L potassium hydrogen phthalate (KHP), 0.9 g/L sodium hydroxide, 0.1 g/L magnesium sulfate, 0.2 g/L potassium hydrogen phosphate, 0.3 g/L ammonium chloride, 10 g/L glucose 0.001 g/L thiamine hydrochloride, 20 g/L agar, 0.25 g/L 5-fluorocytosine, at a pH in the range of 5.0 to 5.2 (see Pore, 1973, App. Microbiology, 26: 648-649). Other suitable media for use with the methods of the invention can be readily identified by consulting the URL identified above, or by consulting other organizations that maintain cultures of microorganisms, such as SAG, CCAP, or CCALA. SAG refers to the Culture Collection of Algae at the University of Göttingen (Göttingen, Germany), CCAP refers to the culture collection of algae and protozoa managed by the Scottish Association for Marine Science (Scotland, United Kingdom), and CCALA refers to the culture collection of algal laboratory at the Institute of Botany (Třeboň, Czech Republic). Additionally, U.S. Pat. No. 5,900,370 describes media formulations and conditions suitable for heterotrophic fermentation of *Prototheca* species.

For oil production, selection of a fixed carbon source is important, as the cost of the fixed carbon source must be sufficiently low to make oil production economical. Thus, while suitable carbon sources include, for example, acetate, floridoside, fructose, galactose, glucuronic acid, glucose, glycerol, lactose, mannose, N-acetylglucosamine, rhamnose, sucrose, and/or xylose, selection of feedstocks containing those compounds is an important aspect of the methods of the invention. Suitable feedstocks useful in accordance with the methods of the invention include, for example, black liquor, corn starch, depolymerized cellulosic material, milk whey, molasses, potato, sorghum, sucrose, sugar beet, sugar cane, rice, and wheat. Carbon sources can also be provided as a mixture, such as a mixture of sucrose and depolymerized sugar beet pulp. The one or more carbon source(s) can be supplied at a concentration of at least about 50 μM, at least about 100 μM, at least about 500 μM, at least about 5 mM, at least about 50 mM, and at least about 500 mM, of one or more exogenously provided fixed carbon source(s). Carbon sources of particular interest for purposes of the present invention include cellulose (in a depolymerized form), glycerol, sucrose, and sorghum, each of which is discussed in more detail below.

In accordance with the present invention, microorganisms can be cultured using depolymerized cellulosic biomass as a feedstock. Cellulosic biomass (e.g., stover, such as corn stover) is inexpensive and readily available; however, attempts to use this material as a feedstock for yeast have failed. In particular, such feedstocks have been found to be inhibitory to yeast growth, and yeast cannot use the 5-carbon sugars produced from cellulosic materials (e.g., xylose from hemi-cellulose). By contrast, microalgae can grow on processed cel-

lulosic material. Cellulosic materials generally include about 40-60% cellulose; about 20-40% hemicellulose; and 10-30% lignin.

Suitable cellulosic materials include residues from herbaceous and woody energy crops, as well as agricultural crops, i.e., the plant parts, primarily stalks and leaves, not removed from the fields with the primary food or fiber product. Examples include agricultural wastes such as sugarcane bagasse, rice hulls, corn fiber (including stalks, leaves, husks, and cobs), wheat straw, rice straw, sugar beet pulp, citrus pulp, citrus peels; forestry wastes such as hardwood and softwood thinnings, and hardwood and softwood residues from timber operations; wood wastes such as saw mill wastes (wood chips, sawdust) and pulp mill waste; urban wastes such as paper fractions of municipal solid waste, urban wood waste and urban green waste such as municipal grass clippings; and wood construction waste. Additional cellulose include dedicated cellulosic crops such as switchgrass, hybrid poplar wood, and *miscanthus*, fiber cane, and fiber sorghum. Five-carbon sugars that are produced from such materials include xylose.

Cellulosic materials are treated to increase the efficiency with which the microbe can utilize the sugar(s) contained within the materials. The invention provides novel methods for the treatment of cellulosic materials after acid explosion so that the materials are suitable for use in a heterotrophic culture of microbes (e.g., microalgae and oleaginous yeast). As discussed above, lignocellulosic biomass is comprised of various fractions, including cellulose, a crystalline polymer of beta 1,4 linked glucose (a six-carbon sugar), hemicellulose, a more loosely associated polymer predominantly comprised of xylose (a five-carbon sugar) and to a lesser extent mannose, galactose, arabinose, lignin, a complex aromatic polymer comprised of sinapyl alcohol and its derivatives, and pectins, which are linear chains of an alpha 1,4 linked polygalacturonic acid. Because of the polymeric structure of cellulose and hemicellulose, the sugars (e.g., monomeric glucose and xylose) in them are not in a form that can be efficiently used (metabolized) by many microbes. For such microbes, further processing of the cellulosic biomass to generate the monomeric sugars that make up the polymers can be very helpful to ensuring that the cellulosic materials are efficiently utilized as a feedstock (carbon source).

Cellulose or cellulosic biomass is subjected to a process, termed "explosion", in which the biomass is treated with dilute sulfuric (or other) acid at elevated temperature and pressure. This process conditions the biomass such that it can be efficiently subjected to enzymatic hydrolysis of the cellulosic and hemicellulosic fractions into glucose and xylose monomers. The resulting monomeric sugars are termed cellulosic sugars. Cellulosic sugars can subsequently be utilized by microorganisms to produce a variety of metabolites (e.g., lipid). The acid explosion step results in a partial hydrolysis of the hemicellulose fraction to constituent monosaccharides. These sugars can be completely liberated from the biomass with further treatment. In some embodiments, the further treatment is a hydrothermal treatment that includes washing the exploded material with hot water, which removes contaminants such as salts. This step is not necessary for cellulosic ethanol fermentations due to the more dilute sugar concentrations used in such processes. In other embodiments, the further treatment is additional acid treatment. In still other embodiments, the further treatment is enzymatic hydrolysis of the exploded material. These treatments can also be used in any combination. The type of treatment can affect the type of sugars liberated (e.g., five carbon sugars versus six carbon sugars) and the stage at which they are liberated in the pro-

cess. As a consequence, different streams of sugars, whether they are predominantly five-carbon or six-carbon, can be created. These enriched five-carbon or six-carbon streams can thus be directed to specific microorganisms with different carbon utilization capabilities.

The methods of the present invention typically involve fermentation to higher cell densities than what is achieved in ethanol fermentation. Because of the higher densities of the cultures for heterotrophic cellulosic oil production, the fixed carbon source (e.g., the cellulosic derived sugar stream(s)) is preferably in a concentrated form. The glucose level of the depolymerized cellulosic material is preferably at least 300 g/liter, at least 400 g/liter, at least 500 g/liter or at least 600 g/liter prior to the cultivation step, which is optionally a fed batch cultivation in which the material is fed to the cells over time as the cells grow and accumulate lipid. Cellulosic sugar streams are not used at or near this concentration range in the production of cellulosic ethanol. Thus, in order to generate and sustain the very high cell densities during the production of lignocellulosic oil, the carbon feedstock(s) must be delivered into the heterotrophic cultures in a highly concentrated form. However, any component in the feedstream that is not a substrate for, and is not metabolized by, the oleaginous microorganism will accumulate in the bioreactor, which can lead to problems if the component is toxic or inhibitory to production of the desired end product. While ligin and lignin-derived by-products, carbohydrate-derived byproducts such as furfurals and hydroxymethyl furfurals and salts derived from the generation of the cellulosic materials (both in the explosion process and the subsequent neutralization process), and even non-metabolized pentose/hexose sugars can present problems in ethanolic fermentations, these effects are amplified significantly in a process in which their concentration in the initial feedstock is high. To achieve sugar concentrations in the 300 g/L range (or higher) for six-carbon sugars that may be used in large scale production of lignocellulosic oil described in the present invention, the concentration of these toxic materials can be 20 times higher than the concentrations typically present in ethanolic fermentations of cellulosic biomass.

The explosion process treatment of the cellulosic material utilizes significant amounts of sulfuric acid, heat and pressure, thereby liberating by-products of carbohydrates, namely furfurals and hydroxymethyl furfurals. Furfurals and hydroxymethyl furfurals are produced during hydrolysis of hemicellulose through dehydration of xylose into furfural and water. In some embodiments of the present invention, these by-products (e.g., furfurals and hydroxymethyl furfurals) are removed from the saccharified lignocellulosic material prior to introduction into the bioreactor. In certain embodiments of the present invention, the process for removal of the by-products of carbohydrates is hydrothermal treatment of the exploded cellulosic materials. In addition, the present invention provides methods in which strains capable of tolerating compounds such as furfurals or hydroxymethyl furfurals are used for lignocellulosic oil production. In another embodiment, the present invention also provides methods and microorganisms that are not only capable of tolerating furfurals in the fermentation media, but are actually able to metabolize these by-products during the production of lignocellulosic oil.

The explosion process also generates significant levels of salts. For example, typical conditions for explosion can result in conductivities in excess of 5 mS/cm when the exploded cellulosic biomass is resuspended at a ratio of 10:1 water: solids (dry weight). In certain embodiments of the present invention, the diluted exploded biomass is subjected to enzy-

matic saccharification, and the resulting supernatant is concentrated up to 25 fold for use in the bioreactor. The salt level (as measured by conductivity) in the concentrated sugar stream(s) can be unacceptably high (up to 1.5 M Na<sup>+</sup> equivalents). Additional salts are generated upon neutralization of the exploded materials for the subsequent enzymatic saccharification process as well. The present invention provides methods for removing these salts so that the resulting concentrated cellulosic sugar stream(s) can be used in heterotrophic processes for producing lignocellulosic oil. In some embodiments, the method of removing these salts is deionization with resins, such as, but not limited to, DOWEX Marathon MR3. In certain embodiments, the deionization with resin step occurs before sugar concentration or pH adjustment and hydrothermal treatment of biomass prior to saccharification, or any combination of the preceding; in other embodiments, the step is conducted after one or more of these processes. In other embodiments, the explosion process itself is changed so as to avoid the generation of salts at unacceptably high levels. For example, a suitable alternative to sulfuric acid (or other acid) explosion of the cellulosic biomass is mechanical pulping to render the cellulosic biomass receptive to enzymatic hydrolysis (saccharification). In still other embodiments, native strains of microorganisms resistant to high levels of salts or genetically engineered strains with resistance to high levels of salts are used.

A preferred embodiment for the process of preparing of exploded cellulosic biomass for use in heterotrophic lignocellulosic oil production using oleaginous microbes is diagrammed in FIG. 10. Step I. comprises adjusting the pH of the resuspended exploded cellulosic biomass to the range of 5.0-5.3 followed by washing the cellulosic biomass three times. This washing step can be accomplished by a variety of means including the use of desalting and ion exchange resins, reverse osmosis, hydrothermal treatment (as described above), or just repeated re-suspension and centrifugation in deionized water. This wash step results in a cellulosic stream whose conductivity is between 100-300  $\mu$ S/cm and the removal of significant amounts of furfurals and hydroxymethyl furfurals. Decants from this wash step can be saved to concentrate five-carbon sugars liberated from the hemicellulose fraction. Step II comprises enzymatic saccharification of the washed cellulosic biomass. In a preferred embodiment, Accellerase (Genencor) is used. Step III comprises the recovery of sugars via centrifugation or decanting and rinsing of the saccharified biomass. The resulting biomass (solids) is an energy dense, lignin rich component that can be used as fuel or sent to waste. The recovered sugar stream in the centrifugation/decanting and rinse process is collected. Step IV comprises microfiltration to remove contaminating solids with recovery of the permeate. Step V comprises a concentration step which can be accomplished using a vacuum evaporator. This step can optionally include the addition of antifoam agents such as P2000 (Sigma/Fluka), which is sometimes necessary due to the protein content of the resulting sugar feedstock.

In another embodiment of the methods of the invention, the carbon source is glycerol, including acidulated and non-acidulated glycerol byproduct from biodiesel transesterification. In one embodiment, the carbon source includes glycerol and at least one other carbone source. In some cases, all of the glycerol and the at least one other fixed carbon source are provided to the microorganism at the beginning of the fermentation. In some cases, the glycerol and the at least one other fixed carbon source are provided to the microorganism simultaneously at a predetermined ratio. In some cases, the

glycerol and the at least one other fixed carbon source are fed to the microbes at a predetermined rate over the course of fermentation.

Some microalgae undergo cell division faster in the presence of glycerol than in the presence of glucose (see PCT Pub. No. 2008/151149). In these instances, two-stage growth processes in which cells are first fed glycerol to rapidly increase cell density, and are then fed glucose to accumulate lipids can improve the efficiency with which lipids are produced. The use of the glycerol byproduct of the transesterification process provides significant economic advantages when put back into the production process. Other feeding methods are provided as well, such as mixtures of glycerol and glucose. Feeding such mixtures also captures the same economic benefits. In addition, the invention provides methods of feeding alternative sugars to microalgae such as sucrose in various combinations with glycerol.

In another embodiment of the methods of the invention, the carbon source is sucrose, including a complex feedstock containing sucrose, such as thick cane juice from sugar cane processing. In one embodiment, the culture medium further includes at least one sucrose utilization enzyme. In some cases, the culture medium includes a sucrose invertase. In one embodiment, the sucrose invertase enzyme is a secretable sucrose invertase enzyme encoded by an exogenous sucrose invertase gene expressed by the population of microorganisms. Thus, in some cases, as described in more detail in Section IV, below, the microalgae has been genetically engineered to express a sucrose utilization enzyme, such as a sucrose transporter, a sucrose invertase, a hexokinase, a glucokinase, or a fructokinase.

Complex feedstocks containing sucrose include waste molasses from sugar cane processing; the use of this low-value waste product of sugar cane processing can provide significant cost savings in the production of hydrocarbons and other oils. Another complex feedstock containing sucrose that is useful in the methods of the invention is sorghum, including sorghum syrup and pure sorghum. Sorghum syrup is produced from the juice of sweet sorghum cane. Its sugar profile consists of mainly glucose (dextrose), fructose and sucrose.

#### 4. Oil Production

For the production of oil in accordance with the methods of the invention, it is preferable to culture cells in the dark, as is the case, for example, when using extremely large (40,000 liter and higher) fermentors that do not allow light to strike the culture. *Prototheca* species are grown and propagated for the production of oil in a medium containing a fixed carbon source and in the absence of light; such growth is known as heterotrophic growth.

As an example, an inoculum of lipid-producing microalgal cells are introduced into the medium; there is a lag period (lag phase) before the cells begin to propagate. Following the lag period, the propagation rate increases steadily and enters the log, or exponential, phase. The exponential phase is in turn followed by a slowing of propagation due to decreases in nutrients such as nitrogen, increases in toxic substances, and quorum sensing mechanisms. After this slowing, propagation stops, and the cells enter a stationary phase or steady growth state, depending on the particular environment provided to the cells. For obtaining lipid rich biomass, the culture is typically harvested well after then end of the exponential phase, which may be terminated early by allowing nitrogen or another key nutrient (other than carbon) to become depleted, forcing the cells to convert the carbon sources, present in excess, to lipid. Culture condition parameters can be manipu-

lated to optimize total oil production, the combination of lipid species produced, and/or production of a specific oil.

As discussed above, a bioreactor or fermentor is used to allow cells to undergo the various phases of their growth cycle. As an example, an inoculum of lipid-producing cells can be introduced into a medium followed by a lag period (lag phase) before the cells begin growth. Following the lag period, the growth rate increases steadily and enters the log, or exponential, phase. The exponential phase is in turn followed by a slowing of growth due to decreases in nutrients and/or increases in toxic substances. After this slowing, growth stops, and the cells enter a stationary phase or steady state, depending on the particular environment provided to the cells. Lipid production by cells disclosed herein can occur during the log phase or thereafter, including the stationary phase wherein nutrients are supplied, or still available, to allow the continuation of lipid production in the absence of cell division.

Preferably, microorganisms grown using conditions described herein and known in the art comprise at least about 20% by weight of lipid, preferably at least about 40% by weight, more preferably at least about 50% by weight, and most preferably at least about 60% by weight. Process conditions can be adjusted to increase the yield of lipids suitable for a particular use and/or to reduce production cost. For example, in certain embodiments, a microalgae is cultured in the presence of a limiting concentration of one or more nutrients, such as, for example, nitrogen, phosphorous, or sulfur, while providing an excess of fixed carbon energy such as glucose. Nitrogen limitation tends to increase microbial lipid yield over microbial lipid yield in a culture in which nitrogen is provided in excess. In particular embodiments, the increase in lipid yield is at least about: 10%, 50%, 100%, 200%, or 500%. The microbe can be cultured in the presence of a limiting amount of a nutrient for a portion of the total culture period or for the entire period. In particular embodiments, the nutrient concentration is cycled between a limiting concentration and a non-limiting concentration at least twice during the total culture period. Lipid content of cells can be increased by continuing the culture for increased periods of time while providing an excess of carbon, but limiting or no nitrogen.

In another embodiment; lipid yield is increased by culturing a lipid-producing microbe (e.g., microalgae) in the presence of one or more cofactor(s) for a lipid pathway enzyme (e.g., a fatty acid synthetic enzyme). Generally, the concentration of the cofactor(s) is sufficient to increase microbial lipid (e.g., fatty acid) yield over microbial lipid yield in the absence of the cofactor(s). In a particular embodiment, the cofactor(s) are provided to the culture by including in the culture a microbe (e.g., microalgae) containing an exogenous gene encoding the cofactor(s). Alternatively, cofactor(s) may be provided to a culture by including a microbe (e.g., microalgae) containing an exogenous gene that encodes a protein that participates in the synthesis of the cofactor. In certain embodiments, suitable cofactors include any vitamin required by a lipid pathway enzyme, such as, for example: biotin, pantothenate. Genes encoding cofactors suitable for use in the invention or that participate in the synthesis of such cofactors are well known and can be introduced into microbes (e.g., microalgae), using constructs and techniques such as those described above.

The specific examples of bioreactors, culture conditions, and heterotrophic growth and propagation methods described herein can be combined in any suitable manner to improve efficiencies of microbial growth and lipid and/or protein production.

Microalgal biomass with a high percentage of oil/lipid accumulation by dry weight has been generated using different methods of culture, which are known in the art (see PCT Pub. No. 2008/151149). Microalgal biomass generated by the culture methods described herein and useful in accordance with the present invention comprises at least 10% microalgal oil by dry weight. In some embodiments, the microalgal biomass comprises at least 25%, at least 50%, at least 55%, or at least 60% microalgal oil by dry weight. In some embodiments, the microalgal biomass contains from 10-90% microalgal oil, from 25-75% microalgal oil, from 40-75% microalgal oil, or from 50-70% microalgal oil by dry weight.

The microalgal oil of the biomass described herein, or extracted from the biomass for use in the methods and compositions of the present invention can comprise glycerolipids with one or more distinct fatty acid ester side chains. Glycerolipids are comprised of a glycerol molecule esterified to one, two or three fatty acid molecules, which can be of varying lengths and have varying degrees of saturation. The length and saturation characteristics of the fatty acid molecules (and the microalgal oils) can be manipulated to modify the properties or proportions of the fatty acid molecules in the microalgal oils of the present invention via culture conditions or via lipid pathway engineering, as described in more detail in Section IV, below. Thus, specific blends of algal oil can be prepared either within a single species of algae by mixing together the biomass or algal oil from two or more species of microalgae, or by blending algal oil of the invention with oils from other sources such as soy, rapeseed, canola, palm, palm kernel, coconut, corn, waste vegetable, Chinese tallow, olive, sunflower, cottonseed, chicken fat, beef tallow, porcine tallow, microalgae, macroalgae, microbes, *Cuphea*, flax, peanut, choice white grease, lard, *Camelina sativa*, mustard seed, cashew nut, oats, lupine, kenaf, *calendula*, hemp, coffee, linseed (flax), hazelnut, *euphorbia*, pumpkin seed, coriander, *camellia*, sesame, safflower, rice, tung tree, cocoa, copra, pium poppy, castor beans, pecan, jojoba, *macadamia*, Brazil nuts, avocado, petroleum, or a distillate fraction of any of the preceding oils.

The oil composition, i.e., the properties and proportions of the fatty acid constituents of the glycerolipids, can also be manipulated by combining biomass or oil from at least two distinct species of microalgae. In some embodiments, at least two of the distinct species of microalgae have different glycerolipid profiles. The distinct species of microalgae can be cultured together or separately as described herein, preferably under heterotrophic conditions, to generate the respective oils. Different species of microalgae can contain different percentages of distinct fatty acid constituents in the cell's glycerolipids.

Generally, *Prototheca* strains have very little or no fatty acids with the chain length C8-C14. For example, *Prototheca moriformis* (UTEX 1435), *Prototheca krugani* (UTEX 329), *Prototheca stagnora* (UTEX 1442) and *Prototheca zopfii* (UTEX 1438) contains no (or undetectable amounts) C8 fatty acids, between 0-0.01% C10 fatty acids, between 0.03-2.1% C12 fatty acids and between 1.0-1.7% C14 fatty acids.

In some cases, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain lengths C8-10 has at least 0.3%, at least 0.8%, at least 1.5% or more fatty acids of chain length C8 and at least 0.3%, at least 1.0%, at least 3.0%, at least 5% or more fatty acids of chain length C10. In other instances, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C12 has at least 3.0%, at least 5%, at least 7%, at least 10%, at least 13%

or more fatty acids of the chain length C12 and at least 1.5%, at least 2%, or at least 3% or more fatty acids of the chain length C14. In other cases, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C14 has at least 4.0%, at least 7%, at least 10%, at least 15%, at least 20%, at least 25% or more fatty acids of the chain length C14, and at least 0.4%, at least 1%, at least 1.5%, or more fatty acids of the chain length C12.

In non-limiting examples, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C8 and C10 has between 0.3-1.58% fatty acids of chain length C8 and between 0.35-6.76% fatty acids of the chain length C10. In other non-limiting examples, *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C12 has between 3.9-14.11% fatty acids of the chain length C12 and between 1.95-3.05% fatty acids of the chain length C14. In other non-limiting examples, *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C14 has between 4.40-17.35% fatty acids of the chain length C14 and between 0.4-1.83 Area % fatty acids of the chain length C12. In some cases, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain lengths between C8 and C14 have between 3.5-20% medium chain (C8-C14) fatty acids. In some instances, keeping the transgenic *Prototheca* strains under constant and high selective pressure to retain exogenous genes is advantageous due to the increase in the desired fatty acid of a specific chain length. In a non-limiting example, Example 5 demonstrates a two fold increase in C14 chain length fatty acids (more than 30% C8-C14 chain length fatty acids) when the culture of *Prototheca moriformis* containing a C14 preferring thioesterase exogenous gene is retained. High levels of exogenous gene retention can also be achieved by inserting exogenous genes into the nuclear chromosomes of the cells using homologous recombination vectors and methods disclosed herein. Recombinant cells containing exogenous genes integrated into nuclear chromosomes are an object of the invention.

Microalgal oil can also include other constituents produced by the microalgae, or incorporated into the microalgal oil from the culture medium. These other constituents can be present in varying amount depending on the culture conditions used to culture the microalgae, the species of microalgae, the extraction method used to recover microalgal oil from the biomass and other factors that may affect microalgal oil composition. Non-limiting examples of such constituents include carotenoids, present from 0.1-0.4 micrograms/ml, chlorophyll present from 0-0.02 milligrams/kilogram of oil, gamma tocopherol present from 0.4-0.6 milligrams/100 grams of oil, and total tocotrienols present from 0.2-0.5 milligrams/gram of oil.

The other constituents can include, without limitation, phospholipids, tocopherols, tocotrienols, carotenoids (e.g., alpha-carotene, beta-carotene, lycopene, etc.), xanthophylls (e.g., lutein, zeaxanthin, alpha-cryptoxanthin and beta-cryptoxanthin), and various organic or inorganic compounds.

In some cases, the oil extracted from *Prototheca* species comprises no more than 0.02 mg/kg chlorophyll. In some cases, the oil extracted from *Prototheca* species comprises no more than 0.4 mcg/ml total carotenoids. In some cases the *Prototheca* oil comprises between 0.40-0.60 milligrams of

gamma tocopherol per 100 grams of oil. In other cases, the *Prototheca* oil comprises between 0.2-0.5 milligrams of total tocotrienols per gram of oil.

### III. GENETIC ENGINEERING METHODS AND MATERIALS

The present invention provides methods and materials for genetically modifying *Prototheca* cells and recombinant host cells useful in the methods of the present invention, including but not limited to recombinant *Prototheca moriformis*, *Prototheca zopfii*, *Prototheca krugani*, and *Prototheca stagnora* host cells. The description of these methods and materials is divided into subsections for the convenience of the reader. In subsection 1, transformation methods are described. In subsection 2, genetic engineering methods using homologous recombination are described. In subsection 3, expression vectors and components are described.

#### 1. Engineering Methods—Transformation

Cells can be transformed by any suitable technique including, e.g., biolistics, electroporation (see Maruyama et al. (2004), *Biotechnology Techniques* 8:821-826), glass bead transformation and silicon carbide whisker transformation. Another method that can be used involves forming protoplasts and using  $\text{CaCl}_2$  and polyethylene glycol (PEG) to introduce recombinant DNA into microalgal cells (see Kim et al. (2002), *Mar. Biotechnol.* 4:63-73, which reports the use of this method for the transformation of *Chorella ellipsoidea*). Co-transformation of microalgae can be used to introduce two distinct vector molecules into a cell simultaneously (see for example *Protist* 2004 December; 155(4):381-93).

Biolistic methods (see, for example, Sanford, *Trends In Biotech.* (1988) 6:299-302, U.S. Pat. No. 4,945,050; electroporation (Fromm et al., *Proc. Nat'l. Acad. Sci. (USA)* (1985) 82:5824-5828); use of a laser beam, microinjection or any other method capable of introducing DNA into a microalga can also be used for transformation of a *Prototheca* cell.

#### 2. Engineering Methods—Homologous Recombination

Homologous recombination is the ability of complementary DNA sequences to align and exchange regions of homology. Transgenic DNA (“donor”) containing sequences homologous to the genomic sequences being targeted (“template”) is introduced into the organism and then undergoes recombination into the genome at the site of the corresponding genomic homologous sequences. The mechanistic steps of this process, in most cases, include: (1) pairing of homologous DNA segments; (2) introduction of double-stranded breaks into the donor DNA molecule; (3) invasion of the template DNA molecule by the free donor DNA ends followed by DNA synthesis; and (4) resolution of double-strand break repair events that result in final recombination products.

The ability to carry out homologous recombination in a host organism has many practical implications for what can be carried out at the molecular genetic level and is useful in the generation of an oleaginous microbe that can produce tailored oils. By its very nature homologous recombination is a precise gene targeting event, hence, most transgenic lines generated with the same targeting sequence will be essentially identical in terms of phenotype, necessitating the screening of far fewer transformation events. Homologous recombination also targets gene insertion events into the host chromosome, resulting in excellent genetic stability, even in the absence of genetic selection. Because different chromosomal loci will likely impact gene expression, even from heterologous promoters/UTRs, homologous recombination

can be a method of querying loci in an unfamiliar genome environment and to assess the impact of these environments on gene expression.

Particularly useful genetic engineering applications using homologous recombination is to co-opt specific host regulatory elements such as promoters/UTRs to drive heterologous gene expression in a highly specific fashion. For example, precise ablation of the endogenous stearyl ACP desaturase gene with a heterologous C12:0 specific FATB (thioesterase) gene cassette and suitable selective marker, might be expected to dramatically decrease endogenous levels of C18:1 fatty acids concomitant with increased levels of the C12:0 fatty acids. Example 13 describes the homologous recombination targeting construct that is suitable for the ablation of an endogenous *Prototheca moriformis* stearyl ACP desaturase gene.

Because homologous recombination is a precise gene targeting event, it can be used to precisely modify any nucleotide(s) within a gene or region of interest, so long as sufficient flanking regions have been identified. Therefore, homologous recombination can be used as a means to modify regulatory sequences impacting gene expression of rRNA and/or proteins. It can also be used to modify protein coding regions in an effort to modify enzyme activities such as substrate specificity, affinities and  $K_m$ , and thus affecting the desired change in metabolism of the host cell. Homologous recombination provides a powerful means to manipulate the host genome resulting in gene targeting, gene conversion, gene deletion, gene duplication, gene inversion and exchanging gene expression regulatory elements such as promoters, enhancers and 3'UTRs.

Homologous recombination can be achieved by using targeting constructs containing pieces of endogenous sequences to “target” the gene or region of interest within the endogenous host cell genome. Such targeting sequences can either be located 5' of the gene or region of interest, 3' of the gene/region of interest or even flank the gene/region of interest. Such targeting constructs can be transformed into the host cell either as a supercoiled plasmid DNA with additional vector backbone, a PCR product with no vector backbone, or as a linearized molecule. In some cases, it may be advantageous to first expose the homologous sequences within the transgenic DNA (donor DNA) with a restriction enzyme. This step can increase the recombination efficiency and decrease the occurrence of undesired events. Other methods of increasing recombination efficiency include using PCR to generate transforming transgenic DNA containing linear ends homologous to the genomic sequences being targeted.

#### 3. Vectors and Vector Components

Vectors for transformation of microorganisms in accordance with the present invention can be prepared by known techniques familiar to those skilled in the art in view of the disclosure herein. A vector typically contains one or more genes, in which each gene codes for the expression of a desired product (the gene product) and is operably linked to one or more control sequences that regulate gene expression or target the gene product to a particular location in the recombinant cell. To aid the reader, this subsection is divided into subsections. Subsection A describes control sequences typically contained on vectors as well as novel control sequences provided by the present invention. Subsection B describes genes typically contained in vectors as well as novel codon optimization methods and genes prepared using them provided by the invention.

##### A. Control Sequences

Control sequences are nucleic acids that regulate the expression of a coding sequence or direct a gene product to a

particular location in or outside a cell. Control sequences that regulate expression include, for example, promoters that regulate transcription of a coding sequence and terminators that terminate transcription of a coding sequence. Another control sequence is a 3' untranslated sequence located at the end of a coding sequence that encodes a polyadenylation signal. Control sequences that direct gene products to particular locations include those that encode signal peptides, which direct the protein to which they are attached to a particular location in or outside the cell.

Thus, an exemplary vector design for expression of an exogenous gene in a microalgae contains a coding sequence for a desired gene product (for example, a selectable marker, a lipid pathway modification enzyme, or a sucrose utilization enzyme) in operable linkage with a promoter active in microalgae. Alternatively, if the vector does not contain a promoter in operable linkage with the coding sequence of interest, the coding sequence can be transformed into the cells such that it becomes operably linked to an endogenous promoter at the point of vector integration. The promoterless method of transformation has been proven to work in microalgae (see for example Plant Journal 14:4, (1998), pp. 441-447).

Many promoters are active in microalgae, including promoters that are endogenous to the algae being transformed, as well as promoters that are not endogenous to the algae being transformed (i.e., promoters from other algae, promoters from higher plants, and promoters from plant viruses or algae viruses). Illustrative exogenous and/or endogenous promoters that are active in microalgae (as well as antibiotic resistance genes functional in microalgae) are described in PCT Pub. No. 2008/151149 and references cited therein.

The promoter used to express an exogenous gene can be the promoter naturally linked to that gene or can be a heterologous gene. Some promoters are active in more than one species of microalgae. Other promoters are species-specific. Illustrative promoters include promoters such as  $\beta$ -tubulin from *Chlamydomonas reinhardtii*, used in the Examples below, and viral promoters, such as cauliflower mosaic virus (CMV) and *Chlorella* virus, which have been shown to be active in multiple species of microalgae (see for example Plant Cell Rep. 2005 March; 23(10-11):727-35; J Microbiol. 2005 August; 43(4):361-5; Mar Biotechnol (NY). 2002 January; 4(1):63-73). Another promoter that is suitable for use for expression of exogenous genes in *Prototheca* is the *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR (SEQ ID NO: 69). Optionally, at least 10, 20, 30, 40, 50, or 60 nucleotides or more of these sequences containing a promoter are used. Illustrative promoters useful for expression of exogenous genes in *Prototheca* are listed in the sequence listing of this application, such as the promoter of the *Chlorella* HUP1 gene (SEQ ID NO:1) and the *Chlorella ellipsoidea* nitrate reductase promoter (SEQ ID NO:2). *Chlorella* virus promoters can also be used to express genes in *Prototheca*, such as SEQ ID NOs: 1-7 of U.S. Pat. No. 6,395,965. Additional promoters active in *Prototheca* can be found, for example, in Biochem Biophys Res Commun. 1994 Oct. 14; 204(1):187-94; Plant Mol. Biol. 1994 October; 26(1):85-93; Virology. 2004 Aug. 15; 326(1):150-9; and Virology. 2004 Jan. 5; 318(1):214-23.

A promoter can generally be characterized as either constitutive or inducible. Constitutive promoters are generally active or function to drive expression at all times (or at certain times in the cell life cycle) at the same level. Inducible promoters, conversely, are active (or rendered inactive) or are significantly up- or down-regulated only in response to a stimulus. Both types of promoters find application in the

methods of the invention. Inducible promoters useful in the invention include those that mediate transcription of an operably linked gene in response to a stimulus, such as an exogenously provided small molecule (e.g. glucose, as in SEQ ID NO:1), temperature (heat or cold), lack of nitrogen in culture media, etc. Suitable promoters can activate transcription of an essentially silent gene or upregulate, preferably substantially, transcription of an operably linked gene that is transcribed at a low level.

Inclusion of termination region control sequence is optional, and if employed, then the choice is be primarily one of convenience, as the termination region is relatively interchangeable. The termination region may be native to the transcriptional initiation region (the promoter), may be native to the DNA sequence of interest, or may be obtainable from another source. See, for example, Chen and Orozco, Nucleic Acids Res. (1988) 16:8411.

The present invention also provides control sequences and recombinant genes and vectors containing them that provide for the compartmentalized expression of a gene of interest. Organelles for targeting are chloroplasts, plastids, mitochondria, and endoplasmic reticulum. In addition, the present invention provides control sequences and recombinant genes and vectors containing them that provide for the secretion of a protein outside the cell.

Proteins expressed in the nuclear genome of *Prototheca* can be targeted to the plastid using plastid targeting signals. Plastid targeting sequences endogenous to *Chlorella* are known, such as genes in the *Chlorella* nuclear genome that encode proteins that are targeted to the plastid; see for example GenBank Accession numbers AY646197 and AF499684, and in one embodiment, such control sequences are used in the vectors of the present invention to target expression of a protein to a *Prototheca* plastid.

The Examples below describe the use of algal plastid targeting sequences to target heterologous proteins to the correct compartment in the host cell. cDNA libraries were made using *Prototheca moriformis* and *Chlorella protothecoides* cells and are described in Examples 12 and Example 11 below. Sequences were BLASTed and analyzed for homology to known proteins that traffic to the plastid/chloroplast. The cDNAs encoding these proteins were cloned and plastid targeting sequences were isolated from these cDNAs. The amino acid sequences of the algal plastid targeting sequences identified from the cDNA libraries and the amino acid sequences of plant fatty acyl-ACP thioesterases that are used in the heterologous expression Examples below are listed in SEQ ID NOs: 127-133.

In another embodiment of the present invention, the expression of a polypeptide in *Prototheca* is targeted to the endoplasmic reticulum. The inclusion of an appropriate retention or sorting signal in an expression vector ensure that proteins are retained in the endoplasmic reticulum (ER) and do not go downstream into Golgi. For example, the IMPACTVECTOR1.3 vector, from Wageningen UR-Plant Research International, includes the well known KDEL retention or sorting signal. With this vector, ER retention has a practical advantage in that it has been reported to improve expression levels 5-fold or more. The main reason for this appears to be that the ER contains lower concentrations and/or different proteases responsible for post-translational degradation of expressed proteins than are present in the cytoplasm. ER retention signals functional in green microalgae are known. For example, see Proc Natl Acad Sci USA. 2005 Apr. 26; 102(17):6225-30.

In another embodiment of the present invention, a polypeptide is targeted for secretion outside the cell into the culture



media. See Hawkins et al., Current Microbiology Vol. 38 (1999), pp. 335-341 for examples of secretion signals active in *Chlorella* that can be used, in accordance with the methods of the invention, in *Prototheca*.

#### B. Genes and Codon Optimization

Typically, a gene includes a promoter, coding sequence, and termination control sequences. When assembled by recombinant DNA technology, a gene may be termed an expression cassette and may be flanked by restriction sites for convenient insertion into a vector that is used to introduce the recombinant gene into a host cell. The expression cassette can be flanked by DNA sequences from the genome or other nucleic acid target to facilitate stable integration of the expression cassette into the genome by homologous recombination. Alternatively, the vector and its expression cassette may remain unintegrated, in which case, the vector typically includes an origin of replication, which is capable of providing for replication of the heterologous vector DNA.

A common gene present on a vector is a gene that codes for a protein, the expression of which allows the recombinant cell containing the protein to be differentiated from cells that do not express the protein. Such a gene, and its corresponding gene product, is called a selectable marker. Any of a wide variety of selectable markers can be employed in a transgene construct useful for transforming *Prototheca*. Examples of suitable selectable markers include the G418 resistance gene, the nitrate reductase gene (see Dawson et al. (1997), Current Microbiology 35:356-362), the hygromycin phosphotransferase gene (HPT; see Kim et al. (2002), Mar. Biotechnol. 4:63-73), the neomycin phosphotransferase gene, and the ble gene, which confers resistance to phleomycin (Huang et al. (2007), Appl. Microbiol. Biotechnol. 72:197-205). Methods of determining sensitivity of microalgae to antibiotics are well known. For example, Mol Gen Genet. 1996 Oct. 16; 252(5):572-9.

For purposes of the present invention, the expression vector used to prepare a recombinant host cell of the invention will include at least two, and often three, genes, if one of the genes is a selectable marker. For example, a genetically engineered *Prototheca* of the invention can be made by transformation with vectors of the invention that comprise, in addition to a selectable marker, one or more exogenous genes, such as, for example, sucrose invertase gene or acyl ACP-thioesterase gene. One or both genes can be expressed using an inducible promoter, which allows the relative timing of expression of these genes to be controlled to enhance the lipid yield and conversion to fatty acid esters. Expression of the two or more exogenous genes may be under control of the same inducible promoter or under control of different inducible (or constitutive) promoters. In the latter situation, expression of a first exogenous gene can be induced for a first period of time (during which expression of a second exogenous gene may or may not be induced) and expression of a second exogenous gene can be induced for a second period of time (during which expression of a first exogenous gene may or may not be induced).

In other embodiments, the two or more exogenous genes (in addition to any selectable marker) are: a fatty acyl-ACP thioesterase and a fatty acyl-CoA/aldehyde reductase, the combined action of which yields an alcohol product. Further provided are other combinations of exogenous genes, including without limitation, a fatty acyl-ACP thioesterase and a fatty acyl-CoA reductase to generate aldehydes. In one embodiment, the vector provides for the combination of a fatty acyl-ACP thioesterase, a fatty acyl-CoA reductase, and a fatty aldehyde decarboxylase to generate alkanes. In each of

these embodiments, one or more of the exogenous genes can be expressed using an inducible promoter.

Other illustrative vectors of the invention that express two or more exogenous genes include those encoding both a sucrose transporter and a sucrose invertase enzyme and those encoding both a selectable marker and a secreted sucrose invertase. The recombinant *Prototheca* transformed with either type of vector produce lipids at lower manufacturing cost due to the engineered ability to use sugar cane (and sugar cane-derived sugars) as a carbon source. Insertion of the two exogenous genes described above can be combined with the disruption of polysaccharide biosynthesis through directed and/or random mutagenesis, which steers ever greater carbon flux into lipid production. Individually and in combination, trophic conversion, engineering to alter lipid production and treatment with exogenous enzymes alter the lipid composition produced by a microorganism. The alteration can be a change in the amount of lipids produced, the amount of one or more hydrocarbon species produced relative to other lipids, and/or the types of lipid species produced in the microorganism. For example, microalgae can be engineered to produce a higher amount and/or percentage of TAGs.

For optimal expression of a recombinant protein, it is beneficial to employ coding sequences that produce mRNA with codons preferentially used by the host cell to be transformed. Thus, proper expression of transgenes can require that the codon usage of the transgene matches the specific codon bias of the organism in which the transgene is being expressed. The precise mechanisms underlying this effect are many, but include the proper balancing of available aminoacylated tRNA pools with proteins being synthesized in the cell, coupled with more efficient translation of the transgenic messenger RNA (mRNA) when this need is met. When codon usage in the transgene is not optimized, available tRNA pools are not sufficient to allow for efficient translation of the heterologous mRNA resulting in ribosomal stalling and termination and possible instability of the transgenic mRNA.

The present invention provides codon-optimized nucleic acids useful for the successful expression of recombinant proteins in *Prototheca*. Codon usage in *Prototheca* species was analyzed by studying cDNA sequences isolated from *Prototheca moriformis*. This analysis represents the interrogation over 24,000 codons and resulted in Table 1 below.

TABLE 1

Preferred codon usage in <i>Prototheca</i> strains.		
Ala	GCG	345 (0.36)
	GCA	66 (0.07)
	GCT	101 (0.11)
	GCC	442 (0.46)
Cys	TGT	12 (0.10)
	TGC	105 (0.90)
Asp	GAT	43 (0.12)
	GAC	316 (0.88)
Glu	GAG	377 (0.96)
	GAA	14 (0.04)
Phe	TTT	89 (0.29)
	TTC	216 (0.71)
Gly	GGG	92 (0.12)
	GGA	56 (0.07)
	GGT	76 (0.10)
	GGC	559 (0.71)

TABLE 1-continued

Preferred codon usage in <i>Prototheca</i> strains.		
His	CAT	42 (0.21)
	CAC	154 (0.79)
Ile	ATA	4 (0.01)
	ATT	30 (0.08)
	ATC	338 (0.91)
Lys	AAG	284 (0.98)
	AAA	7 (0.02)
Leu	TTG	26 (0.04)
	TTA	3 (0.00)
	CTG	447 (0.61)
	CTA	20 (0.03)
	CTT	45 (0.06)
Met	ATG	191 (1.00)
	ATC	191 (1.00)
Asn	AAT	8 (0.04)
	AAC	201 (0.96)
Pro	CCG	161 (0.29)
	CCA	49 (0.09)
	CCT	71 (0.13)
	CCC	267 (0.49)
Gln	CAG	226 (0.82)
	CAA	48 (0.18)
Arg	AGG	33 (0.06)
	AGA	14 (0.02)
	CGG	102 (0.18)
	CGA	49 (0.08)
	CGT	51 (0.09)
Ser	AGT	16 (0.03)
	AGC	123 (0.22)
	TCC	152 (0.28)
	TCA	31 (0.06)
	TCT	55 (0.10)
Thr	ACC	173 (0.31)
	ACG	184 (0.38)
	ACA	24 (0.05)
	ACT	21 (0.05)
Val	ACC	249 (0.52)
	GTG	308 (0.50)
	GTA	9 (0.01)
	GTT	35 (0.06)
Trp	GTC	262 (0.43)
	TGG	107 (1.00)
Tyr	TAT	10 (0.05)
	TAC	180 (0.95)
Stop	TGA/TAG/TAA	

In other embodiments, the gene in the recombinant vector has been codon-optimized with reference to a microalgal strain other than a *Prototheca* strain. For example, methods of recoding genes for expression in microalgae are described in U.S. Pat. No. 7,135,290. Additional information for codon optimization is available, e.g., at the codon usage database of GenBank.

While the methods and materials of the invention allow for the introduction of any exogenous gene into *Prototheca*, genes relating to sucrose utilization and lipid pathway modification are of particular interest, as discussed in the following sections.

## IV. SUCROSE UTILIZATION

In embodiment, the recombinant *Prototheca* cell of the invention further contains one or more exogenous sucrose utilization genes. In various embodiments, the one or more genes encode one or more proteins selected from the group consisting of a fructokinase, a glucokinase, a hexokinase, a sucrose invertase, a sucrose transporter. For example, expression of a sucrose transporter and a sucrose invertase allows *Prototheca* to transport sucrose into the cell from the culture media and hydrolyze sucrose to yield glucose and fructose. Optionally, a fructokinase can be expressed as well in instances where endogenous hexokinase activity is insufficient for maximum phosphorylation of fructose. Examples of suitable sucrose transporters are Genbank accession numbers CAD91334, CAB92307, and CAA53390. Examples of suitable fructokinases are Genbank accession numbers P26984, P26420 and CAA43322.

In one embodiment, the present invention provides a *Prototheca* host cell that secretes a sucrose invertase. Secretion of a sucrose invertase obviates the need for expression of a transporter that can transport sucrose into the cell. This is because a secreted invertase catalyzes the conversion of a molecule of sucrose into a molecule of glucose and a molecule of fructose, both of which can be transported and utilized by microbes provided by the invention. For example, expression of a sucrose invertase (such as SEQ ID NO:3) with a secretion signal (such as that of SEQ ID NO: 4 (from yeast), SEQ ID NO: 5 (from higher plants), SEQ ID NO: 6 (eukaryotic consensus secretion signal), and SEQ ID NO: 7 (combination of signal sequence from higher plants and eukaryotic consensus) generates invertase activity outside the cell. Expression of such a protein, as enabled by the genetic engineering methodology disclosed herein, allows cells already capable of utilizing extracellular glucose as an energy source to utilize sucrose as an extracellular energy source.

*Prototheca* species expressing an invertase in media containing sucrose are a preferred microalgal species for the production of oil. Example 3 illustrates how the methods and reagents of the invention can be used to express a recombinant yeast invertase and secrete it from a recombinant *Prototheca* cell. The expression and extracellular targeting of this fully active protein allows the resulting host cells to grow on sucrose, whereas their non-transformed counterparts cannot. Thus, the present invention provides *Prototheca* recombinant cells with a codon-optimized invertase gene, including but not limited to the yeast invertase gene, integrated into their genome such that the invertase gene is expressed as assessed by invertase activity and sucrose hydrolysis. The present invention also provides invertase genes useful as selectable markers in *Prototheca* recombinant cells, as such cells are able to grow on sucrose, while their non-transformed counterparts cannot; and methods for selecting recombinant host cells using an invertase as a powerful, selectable marker for algal molecular genetics.

The successful expression of a sucrose invertase in *Prototheca* also illustrates another aspect of the present invention in that it demonstrates that heterologous (recombinant) proteins can be expressed in the algal cell and successfully transit outside of the cell and into the culture medium in a fully active and functional form. Thus, the present invention provides methods and reagents for expressing a wide and diverse array of heterologous proteins in microalgae and secreting them outside of the host cell. Such proteins include, for example, industrial enzymes such as, for example, lipases, proteases, cellulases, pectinases, amylases, esterases, oxidoreductases, transferases, lactases, isomerases, and invertases, as well as

therapeutic proteins such as, for example, growth factors, cytokines, full length antibodies comprising two light and two heavy chains, Fabs, scFvs (single chain variable fragment), camellid-type antibodies, antibody fragments, antibody fragment-fusions, antibody-receptor fusions, insulin, interferons, and insulin-like growth factors.

The successful expression of a sucrose invertase in *Prototheca* also illustrates another aspect of the present invention in that it provides methods and reagents for the use of fungal transit peptides in algae to direct secretion of proteins in *Prototheca*; and methods and reagents for determining if a peptide can function, and the ability of it to function, as a transit peptide in *Prototheca* cells. The methods and reagents of the invention can be used as a tool and platform to identify other transit peptides that can successfully traffic proteins outside of a cell, and that the yeast invertase has great utility in these methods. As demonstrated in this example, removal of the endogenous yeast invertase transit peptide and its replacement by other transit peptides, either endogenous to the host algae or from other sources (eukaryotic, prokaryotic and viral), can identify whether any peptide of interest can function as a transit peptide in guiding protein egress from the cell.

Examples of suitable sucrose invertases include those identified by Genbank accession numbers CAB95010, NP\_012104 and CAA06839. Non-limiting examples of suitable invertases are listed below in Table 2. Amino acid sequences for each listed invertase are included in the Sequence Listing below. In some cases, the exogenous sucrose utilization gene suitable for use in the methods and vectors of the invention encodes a sucrose invertase that has at least 40, 50, 60, 75, or 90% or higher amino acid identity with a sucrose invertase selected from Table 2.

TABLE 2

Sucrose invertases.			
Description	Organism	GenBank Accession No.	SEQ ID NO:
Invertase	<i>Chicorium intybus</i>	Y11124	SEQ ID NO: 20
Invertase	<i>Schizosaccharomyces pombe</i>	AB011433	SEQ ID NO: 21
beta-fructofuranosidase (invertase)	<i>Pichia anomala</i>	X80640	SEQ ID NO: 22
Invertase	<i>Debaryomyces occidentalis</i>	X17604	SEQ ID NO: 23
Invertase	<i>Oryza sativa</i>	AF019113	SEQ ID NO: 24
Invertase	<i>Allium cepa</i>	AJ006067	SEQ ID NO: 25
Invertase	<i>Beta vulgaris</i> subsp. <i>Vulgaris</i>	AJ278531	SEQ ID NO: 26
beta-fructofuranosidase (invertase)	<i>Bifidobacterium breve</i> UCC2003	AAT28190	SEQ ID NO: 27
Invertase	<i>Saccharomyces cerevisiae</i>	NP_012104	SEQ ID NO: 8 (nucleotide) SEQ ID NO: 28 (amino acid)
Invertase A	<i>Zymomonas mobilis</i>	AAO38865	SEQ ID NO: 29

The secretion of an invertase to the culture medium by *Prototheca* enable the cells to grow as well on waste molasses from sugar cane processing as they do on pure reagent-grade glucose; the use of this low-value waste product of sugar cane processing can provide significant cost savings in the production of lipids and other oils. Thus, the present invention provides a microbial culture containing a population of *Prototheca* microorganisms, and a culture medium comprising (i) sucrose and (ii) a sucrose invertase enzyme. In various embodiments the sucrose in the culture comes from sorghum, sugar beet, sugar cane, molasses, or depolymerized cellulosic material (which may optionally contain lignin). In another aspect, the methods and reagents of the invention significantly increase the number and type of feedstocks that can be utilized by recombinant *Prototheca*. While the microbes

exemplified here are altered such that they can utilize sucrose, the methods and reagents of the invention can be applied so that feedstocks such as cellulose are utilizable by an engineered host microbe of the invention with the ability to secrete cellulases, pectinases, isomerases, or the like, such that the breakdown products of the enzymatic reactions are no longer just simply tolerated but rather utilized as a carbon source by the host.

## V. LIPID PATHWAY ENGINEERING

In addition to altering the ability of *Prototheca* to utilize feedstocks such as sucrose-containing feedstocks, the present invention also provides recombinant *Prototheca* that have been modified to alter the properties and/or proportions of lipids produced. The pathway can further, or alternatively, be modified to alter the properties and/or proportions of various lipid molecules produced through enzymatic processing of lipids and intermediates in the fatty acid pathway. In various embodiments, the recombinant *Prototheca* cells of the invention have, relative to their untransformed counterparts, optimized lipid yield per unit volume and/or per unit time, carbon chain length (e.g., for renewable diesel production or for industrial chemicals applications requiring lipid feedstock), reduced number of double or triple bonds, optionally to zero, and increasing the hydrogen:carbon ratio of a particular species of lipid or of a population of distinct lipid.

In particular embodiments, one or more key enzymes that control branch points in metabolism to fatty acid synthesis have been up-regulated or down-regulated to improve lipid production. Up-regulation can be achieved, for example, by transforming cells with expression constructs in which a gene encoding the enzyme of interest is expressed, e.g., using a

strong promoter and/or enhancer elements that increase transcription. Such constructs can include a selectable marker such that the transformants can be subjected to selection, which can result in amplification of the construct and an increase in the expression level of the encoded enzyme. Examples of enzymes suitable for up-regulation according to the methods of the invention include pyruvate dehydrogenase, which plays a role in converting pyruvate to acetyl-CoA (examples, some from microalgae, include Genbank accession numbers NP\_415392; AAA53047; Q1XDM1; and CAF05587). Up-regulation of pyruvate dehydrogenase can increase production of acetyl-CoA, and thereby increase fatty acid synthesis. Acetyl-CoA carboxylase catalyzes the initial step in fatty acid synthesis. Accordingly, this enzyme can be up-regulated to increase production of fatty acids (examples,

some from microalgae, include Genbank accession numbers BAA94752; AAA75528; AAA81471; YP\_537052; YP\_536879; NP\_045833; and BAA57908). Fatty acyl production can also be increased by up-regulation of acyl carrier protein (ACP), which carries the growing acyl chains during fatty acid synthesis (examples, some from microalgae, include Genbank accession numbers A0TOF8; P51280; NP\_849041; YP\_874433). Glycerol-3-phosphate acyltransferase catalyzes the rate-limiting step of fatty acid synthesis. Up-regulation of this enzyme can increase fatty acid production (examples, some from microalgae, include Genbank accession numbers AAA74319; AAA33122; AAA37647; P44857; and ABO94442).

Up- and/or down-regulation of genes can be applied to global regulators controlling the expression of the genes of the fatty acid biosynthetic pathways. Accordingly, one or more global regulators of fatty acid synthesis can be up- or down-regulated, as appropriate, to inhibit or enhance, respectively, the expression of a plurality of fatty acid synthetic genes and, ultimately, to increase lipid production. Examples include sterol regulatory element binding proteins (SREBPs), such as SREBP-1a and SREBP-1c (for examples see Genbank accession numbers NP\_035610 and Q9WTN3).

The present invention also provides recombinant *Prototheca* cells that have been modified to contain one or more exogenous genes encoding lipid modification enzymes such as, for example, fatty acyl-ACP thioesterases (see Table 3), fatty acyl-CoA/aldehyde reductases (see Table 4), fatty acyl-CoA reductases (see Table 5), fatty aldehyde decarboxylase (see Table 6), fatty aldehyde reductases, and squalene synthases (see GenBank Accession number AF205791). In some embodiments, genes encoding a fatty acyl-ACP thioesterase and a naturally co-expressed acyl carrier protein are transformed into a *Prototheca* cell, optionally with one or more genes encoding other lipid modification enzymes. In other embodiments, the ACP and the fatty acyl-ACP thioesterase may have an affinity for one another that imparts an advantage when the two are used together in the microbes and methods of the present invention, irrespective of whether they are or are not naturally co-expressed in a particular tissue or organism. Thus, the present invention contemplates both naturally co-expressed pairs of these enzymes as well as those that share an affinity for interacting with one another to facilitate cleavage of a length-specific carbon chain from the ACP.

In still other embodiments, an exogenous gene encoding a desaturase is transformed into the *Prototheca* cell in conjunction with one or more genes encoding other lipid modification enzymes to provide modifications with respect to lipid saturation. Stearoyl-ACP desaturase (see, e.g., GenBank Accession numbers AAF15308; ABM45911; and AAY86086), for example, catalyzes the conversion of stearoyl-ACP to oleoyl-ACP. Up-regulation of this gene can increase the proportion of monounsaturated fatty acids produced by a cell; whereas down-regulation can reduce the proportion of monounsaturates. Similarly, the expression of one or more glycerolipid desaturases can be controlled to alter the ratio of unsaturated to saturated fatty acids such as  $\omega$ -6 fatty acid desaturase,  $\omega$ -3 fatty acid desaturase, or  $\omega$ -6-oleate desaturase. In some embodiments, the desaturase can be selected with reference to a desired carbon chain length, such that the desaturase is capable of making location specific modifications within a specified carbon-length substrate, or substrates having a carbon-length within a specified range.

Thus, in particular embodiments, microbes of the present invention are genetically engineered to express one or more exogenous genes selected from an acyl-ACP thioesterase, an acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a

fatty aldehyde reductase, a fatty aldehyde decarboxylase, or a naturally co-expressed acyl carrier protein. Suitable expression methods are described above with respect to the expression of a lipase gene, including, among other methods, inducible expression and compartmentalized expression. A fatty acyl-ACP thioesterase cleaves a fatty acid from an acyl carrier protein (ACP) during lipid synthesis. Through further enzymatic processing, the cleaved fatty acid is then combined with a coenzyme to yield an acyl-CoA molecule. This acyl-CoA is the substrate for the enzymatic activity of a fatty acyl-CoA reductase to yield an aldehyde, as well as for a fatty acyl-CoA/aldehyde reductase to yield an alcohol. The aldehyde produced by the action of the fatty acyl-CoA reductase identified above is the substrate for further enzymatic activity by either a fatty aldehyde reductase to yield an alcohol, or a fatty aldehyde decarboxylase to yield an alkane or alkene.

In some embodiments, fatty acids, glycerolipids, or the corresponding primary alcohols, aldehydes, alkanes or alkenes, generated by the methods described herein, contain 8, 10, 12, or 14 carbon atoms. Preferred fatty acids for the production of diesel, biodiesel, renewable diesel, or jet fuel, or the corresponding primary alcohols, aldehydes, alkanes and alkenes, for industrial applications contain 8 to 14 carbon atoms. In certain embodiments, the above fatty acids, as well as the other corresponding hydrocarbon molecules, are saturated (with no carbon-carbon double or triple bonds); mono unsaturated (single double bond); poly unsaturated (two or more double bonds); are linear (not cyclic) or branched. For fuel production, greater saturation is preferred.

The enzymes described directly above have a preferential specificity for hydrolysis of a substrate containing a specific number of carbon atoms. For example, a fatty acyl-ACP thioesterase may have a preference for cleaving a fatty acid having 12 carbon atoms from the ACP. In some embodiments, the ACP and the length-specific thioesterase may have an affinity for one another that makes them particularly useful as a combination (e.g., the exogenous ACP and thioesterase genes may be naturally co-expressed in a particular tissue or organism from which they are derived). Therefore, in various embodiments, the recombinant *Prototheca* cell of the invention can contain an exogenous gene that encodes a protein with specificity for catalyzing an enzymatic activity (e.g., cleavage of a fatty acid from an ACP, reduction of an acyl-CoA to an aldehyde or an alcohol, or conversion of an aldehyde to an alkane) with regard to the number of carbon atoms contained in the substrate. The enzymatic specificity can, in various embodiments, be for a substrate having from 8 to 34 carbon atoms, preferably from 8 to 18 carbon atoms, and more preferably from 8 to 14 carbon atoms. A preferred specificity is for a substrate having fewer, i.e., 12, rather than more, i.e., 18, carbon atoms.

In non-limiting but illustrative examples, the present invention provides vectors and *Prototheca* host cells that express an exogenous thioesterase and accordingly produce lipid enriched, relative to the lipid profile of untransformed *Prototheca* cells, in the chain length for which the thioesterase is specific. The thioesterases illustrated are (i) *Cinnamomum camphorum* FatB1 (GenBank Accession No. Q39473, amino acid sequence is in SEQ ID NO: 59, amino acid sequence without plastid targeting sequence (PTS) is in SEQ ID NO: 139, and codon optimized cDNA sequence based on Table 1 is in SEQ ID NO: 60), which has a preference for fatty acyl-ACP substrate with a carbon chain length of 14; (ii) *Cuphea hookeriana* FatB2 (GenBank Accession No. AAC49269, amino acid sequence is in SEQ ID NO: 61, amino acid sequence without PTS is in SEQ ID NO: 138, and codon optimized cDNA sequence based on Table 1 is in SEQ

ID NO: 62), which has a preference for a fatty acyl-ACP substrate with a carbon chain length of 8-10; and (iii) *Umbellularia* Fat B1 (GenBank Accession No. Q41635, amino acid sequence is included in SEQ ID NO: 63, amino acid sequence without PTS is in SEQ ID NO: 139, and codon optimized cDNA sequence based on Table 1 is included in SEQ ID NO: 64), which has a preference for a fatty acyl-ACP substrate with a carbon chain length of 12.

Other fatty acyl-ACP thioesterases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 3.

TABLE 3

Fatty acyl-ACP thioesterases and GenBank accession numbers.
<i>Umbellularia californica</i> fatty acyl-ACP thioesterase (GenBank #AAC49001)
<i>Cinnamomum camphora</i> fatty acyl-ACP thioesterase (GenBank #Q39473)
<i>Umbellularia californica</i> fatty acyl-ACP thioesterase (GenBank #Q41635)
<i>Myristica fragrans</i> fatty acyl-ACP thioesterase (GenBank #AAB71729)
<i>Myristica fragrans</i> fatty acyl-ACP thioesterase (GenBank #AAB71730)
<i>Elaeis guineensis</i> fatty acyl-ACP thioesterase (GenBank #ABD83939)
<i>Elaeis guineensis</i> fatty acyl-ACP thioesterase (GenBank #AAD42220)
<i>Populus tomentosa</i> fatty acyl-ACP thioesterase (GenBank #ABC47311)
<i>Arabidopsis thaliana</i> fatty acyl-ACP thioesterase (GenBank #NP_172327)
<i>Arabidopsis thaliana</i> fatty acyl-ACP thioesterase (GenBank #CAA85387)
<i>Arabidopsis thaliana</i> fatty acyl-ACP thioesterase (GenBank #CAA85388)
<i>Gossypium hirsutum</i> fatty acyl-ACP thioesterase (GenBank #Q9S0I3)
<i>Cuphea lanceolata</i> fatty acyl-ACP thioesterase (GenBank #CAA54060)
<i>Cuphea hookeriana</i> fatty acyl-ACP thioesterase (GenBank #AAC72882)
<i>Cuphea calophytta</i> subsp. <i>mesostemon</i> fatty acyl-ACP thioesterase (GenBank #ABB71581)
<i>Cuphea lanceolata</i> fatty acyl-ACP thioesterase (GenBank #CAC19933)
<i>Elaeis guineensis</i> fatty acyl-ACP thioesterase (GenBank #AAL15645)
<i>Cuphea hookeriana</i> fatty acyl-ACP thioesterase (GenBank #Q39513)
<i>Gossypium hirsutum</i> fatty acyl-ACP thioesterase (GenBank #AAD01982)
<i>Vitis vinifera</i> fatty acyl-ACP thioesterase (GenBank #CAN81819)
<i>Garcinia mangostana</i> fatty acyl-ACP thioesterase (GenBank #AAB51525)
<i>Brassica juncea</i> fatty acyl-ACP thioesterase (GenBank #ABI18986)
<i>Madhuca longifolia</i> fatty acyl-ACP thioesterase (GenBank #AA051637)
<i>Brassica napus</i> fatty acyl-ACP thioesterase (GenBank #ABH11710)
<i>Oryza sativa</i> ( <i>indica</i> cultivar-group) fatty acyl-ACP thioesterase (GenBank #EAY86877)
<i>Oryza sativa</i> ( <i>japonica</i> cultivar-group) fatty acyl-ACP thioesterase (GenBank #NP_001068400)
<i>Oryza sativa</i> ( <i>indica</i> cultivar-group) fatty acyl-ACP thioesterase (GenBank #EAY99617)
<i>Cuphea hookeriana</i> fatty acyl-ACP thioesterase (GenBank #AAC49269)
<i>Ulmus Americana</i> fatty acyl-ACP thioesterase (GenBank #AAB71731)
<i>Cuphea lanceolata</i> fatty acyl-ACP thioesterase (GenBank #CAB60830)
<i>Cuphea palustris</i> fatty acyl-ACP thioesterase (GenBank #AAC49180)
<i>Iris germanica</i> fatty acyl-ACP thioesterase (GenBank #AAG43858)
<i>Cuphea palustris</i> fatty acyl-ACP thioesterase (GenBank #AAC49179)
<i>Myristica fragrans</i> fatty acyl-ACP thioesterase (GenBank #AAB71729)
<i>Cuphea hookeriana</i> fatty acyl-ACP thioesterase (GenBank #U39834)
<i>Umbellularia californica</i> fatty acyl-ACP thioesterase (GenBank #M94159)
<i>Cinnamomum camphora</i> fatty acyl-ACP thioesterase (GenBank #U31813)

The Examples below describe the successful targeting and expression of heterologous fatty acyl-ACP thioesterases from *Cuphea hookeriana*, *Umbellularia californica*, *Cinnamomum camphora* in *Prototheca* species. Additionally, alterations in fatty acid profiles were confirmed in the host cells expression these heterologous fatty acyl-ACP thioesterases. These results were quite unexpected given the lack of sequence identity between algal and higher plant thioesterases in general, and between *Prototheca moriformis* fatty acyl-ACP thioesterase and the above listed heterologous fatty acyl-ACP thioesterases. Two *Prototheca moriformis* acyl-ACP thioesterases were isolated and sequenced. The sequences of the two cDNAs showed a high degree of identity between each other, differing in only 12 positions at the nucleotide level and five positions at the amino acid level, four of these in the plastid transit peptide. Further analysis of genomic sequence from *Prototheca moriformis* confirmed that these two cDNAs were indeed encoded on separate contigs, and although highly homologous, are encoded by two distinct genes. The cDNA and amino acid sequence of the two *Prototheca moriformis* fatty acyl-ACP thioesterase, *P. moriformis* fatty acyl-ACP thioesterase-1 and *P. moriformis* fatty acyl-ACP thioesterase-2, are listed as SEQ ID NOs: 134-137.

When the amino acid sequences of these two cDNAs were BLASTed against the NCBI database, the two most homologous sequences were fatty acyl-ACP thioesterases from *Chlamydomonas reinhardtii* and *Arabidopsis thaliana*. Surprisingly, the level of amino acid identity between the *Prototheca moriformis* fatty acyl-ACP thioesterases and higher plant thioesterases was fairly low, at only 49 and 37% identity. In addition, there also is a subtle difference in the sequences surrounding the amino terminal portion of the catalytic triad (NXHX<sub>3,6</sub>C) among these fatty acyl-ACP thioesterases. Thirty nine of forty higher plant fatty acyl-ACP thioesterases surveyed showed the sequence LD MNQH surrounding the N and H residues at the amino terminus of the triad, while all of the algal sequences identified had the sequence MD MN GH. Given the low amino acid sequence identity and the differences surrounding the catalytic triad of the thioesterases, the successful results of expression of exogenous fatty acyl-ACP thioesterases obtained and described in the Examples were unexpected, particularly given the fact that activity of the exogenous fatty acyl-ACP thioesterases was dependent on a functional protein-protein interaction with the endogenous *Prototheca* acyl carrier protein.

Fatty acyl-CoA/aldehyde reductases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 4.

TABLE 4

Fatty acyl-CoA/aldehyde reductases listed by GenBank accession numbers.
AAC45217, YP_047869, BAB85476, YP_001086217, YP_580344, YP_001280274, YP_264583, YP_436109, YP_959769, ZP_01736962, ZP_01900335, ZP_01892096, ZP_01103974, ZP_01915077, YP_924106, YP_130411, ZP_01222731, YP_550815, YP_983712, YP_001019688, YP_524762, YP_856798, ZP_01115500, YP_001141848, NP_336047, NP_216059, YP_882409, YP_706156, YP_001136150, YP_952365, ZP_01221833, YP_130076, NP_567936, AAR88762, ABK28586, NP_197634, CAD30694, NP_001063962, BAD46254, NP_001030809, EAZ10132, EAZ43639, EAZ07989, NP_001062488, CAB88537, NP_001052541, CAH66597, CAE02214, CAH66590, CAB88538, EAZ39844, AAZ06658, CAA68190, CAA52019, and BAC84377

Fatty acyl-CoA reductases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 5.

TABLE 5

Fatty acyl-CoA reductases listed by GenBank accession numbers.
NP_187805, ABO14927, NP_001049083, CAN83375, NP_191229, EAZ42242, EAZ06453, CAD30696, BAD31814, NP_190040, AAD38039, CAD30692, CAN81280, NP_197642, NP_190041, AAL15288, and NP_190042

Fatty aldehyde decarboxylases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 6.

TABLE 6

Fatty aldehyde decarboxylases listed by GenBank accession numbers.
NP_850932, ABN07985, CAN60676, AAC23640, CAA65199, AAC24373, CAE03390, ABD28319, NP_181306, EAZ31322, CAN63491, EAY94825, EAY86731, CAL55686, XP_001420263, EAZ23849, NP_200588, NP_001063227, CAN83072, AAR90847, and AAR97643

Combinations of naturally co-expressed fatty acyl-ACP thioesterases and acyl carrier proteins are suitable for use with the microbes and methods of the invention.

Additional examples of hydrocarbon or lipid modification enzymes include amino acid sequences contained in, referenced in, or encoded by nucleic acid sequences contained or referenced in, any of the following U.S. Pat. Nos. 6,610,527; 6,451,576; 6,429,014; 6,342,380; 6,265,639; 6,194,185; 6,114,160; 6,083,731; 6,043,072; 5,994,114; 5,891,697; 5,871,988; 6,265,639, and further described in GenBank Accession numbers: AAO18435; ZP\_00513891; Q38710; AAK60613; AAK60610; AAK60611; NP\_113747; CAB75874; AAK60612; AAF20201; BAA11024; AF205791; and CAA03710.

Other suitable enzymes for use with the microbes and the methods of the invention include those that have at least 70% amino acid identity with one of the proteins listed in Tables 3-6, and that exhibit the corresponding desired enzymatic activity (e.g., cleavage of a fatty acid from an acyl carrier protein, reduction of an acyl-CoA to an aldehyde or an alcohol, or conversion of an aldehyde to an alkane). In additional embodiments, the enzymatic activity is present in a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% identity with one of the above described sequences, all of which are hereby incorporated by reference as if fully set forth.

By selecting the desired combination of exogenous genes to be expressed, one can tailor the product generated by the microbe, which may then be extracted from the aqueous biomass. For example, the microbe can contain: (i) an exogenous gene encoding a fatty acyl-ACP thioesterase; and, optionally, (ii) a naturally co-expressed acyl carrier protein or an acyl carrier protein otherwise having affinity for the fatty acyl-ACP thioesterase (or conversely); and, optionally, (iii) an exogenous gene encoding a fatty acyl-CoA/aldehyde reductase or a fatty acyl-CoA reductase; and, optionally, (iv) an exogenous gene encoding a fatty aldehyde reductase or a fatty aldehyde decarboxylase. The microbe, under culture conditions described herein, synthesizes a fatty acid linked to an ACP and the fatty acyl-ACP thioesterase catalyzes the

cleavage of the fatty acid from the ACP to yield, through further enzymatic processing, a fatty acyl-CoA molecule. When present, the fatty acyl-CoA/aldehyde reductase catalyzes the reduction of the acyl-CoA to an alcohol. Similarly, the fatty acyl-CoA reductase, when present, catalyzes the reduction of the acyl-CoA to an aldehyde. In those embodiments in which an exogenous gene encoding a fatty acyl-CoA reductase is present and expressed to yield an aldehyde product, a fatty aldehyde reductase, encoded by the third exogenous gene, catalyzes the reduction of the aldehyde to an alcohol. Similarly, a fatty aldehyde decarboxylase catalyzes the conversion of the aldehyde to an alkane or an alkene, when present.

Genes encoding such enzymes can be obtained from cells already known to exhibit significant lipid production such as *Chlorella protothecoides*. Genes already known to have a role in lipid production, e.g., a gene encoding an enzyme that saturates double bonds, can be transformed individually into recipient cells. However, to practice the invention it is not necessary to make a priori assumptions as to which genes are required. Methods for identifying genes that can alter (improve) lipid production in microalgae are described in PCT Pub. No. 2008/151149.

Thus, the present invention provides a *Prototheca* cell that has been genetically engineered to express a lipid pathway enzyme at an altered level compared to a wild-type cell of the same species. In some cases, the cell produces more lipid compared to the wild-type cell when both cells are grown under the same conditions. In some cases, the cell has been genetically engineered and/or selected to express a lipid pathway enzyme at a higher level than the wild-type cell. In some cases, the lipid pathway enzyme is selected from the group consisting of pyruvate dehydrogenase, acetyl-CoA carboxylase, acyl carrier protein, and glycerol-3 phosphate acyltransferase. In some cases, the cell has been genetically engineered and/or selected to express a lipid pathway enzyme at a lower level than the wild-type cell. In at least one embodiment in which the cell expresses the lipid pathway enzyme at a lower level, the lipid pathway enzyme comprises citrate synthase.

In some embodiments, the cell has been genetically engineered and/or selected to express a global regulator of fatty acid synthesis at an altered level compared to the wild-type cell, whereby the expression levels of a plurality of fatty acid synthetic genes are altered compared to the wild-type cell. In some cases, the lipid pathway enzyme comprises an enzyme that modifies a fatty acid. In some cases, the lipid pathway enzyme is selected from a stearoyl-ACP desaturase and a glycerolipid desaturase.

In other embodiments, the present invention is directed to an oil-producing microbe containing one or more exogenous genes, wherein the exogenous genes encode protein(s) selected from the group consisting of a fatty acyl-ACP thioesterase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty acyl-CoA/aldehyde reductase, a fatty aldehyde decarboxylase, and an acyl carrier protein. In one embodiment, the exogenous gene is in operable linkage with a promoter, which is inducible or repressible in response to a stimulus. In some cases, the stimulus is selected from the group consisting of an exogenously provided small molecule, heat, cold, and limited or no nitrogen in the culture media. In some cases, the exogenous gene is expressed in a cellular compartment. In some embodiments, the cellular compartment is selected from the group consisting of a chloroplast, a plastid and a mitochondrion. In some embodiments the microbe is *Prototheca moriformis*, *Prototheca krugani*, *Prototheca stagnora* or *Prototheca zopfii*.

In one embodiment, the exogenous gene encodes a fatty acid acyl-ACP thioesterase. In some cases, the thioesterase encoded by the exogenous gene catalyzes the cleavage of an 8 to 18-carbon fatty acid from an acyl carrier protein (ACP). In some cases, the thioesterase encoded by the exogenous gene catalyzes the cleavage of a 10 to 14-carbon fatty acid from an ACP. In one embodiment, the thioesterase encoded by the exogenous gene catalyzes the cleavage of a 12-carbon fatty acid from an ACP.

In one embodiment, the exogenous gene encodes a fatty acyl-CoA/aldehyde reductase. In some cases, the reductase encoded by the exogenous gene catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding primary alcohol. In some cases, the reductase encoded by the exogenous gene catalyzes the reduction of a 10 to 14-carbon fatty acyl-CoA to a corresponding primary alcohol. In one embodiment, the reductase encoded by the exogenous gene catalyzes the reduction of a 12-carbon fatty acyl-CoA to dodecanol.

The present invention also provides a recombinant *Prototheca* cell containing two exogenous genes, wherein a first exogenous gene encodes a fatty acyl-ACP thioesterase and a second exogenous gene encodes a protein selected from the group consisting of a fatty acyl-CoA reductase, a fatty acyl-CoA/aldehyde reductase, and an acyl carrier protein. In some cases, the two exogenous genes are each in operable linkage with a promoter, which is inducible in response to a stimulus. In some cases, each promoter is inducible in response to an identical stimulus, such as limited or no nitrogen in the culture media. Limitation or complete lack of nitrogen in the culture media stimulates oil production in some microorganisms such as *Prototheca* species, and can be used as a trigger to induce oil production to high levels. When used in combination with the genetic engineering methods disclosed herein, the lipid as a percentage of dry cell weight can be pushed to high levels such as at least 30%, at least 40%, at least 50%, at least 60%, at least 70% and at least 75%; methods disclosed herein provide for cells with these levels of lipid, wherein the lipid is at least 4% C8-C14, at least 0.3% C8, at least 2% C10, at least 2% C12, and at least 2% C14. In some embodiments the cells are over 25% lipid by dry cell weight and contain lipid that is at least 10% C8-C14, at least 20% C8-C14, at least 30% C8-C14, 10-30% C8-C14 and 20-30% C8-C14.

The novel oils disclosed herein are distinct from other naturally occurring oils that are high in mic-chain fatty acids, such as palm oil, palm kernel oil, and coconut oil. For example, levels of contaminants such as carotenoids are far higher in palm oil and palm kernel oil than in the oils of the invention. Palm and palm kernel oils in particular contain alpha and beta carotenes and lycopene in much higher amounts than is in the oils of the invention. In addition, over 20 different carotenoids are found in palm and palm kernel oil, whereas the Examples demonstrate that the oils of the invention contain very few carotenoids species and very low levels. In addition, the levels of vitamin E compounds such as tocotrienols are far higher in palm, palm kernel, and coconut oil than in the oils of the invention.

In one embodiment, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of an 8 to 18-carbon fatty acid from an ACP. In some embodiments, the second exogenous gene encodes a fatty acyl-CoA/aldehyde reductase which catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding primary alcohol. In some cases, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of a 10 to 14-carbon fatty acid from an ACP, and the reductase encoded by the second exogenous gene catalyzes the reduction of a 10 to 14-carbon fatty acyl-CoA to

the corresponding primary alcohol, wherein the thioesterase and the reductase act on the same carbon chain length. In one embodiment, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of a 12-carbon fatty acid from an ACP, and the reductase encoded by the second exogenous gene catalyzes the reduction of a 12-carbon fatty acyl-CoA to dodecanol. In some embodiments, the second exogenous gene encodes a fatty acyl-CoA reductase which catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding aldehyde. In some embodiments, the second exogenous gene encodes an acyl carrier protein that is naturally co-expressed with the fatty acyl-ACP thioesterase.

In some embodiments, the second exogenous gene encodes a fatty acyl-CoA reductase, and the microbe further contains a third exogenous gene encoding a fatty aldehyde decarboxylase. In some cases, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of an 8 to 18-carbon fatty acid from an ACP, the reductase encoded by the second exogenous gene catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding-fatty aldehyde, and the decarboxylase encoded by the third exogenous gene catalyzes the conversion of an 8 to 18-carbon fatty aldehyde to a corresponding alkane, wherein the thioesterase, the reductase, and the decarboxylase act on the same carbon chain length.

In some embodiments, the second exogenous gene encodes an acyl carrier protein, and the microbe further contains a third exogenous gene encoding a protein selected from the group consisting of a fatty acyl-CoA reductase and a fatty acyl-CoA/aldehyde reductase. In some cases, the third exogenous gene encodes a fatty acyl-CoA reductase, and the microbe further contains a fourth exogenous gene encoding a fatty aldehyde decarboxylase.

The present invention also provides methods for producing an alcohol comprising culturing a population of recombinant *Prototheca* cells in a culture medium, wherein the cells contain (i) a first exogenous gene encoding a fatty acyl-ACP thioesterase, and (ii) a second exogenous gene encoding a fatty acyl-CoA/aldehyde reductase, and the cells synthesize a fatty acid linked to an acyl carrier protein (ACP), the fatty acyl-ACP thioesterase catalyzes the cleavage of the fatty acid from the ACP to yield, through further processing, a fatty acyl-CoA, and the fatty acyl-CoA/aldehyde reductase catalyzes the reduction of the acyl-CoA to an alcohol.

The present invention also provides methods of producing a lipid molecule in a *Prototheca* cell. In one embodiment, the method comprises culturing a population of *Prototheca* cells in a culture medium, wherein the cells contain (i) a first exogenous gene encoding a fatty acyl-ACP thioesterase, and (ii) a second exogenous gene encoding a fatty acyl-CoA reductase, and wherein the microbes synthesize a fatty acid linked to an acyl carrier protein (ACP), the fatty acyl-ACP thioesterase catalyzes the cleavage of the fatty acid from the ACP to yield, through further processing, a fatty acyl-CoA, and the fatty acyl-CoA reductase catalyzes the reduction of the acyl-CoA to an aldehyde.

The present invention also provides methods of producing a fatty acid molecule having a specified carbon chain length in a *Prototheca* cell. In one embodiment, the method comprises culturing a population of lipid-producing *Prototheca* cells in a culture medium, wherein the microbes contain an exogenous gene encoding a fatty acyl-ACP thioesterase having an activity specific or preferential to a certain carbon chain length, such as 8, 10, 12 or 14 carbon atoms, and wherein the microbes synthesize a fatty acid linked to an acyl carrier protein (ACP) and the thioesterase catalyzes the cleavage of

the fatty acid from the ACP when the fatty acid has been synthesized to the specific carbon chain length.

In the various embodiments described above, the *Prototheca* cell can contain at least one exogenous gene encoding a lipid pathway enzyme. In some cases, the lipid pathway enzyme is selected from the group consisting of a stearyl-ACP desaturase, a glycerolipid desaturase, a pyruvate dehydrogenase, an acetyl-CoA carboxylase, an acyl carrier protein, and a glycerol-3 phosphate acyltransferase. In other cases, the *Prototheca* cell contains a lipid modification enzyme selected from the group consisting of a fatty acyl-ACP thioesterase, a fatty acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty aldehyde decarbonylase, and/or an acyl carrier protein.

## VI. FUELS AND CHEMICALS PRODUCTION

For the production of fuel in accordance with the methods of the invention lipids produced by cells of the invention are harvested, or otherwise collected, by any convenient means. Lipids can be isolated by whole cell extraction. The cells are first disrupted, and then intracellular and cell membrane/cell wall-associated lipids as well as extracellular hydrocarbons can be separated from the cell mass, such as by use of centrifugation as described above. Intracellular lipids produced in microorganisms are, in some embodiments, extracted after lysing the cells of the microorganism. Once extracted, the lipids are further refined to produce oils, fuels, or oleochemicals.

After completion of culturing, the microorganisms can be separated from the fermentation broth. Optionally, the separation is effected by centrifugation to generate a concentrated paste. Centrifugation does not remove significant amounts of intracellular water from the microorganisms and is not a drying step. The biomass can then optionally be washed with a washing solution (e.g., DI water) to get rid of the fermentation broth and debris. Optionally, the washed microbial biomass may also be dried (oven dried, lyophilized, etc.) prior to cell disruption. Alternatively, cells can be lysed without separation from some or all of the fermentation broth when the fermentation is complete. For example, the cells can be at a ratio of less than 1:1 v:v cells to extracellular liquid when the cells are lysed.

Microorganisms containing a lipid can be lysed to produce a lysate. As detailed herein, the step of lysing a microorganism (also referred to as cell lysis) can be achieved by any convenient means, including heat-induced lysis, adding a base, adding an acid, using enzymes such as proteases and polysaccharide degradation enzymes such as amylases, using ultrasound, mechanical lysis, using osmotic shock, infection with a lytic virus, and/or expression of one or more lytic genes. Lysis is performed to release intracellular molecules which have been produced by the microorganism. Each of these methods for lysing a microorganism can be used as a single method or in combination simultaneously or sequentially. The extent of cell disruption can be observed by microscopic analysis. Using one or more of the methods described herein, typically more than 70% cell breakage is observed. Preferably, cell breakage is more than 80%, more preferably more than 90% and most preferred about 100%.

In particular embodiments, the microorganism is lysed after growth, for example to increase the exposure of cellular lipid and/or hydrocarbon for extraction or further processing. The timing of lipase expression (e.g., via an inducible promoter) or cell lysis can be adjusted to optimize the yield of

lipids and/or hydrocarbons. Below are described a number of lysis techniques. These techniques can be used individually or in combination.

In one embodiment of the present invention, the step of lysing a microorganism comprises heating of a cellular suspension containing the microorganism. In this embodiment, the fermentation broth containing the microorganisms (or a suspension of microorganisms isolated from the fermentation broth) is heated until the microorganisms, i.e., the cell walls and membranes of microorganisms degrade or breakdown. Typically, temperatures applied are at least 50° C. Higher temperatures, such as, at least 30° C. at least 60° C., at least 70° C., at least 80° C., at least 90° C., at least 100° C., at least 110° C., at least 120° C., at least 130° C. or higher are used for more efficient cell lysis. Lysing cells by heat treatment can be performed by boiling the microorganism. Alternatively, heat treatment (without boiling) can be performed in an autoclave. The heat treated lysate may be cooled for further treatment. Cell disruption can also be performed by steam treatment, i.e., through addition of pressurized steam. Steam treatment of microalgae for cell disruption is described, for example, in U.S. Pat. No. 6,750,048. In some embodiments, steam treatment may be achieved by sparging steam into the fermentor and maintaining the broth at a desired temperature for less than about 90 minutes, preferably less than about 60 minutes, and more preferably less than about 30 minutes.

In another embodiment of the present invention, the step of lysing a microorganism comprises adding a base to a cellular suspension containing the microorganism. The base should be strong enough to hydrolyze at least a portion of the proteinaceous compounds of the microorganisms used. Bases which are useful for solubilizing proteins are known in the art of chemistry. Exemplary bases which are useful in the methods of the present invention include, but are not limited to, hydroxides, carbonates and bicarbonates of lithium, sodium, potassium, calcium, and mixtures thereof. A preferred base is KOH. Base treatment of microalgae for cell disruption is described, for example, in U.S. Pat. No. 6,750,048.

In another embodiment of the present invention, the step of lysing a microorganism comprises adding an acid to a cellular suspension containing the microorganism. Acid lysis can be effected using an acid at a concentration of 10-500 mN or preferably 40-160 nM. Acid lysis is preferably performed at above room temperature (e.g., at 40-160°, and preferably a temperature of 50-130°. For moderate temperatures (e.g., room temperature to 100° C. and particularly room temperature to 65°, acid treatment can usefully be combined with sonication or other cell disruption methods.

In another embodiment of the present invention, the step of lysing a microorganism comprises lysing the microorganism by using an enzyme. Preferred enzymes for lysing a microorganism are proteases and polysaccharide-degrading enzymes such as hemicellulase (e.g., hemicellulase from *Aspergillus niger*; Sigma Aldrich, St. Louis, Mo.; #H2125), pectinase (e.g., pectinase from *Rhizopus* sp.; Sigma Aldrich, St. Louis, Mo.; #P2401), Mannaway 4.0 L (Novozymes), cellulase (e.g., cellulase from *Trichodeima viride*; Sigma Aldrich, St. Louis, Mo.; #C9422), and driselase (e.g., driselase from *Basidiomycetes* sp.; Sigma Aldrich, St. Louis, Mo.; #D9515).

In other embodiments of the present invention, lysis is accomplished using an enzyme such as, for example, a cellulase such as a polysaccharide-degrading enzyme, optionally from *Chlorella* or a *Chlorella* virus, or a proteases, such as *Streptomyces griseus* protease, chymotrypsin, proteinase K, proteases listed in Degradation of Polylactide by Commercial Proteases, Oda Y et al., Journal of Polymers and the



Environment, Volume 8, Number 1, January 2000, pp. 29-32 (4), Alcalase 2.4 FG (Novozymes), and Flavourzyme 100 L (Novozymes). Any combination of a protease and a polysaccharide-degrading enzyme can also be used, including any combination of the preceding proteases and polysaccharide-degrading enzymes.

In another embodiment, lysis can be performed using an expeller press. In this process, biomass is forced through a screw-type device at high pressure, lysing the cells and causing the intracellular lipid to be released and separated from the protein and fiber (and other components) in the cell.

In another embodiment of the present invention, the step of lysing a microorganism is performed by using ultrasound, i.e., sonication. Thus, cells can also be lysed with high frequency sound. The sound can be produced electronically and transported through a metallic tip to an appropriately concentrated cellular suspension. This sonication (or ultrasonication) disrupts cellular integrity based on the creation of cavi-

ties in cell suspension.

In another embodiment of the present invention, the step of lysing a microorganism is performed by mechanical lysis. Cells can be lysed mechanically and optionally homogenized to facilitate hydrocarbon (e.g., lipid) collection. For example, a pressure disrupter can be used to pump a cell containing slurry through a restricted orifice valve. High pressure (up to 1500 bar) is applied, followed by an instant expansion through an exiting nozzle. Cell disruption is accomplished by three different mechanisms: impingement on the valve, high liquid shear in the orifice, and sudden pressure drop upon discharge, causing an explosion of the cell. The method releases intracellular molecules. Alternatively, a ball mill can be used. In a ball mill, cells are agitated in suspension with small abrasive particles, such as beads. Cells break because of shear forces, grinding between beads, and collisions with beads. The beads disrupt the cells to release cellular contents. Cells can also be disrupted by shear forces, such as with the use of blending (such as with a high speed or Waring blender as examples), the french press, or even centrifugation in case of weak cell walls, to disrupt cells.

In another embodiment of the present invention, the step of lysing a microorganism is performed by applying an osmotic shock.

In another embodiment of the present invention, the step of lysing a microorganism comprises infection of the microorganism with a lytic virus. A wide variety of viruses are known to lyse microorganisms suitable for use in the present invention, and the selection and use of a particular lytic virus for a particular microorganism is within the level of skill in the art. For example, *paramecium bursaria chlorella* virus (PBCV-1) is the prototype of a group (family Phycodnaviridae, genus *Chlorovirus*) of large, icosahedral, plaque-forming, double-stranded DNA viruses that replicate in, and lyse, certain unicellular, eukaryotic *chlorella*-like green algae. Accordingly, any susceptible microalgae can be lysed by infecting the culture with a suitable *chlorella* virus. Methods of infecting species of *Chlorella* with a *chlorella* virus are known. See for example *Adv. Virus Res.* 2006; 66:293-336; *Virology*, 1999 Apr. 25; 257(1):15-23; *Virology*, 2004 Jan. 5; 318(1):214-23; *Nucleic Acids Symp. Ser.* 2000; (44):161-2; *J. Virol.* 2006 March; 80(5):2437-44; and *Annu. Rev. Microbiol.* 1999; 53:447-94.

In another embodiment of the present invention, the step of lysing a microorganism comprises autolysis. In this embodiment, a microorganism according to the invention is genetically engineered to produce a lytic protein that will lyse the microorganism. This lytic gene can be expressed using an inducible promoter so that the cells can first be grown to a

desirable density in a fermentor, followed by induction of the promoter to express the lytic gene to lyse the cells. In one embodiment, the lytic gene encodes a polysaccharide-degrading enzyme. In certain other embodiments, the lytic gene is a gene from a lytic virus. Thus, for example, a lytic gene from a *Chlorella* virus can be expressed in an algal cell; see *Virology* 260, 308-315 (1999); *FEMS Microbiology Letters* 180 (1999) 45-53; *Virology* 263, 376-387 (1999); and *Virology* 230, 361-368 (1997). Expression of lytic genes is preferably done using an inducible promoter, such as a promoter active in microalgae that is induced by a stimulus such as the presence of a small molecule, light, heat, and other stimuli.

Various methods are available for separating lipids from cellular lysates produced by the above methods. For example, lipids and lipid derivatives such as fatty aldehydes, fatty alcohols, and hydrocarbons such as alkanes can be extracted with a hydrophobic solvent such as hexane (see Frenz et al. 1989, *Enzyme Microb. Technol.*, 11:717). Lipids and lipid derivatives can also be extracted using liquefaction (see for example Sawayama et al. 1999, *Biomass and Bioenergy* 17:33-39 and Inoue et al. 1993, *Biomass Bioenergy* 6(4):269-274); oil liquefaction (see for example Minowa et al. 1995, *Fuel* 74(12): 1735-1738); and supercritical CO<sub>2</sub> extraction (see for example Mendes et al. 2003, *Inorganica Chimica Acta* 356: 328-334). Miao and Wu describe a protocol of the recovery of microalgal lipid from a culture of *Chlorella protothecoides* in which the cells were harvested by centrifugation, washed with distilled water and dried by freeze drying. The resulting cell powder was pulverized in a mortar and then extracted with n-hexane. Miao and Wu, *Biosource Technology* (2006) 97:841-846.

Thus, lipids, lipid derivatives and hydrocarbons generated by the microorganisms of the present invention can be recovered by extraction with an organic solvent. In some cases, the preferred organic solvent is hexane. Typically, the organic solvent is added directly to the lysate without prior separation of the lysate components. In one embodiment, the lysate generated by one or more of the methods described above is contacted with an organic solvent for a period of time sufficient to allow the lipid and/or hydrocarbon components to form a solution with the organic solvent. In some cases, the solution can then be further refined to recover specific desired lipid or hydrocarbon components. Hexane extraction methods are well known in the art.

Lipids and lipid derivatives such as fatty aldehydes, fatty alcohols, and hydrocarbons such as alkanes produced by cells as described herein can be modified by the use of one or more enzymes, including a lipase, as described above. When the hydrocarbons are in the extracellular environment of the cells, the one or more enzymes can be added to that environment under conditions in which the enzyme modifies the hydrocarbon or completes its synthesis from a hydrocarbon precursor. Alternatively, the hydrocarbons can be partially, or completely, isolated from the cellular material before addition of one or more catalysts such as enzymes. Such catalysts are exogenously added, and their activity occurs outside the cell or in vitro.

Thus, lipids and hydrocarbons produced by cells in vivo, or enzymatically modified in vitro, as described herein can be optionally further processed by conventional means. The processing can include "cracking" to reduce the size, and thus increase the hydrogen:carbon ratio, of hydrocarbon molecules. Catalytic and thermal cracking methods are routinely used in hydrocarbon and triglyceride oil processing. Catalytic methods involve the use of a catalyst, such as a solid acid catalyst. The catalyst can be silica-alumina or a zeolite, which result in the heterolytic, or asymmetric, breakage of a carbon-

carbon bond to result in a carbocation and a hydride anion. These reactive intermediates then undergo either rearrangement or hydride transfer with another hydrocarbon. The reactions can thus regenerate the intermediates to result in a self-propagating chain mechanism. Hydrocarbons can also be processed to reduce, optionally to zero, the number of carbon-carbon double, or triple, bonds therein. Hydrocarbons can also be processed to remove or eliminate a ring or cyclic structure therein. Hydrocarbons can also be processed to increase the hydrogen:carbon ratio. This can include the addition of hydrogen (“hydrogenation”) and/or the “cracking” of hydrocarbons into smaller hydrocarbons.

Thermal methods involve the use of elevated temperature and pressure to reduce hydrocarbon size. An elevated temperature of about 800° C. and pressure of about 700 kPa can be used. These conditions generate “light,” a term that is sometimes used to refer to hydrogen-rich hydrocarbon molecules (as distinguished from photon flux), while also generating, by condensation, heavier hydrocarbon molecules which are relatively depleted of hydrogen. The methodology provides homolytic, or symmetrical, breakage and produces alkenes, which may be optionally enzymatically saturated as described above.

Catalytic and thermal methods are standard in plants for hydrocarbon processing and oil refining. Thus hydrocarbons produced by cells as described herein can be collected and processed or refined via conventional means. See Hillen et al. (Biotechnology and Bioengineering, Vol. XXIV:193-205 (1982)) for a report on hydrocracking of microalgae-produced hydrocarbons. In alternative embodiments, the fraction is treated with another catalyst, such as an organic compound, heat, and/or an inorganic compound. For processing of lipids into biodiesel, a transesterification process is used as described in Section IV herein.

Hydrocarbons produced via methods of the present invention are useful in a variety of industrial applications. For example, the production of linear alkylbenzene sulfonate (LAS), an anionic surfactant used in nearly all types of detergents and cleaning preparations, utilizes hydrocarbons generally comprising a chain of 10-14 carbon atoms. See, for example, U.S. Pat. Nos. 6,946,430; 5,506,201; 6,692,730; 6,268,517; 6,020,509; 6,140,302; 5,080,848; and 5,567,359. Surfactants, such as LAS, can be used in the manufacture of personal care compositions and detergents, such as those described in U.S. Pat. Nos. 5,942,479; 6,086,903; 5,833,999; 6,468,955; and 6,407,044.

Increasing interest is directed to the use of hydrocarbon components of biological origin in fuels, such as biodiesel, renewable diesel, and jet fuel, since renewable biological starting materials that may replace starting materials derived from fossil fuels are available, and the use thereof is desirable. There is an urgent need for methods for producing hydrocarbon components from biological materials. The present invention fulfills this need by providing methods for production of biodiesel, renewable diesel, and jet fuel using the lipids generated by the methods described herein as a biological material to produce biodiesel, renewable diesel, and jet fuel.

Traditional diesel fuels are petroleum distillates rich in paraffinic hydrocarbons. They have boiling ranges as broad as 370° to 780° F., which are suitable for combustion in a compression ignition engine, such as a diesel engine vehicle. The American Society of Testing and Materials (ASTM) establishes the grade of diesel according to the boiling range, along with allowable ranges of other fuel properties, such as cetane number, cloud point, flash point, viscosity, aniline point, sulfur content, water content, ash content, copper strip corro-

sion, and carbon residue. Technically, any hydrocarbon distillate material derived from biomass or otherwise that meets the appropriate ASTM specification can be defined as diesel fuel (ASTM D975), jet fuel (ASTM D1655), or as biodiesel if it is a fatty acid methyl ester (ASTM D6751).

After extraction, lipid and/or hydrocarbon components recovered from the microbial biomass described herein can be subjected to chemical treatment to manufacture a fuel for use in diesel vehicles and jet engines.

Biodiesel is a liquid which varies in color—between golden and dark brown—depending on the production feedstock. It is practically immiscible with water, has a high boiling point and low vapor pressure. Biodiesel refers to a diesel-equivalent processed fuel for use in diesel-engine vehicles. Biodiesel is biodegradable and non-toxic. An additional benefit of biodiesel over conventional diesel fuel is lower engine wear. Typically, biodiesel comprises C14-C18 alkyl esters. Various processes convert biomass or a lipid produced and isolated as described herein to diesel fuels. A preferred method to produce biodiesel is by transesterification of a lipid as described herein. A preferred alkyl ester for use as biodiesel is a methyl ester or ethyl ester.

Biodiesel produced by a method described herein can be used alone or blended with conventional diesel fuel at any concentration in most modern diesel-engine vehicles. When blended with conventional diesel fuel (petroleum diesel), biodiesel may be present from about 0.1% to about 99.9%. Much of the world uses a system known as the “B” factor to state the amount of biodiesel in any fuel mix. For example, fuel containing 20% biodiesel is labeled B20. Pure biodiesel is referred to as B100.


Biodiesel can also be used as a heating fuel in domestic and commercial boilers. Existing oil boilers may contain rubber parts and may require conversion to run on biodiesel. The conversion process is usually relatively simple, involving the exchange of rubber parts for synthetic parts due to biodiesel being a strong solvent. Due to its strong solvent power, burning biodiesel will increase the efficiency of boilers. Biodiesel can be used as an additive in formulations of diesel to increase the lubricity of pure Ultra-Low Sulfur Diesel (ULSD) fuel, which is advantageous because it has virtually no sulfur content. Biodiesel is a better solvent than petrodiesel and can be used to break down deposits of residues in the fuel lines of vehicles that have previously been run on petrodiesel.

Biodiesel can be produced by transesterification of triglycerides contained in oil-rich biomass. Thus, in another aspect of the present invention a method for producing biodiesel is provided. In a preferred embodiment, the method for producing biodiesel comprises the steps of (a) cultivating a lipid-containing microorganism using methods disclosed herein (b) lysing a lipid-containing microorganism to produce a lysate, (c) isolating lipid from the lysed microorganism, and (d) transesterifying the lipid composition, whereby biodiesel is produced. Methods for growth of a microorganism, lysing a microorganism to produce a lysate, treating the lysate in a medium comprising an organic solvent to form a heterogeneous mixture and separating the treated lysate into a lipid composition have been described above and can also be used in the method of producing biodiesel.

The lipid profile of the biodiesel is usually highly similar to the lipid profile of the feedstock oil. Other oils provided by the methods and compositions of the invention can be subjected to transesterification to yield biodiesel with lipid profiles including (a) at least 4% C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

Lipid compositions can be subjected to transesterification to yield long-chain fatty acid esters useful as biodiesel. Preferred transesterification reactions are outlined below and include base catalyzed transesterification and transesterification using recombinant lipases. In a base-catalyzed transesterification process, the triacylglycerides are reacted with an alcohol, such as methanol or ethanol, in the presence of an alkaline catalyst, typically potassium hydroxide. This reaction forms methyl or ethyl esters and glycerol (glycerol) as a byproduct.

Animal and plant oils are typically made of triglycerides which are esters of free fatty acids with the trihydric alcohol, glycerol. In transesterification, the glycerol in a triacylglyceride (TAG) is replaced with a short-chain alcohol such as methanol or ethanol. A typical reaction scheme is as follows:



In this reaction, the alcohol is deprotonated with a base to make it a stronger nucleophile. Commonly, ethanol or methanol is used in vast excess (up to 50-fold). Normally, this reaction will proceed either exceedingly slowly or not at all. Heat, as well as an acid or base can be used to help the reaction proceed more quickly. The acid or base are not consumed by the transesterification reaction, thus they are not reactants but catalysts. Almost all biodiesel has been produced using the base-catalyzed technique as it requires only low temperatures and pressures and produces over 98% conversion yield (provided the starting oil is low in moisture and free fatty acids).

Transesterification has also been carried out, as discussed above, using an enzyme, such as a lipase instead of a base. Lipase-catalyzed transesterification can be carried out, for example, at a temperature between the room temperature and 80° C., and a mole ratio of the TAG to the lower alcohol of greater than 1:1, preferably about 3:1. Lipases suitable for use in transesterification include, but are not limited to, those listed in Table 7. Other examples of lipases useful for transesterification are found in, e.g. U.S. Pat. Nos. 4,798,793; 4,940,845 5,156,963; 5,342,768; 5,776,741 and WO89/01032. Such lipases include, but are not limited to, lipases produced by microorganisms of *Rhizopus*, *Aspergillus*, *Candida*, *Mucor*, *Pseudomonas*, *Rhizomucor*, *Candida*, and *Humicola* and pancreas lipase.

TABLE 7

Lipases suitable for use in transesterification.

*Aspergillus niger* lipase ABG73614, *Candida antarctica* lipase B (novozym-435) CAA83122, *Candida cylindracea* lipase AAR24090, *Candida lipolytica* lipase (Lipase L; Amano Pharmaceutical Co., Ltd.), *Candida rugosa* lipase (e.g., Lipase-OF; Meito Sangyo Co., Ltd.), *Mucor miehei* lipase (Lipozyme IM 20), *Pseudomonas fluorescens* lipase AAA25882, *Rhizopus japonicus* lipase (Lilipase A-10FG) Q7M4U7\_1, *Rhizomucor miehei* lipase B34959, *Rhizopus oryzae* lipase (Lipase F) AAF32408, *Serratia marcescens* lipase (SM Enzyme) ABI13521, *Thermomyces lanuginosa* lipase CAB58509, Lipase P (Nagase ChemteX Corporation), and Lipase QLM (Meito Sangyo Co., Ltd., Nagoya, Japan)

One challenge to using a lipase for the production of fatty acid esters suitable for biodiesel is that the price of lipase is much higher than the price of sodium hydroxide (NaOH) used by the strong base process. This challenge has been

addressed by using an immobilized lipase, which can be recycled. However, the activity of the immobilized lipase must be maintained after being recycled for a minimum number of cycles to allow a lipase-based process to compete with the strong base process in terms of the production cost. Immobilized lipases are subject to poisoning by the lower alcohols typically used in transesterification. U.S. Pat. No. 6,398,707 (issued Jun. 4, 2002 to Wu et al.) describes methods for enhancing the activity of immobilized lipases and regenerating immobilized lipases having reduced activity. Some suitable methods include immersing an immobilized lipase in an alcohol having a carbon atom number not less than 3 for a period of time, preferably from 0.5-48 hours, and more preferably from 0.5-1.5 hours. Some suitable methods also include washing a deactivated immobilized lipase with an alcohol having a carbon atom number not less than 3 and then immersing the deactivated immobilized lipase in a vegetable oil for 0.5-48 hours.

In particular embodiments, a recombinant lipase is expressed in the same microorganisms that produce the lipid on which the lipase acts. Suitable recombinant lipases include those listed above in Table 7 and/or having GenBank Accession numbers listed above in Table 7, or a polypeptide that has at least 70% amino acid identity with one of the lipases listed above in Table 7 and that exhibits lipase activity. In additional embodiments, the enzymatic activity is present in a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% identity with one of the above described sequences, all of which are hereby incorporated by reference as if fully set forth. DNA encoding the lipase and selectable marker is preferably codon-optimized cDNA. Methods of recoding genes for expression in microalgae are described in U.S. Pat. No. 7,135,290.

The common international standard for biodiesel is EN 14214. ASTM D6751 is the most common biodiesel standard referenced in the United States and Canada. Germany uses DIN EN 14214 and the UK requires compliance with BS EN 14214. Basic industrial tests to determine whether the products conform to these standards typically include gas chromatography, HPLC, and others. Biodiesel meeting the quality standards is very non-toxic, with a toxicity rating (LD<sub>50</sub>) of greater than 50 mL/kg.

Although biodiesel that meets the ASTM standards has to be non-toxic, there can be contaminants which tend to crystallize and/or precipitate and fall out of solution as sediment. Sediment formation is particularly a problem when biodiesel is used at lower temperatures. The sediment or precipitates may cause problems such as decreasing fuel flow, clogging fuel lines, clogging filters, etc. Processes are well-known in the art that specifically deal with the removal of these con-

taminants and sediments in biodiesel in order to produce a higher quality product. Examples for such processes include, but are not limited to, pretreatment of the oil to remove contaminants such as phospholipids and free fatty acids (e.g.,

degumming, caustic refining and silica adsorbent filtration) and cold filtration. Cold filtration is a process that was developed specifically to remove any particulates and sediments that are present in the biodiesel after production. This process cools the biodiesel and filters out any sediments or precipitates that might form when the fuel is used at a lower temperature. Such a process is well known in the art and is described in US Patent Application Publication No. 2007-0175091. Suitable methods may include cooling the biodiesel to a temperature of less than about 38° C. so that the impurities and contaminants precipitate out as particulates in the biodiesel liquid. Diatomaceous earth or other filtering material may then added to the cooled biodiesel to form a slurry, which may then filtered through a pressure leaf or other type of filter to remove the particulates. The filtered biodiesel may then be run through a polish filter to remove any remaining sediments and diatomaceous earth, so as to produce the final biodiesel product.

Example 14 described the production of biodiesel using triglyceride oil from *Prototheca moriformis*. The Cold Soak Filterability by the ASTM D6751 A1 method of the biodiesel produced in Example 14 was 120 seconds for a volume of 300 ml. This test involves filtration of 300 ml of B100, chilled to 40° F. for 16 hours, allowed to warm to room temp, and filtered under vacuum using 0.7 micron glass fiber filter with stainless steel support. Oils of the invention can be transesterified to generate biodiesel with a cold soak time of less than 120 seconds, less than 100 seconds, and less than 90 seconds.

Subsequent processes may also be used if the biodiesel will be used in particularly cold temperatures. Such processes include winterization and fractionation. Both processes are designed to improve the cold flow and winter performance of the fuel by lowering the cloud point (the temperature at which the biodiesel starts to crystallize). There are several approaches to winterizing biodiesel. One approach is to blend the biodiesel with petroleum diesel. Another approach is to use additives that can lower the cloud point of biodiesel. Another approach is to remove saturated methyl esters indiscriminately by mixing in additives and allowing for the crystallization of saturates and then filtering out the crystals. Fractionation selectively separates methyl esters into individual components or fractions, allowing for the removal or inclusion of specific methyl esters. Fractionation methods include urea fractionation, solvent fractionation and thermal distillation.

Another valuable fuel provided by the methods of the present invention is renewable diesel, which comprises alkanes, such as C10:0, C12:0, C14:0, C16:0 and C18:0 and thus, are distinguishable from biodiesel. High quality renewable diesel conforms to the ASTM D975 standard. The lipids produced by the methods of the present invention can serve as feedstock to produce renewable diesel. Thus, in another aspect of the present invention, a method for producing renewable diesel is provided. Renewable diesel can be produced by at least three processes: hydrothermal processing (hydrotreating); hydroprocessing; and indirect liquefaction. These processes yield non-ester distillates. During these processes, triacylglycerides produced and isolated as described herein, are converted to alkanes.

In one embodiment, the method for producing renewable diesel comprises (a) cultivating a lipid-containing microorganism using methods disclosed herein (b) lysing the microorganism to produce a lysate, (c) isolating lipid from the lysed microorganism, and (d) deoxygenating and hydrotreating the lipid to produce an alkane, whereby renewable diesel is produced. Lipids suitable for manufacturing renewable diesel can be obtained via extraction from microbial biomass using

an organic solvent such as hexane, or via other methods, such as those described in U.S. Pat. No. 5,928,696. Some suitable methods may include mechanical pressing and centrifuging.

In some methods, the microbial lipid is first cracked in conjunction with hydrotreating to reduce carbon chain length and saturate double bonds, respectively. The material is then isomerized, also in conjunction with hydrotreating. The naphtha fraction can then be removed through distillation, followed by additional distillation to vaporize and distill components desired in the diesel fuel to meet an ASTM D975 standard while leaving components that are heavier than desired for meeting the D975 standard. Hydrotreating, hydrocracking, deoxygenation and isomerization methods of chemically modifying oils, including triglyceride oils, are well known in the art. See for example European patent applications EP1741768 (A1); EP1741767 (A1); EP1682466 (A1); EP1640437 (A1); EP1681337 (A1); EP1795576 (A1); and U.S. Pat. Nos. 7,238,277; 6,630,066; 6,596,155; 6,977,322; 7,041,866; 6,217,746; 5,885,440; 6,881,873.

In one embodiment of the method for producing renewable diesel, treating the lipid to produce an alkane is performed by hydrotreating of the lipid composition. In hydrothermal processing, typically, biomass is reacted in water at an elevated temperature and pressure to form oils and residual solids. Conversion temperatures are typically 300° to 660° F., with pressure sufficient to keep the water primarily as a liquid, 100 to 170 standard atmosphere (atm). Reaction times are on the order of 15 to 30 minutes. After the reaction is completed, the organics are separated from the water. Thereby a distillate suitable for diesel is produced.

In some methods of making renewable diesel, the first step of treating a triglyceride is hydroprocessing to saturate double bonds, followed by deoxygenation at elevated temperature in the presence of hydrogen and a catalyst. In some methods, hydrogenation and deoxygenation occur in the same reaction. In other methods deoxygenation occurs before hydrogenation. Isomerization is then optionally performed, also in the presence of hydrogen and a catalyst. Naphtha components are preferably removed through distillation. For examples, see U.S. Pat. Nos. 5,475,160 (hydrogenation of triglycerides); 5,091,116 (deoxygenation, hydrogenation and gas removal); 6,391,815 (hydrogenation); and 5,888,947 (isomerization).

One suitable method for the hydrogenation of triglycerides includes preparing an aqueous solution of copper, zinc, magnesium and lanthanum salts and another solution of alkali metal or preferably, ammonium carbonate. The two solutions may be heated to a temperature of about 20° C. to about 85° C. and metered together into a precipitation container at rates such that the pH in the precipitation container is maintained between 5.5 and 7.5 in order to form a catalyst. Additional water may be used either initially in the precipitation container or added concurrently with the salt solution and precipitation solution. The resulting precipitate may then be thoroughly washed, dried, calcined at about 300° C. and activated in hydrogen at temperatures ranging from about 100° C. to about 400° C. One or more triglycerides may then be contacted and reacted with hydrogen in the presence of the above-described catalyst in a reactor. The reactor may be a trickle bed reactor, fixed bed gas-solid reactor, packed bubble column reactor, continuously stirred tank reactor, a slurry phase reactor, or any other suitable reactor type known in the art. The process may be carried out either batchwise or in continuous fashion. Reaction temperatures are typically in the range of from about 170° C. to about 250° C. while reaction pressures are typically in the range of from about 300 psig to about 2000 psig. Moreover, the molar ratio of hydro-

gen to triglyceride in the process of the present invention is typically in the range of from about 20:1 to about 700:1. The process is typically carried out at a weight hourly space velocity (WHSV) in the range of from about 0.1 hr<sup>-1</sup> to about 5 hr<sup>-1</sup>. One skilled in the art will recognize that the time period required for reaction will vary according to the temperature used, the molar ratio of hydrogen to triglyceride, and the partial pressure of hydrogen. The products produced by the such hydrogenation processes include fatty alcohols, glycerol, traces of paraffins and unreacted triglycerides. These products are typically separated by conventional means such as, for example, distillation, extraction, filtration, crystallization, and the like.

Petroleum refiners use hydroprocessing to remove impurities by treating feeds with hydrogen. Hydroprocessing conversion temperatures are typically 300° to 700° F. Pressures are typically 40 to 100 atm. The reaction times are typically on the order of 10 to 60 minutes. Solid catalysts are employed to increase certain reaction rates, improve selectivity for certain products, and optimize hydrogen consumption.

Suitable methods for the deoxygenation of an oil includes heating an oil to a temperature in the range of from about 350° F. to about 550° F. and continuously contacting the heated oil with nitrogen under at least pressure ranging from about atmospheric to above for at least about 5 minutes.

Suitable methods for isomerization includes using alkali isomerization and other oil isomerization known in the art.

Hydrotreating and hydroprocessing ultimately lead to a reduction in the molecular weight of the triglyceride feed. The triglyceride molecule is reduced to four hydrocarbon molecules under hydroprocessing conditions: a propane molecule and three heavier hydrocarbon molecules, typically in the C8 to C18 range.

Thus, in one embodiment, the product of one or more chemical reaction(s) performed on lipid compositions of the invention is an alkane mixture that comprises ASTM D975 renewable diesel. Production of hydrocarbons by microorganisms is reviewed by Metzger et al. *Appl Microbiol Biotechnol* (2005) 66: 486-496 and A Look Back at the U.S. Department of Energy's Aquatic Species Program: Biodiesel from Algae, NREL/TP-580-24190, John Sheehan, Terri Dunahay, John Benemann and Paul Roessler (1998).

The distillation properties of a diesel fuel is described in terms of T10-T90 (temperature at 10% and 90%, respectively, volume distilled). Renewable diesel was produced from *Prototheca moriformis* triglyceride oil and is described in Example 14. The T10-T90 of the material produced in Example 14 was 57.9° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other T10-T90 ranges, such as 20, 25, 30, 35, 40, 45, 50, 60 and 65° C. using triglyceride oils produced according to the methods disclosed herein.

The T10 of the material produced in Example 14 was 242.1° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other T10 values, such as T10 between 180 and 295, between 190 and 270, between 210 and 250, between 225 and 245, and at least 290.

The T90 of the material produced in Example 14 was 300° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) dis-

closed herein can be employed to generate renewable diesel compositions with other T90 values, such as T90 between 280 and 380, between 290 and 360, between 300 and 350, between 310 and 340, and at least 290.

The FBP of the material produced in Example 14 was 300° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other FBP values, such as FBP between 290 and 400, between 300 and 385, between 310 and 370, between 315 and 360, and at least 300.

Other oils provided by the methods and compositions of the invention can be subjected to combinations of hydrotreating, isomerization, and other covalent modification including oils with lipid profiles including (a) at least 4% C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

A traditional ultra-low sulfur diesel can be produced from any form of biomass by a two-step process. First, the biomass is converted to a syngas, a gaseous mixture rich in hydrogen and carbon monoxide. Then, the syngas is catalytically converted to liquids. Typically, the production of liquids is accomplished using Fischer-Tropsch (FT) synthesis. This technology applies to coal, natural gas, and heavy oils. Thus, in yet another preferred embodiment of the method for producing renewable diesel, treating the lipid composition to produce an alkane is performed by indirect liquefaction of the lipid composition.

The present invention also provides methods to produce jet fuel. Jet fuel is clear to straw colored. The most common fuel is an unleaded/paraffin oil-based fuel classified as Aeroplane A-1, which is produced to an internationally standardized set of specifications. Jet fuel is a mixture of a large number of different hydrocarbons, possibly as many as a thousand or more. The range of their sizes (molecular weights or carbon numbers) is restricted by the requirements for the product, for example, freezing point or smoke point. Kerosene-type Aeroplane fuel (including Jet A and Jet A-1) has a carbon number distribution between about 8 and 16 carbon numbers. Wide-cut or naphta-type Aeroplane fuel (including Jet B) typically has a carbon number distribution between about 5 and 15 carbons.

Both Aeroplanes (Jet A and Jet B) may contain a number of additives. Useful additives include, but are not limited to, antioxidants, antistatic agents, corrosion inhibitors, and fuel system icing inhibitor (FSII) agents. Antioxidants prevent gumming and usually, are based on alkylated phenols, for example, AO-30, AO-31, or AO-37. Antistatic agents dissipate static electricity and prevent sparking. Stadis 450 with dinonylnaphthylsulfonic acid (DINNSA) as the active ingredient, is an example. Corrosion inhibitors, e.g., DCI-4A is used for civilian and military fuels and DCI-6A is used for military fuels. FSII agents, include, e.g., Di-EGME.

In one embodiment of the invention, a jet fuel is produced by blending algal fuels with existing jet fuel. The lipids produced by the methods of the present invention can serve as feedstock to produce jet fuel. Thus, in another aspect of the present invention, a method for producing jet fuel is provided. Herewith two methods for producing jet fuel from the lipids produced by the methods of the present invention are provided: fluid catalytic cracking (FCC); and hydrodeoxygenation (HDO).

Fluid Catalytic Cracking (FCC) is one method which is used to produce olefins, especially propylene from heavy crude fractions. The lipids produced by the method of the present invention can be converted to olefins. The process

involves flowing the lipids produced through an FCC zone and collecting a product stream comprised of olefins, which is useful as a jet fuel. The lipids produced are contacted with a cracking catalyst at cracking conditions to provide a product stream comprising olefins and hydrocarbons useful as jet fuel.

In one embodiment, the method for producing jet fuel comprises (a) cultivating a lipid-containing microorganism using methods disclosed herein, (b) lysing the lipid-containing microorganism to produce a lysate, (c) isolating lipid from the lysate, and (d) treating the lipid composition, whereby jet fuel is produced. In one embodiment of the method for producing a jet fuel, the lipid composition can be flowed through a fluid catalytic cracking zone, which, in one embodiment, may comprise contacting the lipid composition with a cracking catalyst at cracking conditions to provide a product stream comprising C<sub>2</sub>-C<sub>5</sub> olefins.

In certain embodiments of this method, it may be desirable to remove any contaminants that may be present in the lipid composition. Thus, prior to flowing the lipid composition through a fluid catalytic cracking zone, the lipid composition is pretreated. Pretreatment may involve contacting the lipid composition with an ion-exchange resin. The ion exchange resin is an acidic ion exchange resin, such as Amberlyst™-15 and can be used as a bed in a reactor through which the lipid composition is flowed, either upflow or downflow. Other pretreatments may include mild acid washes by contacting the lipid composition with an acid, such as sulfuric, acetic, nitric, or hydrochloric acid. Contacting is done with a dilute acid solution usually at ambient temperature and atmospheric pressure.

The lipid composition, optionally pretreated, is flowed to an FCC zone where the hydrocarbonaceous components are cracked to olefins. Catalytic cracking is accomplished by contacting the lipid composition in a reaction zone with a catalyst composed of finely divided particulate material. The reaction is catalytic cracking, as opposed to hydrocracking, and is carried out in the absence of added hydrogen or the consumption of hydrogen. As the cracking reaction proceeds, substantial amounts of coke are deposited on the catalyst. The catalyst is regenerated at high temperatures by burning coke from the catalyst in a regeneration zone. Coke-containing catalyst, referred to herein as "coked catalyst", is continually transported from the reaction zone to the regeneration zone to be regenerated and replaced by essentially coke-free regenerated catalyst from the regeneration zone. Fluidization of the catalyst particles by various gaseous streams allows the transport of catalyst between the reaction zone and regeneration zone. Methods for cracking hydrocarbons, such as those of the lipid composition described herein, in a fluidized stream of catalyst, transporting catalyst between reaction and regeneration zones, and combusting coke in the regenerator are well known by those skilled in the art of FCC processes. Exemplary FCC applications and catalysts useful for cracking the lipid composition to produce C<sub>2</sub>-C<sub>5</sub> olefins are described in U.S. Pat. Nos. 6,538,169, 7,288,685, which are incorporated in their entirety by reference.

Suitable FCC catalysts generally comprise at least two components that may or may not be on the same matrix. In some embodiments, both two components may be circulated throughout the entire reaction vessel. The first component generally includes any of the well-known catalysts that are used in the art of fluidized catalytic cracking, such as an active amorphous clay-type catalyst and/or a high activity, crystalline molecular sieve. Molecular sieve catalysts may be preferred over amorphous catalysts because of their much-improved selectivity to desired products. IN some preferred embodiments, zeolites may be used as the molecular sieve in

the FCC processes. Preferably, the first catalyst component comprises a large pore zeolite, such as an Y-type zeolite, an active alumina material, a binder material, comprising either silica or alumina and an inert filler such as kaolin.

In one embodiment, cracking the lipid composition of the present invention, takes place in the riser section or, alternatively, the lift section, of the FCC zone. The lipid composition is introduced into the riser by a nozzle resulting in the rapid vaporization of the lipid composition. Before contacting the catalyst, the lipid composition will ordinarily have a temperature of about 149° C. to about 316° C. (300° F. to 600° F.). The catalyst is flowed from a blending vessel to the riser where it contacts the lipid composition for a time of about 2 seconds or less.

The blended catalyst and reacted lipid composition vapors are then discharged from the top of the riser through an outlet and separated into a cracked product vapor stream including olefins and a collection of catalyst particles covered with substantial quantities of coke and generally referred to as "coked catalyst." In an effort to minimize the contact time of the lipid composition and the catalyst which may promote further conversion of desired products to undesirable other products, any arrangement of separators such as a swirl arm arrangement can be used to remove coked catalyst from the product stream quickly. The separator, e.g. swirl arm separator, is located in an upper portion of a chamber with a stripping zone situated in the lower portion of the chamber. Catalyst separated by the swirl arm arrangement drops down into the stripping zone. The cracked product vapor stream comprising cracked hydrocarbons including light olefins and some catalyst exit the chamber via a conduit which is in communication with cyclones. The cyclones remove remaining catalyst particles from the product vapor stream to reduce particle concentrations to very low levels. The product vapor stream then exits the top of the separating vessel. Catalyst separated by the cyclones is returned to the separating vessel and then to the stripping zone. The stripping zone removes adsorbed hydrocarbons from the surface of the catalyst by counter-current contact with steam.

Low hydrocarbon partial pressure operates to favor the production of light olefins. Accordingly, the riser pressure is set at about 172 to 241 kPa (25 to 35 psia) with a hydrocarbon partial pressure of about 35 to 172 kPa (5 to 25 psia), with a preferred hydrocarbon partial pressure of about 69 to 138 kPa (10 to 20 psia). This relatively low partial pressure for hydrocarbon is achieved by using steam as a diluent to the extent that the diluent is 10 to 55 wt-% of lipid composition and preferably about 15 wt-% of lipid composition. Other diluents such as dry gas can be used to reach equivalent hydrocarbon partial pressures.

The temperature of the cracked stream at the riser outlet will be about 510° C. to 621° C. (950° F. to 1150° F.). However, riser outlet temperatures above 566° C. (1050° F.) make more dry gas and more olefins. Whereas, riser outlet temperatures below 566° C. (1050° F.) make less ethylene and propylene. Accordingly, it is preferred to run the FCC process at a preferred temperature of about 566° C. to about 630° C., preferred pressure of about 138 kPa to about 240 kPa (20 to 35 psia). Another condition for the process is the catalyst to lipid composition ratio which can vary from about 5 to about 20 and preferably from about 10 to about 15.

In one embodiment of the method for producing a jet fuel, the lipid composition is introduced into the lift section of an FCC reactor. The temperature in the lift section will be very hot and range from about 700° C. (1292° F.) to about 760° C. (1400° F.) with a catalyst to lipid composition ratio of about 100 to about 150. It is anticipated that introducing the lipid

composition into the lift section will produce considerable amounts of propylene and ethylene.

In another embodiment of the method for producing a jet fuel using the lipid composition or the lipids produced as described herein, the structure of the lipid composition or the lipids is broken by a process referred to as hydrodeoxygenation (HDO). HDO means removal of oxygen by means of hydrogen, that is, oxygen is removed while breaking the structure of the material. Olefinic double bonds are hydrogenated and any sulphur and nitrogen compounds are removed. Sulphur removal is called hydrodesulphurization (HDS). Pretreatment and purity of the raw materials (lipid composition or the lipids) contribute to the service life of the catalyst.

Generally in the HDO/HDS step, hydrogen is mixed with the feed stock (lipid composition or the lipids) and then the mixture is passed through a catalyst bed as a co-current flow, either as a single phase or a two phase feed stock. After the HDO/MDS step, the product fraction is separated and passed to a separate isomerization reactor. An isomerization reactor for biological starting material is described in the literature (FI 100 248) as a co-current reactor.

The process for producing a fuel by hydrogenating a hydrocarbon feed, e.g., the lipid composition or the lipids herein, can also be performed by passing the lipid composition or the lipids as a co-current flow with hydrogen gas through a first hydrogenation zone, and thereafter the hydrocarbon effluent is further hydrogenated in a second hydrogenation zone by passing hydrogen gas to the second hydrogenation zone as a counter-current flow relative to the hydrocarbon effluent. Exemplary HDO applications and catalysts useful for cracking the lipid composition to produce C<sub>2</sub>-C<sub>5</sub> olefins are described in U.S. Pat. No. 7,232,935, which is incorporated in its entirety by reference.

Typically, in the hydrodeoxygenation step, the structure of the biological component, such as the lipid composition or lipids herein, is decomposed, oxygen, nitrogen, phosphorus and sulphur compounds, and light hydrocarbons as gas are removed, and the olefinic bonds are hydrogenated. In the second step of the process, i.e. in the so-called isomerization step, isomerization is carried out for branching the hydrocarbon chain and improving the performance of the paraffin at low temperatures.

In the first step, i.e. HDO step, of the cracking process, hydrogen gas and the lipid composition or lipids herein which are to be hydrogenated are passed to a HDO catalyst bed system either as co-current or counter-current flows, said catalyst bed system comprising one or more catalyst bed(s), preferably 1-3 catalyst beds. The HDO step is typically operated in a co-current manner. In case of a HDO catalyst bed system comprising two or more catalyst beds, one or more of the beds may be operated using the counter-current flow principle. In the HDO step, the pressure varies between 20 and 150 bar, preferably between 50 and 100 bar, and the temperature varies between 200 and 500° C., preferably in the range of 300-400° C. In the HDO step, known hydrogenation catalysts containing metals from Group VII and/or VIB of the Periodic System may be used. Preferably, the hydrogenation catalysts are supported Pd, Pt, Ni, NiMo or a CoMo catalysts, the support being alumina and/or silica. Typically, NiMo/Al<sub>2</sub>O<sub>3</sub> and CoMo/Al<sub>2</sub>O<sub>3</sub> catalysts are used.

Prior to the HDO step, the lipid composition or lipids herein may optionally be treated by prehydrogenation under milder conditions thus avoiding side reactions of the double bonds. Such prehydrogenation is carried out in the presence of a prehydrogenation catalyst at temperatures of 50-400° C. and at hydrogen pressures of 1-200 bar, preferably at a temperature between 150 and 250° C. and at a hydrogen pressure

between 10 and 100 bar. The catalyst may contain metals from Group VIII and/or VIE of the Periodic System. Preferably, the prehydrogenation catalyst is a supported Pd, Pt, Ni, NiMo or a CoMo catalyst, the support being alumina and/or silica.

A gaseous stream from the HDO step containing hydrogen is cooled and then carbon monoxide, carbon dioxide, nitrogen, phosphorus and sulphur compounds, gaseous light hydrocarbons and other impurities are removed therefrom. After compressing, the purified hydrogen or recycled hydrogen is returned back to the first catalyst bed and/or between the catalyst beds to make up for the withdrawn gas stream. Water is removed from the condensed liquid. The liquid is passed to the first catalyst bed or between the catalyst beds.

After the HDO step, the product is subjected to an isomerization step. It is substantial for the process that the impurities are removed as completely as possible before the hydrocarbons are contacted with the isomerization catalyst. The isomerization step comprises an optional stripping step, wherein the reaction product from the HDO step may be purified by stripping with water vapour or a suitable gas such as light hydrocarbon, nitrogen or hydrogen. The optional stripping step is carried out in counter-current manner in a unit upstream of the isomerization catalyst, wherein the gas and liquid are contacted with each other, or before the actual isomerization reactor in a separate stripping unit utilizing counter-current principle.

After the stripping step the hydrogen gas and the hydrogenated lipid composition or lipids herein, and optionally an n-paraffin mixture, are passed to a reactive isomerization unit comprising one or several catalyst bed(s). The catalyst beds of the isomerization step may operate either in co-current or counter-current manner.

It is important for the process that the counter-current flow principle is applied in the isomerization step. In the isomerization step this is done by carrying out either the optional stripping step or the isomerization reaction step or both in counter-current manner. In the isomerization step, the pressure varies in the range of 20-150 bar, preferably in the range of 20-100 bar, the temperature being between 200 and 500° C., preferably between 300 and 400° C. In the isomerization step, isomerization catalysts known in the art may be used. Suitable isomerization catalysts contain molecular sieve and/or a metal from Group VII and/or a carrier. Preferably, the isomerization catalyst contains SAPO-11 or SAPO-41 or ZSM-22 or ZSM-23 or ferrierite and Pt, Pd or Ni and Al<sub>2</sub>O<sub>3</sub> or SiO<sub>2</sub>. Typical isomerization catalysts are, for example, Pt/SAPO-11/Al<sub>2</sub>O<sub>3</sub>, Pt/ZSM-22/Al<sub>2</sub>O<sub>3</sub>, Pt/ZSM-23/Al<sub>2</sub>O<sub>3</sub> and Pt/SAPO-11/SiO<sub>2</sub>. The isomerization step and the HDO step may be carried out in the same pressure vessel or in separate pressure vessels. Optional prehydrogenation may be carried out in a separate pressure vessel or in the same pressure vessel as the HDO and isomerization steps.

Thus, in one embodiment, the product of the one or more chemical reactions is an alkane mixture that comprises ASTM D1655 jet fuel. In some embodiments, the composition conforming to the specification of ASTM 1655 jet fuel has a sulfur content that is less than 10 ppm. In other embodiments, the composition conforming to the specification of ASTM 1655 jet fuel has a T10 value of the distillation curve of less than 205° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a final boiling point (FBP) of less than 300° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a flash point of at least 38° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a density between

775K/M<sup>3</sup> and 840K/M<sup>3</sup>. In yet another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a freezing point that is below -47° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a net Heat of Combustion that is at least 42.8 MJ/K. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a hydrogen content that is at least 13.4 mass %. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a thermal stability, as tested by quantitative gravimetric JFTOT at 260° C., that is below 3 mm of Hg. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has an existent gum that is below 7 mg/dl.

Thus, the present invention discloses a variety of methods in which chemical modification of microalgal lipid is undertaken to yield products useful in a variety of industrial and other applications. Examples of processes for modifying oil produced by the methods disclosed herein include, but are not limited to, hydrolysis of the oil, hydroprocessing of the oil, and esterification of the oil. The modification of the microalgal oil produces basic oleochemicals that can be further modified into selected derivative oleochemicals for a desired function. In a manner similar to that described above with reference to fuel producing processes, these chemical modifications can also be performed on oils generated from the microbial cultures described herein. Examples of basic oleochemicals include, but are not limited to, soaps, fatty acids, fatty acid methyl esters, and glycerol. Examples of derivative oleochemicals include, but are not limited to, fatty nitriles, esters, dimer acids, quats, surfactants, fatty alkanolamides, fatty alcohol sulfates, resins, emulsifiers, fatty alcohols, olefins, and higher alkanes.

Hydrolysis of the fatty acid constituents from the glycerolipids produced by the methods of the invention yields free fatty acids that can be derivatized to produce other useful chemicals. Hydrolysis occurs in the presence of water and a catalyst which may be either an acid or a base. The liberated free fatty acids can be derivatized to yield a variety of products, as reported in the following: U.S. Pat. Nos. 5,304,664 (Highly sulfated fatty acids); 7,262,158 (Cleansing compositions); 7,115,173 (Fabric softener compositions); 6,342,208 (Emulsions for treating skin); 7,264,886 (Water repellent compositions); 6,924,333 (Paint additives); 6,596,768 (Lipid-enriched ruminant feedstock); and 6,380,410 (Surfactants for detergents and cleaners).

With regard to hydrolysis, in one embodiment of the invention, a triglyceride oil is optionally first hydrolyzed in a liquid medium such as water or sodium hydroxide so as to obtain glycerol and soaps. There are various suitable triglyceride hydrolysis methods, including, but not limited to, saponification, acid hydrolysis, alkaline hydrolysis, enzymatic hydrolysis (referred herein as splitting), and hydrolysis using hot-compressed water. One skilled in the art will recognize that a triglyceride oil need not be hydrolyzed in order to produce an oleochemical; rather, the oil may be converted directly to the desired oleochemical by other known process. For example, the triglyceride oil may be directly converted to a methyl ester fatty acid through esterification.

In some embodiments, catalytic hydrolysis of the oil produced by methods disclosed herein occurs by splitting the oil into glycerol and fatty acids. As discussed above, the fatty acids may then be further processed through several other modifications to obtain derivative oleochemicals. For example, in one embodiment the fatty acids may undergo an amination reaction to produce fatty nitrogen compounds. In

another embodiment, the fatty acids may undergo ozonolysis to produce mono- and dibasic-acids.

In other embodiments hydrolysis may occur via the, splitting of oils produced herein to create oleochemicals. In some preferred embodiments of the invention, a triglyceride oil may be split before other processes is performed. One skilled in the art will recognize that there are many suitable triglyceride splitting methods, including, but not limited to, enzymatic splitting and pressure splitting.

Generally, enzymatic oil splitting methods use enzymes, lipases, as biocatalysts acting on a water/oil mixture. Enzymatic splitting then splits the oil or fat, respectively, is into glycerol and free fatty acids. The glycerol may then migrates into the water phase whereas the organic phase enriches with free fatty acids.

The enzymatic splitting reactions generally take place at the phase boundary between organic and aqueous phase, where the enzyme is present only at the phase boundary. Triglycerides that meet the phase boundary then contribute to or participate in the splitting reaction. As the reaction proceeds, the occupation density or concentration of fatty acids still chemically bonded as glycerides; in comparison to free fatty acids, decreases at the phase boundary so that the reaction is slowed down. In certain embodiments, enzymatic splitting may occur at room temperature. One of ordinary skill in the art would know the suitable conditions for splitting oil into the desired fatty acids.

By way of example, the reaction speed can be accelerated by increasing the interface boundary surface. Once the reaction is complete, free fatty acids are then separated from the organic phase freed from enzyme, and the residue which still contains fatty acids chemically bonded as glycerides is fed back or recycled and mixed with fresh oil or fat to be subjected to splitting. In this manner, recycled glycerides are then subjected to a further enzymatic splitting process. In some embodiments, the free fatty acids are extracted from an oil or fat partially split in such a manner. In that way, if the chemically bound fatty acids (triglycerides) are returned or fed back into the splitting process, the enzyme consumption can be drastically reduced.

The splitting degree is determined as the ratio of the measured acid value divided by the theoretically possible acid value which can be computed for a given oil or fat. Preferably, the acid value is measured by means of titration according to standard common methods. Alternatively, the density of the aqueous glycerol phase can be taken as a measure for the splitting degree.

In one embodiment, the slitting process as described herein is also suitable for splitting the mono-, di- and triglyceride that are contained in the so-called soap-stock from the alkali refining processes of the produced oils. In this manner, the soap-stock can be quantitatively converted without prior saponification of the neutral oils into the fatty acids. For this purpose, the fatty acids being chemically bonded in the soaps are released, preferably before splitting, through an addition of acid. In certain embodiments, a buffer solution is used in addition to water and enzyme for the splitting process.

In one embodiment, oils produced in accordance with the methods of the invention can also be subjected to saponification as a method of hydrolysis. Animal and plant oils are typically made of triacylglycerols (TAGs), which are esters of fatty acids with the trihydric alcohol, glycerol. In an alkaline hydrolysis reaction, the glycerol in a TAG is removed, leaving three carboxylic acid anions that can associate with alkali metal cations such as sodium or potassium to produce fatty acid salts. In this scheme, the carboxylic acid constituents are cleaved from the glycerol moiety and replaced with hydroxyl



groups. The quantity of base (e.g., KOH) that is used in the reaction is determined by the desired degree of saponification. If the objective is, for example, to produce a soap product that comprises some of the oils originally present in the TAG composition, an amount of base insufficient to convert all of the TAGs to fatty acid salts is introduced into the reaction mixture. Normally, this reaction is performed in an aqueous solution and proceeds slowly, but may be expedited by the addition of heat. Precipitation of the fatty acid salts can be facilitated by addition of salts, such as water-soluble alkali metal halides (e.g., NaCl or KCl), to the reaction mixture. Preferably, the base is an alkali metal hydroxide, such as NaOH or KOH. Alternatively, other bases, such as alkanolamines, including for example triethanolamine and aminomethylpropanol, can be used in the reaction scheme. In some cases, these alternatives may be preferred to produce a clear soap product.

In some methods, the first step of chemical modification may be hydroprocessing to saturate double bonds, followed by deoxygenation at elevated temperature in the presence of hydrogen and a catalyst. In other methods, hydrogenation and deoxygenation may occur in the same reaction. In still other methods deoxygenation occurs before hydrogenation. Isomerization may then be optionally performed, also in the presence of hydrogen and a catalyst. Finally, gases and naphtha components can be removed if desired. For example, see U.S. Pat. Nos. 5,475,160 (hydrogenation of triglycerides); 5,091,116 (deoxygenation, hydrogenation and gas removal); 6,391,815 (hydrogenation); and 5,888,947 (isomerization).

In some embodiments of the invention, the triglyceride oils are partially or completely deoxygenated. The deoxygenation reactions form desired products, including, but not limited to, fatty acids, fatty alcohols, polyols, ketones, and aldehydes. In general, without being limited by any particular theory, the deoxygenation reactions involve a combination of various different reaction pathways, including without limitation: hydrogenolysis, hydrogenation, consecutive hydrogenation-hydrogenolysis, consecutive hydrogenolysis-hydrogenation, and combined hydrogenation-hydrogenolysis reactions, resulting in at least the partial removal of oxygen from the fatty acid or fatty acid ester to produce reaction products, such as fatty alcohols, that can be easily converted to the desired chemicals by further processing. For example, in one embodiment, a fatty alcohol may be converted to olefins through FCC reaction or to higher alkanes through a condensation reaction.

One such chemical modification is hydrogenation, which is the addition of hydrogen to double bonds in the fatty acid constituents of glycerolipids or of free fatty acids. The hydrogenation process permits the transformation of liquid oils into semi-solid or solid fats, which may be more suitable for specific applications.

Hydrogenation of oil produced by the methods described herein can be performed in conjunction with one or more of the methods and/or materials provided herein, as reported in the following: U.S. Pat. Nos. 7,288,278 (Food additives or medicaments); 5,346,724 (Lubrication products); 5,475,160 (Fatty alcohols); 5,091,116 (Edible oils); 6,808,737 (Structural fats for margarine and spreads); 5,298,637 (Reduced-calorie fat substitutes); 6,391,815 (Hydrogenation catalyst and sulfur adsorbent); 5,233,099 and 5,233,100 (Fatty alcohols); 4,584,139 (Hydrogenation catalysts); 6,057,375 (Foam suppressing agents); and 7,118,773 (Edible emulsion spreads).

One skilled in the art will recognize that various processes may be used to hydrogenate carbohydrates. One suitable method includes contacting the carbohydrate with hydrogen or hydrogen mixed with a suitable gas and a catalyst under

conditions sufficient in a hydrogenation reactor to form a hydrogenated product. The hydrogenation catalyst generally can include Cu, Re, Ni, Fe, Co, Ru, Pd, Rh, Pt, Os, Ir, and alloys or any combination thereof, either alone or with promoters such as W, Mo, Au, Ag, Cr, Zn, Mn, Sn, B, P, Bi, and alloys or any combination thereof. Other effective hydrogenation catalyst materials include either supported nickel or ruthenium modified with rhenium. In an embodiment, the hydrogenation catalyst also includes any one of the supports, depending on the desired functionality of the catalyst. The hydrogenation catalysts may be prepared by methods known to those of ordinary skill in the art.

In some embodiments the hydrogenation catalyst includes a supported Group VIII metal catalyst and a metal sponge material (e.g., a sponge nickel catalyst). Raney nickel provides an example of an activated sponge nickel catalyst suitable for use in this invention. In other embodiment, the hydrogenation reaction in the invention is performed using a catalyst comprising a nickel-rhenium catalyst or a tungsten-modified nickel catalyst. One example of a suitable catalyst for the hydrogenation reaction of the invention is a carbon-supported nickel-rhenium catalyst.

In an embodiment, a suitable Raney nickel catalyst may be prepared by treating an alloy of approximately equal amounts by weight of nickel and aluminum with an aqueous alkali solution, e.g., containing about 25 weight % of sodium hydroxide. The aluminum is selectively dissolved by the aqueous alkali solution resulting in a sponge shaped material comprising mostly nickel with minor amounts of aluminum. The initial alloy includes promoter metals (i.e., molybdenum or chromium) in the amount such that about 1 to 2 weight % remains in the formed sponge nickel catalyst. In another embodiment, the hydrogenation catalyst is prepared using a solution of ruthenium (III) nitrosylnitrate, ruthenium (III) chloride in water to impregnate a suitable support material. The solution is then dried to form a solid having a water content of less than about 1% by weight. The solid may then be reduced at atmospheric pressure in a hydrogen stream at 300° C. (uncalcined) or 400° C. (calcined) in a rotary ball furnace for 4 hours. After cooling and rendering the catalyst inert with nitrogen, 5% by volume of oxygen in nitrogen is passed over the catalyst for 2 hours.

In certain embodiments, the catalyst described includes a catalyst support. The catalyst support stabilizes and supports the catalyst. The type of catalyst support used depends on the chosen catalyst and the reaction conditions. Suitable supports for the invention include, but are not limited to, carbon, silica, silica-alumina, zirconia, titania, ceria, vanadia, nitride, boron nitride, heteropolyacids, hydroxyapatite, zinc oxide, chromia, zeolites, carbon nanotubes, carbon fullerene and any combination thereof.

The catalysts used in this invention can be prepared using conventional methods known to those in the art. Suitable methods may include, but are not limited to, incipient wetting, evaporative impregnation, chemical vapor deposition, wash-coating, magnetron sputtering techniques, and the like.

The conditions for which to carry out the hydrogenation reaction will vary based on the type of starting material and the desired products. One of ordinary skill in the art, with the benefit of this disclosure, will recognize the appropriate reaction conditions. In general, the hydrogenation reaction is conducted at temperatures of 80° C. to 250° C., and preferably at 90° C. to 200° C., and most preferably at 100° C. to 150° C. In some embodiments, the hydrogenation reaction is conducted at pressures from 500 KPa to 14000 KPa.

The hydrogen used in the hydrogenolysis reaction of the current invention may include external hydrogen, recycled

hydrogen, in situ generated hydrogen, and any combination thereof. As used herein, the term “external hydrogen” refers to hydrogen that does not originate from the biomass reaction itself, but rather is added to the system from another source.

In some embodiments of the invention, it is desirable to convert the starting carbohydrate to a smaller molecule that will be more readily converted to desired higher hydrocarbons. One suitable method for this conversion is through a hydrogenolysis reaction. Various processes are known for performing hydrogenolysis of carbohydrates. One suitable method includes contacting a carbohydrate with hydrogen or hydrogen mixed with a suitable gas and a hydrogenolysis catalyst in a hydrogenolysis reactor under conditions sufficient to form a reaction product comprising smaller molecules or polyols. As used herein, the term “smaller molecules or polyols” includes any molecule that has a smaller molecular weight, which can include a smaller number of carbon atoms or oxygen atoms than the starting carbohydrate. In an embodiment, the reaction products include smaller molecules that include polyols and alcohols. Someone of ordinary skill in the art would be able to choose the appropriate method by which to carry out the hydrogenolysis reaction.

In some embodiments, a 5 and/or 6 carbon sugar or sugar alcohol may be converted to propylene glycol, ethylene glycol, and glycerol using a hydrogenolysis catalyst. The hydrogenolysis catalyst may include Cr, Mo, W, Re, Mn, Cu, Cd, Fe, Co, Ni, Pt, Pd, Rh, Ru, Ir, Os, and alloys or any combination thereof, either alone or with promoters such as Au, Ag, Cr, Zn, Mn, Sn, Bi, B, O, and alloys or any combination thereof. The hydrogenolysis catalyst may also include a carbonaceous pyropolymer catalyst containing transition metals (e.g., chromium, molybdenum, tungsten, rhenium, manganese, copper, cadmium) or Group VIII metals (e.g., iron, cobalt, nickel, platinum, palladium, ruthenium, rhodium, iridium, and osmium). In certain embodiments, the hydrogenolysis catalyst may include any of the above metals combined with an alkaline earth metal oxide or adhered to a catalytically active support. In certain embodiments, the catalyst described in the hydrogenolysis reaction may include a catalyst support as described above for the hydrogenation reaction.

The conditions for which to carry out the hydrogenolysis reaction will vary based on the type of starting material and the desired products. One of ordinary skill in the art, with the benefit of this disclosure, will recognize the appropriate conditions to use to carry out the reaction. In general, the hydrogenolysis reaction is conducted at temperatures of 110° C. to 300° C., and preferably at 170° C. to 220° C., and most preferably at 200° C. to 225° C. In some embodiments, the hydrogenolysis reaction is conducted under basic conditions, preferably at a pH of 8 to 13, and even more preferably at a pH of 10 to 12. In some embodiments, the hydrogenolysis reaction is conducted at pressures in a range between 60 KPa and 16500 KPa, and preferably in a range between 1700 KPa and 14000 KPa, and even more preferably between 4800 KPa and 11000 KPa.

The hydrogen used in the hydrogenolysis reaction of the current invention can include external hydrogen, recycled hydrogen, in situ generated hydrogen, and any combination thereof.

In some embodiments, the reaction products discussed above may be converted into higher hydrocarbons through a condensation reaction in a condensation reactor (shown schematically as condensation reactor 110 in FIG. 1). In such embodiments, condensation of the reaction products occurs in the presence of a catalyst capable of forming higher hydrocarbons. While not intending to be limited by theory, it is

believed that the production of higher hydrocarbons proceeds through a stepwise addition reaction including the formation of carbon-carbon, or carbon-oxygen bond. The resulting reaction products include any number of compounds containing these moieties, as described in more detail below.

In certain embodiments, suitable condensation catalysts include an acid catalyst, a base catalyst, or an acid/base catalyst. As used herein, the term “acid/base catalyst” refers to a catalyst that has both an acid and a base functionality. In some embodiments the condensation catalyst can include, without limitation, zeolites, carbides, nitrides, zirconia, alumina, silica, aluminosilicates, phosphates, titanium oxides, zinc oxides, vanadium oxides, lanthanum oxides, yttrium oxides, scandium oxides, magnesium oxides, cerium oxides, barium oxides, calcium oxides, hydroxides, heteropolyacids, inorganic acids, acid modified resins, base modified resins, and any combination thereof. In some embodiments, the condensation catalyst can also include a modifier. Suitable modifiers include La, Y, Sc, P, B, Bi, Li, Na, K, Rb, Cs, Mg, Ca, Sr, Ba, and any combination thereof. In some embodiments, the condensation catalyst can also include a metal. Suitable metals include Cu, Ag, Au, Pt, Ni, Fe, Co, Ru, Zn, Cd, Ga, In, Rh, Pd, Ir, Re, Mn, Cr, Mo, W, Sn, Os, alloys, and any combination thereof.

In certain embodiments, the catalyst described in the condensation reaction may include a catalyst support as described above for the hydrogenation reaction. In certain embodiments, the condensation catalyst is self-supporting. As used herein, the term “self-supporting” means that the catalyst does not need another material to serve as support. In other embodiments, the condensation catalyst is used in conjunction with a separate support suitable for suspending the catalyst. In an embodiment, the condensation catalyst support is silica.

The conditions under which the condensation reaction occurs will vary based on the type of starting material and the desired products. One of ordinary skill in the art, with the benefit of this disclosure, will recognize the appropriate conditions to use to carry out the reaction. In some embodiments, the condensation reaction is carried out at a temperature at which the thermodynamics for the proposed reaction are favorable. The temperature for the condensation reaction will vary depending on the specific starting polyol or alcohol. In some embodiments, the temperature for the condensation reaction is in a range from 80° C. to 500° C., and preferably from 125° C. to 450° C., and most preferably from 125° C. to 250° C. In some embodiments, the condensation reaction is conducted at pressures in a range between 0 KPa to 9000 KPa, and preferably in a range between 0 KPa and 7000 KPa, and even more preferably between 0 KPa and 5000 KPa.

The higher alkanes formed by the invention include, but are not limited to, branched or straight chain alkanes that have from 4 to 30 carbon atoms, branched or straight chain alkenes that have from 4 to 30 carbon atoms, cycloalkanes that have from 5 to 30 carbon atoms, cycloalkenes that have from 5 to 30 carbon atoms, aryls, fused aryls, alcohols, and ketones. Suitable alkanes include, but are not limited to, butane, pentane, pentene, 2-methylbutane, hexane, hexene, 2-methylpentane, 3-methylpentane, 2,2-dimethylbutane, 2,3-dimethylbutane, heptane, heptene, octane, octene, 2,2,4-trimethylpentane, 2,3-dimethylhexane, 2,3,4-trimethylpentane, 2,3-dimethylpentane, nonane, nonene, decane, decene, undecane, undecene, dodecane, dodecene, tridecane, tridecene, tetradecane, tetradecene, pentadecane, pentadecene, nonyldecane, nonyldecene, eicosane, eicosene, uneicosane, uneicosene, doieicosane, doieicosene, trieicosane,

tricosene, tetraicosane, tetraicosene, and isomers thereof. Some of these products may be suitable for use as fuels.

In some embodiments, the cycloalkanes and the cycloalkenes are unsubstituted. In other embodiments, the cycloalkanes and cycloalkenes are mono-substituted. In still other 5 embodiments, the cycloalkanes and cycloalkenes are multi-substituted. In the embodiments comprising the substituted cycloalkanes and cycloalkenes, the substituted group includes, without limitation, a branched or straight chain alkyl having 1 to 12 carbon atoms, a branched or straight 10 chain alkylene having 1 to 12 carbon atoms, a phenyl, and any combination thereof. Suitable cycloalkanes and cycloalkenes include, but are not limited to, cyclopentane, cyclopentene, cyclohexane, cyclohexene, methyl-cyclopentane, methyl-cyclopentene, ethyl-cyclopentane, ethyl-cyclopentene, ethyl-cyclohexane, ethyl-cyclohexene, isomers and any combination 15 thereof.

In some embodiments, the aryls formed are unsubstituted. In another embodiment, the aryls formed are mono-substituted. In the embodiments comprising the substituted aryls, the substituted group includes, without limitation, a branched or straight chain alkyl having 1 to 12 carbon atoms, a branched or straight chain alkylene having 1 to 12 carbon 20 atoms, a phenyl, and any combination thereof. Suitable aryls for the invention include, but are not limited to, benzene, toluene, xylene, ethyl benzene, para xylene, meta xylene, and any combination thereof.

The alcohols produced in the invention have from 4 to 30 carbon atoms. In some embodiments, the alcohols are cyclic. In other embodiments, the alcohols are branched. In another 30 embodiment, the alcohols are straight chained. Suitable alcohols for the invention include, but are not limited to, butanol, pentanol, hexanol, heptanol, octanol, nonanol, decanol, undecanol, dodecanol, tridecanol, tetradecanol, pentadecanol, hexadecanol, heptyldecanol, octyldecanol, nonyldecanol, eicosanol, uneicosanol, doeicosanol, triecosanol, tetraeicosanol, and isomers thereof.

The ketones produced in the invention have from 4 to 30 carbon atoms. In an embodiment, the ketones are cyclic. In another embodiment, the ketones are branched. In another 40 embodiment, the ketones are straight chained. Suitable ketones for the invention include, but are not limited to, butanone, pentanone, hexanone, heptanone, octanone, nonanone, decanone, undecanone, dodecanone, tridecanone, tetradecanone, pentadecanone, hexadecanone, heptyldecanone, octyldecanone, nonyldecanone, eicosanone, uneicosanone, doeicosanone, triecosanone, tetraeicosanone, and isomers thereof.

Another such chemical modification is interesterification. 50 Naturally produced glycerolipids do not have a uniform distribution of fatty acid constituents. In the context of oils, interesterification refers to the exchange of acyl radicals between two esters of different glycerolipids. The interesterification process provides a mechanism by which the fatty acid constituents of a mixture of glycerolipids can be rearranged to modify the distribution pattern. Interesterification is a well-known chemical process, and generally comprises heating (to about 200° C.) a mixture of oils for a period (e.g., 30 minutes) in the presence of a catalyst, such as an alkali metal or alkali metal alkylate (e.g., sodium methoxide). This process can be used to randomize the distribution pattern of the fatty acid constituents of an oil mixture, or can be directed to produce a desired distribution pattern. This method of chemical modification of lipids can be performed on materials provided 65 herein, such as microbial biomass with a percentage of dry cell weight as lipid at least 20%.

Directed interesterification, in which a specific distribution pattern of fatty acids is sought, can be performed by maintaining the oil mixture at a temperature below the melting point of some TAGs which might occur. This results in selective crystallization of these TAGs, which effectively removes them from the reaction mixture as they crystallize. The process can be continued until most of the fatty acids in the oil have precipitated, for example. A directed interesterification process can be used, for example, to produce a product with a lower calorie content via the substitution of longer-chain fatty acids with shorter-chain counterparts. Directed interesterification can also be used to produce a product with a mixture of fats that can provide desired melting characteristics and structural features sought in food additives or products (e.g., margarine) without resorting to hydrogenation, which can produce unwanted trans isomers.

Interesterification of oils produced by the methods described herein can be performed in conjunction with one or more of the methods and/or materials, or to produce products, as reported in the following: U.S. Pat. Nos. 6,080,853 (Non-digestible fat substitutes); 4,288,378 (Peanut butter stabilizer); 5,391,383 (Edible spray oil); 6,022,577 (Edible fats for food products); 5,434,278 (Edible fats for food products); 5,268,192 (Low calorie nut products); 5,258,197 (Reduce calorie edible compositions); 4,335,156 (Edible fat product); 7,288,278 (Food additives or medicaments); 7,115,760 (Fractionation process); 6,808,737 (Structural fats); 5,888,947 (Engine lubricants); 5,686,131 (Edible oil mixtures); and 4,603,188 (Curable urethane compositions).

In one embodiment in accordance with the invention, transesterification of the oil, as described above, is followed by reaction of the transesterified product with polyol, as reported in U.S. Pat. No. 6,465,642, to produce polyol fatty acid polyesters. Such an esterification and separation process may comprise the steps as follows: reacting a lower alkyl ester with polyol in the presence of soap; removing residual soap from the product mixture; water-washing and drying the product mixture to remove impurities; bleaching the product mixture for refinement; separating at least a portion of the unreacted lower alkyl ester from the polyol fatty acid polyester in the product mixture; and recycling the separated unreacted lower alkyl ester.

Transesterification can also be performed on microbial biomass with short chain fatty acid esters, as reported in U.S. Pat. No. 6,278,006. In general, transesterification may be performed by adding a short chain fatty acid ester to an oil in the presence of a suitable catalyst and heating the mixture. In some embodiments, the oil comprises about 5% to about 90% of the reaction mixture by weight. In some embodiments, the short chain fatty acid esters can be about 10% to about 50% of the reaction mixture by weight. Non-limiting examples of catalysts include base catalysts, sodium methoxide, acid catalysts including inorganic acids such as sulfuric acid and acidified clays, organic acids such as methane sulfonic acid, benzenesulfonic acid, and toluenesulfonic acid, and acidic resins such as Amberlyst 15. Metals such as sodium and magnesium, and metal hydrides also are useful catalysts.

Another such chemical modification is hydroxylation, which involves the addition of water to a double bond resulting in saturation and the incorporation of a hydroxyl moiety. The hydroxylation process provides a mechanism for converting one or more fatty acid constituents of a glycerolipid to a hydroxy fatty acid. Hydroxylation can be performed, for example, via the method reported in U.S. Pat. No. 5,576,027. Hydroxylated fatty acids, including castor oil and its derivatives, are useful as components in several industrial applications, including food additives, surfactants, pigment wetting

agents, defoaming agents, water proofing additives, plasticizing agents, cosmetic emulsifying and/or deodorant agents, as well as in electronics, pharmaceuticals, paints, inks, adhesives, and lubricants. One example of how the hydroxylation of a glyceride may be performed is as follows: fat may be heated, preferably to about 30-50° C. combined with heptane and maintained at temperature for thirty minutes or more; acetic acid may then be added to the mixture followed by an aqueous solution of sulfuric acid followed by an aqueous hydrogen peroxide solution which is added in small increments to the mixture over one hour; after the aqueous hydrogen peroxide, the temperature may then be increased to at least about 60° C. and stirred for at least six hours; after the stirring, the mixture is allowed to settle and a lower aqueous layer formed by the reaction may be removed while the upper heptane layer formed by the reaction may be washed with hot water having a temperature of about 60° C.; the washed heptane layer may then be neutralized with an aqueous potassium hydroxide solution to a pH of about 5 to 7 and then removed by distillation under vacuum; the reaction product may then be dried under vacuum at 100° C. and the dried product steam-deodorized under vacuum conditions and filtered at about 50° to 60° C. using diatomaceous earth.

Hydroxylation of microbial oils produced by the methods described herein can be performed in conjunction with one or more of the methods and/or materials, or to produce products, as reported in the following: U.S. Pat. Nos. 6,590,113 (Oil-based coatings and ink); 4,049,724 (Hydroxylation process); 6,113,971 (Olive oil butter); 4,992,189 (Lubricants and lube additives); 5,576,027 (Hydroxylated milk); and 6,869,597 (Cosmetics).

Hydroxylated glycerolipids can be converted to estolides. Estolides consist of a glycerolipid in which a hydroxylated fatty acid constituent has been esterified to another fatty acid molecule. Conversion of hydroxylated glycerolipids to estolides can be carried out by warming a mixture of glycerolipids and fatty acids and contacting the mixture with a mineral acid, as described by Isbell et al., *JAOCs* 71(2):169-174 (1994). Estolides are useful in a variety of applications, including without limitation those reported in the following: U.S. Pat. Nos. 7,196,124 (Elastomeric materials and floor coverings); 5,458,795 (Thickened oils for high-temperature applications); 5,451,332 (Fluids for industrial applications); 5,427,704 (Fuel additives); and 5,380,894 (Lubricants, greases, plasticizers, and printing inks).

Other chemical reactions that can be performed on microbial oils include reacting triacylglycerols with a cyclopropanating agent to enhance fluidity and/or oxidative stability, as reported in U.S. Pat. No. 6,051,539; manufacturing of waxes from triacylglycerols, as reported in U.S. Pat. No. 6,770,104; and epoxidation of triacylglycerols, as reported in "The effect of fatty acid composition on the acrylation kinetics of epoxidized triacylglycerols", *Journal of the American Oil Chemists' Society*, 79:1, 59-63, (2001) and *Free Radical Biology and Medicine*, 37:1, 104-114 (2004).

The generation of oil-bearing microbial biomass for fuel and chemical products as described above results in the production of delipidated biomass meal. Delipidated meal is a byproduct of preparing algal oil and is useful as animal feed for farm animals, e.g., ruminants, poultry, swine and aquaculture. The resulting meal, although of reduced oil content, still contains high quality proteins, carbohydrates, fiber, ash, residual oil and other nutrients appropriate for an animal feed. Because the cells are predominantly lysed by the oil separation process, the delipidated meal is easily digestible by such animals. Delipidated meal can optionally be combined with other ingredients, such as grain, in an animal feed. Because delipidated meal has a powdery consistency, it can be pressed

into pellets using an extruder or expander or another type of machine, which are commercially available.

The invention, having been described in detail above, is exemplified in the following examples, which are offered to illustrate, but not to limit, the claimed invention.

## VII. EXAMPLES

### Example 1

#### Methods for Culturing *Prototheca*

*Prototheca* strains were cultivated to achieve a high percentage of oil by dry cell weight. Cryopreserved cells were thawed at room temperature and 500 ul of cells were added to 4.5 ml of medium (4.2 g/L K<sub>2</sub>HPO<sub>4</sub>, 3.1 g/L NaH<sub>2</sub>PO<sub>4</sub>, 0.24 g/L MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.25 g/L Citric Acid monohydrate, 0.025 g/L CaCl<sub>2</sub>·2H<sub>2</sub>O, 2 g/L yeast extract) plus 2% glucose and grown for 7 days at 28° C. with agitation (200 rpm) in a 6-well plate. Dry cell weights were determined by centrifuging 1 ml of culture at 14,000 rpm for 5 min in a pre-weighed Eppendorf tube. The culture supernatant was discarded and the resulting cell pellet washed with 1 ml of deionized water. The culture was again centrifuged, the supernatant discarded, and the cell pellets placed at -80° C. until frozen. Samples were then lyophilized for 24 hrs and dry cell weights calculated. For determination of total lipid in cultures, 3 ml of culture was removed and subjected to analysis using an Ankom system (Ankom Inc., Macedon, N.Y.) according to the manufacturer's protocol. Samples were subjected to solvent extraction with an Amkom XT10 extractor according to the manufacturer's protocol. Total lipid was determined as the difference in mass between acid hydrolyzed dried samples and solvent extracted, dried samples. Percent oil dry cell weight measurements are shown in Table 8.

TABLE 8

Percent oil by dry cell weight		
Species	Strain	% Oil
<i>Prototheca stagnora</i>	UTEX 327	13.14
<i>Prototheca moriformis</i>	UTEX 1441	18.02
<i>Prototheca moriformis</i>	UTEX 1435	27.17

Microalgae samples from the strains listed in Table 22 above were genotyped. Genomic DNA was isolated from algal biomass as follows. Cells (approximately 200 mg) were centrifuged from liquid cultures 5 minutes at 14,000×g. Cells were then resuspended in sterile distilled water, centrifuged 5 minutes at 14,000×g and the supernatant discarded. A single glass bead ~2 mm in diameter was added to the biomass and tubes were placed at -80° C. for at least 15 minutes. Samples were removed and 150 µl of grinding buffer (1% Sarkosyl, 0.25 M Sucrose, 50 mM NaCl, 20 mM EDTA, 100 mM Tris-HCl, pH 8.0, RNase A 0.5 ug/ul) was added. Pellets were resuspended by vortexing briefly, followed by the addition of 40 ul of 5M NaCl. Samples were vortexed briefly, followed by the addition of 66 µl of 5% CTAB (Cetyl trimethylammonium bromide) and a final brief vortex. Samples were next incubated at 65° C. for 10 minutes after which they were centrifuged at 14,000×g for 10 minutes. The supernatant was transferred to a fresh tube and extracted once with 300 µl of Phenol:Chloroform:Isoamyl alcohol 12:12:1, followed by centrifugation for 5 minutes at 14,000×g. The resulting aqueous phase was transferred to a fresh tube containing 0.7 vol of isopropanol (~190 µl), mixed by inversion and incubated at room temperature for 30 minutes or overnight at 4° C. DNA was recovered via centrifugation at 14,000×g for 10 minutes. The resulting pellet was then washed twice with 70% ethanol,

followed by a final wash with 100% ethanol. Pellets were air dried for 20-30 minutes at room temperature followed by resuspension in 50  $\mu$ l of 10 mM TrisCl, 1 mM EDTA (pH 8.0).

Five  $\mu$ l of total algal DNA, prepared as described above, was diluted 1:50 in 10 mM Tris, pH 8.0. PCR reactions, final volume 20  $\mu$ l, were set up as follows. Ten  $\mu$ l of 2 $\times$  iProof HF master mix (BIO-RAD) was added to 0.4  $\mu$ l primer SZ02613 (5'-TGTTGAAGAATGAGCCGGCGAC-3' (SEQ ID NO:9) at 10 mM stock concentration). This primer sequence runs from position 567-588 in Gen Bank accession no. L43357 and is highly conserved in higher plants and algal plastid genomes. This was followed by the addition of 0.4  $\mu$ l primer SZ02615 (5'-CAGTGAGCTATTACGCACTC-3' (SEQ ID NO:10) at 10 mM stock concentration). This primer sequence is complementary to position 1112-1093 in Gen Bank accession no. L43357 and is highly conserved in higher plants and algal plastid genomes. Next, 5  $\mu$ l of diluted total DNA and 3.2  $\mu$ l dH<sub>2</sub>O were added. PCR reactions were run as follows: 98° C., 45"; 98° C., 8"; 53° C., 12"; 72° C., 20" for 35 cycles followed by 72° C. for 1 min and holding at 25° C. For purification of PCR products, 20  $\mu$ l of 10 mM Tris, pH 8.0, was added to each reaction, followed by extraction with 40  $\mu$ l of Phenol:Chloroform:isoamyl alcohol 12:12:1, vortexing and centrifuging at 14,000 $\times$ g for 5 minutes. PCR reactions were applied to S-400 columns (GE Healthcare) and centrifuged for 2 minutes at 3,000 $\times$ g. Purified PCR products were subsequently TOPO cloned into PCR8/GW/TOPO and positive clones selected for on LB/Spec plates. Purified plasmid DNA was sequenced in both directions using M13 forward and reverse primers. In total, twelve *Prototheca* strains were selected to have their 23S rRNA DNA sequenced and the sequences are listed in the Sequence Listing. A summary of the strains and Sequence Listing Numbers is included below. The sequences were analyzed for overall divergence from the UTEX 1435 (SEQ ID NO: 15) sequence. Two pairs emerged (UTEX 329/UTEX 1533 and UTEX 329/UTEX 1440) as the most divergent. In both cases, pairwise alignment resulted in 75.0% pairwise sequence identity. The percent sequence identity to UTEX 1435 is also included below.

Species	Strain	% nt identity	SEQ ID NO.
<i>Prototheca kruegani</i>	UTEX 329	75.2	SEQ ID NO: 11
<i>Prototheca wickerhamii</i>	UTEX 1440	99	SEQ ID NO: 12
<i>Prototheca stagnora</i>	UTEX 1442	75.7	SEQ ID NO: 13
<i>Prototheca moriformis</i>	UTEX 288	75.4	SEQ ID NO: 14
<i>Prototheca moriformis</i>	UTEX 1439; 1441; 1435; 1437	100	SEQ ID NO: 15
<i>Prototheca wickerhamii</i>	UTEX 1533	99.8	SEQ ID NO: 16
<i>Prototheca moriformis</i>	UTEX 1434	75.9	SEQ ID NO: 17
<i>Prototheca zopfii</i>	UTEX 1438	75.7	SEQ ID NO: 18
<i>Prototheca moriformis</i>	UTEX 1436	88.9	SEQ ID NO: 19

Lipid samples from a subset of the above-listed strains were analyzed for lipid profile using HPLC. Results are shown below in Table 9.

TABLE 9

Diversity of lipid chains in microalgal species									
Strain	C14:0	C16:0	C16:1	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1
UTEX 327	0	12.01	0	0	50.33	17.14	0	0	0
UTEX 1441	1.41	29.44	0.70	3.05	57.72	12.37	0.97	0.33	0
UTEX 1435	1.09	25.77	0	2.75	54.01	11.90	2.44	0	0

Algal plastid transit peptides were identified through the analysis of UTEX 1435 (*Prototheca moriformis*) or UTEX 250 (*Chlorella protothecoides*) cDNA libraries as described in Examples 12 and Example 11 below. cDNAs encoding potentially plastid targeted proteins based upon BLAST hit homology to other known plastid targeted proteins were subjected to further analysis by the software programs PSORT (<http://psort.ims.u-tokyo.ac.jp/form.html>), ChloroP (<http://www.cbs.dtu.dk/services/ChloroP/>) are TargetP (<http://www.cbs.dtu.dk/services/TargetP/>). Candidate plastid transit peptides identified through at least one of these three programs were then PCR amplified from the appropriate genomic DNA. Below is a summary of the amino acid sequences algal plastid targeting sequences (PTS) that were identified from this screen. Also included are the amino acid sequences of plant fatty acyl-ACP thioesterases that are used in the heterologous expression Examples below.

cDNA	SEQ ID NO.
<i>P. moriformis</i> isopentenyl diphosphate synthase PTS	SEQ ID NO: 127
<i>P. moriformis</i> delta 12 fatty acid desaturase PTS	SEQ ID NO: 128
<i>P. moriformis</i> stearyl ACP desaturase PTS	SEQ ID NO: 129
<i>C. protothecoides</i> stearyl ACP desaturase PTS	SEQ ID NO: 130
<i>Cuphea hookeriana</i> fatty acyl-ACP thioesterase (C8-10)	SEQ ID NO: 131
<i>Umbellularia californica</i> fatty acyl-ACP thioesterase (C12)	SEQ ID NO: 132
<i>Cinnamomum camphora</i> fatty acyl-ACP thioesterase (C14)	SEQ ID NO: 133

## Example 2

### Culturing *Prototheca* on Various Feedstocks

#### A. Sorghum

The following strains were shown to be capable of utilizing sorghum as a sole carbon source: *Prototheca moriformis* strains UTEX 1435, UTEX 1437, UTEX 288, UTEX 1439, UTEX 1441 and UTEX 1434, and *Prototheca stagnora* strain UTEX 1442. The "UTEX" designation indicates the strain number from the algal culture collection of the University of Texas, 1 University State A6700, Austin, Tex. 78712-0183.

Pure sorghum was purchased from Maasdam Sorghumi Mills (Lynnville, Iowa) with a sugar profile of fructose 21.0% w/w, dextrose 28.0% w/w, sucrose 16.0% w/w and maltose <0.5% w/w. The cultures were grown in liquid medium containing 2%, 5%, or 7% (v/v) pure sorghum (diluted from the pure stock) as the sole carbon source and the cultures were grown heterotrophically in the dark, agitating at ~350 rpm. Samples from the cultures were pulled at 24, 40, 48, 67 and 89 hours and growth was measured using A750 readings on a spectrophotometer. Growth was observed for each of the strains tested as shown in FIGS. 1-2.

## B. Cellulose

Wet, exploded corn stover, *Miscanthus*, forage sorghum, beet pulp and sugar cane bagasse were prepared by The National Renewable Energy Laboratory (Golden, Colo.) by cooking in a 1.4% sulfuric acid solution and dewatering the resultant slurry. Percent solids were determined gravimetrically by drying and were as follows: corn stover, 25% solids; *Miscanthus*, 28.7% solids; forage sorghum, 26.7% solids; and sugar cane bagasse, 26% solids.

100 gram wet samples of exploded cellulosic materials (corn stover or switch grass) were resuspended in deionized water to a final volume of 420 mL and the pH was adjusted to 4.8 using 10N NaOH. For beet pulp, 9.8 grams dry solids were brought to 350 mL with deionized water and pH was adjusted to 4.8 with 10 N NaOH. For all of the above feedstocks, Accellerase 1000 (Genencor, N.Y.) was used at a ratio of 0.25 ml enzyme per gram of dry biomass for saccharification of the cellulosic materials. Samples were incubated with agitation (110 rpm) at 50° C. for 72 hours. The pH of each of the samples was adjusted to 7.0 with NaOH (with negligible volume change), filter sterilized through a 0.22 µm filter and used in the processes detailed below. For larger scale processes, the same procedure for saccharification was followed except an additional step of tangential flow filtration (TFF) or microfiltration step was performed to aid in filter sterilization of feedstocks. A sample from each of the feedstocks prepared was reserved for determination of glucose and xylose concentration using an HPLC/ELSD-based system or a hexokinase-based kit (Sigma). Additionally, for beet pulp, the material was initially brought to volume as with the other feedstocks, the pH was then adjusted to 4.0 and a pectinase treatment was carried out at 50° C. for 24 hours. The pH was then adjusted to 4.8 if no washing steps were conducted or 5.3 if washing steps were conducted. Enzymatic saccharification was then performed with the same procedure used for the other feedstocks as described above.

Microalgae *Prototheca moriformis* strain UTEX 1435 was assessed for its ability to grow on a series of cellulosic feedstocks prepared as described above (corn stover, beet pulp, sorghum cane, *Miscanthus* and glucose control). The microalgae culture was grown in conditions described in Example 1 above with the exception of the carbon source. The carbon source was either 4% glucose (for control conditions) or 4% glucose as measured by available glucose in the cellulosic materials. Growth was assessed by A750 readings and the culturing time was 168 hours, with A750 readings at 48, 72, 96, 120, 144 and 168 hours after initiation of the culture. As can be seen in FIG. 7a, the *Prototheca moriformis* culture grew best in corn stover. The other cellulosic feedstocks used, *Miscanthus*, sorghum cane and beet pulp, all exhibited inhibition of growth.

Based on the above results with corn stover derived cellulosic sugars, lipid accumulation was also assessed in *Prototheca moriformis* using different levels of corn stover derived cellulosic sugars and reagent glucose as a control. Cultures were grown in 18 g/L glucose that was completely from corn stover derived cellulosic sugars (100% corn stover condition in FIG. 7b), 9 g/L glucose from corn stover derived cellulosic sugars supplemented with 9 g/L reagent glucose (50% corn stover supplemented with glucose to 18 g/L condition in FIG. 7b), 9 g/L glucose from corn stover derived cellulosic sugars (50% corn stover, not supplemented; glucose at 9 g/L condition in FIG. 7b) and a control culture of 42 g/L reagent glucose and 13 g/L reagent xylose for osmolarity control. All cultures were fed with cellulosic sugars to maintain the glucose concentration at 20 g/L, except for the control culture, which was fed with reagent glucose to maintain the glucose

concentration at 20 g/L. Growth was measured based on the dry cell weight of the culture and lipid productivity was determined as a percent dry cell weight. Total lipids were determined gravimetrically using an Ankom acid hydrolysis/solvent extraction system as described in Example 1 above.

As can be seen in FIG. 7b, based on biomass accumulation (as measured by DCW), all concentrations of the corn stover derived cellulosics out-performed (higher DCW) the control media that was fed glucose alone. Lipid production as a percentage of DCW was also calculated for all of the conditions. In addition to the higher biomass accumulation seen for growth on corn stover, lipid accumulation was also higher in the corn stover derived cellulosics conditions as compared to the glucose control condition. These data demonstrate that, in addition to providing cellulosic derived sugars, corn stover provides additional nutrients/components that contribute to an increased biomass accumulation (growth) and increased product yield.

Because the cellulosic feedstocks contain components in addition to glucose, some of these additional components can accumulate to undesirable levels during culture as more cellulosic derived sugars are fed into the culture as the main carbon source (usually, but not limited to, glucose) is consumed. For example, the xylose present in the cellulosic derived sugar feedstock may build up during the high density cultivation of microalgae to levels inhibitory to growth and end product production. To test the effects of xylose build up during *Prototheca moriformis* cultivation, cultures were grown with 4% glucose in the media and supplemented with 0, 10 g/L, 25 g/L, 50 g/L and 100 g/L xylose. After 6 days of culture, growth and lipid accumulation were assessed using the methods described above. As seen in FIG. 7c, surprisingly, the highest concentrations of xylose tested were not inhibitory to *Prototheca moriformis*' ability to grow and accumulate lipid, and the culture actually grew better and accumulated more lipids at the highest xylose concentrations. To explore this phenomenon, a similar experiment was carried out with sucrose, a carbon source which wild type *Prototheca moriformis* is unable to metabolize. No positive impact was observed with sucrose, suggesting that the increased growth and lipid accumulation seen with xylose is attributable to a mechanism other than the osmotic stress from high concentrations of unmetabolized components in the media and is xylose-specific.

In addition to non-metabolized sugars, salts may accumulate to inhibitory levels as a result of concentrating lignocellulosic derived sugars. Due to the acid hydrolysis step with H<sub>2</sub>SO<sub>4</sub> during the typical preparation of cellulosic materials followed by neutralization of the acid with NaOH, Na<sub>2</sub>SO<sub>4</sub> is formed during the generation of lignocellulosic sugars. To assess the impact of salt concentration on growth and lipid production, *Prototheca moriformis* cultures were grown at Na<sub>2</sub>SO<sub>4</sub> concentrations ranging from 0-700 mM in media supplemented with 4% glucose. As shown in FIG. 7d, a significant inhibition of growth was observed, as measured by DCW accumulation, where Na<sub>2</sub>SO<sub>4</sub> concentrations exceeded 25 mM, specifically at the 80 mM, 240 mM and 700 mM concentrations. In addition, the impact of antifoam P2000 was assessed in the same test. The antifoam compound had a significant, positive impact on biomass productivity. Lipid productivity was also assessed for each condition, and Na<sub>2</sub>SO<sub>4</sub> concentrations above 80 mM, specifically 240 mM and 700 mM, were inhibitory while the addition of antifoam P2000 significantly increased lipid productivity. Thus, in one embodiment, the culturing steps of the methods of the present invention include culturing in media containing an antifoaming agent.

Based on the results discussed above and summarized in FIG. 7a, inhibitors were likely present in the cellulosic feedstocks exhibiting poor growth. The present invention provides means of removing such compounds by washing the materials with hot water (hydrothermal treatment). FIG. 8 summarizes the growth results, as measured by A750, using sugar derived from cellulosic feedstock with a single hot water wash. The culture conditions were identical to those used in the processes summarized in FIG. 7a. Compared to the results shown in FIG. 7a, after just one hot water wash, *Prototheca moriformis* cultures grew better in all cellulosic feedstocks tested, specifically sugar cane bagasse, sorghum

through feeding of the cellulosic sugars. At the conclusion of the culturing, microalgae biomass from each condition was analyzed for lipid profile using the methods described in Example 1. The results of the lipid profile analysis (expressed in Area %) are summarized in Table 10 below. Each condition was tested in duplicates, and the results from each of the duplicate test conditions are included. Growth on cellulosic feedstocks resulted in a significant re-distribution in the lipid profile as compared to the glucose control. For example, there was a significant increase in C18:0 Area % in all of the cellulosic feedstock conditions as compared to the glucose control condition.

TABLE 10

	glucose 1 (ctrl)	glucose 2 (ctrl)	bagasse 1	bagasse 2	sorgh 1	sorgh 2	Miscan 1	Miscan 2
C10:0	n.d.	n.d.	0.03	0.02	n.d.	n.d.	n.d.	n.d.
C12:0	0.04	0.05	0.04	0.04	0.05	0.04	0.04	0.04
C14:0	1.64	1.64	1.07	1.10	1.17	1.14	1.08	1.12
C14:1	0.03	0.04	0.04	0.04	0.06	0.06	0.03	0.03
C15:0	0.04	0.05	0.07	0.05	0.08	0.08	0.06	0.06
C16:0	26.80	26.81	22.32	22.81	22.09	22.19	23.45	23.62
C16:1	0.75	0.82	1.68	1.70	1.92	2.12	1.38	1.23
C17:0	0.14	0.16	0.28	0.17	0.29	0.27	0.21	0.19
C17:1	0.07	0.06	0.10	0.10	0.13	0.12	0.10	0.09
C18:0	3.56	3.64	15.88	10.40	15.30	12.37	10.15	8.69
C18:1	54.22	54.01	49.87	53.87	49.35	50.80	54.05	55.26
C18:2	11.23	11.11	6.54	7.91	7.47	8.80	7.71	7.88
C18:3	0.84	0.85	0.39	0.56	0.47	0.53	0.56	0.60
alpha								
C20:0	0.31	0.30	0.85	0.63	0.76	0.69	0.63	0.56
C20:1	0.15	0.15	0.33	0.28	0.32	0.32	0.27	0.25
C20:3	0.06	0.06	0.13	0.12	0.14	0.12	0.11	0.11
C24:0	0.12	0.12	0.22	0.19	0.22	0.20	0.18	0.15

35

cane, *Miscanthus* and beet pulp, as compared to glucose control. Lipid productivity was also assessed in each of the conditions. Except for the beet pulp condition, which was comparable to the glucose control, cultures grown in sugars derived from cellulosic materials subjected to one hot water wash exhibited better lipid productivity than the glucose control.

One potential impact of hydrothermal treatment (hot water washing) of cellulosic biomass is the removal of furfurals and hydroxymethyl furfurals released by acid explosion of the material. The presence of furfurals and hydroxymethyl furfurals may have contributed to limited growth observed in some of the processes summarized in FIG. 7a. To assess how hydrothermal treatment affected the levels of furfurals (FA) and hydroxymethyl furfurals (HMF), supernatants resulting from one to three washes of cellulosic biomass derived from sugarcane bagasse (B), sorghum cane (S), *Miscanthus* (M) or beet pulp (BP) were assayed for FA and HMF by HPLC. As shown in FIG. 8, FA and RMF levels decrease significantly with each washing step. This result is consistent with the observation that FA and HMF can be inhibitory to microalgal growth (as seen in FIG. 7a) and that hydrothermal treatment removes these compounds and results in improved microalgal growth, even better than the growth in the control glucose conditions (as seen in FIG. 8).

The impact on the lipid profile of *Prototheca moriformis* cultures grown on the various hydrothermally treated lignocellulosic derived sugars was assessed. *Prototheca moriformis* cultures were grown on the following 4x-washed cellulosic feedstocks: *Miscanthus*, sugar cane bagasse and sorghum cane, with glucose levels maintained at 20 g/L

n.d. denotes none detected

Cellulosic sugar stream was generated from exploded corn stover, saccharified using Accellerase enzyme and concentrated using vacuum evaporation. This sugar stream was tested in *Prototheca moriformis* growth assays at a 4% glucose concentration. The results of the growth assays showed very poor growth and the cellulosic sugar stream was tested for conductivity (salt content). The conductivity was very high, far greater than 700 mM sodium equivalents, a level that was shown to be inhibitory to growth as described above and shown in FIG. 7d. Methods of the invention include methods in which salt is reduced or removed from lignocellulosic derived sugars prior to utilizing these feedstocks in the production of lignocellulosic derived microalgal oil. Surprisingly, however, one cannot use resins to desalt concentrated sugar streams, one must first dilute the concentrated sugar stream. To demonstrate this embodiment of the invention, cellulosic sugars derived from corn stover material were diluted eight-fold prior to removing contaminating salts with the resin. The initial conductivity of the concentrated starting material was 87 mS/cm while that of the eight-fold diluted stream was 10990  $\mu$ S/cm at a pH of 5.61. Previous studies had indicated that failure to dilute the concentrated sugar stream prior to de-ionization resulted in an inability to remove salts quantitatively as well as a significant loss of glucose from the sugar stream. Three different bed volumes of IEX resin (DOWEX Marathon MR3) were used (1:2, 1:4 and 1:10). Table 11 summarize results demonstrating the ability of a mixed bed ion exchange (IEX) resin to reduce salts (as measured by conductivity) significantly in a previously concentrated corn stover derived cellulosic sugar stream in diluted feedstocks.

40

50

60

65

TABLE 11

Ability of IEX resin to reduce salts.				
Bed volume resin: cellulosics	pH post- deionization	Conductivity post- deionization ( $\mu\text{S}/\text{cm}$ )	Calculated conductivity post- deionization and 8x re-concentration ( $\mu\text{S}/\text{cm}$ )	Na <sup>+</sup> equivalents (based on std curve) in mM
1:2	3.1	74	592	7.42
1:4	3.1	97	776	9.7
1:10	5.25	6320	50560	634

A process employing a 1:4 bed volume:cellulosic feedstock and re-concentration of the material eight-fold would result in a sodium concentration is well within the range for normal biomass and lipid accumulation. Alternatively, deionization or salt removal can be performed prior to saccharification or after saccharification, but before concentration of the sugar stream. If salt removal is performed before the concentration of the sugar stream, a dilution step of the sugar stream before salt removal would likely not be necessary.

This example demonstrates the efficacy of washing of exploded cellulosic material for the use in cellulosic oil production. As described above, concentration of cellulosically derived sugars without the removal of salts (inherent to the production of exploded cellulosic material and subsequent treatment) results in less than optimal fermentations. The materials treated in the process described below were of the appropriate pH for subsequent saccharification. In addition, the conductivity of this material was significantly reduced (over 100 fold) from the starting feedstock. Therefore, the subsequent concentrated sugars to be used in fermentations were not inhibitory due to the presence of excessive salts. An additional advantage is seen by the removal of furfurals from the cellulosic material. Any xylose or glucose removed in the hemicellulosic fraction can either be discarded or preferably re-concentrated to be used in fermentations.

Wet, exploded sugar cane bagasse (NREL, Colorado) with an initial starting mass of 65 kg wet weight and conductivity of 15,000  $\mu\text{S}/\text{cm}$ , pH 2.4 was brought to 128 kg with deionized water and the pH adjusted to 4.6 with 10 N NaOH, making the resulting conductivity 6,800  $\mu\text{S}/\text{cm}$ . The percent solids were assessed by removal of an aliquot of the suspended materials to a tared (weight=t) aluminum pan, record-

ing the wet weight (weight=w) followed by drying for three hours at 110° C. After drying samples were removed to a desiccator and allowed to come to room temperature (25° C.) at which point, they were weighed again (weight=d). Percent solids were calculated as: % solids=[(d-t/w-t)] $\times$ 100. Conductivities were measured on a Thermo Electron Orion 3 Star Conductivity meter.

The sugar cane bagasse was washed in a semi-continuous fashion by continuously mixing the cellulosic slurry (initial percent solids of 8.2%) at a temperature of 50° C. in a stainless steel reactor (150 L capacity). Cellulosics were discharged from the reactor vessel via a rotary load pump at a flow rate of 1.9-3.8 kg/min to a Sharples Model 660 decanter centrifuge. Liquid permeate was retained batch wise (ca. 35-175 kg aliquots, see Table 12 below) and homogenous aliquots removed for assessment of total sugars (glucose and xylose) and percent solids as described in Table 12. Conductivity and pH of the cellulosic material were controlled via the addition of de-ionized water and 10 N NaOH, respectively. Samples 1-10 in Table 12 represent decanted centrifuge permeate, and as such, solids and sugars present in these fractions are removed from the final, washed cellulosic materials. A mass balance calculation of total solids compared to solids removed minus solids lost plus final solids for saccharification, resulted in a 99% recovery in the above process. FIG. 8 summarizes the furfural and hydroxymethyl furfurals concentration (mg/L) in each of the 11 centrifuge permeates collected and described in Table 12. These data demonstrate a clear removal of furfurals and hydroxymethyl furfurals from the sugar cane bagasse.

TABLE 12

Mass balance for semi-continuous hydrothermal treatment of sugar cane bagasse.						
Sample	kg (wet)	kg (dry)	pH	Conductivity $\mu\text{S}/\text{cm}$	total xylose removed (g)	total glucose removed (g)
1 (initial material)	128	10.50	4.60	6,880	0	0
2	81.8	2.03		3,280	1030.68	286.3
3	76.5	0.49		2,500	298.35	76.50
4	106	0.41			254.40	63.60
5	173.9	0.30	3.74	1,260	226.07	69.56
6	101.8	0.08	4.40	791	71.26	20.36
7	110.6	0.04	4.86	327	44.24	0
8	77.2	0			0	0
9	108.6	0.02	4.7	221	0	0
10	101.5	0			0	0
11	34.8	0	4.7	146	0	0
Solids removed (samples 1-10) lost in process		3.37				
Total xylose removed					1925.00	



TABLE 12-continued

Mass balance for semi-continuous hydrothermal treatment of sugar cane bagasse.						
Sample	kg (wet)	kg (dry)	pH	Conductivity $\mu\text{S}/\text{cm}$	total xylose removed (g)	total glucose removed (g)
Total glucose removed						516.32
Final solids for saccharification	7.03					

In another demonstration of the ability of *Prototheca* to utilize cellulosic-derived feedstock, *Prototheca moriformis* (UTEX 1435) was cultivated in three-liter bioreactors using cellulosic derived sugar as a fixed carbon feedstock. The inoculum was prepared from cryopreserved cells, which were thawed at room temperature and 1 mL of cells were added to 300 mL of inoculum medium based on the basal microalgae medium described in Example 1 with 1 g/L  $(\text{NH}_4)_2\text{SO}_4$ , 4 g/L yeast extract and a trace element solution, plus 4% glucose and grown for 1 day at 28° C. with agitation (200 rpm). This culture was used to inoculate a three-liter bioreactor containing 1 L medium plus 0.26 mL of Antifoam 204 (Sigma, USA). The fermentor was controlled at 28° C. and pH was maintained at 6.8 by addition of KOH. Dissolved oxygen was maintained at 30% saturation by cascading agitation and airflow. Cellulosic sugar feedstock from corn stover was fed to the culture to maintain 0-10 g/L glucose. Desalination of cellulosic sugar feedstocks to less than 300 mM salt was essential to assure similar dry cell weight and lipid accumulation performance as compared to purified sugar feedstock controls. Desalination of the cellulosic sugar feedstock was performed using the methods described above. Fermentor samples were removed to monitor fermentation performance. Cell mass accumulation was monitored by optical density and dry cell weight. Glucose, xylose, ammonia, potassium, sodium and furfural concentrations were also determined and monitored throughout the fermentation time course. Lipid concentration was determined by gravimetric methods discussed above.

### Example 3

#### Methods for Transforming *Prototheca*

##### A. General Method for Biolistic transformation of *Prototheca*

S550d gold carriers from Seashell Technology were prepared according to the protocol from manufacturer. Linearized plasmid (20  $\mu\text{g}$ ) was mixed with 50  $\mu\text{l}$  of binding buffer and 60  $\mu\text{l}$  (30 mg) of S550d gold carriers and incubated in ice for 1 min. Precipitation buffer (100  $\mu\text{l}$ ) was added, and the mixture was incubated in ice for another 1 min. After vortexing, DNA-coated particles were pelleted by spinning at 10,000 rpm in an Eppendorf 5415C microfuge for 10 seconds. The gold pellet was washed once with 500  $\mu\text{l}$  of cold 100% ethanol, pelleted by brief spinning in the microfuge, and resuspended with 50  $\mu\text{l}$  of ice-cold ethanol. After a brief (1-2 sec) sonication, 10  $\mu\text{l}$  of DNA-coated particles were immediately transferred to the carrier membrane.

*Prototheca* strains were grown in proteose medium (2 g/L yeast extract, 2.94 mM  $\text{NaNO}_3$ , 0.17 mM  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.3 mM  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.4 mM  $\text{K}_2\text{HPO}_4$ , 1.28 mM  $\text{KH}_2\text{PO}_4$ , 0.43 mM  $\text{NaCl}$ ) on a gyratory shaker until it reaches a cell density of  $2 \times 10^6$  cells/ml. The cells were harvested, washed once with sterile distilled water, and resuspended in 500 of

medium.  $1 \times 10^7$  cells were spread in the center third of a non-selective proteose media plate. The cells were bombarded with the PDS-1000/He Biolistic Particle Delivery system (Bio-Rad). Rupture disks (1100 and 1350 psi) were used, and the plates are placed 9 and 12 cm below the screen/macrocarrier assembly. The cells were allowed to recover at 25° C. for 12-24 h. Upon recovery, the cells were scraped from the plates with a rubber spatula, mixed with 100  $\mu\text{l}$  of medium and spread on plates containing the appropriate antibiotic selection. After 7-10 days of incubation at 25° C., colonies representing transformed cells were visible on the plates from 1100 and 1350 psi rupture discs and from 9 and 12 cm distances. Colonies were picked and spotted on selective agar plates for a second round of selection.

##### B. Transformation of *Prototheca* with G418 Resistance Gene

*Prototheca moriformis* and other *Prototheca* strains sensitive to G418 can be transformed using the methods described below. G418 is an aminoglycoside antibiotic that inhibits the function of 80S ribosomes and thereby inhibits protein synthesis. The corresponding resistance gene functions through phosphorylation, resulting in inactivation of G418. *Prototheca* strains UTEX 1435, UTEX 1439 and UTEX 1437 were selected for transformation. All three *Prototheca* strains were genotyped using the methods described above. All three *Prototheca* strains had identical 23s rRNA genomic sequences (SEQ ID NO:15).

All transformation cassettes were cloned as EcoRI-SacI fragments into pUC19. Standard molecular biology techniques were used in the construction of all vectors according to Sambrook and Russell, 2001. The *C. reinhardtii* beta-tubulin promoter/5'UTR was obtained from plasmid pHyg3 (Berthold et al., (2002) Protist: 153(4), pp 401-412) by PCR as an EcoRI-AscI fragment. The *Chlorella vulgaris* nitrate reductase 3'UTR was obtained from genomic DNA isolated from UTEX strain 1803 via PCR using the following primer pairs:

Forward: (SEQ ID NO: 35)  
5' TGACCTAGGTGATTAATTAACCTCGAGGCAGCAGCTCGGATAGT

ATCG 3'

Reverse: (SEQ ID NO: 36)  
5' CTACGAGCTCAAGCTTTCATTGTGTTCCATCCACTACTT

CC 3'

The *Chlorella sorokiniana* glutamate dehydrogenase promoter/UTR was obtained via PCR of genomic DNA isolated

from UTEX strain 1230 via PCR using the following primer pairs:

(SEQ ID NO: 37) 5  
Forward: 5' GATCAGAATTCCGCCTGCAACGCAAGG GCAGC 3'

(SEQ ID NO: 38)  
Reverse: 5' GCATACTAGTGGCGGGACGGAGAGA GGGCG 3'

Codon optimization was based on the codons in Table 1 for *Prototheca moriformis*. The sequence of the non-codon optimized neomycin phosphotransferase (nptII) cassette was synthesized as an AscI-XhoI fragment and was based on upon the sequence of Genbank Accession No. YP\_788126. The codon optimized nptII cassette was also based on this Genbank Accession number.

The three *Prototheca* strains were transformed using biolistic methods described above. Briefly, the *Prototheca* strains were grown heterophically in liquid medium containing 2% glucose until they reached the desired cell density ( $1 \times 10^7$  cells/mL to  $5 \times 10^7$  cells/mL). The cells were harvested, washed once with sterile distilled water and resuspended at  $1 \times 10^8$  cells/mL. 0.5 mL of cells were then spread out on a non-selective solid media plate and allowed to dry in a sterile hood. The cells were bombarded with the PDS-1000/He Biolistic Particle Delivery System (BioRad). The cells were allowed to recover at 25° C. for 24 hours. Upon recovery, the cells were removed by washing plates with 1 mL of sterile media and transferring to fresh plates containing 100 µg/mL G418. Cells were allowed to dry in a sterile hood and colonies were allowed to form on the plate at room temperature for up to three weeks. Colonies of UTEX 1435, UTEX 1439 and UTEX 1437 were picked and spotted on selective agar plates for a second round of selection.

A subset of colonies that survived a second round of selection described above, were cultured in small volume and genomic DNA and RNA were extracted using standard molecular biology methods. Southern blots were done on genomic DNA extracted from untransformed (WT), the transformants and plasmid DNA. DNA from each sample was run on 0.8% agarose gels after the following treatments: undigested (U), digested with AvrII (A), digested with NcoI (N), digested with SacI (S). DNA from these gels was blotted on Nylon+ membranes (Amersham). These membranes were probed with a fragment corresponding to the entire coding region of the nptII gene (NeoR probe). FIG. 4 shows maps of the cassettes used in the transformations. FIG. 5 shows the results of Southern blot analysis on three transformants (all generated in UTEX strain 1435) (1, 2, and 3) transformed with either the beta-tubulin::neo::nit (SEQ ID NO: 39) (transformants 1 and 2) or glutamate dehydrogenase::neo::nit (SEQ ID NO: 40) (transformant 3). The glutamate dehydrogenase::neo::nit transforming plasmid was run as a control and cut with both NcoI and SacI. AvrII does not cut in this plasmid. Genomic DNA isolated from untransformed UTEX strain 1435 shows no hybridization to the NeoR probe.

Additional transformants containing the codon-optimized glutamate dehydrogenase::neo::nit (SEQ ID NO: 41) and codon-optimized β-tubulin::neo::nit (SEQ ID NO:42) constructs were picked and analyzed by Southern blot analysis. As expected, only digests with SacI show linearization of the transforming DNA. These transformation events are consistent with integration events that occur in the form of oligomers of the transforming plasmid. Only upon digestion with restriction enzymes that cut within the transforming plasmid DNA do these molecules collapse down the size of the transforming plasmid.

Southern blot analysis was also performed on transformants generated upon transformation of *Prototheca* strains UTEX 1437 and UTEX 1439 with the glutamate dehydrogenase::neo::nit cassette. The blot was probed with the NeoR probe and the results are similar to the UTEX 1435 transformants. The results are indicative of integration events characterized by oligomerization and integration of the transforming plasmid. This type of integration event is known to occur quite commonly in *Dictyostelium discoideum* (see, for example, Kuspa, A. and Loomis, W. (1992) *PNAS*, 89:8803-8807 and Morio et al., (1995) *J. Plant Res.* 108:111-114).

To further confirm expression of the transforming plasmid, Northern blot analysis and RT-PCR analysis were performed on selected transformants. RNA extraction was performed using Trizol Reagent according to manufacturer's instructions. Northern blot analysis were run according to methods published in Sambrook and Russel, 2001. Total RNA (15 µg) isolated from five UTEX 1435 transformants and untransformed UTEX 1435 (control lanes) was separated on 1% agarose-formaldehyde gel and blotted on nylon membrane. The blot was hybridized to the neo-non-optimized probe specific for transgene sequences in transformants 1 and 3. The two other transformants RNAs express the codon-optimized version of the neo-transgene and, as expected, based on the sequence homology between the optimized and non-optimized neo genes, showed significantly lower hybridization signal.

RNA (1 µg) was extracted from untransformed *Prototheca* strain UTEX 1435 and two representative UTEX 1435 transformants and reverse transcribed using an oligo dT primer or a gene specific primer. Subsequently these cDNAs (in duplicate) were subjected to qPCR analysis on ABI Veriti Thermocycler using SYBR-Green qPCR chemistry using the following primers (nptII):

(SEQ ID NO: 43)  
Forward: 5' GCCGCGACTGGCTGCTGCTGG 3'

(SEQ ID NO: 44)  
Reverse: 5' AGGTCCTCGCGTCGGGCATG 3'

Possible genomic DNA contamination was ruled out by a no reverse transcriptase negative control sample. The results indicated that the NeoR genes used to transform these strains is actively transcribed in the transformants.

#### C. Transformation of *Prototheca* with Secreted Heterologous Sucrose Invertase

All of the following experiments were performed using liquid medium/agar plates based on the basal medium described in Ueno et al., (2002) *J Bioscience and Bioengineering* 94(2):160-65, with the addition of trace minerals described in U.S. Pat. Nos. 5,900,370, and 1x DAS Vitamin Cocktail (1000x solution): tricine: 9 g, thiamine HCL: 0.67 g, biotin: 0.01 g, cyannocobalamin (vitamin B12): 0.008 g, calcium pantothenate: 0.02 g and p-aminobenzoic acid: 0.04 g).

Two plasmid constructs were assembled using standard recombinant DNA techniques. The yeast sucrose invertase genes (one codon optimized and one non-codon optimized), suc2, were under the control of the *Chlorella reinhardtii* beta-tubulin promoter/5'UTR and had the *Chlorella vulgaris* nitrate reductase 3'UTR. The sequences (including the 5'UTR and 3'UTR sequences) for the non-codon optimized (Crβ-tub::NCO-suc2::CvNitRed) construct, SEQ ID NO: 57, and codon optimized (Crβ-tub::CO-suc2::CvNitRed) construct, SEQ ID NO: 58, are listed in the Sequence Listing. Codon optimization was based on Table 1 for *Prototheca* sp. FIG. 6 shows a schematic of the two constructs with the relevant

restriction cloning sites and arrows indicating the direction of transcription. Selection was provided by Neo R (codon optimized using Table 1).

Preparation of the DNA/Gold Microcarrier: DNA/Gold Microcarriers were Prepared immediately before use and stored on ice until applied to macrocarriers. The plasmid DNA (in TE buffer) was added to 50  $\mu$ l of binding buffer. Saturation of the gold beads was achieved at 15  $\mu$ g plasmid DNA for 3 mg gold carrier. The binding buffer and DNA were mixed well via vortexing. The DNA and binding buffer should be pre-mix prior to gold addition to ensure uniformed plasmid binding to gold carrier particles. 60  $\mu$ l of S550d (SeaShell Technologies, San Diego, Calif.) gold carrier was added to the DNA/binding buffer mixture. For a gold stock at 50 mg/ml, addition of 60  $\mu$ l results in an optimal ratio of 15  $\mu$ g DNA/3 mg gold carrier. The gold carrier/DNA mixture was allowed to incubate on ice for 1 minute and then 100  $\mu$ l of precipitation buffer was added. The mixture was allowed to incubate again on ice for 1 minute and then briefly vortexed and centrifuged at 10,000 rpm at room temperature for 10 seconds to pellet the gold carrier. The supernatant was carefully removed with a pipette and the pellet was washed with 500  $\mu$ l of ice cold 100% ethanol. The gold particles were re-pelleted by centrifuging again at 10,000 rpm for 10 seconds. The ethanol was removed and 50  $\mu$ l of ice cold ethanol was added to the gold mixture. Immediately prior to applying the gold to macrocarriers, the gold/ethanol was resuspended with a brief 1-2 second pulse at level 2 on a MISONIX sonicator using the micro tip. Immediately after resuspension, 10  $\mu$ l of the dispersed gold particles was transferred to the macrocarrier and allowed to dry in a sterile hood.

The two *Prototheca moriformis* strains (UTEX 1435 and 1441) were grown heterotrophically in liquid medium containing 2% glucose from cryopreserved vials. Each strain was grown to a density of  $10^7$  cells/ml. This seed culture was then diluted with fresh media to a density of  $10^5$  cells/ml and allowed to grow for 12-15 hours to achieve a final cell density of approximately  $10^6$  cells/ml. The microalgae were aliquoted into 50 ml conical tubes and centrifuged for 10 minutes at 3500 rpm. The cells were washed with fresh medium and centrifuged again for 10 minutes at 3500 rpm. The cells were then resuspended at a density of  $1.25 \times 10^8$  cells/ml in fresh medium.

In a sterile hood, 0.4 ml of the above-prepared cells were removed and placed directly in the center of an agar plate (without selection agent). The plate was gently swirled with a level circular motion to evenly distribute the cells to a diameter of no more than 3 cm. The cells were allowed to dry onto the plates in the sterile hood for approximately 30-40 minutes and then were bombarded at a rupture disk pressure of 1350 psi and a plate to macrocarrier distance of 6 cm. The plates were then covered and wrapped with parafilm and allowed to incubate under low light for 24 hours.

After the 24 hour recovery, 1 ml of sterile medium (with no glucose) was added to the lawn of cells. The cells were resuspended using a sterile loop, applied in a circular motion to the lawn of cells and the resuspended cells were collected using a sterile pipette. The cells were then plated onto a fresh agar plate with 2% glucose and 100  $\mu$ g/ml G418. The appearance of colonies occurred 7-12 days after plating. Individual colonies were picked and grown in selective medium with 2% glucose and 100  $\mu$ g/ml G418. The wildtype (untransformed) and transgenic cells were then analyzed for successful introduction, integration and expression of the transgene.

Genomic DNA from transformed *Prototheca moriformis* UTEX 1435 and 1441 and their wildtype (untransformed) counterparts were isolated using standard methods. Briefly,

the cells were centrifuged for 5 minutes at 14,000 rpm in a standard table top Eppendorf centrifuge (model 5418) and flash frozen prior to DNA extraction. Cell pellets were lysed by adding 200  $\mu$ l of Lysis buffer (100 mM Tris HCl, pH 8.0, 1% Lauryl Sarcosine, 50 mM NaCl, 20 mM EDTA, 0.25 M sucrose, 0.5 mg/ml RNase A) for every 100-200 mg of cells (wet weight) and vortexing for 30-60 seconds. Cetyl trimethylammonium bromide (CTAB) and NaCl were brought to 1% and 1 M, respectively, and cell extracts were incubated at 60-65 $^{\circ}$  C. for 10 minutes. Subsequently, extracts were clarified via centrifugation at 14,000 rpm for 10 minutes and the resulting supernatant was extracted with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1). Samples were then centrifuged for 5 minutes at 14,000 rpm and the aqueous phase removed. DNA was precipitated with 0.7 volumes of isopropanol. DNA was pelleted via centrifugation at 14,000 rpm for 10 minutes and washed twice with 80% ethanol, and once with ethanol. After drying, DNA was resuspended in 10 mM Tris HCl, pH 8.0 and DNA concentrations were determined by using PicoGreen fluorescence quantification assay (Molecular Probes).

RNA from transformed *Prototheca moriformis* UTEX 1435 and 1441 and their wildtype (untransformed) counterparts were isolated using standard methods. Briefly, the cells were centrifuged for 5 minutes at 14,000 rpm in a standard table top Eppendorf centrifuge (model 5418) and flash frozen before RNA extraction. Cell pellets were lysed by addition of 1 mL of Trizol reagent (Sigma) for every 100 mg of cells (wet weight) and by vortexing for 1-2 minutes. Samples were incubated at room temperature for 5 minutes and subsequently adjusted with 200  $\mu$ l of chloroform per 1 mL of Trizol reagent. After extensive shaking, cells were incubated at room temperature for 15 minutes and then subjected to centrifugation at 14000 rpm for 15 minutes in a refrigerated table top microcentrifuge. RNA partitioning to the upper aqueous phase was removed and precipitated by addition of isopropanol (500  $\mu$ l per 1 ml of Trizol reagent). RNA was collected by centrifugation for 10 minutes and the resulting pellet washed twice with 1 mL of 80% ethanol, dried, and resuspended in RNase free water. RNA concentration was estimated by RiboGreen fluorescence quantification assay (Molecular Probes).

Expression of neomycin phosphotransferase gene conferring G418 antibiotic resistance and yeast invertase was assayed in non-transformed *Prototheca moriformis* UTEX 1435 and 1441 and transformants T98 (UTEX 1435 transformant) and T97 (UTEX 1441 transformant) using reverse transcription quantitative PCR analysis (RT-qPCR). 20 ng total RNA (isolated as described above) was subjected to one step RT-qPCR analysis using Script SYBR Green RT-PCR kit (BioRad Laboratories) and primer pairs targeting the neomycin resistance gene (forward primer 5'CCGCCGTGCTG-GACGTGGTG 3' and reverse primer 5' GGTG-GCGGGTCCAGGGTGT 3'; SEQ ID NOS: 65 and 66, respectively) and suc2 invertase transcripts (forward primer 5' CGGCCGCGGCTCCTTCAAC 3' and reverse primer 5' GGCGTCCCCGTAGGTCGGGT 3'; SEQ ID NO: 67 and 68, respectively). Endogenous beta-tubulin transcripts served as an internal positive control for PCR amplification and as a normalization reference to estimate relative transcript levels.

Both codon optimized and non-codon optimized constructs were transformed into UTEX 1435 and 1441 *Prototheca moriformis* cells as described above. Initially, transformants were obtained with both constructs and the presence of the transgene was verified by Southern blot analysis followed by RTPCR to confirm the presence of the DNA and mRNA from the transgene. For the Southern blot analysis, genomic

DNA isolated as described above was electrophoresed on 0.7% agarose gels in 1×TAE buffer. Gells were processed as described in Sambrook et al. (Molecular Cloning; A Laboratory Manual, 2<sup>nd</sup> Edition. Cold Spring Harbor Laboratory Press, 1989). Probes were prepared by random priming and hybridizations carried out as described in Sambrook et al. Transformants from both the codon optimized and the non-codon optimized constructs showed the presence of the invertase cassette, while the non-transformed control was negative. Invertase mRNA was also detected in transformants with both the codon optimized and non-codon optimized constructs.

To confirm that the transformants were expressing an active invertase protein, the transformants were plated on sucrose plates. The transformants containing the non-codon optimized cassette failed to grow on the sucrose containing plates, indicating that, while the gene and the mRNA encoding the SUC2 protein were present, the protein was either (1) not being translated, or (2) being translated, but not accumulating to levels sufficient to allow for growth on sucrose as the sole carbon source. The transformants with the codon optimized cassette grew on the sucrose containing plates. To assess the levels of invertase being expressed by these transformants, two clones (T98 and T97) were subjected to an invertase assay of whole cells scraped from solid medium and direct sampling and quantitation of sugars in the culture supernatants after 48 hours of growth in liquid medium containing 2% sucrose as the sole carbon source.

For the invertase assay, the cells (T98 and T97) were grown on plates containing 2% sucrose, scraped off and assayed for invertase activity. 10 µl of the scraped cells was mixed with 40 µl of 50 mM NaOAc pH 5.1. 12.5 µl of 0.5M sucrose was added to the cell mixture and incubated at 37° C. for 10-30 minutes. To stop the reaction, 75 µl of 0.2M K<sub>2</sub>HPO<sub>4</sub> was added. To assay for glucose liberated, 500 µl of reconstituted reagent (glucose oxidase/peroxidase+o-Dianisidine) from Sigma (GAGO-20 assay kit) was added to each tube and incubated at 37° C. for 30 minutes. A glucose standard curve was also created at this time (range: 25 µg to 0.3 µg glucose). After incubation, 500 µl of 6N HCl was added to stop the reaction and to develop the color. The samples were read at 540 nm. The amount of glucose liberated was calculated from the glucose standard curve using the formula  $y=mx+c$ , where y is the 540 nm reading, and x is µg of glucose. Weight of glucose was converted to moles of glucose, and given the equimolar relationship between moles of sucrose hydrolyzed to moles of glucose generated, the data was expressed as nmoles of sucrose hydrolyzed per unit time. The assay showed that both T98 and T97 clones were able to hydrolyze sucrose, indicating that a functional sucrose invertase was being produced and secreted by the cells.

For the sugar analysis on liquid culture media after 48 hours of algal growth, T97 and T98 cells were grown in 2% sucrose containing medium for 48 hours and the culture media were processed for sugar analysis. Culture broths from each transformant (and negative non-transformed cell control) were centrifuged at 14,000 rpm for 5 minutes. The resulting supernatant was removed and subjected to HPLC/ELSD (evaporative light scattering detection). The amount of sugar in each sample was determined using external standards and liner regression analysis. The sucrose levels in the culture media of the transformants were very low (less than 1.2 g/L, and in most cases 0 g/L). In the negative controls, the sucrose levels remained high, at approximately 19 g/L after 48 hours of growth.

These results were consistent with the invertase activity results, and taken together, indicated that the codon optimized

transformants, T97 and T98, secreted an active sucrose invertase that allowed the microalgae to utilize sucrose as the sole carbon source in contrast to (1) the non-codon optimized transformants and (2) the non-transformed wildtype microalgae, both of which could not utilize sucrose as the sole carbon source in the culture medium.

*Prototheca moriformis* strains, T98 and T97, expressing a functional, secreted sucrose invertase (SUC2) transgene were assayed for growth and lipid production using sucrose as the sole carbon source.

Wild type (untransformed), T98 and T97 strains were grown in growth media (as described above) containing either 4% glucose or 4% sucrose as the sole carbon source under heterotrophic conditions for approximately 6 days. Growth, as determined by A750 optical density readings were taken of all four samples every 24 hours and the dry cell weight of the cultures and lipid profiles were determined after the 6 days of growth. The optical density readings of the transgenic strains grown in both the glucose and sucrose conditions were comparable to the wildtype strains grown in the glucose conditions. These results indicate that the transgenic strains were able to grow on either glucose or sucrose as the sole carbon source at a rate equal to wildtype strains in glucose conditions. The non-transformed, wildtype strains did not grow in the sucrose-only condition.

The biomass for the wildtype strain grown on glucose and T98 strain grown on sucrose was analyzed for lipid profile. Lipid samples were prepared from dried biomass (lyophilized) using an Acid Hydrolysis System (Ankom Technology, N.Y.) according to manufacturer's instructions. Lipid profile determinations were carried as described in Example 4. The lipid profile for the non-transformed *Prototheca moriformis* UTEX 1435 strain, grown on glucose as the sole carbon source and two clonal T98 strains (UTEX 1435 transformed with a sucrose invertase transgene), grown on sucrose as the sole carbon source, are disclosed in Table 13 (wildtype UTEX 1435 and T98 clone 8 and clone 11 below. C:19:0 lipid was used as an internal calibration control.

TABLE 13

Lipid profile of wildtype UTEX 1435 and UTEX 1435 clones with <i>suc2</i> transgene.			
Name	wildtype (Area % - ISTD)	T98 clone 11 (Area % - ISTD)	T98 clone 8 (Area % - ISTD)
C 12:0	0.05	0.05	0.05
C 14:0	1.66	1.51	1.48
C 14:1	0.04	nd	nd
C 15:0	0.05	0.05	0.04
C 16:0	27.27	26.39	26.50
C 16:1	0.86	0.80	0.84
C 17:0	0.15	0.18	0.14
C 17:1	0.05	0.07	0.05
C 18:0	3.35	4.37	4.50
C 18:1	53.05	54.48	54.50
C 18:2	11.79	10.33	10.24
C 19:0 (ISTD)	—	—	—
C 18:3 alpha	0.90	0.84	0.81
C 20:0	0.32	0.40	0.38
C 20:1	0.10	0.13	0.12
C 20:1	0.04	0.05	0.04
C 22:0	0.12	0.16	0.12
C 20:3	0.07	0.08	0.07
C 24:0	0.12	0.11	0.10

nd—denotes none detected

Oil extracted from wildtype *Prototheca moriformis* UTEX 1435 (via solvent extraction or using an expeller press (see methods in Example 44 above) was analyzed for carotenoids,

chlorophyll, tocopherols, other sterols and tocotrienols. The results are summarized below in Table 14.

TABLE 14

Carotenoid, chlorophyll, tocopherol/sterols and tocotrienol analysis in oil extracted from <i>Prototheca moriformis</i> (UTEX 1435).		
	Pressed oil (mcg/ml)	Solvent extracted oil (mcg/ml)
cis-Lutein	0.041	0.042
trans-Lutein	0.140	0.112
trans-Zeaxanthin	0.045	0.039
cis-Zeaxanthin	0.007	0.013
t-alpha-Crytoxanthin	0.007	0.010
t-beta-Crytoxanthin	0.009	0.010
t-alpha-Carotene	0.003	0.001
c-alpha-Carotene	none detected	none detected
t-beta-Carotene	0.010	0.009
9-cis-beta-Carotene	0.004	0.002
Lycopene	none detected	none detected
Total Carotenoids	0.267	0.238
Chlorophyll	<0.01 mg/kg	<0.01 mg/kg

Tocopherols and Sterols		
	Pressed oil (mg/100 g)	Solvent extracted oil (mg/100 g)
gamma Tocopherol	0.49	0.49
Campesterol	6.09	6.05
Stigmasterol	47.6	47.8
Beta-sitosterol	11.6	11.5
Other sterols	445	446

Tocotrienols		
	Pressed oil (mg/g)	Solvent extracted oil (mg/g)
alpha Tocotrienol	0.26	0.26
beta Tocotrienol	<0.01	<0.01
gamma Tocotrienol	0.10	0.10
delta Tocotrienol	<0.01	<0.01
Total Tocotrienols	0.36	0.36

The ability of using sucrose as the sole carbon source as the selection factor for clones containing the *suc2* transgene construct instead of G418 (or another antibiotic) was assessed using the positive *suc2* gene transformants. A subset of the positive transformants was grown on plates containing sucrose as the sole carbon source and without antibiotic selection for 24 doublings. The clones were then challenged with plates containing glucose as the sole carbon source and G418. There was a subset of clones that did not grow on the glucose+G418 condition, indicating a loss of expression of the transgene. An additional experiment was performed using a plate containing sucrose as the sole carbon source and no G418 and streaking out a *suc2* transgene expressing clone on one half of the plate and wild-type *Prototheca moriformis* on the other half of the plate. Growth was seen with both the wild-type and transgene-containing *Prototheca moriformis* cells. Wild-type *Prototheca moriformis* has not demonstrated the ability to grow on sucrose, therefore, this result shows that unlike antibiotic resistance, the use of sucrose/invertase selection is not cell-autonomous. It is very likely that the transformants were secreting enough sucrose invertase into the plate/media to support wildtype growth as the sucrose was hydrolyzed into fructose and glucose.

#### Example 4

##### Recombinant *Prototheca* with Exogenous TE Gene

As described above, *Prototheca* strains can be transformed with exogenous genes. *Prototheca moriformis* (UTEX 1435)

was transformed, using methods described above, with either *Umbellularia californica* C12 thioesterase gene or *Cinnamomum camphora* C14 thioesterase gene (both codon optimized according to Table 1). Each of the transformation constructs contained a *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR region (SEQ ID NO: 69) to drive expression of the thioesterase transgene. The thioesterase transgenes coding regions of *Umbellularia californica* C12 thioesterase (SEQ ID NO: 70) or *Cinnamomum camphora* C14 thioesterase (SEQ ID NO: 71), each with the native putative plastid targeting sequence. Immediately following the thioesterase coding sequence is the coding sequence for a c-terminal 3x-FLAG tag (SEQ ID NO: 72), followed by the *Chlorella vulgaris* nitrate reductase 3'UTR (SEQ ID NO: 73). A diagram of the thioesterase constructs that were used in the *Prototheca moriformis* transformations is shown in FIG. 9.

Preparation of the DNA, Gold Microcarrier and *Prototheca Moriformis* (Utex 1435) cells were performed using the methods described above in Example 3. The microalgae were bombarded using the gold microcarrier—DNA mixture and plated on selection plates containing 2% glucose and 100 µg/ml G418. The colonies were allowed to develop for 7 to 12 days and colonies were picked from each transformation plate and screened for DNA construct incorporation using Southern blots assays and expression of the thioesterase constructs were screened using RT-PCR.

Positive clones were picked from both the C12 and C14 thioesterase transformation plates and screened for construct incorporation using Southern blot assays. Southern blot assays were carried out using standard methods (and described above in Example 3) using an optimized c probes, based on the sequence in SEQ ID NO: 70 and SEQ ID NO: 71. Transforming plasmid DNA was run as a positive control. Out of the clones that were positive for construct incorporation, a subset was selected for reverse transcription quantitative PCR (RT-qPCR) analysis for C12 thioesterase and C14 thioesterase expression.

RNA isolation was performed using methods described in Example 3 above and RT-qPCR of the positive clones were performed using 20 ng of total RNA from each clone using the below-described primer pair and iScript SYBR Green RT-PCR kit (Bio-Rad Laboratories) according to manufacturer's protocol. Wildtype (non-transformed) *Prototheca moriformis* total RNA was included as a negative control. mRNA expression was expressed as relative fold expression (RFE) as compared to negative control. The primers that were used in the C12 thioesterase transformation RT-qPCR screening were:

*U. californica* C12 thioesterase PCR primers:

(SEQ ID NO: 74)  
Forward: 5' CTGGCGACGGCTTCGGCAC 3'

(SEQ ID NO: 75)  
Reverse: 5' AAGTCGCGGCGCATGCCGTT 3'

The primers that were used in the C14 thioesterase transformation RT-qPCR screening were:

*Cinnamomum camphora* C14 thioesterase PCR primers:

(SEQ ID NO: 76)  
Forward: 5' TACCCCGCTGGGGCGACAC 3'

(SEQ ID NO: 77)  
Reverse: 5' CTTGCTCAGGCGGGTGC 3'

RT-qPCR results for C12 thioesterase expression in the positive clones showed an increased RFE of about 40 fold to over 2000 fold increased expression as compared to negative



TABLE 16-continued

Summary of total lipid profile of the <i>Prototheca moriformis</i> C14 thioesterase transformants.								
	Wildtype	C14-1	C14-2	C14-3	C14-4	C14-5	C14-6	C14-7
C20:0	0.15	0.16	0.19	0.17	0.17	0.14	0.18	0.16
C20:1	0.22	0.20	0.12	.019	0.19	0.19	0.17	0.20
C20:2	0.05	0.04	0.02	0.03	0.04	0.05	0.03	0.04
C22:0	nd	nd	nd	nd	0.02	0.01	nd	nd
C22:1	nd	0.01	nd	nd	nd	nd	nd	0.01
C20:3	0.05	0.08	0.03	0.06	0.09	0.05	0.05	0.07
C20:4	nd	0.01	nd	nd	nd	nd	0.02	nd
C24:0	nd	0.17	0.14	0.19	0.20	0.16	0.22	0.17

The above-described experiments indicate the successful transformation of *Prototheca moriformis* (UTEX 1435) with transgene constructs of two different thioesterases (C12 and C14), which involved not only the successful expression of the transgene, but also the correct targeting of the expressed protein to the plastid and a functional effect (the expected change in lipid profile) as a result of the transformation. The same transformation experiment was performed using an expression construct containing a codon-optimized (according to Table 1) *Cuphea hookeriana* C8-10 thioesterase coding region with the native plastid targeting sequence (SEQ ID NO: 78) yielded no change in lipid profile. While the introduction of the *Cuphea hookeriana* C8-10 transgene into *Prototheca moriformis* (UTEX 1435) was successful and confirmed by Southern blot analysis, no change in C8 or C10 fatty acid production was detected in the transformants compared to the wildtype strain.

#### Example 5

##### Generation of *Prototheca moriformis* Strain with Exogenous Plant TE with Algal Plastid Targeting Sequence

In order to investigate whether the use of algal chloroplast/plastid targeting sequences would improve medium chain

(C8-C14) thioesterase expression and subsequent medium chain lipid production in *Prototheca moriformis* (UTEX 1435), several putative algal plastid targeting sequences were cloned from *Chlorella protothecoides* and *Prototheca moriformis*. Thioesterase constructs based on *Cuphea hookeriana* C8-10 thioesterase, *Umbellularia californica* C12 thioesterase, and *Cinnamomum camphora* C14 thioesterase were made using made with a *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR and a *Chlorella vulgaris* nitrate reductase 3'UTR. The thioesterase coding sequences were modified by removing the native plastid targeting sequences and replacing them with plastid targeting sequences from the *Chlorella protothecoides* and the *Prototheca moriformis* genomes. The thioesterase expression constructs and their corresponding sequence identification numbers are listed below. Each transformation plasmid also contained a Neo resistance construct that was identical to the ones described in Example 3 above. Additionally, another algal-derived promoter, the *Chlamydomonas reinhardtii*  $\beta$ -tubulin promoter, was also tested in conjunction with the thioesterase constructs. "Native" plastid targeting sequence refers to the higher plant thioesterase plastid targeting sequence. A summary of the constructs used in these experiments is provided below:

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 1	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 79
Construct 2	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>P. moriformis</i> delta 12 fatty acid desaturase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 80
Construct 3	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>P. moriformis</i> isopentenyl diphosphate synthase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 81
Construct 4	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>P. moriformis</i> isopentenyl diphosphate synthase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 82
Construct 5	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>P. moriformis</i> stearyl ACP desaturase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 83
Construct 6	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 84
Construct 7	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>P. moriformis</i> delta 12 fatty acid desaturase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 85
Construct 8	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 86

-continued

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 9	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	Native	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 113
Construct 10	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> isopentenyl diphosphate synthase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 114
Construct 11	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> delta 12 fatty acid desaturase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 115
Construct 12	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 116
Construct 13	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> stearyl ACP desaturase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 117
Construct 14	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	Native	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 118
Construct 15	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> isopentenyl diphosphate	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 119
Construct 16	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> delta 12 fatty acid desaturase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 120
Construct 17	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 121
Construct 18	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> stearyl ACP desaturase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 122
Construct 19	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	Native	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 123
Construct 20	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> isopentenyl diphosphate synthase	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 124
Construct 21	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> delta 12 fatty acid desaturase	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO:
Construct 22	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 87
Construct 23	<i>Chlamydomonas reinhardtii</i> $\beta$ -tubulin	<i>P. moriformis</i> stearyl ACP desaturase	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 88

Each construct was transformed into *Prototheca moriformis* (UTEX 1435) and selection was performed using G418 using the methods described in Example 4 above. Several positive clones from each transformation were picked and screened for the presence thioesterase transgene using Southern blotting analysis. Expression of the thioesterase transgene was confirmed using RT-PCR. A subset of the positive clones (as confirmed by Southern blotting analysis and RT-PCR) from each transformation was selected and grown for lipid profile analysis. Lipid samples were prepared from dried biomass samples of each clone and lipid profile analysis was performed using acid hydrolysis methods described in Example 4. Changes in area percent of the fatty acid corresponding to the thioesterase transgene were compared to wildtype levels, and clones transformed with a thioesterase with the native plastid targeting sequence.

As mentioned in Example 4, the clones transformed with *Cuphea hookeriana* C8-10 thioesterase constructs with the native plastid targeting sequence had the same level of C8 and C10 fatty acids as wildtype. The clones transformed with *Cuphea hookeriana* C8-10 thioesterase constructs (Constructs 1-3) with algal plastid targeting sequences had over a

10-fold increase in C10 fatty acids for Construct 3 and over 40-fold increase in C10 fatty acids for Constructs 1 and 2 (as compared to wildtype). The clones transformed with *Umbellularia californica* C12 thioesterase constructs with the native plastid targeting sequence had a modest 6-8 fold increase in C12 fatty acid levels as compared to wildtype. The clones transformed with the *Umbellularia californica* C12 thioesterase constructs with the algal plasmid targeting constructs (Constructs 4-7) had over an 80-fold increase in C12 fatty acid level for Construct 4, about a 20-fold increase in C12 fatty acid level for Construct 6, about a 10-fold increase in C12 fatty acid level for Construct 7 and about a 3-fold increase in C12 fatty acid level for Construct 5 (all compared to wildtype). The clones transformed with *Cinnamomum camphora* C14 thioesterase with either the native plastid targeting sequence or the construct 8 (with the *Chlorella protothecoides* stearyl ACP desaturase plastid targeting sequence) had about a 2-3 fold increase in C14 fatty acid levels as compared to wildtype. In general clones transformed with an algal plastid targeting sequence thioesterase constructs had a higher fold increase in the corresponding chain-length fatty acid levels than when using the native higher plant targeting sequence.



A. *Chlamydomonas reinhardtii*  $\beta$ -Tubulin Promoter

Additional heterologous thioesterase expression constructs were prepared using the *Chlamydomonas reinhardtii*  $\beta$ -tubulin promoter instead of the *C. sorokinana* glutamate dehydrogenase promoter. The construct elements and sequence of the expression constructs are listed above. Each construct was transformed into *Prototheca moriformis* UTEX 1435 host cells using the methods described above. Lipid profiles were generated from a subset of positive clones for each construct in order to assess the success and productivity of each construct. The lipid profiles compare the fatty acid levels (expressed in area %) to wildtype host cells. The "Mean" column represents the numerical average of the subset of positive clones. The "Sample" column represents the best positive clone that was screened (best being defined as the sample that produced the greatest change in area % of the corresponding chain-length fatty acid production). The "low-high" column represents the lowest area % and the highest area % of the fatty acid from the clones that were screened. The lipid profiles results of Constructs 9-23 are summarized below.

Construct 9. *Cuphea hookeriana* C8-10 TE

Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.05	0.30	0-0.29
C 10:0	0.01	0.63	2.19	0-2.19
C 12:0	0.03	0.06	0.10	0-0.10
C 14:0	1.40	1.50	1.41	1.36-3.59
C 16:0	24.01	24.96	24.20	
C 16:1	0.67	0.80	0.85	
C 17:0	0	0.16	0.16	
C 17:1	0	0.91	0	
C 18:0	4.15	17.52	3.19	
C 18:1	55.83	44.81	57.54	
C 18:2	10.14	7.58	8.83	
C 18:3 $\alpha$	0.93	0.68	0.76	
C 20:0	0.33	0.21	0.29	
C 24:0	0	0.05	0.11	

Construct 10. *Cuphea hookeriana* C8-10 TE

Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.01	0.02	0-0.03
C 10:0	0	0.16	0.35	0-0.35
C 12:0	0.04	0.05	0.07	0-0.07
C 14:0	1.13	1.62	1.81	0-0.05
C 14:1	0	0.04	0.04	
C 15:0	0.06	0.05	0.05	
C 16:0	19.94	26.42	28.08	
C 16:1	0.84	0.96	0.96	
C 17:0	0.19	0.14	0.13	
C 17:1	0.10	0.06	0.05	
C 18:0	2.68	3.62	3.43	
C 18:1	63.96	54.90	53.91	
C 18:2	9.62	9.83	9.11	
C 18:3 $\gamma$	0	0.01	0	
C 18:3 $\alpha$	0.63	0.79	0.73	
C 20:0	0.26	0.35	0.33	
C 20:1	0.06	0.08	0.09	
C 20:1	0.08	0.06	0.07	
C 22:0	0	0.08	0.09	
C 24:0	0.13	0.13	0.11	

Construct 11. *Cuphea hookeriana* C8-10 TE

Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.82	1.57	0-1.87
C 10:0	0	3.86	6.76	0-6.76
C 12:0	0.04	0.13	0.20	0.03-0.20
C 14:0	1.13	1.80	1.98	1.64-2.05
C 14:1	0	0.04	0.04	
C 15:0	0.06	0.06	0.06	
C 16:0	19.94	25.60	25.44	
C 16:1	0.84	1.01	1.02	
C 17:0	0.19	0.13	0.11	
C 17:1	0.10	0.06	0.05	
C 18:0	2.68	2.98	2.38	
C 18:1	63.96	51.59	48.85	
C 18:2	9.62	9.85	9.62	
C 18:3 $\gamma$	0	0.01	0	
C 18:3 $\alpha$	0.63	0.91	0.92	
C 20:0	0.26	0.29	0.26	
C 20:1	0.06	0.06	0	
C 20:1	0.08	0.06	0.03	
C 22:0	0	0.08	0.08	
C 24:0	0.13	0.06	0	

Construct 12. *Cuphea hookeriana* C8-10 TE

Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.31	0.85	0-0.85
C 10:0	0	2.16	4.35	0.20-4.35
C 12:0	0.04	0.10	0.15	0-0.18
C 14:0	1.13	1.96	1.82	1.66-2.97
C 14:1	0	0.03	0.04	
C 15:0	0.06	0.07	0.07	
C 16:0	19.94	26.08	25.00	
C 16:1	0.84	1.04	0.88	
C 17:0	0.19	0.16	0.16	
C 17:1	0.10	0.05	0.07	
C 18:0	2.68	3.02	3.19	
C 18:1	63.96	51.08	52.15	
C 18:2	9.62	11.44	9.47	
C 18:3 $\gamma$	0	0.01	0	
C 18:3 $\alpha$	0.63	0.98	0.90	
C 20:0	0.26	0.30	0.28	
C 20:1	0.06	0.06	0.05	
C 20:1	0.08	0.04	0	
C 22:0	0	0.07	0	
C 24:0	0.13	0.05	0	

Construct 14. *Umbellularia californica* C12 TE

Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.02	0.03	0.02-0.03
C 12:0	0.03	2.62	3.91	0.04-3.91
C 14:0	1.40	1.99	2.11	1.83-2.19
C 16:0	24.01	27.64	27.01	
C 16:1	0.67	0.92	0.92	
C 18:0	4.15	2.99	2.87	
C 18:1	55.83	53.22	52.89	
C 18:2	10.14	8.68	8.41	
C 18:3 $\alpha$	0.93	0.78	0.74	
C 20:0	0.33	0.29	0.27	

Construct 15. <i>Umbellularia californica</i> C12 TE					Construct 18. <i>Umbellularia californica</i> C12 TE				
Fatty Acid	wildtype	Mean	Sample	low/high	Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.05	0.08	0-0.08	C 10:0	0	0.03	0.05	0.01-0.05
C 12:0	0.04	8.12	12.80	4.35-12.80	C 12:0	0.04	5.06	7.77	0.37-7.77
C 13:0	0	0.02	0.03	0-0.03	C 13:0	0	0.02	0	0-0.03
C 14:0	1.13	2.67	3.02	2.18-3.37	C 14:0	1.13	2.11	2.39	1.82-2.39
C 14:1	0	0.04	0.03	0.03-0.10	C 14:1	0	0.03	0.03	0.02-0.05
C 15:0	0.06	0.07	0.06		C 15:0	0.06	0.06	0.06	
C 16:0	19.94	25.26	23.15		C 16:0	19.94	24.60	23.95	
C 16:1	0.84	0.99	0.86		C 16:1	0.84	0.86	0.83	
C 17:0	0.19	0.14	0.14		C 17:0	0.19	0.15	0.14	
C 17:1	0.10	0.05	0.05		C 17:1	0.10	0.06	0.05	
C 18:0	2.68	2.59	2.84		C 18:0	2.68	3.31	2.96	
C 18:1	63.96	46.91	44.93		C 18:1	63.96	51.26	49.70	
C 18:2	9.62	10.59	10.01		C 18:2	9.62	10.18	10.02	
C 18:3 $\alpha$	0.63	0.92	0.83		C 18:3 $\gamma$	0	0.01	0.02	
C 20:0	0.26	0.27	0.24		C 18:3 $\alpha$	0.63	0.86	0.86	
C 20:1	0.06	0.06	0.06		C 20:0	0.26	0.32	0.29	
C 20:1	0.08	0.05	0.04		C 20:1	0.06	0.05	0.05	
C 22:0	0	0.07	0.09		C 20:1	0.08	0.07	0.04	
C 24:0	0.13	0.13	0.12		C 22:0	0	0.08	0.08	
					C 24:0	0.13	0.13	0.13	
Construct 16. <i>Umbellularia californica</i> C12 TE					Construct 19. <i>Cinnamomum camphora</i> C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high	Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.03	0.04	0.02-0.04	C 10:0	0.02	0.01	0.01	0.01-0.02
C 12:0	0.04	2.43	5.32	0.98-5.32	C 12:0	0.05	0.27	0.40	0.08-0.41
C 13:0	0	0.01	0.02	0-0.02	C 14:0	1.52	4.47	5.81	2.10-5.81
C 14:0	1.13	1.77	1.93	1.62-1.93	C 16:0	25.16	28.14	28.55	
C 14:1	0	0.03	0.02	0.02-0.04	C 16:1	0.72	0.84	0.82	
C 15:0	0.06	0.06	0.05		C 18:0	3.70	3.17	2.87	
C 16:0	19.94	24.89	22.29		C 18:1	54.28	51.89	51.01	
C 16:1	0.84	0.91	0.82		C 18:2	12.24	9.36	8.62	
C 17:0	0.19	0.16	0.15		C 18:3 $\alpha$	0.87	0.74	0.75	
C 17:1	0.10	0.06	0.06		C 20:0	0.33	0.33	0.31	
C 18:0	2.68	3.81	3.67						
C 18:1	63.96	53.19	52.82						
C 18:2	9.62	10.38	10.57						
C 18:3 $\alpha$	0.63	0.80	0.77						
C 20:0	0.26	0.35	0.32						
C 20:1	0.06	0.06	0.07						
C 20:1	0.08	0.07	0.08						
C 22:0	0	0.08	0.07						
C 24:0	0.13	0.15	0.14						
Construct 17. <i>Umbellularia californica</i> C12 TE					Construct 20. <i>Cinnamomum camphora</i> C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high	Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.04	0.07	0.03-0.08	C 10:0	0.01	0.01	0.02	0.01-0.02
C 12:0	0.04	7.02	14.11	4.32-14.11	C 12:0	0.03	0.39	0.65	0.08-0.65
C 13:0	0	0.03	0.04	0.01-0.04	C 13:0	0	0.01	0.01	0.01-0.02
C 14:0	1.13	2.25	3.01	1.95-3.01	C 14:0	1.40	5.61	8.4	2.1-8.4
C 14:1	0	0.03	0.03	0.02-0.03	C 14:1	0	0.03	0.03	0.02-0.03
C 15:0	0.06	0.06	0.06		C 15:0	0	0.06	0.07	
C 16:0	19.94	23.20	21.46		C 16:0	24.01	25.93	25.57	
C 16:1	0.84	0.82	0.77		C 16:1	0.67	0.75	0.71	
C 17:0	0.19	0.15	0.14		C 17:0	0	0.13	0.12	
C 17:1	0.10	0.06	0.06		C 17:1	0	0.05	0.05	
C 18:0	2.68	3.47	2.93		C 18:0	4.15	3.30	3.23	
C 18:1	63.96	50.30	45.17		C 18:1	55.83	51.00	48.48	
C 18:2	9.62	10.33	9.98		C 18:2	10.14	10.38	10.35	
C 18:3 $\gamma$	0	0.01	0		C 18:3 $\alpha$	0.93	0.91	0.88	
C 18:3 $\alpha$	0.63	0.84	0.86		C 20:0	0.33	0.35	0.32	
C 20:0	0.26	0.32	0.27		C 20:1	0	0.08	0.08	
C 20:1	0.06	0.07	0.06		C 20:1	0	0.07	0.07	
C 20:1	0.08	0.06	0.06		C 22:0	0	0.08	0.08	
C 22:0	0	0.08	0.09		C 24:0	0	0.14	0.13	
C 24:0	0.13	0.14	0.13						

Construct 21. <i>Cinnamomum camphora</i> C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.01	0.01	0-0.01
C 12:0	0.03	0.10	0.27	0.04-0.27
C 14:0	1.40	2.28	4.40	1.47-4.40
C 16:0	24.01	26.10	26.38	
C 16:1	0.67	0.79	0.73	
C 17:0	0	0.15	0.16	
C 17:1	0	0.06	0.06	
C 18:0	4.15	3.59	3.51	
C 18:1	55.83	53.53	50.86	
C 18:2	10.14	10.83	11.11	
C 18:3 $\alpha$	0.93	0.97	0.87	
C 20:0	0.33	0.36	0.37	
C 20:1	0	0.09	0.08	
C 20:1	0	0.07	0.07	
C 22:0	0	0.09	0.09	

Construct 22. <i>Cinnamomum camphora</i> C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.02	0.02	0.02-0.02
C 12:0	0.03	1.22	1.83	0.59-1.83
C 13:0	0	0.02	0.03	0.01-0.03
C 14:0	1.40	12.77	17.33	7.97-17.33
C 14:1	0	0.02	0.02	0.02-0.04
C 15:0	0	0.07	0.08	
C 16:0	24.01	24.79	24.22	
C 16:1	0.67	0.64	0.58	
C 17:0	0	0.11	0.10	
C 17:1	0	0.04	0.04	
C 18:0	4.15	2.85	2.75	
C 18:1	55.83	45.16	41.23	
C 18:2	10.14	9.96	9.65	
C 18:3 $\alpha$	0.93	0.91	0.85	
C 20:0	0.33	0.30	0.30	
C 20:1	0	0.07	0.06	
C 20:1	0	0.06	0.05	
C 22:0	0	0.08	0.08	

Construct 23. <i>Cinnamomum camphora</i> C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.01	0.02	0-0.02
C 12:0	0.05	0.57	1.08	0.16-1.08
C 13:0	0	0.02	0.02	0-0.02
C 14:0	1.45	7.18	11.24	2.96-11.24
C 14:1	0.02	0.03	0.03	0.02-0.03
C 15:0	0.06	0.07	0.07	
C 16:0	24.13	25.78	25.21	
C 16:1	0.77	0.72	0.66	
C 17:0	0.19	0.13	0.11	
C 17:1	0.08	0.05	0.04	
C 18:0	3.53	3.35	3.12	
C 18:1	56.15	49.65	46.35	
C 18:2	11.26	10.17	9.72	
C 18:3 $\alpha$	0.84	0.95	0.83	
C 20:0	0.32	0.34	0.32	
C 20:1	0.09	0.08	0.09	
C 20:1	0.07	0.05	0.06	
C 22:0	0.07	0.08	0.08	
C 24:0	0.13	0.13	0.12	

Constructs 9-13 were expression vectors containing the *Cuphea hookeriana* C8-10 thioesterase construct. As can be seen in the data summaries above, the best results were seen with Construct 11, with the Sample C8 fatty acid being 1.57

Area % (as compared to 0 in wildtype) and C10 fatty acid being 6.76 Area % (as compared to 0 in wildtype). There was also a modest increase in C12 fatty acids (approximately 2-5 fold increase). While the native plastid targeting sequence produced no change when under the control of the *C. sorokiniana* glutamate dehydrogenase promoter, the same expression construct driven by the *C. reinhardtii*  $\beta$ -tubulin promoter produced significant changes in C8-10 fatty acids in the host cell. This is further evidence of the idiosyncrasies of heterologous expression of thioesterases in *Prototheca* species. All of the clones containing the *C. reinhardtii*  $\beta$ -tubulin promoter C8-10 thioesterase construct had greater increases in C8-10 fatty acids than the clones containing the *C. sorokiniana* glutamate dehydrogenase promoter C8-10 thioesterase construct. Lipid profile data for Construct 13 was not obtained and therefore, not included above.

Constructs 14-18 were expression vectors containing the *Umbellularia californica* C12 thioesterase construct. As can be seen in the data summaries above, the best results were seen with Constructs 15 (*P. moriformis* isopentenyl diphosphate synthase plastid targeting sequence) and 17 (*C. protothecoides* stearoyl ACP desaturase plastid targeting sequence). The greatest change in C12 fatty acid production was seen with Construct 17, with C12 fatty acids levels of up to 14.11 area %, as compared to 0.04 area % in wildtype. Modest changes (about 2-fold) were also seen with C14 fatty acid levels. When compared to the same constructs with the *C. sorokiniana* glutamate dehydrogenase promoter, the same trends were true with the *C. reinhardtii*  $\beta$ -tubulin promoter—the *C. protothecoides* stearoyl ACP desaturase and *P. moriformis* isopentenyl diphosphate synthase plastid targeting sequences produced the greatest change in C12 fatty acid levels with both promoters.

Constructs 19-23 were expression vectors containing the *Cinnamomum camphora* C14 thioesterase construct. As can be seen in the data summaries above, the best results were seen with Constructs 22 and Construct 23. The greatest change in C14 fatty acid production was seen with Construct 22, with C14 fatty acid levels of up to 17.35 area % (when the values for C140 and C141 are combined), as compared to 1.40% in wildtype. Changes in C12 fatty acids were also seen (5-60 fold). When compared to the same constructs with the *C. sorokiniana* glutamate dehydrogenase promoter, the same trends were true with the *C. reinhardtii*  $\beta$ -tubulin promoter the *C. protothecoides* stearoyl ACP desaturase and *P. moriformis* stearoyl ACP desaturase plastid targeting sequences produced the greatest change in C14 fatty acid levels with both promoters. Consistently with all thioesterase expression constructs, the *C. reinhardtii*  $\beta$ -tubulin promoter constructs produced greater changes in C8-14 fatty acid levels than the *C. sorokiniana* glutamate dehydrogenase

Two positive clones from the Construct 22 were selected and grown under high selective pressure (50 mg/L G418). After 6 days in culture, the clones were harvested and their lipid profile was determined using the methods described above. The lipid profile data is summarized below and is expressed in area %.

Construct 22 clones + 50 mg/L G418		
Fatty Acid	Construct 22 A	Construct 22 B
C 12:0	3.21	3.37
C 14:0	27.55	26.99
C 16:0	25.68	24.37

-continued

Construct 22 clones + 50 mg/L G418		
Fatty Acid	Construct 22 A	Construct 22 B
C 16:1	0.99	0.92
C 18:0	1.37	1.23
C 18:1	28.35	31.07
C 18:2	11.73	11.05
C 18:3 $\alpha$	0.92	0.81
C 20:0	0.16	0.17

Both clones, when grown under constant, high selective pressure, produced an increased amount of C14 and C12 fatty acids, about double the levels seen with Construct 22 above. These clones yielded over 30 area % of C12-14 fatty acids, as compared to 1.5 area % of C12-14 fatty acids seen in wildtype cells.

## Example 6

Heterologous Expression of *Cuphea palustris* and *Ulmus americana* Thioesterase in *Prototheca*

Given the success of the above-described heterologous expression thioesterases in *Prototheca* species, expression cassettes containing codon-optimized (according to Table 1) sequences encoding fatty acyl-ACP thioesterases from *Cuphea palustris* and *Ulmus americana* were constructed and described below.

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 27	<i>C. reinhardtii</i> $\beta$ -tubulin	<i>C. protothecoides</i> stearoyl ACP desaturase	<i>Cuphea palustris</i> thioesterase	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 107

The *Ulmus americana* (codon-optimized coding sequence) can be inserted into the expression cassette. The codon-optimized coding sequence without the native plastid targeting sequence for the *Ulmus americana* thioesterase is listed as SEQ ID NO: 108 and can be fused any desired plastid targeting sequence and expression element (i.e., promoter/ 5'UTR and 3'UTR).

These expression cassettes can be transformed in to *Prototheca* species using the methods described above. Positive clones can be screened with the inclusion of an antibiotic resistance gene (e.g, neoR) on the expression construct and screened on G418-containing plates/media. Positive clones can be confirmed using Southern blot assays with probes specific to the heterologous thioesterase coding region and expression of the construct can also be confirmed using RT-PCR and primers specific to the coding region of the heterologous thioesterase. Secondary confirmation of positive clones can be achieved by looking for changes in levels of fatty acids in the host cell's lipid profile. As seen in the above Examples, heterologous expression in *Prototheca* species of

thioesterase can be idiosyncratic to the particular thioesterase. Promoter elements and plastid targeting sequences (and other expression regulatory elements) can be interchanged until the expression of the thioesterase (and the subsequent increase in the corresponding fatty acid) reaches a desired level.

## Example 7

## Dual Transformants—Simultaneous Expression of Two Heterologous Proteins

Microalgae strain *Prototheca moriformis* (UTEX 1435) was transformed using the above disclosed methods with a expression construct containing the yeast sucrose invertase *suc2* gene encoding the secreted form of the *S. cerevisiae* invertase. Successful expression of this gene and targeting to the periplasm results in the host cell's ability to grow on (and utilize) sucrose as a sole carbon source in heterotrophic conditions (as demonstrated in Example 3 above). The second set of genes expressed are thioesterases which are responsible for the cleavage of the acyl moiety from the acyl carrier protein. Specifically, thioesterases from *Cuphea hookeriana* (a C8-10 preferring thioesterase), *Umbellularia californica* (a C12 preferring thioesterase), and *Cinnamomum camphora* (a C14 preferring thioesterase). These thioesterase expression cassettes were cloned as fusions with N-terminal microalgal plastid targeting sequences from either *Prototheca moriformis*

*mis* or *Chlorella protothecoides*, which have been shown (in the above Examples) to be more optimal than the native higher plant plastid targeting sequences. The successful expression of the thioesterase genes and the targeting to the plastid resulted in measurable changes in the fatty acid profiles within the host cell. These changes in profiles are consistent with the enzymatic specificity or preference of each thioesterase. Below is a summary of dual expression constructs that were assembled and transformed into *Prototheca moriformis* (UTEX 1435). Each construct contained the yeast *suc2* gene under the control of the *C. reinhardtii*  $\beta$ -tubulin 5'UTR/promoter and contained the *C. vulgaris* nitrate reductase 3'UTR and a higher plant thioesterase with a microalgal plastid targeting sequence replacing the native sequence under the control of *C. sorokinana* glutamate dehydrogenase 5'UTR and contained the *C. vulgaris* nitrate reductase 3'UTR. Below is a summary of the thioesterase portion of the constructs that were assembled and transformed into *Prototheca moriformis* (UTEX 1435). The entire dual expression cassette with the *suc2* gene and the thioesterase gene and the is listed in the Sequence Identification Listing.

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 24	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>C. protothecoides</i> stearoyl ACP desaturase	<i>Cuphea hookeriana</i> C8-10 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 109
Construct 25	<i>C. sorokiniana</i> glutamate dehydrogenase	<i>P. moriformis</i> isopentenyl diphosphate synthase	<i>Umbellularia californica</i> C12 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 110

-continued

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 26	<i>C. sorokinana</i> glutamate dehydrogenase	<i>C. protothecoides</i> stearyl ACP desaturase	<i>Cinnamomum camphora</i> C14 TE	<i>C. vulgaris</i> nitrate reductase	SEQ ID NO: 111

Similar dual expression constructs with the thioesterase cassettes described in Example 5 (e.g., under the control of a different promoter such as *C. reinhardtii*  $\beta$ -tubulin promoter/ 5'UTR) can also be generated using standard molecular biology methods and methods described herein.

Positive clones containing each of expression constructs were screened using their ability to grow on sucrose-containing plates, where sucrose is the sole-carbon source, as the selection factor. A subset of these positive clones from each construct transformation was selected and the presence of the expression construct was confirmed using Southern blot assays. The function of the yeast sucrose invertase was also confirmed using a sucrose hydrolysis assay. Positive clones were selected and grown in media containing sucrose as the sole carbon source at a starting concentration of 40 g/L. A negative control of wildtype *Prototheca moriformis* (UTEX 1435) grown in media containing glucose as the sole carbon source at the same 40 g/L starting concentration was also included. Utilization of sucrose was measured throughout the course of the experiment by measuring the level of sucrose in the media using a YSI 2700 Biochemistry Analyzer with a sucrose-specific membrane. After six days in culture, the cultures were harvested and processed for lipid profile using the same methods as described above. The lipid profile results are summarized below in Table 17 and are show in area %.

TABLE 17

Lipid profiles of dual transformants with suc2 sucrose invertase and thioesterase.											
Fatty Acid	Wt	C24 A	C24 B	C24 C	C25 A	C25 B	C25 C	C26 A	C26 B	C26 C	
C 10:0	0.01	0.03	0.04	0.08	0.01	0.01	0.01	0.01	0.01	0.01	0.0
C 12:0	0.04	0.04	0.04	0.04	0.28	0.40	0.10	0.04	0.04	0.13	
C 14:0	1.6	1.55	1.53	1.56	1.59	1.59	1.60	1.65	1.56	2.69	
C 14:1	0.03	0.03	0.03	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
C 15:0	0.04	0.03	0.03	0.04	0.04	0.03	0.03	0.03	0.03	0.04	
C 16:0	29.2	29.1	29.0	28.6	28.9	28.6	29.0	28.8	29.5	27.5	
C 16:1	0.86	0.80	0.79	0.82	0.77	0.81	0.82	0.79	0.79	0.86	
C 17:0	0.1	0.08	0.08	0.09	0.09	0.08	0.09	0.08	0.08	0.09	
C 17:1	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.03	0.04	
C 18:0	3.26	3.33	3.37	3.27	3.36	3.28	3.18	3.33	3.36	3.03	
C 18:1	54.5	53.9	54.1	53.9	53.5	53.7	53.5	54.2	53.9	52.7	
C 18:2	8.72	9.35	9.22	9.45	9.68	9.65	9.87	9.31	9.06	10.8	
C 18:3	0.63	0.71	0.69	0.73	0.74	0.73	0.75	0.71	0.66	0.83	
alpha											
C 20:0	0.29	0.31	0.31	0.31	0.32	0.32	0.31	0.32	0.31	0.29	

All of the positive clones selected for the sucrose utilization assay were able to hydrolyze the sucrose in the media and at the end of the 6 day culture period, there were no measurable levels of sucrose in the media. This data, in addition to the successful use of sucrose as a selection tool for positive clones, indicates that the exogenous yeast suc2 sucrose invertase gene was targeted correctly and expressed in the transformants. As show in Table 17 above, the clones expressing Construct 24 (C8-10 thioesterase) had a measurable increase in C10 fatty acids (as high as an eight-fold increase). Likewise there were measurable increases in clones expressing Construct 25 (C12 thioesterase) and Construct 26 (C14

thioesterase) in the corresponding medium chain fatty acids. Taken together, the data shows the successful simultaneous expression in *Prototheca moriformis* two recombinant proteins (e.g., sucrose invertase and a fatty acid acyl-ACP thioesterase), both of which confer useful and quantifiable phenotypic changes on the host organism.

## Example 8

## Effects of Glycerol on C10-C14 Fatty Acid Production in C14 Thioesterase Transformants

Clones from all the thioesterase transformations were selected and further evaluated. One clone expressing Construct 8 (*Cinnamomum camphora* C14 TE) was grown heterotrophically using different carbon sources: glucose only, fructose only and glycerol only. The glucose only condition resulted in higher cell growth and total lipid production when compared to the fructose only and glycerol only conditions. However, the proportion of C12-14 fatty acids produced in the glycerol only condition was two-fold higher than that attained in the glucose only condition.

## Example 9

Expression of *Arabidopsis thaliana* Invertase in *Prototheca moriformis*

Microalgae strain *Prototheca moriformis* (UTEX 1435) was transformed using methods described above, with an expression construct containing a codon-optimized (according to Table 1) cell wall associated invertase from *Arabidopsis thaliana*. The *Arabidopsis* invertase sequence was modified to include the N-terminal 39 amino acids from yeast invertase (SUC2 protein) to ensure efficient targeting to the ER and

ultimately the periplasm. To aid detection, a Flag epitope was added to the C-terminus of the recombinant protein. The transgene was cloned into an expression vector with a *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR region and a *Chlorella vulgaris* nitrate reductase 3'UTR region. The DNA sequence of this transgene cassette is listed as SEQ ID NO: 89 and the translated amino acid sequence is listed as SEQ ID NO: 90. Positive clones were screened and selected using sucrose-containing media/plates. A subset of the positive clones were confirmed for the presence of the transgene and expression of invertase using Southern blot analysis and Western blot analysis for the Flag-tagged invertase. From these screens, 10 positive clones were chosen for lipid productivity and sucrose utilization assays. All 10 clones were grown on media containing sucrose as the sole carbon source and a positive control suc2 invertase transformant was also included. The negative control, wildtype *Prototheca moriformis*, was also grown but on glucose containing media. After six days, the cells were harvested and dried and the total percent lipid by dry cell weight was determined. The media was also analyzed for total sucrose consumption.

All ten positive clones were able to hydrolyze sucrose, however, most clones grew about half as well as either wildtype or the positive control suc2 yeast invertase transformant as determined by dry cell weight at the end of the experiment. Similarly, all ten positive clones produced about half as much total lipid when compared to wildtype or the positive control transformant. This data demonstrate the successful heterologous expression of diverse sucrose invertases in *Prototheca*.

#### Example 10

##### Heterologous Expression of Yeast Invertase (suc2) in *Prototheca krugani*, *Prototheca stagnora* and *Prototheca zopfii*

To test the general applicability of the transformation methods for use in species of the genus *Prototheca*, three other *Prototheca* species were selected: *Prototheca krugani* (UTEX 329), *Prototheca stagnora* (UTEX 1442) and *Prototheca zopfii* (UTEX 1438). These three strains were grown in the media and conditions described in Example 1 and their lipid profiles were determined using the above described methods. A summary of the lipid profiles from the three *Prototheca* strains are summarized below in Area %.

Fatty Acid	<i>P. krugani</i> (UTEX 329)	<i>P. stagnora</i> (UTEX 1442)	<i>P. zopfii</i> (UTEX 1438)
C 10:0	0.0	0.0	0.0
C 10:1	0.0	0.0	0.0
C 12:0	1.5	0.8	2.1
C 14:0	1.2	0.9	1.7
C 16	15.1	17.1	19.7
C 18:0	3.3	4.1	5.4
C 18:1	66.0	61.5	53.8
C 18:2	12.9	15.6	17.3

These three strains were transformed with a yeast invertase (suc2) expression cassette (SEQ ID NO: 58) using the methods described in Example 3 above. This yeast invertase (suc2) expression cassette has been demonstrated to work in *Prototheca moriformis* (UTEX 1435) above in Example 3. The transformants were screened using sucrose containing plates/media. A subset of the positive clones for each *Prototheca* species was selected and the presence of the transgene was

confirmed by Southern blot analysis. Ten of confirmed positive clones from each species were selected for sucrose hydrolysis analysis and lipid productivity. The clones were grown in media containing sucrose as the sole carbon source and compared to its wildtype counterpart grown on glucose. After 6 days, the cultures were harvested and dried and total percent lipid and dry cell weight was assessed. The media from each culture was also analyzed for sucrose hydrolysis using a YSI2700 Biochemistry Analyzer for sucrose content over the course of the experiment. Clones from all three species were able to hydrolyze sucrose, with *Prototheca stagnora* and *Prototheca zopfii* transformants being able to hydrolyze sucrose more efficiently than *Prototheca krugani*. Total lipid production and dry cell weight of the three species of transformants were comparable to their wildtype counterpart grown on glucose. This data demonstrates the successful transformation and expression exogenous genes in multiple species of the genus *Prototheca*.

#### Example 11

##### Algal-Derived Promoters and Genes for Use in Microalgae

###### A. 5'UTR and Promoter Sequences from *Chlorella protothecoides*

A cDNA library was generated from mixotrophically grown *Chlorella protothecoides* (UTEX 250) using standard techniques. Based upon the cDNA sequences, primers were designed in certain known housekeeping genes to "walk" upstream of the coding regions using Seegene's DNA Walking kit (Rockville, Md.). Sequences isolated include an actin (SEQ ID NO:31) and elongation factor-1a (EF1a) (SEQ ID NO:32) promoter/UTR, both of which contain introns (as shown in the lower case) and exons (upper case italicized) and the predicted start site (in bold) and two beta-tubulin promoter/UTR elements: Isoform A (SEQ ID NO:33) and Isoform B (SEQ ID NO:34).

###### B. Lipid Biosynthesis Enzyme and Plastid Targeting Sequences from *C. protothecoides*

From the cDNA library described above, three cDNAs encoding proteins functional in lipid metabolism in *Chlorella protothecoides* (UTEX 250) were cloned using the same methods as described above. The nucleotide and amino acid sequences for an acyl ACP desaturase (SEQ ID NOs: 45 and 46) and two geranyl geranyl diphosphate synthases (SEQ ID NOs:47-50) are included in the Sequence Listing below. Additionally, three cDNAs with putative signal sequences targeting to the plastid were also cloned. The nucleotide and amino acid sequences for a glyceraldehyde-3-phosphate dehydrogenase (SEQ ID NOs:51 and 52), an oxygen evolving complex protein OEE33 (SEQ ID NOs:53 and 54) and a Clp protease (SEQ ID NOs:55 and 56) are included in the Sequence Listing below. The putative plastid targeting sequence has been underlined in both the nucleotide and amino acid sequence. The plastid targeting sequences can be used to target the product of transgenes to the plastid of microbes, such as lipid modification enzymes.

#### Example 12

##### 5'UTR/Promoters that are Nitrogen Responsive from *Prototheca moriformis*

A cDNA library was generated from *Prototheca moriformis* (UTEX 1435) using standard techniques. The *Prototheca moriformis* cells were grown for 48 hours under nitrogen

replete conditions. Then a 5% innoculum (v/v) was then transferred to low nitrogen and the cells were harvested every 24 hours for seven days. After about 24 hours in culture, the nitrogen supply in the media was completely depleted. The collected samples were immediately frozen using dry ice and isopropanol. Total RNA was subsequently isolated from the frozen cell pellet samples and a portion from each sample was held in reserve for RT-PCR studies. The rest of the total RNA harvested from the samples was subjected to polyA selection. Equimolar amounts of polyA selected RNA from each condition was then pooled and used to generate a cDNA library in vector pcDNA 3.0 (Invitrogen). Roughly 1200 clones were randomly picked from the resulting pooled cDNA library and subjected to sequencing on both strands. Approximately 68

ID NO: 99), was cloned into the *Cinnamomum camphora* C14 thioesterase construct with the *Chlorella protothecoides* stearyl ACP desaturase transit peptide described in Example 5 above, replacing the *C. sorokinana* glutamate dehydrogenase promoter. This construct is listed as SEQ ID NO: 112. To test the putative promoter, the thioesterase construct is transformed into *Prototheca moriformis* cells to confirm actual promoter activity by screening for an increase in C14/C12 fatty acids under low/no nitrogen conditions, using the methods described above. Similar testing of the putative nitrogen-regulated promoters isolated from the cDNA/genomic screen can be done using the same methods.

Other putative nitrogen-regulated promoters/5'UTRs that were isolated from the cDNA/genomic screen were:

Promoter/5'UTR	SEQ ID NO.	Fold increased
FatB/A promoter/5'UTR	SEQ ID NO: 91	n/a
NRAMP metal transporter promoter/5'UTR	SEQ ID NO: 92	9.65
Flap Flagellar-associated protein promoter/5'UTR	SEQ ID NO: 93	4.92
SulfRed Sulfite reductase promoter/5'UTR	SEQ ID NO: 94	10.91
SugT Sugar transporter promoter/5'UTR	SEQ ID NO: 95	17.35
Amt03-Ammonium transporter 03 promoter/5'UTR	SEQ ID NO: 96	10.1
Amt02-Ammonium transporter 02 promoter/5'UTR	SEQ ID NO: 97	10.76
Aat01-Amino acid transporter 01 promoter/5'UTR	SEQ ID NO: 98	6.21
Aat02-Amino acid transporter 02 promoter/5'UTR	SEQ ID NO: 99	6.5
Aat03-Amino acid transporter 03 promoter/5'UTR	SEQ ID NO: 100	7.87
Aat04-Amino acid transporter 04 promoter/5'UTR	SEQ ID NO: 101	10.95
Aat05-Amino acid transporter 05 promoter/5'UTR	SEQ ID NO: 102	6.71

Fold increase refers to the fold increase in cDNA abundance after 24 hours of culture in low nitrogen medium.

different cDNAs were selected from among these 1200 sequences and used to design cDNA-specific primers for use in real-time RT-PCR studies.

RNA isolated from the cell pellet samples that were held in reserve was used as substrate in the real time RT-PCR studies using the cDNA-specific primer sets generated above. This reserved RNA was converted into cDNA and used as substrate for RT-PCR for each of the 68 gene specific primer sets. Threshold cycle or  $C_T$  numbers were used to indicate relative transcript abundance for each of the 68 cDNAs within each RNA sample collected throughout the time course. cDNAs showing significant increase (greater than three fold) between nitrogen replete and nitrogen-depleted conditions were flagged as potential genes whose expression was up-regulated by nitrogen depletion. As discussed in the specification, nitrogen depletion/limitation is a known inducer of lipogenesis in oleaginous microorganisms.

In order to identify putative promoters/5'UTR sequences from the cDNAs whose expression was upregulated during nitrogen depletion/limitation, total DNA was isolated from *Prototheca moriformis* (UTEX 1435) grown under nitrogen replete conditions and were then subjected to sequencing using 454 sequencing technology (Roche). cDNAs flagged as being up-regulated by the RT-PCR results above were compared using BLAST against assembled contigs arising from the 454 genomic sequencing reads. The 5' ends of cDNAs were mapped to specific contigs, and where possible, greater than 500 bp of 5' flanking DNA was used to putatively identify promoters/UTRs. The presence of promoters/5'UTR were subsequently confirmed and cloned using PCR amplification of genomic DNA. Individual cDNA 5' ends were used to design 3' primers and 5' end of the 454 contig assemblies were used to design 5' gene-specific primers.

As a first screen, one of the putative promoter, the 5'UTR/promoter isolated from Aat2 (Ammonium transporter, SEQ

### Example 13

#### Homologous Recombination in *Prototheca* Species

Homologous recombination of transgenes has several advantages over the transformation methods described in the above Examples. First, the introduction of transgenes without homologous recombination can be unpredictable because there is no control over the number of copies of the plasmid that gets introduced into the cell. Also, the introduction of transgenes without homologous recombination can be unstable because the plasmid may remain episomal and is lost over subsequent cell divisions. Another advantage of homologous recombination is the ability to "knock-out" gene targets, introduce epitope tags, switch promoters of endogenous genes and otherwise alter gene targets (e.g., the introduction of point mutations).

Two vectors were constructed using a specific region of the *Prototheca moriformis* (UTEX 1435) genome, designated KE858. KE858 is a 1.3 kb, genomic fragment that encompasses part of the coding region for a protein that shares homology with the transfer RNA (tRNA) family of proteins. Southern blots have shown that the KE858 sequence is present in a single copy in the *Prototheca moriformis* (UTEX 1435) genome. The first type of vector that was constructed, designated SZ725 (SEQ ID NO: 103), consisted of the entire 1.3 kb KE858 fragment cloned into a pUC19 vector backbone that also contains the optimized yeast invertase (*suc2*) gene used in Example 3 above. The KE858 fragment contains a unique *Sna*B1 site that does not occur anywhere else in the targeting construct. The second type of vector that was constructed, designated SZ726 (SEQ ID NO: 126), consisted of the KE858 sequence that had been disrupted by the insertion of the yeast invertase gene (*suc2*) at the *Sna*B1 site within the KE858 genomic sequence. The entire DNA fragment con-

taining the KE858 sequences flanking the yeast invertase gene can be excised from the vector backbone by digestion with EcoRI, which cuts at either end of the KE858 region.

Both vectors were used to direct homologous recombination of the yeast invertase gene (*suc2*) into the corresponding KE858 region of the *Prototheca moriformis* (UTEX 1435) genome. The linear DNA ends homologous to the genomic region that was being targeted for homologous recombination were exposed by digesting the vector construct SZ725 with SnaBI and vector construct SZ726 with EcoRI. The digested vector constructs were then introduced into *Prototheca moriformis* cultures using methods described above in Example 3. Transformants from each vector construct were then selected using sucrose plates. Ten independent, clonally pure transformants from each vector transformation were analyzed for successful recombination of the yeast invertase gene into the desired genomic location (using Southern blots) and for transgene stability.

Southern blot analysis of the SZ725 transformants showed that 4 out of the 10 transformants picked for analysis contained the predicted recombinant bands, indicating that a single crossover event had occurred between the KE858 sequences on the vector and the KE858 sequences in the genome. In contrast, all ten of the SZ726 transformants contained the predicted recombinant bands, indicating that double crossover events had occurred between the EcoRI fragment of pSZ726 carrying KE858 sequence flanking the yeast invertase transgene and the corresponding KE858 region of the genome.

Sucrose invertase expression and transgene stability were assessed by growing the transformants for over 15 generations in the absence of selection. The four SZ725 transformants and the ten SZ726 transformants that were positive for the transgene by Southern blotting were selected and 48 single colonies from each of the transformants were grown serially: first without selection in glucose containing media and then with selection in media containing sucrose as the sole carbon source. All ten SZ726 transformants (100%) retained their ability to grow on sucrose after 15 generations, whereas about 97% of the SZ725 transformants retained their ability to grow on sucrose after 15 generations. Transgenes introduced by a double crossover event (SZ726 vector) have extremely high stability over generation doublings. In contrast, transgenes introduced by a single cross over event (SZ725 vector) can result in some instability over generation doublings because in tandem copies of the transgenes were introduced, the repeated homologous regions flanking the transgenes may recombine and excise the transgenic DNA located between them.

These experiments demonstrate the successful use of homologous recombination to generate *Prototheca* transformants containing a heterologous sucrose invertase gene that is stably integrated into the nuclear chromosomes of the organism. The success of the homologous recombination enables other genomic alterations in *Prototheca*, including gene deletions, point mutations and epitope tagging a desired gene product. These experiments also demonstrate the first documented system for homologous recombination in the nuclear genome of an eukaryotic microalgae.

#### A. Use of Homologous Recombination to Knock-Out an Endogenous *Prototheca moriformis* Gene

In the *Prototheca moriformis* cDNA/genomic screen described in Example 11 above, an endogenous stearoyl ACP desaturase (SAPD) cDNA was identified. Stearoyl ACP desaturase enzymes are part of the lipid synthesis pathway and they function to introduce double bonds into the fatty acyl chains. In some cases, it may be advantageous to knock-out or

reduce the expression of lipid pathway enzymes in order to alter a fatty acid profile. A homologous recombination construct was created to assess whether the expression of an endogenous stearoyl ACP desaturase enzyme can be reduced (or knocked out) and if a corresponding reduction in unsaturated fatty acids can be observed in the lipid profile of the host cell. An approximately 1.5 kb coding sequence of a stearoyl ACP desaturase gene from *Prototheca moriformis* (UTEX 1435) was identified and cloned (SEQ ID NO: 104). The homologous recombination construct was constructed using 0.5 kb of the SAPD coding sequence at the 5' end (5' targeting site), followed by the *Chlamydomonas reinhardtii*  $\beta$ -tubulin promoter driving a codon-optimized yeast sucrose invertase *suc2* gene with the *Chlorella vulgaris* 3'UTR. The rest (~1 kb) of the *Prototheca moriformis* SAPD coding sequence was then inserted after the *C. vulgaris* 3'UTR to make up the 3' targeting site. The sequence for this homologous recombination cassette is listed in SEQ ID NO: 105. As shown above, the success-rate for integration of the homologous recombination cassette into the nuclear genome can be increased by linearizing the cassette before transforming the microalgae, leaving exposed ends. The homologous recombination cassette targeting an endogenous SAPD enzyme in *Prototheca moriformis* is linearized and then transformed into the host cell (*Prototheca moriformis*, UTEX 1435). A successful integration will eliminate the endogenous SAPD enzyme coding region from the host genome via a double reciprocal recombination event, while expression of the newly inserted *suc2* gene will be regulated by the *C. reinhardtii*  $\beta$ -tubulin promoter. The resulting clones can be screened using plates/media containing sucrose as the sole carbon source. Clones containing a successful integration of the homologous recombination cassette will have the ability to grow on sucrose as the sole carbon source and changes in overall saturation of the fatty acids in the lipid profile will serve as a secondary confirmation factor. Additionally, Southern blotting assays using a probe specific for the yeast sucrose invertase *suc2* gene and RT-PCR can also confirm the presence and expression of the invertase gene in positive clones. As an alternative, the same construct without the  $\beta$ -tubulin promoter can be used to excise the endogenous SAPD enzyme coding region. In this case, the newly inserted yeast sucrose invertase *suc2* gene will be regulated by the endogenous SAPD promoter/5'UTR.

### Example 14

#### Fuel Production

##### A. Extraction of Oil from Microalgae Using an Expeller Press and a Press Aid

Microalgal biomass containing 38% oil by DCW was dried using a drum dryer resulting in resulting moisture content of 5-5.5%. The biomass was fed into a French L250 press. 30.4 kg (67 lbs.) of biomass was fed through the press and no oil was recovered. The same dried microbial biomass combined with varying percentage of switchgrass as a press aid was fed through the press. The combination of dried microbial biomass and 20% w/w switchgrass yielded the best overall percentage oil recovery. The pressed cakes were then subjected to hexane extraction and the final yield for the 20% switchgrass condition was 61.6% of the total available oil (calculated by weight). Biomass with above 50% oil dry cell weight did not require the use of a pressing aid such as switchgrass in order to liberate oil.



B. Monosaccharide Composition of Delipidated *Prototheca moriformis* Biomass

*Prototheca moriformis* (UTEX 1435) was grown in conditions and nutrient media (with 4% glucose) as described in Example 45 above. The microalgal biomass was then harvested and dried using a drum dryer. The dried algal biomass was lysed and the oil extracted using an expeller press as described in Example 44 above. The residual oil in the pressed biomass was then solvent extracted using petroleum ether. Residual petroleum ether was evaporated from the delipidated meal using a Rotovapor (Buchi Labortechnik AG, Switzerland). Glycosyl (monosaccharide) composition analysis was then performed on the delipidated meal using combined gas chromatography/mass spectrometry (GC/MS) of the per-O-trimethylsilyl (TMS) derivatives of the monosaccharide methyl glycosides produced from the sample by acidic methanolysis. A sample of delipidated meal was subjected to methanolysis in 1M HCl in methanol at 80° C. for approximately 20 hours, followed by re-N-acetylation with pyridine and acetic anhydride in methanol (for detection of amino sugars). The samples were then per-O-trimethylsilylated by treatment with Tri-Sil (Pierce) at 80° C. for 30 minutes (see methods in Merkle and Poppe (1994) *Methods Enzymol.* 230:1-15 and York et al., (1985) *Methods Enzymol.* 118:3-40). GC/MS analysis of the TMS methyl glycosides was performed on an HP 6890 GC interfaced to a 5975b MSD, using a All Tech EC-1 fused silica capillary column (30m×0.25 mm ID). The monosaccharides were identified by their retention times in comparison to standards, and the carbohydrate character of these are authenticated by their mass spectra. 20 micrograms per sample of inositol was added to the sample before derivatization as an internal standard. The monosaccharide profile of the delipidated *Prototheca moriformis* (UTEX 1435) biomass is summarized in Table 18 below. The total percent carbohydrate from the sample was calculated to be 28.7%.

TABLE 18

Monosaccharide (glycosyl) composition analysis of <i>Prototheca moriformis</i> (UTEX 1435) delipidated biomass.		
	Mass (µg)	Mole % (of total carbohydrate)
Arabinose	0.6	1.2
Xylose	n.d.	n.d.
Galacturonic acid (GalUA)	n.d.	n.d.

TABLE 18-continued

Monosaccharide (glycosyl) composition analysis of <i>Prototheca moriformis</i> (UTEX 1435) delipidated biomass.		
	Mass (µg)	Mole % (of total carbohydrate)
Mannose	6.9	11.9
Galactose	14.5	25.2
Glucose	35.5	61.7
N Acetyl Galactosamine (GalNAc)	n.d.	n.d.
N Acetyl Glucosamine (GlcNAc)	n.d.	n.d.
Heptose	n.d.	n.d.
3 Deoxy-2-manno-2 Octulsonic acid (KDO)	n.d.	n.d.
Sum	57	100

n.d. = none detected

The carbohydrate content and monosaccharide composition of the delipidated meal makes it suitable for use as an animal feed or as part of an animal feed formulation. Thus, in one aspect, the present invention provides delipidated meal having the product content set forth in the table above.

C. Production of Biodiesel from *Prototheca* Oil

Degummed oil from *Prototheca moriformis* UTEX 1435, produced according to the methods described above, was subjected to transesterification to produce fatty acid methyl esters. Results are shown below:

The lipid profile of the oil was:

C10:0	0.02
C12:0	0.06
C14:0	1.81
C14:1	0.07
C16:0	24.53
C16:1	1.22
C18:0	2.34
C18:1	59.21
C18:2	8.91
C18:3	0.28
C20:0	0.23
C20:1	0.10
C20:1	0.08
C21:0	0.02
C22:0	0.06
C24:0	0.10

TABLE 19

Biodiesel profile from <i>Prototheca moriformis</i> triglyceride oil.				
Method	Test	Result	Units	
ASTM D6751 A1	Cold Soak Filterability of Biodiesel Blend Fuels	Filtration Time	120	sec
		Volume Filtered	300	ml
ASTM D93	Pensky-Martens Closed Cup Flash Point	Procedure Used	A	
		Corrected Flash Point	165.0	° C.
ASTM D2709	Water and Sediment in Middle Distillate Fuels (Centrifuge Method)	Sediment and Water	0.000	Vol %
EN 14538	Determination of Ca and Mg Content by ICP OES	Sum of (Ca and Mg)	<1	mg/kg
EN 14538	Determination of Ca and Mg Content by ICP OES	Sum of (Na and K)	<1	mg/kg
ASTM D445	Kinematic/Dynamic Viscosity	Kinematic Viscosity @ 104° F./40° C.	4.873	mm <sup>2</sup> /s
ASTM D874	Sulfated Ash from Lubricating Oils and Additives	Sulfated Ash	<0.005	Wt %
ASTM D5453	Determination of Total Sulfur in Light Hydrocarbons, Spark Ignition Engine Fuel, Diesel Engine Fuel, and Engine Oil by Ultraviolet Fluorescence.	Sulfur, mg/kg	1.7	mg/kg



113

The T10 of the material produced was 242.1° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other T10 values, such as T10 between 180 and 295, between 190 and 270, between 210 and 250, between 225 and 245, and at least 290.

The T90 of the material produced was 300° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein can be employed to generate renewable diesel compositions with other T90 values, such as T90 between 280 and 380, between 290 and 360, between 300 and 350, between 310 and 340, and at least 290.

The FBP of the material produced was 300° C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other FBP values, such as FBP between 290 and 400, between 300 and 385, between 310 and 370, between 315 and 360, and at least 300.

Other oils provided by the methods and compositions of the invention can be subjected to combinations of hydrotreating, isomerization, and other covalent modification including oils with lipid profiles including (a) at least 4% C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

#### Example 15

##### Utilization of Sucrose by *Chlorella luteoviridis*

###### A. SAG 2214 Growth on Glucose and Sucrose

SAG 2214 (designated as *Chlorella luteoviridis*) was tested for growth in the dark on media containing either glucose or sucrose. Heterotrophic liquid cultures were initiated using inoculum from a frozen vial in either media containing 4% glucose or 4% sucrose as the sole carbon source. Cultures were grown in the dark, shaking at 200 rpm. Samples from the cultures were taken at 0, 24, 48 and 72 hour time-points and growth was measured by relative absorbance at 750 nm (UV Mini1240, Shimadzu). SAG 2214 grew equally well on glucose as on sucrose, showing that this microalgae can utilize sucrose as effectively as glucose as a sole carbon source. The result of this experiment is represented graphically in FIG. 3.

###### B. Lipid Productivity and Fatty Acid Profile for SAG 2214

Microalgal strain SAG 2214 was cultivated in liquid medium containing either glucose or sucrose as the sole carbon source in similar conditions as described in Example 32 above. After 7 days, cells were harvested for dry cell weight calculation. Cells were centrifuged and lyophilized for 24 hours. The dried cell pellets were weighed and the dry cell weight per liter was calculated. Cells for lipid analysis were also harvested and centrifuged at 4000×g for 10 minutes at room temperature. The supernatant was discarded and the samples were processed for lipid analysis and fatty acid profile using standard gas chromatography (GC/FID) procedures. The results are summarized below in Tables 21 and 22.

114

TABLE 21

Lipid productivity and DCW for SAG 2214.			
Sample	Lipid (g/L)	DCW (g/L)	% Lipid DCW
SAG 2214 glucose	2.43	5.73	42.44%
SAG 2214 sucrose	0.91	2.00	45.56%

TABLE 22

Fatty acid profile for SAG 2214.	
Fatty Acid	Percent (w/w)
C:16:0	21
C:18:1	38
C:18:2	41

##### C. Genomic Comparison of SAG 2214 to Other *Chlorella luteoviridis* Strains

Microalgal strain SAG 2214 proved to be of general interest due to its ability to grow on sucrose as a carbon source (illustrated above). In addition to the growth characteristics of this strain, its taxonomic relationship to other microalgal species was also of interest. Designated by the SAG collection as a *Chlorella luteoviridis* strain, the 23s rRNA gene of SAG 2214 was sequenced and compared to the 23s rRNA genomic sequence of nine other strains also identified by the SAG and UTEX collections as *Chlorella luteoviridis*. These strains were UTEX 21, 22, 28, 257 and 258, and SAG strains 2133, 2196, 2198 and 2203. The DNA genotyping methods used were the same as the methods described above in Example 1. Sequence alignments and unrooted trees were generated using Geneious DNA analysis software. Out of the nine other strains that were genotypes, UTEX 21, 22, 28 and 257 had identical 23s rRNA DNA sequence (SEQ ID NO: 106). The other five *Chlorella luteoviridis* strains had 23s rRNA sequences that were highly homologous to UTEX 21, 22, 28, and 257.

The 23s rRNA gene sequence from SAG 2214 (SEQ ID NO: 30) is decidedly different from that of the other nine *C. luteoviridis* strains, having a large insertion that was not found in the other strains. Further analysis of this 23s rRNA gene sequence using BLAST indicated that it shared the greatest homology with members of the genus *Leptosira* and *Trebouxia* (members of phycobiont portion of lichens). These results indicate that SAG 2214 may not be *Chlorella luteoviridis* strain as categorized by the strain collection, but instead shares significant 23S rRNA nucleotide identity to algal symbionts found in lichen. The genomic analysis along with the growth characteristics indicate that SAG 2214 may be a source for genes and proteins involved in the metabolism of sucrose, as well as signaling and transit peptides responsible for the correct localization of such enzymes. SAG 2214 and other strains with a high degree of genomic similarity may also be strains useful for oil production using sucrose as a source of fixed carbon.

Although this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth.

All references cited herein, including patents, patent applications, and publications, are hereby incorporated by reference in their entireties, whether previously specifically incor-

porated or not. The publications mentioned herein are cited for the purpose of describing and disclosing reagents, methodologies and concepts that may be used in connection with the present invention. Nothing herein is to be construed as an admission that these references are prior art in relation to the inventions described herein. In particular, the following patent applications are hereby incorporated by reference in their entireties for all purposes: U.S. Provisional Application No. 60/941,581, filed Jun. 1, 2007, entitled "Production of Hydrocarbons in Microorganisms"; U.S. Provisional Application No. 60/959,174, filed Jul. 10, 2007, entitled "Production of Hydrocarbons in Microorganisms"; U.S. Provisional Application No. 60/968,291, filed Aug. 27, 2007, entitled "Production of Hydrocarbons in Microorganisms"; U.S. Provisional Application No. 61/024,069, filed Jan. 28, 2008, entitled "Production of Hydrocarbons in Microorganisms"; PCT Application No. PCT/US08/65563, filed Jun. 2, 2008, entitled "Production of Oil in Microorganisms"; U.S. patent application Ser. No. 12/131,783, filed Jun. 2, 2008, entitled "Use of Cellulosic Material for Cultivation of Microorganisms"; U.S. patent application Ser. No. 12/131,773, filed Jun. 2, 2008, entitled "Renewable Diesel and Jet Fuel from Microbial Sources"; U.S. patent application Ser. No. 12/131,793, filed Jun. 2, 2008, entitled "Sucrose Feedstock Utilization for

Oil-Based Fuel Manufacturing"; U.S. patent application Ser. No. 12/131,766, filed Jun. 2, 2008, entitled "Glycerol Feedstock Utilization for Oil-Based Fuel Manufacturing"; U.S. patent application Ser. No. 12/131,804, filed Jun. 2, 2008, entitled "Lipid Pathway Modification in Oil-Bearing Microorganisms"; U.S. Patent Application No. 61/118,590, filed Nov. 28, 2008, entitled "Production of Oil in Microorganisms"; U.S. Provisional Patent Application No. 61/118,994, filed Dec. 1, 2008, entitled "Production of Oil in Microorganisms"; U.S. Provisional Patent Application No. 61/174,357, filed Apr. 3, 2009, entitled "Production of Oil in Microorganisms"; U.S. Provisional Patent Application No. 61/219,525, filed Jun. 23, 2009, entitled "Production of Oil in Microorganisms"; U.S. patent application Ser. No. 12/628,140, filed Nov. 30, 2009, entitled "Novel Triglyceride and Fuel Compositions"; U.S. patent application Ser. No. 12/628,144, filed Nov. 30, 2009, entitled "Cellulosic Cultivation of Oleaginous Microorganisms"; U.S. patent application Ser. No. 12/628,147, filed Nov. 30, 2009, entitled "Nucleic Acids Useful in the Manufacture of Oil"; U.S. patent application Ser. No. 12/628,149, filed Nov. 30, 2009, entitled, "Renewable Chemical Production from Novel Fatty Acid Feedstocks", and U.S. patent application Ser. No. 12/628,120, filed Nov. 30, 2009, entitled "Recombinant Microalgae Cells Producing Novel Oils".

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 143

<210> SEQ ID NO 1

<211> LENGTH: 1187

<212> TYPE: DNA

<213> ORGANISM: *Chlorella* sp.

<400> SEQUENCE: 1

```

gatcagacgg gcctgacctg cgagataatc aagtgtctgt aggcaaccaa ctcagcagct      60
gcttgggtgtt ggggtctgag gatagtgttg cagggcccca aggacagcag gggaaacttac      120
accttgtccc cgaccaggtt ttatggagtg cattgcctca agagcctagc cggagcgcta      180
ggctacatac ttgccgcacc ggtatgaggg gatatagtac tcgcactgcg ctgtctagtg      240
agatgggcag tgctgcccatt aaacaactgg ctgctcagcc atttgttggc ggaccattct      300
gggggggcca gcaatgcctg actttcgggt agggtgaaaa ctgaacaaaag actacaaaaa      360
cagaatttct tcctccttgg aggtaagcgc aggccggccc gctcgcgccc acatggcgct      420
ccgaacacct ccatagctgt aagggcgcaa acatggccgg actggtgtca gcactctttc      480
atggccatac aaggtcatgt cgagattagt gctgagtaag acactatcac cccatgttcg      540
attgaagcgg tgacttcatg ccaacctgcc cctgggcgta gcagacgtat gccatcatga      600
ccactagccg acatgcgctg tcttttgcca caaaaacaac tggtaacacg ctcgaagtgc      660
tgccgcacac ctccgggagt gagtcggcg actcctcccc ggccggccgc ggcctacct      720
gggtagggtc gccatacgc cagcaccaaa cgacgcagga ggggattggg gtagggaatc      780
ccaaccagcc taaccaagac ggcacctata ataataggtg gggggactaa cagccctata      840
tcgcaagctt tgggtgccta tcttgagaag cagagtttg agtggtctgt tacggtcgac      900
cctaaggctg gtgtgcccga gcctgaaaaca aagcgtctag cagctgcttc tataatgtgt      960
cagccgttgt gtttcagtta tattgtatgc tattgtttgt tcgtgctagg gtggcgcagg      1020
cccacctact gtggggggcc attggttggg gcttgaattg cctcaccatc taaggtctga      1080
acgctcactc aaacgccttt gtacaactgc agaactttcc ttggcgctgc aactacagtg      1140

```

-continued

---

 tgcaaaccag cacatagcac tcccttaccat caccagcag tacaaca 1187

<210> SEQ ID NO 2  
 <211> LENGTH: 1414  
 <212> TYPE: DNA  
 <213> ORGANISM: Chlorella ellipsoidea

<400> SEQUENCE: 2

```

cgctgcgcac cagggccgcc agctcgctga tgtcgctcca aatgcgggtcc cccgattttt    60
tgttcttcat cttctccacc ttggtggcct tcttgccag gcccttcagc tgcattgcga    120
cagaccgttg agctcctgat cagcatcctc aggaggcctt ttgacaagca agccccctgtg    180
caagcccatt cacggggtag cagtgggtgt gaggtagatg ggtttgaaaa ggattgctcg    240
gtcgattgct gctcatggaa ttggcatgtg catgcatggt cacaatatgc caccaggcct    300
tggagcaaga gagcatgaat gccttcaggc aggttgaag ttctggggg tgaagaggca    360
gggcccagga ttggaggagg aaagcatcaa gtcgctcgtc atgctcatgt ttcagtcag    420
agtttgccaa gctcacagga gcagagacaa gactggctgc tcagggttg catcgtgtgt    480
gtggtggggg ggggggggtt aatacggtag gaaatgact tggaaatccc acctcatgcc    540
agcggacca catgcttgaa ttcgaggcct gtggggtag aatgctcac tctgcctcg    600
ttgctgaggt acttcaggcc gctgagctca aagtcgatgc cctgctcgtc tatcagggcc    660
tgcacctctg ggctgaccgg ctcagcctcc ttcgagggca tggagtaggc gccggcagcg    720
ttcatgtccg ggcccagggc agcgggtgtg ccataaatgt cggtagatgt ggggaggggg    780
gccgtcgcca caccattgcc gttgctggct gacgcatgca catgtggcct ggctggcacc    840
ggcagcactg gtctccagcc agccagcaag ttgctgttca ggaaagcggc catgttgttg    900
gtcctcgctc atgtaattcc ccagatcaaa ggagggaaca gcttgattt gatgtagtgc    960
ccaaccggac tgaatgtgag atggcaggtc ccttgagtc tcccgaatta ctgacagggc   1020
actgtgacct aacgcagcat gccaaaccgca aaaaaatgat tgacagaaaa tgaagcggtg   1080
tgtaaatatt tgctgtattt atctgtttta atcagcaacc aagttcgaaa cgcaactatc   1140
gtggtgatca agtgaacctc atcagactta cctcgttcgg caaggaaacg gaggcaccaa   1200
attccaattt gatattatcg cttgccaagc tagagctgat ctttgggaaa ccaactgcca   1260
gacagtggac tgtatggag tgcctcagc ggtggagcct cttcgattcg gttagtcatt   1320
actaacgtga accctcagtg aagggacat cagaccagaa agaccagatc tcctcctcga   1380
caccgagaga gtgttcgggc agtaggacga caag                               1414
  
```

<210> SEQ ID NO 3  
 <211> LENGTH: 512  
 <212> TYPE: PRT  
 <213> ORGANISM: Unknown  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Unknown: Yeast sequence

<400> SEQUENCE: 3

```

Met Thr Asn Glu Thr Ser Asp Arg Pro Leu Val His Phe Thr Pro Asn
 1                               5 10 15
Lys Gly Trp Met Asn Asp Pro Asn Gly Leu Trp Tyr Asp Glu Lys Asp
 20 25 30
Ala Lys Trp His Leu Tyr Phe Gln Tyr Asn Pro Asn Asp Thr Val Trp
 35 40 45
Gly Thr Pro Leu Phe Trp Gly His Ala Thr Ser Asp Asp Leu Thr Asn
 50 55 60
  
```

-continued

Trp	Glu	Asp	Gln	Pro	Ile	Ala	Ile	Ala	Pro	Lys	Arg	Asn	Asp	Ser	Gly	65	70	75	80
Ala	Phe	Ser	Gly	Ser	Met	Val	Val	Asp	Tyr	Asn	Asn	Thr	Ser	Gly	Phe	85	90	95	
Phe	Asn	Asp	Thr	Ile	Asp	Pro	Arg	Gln	Arg	Cys	Val	Ala	Ile	Trp	Thr	100	105	110	
Tyr	Asn	Thr	Pro	Glu	Ser	Glu	Glu	Gln	Tyr	Ile	Ser	Tyr	Ser	Leu	Asp	115	120	125	
Gly	Gly	Tyr	Thr	Phe	Thr	Glu	Tyr	Gln	Lys	Asn	Pro	Val	Leu	Ala	Ala	130	135	140	
Asn	Ser	Thr	Gln	Phe	Arg	Asp	Pro	Lys	Val	Phe	Trp	Tyr	Glu	Pro	Ser	145	150	155	160
Gln	Lys	Trp	Ile	Met	Thr	Ala	Ala	Lys	Ser	Gln	Asp	Tyr	Lys	Ile	Glu	165	170	175	
Ile	Tyr	Ser	Ser	Asp	Asp	Leu	Lys	Ser	Trp	Lys	Leu	Glu	Ser	Ala	Phe	180	185	190	
Ala	Asn	Glu	Gly	Phe	Leu	Gly	Tyr	Gln	Tyr	Glu	Cys	Pro	Gly	Leu	Ile	195	200	205	
Glu	Val	Pro	Thr	Glu	Gln	Asp	Pro	Ser	Lys	Ser	Tyr	Trp	Val	Met	Phe	210	215	220	
Ile	Ser	Ile	Asn	Pro	Gly	Ala	Pro	Ala	Gly	Gly	Ser	Phe	Asn	Gln	Tyr	225	230	235	240
Phe	Val	Gly	Ser	Phe	Asn	Gly	Thr	His	Phe	Glu	Ala	Phe	Asp	Asn	Gln	245	250	255	
Ser	Arg	Val	Val	Asp	Phe	Gly	Lys	Asp	Tyr	Tyr	Ala	Leu	Gln	Thr	Phe	260	265	270	
Phe	Asn	Thr	Asp	Pro	Thr	Tyr	Gly	Ser	Ala	Leu	Gly	Ile	Ala	Trp	Ala	275	280	285	
Ser	Asn	Trp	Glu	Tyr	Ser	Ala	Phe	Val	Pro	Thr	Asn	Pro	Trp	Arg	Ser	290	295	300	
Ser	Met	Ser	Leu	Val	Arg	Lys	Phe	Ser	Leu	Asn	Thr	Glu	Tyr	Gln	Ala	305	310	315	320
Asn	Pro	Glu	Thr	Glu	Leu	Ile	Asn	Leu	Lys	Ala	Glu	Pro	Ile	Leu	Asn	325	330	335	
Ile	Ser	Asn	Ala	Gly	Pro	Trp	Ser	Arg	Phe	Ala	Thr	Asn	Thr	Thr	Leu	340	345	350	
Thr	Lys	Ala	Asn	Ser	Tyr	Asn	Val	Asp	Leu	Ser	Asn	Ser	Thr	Gly	Thr	355	360	365	
Leu	Glu	Phe	Glu	Leu	Val	Tyr	Ala	Val	Asn	Thr	Thr	Gln	Thr	Ile	Ser	370	375	380	
Lys	Ser	Val	Phe	Ala	Asp	Leu	Ser	Leu	Trp	Phe	Lys	Gly	Leu	Glu	Asp	385	390	395	400
Pro	Glu	Glu	Tyr	Leu	Arg	Met	Gly	Phe	Glu	Val	Ser	Ala	Ser	Ser	Phe	405	410	415	
Phe	Leu	Asp	Arg	Gly	Asn	Ser	Lys	Val	Lys	Phe	Val	Lys	Glu	Asn	Pro	420	425	430	
Tyr	Phe	Thr	Asn	Arg	Met	Ser	Val	Asn	Asn	Gln	Pro	Phe	Lys	Ser	Glu	435	440	445	
Asn	Asp	Leu	Ser	Tyr	Tyr	Lys	Val	Tyr	Gly	Leu	Leu	Asp	Gln	Asn	Ile	450	455	460	
Leu	Glu	Leu	Tyr	Phe	Asn	Asp	Gly	Asp	Val	Val	Ser	Thr	Asn	Thr	Tyr	465	470	475	480
Phe	Met	Thr	Thr	Gly	Asn	Ala	Leu	Gly	Ser	Val	Asn	Met	Thr	Thr	Gly	485	490	495	

-continued

---

Val Asp Asn Leu Phe Tyr Ile Asp Lys Phe Gln Val Arg Glu Val Lys  
500 505 510

<210> SEQ ID NO 4  
<211> LENGTH: 20  
<212> TYPE: PRT  
<213> ORGANISM: Unknown  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Unknown: Yeast sequence

<400> SEQUENCE: 4

Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys  
1 5 10 15

Ile Ser Ala Ser  
20

<210> SEQ ID NO 5  
<211> LENGTH: 18  
<212> TYPE: PRT  
<213> ORGANISM: Unknown  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Unknown: Higher plant secretion  
signal

<400> SEQUENCE: 5

Met Ala Asn Lys Ser Leu Leu Leu Leu Leu Leu Gly Ser Leu Ala  
1 5 10 15

Ser Gly

<210> SEQ ID NO 6  
<211> LENGTH: 10  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
consensus sequence

<400> SEQUENCE: 6

Met Ala Arg Leu Pro Leu Ala Ala Leu Gly  
1 5 10

<210> SEQ ID NO 7  
<211> LENGTH: 20  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
peptide

<400> SEQUENCE: 7

Met Ala Asn Lys Leu Leu Leu Leu Leu Leu Leu Leu Pro Leu  
1 5 10 15

Ala Ala Ser Gly  
20

<210> SEQ ID NO 8  
<211> LENGTH: 2615  
<212> TYPE: DNA  
<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 8

gaattcccca acatggtgga gcaagacact ctcgtctact ccaagaatat caaagataca 60  
gtctcagaag accaaagggc tattgagact tttcaacaaa gggtaatatc gggaaacctc 120  
ctcggattcc attgcccagc tatctgtcac ttcatacaaaa ggacagtaga aaaggaaggt 180

-continued

---

ggcacctaca aatgccatca ttgcgataaa ggaaaggcta tcgttcaaga tgcctctgcc	240
gacagtggtc ccaaagatgg acccccaccc acgaggagca tcgtggaaaa agaagacgtt	300
ccaaccacgt cttcaaagca agtggattga tgtgaacatg gtggagcacg acactctcgt	360
ctactccaag aatatcaaag atacagtctc agaagaccaa agggctattg agacttttca	420
acaaagggta atatcgggaa acctcctcgg attccattgc ccagctatct gtcacttcat	480
caaaaggaca gtagaaaagg aaggtggcac ctacaaatgc catcattgcg ataaaggaaa	540
ggctatcgtt caagatgcct ctgccgacag tgggtccaaa gatggacccc caccacgag	600
gagcatcgtg gaaaaagaag acgttccaac cacgtcttca aagcaagtgg attgatgtga	660
tatctcact gacgtaagg atgacgcaca atcccactat ccttcgcaag accttctctc	720
tatataagga agttcatttc atttggagag gacacgctga aatcaccagt ctctctctac	780
aaatctatct ctggcgcgcc atataaatgc ttcttcaggc ctttctttt cttcttgctg	840
gttttgetgc caagatcagc gcctctatga cgaacgaaac ctccgataga ccacttgctg	900
actttacacc aaacaagggc tggatgaatg accccaatgg actgtggtac gacgaaaaag	960
atgccaagtg gcatctgtac tttcaataca acccgaacga tactgtctgg gggacgccat	1020
tgttttggg ccacgccacg tccgacgacc tgaccaattg ggaggaccaa ccaatagcta	1080
tcgctccgaa gaggaacgac tccggagcat tctcgggttc catggtggtt gactacaaca	1140
atacttccgg cttttcaac gataccattg acccgagaca acgctgcgtg gccatatgga	1200
cttacaacac accggagtcc gaggagcagt acatctcgta tagcctggac ggtggatata	1260
cttttacaga gtatcagaag aacctgtgc ttgctgcaaa ttcgactcag ttcgagatc	1320
cgaaggtcct ttggtacgag cctcgcaga agtggatcat gacagcggca aagtcacagg	1380
actacaagat cgaatattac tcgtctgacg acctaaatc ctggaagctc gaatccgct	1440
tcgaaaacga gggctttctc ggctaccaat acgaatgcc aggcctgata gaggtcccaa	1500
cagagcaaga tcccagcaag tctactggg tgatgtttat ttccattaat ccaggagcac	1560
cggcaggagg ttcttttaat cagtacttcg tcggaagctt taacggaact catttcgagg	1620
catttgataa ccaatcaaga gtagtgtatt ttggaaagga ctactatgcc ctgcagactt	1680
tcttcaatac tgaccgacc tatgggagcg ctcttgcat tcgctgggct tctaactggg	1740
agtattccgc attcgttctc acaaaccctt ggaggctctc catgtcgtc gtgaggaaat	1800
tctctctcaa cactgagtac caggccaacc cggaaaccga actcataaac ctgaaagccg	1860
aaccgatcct gaacattagc aacgtggcc cctggagccg gtttgcaacc aacaccacgt	1920
tgacgaaagc caacagctac aacgtcagc tttcgaatag caccggtaca cttgaatttg	1980
aactggtgta tgcctcaat accacccaaa cgatctcgaa gtcggtgttc gcgacctct	2040
ccctctgggt taaagccctg gaagaccctg aggagtacct cagaatgggt ttcgaggtt	2100
ctcgtcctc cttcttctt gatcgcggga acagcaaagt aaaatttgtt aaggagaacc	2160
catattttac caacaggatg agcgttaaca accaaccatt caagagcgaa aacgacctgt	2220
cgtaactaca agtgtatggt ttgcttgatc aaaatatct ggaactctac ttcaacgatg	2280
gtgatgtcgt gtccaccaac acataactca tgacaaccgg gaacgcactg ggctccgtga	2340
acatgacgac ggggtgtgat aacctgtct acatcgacaa attccaggtg agggaagtca	2400
agtgagatct gtcgatcgac aagctcagat ttctccataa taatgtgtga gtagttccca	2460
gataagggaa ttagggttcc tatagggttt cgctcatgtg ttgagcatat aagaaacct	2520
tagtatgtat ttgtatttgt aaaatacttc tatcaataaa atttctaatt cctaaaacca	2580



-continued

---

```
aaatccagta ctaaaatcca gatccccga attaa 2615
```

```
<210> SEQ ID NO 9
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
```

```
<400> SEQUENCE: 9
```

```
tgttgaagaa tgagccggcg ac 22
```

```
<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer
```

```
<400> SEQUENCE: 10
```

```
cagtgagcta ttacgcactc 20
```

```
<210> SEQ ID NO 11
<211> LENGTH: 541
<212> TYPE: DNA
<213> ORGANISM: Prototheca kruegani
```

```
<400> SEQUENCE: 11
```

```
tgttgaagaa tgagccggcg agttaaaaag agtggcatgg ttaagaaaa tactctggag 60
```

```
ccatagcgaa agcaagtta gtaagcttag gtcattcttt ttagaccga aaccgagtga 120
```

```
tctaccatg atcaggggta agtgtagta aaataacatg gagggccgaa ccgactaatg 180
```

```
ttgaaaatt agcggatgaa ttgtggtag gggcgaaaa ccaatcgaac tcggagttag 240
```

```
ctggttctcc ccgaaatgcg tttagggcga gcagtagcag tacaataga ggggtaaagc 300
```

```
actgtttctt ttgtgggctt cgaaagtgt acctcaaagt ggcaaactct gaatactcta 360
```

```
tttagatata tactagttag accttggggg ataagctcct tggtaaaaag ggaacagcc 420
```

```
cagatcacca gttaaggccc caaaatgaaa atgatagtag ctaaggatgt gggtagtca 480
```

```
aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540
```

```
g 541
```

```
<210> SEQ ID NO 12
<211> LENGTH: 573
<212> TYPE: DNA
<213> ORGANISM: Prototheca wickerhamii
```

```
<400> SEQUENCE: 12
```

```
tgttgaagaa tgagccggcg acttaaaata aatggcaggc taagagattt aataactcga 60
```

```
aaactaagcg aaagcaagtc ttaatagggc gtcaatttaa caaaacttta aataaattat 120
```

```
aaagtcatat atttttagacc cgaacctgag tgatctaacc atggtcagga tgaacttg 180
```

```
gtgacaccaa gtggaagtcc gaaccgaccg atggtgaaaa atcggcggat gaactgtggt 240
```

```
tagtggtgaa ataccagtcg aactcagagc tagctggttc tccccgaaat gcgttgaggc 300
```

```
gcagcaatat atctcgtcta tctaggggta aagcactggt tcggtgcggg ctatgaaaat 360
```

```
ggtagcaaat cgtggcaaac tctgaatact agaaatgacg atatattagt gagactatgg 420
```

```
gggataagct ccatagtcga gagggaaaca gccagacca ccagttaagg ccccaaatg 480
```

-continued

ataatgaagt ggtaaaggag gtgaaaatgc aaatacaacc aggaggttgg cttagaagca 540  
gccatccttt aaagagtgcg taatagctca ctg 573

<210> SEQ ID NO 13  
<211> LENGTH: 541  
<212> TYPE: DNA  
<213> ORGANISM: *Prototheca stagnora*

<400> SEQUENCE: 13

tgttgaagaa tgagccggcg agttaaaaa aatggcatgg ttaagatat ttctctgaag 60  
ccatagcgaa agcaagtttt acaagctata gtcatttttt ttagaccgga aaccgagtga 120  
tctaccatg atcaggggtga agtgttggtc aaataacatg gaggcccgaa ccgactaatg 180  
gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240  
ctggttctcc ccgaaatgcg tttagggcga gcagtagcaa cacaaataga ggggtaaagc 300  
actgtttctt ttgtgggctt cgaaagtgt acctcaaagt ggcaaactct gaatactcta 360  
tttagatata tactagttag accttggggg ataagctcct tggtaaaaag ggaacagcc 420  
cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggacgt gagtatgtca 480  
aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540  
g 541

<210> SEQ ID NO 14  
<211> LENGTH: 541  
<212> TYPE: DNA  
<213> ORGANISM: *Prototheca moriformis*

<400> SEQUENCE: 14

tgttgaagaa tgagccggcg agttaaaaag agtggcatgg ttaagataa ttctctggag 60  
ccatagcgaa agcaagttta acaagctaaa gtcacccttt ttagaccgga aaccgagtga 120  
tctaccatg atcaggggtga agtgttggtc aaataacatg gaggcccgaa ccgactaatg 180  
gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240  
ctggttctcc ccgaaatgcg tttagggcga gcagtagcaa cacaaataga ggggtaaagc 300  
actgtttctt ttgtgggctt cgaaagtgt acctcaaagt ggcaaactct gaatactcta 360  
tttagatata tactagttag accttggggg ataagctcct tggtaaaaag ggaacagcc 420  
cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggatgt gggtatgtta 480  
aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540  
g 541

<210> SEQ ID NO 15  
<211> LENGTH: 573  
<212> TYPE: DNA  
<213> ORGANISM: *Prototheca moriformis*

<400> SEQUENCE: 15

tgttgaagaa tgagccggcg acttaaaata aatggcaggc taagagaatt aataactcga 60  
aacctaagcg aaagcaagtc ttaatagggc gctaatttaa caaacatta aataaaatct 120  
aaagtcattt attttagacc cgaacctgag tgatctaacc atggtcagga tgaacttgg 180  
gtgacaccaa gtggaagtcc gaaccgaccg atgttgaaaa atcggcggat gaactgtggt 240  
tagtggtgaa ataccagtcg aactcagagc tagctggttc tccccgaaat gcgttgagcc 300  
gcagcaatat atctcgtcta tctaggggta aagcactggt tcggtgcggg ctatgaaaat 360

-continued

---

```

ggtaccaa at cgtggcaaac tctgaatact agaaatgacg atatattagt gagactatgg 420
gggataagct ccatagtcga gagggaaaca gcccagacca ccagttaagg ccccaaatg 480
ataatgaagt ggtaaaggag gtgaaaatgc aaatacaacc aggaggttgg cttagaagca 540
gccatccttt aaagagtgcg taatagctca ctg 573

```

```

<210> SEQ ID NO 16
<211> LENGTH: 573
<212> TYPE: DNA
<213> ORGANISM: Prototheca wickerhamii

```

```

<400> SEQUENCE: 16

```

```

tgttgaagaa tgagccgtcg acttaaaata aatggcaggc taagagaatt aataactcga 60
aacctaagcg aaagcaagtc ttaatagggc gctaatttaa caaacatta aataaaatct 120
aaagtcattt attttagacc cgaacctgag tgatctaacc atggtcagga tgaacttgg 180
gtgacaccaa gtggaagtcc gaaccgacg atgttgaaaa atcggcggat gaactgtggt 240
tagtggtgaa ataccagtcg aactcagagc tagctggttc tccccgaaat gcgttgaggc 300
gcagcaatat atctcgtcta tctaggggta aagcactggt tcggtgcggg ctatgaaat 360
ggtaccaa at cgtggcaaac tctgaatact agaaatgacg atatattagt gagactatgg 420
gggataagct ccatagtcga gagggaaaca gcccagacca ccagttaagg ccccaaatg 480
ataatgaagt ggtaaaggag gtgaaaatgc aaatacaacc aggaggttgg cttagaagca 540
gccatccttt aaagagtgcg taatagctca ctg 573

```

```

<210> SEQ ID NO 17
<211> LENGTH: 541
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 17

```

```

tgttgaagaa tgagccggcg agttaaaaag agtggcgtgg ttaaagaaaa ttctctggaa 60
ccatagcgaa agcaagttta acaagcttaa gtcacttttt ttagaccoga aaccgagtga 120
tctaccatg atcaggggtga agtgttggtg aaataacatg gaggccgaa ccgactaatg 180
gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240
ctggttctcc ccgaaatgcg tttagggcga gcagtagcaa cacaaataga ggggtaaagc 300
actgtttctt ttgtgggctc cgaaagttgt acctcaaagt ggcaaactct gaatactcta 360
ttgatatac tactagttag accttggggg ataagctcct tggtcgaaag ggaacagcc 420
cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggatgt gagtatgtca 480
aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540
g 541

```

```

<210> SEQ ID NO 18
<211> LENGTH: 541
<212> TYPE: DNA
<213> ORGANISM: Prototheca zopfii

```

```

<400> SEQUENCE: 18

```

```

tgttgaagaa tgagccggcg agttaaaaag agtggcatgg ttaaagaaaa ttctctggag 60
ccatagcgaa agcaagttta acaagcttaa gtcacttttt ttagaccoga aaccgagtga 120
tctaccatg atcaggggtga agtgttggtg aaataacatg gaggccgaa ccgactaatg 180
gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240

```

-continued

---

```

ctggttctcc ccgaaatgcg tttagggcga gcagtagcaa cacaaataga ggggtaaagc 300
actgtttctt tcgtgggctt cgaaagtgt acctcaaagt ggcaaactct gaatactcta 360
tttagatate tactagttag accttggggg ataagctect tggtcaaaag ggaacagcc 420
cagatcacca gtttaaggccc caaaatgaaa atgatagtga ctaaggatgt gagtatgtca 480
aaacctccag caggtttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540
g 541

```

```

<210> SEQ ID NO 19
<211> LENGTH: 565
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 19

```

```

tgttgaagaa tgagccggcg acttagaaaa ggtggcatgg ttaaggaat attccgaagc 60
cgtagcaaaa gcgagtctga atagggcgat aaaatatatt aatatttaga atctagtcat 120
ttttctaga cccgaaccgg ggtgatctaa ccatgaccag gatgaagctt gggtgatacc 180
aagtgaaggt ccgaaccgac cgatgttgaa aaatcggcgg atgagttgtg gttagcggtg 240
aaataccagt cgaaccggga gctagctggt tctccccgaa atgcggtgag ggcagcagc 300
acatctagtc tatctagggg taaagcactg ttctcggcgg ggctgtgaga acggtaccaa 360
atcgtggcaa actctgaata ctagaatga cgatgtagta gtgagactgt gggggataag 420
ctccattgtc aagagggaaa cagcccagac caccagctaa ggccccaaaa tggtaatgta 480
gtgaaaagg aggtgaaat gcaaatacaa ccaggaggtt ggcttagaag cagccatcct 540
ttaaagagtg cgtaatagct cactg 565

```

```

<210> SEQ ID NO 20
<211> LENGTH: 550
<212> TYPE: PRT
<213> ORGANISM: Cichorium intybus

```

```

<400> SEQUENCE: 20

```

```

Met Ser Asn Ser Ser Asn Ala Ser Glu Ser Leu Phe Pro Ala Thr Ser
1          5          10          15
Glu Gln Pro Tyr Arg Thr Ala Phe His Phe Gln Pro Pro Gln Asn Trp
20          25          30
Met Asn Asp Pro Asn Gly Pro Met Cys Tyr Asn Gly Val Tyr His Leu
35          40          45
Phe Tyr Gln Tyr Asn Pro Phe Gly Pro Leu Trp Asn Leu Arg Met Tyr
50          55          60
Trp Ala His Ser Val Ser His Asp Leu Ile Asn Trp Ile His Leu Asp
65          70          75          80
Leu Ala Phe Ala Pro Thr Glu Pro Phe Asp Ile Asn Gly Cys Leu Ser
85          90          95
Gly Ser Ala Thr Val Leu Pro Gly Asn Lys Pro Ile Met Leu Tyr Thr
100         105         110
Gly Ile Asp Thr Glu Asn Arg Gln Val Gln Asn Leu Ala Val Pro Lys
115         120         125
Asp Leu Ser Asp Pro Tyr Leu Arg Glu Trp Val Lys His Thr Gly Asn
130         135         140
Pro Ile Ile Ser Leu Pro Glu Glu Ile Gln Pro Asp Asp Phe Arg Asp
145         150         155         160
Pro Thr Thr Thr Trp Leu Glu Glu Asp Gly Thr Trp Arg Leu Leu Val

```

-continued

	165		170		175
Gly Ser Gln Lys Asp Lys Thr Gly Ile Ala Phe Leu Tyr His Ser Gly	180		185		190
Asp Phe Val Asn Trp Thr Lys Ser Asp Ser Pro Leu His Lys Val Ser	195		200		205
Gly Thr Gly Met Trp Glu Cys Val Asp Phe Phe Pro Val Trp Val Asp	210		215		220
Ser Thr Asn Gly Val Asp Thr Ser Ile Ile Asn Pro Ser Asn Arg Val	225	230		235	240
Lys His Val Leu Lys Leu Gly Ile Gln Asp His Gly Lys Asp Cys Tyr	245		250		255
Leu Ile Gly Lys Tyr Ser Ala Asp Lys Glu Asn Tyr Val Pro Glu Asp	260		265		270
Glu Leu Thr Leu Ser Thr Leu Arg Leu Asp Tyr Gly Met Tyr Tyr Ala	275		280		285
Ser Lys Ser Phe Phe Asp Pro Val Lys Asn Arg Arg Ile Met Thr Ala	290	295		300	
Trp Val Asn Glu Ser Asp Ser Glu Ala Asp Val Ile Ala Arg Gly Trp	305	310		315	320
Ser Gly Val Gln Ser Phe Pro Arg Ser Leu Trp Leu Asp Lys Asn Gln	325		330		335
Lys Gln Leu Leu Gln Trp Pro Ile Glu Glu Ile Glu Met Leu His Gln	340		345		350
Asn Glu Val Ser Phe His Asn Lys Lys Leu Asp Gly Gly Ser Ser Leu	355		360		365
Glu Val Leu Gly Ile Thr Ala Ser Gln Ala Asp Val Lys Ile Ser Phe	370		375		380
Lys Leu Ala Asn Leu Glu Glu Ala Glu Glu Leu Asp Pro Ser Trp Val	385	390		395	400
Asp Pro Gln Leu Ile Cys Ser Glu Asn Asp Ala Ser Lys Lys Gly Lys	405		410		415
Phe Gly Pro Phe Gly Leu Leu Ala Leu Ala Ser Ser Asp Leu Arg Glu	420		425		430
Gln Thr Ala Ile Phe Phe Arg Val Phe Arg Lys Asn Gly Arg Tyr Val	435		440		445
Val Leu Met Cys Ser Asp Gln Ser Arg Ser Ser Met Lys Asn Gly Ile	450		455		460
Glu Lys Arg Thr Tyr Gly Ala Phe Val Asp Ile Asp Pro Gln Gln Asp	465	470		475	480
Glu Ile Ser Leu Arg Thr Leu Ile Asp His Ser Ile Val Glu Ser Phe	485		490		495
Gly Gly Arg Gly Lys Thr Cys Ile Thr Thr Arg Val Tyr Pro Thr Leu	500		505		510
Ala Ile Gly Glu Gln Ala Arg Leu Phe Ala Phe Asn His Gly Thr Glu	515		520		525
Ser Val Glu Ile Ser Glu Leu Ser Ala Trp Ser Met Lys Lys Ala Gln	530		535		540
Met Lys Val Glu Glu Pro	545		550		

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 581

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Schizosaccharomyces pombe

-continued

&lt;400&gt; SEQUENCE: 21

Met Phe Leu Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu  
 1 5 10 15  
 Leu Ser Ser Thr Asn Ala Ala Pro Arg His Leu Tyr Val Lys Arg Tyr  
 20 25 30  
 Pro Val Ile Tyr Asn Ala Ser Asn Ile Thr Glu Val Ser Asn Ser Thr  
 35 40 45  
 Thr Val Pro Pro Pro Phe Val Asn Thr Thr Ala Pro Asn Gly Thr  
 50 55 60  
 Cys Leu Gly Asn Tyr Asn Glu Tyr Leu Pro Ser Gly Tyr Tyr Asn Ala  
 65 70 75 80  
 Thr Asp Arg Pro Lys Ile His Phe Thr Pro Ser Ser Gly Phe Met Asn  
 85 90 95  
 Asp Pro Asn Gly Leu Val Tyr Thr Gly Gly Val Tyr His Met Phe Phe  
 100 105 110  
 Gln Tyr Ser Pro Lys Thr Leu Thr Ala Gly Glu Val His Trp Gly His  
 115 120 125  
 Thr Val Ser Lys Asp Leu Ile His Trp Glu Asn Tyr Pro Ile Ala Ile  
 130 135 140  
 Tyr Pro Asp Glu His Glu Asn Gly Val Leu Ser Leu Pro Phe Ser Gly  
 145 150 155 160  
 Ser Ala Val Val Asp Val His Asn Ser Ser Gly Leu Phe Ser Asn Asp  
 165 170 175  
 Thr Ile Pro Glu Arg Ile Val Leu Ile Tyr Thr Asp His Trp Thr  
 180 185 190  
 Gly Val Ala Glu Arg Gln Ala Ile Ala Tyr Thr Thr Asp Gly Gly Tyr  
 195 200 205  
 Thr Phe Lys Lys Tyr Ser Gly Asn Pro Val Leu Asp Ile Asn Ser Leu  
 210 215 220  
 Gln Phe Arg Asp Pro Lys Val Ile Trp Asp Phe Asp Ala Asn Arg Trp  
 225 230 235 240  
 Val Met Ile Val Ala Met Ser Gln Asn Tyr Gly Ile Ala Phe Tyr Ser  
 245 250 255  
 Ser Tyr Asp Leu Ile His Trp Thr Glu Leu Ser Val Phe Ser Thr Ser  
 260 265 270  
 Gly Tyr Leu Gly Leu Gln Tyr Glu Cys Pro Gly Met Ala Arg Val Pro  
 275 280 285  
 Val Glu Gly Thr Asp Glu Tyr Lys Trp Val Leu Phe Ile Ser Ile Asn  
 290 295 300  
 Pro Gly Ala Pro Leu Gly Gly Ser Val Val Gln Tyr Phe Val Gly Asp  
 305 310 315 320  
 Trp Asn Gly Thr Asn Phe Val Pro Asp Asp Gly Gln Thr Arg Phe Val  
 325 330 335  
 Asp Leu Gly Lys Asp Phe Tyr Ala Ser Ala Leu Tyr His Ser Ser Ser  
 340 345 350  
 Ala Asn Ala Asp Val Ile Gly Val Gly Trp Ala Ser Asn Trp Gln Tyr  
 355 360 365  
 Thr Asn Gln Ala Pro Thr Gln Val Phe Arg Ser Ala Met Thr Val Ala  
 370 375 380  
 Arg Lys Phe Thr Leu Arg Asp Val Pro Gln Asn Pro Met Thr Asn Leu  
 385 390 395 400  
 Thr Ser Leu Ile Gln Thr Pro Leu Asn Val Ser Leu Leu Arg Asp Glu  
 405 410 415

-continued

Thr Leu Phe Thr Ala Pro Val Ile Asn Ser Ser Ser Ser Leu Ser Gly  
 420 425 430  
 Ser Pro Ile Thr Leu Pro Ser Asn Thr Ala Phe Glu Phe Asn Val Thr  
 435 440 445  
 Leu Ser Ile Asn Tyr Thr Glu Gly Cys Thr Thr Gly Tyr Cys Leu Gly  
 450 455 460  
 Arg Ile Ile Ile Asp Ser Asp Asp Pro Tyr Arg Leu Gln Ser Ile Ser  
 465 470 475 480  
 Val Asp Val Asp Phe Ala Ala Ser Thr Leu Val Ile Asn Arg Ala Lys  
 485 490 495  
 Ala Gln Met Gly Trp Phe Asn Ser Leu Phe Thr Pro Ser Phe Ala Asn  
 500 505 510  
 Asp Ile Tyr Ile Tyr Gly Asn Val Thr Leu Tyr Gly Ile Val Asp Asn  
 515 520 525  
 Gly Leu Leu Glu Leu Tyr Val Asn Asn Gly Glu Lys Thr Tyr Thr Asn  
 530 535 540  
 Asp Phe Phe Phe Leu Gln Gly Ala Thr Pro Gly Gln Ile Ser Phe Ala  
 545 550 555 560  
 Ala Phe Gln Gly Val Ser Phe Asn Asn Val Thr Val Thr Pro Leu Lys  
 565 570 575  
 Thr Ile Trp Asn Cys  
 580

<210> SEQ ID NO 22  
 <211> LENGTH: 550  
 <212> TYPE: PRT  
 <213> ORGANISM: Pichia anomala

<400> SEQUENCE: 22

Met Ile Gln Leu Ser Pro Leu Leu Leu Leu Pro Leu Phe Ser Val Phe  
 1 5 10 15  
 Asn Ser Ile Ala Asp Ala Ser Thr Glu Tyr Leu Arg Pro Gln Ile His  
 20 25 30  
 Leu Thr Pro Asp Gln Gly Trp Met Asn Asp Pro Asn Gly Met Phe Tyr  
 35 40 45  
 Asp Arg Lys Asp Lys Leu Trp His Val Tyr Phe Gln His Asn Pro Asp  
 50 55 60  
 Lys Lys Ser Ile Trp Ala Thr Pro Val Thr Trp Gly His Ser Thr Ser  
 65 70 75 80  
 Lys Asp Leu Leu Thr Trp Asp Tyr His Gly Asn Ala Leu Glu Pro Glu  
 85 90 95  
 Asn Asp Asp Glu Gly Ile Phe Ser Gly Ser Val Val Val Asp Arg Asn  
 100 105 110  
 Asn Thr Ser Gly Phe Phe Asn Asp Ser Thr Asp Pro Glu Gln Arg Ile  
 115 120 125  
 Val Ala Ile Tyr Thr Asn Asn Ala Gln Leu Gln Thr Gln Glu Ile Ala  
 130 135 140  
 Tyr Ser Leu Asp Lys Gly Tyr Ser Phe Ile Lys Tyr Asp Gln Asn Pro  
 145 150 155 160  
 Val Ile Asn Val Asn Ser Ser Gln Gln Arg Asp Pro Lys Val Leu Trp  
 165 170 175  
 His Asp Glu Ser Asn Gln Trp Ile Met Val Val Ala Lys Thr Gln Glu  
 180 185 190  
 Phe Lys Val Gln Ile Tyr Gly Ser Pro Asp Leu Lys Lys Trp Asp Leu  
 195 200 205

-continued

---

Lys Ser Asn Phe Thr Ser Asn Gly Tyr Leu Gly Phe Gln Tyr Glu Cys  
 210 215 220  
 Pro Gly Leu Phe Lys Leu Pro Ile Glu Asn Pro Leu Asn Asp Thr Val  
 225 230 235 240  
 Thr Ser Lys Trp Val Leu Leu Leu Ala Ile Asn Pro Gly Ser Pro Leu  
 245 250 255  
 Gly Gly Ser Ile Asn Glu Tyr Phe Ile Gly Asp Phe Asp Gly Thr Thr  
 260 265 270  
 Phe His Pro Asp Asp Gly Ala Thr Arg Phe Met Asp Ile Gly Lys Asp  
 275 280 285  
 Phe Tyr Ala Phe Gln Ser Phe Asp Asn Thr Glu Pro Glu Asp Gly Ala  
 290 295 300  
 Leu Gly Leu Ala Trp Ala Ser Asn Trp Gln Tyr Ala Asn Thr Val Pro  
 305 310 315 320  
 Thr Glu Asn Trp Arg Ser Ser Met Ser Leu Val Arg Asn Tyr Thr Leu  
 325 330 335  
 Lys Tyr Val Asp Val Asn Pro Glu Asn Tyr Gly Leu Thr Leu Ile Gln  
 340 345 350  
 Lys Pro Val Tyr Asp Thr Lys Glu Thr Arg Leu Asn Glu Thr Leu Lys  
 355 360 365  
 Thr Leu Glu Thr Ile Asn Glu Tyr Glu Val Asn Asp Leu Lys Leu Asp  
 370 375 380  
 Lys Ser Ser Phe Val Ala Thr Asp Phe Asn Thr Glu Arg Asn Ala Thr  
 385 390 395 400  
 Gly Val Phe Glu Phe Asp Leu Lys Phe Thr Gln Thr Asp Leu Lys Met  
 405 410 415  
 Gly Tyr Ser Asn Met Thr Thr Gln Phe Gly Leu Tyr Ile His Ser Gln  
 420 425 430  
 Thr Val Lys Gly Ser Gln Glu Thr Leu Gln Leu Val Phe Asp Thr Leu  
 435 440 445  
 Ser Thr Thr Trp Tyr Ile Asp Arg Thr Thr Gln His Ser Phe Gln Arg  
 450 455 460  
 Asn Ser Pro Val Phe Thr Glu Arg Ile Ser Thr Tyr Val Glu Lys Ile  
 465 470 475 480  
 Asp Thr Thr Asp Gln Gly Asn Val Tyr Thr Leu Tyr Gly Val Val Asp  
 485 490 495  
 Arg Asn Ile Leu Glu Leu Tyr Phe Asn Asp Gly Ser Ile Ala Met Thr  
 500 505 510  
 Asn Thr Phe Phe Phe Arg Glu Gly Lys Ile Pro Thr Ser Phe Glu Val  
 515 520 525  
 Val Cys Asp Ser Glu Lys Ser Phe Ile Thr Ile Asp Glu Leu Ser Val  
 530 535 540  
 Arg Glu Leu Ala Arg Lys  
 545 550

<210> SEQ ID NO 23  
 <211> LENGTH: 533  
 <212> TYPE: PRT  
 <213> ORGANISM: Debaryomyces occidentalis

<400> SEQUENCE: 23

Met Val Gln Val Leu Ser Val Leu Val Ile Pro Leu Leu Thr Leu Phe  
 1 5 10 15  
 Phe Gly Tyr Val Ala Ser Ser Ser Ile Asp Leu Ser Val Asp Thr Ser  
 20 25 30





-continued

Asp Tyr Asp Gln Asp Leu Arg Val Phe Ser Leu Tyr Gly Ile Val Asp  
 465 470 475 480  
 Lys Asn Ile Ile Glu Leu Tyr Phe Asn Asp Gly Thr Val Ala Met Thr  
 485 490 495  
 Asn Thr Phe Phe Met Gly Glu Gly Lys Tyr Pro His Asp Ile Gln Ile  
 500 505 510  
 Val Thr Asp Thr Glu Glu Pro Leu Phe Glu Leu Glu Ser Val Ile Ile  
 515 520 525  
 Arg Glu Leu Asn Lys  
 530

<210> SEQ ID NO 24  
 <211> LENGTH: 654  
 <212> TYPE: PRT  
 <213> ORGANISM: *Oryza sativa*

<400> SEQUENCE: 24

Met Ala Thr Ser Arg Leu Thr Pro Ala Tyr Asp Leu Lys Asn Ala Ala  
 1 5 10 15  
 Ala Ala Val Tyr Thr Pro Leu Pro Glu Gln Pro His Ser Ala Glu Val  
 20 25 30  
 Glu Ile Arg Asp Arg Lys Pro Phe Lys Ile Ile Ser Ala Ile Ile Leu  
 35 40 45  
 Ser Ser Leu Leu Leu Leu Ala Leu Ile Leu Val Ala Val Asn Tyr Gln  
 50 55 60  
 Ala Pro Pro Ser His Ser Ser Gly Asp Asn Ser Gln Pro Ala Ala Val  
 65 70 75 80  
 Met Pro Pro Ser Arg Gly Val Ser Gln Gly Val Ser Glu Lys Ala Phe  
 85 90 95  
 Arg Gly Ala Ser Gly Ala Gly Asn Gly Val Ser Phe Ala Trp Ser Asn  
 100 105 110  
 Leu Met Leu Ser Trp Gln Arg Thr Ser Tyr His Phe Gln Pro Val Lys  
 115 120 125  
 Asn Trp Met Asn Asp Pro Asn Gly Pro Leu Tyr Tyr Lys Gly Trp Tyr  
 130 135 140  
 His Leu Phe Tyr Gln Tyr Asn Pro Asp Ser Ala Val Trp Gly Asn Ile  
 145 150 155 160  
 Thr Trp Gly His Ala Val Ser Thr Asp Leu Ile Asn Trp Leu His Leu  
 165 170 175  
 Pro Phe Ala Met Val Pro Asp Gln Trp Tyr Asp Val Asn Gly Val Trp  
 180 185 190  
 Thr Gly Ser Ala Thr Ile Leu Pro Asp Gly Arg Ile Val Met Leu Tyr  
 195 200 205  
 Thr Gly Asp Thr Asp Asp Tyr Val Gln Asp Gln Asn Leu Ala Phe Pro  
 210 215 220  
 Ala Asn Leu Ser Asp Pro Leu Leu Val Asp Trp Val Lys Tyr Pro Asn  
 225 230 235 240  
 Asn Pro Val Ile Tyr Pro Pro Pro Gly Ile Gly Val Lys Asp Phe Arg  
 245 250 255  
 Asp Pro Thr Thr Ala Gly Thr Ala Gly Met Gln Asn Gly Gln Arg Leu  
 260 265 270  
 Val Thr Ile Gly Ser Lys Val Gly Lys Thr Gly Ile Ser Leu Val Tyr  
 275 280 285  
 Glu Thr Thr Asn Phe Thr Thr Phe Lys Leu Leu Tyr Gly Val Leu His  
 290 295 300

-continued

Ala Val Pro Gly Thr Gly Met Trp Glu Cys Val Asp Leu Tyr Pro Val  
 305 310 315 320  
 Ser Thr Thr Gly Glu Asn Gly Leu Asp Thr Ser Val Asn Gly Leu Gly  
 325 330 335  
 Val Lys His Val Leu Lys Thr Ser Leu Asp Asp Asp Lys His Asp Tyr  
 340 345 350  
 Tyr Ala Leu Gly Thr Tyr Asp Pro Val Lys Asn Lys Trp Thr Pro Asp  
 355 360 365  
 Asn Pro Asp Leu Asp Val Gly Ile Gly Leu Arg Leu Asp Tyr Gly Lys  
 370 375 380  
 Tyr Tyr Ala Ala Arg Thr Phe Tyr Asp Gln Asn Lys Gln Arg Arg Ile  
 385 390 395 400  
 Leu Trp Gly Trp Ile Gly Glu Thr Asp Leu Glu Ala Val Asp Leu Met  
 405 410 415  
 Lys Gly Trp Ala Ser Leu Gln Ala Ile Pro Arg Thr Ile Val Phe Asp  
 420 425 430  
 Lys Lys Thr Gly Thr Asn Val Leu Gln Arg Pro Glu Glu Glu Val Glu  
 435 440 445  
 Ser Trp Ser Ser Gly Asp Pro Ile Thr Gln Arg Arg Ile Phe Glu Pro  
 450 455 460  
 Gly Ser Val Val Pro Ile His Val Ser Gly Ala Thr Gln Leu Asp Ile  
 465 470 475 480  
 Thr Ala Ser Phe Glu Val Asp Glu Thr Leu Leu Glu Thr Thr Ser Glu  
 485 490 495  
 Ser His Asp Ala Gly Tyr Asp Cys Ser Asn Ser Gly Gly Ala Gly Thr  
 500 505 510  
 Arg Gly Ser Leu Gly Pro Phe Gly Leu Leu Val Val Ala Asp Glu Lys  
 515 520 525  
 Leu Ser Glu Leu Thr Pro Val Tyr Leu Tyr Val Ala Lys Gly Gly Asp  
 530 535 540  
 Gly Lys Ala Lys Ala His Leu Cys Ala Tyr Gln Thr Arg Ser Ser Met  
 545 550 555 560  
 Ala Ser Gly Val Glu Lys Glu Val Tyr Gly Ser Ala Val Pro Val Leu  
 565 570 575  
 Asp Gly Glu Asn Tyr Ser Ala Arg Ile Leu Ile Asp His Ser Ile Val  
 580 585 590  
 Glu Ser Phe Ala Gln Ala Gly Arg Thr Cys Val Arg Ser Arg Asp Tyr  
 595 600 605  
 Pro Thr Lys Asp Ile Tyr Gly Ala Ala Arg Cys Phe Phe Phe Asn Asn  
 610 615 620  
 Ala Thr Glu Ala Ser Val Arg Ala Ser Leu Lys Ala Trp Gln Met Lys  
 625 630 635 640  
 Ser Phe Ile Arg Pro Tyr Pro Phe Ile Pro Asp Gln Lys Ser  
 645 650

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 690

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Allium cepa

&lt;400&gt; SEQUENCE: 25

Met Ser Ser Asp Asp Leu Glu Ser Pro Pro Ser Ser Tyr Leu Pro Ile  
 1 5 10 15  
 Pro Pro Ser Asp Glu Phe His Asp Gln Pro Pro Pro Leu Arg Ser Trp  
 20 25 30

-continued

---

Leu Arg Leu Leu Ser Ile Pro Leu Ala Leu Met Phe Leu Leu Phe Leu  
           35                                  40                                  45  
 Ala Thr Phe Leu Ser Asn Leu Glu Ser Pro Pro Ser Asp Ser Gly Leu  
           50                                  55                                  60  
 Val Ser Asp Pro Val Thr Phe Asp Val Asn Pro Ala Val Val Arg Arg  
   65                                  70                                  75                                  80  
 Gly Lys Asp Ala Gly Val Ser Asp Lys Thr Ser Gly Val Asp Ser Gly  
                                   85                                  90  
 Phe Val Leu Asp Pro Val Ala Val Asp Ala Asn Ser Val Val Val His  
                                   100                                  105                                  110  
 Arg Gly Lys Asp Ala Gly Val Ser Asp Lys Thr Ser Gly Val Asp Ser  
                                   115                                  120                                  125  
 Gly Leu Leu Lys Asp Ser Pro Leu Gly Pro Tyr Pro Trp Thr Asn Gln  
   130                                  135                                  140  
 Met Leu Ser Trp Gln Arg Thr Gly Phe His Phe Gln Pro Val Lys Asn  
   145                                  150                                  155                                  160  
 Trp Met Asn Asp Pro Asn Gly Pro Leu Tyr Tyr Lys Gly Trp Tyr His  
                                   165                                  170                                  175  
 Phe Phe Tyr Gln Tyr Asn Pro Glu Gly Ala Val Trp Gly Asn Ile Ala  
                                   180                                  185                                  190  
 Trp Gly His Ala Val Ser Arg Asp Leu Val His Trp Thr His Leu Pro  
   195                                  200                                  205  
 Leu Ala Met Val Pro Asp Gln Trp Tyr Asp Ile Asn Gly Val Trp Thr  
   210                                  215                                  220  
 Gly Ser Ala Thr Ile Leu Pro Asp Gly Gln Ile Val Met Leu Tyr Thr  
   225                                  230                                  235                                  240  
 Gly Ala Thr Asn Glu Ser Val Gln Val Gln Asn Leu Ala Val Pro Ala  
                                   245                                  250                                  255  
 Asp Gln Ser Asp Thr Leu Leu Leu Arg Trp Lys Lys Ser Glu Ala Asn  
                                   260                                  265                                  270  
 Pro Ile Leu Val Pro Pro Pro Gly Ile Gly Asp Lys Asp Phe Arg Asp  
   275                                  280                                  285  
 Pro Thr Thr Ala Trp Tyr Glu Pro Ser Asp Asp Thr Trp Arg Ile Val  
   290                                  295                                  300  
 Ile Gly Ser Lys Asp Ser Ser His Ser Gly Ile Ala Ile Val Tyr Ser  
   305                                  310                                  315                                  320  
 Thr Lys Asp Phe Ile Asn Tyr Lys Leu Ile Pro Gly Ile Leu His Ala  
                                   325                                  330                                  335  
 Val Glu Arg Val Gly Met Trp Glu Cys Val Asp Phe Tyr Pro Val Ala  
                                   340                                  345                                  350  
 Thr Ala Asp Ser Ser His Ala Asn His Gly Leu Asp Pro Ser Ala Arg  
                                   355                                  360                                  365  
 Pro Ser Pro Ala Val Lys His Val Leu Lys Ala Ser Met Asp Asp Asp  
   370                                  375                                  380  
 Arg His Asp Tyr Tyr Ala Ile Gly Thr Tyr Asp Pro Ala Gln Asn Thr  
   385                                  390                                  395                                  400  
 Trp Val Pro Asp Asp Ala Ser Val Asp Val Gly Ile Gly Leu Arg Tyr  
                                   405                                  410                                  415  
 Asp Trp Gly Lys Phe Tyr Ala Ser Lys Thr Phe Tyr Asp His Ala Lys  
                                   420                                  425                                  430  
 Lys Arg Arg Ile Leu Trp Ser Trp Ile Gly Glu Thr Asp Ser Glu Thr  
                                   435                                  440                                  445  
 Ala Asp Ile Ala Lys Gly Trp Ala Ser Leu Gln Gly Val Pro Arg Thr

-continued

---

450		455		460
Val Leu Leu Asp Val Lys Thr Gly Ser Asn Leu Ile Thr Trp Pro Val				
465		470		475
Val Glu Ile Glu Ser Leu Arg Thr Arg Pro Arg Asp Phe Ser Gly Ile				
		485		490
				495
Thr Val Asp Ala Gly Ser Thr Phe Lys Leu Asp Val Gly Gly Ala Ala				
		500		505
				510
Gln Leu Asp Ile Glu Ala Glu Phe Lys Ile Ser Ser Glu Glu Leu Glu				
		515		520
				525
Ala Val Lys Glu Ala Asp Val Ser Tyr Asn Cys Ser Ser Ser Gly Gly				
		530		535
				540
Ala Ala Glu Arg Gly Val Leu Gly Pro Phe Gly Leu Leu Val Leu Ala				
		545		550
				555
Asn Gln Asp Leu Thr Glu Gln Thr Ala Thr Tyr Phe Tyr Val Ser Arg				
		565		570
				575
Gly Met Asp Gly Glu Leu Asn Thr His Phe Cys Gln Asp Glu Lys Arg				
		580		585
				590
Ser Ser Lys Ala Ser Asp Ile Val Lys Arg Ile Val Gly His Ser Val				
		595		600
				605
Pro Val Leu Asp Gly Glu Ser Phe Ala Leu Arg Ile Leu Val Asp His				
		610		615
				620
Ser Ile Val Glu Ser Phe Ala Gln Gly Gly Arg Ala Ser Ala Thr Ser				
		625		630
				635
Arg Val Tyr Pro Thr Glu Ala Ile Tyr Asn Asn Ala Arg Val Phe Val				
		645		650
				655
Phe Asn Asn Ala Thr Gly Ala Lys Val Thr Ala Gln Ser Leu Lys Val				
		660		665
				670
Trp His Met Ser Thr Ala Ile Asn Glu Ile Tyr Asp Pro Ala Thr Ser				
		675		680
				685
Val Met				
690				

<210> SEQ ID NO 26  
 <211> LENGTH: 501  
 <212> TYPE: PRT  
 <213> ORGANISM: Beta vulgaris  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (382)..(382)  
 <223> OTHER INFORMATION: Any amino acid

<400> SEQUENCE: 26

Leu Phe Tyr Gln Tyr Asn Pro Asn Gly Val Ile Trp Gly Pro Pro Val				
1		5		10
				15
Trp Gly His Ser Thr Ser Lys Asp Leu Val Asn Trp Val Pro Gln Pro				
		20		25
				30
Leu Thr Met Glu Pro Glu Met Ala Ala Asn Ile Asn Gly Ser Trp Ser				
		35		40
				45
Gly Ser Ala Thr Ile Leu Pro Gly Asn Lys Pro Ala Ile Leu Phe Thr				
		50		55
				60
Gly Leu Asp Pro Lys Tyr Glu Gln Val Gln Val Leu Ala Tyr Pro Lys				
		65		70
				75
Asp Thr Ser Asp Pro Asn Leu Lys Glu Trp Phe Leu Ala Pro Gln Asn				
		85		90
				95
Pro Val Met Phe Pro Thr Pro Gln Asn Gln Ile Asn Ala Thr Ser Phe				
		100		105
				110

-continued

---

Arg Asp Pro Thr Thr Ala Trp Arg Leu Pro Asp Gly Val Trp Arg Leu  
 115 120 125  
 Leu Ile Gly Ser Lys Arg Gly Gln Arg Gly Leu Ser Leu Leu Phe Arg  
 130 135 140  
 Ser Arg Asp Phe Val His Trp Val Gln Ala Lys His Pro Leu Tyr Ser  
 145 150 155 160  
 Asp Lys Leu Ser Gly Met Trp Glu Cys Pro Asp Phe Phe Pro Val Tyr  
 165 170 175  
 Ala Asn Gly Asp Gln Met Gly Val Asp Thr Ser Ile Ile Gly Ser His  
 180 185 190  
 Val Lys His Val Leu Lys Asn Ser Leu Asp Ile Thr Lys His Asp Ile  
 195 200 205  
 Tyr Thr Ile Gly Asp Tyr Asn Ile Lys Lys Asp Ala Tyr Thr Pro Asp  
 210 215 220  
 Ile Gly Tyr Met Asn Asp Ser Ser Leu Arg Tyr Asp Tyr Gly Lys Tyr  
 225 230 235 240  
 Tyr Ala Ser Lys Thr Phe Phe Asp Asp Ala Lys Lys Glu Arg Ile Leu  
 245 250 255  
 Leu Gly Trp Ala Asn Glu Ser Ser Ser Val Glu Asp Asp Ile Lys Lys  
 260 265 270  
 Gly Trp Ser Gly Ile His Thr Ile Pro Arg Lys Ile Trp Leu Asp Lys  
 275 280 285  
 Leu Gly Lys Gln Leu Ile Gln Trp Pro Ile Ala Asn Ile Glu Lys Leu  
 290 295 300  
 Arg Gln Lys Pro Val Asn Ile Tyr Arg Lys Val Leu Lys Gly Gly Ser  
 305 310 315 320  
 Gln Ile Glu Val Ser Gly Ile Thr Ala Ala Gln Ala Asp Val Glu Ile  
 325 330 335  
 Ser Phe Lys Ile Lys Asp Leu Lys Asn Val Glu Lys Phe Asp Ala Ser  
 340 345 350  
 Trp Thr Ser Pro Gln Leu Leu Cys Ser Lys Lys Gly Ala Ser Val Lys  
 355 360 365  
 Gly Gly Leu Gly Pro Phe Gly Leu Leu Thr Leu Ala Ser Xaa Gly Leu  
 370 375 380  
 Glu Glu Tyr Thr Ala Val Phe Phe Arg Ile Phe Lys Ala Tyr Asp Asn  
 385 390 395 400  
 Lys Phe Val Val Leu Met Cys Ser Asp Gln Ser Arg Ser Ser Leu Asn  
 405 410 415  
 Pro Thr Asn Asp Lys Thr Thr Tyr Gly Thr Phe Val Asp Val Asn Pro  
 420 425 430  
 Ile Arg Glu Gly Leu Ser Leu Arg Val Leu Ile Asp His Ser Val Val  
 435 440 445  
 Glu Ser Phe Gly Ala Lys Gly Lys Asn Val Ile Thr Ala Arg Val Tyr  
 450 455 460  
 Pro Thr Leu Ala Ile Asn Glu Lys Ala His Leu Tyr Val Phe Asn Arg  
 465 470 475 480  
 Gly Thr Ser Asn Val Glu Ile Thr Gly Leu Thr Ala Trp Ser Met Lys  
 485 490 495  
 Lys Ala Asn Ile Ala  
 500

&lt;210&gt; SEQ ID NO 27

&lt;211&gt; LENGTH: 518

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bifidobacterium breve

-continued

&lt;400&gt; SEQUENCE: 27

Met Thr Asp Phe Thr Pro Glu Thr Pro Val Leu Thr Pro Ile Arg Asp  
 1 5 10 15  
 His Ala Ala Glu Leu Ala Lys Ala Glu Ala Gly Val Ala Glu Met Ala  
 20 25 30  
 Ala Lys Arg Asn Asn Arg Trp Tyr Pro Lys Tyr His Ile Ala Ser Asn  
 35 40 45  
 Gly Gly Trp Ile Asn Asp Pro Asn Gly Leu Cys Phe Tyr Lys Gly Arg  
 50 55 60  
 Trp His Val Phe Tyr Gln Leu His Pro Tyr Gly Thr Gln Trp Gly Pro  
 65 70 75 80  
 Met His Trp Gly His Val Ser Ser Thr Asp Met Leu Asn Trp Lys Arg  
 85 90 95  
 Glu Pro Ile Met Phe Ala Pro Ser Leu Glu Gln Glu Lys Asp Gly Val  
 100 105 110  
 Phe Ser Gly Ser Ala Val Ile Asp Asp Asn Gly Asp Leu Arg Phe Tyr  
 115 120 125  
 Tyr Thr Gly His Arg Trp Ala Asn Gly His Asp Asn Thr Gly Gly Asp  
 130 135 140  
 Trp Gln Val Gln Met Thr Ala Leu Pro Asp Asn Asp Glu Leu Thr Ser  
 145 150 155 160  
 Ala Thr Lys Gln Gly Met Ile Ile Asp Cys Pro Thr Asp Lys Val Asp  
 165 170 175  
 His His Tyr Arg Asp Pro Lys Val Trp Lys Thr Gly Asp Thr Trp Tyr  
 180 185 190  
 Met Thr Phe Gly Val Ser Ser Glu Asp Lys Arg Gly Gln Met Trp Leu  
 195 200 205  
 Phe Ser Ser Lys Asp Met Val Arg Trp Glu Tyr Glu Arg Val Leu Phe  
 210 215 220  
 Gln His Pro Asp Pro Asp Val Phe Met Leu Glu Cys Pro Asp Phe Phe  
 225 230 235 240  
 Pro Ile Lys Asp Lys Asp Gly Asn Glu Lys Trp Val Ile Gly Phe Ser  
 245 250 255  
 Ala Met Gly Ser Lys Pro Ser Gly Phe Met Asn Arg Asn Val Asn Asn  
 260 265 270  
 Ala Gly Tyr Met Ile Gly Thr Trp Glu Pro Gly Gly Glu Phe Lys Pro  
 275 280 285  
 Glu Thr Glu Phe Arg Leu Trp Asp Cys Gly His Asn Tyr Tyr Ala Pro  
 290 295 300  
 Gln Ser Phe Asn Val Asp Gly Arg Gln Ile Val Tyr Gly Trp Met Ser  
 305 310 315 320  
 Pro Phe Val Gln Pro Ile Pro Met Glu Asp Asp Gly Trp Cys Gly Gln  
 325 330 335  
 Leu Thr Leu Pro Arg Glu Ile Thr Leu Asp Asp Asp Gly Asp Val Val  
 340 345 350  
 Thr Ala Pro Val Ala Glu Met Glu Gly Leu Arg Glu Asp Thr Leu Asp  
 355 360 365  
 His Gly Ser Ile Thr Leu Asp Met Asp Gly Glu Gln Val Ile Ala Asp  
 370 375 380  
 Asp Ala Glu Ala Val Glu Ile Glu Met Thr Ile Asp Leu Ala Ala Ser  
 385 390 395 400  
 Thr Ala Asp Arg Ala Gly Leu Lys Ile His Ala Thr Glu Asp Gly Ala  
 405 410 415

-continued

Tyr Thr Tyr Val Ala Tyr Asp Asp Gln Ile Gly Arg Val Val Val Asp  
                   420                                  425                                  430  
 Arg Gln Ala Met Ala Asn Gly Asp His Gly Tyr Arg Ala Ala Pro Leu  
                   435                                  440                                  445  
 Thr Asp Ala Glu Leu Ala Ser Gly Lys Leu Asp Leu Arg Val Phe Val  
                   450                                  455                                  460  
 Asp Arg Gly Ser Val Glu Val Tyr Val Asn Gly Gly His Gln Val Leu  
                   465                                  470                                  475                                  480  
 Ser Ser Tyr Ser Tyr Ala Ser Glu Gly Pro Arg Ala Ile Lys Leu Val  
                                   485                                  490                                  495  
 Ala Glu Phe Gly Asn Leu Lys Val Glu Ser Leu Lys Leu His His Met  
                   500                                  505                                  510  
 Lys Ser Ile Gly Leu Glu  
                   515

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 532

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

&lt;400&gt; SEQUENCE: 28

Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys  
 1                  5                                  10                                  15  
 Ile Ser Ala Ser Met Thr Asn Glu Thr Ser Asp Arg Pro Leu Val His  
                   20                                  25                                  30  
 Phe Thr Pro Asn Lys Gly Trp Met Asn Asp Pro Asn Gly Leu Trp Tyr  
                   35                                  40                                  45  
 Asp Glu Lys Asp Ala Lys Trp His Leu Tyr Phe Gln Tyr Asn Pro Asn  
                   50                                  55                                  60  
 Asp Thr Val Trp Gly Thr Pro Leu Phe Trp Gly His Ala Thr Ser Asp  
 65                  70                                  75                                  80  
 Asp Leu Thr Asn Trp Glu Asp Gln Pro Ile Ala Ile Ala Pro Lys Arg  
                   85                                  90                                  95  
 Asn Asp Ser Gly Ala Phe Ser Ser Gly Ser Met Val Val Asp Tyr Asn Asn  
                   100                                  105                                  110  
 Thr Ser Gly Phe Phe Asn Asp Thr Ile Asp Pro Arg Gln Arg Cys Val  
                   115                                  120                                  125  
 Ala Ile Trp Thr Tyr Asn Thr Pro Glu Ser Glu Glu Gln Tyr Ile Ser  
                   130                                  135                                  140  
 Tyr Ser Leu Asp Gly Gly Tyr Thr Phe Thr Glu Tyr Gln Lys Asn Pro  
 145                  150                                  155                                  160  
 Val Leu Ala Ala Asn Ser Thr Gln Phe Arg Asp Pro Lys Val Phe Trp  
                   165                                  170                                  175  
 Tyr Glu Pro Ser Gln Lys Trp Ile Met Thr Ala Ala Lys Ser Gln Asp  
                   180                                  185                                  190  
 Tyr Lys Ile Glu Ile Tyr Ser Ser Asp Asp Leu Lys Ser Trp Lys Leu  
                   195                                  200                                  205  
 Glu Ser Ala Phe Ala Asn Glu Gly Phe Leu Gly Tyr Gln Tyr Glu Cys  
                   210                                  215                                  220  
 Pro Gly Leu Ile Glu Val Pro Thr Glu Gln Asp Pro Ser Lys Ser Tyr  
 225                  230                                  235                                  240  
 Trp Val Met Phe Ile Ser Ile Asn Pro Gly Ala Pro Ala Gly Gly Ser  
                   245                                  250                                  255  
 Phe Asn Gln Tyr Phe Val Gly Ser Phe Asn Gly Thr His Phe Glu Ala  
                   260                                  265                                  270



-continued

Phe Asp Asn Gln Ser Arg Val Val Asp Phe Gly Lys Asp Tyr Tyr Ala  
 275 280 285  
 Leu Gln Thr Phe Phe Asn Thr Asp Pro Thr Tyr Gly Ser Ala Leu Gly  
 290 295 300  
 Ile Ala Trp Ala Ser Asn Trp Glu Tyr Ser Ala Phe Val Pro Thr Asn  
 305 310 315 320  
 Pro Trp Arg Ser Ser Met Ser Leu Val Arg Lys Phe Ser Leu Asn Thr  
 325 330 335  
 Glu Tyr Gln Ala Asn Pro Glu Thr Glu Leu Ile Asn Leu Lys Ala Glu  
 340 345 350  
 Pro Ile Leu Asn Ile Ser Asn Ala Gly Pro Trp Ser Arg Phe Ala Thr  
 355 360 365  
 Asn Thr Thr Leu Thr Lys Ala Asn Ser Tyr Asn Val Asp Leu Ser Asn  
 370 375 380  
 Ser Thr Gly Thr Leu Glu Phe Glu Leu Val Tyr Ala Val Asn Thr Thr  
 385 390 395 400  
 Gln Thr Ile Ser Lys Ser Val Phe Ala Asp Leu Ser Leu Trp Phe Lys  
 405 410 415  
 Gly Leu Glu Asp Pro Glu Glu Tyr Leu Arg Met Gly Phe Glu Val Ser  
 420 425 430  
 Ala Ser Ser Phe Phe Leu Asp Arg Gly Asn Ser Lys Val Lys Phe Val  
 435 440 445  
 Lys Glu Asn Pro Tyr Phe Thr Asn Arg Met Ser Val Asn Asn Gln Pro  
 450 455 460  
 Phe Lys Ser Glu Asn Asp Leu Ser Tyr Tyr Lys Val Tyr Gly Leu Leu  
 465 470 475 480  
 Asp Gln Asn Ile Leu Glu Leu Tyr Phe Asn Asp Gly Asp Val Val Ser  
 485 490 495  
 Thr Asn Thr Tyr Phe Met Thr Thr Gly Asn Ala Leu Gly Ser Val Asn  
 500 505 510  
 Met Thr Thr Gly Val Asp Asn Leu Phe Tyr Ile Asp Lys Phe Gln Val  
 515 520 525  
 Arg Glu Val Lys  
 530

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 512

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Zymomonas mobilis*

&lt;400&gt; SEQUENCE: 29

Met Glu Ser Pro Ser Tyr Lys Asn Leu Ile Lys Ala Glu Asp Ala Gln  
 1 5 10 15  
 Lys Lys Ala Gly Lys Arg Leu Leu Ser Ser Glu Trp Tyr Pro Gly Phe  
 20 25 30  
 His Val Thr Pro Leu Thr Gly Trp Met Asn Asp Pro Asn Gly Leu Ile  
 35 40 45  
 Phe Phe Lys Gly Glu Tyr His Leu Phe Tyr Gln Tyr Tyr Pro Phe Ala  
 50 55 60  
 Pro Val Trp Gly Pro Met His Trp Gly His Ala Lys Ser Arg Asp Leu  
 65 70 75 80  
 Val His Trp Glu Thr Leu Pro Val Ala Leu Ala Pro Gly Asp Leu Phe  
 85 90 95  
 Asp Arg Asp Gly Cys Phe Ser Gly Cys Ala Val Asp Asn Asn Gly Val  
 100 105 110



-continued

---

<213> ORGANISM: *Chlorella luteoviridis*

<400> SEQUENCE: 30

```

tgttgaagaa tgagccggcg acttatagga agtggcttgg ttaaggatac tttccgaagc   60
ctaagcgaaa gcaagttgta acaatagcga tatacctctt ttaggtcag tcactttcta   120
tggacccgaa cccgggtgat ctaaccatga ccaggatgaa gcttgggtaa caccaagtga   180
aggctccgaac tcttcgatct ttaaaaatcg tgagatgagt tatggtagg ggtaaatctg   240
gcagttttgc cccgcaaaag ggtaaccttt tgtaattact gactcataac ggtgaagcct   300
aaggcgttag ctatggtaat accgtgggaa gtttcaatac cttcttgcac attttttatt   360
tgcaccttta gtgcaaacag tgtaaagaaa gcgttttgaa acccctaac gactaatttt   420
ttgcttttgc aagaactgca gcaactacca atacacttcc cgtttttttc ttttattaat   480
taaagcaaca taaaaatata ttttatagct ttaatcataa aactatgta gcacttcgtg   540
ctaatgtgct aatgtgctaa tcaaatgaaa agtgttctta aaagtgagtt gaaggtagag   600
tctaactctg cctgaaaggg caagctgcac attttttttt gaatgtgcaa caatggaaat   660
gccaatcgaa ctcgagacta gctggttctc cccgaaatgt gttgaggcgc agcgattcat   720
gattagtacg gtgtaggggt aaagcactgt ttcggtgcgg gctgtgaaaa cggtagcaaaa   780
tcgtggcaaa ctaagaatac tacgcttgta taccatggat cagtgagact atgggggata   840
agctccatag tcaagagggg aacagcccag atcaccagtt aaggcccaaa aatgacagct   900
aagtggcaaa ggagtgtaaa gtgcagaaac aaccaggagg tttgccaga agcagccatc   960
ctttaaagag tgcgtaatag ctcaactg   987

```

<210> SEQ ID NO 31

<211> LENGTH: 1412

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 31

```

gaattcgagt ttaggtccag cgtccgtggg gggggacggg ctgggagctt gggccgggaa   60
gggcaagacg atgcagtcct tctggggagt cacagccgac tgtgtgtggt gcactgtgcg   120
gcccgcagca ctcacacgca aaatgctctg ccgacaggca ggcctctgtc agtgaacat   180
ccacgggtccc tctcatcagg ctcaccttgc tcattgacat aacggaatgc gtaccgctct   240
ttcagatctg tccatccaga gaggggagca ggctccccac cgacgctgtc aaacttgctt   300
cctgccaac cgaaaacatt attgtttgag gggggggggg ggggggcaga ttgcatggcg   360
ggatatctcg tgaggaacat cactgggaca ctgtggaaca cagtgagtgc agtatgcaga   420
gcatgtatgc taggggtcag cgcaggaagg gggcctttcc cagtctccca tgccactgca   480
ccgatccac gactcaccag gaccagcttc ttgatcggct tccgctcccg tggacaccag   540
tgtgtagcct ctggactcca ggtatgctgt caccgcaaag gccagccgat cgtgccgatt   600
cctgggggtg aggatatgag tcagccaact tggggctcag agtgcacact ggggcacgat   660
acgaaacaac atctacaccg tgtcctccat gctgacacac cacagcttcg ctccacctga   720
atgtgggctc atggggccga atcacagcca atgtcgctgc tgccataatg tgatccagac   780
cctctccgcc cagatgccga gcggatcgtg ggcgctgaat agattcctgt ttcgatcaact   840
gtttgggtcc tttccttttc gtctcggatg cgcgtctcga aacaggctgc gtcgggcttt   900
cggatccctt ttgctccctc cgtcaaccatc ctgcccggcg gcaagttgct tgaccctggg   960

```

-continued

---

```

ctggtaccag ggttgagg gttattaccgcg tcaggccatt cccagcccgg attcaattca 1020
aagtctgggc caccaccctc cgccgctctg tctgatcact ccacattcgt gcatacacta 1080
cgttcaagtc ctgatccagg cgtgtctcgg gacaagggtg gcttgagttt gaatctcaag 1140
gaccactcc agcacagctg ctggttgacc cgcacctcgc aactccctac catgtctgct 1200
ggtaggtcca gggatctttg ccatgcacac aggaccccg tttgtggggg ccccggtgca 1260
tgctgtcgt gtgcaggcgc cgggtgggg cctgggccc gcgggagctc aactcctcc 1320
catatgcctg ccgtccctcc caccaccgc gacctggccc cctttgcaga ggaaggcgaa 1380
gtcagcgcca tcgtgtgcga taatggatcc gg 1412

```

```

<210> SEQ ID NO 32
<211> LENGTH: 1627
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 32
gaattcgccc ttgagtttag gtccagcgtc cgtggggggg gcgtgagact cccccctgac 60
cttcgtatgg cagggactcc tacttgccaa gtaatcagtt gacaatgcca cttcaatgct 120
cgttggtgta cactgacgcg ggtctaacat actgggaagc atgaattgcc gacatggact 180
cagttggaga cagtaacage tctttgtgtt ctatcttcag gaacacattt ggcagcgcac 240
ccatacagtg gcgcacacgc agctgtacct gatgtggctc tattcccaca tgtttcaact 300
tgatccaaaa gtcaactcaga ctctcagcag ctagacttga tcgcatcttt ggccatgaag 360
atgcttgccg aactctagga atgggacgag aaaagagcct gctctgatcg gatatttcca 420
ttctctggat gggactgaga tgattctgaa gaaatgctgc tcgacttatt tggaagaaca 480
gcacctgacg catgctttga ggctgctgtg gctgggatgt gctgtatttg tcagcattga 540
gcatctacgg gtagatggcc ataaccacgc gctgcctatc atgcggtggg ttgtgtggaa 600
aacgtacaat ggacagaaat caatcccatt cgcagcctag cgtgcagcca tgcgctccct 660
ctgtagcccc gctccaagac aaagccagcc aatgccagaa cccacataga gagggatatc 720
tcctaatagac ctgcgccatc atttctccca aattaactat aatgccttga ttgtggagtt 780
ggctttggct tgcagctgct cgcgctggca cttttgtagg cagcacaggg tatgccagcg 840
ccgaactttg tgcccttgag caggccacaa gggcacaaga ctacaccatg cagctgggat 900
acttggaaact gataccatc ttaccaagca aggcacagca cagcctgcac cgactcactt 960
tgcttgagcg gggcacagcg ccgagactga tcctgcgagc tgtggggagt tccgactggt 1020
ctggacctcg gtctctgaaa gatgtgtacg atgggatcaa gtcattcaag tatgctcttc 1080
acatgagcaa tcgggggaga cacggtggcc ctaaagggtg tcactctgatt caagtgtagt 1140
gggggggtgc tgtttgtccc ggggcgcccc ccgctccccg acccgggaga agggccccag 1200
aggactcggc cgcacacaga ggaataaccg ggcgtggctc ggcctcgcgc ctccctcttt 1260
caatatttca cctggtgttc agtgcacgga cacgtaaaga actagataca atggccgagg 1320
gaaagacggt gagagcttgg cgttggtgga ccgggcagca tcagaaactc ctcttccccg 1380
cccgccttga aactcactgt aactccctcc tcttccccct cgcagcatct gtctatcgtt 1440
atcgtgagtg aaagggactg ccatgtgtcg ggtcgttgac cacggtcggc tcgggcgctg 1500
ctgccccgct cgcgaacggt ccctgcaaac gccgcgcagc cgtccctttt tctgcccgcg 1560

```

-continued

---

```

ccccaccccc tcgctccccc cttcaatcac accgcagtgc ggacatgtcg attccggcaa 1620
gtccacc 1627

```

```

<210> SEQ ID NO 33
<211> LENGTH: 570
<212> TYPE: DNA
<213> ORGANISM: Chlorella protothecoides

```

```

<400> SEQUENCE: 33

```

```

gaattccctg caggaagaag gccggcagca gctggtactt gtccttcacc tccttgatcg 60
gctgggtgag cttggccggg tcgcagtcgt cgatgccggc atcgcccagc acgctgtgcg 120
gggagccggc atcgacaacc ttggcactgc tcaccttggg caccggcatg gggtcattggc 180
gctgcagacc agcggcctgt cagcatgctg caggcatctg tgtttttag tagatacttt 240
ctgatgcate accacacggt tggaaagtcc ccaagccct tcaacagtct cgacatatga 300
cactcgcgcc ctcttctcgc tcccgtggcc tgatgagggt acgcaggtac cgcagctgcg 360
ccccgtcccc ccagttgccc tggccccgcc gggcccaatc tgttcattgc cgctccctgg 420
cagccgtgaa cttcacacta ccgctctctg tgacctcag cacagcagga atcgccattt 480
cacggcgggt cgttgctgcg gacgctcagc tgatctcgcc tgcgagacc cacagtttga 540
atttgcggtc cccacacaac ctctgacgcc 570

```

```

<210> SEQ ID NO 34
<211> LENGTH: 568
<212> TYPE: DNA
<213> ORGANISM: Chlorella protothecoides
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (133)..(133)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

```

```

<400> SEQUENCE: 34

```

```

gaattccctc aggaagaagg ccggcagcag ctggtacttg tccttcacct ccttgatcgg 60
ctgggtgagc ttcgcaggat cgcagtcgtc gatgccgca tcgcccagca cgctgtgctg 120
ggagccggca tcnacaacct tggcactgct ccccttggtc accggcatgg ggtcatggcg 180
ctgcagccca gccgcctgtc agcatgctgc aggcactctg gtatttagt aggtacttcc 240
tgatgcatca acacacggtt ggaagctccc caagcccctt caacagtctc gacgatgac 300
actcgcgcc tcttctcgc cccgtggcct gatgaggta cgcaggtacc acagctgccc 360
cccgtcccgc cagttgcctt ggccccggcg ggcccactct gttcattgcc gctccctggt 420
agccgtgaac tcacattacc gctctctgtg accttcagca cagcaggaat cgccatttca 480
ccggcggctg ttgctgcgga gcctcagctg atctcgctg cgagacccca cagtttgaat 540
ttgcgggtccc cacacaacct ctgacgcc 568

```

```

<210> SEQ ID NO 35
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

```

```

<400> SEQUENCE: 35

```

```

tgacctaggt gattaattaa ctcgaggcag cagcagctcg gatagtatcg 50

```

```

<210> SEQ ID NO 36
<211> LENGTH: 45

```

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 36  
  
 ctacgagctc aagctttcca ttgtgttcc catcccacta cttec 45

<210> SEQ ID NO 37  
 <211> LENGTH: 32  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 37  
  
 gatcagaatt cgcctgcaa cgcaagggca gc 32

<210> SEQ ID NO 38  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 38  
  
 gcatactagt ggcgggacgg agagagggcg 30

<210> SEQ ID NO 39  
 <211> LENGTH: 1568  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide  
  
 <400> SEQUENCE: 39  
  
 gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60  
 tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120  
 atgggcgctc cgatgccgt ccaggggcag cgctgtttaa atagccaggc ccccgattgc 180  
 aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgact accacttcta 240  
 cacaggccac tcgagcttgt gatcgcactc cgctaagggg ggcctcttc ctcttcgttt 300  
 cagtcaaac ccgcaaacgg cgcgccatat caatgattga acaagatgga ttgcacgcag 360  
 gttctccggc cgcttggtg gagaggtat tcggctatga ctgggcacaa cagacaatcg 420  
 gctgctctga tgccgcctg ttccggctgt cagcgcaggg ggcgccggtt cttttgtca 480  
 agaccgacct gtccggtgcc ctgaatgaac tgcaggacga ggcagcgcgg ctatcgtggc 540  
 tggccacgac gggcgttctc tgccgagctg tgctcgactg tgctcactgaa gcggaagg 600  
 actggctgct attggcgcaa gtgccggggc aggatctcct gtcactctac cttgctcctg 660  
 ccgagaaagt atccatcatg gctgatgcaa tgccggcgct gcatacgctt gatccggcta 720  
 cctgcccatt cgaccaccaa gcgaaacatc gcatcgagcg agcacgtact cggatggaag 780  
 ccggtcttgt cgatcaggat gatctggacg aagagcatca ggggctcgcg ccagccgaac 840  
 tgttcgccag gctcaaggcg cgcattgccg acggcgagga tctcgtcgtg acccatggcg 900  
 atgctgctt gccgaatata atggtgaaa atggccgctt ttctggattc atcgaactgtg 960  
 gccgctggg tgtggcgac cgctatcagg acatagcgtt ggctaccogt gatattgctg 1020

-continued

---

```

aagagcttgg cggcgaatgg gctgaccgct tcctcgtgct ttacggatc gccgctccc 1080
attcgcagcg catcgccctc tatcgccctc ttgacgagtt cttctaagat ctgtcgatcg 1140
acaagtgact cgaggcagca gcagctcggg tagtatcgac acaactctgga cgctggctgt 1200
gtgatggact gttgccgcca cacttctgct cttgacctgt gaatatccct gccgctttta 1260
tcaaacagcc tcagtgtggt tgatcttctg tgtacgcgct tttgcgagtt gctagctgct 1320
tgtgctatctt gcgaatacca ccccagcat ccccttcct cgtttcatat cgettgcatc 1380
ccaaccgcaa cttatctacg ctgtcctgct atccctcagc gctgctcctg ctctgctca 1440
ctgcccctcg cacagccttg gtttgggctc cgctgtatt ctctggtagc tgcaacctgt 1500
aaaccagcac tgcaatgctg atgcacggga agtagtggga tgggaacaca aatggaaagc 1560
ttgagctc 1568

```

```

<210> SEQ ID NO 40
<211> LENGTH: 2571
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (997)..(999)
<223> OTHER INFORMATION: a, c, t, g, unknown or other

```

```

<400> SEQUENCE: 40

```

```

gaattccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgcccgc tgcctgccgc atgcttctgc tggtaggct gggcagtgct 120
gccatgctga ttgagcttg gttcatcggg tggaaactta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgcatgggt gcaagagggc ggcagcactt gctggacgtg ccgctgtgcc 240
tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgctg acaggttgag 300
cagcagtgct agcaaagga gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgttcc agctacaacg 480
gcaggagtca tccaactaac catagctgat caacactgca atcatcggcg gctgatgcaa 540
gcacactgca agacacatgc tgtgcgatgc tgcctgctg cctgctgctg acgcccttga 600
ggtggcagca gctcagccat gcaactggatc aggtgggct gccactgcaa tgtggtggat 660
aggatgcaag tggagcgaat accaaaccct ctggctgctt gctgggttgc atggcatcgc 720
accatcagca ggagcgcagc cgaagggact ggccccatgc acgccatgcc aaaccggagc 780
gcaccgagtg tccacactgt caccaggccc gcaagctttg cagaacctatg ctcatggacg 840
catgtagcgc tgacgtccct tgacggcgct cctctcgggt gtgggaaacg caatgcagca 900
caggcagcag agggcgccgc agcagagcgg cggcagcagc ggcgggggccc acccttcttg 960
cggggtcgcg ccccagccag cgggtgatgc ctgatcnnc caaacgagtt cacattcatt 1020
tgcagcctgg agaagcaggg ctggggcctt tgggctggtg cagcccgcaa tggaaatcgg 1080
gaccgccagg ctagcagcaa agggcgcctc cctactcgcg atcgatgttc catagtgcac 1140
tggactgcat ttgggtgggg cggccggctg tttcttctgt gttgcaaac gcgccacgctc 1200
agcaacctgt cccgtgggtc ccccgtgcgc atgaaatcgt gtgcacgccg atcagctgat 1260
tgcccggctc gegaagtagg cgcctctctt ctgctcgcgc tctctcgcgc ccgccactag 1320

```

-continued

tggcgcgcca	tatcaatgat	tgaacaagat	ggattgcacg	caggttctcc	ggcgcgttg	1380
gtggagagge	tattcggtta	tgactgggca	caacagacaa	tcggctgctc	tgatgcccgc	1440
gtgttccggc	tgtagcgca	ggggcggccg	gttctttttg	tcaagaccga	cctgtccggg	1500
gcccgaatg	aactgcagga	cgaggcagcg	eggctatcgt	ggctggccac	gacgggcggt	1560
ccttgccgag	ctgtgctega	cgttgtcact	gaagcgggaa	gggactggct	gctattgggc	1620
gaagtgccgg	ggcaggatct	cctgtcatct	caccttgctc	ctgccagaaa	agtatccatc	1680
atggctgatg	caatgcggcg	gctgcatacg	cttgatccgg	ctacctgccc	attcgaccac	1740
caagcgaaac	atcgcatcga	gcgagcacgt	actcggatgg	aagccggctc	tgtcgatcag	1800
gatgatctgg	acgaagagca	tcaggggctc	gcgccagccg	aactgttcgc	caggetcaag	1860
gcgcgcatgc	ccgacggcga	ggatctcctc	gtgacccatg	gcgatgcctg	cttgccgaat	1920
atcatgggtg	aaaatggccg	ctttcttgga	ttcatcgact	gtggccggct	gggtgtggcg	1980
gaccgctatc	aggacatagc	gttggtatcc	cgtgatattg	ctgaagagct	tggcggcgaa	2040
tgggctgacc	gcttccctct	gctttacggt	atcgccgctc	ccgattcgca	gcgcatcgcc	2100
ttctatcgcc	ttcttgacga	gttcttctaa	gatctgtcga	tcgacaagtg	actcgaggca	2160
gcagcagctc	ggatagtatc	gacacactct	ggacgctggg	cgtgtgatgg	actggtgccc	2220
ccacacttgc	tgccctgacc	tgtgaatata	cctgccgctt	ttatcaaaaca	gcctcagtgt	2280
gtttgatctt	gtgtgtacgc	gcttttgcca	gttgetagct	gcttgtgcta	tttgccaata	2340
ccacccccag	catccccctc	cctcgtttca	tatcgcttgc	atcccaaccg	caatttatct	2400
acgctgtcct	gctatccctc	agcgtgctc	ctgctcctgc	tactgcccc	tcgcacagcc	2460
ttggtttggg	ctccgcctgt	attctcctgg	tactgcaacc	tgtaaaccag	cactgcaatg	2520
ctgatgcacg	ggaagtatgt	ggatgggaa	acaaatggaa	agcttgagct	c	2571

&lt;210&gt; SEQ ID NO 41

&lt;211&gt; LENGTH: 2550

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: modified\_base

&lt;222&gt; LOCATION: (997)..(999)

&lt;223&gt; OTHER INFORMATION: a, c, t, g, unknown or other

&lt;400&gt; SEQUENCE: 41

gaattccgcc	tgcaacgcaa	gggcagccac	agccgctccc	accgcccgt	gaaccgacac	60
gtgcttgggc	gcctgcccgc	tgccctgccg	atgcttgtgc	tggtagggct	gggcagtgtc	120
gccatgtcga	ttgaggcttg	gttcatcggg	tggaagctta	tgtgtgtgct	ggccttgcac	180
gccgggcaat	gcgcatgggt	gcaagagggc	ggcagcactt	gctggacgtg	ccgcggtgcc	240
tccagggtgt	tcaatcgccg	cagccagagg	gatttcagat	gatcgcgctg	acaggttgag	300
cagcagtgtc	agcaaaggta	gcagtttgcc	agaatgatcg	gttcagctgt	taatcaatgc	360
cagcaagaga	aggggtcaag	tgcaaacacg	ggcatgccac	agcacgggca	ccggggagtg	420
gaatggcacc	accaagtgtg	tgcgagccag	catcgcgcc	tggctgtttc	agctacaacg	480
gcaggagtca	tccaactaac	catagctgat	caacactgca	atcatcggcg	gctgatgcaa	540
gcacctcgtc	agacacatgc	tgtgcatgct	tgccctgctg	cctgctgcgc	acgccgttga	600
gttggcagca	gctcagccat	gcaactggatc	aggctgggct	gccactgcaa	tgtggtggat	660
aggatgcaag	tggagcgaat	accaaaccct	ctggctgctt	gctgggttgc	atggcatcgc	720



-continued

---

```

accatcagca ggagcgcgatg cgaagggact ggcccatgc acgccatgcc aaaccggagc 780
gcaccgagtg tccacactgt caccaggccc gcaagctttg cagaacctatg ctcatggacg 840
catgtagcgc tgacgtccct tgacggcgct cctctcgggt gtgggaaacg caatgcagca 900
caggcagcag agggcggcgc agcagagcgg cggcagcagc ggcgggggccc acccttcttg 960
cggggtcgcg cccagccag cggatgatgc ctgatcnnc caaacgagtt cacattcatt 1020
tgagcctgg agaagcgagg ctggggcctt tgggtggtg cagcccgcaa tggaatgctg 1080
gaccgccagg ctagcagcaa aggcgcctcc cctactccgc atcgatgttc catagtgcac 1140
tggactgcat ttgggtgggg cggccggctg tttcttctgt gttgcaaac gcgccacgtc 1200
agcaacctgt cccgtgggtc cccctgccc atgaaatcgt gtgcacgccc atcagctgat 1260
tgcccggctc ggaagtagg cgcctcttt ctgctcggcc tctctcggtc ccgccactag 1320
tggcgcgcca tatcaatgat cgagcaggac ggcctccac cggctcccc cgcgcctgg 1380
gtggagcgcc tgttcggcta cgaactggcc cagcagacca tggctgctc cgacgcccgc 1440
gtgttcggcc tgtccgcca gggccgccc gtgctgttc tgaagaccga cctgtccggc 1500
gccctgaacg agctgcagga cgaggcccgc cgcctgtcct ggctggccac caccgcgctg 1560
ccctgcgcgc cgtgctgga cgtggtgacc gaggcggccc gcgactggtc gctgctgggc 1620
gaggtgcccc gccaggacct gctgtcctcc cacctggccc ccgcccagaa ggtgtccatc 1680
atggccgacg ccatgcgcgc cctgcacacc ctggacccc ccacctgccc cttegaccac 1740
caggccaagc accgcatcga gcgcgcccgc acccgcatgg aggcgggccc ggtggaccag 1800
gacgacctgg acgaggagca ccagggcctg gccccgcgc agctgttcgc ccgctgaag 1860
gcccgcctgc ccgacggcga ggacctggtg gtgacccacg gcgacgctg cctgcccacc 1920
atcatggtgg agaacggcgc cttctcgggc ttcactgact gcggccgccc gggcgtggcc 1980
gaccgctacc aggacatcgc cctggccacc cgcgacatcg ccgaggagct gggcggcgag 2040
tgggcccacc gcttctggt gctgtacgac atcgcgccc ccgactccca gcgcatcgcc 2100
ttctaccgcc tgctggacga gttcttctga ctcgaggcag cagcagctcg gatagtatcg 2160
acacactctg gacgctggtc gtgtgatgga ctggtgccc cacttctgct gccttgacct 2220
gtgaatatcc ctgcccttt tatcaaacag cctcagtggt tttgatcttg tgtgtacgcg 2280
cttttgcgag ttgctagctg cttgtgctat ttgcgaatac caccccagc atccccttc 2340
ctcgtttcat atcgttgca tcccaccgc aacttatcta cgtgtcctg ctatccctca 2400
gcgctgctcc tgctcctgct cactgcccct cgcacagcct tggtttgggc tccgctgta 2460
ttctcctggt actgcaacct gtaaaccagc actgcaatgc tgatgcacgg gaagtagtgg 2520
gatgggaaca caaatggaac gcttgagctc 2550

```

&lt;210&gt; SEQ ID NO 42

&lt;211&gt; LENGTH: 1547

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 42

```

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc 180

```

-continued

---

```

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg ggcctcttc ctcttcgttt 300
cagtcaaac ccgcaaacgg cgcgccatat caatgatcga gcaggacggc ctccacgccg 360
gctccccgc cgctgggtg gagcgctgt tcggctacga ctgggccag cagaccatcg 420
gctgtccga cgccgcctg ttccgcctgt ccgccaggg ccgccctg ctgttcgtga 480
agaccgacct gtccgggcc ctgaacgagc tgcaggacga gcccgccgc ctgtcctggc 540
tggccaccac cggcgtgccc tgcgcgccg tgetggacgt ggtgaccgag gccggccgcg 600
actggctget gctggggcag gtgccggcc aggacctgt gtccctccac ctggccccg 660
ccgagaaggt gtccatcatg gccgaogcca tgcgccgct gcacacctg gacccccca 720
cctgccccct cgaccaccag gccaaagcacc gcatcgagcg cgcgccacc ccatggagg 780
ccggcctggt ggaccaggac gacctggacg aggagacca gggcctggcc cccgccgagc 840
tgttcgccc cctgaagggc cgcagcccg acggcgagga cctggtggtg acccacggcg 900
acgcctgcct gcccaacatc atggtggaga acggccgctt ctccggttc atcgactgcg 960
gccgcctggg cgtggccgac cgctaccagg acatcgccct ggccaccgc gacatcgccg 1020
aggagctggg cggcgagtgg gccgaocgct tcttgggtgt gtacggcatc gccgccccg 1080
actcccagc catcgccctc tacccctgc tggacgagtt cttctgactc gaggcagcag 1140
cagctcggat agtategaca cactctggac gctggctgtg tgatggactg ttgccgccac 1200
acttctgccc ttgacctgtg aataccctg ccgctttat caaacagct cagtgtgttt 1260
gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg cgaataccac 1320
ccccagcacc ccctccctc gtttcatac gcttgcacc caaccgcaac ttatctacgc 1380
tgtctgcta tcctcagcg ctgctcctgc tctgctcac tgccctcgc acagccttg 1440
tttggctcc gctgtatcc tctgtgtact gcaacctgta aaccagcact gcaatgctga 1500
tgcacgggaa gtagtgggat gggaacacaa atggaaagct tgagctc 1547

```

```

<210> SEQ ID NO 43
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```
<400> SEQUENCE: 43
```

```
gccgcgactg gctgctgctg g 21
```

```

<210> SEQ ID NO 44
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```
<400> SEQUENCE: 44
```

```
aggtcctcgc cgtcgggcat g 21
```

```

<210> SEQ ID NO 45
<211> LENGTH: 1292
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

-continued

&lt;400&gt; SEQUENCE: 45

```

atcaaaggca tagattcaca tttgttgca ttgcagagca atcatcgcgc aggacgaaca      60
tcgctcacca agcacgtact gggcatccgg aggctcgcgc aaattcctgc aacaggactc     120
gctgatcagt tcgccaagg tctacgacgc tcctatcgg cgctagactt caacacatat     180
ttcactgtca cagcctcgcgc atgcatcagg cctcagtctc caccatgaag accatccagt     240
ctcggcacgc cggctccatc ggacatgtgc agtcgggtcg ccgatcggcg gggcgcgcg     300
gatcccgcgc ggcgaccccc gtggcgcgag ctaccgtcgc agcccctcgc tcggccctca     360
acctctcccc caccatcatt cgacaggagg tgctccactc cgccagcgcc cagcaactag     420
actgctgggc ctcccggcg cccgtcttcg agtcccagat cctcccctc ctgacgccc     480
tggacgagat gtggcagccc accgacttcc tcccgcctc gaactcggag gcattcttcg     540
accagatcgg cgacctgccc gcgcatcgg cggccatccc cgacgacctg ctggtctgccc     600
tgggtggggga catgatcacg gaggaggccc tgcccaccta catggccatg ctgaacaccc     660
tggacgtcgt gcgcatgag acagggcaca gccagcacc ctacgccaag tggaccaggg     720
cttggatcgc ggaggagaac cgccatggcg acctgctgaa caagtacatg tggctgacgg     780
ggcggtggg acatgctggc ggtggagcgc accatccagc catgctggcg gtggagcgca     840
ccatccagcg cctcatctca tcgggcatgg acccgggcac ggagaaccac cctaccacg     900
cctttgtgtt caccagcttc caggagcgcg ccaccaagct gagccacggc tccaccgcc     960
gcctggcggt cgccgcccgg gacgaggccc tggccaagat ctgcccggacc attgcccggg    1020
acgagtcgcg ccacgaggcg gcgtacacgc ggaccatgga tgccatcttc cagcgcgacc    1080
ccagcggggc catggtggcg tttgcgcaca tgatgatgcg caagatcacc atgcccggcc    1140
acctcatgga cgacggccag cacggcgcgc gcaacggggg ggcgcaactt gttcgacgac    1200
tttgcggcag tggcggagcg ggcaggggtg tacaccgccc gcgactacat cggcatcctg    1260
cgccacctca tccggcgtg ggacgtggag gg                                     1292

```

&lt;210&gt; SEQ ID NO 46

&lt;211&gt; LENGTH: 364

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 46

```

Met His Gln Ala Ser Val Ser Thr Met Lys Thr Ile Gln Ser Arg His
1           5           10           15
Ala Gly Pro Ile Gly His Val Gln Ser Gly Arg Arg Ser Ala Gly Arg
20           25           30
Ala Gly Ser Arg Met Ala Thr Pro Val Ala Ala Ala Thr Val Ala Ala
35           40           45
Pro Arg Ser Ala Leu Asn Leu Ser Pro Thr Ile Ile Arg Gln Glu Val
50           55           60
Leu His Ser Ala Ser Ala Gln Gln Leu Asp Cys Val Ala Ser Leu Ala
65           70           75           80
Pro Val Phe Glu Ser Gln Ile Leu Pro Leu Leu Thr Pro Val Asp Glu
85           90           95
Met Trp Gln Pro Thr Asp Phe Leu Pro Ala Ser Asn Ser Glu Ala Phe
100          105          110
Phe Asp Gln Ile Gly Asp Leu Arg Ala Arg Ser Ala Ala Ile Pro Asp

```

-continued

115	120	125
Asp Leu Leu Val Cys Leu Val Gly Asp Met Ile Thr Glu Glu Ala Leu 130 135 140		
Pro Thr Tyr Met Ala Met Leu Asn Thr Leu Asp Val Val Arg Asp Glu 145 150 155 160		
Thr Gly His Ser Gln His Pro Tyr Ala Lys Trp Thr Arg Ala Trp Ile 165 170 175		
Ala Glu Glu Asn Arg His Gly Asp Leu Leu Asn Lys Tyr Met Trp Leu 180 185 190		
Thr Gly Arg Val Gly His Ala Gly Gly Ala His His Pro Ala Met 195 200 205		
Leu Ala Val Glu Arg Thr Ile Gln Arg Leu Ile Ser Ser Gly Met Asp 210 215 220		
Pro Gly Thr Glu Asn His Pro Tyr His Ala Phe Val Phe Thr Ser Phe 225 230 235 240		
Gln Glu Arg Ala Thr Lys Leu Ser His Gly Ser Thr Ala Arg Leu Ala 245 250 255		
Val Ala Ala Gly Asp Glu Ala Leu Ala Lys Ile Cys Gly Thr Ile Ala 260 265 270		
Arg Asp Glu Ser Arg His Glu Ala Ala Tyr Thr Arg Thr Met Asp Ala 275 280 285		
Ile Phe Gln Arg Asp Pro Ser Gly Ala Met Val Ala Phe Ala His Met 290 295 300		
Met Met Arg Lys Ile Thr Met Pro Ala His Leu Met Asp Asp Gly Gln 305 310 315 320		
His Gly Ala Arg Asn Gly Gly Ala Gln Leu Val Arg Arg Leu Cys Gly 325 330 335		
Ser Gly Gly Ala Gly Arg Gly Val His Arg Arg Arg Leu His Arg His 340 345 350		
Pro Ala Pro Pro His Pro Ala Leu Gly Arg Gly Gly 355 360		

&lt;210&gt; SEQ ID NO 47

&lt;211&gt; LENGTH: 1395

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 47

```

attatacatc ggcacgtct caggtttcac gatctgcatg ctatctatgg gactgtgact      60
ccgcgggcca ggttgtggtg cgcgagaatc ctccccgctc ctgccttctc atttcctga    120
cgggagtgcg cgctgagcac cgggaggatc atgggcgtcg gcacactcca aaccatcat      180
acatgtggtc gtgcattcac gcatagcgca cggtatgtcc cgcgacgcgc ggctcgaagc    240
cgtggccatc cgacgcgctg cacggccgag gtgagggcac gccctccgc caatggcgcg     300
cagcccatga ccgccttoga cttccggcag tacatgcagc agcgcgccgc gctggtggac    360
gcagcgtcgg acctggcagt gccgtgcagc taccocgaga agatcaacga ggccatgcgg    420
tacagcctgc tggccggggg caagecgcgtg cgcgccgcgc tctgcctcgc tgectgcgag   480
ctcgtggggc gccccctgga ggccggccatg cccgccgctc gcgccatgga gatgatccac   540
accatgagcc tcatccacga cgacctcccc gccatggaca acgacgactt ccggcgcggc    600
cagcccccca accacaaggc ctatggcgag gagattgcga tcctggcggg cgacgcgctg   660

```

-continued

---

```

ctgtcgtga gctttgagca catcgcgcgc gagacgcgag gcgtggaccc ggtgcgcgtc 720
ctggccgcca tctcggagtg gcgcgcgggtg ggcagccgcg ggctgggtggc ggggcaggtg 780
gtggacctgg gtttcgaggg cggcggcggtg gggctggccc cgctgcgcta catccacgag 840
cacaaaaccg cggcgcgtgt ggaggcggcg gtggtgtccg gcgcgctgct gggcggcgcg 900
gaggaggcgg acctggagcg cctgcgcacc tacaaccgcg ccctcggcct cgctttccag 960
gtggtggggg acatcctgga catccggggg accagcgagg agctgggcaa gaccgcgggc 1020
aaggacctga gctcccccaa aaccctctac cgtcccttg tggggctggc caggtccaaa 1080
aaaaattcgg acgaactgat tgaggacgag aaaacccaac tcaccagta cgagccggcc 1140
cgagcggcgc ccctcgtaac cctggccgaa aacatttgaa accggaagaa ctgactgggg 1200
gccccccctg cccccagata cggcggggct cctccatcca gttttgggat gggaggagcg 1260
acaaccgacc ccgtaacct gtgacgcgtt tgccttgcac acgtacgcat gccttgaaac 1320
ccatccatga ccctcaacaa tacctggttg tgtgtagctt ggtcctgaaa aaaaaaaaaa 1380
aaaaaaaaa aaaaa 1395

```

&lt;210&gt; SEQ ID NO 48

&lt;211&gt; LENGTH: 342

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 48

```

Met Gly Val Gly Thr Leu Gln Thr Pro Tyr Thr Cys Gly Arg Ala Phe
1           5           10           15

Thr His Ser Ala Arg Tyr Val Pro Arg Arg Ala Ala Arg Ser Arg Gly
20          25          30

His Pro Thr Arg Cys Thr Ala Glu Val Arg Ala Arg Pro Ser Ala Asn
35          40          45

Gly Ala Gln Pro Met Thr Ala Phe Asp Phe Arg Gln Tyr Met Gln Gln
50          55          60

Arg Ala Ala Leu Val Asp Ala Ala Leu Asp Leu Ala Val Pro Leu Gln
65          70          75          80

Tyr Pro Glu Lys Ile Asn Glu Ala Met Arg Tyr Ser Leu Leu Ala Gly
85          90          95

Gly Lys Arg Val Arg Pro Ala Leu Cys Leu Ala Ala Cys Glu Leu Val
100         105         110

Gly Gly Pro Leu Glu Ala Ala Met Pro Ala Ala Cys Ala Met Glu Met
115         120         125

Ile His Thr Met Ser Leu Ile His Asp Asp Leu Pro Ala Met Asp Asn
130         135         140

Asp Asp Phe Arg Arg Gly Gln Pro Ala Asn His Lys Ala Tyr Gly Glu
145         150         155         160

Glu Ile Ala Ile Leu Ala Gly Asp Ala Leu Leu Ser Leu Ser Phe Glu
165         170         175

His Ile Ala Arg Glu Thr Arg Gly Val Asp Pro Val Arg Val Leu Ala
180         185         190

Ala Ile Ser Glu Trp Arg Ala Val Gly Ser Arg Gly Leu Val Ala Gly
195         200         205

Gln Val Val Asp Leu Gly Phe Glu Gly Gly Gly Val Gly Leu Ala Pro
210         215         220

Leu Arg Tyr Ile His Glu His Lys Thr Ala Ala Leu Leu Glu Ala Ala

```

-continued

---

225	230	235	240
Val Val Ser Gly Ala Leu Leu Gly Gly Ala Glu Glu Ala Asp Leu Glu	245	250	255
Arg Leu Arg Thr Tyr Asn Arg Ala Ile Gly Leu Ala Phe Gln Val Val	260	265	270
Gly Asp Ile Leu Asp Ile Pro Gly Thr Ser Glu Glu Leu Gly Lys Thr	275	280	285
Ala Gly Lys Asp Leu Ser Ser Pro Lys Thr Pro Tyr Pro Ser Leu Val	290	295	300
Gly Leu Ala Arg Ser Lys Lys Ile Ala Asp Glu Leu Ile Glu Asp Ala	305	310	315
Lys Thr Gln Leu Thr Gln Tyr Glu Pro Ala Arg Ala Ala Pro Leu Val	325	330	335
Thr Leu Ala Glu Asn Ile	340		

<210> SEQ ID NO 49  
 <211> LENGTH: 833  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 49

cgatgccat ggcgccctcgg gccgcgggcc tgagggtcca cgcagcgtcc tcggtggccc	60
agacgcacca ggccgcccc ccggcggaca ggaggttcga cgactaccag ccccgaccg	120
ccatcctctt ccccgcccag ggccgcgaca gcgtgggcat ggccggagag ctggcgaagg	180
ccgtccccgc cgcgcggcgg ctgttcgacg ccgcctccga ccagctcggc tatgacctgc	240
tccgcgtgtg cgttgaggcc cccaaggcgc gcctggacag caccgcccgc agccagcccg	300
ccatctacgt ggccagcctg gcggcgggtg agaagctgcy cgcggagggc ggggaggagg	360
cactggccgc catcgacgtc gctgcccgtc tgccttggg cgagtacacc gcgctggcct	420
ttgcggcgc cttctcttc gccgacgggc tgcgcctggt ggccctgcgc ggcgccagca	480
tgcaggccgc cgcgcagccc gcacctcgg gcatggtctc cgtcatcggc ctgccctccg	540
acgcgggtgc cgcgctgtgc gaggccgcca acgcgcaggt ggccccgac caggccgtgc	600
gcatcgccaa ctacctctgc gacggcaact acgcgctcag cgtgggctg gagggtcgcg	660
cgccgggtga gggcctggcc aaggcccaaca aggcgcgcat gacggtgccg ctggcgggtg	720
cgccgcctt ccacaccccc ttcctgcagc cggcgggtga ggcgctgagc gcgggcgctg	780
gcggacacgc cgctggtcgc gccgcgcac cccgtggtca gcaacgggac gcc	833

<210> SEQ ID NO 50  
 <211> LENGTH: 275  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 50

Met Arg Pro Arg Ala Ala Gly Leu Arg Val His Ala Ala Ser Ser Val	1	5	10	15
Ala Gln Thr His Gln Ala Ala Pro Pro Ala Asp Arg Arg Phe Asp Asp	20	25	30	
Tyr Gln Pro Arg Thr Ala Ile Leu Phe Pro Gly Gln Gly Ala Gln Ser				

-continued

35				40				45							
Val	Gly	Met	Ala	Gly	Glu	Leu	Ala	Lys	Ala	Val	Pro	Ala	Ala	Ala	Ala
	50					55					60				
Leu	Phe	Asp	Ala	Ala	Ser	Asp	Gln	Leu	Gly	Tyr	Asp	Leu	Leu	Arg	Val
65				70						75				80	
Cys	Val	Glu	Gly	Pro	Lys	Ala	Arg	Leu	Asp	Ser	Thr	Ala	Val	Ser	Gln
				85					90					95	
Pro	Ala	Ile	Tyr	Val	Ala	Ser	Leu	Ala	Ala	Val	Glu	Lys	Leu	Arg	Ala
			100				105							110	
Glu	Gly	Gly	Glu	Glu	Ala	Leu	Ala	Ile	Asp	Val	Ala	Ala	Gly	Leu	
		115				120					125				
Ser	Leu	Gly	Glu	Tyr	Thr	Ala	Leu	Ala	Phe	Ala	Gly	Ala	Phe	Ser	Phe
	130					135					140				
Ala	Asp	Gly	Leu	Arg	Leu	Val	Ala	Leu	Arg	Gly	Ala	Ser	Met	Gln	Ala
145				150						155				160	
Ala	Ala	Asp	Ala	Ala	Pro	Ser	Gly	Met	Val	Ser	Val	Ile	Gly	Leu	Pro
				165					170					175	
Ser	Asp	Ala	Val	Ala	Ala	Leu	Cys	Glu	Ala	Ala	Asn	Ala	Gln	Val	Ala
		180					185						190		
Pro	Asp	Gln	Ala	Val	Arg	Ile	Ala	Asn	Tyr	Leu	Cys	Asp	Gly	Asn	Tyr
		195				200						205			
Ala	Val	Ser	Gly	Gly	Leu	Glu	Gly	Cys	Ala	Ala	Val	Glu	Gly	Leu	Ala
	210					215					220				
Lys	Ala	His	Lys	Ala	Arg	Met	Thr	Val	Arg	Leu	Ala	Val	Ala	Gly	Ala
225				230						235				240	
Phe	His	Thr	Pro	Phe	Met	Gln	Pro	Ala	Val	Glu	Ala	Leu	Ser	Ala	Gly
				245					250					255	
Ala	Gly	Gly	His	Ala	Ala	Gly	Arg	Ala	Ala	His	Pro	Arg	Gly	Gln	Gln
			260				265						270		
Arg	Asp	Ala													
		275													

&lt;210&gt; SEQ ID NO 51

&lt;211&gt; LENGTH: 787

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 51

```

tgtccatctc cccccaccct ccaccaacc atcgctgacg gcatgcaggc gctgtgttct    60
cacccecgct ccctcagcgc gcgtgcggtg ccccatgggc gggccagccc agcacagcgg    120
gtgtccagcg ccggcccggc ctacaccggc ctgtcccggc acaccctggg ctgccccage    180
acccccacc tccagtcctc gcctcggtc cagaccggc gctcctctc cggtccacc    240
acgcgcatga ccaccaccgc ccagcgcaag atcaaggtgg ccatcaacgg gttcggccgc    300
atcgccgcgc agttcctgcg ctgcgtggag gggcgcgagg actcgctgct ggagatcgtg    360
gccgtgaacg actccggcgg cgtgaagcag gccagccacc tgctcaagta cgactccacc    420
atgggacact tcaacgccga catcaagatc tcgggcgagg gcaccttctc cgtcaacggc    480
cgcgacatcc gcgtgctctc ctcccgcgac cccctggccc tgcctggggg cgagctgggc    540
gtggacctgg tgatcgaggg gacgggagtg tttgtggacc gcaaggggtg cagcaagcac    600
ctgcaggcgg gggccaagaa ggtcatcacc accgcgcggg ccaagggctc cgacgtgccc    660

```

-continued

---

```

acctacgtca tgggcgtgaa cgcggaccag tactccaact cgcacgacat catctccaac 720
gcctcctgca ccaccaactg cctggcgccc tttgtcaagg tgctcaacga cgccttcggc 780
atcgtga 787

```

```

<210> SEQ ID NO 52
<211> LENGTH: 248
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 52

```

```

Met Gln Ala Leu Cys Ser His Pro Ala Ser Leu Thr Ala Arg Ala Val
 1          5          10          15
Pro His Gly Arg Ala Ser Pro Ala Gln Arg Val Ser Ser Ala Gly Pro
      20          25          30
Ala Tyr Thr Gly Leu Ser Arg His Thr Leu Gly Cys Pro Ser Thr Pro
      35          40          45
Thr Leu Gln Ser Arg Ala Ala Val Gln Thr Arg Gly Ser Ser Ser Gly
      50          55          60
Ser Thr Thr Arg Met Thr Thr Ala Gln Arg Lys Ile Lys Val Ala
      65          70          75          80
Ile Asn Gly Phe Gly Arg Ile Gly Arg Gln Phe Leu Arg Cys Val Glu
      85          90          95
Gly Arg Glu Asp Ser Leu Leu Glu Ile Val Ala Val Asn Asp Ser Gly
      100         105         110
Gly Val Lys Gln Ala Ser His Leu Leu Lys Tyr Asp Ser Thr Met Gly
      115         120         125
Thr Phe Asn Ala Asp Ile Lys Ile Ser Gly Glu Gly Thr Phe Ser Val
      130         135         140
Asn Gly Arg Asp Ile Arg Val Val Ser Ser Arg Asp Pro Leu Ala Leu
      145         150         155         160
Pro Trp Gly Glu Leu Gly Val Asp Leu Val Ile Glu Gly Thr Gly Val
      165         170         175
Phe Val Asp Arg Lys Gly Ala Ser Lys His Leu Gln Ala Gly Ala Lys
      180         185         190
Lys Val Ile Ile Thr Ala Pro Ala Lys Gly Ser Asp Val Pro Thr Tyr
      195         200         205
Val Met Gly Val Asn Ala Asp Gln Tyr Ser Asn Ser Asp Asp Ile Ile
      210         215         220
Ser Asn Ala Ser Cys Thr Thr Asn Cys Leu Ala Pro Phe Val Lys Val
      225         230         235         240
Leu Asn Asp Arg Phe Gly Ile Val
      245

```

```

<210> SEQ ID NO 53
<211> LENGTH: 860
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

```

```

<400> SEQUENCE: 53

```

```

gatgttgaga atagtagctt gctgccttgt cgccatgcag agcgtgtgcg cgcagtcggt 60
ttcatgcaag ggggccttca cccagtcctt gcggaccccc cgatgcagca ggagccagct 120

```



-continued

---

```

cgtctgccgg gctgatggca aggccgggagc cttcatcaag accgtaaaga gcggtgctgc 180
cgctctgggt gctcctctcc tctgtcttgg gggtgccgggc gcactgacct ttgatgagct 240
gcagggcctg acctacctgc aggtgaaggg ctctggcatc gccaacacct gccccaccct 300
gtctggcggc tctccaaca tcaaggacct gaagagcggg acctactccg tcaacaagat 360
gtgctgggag cccacgtcct tcaaggtcaa ggaggaggca cagttcaaga acggcgaggc 420
cgactttgtg cccaccaagc tcgtcacgcg tctgacctac acctggagc agatctctgg 480
ccagatgaag atcgacggca gcggcggcgt ggagtccaag gaggaggatg gcatcgacta 540
tgctgcagtc accgtgcagc ttccggggcg ggagcgcgtg cccttcctct tcaccatcaa 600
ggagcttgac gccaaagggg ctgccagcgg cttcaagggc gaggttaccg tgccctccta 660
ccgtgggtcc tcttctctgg accccaaggg cgcggcgccc tccaccggtc acgacaacgc 720
cgtggccctg cccgcgcgcg gcgattccga ggagttggag aaggagaaca acaagtccac 780
caaggtctg aagggggagg ccatcttctc catcgccaag gtggacgccg ggacagggga 840
ggtgccgggc atctttgagt 860

```

&lt;210&gt; SEQ ID NO 54

&lt;211&gt; LENGTH: 275

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 54

```

Met Gln Ser Val Cys Ala Gln Ser Val Ser Cys Lys Gly Ala Phe Thr
1           5           10           15
Gln Ser Leu Arg Thr Pro Arg Cys Ser Arg Ser Gln Leu Val Cys Arg
20           25           30
Ala Asp Gly Lys Ala Gly Ala Phe Ile Lys Thr Val Lys Ser Gly Ala
35           40           45
Ala Ala Leu Ala Ala Ser Leu Leu Leu Ser Gly Gly Ala Gly Ala Leu
50           55           60
Thr Phe Asp Glu Leu Gln Gly Leu Thr Tyr Leu Gln Val Lys Gly Ser
65           70           75           80
Gly Ile Ala Asn Thr Cys Pro Thr Leu Ser Gly Gly Ser Ser Asn Ile
85           90           95
Lys Asp Leu Lys Ser Gly Thr Tyr Ser Val Asn Lys Met Cys Leu Glu
100          105          110
Pro Thr Ser Phe Lys Val Lys Glu Glu Ala Gln Phe Lys Asn Gly Glu
115          120          125
Ala Asp Phe Val Pro Thr Lys Leu Val Thr Arg Leu Thr Tyr Thr Leu
130          135          140
Asp Glu Ile Ser Gly Gln Met Lys Ile Asp Gly Ser Gly Gly Val Glu
145          150          155          160
Phe Lys Glu Glu Asp Gly Ile Asp Tyr Ala Ala Val Thr Val Gln Leu
165          170          175
Pro Gly Gly Glu Arg Val Pro Phe Leu Phe Thr Ile Lys Glu Leu Asp
180          185          190
Ala Lys Gly Thr Ala Asp Gly Phe Lys Gly Glu Phe Thr Val Pro Ser
195          200          205
Tyr Arg Gly Ser Ser Phe Leu Asp Pro Lys Gly Arg Gly Ala Ser Thr
210          215          220
Gly Tyr Asp Asn Ala Val Ala Leu Pro Ala Ala Gly Asp Ser Glu Glu

```

-continued

225	230	235	240
Leu Glu Lys Glu Asn Asn Lys Ser Thr Lys Ala Leu Lys Gly Glu Ala	245	250	255
Ile Phe Ser Ile Ala Lys Val Asp Ala Gly Thr Gly Glu Val Ala Gly	260	265	270
Ile Phe Glu	275		

<210> SEQ ID NO 55  
 <211> LENGTH: 818  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 55

ataatcgga	cccagctgca	cgcaccatca	gtgcggcagc	atgcagaccg	tcgagccag	60
ctatggcgta	ttggcgccct	ccggctccag	cgtgaccceg	ggctcgacca	gcagcaagca	120
gcacttcacc	accctcactc	ccttttccgg	cttcaggcgc	ctgaatcatg	tggatcgggc	180
ggggcaggcg	gggtctggga	gccccagac	cctgcagcag	gccgtgggca	aggccgtgcg	240
ccggtcgcgg	ggccgcacca	ccagcgcctg	gcgcgtgacc	cgcatgatgt	ttgagcggtt	300
caccgagaag	gccatcaagg	tggtcatgct	cgcgcaggag	gaggctcgcc	gtctgggcca	360
caacttcgtg	gggacggagc	aaatcctgct	ggggttgatt	ggggagtcca	caggcatcgc	420
cgccaaggtc	ctcaagtcca	tgggcgtcac	gctgaaagat	gcgcgtgtgg	aggtcgagaa	480
gatcatcggc	cgggggagcg	gctttgtggc	cgtggagatc	cccttcaccc	cccgcgccaa	540
gcgtgtgctg	gagctgtccc	tggaggaggc	tcgccagctc	ggccacaact	acattggcac	600
ggagcacatc	ctgctggggc	tgctgcgcga	gggtgagggc	gtggcctccc	gcgtgctgga	660
gaccttgggc	gcccagcccc	agaagatccg	cactcagggtg	gtacgcatgg	tgggtgagtc	720
gcaggagccc	gtgggcacca	cggtgggccc	agggtccacc	ggctccaaca	agatgcccac	780
cctggaggag	tacggcacca	acctgaccgc	ccaggccc			818

<210> SEQ ID NO 56  
 <211> LENGTH: 259  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 56

Met Gln Thr Val Ala Ala Ser Tyr Gly Val Leu Ala Pro Ser Gly Ser	1	5	10	15
Ser Val Thr Arg Gly Ser Thr Ser Ser Lys Gln His Phe Thr Thr Leu	20	25	30	
Thr Pro Phe Ser Gly Phe Arg Arg Leu Asn His Val Asp Arg Ala Gly	35	40	45	
Gln Ala Gly Ser Gly Ser Pro Gln Thr Leu Gln Gln Ala Val Gly Lys	50	55	60	
Ala Val Arg Arg Ser Arg Gly Arg Thr Thr Ser Ala Val Arg Val Thr	65	70	75	80
Arg Met Met Phe Glu Arg Phe Thr Glu Lys Ala Ile Lys Val Val Met	85	90	95	
Leu Ala Gln Glu Glu Ala Arg Arg Leu Gly His Asn Phe Val Gly Thr				

-continued

100	105	110
Glu Gln Ile Leu Leu Gly Leu Ile Gly Glu Ser Thr Gly Ile Ala Ala 115 120 125		
Lys Val Leu Lys Ser Met Gly Val Thr Leu Lys Asp Ala Arg Val Glu 130 135 140		
Val Glu Lys Ile Ile Gly Arg Gly Ser Gly Phe Val Ala Val Glu Ile 145 150 155 160		
Pro Phe Thr Pro Arg Ala Lys Arg Val Leu Glu Leu Ser Leu Glu Glu 165 170 175		
Ala Arg Gln Leu Gly His Asn Tyr Ile Gly Thr Glu His Ile Leu Leu 180 185 190		
Gly Leu Leu Arg Glu Gly Glu Gly Val Ala Ser Arg Val Leu Glu Thr 195 200 205		
Leu Gly Ala Asp Pro Gln Lys Ile Arg Thr Gln Val Val Arg Met Val 210 215 220		
Gly Glu Ser Gln Glu Pro Val Gly Thr Thr Val Gly Gly Gly Ser Thr 225 230 235 240		
Gly Ser Asn Lys Met Pro Thr Leu Glu Glu Tyr Gly Thr Asn Leu Thr 245 250 255		
Ala Gln Ala		

&lt;210&gt; SEQ ID NO 57

&lt;211&gt; LENGTH: 2357

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 57

```

ctttcttgcg ctatgacact tccagcaaaa ggtagggcgg gctgcgagac ggcttcccgg    60
cgctgcagtc aacaccgatg atgcttcgac cccccgaagc tccttcgggg ctgcatgggc    120
gctccgatgc cgctccaggg cgagcgtgt ttaaatagcc aggccccga ttgcaaagac    180
attatagcga gtaccaaag ccatattcaa acacctagat cactaccact tctacacagg    240
cactcgagc ttgtgatcgc actccgctaa gggggcgect ctctctcttc gtttcagtea    300
caaccgcaa acggcgcgcc atatcaatgc ttcttcaggc ctttctttt cttcttgctg    360
gttttgctgc caagatcagc gcctctatga cgaacgaaac ctgggataga ccacttgctc    420
actttacacc aaacaagggc tggatgaatg accccaatgg actgtggtac gacgaaaaag    480
atgccaagtg gcatctgtac tttcaataca acccgaacga tactgtctgg gggacgccat    540
tgttttgggg ccacgccacg tccgacgacc tgaccaattg ggaggaccaa ccaatagcta    600
tcgctccgaa gaggaacgac tccggagcat tctcgggttc catggtggtt gactacaaca    660
atacttccgg ctttttcaac gataccattg acccgagaca acgctgcgtg gccatatgga    720
cttacaacac accggagtcc gaggagcagt acatctcgta tagcctggac ggtggatata    780
cttttacaga gtatcagaag aacctgtgct ttgctgcaaa ttcgactcag ttccgagatc    840
cgaaggtcct ttggtacgag ccctcgcaga agtggatcat gacagcggca aagtacacag    900
actacaagat cgaattttac tcgtctgacg accttaaatc ctggaagctc gaatccgcgt    960
tcgcaaacga gggctttctc ggctaccaat acgaatgccc aggcctgata gaggtcccaa    1020
cagagcaaga tcccagcaag tcctactggg tgatgtttat ttccattaat ccaggagcac    1080
cggcaggagg ttcttttaat cagtacttcg tcggaagcct taacggaact catttcgagg    1140

```

-continued

---

```

catttgataa ccaatcaaga gtagtgtgatt ttggaagga ctactatgcc ctgcagactt 1200
tcttcaatac tgacccgacc tatgggagcg ctcttggeat tgcgtgggct tctaactggg 1260
agtattccgc attcgttctt acaaacctt ggaggctctc catgtcgtc gtgaggaaat 1320
tctctctcaa cactgagtac caggccaacc cggaaaccga actcataaac ctgaaagccg 1380
aacgatcct gaacattagc aacgtggcc cctggagccg gtttgcaacc aacaccacgt 1440
tgacgaaagc caacagctac aacgtcgatc ttctgaatag caccggtaca cttgaatttg 1500
aactggtgta tgccgtcaat accacccaaa cgatctcgaa gtcgggtgtc gcgacacctt 1560
ccctctgggt taaagccctg gaagaccctg aggagtacct cagaatgggt ttcgaggttt 1620
ctggtcctc cttcttctt gatcgcggga acagcaaagt aaaatttgtt aaggagaacc 1680
catattttac caacaggatg agcgttaaca accaaccatt caagagcga aacgacctgt 1740
cgtactacaa agtgtatggt ttgcttgatc aaaatatcct ggaactctac tccaacgatg 1800
gtgatgtcgt gtccaccaac acatacttca tgacaaccgg gaacgcactg ggctccgtga 1860
acatgacgac ggggtgtggat aacctgttct acatcgacaa attccagggt agggaagtca 1920
agtgagatct gtcgatcgac aagctcgagg cagcagcagc tcggatagta tcgacacact 1980
ctggacgctg gtcgtgtgat ggactgttgc cgccacctt gctgcctga cctgtgaata 2040
tccctgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac gcgcttttgc 2100
gagttgctag ctgcttgtgc tatttgcgaa taccacccc agcateccct tccctcgttt 2160
catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgctatccc tcagcgtgac 2220
tctgtctcct gctactgccc cctcgcacag ccttggtttg ggctccgctt gtattctcct 2280
ggtactgcaa cctgtaaacc agcaactgca tgctgatgca cgggaagtag tgggatggga 2340
acacaaatgg aaagctt 2357

```

&lt;210&gt; SEQ ID NO 58

&lt;211&gt; LENGTH: 2335

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 58

```

ctttcttgcg ctatgacact tccagcaaaa ggtagggcgg gctgagagac ggttcccgg 60
cgctgcatgc aacaccgatg atgcttcgac cccccgaagc tccttcgggg ctgcatgggc 120
gctccgatgc cgctccaggc cgagcgtgtg ttaaatagcc aggccccga ttgcaaagac 180
attatagcga gctacaaaag ccatattcaa acacctagat cactaccact tctacacagg 240
cactcgagc ttgtgatcgc actccgctaa gggggcgctt cttcctcttc gtttcagtca 300
caaccggcaa acggcgcgcc atgctgctgc aggccttctt gttcctgctg gccggcttcg 360
ccgccaagat cagcgctccc atgacgaacg agacgtccga ccgccccctg gtgcacttca 420
cccccaaaa gggctggatg aacgacccca acggcctgtg gtacgacgag aaggacgcca 480
agtggcacct gtacttccag tacaaccgca acgacaccgt ctgggggacg cccttgttct 540
ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccctc gccatcgccc 600
cgaagcgcaa cgactccggc gccttctccg gctccatggt ggtggactac aacaacacct 660
ccggcttctt caacgacacc atcgaccgac gccagcgtcg cgtggccatc tggacctaca 720
acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc tacaccttca 780
ccgagtacca gaagaacccc gtgctggcgg ccaactccac ccagttccgc gacccgaagg 840

```

-continued

---

```

tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc caggactaca 900
agatcgagat ctactcctcc gacgacctga agtcctggaa gctggagtcc gcgttcgcca 960
acgagggcct cctcggctac cagtaacgag gccccggcct gatcgaggtc cccaccgagc 1020
aggaccccaag caagtctac tgggtgatgt tcatctccat caacccccgc gccccggccg 1080
gcggctcctt caaccagtac ttcgtcggca gcttcaacgg cacccacttc gaggccttcg 1140
acaaccagtc ccgcgtgggt gacttcggca aggactacta cgccctgcag acctcttca 1200
acaccgaccc gacctacggg agcgcctgg gcatecgtg ggctccaac tgggagtact 1260
ccgccttcgt gccaccaaac ccttggcgt cctccatgtc cctcgtgcgc aagttctccc 1320
tcaaacccga gtaccaggcc aacccggaga cggagctgat caacctgaag gccgagccga 1380
tctgaacat cagcaacgcc ggcccctgga gccggttcgc caccaacacc acgttgacga 1440
aggccaacag ctacaacgtc gacctgtcca acagcaccgg cacccctggag ttcgagctgg 1500
tgtacgccgt caacaccacc cagacgatct ccaagtccgt gttcgcggac ctctccctct 1560
ggttcaaggg cctggaggac cccgaggagt acctccgcat gggcttcgag gtgtccgct 1620
cctccttctt cctggaccgc gggaacagca aggtgaagtt cgtgaaggag aaccctact 1680
tcaccaaccg catgagcgtg aacaaccagc cctcaagag cgagaacgac ctgtcctact 1740
acaagtgta cggttctgt gaccagaaca tcttgagct gtacttcaac gacggcgacg 1800
tcgtgtccac caacacctac ttcattgacca cgggaaacgc cctgggctcc gtgaacatga 1860
cgacgggggt ggacaacctg ttctacatcg acaagttcca ggtcgcgcgag gtcaagtgat 1920
taattaactc gaggcagcag cagctcggat agtatcgaca cactctggac gctggctgtg 1980
tgatggactg ttgccgccac acttgctgcc ttgacctgtg aatatccctg ccgcttttat 2040
caaacagcct cagtgtgttt gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt 2100
gtgtatttg cgaataccac cccagcacc ccttccctc gtttcatatc gcttgcaccc 2160
caaccgcaac ttatctacgc tgcctcgtc tccctcagcg ctgctcctgc tctgctcac 2220
tgcccctcgc acagccttg tttgggctcc gctgtattc tctgggtact gcaacctgta 2280
aaccagcact gcaatgctga tgcacgggaa gtagtgggat gggaacacaa atgga 2335

```

&lt;210&gt; SEQ ID NO 59

&lt;211&gt; LENGTH: 382

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Cinnamomum camphorum

&lt;400&gt; SEQUENCE: 59

```

Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val
1           5           10           15
Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu
20           25           30
Gln Leu Arg Ala Gly Asn Ala Gln Thr Ser Leu Lys Met Ile Asn Gly
35           40           45
Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Lys Leu Pro Asp Trp Ser
50           55           60
Met Leu Phe Ala Val Ile Thr Thr Ile Phe Ser Ala Ala Glu Lys Gln
65           70           75           80
Trp Thr Asn Leu Glu Trp Lys Pro Lys Pro Asn Pro Pro Gln Leu Leu
85           90           95
Asp Asp His Phe Gly Pro His Gly Leu Val Phe Arg Arg Thr Phe Ala
100          105          110

```

-continued

---

Ile Arg Ser Tyr Glu Val Gly Pro Asp Arg Ser Thr Ser Ile Val Ala  
 115 120 125

Val Met Asn His Leu Gln Glu Ala Ala Leu Asn His Ala Lys Ser Val  
 130 135 140

Gly Ile Leu Gly Asp Gly Phe Gly Thr Thr Leu Glu Met Ser Lys Arg  
 145 150 155 160

Asp Leu Ile Trp Val Val Lys Arg Thr His Val Ala Val Glu Arg Tyr  
 165 170 175

Pro Ala Trp Gly Asp Thr Val Glu Val Glu Cys Trp Val Gly Ala Ser  
 180 185 190

Gly Asn Asn Gly Arg Arg His Asp Phe Leu Val Arg Asp Cys Lys Thr  
 195 200 205

Gly Glu Ile Leu Thr Arg Cys Thr Ser Leu Ser Val Met Met Asn Thr  
 210 215 220

Arg Thr Arg Arg Leu Ser Lys Ile Pro Glu Glu Val Arg Gly Glu Ile  
 225 230 235 240

Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp Glu Glu Ile Lys  
 245 250 255

Lys Pro Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr Ile Gln Gly Gly  
 260 265 270

Leu Thr Pro Arg Trp Asn Asp Leu Asp Ile Asn Gln His Val Asn Asn  
 275 280 285

Ile Lys Tyr Val Asp Trp Ile Leu Glu Thr Val Pro Asp Ser Ile Phe  
 290 295 300

Glu Ser His His Ile Ser Ser Phe Thr Ile Glu Tyr Arg Arg Glu Cys  
 305 310 315 320

Thr Met Asp Ser Val Leu Gln Ser Leu Thr Thr Val Ser Gly Gly Ser  
 325 330 335

Ser Glu Ala Gly Leu Val Cys Glu His Leu Leu Gln Leu Glu Gly Gly  
 340 345 350

Ser Glu Val Leu Arg Ala Lys Thr Glu Trp Arg Pro Lys Leu Thr Asp  
 355 360 365

Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Ser Ser Val  
 370 375 380

&lt;210&gt; SEQ ID NO 60

&lt;211&gt; LENGTH: 1240

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Cinnamomum camphorum

&lt;400&gt; SEQUENCE: 60

```

ggcgcgccat ggccaccacc tccctggcct cgccttctg cagcatgaag gccgtgatgc   60
tggccccgca cgcccgcggc atgaagcccc gctccagcga cctgcagctg cgcgccggca   120
acgcccagac ctcccgaag atgatcaacg gcaccaagtt ctctacacc gagagcctga   180
agaagctgcc cgactggtcc atgctgttcg ccgtgatcac caccatcttc tccgccgccc   240
agaagcagtg gaccaacctg gagtgggaag ccaagcccaa cccccccag ctgctggacg   300
accacttcgg ccccccacgc ctggtgttcc gccgcacctt cgccatccgc agtaacgagg   360
tgggccccga ccgctccacc agcatcgtgg ccgtgatgaa ccacctgcag gaggccgccc   420
tgaaccacgc caagtccgtg ggcacacctg gcgacggctt cggcaccacc ctggagatgt   480
ccaagcgcca cctgatctgg gtggtgaagc gcacccacgt ggccgtggag cgctaccccc   540
cctggggcga caccgtggag gtggagtgtt ggggtggcgc ctccggcaac aacggccgcc   600
gccacgactt cctggtgcgc gactgcaaga ccggcgagat cctgaccgcg tgcacctccc   660

```

-continued

---

```

tgagcgtgat gatgaacacc cgcacccgcc gectgagcaa gatccccgag gaggtgcgcg 720
gcgagatcgg ccccccttc atcgacaacg tggccgtgaa ggacgaggag atcaagaagc 780
cccagaagct gaacgactcc accgcgact acatccaggg cggcctgacc ccccgctgga 840
acgacctgga catcaaccag cacgtgaaca acatcaagta cgtggactgg atcctggaga 900
ccgtgccccg cagcatcttc gagagccacc acatctcctc cttaccatc gactaccgcc 960
gcgagtgcac catggacagc gtgctgcagt cctgaccac cgtgagcggc ggetcctccg 1020
aggccggcct ggtgtgcgag cacctgctgc agctggaggg cggcagcgag gtgctgcgcg 1080
ccaagaccga gtggcgcgcc aagctgaccg actccttcg cggcatcagc gtgatccccg 1140
ccgagtccag cgtgatggac tacaaggacc acgacggcga ctacaaggac cagacatcg 1200
actacaagga cgacgacgac aagtgactcg agttaattaa 1240

```

```

<210> SEQ ID NO 61
<211> LENGTH: 415
<212> TYPE: PRT
<213> ORGANISM: Cuphea hookeriana

```

```

<400> SEQUENCE: 61

```

```

Met Val Ala Ala Ala Ala Ser Ser Ala Phe Phe Pro Val Pro Ala Pro
1          5          10         15
Gly Ala Ser Pro Lys Pro Gly Lys Phe Gly Asn Trp Pro Ser Ser Leu
20         25         30
Ser Pro Ser Phe Lys Pro Lys Ser Ile Pro Asn Gly Gly Phe Gln Val
35         40         45
Lys Ala Asn Asp Ser Ala His Pro Lys Ala Asn Gly Ser Ala Val Ser
50         55         60
Leu Lys Ser Gly Ser Leu Asn Thr Gln Glu Asp Thr Ser Ser Ser Pro
65         70         75         80
Pro Pro Arg Thr Phe Leu His Gln Leu Pro Asp Trp Ser Arg Leu Leu
85         90         95
Thr Ala Ile Thr Thr Val Phe Val Lys Ser Lys Arg Pro Asp Met His
100        105        110
Asp Arg Lys Ser Lys Arg Pro Asp Met Leu Val Asp Ser Phe Gly Leu
115        120        125
Glu Ser Thr Val Gln Asp Gly Leu Val Phe Arg Gln Ser Phe Ser Ile
130        135        140
Arg Ser Tyr Glu Ile Gly Thr Asp Arg Thr Ala Ser Ile Glu Thr Leu
145        150        155        160
Met Asn His Leu Gln Glu Thr Ser Leu Asn His Cys Lys Ser Thr Gly
165        170        175
Ile Leu Leu Asp Gly Phe Gly Arg Thr Leu Glu Met Cys Lys Arg Asp
180        185        190
Leu Ile Trp Val Val Ile Lys Met Gln Ile Lys Val Asn Arg Tyr Pro
195        200        205
Ala Trp Gly Asp Thr Val Glu Ile Asn Thr Arg Phe Ser Arg Leu Gly
210        215        220
Lys Ile Gly Met Gly Arg Asp Trp Leu Ile Ser Asp Cys Asn Thr Gly
225        230        235        240
Glu Ile Leu Val Arg Ala Thr Ser Ala Tyr Ala Met Met Asn Gln Lys
245        250        255
Thr Arg Arg Leu Ser Lys Leu Pro Tyr Glu Val His Gln Glu Ile Val
260        265        270

```

-continued

---

Pro Leu Phe Val Asp Ser Pro Val Ile Glu Asp Ser Asp Leu Lys Val  
 275 280 285

His Lys Phe Lys Val Lys Thr Gly Asp Ser Ile Gln Lys Gly Leu Thr  
 290 295 300

Pro Gly Trp Asn Asp Leu Asp Val Asn Gln His Val Ser Asn Val Lys  
 305 310 315 320

Tyr Ile Gly Trp Ile Leu Glu Ser Met Pro Thr Glu Val Leu Glu Thr  
 325 330 335

Gln Glu Leu Cys Ser Leu Ala Leu Glu Tyr Arg Arg Glu Cys Gly Arg  
 340 345 350

Asp Ser Val Leu Glu Ser Val Thr Ala Met Asp Pro Ser Lys Val Gly  
 355 360 365

Val Arg Ser Gln Tyr Gln His Leu Leu Arg Leu Glu Asp Gly Thr Ala  
 370 375 380

Ile Val Asn Gly Ala Thr Glu Trp Arg Pro Lys Asn Ala Gly Ala Asn  
 385 390 395 400

Gly Ala Ile Ser Thr Gly Lys Thr Ser Asn Gly Asn Ser Val Ser  
 405 410 415

&lt;210&gt; SEQ ID NO 62

&lt;211&gt; LENGTH: 1339

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Cuphea hookeriana*

&lt;400&gt; SEQUENCE: 62

```

ggcgcgccat ggtggcgccc gccgcctcca gcgccttett ccccggtccc gcccccgggc 60
cctcccccaa gcccggcaag ttcggcaact ggccctccag cctgagcccc tccttcaagc 120
ccaagtccat ccccaacggc ggcttccagg tgaaggccaa cgacagcgcc caccccaagg 180
ccaacggctc cgccgtgagc ctgaagagcg gcagcctgaa caccagggag gacacctcct 240
ccagcccccc cccccgcacc ttctgcacc agtgcccga ctggagccgc ctgctgaccg 300
ccatcaccac cgtgttcgtg aagtccaagc gcccgcacat gcacgaccgc aagtccaagc 360
gccccgacat gctggtggac agcttcggcc tggagtccac cgtgcaggac ggctggtgt 420
tccgccagtc cttctccate cgctcctacg agatcggcac cgaccgcacc gccagcatcg 480
agacctgat gaaccacctg caggagacct ccctgaacca ctgcaagagc accggcatcc 540
tgctggacgg cttcggccgc accctggaga tgtgcaagcg cgacctgatc tgggtggtga 600
tcaagatgca gatcaaggtg aaccgctacc ccgctgggg cgacaccgtg gagatcaaca 660
cccgttccag ccgcctgggc aagatcggca tgggccgcca ctggctgatc tccgactgca 720
acaccggcga gatcctggtg cgcgccacca gcgcctacgc catgatgaac cagaagacce 780
ggccctgtc caagctgccc tacgaggtgc accaggagat cgtgcccctg ttcgtggaca 840
gccccgtgat cgaggactcc gacctgaagg tgcacaagtt caaggtgaag accggcgaca 900
gcatccagaa gggcctgacc cccggctgga acgacctgga cgtgaaccag cacgtgtcca 960
acgtgaagta catcggctgg atcctggaga gcatgcccac cgagggtctg gagaccagg 1020
agctgtgctc cctggccctg gactaccgcc gcgagtgcgg ccgcgactcc gtgctggaga 1080
gcgtgaccgc catggacccc agcaaggtgg gcgtgcgctc ccagtaccag cacctgctgc 1140
gcctggagga cggcaccgcc atcgtgaacg gcgccaccga gtggcgcccc aagaacgccg 1200
gcgccaacgg cgccatctcc accggcaaga ccagcaacgg caactccgtg tccatggact 1260
acaaggacca cgacggcgac tacaaggacc acgacatcga ctacaaggac gacgacgaca 1320
agtgactcga gttaattaa 1339

```



-continued

<210> SEQ ID NO 63  
 <211> LENGTH: 382  
 <212> TYPE: PRT  
 <213> ORGANISM: Umbellularia sp.

<400> SEQUENCE: 63

Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val  
 1 5 10 15

Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu  
 20 25 30

Gln Leu Arg Ala Gly Asn Ala Pro Thr Ser Leu Lys Met Ile Asn Gly  
 35 40 45

Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Arg Leu Pro Asp Trp Ser  
 50 55 60

Met Leu Phe Ala Val Ile Thr Thr Ile Phe Ser Ala Ala Glu Lys Gln  
 65 70 75 80

Trp Thr Asn Leu Glu Trp Lys Pro Lys Pro Lys Leu Pro Gln Leu Leu  
 85 90 95

Asp Asp His Phe Gly Leu His Gly Leu Val Phe Arg Arg Thr Phe Ala  
 100 105 110

Ile Arg Ser Tyr Glu Val Gly Pro Asp Arg Ser Thr Ser Ile Leu Ala  
 115 120 125

Val Met Asn His Met Gln Glu Ala Thr Leu Asn His Ala Lys Ser Val  
 130 135 140

Gly Ile Leu Gly Asp Gly Phe Gly Thr Thr Leu Glu Met Ser Lys Arg  
 145 150 155 160

Asp Leu Met Trp Val Val Arg Arg Thr His Val Ala Val Glu Arg Tyr  
 165 170 175

Pro Thr Trp Gly Asp Thr Val Glu Val Glu Cys Trp Ile Gly Ala Ser  
 180 185 190

Gly Asn Asn Gly Met Arg Arg Asp Phe Leu Val Arg Asp Cys Lys Thr  
 195 200 205

Gly Glu Ile Leu Thr Arg Cys Thr Ser Leu Ser Val Leu Met Asn Thr  
 210 215 220

Arg Thr Arg Arg Leu Ser Thr Ile Pro Asp Glu Val Arg Gly Glu Ile  
 225 230 235 240

Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp Asp Glu Ile Lys  
 245 250 255

Lys Leu Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr Ile Gln Gly Gly  
 260 265 270

Leu Thr Pro Arg Trp Asn Asp Leu Asp Val Asn Gln His Val Asn Asn  
 275 280 285

Leu Lys Tyr Val Ala Trp Val Phe Glu Thr Val Pro Asp Ser Ile Phe  
 290 295 300

Glu Ser His His Ile Ser Ser Phe Thr Leu Glu Tyr Arg Arg Glu Cys  
 305 310 315 320

Thr Arg Asp Ser Val Leu Arg Ser Leu Thr Thr Val Ser Gly Gly Ser  
 325 330 335

Ser Glu Ala Gly Leu Val Cys Asp His Leu Leu Gln Leu Glu Gly Gly  
 340 345 350

Ser Glu Val Leu Arg Ala Arg Thr Glu Trp Arg Pro Lys Leu Thr Asp  
 355 360 365

Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Pro Arg Val  
 370 375 380

-continued

<210> SEQ ID NO 64  
 <211> LENGTH: 1240  
 <212> TYPE: DNA  
 <213> ORGANISM: Umbellularia sp.

<400> SEQUENCE: 64

```

ggcgcgccat ggccaccacc agcctggcct cgccttctg ctccatgaag gccgtgatgc      60
tggcccgcga cggcccgggc atgaagcccc gcagctccga cctgcagctg cgcgccggca    120
acgcccccac ctccctgaag atgatcaacg gcaccaagtt cagctacacc gagagcctga    180
agcgcctgcc cgactggtcc atgctggttc cgtgatcac caccatcttc agcgcgcccg    240
agaagcagtg gaccaacctg gagtggaagc ccaagcccaa gctgccccag ctgctggacg    300
accattcgg cctgcacggc ctggtgttcc gccgcacctt cgccatccgc tcttacgagg    360
tgggccccga ccgcagcacc tccatcctgg ccgtgatgaa ccacatgcag gaggccacc    420
tgaaccacgc caagagcgtg ggcatcctgg gcgacggctt cggcaccacc ctggagatgt    480
ccaagcgcga cctgatgtgg gtggtgcgcc gcaccacgt gccgtggag cgctaccca    540
cctggggcga caccgtggag gtggagtgtt ggatcggcgc cagcggcaac aacggcatgc    600
gccgcgactt cctggtgcgc gactgcaaga ccggcgagat cctgaccgcg tgcacctccc    660
tgagcgtgct gatgaacacc cgcacccgcc gctgagcac catccccgac gaggtgcgcg    720
ggcagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgacgag atcaagaagc    780
tgcagaaget gaacgactcc accgccgact acatccaggg cggcctgacc ccccgtgga    840
acgacctgga cgtgaaccag cacgtgaaca acctgaagta cgtggcctgg gtgttcgaga    900
ccgtgcccga cagcatcttc gagtcccacc acatcagctc cttaccctg gagtacgcc    960
gcgagtgcac ccgcgactcc gtgctgcgca gctgaccac cgtgagcggc ggcagctccg   1020
aggccggcct ggtgtgcgac cacctgctgc agctggaggg cggcagcagc gtgctgcgcg   1080
cccgcaccga gtggcgcccc aagctgaccg actccttcg cgccatcagc gtgatccccg   1140
ccgagccccg cgtgatggac tacaaggacc acgacggcga ctacaaggac cagacatcg   1200
actacaagga cgacgacgac aagtgactcg agttaattaa                               1240

```

<210> SEQ ID NO 65  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 65

```

ccgcccgtgt ggacgtggtg      20

```

<210> SEQ ID NO 66  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 66

```

ggtgccgggg tccaggtgt      20

```

<210> SEQ ID NO 67  
 <211> LENGTH: 20

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 67  
  
 cggccggcgg ctccttcaac 20  
  
  
 <210> SEQ ID NO 68  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 68  
  
 ggcgctcccg taggtcgggt 20  
  
  
 <210> SEQ ID NO 69  
 <211> LENGTH: 1335  
 <212> TYPE: DNA  
 <213> ORGANISM: Chlorella sorokiniana  
  
 <400> SEQUENCE: 69  
  
 cgctgcaac gcaagggcag ccacagccgc tcccaccgc cgctgaaccg acaagtgtt 60  
 gggcgctgc cgctgcctg ccgcatgtt gtgctggtga ggctgggcag tgctgccaatg 120  
 ctgattgagg cttggttcat cgggtggaag cttatgtgtg tgctgggctt gcatgccggg 180  
 caatgcat ggtggcaaga gggcggcagc acttgctgga gctgccggcg tgctccagg 240  
 tggttcaatc gggcagccca gagggatttc agatgatcgc gcgtacaggt tgagcagcag 300  
 tgtcagcaaa ggtagcagtt tgccagaatg atcggttcag ctgttaatca atgccagcaa 360  
 gagaaggggt caagtcaaa cacgggcatg ccacagcacg ggcaccgggg agtggaaatgg 420  
 caccaccaag tgtgtgcgag ccagcagcgc cgctggctg tttcagctac aacggcagga 480  
 gtcacccaac gtaaccatga gctgatcaac actgcaatca tcgggcgggc gtgatgcaag 540  
 catgcctggc gaagacacat ggtgtgcgga tgctgccggc tgctgcctgc tgcgcacgcc 600  
 gttgagtgg cagcaggctc agccatgcac tggatggcag ctgggctgcc actgcaatgt 660  
 ggtggatagg atgcaagtgg agcgaatacc aaaccctctg gctgcttctt gggttgcatg 720  
 gcatgcacc atcagcagga gcgcatgcga agggactggc cccatgcacg ccatgccaaa 780  
 ccggagcgc cagagtgtcc aactgtcac caggcccgc agctttgcag aaccatgctc 840  
 atggacgcat gtagecctga cgtcccttga cggcgtctct ctggggtgtg ggaaacgcaa 900  
 tgcagcacag gcagcagagg cggcggcagc agagcggcgg cagcagcggc gggggccacc 960  
 cttcttgccg ggtcgcgccc cagccagcgg tgatgcgctg atcccaaacg agttcacatt 1020  
 catttgcatg cctggagaag cgaggctggg gcctttgggc tggatgcagcc cgcaatggaa 1080  
 tgcgggaccg ccaggctagc agcaaaggcg cctcccctac tccgcatcga tgttccatag 1140  
 tgcattggac tgcatttggg tggggcggcc ggctgtttct ttcgtgttgc aaaacgcgcc 1200  
 agctcagcaa cctgtcccgt gggccccccg tgccgatgaa atcgtgtgca cgcgatcag 1260  
 ctgattgccc ggctcgcgaa gtaggcgccc tcctttctgc tcgccctctc tccgtcccgc 1320  
 cactagtggc gcgcc 1335

<210> SEQ ID NO 70  
 <211> LENGTH: 1146

-continued

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Umbellularia californica*

&lt;400&gt; SEQUENCE: 70

```

atggccacca ccagcctggc ctccgccttc tgcctcatga aggcctgat gctggcccgc   60
gacggcccgc gcatgaagcc ccgcagctcc gacctgcagc tgcgcgccgg caacgcccc   120
acctccctga agatgatcaa cggcaccaag ttcagctaca ccgagagcct gaagcgctg   180
cccgactggt ccatgctggt cgccgtgatc accaccatct tcagcgccgc cgagaagcag   240
tggaccaaac tggagtggaa gcccaagccc aagctgcccc agctgctgga cgaccacttc   300
ggcctgcacg gctggtggtt ccgcccgcacc ttcgcatcc gctcctacga ggtgggcccc   360
gaccgcagca cctccatcct ggccgtgatg aaccacatgc aggaggccac cctgaaccac   420
gccaaagagc tgggcatcct gggcgacggc ttcggcacca ccctggagat gtccaagcgc   480
gacctgatgt ggggtgtgcg ccgcacccac gtggccgtgg agcgctaccc cacctggggc   540
gacaccgtgg aggtggagtg ctggatcggc gccagcggca acaacggcat gcgcccgcgac   600
ttctggtgic ggcactgcaa gaccggcgag atcctgaccc gctgcacctc cctgagcgtg   660
ctgatgaaca ccgcacccc cgccctgagc accatccccg acgaggtgcg cggcgagatc   720
ggccccgcct tcatcgacaa cgtggccgtg aaggacgagc agatcaagaa gctgcagaag   780
ctgaacgact ccaccgccga ctacatccag ggcggcctga ccccccgctg gaacgacctg   840
gacgtgaacc agcacgtgaa caacctgaag tacgtggcct ggggtgttoga gaccgtgccc   900
gacagcatct tcgagtccca ccacatcagc tccttcaccc tggagtaccg ccgcgagtgc   960
accgcgact ccgtgctgcg cagcctgacc accgtgagcg gcggcagctc cgaggccggc  1020
ctggtgtgic accacctgct gcagctggag ggcggcagcg aggtgctgcg cgcccgcacc  1080
gagtggcgcc ccaagctgac cgactccttc cgcggcatca gcgtgatccc cgccgagccc  1140
cgcgctg                                           1146

```

&lt;210&gt; SEQ ID NO 71

&lt;211&gt; LENGTH: 1146

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Cinnamomum camphora*

&lt;400&gt; SEQUENCE: 71

```

atggccacca cctccctggc ctccgccttc tgcagcatga aggcctgat gctggcccgc   60
gacggcccgc gcatgaagcc ccgctccagc gacctgcagc tgcgcgccgg caacgcccag   120
acctccctga agatgatcaa cggcaccaag ttctcctaca ccgagagcct gaagaagctg   180
cccgactggt ccatgctggt cgccgtgatc accaccatct tctccgcccgc cgagaagcag   240
tggaccaaac tggagtggaa gcccaagccc aacccccccc agctgctgga cgaccacttc   300
ggcccccaag gctggtggtt ccgcccgcacc ttcgcatcc gcagctacga ggtgggcccc   360
gaccgctcca ccagcatcgt ggccgtgatg aaccacctgc aggaggccgc cctgaaccac   420
gccaaagtcc tgggcatcct gggcgacggc ttcggcacca ccctggagat gtccaagcgc   480
gacctgatct ggggtgtgaa gcgcacccac gtggccgtgg agcgctaccc cgctggggc   540
gacaccgtgg aggtggagtg ctgggtgggc gctccggca acaacggccg ccgccacgac   600
ttctggtgic ggcactgcaa gaccggcgag atcctgaccc gctgcacctc cctgagcgtg   660
atgatgaaca ccgcacccc cgccctgagc aagatccccg aggaggtgcg cggcgagatc   720
ggccccgcct tcatcgacaa cgtggccgtg aaggacgagg agatcaagaa gccccagaag   780
ctgaacgact ccaccgccga ctacatccag ggcggcctga ccccccgctg gaacgacctg   840

```

-continued

---

```

gacatcaacc agcacgtgaa caacatcaag tacgtggact ggatcctgga gaccgtgccc 900
gacagcatct tcgagagcca ccacatctcc tcttcacca tcgagtaccg ccgcgagtgc 960
accatggaca gcgtgctgca gtccctgacc accgtgagcg gcggtcctc cgaggccggc 1020
ctggtgtgcg agcacctgct gcagctggag ggcggcagcg aggtgctgcg cgccaagacc 1080
gagtggcgcc ccaagctgac cgactcctc cgcgcatca gcgtgatccc cgccgagtcc 1140
agcgtg 1146

```

```

<210> SEQ ID NO 72
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

```

```

<400> SEQUENCE: 72

```

```

atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 60
gacgacaagt ga 72

```

```

<210> SEQ ID NO 73
<211> LENGTH: 408
<212> TYPE: DNA
<213> ORGANISM: Chlorella vulgaris

```

```

<400> SEQUENCE: 73

```

```

ctcgaggcag cagcagctcg gatagtatcg acacactctg gacgctggtc gtgtgatgga 60
ctgttgccgc cacacttget gccttgacct gtgaatatcc ctgccgcttt tatcaaacag 120
cctcagtgctg tttgatcttg tgtgtacgcg cttttgcgag ttgctagctg cttgtgctat 180
ttggaatac cccccccagc atccccctcc ctcgtttcat atcgcttgca tcccaaccgc 240
aaattatcta cgctgtcctg ctatccctca gcgctgctcc tgetcctgct cactgcccct 300
cgcacagcct tggtttgggc tccgcctgta ttctcctggt actgcaacct gtaaaccagc 360
actgcaatgc tgatgcacgg gaagtagtgg gatgggaaca caaatgga 408

```

```

<210> SEQ ID NO 74
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

```

```

<400> SEQUENCE: 74

```

```

ctgggcgacg gcttcggcac 20

```

```

<210> SEQ ID NO 75
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

```

```

<400> SEQUENCE: 75

```

```

aagtcgcggc gcatgccgtt 20

```

```

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: DNA

```

-continued

---

<213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 76  
 taccccgcct ggggcgacac 20  
  
 <210> SEQ ID NO 77  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer  
  
 <400> SEQUENCE: 77  
 cttgctcagg cggcgggtgc 20  
  
 <210> SEQ ID NO 78  
 <211> LENGTH: 1317  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide  
  
 <400> SEQUENCE: 78  
 atggtggccg ccgcccctc cagcgccttc tccccctgc ccgcccccg cgctcccc 60  
 aagcccggca agttcggcaa ctggccctcc agcctgagcc cctcctcaa gcccaagtc 120  
 atccccaacg gcggcttcca ggtgaaggcc aacgacagcg cccaccccaa ggccaacggc 180  
 tccgcccgtg gcctgaagag cggcagcctg aacacccagg aggacacctc ctccagcccc 240  
 cccccccgca cttcctgca ccagctgccc gactggagcc gcctgctgac cgccatcacc 300  
 accgtgttcg tgaagtcaa gcgccccgac atgcacgacc gcaagtcaa gcgccccgac 360  
 atgctggtgg acagcttcgg cctggagtcc accgtgcagg acggcctggt gttccgcccag 420  
 tccttctcca tccgctccta cgagatcggc accgaccgca ccgcccagcat cgagaccctg 480  
 atgaaccacc tgcaggagac ctcccgtaac cactgcaaga gcaccggcat cctgctggac 540  
 ggcttcggcc gcacctgga gatgtgcaag cgcgacctga tctgggtggt gatcaaatg 600  
 cagatcaagg tgaaccgcta ccccgcctgg ggcgacaccg tggagatcaa caccgccttc 660  
 agccgcctgg gcaagatcgg catgggcccg cactggctga tctccgactg caacaccggc 720  
 gagatcctgg tgcgcgccac cagcgcctac gccatgatga accagaagac ccgcccctg 780  
 tccaagctgc cctacgaggt gcaccaggag atcgtgcccc tgttcgtgga cagccccgtg 840  
 atcgaggact ccgacctgaa ggtgcacaag ttcaaggtga agaccggcga cagcatccag 900  
 aagggcctga cccccgctg gaacgacctg gacgtgaacc agcacgtgtc caactggaag 960  
 tacatcggct ggatcctgga gagcatgccc accgaggtgc tggagaccca ggagctgtgc 1020  
 tccctggccc tggagtacc ccgagctgc ggccgagact ccgtgctgga gagcgtgacc 1080  
 gccatggacc ccagcaaggt gggcgtgcgc tcccagtacc agcacctgct gcgcctggag 1140  
 gacggcaccg ccatcgtgaa cggcgcacc gagtggcgc ccaagaacgc cggcgcacc 1200  
 ggcgccatct ccaccggcaa gaccagcaac ggcaactccg tgtccatgga ctacaaggac 1260  
 cagcagggcg actacaagga ccacgacatc gactacaagg acgacgacga caagtga 1317  
  
 <210> SEQ ID NO 79  
 <211> LENGTH: 1170

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 79

atggccaccg catccacttt ctcgcggttc aatgcccgtt gcgcgacct gcgtcgctcg      60
gcgggctccg ggccccggcg cccagcgagg cccctccccg tgcgggggcg cgcccagctg      120
cccgactgga gccgcctgct gaccgccatc accaccgtgt tcgtgaagtc caagcgcccc      180
gacatgcacg accgcaagtc caagcgcccc gacatgctgg tggacagctt cggcctggag      240
tccaccgtgc aggacggcct ggtgttccgc cagtccttct ccatccgctc ctacgagatc      300
ggcaccgacc gcaccgccag catcgagacc ctgatgaacc acctgcagga gacctcctcg      360
aaccactgca agagcaccgg catcctgctg gacggcttcg gccgcaccct ggagatgtgc      420
aagcgcgacc tgatctgggt ggtgatcaag atgcagatca aggtgaaccg ctaccccgcc      480
tggggcgaca ccgtggagat caacacccgc ttcagccgcc tgggcaagat cggcatgggc      540
cgcgactggc tgatctccga ctgcaacacc ggcgagatcc tgggtgcgcgc caccagcgcc      600
tacgccatga tgaaccagaa gacccgccgc ctgtccaagc tgcctacga ggtgcaccag      660
gagatcgtgc ccctgttctg ggacgcccc gtgatcgagg actccgacct gaaggtgcac      720
aagttcaagg tgaagaccgg cgacagcatc cagaagggcc tgacccccgg ctggaacgac      780
ctggacgtga accagcacgt gtccaacgtg aagtacatcg gctggatcct ggagagcatg      840
cccaccgagg tgctggagac ccaggagctg tgctccctgg ccctggagta ccgcccgag      900
tgcggccgcg actccgtgct ggagagcgtg accgccatgg accccagcaa ggtgggcgtg      960
cgctcccagt accagcacct gctgcgcctg gaggacggca ccgccatcgt gaacggcgcc      1020
accgagtggc gccccaaгаа cgcggcgccc aacggcgcca tctccaccgg caagaccagc      1080
aacggcaact ccgtgtccat ggactacaag gaccacgacg gcgactacaa ggaccacgac      1140
atcgactaca aggacgacga cgacaagtga      1170

```

```

<210> SEQ ID NO 80
<211> LENGTH: 1170
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 80

atggctatca agacgaacag gcagcctgtg gagaagcctc cgttcacgat cgggacgctg      60
cgcaaggcca tccccgcgca ctgtttcgag cgctcggcgc ttcgtggggcg cgcccagctg      120
cccgactgga gccgcctgct gaccgccatc accaccgtgt tcgtgaagtc caagcgcccc      180
gacatgcacg accgcaagtc caagcgcccc gacatgctgg tggacagctt cggcctggag      240
tccaccgtgc aggacggcct ggtgttccgc cagtccttct ccatccgctc ctacgagatc      300
ggcaccgacc gcaccgccag catcgagacc ctgatgaacc acctgcagga gacctcctcg      360
aaccactgca agagcaccgg catcctgctg gacggcttcg gccgcaccct ggagatgtgc      420
aagcgcgacc tgatctgggt ggtgatcaag atgcagatca aggtgaaccg ctaccccgcc      480
tggggcgaca ccgtggagat caacacccgc ttcagccgcc tgggcaagat cggcatgggc      540
cgcgactggc tgatctccga ctgcaacacc ggcgagatcc tgggtgcgcgc caccagcgcc      600
tacgccatga tgaaccagaa gacccgccgc ctgtccaagc tgcctacga ggtgcaccag      660

```

-continued

---

```

gagatcgtgc ccctgttcgt ggacagcccc gtgatcgagg actccgacct gaaggtgcac 720
aagttcaagg tgaagaccgg cgacagcatc cagaagggcc tgacccccgg ctggaacgac 780
ctggacgtga accagcacgt gtccaacgtg aagtacatcg gctggatcct ggagagcatg 840
cccaccgagg tgctggagac ccaggagctg tgctccctgg ccctggagta ccgcccgag 900
tgccggccgg actccgtgct ggagagcgtg accgccatgg accccagcaa ggtgggcgtg 960
cgctcccagt accagcacct gctgcgcctg gaggacggca ccgccatcgt gaacggcgcc 1020
accgagtggc gccccaaaga cgcggcgccc aacggcgcca tctccaccgg caagaccagc 1080
aacggcaact ccgtgtccat ggactacaag gaccacgacg gcgactacaa ggaccacgac 1140
atcgactaca aggacgacga cgacaagtga 1170

```

```

<210> SEQ ID NO 81
<211> LENGTH: 1167
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 81

```

```

atgacgttcg gggtcgacct cccggccatg ggccgcttg tctccctcc ccggcccagg 60
gtcgcgggtg gcgccagtc ggcgagtcag gttttggaga gccggcgctc ccagctgccc 120
gactggagcc gcctgtgtac cgccatcacc accgtgttcg tgaagtcaa gcgcccgcac 180
atgcacgacc gcaagtccaa gcgccccgac atgctggtgg acagcttcgg cctggagtcc 240
accgtgcagg acggcctggt gttccgccag tccttctcca tccgctccta cgagatcggc 300
accgaccgca ccgccagcat cgagaccctg atgaaccacc tgcaggagac ctccctgaac 360
cactgcaaga gcaccggcat cctgtggac ggcttcggcc gcaccctgga gatgtgcaag 420
cgcgacctga tctgggtggt gatcaagatg cagatcaagg tgaaccgcta ccccgctgg 480
ggcgacaccg tggagatcaa cccccgcttc agccgcttg gcaagatcg catgggccc 540
gactggctga tctccgactg caacaccggc gagatcctgg tgccgcccac cagcgcctac 600
gccatgatga accagaagac ccgccgctg tccaagctgc cctacgaggt gcaccaggag 660
atcgtgcccc tgttcgtgga cagccccgtg atcgaggact ccgacctgaa ggtgcacaag 720
ttcaaggtga agaccggcga cagcatccag aagggcctga cccccggctg gaacgacctg 780
gacgtgaacc agcacgtgtc caacgtgaag tacatcggct ggatcctgga gagcatgccc 840
accgaggtgc tggagaccca ggagctgtgc tccctggccc tggagtaacc ccgcgagtg 900
ggccgcgact ccgtgtgga gagcgtgacc gccatggacc ccagcaaggt gggcgtgccc 960
tcccagtacc agcacctgct gcgctggag gacggcaccg ccatcgtgaa cggcgccacc 1020
gagtggcgcc ccaagaacgc cggcgccaac ggcgccatct ccaccgcaa gaccagcaac 1080
ggcaactccg tgtccatgga ctacaaggac cacgacggcg actacaagga ccacgacatc 1140
gactacaagg acgacgacga caagtga 1167

```

```

<210> SEQ ID NO 82
<211> LENGTH: 1149
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 82

```



-continued

---

```

atgacgttcg gggctgcacct cccggccatg ggccgcggtg tctcccttcc ccggcccagg 60
gtcgcggtgc gcgcccagtc ggcgagtcag gttttggaga gcgggcgcgc ccccgactgg 120
tccatgctgt tcgcccgtgat caccaccatc ttcagcgccg ccgagaagca gtggaccaac 180
ctggagtgga agcccaagcc caagctgccc cagctgctgg acgaccactt cggcctgcac 240
ggcctggtgt tccgcccagc cttcgcctac cgctcctacg aggtgggccc cgaccgcagc 300
acctccatcc tggccgtgat gaaccacatg caggaggcca ccctgaacca cgccaagagc 360
gtgggcatcc tgggcgacgg cttcggcacc accctggaga tgtccaagcg cgacctgatg 420
tgggtggtgc gccgcaccca cgtggccgtg gagcgctacc ccacctgggg cgacaccgtg 480
gaggtggagt gctggatcgg cgccagcggc aacaacggca tgcgccgga cttcctggtg 540
cgcgactgca agaccggcga gatcctgacc cgctgcacct ccctgagcgt gctgatgaac 600
accgcacccc gccgcctgag caccatcccc gacgaggtgc gcggcgagat cggccccgcc 660
ttcatcgaca acgtggccgt gaaggacgac gagatcaaga agctgcagaa gctgaacgac 720
tccaccgccg actacatcca gggcgccctg accccccctg ggaacgacct ggacgtgaac 780
cagcacgtga acaacctgaa gtacgtggcc tgggtgttcg agaccgtgcc cgacagcatc 840
ttcaggtccc accacatcag ctccttcacc ctggagtacc gccgcgagtg caccgcgac 900
tccgtgctgc gcagcctgac caccgtgagc ggcggcagct ccgaggccgg cctggtgtgc 960
gaccacctgc tgcagctgga gggcggcagc gaggtgctgc gcgccgcac cgagtggcgc 1020
cccaagctga ccgactcctt ccgcgccatc agcgtgatcc ccgccgagcc ccgctgatg 1080
gactacaagg accacgacgg cgactacaag gaccacgaca tcgactaaa ggacgacgac 1140
gacaagtga 1149

```

&lt;210&gt; SEQ ID NO 83

&lt;211&gt; LENGTH: 1146

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 83

```

atggcttccc cggcattcac catgtcggcg tgccccgca tgactggcag gggcccctggg 60
gcacgtcgtc ccggacggcc agtcgccacc cgcctgaggg ggcgcgcccc cgactggtcc 120
atgctgttcg ccgtgatcac caccatcttc agcgcgcccg agaagcagtg gaccaacctg 180
gagtggaagc ccaagcccaa gctgccccag ctgctggacg accacttcgg cctgcacggc 240
ctggtgttcc gccgcacctt cgccatccgc tcctacgagg tgggccccga ccgcagcacc 300
tccatcctgg ccgtgatgaa ccacatgcag gaggccaccc tgaaccacgc caagagcgtg 360
ggcatcctgg gcgacggcct cggcaccacc ctggagatgt ccaagcgcga cctgatgtgg 420
gtggtgcgcc gcacccacgt ggccgtggag cgctacccca cctggggcga caccgtggag 480
gtggagtget ggatcggcgc cagcggcaac aacggcatgc gccgcgactt cctggtgcgc 540
gactgcaaga ccggcgagat cctgaccgcc tgcacctccc tgagcgtgct gatgaacacc 600
cgaccccgcc gcctgagcac catccccgac gaggtgcgcy gcgagatcgg ccccgccttc 660
atcgacaacy tggccgtgaa ggacgacgag atcaagaagc tgcagaagct gaacgactcc 720
accgcccact acatccaggg cggcctgacc ccccgtgga acgacctgga cgtgaaccag 780
cacgtgaaca acctgaagta cgtggcctgg gtgttcgaga ccgtgcccga cagcatcttc 840

```

-continued

---

```

gagtcccacc acatcagctc cttcaccctg gagtaccgcc gcgagtgcac ccgcgactcc 900
gtgtgctgca gacctgaccac cgtgagcggc ggcagctccg aggccggcct ggtgtgctgac 960
cacctgtctg agctggaggg cggcagcggg gtgtgctgctg cccgcaccga gtggcgcccc 1020
aagctgaccg actccttccg cggcatcagc gtgatccccg ccgagccccg cgtgatggac 1080
tacaaggacc acgacggcga ctacaaggac cacgacatcg actacaagga cgacgacgac 1140
aagtga 1146

```

```

<210> SEQ ID NO 84
<211> LENGTH: 1155
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 84

```

```

atggccaccg catccacttt ctccggcttc aatgcccgtt gggcgacct gcgtcctctg 60
gctggctccg ggcctccggc cccagcggag cccctcccgc tgcgctggcg cgcctccgac 120
tggtccatgc tgttcgcctg gatcaccacc atcttcagcg ccgccgagaa gcagtggacc 180
aacctggagt ggaagcccaa gcccaagctg cccagctgct tggacgacca cttcggcctg 240
cacggcctgg tgttccgcgc caccttgcgc atccgctcct acgaggtggg ccccgaccgc 300
agcacctcca tcttggcctg gatgaaccac atgcaggagg ccaccctgaa ccacgccaag 360
agcgtgggca tcctggggca cggtctggcc accaccctgg agatgtccaa gcgcgacctg 420
atgtgggtgg tgcgcccga ccaactggcc gtggagcgtt accccacctg gggcgacacc 480
gtggaggtgg agtgctggat cggcgccagc ggcaacaacg gcatgctgctg cgacttctg 540
gtgctgctg gcaagaccgg cgagatcctg acccgtgca cctccctgag cgtgctgatg 600
aacaccgcga cccgcccctt gagcaccatc cccgacgagg tgcgctggca gatcggcccc 660
gccttcctg acaactgtgc cgtgaaggac gacgagatca agaagctgca gaagctgaac 720
gactccaccg ccgactacat ccaggggcgc ctgaccccc gctggaacga cctggacgtg 780
aaccagcagc tgaacaacct gaagtacgtg gcctgggtgt tgcgacccgt gcccgacagc 840
atcttcgagt cccaccacat cagctccttc accctggagt accgctgca gtgcacccgc 900
gactcctgct tgcgagcctt gaccaccgtg agcggcggca gctccgaggc cggcctggtg 960
tgcgaccacc tctctgagct ggagggcggc agcagaggtg tgcgctccc caccgagttg 1020
cgccccaaag tgaccgactc cttccgcggc atcagcgtga tccccgcga gccccgctg 1080
atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 1140
gacgacaagt gatga 1155

```

```

<210> SEQ ID NO 85
<211> LENGTH: 1152
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 85

```

```

atggctatca agacgaacag gcagcctgtg gagaagcctc cgttcacgat cgggacgctg 60
cgcaaggcca tccccgcga ctgtttcagc cgctcggcgc ttcgtgggcg cgcctccgac 120
tggtccatgc tgttcgcctg gatcaccacc atcttcagcg ccgccgagaa gcagtggacc 180

```

-continued

---

```

aacctggagt ggaagcccaa gcccaagctg ccccagctgc tggacgacca cttcggcctg 240
cacggcctgg tgttcgcgcg caccttcgcc atccgctcct acgaggtggg ccccgaccgc 300
agcactcca tcttgccgtg gatgaaccac atgcaggagg ccacctgaa ccacgccaag 360
agcgtgggca tcttgccgca cggcttcggc accaccctgg agatgtccaa gcgcgacctg 420
atgtgggtgg tgcgcgcgac ccacgtggcc gtggagcgct accccacctg gggcgacacc 480
gtggaggtgg agtgctggat cggcgccagc ggcaacaacg gcatgcgccg cgacttctctg 540
gtgcgcgact gcaagaccgg cgagatcctg acccgctgca cctccctgag cgtgctgatg 600
aacaccgca cccgcgcgct gagcaaccac cccgacgagg tgcgcggcga gatcgcccc 660
gccttcacg acaacgtggc cgtgaaggac gacgagatca agaagctgca gaagctgaac 720
gactccaccg ccgactacat ccagggcggc ctgaccccc gctggaacga cctggacctg 780
aaccgacag tgaacaacct gaagtacgtg gcctgggtgt tcgagaccgt gcccgacagc 840
atcttcgagt cccaccacat cagctccttc accctggagt accgcccga gtgcaccgcg 900
gactccgtgc tgcgcgcgct gaccaccgtg agcggcggca gctccgaggc cggcctggtg 960
tgcgaccacc tgctgcagct ggagggcggc agcgaggtgc tgcgcgcccg caccgagtgg 1020
cgccccaaag tgaccgactc cttccgcggc atcagcgtga tccccgcga gccccgcgtg 1080
atggactaca aggaccagca cggcgactac aaggaccagc acatcgacta caaggacgac 1140
gacgacaagt ga 1152

```

&lt;210&gt; SEQ ID NO 86

&lt;211&gt; LENGTH: 1155

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 86

```

atggccaccg catccacttt ctgcgcttc aatgcccgtc gcggcgacct gcgtcgctcg 60
gcgggctccg ggccccggcg cccagcgagg cccctccccg tgcgcggggc cgccccgac 120
tggtccatgc tgttcgcgct gatcaccacc atcttctccg ccgcccagaa gcagtggacc 180
aacctggagt ggaagcccaa gcccaacccc ccccagctgc tggacgacca cttcggcctc 240
cacggcctgg tgttcgcgcg caccttcgcc atccgagct acgaggtggg ccccgaccgc 300
tccaccagca tcgtggccgt gatgaaccac ctgcaggagg ccgcccgtaa ccacgccaag 360
tccgtgggca tcttgccgca cggcttcggc accaccctgg agatgtccaa gcgcgacctg 420
atctgggtgg tgaagcgcac ccacgtggcc gtggagcgct accccgctg gggcgacacc 480
gtggaggtgg agtgctgggt gggcgccctc ggcaacaacg gccgcccga cgacttctctg 540
gtgcgcgact gcaagaccgg cgagatcctg acccgctgca cctccctgag cgtgatgatg 600
aacaccgca cccgcgcgct gagcaagatc cccgaggagg tgcgcggcga gatcgcccc 660
gccttcacg acaacgtggc cgtgaaggac gaggagatca agaagcccca gaagctgaac 720
gactccaccg ccgactacat ccagggcggc ctgaccccc gctggaacga cctggacatc 780
aaccgacag tgaacaacat caagtacgtg gactggatcc tggagaccgt gcccgacagc 840
atcttcgaga gccaccacat ctcctccttc accatcgagt accgcccga gtgcaccatg 900
gacagcgtgc tgcagtcctc gaccaccgtg agcggcggct cctccgaggc cggcctggtg 960
tgcgagcacc tgctgcagct ggagggcggc agcgaggtgc tgcgcgccaa gaccgagtgg 1020
cgccccaaag tgaccgactc cttccgcggc atcagcgtga tccccgcga gtcaccgctg 1080

```

-continued

---

```

atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 1140
gacgacaagt gatga 1155

```

```

<210> SEQ ID NO 87
<211> LENGTH: 1893
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 87
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc 180
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300
cagtcacaac ccgcaaacac tagtatggcc accgcatcca ctttctcggc gttcaatgcc 360
cgctgcggcg acctgcgctg ctcggcgggc tccgggcccc ggcgccccagc gaggccctc 420
cccgtgcgcg ggcgcgcccc cgactggctc atgctgttcg ccgtgatcac caccatcttc 480
tccgccgccc agaagcagtg gaccaacctg gagtggaagc ccaagcccaa cccccccag 540
ctgctggacg accacttcgg cccccacggc ctggtgttcc gccgcacctt cgccatccgc 600
agctacgagg tgggccccga ccgctccacc agcatcgtgg ccgtgatgaa ccacctgcag 660
gaggccgccc tgaaccacgc caagtccgtg ggcatcctgg gcgacggctt cggcaccacc 720
ctggagatgt ccaagcgcga cctgatctgg gtggtgaagc gcacccacgt ggcctggag 780
cgctaccccc cctggggcga cacctggag gtggagtct gggtgggcgc ctccgcaac 840
aacggccgcc gccacgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 900
tgcacctccc tgagcgtgat gatgaacacc cgcacccgcc gcctgagcaa gatccccgag 960
gaggtgcgcg gcgagatcgg ccccgcttc atcgacaacg tggccgtgaa ggacgaggag 1020
atcaagaagc cccagaagct gaacgactcc accgccgact acatccaggg cggcctgacc 1080
ccccgctgga acgacctgga catcaaccag cacgtgaaca acatcaagta cgtggactgg 1140
atcctggaga ccgtgcccga cagcatcttc gagagccacc acatctcttc cttcaccatc 1200
gagtaccgcc gcgagtgcac catggacagc gtgctgcagt ccctgaccac cgtgagcggc 1260
ggctcctccc aggccggcct ggtgtgcgag cacctgctgc agctggaggg cggcagcgag 1320
gtgctgcgcg ccaagaccga gtggcgcgcc aagctgaccg actccttcg cggcatcagc 1380
gtgatccccg ccgagtccag cgtgatggac tacaaggacc acgacggcga ctacaaggac 1440
cacgacatcg actacaagga cgacgacgac aagtgatgac tcgaggcagc agcagctcgg 1500
atagtatcga cacactctgg acgctggctg tgtgatggac tgttcccgc acacttctctg 1560
ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 1620
gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tggaatacc acccccagca 1680
tccccttccc tcgtttcata tcgcttgcac cccaaccgca acttatctac gctgtcctgc 1740
tatacctcag cgctgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 1800
ccgctgtgat tctcctggta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 1860
aagtagtggg atgggaacac aaatggaaaag ctt 1893

```

-continued

---

```

<210> SEQ ID NO 88
<211> LENGTH: 1887
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 88

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacgget      60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc      120
atgggcgctc cgatgccgct ccaggggcgag cgctgtttta atagccaggc ccccgattgc      180
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta      240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg ggcctcttc ctcttcgttt      300
cagtcacaac ccgcaaacac tagtatggct tccgcggcat tcaccatgtc ggcgtgcccc      360
gcgatgactg gcaggggccc tggggcacgt cgctccggac ggccagtcgc caccgcctg      420
agggggcgcg cccccgactg gtccatgctg ttcgccgtga tcaccacat cttctccgcc      480
gccgagaagc agtggaccaa cctggagtgg aagcccaagc ccaaccccc ccagctgctg      540
gacgaccact tcggccccca cggcctgggt ttccgccgca ccttcgccat ccgcagetac      600
gaggtgggccc ccgaccgctc caccagcatc gtggccgtga tgaaccaact gcaggaggcc      660
gccctgaacc acgccaagtc cgtgggcatc ctgggcgacg gcttcggcac caccctggag      720
atgtccaagc gcgacctgat ctgggtgggt aagcgcaccc acgtggcctg ggagcgetac      780
cccgcctggg gcgacaccgt ggaggtggag tgetgggtgg ggcctccgg caacaacggc      840
cgcgccacag acttctctgt gcgcgactgc aagaccggcg agatcctgac ccgctgcacc      900
tccttgagcg tgatgatgaa caccgcacc cgccgcctga gcaagatccc cgaggagggtg      960
cgcgccgaga tcggccccgc cttcatcgac aacgtggcgg tgaaggacga ggagatcaag     1020
aagccccaga agctgaacga ctccaccgcc gactacatcc agggcggcct gaacccccgc     1080
tggaacgacc tggacatcaa ccagcacgtg aacaacatca agtacgtgga ctggatcctg     1140
gagaccgtgc ccgacagcat cttcgagagc caccacatct cctccttcac catcgagtac     1200
cgccgcgagt gcaccatgga cagcgtgctg cagtccctga ccaccgtgag cggcggctcc     1260
tccgagggcg gcctggtgtg cgagcacctg ctgcagetgg agggcggcag cgagggtgctg     1320
cgcgccaaga ccgagtggcg ccccaagctg accgactcct tccgcggcat cagcgtgatc     1380
cccgcggagt ccagcgtgat ggactacaag gaccacgacg gcgactacaa ggaccacgac     1440
atcgactaca aggacgacga cgacaagtga tgactcgagg cagcagcagc tcggatagta     1500
tcgacacact ctggacgctg gtcgtgtgat ggactgttgc cgccacactt gctgccttga     1560
cctgtgaata tccttgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac     1620
gcgcttttgc gagttgctag ctgcttgtgc tatttgcgaa taccaccccc agcatcccct     1680
tccttcgttt catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgetatccc     1740
tcagcgtgct tectgctcct gctcactgcc cctcgcacag ccttggtttg ggetccgcct     1800
gtattctcct ggtactgcaa cctgtaaacc agcactgcaa tgctgatgca cgggaagtag     1860
tgggatggga acacaaatgg aaagctt                                     1887

```

```

<210> SEQ ID NO 89
<211> LENGTH: 3631
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 89

```

gaattccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac    60
gtgcttgggc gcctgcccgc tgcctgccgc atgcttgtgc tggtaggct gggcagtget    120
gccatgctga ttgagcttg gttcatcggg tggaaactta tgtgtgtgct gggcttgcac    180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc    240
tccaggtggt tcaatcgccg cagccagagg gatttcagat gatcgccgct acaggttgag    300
cagcagtgtc agcaaagga gcagtttggc agaatgatcg gttcagctgt taatcaatgc    360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg    420
gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgttcc agctacaacg    480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta    540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcy    600
cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg    660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt    720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat    780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc    840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa    900
acgcaatgca gcacagcgag cagagggcgc ggcagcagag cggcggcagc agcggcgggg    960
gccacccttc ttgcccgggc gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt   1020
cacattcatt tgcattgctg gagaagcgag gctggggcct ttgggctggt gcagcccgca   1080
atggaatgcy ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt   1140
ccatagtgca ttggactgca tttgggtggg gcggccggct gtttcttctg tgttgcaaaa   1200
cgcgccagct cagcaacctg tcccgtgggt ccccctgccc gatgaaatcg tgtgcacgcc   1260
gatcagctga ttgcccggct cgcgaagtag gcgcctcctt ttctgctcgc cctctctccg   1320
tcccgcact agtatgctgc tgcaggcctt cctgttctg ctggccggct tcgcccgcaa   1380
gatcagcgc tccatgacga acgagacgtc cgaccgccc ctggtgact tcacccccaa   1440
caagggctgy gggcgcgcca gccaccacgt gtacaagcgc ctgaccaga gcaccaacac   1500
caagtcccc agcgtgaacc agccctaccg caccggcttc cacttccagc cccccaagaa   1560
ctgatgaac gaacccaacg gccccatgat ctacaagggc atctaccacc tgttctacca   1620
gtggaacccc aagggcgcgc tgtggggcaa catcgtgtgg gccactcca ccagcaccga   1680
cctgatcaac tgggaacccc accccccgc catcttccc agcggcccct tcgacatcaa   1740
cggctgctgy tccggcagcg ccaccatcct gcccaacggc aagcccgtga tcctgtacac   1800
cggcatcgac cccaagaacc agcaggtgca gaacatcgcc gagccaaga acctgtccga   1860
cccctacctg cgcgagtgga agaagagccc cctgaacccc ctgatggccc ccgacgccgt   1920
gaacggcatc aacgcctcca gcttcgcgca ccccaccacc gcctggctgg gccaggacaa   1980
gaagtggcgc gtgateatcg gctccaagat ccaccgccc ggctggcca tcacctacac   2040
cagcaaggac ttctgaagt gggagaagtc ccccagagccc ctgcaactac acgacggcag   2100
cggcatgtgy gagtgcctcg acttcttccc cgtgaccgcg ttggcagca acggcgtgga   2160
gacctccagc ttcggcagc ccaacgagat cctgaagcac gtgctgaaga tctccctgga   2220

```

-continued

---

```

cgacaccaag cagcactact acaccatcgg cacctacgac cgcgtgaagg acaagttcgt 2280
gcccgacaac ggcttcaaga tggacggcac cgccccccgc tacgactacg gcaagtacta 2340
cgccagcaag accttcttcg actccgccaa gaaccgccgc atcctgtggg gctggaccaa 2400
cgagtcctcc agcgtggagg acgacgtgga gaagggtgg tccggcatcc agaccatccc 2460
ccgcaagate tggctggacc gcagcggcaa gcagctgatc cagtggcccg tgcgcgaggt 2520
ggagcgctg cgcaccaagc aggtgaagaa cctgcgcaac aagtgctga agtccggcag 2580
ccgctggag gtgtacggcg tgaccgccgc ccaggccgac gtggaggtgc tgttcaaggt 2640
gcgagacctg gagaaggccg acgtgatcga gccctcctgg accgaccccc agctgatctg 2700
cagcaagatg aacgtgtccg tgaagtccgg cctgggcccc ttcggcctga tgggtctggc 2760
cagcaagaac ctggaggagt acacctccgt gtacttccgc atcttcaagg cccgccagaa 2820
cagcaacaag tacgtggtgc tgatgtgctc cgaccagtcc cgcagctccc tgaaggagga 2880
caacgacaag accacctacg gcgccttcgt ggacatcaac ccccaccagc ccctgagcct 2940
gcgcgccctg atcgaccact ccgtggtgga gagcttcggc ggcaagggcc gcgcctgcat 3000
cacctcccgc gtgtacccca agctggccat cggcaagtcc agccacctgt tcgccttcaa 3060
ctacggctac cagtccgtgg acgtgtgaa cctgaacgcc tggagcatga actccgcca 3120
gatcagcatg gactacaagg accacgacgg cgactacaag gaccacgaca tcgactaaa 3180
ggacgacgac gacaagtgat taattaaccg gctcgaggca gcagcagctc ggatagtatc 3240
gacacactct ggacgctggt cgtgtgatgg actgttgcg ccacacttgc tgccttgacc 3300
tgtgaatata cctgcccgtt ttatcaaaac gcctcagtgt gttgatctt gtgtgtacgc 3360
gcttttgca gttgctagct gcttgtgcta tttgcaata ccacccccag catccccttc 3420
cctcgtttca tatcgcttgc atcccaaccg caacttatct acgtgttct gctatccctc 3480
agcgtgctc ctgctctcgc tcaactgccc tcgcacagcc ttggttggg ctcgcctgt 3540
attctcctgg tactgcaacc tgtaaaccag cactgcaatg ctgatgcacg ggaagtagtg 3600
ggatgggaac acaaatggaa agcttgagct c 3631

```

&lt;210&gt; SEQ ID NO 90

&lt;211&gt; LENGTH: 621

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 90

```

Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys
1           5           10          15
Ile Ser Ala Ser Met Thr Asn Glu Thr Ser Asp Arg Pro Leu Val His
20          25          30
Phe Thr Pro Asn Lys Gly Trp Gly Arg Ala Ser His His Val Tyr Lys
35          40          45
Arg Leu Thr Gln Ser Thr Asn Thr Lys Ser Pro Ser Val Asn Gln Pro
50          55          60
Tyr Arg Thr Gly Phe His Phe Gln Pro Pro Lys Asn Trp Met Asn Asp
65          70          75          80
Pro Asn Gly Pro Met Ile Tyr Lys Gly Ile Tyr His Leu Phe Tyr Gln
85          90          95
Trp Asn Pro Lys Gly Ala Val Trp Gly Asn Ile Val Trp Ala His Ser
100         105         110

```

-continued

---

Thr	Ser	Thr	Asp	Leu	Ile	Asn	Trp	Asp	Pro	His	Pro	Pro	Ala	Ile	Phe
		115					120					125			
Pro	Ser	Ala	Pro	Phe	Asp	Ile	Asn	Gly	Cys	Trp	Ser	Gly	Ser	Ala	Thr
	130					135					140				
Ile	Leu	Pro	Asn	Gly	Lys	Pro	Val	Ile	Leu	Tyr	Thr	Gly	Ile	Asp	Pro
145					150					155					160
Lys	Asn	Gln	Gln	Val	Gln	Asn	Ile	Ala	Glu	Pro	Lys	Asn	Leu	Ser	Asp
				165					170						175
Pro	Tyr	Leu	Arg	Glu	Trp	Lys	Lys	Ser	Pro	Leu	Asn	Pro	Leu	Met	Ala
			180					185					190		
Pro	Asp	Ala	Val	Asn	Gly	Ile	Asn	Ala	Ser	Ser	Phe	Arg	Asp	Pro	Thr
		195					200					205			
Thr	Ala	Trp	Leu	Gly	Gln	Asp	Lys	Lys	Trp	Arg	Val	Ile	Ile	Gly	Ser
	210					215					220				
Lys	Ile	His	Arg	Arg	Gly	Leu	Ala	Ile	Thr	Tyr	Thr	Ser	Lys	Asp	Phe
225					230					235					240
Leu	Lys	Trp	Glu	Lys	Ser	Pro	Glu	Pro	Leu	His	Tyr	Asp	Asp	Gly	Ser
				245					250					255	
Gly	Met	Trp	Glu	Cys	Pro	Asp	Phe	Phe	Pro	Val	Thr	Arg	Phe	Gly	Ser
			260					265						270	
Asn	Gly	Val	Glu	Thr	Ser	Ser	Phe	Gly	Glu	Pro	Asn	Glu	Ile	Leu	Lys
		275					280						285		
His	Val	Leu	Lys	Ile	Ser	Leu	Asp	Asp	Thr	Lys	His	Asp	Tyr	Tyr	Thr
	290					295					300				
Ile	Gly	Thr	Tyr	Asp	Arg	Val	Lys	Asp	Lys	Phe	Val	Pro	Asp	Asn	Gly
305					310					315					320
Phe	Lys	Met	Asp	Gly	Thr	Ala	Pro	Arg	Tyr	Asp	Tyr	Gly	Lys	Tyr	Tyr
				325					330					335	
Ala	Ser	Lys	Thr	Phe	Phe	Asp	Ser	Ala	Lys	Asn	Arg	Arg	Ile	Leu	Trp
			340					345					350		
Gly	Trp	Thr	Asn	Glu	Ser	Ser	Ser	Val	Glu	Asp	Asp	Val	Glu	Lys	Gly
		355					360					365			
Trp	Ser	Gly	Ile	Gln	Thr	Ile	Pro	Arg	Lys	Ile	Trp	Leu	Asp	Arg	Ser
	370					375					380				
Gly	Lys	Gln	Leu	Ile	Gln	Trp	Pro	Val	Arg	Glu	Val	Glu	Arg	Leu	Arg
385					390					395					400
Thr	Lys	Gln	Val	Lys	Asn	Leu	Arg	Asn	Lys	Val	Leu	Lys	Ser	Gly	Ser
				405					410					415	
Arg	Leu	Glu	Val	Tyr	Gly	Val	Thr	Ala	Ala	Gln	Ala	Asp	Val	Glu	Val
			420					425					430		
Leu	Phe	Lys	Val	Arg	Asp	Leu	Glu	Lys	Ala	Asp	Val	Ile	Glu	Pro	Ser
		435					440					445			
Trp	Thr	Asp	Pro	Gln	Leu	Ile	Cys	Ser	Lys	Met	Asn	Val	Ser	Val	Lys
	450					455					460				
Ser	Gly	Leu	Gly	Pro	Phe	Gly	Leu	Met	Val	Leu	Ala	Ser	Lys	Asn	Leu
465					470					475					480
Glu	Glu	Tyr	Thr	Ser	Val	Tyr	Phe	Arg	Ile	Phe	Lys	Ala	Arg	Gln	Asn
				485					490					495	
Ser	Asn	Lys	Tyr	Val	Val	Leu	Met	Cys	Ser	Asp	Gln	Ser	Arg	Ser	Ser
			500					505					510		
Leu	Lys	Glu	Asp	Asn	Asp	Lys	Thr	Thr	Tyr	Gly	Ala	Phe	Val	Asp	Ile
		515					520					525			
Asn	Pro	His	Gln	Pro	Leu	Ser	Leu	Arg	Ala	Leu	Ile	Asp	His	Ser	Val
						535						540			



-continued

Val Glu Ser Phe Gly Gly Lys Gly Arg Ala Cys Ile Thr Ser Arg Val  
 545 550 555 560

Tyr Pro Lys Leu Ala Ile Gly Lys Ser Ser His Leu Phe Ala Phe Asn  
 565 570 575

Tyr Gly Tyr Gln Ser Val Asp Val Leu Asn Leu Asn Ala Trp Ser Met  
 580 585 590

Asn Ser Ala Gln Ile Ser Met Asp Tyr Lys Asp His Asp Gly Asp Tyr  
 595 600 605

Lys Asp His Asp Ile Asp Tyr Lys Asp Asp Asp Asp Lys  
 610 615 620

&lt;210&gt; SEQ ID NO 91

&lt;211&gt; LENGTH: 997

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 91

```
cctgtcgatc gaagagaagg agacatgtgt acattattgg tgtgagggcg ctgaatcggc 60
cattttttaa aatgatcacg ctcacgcaa tagacgggc acataacgac gttcaaacc 120
ccgcaaaagc cgcggacaac cccatccctc cacaccccc acacaaagaa cccgccaccg 180
cttaccttgc ccacgaggta ggcctttcgt tgcgcaaaac cggcctcggg gatgaatgca 240
tgcccgttcc tgacgagcgc tgcccggggc aacacgctct tttgctcggc ctccctcaggc 300
ttgggggctt ccttgggctt ggggtgccgc atgatctcgc cgcacagag aacgcttgc 360
ggtaaaaagg agcgcgccgc tgcgcaatat atatataggc atgccaacac agcccaacct 420
cactcgggag cccgtcccac caccoccaa tgcgctgctc tgacggcata ctgctgcaga 480
agcttcatga gaatgatgcc gaacaagagg ggcacgagga cccaatcccg gacatccttg 540
tcgataatga tctcgtgagt ccccatcgtc cgcccagcgc tccggggagc ccgccgatgc 600
tcaagacgag agggccctcg accaggaggg gctggcccgg gcgggcactg gcgtcgaagg 660
tgcgcccgtc gttcgcctgc agtccctatg cacaaaacaa gtcttctgac ggggtgcgct 720
tgctcccgtg cgggcaggca acagaggtat tcaccctggt catggggaga tcggcgatcg 780
agctgggata agagatactt ctggcaagca atgacaactt gtcaggaccg gaccgtgcca 840
tatatttctc acctagcgcc gcaaaaccta acaatttggg agtcaactgtg ccaactgagtt 900
cgactggtag ctgaatggag tcgctgctcc actaaacgaa ttgtcagcac cgcagccgg 960
ccgaggaccc gagtcatagc gagggtagta gcgcgcc 997
```

&lt;210&gt; SEQ ID NO 92

&lt;211&gt; LENGTH: 753

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 92

```
actaattgca atcgtgcagt aatcatcgat atggtcacia gtagatcccc tactgacacc 60
ctctcgtaca tgtaggcaat gtcacgccc cgcctcctgt gaccgatgcc gacgtagcag 120
agcagacccc ggccgatctg ggatacagac cggccctcca cctgcgctcg aggtggaatc 180
aagtaataaa ccaatacact ttctgacacc acacagagtt gcacggacgg tggcgtacct 240
ctacgctcgc gctcttcacg cgctggagca ccgcacgcat gagcccgggt ggcttggctc 300
gggctgcaaa aatgcacaac aaacaagat cagacgctca tggatgcaca cgcgctcca 360
agcagctca gactaaatat tacagtagct cgtatctgat aagatatcga gacataccgc 420
```

-continued

---

```

tcaactcacc cgcaaactgc gccccgccag gtgatgcgca cagggcccca ccatgcgatc 480
catcgcatcg ctctctgagg gcgctatcac gtggccggag agcgttcaca gegtacgcca 540
ctgtatctgg gcggtatgcg gtccgtcaac atggagacag ataccgcac caccaccttg 600
caagctcttc catattggaa gtagaaaatt gtaattgtat catcgcacga ggggccaact 660
tgccgtcggc gagctggggc acgaacacca cctggacgtt gtcgagactc gctcgtgccc 720
tgccgccggc cgctgggtat ccagaccgtc gcc 753

```

```

<210> SEQ ID NO 93
<211> LENGTH: 1122
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 93

```

```

caacgacaac cagcaggcaa ctccgtcagc gaccaaacac gcgagtcaaa ttgttgctg 60
ttcttgctt gtctatttac tgtgatagca agactgtcgg tcagtcaata ccgggtgctg 120
cacgtcgggg tgccaagcct agcagagcac gggacggctg gtgctgtgcg ccagctcagc 180
tcgcttcgcg accaattgta ggaccggcaa agtcacaaa acatgccagc ggtgctgctc 240
aattggtcat gagctctaca aaattgtttt gtgcgtcgcg caggtatcca acggcgcggc 300
agagaaagt tgacagctct cgatttcac tcgaaaaat ggggagaatt tatgacacac 360
aagtgcgcag gcggcccagg cgcccagcat attctggcgt gacctgggccc gcccaaaaa 420
tgcttgatg cactctaaaa taattatatt tgccatgaac aagggaagag ttaccgcacc 480
cagccctaga cttgggcgcc cgagcaaggt tacgtcaagc caccttcgcc catcgcccaa 540
ctccgtatc cccgacagcc gcacgtggcc ctgcgccgaa tgaacctga atcgccatca 600
cgccacgcgt tcgccaatcg ttccgctctc tggttctcgc ggctgcgccc ttcacgtcgt 660
ggtcacgaca gtgcattcat acttccattt gcacctcggc acacactttt acgcatcgcc 720
tacccttget gcggcagctt agggtcactt tgcagccatg ggacagtgct acaccaccgt 780
cgggtgcgca agctatttca agtgaaccgt gggcgaaaa aaggaatgta cactgtctca 840
accgactcct acaattgttt accatgcaga tcagagctcg acggccatca tcgagcaggt 900
gtggggcctt ggtggcgcgg cgccggggccc cagggcgtcg caggcattga tggcactctg 960
gaactttcgc acgcgcatga gggaccccat caagagaaga gtgtgtcttt atgtcccat 1020
tcatgatgat gtatcttgtg attgtcgcag tttggcaagt ttaaccggat cgccgctcca 1080
ggtgtggcgt ggcggatttt tctaggggtg cttgagcagt cg 1122

```

```

<210> SEQ ID NO 94
<211> LENGTH: 574
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 94

```

```

ggcccagggc cctgcgatg gcccacacca gatctagcct ctcttatgcc atgcccgcct 60
cgctgcccgt cgtatcccc cgccgatccg cgcgtagggg acccgggcct gaccacgccc 120
acgaaagagc tttgctcctc aattttctgc caacagaacc gtatcaaacg ctcaacgcct 180
atcccgaaca atccgtatcc acaccaaacc gagtataccg gactggtttg cctagtcttg 240
aaggaaatga tcccgtccat gctcgggaag gggagcgggc ggaggatcct actcatctct 300
gaaatgggat tggtcggaag atgggttggg caagcacgtg ccaaacccca gcgagttgct 360
gacgagcagg ctcatccaat cccccggcga atcctccctc acgcccgcga tgcatacaag 420

```

-continued

---

tcctcccac acgccccctc ccattccattt tcgctgggc cgaacgcgag cggcgtcgag	480
geggaccaact tgcctccgag cgcctctgg gtctccacc cacagcggct ttgctgccag	540
aggaaccccc cttgccccac ctctcttgc agcc	574

<210> SEQ ID NO 95  
 <211> LENGTH: 1096  
 <212> TYPE: DNA  
 <213> ORGANISM: Prototheca moriformis

<400> SEQUENCE: 95

ccaggcaggc ggtaggggtg cggattgctt gagcgaattg gaagatataa tttttgtgg	60
tgctccctgga cgctgtttgt ggcgctcctt tttggagaag attgcgtggg ggagctttcc	120
atgtaccacg cttccttctg aaaggattct ggccgagtcc tgatgagccc aaagaaaaca	180
ctcgccttcc agtgcgggca ctctgaaaac gtcaacagat gattatacat gtcacaaaag	240
gcagccgatt aggaacggga gctctggccg ttcgtttggc tgctgggct gattgaagtg	300
atccaccctg ttcgaatgaa ggcggtcgag tcgaattatc gaccggagct gtcgggaagg	360
cgccgggggc agagtggagt gctgcggcct ggttgcgctt caaaaagacc ccggtagccc	420
aacatcacg aacgaagga atataattgc tgcatacta tacattcagt ttctatgtgg	480
cggttagaca agtctcatgg gcttctaaag gctgtccctt gaaggctact tataaaaact	540
tgctgcgcca tggcacggat cgcgcttgcg caggctgcaa ccctgcgcgc aaggcacaat	600
acacagcaaa agatactaac agaatttcta aaaacattta aatattgtt tcgaccagcc	660
aattgtggtc gtaggcacgc aaaagacttt gttttgcgcc caccgagcat ccacgctggc	720
agtcaagcca gtccgatgtg cattgcgtgg cagcatcgag gagcatcaaa aacctcgtgc	780
acgcttttct gtcaatcate atcaaccact ccaccatgta taccgatgc atcgcggtgc	840
gcagcgcgcc acgcgtccca gacccgccca aaaaccagc agcggcgaaa gcaaatcttc	900
acttgcccga aaccccgagc agcggcattc acacgtgggc gaaaaccoca cttgcctaa	960
caggcgtatg tctgctgtca cgatgcctga caacggtatt atagatatac actgattaat	1020
gtttgagtgt gtgcgagtcg cgaatcagga atgaattgct agtaggcact ccgaccgggc	1080
gggggcccag ggacca	1096

<210> SEQ ID NO 96  
 <211> LENGTH: 1075  
 <212> TYPE: DNA  
 <213> ORGANISM: Prototheca moriformis

<400> SEQUENCE: 96

ggccgacagg acgcgcgtca aagggtctgg gcgtgatgc cctggtcggc aggtcgttgc	60
tggtgctgcy ctcgtgggtc cgcaaccctg attttggcgt cttattctgg cgtggcaagc	120
gctgacgccc gcgagccggg ccggcggcga tgcggtgtct cacggctgcc gagctccaag	180
ggaggcaaga gcgccggat cagctgaagg gctttacacg caaggtagc ccgctcctgc	240
aaggctcgtt ggtggacttg aacctgtagg tcctctgctg aagttcctcc actacctcac	300
caggcccagc agaccaaagc acaggctttt caggctcgtg tcattcactc taaaacactc	360
gactacgacc tactgatgac ctagattct tcatacaaa tgctgagac acttgctcag	420
aattgaaact ccctgaaggg accaccagag gccctgagtt gttccttccc cccgtggcga	480
gctgccagcc aggctgtacc tgtgatcgag gctggcggga aaataggctt cgtgtgctca	540
ggctcatggga ggtgcaggac agctcatgaa acgccaacaa tcgcacaatt catgtcaagc	600

-continued

---

taatcagcta tttcctcttc acgagctgta attgtcccaa aattctggtc taccgggggt	660
gatocttcgt gtacgggccc ttccctcaac cctaggtatg cgcgcattgc gtegccgcgc	720
aactcgcgcg agggccgagg gtttgggacg ggcctgccg aatgcagtt gcacccggat	780
gcgcggcgcc tttcttgcga taatttatgc aatggactgc tctgcaaatt tctgggtctg	840
tcgcaacccc taggatcagc ggcgtaggat ttcgtaatca ttcgtcctga tggggagcta	900
ccgactaccc taatatcagc ccggctgcct gacgccagcg tccacttttg cgtacacatt	960
ccattcgtgc ccaagacatt tcattgtggt gcgaagcgtc ccagttacg ctcacctgtt	1020
tcccgaacct cttactgttc tgcgacaga gcgggcccac aggcgggtcg cagcc	1075

<210> SEQ ID NO 97  
 <211> LENGTH: 772  
 <212> TYPE: DNA  
 <213> ORGANISM: Prototheca moriformis

<400> SEQUENCE: 97

tcaccagcgg acaaagcacc ggtgtatcag gtccgtgtca tccactctaa agagctcgac	60
tacgacctac tgatggccct agattcttca tcaaaaacgc ctgagacact tgcccaggat	120
tgaactccc tgaagggacc accaggggcc ctgagttggt cctccccccc gtggcgagct	180
gccagccagg ctgtacctgt gatcggggct ggcgggaaaa caggcttcgt gtgctcaggt	240
tatgggaggt gcaggacagc tcattaaaacg ccaacaatcg cacaaatcat ggcaagctaa	300
tcagttatct cccattaacg agctataatt gtcccaaat tctggtctac cgggggtgat	360
ccttcgtgta cgggcccctc cctcaacctc aggtatgcgc acatgcggtc gccgcgcaac	420
gcgcgcgagg gccgagggtt tgggacgggc cgtcccgaat tgcagttgca cccggatgcg	480
tggcaccttt tttgcgataa tttatgcaat ggactgctct gcaaaatctt ggctctgctg	540
ccaaccttag gatcagcggg gtaggatttc gtaatcattc gtctgatgg ggagctaccg	600
actgccttag tatcagcccg actgcctgac gccagcgtcc acttttgtgc acacattcca	660
ttcgtgcccc agacatttca ttgtggtgcg aagcgtcccc agttacgctc acctgatccc	720
caacctcctt attgttctgt cgacagagtg ggcccagagg ccggtcgcag cc	772

<210> SEQ ID NO 98  
 <211> LENGTH: 991  
 <212> TYPE: DNA  
 <213> ORGANISM: Prototheca moriformis

<400> SEQUENCE: 98

cgaaggggtc tgcacgatt cgcgcggctc ggaggccagc gtgactgctc gcgaaaatgc	60
tctgccgtgt cgggctctgg ctggggcggc cagagatctc accgtgccac acgcaactgc	120
cgcactctgt gcccgccacc tggcgcgcac atgcgacctc ttccccgta taccctctcc	180
tcattgtatc tttccacacg agtgacgcag gtgcgcggag tggagggaaat caggacgttt	240
tcaaggtaac tgctcagacc gtaccaacag ctgcgcgccg gcaaggaaga gatcgaggca	300
gagattgccc ggctggaggc ccggataacg gagctcaaga gcaagctgct cgagtgcgac	360
cgcccagggt cacgtgtcga ctcgctatga catgtactcg acacaacatg aggaattcat	420
cgaatttgta ggaagcgggc attgttacgg gagtgggaaa gcgaaaaaac ctccctccgg	480
cagtgccttc tgccggagtc gaacgttgat agggttctcg tgacaggggtg tgacctctca	540
gccttgcatc aattaacgc tatagacatt atcagtaacc gtgaatcccc cattggatgc	600
caccgcgcgc accattgggg acctgcatta cagatctagg tgagatgaca gcgaggcaac	660

-continued

---

```

ttcgccccgc ggcccagctt gggcgccacc aatattggtc acgggaagcc acacaccgac 720
cataaatgaa tacttgtaag ctatgtcaac cgatcaatgg cgtcgaaagt gtgccacgag 780
gatccatctg gggggggcgc gtggcgcaca agcgcagtcg caatttctcg gacccatctg 840
acctaggccc agcgcgcgcg gagaaatccc eggcgggtcc tccacgcagt aaccctaag 900
agtatcgagc gccgaccatt tacaccatcg cccccgaaat ccttcgcaca ttattattat 960
cttttagatc ttggaacaga ctctgccaac c 991

```

```

<210> SEQ ID NO 99
<211> LENGTH: 1347
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 99

```

```

agagagcggg ggtggggttg tgaggtgggg ttgctgacca ggagctcgcg tcgccgagcg 60
cgactcgcac acggtccagt tcccccccc tccgcccaaa cgcaagcctc ccatcttgat 120
gcctttcccg ccacctatac tatttcttag ttcgctgtaa catccagacc gtctctgaata 180
ataacaatgc cctgtgtcaa gtgcattcct aaaaaaatc tgtccaacc aacaatccca 240
cctgaaatac caccagccct gccagtcaca ctcttccaat accatctccc tacctccacg 300
cgcaagcgac ccccatgcgc gaccaggctc gaaagtgatt tatgactga gacgagcgag 360
tggggcgcg gtgcactgcc ttttcatcac gtgccgtacg tcggcgaccg ctagggttt 420
gcacggcaac gcacggcttc gccaaaccga ccagccagga cctcgactac tctaccgca 480
attcgctca agaagtcgcc aaatgtgcca tacaccattc cttacagcac tgttcaaact 540
tgatgccaat tttgacattc ggggtgctcg ttggctgcgc ccacatcggc cgtgagtga 600
gcagggcgga tcggacacgg aggacgcggc gtcacgcccc gaacgcagcc cgtaaactca 660
catcaacacg acgtgttgcg taatcccgcc eggctgcgca tcgtgccaac ccattcgcga 720
tggatggtcg gaaaatggtg tgccaactgc cctgaggag gctctcgcga aacgggcaag 780
tccctgaaac cgaaactgtg gccttgctgt cggccacgca agcacgtgga ccctaaacac 840
caagaaaatc agtaacaag gttgacatcc tctacgggcg aattgtttgc ccaaccctc 900
atcgcaact gccattataa tgcattctagc tcggcgacaa gtttagaaaa ggcaggctgc 960
attgttccat ttcccgctgg cggcgtgggt gccatttta cgaggtttgg gctcccgggc 1020
agcgaccgag ccaggtcgag tccctctcgc ccgctgacaa tgttgcaaac cccacaagcg 1080
gctaacaaca acttgatggt acctgtacac tgccaattcc ttcttccccg gccgaggttt 1140
acagtgatg gccatggctt cgcattcagg ccgacttccc attccgactt tccagaggtt 1200
ccgcggaagc tgggggttgg ctgcctgagg cccacccttt gttccccgcg tcccgacaaa 1260
cacaattgag ttacataagg gggagccgcc cccgttcaga gtgcagaaat ctttactat 1320
atthccagc cgtcagcga atcaagt 1347

```

```

<210> SEQ ID NO 100
<211> LENGTH: 1180
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 100

```

```

gatggtgggg tgtctgcctt gggctgggtg atggaggctg gtggtgcgcg ggtttcctga 60
tgcattctat ctacgcagtg tcatggtgct cattccacac accagtacac ctttacta 120
aggatccatc cctccttccc tcttcaggac tacatggacc ccacgagcta ccgaccgggc 180

```

-continued

---

tttctcaaaa acgtcaaggt catgtttgac atgcgggacg tggaggacga cgtgcaaggt	240
gcgtccggag tgcgcgcaaa tgagcaagtc gggcaatgtg tcgggggtggg caccgggggt	300
ggagatccgc gatccccgag aaaacgccgt accaccccc gcgctatcc ctcgattgcg	360
cgcagatgtg gtgaccgaca cgggggacaa cctggcggac atggggcgcc ggacctgaa	420
gcacgccaaag tgcacacagg ggaggctcgt gcagtccccc ccctcgtacc tcaagggtct	480
ctttggtcgc gatccaaagt acgctggtgg catggcatgc ccgaaatgaa catcatgtgt	540
gatctccgat tgccaatgac cacctccacg gaccaccttg caggcgggaag cgcaatccag	600
ggcccagacc tgacgaggac ggagactcct cgtccagcgc ggggtccccc acccgacgca	660
gcagccgacc cctgctaacc cggcaacgat cggaccagca acctgtctgt agtccgatc	720
cgtgatgacg ggcaattgcc ccgctcagtc cgtttgatg actgtctatt atttgcgcg	780
agccccctcg gaacctacc ccgctcttgc aagccccctg catcggagat cctcgtgcgc	840
ccgcgatgac cccactggat tgcccacat ccttctttat cgtgtaaaat gtgattcctc	900
gggtgcaatc gactggcctt cgcttctggc cccaagaggg ctgcaacgtg cggcagcgag	960
ggcgtgaca cacccaagcc ctagggttt caacgtcggc tgccaggccg gataggggga	1020
tcgctcctt tccaccacc acctaogagg gattcgagtc ggcttcagc tcagctattc	1080
ggcgcgccc ccggccctgc agacgtcctc cagtttccga acaggtcgct ctcagaacac	1140
ctgcccgggc tgcgatacgg caggetctca aagcgtcgac	1180

&lt;210&gt; SEQ ID NO 101

&lt;211&gt; LENGTH: 1263

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 101

cgcgaggagc ggtgcgtgcg gatgcgcgc gcctgccaaag gccttttcta tgctggcct	60
gggaagtttc ctgactgaag catcttcaag atgctctctc acgaccagcg acaccaaac	120
cgtaactttt tgcctcctc gccgcagggt ccaactttcta cttgacgtc ttctccaggc	180
ggtacattgc gggactgagc gccaatcgg ccaagaacag cgtcgtcgcac ttgaggaggc	240
aggggtccgt cgaactctgc gagtgacacg ccttcgacc gactgtacta cggcctgctg	300
aagagtgggt ctgcggccc gccgtgacc gccctgtgcc cacaatcgac catctattcg	360
ctccttgcac tctggcgccg tcaattgccc cgcacttgac ggcaactggc tcgatcgagt	420
cgtattgaaa aagcacggtt tgtcctacag ggccgcggtc cgttaccaac gtggttctcg	480
ttaggttttc gtcggggcgt ggtgcgcgaa ctgtccgatg ccacccggc aaacccagc	540
aaggtcgcca gtctggttct gacgcaatag agtgcgtttt gggccagtct aaaaattcgt	600
ctggcatgac gtggctccac atcgtaccgg gagcctgect tggtaatgtg aggcaccggt	660
gccaaactcca ttatggcagg catcgagcgc gcaggtgagt acatgacctt ccgtgaattg	720
ggaaggcgag cttgtgtaac gcctgcgac gtgcccagtg ggcatcgtaa actcaaaata	780
ttttgtagaa agtgtctgat gcctggtgag gctgcgtagg gcaagggcaa gcccttggca	840
gatgggtaat gggctccgac ctcaaacag caaccccgcg tcccccttag ggcccctgag	900
gctcgtggc agggccagcg agcccggcg caaagggcgc catcccacgg tcgcccacg	960
actccacggg tcctatacct catcttgaat ggcaactaaa actatagaat atcgggcaact	1020
ggtggcgctc tggggtacag ctggccgagc gcagtgcaa accctaggtc ccgctcaag	1080
ggcgattccc gggccaatga cacgcaagca agatcacatg gcgcgggtccg cctcgggct	1140

-continued

---

```

ccacaccag gcctagttt cgcaaccat aaatatcgcc ccgataccat cataagccag 1200
caaataattt ttatcagag ttccaaacct cctcagctgt gggaaaccag cccactctga 1260
acg 1263

```

```

<210> SEQ ID NO 102
<211> LENGTH: 1400
<212> TYPE: DNA
<213> ORGANISM: Prototheca moriformis

```

```

<400> SEQUENCE: 102

```

```

ccgagcagtt catggccaag tacaaggact agagaccgga ggtcggtagg ctgaatggag 60
ctggcgctgt cgtgcgcgac gtgcaocgca tgcgatacta cgaccccaca aacgcctgcc 120
tcccattctg atgcctttcc ggccatttat actattttctc atttcgctgt aacatcttga 180
ataatagaat tgcctgtgt caagtggatt ccaagaaata ttctgtccca acaaaacaac 240
ccaacctgaa aacaacctca aataccacca gccctgcccga cctgcccagt acacttttcc 300
aataccatct ccctaccttc acgcycaagc ggcacccatg cgcgaccagg ctgaaagga 360
tttcacgact caggacgagc gagtggcggc gcgaccgctt gcctgttctg cactgcccgt 420
acgtcggcga ccgctagagc tttgcctggc aacccccggc ttcgtcaacc cggccagcca 480
ggatctcgac cactctaccg cgaaatcgcc tcaagaagtc gccaaaagtg ccgtacacca 540
tgcttcgcag cgctgttcaa acttgatgcc aatcttgaca atcaggttgc tcgttgctg 600
cgtccacatc gccctgtgatt gcagcaggcg gggatcggac acggaggacg cggcgtcacg 660
ccggaacgcg agcccgtaac tctacatcaa cgcgatatgt tgcgtaatcc cgcccggctg 720
cgcatgtga caaccatct gcgatggatg gtcggaaaat ggtgtgcca ctgccctgag 780
ggactctctc gcgaaacggg cacgtccctg tatccgaaac tgtggcatgg cctgtctgac 840
cacgcaagca cgtggacctt aacaccacga aaataagtaa aaaaggttga catcctctac 900
gagcgaattg tttgctcgac ccttcactgc aactgtcat tataatgcat ctagctcggc 960
gacaagtta aaaaaggcag gctgcattat tccattttgc cgtggcggca tgggtgcca 1020
ttttatgagg tttggctctt tgggcagcga ccgagccagg ttgagtcctt ctgcccctc 1080
gacaacgttc caaagcccat aagtggctaa taaacaactt gatggtacct gtacactgac 1140
agttccttct tccccggcgg aggtttacac gtgatggcca tggcttcgcg tttcaggctg 1200
acttccatt ccgactttcc agagggtccg cggacgcggg ggtttggctg cgtgaggccc 1260
acccttggtt ccccgcgtcc cgacaaacac aattgcgtta cataaggggg aagccgcccc 1320
ccgttcagag tgcaaacatc tttcattata tttttcagtc gtcagcgaat tcaagtatgt 1380
cgctgacagg catgaaggcc 1400

```

```

<210> SEQ ID NO 103
<211> LENGTH: 3681
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide

```

```

<400> SEQUENCE: 103

```

```

gccctttgtc atcgttggca tgctttttgc gtatgtacca tatgttgaat gtataatag 60
aacggttgac cgtctgagat gcgagctttg ggtcttgcca aatgcgtggc cgcacggctc 120
cctcgcaccc agccccgagg cgtcgcgcac ctggcgagga gcagaccac gccaaagaa 180
tctagtccag catgtaacaa catcaggcaa tgtgacgttt tcggttcccg atttctctgc 240

```

-continued

---

cgctctttga	cggcaggcac	gggcgagcaa	ccggcggcgc	tcgcgtcagg	cacgatggat	300
gcggcgctgc	ccacctgtca	atgtacccca	ccagtctgtc	gatcgctaca	agcaaccttg	360
tgctccacat	tcccacttgc	agacagtcta	gtcgattttg	ccaagctgga	tgtgaggatt	420
ggccatatct	tggaggccaa	gattcaccgc	gatgctgatg	ggtacgtacg	cgagccaggc	480
aggcagctgc	gttgactttc	tgattggcac	aaagctttgg	ctactctcaa	taccaaccac	540
gtgccccttc	tgcacacctg	cttcctttcg	atgaccactc	gccacgcattg	tcgcagtctg	600
tacgtcgagc	agatcgacct	cggcgaggag	gggggcccctc	gcaccatcgt	gagtggcctg	660
gtccggcacg	tgacctgga	ggacctgttc	ggccggcggg	tggtggtgct	ggccaacctc	720
aagcctcgga	gcatgcgcgg	ggtcaaatcg	gctgggatgc	tgtctcgcgc	cgccaacgcg	780
gatcacaccg	cggtggagcc	gctgcgggtc	ccggacgccg	ccgtgacggg	ggagcgggtc	840
tggcgggggg	acgaggcact	cctgtccacg	gagcctgcca	cacccaatca	ggtaaggaca	900
cgttattggt	gcgatgggtg	cgaatgcgtg	gtctgacctg	ctgtgggtat	gtgttggtggg	960
attggaaac	gaatgagggc	cgttcaggat	tgagcccttg	gccccacct	gctcatcctc	1020
tcacgcccgc	aggtccagaa	gaagaaaatc	tgggaggcag	tacagccgct	gctgagagtg	1080
aacgcccagg	ggatcgctac	tgtggcagga	gaggetatgg	tgaccagtgc	ggggccactg	1140
accgcccaca	cgctggttga	cgcccgatt	tctgacgcg	agcgaactgat	tcttgacctt	1200
tgagaagcca	ccacagcacc	atcttcattg	ttcatccttg	atctcagtac	gacttctcac	1260
catttcagta	ctgtaggacc	ccccaaatag	tgtgatcacg	ctcgcaagge	acctgtgtga	1320
tcacggggaa	ggcgaaatc	ctttctgtcg	ctatgacct	tccagcaaaa	ggtaggggcg	1380
gctgcgagac	ggcttcccgg	cgctgcatgc	aacaccgatg	atgcttcgac	ccccgaagc	1440
tccttcgggg	ctgcatgggc	gctccgatgc	cgctccaggg	cgagcgetgt	ttaaatagcc	1500
aggccccga	ttgcaagac	attatagcga	gctaccaaag	ccatattcaa	acacctagat	1560
cactaccact	tctacacagg	ccactcgagc	ttgtgatcgc	actccgctaa	gggggcgcct	1620
cttctcttc	gtttcagtea	caaccogcaa	acggcgcgcc	atgctgctgc	aggccttct	1680
gttctgctg	gcccgttctg	ccgccaagat	cagcgcctcc	atgacgaacg	agcgtccga	1740
ccgccccctg	gtgcacttca	cccccaacaa	gggctggatg	aacgacccca	acggcctgtg	1800
gtacgacgag	aaggacgcca	agtggcacct	gtacttccag	tacaacccga	acgacaccgt	1860
ctgggggacg	cccttgttct	ggggccacgc	cacgtccgac	gacctgacca	actgggagga	1920
ccagcccac	gcccctgcgc	cgaagcgcaa	cgactccggc	gccttctccg	gctccatggt	1980
ggtggactac	aacaacacct	ccggettctt	caacgacacc	atcgaccgcg	gccagcctg	2040
cgtggccac	tggacctaca	acaccccgga	gtccgaggag	cagtacatct	cctacagcct	2100
ggacggcggc	tacaccttca	ccgagtacca	gaagaacccc	gtgctggcgg	ccaactccac	2160
ccagttccgc	gacccgaagg	tcttctggta	cgagccctcc	cagaagtgga	tcatgaccgc	2220
ggccaagtcc	caggactaca	agatcgagat	ctactcctcc	gacgacctga	agtccctgaa	2280
gctggagtcc	gcttccgcca	acgagggctt	cctcggttac	cagtacgagt	gccccggcct	2340
gatcgaggtc	cccaccgagc	aggaccccag	caagtcctac	tgggtgatgt	tcacttccat	2400
caaccccggc	gccccggcgc	gcggetcctt	caaccagtac	ttcgtcggca	gcttcaacgg	2460
caccaccttc	gaggccttcg	acaaccagtc	ccgcgtggtg	gacttcggca	aggactacta	2520
cgccctgcag	accttcttca	acaccgaccc	gacctacggg	agcgcctctg	gcatcgctgt	2580
ggcctccaac	tgggagtact	ccgccttctg	gcccaccaac	ccctggcgtc	cctccatgtc	2640



-continued

---

```

cctcgtgccc aagttctccc tcaacaccga gtaccaggcc aaccgggaga cggagctgat 2700
caacctgaag gccgagccga tcctgaacat cagcaacgcc ggcccctgga gccggttcgc 2760
caccaacacc acgttgacga aggccaacag ctacaacgtc gacctgtcca acagcaccgg 2820
cacccctggag ttcgagctgg tgtacgccgt caacaccacc cagacgatct ccaagtccgt 2880
ggtcgcggac ctctccctct ggttcaaggc cctggaggac cccgaggagt acctccgcat 2940
gggcttcgag gtgtcccgct cctccttctt cctggaccgc gggaacagca aggtgaagtt 3000
cgtgaaggag aaccctact tcaccaaccg catgagcgtg aacaaccagc ccttcaagag 3060
cgagaacgac ctgtcctact acaagggtga cggcttgctg gaccagaaca tcctggagct 3120
gtacttcaac gacggcgacg tcgtgtccac caacacctac ttcattgacca ccgggaacgc 3180
cctgggctcc gtgaacatga cgacgggggt ggacaacctg ttctacatcg acaagtcca 3240
ggtgcccgag gtcaagtgat taattaactc gaggcagcag cagctcggat agtatcgaca 3300
cactctggac getggtcgtg tgatggactg ttgccgccac acttgetgce ttgacctgtg 3360
aatatccctg ccgcttttat caaacagcct cagtgtgttt gatcttgtgt gtaocgcctt 3420
ttgcgagttg ctagctgctt gtgctatttg cgaataccac ccccgacatc cccttccctc 3480
gtttcatatc gcttgcattc caaccgcaac ttatctacgc tgcctgcta tcctcagcg 3540
ctgctcctgc tcctgtctac tgcccctcgc acagccttgg tttgggctcc gcctgtattc 3600
tcctggtact gcaacctgta aaccagcact gcaatgctga tgcacgggaa gtagtgggat 3660
gggaacacaa atggaaagct t 3681

```

&lt;210&gt; SEQ ID NO 104

&lt;211&gt; LENGTH: 3850

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 104

```

ttggccccc ctttccagct ccgcatctgc tggcgtccgc cgcgagacgt gacatcgcac 60
gtcgcgggga gccccagctt gatcacttgg cagggggccg tgctctacaa ataccaggcc 120
ccgcggcggc cagttcgcac atccaatacc tgccgagcca tcttgctac actttttatc 180
gactcctcta ctctgttccg gagagcgcct ggtccaggct tggaattcgc cgaattcagc 240
tcgatcagtc gcttctgcaa ctgatctcgg ccgttcgcag actgcctttt ctcagcttgt 300
caggtagcga gttgtgtttt tatatttatt cgatttcac tgtgttgcac gtcttgttcg 360
tgctgtgcgt tctttctggg ccgcgctgtc gggtcgcatg ggctagctgt actcatgtta 420
gtcatgccgg tccgaacctg ttcgaggaag gcccacact gagcgtgccc tctttctaca 480
ccccctgtgc agaaattaga tagaaagcag aattccttcc ttgcgctatg acacttccag 540
caaaaggtag gccgggctgc gagacggctt cccggcctcg catgcaacac cgatgatgct 600
tcgaccccc gaagctcctt cggggctgca tgggcgctcc gatgccgctc cagggcgagc 660
gctgtttaaa tagccaggcc cccgattgca aagacattat agcgagctac caaagccata 720
ttcaaacacc tagataacta ccacttttac acaggccact cgagcttctg atcgactcc 780
gctaaggggg cgctcttcc tcttcgttcc agtcacaacc cgcaaacggc gcgccatgct 840
gctgcaggcc tcctgttcc tgctggccgg ctccgccgcc aagatcagcg cctccatgac 900
gaacgagacg tccgaccgcc ccctggtgca ctccacccc aacaagggct ggatgaacga 960

```

-continued

---

ccccaacggc	ctgtggtacg	acgagaagga	cgccaagtgg	cacctgtact	tccagtacaa	1020
cccgaacgac	accgtctggg	ggacgcccctt	gttctggggc	cacgccacgt	ccgacgacct	1080
gaccaactgg	gaggaccagc	ccatcgccat	cgccccgaag	cgcaacgact	ccggcgccct	1140
ctccggctcc	atgggtggtg	actacaacaa	cacctccggc	ttcttcaacg	acaccatcga	1200
cccgcgccag	cgctgcgtgg	ccatctggac	ctacaacacc	ccggagtccg	aggagcagta	1260
catctcttac	agcctggacg	gcggtacac	cttaccgag	taccagaaga	accccgtgct	1320
ggccgccaac	tccaccagct	tccgcgaccc	gaaggtcttc	tggtacgagc	cctcccagaa	1380
gtggatcatg	accgcggcca	agtcccagga	ctacaagatc	gagatctact	cctccgacga	1440
cctgaagtcc	tggaaagtgg	agtcccgctt	cgccaacgag	ggcttctctg	gctaccagta	1500
cgagtgcgcc	ggcctgatcg	aggctccccc	cgagcaggac	cccagcaagt	cctactgggt	1560
gatgttcate	tccatcaacc	ccggcgcccc	ggccggcggc	tccttcaacc	agtacttctg	1620
cggcagcttc	aacggcacc	acttcgaggc	cttcgacaac	cagtcgcccg	tggtggactt	1680
cggaaggac	tactacgccc	tgcagacctt	cttcaacacc	gacccgacct	acgggagcgc	1740
cctgggcatc	gcgtgggct	ccaactggga	gtactccgcc	ttcgtgcccc	ccaaccctg	1800
gcgctctcc	atgtccctcg	tgcgcaagtt	ctccctcaac	accgagtacc	aggccaacc	1860
ggagacggag	ctgatcaacc	tgaaggccga	gccgatcctg	aacatcagca	acgccggccc	1920
ctggagccgg	ttcgccacca	acaccagctt	gacgaaggcc	aacagctaca	acgtcgacct	1980
gtccaacagc	accggcacc	tggagtccga	gctgggttac	gccgtcaaca	ccaccagac	2040
gatctccaag	tccgtgttcg	cggacctctc	cctctggttc	aagggcctgg	aggaccccga	2100
ggagtacctc	cgcattgggt	tgcaggtgtc	cgcgtcctcc	ttcttctctg	accgcgggaa	2160
cagcaagggt	aagtctctga	aggagaacc	ctacttcacc	aaccgcatga	gcgtgaacaa	2220
ccagcccttc	aagagcgaga	acgacctgtc	ctactacaag	gtgtacggct	tgctggacca	2280
gaacatcctg	gagctgtact	tcaacgacgg	cgacgtctgt	tccaccaaca	cctacttcat	2340
gaccaccggg	aacgcccctg	gctccgtgaa	catgacgacg	ggggtggaca	acctgttcta	2400
catcgacaag	ttccaggtgc	gcgaggtcaa	gtgattaatt	aactcgaggc	agcagcagct	2460
cggatagtat	cgacacactc	tggacgctgg	tcggtgatg	gactggtgcc	gccacacttg	2520
ctgccttgac	ctgtgaatat	ccctgccgct	tttatcaaac	agcctcagtg	tgtttgatct	2580
tgtgtgtacg	cgcttttgcg	agttgctagc	tgcttgtgct	atttgcgaat	accaccccga	2640
gcatcccctt	ccctcgttct	atatcgcttg	catcccacc	gcaacttate	tacgctgtcc	2700
tgctatccct	cagcgtgctg	cctgctcctg	ctcaactgcc	ctcgcacagc	cttggtttgg	2760
gctccgcctg	tattctcctg	gtactgcaac	ctgtaaacca	gcactgcaat	gctgatgcac	2820
gggaagtagt	gggatgggaa	cacaaatgga	ccgacacgcc	cccggcccag	gtccagttct	2880
cctgggtctt	ccagaggccc	gtcgcacatg	aaagtggcag	agattggcgc	ctgattcgat	2940
ttggatccaa	ggatctccaa	tccgtgatgg	ggactgagtg	cccaactacc	acccttgcac	3000
tatcgtcctc	gcactattta	ttcccacctt	ctgctcgccc	tgccgggcca	ttgcgggctg	3060
ttctgcccct	gacgtatcaa	tttcgcccct	gctggcgcga	ggattcttca	ttctaataag	3120
aactcactcc	cgccagctct	gtacttttcc	tgccggggccc	ctgcatggct	tgttcccaat	3180
gcttgcctga	tgcagcggcg	ccattgcccc	cggcgctgcc	gcatccatgt	gaagaaacac	3240
ggaagagtgc	gaagactgga	agtgaattaa	gagtataaga	agaggtaacca	agggattctc	3300
aggtgctctt	aggaacggct	tttccttcgc	caagagaaac	tgctactgct	cgtgtcgcca	3360

-continued

---

cgggtggtcaa ggcgccccat ctgcatcca ccaggccccat ccgcggaactc gcatcagcc	3420
tgctggatcc ggactgccga cctgaccgct cgcattccacc attacaaccc tccaattgga	3480
caccactccc acgtcctaaa gttcaccatg caagctgac gatcgattc gccgatgcac	3540
tcgctgcca cagaggtgtg cgcttcggac tagcgtgcag ggcgccccgag gccaccagca	3600
tgaccgatg gaagcgggca cggccgctgc tccaggctgc tggtctgctc agaccatag	3660
caacctccgc tgcgtcccta aatgtcacac agagcgtctt tgatgggtac ggatgggaga	3720
gaatctgatt gggcattgct ggtgcagtgc aggaagatgg caagtgcaca gtcagtcag	3780
ctgtacaaa tggtgctcgc tagtattgac tcgtatagtg catagatca tgcattgctg	3840
ttacttgc	3850

&lt;210&gt; SEQ ID NO 105

&lt;211&gt; LENGTH: 3108

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 105

tttgccccg ctttccagct ccgcatctgc tggcgtccgc cgcgagacgt gacatcgac	60
gtcgcgggga ggcaccgctt gatcaattgg cagggggccg tgctctaaa ataccaggcc	120
ccgcgccggt cagttccgac atccaatacc tgcgagcca tcttgcctac actttttatc	180
gactcctcta ctctgttcgc gagagcctc ggtccaggct tggaaatcgc cgaattcagc	240
tcgatcagtc gcttctgcaa ctgatctcgc ccgcttcgag actgcctttt ctccagctgt	300
caggtagcga gttgtgtgtt tatatttatt cgatttcac tgtgttgcac gtctgttcg	360
tgctgtgctt tctttctggg ccgctctgc gggctgcag ggctagctgt actcatgta	420
gtcatgccg tccgacctg ttcgaggaag gcccacact gagcgtgcc tctttctaca	480
ccccctgtgc agaaattaga tagaaagcaa tgctgctgca ggccttctc ttcctgctg	540
ccggcttgc cccaagatc agcgcctcca tgacgaacga gacgtccgac cgcacctg	600
tgcaactcac cccaacaag ggctggatga acgacccca cggcctgtg tacgacgaga	660
aggacgcca gtggcacctg tacttccagt acaaccgaa cgacaccgctc tgggggacgc	720
ccttgttctg gggccacgcc acgtccgacg acctgaccaa ctgggaggac cagcccatcg	780
ccatcgcgcc gaagcgaac gactccggcg ccttctccgg ctccatggtg gtggactaca	840
acaacacctc cggtctctc aacgacacca tcgacccgcg ccagcgtgc gtggccatct	900
ggacctaca cccccggag tccgaggagc agtacatctc ctacagctg gacggcggct	960
acacctcac cgagtaccag aagaacccc tgctggccgc caactccacc cagttccgcg	1020
accgaaggt cttctggtac gagccctccc agaagtggat catgaccgcg gccaaagtc	1080
aggactaca gatcgagatc tactcctccg acgacctgaa gtctggaag ctggagtcg	1140
cgctcgcaca cgaggcttc ctcgctacc agtacagtg ccccgccctg atcgaggtcc	1200
ccaccgagca ggacccagc aagtcctact gggatgatgt catctccatc aaccccgcg	1260
ccccggccg cggtccttc aaccagtact tcgtcggcag cttcaacggc acccacttcg	1320
aggccttca caaccagtc cgcgtgggtg acttcggcaa ggactactac gccctgcaga	1380
ccttcttcaa caccgaccg acctacggga gcgcctggg catcgcgtgg gcctccaact	1440
gggagtactc cgcctctgt cccaccaacc cctggcctc ctccatgtcc ctctgctgca	1500
agttctccct caacaccgag taccaggcca acccgagac ggagctgatc aacctgaag	1560

-continued

---

```

ccgagccgat cctgaacatc agcaacgccg gcccttgag cgggttcgcc accaacacca 1620
cgttgacgaa ggccaacagc tacaacgtcg acctgtccaa cagcaccggc acctggagt 1680
tcgagctggt gtacgccgtc aacaccacc agacgatctc caagtccgtg ttcgaggacc 1740
tctccctctg gttcaagggc ctggaggacc cggaggagta cctccgatg ggcttcgagg 1800
tgtcccgctc ctcttcttc ctggaccgcg ggaacagcaa ggtgaagttc gtgaaggaga 1860
accctactt caccaaccgc atgagcgtga acaaccagcc cttcaagagc gagaacgacc 1920
tgtcctacta caaggtgtac ggcttctgg accagaacat cctggagctg tacttcaacg 1980
acggcgacgt cgtgtccacc aacacctact tcatgaccac cgggaacgcc ctgggtccg 2040
tgaacatgac gacgggggtg gacaacctgt tctacatcga caagttccag gtgcgagagg 2100
tcaagtgacc gacacgcccc cggcccaggt ccagttctcc tgggtcttcc agaggcccgt 2160
cgccatgtaa agtggcagag attggcgctt gattcgattt ggatccaagg atctccaatc 2220
ggtgatgggg actgagtgcc caactaccac ccttgcaacta tctcctctgc actatttatt 2280
cccactctt getcgcctg cggggcgatt gggggcgtt ctgccctga cgtatcaatt 2340
tcgcccctgc tggcgcgagg attcttcatt ctaataagaa ctactcccg ccagctctgt 2400
acttttctg cggggcccct gcattgcttg ttcccaatgc ttgctcgatc gacggcgccc 2460
attgcccacg gcgctgccgc atccatgtga agaaacacgg aagagtgcga agactggaag 2520
tgaattaaga gtataagaag aggtaccaag ggattctcag gtgctcttag gaacggcttt 2580
tccttcgcca agagaaactg ctactgctcg tctcgcacg gtggtcaagc cgccccatct 2640
gcatccacc aggccatcc gcggactcgc gatcagcctg ctggatccgg actgccgacc 2700
tgaccgctcg catccaccat tacaaccctc caattggaca ccactcccac gtctctaaagt 2760
tcaccatgca agctgatcga tgcattcgc cgatgcactc gcctgccaca gaggtgtgag 2820
cttggacta gcgtgcaggc gccccaggc caccagcatg caccgatgga agcgggcacg 2880
gcccgtgctc caggctcgtg gctcgtcag acctagatga acctccgctg cgtccctaaa 2940
tgtcacacag agcgtctttg atgggtacgg atgggagaga atctgattgg gcattgctgg 3000
tgcagtgcag gaagtggca agtgcacagt cagtcatgct gtacaaactg gtgcctcgta 3060
gtattgactc gtatagtgca tagtatcatg catggctggt acttgcaa 3108

```

&lt;210&gt; SEQ ID NO 106

&lt;211&gt; LENGTH: 559

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Chlorella luteoviridis*

&lt;400&gt; SEQUENCE: 106

```

tgttgaagaa tgagccggcg acctatagga agtggcgtgg ttaaggaatt tccgaagcc 60
caagcgaag caagttttaa aaatagcgt atttgtcact ttttatggac ccgaaccgg 120
gtgatctaac cgtgaccagg atgaagcttg ggtaacacca agtgaaggtc cgaactcttc 180
gatctttaa aatcgtgaga tgagtgcgg ttagtaggtg aaatgccaat cgaactcgga 240
gctagctggt tctcccga atgtgttag gcgcagcgt gaatgacaaa acaaatagta 300
cgggtgtagg gtaaaact gtttcggtgc gggctgcgaa agcggtagca aatcgtggca 360
aactcagaat actacgcttg tataccatc atcagtgaga ctgtggggga taagctccat 420
agtcaagagg gaaacagccc agatcaccag ttaaggcccc aaaatgacag ctaagtggca 480
aaggaggtga aagtgcagaa acaaccagga ggtttgccca gaagcagcca tcctttaaag 540
agtgcgtaat agctcactg 559

```

-continued

---

```

<210> SEQ ID NO 107
<211> LENGTH: 1841
<212> TYPE: DNA
<213> ORGANISM: Cuphea palustris

<400> SEQUENCE: 107
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct    60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc    120
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc    180
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta    240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgectcttc ctcttcgttt    300
cagtcacaac ccgcaaacac tagtatggcc accgcatcca cttctcggc gttcaatgcc    360
cgctgcggcg acctgcgctg ctcggcgggc tccgggcccc ggcgcccagc gaggccctc    420
cccgtgcgcg ggcgcgccag catgctgctg tcggcgggtga ccacggctct cggcgtggcc    480
gagaagcagt ggcccagctt ggaccgcaag tccaagcgcc ccgacatgct ggtcgagccc    540
ctgggcgtgg accgcatcgt ctacgacggc gtgagcttcc gccagtcggt ctccatccgc    600
agctacgaga tcggcgccga ccgcaccgcc tcgatcgaga cgctgatgaa catgttccag    660
gagacctccc tgaaccactg caagatcctc ggctgctga acgacggctt cggccgcacg    720
cccgagatgt gcaagcgcca cctgatctgg gtcgtgacca agatgcagat cgaggtgaac    780
cgctaccccc cgtggggcga caccatcgag gtcaaacagt gggtgagcgc ctggggcaag    840
cacggcatgg gccgcgactg gctgatctcc gactgccaca ccggcgagat cctgatccgc    900
gcgacgagcg tctgggcatg gatgaaccag aagaccgcc gcctgtcgaa gatcccctac    960
gaggtgcgcc aggagatcga gccccagttc gtcgactccg cccccgtgat cgtggacgac   1020
cgcaagtccc acaagctgga cctgaagcgc ggcgacagca tctgcaacgg cctgaccccc   1080
cgctggacgc acctggacgt gaaccagcac gtcaacaacg tgaagtacat cggctggatc   1140
ctgcagtcgg tccccaccga ggtgttcgag acgcaggagc tgtcgggctt gaccctggag   1200
taccgccgcg agtgccgccc cgactccgtg ctggagagcg tcacggccat ggaccctcgc   1260
aaggaggggc accgctccct gtaccagcac ctgctgcgcc tggaggacgg cgcggacatc   1320
gtgaagggcc gcaccgagtg gcgccccaaq aacgcggcgc ccaagggcgc catcctgacg   1380
ggcaagacca gcaacggcaa ctcgatctcc tgactcgagt taattaactc gaggcagcag   1440
cagctcggat agtatcgaca cactctggac gctggctcgt tgatggactg ttgccgccac   1500
acttctgctc ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt   1560
gatcttctgt gtacgcgctt ttgcgagttg ctactgctt gtgctatttg cgaataccac   1620
ccccagcatc ccttccctc gtttcaatc gcttgcctcc caaccgcaac ttatctacgc   1680
tgtctgcta tccctcagcg ctgctcctgc tctgctcac tgcccctcgc acagccttgg   1740
tttgggctcc gcctgtatc tcctggtact gcaacctgta aaccagcact gcaatgctga   1800
tgcaagggaa gtagtgggat gggaacacaa atggaagct t                               1841

```

```

<210> SEQ ID NO 108
<211> LENGTH: 1010
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
polynucleotide

```

-continued

&lt;400&gt; SEQUENCE: 108

```

ggcgcgccca gctgcccagc tggagcatgc tgctggccgc gateaccacc ctgttcctgg    60
cggccgagaa gcagtggatg atgctggact ggaagcccaa cgcccccgac atgctggtgg    120
accctctcgg cctgggccgc ttcgtgcagg acggcctggt gttccgcaac aacttcagca    180
tccgagccta cgagatcggc gcggaccgca ccgccagcat cgagaccctg atgaaccacc    240
tgcaggagac ccccctgaac cacgtgaaga gcgtgggcct gctggaggac ggcctgggca    300
gcacccgcga gatgagcctg cgcaacctga tctgggtggt gaccaagatg caggtggcgg    360
tggaccgcta ccccactcgg ggcgacgagg tgcaggtgag cagctgggcg accgccatcg    420
gcaagaacgg catgcgccgc gactggatcg tgaccgactt ccgcaccggc gagaccctgc    480
tgcgcgccac cagcgtgtgg gtgatgatga acaagctgac ccgccgcatc agcaagatcc    540
ccgaggaggt gtggcacgag atcggcccca gcttcatega cgcgcccccc ctgcccaccg    600
tggaggacga cggccgcaag ctgacccgct tcgacgagag cagcggccgac ttcacccgca    660
agggcctgac cccccctcgg agcgcactgg acatcaacca gcacgtgaac aacgtgaagt    720
acatcggtcty gctgctggag agcgcgcccc ccgagatcca cgagagccac gagatcgcca    780
gcctgacctt ggagtaccgc cgcgagtgcg gccgcgacag cgtgctgaac agcggcacca    840
aggtgagcga cagcagccag ctgggcaaga gcgccgtgga gtgcaaccac ctggtgcgcc    900
tgcagaacgg cggcgagatc gtgaaggggc gcaccgtgtg gcgccccaaag cggcccctgt    960
acaacgacgg cgcctgtggt gacgtgcccg ccaagaccag ctgactcgag    1010

```

&lt;210&gt; SEQ ID NO 109

&lt;211&gt; LENGTH: 5472

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 109

```

ggtacccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct    60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc    120
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc    180
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta    240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt    300
cagtcacaac ccgcaaaactc tagaataatc atgctgctgc aggccctcct gttcctgctg    360
gccggtctcg ccgccaagat cagcgcctcc atgacgaacg agacgtccga ccgccccctg    420
gtgcacttca cccccaaaaa gggctggatg aacgacccca acggcctgtg gtacgacgag    480
aaggacgcca agtggcacct gtacttccag tacaaccgca acgacaccgt ctgggggacg    540
cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc    600
gccatcgccc cgaagcgcga cgactccggc gccttctccg gctccatggt ggtggactac    660
aacaacacct ccggttctct caacgacacc atcgaccgc gccagcgtg cgtggccatc    720
tggacctaca acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc    780
tacacettca ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc    840
gaccgcaagg tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc    900
caggactaca agatcgagat ctactcctcc gacgacctga agtcctggaa gctggagtcc    960
gcgttcgcca acgagggtct cctcggctac cagtacgagt gccccggcct gatcgaggtc    1020

```

-continued

---

cccaccgagc	aggaccccag	caagtcctac	tgggtgatgt	tcatttccat	caaccccggc	1080
gccccggccg	geggctcctt	caaccagtac	ttcgtcggca	gcttcaacgg	cacccacttc	1140
gaggccttcg	acaaccagtc	cgcgctgggtg	gacttcggca	aggactacta	cgccttcgag	1200
acctttctca	acaccgacce	gacctacggg	agcgcctcgg	gcacgcgctg	ggcctccaac	1260
tgggagtact	cgccttcctg	gcccaccaac	ccctggcgct	cctccatgtc	cctcgtgcgc	1320
aagttctccc	tcaacaccga	gtaccaggcc	aaccgggaga	cggagctgat	caacctgaag	1380
gccgagccga	tctgaacat	cagcaacgcc	ggccctgga	gccggttcgc	caccaacacc	1440
acgttgacga	aggccaacag	ctacaacgtc	gacctgtcca	acagcacccg	caccctggag	1500
ttcagctcgg	tgtacgccgt	caacaccacc	cagacgatct	ccaagtccgt	ggtcgcggac	1560
ctctccctct	ggttcaaggg	cctggaggac	cccaggaggt	acctccgcat	gggcttcgag	1620
gtgtccgcgt	cctccttctt	cctggaccgc	gggaacagca	aggtgaagtt	cgtgaaggag	1680
aaccctact	tcaccaaccg	catgagcgtg	aacaaccagc	ccttcaagag	cgagaacgac	1740
ctgtcctact	acaaggtgta	cggcttgctg	gaccagaaca	tcctggagct	gtacttcaac	1800
gacggcgacg	tcgtgtccac	caacacctac	ttcatgacca	ccgggaacgc	cctgggctcc	1860
gtgaacatga	cgacgggggt	ggacaacctg	ttctacatcg	acaagttcca	ggtgcgcgag	1920
gtcaagtgac	aattggcagc	agcagctcgg	atagtatcga	cacactctgg	acgctggctg	1980
tgtgatggac	tgttgccgcc	acacttgctg	ccttgacctg	tgaatatccc	tgccgctttt	2040
atcaaacagc	ctcagtgtgt	ttgatcttgt	gtgtacgcgc	ttttgcgagt	tgetagetgc	2100
ttgtgctatt	tgcaataacc	accccagca	tccccttccc	tcgtttcata	tcgcttgcac	2160
cccaaccgca	acttatctac	gctgtcctgc	tatccctcag	cgctgctcct	gctcctgctc	2220
actgccctc	gcacagcctt	ggtttgggct	ccgcctgtat	tctcctggta	ctgcaacctg	2280
taaacaccga	ctgcaatgct	gatgcacggg	aagtagtggg	atgggaacac	aatggaggga	2340
tcccgcgtct	cgaacagagc	gcgcagagga	acgctgaagg	tctcgcctct	gtcgcacctc	2400
agcgcggcat	acaccacaat	aaccaacctga	cgaatgcgct	tggttctctg	tccattagcg	2460
aagcgtccgg	ttcacacacg	tgccaagtgg	gcgaggtggc	aggtgacaat	gatcggtgga	2520
gctgatggtc	gaaacgttca	cagcctaggg	atatacaatt	ccgcctgcaa	cgcaagggca	2580
gccacagccg	ctcccacccg	ccgctgaacc	gacacgtgct	tgggcgcctg	ccgcctgcct	2640
gccgcatgct	tgtgctgggt	aggctgggca	gtgctgccat	gctgattgag	gcttggttca	2700
tcgggtggaa	gcttatgtgt	gtgctgggct	tgcatgccgg	gcaatgcgca	tgggtggcaag	2760
agggcggcag	cacttgctgg	agctgcccg	gtgcctccag	gtggttcaat	cgcggcagcc	2820
agagggattt	cagatgatcg	cgcgtacagg	ttgagcagca	gtgtcagcaa	aggtagcagt	2880
ttgccagaat	gatcggttca	gctgttaatc	aatgccagca	agagaagggg	tcaagtgcaa	2940
acacgggcat	gccacagcac	gggcacccgg	gagtggaaatg	gcaccaccaa	gtgtgtgcga	3000
gccagcatcg	cgcctgggct	gtttcagcta	caacggcagg	agtcaccaa	cgtaacctatg	3060
agctgatcaa	cactgcaatc	atcgggcccgg	cgtgatgcaa	gcacgcctgg	cgaagacaca	3120
tgggtgtcgg	atgctgccgg	ctgctgcctg	ctgcgcacgc	cgttgagttg	gcagcaggct	3180
cagccatgca	ctggatggca	gctgggctgc	cactgcaatg	tgggtgatag	gatgcaagtg	3240
gagcgaatac	caaacctctt	ggctgcttgc	tgggttgcat	ggcatcgcac	catcagcagg	3300
agcgcctcgc	aagggactgg	ccccatgcac	gccatgcca	accggagcgc	accgagtgct	3360
cacactgtca	ccaggcccgc	aagctttgca	gaacctgct	catggacgca	tgtagcgtg	3420

-continued

---

```

acgtcccttg acggcgctcc tctcgggtgt gggaaacgca atgcagcaca ggcagcagag 3480
gcggcgccag cagagcggcg gcagcagcgg cggggggccac ccttcttgcg gggtcgcgcc 3540
ccagccagcg gtgatgcgct gatcccaaac gagttccat tcatttgcat gctgggagaa 3600
gcgaggctgg ggcttttggg ctggtgcagc cgcgaatgga atgcgggacc gccaggctag 3660
cagcaaaggc gcctccccta ctccgcatcg atgttccata gtgcattgga ctgcatttgg 3720
gtggggcgcg cggctgtttc tttcgtgttg caaaacgcgc cagctcagca acctgtcccg 3780
tgggtccccc gtgccgatga aatcgtgtgc acgccgatca gctgattgcc cggtcgcga 3840
agtaggcgcc ctccctttctg ctgcgccctc ctccgtcccg ccactagtat ggccaccgca 3900
tccactttct cggcgttcaa tgcccgtgc ggcgacctgc gtcgctcggc gggctccggg 3960
ccccggcgcc cagcagggcc cctcccctg cgcgggcgcg ccagctgcc cgactggagc 4020
cgctgctga ccgccatcac caccgtgttc gtgaagtcca agcgcgccga catgcacgac 4080
cgcaagtcca agcgcgccga catgctggtg gacagcttcg gcctggagtc caccgtgcag 4140
gacggcctgg tgttccgccca gtccttctcc atccgctcct acgagatcgg caccgaccgc 4200
accgccagca tcgagaccct gatgaaccac ctgcaggaga cctccctgaa ccactgcaag 4260
agcaccggca tcctgctgga cggttcggc cgcaccctgg agatgtgcaa gcgcgacctg 4320
atctgggtgg tgatcaagat gcagatcaag gtgaaccgct accccgcctg gggcgacacc 4380
gtggagatca acaccgcctt cagccgcctg ggcaagatcg gcatgggccc cgactggctg 4440
atctccgact gcaacaccgg cgagatcctg gtgcgcgcca ccagcgcta cgccatgatg 4500
aaccagaaga cccgccgcct gtccaagctg ccctacgagg tgcaccagga gatcgtgccc 4560
ctgttcgtgg acagcccctg gatcgaggac tccgacctga aggtgcacaa gttcaagggtg 4620
aagaccggcg acagcatcca gaagggcctg acccccggct ggaacgacct ggaagtgaac 4680
cagcacgtgt ccaacgtgaa gtacatcggc tggatcctgg agagcatgcc caccgaggtg 4740
ctggagacc caggagctgt ctcccctg ccctggagtacc gccgcgagtg cggcccgcgac 4800
tccgtgctgg agagcgtgac cgccatggac ccagcaagg tgggcgtgcg ctcccagtac 4860
cagcacctgc tgcgcctgga ggacggcacc gccatcgtga acggcgccac cgagtggcgc 4920
cccaagaacg cggcgcccaa cggcgccatc tccaccgca agaccagcaa cggcaactcc 4980
gtgtccatgg actacaagga ccacgacggc gactacaagg accacgacat cgactacaag 5040
gacgacgacg acaagtgact cgaggcagca gcagctcggg tagtatcgac aactctgga 5100
cgctggctgt gtgatggact gttgccgcca cacttctgct cttgacctgt gaatatccct 5160
gccgctttta tcaaacagcc tcagtgtgtt tgatcttctg tgtacgcgct tttgcgagtt 5220
gctagctgct tgtgctatct gcgaatacca ccccagcat ccccttcct cgtttcatat 5280
cgcttgcac ccaaccgcaa cttatctacg ctgtcctgct atccctcagc gctgctcctg 5340
ctcctgctca ctgcccctcg cacagccttg gtttgggctc cgctgtatt ctccctgtac 5400
tgcaacctgt aaaccagcac tgcaatgctg atgcacggga agtagtggga tgggaacaca 5460
aatgaaagc tt 5472

```

&lt;210&gt; SEQ ID NO 110

&lt;211&gt; LENGTH: 5451

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide



-continued

&lt;400&gt; SEQUENCE: 110

```

ggtagcccttt cttgcgctat gacacttcca gcaaaaagta gggcgggctg cgagacgget      60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc      120
atgggcgctc cgatgcgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc      180
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta      240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt      300
cagtcacaac ccgcaaaactc tagaataatca atgctgctgc aggccttctt gttcctgctg      360
gccggcttcg ccgccaagat cagcgcctcc atgacgaacg agacgtccga ccgcccctg      420
gtgcacttca ccccaacaa gggctggatg aacgacccca acggcctgtg gtaoacagag      480
aaggacgcca agtggcacct gtacttccag tacaaccga acgacaccgt ctgggggacg      540
cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc      600
gccatcgccc cgaagcga cgaactcggc gccttctccg gctccatggt ggtggactac      660
aacaacacct ccggcttctt caacgacacc atcgaccgc gccagcgtg cgtggccatc      720
tggacctaca acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc      780
tacaccttca ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc      840
gaccogaagg tcttctgcta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc      900
caggactaca agatcgagat ctactcctcc gacgacctga agtccctgaa getggagtec      960
gcgttcgcca acgagggctt cctcggctac cagtacgagt gcccggcct gatcgaggtc     1020
cccaccgagc aggaccccg caagtccctac tgggtgatgt tcatctccat caaccccggc     1080
gccccggcgc gcggctcctt caaccagtac ttcgtcggca gcttcaacgg cacccacttc     1140
gaggccttcg acaaccagtc ccgctgggtg gacttcggca aggactacta cgcctcagag     1200
accttcttca acaccgaccc gacctacggg agcgcctcgg gcctcgcgtg ggctccaac     1260
tgggagtact ccgcctctgt gcccaccaac cctggcgct cctccatgtc cctcgtcgcg     1320
aagtctctcc tcaacaccga gtaccaggcc aaccgggaga cggagctgat caacctgaag     1380
gccgagccga tcttgaacat cagcaacgcc ggcccctgga gccggttcgc caccaacacc     1440
acgttgacga aggccaacag ctacaacgtc gacctgtcca acagcaccgg cacccctggag     1500
ttcgagctgg tgtacccgt caacaccacc cagacgatct ccaagtccgt gttcgcggac     1560
ctctccctct ggttcaaggg cctggaggac cccgaggagt acctccgcat gggcttcgag     1620
gtgtcccgct cctccttctt cctggaccgc gggaacagca aggtgaagtt cgtgaaggag     1680
aaccctact tcaccaaccg catgagcgtg aacaaccagc cctcaagag cgagaacgac     1740
ctgtcctact acaagggtga cggcttctg gaccagaaca tcttgagct gtacttcaac     1800
gacggcgacg tcgtgtccac caacacctac ttcattgacca ccgggaacgc cctgggctcc     1860
gtgaacatga cgacgggggt ggacaacctg ttctacatcg acaagttcca ggtgcccagag     1920
gtcaagtgac aattggcagc agcagctcgg atagtatcga cacactctgg acgctggctg     1980
tgtgatggac tgttgccgcc acacttctgt ccttgacctg tgaatatccc tgccgctttt     2040
atcaaacagc ctcaagtgtt ttgatcttgt gtgtacgcgc ttttcgagat tgctagctgc     2100
ttgtctatt tgcaataacc acccccgaca tccccttccc tcgtttcata tcgcttgcat     2160
cccaaccgca acttatctac gctgtcctgc tatccctcag cgetgctcct gctcctgctc     2220
actgcccctc gcacagcctt ggtttgggct ccgcctgtat tctcctgta ctgcaacctg     2280
taaaccagca ctgcaatgct gatgcacggg aagtagtggg atgggaacac aaatggagga     2340

```

-continued

---

tcccgcgtct	cgaacagagc	gcgcagagga	acgctgaagg	tctcgcctct	gtcgcacctc	2400
agcgcggcat	acaccacaat	aaccacctga	cgaatgcgct	tggttcttcg	tccattagcg	2460
aagcgtcccg	ttcacacacg	tgccacgttg	gcgaggtggc	aggtagacaat	gatcggtgga	2520
gctgatggtc	gaaacgttca	cagcctaggg	atatcgaatt	ccgcctgcaa	cgcaagggca	2580
gccacagccg	ctcccaccgg	ccgctgaacc	gacacgtgct	tggggcctcg	ccgcctgcct	2640
gccgcagctc	tgtgctgggt	aggctgggca	gtgctgccat	gctgattgag	gcttggttca	2700
tccgggtgaa	gcttatgtgt	gtgctgggct	tgcatgccgg	gcaatgcgca	tggtggcaag	2760
agggcggcag	cacttgctgg	agctgcccg	gtgcctccag	gtggttcaat	cgcggcagcc	2820
agagggattt	agatgatcgc	cgcgtacagg	ttgagcagca	gtgtcagcaa	aggtagcagt	2880
ttgcagaat	gatcggttca	gctgttaatc	aatgccagca	agagaagggg	tcaagtgcaa	2940
acacgggcat	gccacagcac	gggcaccggg	gagtggaaatg	gcaccaccaa	gtgtgtgcga	3000
gccagcatcg	ccgcctggct	gtttcagcta	caacggcagg	agtcaccaa	cgtaaccatg	3060
agctgatcaa	cactgcaatc	atcgggcccgg	cgtgatgcaa	gcatgcctgg	cgaagacaca	3120
tggtgtgccc	atgctgccgg	ctgctgcctg	ctgcgcacgc	cgttgagttg	gcagcaggct	3180
cagccatgca	ctggatggca	gctgggctgc	cactgcaatg	tggtggatag	gatgcaagtg	3240
gagcgaatac	caaacctctc	ggctgcttgc	tgggttgcat	ggcatcgcac	catcagcagg	3300
agcgcagcgc	aagggactgg	ccccatgcac	gccatgcaa	accggagcgc	accgagtgtc	3360
cacactgtca	ccaggcccgc	aagctttgca	gaaccatgct	catggacgca	tgtagcctg	3420
acgtcccttg	acggcgctcc	tctcgggtgt	gggaaacgca	atgcagcaca	ggcagcagag	3480
gcggcggcag	cagagcggcg	gcagcagcgg	cgggggcccac	ccttcttgcg	gggtcgcgcc	3540
ccagccagcg	gtgatgcgct	gatccccaac	gagttcacat	tcatttgcat	gcctggagaa	3600
gcgaggctgg	ggcctttggg	ctggtgcagc	ccgcaatgga	atgcgggacc	gccaggtag	3660
cagcaaaagg	gcctccccta	ctccgcatcg	atgttccata	gtgcattgga	ctgcatttgg	3720
gtggggcggc	cggctgtttc	tttctgtgtg	caaaacgcgc	cagctcagca	acctgtcccg	3780
tgggtccccc	gtgccgatga	aatcgtgtgc	acgccgatca	gctgattgcc	cggctcgcga	3840
agtaggcgcc	ctcctttctg	ctcgcctctc	ctccgtcccg	ccactagtat	gacgttcggg	3900
gtcgcctccc	cggccatggg	ccgcgggtgc	tcccttcccc	ggcccagggt	cgcgggtgcgc	3960
gcccagtcgg	cgagtcaggt	tttgagagc	gggcgcgccc	ccgactggtc	catgctgttc	4020
gccgtgatca	ccaccatctt	cagcgcgcc	gagaagcagt	ggaccaacct	ggagtggaaag	4080
cccaagccca	agctgcccc	gctgctggac	gaccacttcg	gcctgcacgg	cctggtgttc	4140
cgcgcacctc	tgcctatccg	ctcctacgag	gtgggccccg	accgcagcac	ctccatcctg	4200
gccgtgatga	accacatgca	ggaggccacc	ctgaaccacg	ccaagagcgt	gggcacccctg	4260
ggcgacggct	tgggcaccac	cctggagatg	tccaagcgcg	acctgatgtg	ggtggtgcgc	4320
cgcaccacag	tggccgtgga	gcgctacccc	acctggggcg	acaccgtgga	ggtggagtgc	4380
tggatcggcg	ccagcggcaa	caacggcatg	cgccgcgact	tcctggtgcg	cgactgcaag	4440
accggcgaga	tcctgaccgg	ctgcaacctc	ctgagcgtgc	tgatgaaac	ccgcacccgc	4500
cgcctgagca	ccatccccga	cgaggtgcgc	ggcgagatcg	gccccgcctt	catcgacaac	4560
gtggccgtga	aggacgacga	gatcaagaag	ctgcagaagc	tgaacgactc	caccgcccgc	4620
tacatccagg	ggggcctgac	ccccgcctgg	aacgacctgg	acgtgaaaca	gcacgtgaac	4680
aaactgaagt	acgtggcctg	ggtgttcgag	accgtgcccc	acagcatctt	cgagtcccac	4740

-continued

---

```

cacatcagct ccttcaccct ggagtaccgc cgcgagtgca cccgcgactc cgtgctgcgc 4800
agcctgacca ccgtgagcgg cggcagctcc gaggccggcc tgggtgtgca ccacctgctg 4860
cagctggagg gccgcagcga ggtgtgcgc gcccgaccg agtggcgccc caagctgacc 4920
gactccttcc gggcatcag cgtgatcccc gccgagcccc gcgtgatgga ctacaaggac 4980
cacgacggcg actacaagga ccacgacatc gactacaagg acgacgacga caagtgactc 5040
gaggcagcag cagctcggat agtatcgaca cactctggac gctggctgtg tgatggactg 5100
ttgcgcccac acttgcctgc ttgacctgtg aatatccctg ccgcttttat caaacagcct 5160
cagtggtgtt gatcttgtgt gtacgcgctt ttgcgagtgt ctagctgctt gtgctatttg 5220
cgaataccac ccccagcacc cccttccttc gtttcataac gcttgcaccc caaccgcaac 5280
ttatctacgc tgcctgcta tccctcagcg ctgctcctgc tcctgctcac tgcccctcgc 5340
acagccttgg tttgggctcc gcctgtattc tcctggctact gcaacctgta aaccagcact 5400
gcaatgctga tgcacgggaa gtagtgggat gggaaacaaa atggaaagct t 5451

```

&lt;210&gt; SEQ ID NO 111

&lt;211&gt; LENGTH: 5454

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 111

```

ggtacccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc 180
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgatcact accacttcta 240
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300
cagtcacaac ccgcaaaactc tagaataatca atgctgctgc aggccttccct gttcctgctg 360
gccggtctcg ccgcccaagat cagcgcctcc atgacgaacg agacgtccga ccgcccctg 420
gtgcacttca cccccaaaaa gggctggatg aacgacccca acggcctgtg gtacgacgag 480
aaggacgcca agtggcacct gtacttccag tacaaccgga acgacaccgt ctgggggacg 540
cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc 600
gccatcgccc cgaagcgcga cgactccggc gccttctccg gctccatggt ggtggactac 660
aacaacacct ccggcttctt caacgacacc atcgaccgc gccagcctg cgtggccatc 720
tggacctaca acaccocgga gtccgaggag cagtacatct cctacagcct ggacggcggc 780
tacaccttca ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc 840
gaccgaaggt tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc 900
caggactaca agatcgagat ctactcctcc gacgacctga agtccctggaa gctggagctc 960
gcgttcgccca acgagggctt cctcggctac cagtacgagt gccccggcct gatcgaggtc 1020
cccaccgagc aggaccccag caagtccctac tgggtgatgt tcactccat caaccocggc 1080
gccccggcgg gcggctcctt caaccagtac ttcgtcggca gcttcaacgg caccacttc 1140
gaggccttcc acaaccagtc ccgcgtgggt gacttcggca aggactacta cgcctgcag 1200
accttcttca acaccgaccc gacctacggg agcgcctcgg gcctcgcgtg ggccctcaac 1260
tgggagtaet ccgcctctgt gccaccaaac ccctggcgtc cctccatgct cctcgtgcgc 1320
aagttctccc tcaacaccga gtaccaggcc aaccocggaga cggagctgat caacctgaag 1380

```

-continued

---

gccgagccga	tctgaacat	cagcaacgcc	ggccctgga	gccggttcgc	caccaacacc	1440
acgttgacga	aggccaacag	ctacaacgtc	gacctgtcca	acagcacccg	caccctggag	1500
ttcgagctgg	tgtacgccgt	caacaaccacc	cagacgatct	ccaagtccgt	gttcgaggac	1560
ctctccctct	ggttcaaggg	cctggaggac	cccaggaggt	acctccgcat	gggcttcgag	1620
gtgtccgcgt	cctccttctt	cctggaccgc	gggaacagca	aggtgaagtt	cgtgaaggag	1680
aaccctact	tcaccaaccg	catgagcgtg	aacaaccagc	ccttcaagag	cgagaacgac	1740
ctgtcctact	acaaggtgta	cggtctgtg	gaccagaaca	tcctggagct	gtacttcaac	1800
gacggcgacg	tcgtgtccac	caacacctac	ttcatgacca	ccgggaacgc	cctgggctcc	1860
gtgaacatga	cgacgggggt	ggacaacctg	ttctacatcg	acaagttcca	ggtgcgagag	1920
gtcaagtgac	aattggcagc	agcagctcgg	atagtatcga	cacactctgg	acgtcggctg	1980
tgtgatggac	tgttgcgcc	acacttgctg	ccttgacctg	tgaatatccc	tgccgctttt	2040
atcaaacagc	ctcagtgtgt	ttgatcttgt	gtgtacgcgc	ttttgagagt	tgetagetgc	2100
ttgtgctatt	tggaataacc	accccagca	tccccttccc	tcgtttcata	tcgcttgcac	2160
cccaaccgca	acttatctac	gctgtcctgc	tatccctcag	cgctgctcct	gctcctgctc	2220
actgccctc	gcacagcctt	ggtttgggct	ccgcctgtat	tctcctggta	ctgcaacctg	2280
taaaccagca	ctgcaatgct	gatgcacggg	aagtagtggg	atgggaacac	aatggagga	2340
tcccgcgtct	cgaacagagc	gcgcagagga	acgtgaagg	tctcgcctct	gtcgcacctc	2400
agcgcggcat	acaccacaat	aaccacctga	cgaatgcgct	tggttctctg	tccattagcg	2460
aagcgtccgg	ttcacacacg	tgccaogttg	gogaggtggc	aggtgacaat	gatcggtgga	2520
gctgatggtc	gaaacgttca	cagcctaggg	atatcgaatt	ccgcctgcaa	cgcaagggca	2580
gccacagccg	ctcccaccgg	ccgctgaacc	gacacgtgct	tgggcccctg	ccgctgcct	2640
gccgatgct	tgtgctgggt	aggctgggca	gtgctgccat	gctgattgag	gcttggttca	2700
tcgggtggaa	gcttatgtgt	gtgctgggct	tgcatgccgg	gcaatgcgca	tggtggcaag	2760
agggcgagc	cacttgctgg	agctgcccgg	gtgcctccag	gtggttcaat	cgcggcagcc	2820
agagggattt	cagatgatcg	cgcgtacagg	ttgagcagca	gtgtcagcaa	aggtagcagt	2880
ttgccagaat	gatcggttca	gctgttaate	aatgccagca	agagaagggg	tcaagtgcaa	2940
acacgggcat	gccacagcac	gggcaccggg	gagtggaatg	gcaccaccaa	gtgtgtgcca	3000
gccagcatcg	ccgctgggct	gtttcagcta	caacggcagg	agtcaccaa	cgtaacctag	3060
agctgatcaa	cactgcaate	atcgggcccgg	cgtgatgcaa	gcctgcctgg	cgaagacaca	3120
tggtgtgccc	atgctgccgg	ctgctgcctg	ctgcgcacgc	cgttgagttg	gcagcaggct	3180
cagccatgca	ctggatggca	gctgggctgc	cactgcaatg	tggtggatag	gatgcaagtg	3240
gagcgaatac	caaacctctt	ggctgcttgc	tgggttgcat	ggcatcgcac	catcagcagg	3300
agcgcagcgc	aagggactgg	ccccatgcac	gccatgcca	accggagcgc	accgagtgtc	3360
cacactgtca	ccaggcccgc	aagctttgca	gaacctgct	catggacgca	tgtagcgtg	3420
acgtcccttg	acggcgctcc	tctcgggtgt	gggaaacgca	atgcagcaca	ggcagcagag	3480
gcggcggcag	cagagcggcg	gcagcagcgg	cgggggccac	ccttcttgcg	gggtcgcgcc	3540
ccagccagcg	gtgatgcgct	gatcccaaac	gagttccat	tcatttgcat	gcctggagaa	3600
gcgaggtcgg	ggcctttggg	ctggtgcagc	ccgcaatgga	atgcgggacc	gccaggtag	3660
cagcaaaggc	gcctccccta	ctccgcatcg	atgttccata	gtgcattgga	ctgcatttgg	3720
gtggggcggc	cggtgtttc	ttctgtgttg	caaaacgcgc	cagctcagca	acctgtccc	3780

-continued

---

```

tgggtcccc gtgccgatga aatcgtgtgc acgccgatca gctgattgcc cggtcgcga 3840
agtaggcgcc ctcccttctg ctgcgccctct ctccgtcccg ccactagtat ggccaccgca 3900
tccactttct cggcgttcaa tgcccgtgc ggcgacctgc gtcgctcggc gggctccggg 3960
ccccggcgcc cagcgaggcc cctccccgtg cgcgggcgcg cccccgactg gtccatgctg 4020
ttcgccgtga tcaccaccat cttctccgcc gccgagaagc agtggaccaa cctggagtgg 4080
aagcccaagc ccaaccccc ccagctgctg gacgaccact tcggccccca cggcctgggtg 4140
ttccgccgca ccttcgcat ccgcagctac gaggtgggcc ccgaccgctc caccagcatc 4200
gtggccgtga tgaaccacct gcaggaggcc gccctgaacc acgccaagtc cgtgggcatc 4260
ctgggcgacg gcttcggcac caccctggag atgtccaagc gcgacctgat ctgggtgggtg 4320
aagcgcaccc acgtggccgt ggagcgtac cccgctggg gcgacaccgt ggaggtggag 4380
tgctgggtgg gcgcctccgg caacaacggc cgcgccacg acttcctggt gcgcgactgc 4440
aagaccggcg agatcctgac ccgctgcacc tccttgagcg tgatgatgaa cccccgacc 4500
cgccgcctga gcaagatccc cgaggaggtg cgcggcgaga tcggccccgc cttcatcgac 4560
aacgtggccg tgaaggacga ggagatcaag aagccccaga agctgaacga ctccaccgcc 4620
gactacatcc agggcggcct gacccccgc tggaacgacc tggacatcaa ccagcacgtg 4680
aacaacatca agtacgtgga ctggatcctg gagaccgtgc ccgacagcat cttcgagagc 4740
caccacatct cctccttcac catcgagtac cgcgcgagt gcaccatgga cagcgtgctg 4800
cagtccctga ccaccgtgag cggcggctcc tccgaggccg gcctggtgtg cgagcacctg 4860
ctgcagctgg agggcggcag cgaggtgctg cgcgccaaga ccgagtggcg ccccaagctg 4920
accgactcct tcgcggcgt cagcgtgatc cccgcccagt ccagcgtgat ggactacaag 4980
gaccacgacg gcgactacaa ggaccacgac atcgactaca aggacgacga cgacaagtga 5040
ctcgaggcag cagcagctcg gatagtatcg acacactctg gacgctggtc gttgatgga 5100
ctgttgccg cacacttget gccttgacct gtgaatatcc ctgccgttt tatcaaacag 5160
cctcagtggt tttgatcttg tgtgtacgag cttttgcgag ttgctagctg cttgtgctat 5220
ttggaatac caccccage atccccctcc ctcgtttcat atcgcttga tcccaaccgc 5280
aacttatcta cgtgtcctg ctatccctca gcgctgctcc tgctcctgct cactgcccct 5340
cgcacagcct tggtttgggc tccgctgta ttctcctggt actgcaacct gtaaacccagc 5400
actgcaatgc tgatgcacgg gaagtgtgg gatgggaaca caaatggaaa gctt 5454

```

&lt;210&gt; SEQ ID NO 112

&lt;211&gt; LENGTH: 2933

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 112

```

agagagcgga ggtggggttg tgaggtgggg ttgctgacca ggagctcgcg tcgccgagcg 60
cgactcgcac acggtccagt tcccccccc tccgccaaa cgcaagctc ccatcttgat 120
gcctttccgg ccacctatac tatttcttag ttcgctgtaa catccagacc gtectgaata 180
ataacaatgc cctgtgtcaa gtgcattcct aaaaaaatc tgtccaacc aacaatccca 240
cctgaaatac caccagccct gccagctaca ctcttccat accatctccc tacctccacg 300
cgcaagcgac cccatgcgc gaccaggctc gaaagtgatt tatgactga gacgagcgag 360

```

-continued

---

tggcggcgcg gtcgactgcc ttttcatcac gtgccgtacg tcggcgcaccg ctagggtttt	420
gcacggcaac gcacggcttc gccaaaccga ccagccagga cctcgactac tctaccgcga	480
attcgcctca agaagtgcgc aaatgtgcc aacaccattc cttacagcac tgttcaaact	540
tgatgccaat tttgacattc gggttgctcg ttggctgcgc ccacatcggc cgtgagtgca	600
gcaggcggga tcggacacgg aggacgcggc gtcacgcccc gaacgcagcc cgtaaactca	660
catcaacacg acgtgttgcg taatcccgcc cggctgcgca tcgtgccaac ccattcgcga	720
tgatggtcgc gaaaatggtg tgccaactgc cctgaggag gctctcgcga aacgggcacg	780
tcctgaaac cgaaactgtg gccttgctcg cggccacgca agcacgtgga ccctaaacac	840
caagaaaatc agtaaacag gttgacatcc tctacggcg aattgtttgc ccaacccttc	900
atcgcacact gccattataa tgcattctagc tcggcgacaa gtttagaaaa ggcaggtcgc	960
attgttccat ttccgcgtgg cggcgtgggt gccatttta cgaggtttgg gctcccgggc	1020
agcgaccgag ccaggtcgag tccctctcgc ccgtcgacaa tgttgcaaac cccacaagcg	1080
gctaacaaca acttgatggt acctgtacac tgccaattcc ttcttccccg gccgaggttt	1140
acacgtgatg gccatggctt cgcattcagg ccgacttccc attccgactt tccagagggt	1200
cccgggacgc tgggggttgg ctgcctgagg cccacccttt gttccccgcg tcccgacaaa	1260
cacaattgcy ttacataagg gggagccgcc cccgttcaga gtgcagaaat ctttactat	1320
atthtccagt cgtcagcgaa atcaagtact agtatggcca ccgcatccac tttctcggcg	1380
ttcaatgcc gctgcggcga cctgcctcgc tcggcgggct ccgggccccg gcgcccagcg	1440
aggccccctc ccgtgcgcgg gcgcgcccc gactggteca tgctgttcgc cgtgatcacc	1500
accatcttct ccgcccgcga gaagcagtg accaacctgg agtggaagcc caagcccaac	1560
cccccccagc tgetggacga ccaactcggc ccccacggcc tgggtttccg ccgcaccctc	1620
gccatccgca gctacagagg gggccccgac cgctccacca gcatcgtggc cgtgatgaac	1680
caactgcagg aggcgcctct gaaccacgcc aagtccgtgg gcatcctggg cgacggcttc	1740
ggcaccaccc tggagatgct caagcgcgac ctgatctggg tgggaaagcg caccacgtg	1800
gccgtggagc gctaccccgc ctggggcgac accgtggagg tggagtgctg ggtgggcgcc	1860
tccggcaaca acggccgcgc ccacgacttc ctggtgcgcg actgcaagac cggcgagatc	1920
ctgaccgcct gcacctccct gagcgtgatg atgaacaccc gcaccccgcg cctgagcaag	1980
atccccgagg aggtgcgcgg cgagatcggc cccgccttca tcgacaacgt ggcctggaag	2040
gacgaggaga tcaagaagcc ccagaagctg aacgactcca ccgcccacta catccagggc	2100
ggcctgacct cccgctggaa cgacctggac atcaaccagc acgtgaacaa catcaagtac	2160
gtggactgga tccctggagac cgtgccccgac agcatcttcg agagccacca catctcctcc	2220
ttaccatcgy agtaccgcgc cgagtgcacc atggacagcg tgctgcagtc cctgaccacc	2280
gtgagcggcg gctcctccga ggcggcctg gtgtgcgagc acctgctgca gctggagggc	2340
ggcagcgagg tgctgcgcgc caagaccgag tggcgcceca agctgaccga ctcttccgc	2400
ggcatcagcg tgatccccgc cgagtccagc gtgatggact acaaggacca cgacggcgac	2460
tacaaggacc acgacatcga ctacaaggac gacgacgaca agtgactcga gtttaattaac	2520
tcgaggcagc agcagctcgg atagtatcga cacactctgg acgctggtcg tgtgatggac	2580
tgttgccgcc acacttgctg ccttgacctg tgaatatccc tgccgctttt atcaaacagc	2640
ctcagtggtg ttgatcttgt gtgtaocgcg ttttgcgagt tgctagctgc ttgtgtatt	2700
tgcaataacc acccccagca tccccttccc tcgtttcata tcgcttgcac cccaaccgca	2760

-continued

---

```

acttatctac gctgtcctgc tatccctcag cgctgctect gctcctgctc actgcccctc 2820
gcacagcctt ggtttgggct ccgcctgtat tctcctggta ctgcaacctg taaaccagca 2880
ctgcaatgct gatgcacggg aagtagtggg atgggaacac aatggaaag ctt 2933

```

```

<210> SEQ ID NO 113
<211> LENGTH: 4817
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    polynucleotide

```

```

<400> SEQUENCE: 113

```

```

ggtagccgce tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgcccgc tgccctgccc atgcttgtgc tggtaggct gggcagtgtc 120
gccatgctga ttgaggcttg gttcatcggy tggaaactta tgtgtgtgct gggcttgcat 180
gccgggcaat gcgatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240
tccaggttgt tcaatcgccg cagccagagg gatttcagat gatcgccgct acaggttgag 300
cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgcccgc tggctgtttc agctacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gctgctgctg 600
cacgcccgtt agttggcagc aggcctcagc atgcaactga tggcagctgg gctgccactg 660
caatgtggtg gataggtatc aagtggagcg aataccaaac cctctggctg cttgctgggt 720
tgcattggat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatg acgatgtag cgtgacgctc cctgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacagcgag cagagggggc ggcagcagag cggcggcagc agcggcgggg 960
gccacccttc ttgcccgggc gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcattgctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080
atggaatgcy ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatggt 1140
ccatagtgca ttggactgca tttgggtggg gcggccgctt gtttctttcg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt ccccgtgcc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgccaagtag gcgcccctct ttctgctcgc cctctctccg 1320
tcccgcctct agaatatcaa tgatcgagca ggaaggcctc cacgcccggct ccccgcggc 1380
ctgggtggag cgctgttctg gctacgactg ggcccagcag accatcggct gctccgacgc 1440
cgccgtgttc cgctgttccg cccagggccg ccccgtgctg ttctggaaga ccgacctgtc 1500
cgggccctgt aacgagctgc agaacagggc cgcccgcctg tcctggctgg ccaccaccgg 1560
cgtgcctcgc gccgcctgct tggacgtggt gaccgaggcc ggcccgact ggetgctgct 1620
gggagaggtg cccggccagg acctgctgtc ctcccactg gccccgcggc agaaggtgtc 1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccttcga 1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgacccgc atggaggccg gctggtgga 1800
ccaggacgac ctggacgagg agcaccaggc cctggccccg gccgagctgt tcgcccgcct 1860
gaagggccgc atggccgagc gcgaggacct ggtggtgacc cacggcagc cctgctgcc 1920

```

-continued

---

caacatcatg	gtggagaacg	gcccgttctc	cggttctatc	gactgcggcc	gcctgggcgt	1980
ggccgaccgc	taccaggaca	tcgccttggc	caccgcgcac	atcgccgagg	agctgggcgg	2040
cgagtgggcc	gaccgcttcc	tggtgctgta	eggcatcgcc	gccccgact	cccagcgcac	2100
cgctctctac	cgctgctgg	acgagttctt	ctgacaattg	gcagcagcag	ctcggatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgtgccttg	2220
acctgtgaat	atcccgtccg	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcgcttttg	cgagttgcta	gctgcttgtg	ctatttgcca	ataccacccc	cagcatcccc	2340
ttccctcgtt	tcatatcgct	tgcaccccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcgctg	ctcctgctcc	tgctcactgc	ccctcgcaaa	gccttggttt	gggctccgcc	2460
tgtattctcc	tggtactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccc	cgtctcgaac	agagcgcgca	gaggaacgct	2580
gaaggtctcg	cctctgtcgc	acctcagcgc	ggcatacacc	acaataacca	cctgacgaat	2640
gcgcttgggt	cttcgtccat	tagcgaagcg	tccggttcac	acacgtgcca	cgttggcgag	2700
gtggcaggtg	acaatgatcg	gtggagctga	tggtcgaaac	gttcacagcc	tagggatata	2760
gaattccttt	cttgcgctat	gacacttcca	gcaaaaggta	gggcgggctg	cgagacggct	2820
tcccgcgctc	gcatgcaaca	ccgatgatgc	ttcgaccccc	cgaagctcct	tcggggctgc	2880
atgggcgctc	cgatgcgct	ccagggcgag	cgctgtttaa	atagccaggc	ccccgattgc	2940
aaagacatta	tagcgagcta	ccaaagccat	attcaaacac	ctagatcact	accacttcta	3000
cacaggccac	tcgagcttgt	gatcgcactc	cgctaagggg	gcgcctcttc	ctcttcgttt	3060
cagtcacaac	ccgcaaacgg	cgcgccatgg	tggccgcgcg	cgctccagc	gccttcttcc	3120
ccgtgccccg	ccccggcgcc	tcccccaagc	ccggcaagtt	cggcaactgg	ccctccagcc	3180
tgagccctc	cttcaagccc	aagtcacatc	ccaacggcgg	cttccaggtg	aaggccaacg	3240
acagcgccca	ccccaaagcc	aacggctccg	ccgtgagcct	gaagagcggc	agcctgaaca	3300
cccaggagga	cacctctctc	agcccccccc	ccgcacactt	cctgcaccag	ctgcccgact	3360
ggagccgctc	gctgaccgcc	atcaccaccg	tgttcgtgaa	gtccaagcgc	cccgacatgc	3420
acgaccgcaa	gtccaagcgc	cccgacatgc	tggtggaacg	cttcggcctg	gagtccaccg	3480
tgcaggacgg	cctggtgttc	cgccagctct	tctccatccg	ctcctacgag	atcggcaccg	3540
accgcaaccg	cagcatcgag	accctgatga	accactgca	ggagacctcc	ctgaaccact	3600
gcaagagcac	cggcatcctg	ctggacggct	tcggccgcac	cctggagatg	tgcaagcgcg	3660
acctgatctg	ggtggtgatc	aagatgcaga	tcaaggtgaa	ccgtacccc	gcctggggcg	3720
acaccgtgga	gatcaacacc	cgcttcagcc	gcctgggcaa	gatcggcatg	ggccgcgact	3780
ggctgatctc	cgactgcaac	accggcgaga	tcctggtgcg	cgccaccagc	gcctacgcca	3840
tgatgaacca	gaagaccgcg	cgctgtcca	agctgcctca	cgaggtgcac	caggagatcg	3900
tgcccctgtt	cgtggacagc	cccgtgatcg	aggactccga	cctgaaggtg	cacaagttca	3960
aggtgaagac	cggcgacagc	atccagaagg	gcctgacccc	cggctggaac	gacctggacg	4020
tgaaccagca	cgtgtccaac	gtgaagtaca	tcggctggat	cctggagagc	atgcccaccg	4080
aggtgctgga	gaccaggag	ctgtgctccc	tggccctgga	gtaccgcgcg	gagtgcggcc	4140
gcgactccgt	gctggagagc	gtgaccgcca	tggaccccag	caaggtgggc	gtgcgctccc	4200
agtaccagca	cctgctgcgc	ctggaggacg	gcaccgccat	cgtgaacggc	gccaccgagt	4260
ggcgcccaaa	gaacgcggcg	gccaacggcg	ccatctccac	cggcaagacc	agcaacggca	4320



-continued

---

```

actccgtgtc catggactac aaggaccacg acggcgacta caaggaccac gacatcgact 4380
acaaggacga cgacgacaag tgactcgagg cagcagcagc tcggatagta tcgacacact 4440
ctggacgctg gtcgtgtgat ggactgttgc cgccacactt gctgccttga cctgtgaata 4500
tccctgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac gcgcttttgc 4560
gagttgctag ctgcttgtgc tatttgcgaa taccacccc agcatcccct tccctcgttt 4620
catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgetatccc tcagcgctgc 4680
tctgtcctct gctcactgcc cctcgcacag ccttggtttg ggctccgect gtattctcct 4740
ggtaactgcaa cctgtaaacc agcactgcaa tgctgatgca cgggaagtag tgggatggga 4800
acacaaatgg aaagctt 4817

```

&lt;210&gt; SEQ ID NO 114

&lt;211&gt; LENGTH: 4665

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 114

```

ggtaccgcgc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtget 120
gccatgctga ttgaggcttg gttcatcggg tggaaagctta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240
tccagggtgt tcaatcgccg cagccagagg gatttcagat gatcgccgct acaggttgag 300
cagcagtgtc agcaaagga gcagtttggc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgttcc agctacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gctgtgtgctg 600
cacgcccgtt agttggcagc aggctcagcc atgcaactgga tggcagctgg gctgccactg 660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacaggcag cagagggcgc ggcagcagag cggcggcagc agcggcgggg 960
gccacccttc ttgcccggtc gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcattgctg gagaagcag gctggggcct ttgggctggt gcagcccgca 1080
atggaatgct ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140
ccatagtgca ttggactgca tttgggtggg gcggccggct gtttcttctg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt ccccctgccc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgtctgc cctctctccg 1320
tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380
ctgggtggag cgcctgttgc gctacgaact ggcccagcag accatcggct gctccgacgc 1440
cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttctgtaaga ccgacctgtc 1500

```

-continued

---

cggcgcctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg	1560
cgtgcctcgc gccgcctgtc tggacgtggt gaccgaggcc ggccgcgact ggtgctgct	1620
ggcgaggtg cccggccagg acctgctgtc ctcccactg gccccgcgc agaaggtgtc	1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccttcga	1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcaccgc atggaggccg gectggtgga	1800
ccaggacgac ctggacgagg agcaccaggg cctggcccc gccgagctgt tcgcccgcct	1860
gaaggcccc atgcccgaag gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc	1920
caacatcatg gtggagaacg gccgcttctc cggtctcacc gactgcggcc gectgggct	1980
ggccgaccgc taccaggaca tcgccctggc caccgcgac atcgcgagg agctgggagg	2040
cgagtgggcc gaccgcttc tggctgtgta cggcatcgcc gccccgact cccagcgcat	2100
cgctcttac cgctgctgg acgagttctt ctgacaattg gcagcagcag ctgggatagt	2160
atcgacacac tetggacgct ggtcgtgtga tggactgttg ccgccacct tgctgccttg	2220
acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta	2280
cgctctttg cgagttgcta gctgcttgtg ctatttgca ataccacccc cagcatcccc	2340
ttcctcgtt tcatatcgtc tgcacccaa ccgcaactta tctacgctgt cctgctatcc	2400
ctcagcctg ctctgctcc tgctcactgc ccctcgaca gccttggtt gggctccgcc	2460
tgtattctcc tggactgca acctgtaaac cagcactgca atgctgatgc acgggaagta	2520
gtgggatggg aacacaaatg gaggatccc cgtctcgaac agagcgcgca gaggaacgct	2580
gaaggtctcg cctctgctgc acctcagcgc ggcatacacc acaataacca cctgacgaat	2640
gccttggtt cttcgtccat tagcgaagcg tccggttcc acacgtgcca cgttggcgag	2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatata	2760
gaattccttt cttgcctat gacacttcca gcaaaaggta gggcgggctg cgagacggct	2820
tcccgcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc	2880
atggcgctc cgatccgct ccaggcgag cgctgtttaa atagccaggc ccccgattgc	2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgact accacttcta	3000
cacaggccac tcgagcttgt gatcgactc cgctaagggg gcgcctcttc ctcttcttt	3060
cagtcacaac ccgcaaacac tagtatgacg ttcggggtcg ccctccggc catgggccc	3120
ggtgtctccc tccccggcc cagggtcgcg gtgcgcgcc agtcggcgag tcaggttttg	3180
gagagcgggc gcccccagct gcccgactgg agccgcctgc tgaccgccat caccaccgtg	3240
ttcgtgaagt ccaagcgcgc cgacatgac gaccgcaagt ccaagcgcgc cgacatgctg	3300
gtggacagct tcggcctgga gtccaccgtg caggacggcc tgggttccg ccagtccttc	3360
tccatccgct cctacgagat cggcaccgac cgcaccgcca gcatcgagac cctgatgaac	3420
cacctgcagg agacctccct gaaccactgc aagagcaccg gcatcctgct ggacggcttc	3480
ggccgcaccc tggagatgtg caagcgcgac ctgatctggg tggatgata gatgcagatc	3540
aaggtgaacc gctaccccgc ctggggcgac accgtggaga tcaacaccgc cttcagccgc	3600
ctgggcaaga tcggcatggg ccgcgactgg ctgatctccg actgcaaac ccgagagatc	3660
ctggtgcgag ccaccagcgc ctacgccatg atgaaccaga agaccgcgc cctgtccaag	3720
ctgccctacg aggtgcacca ggagatcgtg cccctgttcg tggacagccc cgtgatcgag	3780
gactccgacc tgaaggtgca caagttcaag gtgaagaccg gcgacagcat ccagaaggcc	3840
ctgacccccg gctggaacga cctggacgtg aaccagcacg tgtccaacgt gaagtacatc	3900

-continued

---

```

ggctggatcc tggagagcat gccaccgag gtgctggaga cccaggagct gtgctcctg 3960
gcectggagt accgcccga gtgcccgc gactccgtgc tggagagcgt gaccgceatg 4020
gaccccagca aggtggcggt gcgctcccag taccagcacc tgctgcgect ggaggacggc 4080
accgceatcg tgaacggcgc caccgagtgg cgcccaaga acgcccgc caacggcgc 4140
atctccaccg gcaagaccag caacggcaac tccgtgtcca tggactaaa ggaccacgac 4200
ggcgactaca aggaccacga catcgactac aaggacgacg acgacaagtg actcgaggca 4260
gcagcagctc ggatagtatc gacacactct ggacgctggt cgtgtgatgg actggtgccc 4320
ccacacttgc tgcttgacc tgtgaatata cctgcccgtt ttatcaaaaca gcctcagtgt 4380
gtttgatctt gtgtgtacgc gcttttgcga gttgctaget gcttgtgcta tttgcgaata 4440
ccacccccag catccccctc cctcgtttca tatcgcttgc atcccaaccg caacttatct 4500
acgctgtcct gctatccctc agcgtgtctc ctgctcctgc tcactgccc tcgcacagcc 4560
ttggttggg ctccgctgt attctcctgg tactgcaacc tgtaaaccag cactgcaatg 4620
ctgatgcacg ggaagtagtg ggatgggaac acaaatggaa agctt 4665

```

&lt;210&gt; SEQ ID NO 115

&lt;211&gt; LENGTH: 4668

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 115

```

ggtaccgcc tgcacgcaa gggcagccac agccgctccc accgcccgt gaaccgacac 60
gtgcttgggc gcctgcccgc tgcctgccc atgcttctgc tggtaggct gggcagtgct 120
gcatgctga ttgagcctg gttcatcggg tggaaactta tgtgtgtgct gggcttgcat 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcccgtgc 240
tccagtggt tcaatcggc cagccagagg gatttcagat gatcgcgctg acaggttgag 300
cagcagtgtc agcaaaggta gcagtttgc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccc catcgcgcc tggctgttcc agtacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgga 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgctg 600
cacgcccgtt agttggcagc aggctcagcc atgactgga tggcagctgg gctgccactg 660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacagggcag cagagggcgc ggcagcagag cggcggcagc agcggcggg 960
gccacccttc ttgcccggtc gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcagcctg gagaagcag gctggggcct ttgggctggt gcagcccgca 1080
atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatggt 1140
ccatagtgca ttggactgca tttgggtggg gcgcccggct gttcttctcg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt ccccgtgcc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgcgaagtag ggcctcctc tctgctcgc cctctctccg 1320

```

-continued

---

tcccgctct	agaatatcaa	tgatcgagca	ggacggcctc	cacgccgget	ccccgcgcg	1380
ctgggtggag	cgectgttcg	gctacgactg	ggcccagcag	accatcgget	gctccgacgc	1440
cgccgtgttc	cgectgtccg	cccagggcgc	ccccgtgctg	ttcgtgaaga	ccgacctgtc	1500
cgggccctgc	aacgagctgc	aggacgaggc	cgcccgcctg	tcctggctgg	ccaccaccgg	1560
cgtgccctgc	gcccgcctgc	tggacgtggt	gaccgaggcc	ggccgcgact	ggctgctgct	1620
ggcgaggtg	cccggccagg	acctgctgtc	ctcccactcg	gccccgcgcg	agaaggtgtc	1680
catcatggcc	gacgccatgc	gcccgcctgca	caccttgac	cccgccacct	gccccttcga	1740
ccaccaggcc	aagcaccgca	tcgagcgcgc	cgcacccgc	atggaggccg	gctggtgga	1800
ccaggacgac	ctggacgagg	agcaccaggg	cctggcccc	gccgagctgt	tcgcccgcct	1860
gaaggcccg	atgcccgacg	gagaggacct	ggtggtgacc	cacggcgacg	cctgcctgcc	1920
caacatcatg	gtggagaacg	gcccgttctc	cggcttcac	gactgcggcc	gctgggctgt	1980
ggccgaccgc	taccaggaca	tcgcccctgg	caccgcgac	atcgcgagg	agctggggcg	2040
cgagtgggcc	gaccgcttc	tggtgctgta	cggcatcgcc	gccccgact	cccagcgc	2100
cgcttctac	cgectgtg	acgagttctt	ctgacaattg	gcagcagcag	ctcgatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgtgccttg	2220
acctgtgaat	atccctgcgc	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcttcttg	cgagttgcta	gctgcttg	ctatttgcga	ataccacccc	cagcatcccc	2340
ttccctcgtt	tcatatcgct	tgcatcccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcgtg	ctcctgctcc	tgctcactgc	ccctcgaca	gccttggtt	gggctccgcc	2460
tgtattctcc	tggtactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccg	cgtctcgaac	agagcgcgca	gaggaaacgct	2580
gaaggtctcg	cctctgtcgc	acctcagcgc	ggcatacacc	acaataacca	cctgacgaat	2640
gcgcttggtt	cttcgtccat	tagcgaagcg	tccggttcac	acacgtgcc	cgttgccgag	2700
gtggcaggtg	acaatgatcg	gtggagctga	tggtcgaaac	gttcacagcc	tagggatate	2760
gaattccttt	cttgcgctat	gacacttcca	gcaaaagta	gggcccgtg	cgagacggct	2820
tcccgcgct	gcatgcaaca	ccgatgatgc	ttcgacccc	cgaagctcct	tcggggctgc	2880
atgggcgctc	cgatgcgct	ccagggcgag	cgctgtttaa	atagccaggc	ccccgattgc	2940
aaagacatta	tagcgagcta	ccaaagccat	attcaaacac	ctagatcact	accacttcta	3000
cacaggccac	tcgagcttgt	gatcgccactc	cgctaagggg	gcgctcttc	ctcttcgttt	3060
cagtcacaac	ccgcaaacac	tagtatggct	atcaagacga	acaggcagcc	tgtggagaag	3120
cctccgttca	cgatcgggac	gctgcgcaag	gccatccccg	cgcactgttt	cgagcgcctg	3180
gcgcttcgtg	ggcgccccca	gctgcccagc	tggagccgcc	tgctgaccgc	catcaccacc	3240
gtgttcgtga	agtccaagcg	ccccgacatg	cacgaccgca	agtccaagcg	ccccgacatg	3300
ctggtggaca	gcttcggcct	ggagtccacc	gtgcaggacg	gcctggtgtt	ccgccagctc	3360
ttctccatcc	gctcctacga	gatcggcacc	gaccgcaccg	ccagcatcga	gaccctgatg	3420
aaccacctgc	aggagacctc	cctgaaccac	tgcaagagca	ccggcatcct	gctggacggc	3480
ttcgcccgca	ccctggagat	gtgcaagcgc	gacctgatct	gggtggtgat	caagatgcag	3540
atcaaggtga	accgctaccc	cgcctggggc	gacaccgtgg	agatcaaac	ccgcttcagc	3600
cgctgggca	agatcggcat	ggcccgcgac	tggtgatct	ccgactgcaa	caccggcgag	3660
atcctggtgc	gcccaccag	cgcctacgcc	atgatgaacc	agaagaccgc	ccgctgtcc	3720

-continued

---

```

aagctgccct acgaggtgca ccaggagatc gtgccctgt tegtggacag ccccgatgc 3780
gaggactccg acctgaaggt gcacaagttc aaggtgaaga ccggcgacag catccagaag 3840
ggcctgacct ccggctggaa cgacctggac gtgaaccagc acgtgtccaa cgtgaagtac 3900
atcggtgga tcttgagag catgcccacc gaggtgctgg agaccagga gctgtgctcc 3960
ctggccctgg agtaccgcc cgagtgcggc cgcgactccg tgctggagag cgtgaccgcc 4020
atggacccca gcaaggtggg cgtgcgctcc cagtaccagc acctgctgcg cctggaggac 4080
ggcaccgcca tcgtgaacgg cgccaccgag tggcgcccca agaacgccgg cgccaacggc 4140
gccatctcca ccggcaagac cagcaacggc aactccgtgt ccatggacta caaggaccac 4200
gacggcgact acaaggacca cgacatcgac tacaaggacg acgacgaaa gtgactcgag 4260
gcagcagcag ctcgatagt atcgacacac tctggacgct ggtcgtgtga tggactgttg 4320
ccgccacact tgctgccttg acctgtgaat atccctgccg cttttatcaa acagcctcag 4380
tgtgtttgat cttgtgtgta cgcgcttttg cgagttgcta gctgcttggt ctatttgga 4440
ataccacccc cagcatcccc ttcctcgtt tcatatcget tgcacccaa ccgcaactta 4500
tctacgctgt cctgctatcc ctacagcgtg ctctgctcc tgctcactgc cctcgcaca 4560
gccttggttt gggctccgcc tgtattctcc tggactgca acctgtaaac cagcactgca 4620
atgctgatgc acgggaagta gtgggatggg aacacaaatg gaaagctt 4668

```

&lt;210&gt; SEQ ID NO 116

&lt;211&gt; LENGTH: 4668

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 116

```

ggtaccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtaggct gggcagtgtc 120
gccatgtgta ttgagcttg gttcatcggg tggaaactta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgatggtg gcaagagggc ggcagcactt gctggagctg ccgctgtgcc 240
tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgctg acaggttgag 300
cagcagtgtc agcaaaggta gcagtttgc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcattgccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgcgcc tggctgtttc agtacaacg 480
gcaggagtca tccaactgaa ccatgagctg atcaacctg caatcatcgg gcgggctgta 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gctgctgctg 600
cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660
caatgtgggt gataggatgc aagttggagc aataccaaac cctctggctg cttgctgggt 720
tgcattggat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaacccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatgg acgatgtgag cgctgacgct ccttgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacaggcag cagagggcggc ggcagcagag cggcggcagc agcggcgggg 960
gccacccttc ttgcccgggtc gcgccccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

```

-continued

---

atggaatgcg	ggaccgcccag	gctagcagca	aaggcgccctc	ccctactccg	catcgatggt	1140
ccatagtgca	ttggactgca	tttgggtggg	gcgccgggct	gtttctttcg	tgttgcaaaa	1200
cgcgccagct	cagcaacctg	tcccgtgggt	ccccgtgcc	gatgaaatcg	tgtgcacgcc	1260
gatcagctga	ttgcccggct	cgcaagtag	gcgccctect	ttctgctcgc	cctctctccg	1320
tcccgcctct	agaatatcaa	tgatecgagca	ggacggcctc	cacgcccggct	ccccgcgcg	1380
ctgggtggag	cgctgtttcg	gctacgactg	ggcccagcag	accatcggt	gctccgacgc	1440
cgccgtgttc	cgctgtccg	cccagggccg	ccccgtgctg	ttcgtgaaga	ccgacctgtc	1500
cgcgccctg	aacgagctgc	aggacgaggc	cgcccgcctg	tcctggctgg	ccaccaccgg	1560
cggtccctgc	gcccgcctgc	tggacgtggg	gaccgaggcc	ggccgcgact	ggctgctgct	1620
ggcgagggty	cccggcccagg	acctgctgtc	ctcccacctg	gccccgcgcg	agaaggtgtc	1680
catcatggcc	gacgccatgc	gcccgcctgca	cacctggac	cccgccacct	gccccttcca	1740
ccaccaggcc	aagcaccgca	tcgagcgcgc	ccgcaccgcg	atggaggccg	gcctggtgga	1800
ccaggacgac	ctggacgagg	agcaccaggg	cctggccccc	gcccagctgt	tcgcccgcct	1860
gaagcccgcg	atgcccagcg	gcgaggacct	ggtggtgacc	cacggcgacg	cctgcctgcc	1920
caacatcatg	gtggagaacg	gcccgtttct	cggtttcatc	gactgcggcc	gcctgggctg	1980
ggcgaccgcg	taccaggaca	tcgcccctgg	caccgcgac	atcgcgagg	agctgggctg	2040
cgagtgggcc	gaccgcttcc	tgggtctgta	cgccatgcc	gcccccgact	cccagcgcct	2100
cgcttctac	cgctgctgg	acgagttctt	ctgacaattg	gcagcagcag	ctcggatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgctgccttg	2220
acctgtgaat	atcccctgcg	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcgcttttg	cgagttgcta	gctgcttgtg	ctatttgcga	ataccacccc	cagcatcccc	2340
ttccctcgtt	tcatatcgct	tgcataccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcctg	ctcctgctcc	tgctcactgc	ccctcgcaca	gccttggttt	gggctccgcc	2460
tgtattctcc	tggactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccc	cgtctcgaac	agagcgcgca	gaggaaacgct	2580
gaaggtctcg	cctctgtcgc	acctcagcgc	ggcatacacc	acaataacca	cctgacgaat	2640
gcgcttggtt	cttcgtccat	tagcgaagcg	tccggttcac	acacgtgcc	cgttggcggag	2700
gtggcaggtg	acaatgatcg	gtggagctga	tggtcgaaac	gttcacagcc	tagggatatac	2760
gaattccttt	cttgcgctat	gacacttcca	gcaaaaggta	ggcggggctg	cgagacggct	2820
tcccggcgct	gcatgcaaca	ccgatgatgc	ttcgaccccc	cgaagctcct	tcggggctgc	2880
atggcgctc	cgatgccgct	ccaggcgag	cgctgtttaa	atagccaggc	ccccgattgc	2940
aaagacatta	tagcgagcta	ccaaagccat	attcaaacac	ctagatcact	accacttcta	3000
cacaggccac	tcgagcttgt	gatcgcactc	cgtaaggggg	gcgctcttc	ctctctgttt	3060
cagtcacaac	ccgcaaacac	tagtatggcc	accgcatacca	ctttctcggc	gttcaatgcc	3120
cgctgcggcg	acctgcgtcg	ctcggcgggc	tccgggcccc	ggcgcccagc	gaggcccctc	3180
cccgtgcgcg	ggcgcccca	gctgcccagc	tggagccgcc	tgctgaccgc	catcaccacc	3240
gtgtctgtga	agtccaagcg	cccgcacatg	cacgaccgca	agtccaagcg	ccccgacatg	3300
ctgggtggaca	gcttcggcct	ggagtccacc	gtgcaggacg	gcctggtggt	ccgccagctc	3360
ttctccatcc	gctcctacga	gatcggcacc	gaccgcaccg	ccagcatcga	gaccctgatg	3420
aaccacctgc	aggagacctc	cctgaaccac	tgcaagagca	ccggcatcct	gctggacggc	3480

-continued

---

ttcggccgca ccttggagat gtgcaagcgc gacctgatct ggggtggtgat caagatgcag	3540
atcaaggtga accgctaccg cgcctggggc gacaccgtgg agatcaacac ccgcttcagc	3600
cgcttgggca agatcgccat gggcccgagc tggctgatct ccgactgcaa cacggcgag	3660
atcctggtgc gcgccaccag cgcctacgcc atgatgaacc agaagaccgg ccgctgtcc	3720
aagctgccct acgaggtgca ccaggagatc gtgcccctgt tcgtggacag ccccgatgc	3780
gaggactccg acctgaaggt gcacaagttc aaggtgaaga ccggcgacag catccagaag	3840
ggcctgaccg ccggctggaa cgacctggac gtgaaccagc acgtgtccaa cgtgaagtac	3900
atcgctgga tcctggagag catgcccacc gaggtgctgg agaccagga gctgtgctcc	3960
ctggccctgg agtaccggcg cgagtggggc cggactccg tgctggagag cgtgaccgcc	4020
atggacccca gcaaggtggg cgtgcgctcc cagtaccagc acctgctgcg cctggaggac	4080
ggcaccgcca tcgtgaacgg cgccaccgag tggcgcccca agaacggcgg cgccaacggc	4140
gccatctcca ccggcaagac cagcaacggc aactccgtgt ccatggacta caaggaccac	4200
gacggcgact acaaggacca cgacatcgac tacaaggacg acgacgaaa gtgactcgag	4260
gcagcagcag ctcgatagat atcgacacac tctggacgct ggtcgtgtga tggactgttg	4320
ccgccacact tgctgccttg acctgtgaat atccctgccc cttttatcaa acagcctcag	4380
tgtgtttgat cttgtgtgta cgcgcttttg cgagttgcta gctgcttggt ctatttgca	4440
ataccacccc cagcatcccc ttccctcggt tcatatcgct tgcateccaa ccgcaactta	4500
tctacgctgt cctgctatcc ctcagcgctg ctctgctcc tgctcactgc cctcgcaca	4560
gccttggttt gggtccggcg tgtattctcc tggactgca acctgtaaac cagcactgca	4620
atgctgatgc acgggaagta gtgggatggg aacacaaatg gaaagctt	4668

&lt;210&gt; SEQ ID NO 117

&lt;211&gt; LENGTH: 4656

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 117

ggtaaccgccc tgcaacgcaa gggcagccc acgcccctccc acccgccgct gaaccgacac	60
gtgcttgggc gcctgcccgc tgcctgccgc atgcttgtgc tggtaggct gggcagtgct	120
gccatgctga ttgagcttg gttcatcggg tggaaagctta tgtgtgtgct gggcttgcac	180
gcccggcaat gcgcatgggt gcaagagggc ggcagcactt gctggagctg ccgctgtgcc	240
tccaggtggt tcaatcgccg cagccagagg gatttcagat gatcgccgct acaggttgag	300
cagcagtgtc agcaaaggtg gcagtttgcc agaatgatcg gttcagctgt taatcaatgc	360
cagcaagaga aggggtcaag tgcaaacacg ggcagcccac agcacgggca ccggggagtg	420
gaatggcacc accaagtgtg tgcgagccag catcgcccgc tggctgttcc agctacaacg	480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta	540
tgcaagcatg cctggcgaag acacatgggt tgcggatgct gccggctgct gcctgctgcg	600
cacgcccgtt agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg	660
caatgtggtg gataggtatg aagtggagcg aataccaaac cctctggctg cttgctgggt	720
tgcattggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat	780
gccccaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc	840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa	900

-continued

---

acgcaatgca gcaacaggcag cagagggcggc ggcagcagag cggcggcagc agcggcgggg	960
gccacccttc ttgcggggtc gcgccccagc cagcgggtgat gcgctgatcc caaacgagtt	1020
cacattcatt tgcattgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca	1080
atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt	1140
ccatagtgca ttggactgca tttgggtggg gcggccggct gtttcttctg tgttgcaaaa	1200
cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc	1260
gatcagctga ttgcccggct cgcgaagtag gcgcctcct ttctgctcgc cctctctccg	1320
tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccg	1380
ctgggtggag cgctgttctg gctacgactg ggcccagcag accatcggct gctccgacgc	1440
cgcctgttcc cgctgtcgc ccagggccg ccccgctgctg ttcgtgaaga ccgacctgtc	1500
cggcgccttg aacgagctgc aggacgaggc gcgccgcctg tcctggctgg ccaccaccgg	1560
cgtgccctgc gcgcgcctgc tggacgtggt gaccgaggcc ggcccgact ggtgctgct	1620
gggcgaggtg cccggccagg acctgctgtc ctcccactg gccccgcgcg agaaggtgtc	1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccttcga	1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcaccgc atggaggccg gctggtgga	1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgccgcct	1860
gaaggcccg atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc	1920
caacatcatg gtggagaacg gccgcttctc cggttctac gactgcggcc gctgggcgt	1980
ggccgaccgc taccaggaca tcgccctggc caccgcgac atcgcgcagg agctgggcgg	2040
cgagtgggcc gaccgcttcc tgggtctgta cggcatcgcc gccccgact cccagcgc	2100
cgcttctac cgctgctgg acgagttctt ctgacaattg gcagcagcag ctcgatagt	2160
atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacct tgcctcctg	2220
acctgtgaat atccctgcgc cttttatcaa acagcctcag tgtgtttgat cttgtgtgta	2280
cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc	2340
ttccctcgtt tcatatcgct tgcattccaa ccgcaactta tctacgctgt cctgctatcc	2400
ctcagcctg ctccctgctc tgctcactgc ccctcgaca gccttggtt gggtccgcc	2460
tgtattctcc tggactgca acctgtaaac cagcactgca atgctgatgc acgggaagta	2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct	2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat	2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag	2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatc	2760
gaattccttt cttgcctat gacacttcca gcaaaaggta gggcgggctg cgagacggct	2820
tcccgccgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc	2880
atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc	2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta	3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcggtt	3060
cagtcaaac ccgcaaacac tagtatggct tccgcggcat tcaccatgct ggcgtgcccc	3120
gcgatgactg gcagggcccc tggggcacgt cgctccggac ggccagtcgc cacccgcctg	3180
agggggcgcg cccccgactg gagccgctg ctgaccgcca tcaccaccgt gttcgtgaag	3240
tccaagcgcc ccgacatgca cgaccgcaag tccaagcgcc ccgacatgct ggtggacagc	3300



-continued

---

```

ttcgccctgg agtccaccgt gcaggacggc ctggtgttcc gccagtcctt ctccatccgc 3360
tcctacgaga tcggcaccga ccgcaccgcc agcatcgaga ccctgatgaa ccacctgcag 3420
gagacctccc tgaaccactg caagagcacc ggcattcctgc tggacggctt cggccgcacc 3480
ctggagatgt gcaagcgcga cctgatctgg gtggtgatca agatgcagat caaggtgaac 3540
cgctaccccc cctggggcga caccgtggag atcaacaccc gcttcagccg cctgggcaag 3600
atcggcattg gccgcgactg gctgatctcc gactgcaaca ccggcgagat cctggtgcgc 3660
gccaccagcg cctacgccat gatgaaccag aagacccgcc gcctgtccaa gctgccctac 3720
gaggtgcacc aggagatcgt gcccctgttc gtggacagcc ccgtgatcga ggactccgac 3780
ctgaaggtgc acaagttcaa ggtgaagacc ggcgacagca tccagaaggg cctgaccccc 3840
ggctggaacg acctggacgt gaaccagcac gtgtccaacg tgaagtacat cggctggatc 3900
ctggagagca tgcccaccga ggtgctggag acccaggagc tgtgctccct ggccctggag 3960
taccgcgcgc agtgccggcgc cgactccgtg ctggagagcg tgaccgccat ggaccccagc 4020
aagtggggcg tgcgctccca gtaccagcac ctgctgcgcc tggaggacgg cacccccatc 4080
gtgaacggcg ccaccagatg gcgccccaa gacgcccggc ccaacggcgc catctccacc 4140
ggcaagacca gcaacggcaa ctccctgtcc atggactaca aggaccacga cggcgactac 4200
aaggaccacg acatcgacta caaggacgac gacgacaagt gactcgaggc agcagcagct 4260
cggatagtat cgacacactc tggacgctgg tcgtgtgatg gactgttgcc gccacacttg 4320
ctgccttgac ctgtgaatat ccctgcccgt tttatcaaac agcctcagtg tgtttgatct 4380
tgtgtgtacg cgcttttgcg agttgttagc tgettgtgct atttgcgaat accaccccc 4440
gcatccccct ccctcgttcc atatcgcttg catcccaacc gcaacttacc tacgctgtcc 4500
tgetatccct cagcgtgct cctgctcctg ctcaactgcc ctgcacagc cttggtttgg 4560
gctccgctg tattctcctg gtactgcaac ctgtaaacca gcaactgcaat gctgatgcac 4620
gggaagtagt gggatgggaa cacaaatgga aagctt 4656

```

&lt;210&gt; SEQ ID NO 118

&lt;211&gt; LENGTH: 4721

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 118

```

ggtaaccgcc tgcaacgcaa gggcagccac agccgctccc acccggcctt gaaccgacac 60
gtgcttgggc gcctgcccgc tgcctgccgc atgcttgtgc tggtaggctt gggcagtgct 120
gccatgctga ttgaggcttg gttcatcggg tggaagetta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgcatgggt gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240
tccaggtggt tcaatcgccg cagccagagg gatttcagat gatcgcgctt acaggttgag 300
cagcagtgct agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcattgccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgttcc agtacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcggggctga 540
tgcaagcatg cctggcgaag acacatgggt tgcggatgct gccggctgct gcctgctgcg 600
cacgcccgtt agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

```

-continued

---

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt	720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat	780
gccaaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc	840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa	900
acgcaatgca gcacaggcag cagaggcggc ggacgacag cgcgggcagc agcggcgggg	960
gccacccttc ttgctgggtc gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt	1020
cacattcatt tgcatgcctg gagaagcag gctggggcct ttgggctggt gcagcccga	1080
atggaatgcg ggaccgccag gctagcagca aaggcgctc ccctactccg catcgatgtt	1140
ccatagtgca ttggactgca tttgggtggg gcggccggt gtttctttcg tgtgcaaaa	1200
cgcgccagct cagcaacctg tcccgtgggt ccccgtgcc gatgaaatcg tgtgcacgcc	1260
gatcagctga ttgccggct cgcgaaagtag gcgcccctct ttctgctcgc cctctctccg	1320
tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgcccgtt ccccgcgcg	1380
ctgggtggag cgctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc	1440
cgcgctgttc cgctgtccg cccagggcgg ccccgctgctg ttctggaaga cgcacctgtc	1500
cgggccctg aacgagctgc aggacgaggc cgcccgcctg tcttggttg ccaccaccgg	1560
cgtgccctgc gcgcccgtgc tggacgtggt gaccgaggcc ggccgcgact ggtgctgct	1620
ggcgagaggt cccggccagg acctgctgtc ctcccactg gccccgcgcg agaaggtgtc	1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccact gcccttcga	1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcaccgc atggaggccg gcttgggtga	1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct	1860
gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgctgccc	1920
caacatcatg gtggagaacg gccgcttctc cggcttcac gactgcgccc gcttggcgct	1980
ggccgaccgc taccaggaca tcgcccctggc caccgcgac atcgccgagg agctggggcg	2040
cgagtgggcc gaccgcttc ttggtgctga cggcatcgcc gccccgact cccagcgcct	2100
cgctctctac cgctgctgg acgagttctt ctgacaattg gcagcagcag ctcgatagt	2160
atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tctgctctg	2220
acctgtgaat atccctgcgc cttttatcaa acagcctcag tgtgtttgat cttgtgtgta	2280
cgcgcttttg cgagttgcta gctgctgtg ctatttgcga ataccacccc cagcatcccc	2340
ttccctcggt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc	2400
ctcagcgtg ctctgctcc tgctcactgc ccctcgaca gccttggttt gggtccgcc	2460
tgtattctcc tggactgca acctgtaaac cagcactgca atgctgatgc acgggaagta	2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct	2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat	2640
gcgcttgggt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag	2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc taggatatac	2760
gaattccttt cttgctgat gacacttcca gcaaaagta gggcgggctg cgagacggct	2820
tcccggcgtc gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc	2880
atgggcgctc cgatgccgt ccaggggcag cgctgtttaa atagccaggc ccccgattgc	2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgactact accacttcta	3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgccccttc ctctctggtt	3060

-continued

---

```

cagtcacaac ccgcaaacgg cgcgccatgg ccaccaccag cctggcctcc gccttctgct 3120
ccatgaagge cgtgatgctg gcccgcgacg gcccgggcat gaagccccgc agctccgacc 3180
tgcagctgcg cgcggcaaac gccccacct cctgaagat gatcaacggc accaagttca 3240
gctacaccga gagcctgaag cgcctgcccc actggtccat gctgttcgcc gtgatacca 3300
ccatcttcag cgcggcggag aagcagtgga ccaacctgga gtggaagccc aagccccagc 3360
tgccccagct gctggacgac cacttcggcc tgcacggcct ggtgttcgcg cgcaccttcg 3420
ccatccgctc ctacgaggtg ggccccgacc gcagcacctc catcctggcc gtgatgaacc 3480
acatgcagga ggccaccctg aaccacgcca agagcgtggg catcctgggc gacggcttcg 3540
gcaccacctt ggagatgtcc aagcgcgacc tgatgtgggt ggtgcgccc acccacgtgg 3600
ccgtggagcg ctacccccacc tggggcgaca ccgtggaggt ggagtgetgg atcggcgcca 3660
gcggcaacaa cggcatgcgc cgcgacttcc tggtgcgcga ctgcaagacc ggcgagatcc 3720
tgaccgctg cactccctg agcgtgctga tgaacaccgg caccgcgccc ctgagacca 3780
tccccgacga ggtgcgcggc gagatcggcc ccgccttcat cgacaacgtg gccgtgaagg 3840
acgacgagat caagaagctg cagaagctga acgactccac cgcgactac atccagggcg 3900
gcctgacccc ccgctggaac gacctggacg tgaaccagca cgtgaacaac ctgaagtacg 3960
tggcctgggt gttcgagacc gtgccccgaca gcattctcga gtcccaccac atcagctcct 4020
tcacctgga gtaccgccc gagtgcaccc gcgactccgt gctgcgcagc ctgaccaccg 4080
tgagcggcgg cagctccgag gccggcctgg tgtgcgacca cctgctgcag ctggagggcg 4140
gcagcgaggt gctgcgcgcc cgcacagagt ggcgccccaa gctgaccgac tccttcgcg 4200
gcatcagcgt gatccccgcc gagccccgcg tgatggacta caaggaccac gacggcgcact 4260
acaaggacca cgacatcgac tacaaggacg acgacgacaa gtgatgactc gaggcagcag 4320
cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg ttgccgccac 4380
acttgetgce ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt 4440
gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg cgaataccac 4500
ccccagcatc ccttccctc gtttcatac gcttgcaccc caaccgcaac ttatctacgc 4560
tgtctgcta tccctcagcg ctgctcctgc tctgctcac tgcccctcgc acagccttgg 4620
tttgggtccc gcctgtatct tcctggtact gcaacctgta aaccagcact gcaatgctga 4680
tgacgggaa gtagtgggat gggaacacaa atggaagct t 4721

```

&lt;210&gt; SEQ ID NO 119

&lt;211&gt; LENGTH: 4650

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 119

```

ggtaccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gectgcgcc tgcctgccc atgcttctgc tggtaggct gggcagtgct 120
gccatgctga ttgagcttg gttcatcggg tggaagetta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240
tccaggtggt tcaatcgcg cagccagagg gatttcagat gatcgcgctg acaggttgag 300
cagcagtgtc agcaaaggtg gcagtttccc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420

```

-continued

---

gaatggcacc	accaagtgtg	tgcgagccag	catcgccgcc	tggctgtttc	agctacaacg	480
gcaggagtca	tccaacgtaa	ccatgagctg	atcaaacctg	caatcatcgg	gcgggctgga	540
tgcaagcatg	cctggcgaag	acacatggtg	tgcggatgct	gccggctgct	gectgctgag	600
cacgcccgtt	agttggcagc	aggctcagcc	atgcactgga	tggcagctgg	gctgccactg	660
caatgtggtg	gataggatgc	aagtggagcg	aataccaaac	cctctggctg	cttgcctgggt	720
tgcatggcat	cgcaccatca	gcaggagcgc	atgcgaaggg	actggcccca	tgcacgccat	780
gccaaaccgg	agcgcaccga	gtgtccacac	tgtcaccagg	cccgaagct	ttgcagaacc	840
atgctcatgg	acgcatgtag	cgctgacgtc	ccttgacggc	gctcctctcg	ggtgtgggaa	900
acgcaatgca	gcacaggcag	cagagggcgc	ggcagcagag	cggcggcagc	agcggcgggg	960
gccacccttc	ttgcccggtc	gcgccccagc	cagcggatgat	gcgctgatcc	caaacgagtt	1020
cacattcatt	tgcatgcctg	gagaagcgag	gctggggcct	ttgggctggt	gcagcccgca	1080
atggaatgcg	ggaccgccag	gctagcagca	aaggcgcttc	ccctactcgg	catcgatggt	1140
ccatagtgca	ttggactgca	tttgggtggg	gcggccggct	gtttctttcg	tgttgcaaaa	1200
cgcgccagct	cagcaacctg	tcccgtgggt	ccccctgccc	gatgaaatcg	tgtgcacgcc	1260
gatcagctga	ttgccggctt	cgcgaagtag	gcgcccctct	ttctgctcgc	cctctctcgg	1320
tcccgcctct	agaatatcaa	tgatcgagca	ggcggccttc	cacgccggct	ccccgcgcgc	1380
ctgggtggag	cgctgttctg	gtacgactg	ggcccagcag	accatcggtt	gctccgacgc	1440
cgcctgttct	cgctgtctcg	cccagggcgg	ccccgtgctg	ttcgtgaaga	cgcacctgtc	1500
cggcgcctct	aacgagctgc	aggacgaggc	cgcccgcctg	tcctggctgg	ccaccaccgg	1560
cgtgccctgc	gcgcctgtgc	tggacgtggt	gaccgaggcc	ggccgcgact	ggctgctgct	1620
ggggcaggtg	cccggccagg	acctgctgtc	ctcccacctg	gccccgcggc	agaagggtgtc	1680
catcatggcc	gacgccatgc	gccgcctgca	cacctggac	cccgccacct	gcccttcga	1740
ccaccaggcc	aagcaccgca	tcgagcgcgc	ccgcaccgcg	atggaggcgg	gcctggtgga	1800
ccaggacgac	ctggacgagg	agcaccaggg	cctggccccc	gccgagctgt	tgcgccgctt	1860
gaaggcccgc	atgcccgacg	gcgaggacct	ggtggtgacc	cacggcgacg	cctgctgccc	1920
caacatcatg	gtggagaacg	gccgcttctc	cggttctatc	gactgcggcc	gcctgggctg	1980
ggccgaccgc	taccaggaca	tgcctctggc	caccgcggac	atcgcggagg	agctgggctg	2040
cgagtgggcc	gaccgcttcc	tgggtgctga	cggcatcgcc	gcccccgact	cccagcgcct	2100
cgcctctctc	cgctgctggg	acgagttctt	ctgacaattg	gcagcagcag	ctcggatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgtgccttgg	2220
acctgtgaat	atccctgcgg	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcgcttttg	cgagttgcta	gctgcttggg	ctatttgcga	ataccacccc	cagcatcccc	2340
ttccctcggt	tcatatcgct	tgcaccccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcctg	ctcctgctcc	tgtcactgct	ccctcgcaca	gccttggttt	gggctccgcc	2460
tgtattctcc	tggactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccg	cgtctcgaac	agagcgcgca	gaggaaacgct	2580
gaaggtctcg	cctctgtcgc	acctcagcgc	ggcatacacc	acaataacca	cctgacgcaat	2640
gcgcttgggt	cttgcctcat	tagcgaagcg	tccggttcac	acacgtgcca	cgttggcggag	2700
gtggcaggtg	acaatgatcg	gtggagctga	tggtcgaaac	gttcacagcc	tagggatatac	2760
gaattccttt	cttgcgctat	gacacttcca	gcaaaaggta	gggcgggctg	cgagacggct	2820

-continued

---

```

tcccgcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgcgcgt ccaggggcgag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcggagcta ccaaagccat attcaaacac ctgatcactt accacttcta 3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg ggcctcttc ctcttcgttt 3060
cagtcacaac ccgcaaacac tagtatgacg ttcggggctc ccctccggc catgggccc 3120
ggtgtctccc ttcccggcc cagggctcgc gtgcgcgcc agtcggcgag tcaggttttg 3180
gagagcgggc gcgccccga ctggtccatg ctgttcgcc tgatcaccac catcttcagc 3240
gccgcccaga agcagtgagc caacctggag tggaaagcca agcccaagct gccccagctg 3300
ctggacgacc acttcggcct gcacggcctg gtgttcgcc gcaccttcgc catccgctcc 3360
tacgaggtgg gccccagcc cagcaacctc atcctggccg tgatgaacca catgcaggag 3420
gccacctga accacgccc gagcgtgggc atcctggcg acggcttcgg caccacctg 3480
gagatgtcca agcgcgacct gatgtgggtg gtgcgcgcc cccacgtggc cgtggagcgc 3540
taccacctc gggcgacac cgtggaggtg gagtgtgga tcggcgccag cggcaacaac 3600
ggcatgcgcc gcgacttctt ggtgcgcgac tgcaagaccg gcgagatcct gacctcctgc 3660
acctccctga gcgtgctgat gaacacctgc acccgccgcc tgagcaccat ccccgacgag 3720
gtgcgcggcg agatcgccc cgccttcac gacaacctgg ccgtgaagga cgacgagatc 3780
aagaagctgc agaagctgaa cgactccacc gccgactaca tccagggcgg cctgaccccc 3840
cgctggaacg acctggacgt gaaccagcac gtgaacaacc tgaagtacgt ggctgggtg 3900
ttcgagaccg tgcccagac catcttcgag tcccaccaca tcagctcctt cacctggag 3960
taccgcccgc agtgcacccc cgactccgtg ctgcgcagcc tgaccaccgt gagcggcggc 4020
agetccgagg ccggcctggt gtgcgaccac ctgctgcagc tggagggcgg cagcgagggt 4080
ctgcgcgcc gcaccgagtg gcgccccaa ctgaccgact ccttcgcgg catcagcgtg 4140
atcccgcgc agccccgcgt gatggactac aaggaccacg acggcgacta caaggaccac 4200
gacatcgact acaaggacga cgacgacaag tgatgactcg aggcagcagc agctcggata 4260
gtatcgacac actctggacg ctggtcgtgt gatggactgt tgccgccaca cttgctgcct 4320
tgacctgta atacctctc cgcttttata aacagcctc agtgtgtttg atcttgtgtg 4380
tacgcgcttt tcgagttgc tagctgcttg tgctatttgc gaataccacc cccagcatcc 4440
ccttccctcg tttcatatcg cttgcatccc aaccgcaact tatctacgt gtctgctat 4500
ccctcagcgc tgctcctgct cctgctcact gccctcgcga cagccttggg ttgggctccg 4560
cctgtattct cctggtactg caacctgtaa accagcactg caatgctgat gcacgggaag 4620
tagtgggatg ggaacacaaa tggaaagctt 4650

```

&lt;210&gt; SEQ ID NO 120

&lt;211&gt; LENGTH: 4653

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 120

```

ggtaccgcgc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgcgcgc tgcctgcgcg atgcttgtgc tggtaggct gggcagtgct 120
gccatgctga ttgaggcttg gttcatcggg tggaaagctta tgtgtgtgct gggcttgcac 180

```

-continued

---

gccgggcaat	gcgcatggtg	gcaagagggc	ggcagcactt	gctggagctg	ccgcggtgcc	240
tccaggtggt	tcaatcgcg	cagccagagg	gatttcagat	gatcgcgctg	acaggttgag	300
cagcagtgtc	agcaaaggta	gcagtttgcc	agaatgatcg	gttcagctgt	taatcaatgc	360
cagcaagaga	aggggtcaag	tgcaaacacg	ggcatgccac	agcacgggca	ccggggagtg	420
gaatggcacc	accaagtgtg	tgcgagccag	catcgccgcc	tggtgtttc	agctacaacg	480
gcaggagtca	tccaacgtaa	ccatgagctg	atcaaacctg	caatcatcgg	gcgggctgta	540
tgcaagcatg	cctggcgaag	acacatggtg	tgcggatgct	gccggctgct	gcctgctgcg	600
cacgccgttg	agttggcagc	aggctcagcc	atgcactgga	tggcagctgg	gctgccactg	660
caatgtggtg	gataggatgc	aagtggagcg	aataccaaac	cctctggctg	cttgtctgggt	720
tgcatggcat	cgcaccatca	gcaggagcgc	atgcgaaggg	actggcccca	tgcacgccat	780
gccaaaaccg	agcgcaccga	gtgtccacac	tgtcaccagg	cccgaagct	ttgcagaacc	840
atgtctatgg	acgcatgtag	cgctgacgtc	ccttgacggc	gctcctctcg	ggtgtgggaa	900
acgcaatgca	gcacaggcag	cagagggcgc	ggcagcagag	cggcggcagc	agcggcgggg	960
gccacccttc	ttgcggggtc	gcgccccagc	cagcgggtgat	gcgctgatcc	caaacgagtt	1020
cacattcatt	tgcatgcctg	gagaagcgag	gctggggcct	ttgggctggt	gcagcccgc	1080
atggaatgcg	ggaccgccag	gctagcagca	aaggcgcctc	ccctactccg	catcgatggt	1140
ccatagtgca	ttggactgca	tttgggtggg	gcggccggct	gtttctttcg	tgttgcaaaa	1200
cgcgccagct	cagcaacctg	tcccgtgggt	ccccctgcc	gatgaaatcg	tgtgcacgcc	1260
gatcagctga	ttgcccggct	cgcgaagtag	gcgcctcct	ttctgctcgc	cctctctccg	1320
tcccgcctct	agaatatcaa	tgatcgagca	ggacggcctc	cacgccggct	ccccgcgcgc	1380
ctgggtggag	cgctgtttcg	gctacgactg	ggcccagcag	accatcggt	gctccgacgc	1440
cgcctgttcc	cgctgtccg	cccagggccg	ccccgtgctg	ttcgtgaaga	ccgacctgtc	1500
cgggcccttg	aacgagctgc	aggacgaggg	cgcccgcctg	tcctggctgg	ccaccaccgg	1560
cgtgccctgc	gcccgcctgc	tggacgtggt	gaccgaggcc	ggccgcgact	ggctgctgct	1620
gggcgaggtg	cccggccagg	acctgctgtc	ctcccacctg	gccccgcgcg	agaaggtgtc	1680
catcatggcc	gacgccatgc	gcccgcctgca	caccctggac	cccgccacct	gccccttcca	1740
ccaccaggcc	aagcaccgca	tcgagcgcgc	ccgcaccgcg	atggaggccg	gcctggtgga	1800
ccaggacgac	ctggacgagg	agcaccaggg	cctggccccc	gccgagctgt	tcgcccgcct	1860
gaaggcccgc	atgcccgacg	gcgaggacct	ggtggtgacc	cacggcgacg	cctgcctgce	1920
caacatcatg	gtggagaacg	gccgcttctc	cggtctcatc	gactgcggcc	gcctgggctg	1980
ggccgaccgc	taccaggaca	tcgccttggc	caccgcgcac	atcgccgagg	agctgggctg	2040
cgagtgggce	gaccgcttcc	tggtgctgta	cggcctcgcc	gccccgact	cccagcgc	2100
cgcttctac	cgctgctgg	acgagttctt	ctgacaattg	gcagcagcag	ctcgatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgtgccttg	2220
acctgtgaat	atccctgccg	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcgcttttg	cgagttgcta	gctgctgtg	ctatttgcga	ataccacccc	cagcatcccc	2340
ttccctcgtt	tcatatcgct	tgcatcccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcgtg	ctcctgctcc	tgctcactgc	ccctcgcaca	gccttggttt	gggctccgce	2460
tgtattctcc	tggtactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccc	cgtctcgaac	agagcgcgca	gaggaacgct	2580

-continued

---

```

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcc a cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatac 2760
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tegggtctgc 2880
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgactact accacttcta 3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060
cagtcacaac ccgcaaacac tagtatggct atcaagacga acaggcagcc tgtggagaag 3120
cctccgttca cgatcgggac gctgcgcaag gccatccccg cgcactgttt cgagcgctcg 3180
gcgcttcgtg ggcgcgcccc cgactggctc atgctgttcg ccgtgatcac caccatcttc 3240
agcgcgcccc agaagcagtg gaccaacctg gactggaagc ccaagcccaa gctgccccag 3300
ctgctggacg accacttcg cctgcacggc ctggtgttcc gccgcacctt cgccatccgc 3360
tcctacgagg tgggccccga ccgcagcacc tccatcctgg ccgtgatgaa ccacatgcag 3420
gaggccaccc tgaaccacgc caagagcgtg ggcacacctg gcgacggctt cggcaccacc 3480
ctggagatgt ccaagcgcg cctgatgtgg gtggtgcgcc gcaccacgt gcccgaggag 3540
cgctaccccc cctgggggga caccgtggag gtggagtgtt ggatcggcgc cagcggcaac 3600
aacggcatgc gccgcgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 3660
tgcaacctcc tgagcgtgct gatgaacacc cgcacccgcc gcctgagcac catccccgac 3720
gaggtgcgcg gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggaocgagag 3780
atcaagaagc tcgagaagct gaacgactcc accgccgact acatccaggg cggcctgacc 3840
ccccgctgga acgacctgga cgtgaaccag cacgtgaaca acctgaagta cgtggcctgg 3900
gtgttcgaga ccgtgccccg cagcatcttc gactcccacc acatcagctc cttcaccttg 3960
gagtaccgcc gcgagtgca cccgcactcc gtgctgcgca gcctgaccac cgtgagcggc 4020
ggcagctccg aggcggcctt ggtgtgcgac cacctgctgc agctggaggg cggcagcgag 4080
gtgctgcgcg cccgcaccga gtggcgcccc aagctgaccg actccttcgg cggcatcagc 4140
gtgatcccc cggagccccg cgtgatggac tacaaggacc acgacggcga ctacaaggac 4200
cacgacatcg actacaagga cgcgacgac aagtgatgac tcgaggcagc agcagctcgg 4260
atagtatcga cacactctgg acgctggctg tgtgatggac tgttgccgcc acacttctgt 4320
ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 4380
gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcaataacc acccccagca 4440
tccccttccc tcgtttcata tcgcttgcat cccaaccgca acttatctac gctgtcctgc 4500
tatccctcag cgtgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 4560
ccgcctgtat tctcctgcta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 4620
aagtagtggg atgggaacac aaatggaaag ctt 4653

```

&lt;210&gt; SEQ ID NO 121

&lt;211&gt; LENGTH: 4653

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 121

-continued

---

ggtaccgccc	tgcaacgcaa	gggcagccac	agccgctccc	accgcccgt	gaaccgacac	60
gtgcttgggc	gctgcccgc	tgccctgccc	atgcttgtgc	tggtaggct	gggcagtgt	120
gccatgctga	ttgaggcttg	gttcatcggg	tggaagctta	tgtgtgtgct	gggcttgcac	180
gccgggcaat	gdcgatggg	gcaagaggg	ggcagcactt	gctggagctg	ccgcggtgcc	240
tccaggtggt	tcaatcgccg	cagccagagg	gatttcagat	gatcgccgct	acaggttgag	300
cagcagtgtc	agcaaaggta	gcagtttgcc	agaatgatcg	gttcagctgt	taatcaatgc	360
cagcaagaga	aggggtcaag	tgcaaacacg	ggcatgccac	agcacgggca	ccggggagtg	420
gaatggcacc	accaagtgtg	tgccagccag	catcgccgcc	tggctgtttc	agctacaacg	480
gcaggagtca	tccaacgtaa	ccatgagctg	atcaaacctg	caatcatcgg	gcgggctgta	540
tgaagcatg	cctggcgaag	acacatggtg	tgcggatgct	gccggctgct	gcctgctgcg	600
cacgcccgtt	agttggcagc	aggctcagcc	atgcactgga	tggcagctgg	gctgccactg	660
caatgtggtg	gataggatgc	aagtggagcg	aataccaaac	cctctggctg	cttctctgggt	720
tgcattggat	cgccaccatca	gcaggagcgc	atgcgaaggg	actggcccca	tgccacccat	780
gccaaaccgg	agcgcaccga	gtgtccacac	tgccaccagg	cccgaagct	ttgcagaacc	840
atgctcatgg	acgcattgat	cgctgacgtc	ccttgacggc	gctcctctcg	ggtgtgggaa	900
acgcaatgca	gcacagcgag	cagagggcgg	ggcagcagag	cgccggcagc	agcggcgggg	960
gccacccttc	ttgcggggtc	gcgccccagc	cagcggatgat	gcgctgatcc	caaaccgagtt	1020
cacattcatt	tgcatgcctg	gagaagcgag	gctggggcct	ttgggtggtg	gcagcccgca	1080
atggaatgcg	ggaccgccag	gctagcagca	aaggcgcctc	ccctactccg	catcgatggt	1140
ccatagtgca	ttggactgca	tttgggtggg	gcggccggct	gtttctttcg	tgttgcaaaa	1200
cgccgacgct	cagcaacctg	tcccgtgggt	ccccgctgcc	gatgaaatcg	tgtgcacgcc	1260
gatcagctga	ttgccggcct	cgccgaagtag	gcgcccctct	ttctgctcgc	cctctctccg	1320
tcccgcctct	agaatatcaa	tgatcgagca	ggaaggcctc	cacgcgggct	ccccgcggc	1380
ctgggtggag	cgctctttcg	gctacgactg	ggcccagcag	accatcggct	gctccgacgc	1440
cgccgtgttc	cgctgtcccg	cccagggccg	ccccgtgctg	ttcgtgaaga	ccgacctgtc	1500
cgccgcccct	aacgagctgc	aggacgaggc	cgcccgcctg	tcctggtggt	ccaccaccgg	1560
cgtgcccctg	gccgcccgtg	tggacgtggt	gaccgaggcc	ggccgcgact	ggtgctgct	1620
ggcgagaggt	cccggccagg	acctgctgtc	ctcccactcg	gcccccccg	agaaggtgtc	1680
catcatggcc	gacgcccagc	gcccgcctgca	caccttgac	cccgccacct	gccccttcca	1740
ccaccaggcc	aagcaccgca	tccgagcgcg	ccgcaccgcc	atggaggccg	gctggtgga	1800
ccaggacgac	ctggacgagg	agcaccaggg	cctggccccc	gccgagctgt	tcgcccgcct	1860
gaagggcccg	atgcccgcag	gcgaggacct	ggtggtgacc	cagggcgacg	cctgcccgtc	1920
caacatcatg	gtggagaacg	gccgcttctc	cggttctatc	gactgcggcc	gctggggcgt	1980
ggccgaccgc	taccaggaca	tcgcccctgg	caccgcgac	atcggccagg	agctggggcg	2040
cgagtgggcc	gaccgcttcc	tgggtgctgta	cgccatcgcc	gcccccgact	cccagcgcac	2100
cgccctctac	cgctgctgg	acgagttctt	ctgacaattg	gcagcagcag	ctcggatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgtgccttg	2220
acctgtgaat	atcccgtccc	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcccttttg	cgagttgcta	gctgcttggt	ctatttgcca	ataccacccc	cagcatcccc	2340
ttccctcggt	tcatatcgct	tgcatcccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400



-continued

---

```

ctcagcgctg ctctgctcc tgctcactgc cctcgcaca gccttggtt gggtccgce 2460
tgtattctcc tggactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatata 2760
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccgcgctg gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgcgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgactact accacttcta 3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060
cagtcacaac ccgcaaacac tagtatggcc accgcatcca ctttctcggc gttcaatgcc 3120
cgctgcggcg acctgcgctg ctccgggggc tccgggcccc ggcgcccagc gaggccctc 3180
cccgtgcgcg ggcgcccc cgaactggtc atgctgttcg ccgtgatcac caccatcttc 3240
agcgcgcccg agaagcagtg gaccaacctg gaggtggaagc ccaagcccaa gctgccccag 3300
ctgctggagc accacttcgg cctgcacggc ctggtgttcc gccgcacctt cgccatccgc 3360
tcctacgagg tgggccccga ccgcagcacc tccatcctgg ccgtgatgaa ccacatgcag 3420
gaggccaccc tgaaccacgc caagagcgtg ggcatcctgg gcgacggctt cggcaccacc 3480
ctggagatgt ccaagcgcga cctgatgtgg gtggtgcgcc gcacccacgt ggcctggag 3540
cgctacccca cctggggcga cacctggag gtggagtgtt ggatcggcgc cagcggcaac 3600
aacggcatgc gccgcgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 3660
tgcacctccc tgagcgtgct gatgaacacc cgcacccgcc gcctgagcac catccccgac 3720
gaggtgcgcg gcgagatcgg ccccgcttc atcgacaacg tggccgtgaa ggacgacgag 3780
atcaagaage tgcagaaget gaacgactcc accgcccact acatccaggg cggcctgacc 3840
ccccgctgga acgacctgga cgtgaaccag cacgtgaaca acctgaagta cgtggcctgg 3900
gtgttcgaga ccgtgcccga cagcatcttc gagtcccacc acatcagctc cttcacctg 3960
gagtaccgcc gcgagtgcac ccgcgactcc gtgctgcgca gcctgaccac cgtgagcggc 4020
ggcagctccc aggccggcct ggtgtgcgac cacctgctgc agctggaggg cggcagcag 4080
gtgctgcgcg cccgcaccga gtggcgcgcc aagctgaccg actccttcg cggcatcagc 4140
gtgatccccg ccgagccccg cgtgatggac tacaaggacc acgacggcga ctacaaggac 4200
cacgacatcg actacaagga cgcagcagc aagtgatgac tcgaggcagc agcagctcgg 4260
atagtatcga cacactctgg acgctggtcg tgtgatggac tgttgcgcc acacttctg 4320
ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatctgt 4380
gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcgaatacc acccccagca 4440
tccccttccc tegtctcata tcgcttgcac cccaaccgca acttatctac gctgtcctgc 4500
tatccctcag cgctgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 4560
ccgctgtat tctcctgcta ctgcaacctg taaaccagca ctgcaatgct gatgacggg 4620
aagtagtggg atgggaacac aaatggaaag ctt 4653

```

&lt;210&gt; SEQ ID NO 122

&lt;211&gt; LENGTH: 4647

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 122

ggtaccgcgc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac   60
gtgcttgggc gcctgccgcc tgccctgccgc atgcttgtgc tggtaggct gggcagtgct   120
gccatgctga ttgaggcttg gttcatcggg tggaaactta tgtgtgtgct gggcttgcat   180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc   240
tccagtggtg tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag   300
cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc   360
cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg   420
gaatggcacc accaagtgtg tgcgagccag catcgcgcc ccgctgttcc agctacaacg   480
gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggctgga   540
tgcaagcatg cctggcgaa acacatggtg tgcggatgct gccggctgct gctgctgctg   600
cacgcccgtt agttggcagc aggcctcagcc atgcactgga tggcagctgg gctgccactg   660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt   720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat   780
gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc   840
atgctcatg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa   900
acgcaatgca gcacagcgag cagagggcgc ggcagcagag cggcggcagc agcggcgggg   960
gccacccttc ttgcggggtc gcgccccagc cagcggatgat gcgctgatcc caaacgagtt  1020
cacattcatt tgcatgcctg gagaagcag gctggggcct ttgggctggt gcagcccgca  1080
atggaatgcg ggaccgccag gctagcagca aaggcgccct ccctactccg catcgatggt  1140
ccatagtgca ttggactgca tttgggtggg gcggccgctt gtttcttctg tgttgcaaaa  1200
cgcgccagct cagcaacctg tcccgtgggt ccccgtgcc gatgaaatcg tgtgcacgcc  1260
gatcagctga ttgccggct cgcgaaagtag gcgcccctct ttctgctcgc cctctctccg  1320
tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccgct cccccgccgc  1380
ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc  1440
cgcctgttcc cgcctgtccg cccagggccg ccccgtgctg ttctggaaga ccgacctgtc  1500
cggcgcctct aacgagctgc aggacagagg cgcgccctg tcctggctgg ccaccaccgg  1560
cgtgccctgc gccgcctgtc tggacgtggt gaccgaggcc ggcccgact ggetgctgct  1620
gggagaggtg cccggccagg acctgctgtc ctcccactg gcccccgcg agaaggtgtc  1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccact gcccttcga  1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gctggtgga  1800
ccaggacgac ctggacgagg agcaccaggg cctggcccc gccgagctgt tcgcccgcct  1860
gaaggccccc atgcccagc gcgaggacct ggtggtgacc cacggcgacg cctgctgccc  1920
caacatcatg gtggagaacg gcccttctc eggcttctc gactgcccgc gctgggctg  1980
ggccgaccgc taccaggaca tcgcccctgg caccgcgac atcgcgagg agctgggctg  2040
cgagtgggcc gaccgcttcc tgggtctgta cggcatcgcc gcccccgact cccagcgcct  2100
cgcttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcgatagtt  2160

```

-continued

---

atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacact	tgtgacctg	2220
acctgtgaat	atccctgcgc	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcgcttttg	cgagttgcta	gctgcttgtg	ctatttgcca	ataccacccc	cagcatcccc	2340
ttccctcgtt	tcatatcgct	tgcataccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcgctg	ctcctgctcc	tgtcactgac	ccctcgcaca	gccttggttt	gggctccgce	2460
tgtattctcc	tggactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccc	cgtctcgaac	agagcgcgca	gaggaacgct	2580
gaaggtctcg	cctctgctgc	acctcagcgc	ggcatacacc	acaataacca	cctgacgaat	2640
gcgcttggtt	cttcgtccat	tagcgaagcg	tccggttcac	acacgtgcca	cgttgcgag	2700
gtggcaggtg	acaatgatcg	gtggagctga	tggtcgaaac	gttcacagcc	tagggatatac	2760
gaattccttt	cttgcgctat	gacacttcca	gcaaaagta	gggcgggctg	cgagacggct	2820
tcccgcgctg	gcatgcaaca	ccgatgatgc	ttcgaccccc	cgaagctcct	tccgggctgc	2880
atgggcgctc	cgatgccgct	ccagggcgag	cgtgttttaa	atagccaggc	ccccgattgc	2940
aaagacatta	tagcgagcta	ccaaagccat	attcaaacac	ctagatcact	accacttcta	3000
cacaggccac	tcgagcttgt	gatcgcactc	cgctaagggg	gcgcctcttc	ctcttcgttt	3060
cagtcacaac	ccgcaaacac	tagtatggct	tccgcggcat	tcaccatgtc	ggcgtgcccc	3120
gcgatgactg	gcagggcccc	tggggcacgt	cgtcccgac	ggccagtcgc	cacccgectg	3180
agggggcgcg	cccccgactg	gtccatgctg	ttcgccgtga	tcaccaccat	cttcagcgcc	3240
gccgagaagc	agtggaccaa	cctggagtgg	aagcccaagc	ccaagctgcc	ccagctgctg	3300
gacgaccact	tggcctgca	cggcctgggtg	ttccgcccga	ccttcgccat	ccgctcctac	3360
gaggtgggccc	ccgaccgcag	cacctccatc	ctggccgtga	tgaaccacat	gcaggaggcc	3420
acctgaaacc	acgccaagag	cgtgggcatac	ctgggcgacg	gcttcggcac	cacctggag	3480
atgtccaagc	gcgacctgat	gtgggtgggtg	cgccgcaccc	acgtggccgt	ggagcgtac	3540
cccacctggg	gcgacaccgt	ggaggtggag	tgtgggatcg	gcgccagcgg	caacaacggc	3600
atgcgcgcg	acttctcgtt	gcgcgactgc	aagaccggcg	agatcctgac	ccgctgcacc	3660
tccctgagcg	tgtgatgaa	cacccgcacc	cgccgcctga	gcaccatccc	cgacgaggtg	3720
cgcggcgaga	tggccccgc	cttcatcgac	aacgtggccg	tgaaggacga	cgagatcaag	3780
aagctgcaga	agctgaacga	ctccaccgcc	gactacatcc	agggcgccct	gacccccgc	3840
tggaacgacc	tggacgtgaa	ccagcacgtg	aacaacctga	agtacgtggc	ctgggtgttc	3900
gagaccgtgc	ccgacagcat	cttcgagtcc	caccacatca	gctccttcac	cctggagtac	3960
cgccgcgagt	gcacccgcga	ctccgtgctg	cgccgcctga	ccaccgtgag	cgccgcgacg	4020
tccgagggcg	gcctggtgtg	cgaccacctg	ctgcagctgg	agggcggcag	cgaggtgctg	4080
cgcccccgca	ccgagtggcg	ccccaaagctg	accgactcct	tccgcggcat	cagcgtgatc	4140
cccgcgagc	cccgcgtgat	ggactacaag	gaccacgacg	gcgactacaa	ggaccacgac	4200
atcgactaca	aggacgacga	cgacaagtga	tgactcgagg	cagcagcagc	tccgatagta	4260
tgcacacact	ctggacgctg	gtcgtgtgat	ggactgttgc	cgccacactt	gctgccttga	4320
cctgtgaata	tccctgccgc	ttttatcaaa	cagcctcagt	gtgtttgatc	ttgtgtgtac	4380
gcgcttttgc	gagttgctag	ctgcttgtgc	tatttgcgaa	taccaccccc	agcatcccct	4440
tccctcgttt	catatcgctt	gcatcccaac	cgcaacttat	ctacgctgtc	ctgctatccc	4500
tcagcgtgc	tctgctcct	gctcactgcc	cctcgcaacg	ccttggtttg	ggctccgct	4560

-continued

---

```
gtattctcct ggtactgcaa cctgtaaacc agcactgcaa tgctgatgca cggaagtag 4620
tgggatggga acacaaatgg aaagctt 4647
```

```
<210> SEQ ID NO 123
<211> LENGTH: 4721
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide
```

```
<400> SEQUENCE: 123
```

```
ggtaccgccc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgcccgc tgcctgcccgc atgcttgtgc tggtaggct gggcagtgt 120
gccatgctga ttgagcttg gttcatcggg tggaaactta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcccgtgcc 240
tccaggtggt tcaatcgccg cagccagagg gatttcagat gatcgccgct acaggttgag 300
cagcagtgtc agcaaagga gcagtttggc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgcccgc tggctgttcc agctacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600
cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacaggcag cagagggcgc ggcagcagag cggcggcagc agcggcgggg 960
gccacccttc ttgcccgggc gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcattgctg gagaagcag gctggggcct ttgggctggt gcagcccgca 1080
atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140
ccatagtgca ttggactgca tttgggtggg gcggccggct gtttcttctg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt ccccctgccc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgcgaagtag gcgcctcctc ttctgctcgc cctctctccg 1320
tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct ccccgcgcgc 1380
ctgggtggag cgctgttctg gctacgactg ggcccagcag accatcggct gctccgacgc 1440
cgccgtgttc cgctgtccc cccagggcgc ccccgtgctg ttctggaaga ccgacctgtc 1500
cggcgccctg aacgagctgc aggacgaggc cggcccctg tcctggctgg ccaccaccgg 1560
cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620
gggcgaggtg cccggccagg acctgctgtc ctcccacctg gccccgcgcg agaaggtgtc 1680
catcatggcc gaccgatgac gccgcctgca caccctggac cccgccacct gcccttctga 1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcaccgcg atggaggccg gcctggtgga 1800
ccaggacgac ctggacgagg agcaccaggg cctggcccc gcccagctgt tcgcccgcct 1860
gaaggcccgc atgcccagc gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920
caacatcatg tgggagaacg gccgcttctc cggttctatc gactgcggcc gcctgggctg 1980
```

-continued

---

ggccgaccgc taccaggaca tcgcctggc caccgcgac atcgcgagg agctgggagg 2040
cgagtggggc gaccgcttcc tgggtctgta cggcatcgcc gccccgact cccagcgcat 2100
cgcttctac cgctgctgg acgagttctt ctgacaattg gcagcagcag ctgggatagt 2160
atcgacacac tctggacgct ggtcgtgta tggactgtg cgcacacact tgctgccttg 2220
acctgtgaat atccctgccc cttttatcaa acagcctcag tgtgttgat cttgtgtgta 2280
cgcgcttttg cgagttgcta gctgcttggt ctatttgcga ataccacccc cagcatcccc 2340
ttcctcgtt tcatatcgct tgcacccaa cgcgaactta tctacgctgt cctgctatcc 2400
ctcagcgctg ctctgctcc tgctcactgc cctcgcaca gccttggttt gggctccgcc 2460
tgtattctcc tggactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520
gtgggatggg aacacaaatg gaggatccc cgtctcgaac agagcgcgca gaggaacgct 2580
gaaggtctcg cctctgctgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttgggt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatata 2760
gaattccttt cttgctctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccggcget gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgactact accacttcta 3000
cacaggccac tcgagcttgt gatcgactc cgctaagggg gcgctcttc ctcttctgtt 3060
cagtcacaac ccgcaaacgg cgcgcatgg ccaccacctc cctggcctcc gccttctgca 3120
gcatgaaggc cgtgatgctg gcccgcgacg gccgcggcat gaagccccgc tccagcgacc 3180
tgcagctgcg cgcggcaac gcccgacct cctgaagat gatcaacggc accaagttct 3240
cctacaccga gagcctgaag aagctgccc actggtccat gctgttcgcc gtgatcacca 3300
ccatcttctc cgcgcggcag aagcagtgga ccaacctgga gtggaagccc aagcccacc 3360
ccccccagct gctggacgac cacttggcc cccacggcct ggtgttcgc cgcacctcg 3420
ccatccgag ctacgaggtg gccccgacc gctccaccag catcgtggcc gtgatgaacc 3480
acctgcagga ggccgctg aaccacgcca agtccgtggg catcctgggc gacggcttcg 3540
gcaccacctt ggagatgtcc aagcgcgacc tgatctgggt ggtgaagcgc acccacgtgg 3600
ccgtggagcg ctaccccgcc tggggcgaca ccgtggaggt ggagtgtctg gtgggcgctt 3660
ccggcaacaa cggccgccc caccgacttc tggtgccgca ctgcaagacc ggcgagatcc 3720
tgaccgctg cacctccctg agcgtgatga tgaacaccgc caccgcccgc ctgagcaaga 3780
tccccgagga ggtgcgcgcc gagatcgcc ccgccttcat cgacaactg gccgtgaagg 3840
acgaggagat caagaagccc cagaagctga acgactccac cgcgactac atccagggcg 3900
gcctgacccc ccgctggaac gacctggaca tcaaccagca cgtgaacaac atcaagtacg 3960
tggactggat cctggagacc gtgcccgaca gcatcttcca gagccaccac atctctcct 4020
tcaccatcga gtaccgccc gagtgacca tggacagcgt gctgcagtcc ctgaccaccg 4080
tgagcggcgg ctctccgag gccggcctgg tgtgcgagca cctgctgcag ctggagggcg 4140
gcagcgaggt gctgcgccc aagaccgagt ggcgccccaa gctgaccgac tccttccgcg 4200
gcatcagcgt gatccccgc gagtccagc tgatggacta caaggaccac gacggcgact 4260
acaaggacca cgacatcgac tacaaggacg acgacgaca gtgatgactc gaggcagcag 4320
cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg ttgcccacc 4380

-continued

---

```

acttgetgccc ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt 4440
gatcttgtgt gtacgcgctt ttgcgagtgt ctagnetgctt gtgctatttg cgaataccac 4500
ccccagcate cccttcctc gtttcataac gcttgcaccc caaccgcaac ttatctacgc 4560
tgtctgcta tcctcagcgc ctgctcctgc tctgtctcac tgccctcgc acagccttgg 4620
tttgggtccc gcctgtatcc tctgtgtact gcaacctgta aaccagcact gcaatgctga 4680
tgcacgggaa gtagtgggat gggaaacaaa atggaaagct t 4721

```

&lt;210&gt; SEQ ID NO 124

&lt;211&gt; LENGTH: 4650

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 124

```

ggtaaccgccc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgcgcc tgcctgccgc atgcttgtgc tggtaggct gggcagtgt 120
gccatgctga ttgagcttg gttcatcggg tggaaactta tgtgtgtgt gggcttgcac 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgctgtgccc 240
tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300
cagcagtgtc agcaaagga gcagtttggc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgcgcc tggctgttcc agctacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gctgtgtgctg 600
cacgcccgtt agttggcagc aggctcagcc atgcactgga tggcagctgg gctgcccactg 660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggtg cttgctgggt 720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacaggcag cagagggcgc ggcagcagag cggcggcagc agcggcgggg 960
gccacccttc ttgcccgggt gcgcccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcattcctg gagaagcag gctggggcct ttgggctggt gcagcccgc 1080
atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcagatgt 1140
ccatagtgca ttggactgca tttgggtggg gcggccggct gtttcttctg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt ccccctgccc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgcgaagtag gcgcctcct ttctgctcgc cctctctccg 1320
tcccgcctct agaatatcaa tgatcagaca ggacggcctc cacgccggct cccccgccgc 1380
ctgggtggag cgctgttctg gctacgactg ggcccagcag accatcggct gctccgacgc 1440
cgccgtgttc cgctgtcgc ccagggcgc cccctgctg ttctggaaga ccgacctgtc 1500
cggcgccttg aacgagctgc aggacgaggc cgcgccctg tcctggctgg ccaccaccgg 1560
cgtgcccctg gcgcgcctg tggacgtggt gaccgagcc ggccgcgact ggctgtgtgt 1620
gggcgaggtg ccggccagc acctgctgtc ctcccacctg gccccgcgc agaaggtgtc 1680

```

-continued

---

catcatggcc	gacgccatgc	gcccgcctgca	caccctggac	cccgccacct	gccccttcga	1740
ccaccaggcc	aagcaccgca	tcgagcgcgc	cgcacccgc	atggaggccg	gectggtgga	1800
ccaggacgac	ctggacgagg	agcaccaggg	cctggccccc	gccgagctgt	tcgcccgcct	1860
gaaggcccg	atgcccga	gcgaggacct	ggtggtgacc	cacggcgacg	cctgcctgcc	1920
caacatcatg	gtggagaacg	gcccgttctc	eggcttcac	gactgcggcc	gectgggctg	1980
ggccgaccgc	taccaggaca	tcgccctggc	caccgcgac	atgcccgagg	agctgggctg	2040
cgagtgggcc	gaccgcttcc	tggtgctgta	cggcacgccc	gccccgact	cccagcgcat	2100
cgcttctac	cgctgctgg	acgagttctt	ctgacaattg	gcagcagcag	ctcggatagt	2160
atcgacacac	tctggacgct	ggtcgtgtga	tggactgttg	ccgccacct	tgtgccttg	2220
acctgtgaat	atccctgccg	cttttatcaa	acagcctcag	tgtgtttgat	cttgtgtgta	2280
cgcgcttttg	cgagtgtcta	gctgcttgtg	ctatttgcca	ataccacccc	cagcatcccc	2340
ttccctcgtt	tcatatcgct	tgcatcccaa	ccgcaactta	tctacgctgt	cctgctatcc	2400
ctcagcgcctg	ctcctgctcc	tgctcactgc	ccctcgcaca	gccttggttt	gggctccgcc	2460
tgtattctcc	tggtactgca	acctgtaaac	cagcactgca	atgctgatgc	acgggaagta	2520
gtgggatggg	aacacaaatg	gaggatcccg	cgtctcgaac	agagcgcgca	gaggaacgct	2580
gaaggtctcg	cctctgtcgc	acctcagcgc	ggcatacacc	acaataacca	cctgacgaat	2640
gcgcttggtt	cttcgtccat	tagcgaagcg	tccggttcac	acacgtgcca	cgttggcgag	2700
gtggcaggtg	acaatgatcg	gtggagctga	tggtcgaaac	gttcacagcc	tagggatata	2760
gaattccttt	cttgcgctat	gacacttcca	gcaaaagta	gggcgggctg	cgagacggct	2820
tcccggcgct	gcatgcaaca	ccgatgatgc	ttcgaccccc	cgaagctcct	tcggggctgc	2880
atgggcgctc	cgatgccgct	ccaggggcag	cgctgtttaa	atagccaggc	ccccgatgct	2940
aaagacatta	tagcgagcta	ccaaagccat	attcaaacac	ctagatcact	accacttcta	3000
cacaggccac	tcgagcttgt	gatcgcactc	cgctaagggg	gcgcctcttc	ctcttcgttt	3060
cagtcacaac	ccgcaaacac	tagtatgacg	ttcggggctg	ccctcccggc	catgggccgc	3120
ggtgtctccc	ttcccggccc	cagggctcgc	gtgcgcgccc	agtcggcgag	tcaggttttg	3180
gagagcgggc	gcccctccga	ctgggtccatg	ctggtcgcgg	tgatcaccac	catcttctcc	3240
gcccgcgaga	agcagtgga	caacctggag	tggaagccca	agcccaaccc	ccccagctg	3300
ctggacgacc	acttcggccc	ccacggcctg	gtgttcggcc	gcaccttcgc	catccgcagc	3360
tacgaggtgg	gccccgaccg	ctccaccagc	atcgtggccg	tgatgaacca	cctgcaggag	3420
gcccgcctga	accacgcaa	gtccgtgggc	atcctggggc	acggcttcgg	caccacctg	3480
gagatgtcca	agcgcgacct	gatctgggtg	gtgaagcgca	cccacgtggc	cgtggagcgc	3540
taccccgcct	ggggcgacac	cgtggaggtg	gagtgctggg	tgggcgcctc	cggaacaac	3600
ggccgcgccc	acgacttctc	ggtgcgcgac	tgcaagaccg	gcgagacct	gacccgctgc	3660
acctccctga	gcgtgatgat	gaacacccgc	acccgcgccc	tgagcaagat	ccccgaggag	3720
gtgcgcggcg	agatcggccc	cgcttctac	gacaacgtgg	ccgtgaagga	cgaggagatc	3780
agaagcccc	agaagctgaa	cgactccacc	gccgactaca	tccagggcgg	cctgaccccc	3840
cgctggaacg	acctggacat	caaccagcac	gtgaacaaca	tcaagtacgt	ggactggatc	3900
ctggagaccg	tgcccagacg	catcttcgag	agccaccaca	tctcctcctt	caccatcgag	3960
taccgcgcg	agtgacccat	ggacagcgtg	ctgcagctcc	tgaccacctg	gagcggcggc	4020
tcctccgagg	ccggcctggt	gtgcgagcac	ctgctgcagc	tggagggcgg	cagcaggtg	4080

-continued

---

```

ctgcgcgcca agaccgagtg gcgccccaaag ctgaccgact ccttccgcgg catcagcgtg 4140
atccccgccc agtccagcgt gatggactac aaggaccacg acggcgacta caaggaccac 4200
gacatcgact acaaggacga cgacgacaag tgatgactcg aggcagcagc agctcggata 4260
gtatcgacac actctggacg ctggtcgtgt gatggactgt tgccgcccaca cttgctgcct 4320
tgacctgtga atatccctgc cgcttttata aaacagcctc agtgtgtttg atcttgtgtg 4380
tacgcgcttt tgcgagttag tagctgcttg tgctatttgc gaataccacc cccagcatcc 4440
ccttccctcg tttcatatcg cttgcatccc aaccgcaact tatctacgct gtectgctat 4500
ccctcagcgc tgctcctgct cctgctcact gccctcgcga cagccttggg ttgggctccg 4560
cctgtattct cctggtagct caacctgtaa accagcactg caatgctgat gcacgggaag 4620
tagtgggatg ggaacacaaa tggaaagctt 4650

```

&lt;210&gt; SEQ ID NO 125

&lt;211&gt; LENGTH: 4653

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 125

```

ggtaccggcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgcccgc tgcctgccgc atgcttgtgc tggtaggct gggcagtgct 120
gccatgctga ttgaggcttg gttcatcggg tggaagetta tgtgtgtgct gggcttgcac 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240
tccaggtggt tcaatcgccg cagccagagg gatttcagat gatcgcgctg acaggttgag 300
cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcagccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agtacaacg 480
gcaggagtca tccaacgtaa ccatgagctg atcaaacctg caatcatcgg gcgggctgta 540
tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgctg 600
cacgcccgtt agttggcagc aggctcagcc atgcaactga tggcagctgg gctgccactg 660
caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720
tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780
gccaaaaccg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840
atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900
acgcaatgca gcacaggcag cagagggcgc ggcagcagag cggcggcagc agcggcgggg 960
gccacccttc ttgcccggtc gcgccccagc cagcgggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcattgctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080
atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatggt 1140
ccatagtgca ttggactgca tttgggtggg gcgcccggct gtttctttcg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgcgaagtag gcgcctcct ttctgctcgc cctctctccg 1320
tcccgcctct agaatatcaa tgatecagca ggacggcctc cacgcccggct cccccgcgcg 1380
ctgggtggag cgctgttctg gctacgactg ggcccagcag accatcggtg gctccgacgc 1440
cgccgtgttc cgctgtctcg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgct 1500

```



-continued

---

cgggccctg aacgagctgc aggacgaggc cgcccgcctg tcttggctgg ccaccaccgg	1560
cgtgccctgc gccgcccgtgc tggacgtggt gaccgaggcc ggccgcgact ggetgctgct	1620
gggagaggtg cccggcccagg acctgctgtc ctcccacctg gccccgcgag agaaggtgtc	1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccttcga	1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcttgggtga	1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct	1860
gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc	1920
caacatcatg gtggagaacg gccgcttctc cggcttcacg gactgcggcc gcttgggctg	1980
ggccgaccgc taccaggaca tcgccctggc caccgcgacg atcgcgagg agctgggctg	2040
cgagtgggcc gaccgcttcc tgggtctgta cggcatcgcc gccccgact cccagcgc	2100
cgcttctac cgctgctgg acgagttctt ctgacaattg gcagcagcag ctgggatagt	2160
atcgacacac tctggacgct ggtctgtgta tggactgttg ccgccacct tgetgccttg	2220
acctgtgaat atccctgcgc cttttatcaa acagcctcag tgtgtttgat cttgtgtgta	2280
cgcttcttg cgagttgcta gctgcttggt ctatttgcga ataccacccc cagcatcccc	2340
ttccctcgtt tcatatcgct tgcattccaa ccgcaactta tctacgctgt cctgctatcc	2400
ctcagcctgc ctctgctcc tgctcactgc cctcgcaca gccttggttt gggctccgcc	2460
tgtattctcc tggactgca acctgtaaac cagcactgca atgctgatgc acgggaagta	2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaaagct	2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat	2640
gcgcttgggt cttcgtccat tagcgaagcg tccggttcac acacgtgccg cgttggcgag	2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatc	2760
gaattccttt cttgcctat gacacttcca gcaaaagta gggcgggctg cgagacggct	2820
tcccgccgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc	2880
atgggcgctc cgatgccgct ccaggggcag cgctgtttaa atagccaggc ccccgattgc	2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctatgact accacttcta	3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgctcttc ctcttcgttt	3060
cagtcaaac ccgcaaacac tagtatggct atcaagacga acagggcagc tgtggagaag	3120
cctccgttca cgatcgggac gctgcgcaag gccatccccg cgcactgttt cgagcgcctg	3180
gcgcttcgtg ggccgcccc cgactggctc atgctgttcg ccgtgatcac caccatcttc	3240
tccgcccgcg agaagcagtg gaccaacctg gagtggaagc ccaagcccaa cccccccag	3300
ctgctggacg accacttcgg cccccacggc ctggtgttcc gccgcacctt cgcctccgc	3360
agctacgagg tgggccccga ccgctccacc agcatcgtgg ccgtgatgaa ccacctgcag	3420
gaggccgccc tgaaccacgc caagtccgtg ggcatcctgg gcgacggctt cggcaccacc	3480
ctggagatgt ccaagcgcga cctgatctgg gtggtgaagc gcacccacgt ggcctggag	3540
cgtaacccgc cctggggcga caccgtggag gtggagtgtt gggggggcgc ctccggcaac	3600
aacggccgcc gccacgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc	3660
tgacacctcc tgagcgtgat gatgaacacc cgcacccgcc gcctgagcaa gatccccgag	3720
gaggtcgag gcgagatcgg cccgccttc atcgacaacg tggccgtgaa ggacgaggag	3780
atcaagaagc cccagaagct gaacgactcc accgcccact acatccaggg cggcctgacc	3840
ccccgctgga acgacctgga catcaaccag cacgtgaaca acatcaagta cgtggactgg	3900

-continued

---

```

atcctggaga ccgtgcccga cagcatcttc gagagccacc acatctcttc cttcaccatc 3960
gagtaccgcc gcgagtgcac catggacagc gtgctgcagt ccctgaccac cgtgagcggc 4020
ggctcctccg aggccggcct ggtgtgagag cacctgctgc agctggaggg cggcagcgag 4080
gtgctgcgcg ccaagaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc 4140
gtgatccccg ccgagtccag cgtgatggac tacaaggacc acgacggcga ctacaaggac 4200
cacgacatcg actacaagga cgcgacgac aagtgatgac tgcaggcagc agcagctcgg 4260
atagtatcga cacactctgg acgctggtcg tgtgatggac tgttgccgcc acacttctgt 4320
ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 4380
gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcaataacc acccccagca 4440
tccccttccc tcgtttcata tcgcttgcac cccaaccgca acttatctac gctgtcctgc 4500
tatocctcag cgtgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 4560
ccgctgtat tctcctggta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 4620
aagtagtggg atgggaacac aatggaaaag ctt 4653

```

&lt;210&gt; SEQ ID NO 126

&lt;211&gt; LENGTH: 3669

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 126

```

cccgtgatca cacaggtgcc ttgagagcgt gatcacacta ttttgggggt cctacagtac 60
tgaatgggtg agaagtcgta ctgaaatcaa ggatgaacaa tgaatgggtg gctgtgggtg 120
cttctcaaaag gtcaagaatc agtcgctcgc gtcaggaaat cgcggcgctc accagcgtgg 180
gcgcggtcag tggccccgca ctggtcacca tagcctctcc tgccacagta gcgatcccc 240
ggggtttcac tctcagcagc ggctgtactg cctcccagat tttcttcttc tggacctgcg 300
ggcgtgagag gatgagcagg gtggggccaa gggctcaatc ctgaacggcc ctcattcggg 360
ttccaatccc acaacacata cccacagcag gtcagaccac gcattcgcac catgcgcacc 420
aaataacgtg tccttacctg attgggtgtg gcaggctcgg tggacaggag tgctcgtcc 480
cccgccaga cccgctcccc cgtcacggcg gcgtccggga cccgcagcgg ctccaccgcg 540
gtgtgatccg cgttggcggc gcagagcagc atcccagccg atttgacccc gcgcatgctc 600
cgaggcttga ggttggccag caccaccacc cgcgggccga caaggtcctc cagggtcacc 660
tgccggacca ggccactcac gatggtgcga gggccccctt cctcgccgag gtcgatctgc 720
tcgacgtaca gactgcgaca tgcgtggcga gtggtcatca gaaggaagca ggtgtgcaga 780
aggggcacgt ggttgggtatt gagagtagcc aaagctttgt gccaatcaga aagtcaacgc 840
agctgcctgc ctggctgcgc tacaattcct ttcttgctct atgacacttc cagcaaaagg 900
tagggcgggc tgcgagacgg cttcccggcg ctgcatgcaa caccgatgat gcttcgacce 960
cccgaagctc cttcggggct gcatggggcg tccgatgccc ctcaggggcg agcgtggtt 1020
aaatagccag gccccgatt gcaaagacat tatagcgagc taccaaagcc atattcaaac 1080
acctagatca ctaccacttc tacacaggcc actcgagctt gtgatcgcac tccgctaagg 1140
gggcgcctct tctcttctgt ttcagtcaca acccgcaaac ggcgcgcat gctgctgcag 1200
gccttctctg tctgctggc cggcttcgcc gccaaagatca gcgcctccat gacgaacgag 1260

```

-continued

---

acgtccgacc	gccccctggt	gcacttcacc	cccaacaagg	gctggatgaa	cgaccccaac	1320
ggcctgtggt	acgacgagaa	ggacgccaag	tggcacctgt	acttccagta	caacccgaac	1380
gacaccgtct	gggggacgcc	cttgttctgg	ggccacgcca	cgtccgacga	cctgaccaac	1440
tgggaggacc	agcccatcgc	catcgccccg	aagcgcaacg	actccggcgc	cttctccggc	1500
tccatggtgg	tggactaaaa	caacacctcc	ggcttcttca	acgacaccat	cgacccgcgc	1560
cagcgtcgcg	tggccatctg	gacctacaac	accccgaggt	ccgaggagca	gtacatctcc	1620
tacagcctgg	acggcggcta	caccttcacc	gagtaccaga	agaaccccgt	gctggccgcc	1680
aactccacce	agttcccgca	cccgaaggtc	ttctggtagc	agcctcccca	gaagtggatc	1740
atgaccgcgg	ccaagtccca	ggactacaag	atcgagatct	actcctccga	cgacctgaag	1800
tcctggaagc	tggagtcctc	gttcgccaac	gagggcttcc	tcggctacca	gtacgagtgc	1860
cccgccctga	tcgaggtccc	caccgagcag	gaccccagca	agtcctactg	ggtgatgttc	1920
atctccatca	accccgccgc	cccggccggc	ggctccttca	accagtactt	cgtccgcagc	1980
ttcaacggca	cccactctga	ggccttcgac	aaccagtccc	gcgtggtgga	cttcggcaag	2040
gactactacg	ccctgcagac	cttcttcaac	accgacccga	cctacgggag	cgccctgggc	2100
atcgctggg	cctccaactg	ggagtactcc	gccttcgtgc	ccaccaaccc	ctggcgctcc	2160
tccatgtccc	tcgtgcgcaa	gttctccctc	aacaccgagt	accaggccaa	cccggagacg	2220
gagctgatca	acctgaaggc	cgagccgatc	ctgaacatca	gcaacgccgg	ccctggagc	2280
cggttcgcc	ccaacaccac	gttgacgaag	gccaacagct	acaacgtcga	cctgtccaac	2340
agcaccggca	ccctggaggt	cgagctggtg	tacgcccgtc	acaccacca	gacgatctcc	2400
aaatccgtgt	tcgcccagct	ctccctctgg	ttcaagggcc	tggaggaccc	cgaggagtac	2460
ctccgcctgg	gcttcgaggt	gtcccgctcc	tccttcttcc	tggaccgcgg	gaacagcaag	2520
gtgaagtctg	tgaaggagaa	cccctacttc	accaaccgca	tgagcgtgaa	caaccagccc	2580
ttcaagagcg	agaacgacct	gtcctactac	aagggtgtacg	gcttgcctgga	ccagaacatc	2640
ctggagctgt	acttcaacga	cggcgacgtc	gtgtccacca	acacctactt	catgaccacc	2700
gggaacgccc	tgggctccgt	gaacatgacg	acgggggtgg	acaacctggt	ctacatcgac	2760
aaatccaggg	tgccgcaggt	caagtgatta	attaactcga	ggcagcagca	gctcggatag	2820
tatcgacaca	ctctggacgc	tggctggtgt	atggactgtt	gccgccacac	ttgtgcctt	2880
gacctgtgaa	tatccctgcc	gcttttatca	aacagcctca	gtgtgtttga	tcttgtgtgt	2940
acgcgctttt	gagagttgct	agctgcttgt	gctatcttgcg	aataccaccc	ccagcatccc	3000
cttccctcgt	ttcatatcgc	ttgcatccca	accgcaactt	atctacgtcg	tcctgctatc	3060
cctcagcgtc	gctcctgctc	ctgctcactg	cccctcgcac	agccttgggt	tgggctccgc	3120
ctgtattctc	ctggactactg	aacctgtaaa	ccagcactgc	aatgctgatg	cacgggaagt	3180
agtgggatgg	gaacacaaat	ggaaagcttg	agctcggtag	ccgtacccat	cagcatccgg	3240
gtgaatcttg	gcctccaaga	tatggccaat	cctcacatcc	agcttgccaa	aatcgactag	3300
actgtctgca	agtgggaatg	tggagcacia	ggttgcttgt	agcgatcgac	agactgggtg	3360
ggtaattga	caggtgggca	gcgccgcatc	catcgtgcct	gacgcgagcg	ccgcccgttg	3420
ctcgcctcgt	cctgcctgca	aagagcggca	gagaaatcgg	gaaccgaaaa	cgtcacattg	3480
cctgatgttg	ttacatgctg	gactagactt	tcttggcgtg	ggtctgctcc	tcgccaggtg	3540
cgcgacgcct	cggggctggg	tgcgagggag	ccgtgcggcc	acgcatttga	caagacccaa	3600
agctcgcctc	tcagacggtc	aaccgttctg	attatacatt	caacatattg	tacatacgca	3660

-continued

aaaagcatg

3669

<210> SEQ ID NO 127  
 <211> LENGTH: 39  
 <212> TYPE: PRT  
 <213> ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 127

Met Thr Phe Gly Val Ala Leu Pro Ala Met Gly Arg Gly Val Ser Leu  
 1 5 10 15

Pro Arg Pro Arg Val Ala Val Arg Ala Gln Ser Ala Ser Gln Val Leu  
 20 25 30

Glu Ser Gly Arg Ala Gln Leu  
 35

<210> SEQ ID NO 128  
 <211> LENGTH: 40  
 <212> TYPE: PRT  
 <213> ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 128

Met Ala Ile Lys Thr Asn Arg Gln Pro Val Glu Lys Pro Pro Phe Thr  
 1 5 10 15

Ile Gly Thr Leu Arg Lys Ala Ile Pro Ala His Cys Phe Glu Arg Ser  
 20 25 30

Ala Leu Arg Gly Arg Ala Gln Leu  
 35 40

<210> SEQ ID NO 129  
 <211> LENGTH: 36  
 <212> TYPE: PRT  
 <213> ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 129

Met Ala Ser Ala Ala Phe Thr Met Ser Ala Cys Pro Ala Met Thr Gly  
 1 5 10 15

Arg Ala Pro Gly Ala Arg Arg Ser Gly Arg Pro Val Ala Thr Arg Leu  
 20 25 30

Arg Gly Arg Ala  
 35

<210> SEQ ID NO 130  
 <211> LENGTH: 40  
 <212> TYPE: PRT  
 <213> ORGANISM: Chlorella protothecoides

&lt;400&gt; SEQUENCE: 130

Met Ala Thr Ala Ser Thr Phe Ser Ala Phe Asn Ala Arg Cys Gly Asp  
 1 5 10 15

Leu Arg Arg Ser Ala Gly Ser Gly Pro Arg Arg Pro Ala Arg Pro Leu  
 20 25 30

Pro Val Arg Gly Arg Ala Gln Leu  
 35 40

<210> SEQ ID NO 131  
 <211> LENGTH: 87  
 <212> TYPE: PRT  
 <213> ORGANISM: Cuphea hookeriana

&lt;400&gt; SEQUENCE: 131

Met Val Ala Ala Ala Ala Ser Ser Ala Phe Phe Pro Val Pro Ala Pro  
 1 5 10 15

-continued

Gly Ala Ser Pro Lys Pro Gly Lys Phe Gly Asn Trp Pro Ser Ser Leu  
 20 25 30  
 Ser Pro Ser Phe Lys Pro Lys Ser Ile Pro Asn Gly Gly Phe Gln Val  
 35 40 45  
 Lys Ala Asn Asp Ser Ala His Pro Lys Ala Asn Gly Ser Ala Val Ser  
 50 55 60  
 Leu Lys Ser Gly Ser Leu Asn Thr Gln Glu Asp Thr Ser Ser Ser Pro  
 65 70 75 80  
 Pro Pro Arg Thr Phe Leu His  
 85

<210> SEQ ID NO 132  
 <211> LENGTH: 60  
 <212> TYPE: PRT  
 <213> ORGANISM: Umbellularia californica

<400> SEQUENCE: 132

Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val  
 1 5 10 15  
 Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu  
 20 25 30  
 Gln Leu Arg Ala Gly Asn Ala Pro Thr Ser Leu Lys Met Ile Asn Gly  
 35 40 45  
 Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Arg Leu  
 50 55 60

<210> SEQ ID NO 133  
 <211> LENGTH: 60  
 <212> TYPE: PRT  
 <213> ORGANISM: Cinnamomum camphora

<400> SEQUENCE: 133

Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val  
 1 5 10 15  
 Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu  
 20 25 30  
 Gln Leu Arg Ala Gly Asn Ala Gln Thr Ser Leu Lys Met Ile Asn Gly  
 35 40 45  
 Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Lys Leu  
 50 55 60

<210> SEQ ID NO 134  
 <211> LENGTH: 1104  
 <212> TYPE: DNA  
 <213> ORGANISM: Prototheca moriformis

<400> SEQUENCE: 134

atggcaccga ccagcctgct tgccagtact ggcgtctctt ccgcttctct gtggctctct 60  
 gcgcgctcca gcgcgtgcgc ttttccggtg gatcatgcgg tccgtggcgc accgcagcgg 120  
 ccgctgcccc tgcagcgcgc ctgcttccga acagtggcgg tcagggcgc acccgcggta 180  
 gccgtccgct cggaaccgc ccaagagttt tgggagcagc ttgagccctg caagatggcg 240  
 gaggacaagc gcatcttctt ggaggagcac cgcattcggg gcaacgaggt gggcccctcg 300  
 cagcggctga cgatcacggc ggtggccaac atcctgcagg aggcggcggg caaccacgcg 360  
 gtggccatgt ggggccggag ctcggagggt ttcgcgacgg acccggagct gcaggaggcg 420  
 ggtctcatct ttgtgatgac gcgcatgcag atccaaatgt accgctaccc gcgctggggc 480

-continued

---

```

gacctgatgc aggtggagac ctggttccag acggcgggca agctaggcgc gcagcgcgag 540
tgggtgctgc gcgacaagct gaccggcgag gcgctgggcg cggccacctc cagctgggtc 600
atgatcaaca tccgcacgcg ccggccgtgc cgcattgccg agctcgtccg cgtcaagtgc 660
gccttcttcg cgcgcgagcc gccgcgcctg gcgctgccgc ccacggtcac gcgcgccaaag 720
ctgccaaca tgcgcagccc gccgcgcctg cgcgggcacc gccaggtcgc gcgccgcacc 780
gacatggaca tgaacgggca cgtgaacaac gtggcctacc tggcctggtg cctggaggcc 840
gtgcccgagc acgtcttcag cgactaccac ctctaccaga tggagatcga cttcaaggcc 900
gagtgcacg cgggcgacgt catctcctcc caggccgagc agatcccgcc ccaggaggcg 960
ctcagcaca acggcgcggc ccgcaacccc tctgcttcg tccatagcat tctgcgcgcc 1020
gagaccgagc tcgtccgcgc gcgaaccaca tggtcggccc ccacgcagc gcccgccgcc 1080
aagccgcca aggcgagcca ctga 1104

```

&lt;210&gt; SEQ ID NO 135

&lt;211&gt; LENGTH: 1104

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Prototheca moriformis*

&lt;400&gt; SEQUENCE: 135

```

atggcaccga ccagcctgct tgcccgtact ggcgtctctt ccgcttctct gtgctcctct 60
acgcgctccg gcgcgtgcgc ttttcgggtg gatcatgegg tccgtggcgc accgcagcgg 120
ccgctgcccc tgcagcgcgc ctgcttccga acagtggctg tcaggggcgc acccgcagta 180
gccgtccgtc cggaaaccgc ccaagagttt tgggagcagc ttgagccctg caagatggcg 240
gaggacaagc gcatcttctt ggaggagcac cgcattcgtg gcaacgaggt gggcccctcg 300
cagcggctga cgatcacggc ggtggccaac atcctgcagg aggcggcggg caaccacgcg 360
gtggccatgt ggggtcggag ctcggagggt ttcgcgacgg acccgagct gcaggaggcg 420
ggcctcatct ttgtgatgac gcgcatgcag atccaaatgt accgctacc gcgctggggc 480
gacctgatgc aggtggagac ctggttccag acggcgggca agctaggcgc gcagcgcgag 540
tgggtgctgc gcgacaagct gaccggcgag gcgctgggcg cggccacctc cagctgggtc 600
atgatcaaca tccgcacgcg ccggccgtgc cgcattgccg agctcgtccg cgtcaagtgc 660
gccttcttcg cgcgcgagcc gccgcgcctg gcgctgccgc ccgcggtcac gcgtgccaaag 720
ctgccaaca tgcgcagccc gccgcgcctg cgcgggcacc gccaggtcgc gcgccgcacc 780
gacatggaca tgaacgggca cgtgaacaac gttgcctacc tggcctggtg cctggaggcc 840
gtgcccgagc acgtcttcag cgactaccac ctctaccaga tggagatcga cttcaaggcc 900
gagtgcacg cgggcgacgt catctcctcc caggccgagc agatcccgcc ccaggaggcg 960
ctcagcaca acggcgcggc ccgcaacccc tctgcttcg tccatagcat tctgcgcgcc 1020
gagaccgagc tcgtccgcgc gcgaaccaca tggtcggccc ccacgcagc gcccgccgcc 1080
aagccgcca aggcgagcca ctga 1104

```

&lt;210&gt; SEQ ID NO 136

&lt;211&gt; LENGTH: 367

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Prototheca moriformis*

&lt;400&gt; SEQUENCE: 136

```

Met Ala Pro Thr Ser Leu Leu Ala Ser Thr Gly Val Ser Ser Ala Ser
1           5             10           15

```

```

Leu Trp Ser Ser Ala Arg Ser Ser Ala Cys Ala Phe Pro Val Asp His

```

-continued

20					25					30					
Ala	Val	Arg	Gly	Ala	Pro	Gln	Arg	Pro	Leu	Pro	Met	Gln	Arg	Arg	Cys
		35					40					45			
Phe	Arg	Thr	Val	Ala	Val	Arg	Ala	Ala	Pro	Ala	Val	Ala	Val	Arg	Pro
		50					55					60			
Glu	Pro	Ala	Gln	Glu	Phe	Trp	Glu	Gln	Leu	Glu	Pro	Cys	Lys	Met	Ala
		65					70					75			80
Glu	Asp	Lys	Arg	Ile	Phe	Leu	Glu	Glu	His	Arg	Ile	Arg	Gly	Asn	Glu
				85					90					95	
Val	Gly	Pro	Ser	Gln	Arg	Leu	Thr	Ile	Thr	Ala	Val	Ala	Asn	Ile	Leu
				100					105					110	
Gln	Glu	Ala	Ala	Gly	Asn	His	Ala	Val	Ala	Met	Trp	Gly	Arg	Ser	Ser
				115					120					125	
Glu	Gly	Phe	Ala	Thr	Asp	Pro	Glu	Leu	Gln	Glu	Ala	Gly	Leu	Ile	Phe
				130					135					140	
Val	Met	Thr	Arg	Met	Gln	Ile	Gln	Met	Tyr	Arg	Tyr	Pro	Arg	Trp	Gly
				145					150					155	
Asp	Leu	Met	Gln	Val	Glu	Thr	Trp	Phe	Gln	Thr	Ala	Gly	Lys	Leu	Gly
				165					170					175	
Ala	Gln	Arg	Glu	Trp	Val	Leu	Arg	Asp	Lys	Leu	Thr	Gly	Glu	Ala	Leu
				180					185					190	
Gly	Ala	Ala	Thr	Ser	Ser	Trp	Val	Met	Ile	Asn	Ile	Arg	Thr	Arg	Arg
				195					200					205	
Pro	Cys	Arg	Met	Pro	Glu	Leu	Val	Arg	Val	Lys	Ser	Ala	Phe	Phe	Ala
				210					215					220	
Arg	Glu	Pro	Pro	Arg	Leu	Ala	Leu	Pro	Pro	Thr	Val	Thr	Arg	Ala	Lys
				225					230					235	
Leu	Pro	Asn	Ile	Ala	Thr	Pro	Ala	Pro	Leu	Arg	Gly	His	Arg	Gln	Val
				245					250					255	
Ala	Arg	Arg	Thr	Asp	Met	Asp	Met	Asn	Gly	His	Val	Asn	Asn	Val	Ala
				260					265					270	
Tyr	Leu	Ala	Trp	Cys	Leu	Glu	Ala	Val	Pro	Glu	His	Val	Phe	Ser	Asp
				275					280					285	
Tyr	His	Leu	Tyr	Gln	Met	Glu	Ile	Asp	Phe	Lys	Ala	Glu	Cys	His	Ala
				290					295					300	
Gly	Asp	Val	Ile	Ser	Ser	Gln	Ala	Glu	Gln	Ile	Pro	Pro	Gln	Glu	Ala
				305					310					315	
Leu	Thr	His	Asn	Gly	Ala	Gly	Arg	Asn	Pro	Ser	Cys	Phe	Val	His	Ser
				325					330					335	
Ile	Leu	Arg	Ala	Glu	Thr	Glu	Leu	Val	Arg	Ala	Arg	Thr	Thr	Trp	Ser
				340					345					350	
Ala	Pro	Ile	Asp	Ala	Pro	Ala	Ala	Lys	Pro	Pro	Lys	Ala	Ser	His	
				355					360					365	

&lt;210&gt; SEQ ID NO 137

&lt;211&gt; LENGTH: 367

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Prototheca moriformis

&lt;400&gt; SEQUENCE: 137

Met	Ala	Pro	Thr	Ser	Leu	Leu	Ala	Arg	Thr	Gly	Val	Ser	Ser	Ala	Ser
				5					10					15	
Leu	Cys	Ser	Ser	Thr	Arg	Ser	Gly	Ala	Cys	Ala	Phe	Pro	Val	Asp	His
				20				25						30	
Ala	Val	Arg	Gly	Ala	Pro	Gln	Arg	Pro	Leu	Pro	Met	Gln	Arg	Arg	Cys

-continued

35				40				45							
Phe	Arg	Thr	Val	Ala	Val	Arg	Ala	Ala	Pro	Ala	Val	Ala	Val	Arg	Pro
50						55				60					
Glu	Pro	Ala	Gln	Glu	Phe	Trp	Glu	Gln	Leu	Glu	Pro	Cys	Lys	Met	Ala
65				70						75					80
Glu	Asp	Lys	Arg	Ile	Phe	Leu	Glu	Glu	His	Arg	Ile	Arg	Gly	Asn	Glu
				85					90					95	
Val	Gly	Pro	Ser	Gln	Arg	Leu	Thr	Ile	Thr	Ala	Val	Ala	Asn	Ile	Leu
			100					105					110		
Gln	Glu	Ala	Ala	Gly	Asn	His	Ala	Val	Ala	Met	Trp	Gly	Arg	Ser	Ser
		115					120					125			
Glu	Gly	Phe	Ala	Thr	Asp	Pro	Glu	Leu	Gln	Glu	Ala	Gly	Leu	Ile	Phe
	130					135					140				
Val	Met	Thr	Arg	Met	Gln	Ile	Gln	Met	Tyr	Arg	Tyr	Pro	Arg	Trp	Gly
145				150					155						160
Asp	Leu	Met	Gln	Val	Glu	Thr	Trp	Phe	Gln	Thr	Ala	Gly	Lys	Leu	Gly
				165					170					175	
Ala	Gln	Arg	Glu	Trp	Val	Leu	Arg	Asp	Lys	Leu	Thr	Gly	Glu	Ala	Leu
			180					185					190		
Gly	Ala	Ala	Thr	Ser	Ser	Trp	Val	Met	Ile	Asn	Ile	Arg	Thr	Arg	Arg
		195					200					205			
Pro	Cys	Arg	Met	Pro	Glu	Leu	Val	Arg	Val	Lys	Ser	Ala	Phe	Phe	Ala
	210					215					220				
Arg	Glu	Pro	Pro	Arg	Leu	Ala	Leu	Pro	Pro	Ala	Val	Thr	Arg	Ala	Lys
225				230						235					240
Leu	Pro	Asn	Ile	Ala	Thr	Pro	Ala	Pro	Leu	Arg	Gly	His	Arg	Gln	Val
				245					250					255	
Ala	Arg	Arg	Thr	Asp	Met	Asp	Met	Asn	Gly	His	Val	Asn	Asn	Val	Ala
			260					265					270		
Tyr	Leu	Ala	Trp	Cys	Leu	Glu	Ala	Val	Pro	Glu	His	Val	Phe	Ser	Asp
		275				280					285				
Tyr	His	Leu	Tyr	Gln	Met	Glu	Ile	Asp	Phe	Lys	Ala	Glu	Cys	His	Ala
		290				295				300					
Gly	Asp	Val	Ile	Ser	Ser	Gln	Ala	Glu	Gln	Ile	Pro	Pro	Gln	Glu	Ala
305					310					315					320
Leu	Thr	His	Asn	Gly	Ala	Gly	Arg	Asn	Pro	Ser	Cys	Phe	Val	His	Ser
				325					330					335	
Ile	Leu	Arg	Ala	Glu	Thr	Glu	Leu	Val	Arg	Ala	Arg	Thr	Thr	Trp	Ser
			340					345					350		
Ala	Pro	Ile	Asp	Ala	Pro	Ala	Ala	Lys	Pro	Pro	Lys	Ala	Ser	His	
		355					360					365			

&lt;210&gt; SEQ ID NO 138

&lt;211&gt; LENGTH: 328

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Cuphea hookeriana

&lt;400&gt; SEQUENCE: 138

Gln	Leu	Pro	Asp	Trp	Ser	Arg	Leu	Leu	Thr	Ala	Ile	Thr	Thr	Val	Phe
1				5					10					15	
Val	Lys	Ser	Lys	Arg	Pro	Asp	Met	His	Asp	Arg	Lys	Ser	Lys	Arg	Pro
			20					25					30		
Asp	Met	Leu	Val	Asp	Ser	Phe	Gly	Leu	Glu	Ser	Thr	Val	Gln	Asp	Gly
		35					40					45			
Leu	Val	Phe	Arg	Gln	Ser	Phe	Ser	Ile	Arg	Ser	Tyr	Glu	Ile	Gly	Thr





-continued

100					105					110					
Val	Glu	Arg	Tyr	Pro	Thr	Trp	Gly	Asp	Thr	Val	Glu	Val	Glu	Cys	Trp
			115				120						125		
Ile	Gly	Ala	Ser	Gly	Asn	Asn	Gly	Met	Arg	Arg	Asp	Phe	Leu	Val	Arg
			130				135						140		
Asp	Cys	Lys	Thr	Gly	Glu	Ile	Leu	Thr	Arg	Cys	Thr	Ser	Leu	Ser	Val
			145				150								160
Leu	Met	Asn	Thr	Arg	Thr	Arg	Arg	Leu	Ser	Thr	Ile	Pro	Asp	Glu	Val
				165											175
Arg	Gly	Glu	Ile	Gly	Pro	Ala	Phe	Ile	Asp	Asn	Val	Ala	Val	Lys	Asp
				180											190
Asp	Glu	Ile	Lys	Lys	Leu	Gln	Lys	Leu	Asn	Asp	Ser	Thr	Ala	Asp	Tyr
				195											205
Ile	Gln	Gly	Gly	Leu	Thr	Pro	Arg	Trp	Asn	Asp	Leu	Asp	Val	Asn	Gln
				210											220
His	Val	Asn	Asn	Leu	Lys	Tyr	Val	Ala	Trp	Val	Phe	Glu	Thr	Val	Pro
				225											240
Asp	Ser	Ile	Phe	Glu	Ser	His	His	Ile	Ser	Ser	Phe	Thr	Leu	Glu	Tyr
				245											255
Arg	Arg	Glu	Cys	Thr	Arg	Asp	Ser	Val	Leu	Arg	Ser	Leu	Thr	Thr	Val
				260											270
Ser	Gly	Gly	Ser	Ser	Glu	Ala	Gly	Leu	Val	Cys	Asp	His	Leu	Leu	Gln
				275											285
Leu	Glu	Gly	Gly	Ser	Glu	Val	Leu	Arg	Ala	Arg	Thr	Glu	Trp	Arg	Pro
				290											300
Lys	Leu	Thr	Asp	Ser	Phe	Arg	Gly	Ile	Ser	Val	Ile	Pro	Ala	Glu	Pro
				305											320
Arg	Val														

&lt;210&gt; SEQ ID NO 140

&lt;211&gt; LENGTH: 345

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Cinnamomum camphora

&lt;400&gt; SEQUENCE: 140

Pro	Asp	Trp	Ser	Met	Leu	Phe	Ala	Val	Ile	Thr	Thr	Ile	Phe	Ser	Ala
1				5						10				15	
Ala	Glu	Lys	Gln	Trp	Thr	Asn	Leu	Glu	Trp	Lys	Pro	Lys	Pro	Asn	Pro
			20					25					30		
Pro	Gln	Leu	Leu	Asp	Asp	His	Phe	Gly	Pro	His	Gly	Leu	Val	Phe	Arg
		35					40					45			
Arg	Thr	Phe	Ala	Ile	Arg	Ser	Tyr	Glu	Val	Gly	Pro	Asp	Arg	Ser	Thr
		50				55					60				
Ser	Ile	Val	Ala	Val	Met	Asn	His	Leu	Gln	Glu	Ala	Ala	Leu	Asn	His
		65			70					75				80	
Ala	Lys	Ser	Val	Gly	Ile	Leu	Gly	Asp	Gly	Phe	Gly	Thr	Thr	Leu	Glu
				85					90					95	
Met	Ser	Lys	Arg	Asp	Leu	Ile	Trp	Val	Val	Lys	Arg	Thr	His	Val	Ala
			100					105						110	
Val	Glu	Arg	Tyr	Pro	Ala	Trp	Gly	Asp	Thr	Val	Glu	Val	Glu	Cys	Trp
			115				120						125		
Val	Gly	Ala	Ser	Gly	Asn	Asn	Gly	Arg	Arg	His	Asp	Phe	Leu	Val	Arg
			130				135					140			
Asp	Cys	Lys	Thr	Gly	Glu	Ile	Leu	Thr	Arg	Cys	Thr	Ser	Leu	Ser	Val
				145			150				155				160

-continued

---

Met Met Asn Thr Arg Thr Arg Arg Leu Ser Lys Ile Pro Glu Glu Val  
165 170 175

Arg Gly Glu Ile Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp  
180 185 190

Glu Glu Ile Lys Lys Pro Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr  
195 200 205

Ile Gln Gly Gly Leu Thr Pro Arg Trp Asn Asp Leu Asp Ile Asn Gln  
210 215 220

His Val Asn Asn Ile Lys Tyr Val Asp Trp Ile Leu Glu Thr Val Pro  
225 230 235 240

Asp Ser Ile Phe Glu Ser His His Ile Ser Ser Phe Thr Ile Glu Tyr  
245 250 255

Arg Arg Glu Cys Thr Met Asp Ser Val Leu Gln Ser Leu Thr Thr Val  
260 265 270

Ser Gly Gly Ser Ser Glu Ala Gly Leu Val Cys Glu His Leu Leu Gln  
275 280 285

Leu Glu Gly Gly Ser Glu Val Leu Arg Ala Lys Thr Glu Trp Arg Pro  
290 295 300

Lys Leu Thr Asp Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Ser  
305 310 315 320

Ser Val Met Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp  
325 330 335

Ile Asp Tyr Lys Asp Asp Asp Asp Lys  
340 345

<210> SEQ ID NO 141  
<211> LENGTH: 4  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
peptide

<400> SEQUENCE: 141

Lys Asp Glu Leu  
1

<210> SEQ ID NO 142  
<211> LENGTH: 6  
<212> TYPE: PRT  
<213> ORGANISM: Unknown  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Unknown: Higher plant fatty  
acyl-ACP thioesterase sequence

<400> SEQUENCE: 142

Leu Asp Met Asn Gln His  
1 5

<210> SEQ ID NO 143  
<211> LENGTH: 6  
<212> TYPE: PRT  
<213> ORGANISM: Unknown  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Unknown: Algal fatty acyl-ACP  
thioesterase sequence

<400> SEQUENCE: 143

Met Asp Met Asn Gly His  
1 5

---

355

What is claimed is:

1. A method of manufacturing a chemical, the method comprising:
  - (a) cultivating a cell of an oleaginous microorganism that comprises an exogenous fatty acyl-ACP thioesterase gene that encodes a fatty acyl-ACP thioesterase having hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12, or C14 and an exogenous invertase gene, wherein the cell is cultured in a medium comprising sucrose and the cell produces triglyceride oil that has a lipid profile of at least 4% C8-C14;
  - (b) performing one or more chemical reactions on the triglyceride oil to produce the chemical; and
  - (c) separating the chemical from the reaction mixture after performing step (b).
2. A method of manufacturing a chemical, the method comprising:
  - (a) cultivating a cell of an oleaginous microorganism that comprises an exogenous fatty acyl-ACP thioesterase gene that encodes a fatty acyl-ACP thioesterase having hydrolysis activity toward one or more fatty acyl-ACP substrates of chain length C8, C10, C12, or C14, wherein the cell produces triglyceride oil that has a lipid profile of at least 4% C8-C14;
  - (b) performing one or more chemical reactions on the triglyceride oil, wherein the one or more chemical reactions comprise an amination reaction to produce the chemical, which comprises a fatty nitrogen compound; and
  - (c) separating the chemical from the reaction mixture after performing step (b).
3. The method of claim 1, wherein the fatty acyl-ACP thioesterase gene is from *Umbellularia californica*, *Cinnamomum camphora*, *Cuphea hookeriana*, *Cuphea palustris*, or *Ulmus americana*.
4. The method of claim 1, wherein the oleaginous microorganism is an obligate heterotroph.
5. The method of claim 4, wherein the triglyceride oil is blended with at least one of the following oils before performing the one or more chemical reactions: soy, rapeseed, canola, palm, palm kernel, coconut, corn, waste vegetable, Chinese tallow, olive, sunflower, cotton seed, chicken fat, beef tallow, porcine tallow, microalgae, macroalgae, *Cuphea*, flax, peanut, choice white grease, lard, *Camelina sativa*, mustard seed, cashew nut, oats, lupine kenaf, calendula, hemp, coffee, hazelnut, euphorbia, pumpkin seed, coriander, camellia, sesame, safflower, rice, tung tree, cocoa, copra, plum poppy, castor bean, pecan, jojoba, macademia, Brazil nut, avocado, petroleum, or a distillate fraction of any of the proceeding oils.

356

6. A method of manufacturing a chemical, the method comprising:
  - (a) cultivating a cell of an oleaginous microorganism that comprises an exogenous fatty acyl-ACP thioesterase gene that encodes a fatty acyl-ACP thioesterase having hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12, or C14, wherein the cell produces triglyceride oil that has a lipid profile of at least 4% C8-C14, wherein the cell further comprises an endogenous desaturase gene that is knocked out or down-regulated;
  - (b) performing one or more chemical reactions on the triglyceride oil to produce the chemical; and
  - (c) separating the chemical from the reaction mixture after performing step (b).
7. The method of claim 1, wherein the one or more chemical reactions comprise a transesterification, hydrogenation, hydrocracking, hydroprocessing, hydrotreating, fluid catalytic cracking, hydrodeoxygenation, hydrodesulfurization, epoxidation, deoxygenation, isomerization, interestification, directed interestification, hydroxylation and/or hydrolysis.
8. The method of claim 1, wherein the oleaginous microorganism is a microalga.
9. The method of claim 8, wherein the microalga is of the genus *Prototheca*.
10. The method of claim 1, wherein the chemical comprises a fatty acid, a fatty acid methyl ester, a fatty nitrile, an ester, a dimer acid, a quat, a surfactant, a fatty alkanolamide, a fatty alcohol sulfate, a resin, an emulsifier, a fatty alcohol, an olefin, a higher alkane, a linear alkylbenzyl sulfonate, or a sulfated fatty acid.
11. The method of claim 2, wherein the oleaginous microorganism is an obligate heterotroph.
12. The method of claim 2, wherein the oleaginous microorganism is a microalga.
13. The method of claim 12, wherein the microalga is an obligate heterotroph.
14. The method of claim 12, wherein the microalga is of the genus *Prototheca* or *Chlorella*.
15. The method of claim 6, wherein the oleaginous microorganism is an obligate heterotroph.
16. The method of claim 6, wherein the oleaginous microorganism is a microalga.
17. The method of claim 16, wherein the microalga is an obligate heterotroph.
18. The method of claim 16, wherein the microalga is of the genus *Prototheca* or *Chlorella*.
19. The method of claim 8, wherein the microalga is of the species *Prototheca moriformis*.

\* \* \* \* \*