

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 April 2011 (14.04.2011)

PCT

(10) International Publication Number
WO 2011/044187 A1

- (51) **International Patent Classification:**
A01N 43/54 (2006.01) *A61K 31/505* (2006.01)
- (21) **International Application Number:**
PCT/US2010/051560
- (22) **International Filing Date:**
6 October 2010 (06.10.2010)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/249,684 8 October 2009 (08.10.2009) US
- (71) **Applicant (for all designated States except US):**
SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **SCOTT, Jack, D.** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **STAMFORD, Andrew, W.** [AU/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **GILBERT, Eric, J.** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **CUMMING, Jared, N.** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **ISERLOH, Ulrich** [DE/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **WANG, Lingyan** [CN/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **LI, Wei** [CN/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US).
- (74) **Common Representative:** **SCHERING CORPORATION**; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US).

- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) **Title:** IMINOTHIADIAZINE DIOXIDE COMPOUNDS AS BACE INHIBITORS, COMPOSITIONS, AND THEIR USE



(57) **Abstract:** In its many embodiments, the present invention provides provides certain iminothiadiazone dioxide compounds, including compounds Formula (a) and include tautomers, solvates, prodrugs, esters, and deuterates thereof, and pharmaceutically acceptable salts of said compounds, tautomers, solvates, prodrugs, esters, and deuterates, wherein each of R¹, R², R³, R⁴, R⁵, R⁶, ring A, ring B, ring C, m, n, p, q, -L₁-, -L₂-, -L₃-, and -L₄- is selected independently and as defined herein. The compounds of the invention have, surprisingly and advantageously, improved solution stability. Pharmaceutical compositions comprising one or more such compounds (alone and in combination with one or more other active agents), and methods for their preparation and use in treating pathologies associated with amyloid beta (A β) protein, including Alzheimers Disease, are also disclosed.

WO 2011/044187 A1

- 2 -

protease responsible for the production of A β by β -secretase activity. The inhibition of BACE-1 has been shown to inhibit the production of A β .

AD is estimated to afflict more than 20 million people worldwide and is believed to be the most common cause of dementia. AD is a disease characterized by degeneration and loss of neurons and also by the formation of senile plaques and neurofibrillary tangles. Presently, treatment of Alzheimer's disease is limited to the treatment of its symptoms rather than the underlying causes. Symptom-improving agents approved for this purpose include, for example, N-methyl-D-aspartate receptor antagonists such as memantine (Namenda®, Forrest Pharmaceuticals, Inc.), cholinesterase inhibitors such as donepezil (Aricept®, Pfizer), rivastigmine (Exelon®, Novartis), galantamine (Razadyne Reminyl®), and tacrine (Cognex®).

In AD, A β peptides, formed through β -secretase and *gamma*-secretase activity, can form tertiary structures that aggregate to form amyloid fibrils. A β peptides have also been shown to form A β oligomers (sometimes referred to as "A β aggregates" or "Abeta oligomers"). A β oligomers are small multimeric structures composed of 2 to 12 A β peptides that are structurally distinct from A β fibrils. Amyloid fibrils can deposit outside neurons in dense formations known as senile plaques, neuritic plaques, or diffuse plaques in regions of the brain important to memory and cognition. A β oligomers are cytotoxic when injected in the brains of rats or in cell culture. This A β plaque formation and deposition and/or A β oligomer formation, and the resultant neuronal death and cognitive impairment, are among the hallmarks of AD pathophysiology. Other hallmarks of AD pathophysiology include intracellular neurofibrillary tangles comprised of abnormally phosphorylated tau protein, and neuroinflammation.

Evidence suggests that A β , A β fibrils, aggregates, oligomers, and/or plaque play a causal role in AD pathophysiology. (Ohno et al., *Neurobiology of Disease*, No. 26 (2007), 134-145). Mutations in the genes for APP and presenilins 1/2 (PS1/2) are known to cause familial AD and an increase in the production of the 42-amino acid form of A β is regarded as causative. A β has been shown to be neurotoxic in culture and in vivo. For example, when injected into the brains of aged primates, fibrillar A β causes neuronal cell death around the injection site. Other direct and circumstantial evidence of the role of A β in Alzheimer etiology has also been published.

BACE-1 has become an accepted therapeutic target for the treatment of Alzheimer's disease. For example, McConlogue et al., *J. Bio. Chem.*, Vol. 282, No. 36 (Sept. 2007), have shown that partial reductions of BACE-1 enzyme activity and concomitant reductions of A β levels lead to a dramatic inhibition of A β -driven AD-like pathology, making β -secretase a target for therapeutic intervention in AD. Ohno et al. *Neurobiology of Disease*, No. 26 (2007), 134-

- 3 -

145, report that genetic deletion of BACE-1 in 5XFAD mice abrogates A β generation, blocks amyloid deposition, prevents neuron loss found in the cerebral cortex and subiculum (brain regions manifesting the most severe amyloidosis in 5XFAD mice), and rescues memory deficits in 5XFAD mice. The group also reports that A β is ultimately responsible for neuron death in AD and concludes that BACE-1 inhibition has been validated as an approach for the treatment of AD. Roberds et al., *Human Mol. Genetics*, 2001, Vol. 10, No. 12, 1317-1324, established that inhibition or loss of β -secretase activity produces no profound phenotypic defects while inducing a concomitant reduction in A β . Luo et al., *Nature Neuroscience*, Vol. 4, No. 3, March 2001, report that mice deficient in BACE-1 have normal phenotype and abolished β -amyloid generation.

BACE-1 has also been identified or implicated as a therapeutic target for a number of other diverse pathologies in which A β or A β fragments have been identified to play a causative role. One such example is in the treatment of AD-type symptoms of patients with Down's syndrome. The gene encoding APP is found on chromosome 21, which is also the chromosome found as an extra copy in Down's syndrome. Down's syndrome patients tend to acquire AD at an early age, with almost all those over 40 years of age showing Alzheimer's-type pathology. This is thought to be due to the extra copy of the APP gene found in these patients, which leads to overexpression of APP and therefore to increased levels of A β causing the prevalence of AD seen in this population. Furthermore, Down's patients who have a duplication of a small region of chromosome 21 that does not include the APP gene do not develop AD pathology. Thus, it is thought that inhibitors of BACE-1 could be useful in reducing Alzheimer's type pathology in Down's syndrome patients.

Another example is in the treatment of glaucoma (Guo et al., *PNAS*, Vol. 104, No. 33, August 14, 2007). Glaucoma is a retinal disease of the eye and a major cause of irreversible blindness worldwide. Guo et al. report that A β colocalizes with apoptotic retinal ganglion cells (RGCs) in experimental glaucoma and induces significant RGC cell loss in vivo in a dose- and time-dependent manner. The group report having demonstrated that targeting different components of the A β formation and aggregation pathway, including inhibition of β -secretase alone and together with other approaches, can effectively reduce glaucomatous RGC apoptosis in vivo. Thus, the reduction of A β production by the inhibition of BACE-1 could be useful, alone or in combination with other approaches, for the treatment of glaucoma.

Another example is in the treatment of olfactory impairment. Getchell et al., *Neurobiology of Aging*, 24 (2003), 663-673, have observed that the olfactory epithelium, a

- 4 -

neuroepithelium that lines the posterior-dorsal region of the nasal cavity, exhibits many of the same pathological changes found in the brains of AD patients, including deposits of A β , the presence of hyperphosphorylated tau protein, and dystrophic neurites among others. Other evidence in this connection has been reported by Bacon AW, et al., *Ann NY Acad Sci* 2002; 855:723–31; Crino PB, Martin JA, Hill WD, et al., *Ann Otol Rhinol Laryngol*, 1995;104:655–61; Davies DC, et al., *Neurobiol Aging*, 1993;14:353-7; Devanand DP, et al., *Am J Psychiatr*, 2000;157:1399-405; and Doty RL, et al., *Brain Res Bull*, 1987;18:597–600. It is reasonable to suggest that addressing such changes by reduction of A β by inhibition of BACE-1 could help to restore olfactory sensitivity in patients with AD.

For compounds which are inhibitors of BACE-2, another example is in the treatment of type-II diabetes, including diabetes associated with amyloidogenesis. BACE-2 is expressed in the pancreas. BACE-2 immunoreactivity has been reported in secretory granules of beta cells, co-stored with insulin and IAPP, but lacking in the other endocrine and exocrine cell types. Stoffel et al., WO2010/063718, disclose the use of BACE-2 inhibitors in the treatment of metabolic diseases such as Type-II diabetes. The presence of BACE-2 in secretory granules of beta cells suggests that it may play a role in diabetes-associated amyloidogenesis. (Finzi, G. Franzi, et al., *Ultrastruct Pathol*. 2008 Nov-Dec;32(6):246-51.)

Other diverse pathologies characterized by the formation and deposition of A β or fragments thereof, and/or by the presence of amyloid fibrils, oligomers, and/or plaques, include neurodegenerative diseases such as scrapie, bovine spongiform encephalitis, traumatic brain injury (“TBI”), Creutzfeld-Jakob disease and the like, type II diabetes (which is characterized by the localized accumulation of cytotoxic amyloid fibrils in the insulin producing cells of the pancreas), and amyloid angiopathy. In this regard reference can be made to the patent literature. For example, Kong et al., US2008/0015180, disclose methods and compositions for treating amyloidosis with agents that inhibit A β peptide formation. As another example, Loane, et al. report the targeting of amyloid precursor protein secretases as therapeutic targets for traumatic brain injury. (Loane et al., “Amyloid precursor protein secretases as therapeutic targets for traumatic brain injury”, *Nature Medicine*, Advance Online Publication, published online March 15, 2009.) Still other diverse pathologies characterized by the inappropriate formation and deposition of A β or fragments thereof, and/or by the presence of amyloid fibrils, and/or for which inhibitor(s) of BACE-1 is expected to be of therapeutic value are discussed further hereinbelow.

- 5 -

The therapeutic potential of inhibiting the deposition of A β has motivated many groups to characterize BACE-1 and to identify inhibitors of BACE-1 and of other secretase enzyme inhibitors. Examples from the patent literature are growing and include WO2006009653, WO2007005404, WO2007005366, WO2007038271, WO2007016012, US2005/0282826, 5 US2007072925, WO2007149033, WO2007145568, WO2007145569, WO2007145570, WO2007145571, WO2007114771, US20070299087, WO2005/016876, WO2005/014540, WO2005/058311, WO2006/065277, WO2006/014762, WO2006/014944, WO2006/138195, WO2006/138264, WO2006/138192, WO2006/138217, WO2007/050721, WO2007/053506, WO2007/146225, WO2006/138230, WO2006/138265, WO2006/138266, 10 WO2007/053506, WO2007/146225, WO2008/073365, WO2008/073370, WO2008/103351, US2009/041201, US2009/041202, and WO2010/047372.

SUMMARY OF THE INVENTION


The present invention provides certain iminothiadiazine dioxide compounds 15 which are collectively or individually referred to herein as "compound(s) of the invention", as described herein. The compounds of the present invention, which contain an iminothiadiazine dioxide, have been found, surprisingly and advantageously, to exhibit improved solution stability compared to structurally similar compounds.

In each of the various embodiments of the compounds of the invention described herein, 20 each variable including those in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2), and the various embodiments thereof, each variable is selected independently of the others unless otherwise indicated.

In each of the various embodiments of the compounds of the invention described herein, 25 including those in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2), and the various embodiments thereof and the compounds of the examples, such formulas and examples are intended to encompass all forms of the compounds such as, for example, any solvates, hydrates, stereoisomers, and tautomers of said compounds and of any pharmaceutically acceptable salts thereof.

- 6 -

In one embodiment, the compounds of the invention have the structural Formula (a):



(a)

wherein:

5 -L₁- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

 -L₂- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

10 -L₃- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

 each -L₄- is independently present or absent and when present independently represents a divalent moiety independently selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, -alkynyl-, -N(R⁸)-, -N(R⁸)C(O)-, and -C(O)N(R⁸)-;

 m, n, p and q are each independently selected integers, wherein:

15 m is 0 or more,

 n is 0 or more,

 p is 0 or more,

 q is 0 or more,

20 wherein the maximum value of the sum of m and q is the maximum number of available substitutable hydrogen atoms on ring A,

 wherein the maximum value of n is the maximum number of available substitutable hydrogen atoms on ring B, and

- 7 -

wherein the maximum value of p is the maximum number of available substitutable hydrogen atoms on ring C;

R¹ is selected from the group consisting of: H, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and heteroarylalkyl-,

wherein each of said alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and heteroarylalkyl- of R¹ is unsubstituted or substituted with from 1 to 5 independently selected R¹⁰ groups;

R² is selected from the group consisting of H, alkyl, halo, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

wherein each of said alkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R² is unsubstituted or substituted with from 1 to 5 independently selected R¹⁰ groups;

R³ is selected from the group consisting of H, alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

wherein each of said alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R³ is unsubstituted or substituted with from 1 to 5 independently selected R¹⁰ groups;

ring A is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R⁴ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -NR⁸C(O)N(R⁸)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁸)₂, -S(O)R⁷, -S(O)₂R⁷, -S(O)₂N(R⁸)₂, -OR⁷, -SR⁷, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, aryl, and heteroaryl of R⁴ (when present) is optionally

- 8 -

independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, -CN, -SF₅, -OSF₅, -NO₂, -N(R⁸)₂, -OR⁷, -C(O)N(R⁸)₂, and cycloalkyl;

5 ring B is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷,
 10 -NR⁸C(O)N(R⁸)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁸)₂, -S(O)R⁷, -S(O)₂R⁷, -S(O)₂N(R⁸)₂, -OR⁷, -SR⁷, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, aryl, and heteroaryl of R⁵ (when present) is optionally
 15 independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, -CN, -SF₅, -OSF₅, -NO₂, -N(R⁸)₂, -OR⁷, -C(O)N(R⁸)₂, and cycloalkyl;

each ring C (when present) is independently selected from the group consisting of
 20 monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R⁶ (when present) is independently selected from the group consisting of alkyl, aryl, arylalkyl-, haloalkyl, cycloalkyl, cycloalkylalkyl-, heteroaryl, and heteroarylalkyl-;

each R⁷ (when present) is independently selected from the group consisting of H, alkyl,
 25 alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

each R⁸ (when present) is independently selected from the group consisting of H, alkyl, alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

each R⁹ (when present) is independently selected from the group consisting of: halogen, -
 30 CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -NR⁸C(O)N(R⁸)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁸)₂, -S(O)R⁷, -

- 9 -

$S(O)_2R^7$, $-S(O)_2N(R^8)_2$, $-OR^7$, $-SR^7$, alkyl, haloalkyl, heteroalkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, heteroaryl, heteroarylalkyl-, and heterocycloalkyl;

and

each R^{10} (when present) is independently selected from the group consisting of halo, -CN,
 5 $-NO_2$, $-Si(R^6)_3$, $-P(O)(OR^7)_2$, $-P(O)(OR^7)(R^7)$, $-N(R^8)_2$, $-NR^8C(O)R^7$, $-NR^8S(O)_2R^7$, $-$
 $NR^8C(O)N(R^8)_2$, $-NR^8C(O)OR^7$, $-C(O)R^7$, $-C(O)_2R^7$, $-C(O)N(R^8)_2$, $-S(O)R^7$, $-S(O)_2R^7$,
 $-S(O)_2N(R^8)_2$, $-OR^7$, $-SR^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
 heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
 10 heterocycloalkyl, aryl, and heteroaryl of R^{10} (when present) is optionally independently
 unsubstituted or further substituted with one or more independently selected groups
 selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower
 heteroalkyl, halo, -CN, $-NO_2$, $-N(R^8)_2$, $-OR^7$, $-C(O)N(R^8)_2$, and cycloalkyl.

15 In other embodiments, the invention provides various methods of treating, preventing,
 ameliorating, and/or delaying the onset of an amyloid β pathology ($A\beta$ pathology) and/or a
 symptom or symptoms thereof, comprising administering a composition comprising an effective
 amount of one or more compounds of the invention, or a tautomer thereof, or pharmaceutically
 acceptable salt or solvate of said compound(s) and/or said tautomer(s), to a patient in need
 20 thereof. Such methods optionally additionally comprise administering an effective amount of one
 or more additional therapeutic agents suitable for treating the patient being treated.

These and other embodiments of the invention, which are described in detail below or
 will become readily apparent to those of ordinary skill in the art, are included within the scope of
 the invention.

25


DETAILED DESCRIPTION:

In one embodiment, the compounds of the invention have the structural Formula (a) as
 described above.

In one embodiment, in Formula (a), q is 0 or 1.

- 10 -

In one embodiment, in Formula (a), q is 0 and the compounds of Formula (a) have a structure according to Formula (I):



5 wherein:

-L₁- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

-L₂- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

10 -L₃- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

m and n are each independently selected integers, wherein:

m is 0 or more,

n is 0 or more,

15 wherein the maximum value of m is the maximum number of available substitutable hydrogen atoms on ring A,

wherein the maximum value of n is the maximum number of available substitutable hydrogen atoms on ring B, and

20 R¹ is selected from the group consisting of: H, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and heteroarylalkyl-,

wherein each of said alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and

- 11 -

heteroarylalkyl- of R^1 is unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups;

R^2 is selected from the group consisting of H, alkyl, halo, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

wherein each of said alkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R^2 is unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups;

R^3 is selected from the group consisting of H, alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

wherein each of said alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R^3 is unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups;

ring A is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R^4 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R^6)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R^8)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -NR⁸C(O)N(R^8)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R^8)₂, -S(O)R⁷, -S(O)₂R⁷, -S(O)₂N(R^8)₂, -OR⁷, -SR⁷, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl of R^4 (when present) is optionally independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, -CN, -SF₅, -OSF₅, -NO₂, -N(R^8)₂, -OR⁷, -C(O)N(R^8)₂, and cycloalkyl;

ring B is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R^5 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R^6)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R^8)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷,

- 12 -

$-\text{NR}^8\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$,
 $-\text{S}(\text{O})_2\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{SR}^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl,

5 wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl of R^5 (when present) is optionally independently
unsubstituted or further substituted with one or more independently selected groups
selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower
heteroalkyl, halo, $-\text{CN}$, $-\text{SF}_5$, $-\text{OSF}_5$, $-\text{NO}_2$, $-\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, and cycloalkyl;
each R^6 (when present) is independently selected from the group consisting of alkyl, aryl,
10 arylalkyl-, haloalkyl, cycloalkyl, cycloalkylalkyl-, heteroaryl, and heteroarylalkyl-;

each R^7 (when present) is independently selected from the group consisting of H, alkyl,
alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl,
cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

15 each R^8 (when present) is independently selected from the group consisting of H, alkyl,
alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl,
cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;


and

20 each R^{10} (when present) is independently selected from the group consisting of halo, $-\text{CN}$,
 $-\text{NO}_2$, $-\text{Si}(\text{R}^6)_3$, $-\text{P}(\text{O})(\text{OR}^7)_2$, $-\text{P}(\text{O})(\text{OR}^7)(\text{R}^7)$, $-\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{R}^7$, $-\text{NR}^8\text{S}(\text{O})_2\text{R}^7$, $-\text{NR}^8\text{C}(\text{O})\text{N}(\text{R}^8)_2$,
 $-\text{NR}^8\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$,
 $-\text{S}(\text{O})_2\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{SR}^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl,

25 wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl of R^{10} (when present) is optionally independently
unsubstituted or further substituted with one or more independently selected groups
selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower
heteroalkyl, halo, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, and cycloalkyl.

- 13 -

In one embodiment, the compounds of the invention have the structural Formula (IA):




(IA)

wherein L₁, L₂, L₃, R¹, R², R³, R⁴, R⁵, ring A, ring B, m, and n are as defined in

5 Formula (I).

In one embodiment, the compounds of the invention have the structural Formula (IA-1):




(IA-1)

10 wherein L₁, L₂, L₃, R¹, R², R³, R⁴, R⁵, ring A, ring B, m, and n are each as defined in Formula (I).

- 14 -

In one embodiment, the compounds of the invention have the structural Formula (IA-2):



(IA-2)

5 wherein L₁, L₂, L₃, R¹, R², R³, R⁴, R⁵, ring A, ring B, m, and n are each as defined in Formula (I).

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R¹ is selected from the group consisting of H, lower alkyl, and cyclopropyl.

10 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R¹ is selected from the group consisting of H and methyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R¹ is H.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R¹ is methyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R² is H.

15 In one embodiment, in each of Formulas (I), (IA), (IA-1), and (IA-2):

R¹ is selected from the group consisting of H, lower alkyl, and cyclopropyl; and

R² is H.

20 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), -L₂- is absent or a -CH₂- group.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), -L₂- is absent.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), -L₂- is a -CH₂- group.

- 15 -

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R^3 is selected from the group consisting H, alkyl, haloalkyl, heteroalkyl, cycloalkyl, and cycloalkylalkyl-

5 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R^3 is lower alkyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), R^3 is methyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), $-L_2-$ is absent and R^3 is methyl.

10 In one embodiment, in each of Formulas (I), (IA), (IA-1), and (IA-2):

R^1 is selected from the group consisting of H, lower alkyl, and cyclopropyl;


R^2 is H;

$-L_2-$ is absent; and

R^3 is methyl.

15

In one embodiment, the compounds of the invention have the structural Formula (II):




(II)

wherein $-L_1-$, $-L_3-$, ring A, ring B, R^4 , R^5 , m and n are each as defined in Formula (I).

20

In one embodiment, the compounds of the invention have the structural Formula (IIA):


- 16 -



(IIA)

wherein $-L_1-$, $-L_3-$, ring A, ring B, R^4 , R^5 , m and n are each as defined in Formula (I).

5 In one embodiment, the compounds of the invention have the structural Formula (IIA-1):




(IIA-1)

wherein $-L_1-$, $-L_3-$, ring A, ring B, R^4 , R^5 , m and n are each as defined in Formula (I).

10 In one embodiment, the compounds of the invention have the structural Formula (IIA-2):

- 17 -



(IIA-2)

wherein $-L_1-$, $-L_3-$, ring A, ring B, R^4 , R^5 , m and n are each as defined in Formula (I).

- 5 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- $-L_1-$ is absent or a divalent $-alkyl-$ group.
- In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- 10 $-L_1-$ is absent.
- In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- $-L_1-$ is a divalent $-alkyl-$ group.
- In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- 15 $-L_1-$ is a divalent $-CH_2-$ group.
- In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- $-L_1-$ is a divalent $-CH_2CH_2-$ group.
- 20 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

- 18 -

m is 0 or more and ring A is selected from the group consisting of phenyl and monocyclic heteroaryl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

5 m is 0 or more and ring A is selected from the group consisting of phenyl, pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, thiazolyl, oxazolyl, pyrazolyl, imidazolyl, isoxazolyl, and isothiazolyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

10 m is 0 or more and ring A is selected from the group consisting of phenyl, pyridyl, and thienyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is selected from the group consisting of phenyl and thienyl;

15 wherein, when ring A is phenyl, m is 0 to 5, and
when ring A is thienyl, m is 0 to 3.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is selected from the group consisting of phenyl and thienyl;

20 wherein, when ring A is phenyl, m is 0 to 3, and
when ring A is thienyl, m is 0 to 2.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is selected from the group consisting of phenyl and thienyl;

25 wherein, when ring A is phenyl, m is 0 to 2, and
when ring A is thienyl, m is 0 to 1.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is selected from the group consisting of phenyl and thienyl;

30 wherein, when ring A is phenyl, m is 0 to 1, and
when ring A is thienyl, m is 0.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

- 19 -

ring A is selected from the group consisting of phenyl and thienyl;
wherein, when ring A is phenyl, m is 2 to 3, and
when ring A is thienyl, m is 1 to 2.

5 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is selected from the group consisting of phenyl and thienyl;
wherein, when ring A is phenyl, m is 0 to 3, and
when ring A is thienyl, m is 0 to 2.

10

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is phenyl and m is 0 to 3.

15 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is phenyl and m is 2 to 3.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

20 ring A is thienyl and m is 0 to 2.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

ring A is thienyl and m is 1 to 2.

25 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R^4 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -NO₂, -N(R⁸)₂, -NR⁸C(O)R⁷, -C(O)N(R⁸)₂, -OR⁷, alkyl, haloalkyl, heteroalkyl, and alkynyl.

30 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R^4 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -NO₂, -N(R⁸)₂, -NR⁸C(O)R⁷, -C(O)N(R⁸)₂, -OR⁷, alkyl, haloalkyl,

heteroalkyl, and alkynyl, wherein each R⁷ and each R⁸ (when present) is independently selected from H and lower alkyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R⁴ (when present) is independently selected from the group consisting of halo.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R⁴ (when present) is independently selected from the group consisting of fluoro, chloro, and bromo.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):


each R⁴ (when present) is independently selected from the group consisting of fluoro and bromo.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):


each R⁴ (when present) is fluoro.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

-L₁- is absent or -CH₂-;



- 21 -



- 5 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
 -L₃- is absent or a -alkyl- group.
 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- 10 -L₃- is absent.
 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
 -L₃- is a -alkyl- group.
- 15 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
 -L₃- is absent, or a divalent -CH₂- group, or a divalent -CH₂-CH₂- group.
 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):
- 20 -L₃- is a -CH₂- group.
 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

- 22 -

-L₁- is a divalent -CH₂CH₂- group.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

5 n is 0 or more and ring B is selected from the group consisting of phenyl, pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, thiazolyl, oxazolyl, benzofuranyl, benzimidazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, naphthyl, benzothienyl, benzothiazolyl, indazolyl, indolyl, benzocyclobutanyl, and difluorodioxolanyl.

10 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

n is 0 or more and ring B is selected from the group consisting of phenyl, pyridyl, thienyl, naphthyl, quinolinyl, isoquinolinyl, benzothienyl, benzocyclobutanyl, and difluorodioxolanyl.

15 In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -C(O)R⁷, -S(O)R⁷, -S(O)₂R⁷, -OR⁷, -SR⁷, alkyl, haloalkyl, heteroalkyl, -O-heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and monocyclic heteroaryl,

20 wherein each said alkyl, said alkenyl, said alkynyl, said cycloalkyl, said heterocycloalkyl, said aryl, and said monocyclic heteroaryl of R⁵ (when present) is optionally and independently further substituted with one or more groups independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R⁸)₂, and -OR⁷.

25

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

30 each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -C(O)R⁷, -S(O)R⁷, -S(O)₂R⁷, -OR⁷, -SR⁷, alkyl, haloalkyl, heteroalkyl, -O-heteroalkyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl,

- 23 -

oxetanyl, pyrrolyl, oxadiazolyl, pyrrolidinyl, furanyl, tetrahydrofuranyl, piperidinyl, and tetrahydropyranyl;

wherein each R^7 and each R^8 (when present) is independently selected from the group consisting of H and lower alkyl,

and

wherein each said alkyl, -alkoxy, haloalkyl, heteroalkyl, -O-heteroalkyl, alkenyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, furanyl, tetrahydrofuranyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl, oxetanyl, and pyrrolyl of R^5 (when present) is optionally and independently further substituted with one or more groups independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R^8)₂, and -OH.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R^5 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -N(R^8)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -C(O)R⁷, -S(O)R⁷, -S(O)₂R⁷, -OR⁷, -SR⁷, alkyl, haloalkyl, heteroalkyl, -O-heteroalkyl, alkenyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, furanyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl, oxetanyl, and pyrrolyl,

wherein each said alkyl, -alkoxy, haloalkyl, heteroalkyl, -O-heteroalkyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, furanyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl, oxetanyl, and pyrrolyl of R^5 (when present) is optionally and independently further substituted with one or more groups independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R^8)₂, and -OH, and

wherein each R^7 and each R^8 (when present) is independently selected from the group consisting of H and lower alkyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

each R^5 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, lower alkyl, lower alkenyl, lower haloalkyl, -C(O)-cyclopropyl, oxetanyl, lower alkyl-

- 24 -

substituted oxetanyl, cyclopropyl, lower heteroalkyl substituted cyclopropyl, lower alkyl-CN, lower heteroalkyl, and phenyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

-L₁- is absent or a divalent -alkyl- group;

m is 0 or more and ring A is selected from the group consisting of phenyl, pyridyl, pyrazinyl, pyrazolyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, thiazolyl, and oxazolyl.

each R⁴ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -NO₂, -N(R⁸)₂, -NR⁸C(O)R⁷, -C(O)N(R⁸)₂, -OR⁷, alkyl, haloalkyl, heteroalkyl, and alkynyl;

-L₃- is absent or a divalent -CH₂- group;

n is 0 or more and ring B is selected from the group consisting of phenyl, indazolyl, pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, thiazolyl, oxazolyl, naphthyl, quinolinyl, isoquinolinyl, benzothienyl, benzocyclobutanyl, and difluorodioxolanyl;

and

each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -C(O)R⁷, -S(O)R⁷, -S(O)₂R⁷, -OR⁷, -SR⁷, alkyl, haloalkyl, heteroalkyl, -O-heteroalkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, and monocyclic heteroaryl,

wherein each said alkyl, said alkenyl, said alkenyl, said cycloalkyl, said heterocycloalkyl, said aryl, and said monocyclic heteroaryl of R⁵ (when present) is optionally and independently further substituted with one or more groups independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R⁸)₂, and -OH.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

-L₁- is absent or a divalent -CH₂- group;

ring A is selected from the group consisting of phenyl and thienyl;

wherein, when ring A is phenyl, m is 0 to 5, and

when ring A is thienyl, m is 0 to 3;

- 25 -

each R⁴ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -NO₂, -N(R⁸)₂, -NR⁸C(O)R⁷, -C(O)N(R⁸)₂, -OR⁷, alkyl, haloalkyl, heteroalkyl, and alkynyl;

-L₃- is absent or a divalent -alkyl- group;

5 ring B is selected from the group consisting of phenyl, indazolyl, pyridyl, thienyl, naphthyl, quinolinyl, isoquinolinyl, benzothienyl, benzocyclobutanyl, and difluorodioxolanyl;

n is 0 or more;

10 each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -C(O)R⁷, -S(O)R⁷, -S(O)₂R⁷, -OR⁷, -SR⁷, alkyl, haloalkyl, heteroalkyl, -O-heteroalkyl, alkenyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, furanyl, thienyl, pyridazinyl, oxazolyl, oxadiazolyl, isoxazolyl, oxetanyl, and pyrrolyl,



15 wherein each said alkyl, -alkoxy, haloalkyl, heteroalkyl, -O-heteroalkyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, furanyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl, oxetanyl, and pyrrolyl of R⁵ (when present) is optionally and independently further substituted with one or more groups independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R⁸)₂, and -OH;

and

20 each R⁷ and each R⁸ (when present) is independently selected from the group consisting of H and lower alkyl.

In one embodiment, in each of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2):

25 -L₁- is absent or a divalent -CH₂- group;



n is 0, 1, 2, or 3;

ring B is selected from the group consisting of phenyl, indazolyl, pyridyl, thienyl, naphthyl, quinolinyl, isoquinolinyl, benzothienyl, benzocyclobutanyl, and difluorodioxolanyl;

10 and

each R^5 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, lower alkyl, lower alkenyl, lower haloalkyl, -C(O)-cyclopropyl, oxetanyl, lower alkyl-substituted oxetanyl, cyclopropyl, lower heteroalkyl substituted cyclopropyl, lower alkyl-CN, lower heteroalkyl, and phenyl.

- 28 -

For example, in one non-limiting embodiment, in Formula (I^d), R¹ is D and each of R², R³, R⁴, R⁵, R⁹, -L₁-, -L₂-, -L₃-, ring A, ring B, ring C m, n, and p are as defined in Formula (I) or as in any one of (IA), (IA-1), (IA-2), (II), (II-A), (II-A1), or (II-A2), or the various embodiments described herein.

5 As another example, in another non-limiting embodiment, in Formula (I^d), R² is D and each of R¹, R³, R⁴, R⁵, R⁹, -L₁-, -L₂-, -L₃-, ring A, ring B, ring C m, n, and p are as defined in Formula (I) or as in any one of (IA), (IA-1), (IA-2), (II), (II-A), (II-A1), or (II-A2), or the various embodiments described herein.

10 As another example, in another non-limiting embodiment, in Formula (I^d), R³ is D and each of R¹, R², R⁴, R⁵, R⁹, -L₁-, -L₂-, -L₃-, ring A, ring B, ring C m, n, and p are as defined in Formula (I) or as in any one of (IA), (IA-1), (IA-2), (II), (II-A), (II-A1), or (II-A2), or the various embodiments described herein.


15 As another example, in another non-limiting embodiment, in Formula (I^d), R⁴ is D and each of R¹, R², R³, R⁵, R⁹, -L₁-, -L₂-, -L₃-, ring A, ring B, ring C m, n, and p are as defined in Formula (I) or as in any one of (IA), (IA-1), (IA-2), (II), (II-A), (II-A1), or (II-A2), or the various embodiments described herein.

20 As another example, in another non-limiting embodiment, in Formula (I^d), R⁵ is D and each of R¹, R², R³, R⁴, R⁹, -L₁-, -L₂-, -L₃-, ring A, ring B, ring C m, n, and p are as defined in Formula (I) or as in any one of (IA), (IA-1), (IA-2), (II), (II-A), (II-A1), or (II-A2), or the various embodiments described herein.

25 As another example, in another non-limiting embodiment, in Formula (I^d), R⁹ is D and each of R¹, R², R³, R⁴, R⁵, -L₁-, -L₂-, -L₃-, ring A, ring B, ring C m, n, and p are as defined in Formula (I) or as in any one of (IA), (IA-1), (IA-2), (II), (II-A), (II-A1), or (II-A2), or the various embodiments described herein.

- 29 -


By way of further illustration, in another non-limiting embodiment, deuterated compounds of the invention have the structural Formula (II^d):



- 5 wherein:
- the moiety $-CD_3$ represents a deuterated form of the moiety $-CH_3$; and
- each of the remaining variables is as defined in Formula (I), or as described in any of the
- embodiments described herein, e.g., those of formulas (IA), (IA-1), (IA-2), (II), (II-A), (II-A1),
- and (II-A2), and the various embodiments thereof, are also within the scope of the compounds of
- 10 Formula (II^d).

- 30 -

By way of further illustration, in another non-limiting embodiment, deuterated compounds of the invention have the structural Formula (III^d):



5 wherein:
 the moiety -D represents a deuterated form of hydrogen; and
 each of the remaining variables is as defined in Formula (I), or as described in any of the
 10 embodiments described herein, e.g., those of formulas (IA), (IA-1), (IA-2), (II), (II-A), (II-A1),
 and (II-A2), and the various embodiments thereof, are also within the scope of the compounds of
 15 Formula (III^d).

 In another embodiment, the present invention encompasses a stereoisomer or racemic
 mixture of a compound of the invention, or a tautomer thereof, or a pharmaceutically acceptable
 salt of said compound or said tautomer. It shall be appreciated that, while the present invention
 encompasses all stereoisomers and racemic mixtures of the compounds of the invention, the
 15 stereoconfiguration shown in the structural formulas and in the examples are also contemplated as
 being within the scope of the invention.

 In another embodiment, 1 to 3 carbon atoms of the compounds of the invention may be
 replaced with 1 to 3 silicon atoms so long as all valency requirements are satisfied.

 In another embodiment, the compounds of the invention are each of the compounds of the
 20 tables below and have a structure shown for the corresponding example in the preparative
 examples below.

 The present invention includes tautomers and stereoisomers of each of the compounds of
 the invention, and pharmaceutically acceptable salts and solvates of said compounds, said

- 31 -

stereoisomers, and/or said tautomers. Such tautomers and stereoisomers of each of the example compounds of the invention, and pharmaceutically and solvates of said compounds, said stereoisomers, and/or said tautomers, each represent additional embodiments of the invention.

In another embodiment, the invention provides a composition comprising at least one
5 compound of the invention, or a tautomer or stereoisomer thereof, or salt or solvate of said compound, said stereoisomer, or said tautomer, and a suitable carrier or diluent.

In another embodiment, the invention provides a pharmaceutical composition comprising at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said
10 tautomer, and a pharmaceutically acceptable carrier or diluent.

In another embodiment, the invention provides a pharmaceutical composition comprising at least one solvate of a compound of the invention, or a tautomer or isomer thereof, or pharmaceutically acceptable salt or solvate of said compound or said tautomer, and a pharmaceutically acceptable carrier or diluent.

In another embodiment, the invention provides a pharmaceutical composition comprising at least one pharmaceutically acceptable salt of a compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said
15 stereoisomer, or said tautomer, and a pharmaceutically acceptable carrier or diluent.

In another embodiment, the invention provides a pharmaceutical composition comprising at least one tautomer of a compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said
20 tautomer, and a pharmaceutically acceptable carrier or diluent.

In another embodiment, the invention provides a pharmaceutical composition comprising at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said
25 tautomer, together with at least one additional therapeutic agent, and a pharmaceutically acceptable carrier or diluent.

Non-limiting examples of additional therapeutic agents for use in combination with the compounds of the invention include drugs selected from the group consisting of: (a) drugs useful
30 for the treatment of Alzheimer's disease and/or drugs useful for treating one or more symptoms of Alzheimer's disease, (b) drugs useful for inhibiting the synthesis A β , and (c) drugs useful for treating neurodegenerative diseases.

- 32 -

Additional non-limiting examples of additional therapeutic agents for use in combination with the compounds of the invention include drugs useful for the treatment, prevention, delay of onset, amelioration of any pathology associated with A β and/or a symptom thereof. Non-limiting examples of pathologies associated with A β include: Alzheimer's disease, Down's syndrome, 5 Parkinson's disease, memory loss, memory loss associated with Alzheimer's disease, memory loss associated with Parkinson's disease, attention deficit symptoms, attention deficit symptoms associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, dementia, stroke, microgliosis and brain inflammation, pre-senile dementia, senile dementia, dementia associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, progressive 10 supranuclear palsy, cortical basal degeneration, neurodegeneration, olfactory impairment, olfactory impairment associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, β -amyloid angiopathy, cerebral amyloid angiopathy, hereditary cerebral hemorrhage, mild cognitive impairment ("MCI"), glaucoma, amyloidosis, type II diabetes, hemodialysis complications (from β_2 microglobulins and complications arising therefrom in hemodialysis 15 patients), scrapie, bovine spongiform encephalitis, traumatic brain injury ("TBI"), and Creutzfeldt-Jakob disease, comprising administering to said patient at least one compound of the invention, or a tautomer or isomer thereof, or pharmaceutically acceptable salt or solvate of said compound or said tautomer, in an amount effective to inhibit said pathology or pathologies.

In embodiments of the invention comprising at least one additional therapeutic agent, 20 additional non-limiting examples of additional therapeutic agents for use in combination with compounds of the invention include: muscarinic antagonists (e.g., m₁ agonists (such as acetylcholine, oxotremorine, carbachol, or McNa343), or m₂ antagonists (such as atropine, dicycloverine, tolterodine, oxybutynin, ipratropium, methoctramine, triptamine, or gallamine)); cholinesterase inhibitors (e.g., acetyl- and/or butyrylcholinesterase inhibitors such as donepezil 25 (Aricept®), galantamine (Razadyne®), and rivastigimine (Exelon®); N-methyl-D-aspartate receptor antagonists (e.g., Namenda® (memantine HCl, available from Forrest Pharmaceuticals, Inc.); combinations of cholinesterase inhibitors and N-methyl-D-aspartate receptor antagonists; gamma secretase modulators; gamma secretase inhibitors; non-steroidal anti-inflammatory agents; anti-inflammatory agents that can reduce neuroinflammation; anti-amyloid antibodies 30 (such as bapineuzemab, Wyeth/Elan); vitamin E; nicotinic acetylcholine receptor agonists; CB1 receptor inverse agonists or CB1 receptor antagonists; antibiotics; growth hormone secretagogues; histamine H3 antagonists; AMPA agonists; PDE4 inhibitors; GABA_A inverse agonists; inhibitors of amyloid aggregation; glycogen synthase kinase beta inhibitors; promoters

- 33 -

of alpha secretase activity; PDE-10 inhibitors; Tau kinase inhibitors (e.g., GSK3beta inhibitors, cdk5 inhibitors, or ERK inhibitors); Tau aggregation inhibitors (e.g., Rember®); RAGE inhibitors (e.g., TTP 488 (PF-4494700)); anti-Abeta vaccine; APP ligands; agents that upregulate insulin, cholesterol lowering agents such as HMG-CoA reductase inhibitors (for example, statins such as Atorvastatin, Fluvastatin, Lovastatin, Mevastatin, Pitavastatin, Pravastatin, Rosuvastatin, Simvastatin) and/or cholesterol absorption inhibitors (such as Ezetimibe), or combinations of HMG-CoA reductase inhibitors and cholesterol absorption inhibitors (such as, for example, Vytorin®); fibrates (such as, for example, clofibrate, Clofibrade, Etofibrate, and Aluminium Clofibrate); combinations of fibrates and cholesterol lowering agents and/or cholesterol absorption inhibitors; nicotinic receptor agonists; niacin; combinations of niacin and cholesterol absorption inhibitors and/or cholesterol lowering agents (e.g., Simcor® (niacin/simvastatin, available from Abbott Laboratories, Inc.); LXR agonists; LRP mimics; H3 receptor antagonists; histone deacetylase inhibitors; hsp90 inhibitors; 5-HT4 agonists (e.g., PRX-03140 (Epix Pharmaceuticals)); 5-HT6 receptor antagonists; mGluR1 receptor modulators or antagonists; mGluR5 receptor modulators or antagonists; mGluR2/3 antagonists; Prostaglandin EP2 receptor antagonists; PAI-1 inhibitors; agents that can induce Abeta efflux such as gelsolin; Metal-protein attenuating compound (e.g., PBT2); and GPR3 modulators; and antihistamines such as Dimebolin (e.g., Dimebon®, Pfizer).

In another embodiment, the invention provides a pharmaceutical composition comprising an effective amount of one or more (e.g., one) compounds of the invention, and effective amount of one or more cholinesterase inhibitors (e.g., acetyl- and/or butyrylcholinesterase inhibitors), and a pharmaceutically acceptable carrier.

In another embodiment, the invention provides a pharmaceutical composition comprising an effective amount of one or more (e.g., one) compounds of the invention, and effective amount of one or more muscarinic antagonists (e.g., m₁ agonists or m₂ antagonists), and a pharmaceutically acceptable carrier.

In one embodiment, the invention provides combinations comprising an effective (i.e., therapeutically effective) amount of one or more compounds of the invention, in combination with an effective (i.e., therapeutically effective) amount of one or more compounds selected from the group consisting of cholinesterase inhibitors (such as, for example, (±)-2,3-dihydro-5,6-dimethoxy-2-[[1-(phenylmethyl)-4-piperidiny]methyl]-1 H -inden-1-one hydrochloride, i.e., donepezil hydrochloride, available as the Aricept® brand of donepezil hydrochloride), N-methyl-D-aspartate receptor inhibitors (such as, for example, Namenda® (memantine HCl)); anti-

- 34 -

amyloid antibodies (such as bapineuzumab, Wyeth/Elan), gamma secretase inhibitors, gamma secretase modulators, and beta secretase inhibitors other than the compounds of the invention.

In one embodiment, the invention provides combinations comprising an effective (i.e., therapeutically effective) amount of one or more compounds of the invention, in combination
5 with an effective (i.e., therapeutically effective) amount of one or more compounds selected from the group consisting of cholinesterase inhibitors (such as, for example, (\pm)-2,3-dihydro-5,6-dimethoxy-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1 H -inden-1-one hydrochloride, i.e., donepezil hydrochloride, available as the Aricept $\text{\textcircled{R}}$ brand of donepezil hydrochloride), N-methyl-D-aspartate receptor inhibitors (such as, for example, Namenda $\text{\textcircled{R}}$ (memantine HCl)).

10 In one embodiment, the invention provides combinations comprising an effective (i.e., therapeutically effective) amount of one or more compounds of the invention, in combination with an effective (i.e., therapeutically effective) amount of one or more gamma secretase inhibitors.

15 In one embodiment, the invention provides combinations comprising an effective (i.e., therapeutically effective) amount of one or more compounds of the invention, in combination with an effective (i.e., therapeutically effective) amount of one or more gamma secretase modulators.

20 In one embodiment, the invention provides combinations comprising an effective (i.e., therapeutically effective) amount of one or more compounds of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in combination with an effective (i.e., therapeutically effective) amount of one or more gamma secretase inhibitors and in further combination with one or more gamma secretase modulators.

25 In another embodiment, the invention provides a compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in pure form.

In another embodiment, the invention provides a compound of the invention or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in isolated form.

30 In another embodiment, the invention provides a compound of the invention or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in pure and isolated form.

- 35 -

Esters and prodrugs of the compounds of the invention, or tautomers or stereoisomers thereof, or pharmaceutically acceptable salts or solvates of said compounds, said stereoisomers, and/or said tautomers, are also contemplated as being included within the scope of the invention, and are described more fully below.

5 Deuterates of the compounds of the invention, or tautomers or stereoisomers of said deuterates, or pharmaceutically acceptable salts or solvates of said deuterates, said stereoisomers, and/or said tautomers, are also contemplated as being included within the scope of the invention, and are described more fully above.

10 In another embodiment, the invention provides a method of preparing a pharmaceutical composition comprising the step of admixing at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, and a pharmaceutically acceptable carrier or diluent.

15 In another embodiment, the invention provides a method of inhibiting β -secretase comprising exposing a population of cells expressing β -secretase to at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in an amount effective to inhibit β -secretase.

20 In another embodiment, the invention provides a method of inhibiting β -secretase in a patient in need thereof comprising administering at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in a therapeutically effective amount to inhibit β -secretase in said patient.

25 In another embodiment, the invention provides a method of inhibiting BACE-1 comprising exposing a population of cells expressing BACE-1 to at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound or said tautomer, in an amount effective to inhibit BACE-1 in said cells. In one such embodiment, said population of cells is *in vivo*. In another such embodiment, said population of cells is *ex vivo*. In another such embodiment, said population of cells is *in vitro*.

30 In another embodiment, the invention provides a method of inhibiting BACE-2 comprising exposing a population of cells expressing BACE-2 to at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound or said tautomer, in an amount effective to inhibit BACE-2 in said cells. In one

- 36 -

such embodiment, said population of cells is *in vivo*. In another such embodiment, said population of cells is *ex vivo*. In another such embodiment, said population of cells is *in vitro*.

In another embodiment, the invention provides a method of inhibiting BACE-1 in a patient in need thereof comprising administering to said patient at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of
5 said compound, said stereoisomer, or said tautomer, in a therapeutically effective amount to inhibit BACE-1 in said patient.

In another embodiment, the invention provides a method of inhibiting BACE-2 in a patient in need thereof comprising administering to said patient at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of
10 said compound, said stereoisomer, or said tautomer, in a therapeutically effective amount to inhibit BACE-2 in said patient.

In another embodiment, the invention provides a method of inhibiting the formation of A β from APP in a patient in need thereof, comprising administering to said patient at least one
15 compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in an amount effective to inhibit said A β formation.

In another embodiment, the invention provides a method of inhibiting the formation of A β plaque in a patient in need thereof, comprising administering to said patient at least one
20 compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in an amount effective to inhibit said A β plaque formation.

In another embodiment, the invention provides a method of inhibiting the formation of A β fibrils in a patient in need thereof, comprising administering to said patient at least one compound
25 of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in an amount effective to inhibit said A β fibril formation.

In another embodiment, the invention provides a method of inhibiting the formation of A β oligomers in a patient in need thereof, comprising administering to said patient at least one
30 compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in an amount effective to inhibit said A β fibril formation.

- 37 -

In another embodiment, the invention provides a method of inhibiting the formation of A β fibrils and A β oligomers in a patient in need thereof, comprising administering to said patient at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in an amount
5 effective to inhibit said A β fibril formation.

In another embodiment, the invention provides a method of inhibiting the formation of senile plaques and/or neurofibrillary tangles in a patient in need thereof, comprising administering to said patient at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said
10 stereoisomer, or said tautomer, in an amount effective to inhibit said A β fibril formation.

In another embodiment, the invention provides a method of treating, preventing, and/or delaying the onset of an amyloid β pathology ("A β pathology") and/or one or more symptoms of said pathology comprising administering at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said
15 stereoisomer, or said tautomer, to a patient in need thereof in an amount effective to treat said pathology.

In another embodiment, the invention provides a method of treating, preventing, and/or delaying the onset of one or more pathologies associated with A β and/or one or more symptoms of one or more pathologies associated with A β . Non-limiting examples of pathologies associated
20 with A β include: Alzheimer's disease, Down's syndrome, Parkinson's disease, memory loss, memory loss associated with Alzheimer's disease, memory loss associated with Parkinson's disease, attention deficit symptoms, attention deficit symptoms associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, dementia, stroke, microgliosis and brain inflammation, pre-senile dementia, senile dementia, dementia associated with Alzheimer's
25 disease, Parkinson's disease, and/or Down's syndrome, progressive supranuclear palsy, cortical basal degeneration, neurodegeneration, olfactory impairment, olfactory impairment associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, β -amyloid angiopathy, cerebral amyloid angiopathy, hereditary cerebral hemorrhage, mild cognitive impairment ("MCI"), glaucoma, amyloidosis, type II diabetes, diabetes-associated amyloidogenesis,
30 hemodialysis complications (from β_2 microglobulins and complications arising therefrom in hemodialysis patients), scrapie, bovine spongiform encephalitis, traumatic brain injury ("TBI") and Creutzfeld-Jakob disease, comprising administering to said patient at least one compound of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or

- 38 -

solvate of said compound, said stereoisomer, or said tautomer, in an amount effective to inhibit said pathology or pathologies.

In one embodiment, the invention provides a method of treating one or more neurodegenerative diseases, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of inhibiting the deposition of amyloid protein (e.g., amyloid beta protein) in, on or around neurological tissue (e.g., the brain), comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of inhibiting the deposition of amyloid protein (e.g., amyloid beta protein) in, on or around neurological tissue (e.g., the brain), comprising administering an effective (i.e., therapeutically effective) amount of a compound of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) in combination with an effective (i.e., therapeutically effective) amount of one or more additional therapeutic agents useful for treating Alzheimer's disease to a patient in need of treatment.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination

with an effective (i.e., therapeutically effective) amount of one or more cholinesterase inhibitors (such as, for example, (\pm)-2,3-dihydro-5,6-dimethoxy-2-[[1-(phenylmethyl)-4-piperidiny]methyl]-1 H -inden-1-one hydrochloride, i.e, donepezil hydrochloride, available as the Aricept® brand of donepezil hydrochloride), to a patient in need of treatment.

5 In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination
10 with an effective (i.e., therapeutically effective) amount of one or more compounds selected from the group consisting of A β antibody inhibitors, gamma secretase inhibitors, gamma secretase modulators, and beta secretase inhibitors other than a compound of the invention.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of
15 said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more compounds selected from the group consisting of A β antibody inhibitors, gamma secretase inhibitors, gamma secretase modulators, and beta secretase inhibitors.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more (e.g., one) compounds of the
20 invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more BACE inhibitors.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
25 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of Exelon (rivastigmine).

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
30 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of Cognex (tacrine).

- 40 -

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of a
5 Tau kinase inhibitor.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one
10 or more Tau kinase inhibitor (e.g., GSK3beta inhibitor, cdk5 inhibitor, ERK inhibitor).

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one
15 anti-Abeta vaccination (active immunization).

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one
20 or more APP ligands.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one
25 or more agents that upregulate insulin degrading enzyme and/or neprilysin.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one
30 or more cholesterol lowering agents (for example, statins such as Atorvastatin, Fluvastatin, Lovastatin, Mevastatin, Pitavastatin, Pravastatin, Rosuvastatin, Simvastatin, and cholesterol absorption inhibitor such as Ezetimibe).

- 41 -

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more fibrates (for example, clofibrate, Clofibrade, Etofibrate, Aluminium Clofibrate).

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more LXR agonists.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more LRP mimics.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more 5-HT₆ receptor antagonists.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more nicotinic receptor agonists.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more H₃ receptor antagonists.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said

- 42 -

compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more histone deacetylase inhibitors.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
5 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more hsp90 inhibitors.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
10 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more m1 muscarinic receptor agonists.

Another embodiment of this invention is directed to a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the
15 invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more 5-HT6 receptor antagonists, or mGluR1, or mGluR5 positive allosteric modulators or agonists.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
20 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more mGluR2/3 antagonists.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
25 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more anti-inflammatory agents that can reduce neuroinflammation.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a
30 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more Prostaglandin EP2 receptor antagonists.

- 43 -

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more PAI-1 inhibitors.

In one embodiment, the invention provides a method of treating Alzheimer's disease, comprising administering an effective amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective amount of one or more agents that can induce Abeta efflux such as gelsolin.

In one embodiment, the invention provides a method of treating Down's syndrome, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating Down's syndrome, comprising administering an effective (i.e., therapeutically effective) amount of one or more compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), in combination with an effective (i.e., therapeutically effective) amount of one or more cholinesterase inhibitors (such as, for example, (\pm)-2,3-dihydro-5,6-dimethoxy-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1 *H* -inden-1-one hydrochloride, i.e. donepezil hydrochloride, available as the Aricept ® brand of donepezil hydrochloride), to a patient in need of treatment.

In one embodiment, the invention provides a method of treating mild cognitive impairment, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating mild cognitive impairment, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or

- 44 -

more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating glaucoma, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
5 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating glaucoma, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
10 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating cerebral amyloid angiopathy, comprising administering an effective amount of one or more (e.g., one) compounds
15 of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating cerebral amyloid angiopathy, comprising administering an effective amount of one or more (e.g., one) compounds
20 of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating stroke, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
25 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating stroke, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
30 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating dementia, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a

tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating dementia, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
5 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating microgliosis, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
10 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating microgliosis, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a
15 tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating brain inflammation, comprising administering an effective amount of one or more (e.g., one) compounds of the
20 invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

In one embodiment, the invention provides a method of treating brain inflammation, comprising administering an effective amount of one or more (e.g., one) compounds of the
25 invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a method of treating traumatic brain injury, comprising administering an effective amount of one or more (e.g., one) compounds of the
invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of
said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

30 In one embodiment, the invention provides a method of treating olfactory function loss, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of
said compound, said stereoisomer, or said tautomer) to a patient in need of treatment.

- 46 -

In one embodiment, the invention provides a method of treating olfactory function loss, comprising administering an effective amount of one or more (e.g., one) compounds of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer), and an effective amount of one or more
5 additional therapeutic agents suitable for use in such patients, to a patient in need of treatment.

In one embodiment, the invention provides a kit comprising, in separate containers, in a single package, pharmaceutical compositions for use in combination, wherein one container comprises an effective amount of a compound of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or
10 said tautomer) in a pharmaceutically acceptable carrier, and another container (i.e., a second container) comprises an effective amount of another pharmaceutically active ingredient, the combined quantities of the compound of the invention and the other pharmaceutically active ingredient being effective to: (a) treat Alzheimer's disease, or (b) inhibit the deposition of amyloid protein in, on or around neurological tissue (e.g., the brain), or (c) treat
15 neurodegenerative diseases, or (d) inhibit the activity of BACE-1.

In one embodiment, the invention provides a kit comprising, in separate containers, in a single package, pharmaceutical compositions for use in combination, wherein one container comprises an effective amount of a compound of the invention (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or
20 said tautomer) in a pharmaceutically acceptable carrier, and another container (i.e., a second container) comprises an effective amount of another pharmaceutically active ingredient (as described below), the combined quantities of the compound of the invention and the other pharmaceutically active ingredient being effective to: (a) treat Alzheimer's disease, or (b) inhibit the deposition of amyloid protein (e.g., amyloid beta protein) in, on or around neurological tissue
25 (e.g., the brain), or (c) treat neurodegenerative diseases, or (d) inhibit the activity of BACE-1.

In various embodiments, the invention provides any one of the methods disclosed above and below wherein the compound(s) of the invention is a compound or compounds selected from the group consisting of the exemplary compounds of the invention described below.

In various embodiments, the invention provides any one of the pharmaceutical
30 compositions disclosed above and below wherein the compound(s) of the invention is a compound or compounds selected from the group consisting of the exemplary compounds of the invention described below.

- 47 -

Other embodiments of this invention are directed to any one of the embodiments above or below that are directed to compounds of the invention, or the use of compounds of the invention (e.g. the embodiments directed to methods of treatment, pharmaceutical compositions and kits).

In another embodiment, the invention provides for the use of a compound of the
5 invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, in the manufacture of a medicament for use in the treatment, the delay of onset, and/or the prevention of one or more A β pathologies and/or in the treatment, the delay of onset, and/or the prevention of one or more symptoms of one or more A β pathologies.

10 In another embodiment, the invention provides a kit comprising: (a) one or more compounds of the invention, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, preferably provided as a pharmaceutical composition and in a suitable container or containers and/or with suitable packaging; (b) optionally one or more additional active agents, which if present are
15 preferably provided as a pharmaceutical composition and in a suitable container or containers and/or with suitable packaging; and (c) instructions for use, for example written instructions on how to administer the compound or compositions.

20 In another embodiment, the invention provides a kit comprising a single container or multiple containers: (a) a pharmaceutically acceptable composition comprising one or more compounds of claim 1, or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer, (b) optionally pharmaceutically acceptable composition comprising one or more additional therapeutic agents; and (c) instructions for use their use. Said kit may optionally comprise labeling appropriate to the intended use or uses.

- 48 -

DEFINITIONS

The terms used herein have their ordinary meaning and the meaning of such terms is independent at each occurrence thereof. That notwithstanding and except where stated otherwise, the following definitions apply throughout the specification and claims. Chemical names, common names and chemical structures may be used interchangeably to describe that same structure. These definitions apply regardless of whether a term is used by itself or in combination with other terms, unless otherwise indicated. Hence the definition of "alkyl" applies to "alkyl" as well as the "alkyl" portion of "hydroxyalkyl", "haloalkyl", arylalkyl-, alkylaryl-, "alkoxy" etc.

It shall be understood that, in the various embodiments of the invention described herein, any variable not specifically defined in the context of the embodiment is as defined in Formula (I). All valences not explicitly filled are assumed to be filled by hydrogen.


As described herein, the "example compounds of the invention" (or "example compounds" or "examples") include, collectively and individually, each of the compounds set forth with example numbers in the preparative examples.


In each of the various embodiments of the compounds of the invention described herein, including those of Formulas (a), (I), (IA), (IA-1), and (IA-2), (II), (IIA), (IIA-1), and (IIA-2), and the various embodiments thereof, each variable is selected independently of the others unless otherwise noted.


As described herein, variables such as R^1 , R^2 , and R^3 may be unsubstituted or substituted with one or more R^{10} groups. It shall be understood that the upper limit of the number of substituents (referred to in the phrase "one or more substituents") is the number of available hydrogen atoms on the relevant moiety (R^1 , R^2 , or R^3) that are available for replacement by a substituent which will result in a chemically stable moiety. If an upper number of a range is given (e.g., in Formula (a), variables such as R^1 may be unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups), the maximum number of substitutable positions is the lessor of the upper number of the range (e.g., 5) or the maximum number of available substitutable hydrogen atoms on the substituted moiety.

As described herein, one or more of the variables of the general formulae representing the various embodiments of the compounds of the invention (e.g., variables $-L_1-$, $-L_2-$, $-L_3-$, and $-L_4-$) optionally independently is absent. It shall be understood that where such a variable is absent, the moieties which are shown connected by that variable are directly attached by bond. Thus, by way of non-limiting illustration only, a compound of Formula (I) wherein $-L_1-$, $-L_2-$, $-L_3-$ and $-L_4-$ each independently is absent, such compounds are understood to be depicted as:


- 49 -



The moiety , which may be optionally substituted as described herein, represents a ring referred to herein as "ring A."

The moiety , which may be optionally substituted as described herein, represents a ring referred to herein as "ring B."

5

The moiety , which may be optionally substituted as described herein, represents a ring referred to herein as "ring C."

In the various Formulas of the compounds of the invention, e.g., in Formula (I), m, n, p and q are each independently selected integers, wherein:

- 10
- m is 0 or more,
 - n is 0 or more,
 - p is 0 or more, and
 - q is 0 or more.


Where the upper limit on such variables is indicated by the phrase "or more", it shall be understood that the maximum value of that variable is the maximum number of available substitutable hydrogen atoms on the moiety to which that variable is attached. For example, the maximum value of the sum of m and q in the phrase "m is 0 or more" and "q is 0 or more" is the maximum number of available substitutable hydrogen atoms on ring A. The maximum value of n is the phrase "n is 0 or more" is the maximum number of available substitutable hydrogen atoms


15

- 50 -


on ring B. The maximum value of p in the phrase “p is 0 or more” is the maximum number of available substitutable hydrogen atoms on ring C. Except for salt forms, the maximum number of substitutable hydrogen atoms is understood to be the maximum number that will result in a neutral molecule.

5 By way of non-limiting illustration, when ring A is a phenyl group, the maximum value of





m is 5. When ring A is a  group, the maximum value of m is 4.

By way of further non-limiting illustration, and in one embodiment, in the various formulas of the compounds of the invention, e.g., in Formula (I) wherein ring A is a multicyclic



10 group, the minimum value of the sum of m and q is 0 and the maximum value of the sum of m and q is 17.

When ring B is a multicyclic  group, the minimum value of the sum of n is 0 and the maximum value of n is 17.

When ring C is a multicyclic  group, the minimum value of the sum of p is 0 and the maximum value of p is 17.

15 Thus, in one embodiment, in Formula (a):

m, n, p and q are each independently selected integers, wherein:

the minimum value of the sum of m and q is 0 and the maximum value of the sum of m and q is 17;

n is 0 to 17; and

20 p is 0 to 17.

In the compounds of the invention, e.g., in Formula (I), each of ring A, ring B, and ring C (when present) is independently selected from the group consisting of a monocyclic aryl, a monocyclic heteroaryl, a monocyclic cycloalkyl, a monocyclic cycloalkenyl, a monocyclic heterocycloalkyl, a monocyclic heterocycloalkenyl, and a multicyclic group, each of which

25 groups may be unsubstituted or optionally further substituted as shown.

As used herein, the term “monocyclic aryl” refers to phenyl.

- 51 -


As used herein, the term “monocyclic heteroaryl” refers to a 4- to 7-membered monocyclic heteroaryl group comprising from 1 to 4 ring heteroatoms, said ring heteroatoms being independently selected from the group consisting of N, O, and S, and oxides thereof. The point of attachment to the parent moiety is to any available ring carbon or ring heteroatom. Non-limiting examples of monocyclic heteroaryl moieties include pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, pyridone, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, thiadiazolyl (e.g., 1,2,4-thiadiazolyl), pyrazinyl, pyridazinyl, imidazolyl, and triazinyl (e.g., 1,2,4-triazinyl), and oxides thereof.

As used herein, the term “monocyclic cycloalkyl” refers to a 3- to 7-membered monocyclic cycloalkyl group. Non-limiting examples of monocyclic cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

As used herein, the term “monocyclic cycloalkenyl” refers to a non-aromatic 3- to 7-membered cycloalkyl group which contains one or more carbon-carbon double bonds. Non-limiting examples include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and cycloheptenyl.

As used herein, the term “monocyclic heterocycloalkyl” refers to a 4- to 7-membered monocyclic heterocycloalkyl group comprising from 1 to 4 ring heteroatoms, said ring heteroatoms being independently selected from the group consisting of N, N-oxide, O, S, S-oxide, S(O), and S(O)₂. The point of attachment to the parent moiety is to any available ring carbon or ring heteroatom. Non-limiting examples of monocyclic heterocycloalkyl groups include piperidyl, oxetanyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, *beta* lactam, *gamma* lactam, *delta* lactam, *beta* lactone, *gamma* lactone, *delta* lactone, and pyrrolidinone, and oxides thereof.

Non-limiting examples of lower alkyl-substituted oxetanyl include the moiety:



As used herein, the term “monocyclic heterocycloalkenyl” refers to a 4- to 7-membered monocyclic heterocycloalkenyl group comprising from 1 to 4 ring heteroatoms, said ring heteroatoms being independently selected from the group consisting of N, N-oxide, O, S, S-oxide, S(O), and S(O)₂. The point of attachment to the parent moiety is to any available ring carbon or ring heteroatom. Non-limiting examples of monocyclic heterocycloalkenyl groups include 1,2,3,4-

30

tetrahydropyridinyl, 1,2-dihydropyridinyl, 1,4-dihydropyridinyl, 1,2,3,6-tetrahydropyridinyl, 1,4,5,6-tetrahydropyrimidinyl, 2-pyrrolinyl, 3-pyrrolinyl, 2-imidazoliny, 2-pyrazolinyl, dihydroimidazolyl, dihydrooxazolyl, dihydrooxadiazolyl, dihydrothiazolyl, 3,4-dihydro-2H-pyranyl, dihydrofuranyl, fluorodihydrofuranyl, dihydrothiophenyl, and dihydrothiopyranyl, and oxides thereof.

5


As used herein, the term "multicyclic group" refers to a fused ring system comprising two (bicyclic), three (tricyclic), or more fused rings, wherein each ring of the fused ring system is independently selected from the group consisting of phenyl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, and monocyclic heterocycloalkenyl. The point of attachment to the parent moiety is to any available ring carbon or (if present) ring heteroatom on any of the fused rings.

10

It shall be understood that each of the following multicyclic groups pictured may be unsubstituted or substituted, as described herein. Only the point of attachment to the parent moiety is shown by the wavy line.

15

The term multicyclic groups includes bicyclic aromatic groups. Non-limiting examples of


multicyclic groups which are bicyclic aromatic groups include: 


The term multicyclic groups includes bicyclic heteroaromatic groups comprising from 1 to 3 or more ring heteroatoms, each said ring heteroatom being independently selected from the group consisting of N, O, and S, S(O), S(O)₂, and oxides of N, O, and S, and oxides thereof.

20


Non-limiting examples of multicyclic groups which are bicyclic heteroaromatic groups comprising from 1 to 3 ring heteroatoms, each said ring heteroatom being independently selected from N, O, and S include the following, and oxides thereof:

25






10 The term multicyclic groups includes saturated bicyclic cycloalkyl groups. Non-limiting examples of multicyclic groups which are saturated bicyclic cycloalkyl groups include the following:








15 The term multicyclic group includes partially unsaturated bicyclic cycloalkyl groups. Non-limiting examples of multicyclic groups which comprise partially unsaturated bicyclic cycloalkyl groups include the following:








20 The term multicyclic groups includes partially or fully saturated bicyclic groups comprising from 1 to 3 ring heteroatoms, each said ring heteroatom is independently selected from the group consisting of N, O, and S, S(O), S(O)₂, and oxides of N and S. Such rings may

also optionally contain one or more oxo groups, as defined herein. Non-limiting examples of multicyclic groups which are partially or fully saturated bicyclic groups comprising from 1 to 3 ring heteroatoms, each said ring heteroatom being independently selected from N, O, and S include the following, and oxides thereof:








5





10





The term multicyclic groups includes aromatic tricyclic groups, cycloalkyl tricyclic groups, as well as heteroaromatic and partially and fully saturated tricyclic groups. For tricyclic groups comprising ring heteroatoms, said tricyclic groups comprise one or more (e.g., from 1 to 5) ring heteroatoms, wherein each said ring heteroatom is independently selected from N, O, and S, S(O), S(O)₂, and oxides of N, O, and S: Non-limiting examples of tricyclic multicyclic groups include the following, and, where possible, oxides thereof:





"Patient" includes both human and non-human animals. Non-human animals include those research animals and companion animals such as mice, primates, monkeys, great apes, canine (*e.g.*, dogs), and feline (*e.g.*, house cats).

"Pharmaceutical composition" (or "pharmaceutically acceptable composition") means a composition suitable for administration to a patient. Such compositions may contain the neat compound (or compounds) of the invention or mixtures thereof, or salts, solvates, prodrugs,

- 58 -

isomers, or tautomers thereof, or they may contain one or more pharmaceutically acceptable carriers or diluents. The term "pharmaceutical composition" is also intended to encompass both the bulk composition and individual dosage units comprised of more than one (e.g., two) pharmaceutically active agents such as, for example, a compound of the present invention and an additional agent selected from the lists of the additional agents described herein, along with any pharmaceutically inactive excipients. The bulk composition and each individual dosage unit can contain fixed amounts of the afore-said "more than one pharmaceutically active agents". The bulk composition is material that has not yet been formed into individual dosage units. An illustrative dosage unit is an oral dosage unit such as tablets, pills and the like. Similarly, the herein-described method of treating a patient by administering a pharmaceutical composition of the present invention is also intended to encompass the administration of the afore-said bulk composition and individual dosage units.

"Halogen" means fluorine, chlorine, bromine, or iodine. Preferred are fluorine, chlorine and bromine.

"Alkyl" means an aliphatic hydrocarbon group which may be straight or branched and comprising about 1 to about 20 carbon atoms in the chain. Preferred alkyl groups contain about 1 to about 12 carbon atoms in the chain. More preferred alkyl groups contain about 1 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. "Lower alkyl" means a group having about 1 to about 6 carbon atoms in the chain which may be straight or branched. "Alkyl" may be unsubstituted or optionally substituted by one or more substituents which may be the same or different, each substituent being as described herein or independently selected from the group consisting of halo, alkyl, haloalkyl, spirocycloalkyl, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -O-C(O)-alkyl, -O-C(O)-aryl, -O-C(O)-cycloalkyl, carboxy and -C(O)O-alkyl. Non-limiting examples of suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl and t-butyl.


"Haloalkyl" means an alkyl as defined above wherein one or more hydrogen atoms on the alkyl is replaced by a halo group defined above.

"Heteroalkyl" means an alkyl moiety as defined above, having one or more carbon atoms, for example one, two or three carbon atoms, replaced with one or more heteroatoms, which may be the same or different, where the point of attachment to the remainder of the molecule is through a carbon atom of the heteroalkyl radical. Suitable such heteroatoms include O, S, S(O), S(O)₂, and -NH-, -N(alkyl)-. Non-limiting examples include ethers, thioethers, amines,

- 59 -

hydroxymethyl, 3-hydroxypropyl, 1,2-dihydroxyethyl, 2-methoxyethyl, 2-aminoethyl, 2-dimethylaminoethyl, and the like.

"Alkenyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkenyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkenyl chain. "Lower alkenyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. "Alkenyl" may be unsubstituted or optionally substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halo, alkyl, aryl, cycloalkyl, cyano, alkoxy and -S(alkyl). Non-limiting examples of suitable alkenyl groups include ethenyl, propenyl, n-butenyl, 3-methylbut-2-enyl, n-pentenyl, octenyl and decenyl.

"Alkylene" means a difunctional group obtained by removal of a hydrogen atom from an alkyl group that is defined above. Non-limiting examples of alkylene include methylene, ethylene and propylene. More generally, the suffix "ene" on alkyl, aryl, heterocycloalkyl, etc. indicates a divalent moiety, e.g., -CH₂CH₂- is ethylene, and  is para-phenylene.

"Alkynyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkynyl chain. "Lower alkynyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butyne-1-yl and 3-methylbutynyl. "Alkynyl" may be unsubstituted or optionally substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of alkyl, aryl and cycloalkyl.

"Alkenylene" means a difunctional group obtained by removal of a hydrogen from an alkenyl group that is defined above. Non-limiting examples of alkenylene include -CH=CH-, -C(CH₃)=CH-, and -CH=CHCH₂-.

"Aryl" means an aromatic monocyclic or polycyclic ring system comprising about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be

- 60 -


optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl.

"Heteroaryl" means an aromatic monocyclic or multicyclic ring system comprising about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the ring atoms is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. Preferred heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. A nitrogen atom of a heteroaryl can be optionally oxidized to the corresponding N-oxide. "Heteroaryl" may also include a heteroaryl as defined above fused to an aryl as defined above. Non-limiting examples of suitable heteroaryls include pyridyl, pyrazinyl, furanyl, thienyl (alternatively referred to herein as thiophenyl), pyrimidinyl, pyridone (including N-substituted pyridones), isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, 1,2,4-thiadiazolyl, pyrazinyl, pyridazinyl, quinoxalyl, phthalazinyl, oxindolyl, imidazo[1,2-a]pyridinyl, imidazo[2,1-b]thiazolyl, benzofurazanyl, indolyl, azaindolyl, benzimidazolyl, benzothienyl, quinolinyl, imidazolyl, thienopyridyl, quinazolinyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, isoquinolinyl, benzoazaindolyl, 1,2,4-triazinyl, benzothiazolyl and the like. The term "heteroaryl" also refers to partially saturated heteroaryl moieties such as, for example, tetrahydroisoquinolyl, tetrahydroquinolyl and the like.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms. The cycloalkyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable monocyclic cycloalkyls include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable multicyclic cycloalkyls include 1-decalinyl, norbornyl, adamantyl and the like. Further non-limiting examples of cycloalkyl include the following:

30


- 61 -



"Cycloalkenyl" means a non-aromatic mono or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms which contains at least one carbon-carbon double bond. Preferred cycloalkenyl rings contain about 5 to about 7 ring atoms. The cycloalkenyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkenyls include cyclopentenyl, cyclohexenyl, cyclohepta-1,3-dienyl, and the like. Non-limiting example of a suitable multicyclic cycloalkenyl is norbornenyl.

- 62 -


“Heterocycloalkyl” (or “heterocyclyl”) means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. Any –NH in a heterocyclyl ring may exist protected such as, for example, as an -N(Boc), -N(CBz), -N(Tos) group and the like; such protections are also considered part of this invention. The heterocyclyl can be optionally substituted by one or more “ring system substituents” which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Thus, the term “oxide,” when it appears in a definition of a variable in a general structure described herein, refers to the corresponding N-oxide, S-oxide, or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, lactam, lactone, and the like. “Heterocyclyl” also includes rings wherein =O replaces two available hydrogens on the same carbon atom (i.e., heterocyclyl includes rings having a carbonyl group in the ring). Such =O groups may be referred to herein as “oxo.” An example of such a moiety is pyrrolidinone (or pyrrolidone):




“Heterocycloalkenyl” (or “heterocyclenyl”) means a non-aromatic monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur atom, alone or in combination, and which contains at least one carbon-carbon double bond or carbon-nitrogen double bond. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocyclenyl rings contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclenyl root name means that at

- 63 -

least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. The heterocyclenyl can be optionally substituted by one or more ring system substituents, wherein "ring system substituent" is as defined above. The nitrogen or sulfur atom of the heterocyclenyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable heterocyclenyl groups include 1,2,3,4- tetrahydropyridinyl, 1,2-dihydropyridinyl, 1,4-dihydropyridinyl, 1,2,3,6-tetrahydropyridinyl, 1,4,5,6-tetrahydropyrimidinyl, 2-pyrrolinyl, 3-pyrrolinyl, 2-imidazoliny, 2-pyrazoliny, dihydroimidazolyl, dihydrooxazolyl, dihydrooxadiazolyl, dihydrothiazolyl, 3,4-dihydro-2H-pyranyl, dihydrofuranyl, fluorodihydrofuranyl, 7-oxabicyclo[2.2.1]heptenyl, dihydrothiophenyl, dihydrothiopyranyl, and the like. "Heterocyclenyl" also includes rings wherein =O replaces two available hydrogens on the same carbon atom (i.e., heterocyclenyl includes rings having a carbonyl group in the ring). Example of such moiety is pyrrolidenone (or pyrrolone):



It should be noted that in hetero-atom containing ring systems of this invention, there are no hydroxyl groups on carbon atoms adjacent to a N, O or S, as well as there are no N or S groups on carbon adjacent to another heteroatom. Thus, for example, in the ring:




there is no -OH attached directly to carbons marked 2 and 5.

"Arylcycloalkyl" (or "arylfused cycloalkyl") means a group derived from a fused aryl and cycloalkyl as defined herein. Preferred arylcycloalkyls are those wherein aryl is phenyl (which may be referred to as "benzofused") and cycloalkyl consists of about 5 to about 6 ring atoms. The arylcycloalkyl can be optionally substituted as described herein. Non-limiting examples of suitable arylcycloalkyls include indanyl (a benzofused cycloalkyl) and 1,2,3,4-tetrahydronaphthyl and the like. The bond to the parent moiety is through a non-aromatic carbon atom.

- 64 -

“Arylheterocycloalkyl” (or “arylfused heterocycloalkyl”) means a group derived from a fused aryl and heterocycloalkyl as defined herein. Preferred arylheterocycloalkyls are those wherein aryl is phenyl (which may be referred to as “benzofused”) and heterocycloalkyl consists of about 5 to about 6 ring atoms. The arylheterocycloalkyl can be optionally substituted, and/or
 5 contain the oxide or oxo, as described herein. Non-limiting examples of suitable arylfused heterocycloalkyls include:



The bond to the parent moiety is through a non-aromatic carbon atom.

It is also understood that the terms “arylfused aryl”, “arylfused cycloalkyl”, “arylfused
 10 cycloalkenyl”, “arylfused heterocycloalkyl”, arylfused heterocycloalkenyl”, “arylfused heteroaryl”, “cycloalkylfused aryl”, “cycloalkylfused cycloalkyl”, “cycloalkylfused cycloalkenyl”, “cycloalkylfused heterocycloalkyl”, “cycloalkylfused heterocycloalkenyl”, “cycloalkylfused heteroaryl”, “cycloalkenylfused aryl”, “cycloalkenylfused cycloalkyl”, “cycloalkenylfused cycloalkenyl”, “cycloalkenylfused heterocycloalkyl”, “cycloalkenylfused heterocycloalkenyl”, “cycloalkenylfused heteroaryl”, “heterocycloalkylfused aryl”, “heterocycloalkylfused cycloalkyl”, “heterocycloalkylfused cycloalkenyl”, “heterocycloalkylfused heterocycloalkyl”, “heterocycloalkylfused heterocycloalkenyl”, “heterocycloalkylfused heteroaryl”, “heterocycloalkenylfused aryl”, “heterocycloalkenylfused cycloalkyl”, “heterocycloalkenylfused cycloalkenyl”, “heterocycloalkenylfused heterocycloalkyl”, “heterocycloalkenylfused heterocycloalkenyl”, “heterocycloalkenylfused heteroaryl”, “heteroaryl fused aryl”, “heteroaryl fused cycloalkyl”, “heteroaryl fused cycloalkenyl”, “heteroaryl fused heterocycloalkyl”, “heteroaryl fused heterocycloalkenyl”, and “heteroaryl fused heteroaryl” are similarly represented by the combination of the groups aryl, cycloalkyl,
 20 cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, and heteroaryl, as previously described. Any such groups may be unsubstituted or substituted with one or more ring system substituents at any available position as described herein.

“Aralkyl” or “arylalkyl” means an aryl-alkyl- group in which the aryl and alkyl are as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, 2-phenethyl and naphthalenylmethyl. The bond to the

parent moiety is through the alkyl. The term (and similar terms) may be written as "arylalkyl-" to indicate the point of attachment to the parent moiety.

Similarly, "heteroarylalkyl", "cycloalkylalkyl", "cycloalkenylalkyl", "heterocycloalkylalkyl", "heterocycloalkenylalkyl", etc., mean a heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, etc. as described herein bound to a parent moiety through an alkyl group. Preferred groups contain a lower alkyl group. Such alkyl groups may be straight or branched, unsubstituted and/or substituted as described herein.

Similarly, "arylfused arylalkyl-", arylfused cycloalkylalkyl-, etc., means an arylfused aryl group, arylfused cycloalkyl group, etc. linked to a parent moiety through an alkyl group. Preferred groups contain a lower alkyl group. Such alkyl groups may be straight or branched, unsubstituted and/or substituted as described herein.

"Alkylaryl" means an alkyl-aryl- group in which the alkyl and aryl are as previously described. Preferred alkylaryls comprise a lower alkyl group. Non-limiting example of a suitable alkylaryl group is tolyl. The bond to the parent moiety is through the aryl.

"Cycloalkylether" means a non-aromatic ring of 3 to 7 members comprising an oxygen atom and 2 to 7 carbon atoms. Ring carbon atoms can be substituted, provided that substituents adjacent to the ring oxygen do not include halo or substituents joined to the ring through an oxygen, nitrogen or sulfur atom.

"Cycloalkylalkyl" means a cycloalkyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable cycloalkylalkyls include cyclohexylmethyl, adamantylmethyl, adamantylpropyl, and the like.

"Cycloalkenylalkyl" means a cycloalkenyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable cycloalkenylalkyls include cyclopentenylmethyl, cyclohexenylmethyl and the like.

"Heteroarylalkyl" means a heteroaryl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable heteroaryls include 2-pyridinylmethyl, quinolinylmethyl and the like.

"Heterocyclalkyl" (or "heterocycloalkyl") means a heterocyclalkyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of suitable heterocyclalkyls include piperidinylmethyl, piperazinylmethyl and the like.

"Heterocyclenylalkyl" means a heterocyclenyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core.

"Alkynylalkyl" means an alkynyl-alkyl- group in which the alkynyl and alkyl are as previously described. Preferred alkynylalkyls contain a lower alkynyl and a lower alkyl group. The bond to the parent moiety is through the alkyl. Non-limiting examples of suitable alkynylalkyl groups include propargylmethyl.

5 "Heteroaralkyl" means a heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain a lower alkyl group. Non-limiting examples of suitable aralkyl groups include pyridylmethyl, and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkyl.

10 "Hydroxyalkyl" means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

"Cyanoalkyl" means a NC-alkyl- group in which alkyl is as previously defined. Preferred cyanoalkyls contain lower alkyl. Non-limiting examples of suitable cyanoalkyl groups include
15 cyanomethyl and 2-cyanoethyl.

"Acyl" means an H-C(O)-, alkyl-C(O)- or cycloalkyl-C(O)-, group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl and propanoyl.

20 "Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1-naphthoyl.

"Heteroaroyl" means a heteroaryl-C(O)- group in which the heteroaryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting
25 examples of suitable groups include pyridoyl.

"Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy, *n*-propoxy, isopropoxy and *n*-butoxy. The bond to the parent moiety is through the ether oxygen.

"Alkyoxyalkyl" means a group derived from an alkoxy and alkyl as defined herein. The
30 bond to the parent moiety is through the alkyl.

"Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond to the parent moiety is through the ether oxygen.

"Aralkyloxy" (or "arylalkyloxy") means an aralkyl-O- group (an arylalkyl-O- group) in which the aralkyl group is as previously described. Non-limiting examples of suitable aralkyloxy groups include benzyloxy and 1- or 2-naphthalenemethoxy. The bond to the parent moiety is through the ether oxygen.

5 "Arylalkenyl" means a group derived from an aryl and alkenyl as defined herein. Preferred arylalkenyls are those wherein aryl is phenyl and the alkenyl consists of about 3 to about 6 atoms. The arylalkenyl can be optionally substituted by one or more substituents. The bond to the parent moiety is through a non-aromatic carbon atom.

10 "Arylalkynyl" means a group derived from a aryl and alkynyl as defined herein. Preferred arylalkynyls are those wherein aryl is phenyl and the alkynyl consists of about 3 to about 6 atoms. The arylalkynyl can be optionally substituted by one or more substituents. The bond to the parent moiety is through a non-aromatic carbon atom.

15 "Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkylthio groups include methylthio and ethylthio. The bond to the parent moiety is through the sulfur.

"Arylthio" means an aryl-S- group in which the aryl group is as previously described. Non-limiting examples of suitable arylthio groups include phenylthio and naphthylthio. The bond to the parent moiety is through the sulfur.

20 "Aralkylthio" means an aralkyl-S- group in which the aralkyl group is as previously described. Non-limiting example of a suitable aralkylthio group is benzylthio. The bond to the parent moiety is through the sulfur.

"Alkoxy carbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxy carbonyl groups include methoxy carbonyl and ethoxy carbonyl. The bond to the parent moiety is through the carbonyl.

25 "Aryloxy carbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxy carbonyl groups include phenoxy carbonyl and naphthoxy carbonyl. The bond to the parent moiety is through the carbonyl.

30 "Aralkoxy carbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxy carbonyl group is benzyloxy carbonyl. The bond to the parent moiety is through the carbonyl.

"Alkylsulfonyl" means an alkyl-S(O₂)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

- 68 -

"Arylsulfonyl" means an aryl-S(O₂)- group. The bond to the parent moiety is through the sulfonyl.

5 "Spirocycloalkyl" means a cycloalkyl group attached to a parent moiety by replacement of two available hydrogen atoms at a single carbon atom. Non-limiting examples of spirocycloalkyl wherein the parent moiety is a cycloalkyl include spiro [2.5] octane, spiro [2.4] heptane, etc. The moiety may optionally be substituted as described herein. Non-limiting spirocycloalkyl groups include spirocyclopropyl, spirocyclobutyl, spirocycloheptyl, and spirocyclohexyl.

10 The term "substituted" means that one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. By "stable compound" or "stable structure" is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

15 The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties.

20 Substitution on a cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, heteroarylalkyl, arylfused cycloalkylalkyl- moiety or the like includes substitution on any ring portion and/or on the alkyl portion of the group.

When a variable appears more than once in a group, e.g., R⁸ in -N(R⁸)₂, or a variable appears more than once in a structure presented herein, the variables can be the same or different.


25 With reference to the number of moieties (e.g., substituents, groups or rings) in a compound, unless otherwise defined, the phrases "one or more" and "at least one" mean that there can be as many moieties as chemically permitted, and the determination of the maximum number of such moieties is well within the knowledge of those skilled in the art. With respect to the compositions and methods comprising the use of "at least one compound of the invention, e.g., of Formula (II)," one to three compounds of the invention, e.g., of Formula (II) can be administered at the same time, preferably one.

30 Compounds of the invention may contain one or more rings having one or more ring system substituents. "Ring system substituent" means a substituent attached to an aromatic or non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being as described herein or

- 69 -


independently selected from the group consisting of alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, aryl, heteroaryl, aralkyl, alkylaryl, heteroaralkyl, heteroarylalkenyl, heteroarylalkynyl, alkylheteroaryl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halo, nitro, cyano, carboxy, alkoxy-carbonyl, aryloxy-carbonyl, aralkoxy-carbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, heterocyclyl, -O-C(O)-alkyl, -O-C(O)-aryl, -O-C(O)-cycloalkyl, -C(=N-CN)-NH₂, -C(=NH)-NH₂, -C(=NH)-NH(alkyl), Y₁Y₂N-, Y₁Y₂N-alkyl-, Y₁Y₂NC(O)-, Y₁Y₂NSO₂- and -SO₂NY₁Y₂, wherein Y₁ and Y₂ can be the same or different and are independently selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl, and aralkyl. "Ring system substituent" may also mean a single moiety which simultaneously replaces two available hydrogens on two adjacent carbon atoms (one H on each carbon) on a ring system. Examples of such moieties are rings such as heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl rings. Additional non-limiting examples include methylene dioxy, ethylenedioxy, -C(CH₃)₂- and the like which form moieties such as, for example:

15




As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

20 The line ----, as a bond generally indicates a mixture of, or either of, the possible isomers, e.g., containing (R)- and (S)- stereochemistry. For example:



25 The wavy line ~~~, as used herein, indicates a point of attachment to the rest of the compound. For example, each wavy line in the following structure:




2

indicates a point of attachment to the core structure, as described herein.


- 70 -

Lines drawn into the ring systems, such as, for example:



indicate that the indicated line (bond) may be attached to any of the substitutable ring carbon atoms.


- 5 "Oxo" is defined as a oxygen atom that is double bonded to a ring carbon in a cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, or other ring described herein, e.g.,



In this specification, where there are multiple oxygen and/or sulfur atoms in a ring system, there cannot be any adjacent oxygen and/or sulfur present in said ring system.

- 10 It is noted that the carbon atoms for compounds of the invention may be replaced with 1 to 3 silicon atoms so long as all valency requirements are satisfied.

As well known in the art, a bond drawn from a particular atom wherein no moiety is depicted at the terminal end of the bond indicates a methyl group bound through that bond to the atom, unless stated otherwise. For example:



- 15 The term "purified", "in purified form" or "in isolated and purified form" for a compound refers to the physical state of said compound after being isolated from a synthetic process (e.g. from a reaction mixture), or natural source or combination thereof. Thus, the term "purified", "in purified form" or "in isolated and purified form" for a compound refers to the physical state of
- 20 said compound (or a tautomer or stereoisomer thereof, or pharmaceutically acceptable salt or solvate of said compound, said stereoisomer, or said tautomer) after being obtained from a purification process or processes described herein or well known to the skilled artisan (e.g., chromatography, recrystallization and the like), in sufficient purity to be suitable for in vivo or medicinal use and/or characterizable by standard analytical techniques described herein or well
- 25 known to the skilled artisan.

- 71 -

It should also be noted that any carbon as well as heteroatom with unsatisfied valences in the text, schemes, examples and Tables herein is assumed to have the sufficient number of hydrogen atom(s) to satisfy the valences.

When a functional group in a compound is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for example, T. W. Greene *et al*, *Protective Groups in organic Synthesis* (1991), Wiley, New York.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

Prodrugs and solvates of the compounds of the invention are also contemplated herein. A discussion of prodrugs is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems* (1987) 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, (1987) Edward B. Roche, ed., American Pharmaceutical Association and Pergamon Press. The term "prodrug" means a compound (e.g, a drug precursor) that is transformed *in vivo* to yield a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate of the compound. The transformation may occur by various mechanisms (e.g., by metabolic or chemical processes), such as, for example, through hydrolysis in blood. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

For example, if a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate of the compound contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as, for example, (C₁-C₈)alkyl, (C₂-C₁₂)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxy-carbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxy-carbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxy-carbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxy-carbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxy-carbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁-C₂)alkylamino(C₂-C₃)alkyl (such as β-

- 72 -

dimethylaminoethyl), carbamoyl-(C₁-C₂)alkyl, N,N-di (C₁-C₂)alkylcarbamoyl-(C₁-C₂)alkyl and piperidino-, pyrrolidino- or morpholino(C₂-C₃)alkyl, and the like.

Similarly, if a compound of the invention contains an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as, for example, (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl, (C₁-C₆)alkoxycarbonyloxymethyl, N-(C₁-C₆)alkoxycarbonylaminomethyl, succinoyl, (C₁-C₆)alkanoyl, α -amino(C₁-C₄)alkanyl, arylacyl and α -aminoacyl, or α -aminoacyl, where each α -aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate), and the like.

If a compound of the invention incorporates an amine functional group, a prodrug can be formed by the replacement of a hydrogen atom in the amine group with a group such as, for example, R-carbonyl, RO-carbonyl, NRR'-carbonyl where R and R' are each independently (C₁-C₁₀)alkyl, (C₃-C₇) cycloalkyl, benzyl, or R-carbonyl is a natural α -aminoacyl or natural α -aminoacyl, —C(OH)C(O)OY¹ wherein Y¹ is H, (C₁-C₆)alkyl or benzyl, —C(OY²)Y³ wherein Y² is (C₁-C₄) alkyl and Y³ is (C₁-C₆)alkyl, carboxy (C₁-C₆)alkyl, amino(C₁-C₄)alkyl or mono-N— or di-N,N-(C₁-C₆)alkylaminoalkyl, —C(Y⁴)Y⁵ wherein Y⁴ is H or methyl and Y⁵ is mono-N— or di-N,N-(C₁-C₆)alkylamino morpholino, piperidin-1-yl or pyrrolidin-1-yl, and the like.

One or more compounds of the invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. "Solvate" means a physical association of a compound of this invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanlates, methanlates, and the like. "Hydrate" is a solvate wherein the solvent molecule is H₂O.

One or more compounds of the invention may optionally be converted to a solvate. Preparation of solvates is generally known. Thus, for example, M. Caira *et al*, *J. Pharmaceutical Sci.*, 93(3), 601-611 (2004) describe the preparation of the solvates of the antifungal fluconazole in ethyl acetate as well as from water. Similar preparations of solvates, hemisolvate, hydrates and the like are described by E. C. van Tonder *et al*, *AAPS PharmSciTech.*, 5(1), article 12 (2004);

- 73 -

and A. L. Bingham *et al*, *Chem. Commun.*, 603-604 (2001). A typical, non-limiting, process involves dissolving the inventive compound in desired amounts of the desired solvent (organic or water or mixtures thereof) at a higher than ambient temperature, and cooling the solution at a rate sufficient to form crystals which are then isolated by standard methods. Analytical techniques such as, for example I. R. spectroscopy, show the presence of the solvent (or water) in the crystals as a solvate (or hydrate).

"Effective amount" or "therapeutically effective amount" is meant to describe an amount of compound or a composition of the present invention effective in inhibiting the above-noted diseases and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect.

The compounds of the invention can form salts which are also within the scope of this invention. Reference to a compound of the invention herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of the invention contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compounds of the invention may be formed, for example, by reacting a compound of the invention with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides, lactates, maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates,) and the like. Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by P. Stahl *et al*, Camille G. (eds.) *Handbook of Pharmaceutical Salts. Properties, Selection and Use*. (2002) Zurich: Wiley-VCH; S. Berge *et al*, *Journal of Pharmaceutical Sciences* (1977) 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33 201-217; Anderson *et al*, *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug

Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with
5 organic bases (for example, organic amines) such as dicyclohexylamines, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, and dibutyl sulfates), long chain
10 halides (e.g. decyl, lauryl, and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

All such acid salts and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.

Pharmaceutically acceptable esters of the present compounds include the following
15 groups: (1) carboxylic acid esters obtained by esterification of the hydroxy groups, in which the non-carbonyl moiety of the carboxylic acid portion of the ester grouping is selected from straight or branched chain alkyl (for example, acetyl, n-propyl, t-butyl, or n-butyl), alkoxyalkyl (for example, methoxymethyl), aralkyl (for example, benzyl), aryloxyalkyl (for example, phenoxymethyl), aryl (for example, phenyl optionally substituted with, for example, halogen, C₁-
20 4alkyl, or C₁₋₄alkoxy or amino); (2) sulfonate esters, such as alkyl- or aralkylsulfonyl (for example, methanesulfonyl); (3) amino acid esters (for example, L-valyl or L-isoleucyl); (4) phosphonate esters and (5) mono-, di- or triphosphate esters. The phosphate esters may be further esterified by, for example, a C₁₋₂₀ alcohol or reactive derivative thereof, or by a 2,3-di (C₆₋₂₄)acyl
glycerol.


25 Compounds of the invention, and salts, solvates, esters and prodrugs thereof, may exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

The compounds of the invention may contain asymmetric or chiral centers, and, therefore, exist in different stereoisomeric forms. It is intended that all stereoisomeric forms of the
30 compounds of the invention as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, the present invention embraces all geometric and positional isomers. For example, if a compound of the invention incorporates a double bond or a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within the scope of the invention.

- 75 -


Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of the invention may be atropisomers (e.g., substituted biaryls) and are considered as part of this invention. Enantiomers can also be separated by use of chiral HPLC column.

It is also possible that the compounds of the invention may exist in different tautomeric forms, and all such forms are embraced within the scope of the invention. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the invention. Thus, for



example, the compounds of the invention conforming to the formula:

, and



their tautomers:

are both contemplated as being within the scope of the

compounds of the invention.

All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms,

- 76 -

are contemplated within the scope of this invention, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). (For example, if a compound of the invention incorporates a double bond or a fused ring, both the cis- and trans-forms, as well as mixtures, are embraced within the scope of the invention. Also, for example, all keto-enol and imine-enamine forms of the compounds are included in the invention.)

Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the *IUPAC* 1974 Recommendations. The use of the terms "salt", "solvate", "ester", "prodrug" and the like, is intended to equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the inventive compounds.

The present invention also embraces isotopically-labelled compounds of the present invention which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively.

Certain isotopically-labelled compounds of the invention (e.g., those labeled with ^3H and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (i.e., ^3H) and carbon-14 (i.e., ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (e.g., increased in vivo half-life or reduced dosage requirements) and hence may be preferred in some circumstances. Isotopically labelled compounds of the invention can generally be prepared by following procedures analogous to those disclosed in the Schemes and/or in the Examples hereinbelow, by substituting an appropriate isotopically labelled reagent for a non-isotopically labelled reagent. Non-limiting examples of deuterated compounds of the invention are described hereinbelow.

Polymorphic forms of the compounds of the invention, and of the salts, solvates, esters and prodrugs of the compounds of the invention, are intended to be included in the present invention.

- 77 -

Suitable doses for administering compounds of the invention to patients may readily be determined by those skilled in the art, e.g., by an attending physician, pharmacist, or other skilled worker, and may vary according to patient health, age, weight, frequency of administration, use with other active ingredients, and/or indication for which the compounds are administered. Doses may range from about 0.001 to 500 mg/kg of body weight/day of the compound of the invention. In one embodiment, the dosage is from about 0.01 to about 25 mg/kg of body weight/day of a compound of the invention, or a pharmaceutically acceptable salt or solvate of said compound. In another embodiment, the quantity of active compound in a unit dose of preparation may be varied or adjusted from about 1 mg to about 100 mg, preferably from about 1 mg to about 50 mg, more preferably from about 1 mg to about 25 mg, according to the particular application. In another embodiment, a typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 500 mg/day, preferably 1 mg/day to 200 mg/day, in two to four divided doses.

As discussed above, the amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated.

When used in combination with one or more additional therapeutic agents, the compounds of this invention may be administered together or sequentially. When administered sequentially, compounds of the invention may be administered before or after the one or more additional therapeutic agents, as determined by those skilled in the art or patient preference.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically active agent or treatment within its dosage range.

Accordingly, in an aspect, this invention includes combinations comprising an amount of at least one compound of the invention, or a pharmaceutically acceptable salt, solvate, ester or prodrug thereof, and an effective amount of one or more additional agents described above.

The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological assays. Certain assays are exemplified elsewhere in this document.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient.

- 78 -

Suitable solid carriers are known in the art, e.g., magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), *Remington's*
5 *Pharmaceutical Sciences*, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

10 Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms
15 include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

20 The compounds of this invention may also be delivered subcutaneously.

In one embodiment, the compound is administered orally.

In some embodiments, it may be advantageous for the pharmaceutical preparation comprising one or more compounds of the invention be prepared in a unit dosage form. In such forms, the preparation is subdivided into suitably sized unit doses containing appropriate
25 quantities of the active component, e.g., an effective amount to achieve the desired purpose.

PREPARATIVE EXAMPLES

Compounds of the invention can be made using procedures known in the art. The following reaction schemes show typical procedures, but those skilled in the art will recognize that other procedures can also be suitable.

30 Where NMR data are presented, spectra were obtained on either a Varian VXR-200 (200 MHz, ¹H), Varian Gemini-300 (300 MHz) or XL-400 (400 MHz), or Bruker AVANCE 300 or 500 MHz spectrometers and are reported as ppm (δ) down field from Me₄Si with number of

- 79 -

protons, multiplicities, and coupling constants in Hertz indicated parenthetically. Optical rotation data was obtained on a Perkin Elmer 341 polarimeter and substrate concentration *c* is reported in mg/mL.

Techniques, solvents and reagents may be referred to by their following abbreviations:

- 5 Thin layer chromatography: TLC
High performance liquid chromatography: HPLC
ethyl acetate: AcOEt or EtOAc
methanol: MeOH
ethanol: EtOH
- 10 ether or diethyl ether: Et₂O
tetrahydrofuran: THF
Acetonitrile: MeCN
1,2-dimethoxyethane: DME
Trifluoroacetic acid: TFA
- 15 Deoxofluor: bis-(2-methoxyethyl) aminosulfur trifluoride
Dimethylacetamide: DMA
Dimethylformamide: DMF
Dimethylsulfoxide: DMSO
triethylamine: Et₃N or TEA
- 20 tert-Butoxycarbonyl: t-Boc or Boc
2-(Trimethylsilyl)ethoxycarbonyl: Teoc
nuclear magnetic resonance spectroscopy: NMR
liquid chromatography mass spectrometry: LCMS
high resolution mass spectrometry: HRMS
- 25 liters : L
milliliters: mL
millimoles: mmol
microliters: μ l (or μ L)
grams: g
- 30 milligrams: mg
centimeters: cm
room temperature (ambient, about 25°C): rt (or RT)


- 80 -

- minutes: min
Retention time: t_R
hours: h (or hr)
N-bromosuccinimide: NBS
- 5 Methyl magnesium bromide: MeMgBr
iron(III) acetylacetonate: Fe(acac)₃
Diphenylphosphoryl azide: DPPA
1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride: EDCI
Diisopropylethylamine: DIEA or iPr_2NEt
- 10 Diisopropylamine: iPr_2NH
2-(Trimethylsilyl)ethanol: TMSethanol
3-Chloroperoxybenzoic acid: mCPBA
n-Butyllithium: nBuLi
lithium diisopropylamide: LDA
- 15 [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II): PdCl₂dppf
Palladium(II) acetate: Pd(OAc)₂
Methanesulfonyl chloride: MeSO₂Cl
Benzyl: Bn
saturated: Sat.
- 20 round bottom flask: RBF
acetonitrile: MeCN
butyl: Bu
4-methoxy benzyl: PMB
Sodium methoxide: NaOMe
- 25 Hexane: hex (or hex.)
Molar: M
aqueous: aq.
acetic acid: AcOH (or HOAc)
methylene chloride: DCM
- 30 reverse phase: RP
dichloro ethane: DCE
phenyl: Ph
preparative: prep (or prep.)

- 81 -

XPhos: 2-dicyclohexylphosphino-2',4',6'-trisopropylbiphenyl

Scheme 1



5 Step 1:

To a solution of 2,4-difluoroacetophenone (15.0 g, 96 mmol) in THF (100 mL) was added (R)-2-methyl-2-propanesulfonamide (12.8 g, 106 mmol) and $\text{Ti}(\text{OEt})_4$ (32.0 g, 120 mmol). The resultant solution was heated to reflux overnight. After that time, the solution was cooled to RT and poured onto ice. To this mixture was added CH_2Cl_2 and the resultant mixture was stirred at
 10 RT for 10 min. The mixture was then filtered through Celite. The filter cake was washed with CH_2Cl_2 . The layers were separated. The aqueous layer was extracted with CH_2Cl_2 (2x). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The crude product was purified via flash chromatography (SiO_2 ; gradient elution 100:0 to 45:55 hexanes:EtOAc) to afford the ketimine (12.3 g).

15

Step 2:

To a stirred solution of 4-methoxybenzylamine (198.9 g, 1.45 mol) in anhydrous pyridine (400 mL) at 0°C was added dropwise via an addition funnel methanesulfonyl chloride (116 mL, 1.45 mol) over 45 min. After the addition was complete, the cooling bath was removed and the

- 82 -

resultant solution was stirred at RT overnight. The reaction was concentrated *in vacuo* (water bath 60-65°C) to remove most of the pyridine. The brown slurry was taken up in CH₂Cl₂ (1 L). The organic solution was washed with 1 N HCl_(aq.) (2 x 1 L), sat. NaHCO₃ (aq) (2 x 1 L) and brine (1 x 500 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated to afford
5 a crude solid. This solid was dissolved in 95% EtOH (430 mL) using a steam bath to warm the solution. The solution was allowed to cool, causing the product to solidify out of solution. The product was removed by filtration and the solid was washed with cold EtOH (3 x 150 mL). A second crop was obtained after allowing the mother liquor to stir at RT overnight. The overall yield of the product was 246.5 g (79% yield) as a pale orange crystalline solid.

10 This product was dissolved in anhydrous DMF (3.0 L), cooled to 0°C and placed under an atmosphere of N₂. To this solution was added in small portions sodium hydride (60% in mineral oil, 60.2 g, 1.51 mol, 1.3 eq.). After the addition was complete, the mixture was stirred for an additional 10 min. To this mixture was added dropwise via an addition funnel methyl iodide (250
15 g, 1.76 mol, 1.5 eq.). After the addition was complete, the cooling bath was removed and the mixture was allowed to stir at RT overnight. The mixture was then concentrated *in vacuo* (pressure = 10 torr, bath temp = 55-60°C) to remove ca. 2.5 L of DMF. The product was partitioned between 5 L ice water, 5 L Et₂O and 500 mL of EtOAc. The organic layer was separated. The aqueous layer was extracted with Et₂O (2 x 1 L). The combined organic layers were washed with brine (2 x 1 L), dried over Na₂SO₄, filtered and concentrated. The oily solid
20 was stirred with hexanes using a wire stir blade to powderize the solid. The solid was removed by filtration and washed with hexanes (2 x 250 mL). The solid was dissolved in hexanes/EtOAc (1:1, 450 mL) using a steam bath to warm the mixture. An off white precipitate formed on cooling and was filtered off (182 g). The remaining mother liquor was purified via flash chromatography (SiO₂: 1:1 hexanes:EtOAc) to afford additional product (51.8 g) for a total yield
25 of 233.8g (89% yield).

Step 3:

To a solution of the sulfonamide from step 2 (4.18 g, 18.2 mmol) in anhydrous THF (50 mL) at -78°C under an atmosphere of N₂ was added dropwise a solution of *n*-BuLi (1.6 M in hexanes, 11.4 mL, 18.2 mmol). The resultant solution was stirred at -78°C for 30 min. After that
30 time, a solution of the ketimine from step 1 (3.15 g, 12.1 mmol) in THF (50 mL) precooled to -78°C in a separate round bottom flask was transferred via cannula into the solution above. The resultant solution was stirred at -78°C for 3.5 hours. After that time, water was added and the mixture was allowed to warm to RT. The aqueous layer was extracted with EtOAc (3x). The

- 83 -

combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated. The crude product was purified via flash chromatography (SiO_2 : gradient elution 100:0 to 40:60 hexanes:EtOAc) to afford the sulfinamide (3.95 g, 67% yield).

Step 4:

5 To a solution of the sulfinamide from step 3 (3.80 g, 7.6 mmol) in $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (3:1 80 mL) was added a solution of 4 M HCl (dioxane) (11.4 mL, 45.4 mmol). The resultant solution was stirred at RT for 1.5 hours. The solution was concentrated. The residue was reconcentrated from toluene (1x). The residue was then taken up in CHCl_3 and TFA (26 mL, 1:1). To this solution was added 1,3-dimethoxybenzene (6.5 mL, 50 mmol). The resultant solution was stirred at RT
10 overnight. The resultant dark pink solution was concentrated. The oil was partitioned between Et_2O and 1 M HCl (aq.). The aqueous layer was extracted with Et_2O (2x). The aqueous layer was then adjusted to pH 10 with the addition of sat. Na_2CO_3 (aq.). The aqueous layer was extracted with CH_2Cl_2 (3x). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated to afford the amine (1.88 g, 85%) as a clear oil.

15 Step 5:

To a solution of the amine from step 4 (1.80 g, 6.8 mmol) in CH_2Cl_2 (30 mL) was added benzoyl isothiocyanate (1.01 mL, 7.49 mmol). The resultant solution was stirred at RT overnight. After that time, the solution was concentrated. The residue was redissolved in MeOH (20 mL). To this solution was added a solution of NaOMe in MeOH (25%, 3.9 mL). The resultant solution
20 was stirred at RT for 45 min. The solution was then concentrated. The residue was partitioned between CH_2Cl_2 and water. The pH of the aqueous layer was adjusted to ca 8 with the addition of NaHCO_3 (aq.). The aqueous layer was extracted with CH_2Cl_2 (3x). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated to afford the thiourea (1.90 g, 86%).

Step 6:

25 To the thiourea from step 5 (1.90 g, 5.88 mmol) in EtOH (40 mL) was added methyl iodide (0.42 mL, 6.7 mmol). The resultant solution was heated to reflux for 3 hours. The solution was cooled to RT and concentrated *in vacuo*. The residue was partitioned between EtOAc and Na_2CO_3 (aq.). The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated. The crude
30 product was purified via flash chromatography (SiO_2 : gradient elution 100:0 to 92:8 CH_2Cl_2 :MeOH) to afford the thiadiazine dioxide (1.12 g, 66% yield).


- 84 -

Table I: The following imines were prepared using a method similar to that described in Scheme 1 Step 1.

Entry	Ketone	Imine
1		
2		
3		

5 **Table II:** The following thiadiazine dioxides were prepared using methods similar to that described in Scheme 1.

Entry	Ketone	Thiadiazine dioxide
1		
2		

Scheme 2:

- 85 -

To a solution of N-(4-methoxybenzyl)-N-methylmethanesulfonamide (26.8 g, 117 mmol) in THF (200 mL) at -78 °C was added n-butyllithium (2.5 M in hexanes, 47 mL, 118 mmol) over 10 minutes. After the addition was complete, the mixture was allowed to stir at -78 °C for 1h.


To this mixture was then added a solution of (S)-2-methyl-N-(1-(2,4,6-trifluorophenyl)ethylidene)propane-2-sulfinamide (21.6 g, 77.9 mmol, prepared from 2,4,6-trifluoroacetophenone and (S)-2-methyl-2-propanesulfinamide according to Scheme 1, Step 1) in THF (150 mL) at -78 °C. The resulting mixture was allowed to stir at -78 °C for 4h. At that time, the reaction was quenched by rapid dilution with water (~ 400 mL). The mixture was then warmed to RT, further diluted with EtOAc and brine. The phases were separated, and the aqueous layer was extracted with EtOAc (4X). The organic portions were combined, washed with brine, dried over MgSO₄, filtered and concentrated. This crude residue was subjected to column chromatography (600g silica, 100 mL/min, 0% to 60% EtOAc/hexanes) to give (R)-2-((S)-1,1-dimethylethylsulfinamido)-N-(4-methoxybenzyl)-N-methyl-2-(2,4,6-trifluorophenyl)propane-1-sulfonamide as a 4:1 mixture with its diastereomer (14.5 g total mass, 37%).

This material was further subjected to SFC chromatography (TharSFC80, Chiralpak OJ-H, 21 x 250 mm, 5 mm, 200 bar with 5% MeOH, 55 g/min, 35 °C) to give (R)-2-((S)-1,1-dimethylethylsulfinamido)-N-(4-methoxybenzyl)-N-methyl-2-(2,4,6-trifluorophenyl)propane-1-sulfonamide).

The above material was treated according to Scheme 1, Steps 4–6 to afford the thiadiazine dioxide A.

Scheme 3:

- 86 -



Step 1:

To a solution of 4-methoxy-N-methylbenzylamine (2.04 g, 13.5 mmol) in CH_2Cl_2 (30 mL) at 0°C was added Et_3N (2.44 mL, 17.5 mmol) followed by the dropwise addition of benzylsulfonyl chloride (2.96g, 15.5 mmol). The solution was stirred at 0°C for 30 minutes. The solution was then warmed to RT and stirred overnight. After that time, the mixture was diluted with CH_2Cl_2 , washed with 1 M $\text{HCl}_{(\text{aq})}$ and 1/2 saturated $\text{NaHCO}_3_{(\text{aq})}$. The organic layer was then dried over Na_2SO_4 , filtered and concentrated. The crude residue was purified via flash chromatography (SiO_2 : gradient elution 100:0 to 1:1 hexanes:EtOAc) to afford the sulfonamide (3.5g, 85%) as an off white solid.

Step 2:

To a solution of the sulfonamide from step 1 (1.95 g, 6.38 mmol) in toluene (40 mL) at -78°C was added dropwise a solution of n-BuLi in hexane (1.6 M, 4.0 mL). The resultant solution was stirred at -78°C for 30 minutes. To a separate round bottom flask containing the ketimine from Entry 3 Table I (1.30 g, 4.21 mmol) in toluene (30 mL) at -78°C was added dropwise a solution of trimethylaluminum in toluene (2.0 M, 2.32 mL). The resultant solution was stirred at -78°C for 5 min. This solution was then transferred via cannula to the solution of the sulfonamide anion. The resultant solution was stirred at -78°C for 2.25 hours. Water was added and the mixture was warmed to RT. The mixture was diluted with EtOAc and filtered through Celite. The layers were separated. The organic layer was dried over Na_2SO_4 , filtered and concentrated. The crude product was purified via flash chromatography (SiO_2 : gradient elution: 100:0 to 66:34


- 87 -

hexanes:EtOAc) to afford 510 mg of the faster eluting isomer **B** and 320 mg of the slower eluting stereoisomer **C**.

Example 1 was prepared from compound **B** using methods similar to that described in Scheme 1 steps 4-6. LCMS data: Obs. MH^+ : 416.2, Ret. Time: 3.35 min, LCMS method: A.

Example 2 was prepared from compound **C** using methods similar to that described in Scheme 1 steps 4-6. LCMS data: Obs. MH^+ : 416.2, Ret. Time: 2.77 min, LCMS method: A.

10 Scheme 4:



Step 1:

To a solution of 4-(trifluoromethyl)benzylsulfonyl chloride (10.0g, 38.7 mmol) in CH_2Cl_2 (100 mL) at $0^\circ C$ was added dropwise a solution of methylamine (2 M in THF, 116 mL). The solution was allowed to slowly warm to RT over 2 hours. After that time, the mixture was concentrated. The residue was partitioned between CH_2Cl_2 and $\frac{1}{2}$ saturated $NaHCO_3$ (aq.). The aqueous layer was extracted with CH_2Cl_2 (2x). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The product was recrystallized from EtOAc/hexanes to afford the sulfonamide (7.71 g, 79% yield) as a white solid.

20 Step 2:

To a solution of the above sulfonamide (9.80 g, 38.7 mmol) in anhydrous THF (200 mL) at $-78^\circ C$ under an atmosphere of N_2 was added a solution of n-BuLi (1.6 M in hexanes, 48.8

- 88 -

mL, 78 mmol). The resultant solution was stirred at -78°C for 30 min. After that time, a precooled solution (-78°C) of the ketimine (7.22 g, 27.9 mmol) in THF (100 mL) was transferred via cannula to the solution of the sulfonamide anion. The resultant solution was stirred at -78°C for 2.5 hours. Water was added to the solution and the mixture was warmed to RT. The aqueous layer was adjusted to ca. pH 8 with the addition of 1 M HCl_(aq.) and sat. NaHCO_{3(aq.)}. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude residue was purified via flash chromatography (SiO₂; gradient elution 100:0 to 30:70 hexanes:EtOAc) to afford **D** (4.64 g, 33% yield) as a mixture of diastereomers.

10


Step 3:

To a solution of **D** (4.60 g, 8.97 mmol) in 5:2 CH₂Cl₂:MeOH (140 mL) was added a solution of HCl (4 M in dioxane, 13.5 mL, 53.8 mmol). The solution was stirred at RT for 40 min. After that time, the solution was concentrated. The residue was partitioned between CH₂Cl₂ and sat. Na₂CO_{3(aq.)}. The aqueous layer was extracted with CH₂Cl₂ (3x). The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The crude residue was purified via flash chromatography (SiO₂; gradient elution 100:0 to 40:60 hexanes:EtOAc) to afford **E** (3.34 g, 91% yield) as a mixture of diastereomers.

20 **Example 3** and **Example 4** were prepared from **E** using methods similar to that described in Scheme 1 steps 5 and 6. LCMS data (Ex. 3): Obs. MH⁺: 434.2, Ret. Time: 3.17 min, LCMS method: A. LCMS data (Ex. 4): Obs. MH⁺: 434.2, Ret. Time: 3.29 min, LCMS method: A.


Scheme 5:

25




- 89 -

Sodium hydride (60% in oil, 1.5 g, 37.5 mmol, 1.2 equiv) was added to a solution of 5-bromoindazole **F** (6 g, 30.6 mmol, 1 equiv) in DMF (60 mL) at RT. After stirring for 30 min, methyl iodide (2.83 mL, 45.9 mmol, 1.5 equiv) was added and the reaction stirred for another 2 h at RT. The reaction was quenched with sat. NaHCO₃, extracted with EtOAc (1x), dried over MgSO₄, filtered, and concentrated under reduced pressure to give a mixture of *N*-1 and *N*-2 methylated 5-bromoindazoles **G** and **H**, which were separated by silica-gel chromatography using 0→30 % EtOAc/hexanes as eluent. The *N*1-alkylated regioisomer **G** elutes first, followed by the *N*2-methyl regioisomer **H**. Other *N*-1-alkylated 5-bromoindazoles (**I**, **K**, **M**) were prepared by the same procedure, substituting the appropriate electrophile for methyl iodide (ethyl iodide, *i*-propyl iodide, *n*-propyl iodide).

Scheme 6:

5-Bromoindazole (3 g, 15.5 mmol, 1 equiv), cyclopropyl boronic acid (2.66 g, 31 mmol, 2 equiv), Cu(OAc)₂ (2.81 g, 15.5 mmol, 1 equiv), Na₂CO₃ (3.29 g, 31 mmol, 2 equiv), bipyridine (2.42 g, 15.5 mmol, 1 equiv) were suspended in DCE (150 mL) and stirred for 4 h at RT. The reaction was quenched with sat. aqueous NH₄Cl, and extracted with CH₂Cl₂ (2x). The combined organic layers were washed with brine (1x), then dried over Na₂SO₄, filtered, and concentrated under reduced pressure. 5-bromoindazoles **O** and **P** were separated by silica-gel chromatography using 0→30 % EtOAc/hexanes as eluent to give *N*-1-cyclopropyl regioisomer **O** in 68% yield (2.5 g, 10.5 mmol, first compound to elute).



Scheme 7:

LiHMDS (1 M in THF, 21 mL, 21 mmol, 1.3 equiv) was added to a -20°C solution of sulfonamide (Scheme 1, Step 2) (3.7 g, 16.3 mmol, 1 equiv) in THF (20 mL) in a flame-dried round-bottom flask. After stirring for 60 min, a ZnCl₂-solution (1.2 M in THF, 21.5 mL, 17.9


- 90 -

mmol, 1.1 equiv) was added and the reaction warmed to RT over 45 min. *N*-1-methyl-5-bromoindazole **G** (3.2 g, 16.3 mmol, 1 equiv), Pd(OAc)₂ (183 mg, 0.81 mmol, 0.05 equiv), X-Phos (777 mg, 1.63 mmol, 0.1 equiv) in THF (15 mL) was added, and the reaction degassed with three cycles of vacuum/N₂, then placed in a preheated 65°C oil bath. After stirring for 18 h, the reaction mixture was cooled to RT, diluted with EtOAc and sat. aqueous NH₄Cl, extracted with EtOAc (1x), dried over MgSO₄, filtered, and concentrated under reduced pressure to give a residue, which was subjected to silica-gel chromatography using 0→50 % EtOAc/hexanes as eluent to give sulfonamide **Q** as a solid in 63% yield (3.7 g, 10.3 mmol).

Table III: The following arylated sulfonamide was prepared using methods similar to those described in Scheme 7.

Aryl bromide	Product
	

Scheme 8:



n-BuLi (1.6 M in hexanes, 9.3 mL, 15 mmol, 1.5 equiv) was slowly added to a solution of sulfonamide **Q** (3.59 g, 10 mmol, 1 equiv) in THF (70 mL) at -78°C in a flame-dried 250 mL RBF. After 30 min, a solution of the ketimine (Entry 2, Table I) (2.77 g, 10 mmol, 1 equiv) in THF (25 mL) was slowly added via cannula. After stirring for 2.5 h, the reaction mixture was quenched with sat. aqueous NH₄Cl, extracted with EtOAc (3x), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give a residue, which was subjected to silica-gel




- 91 -

chromatography using 0→80 % EtOAc/hexanes as eluent to give aldol adduct **R** as a solid in 52% yield (3.3 g, 5.18 mmol).


Examples 5 and 6 were prepared from **R** using methods similar to that described in Scheme 1 Steps 4-6. LCMS data (Ex. 5): Obs. MH^+ : 438.2, Ret. Time: 1.84 min, LCMS method: C.

5 LCMS data (Ex. 6): Obs. MH^+ : 438.2, Ret. Time: 1.87 min, LCMS method: C.

Table IV: The following example was prepared using methods similar to those described in Scheme 8.

Sulfonamide	Ketimine	Example		LCMS Obs. MH^+	LCMS Ret. Time (min)	LCMS method
 Table 3	 Scheme 2	7		425.2	3.11	B

Scheme 9:



10

Sulfonamide **H** was prepared using methods described in Zhou, G. *et al.* *Org. Lett.* **2008**, 10, 2517.





15


Table Va: The following sulfonamides were prepared using methods similar to those described in Scheme 4 step 1.

Sulfonyl Chloride	Sulfonamide

Table Vb: The following examples were prepared using methods similar to those described in Scheme 4.



Sulfonamide	Ketimine	Example	LCMS Obsv. MH ⁺	LCMS Ret. Time (min)	LCMS method
 Table Va	 Table I		446.2	3.33	A
 Scheme 4	 Table I		434.2	3.33	A
			434.2	3.35	A
 SF ₅	 Table I		492.3	3.53	A

Scheme 9		12		492.3	3.50	A
 Table Va	 Table I	13		366.2	2.71	A





Scheme 10:

To a solution of the thiazidine dioxide (Table II, Entry 1) (3.8 g, 12.2 mmol) in MeCN
 5 (40 mL) was added 4-methoxybenzyl chloride (4.6 g, 29 mmol), Cs₂CO₃ (9.9 g, 31 mmol) and n-Bu₄NI (450 mg, 1.2 mmol). The resultant mixture was heated to reflux for 16 hours. After that time, additional 4-methoxybenzyl chloride (1.9 g, 12 mmol) and Cs₂CO₃ (4.4 g, 12 mmol) were added and the mixture was heated to reflux for an additional 4 hours. The mixture was then concentrated *in vacuo* at RT. The residue was partitioned between water and CH₂Cl₂. The
 10 aqueous layer was extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄, filtered and concentrated. The crude residue was purified via flash chromatography (SiO₂: gradient elution 100:0 to 80:20 hexanes:EtOAc) to afford the bis-PMB compound **T** (4.9 g, 73%).


Table VI: The following compounds were prepared using a method similar to that described in
 15 Scheme 10.

Entry	Iminothiadiazine dioxide	bis-PMB core
1		

- 94 -

2		
3		 Note 1

Note 1: Entry 3 was prepared as described in Scheme 10 with the following exception: excess diethylamine was added to the mixture. The mixture was stirred at RT overnight and filtered. The filtrate was concentrated and the residue was subjected directly to flash chromatography.

5 **Scheme 11:**

Step 1:

To a solution of ethyl 4-bromophenylacetate (4.5 g, 18.5 mmol) in 40 mL of DMF was added iodomethane (4.03 mL, 64.8 mmol) and sodium t-butoxide (4.45 g, 46.3 mmol). The resulting mixture was stirred at room temperature for 2 h. The reaction was quenched with water, and the mixture was extracted with EtOAc and hexane. The organic layer was separated and washed with saturated sodium bicarbonate solution. The organic layer was separated, dried over MgSO₄, and concentrated to give ethyl 2-(4-bromophenyl)-2-methylpropanoate (3.64 g, 73%).

15 Step 2:

To a solution of the material from Step 1 (2.4 g, 8.9 mmol) in 30 mL of THF at -78°C was added LiAlH₄ (337mg, 8.9mmol). The mixture was stirred at or below -60°C for 2 hr, and then stirred at room temperature for 1 hr. The reaction was quenched with 10% NaOH (aq) solution, and the mixture was extracted with EtOAc. The organic layer was separated, dried over MgSO₄, and concentrated to get 2-(4-bromophenyl)-2-methylpropan-1-ol (2.03g, 100%).

Step 3:


- 95 -

To a solution of the material from Step 2 (500 mg, 2.18 mmol) in 8 mL of DMF was added iodomethane (0.68 mL, 10.9 mmol) and NaH (60 % in oil, 131 mg, 3.28 mmol). The mixture was stirred at room temperature for 2 h. The reaction was quenched with water, and extracted with EtOAc and hexane. The organic layer was separated, washed with saturated sodium bicarbonate, dried over MgSO₄, and concentrated. The crude was purified by flash silica column (eluting with 5% EtOAc in hexane) to get 1-bromo-4-(1-methoxy-2-methylpropan-2-yl)benzene (460 mg, 87%).

- 10 **Table VII:** The following aryl bromides were prepared from ethyl 4-bromophenylacetate using methods similar to that described in Scheme 11.

Step 1 conditions	Step 3 conditions	Product
NaOtBu, CH ₃ I DMF	EtBr, NaH DMF	
Br-CH ₂ -CH ₂ -Br NaH, DMF 0 °C → RT	EtBr, NaH DMF	

Scheme 12:




15

To a pressure tube containing 1-(4-bromophenyl)propan-1-one (5.0 g, 24 mmol) was added Deoxyfluor® (7.8 g, 36 mmol). The tube was sealed and the mixture was heated to 85°C with stirring overnight. After that time, the mixture was cooled to RT and poured onto ice water. The aqueous layer was adjusted to pH ~8. The aqueous layer was extracted with CH₂Cl₂. The organic layer was then dried over Na₂SO₄, filtered and concentrated. The crude residue was purified via flash chromatography (SiO₂; gradient elution: 100:0 to 92:8 hexanes:EtOAc) to afford 1-bromo-4-(1,1-difluoropropyl)benzene (0.8 g).

20

- 96 -

Scheme 13:



Step 1:

To a capped, flame dried microwave vial containing the bis-PMB thiadiazine dioxide (Table VI, entry 2) (159 mg, 0.300 mmol) in dioxane under an atmosphere of N₂ was added a solution of NaHMDS (1 M in THF, 0.35 mL). The resultant mixture was stirred at RT for 30 min. To the mixture was added a freshly prepared solution of ZnCl₂ (1.2 M in THF, 0.313 mL). The resultant mixture was stirred for an additional 30 min. To the mixture was added Pd(OAc)₂ (13.5 mg, 0.0600 mmol), X-Phos (57.2 mg, 0.120 mmol) and the aryl bromide (131 mg, 0.540 mmol). The mixture was then degassed by bubbling N₂ through the mixture for 5 min. The vial was then placed into a preheated oil bath (100°C) and stirred at that temperature for 3 hours. After that time, the mixture was diluted with water and EtOAc. The mixture was then filtered through Celite®. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude residue was purified via reverse phase flash chromatography (C18; gradient elution 90:10:0.1 to 0:100:0.1 water:MeCN:formic acid) to afford the arylated product **U** (87 mg, 43% yield) as a light yellow foam.

Step 2:






















To a solution of **U** (87 mg, 0.12 mmol) in MeCN (5.9 mL) at 75°C was added a heated and fully dissolved solution of sodium persulfate (435 mg, 1.61 mmol) and potassium phosphate dibasic (153 mg, 0.880 mmol) in water (2.9 mL). The resultant mixture was stirred at 75°C for 45 min. After that time, the mixture was cooled to RT and diluted with water and EtOAc. The pH of the aqueous layer was adjusted to ca. 10 with the addition of sat. Na₂CO₃ (aq). The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated. The crude product was purified via flash chromatography [SiO₂; gradient elution 100:0:0 to 94:6:0.5 CH₂Cl₂:MeOH: 7N NH₃ (MeOH)] to afford a semi-crude product that was repurified via preparative TLC (SiO₂; 95:5 CH₂Cl₂:MeOH) to afford **Ex. 14** that was converted to the TFA salt with the addition of a slight excess of TFA






















- 97 -






















in DCM followed by concentration under reduced pressure (38 mg, 53% yield). LCMS data (Ex. 14): Obs. MH^+ : 452.2, Ret. Time: 3.10 min, LCMS method: A.



















5 **Table VIII:** The following examples were prepared from the thiadiazine dioxides described in Scheme 10 or Table VI using methods similar to that described in Scheme 13.



















Thiadiazine dioxide	Aryl bromide	Example	LCMS Obs. MH^+	LCMS Ret. Time (min)	LCMS method
		15 	380.2	3.76	A
		16 	430.2	4.02	A
	 WO2006106416	17 	436.2	3.52	A
		18 	434.2	3.01	A
		19 	446.2	2.41	A



















	 <p>Scheme 5</p>	<p>20</p>		<p>434.2</p>	<p>1.84</p>	<p>C</p>
	 <p>Scheme 5</p>	<p>21</p>		<p>420.2</p>	<p>1.90</p>	<p>C</p>
	 <p>Scheme 6</p>	<p>22</p>		<p>446.2</p>	<p>1.98</p>	<p>C</p>
	 <p>Scheme 5</p>	<p>23</p>		<p>448.2</p>	<p>2.02</p>	<p>C</p>
	 <p>Scheme 12</p>	<p>24</p>		<p>444.0</p>	<p>1.99</p>	<p>C</p>
		<p>25</p>		<p>422.2</p>	<p>4.78</p>	<p>A</p>
		<p>26</p>		<p>392.2</p>	<p>4.23</p>	<p>A</p>

	 <p>Scheme 5</p>	<p>27</p>		<p>448.2</p>	<p>1.97</p>	<p>C</p>
		<p>28</p>		<p>433.2</p>	<p>4.04</p>	<p>A</p>
	 <p>Table VII</p>	<p>29</p>		<p>466.3</p>	<p>4.57</p>	<p>A</p>
	 <p>Table VII</p>	<p>30</p>		<p>464.3</p>	<p>4.39</p>	<p>A</p>
		<p>31</p>		<p>431.0</p>	<p>1.84</p>	<p>C</p>
		<p>32</p>		<p>398.2</p>	<p>2.00</p>	<p>C</p>
		<p>33</p>		<p>442.2</p>	<p>0.95</p>	<p>C</p>

		<p>34</p>		<p>435.2</p>	<p>1.95</p>	<p>C</p>
		<p>35</p>		<p>422.2</p>	<p>4.95</p>	<p>A</p>
		<p>36</p>		<p>492.0</p>	<p>2.10</p>	<p>C</p>
		<p>37</p>		<p>430.0</p>	<p>1.92</p>	<p>C</p>
	 <p>WO2006106416</p>	<p>38</p>		<p>446.0</p>	<p>1.95</p>	<p>C</p>
		<p>39</p>		<p>436.2</p>	<p>1.83</p>	<p>C</p>
	 <p>Scheme 5</p>	<p>40</p>		<p>434.0</p>	<p>1.84</p>	<p>C</p>


		<p>41</p>		<p>452.0</p>	<p>1.91</p>	<p>C</p>
		<p>42</p>		<p>434.0</p>	<p>1.88</p>	<p>C</p>
	 <p>Scheme 12</p>	<p>43</p>		<p>444.0</p>	<p>1.98</p>	<p>C</p>
	 <p>Scheme 5</p>	<p>44</p>		<p>448.2</p>	<p>3.37</p>	<p>B</p>
	 <p>Scheme 5</p>	<p>45</p>		<p>420.2</p>	<p>1.87</p>	<p>C</p>
	 <p>Scheme 5</p>	<p>46</p>		<p>448.2</p>	<p>1.99</p>	<p>C</p>

		<p>47</p>		<p>468.0</p>	<p>2.01</p>	<p>C</p>
		<p>48</p>		<p>450.2</p>	<p>1.98</p>	<p>C</p>
	 <p>Table VII</p>	<p>49</p>		<p>466.3</p>	<p>4.64</p>	<p>A</p>
	 <p>Scheme 11</p>	<p>50</p>		<p>452.2</p>	<p>4.33</p>	<p>A</p>
		<p>51</p>		<p>431.2</p>	<p>1.86</p>	<p>C</p>
		<p>52</p>		<p>380.2</p>	<p>1.98</p>	<p>C</p>

		<p>53</p>		<p>444.2</p>	<p>0.94</p>	<p>C</p>
		<p>54</p>		<p>440.2</p>	<p>3.58</p>	<p>A</p>
		<p>55</p>		<p>452.0</p>	<p>1.98</p>	<p>C</p>
	 <p>WO2006106416</p>	<p>56</p>		<p>454.2</p>	<p>1.88</p>	<p>C</p>
	 <p>Scheme 12</p>	<p>57</p>		<p>462.0</p>	<p>2.03</p>	<p>C</p>
		<p>58</p>		<p>448.2</p>	<p>4.29</p>	<p>A</p>

Scheme 14:

- 104 -



Step 1:

A flame-dried microwave tube was charged with 5R-(2,4,6-trifluorophenyl)-5,6-dihydro-N,N-bis[(4-methoxyphenyl)methyl]-2,5-dimethyl-2H-1,2,4-thiadiazin-3-amine-1,1-dioxide (0.50 g, 0.91 mmol) and dioxane (2 mL). To this mixture at RT was added sodium hexamethyldisilazane (1.0M in THF, 2.3 mL, 2.3 mmol) dropwise via syringe. After 30 min, a zinc dichloride solution (1.2 M in THF, 2.0 mL, 2.4 mmol) was added via syringe. After an additional 30 min, 1-bromo-4-trifluoromethylbenzene (0.23 mL, 1.6 mmol), palladium(II) acetate (0.041 g, 0.18 mmol) and X-Phos (0.17 g, 0.36 mmol) were added, and the mixture was degassed by evacuation and back-fill with N₂ (5X). The nitrogen line was removed, and tube was immersed in an oil bath at 100 °C. After 3h, the reaction was cooled and diluted with 10% w/v citric acid and EtOAc. The mixture was stirred vigorously 5 minutes. The phases were then separated and the aqueous layer was extracted 2X with EtOAc. The organic portions were combined, dried over MgSO₄, filtered and concentrated. This crude sample was subjected to column chromatography (80 g silica, 65 mL/min, 0% to 50% EtOAc/hexanes) to give arylated product 6(R)-[4-(trifluoromethyl)phenyl]-5R-(2,4,6-trifluorophenyl)-5,6-dihydro-N,N-bis[(4-methoxyphenyl)methyl]-2,5-dimethyl-2H-1,2,4-thiadiazin-3-amine-1,1-dioxide (330 mg, 52%).













Step 2:

A large microwave tube was charged with the product of step 1 (0.33 g, 0.48 mmol) and MeCN (25 mL). This mixture was immersed in an oil bath at 85 °C with stirring. After 5 minutes, a solution of potassium persulfate (0.77 g, 2.9 mmol), potassium phosphate monobasic (97 mg, 0.72 mmol), and potassium phosphate dibasic (120 mg, 0.72 mmol) in water (12 mL) also at 85 °C was added. The resulting mixture was heated at 85 °C under N₂. After 1h, the reaction was cooled and then diluted with EtOAc and sat. aq. NaHCO₃ and stirred vigorously for 5 min. The phases were separated and the aqueous layer was extracted 2X with EtOAc. The organic portions were combined, dried over MgSO₄, filtered and concentrated. This crude sample was subjected first to column chromatography (40 g silica, 45 mL/min, 0% to 5% 7N NH₃/MeOH in DCM) to

- 105 -

give a product. This material was further subjected to RP-HPLC (C_{18} radial compression, 35 mL/min, 10% to 95% MeCN/H₂O with 0.1% TFA) to give **Example 59** (76 mg; 28%). LCMS data (Ex. 59): Obs. MH^+ : 452.0, Ret. Time: 3.55 min, LCMS method: B.


5 **Table IX:** The following examples were prepared using methods similar to that described in Scheme 14.

Thiadiazine dioxide	Aryl bromide	Example		LCMS Obs. MH^+	LCMS Ret. Time (min)	LCMS method
	 WO2006106416	60		454.2	3.11	B
		61		453.0	3.11	B
		62		448.2	3.48	B
	 Scheme 11	63		470.2	3.75	B

10

Scheme 15:

- 106 -




Step 1:

A flame dried flask was charged with anhydrous ZnCl_2 (5.13 g, 37.6 mmol) and THF (29 mL). Once a clear solution was obtained, the flask was immersed in a cooling bath at -20°C . To this mixture was added LHMDS (1.0 M in THF, 34.2 mL, 34.2 mmol) via syringe. The resulting mixture was stirred for ~ 1 h while the bath was kept at -20°C .

Meanwhile, another flame-dried flask was charged with 5R-(2,4,6-trifluorophenyl)-5,6-dihydro-N,N-bis[(4-methoxyphenyl)methyl]-2,5-dimethyl-2H-1,2,4-thiadiazin-3-amine-1,1-dioxide (0.50 g, 0.91 mmol), 2-fluoro-5-(trifluoromethyl)pyridine (1.51 g, 9.13 mmol), and THF (1 mL). The resulting mixture was stirred for 5 min, then a portion of the above base solution (0.54 M, 1.82 mmol, 3.4 mL) was added. After 1h, sodium hexamethyldisilazane (1.0 M in THF, 0.91 mL) was added. After an additional 3h, lithium hexamethyldisilazide (1.0 M in toluene, 1.4 mL) was added. After an additional hour, a second aliquot of lithium hexamethyldisilazide (1.0 M in toluene, 1.4 mL) was added. After one additional hour, the reaction was diluted with 10% w/v citric acid and EtOAc and stirred vigorously until both phases cleared. The phases were separated and the aqueous layer was extracted 2X with EtOAc. The organic portions were combined, washed with brine, dried over MgSO_4 , filtered and concentrated. This crude sample was subjected to column chromatography (120 g silica, 85 mL/min, 0% to 20% EtOAc/hexanes) to give 6(R)-[5-(trifluoromethyl)-2-pyridyl]-5R-(2,4,6-trifluorophenyl)-5,6-dihydro-N,N-bis[(4-methoxyphenyl)methyl]-2,5-dimethyl-2H-1,2,4-thiadiazin-3-amine-1,1-dioxide (180 mg, 28%). The above product of step 1 was treated according to Scheme 14, step 2 to give **Example 64**. LCMS data (Ex. 64): Obs. MH^+ : 453.0, Ret. Time: 3.19 min, LCMS method: B.

Scheme 16:

- 107 -



Step 1:

A solution of Ex. 8 (250 mg, 0.56 mmol) and potassium-2-propenetrifluoroborate (200 mg, 1.35 mmol) in EtOH (10 mL) in a pressure tube was degassed by bubbling N_2 through it for 10 min. To this solution was then added Et_3N (102 mg, 1.0 mmol) and $\text{Pd}(\text{dppf})\text{Cl}_2$ (19 mg, 0.023 mmol). The tube was sealed and heated to 100°C with stirring for 4.5 hours. The mixture was cooled to RT. To the mixture was added water and CH_2Cl_2 . The aqueous layer was extracted with CH_2Cl_2 (3x). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated to afford V.

10

Step 2:

A portion of the crude V was purified via prep TLC (SiO_2 ; 95:5:0.5 CH_2Cl_2 :MeOH: conc NH_4OH (aq.)) to afford Ex. 65 and Ex. 66. LCMS data (Ex. 65): Obs. MH^+ : 406.2, Ret. Time: 3.49 min, LCMS method: A. LCMS data (Ex. 66): Obs. MH^+ : 406.2, Ret. Time: 3.45 min, LCMS method:

15 A.

Step 3:

20

To a solution of V from step 1 (227 mg, 0.56 mmol) in CH_2Cl_2 was added Et_3N (68 mg, 0.67 mmol) and di-*tert*-butyldicarbonate (146 mg, 0.67 mmol). The resultant solution was stirred

- 108 -

at RT overnight. The solution was concentrated and partitioned between CH_2Cl_2 and water. The aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The crude material was purified via flash chromatography (SiO_2 ; gradient elution 100:0 to 75:25 hexanes:EtOAc) to afford the carbamate **W** (189 mg, 67% yield).

5


Step 4:

To a solution of **W** (30 mg, 0.06 mmol) in EtOH (2 mL) under an atmosphere of nitrogen was added Pd/C (10% Pd w/w, 5 mg, 0.003 mmol). The atmosphere was replaced with hydrogen and the mixture was stirred at RT under a hydrogen balloon for 3 hours. After that time, the mixture was filtered and the solvent was concentrated. The crude material was purified via preparative TLC (SiO_2 ; 75:25 hexanes:EtOAc) to afford a carbamate intermediate (21 mg). To a solution of the carbamate in CH_2Cl_2 (0.5 mL) was added TFA (0.5 mL). The resultant solution was stirred at RT for 30 min. The solution was concentrated. The residue was partitioned between CH_2Cl_2 and sat. Na_2CO_3 (aq.). The organic layer was separated, dried over Na_2SO_4 , filtered and concentrated. The crude product was purified via prep TLC (SiO_2 ; 95:5:0.5 CH_2Cl_2 :MeOH: conc NH_4OH (aq.)) to afford **Ex. 67** and **Ex. 68**. LCMS data (Ex. 67): Obs. MH^+ : 408.2, Ret. Time: 4.28 min, LCMS method: A. LCMS data (Ex. 68): Obs. MH^+ : 408.2, Ret. Time: 4.27 min, LCMS method: A.

10

15

Scheme 17:



20

Step 1:


A solution of **W** (Scheme 16) (67 mg) in CH_2Cl_2 was cooled to -78°C . To this solution was bubbled O_3 until the solution turned blue. After the color change, the solution was degassed by bubbling N_2 through it for 5 min. Excess Me_2S was added and the solution was warmed to RT with stirring overnight. After that time, the solution was concentrated and the crude residue was purified via prep TLC (SiO_2 ; 3:1 hexanes:EtOAc) to afford the ketone (32 mg).

25

Step 2:

- 109 -

To a solution of the ketone in CH_2Cl_2 (0.5 mL) was added TFA (0.5 mL). The resultant solution was stirred at RT for 30 min. The solution was concentrated. The residue was partitioned between CH_2Cl_2 and sat. Na_2CO_3 (aq.) and separated. The organic layer was separated, dried over Na_2SO_4 , filtered and concentrated. The crude product was purified via prep TLC [SiO_2 ; 95:5:0.5 CH_2Cl_2 :MeOH: conc NH_4OH (aq.)] to afford **Ex. 69** and **Ex. 70**. LCMS data (Ex. 69): Obs. MH^+ : 408.2, Ret. Time: 2.94 min, LCMS method: A. LCMS data (Ex. 70): Obs. MH^+ : 408.2, Ret. Time: 2.78 min, LCMS method: A.

Scheme 18:

10 Step 1:

To a solution of **Example 13** (213 mg, 0.58 mmol) in CH_2Cl_2 was added Et_3N (0.097 mL, 0.7 mmol) and di-*tert*-butyldicarbonate (445 mg, 2 mmol). The solution was stirred at RT overnight. After that time, the solution was concentrated. The crude residue was purified via flash chromatography (SiO_2 : gradient elution 100:0 to 75:25 hexanes:EtOAc) to afford the carbamate intermediate (237 mg, 88% yield).

15


Step 2:

The carbamate from step 1 (20 mg, 0.043 mmol) was dissolved in TFA (0.7 mL). To the solution was added H_2SO_4 (0.07 mL) and the resultant solution was cooled to 0°C . To this solution was added NBS (7.7 mg, 0.043 mmol) and the solution was stirred at 0°C for 45 min. After that time, additional NBS (7.7 mg) was added and the solution was stirred at 0°C for 20 min. At that time, additional NBS (5 mg) was added to the solution. The solution was stirred for an additional 30 min. To the solution was added sat. Na_2CO_3 (aq.) and Na_2SO_5 (s). The aqueous layer was extracted with CH_2Cl_2 (3x). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated. The crude residue was purified via prep TLC (SiO_2 ; 50:50:1 EtOAc:hexanes: Et_3N) to afford **Example 71** (6 mg). LCMS data (Ex. 71): Obs. MH^+ : 446.2, Ret. Time: 3.15 min, LCMS method: A.

25

Scheme 19:

- 110 -



Step 1:

To a solution of the iminothiadiazine dioxide (entry 2, table II) (377 mg, 1.3 mmol) in CH_2Cl_2 (15 mL) was added triethylamine (0.22 mL, 1.56 mmol) and di-*tert*-butyldicarbonate (340 mg, 1.56 mmol). The resultant solution was stirred at RT overnight. After that time, the solution was diluted with CH_2Cl_2 (50 mL) and washed with $\frac{1}{2}$ sat. NaHCO_3 (aq.). The aqueous layer was extracted with CH_2Cl_2 (3x). The combined organic layers were dried and concentrated. The crude residue was purified via flash chromatography (SiO_2 : gradient elution 100:0 to 70:30 heptane:EtOAc) to afford the carbamate (331 mg, 65%) as a white solid.

Step 2:

To a solution of the carbamate from step 1 (263 mg, 0.675 mmol) in THF (3.0 mL) at -78°C was added a solution of NaHMDS (1 M in THF, 1.49 mL, 1.49 mmol). The resultant solution was stirred at -78°C for 1 hour. After that time, a solution of benzyl bromide (0.5 M in THF, 1.35 mL, 0.675 mmol) was added. The resultant solution was stirred at -78°C for 30 min. To the solution was added sat. NH_4Cl (aq.). The mixture was warmed to RT and partitioned between water and EtOAc. The aqueous layer was extracted with EtOAc (3x). The combined organic layers were washed with brine, dried and concentrated. The crude residue was purified via flash chromatography (SiO_2 : gradient elution 100:0 to 75:25 heptane:EtOAc) to afford X (21 mg, 7%) and Y (48 mg, 15%).

Step 3 (for Example 72):

To a solution of X (49 mg, 0.103 mmol) in CH_2Cl_2 (4.0 mL) was added TFA (1.0 mL). The resultant solution was stirred at RT for 1.5 hours. After that time, the solvents were removed. The crude product was purified via preparative reverse phase HPLC [C_{18} : gradient elution 90:10

- 111 -

to 0:100 H₂O (w/ 0.025% HCl): MeCN] to afford **Ex. 72** (33 mg, 79%) as a white solid. LCMS data (Ex. 72): Obs. MH⁺: 380.0, Ret. Time: 2.25 min, LCMS method: D.

Step 3: (for Example 73):


- 5 **Example 73** was prepared using a procedure similar to that described for the preparation of **Ex. 72** except **Y** was used as the starting material. LCMS data (Ex. 73): Obs. MH⁺: 380.1, Ret. Time: 2.39 min, LCMS method: D.

- 10 **Example 74** was prepared using a procedure similar to that described for the preparation of **Ex. 72** with the intermediates indicated in Table X.

Table X			
Core	Alkyl halide	Ex.	
 Table X, entry 1		74	

LCMS data (Ex. 74): Obs. MH⁺: 344.1, Ret. Time: 2.33 min, LCMS method: B.

Scheme 20:



15 Step 1:

- 112 -

To a solution of 1-(4-fluoro-3-nitrophenyl)ethanone **AA** (20 g, 109 mmol) in DMF (160 mL) was added potassium carbonate (45.28 g, 327 mmol) and 4-methoxybenzylamine (PMBNH₂, 24.92 g, 240 mmol). The reaction was heated for 2 h, then filtered while hot. The filtrate was cooled to RT, diluted with EtOAc and washed with 1 M HCl (500 mL) upon which a yellow solid precipitated out. The solid was filtered off, dried under a stream of air and further dried under vacuum overnight to give intermediate **AB** (29.7 g, 100 mmol, 91.7%) which was used directly in the next step.

Step 2:

Intermediate **AB** (29.7 g, 100 mmol) was dissolved in THF/MeOH/water (600 mL/150 mL/60 mL), and zinc powder (65 g, 1 mol) and solid NH₄Cl (26.75 g, 500 mmol) were added to the mechanically stirred reaction. After heating to 85 C for 30 min, the reaction was filtered through celite, and the residue washed with MeOH. The combined filtrate was concentrated under reduced pressure, then diluted with EtOAc and water. The aqueous layer was extracted with EtOAc (2x) and the combined organic layers concentrated under reduced pressure to give intermediate **AC** as a green solid (22.2 g, 82 mmol, 82%) and used as is in the next step.

Step 3:

A solution of intermediate **AC** (22.2 g, 82 mmol) in propionic acid (300 mL) and 4 N HCl (80 mL) was heated to reflux for 4 h. The volatiles were removed under reduced pressure, and the resulting residue subjected to flash chromatography (SiO₂: gradient elution 50:50 to 30:70 hexanes:EtOAc, then 100% EtOAc) to afford benzimidazole **AD** (11.1 g, 36 mmol, 44%).

Step 4:

To a solution of benzimidazole **AD** (6.00 g, 20 mmol) in THF (50 mL) was added (R)-2-methylpropane-2-sulfinamide **AE** (3.63 g, 30 mmol) and Ti(OEt)₄ (6.13 mL, 30 mmol). After heating at 80 C overnight, the reaction was cooled to RT, then poured into ice water. The mixture was diluted with EtOAc and filtered over celite. The organic layer of the filtrate was washed with water, brine, then dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was subjected to flash chromatography (SiO₂: gradient elution 30:70 to 0:100 hexanes:EtOAc) to afford ketimine **AF** (2.3 g, 5.58 mmol, 28%) as a viscous liquid.

Step 5:

Intermediate **AG** was prepared as a mixture of diastereomers from ketimine **AF** (600 mg, 1.46 mmol) and N-methyl-1-(4-(trifluoromethyl)phenyl)methanesulfonamide (554 mg, 2.19 mmol, Scheme 4, step 1) using methods similar to that described in Scheme 4 steps 2 and 3.

- 113 -

Step 6:

Example 75 was prepared from intermediate **AG** using methods similar to that described in Scheme 1 steps 5 and 6. The desired diastereomer was obtained from RP-HPLC (C₁₈ radial compression, 35 mL/min, 10% to 95% MeCN/H₂O with 0.1% TFA) to give **Example 75** as TFA salt (1.9 mg). LCMS data (Ex. 75): Obs. MH⁺: 586.2, Ret. Time: 1.93 min, LCMS method: C.

LC/MS Conditions

Method A:

Column: Gemini C-18, 50 x 4.6 mm, 5 micron, obtained from Phenomenex.

Mobile phase: A: 0.05% Trifluoroacetic acid in water
B: 0.05% Trifluoroacetic acid in acetonitrile

Gradient: 90:10 to 5:95 (A:B) over 5 min.

Flow rate: 1.0 mL/min
UV detection: 254 nm

ESI-MS: Electro Spray Ionization Liquid chromatography-mass spectrometry (ESI-LC/MS) was performed on a PE SCIEX API-150EX, single quadrupole mass spectrometer.

Method B:

Column: Agilent Zorbax SB-C18 (3.0 x 50 mm) 1.8 μM

Mobile phase: A: 0.05% Trifluoroacetic acid in water
B: 0.05% Trifluoroacetic acid in acetonitrile

Gradient: 90:10 (A:B) for 0.3 min, 90:10 to 5:95 (A:B) over 5.1 min, 5:95 (A:B) for 1.2 min.

Flow rate: 1.0 mL/min
UV detection: 254 and 220 nm

Mass spectrometer: Agilent 6140 quadrupole.

Method C:

Column: Agilent Zorbax SB-C18 (3.0 x 50 mm) 1.8 μM

Mobile phase: A: 0.05% Trifluoroacetic acid in water
B: 0.05% Trifluoroacetic acid in acetonitrile

- 114 -

Gradient: 90:10 (A:B) for 0.3 min, 90:10 to 5:95 (A:B) over 1.2 min, 5:95 (A:B) for 1.2 min.

5 Flow rate: 1.0 mL/min
UV detection: 254 and 220 nm

Mass spectrometer: Agilent 6140 quadrupole.

10 **Method D:**

Column: Waters SunFire C-18 4.6mm x 50 mm

15 Mobile phase: A: 0.05% Trifluoroacetic acid in water
B: 0.05% Trifluoroacetic acid in acetonitrile

Gradient: 90:10 (A:B) for 1 min, 90:10 to 0:100 (A:B) over 4 min, 0:100 (A:B) for 2 min.

20 Flow rate: 1.0 mL/min
UV detection: 254 nm

Mass spectrometer: Finnigan LCQ Duo electrospray.

ASSAYS

25 The protocol that was used to determine the recited values is described as follows.

BACE1 HTRF FRET Assay

Reagents

Na⁺-Acetate pH 5.0

1% Brij-35

30 Glycerol

Dimethyl Sulfoxide (DMSO)

Recombinant human soluble BACE1 catalytic domain (>95% pure)

APP Swedish mutant peptide substrate (QSY7-APP^{swc}-Eu): QSY7-EISEVNLDAEFC-Europium-amide

35 A homogeneous time-resolved FRET assay was used to determine IC₅₀ values for inhibitors of the soluble human BACE1 catalytic domain. This assay monitored the increase of 620 nm fluorescence that resulted from BACE1 cleavage of an APPswedish APP^{swc} mutant peptide FRET substrate (QSY7-EISEVNLDAEFC-Europium-amide). This substrate contained an N-terminal QSY7 moiety that served as a quencher of the C-terminal Europium fluorophore

- 115 -

(620nm Em). In the absence of enzyme activity, 620 nm fluorescence was low in the assay and increased linearly over 3 hours in the presence of uninhibited BACE1 enzyme. Inhibition of BACE1 cleavage of the QSY7-APP^{swc}-Eu substrate by inhibitors was manifested as a suppression of 620 nm fluorescence.

5 Varying concentrations of inhibitors at 3x the final desired concentration in a volume of 10ul were preincubated with purified human BACE1 catalytic domain (3 nM in 10 μ l) for 30 minutes at 30° C in reaction buffer containing 20 mM Na-Acetate pH 5.0, 10% glycerol, 0.1% Brij-35 and 7.5% DMSO. Reactions were initiated by addition of 10 μ l of 600 nM QSY7-APP^{swc}-Eu substrate (200 nM final) to give a final reaction volume of 30 μ l in a 384 well Nunc
10 HTRF plate. The reactions were incubated at 30° C for 1.5 hours. The 620nm fluorescence was then read on a Rubystar HTRF plate reader (BMG Labtechnologies) using a 50 μ s delay followed by a 400 millisecond acquisition time window. Inhibitor IC₅₀ values were derived from non-linear regression analysis of concentration response curves. K_i values were then calculated from IC₅₀ values using the Cheng-Prusoff equation using a previously determined μ M value of 8 μ M
15 for the QSY7-APP^{swc}-Eu substrate at BACE1.

All of the example compounds of the invention exhibited K_i values of less than about 6 μ M and greater than about 1 nM in this assay. All of the example compounds of the invention except for examples 2, 73, and 74 exhibited K_i values of less than about 3 μ M in this assay. All of the example compounds of the invention except for examples 1, 2, 65, 72, 73, 74, and 75
20 exhibited K_i values of less than about 1 μ M in this assay. Some of the example compounds of the invention exhibited K_i values of less than about 300 nM in this assay; others less than about 200 nM in this assay; others less than about 100 nM in this assay; others less than about 50 nM in this assay; others less than about 10 nM in this assay; others less than about 5 nM in this assay. The compound of example 3 exhibited a K_i value of about 9 nM in this assay.

25 BACE-2

Inhibitor IC₅₀ values at purified human autoBACE-2 were determined in a time-resolved endpoint proteolysis assay that measures hydrolysis of the QSY7-EISEVNLD~~AE~~FC-Eu-amide FRET peptide substrate (BACE-HTRF assay). BACE-mediated hydrolysis of this peptide results in an increase in relative fluorescence (RFU) at 620 nm after excitation with 320 nm light.
30 Inhibitor compounds, prepared at 3x the desired final concentration in 1x BACE assay buffer (20 mM sodium acetate pH 5.0, 10% glycerol, 0.1% Brij-35) supplemented with 7.5% DMSO were pre-incubated with an equal volume of autoBACE-2 enzyme diluted in 1x BACE assay buffer

- 116 -

(final enzyme concentration 1 nM) in black 384-well NUNC plates for 30 minutes at 30°C. The assay was initiated by addition of an equal volume of the QSY7-EISEVNLDAEFC-Eu-amide substrate (200 nM final concentration, $K_m=8 \mu\text{M}$ for 4 μM for autoBACE-2) prepared in 1x BACE assay buffer supplemented with 7.5% DMSO and incubated for 90 minutes at 30°C.

5 DMSO was present at 5% final concentration in the assay. Following laser excitation of sample wells at 320 nm, the fluorescence signal at 620 nm was collected for 400 ms following a 50 μs delay on a RUBYstar HTRF plate reader (BMG Labtechnologies). Raw RFU data was normalized to maximum (1.0 nM BACE/DMSO) and minimum (no enzyme/DMSO) RFU values. IC_{50} values were determined by nonlinear regression analysis (sigmoidal dose response, variable
10 slope) of percent inhibition data with minimum and maximum values set to 0 and 100 percent respectively. Similar IC_{50} values were obtained when using raw RFU data. The K_i values were calculated from the IC_{50} using the Cheng-Prusoff equation.

All of the example compounds of the invention were assayed for BACE 2 except the following: Example Nos. 1,2,6,10,11,12,15,33,53,65,67,68,69,70,72,73, and 74. All of the
15 remaining example compounds of the invention were assayed for BACE-2 inhibition and exhibited K_i values of less than about 500 nM and greater than about 0.5 nM in this assay. All of these compounds except for example 71 exhibited K_i values of less than about 500 nM in this assay. All of these example compounds except for examples 32, 52, and 71 exhibited K_i values of less than about 200 nM in this assay. The compound of example 3 exhibited a K_i value of
20 about 11 nM in this assay.

Solution Stability

Substituted iminopyrimidinones are known in the art to be useful as aspartyl protease (e.g., BACE) inhibitors and for the treatment of Alzheimer's disease and other indications. *See*, e.g., Zhu, et al, PCT publication Nos. WO2005/058311, published June 30, 2005;
25 WO2006/065277, published June 22, 2006, and WO2008103351, published 28 August 2008. Applicants have found that the compounds of the invention exhibit properties that are both unexpected and advantageous for their use as BACE inhibitors and for the indications described herein. For instance, it has been found that the compounds of the invention, each of which contains an iminothiadiazine moiety according to Formula I, exhibit superior resistance to
30 hydrolysis, and hence improved solution stability, than is exhibited by compounds having an iminopyrimidinone moiety which are otherwise structurally identical.

- 117 -

The following procedures were used to measure solution stability, and to compare the solution stability of the compounds of the invention to that of otherwise structurally identical iminopyrimidinones. Results are reported as Example A below.

Stock solutions of the tested compounds were prepared by dissolving about 3 mg of each compound in 3 mL of acetonitrile. Standards for test compounds were prepared by diluting 1 mL of the stock solution with an additional 4 mL of acetonitrile. These standards were stored at 4°C. Samples were prepared by diluting 1 mL of the stock solution with 4 mL of 50 mM pH 7.4 phosphate buffer. These samples were stored at the appropriate temperature in the absence of light. Standards and samples were analyzed by LC/MS initially and at day 1, day 4, and day 6.

HPLC conditions:

Mobile phase A: 10 mM pH 5 ammonium acetate buffer : methanol (90 : 10)

Mobile phase B: 10 mM pH 5 ammonium acetate buffer : methanol (10 : 90)

Column: Zorbax SB-Phenyl 4.6 x 50 mm, 1.8 μm

Column temperature: 40°C

Flow: 0.8 mL/min.

Gradient:

Time (min.)	% B
0	40
9	100
11	100

Detectors: UV at 220 nm and 236 nm

MS, ES ionization, positive mode, for identification only at final time point.

The terms reported in the tables below have the following meanings:

Area % is the integration of peak from HPLC as reported by Waters Empower II software.

RRT is the relative retention time of new product compared to the standard of the test compound.

Formula for RRT is:

$$\frac{\text{Retention time of new product}}{\text{Retention time of standard}}$$

M + 1 is the mass observed including protonation (+ 1 mass unit).

ND stands for no peak detected by the UV detector.



* stands for no ion detected by the mass spectrometer.

Example A: Stability Studies Comparing Example 3 With Compound Y:

In the following study, the solution stability of the compound of Example 3 was measured and compared to that of Compound Y. The compound of Example 3 is an iminothiadiazine dioxide compound of the invention. Compound Y is the corresponding iminopyrimidinone compound, disclosed in WO2008/103351. The structures of the compound of Example 3 and of

- 118 -

Compound Y are shown in the table below. Studies were performed at 25°C, 40°C, and 60°C. The iminothiadiazine dioxide of Example 3 exhibited no hydrolysis and the iminopyrimidinone of Compound Y showed 3.36% hydrolysis at 25°C over 6 days. At 40°C, no hydrolysis of Example 3 was observed over 4 days while the hydrolysis product of Compound Y accounted for 31.4% of the sample. At 60°C, no hydrolysis was observed for Example 3 while the hydrolysis product of Compound Y accounted for 28.5% of the sample.

Compound of the Invention (iminothiadiazine dioxide)	Comparator Compound (iminopyrimidinone)
 <p style="text-align: center;">Example 3</p>	 <p style="text-align: center;">Compound Y (WO2008103351)</p>

Studies run at 25°C:

10 **Example 3:** free base M.W. = 433.1

	Peak Description	RRT	M + 1	Area %, Initial	Area %, Day 1	Area %, Day 4	Area %, Day 6
Standard	Example 3	1.00	434.1	100.0	100.0	100.0	100.0
Sample at pH 7.4	Example 3	1.00	434.1	100.0	100.0	100.0	100.0

Compound Y: free base M.W. = 397.2

	Peak Description	RRT	M + 1	Area %, Initial	Area %, Day 1	Area %, Day 4	Area %, Day 6
Standard	Compound Y	1.00	398.2	100.0	100.0	100.0	100.0
Sample at pH 7.4	Compound Y	1.00	398.2	100.0	100.0	100.0	89.79
	Hydrolysis product	0.63	416.2	ND	ND	ND	3.36

15 **Studies run at 40°C and 60°C:**

Example 3: free base M.W. = 433.1

	Peak Description	RRT	M + 1	Area %, Initial	Area %, Day 1	Area %, Day 4
Standard	Example 3	1.00	434.1	100.0	100.0	100.0
Sample initial	Example 3	1.00	434.1	100.0		
pH 7.4 at 40°C	Example 3	1.00	434.1		100.0	100.0
pH 7.4 at 60°C	Example 3	1.00	434.1			87.3
		0.52	*			12.7

- 119 -

Compound Y: free base M.W. = 397.2


	Peak Description	RRT	M + 1	Area %, Initial	Area %, Day 1	Area %, Day 4
Standard	Compound Y	1.00	398.2	98.8	99.4	100.0
Sample initial	Compound Y	1.00	398.2	98.2		
	Hydrolysis product	0.68	416.2	0.7		
pH 7.4 at 40°C	Compound Y	1.00	398.2		86.3	55.8
	Hydrolysis product	0.69	416.2		10.5	31.4
		0.87	*			5.8
pH 7.4 at 60°C	Compound Y	1.00	398.2		56.7	
	Hydrolysis product	0.68	416.2		28.5	
		0.87	*		6.6	

While the present invention has been described in view of the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

- 120 -

WE CLAIM:

1. A compound, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, said compound having the structural Formula (a):



(a)

5

wherein:

-L₁- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

-L₂- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

-L₃- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

each -L₄- is independently present or absent and when present independently represents a divalent moiety independently selected from the group consisting of -alkyl-, -haloalkyl-,

-heteroalkyl-, -alkenyl-, -alkynyl-, -N(R⁸)-, -NR⁸C(O)-, and -C(O)NR⁸-;

m, n, p and q are each independently selected integers, wherein:

m is 0 or more,

n is 0 or more,

p is 0 or more,

q is 0 or more,

20

wherein the maximum value of the sum of m and q is the maximum number of available substitutable hydrogen atoms on ring A,

- 121 -

wherein the maximum value of n is the maximum number of available substitutable hydrogen atoms on ring B, and

wherein the maximum value of p is the maximum number of available substitutable hydrogen atoms on ring C;

5 R^1 is selected from the group consisting of: H, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and heteroarylalkyl-,

wherein each of said alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and
10 heteroarylalkyl- of R^1 is unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups;

R^2 is selected from the group consisting of H, alkyl, halo, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and
15 heterocycloalkenyl,

wherein each of said alkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R^2 is unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups;

20 R^3 is selected from the group consisting of H, alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

wherein each of said alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R^3 is unsubstituted or substituted with from 1 to 5 independently selected R^{10} groups;

25 ring A is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R^4 (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -NR⁸C(O)N(R⁸)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁸)₂, -S(O)R⁷, -S(O)₂R⁷, -S(O)₂N(R⁸)₂, -OR⁷, -SR⁷, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, aryl, and heteroaryl,

- 122 -

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, aryl, and heteroaryl of R⁴ (when present) is optionally independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, -CN, -SF₅, -OSF₅, -NO₂, -N(R⁸)₂, -OR⁷, -C(O)N(R⁸)₂, and cycloalkyl;

ring B is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -NR⁸C(O)N(R⁸)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁸)₂, -S(O)R⁷, -S(O)₂R⁷, -S(O)₂N(R⁸)₂, -OR⁷, -SR⁷, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl of R⁵ (when present) is optionally independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, -CN, -SF₅, -OSF₅, -NO₂, -N(R⁸)₂, -OR⁷, -C(O)N(R⁸)₂, and cycloalkyl;

each ring C (when present) is independently selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R⁶ (when present) is independently selected from the group consisting of alkyl, aryl, arylalkyl-, haloalkyl, cycloalkyl, cycloalkylalkyl-, heteroaryl, and heteroarylalkyl-;

each R⁷ (when present) is independently selected from the group consisting of H, alkyl, alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

each R⁸ (when present) is independently selected from the group consisting of H, alkyl, alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

each R⁹ (when present) is independently selected from the group consisting of: halogen, -CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -NR⁸C(O)N(R⁸)₂, -NR⁸C(O)OR⁷, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁸)₂, -S(O)R⁷, -

- 123 -


$S(O)_2R^7$, $-S(O)_2N(R^8)_2$, $-OR^7$, $-SR^7$, alkyl, haloalkyl, heteroalkyl, alkenyl, alkynyl, aryl, arylalkyl, cycloalkyl, heteroaryl, heteroarylalkyl-, and heterocycloalkyl;

and

each R^{10} (when present) is independently selected from the group consisting of halo, $-CN$,
 5 $-NO_2$, $-Si(R^6)_3$, $-P(O)(OR^7)_2$, $-P(O)(OR^7)(R^7)$, $-N(R^8)_2$, $-NR^8C(O)R^7$, $-NR^8S(O)_2R^7$, $-NR^8C(O)N(R^8)_2$, $-NR^8C(O)OR^7$, $-C(O)R^7$, $-C(O)_2R^7$, $-C(O)N(R^8)_2$, $-S(O)R^7$, $-S(O)_2R^7$, $-S(O)_2N(R^8)_2$, $-OR^7$, $-SR^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
 10 heterocycloalkyl, aryl, and heteroaryl of R^{10} (when present) is optionally independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, $-CN$, $-NO_2$, $-N(R^8)_2$, $-OR^7$, $-C(O)N(R^8)_2$, and cycloalkyl.

15 2. A compound, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, said compound having the structural Formula (I):



wherein:

20 $-L_1-$ is present or absent and when present represents a divalent moiety selected from the group consisting of $-alkyl-$, $-haloalkyl-$, $-heteroalkyl-$, $-alkenyl-$, and $-alkynyl-$;

$-L_2-$ is present or absent and when present represents a divalent moiety selected from the group consisting of $-alkyl-$, $-haloalkyl-$, $-heteroalkyl-$, $-alkenyl-$, and $-alkynyl-$;

- 124 -

-L₃- is present or absent and when present represents a divalent moiety selected from the group consisting of -alkyl-, -haloalkyl-, -heteroalkyl-, -alkenyl-, and -alkynyl-;

m and n are each independently selected integers, wherein:

m is 0 or more,

5 n is 0 or more,

wherein the maximum value of m is the maximum number of available substitutable hydrogen atoms on ring A,

wherein the maximum value of n is the maximum number of available substitutable hydrogen atoms on ring B, and

10 R¹ is selected from the group consisting of: H, alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and heteroarylalkyl-,

wherein each of said alkyl, haloalkyl, heteroalkyl, heterohaloalkyl, cycloalkyl, cycloalkylalkyl-, heterocycloalkyl, heterocycloalkylalkyl-, aryl, arylalkyl-, heteroaryl, and
15 heteroarylalkyl- of R¹ is unsubstituted or substituted with from 1 to 5 independently selected R¹⁰ groups;

R² is selected from the group consisting of H, alkyl, halo, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

20 wherein each of said alkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl, cycloalkylalkyl-, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R² is unsubstituted or substituted with from 1 to 5 independently selected R¹⁰ groups;

R³ is selected from the group consisting of H, alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl,

25 wherein each of said alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycloalkyl, and heterocycloalkenyl of R³ is unsubstituted or substituted with from 1 to 5 independently selected R¹⁰ groups;

ring A is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl, monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic heterocycloalkenyl, and a multicyclic group;

each R⁴ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -OSF₅, -NO₂, -Si(R⁶)₃, -P(O)(OR⁷)₂, -P(O)(OR⁷)(R⁷), -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷,

- 125 -

$-\text{NR}^8\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$,
 $-\text{S}(\text{O})_2\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{SR}^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl,

5 wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl of R^4 (when present) is optionally independently
unsubstituted or further substituted with one or more independently selected groups
selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower
heteroalkyl, halo, $-\text{CN}$, $-\text{SF}_5$, $-\text{OSF}_5$, $-\text{NO}_2$, $-\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, and cycloalkyl;

10 ring B is selected from the group consisting of monocyclic aryl, monocyclic heteroaryl,
monocyclic cycloalkyl, monocyclic cycloalkenyl, monocyclic heterocycloalkyl, monocyclic
heterocycloalkenyl, and a multicyclic group;

each R^5 (when present) is independently selected from the group consisting of halo, $-\text{CN}$,
 $-\text{SF}_5$, $-\text{OSF}_5$, $-\text{NO}_2$, $-\text{Si}(\text{R}^6)_3$, $-\text{P}(\text{O})(\text{OR}^7)_2$, $-\text{P}(\text{O})(\text{OR}^7)(\text{R}^7)$, $-\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{R}^7$, $-\text{NR}^8\text{S}(\text{O})_2\text{R}^7$,
15 $-\text{NR}^8\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$,
 $-\text{S}(\text{O})_2\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{SR}^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl,
heterocycloalkyl, aryl, and heteroaryl of R^5 (when present) is optionally independently
20 unsubstituted or further substituted with one or more independently selected groups
selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower
heteroalkyl, halo, $-\text{CN}$, $-\text{SF}_5$, $-\text{OSF}_5$, $-\text{NO}_2$, $-\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, and cycloalkyl;

each R^6 (when present) is independently selected from the group consisting of alkyl, aryl,
arylalkyl-, haloalkyl, cycloalkyl, cycloalkylalkyl-, heteroaryl, and heteroarylalkyl-;

25 each R^7 (when present) is independently selected from the group consisting of H, alkyl,
alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl,
cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

each R^8 (when present) is independently selected from the group consisting of H, alkyl,
alkenyl, heteroalkyl, haloalkyl, aryl, arylalkyl-, heteroaryl, heteroarylalkyl-, cycloalkyl,
30 cycloalkylalkyl-, heterocycloalkyl, and heterocycloalkylalkyl-;

and

each R^{10} (when present) is independently selected from the group consisting of halo, $-\text{CN}$,
 $-\text{NO}_2$, $-\text{Si}(\text{R}^6)_3$, $-\text{P}(\text{O})(\text{OR}^7)_2$, $-\text{P}(\text{O})(\text{OR}^7)(\text{R}^7)$, $-\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{R}^7$, $-\text{NR}^8\text{S}(\text{O})_2\text{R}^7$, -

- 126 -

$\text{NR}^8\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{NR}^8\text{C}(\text{O})\text{OR}^7$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})_2\text{R}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, $-\text{S}(\text{O})\text{R}^7$, $-\text{S}(\text{O})_2\text{R}^7$, $-\text{S}(\text{O})_2\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{SR}^7$, alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl,

wherein each said alkyl, haloalkyl, haloalkoxy, heteroalkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl of R^{10} (when present) is optionally independently unsubstituted or further substituted with one or more independently selected groups selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, halo, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}(\text{R}^8)_2$, $-\text{OR}^7$, $-\text{C}(\text{O})\text{N}(\text{R}^8)_2$, and cycloalkyl.

10

3. A compound of claim 1, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, wherein:

R^1 is selected from the group consisting of H, lower alkyl, and cyclopropyl.

15

4. A compound of claim 1, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, wherein:

R^2 is H.

20

5. A compound of claim 1, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, wherein:


$-\text{L}_2-$ is absent or a $-\text{alkyl}-$ group; and

R^3 is selected from the group consisting H, alkyl, haloalkyl, heteroalkyl, cycloalkyl, and cycloalkylalkyl-.

25

6. A compound of claim 1, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, said compound having a structural Formula (II):

- 127 -



7. A compound of claim 6, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, wherein:

$-L_1-$ is absent or a divalent $-alkyl-$ group;

m is 0 or more and ring A is selected from the group consisting of phenyl, pyridyl, pyrazinyl, pyrazolyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, thiazolyl, and oxazolyl.

10 each R^4 (when present) is independently selected from the group consisting of halo, $-CN$, $-SF_5$, $-NO_2$, $-N(R^8)_2$, $-NR^8C(O)R^7$, $-C(O)N(R^8)_2$, $-OR^7$, alkyl, haloalkyl, heteroalkyl, and alkynyl;

$-L_3-$ is absent or a divalent $-CH_2-$ group;

15 n is 0 or more and ring B is selected from the group consisting of phenyl, pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridazinyl, thiazolyl, oxazolyl, benzofuranyl, benzimidazolyl, benzoxazolyl, quinoliny, isoquinoliny, naphthyl, benzothienyl, benzothiazolyl, indazolyl, indolyl, benzocyclobutanyl, and difluorodioxolanyl; and

20 each R^5 (when present) is independently selected from the group consisting of halo, $-CN$, $-SF_5$, $-N(R^8)_2$, $-NR^8C(O)R^7$, $-NR^8S(O)_2R^7$, $-C(O)R^7$, $-S(O)R^7$, $-S(O)_2R^7$, $-OR^7$, $-SR^7$, alkyl, haloalkyl, heteroalkyl, $-O-$ heteroalkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, and monocyclic heteroaryl,

wherein each said alkyl, said alkenyl, said alkenyl, said cycloalkyl, said heterocycloalkyl, said aryl, and said monocyclic heteroaryl of R^5 (when present) is optionally and independently further substituted with one or more groups


- 128 -


independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R⁸)₂, and -OH.

8. A compound of claim 6, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, wherein:
- 5 -L₁- is absent or a divalent -CH₂- group;
ring A is selected from the group consisting of phenyl and thienyl;
wherein, when ring A is phenyl, m is 0 to 5, and
when ring A is thienyl, m is 0 to 3;
- 10 each R⁴ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -NO₂, -N(R⁸)₂, -NR⁸C(O)R⁷, -C(O)N(R⁸)₂, -OR⁷, alkyl, haloalkyl, heteroalkyl, and alkynyl;
- L₃- is absent or a divalent -alkyl- group;
ring B is selected from the group consisting of phenyl, indazolyl, pyridyl, thienyl,
15 naphthyl, quinolinyl, isoquinolinyl, benzothienyl, benzocyclobutanyl, and difluorodioxolanyl;
n is 0 or more;
- each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, -N(R⁸)₂, -NR⁸C(O)R⁷, -NR⁸S(O)₂R⁷, -C(O)R⁷, -S(O)R⁷, -S(O)₂R⁷, -OR⁷, -SR⁷, alkyl, haloalkyl, heteroalkyl, -O-heteroalkyl, alkenyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl,
20 cyclohexyl, phenyl, pyridyl, furanyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl, oxetanyl, and pyrrolyl,
- wherein each said alkyl, -alkoxy, haloalkyl, heteroalkyl, -O-heteroalkyl, alkynyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, furanyl, thienyl, pyridazinyl, oxazolyl, isoxazolyl, oxetanyl, and pyrrolyl of R⁵ (when present) is
25 optionally and independently further substituted with one or more groups independently selected from the group consisting of halo, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, -CN, -SF₅, -NO₂, -N(R⁸)₂, and -OH;
and
each R⁷ and each R⁸ (when present) is independently selected from the group consisting
30 of H and lower alkyl.





9. A compound of claim 6, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, wherein:

-L₁- is absent or a divalent -CH₂- group;



the moiety, $(R^{11})_s$, is selected from the group consisting of Br-,

5



n is 0, 1, 2, or 3;

10 ring B is selected from the group consisting of phenyl, indazolyl, pyridyl, thienyl, naphthyl, quinolinyl, isoquinolinyl, benzothienyl, benzocyclobutanyl, and difluorodioxolanyl; and







each R⁵ (when present) is independently selected from the group consisting of halo, -CN, -SF₅, lower alkyl, lower alkenyl, lower haloalkyl, -C(O)-cyclopropyl, oxetanyl, lower alkyl-







substituted oxetanyl, cyclopropyl, lower heteroalkyl substituted cyclopropyl, lower alkyl-CN, lower heteroalkyl, and phenyl.

10. A compound, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, said compound being selected from the group consisting of:

Ex.	Structure
1	
2	
3	
4	
5	

Ex.	Structure
6	
7	
8	
9	

Ex.	Structure
10	
11	
12	
13	
14	
15	

Ex.	Structure
16	
17	
18	
19	
20	
21	

Ex.	Structure
22	
23	
24	
25	
26	
27	

Ex.	Structure
28	
29	
30	
31	
32	
33	

Ex.	Structure
34	
35	
36	
37	
38	

Ex.	Structure
39	
40	
41	
42	
43	
44	

Ex.	Structure
45	
46	
47	
48	
49	

Ex.	Structure
50	
51	
52	
53	
54	

Ex.	Structure
55	
56	
57	
58	
59	
60	

Ex.	Structure
61	
62	
63	
64	
65	
66	

- 136 -

Ex.	Structure	Ex.	Structure
67		72	
68		73	
69		74	
70		75	
71			

11. A pharmaceutical composition comprising at least one compound of any one of claims 1-10, or a tautomer or stereoisomer thereof, or a pharmaceutically acceptable salt of said compound, tautomer, or stereoisomer, and a pharmaceutically acceptable carrier or diluent.

12. A pharmaceutical composition comprising a compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, together with at least one additional therapeutic agent, and a pharmaceutically acceptable carrier or diluent.

- 137 -

13. A pharmaceutical composition of claim 12, wherein said at least one additional therapeutic agent is at least one agent selected from:

5 m_1 agonists; m_2 antagonists; cholinesterase inhibitors; galantamine; rivastigimine; N-methyl-D-aspartate receptor antagonists; combinations of cholinesterase inhibitors and N-methyl-D-aspartate receptor antagonists; gamma secretase modulators; gamma secretase inhibitors; non-steroidal anti-inflammatory agents; anti-inflammatory agents that can reduce neuroinflammation; anti-amyloid antibodies; vitamin E; nicotinic acetylcholine receptor agonists; CB1 receptor inverse agonists; CB1 receptor antagonists; antibiotics; growth hormone secretagogues; histamine
10 H3 antagonists; AMPA agonists; PDE4 inhibitors; GABA_A inverse agonists; inhibitors of amyloid aggregation; glycogen synthase kinase beta inhibitors; promoters of alpha secretase activity; PDE-10 inhibitors; Tau kinase inhibitors; Tau aggregation inhibitors; RAGE inhibitors; anti-Abeta vaccine; APP ligands; agents that upregulate insulin, cholesterol lowering agents; cholesterol absorption inhibitors; combinations of HMG-CoA reductase inhibitors and cholesterol
15 absorption inhibitors; fibrates; combinations of fibrates and cholesterol lowering agents and/or cholesterol absorption inhibitors; nicotinic receptor agonists; niacin; combinations of niacin and cholesterol absorption inhibitors and/or cholesterol lowering agents; LXR agonists; LRP mimics; H3 receptor antagonists; histone deacetylase inhibitors; hsp90 inhibitors; 5-HT4 agonists; 5-HT6 receptor antagonists; mGluR1 receptor modulators or antagonists; mGluR5 receptor modulators
20 or antagonists; mGluR2/3 antagonists; Prostaglandin EP2 receptor antagonists; PAI-1 inhibitors; agents that can induce Abeta efflux; Metal-protein attenuating compound; GPR3 modulators; and antihistamines.

14. A method of treating, preventing, and/or delaying the onset of an amyloid β pathology
25 ("A β pathology") and/or one or more symptoms of said pathology comprising administering at least one compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, to a patient in need thereof in an amount effective to treat said pathology.

15. A method of claim 14, wherein said A β pathology is selected from Alzheimer's Disease,
30 Down's syndrome, Parkinson's disease, memory loss, memory loss associated with Alzheimer's disease, memory loss associated with Parkinson's disease, attention deficit symptoms, attention deficit symptoms associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, dementia, stroke, microgliosis and brain inflammation, pre-senile dementia, senile

- 138 -

dementia, dementia associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, progressive supranuclear palsy, cortical basal degeneration, neurodegeneration, olfactory impairment, olfactory impairment associated with Alzheimer's disease, Parkinson's disease, and/or Down's syndrome, β -amyloid angiopathy, cerebral amyloid angiopathy, hereditary cerebral hemorrhage, mild cognitive impairment ("MCI"), glaucoma, amyloidosis, type II diabetes, diabetes-associated amyloidogenesis, hemodialysis complications (from β_2 microglobulins and complications arising therefrom in hemodialysis patients), scrapie, bovine spongiform encephalitis, traumatic brain injury ("TBI"), Creutzfeld-Jakob disease, and traumatic brain injury.

5
10

16. A method of claim 15, wherein said $A\beta$ pathology is Alzheimer's disease.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 10/51560

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(8) - A01N 43/54; A61K 31/505 (2010.01)
 USPC - 514/272
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 USPC: 514/272

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 USPC: 514/256-257; 514/259.5; 514/269; 514/274 (see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 WEST:PGPB, USPT, USOC, EPAB, JPAB
 Google: Scholar/Patents: bace1 inhibitors pentafluoro alzheimer's amyloid beta galantamine beta-secretase memantine asp2 inhibitors sulfur pentafluoride cyclic imine imino amide sulfonamide

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- Y	US 2007/0287692 A1 (WU et al.) 13 December 2007 (13.12.2007) para [0006]-[0009], [0018]-[0024], [0031], [0034], [0040]-[0042], [0048], [0098]-[0099], [0133], [0147], [0164], [0220]-[0228], [0239]	1-9, 11-16 ----- 10
Y	STACHEL et al. Structure-Based Design of Potent and Selective Cell-Permeable Inhibitors of Human Beta-Secretase (BACE-1). Journal of Medicinal Chemistry, 2004, Vol 47, pp 6447-6450; pg 6447, col 1, para 2; pg 6448, col 1, para 3; col 2, para 1-2; pg 6448, Fig 1; pg 6449, col 1, para 2-3; col 2, para 1	10

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 14 November 2010 (14.11.2010)	Date of mailing of the international search report 22 NOV 2010
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201	Authorized officer: Lee W. Young PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774