

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 February 2002 (28.02.2002)

PCT

(10) International Publication Number  
**WO 02/16363 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 487/04**,  
A61K 31/50, A61P 25/28

**STERNFELD, Francine** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).

(21) International Application Number: PCT/GB01/03166

(74) Agent: **HORGAN, James, Michael, Frederic**; European Patent Department, Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).

(22) International Filing Date: 13 July 2001 (13.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0017543.0 17 July 2000 (17.07.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **MERCK SHARP & DOHME LIMITED** [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CARLING, William, Robert** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **LADDUWAHETTY, Tamara** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **MACLEOD, Angus, Murray** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). **REEVE, Austin** [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: IMIDAZOLOPHTHALAZINE DERIVATIVES AS LIGANDS FOR GABA<sub>A</sub> RECEPTORS

(57) Abstract: A class of substituted imadazolo[2,1-a]phthalazine derivatives as ligands for GABAA receptors which are partial or full inverse agonists of an  $\alpha 5$  receptor subunit while being relatively free of activity at  $\alpha 1$  and/or  $\alpha 2$  and/or  $\alpha 3$  receptor subunit binding sites; they are therefore of benefit as a medicament for enhancing cognition but with the reduction or elimination of proconvulsant activity.



WO 02/16363 A1

**IMIDAZOLOPHTHALAZINE DERIVATIVES AS**  
**LIGANDS FOR GABA<sub>A</sub> RECEPTORS**

The present invention relates to a class of substituted  
5 imidazolophthalazine derivatives and to their use in therapy. More particularly, this invention is concerned with substituted imidazolo[2,1-a]phthalazine derivatives which are ligands for GABA<sub>A</sub> receptors containing the  $\alpha 5$  subunit and are therefore useful in therapy where cognition enhancement is required.

10 Receptors for the major inhibitory neurotransmitter, gamma-aminobutyric acid (GABA), are divided into two main classes: (1) GABA<sub>A</sub> receptors, which are members of the ligand-gated ion channel superfamily; and (2) GABA<sub>B</sub> receptors, which may be members of the G-protein linked receptor superfamily. Since the first cDNAs encoding individual GABA<sub>A</sub>  
15 receptor subunits were cloned the number of known members of the mammalian family has grown to thirteen (six  $\alpha$  subunits, three  $\beta$  subunits, three  $\gamma$  subunits and one  $\delta$  subunit). It may be that further subunits remain to be discovered; however, none has been reported since 1993.

Although knowledge of the diversity of the GABA<sub>A</sub> receptor gene  
20 family represents a huge step forward in our understanding of this ligand-gated ion channel, insight into the extent of subtype diversity is still at an early stage. It has been indicated that an  $\alpha$  subunit, a  $\beta$  subunit and a  $\gamma$  subunit constitute the minimum requirement for forming a fully functional GABA<sub>A</sub> receptor expressed by transiently transfecting cDNAs  
25 into cells. As indicated above, a  $\delta$  subunit also exists, but is apparently uncommon in the native receptor.

Studies of receptor size and visualisation by electron microscopy conclude that, like other members of the ligand-gated ion channel family, the native GABA<sub>A</sub> receptor exists in pentameric form. The selection of at  
30 least one  $\alpha$ , one  $\beta$  and one  $\gamma$  subunit from a repertoire of thirteen allows for the possible existence of more than 10,000 pentameric subunit

combinations. Moreover, this calculation overlooks the additional permutations that would be possible if the arrangement of subunits around the ion channel had no constraints (i.e. there could be 120 possible variants for a receptor composed of five different subunits).

5           Receptor subtype assemblies which do exist include  $\alpha 1\beta 2\gamma 2$ ,  $\alpha 2\beta 2/3\gamma 2$ ,  $\alpha 3\beta 2/3$ ,  $\alpha 2\beta 1$ ,  $\alpha 5\beta 3\gamma 2/3$ ,  $\alpha 6\beta 2$ ,  $\alpha 6\beta \delta$  and  $\alpha 4\beta \delta$ . Subtype assemblies containing an  $\alpha 1$  subunit are present in most areas of the brain and account for over 40% of GABA<sub>A</sub> receptors in the rat. Subtype assemblies containing  $\alpha 2$  and  $\alpha 3$  subunits respectively account for about  
10   25% and 17% of GABA<sub>A</sub> receptors in the rat. Subtype assemblies containing an  $\alpha 5$  subunit are primarily hippocampal and represent about 4% of receptors in the rat.

A characteristic property of some GABA<sub>A</sub> receptors is the presence of a number of modulatory sites, of which the most explored is the  
15   benzodiazepine (BZ) binding site through which anxiolytic drugs such as diazepam and temazepam exert their effect. Before the cloning of the GABA<sub>A</sub> receptor gene family, the benzodiazepine binding site was historically subdivided into two subtypes, BZ1 and BZ2, on the basis of radioligand binding studies. The BZ1 subtype has been shown to be  
20   pharmacologically equivalent to a GABA<sub>A</sub> receptor comprising the  $\alpha 1$  subunit in combination with  $\beta 2$  and  $\gamma 2$ . This is the most abundant GABA<sub>A</sub> receptor subtype, representing almost half of all GABA<sub>A</sub> receptors in the brain.

A number of dementing illnesses such as Alzheimer's disease are  
25   characterised by a progressive deterioration in cognition in the sufferer. It would clearly be desirable to enhance cognition in subjects desirous of such treatment, for example for subjects suffering from a dementing illness.


It has been reported by McNamara and Skelton in Psychobiology, 21:101-108, that the benzodiazepine receptor inverse agonist  $\beta$ -CCM  
30   enhanced spatial learning in the Morris watermaze. However,  $\beta$ -CCM and other conventional benzodiazepine receptor inverse agonists are

proconvulsant which makes it clear that they cannot be used as cognition enhancing agents in humans.

However, we have now discovered that it is possible to obtain medicaments which have cognition enhancing effects which may be employed with less risk of proconvulsant effects previously described with benzodiazepine receptor partial or full inverse agonists.

It has now been discovered that use of an  $\alpha 5$  receptor partial or full inverse agonist which is relatively free of activity at  $\alpha 1$  and/or  $\alpha 2$  and/or  $\alpha 3$  receptor binding sites can be used to provide a medicament which is useful for enhancing cognition but in which proconvulsant activity is reduced or eliminated. Inverse agonists at  $\alpha 5$  which are not free of activity at  $\alpha 1$  and/or  $\alpha 2$  and/or  $\alpha 3$  but which are functionally selective for  $\alpha 5$  can also be used. Inverse agonists which are both selective for  $\alpha 5$  and are relatively free of activity at  $\alpha 1$ ,  $\alpha 2$  and  $\alpha 3$  receptor binding sites are preferred.

The present invention provides a compound of formula I, or a pharmaceutically acceptable salt thereof:



(I)

wherein:

L is O, S or  $\text{NR}^n$  where  $\text{R}^n$  is H,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{3-6}$ cycloalkyl;

X is a 5-membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently chosen from oxygen, nitrogen and sulphur, at most one of the heteroatoms being oxygen or sulphur, or a 6-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms, the 5- or

6-membered heteroaromatic ring being optionally fused to a benzene or pyridine ring and the heteroaromatic ring being optionally substituted by  $R^x$  and/or  $R^y$  and/or  $R^z$ , where  $R^x$  is halogen,  $R^3$ ,  $OR^3$ ,  $OCOR^3$ ,  $NR^4R^5$ ,  $NR^4COR^5$ , OH, tri( $C_{1-6}$ alkyl)silyl $C_{1-6}$ alkoxy $C_{1-4}$ alkyl, CN or  $R^6$ ,  $R^y$  is  
5 halogen,  $R^3$ ,  $OR^3$ ,  $OCOR^3$ ,  $NR^4R^5$ ,  $NR^4COR^5$ ,  $NR^4R^5(C_{1-6})$ alkyl or CN and  $R^z$  is  $R^3$ ,  $OR^3$  or  $OCOR^3$ , providing that when X is a pyridine derivative, the pyridine ring is optionally in the form of the N-oxide, and providing that when X is a tetrazole derivative it is protected by a  $C_{1-4}$ alkyl group; or X is phenyl optionally substituted by one, two or three groups  
10 independently selected from halogen, cyano,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl and  $C_{3-6}$ cycloalkyl;

Y is optionally branched  $C_{1-4}$ alkylene optionally substituted by an oxo group or Y is a group  $(CH_2)_jO$  wherein the oxygen atom is nearest the group X and j is 2, 3 or 4;

15 Z represents a phenyl ring, a 5-membered heteroaromatic ring containing 1, 2 or 3 heteroatoms independently selected from oxygen, nitrogen and sulphur, at most one of the heteroatoms being oxygen or sulphur, or a 6-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms, each of which rings may be optionally substituted with  
20 one or more substituents selected from halogen,  $R^3$ ,  $OR^3$ ,  $OCOR^3$ ,  $NR^4R^5$ ,  $NR^4R^5(C_{1-6})$ alkyl,  $NR^4R^5CO$ ,  $NR^4R^5CO(C_{1-6})$ alkyl, CN, cyano( $C_{1-6})$ alkyl or  $R^6$ ;

$R^3$  represents  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-6}$ cycloalkyl,  $C_{3-6}$ cycloalkyl( $C_{1-6})$ alkyl, cyano( $C_{1-6})$ alkyl, hydroxy $C_{1-6}$ alkyl and  $R^3$  is  
25 optionally mono, di- or tri-fluorinated;

$R^4$  and  $R^5$  are each independently hydrogen,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-6}$ cycloalkyl or  $CF_3$  or  $R^4$  and  $R^5$ , together with a nitrogen atom to which they are commonly attached, form a 4-7 membered heteroaliphatic ring containing the said nitrogen atom and optionally one  
30 other heteroatom selected from O, N and S; and

R<sup>6</sup> is C<sub>6-10</sub>aryl, C<sub>6-10</sub>aryl (C<sub>1-6</sub>)alkyl, heteroaryl or heteroaryl(C<sub>1-6</sub>)alkyl, where heteroaryl is a five-membered aromatic ring containing one, two, three or four nitrogen atoms or one oxygen atom or sulphur atom and optionally one or two nitrogen atoms or a six-membered aromatic ring containing one, two or three nitrogen atoms, and R<sup>6</sup> is optionally substituted by one, two or three substituents independently chosen from halogen atoms and C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>2-4</sub>alkenyloxy and C<sub>2-4</sub>alkynyloxy groups each of which groups is unsubstituted or substituted by one, two or three halogen atoms.

As used herein, the expression "C<sub>1-6</sub>alkyl" includes methyl and ethyl groups, and straight-chained and branched propyl, butyl, pentyl and hexyl groups. Particular alkyl groups are methyl, ethyl, n-propyl, isopropyl and t-butyl. Derived expressions such as "C<sub>1-4</sub>alkyl", "C<sub>2-4</sub>alkenyl", "C<sub>2-6</sub>alkenyl", "hydroxyC<sub>1-6</sub>alkyl", "C<sub>2-4</sub>alkyl" and "C<sub>2-6</sub>alkynyl" are to be construed in an analogous manner.

The expression "C<sub>3-6</sub>cycloalkyl" as used herein includes cyclic propyl, butyl, pentyl and hexyl groups such as cyclopropyl and cyclohexyl.

The expression "C<sub>3-6</sub> cycloalkyl(C<sub>1-6</sub>)alkyl" as used herein includes cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

Typical C<sub>4-7</sub> cycloalkenyl groups include cyclobutenyl, cyclopentenyl and cyclohexenyl.

The expression "C<sub>1-4</sub>alkylene" as used herein refers to alkanediyl groups of up to 4 carbon atoms in which the unsatisfied valencies reside on the same carbon atom or on different carbon atoms.

Typical C<sub>6-8</sub> bicycloalkyl groups include bicyclo[2.1.1]hexyl and bicyclo[2.2.1]heptyl.

Typical aryl groups include phenyl and naphthyl, preferably phenyl.

The expression "aryl(C<sub>1-6</sub>)alkyl" as used herein includes benzyl, phenylethyl, phenylpropyl and naphthylmethyl.

Suitable heterocycloalkyl groups include azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl and thiomorpholinyl groups.

Unless otherwise specified, 5- and 6-membered heteroaromatic rings shall include pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, furyl, 5 thienyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, oxadiazolyl, triazolyl and thiadiazolyl groups and benzo-fused analogues thereof. A suitable 5-membered heteroaromatic ring containing four nitrogen atoms is tetrazolyl. Suitable 6-membered heteroaromatic rings containing three nitrogen atoms include 1,2,4-triazine and 1,3,5-triazine. 10 When a heteroaromatic ring comprises a hydroxy group as a substituent, and keto-enol tautomerism is possible, both tautomers are included within the scope of the invention. Thus, for example, a 3-hydroxy-1,2,4-triazole ring will be considered equivalent to the 3-keto tautomer.

The expression "heteroaryl(C<sub>1-6</sub>)alkyl" as used herein includes 15 furylmethyl, furylethyl, thienylmethyl, thienylethyl, pyrazolylmethyl, oxazolylmethyl, oxazolylethyl, isoxazolylmethyl, thiazolylmethyl, thiazolylethyl, imidazolylmethyl, imidazolylethyl, benzimidazolylmethyl, oxadiazolylmethyl, oxadiazolylethyl, thiadiazolylmethyl, thiadiazolylethyl, triazolylmethyl, triazolylethyl, tetrazolylmethyl, tetrazolylethyl, 20 pyridinylmethyl, pyridinylethyl, pyridazinylmethyl, pyrimidinylmethyl, pyrazinylmethyl, quinolinylmethyl, isoquinolinylmethyl and quinoxalinylmethyl.

The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, of which fluorine and chlorine are preferred.

25 As used herein the term "C<sub>1-6</sub>alkoxy" includes methoxy and ethoxy groups, and straight-chained, branched and cyclic propoxy, butoxy, pentoxy and hexoxy groups, including cyclopropylmethoxy. Derived expressions such as "C<sub>2-6</sub>alkenyloxy", "C<sub>2-6</sub>alkynyloxy", "C<sub>1-4</sub>alkoxy", "C<sub>2-4</sub>alkenyloxy" and "C<sub>2-4</sub>alkyloxy" should be construed in an analogous 30 manner.

For use in medicine, the salts of the compounds of formula I will be pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts.

Where the compounds according to the invention have at least one asymmetric centre, they may accordingly exist as enantiomers. Where the compounds according to the invention possess two or more asymmetric centres, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

Preferably L is an oxygen atom. L may also be S or  $\text{NR}^n$  in which  $\text{R}^n$  is preferably hydrogen or methyl.  $\text{R}^n$  may be hydrogen.

X is generally: pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl optionally substituted by a halogen atom or a group  $\text{R}^3$ ,  $\text{OR}^3$ ,  $\text{NR}^4\text{R}^5$  or a five membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms, and X is optionally fused to a benzene ring; a 5-membered heteroaromatic ring containing 2 or 3 heteroatoms chosen from oxygen, sulphur and nitrogen, at most one of the heteroatoms being oxygen or sulphur, which is unsubstituted or substituted by one, two or three groups independently chosen from halogen and  $\text{R}^3$ , or which is substituted by a pyridyl, phenyl

or benzyl ring which ring is optionally independently substituted by one, two or three halogen atoms or C<sub>1-6</sub>alkyl or CF<sub>3</sub> groups; or phenyl optionally substituted by one, two or three independently chosen halogen atoms.

Alternatively, X may represent a 5-membered heteroaromatic ring  
5 containing 2 or 3 heteroatoms chosen from oxygen, sulphur and nitrogen, at most one of the heteroatoms being oxygen or sulphur, which is fused to a benzene or pyridine ring, or which is substituted by NH<sub>2</sub> or by OH which may exist as the keto tautomer; or X may represent a tetrazole ring bearing a C<sub>1-4</sub>alkyl substituent.

10 When X is a substituted 6-membered heteroaromatic ring:

R<sup>x</sup> is preferably halogen, R<sup>3</sup>, OR<sup>3</sup>, NR<sup>4</sup>R<sup>5</sup> or a five-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms and more preferably methyl, CF<sub>3</sub>, methoxy, bromine, chlorine, isopropoxy, dimethylamino or a five-membered heterocyclic ring containing 1, 2 or 3  
15 nitrogen atoms; and R<sup>y</sup> and R<sup>z</sup> are preferably absent.

When X is a substituted 5-membered heteroaromatic ring:

R<sup>x</sup> is preferably halogen, R<sup>3</sup> or a pyridyl, phenyl or benzyl ring, which ring is optionally independently substituted by one, two or three halogen atoms or C<sub>1-6</sub>alkyl or CF<sub>3</sub> groups, and more preferably R<sup>x</sup> is methyl, CF<sub>3</sub>, chlorine  
20 or a phenyl, pyridyl or benzyl ring which ring is unsubstituted or substituted by chlorine or CF<sub>3</sub>; and R<sup>y</sup> and R<sup>z</sup> are preferably halogen or R<sup>3</sup> or are absent, and more preferably are methyl, CF<sub>3</sub> or chlorine, or are absent.

Specific values of X are 2-pyridyl, 1-methyl-1,2,4-triazol-3-yl, 2-  
25 methyl-1,2,4-triazol-3-yl, 1-methyl-1,2,3-triazol-4-yl, 5-methylisoxazol-3-yl, 1,2,4-oxadiazol-3-yl, and 2-methylthiazol-4-yl, particularly 1-methyl-1,2,3-triazol-4-yl.

Apt values for Y include CH<sub>2</sub>, CH(CH<sub>3</sub>), CH<sub>2</sub>CH<sub>2</sub> and CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub> optionally substituted by an oxo group, and CH<sub>2</sub>CH<sub>2</sub>O and CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O.  
30 For example, Y can be CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>O or CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O. Preferably Y is CH<sub>2</sub> or CH<sub>2</sub>CH<sub>2</sub> and most preferably CH<sub>2</sub>.

From the foregoing it will be understood that particularly suitable groups L-Y-X are OCH<sub>2</sub>X groups where X is pyridyl or triazolyl, particularly 1,2,4-triazol-3-yl substituted with methyl in the 1- or 2-position.

5            Suitable values for Z include optionally substituted pyrimidinyl, triazinyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl and thiadiazolyl groups.

             Z is very aptly an optionally substituted 5-membered heteroaromatic ring containing one oxygen and one or two nitrogen ring  
10       atoms.

             Favoured values for Z include optionally substituted isoxazoles and oxadiazoles, particularly isoxazoles.


             Typical substituents on Z include C<sub>1-6</sub>alkyl, C<sub>3-6</sub>cycloalkyl, hydroxyC<sub>1-6</sub>alkyl, pyridyl, thienyl or amino, particularly methyl, ethyl,  
15       isopropyl, cyclopropyl, thienyl or pyridyl.

             Z may be unsubstituted.

             Z may very aptly be substituted by methyl.

             Particular values of Z include 5-methylisoxazol-3-yl.

             A particular sub-class of compounds according to the invention is  
20       represented by the compounds of formula IIA, and pharmaceutically acceptable salts thereof:



(IIA)

25       wherein

X' represents phenyl, pyrazolyl, isoxazolyl, thiazolyl, imidazolyl, thiadiazolyl, benzimidazolyl, oxadiazolyl, triazolyl, tetrazolyl, pyridinyl, pyridazinyl, oxazolopyridinyl, pyrimidinyl, pyrazinyl, quinolinyl, isoquinolinyl or quinoxalinyl, any of which groups may be optionally substituted by one or more of C<sub>1-6</sub> alkyl, amino, pyridyl, CF<sub>3</sub>, aryl(C<sub>1-6</sub>)alkyl, pyridyl(C<sub>1-6</sub>)alkyl, halogen, cyano, cyano(C<sub>1-6</sub>)alkyl, hydroxymethyl, C<sub>1-6</sub>alkoxy, C<sub>3-6</sub>cycloalkyl(C<sub>1-6</sub>)alkoxy, hydroxy or the keto tautomer thereof, di(C<sub>1-6</sub>)alkylamino(C<sub>1-6</sub>)alkyl, amino(C<sub>1-6</sub>)alkyl, N-(C<sub>1-6</sub>)alkylpiperidinyl, pyrrolidinyl(C<sub>1-6</sub>)alkyl, piperazinyl(C<sub>1-6</sub>)alkyl and morpholinyl(C<sub>1-6</sub>)alkyl; and

Z' represents a 5-membered heteroaromatic ring containing one oxygen and one or two nitrogen ring atoms which is optionally substituted with C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkoxy.

In a subset of the compound of Formula IIA, X' represents phenyl, pyrazolyl, isoxazolyl, thiazolyl, imidazolyl, benzimidazolyl, triazolyl, tetrazolyl, pyridinyl, pyridazinyl, oxazolopyridinyl, pyrimidinyl, pyrazinyl, quinolinyl, isoquinolinyl or quinoxalinyl, any of which groups may be optionally substituted by one or more of C<sub>1-6</sub> alkyl, CF<sub>3</sub>, aryl(C<sub>1-6</sub>)alkyl, pyridyl(C<sub>1-6</sub>)alkyl, halogen, cyano, cyano(C<sub>1-6</sub>)alkyl, hydroxymethyl, C<sub>1-6</sub>alkoxy, C<sub>3-6</sub>cycloalkyl(C<sub>1-6</sub>)alkoxy, hydroxy or the keto tautomer thereof, di(C<sub>1-6</sub>)alkylamino(C<sub>1-6</sub>)alkyl, amino(C<sub>1-6</sub>)alkyl, N-(C<sub>1-6</sub>)alkylpiperidinyl, pyrrolidinyl(C<sub>1-6</sub>)alkyl, piperazinyl(C<sub>1-6</sub>)alkyl and morpholinyl(C<sub>1-6</sub>)alkyl.

Illustrative values of specific substituents on the group X' include methyl, ethyl, *n*-propyl, isopropyl, benzyl, pyridinylmethyl, chloro, cyano, cyanomethyl, hydroxymethyl, ethoxy, cyclopropylmethoxy, dimethylaminomethyl, aminoethyl, dimethylaminoethyl, N-methylpiperidinyl, pyrrolidinyethyl, piperazinylethyl and morpholinylmethyl.

Further illustrative values of specific substituents on the group X' include trifluoromethyl, amino, pyridyl and hydroxy or its keto tautomer.

Selected substituents for the group X' include methyl, ethyl,

*n*-propyl, and isopropyl, especially methyl.

Further selected substituents for the group X' include trifluoromethyl, amino, pyridyl and hydroxy or its keto tautomer.

Specific values of X' include pyridyl, methyl-triazolyl, ethyl-  
5 triazolyl, propyl-triazolyl and isopropyl-triazolyl.

Further specific values of X' include 1,2,4-oxadiazolyl, methyl-thiazolyl, 5-hydroxy-1,2,4-triazolyl (which is equivalent to 5-oxo-1H, 4H-1,2,4-triazolyl), trifluoromethylpyridyl, methyl-imidazolyl, thiazolyl, amino-thiazolyl, amino-1,2,4-thiadiazolyl, pyridazinyl, pyrazinyl,  
10 quinoxalinyl, pyrimidinyl, methyl-tetrazolyl, pyridyl-triazolyl, oxazolopyridinyl and methyl-pyrazolyl.

A favoured value of X' is methyltriazolyl, especially 1-methyl-1,2,3-triazol-4-yl.

A particular compound within the scope of the present invention is:  
15 3-(5-methylisoxazol-3-yl)-6-[(1-methyl-1,2,3-triazol-4-yl)methoxy]imidazo[2,1-a]phthalazine  
and pharmaceutically acceptable salts thereof.

The compounds of the present invention have a good binding affinity ( $K_i$ ) for the  $\alpha 5$  subunit of the GABA<sub>A</sub> receptor. In a preferred  
20 embodiment the compounds of the invention are binding selective for the  $\alpha 5$  subunit relative to the  $\alpha 1$ ,  $\alpha 2$  and  $\alpha 3$  subunits. In another preferred embodiment the compounds are functionally selective for the  $\alpha 5$  subunit as partial or full inverse agonists whilst substantially being antagonists at the  $\alpha 1$ ,  $\alpha 2$  and  $\alpha 3$  subunits.

25 The invention also provides pharmaceutical compositions comprising one or more compounds of this invention and a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid  
30 sprays, drops, ampoules, transdermal patches, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal

administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, 5 stearic acid, magnesium stearate, dicalcium phosphate or gums or surfactants such as sorbitan monooleate, polyethylene glycol, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When 10 referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage 15 forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the 20 advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the 25 duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The present invention also provides a compound of the invention for 30 use in a method of treatment of the human body. Preferably the treatment is for a condition associated with GABA<sub>A</sub> receptors comprising

the  $\alpha 5$  subunit and/or for the enhancement of cognition. Preferably the condition is a neurological deficit with an associated cognitive disorder such as a dementing illness such as Alzheimer's disease. Other conditions to be treated include cognition deficits due to traumatic injury, stroke,  
5 Parkinson's disease, Downs syndrome, age related memory deficits, attention deficit disorder and the like.

The present invention further provides the use of a compound of the present invention in the manufacture of a medicament for the enhancement of cognition, preferably in a human suffering from a  
10 dementing illness such as Alzheimer's disease.

Also disclosed is a method of treatment of a subject suffering from a cognition deficit, such as that resulting from a dementing illness such as Alzheimer's disease, which comprises administering to that subject an effective amount of a compound according to the present invention.



15 The compounds of the present invention can be used in a variety of disorders of the central nervous system. Such disorders include delirium, dementia and amnestic and other cognitive disorders. Examples of delirium are delirium due to substance intoxication or substance withdrawal, delirium due to multiple etiologies and delirium NOS (not  
20 otherwise specified). Examples of dementia are: dementia of the Alzheimer's type with early onset which can be uncomplicated or with delirium, delusions or depressed mood; dementia of the Alzheimer's type, with late onset, which can be uncomplicated or with delirium, delusions or depressed mood; vascular dementia which can be uncomplicated or with  
25 delirium, delusions or depressed mood; dementia due to HIV disease; dementia due to head trauma; dementia due to Parkinson's disease; dementia due to Huntington's disease; dementia due to Pick's disease; dementia due to Creutzfeld-Jakob disease; dementia which is substance-induced persisting or due to multiple etiologies; and dementia NOS.  
30 Examples of amnestic disorders are amnestic disorder due to a particular

medical condition or which is substance-induced persisting or which is amnesic disorder NOS.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

For the enhancement of cognition, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.01 to 100 mg/kg per day, and especially about 0.01 to 5 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day. In some cases, however, dosage outside these limits may be used.


The present invention also provides a process for producing a compound of formula I which comprises reacting a compound of formula II with a compound of formula III:


$$(II)$$


(III)

wherein X, Y and Z are as defined above, generally in the presence of a base such as sodium hydride optionally followed by an alkali such as potassium carbonate, and in a solvent such as dimethylformamide.

The compound of formula II can be prepared from a compound of formula IV:



5


wherein Z is as defined above by reaction with

(1) a diazotising reagent such as trimethylsilyldiazomethane generally in a solvent such as diethyl ether, to give the diazomethyl derivative;

10 (2) then reacting with an acid such as trifluoroacetic acid generally in a solvent such as diethylether to give the hydroxy derivative;

(3) and finally reacting with 4-amino-1-phthalazine (Angew. (1967) 6(2) 173-4) generally in the presence of an acid such as trifluoroacetic acid and a solvent such as toluene.

15 The compound of formula III can be made by reacting a compound of formula V with a chlorinating agent such as thionyl chloride:



20 wherein X and Y are as defined above.

Where they are not commercially available, the starting materials of formulae III and V may be prepared by methods analogous to those described in the accompanying Example, or by standard methods well known from the art.

25 It will be understood that any compound of formula I initially obtained from any of the above processes may, where appropriate,

subsequently be elaborated into a further compound of formula I by techniques known from the art.

It will also be appreciated that where more than one isomer can be obtained from a reaction then the resulting mixture of isomers can be  
5 separated by conventional means.

Where the above-described process for the preparation of the compounds according to the invention gives rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The novel compounds may be  
10 prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques such as preparative HPLC, or the formation of diastereomeric pairs by salt formation with an optically active acid, such  
15 as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

20 During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene &  
25 P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The compounds in accordance with this invention potently inhibit the binding of [3H]-flumazenil to the benzodiazepine binding site of  
30 human GABA<sub>A</sub> receptors containing the  $\alpha 5$  subunit stably expressed in Ltk- cells.

### Reagents

- Phosphate buffered saline (PBS).
- Assay buffer: 10 mM  $\text{KH}_2\text{PO}_4$ , 100 mM KCl, pH 7.4 at room temperature.
- 5 • [3H]-Flumazenil (18 nM for  $\alpha 1\beta 3\gamma 2$  cells; 18 nM for  $\alpha 2\beta 3\gamma 2$  cells; 10 nM for  $\alpha 3\beta 3\gamma 2$  cells; 10 nM for  $\alpha 5\beta 3\gamma 2$  cells) in assay buffer.
- Flunitrazepam 100  $\mu\text{M}$  in assay buffer.
- Cells resuspended in assay buffer (1 tray to 10 ml).

### 10 Harvesting Cells

Supernatant is removed from cells. PBS (approximately 20 ml) is added. The cells are scraped and placed in a 50 ml centrifuge tube. The procedure is repeated with a further 10 ml of PBS to ensure that most of the cells are removed. The cells are pelleted by centrifuging for 20 min at  
15 3000 rpm in a benchtop centrifuge, and then frozen if desired. The pellets are resuspended in 10 ml of buffer per tray (25 cm x 25 cm) of cells.

### Assay

Can be carried out in deep 96-well plates or in tubes. Each tube  
20 contains:

- 300  $\mu\text{l}$  of assay buffer.
- 50  $\mu\text{l}$  of [3H]-flumazenil (final concentration for  $\alpha 1\beta 3\gamma 2$ : 1.8 nM; for  $\alpha 2\beta 3\gamma 2$ : 1.8 nM; for  $\alpha 3\beta 3\gamma 2$ : 1.0 nM; for  $\alpha 5\beta 3\gamma 2$ : 1.0 nM).
- 50  $\mu\text{l}$  of buffer or solvent carrier (e.g. 10% DMSO) if compounds are  
25 dissolved in 10% DMSO (total); test compound or flunitrazepam (to determine non-specific binding), 10  $\mu\text{M}$  final concentration.
- 100  $\mu\text{l}$  of cells.

Assays are incubated for 1 hour at 40°C, then filtered using either a Tomtec or Brandel cell harvester onto GF/B filters followed by 3 x 3 ml  
30 washes with ice cold assay buffer. Filters are dried and counted by liquid scintillation counting. Expected values for total binding are 3000-4000

dpm for total counts and less than 200 dpm for non-specific binding if using liquid scintillation counting, or 1500-2000 dpm for total counts and less than 200 dpm for non-specific binding if counting with meltilex solid scintillant. Binding parameters are determined by non-linear least squares regression analysis, from which the inhibition constant  $K_i$  can be calculated for each test compound.

The compounds of the accompanying Examples were tested in the above assay, and all were found to possess a  $K_i$  value for displacement of [3H]Ro 15-1788 from the  $\alpha 5$  subunit of the human GABA<sub>A</sub> receptor of 100 nM or less, most were at 50 nM or less, many were at 10 nM or less and some were at 1 nM or less.

The compounds of the present invention have been shown to enhance cognition in the rat water maze test (Morris, Learning and Motivation, 1981, 12, 239ff). Further details of methodology for demonstrating that the present compounds enhance cognition can be found in WO-A-9625948.

The following Example illustrates the present invention:

### EXAMPLE 1

3-(5-Methylisoxazol-3-yl)-6-[(1-methyl-1,2,3-triazol-4-yl)methoxy]imidazo[2,1-a]phthalazine

#### a) 2-Diazo-1-(5-methylisoxazol-3-yl)ethanone

5-Methylisoxazole-3-carboxylic acid chloride (15.7g, 0.108 mol) was dissolved in diethyl ether (100 ml) and added dropwise to a cooled solution of trimethylsilyldiazomethane (2M in hexane, 56.3 ml, 0.112 mol). The solution was stirred at 0°C for 1 hour, then allowed to warm up to room temperature and stirred for 1 hour. Acetic acid (1.5 ml) was added and the solution was stirred for 15 min. The solution was then washed with water and brine, and dried ( $\text{Na}_2\text{SO}_4$ ). The solution was evaporated to dryness

and the residue purified by chromatography on silica gel, eluting with ethyl acetate/hexane (1:2), to yield the required diazoketone (7.4 g, 45%).  
<sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>) 2.48 (3H, s), 6.19 (1H, s), 6.42 (1H, s).

5    b) 2-Hydroxy-1-(5-methylisoxazol-3-yl)ethanone

2-Diazo-1-(5-methylisoxazol-3-yl)ethanone (7.0 g, 46 mmol) was dissolved in diethyl ether (100 ml) and trifluoroacetic acid (25 ml) and water (100 ml) were added. The solution was stirred at room temperature for 2 hours, then the diethyl ether was removed *in vacuo* and the aqueous  
10    solution was stirred at room temperature for 18 hours. Further water (100 ml) was added and the aqueous solution was extracted twice with ethyl acetate. The organic extracts were washed successively with water, 10% aqueous potassium carbonate solution and brine, and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent yielded the title-compound as a pale solid  
15    (4.0 g, 61%), <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>) 2.52 (3H, s), 4.90 (2H, s), 6.44 (1H, s).

c) 3-(5-Methylisoxazol-3-yl)imidazo[2,1-a]phthalazin-6-one

4-Amino-1-phthalazinone (2.28 g, 14.2 mmol) [Angew. (1967) 6(2)  
20    173-4] and 2-hydroxy-1-(5-methylisoxazol-3-yl)ethanone (2.0 g, 14.2 mmol) were suspended in toluene (100 ml) and trifluoroacetic acid (5 ml) was added. The suspension was heated at reflux under Dean & Stark conditions, for 24 hours. The solvent was then removed *in vacuo* and the residue was purified by chromatography on silica gel eluting with 10%  
25    methanol:dichloromethane, to provide the title product (1.62 g, 43%),  
<sup>1</sup>H NMR (360 MHz, d<sup>6</sup>-DMSO) 2.50 (3H, s), 5.96 (2H, br s), 7.79 – 8.22 (4H, m) 11.48 (1H, s).

d) 4-Chloromethyl-1-methyl-1,2,3-triazole hydrochloride

1-Methyl-4-hydroxymethyl-1,2,3-triazole hydrochloride hydrate (1.0 g, 6.08 mmol) was suspended in 20 ml thionyl chloride and the mixture was refluxed for 1 hour. Excess thionyl chloride was evaporated off and  
5 the residue was triturated with diethyl ether (25 ml). The white solid produced was washed by decantation with diethyl ether and dried *in vacuo* to give the required product (0.52 g, 51%). <sup>1</sup>H NMR (360 MHz, d<sup>6</sup>-DMSO) 4.04 (3H, s), 4.82 (2H, s), 8.16 (1H, s); MS<sup>+</sup> 131 & 133.


10 e) 3-(5-Methylisoxazol-3-yl)-6-[(1-methyl-1,2,3-triazol-4-yl)methoxy]imidazo[2,1-a]phthalazine

3-(5-Methylisoxazol-3-yl)imidazo[2,1-a]phthalazin-6-one (100 mg, 0.375 mmol) was suspended in DMF (5 ml) and sodium hydride (60% dispersion in oil, 33 mg, 0.83 mmol) was added. After stirring at room  
15 temperature for 10 minutes potassium carbonate (50 mg, 0.36 mmol) was added, followed by 4-chloromethyl-1-methyl triazole hydrochloride (69 mg, 0.41 mmol). The solution was heated at 80°C for 4 hours, then cooled and poured into water. The solid product was filtered off and washed with water and hexane. The solid was recrystallised from ethyl acetate/hexane  
20 to yield the title compound (83 mg, 62%), <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>) 2.55 (3H, s), 4.11 (3H, s), 5.75 (2H, s), 6.72 (1H, s), 7.67 (1H, t), 7.86 (1H, t), 7.99 (1H, s), 8.17 (2H, d), 8.51 (1H, d); MS<sup>+</sup> 362 [MH]<sup>+</sup>.

CLAIMS

1. A compound of formula I, or a pharmaceutically acceptable salt thereof:

5



(I)

wherein:

L is O, S or  $\text{NR}^n$  where  $\text{R}^n$  is H,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{3-6}$ cycloalkyl;

10 X is a 5-membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently chosen from oxygen, nitrogen and sulphur, at most one of the heteroatoms being oxygen or sulphur, or a 6-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms, the 5- or 6-membered heteroaromatic ring being optionally fused to a benzene or  
 15 pyridine ring and the heteroaromatic ring being optionally substituted by  $\text{R}^x$  and/or  $\text{R}^y$  and/or  $\text{R}^z$ , where  $\text{R}^x$  is halogen,  $\text{R}^3$ ,  $\text{OR}^3$ ,  $\text{OCOR}^3$ ,  $\text{NR}^4\text{R}^5$ ,  $\text{NR}^4\text{COR}^5$ , OH,  $\text{tri}(\text{C}_{1-6}\text{alkyl})\text{silylC}_{1-6}\text{alkoxyC}_{1-4}\text{alkyl}$ , CN or  $\text{R}^6$ ,  $\text{R}^y$  is halogen,  $\text{R}^3$ ,  $\text{OR}^3$ ,  $\text{OCOR}^3$ ,  $\text{NR}^4\text{R}^5$ ,  $\text{NR}^4\text{COR}^5$ ,  $\text{NR}^4\text{R}^5(\text{C}_{1-6})\text{alkyl}$  or CN and  $\text{R}^z$  is  $\text{R}^3$ ,  $\text{OR}^3$  or  $\text{OCOR}^3$ , providing that when X is a pyridine derivative,  
 20 the pyridine ring is optionally in the form of the N-oxide, and providing that when X is a tetrazole derivative it is protected by a  $\text{C}_{1-4}\text{alkyl}$  group; or X is phenyl optionally substituted by one, two or three groups independently selected from halogen, cyano,  $\text{C}_{1-6}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$  and  $\text{C}_{3-6}\text{cycloalkyl}$ ;

Y is optionally branched C<sub>1-4</sub>alkylene optionally substituted by an oxo group or Y is a group (CH<sub>2</sub>)<sub>j</sub>O wherein the oxygen atom is nearest the group X and j is 2, 3 or 4;


Z represents a phenyl ring, a 5-membered heteroaromatic ring  
5 containing 1, 2 or 3 heteroatoms independently selected from oxygen, nitrogen and sulphur, at most one of the heteroatoms being oxygen or sulphur, or a 6-membered heteroaromatic ring containing 1, 2 or 3 nitrogen atoms, each of which rings may be optionally substituted with one or more substituents selected from halogen, R<sup>3</sup>, OR<sup>3</sup>, OCOR<sup>3</sup>, NR<sup>4</sup>R<sup>5</sup>,  
10 NR<sup>4</sup>R<sup>5</sup>(C<sub>1-6</sub>)alkyl, NR<sup>4</sup>R<sup>5</sup>CO, NR<sup>4</sup>R<sup>5</sup>CO(C<sub>1-6</sub>)alkyl, CN, cyano(C<sub>1-6</sub>)alkyl or R<sup>6</sup>;

R<sup>3</sup> represents C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-6</sub>cycloalkyl, C<sub>3-6</sub>cycloalkyl(C<sub>1-6</sub>)alkyl, cyano(C<sub>1-6</sub>)alkyl, hydroxyC<sub>1-6</sub>alkyl and R<sup>3</sup> is optionally mono, di- or tri-fluorinated;

15 R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-6</sub>cycloalkyl or CF<sub>3</sub> or R<sup>4</sup> and R<sup>5</sup>, together with a nitrogen atom to which they are commonly attached, form a 4-7 membered heteroaliphatic ring containing the said nitrogen atom and optionally one other heteroatom selected from O, N and S; and

20 R<sup>6</sup> is C<sub>6-10</sub>aryl, C<sub>6-10</sub>aryl (C<sub>1-6</sub>)alkyl, heteroaryl or heteroaryl(C<sub>1-6</sub>)alkyl, where heteroaryl is a five-membered aromatic ring containing one, two, three or four nitrogen atoms or one oxygen atom or sulphur atom and optionally one or two nitrogen atoms or a six-membered aromatic ring containing one, two or three nitrogen atoms, and R<sup>6</sup> is optionally  
25 substituted by one, two or three substituents independently chosen from halogen atoms and C<sub>1-4</sub>alkyl, C<sub>2-4</sub>alkenyl, C<sub>2-4</sub>alkynyl, C<sub>1-4</sub>alkoxy, C<sub>2-4</sub>alkenyloxy and C<sub>2-4</sub>alkynyloxy groups each of which groups is unsubstituted or substituted by one, two or three halogen atoms.

30 2. A compound as claimed in claim 1 represented by a compound of formula IIA, or a pharmaceutically acceptable salt thereof:



(IIA)

wherein

- 5           X' represents phenyl, pyrazolyl, isoxazolyl, thiazolyl, imidazolyl, thiadiazolyl, benzimidazolyl, oxadiazolyl, triazolyl, tetrazolyl, pyridinyl, pyridazinyl, oxazolopyridinyl, pyrimidinyl, pyrazinyl, quinolinyl, isoquinolinyl or quinoxalinyl, any of which groups may be optionally substituted by one or more of C<sub>1-6</sub> alkyl, amino, pyridyl, CF<sub>3</sub>, aryl(C<sub>1-6</sub>)alkyl, pyridyl(C<sub>1-6</sub>)alkyl, halogen, cyano, cyano(C<sub>1-6</sub>)alkyl, hydroxymethyl, C<sub>1-6</sub>alkoxy, C<sub>3-6</sub>cycloalkyl(C<sub>1-6</sub>)alkoxy, hydroxy or the keto tautomer thereof, di(C<sub>1-6</sub>)alkylamino(C<sub>1-6</sub>)alkyl, amino(C<sub>1-6</sub>)alkyl, N-(C<sub>1-6</sub>)alkylpiperidinyl, pyrrolidinyl(C<sub>1-6</sub>)alkyl, piperazinyl(C<sub>1-6</sub>)alkyl and morpholinyl(C<sub>1-6</sub>)alkyl; and
- 15           Z' represents a 5-membered heteroaromatic ring containing one oxygen and one or two nitrogen ring atoms which is optionally substituted with C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkoxy.

3.       The compound:


- 20       3-(5-methylisoxazol-3-yl)-6-[(1-methyl-1,2,3-triazol-4-yl)methoxy]imidazo[2,1-a]phthalazine  
and pharmaceutically acceptable salts thereof.

4.       A pharmaceutical composition comprising a compound of formula I  
25       as defined in claim 1, 2 or 3 or a pharmaceutically acceptable salt thereof  
in combination with a pharmaceutically acceptable carrier.


5. The use of a compound as claimed in any one of claims 1 to 3 for the manufacture of a medicament for the enhancement of cognition.

5 6. A process for the preparation of a compound as claimed in claim 1 which comprises:

(A) reacting a compound of formula II with a compound of formula III:



(II)



(III)

wherein X, Y and Z are as defined in claim 1; and

(B) subsequently, if desired, converting a compound of formula I  
15 initially obtained into a further compound of formula I by techniques known from the art.

7. A method of treatment of a subject suffering from a cognition deficit which comprises administering to that subject an effective amount of a  
20 compound of formula I as defined in claim 1, or a pharmaceutically acceptable salt thereof.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/03166

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/50 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                 | Relevant to claim No. |
|------------|--------------------------------------------------------------------------------------------------------------------|-----------------------|
| Y          | WO 99 37644 A (MERCK SHARP & DOHME)<br>29 July 1999 (1999-07-29)<br>page 3, line 17 - line 31; claims;<br>examples | 1-7                   |
| Y          | WO 00 12505 A (MERCK SHARP & DOHME)<br>9 March 2000 (2000-03-09)<br>page 3, line 17 - line 28; claims;<br>examples | 1-7                   |
| Y          | US 6 013 799 A (SHAW ET. AL.)<br>11 January 2000 (2000-01-11)<br>claims; example 19                                | 1-7                   |
|            | ---<br>-/--                                                                                                        |                       |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

11 October 2001

Date of mailing of the international search report

24/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

## INTERNATIONAL SEARCH REPORT

Inter Application No

PCT/GB 01/03166

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                                                                                                                                                                                                                                                                                                                        | Relevant to claim No. |
|------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| Y          | CHEMICAL ABSTRACTS, vol. 119, no. 15,<br>11 October 1993 (1993-10-11)<br>Columbus, Ohio, US;<br>abstract no. 151635r,<br>CATARZI, D. ET. AL.: "Reaction of<br>1-aminophthalazines with<br>alpha-halocarbonyl compounds.<br>Imidazo[2,1-a]phthalazines and their<br>benzodiazepine receptor activities."<br>page 22; column 1;<br>XP002179824<br>abstract<br>& IL FARMACO,<br>vol. 48, no. 4, 1993, pages 447-57,<br>----- | 1-7                   |
| A          | DE 25 23 876 A (ASPRO NICHOLAS LTD.)<br>15 January 1976 (1976-01-15)<br>claims; example 3<br>-----                                                                                                                                                                                                                                                                                                                        | 1-7                   |

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern Application No  
PCT/GB 01/03166

| Patent document<br>cited in search report |   | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|-------------------------------------------|---|---------------------|----------------------------|---------------------|
| WO 9937644                                | A | 29-07-1999          | AU 2065799 A               | 09-08-1999          |
|                                           |   |                     | BG 104671 A                | 31-05-2001          |
|                                           |   |                     | BR 9906905 A               | 10-10-2000          |
|                                           |   |                     | CN 1288465 T               | 21-03-2001          |
|                                           |   |                     | EP 1049696 A1              | 08-11-2000          |
|                                           |   |                     | WO 9937644 A1              | 29-07-1999          |
|                                           |   |                     | HR 20000492 A1             | 31-12-2000          |
|                                           |   |                     | HU 0100575 A2              | 28-09-2001          |
|                                           |   |                     | NO 20003730 A              | 20-09-2000          |
|                                           |   |                     | PL 341652 A1               | 23-04-2001          |
|                                           |   |                     | TR 200002100 T2            | 21-12-2000          |
| WO 0012505                                | A | 09-03-2000          | US 6297235 B1              | 02-10-2001          |
|                                           |   |                     | AU 5435599 A               | 21-03-2000          |
|                                           |   |                     | EP 1107967 A2              | 20-06-2001          |
|                                           |   |                     | WO 0012505 A2              | 09-03-2000          |
| US 6013799                                | A | 11-01-2000          | US 5696260 A               | 09-12-1997          |
|                                           |   |                     | US 5426186 A               | 20-06-1995          |
| DE 2523876                                | A | 15-01-1976          | AU 8162975 A               | 02-12-1976          |
|                                           |   |                     | BE 829522 A1               | 27-11-1975          |
|                                           |   |                     | DE 2523876 A1              | 15-01-1976          |
|                                           |   |                     | DK 236175 A                | 01-12-1975          |
|                                           |   |                     | FR 2272661 A1              | 26-12-1975          |
|                                           |   |                     | JP 51006997 A              | 20-01-1976          |
|                                           |   |                     | NL 7506327 A               | 02-12-1975          |
|                                           |   |                     | ZA 7503363 A               | 26-05-1976          |