




(86) Date de dépôt PCT/PCT Filing Date: 2008/04/10
(87) Date publication PCT/PCT Publication Date: 2008/10/23
(45) Date de délivrance/Issue Date: 2020/01/07
(85) Entrée phase nationale/National Entry: 2009/10/09
(86) N° demande PCT/PCT Application No.: US 2008/059949
(87) N° publication PCT/PCT Publication No.: 2008/127996
(30) Priorité/Priority: 2007/04/13 (US60/911,773)

(51) Cl.Int./Int.Cl. *A61K 38/10* (2006.01),
A61P 35/00 (2006.01)
(72) Inventeurs/Inventors:
GULATI, ANIL, US;
GURU, REDDY, US;
LENAZ, LUIGI, US
(73) Propriétaires/Owners:
SPECTRUM PHARMACEUTICALS, INC., US;
THE BOARD OF TRUSTEES OF THE UNIVERSITY OF
ILLINOIS, US
(74) Agent: SMART & BIGGAR LLP

(54) Titre : PROCEDES ET COMPOSITIONS SUSCEPTIBLES DE CONTRIBUER AU TRAITEMENT DE CANCERS
(54) Title: METHODS AND COMPOSITIONS FOR CONTRIBUTING TO THE TREATMENT OF CANCERS



(57) **Abrégé/Abstract:**

Methods and compositions for contributing to the treatment of cancers, especially ovarian tumors, are disclosed. The methods and compositions utilize an endothelin B agonist (ET_B) to enhance the delivery and resulting efficacy of chemotherapeutic agent(s) (e.g., cisplatin and/or cyclophosphamide).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 October 2008 (23.10.2008)

PCT

(10) International Publication Number
WO 2008/127996 A1

(51) International Patent Classification:

A61K 31/337 (2006.01) A61K 38/20 (2006.01)
 A61K 31/513 (2006.01) A61K 45/06 (2006.01)
 A61K 31/704 (2006.01) A61P 35/00 (2006.01)
 A61K 38/03 (2006.01) A61K 31/675 (2006.01)
 A61K 38/10 (2006.01) A61K 33/24 (2006.01)
 A61K 38/16 (2006.01)

(21) International Application Number:

PCT/US2008/059949

(22) International Filing Date: 10 April 2008 (10.04.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/911,773 13 April 2007 (13.04.2007) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:

US 60/911,773 (CIP)
 Filed on 13 April 2007 (13.04.2007)

(71) Applicants (for all designated States except US): SPECTRUM PHARMACEUTICALS, INC. [US/US]; 157

Technology Drive, Irvine, CA 92618 (US). **THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ILLINOIS** [US/US]; 352 Administration Building, 506 South Wright Street, Urbana, IL 61801 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GULATI, Anil** [US/US]; 1929 Lisson Road, Naperville, IL 60665 (US). **GURU, Reddy** [US/US]; 120 Arden, Irvine, CA 92620 (US). **LENAZ, Luigi** [US/US]; 11 Planetree Court, Newton, PA 18940 (US).

(74) Agents: **CULLMAN, Louis, C.** et al.; Kirkpatrick & Lockhart Preston Gates Ellis LLP, 1900 Main Street, Sixth Floor, Irvine, CA 92614-7319 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US (patent), UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]

(54) Title: METHODS AND COMPOSITIONS FOR CONTRIBUTING TO THE TREATMENT OF CANCERS

FIG. 1

Effect of IRL-1620 on the efficacy of cisplatin (CIS)
 in ovarian tumor bearing athymic mice



(57) Abstract: Methods and compositions for contributing to the treatment of cancers, especially ovarian tumors, are disclosed. The methods and compositions utilize an endothelin B agonist (ET_B) to enhance the delivery and resulting efficacy of chemotherapeutic agent(s) (e.g., cisplatin and/or cyclophosphamide).

WO 2008/127996 A1



(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,

NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

51432-75

METHODS AND COMPOSITIONS FOR CONTRIBUTING TO THE TREATMENT OF CANCERS

[0001]

FIELD OF INVENTION

[0002] The present invention relates to methods and compositions for the treatment of cancers, including solid tumors, with an endothelin agonist and at least one chemotherapeutic agent such as cyclophosphamide or cisplatin.

BACKGROUND OF INVENTION

[0003] Chemotherapy with most anticancer agents is complicated by limitations, such as low therapeutic index and host tissue toxicity, which leads to life-threatening situations and decreases the quality of life for many patients. Lack of efficacy and significant potential side effects of standard chemotherapy regimens are driving the search for novel approaches to anticancer therapy. One approach to increase the effectiveness of existing anticancer drugs for cancer treatment is to increase the uptake of drugs into the tumor thereby increasing the therapeutic tumor concentration.

[0004] Successful treatment of cancers, including solid tumors, remains an unfulfilled medical goal, despite increased understanding of the molecular biology of tumor cells and the availability of an increased number of potential therapeutic agents. For example, ovarian cancer is the fifth leading cause of cancer deaths in

women, the leading cause of death from gynecological malignancy, and the second most commonly diagnosed gynecologic malignancy.

[0005] One problem in the treatment of cancers is that an effective dose of a wide variety of potential chemotherapeutic agents is restricted by these agents' non-selective, highly toxic effect on normal tissues. As a result, many patients suffer from the side effects of chemotherapy without reaping the benefits of the treatment. For example, the chemotherapeutic agents cyclophosphamide and cisplatin inhibit cellular proliferation and induces apoptosis of tumor cells. The clinical utility of cyclophosphamide and cisplatin have been hampered, however, by dose limiting toxicities. Thus, there is a necessity to develop more specific and less toxic cancer therapies.

[0006] Targeted delivery of chemotherapeutic agents to tumors could have the advantage of enhancing the benefit of chemotherapeutic agents while minimizing their systemic toxic effects. Such targeted delivery could also serve to lower the required dose of chemotherapeutic agents thus potentially reducing the unacceptable adverse effects of these agents. One possible way to achieve targeted delivery of chemotherapeutic agents is to utilize the distinctive features of tumor vasculature.

[0007] Tumors greater than a few millimeters in size require a constant nutrient supply, and, therefore, develop their own vascular bed and blood flow (Folkman, Cancer Res, 46:467 (1986)). Without constant nourishment from these developing blood vessels, the tumors become hypoxic and subsequently die. Recruitment of new vasculature from preexisting blood vessels is termed "angiogenesis."

[0008] During angiogenesis, tumor blood vessels develop substantially differently from normal vasculature, and have different properties. Single layered epithelial cells are the first hastily formed tumor blood vessels. These newly formed tumor blood vessels do not have a smooth muscle layer or innervation. Tumors also incorporate mature blood vessels that possess all their autoregulatory functions. Mattsson *et al.*, Tumor Blood Circulation, CRC Press, Boca Raton, pg. 129 (1979); Reinhold, Tumor Blood Circulation, CRC Press, Boca Raton, pg. 115 (1979); Warren, Tumor Blood Circulation, CRC Press, Boca Raton, pg. 26 (1979).

[0009] Vascular tone (the degree to which blood vessels are dilated or constricted) is governed by a host of endogenous factors including H^+ , K^+ , Ca^{2+} , pO_2 , pCO_2 and nitric oxide (NO), as well as other regulatory substances such as endothelin (ET-1). Secombe *et al.*, Landes, Austin, pg. 40 (1994); Luscher *et al.*, The endothelium: modulator of cardiovascular function, CRC Press, Boca Raton, pg. 61 (1990). ET-1 contributes significantly to regulating vascular tone (Yanagisawa *et al.*, Nature, 332:411 (1988)) and investigators have shown an increase in ET1 and ET_B receptor expression in solid tumors including breast carcinomas. Alanen *et al.*, Histopathology, 36:161 (2000); Nelson *et al.*, Cancer Res, 56:663 (1996); Kar *et al.*, Biochem Biophys Res Commun 216:514 (1995); Pagotto *et al.*, J Clin Invest, 96:2017 (1995); Yamashita *et al.*, Cancer Res, 52:4046 (1992); Yamashita *et al.*, Res Commun Chem Pathol Pharmacol, 74:363 (1991). Further, stimulation of ET_B receptors causes an increase in blood supply to tumors through vasodilation of tumor blood vessels. The present disclosure takes advantage of this fact by using ET_B receptor agonists to selectively increase blood flow to tumors to enhance the targeted delivery of chemotherapeutic agents

SUMMARY OF INVENTION

[0010] The present disclosure is related to selectively-increasing the uptake of anticancer agents such as cyclophosphamide and cisplatin, into tumors such as ovarian tumors by administering an ET_B agonist with at least one anti-cancer agent. A treatment regimen in accordance with the present invention reduces tumor volume, such as ovarian tumor volume, in animals.

[0011] The methods of the present invention contribute to the treatment of cancer. The methods include administering to an animal in need of treatment, at least one endothelin B (ET_B) agonist and at least one chemotherapeutic agent. The cancers treatable in accordance with the present invention include, but are not limited to, solid tumors like ovarian tumors, colon tumors, Kaposi's sarcoma, breast tumors, melanomas, prostate tumors, meningiomas, liver tumors, breast phyllode tumors, brain tumors, neck tumors, lung tumors, and combinations of tumors. The methods of the present invention include mechanisms whereby the ET_B agonist selectively increases the blood supply to the tumor thereby increasing the delivery of

chemotherapeutic agent(s) to the tumor. The ET_B agonist and the chemotherapeutic agents may be administered substantially simultaneously (e.g. as a single composition) or sequentially (e.g. the chemotherapeutic agent is administered prior to the ET_B agonist or vice versa).

5 **[0012]** The present invention is also directed to compositions having at least one chemotherapeutic agent, at least one ET_B agonist, and, optionally, at least one pharmaceutically acceptable excipient. The ET_B agonist(s) and chemotherapeutic agent(s) may be part of the same composition, are provided as separate compositions, or both.

10 **[0013]** ET_B agonists suitable for use in either the methods or the compositions of the present invention include, but are not limited to, ET-1, ET-2, ET-3, BQ3020, IRL1620 (N-SUC-[Glu⁹, Ala^{11, 15}]ET-1 (8-21)), sarafotoxin 56c, [Ala^{1, 3, 11, 15}]ET-1, and combinations. IRL-1620 is a preferred ET_B agonist.

[0014] Chemotherapeutic agents suitable for use in either the methods or the
15 compositions of the present invention include, but are not limited to, adriamycin, camptothecin, carboplatin, cisplatin, cyclophosphamide, daunorubicin, doxorubicin, alpha interferon, beta interferon, gamma interferon, interleukin 2, irinotecan, docetaxel, paclitaxel, topotecan, 5-fluorouracil, and combinations. Cisplatin, cyclophosphamide, and their combination are preferred chemotherapeutic agents.

20 **[0014a]** In one use aspect, the invention relates to use of an endothelin B (ET_B) receptor agonist in combination with a chemotherapeutic agent for the treatment of an ovarian tumor wherein the ET_B receptor agonist is IRL-1620, wherein said chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.

[0014b] In one composition aspect, the invention relates to a composition for
25 use in the treatment of an ovarian cancer comprising at least one chemotherapeutic agent, an IRL-1620, and an optional excipient, wherein said chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.

[0014c] In one kit aspect, the invention relates to a kit comprising a composition comprising IRL-1620 and an excipient and instructional information directing the administration of said composition comprising said IRL-1620 in combination with a chemotherapeutic agent to treat an ovarian tumor, wherein the kit further comprises a
5 composition comprising a chemotherapeutic agent and an excipient, wherein said chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.

[0014d] In a further use aspect, the invention relates to use of an endothelin B (ET_B) receptor agonist in the manufacture of a medicament to be used in conjunction with a chemotherapeutic agent for the treatment of an ovarian tumor wherein said
10 (ET_B) receptor agonist is IRL-1620, wherein said chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] Figure 1 depicts the tumor volume of ovarian tumor mice (N=6) separately given saline, IRL-1620 (3 nmol/kg), saline + cisplatin (2.5 mg/kg), or
15 IRL-1620 (3 nmol/kg) + cisplatin (2.5 mg/kg) once every three days for a total of 4 doses;

[0016] Figure 2 depicts the body weight of ovarian tumor mice (N=6) separately given saline, IRL-1620 (3 nmol/kg), saline + cisplatin (2.5 mg/kg), or IRL-1620 (3 nmol/kg) + cisplatin (2.5 mg/kg) once every three days for a total
20 of 4 doses;

[0017] Figure 3 depicts the tumor volume of ovarian tumor mice (N=6) separately given saline, IRL-1620 (3 nmol/kg), saline + cyclophosphamide (75 mg/kg), or IRL-1620 (3 nmol/kg) + cyclophosphamide (75 mg/kg) once every three days for a total of 4 doses; and,

[0018] Figure 4 depicts the body weight of ovarian tumor mice (N=6) separately given saline, IRL-1620 (3 nmol/kg), saline + cyclophosphamide (75 mg/kg), or IRL-1620 (3 nmol/kg) + cyclophosphamide (75 mg/kg) once every three days for a total of 4 doses.

DETAILED DESCRIPTION OF THE INVENTION

[0019] The methods of the present invention contribute to the treatment of cancer. In one embodiment, the present disclosure relates to a method of contributing to the treatment of cancer comprising administering an endothelin B (ET_B) agonist and at least one chemotherapeutic agent.

[0020] As used herein, the terms "treat", "treatment" and "contributing to the treatment of" shall mean preventing, retarding the progression or growth of, shrinking, or eliminating a cancer including a solid tumor. As such, these terms include both medical therapeutic and/or prophylactic administration, as appropriate.

[0021] Most chemotherapeutic agents have cytotoxic properties that are targeted to destroy cancer cells, but in the process inflict considerable damage to the body's normal physiological systems. It would be of great advantage, therefore, to selectively deliver chemotherapeutic agents to solid tumors thus helping to avoid these negative effects of cancer treatment.

[0022] The angioarchitecture of tumor blood vessels is different from that of normal blood vessels. Carmeliet & Jain, *Nature*, 407:249 (2000). Therefore, the vascular reactivity of tumors differs from that of normal tissue. For example, the administration of nitric oxide donors, nicotinamide and bradykinin agonists modulate blood flow to tumors. Jordan *et al.*, *Int J Radiat Oncol Biol Phys*, 48:565 (2000); Fukumura *et al.*, *Am J Pathol*, 150:713 (1997); Hirst *et al.*, *Br J Radiol*, 67: 795 (1994).

[0023] Endothelin is a vasoactive substance that modulates blood flow and is present in large concentrations in breast carcinoma tissues compared to normal breast tissue (specifically, endothelin can be present in an amount of about 12 pg/mg in breast carcinoma tissues as compared to about 0.12 pg/mg in normal breast tissue). Kojima *et al.*, *Surg Oncol*, 4(6):309 (1995); Kurbel *et al.*, *Med Hypotheses*,

52(4):329 (1999); Patel *et al.*, Mol Cell Endocrinol, 126(2):143 (1997); Yamashita *et al.*, Cancer Res, 52(14):4046 (1992); Yamashita *et al.*, Res Commun Chem Pathol Pharmacol, 74(3):363 (1991). Endothelins are a family of cyclic peptides with 21 amino acids, comprising three isoforms in mammals, ET-1, ET-2 and ET-3. Inoue *et al.*, Proc Natl Acad Sci USA 86:2863 (1989); Yanagisawa *et al.*, Nature, 332:411 (1988). Endothelins exert their effects by binding to two distinct cell surface receptors, ET_A and ET_B. The ET_B receptor binds the three peptide isotypes with equal affinity. In contrast, the ET_A receptor binds ET-1 with higher affinity than the other isoforms. Both receptors belong to the G protein-coupled receptor system and mediate biological responses from a variety of stimuli, including growth factors, vasoactive polypeptides, neurotransmitters and hormones. Masaki, J Cardiovasc Pharmacol, 35:S3 (2000); Gulati, Preface. Adv Drug Deliv Rev, 40:129 (2000); Gulati *et al.*, Am J Physiol, 273:H827 (1997); Levin, N Engl J Med, 333:356 (1995). ET_B receptors, a focus of the present invention, are present on both endothelial cells (ECs) and vascular smooth muscle cells (VSMCs) and are increased in breast cancer tissue (including in invasive as well as in ductal and lobular breast carcinoma tissue in humans) when compared to normal breast tissue. Wulfing *et al.*, Oncol Rep, 11:791 (2004); Wulfing *et al.*, Clin Cancer Res, 9:4125 (2003); Alanen *et al.*, Histopathology, 36(2):161 (2000). Endothelin acts on ET_B receptors to produce vascular dilation and increase blood flow to breast tumor tissue. ET_B receptors predominating on ECs, produce vasodilatation via the release of factors such as prostacyclin and nitric oxide. de Nucci *et al.*, Proc Natl Acad Sci USA, 85:9797 (1988). Because ET-1 produces an increase in blood flow to tumors by stimulating ET_B receptors, an ET_B receptor agonist can be used to selectively increase blood supply to tumors, thus increasing the targeted delivery and resulting efficacy of chemotherapeutic agents.

[0024] ET_B receptors have been shown in, for example and without limitation, ovarian cancers, myofibroblasts, Kaposi's sarcoma tumor and intratumoral vessels, breast cancers and melanomas. Bagnato *et al.*, Am J Pathol, 158:841 (2001); Alanen *et al.*, Histopathology, 36(2):161 (2000); Bagnato *et al.*, Cancer Res, 59:720 (1999); Kikuchi *et al.*, Biochem Biophys Res Comm, 219:734 (1996). Therefore, administration of an ET_B receptor agonist in combination with a chemotherapeutic agent can be used to contribute to the treatment of solid tumors, including, without

limitation, ovarian cancer, colon carcinoma, Kaposi's sarcoma, breast cancer, and melanomas.

[0025] ET_B agonists useful in accordance with the present invention include, without limitation, ET-1, ET-2, ET-3, BQ3020, IRL1620 (N-suc-[Glu⁹, Ala^{11,15}]ET-1 (8-21)), sarafotoxin 56c, [Ala^{1,3,11,15}]ET-1, and combinations thereof. [Ala^{1,3,11,15}]ET-1 is a linear analog of ET-1 in which the disulfide bridges have been removed by substitution of Ala for Cys residues. Saeki et al., *Biochem Biophys Res Commun*, 179:286 (1991). BQ3020 and IRL1620 are truncated linear synthetic analogs of ET-1 and are the most widely used selective synthetic agonists. IRL-1620 is a linear ET-analog whose structure is based on the carboxy terminal end of ET-1 and has 120,000 fold selectivity for the ET_B receptors. Okada & Nishikibe, *Cardiovasc Drug Rev*, 20:53 (2002); Douglas et al., *Br J Pharmacol*, 114:1529 (1995). IRL-1620 is a highly selective and potent ET_B agonist, with evidence being reported of its selectivity for the ET_{B1} receptor subtype in preference over the ET_{B2} subtype. Brooks et al., *J Cardiovasc Pharmacol*, 26 Suppl 3:S322 (1995).

[0026] Chemotherapeutic agents useful in accordance with the present invention include, for example and without limitation, alkylating agents, antimetabolites, hormones and antagonists thereof, radioisotopes, antibodies, as well as natural products, and combinations thereof. For example, an ET_B agonist can be administered with antibiotics, such as doxorubicin and other anthracycline analogs, nitrogen mustards, such as, without limitation, cyclophosphamide, pyrimidine analogs such as, without limitation, 5-fluorouracil, cisplatin, hydroxyurea, and its natural and synthetic derivatives, and the like. As another example, in the case of mixed tumors, such as adenocarcinoma of the breast, where the tumors include gonadotropin-dependent and gonadotropin-independent cells, the ET_B agonist can be administered in conjunction with, without limitation, leuprolide or goserelin (synthetic peptide analogs of LH-RH). Additional non-limiting examples of chemotherapeutic agents that can be used with the present invention include cyclophosphamide, cisplatin, adriamycin, camptothecin, carboplatin, daunorubicin, doxorubicin, interferon (alpha, beta, and/or gamma), interleukin 2, irinotecan, docetaxel, paclitaxel, topotecan, and therapeutically effective analogs and derivatives of the same.

[0027] It is theorized, but not relied upon herein, that endothelin agonists stimulate ET_B receptors to dilate tumor blood vessels, thereby increasing blood flow and the resultant delivery of chemotherapeutic agents to the tumor. The increased blood perfusion of tumors caused by endothelin agonists also increases oxygenation of the tissue. Improved oxygenation can enhance the therapeutic action of chemotherapeutic agents. Endothelin also can have mitogenic properties. The mitogenic actions of endothelin can help increase the action of chemotherapeutic agents, when administered together. The mitogenic action of an endothelin agonist can increase the action of chemotherapeutic agents by improving their incorporation into dividing cells, thus increasing their efficacy.

[0028] Chemotherapy is frequently indicated as an adjuvant to surgery in the treatment of a cancer. The goal of chemotherapy in the adjuvant setting is to reduce the risk of recurrence and enhance disease-free survival when the primary tumor has been controlled. Chemotherapy is utilized as a treatment adjuvant for a cancer, frequently when the disease is metastatic. An ET_B agonist, therefore, is particularly useful before or following surgery in the treatment of a solid tumor in combination with chemotherapy.

[0029] In another embodiment of the present method, the cancer is a solid tumor. In another embodiment, the solid tumor is selected from the group consisting of an ovarian tumor, a colon tumor, Kaposi's sarcoma, a breast tumor, a melanoma, a prostate tumor, a meningioma, a liver tumor, a breast phyllode tumor, brain tumor, neck tumor, lung tumor, and combinations thereof. In another embodiment, the solid tumor is ovarian tumor.

[0030] In another embodiment, the ET_B agonist is selected from the group consisting of ET-1, ET-2, ET-3, BQ3020, IRL1620 (N-suc-[Glu⁹, Ala^{11,15}]ET-1 (8-21)), sarafotoxin 56c, [Ala^{1, 3, 11, 15}]ET-1, and combinations thereof. In another embodiment, the ET_B agonist is IRL-1620.

[0031] In another embodiment, the chemotherapeutic agent is selected from the group consisting of adriamycin, camptothecin, carboplatin, cisplatin, cyclophosphamide, daunorubicin, alpha interferon, beta interferon, gamma interferon, interleukin 2, irinotecan, docetaxel, paclitaxel, topotecan, 5-fluorouracil,

and combinations thereof. In another embodiment, the chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.

[0032] In another embodiment, the increase in the blood supply to the tumor increases the delivery of the chemotherapeutic agent to the solid tumor. In another embodiment, the ET_B agonist and the chemotherapeutic agent are administered substantially simultaneously. In another embodiment, the ET_B agonist and the chemotherapeutic agent are administered as a single composition. In another embodiment, the ET_B agonist and the chemotherapeutic agent are administered sequentially. The chemotherapeutic agent may be administered prior to the ET_B agonist or the ET_B agonist may be administered prior the chemotherapeutic agent.

[0033] The present disclosure also relates to a composition comprising a chemotherapeutic agent, and ET_B agonist, and an optional excipient. The ET_B agonist and the chemotherapeutic agent are part of the same composition, are provided as separate compositions, or both. The ET_B agonist may be selected from the group consisting of ET-1, ET-2, ET-3, BQ3020, IRL1620 (N-suc-[Glu⁹, Ala^{11,15}]ET-1 (8-21)), sarafotoxin 56c, [Ala^{1, 3, 11, 15}]ET-1, and combinations thereof. The ET_B agonist may be IRL-1620. The chemotherapeutic agent maybe selected from the group consisting of adriamycin, camptothecin, carboplatin, cisplatin, cyclophosphamide, daunorubicin, doxorubicin, alpha interferon, beta interferon, gamma interferon, interleukin 2, irinotecan, docetaxel, paclitaxel, topotecan, 5-fluorouracil, and combinations thereof.

[0034] In another embodiment, the chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof and the ET_B agonist is IRL-1620.

EXAMPLE

[0035] It surprisingly has been discovered that the administration of IRL-1620 and at least one anti-cancer agent selectively-increases the uptake of anticancer agents, e.g. cyclophosphamide and cisplatin, into ovarian tumors. A treatment regimen in accordance with the present invention reduces ovarian tumor volume in animals.

[0036] A preclinical efficacy study was conducted on the human ovarian tumor model in mice. The tumor cells used were SK-OV3 cells purchased from ATCC. The mice were 4–6 week old Athymic (nu+/nu+) female mice. The IRL-1620 was intravenously-infused slowly (over 30 seconds). The IRL-1620 was obtained from American Peptides, CA. The cyclophosphamide was obtained from Cadila Pharmaceuticals, Ahmedabad (Batch No. GW-4005). The cisplatin also was obtained from Cadila Pharmaceuticals, Ahmedabad (Batch No. KL-5005).

[0037] A solid tumor was induced in each animal using 3 million viable SK-OV3 cells suspended in 200 μ l of PBS per animal. After 10 days, when established tumors of 100 mm³ in diameter were detectable, the mice randomly were divided into six groups of 6 animals per group with each group receiving different treatments. The treatment regimens were administered via the tail vein every third day for a total of 4 doses. The groups were treated as follows: (1) tumor bearing mice received saline (N=6); (2) tumor bearing mice received IRL-1620 (3 nmol/kg) (N=6); (3) tumor bearing mice received vehicle (saline) + cyclophosphamide (75 mg/kg) (N=6); (4) tumor bearing mice received IRL-1620 (3 nmol/kg) + cyclophosphamide (75 mg/kg) (N=6); (5) tumor bearing mice received saline + cisplatin (2.5 mg/kg) (N=6); and, (6) tumor bearing mice received IRL-1620 (3 nmol/kg) + cisplatin (2.5 mg/kg) (N=6).

[0038] The effect of these treatments on the tumor size was measured using a digital caliper for a total of 12 days after drug administration. The mice were monitored and weighed on every third day. The administered doses were half of the maximum tolerated dose of cyclophosphamide and cisplatin for the human ovarian xenograft model in nude mice reported in published literature (*British Journal of Cancer* (2000) 83, 921-927). The animals were euthanized with pentobarbital 150-200 mg/kg for each mouse. Progression, stasis, partial regression or complete regression of tumor growth were measured by tumor load measurements with calipers and was calculated using the following formula: $(3.14/6) \times \text{larger diameter} \times \text{smaller diameter}$.

[0039] The results of the study are shown in the figures. Figure 1 depicts tumor volume of ovarian tumor bearing mice given IRL-1620 (3 nmol/kg) and cisplatin (2.5 mg/kg). Six ovarian tumor bearing mice (N=6) were given saline, IRL-1620 (3 nmol/kg), saline + cisplatin (2.5 mg/kg), or IRL-1620 (3 nmol/kg) + cisplatin (2.5

mg/kg). The dosing schedule was once every three days for a total of 4 doses. Body weight and tumor volumes were measured on every third day through the 22nd day. IRL-1620/cisplatin treated mice showed more reduction in tumor volume compared to saline/cisplatin treated mice.

[0040] Figure 2 depicts body weight of ovarian tumor bearing mice given IRL-1620 (3 nmol/kg) and cisplatin (2.5 mg/kg). Six ovarian tumor bearing mice (N=6) were given saline, IRL-1620 (3 nmol/kg), saline + cisplatin (2.5 mg/kg), or IRL-1620 (3 nmol/kg) + cisplatin (2.5 mg/kg). The dosing schedule was once every three days for a total of 4 doses. Body weight and tumor volumes were measured on every third day through the 22nd day. The body weight in all groups was similar.

[0041] Figure 3 depicts tumor volume of ovarian tumor bearing mice given IRL-1620 (3 nmol/kg) and cyclophosphamide (75 mg/kg). Six ovarian tumor bearing mice (N=6) were given saline, IRL-1620 (3 nmol/kg), saline + cyclophosphamide (75 mg/kg), or IRL-1620 (3 nmol/kg) + cyclophosphamide (75 mg/kg). The dosing schedule was once every three days for a total of 4 doses. Body weight and tumor volumes were measured on every third day through the 22nd day. IRL-1620/cyclophosphamide treated mice showed more reduction in tumor volume compared to saline/cyclophosphamide treated mice.

[0042] Figure 4 depicts body weight of ovarian tumor bearing mice given IRL-1620 (3 nmol/kg) and cyclophosphamide (75 mg/kg). Six ovarian tumor bearing mice (N=6) were given saline, IRL-1620 (3 nmol/kg), saline + cyclophosphamide (75 mg/kg), IRL-1620 (3 nmol/kg) + cyclophosphamide (75 mg/kg). Dosing schedule was once every three days for a total of 4 doses. Body weight and tumor volumes were measured on every third day through the 22nd day. The body weight in all the groups was similar.

[0043] Based on these results, it is apparent that the administration of IRL-1620 alone with, *inter alia*, cyclophosphamide or cisplatin contributes to the treatment of ovarian cancer tumors by reducing tumor size. There was a significant increase in the tumor volume in saline only treated ovarian tumor bearing mice. In this study, IRL-1620 alone did not significantly affect tumor volume when compared to controls. Animals treated with saline/cisplatin showed a significant reduction in tumor volume. However, IRL-1620/cisplatin treated mice showed a greater reduction in tumor

volume than saline/cisplatin treated mice. Animals treated with saline/cyclophosphamide showed a significant reduction in tumor volume. However, IRL-1620/cyclophosphamide treated mice showed a greater reduction in tumor volume than saline/cyclophosphamide treated mice.

[0100] Pharmaceutical compositions containing the described active ingredients are suitable for administration to humans or other mammals. Typically, the pharmaceutical compositions are sterile, and contain no toxic, carcinogenic, or mutagenic compounds that would cause an adverse reaction when administered. Administration of the pharmaceutical composition can be performed before, during, or after the onset of solid tumor growth.

[0101] A method of the present invention can be accomplished using active ingredients as described above, or as a physiologically acceptable salt, derivative, prodrug, or solvate thereof. The active ingredients can be administered as the neat compound, or as a pharmaceutical composition containing either or both entities.

[0102] As used herein, the term "prodrug" shall mean compounds that transform rapidly *in vivo* to a compound useful in the invention, for example, by hydrolysis. A thorough discussion of prodrugs is provided in Higuchi *et al.*, Prodrugs as Novel Delivery Systems, Vol. 14, of the A.C.S.D. Symposium Series, and in Roche (ed.), Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987.

[0103] The pharmaceutical compositions include those wherein the active ingredients are administered in an effective amount to achieve their intended purpose. More specifically, a "therapeutically effective amount" means an amount effective to prevent development of, to eliminate, to retard the progression of, or to reduce the size of a solid tumor. Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

[0104] A "therapeutically effective dose" refers to that amount of the active ingredients that results in achieving the desired effect. Toxicity and therapeutic efficacy of such active ingredients can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., determining the LD₅₀ (the

dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, which is expressed as the ratio between LD₅₀ and ED₅₀. A high therapeutic index is preferred. The data obtained can be used in formulating a range of dosage for use in humans. The dosage of the active ingredients preferably lies within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed, and the route of administration utilized.

[0105] The exact formulation and dosage is determined by an individual physician in view of the patient's condition. Dosage amount and interval can be adjusted individually to provide levels of the active ingredients that are sufficient to maintain therapeutic or prophylactic effects.

[0106] The amount of pharmaceutical composition administered can be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration, and the judgment of the prescribing physician.

[0107] The active ingredients can be administered alone, or in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. Pharmaceutical compositions for use in accordance with the present invention thus can be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries that facilitate processing of the active ingredients into preparations which can be used pharmaceutically.

[0108] As used herein, excipients may be pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral or intranasal application which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary

agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds.

[0109] When a therapeutically effective amount of the active ingredients is administered, the composition can be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred composition for intravenous injection typically will contain an isotonic vehicle although this characteristic is not required.

[0110] For veterinary use, the active ingredients are administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen that is most appropriate for a particular animal.

[0111] Various adaptations and modifications of the embodiments can be made and used without departing from the scope and spirit of the present invention which can be practiced other than as specifically described herein. The above description is intended to be illustrative, and not restrictive. The scope of the present invention is to be determined only by the claims.

[0044] The terms and expressions which have been employed herein are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding equivalents of the features shown and described, or portions thereof, it being recognized that various modifications are possible within the scope of the present invention claimed. Moreover, any one or more features of any embodiment of the present invention can be combined with any one or more other features of any other embodiment of the present invention, without departing from the scope of the present invention.

[0045] Unless otherwise indicated, all numbers expressing quantities of ingredients, properties such as molecular weight, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are

approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0046] The terms “a,” “an,” “the” and similar referents used in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. Recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of the invention.

[0047] Groupings of alternative elements or embodiments of the invention disclosed herein are not to be construed as limitations. Each group member may be referred to and claimed individually or in any combination with other members of the group or other elements found herein. It is anticipated that one or more members of a group may be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

51432-75

[0048] Certain embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Of course, variations on these described embodiments will become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventor expects skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

[0049] Furthermore, numerous references have been made to patents and printed publications throughout this specification.

[0050] In closing, it is to be understood that the embodiments of the invention disclosed herein are illustrative of the principles of the present invention. Other modifications that may be employed are within the scope of the invention. Thus, by way of example, but not of limitation, alternative configurations of the present invention may be utilized in accordance with the teachings herein. Accordingly, the present invention is not limited to that precisely as shown and described.


CLAIMS:

1. Use of an endothelin B (ET_B) receptor agonist in combination with a chemotherapeutic agent for the treatment of an ovarian tumor wherein the ET_B receptor agonist is IRL-1620, wherein said chemotherapeutic agent is cisplatin,
5 cyclophosphamide, or a combination thereof.
2. The use according to claim 1, wherein said chemotherapeutic agent is cisplatin.
3. The use according to claim 1, wherein said IRL-1620 selectively increases blood supply to said ovarian tumor.
- 10 4. The use according to claim 3, wherein said increase in said blood supply to said ovarian tumor increases the delivery of said chemotherapeutic agent to said ovarian tumor.
5. The use according to claim 1, wherein said IRL-1620 and said chemotherapeutic agent are for administration substantially simultaneously.
- 15 6. The use according to claim 5, wherein said IRL-1620 and said chemotherapeutic agent are for administration as a single composition.
7. The use according to claim 1, wherein said IRL-1620 and said chemotherapeutic agent are for administration sequentially.
8. The use according to claim 7, wherein said chemotherapeutic agent is
20 for administration prior to said IRL-1620.
9. The use according to claim 7, wherein said IRL-1620 is for administration prior to said chemotherapeutic agent.
10. The use according to claim 1, wherein said chemotherapeutic agent is cyclophosphamide.

11. A composition for use in the treatment of an ovarian cancer comprising at least one chemotherapeutic agent, an IRL-1620, and an optional excipient, wherein said chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.
- 5 12. A kit comprising a composition comprising IRL-1620 and an excipient and instructional information directing the administration of said composition comprising said IRL-1620 in combination with a chemotherapeutic agent to treat an ovarian tumor, wherein the kit further comprises a composition comprising a chemotherapeutic agent and an excipient, wherein said chemotherapeutic agent is
10 cisplatin, cyclophosphamide, or a combination thereof.
13. The kit according to claim 12, wherein said IRL-1620 and said chemotherapeutic agent are part of the same composition, are provided as separate compositions, or both.
14. The kit according to claim 12, wherein said chemotherapeutic agent is
15 cisplatin.
15. The kit according to claim 12, wherein said chemotherapeutic agent is cyclophosphamide.
16. Use of an endothelin B (ET_B) receptor agonist in the manufacture of a medicament to be used in conjunction with a chemotherapeutic agent for the
20 treatment of an ovarian tumor wherein said (ET_B) receptor agonist is IRL-1620, wherein said chemotherapeutic agent is cisplatin, cyclophosphamide, or a combination thereof.
17. The use according to claim 16, wherein said chemotherapeutic agent is cisplatin.
- 25 18. The use according to claim 16, wherein said chemotherapeutic agent is cyclophosphamide.


1/4

FIG. 1



2/4


FIG. 2



3/4


FIG. 3

Effect of IRL 1620 on the efficacy of cyclophosphamide (CYP)
in ovarian tumor bearing athymic mice



4/4

FIG. 4



**Effect of IRL-1620 on the efficacy of cisplatin (CIS)
in ovarian tumor bearing athymic mice**

