



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2799270 C 2013/10/08

(11)(21) **2 799 270**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) Date de dépôt/Filing Date: 2008/12/12
(41) Mise à la disp. pub./Open to Public Insp.: 2009/06/18
(45) Date de délivrance/Issue Date: 2013/10/08
(62) Demande originale/Original Application: 2 706 142
(30) Priorité/Priority: 2007/12/12 (US PCT/US2007/087145)

(51) Cl.Int./Int.Cl. *A61K 8/26* (2006.01),
A61K 8/28 (2006.01), *A61K 8/44* (2006.01),
A61Q 15/00 (2006.01)
(72) Inventeur/Inventor:
PAN, LONG, US
(73) Propriétaire/Owner:
COLGATE-PALMOLIVE COMPANY, US
(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSITIONS D'ACTIF ANTISUDORIFIQUE PRESENTANT UN CHROMATOGRAMME SEC
PRESENTANT UNE INTENSITE ELEVEE DE PIC 4 SEC
(54) Title: ANTIPERSPIRANT ACTIVE COMPOSITIONS HAVING SEC CHROMATOGRAM EXHIBITING HIGH SEC
PEAK 4 INTENSITY

(57) **Abrégé/Abstract:**

An antiperspirant active composition comprising an aluminum salt having an aluminum to chloride molar ratio of about 0.3:1 to about 3:1, exhibiting a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a Peak 4 intensity greater than a Peak 5 intensity in aqueous solution. The composition can optionally include zirconium. Also, methods and processes of making the same.



62301-2944D

Abstract

An antiperspirant active composition comprising an aluminum salt having an aluminum to chloride molar ratio of about 0.3:1 to about 3:1, exhibiting a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a Peak 4 intensity greater than a Peak 5 intensity in aqueous solution. The composition can optionally include zirconium. Also, methods and processes of making the same.

62301-2944D

TITLE

ANTIPERSPIRANT ACTIVE COMPOSITIONS HAVING SEC CHROMATOGRAM
EXHIBITING HIGH SEC PEAK 4 INTENSITY

5 This is a divisional application of Canadian Patent Application No. 2,706,142
filed on December 12, 2008. It should be understood that the expression “present invention”,
or the like, encompasses the subject matters of both this divisional application and the parent
application.

62301-2944D

BACKGROUND OF THE INVENTION

[0002] Antiperspirant salts, such as aluminum chlorohydrate (also called aluminum chlorohydrate polymeric salts and abbreviated here as "ACH") and aluminum zirconium glycine salts (abbreviated here as "ZAG", "ZAG complexes" or "AZG"), are known to contain a variety of polymeric and oligomeric species with molecular weights (MW) of 100-500,000. It has been clinically shown that, in general, the smaller the species, the higher the efficacy for reducing sweat.

[0003] In an attempt to increase the quality and quantity of smaller aluminum and/or zirconium species, a number of efforts have focused on: (1) how to select the components of ACH and ZAG that affect the performance of these materials as antiperspirants; and (2) how to manipulate these components to obtain and/or maintain the presence of smaller types of these components. These attempts have included the development of analytical techniques to identify the components. Size exclusion chromatography ("SEC") or gel permeation chromatography ("GPC") are methods frequently used for obtaining information on polymer distribution in antiperspirant salt solutions. With appropriate chromatographic columns, generally five distinctive groups of polymer species can be detected in commercial ACH and ZAG complexes appearing in a chromatogram as peaks 1, 2, 3, 4 and a peak known as "5,6". Peak 1 is the larger Zr species (greater than 60 Angstroms). Peaks 2 and 3 are larger aluminum species. Peak 4 is smaller aluminum species (aluminum oligomers, or small aluminum cluster) and has been correlated with enhanced efficacy for both Al and Al/Zr salts. Peak 5, 6 is the smallest aluminum species. Various analytical approaches for characterizing the peaks of ACH and various types of ZAG actives are found in "Antiperspirant Actives--Enhanced Efficacy Aluminum-Zirconium-Glycine (AZG) Salts" by Dr. Allan H. Rosenberg (Cosmetics and Toiletries Worldwide, Fondots, D. C. ed., Hartfordshire, UK: Aston Publishing Group, 1993, pages 252, 254-256).

[0004] Attempts to activate antiperspirant salts to produce materials having improved efficacy have included developing processes for obtaining composition having large amounts

62301-2944D(S)

of Peak 4 species. None of these efforts, however, have resulted in an antiperspirant composition having a composition with little or no Peak 3 and optionally little or no Peak 5.

BRIEF SUMMARY OF THE INVENTION

[0005] The present invention provides for an antiperspirant active composition comprising an aluminum salt having an aluminum to chloride molar ratio of about 0.3:1 to about 3:1, exhibiting a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a Peak 4 intensity greater than a Peak 5 intensity in aqueous solution, and optionally including zirconium.

[0006] The present invention also provides for a method of making an antiperspirant active composition. In one aspect, the method relates to making an antiperspirant active that exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a Peak 4 intensity greater than a Peak 5 intensity in aqueous solution comprising:

I) heating an aqueous solution containing an aluminum salt having an aluminum to chloride molar ratio of about 0.3:1 to about 3:1, optionally with a buffer agent, at a temperature of about 50°C to about 95°C to reflux for a period of time of about 1 hour to about 5 hours to obtain an aluminum salt solution;

II) adding an aqueous solution of an inorganic base to obtain an aluminum salt solution having an OH:Al molar ratio of about 2:1 to about 2.6:1 to obtain a pH adjusted aluminum salt solution having a pH of about 2 to about 5; and

III) optionally adding an aqueous solution containing a zirconium compound to the pH adjusted aluminum salt solution to thereby obtain an aluminum-zirconium salt solution having a molar ratio of aluminum to zirconium of about 5:1 to about 10:1.

In another related aspect, the invention relates to a method of making an antiperspirant active composition comprising the following sequence of steps:

(i) heating an aqueous solution containing an aluminum chloride compound having an aluminum to chloride molar ratio of 0.3:1 to 3:1, optionally with a buffer agent, at a

62301-2944D(S)

temperature of 50°C to 95°C to reflux for a period of time of 1 hour to 5 hours to obtain an aluminum salt solution;

(ii) after the heating step, adding an aqueous solution of an inorganic base to obtain an aluminum salt solution having an OH:Al molar ratio of 2:1 to 2.6:1 to obtain a pH
5 adjusted aluminum salt solution having a pH of 2 to 5; and

(iii) optionally adding an aqueous solution containing a zirconium compound to the pH adjusted aluminum salt solution to thereby obtain an aluminum-zirconium salt solution having a molar ratio of aluminum to zirconium of 5:1 to 10:1.

BRIEF DESCRIPTION OF THE DRAWINGS

10 [0007] The accompanying drawings, which are included to provide further understanding of the disclosure and are incorporated in and constitute a part of this specification, illustrate embodiments of the disclosure and, together with the description, serve to explain the principles of the disclosure.

[0008] Figure 1 illustrates an SEC chromatogram of a prior art antiperspirant active
15 composition.

[0009] Figure 2 illustrates an SEC chromatogram having exclusive peaks 4 and 5 for an inventive product, Example 1, of the present invention.

[0010] Figure 3 illustrates an SEC chromatogram having exclusive peak 4 for an inventive product, Example 2, of the present invention.

[0011] **Figure 4** illustrates an SEC chromatogram having exclusive peaks 4 and 5 for an inventive product, Example 3, of the present invention.

[0012] **Figure 5** illustrates an SEC chromatogram having exclusive peaks 4 and 5 for an inventive product, Example 4, of the present invention.

[0013] **Figure 6** illustrates an SEC chromatogram having exclusive peak 4 for an inventive product, Example 5, of the present invention.

[0014] **Figure 7** illustrates an SEC chromatogram having exclusive peaks 4 and 5 for an inventive product, Example 6, of the present invention.

[0015] **Figure 8** illustrates an SEC chromatogram of an inventive product from an ACH batch scale-up, Example 7, of the present invention.

[0016] **Figure 9** illustrates an SEC chromatogram for inventive products, Examples 8-10, of the present invention produced within an optimal reaction temperature range.

[0017] **Figure 10** illustrates an SEC chromatogram for inventive products, Examples 8, 10, and 11, of the present invention produced within an optimal reaction time range.

[0018] **Figure 11** illustrates an SEC chromatogram for an inventive product, Example 12, of the present invention produced within an optimal reaction time range.

[0019] **Figure 12** illustrates an SEC chromatogram for inventive products, Examples 13-15, of the present invention produced within an optimal Ca(OH)_2 :glycine molar ratio range.

[0020] **Figure 13** illustrates an SEC chromatogram for inventive products, Examples 16 and 17, of the present invention produced at different revolutions per minute.

[0021] **Figure 14** illustrates an SEC chromatogram for inventive products, Examples 18 and 19, of the present invention produced within an optimal revolutions per minute range.

[0022] **Figure 15** illustrates an SEC chromatogram for inventive products, Examples 20 and 21, of the present invention produced by different methods of adding Ca(OH)_2 .

DETAILED DESCRIPTION OF THE INVENTION

[0023] As used throughout, ranges are used as a shorthand for describing each and every value that is within the range. Any value within the range can be selected as the terminus of the range.

[0024] The present invention is directed to an antiperspirant active composition having a high SEC peak 4 in aqueous solution. The composition is obtained by a stepwise procedure to neutralize aluminum chloride in a solution (optionally buffered) using inorganic bases. In some embodiments, the antiperspirant active compositions obtained by this stepwise procedure include aluminum salts having an aluminum to chloride molar ratio of about 0.3:1 to about 3:1, the aluminum salt has a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a

Peak 4 intensity greater than a Peak 5 intensity in aqueous solution. The composition may optionally include zirconium.

[0025] Optionally, a buffer can be included. Buffers that can be used can be chosen from amino acids, glycine, and betaine. The buffer to aluminum molar ratio in certain embodiments can be about 0.1:1 to about 3:1. In another embodiment, the buffer to aluminum molar ratio is about 0.5:1 to about 2:1. In another embodiment, the buffer to aluminum molar ratio is about 1:1 to about 1.5:1.

[0026] The compositions may be made in a variety of ways involving a stepwise procedure to neutralize aluminum chloride in solution (optionally buffered) using inorganic basic salts. The procedure generally includes the step of heating an aqueous solution containing an aluminum chloride compound (optionally with a buffer agent) at a temperature of about 50°C to about 95°C to reflux for a period of time of about 1 hour to about 5 hours. In one such embodiment, an aqueous solution containing an aluminum chloride compound is heated at a temperature of about 75°C to about 95°C to reflux for a period of time of about 3 hours to about 4 hours. In another such embodiment, an aqueous solution containing an aluminum chloride compound and a buffer agent is heated at a temperature of about 75°C to about 95°C to reflux for a period of time of about 3 hours to about 4 hours. In one embodiment, the temperature is about 85°C.

[0027] In some embodiments, the solution has a buffer agent to aluminum molar ratio of about 0.1:1 to about 3:1. To adjust the pH of the aluminum salt solution, an aqueous solution of an inorganic base is added to the heated solution to thereby obtain a pH adjusted aluminum salt solution having a hydroxide to aluminum molar ratio of about 1:1 to about 4:1, and a pH of about 2 to about 5. In one such embodiment, the hydroxide to aluminum molar ratio of about 2:1 to about 3:1. In another such embodiment, the hydroxide to aluminum molar ratio is about 2.1:1 to about 2.6:1.

[0028] In some embodiments, a zirconium salt may also be added to the pH adjusted aluminum salt solution. In one other such embodiment, the molar ratio of Al:Zr is about 5:1 to about 10:1. The antiperspirant active composition has a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a Peak 4 intensity greater than a Peak 5 intensity in aqueous solution.

[0029] In one embodiment, an aqueous aluminum chloride salt solution is buffered with betaine monohydrate and held at about 50°C to about 95°C to reflux for a period time of about 1 to about 6 hours. To the heated solution, an aqueous solution of an inorganic base is added dropwise over a period of time of about 1 to about 3 hours while maintaining the aluminum-betaine solution at about 50°C to about 95°C to reflux. In one such embodiment, the solution

has a betaine to aluminum molar ratio of about 1.1. In another such embodiment, the solution has a betaine to aluminum molar ratio of about 1.25.

[0030] In one embodiment, an aqueous solution containing an aluminum chloride compound is buffered with betaine monohydrate and held at about 75°C to about 95°C to reflux for a period of time of about 3 hours to about 4 hours. In another such embodiment, an aqueous solution of an inorganic base is added dropwise over a period of time of about 1 to about 3 hours while maintaining the aluminum-betaine solution at about 75°C to about 95°C to reflux. In another embodiment, an aqueous solution of an inorganic base is added over a period of time in a series of additions while maintaining the aluminum-betaine solution at about 75°C to about 95°C to reflux. In one such embodiment, the inorganic base is added in at least 3 additions. In another such embodiment, the inorganic base is added in at least 5 additions. In another embodiment, a ZrOCl_2 solution is added to the pH adjusted aluminum-betaine solution. In one such embodiment, the molar ratio of Al:Zr is about 8. In another such embodiment, the molar ratio of Al:Zr is about 7. In one other such embodiment, the molar ratio of Al:Zr is about 9.

[0031] In another embodiment, an aqueous aluminum chloride solution is buffered with glycine and held at about 50°C to about 95°C to reflux for a period time of about 1 to about 6 hours. To the heated solution, an aqueous solution of an inorganic base is added dropwise over a period of time of about 1 to about 3 hours while maintaining the aluminum-glycine solution at about 50°C to about 95°C to reflux. In one such embodiment, the solution has an aluminum to glycine molar ratio of about 0.4. In another such embodiment, the solution has an aluminum to glycine molar ratio of about 0.8.

[0032] In another embodiment, an aqueous solution containing an aluminum chloride compound is buffered with glycine and held at about 75°C to about 95°C to reflux for a period of time of about 3 hours to about 4 hours. In another such embodiment, an aqueous solution of an inorganic base is added dropwise over a period of time of about 1 to about 3 hours while maintaining the aluminum-glycine solution at about 75°C to about 95°C to reflux. In another embodiment, an aqueous solution of an inorganic base is added over a period of time in a series of additions while maintaining the aluminum-glycine solution at about 75°C to about 95°C to reflux. In one such embodiment, the inorganic base is added in at least 3 additions. In another such embodiment, the inorganic base is added in at least 5 additions. In one embodiment, the inorganic base is calcium hydroxide. In one such embodiment, the addition of calcium hydroxide provides an aqueous solution having a Ca(OH)_2 :glycine molar ratio of about 1.25:1 to about 1:1.

[0033] In another embodiment, a ZrOCl_2 solution is added to the pH adjusted aluminum-glycine solution. In one such embodiment, the molar ratio of Al:Zr is about 8. In another such embodiment, the molar ratio of Al:Zr is about 7. In one other such embodiment, the molar ratio of Al:Zr is about 9.

[0034] For the above methods, the aluminum chloride salt and inorganic base may be obtained from a variety of sources. In one embodiment, the aluminum chloride salt includes aluminum trichloride, aluminum chlorohexahydrate and aluminum dichlorohydrate. In one such embodiment, the aluminum chloride salt is aluminum chlorohexahydrate.

[0035] In one embodiment, the inorganic base can be at least one base chosen from metal hydroxides, calcium hydroxide, strontium hydroxide, sodium hydroxide, barium hydroxide, metal oxides, calcium oxide, strontium oxide, and barium oxide.

[0036] The present invention provides for aluminum antiperspirant active compositions and/or aluminum-zirconium antiperspirant active compositions having high levels of low molecular weight Al and Zr species. As illustrated in **Figures 2 to 7**, the high levels of low molecular weight Al and Zr species is reflected in a SEC trace that has an intense Peak 4, low Peaks 1, 2, 3 and 5. The polymerization of the antiperspirant actives in aqueous solutions and the correspondent gelation process were followed by monitoring the molecular weight profile of the polyoxohalides in time by SEC. The relative retention time ("Kd") for each of these peaks varies depending on the experimental conditions, but the peaks remain relative to each other. Data for Tables in the examples was obtained using an SEC chromatogram using the following parameters: Waters®600 analytical pump and controller, Rheodyne® 7725I injector, Protein-Pak® 125 (Waters) column, Waters 2414 Refractive Index Detector. 5.56mM nitric acid mobile phase, 0.50ml/min flow rate, 2.0 microliter injection volume. Data was analyzed using Water® Empower software (Waters Corporation, Milford, Mass.). The concentration of the antiperspirant in solution does not affect the retention time in the machine.

[0037] The design of modern AP salts aims at actives with high levels of low molecular weight Al and Zr species, which is reflected in a SEC trace that has intense Peak 4 and low Peaks 1, 2, and 3. Throughout the present study, the levels of the species corresponding to these peaks are estimated based on the following ratios (or percentages):

$$f_{Pi} = \frac{Pi}{\sum Pj} \quad i = 1, 2, 3, 4, 5; \quad j = 2, 3, 4, 5$$

[0038] where f_{p_i} is the fraction of peak i , and P_i or P_j are the intensity of peaks P_i or P_j , respectively. The amount of low molecular weight Al species will be correlated with the fraction, f_{p_4} , or percentage, $f_{p_4} \times 100$, of SEC-Peak 4. In brief, a preferred antiperspirant salt would have a very low f_{p_1} , f_{p_2} , f_{p_3} , and/or f_{p_5} , and a high f_{p_4} .

[0039] In certain embodiments, the ratio of Peak 4 to Peak 3 is at least 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, or any number up to infinity.

[0040] In one embodiment, an aluminum salt and/or aluminum-zirconium salt, in aqueous solution, exhibit a SEC profile wherein the SEC Peak 4 to Peak 3 intensity ratio is at least 7. In such embodiments, the percentage of SEC Peak 4 of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram is: at least 50%; at least 60%; at least 70%; at least 80%; at least 90%, or 95 to 100%. In another such embodiment, the SEC Peak 4 area is 100%.

[0041] In another embodiment, the aluminum salt and/or the aluminum-zirconium salt, in aqueous solution, exhibits a SEC profile wherein the SEC Peak 4 to Peak 3 intensity ratio is at least 7 and exhibits low percentage of SEC Peak 3. In such embodiments, the composition has the percentage of SEC Peak 3 area of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram is: less than about 10 %; less than about 5 %; less than about 2 %; less than about 1 %; less than about 0.9 %; less than about 0.8 %; less than about 0.7 %; less than about 0.6 %; of less than about 0.5 %; less than about 0.4 %; less than about 0.3 %; less than about 0.2 %; or less than about 0.1 %. In another such embodiment, the composition has no SEC Peak 3 area.

[0042] In another embodiment, the aluminum salt and/or the aluminum-zirconium salt, in aqueous solution, exhibits a SEC profile wherein the SEC Peak 4 to Peak 3 intensity ratio is at least 7 and exhibits low percentages of SEC Peak 5. In such embodiments, the percentage of SEC Peak 5 area of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram is: less than about 30 %; less than about 20 %; less than about 10 %; less than about 5 %; or less than about 1 %. In another such embodiment, the composition has no SEC Peak 5 area.

[0043] In another embodiment, the aluminum salt and/or the aluminum-zirconium salt, in aqueous solution, exhibits a SEC profile wherein the SEC Peak 4 to Peak 3 ratio is at least 7, and exhibits a low percentage of SEC Peak 1 and a low percentage of SEC Peak 2. In such embodiment, the percentage of SEC Peak 1 area of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram is: less than about 10 %; a SEC Peak 1 area less than about 5 %; less than about 2 %; less than about 1 %; less than about 0.9 %; less than about 0.8 %; of less than about 0.7 %; less than about 0.6 %; less than about 0.5 %; less than about 0.4 %; less than

about 0.3 %; less than about 0.2 %; or less than about 0.1 %. In another embodiment, the complex has no SEC Peak 1 area. In another embodiment, the percentage of SEC Peak 2 area of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram is: less than about 10 %; less than about 5 %; less than about 2 %; less than about 1 %; less than about 0.9 %; less than about 0.8 %; less than about 0.7 %; less than about 0.6 %; less than about 0.5 %; less than about 0.4 %; less than about 0.3 %; less than about 0.2 %; or less than about 0.1 %. In another embodiment, the composition has no SEC Peak 2 area.

[0044] The aluminum antiperspirant active compositions and/or aluminum-zirconium antiperspirant active compositions may be used in a variety of antiperspirant products. If the product is used as a solid powder, the size of the particles of antiperspirant active of the invention can be any desired size, and may include conventional sizes such as in the range of 2 to 100 microns, with selected grades having an average particle size of 30-40 microns; finer sized grades having an average particle size distribution of 2-10 microns with an average size of about 7 microns as made by a suitable dry-grinding method; and micronized grades having an average particle size of less than about or equal to 2 microns, or less than about or equal to 1.5 microns.

[0045] The compositions of this invention may be used to formulate antiperspirants having improved efficacy. Such antiperspirants include solids such as sticks and creams (creams sometimes being included in the term "soft solid"), gels, liquids (such as are suitable for roll-on products), and aerosols. The forms of these products may be suspensions or emulsions. These antiperspirant actives can be used as the antiperspirant active in any antiperspirant composition.

[0046] Examples of Suitable Formulations

[0047] Sticks

Stick products may be made with conventional gelling agents such as stearyl alcohol and dibenzylidene sorbitol. A sample formulation is as follows:

40-55% (particularly 45%);
cyclomethicone (especially D5 cyclomethicone);
20-30% (particularly 21%);
stearyl alcohol 7-15% (particularly 10%);
talc 15-22% (particularly 22%);
antiperspirant active of the invention in particle form; and
1-3% (particularly 2%) fragrance.

[0048] Roll Ons

Roll Ons having a sample formulation:

45-65% (particularly 55%) cyclomethicone (especially D5 cyclomethicone);
 0.1-10% (particularly 3%) cyclomethicone/dimethicone copolyol (such as Dow Corning 2-5185C) 10-25% (particularly 20%);
 antiperspirant active of the invention in solution form (25-45% actives on an anhydrous basis in water);
 5-30% (particularly 20%) water; and
 1-3% (particularly 2%) fragrance.

[0049] Soft solids

Soft solids may be made with formulations described in U.S. Patent No. 6,682,749. A sample formulation is as follows:

40-70% (particularly 50%) elastomer in cyclomethicone (KSG-15 from Shin-Etsu);
 5-15% (particularly 6%) polyethylene (for example, beads having a density in the range of 0.91-0.98 g/cm³ and an average particle size in the range of 5-40 microns);
 10-20% (particularly 15%) C12-15 alkylbenzoate (FINSOLV™ TN from Finetex);
 0.1-25%% (particularly 22%) antiperspirant active of the invention in powder form;
 1-15% (particularly 5%) dimethicone (particularly with a viscosity of 100 centistokes); and
 1-3% (particularly 2%) fragrance.

[0050] Gels

Gels may be made with a variety of formulations such as:

5-50% (particularly 29%) cyclomethicone (particularly D5);
 0.1-10% (particularly 3%) cyclomethicone/dimethicone copolyol (such as Dow Corning 2-5185C);
 0-10% (particularly 5%) hydrogenated polyisobutene 250;
 0-10% (particularly 5%) C12-15 alkylbenzoate (FINSOLV™ TN from Finetex);
 0-10% (particularly 5%) dimethicone (particularly with a viscosity of 100 centistokes);
 0.1-25% (particularly 20%) antiperspirant active of the invention in powder form or
 10-25% (particularly 20%) of active in solution (25-45% actives on an anhydrous basis);
 5-50% (particularly 30%) water; and
 1-3% (particularly 2%) fragrance.

[0051] Note that in the explanation of the invention, where water is listed it is intended to count the contribution of the water present in the antiperspirant solution as part of the overall water content. Thus, water is sometimes listed as part of the actives solution or sometimes listed separately.

[0052] In one embodiment the refractive indices of the external and internal phases are matched within 0.005 to obtain a clear product.

[0053] Other Formulations of Interest

[0054] Formulation A

0.5-2.5% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
55-65% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
1-10% PPG-3 myristyl ether;
10-25% antiperspirant active of the invention;
10-25% water; and
0.5-1.5% fragrance.

[0055] Formulation B

1.0-3.0% dimethicone copolyol (for example, Dow Corning 2-5185C (48%))
40-60% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
1-5% cyclomethicone (in addition to that found in the elastomer);
4-12% PPG-3 myristyl ether;
15-30% antiperspirant active of the invention;
15-35% water; and
0.5-1.5% fragrance.

[0056] Formulation C

1.0-3.0% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
1-10% hydrogenated polyisobutene (for example, Fancol™. Polyiso 250);
40-55% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
3-8% PPG-3 myristyl ether;
15-20% antiperspirant active of the invention;

WO 2009/076591

PCT/US2008/086556

20-30% water; and
1.0-3.0% fragrance.

[0057] Formulation D

1.0-3.0% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
40-60% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
3-8% PPG-3 myristyl ether;
15-30% antiperspirant active of the invention;
15-30% water;
0.5-1.5% fragrance; and
1-10% diethylhexyl naphthalate

[0058] Formulation E

0.5-2.5% dimethicone copolyol (for example, Dow Corning 2-5185C (48%));
60-70% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
7-10% antiperspirant active of the invention;
25-35% water;
1-10% methylpropylene diol (MPDiol) ; and
0.5-1.5% fragrance

[0059] Formulation F

1.0-3.0% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
6-10% hydrogenated polyisobutene (for example, FANCOL™ Polyiso 250);
35-45% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
6-10% PPG-3 myristyl ether;
40-50% antiperspirant active of the invention as 43% active in water no additional water; and
0.5-1.0% fragrance.

[0060] Formulation G

0.1-0.6% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
4-7% hydrogenated polyisobutene (for example, FANCOL™ Polyiso 250);
40-50% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
4-7% PPG-3 myristyl ether;
40-50% antiperspirant active of the invention as 43% active in water no additional water; and
0.5-1.0% fragrance.

[0061] Formulation H

0.5-2.0% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
1-7% hydrogenated polyisobutene (for example, FANCOL™ Polyiso 250);
40-50% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
45-55% antiperspirant active as 43% active of the invention in water no additional water; and
0.5-1.5% fragrance.

[0062] Formulation I

2-7% dimethicone copolyol (for example, Dow Corning 2-5185 C (48%));
0.1-1% Oleath-20 1-5% C12-15 alkyl benzoate (FINSOLV™ TN);
15-25% elastomer in cyclomethicone (for example, DC-9040 from Dow Corning Corporation (Midland, Mich.) or KSG-15 from Shin-Etsu Silicones of America (Akron, Ohio));
15-25% antiperspirant active of the present invention;
15-30% water; and
0.5-1.5% fragrance

[0063] The cosmetic composition according to the present invention can be packaged in conventional containers, using conventional techniques. Where a gel, cream or soft-solid cosmetic composition is produced, the composition can be introduced into a dispensing package (for example, conventional packages for gels with glide on applicators, jars where the gel or cream is applied by hand, and newer style packages having a top surface with pores) as conventionally done in the art. Thereafter, the product can be dispensed from the

dispensing package as conventionally done in the art, to deposit the active material, for example, on the skin. For sticks, sprays, aerosols and roll-ons the compositions can be placed in a conventional types of container (with the inclusion of propellants in aerosols). This provides good deposition of the active material on the skin.

[0064] Compositions of the present invention can be formulated as clear, translucent or opaque products. A desired feature of the present invention is that a clear, or transparent, cosmetic composition, (for example, a clear or transparent deodorant or antiperspirant composition) can be provided. The term clear or transparent according to the present invention is intended to connote its usual dictionary definition; thus, a clear liquid or gel antiperspirant composition of the present invention allows ready viewing of objects behind it. By contrast, a translucent composition, although allowing light to pass through, causes the light to be scattered so that it will be impossible to see clearly objects behind the translucent composition. An opaque composition does not allow light to pass there through. Within the context of the present invention, a gel or stick is deemed to be transparent or clear if the maximum transmittance of light of any wavelength in the range 400-800 nm through a sample 1 cm thick is at least 35%, or at least 50%. The gel or liquid is deemed translucent if the maximum transmittance of such light through the sample is between 2% and less than about 35%. A gel or liquid is deemed opaque if the maximum transmittance of light is less than about 2%. The transmittance can be measured by placing a sample of the aforementioned thickness into a light beam of a spectrophotometer whose working range includes the visible spectrum, such as a Bausch & Lomb Spectronic 88 Spectrophotometer. As to this definition of clear, see European Patent Application Publication No. 291,334 A2. Thus, according to the present invention, there are differences between transparent (clear), translucent and opaque compositions.

EXAMPLES

[0065] Comparative Examples

[0066] A 0.72 M $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (18 mmol) is held at 90°C and stirred vigorously. To this solution, a 4 N $\text{Ca}(\text{OH})_2$ (20 mmol) is added dropwise over a 1 hour 30 minute period. A ratio of OH:Al of 2.22 is employed in an attempt to prevent the formation of larger unwanted Al species. The pH after the reaction was 2.36 due to the low OH: Al ratio. The SEC chromatogram, illustrated in **Figure 1**, exhibits multiple peaks including, SEC-Peak 4, and SEC-Peak 5 indicating multiple Al species are present in solution. At a retention time of

approximately 15.5 minutes, SEC-Peak 3 is observed due to no buffer (i.e., betaine or glycine) as control.

[0067] Also for comparison, 10% solutions are prepared from commercially available antiperspirants. The solutions are prepared by adding 1g of antiperspirant to 9g of water and mixing. The antiperspirant salts were Reach™ 103, Reach™ 301 from Reheis, and Summit™ Z576 from Summit Research Labs.

[0068] Example 1

[0069] A 0.72M $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (18 mmol) is buffered with 20 mmol betaine monohydrate, held at 90°C, and stirred vigorously. To this solution, a 4 N $\text{Ca}(\text{OH})_2$ (20 mmol) is added dropwise over a 1 hour 30 minute period. A ratio of OH:Al of 2.22 is employed in an attempt to prevent the formation of large Al species. The pH after the reaction is 2.56 due the low OH:Al ratio. As illustrated in **Figure 2**, the SEC chromatogram shows exclusively SEC-Peak 4 and SEC-Peak 5, which are known to represent active anti-perspirant species. Substantially no SEC-Peak 3 species is observed at a retention time of approximately 15.5 minutes.

[0070] Example 2

[0071] A 0.72M $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (16.26 mmol) was buffered with 20 mmol anhydrous betaine, held at 90°C, and stirred vigorously. To this solution, a 4 N $\text{Ca}(\text{OH})_2$ (20mmol) was added dropwise over a 2 hour period. A ratio of OH:Al of 2.46 was employed in an attempt increase the final pH and to reduce SEC-Peak 5 species. Because a higher OH:Al ratio was used, the addition of the base was extended over a 2 hour period. The pH after the reaction was 4.8. As illustrated in **Figure 3**, the SEC chromatogram indicated that the solution contained exclusively SEC-Peak 4 antiperspirant active species. Substantially no SEC-Peak 3 species was observed at a retention time of approximately 15.5 minutes.

[0072] Example 3

[0073] A small portion of the solution from Example 2 is taken to determine the effects of Zr on the peak distribution. $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$ is added to achieve a molar ratio of 8:1, Al:Zr. The pH after adding Zr reduces to 3.7. As shown in **Figure 4**, the SEC chromatogram of this zirconium, aluminum solution shows two notable features. Firstly, the SEC-Peak 4 remains the predominate peak, and the SEC-Peak 5 intensity increased to 1% - as expected by the reduced pH. Secondly, the SEC chromatogram does not show any peaks with retention time of 12.5 minute indicating the absence of undesirable Zr polymer species. The absence of this SEC-Peak indicates that the pure SEC-Peak 4 solution of Example 2 did not promote the

WO 2009/076591

PCT/US2008/086556

aggregation of Zr into larger, less efficacious species. Also substantially no SEC-Peak 3 species is observed at a retention time of approximately 15.5 minutes.

[0074] Example 4

[0075] A 0.5M $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (25 mmol) is buffered with 31.25 mmol glycine, held at 95°C, and stirred vigorously. To this buffered solution, a 1.0 N $\text{Ca}(\text{OH})_2$ (31.25 mmol) is added dropwise over a 1 hour period. A ratio of OH:Al of 2.5 is employed in an attempt to increase the final pH and to reduce SEC-Peak 5 species. The pH after the reaction is 4.52. The SEC chromatogram shown in **Figure 5** exhibits primarily SEC-Peak 4 and a smaller SEC-Peak 5 (520). Substantially no SEC-Peak 3 species is observed at a retention time of approximately 15.5 minutes.

[0076] Example 5

[0077] A 0.5M $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (25 mmol) is buffered with 62.5 mmol glycine, held at 95°C, and stirred vigorously. To this buffered solution a 1.0 N $\text{Ca}(\text{OH})_2$ (31.25 mmol) is added dropwise over a 1 hour period. A ratio of OH:Al of 2.5 is employed in an attempt to increase the final pH and to reduce the SEC-Peak 5 species. The pH after the reaction is 4.52. The SEC chromatogram shown in **Figure 6** exhibits exclusively SEC-Peak 4 and no SEC-Peak 5. Substantially no SEC-Peak 3 species is observed at a retention time of approximately 15.5 minutes.

[0078] Example 6

[0079] A small portion of the solution from Example 5 is taken to determine the effects of Zr on the peak distribution. $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$ is added to achieve an Al:Zr molar ratio of 8:1. The pH after adding Zr reduces to 3.3. The SEC chromatogram shown in **Figure 7** exhibits primarily SEC-Peak 4 and substantially no SEC-Peak 5 (720). This data indicates that the pure SEC-Peak 4 solution of Example 5 does not promote the aggregation of Zr into larger, less efficacious species. Also substantially no SEC-Peak 3 species is observed at a retention time of approximately 15.5 minutes.

Table 1. Comparison of the Examples

Example	Ratio OH:Al	pH	Comparable ACH Solution	Relative Peak Distribution after Reaction (%)				
				Peak 2	Peak 3	Peak 4	Peak 5	Peak 4/ Peak 3
Summit™ Z576				3.1	34.1	40	22.6	1.2
Reach™ 103	n/a		10% ACH		63	34	3	0.54
Reach™ 301	n/a		10% ACH	7	65	12	16	0.18
Comparative	2.2	2.36	21% ACH	0	42	42	16	1.0
Example 1	2.2	2.56	21% ACH	0	0	75	25	∞
Example 2	2.5	4.8	22% ACH	0	0	100	0	∞
Example 3	2.5	3.7	22% ACH	0	0	99	1	∞
Example 4	2.5	4.5	4% ACH	0	0	93	7	∞
Example 5	2.5	4.52	4% ACH	0	0	100	0	∞
Example 6		3.32	4% ACH	0	0	98	2	∞

[0080] Large Scale Production of Antiperspirant Active Compositions**[0081] Example 7****[0082] Process:**

1. Aluminum chloride hexahydrate (3.1055 kg) and anhydrous glycine (1.1863 kg) were combined in a 316 stainless steel, 25 gallon batch vessel provided with two level hydrofoil blade agitation using medium (60-80 rpm) agitation. Distilled water (19.8192 kg) was added to the mixture and the solution was heated to 85°C with vigorous (100-125 rpm) agitation. Temperature control was used to maintain effective temperature rate increase and target level during the trial. The heating source was a steam jacket provided with 3 bar/42 psig steam throughout the trial.
2. In a separate reaction vessel, calcium hydroxide (1.1725 kg) was dissolved in 4.7165 kg of distilled water.

3. When the aluminum chloride hexahydrate/glycine solution reached 85°C, the calcium hydroxide solution was added over a 1 hour 30 minute period. The reaction vessel was vigorously agitated throughout the addition and care was taken to ensure that the calcium hydroxide residue did not form at the top of the vessel.
4. After the addition of calcium hydroxide, the solution was held at 85°C under vigorous agitation for an additional three hours. The reaction yielded 30.18 kg (100.4%) of ACH.

[0083] Analysis:

[0084] SEC analysis showed that the scaled-up synthesis of ACH was successful in replicating laboratory results: the resultant active solution was void of Peaks 1-3 and contained a very small Peak 5. **Figure 8** shows the SEC profile of the product from the ACH batch scale-up in which Peak 4 elutes at 14.5 minutes in both SEC profiles. Visual analysis clearly indicates that Peaks 1-3 are absent from the ACH solution. There is also a strong Peak 4 and a minimal Peak 5. The results of Peak distribution are summarized in **Table 2** below.

Table 2. Comparison of Peak Distribution of the Scaled-up Batch (ACH) vs. an Activated ACH (Reach™ 103)

Solution	Relative Peak Distribution after Reaction (%)			Peak 4 / Peak 3	pH
	Peak 3	Peak 4	Peak 5		
Reach™ 103	61.00	35.7	3.3	0.585	4.07
Example 7	0	96.74	3.26	∞	3.89

[0085] Conclusion:

[0086] The result of scaled-up batching in the pilot shows its uniform aluminum species under Peak 4 can be obtained in contrast to un-removable larger species in current activated ACH (Reach™ 103). The process of the present invention can be successfully adopted into any large facility antiperspirant manufacturers.

[0087] Optimal Reaction Parameters for Manufacturing Process

[0088] Optimal Reaction Temperature

[0089] The reactions for the following Examples 8-10 were all performed with a Ca(OH)₂:glycine molar ratio of 1:1 and a OH:Al molar ratio of 2.46:1. In all three reactions, a 0.65 M AlCl₃·6H₂O (19 mmol) aqueous solution was buffered with 23 mmol glycine and heated to 90 °C with stirring. A Ca(OH)₂ suspension was added drop-wise by hand to the

aqueous aluminum chloride salt solution over 1 hour and 40 minutes, with a total reaction time of 3.5 to 4 hours. The SEC chromatogram in **Figure 9** illustrates large Peak 4 with negligible Peak 3 for Examples 8-10. There is little to no increase in Peak 3 formation upon lowering the reaction temperature from 90°C to 75°C. There is only a small increase of 3.9%-6.9% for Peak 5 upon lowering the reaction temperatures. A small Peak 5 will add to the long term stability and efficacy of the active product. Therefore, the optimal reaction temperature is between 75°C and 90°C. According to the SEC peak areas in **Figure 9**, the solutions of Examples 8-10 are approximately comparable to a ~5% ACH solution.

Table 3. Comparison of the Examples (75°C vs. 90°C)

Solution	Basic Source	Relative Peak Distribution after Reaction (%)			Comparable ACH (%)	Temperature (°C)
		Peak 3	Peak 4	Peak 5		
Example 8	Ca(OH) ₂	0	92.0	8.0	5.6	75
Example 9	Ca(OH) ₂	0.2	94.8	5.0	5.2	75
Example 10	Ca(OH) ₂	0	98.9	1.1	4.9	90

[0090] Optimal Reaction Time For Synthesis of ACH

[0091] The reactions for the following Examples 8, 10, and 11, were all performed with a Ca(OH)₂:glycine molar ratio of 1:1 and a OH:Al molar ratio of 2.46:1. The total reaction time for all three examples was 3-4 hours. The SEC chromatogram in **Figure 10** shows only a 0.2% increase in Peak 3 upon lowering the reaction time to 3 hours at 70°C. According to the SEC peak area, the solutions for Examples 8, 10, and 11 are comparable to a ~5% ACH solution. **Figure 10** further shows exclusively Peaks 4 and 5 in reactions performed at 75°C and 90°C for 4 hours (respectively, Examples 8 and 10).

[0092] The reaction for Example 12 was performed with a Ca(OH)_2 :glycine molar ratio of 1.25:1 and a OH:Al molar ratio of 2.46:1. The SEC chromatogram in **Figure 11** shows favorable results when the reaction was performed using optimal parameters and reacted at 75°C for 3 hours and 30 minutes. **Figure 11** also shows an increase in Peak 3 of 0.8% and an increase of 2.4% for Peak 5 upon lowering the reaction time to 3 hours and 30 minutes at 75°C. According to the SEC peak areas in **Figure 11**, the solution of Example 12 is approximately comparable to a ~5% ACH solution. Therefore, the optimal reaction time is between 3 and 4 hours.

Table 4. Comparison of Example Reaction Times

Solution	Basic Source	Relative Peak Distribution after Reaction (%)			Comparable ACH (%)	Temperature (°C)	Reaction Time (hrs)
		Peak 3	Peak 4	Peak 5			
Example 8	Ca(OH)_2	0	92.0	8.0	5.6	75	4
Example 10	Ca(OH)_2	0	98.9	1.1	4.9	90	4
Example 11	Ca(OH)_2	0.2	96.8	3.0	4.7	70	3
Example 12	Ca(OH)_2	0.8	96.7	2.5	5.2	75	3.5

[0093] Optimal Ca(OH)_2 :Glycine Molar Ratio

[0094] The reactions for the following Examples 13-15 were all performed using an OH:Al molar ratio of 2.46:1 and were reacted for 2-4 hours. For Examples 13-15, a 0.235 M $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (47 mmol) aqueous solution was buffered with 23 mmol glycine and heated to 90 °C with stirring. **Figure 12** and **Table 5** illustrate that the closer the Ca(OH)_2 :glycine molar ratio is to a 1:1 value, the smaller the relative distribution percentage of Peak 3. A Ca(OH)_2 :glycine molar ratio of 2.5:1 (Example 13) produced an increase in Peak 3 by 7.8% compared to the standard reaction (Example 10). A Ca(OH)_2 :glycine molar ratio of 2:1 (Example 14) was used, there was a 1.4% increase in Peak 3. When the Ca(OH)_2 :glycine molar ratio was lowered to 1.25:1 (Example 15), there was only a 0.4% increase in Peak 3. Using a Ca(OH)_2 :glycine molar ratio of 1.25:1 produces a high Peak 4/Peak 3 ratio. Example 12 illustrates that an increase of only 0.8% in Peak 3 when using a Ca(OH)_2 :glycine molar ratio of 1.25:1 at 75°C. Following the optimal reaction parameters, there was only a 0.4% increase in Peak 3 compared to Example 15 done at 90°C. Therefore, the optimal Ca(OH)_2 :glycine molar ratio is between 1.25:1 and 1:1.

Table 5. Comparison of Example Ca(OH)₂:Glycine Molar Ratios

Solution	Basic Source	Relative Peak Distribution after Reaction (%)			Comparable ACH (%)	Temperature (°C)	Ca(OH) ₂ :glycine
		Peak 3	Peak 4	Peak 5			
Example 12	Ca(OH) ₂	0.8	96.7	2.5	5.2	75	1.25:1
Example 13	Ca(OH) ₂	7.8	92.2	0	4.7	90	2.5:1
Example 14	Ca(OH) ₂	1.4	93.0	5.6	5.0	90	2:1
Example 15	Ca(OH) ₂	0.4	94.8	4.8	5.1	90	1.25:1

[0095] Optimal Revolutions Per Minute (RPM)

[0096] The ACH synthesis of Example 10 at 90°C over 4 hours used a magnetic stir bar for stirring. This method does not provide for a way to specifically control the revolutions per minute (rpm) of the stirring and results in a product with a great deal of unreacted calcium hydroxide present. The first two reactions were performed at 200 rpm and 450 rpm, without covering the top of the reaction flask. This resulted in water loss and turned the solution into a gel which made analysis impossible. Thus, it is clear that water loss must be minimized while stirring the reaction to ensure favorable results. **Figure 13** illustrates the success of reducing water loss and stirring the mixture at a high rpm. Examples 16 and 17 below were performed using an Erlenmeyer flask fitted with a rubber stopper to help minimize water loss and at 750 rpm and 250 rpm, respectively. Examples 18 and 19 were performed at 600 rpm and 400 rpm, respectively. It is clear that a higher rpm produces a more favorable product by reducing the formation of Peak 3. **Figure 14** illustrates that running the reaction at 500 rpm produced a 0.8% increase in Peak 3 and a slight increase in Peak 5. Therefore, the optimal revolutions per minute are between 500 and 600 rpm for the method of this example.

Table 6. Comparison of Example Revolutions Per Minute (RPM)

Solution	Basic Source	Relative Peak Distribution after Reaction (%)			Comparable ACH (%)	RPM
		Peak 3	Peak 4	Peak 5		
Example 16	Ca(OH) ₂	0.2	89.9	9.9	2.7	750
Example 17	Ca(OH) ₂	0.6	96.2	3.2	4.0	250
Example 18	Ca(OH) ₂	0.6	94.3	5.1	5.3	600
Example 19	Ca(OH) ₂	1.2	96.6	2.2	5.2	400

[0097] Optimal Method for the Addition of Ca(OH)_2

[0098] Simultaneous addition of aluminum chloride, glycine, and calcium hydroxide then mixing and heating yields unfavorable results: 25.5% Peak 3 and 16.7 Peak 5 which are poor for the formation of a pure Peak 4 complex. Also studied was the addition of a calcium hydroxide power to an aqueous aluminum chloride salt solution once the reaction was at 90°C . This produced increases in Peak 3 by 4.3% to 7.3%, and increases in Peak 5 by 13.3% to 17%. **Figure 15** and **Table 7** below illustrate the benefit of using a calcium hydroxide solution compared to a powder. Using a calcium hydroxide solution (Example 20) produced a 1% smaller Peak 3 and a 3.9% smaller Peak 5 compared to using a calcium hydroxide powder (Example 21). The calcium hydroxide in Examples 20 and 21 were added to the aluminum chloride salt solution 4 times over 1.5 hours. Further, the addition of a calcium hydroxide suspension initially occurred drop-wise by hand over 1 hour and 45 minutes. However, Example 10 and **Figure 11** illustrates favorable results obtained from adding a calcium hydroxide solution 5 times over 1 hour and 45 minutes, following all other optimal reaction conditions. Therefore the optimal way to add Ca(OH)_2 is in solution form over several additions.

Table 7. Comparison of Ca(OH)_2 solution vs. Ca(OH)_2 powder

Solution	Basic Source	Relative Peak Distribution after Reaction (%)			Comparable ACH (%)	No. of Additions
		Peak 3	Peak 4	Peak 5		
Example 20	Ca(OH)_2 solution	0.6	94.7	4.7	4.9	4 additions over 1.5 hours
Example 21	Ca(OH)_2 powder	1.6	89.8	8.6	2.7	4 additions over 1.5 hours

62301-2944D(S)

CLAIMS:

1. A method of making an antiperspirant active composition comprising the following sequence of steps:

(i) heating an aqueous solution containing an aluminum chloride compound
 5 having an aluminum to chloride molar ratio of 0.3:1 to 3:1, optionally with a buffer agent, at a temperature of 50°C to 95°C to reflux for a period of time of 1 hour to 5 hours to obtain an aluminum salt solution;

(ii) after the heating step, adding an aqueous solution of an inorganic base to obtain an aluminum salt solution having an OH:Al molar ratio of 2:1 to 2.6:1 to obtain a pH
 10 adjusted aluminum salt solution having a pH of 2 to 5; and

(iii) optionally adding an aqueous solution containing a zirconium compound to the pH adjusted aluminum salt solution to thereby obtain an aluminum-zirconium salt solution having a molar ratio of aluminum to zirconium of 5:1 to 10:1.

2. The method of claim 1, wherein the buffer agent is present in a molar ratio of
 15 buffer to aluminum is about 0.1:1 to about 3:1.

3. The method of claim 1 or 2, wherein the buffer agent is present and is at least one buffer selected from the group consisting of an amino acid, and betaine.

4. The method of claim 3, wherein the amino acid is glycine.

5. The method of any one of claims 1 to 4, wherein the inorganic base is selected
 20 from the group consisting of metal hydroxides and metal oxides.

6. The method of claim 5, wherein the inorganic base is selected from the group consisting of calcium hydroxide, strontium hydroxide, sodium hydroxide, barium hydroxide, calcium oxide, strontium oxide, and barium oxide.

62301-2944D(S)

7. The method of any one of claims 1 to 6, wherein the aluminum chloride compound is selected from the group consisting of aluminum trichloride, aluminum chlorohexahydrate, and aluminum dichlorohydrate.
8. The method of any one of claims 1 to 6, wherein the aluminum chloride
5 compound is aluminum trichloride.
9. The method of any one of claims 1 to 8, wherein the aqueous solution containing zirconium compound is added.
10. The method of claim 9, wherein the zirconium compound is $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$.
11. The method of any one of claims 1 to 10, wherein the antiperspirant active
10 composition exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 7 and a Peak 4 intensity greater than a Peak 5 intensity in aqueous solution.
12. The method of any one of claims 1 to 11, wherein the antiperspirant active composition has a SEC Peak 4 area of at least 50% of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram.
- 15 13. The method of any one of claims 1 to 11, wherein the antiperspirant active composition has a SEC Peak 4 area of at least 60% of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram.
14. The method of any one of claims 1 to 11, wherein the antiperspirant active composition has a SEC Peak 4 area of at least 70% of a total area of Peaks 1, 2, 3, 4, 5, and 6
20 in the SEC chromatogram.
15. The method of any one of claims 1 to 10, wherein the antiperspirant active composition has a SEC Peak 4 area of 95 to 100% of the total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram.

62301-2944D(S)

16. The method of any one of claims 1 to 10, wherein the antiperspirant active composition has a SEC Peak 3 area of less than about 10% of the total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram.

17. The method of any one of claims 1 to 10, wherein the antiperspirant active
5 composition has no SEC Peak 3 area.

18. The method of any one of claims 1 to 10, wherein the antiperspirant active composition has a SEC Peak 5 area of less than about 30% of the total area of Peaks 1, 2, 3, 4, 5, and 6.

19. The method of any one of claims 1 to 10, wherein the antiperspirant active
10 composition has no SEC Peak 5 area.

20. The method of any one of claims 1 to 10, wherein the antiperspirant active composition has a SEC Peak 1 area of less than about 10% and a SEC Peak 2 area of less than about 10% of the total area of Peaks 1, 2, 3, 4, 5, and 6.

21. The method of any one of claims 1 to 10, wherein the composition has a SEC
15 Peak 4 area of 95 to 100%, no SEC Peak 3 area, and no SEC Peak 5 area of a total area of Peaks 1, 2, 3, 4, 5, and 6 in the SEC chromatogram.

22. The method of any one of claims 1 to 10, wherein the antiperspirant active composition exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 10.

20 23. The method of any one of claims 1 to 10, wherein the antiperspirant active composition exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 17.

24. The method of any one of claims 1 to 10, wherein the antiperspirant active composition exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at
25 least 20.

62301-2944D(S)

25. The method of any one of claims 1 to 10, wherein the antiperspirant active composition exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 50.

26. The method of any one of claims 1 to 10, wherein the antiperspirant active
5 composition exhibits a SEC chromatogram having a SEC Peak 4 to Peak 3 intensity ratio of at least 100.

1/15




Figure 1
(Prior Art)




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15