




(86) Date de dépôt PCT/PCT Filing Date: 2007/07/13
 (87) Date publication PCT/PCT Publication Date: 2008/02/14
 (85) Entrée phase nationale/National Entry: 2009/02/06
 (86) N° demande PCT/PCT Application No.: EP 2007/057224
 (87) N° publication PCT/PCT Publication No.: 2008/017568
 (30) Priorité/Priority: 2006/08/10 (EP06118717.5)

(51) Cl.Int./Int.Cl. *A61K 39/145* (2006.01),
A61K 48/00 (2006.01), *A61P 31/00* (2006.01),
A61P 35/00 (2006.01), *C12N 15/79* (2006.01)
 (71) Demandeur/Applicant:
 INTERNATIONAL INVESTMENT AND PATENTS SA, LU
 (72) Inventeur/Inventor:
 GERLONI, MARA, US
 (74) Agent: RICHES, MCKENZIE & HERBERT LLP

(54) Titre : PLASMIDES AYANT UNE ACTION IMMUNOLOGIQUE
 (54) Title: PLASMIDS WITH IMMUNOLOGICAL ACTION

Map of plasmid 1



1-2483bp Segment encoding the variable region Vh62
 2484-5077 bp Segment encoding the constant region γ1
 6943-7737bp Segment encoding the gene for resistance to neomycin
 9324-9267bp *E. coli* origin of replication

(57) Abrégé/Abstract:

Recombinant plasmids usable for the transfection of eukaryotic and prokaryotic cells are described; such plasmids have a length comprised between 7 and 12 kbases and comprise a sequence encoding the heavy chain of an immunoglobulin; in particular, they may be used: -in a process of transfection of prokaryotic or eukaryotic cells (ex vivo) which can be inoculated into higher organisms in order to induce a prophylactic or therapeutic immune response; -in a protocol of direct inoculation (in vivo) in higher organisms in genie immunization methodologies with the aim of evoking prophylactic or therapeutic immune responses.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 February 2008 (14.02.2008)

PCT

(10) International Publication Number
WO 2008/017568 A1

(51) International Patent Classification:

A61K 39/145 (2006.01) A61P 35/00 (2006.01)
C12N 15/79 (2006.01) A61P 31/00 (2006.01)
A61K 48/00 (2006.01)

(21) International Application Number:

PCT/EP2007/057224

(22) International Filing Date: 13 July 2007 (13.07.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

06118717.5 10 August 2006 (10.08.2006) EP

(71) Applicant (for all designated States except US): INTERNATIONAL INVESTMENT AND PATENTS SA; 18 Avenue de la Porte-Neuve, L-2227 Luxembourg (LU).

(72) Inventor; and

(75) Inventor/Applicant (for US only): GERLONI, Mara [IT/US]; 2933 Arianne Drive, San Diego, US92117 (US).

(74) Agent: PISTOLESI, Roberto; Via Turati 32, I-20121 Milano, (IT).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: PLASMIDS WITH IMMUNOLOGICAL ACTION

Map of plasmid 1



1-2483bp Segment encoding the variable region Vh62
2484-5077 bp Segment encoding the constant region γ 1
6943-7737bp Segment encoding the gene for resistance to neomycin
9324-9267bp *E. coli* origin of replication

(57) Abstract: Recombinant plasmids usable for the transfection of eukaryotic and prokaryotic cells are described; such plasmids have a length comprised between 7 and 12 kbases and comprise a sequence encoding the heavy chain of an immunoglobulin; in particular, they may be used: -in a process of transfection of prokaryotic or eukaryotic cells (ex vivo) which can be inoculated into higher organisms in order to induce a prophylactic or therapeutic immune response; -in a protocol of direct inoculation (in vivo) in higher organisms in gene immunization methodologies with the aim of evoking prophylactic or therapeutic immune responses.

WO 2008/017568 A1

WO 2008/017568 A1



Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

Published:

— *with international search report*

— *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

Title**Plasmids with immunological action****Description**

5

The object of the present invention is represented by a recombinant plasmid usable for the transfection of eukaryotic and prokaryotic cells, having a length comprised between 7 and 12 kbases and comprising a sequence encoding the heavy chain of the immunoglobulin. A further object of the invention is represented by the use of the aforesaid plasmid for the preparation of a pharmaceutical formulation, or of a vaccine or a therapeutic treatment, for inducing an immune response in a human or animal organism.

15

In particular, described and claimed is a sequence of bases that are used for the construction of plasmids which may be used:

20

- in a process of transfection of prokaryotic or eukaryotic cells (*ex vivo*) which can be inoculated into higher organisms in order to induce a prophylactic or therapeutic immune response (in particular by means of the plasmid 1 having the sequence SEQ ID NO. 1);

25

- in a protocol of direct inoculation (*in vivo*) into higher organisms in genic immunization methodologies in order to evoke prophylactic or therapeutic immune responses (in particular by means of the plasmid 2 having the sequence SEQ ID NO.2).

30

A description is also given of a profile of pathologies which may be treated with the plasmids described.

STATE OF THE ART

DNA of plasmidic origin may be used for the transfection of prokaryotic and eukaryotic cells through known methods. The plasmids constructed for this purpose are generally constituted by a skeleton which has inserted units of genetic material encoding a certain protein which may or may not be provided with its own biological activity.

The plasmids may be of commercial origin, into which a part bearing the specificity is introduced into a structural construct that is already known and used, or generated autonomously, that is, by assembling fragments of selected genetic material on the basis of a certain profile to be reconstructed at the organism for which the plasmid is destined.

The synthesis of a plasmid is an operation of fundamental importance for the purpose of obtaining the desired cellular properties. The plasmids determine the efficiency of the transfection and of the synthesis of the transgenic protein, and also the safety of the transfection, and therefore in the final analysis the resultant efficacy of the proteic expression of the transfected cell.

A plasmid is a vector of genetic data which influence the cellular cycle of the host cell and, consequently, the life cycle of the organism which hosts these transfected cells: the more data introduced into the plasmid, the more risks are run in the course of the transfection.

A plasmid is characterized by the specificity of the data contained: the less complex it is, the easier the synthesis thereof and the safer the use thereof.

A plasmid exercises the capacity of transfecting a cellular population more or less spontaneously

depending on the type of cell and the experimental conditions of contact/incubation in which transfection is carried out. The more selective a plasmid is for a specific cellular population, the more it is usable in conditions of safety.

The conditions in which transfection is carried out are decisive for the success thereof: the homogeneity and the concentration of the cells to be transfected, the incubation time and conditions, the possibility of monitoring the phenomenon with specific and selective methods constitute an extremely important corollary for the success of the operation.

In the specific case in which a plasmid is used for transfecting cells to be injected *in vivo* in patients, such cells are to be handled with extreme caution inasmuch as they must be re-inserted into the patient, and risky and fatal collateral phenomena cannot be risked: the less handling is necessary, the greater the safety of the method and the more reproducible the result.

From the above it inevitably follows that the use of a plasmid of reduced dimensions, especially if combined with a method with limited handling, is a condition to be preferred in transfection used for prophylactic and therapeutic purposes.

The use of a plasmid encoding immunogenic epitopes in an autologous transfection method is one of the systems that can be used for inducing a specific immune response in some pathologies characterized by the appearance of infecting agents, or by spontaneous or induced cellular mutation (carcinogenesis), with modification of the apoptotic course of a selected cell or cellular line.

WO 90/09804 describes immunoglobulins genetically engineered for expressing a predefined peptide epitope in the variable region or in the bond domain of the immunoglobulin.

5 WO 00/61766 describes tumoral antigens derived from telomerasis that can be used for generating a response mediated by T-cells against telomerasis and consequently against the tumour itself.

WO 00/64488 describes a plasmid encoding chimeric heavy chain of an immunoglobulin, the pN γ_1 V_H62 plasmid, which is obtained by subcloning the murine V_H62 gene into the pN γ_1 plasmid, containing a sequence encoding a human γ_1 constant region. This plasmid can be modified by introduction of heterologous epitopes in any of the complementarity determining regions of the variable region. Therefore, its use in a method for inducing an immunoresponse is disclosed.

However, the pN γ_1 V_H62 plasmid contains portions that will be harmful if this plasmid would be injected into humans. Furthermore, the dimensions of this plasmid are such that a very low transfection yield is obtained.

DESCRIPTION OF THE INVENTION

The present inventors have now developed plasmids comprising a sequence encoding the heavy chain of immunoglobulin that do not present the drawbacks of the pN γ_1 V_H62 plasmid when used for inducing an immunoresponse *in vivo* or *ex vivo* in a human or animal organism. In particular, the plasmids developed have a better safety profile and show an increase yield when transfected into cells.

30 These plasmids have a length comprised between 7 and 12 kbases, preferably between 8 and 12 kbases, more preferably between 9 and 11 kbases and, even more

preferably, between 9 and 10 kbases and are able to express the heavy chain of immunoglobulin when they are transfected into lymphocytes.

According to a preferred embodiment, these plasmids express a chimeric heavy chain of immunoglobulin, preferably comprising a murine variable region and a human constant region, preferably Ig γ 1.

The murine variable region is preferably the V_H region from hybridoma 62 derived from splenocytes of an adult hyperimmunised mouse (Zanetti et al. *J. Immunol.*, 1983, 131:2452), hereinafter referred to as V_H62 region. The V_H62 hybridoma 62 secretes a monoclonal antibody with anti-tyroglobulin activity

The Ig γ 1 region gene is preferably cloned from the vector pN γ 1.

The plasmids may further contain a promoter specific for lymphocytic cells, preferably of around 50 bp, or of viral origin, preferably the CMV promoter.

The plasmids of the invention preferably also comprise the polyadenylation sequence AATAAA.

It is preferred that the plasmids of the invention do not express resistance to betalactamic antibiotics and in particular to ampicillin, and/or do not comprise a replication origin of SV40.

Accordingly, the plasmids preferably contain a replication origin of *Escherichia coli*, preferably PBR322. Furthermore, the plasmids preferably express resistance to neomycin.

Preferred plasmids of the invention are plasmids 1 and 2.

The plasmid 1 has a length of 9727 bp (SEQ ID NO. 1 and Figure 1) and the plasmid 2 of 10004 bp (SEQ ID NO. 2

and Figure 2). Both the plasmids described encode a chimeric heavy chain of immunoglobulin.

The sole structural difference between the two plasmids is determined by the specific promoter, which
5 in the case of the plasmid 1 is a specific promoter for lymphocytic cells of 50 bp, and in the case of the plasmid 2 is a viral promoter with dimension of 742 bp.

The skeleton of the plasmids is represented by pSV2neo, a DNA of bacterial origin containing the gene
10 for resistance to neomycin and the origin of replication PBR322.

The genetic sequence encoding the heavy immunoglobulinic chain is composed of:

- a murine variable region (V_H62) of around 2.5 kb,
15 originally cloned from a mouse hybridoma (hybridoma 62) secreting a monoclonal antibody with anti-tyroglobulin activity (Sollazzo, et. al., *Eur. J. Immunol.*, 1989);
- a human Igγ1 constant region deriving from the pNγ1 vector (Hybritech Corporation, San Diego, CA).

20 The immunoglobulinic promoter of the plasmid 1 is an integral part of V_H62, while the viral promoter of the plasmid 2 was derived from the plasmid pMGFP (Promega, WI, USA). The gene for resistance to neomycin also confers a resistance to kanamycin for
25 selective growth in prokaryotic cells.

The polyadenylation sequence is in position 5178:5183 of the plasmid and follows the human constant region. The sequence is as follows: AATAAA. The origin of bacterial replication is of *Escherichia coli*
30 and is situated in position 8324:9264.

The particular feature of the plasmids of the invention is that the regions determining the complementarity (CDR) of the encoded protein may be mutagenized for the purpose of introducing therein

epitopes (from 5 to 25 amino acid residues) with antigenic properties (Gerloni et al *Nature Biotechnology* 1997). Such properties make it possible to use the plasmid in question for inducing a specific immune response either by means of a transfection *ex vivo* with the inoculum of cells transfected spontaneously or by means of direct inoculation *in vivo* (genetic immunization).

Accordingly, an object of the present invention is the use of the plasmids of the invention for the preparation of a pharmaceutical formulation for DNA vaccination, in particular for inducing an immunoresponse in a human or animal organism. Furthermore, another object of the invention is a formulation containing at least one plasmid according to any one of the preceding claims together with pharmaceutically acceptable excipients and/or adjuvants.

The formulation of the invention may be used for prophylactic or therapeutic purposes.

The formulation of the invention may be used in particular in a human or animal organism that is or was affected by:

-tumours belonging to the family of carcinomas and/or adenomas and/or sarcomas and/or lipomas and/or solid and/or ascitic tumours, by prostate or pancreatic, renal or pulmonary carcinoma. Preferably, in this case said organism is or was affected by the presence of tumour cells having on the surface at least one antigenic epitope, the encoding sequence of which is contained in the plasmid.

-bacterial, viral, fungal and/or parasitic infections.

A further object of the present invention is a method for inducing an immunoresponse in a human or animal

organism. Said method comprised transfection of prokaryotic and/or eukaryotic cells *ex vivo* and the subsequent inoculation of said prokaryotic and/or eukaryotic cells into said human or animal organism. Preferably, said transfected cells belong to the family of lymphocytes and are preferably taken from the peripheral vessels of said human or animal organism. Alternatively, said method comprises inoculation of the plasmid *in vivo* into said human human or animal organism.

Inoculation of the plasmid or of the prokaryotic and/or eukaryotic cells into said human or animal organism is preferably carried out by means of injective or transmucosal administration.

The protein encoded by the plasmids of the invention, like all the immunoglobulins, possesses 3 CDRs, CDR1 with a restriction site usable for the insertion of peptide sequences AfiIII, CDR2 with a NcoI site and CDR3 with an Acc65I site.

It is therefore possible to insert into them various peptide sequences capable of evoking various immune responses both of the humoral type (mediated by B cells) and of the cellular mediated type (mediated by T-cells CD4 and CD8); for example, it is possible to insert at least a single sequence on each CDR, for a total of three sequences capable of evoking various immune responses; it is further possible to insert one or more sequences, optionally fused with one another, on each CDR.

An example of insertion of antigenic epitopes into the CDRs is shown in Figure 3.

The antigenic epitopes preferred for the purposes of the present invention are:

-tumoral antigens, such as for example that of telomerasis and, even more preferably, p540 (ILAKFLHWL), p572 (RLFFYRKS SV), pY572 (YLFFYRKS SV) and p865 (RLVDDFLLV);

- 5 -antigens deriving from infective microorganisms, such as, for example, that of influenza and, preferably, the epitope pNP (ASNENMETM).

The transgenic product encoded plasmids 1 and 2 is a protein with a molecular weight of around 156.000 daltons (Figure 4). The heavy region encoded is chimeric in nature: part human (the constant region) and part murine (the variable region). Nevertheless, the murine part contains sequences 80% homologous with the human variable regions.

15 Important characteristics of the aforesaid plasmids are the absence of the gene for resistance to ampicillin and the presence of the gene for resistance to kanamycin which provides for the safe use thereof in subjects with potential allergies to the betalactamic antibiotics. Another advantageous factor is that kanamycin is an antibiotic stable at 37°C (conditions of culture of the plasmid) for 24-48 hours, while ampicillin is stable only for 3-4 hours, consequently allowing a culture yield of the plasmid which is greater (less expensive production) and more stable.

25 The aforesaid plasmids are also devoid of "useless" sequences (for example the sequence SV40 or pieces of genomic material) which would represent greater risks of homology with the genome of the host cell and therefore greater risks of integration in the cell itself.

Analysis of the restriction map of the plasmids

30 The restriction map obtained by digestion with restriction enzymes is the first criterion to be

considered in order to define the identity of a plasmid. In particular, the map shown in Figure 5 identifies in an exclusive manner the plasmids 1 and 2. There is also shown in succession an image of the fragments of the plasmid 1 after digestion, run on agarose gel in parallel with a standard of known dimension (1Kb ladder) in order to determine the exact dimension thereof. The sequence of the plasmid 1 is shown in SEQ ID NO. 1, while the sequence of the plasmid 2 is shown in SEQ ID NO. 2.

Biological characterization of the plasmids

It has already been indicated previously that a plasmid encoding immunogenic epitopes in a method of transfection is one of the systems that can be used for inducing a specific immune response in some pathologies characterized by the appearance of infecting agents, or by spontaneous or induced cellular mutation (carcinogenesis), with modification of the apoptic course of a selected cell or cellular line.

A further use for such a plasmid is the direct injection *in vivo* into immuno-competent organisms in order to induce an immune response against proteins of a foreign nature and pathogenic microorganisms (Tang et al., *Nature* 1992, Ulmer et al., *Science* 1993, Gerloni et al., *Nature Biotechnology* 1997). In this field with the inoculation of functional genes the induction of humoral responses (mediated by antibodies) and cellular mediated responses (mediated by T-lymphocytes of type CD4 and CD8) effective in the treatment or prevention of pathologies of infective and cancerous origin was demonstrated.

Consequently, the concept of genic immunization is now adopted by vaccinologists all over the world, who use plasmids encoding antigens deriving from bacteria,

viruses and parasites and also from various types of tumour in order to evoke specific and protective immune responses. Clinical trials are currently under way for the therapy or prophylaxis of HIV, herpes, influenza, avian influenza, SARS, hepatitis B and C and carcinomas of various kinds.

The essential components of a plasmid to be used *in vivo* are the gene encoding the antigen (or pieces thereof) of interest, a promoter sequence (normally derived from cytomegalovirus, CMV) which guides the transcription of the antigen, a region of polyadenylation which ensures the translation thereof.

Furthermore, together with the origin of replication for the amplification of the plasmid in bacterial cells there is also a gene which encodes antibiotic resistance in order to ensure the selection of the bacterial population and to eliminate contamination during culture.

Another intrinsic property of the DNA vaccines is that plasmids of bacterial origin contain sequences of non-methylated cytosine together with residues of guanosine (CpG). These CpG units have the capacity of increasing the immunogenic capacity of the plasmids themselves and therefore function as adjuvants.

The direct inoculation of nucleic acids into somatic cells appears to mimic the immunity induced by natural infections and offers various advantages, including the possibility of producing and testing such plasmids in an inexpensive, easy and rapid manner. Moreover, the plasmids are much more stable than conventional vaccines and may be preserved as lyophilisates.

The plasmid is usually inoculated *in vivo* by the intramuscular or intradermal route, although other

routes such as the oral, vaginal, endovenous, intraperitoneal and subcutaneous routes are applicable. The plasmids are administered in a variety of diluents which include distilled water, saline or sugar solutions, physiological buffers, isotonicising compounds, preservative or cryoprotective substances in case the processes of lyophilisation are necessary.

The dose of plasmid used in the immunization protocols varies from case to case but, as a rule, amounts of from 25 to 200 µg per dose are used with 3 doses/injections at intervals of three weeks.

A further object of the present invention is therefore constituted by two recombinant plasmids characterized by a sequence corresponding to SEQ ID NO. 1 and SEQ ID NO.2, respectively or a sequence at least 90% homologous, preferably 95% homologous to SEQ ID NO. 1 and SEQ ID NO.2. These plasmids have dimensions that are reduced but suitable for the purpose and a transfection method that is suitable, reproducible, safe and effective.

The use of such plasmids has specific characteristics:

-optimum yield in the production process, inasmuch as a plasmid of measured content is simpler to produce and gives rise to better production yields if reduced in amplitude;

-optimum stability of the cellular culture, since the plasmids contains the gene for resistance to antibiotic kanamycin stable at 37°C for 24-48 hours (as opposed to ampicillin stable only for 4-6 hours at the same temperature) the stability of the bacterial culture is consequently significantly increased, with undoubted advantages in terms of yield;

-high efficiency of spontaneous transfection (capacity of penetration into the cell) because of contained molecular dimensions and lesser steric bulk, a not negligible detail in the case of protocols in which the use of spontaneous transfection is envisaged;

-the absence of the possibility of anaphylactic reactions that can be induced in the subjects treated, if predisposed to an allergic reaction to ampicillin, inasmuch as the plasmid, not possessing the gene for resistance to ampicillin, is not grown in the presence of that antibiotic;

-low, if not zero, possibility of integration in the genome of the host cell inasmuch as the minimum quantity of plasmid used for transfection renders practically negligible the risk of integration (consequently reduced possibility of induction of oncogenic mutations in the host cell);

-low, if not zero, possibility of direct integration in the genome of the host cell inasmuch as the plasmid does not possess extraneous sequences (such as SV40) which could have homologies with the genome of the cell itself and represent greater risks of integration;

-high flexibility of use of the plasmid also for direction injection *in vivo* in genic immunization protocols. In fact, by substituting in the nucleotidic sequence the specific promoter for B cells with a viral promoter (CMV) with wider expression, the use of the plasmid is extended by simple transfection *ex vivo* to the use of direct inoculation in higher organisms.

A further object of the present invention is constituted by the use of a method of transfection *ex vivo* without the use of any physical or chemical means which might facilitate the process. Such a method does not induce disturbances in the functionality of the

transfected cells and does not induce genetic transformation thereof. The method is characterized by separation of the specific cellular material to be transfected from the remainder of the corpuscular and fluid part of the peripheral blood, obtainable with processes with less handling than the normal centrifuging methods: the use of apheresis makes it possible to separate a large number of lymphocytic cells on which to carry out transfection in a manner which is painless for the patient and very much more useful for experimental purposes. In fact:

- it does not disturb the functionality of the cells with gravitational or mechanical shocks;
- it allows the harvesting of a large number of cells, to be used for the process and to keep stored as reference for the subsequent phase of therapeutic monitoring;
- it does not impoverish the functional and structural resources of the patient or his coagulative or reparative processes;
- it does not add any risk of contamination of the biological material and/or of the patient.

An object of the present invention is constituted by the use of a plasmid as described for effecting the transfection *ex vivo* of selected cells by means of a spontaneous process with the adoption of the following modalities:

1. transfer of the peripheral blood originating from a patient into an instrument capable of directly separating the family of lymphocytes from the rest of the corpuscular fraction and from the serum (apheresis process);
2. isolation of a quantity of lymphocytic cells suitable for the application of the treatment

- described hereinafter; then transfer of the isolated lymphocytes and washing with PBS (without Ca^{++} and Mg^{++}) and further centrifuging;
3. dilution 1:1 with tryptan blue and counting of the lymphocytes via haemocytometer and microscope in order to verify that they have preserved at least 90% of vitality;
 4. re-suspension of the lymphocytes at a concentration of around 20×10^6 cells/ml in PBS (without Ca^{++} and Mg^{++}) and re-division of an aliquot into plates with U-shaped wells, where 25 μg of a plasmid having the sequence shown in SEQ ID NO. 1 are added;
 5. incubation of the plates with the transfection wells in an incubator for 30-90 minutes at 37°C and 5% of CO_2 ;
 6. transfer of a suitable aliquot of cells treated with plasmid into a suitable culture medium and leaving to incubate for one night;
 7. verifying that the cellular vitality is maintained above a threshold judged to be appropriate (typically 70%) and calculating the dose to be transferred to the patient;
 8. transfer into a phlebotomy bag of a suitable volume of transfected cells containing the dose to be re-administered to the patient (typically variable from a few thousands to a few tens of millions of cells);
 9. within the scope of the treatment intended to arouse an immune response against the cells which express at the surface the proteins of which the epitopes are contained in the plasmid, inoculation into the patient of the blood contained in the

bag, transgenized by means of the use of the plasmid having the sequence shown in SEQ ID NO. 1.

As an alternative to the method described above, the isolation of the lymphocytes to be transfected may also be carried out by a classic method based on centrifuging; in that case the treatment of apheresis described above in item 1 and item 2 may be substituted by the following:

1. transfer of the peripheral blood originating from a patient into test tubes, addition of buffer with Ficoll and centrifuging until the lymphocytes stratify into an unmistakable band;
2. transfer of the isolated lymphocytes and washing with PBS (without Ca^{++} and Mg^{++}) and further centrifuging.

The use of the plasmid of reduced dimensions and a selected population of cells as described above makes it possible to obtain advantages from the process, such as a reduced time for handling of the blood and of its lymphocytic fraction, a reduced incubation time, a substantially reproducible process yield, the possibility of applying a high degree of automation to the various steps of the process and the possibility of performing the transfection process directly inside an isolated device which does not require the adoption of sterile conditions of the working ambience outside the device itself, which are required by the handling of biological fluids intended for human administration.

A object of the present invention is constituted by the use of a plasmid of contained dimension and characterized by the sequence described, with a method of isolation of the material to be transfected with reduced handling, with transfection conditions

described for inducing immune responses in the treated patients against the agent responsible for the infection or the mutation, of which the specific imprint at the sequential and conformational level is contained in the genetic material represented in the plasmid.

A further object of the present invention is constituted by the use of a device in which to carry out the incubation of the plasmid with the cells, characterized by the presence of a space in which to accommodate the desired volume of cells, a port through which to insert a needle in order to deposit a solution of the plasmid and an optional other port through which to effect the sampling of the transfected cells.

In a typical application of the present invention, the plasmid with the sequence shown in SEQ ID NO. 1 is used on a population of lymphocytic cells separated from the blood by the method of apheresis; these transfected cells, after incubation under the normal conditions used for biological fluids, are injected back into a patient in order to induce an immune type of response.

In a further application of the present invention, the plasmid constructed as described is used under the conditions described for transfecting a population of lymphocytic cells harvested by means of apheresis or centrifuging from a patient affected by a tumour of the pancreas, or prostate, lung, kidney, or skin, or affected by some other type of adenomatous or carcinomatous or other tumoral form, either solid or acytic in form, in order to induce in the patient himself a selective immune response against only the cancerous cells, expressing superficially the proteic fraction contained in the plasmid and by these caused

to express the lymphocytes, used as APC (cells presenting the antigen).

In a further application of the present invention, the plasmid shown in SEQ ID NO. 1 is modified by the substitution of the promoter for lymphocytic cells with a promoter of viral origin (CMV). Such substitution gives rise to a plasmid of slightly larger dimensions (300 bp), shown in SEQ ID NO. 2, and used for inducing an immune response in an organism by means of direct parenteral (intramuscular, intradermal, subcutaneous or endovenous) administration or transmucosal administration (via the nasal, oral, intestinal or vaginal mucosa).

The plasmids described were evaluated for:

1. the capacity for spontaneously transfecting human lymphocytes;
2. the capacity for inducing an immune response in laboratory mice when injected with transfected cells *ex vivo*;
3. the capacity for inducing an immune response in laboratory mice when injected directly with the plasmid 2 in a genic immunization protocol.

The following examples are intended purely as non-limiting illustrations of the invention.

The plasmid 1 was constructed according to the following protocol:

1. Modification of the plasmid PSV2 neo
 - The plasmid PSV2 neo was digested with the restriction enzymes AhdI and XmnI in order to remove the segment for resistance to ampicillin (amp^r)

- PSV2 neo (amp^r minus) was digested with BsmBI and HindIII in order to remove the origin of replication SV40 (SV40)
 - PSV2 neo (amp^r and SV40 minus) was digested with AatII and BstBI. At the end of this process a fragment of 2737 pairs of bases containing the origin of replication of *E. Coli* and the gene for resistance to neomycin/kanamycin was obtained. This fragment was isolated from agarose gel, purified on a column and maintained at 4°C until subsequent use.
2. Modification of the plasmid γ 1Vh62
- The plasmid γ 1V_H62 was digested with the restriction enzymes FseI and BamHI in order to remove the segment of 4787 pairs of bases encoding human genomic DNA
 - The plasmid thus derived (DNA^{genomic}minus) was circularised with the new dimension of 10721 pairs of bases. It was digested with AatII and BstBI and the fragment containing the variable and constant region was purified for the subsequent ligation reaction
3. Construction of the final plasmid 1
- The plasmid PSV2 neo (amp^r and SV40 minus) deriving from the passage 1 was digested with AatII and BstBI.
 - The modified plasmid γ 1V_H62 (DNA^{genomic}minus) deriving from the passage 2 was digested with AatII and BstBI.
 - The two modified plasmids thus digested were bonded together and used for transfecting competent cells of *E. coli*.

- The plasmidic DNA extracted from the cells of *E. coli* was then used for the tests of identification by means of PCR, restriction map and sequencing.
- 5 4. Construction of the final plasmid 2
- The plasmid 1 was digested with DraII and BclI in order to remove the immunoglobulinic promoter contained therein
 - From the plasmid pMGFP, by the same
10 digestion, the viral promoter CMV was removed
 - To the plasmid 1, without promoter, by means of PCR, the promoter CMV was added so as to obtain the plasmid 2.

15 The following examples are intended purely as a non-limiting illustration of the invention.

Example 1.

An aliquot of around 50 ml of peripheral blood originating from a patient, with the addition of a suitable aliquot of anticoagulant, is transferred into
20 50 ml tubes, diluted 1:1 with buffer, 20 ml of Ficoll-Paque™ are added and centrifuging is carried out for 20 minutes at 2000 rpm.

After centrifuging, the lymphocytes, contained in the interface band, are harvested and transferred into
25 a new tube and washed 3 times in PBS (without Ca⁺⁺ and Mg⁺⁺) with recovery by centrifuging.

An aliquot of lymphocytes is diluted 1:1 in tryptan blue and counted in a haemocytometer with the microscope in order to determine the vitality thereof
30 (at least 90%).

After the count, the lymphocytes are re-suspended at a concentration of 20×10^6 cells per ml in PBS and aliquoted into plates provided with wells with U-shaped bases. There are then added 25 µg of a plasmid, having

the sequence shown in SEQ ID NO. 1, and the plate is placed in an incubator for 60-90 minutes at 37°C and 5% of CO₂.

The cells are then diluted to a concentration of 1X10⁶ cells per ml in suitable culture medium, placed in flasks inside an incubator and left for one night at 37°C and 5% of CO₂. After having verified that the cellular vitality is above 70%, the cells are washed twice in saline solution, the number of total cells containing the dose to be inoculated is then calculated, corresponding to a number of cells variable from 10,000 to 100 million and the latter are then re-suspended in an endovenous administration bag, which is then used for inoculation into the patient.

An aliquot of the transfected lymphocytes is tested before use by means of the extraction of DNA and mRNA in a nested PCR test in such a way as to evaluate the transfection of lymphocytes that has occurred and give a quantification thereof, even approximate.

Figure 6 shows the amplification with a PCR test of the DNA from human lymphocytes transfected with plasmidic DNA. Ladder is the reference for the determination of the dimension of the amplified fragments. The numbers refer to the number of cells amplified, and Naïve means lymphocytes not transfected with DNA (negative control). The figure demonstrates the transfection of the patient's lymphocytes which has taken place, and above all denotes its specificity, inasmuch as no fragment is amplified from lymphocytes not subjected to transfection.

Example 2.

Proceeding from the assumption that the lymphocytes transfected with a DNA encoding antigenic sequences are capable of evoking an immune response

after inoculation into laboratory animals and that the plasmid of the present invention, shown in SEQ ID NO. 1, encodes an epitope capable of evoking a cellular response on the part of the T CD4 lymphocytes, in order to verify the biological activity of the lymphocytes transfected by DNA, an experiment was planned for the verification of the immunogenicity of such lymphocytes *in vivo*, devised as described hereinafter.

- 4 C57/B16 mice were inoculated with 5,000 murine lymphocytes transfected with the plasmid as described in the protocol contained in the present invention. Inoculation was carried out endovenously in the tail vein;

- 14 days after inoculation, the mice were sacrificed and the cells of the spleen were used in a test of proliferation in the presence of the epitope encoded by the plasmid used for transfection;

- the splenocytes were cultured for 72 hours with the epitope of reference, and then tritiated thymidine (a radioisotope capable of showing the cellular proliferation) was added thereto;

- after 18 hours, the cells were harvested and the radioactivity measured with a beta counter instrument. The proliferation was expressed as stimulation index (SI), which is calculated by dividing the radioactive counts in the presence of the specific epitope by the radioactive counts in the presence of an uncorrelated epitope (non-specific proliferation). Conventionally, an SI of more than 3 is considered positive.

Table I : example of immune response induced by the inoculation into laboratory animals of lymphocytes transfected with the plasmid in question. The specific peptide refers to the epitope encoded by the DNA

incorporated in the plasmid of the invention, the control peptide has an uncorrelated sequence.

5	Mouse	SI Control peptide	SI Specific peptide
	#1	0.8	11.8
10	#2	1.3	14.3
	#3	1	13.3
15	#4	1.1	16.3

The experiment described above shows a specific immune response induced in mice when immunized with lymphocytes transfected with the plasmidic DNA, proving a clear biological property thereof when used in a test *in vivo*.

Example 3.

- Using a plasmid as having the sequence shown in SEQ ID NO. 2, 10 C57/B16 mice were inoculated intramuscularly, in the quadriceps, with 100 µg of DNA diluted in 100 µl of saline solution (group A). The immunization protocol consisted of a total of 3 injections separated by three weeks' interval. Another group of 10 C57/B16 were immunized in the same way, using the same plasmid with the CDRs empty (without any epitope) (group B) as control of specificity of the vaccination;

- 14 days after the last inoculation, the mice were sacrificed and the cells of the spleen were used in a

test of cytotoxicity in comparison with cells pulsed with epitope (NP) encoded by the plasmid used for transfection. The specific epitope is a cytotoxic sequence of the nucleoprotein of Influenza Virus A/PR8;

5 - the splenocytes were cultured for 5 days with the epitope of reference and then used as effectric cells in a cytotoxicity test;

- the cytotoxicity test consists in measuring the radioactivity (^{51}Cr) released by cells (EL-4) pulsed

10 with epitope NP which are killed by the effectric cells from the immunized mice; the cytotoxicity is expressed as a percentage of lysis at a certain ratio of Effectrics:Target (E:T), which is calculated by dividing the amount of radioactivity released by the

15 cells pulsed with the peptide by the amount of radioactivity released by the non-pulsed cells (specific cytotoxicity).

Table II: Example of cytotoxicity mediated by specific T CD8 cells in animals immunized with the plasmidic DNA

20 #2.

The experiment described above shows the specific immune response induced in mice when immunized with

Group	% of specific lysis
A	50% <u>+12</u>
B (control)	3% <u>+2</u>

plasmidic DNA, proving a clear biological property thereof when used in a test *in vivo*.


C L A I M S

1. A recombinant plasmid having a length of between 7 and 12 kbases and comprising a sequence encoding the heavy chain of immunoglobulin, characterized in that it contains at least one sequence encoding an antigenic epitope and in that it does not express resistance to betalactamic antibiotics.
2. A plasmid according to claim 1, characterized in that said betalactamic antibiotic is ampicillin.
3. A plasmid according to claim 1, characterized in that it has a length of between 8 and 12 kbases.
4. A plasmid according to claim 1, characterized in that it has a length of between 9 and 11 kbases, preferably of between 9 and 10 kbases.
5. A plasmid according to claim 1, characterized in that said heavy chain of the immunoglobulin is chimeric.
6. A plasmid according to claim 5, characterized in that said heavy chain of the immunoglobulin comprises a murine variable region and a human constant region.
7. A plasmid according to claim 6, characterized in that said murine variable region is V_H62.
8. A plasmid according to claim 7, characterized in that said murine variable region V_H62 is cloned from a mouse hybridoma secreting a monoclonal antibody with anti-tyroglobulin activity.
9. A plasmid according to claim 6, characterized in that said human constant region is Ig_γ1.
10. A plasmid according to claim 9, characterized in that said human constant region Ig_γ1 is cloned from the vector pN_γ1.
11. A plasmid according to claim 1, characterized in that it comprises a promoter for lymphocytic cells, preferably of around 50 bp, or a viral promoter, preferably of around 742 bp.
12. A plasmid according to claim 1, characterized in that it comprises a polyadenylation sequence AATAAA.

13. A plasmid according to claim 1, characterized in that it comprises a bacterial replication origin of *Escherichia coli*.
14. A plasmid according to claim 1, characterized in that it expresses resistance to neomycin and/or that it comprises the replication origin PBR322.
15. A plasmid according to claim 1, characterized in that it does not comprise a replication origin of SV40.
16. A plasmid according to any one of the preceding claims, characterized in that said antigenic epitope is selected from among p540 (ILAKFLHWL), p572 (RLFFYRKS SV), pY572 (YLFFYRKS SV), p865 (RLVDDFLLV) and pNP (ASNENMETM).
17. A recombinant plasmid characterized in that it has the sequence SEQ ID NO. 1 or a sequence at least 90% homologous, preferably at least 95%, to SEQ ID NO. 1.
18. A recombinant plasmid having the sequence SEQ ID NO. 2 or a sequence at least 90% homologous, preferably at least 95%, to SEQ ID NO. 2.
19. A formulation containing at least one plasmid according to any one of the preceding claims together with pharmaceutically acceptable excipients and/or adjuvants.
20. The use of a plasmid according to claims 1-18 for the preparation of a pharmaceutical formulation for inducing an immune response in a human or animal organism.
21. The use according to claim 20, characterized in that said human or animal organism is or was affected by tumours belonging to the family of carcinomas and/or adenomas and/or sarcomas and/or lipomas and/or solid and/or ascitic tumours, by prostate or pancreatic, renal or pulmonary carcinoma.
22. The use according to claim 20, characterized in that said human or animal organism is or was affected by bacterial, viral, fungal and/or parasitic infections.
23. The use according to claim 20, characterized in that such a formulation is usable for prophylactic purposes.
24. The use according to claim 20, characterized in that said organism is or was affected by the presence of tumour cells


- having on the surface at least one antigenic epitope, the encoding sequence of which is contained in said plasmid.
25. Method for inducing an immunoresponse in a human or animal organism comprising the transfection of prokaryotic and/or eukaryotic cells *ex vivo* by means of a plasmid according to claims 1-18 and/or a formulation according to claim 19 and the subsequent inoculation of said prokaryotic and/or eukaryotic cells into said human or animal organism.
 26. The method according to claim 25, characterized in that the cells to be transfected belong to the family of lymphocytes.
 27. The method according to claim 26, characterized in that said cells are taken from the peripheral vessels of said human or animal organism.
 28. The method according to claim 25, characterized in that said plasmid has the sequence SEQ ID NO. 1 or a sequence at least 90% homologous, preferably at least 95%, with respect to SEQ ID NO. 1.
 29. Method for inducing an immunoresponse in a human or animal organism comprising the inoculation *in vivo* into said human or animal organism of a plasmid according to claims 1-18 and/or of a formulation according to claim 19.
 30. The method according to claim 29, characterized in that said plasmid has the sequence SEQ ID NO. 2 or a sequence at least 90% homologous, preferably at least 95%, to SEQ ID NO.2.
 31. The method according to claims 25 to 30, wherein the inoculation of the plasmid or of the prokaryotic and/or eukaryotic cells into said human or animal organism is carried out by means of injective or transmucosal administration.

Figure 1. Map of plasmid 1



1-2483bp	Segment encoding the variable region Vh62
2484-5077 bp	Segment encoding the constant region γ 1
6943-7737bp	Segment encoding the gene for resistance to neomycin
9324-9267bp	<i>E. coli</i> origin of replication

Figure 2. Map of plasmid 2



- | | |
|------------------|--|
| 1 to 2775 | Segment encoding the variable region Vh62 |
| 161 to 907 | Viral promoter CMV |
| 2776 to 5366bp | Segment encoding the constant region γ1 |
| 7232 to 8026bp | Segment encoding the gene for resistance to neomycin |
| 8613bp to 9553bp | <i>E. coli</i> origin of replication |

Figure 3. Diagrammatic representation of a model of insertion of epitopes into the CDRs of the plasmid




Figure 4. Hypothethised structure of a protein with antigenic epitopes that is encoded by the plasmid described




Figure 5. Restriction map of the plasmids

Restriction enzyme	Originated fragments	
	Plasmid 1	Plasmid 2
EcoRI	2.5kb + 7.2kb	2.6kb + 7.2kb
XbaI	1kb + 8.7kb	1kb + 8.8kb
BamHI	9.7kb	9.8kb
HindIII	0.5kb + 2.4kb + 4.9kb + 1.9kb	0.5kb + 2.3kb + 7.0kb + 1.9kb






Figure 6. Amplification by means of PCR test of the plasmid from lymphocytes transfected *ex vivo*



Map of plasmid 1



1-2483bp

Segment encoding the variable region Vh62

2484-5077 bp

Segment encoding the constant region γ 1

6943-7737bp

Segment encoding the gene for resistance to neomycin

9324-9267bp

E. coli origin of replication