

(19) **DANMARK**

(10) **DK/EP 3936153 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 48/00 (2006.01)** **A 61 K 9/127 (2006.01)** **A 61 K 31/4745 (2006.01)**
A 61 K 31/496 (2006.01) **A 61 K 31/4965 (2006.01)** **A 61 K 31/497 (2006.01)**
A 61 K 31/555 (2006.01) **A 61 K 31/7068 (2006.01)** **A 61 P 35/00 (2006.01)**
C 07 D 403/12 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2024-11-25**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-08-21**
- (86) Europæisk ansøgning nr.: **21182826.4**
- (86) Europæisk indleveringsdag: **2017-01-11**
- (87) Den europæiske ansøgnings publiceringsdag: **2022-01-12**
- (30) Prioritet: **2016-01-11 US 201662277262 P** **2016-11-10 US 201662420258 P**
 2017-01-09 US 201762444172 P
- (62) Stamansøgningsnr: **17738831.1**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Celator Pharmaceuticals, Inc., 3170 Porter Drive, Palo Alto, CA 94304, USA**
- (72) Opfinder: **DRUMMOND, Daryl C., , Palo Alto, CA 94304, USA**
GENG, Bolin, , Palo Alto, CA 94304, USA
KIRPOTIN, Dmitri B., , Palo Alto, CA 94304, USA
TIPPARAJU, Suresh K., , Palo Alto, CA 94304, USA
KOSHKARYEV, Alexander, , Palo Alto, CA 94304, USA
ALKAN, Ozan, , Palo Alto, CA 94304, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **HÆMNING AF ATAXIA TELANGIECTASIA OG RAD3-RELATERET PROTEIN (ATR)**
- (56) Fremdragne publikationer:
WO-A1-2013/049722
US-B2- 8 147 867
US-B2- 8 841 449
US-B2- 8 912 198

DESCRIPTION

Description

Related Applications

[0001] This application claims priority to U.S. Provisional Patent Application No. 62/277,262, filed January 11, 2016, U.S. Provisional Patent Application No. 62/420,258, filed November 10, 2016, and U.S. Provisional Patent Application No. 62/444,172, filed January 9, 2017.

Field

[0002] This disclosure relates to compounds and related methods of inhibiting ataxia telangiectasia and Rad3-related protein (ATR), including methods and compounds useful for the treatment of cancer.

Background

[0003] The ataxia-telangiectasia and Rad3-related (ATR) kinase is a serine/threonine protein kinase believed to be involved in the cellular DNA damage repair processes and cell cycle signaling. ATR kinase acts with ATM ("ataxia telangiectasia mutated") kinase and other proteins to regulate a cell's response to DNA damage, commonly referred to as the DNA Damage Response ("DDR"). The DDR is believed to stimulate DNA repair, promote survival and stalls cell cycle progression by activating cell cycle checkpoints, which provide time for repair. Without the DDR, cells are much more sensitive to DNA damage and readily die from DNA lesions induced by endogenous cellular processes such as DNA replication or exogenous DNA damaging agents commonly used in cancer therapy.

[0004] The disruption of ATR function (e.g. by gene deletion) has been shown to promote cancer cell death both in the absence and presence of DNA damaging agents. Mutations of ATR have been linked to cancers of the stomach and endometrium, and lead to increased sensitivity to ionizing radiation and abolished cell cycle checkpoints. ATR is essential for the viability of somatic cells, and deletion of ATR has been shown to result in loss of damage checkpoint responses and cell death. See Cortez et al., *Science* 294: 1713-1716 (2001). ATR is also essential for the stability of fragile sites, and low ATR expression in Seckel syndrome patients results in increased chromosomal breakage following replication stress. See Casper et al., *Am. J. Hum. Genet* 75: 654-660 (2004). The replication protein A (RPA) complex recruits

ATR, and its interacting protein ATRIP, to sites of DNA damage, and ATR itself mediates the activation of the CHK1 signaling cascade. See Zou et al., *Science* 300:1542-1548 (2003). ATR, like its related checkpoint kinase ATM, phosphorylates RAD17 early in a cascade that is critical to for checkpoint signaling in DNA-damaged cells. See Bao et al., *Nature* 411: 969-974 (2001). It is believed that ATR is particularly essential in the early mammalian embryo, to sense incomplete DNA replication and prevent mitotic catastrophe.


[0005] However, while DNA-damaging chemotherapy agents and ionizing radiation (IR) therapy have provided initial therapeutic benefits to cancer patients, existing therapies have lost clinical efficacy (e.g., due to tumor cell DNA repair responses). *In vivo* effects of an ATR inhibitor and a DNA damaging agent have shown some promise in the selective treatment of cancer compared to normal cells, particularly in treating tumor cells deficient in the G1 check point control (which may depend more on the ATR for survival). US8912198 describes compounds useful as inhibitors of ATR protein kinase.

[0006] There remains a need for the development of potent and selective therapies to deliver ATR inhibitors for the treatment of cancer, either as single agents or as part of combination therapies (e.g., in combination with chemotherapy and/or radiation therapy).

Summary

[0007] Applicants have discovered novel chemical compounds useful for inhibiting ataxia-telangiectasia and Rad3-related (ATR) kinase and the treatment of cancer, and liposome formulations of certain inhibitors of ATR protein kinase having desirable properties (e.g., extended half-life in blood circulation and efficacy in treating tumors). The inventions are based in part on the discovery of certain novel compounds for inhibiting ATR protein kinase, as well as extended plasma half-lives and enhanced antitumor efficacy of certain liposomal formulations of ATR protein kinase inhibitor compounds.


[0008] In a first embodiment is provided a compound of formula (Ia), or a pharmaceutically acceptable salt thereof:



(Ia),


wherein R' is an alkyl-amino moiety comprising a tertiary alkyl-substituted amine having a pKa of 9.5 or greater.

[0009] In a second embodiment is provided a compound of the formula of Compound 6, or a pharmaceutically acceptable salt thereof:



(Compound 6).


[0010] In a third embodiment is provided a liposome composition comprising an ATR protein kinase inhibitor of formula (I), or a pharmaceutically acceptable salt thereof



(I),

wherein R is:

1. i)




wherein A¹ is either absent or C₁-C₄ alkyl, and R¹ is C₁-C₄alkylamino;

2. ii) -N(H)(C₁-C₄ alkyl)-NR^aR^b, wherein R^a and R^b are each independently C₁-C₄ alkyl;

3. iii) -(G)-NR^aR^b; wherein R^a and R^b are each independently C₁-C₄ alkyl; and G is C₁-C₄ alkyl, wherein G can be optionally substituted with C₁-C₄ alkyl; or

4. iv)





wherein R^c and R^d are each independently C₁-C₄ alkyl.

[0011] Further embodiments of the invention are set out in the appended claims.

Description of the Drawings

[0012]

FIG. 1 is a first chemical reaction scheme useful in manufacturing certain compounds disclosed herein.

FIG. 2 is a second chemical reaction scheme useful in manufacturing certain compounds disclosed herein.

FIG. 3 is a third chemical reaction scheme useful in manufacturing certain compounds disclosed herein.

FIG. 4 is a fourth chemical reaction scheme useful in manufacturing certain compounds disclosed herein.

FIG. 5 is a graph showing the blood pharmacokinetics of liposomal ATR inhibitors, as discussed in Example 8.

FIG. 6 is a graph showing the antitumor efficacy of liposomal Compound A in combinations with MM-398 in cervical MS751 xenograft model, as described in Example 9A.

FIG. 7 is a graph showing the antitumor efficacy of liposomal Compound A in combinations with MM-398 in cervical C33A xenograft model, as described in Example 9A.

FIG. 8 is a graph showing the antitumor efficacy of liposomal Compound A in combinations with MM-398 in cervical C33A xenograft model, according to Example 9A.

FIG. 9 is a graph showing the tolerability of liposomal Compound A in combination with MM-398 in cervical MS751, according to Example 9A.

FIG. 10A and **FIG. 10B** are each a graph, showing the efficacy of liposomal Compound 5 in combination with MM-398 in NCI-H2170 (**FIG. 10A**) or DMS-114 (**FIG. 10B**) mice xenograft models, as discussed in Example 10.

FIG. 11A and **FIG. 11B** are each a graph showing the Kaplan-Meier survival curves representing efficacy of liposomal Compound 5 in combination with MM-398 in NCI-H2170

(FIG. 11A) and DMS-114 (FIG. 11B) a mouse xenograft model, as discussed in Example 10.

FIG. 12A and **FIG. 12B** are each a graph showing Tolerability of liposomal Compound 5 in combination with MM-398 in NCI-H2170 (FIG. 12A) or DMS-114 (FIG. 12B) in a mouse xenograft model.

FIG. 13A and **FIG. 13B** are graphs showing the efficacy of liposomal Compound 5 in combination with MM-398 in Calu-6 (FIG. 13A) or COLO-699 (FIG. 13B) mice xenograft models, as described in Example 11.

FIG. 14A is a graph illustrating the in vitro monotherapy cell kill of Compound 6 and Compound 5 in a panel of lung cancer cell lines.

FIG. 14B is a graph illustrating the effect of Compound 5 and Compound 6 in combination with three chemotherapeutic agents (Carboplatin, Gemcitabine, Compound B).

FIG. 15 is a graph showing the IC_{50} shift values measured in Sum190PT cell line (Triple Negative Breast Cancer, TNBC) for the ATR protein kinase inhibitor Compound 6 of Example 2, with and without various concentrations of Compound B.

FIG. 16A is a graph showing the IC_{50} shift values of a combination of ATR protein kinase inhibitor Compound 6 of Example 2, measured in MDA-MB-453 TNBC cancer cell line.

FIG. 16B is a graph showing the IC_{50} shift values of a combination of ATR protein kinase inhibitor Compound A, measured in MDA-MB-453 TNBC cancer cell line.

FIG. 17 is a western blot analysis of obtained from lung cancer cells DMS-114 exposed either to Gemcitabine [16nM] or ATR inhibitor (Compound A or Compound 5 [1uM]) alone or in combination in-vitro.

FIG. 18A-B is cell death and growth assay performed combining Compound A and Gemcitabine at various concentrations in U2OS cells. The results are compared back to a prediction of cell growth and death with the combination of the two compounds (Fig. 18A). The number of living and apoptotic cells is also determined at set concentrations of Compound A (1 μ M) and Gemcitabine (0.04 μ M) in USO2 cells (Fig. 18B).

FIG. 19 shows Compound A being tested alone and in combination with SN38 at set concentrations (1 μ M and 0.2 μ M, respectively) in several lung cancer cell lines (NCI-H520 and NCI-H596) and U2OS cells, with the number of cells monitored over time.

FIG. 20 is a graph illustrating the effect of Compound A and SN38 in combination and alone in the cervical cancer cell line MS751.

FIG. 21A-C demonstrates the effects of combining Compound A or Compound 5 with Gemcitabine. Fig. 21A is a heat map illustrating the effect of combining Compound A or Compound 5 with Gemcitabine at various concentrations in U2OS, H358, and A549 cell lines. Microscopy images of cells treated with set concentrations of Compound 5 and Gemcitabine or Compound A and Gemcitabine in cells are shown (Fig. 21B). Proliferation assays of USO2 and

H358 cells with set concentrations are shown as well (Fig. 21C).

FIG. 22A-B illustrate growth curves of A549 cells with Compound A or Compound 5 in combination with SN38 (Fig. 22A) or alone (Fig. 22B).

FIG. 23 shows IC₅₀ values (μM) in several cell lines using a set concentration of Gemcitabine with varying concentrations of Compound A or Compound 5.

FIG. 24 shows a summary of the lung cancer cell lines that are responsive to Compound A or Compound 5 in combination with Gemcitabine or SN38.

FIG. 25A-B show various ATR inhibitors tested for their ability to inhibit ATR (on-target) and ATM (off-target). Inhibition is reported as IC₅₀ in nM (Fig. 25A). Additional "off-target" kinases are tested with Compound A or Compound 5 as well (Fig. 25B).

FIG. 26A-F shows on-target results with Compound A or Compound 5 in A549 lung cancer cells (Fig. 26A-B), H23 lung cancer cells (Fig. 26C-D), and DMS-114 cells (Fig. 26E-F). CHK1 S345 phosphorylation, a readout of ATR inhibition, is measured by western blot. Each compound is used with a fixed concentration of Gemcitabine as well.

FIG. 27A-B shows further on-target analysis on the HCC-70 TNBC, MDA-MB-468 TNBC, and DMS-114 cell lines. A set concentration of SN38 is used with a range of Compound A or Compound 5 concentrations. Various on-target activity parameters are tested and measured by western blot (Fig. 27A-B)

FIG. 28 shows cell cycle stage profiles of SUM149 cells 24 hours after addition of SN38 and Compound A or Compound 5.

FIG. 29 shows DMS-114 lung xenograft models used to measure the effects of Ls-Compound A or Ls-Compound 5 in combination with MM398. A set dosage of MM398 is used (5mpk) with two different dosages of Compound A or Compound 5 (20mpk or 80mpk). The effects of treatment are assayed by measuring the levels of CHK1 S345 phosphorylation.

FIG. 30A-C illustrates the effects of Ls-Compound A with MM398 in the SUM-149 cell line. The effects of the treatment are assayed by measuring phosphorylated levels of RPA2 (Fig. 30A), DNAPK, CHK1, and γH2AX (Fig. 30B). Ls-Compound 5 is also tested without the combination of MM398 (Fig. 30C).

FIG. 31 is a graph showing the efficacy of liposomal Compound 5 in combination with MM-398 in SUM-149 mice xenograft model.

FIG. 32 is a graph showing the tolerability of liposomal Compound 5 in combination with MM-398 in SUM-149 mice xenograft models

FIG. 33 Basal levels of various PD markers associated with the DNA damage response pathway, quantified by Western blot over a panel of cell lines.

FIG. 34 A schematic illustrating how the integral score is calculated for each well in the

dynamic cell viability assay.

FIG. 35 Correlation of basal MRE11 protein expression (quantified by Western blot) and the integral score, a measure of dynamic cell viability, across lung cancer cell lines exposed to the ATR inhibitor Compound 5 and/or SN38.

FIG. 36 Correlation of basal ATM protein expression (quantified by Western blot) and the integral score, a measure of dynamic cell viability, across lung cancer cell lines exposed to the ATR inhibitor Compound 5 and/or SN38.

FIG. 37 Correlation of basal NBS protein expression (quantified by Western blot) and the integral score, a measure of dynamic cell viability, across lung cancer cell lines exposed to the ATR inhibitor Compound 5 and/or SN38.

FIG. 38 Correlation of basal NBS protein expression (quantified by Western blot) and the integral score, a measure of dynamic cell viability, across p53 functionally impaired lung cancer cell lines exposed to the ATR inhibitor Compound 5 and/or SN38.

FIG. 39 Correlation of basal NBS protein expression (quantified by Western blot) and the integral score, a measure of dynamic cell viability, across p53 functionally impaired lung cancer cell lines exposed to the ATR inhibitor Compound 5 and/or SN38.

FIG. 40A-F Fold change in cancer cell line NCIH1299 pharmacodynamics markers after exposure to ATR inhibition and/or SN38.

FIG. 40A-F Fold change in cancer cell line NCIH460 pharmacodynamics markers after exposure to ATR inhibition and/or SN38.

FIG. 42A-F Fold change in cancer cell line DMS114 pharmacodynamics markers after exposure to ATR inhibition and/or SN38.

FIG. 43A-F Fold change in cancer cell line HCC70 pharmacodynamics markers after exposure to ATR inhibition and/or SN38.

FIG. 44A-F Fold change in cancer cell line MDAMB468 pharmacodynamics markers after exposure to ATR inhibition and/or SN38.

FIG. 45 Western blots of pharmacodynamics markers in cancer cell line A549 after 6 or 18 hours exposure to ATR inhibition and/or gemcitabine.

FIG. 46 Western blots of pharmacodynamics markers in cancer cell line NCIH23 after 6 or 18 hours exposure to ATR inhibition and/or gemcitabine.

FIG. 47 Western blots of pharmacodynamics markers in cancer cell line DMS114 after 6 or 18 hours exposure to ATR inhibition and/or gemcitabine.

FIG. 48 Western blots of pharmacodynamics markers in cancer cell line U20S after 6 or 18 hours exposure to ATR inhibition and/or gemcitabine.

FIG. 49 Western blots of pharmacodynamics markers in cancer cell line NCIH460 after 6 or 18 hours exposure to ATR inhibition and/or gemcitabine.

FIG. 50 Western blots of pharmacodynamics markers in cancer cell line HCC827 after 6 or 18 hours exposure to ATR inhibition and/or gemcitabine.

FIG. 51 Western blots of pharmacodynamics markers in a panel of colorectal cancer cell lines after 18 hours exposure to Compound 5 and/or SN38.


FIG. 52 Normalized quantification of phosphorylated Chk1 levels in a panel of colorectal cancer cell lines after 18 hours exposure to Compound 5 and/or SN38 (the signal for each cell line is normalized to the signal in the presence of SN38 alone).

FIG. 53 Normalized quantification of phosphorylated RPA2 levels in a panel of colorectal cancer cell lines after 18 hours exposure to Compound 5 and/or SN38 (the signal for each cell line is normalized to the signal in the presence of SN38 alone).

FIG. 54 Normalized quantification of γ H2AX levels in a panel of colorectal cancer cell lines after 18 hours exposure to Compound 5 and/or SN38 (the signal for each cell line is normalized to the signal in the presence of SN38 alone).


Detailed Description

[0013] Novel compounds for inhibiting ATR protein kinase are described by formula (I):




or a pharmaceutically acceptable salt thereof, wherein R is a moiety comprising an amine with a pK_a of greater than 7.0 (preferably greater than 8.0, and most preferably at least about 9.5), selected to provide a plasma half-life of at least about 5 hours in mice (obtained according to Example 7). Preferably, R includes an amine-substituted alkyl moiety with 4-12 carbons. R can be selected to include only a combination of tertiary-substituted amine and hydrogenated alkyl groups. R preferably further includes a tertiary-alkyl substituted amine having a pK_a of at least 7, but most preferably at least about 9.5 (e.g., a pK_a of about 9.5-10.5). Examples of compounds of formula (I), or pharmaceutically acceptable salts thereof, include Compounds 1-6 (see Examples 1-6):






Compound 1

MW: 439.53




Compound 2

MW: 453.56




Compound 3

MW: 480.59




Compound 4

MW: 440.52



Compound 5


MW: 536.70



Compound 6

MW: 536.70


[0014] The compounds of formula (I) preferably include one or more tertiary amine moieties at R selected to provide desired inhibition of ATR and/or liposome formation and stability characteristics. In some examples, R is a heterocyclic moiety comprising a first tertiary substituted nitrogen, preferably substituted with an alkylamino moiety comprising a second tertiary substituted nitrogen. In particular, compounds of formula (I) include those where R can be a moiety of the formula:



wherein A^1 is either absent or is an alkyl (e.g., C_1 - C_4 alkyl (preferably $-(CH_2)_2-$), and R^1 is lower (e.g., C_1 - C_4) alkylamino. In one example, R^1 is $(C_1$ - C_4 alkyl)- NR^aR^b , wherein R^a and R^b are each independently C_1 - C_4 alkyl, e.g., R^1 is $-(CH_2)_2-N(CH_3)(CH_3)$. In another example, R^1 is NR^aR^b , wherein R^a and R^b are each independently C_1 - C_4 alkyl, e.g., R^1 is $-N(CH_2CH_3)(CH_2CH_3)$.

[0015] In another example of formula (I), R is $-N(H)(C_1$ - C_4 alkyl)- NR^aR^b , wherein R^a and R^b are each independently C_1 - C_4 alkyl, or R is $-(G)-NR^aR^b$, wherein R^a and R^b are each independently C_1 - C_4 alkyl, wherein G is C_1 - C_4 alkyl, and wherein G can be further substituted with C_1 - C_4 alkyl.


[0016] In another example of formula (I), R can be a moiety of the formula:



wherein R^c and R^d are each independently C₁-C₄ alkyl.

[0017] Preferred examples include liposomes comprising compounds selected from the group consisting of compounds 1, 2, 3, 4, 5, or 6 above. In some examples, the compound is compound 5 or compound 6.

[0018] The compound of formula (I), in embodiments of the invention, has the chemical structure of formula (Ia), or a pharmaceutically acceptable salt thereof, wherein R' is a tertiary alkyl substituted amine having a pK_a of about 9.5 or greater:



(Ia).

Examples of compounds of formula (Ia) include compound 5 disclosed herein (e.g., Examples 1). In one embodiment of formula (Ia), R' is NR^aR^b, wherein R^a and R^b are each independently C₁-C₄ alkyl.


[0019] Liposome formulations of ATR protein kinase inhibitor compounds (e.g., as described in Example 7) can provide desirable pharmacokinetic properties such as enhanced plasma half-life of 5 hours or more in the mouse model described in Example 8. The liposomes typically comprise vesicles containing one or more lipid bilayers enclosing an aqueous interior. Liposome compositions usually include liposomes in a medium, such as an aqueous fluid exterior to the liposome. Liposome lipids can include amphiphilic lipid components that, upon contact with aqueous medium, spontaneously form bilayer membranes, such as phospholipids, for example, phosphatidylcholines. Liposomes also can include membrane-rigidifying components, such as sterols, for example, cholesterol. In some cases, liposomes also include lipids conjugated to hydrophilic polymers, such as, polyethyleneglycol (PEG) lipid derivatives that may reduce the tendency of liposomes to aggregate and also have other beneficial effects.

[0020] The liposome formulation can include a compound of formula (I) encapsulated with a polyanion (e.g., a polyanionized sugar such as sucrose octasulfate, or a suitable polyanionized

polyol) in a unilamellar vesicle formed from one or more liposome-forming lipids (e.g., hydrogenated soy phosphatidylcholine (HSPC)), cholesterol and a polymer-conjugated lipid (e.g., methoxy-poly(ethylene glycol)-1,2-distearoyl-sn-glycerol (PEG2000-DSG)). The liposome-forming lipid preferably comprises one or more phospholipids, with the ratio of the phospholipid(s) and the cholesterol selected to provide a desired amount of liposome membrane rigidity while maintaining a sufficiently reduced amount of leakage of the compound of formula (I) from the liposome.

[0021] Liposomes typically have the size in a micron or submicron range and are well recognized for their capacity to carry pharmaceutical substances, including anticancer drugs, such as irinotecan, and to change their pharmaceutical properties in various beneficial ways. Methods of preparing and characterizing pharmaceutical liposome compositions are known in the field (see, e.g., Lasic D. *Liposomes: From physics to applications*, Elsevier, Amsterdam 1993; G. Greroriadis (ed.), *Liposome Technology*, 3rd edition, vol. 1-3, CRC Press, Boca Raton, 2006; Hong et al., US Pat. 8,147,867).

[0022] In some examples (e.g., Example 7), ATR protein kinase inhibitor compositions can include a liposome comprising a ATR protein kinase inhibitor compound encapsulated in a liposome with polyanion such as a polysulfated sugar (e.g., sucrose octasulfate). Sucrosolate, a fully substituted sulfate ester of sucrose having, in its fully protonated form, the following structure:



Sucrosolate is also referred to as sucrose octasulfate or sucrooctasulfate (SOS). Methods of preparing sucrosolate in the form of various salts, e.g., ammonium, sodium, or potassium salts, are well known in the field (e.g., US Pat. 4,990,610).

[0023] The ATR protein kinase inhibitor liposomes can be prepared in multiple steps comprising the formation of a TEA containing liposome, followed by loading of an ATR protein kinase inhibitor compound (e.g., Compound A or a compound of formula (I)) into the liposome as the TEA leaves the liposome. For example, the ATR protein kinase inhibitor liposomes can be prepared by a process that includes the steps of (a) preparing a liposome containing triethylamine (TEA) as a triethylammonium salt of sucrosolate (TEA-SOS), and (b) subsequently contacting the TEA-SOS liposome with irinotecan under conditions effective for the irinotecan to enter the liposome and to permit a corresponding amount of TEA to leave the liposome (thereby exhausting or reducing the concentration gradient of TEA across the resulting liposome).

[0024] The first step can include forming the TEA-sucrosolate containing liposome by

hydrating and dispersing the liposome lipids in the solution of TEA sucrososfate. This can be performed, for example, by dissolving the lipids, including HSPC and cholesterol, in heated ethanol, and dispersing the dissolved and heated lipid solution in the TEA-sucrososfate aqueous solution at the temperature above the transition temperature (T_m) of the liposome lipid, e.g., 60 °C or greater. The lipid dispersion can be formed into liposomes having the average size of 75-125 nm (such as 80-120 nm, or in some embodiments, 90-115 nm), by extrusion through track-etched polycarbonate membranes with the defined pore size, e.g., 100 nm. The TEA-sucrososfate can include at least 8 molar equivalents of TEA to each molar equivalent of sucrososfate to obtain a solution that can have a concentration of about 0.40-0.50 N, and a pH (e.g., about 6.5) that is selected to prevent unacceptable degradation of the liposome phospholipid during the dispersion and extrusion steps (e.g., a pH selected to minimize the degradation of the liposome phospholipid during these steps). Then, the non-entrapped TEA-SOS can be removed from the liposome dispersion, e.g., by dialysis, gel chromatography, ion exchange or ultrafiltration prior to the drug encapsulation. The resulting liposomes can contain ATR protein kinase inhibitor sucrososfate. These ATR inhibitor liposomes can be stabilized by loading enough drug into the liposomes to reduce the amount of TEA in the resulting liposome composition to a level that results in less than a given maximum level of lyso-PC formation after 180 days at 4 °C, or less than a given maximum level of lyso-PC accumulation rate in the liposome composition during storage in a refrigerator at about 4 °C, or, more commonly, at 5 ± 3 °C, measured, e.g., in mg/mL/month, or % PC conversion into a lyso-PC over a unit time, such as, mol% lyso-PC/ month. Next, the TEA exchanged from the liposomes into the external medium during the loading process, along with any untrapped ATR inhibitor, is typically removed from the liposomes by any suitable known process(es) (e.g., by gel chromatography, dialysis, diafiltration, ion exchange or ultrafiltration). The liposome external medium can be exchanged for an injectable isotonic fluid (e.g. isotonic solution of sodium chloride), buffered at a desired pH.

[0025] The antitumor efficacy of various liposome formulations comprising liposome encapsulated ATR protein kinase inhibitor compounds was tested in human cervical cancer cell lines (e.g., MS751, C33A and SiHa cell lines, as shown in Example 9), and various lung cancer cell lines including lung squamous cell carcinoma cell line (e.g., NCI-H2170 cell line in Example 10), small cell lung carcinoma cell line (e.g., DMS-114 cell line in Example 10), and human Calu-6 and COLO-699 cell lines (Example 11).

[0026] Referring to FIGs. 6-9 and Example 9, a liposome formulation of the ATR inhibitor Compound A (Example 7) was tested against three human cervical cancer cell lines in mouse xenograph model (Example 9A), alone and in combination with the irinotecan liposome formulation MM398 (Example 9B). Greater tumor volume was observed over time for the liposomal Compound A formulation of Example 7 compared to the control experiment for 2 of the 3 cervical cancer cell lines (MS571 and C33A). However, administering the irinotecan liposome MM398 (Example 9B) in combination with the Compound A liposome formulation (Example 7) resulted in greater suppression of tumor volume in all three cervical cancer cell lines than either administration of MM398 alone or liposomal Compound A alone.

[0027] Referring to FIGs. 10A-10B and FIGs. 11A-11B, a liposome formulation of the ATR inhibitor Compound 5 of formula (I) and formula (Ia) (the compound of Example 1 formulated as a liposome as described in Example 7) was tested against two lung cancer cell lines in a mouse xenograph model (Example 10), alone and in combination with the irinotecan liposome formulation MM398 (Example 9B). Referring to FIG. 10A and FIG. 10B, the administration of the liposome formulation of Compound 5 reduced tumor volume in each cell line tested compared to the control experiment in Example 10, the combination of MM398 and the Compound 5 liposome composition of Example 7 reduced the tumor volume in the mouse model to a greater degree than either compound administered independently of the other. Similarly, the Kaplan-Meier survival curves presented in Example 10 (FIG. 11A and FIG. 11B) demonstrate increased survival in mouse lung cancer xenograph testing when a combination of both the irinotecan liposome MM398 of Example 9B was administered in combination with the Compound 5 liposome formulation of Example 7 using two different cell lines.

[0028] Referring to FIG. 12A and FIG. 12B, the tolerability of various liposome formulations of ATR protein kinase inhibitor compounds was assessed in Example 10. Referring to FIG. 12A, the decline in mouse bodyweight tested in the NCI-H2170 mouse xenograph model was lowest over time for the liposome formulation of Compound 5 (Example 7), compared to the irinotecan liposome MM398 (Example 9B), the control or the combination of the liposome formulation of Compound 5 in combination with MM398. Referring to FIG. 12B, the decline in mouse bodyweight tested in the DMS-114 mouse xenograph model was lowest over time for the combination of the liposome formulation of Compound 5 in combination with MM398, compared to the liposome formulation of Compound 5 (Example 7), or the irinotecan liposome MM398 (Example 9B) administered independently.

[0029] Referring to FIG. 13A and FIG. 13B, administering a combination of the irinotecan liposome MM398 (Example 9B) with the liposome formulation of ATR protein kinase inhibitor Compound 5 resulted in the greatest reduction in tumor volume in both the Calu-6 and COLO699 cell lines in mice xenograft models, compared to the control, the administration of MM398 irinotecan liposome alone, administration of the Compound A liposome formulation (Example 7) or the combination of the MM398 irinotecan liposome (Example 9B) with the Compound A liposome formulation (Example 7).


Examples

[0030] The following examples illustrate some embodiments of the invention and examples of the disclosure. The examples and preparations which follow are provided to enable those skilled in the art to more clearly understand and to practice these and other embodiments present invention and examples of the disclosure. Comparative subject matter is marked *.

[0031] ATR peptide can be expressed and isolated using a variety of methods known in the literature (see e.g., Ünsal-Kaçmaz et al, PNAS 99: 10, pp 6673-6678, May 14, 2002; see also Kumagai et al. Cell 124, pp 943-955, Mar. 10, 2006; Ünsal-Kacmaz et al. Molecular and

Cellular Biology, February 2004, p 1292-1300; and Hall-Jackson et al. Oncogene 1999, 18, 6707-6713).

[0032] Comparative compound A can be obtained by methods disclosed (for example) in publication WO2010/071827A1 (published June 24, 2010). The structure of Compound A is as follows:







Compound A*.


[0033] Various compounds of formula (I) can be prepared as described herein, and summarized in the table below.

Table 1: Selected Compounds and Comparative Compounds (marked *) of Formula (I)

| <p>(I)</p> | | |
|-------------|---|---------|
| Compound | R | Example |
| Compound 5 | | 1 |
| Compound 6 | | 2 |
| Compound 2* | | 3 |


| Compound | R | Example |
|-------------|---|---------|
| |  | |
| Compound 4* |  | 4 |
| Compound 3* |  | 5 |
| Compound 1* |  | 6 |

[0034] The examples 1, 2, 3 and 6 were prepared in a one-pot Suzuki cross-coupling using boronic ester generated in-situ shown in Scheme 1 in FIG. 1. Referring to FIG. 1, the synthesis of Intermediate 3: 1-bromo-4-(2-bromoethylsulfonyl)benzene can be obtained as described below.




[0035] To a solution of Intermediate 2 (35 g, 133 mmol) in DCM (400 mL) was added PBr_3 (40 g, 146 mmol) at 0 °C dropwise. Then the mixture was stirred at rt overnight. Water (15 mL) was added to quench the reaction. Then the resultant was washed with water (120 mL) and brine (120 mL). The organic phase was concentrated to afford 20 g of crude 3 as yellow oil, which was used in the next step without further purification.

[0036] Referring to FIG. 1, the synthesis of intermediate 2 can be performed as described below:




[0037] To a solution of Intermediate 1 (45 g, 194 mmol) in DCM (500 ml) was added m-CPBA (134 g, 776 mmol) at rt in several batches. Then the mixture was stirred at rt overnight. The reaction mixture was filtered and DCM (500 ml) was added to wash the solid. The filtrate was washed with NaOH (1M, 300 mL X 3) and brine (300 mL). The organic layer was concentrated to dryness to afford 36 g of 2 (70 %) as white solid.

[0038] Referring to FIG. 1, the synthesis of intermediate 1 can be performed as described below.



[0039] To a solution of 4-bromobenzenethiol (45 g, 238 mmol) in MeCN (600 ml) was added K_2CO_3 (60 g, 476 mmol) and NaI (36 g, 238 mmol). The mixture was stirred at rt for 10 minutes. Then 2-bromoethanol was added dropwise. After addition, the mixture was stirred at rt overnight. The reaction mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica-gel column to afford 45 g of 1 (81%) as light yellow oil.

[0040] Block B can be prepared according to scheme 2 in FIG. 2. Referring to FIG. 2, the synthesis of Block B can be performed as described below.




[0041] To a solution of Intermediate 6 (6.0 g, 27.4 mmol) in DMSO (30 mL) was added CDI (8.9 g, 54.8 mmol), DIPEA (3.8 g, 30.1 mmol) and DMAP (0.17 g, 1.37 mmol). The solution was stirred at rt for 4 hours. Aniline (2.5 g, 27.4 mmol) was added and the mixture was stirred at rt overnight. Water was added and the formed solid was collected by filtration. The crude product was purified by silica-gel column to afford 2.5 g of Block B (31 %) as yellow solid.

[0042] LC-MS (M+1): 293.2; 1H NMR (400 MHz, DMSO- d_6) δ 10.28 (s, 1H), 8.42 (s, 1H), 7.78 (d, J = 8.0 Hz, 2H), 7.74 (s, 2H), 7.36 (t, J = 8.0 Hz, 2H), 7.13 (t, J = 7.6 Hz, 1H).

[0043] Referring again to FIG. 2, the synthesis of Intermediate 6 can be performed as described below.






Intermediate 6

[0044] To the solution of methyl 3-amino-6-bromopyrazine-2-carboxylate (10.0 g 43.1 mmol) in MeOH (70 mL) was added a solution of LiOH (9.0 g, 215 mmol) in water (70 mL). The mixture was stirred at 90 °C for 3 hours. The reaction mixture was cooled to rt and acidified to PH = 4~5 with HCl (2 M). The mixture was filtered to afford 7.4 g of 6 (79 %) as yellow solid.

[0045] LC-MS (M+1): 218.0; ¹H NMR (400 MHz, DMSO-d₆) δ 8.39 (s, 1H), 7.59 (br, 2H).

Example 1: Synthesis of Compound 5 (3-amino-6-(4-((2-(4-(2-(dimethylamino)ethyl)piperidin-1-yl)ethyl)sulfonyl)phenyl)-N-phenylpyrazine-2-carboxamide)

[0046]



Exact Mass: 536.26; Molecular Weight: 536.70; Compound 5; More basic; 143 mg; Yield 8.2%; pK_a 10.00.


[0047] To a solution of 2-(1-(2-((4-bromophenyl)sulfonyl)ethyl)piperidin-4-yl)-N,N-dimethylethan-1-amine (Block A1) (261 mg, 0.648 mmol) in anhydrous dioxane (3 ml) was added potassium acetate (191 mg, 1.944 mmol) and Bis(pinacolato) diborane (246mg, 0.971 mmol), the reaction vessel was degassed by repeating vacuum/ nitrogen cycle and then was added Pd(dppf)₂Cl₂. CH₂Cl₂ (53mg, 0.0648mmol), again was degassed and the reaction was heated at 90 °C for 2 hours under nitrogen. The reaction was then cooled down to room temperature and was added 3-amino-6-bromo-N-phenylpyrazine-2-carboxamide (Block B), 2M K₂CO₃ (1 ml) and degassed and purged with nitrogen. The Pd(PPh₃)₄ (75 mg, 0.0648 mmol) was added. The reaction was heated at 100 °C for 4 hrs. The reaction was cooled to room temperature and diluted with ethyl acetate and washed with brine three times and organic layer was dried over Na₂SO₄. Upon removal of solvent on the rotovap, a dark oil residual crude product was obtained and it was purified on silica-gel column chromatography (Reveleris Flash Chromatography System) using 0-15% methanol in dichloromethane as an eluent. Desired

product was obtained as a yellow solid (149 mg, yield 43%). MS (M+H)⁺ 537; ¹H NMR (400MHz, DMSO-d₆): δ 10.45 (s, 1H), 9.03 (s, 1H), 8.49 (d, 2H, 6.8Hz), 7.95 (d, 2H, 6.8Hz), 7.88 (s, br, 2H), 7.81 (d, 2H, 8.8Hz), 7.40 (t, 2H, 7.2Hz), 7.18 (t, 1H, 7.2Hz), 3.53 (t, 2H, 7.2Hz), 2.64 (d, 2H, 11.6 Hz), 2.55 (t, 2H, 7.2Hz), 2.05 (m, 2H), 1.98 (s, 6H), 1.17 (m, 2H), 1.42 (d, 2H, 12.0Hz), 1.18 (m, 3H), 0.78 (m, 2H).

[0048] High resolution mass (Thermo Scientific™ Q Exactive™ hybrid quadrupole-Orbitrap mass spectrometer): Calculated for C₂₈H₃₆N₆O₃S + Proton (1.00728) = 537.2642; theoretical m/z of single charged ion: 537.2642; Found: 537.2636.

Example 2: Synthesis of Compound 6 (3-amino-6-(4-((2-(4-(diethylamino)piperidin-1-yl)ethyl)sulfonyl)phenyl)-N-phenylpyrazine-2-carboxamide)

[0049]




Exact Mass: 536.26; Molecular Weight: 536.70; Compound 6; More basic; 53 mg; Yield 11.1%; pK_a 9.81.

[0050] The example 2 was prepared in a similar fashion using Block A2 (1-(2-((4-bromophenyl)sulfonyl)ethyl)-N,N-diethylpiperidin-4-amine), a yellow solid was obtained (52mg, yield 24%). MS (M+H)⁺ 537; ¹H NMR (400MHz, DMSO-d₆): δ 10.45 (s, 1H), 9.05 (s, 1H), 8.51 (d, 2H, 8.4Hz), 7.94 (d, 2H, 8.8Hz), 7.87 (s, br, 2H), 7.79 (d, 2H, 8.8Hz), 7.42 (t, 2H, 8.4Hz), 7.18 (t, 1H, 7.2Hz), 3.54 (t, 2H, 6.4Hz), 2.65 (d, 2H, 11.2 Hz), 2.56 (t, 2H, 6.4Hz), 2.20 (q, 4H, 6.8 Hz), 1.71 (t, 2, 10.4 Hz), 1.33 (d, 2H, 12.4Hz), 0.85 (qd, 2H, 12.4 Hz), 0.74(t, 6H, 7.2 Hz).

[0051] 1-(2-((4-bromophenyl)sulfonyl)ethyl)-N,N-diethylpiperidin-4-amine (Block A2) was prepared in a similar fashion using corresponding 4-diethylaminopiperidine. A colorless oil was obtained (661mg, 47% yield), MS (M+H)⁺ 403, 405.

Example 3: Synthesis of Comparative Compound 2* (3-amino-6-(4-((2-(diethylamino)ethyl)sulfonyl)phenyl)-N phenylpyrazine-2-carboxamide)

[0052]




Exact Mass: 453.18; Molecular Weight: 453.56; Compound 2; Less basic; 98 mg; Yield 7.7%; pK_a 7.46.

[0053] The compound of example 3 was prepared in a similar fashion using intermediate Block A3, 2-((4-bromophenyl)sulfonyl)-N,N-diethylethan-1-amine, a yellow solid was obtained (98 mg, yield 11%). MS (M+H)⁺ 454; ¹H NMR (400MHz, DMSO-d₆): δ 10.46 (s, 1H), 9.05 (s, 1H), 8.51 (d, 2H, 6.8Hz), 7.98 (d, 2H, 6.8Hz), 7.88 (s, br, 2H), 7.81 (d, 2H, 8.8Hz), 7.41 (t, 2H, 7.2Hz), 7.17 (t, 1H, 7.2Hz), 3.48 (dd, 2H, 6.8Hz), 2.73 (m, 2H), 2.33 (q, 4H, 6.8Hz), 0.81 (t, 6H, 6.8Hz).

[0054] Block A3 (2-((4-bromophenyl)sulfonyl)-N,N-diethylethan-1-amine) was prepared in a similar fashion using corresponding 4-diethylamine. A colorless oil was obtained (1.42g, 73% yield), MS (M+H)⁺ 320, 322

Example 4: Synthesis of Comparative Compound 4* (3-amino-6-(4-(((2-(dimethylamino)ethyl)- λ^2 -azanyl)sulfonyl)phenyl)-N-phenylpyrazine-2-carboxamide)

[0055]




Exact Mass: 439.16; Molecular Weight: 439.51; Compound 4; pK_a 8.36.

[0056] Compound 4 can be prepared as shown in scheme 3 in FIG. 3. To a mixture of Block B (150 mg, 0.51 mmol), Block F (153 mg, 0.56 mmol) and Na₂CO₃ (216 mg, 2.0 mmol) in toluene / ethanol / water (2 mL / 2 mL / 2 mL) was added Pd(dppf)Cl₂ (30 mg). The mixture was stirred at 75 °C under argon atmosphere for 4 hours. The reaction mixture was

concentrated to dryness. The residue was purified by prep-HPLC to afford 100 mg of TM4 (45 %) as white solid.

[0057] LC-MS (M+1): 441.4; $^1\text{H NMR}$ (400 MHz, CD_3OD) δ 8.88 (s, 1H), 8.33 (dd, $J = 6.8$ Hz, 1.6 Hz, 2H), 8.00 (dd, $J = 6.8$ Hz, 1.6 Hz, 2H), 7.80 (dd, $J = 8.4$ Hz, 1.2 Hz, 2H), 7.41 (t, $J = 7.6$ Hz, 2H), 7.19 (t, $J = 7.6$ Hz, 1H), 3.10 (t, $J = 6.4$ Hz, 2H), 2.73 (t, $J = 6.4$ Hz, 2H), 2.46 (s, 6H).

[0058] Referring again to Scheme 3 in FIG. 3, the synthesis of Block F can be performed as described below.




Intermediate 7

Block F

To a solution of Intermediate 7 (10.0 g, 32.6 mmol) in THF (200 mL) at -78 °C under argon atmosphere was added $\text{B}(\text{i-Pr})_3$ (30.6 g, 163 mmol). Then $n\text{-BuLi}$ (2.5 M, 65 mL) was added dropwise. The mixture was stirred at -78 °C for 2 hours and at then rt for another 16 hours. Water was added to quench the reaction. The mixture was concentrated to dryness. The residue was purified by prep-HPLC to afford 5.2 g of **Block F** (59 %) as white solid.

[0059] LC-MS (M+1): 273.4; $^1\text{H NMR}$ (400 MHz, DMSO-d_6) δ 7.90-7.45 (m, 4H), 2.91 (s, 2H), 2.69 (t, $J = 6.4$ Hz, 2H), 2.18 (t, $J = 6.8$ Hz, 2H), 2.03 (s, 6H).

[0060] Referring again to Scheme 3 in FIG. 3, the synthesis of Intermediate 7 can be performed as described below.




Intermediate 7

To a solution of 4-bromobenzene-1-sulfonyl chloride (20 g, 78.3 mmol) in DCM (300 mL) at 0 °C was added TEA (22 mL, 158 mmol), followed by N,N' -dimethylethane-1,2-diamine (8.3 g, 94.0 mmol). The resulting solution was stirred at rt for 1 hour and then diluted with DCM (300 mL). The solution was washed with water (200 mL) and brine (200 mL). The organic layer was concentrated to dryness. The residue was purified by silica-gel column to afford 17.0 g of 7 (71 %) as off-white solid.


Example 5: Synthesis of Comparative Compound 3* (3-amino-6-(4-((2-(4-methylpiperazin-1-yl)ethyl)sulfonyl)phenyl)-N-phenylpyrazine-2-carboxamide)

[0061]




Exact Mass: 480.19; Molecular Weight: 480.59; Compound 3; pK_a 7.73.

[0062] The compound of Example 5 can be obtained by Scheme 4 shown in FIG. 4. Intermediate 5 in Scheme 4 can be obtained as described below.



[0063] To a solution of Intermediate 4 (1.7 g, 5.0 mmol) in THF (30 mL) at -78 °C under argon atmosphere was added B(i-Pr)_3 (4.7 g, 25 mmol). Then n-BuLi (2.5 M, 10 mL) was added dropwise. The mixture was stirred at -78 °C for 2 hours and at then rt for another 16 hours. Water was added to quench the reaction. The mixture was concentrated to dryness. The residue was purified by prep-HPLC to afford 300 mg of 5 (19 %) as white solid.


[0064] Referring again to FIG. 4, Intermediate 4 in Scheme 4 can be obtained as described below.



[0065] To a solution of Intermediate 3 (20 g, 60 mmol) in MeCN (300 mL) was added 1-methylpiperazine (9.0 g, 90 mmol) and K_2CO_3 (16.6 g, 120 mmol). The mixture was stirred at rt overnight. The reaction mixture was filtered and the filtrate was concentrated to dryness. The residue was purified by silica-gel column to afford 15 g of 4 (71 %) as pale solid.

Example 6: Synthesis of Comparative Compound 1* (3-amino-6-(4-((1-(dimethylamino)propan-2-yl)sulfonyl)phenyl)-N-phenylpyrazine-2-carboxamide)

[0066]



Exact Mass: 439.17; Molecular Weight: 439.53; Compound 1; Less basic; 427 mg; Yield 12.9%; pK_a 7.04.

[0067] The compound of Example 6 was prepared according to the procedure for preparing Compound 1 given in J. Med. Chem. 2011, 54, 2320 (supplementary materials) except using 1-bromo-4-(2-bromoethylsulfonyl)benzene. A yellow solid was obtained (427 mg, yield 11%).

MS (M+H)⁺ 440; ¹H NMR (400MHz, DMSO-d₆): δ 10.46 (s, 1H), 9.05 (s, 1H), 8.52 (d, 2H, 6.8Hz), 7.93 (d, 2H, 6.8Hz), 7.89 (s, br, 2H), 7.82 (d, 2H, 8.0Hz), 7.40 (t, 2H, 7.2Hz), 7.18 (t, 1H, 7.2Hz), 3.53 (t, 2H, 7.2Hz), 2.64 (d, 2H, 11.6 Hz), 2.55 (t, 2H, 7.2Hz), 2.05 (m, 2H), 1.98 (s, 6H), 1.17 (m, 2H), 1.42 (d, 2H, 12.0Hz), 1.18 (m, 3H), 0.78 (m, 2H).

Example 7: Preparation of liposomes with entrapped triethylammonium SOS salts and loading of ATRi into the liposome.

[0068] Sodium sucrose octasulfate (equivalent weight 144.8) is a sodium salt of sucrose derivate in which all hydroxyl groups have formed sulfuric acid esters. Sixty gram of sucrose octasulfate (SOS) sodium salt were dissolved in in 150 ml of di water, heated with shaking (swirling) of a 50 °C water bath. The solution was passed through a column packed with sulfonated polystyrene divinylbenzene copolymer cation exchange resin beads (Dowex 50Wx8-100-200 mesh, Dow Chemical Co.). The column was pre-equilibrated with aqueous 3-3.6 M HCl to bring the resin in the hydrogen form and washed with deionized water until the outflow shows conductivity of <1 μS/cm. The eluent was monitored using a conductivity detector. The SOS fraction was collected corresponding to conductivity peak and immediately titrated with neat triethylamine (TEA) solution to pH 6-6.5. The solution was analyzed for residual sodium by potentiometry using a sodium-sensitive electrode and for SOS concentration using a refractometer. The solution having residual sodium less than 0.25 % was diluted with di water to final SOS concentration 1.1 M and then sterile-filtered using Millipore 0.22 μm Steri-Top filter.

[0069] Cholesterol (Chol) was purchased from Avanti Polar Lipids, Alabaster, Ala, USA, hydrogenated soy phosphocholine (HSPC) and methoxy-poly(ethylene glycol)-1,2-distearoyl-sn-glycerol (PEG2000-DSG) were obtained from Lipoid GmbH, Ludwigshafen, Germany.

[0070] Chol, HSPC and PEG2000-DSG were co-dissolved in 100% ethanol (200-proof, Sigma cat#: 459828) at a molar ratio of 3:2:0.15 at 65 °C. The solution of TEA-SOS (10 times the volume of added ethanol) was mixed with the lipid solution at 60-65 °C and stirred at this temperature until a homogeneous milky suspension of multilamellar vesicles was formed. This suspension was extruded 3 times through 5 stacked polycarbonate track-etched filters (Corning Nuclepore) with the pore size of 100 nm using argon pressure extruder (Lipex Biomembranes) at 60-65 °C, and resulting unilamellar liposomes were quickly chilled in ice and then stored at 4-6 °C before use. Concentration of phospholipids was measured by phosphate assay and particle diameter was recorded on a Malvern Nanosizer.

[0071] Prior to drug loading, the TEA-SOS gradient was created by removing excess of untrapped TEA-SOS using gel-chromatography (Sephacrose CL-4B, Pharmacia). Osmolarity of liposomes was balanced using 50% dextrose solution. Final dextrose concentration was 15%.

[0072] ATR inhibitors were dissolved in 15% dextrose solutions in di water by titration with 1 M HCl and heating at 45 °C and then filtered with 0.2 micron NALGENE 13mm Syringe Filters. Drug concentration in the solution was detected by HPLC. A stock solution of ATR inhibitors containing 9-10 mg/mL of drugs was added to the liposomes at a drug/lipid ratio of 800 mg/mmol phospholipids and pH was adjusted to pH 6.5 with 1M HEPES buffer and 0.1 N NaOH.

[0073] The liposome-drug mixture was incubated with occasional agitation for 30 minutes at 65 °C. The incubation mixture was quickly cooled down and incubated for 10 minutes at 0 °C, then allowed to attain ambient temperature. Un-encapsulated drug was removed by gel chromatography on Sephadex G-25 (Amersham Pharmacia) eluted with HBS-6.5 buffer (5 mM 2-(4-(2-hydroxyethyl)-piperazino)-ethylsulfonic acid (HEPES), 144 mM NaCl, pH 6.5). Liposome fractions eluted in the void volume were combined, sterilized by 0.2 micron filtration, and stored at 4-6 °C before use. The liposomes were characterized by lipid concentration, drug concentration, and particle size (Table 2). All ATR inhibitors showed good loading efficacy except Compound 1, which at drug to lipid ration higher than 400 g/mol formed aggregates and participated.

Table 2. Characterization of Liposome loaded with ATR inhibitors.

| Example 8: General description of PK study of liposomal ATRi | | | | |
|---|---------------------------------|--------------------------------|------------------------|-------------------------------|
| ATR inhibitor | Drug/lipid ratio before loading | Drug/lipid ratio after loading | Loading efficiency (%) | Liposome size, (mean ± SD) nm |
| Compound 6 | 743 ± 89 | 550 ± 64 | 74 ± 0.3 | 119 ± 28 |
| Compound 2 | 642 ± 10 | 639 ± 22 | 100 ± 5 | 130 ± 34 |

| Example 8: General description of PK study of liposomal ATRi | | | | |
|---|---------------------------------|--|------------------------|-----------------------------------|
| ATR inhibitor | Drug/lipid ratio before loading | Drug/lipid ratio after loading | Loading efficiency (%) | Liposome size, (mean \pm SD) nm |
| Compound 5 | 732 \pm 58 | 619 \pm 110 | 84 \pm 8 | 120 \pm 29 |
| Compound 3 | 646 \pm 28 | 628 \pm 6 | 97 \pm 5 | 120 \pm 27 |
| Compound A | 736 \pm 57 | 716 \pm 41 | 97 \pm 2 | 121 \pm 26 |
| Compound 1 | 600-800 | Not loadable. Formed aggregates and precipitated | | |
| | 379 | 369 | 98 | 122 \pm 24 |

[0074] FIG. 5 is a graph showing the blood pharmacokinetics of liposomal ATR inhibitors. Liposomal formulations of ATR inhibitors were prepared as described in Example 7. The liposomes were administered intravenously at a dose of 20 mg drug/kg to three 7-9-week-old female CD-1 mice (Charles River) (body weight about 25 g). Blood samples were collected into lithium heparin tubes by bleeding from saphenous vein at 0.08, 1.5, 4, 8 and 24 h time points. Plasma was separated from the cell fraction by centrifugation at 10000 rpm for 5 min. Drugs were extracted by incubation of plasma samples with 200 μ l of 1% acidic acid in methanol (1%Ac/MeOH) to at least 2 hours at -80 $^{\circ}$ C. Plasma proteins were spun down by centrifugation at 15000 rpm for 20 min. Then 75 μ l of supernatant was transferred to HPLC vials (Thermo Scientific, Cat# C4011-LV1) and additional 75 μ l of 1% Ac/MeOH were added. Drug content was analyzed by HPLC with each sample measured in duplicate. The data were expressed as the % injected dose plotted against post injection time. As shown in FIG. 5, liposomal formulation of Compound 1, Compound 3 and Compound 2 were unstable in the circulation with liposomal Compound 1 and Compound 3 un-detectable by HPLC analysis at 24 hour time point and liposomal Compound 2 un-detectable already at 8 h time point. Two liposomal formulations Compound 6 and Compound 5 have good circulation longevity with above 16% of the initial injected dose after 24 hours. Table 2 below summarizes the blood PK curves. Liposomal Compound 6 and Compound 5 show the highest plasma half-lives compare to other liposomal examples.

Table 3. Pharmacokinetics parameters of liposomal ATR inhibitors

| Drug | C _{max} (mg/ml) | AUC (mg/ml*h) | V _d (ml) | Cl (ml/h) | T _{1/2} (h) | % of ID after 24 h |
|---------------|--------------------------|---------------|---------------------|-----------|----------------------|--------------------|
| Ls-Compound 3 | 0.271 | 1.659 | 2.020 | 0.330 | 4.24 | 0.0 |
| Ls-Compound 6 | 0.303 | 3.021 | 1.655 | 0.166 | 6.91 | 16.4 |
| Ls-Compound 2 | 0.080 | 0.160 | 5.806 | 2.890 | 1.39 | 0.0 |

| Drug | C _{max} (mg/ml) | AUC (mg/ml*h) | V _d (ml) | Cl (ml/h) | T _{1/2} (h) | % of ID after 24 h |
|-------------------|-----------------------------|------------------|------------------------|--------------|-------------------------|-----------------------|
| Ls- Compound 5 | 0.227 | 2.504 | 2.012 | 0.182 | 7.65 | 17.2 |
| Ls- Compound A | 0.237 | 1.902 | 0.712 | 0.089 | 5.56 | 9.0 |
| Ls- Compound 1 | 0.053 | 0.043 | 6.358 | 7.841 | 0.56 | 0.0 |

Example 9A: *In vivo* antitumor efficacy and tolerability of LS-Compound A prepared using TEA.SOS against cervical cancer xenografts in mice

[0075] FIG. 6 is a graph showing the antitumor efficacy of liposomal Compound A in combinations with MM-398 in cervical MS571 xenograft model, as described in Example 9A.

[0076] FIG. 7 is a graph showing the antitumor efficacy of liposomal Compound A in combinations with MM-398 in cervical C33A xenograft model, as described in Example 9A.

[0077] FIG. 8 is a graph showing the antitumor efficacy of liposomal Compound A in combinations with MM-398 in cervical C33A xenograft model, according to Example 9A.

[0078] FIG. 9 is a graph showing the tolerability of liposomal Compound A in combination with MM-398 in cervical MS751, according to Example 9A.

[0079] Antitumor efficacy of liposomes loaded with an ATR inhibitor Compound A (Ls-Compound A) in combinations with MM-398 (liposomal Irinotecan) was studied in the model of human cervical MS751, C33A and SiHa cell line. The cells were obtained from American Type Culture Collection (Rockville, MD) and propagated in RPMI medium supplemented with 10% fetal calf serum, 50 U/mL penicillin G, and 50 µg/mL of streptomycin sulfate at 37 °C, 5% CO₂ as recommended by the supplier. NCR nu/nu homozygous athymic male nude mice (4-5 week old, weight at least 16 g) were obtained from Charles River. The mice were inoculated subcutaneously in the right flank with 0.1 mL of the suspension containing 5 x 10⁶ cells suspended in PBS supplemented with 30% Matrigel. When tumors achieved the size between 150 mm³ and 350 mm³ the animals were assigned to the treatment groups according to the following method. The animals were ranked according to the tumor size, and divided into 6 categories of decreasing tumor size. Four treatment groups of 10 animals/group were formed by randomly selecting one animal from each size category, so that in each treatment group all tumor sizes were equally represented.

[0080] The animals received four tail vein injections, at the intervals of 7 days, of the following preparations: 1) Control (HEPES-buffered saline pH 6.5); 2) MM-398 at dose 2 or 5 mg/kg per

injection; 3) Liposomal Compound A at 20 or 60 mg/kg per injection; 4) MM-398 followed by injections of liposomal Compound A with a 24 h interval. Liposomes for injections were prepared as described in Example 7. The animal weight and tumor size were monitored twice weekly. The tumor progression was monitored by palpation and caliper measurements of the tumors along the largest (length) and smallest (width) axis twice a week. The tumor sizes were determined twice weekly from the caliper measurements using the formula (Geran, R.I., et al., 1972 Cancer Chemother. Rep. 3:1-88):

$$\text{Tumor volume} = [(\text{length}) \times (\text{width})^2] / 2$$

[0081] To assess treatment-related toxicity, the animals were also weighted twice weekly. The animals were observed for 60 days following tumor inoculation. When the tumors in the group reached 10% of the mouse body weight, the animals in the group were euthanized. Average tumor volumes across the groups were plotted together and compared over time. As shown in FIGs. 6, 7, and 8, combination of liposomal ATR inhibitor Compound A with MM-398 has a significantly stronger antitumor effect compare to MM-398 and liposomal Compound A alone in all three xenograft models. The treatment related toxicity was assessed by the dynamics of animals' body weight (FIG. 9). Neither group revealed any significant toxicity. The weight of the animals in all treated groups was comparable to the control group and was consistently increasing. Thus, the liposome formulation of ATR inhibitor Compound A showed increased antitumor activity in the studied tumor models without an appreciable increase in toxicity.

Example 9B: MM-398 irinotecan liposome manufacturing

[0082] The MM398 used in Example 9A and elsewhere herein is an irinotecan liposome that can be prepared in a multi-step process. First, lipids are dissolved in heated ethanol. The lipids can include DSPC, cholesterol and MPEG-2000-DSPE combined in a 3:2:0.015 molar ratio. Preferably, the liposomes can encapsulate irinotecan sucrose octasulfate (SOS) encapsulated in a vesicle consisting of DSPC, cholesterol and MPEG-2000-DSPE combined in a 3:2:0.015 molar ratio. The resulting ethanol-lipid solution is dispersed in an aqueous medium containing substituted amine and polyanion under conditions effective to form a properly sized (e.g. 80-120 nm) essentially unilamellar liposome containing the substituted amine (in the ammonium form) and polyanion encapsulated within a vesicle formed from the dissolved lipids. The dispersing can be performed, e.g., by mixing the ethanolic lipid solution with the aqueous solution containing a substituted amine and polyanion at the temperature above the lipid transition temperature, e.g., 60-70 °C, and extruding the resulting hydrated lipid suspension (multilamellar liposomes) under pressure through one or more track-etched, e.g. polycarbonate, membrane filters with defined pore size, e.g. 50 nm, 80 nm, 100 nm, or 200 nm. The substituted amine can be triethylamine (TEA) and the polyanion can be sucrose octasulfate (SOS) combined in a stoichiometric ratio (e.g., TEAsOS) at a concentration of about 0.4-0.5N. All or substantially all non-entrapped TEA or SOS is then removed (e.g., by gel-filtration, dialysis or ultrafiltration) prior to contacting the liposome with irinotecan under conditions effective to allow the irinotecan to enter the liposome in exchange with TEA leaving

the liposome. The conditions can include one or more conditions selected from the group consisting of: addition of the osmotic agent (e.g., 5% dextrose) to the liposome external medium to balance the osmolality of the entrapped TEA-SOS solution and/or prevent osmotic rupture of the liposomes during the loading, adjustment and/or selection of the pH (e.g. to 6.5) to reduce the drug and/or lipid degradation during the loading step, and increase of the temperature above the transition temperature of the liposome lipids (e.g., to 60-70 °C) to accelerate the transmembrane exchange of TEA and irinotecan. The loading of irinotecan by exchange with TEA across the liposome preferably continues until all or substantially all of the TEA is removed from the liposome, thereby exhausting its concentration gradient across the liposome. Preferably, the irinotecan liposome loading process continues until the gram-equivalent ratio of irinotecan to sucroctasulfate is at least 0.9, at least 0.95, 0.98, 0.99 or 1.0 (or ranges from about 0.9-1.0, 0.95-1.0, 0.98-1.0 or 0.99-1.0). Preferably, the irinotecan liposome loading process continues until the TEA is at least 90%, at least 95%, at least 98%, at least 99% or more of the TEA is removed from the liposome interior. The irinotecan can form irinotecan sucrosolate within the liposome, such as irinotecan and sucrose octasulfate in a molar ratio of about 8:1. Next, any remaining extra-liposomal irinotecan and TEA is removed to obtain the irinotecan liposome using, e.g., gel (size exclusion) chromatography, dialysis, ion exchange, or ultrafiltration methods. The liposome external medium is replaced with injectable, pharmacologically acceptable fluid, e.g., buffered isotonic saline. Finally, the liposome composition is sterilized, e.g., by 0.2-micron filtration, dispensed into dose vials, labeled and stored, e.g., upon refrigeration at 2-8 °C, until use. The liposome external medium can be replaced with pharmacologically acceptable fluid at the same time as the remaining extra-liposomal irinotecan and TEA is removed. The extra-liposomal pH of the composition can be adjusted or otherwise selected to provide a desired storage stability property (e.g., to reduce formation of the lyso-PC within the liposome during storage at 4 °C over 180 days), for example by preparing the composition at a pH of about 6.5-8.0, or any suitable pH value there between (including, e.g., 7.0-8.0, and 7.25).

[0083] DSPC, cholesterol (Chol), and PEG-DSPE were weighed out in amounts that corresponded to a 3:2:0.015 molar ratio, respectively (e.g., 1264 mg/412.5 mg/22.44 mg). The lipids were dissolved in chloroform/methanol (4/1 v/v), mixed thoroughly, and divided into 4 aliquots (A-D). Each sample was evaporated to dryness using a rotary evaporator at 60 °C. Residual chloroform was removed from the lipids by placing under vacuum (180 µtorr) at room temperature for 12 h. The dried lipids were dissolved in ethanol at 60 °C, and pre-warmed TEAsOS of appropriate concentration was added so that the final alcohol content was 10% (v/v). The lipid concentration was 75 mM. The lipid dispersion was extruded at about 65 °C through 2 stacked 0.1 µm polycarbonate membranes (Nucleopore) 10 times using Lipex thermobarrel extruder (Northern Lipids, Canada), to produce liposomes with a typical average diameter of 95-115 nm (determined by quasielastic light scattering). The pH of the extruded liposomes was adjusted with 1 N NaOH to pH 6.5 as necessary. The liposomes were purified by a combination of ionexchange chromatography and size-exclusion chromatography. First, DOWEX IRA 910 resin was treated with 1 N NaOH, followed by 3 washes with deionized water and then followed by 3 washes of 3 N HCl, and then multiple washes with water. The liposomes were passed through the prepared resin, and the conductivity of the eluted fractions

was measured by using a flow-cell conductivity meter (Pharmacia, Upsalla, Sweden). The fractions were deemed acceptable for further purification if the conductivity was less than 15 $\mu\text{S}/\text{cm}$. The liposome eluate was then applied to a Sephadex G-75 (Pharmacia) column equilibrated with deionized water, and the collected liposome fraction was measured for conductivity (typically less than 1 $\mu\text{S}/\text{cm}$). Cross-membrane isotonicity was achieved by addition of 40% dextrose solution to a final concentration of 5% (w/w) and the buffer (Hepes) added from a stock solution (0.5 M, pH 6.5) to a final concentration of 10 mM.

[0084] A stock solution of irinotecan was prepared by dissolving irinotecan·HCl trihydrate powder in deionized water to 15 mg/mL of anhydrous irinotecan-HCl, taking into account water content and levels of impurities obtained from the certificate of analysis of each batch. Drug loading was initiated by adding irinotecan at 500g/mol liposome phospholipid and heating to 60 ± 0.1 °C for 30 min in a hot water bath. The solutions were rapidly cooled upon removal from the water bath by immersing in ice cold water. Extraliposomal drug was removed by size exclusion chromatography, using Sephadex G75 columns equilibrated and eluted with Hepes buffered saline (10 mM Hepes, 145 mM NaCl, pH 6.5). The samples were analyzed for irinotecan by HPLC and phosphate by the method of Bartlett (see Phosphate Determination).

[0085] One preferred example of a storage stable irinotecan liposome described herein is the product that will be marketed as ONIVYDE (irinotecan liposome injection). ONIVYDE is a topoisomerase inhibitor, formulated with irinotecan hydrochloride trihydrate into a liposomal dispersion, for intravenous use. ONIVYDE indicated for the treatment of metastatic adenocarcinoma of the pancreas after disease progression following gemcitabine-based therapy.

[0086] ONIVYDE is a storage stabilized liposome having a pH of about 7.25. The ONIVYDE product contains irinotecan sucrosolate encapsulated in a liposome, obtained from an irinotecan hydrochloride trihydrate starting material. The chemical name of irinotecan is (S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo 1H-pyrano[3',4':6,7]-indolizino[1,2-b]quinolin-9-yl-[1,4'bipiperidine]-1'-carboxylate. The dosage of ONIVYDE can be calculated based on the equivalent amount of irinotecan trihydrate hydrochloride starting material used to prepare the irinotecan liposomes, or based on the amount of irinotecan in the liposome. There are about 866 mg of irinotecan per gram of irinotecan trihydrate hydrochloride. For example, an ONIVYDE dose of 80 mg based on the amount of irinotecan hydrochloride trihydrate starting material actually contains about $0.866 \times (80\text{mg})$ of irinotecan in the final product (i.e., a dose of $80 \text{ mg}/\text{m}^2$ of ONIVYDE based on the weight of irinotecan hydrochloride starting material is equivalent to about $70 \text{ mg}/\text{m}^2$ of irinotecan in the final product). ONIVYDE is a sterile, white to slightly yellow opaque isotonic liposomal dispersion. Each 10 mL single-dose vial contains 43 mg irinotecan free base at a concentration of 4.3 mg/mL. The liposome is a unilamellar lipid bilayer vesicle, approximately 110 nm in diameter, which encapsulates an aqueous space containing irinotecan in a gelated or precipitated state as the sucrose octasulfate salt. The vesicle is composed of 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC) 6.81 mg/mL, cholesterol 2.22 mg/mL, and methoxy-terminated polyethylene glycol (MW 2000)-distearoylphosphatidyl ethanolamine (MPEG-2000-DSPE) 0.12 mg/mL. Each mL

also contains 2-[4-(2-hydroxyethyl) piperazin-1-yl]ethanesulfonic acid (HEPES) as a buffer 4.05 mg/mL and sodium chloride as an isotonicity reagent 8.42 mg/mL. Each vial of ONIVYDE contains 43 mg/10 mL irinotecan free base as a white to slightly yellow, opaque, liposomal dispersion in a single-dose vial.

[0087] In one example of the invention, an ONIVYDE unit dosage form is a pharmaceutical composition comprising an amount of irinotecan encapsulated in a liposome that provides a total amount of about 70 mg/m² of irinotecan, providing an amount of irinotecan equivalent to 80 mg/m² irinotecan hydrochloride trihydrate, and less than about 20% lyso-PC. The unit dosage form can be an intravenous formulation having a total volume of about 500 mL. ONIVYDE is prepared for administering by diluting the isotonic liposomal dispersion from the vial as follows: withdraw the calculated volume of ONIVYDE from the vial. ONIVYDE is diluted in 500 mL 5% Dextrose Injection, USP or 0.9% Sodium Chloride Injection, USP and mix diluted solution by gentle inversion; protect diluted solution from light and administer diluted solution within 4 hours of preparation when stored at room temperature or within 24 hours of preparation when stored under refrigerated conditions [2°C to 8°C (36°F to 46°F)].

[0088] ONIVYDE (irinotecan liposome injection) is indicated, in combination with 5-fluorouracil and leucovorin, for the treatment of patients with metastatic adenocarcinoma of the pancreas that has progressed following gemcitabine-based therapy. Administer ONIVYDE prior to leucovorin and fluorouracil. The recommended dose of ONIVYDE is 70 mg/m² irinotecan administered by intravenous infusion over 90 minutes every 2 weeks. The recommended starting dose of ONIVYDE in patients known to be homozygous for the UGT1A1*28 allele is 50 mg/m² irinotecan administered by intravenous infusion over 90 minutes. The dose of ONIVYDE can be increased to 70 mg/m² as tolerated in subsequent cycles. There is no recommended dose of ONIVYDE for patients with serum bilirubin above the upper limit of normal. ONIVYDE is infused as a diluted solution intravenously over 90 minutes.

[0089] Suitable treatment regimens include ONIVYDE 70 mg/m² with (1+d racemic form) leucovorin 400 mg/m² (or 200 mg/m² of the active I form of leucovorin) and fluorouracil 2,400 mg/m² over 46 hours every 2 weeks (ONIVYDE/5-FU/LV; n=117), ONIVYDE 100 mg/m² every 3 weeks (n=147), or leucovorin 200 mg/m² and fluorouracil 2000 mg/m² over 24 hours weekly for 4 weeks followed by 2 week rest (5-FU/LV; n=134).

Example 10: In vivo antitumor efficacy and tolerability of Ls-Compound 5 prepared using TEA.SOS against lung cancer xenografts in mice

[0090] FIG. 10A and FIG. 10B are each a graph, showing the efficacy of liposomal Compound 5 in combination with MM-398 in NCI-H2170 (FIG. 10A) or DMS-114 (FIG. 10B) mice xenograft models, as discussed in Example 10.

[0091] FIG. 11A and FIG. 11B are each a graph showing the Kaplan-Meyer survival curves representing efficacy of liposomal Compound 5 in combination with MM-398 in NCI-H2170 (FIG. 11A) and DMS-114 (FIG. 11B) a mouse xenograft model, as discussed in Example 10.

[0092] FIG. 12A and FIG. 12B are each a graph showing Tolerability of liposomal Compound 5 in combination with MM-398 in NCI-H2170 (FIG. 12A) or DMS-114 (FIG. 12B) in a mouse xenograft model.

[0093] Antitumor efficacy of liposomes loaded with an ATR inhibitor Compound 5 in combinations with MM-398 (liposomal Irinotecan) and was studied in the model of human NCI-H2170 (lung squamous cell carcinoma) and DMS-114 (small cells lung carcinoma) lung cell lines.

[0094] The cells were obtained from American Type Culture Collection (Rockville, MD) and propagated in RPMI medium supplemented with 10% fetal calf serum, 50 U/mL penicillin G, and 50 µg/mL of streptomycin sulfate at 37 °C, 5% CO₂ as recommended by the supplier. NCR nu/nu homozygous athymic male nude mice (4-5 week old, weight at least 16 g) were obtained from Charles River. The mice were inoculated subcutaneously in the right flank with 0.1 mL of the suspension containing 5 × 10⁶ cells suspended in PBS supplemented with 30% Matrigel. When tumors achieved the size between 150 mm³ and 350 mm³ the animals were assigned to the treatment groups according to the following method. The animals were ranked according to the tumor size, and divided into 6 categories of decreasing tumor size. Four treatment groups of 10 animals/group were formed by randomly selecting one animal from each size category, so that in each treatment group all tumor sizes were equally represented. The animals received four tail vein injections, at the intervals of 7 days, of the following preparations: 1) Control (HEPES-buffered saline pH 6.5); 2) MM-398 at dose 5 mg/kg per injection; 3) Liposomal Compound 5 at 80 mg/kg per injection; 4) MM-398 followed by injections of liposomal Compound 5 with a 24 h interval. Liposomes for injections were prepared as described in Example 7. MM-398 is described in Example 9B.

[0095] The animal weight and tumor size were monitored twice weekly. The tumor progression was monitored by palpation and caliper measurements of the tumors along the largest (length) and smallest (width) axis twice a week. The tumor sizes were determined twice weekly from the caliper measurements using the formula:

$$\text{Tumor volume} = [(\text{length}) \times (\text{width})^2] / 2$$

[0096] To assess treatment-related toxicity, the animals were also weighted twice weekly. When the tumors in the group reached 10% of the mouse body weight, the animals in the group were euthanized. Average tumor volumes across the groups were plotted together and compared over time.

[0097] As demonstrated in FIGs. 10A and 10B and FIGs. 11A and 11B, liposomal ATR inhibitor Compound 5 significantly improved antitumor efficacy of MM-398 in both lung xenograft

models. The combinational treatment of liposomal ATR inhibitor and MM-398 did not affect the animals' body weight (FIGs 12A and 12B).

Example 11: Comparison of In vivo antitumor efficacy of liposomal inhibitors Ls-Compound 5 and LS-Compound A in combination with MM-398 against lung cancer xenografts in mice

[0098] FIGs. 13A and 13B are graphs showing the efficacy of liposomal Compound 5 in combination with MM-398 in Calu-6 (FIG. 13A) or COLO-699 (FIG. 13B) mice xenograft models. For example, the data in FIGs. 13A and 13B show that while liposomal ATR inhibitor Compound 5 significantly improved the antitumor efficacy of MM-398 in both models, Compound A formulated in liposomes was active only in COLO-699 xenograft model.

[0099] Antitumor efficacy of liposomes loaded with an ATR inhibitor Compound 5 in combinations with MM-398 (liposomal Irinotecan) was compared with liposomal formulation of Compound A in the xenograft model of human Calu-6 and COLO-699 lung cell lines.

[0100] The cells were obtained from American Type Culture Collection (Rockville, MD) and propagated in RPMI medium supplemented with 10% fetal calf serum, 50 U/mL penicillin G, and 50 µg/mL of streptomycin sulfate at 37 °C, 5% CO₂ as recommended by the supplier. NCR nu/nu homozygous athymic male nude mice (4-5 week old, weight at least 16 g) were obtained from Charles River. The mice were inoculated subcutaneously in the right flank with 0.1 mL of the suspension containing 5 x 10⁶ cells suspended in PBS supplemented with 30% Matrigel. When tumors achieved the size between 150 mm³ and 350 mm³ the animals were assigned to the treatment groups according to the following method. The animals were ranked according to the tumor size, and divided into 6 categories of decreasing tumor size. Four treatment groups of 10 animals/group were formed by randomly selecting one animal from each size category, so that in each treatment group all tumor sizes were equally represented. The animals received four tail vein injections, at the intervals of 7 days, of the following preparations: 1) Control (HEPES-buffered saline pH 6.5); 2) MM-398 at dose 10 or 20 mg/kg per injection; 3) Liposomal Compound 5 at 80 mg/kg per injection; 4) Liposomal Compound A at 80 mg/kg per injection; 5) MM-398 followed by injections of liposomal Compound 5 with a 24 h interval. 6) MM-398 followed by injections of liposomal Compound A with a 24 h interval. Liposomes for injections were prepared as described in Example 7.

[0101] The animal weight and tumor size were monitored twice weekly. The tumor progression was monitored by palpation and caliper measurements of the tumors along the largest (length) and smallest (width) axis twice a week. The tumor sizes were determined twice weekly from the caliper measurements using the formula:

$$\text{Tumor volume} = [(\text{length}) \times (\text{width})^2] / 2$$

[0102] To assess treatment-related toxicity, the animals were also weighted twice weekly. When the tumors in the group reached 10% of the mouse body weight, the animals in the group were euthanized. Average tumor volumes across the groups were plotted together and compared over time.

Example 12: Free drug Combination screening study


[0103] Fig 14A. is a graph showing the in vitro monotherapy cell kill of Compound 6 and Compound 5 in a panel of lung cancer cell lines. Compound 5 is more efficacious with $IC_{50} \sim 3$ ($=100.5$) folds lower than Compound 6. FIG. 14B is a graph showing the effect of Compound 5 vs. Compound 6 in combination with three chemotherapeutic agents (Carboplatin, Gemcitabine, Compound B). The figure compares the IC_{50} in log (μM) of the combination of the chemotherapeutic agents with 1 $\mu g/ml$ of Compound 6 or Compound 5. The addition of Compound 5 to the chemotherapeutic agents was more potent (lower IC_{50}) than Compound 6 in all tested cytotoxic agents and in all cell lines except one.

[0104] FIG. 15 is a graph showing the combination (Example 2 + Compound B) IC_{50} shift in Sum190PT cell line (TNBC).

[0105] FIG. 16A and FIG. 16B are graphs showing a comparison of IC_{50} shifts of the compound of Example 2 with Compound B vs combination of Compound A with Compound B in MDA-MB-453 cell line (TNBC).

Culture/Treatment Condition

[0106] In vitro efficacy study was done using CELLTITER-GLO Luminescent Cell Viability Assay (Promega) with Corning Cat #3707 384well White Clear bottom plates. Cells were plated (1000 cells/well) in 384 well format and allowed to incubate at 37 °C for 24 hours. Monotherapy drugs were added at the 24hr time point and then allowed to incubate at 37 °C for 24 hours. At the 48hr time point the drugs in media were removed, washed with PBS, and fresh media was added. Cells were then allowed to incubate at 37 °C for 72 hours. For combination studies, cells were exposed to Carboplatin or Compound B or Gemcitabine for 24 hours, then the chemotherapeutic agent was removed and cells were exposed to the secondary compound (ATR inhibitor) for 24 hours. Cells were cultured in fresh media for an additional 48 hours.



with the addition of ATP to a final concentration of 10 μ M.

[0113] The individual replicates were expressed in terms of the % of the DMSO positive control activity. The mean activity (% control) of each inhibitor concentration was plotted against the inhibitor concentration, and EC₅₀ values were determined using GRAPHPAD PRISM.

[0114] The data expressed as % control activity were plotted against inhibitor concentration, and fitted to a four parameter logistic using GRAPHPAD PRISM. Graphs for each compound are shown alongside the data in the accompanying Excel report. A summary of the compound potencies is shown below.

Table 4

| Compound ID | Kinase | ATP μ M | EC ₅₀ * nM | Top | Bottom | Hill | r2 |
|-------------|-----------------|-------------|-----------------------|-----|--------|--------|-------|
| Compound 6 | ATR ATRIP(h) | 10 | 470.8 | 101 | -2 | -0.768 | 0.998 |
| Compound 2 | ATR ATRIP(h) | 10 | 139.8 | 94 | 1 | -0.911 | 0.998 |
| Compound 5 | ATR ATRIP(h) | 10 | 233.9 | 95 | 1 | -0.819 | 0.996 |
| Compound 1 | ATR ATRIP(h) | 10 | 49.6 | 95 | -1 | -0.772 | 0.994 |
| Compound 4 | ATR ATRIP(h) | 10 | 353.4 | 97 | 1 | -0.860 | 0.997 |
| Compound 3 | ATR ATRIP(h) | 10 | 196.1 | 98 | -3 | -0.776 | 0.999 |
| Compound A | ATR ATRIP(h) | 10 | <1 | ND | ND | ND | ND |

Example 14: ATM activity determination

[0115] Estimated IC₅₀ values are as follows (obtained using STANDARD KINASEPROFII,ER):

Table 5

| Compound | Kinase | IC ₅₀ (nM) |
|------------|--------|-----------------------|
| Compound 6 | ATM(h) | >10,000 |
| Compound 2 | ATM(h) | 4817 |
| Compound 5 | ATM(h) | >10,000 |
| Compound 1 | ATM(h) | 3864 |
| Compound 4 | ATM(h) | 6869 |
| Compound 3 | ATM(h) | >10,000 |

| Compound | Kinase | IC ₅₀ (nM) |
|------------|--------|-----------------------|
| Compound A | ATM(h) | 42 |

Example 15: Detecting total and phospho-ATR with Western Blot Analysis

[0116] Referring to FIG. 17: Lung cancer cells DMS-114 were exposed either to Gemcitabine [16nM] or ATR inhibitor (Compound A or Compound 5 [1 μM]) alone or in combination *in vitro*. Whole cell protein lysates were generated after 1, 3, 6 and 24 hours using a 2% SDS containing cell lysis buffer and stored at -80 °C until all samples were collected and run together at the same time for the WesternBlot analysis. The following proteins and phosphoproteins of interest were detected by antibodies purchased from Cell Signaling Technology (see methods and materials part below): total and phospho-ATR (S428), phospho-CHK1 (S317 and S345), γH2AX, beta-actin.

[0117] Results in FIG. 17: Total and phospho-ATR signal is strongly reduced over time with the addition of either of the two ATR inhibitors (Compound A and Compound 5) and not in control or Gemcitabine alone treatment (compare e.g. lane 1, 2 with 3, 4 or 5, 6 at 24 hrs). In response to Gemcitabine, the phospho-CHK1 (S317 and S345) signaling is significantly increased over the period of 24 hrs. The addition of any of the used ATR inhibitors (Compound A or Compound 5) abrogates the phosphorylation signal of CHK1. The reduction of the ATR protein and downstream CHK1 signaling by both inhibitors confirms the specific on-target effects of both molecules. More importantly, our hypothesis is that the loss of the cell cycle check-point, driven by the CHK1 phosphorylation signaling that supposed to lead to cell cycle arrest and repair of DNA damage, induces cell death by accumulation of DNA damage. The increased γH2AX signal can be seen as a surrogate for the measurement of increased toxicity and accumulation of DNA damage.

Cell culture

[0118] U2OS and all other cells were grown in (RPMI) supplemented with 10% fetal bovine serum (FBS) and antibiotics. Imaging and time-lapse compatible NuLight red cell lines were generated as recommend by Essen Bioscience Inc. using lentiviral particles, infection and puromycin selection protocols.

Antibodies and Western-Blots

[0119] All target and phospho-specific antibodies were purchased from Cell Signaling and/or Epitomics and were used at 1:1000 dilutions. A list of all antibodies is available in Table 6.

Standard antibody based Western blot and immunohistochemistry protocols were used. Briefly, 2% SDS based cell lysis buffer was used to collect whole cell lysate for Western Blotting. Secondary antibodies and staining protocols were purchased and followed regarding LiCor Biosciences and/or BD Bioscience.

Table 6

| Target | Company | Catalog # |
|---------------------|---------------------------|-----------|
| ATR | Cell Signaling Technology | 2790 |
| ATR phospho (S428) | Cell Signaling Technology | 2853 |
| CHK1 phospho (S317) | Cell Signaling Technology | 12302 |
| CHK1 phospho (S345) | Cell Signaling Technology | 2348 |
| H2AX phospho (S139) | Cell Signaling Technology | 9718 |
| beta-actin | Cell Signaling Technology | 4970 |

Whole cell lysis protocol

[0120] Cells were grown and treated in a 6 cm dish scale. To harvest cells, medium was removed and quickly replaced by ice cold PBS. PBS was then replaced with 250 μ L of 2% SDS lysis buffer. After five minutes of incubation, the lysed cells were scraped off and loaded onto a cell lysate homogenizer microcentrifuge spin-column (QIAshredder, Qiagen, Cat.-Nr. 79656). The filtrate was then loaded onto a 0.2 μ m centrifugal filter column (Nanosep MF 0.2_m, Pall, Cat.-Nr. ODM02C34). The lysates were then stored at -80 °C.

[0121] 2% SDS containing lysis buffer (Table 7) was modified according to Steven et al., "Protein microarrays for multiplex analysis of signal transduction pathways," Nat Med 10, no. 12 (December 2004): 1390-1396.

Table 7

| Titrated pH=6.8 | Final Conc. |
|-----------------|-------------|
| Trizma-Base | 50mM |
| SDS | 2% |
| Glycerol | 5% |
| EDTA | 5mM |
| NaF | 1mM |

Example 16: Broad Kinase Panel Screening (359)

[0122] Major hits out of 359 were: ALK, ARK, c-MER, CLK1, DYRK, GSK3a, GSK3b, FLT2,

FLT3, MLK1, SIK2, TNIK, and YSK4.

Table 8

| Kinase: | Compound 5 (M) | Log IC ₅₀ | Kinase: | nM |
|--------------|----------------|----------------------|--------------|------|
| ALK1/ACVRL1 | 3.57E-08 | -7.45 | ALK1/ACVRL1 | 35.7 |
| ALK2/ACVR1 | 1.66E-08 | -7.78 | ALK2/ACVR1 | 16.6 |
| ARK5/NUAK1 | 1.68E-08 | -7.78 | ARK5/NUAK1 | 16.8 |
| CLK1 | 1.34E-07 | -6.87 | CLK1 | 134 |
| CLK4 | 8.86E-08 | -7.05 | CLK4 | 88.6 |
| DDR1 | 1.91E-07 | -6.72 | DDR1 | 191 |
| DYRK2 | 7.18E-08 | -7.14 | DYRK2 | 71.8 |
| FLT4/VEGFR3 | 6.34E-08 | -7.20 | FLT4/VEGFR3 | 63.4 |
| GSK3a | 5.73E-09 | -8.24 | GSK3a | 5.73 |
| GSK3b | 6.30E-08 | -7.20 | GSK3b | 63 |
| MLK1/MAP3K9 | 1.13E-08 | -7.95 | MLK1/MAP3K9 | 11.3 |
| MLK2/MAP3K10 | 9.58E-09 | -8.02 | MLK2/MAP3K10 | 9.58 |
| MLK3/MAP3K11 | 5.90E-09 | -8.23 | MLK3/MAP3K11 | 5.9 |
| PIM1 | 7.26E-08 | -7.14 | PIM1 | 72.6 |
| PKCmu/PRKD1 | 1.34E-07 | -6.87 | PKCmu/PRKD1 | 134 |
| PKD2/PRKD2 | 1.57E-07 | -6.80 | PKD2/PRKD2 | 157 |
| RET | 7.74E-08 | -7.11 | RET | 77.4 |
| SIK2 | 1.49E-07 | -6.83 | SIK2 | 149 |
| TNIK | 1.10E-07 | -6.96 | TNIK | 110 |
| YSK4/MAP3K19 | 2.92E-08 | -7.53 | YSK4/MAP3K19 | 29.2 |

Example 17: Data using Compound A as a comparator.

[0123] A cell death and growth assay was performed combining Compound A and Gemcitabine at various concentrations in U2OS cells. The results were compared back to a prediction of cell growth and death with the combination of the two compounds (Fig. 18A). The number of living and apoptotic cells were also determined at set concentrations of Compound A (1 μ M) and Gemcitabine (0.04 μ M) in USO2 cells (Fig. 18B). The results show that the combination of Compound A and Gemcitabine are better at promoting cell death than either compound alone.

[0124] Compound A was also tested alone and in combination with SN38 at set concentrations (1 μ M and 0.2 μ M, respectively) in several lung cancer cell lines (NCI-H520 and NCI-H596)

and U2OS cells, with the number of cells monitored over time. The results indicate that the combination is more effective at maintaining or reducing cell numbers over time than either Compound A or SN38 in isolation (Fig. 19).

[0125] *In vitro* growth assays were also performed with Compound A and SN38 in combination and alone in the cervical cancer cell line MS-751. The *in vitro* assay shows a reduction in cell number over time with the combination as compared to either compound alone (Fig. 20).

Example 18: Comparison of Compound A and Compound 5 in combination with Gemcitabine or SN38.

[0126] *In vitro* cell death and growth assays were performed combining Compound A or Compound 5 with Gemcitabine at various concentrations in U2OS, H358, and A549 cell lines. The results indicate the concentrations of each compound and Gemcitabine that are required to trigger cell death (Fig. 21A). Set concentrations of Compound 5 and Gemcitabine or Compound A and Gemcitabine were tested in U2OS and H358 cells as well. In all cases, relative cell proliferation is reduced with the combinations compared to the drugs alone (Fig. 21B-C).

[0127] Compound A or Compound 5 was used in combination with SN38 in A549 cells. Cell growth was measured over time and at a range of concentrations. The growth curve at the optimal Compound/SN38 concentration is highlighted (Fig. 22A). Relative proliferation of A549 cells was measured with either Compound A or Compound 5 alone at a range of concentrations (Fig. 22B).

[0128] The IC₅₀ values were determined in several cell lines using a set concentration of Gemcitabine with varying concentrations of Compound A or Compound 5 (Fig. 23).

[0129] A summary of the lung cancer cell lines that are responsive to Compound A or Compound 5 in combination with Gemcitabine or SN38 is provided in Fig. 24.

Example 19: On-target and Off-target comparisons of Compound A and Compound 5.

[0130] Various ATR inhibitors were tested for their ability to inhibit ATR (on-target) and ATM (off-target). Inhibition is reported as IC₅₀ in nM (Fig. 25A). Additional "off-target" kinases were tested with Compound A or Compound 5 as well (Fig. 25B).

[0131] On-target testing was performed with Compound A or Compound 5 in A549 lung cancer cells (Fig. 26A-B), H23 lung cancer cells (Fig. 26C-D), and DMS-114 cells (Fig. 26E-F). CHK1 S345 phosphorylation, a readout of ATR inhibition, was measured by western blot. Each compound was used with a fixed concentration of Gemcitabine as well.

[0132] Further on-target studies were performed on the HCC-70 TNBC, MDA-MB-468 TNBC, and DMS-114 cell line. A set concentration of SN38 was used with a range of Compound A or Compound 5 concentrations. Various on-target activity parameters were tested and measured by western blot (Fig. 27A-B)

[0133] Cell cycle stage profiles of SUM149 cells were determined 24 hours after addition of SN38 and Compound A or Compound 5 (Fig. 28). A larger percentage of cells are arrested in G2 phase when receiving the combination of drugs as compared to controls.

Example 20: Comparison of liposomal formulations of Compound A and Compound 5 combined with MM398.

[0134] DMS-114 lung xenograft models were used to measure the effects of Ls-Compound A or Ls-Compound 5 in combination with MM398. A set dosage of MM398 was used (5mpk) with two different dosages of Compound A or Compound 5 (20mpk or 80mpk). The effects of treatment were assayed by measuring the levels of CHK1 S345 phosphorylation (Fig. 29).

[0135] The effects of the Ls-Compound A with MM398 was also tested in the SUM-149 cell line. The effects of the treatment were assayed by measuring phosphorylated levels of RPA2, DNAPK, CHK1, and γ H2AX. Ls-Compound 5 was also tested without the combination of MM398 (Fig.30A-C).

Example 21: In vivo antitumor efficacy and tolerability of Ls- Compound 5 prepared using TEA.SOS against triple negative breast cancer xenografts in mice

[0136] Antitumor efficacy of liposomes loaded with an ATR inhibitor Compound 5 in combinations with MM-398 (liposomal Irinotecan) and was studied in the model of human SUM-149 (triple negative breast cancer) cell line.

[0137] The cells were obtained from American Type Culture Collection (Rockville, MD) and propagated in RPMI medium supplemented with 10% fetal calf serum, 50 U/ml penicillin G, and 50 μ g/mL of streptomycin sulfate at 37°C, 5% CO₂ as recommended by the supplier. NCR nu/nu homozygous athymic male nude mice (4-5 week old, weight at least 16 g) were obtained from Charles River. The mice were inoculated subcutaneously in the right flank with 0.1 mL of the suspension containing 10⁷ cells suspended in PBS supplemented with 30% Matrigel. When tumors achieved the size between 150 mm³ and 350 mm³ the animals were assigned to the treatment groups according to the following method. The animals were ranked according to the tumor size, and divided into 6 categories of decreasing tumor size. Four treatment groups of 10 animals/group were formed by randomly selecting one animal from each size category, so that in each treatment group all tumor sizes were equally represented. The animals received five tail vein injections, at the intervals of 7 days, of the following preparations: 1) Control

(HEPES-buffered saline pH 6.5); 2) MM-398 at dose 5 mg/kg per injection; 3) Liposomal Compound 5 at 80 mg/kg per injection; 4) MM-398 followed by injections of liposomal Compound 5 with a 24 h interval; 5)) MM-398 followed by injections of free un-capsulated ATR inhibitor Compound A. Liposomes for injections were prepared as described in Example 10.

[0138] The animal weight and tumor size were monitored twice weekly. The tumor progression was monitored by palpation and caliper measurements of the tumors along the largest (length) and smallest (width) axis twice a week. The tumor sizes were determined twice weekly from the caliper measurements using the formula:

$$\text{Tumor volume} = [(\text{length}) \times (\text{width})^2] / 2$$

[0139] To assess treatment-related toxicity, the animals were also weighted twice weekly. When the tumors in the group reached 10% of the mouse body weight, the animals in the group were euthanized. Average tumor volumes across the groups were plotted together and compared over time.

[0140] As demonstrated in Figure 31, liposomal ATR inhibitor Compound 5 significantly improved antitumor efficacy of MM-398 in the xenograft model. The combinational treatment of liposomal ATR inhibitor and MM-398 did not affect the animals' body weight (Figure 32).

Example 22: Cell line profiling and comparison with Incucyte data

[0141] A panel of fluorescently labeled cell lines was profiled for various proteins involved in the DNA damage pathway. Parental cancer cell lines were transduced with a NucLight Red lentivirus and selected with puromycin. Basal protein levels were measured and quantified by Western blot analysis. When quantification was performed, the PD marker signal was normalized to the beta actin signal. Protein levels were correlated with the cell line's "integral score", a measure of the in vitro cellular response to Compound 5 and chemotherapy combination treatment. The integral score can be calculated as follows:

[0142] Referring to Figures 33 and 34, cell lines were seeded in 96 well plates and each well was exposed to a dose matrix of Compound 5 and chemotherapy for 4 days. Dynamic cell viability was measured using an Incucyte assay. For each drug combination in the dose matrix, a dose combination-specific integral score was calculated by summing the area under normalized cell proliferation curve. A cell line specific integral score was computed by averaging the four largest scores over the dose matrix.

[0143] Referring to Figures 35 and 36, it was observed that in lung cancer cell lines exposed to Compound 5 and SN38, basal MRE11 protein levels and basal ATM levels both correlated significantly with the cell line integral scores. In lung cancer cells that have a p53 compromised background and were exposed to Compound 5 and SN38, the integral score correlates with NBS, MRE11 and RAD50 protein levels. "Compromised p53" cells are those that have non-

zero basal p53 protein levels (in most cells, p53 levels are only visible in Western blots after treatment). (See Figure 37, Figure 38 and Figure 39).

Example 23: Signaling experiment: Compound 5 vs. Compound A in combination with SN38: 6 different PD markers

[0144] Referring to Figures 40-44: Pharmacodynamics markers pChk1, pRPA2, pATR, pDNAPK, pChk2 and γ H2AX were quantified by Western blot after cancer cell lines were exposed to various doses of Compound 5 or Compound A in combination with SN38. HCC70, DMS 114, MDAMB468, NCIH1299 and NCIH460 cells were seeded in 12 well plates, allowed to incubate overnight then exposed to SN38 and/or the ATR inhibitor. After 24 hours of drug exposure, cells were lysed for Western blotting. When quantification was performed, the PD marker signal was normalized to the beta actin signal.

Example 24: Signaling experiment: Compound 5 vs. Compound A in combination with gemcitabine: 3 different PD markers

[0145] Referring to Figures 45-50: Six fluorescently labeled cell lines were profiled for various proteins involved in the DNA damage pathway after exposure to an ATR inhibitor and/or gemcitabine. Parental cancer cell lines were transduced with a NucLight Red lentivirus and selected with puromycin. Pharmacodynamics markers pChk1, pATR, and γ H2AX were quantified by Western blot after cancer cell lines were exposed to various doses of Compound 5 or Compound A in combination with 16 nM gemcitabine. A549, NCIH23, DMS114, U20S, HCC827 and NCIH460 cells were seeded in 12 well plates, allowed to incubate overnight then exposed to SN38 and/or the ATR inhibitor. After 6 or 18 hours of drug exposure, cells were lysed for Western blotting. When quantification was performed, the PD marker signal was normalized to the beta actin signal.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [US62277262](#) [0001]
- [US62420258](#) [0001]
- [US62444172](#) [0001]
- [US8912198B](#) [0005]
- [US8147867B](#) [0021]
- [US4990610A](#) [0022]
- [WO2010071827A1](#) [0032]

Non-patent literature cited in the description

- **CORTEZ et al.** Science, 2001, vol. 294, 1713-1716 [0004]
- **CASPER et al.** Am. J. Hum. Genet, 2004, vol. 75, 654-660 [0004]
- **ZOU et al.** Science, 2003, vol. 300, 1542-1548 [0004]
- **BAO et al.** Nature, 2001, vol. 411, 969-974 [0004]
- **LASIC D.** Liposomes: From physics to applications Elsevier 19930000 [0021]
- Liposome Technology CRC Press 20060000 vol. 1-3, [0021]
- **ÜNSAL-KAÇMAZ et al.** PNAS, 2002, vol. 99, 106673-6678 [0031]
- **KUMAGAI et al.** Cell, 2006, vol. 124, 943-955 [0031]
- **UNSAL-KACMAZ et al.** Molecular and Cellular Biology, 2004, 1292-1300 [0031]
- **HALL-JACKSON et al.** Oncogene, 1999, vol. 18, 6707-6713 [0031]
- J. Med. Chem., 2011, vol. 54, 2320- [0067]
- **GERAN, R.I. et al.** Cancer Chemother. Rep., 1972, vol. 3, 1-88 [0080]
- **STEVEN et al.** Protein microarrays for multiplex analysis of signal transduction pathways Nat Med, 2004, vol. 10, 121390-1396 [0121]

Patentkrav

1. Forbindelse med formel (Ia) eller et farmaceutisk acceptabelt salt deraf:




5 (Ia),

hvor R' er en alkylaminodel, der omfatter en tertiær alkylsubstitueret amin, der har en pKa på 9,5 eller derover.


2. Forbindelse ifølge krav 1, hvor R' er NR^aR^b, hvor R^a og R^b
10 hver for sig er C₁-C₄-alkyl.

3. Forbindelse ifølge krav 1 eller krav 2, som har formlen ifølge forbindelse 5:




15 (Forbindelse 5)
eller et farmaceutisk acceptabelt salt deraf.

4. Forbindelse med formlen ifølge forbindelse 6 eller et farmaceutisk acceptabelt salt deraf:




(Forbindelse 6).

5. Liposomsammensætning, som omfatter en ATR-
5 proteinkinasehæmmer med formel (I) eller et farmaceutisk acceptabelt salt deraf:




(I),

hvor R er:




- i) og R¹ er C₁-C₄-alkylamino;
10 ii) -N(H)(C₁-C₄-alkyl)-NR^aR^b, hvor R^a og R^b hver for sig er C₁-C₄-alkyl;
iii) -(G)-NR^aR^b; hvor R^a og R^b hver for sig er C₁-C₄-alkyl; og G er C₁-C₄-alkyl, hvor G eventuelt kan være substitueret med C₁-
15 C₄-alkyl; eller



- iv) hvor R^c og R^d hver for sig er C₁-C₄-alkyl.

6. Liposomsammensætning ifølge krav 4, hvor forbindelsen er

forbindelse 5:




(Forbindelse 5)

eller et farmaceutisk acceptabelt salt deraf.

5

7. Liposomsammensætning ifølge krav 4, hvor forbindelsen er forbindelse 6:



(Forbindelse 6)

10 eller et farmaceutisk acceptabelt salt deraf.

8. Liposomsammensætning ifølge et hvilket som helst af kravene 4-6, hvor liposomet omfatter et phospholipid og kolesterol.

15 9. Liposomsammensætning ifølge et hvilket som helst af kravene 4-6, hvor liposomet omfatter et polyethylenglycol (PEG)-lipidderivat og kolesterol.

20 10. Liposomsammensætning ifølge krav 8, hvor liposomet yderligere omfatter methoxy-poly(ethylenglycol)-1,2-distearoyl-sn-glycerol (PEG2000-DSG).



11. Liposomsammensætning ifølge krav 10, hvor liposomet omfatter HSPC, kolesterol og PEG2000-DSG i et molært forhold på 3:2:0,15.

5 12. Liposomsammensætning ifølge krav 5, hvor R er en del, der omfatter en amin med en pK_a over 7,0.

13. Liposomsammensætning ifølge krav 5, hvor R er en del, der omfatter en amin med en pK_a over 8,0.

DRAWINGS

Drawing



BlockA1: R₁R₂N= 4-diethylaminopiperidine
 BlockA2: R₁R₂N= 4-(2-(diethyl amino)ethyl)piperidine
 BlockA3: R₁R₂N= Diethylamine




Fig. 1

Scheme 2




Fig. 2






Fig. 3

**Fig. 4****Fig. 5**




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10A




Fig. 10B




Fig. 11A




Fig. 11B




Fig. 12A




Fig. 12B




Fig. 13A




Fig. 13B




Fig. 14A




Fig. 14B






Fig. 15

**Fig. 16A****Fig. 16B**




Fig. 17





Fig. 18A






Fig. 18B

**Fig. 19**

In vitro growth of MS-751





Fig. 20



Gemcitabine [0.128 to 16nM]

Fig. 21A





Fig. 21B

Compound 5 Gem combo

Compound A Gem combo



H358 Compound 5

H358 Compound A






Fig. 21C



Fixed Gemcitabine conc. [3.2nM]

ATRI conc. Serial dilutions: 3, 1, 0.11, 0.33, 0.004uM




Fig. 23

| Gemcitabine | Compound A | Compound 5 |
|--------------------|---------------------------------|---------------------------------|
| responder | A549, HCC827, H2170, H358, H460 | A549, HCC827, H2170, H358, H460 |
| non-responder | H2170, H358, H596 | H2170, H358, H596 |

| SN38 | Compound A | Compound 5 |
|---------------|---|---|
| responder | A549, DMS-114, H1299, H23, H460, Calu-6 | A549, DMS-114, H1299, H23, H460, Calu-6 |
| non-responder | Calu-6, H1993, H23, H596 | Calu-6, H1993, H23, H596 |

Fig. 24

| Kinase | ATR IC ₅₀ (nM) | ATM IC ₅₀ (nM) |
|-----------------------|-------------------------------------|-------------------------------------|
| Compound 6 | 470.8 | > 10,000 |
| Compound 2 | 139.8 | 4817 |
| Compound 5 | 233.9 | > 10,000 |
| Compound 4 | 353.4 | 6869 |
| Compound 3 | 196.1 | > 10,000 |
| Compound A | < 1 * | 42 |
| Compound 1 *** | 49.6 ** | 3864 |

Lit. ref.

EuroFin, UK, cell freesystem

* KI < 0.2nM

** KI = 6 nM

IC₅₀ = 120nM

*** discontinued

Fig. 25A

| Kinase | Compound 5 IC ₅₀ (nM) | Compound A IC ₅₀ (nM) |
|---------------|--|--|
| GSK3 α | 206 | 126 |
| GSK3 β | 876 | 210 |
| MLK1 | 74 | 59 |
| MLK2 | 91 | 76 |

EuroFin, UK, cell freesystem

Fig. 25B

A549 Cells

Western-Blot based analysis of inhibition of CHK1 S345 phosphorylation by ATRi (dose and time-course)




Fig. 26A

IC50 of inhibition of CHK1 S345 phosphorylation




Fig. 26B

IC50 of inhibition of CHK1 S345 phosphorylation




Fig. 26D




Fig. 26F




Fig. 27A





Fig. 27B



| | G1 | S | G2 |
|-----------------------------|-----|-----|-----|
| Control | 67% | 14% | 19% |
| Cmpd 5 alone [1uM] | 66% | 14% | 20% |
| SN38 alone [25nM] | 35% | 51% | 14% |
| SN38 [25nM] Cmpd 5 {3uM} | 44% | 22% | 34% |
| SN38 [25nM] Cmpd 5 [1uM] | 34% | 31% | 35% |
| SN38 [25nM] Cmpd 5 [0.33uM] | 40% | 31% | 29% |
| SN38 [25nM] Cmpd 5 [0.11uM] | 54% | 39% | 7% |

| | G1 | S | G2 |
|-----------------------------|-----|-----|-----|
| Control | 67% | 14% | 19% |
| Cmpd A alone [1uM] | 69% | 17% | 14% |
| SN38 alone [25nM] | 35% | 51% | 14% |
| SN38 [25nM] Cmpd A {3uM} | 50% | 16% | 35% |
| SN38 [25nM] Cmpd A [1uM] | 41% | 29% | 30% |
| SN38 [25nM] Cmpd A [0.33uM] | 37% | 32% | 31% |
| SN38 [25nM] Cmpd A [0.11uM] | 37% | 40% | 23% |

Fig. 28







Fig. 29





Fig. 30B





Fig. 30C




Fig. 31




Fig. 32




Fig. 33

Integral Score




Fig. 34

MRE11 protein expression and SN38 Integral Score correlation
(ATRI/SN38 combo; Lung only)




Fig. 35

ATM protein expression and SN38 Integral Score correlation
(ATRI/SN38 combo; Lung only)




Fig. 36

NBS protein expression and SN38 Integral Score correlation
(ATR/IN38 combo; p53 compromised; Lung only)




Fig. 37

MRE11 protein expression and SN38 Integral Score correlation
(ATR/ SN38 combo; p53 compromised; Lung only)




Fig. 38

MRE11 protein expression and SN38 Integral Score correlation
(ATR/SN38 combo;p53 compromised; Lung only)




Fig. 39

H1299 SN38 Compound 5 or Compound A combo
pATR S426 dynamics




Fig. 40A

H1299 SN38 Compound 5 or Compound A combo
pDNAPK S2065 dynamics





Fig. 40B

H1299 SN38 Compound 5 or Compound A combo
pCHK1 S345 dynamics





Fig. 40C

H1299 SN38 Compound 5 or Compound A combo
pRPA2 S33 dynamics





Fig. 40D

H1299 SN38 Compound 5 or Compound A combo
pCHK2 T68 dynamics





Fig. 40E

H1299 SN38 Compound 5 or Compound A combo
yH2AX S2065 dynamics





Fig. 40F

H460 SN38 Compound 5 or Compound A combo
pDNAPK S2065 dynamics




Fig. 41B

H460 SN38 Compound 5 or Compound A combo
pCHK1 S345 dynamics





Fig. 41C

H460 SN38 Compound 5 or Compound A combo
pRPA2 S33 dynamics





Fig. 41D

H460 SN38 Compound 5 or Compound A combo
pCHK1 T68 dynamics





Fig. 41E

H460 SN38 Compound 5 or Compound A combo
yH2AX S2065 dynamics




Fig. 41F

DMS-114 SN38 Compound 5 or Compound A combo
pATR S428 dynamics




Fig. 42A

DMS-114 SN38 Compound 5 or Compound A combo
pDNAPK S2065 dynamics




Fig. 42B

DMS-114 SN38 Compound 5 or Compound A combo
pCHK1 S345 dynamics




Fig. 42C

DMS-114 SN38 Compound 5 or Compound A combo
pRPA2 S33 dynamics





Fig. 42D

DMS-114 SN38 Compound 5 or Compound A combo
pCHK2 T68 dynamics





Fig. 42E

DMS-114 SN38 Compound 5 or Compound A combo
yH2AX dynamics




Fig. 42F

HCC-70 SN38 Compound 5 or Compound A combo
pATR S428 dynamics




Fig. 43A

HCC-70 SN38 Compound 5 or Compound A combo
pDNAPK S2065 dynamics





Fig. 43B

HCC-70 SN38 Compound 5 or Compound A combo
pCHK1 S345 dynamics




Fig. 43C

HCC-70 SN38 Compound 5 or Compound A combo
pRPA2 S33 dynamics





Fig. 43D

HCC-70 SN38 Compound 5 or Compound A combo
pCHK2 T68 dynamics





Fig. 43E

HCC-70 SN38 Compound 5 or Compound A combo
yH2AX dynamics





Fig. 43F

MDA-MB468 SN38 Compound 5 or Compound A combo
pATR S428 dynamics




Fig. 44A

MDA-MB468 SN38 Compound 5 or Compound A combo
pDNAPK S2065 dynamics




Fig. 44B




Fig. 44C




Fig. 44D

MDA-MB468 SN38 Compound 5 or Compound A combo
pCHK2 T68 dynamics





Fig. 44E

MDA-MB468 SN38 Compound 5 or Compound A combo
yH2AX dynamics




Fig. 44F




Fig. 45





Fig. 47

**Fig. 51**





Fig. 52

inhibition of RPA2



fold changes in phospho signaling

Fig. 53




Fig. 54