



(22) Date de dépôt/Filing Date: 2000/08/24
(41) Mise à la disp. pub./Open to Public Insp.: 2001/03/08
(45) Date de délivrance/Issue Date: 2011/08/09
(62) Demande originale/Original Application: 2 380 355
(30) Priorité/Priority: 1999/09/01 (US PCT/US99/20111)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
C07K 14/705 (2006.01), *C07K 16/28* (2006.01),
C07K 19/00 (2006.01), *C12P 21/02* (2006.01)

(72) Inventeurs/Inventors:
EATON, DAN L., US;
FILVAROFF, ELLEN, US;
GERRITSEN, MARY E., US;
GODDARD, AUDREY, US;
GODOWSKI, PAUL J., US;
GRIMALDI, CHRISTOPHER J., US;
...

(73) Propriétaire/Owner:
GENENTECH, INC., US

(74) Agent: DENNISON ASSOCIATES

(54) Titre : POLYPEPTIDES SECRETES ET TRANSMEMBRANAIRES ET ACIDES NUCLEIQUES CODANT POUR CEUX-CI

(54) Title: SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME

GGGGCTTCGGCGCCAGCGGCCAGCGCTAGTCGGTCTGGTAAGGATTTACAAAAGGTGCAGGTA
TGAGCAGGTCTGAAGACTAACATTTTGTGAAGTTGTAAAACAGAAAACCTGTTAGAAATGTTGG
TGGTTTCAGCAAGGCCTCAGTTTCCTTCCTTCAGCCCTTGTAATTTGGACATCTGCTGCTTTC
ATATTTTCATACATTACTGCAGTAACACTCCACCATATAGACCCGGCTTTACCTTATATCAGT
GACACTGGTACAGTAGCTCCAGAAAAATGCTTATTTGGGGCAATGCTAAATATTGCGGCAGTT
TTATGCATTGCTACCATTTATGTTTCGTTATAAGCAAGTTCATGCTCTGAGTCCTGAAGAGAAC
GTTATCATCAAATTAAACAAGGCTGGCCTTGTAAGTACTGAGTTGTTTAGGACTTTCT
ATTGTGGCAAACCTCCAGAAAACAACCCTTTTGTGCTGCACATGTAAGTGGAGCTGTGCTTACC
TTTGGTATGGGCTCATTATATATGTTTGTTCAGACCATCCTTTCCTACCAAATGCAGCCCAA
ATCATGGCAAACAAGTCTTCTGGATCAGACTGTTGTTGGTTATCTGGTGTGGAGTAAGTGCA
CTTAGCATGCTGACTTGCTCATCAGTTTTCACAGTGGCAATTTTGGGACTGATTTAGAACAG
AAACTCCATTGGAACCCCGAGGACAAAGGTTATGTGCTTCACATGATCACTACTGCAGCAGAA
TGGTCTATGTCATTTTCCTTCTTTGGTTTTTTCCTGACTTACATTCGTGATTTTCAGAAAATT
TCTTTACGGGTGGAAGCCAATTTACATGGATTAACCCTCTATGACACTGCACCTTGCCCTATT
AACAAATGAACGAACACGGCTACTTTCAGAGATATTGATGAAAGGATAAAATATTTCTGTAA
TGATTATGATTCTCAGGGATTGGGGAAAGGTTACAGAAAGTTGCTTATTCTTCTCTGAAATTT
TCAACCACTTAATCAAGGCTGACAGTAACACTGATGAATGCTGATAATCAGGAAACATGAAAG
AAGCCATTTGATAGATTATTCTAAAGGATATCATCAAGAAGACTATTAAAAACACCTATGCCT
ATACTTTTTTATCTCAGAAAATAAAGTCAAAAGACTATG

(57) Abrégé/Abstract:

The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.



(72) Inventeurs(suite)/Inventors(continued): GURNEY, AUSTIN L., US; WATANABE, COLIN K., US; WOOD, WILLIAM I., US

(57) Abrégé/Abstract:

The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME

FIELD OF THE INVENTION

5 The present invention relates generally to the identification and isolation of novel DNA and to the recombinant production of novel polypeptides.

BACKGROUND OF THE INVENTION

10 Extracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of
15 action in the extracellular environment.

20 Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., Proc. Natl. Acad. Sci. 93:7108-7113 (1996); U.S. Patent No. 5,536,637)].

25 Membrane-bound proteins and receptors can play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins.
30 Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and

nerve growth factor receptor.

Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

SUMMARY OF THE INVENTION

In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94%

nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length,

alternatively at least about 150 nucleotides in length, alternatively at least about 160 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700 nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83%

amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By

way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules
5 comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In another embodiment, the invention provides an antibody which binds, preferably specifically, to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

10 In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences.

In yet other embodiments, the present invention is directed to methods of using the PRO polypeptides of the present invention for a variety of uses based upon the functional biological assay data presented in the
15 Examples below.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO180 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA26843-1389".

20 Figure 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in Figure 1.

Figure 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO218 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA30867-1335".

25 Figure 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in Figure 3.

Figure 5 shows a nucleotide sequence (SEQ ID NO:5) of a native sequence PRO263 cDNA, wherein SEQ ID NO:5 is a clone designated herein as "DNA34431-1177".

Figure 6 shows the amino acid sequence (SEQ ID NO:6) derived from the coding sequence of SEQ ID NO:5 shown in Figure 5.

30 Figure 7 shows a nucleotide sequence (SEQ ID NO:7) of a native sequence PRO295 cDNA, wherein SEQ ID NO:7 is a clone designated herein as "DNA38268-1188".

Figure 8 shows the amino acid sequence (SEQ ID NO:8) derived from the coding sequence of SEQ ID NO:7 shown in Figure 7.

35 Figure 9 shows a nucleotide sequence (SEQ ID NO:9) of a native sequence PRO874 cDNA, wherein SEQ ID NO:9 is a clone designated herein as "DNA40621-1440".

Figure 10 shows the amino acid sequence (SEQ ID NO:10) derived from the coding sequence of SEQ ID NO:9 shown in Figure 9.

Figure 11 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO300 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA40625-1189".

Figure 12 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in Figure 11.

5 Figure 13 shows a nucleotide sequence (SEQ ID NO:13) of a native sequence PRO1864 cDNA, wherein SEQ ID NO:13 is a clone designated herein as "DNA45409-2511".

Figure 14 shows the amino acid sequence (SEQ ID NO:14) derived from the coding sequence of SEQ ID NO:13 shown in Figure 13.

Figure 15 shows a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO1282 cDNA, wherein SEQ ID NO:15 is a clone designated herein as "DNA45495-1550".

10 Figure 16 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in Figure 15.

Figure 17 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO1063 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA49820-1427".

15 Figure 18 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in Figure 17.

Figure 19 shows a nucleotide sequence (SEQ ID NO:19) of a native sequence PRO1773 cDNA, wherein SEQ ID NO:19 is a clone designated herein as "DNA56406-1704".

Figure 20 shows the amino acid sequence (SEQ ID NO:20) derived from the coding sequence of SEQ ID NO:19 shown in Figure 19.

20 Figure 21 shows a nucleotide sequence (SEQ ID NO:21) of a native sequence PRO1013 cDNA, wherein SEQ ID NO:21 is a clone designated herein as "DNA56410-1414".

Figure 22 shows the amino acid sequence (SEQ ID NO:22) derived from the coding sequence of SEQ ID NO:21 shown in Figure 21.

25 Figure 23 shows a nucleotide sequence (SEQ ID NO:23) of a native sequence PRO937 cDNA, wherein SEQ ID NO:23 is a clone designated herein as "DNA56436-1448".

Figure 24 shows the amino acid sequence (SEQ ID NO:24) derived from the coding sequence of SEQ ID NO:23 shown in Figure 23.

Figure 25 shows a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO842 cDNA, wherein SEQ ID NO:25 is a clone designated herein as "DNA56855-1447".

30 Figure 26 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in Figure 25.

Figure 27 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO1180 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA56860-1510".

35 Figure 28 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in Figure 27.

Figure 29 shows a nucleotide sequence (SEQ ID NO:29) of a native sequence PRO831 cDNA, wherein SEQ ID NO:29 is a clone designated herein as "DNA56862-1343".

Figure 30 shows the amino acid sequence (SEQ ID NO:30) derived from the coding sequence of SEQ ID NO:29 shown in Figure 29.

Figure 31 shows a nucleotide sequence (SEQ ID NO:31) of a native sequence PRO1115 cDNA, wherein SEQ ID NO:31 is a clone designated herein as "DNA56868-1478".

5 Figure 32 shows the amino acid sequence (SEQ ID NO:32) derived from the coding sequence of SEQ ID NO:31 shown in Figure 31.

Figure 33 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO1277 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA56869-1545".

Figure 34 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in Figure 33.

10 Figure 35 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO1074 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA57704-1452".

Figure 36 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in Figure 35.

15 Figure 37 shows a nucleotide sequence (SEQ ID NO:37) of a native sequence PRO1344 cDNA, wherein SEQ ID NO:37 is a clone designated herein as "DNA58723-1588".

Figure 38 shows the amino acid sequence (SEQ ID NO:38) derived from the coding sequence of SEQ ID NO:37 shown in Figure 37.

Figure 39 shows a nucleotide sequence (SEQ ID NO:39) of a native sequence PRO1136 cDNA, wherein SEQ ID NO:39 is a clone designated herein as "DNA57827-1493".

20 Figure 40 shows the amino acid sequence (SEQ ID NO:40) derived from the coding sequence of SEQ ID NO:39 shown in Figure 39.

Figure 41 shows a nucleotide sequence (SEQ ID NO:41) of a native sequence PRO1109 cDNA, wherein SEQ ID NO:41 is a clone designated herein as "DNA58737-1473".

25 Figure 42 shows the amino acid sequence (SEQ ID NO:42) derived from the coding sequence of SEQ ID NO:41 shown in Figure 41.

Figure 43 shows a nucleotide sequence (SEQ ID NO:43) of a native sequence PRO1003 cDNA, wherein SEQ ID NO:43 is a clone designated herein as "DNA58846-1409".

Figure 44 shows the amino acid sequence (SEQ ID NO:44) derived from the coding sequence of SEQ ID NO:43 shown in Figure 43.

30 Figure 45 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO1138 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA58850-1495".

Figure 46 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in Figure 45.

35 Figure 47 shows a nucleotide sequence (SEQ ID NO:47) of a native sequence PRO994 cDNA, wherein SEQ ID NO:47 is a clone designated herein as "DNA58855-1422".

Figure 48 shows the amino acid sequence (SEQ ID NO:48) derived from the coding sequence of SEQ ID NO:47 shown in Figure 47.

Figure 49 shows a nucleotide sequence (SEQ ID NO:49) of a native sequence PRO1069 cDNA, wherein SEQ ID NO:49 is a clone designated herein as "DNA59211-1450".

Figure 50 shows the amino acid sequence (SEQ ID NO:50) derived from the coding sequence of SEQ ID NO:49 shown in Figure 49.

5 Figure 51 shows a nucleotide sequence (SEQ ID NO:51) of a native sequence PRO1411 cDNA, wherein SEQ ID NO:51 is a clone designated herein as "DNA59212-1627".

Figure 52 shows the amino acid sequence (SEQ ID NO:52) derived from the coding sequence of SEQ ID NO:51 shown in Figure 51.

Figure 53 shows a nucleotide sequence (SEQ ID NO:53) of a native sequence PRO1129 cDNA, wherein SEQ ID NO:53 is a clone designated herein as "DNA59213-1487".

10 Figure 54 shows the amino acid sequence (SEQ ID NO:54) derived from the coding sequence of SEQ ID NO:53 shown in Figure 53.

Figure 55 shows a nucleotide sequence (SEQ ID NO:55) of a native sequence PRO1027 cDNA, wherein SEQ ID NO:55 is a clone designated herein as "DNA59605-1418".

15 Figure 56 shows the amino acid sequence (SEQ ID NO:56) derived from the coding sequence of SEQ ID NO:55 shown in Figure 55.

Figure 57 shows a nucleotide sequence (SEQ ID NO:57) of a native sequence PRO1106 cDNA, wherein SEQ ID NO:57 is a clone designated herein as "DNA59609-1470".

Figure 58 shows the amino acid sequence (SEQ ID NO:58) derived from the coding sequence of SEQ ID NO:57 shown in Figure 57.

20 Figure 59 shows a nucleotide sequence (SEQ ID NO:59) of a native sequence PRO1291 cDNA, wherein SEQ ID NO:59 is a clone designated herein as "DNA59610-1556".

Figure 60 shows the amino acid sequence (SEQ ID NO:60) derived from the coding sequence of SEQ ID NO:59 shown in Figure 59.

25 Figure 61 shows a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO3573 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA59837-2545".

Figure 62 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in Figure 61.

Figure 63 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO3566 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA59844-2542".

30 Figure 64 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in Figure 63.

Figure 65 shows a nucleotide sequence (SEQ ID NO:65) of a native sequence PRO1098 cDNA, wherein SEQ ID NO:65 is a clone designated herein as "DNA59854-1459".

35 Figure 66 shows the amino acid sequence (SEQ ID NO:66) derived from the coding sequence of SEQ ID NO:65 shown in Figure 65.

Figure 67 shows a nucleotide sequence (SEQ ID NO:67) of a native sequence PRO1158 cDNA, wherein SEQ ID NO:67 is a clone designated herein as "DNA60625-1507".

Figure 68 shows the amino acid sequence (SEQ ID NO:68) derived from the coding sequence of SEQ ID NO:67 shown in Figure 67.

Figure 69 shows a nucleotide sequence (SEQ ID NO:69) of a native sequence PRO1124 cDNA, wherein SEQ ID NO:69 is a clone designated herein as "DNA60629-1481".

5 Figure 70 shows the amino acid sequence (SEQ ID NO:70) derived from the coding sequence of SEQ ID NO:69 shown in Figure 69.

Figure 71 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO1287 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA61755-1554".

Figure 72 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in Figure 71.

10 Figure 73 shows a nucleotide sequence (SEQ ID NO:73) of a native sequence PRO1335 cDNA, wherein SEQ ID NO:73 is a clone designated herein as "DNA62812-1594".

Figure 74 shows the amino acid sequence (SEQ ID NO:74) derived from the coding sequence of SEQ ID NO:73 shown in Figure 73.

15 Figure 75 shows a nucleotide sequence (SEQ ID NO:75) of a native sequence PRO1315 cDNA, wherein SEQ ID NO:75 is a clone designated herein as "DNA62815-1576".

Figure 76 shows the amino acid sequence (SEQ ID NO:76) derived from the coding sequence of SEQ ID NO:75 shown in Figure 75.

Figure 77 shows a nucleotide sequence (SEQ ID NO:77) of a native sequence PRO1357 cDNA, wherein SEQ ID NO:77 is a clone designated herein as "DNA64881-1602".

20 Figure 78 shows the amino acid sequence (SEQ ID NO:78) derived from the coding sequence of SEQ ID NO:77 shown in Figure 77.

Figure 79 shows a nucleotide sequence (SEQ ID NO:79) of a native sequence PRO1356 cDNA, wherein SEQ ID NO:79 is a clone designated herein as "DNA64886-1601".

25 Figure 80 shows the amino acid sequence (SEQ ID NO:80) derived from the coding sequence of SEQ ID NO:79 shown in Figure 79.

Figure 81 shows a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO1557 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA64902-1667".

Figure 82 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in Figure 81.

30 Figure 83 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO1347 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA64950-1590".

Figure 84 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID NO:83 shown in Figure 83.

35 Figure 85 shows a nucleotide sequence (SEQ ID NO:85) of a native sequence PRO1302 cDNA, wherein SEQ ID NO:85 is a clone designated herein as "DNA65403-1565".

Figure 86 shows the amino acid sequence (SEQ ID NO:86) derived from the coding sequence of SEQ ID NO:85 shown in Figure 85.

Figure 87 shows a nucleotide sequence (SEQ ID NO:87) of a native sequence PRO1270 cDNA, wherein SEQ ID NO:87 is a clone designated herein as "DNA66308-1537".

Figure 88 shows the amino acid sequence (SEQ ID NO:88) derived from the coding sequence of SEQ ID NO:87 shown in Figure 87.

5 Figure 89 shows a nucleotide sequence (SEQ ID NO:89) of a native sequence PRO1268 cDNA, wherein SEQ ID NO:89 is a clone designated herein as "DNA66519-1535".

Figure 90 shows the amino acid sequence (SEQ ID NO:90) derived from the coding sequence of SEQ ID NO:89 shown in Figure 89.

Figure 91 shows a nucleotide sequence (SEQ ID NO:91) of a native sequence PRO1327 cDNA, wherein SEQ ID NO:91 is a clone designated herein as "DNA66521-1583".

10 Figure 92 shows the amino acid sequence (SEQ ID NO:92) derived from the coding sequence of SEQ ID NO:91 shown in Figure 91.

Figure 93 shows a nucleotide sequence (SEQ ID NO:93) of a native sequence PRO1328 cDNA, wherein SEQ ID NO:93 is a clone designated herein as "DNA66658-1584".

15 Figure 94 shows the amino acid sequence (SEQ ID NO:94) derived from the coding sequence of SEQ ID NO:93 shown in Figure 93.

Figure 95 shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO1329 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA66660-1585".

Figure 96 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in Figure 95.

20 Figure 97 shows a nucleotide sequence (SEQ ID NO:97) of a native sequence PRO1340 cDNA, wherein SEQ ID NO:97 is a clone designated herein as "DNA66663-1598".

Figure 98 shows the amino acid sequence (SEQ ID NO:98) derived from the coding sequence of SEQ ID NO:97 shown in Figure 97.

25 Figure 99 shows a nucleotide sequence (SEQ ID NO:99) of a native sequence PRO1342 cDNA, wherein SEQ ID NO:99 is a clone designated herein as "DNA66674-1599".

Figure 100 shows the amino acid sequence (SEQ ID NO:100) derived from the coding sequence of SEQ ID NO:99 shown in Figure 99.

Figure 101 shows a nucleotide sequence (SEQ ID NO:101) of a native sequence PRO3579 cDNA, wherein SEQ ID NO:101 is a clone designated herein as "DNA68862-2546".

30 Figure 102 shows the amino acid sequence (SEQ ID NO:102) derived from the coding sequence of SEQ ID NO:101 shown in Figure 101.

Figure 103 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO1472 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA68866-1644".

35 Figure 104 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in Figure 103.

Figure 105 shows a nucleotide sequence (SEQ ID NO:105) of a native sequence PRO1461 cDNA, wherein SEQ ID NO:105 is a clone designated herein as "DNA68871-1638".

Figure 106 shows the amino acid sequence (SEQ ID NO:106) derived from the coding sequence of SEQ ID NO:105 shown in Figure 105.

Figure 107 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO1568 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA68880-1676".

5 Figure 108 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in Figure 107.

Figure 109 shows a nucleotide sequence (SEQ ID NO:109) of a native sequence PRO1753 cDNA, wherein SEQ ID NO:109 is a clone designated herein as "DNA68883-1691".

Figure 110 shows the amino acid sequence (SEQ ID NO:110) derived from the coding sequence of SEQ ID NO:109 shown in Figure 109.

10 Figure 111 shows a nucleotide sequence (SEQ ID NO:111) of a native sequence PRO1570 cDNA, wherein SEQ ID NO:111 is a clone designated herein as "DNA68885-1678".

Figure 112 shows the amino acid sequence (SEQ ID NO:112) derived from the coding sequence of SEQ ID NO:111 shown in Figure 111.

15 Figure 113 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO1446 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA71277-1636".

Figure 114 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in Figure 113.

Figure 115 shows a nucleotide sequence (SEQ ID NO:115) of a native sequence PRO1565 cDNA, wherein SEQ ID NO:115 is a clone designated herein as "DNA73727-1673".

20 Figure 116 shows the amino acid sequence (SEQ ID NO:116) derived from the coding sequence of SEQ ID NO:115 shown in Figure 115.

Figure 117 shows a nucleotide sequence (SEQ ID NO:117) of a native sequence PRO1572 cDNA, wherein SEQ ID NO:117 is a clone designated herein as "DNA73734-1680".

25 Figure 118 shows the amino acid sequence (SEQ ID NO:118) derived from the coding sequence of SEQ ID NO:117 shown in Figure 117.

Figure 119 shows a nucleotide sequence (SEQ ID NO:119) of a native sequence PRO1573 cDNA, wherein SEQ ID NO:119 is a clone designated herein as "DNA73735-1681".

Figure 120 shows the amino acid sequence (SEQ ID NO:120) derived from the coding sequence of SEQ ID NO:119 shown in Figure 119.

30 Figure 121 shows a nucleotide sequence (SEQ ID NO:121) of a native sequence PRO1550 cDNA, wherein SEQ ID NO:121 is a clone designated herein as "DNA76393-1664".

Figure 122 shows the amino acid sequence (SEQ ID NO:122) derived from the coding sequence of SEQ ID NO:121 shown in Figure 121.

35 Figure 123 shows a nucleotide sequence (SEQ ID NO:123) of a native sequence PRO1693 cDNA, wherein SEQ ID NO:123 is a clone designated herein as "DNA77301-1708".

Figure 124 shows the amino acid sequence (SEQ ID NO:124) derived from the coding sequence of SEQ ID NO:123 shown in Figure 123.

Figure 125 shows a nucleotide sequence (SEQ ID NO:125) of a native sequence PRO1566 cDNA, wherein SEQ ID NO:125 is a clone designated herein as "DNA77568-1626".

Figure 126 shows the amino acid sequence (SEQ ID NO:126) derived from the coding sequence of SEQ ID NO:125 shown in Figure 125.

5 Figure 127 shows a nucleotide sequence (SEQ ID NO:127) of a native sequence PRO1774 cDNA, wherein SEQ ID NO:127 is a clone designated herein as "DNA77626-1705".

Figure 128 shows the amino acid sequence (SEQ ID NO:128) derived from the coding sequence of SEQ ID NO:127 shown in Figure 127.

Figure 129 shows a nucleotide sequence (SEQ ID NO:129) of a native sequence PRO1928 cDNA, wherein SEQ ID NO:129 is a clone designated herein as "DNA81754-2532".

10 Figure 130 shows the amino acid sequence (SEQ ID NO:130) derived from the coding sequence of SEQ ID NO:129 shown in Figure 129.

Figure 131 shows a nucleotide sequence (SEQ ID NO:131) of a native sequence PRO1865 cDNA, wherein SEQ ID NO:131 is a clone designated herein as "DNA81757-2512".

15 Figure 132 shows the amino acid sequence (SEQ ID NO:132) derived from the coding sequence of SEQ ID NO:131 shown in Figure 131.

Figure 133 shows a nucleotide sequence (SEQ ID NO:133) of a native sequence PRO1925 cDNA, wherein SEQ ID NO:133 is a clone designated herein as "DNA82302-2529".

Figure 134 shows the amino acid sequence (SEQ ID NO:134) derived from the coding sequence of SEQ ID NO:133 shown in Figure 133.

20 Figure 135 shows a nucleotide sequence (SEQ ID NO:135) of a native sequence PRO1926 cDNA, wherein SEQ ID NO:135 is a clone designated herein as "DNA82340-2530".

Figure 136 shows the amino acid sequence (SEQ ID NO:136) derived from the coding sequence of SEQ ID NO:135 shown in Figure 135.

25 Figure 137 shows a nucleotide sequence (SEQ ID NO:137) of a native sequence PRO1801 cDNA, wherein SEQ ID NO:137 is a clone designated herein as "DNA83500-2506".

Figure 138 shows the amino acid sequence (SEQ ID NO:138) derived from the coding sequence of SEQ ID NO:137 shown in Figure 137.

Figure 139 shows a nucleotide sequence (SEQ ID NO:139) of a native sequence PRO4405 cDNA, wherein SEQ ID NO:139 is a clone designated herein as "DNA84920-2614".

30 Figure 140 shows the amino acid sequence (SEQ ID NO:140) derived from the coding sequence of SEQ ID NO:139 shown in Figure 139.

Figure 141 shows a nucleotide sequence (SEQ ID NO:141) of a native sequence PRO3435 cDNA, wherein SEQ ID NO:141 is a clone designated herein as "DNA85066-2534".

35 Figure 142 shows the amino acid sequence (SEQ ID NO:142) derived from the coding sequence of SEQ ID NO:141 shown in Figure 141.

Figure 143 shows a nucleotide sequence (SEQ ID NO:143) of a native sequence PRO3543 cDNA, wherein SEQ ID NO:143 is a clone designated herein as "DNA86571-2551".

Figure 144 shows the amino acid sequence (SEQ ID NO:144) derived from the coding sequence of SEQ ID NO:143 shown in Figure 143.

Figure 145 shows a nucleotide sequence (SEQ ID NO:145) of a native sequence PRO3443 cDNA, wherein SEQ ID NO:145 is a clone designated herein as "DNA87991-2540".

5 Figure 146 shows the amino acid sequence (SEQ ID NO:146) derived from the coding sequence of SEQ ID NO:145 shown in Figure 145.

Figure 147 shows a nucleotide sequence (SEQ ID NO:147) of a native sequence PRO3442 cDNA, wherein SEQ ID NO:147 is a clone designated herein as "DNA92238-2539".

Figure 148 shows the amino acid sequence (SEQ ID NO:148) derived from the coding sequence of SEQ ID NO:147 shown in Figure 147.

10 Figure 149 shows a nucleotide sequence (SEQ ID NO:149) of a native sequence PRO5990 cDNA, wherein SEQ ID NO:149 is a clone designated herein as "DNA96042-2682".

Figure 150 shows the amino acid sequence (SEQ ID NO:150) derived from the coding sequence of SEQ ID NO:149 shown in Figure 149.

15 Figure 151 shows a nucleotide sequence (SEQ ID NO:151) of a native sequence PRO4342 cDNA, wherein SEQ ID NO:151 is a clone designated herein as "DNA96787-2534".

Figure 152 shows the amino acid sequence (SEQ ID NO:152) derived from the coding sequence of SEQ ID NO:151 shown in Figure 151.

Figure 153 shows a nucleotide sequence (SEQ ID NO:153) of a native sequence PRO10096 cDNA, wherein SEQ ID NO:153 is a clone designated herein as "DNA125185-2806".

20 Figure 154 shows the amino acid sequence (SEQ ID NO:154) derived from the coding sequence of SEQ ID NO:153 shown in Figure 153.

Figure 155 shows a nucleotide sequence (SEQ ID NO:155) of a native sequence PRO10272 cDNA, wherein SEQ ID NO:155 is a clone designated herein as "DNA147531-2821".

25 Figure 156 shows the amino acid sequence (SEQ ID NO:156) derived from the coding sequence of SEQ ID NO:155 shown in Figure 155.

Figure 157 shows a nucleotide sequence (SEQ ID NO:157) of a native sequence PRO5801 cDNA, wherein SEQ ID NO:157 is a clone designated herein as "DNA115291-2681".

Figure 158 shows the amino acid sequence (SEQ ID NO:158) derived from the coding sequence of SEQ ID NO:157 shown in Figure 157.

30 Figure 159 shows a nucleotide sequence (SEQ ID NO:159) of a native sequence PRO20110 cDNA, wherein SEQ ID NO:159 is a clone designated herein as "DNA166819".

Figure 160 shows the amino acid sequence (SEQ ID NO:160) derived from the coding sequence of SEQ ID NO:159 shown in Figure 159.

35 Figure 161 shows a nucleotide sequence (SEQ ID NO:161) of a native sequence PRO20040 cDNA, wherein SEQ ID NO:161 is a clone designated herein as "DNA164625-2890".

Figure 162 shows the amino acid sequence (SEQ ID NO:162) derived from the coding sequence of SEQ ID NO:161 shown in Figure 161.

Figure 163 shows a nucleotide sequence (SEQ ID NO:163) of a native sequence PRO20233 cDNA, wherein SEQ ID NO:163 is a clone designated herein as "DNA165608".

Figure 164 shows the amino acid sequence (SEQ ID NO:164) derived from the coding sequence of SEQ ID NO:163 shown in Figure 163.

5 Figure 165 shows a nucleotide sequence (SEQ ID NO:165) of a native sequence PRO19670 cDNA, wherein SEQ ID NO:165 is a clone designated herein as "DNA131639-2874".

Figure 166 shows the amino acid sequence (SEQ ID NO:166) derived from the coding sequence of SEQ ID NO:165 shown in Figure 165.

Figure 167 shows a nucleotide sequence (SEQ ID NO:167) of a native sequence PRO1890 cDNA, wherein SEQ ID NO:167 is a clone designated herein as "DNA79230-2525".

10 Figure 168 shows the amino acid sequence (SEQ ID NO:168) derived from the coding sequence of SEQ ID NO:167 shown in Figure 167.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. Definitions

15 The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO
20 polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The term "PRO polypeptide" refers to each individual PRO/number polypeptide disclosed herein. All disclosures in this specification which refer to the "PRO polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the preparation of, purification of, derivation of, formation of antibodies to or against,
25 administration of, compositions containing, treatment of a disease with, etc., pertain to each polypeptide of the invention individually. The term "PRO polypeptide" also includes variants of the PRO/number polypeptides disclosed herein.

A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be
30 isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence
35 polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino

acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1 % of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5 % of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

The approximate location of the "signal peptides" of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., Prot. Eng. 10:1-6 (1997) and von Heinje et al., Nucl. Acids. Res. 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

"PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid

sequence identity, alternatively at least about 91 % amino acid sequence identity, alternatively at least about 92 % amino acid sequence identity, alternatively at least about 93 % amino acid sequence identity, alternatively at least about 94 % amino acid sequence identity, alternatively at least about 95 % amino acid sequence identity, alternatively at least about 96 % amino acid sequence identity, alternatively at least about 97 % amino acid sequence identity, alternatively at least about 98 % amino acid sequence identity and alternatively at least about 99 % amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20 amino acids in length, alternatively at least about 30 amino acids in length, alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length, alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X," "Y" and "Z" each represent different hypothetical amino acid residues.

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an the amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

"PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 210 nucleotides in length, alternatively at least about 240 nucleotides in length, alternatively at least about 270 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA", wherein "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides.

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and

WO 01/16318

PCT/US00/23328

scoring matrix = BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO polypeptide as disclosed herein. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous

solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the PRO polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" PRO polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "antibody" is used in the broadest sense and specifically covers, for example, single anti-PRO monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO antibody compositions with polyepitopic specificity, single chain anti-PRO antibodies, and fragments of anti-PRO antibodies (see below). The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when

complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" may be identified as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any

immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

"Active" or "activity" for the purposes herein refers to form(s) of a PRO polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO polypeptide may comprise contacting a PRO polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO polypeptide.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-

forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata et al., Protein Eng. 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains

on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993).

5 An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or 10 nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

15 An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

20 By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes 25 a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

30 A "small molecule" is defined herein to have a molecular weight below about 500 Daltons.

Table 1

```

/*
*
* C-C increased from 12 to 15
* Z is average of EQ
5  * B is average of ND
* match with stop is _M; stop-stop = 0; J (joker) match = 0
*/
#define _M      -8      /* value of a match with a stop */

10 int  _day[26][26] = {
/*   A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */   { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */   { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */   {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
15 /* D */   { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */   { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */   {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */   { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */   {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
20 /* I */   {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */   { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */   {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */   {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */   {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
25 /* N */   { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */   {_M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, 0, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */   { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */   { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */   {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
30 /* S */   { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */   { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */   { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */   { 0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */   {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
35 /* X */   { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */   {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */   { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

Table 1 (cont')

```

/*
*/
#include <stdio.h>
#include <ctype.h>

5
#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

10
#define DMAT         3      /* value of matching bases */
#define DMIS         0      /* penalty for mismatched bases */
#define DINS0        8      /* penalty for a gap */
#define DINS1        1      /* penalty per base */
15
#define PINS0        8      /* penalty for a gap */
#define PINS1        4      /* penalty per residue */

struct jmp {
20
    short            n[MAXJMP];    /* size of jmp (neg for dely) */
    unsigned short   x[MAXJMP];    /* base no. of jmp in seq x */
};                                /* limits seq to 2^16 -1 */

struct diag {
25
    int              score;        /* score at last jmp */
    long             offset;       /* offset of prev block */
    short            jmp;         /* current jmp index */
    struct jmp        jp;         /* list of jmps */
};

30
struct path {
    int              spc;          /* number of leading spaces */
    short            n[JMP];      /* size of jmp (gap) */
    int              x[JMP];      /* loc of jmp (last elem before gap) */
};

35
char               *ofile;        /* output file name */
char               *namex[2];     /* seq names: getseqs() */
char               *prog;         /* prog name for err msgs */
char               *seqx[2];      /* seqs: getseqs() */
40
int                dmax;          /* best diag: nw() */
int                dmax0;         /* final diag */
int                dna;           /* set if dna: main() */
int                endgaps;       /* set if penalizing end gaps */
int                gapx, gapy;    /* total gaps in seqs */
45
int                len0, len1;     /* seq lens */
int                ngapx, ngapy;   /* total size of gaps */
int                smax;          /* max score: nw() */
int                *xbm;          /* bitmap for matching */
long               offset;        /* current offset in jmp file */
50
struct             diag           /* holds diagonals */
struct             path          /* holds path for seqs */
    *dx;
    pp[2];

char               *calloc(), *malloc(), *index(), *strcpy();
55
char               *getseq(), *g_calloc();

```

Table 1 (cont')

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
5 * The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
10 *
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
15 #include "day.h"

static _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};
20
static _pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
25 1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)
30     int      ac;
     char      *av[ ];
{
    prog = av[0];
    if (ac != 3) {
35         fprintf(stderr, "usage: %s file1 file2\n", prog);
         fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
         fprintf(stderr, "The sequences can be in upper- or lower-case\n");
         fprintf(stderr, "Any lines beginning with ';' or '<' are ignored\n");
         fprintf(stderr, "Output is in the file \"align.out\"\n");
         exit(1);
40     }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
45     xbm = (dna)? _dbval : _pbval;

    endgaps = 0;                /* 1 to penalize endgaps */
    ofile = "align.out";        /* output file */

50     nw();                    /* fill in the matrix, get the possible jmps */
    readjmps();                /* get the actual jmps */
    print();                    /* print stats, alignment */

55     cleanup(0);              /* unlink any tmp files */
}

```

main

Table 1 (cont')

```

/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/

```

```

nw()

```

```

nw

```

```

{
    char      *px, *py;          /* seqs and ptrs */
    int        *ndely, *dely;    /* keep track of dely */
    int        ndelx, delx;      /* keep track of delx */
    int        *tmp;            /* for swapping row0, row1 */
    int        mis;             /* score for each type */
    int        ins0, ins1;       /* insertion penalties */
    register   id;              /* diagonal index */
    register   ij;              /* jmp index */
    register   *col0, *col1;     /* score for curr, last row */
    register   xx, yy;          /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

Table 1 (cont')

...nw

```

5   for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
        mis = col0[yy-1];
        if (dna)
            mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
        else
            mis += _day[*px-'A'][*py-'A'];

10      /* update penalty for del in x seq;
        * favor new del over ongong del
        * ignore MAXGAP if weighting endgaps
        */
        if (endgaps || ndely[yy] < MAXGAP) {
            if (col0[yy] - ins0 >= dely[yy]) {
15                dely[yy] = col0[yy] - (ins0+ins1);
                ndely[yy] = 1;
            } else {
                dely[yy] -= ins1;
                ndely[yy]++;
20            }
        } else {
            if (col0[yy] - (ins0+ins1) >= dely[yy]) {
                dely[yy] = col0[yy] - (ins0+ins1);
                ndely[yy] = 1;
25            } else
                ndely[yy]++;
        }

30      /* update penalty for del in y seq;
        * favor new del over ongong del
        */
        if (endgaps || ndelx < MAXGAP) {
            if (col1[yy-1] - ins0 >= delx) {
35                delx = col1[yy-1] - (ins0+ins1);
                ndelx = 1;
            } else {
                delx -= ins1;
                ndelx++;
40            }
        } else {
            if (col1[yy-1] - (ins0+ins1) >= delx) {
                delx = col1[yy-1] - (ins0+ins1);
                ndelx = 1;
45            } else
                ndelx++;
        }

50      /* pick the maximum score; we're favoring
        * mis over any del and delx over dely
        */
55
60

```

Table 1 (cont')

...nw

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    coll[yy] = mis;
5   else if (delx >= dely[yy]) {
        coll[yy] = delx;
        ij = dx[id].ijmp;
        if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
10      && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
            dx[id].ijmp++;
            if (++ij >= MAXJMP) {
                writejumps(id);
                ij = dx[id].ijmp = 0;
                dx[id].offset = offset;
15      offset += sizeof(struct jmp) + sizeof(offset);
            }
        }
        dx[id].jp.n[ij] = ndelx;
        dx[id].jp.x[ij] = xx;
20      dx[id].score = delx;
    }
    else {
        coll[yy] = dely[yy];
        ij = dx[id].ijmp;
25      if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
            && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
                dx[id].ijmp++;
                if (++ij >= MAXJMP) {
                    writejumps(id);
                    ij = dx[id].ijmp = 0;
                    dx[id].offset = offset;
30      offset += sizeof(struct jmp) + sizeof(offset);
                }
            }
        dx[id].jp.n[ij] = -ndely[yy];
        dx[id].jp.x[ij] = xx;
        dx[id].score = dely[yy];
    }
40      if (xx == len0 && yy < len1) {
            /* last col
            */
            if (endgaps)
                coll[yy] -= ins0+ins1*(len1-yy);
            if (coll[yy] > smax) {
45      smax = coll[yy];
                dmax = id;
            }
        }
    }
50      if (endgaps && xx < len0)
        coll[yy-1] -= ins0+ins1*(len0-xx);
        if (coll[yy-1] > smax) {
            smax = coll[yy-1];
            dmax = id;
55      }
    }
    tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
60  (void) free((char *)col0);
    (void) free((char *)coll);
    }

```

Table 1 (cont')

```

/*
 *
 * print() -- only routine visible outside this module
 *
5  * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[ ]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
10 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

15 #include "nw.h"

#define SPC      3
#define P_LINE  256    /* maximum output line */
#define P_SPC    3      /* space between name or num and seq */

20 extern _day[26][26];
int olen;             /* set output line length */
FILE *fx;             /* output file */

25 print()                                     print
{
    int    lx, ly, firstgap, lastgap;    /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
30         fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "< first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "< second sequence: %s (length = %d)\n", namex[1], len1);
35     olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) {    /* leading gap in x */
40         pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
45         lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) {    /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
50     }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
55     getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

60

```

Table 1 (cont')

getmat

```

/*
 * trace back the best path, count matches
 */
static
5 getmat(lx, ly, firstgap, lastgap)
    int    lx, ly;          /* "core" (minus endgaps) */
    int    firstgap, lastgap; /* leading trailing overlap */
{
    int    nm, i0, i1, siz0, siz1;
10 char    outx[32];
    double pct;
    register n0, n1;
    register char *p0, *p1;

15 /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
20 n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
25         if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
30         else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
35         else {
            if (xbm[*p0-'A'] & xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
40             if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
45     }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
50     if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
55     pct = 100.*((double)nm)/((double)lx);
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);
60

```

Table 1 (cont')

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
        ngapx, (dna)? "base":"residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
            ngapy, (dna)? "base":"residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "< endgaps not penalized\n");
}

static nm; /* matches in core -- for checking */
static lmax; /* lengths of stripped file names */
static ij[2]; /* jmp index for a path */
static nc[2]; /* number at start of current line */
static ni[2]; /* current elem number -- for gapping */
static siz[2];
static char *ps[2]; /* ptr to current element */
static char *po[2]; /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp [ ]
 */
static
pr_align()
{
    int nn; /* char count */
    int more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(nameex[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

...getmat

pr_align

Table 1 (cont')

...pr_align

```

for (nn = nm = 0, more = 1; more; ) {
    for (i = more = 0; i < 2; i++) {
        /*
5         * do we have more of this sequence?
        */
        if (!*ps[i])
            continue;

10        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
15        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
20        }
        else { /* we're putting a seq element
            */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);
25            po[i]++;
            ps[i]++;

            /*
            * are we at next gap for this seq?
            */
30            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                * we need to merge all gaps
                * at this location
                */
35                siz[i] = pp[i].n[ij[i] + +];
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i] + +];

                }
                ni[i]++;
40            }
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nm = 0;
    }
50 }

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
55 static
dumpblock()
{
    register i;

60    for (i = 0; i < 2; i++)
        *po[i]-- = '\0';

```

dumpblock

Table 1 (cont')

...dumpblock

```

5      (void) putc('\n', fx);
      for (i = 0; i < 2; i++) {
          if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
              if (i == 0)
                  nums(i);
              if (i == 0 && *out[1])
                  stars();
10             putline(i);
              if (i == 0 && *out[1])
                  fprintf(fx, star);
              if (i == 1)
                  nums(i);
15             }
          }
      }

20  /*
   * put out a number line: dumpblock()
   */
   static
   nums(ix)
25  {
       int      ix;      /* index in out[] holding seq line */

       char      nline[P_LINE];
       register  i, j;
       register char *pn, *px, *py;

30     for (pn = nline, i = 0; i < lmax + P_SPC; i++, pn++)
         *pn = ' ';
       for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
           if (*py == ' ' || *py == '-')
               *pn = ' ';
35           else {
               if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                   j = (i < 0)? -i : i;
                   for (px = pn; j; j /= 10, px--)
                       *px = j%10 + '0';
40                   if (i < 0)
                       *px = '-';

                   }
               else
                   *pn = ' ';
45               i++;
           }
       }
       *pn = '\0';
       nc[ix] = i;
       for (pn = nline; *pn; pn++)
           (void) putc(*pn, fx);
       (void) putc('\n', fx);
50   }

55  /*
   * put out a line (name, [num], seq, [num]): dumpblock()
   */
   static
   putline(ix)
60  {
       int      ix;

```

nums

putline

Table 1 (cont')

...putline

```

5      int          i;
      register char *px;

      for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
          (void) putc(*px, fx);
      for (; i < lmax+P_SPC; i++)
          (void) putc(' ', fx);

10     /* these count from 1:
       * ni[] is current element (from 1)
       * nc[] is number at start of current line
       */
15     for (px = out[ix]; *px; px++)
          (void) putc(*px&0x7F, fx);
      (void) putc('\n', fx);
  }

20  /*
   * put a line of stars (seqs always in out[0], out[1]): dumpblock()
   */
   static
25  stars()
   {
       int          i;
       register char *p0, *p1, cx, *px;

30     if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
          !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
          return;
       px = star;
       for (i = lmax+P_SPC; i; i--)
35         *px++ = ' ';

       for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
           if (isalpha(*p0) && isalpha(*p1)) {

40                 if (xbm[*p0-'A']&xbm[*p1-'A']) {
                     cx = '*';
                     nm++;
                 }
                 else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
45                     cx = '.';
                 else
                     cx = ' ';
           }
           else
50             cx = ' ';
           *px++ = cx;
       }
       *px++ = '\n';
       *px = '\0';
55  }

```

stars

Table 1 (cont')

```
/*  
 * strip path or prefix from pn, return len: pr_align()  
 */
```

```
static
```

stripname

```
stripname(pn)
```

```
    char    *pn;    /* file name (may be path) */
```

```
{
```

```
    register char    *px, *py;
```

```
    py = 0;
```

```
    for (px = pn; *px; px++)
```

```
        if (*px == '/')
```

```
            py = px + 1;
```

```
    if (py)
```

```
        (void) strcpy(pn, py);
```

```
    return(strlen(pn));
```

```
}
```

Table 1 (cont')

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_calloc() -- calloc() with error checkin
5  * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>

10 char    *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE    *fj;

int      cleanup();                          /* cleanup tmp file */
15 long    lseek();

/*
 * remove any tmp file if we blow
 */
20 cleanup(i)                                cleanup
    int    i;
{
    if (fj)
        (void) unlink(jname);
25    exit(i);
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
30 char    *
getseq(file, len)                                getseq
35     char    *file;    /* file name */
    int      *len;      /* seq len */
{
    char      line[1024], *pseq;
    register char *px, *py;
    int      natgc, tlen;
    FILE      *fp;

    if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
45     exit(1);
    }
    tlen = natgc = 0;
    while (fgets(line, 1024, fp)) {
        if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
                tlen++;
    }
50     if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
    }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';
60

```

Table 1 (cont')

...getseq

```

py = pseq + 4;
*len = tlen;
rewind(fp);
5
while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
10        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
15            natgc++;
    }
}
*py++ = '\0';
*py = '\0';
20 (void) fclose(fp);
dna = natgc > (tlen/3);
return(pseq+4);
}

25 char *
g_alloc(msg, nx, sz)
    char *msg;          /* program, calling routine */
    int nx, sz;          /* number and size of elements */
{
30     char *px, *calloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
35             fprintf(stderr, "%s: g_alloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}
40
/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
 */
readjmps()
45 {
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

50     if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
55         }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
        while (1) {
60             for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)

```

g_alloc

readjmps

Table 1 (cont')

...readjumps

```

5      if (j < 0 && dx[dmax].offset && fj) {
        (void) lseek(fd, dx[dmax].offset, 0);
        (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
        (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
        dx[dmax].ijmp = MAXJMP-1;
      }
      else
        break;
10     }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
    }
15     if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
        if (siz < 0) { /* gap in second seq */
20             pp[1].n[i1] = -siz;
             xx += siz;
             /* id = xx - yy + len1 - 1
              */
             pp[1].x[i1] = xx - dmax + len1 - 1;
25             gapy++;
             ngapy -= siz;
        /* ignore MAXGAP when doing endgaps */
             siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
             i1++;
30         }
        else if (siz > 0) { /* gap in first seq */
             pp[0].n[i0] = siz;
             pp[0].x[i0] = xx;
             gapx++;
             ngapx += siz;
35         /* ignore MAXGAP when doing endgaps */
             siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
             i0++;
40         }
    }
    else
        break;
}

45     /* reverse the order of jumps
    */
    for (j = 0, i0--; j < i0; j++, i0--) {
        i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
        i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
50    }
    for (j = 0, i1--; j < i1; j++, i1--) {
        i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
        i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
55    }
    if (fd >= 0)
        (void) close(fd);
    if (fj) {
        (void) unlink(jname);
        fj = 0;
        offset = 0;
60    }
}

```

Table 1 (cont')

```

/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
5 writejumps(ix)                                     writejumps
    int    ix;
    {
        char    *mktemp();
10         if (!fj) {
            if (mktemp(jname) < 0) {
                fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
                cleanup(1);
            }
15         if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
20         (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
        (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
    }
}
25
30
35
40
45
50
55
60

```

Table 2

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXYYYYYYYYY	(Length = 12 amino acids)

5 % amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

10 5 divided by 15 = 33.3%

Table 3

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXYYYYYYZZYZ	(Length = 15 amino acids)

5 % amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

10 5 divided by 10 = 50%

Table 4

PRO-DNA	NNNNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLLLLLL	(Length = 16 nucleotides)

5 % nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

10 6 divided by 14 = 42.9%

Table 5

PRO-DNA	NNNNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLL	(Length = 9 nucleotides)

5 % nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

10 4 divided by 12 = 33.3%

II. Compositions and Methods of the Invention

A. Full-Length PRO Polypeptides

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding various PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

As disclosed in the Examples below, various cDNA clones have been deposited with the ATCC. The actual nucleotide sequences of those clones can readily be determined by the skilled artisan by sequencing of the deposited clone using routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

B. PRO Polypeptide Variants

In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence PRO or in various domains of the PRO described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO that results in a change in the amino acid sequence of the PRO as compared with the native sequence PRO. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

PRO polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO polypeptide.

5 PRO fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired
10 termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO polypeptide fragments share at least one biological and/or immunological activity with the native PRO polypeptide disclosed herein.

In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial
15 changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table 6

	Original Residue	Exemplary Substitutions	Preferred Substitutions
5	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
10	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
	Ile (I)	leu; val; met; ala; phe;	
15		norleucine	leu
	Leu (L)	norleucine; ile; val;	
		met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
20	Phe (F)	leu; val; ile; ala; tyr	leu
	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr; phe	tyr
25	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe;	
		ala; norleucine	leu

30 Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- 35 (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

40 Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA,

45

317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

C. Modifications of PRO

Covalent modifications of PRO are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the PRO. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the

desired amino acids.

Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

5 Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety
10 of endo- and exo-glycosidases as described by Thotakura et al., Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of PRO comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

15 The PRO of the present invention may also be modified in a way to form a chimeric molecule comprising PRO fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO. The presence of such epitope-tagged forms of the PRO can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO to
20 be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the
25 Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

30 In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred
35 embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

D. Preparation of PRO

The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

1. Isolation of DNA Encoding PRO

DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., *supra*; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ³²P-labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan'*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7*

ilvG kan^r; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., Bio/Technology, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906; Van den Berg et al., Bio/Technology, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol., 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983]; Tilburn et al., Gene, 26:205-221 [1983]; Yelton et al., Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4:475-479 [1985]). Methylotropic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylotrophs, 269 (1982).

Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol., 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally

include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

5 The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g.,
10 the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral
15 secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for
20 cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

25 An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene
30 provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well
35 known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid

promoters such as the tac promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

5 Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

10 Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

15 PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

20 Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 25 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

30 Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

35 Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

5. Purification of Polypeptide

Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

E. Uses for PRO

Nucleotide sequences (or their complement) encoding PRO have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. PRO nucleic acid will also be useful for the preparation of PRO polypeptides by the recombinant techniques described herein.

The full-length native sequence PRO gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length PRO cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO or PRO from other species) which have a desired sequence identity to the native PRO sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence PRO. By way of example, a screening method will comprise isolating the coding region of the PRO gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ^{32}P or ^{35}S , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

Other useful fragments of the PRO nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target PRO mRNA (sense) or PRO DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of PRO DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (Cancer Res. 48:2659, 1988) and van der Krol et al. (BioTechniques 6:958, 1988).

Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of PRO proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable *in vivo* (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO_4 -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

Antisense or sense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO coding sequences.

Nucleotide sequences encoding a PRO can also be used to construct hybridization probes for mapping the gene which encodes that PRO and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as *in situ* hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

When the coding sequences for PRO encode a protein which binds to another protein (example, where the PRO is a receptor), the PRO can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO or a receptor for PRO. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small

molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

Nucleic acids which encode PRO or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of PRO can be used to construct a PRO "knock out" animal which has a defective or altered gene encoding PRO as a result of homologous recombination between the endogenous gene encoding PRO and altered genomic DNA encoding PRO introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques. A portion of the genomic DNA encoding PRO can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of

the PRO polypeptide.

Nucleic acid encoding the PRO polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve *in vivo* synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. "Gene therapy" includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes *in vivo*. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik *et al.*, Proc. Natl. Acad. Sci. USA 83:4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred *in vivo* gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau *et al.*, Trends in Biotechnology 11, 205-210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, J. Biol. Chem. 262, 4429-4432 (1987); and Wagner *et al.*, Proc. Natl. Acad. Sci. USA 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson *et al.*, Science 256, 808-813 (1992).

The PRO polypeptides described herein may also be employed as molecular weight markers for protein electrophoresis purposes and the isolated nucleic acid sequences may be used for recombinantly expressing those markers.

The nucleic acid molecules encoding the PRO polypeptides or fragments thereof described herein are useful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each PRO nucleic acid molecule of the present invention can be used as a chromosome marker.

The PRO polypeptides and nucleic acid molecules of the present invention may also be used diagnostically for tissue typing, wherein the PRO polypeptides of the present invention may be differentially expressed in one tissue as compared to another, preferably in a diseased tissue as compared to a normal tissue of the same tissue type. PRO nucleic acid molecules will find use for generating probes for PCR, Northern

analysis, Southern analysis and Western analysis.

The PRO polypeptides described herein may also be employed as therapeutic agents. The PRO polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the PRO product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient
5 having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins,
10 such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, PLURONICS™ or PEG.

15 The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

20 The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician. Animal experiments provide reliable guidance
25 for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" In *Toxicokinetics and New Drug Development*, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42-96.

30 When *in vivo* administration of a PRO polypeptide or agonist or antagonist thereof is employed, normal dosage amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 µg/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may
35 necessitate delivery in a manner different from that to another organ or tissue.

Where sustained-release administration of a PRO polypeptide is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of the PRO

polypeptide, microencapsulation of the PRO polypeptide is contemplated. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon- (rhIFN-), interleukin-2, and MN rgp120. Johnson et al., Nat. Med., 2:795-799 (1996); Yasuda, Biomed. Ther., 27:1221-1223 (1993); Hora et al., Bio/Technology, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems," in Vaccine Design: The Subunit and Adjuvant Approach, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

The sustained-release formulations of these proteins were developed using poly-lactic-coglycolic acid (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), Biodegradable Polymers as Drug Delivery Systems (Marcel Dekker: New York, 1990), pp. 1-41.

This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, Nature (London), 340:245-246 (1989); Chien et al., Proc. Natl. Acad. Sci. USA, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, Proc. Natl. Acad. Sci. USA, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

To assay for antagonists, the PRO polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., Current Protocols in Immun., 1(2): Chapter 5 (1991).

Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res., 6:3073 (1979); Cooney et al., Science, 241: 456 (1988); Dervan et al., Science, 251:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the PRO polypeptide (antisense - Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, FL, 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of the PRO

polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, Current Biology, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, *supra*.

These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

Diagnostic and therapeutic uses of the herein disclosed molecules may also be based upon the positive functional assay hits disclosed and described below.

F. Anti-PRO Antibodies

The present invention further provides anti-PRO antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

1. Polyclonal Antibodies

The anti-PRO antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

2. Monoclonal Antibodies

The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression
5 vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison et al., *supra*] or by covalently joining to the immunoglobulin coding
10 sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well
15 known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to
20 produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

3. Human and Humanized Antibodies

The anti-PRO antibodies of the invention may further comprise humanized antibodies or human
25 antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human
30 species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR
35 regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones

et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985) and Boerner et al., J. Immunol., 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks *et al.*, Bio/Technology 10, 779-783 (1992); Lonberg *et al.*, Nature 368 856-859 (1994); Morrison, Nature 368, 812-13 (1994); Fishwild *et al.*, Nature Biotechnology 14, 845-51 (1996); Neuberger, Nature Biotechnology 14, 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol. 13 65-93 (1995).

The antibodies may also be affinity matured using known selection and/or mutagenesis methods as described above. Preferred affinity matured antibodies have an affinity which is five times, more preferably 10 times, even more preferably 20 or 30 times greater than the starting antibody (generally murine, humanized or human) from which the matured antibody is prepared.

4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain

pairs, where the two heavy chains have different specificities [Milstein and Cuello, Nature, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., EMBO J., 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby *et al.*, J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers.

5 This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with

10 the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber *et al.*, J. Immunol. 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.*, J. Immunol. 147:60 (1991).

15 Exemplary bispecific antibodies may bind to two different epitopes on a given PRO polypeptide herein. Alternatively, an anti-PRO polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO polypeptide. Bispecific antibodies may also be used to localize

20 cytotoxic agents to cells which express a particular PRO polypeptide. These antibodies possess a PRO-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO polypeptide and further binds tissue factor (TF).

25 5. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared *in vitro*

30 using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

35 6. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, *e.g.*, the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be

introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, J. Exp Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.* Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.*, Anti-Cancer Drug Design, 3: 219-230 (1989).

7. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ²¹²Bi, ¹³¹I, ¹³¹In, ⁹⁰Y, and ¹⁸⁶Re.

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (*e.g.*, avidin) that is conjugated to a cytotoxic agent (*e.g.*, a radionucleotide).

8. Immunoliposomes

The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos.

4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem., **257**: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon *et al.*, J. National Cancer Inst., **81**(19): 1484 (1989).

9. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, *e.g.*, Marasco *et al.*, Proc. Natl. Acad. Sci. USA, **90**: 7889-7893 (1993). The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl

WO 01/16318

PCT/US00/23328

acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

G. Uses for anti-PRO Antibodies

The anti-PRO antibodies of the invention have various utilities. For example, anti-PRO antibodies may be used in diagnostic assays for PRO, *e.g.*, detecting its expression (and in some cases, differential expression) in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, Monoclonal Antibodies: A Manual of Techniques, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ³H, ¹⁴C, ³²P, ³⁵S, or ¹²⁵I, a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., Nature, 144:945 (1962); David et al., Biochemistry, 13:1014 (1974); Pain et al., J. Immunol. Meth., 40:219 (1981); and Nygren, J. Histochem. and Cytochem., 30:407 (1982).

Anti-PRO antibodies also are useful for the affinity purification of PRO from recombinant cell culture or natural sources. In this process, the antibodies against PRO are immobilized on a suitable support, such a Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO from the antibody.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA.

5

EXAMPLE 1: Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (e.g., Dayhoff, GenBank), and proprietary databases (e.g. LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST-2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, WA).

15

Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST or BLAST-2 and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

20

25

Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., Current Protocols in Molecular Biology, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

30

The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

35

EXAMPLE 2: Isolation of cDNA clones by Amylase Screening**1. Preparation of oligo dT primed cDNA library**

mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, CA (Fast Track^{*} 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, MD (Super Script Plasmid System^{*}).

5 In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the Sall/NotI linker cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

2. Preparation of random primed cDNA library

10 A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, linker with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI

15 cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

3. Transformation and Detection

20 DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was

25 added and the mixture was incubated at 37°C for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37°C). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, e.g. CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

30 The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

35 The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL⁺, SUC⁺, GAL⁺. Preferably, yeast mutants can be employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in *sec71*, *sec72*, *sec62*, with truncated *sec71* being most preferred. Alternatively, antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins

implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

Transformation was performed based on the protocol outlined by Gietz et al., Nucl. Acid. Res., 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30°C. The YEPD broth was prepared as described in Kaiser et al., Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 207 (1994). The overnight culture was then diluted to about 2×10^6 cells/ml (approx. $OD_{600}=0.1$) into fresh YEPD broth (500 ml) and regrown to 1×10^7 cells/ml (approx. $OD_{600}=0.4-0.5$).

The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li_2OOCCH_3), and resuspended into LiAc/TE (2.5 ml).

Transformation took place by mixing the prepared cells (100 μ l) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, MD) and transforming DNA (1 μ g, vol. < 10 μ l) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 μ l, 40% polyethylene glycol-4000, 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li_2OOCCH_3 , pH 7.5) was added. This mixture was gently mixed and incubated at 30°C while agitating for 30 minutes. The cells were then heat shocked at 42°C for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 μ l, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 μ l) were spread onto the selective media previously prepared in 150 mm growth plates (VWR).

Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser et al., Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 208-210 (1994). Transformants were grown at 30°C for 2-3 days.

The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was coupled to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely et al., Anal. Biochem., 172:176-179 (1988). The coupled starch was incorporated into the SCD-Ura agar plates at a final concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final concentration).

The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

4. Isolation of DNA by PCR Amplification

When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30 μ l) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5 μ l) was used as a template for the PCR reaction in a 25 μ l volume containing: 0.5 μ l Klen*taq (Clontech, Palo Alto, CA); 4.0 μ l 10 mM dNTP's (Perkin Elmer-Cetus); 2.5 μ l Kentaq buffer (Clontech); 0.25 μ l forward oligo 1; 0.25 μ l reverse oligo 2; 12.5 μ l distilled water. The sequence of the forward oligonucleotide 1 was:

5'-TGTA*AAACGACGGCCAGTTAAATAGACCTGCAATTATTAATCT-3' (SEQ ID NO:169)

The sequence of reverse oligonucleotide 2 was:

5'-CAGGAAACAGCTATGACCACCTGCACACCTGCAAATCCATT-3' (SEQ ID NO:170)

PCR was then performed as follows:

- | | | |
|----|---------------|------------------|
| a. | Denature | 92°C, 5 minutes |
| b. | 3 cycles of: | |
| | Denature | 92°C, 30 seconds |
| | Anneal | 59°C, 30 seconds |
| | Extend | 72°C, 60 seconds |
| c. | 3 cycles of: | |
| | Denature | 92°C, 30 seconds |
| | Anneal | 57°C, 30 seconds |
| | Extend | 72°C, 60 seconds |
| d. | 25 cycles of: | |
| | Denature | 92°C, 30 seconds |
| | Anneal | 55°C, 30 seconds |
| | Extend | 72°C, 60 seconds |
| e. | Hold | 4°C |

The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

Following the PCR, an aliquot of the reaction (5 μ l) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook et al., *supra*. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, CA).

EXAMPLE 3: Isolation of cDNA Clones Using Signal Algorithm Analysis

Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc. (South San Francisco, CA) upon ESTs as well as clustered and assembled EST fragments from public (e.g., GenBank) and/or private (LIFESEQ®, Incyte Pharmaceuticals, Inc., Palo Alto, CA) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine

codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

EXAMPLE 4: Isolation of cDNA Clones Encoding Human PRO Polypeptides

Using the techniques described in Examples 1 to 3 above, numerous full-length cDNA clones were identified as encoding PRO polypeptides as disclosed herein. These cDNAs were then deposited under the terms of the Budapest Treaty with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC) as shown in Table 7 below.

Table 7

	<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
15	DNA26843-1389	203099	August 4, 1998
	DNA30867-1335	209807	April 28, 1998
	DNA34431-1177	209399	October 17, 1997
	DNA38268-1188	209421	October 28, 1997
20	DNA40621-1440	209922	June 2, 1998
	DNA40625-1189	209788	April 21, 1998
	DNA45409-2511	203579	January 12, 1999
	DNA45495-1550	203156	August 25, 1998
	DNA49820-1427	209932	June 2, 1998
25	DNA56406-1704	203478	November 17, 1998
	DNA56410-1414	209923	June 2, 1998
	DNA56436-1448	209902	May 27, 1998
	DNA56855-1447	203004	June 23, 1998
	DNA56860-1510	209952	June 9, 1998
30	DNA56862-1343	203174	September 1, 1998
	DNA56868-1478	203024	June 23, 1998
	DNA56869-1545	203161	August 25, 1998
	DNA57704-1452	209953	June 9, 1998
	DNA58723-1588	203133	August 18, 1998
35	DNA57827-1493	203045	July 1, 1998
	DNA58737-1473	203136	August 18, 1998
	DNA58846-1409	209957	June 9, 1998
	DNA58850-1495	209956	June 9, 1998
	DNA58855-1422	203018	June 23, 1998
40	DNA59211-1450	209960	June 9, 1998
	DNA59212-1627	203245	September 9, 1998
	DNA59213-1487	209959	June 9, 1998
	DNA59605-1418	203005	June 23, 1998
	DNA59609-1470	209963	June 9, 1998
45	DNA59610-1556	209990	June 16, 1998
	DNA59837-2545	203658	February 9, 1999
	DNA59844-2542	203650	February 9, 1999
	DNA59854-1459	209974	June 16, 1998

Table 7 (cont')

	DNA60625-1507	209975	June 16, 1998
	DNA60629-1481	209979	June 16, 1998
	DNA61755-1554	203112	August 11, 1998
5	DNA62812-1594	203248	September 9, 1998
	DNA62815-1576	203247	September 9, 1998
	DNA64881-1602	203240	September 9, 1998
	DNA64886-1601	203241	September 9, 1998
	DNA64902-1667	203317	October 6, 1998
10	DNA64950-1590	203224	September 15, 1998
	DNA65403-1565	203230	September 15, 1998
	DNA66308-1537	203159	August 25, 1998
	DNA66519-1535	203236	September 15, 1998
	DNA66521-1583	203225	September 15, 1998
15	DNA66658-1584	203229	September 15, 1998
	DNA66660-1585	203279	September 22, 1998
	DNA66663-1598	203268	September 22, 1998
	DNA66674-1599	203281	September 22, 1998
	DNA68862-2546	203652	February 9, 1999
20	DNA68866-1644	203283	September 22, 1998
	DNA68871-1638	203280	September 22, 1998
	DNA68880-1676	203319	October 6, 1998
	DNA68883-1691	203535	December 15, 1998
	DNA68885-1678	203311	October 6, 1998
25	DNA71277-1636	203285	September 22, 1998
	DNA73727-1673	203459	November 3, 1998
	DNA73734-1680	203363	October 20, 1998
	DNA73735-1681	203356	October 20, 1998
	DNA76393-1664	203323	October 6, 1998
30	DNA77301-1708	203407	October 27, 1998
	DNA77568-1626	203134	August 18, 1998
	DNA77626-1705	203536	December 15, 1998
	DNA81754-2532	203542	December 15, 1998
	DNA81757-2512	203543	December 15, 1998
35	DNA82302-2529	203534	December 15, 1998
	DNA82340-2530	203547	December 22, 1998
	DNA83500-2506	203391	October 29, 1998
	DNA84920-2614	203966	April 27, 1999
	DNA85066-2534	203588	January 12, 1999
40	DNA86571-2551	203660	February 9, 1999
	DNA87991-2540	203656	February 9, 1999
	DNA92238-2539	203602	January 20, 1999
	DNA96042-2682	PTA-382	July 20, 1999
	DNA96787-2534	203589	January 12, 1999
45	DNA125185-2806	PTA-1031	December 7, 1999
	DNA147531-2821	PTA-1185	January 11, 2000
	DNA115291-2681	PTA-202	June 8, 1999
	DNA164625-28890	PTA-1535	March 21, 2000
	DNA131639-2874	PTA-1784	April 25, 2000
50	DNA79230-2525	203549	December 22, 1998

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The

deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks

5

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

10

EXAMPLE 5: Use of PRO as a hybridization probe

The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

15

DNA comprising the coding sequence of full-length or mature PRO as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

20

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-derived probe to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO can then be identified using standard techniques known in the art.

25

EXAMPLE 6: Expression of PRO in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

30

The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO coding region, lambda transcriptional terminator, and an argU gene.

35

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

5 Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

10 After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

15 PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110
20 *fuH*A(tonA) *lon* *galE* *rpoH*ts(htpRts) *clpP*(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30°C with shaking until an O.D.600 of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate•2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30°C with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

25 *E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4°C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The
30 clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4°C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

35 The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The

refolding solution is stirred gently at 4°C for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros^{*} R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine^{*} (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 7: Expression of PRO in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook et al., *supra*. The resulting vector is called pRK5-PRO.

In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 µg pRK5-PRO DNA is mixed with about 1 µg DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 µl of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 µl of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 µCi/ml ³⁵S-cysteine and 200 µCi/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in

serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Sompanyrac et al., Proc. Natl. Acad. Sci., 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 μ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 μ g/ml bovine insulin and 0.1 μ g/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO_4 or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ^{35}S -methionine. After determining the presence of PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO can then be concentrated and purified by any selected method.

Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni^{2+} -chelate affinity chromatography.

PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., Current Protocols of Molecular Biology, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., Nucl. Acids Res. 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect* (Quiagen), Dosper* or Fugene* (Boehringer Mannheim). The cells are grown as described in Lucas et al., *supra*. Approximately 3×10^7 cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2 μ m filtered PS20 with 5% 0.2 μ m diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37°C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with 3×10^5 cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2×10^6 cells/mL. On day 0, the cell number pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33°C, and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate was either stored at 4°C or immediately loaded onto columns for purification.

For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 8: Expression of PRO in Yeast

The following method describes recombinant expression of PRO in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 9: Expression of PRO in Baculovirus-Infected Insect Cells

The following method describes recombinant expression of PRO in Baculovirus-infected insect cells.

The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley et al., Baculovirus expression vectors: A Laboratory Manual, Oxford: Oxford University Press (1994).

Expressed poly-his tagged PRO can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., Nature, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 μm

filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching
5 A₂₈₀ baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged PRO are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known
10 chromatography techniques, including for instance, Protein A or protein G column chromatography.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 10: Preparation of Antibodies that Bind PRO

This example illustrates preparation of monoclonal antibodies which can specifically bind PRO.

15 Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

20 Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be
25 boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells
30 which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

35 The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively,

affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

EXAMPLE 11: Purification of PRO Polypeptides Using Specific Antibodies

Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (*e.g.*, high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (*e.g.*, a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

EXAMPLE 12: Drug Screening

This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with

an PRO polypeptide or fragment thereof and assaying (I) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

EXAMPLE 13: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (*i.e.*, a PRO polypeptide) or of small molecules with which they interact, *e.g.*, agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide *in vivo* (*c.f.*, Hodgson, Bio/Technology, 9: 19-21 (1991)).

In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, Biochemistry, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda *et al.*, J. Biochem., 113:742-746 (1993).

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic

antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

EXAMPLE 14: Pericyte c-Fos Induction (Assay 93)

This assay shows that certain polypeptides of the invention act to induce the expression of c-fos in pericyte cells and, therefore, are useful not only as diagnostic markers for particular types of pericyte-associated tumors but also for giving rise to antagonists which would be expected to be useful for the therapeutic treatment of pericyte-associated tumors. Induction of c-fos expression in pericytes is also indicative of the induction of angiogenesis and, as such, PRO polypeptides capable of inducing the expression of c-fos would be expected to be useful for the treatment of conditions where induced angiogenesis would be beneficial including, for example, wound healing, and the like. Specifically, on day 1, pericytes are received from VEC Technologies and all but 5 ml of media is removed from flask. On day 2, the pericytes are trypsinized, washed, spun and then plated onto 96 well plates. On day 7, the media is removed and the pericytes are treated with 100 μ l of PRO polypeptide test samples and controls (positive control = DME+5% serum +/- PDGF at 500 ng/ml; negative control = protein 32). Replicates are averaged and SD/CV are determined. Fold increase over Protein 32 (buffer control) value indicated by chemiluminescence units (RLU) luminometer reading verses frequency is plotted on a histogram. Two-fold above Protein 32 value is considered positive for the assay. ASY Matrix: Growth media = low glucose DMEM = 20% FBS + 1X pen strep + 1X fungizone. Assay Media = low glucose DMEM +5% FBS.

The following polypeptides tested positive in this assay: PRO1347 and PRO1340.

EXAMPLE 15: Ability of PRO Polypeptides to Stimulate the Release of Proteoglycans from Cartilage (Assay 97)

The ability of various PRO polypeptides to stimulate the release of proteoglycans from cartilage tissue was tested as follows.

The metacarpophalangeal joint of 4-6 month old pigs was aseptically dissected, and articular cartilage was removed by free hand slicing being careful to avoid the underlying bone. The cartilage was minced and cultured in bulk for 24 hours in a humidified atmosphere of 95% air, 5% CO₂ in serum free (SF) media (DME/F12 1:1) with 0.1% BSA and 100U/ml penicillin and 100 μ g/ml streptomycin. After washing three times, approximately 100 mg of articular cartilage was aliquoted into micronics tubes and incubated for an additional 24 hours in the above SF media. PRO polypeptides were then added at 1% either alone or in combination with 18 ng/ml interleukin-1 α , a known stimulator of proteoglycan release from cartilage tissue.

The supernatant was then harvested and assayed for the amount of proteoglycans using the 1,9-dimethyl-methylene blue (DMB) colorimetric assay (Farndale and Buttle, Biochem. Biophys. Acta 883:173-177 (1985)). A positive result in this assay indicates that the test polypeptide will find use, for example, in the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis.

When various PRO polypeptides were tested in the above assay, the polypeptides demonstrated a marked ability to stimulate release of proteoglycans from cartilage tissue both basally and after stimulation with interleukin-1 α and at 24 and 72 hours after treatment, thereby indicating that these PRO polypeptides are useful for stimulating proteoglycan release from cartilage tissue. As such, these PRO polypeptides are useful for the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis. The polypeptides testing positive in this assay are: PRO1565, PRO1693, PRO1801 and PRO10096.

EXAMPLE 16: Detection of Polypeptides That Affect Glucose or FFA Uptake in Skeletal Muscle (Assay 106)

This assay is designed to determine whether PRO polypeptides show the ability to affect glucose or FFA uptake by skeletal muscle cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where either the stimulation or inhibition of glucose uptake by skeletal muscle would be beneficial including, for example, diabetes or hyper- or hypo-insulinemia.

In a 96 well format, PRO polypeptides to be assayed are added to primary rat differentiated skeletal muscle, and allowed to incubate overnight. Then fresh media with the PRO polypeptide and +/- insulin are added to the wells. The sample media is then monitored to determine glucose and FFA uptake by the skeletal muscle cells. The insulin will stimulate glucose and FFA uptake by the skeletal muscle, and insulin in media without the PRO polypeptide is used as a positive control, and a limit for scoring. As the PRO polypeptide being tested may either stimulate or inhibit glucose and FFA uptake, results are scored as positive in the assay if greater than 1.5 times or less than 0.5 times the insulin control.

The following PRO polypeptides tested positive as either stimulators or inhibitors of glucose and/or FFA uptake in this assay: PRO4405.

EXAMPLE 17: Identification of PRO Polypeptides That Stimulate TNF- α Release In Human Blood (Assay 128)

This assay shows that certain PRO polypeptides of the present invention act to stimulate the release of TNF- α in human blood. PRO polypeptides testing positive in this assay are useful for, among other things, research purposes where stimulation of the release of TNF- α would be desired and for the therapeutic treatment of conditions wherein enhanced TNF- α release would be beneficial. Specifically, 200 μ l of human blood supplemented with 50mM Hepes buffer (pH 7.2) is aliquotted per well in a 96 well test plate. To each well is then added 300 μ l of either the test PRO polypeptide in 50 mM Hepes buffer (at various concentrations) or 50 mM Hepes buffer alone (negative control) and the plates are incubated at 37°C for 6 hours. The samples are then centrifuged and 50 μ l of plasma is collected from each well and tested for the presence of TNF- α by ELISA assay. A positive in the assay is a higher amount of TNF- α in the PRO polypeptide treated samples as compared to the negative control samples.

The following PRO polypeptides tested positive in this assay: PRO263, PRO295, PRO1282, PRO1063, PRO1356, PRO3543, and PRO5990.

EXAMPLE 18: Tumor Versus Normal Differential Tissue Expression Distribution

Oligonucleotide probes were constructed from some of the PRO polypeptide-encoding nucleotide sequences shown in the accompanying figures for use in quantitative PCR amplification reactions. The oligonucleotide probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human tumor and normal human tissue samples and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO polypeptide-encoding nucleic acid in the various tumor and normal tissues tested. β -actin was used as a control to assure that equivalent amounts of nucleic acid was used in each reaction. Identification of the differential expression of the PRO polypeptide-encoding nucleic acid in one or more tumor tissues as compared to one or more normal tissues of the same tissue type renders the molecule useful diagnostically for the determination of the presence or absence of tumor in a subject suspected of possessing a tumor as well as therapeutically as a target for the treatment of a tumor in a subject possessing such a tumor. These assays provided the following results.

<u>Molecule</u>	<u>is more highly expressed in:</u>	<u>as compared to:</u>
DNA26843-1389	normal lung rectum tumor	lung tumor normal rectum
DNA30867-1335	normal kidney	kidney tumor
DNA40621-1440	normal lung	lung tumor
DNA40625-1189	normal lung	lung tumor
DNA45409-2511	melanoma tumor	normal skin
DNA56406-1704	kidney tumor normal skin	normal kidney melanoma tumor
DNA56410-1414	normal stomach	stomach tumor
DNA56436-1448	normal skin	melanoma tumor
DNA56855-1447	normal esophagus rectum tumor	esophageal tumor normal rectum
DNA56860-1510	normal kidney rectum tumor	kidney tumor normal rectum
DNA56862-1343	kidney tumor normal lung	normal kidney lung tumor

WO 01/16318

PCT/US00/23328

	<u>Molecule</u>	<u>is more highly expressed in:</u>	<u>as compared to:</u>
	DNA56868-1478	normal stomach normal lung	stomach tumor lung tumor
5	DNA56869-1545	normal esophagus normal skin	esophageal tumor melanoma tumor
	DNA57704-1452	normal stomach rectum tumor	stomach tumor normal rectum
10	DNA58723-1588	normal stomach kidney tumor normal skin	stomach tumor normal kidney melanoma tumor
15	DNA57827-1493	normal stomach normal skin	stomach tumor melanoma tumor
	DNA58737-1473	esophageal tumor normal stomach	normal esophagus stomach tumor
20	DNA58846-1409	lung tumor	normal lung
	DNA58850-1495	esophageal tumor kidney tumor	normal esophagus normal kidney
25	DNA58855-1422	normal stomach rectum tumor	stomach tumor normal rectum
	DNA59211-1450	normal kidney	kidney tumor
30	DNA59212-1627	normal skin	melanoma tumor
	DNA59213-1487	normal stomach normal skin	stomach tumor melanoma tumor
35	DNA59605-1418	melanoma tumor	normal skin
	DNA59609-1470	esophageal tumor	normal esophagus
40	DNA59610-1556	esophageal tumor lung tumor normal skin	normal esophagus normal lung melanoma tumor
	DNA59837-2545	normal skin	melanoma tumor
45	DNA59844-2542	normal skin esophageal tumor	melanoma tumor normal esophagus
	DNA59854-1459	normal esophagus stomach tumor normal lung	esophageal tumor normal stomach lung tumor
50	DNA60625-1507	normal lung	lung tumor
55	DNA60629-1481	normal esophagus normal rectum	esophageal tumor rectum tumor

	<u>Molecule</u>	<u>is more highly expressed in:</u>	<u>as compared to:</u>
	DNA61755-1554	normal stomach kidney tumor	stomach tumor normal kidney
5	DNA62812-1594	normal stomach normal lung normal rectum normal skin	stomach tumor lung tumor rectum tumor melanoma tumor
10	DNA62815-1576	esophageal tumor	normal esophagus
	DNA64881-1602	normal stomach normal lung	stomach tumor lung tumor
15	DNA64902-1667	esophageal tumor kidney tumor	normal esophagus normal kidney
	DNA65403-1565	normal esophagus	esophageal tumor
20	DNA66308-1537	normal lung	lung tumor
	DNA66519-1535	kidney tumor	normal kidney
25	DNA66521-1583	normal esophagus normal stomach normal lung normal rectum normal skin	esophageal tumor stomach tumor lung tumor rectum tumor melanoma tumor
30	DNA66658-1584	normal lung melanoma tumor	lung tumor normal skin
	DNA66660-1585	lung tumor	normal lung
35	DNA66674-1599	kidney tumor normal lung	normal kidney lung tumor
	DNA68862-2546	melanoma tumor	normal skin
40	DNA68866-1644	normal stomach	stomach tumor
	DNA68871-1638	lung tumor normal skin	normal lung melanoma tumor
45	DNA68880-1676	normal lung normal skin	lung tumor melanoma tumor
	DNA68883-1691	esophageal tumor	normal esophagus
50	DNA68885-1678	lung tumor	normal lung
	DNA71277-1636	normal stomach	stomach tumor
55	DNA73734-1680	normal lung	lung tumor

	<u>Molecule</u>	<u>is more highly expressed in:</u>	<u>as compared to:</u>
5	DNA73735-1681	esophageal tumor normal kidney lung tumor normal skin	normal esophagus kidney tumor normal lung melanoma tumor
	DNA76393-1664	esophageal tumor stomach tumor lung tumor rectum tumor	normal esophagus normal stomach normal lung normal rectum
10	DNA77568-1626	normal stomach lung tumor	stomach tumor normal lung
	DNA77626-1705	normal rectum	rectum tumor
15	DNA81754-2532	normal skin	melanoma tumor
	DNA81757-2512	esophageal tumor normal stomach melanoma tumor	normal esophagus stomach tumor normal skin
20	DNA82302-2529	normal stomach normal lung	stomach tumor lung tumor
25	DNA82340-2530	normal esophagus	esophageal tumor
	DNA85066-2534	lung tumor normal skin	normal lung melanoma tumor
30	DNA87991-2540	esophageal tumor	normal esophagus
	DNA92238-2539	normal skin	melanoma tumor
35	DNA96787-2534	normal kidney	kidney tumor

EXAMPLE 19: Identification of Receptor/Ligand Interactions

In this assay, various PRO polypeptides are tested for ability to bind to a panel of potential receptor or ligand molecules for the purpose of identifying receptor/ligand interactions. The identification of a ligand for a known receptor, a receptor for a known ligand or a novel receptor/ligand pair is useful for a variety of indications including, for example, targeting bioactive molecules (linked to the ligand or receptor) to a cell known to express the receptor or ligand, use of the receptor or ligand as a reagent to detect the presence of the ligand or receptor in a composition suspected of containing the same, wherein the composition may comprise cells suspected of expressing the ligand or receptor, modulating the growth of or another biological or immunological activity of a cell known to express or respond to the receptor or ligand, modulating the immune response of cells or toward cells that express the receptor or ligand, allowing the preparation of agonists, antagonists and/or antibodies directed against the receptor or ligand which will modulate the growth of or a biological or immunological activity of a cell expressing the receptor or ligand, and various other indications which will be readily apparent to the ordinarily skilled artisan.

The assay is performed as follows. A PRO polypeptide of the present invention suspected of being a ligand for a receptor is expressed as a fusion protein containing the Fc domain of human IgG (an immunoadhesin). Receptor-ligand binding is detected by allowing interaction of the immunoadhesin polypeptide with cells (e.g. Cos cells) expressing candidate PRO polypeptide receptors and visualization of bound immunoadhesin with fluorescent reagents directed toward the Fc fusion domain and examination by microscope.

- 5 Cells expressing candidate receptors are produced by transient transfection, in parallel, of defined subsets of a library of cDNA expression vectors encoding PRO polypeptides that may function as receptor molecules. Cells are then incubated for 1 hour in the presence of the PRO polypeptide immunoadhesin being tested for possible receptor binding. The cells are then washed and fixed with paraformaldehyde. The cells are then incubated with fluorescent conjugated antibody directed against the Fc portion of the PRO polypeptide immunoadhesin (e.g. FITC conjugated goat anti-human-Fc antibody). The cells are then washed again and examined by microscope. A positive interaction is judged by the presence of fluorescent labeling of cells transfected with cDNA encoding a particular PRO polypeptide receptor or pool of receptors and an absence of similar fluorescent labeling of similarly prepared cells that have been transfected with other cDNA or pools of cDNA. If a defined pool of cDNA expression vectors is judged to be positive for interaction with a PRO polypeptide immunoadhesin, the individual cDNA species that comprise the pool are tested individually (the pool is "broken down") to determine the specific cDNA that encodes a receptor able to interact with the PRO polypeptide immunoadhesin.

- 20 In another embodiment of this assay, an epitope-tagged potential ligand PRO polypeptide (e.g. 8 histidine "His" tag) is allowed to interact with a panel of potential receptor PRO polypeptide molecules that have been expressed as fusions with the Fc domain of human IgG (immunoadhesins). Following a 1 hour co-incubation with the epitope tagged PRO polypeptide, the candidate receptors are each immunoprecipitated with protein A beads and the beads are washed. Potential ligand interaction is determined by western blot analysis of the immunoprecipitated complexes with antibody directed towards the epitope tag. An interaction is judged to occur if a band of the anticipated molecular weight of the epitope tagged protein is observed in the western blot analysis with a candidate receptor, but is not observed to occur with the other members of the panel of potential receptors.

Using these assays, the following receptor/ligand interactions have been herein identified:

- (1) PRO10272 binds to PRO5801.
 (2) PRO20110 binds to the human IL-17 receptor (Yao et al., *Cytokine* 9(11):794-800 (1997); also herein designated as PRO1) and to PRO20040.
 30 (3) PRO10096 binds to PRO20233.
 (4) PRO19670 binds to PRO1890.

- The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the

claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

WHAT IS CLAIMED IS:

1. A method for detecting the presence of tumor in a mammal, said method comprising comparing the level of expression of a PRO1865 polypeptide in (i) a test sample of tissue cells obtained from said mammal, and (ii) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower level of expression of said PRO1865 polypeptide in the test sample as compared to the control sample is indicative of the presence of tumor in said mammal, wherein the PRO1865 polypeptide is a polypeptide having at least 80% amino acid sequence identity to:

- (a) the amino acid sequence shown in SEQ ID NO:132;
- (b) the amino acid sequence shown in SEQ ID NO: 132, lacking its associated signal peptide;
- (c) the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132; or
- (d) the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132, lacking its associated signal peptide,

wherein the sequence identity is determined over the full length of the reference sequence and higher expression is indicative of the presence of esophageal tumor or melanoma and lower expression is indicative of the presence of stomach tumor.

- 2. The method of claim 1, wherein the sequence identity is at least 85%.
- 3. The method of claim 1, wherein the sequence identity is at least 90%.
- 4. The method of claim 1, wherein the sequence identity is at least 95%.
- 5. The method of claim 1, wherein the sequence identity is at least 99%.
- 6. The method of claim 1, wherein the polypeptide comprises:

- (a) the amino acid sequence shown in SEQ ID NO:132;
- (b) the amino acid sequence shown in SEQ ID NO: 132, lacking its associated signal peptide;
- (c) the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132; or
- (d) the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132, lacking its associated signal peptide.

7. A method for detecting the presence of tumor in a mammal, said method comprising (i) contacting an antibody which specifically binds to a PRO1865 polypeptide with a test sample of tissue cells obtained from the mammal and with a control sample of known normal tissue cells of the same cell type, and (ii) detecting the formation of a complex between said antibody and said polypeptide in the test sample and control sample, wherein a larger or smaller quantity of complexes

formed in the test sample as compared to the control sample is indicative of the presence of tumor in the mammal, wherein the PRO1865 polypeptide is as defined in claim 1 and wherein a larger quantity of complexes is indicative of the presence of esophageal tumor or melanoma and a smaller quantity of complexes is indicative of the presence of stomach tumor.

8. The method of any one of claims 1 to 7, wherein said PRO1865 polypeptide is detected using an antibody which specifically binds to a PRO1865 polypeptide as defined in claim 1.

9. The method of claim 8, wherein the antibody is detectably labeled.

10. The method of claim 8, wherein the antibody is an antibody fragment.

11. The method of claim 10, wherein the antibody fragment is selected from the group consisting of Fv, Fab, Fab' and F(ab')₂ fragments.

12. A method for detecting the presence of tumor in a mammal, said method comprising comparing the level of expression of a PRO1865-encoding nucleic acid in (i) a test sample of tissue cells obtained from said mammal, and (ii) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower level of expression of said PRO1865-encoding nucleic acid in the test sample as compared to the control sample is indicative of the presence of tumor in said mammal, wherein the PRO1865-encoding nucleic acid is a nucleic acid having at least 80% nucleic acid sequence identity to:

(a) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:132;

(b) the nucleotide sequence shown in SEQ ID NO:131;

(c) the full-length coding sequence of the nucleotide sequence shown in SEQ ID NO:131;

(b) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO: 132, lacking its associated signal peptide;

(c) a nucleotide sequence encoding the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132; or

(d) a nucleotide sequence encoding the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132, lacking its associated signal peptide,

wherein the sequence identity is determined over the full length of the reference sequence and higher expression is indicative of the presence of esophageal tumor or melanoma and lower expression is indicative of the presence of stomach tumor.

13. The method of claim 12, wherein the level of identity is at least 85%.

14. The method of claim 12, wherein the level of identity is at least 90%.

15. The method of claim 12, wherein the level of identity is at least 95%.

16. The method of claim 12, wherein the level of identity is at least 99%.
17. The method of claim 12, wherein the nucleic acid comprises:
 - (a) a nucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:132;
 - (b) the nucleotide sequence shown in SEQ ID NO: 131;
 - (c) a sequence encoding the full-length coding sequence of the nucleotide sequence shown in SEQ ID NO: 131;
 - (b) a sequence encoding the amino acid sequence shown in SEQ ID NO: 132, lacking its associated signal peptide;
 - (c) the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132; or
 - (d) a sequence encoding the amino acid sequence of an extracellular domain of the polypeptide shown in SEQ ID NO: 132, lacking its associated signal peptide.
18. The method of claim 12, wherein said PR01865-encoding nucleic acid is detected using a nucleic acid probe which specifically hybridizes to said PR01865-encoding nucleic acid.
19. The method of claim 18, wherein the nucleic acid probe is detectably labeled.
20. The method of claim 12, said PR01865-encoding nucleic acid is detected using a method selected from the group consisting of PCR, Northern analysis, Southern analysis, and *in situ* hybridization.

1/168

FIGURE 1

GGGGCTTCGGCGCCAGCGGCCAGCGCTAGTCGGTCTGGTAAGGATTTACAAAAGGTGCAGGTA
TGAGCAGGTCTGAAGACTAACATTTTGTGAAGTTGTAAAACAGAAAACCTGTTAGAAATGTGG
TGGTTTCAGCAAGGCCTCAGTTTCCTTCCTTCAGCCCTTGTAATTTGGACATCTGCTGCTTTC
ATATTTTCATACATTACTGCAGTAACACTCCACCATATAGACCCGGCTTTACCTTATATCAGT
GACACTGGTACAGTAGCTCCAGAAAAATGCTTATTTGGGGCAATGCTAAATATTGCGGCAGTT
TTATGCATTGCTACCATTTATGTTTCGTTATAAGCAAGTTCATGCTCTGAGTCCTGAAGAGAAC
GTTATCATCAAATTAAACAAGGCTGGCCTTGTACTTGGAATACTGAGTTGTTTAGGACTTTCT
ATTGTGGCAAACCTTCCAGAAAACAACCCTTTTTGCTGCACATGTAAGTGGAGCTGTGCTTACC
TTTGGTATGGGCTCATTATATATGTTTGTTCAGACCATCCTTTCCTACCAAATGCAGCCCAA
ATCCATGGCAAACAAGTCTTCTGGATCAGACTGTTGTTGGTTATCTGGTGTGGAGTAAGTGCA
CTTAGCATGCTGACTTGCTCATCAGTTTTGCACAGTGGCAATTTTGGGACTGATTTAGAACAG
AAACTCCATTGGAACCCCGAGGACAAAGGTTATGTGCTTCACATGATCACTACTGCAGCAGAA
TGGTCTATGTCATTTTCCTTCTTTGGTTTTTTCCTGACTTACATTCGTGATTTTCAGAAAATT
TCTTTACGGGTGGAAGCCAATTTACATGGATTAACCCTCTATGACACTGCACCTTGCCCTATT
AACAATGAACGAACACGGCTACTTTCCAGAGATATTTGATGAAAGGATAAAATATTTCTGTAA
TGATTATGATTCTCAGGGATTGGGGAAAGGTTACAGAAAGTTGCTTATTCTTCTCTGAAATTT
TCAACCACTTAATCAAGGCTGACAGTAACACTGATGAATGCTGATAATCAGGAAACATGAAAG
AAGCCATTTGATAGATTATTCTAAAGGATATCATCAAGAAGACTATTAAAAACACCTATGCCT
ATACTTTTTTATCTCAGAAAATAAAGTCAAAGACTATG

2/168

FIGURE 2

<subunit 1 of 1, 266 aa, 1 stop

<MW: 29766, pI: 8.39, NX(S/T): 0

MWWFQQGLSFLPSALVIWTSAAFIIFSITAVTLHHIDPALPYISDTGTVAPEKCLFGAMLNIA
AVLCIATIIYVRYKQVHALSPEENVIIKLNKAGLVLGILSCLGLSIVANFQKTTLFAAHVSGAV
LTFGMGSLYMFVQTIILSYQMOPKIHGKQVFWIRLLLVIWCGVSALSMLTCSSVLHSGNFGTDL
EQKLHWNPEDKGYVLHMITTAAEWSMSFSFFGFFLTYYIRDFQKISLRVEANLHGLTLYDTAPC
PINNERTRLSRDI

Important features:**Type II transmembrane domain:**

amino acids 13-33

Other Transmembrane domains:

amino acids 54-73, 94-113, 160-180, 122-141

N-myristoylation sites.

amino acids 57-63, 95-101, 99-105, 124-130, 183-189

3/168

FIGURE 3

CGGACGCGTGGGCGGACGCGTGGGGGAGAGCCGCGAGTCCCGGCTGCAGCACCTGGGAGAAGGC
AGACCGTGTGAGGGGGCCTGTGGCCCCAGCGTGCTGTGGCCTCGGGGAGTGGGAAGTGGAGGC
AGGAGCCTTCCTTACACTTCGCCATGAGTTTCCTCATCGACTCCAGCATCATGATTACCTCCC
AGATACTATTTTTTGGATTTGGGTGGCTTTTCTTCATGCGCCAATTGTTTAAAGACTATGAGA
TACGTCAGTATGTTGTACAGGTGATCTTCTCCGTGACGTTTGCATTTTCTTGCAACCATGTTTG
AGCTCATCATCTTTGAAATCTTAGGAGTATTGAATAGCAGCTCCCGTTATTTTCACTGGAAAA
TGAACCTGTGTGTAATTCTGCTGATCCTGGTTTTTCATGGTGCCTTTTTACATTGGCTATTTTA
TTGTGAGCAATATCCGACTACTGCATAACAACGACTGCTTTTTTCTGTCTCTTATGGCTGA
CCTTTATGTATTTCTTCTGGAACTAGGAGATCCCTTTCCCATTCCTCAGCCCCAAAACATGGGA
TCTTATCCATAGAACAGCTCATCAGCCGGGTGGTGTGATTGGAGTGACTCTCATGGCTCTTC
TTTCTGGATTTGGTGCTGTCAACTGCCCATACACTTACATGTCTTACTTCCTCAGGAATGTGA
CTGACACGGATATTCTAGCCCTGGAACGGCGACTGCTGCAAACCATGGATATGATCATAAGCA
AAAAGAAAAGGATGGCAATGGCACGGGAGAACAATGTTCCAGAAGGGGGGAAGTGCATAACAAAC
CATCAGGTTTCTGGGGAATGATAAAAAGTGTTACCACTTCAGCATCAGGAAGTGAAAATCTTA
CTCTTATTCAACAGGAAGTGGATGCTTTGGAAGAATTAAGCAGGCAGCTTTTTCTGGAAACAG
CTGATCTATATGCTACCAAGGAGAGAATAGAATACTCCAAAACCTTCAAGGGGAAATATTTTA
ATTTTCTTGGTTACTTTTTCTCTATTTACTGTGTTTGGAAAATTTTCATGGCTACCATCAATA
TTGTTTTTGTATCGAGTTGGGAAAACGGATCCTGTCACAAGAGGCATTGAGATCACTGTGAATT
ATCTGGGAATCCAATTTGATGTGAAGTTTTGGTCCCAACACATTTCTTCATTCTTGTGGAA
TAATCATCGTCACATCCATCAGAGGATTGCTGATCACTCTTACCAAGTTCTTTTATGCCATCT
CTAGCAGTAAGTCCTCCAATGTCATTGTCCTGCTATTAGCACAGATAATGGGCATGTACTTTG
TCTCCTCTGTGCTGCTGATCCGAATGAGTATGCCTTTAGAATAACCGCACCATAACTCACTGAAG
TCCTTGGAGAACTGCAGTTCAACTTCTATCACCGTTGGTTTGATGTGATCTTCCTGGTCAGCG
CTCTCTCTAGCATACTCTTCCTCTATTTGGCTCACAAACAGGCACCAGAGAAGCAAATGGCAC
CTTGAACTTAAGCCTACTACAGACTGTTAGAGGCCAGTGGTTTCAAAATTTAGATATAAGAGG
GGGGAAAAATGGAACCAGGGCCTGACATTTTATAAACAACAAAATGCTATGGTAGCATTTTTT
CACCTTCATAGCATACTCCTTCCCCGTCAGGTGATACTATGACCATGAGTAGCATCAGCCAGA
ACATGAGAGGGGAGAACTAACTCAAGACAATACTCAGCAGAGAGCATCCCGTGTGGATATGAGG
CTGGTGTAGAGGCGGAGAGGAGCCAAGAACTAAAGGTGAAAAATACACTGGAACCTCTGGGGC
AAGACATGTCTATGGTAGCTGAGCCAAACACGTAGGATTTCCGTTTTAAGGTTACATGGAAA
AGGTTATAGCTTTGCCTTGAGATTGACTCATTAATAATCAGAGACTGTAACAAAAA
AAAAAAAGGGCGGCCGCGACTCTAGAGTCGACCTGCAGAAGCTTGGCCGCCATGGCCCAACT
TGTTTATTGCAGCTTATAATG

4/168

FIGURE 4

MSFLIDSSIMITSQILFFGFGWLFFMRQLFKDYEIRQYVVQVIFSVTFASFCTMFELIIFEIL
GVLNSSSRYPFWKMNLCVILLILVFMVPFYIGYFIVSNIRLLHKQRLLEFSCLLWLTFMYFFWK
LGDPFPILSPKHGILSIEQLISRVGVIGVTLMALLSGFGAVNCPYTYMSYFLRNVTDTDILAL
ERRLLQTMDMIISKKKRMAMARRTMFQKGEVHNKPSGFWGMIKSVTTSASGSENLTLIQQEVD
ALEELSRQLFLETADLYATKERIEYSKTFKGKYFNFLGYFFSIYCVWKIFMATINIVFDRVGK
TDPVTRGIEITVNYLGIQFDVKFWSQHISFILVGIIIVTSIRGLLITLTKEFFYAISSSKSSNV
IVLLLAQIMGYFVSSVLLIRMSMPLEYRTIITEVLGELQFNFYHRWFDVIFLVSALSSILFL
YLAHKQAPEKQMAP

Important features:**Signal peptide:**

amino acids 1-23

Potential transmembrane domains:amino acids 37-55, 81-102, 150-168, 288-311, 338-356, 375-398,
425-444**N-glycosylation sites.**

amino acids 67-70, 180-183 and 243-246

Eukaryotic cobalamin-binding proteins

amino acids 151-160

5/168

FIGURE 5

AGCAGGGAAATCCGGATGTCTCGGTTATGAAGTGGAGCAGTGAGTGTGAGCCTCAACATAGTT
CCAGAACTCTCCATCCGGACTAGTTATTGAGCATCTGCCTCTCATATCACCAGTGGCCATCTG
AGGTGTTTCCCTGGCTCTGAAGGGGTAGGCACGATGGCCAGGTGCTTCAGCCTGGTGTGCTT
CTCACTTCCATCTGGACCACGAGGCTCCTGGTCCAAGGCTCTTTGCGTGCAGAAGAGCTTTCC
ATCCAGGTGTCATGCAGAATTATGGGGATCACCTTGTGAGCAAAAAGGCGAACCAGCAGCTG
AATTTACAGAAGCTAAGGAGGCCTGTAGGCTGCTGGGACTAAGTTTGGCCGGCAAGGACCAA
GTTGAAACAGCCTTGAAAGCTAGCTTTGAACTTGCAGCTATGGCTGGGTGGAGATGGATTC
GTGGTCATCTCTAGGATTAGCCCAAACCCCAAGTGTGGGAAAAATGGGGTGGGTGTCCTGATT
TGGAAGGTTCAGTGAGCCGACAGTTTGCAGCCTATTGTTACAACATCATCTGATACTTGGACT
AACTCGTGCATTCCAGAAATTATCACCACCAAAGATCCCATATTCAACACTCAAACCTGCAACA
CAAACAACAGAATTTATTGTCAGTGACAGTACCTACTCGGTGGCATCCCCTTACTCTACAATA
CCTGCCCCTACTACTACTCCTCCTGCTCCAGCTTCCACTTCTATTCCACGGAGAAAAAAATTG
ATTTGTGTCACAGAAGTTTTTATGGAACTAGCACCATGTCTACAGAACTGAACCATTTGTT
GAAATAAAGCAGCATTCAGAATGAAGCTGCTGGGTGGAGGTGTCCCCACGGCTCTGCTA
GTGCTTGCTCTCCTCTTCTTTGGTGCTGCAGCTGGTCTTGGATTTTGCTATGTCAAAGGTAT
GTGAAGGCCTTCCCTTTTACAAACAAGAATCAGCAGAAGGAAATGATCGAAACCAAAGTAGTA
AAGGAGGAGAAGGCCAATGATAGCAACCCTAATGAGGAATCAAAGAAAACCTGATAAAAACCCA
GAAGAGTCCAAGAGTCCAAGCAAACTACCGTGCGATGCCTGGAAGCTGAAGTTTAGATGAGA
CAGAAATGAGGAGACACACCTGAGGCTGGTTTTCTTTCATGCTCCTTACCCTGCCCCAGCTGGG
GAAATCAAAGGGCCAAAGAACCAAAGAAGAAAGTCCACCCTTGGTTCCTAACTGGAATCAGC
TCAGGACTGCCATTGGACTATGGAGTGCACCAAAGAGAATGCCCTTCTCCTTATTGTAACCCT
GTCTGGATCCTATCCTCCTACCTCCAAAGCTTCCCACGGCCTTTCTAGCCTGGCTATGTCCTA
ATAATATCCCCTGAGGAGAAAGGAGTTTTGCAAAGTGCAAGGACCTAAACATCTCATCAGTA
TCCAGTGGTAAAAAGGCCTCCTGGCTGTCTGAGGCTAGGTGGGTGAAAGCCAAGGAGTCACT
GAGACCAAGGCTTTCTCTACTGATTCCGCAGCTCAGACCCTTTCTTCAGCTCTGAAAGAGAAA
CACGTATCCCACCTGACATGTCCTTCTGAGCCCGGTAAGAGCAAAAGAATGGCAGAAAAGTTT
AGCCCCTGAAAGCCATGGAGATTCTCATAACTTGAGACCTAATCTCTGTAAAGCTAAAATAAA
GAAATAGAACAAGGCTGAGGATACGACAGTACACTGTCAGCAGGGACTGTAAACACAGACAGG
GTCAAAGTGTTTTCTCTGAACACATTGAGTTGGAATCACTGTTTAGAACACACACACTTACTT
TTTCTGGTCTCTACCACTGCTGATATTTTCTCTAGGAAATATACTTTTACAAGTAACAAAAAT
AAAAACTCTTATAAATTTCTATTTTTTATCTGAGTTACAGAAATGATTACTAAGGAAGATTACT
CAGTAATTTGTTTAAAAAGTAATAAAATTCAACAAACATTTGCTGAATAGCTACTATATGTCA
AGTGCTGTGCAAGGTATTACACTCTGTAATTGAATATTATTCCTCAAAAAATTGCACATAGTA
GAACGCTATCTGGGAAGCTATTTTTTTTCAGTTTTGATATTTCTAGCTTATCTACTTCCAAACT
AATTTTTATTTTTGCTGAGACTAATCTTATTCATTTTCTCTAATATGGCAACCATTATAACCT
TAATTTATTATTAAACATACCTAAGAAGTACATTGTTACCTCTATATACCAAAGCACATTTTAA
AAGTGCCATTAAACAAATGTATCACTAGCCCTCCTTTTTTCCAACAAGAAGGGACTGAGAGATGC
AGAAATATTTGTGACAAAAAATTAAAGCATTTAGAAAACCTT

6/168

FIGURE 6

MARCFSLVLLLSIWTTTRLLVQGSRLAEELSIQVSCRIMGITLVSKKANQQLNFTEAKEACRL
LGLSLAGKDQVETALKASFETCSYGWVGDFVVISRISPNPKCGKNGVGVLIWKVPVSRQFAA
YCYNSSDTWTNSCIPEIITTKDPIFNTQTATQTTEFIVSDSTYSVASPYSTIPAPTTTPPAPA
STSIPRRKKLICVTEVFMETSTMSTETEPFVENKAAFKNEAAGFGGVPTALLVLALLFFGAAA
GLGFCYVKRYVKAFPFNTKNQOKEMIETKVVKEEKANDSNPNEESKKTDKNPESKSPSKTTV
RCLEAEV

Signal sequence:

amino acids 1-16

Transmembrane domain:

amino acids 235-254

N-glycosylation site.

amino acids 53-57, 130-134, 289-293

Casein kinase II phosphorylation site.

amino acids 145-149, 214-218

Tyrosine kinase phosphorylation site.

amino acids 79-88

N-myristoylation site.

amino acids 23-29, 65-71, 234-240, 235-239, 249-255, 253-259

7/168

FIGURE 7

CGCCGCGCTCCCGCACCCGCGGCCCCGCCACCGCGCCGCTCCCGCATCTGCACCCGCGAGCCCG
GCGGCCTCCCGGCGGGAGCGAGCAGATCCAGTCCGGCCCCGAGCGCAACTCGGTCCAGTCGGG
GCGGCGGCTGCGGGCGCAGAGCGGAGATGCGAGCGGCTTGGGGCCACCCTGCTGTGCCTGCTGC
TGGCGGCGGCGGTCCCCACGGCCCCCGCGCCCGCTCCGACGGCGACCTCGGCTCCAGTCAAGC
CCGGCCCCGGCTCTCAGCTACCCGCAGGAGGAGGCCACCCTCAATGAGATGTTCCGCGAGGTTG
AGGAACTGATGGAGGACACGCAGCACAAATTGCGCAGCGCGGTGGAAGAGATGGAGGCAGAAG
AAGCTGCTGCTAAAGCATCATCAGAAGTGAACCTGGCAAACCTTACCTCCCAGCTATCACAATG
AGACCAACACAGACACGAAGGTTGGAAATAATACCATCCATGTGCACCGAGAAATTCACAAGA
TAACCAACAACCAGACTGGACAAATGGTCTTTTCAGAGACAGTTATCACATCTGTGGGAGACG
AAGAAGGCAGAAGGAGCCACGAGTGCATCATCGACGAGGACTGTGGGGCCAGCATGTACTGCC
AGTTTGCCAGCTTCCAGTACACCTGCCAGCCATGCCGGGGGCCAGAGGATGCTCTGCACCCGGG
ACAGTGAGTGCTGTGGAGACCAGCTGTGTGTCTGGGGTCACTGCACCAAAATGGCCACCAGGG
GCAGCAATGGGACCATCTGTGACAACCAGAGGGACTGCCAGCCGGGGCTGTGCTGTGCCTTCC
AGAGAGGCCTGCTGTTCCCTGTGTGCACACCCCTGCCCGTGGAGGGCGAGCTTTGCCATGACC
CCGCCAGCCGGCTTCTGGACCTCATCACCTGGGAGCTAGAGCCTGATGGAGCCTTGGACCGAT
GCCCTTGTGCCAGTGGCCTCCTCTGCCAGCCCCACAGCCACAGCCTGGTGTATGTGTGCAAGC
CGACCTTCGTGGGGAGCCGTGACCAAGATGGGGAGATCCTGCTGCCAGAGAGGTCCCCGATG
AGTATGAAGTTGGCAGCTTCATGGAGGAGGTGCGCCAGGAGCTGGAGGACCTGGAGAGGAGCC
TGACTIONTGAAGAGATGGCGCTGGGGGAGCCTGCGGCTGCCGCCGCTGCACTGCTGGGAGGGGAAG
AGATTTAGATCTGGACCAGGCTGTGGGTAGATGTGCAATAGAAATAGCTAATTTATTTCCCA
GGTGTGTGCTTTAGGCGTGGGCTGACCAGGCTTCTTCCTACATCTTCTTCCCAGTAAGTTTCC
CCTCTGGCTTGACAGCATGAGGTGTTGTGCATTTGTTTCAGCTCCCCCAGGCTGTTCTCCAGGC
TTCACAGTCTGGTGCTTGGGAGAGTCAGGCAGGGTTAAACTGCAGGAGCAGTTTGCCACCCCT
GTCCAGATTATTGGCTGCTTTGCCTCTACCAGTTGGCAGACAGCCGTTTGTCTACATGGCTT
TGATAATTGTTTGAGGGGAGGAGATGGAAACAATGTGGAGTCTCCCTCTGATTGGTTTTGGGG
AAATGTGGAGAAGAGTGCCCTGCTTTGCAAACATCAACCTGGCAAAAATGCAACAAATGAATT
TTCCACGCAGTTCTTTCCATGGGCATAGGTAAGCTGTGCCTTCAGCTGTTGCAGATGAAATGT
TCTGTTACCCCTGCATTACATGTGTTTATTCATCCAGCAGTGTTGCTCAGCTCCTACCTCTGT
GCCAGGGCAGCATTTTCATATCCAAGATCAATTCCCTCTCTCAGCACAGCCTGGGGAGGGGGT
CATTGTTCTCCTCGTCCATCAGGGATCTCAGAGGCTCAGAGACTGCAAGCTGCTTGCCCAAGT
CACACAGCTAGTGAAGACCAGAGCAGTTTCATCTGGTTGTGACTCTAAGCTCAGTGCTCTCTC
CACTACCCACACACCAGCCTTGGTGCCACCAAAAGTGCTCCCCAAAAGGAAGGAGAATGGGATT
TTTCTTGAGGCATGCACATCTGGAATTAAGGTCAAACCTAATTCTCACATCCCTCTAAAAGTAA
ACTACTGTTAGGAACAGCAGTGTTCTCACAGTGTGGGGCAGCCGTCTTCTAATGAAGACAAT
GATATTGACACTGTCCCTCTTTGGCAGTTGCATTAGTAACTTTGAAAGGTATATGACTGAGCG
TAGCATAACAGGTTAACCTGCAGAAACAGTACTTAGGTAATTGTAGGGCGAGGATTATAAATGA
AATTTGCAAAATCACTTAGCAGCAACTGAAGACAATTATCAACCACGTGGAGAAAATCAAACC
GAGCAGGGGCTGTGTGAAACATGGTTGTAATATGCGACTGCGAACACTGAACTCTACGCCACTC
CACAAATGATGTTTTTCAGGTGTCATGGACTGTTGCCACCATGTATTCATCCAGAGTTCTTAAA
GTTTAAAGTTGCACATGATTGTATAAGCATGCTTTCTTTGAGTTTTTAAATTATGTATAAACAT
AAGTTGCATTTAGAAATCAAGCATAAATCACTTCAACTGCAAAAAAAAAAAAAAAAAAAAAA
AAA

8/168

FIGURE 8

MQRLGATLLCLLLAAAVPTAPAPAPTATSAPVKPGPALSYPQEEATLNEMFREVEELMEDTQH
KLRSVEEMEAEAEAAKASSEVNLANLPPSYHNETNTDTKVGNNTIHVHREIHKITNNQTGQM
VFSETVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQPCRGQRM LCTR DSECCGDQL
CVWGHCTKMATRGSNGTICDNQRDCQ PGLCCA FQRGLLFPVCTPLPVEGELCHDPASRLLDLI
TWELEPDGALDRCPCASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILLPREVPDEYEVGSFME
EVRQELEDLERSLTEEMALGEPAAAAAALLGGEI

Signal sequence:

amino acids 1-19

N-glycosylation site.

amino acids 96-100, 106-110, 121-125, 204-208

Casein kinase II phosphorylation site.amino acids 46-50, 67-71, 98-102, 135-139, 206-210, 312-316,
327-331**N-myristoylation site.**

amino acids 202-208, 217-223

Amidation site.

amino acids 140-144

9/168

FIGURE 9

CGGACGCGTGGGCGGACGCGTGGGGGCTGTGAGAAAGTGCCAATAAATACATCATGCAACCCC
ACGGCCCACCTTGTGAACTCCTCGTGCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCC
AAAGGCCTAATCCAACGTTCTGTCTTCAATCTGCAAATCTATGGGGTCCTGGGGCTCTTCTGG
ACCCTTAACTGGGTACTGGCCCTGGGCCAATGCGTCCTCGCTGGAGCCTTTGCCTCCTTCTAC
TGGGCCTTCCACAAGCCCCAGGACATCCCTACCTTCCCCTTAATCTCTGCCTTCATCCGCACA
CTCCGTTACCACACTGGGTCATTGGCATTGAGGCCCTCATCCTGACCCTTGTGCAGATAGCC
CGGGTCATCTTGGAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCCTGTAGCCCGCTGC
ATCATGTGCTGTTTCAAGTGCTGCCTCTGGTGTCTGGAAAAATTTATCAAGTTCCTAAACCGC
AATGCATACATCATGATCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTC
ATGCTACTCATGCGAAACATTGTCAGGGTGGTCGTCCTGGACAAAGTCACAGACCTGCTGCTG
TTCTTTGGGAAGCTGCTGGTGGTCGGAGGCGTGGGGGTCCTGTCCTTCTTTTTTTTCTCCGGT
CGCATCCCGGGGCTGGGTAAAGACTTTAAGAGCCCCCACCTCAACTATTACTGGCTGCCCATC
ATGACCTCCATCCTGGGGGCTATGTCATCGCCAGCGGCTTCTTCAGCGTTTTTCGGCATGTGT
GTGGACACGCTCTTCCTCTGCTTCCTGGAAGACCTGGAGCGGAACAACGGCTCCCTGGACCGG
CCCTACTACATGTCCAAGAGCCTTCTAAAGATTCTGGGCAAGAAGAACGAGGCGCCCCCGGAC
ACAAGAAGAGGAAGAAGTGACAGCTCCGGCCCTGATCCAGGACTGCACCCCACCCCCACCGT
CCAGCCATCCAACCTCACTTCGCCTTACAGGTCTCCATTTTGTGGTAAAAAAGGTTTTAGGC
CAGGCGCCGTGGCTCACGCCTGTAATCCAACACTTTGAGAGGCTGAGGCGGGCGGATCACCTG
AGTCAGGAGTTCGAGACCAGCCTGGCCAACATGGTGAAACCTCCGTCTCTATTAAAAATACAA
AAATTAGCCGAGAGTGGTGGCATGCACCTGTCATCCCAGCTACTCGGGAGGCTGAGGCAGGAG
AATCGCTTGAACCCGGGAGGCAGAGGTTGCAGTGAGCCGAGATCGCGCCACTGCACTCCAACC
TGGGTGACAGACTCTGTCTCCAAAACAAAACAAACAAAAGATTTTATTAAAGATATTT
TGTTAACTC

10/168

FIGURE 10

RTRGRTRGGCEKVPINTSCNPTAHLVNSSCPGLMCVFQGYSSKGLIQRSVFNLQIYGVLGLFW
TLNWVLALGQCVLAGAFASFYWAFHKPQDIPTEPLISAFIRTLRYHTGSLAFGALILTLVQIA
RVILEYIDHKLRGVQNPVARCIMCCFKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAF
MLLMRNIVRVVLDKVTDLLLFFGKLLVVGGVGVLSFFFFSGRIPGLGKDFKSPHLNYYWLPI
MTSILGAYVIASGFFSVFGMCVDTLFLCFLEDLERNNGSLDRPYYSKSLKILGKKNEAPPD
NKKRKK

Important features:**Transmembrane domains:**

amino acids 57-80 (type II), 110-126, 215-231, 254-274

N-glycosylation sites.

amino acids 16-20, 27-31, 289-293

Hypothetical YBR002c family proteins.

amino acids 276-288

Ammonium transporters proteins.

amino acids 204-231

N-myristoylation sites.

amino acids 60-66, 78-84

Amidation site.

amino acids 306-310

11/168

FIGURE 11

GCCCCGCGCCCGGCGCCGGGCGCCCGAAGCCGGGAGCCACCGCCATGGGGGCCTGCCTGGGAG
CCTGCTCCCTGCTCAGCTGCGCGTCCTGCCTCTGCGGCTCTGCCCCCTGCATCCTGTGCAGCT
GCTGCCCCGCCAGCCGCAACTCCACCGTGAGCCGCCTCATCTTCACGTTCTTCCTCTTCCTGG
GGGTGCTGGTGTCCATCATTATGCTGAGCCCGGGCGTGGAGAGTCAGCTCTACAAGCTGCCCT
GGGTGTGTGAGGAGGGGGCCGGGATCCCCACCGTCCTGCAGGGCCACATCGACTGTGGCTCCC
TGCTTGGCTACCGCGCTGTCTACCGCATGTGCTTCGCCACGGCGGCCTTCTTCTTCTTCTTTT
TCACCCTGCTCATGCTCTGCGTGAGCAGCAGCCGGGACCCCCGGGCTGCCATCCAGAATGGGT
TTTGGTTCTTTAAGTTCCTGATCCTGGTGGGCCTCACCGTGGGTGCCTTCTACATCCCTGACG
GCTCCTTCACCAACATCTGGTTCTACTTCGGCGTCGTGGGCTCCTTCCTCTTCATCCTCATCC
AGCTGGTGCTGCTCATCGACTTTGCGCACTCCTGGAACCAGCGGTGGCTGGGCAAGGCCGAGG
AGTGCGATTCCCGTGCCCTGGTACGCAGGCCTCTTCTTCTTCACTCTCCTCTTCTACTTGCTGT
CGATCGCGGCCGTGGCGCTGATGTTTCATGTACTACACTGAGCCCAGCGGCTGCCACGAGGGCA
AGGTCTTCATCAGCCTCAACCTCACCTTCTGTGTCTGCGTGTCCATCGCTGCTGTCCTGCCCA
AGGTCCAGGACGCCCAGCCCAACTCGGGTCTGCTGCAGGCCTCGGTCATCACCTCTACACCA
TGTTTGTACCTGGTCAGCCCTATCCAGTATCCCTGAACAGAAATGCAACCCCCATTGCCAA
CCCAGCTGGGCAACGAGACAGTTGTGGCAGGCCCCGAGGGCTATGAGACCCAGTGGTGGGATG
CCCCGAGCATTGTGGGCCTCATCATCTTCCTCCTGTGCACCCTCTTCATCAGTCTGCGCTCCT
CAGACCACCGGCAGGTGAACAGCCTGATGCAGACCGAGGAGTGCCCACCTATGCTAGACGCCA
CACAGCAGCAGCAGCAGCAGGTGGCAGCCTGTGAGGGCCGGGCCTTTGACAACGAGCAGGACG
GCGTCACCTACAGCTACTCCTTCTTCCACTTCTGCCTGGTGCTGGCCTCACTGCACGTCATGA
TGACGCTCACCAACTGGTACAAGCCCGGTGAGACCCGGAAGATGATCAGCACGTGGACCGCCG
TGTGGGTGAAGATCTGTGCCAGCTGGGCAGGGCTGCTCCTCTACCTGTGGACCCTGGTAGCCC
CACTCCTCCTGCGCAACCGCGACTTCAGCTGAGGCAGCCTCACAGCCTGCCATCTGGTGCCTC
CTGCCACCTGGTGCCTCTCGGCTCGGTGACAGCCAACCTGCCCCCTCCCCACACCAATCAGCC
AGGCTGAGCCCCCACCCTGCCCCAGCTCCAGGACCTGCCCCTGAGCCGGGCCTTCTAGTCGT
AGTGCCTTCAGGGTCCGAGGAGCATCAGGCTCCTGCAGAGCCCCATCCCCCGCCACACCCAC
ACGGTGGAGCTGCCTCTTCCTTCCCCTCCTCCCTGTTGCCCATACTCAGCATCTCGGATGAAA
GGGCTCCCTTGTCCTCAGGCTCCACGGGAGCGGGGCTGCTGGAGAGAGCGGGGAACTCCCACC
ACAGTGGGGCATCCGGCACTGAAGCCCTGGTGTTCCTGGTCACGTCCCCCAGGGGACCCTGCC
CCCTTCCTGGACTTCGTGCCTTACTGAGTCTCTAAGACTTTTTCTAATAACAAGCCAGTGCG
TGTAATAAAAAA

12/168

FIGURE 12

MGACLGACSLSCASCLCGSAPCILCSCCPASRNSTVSRLI FTFFLFLGVLVSIIMLSPGVES
QLYKLPWVCEEAGAGIPTVLQGHIDCGSLLGYRAVYRMCFATAAFFFFFFFFTLLMLCVSSSRDPR
AAIQNGFWFFKFLILVGLTVGAFYIPDGSETNIWFYFGVVGSFLFILIQLVLLIDFAHSWNQR
WLGKAEECDSRAWYAGLFFFTLLFYLLSIAAVALMFMYYTEPSGCHEGKVFISLNLTFVCVVS
IAAVLPKVQDAQPNSGLLQASVITLYTMEVTWSALSSIPEQKCNPHLPTQLGNETVVAGPEGY
ETQWWDAPSIIVGLIIFLLCTLFISLRSSDHRQVNSLMQTEECPPMLDATQQQQQQVAACEGRA
FDNEQDGV TYSYSFFHFCLVLASLHVMMTLTNWYKPGETRKMISTWTAVVWKICASWAGLLLY
LWTLVAPLLLRNRDFS

Signal sequence:

amino acids 1-20

Transmembrane domains:

amino acids 40-58, 101-116, 134-150, 162-178, 206-223, 240-257,
272-283, 324-340, 391-406, 428-444

13/168

FIGURE 13

CGGGCCAGCCTGGGGCGGCCGGCCAGGAACCAACCCGTTAAGGTGTCTTCTCTTTAGGGATGGT
GAGGTTGGAAAAAGACTCCTGTAACCCTCCTCCAGGATGAACCACCTGCCAGAAGACATGGAG
AACGCTCTCACCGGGAGCCAGAGCTCCCATGCTTCTCTGCGCAATATCCATTCCATCAACCCC
ACACAACCTCATGGCCAGGATTGAGTCCTATGAAGGAAGGGAAAAGAAAGGCATATCTGATGTC
AGGAGGACTTTCTGTTTGTTCACCTTTGACCTCTTATTCGTAACATTACTGTGGATAATA
GAGTTAAATGTGAATGGAGGCATTGAGAACACATTAGAGAAGGAGGTGATGCAGTATGACTAC
TATTCTTCATATTTTGATATATTTCTTCTGGCAGTTTTTCGATTTAAAGTGTTAATACTTGCA
TATGCTGTGTGCAGACTGCGCCATTGGTGGGCAATAGCGTTGACAACGGCAGTGACCAGTGCC
TTTTTACTAGCAAAAGTGATCCTTTTGAAGCTTTTCTCTCAAGGGGCTTTTGGCTATGTGCTG
CCCATCATTTTCATTCATCCTTGCCTGGATTGAGACGTGGTTCCTGGATTTCAAAGTGTTACCT
CAAGAAGCAGAAGAAGAAAACAGACTCCTGATAGTTCAGGATGCTTCAGAGAGGGCAGCACTT
ATACCTGGTGGTCTTTCTGATGGTCAGTTTTATTCCCCTCCTGAATCCGAAGCAGGATCTGAA
GAAGCTGAAGAAAAACAGGACAGTGAGAAACCACTTTTAGAACTATGAGTACTACTTTTGTTA
AATGTGAAAAACCCTCACAGAAAGTCATCGAGGCAAAAAGAGGCAGGCAGTGGAGTCTCCCTG
TCGACAGTAAAGTTGAAATGGTGACGTCCACTGCTGGCTTTATTGAACAGCTAATAAAGATTT
ATTTATTGTAATACCTCACAAACGTTGTACCATATCCATGCACATTTAGTTGCCTGCCTGTGG
CTGGTAAGGTAATGTCATGATTCATCCTCTCTTCAGTGAGACTGAGCCTGATGTGTTAACAAA
TAGGTGAAGAAAGTCTTGCTGTATTCCTAATCAAAAGACTTAATATATTGAAGTAACACTT
TTTTAGTAAGCAAGATACCTTTTTATTTCATTCACAGAATGGAATTTTTTTTGTTCATGTCT
CAGATTTATTTTGTATTTCTTTTTTAACACTCTACATTTCCCTTGTTTTTTAACTCATGCACA
TGTGCTCTTTGTACAGTTTTTAAAAAGTGTAATAAAATCTGACATGTCAATGTGGCTAGTTTTA
TTTTTCTTGTTTTGCATTATGTGTATGGCCTGAAGTGTTGGACTTGCAAAAGGGGAAGAAAGG
AATTGCGAATACATGTAAAATGTCACCAGACATTTGTATTATTTTTTATCATGAAATCATGTTT
TTCTCTGATTGTTCTGAAATGTTCTAAATACTCTTATTTTGAATGCACAAAATGACTTAAACC
ATTCATATCATGTTTCCTTTGCGTTCAGCCAATTTCAATTAAAATGAACTAAATTAAAAA

14/168

FIGURE 14

MNHLPEDMENALTGSQSSHASLRNIHSINPTQLMARIESYEGREKKGISDVRRTFCLFVTFDL
LEVTLLWIIELNVNGGIENLEKEVMQYDYYSSYFDIFLLAVFRFKVLILAYAVCRLRHWWAI
ALTTAVTSAFLAKVILSKLFSQGAFGYVLPIISFILAWIETWFLDFKVLPOEAEEENRLLIV
QDASERAALIPGGLSDGQFYSPPESEAGSEEAEEKQDSEKPLLEL

Important features of the protein:

Signal peptide:

amino acids 1-20

Transmembrane domains:

amino acids 54-72, 100-118, 130-144, 146-166

N-myristoylation sites.

amino acids 14-20, 78-84, 79-85, 202-208, 217-223

15/168

FIGURE 15

ACTCGAACGCAGTTGCTTCGGGACCCAGGACCCCCTCGGGCCCGACCCGCCAGGAAAGACTGA
GGCCGCGGCCTGCCCCGCCCCGGCTCCCTGCGCCGCGCGCCCTCCCGGGACAGAAGATGTGCT
CCAGGGTCCCTCTGCTGCTGCCGCTGCTCCTGCTACTGGCCCTGGGGCCTGGGGTGCAGGGCT
GCCCATCCGGCTGCCAGTGCAGCCAGCCACAGACAGTCTTCTGCACTGCCCCGCCAGGGGACCA
CGGTGCCCCGAGACGTGCCACCCGACACGGTGGGGCTGTACGTCTTTGAGAACGGCATCACCA
TGCTCGACGCAGGCAGCTTTGCCGGCCTGCCGGGCCTGCAGCTCCTGGACCTGTCACAGAACC
AGATCGCCAGCCTGCCAGCGGGGTCTTCCAGCCACTCGCCAACCTCAGCAACCTGGACCTGA
CGGCCAACAGGCTGCATGAAATCACCAATGAGACCTTCCGTGGCCTGCGGGCGCCTCGAGCGCC
TCTACCTGGGCAAGAACCGCATCCGCCACATCCAGCCTGGTGCCTTCGACACGCTCGACCGCC
TCCTGGAGCTCAAGCTGCAGGACAACGAGCTGCGGGCACTGCCCCCGCTGCGCCTGCCCCGCC
TGCTGCTGCTGGACCTCAGCCACAACAGCCTCCTGGCCCTGGAGCCCGGCATCCTGGACACTG
CCAACGTGGAGGCGCTGCGGCTGGCTGGTCTGGGGCTGCAGCAGCTGGACGAGGGGCTCTTCA
GCCGCTTGCGCAACCTCCACGACCTGGATGTGTCCGACAACCAGCTGGAGCGAGTGCCACCTG
TGATCCGAGGCCTCCGGGGCCTGACGCGCCTGCGGCTGGCCGGCAACACCCGCATTGCCCAGC
TGCGGGCCCGAGGACCTGGCCGGCCTGGCTGCCCTGCAGGAGCTGGATGTGAGCAACCTAAGCC
TGCAGGCCCTGCCTGGCGACCTCTCGGGCCTCTTCCCCCGCCTGCGGCTGCTGGCAGCTGCCC
GCAACCCCTTCAACTGCGTGTGCCCCCTGAGCTGGTTTGGCCCCCTGGGTGCGCGAGAGCCACG
TCACACTGGCCAGCCCTGAGGAGACGCGCTGCCACTTCCCGCCCAAGAACGCTGGCCGGCTGC
TCCTGGAGCTTGACTACGCCGACTTTGGCTGCCCAGCCACCACCACAGCCACAGTGCCCA
CCACGAGGCCCGTGGTGCGGGAGCCACAGCCTTGTCTTCTAGCTTGGCTCCTACCTGGCTTA
GCCCCACAGCGCCGGCCACTGAGGCCCCAGCCCGCCCTCCACTGCCCCACCGACTGTAGGGC
CTGTCCCCCAGCCCCAGGACTGCCACCGTCCACCTGCCTCAATGGGGGCACATGCCACCTGG
GGACACGGCACCACCTGGCGTGCTTGTGCCCCGAAGGCTTCACGGGCCTGTACTGTGAGAGCC
AGATGGGGCAGGGGACACGGCCCAGCCCTACACCAGTCACGCCGAGGCCACCACGGTCCCTGA
CCCTGGGCATCGAGCCGGTGAGCCCCACCTCCCTGCGCGTGCGGGCTGCAGCGCTACCTCCAGG
GGAGCTCCGTGCAGCTCAGGAGCCTCCGTCTCACCTATCGCAACCTATCGGGCCCTGATAAGC
GGCTGGTGACGCTGCGACTGCCTGCCTCGCTCGCTGAGTACACGGTCACCCAGCTGCGGCCCCA
ACGCCACTTACTCCGTCTGTGTCATGCCTTTGGGGCCCGGGCGGGTGCCGGAGGGGCGAGGAGG
CCTGCGGGGAGGGCCATACACCCCCAGCCGTCCACTCCAACCACGCCCCAGTCACCCAGGCCC
GCGAGGGCAACCTGCCGCTCCTCATTGCGCCCCGCCCTGGCCGCGGTGCTCCTGGCCGCGCTGG
CTGCGGTGGGGGCAGCCTACTGTGTGCGGCGGGGGCGGGCCATGGCAGCAGCGGCTCAGGACA
AAGGGCAGGTGGGGCCAGGGGCTGGGCCCTGGAACCTGGAGGGAGTGAAGGTCCCCTTGGAGC
CAGGCCCGAAGGCAACAGAGGGCGGTGGAGAGGCCCTGCCAGCGGGTCTGAGTGTGAGGTGC
CACTCATGGGCTTCCCAGGGCCTGGCCTCCAGTCACCCCTCCACGCAAAGCCCTACATCTAAG
CCAGAGAGAGACAGGGCAGCTGGGGCCGGGCTCTCAGCCAGTGAGATGGCCAGCCCCCTCCTG
CTGCCACACCACGTAAGTTCTCAGTCCCAACCTCGGGGATGTGTGCAGACAGGGCTGTGTGAC
CACAGCTGGGCCCTGTTCCCTCTGGACCTCGGTCTCCTCATCTGTGAGATGCTGTGGCCCAGC
TGACGAGCCCTAACGTCCCCAGAACCGAGTGCCCTATGAGGACAGTGTCCGCCCTGCCCTCCGC
AACGTGCAGTCCCTGGGCACGGCGGGCCCTGCCATGTGCTGGTAACGCATGCCTGGGTCCCTGC
TGGGCTCTCCCACTCCAGGCGGACCCTGGGGGCCAGTGAAGGAAGCTCCCGGAAAGAGCAGAG
GGAGAGCGGGTAGGCGGCTGTGTGACTCTAGTCTTGGCCCCAGGAAGCGAAGGAACAAAAGAA
ACTGGAAAGGAAGATGCTTTAGGAACATGTTTTGCTTTTTTAAAATATATATATTTATAAGAG
ATCCTTTCCCATTTATTCTGGGAAGATGTTTTTCAAACCTCAGAGACAAGGACTTTGGTTTTTG
TAAGACAAACGATGATATGAAGGCCTTTTGTAAGAAAAAATAAAAGATGAAGTGTGAAA

16/168

FIGURE 16

MCSRVP L L L L L L L L L L A L G P G V Q G C P S G C Q C S Q P Q T V F C T A R Q G T T V P R D V P P D T V G L Y V F E N G
I T M L D A G S F A G L P G L Q L L D L S Q N Q I A S L P S G V F Q P L A N L S N L D L T A N R L H E I T N E T F R G L R R L
E R L Y L G K N R I R H I Q P G A F D T L D R L L E L K L Q D N E L R A L P P L R L P R L L L L D L S H N S L L A L E P G I L
D T A N V E A L R L A G L G L Q Q L D E G L F S R L R N L H D L D V S D N Q L E R V P P V I R G L R G L T R L R L A G N T R I
A Q L R P E D L A G L A A L Q E L D V S N L S L Q A L P G D L S G L F P R L R L L A A A R N P F N C V C P L S W F G P W V R E
S H V T L A S P E E T R C H F P P K N A G R L L L E L D Y A D F G C P A T T T T A T V P T T R P V V R E P T A L S S S L A P T
W L S P T A P A T E A P S P P S T A P P T V G P V P Q P Q D C P P S T C L N G G T C H L G T R H H L A C L C P E G F T G L Y C
E S Q M G Q G T R P S P T P V T P R P P R S L T L G I E P V S P T S L R V G L Q R Y L Q G S S V Q L R S L R L T Y R N L S G P
D K R L V T L R L P A S L A E Y T V T Q L R P N A T Y S V C V M P L G P G R V P E G E E A C G E A H T P P A V H S N H A P V T
Q A R E G N L P L L I A P A L A A V L L A A L A A V G A A Y C V R R G R A M A A A A Q D K G Q V G P G A G P L E L E G V K V P
L E P G P K A T E G G G E A L P S G S E C E V P L M G F P G P G L Q S P L H A K P Y I

Important features:**Signal peptide:**

amino acids 1-23

Transmembrane domain:

amino acids 579-599

EGF-like domain cysteine pattern signature.

amino acids 430-442

Leucine zipper pattern.

amino acids 197-219, 269-291

N-glycosylation sites.

amino acids 101-105, 117-121, 273-277, 500-504, 528-532

Tyrosine kinase phosphorylation sites.

amino acids 124-131, 337-345

N-myristoylation sites.amino acids 23-29, 27-33, 70-76, 142-148, 187-193, 348-354,
594-600, 640-646

17/168

FIGURE 17

GCAGCGGCGAGGCGGCGGTGGTGGCTGAGTCCGTGGTGGCAGAGGCGAAGGCGACAGCTCATG
CGGGTCCGGATAGGGCTGACGCTGCTGCTGTGTGCGGTGCTGCTGAGCTTGGCCTCGGCGTCC
TCGGATGAAGAAGGCAGCCAGGATGAATCCTTAGATTCCAAGACTACTTTGACATCAGATGAG
TCAGTAAAGGACCATACTACTGCAGGCAGAGTAGTTGCTGGTCAAATATTTCTTGATTTCAGAA
GAATCTGAATTAGAATCCTCTATTCAAGAAGAGGAAGACAGCCTCAAGAGCCAAGAGGGGGAA
AGTGTCACAGAAGATATCAGCTTTCTAGAGTCTCCAAATCCAGAAAACAAGGACTATGAAGAG
CCAAAGAAAGTACGGAAACCAGCTTTGACCGCCATTGAAGGCACAGCACATGGGGAGCCCTGC
CACTTCCCTTTTCTTTTCCTAGATAAGGAGTATGATGAATGTACATCAGATGGGAGGGGAAGAT
GGCAGACTGTGGTGTGCTACAACCTATGACTACAAAGCAGATGAAAAGTGGGGCTTTTGTGAA
ACTGAAGAAGAGGCTGCTAAGAGACGGCAGATGCAGGAAGCAGAAATGATGTATCAAACCTGGA
ATGAAAATCCTTAATGGAAGCAATAAGAAAAGCCAAAAAAGAGAAGCATATCGGTATCTCCAA
AAGGCAGCAAGCATGAACCATAACCAAGCCCTGGAGAGAGTGTCATATGCTCTTTTATTTGGT
GATTACTTGCCACAGAATATCCAGGCAGCGAGAGAGATGTTTGAGAAGCTGACTGAGGAAGGC
TCTCCCAAGGGACAGACTGCTCTTGGCTTTCTGTATGCCTCTGGACTTGGTGTTAATTCAAGT
CAGGCAAAGGCTCTTGTATATTATACATTTGGAGCTCTTGGGGGCAATCTAATAGCCCACATG
GTTTTGGTAAGTAGACTTTAGTGGAAGGCTAATAATATTAACATCAGAAGAATTTGTGGTTTA
TAGCGGCCACAACCTTTTTTCAGCTTTCATGATCCAGATTTGCTTGTATTAAGACCAAATATTCA
GTTGAACTTCCTTCAAATTCTTGTTAATGGATATAACACATGGAATCTACATGTAAATGAAAG
TTGGTGGAGTCCACAATTTTTCTTTAAATGATTAGTTTGGCTGATTGCCCCCTAAAAAGAGAG
ATCTGATAAATGGCTCTTTTTTAAATTTTCTCTGAGTTGGAATTGTCAGAATCATTTTTTACAT
TAGATTATCATAATTTTAAAAATTTTTCTTTAGTTTTTCAAATTTTGTAATGGTGGCTATA
GAAAAACAACATGAAATATTATACAATATTTTGCAACAATGCCCTAAGAATTGTTAAATTC
TGGAGTTATTTGTGCAGAATGACTCCAGAGAGCTCTACTTTCTGTTTTTTTACTTTTCATGATT
GGCTGTCTTCCCATTTATTCTGGTCATTTATTGCTAGTGACACTGTGCCTGCTTCCAGTAGTC
TCATTTTCCCTATTTTGCTAATTTGTTACTTTTTCTTTGCTAATTTGGAAGATTAACATTTT
TTAATAAAATTATGTCTAAGATTA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

18/168

FIGURE 18

MRVRIGLTLLLCVLLSLASASSDEEGSQDESLDSKTTLTSDSVKDHTTAGRVVAGQIFLDS
EELSELESSIQEEEDSLKSQEGESVTEDISFLESPNPENKDYEEPKKVRKPALTAIEGTAHGEP
CHFPFLFLDKEYDECTSDGREDGRLWCATTYDYKADEKWGFCETEEEEAAKRRQMQEAEEMMYQT
GMKILNGSNKKSQKREAYRYLQKAASMNHTKALERVSYALLFGDYLPQNIQAAREMFEEKLTEE
GSPKGQTALGFLYASGLGVNSSQAKALVYYTFGALGGNLI AHMVLVSRL

Important features:**Signal peptide:**

amino acids 1-21

N-glycosylation sites.

amino acids 195-199, 217-221, 272-276

Tyrosine kinase phosphorylation site.

amino acids 220-228

N-myristoylation sites.

amino acids 120-126, 253-259, 268-274, 270-274, 285-291, 289-295

Glycosaminoglycan attachment site.

amino acids 267-271

Microbodies C-terminal targeting signal.

amino acids 299-303

Type II fibronectin collagen-binding domain protein.

amino acids 127-169

Fructose-bisphosphate aldolase class-II protein.

amino acids 101-119

19/168

FIGURE 19

AATTCAGATTTTAAGCCCATTTCTGCAGTGGAATTTTCATGAACTAGCAAGAGGACACCATCTTC
TTGTATTATACAAGAAAGGAGTGTACCTATCACACACAGGGGGGAAAAATGCTCTTTTGGGTGC
TAGGCCTCCTAATCCTCTGTGGTTTTCTGTGGACTCGTAAAGGAAAACCTAAAGATTGAAGACA
TCACTGATAAGTACATTTTTATCACTGGATGTGACTCGGGCTTTGGAACTTGGCAGCCAGAA
CTTTTGATAAAAAGGGATTTTCATGTAATCGCTGCCTGTCTGACTGAATCAGGATCAACAGCTT
TAAAGGCAGAAACCTCAGAGAGACTTCGTACTGTGCTTCTGGATGTGACCGACCCAGAGAATG
TCAAGAGGACTGCCCAGTGGGTGAAGAACCAAGTTGGGGAGAAAGGTCTCTGGGGTCTGATCA
ATAATGCTGGTGTTCCCGGCGTGCTGGCTCCCACTGACTGGCTGACACTAGAGGACTACAGAG
AACCTATTGAAGTGAACCTGTTTGGACTCATCAGTGTGACACTAAATATGCTTCCTTTGGTCA
AGAAAGCTCAAGGGAGAGTTATTAATGTCTCCAGTGTTGGAGGTCGCCTTGCAATCGTTGGAG
GGGGCTATACTCCATCCAAATATGCAGTGGAAGGTTTCAATGACAGCTTAAGACGGGACATGA
AAGCTTTTGGTGTGCACGTCTCATGCATTGAACCAGGATTGTTCAAACAAACTTGGCAGATC
CAGTAAAGGTAATTGAAAAAAACTCGCCATTTGGGAGCAGCTGTCTCCAGACATCAAACAAC
AATATGGAGAAGGTTACATTGAAAAAAGTCTAGACAAACTGAAAGGCAATAAATCCTATGTGA
ACATGGACCTCTCTCCGGTGGTAGAGTGCATGGACCACGCTCTAACAAGTCTCTTCCCTAAGA
CTCATTATGCCGCTGGAAAAGATGCCAAAATTTTCTGGATACCTCTGTCTCACATGCCAGCAG
CTTTGCAAGACTTTTTATTGTTGAAACAGAAAGCAGAGCTGGCTAATCCCAAGGCAGTGTGAC
TCAGCTAACCACAAATGTCTCCTCCAGGCTATGAAATTGGCCGATTTCAAGAACACATCTCCT
TTTCAACCCCATTCCTTATCTGCTCCAACCTGGACTCATTTAGATCGTGCTTATTTGGATTGC
AAAAGGGAGTCCCACCATCGCTGGTGGTATCCCAGGGTCCCTGCTCAAGTTTTCTTTGAAAAG
GAGGGCTGGAATGGTACATCACATAGGCAAGTCCTGCCCTGTATTTAGGCTTTGCCTGCTTGG
TGTGATGTAAGGGAAATTGAAAGACTTGCCCATTCAAAATGATCTTTACCGTGGCCTGCCCA
TGCTTATGGTCCCCAGCATTTACAGTAACTTGTGAATGTTAAGTATCATCTCTTATCTAAATA
TTAAAAGATAAGTCAACCCAAAAA

20/168

FIGURE 20

MLFWVLGLLILCGFLWTRKGKLIKIEDITDKYIFITGCDSGFGNLAARTFDKKGFHVIAACLTE
SGSTALKAETSERLRTVLLDVTDPENVKRTAQWVKNQVGEKGLWGLINNAGVPGVLAPTDWLT
LEDYREPIEVNLFGLISVTNLMLPLVKKAQGRVINVSSVGGRLAIVGGGYTPSKYAVEGFNDS
LRRDMKAFGVHVSCIEPGLFKTNLADPVKVIEKKLAIWEQLSPDIKQQYGEgyIEKSLDKLKG
NKSyVNMDLSPVVECMDHALTSLEPKTHYAAGKDAKIFWIPLSHMPAALQDFLLKQKAELAN
PKAV

Important features of the protein:

Signal peptide:

amino acids 1-17

Transmembrane domain:

amino acids 136-152

N-glycosylation sites.

amino acids 161-163, 187-190 and 253-256

Glycosaminoglycan attachment site.

amino acids 39-42

N-myristoylation sites.

amino acids 36-41, 42-47, 108-113, 166-171, 198-203 and 207-212

21/168

FIGURE 21

CTGAGGCGGCGGTAGCATGGAGGGGGAGAGTACGTCGGCGGTGCTCTCGGGCTTTGTGCTCGG
CGCACTCGCTTTCCAGCACCTCAACACGGACTCGGACACGGAAGGTTTTCTTCTTGGGGAAGT
AAAAGGTGAAGCCAAGAACAGCATTACTGATTCCCAAATGGATGATGTTGAAGTTGTTTATAC
AATTGACATTCAGAAATATATTCCATGCTATCAGCTTTTTAGCTTTTATAATTCTTCAGGCGA
AGTAAATGAGCAAGCACTGAAGAAAATATTATCAAATGTCAAAAAGAATGTGGTAGGTTGGTA
CAAATTCCGTCGTCATTTCAGATCAGATCATGACGTTTAGAGAGAGGCTGCTTCACAAAACTT
GCAGGAGCATTTTTCAAACCAAGACCTTGTTTTTCTGCTATTAACACCAAGTATAATAACAGA
AAGCTGCTCTACTCATCGACTGGAACATTCCTTATATAAACCTCAAAAAGGACTTTTTTCACAG
GGTACCTTTAGTGGTTGCCAATCTGGGCATGTCTGAACAACCTGGGTTATAAACTGTATCAGG
TTCCTGTATGTCCACTGGTTTTAGCCGAGCAGTACAAACACACAGCTCTAAATTTTTTTGAAGA
AGATGGATCCTTAAAGGAGGTACATAAGATAAATGAAATGTATGCTTCATTACAAGAGGAATT
AAAGAGTATATGCAAAAAAGTGGAAGACAGTGAACAAGCAGTAGATAAACTAGTAAAGGATGT
AAACAGATTAAAACGAGAAATTGAGAAAAGGAGAGGAGCACAGATTCAGGCAGCAAGAGAGAA
GAACATCCAAAAAGACCCTCAGGAGAACATTTTTCTTTGTCAGGCATTACGGACCTTTTTTCC
AAATTCTGAATTTCTTCATTCATGTGTTATGTCTTTAAAAAATAGACATGTTTCTAAAAGTAG
CTGTAAC TACAACCACCATCTCGATGTAGTAGACAATCTGACCTTAATGGTAGAACACACTGA
CATTCCTGAAGCTAGTCCAGCTAGTACACCACAAATCATTAAAGCATAAAGCCTTAGACTTAGA
TGACAGATGGCAATTCAAGAGATCTCGGTTGTTAGATACACAAGACAAACGATCTAAAGCAAA
TACTGGTAGTAGTAACCAAGATAAAGCATCCAAAATGAGCAGCCCAGAAACAGATGAAGAAAT
TGAAAAGATGAAGGGTTTTGGTGAATATTCACGGTCTCCTACATTTTGATCCTTTTAACCTTA
CAAGGAGATTTTTTTTATTTGGCTGATGGGTAAAGCCAAACATTTCTATTGTTTTTACTATGTT
GAGCTACTTGCAGTAAGTTCATTTGTTTTTACTATGTTACCTGTTTGCAGTAATACACAGAT
AACTCTTAGTGCATTTACTTCACAAAGTACTTTTTCAAACATCAGATGCTTTTATTTCCAAAC
CTTTTTTTCACCTTTCACTAAGTTGTTGAGGGGAAGGCTTACACAGACACATTCTTTAGAATT
GGAAAAGTGAGACCAGGCACAGTGGCTCACACCTGTAATCCCAGCACTTAGGGAAGACAAGTC
AGGAGGATTGATTGAAGCTAGGAGTTAGAGACCAGCCTGGGCAACGTATTGAGACCATGTCTA
TTAAAAAATAAAATGGAAAAGCAAGAATAGCCTTATTTTCAAATATGGAAAGAAATTTATAT
GAAAATTTATCTGAGTCATTAAAATTCTCCTTAAGTGATACTTTTTTAGAAGTACATTATGGC
TAGAGTTGCCAGATAAAATGCTGGATATCATGCAATAAATTTGCAAAACATCATCTAAATTT
AAAAAAAAAAAAAAAAAAAAA

22/168

FIGURE 22

MEGESTSAVLSGFVLGALAFQHLNTDSDTEGFLGGEVKGEAKNSITDSQMDDVEVVYTIDIQK
YIPCYQLFSFYNSSGEVNEQALKKILSNVKKNVVGWYKFRRHSDQIMTFRERLLHKNLQEHFS
NQDLVFLLLTPSIITESCSTHRLEHSLYKPQKGLFHRVPLVVANLGMSEQLGYKTVSGSCMST
GFSRAVQTHSSKFFEEEDGSLKEVHKINEMYASLQEELKSICKKVEDSEQAVDKLVKDVNRLKR
EIEKRRGAQIQAAAREKNIQKDPQENIFLCQALRTFFPNSEFLHSCVMSLKNRHVSKSSCNYNH
HLDVVDNLTLMVEHTDIPEASPASTPQIIKHKALDLDDRWQFKRSRLLDTDKRSKANTGSSN
QDKASKMSSPETDEEIEKMKGFGGEYSRSPTF

Important features:**Signal peptide:**

amino acids 1-19

N-glycosylation sites.

amino acids 75-79, 322-326

N-myristoylation site.

amino acids 184-154

Growth factor and cytokines receptors family.

amino acids 134-150

23/168

FIGURE 23

GGCACAGCCGCGCGGGCGGAGGGCAGAGTCAGCCGAGCCGAGTCCAGCCGGACGAGCGGACCAGCGCAGGGCAGCC
CAAGCAGCGCGCAGCGAACGCCCCGCCGCCGCCACACCCTCTGCGGTCCCCGCGGCGCCTGCCACCCTTCCCTCC
TTCCCCGCGTCCCCGCTCGCCGGCCAGTCAGCTTGCCGGGTTCGCTGCCCCGCGAAACCCCGAGGTCAACAGCC
CGCGCCTCTGCTTCCCTGGGCGCGCGCCGCCCTCCACGCCCTCCTTCTCCCTGGCCCGGCGCCTGGCACCGGGG
ACCGTTGCCTGACGCGAGGCCCAGCTCTACTTTTCGCCCCGCGTCTCCTCCGCCTGCTCGCCTCTTCCACCAACT
CCAATCCTTCTCCCTCCAGCTCCACTCGCTAGTCCCCGACTCCGCCAGCCCTCGGCCCGCTGCCGTAGCGCCGC
TTCCCGTCCGGTCCCAAAGGTGGGAACGCGTCCGCCCCGGCCCGCACCATGGGCACGGTTCCGGCTTGCCCGCGCTT
CTCTGCACCCTGGCAGTGCTCAGCGCCGCGCTGCTGGCTGCCGAGCTCAAGTCGAAAAGTTGCTCGGAAGTGCGA
CGTCTTTACGTGTCCAAAGGCTTCAACAAGAACGATGCCCCCTCCACGAGATCAACGGTGATCATTTGAAGATC
TGTCCCCAGGGTTCTACCTGCTGCTCTCAAGAGATGGAGGAGAAGTACAGCCTGCAAAGTAAAGATGATTTCAA
AGTGTGGTCAGCGAACAGTGCAATCATTTGCAAGCTGTCTTTGCTTCACGTTACAAGAAGTTTGATGAATTCTTC
AAAGAACTACTTGAAAATGCAGAGAAATCCCTGAATGATATGTTTGTGAAGACATATGGCCATTTATACATGCAA
AATTCTGAGCTATTTAAAGATCTCTTCGTAGAGTTGAAACGTTACTACGTGGTGGGAAATGTGAACCTGGAAGAA
ATGCTAAATGACTTCTGGGCTCGCCTCCTGGAGCGGATGTTCCGCCTGGTGAACCTCCAGTACCACTTTACAGAT
GAGTATCTGGAATGTGTGAGCAAGTATACGGAGCAGCTGAAGCCCTTCGGAGATGTCCCTCGCAAATTGAAGCTC
CAGGTTACTCGTGCTTTTGTAGCAGCCCGTACTTTTCGCTCAAGGCTTAGCGGTTGCGGGAGATGTCGTGAGCAAG
GTCTCCGTGGTAAACCCACAGCCCAAGTGTACCCATGCCCTGTTGAAGATGATCTACTGCTCCCACTGCCGGGGT
CTCGTGACTGTGAAGCCATGTTACAACACTACTGCTCAAACATCATGAGAGGCTGTTTGGCCAACCAAGGGGATCTC
GATTTTGAATGGAACAATTTCATAGATGCTATGCTGATGGTGGCAGAGAGGCTAGAGGGTCCTTTCAACATTGAA
TCGGTCATGGATCCCATCGATGTGAAGATTTCTGATGCTATTATGAACATGCAGGATAATAGTGTTCAAGTGTCT
CAGAAGGTTTTCCAGGGATGTGGACCCCCCAAGCCCCCTCCAGCTGGACGAATTTCTCGTTCCATCTCTGAAAGT
GCCTTCAGTGCTCGCTTCAGACCACATCACCCCGAGGAACGCCCAACCACAGCAGCTGGCACTAGTTTGGACCGA
CTGGTTACTGATGTCAAGGAGAACTGAAACAGGCCAAGAAATTCTGGTCCTCCCTTCAGAGCAACGTTTGCAAC
GATGAGAGGATGGCTGCAGGAAACGGCAATGAGGATGACTGTTGGAATGGGAAAGGCAAAGCAGGTACCTGTTT
GCAGTGACAGGAAATGGATTAGCCAACCAGGGCAACAACCCAGAGGTCCAGGTTGACACCAGCAAACCAGACATA
CTGATCCTTCGTCAAATCATGGCTCTTCGAGTGATGACCAGCAAGATGAAGAATGCATACAATGGGAACGACGTG
GACTTCTTTGATATCAGTGATGAAAGTAGTGGAGAAGGAAGTGGAAGTGGCTGTGAGTATCAGCAGTGCCCTTCA
GAGTTTGACTACAATGCCACTGACCATGCTGGGAAGAGTGCCAATGAGAAAGCCGACAGTGCTGGTGTCCGTCCT
GGGGCACAGGCCTACCTCCTCACTGTCTTCTGCATCTTGTTCCCTGGTTATGCAGAGAGAGTGGAGATTAATTCTCA
AACTCTGAGAAAAAGTGTTTCATCAAAAAGTTAAAAGGCACCAGTTATCACTTTTCTACCATCCTAGTGACTTTGC
TTTTTAAATGAATGGACAACAATGTACAGTTTTTACTATGTGGCCACTGGTTTAAGAAGTGCTGACTTTGTTTTCT
TCATTTCAGTTTTGGGAGGAAAAGGGACTGTGCATTGAGTTGGTTCCTGCTCCCCCAAACCATGTTAAACGTGGCT
AACAGTGTAGGTACAGAACTATAGTTAGTTGTGCATTTGTGATTTTATCACTCTATTATTTGTTTGTATGTTTTT
TTCTCATTTTCGTTTGTGGGTTTTTTTTTCCAACTGTGATCTCGCCTTGTTTCTTACAAGCAAACCAGGGTCCCTT
CTTGGCACGTAACATGTACGTATTTCTGAAATATTAAATAGCTGTACAGAAGCAGGTTTTATTATCATGTTATC
TTATTAAAAGAAAAAGCCCAAAAAGC

24/168

FIGURE 24

MARFGLPALLCTLAVLSAALLAAELKSKSCSEVRRLYVSKGFNKNDAPLHEINGDHLKICPQG
STCCSQEMEEKYSLQSKDDFKSVVSEQCNHLQAVFASRYKKFDEFFKELLENAEKSLNDMFVK
TYGHLYMQNSELFKDLFVELKRYVVGNVNLEEMLNDFWARLLERMFRLVNSQYHFTDEYLEC
VSKYTEQLKPFQDVPRKCLKQVTRAFVAARTFAQGLAVAGDVVSKVSVVNPTAQCTHALLKMI
YCSHCRGLVTVKPCYNYCSNIMRGCLANQGDLD FEWNNFIDAMLMVAERLEGPFNIESVMDPI
DVKISDAIMNMQDNSVQVSQKVFQGC GPPKPLPAGRISR SISESAFSARFRPHHPEERPTTAA
GTSLDRLVTDVKEKLKQAKKFWSSLPSNVCNDERMAAGNGNEDDCWNGKGKSRYLFAVTGNGL
ANQGNNPEVQVDTSKPDILILRQIMALRVMTSKMKNAYNGNDVDFFDISDESSGEGSGSGCEY
QQCPSEFDYNATDHAGKSANEKADSAGVRPGAQAYLLTVFCILFLVMQREWR

Important features:**Signal peptide:**

amino acids 1-22

ATP/GTP-binding site motif A (P-loop).

amino acids 515-524

N-glycosylation site.

amino acids 514-518

Glycosaminoglycan attachment sites.

amino acids 494-498, 498-502

N-myristoylation sites.

amino acids 63-69, 224-230, 276-282, 438-444, 497-503, 531-537

Glypicans proteins.

amino acids 54-75, 105-157, 238-280, 309-346, 423-460, 468-506

25/168

FIGURE 25

CTCGCCCTCAAATGGGAACGCTGGCCTGGGACTAAAGCATAGACCACCAGGCTGAGTATCCTG
ACCTGAGTCATCCCCAGGGATCAGGAGCCTCCAGCAGGGGAACCTTCCATTATATTCTTCAAGC
AACTTACAGCTGCACCGACAGTTGCGATGAAAGTTCTAATCTCTTCCCTCCTCCTGTTGCTGC
CACTAATGCTGATGTCCATGGTCTCTAGCAGCCTGAATCCAGGGGTCGCCAGAGGCCACAGGG
ACCGAGGCCAGGCTTCTAGGAGATGGCTCCAGGAAGGCGGCCAAGAATGTGAGTGCAAAGATT
GGTTCCTGAGAGCCCCGAGAAGAAAATTCATGACAGTGTCTGGGCTGCCAAAGAAGCAGTGCC
CCTGTGATCATTTCAGGGCAATGTGAAGAAAACAAGACACCAAAGGCACCACAGAAAGCCAA
ACAAGCATTCCAGAGCCTGCCAGCAATTTCTCAAACAATGTCAGCTAAGAAGCTTTGCTCTGC
CTTTGTAGGAGCTCTGAGCGCCCACTCTTCCAATTAAACATTCTCAGCCAAGAAGACAGTGAG
CACACCTACCAGACACTCTTCTTCTCCACCTCACTCTCCCACTGTACCCACCCCTAAATCAT
TCCAGTGCTCTCAAAAAGCATGTTTTTCAAGATCATTTTGTTTGTTGCTCTCTCTAGTGTCTT
CTTCTCTCGTCAGTCTTAGCCTGTGCCCTCCCCTTACCCAGGCTTAGGCTTAATTACCTGAAA
GATTCCAGGAAACTGTAGCTTCCTAGCTAGTGTCAATTAACCTTAAATGCAATCAGGAAAGTA
GCAAACAGAAGTCAATAAATATTTTTTAAATGTCAAAAAAAAAAAAAAAAAAAAAA

26/168

FIGURE 26

MKVLISLLLLLPLMLMSMVSSSLNPGVARGHRDRGQASRRWLQEGGQECECKDWFLRAPRRK
FMTVSGLPKKQCPCDHFKGNVKKTRHQRHHRKPNKHSRACQQFLKQCQLRSFALPL

Important features:

Signal peptide:

amino acids 1-22

N-myristoylation sites.

amino acids 27-33, 46-52

27/168

FIGURE 27

GGACGCCAGCGCCTGCAGAGGCTGAGCAGGGAAAAAGCCAGTGCCCCAGCGGAAGCACAGCTC
AGAGCTGGTCTGCCATGGACATCCTGGTCCCCTCCTGCAGCTGCTGGTGCTGCTTCTTACCC
TGCCCCTGCACCTCATGGCTCTGCTGGGCTGCTGGCAGCCCCCTGTGCAAAGCTACTTCCCCT
ACCTGATGGCCGTGCTGACTCCCAAGAGCAACCGCAAGATGGAGAGCAAGAAACGGGAGCTCT
TCAGCCAGATAAAGGGGCTTACAGGAGCCTCCGGGAAAGTGGCCCTACTGGAGCTGGGCTGCG
GAACCGGAGCCAACCTTTCAGTTCTACCCACCGGGCTGCAGGGTCACCTGCCTAGACCCAAATC
CCCCTTTGAGAAGTTCCTGACAAAGAGCATGGCTGAGAACAGGCACCTCCAATATGAGCGGT
TTGTGGTGGCTCCTGGAGAGGACATGAGACAGCTGGCTGATGGCTCCATGGATGTGGTGGTCT
GCACTCTGGTGCTGTGCTCTGTGCAGAGCCCAAGGAAGGTCCTGCAGGAGGTCCGGAGAGTAC
TGAGACCGGGAGGTGTGCTCTTTTTCTGGGAGCATGTGGCAGAACCATATGGAAGCTGGGCCT
TCATGTGGCAGCAAGTTTTTCGAGCCCACCTGGAAACACATTGGGGATGGCTGCTGCCTCACCA
GAGAGACCTGGAAGGATCTTGAGAACGCCCAGTTCTCCGAAATCCAAATGGAACGACAGCCCC
CTCCCTTGAAGTGGCTACCTGTTGGGCCCCACATCATGGGAAAGGCTGTCAAACAATCTTTCC
CAAGCTCCAAGGCACTCATTTGCTCCTTCCCCAGCCTCCAATTAGAACAAGCCACCCACCAGC
CTATCTATCTTCCACTGAGAGGGACCTTAGCAGAATGAGAGAAGACATTCATGTACCACCTACT
AGTCCCTCTCTCCCCAACCTCTGCCAGGGCAATCTCTAACTTCAATCCCGCCTTCGACAGTGA
AAAAGCTCTACTTCTACGCTGACCCAGGGAGGAAACACTAGGACCCTGTTGTATCCTCAACTG
CAAGTTTCTGGACTAGTCTCCCAACGTTTGCCTCCCAATGTTGTCCCTTTCCTTCGTTCCCAT
GGTAAAGCTCCTCTCGCTTTCCTCCTGAGGCTACACCCATGCGTCTCTAGGAACTGGTCACAA
AAGTCATGGTGCCTGCATCCCTGCCAAGCCCCCCTGACCCTCTCTCCCCACTACCACCTTCTT
CCTGAGCTGGGGGCACCAGGGAGAATCAGAGATGCTGGGGATGCCAGAGCAAGACTCAAAGAG
GCAGAGGTTTTGTTCTCAAATATTTTTTAATAAATAGACGAAACCACG

28/168

FIGURE 28

MDILVPLLQLLVLLLTLPPLHLMALLGCWQPLCKSYFPYLMAVLTPKSNRKMESKKRELFSSQIK
GLTGASGKVALLELGCGTGANFQFYPPGCRVTCLDPNPHFEKFLTKSMAENRHLQYERFVVAP
GEDMRQLADGSMDVVVCTLVLCVQSPRKVLQEVRRVLRPGGVLFFWEHVAEPYGSWAFMWQQ
VFEPTWKHIGDGCCLTRETWKDLENAQFSEIQMERQPPPLKWLPVGPHIMGKAVKQSFSSKA
LICSEFPSLQLEQATHQPIYLPLRGT

Important features:**Signal peptide:**

amino acids 1-23

Leucine zipper pattern.

amino acids 10-32

N-myristoylation sites.

amino acids 64-70, 78-84, 80-86, 91-97, 201-207

29/168

FIGURE 29

CAATGTTTGCCTATCCACCTCCCCCAAGCCCCTTTACCTATGCTGCTGCTAACGCTGCTGCTG
CTGCTGCTGCTGCTTAAAGGCTCATGCTTGGAGTGGGGACTGGTCGGTGCCCAGAAAGTCTCT
TCTGCCACTGACGCCCCCATCAGGGATTGGGCCTTCTTTCCCCCTTCCTTTCTGTGTCTCCTG
CCTCATCGGCCTGCCATGACCTGCAGCCAAGCCCAGCCCCGTGGGGAAGGGGAGAAAGTGGGG
GATGGCTTAAGAAAGCTGGGAGATAGGGAACAGAAGAGGGTAGTGGGTGGGCTAGGGGGGGCTGC
CTTATTTAAAGTGGTTGTTTATGATTCTTATACTAATTTATACAAAGATATTAAGGCCCTGTT
CATTAAGAAATTGTTCCCTTCCCCTGTGTTCAATGTTTGTAAGATTGTTCTGTGTAAATATG
TCTTTATAATAAACAGTTAAAAGCTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

30/168

FIGURE 30

MLLLTLLLLLLLLLLKGSCLEWGLVGAQKVSSATDAPIRDWAFPPSFLCLLPHRPAMTCSQAQP
RGEGEKVG DG

Important features:

Signal peptide:

amino acids 1-15

Growth factor and cytokines receptors family:

amino acids 3-18

31/168

FIGURE 31

GTTTGAATTCCTTCAACTATACCCACAGTCCAAAAGCAGACTCACTGTGTCCCAGGCTACCAG
TTCCTCCAAGCAAGTCATTTCCCTTATTTAACCGATGTGTCCCTCAAACACCTGAGTGCTACT
CCCTATTTGCATCTGTTTTGATAAATGATGTTGACACCCTCCACCGAATTCTAAGTGGAATCA
TGTCGGGAAGAGATACAATCCTTGGCCTGTGTATCCTCGCATTAGCCTTGTCTTTGGCCATGA
TGTTTACCTTCAGATTCATCACCACCCTTCTGGTTCACATTTTCATTTCAATTGGTTATTTTGG
GATTGTTGTTTGTCTGCGGTGTTTTATGGTGGCTGTATTATGACTATAACCAACGACCTCAGCA
TAGAATTGGACACAGAAAGGGAAAATATGAAGTGCGTGCTGGGGTTTGCTATCGTATCCACAG
GCATCACGGCAGTGCTGCTCGTCTTGATTTTTGTTCTCAGAAAGAGAATAAAATTGACAGTTG
AGCTTTTCCAAATCACAAATAAAGCCATCAGCAGTGCTCCCTTCCTGCTGTTCCAGCCACTGT
GGACATTTGCCATCCTCATTTTCTTCTGGGTCCTCTGGGTGGCTGTGCTGCTGAGCCTGGGAA
CTGCAGGAGCTGCCCAGGTTATGGAAGGCGGCCAAGTGGAATATAAGCCCCTTTCGGGCATTC
GGTACATGTGGTCGTACCATTTAATTGGCCTCATCTGGACTAGTGAATTCATCCTTGCGTGCC
AGCAAATGACTATAGCTGGGGCAGTGGTTACTTGTTATTTCAACAGAAGTAAAAATGATCCTC
CTGATCATCCCATCCTTTTCGTCTCTCTCCATTCTCTTCTTCTACCATCAAGGAACCGTTGTGA
AAGGGTCATTTTAAATCTCTGTGGTGAGGATTCCGAGAATCATTGTCATGTACATGCAAAACG
CACTGAAAGAACAGCAGCATGGTGCATTGTCCAGGTACCTGTTCCGATGCTGCTACTGCTGTT
TCTGGTGTCTTGACAAATACCTGCTCCATCTCAACCAGAATGCATATACTACAACCTGCTATTA
ATGGGACAGATTTCTGTACATCAGCAAAAGATGCATTCAAATCTTGTCCAAGAACTCAAGTC
ACTTTACATCTATTAACCTGCTTTGGAGACTTCATAATTTTTCTAGGAAAGGTGTTAGTGGTGT
GTTTCACTGTTTTTGGAGGACTCATGGCTTTTAACTACAATCGGGCATTCCAGGTGTGGGCAG
TCCCTCTGTTATTGGTAGCTTTTTTTTGCCTACTTAGTAGCCCATAGTTTTTTTATCTGTGTTTG
AAACTGTGCTGGATGCACTTTTCCTGTGTTTTGCTGTTGATCTGGAAACAAATGATGGATCGT
CAGAAAAGCCCTACTTTATGGATCAAGAATTTCTGAGTTTCGTAAAAAGGAGCAACAAATTAA
ACAATGCAAGGGCACAGCAGGACAAGCACTCATTAAAGGAATGAGGAGGGGAACAGAACTCCAGG
CCATTGTGAGATTAGATACCCATTTAGGTATCTGTACCTGGAAAACATTTCTTCTAAGAGCCA
TTTACAGAATAGAAGATGAGACCACTAGAGAAAAGTTAGTGAATTTTTTTTTTAAAGACCTAA
TAAACCCTATTCTTCCTCAAAA

32/168

FIGURE 32

MSGRDTILGLCILALALSLAMMFTFRFITLLVHIFISLVILGLLEFVCGVLWWLYDYDNDLS
IELDTERENMKCVLGFAIVSTGITAVLLVLI FVLRKRIKLTVELFQITNKAISSAPFLLFQPL
WTFAILIFFWVLWVAVLLSLGTAGAAQVMEGGQVEYKPLSGIRYMWSYHLIGLIWTSEFILAC
QQMTIAGAVVTCYFNRSKNDPPDHPILSSLSILFFYHQGTVVKGSFLISVVRIPRIIVMYMQN
ALKEQQHGALSRYLFRCCYCCFWCLDKYLLHLNQNAYTTTAINGTDFCTSAKDAFKILSKNSS
HFTSINCFGDFIIFLGKVLVVCFTVFGGLMAFNYNRAFQVWAVPLLLVAFFAYLVAHSFLSVF
ETVLDALFLCFAVDLETNDGSSEKPYFMDQEFLSFVKRSNKLNNARAQQDKHSLRNEEGTELQ
AIVR

Important features:**Signal peptide:**

amino acids 1-20

Putative transmembrane domains:

amino acids 35-54, 75-97, 126-146, 185-204, 333-350, 352-371

N-glycosylation sites.

amino acids 204-208, 295-299, 313-317

N-myristoylation sites.

amino acids 147-153, 178-184, 196-202, 296-275, 342-348

[illegible]

34/168

FIGURE 34

MRTVVLTMKASVIEMFLVLLVTGVHSNKETAKKIKRPKFTVPQINCDVKAGKIIDPEFIVKCP
AGCQDPKYHVYGTDVYASYSSVCGAAVHSGVLDNSGGKILVRKVAGQSGYKGSYSNGVQSLSL
PRWRESFIVLESKPKKGVTPSALTYSSSKSPAAQAGETTKAYQRPPIPGTTAQPVTLMQLLA
VTVAVATPTTLPRPSPSAASTTSIPRPQSVGHRSEQEMDLWSTATYTSSQNRPRADPGIQRQDP
SGAAFQKPVGADVSLGLVPKEELSTQSLEPVSLGDPNCKIDLSFLIDGSTSIGKRRFRIQKQL
LADVAQALDIGPAGPLMGVVQYGDNPATHFNLKTHNTSRDLKTAIEKITQRGGLSNVGRAISF
VTKNFFSKANGNRSGAPNVVVVMVDGWPTDKVEEASRLARESGINIFFITIEGAAENEKQYVV
EPNFANKAVCRTNGFYSLHVQSWFGLHKTLOPLVKRVCDTDRACSKTCLNSADIGFVIDGSS
SVGTGNFRTVLQFVTNLTKEFEISDTDRIGAVQYTYEQRLFEFGFDKYSSKPDILNAIKRVGY
WSGGTSTGAAINFALEQLFKKSKPNKRKLMILITDGRSYDDVRI PAMAAHLKGVITYAIGVAW
AAQEELEVIATHPARDHSFFVDEFDNLHQYVPRIIQNICTEFNSQPRN

Important features:**Signal peptide:**

amino acids 1-26

Transmembrane domain:

amino acids 181-200

N-glycosylation sites.

amino acids 390-394, 520-524

N-myristoylation sites.amino acids 23-29, 93-99, 115-121, 262-268, 367-373, 389-395,
431-437, 466-472, 509-515, 570-576, 571-577, 575-581, 627-633**Amidation site.**

amino acids 304-308

35/168

FIGURE 35

CCGAGCACAGGAGATTGCCTGCGTTTAGGAGGTGGCTGCGTTGTGGGAAAAGCTATCAAGGAA
GAAATTGCCAAACCATGTCTTTTTTTCTGTTTTTCAGAGTAGTTCACAACAGATCTGAGTGTTT
TAATTAAGCATGGAATACAGAAAACAACAAAAAACTTAAGCTTTAATTTTCATCTGGAATTCCA
CAGTTTTCTTAGCTCCCTGGACCCGGTTGACCTGTTGGCTCTTCCCGCTGGCTGCTCTATCAC
GTGGTGCTCTCCGACTACTCACCCCGAGTGTAAGAACCTTCGGCTCGCGTGCTTCTGAGCTG
CTGTGGATGGCCTCGGCTCTCTGGACTGTCCTTCCGAGTAGGATGTCACTGAGATCCCTCAAA
TGAGCCTCCTGCTGCTGTCACCTGAGTTTCTTTGTGATGTGGTACCTCAGCCTTCCCCAC
TACAATGTGATAGAACGCGTGAACCTGGATGTACTTCTATGAGTATGAGCCGATTTACAGACAA
GACTTTCACCTCACACTTCGAGAGCATTCAAACCTGCTCTCATCAAAATCCATTTCTGGTCATT
CTGGTGACCTCCCACCCTTCAGATGTGAAAGCCAGGCAGGCCATTAGAGTTACTTGGGGTGAA
AAAAAGTCTTGGTGGGGATATGAGGTTCTTACATTTTTCTTATTAGGCCAAGAGGCTGAAAAG
GAAGACAAAATGTTGGCATTGTCCTTAGAGGATGAACACCTTCTTTATGGTGACATAATCCGA
CAAGATTTTTTAGACACATATAATAACCTGACCTTGAAAACCATTATGGCATTTCAGGTGGGTA
ACTGAGTTTTGCCCCAATGCCAAGTACGTAATGAAGACAGACACTGATGTTTTTCATCAATACT
GGCAATTTAGTGAAGTATCTTTTAAACCTAAACCACTCAGAGAAGTTTTTCACAGGTTATCCT
CTAATTGATAATTATTCCTATAGAGGATTTTACCAAAAAACCCATATTTCTTACCAGGAGTAT
CCTTTCAAGGTGTTCCCTCCATACTGCAGTGGGTTGGGTTATATAATGTCCAGAGATTTGGTG
CCAAGGATCTATGAAATGATGGGTCACGTAAAACCCATCAAGTTTGAAGATGTTTATGTCGGG
ATCTGTTTGAATTTATTAAAAGTGAACATTCATATTCCAGAAGACACAAATCTTTTCTTTCTA
TATAGAATCCATTTGGATGTCTGTCAACTGAGACGTGTGATTGCAGCCCATGGCTTTTCTTCC
AAGGAGATCATCACTTTTTTGGCAGGTCATGCTAAGGAACACCACATGCCATTATTAACTTCAC
ATTCTACAAAAGCCTAGAAGGACAGGATACCTTGTGGAAAGTGTTAAATAAAGTAGGTAAGT
TGGAATAATTCATGGGGAGGTCAGTGTGCTGGCTTACACTGAACTGAACTCATGAAAACCCA
GACTGGAGACTGGAGGGTTACACTTGTGATTTATTAGTCAGGCCCTTCAAAGATGATATGTGG
AGGAATTAAATATAAAGGAATTGGAGGTTTTTGCTAAAGAAATTAATAGGACCAAACAATTTG
GACATGTCATTCTGTAGACTAGAATTTCTTAAAAGGGTGTTACTGAGTTATAAGCTCACTAGG
CTGTAAAAACAAAACAATGTAGAGTTTTATTATTGAACAATGTAGTCACTTGAAGGTTTTGT
GTATATCTTATGTGGATTACCAATTTAAAAATATATGTAGTTCTGTGTCAAAAAACTTCTTCA
CTGAAGTTATACTGAACAAAATTTTACCTGTTTTTGGTCATTTATAAAGTACTTCAAGATGTT
GCAGTATTTACAGTTATTATTATTAAAATTACTTCAACTTTGTGTTTTTAAATGTTTTGAC
GATTTCAATACAAGATAAAAAGGATAGTGAATCATTCTTTACATGCAAACATTTTCCAGTTAC
TTAACTGATCAGTTTATTATTGATACATCACTCCATTAATGTAAAGTCATAGGTCATTATTGC
ATATCAGTAATCTCTTGGACTTTGTAAATATTTTACTGTGGTAATATAGAGAAGAATTAAAG
CAAGAAAATCTGAAAA

36/168

FIGURE 36

MASALWTVLPSRMSLRSLKWSLLLLLSLLSFFVMWYLSLPHYNVIERVNWMYFYEYEPPIYRQDF
HFTLREHSNCSHQNPFLVILVTSHPSDVKARQAIRVTWGEKKSWWGYEVLTFLLGQEAEKED
KMLALSLEDEHLLYGDIIRQDFLDTYNNLTTLKTIMAFRWVTEFCPNAKYVMKTDTDVFINTGN
LVKYLLNLNHSEKFFTGYPLIDNYSYRGFYQKTHISYQEYPFKVFPPYCSGLGYIMSRDLVPR
IYEMMGHVKPIKFEDVYVGICLNLLKVNIHIPEDTNLFFLYRIHLDVCQLRRVIAAHGFSKE
IITFWQVMLRNTTCHY

Important features:**Type II transmembrane domain:**

amino acids 20-39

N-glycosylation sites.

amino acids 72-76, 154-158, 198-202, 212-216, 326-330

Glycosaminoglycan attachment site.

amino acids 239-243

Ly-6 / u-PAR domain proteins.

amino acids 23-37

N-myristoylation site.

amino acids 271-277

[illegible]

38/168

FIGURE 38

MELGCWTQLGLTFLQLLLISSLPREYTVINEACPGAENIMCRECCEYDQIECVCPGKREVVG
 YTIPCCRNEENECDSCLIHFGCTIFENCKSCRNGSWGGLDDEFYVKGIFYCAECRAGWYGGDCM
 RCGQVLRAPKGQILLESYPLNAHCEWTIHAKPGFVIQLRFVMLSLEFDYMCQYDYVEVRDGDN
 RDGQIIKRVCGNERPAPIQSIGSSLHVLHSDGSKNFDGFGHAIYEEITACSSSPCFHDGTCVL
 DKAGSYKCACLAGYTGQRCENLLEERNCSDPGGPVNGYQKITGGPGLINGRHAKIGTVVSFFC
 NNSYVLSGNEKRTCQQNGEWSGKQPICIKACREPKISDLVRRRVLPMPVQSRETPLHQLYSAA
 FSKQKLQSAPTKKPALPFGDLPMGYQHLHTQLQYECISPFYRRLGSSRRTCLRTGKWSGRAPS
 CIPICGKIENITAPKTQGLRWPWQAAIYRRTSGVHDGSLHKGAWFLVCSGALVNERTVVVAAH
 CVTDLGKVTMIKTADLKVVVLGKFYRDDDRDEKTIQSLQISAILHPNYDPILLDADIAILKLL
 DKARISTRVQPICLAASRDLSTSFQESHITVAGWNVLADVRSFGKNDTLRSGVVSVDLLC
 EEQHEDHGIPVSVTDNMFCASWEPTAPSDICTAETGGIAAVSFPGRASPEPRWHLMGLVSWSY
 DKTCSHRLSTAFTKVLFPKDWIERNMK

Important features of the protein:**Signal peptide:**

amino acids 1-23

EGF-like domain cysteine pattern signature.

amino acids 260-272

N-glycosylation sites.

amino acids 96-100, 279-283, 316-320, 451-455, 614-618

N-myristoylation sites.amino acids 35-41, 97-103, 256-262, 284-290, 298-304, 308-314,
474-480, 491-497, 638-644, 666-672**Amidation site.**

amino acids 56-60

Serine proteases, trypsin family.

amino acids 489-506

CUB domain proteins profile.

amino acids 150-167

39/168

FIGURE 39

GGTTCCTACATCCTCTCATCTGAGAATCAGAGAGCATAATCTTCTTACGGGCCCCTGATTTATTAACGTGGCTTA
ATCTGAAGGTTCTCAGTCAAATTCTTTGTGATCTACTGATTGTGGGGGCATGGCAAGGTTTGCTTAAAGGAGCTT
GGCTGGTTTGGGGCCCTTGTAGCTGACAGAAGGTGGCCAGGGAGAATGCAGCACACTGCTCGGAGAATGAAGGCGC
TTCTGTTGCTGGTCTTGCCTTGGCTCAGTCCTGCTAACTACATTGACAATGTGGGCAACCTGCACTTCCTGTATT
CAGAACTCTGTAAAGGTGCCTCCCACTACGGCCTGACCAAAGATAGGAAGAGGGCGCTCACAAGATGGCTGTCCAG
ACGGCTGTGCGAGCCTCACAGCCACGGCTCCCTCCCCAGAGGTTTCTGCAGCTGCCACCATCTCCTTAATGACAG
ACGAGCCTGGCCTAGACAACCCTGCCTACGTGTCCTCGGCAGAGGACGGGCAGCCAGCAATCAGCCCAGTGGACT
CTGGCCGGGAGCAACCGAACTAGGGCACGGCCCTTTGAGAGATCCACTATTAGAAGCAGATCATTTAAAAAATAA
ATCGAGCTTTGAGTGTTCTTCGAAGGACAAAGAGCGGGAGTGCAAGTTGCCAACCATGCCGACCAGGGCAGGGAAA
ATTCTGAAAACACCACTGCCCCCTGAAGTCTTTCCAAGGTTGTACCACCTGATTCCAGATGGTGAAATTACCAGCA
TCAAGATCAATCGAGTAGATCCCAGTGAAAGCCTCTCTATTAGGCTGGTGGGAGGTAGCGAAACCCCACTGGTCC
ATATCATTATCCAACACATTTATCGTGATGGGGTGATCGCCAGAGACGGCCGGCTACTGCCAGGAGACATCATTC
TAAAGGTCAACGGGATGGACATCAGCAATGTCCCTCACAACCTACGCTGTGCGTCTCCTGCGGCAGCCCTGCCAGG
TGCTGTGGCTGACTGTGATGCGTGAACAGAAGTTCGCGCAGCAGGAACAATGGACAGGCCCCGGATGCCTACAGAC
CCCGAGATGACAGCTTTCATGTGATTCTCAACAAAAGTAGCCCCGAGGAGCAGCTTGGAATAAACTGGTGCGCA
AGGTGGATGAGCCTGGGGTTTTTCATCTTCAATGTGCTGGATGGCGGTGTGGCATATCGACATGGTCAGCTTGAGG
AGAATGACCGTGTTAGCCATCAATGGACATGATCTTCGATATGGCAGCCAGAAAGTGCGGCTCATCTGATTC
AGGCCAGTGAAAGACGTGTTACCTCGTCGTGTCCCGCCAGGTTCCGGCAGCGGAGCCCTGACATCTTTCAGGAAG
CCGGCTGGAACAGCAATGGCAGCTGGTCCCCAGGGCCAGGGGAGAGGAGCAACACTCCCAAGCCCCTCCATCCTA
CAATTACTTGTGATGAGAAGGTGGTAAATATCCAAAAGACCCCGGTGAATCTCTCGGCATGACCGTCGCAGGGG
GAGCATCACATAGAGAATGGGATTTGCCTATCTATGTGATCAGTGTTGAGCCCGGAGGAGTCATAAGCAGAGATG
GAAGAATAAAACAGGTGACATTTTGTGTAATGTGGATGGGGTCGAACTGACAGAGGTGAGCCGGAGTGAGGCAG
TGGCATTATTGAAAAGAACATCATCCTCGATAGTACTCAAAGCTTTGGAAGTCAAAGAGTATGAGCCCCAGGAAG
ACTGCAGCAGCCAGCAGCCCTGGACTCCAACCACAACATGGCCCCACCCAGTGACTGGTCCCCATCCTGGGTCA
TGTGGCTGGAATTACCACGGTGCTTGTATAACTGTAAAGATATTGTATTACGAAGAAACACAGCTGGAAGTCTGG
GCTTCTGCATTGTAGGAGGTTATGAAGAATACAATGGAAACAAACCTTTTTTTCATCAAATCCATTGTTGAAGGAA
CACCAGCATACAATGATGGAAGAATTAGATGTGGTGATATTCTTCTTGCTGTCAATGGTAGAAGTACATCAGGAA
TGATACATGCTTGCTTGGCAAGACTGCTGAAAGAACTTAAAGGAAGAATTACTCTAACTATTGTTTCTTGGCCTG
GCACTTTTTTATAGAATCAATGATGGGTGAGAGGAAAACAGAAAAATCACAAATAGGCTAAGAAGTTGAAACACT
ATATTTATCTTGTCAGTTTTTATATTTAAAGAAAGAATACATTGTAAAAATGTCAGGAAAAGTATGATCATCTAA
TGAAAGCCAGTTACACCTCAGAAAATATGATTCCAAAAAATTAAACTACTAGTTTTTTTTTTCAGTGTGGAGGAT
TTCTCATTACTCTACAACATTGTTTATATTTTTTCTATTCAATAAAAAGCCCTAAAACAATAAATGATTGATT
TGTATACCCCACTGAATTCAAGCTGATTTAAATTTAAATTTGGTATATGCTGAAGTCTGCCAAGGGTACATTAT
GGCCATTTTTTAATTTACAGCTAAAATATTTTTTAAATGCATTGCTGAGAAACGTTGCTTTCATCAAACAAGAAT
AAATATTTTTTCAGAAGTTAAA

40/168

FIGURE 40

MKALLLLVLPWLSPANYIDNVGNLHFLYSELCKGASHYGLTKDRKRRSQDGCPDGCASLTATA
PSPEVSAAATISLMTDEPGLDNPAYVSSAEDGQPAISPVDSGRSNRTRARPFERSTIRSRSEK
KINRALSVLRRTKSGSAVANHADQGRESENTTAPEVFPRLYHLIPDGEITSIKINRVDPSES
LSIRLVGGSETPLVHIIIQHIYRDGVIARDGRLLPGDIILKVNGMDISNVPHNYAVRLLRQPC
QVLWLTVMREQKFRSRNNGQAPDAYRPRDDSEFHVILNKSSPEEQLGIKLVRKVDEPGVFIENV
LDGGVAYRHGQLEENDRVLAINGHDLRYGSPESA AHLIQASERRVHLVVSQRQVRQSPDIFQE
AGWNSNGSWSPGPGERSNTPKPLHPTITCHEKVVNIQKDPGESLGMTVAGGASHREWDLPIYV
ISVEPGGVISRDGRIKTGDILLNVDGVELTEVSRSEAVALLKRTSSSIVLKALEVKEYEPQED
CSSPAALDSNHNMAPPSDWSPSWVMWLELPRCLYNCKDIVLRRNTAGSLGFCIVGGYEEYNGN
KPFIFKSIVEGTPAYNDGRIRCGDILLAVNGRSTSGMIHACLARLLKELKGRITLTIVSWPGTFL

Important features:**Signal peptide:**

amino acids 1-15

N-glycosylation sites.

amino acids 108-112, 157-161, 289-293, 384-388

Tyrosine kinase phosphorylation sites.

amino acids 433-441, 492-500

N-myristoylation sites.amino acids 51-57, 141-147, 233-239, 344-350, 423-429, 447-453,
467-473, 603-609

41/168

FIGURE 41

ACCAGGCATTGTATCTTCAGTTGTCATCAAGTTCGCAATCAGATTGGAAAAGCTCAACTTGAA
GCTTTCTTGCCTGCAGTGAAGCAGAGAGATAGATATTATTCACGTAATAAAAAACATGGGCTT
CAACCTGACTTTCCACCTTTCCTACAAATTCCGATTACTGTTGCTGTTGACTTTGTGCCTGAC
AGTGGTTGGGTGGGCCACCAGTAACTACTTCGTGGGTGCCATTCAAGAGATTCCCTAAAGCAAA
GGAGTTCATGGCTAATTTCCATAAGACCCTCATTTTGGGGAAGGGAAAAACTCTGACTAATGA
AGCATCCACGAAGAAGGTAGAACTTGACAACCTGTCCTTCTGTGTCTCCTTACCTCAGAGGCCA
GAGCAAGCTCATTTTCAAACCAGATCTCACTTTGGAAGAGGTACAGGCAGAAAATCCCAAAGT
GTCCAGAGGCCGGTATCGCCCTCAGGAATGTAAAGCTTTACAGAGGGTCGCCATCCTCGTTCC
CCACCGGAACAGAGAGAAACACCTGATGTACCTGCTGGAACATCTGCATCCCTTCCTGCAGAG
GCAGCAGCTGGATTATGGCATCTACGTCATCCACCAGGCTGAAGGTAAAAAGTTTAATCGAGC
CAAACCTCTTGAATGTGGGCTATCTAGAAGCCCTCAAGGAAGAAAATTGGGACTGCTTTATATT
CCACGATGTGGACCTGGTACCCGAGAATGACTTTAACCTTTACAAGTGTGAGGAGCATCCCAA
GCATCTGGTGGTTGGCAGGAACAGCACTGGGTACAGGTTACGTTACAGTGGATATTTTGGGGG
TGTTACTGCCCTAAGCAGAGAGCAGTTTTTCAAGGTGAATGGATTCTCTAACAACCTACTGGGG
ATGGGGAGGCGAAGACGATGACCTCAGACTCAGGGTTGAGCTCCAAAGAATGAAAATTTCCCG
GCCCCTGCCTGAAGTGGGTAAATATACAATGGTCTTCCACACTAGAGACAAAGGCAATGAGGT
GAACGCAGAACGGATGAAGCTCTTACACCAAGTGTCACGAGTCTGGAGAACAGATGGGTTGAG
TAGTTGTTCTTATAAATTAGTATCTGTGGAACACAATCCTTTATATATCAACATCACAGTGGA
TTTCTGGTTTGGTGCATGACCCTGGATCTTTTGGTGATGTTTGGGAAGAACTGATTCTTTGTTT
GCAATAATTTTGGCCTAGAGACTTCAAATAGTAGCACACATTAAGAACCTGTTACAGCTCATT
GTTGAGCTGAATTTTTCCTTTTTGTATTTTCTTAGCAGAGCTCCTGGTGATGTAGAGTATAAA
ACAGTTGTAACAAGACAGCTTTCTTAGTCATTTTGATCATGAGGGTTAAATATTGTAATATGG
ATACTTGAAGGACTTTATATAAAAGGATGACTCAAAGGATAAAATGAACGCTATTTGAGGACT
CTGGTTGAAGGAGATTTATTTAAATTTGAAGTAATATATTATGGGATAAAAGGCCACAGGAAA
TAAGACTGCTGAATGTCTGAGAGAACCAGAGTTGTTCTCGTCCAAGGTAGAAAGGTACGAAGA
TACAATACTGTTATTCATTTATCCTGTACAATCATCTGTGAAGTGGTGGTGTCAGGTGAGAAG
GCGTCCACAAAAGAGGGGAGAAAAGGCGACGAATCAGGACACAGTGAACCTTGGGAATGAAGAG
GTAGCAGGAGGGTGGAGTGTCGGCTGCAAAGGCAGCAGTAGCTGAGCTGGTTGCAGGTGCTGA
TAGCCTTCAGGGGAGGACCTGCCCAGGTATGCCTTCCAGTGATGCCCACCAGAGAATACATTC
TCTATTAGTTTTTTAAAGAGTTTTTGTAAAATGATTTTGTACAAGTAGGATATGAATTAGCAGT
TTACAAGTTTACATATTAATAATAATAATATGTCTATCAAATACCTCTGTAGTAAAATGTG
AAAAAGCAAAA

42/168

FIGURE 42

MGFNLT FHLSYKFRL LLLTLCLTVVGWATSNYFVGAIQEIPKAKEFMANFHKTLLILGKGKTL
TNEASTKKVELDNCPSVSPYLRGQSKLIFKPDLTLEEVQAENPKVSRGRYRPQECKALQRVAI
LVPHRNREKHLMYLLEHLHPFLQRQQLDYGIYVIHQAEKGKFNRAKLLNVGYLEALKEENWDC
FIFHDVDLVPENDFNLYKCEEHPKHLVGRNSTGYRLRYSGYFGGVTALSREQFFKVNGFSNN
YWGWGGEDDLRLRVELQRMKISRPLPEVGKYTMVFHTRDKGNEVNAERMKLLHQVSRVWRTD
GLSSCSYKLVSVEHNPLYINITVDFWFGA

Important features:

Signal peptide:

amino acids 1-27

N-glycosylation sites.

amino acids 4-8, 220-224, 335-339

Xylose isomerase proteins.

amino acids 191-202

44/168

FIGURE 44

MALSSQIWAACLLLLLLLLASLTSGSVFPQQTGQLAELQPQDRAGARASWMPMFQRRRRRDTHF
PICIFCCGCCCHRSKCGMCCKT

Important features:

Signal peptide:

amino acids 1-24

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 58-59

N-myristoylation site.

amino acids 44-50

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 1-12

45/168

FIGURE 45

GTGGCTTCATTTTCAGTGGCTGACTTCCAGAGAGCAATATGGCTGGTTCCCCAACATGCCTCAC
CCTCATCTATATCCTTTGGCAGCTCACAGGGTCAGCAGCCTCTGGACCCGTGAAAGAGCTGGT
CGGTTCCGTTGGTGGGGCCGTGACTTTCCCCCTGAAGTCCAAAGTAAAGCAAGTTGACTCTAT
TGTCTGGACCTTCAACACAACCCCTCTTGTCAACATACAGCCAGAAGGGGGCACTATCATAGT
GACCCAAAATCGTAATAGGGAGAGAGTAGACTTCCCAGATGGAGGCTACTCCCTGAAGCTCAG
CAAACCTGAAGAAGAATGACTCAGGGATCTACTATGTGGGGATATACAGCTCATCACTCCAGCA
GCCCTCCACCCAGGAGTACGTGCTGCATGTCTACGAGCACCTGTCAAAGCCTAAAGTCACCAT
GGGTCTGCAGAGCAATAAGAATGGCACCTGTGTGACCAATCTGACATGCTGCATGGAACATGG
GGAAGAGGATGTGATTTATACCTGGAAGGCCCTGGGGCAAGCAGCCAATGAGTCCCATAATGG
GTCCATCCTCCCCATCTCCTGGAGATGGGGAGAAAGTGATATGACCTTCATCTGCGTTGCCAG
GAACCCTGTCAGCAGAACTTCTCAAGCCCCATCCTTGCCAGGAAGCTCTGTGAAGGTGCTGC
TGATGACCCAGATTCTCCATGGTCCTCCTGTGTCTCCTGTTGGTGCCCCCTCCTGCTCAGTCT
CTTTGTACTGGGGCTATTTCTTTGGTTTCTGAAGAGAGAGAGACAAGAAGAGTACATTGAAGA
GAAGAAGAGAGTGGACATTTGTCGGGAAACTCCTAACATATGCCCCCATCTGGAGAGAACAC
AGAGTACGACACAATCCCTCACACTAATAGAACAATCCTAAAGGAAGATCCAGCAAATACGGT
TTACTCCACTGTGGAAATACCGAAAAAGATGGAAAATCCCCACTCACTGCTCACGATGCCAGA
CACACCAAGGCTATTTGCCTATGAGAATGTTATCTAGACAGCAGTGCACTCCCCTAAGTCTCT
GCTCA

46/168

FIGURE 46

MAGSPTCLTLIYILWQLTGSAASGPVKELVGSVGGAVTFPLKSKVKQVDSIVWTFNTTPLVTI
QPEGGTIIVTQNRNRERVDFPDGGYSLKLSKLKKNDSGIYYVGIYSSSLQQPSTQEYVLHVYE
HLSKPKVTMGLQSNKNGTCVTNLTCMEHGEEDVIYTWKALGQAANESHNGSILPISWRWGES
DMTFICVARNPVSRNFSSPILARKLCEGAADDPDSSMVLLCLLLVPLLLSLFVLGLFLWFLKR
ERQEEYIEEKKRVDICRETPNICPHSGENTYDTIPHTNRTILKEDPANTVYSTVEIPKKMEN
PHSLLTMPDTPRLFAYENVI

Important features:

Signal peptide:

amino acids 1-22

Transmembrane domain:

amino acids 224-250

Leucine zipper pattern.

amino acids 229-251

N-glycosylation sites.

amino acids 98-102, 142-146, 148-152, 172-176, 176-180, 204-208,
291-295

47/168

FIGURE 47

GGCTCGAGCGTTTCTGAGCCAGGGGTGACCATGACCTGCTGCGAAGGATGGACATCCTGCAAT
GGATTCAGCCTGCTGGTTCTACTGCTGTTAGGAGTAGTTCTCAATGCGATACCTCTAATTGTC
AGCTTAGTTGAGGAAGACCAATTTTCTCAAAACCCCATCTCTTGCTTTGAGTGGTGGTTCCCA
GGAATTATAGGAGCAGGTCTGATGGCCATTCCAGCAACAACAATGTCCTTGACAGCAAGAAAA
AGAGCGTGCTGCAACAACAGAACTGGAATGTTTCTTTCATCATTTTTTCAGTGTGATCACAGTC
ATTGGTGCTCTGTATTGCATGCTGATATCCATCCAGGCTCTCTTAAAAGGTCCTCTCATGTGT
AATTCTCCAAGCAACAGTAATGCCAATTGTGAATTTTCATTGAAAAACATCAGTGACATTCAT
CCAGAATCCTTCAACTTGCAGTGGTTTTTTCAATGACTCTTGTGCACCTCCTACTGGTTTCAAT
AAACCCACCAGTAACGACACCATGGCGAGTGGCTGGAGAGCATCTAGTTTCCACTTCGATTCT
GAAGAAAACAAACATAGGCTTATCCACTTCTCAGTATTTTTTAGGTCTATTGCTTGTTGGAATT
CTGGAGGTCCTGTTTGGGCTCAGTCAGATAGTCATCGGTTTCCTTGGCTGTCTGTGTGGAGTC
TCTAAGCGAAGAAGTCAAATTGTGTAGTTTAATGGGAATAAAATGTAAGTATCAGTAGTTTGA
AAAAAAAAAA

48/168

FIGURE 48

MTCCEGWTSCNGFSLVLLLLGVVLNAIPLIVSLVEEDQFSQNPISCFEWWFPGIIGAGLMAI
PATMSLTARKKRACCNRTGMFLSSFFSVITVIGALYCMLISIQALLKGPLMCNSPSNSNANC
EESLKNISDIHPESFNLQWFFNDSCAPPTGFNKPTSNDTMASGWRASSFHFDSEENKHRLIHF
SVFLGLLLVGILEVLFGLSQIVIGFLGCLCGVSKRRSQIV

Important features:**Transmembrane domains:**

amino acids 10-31 (type II), 50-72, 87-110, 191-213

N-glycosylation sites.

amino acids 80-84, 132-136, 148-152, 163-167

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 223-227

N-myristoylation sites.

amino acids 22-28, 54-60, 83-89, 97-103, 216-222

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 207-218

TNFR/NGFR family cysteine-rich region protein.

amino acids 4-12

49/168

FIGURE 49

ATCCGTTCTCTGCGCTGCCAGCTCAGGTGAGCCCTCGCCAAGGTGACCTCGCAGGACACTGGT
GAAGGAGCAGTGAGGAACCTGCAGAGTCACACAGTTGCTGACCAATTGAGCTGTGAGCCTGGA
GCAGATCCGTGGGCTGCAGACCCCCGCCCCAGTGCCTCTCCCCCTGCAGCCCTGCCCCCTCGAA
CTGTGACATGGAGAGAGTGACCCTGGCCCTTCTCCTACTGGCAGGCCTGACTGCCTTGGAAGC
CAATGACCCATTTGCCAATAAAGACGATCCCTTCTACTATGACTGGAAAAACCTGCAGCTGAG
CGGACTGATCTGCGGAGGGCTCCTGGCCATTGCTGGGATCGCGGCAGTTCTGAGTGGCAAATG
CAAATACAAGAGCAGCCAGAAGCAGCACAGTCCTGTACCTGAGAAGGCCATCCCACATCAC
TCCAGGCTCTGCCACTACTTGCTTGAGCACAGGACTGGCCTCCAGGGATGGCCTGAAGCCTAAC
ACTGGCCCCCAGCACCTCCTCCCCTGGGAGGCCTTATCCTCAAGGAAGGACTTCTCTCCAAGG
GCAGGCTGTTAGGCCCTTTCTGATCAGGAGGCTTCTTTATGAATTAAACTCGCCCCACCACC
CCCTCA

50/168

FIGURE 50

MERVTLALLLLAGLTALEANDPFANKDDPFYYDWKNLQLSGLICGGLLAIAAGIAAVLSGKCKY
KSSQKQHSPVPEKAIPPLITPGSATTC

Important features:

Signal peptide:

amino acids 1-16

Transmembrane domain:

amino acids 36-59

N-myristoylation sites.

amino acids 41-47, 45-51, 84-90

Extracellular proteins SCP/Tpx-1/Ag5/PR-1/Sc7.

amino acids 54-67

[illegible]

52/168

FIGURE 52

MKFQGPLACLLLLALCLGSGEAGPLQSGEESTGTNIGEALGHGLGDALSEG VGKAIGKEAGGAA
GSKVSEALGQGTREAVGTGVRQVPGFGAADALGNRVGEAAHALGNTGHEIGRQAEDVIRHGAD
AVRGSWQGVPGHSGAWETSGGHGIFGSQGGGLGGQQGNPGLGTPWVHGYPGNSAGSFGMNPQ
GAPWGQGGNGGPPNFGTNTQGAVAQPGYGSVRASNQNEGCTNPPPSGSGGGSSNSGGGSGSQS
GSSGSGSNGDNNNGSSSGGSSSGSSSGSSSGSSSGSSSGSSSGNSGGSRGDSGSESSWGSSTG
SSSGNHGGSGGGNGHKPGCEKPGNEARGSGESGIQGFRGQGVSSNMREISKEGNRLLGGSGDN
YRQQGSSWGSGGGDAVGGVNTVNSETSPGMFNFDTFWKNFKSKLGFINWDINKDQRSSRIP

Signal peptide:

amino acids 1-21

N-glycosylation site.

amino acids 265-269

Glycosaminoglycan attachment site.

amino acids 235-239, 237-241, 244-248, 255-259, 324-328, 388-392

Casein kinase II phosphorylation site.

amino acids 26-30, 109-113, 259-263, 300-304, 304-308

N-myristoylation site.

amino acids 17-23, 32-38, 42-48, 50-56, 60-66, 61-67, 64-70,
74-80, 90-96, 96-102, 130-136, 140-146, 149-155, 152-158,
155-161, 159-165, 163-169, 178-184, 190-196, 194-200, 199-205,
218-224, 236-242, 238-244, 239-245, 240-246, 245-251, 246-252,
249-252, 253-259, 256-262, 266-272, 270-276, 271-277, 275-281,
279-285, 283-289, 284-290, 287-293, 288-294, 291-297, 292-298,
295-301, 298-304, 305-311, 311-317, 315-321, 319-325, 322-328,
323-329, 325-331, 343-349, 354-360, 356-362, 374-380, 381-387,
383-389, 387-393, 389-395, 395-401

Cell attachment sequence.

amino acids 301-304

53/168

FIGURE 53

GGAGAAGAGGTTGTGTGGGACAAGCTGCTCCCGACAGAAGGATGTCGCTGCTGAGCCTGCCCT
GGCTGGGCCTCAGACCGGTGGCAATGTCCCCATGGCTACTCCTGCTGCTGGTTGTGGGCTCCT
GGCTACTCGCCCGCATCCTGGCTTGGACCTATGCCTTCTATAACAACCTGCCGCCGGCTCCAGT
GTTTCCCACAGCCCCCAAACGGAACTGGTTTTGGGGTCACCTGGGCCTGATCACTCCTACAG
AGGAGGGGCTTGAAGGACTCGACCCAGATGTCGGCCACCTATTCCCAGGGCTTTACGGTATGGC
TGGGTCCCATCATCCCCTTCATCGTTTTATGCCACCCTGACACCATCCGGTCTATCACCAATG
CCTCAGCTGCCATTGCACCCAAGGATAATCTCTTCATCAGGTTCTGAAGCCCTGGCTGGGAG
AAGGGATACTGCTGAGTGGCGGTGACAAGTGGAGCCGCCACCGTCGGATGCTGACGCCCCGCCT
TCCATTTCAACATCCTGAAGTCCTATATAACGATCTTCAACAAGAGTGCAAACATCATGCTTG
ACAAGTGGCAGCACCTGGCCTCAGAGGGCAGCAGTCGTCTGGACATGTTTGAGCACATCAGCC
TCATGACCTTGGACAGTCTACAGAAATGCATCTTCAGCTTTGACAGCCATTGTCAGGAGAGGC
CCAGTGAATATATTGCCACCATCTTGGAGCTCAGTGCCCTTGTAGAGAAAAGAAGCCAGCATA
TCCTCCAGCACATGGACTTTCTGTATTACCTCTCCCATGACGGGCGGCGCTTCCACAGGGCCT
GCCGCCTGGTGCATGACTTCACAGACGCTGTCATCCGGGAGCGGCGTCGCACCCTCCCCACTC
AGGGTATTGATGATTTTTTCAAAGACAAAGCCAAGTCCAAGACTTTGGATTTCAATTGATGTGC
TTCTGCTGAGCAAGGATGAAGATGGGAAGGCATTGTCAGATGAGGATATAAGAGCAGAGGCTG
ACACCTTCATGTTTGGAGGCCATGACACCACGGCCAGTGGCCTCTCCTGGGTCCTGTACAACC
TTGCGAGGCACCCAGAATACCAGGAGCGCTGCCGACAGGAGGTGCAAGAGCTTCTGAAGGACC
GCGATCCTAAAGAGATTGAATGGGACGACCTGGCCCAGCTGCCCTTCCTGACCATGTGCGTGA
AGGAGAGCCTGAGGTTACATCCCCCAGCTCCCTTCATCTCCCGATGCTGCACCCAGGACATTG
TTCTCCCAGATGGCCGAGTCATCCCCAAAGGCATTACCTGCCTCATCGATATTATAGGGGTCC
ATCACAACCCAACTGTGTGGCCGGATCCTGAGGTCTACGACCCCTTCCGCTTTGACCCAGAGA
ACAGCAAGGGGAGGTCACCTCTGGCTTTTATTCCTTTCTCCGCAGGGCCCAGGAACTGCATCG
GGCAGGCGTTCGCCATGGCGGAGATGAAAGTGGTCCTGGCGTTGATGCTGCTGCACTTCCGGT
TCCTGCCAGACCACACTGAGCCCCGCAGGAAGCTGGAATTGATCATGCGCGCCGAGGGCGGGC
TTTGGCTGCGGGTGGAGCCCCTGAATGTAGGCTTGCAGTGACTTTCTGACCCATCCACCTGTT
TTTTTGCAGATTGTCATGAATAAAACGGTGCTGTCAA

54/168

FIGURE 54

MSLLSLPWLGLRPVAMSPWLLLLLVVGSWLLARILAWTYAFYNNCRRLQCFPQPPKRNWFWGH
LGLITPTEEGLKDSTQMSATYSQGFTVWLGPIIPFIVLCHPDTIRSITNASAAIAPKDNLFIR
FLKPWLGE GILLSGGDKWSRHRRLTPAFHFENILKSYITIFNKSANIMLDKWQHLASEGSSRL
DMFEHISLMTLDSLQKCI FSFDSHCQERPSEYIATILELSALVEKRSQHILQHMDFLYYLSHD
GRRFHRACRLVHDFTDVIRERRRTLPTQGIDDFKDKAKSKTLDFIDVLLLSKDEDGKALSD
EDIRAEADTFMFGGHDTTASGLSWVLYNLARHPEYQERCRCQEVQELLKDRDPKEIEWDDLAQL
PFLTMCVKESLRLHPPAPFISRCCTQDIVLPDGRVIPKGITCLIDIIGVHHNPTVWPDPEVYD
PFRFDPENSKGRSPLAFIPFSAGPRNCIGQAFAMAEMKVVLALMLLHFRFLPDHTEPRRKLEL
IMRAEGGLWLRVEPLNVGLQ

Important features:**Transmembrane domains:**

amino acids 13-32 (type II), 77-102

Cytochrome P450 cysteine heme-iron ligand signature.

amino acids 461-471

N-glycosylation sites.

amino acids 112-116, 168-172

55/168

FIGURE 55

[illegible]

56/168

FIGURE 56

MGPVKQLKRMFEPTRLIATIMVLLCFALTLCSAFWWHNKGLALIFCILQSLALTWYSLSFIPF
ARDAVKKCFVCLA

Important features:

Signal peptide:

amino acids 1-33

Type II fibronectin collagen-binding domain protein.

amino acids 30-72

57/168

FIGURE 57

CGGCTCGAGCTCGAGCCGAATCGGCTCGAGGGGCGAGTGGAGCACCCAGCAGGCCGCCAACATGCTCTGTCTGTGC
CTGTACGTGCCGGTCATCGGGGAAGCCCAGACCGAGTTCCAGTACTTTGAGTCGAAGGGGCTCCCTGCCGAGCTG
AAGTCCATTTTCAAGCTCAGTGTCTTCATCCCCTCCCAGGAATTCTCCACCTACCGCCAGTGGAAAGCAGAAAATT
GTACAAGCTGGAGATAAGGACCTTGATGGGCAGCTAGACTTTGAAGAATTTGTCCATTATCTCCAAGATCATGAG
AAGAAGCTGAGGCTGGTGTTTAAGATTTTGGACAAAAAGAATGATGGACGCATTGACGCGCAGGAGATCATGCAG
TCCCTGCGGGACTTGGGAGTCAAGATATCTGAACAGCAGGCAGAAAAAATTCTCAAGAGCATGGATAAAAACGGC
ACGATGACCATCGACTGGAACGAGTGGAGAGACTACCACCTCCTCCACCCCGTGGAAAACATCCCCGAGATCATC
CTCTACTGGAAGCATTCCACGATCTTTGATGTGGGTGAGAATCTAACGGTCCCGGATGAGTTCACAGTGGAGGAG
AGGCAGACGGGGATGTGGTGGAGACACCTGGTGGCAGGAGGTGGGGCAGGGGGCGTATCCAGAACCTGCACGGCC
CCCCTGGACAGGCTCAAGGTGCTCATGCAGGTCCATGCCTCCCGCAGCAACAACATGGGCATCGTTGGTGGCTTC
ACTCAGATGATTCGAGAAGGAGGGGGCCAGGTCACTCTGGCGGGGCAATGGCATCAACGTCTCAAAATTGCCCCC
GAATCAGCCATCAAATTCATGGCCTATGAGCAGATCAAGCGCCTTGTTGGTAGTGACCAGGAGACTCTGAGGATT
CAGGAGAGGCTTGTTGGCAGGGTCCTTGGCAGGGGGCCATCGCCCAGAGCAGCATCTACCCAATGGAGGTCTGAAG
ACCCGGATGGCGCTGCGGAAGACAGGCCAGTACTCAGGAATGCTGGACTGCGCCAGGAGGATCCTGGCCAGAGAG
GGGGTGGCCGCTTCTACAAAGGCTATGTCCCAACATGCTGGGCATCATCCCCTATGCCGGCATCGACCTTGCA
GTCTACGAGACGCTCAAGAATGCCTGGCTGCAGCACTATGCAGTGAACAGCGCGGACCCCGGCGTGTGTGTGCTC
CTGGCCTGTGGCACCATGTCCAGTACCTGTGGCCAGCTGGCCAGCTACCCCTGGCCCTAGTCAGGACCCGGATG
CAGGCGCAAGCCTCTATTGAGGGCGCTCCGGAGGTGACCATGAGCAGCCTCTTCAAACATATCCTGCGGACCGAG
GGGGCCTTCGGGCTGTACAGGGGGCTGGCCCCCACTTCATGAAGGTATCCCAGCTGTGAGCATCAGCTACGTG
GTCTACGAGAACCTGAAGATCACCTGGGCGTGCAGTCGCGGTGACGGGGGAGGGCCGCCCGGCAGTGGACTCG
CTGATCCTGGGCGCAGCCTGGGGTGTGCAGCCATCTCATTCTGTGAATGTGCCAACACTAAGCTGTCTCGAGCC
AAGCTGTGAAAACCTAGACGCACCCGCAGGGAGGGTGGGGAGAGCTGGCAGGCCCCAGGGCTTGTCTCTGCTGACC
CCAGCAGACCCTCCTGTTGGTTCCAGCGAAGACCACAGGCATTCCTTAGGGTCCAGGGTCAGCAGGCTCCGGGCT
CACATGTGTAAGGACAGGACATTTTCTGCAGTGCCTGCCAATAGTGAGCTTGGAGCCTGGAGGCCGGCTTAGTTC
TTCCATTTACCCCTTGCAGCCAGCTGTTGGCCACGGCCCCCTGCCCTCTGGTCTGCCGTGCATCTCCCTGTGCCCT
CTTGCTGCCTGCCTGTCTGCTGAGGTAAGGTGGGAGGAGGGCTACAGCCACATCCCACCCCTCGTCCAATCCC
ATAATCCATGATGAAAGGTGAGGTACGTGGCCTCCCAGGCCTGACTTCCCAACCTACAGCATTGACGCCAACTT
GGCTGTGAAGGAAGAGGAAAGGATCTGGCCTTGTGGTCACTGGCATCTGAGCCCTGCTGATGGCTGGGGCTCTCG
GGCATGCTTGGGAGTGCAGGGGGCTCGGGCTGCCTGGCCTGGCTGCACAGAAGGCAAGTGCTGGGGCTCATGGTG
CTCTGAGCTGGCCTGGACCCTGTCAGGATGGGCCCCACCTCAGAACCAACTCACTGTCCCCACTGTGGCATGAG
GGCAGTGGAGCACCATGTTTGAGGGCGAAGGGCAGAGCGTTTGTGTGTTCTGGGGAGGGAAGGAAAAGGTGTTGG
AGGCCTTAATTATGGAAGTGTGGGAAAAGGGTTTTGTCCAGAAGGACAAGCCGGACAAATGAGCGACTTCTGTGC
TTCCAGAGGAAGACGAGGGAGCAGGAGCTTGGCTGACTGCTCAGAGTCTGTTCTGACGCCCTGGGGGTTCCTGTC
CAACCCAGCAGGGGCGCAGCGGGACCAGCCCCACATTCCTTGTGTCACTGCTTGGAACTATTTATTTTGTGTA
TTTATTTGAACAGAGTTATGTCCTAACTATTTTTATAGATTTGTTTAAATTAATAGCTTGTCAATTTTCAAGTTCAT
TTTTTATTCATATTTATGTTTCATGGTTGATTGTACCTTCCCAAGCCCGCCAGTGGGATGGGAGGAGGAGAGAA
GGGGGGCCTTGGGCCGCTGCAGTCACATCTGTCCAGAGAAATTCCTTTTGGGACTGGAGGCAGAAAAGCGGCCAG
AAGGCAGCAGCCCTGGCTCCTTTCTTTGGCAGGTTGGGGAAGGGCTTGCCCCCAGCCTTAGGATTTCAGGGTTT
GACTGGGGGCGTGGAGAGAGAGGGAGGAACCTCAATAACCTTGAAGGTGGAATCCAGTTATTTCTGCGCTGCCA
GGGTTTCTTTATTTCACTCTTTTCTGAATGTCAAGGCAGTGAGGTGCCTCTCACTGTGAATTTGTGGTGGGCGGG
GGCTGGAGGAGAGGGTGGGGGGCTGGCTCCGTCCCTCCCAGCCTTCTGCTGCCCTTGCTTAACAATGCCGGCCAA
CTGGCGACCTCACGGTTGCACTTCCATTCCACCAGAATGACCTGATGAGGAAATCTTCAATAGGATGCAAAGATC
AATGCAAAAATTGTTATATATGAACATATAACTGGAGTCGTCAAAAAGCAAATTAAGAAAGAATTGGACGTTAGA
AGTTGTCAATTTAAAGCAGCCTTCTAATAAAGTTGTTTCAAAGCTGAAAAAAGGAAAAAAGGAAAAAAGGAAAAA
AAAAAAGGAAAAAAGGAAAAAAGGAAAAAAGGAAAAAAGGAAAAAAGGAAAAAAGGAAAAAAGGAAAAAAGGAAAAA

58/168

FIGURE 58

MLCLCLYVPVIGEAQTEFQYFESKGLPAELKSIFKLSVFIPSQEFSTYRQWKQKIVQAGDKDL
DGQLDFEEFVHYLQDHEKKLRLVFKILDKKNDGRIDAQEIMQSLRDLGVKISEQQAEEKILKSM
DKNGTMTIDWNEWRDYHLLHPVENIPEIILYWKHSTIFDVGENLTVPDEFTVEERQTGMWWRH
LVAGGGAGAVSRTCTAPLDRLKVLMOVHASRSNNMGIVGGFTQMIREGGARSLWRGNGINVLK
IAPESAIKFMAYEQIKRLVGSDQETLRIHERLVAGSLAGAIQAQSSIYPMEVLKTRMALRKTGQ
YSGMLDCARRILAREGVAAFYKGYVPNMLGIIIPYAGIDLAVYETLKNAWLQHYAVNSADPGVF
VLLACGTMSSTCGQLASYPLALVRTRMQAQASIEGAPEVTMSSLEFKHILRTEGAFGLYRGLAP
NFMKVIPAVSISYVVYENLKITLGVQSR

Important features:**Signal peptide:**

amino acids 1-16

Putative transmembrane domains:

amino acids 284-304, 339-360, 376-394

Mitochondrial energy transfer proteins signature.

amino acids 206-215, 300-309

N-glycosylation sites.

amino acids 129-133, 169-173

Elongation Factor-hand calcium-binding protein.

amino acids 54-73, 85-104, 121-140

[illegible]

60/168

FIGURE 60

MASLGQILFWSIISIIIIILAGAIALIIGFGISGRHSITVTTVASAGNIGEDGILSCTFEPDIK
LSDIVIQWLKEGVLGLVHEFKEGKDELSEQDEMFRGRTAVFADQVIVGNASLRLKNVQLTDAG
TYKCYIITSKGKGNANLEYKTGAFSMPEVNVDYNASSETLRCEAPRWFPQPTV VWASQVDQGA
NFSEVSNTSFELNSENVTMKVVSVLNVNTINNTYSCMIENDIAKATGDIKVTSESEIKRRSHLQ
LLNSKASLCVSSFFAISWALLPLSPYMLK

Important features:**Signal peptide:**

amino acids 1-28

Transmembrane domain:

amino acids 258-281

N-glycosylation sites.amino acids 112-116, 160-164, 190-194, 196-200, 205-209, 216-220,
220-224**N-myristoylation sites.**

amino acids 52-58, 126-132, 188-194

61/168

FIGURE 61

TGACGTCAGAATCACCCATGGCCAGCTATCCTTACCGGCAGGGCTGCCCAGGAGCTGCAGGACA
AGCACCAGGAGCCCCCTCCGGGTAGCTACTACCCTGGACCCCCCAATAGTGGAGGGCAGTATGG
TAGTGGGCTACCCCCTGGTGGTGGTTATGGGGGTCTTGCCCCTGGAGGGCCTTATGGACCACC
AGCTGGTGGAGGGCCCTATGGACACCCCAATCCTGGGATGTTCCCCTCTGGAACCTCCAGGAGG
ACCATATGGCGGTGCAGCTCCCGGGGGCCCCCTATGGTCAGCCACCTCCAAGTTCCTACGGTGC
CCAGCAGCCTGGGCTTTATGGACAGGGTGGCGCCCCCTCCCAATGTGGATCCTGAGGCCTACTC
CTGGTTCCAGTCGGTGGACTCAGATCACAGTGGCTATATCTCCATGAAGGAGCTAAAGCAGGC
CCTGGTCAACTGCAATTGGTCTTCATTCAATGATGAGACCTGCCTCATGATGATAAACATGTT
TGACAAGACCAAGTCAGGCCGCATCGATGTCTACGGCTTCTCAGCCCTGTGGAAATTCATCCA
GCAGTGGAAGAACCTCTTCCAGCAGTATGACCGGGACCGCTCGGGCTCCATTAGCTACACAGA
GCTGCAGCAAGCTCTGTCCCAAATGGGCTACAACCTGAGCCCCCAGTTCACCCAGCTTCTGGT
CTCCCGCTACTGCCCACGCTCTGCCAATCCTGCCATGCAGCTTGACCGCTTCATCCAGGTGTG
CACCCAGCTGCAGGTGCTGACAGAGGCCTTCCGGGAGAAGGACACAGCTGTACAAGGCAACAT
CCGGCTCAGCTTCGAGGACTTCGTCACCATGACAGCTTCTCGGATGCTATTGACCCAACCATCT
GTGGAGAGTGGAGTGCACCAGGGACCTTTCCTGGCTTCTTAGAGTGAGAGAAGTATGTGGACA
TCTCTTCTTTTCCTGTCCCTCTAGAAGAACATTCTCCCTTGCTTGATGCAACACTGTTCCAAA
AGAGGGTGGAGAGTCCTGCATCATAGCCACCAAATAGTGAGGACCGGGGCTGAGGCCACACAG
ATAGGGGCCTGATGGAGGAGAGGATAGAAGTTGAATGTCCTGATGGCCATGAGCAGTTGAGTG
GCACAGCCTGGCACCAGGAGCAGGTCCTTGTAATGGAGTTAGTGTCCAGTCAGCTGAGCTCCA
CCCTGATGCCAGTGGTGGTGGTTCATCGGCCTGTTACCGTTAGTACCTGTGTTCCCTCACCAG
GCCATCCTGTCAAACGAGCCCATTCTTCTCAAAGTGGAATCTGACCAAGCATGAGAGAGATCT
GTCTATGGGACCAGTGGCTTGGATTCTGCCACACCCATAAATCCTTGTGTGTTAACTTCTAGC
TGCTTGGGGCTGGCCCTGCTCAGACAAATCTGCTCCCTGGGCATCTTTGGCCAGGCTTCTGCC
CCCTGCAGCTGGGACCCCTCACTTGCTGCTGCTGCTCGGCTTCAGTCTCCAGGAGACA
GTGGTCACCTCTCCCTGCCAATACTTTTTTTAATTTGCATTTTTTTTTCATTTGGGGCCAAAAG
TCCAGTGAAATTGTAAGCTTCAATAAAAGGATGAAACTCTGA

62/168

FIGURE 62

MASYPYRQGCPGAAGQAPGAPPGSYYPGPPNSGGQYGSGLPPGGGYGGPAPGGPYGPPAGGGP
YGHPNPGMFPSGTPGGPYGGAAPGGPYGQPPPSSYGAQQPGLYGQGGAPPNVDPEAYSWFQSV
DSDHSGYISMKELKQALVNCNWSSFNDETCLMMINMFDKTKSGRIDVYGFSALWKFIQQWKNL
FQQYDRDRSGSISYTELQQALSQMGYNLSPQFTQLLVSRYCPRSANPAMQLDRFIQVCTQLQV
LTEAFREKDTAVQGNIRLSFEDFVTMTASRML

Important features of the protein:

Signal peptide:

amino acids 1-19

N-glycosylation site.

amino acids 147-150

Casein kinase II phosphorylation sites.

amino acids 135-138, 150-153, 202-205, 271-274

N-myristoylation sites.

amino acids 9-14, 15-20, 19-24, 33-38, 34-39, 39-44, 43-48, 61-
66, 70-75, 78-83, 83-88, 87-92, 110-115

63/168

FIGURE 63

CAGGATGCAGGGCCGCGTGGCAGGGAGCTGCGCTCCTCTGGGCCTGCTCCTGGTCTGTCTTCA
TCTCCCAGGCCTCTTTGCCCCGGAGCATCGGTGTTGTGGAGGAGAAAGTTTCCCAAACCTTCGG
GACCAACTTGCCTCAGCTCGGACAACCTTCCTCCACTGGCCCCCTCTAACTCTGAACATCCGCA
GCCCCGCTCTGGACCCTAGGTCTAATGACTTGGCAAGGGTTCCTCTGAAGCTCAGCGTGCCTCC
ATCAGATGGCTTCCCACCTGCAGGAGGTTCTGCAGTGCAGAGGTGGCCTCCATCGTGGGGGCT
GCCTGCCATGGATTCTTGCCCCCTGAGGATCCTTGGCAGATGATGGCTGCTGCGGCTGAGGA
CCGCCTGGGGGAAGCGCTGCCTGAAGAACTCTCTTACCTCTCCAGTGCTGCGGCCCTCGCTCC
GGGCAGTGGCCCTTTGCCTGGGGAGTCTTCTCCCGATGCCACAGGCCTCTCACCTGAGGCTTC
ACTCCTCCACCAGGACTCGGAGTCCAGACGACTGCCCCGTTCTAATTCACTGGGAGCCGGGGG
AAAAATCCTTTCCCAACGCCCTCCCTGGTCTCTCATCCACAGGGTTCCTGCCTGATCACCCCTG
GGGTACCCTGAATCCCAGTGTGTCCTGGGGAGGTGGAGGCCCTGGGACTGGTTGGGGAACGAG
GCCCATGCCACACCCTGAGGGAATCTGGGGTATCAATAATCAACCCCCAGGTACCAGCTGGGG
AAATATTAATCGGTATCCAGGAGGCAGCTGGGGAAATATTAATCGGTATCCAGGAGGCAGCTG
GGGGAATATTAATCGGTATCCAGGAGGCAGCTGGGGGAATATTCATCTATACCCAGGTATCAA
TAACCCATTTCTCCTGGAGTTCTCCGCCCTCCTGGCTCTTCTTGGAACATCCCAGCTGGCTT
CCCTAATCCTCCAAGCCCTAGGTTCAGTGGGGCTTAGAGCACGATAGAGGGAAACCCAACATT
GGGAGTTAGAGTCCTGCTCCCGCCCCCTTGCTGTGTGGGCTCAATCCAGGCCCTGTTAACATGT
TTCCAGCACTATCCCCACTTTTCAGTGCCTCCCCTGCTCATCTCCAATAAAATAAAAGCACTT
ATGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

64/168

FIGURE 64

MQGRVAGSCAPLGLLLVCLHLPGLFARSIGVVEEKVSQNFGTNLPQLGQPSSTGPSNSEHPQP
ALDPRSNDLARVPLKLSVPPSDGFPPAGGSAVQRWPPSWGLPAMDSWPPEDPWQMMAAAAEDR
LGEALPEELSYLSSAAALAPGSGPLPGESSPDATGLSPEASLLHQDSESRRLPRSNSLGAGGK
ILSQRPPWSLIHRVLPDHPWGTLNPSVSWGGGGPGTGWGTRPMPHPEGIWGINNQPPGTSWGN
INRYPGGSWGNIINRYPGGSWGNIINRYPGGSWGNIHLYPGINNPFPFPGVLRPPGSSWNIPAGFP
NPPSPRLQWG

Important features of the protein:

Signal peptide:

amino acids 1-26

Casein kinase II phosphorylation sites.

amino acids 56-59, 155-158

N-myristoylation sites.

amino acids 48-53, 220-225, 221-226, 224-229, 247-252, 258-263,
259-264, 269-274, 270-275, 280-285, 281-286, 305-310

65/168

FIGURE 65

AAGGAGAGGCCACCGGGACTTCAGTGTCTCCTCCATCCCAGGAGCGCAGTGGCCACTATGGGG
TCTGGGCTGCCCCCTTGTCCTCCTCTTGACCCTCCTTGGCAGCTCACATGGAACAGGGCCGGGT
ATGACTTTGCAACTGAAGCTGAAGGAGTCTTTTCTGACAAATTCCTCCTATGAGTCCAGCTTC
CTGGAATTGCTTGAAAAGCTCTGCCTCCTCCTCCATCTCCCTTCAGGGACCAGCGTCACCCTC
CACCATGCAAGATCTCAACACCATGTTGTCTGCAACACATTGACAGCCATTGAAGCCTGTGTCC
TTCTTGGCCCCGGGCTTTTGGGCCGGGGATGCAGGAGGCAGGCCCCGACCCTGTCTTTCAGCAG
GCCCCCACCCTCCTGAGTGGCAATAAATAAAATTTCGGTATGCTG

66/168

FIGURE 66

MGSGPLVLLLLTLLGSSHGTGPGMTLQLKLKESFLTNSSESSFLELLEKLCLLLHLPSGTSV
TLHHARSQHHVVCNT

Important features:

Signal peptide:

amino acids 1-19

N-glycosylation site.

amino acids 37-41

N-myristoylation sites.

amino acids 15-21, 19-25, 60-66

67/168

FIGURE 67

ACGGACCGAGGGTTCGAGGGAGGGACACGGACCAGGAACCTGAGCTAGGTCAAAGACGCCCCGG
GCCAGGTGCCCCGTCGCAGGTGCCCCTGGCCGGAGATGCGGTAGGAGGGGCGAGCGCGAGAAG
CCCCTTCCTCGGCGCTGCCAACCCGCCACCCAGCCCATGGCGAACCCCGGGCTGGGGCTGCTT
CTGGCGCTGGGCCTGCCGTTCTTGCTGGCCCGCTGGGGCCGAGCCTGGGGGCAAATACAGACC
ACTTCTGCAAATGAGAATAGCACTGTTTTGCCTTCATCCACCAGCTCCAGCTCCGATGGCAAC
CTGCGTCCGGAAGCCATCACTGCTATCATCGTGGTCTTCTCCCTCTTGGCTGCCTTGCTCCTG
GCTGTGGGGCTGGCACTGTTGGTGCGGAAGCTTCGGGAGAAGCGGCAGACGGAGGGCACCTAC
CGGCCCAGTAGCGAGGAGCAGTTCTCCCATGCAGCCGAGGCCCGGGCCCCCTCAGGACTCCAAG
GAGACGGTGCAGGGCTGCCTGCCCATCTAGGTCCCCTCTCCTGCATCTGTCTCCCTTCATTGC
TGTGTGACCTTGGGGAAAGGCAGTGCCCTCTCTGGGCAGTCAGATCCACCCAGTGCTTAATAG
CAGGGAAGAAGGTACTTCAAAGACTCTGCCCCTGAGGTCAAGAGAGGATGGGGCTATTCATT
TTATATATTTATATAAAATTAGTAGTGAGATGTAAAAAAAAAAAAAAAAAAAAA

68/168

FIGURE 68

MANPGLGLLLALGLPFLLARWGRAWGQIQTTSANENSTVLPSSSTSSSSDGNLRPEAITAIIVV
FSLLAALLLAVGLALLVRKLREKRQTEGTYRPSSEEQFSHAAEARAPQDSKETVQGCLPI

Important features:**Signal peptide:**

amino acids 1-19

Transmembrane domain:

amino acids 56-80

N-glycosylation site.

amino acids 36-40

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 86-90

Tyrosine kinase phosphorylation site.

amino acids 86-94

N-myristoylation sites.

amino acids 7-13, 26-32

[illegible]

70/168

FIGURE 70

MGLFRGFVFLLVLCLLHQSNTSFIKLNNGFEDIVIVIDPSVPEDEKIIIEQIEDMVTASTYL
FEATEKRFFFKNVSILIPENWKENPQYKRPKHENHKHADVIVAPPTLPGRDEPYTKQFTECGE
KGEYIHFTPDLLLGKKQNEYGPPGKLFVHEWAHLRWGVFDEYNEDQPFYRAKSKKIEATRCSA
GISGRNRVYKCQGGSCLSRACRIDSTTKLYGKDCQFFPDKVQTEKASIMFMQSIDSVVEFCNE
KTHNQEAPSLQNIKCNFRSTWEVISNSEDFKNTIPMVTPPPPPVFSLLKISQRIVCLVLDKSG
SMGGKDRLNRMNQAAKHFLLOTVENGSWVGMVHFDSTATIVNKLIQIKSSDERNTLMAGLPTY
PLGGTSICSGIKYAFQVIGELHSQLDGSEVLLLLTDGEDNTASSCIDEVKQSGAIVHFIALGRA
ADEAVIEMSKITGGSHFYVSDEAQNNGLIDAFGALTSGNTDLSQKSLQLESKGLTLNSNAWMN
DTVIIDSTVGKDTFFLITWNSLPPSISLWDPSGTIMENFTVDATSKMAYLSIPGTAKVGTWAY
NLQAKANPETLTITVTSRAANSSVPPITVNAKMNKDVNSFPSPMIVYAEILQGYVPVLGANVT
AFIESQNGHTEVLELLDNGAGADSFKNDGVYSRYFTAYTENGRYSLKVRAHGGANTARLKL RP
PLNRAAYIPGWVVNGEIEANPPRPEIDEDTQTTLEDFSRTASGGAFVVSQVPSLPLPDQYPPS
QITDL DATVHEDKIIILTWTAPGDNFDVGKVQRYIIIRISASILDLRDSFDDALQVNTTDLSPKE
ANSKESFAFKPENISEENATHIFIAIKSIDKSNLTSKVSNIAQVTLFIPQANPDDIDPTPTPT
PTPTPDKSHNSGVNISTLVLSVIGSVVIVNFILSTTI

Signal peptide:

amino acids 1-21

Putative transmembrane domains:

amino acids 284-300, 617-633

Leucine zipper pattern.

amino acids 469-491, 476-498

N-glycosylation site.amino acids 20-24, 75-79, 340-344, 504-508, 542-546, 588-592,
628-632, 811-815, 832-836, 837-841, 852-856, 896-900

71/168

FIGURE 71

CTCCTTAGGTGGAAACCCTGGGAGTAGAGTACTGACAGCAAAGACCGGGAAAGACCATACGTCCCCGGGCAGGGG
TGACAAACAGGTGTCATCTTTTTGATCTCGTGTGTGGCTGCCTTCCTATTTCAAGGAAAGACGCCAAGGTAATTTT
GACCCAGAGGAGCAATGATGTAGCCACCTCCTAACCTTECCTTCTTGAACCCCCAGTTATGCCAGGATTTACTAG
AGAGTGTCAACTCAACCAGCAAGCGGCTCCTTCGGCTTAACCTTGTGGTTGGAGGAGAGAACCTTTGTGGGGCTGC
GTTCTCTTAGCAGTGCTCAGAAGTGACTTGCCTGAGGGTGGACCAGAAGAAAGGAAAGGTCCCCTCTTGCTGTTG
GCTGCACATCAGGAAGGCTGTGATGGGAATGAAGGTGAAAACCTTGAGATTTCACTTCAGTCATTGCTTCTGCCT
GCAAGATCATCCTTTAAAAGTAGAGAAGCTGCTCTGTGTGGTGGTTAACTCCAAGAGGCAGAACTCGTTCTAGAA
GGAAATGGATGCAAGCAGCTCCGGGGGGCCCCAAACGCATGCTTCCTGTGGTCTAGCCCAGGGAAGCCCTTCCGTG
GGGGCCCCGGCTTTGAGGGATGCCACCGGTTCTGGACGCATGGCTGATTCTGAATGATGATGGTTCGCCGGGGG
CTGCTTGCGTGGATTTCCCGGGTGGTGGTTTTGCTGGTGTCTCTCTGTCTGTCTATCTCTGTCTGTACATGTTG
GCCTGCACCCCCAAAAGGTGACGAGGAGCAGCTGGCACTGCCAGGGCCAACAGCCCCACGGGGAAGGAGGGGTAC
CAGGCCGTCCTTCAGGAGTGGGAGGAGCAGCACCGCAACTACGTGAGCAGCCTGAAGCGGCAGATCGCACAGCTC
AAGGAGGAGCTGCAGGAGAGGAGTGAGCAGCTCAGGAATGGGCAGTACCAAGCCAGCGATGCTGCTGGCCTGGGT
CTGGACAGGAGCCCCCAGAGAAAACCCAGGCCGACCTCCTGGCCTTCCTGCACTCGCAGGTGGACAAGGCAGAG
GTGAATGCTGGCGTCAAGCTGGCCACAGAGTATGCAGCAGTGCCTTTGATAGCTTTACTCTACAGAAGGTGTAC
CAGCTGGAGACTGGCCTTACCCGCCACCCCGAGGAGAAGCCTGTGAGGAAGGACAAGCGGGATGAGTTGGTGGAA
GCCATTGAATCAGCCTTGAGAGCCCTGAACAATCCTGCAGAGAACAGCCCCAATCACCGTCCTTACACGGCCTCT
GATTTCATAGAAGGGATCTACCGAACAGAAAGGGACAAAGGGACATTGTATGAGCTCACCTCAAAGGGGACCAC
AAACACGAATTCAAACGGCTCATCTTATTTTCGACCATTGAGCCCCATCATGAAAGTGAAAAATGAAAAGCTCAAC
ATGGCCAACACGCTTATCAATGTTATCGTGCCTCTAGCAAAAAGGGTGGACAAGTTCCGGCAGTTCATGCAGAAT
TTCAGGGAGATGTGCATTGAGCAGGATGGGAGAGTCCATCTCACTGTTGTTTACTTTGGGAAAGAAGAAATAAAT
GAAGTCAAAGGAATACTTGAAAACACTTCCAAAGCTGCCAACTTCAGGAACCTTACCTTCATCCAGCTGAATGGA
GAATTTTCTCGGGGAAGGGACTTGATGTTGGAGCCCGCTTCTGGAAGGGAAGCAACGTCCTTCTCTTTTCTGT
GATGTGGACATCTACTTCACATCTGAATTCCTCAATACGTGTAGGCTGAATACACAGCCAGGGAAGAAGGTATTT
TATCCAGTTCTTTTCAGTCAGTACAATCCTGGCATAATATACGGCCACCATGATGCAGTCCCTCCCTTGGAACAG
CAGCTGGTCATAAAGAAGGAAACTGGATTTTGGAGAGACTTTGGATTTGGGATGACGTGTGAGTATCGGTGAGAC
TTCATCAATATAGGTGGGTTTGATCTGGACATCAAAGGCTGGGGCGGAGAGGATGTGCACCTTTATCGCAAGTAT
CTCCACAGCAACCTCATAGTGGTACGGACGCCTGTGCGAGGACTCTTCCACCTCTGGCATGAGAAGCGCTGCATG
GACGAGCTGACCCCCGAGCAGTACAAGATGTGCATGCAGTCCAAGGCCATGAACGAGGCATCCCACGGCCAGCTG
GGCATGCTGGTGTTCAGGCACGAGATAGAGGCTCACCTTCGCAAACAGAAACAGAAAGACAAGTAGCAAAAAACA
TGAACTCCCAGAGAAGGATTGTGGGAGACACTTTTTCTTTCTTTTGCAATTACTGAAAGTGGCTGCAACAGAGA
AAAGACTTCCATAAAGGACGACAAAAGAATTGGACTGATGGGTGAGAGATGAGAAAGCCTCCGATTTCTCTGT
TGGGCTTTTTTACAACAGAAATCAAATCTCCGCTTTGCCTGCAAAAGTAACCCAGTTGCACCCCTGTGAAGTGTCT
GACAAAGGCAGAATGCTTGTGAGATTATAAGCCTAATGGTGTGGAGGTTTTGATGGTGTTTACAATACTGAGA
CCTGTTGTTTTGTGTGCTCATTGAAATATTGATGATTTAAGAGCAGTTTTGTAAAAAATTCATTAGCATGAAAGG
CAAGCATATTTCTCCTCATATGAATGAGCCTATCAGCAGGGCTCTAGTTTCTAGGAATGCTAAAATATCAGAAGG
CAGGAGAGGAGATAGGCTTATTATGATACTAGTGAGTACATTAAGTAAAATAAAATGGACCAGAAAAGAAAAGAA
ACCATAAATATCGTGTGATATTTTCCCCAAGATTAACCAAAAATAATCTGCTTATCTTTTTGGTTGTCCTTTTAA
CTGTCTCCGTTTTTTTTCTTTTATTTAAAAATGCACTTTTTTTCCCTTGTGAGTTATAGTCTGCTTATTTAATTAC
CACTTTGCAAGCCTTACAAGAGAGCACAAGTTGGCCTACATTTTTTATATTTTTTAAGAAGATACTTTGAGATGCA
TTATGAGAACTTTGAGTTCAAAGCATCAAATTGATGCCATATCCAAGGACATGCCAAATGCTGATTCTGTCAGGC
ACTGAATGTCAGGCATTGAGACATAGGGAAGGAATGGTTTGTACTAATACAGACGTACAGATACTTTCTCTGAAG
AGTATTTTTCGAAGAGGAGCAACTGAACACTGGAGGAAAAGAAAATGACACTTTCTGCTTTACAGAAAAGGAAACT
CATTCAGACTGGTGATATCGTGATGTACCTAAAAGTCAGAAACCACATTTTCTCCTCAGAAGTAGGGACCGCTTT
CTTACCTGTTTAAATAAACCAAGTATAACCGTGTGAACCAACAATCTTTTTCAAACAGGGTGCTCCTCCTGG
CTTCTGGCTTCCATAAGAAGAAATGGAGAAAATATATATATATATATATATTGTGAAAGATCAATCCATCTG
CCAGAATCTAGTGGGATGGAAGTTTTTGCTACATGTTATCCACCCAGGCCAGGTGGAAGTAACTGAATTATTTT
TTAAATTAAGCAGTTCTACTCAATCACCAAGATGCTTCTGAAAATTGCATTTTATTACCATTTCAAACTATTTTT
TAAAAATAAATACAGTTAACATAGAGTGGTTTCTTCATTCATGTGAAAATTATTAGCCAGCACCAGATGCATGAG
CTAATTATCTCTTTGAGTCCTTGCTTCTGTTTGCTCACAGTAACTCATTGTTTAAAGCTTCAAGAACATTCAA
GCTGTTGGTGTGTTAAAAAATGCATTGTATTGATTTGTAAGTTTATGAAATTTAATTAAACACAGGCCA
TGAATGGAAGGTGGTATTGCACAGCTAATAAAATATGATTTGTGGATATGAA

72/168

FIGURE 72

MMVRRGLLAWISRVVLLVLLCCAISVLYMLACTPKGDEEQALALPRANSPTGKEGYQAVLQE
WEEQHRNYVSSLKRQIAQLKEELQERSEQLRNGQYQASDAAGLGLDRSPPEKTQADLLAFLHS
QVDKAEVNAGVKLATEYAAVPFDSFTLQKVYQLETGLTRHPEEKPV RKDKRDELVEAIESALE
TLNNPAENSPNHRPYTASDFIEGIYRTERDKGTLYELTFKGDHKHEFKRLILFRPFSPIMKVK
NEKLNMA NTLINVIVPLAKRVDKFRQFMQNFREMCIEQDGRVHLTVVYFGKEEINEVKGILEN
TSKAANFRNFTFIQLNGEFSRGKGLDVGARFWKGSNVLLFFCDVDIYFTSEFLNTCRLNTQPG
KKVFYPVLFSQYNPGIIYGHHDVPPLEQQLVIKKETGFWRDFGFGMTCQYRSDFINIGGFDL
DIKGWGGEDVHLYRKYLHSNLIVVRTPV RGLFHLWHEKRCMDELTPEQYKMCMQSKAMNEASH
GQLGMLVFRHEIEAHLRKQKQKTSSKKT

Important features:**Signal peptide:**

amino acids 1-27

N-glycosylation sites.

amino acids 315-319, 324-328

N-myristoylation sites.

amino acids 96-102, 136-142, 212-218, 311-317, 339-345, 393-399

Amidation site.

amino acids 377-381

73/168

FIGURE 73

GAGACTGCAGAGGGGAGATAAAGAGAGAGGGGCAAAGAGGCAGCAAGAGATTTGTCCTGGGGATC
CAGAAACCCATGATACCCTACTGAACACCGAATCCCCCTGGAAGCCCACAGAGACAGAGACAGC
AAGAGAAGCAGAGATAAATACTCACGCCAGGAGCTCGCTCGCTCTCTCTCTCTCTCTCTCA
CTCCTCCCTCCCTCTCTCTCTGCTGTCCTAGTCCTCTAGTCCTCAAATTCCCAGTCCCCTGC
ACCCCTTCCTGGGACACTATGTTGTTCTCCGCCCTCCTGCTGGAGGTGATTTGGATCCTGGCT
GCAGATGGGGGTCAACACTGGACGTATGAGGGCCCACATGGTCAGGACCATTGGCCAGCCTCT
TACCCTGAGTGTGGAAACAATGCCAGTCGCCCATCGATATTCAGACAGACAGTGTGACATTT
GACCCTGATTTGCCTGCTCTGCAGCCCCACGGATATGACCAGCCTGGCACCGAGCCTTTGGAC
CTGCACAACAATGGCCACACAGTGCAACTCTCTCTGCCCTCTACCCTGTATCTGGGTGGACTT
CCCCGAAAATATGTAGCTGCCAGCTCCACCTGCACTGGGGTCAGAAAGGATCCCCAGGGGGG
TCAGAACACCAGATCAACAGTGAAGCCACATTTGCAGAGCTCCACATTGTACATTATGACTCT
GATTCCTATGACAGCTTGAGTGAGGCTGCTGAGAGGCCTCAGGGCCTGGCTGTCCTGGGCATC
CTAATTGAGGTGGGTGAGACTAAGAATATAGCTTATGAACACATTCTGAGTCACTTGCATGAA
GTCAGGCATAAAGATCAGAAGACCTCAGTGCCTCCCTTCAACCTAAGAGAGCTGCTCCCCAAA
CAGCTGGGGCAGTACTTCCGCTACAATGGCTCGCTCACAACCTCCCCCTTGCTACCAGAGTGTG
CTCTGGACAGTTTTTTATAGAAGGTCCCAGATTTCAATGGAACAGCTGGAAAAGCTTCAGGGG
ACATTGTTCTCCACAGAAGAGGAGCCCTCTAAGCTTCTGGTACAGAACTACCGAGCCCTTCAG
CCTCTCAATCAGCGCATGGTCTTTGCTTCTTTCATCCAAGCAGGATCCTCGTATACCACAGGT
GAAATGCTGAGTCTAGGTGTAGGAATCTTGGTTGGCTGTCTCTGCCTTCTCCTGGCTGTTTAT
TTCATTGCTAGAAAGATTCGGAAGAAGAGGCTGGAAAACCGAAAGAGTGTGGTCTTCACCTCA
GCACAAGCCACGACTGAGGCATTAAATTCCTTCTCAGATACCATGGATGTGGATGACTTCCCTT
CATGCCTATCAGGAAGCCTCTAAAATGGGGTGTAGGATCTGGCCAGAAACACTGTAGGAGTAG
TAAGCAGATGTCCTCCTTCCCCCTGGACATCTCTTAGAGAGGAATGGACCCAGGCTGTCATTCC
AGGAAGAAGTGCAGAGCCTTCAGCCTCTCCAAACATGTAGGAGGAAATGAGGAAATCGCTGTG
TTGTTAATGCAGAGANCAAACTCTGTTTAGTTGCAGGGGAAGTTTGGGATATACCCCAAAGTC
CTCTACCCCTCACTTTTATGGCCCTTCCCTAGATATACTGCGGGATCTCTCCTTAGGATAA
AGAGTTGCTGTTGAAGTTGTATATTTTTGATCAATATATTTGGAAATTAAAGTTTCTGACTTT

74/168

FIGURE 74

MLFSALLLEVIWILAADGGQHWTYEGPHGQDHWPA SYPECGNNAQSPIDIQTDSVT FDPDLPA
LQPHGYDQPGTEPLDLHNNGHTVQLSLPSTLYLGGLPRKYVAAQLHLHWGQKGSPGGSEHQIN
SEATFAELHIVHYDSYDSLSEAAERPQGLAVLGILIEVGETKNIAYEHILSHLHEVRHKDQ
KTSVPPFNLRELLPKQLGQYFRYNGSLTTPPCYQSVLWTVFYRRSQISMEQLEKLQGTLFSTE
EEPSKLLVQNYRALQPLNQRMVFASFIQAGSSYTTGEMLSLGVGILVGCLCLLLAVYFIARKI
RKKRLENRKS VVFTSAQATTEA

Important features of the protein:

Signal peptide:

amino acids 1-15

Transmembrane domain:

amino acids 291-310

N-glycosylation site.

amino acids 213-216

Eukaryotic-type carbonic anhydrases proteins

amino acids 197-245, 104-140, 22-69

75/168

FIGURE 75

TGCCGCTGCCGCCGCTGCTGCTGTTGCTCCTGGCGGGCGCCTTGGGGACGGGCAGTTCCCTGTG
TCTCTGGTGGTTTGCCTAAACCTGCAAACATCACCTTCTTATCCATCAACATGAAGAATGTCC
TACAATGGACTCCACCAGAGGGTCTTCAAGGAGTTAAAGTTACTTACACTGTGCAGTATTTCA
TCACAAATTGGCCCACCAGAGGTGGCACTGACTACAGATGAGAAGTCCATTTCTGTTGTCCTG
ACAGCTCCAGAGAAGTGGAAGAGAAATCCAGAAGACCTTCCTGTTTCCATGCAACAAATATAC
TCCAATCTGAAGTATAACGTGTCTGTGTTGAATACTAAATCAAACAGAACGTGGTCCCAGTGT
GTGACCAACCACACGCTGGTGCTCACCTGGCTGGAGCCGAACACTCTTTACTGCGTACACGTG
GAGTCCTTCGTCCCAGGGCCCCCTCGCCGTGCTCAGCCTTCTGAGAAGCAGTGTGCCAGGACT
TTGAAAGATCAATCATCAGAGTTCAAGGCTAAAATCATCTTCTGGTATGTTTTGCCCATATCT
ATTACCGTGTTTCTTTTTTCTGTGATGGGCTATTCCATCTACCGATATATCCACGTTGGCAAA
GAGAAACACCCAGCAAATTTGATTTTGATTTATGGAAATGAATTTGACAAAAGATTCTTTGTG
CCTGCTGAAAAAATCGTGATTAACTTTATCACCTCAATATCTCGGATGATTCTAAAATTTCT
CATCAGGATATGAGTTTACTGGGAAAAAGCAGTGATGTATCCAGCCTTAATGATCCTCAGCCC
AGCGGGAACCTGAGGCCCCCTCAGGAGGAAGAGGAGGTGAAACATTTAGGGTATGCTTCGCAT
TTGATGGAAATTTTTTGTGACTCTGAAGAAAACACGGAAGGTACTTCTCTCACCCAGCAAGAG
TCCCTCAGCAGAACAAATACCCCCGGATAAAACAGTCATTGAATATGAATATGATGTCAGAACC
ACTGACATTTGTGCGGGGCCTGAAGAGCAGGAGCTCAGTTTGCAGGAGGAGGTGTCCACACAA
GGAACATTATTGGAGTCGCAGGCAGCGTTGGCAGTCTTGGGCCCCGCAAACGTTACAGTACTCA
TACACCCCTCAGCTCCAAGACTTAGACCCCCTGGCGCAGGAGCACACAGACTCGGAGGAGGGG
CCGGAGGAAGAGCCATCGACGACCCTGGTCGACTGGGATCCCCAAACTGGCAGGCTGTGTATT
CCTTCGCTGTCCAGCTTCGACCAGGATTCAGAGGGCTGCGAGCCTTCTGAGGGGGATGGGCTC
GGAGAGGAGGGTCTTCTATCTAGACTCTATGAGGAGCCGGCTCCAGACAGGCCACCAGGAGAA
AATGAAACCTATCTCATGCAATTCATGGAGGAATGGGGGTATATGTGCAGATGGAAAACTGA
TGCCAACACTTCCTTTTGCCTTTTGTTCCTGTGCAAACAAGTGAGTCACCCCTTTGATCCCA
GCCATAAAGTACCTGGGATGAAAGAAGTTTTTTCCAGTTTGTGAGTGTCTGTGAGAATTACTT
ATTTCTTTTCTCTATTCTCATAGCACGTGTGTGATTGGTTCATGCATGTAGGTCTCTTAACAA
TGATGGTGGGCCTCTGGAGTCCAGGGGCTGGCCGGTTGTTCTATGCAGAGAAAGCAGTCAATA
AATGTTTGCCAGACTGGGTGCAGAATTTATTCAGGTGGGTGT

76/168

FIGURE 76

MSYNGLHQRVFKELKLLTLCSISSQIGPPEVALTTDEKSISVVLTAPEKWKRNPEDLPVSMQQ
IYSNLKYNVSVLNTKSNRTWSQCVTNHTLVLTWLEPNTLYCVHVESFVPGPPRAQPSEKQCA
RTLKDQSSEFKAKIIFWYVLPISITVFLFSVMGYSIYRYIHVGKEKHPANLILYGNEDKRF
FVPAEKIVINFITLNISSDDSKISHQDMSLLGKSSDVSSSLNDPQPSGNLRPPQEEEEVKHLGYA
SHLMEIFCDSEENTEGTSLTQQESLSRTIPDKTVIEWEYDVRTTDICAGPEEQELSLQEEVS
TQGTLLSQAALAVLGPOTLQYSYTPQLQDLPLAQEHTDSEEGPEEPPSTTLVDWDPQTGRL
CIPSLSSFDQDSEGCEPSEGDLGEEGLLSRLYEPPAPDRPPGENETYLMQFMEEWGLYVQMEN

Important features:**Signal peptide:**

amino acids 1-28

Transmembrane domain:

amino acids 140-163

N-glycosylation sites.

amino acids 71-74, 80-83, 89-92, 204-207, 423-426

77/168

FIGURE 77

GAGGAGCGGGCCGAGGACTCCAGCGTGCCCAGGTCTGGCATCCTGCACTTGCTGCCCTCTGAC
ACCTGGGAAGATGGCCGGCCCGTGGACCTTCACCCTTCTCTGTGGTTTGCTGGCAGCCACCTT
GATCCAAGCCACCCTCAGTCCCAGTTCATCCTCGGCCCAAAGTCATCAAAGAAAA
GCTGACACAGGAGCTGAAGGACCACAACGCCACCAGCATCCTGCAGCAGCTGCCGCTGCTCAG
TGCCATGCGGGAAAAGCCAGCCGGAGGCATCCCTGTGCTGGGCAGCCTGGTGAACACCGTCCT
GAAGCACATCATCTGGCTGAAGGTCATCACAGCTAACATCCTCCAGCTGCAGGTGAAGCCCTC
GGCCAATGACCAGGAGCTGCTAGTCAAGATCCCCCTGGACATGGTGGCTGGATTCAACACGCC
CCTGGTCAAGACCATCGTGGAGTTCCACATGACGACTGAGGCCCAAGCCACCATCCGCATGGA
CACCAGTGCAAGTGGCCCCACCCGCCTGGTCCTCAGTGACTGTGCCACCAGCCATGGGAGCCT
GCGCATCCAAGTCTGTATAAGCTCTCCTTCCTGGTGAACGCCTTAGCTAAGCAGGTTCATGAA
CCTCCTAGTGCCATCCCTGCCCAATCTAGTGAAAAACCAGCTGTGTCCCGTGATCGAGGCTTC
CTTCAATGGCATGTATGCAGACCTCCTGCAGCTGGTGAAGGTGCCCATTTCCCTCAGCATTGA
CCGTCTGGAGTTTGACCTTCTGTATCCTGCCATCAAGGGTGACACCATTAGCTCTACCTGGG
GGCCAAGTTGTTGGACTCACAGGGAAAGGTGACCAAGTGGTTCAATAACTCTGCAGCTTCCCT
GACAATGCCCACCCTGGACAACATCCCGTTCAGCCTCATCGTGAGTCAGGACGTGGTGAAAGC
TGCAGTGGCTGCTGTGCTCTCTCCAGAAGAATTCATGGTCCTGTTGGACTCTGTGCTTCCTGA
GAGTGCCCATCGGCTGAAGTCAAGCATCGGGCTGATCAATGAAAAGGCTGCAGATAAGCTGGG
ATCTACCCAGATCGTGAAGATCCTAACTCAGGACACTCCCGAGTTTTTTATAGACCAAGGCCA
TGCCAAGGTGGCCCAACTGATCGTGCTGGAAGTGTTTCCCTCCAGTGAAGCCCTCCGCCCTTT
GTTACCCCTGGGCATCGAAGCCAGCTCGGAAGCTCAGTTTTACACCAAAGGTGACCAACTTAT
ACTCAACTTGAATAACATCAGCTCTGATCGGATCCAGCTGATGAACTCTGGGATTGGCTGGTT
CCAACCTGATGTTCTGAAAAACATCATCACTGAGATCATCCACTCCATCCTGCTGCCGAACCA
GAATGGCAAATTAAGATCTGGGGTCCCAGTGTCATTGGTGAAGGCCTTGGGATTCGAGGCAGC
TGAGTCCTCACTGACCAAGGATGCCCTTGTGCTTACTCCAGCCTCCTTGTGGAAACCCAGCTC
TCCTGTCTCCCAGTGAAGACTTGGATGGCAGCCATCAGGGAAGGCTGGGTCCCAGCTGGGAGT
ATGGGTGTGAGCTCTATAGACCATCCCTCTCTGCAATCAATAAACACTTGCCTGTGAAAAA

78/168

FIGURE 78

MAGPWTFTLLCGLLAATLIQATLSPTAVLILGPKVIKEKLTQELKDNATSILQQLPLLSAMR
EKPAGGIPVLGSLVNTVLKHIIWLKVITANILQLQVKPSANDQELLVKIPLDMVAGFNTPLVK
TIVEFHMTTEAQATIRMDTSASGPTRLVLSDCATSHGSLRIQLLYKLSFLVNALAKQVMNLLV
PSLPNLVKNQLCPVIEASFNGMYADLLQLVKVPISLSIDRLEFDLLYPAIKGDITQLYLGAKL
LDSQGKVTKWFNNSAASLTMPITLDNIPFSLIVSQDVVKAAVAVALSPEEFMVLLDSVLPESAH
RLKSSIGLINEKAADKLGSTQIVKILTQDTPEFFIDQGHAKVAQLIVLEVFPSSSEALRPLFTL
GLEASSEAQFYTKGDQLILNLNNISSDRIQLMNSGIGWFQPDVLKNIITEIIHSILLPNQNGK
LRSGVPVSLVKALGFEEAESSLTKDALVLTTPASLWKPSSPVSQ

Important features of the protein:

Signal peptide:

amino acids 1-21

N-glycosylation sites.

amino acids 48-51, 264-267, 401-404

Glycosaminoglycan attachment site.

amino acids 412-415

LBP / BPI / CETP family proteins.

amino acids 407-457

79/168

FIGURE 79

GAGAGAAGTCAGCCTGGCAGAGAGACTCTGAAATGAGGGATTAGAGGTGTTCAAGGAGCAAGA
GCTTCAGCCTGAAGACAAGGGAGCAGTCCCTGAAGACGCTTCTACTGAGAGGTCTGCCATGGC
CTCTCTTGGCCTCCAACCTTGTGGGCTACATCCTAGGCCTTCTGGGGCTTTTGGGCACACTGGT
TGCCATGCTGCTCCCCAGCTGGAAAACAAGTTCTTATGTCGGTGCCAGCATTGTGACAGCAGT
TGGCTTCTCCAAGGGCCTCTGGATGGAATGTGCCACACACAGCACAGGCATCACCCAGTGTGA
CATCTATAGCACCTTCTGGGCCTGCCCCGCTGACATCCAGGCTGCCCAGGCCATGATGGTGAC
ATCCAGTGCAATCTCCTCCCTGGCCTGCATTATCTCTGTGGTGGGCATGAGATGCACAGTCTT
CTGCCAGGAATCCCGAGCCAAAGACAGAGTGGCGGTAGCAGGTGGAGTCTTTTTTCATCCTTGG
AGGCCTCCTGGGATTCATTCCTGTTGCCTGGAATCTTCATGGGATCCTACGGGACTTCTACTC
ACCACTGGTGCCTGACAGCATGAAATTTGAGATTGGAGAGGCTCTTTACTTGGGCATTATTTT
TTCCCTGTTCTCCCTGATAGCTGGAATCATCCTCTGCTTTTCTGCTCATCCCAGAGAAATCG
CTCCAACCTACTACGATGCCTACCAAGCCCAACCTCTTGCCACAAGGAGCTCTCCAAGGCCTGG
TCAACCTCCCAAAGTCAAGAGTGAGTTCAATTCCTACAGCCTGACAGGGTATGTGTGAAAGAAC
CAGGGGCCAGAGCTGGGGGGTGGCTGGGTCTGTGAAAAACAGTGGACAGCACCCCGAGGGCCA
CAGGTGAGGGACACTACCACTGGATCGTGTCAGAAGGTGCTGCTGAGGATAGACTGACTTTGG
CCATTGGATTGAGCAAAGGCAGAAATGGGGGCTAGTGTAACAGCATGCAGGTTGAATTGCCAA
GGATGCTCGCCATGCCAGCCTTTCTGTTTTCTCACCTTGCTGCTCCCCTGCCCTAAGTCCCC
AACCTCAACTTGAAACCCCATTCCTTAAGCCAGGACTCAGAGGATCCCTTTGCCCTCTGGT
TTACCTGGGACTCCATCCCCAAACCCACTAATCACATCCCACTGACTGACCCTCTGTGATCAA
AGACCCTCTCTCTGGCTGAGGTTGGCTCTTAGCTCATTGCTGGGGATGGGAAGGAGAAGCAGT
GGCTTTTGTGGGCATTGCTCTAACCTACTTCTCAAGCTTCCCTCCAAAGAACTGATTGGCCC
TGGAACCTCCATCCCACTCTTGTTATGACTCCACAGTGTCCAGACTAATTTGTGCATGAACTG
AAATAAAACCATCCTACGGTATCCAGGGAACAGAAAGCAGGATGCAGGATGGGAGGACAGGAA
GGCAGCCTGGGACATTTAAAAAAATA

80/168

FIGURE 80

MASLGLQLVGYYILGLLGLLGTLVAMLLPSWKTSSYVGASIVTAVGFSKGLWMECATHSTGITQ
CDIYSTLLGLPADIQAAQAMMVTSSAIISSLACIISVVGMRCTVFCQESRAKDRVAVAGGVFFI
LGLLGFIPVAWNLHGILRDFYSPLVPDSMKFEIGEALYLGIISSLFSLIAGIILCFSCSSQR
NRSNYYDAYQAQPLATRSSPRPGQPPKVKSEFNSYSLTGYV

Important features of the protein:

Signal peptide:

amino acids 1-24

Transmembrane domains:

amino acids 82-102, 117-140, 163-182

N-glycosylation site.

amino acids 190-193

PMP-22 / EMP / MP20 family proteins.

amino acids 46-59

81/168

FIGURE 81

CCCACGCGTCCGCGCCTCTCCCTTCTGCTGGACCTTCCTTCGTCTCTCCATCTCTCCCTCCTT
TCCCCGCGTTCTCTTTCCACCTTTCTCTTCTTCCCACCTTAGACCTCCCTTCCTGCCCTCCTT
TCCTGCCCACCGCTGCTTCCTGGCCCTTCTCCGACCCCGCTCTAGCAGCAGACCTCCTGGGGT
CTGTGGGTTGATCTGTGGCCCCCTGTGCCTCCGTGTCCTTTTCGTCTCCCTTCCTCCCGACTCC
GCTCCCGGACCAGCGGCCTGACCCTGGGGAAAGGATGGTTCCCGAGGTGAGGGTCCTCTCCTC
CTTGCTGGGACTCGCGCTGCTCTGGTTCCCCCTGGACTCCCACGCTCGAGCCCGCCCAGACAT
GTCTGCCTTTTCCATGGGAAGAGATACTCCCCCGGCGAGAGCTGGCACCCCTACTTGGAGCC
ACAAGGCCTGATGTACTGCCTGCGCTGTACCTGCTCAGAGGGCGCCCATGTGAGTTGTTACCG
CCTCCACTGTCCGCCTGTCCACTGCCCCCAGCCTGTGACGGAGCCACAGCAATGCTGTCCCAA
GTGTGTGGAACCTCACACTCCCTCTGGACTCCGGGCCCCACCAAAGTCCTGCCAGCACAAACGG
GACCATGTACCAACACGGAGAGATCTTCAGTGCCCATGAGCTGTTCCCTCCCGCCTGCCCAA
CCAGTGTGTCCTCTGCAGCTGCACAGAGGGCCAGATCTACTGCGGCCTCACAACCTGCCCCGA
ACCAGGCTGCCCAGCACCCCTCCCACTGCCAGACTCCTGCTGCCAAGCCTGCAAAGATGAGGC
AAGTGAGCAATCGGATGAAGAGGACAGTGTGCAGTCGCTCCATGGGGTGAGACATCCTCAGGA
TCCATGTTCCAGTGATGCTGGGAGAAAGAGAGGGCCCGGGCACCCCAAGCCCCACTGGCCTCAG
CGCCCCCTCTGAGCTTCATCCCTCGCCACTTCAGACCCAAGGGAGCAGGCAGCACAACTGTCAA
GATCGTCCTGAAGGAGAAACATAAGAAAGCCTGTGTGCATGGCGGGAAGACGTACTCCACGG
GGAGGTGTGGCACCCGGCCTTCCGTGCCTTCGGCCCCCTTGCCCTGCATCCTATGCACCTGTGA
GGATGGCCGCCAGGACTGCCAGCGTGTGACCTGTCCCACCGAGTACCCCTGCCGTACCCCGA
GAAAGTGGCTGGGAAGTGCTGCAAGATTTGCCCAGAGGACAAAGCAGACCCTGGCCACAGTGA
GATCAGTTCTACCAGGTGTCCCAAGGCACCGGGCCGGGTCCTCGTCCACACATCGGTATCCCC
AAGCCCAGACAACCTGCGTCGCTTTGCCCTGGAACACGAGGCCTCGGACTTGGTGGAGATCTA
CCTCTGGAAGCTGGTAAAAGATGAGGAACTGAGGCTCAGAGAGGTGAAGTACCTGGCCCAAG
GCCACACAGCCAGAATCTTCCACTTGA CT CAGATCAAGAAAGTCAGGAAGCAAGACTTCCAGA
AAGAGGCACAGCACTTCCGACTGCTCGCTGGCCCCCACGAAGGTC ACT GGAACGTCTTCCTAG
CCCAGACCCTGGAGCTGAAGGTCACGGCCAGTCCAGACAAAGTGACCAAGACATAACAAAGAC
CTAACAGTTGCAGATATGAGCTGTATAATTGTTGTTATTATATATTAATAATAAGAAGTTGC
ATTACCCTCAAAAAAAAAAAAAAAAAAAAAA

82/168

FIGURE 82

MVPEVRVLSSLLGLALLWFPLDSHARARPD MFCLFHGKRYSPGESWHPYLEPQGLMYCLRCTC
SEGAHVSCYRLHCPPVHCPQPVTEPQQCCPKCVEPHTPSGLRAPPKSCQHNGTMYQHGEIFSA
HELFPSRLPNQCVLCSCTEGQIYCGLTTCPEPGCPAPLPLPDSCCQACKDEASEQSDEEDSVQ
SLHGVRHPQDPCSSDAGRKRGP GTPAPTGLSAPLSFIPRHFRPKGAGSTTVKIVLKEKHKKAC
VHGGKTYSHGEVWHPAFRAFGPLPCILCTCEDGRQDCQRVTCPT EYPCRHPK VAGKCKICP
EDKADPGHSEISSTRCPKAPGRVLVHTSVSPSPDNLRRFALEHEASDLVEIYLWKLVKDEETE
AQRGEVPGPRPHSQNLPLDSDQESQEARLPERGTALPTARWP PRRSLERLPSPDPGAEGHGQS
RQSDQDITKT

Signal peptide:

amino acids 1-25

83/168

FIGURE 83

GACAGCTGTGTCTCGATGGAGTAGACTCTCAGAACAGCGCAGTTTGCCCTCCGCTCACGCAGA
GCCTCTCCGTGGCTTCCGCACCTTGAGCATTAGGCCAGTTCTCCTCTTCTCTCTAATCCATCC
GTCACCTCTCCTGTCATCCGTTTCCATGCCGTGAGGTCCATTACAGAACACATCCATGGGCTC
TCATGCTCAGTTTGGTTCTGAGTCTCCTCAAGCTGGGATCAGGGCAGTGGCAGGTGTTTGGGC
CAGACAAGCCTGTCCAGGCCTTGGTGGGGGAGGACGCAGCATTCTCCTGTTTCCTGTCTCCTA
AGACCAATGCAGAGGCCATGGAAGTGCGGTTCTTCAGGGGCCAGTTCTCTAGCGTGGTCCACC
TCTACAGGGACGGGAAGGACCAGCCATTTATGCAGATGCCACAGTATCAAGGCAGGACAAAAC
TGGTGAAGGATTCTATTGCGGAGGGGGCGCATCTCTCTGAGGCTGGAAAACATTACTGTGTTGG
ATGCTGGCCTCTATGGGTGCAGGATTAGTTCCCAGTCTTACTACCAGAAGGCCATCTGGGAGC
TACAGGTGTCAGCACTGGGCTCAGTTCCTCTCATTTCATCACGGGATATGTTGATAGAGACA
TCCAGCTACTCTGTCAGTCCTCGGGCTGGTTCCCCCGGCCACAGCGAAGTGGAAGGTCCAC
AAGGACAGGATTTGTCCACAGACTCCAGGACAAACAGAGACATGCATGGCCTGTTTGATGTGG
AGATCTCTCTGACCGTCCAAGAGAACGCCGGGAGCATATCCTGTTCCATGCGGCATGCTCATC
TGAGCCGAGAGGTGGAATCCAGGGTACAGATAGGAGATACCTTTTTTCGAGCCTATATCGTGGC
ACCTGGCTACCAAAGTACTGGGAATACTCTGCTGTGGCCTATTTTTTTGGCATTGTTGGACTGA
AGATTTTCTTCTCCAAATTCCAGTGGAATCCAGGCGGAAGTGGACTGGAGAAGAAAGCACG
GACAGGCAGAATTGAGAGACGCCCCGGAACACGCAGTGGAGGTGACTCTGGATCCAGAGACGG
CTCACCCGAAGCTCTGCGTTTCTGATCTGAAAAGTGTAAACCATAGAAAAGCTCCCCAGGAGG
TGCCTCACTCTGAGAAGAGATTTACAAGGAAGAGTGTGGTGGCTTCTCAGAGTTTCCAAGCAG
GGAAACATTACTGGGAGGTGGACGGAGGACACAATAAAAGGTGGCGCGTGGGAGTGTGCCGGG
ATGATGTGGACAGGAGGAAGGAGTACGTGACTTTGTCTCCCGATCATGGGTACTGGGTCTCA
GACTGAATGGAGAACATTTGTATTTACATTAAATCCCCGTTTTTATCAGCGTCTTCCCCAGGA
CCCCACCTACAAAATAGGGGTCTTCCTGGACTATGAGTGTGGGACCATCTCCTTCTTCAACA
TAAATGACCAGTCCCTTATTTATACCCTGACATGTCGGTTTGAAGGCTTATTGAGGCCCTACA
TTGAGTATCCGTCCTATAATGAGCAAAATGGAAGTCCCATAGTCATCTGCCAGTCACCCAGG
AATCAGAGAAAGAGGCCTCTTGGCAAAGGGCCTCTGCAATCCCAGAGACAAGCAACAGTGAGT
CCTCCTCACAGGCAACCACGCCCTTCCTCCCCAGGGGTGAAATGTAGGATGAATCACATCCCA
CATTCTTCTTTAGGGATATTAAGGTCTCTCTCCCAGATCCAAAGTCCCGCAGCAGCCGGCCAA
GGTGGCTTCCAGATGAAGGGGGACTGGCCTGTCCACATGGGAGTCAGGTGTCATGGCTGCCCT
GAGCTGGGAGGGAAGAAGGCTGACATTACATTTAGTTTGCTCTCACTCCATCTGGCTAAGTGA
TCTTGAAATACCACCTCTCAGGTGAAGAACCGTCAGGAATTCCCATCTCACAGGCTGTGGTGT
AGATTAAGTAGACAAGGAATGTGAATAATGCTTAGATCTTATTGATGACAGAGTGTATCCTAA
TGGTTTGTTTATTATATTACACTTTCAGTAAAAAAA

84/168

FIGURE 84

MALMLSLVLSLLKLGSGQWQVFGPDKPVQALVGEDAAFSCLSPKTNAEAMEVRFFRGQFSSV
VHLYRDGKDQPFMQMPQYQGR TKLVKDSIAEGRISLRLENITVLDAGLYGCRISSQSYQKAI
WELQVSALGSVPLISITGYVDRDIQLLCQSSGWFPRTAKWKGPQGQDLSTDSRTNRDMHGLF
DVEISLTVQENAGSISCSMRHAHLSREVESRVQIGDTFFEPISWHLATKVLGILCCGLFFGIV
GLKIFFSKFQWKIQAE LDWRRKHGQAE LR DARK HAVE VT LDPETAHPKLCVSDLKTVTHR KAP
QEVPHSEKRFRTRKSVVASQS FQAGKHYWEVDGGHNKRWRVGVC RDDVDRRKEYVTLS PDHGYW
VLR L NGEHLYFTLNPRFISVFPRT PPTKIGVFLDYECGTISFFNINDQS LIYTLTCRFEGLLR
PYIEYPSYNEQNGTPIVICPVTQESEKEASWQRASAI PETSNSESSSQATTPFLPRGEM

Signal peptide:

amino acids 1-17

Transmembrane domain:

amino acids 239-255

85/168

FIGURE 85

AACAGACGTTCCCTCGCGGCCCTGGCACCTCTAACCCCAGACATGCTGCTGCTGCTGCTGCTGCCC
CTGCTCTGGGGGAGGGAGAGGGCGGAAGGACAGACAAGTAAACTGCTGACGATGCAGAGTTCC
GTGACGGTGCAGGAAGGCCTGTGTGTCCATGTGCCCTGCTCCTTCTCCTACCCCTCGCATGGC
TGGATTTACCCTGGCCCAGTAGTTCATGGCTACTGGTTCCGGGAAGGGGCCAATACAGACCAG
GATGCTCCAGTGGCCACAAACAACCCAGCTCGGGCAGTGTGGGAGGAGACTCGGGACCGATTC
CACCTCCTTGGGGACCCACATACCAAGAATTGCACCCTGAGCATCAGAGATGCCAGAAGAAGT
GATGCGGGGAGATACTTCTTTCGTATGGAGAAAGGAAGTATAAAATGGAATTATAAACATCAC
CGGCTCTCTGTGAATGTGACAGCCTTGACCCACAGGCCCAACATCCTCATCCCAGGCACCCTG
GAGTCCGGCTGCCCCCAGAATCTGACCTGCTCTGTGCCCTGGGCCTGTGAGCAGGGGACACCC
CCTATGATCTCCTGGATAGGGACCTCCGTGTCCCCCCTGGACCCCTCCACCACCCGCTCCTCG
GTGCTCACCCCTCATCCCACAGCCCCAGGACCATGGCACCAGCCTCACCTGTCAGGTGACCTTC
CCTGGGGCCAGCGTGACCACGAACAAGACCGTCCATCTCAACGTGTCCTACCCGCCTCAGAAC
TTGACCATGACTGTCTTCCAAGGAGACGGCACAGTATCCACAGTCTTGGGAAATGGCTCATCT
CTGTCACTCCCAGAGGGGCCAGTCTCTGCGCCTGGTCTGTGCAGTTGATGCAGTTGACAGCAAT
CCCCCTGCCAGGCTGAGCCTGAGCTGGAGAGGCCTGACCCTGTGCCCCTCACAGCCCTCAAAC
CCGGGGGTGCTGGAGCTGCCTTGGGTGCACCTGAGGGATGCAGCTGAATTACCTGCAGAGCT
CAGAACCCTCTCGGCTCTCAGCAGGTCTACCTGAACGTCTCCCTGCAGAGCAAAGCCACATCA
GGAGTGACTCAGGGGGTGGTCGGGGGAGCTGGAGCCACAGCCCTGGTCTTCCTGTCCTTCTGC
GTCATCTTCGTTGTAGTGAGGTCCTGCAGGAAGAAATCGGCAAGGCCAGCAGCGGGCGTGGA
GATACGGGCATAGAGGATGCAAACGCTGTCAGGGGTTCAGCCTCTCAGGGGGCCCCTGACTGAA
CCTTGGGCAGAAGACAGTCCCCCAGACCAGCCTCCCCCAGCTTCTGCCCCGCTCCTCAGTGGGG
GAAGGAGAGCTCCAGTATGCATCCCTCAGCTTCCAGATGGTGAAGCCTTGGGACTCGCGGGGA
CAGGAGGCCACTGACACCGAGTACTCGGAGATCAAGATCCACAGATTGAGAAACTGCAGAGACT
CACCCCTGATTGAGGGATCACAGCCCCTCCAGGCAAGGGAGAAGTCAGAGGCTGATTCTTGTAG
AATTAACAGCCCTCAACGTGATGAGCTATGATAACACTATGAATTATGTGCAGAGTGAAAAGC
ACACAGGCTTTAGAGTCAAAGTATCTCAAACCTGAATCCACACTGTGCCCTCCCTTTTATTTT
TTTAACTAAAAGACAGACAAATTCCTA

86/168

FIGURE 86

MLLLLLPLLWGRERAEGQTSKLLTMQSSVTVQEGLCVHVPCSFSPSHGWIYPGPVVHGYWFR
EGANTDQDAPVATNNPARAVWEETRDREHLLGDPHTKNCTLSIRDARRSDAGRYFFRMEKGS I
KWNYPKHRLSVNVTALTHRPNILIPGTLESGCPQNLTCVWPWACEQGTPPMISWIGTSVSPLD
PSTTRSSVLTLPQPQDHGTS LTCQVTFPGASVTTNKTVHLNVSYPPQNLTM TVFQGDGTVST
VLGNGSSL LPEGQSLRLVCAVDAVD SNPPARLSLSWRGLTLCPSQPSNPGVLELPWVHLRDA
AEFTCRAQNPLGSQQVYLNVSLSKATSGVTQGVVGGAGATALVFLSFCVIFVVVRSCRK KSA
RPAAGVGDTGIEDANAVRG SASQGPLTEPWAEDSPPDQPPPASARSSVGE GELQYASLSFQMV
KPWDSRGQEATDTEYSEIKIHR

Signal peptide:

amino acids 1-15

Transmembrane domain:

amino acids 351-370

87/168

FIGURE 87

AGAAAGCTGCACTCTGTTGAGCTCCAGGGCGCAGTGGAGGGAGGGAGTGAAGGAGCTCTCTGT
ACCCAAGGAAAGTGCAGCTGAGACTCAGACAAGATTACAATGAACCAACTCAGCTTCCTGCTG
TTTCTCATAGCGACCACCAGAGGATGGAGTACAGATGAGGCTAATACTTACTTCAAGGAATGG
ACCTGTTCTTCGTCTCCATCTCTGCCCAGAAGCTGCAAGGAAATCAAAGACGAATGTCCTAGT
GCATTTGATGGCCTGTATTTTCTCCGCACTGAGAATGGTGTATCTACCAGACCTTCTGTGAC
ATGACCTCTGGGGGTGGCGGCTGGACCCTGGTGGCCAGCGTGCATGAGAATGACATGCGTGGG
AAGTGCACGGTGGGCGATCGCTGGTCCAGTCAGCAGGGCAGCAAAGCAGACTACCCAGAGGGG
GACGGCAACTGGGCCAACTACAACACCTTTGGATCTGCAGAGGCGGCCACGAGCGATGACTAC
AAGAACCCTGGCTACTACGACATCCAGGCCAAGGACCTGGGCATCTGGCACGTGCCCAATAAG
TCCCCCATGCAGCACTGGAGAAACAGCTCCCTGCTGAGGTACCGCACGGACACTGGCTTCCTC
CAGACACTGGGACATAATCTGTTTGGCATCTACCAGAAATATCCAGTGAAATATGGAGAAGGA
AAGTGTTGGACTGACAACGGCCCCGGTGATCCCTGTGGTCTATGATTTTGGCGACGCCCAGAAA
ACAGCATCTTATTACTCACCTATGGCCAGCGGGAATTCAGTGCGGGATTTGTTTCAGTTCAGG
GTATTTAATAACGAGAGAGCAGCCAACGCCTTGTGTGCTGGAATGAGGGTCACCGGATGTAAC
ACTGAGCATCACTGCATTGGTGGAGGAGGATACTTCCAGAGGCCAGTCCCCAGCAGTGTGGA
GATTTTCTGGTTTTGATTGGAGTGGATATGGAACCTCATGTTGGTTACAGCAGCAGCCGTGAG
ATAACTGAGGCAGCTGTGCTTCTATTCTATCGTTGAGAGTTTTGTGGGAGGGAACCCAGACCT
CTCCTCCCAACCATGAGATCCCAAGGATGGAGAACAACCTTACCCAGTAGCTAGAATGTTAATG
GCAGAAGAGAAAACAATAAATCATATTGACTCAAGAAAAAA

88/168

FIGURE 88

MNQLSFLLFLIATTRGWSTDEANTYFKEWTCSSSPSLPRSCKEIKDECPSAFDGLYFLRTENG
VIYQTFCDMTSGGGGWTLVASVHENDMRGKCTVGDRWSSQQGSKADYPEGDGNWANYNTFGSA
EAATSDDYKNPGYYDIQAKDLGIWHVFNKSPMQHWRNSSLLRYRTDTGFLQTLGHNLFGLIYQK
YPVKYGEKGCWTDNGPVI PVVYDFGDAQKTASYYSPIYGQREFTAGFVQFRVFNNERAANALCA
GMRVTGCNTEHHCIGGGGYFPEASPOQCGDFSGFDWSGYGTHVGYSSSREITEAAVLLFYR

Important features:**Signal peptide:**

amino acids 1-16

N-glycosylation site.

amino acids 163-167

Glycosaminoglycan attachment sites.

amino acids 74-78, 289-293

N-myristoylation sites.

amino acids 76-82, 115-121, 124-130, 253-259, 292-298

89/168

FIGURE 89

CTAGATTTGTCGGCTTGCGGGGAGACTTCAGGAGTCGCTGTCTCTGAACTTCCAGCCTCAGAG
ACCGCCGCCCTTGTCCCCGAGGGCCATGGGCCGGGTCTCAGGGCTTGTGCCCTCTCGCTTCCT
GACGCTCCTGGCGCATCTGGTGGTCGTCATCACCTTATTCTGGTCCCGGGACAGCAACATACA
GGCCTGCCTGCCTCTCACGTTACCCCCGAGGAGTATGACAAGCAGGACATTCAGCTGGTGGC
CGCGCTCTCTGTACCCCTGGGCCTCTTTGCAGTGGAGCTGGCCGGTTTCCTCTCAGGAGTCTC
CATGTTCAACAGCACCCAGAGCCTCATCTCCATTGGGGCTCACTGTAGTGCATCCGTGGCCCT
GTCCTTCTTCATATTCGAGCGTTGGGAGTGCACCTACGTATTGGTACATTTTTGTCTTCTGCAG
TGCCCTTCCAGCTGTCACCTGAAATGGCTTTATTCGTCACCGTCTTTGGGCTGAAAAAGAAACC
CTTCTGATTACCTTCATGACGGGAACCTAAGGACGAAGCCTACAGGGGCAAGGGCCGCTTCGT
ATTCCTGGAAGAAGGAAGGCATAGGCTTCGGTTTTCCCCTCGGAAACTGCTTCTGCTGGAGGA
TATGTGTTGGAATAATTACGTCTTGAGTCTGGGATTATCCGCATTGTATTTAGTGCTTTGTAA
TAAATATGTTTTGTAGTAACATTAAGACTTATATACAGTTTTAGGGGACAATTAAAAAAA
AAA

WO 01/16318

PCT/US00/23328

90/168

FIGURE 90

MGRVSGLVPSRELTLLAHLVVVITLFWSRDSNIQACLPLTFTPEEYDKQDIQLVAALSVTLGL
FAVELAGFLSGVSMFNSTQSLISIGAHCSASVALSFFIFERWECTTYWYIFVFCSALPAVTEM
ALFVTVFGLKKKPF

Transmembrane domain:

amino acids 12-28 (type II), 51-66, 107-124

91/168

FIGURE 91

CTGGGACCCCGAAAAGAGAAGGGGAGAGCGAGGGGACGAGAGCGGAGGAGGAAGATGCAACTG
ACTCGCTGCTGCTTCGTGTTCTGGTGCAGGGTAGCCTCTATCTGGTCATCTGTGGCCAGGAT
GATGGTCCTCCCGGCTCAGAGGACCCTGAGCGTGATGACCACGAGGGGCCAGCCCCGGCCCCGG
GTGCCTCGGAAGCGGGGCCACATCTCACCTAAGTCCCGCCCCATGGCCAATTCCACTCTCCTA
GGGCTGCTGGCCCCGCCTGGGGAGGCTTGGGGCATTCTTGGGCAGCCCCCAACCGCCCGAAC
CACAGCCCCCCCACCCTCAGCCAAGGTGAAGAAAATCTTTGGCTGGGGCGACTTCTACTCCAAC
ATCAAGACGGTGGCCCTGAACCTGCTCGTCACAGGGAAGATTGTGGACCATGGCAATGGGACC
TTCAGCGTCCACTTCCAACACAATGCCACAGGCCAGGGAAACATCTCCATCAGCCTCGTGCCC
CCCAGTAAAGCTGTAGAGTTCCACCAGGAACAGCAGATCTTCATCGAAGCCAAGGCCTCCAAA
ATCTTCAACTGCCGGATGGAGTGGGAGAAGGTAGAACGGGGCCGCCGGACCTCGCTTTGCACC
CACGACCCAGCCAAGATCTGCTCCCGAGACCACGCTCAGAGCTCAGCCACCTGGAGCTGCTCC
CAGCCCTTCAAAGTCGTCTGTGTCTACATCGCCTTCTACAGCACGGACTATCGGCTGGTCCAG
AAGGTGTGCCCAGATTACAACCTACCATAGTGATACCCCCTACTACCCATCTGGGTGACCCCGGG
GCAGGCCACAGAGGCCAGGCCAGGGCTGGAAGGACAGGCCTGCCCATGCAGGAGACCATCTGG
ACACCGGGCAGGGAAGGGGTGGGGCCTCAGGCAGGGAGGGGGGTGGAGACGAGGAGATGCCAA
GTGGGGCCAGGGCCAAGTCTCAAGTGGCAGAGAAAGGGTCCCAAGTGCTGGTCCCAACCTGAA
GCTGTGGAGTGACTAGATCACAGGAGCACTGGAGGAGGAGTGGGCTCTCTGTGCAGCCTCACA
GGGCTTTGCCACGGAGCCACAGAGAGATGCTGGGTCCCCGAGGCCTGTGGGCAGGCCGATCAG
TGTGGCCCCAGATCAAGTCATGGGAGGAAGCTAAGCCCTTGGTTCTTGCCATCCTGAGGAAAG
ATAGCAACAGGGAGGGGGGAGATTTTCATCAGTGTGGACAGCCTGTCAACTTAGGATGGATGGCT
GAGAGGGCTTCCTAGGAGCCAGTCAGCAGGGTGGGGTGGGGCCAGAGGAGCTCTCCAGCCCTG
CCTAGTGGGCGCCCTGAGCCCCTTGTCGTGTGCTGAGCATGGCATGAGGCTGAAGTGGCAACC
CTGGGGTCTTTGATGTCTTGACAGATTGACCATCTGTCTCCAGCCAGGCCACCCCTTTCCAAA
ATTCCCTCTTCTGCCAGTACTCCCCCTGTACCACCCATTGCTGATGGCACACCCATCCTTAAG
CTAAGACAGGACGATTGTGGTCCTCCCACACTAAGGCCACAGCCCATCCGCGTGCTGTGTGTC
CCTCTTCCACCCCAACCCCTGCTGGCTCCTCTGGGAGCATCCATGTCCCGGAGAGGGGTCCCT
CAACAGTCAGCCTCACCTGTCAGACCGGGGTCTCCCGGATCTGGATGGCGCCGCCCTCTCAG
CAGCGGGCACGGGTGGGGCGGGGCCGGGCCGAGAGCATGTGCTGGATCTGTTCTGTGTGTCT
GTCTGTGGGTGGGGGGAGGGGAGGGAAGTCTTGTGAAACCGCTGATTGCTGACTTTTGTGTGA
AGAATCGTGTTCTTGGAGCAGGAAATAAAGCTTGCCCCGGGGCA

92/168

FIGURE 92

MQLTRCCFVFLVQGSlyLVICGQDDGPPGSEDPERDDHEGQPRPRVPRKRGHISPKSRPMANS
TLLGLLAPPGEAWGILGQPPNRPNHSPPPSAKVKKIFGWGDFYSNIKTVALNLLVTGKIVDHG
NGTFSVHFQHNATGQGNISISLVPPSKAVEFHQEQQIFIEAKASKIFNCRMWEKVERGRRTS
LCTHDPAKICSRDHAQSSATWSCSQPFKVVCVYIAFYSTDYRLVQKVCPDYNHSDTPYYPSG

Important features of the protein:

Signal peptide:

amino acids 1-14

N-glycosylation sites.

amino acids 62-65, 127-130, 137-140, 143-146

2-oxo acid dehydrogenases acyltransferase

amino acids 61-71

93/168

FIGURE 93

CGGTGGCCATGACTGCGGCCGTGTTCTTCGGCTGCGCCTTCATTGCCTTCGGGCCTGCGCTCG
CCCTTTATGTCTTCACCATCGCCATCGAGCCGTTGCGTATCATCTTCCTCATCGCCGGAGCTT
TCTTCTGGTTGGTGTCTCTACTGATTTTCGTCCCTTGTTTGGTTCATGGCAAGAGTCATTATTG
ACAACAAAGATGGACCAACACAGAAATATCTGCTGATCTTTGGAGCGTTTGTCTCTGTCTATA
TCCAAGAAATGTTCCGATTTGCATATTATAAACTCTTAAAAAAGCCAGTGAAGGTTTGAAGA
GTATAAACCAGGTGAGACAGCACCTCTATGCGACTGCTGGCCTATGTTTCTGGCTTGGGCT
TTGGAATCATGAGTGGAGTATTTTCCTTTGTGAATACCCTATCTGACTCCTTGGGGCCAGGCA
CAGTGGGCATTCATGGAGATTCTCCTCAATTCTTCCTTTATTCAGCTTTCATGACGCTGGTCA
TTATCTTGCTGCATGTATTCTGGGGCATTGTATTTTTTTGATGGCTGTGAGAAGAAAAAGTGGG
GCATCCTCCTTATCGTTCTCCTGACCCACCTGCTGGTGTGAGCCAGACCTTCATAAGTTCTT
ATTATGGAATAAACCTGGCGTCAGCATTATAATCCTGGTGCTCATGGGCACCTGGGCATTCT
TAGCTGCGGGAGGCAGCTGCCGAAGCCTGAAACTCTGCCTGCTCTGCCAAGACAAGAACTTTC
TTCTTTACAACCAGCGCTCCAGATTAACCTCAGGGAACCAGCACTTCCCAAACCGCAGACTACA
TCTTTAGAGGAAGCACAACTGTGCCTTTTTCTGAAAATCCCTTTTTCTGGTGGAATTGAGAAA
GAAATAAACTATGCAGATA

94/168

FIGURE 94

MTAAVFFGCAFIAGFPALALYVFTIAIEPLRIIFLIAGAFFWLVSLLISSLVWFMARVIIDNK
DGPTQKYLLIFGAFVSVYIQEMFRFAYYKLLKKASEGLKSINPGETAPSMRLLAYVSGLGFGI
MSGVFSFVNTLSDSLGP GTVGIHG DSPQFFLYSAFMTLVIILLHVFWGIVFFDGCEKKKWGIL
LIVLLTHLLVSAQTFISSYYGINLASAFIILVLMGTWAF LAAGGSCRS LKLCLLCQDKNFLLY
NQRSR

Important features of the protein:

Signal peptide:

amino acids 1-19

Transmembrane domains:

amino acids 32-51, 119-138, 152-169, 216-235

Glycosaminoglycan attachment site:

amino acids 120-123

Sodium:neurotransmitter symporter family protein

amino acids 31-65

[illegible]

96/168

FIGURE 96

MRSTILLFCLLGSTRSLPQLKPALGLPPTKLAPDQGTLPNQQQSNQVFPSLSLIPLTQM
LTLGPDLHLLNPAAGMTPGTQTHPLTLGGLNVQQQLHPHVLPIFVTQLGAQGTILSSEE
LPQIFTSLSLIHSLEFPGGILPTSQAGANPDVQDGSLPAGGAGVNPATQGTPAGRLPTPSG
TDDDFAVTTPAGIQRSTHAIEEATTESANGIQ

Signal peptide:

amino acids 1-16

97/168

FIGURE 97

GCTCAAGTGCCCTGCCTTGCCCCACCCAGCCCAGCCTGGCCAGAGCCCCCTGGAGAAGGAGCT
CTCTTCTTGCTTGGCAGCTGGACCAAGGGAGCCAGTCTTGGGCGCTGGAGGGCCTGTCCTGAC
CATGGTCCCTGCCTGGCTGTGGCTGCTTTGTGTCTCCGTCCCCCAGGCTCTCCCCAAGGCCCA
GCCTGCAGAGCTGTCTGTGGAAGTTCCAGAAAACCTATGGTGGAAATTTCCCTTTATACCTGAC
CAAGTTGCCGCTGCCCCGTGAGGGGGCTGAAGGCCAGATCGTGCTGTCAGGGGACTCAGGCAA
GGCAACTGAGGGCCCCATTTGCTATGGATCCAGATTCTGGCTTCCTGCTGGTGACCAGGGCCCT
GGACCGAGAGGAGCAGGCAGAGTACCAGCTACAGGTCACCCTGGAGATGCAGGATGGACATGT
CTTGTGGGGTCCACAGCCTGTGCTTGTGCACGTGAAGGATGAGAATGACCAGGTGCCCCATTT
CTCTCAAGCCATCTACAGAGCTCGGCTGAGCCGGGGTACCAGGCCTGGCATCCCCTTCCTCTT
CCTTGAGGCTTCAGACCGGGATGAGCCAGGCACAGCCAACTCGGATCTTCGATTCCACATCCT
GAGCCAGGCTCCAGCCCAGCCTTCCCCAGACATGTTCCAGCTGGAGCCTCGGCTGGGGGCTCT
GGCCCTCAGCCCCAAGGGGAGCACCAGCCTTGACCACGCCCTGGAGAGGACCTACCAGCTGTT
GGTACAGGTCAAGGACATGGGTGACCAGGCCTCAGGCCACCAGGCCACTGCCACCGTGGAAGT
CTCCATCATAGAGAGCACCTGGGTGTCCCTAGAGCCTATCCACCTGGCAGAGAATCTCAAAGT
CCTATACCCGCACCACATGGCCCAGGTACACTGGAGTGGGGGTGATGTGCACTATCACCTGGA
GAGCCATCCCCCGGGACCCTTTGAAGTGAATGCAGAGGGAAACCTCTACGTGACCAGAGAGCT
GGACAGAGAAGCCCAGGCTGAGTACCTGCTCCAGGTGCGGGCTCAGAATTCCCATGGCGAGGA
CTATGCGGCCCCCTCTGGAGCTGCACGTGCTGGTGATGGATGAGAATGACAACGTGCCTATCTG
CCCTCCCCGTGACCCACAGTCAGCATCCCTGAGCTCAGTCCACCAGGTACTGAAGTGACTAG
ACTGTCAGCAGAGGATGCAGATGCCCCCGGCTCCCCCAATTCCACGTTGTGTATCAGCTCCT
GAGCCCTGAGCCTGAGGATGGGGTAGAGGGGAGAGCCTTCCAGGTGGACCCCACTTCAGGCAG
TGTGACGCTGGGGGTGCTCCCACTCCGAGCAGGCCAGAACATCCTGCTTCTGGTGCTGGCCAT
GGACCTGGCAGGCGCAGAGGGTGGCTTCAGCAGCACGTGTGAAGTCGAAGTCGCAGTCACAGA
TATCAATGATCACGCCCCCTGAGTTCATCACTTCCCAGATTGGGCCTATAAGCCTCCCTGAGGA
TGTGGAGCCCGGGACTCTGGTGGCCATGCTAACAGCCATTGATGCTGACCTCGAGCCCGCCTT
CCGCCTCATGGATTTTGCCATTGAGAGGGGAGACACAGAAGGGGACTTTTGGCCTGGATTGGGA
GCCAGACTCTGGGCATGTTAGACTCAGACTCTGCAAGAACCTCAGTTATGAGGCAGCTCCAAG
TCATGAGGTGGTGGTGGTGGTGCAGAGTGTGGCGAAGCTGGTGGGGCCAGGCCCAAGGCCCTGG
AGCCACCGCCACGGTGACTGTGCTAGTGGAGAGAGTGATGCCACCCCCCAAGTTGGACCAGGA
GAGCTACGAGGCCAGTGTCCCCATCAGTGCCCCAGCCGGCTCTTTCCTGCTGACCATCCAGCC
CTCCGACCCCATCAGCCGAACCCTCAGGTTCTCCCTAGTCAATGACTCAGAGGGCTGGCTCTG
CATTGAGAAATTCTCCGGGGAGGTGCACACCGCCCAGTCCCTGCAGGGCGCCCAGCCTGGGGA
CACCTACACGGTGCTTGTGGAGGCCCAGGATACAGCCCTGACTCTTGCCCCTGTGCCCTCCCA
ATACCTCTGCACACCCCGCCAAGACCATGGCTTGATCGTGAGTGGACCCAGCAAGGACCCCGA
TCTGGCCAGTGGGCACGGTCCCTACAGCTTCACCCTTGGTCCCAACCCACGGTGCAACGGGA
TTGGCGCCTCCAGACTCTCAATGGTTCCCATGCCTACCTCACCTTGGCCCTGCATTGGGTGGA
GCCACGTGAACACATAATCCCCGTGGTGGTCAGCCACAATGCCCAGATGTGGCAGCTCCTGGT
TCGAGTGATCGTGTGTCGCTGCAACGTGGAGGGGCGAGTGCATGCGCAAGGTGGGCCGCATGAA
GGGCATGCCACGAAGCTGTGCGCAGTGGGCATCCTTGTAGGCACCCTGGTAGCAATAGGAAT
CTTCCTCATCCTCATTTTCACCCACTGGACCATGTCAAGGAAGAAGGACCCGGATCAACCAGC
AGACAGCGTGCCCCCTGAAGGCGACTGTCTGAATGGCCCAGGCAGCTCTAGCTGGGAGCTTGGC
CTCTGGCTCCATCTGAGTCCCCTGGGAGAGAGCCCAGCACCCCAAGATCCAGCAGGGGGACAGGA
CAGAGTAGAAGCCCCTCCATCTGCCCTGGGGTGGAGGCACCATCACCATCACAGGCATGTCT
GCAGAGCCTGGACACCAACTTTATGGACTGCCCATGGGAGTGCTCCAAATGTCAGGGTGTTTG
CCCAATAATAAAGCCCCAGAGAACTGGGCTGGGCCCTATGGGAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAG

98/168

FIGURE 98

MVPAWLWLLCVSVPQALPKAQPAELSVEVPENYGGNFPLYLTKLPLPREGAEGQIVLSGDSGK
ATEGPFAMDPDSGFLLVTRALDREEQAQYQLQVTLEMQDGHVLWGPQPVLVHVKDENDQVPHF
SQAIYRARLSRGTRPGIPFLFLEASDRDEPGTANSDLRFHILSQAPAQPSDFMFQLEPRLGAL
ALSPKGSTSLDHALERTYQLLVQVKMDQASGHQATATVEVSIESTWVSLEPIHLAENLKV
LYPHHMAQVHWSSGGDVHYHLESHPPGPFVNAEGNLYVTRELDREAQAEYLLQVRAQNSHGED
YAAPLELHVLVMDENDNVPICPPRDPTVSIPELSPPGTEVTRLAEDADAPGSPNSHVYQLL
SPEPEDGVEGRAQVDPTSGSVTLGVLPLRAGQNILLVLAMDLAGAEGGFSSTCEVEVAVTD
INDHAPEFITSQIGPISLPEDVEPGTLVAMLTADLEPAFRLMDFAIERGDTEGTFGLDWE
PDSGHVRLRLCKNLSYEAAPSHEVVVVVQSVAKLVGPGPGGATATVTVLVERVMPPPKLDQE
SYEASVPISAPAGSFLTIQPSDPISRTLRFSLVNDSEGWLCIEKFSGEVHTAQSLQGAQPGD
TYTVLVEAQDTALTAPVPSQYLCTPRQDHGLIVSGPSKDPDLASGHGPYSFTLGNPTVQRD
WRLQTLNGSHAYLTLALHWVEPREHII PVVVSHNAQMWQLLVRVIVCRCNVEGQCMRKVGRMK
GMPTKLSAVGILVGTLLVAIGIFLILIFTHWTMSRKKDPDQPADSVPLKATV

Signal peptide:

amino acids 1-18

Transmembrane domain:

amino acids 762-784

99/168

FIGURE 99

GGCTGACCGTGCTACATTGCCTGGAGGAAGCCTAAGGAACCCAGGCATCCAGCTGCCCCACGCC
TGAGTCCAAGATTCTTCCCAGGAACACAAACGTAGGAGACCCACGCTCCTGGAAGCACCAGCC
TTTATCTCTTCACCTTCAAGTCCCCTTTCTCAAGAATCCTCTGTTCTTTGCCCTCTAAAGTCT
TGGTACATCTAGGACCCAGGCATCTTGCTTTCCAGCCACAAAGAGACAGATGAAGATGCAGAA
AGGAAATGTTCTCCTTATGTTTGGTCTACTATTGCATTTAGAAGCTGCAACAAATTCCAATGA
GACTAGCACCTCTGCCAACACTGGATCCAGTGTGATCTCCAGTGGAGCCAGCACAGCCACCAA
CTCTGGGTCCAGTGTGACCTCCAGTGGGGTCAGCACAGCCACCATCTCAGGGTCCAGCGTGAC
CTCCAATGGGGTCAGCATAGTCAACCACTCTGAGTTCCATACAACCTCCAGTGGGATCAGCAC
AGCCACCAACTCTGAGTTCAGCACAGCGTCCAGTGGGATCAGCATAGCCACCAACTCTGAGTC
CAGCACAACTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTCCAGCACACCCTCCAGTGG
GGCCAGCACAGTCACCAACTCTGGGTCCAGTGTGACCTCCAGTGGAGCCAGCACTGCCACCAA
CTCTGAGTCCAGCACAGTGTCCAGTAGGGCCAGCACTGCCACCAACTCTGAGTCTAGCACACT
CTCCAGTGGGGCCAGCACAGCCACCAACTCTGACTCCAGCACAACTCCAGTGGGGCTAGCAC
AGCCACCAACTCTGAGTCCAGCACAACTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTC
CAGCACAGTGTCCAGTAGGGCCAGCACTGCCACCAACTCTGAGTCCAGCACAACTCCAGTGG
GGCCAGCACAGCCACCAACTCTGAGTCCAGAACGACCTCCAATGGGGCTGGCACAGCCACCAA
CTCTGAGTCCAGCACGACCTCCAGTGGGGCCAGCACAGCCACCAACTCTGACTCCAGCACAGT
GTCCAGTGGGGCCAGCACTGCCACCAACTCTGAGTCCAGCACGACCTCCAGTGGGGCCAGCAC
AGCCACCAACTCTGAGTCCAGCACGACCTCCAGTGGGGCTAGCACAGCCACCAACTCTGACTC
CAGCACAACTCCAGTGGGGCCGGCACAGCCACCAACTCTGAGTCCAGCACAGTGTCCAGTGG
GATCAGCACAGTCACCAATTCTGAGTCCAGCACACCCTCCAGTGGGGCCAACACAGCCACCAA
CTCTGAGTCCAGTACGACCTCCAGTGGGGCCAACACAGCCACCAACTCTGAGTCCAGCACAGT
GTCCAGTGGGGCCAGCACTGCCACCAACTCTGAGTCCAGCACAACTCCAGTGGGGTCCAGCAC
AGCCACCAACTCTGAGTCCAGCACAACTCCAGTGGGGCTAGCACAGCCACCAACTCTGACTC
CAGCACAACTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTCTAGCACAGTGTCCAGTGG
GATCAGCACAGTCACCAATTCTGAGTCCAGCACAACTCCAGTGGGGCCAACACAGCCACCAA
CTCTGGGTCCAGTGTGACCTCTGCAGGCTCTGGAACAGCAGCTCTGACTGGAATGCACACAAC
TTCCCATAGTGCATCTACTGCAGTGAAGTGAAGGCAAAGCCTGGTGGGTCCCTGGTGGCGTGGGA
AATCTTCCTCATCACCTGGTCTCGGTGTGGCGGCGGTGGGGCTCTTTGCTGGGCTCTTCTT
CTGTGTGAGAAACAGCCTGTCCCTGAGAAACACCTTTAACACAGCTGTCTACCACCCTCATGG
CCTCAACCATGGCCTTGGTCCAGGCCCTGGAGGGAATCATGGAGCCCCCACAGGCCCAGGTG
GAGTCCTAACTGGTTCTGGAGGAGACCAGTATCATCGATAGCCATGGAGATGAGCGGGAGGAA
CAGCGGGCCCTGAGCAGCCCCGGAAGCAAGTGCCGCATTCTTCAGGAAGGAAGAGACCTGGGC
ACCAAGACCTGGTTTCTTTTCATTCATCCCAGGAGACCCCTCCCAGCTTTGTTTGAGATCCT
GAAAATCTTGAAGAAGGTATTCTTACCTTTCTTGCCTTTACCAGACACTGGAAAGAGAATAC
TATATTGCTCATTTAGCTAAGAAATAAATACATCTCATCTAACACACACGACAAAGAGAAGCT
GTGCTTGCCCCGGGGTGGGTATCTAGCTCTGAGATGAACTCAGTTATAGGAGAAAACCTCCAT
GCTGGACTCCATCTGGCATTCAAAATCTCCACAGTAAAATCCAAAGACCTCAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

100/168

FIGURE 100

MKMQKGNVLLMFGLLLHLEAATNSNETSTSANTGSSVSSGASTATNSGSSVTSSGVSTATIS
GSSVTSNGVSIVTNSEFHTTSSGISTATNSEFSTASSGISIATNSESSTTSSGASTATNSESS
TPSSGASTVTNSGSSVTSSGASTATNSESSTVSSRASTATNSESSTLSSGASTATNSDSSTTS
SGASTATNSESSTTSSGASTATNSESSTVSSRASTATNSESSTTSSGASTATNSESRTTSNGA
GTATNSESSTTSSGASTATNSDSSTVSSGASTATNSESSTTSSGASTATNSESSTTSSGASTA
TNSDSSTTSSGAGTATNSESSTVSSGISTVTNSESSTPSSGANTATNSESSTTSSGANTATNS
ESSTVSSGASTATNSESSTTSSGVSTATNSESSTTSSGASTATNSDSSTTSSEASTATNSESS
TVSSGISTVTNSESSTTSSGANTATNSGSSVTSAGSGTAALTGMHTTSHSASTAVSEAKPGGS
LVPWEIFLITLVSVVAAVGLFAGLFFCVRNSLSLRNTENTAVYHPHGLNHGLGPGPGGNHGAP
HRPRWSPNWFWRPVS SIAMEMSGRNSGP

Signal peptide:

amino acids 1-20

Transmembrane domain:

amino acids 510-532

101/168

FIGURE 101

GGCCGGACGCCTCCGCGTTACGGGATGAATTAACGGCGGGTTCGCGACGGAGGTTGTGACCCC
TACGGAGCCCCAGCTTGCCACGCACCCCACTCGGCGTCGCGCGGCGTGCCCTGCTTGTCACA
GGTGGGAGGCTGGAACATATCAGGCTGAAAAACAGAGTGGGTACTCTCTTCTGGGAAGCTGGCA
ACAAATGGATGATGTGATATATGCATTCCAGGGGAAGGGAAATTGTGGTGCTTCTGAACCCAT
GGTCAATTAACGAGGCAGTTTCTAGCTACTGCACGTACTTCATAAAGCAGGACTCTAAAAGCT
TTGGAATCATGGTGTCATGGAAAGGGATTTACTTTATACTGACTCTGTTTTGGGGAAGCTTTT
TTGGAAGCATTTCATGCTGAGTCCCTTTTTACCTTTGATGTTTGTAACCCATCTTGGTATC
GCTGGATCAACAACCGCCTTGTGGCAACATGGCTCACCTACCTGTGGCATTATTGGAGACCA
TGTTTGGTGTAAGAGTGATTATAACTGGGGATGCATTTGTTCCCTGGAGAAAGAAGTGTCATTA
TCATGAACCATCGGACAAGAATGGACTGGATGTTCCCTGTGGAATTGCCTGATGCGATATAGCT
ACCTCAGATTGGAGAAAATTTGCCTCAAAGCGAGTCTCAAAGGTGTTCCCTGGATTTGGTTGGG
CCATGCAGGCTGCTGCCTATATCTTCATTCATAGGAAATGGAAGGATGACAAGAGCCATTTTCG
AAGACATGATTGATTACTTTTGTGATATTCACGAACCACTTCAACTCCTCATATTCCCAGAAG
GGACTGATCTCACAGAAAACAGCAAGTCTCGAAGTAATGCATTTGCTGAAAAAATGGACTTC
AGAAATATGAATATGTTTTACATCCAAGAACTACAGGCTTTACTTTTGTGGTAGACCGTCTAA
GAGAAGGTAAGAACCTTGATGCTGTCCATGATATCACTGTGGCGTATCCTCACAACATTCCTC
AATCAGAGAAGCACCTCCTCCAAGGAGACTTTCCCAGGGAAATCCACTTTCACGTCCACCGGT
ATCCAATAGACACCCTCCCCACATCCAAGGAGGACCTTCAACTCTGGTGCCACAAACGGTGGG
AAGAGAAAGAAGAGAGGCTGCGTTCCTTCTATCAAGGGGAGAAGAATTTTTATTTTACCGGAC
AGAGTGTCATTCCACCTTGCAAGTCTGAACTCAGGGTCCTTGTGGTCAAATTGCTCTCTATAC
TGTATTGGACCCTGTTTCAGCCCTGCAATGTGCCTACTCATATATTTGTACAGTCTTGTTAAGT
GGTATTTTATAATCACCATTGTAATCTTTGTGCTGCAAGAGAGAATATTTGGTGGACTGGAGA
TCATAGAACTTGCATGTTACCGACTTTTACACAAACAGCCACATTTAAATTCAAAGAAAAATG
AGTAAGATTATAAGGTTTGCCATGTGAAAACCTAGAGCATATTTTGGAAATGTTCTAAACCTT
TCTAAGCTCAGATGCATTTTTGCATGACTATGTCGAATATTTCTTACTGCCATCATTATTTGT
TAAAGATATTTTGCACCTTAATTTTGTGGGAAAAATATTGCTACAATTTTTTTTAAATCTCTGAA
TGTAATTTTCGATACTGTGTACATAGCAGGGAGTGATCGGGGTGAAATAACTTGGGCCAGAATA
TTATTAAACAATCATCAGGCTTTTAAA

102/168

FIGURE 102

MHSRGREIVVLLNPWSINEAVSSYCTYFIKQDSKSF GIMVSWKGIYFILTLFWGSFFGSIFML
SPFLPLMFVNPSWYRWINNRLVATWLTLPVALLETMFGVKVIITGDAFVPGERSVIIMNHRTR
MDWMFLWNCLMRYSYLRLEKICLKASLKGVPGF GWAMQAAAYIFIHRKWKDDKSHFEDMIDYF
CDIHEPLQLLIFPEGTDLTENSKSRSNAFAEKNG LQKYEYVLHPRTTGFTFVVDRLREGKNLD
AVHDITVAYPHNIPQSEKHLLQGDFPREIH FHVHRYPIDTLPTSKEDLQLWCHKRWEEKEERL
RSFYQGEKNFYFTGQSVIPPCKSELRVLVVKLLS ILYWTLFSPAMCLLIYLYSLVKWYFIITI
VIFVLQERIFGGLEIIEELACYRLLHKQPHLNSK KNE

Important features of the protein:

Signal peptide:

amino acids 1-22

Transmembrane domains:

amino acids 44-63, 90-108, 354-377

103/168

FIGURE 103

CGGCTCGAGCGGCTCGAGTGAAGAGCCTCTCCACGGCTCCTGCGCCTGAGACAGCTGGCCTGA
CCTCCAAATCATCCATCCACCCCTGCTGTCATCTGTTTTTCATAGTGTGAGATCAACCCACAGG
AATATCCATGGCTTTTTGTGCTCATTTTGGTTCTCAGTTTCTACGAGCTGGTGTGAGGACAGTG
GCAAGTCACTGGACCGGGCAAGTTTGTCCAGGCCTTGGTGGGGGAGGACGCCGTGTTCTCCTG
CTCCCTCTTTCCTGAGACCAGTGCAGAGGCTATGGAAGTGCGGTTCTTCAGGAATCAGTTCCA
TGCTGTGGTCCACCTCTACAGAGATGGGGAAGACTGGGAATCTAAGCAGATGCCACAGTATCG
AGGGAGAACTGAGTTTGTGAAGGACTCCATTGCAGGGGGGCGTGTCTCTCTAAGGCTAAAAAA
CATCACTCCCTCGGACATCGGCCTGTATGGGTGCTGGTTCAGTTCCCAGATTTACGATGAGGA
GGCCACCTGGGAGCTGCGGGTGGCAGCACTGGGCTCACTTCCTCTCATTTCATCGTGGGATA
TGTTGACGGAGGTATCCAGTTACTCTGCCTGTCCTCAGGCTGGTTCCCCCAGCCCACAGCCAA
GTGGAAAGGTCCACAAGGACAGGATTTGTCTTCAGACTCCAGAGCAAATGCAGATGGGTACAG
CCTGTATGATGTGGAGATCTCCATTATAGTCCAGGAAAATGCTGGGAGCATATTGTGTTCCAT
CCACCTTGCTGAGCAGAGTCATGAGGTGGAATCCAAGGTATTGATAGGAGAGACGTTTTTCCA
GCCCTCACCTTGGCGCCTGGCTTCTATTTTACTCGGGTTACTCTGTGGTGCCCTGTGTGGTGT
TGTCAATGGGGATGATAATTGTTTTCTTCAAATCCAAAGGGGAAAATCCAGGCGGAACTGGACTG
GAGAAGAAAGCACGGACAGGCAGAATTGAGAGACGCCCCGAAACACGCAGTGGAGGTGACTCT
GGATCCAGAGACGGCTCACCCGAAGCTCTGCGTTTCTGATCTGAAAATGTAACCCATAGAAA
AGCTCCCCAGGAGGTGCCTCACTCTGAGAAGAGATTTACAAGGAAGAGTGTGGTGGCTTCTCA
GGGTTTCCAAGCAGGGAGACATTACTGGGAGGTGGACGTGGGACAAAATGTAGGGTGGTATGT
GGGAGTGTGTCGGGATGACGTAGACAGGGGGAAGAACAATGTGACTTTGTCTCCCAACAATGG
GTATTGGGTCTCAGACTGACAACAGAACATTTGTATTTACATTCAATCCCCATTTTATCAG
CCTCCCCCCCCAGCACCCCTCCTACACGAGTAGGGGTCTTCTGACTATGAGGGTGGGACCAT
CTCCTTCTTCAATACAAATGACCAGTCCCTTATTTATACCCTGCTGACATGTCAGTTTGAAGG
CTTGTTGAGACCCCTATATCCAGCATGCGATGTATGACGAGGAAAAGGGGACTCCCATATTCAT
ATGTCCAGTGTCTGCGGATGAGACAGAGAAGACCCTGCTTAAAGGGCCCCACACCACAGACC
CAGACACAGCCAAGGGAGAGTGCTCCCGACAGGTGGCCCCAGCTTCCTCTCCGGAGCCTGCGC
ACAGAGAGTCACGCCCCCCTCTCCTTTAGGGAGCTGAGGTTCTTCTGCCCTGAGCCCTGCA
GCAGCGGCAGTCACAGCTTCCAGATGAGGGGGGATTGGCCTGACCCTGTGGGAGTCAGAAGCC
ATGGCTGCCCTGAAGTGGGGACGGAATAGACTCACATTAGGTTTAGTTTGTGAAAATCCATC
CAGCTAAGCGATCTTGAACAAGTCACAACCTCCCAGGCTCCTCATTTGCTAGTCACGGACAGT
GATTCCTGCCTCACAGGTGAAGATTAAAGAGACAACGAATGTGAATCATGCTTGCAGGTTTGA
GGGCACAGTGTTTGCTAATGATGTGTTTTTATATTATACATTTTCCCACCATAAACTCTGTTT
GCTTATTCCACATTAATTTACTTTTCTCTATAACCAATCACCCATGGAATAGTTATTGAACAC
CTGCTTTGTGAGGCTCAAAGAATAAAGAGGAGGTAGGATTTTTTCACTGATTCTATAAGCCCAG
CATTACCTGATACCAAACCAGGCAAAGAAAACAGAAGAAGAGGAAGGAAAACACTACAGGTCCA
TATCCCTCATTAACACAGACACAAAATTCTAAATAAAATTTTAAACAATTAACTAAACAAT
ATATTTAAAGATGATATATACTACTCAGTGTGGTTTGTCCCACAAATGCAGAGTTGGTTTAA
TATTTAAATATCAACCAGTGTAATTCAGCACATTAATAAAGTAAAAAAGAAAACCATAAAAAA
AAAAAAA

WO 01/16318

PCT/US00/23328

104/168

FIGURE 104

MAFVLILVLSFYELVSGWQVTGPGKFVQALVGEDAVFSCSLFPETSAEAMEVRFFRNQFHAV
VHLYRDGEDWESKQMPQYRGRTEFVKDSIAGGRVSLRLKNITPSDIGLYGCWFSSQIYDEEAT
WELRVAALGSLPLISIVGYVDGGIQLLCLSSGWFPQPTAKWKGPQGQDLSSDSRANADGYSLY
DVEISIIVQENAGSILCSIHLEQSHVESKVLIGETFFQPSPWRLASILLGLLCGALCGVVM
GMIIVFFKSKGKIQAELEDWRRKHGQAEIRDARKHAVEVTLPETAHPKLCVSDLKTVTHRKAP
QEVPHSEKRFTTRKSVVASQGFQAGRHYWEVDVGQNVGWYVGVCRDDVDRGKNNVTLSPNNGYW
VLRLTTEHLYFTFNPHFISLPPSTPPTRVGVFLDYEGGTISFFNTNDQSLIYTLLTCQFEGLL
RPYIQHAMYDEEKGTPIFICPVSWG

Signal peptide:

amino acids 1-17

Transmembrane domains:

amino acids 131-150, 235-259

105/168

FIGURE 105

CCTTCACAGGACTCTTCATTGCTGGTTGGCAATGATGTATCGGCCAGATGTGGTGAGGGCTAG
GAAAAGAGTTTGTGGGAACCCTGGGTATCGGCCTCGTCATCTTCATATCCCTGATTGTCCT
GGCAGTGTGCATTGGACTCACTGTTTATTATGTGAGATATAATCAAAGAAGACCTACAATTA
CTATAGCACATTGTCATTTACAACCTGACAACTATATGCTGAGTTTGGCAGAGAGGCTTCTAA
CAATTTTACAGAAATGAGCCAGAGACTTGAATCAATGGTGAAAAATGCATTTTATAAATCTCC
ATTAAGGGAAGAATTTGTCAAGTCTCAGGTTATCAAGTTCAGTCAACAGAAGCATGGAGTGTT
GGTCATATGCTGTTGATTTGTAGATTTCACTCTACTGAGGATCCTGAACTGTAGATAAAAT
TGTTCAACTTGTTTTACATGAAAAGCTGCAAGATGCTGTAGGACCCCTAAAGTAGATCCTCA
CTCAGTTAAAATTAAAAAAATCAACAAGACAGAAACAGACAGCTATCTAAACCATTGCTGCGG
AACACGAAGAAGTAAAACTCTAGGTCAGAGTCTCAGGATCGTTGGTGGGACAGAAGTAGAAGA
GGGTGAATGGCCCTGGCAGGCTAGCCTGCAGTGGGATGGGAGTCATCGCTGTGGAGCAACCTT
AATTAATGCCACATGGCTTGTGAGTGCTGCTCACTGTTTTACAACATATAAGAACCCTGCCAG
ATGGACTGCTTCCTTTGGAGTAACAATAAAACCTTCGAAAATGAAACGGGGTCTCCGGAGAAT
AATTGTCCATGAAAAATACAAACACCCATCACATGACTATGATATTTCTCTTGCAGAGCTTTC
TAGCCCTGTTCCCTACACAAATGCAGTACATAGAGTTTGTCTCCCTGATGCATCCTATGAGTT
TCAACCAGGTGATGTGATGTTTGTGACAGGATTTGGAGCACTGAAAAATGATGGTTACAGTCA
AAATCATCTTCGACAAGCACAGGTGACTCTCATAGACGCTACAACCTGCAATGAACCTCAAGC
TTACAATGACGCCATAACTCCTAGAATGTTATGTGCTGGCTCCTTAGAAGGAAAAACAGATGC
ATGCCAGGGTGACTCTGGAGGACCACTGGTTAGTTCAGATGCTAGAGATATCTGGTACCTTGC
TGGAATAGTGAGCTGGGGAGATGAATGTGCGAAACCCAACAAGCCTGGTGTTTATACTAGAGT
TACGGCCTTGCGGGACTGGATTACTTCAAAAACCTGGTATCTTAAGAGACAAAAGCCTCATGGAA
CAGATAACATTTTTTTTTTGTTTTTTGGGTGTGGAGGCCATTTTTAGAGATACAGAATTGGAGA
AGACTTGCAAAACAGCTAGATTTGACTGATCTCAATAAACTGTTTGCTTGATGCATGTATTTT
CTTCCCAGCTCTGTTCCGCACGTAAGCATCCTGCTTCTGCCAGATCAACTCTGTCATCTGTGA
GCAATAGTTGAACTTTATGTACATAGAGAAATAGATAATACAATATTACATTACAGCCTGTA
TTCATTTGTTCTCTAGAAGTTTGTGTCAGAATTTTGAAGTTGTTGACATAAATTTGTAATGCATA
TATACAATTTGAAGCACTCCTTTTCTTCAGTTCCTCAGCTCCTCTCATTTTCAGCAAATATCCA
TTTTCAAGGTGCAGAACAAAGGAGTGAAAGAAAATATAAGAAGAAAAAAATCCCCTACATTTTA
TTGGCACAGAAAAGTATTAGGTGTTTTTCTTAGTGGAATATTAGAAATGATCATATTCATTAT
GAAAGGTCAAGCAAAGACAGCAGAATACCAATCACTTCATCATTTAGGAAGTATGGGAACATA
GTTAAGGAAGTCCAGAAAGAAGCCAAGATATATCCTTATTTTCATTTCCAAACAACCTACTATG
ATAAATGTGAAGAAGATTCTGTTTTTTTTGTGACCTATAATAATTATACAAACTTCATGCAATG
TACTTGTTCTAAGCAAATTAAAGCAAATATTTATTTAACATTGTTACTGAGGATGTCAACATA
TAACAATAAAATATAAATCACCCA

106/168

FIGURE 106

MMYRPDVVRARKRVCWEPWVIGLVIFISLIVLAVCIGLTVHYVRYNQKKTYNYYSTLSFTTDK
LYAEFGREASNNFTEMSQRLESMVKNAFYKSPLREEFVKSQVIKFSQQKHGVLAHMLLICRFH
STEDPETVDKIVQLVLHEKLQDAVGPPKVDPHSVKIKKINKTETDSYLNHCCGTRRSKTLGQS
LRIVGGTEVEEGEWPWQASLQWDGSHRCGATLINATWLVSAAHCFTTYKNPARWTASFGVTIK
PSKMKRGLRRIIVHEKYKHPSHDYDISLAELSSPVYTNVHRVCLPDASYEFQPGDVMFVTG
FGALKNDGYSONHLRQAQVTLIDATTCNEPQAYNDAITPRMLCAGSLEGKTDACQGDSSGGPLV
SSDARDIWYLAGIVSWGDECAKPNKPGVYTRVTALRDWITSKTGI

Transmembrane domain:

amino acids 21-40 (type II)

107/168

FIGURE 107

AGAGAAAGAAGCGTCTCCAGCTGAAGCCAATGCAGCCCTCCGGCTCTCCGCGAAGAAGTTCCC
TGCCCCGATGAGCCCCCGCCGTGCGTCCCCGACTATCCCCAGGCGGGCGTGGGGCACCGGGCC
CAGCGCCGACGATCGCTGCCGTTTTGCCCCTTGGGAGTAGGATGTGGTGAAAGGATGGGGCTTC
TCCCTTACGGGGCTCACAATGGCCCAGAGAAGATTCCGTGAAGTGTCTGCGCTGCCTGCTCTAC
GCCCTCAATCTGCTCTTTTGGTTAATGTCCATCAGTGTGTTGGCAGTTTCTGCTTGGATGAGG
GACTACCTAAATAATGTTCTCACTTTAACTGCAGAAACGAGGGTAGAGGAAGCAGTCATTTTG
ACTTACTTTTCTGTGGTTCATCCGGTCATGATTGCTGTTTGCTGTTTCCTTATCATTGTGGGG
ATGTTAGGATATTGTGGAACGGTGAAAAGAAATCTGTTGCTTCTTGCAATGGTACTTTGGAAGT
TTGCTTGTCAATTTTCTGTGTAGAACTGGCTTGTGGCGTTTGGACATATGAACAGGAACCTTATG
GTTCCAGTACAATGGTCAGATATGGTCACCTTTGAAAGCCAGGATGACAAATTATGGATTACCT
AGATATCGGTGGCTTACTCATGCTTGGAAATTTTTTTTCAGAGAGAGTTTAAAGTGCTGTGGAGTA
GTATATTTCACTGACTGGTTGGAAATGACAGAGATGGACTGGCCCCCAGATTCTCTGCTGTGTT
AGAGAATTCCCAGGATGTTCCAAACAGGCCACCAGGAAGATCTCAGTGACCTTTATCAAGAG
GGTTGTGGGAAGAAAATGTATTCCTTTTTTGAGAGGAACCAAACAACCTGCAGGTGCTGAGGTTT
CTGGGAATCTCCATTGGGGTGACACAAATCCTGGCCATGATTCTCACCATTACTCTGCTCTGG
GCTCTGTATTATGATAGAAGGGAGCCTGGGACAGACCAAATGATGTCCTTGAAGAATGACAAC
TCTCAGCACCTGTCATGTCCCTCAGTAGAACTGTTGAAACCAAGCCTGTCAAGAATCTTTGAA
CACACATCCATGGCAAACAGCTTTAATACACACTTTGAGATGGAGGAGTTATTAAAAAGAAATG
TCACAGAAGAAAACCACAAACCTTGTTTTATTGGACTTGTGAATTTTTTGAGTACATACTATGTG
TTTCAGAAATATGTAGAAATAAAAATGTTGCCATAAAATAACACCTAAGCATATACTATTCTA
TGCTTTTAAATGAGGATGGAAAAGTTTCATGTCATAAGTCACCACCTGGACAATAATTGATGC
CCTTAAATGCTGAAGACAGATGTCATACCCACTGTGTAGCCTGTGTATGACTTTTACTGAAC
ACAGTTATGTTTTGAGGCAGCATGGTTTGATTAGCATTTCCGCATCCATGCAAACGAGTCACA
TATGGTGGGACTGGAGCCATAGTAAAGGTTGATTTACTTCTACCAACTAGTATATAAAGTACT
AATTAAATGCTAACATAGGAAGTTAGAAAATACTAATAACTTTTATTACTCAGCGATCTATTC
TTCTGATGCTAAATAAATTATATATCAGAAAACCTTCAATATTGGTGACTACCTAAATGTGAT
TTTTGCTGGTTACTAAAATATTCTTACCACTTAAAGAGCAAGCTAACACATTGTCTTAAGCT
GATCAGGGATTTTTTGTATATAAGTCTGTGTTAAATCTGTATAATTCAGTCGATTTTCAGTTCT
GATAATGTTAAGAATAACCATTATGAAAAGGAAAATTTGTCCTGTATAGCATCATTATTTTTTA
GCCTTTCCTGTTAATAAAGCTTTACTATTCTGTCCTGGGCTTATATTACACATATAACTGTTA
TTTAAATACTTAACCACTAATTTTGAAAATTACCAGTGTGATACATAGGAATCATTATTCAGA
ATGTAGTCTGGTCTTTAGGAAGTATTAATAAGAAAATTTGCACATAACTTAGTTGATTCAGAA
AGGACTTGATGCTGTTTTTCTCCCAAATGAAGACTCTTTTTTGACACTAAACACTTTTTTAAAA
AGCTTATCTTTGCCTTCTCCAAACAAGAAGCAATAGTCTCCAAGTCAATATAAATTCTACAGA
AAATAGTGTTCTTTTTCTCCAGAAAATGCTTGTGAGAATCATTAAACATGTGACAATTTAG
AGATTCTTTGTTTTATTTCCTGATTAATATACTGTGGCAAATTACACAGATTATTAAATTTT
TTTACAAGAGTATAGTATATTTATTTGAAATGGGAAAAGTGCATTTTACTGTATTTTGTGTAT
TTTGTTTATTTCTCAGAATATGGAAAGAAAATTAAATGTGTCAATAAATATTTTCTAGAGAG
TAA

108/168

FIGURE 108

MAREDSVKCLRCLLYALNLLFWLMSISVLAVSAWMRDYLNNVLTTLTAETRVEEAVILTYFPVV
HPVMIAVCCFLIIVGMLGYCGTVKRNLLLLLAWYFGSLLVIFCVELACGVWTYEQELMVPVQWS
DMVTLKARMTNYGLPRYRWLTHAWNFFQREFKCCGVVYFTDWLEMTMDWPPDSCCVREFPGC
SKQAHQEDLSDLYQEGCGKKMYSFLRGTKQLQVLRFLGISIGVTQILAMILTITLLWALYYDR
REPGTDQMMSLKNDNSQHLSCPSVELLKPSLSRIFEHTSMANSFNTHFEMEEL

Signal peptide:

amino acids 1-33

Transmembrane domains:

amino acids 12-35, 57-86, 94-114, 226-248

109/168

FIGURE 109

CCAAGGCCAGAGCTGTGGACACCTTATCCCACTCATCCTCATCCTCTTCCTCTGATAAAGCCC
CTACCAGTGCTGATAAAGTCTTTCTCGTGAGAGCCTAGAGGCCTTAAAAAAGTGCTTGA
AAGAGAAGGGGACAAAGGAACACCAGTATTAAGAGGATTTTCCAGTGTTTCTGGCAGTTGGTC
CAGAAGGATGCCTCCATTCTGCTTCTCACCTGCCTCTTCATCACAGGCACCTCCGTGTCACC
CGTGGCCCTAGATCCTTGTTCTGCTTACATCAGCCTGAATGAGCCCTGGAGGAACACTGACCA
CCAGTTGGATGAGTCTCAAGGTCCTCCTCTATGTGACAACCATGTGAATGGGGAGTGGTACCA
CTTCACGGGCATGGCGGGAGATGCCATGCCTACCTTCTGCATACCAGAAAACCACTGTGGAAC
CCACGCACCTGTCTGGCTCAATGGCAGCCACCCCTAGAAAGGCGACGGCATTGTGCAACGCCA
GGCTTGTGCCAGCTTCAATGGGAAGTGTGTCTCTGGAACACCACGGTGGAAAGTCAAGGCTTG
CCGTGGAGGCTACTATGTGTATCGTCTGACCAAGCCCAGCGTCTGCTTCCACGTCTACTGTGG
TCATTTTTTATGACATCTGCGACGAGGACTGCCATGGCAGCTGCTCAGATACCAGCGAGTGCAC
ATGCGCTCCAGGAAGTGTGCTAGGCCCTGACAGGCAGACATGCTTTGATGAAAATGAATGTGA
GCAAAACAACGGTGGCTGCAGTGAGATCTGTGTGAACCTCAAAAACCTCCTACCGCTGTGAGTG
TGGGGTTGGCEGTGTGCTAAGAAGTGATGGCAAGACTTGTGAAGACGTTGAAGGATGCCACAA
TAACAATGGTGGCTGCAGCCACTCTTGCTTGGATCTGAGAAAGGCTACCAGTGTGAATGTCC
CCGGGGCCTGGTGCTGTCTGAGGATAACCACACTTGCCAAGTCCCTGTGTTGTGCAAATCAAA
TGCCATTGAAGTGAACATCCCCAGGGAGCTGGTTGGTGGCCTGGAGCTCTTCCTGACCAACAC
CTCCTGCCGAGGAGTGTCCAACGGCACCCATGTCAACATCCTCTTCTCTCAAGACATGTGG
TACAGTGGTCGATGTGGTGAATGACAAGATTGTGGCCAGCAACCTCGTGACAGGTCTACCCAA
GCAGACCCCGGGGAGCAGCGGGGACTTCATCATCCGAACCAGCAAGCTGCTGATCCCGGTGAC
CTGCGAGTTTCCACGCCTGTACACCATTCTGAAGGATACGTTCCCAACCTTCGAAACTCCCC
ACTGGAAATCATGAGCCGAAATCATGGGATCTTCCCATTCCTCTGGAGATCTTCAAGGACAA
TGAGTTTGAAGAGCCTTACCGGGAAGCTCTGCCCACCCTCAAGCTTCGTGACTCCCTCTACTT
TGGCATTGAGCCCGTGGTGCACGTGAGCGGCTTGGAAAGCTTGGTGGAGAGCTGCTTTGCCAC
CCCCACCTCCAAGATCGACGAGGTCTGAAATACTACCTCATCCGGGATGGCTGTGTTTCAGA
TGACTCGGTAAAGCAGTACACATCCCGGGATCACCTAGCAAAGCACTTCCAGGTCCCTGTCTT
CAAGTTTGTGGGCAAAGACCACAAGGAAGTGTCTGCACTGCCGGGTCTTGTCTGTGGAGT
GTTGGACGAGCGTTCCCGCTGTGCCAGGGTTGCCACCGGCGAATGCGTCGTGGGGCAGGAGG
AGAGGACTCAGCCGGTCTACAGGGCCAGACGCTAACAGGCGGCCCCGATCCGCATCGACTGGGA
GGACTAGTTCGTAGCCATACCTCGAGTCCCTGCATTGGACGGCTCTGCTCTTTGGAGCTTCTC
CCCCACCGCCCTCTAAGAACATCTGCCAACAGCTGGGTTCAGACTTCACACTGTGAGTTCAG
ACTCCCAGCACCAACTCACTCTGATTCTGGTCCATTAGTGGGCACAGGTACAGCACTGCTG
AACAATGTGGCCTGGGTGGGGTTTCATCTTTCTAGGGTTGAAAATAAAGTGTCCACCCAGAA
AGACACTCACCCCATTTCCCTCATTTCTTTCTTACACTTAAATACCTCGTGTATGGTGAATC
AGACCACAAAATCAGAAGCTGGGTATAATATTTCAAGTTACAAACCTAGAAAAATTAAACAG
TTACTGAAATTATGACTTAAATACCCAATGACTCCTTAAATATGTAAATTATAGTTATACCTT
GAAATTTCAATTCAAATGCAGACTAATTATAGGGAATTTGGAAGTGTATCAATAAACAGTAT
ATAATTTT

110/168

FIGURE 110

MPPFLLLTCLFITGTSVSPVALDPCSAYISLNEPWRNTDHOQDESQGPPLCDNHVNGEWYHFT
GMAGDAMPTFCIPENHCGTHAPVWLNGSHPLEGDGIVQRQACASFNGNCCLWNTTVEVKACPG
GYVYRLTKPSVCFHVYCGHFYDIEDCHGSCSDTSECTCAPGTVLGPDRQTCFDENECEQN
NGGCSEICVNLKNSYRCECGVGRVLRSDGKTCEDEGCHNNNGGCSHSCLGSEKGYQCECPRG
LVLSEDNHTCQVPVLCKSNAIEVNIPRELVGGLLEFLTNTSCRGVSNHGVNIFSLKTCGTV
VDVVNDKIVASNLVTGLPKQTPGSSGDFIIRTSKLLIPVTCEFPRLYTISEGYVPNLRNSPLE
IMSRNHGIFPFTLEIFKDNEFEPEPYREALPTLKLKRDLSLYFGIEPVVHVSGLESLVESCFATPT
SKIDEVLKYYLIRDGCVSDDSVKQYTSRDHLAKHFQVPVFKEVVGKDHKEVFLHCRVLCVGLD
ERSRCAQGCHRRMRRGAGGEDSAGLQGQTLTGGPIRIDWED

Important features of the protein:

Signal peptide:

amino acids 1-16

N-glycosylation sites.

amino acids 89-93, 116-120, 259-263, 291-295, 299-303

Tyrosine kinase phosphorylation sites.

amino acids 411-418, 443-451

N-myristoylation sites.

amino acids 226-232, 233-239, 240-246, 252-258, 296-302, 300-306,
522-528, 531-537

Aspartic acid and asparagine hydroxylation site.

amino acids 197-209

ZP domain proteins.

amino acids 431-457

Calcium-binding EGF-like proteins.

amino acids 191-212, 232-253

111/168

FIGURE 111

GAGAGAGGCAGCAGCTTGCTCAGCGGACAAGGATGCTGGGCGTGAGGGACCAAGGCCTGCCCT
GCACTCGGGCCTCCTCCAGCCAGTGCTGACCAGGGACTTCTGACCTGCTGGCCAGCCAGGACC
TGTGTGGGGAGGCCCTCCTGCTGCCTTGGGGTGACAATCTCAGCTCCAGGCTACAGGGAGACC
GGGAGGATCACAGAGCCAGCATGTTACAGGATCCTGACAGTGATCAACCTCTGAACAGCCTCG
ATGTCAAACCCCTGCGCAAACCCCGTATCCCCATGGAGACCTTCAGAAAGGTGGGGATCCCCA
TCATCATAGCACTACTGAGCCTGGCGAGTATCATCATTGTGGTTGTCCTCATCAAGGTGATTC
TGGATAAATACTACTTCTCTGCGGGCAGCCTCTCCACTTCATCCCGAGGAAGCAGCTGTGTG
ACGGAGAGCTGGACTGTCCCTTGGGGGAGGACGAGGAGCACTGTGTCAAGAGCTTCCCCGAAG
GGCCTGCAGTGGCAGTCCGCCTCTCCAAGGACCGATCCACACTGCAGGTGCTGGACTCGGCCA
CAGGGAACCTGGTTCTCTGCCTGTTTCGACAACCTTCACAGAAGCTCTCGCTGAGACAGCCTGTA
GGCAGATGGGCTACAGCAGAGCTGTGGAGATTGGCCCAGACCAGGATCTGGATGTTGTTGAAA
TCACAGAAAACAGCCAGGAGCTTCGCATGCGGAACTCAAGTGGGCCCTGTCTCTCAGGCTCCC
TGGTCTCCCTGCACTGTCTTGCCTGTGGGAAGAGCCTGAAGACCCCCCGTGTGGTGGGTGGGG
AGGAGGCCTCTGTGGATTCTTGGCCTTGGCAGGTCAGCATCCAGTACGACAAACAGCACGTCT
GTGGAGGGAGCATCCTGGACCCCCACTGGGTCCTCACGGCAGCCCCTGCTTCAGGAAACATA
CCGATGTGTTCAACTGGAAGGTGCGGGCAGGCTCAGACAAACTGGGCAGCTTCCCATCCCTGG
CTGTGGCCAAGATCATCATCATTGAATTCAACCCCATGTACCCCAAAGACAATGACATCGCCC
TCATGAAGCTGCAGTTCCCCTCACTTTCTCAGGCACAGTCAGGCCCATCTGTCTGCCCTTCT
TTGATGAGGAGCTCACTCCAGCCACCCCCTCTGGATCATTGGATGGGGCTTTACGAAGCAGA
ATGGAGGGAAGATGTCTGACATACTGCTGCAGGCGTCAGTCCAGGTCATTGACAGCACACGGT
GCAATGCAGACGATGCGTACCAGGGGGAAGTCACCGAGAAGATGATGTGTGCAGGCATCCCGG
AAGGGGGTGTGGACACCTGCCAGGGTGACAGTGGTGGGCCCCCTGATGTACCAATCTGACCAGT
GGCATGTGGTGGGCATCGTTAGCTGGGGCTATGGCTGCGGGGGCCCGAGCACCCCAGGAGTAT
ACACCAAGGTCTCAGCCTATCTCAACTGGATCTACAATGTCTGGAAGGCTGAGCTGTAATGCT
GCTGCCCCCTTTGCAGTGCTGGGAGCCGCTTCCTTCCTGCCCTGCCACCTGGGGATCCCCCAA
AGTCAGACACAGAGCAAGAGTCCCCTTGGGTACACCCCTCTGCCACAGCCTCAGCATTTCTT
GGAGCAGCAAAGGGCCTCAATTCCTGTAAGAGACCCTCGCAGCCCAGAGGCGCCCAGAGGAAG
TCAGCAGCCCTAGCTCGGCCACACTTGGTGCTCCCAGCATCCCAGGGAGAGACACAGCCCCT
GAACAAGGTCTCAGGGGTATTGCTAAGCCAAGAAGGAACTTTCCCACACTACTGAATGGAAGC
AGGCTGTCTTGTAAGAGCCAGATCACTGTGGGCTGGAGAGGAGAAGGAAAGGGTCTGCGCCA
GCCCTGTCCGTCTTCACCCATCCCCAAGCCTACTAGAGCAAGAAACCAGTTGTAATATAAAAT
GCACTGCCCTACTGTTGGTATGACTACCGTTACCTACTGTTGTCATTGTTATTACAGCTATGG
CCACTATTATTAAAGAGCTGTGTAACATCTCTGGCAAAAAAAAAAAAA

112/168

FIGURE 112

MLQDPDS DQPLNSLDVKPLRKPRIPMETFRKVGIP IIIALLSLASIIIVVVL IKVILDKYYFL
CGQPLHFIPRKQLCDGELDCPLGEDEEHCVKSFPEGPAVAVRLSKDRSTLQVLDSATGNWFSA
CFDNFTEALAETACRQMGYSRAVEIGPDQDL DVVEITENSQELMRNSSG PCLSGSLVSLHCL
ACGKSLKT PRVVG GEEASVDSWPWQVSIQYDKQHVC GGSILDPHWVLTAAHCFRKHTDVFNWK
VRAGSDKLGSFPSLAVAKIIIIIEFNPMYPKDNDIALMKLQFPLTFSGTVRPICLPFFDEELTP
ATPLWIIIGWGFTKQNGGKMSDILLQASVQVIDSTRCNAD DAYQGEVTEKMMCAGIPEGGVDTC
QGDSGGPLMYQSDQWHVVGIVSWG YGCGGPSTPGVYTKVSAYLNWIYNVWKAEL

Transmembrane domain:

amino acids 32-53 (typeII)

113/168

FIGURE 113

GGCTGGACTGGAACCTCCTGGTCCCAAGTGATCCACCCGCCTCAGCCTCCCAAGGTGCTGTGAT
TATAGGTGTAAGCCACCGTGTCTGGCCTCTGAACAACCTTTTTCAGCAACTAAAAAGCCACAG
GAGTTGAACTGCTAGGATTCTGACTATGCTGTGGTGGCTAGTGCTCCTACTCCTACCTACATT
AAAATCTGTTTTTTGTTCTCTTGTAAGTAGCCTTTACCTTCCTAACACAGAGGATCTGTCACT
GTGGCTCTGGCCCAAACCTGACCTTCACTCTGGAACGAGAACAGAGGTTTCTACCCACACCGT
CCCCTCGAAGCCGGGGACAGCCTCACCTTGCTGGCCTCTCGCTGGAGCAGTGCCCTCACCAAC
TGTCTCACGTCTGGAGGCACTGACTCGGGCAGTGCAGGTAGCTGAGCCTCTTGGTAGCTGCGG
CTTTCAAGGTGGGCCTTGCCCTGGCCGTAGAAGGGATTGACCAAGCCCGAAGATTTCATAGGCG
ATGGCTCCCACTGCCCAGGCATCAGCCTTGCTGTAGTCAATCACTGCCCTGGGGCCAGGACGG
GCCGTGGACACCTGCTCAGAAGCAGTGGGTGAGACATCACGCTGCCCGCCCATCTAACCTTTT
CATGTCCTGCACATCACCTGATCCATGGGCTAATCTGAACTCTGTCCCAAGGAACCCAGAGCT
TGAGTGAGCTGTGGCTCAGACCCAGAAGGGGTCTGCTTAGACCACCTGGTTTATGTGACAGGA
CTTGCAATTCTCCTGGAACATGAGGGAACGCCGGAGGAAAGCAAAGTGGCAGGGAAGGAACTTG
TGCCAAATTATGGGTCAGAAAAGATGGAGGTGTTGGGTTATCACAAGGCATCGAGTCTCCTGC
ATTCAGTGGACATGTGGGGGAAGGGCTGCCGATGGCGCATGACACACTCGGGACTCACCTCTG
GGGCCATCAGACAGCCGTTTCCGCCCCGATCCACGTACCAGCTGCTGAAGGGCAACTGCAGGC
CGATGCTCTCATCAGCCAGGCAGCAGCCAAAATCTGCGATCACCAGCCAGGGGCAGCCGTCTG
GGAAGGAGCAAGCAAAGTGACCATTTCTCCTCCCCTCCTTCCCTCTGAGAGGCCCTCCTATGT
CCCTACTAAAGCCACCAGCAAGACATAGCTGACAGGGGCTAATGGCTCAGTGTTGGCCCAGGA
GGTCAGCAAGGCCTGAGAGCTGATCAGAAGGGCCTGCTGTGCGAACACGGAAATGCCTCCAGT
AAGCACAGGCTGCAAAATCCCCAGGCAAAGGACTGTGTGGCTCAATTTAAATCATGTTCTAGT
AATTGGAGCTGTCCCAAGACCAAAGGAGCTAGAGCTTGGTTCAAATGATCTCCAAGGGCCCT
TATACCCCAGGAGACTTTGATTTGAATTTGAAACCCCAAATCCAAACCTAAGAACCAGGTGCA
TTAAGAATCAGTTATTGCCGGGTGTGGTGGCCTGTAATGCCAACATTTTGGGAGGCCGAGGCG
GGTAGATCACCTGAGGTCAGGAGTTCAAGACCAGCCTGGCCAACATGGTGAAACCCCTGTCTC
TACTAAAAATACAAAAAACTAGCCAGGCATGGTGGTGTGTGCCTGTATCCCAGCTACTCGGG
AGGCTGAGACAGGAGAATTACTTGAACCTGGGAGGTGAAGGAGGCTGAGACAGGAGAATCACT
TCAGCCTGAGCAACACAGCGAGACTCTGTCTCAGAAAAAATAAAAAAAGAATTATGGTTATTT
GTAA

114/168

FIGURE 114

MLWWLVLLLLPTLKSVFCSLVTSLYLPNTEDLSLWLWPKPDLHSGTRTEVSTHTVPSKPGTAS
PCWPLAGAVPSPTVSRLEALTRAVQVAEPLGSCGFQGGPCPGRRRD

Signal peptide:

amino acids 1-15

115/168

FIGURE 115

CAGCAGTGGTCTCTCAGTCCTCTCAAAGCAAGGAAAGAGTACTGTGTGCTGAGAGACCATGGC
AAAGAATCCTCCAGAGAATTGTGAAGACTGTCACATTCTAAATGCAGAAGCTTTTAAATCCAA
GAAAATATGTAAATCACTTAAGATTTGTGGACTGGTGTTTGGTATCCTGGCCCTAACTCTAAT
TGTCCTGTTTTGGGGGAGCAAGCACTTCTGGCCGGAGGTACCCAAAAAAGCCTATGACATGGA
GCACACTTTCTACAGCAATGGAGAGAAGAAGAAGATTTACATGGAAATTGATCCTGTGACCAG
AACTGAAATATTCAGAAGCGGAAATGGCACTGATGAAACATTGGAAGTGCACGACTTTAAAAA
CGGATACACTGGCATCTACTTCGTGGGTCTTCAAAAATGTTTTATCAAACTCAGATTAAAGT
GATTCCTGAATTTTCTGAACCAGAAGAGGAAATAGATGAGAATGAAGAAATTACCACAACCTT
CTTTGAACAGTCAGTGATTTGGGTCCCAGCAGAAAAGCCTATTGAAAACCGAGATTTTCTTAA
AAATTCCAAAATTCTGGAGATTTGTGATAACGTGACCATGTATTGGATCAATCCCCTCTAAT
ATCAGTTTCTGAGTTACAAGACTTTGAGGAGGAGGGGAGAAGATCTTCACTTTCCTGCCAACGA
AAAAAAGGGATTGAACAAAATGAACAGTGGGTGGTCCCTCAAGTGAAAGTAGAGAAGACCCG
TCACGCCAGACAAGCAAGTGAGGAAGAACTTCCAATAAATGACTATACTGAAAATGGAATAGA
ATTTGATCCCATGCTGGATGAGAGAGGTTATTGTTGTATTTACTGCCGTCGAGGCAACCGCTA
TTGCCGCCGCGTCTGTGAACCTTTACTAGGCTACTACCCATATCCATACTGCTACCAAGGAGG
ACGAGTCATCTGTCGTGTCATCATGCCTTGTAACCTGGTGGGTGGCCCGCATGCTGGGGAGGGT
CTAATAGGAGGTTTGAGCTCAAATGCTTAACTGCTGGCAACATATAATAAATGCATGCTATT
CAATGAATTTCTGCCTATGAGGCATCTGGCCCCTGGTAGCCAGCTCTCCAGAATTACTTGTAG
GTAATTCCTCTCTTCATGTTCTAATAAACTTCTACATTATCACCAAAAAAAAAAAAAAAAAAAAA

116/168

FIGURE 116

MAKNPPENCEDCHILNAEAFKSKKICKSLKICGLVFGILALT LIVLFWGSKHFWPEVPPKKAYD
MEHTFYSNGEKKKIYMEIDPVTRTEIFRSGNGTDETLEVHDFKNGYTG IYFVGLQKCFIKTQI
KVIPEFSEPEEEIDENEEITTTFFEQSVIWVPAEKPIENRDFLKN SKILEICDNVTMYWINPT
LISVSELQDFEEEGEDLHF PANEEKKGIEQNEQWVVPQVKVEKTRHARQASEEELPINDYTENG
IEFDPMLDERGYCC IYCRRGNRYCRRVCEPLLGYYPYPYCYQGGRVICRVIMPCNWWVARMLGRV

Important features of the protein:

Signal peptide:

amino acids 1-40

Transmembrane domain:

amino acids 25-47 (type II)

N-glycosylation sites.

amino acids 94-97, 180-183

Glycosaminoglycan attachment sites.

amino acids 92-95, 70-73, 85-88, 133-136, 148-151, 192-195, 239-
242

N-myristoylation sites.

amino acids 33-38, 95-100, 116-121, 215-220, 272-277

Microbodies C-terminal targeting signal.

amino acids 315-317

Cytochrome c family heme-binding site signature.

amino acids 9-14

117/168

FIGURE 117

GAGCTCCCCTCAGGAGCGCGTTAGCTTCACACCTTCGGCAGCAGGAGGGCGGCAGCTTCTCGC
AGGCGGCAGGGCGGGCGGCCAGGATCATGTCCACCACCACATGCCAAGTGGTGGCGTTCCTCC
TGTCCATCCTGGGGCTGGCCGGCTGCATCGCGGCCACCGGGATGGACATGTGGAGCACCCAGG
ACCTGTACGACAACCCCGTCACCTCCGTGTTCCAGTACGAAGGGCTCTGGAGGAGCTGCGTGA
GGCAGAGTTCAGGCTTCACCGAATGCAGGCCCTATTTCACCATCCTGGGACTTCCAGCCATGC
TGCAGGCAGTGCGAGCCCTGATGATCGTAGGCATCGTCCTGGGTGCCATTGGCCTCCTGGTAT
CCATCTTTGCCCTGAAATGCATCCGCATTGGCAGCATGGAGGACTCTGCCAAAGCCAACATGA
CACTGACCTCCGGGATCATGTTTCATTGTCTCAGGTCTTTGTGCAATTGCTGGAGTGTCTGTGT
TTGCCAACATGCTGGTGACTAACTTCTGGATGTCCACAGCTAACATGTACACCGGCATGGGTG
GGATGGTGCAGACTGTTTCAGACCAGGTACACATTTGGTGCGGCTCTGTTTCGTGGGCTGGGTG
CTGGAGGCCTCACACTAATTGGGGGTGTGATGATGTGCATCGCCTGCCGGGGCCTGGCACCAG
AAGAAACCAACTACAAAGCCGTTTCTTATCATGCCTCAGGCCACAGTGTTGCCTACAAGCCTG
GAGGCTTCAAGGCCAGCACTGGCTTTGGGTCCAACACCAAAAACAAGAAGATATACGATGGAG
GTGCCCCGCACAGAGGACGAGGTACAATCTTATCCTTCCAAGCACGACTATGTGTAATGCTCTA
AGACCTCTCAGCACGGGCGGAAGAACTCCCGGAGAGCTCACCCAAAAACAAGGAGATCCCA
TCTAGATTTCTTCTTGCTTTTGACTCACAGCTGGAAGTTAGAAAAGCCTCGATTTTCATCTTTG
GAGAGGCCAAATGGTCTTAGCCTCAGTCTCTGTCTCTAAATATTCCACCATAAAACAGCTGAG
TTATTTATGAATTAGAGGCTATAGCTCACATTTTCAATCCTCTATTTCTTTTTTTAAATATAA
CTTTCTACTCTGATGAGAGAATGTGGTTTTAATCTCTCTCTCACATTTTGATGATTTAGACAG
ACTCCCCCTCTTCCTCCTAGTCAATAAACCCATTGATGATCTATTTCCCAGCTTATCCCCAAG
AAAACTTTTGAAAGGAAAGAGTAGACCCAAAGATGTTATTTTCTGCTGTTTGAATTTTGTCTC
CCCACCCCCCAACTTGGCTAGTAATAAACACTTACTGAAGAAGAAGCAATAAGAGAAAGATATT
TGTAATCTCTCCAGCCCATGATCTCGGTTTTCTTACACTGTGATCTTAAAAGTTACCAAACCA
AAGTCATTTTCAGTTTGAGGCAACCAAACCTTTCTACTGCTGTTGACATCTTCTTATTACAGC
AACACCATTCTAGGAGTTTCCTGAGCTCTCCACTGGAGTCCTCTTTCTGTGCGGGGTCAGAAA
TTGTCCCTAGATGAATGAGAAAATTATTTTTTTTTAATTTAAGTCCTAAATATAGTTAAAATAA
ATAATGTTTTAGTAAAATGATACACTATCTCTGTGAAATAGCCTCACCCCTACATGTGGATAG
AAGGAAATGAAAAAATAATTGCTTTGACATTGTCTATATGGTACTTTGTAAAGTCATGCTTAA
GTACAAATTCCATGAAAAGCTCACACCTGTAATCCTAGCACTTTGGGAGGCTGAGGAGGAAGG
ATCACTTGAGCCCAGAAGTTCGAGACTAGCCTGGGCAACATGGAGAAGCCCTGTCTCTACAAA
ATACAGAGAGAAAAAATCAGCCAGTCATGGTGGCATAACCTGTAGTCCCAGCATTCGGGGAG
GCTGAGGTGGGAGGATCACTTGAGCCCAGGGAGGTTGGGGCTGCAGTGAGCCATGATCACACC
ACTGCACTCCAGCCAGGTGACATAGCGAGATCCTGTCTAAAAAAATAAAAAATAAATAATGGA
ACACAGCAAGTCCTAGGAAGTAGGTTAAACTAATTCTTTAA

118/168

FIGURE 118

MSTTTCQVVAFLLSILGLAGCIAATGMDMWSTQDLYDNPVTSVFQYEGLRSCVRQSSGFTEC
RPYFTILGLPAMLQAVRALMIVGIVLGAIGLLVSIFALKCIRIGSMEDSAKANMTLTSGIMFI
VSGLCAIAGVSVFANMLVTNFWMSTANMYTGMGGMVQTVQTRYTFGAALFVGWVAGGLTLIGG
VMMCIACRGLAPEETNYKAVSYHASGHSVAYKPGGFKASTGFGSNTKNKKIYDGGARTEDEVQ
SYPSKHDYV

Signal peptide:

amino acids 1-23

Transmembrane domains:

amino acids 81-100, 121-141, 173-194

119/168

FIGURE 119

GGAAAACTGTTCTCTTCTGTGGCACAGAGAACCCTGCTTCAAAGCAGAAGTAGCAGTTCCGG
AGTCCAGCTGGCTAAAACTCATCCCAGAGGATAATGGCAACCCATGCCTTAGAAATCGCTGGG
CTGTTTCTTGGTGGTGTGGAATGGTGGGCACAGTGGCTGTCACTGTCATGCCTCAGTGGAGA
GTGTCGGCCTTCATTGAAAACAACATCGTGGTTTTTGAAACTTCTGGGAAGGACTGTGGATG
AATTGCGTGAGGCAGGCTAACATCAGGATGCAGTGCAAAATCTATGATTCCCTGCTGGCTCTT
TCTCCGGACCTACAGGCAGCCAGAGGACTGATGTGTGCTGCTTCCGTGATGTCCTTCTTGGCT
TTCATGATGGCCATCCTTGGCATGAAATGCACCAGGTGCACGGGGGACAATGAGAAGGTGAAG
GCTCACATTCTGCTGACGGCTGGAATCATCTTCATCATCACGGGCATGGTGGTGCTCATCCCT
GTGAGCTGGGTTGCCAATGCCATCATCAGAGATTTCTATAACTCAATAGTGAATGTTGCCCAA
AAACGTGAGCTTGGAGAAGCTCTCTACTTAGGATGGACCACGGCACTGGTGCTGATTGTTGGA
GGAGCTCTGTTCTGCTGCGTTTTTTGTTGCAACGAAAAGAGCAGTAGCTACAGATACTCGATA
CCTTCCCATCGCACAAACCCAAAAAAGTTATCACACCGGAAAGAAGTCACCGAGCGTCTACTCC
AGAAGTCAGTATGTGTAGTGTGTATGTTTTTTTAACTTTACTATAAAGCCATGCAAATGACA
AAAATCTATATTACTTTCTCAAATGGACCCCAAAGAACTTTGATTTACTGTTCTTAACTGC
CTAATCTTAATTACAGGAAGTGTGCATCAGCTATTTATGATTCTATAAGCTATTTTCAGCAGAA
TGAGATATTAAACCCAATGCTTTGATTGTTCTAGAAAGTATAGTAATTTGTTTTCTAAGGTGG
TTCAAGCATCTACTCTTTTTATCATTTACTTCAAATGACATTGCTAAAGACTGCATTATTTT
ACTACTGTAATTTCTCCACGACATAGCATTATGTACATAGATGAGTGTAACATTTATATCTCA
CATAGAGACATGCTTATATGGTTTTATTTAAAATGAAATGCCAGTCCATTACACTGAATAAAT
AGAACTCAACTATTGCTTTTCAGGGAAATCATGGATAGGGTTGAAGAAGGTACTATTAATTG
TTTAAAAACAGCTTAGGGATTAATGTCCTCCATTTATAATGAAGATTAAAATGAAGGCTTTAA
TCAGCATTGTAAAGGAAATTGAATGGCTTTCTGATATGCTGTTTTTTAGCCTAGGAGTTAGAA
ATCCTAACTTCTTTATCCTCTTCTCCCAGAGGCTTTTTTTTTCTTGTGTATTAAATTAACATT
TTTAAAACGCAGATATTTTGTCAAGGGGCTTTGCATTCAAACCTGCTTTTCCAGGGCTATACTC
AGAAGAAAGATAAAAGTGTGATCTAAGAAAAAGTGATGGTTTTAGGAAAGTGAAAATATTTTT
GTTTTTGTATTTGAAGAAGAATGATGCATTTTGACAAGAAATCATATATGTATGGATATATTT
TAATAAGTATTTGAGTACAGACTTTGAGGTTTCATCAATATAAATAAAAGAGCAGAAAAATAT
GTCTTGGTTTTTCATTTGCTTACCAAAAAACAACAACAAAAAAGTTGTCCTTTGAGAACTTC
ACCTGCTCCTATGTGGGTACCTGAGTCAAATTTGTCATTTTTTGTCTGTGAAAAATAAATTTTC
CTTCTTGTACCATTTCTGTTTAGTTTTACTAAAATCTGTAAATACTGTATTTTTCTGTTTATT
CCAAATTTGATGAACTGACAATCCAATTTGAAAGTTTGTGTCGACGTCTGTCTAGCTTAAAT
GAATGTGTTCTATTTGCTTTATACATTTATATTAATAAATTGTACATTTTTCTAATT

120/168

FIGURE 120

MATHALEIAGLFLGGVGMVGTVAVTVMPPQWRVSAFIENNIVVFENFWEGLWMNCVRQANIRMQ
CKIYDSLLALSPDLQAARGLMCAASVMSFLAFMMAILGMKCTRCTGDNEKVKAHILLTAGIIF
IITGMVVLIPVSWVANAIIRDFYNSIVNVAQKRELGEALYLGWTTALVLIVGGALFCCVFCCN
EKSSSYRYSIPSHRTTQKSYHTGKKSPSVYSRSQYV

Signal peptide:

amino acids 1-17

Transmembrane domains:

amino acids 82-101, 118-145, 164-188

121/168

FIGURE 121

GGAGAGAGGCGCGCGGGTGAAGGCGCATTGATGCAGCCTGCGGCGGCCTCGGAGCGCGGCGG
AGCCAGACGCTGACCACGTTCTCTCTCGGTCTCCTCCGCCTCCAGCTCCGCGCTGCCCGGC
AGCCGGGAGCCATGCGACCCCAGGGCCCCGCCGCTCCCCGCAGCGGCTCCGCGGCCTCCTGC
TGCTCCTGCTGCTGCAGCTGCCCGCGCCGTGAGCGCCTCTGAGATCCCCAAGGGGAAGCAAA
AGGCGCAGCTCCGGCAGAGGGAGGTGGTGGACCTGTATAATGGAATGTGCTTACAAGGGCCAG
CAGGAGTGCCTGGTTCGAGACGGGAGCCCTGGGGCCAATGTTATTCCGGGTACACCTGGGATCC
CAGGTCGGGATGGATTCAAAGGAGAAAAGGGGGAATGTCTGAGGGAAAGCTTTGAGGAGTCCT
GGACACCCAACACTACAAGCAGTGTTTCATGGAGTTCATTGAATTATGGCATAGATCTTGGGAAAA
TTGCGGAGTGTACATTTACAAAGATGCGTTCAAATAGTGCTCTAAGAGTTTTGTTCAGTGGCT
CACTTCGGCTAAAATGCAGAAATGCATGCTGTCAGCGTTGGTATTTACATTCAATGGAGCTG
AATGTTTCAGGACCTCTTCCCATTGAAGCTATAATTTATTTGGACCAAGGAAGCCCTGAAATGA
ATTCAACAATTAATATTCATCGCACTTCTTCTGTGGAAGGACTTTGTGAAGGAATTGGTGCTG
GATTAGTGGATGTTGCTATCTGGGTTGGCACTTGTTTCAGATTACCCAAAAGGAGATGCTTCTA
CTGGATGGAATTCAGTTTCTCGCATCATTATTGAAGAACTACCAAAATTAAATGCTTTAATTTT
CATTTGCTACCTCTTTTTTTTATTATGCCTTGGAATGGTTCACTTAAATGACATTTTAAATAAG
TTTATGTATACATCTGAATGAAAAGCAAAGCTAAATATGTTTACAGACCAAAGTGTGATTTC
CACTGTTTTTTAAATCTAGCATTATTCATTTTGCTTCAATCAAAGTGGTTTCAATATTTTTTT
TAGTTGGTTAGAATACTTTCTTCATAGTCACATTCTCTCAACCTATAATTTGGAATATTGTTG
TGGTCTTTTGTTTTTTCTCTTAGTATAGCATTTTTTAAAAAAATATAAAAGCTACCAATCTTG
TACAATTTGTAAATGTTAAGAATTTTTTTTTATATCTGTAAATAAAAATTATTTCCAACA

122/168

FIGURE 122

MRPQGPAASPQRLRGLLLLLLLQLPAPSSASEIPKGKQKAQLRQREVVDLYNGMCLQGPAGVP
GRDGSPGANVIPGTPGIPGRDGFKGEKGECLRESFEESWTPNYKQCSWSSLNYGIDLGKIAEC
TFTKMRSNSALRVLFSGSLRLKCRNACCQRWYFTFNGAECSGPLPIEAIYLDQGSPEMNSTI
NIHRTSSVEGLCEGIGAGLVDVAIWVGTCSDYPKGDASTGWNSVSRIIEELPK

Signal peptide:

amino acids 1-30

Transmembrane domain:

amino acids 195-217

123/168

FIGURE 123

GCTGAGCGTGTGCGCGGTACGGGGCTCTCCTGCCTTCTGGGCTCCAACGCAGCTCTGTGGCTG
AACTGGGTGCTCATCACGGGAAGTGTGGGCTATGGAATACAGATGTGGCAGCTCAGGTAGCC
CCAAATTGCCTGGAAGAATACATCATGTTTTTCGATAAGAAGAAATTGTAGGATCCAGTTTTT
TTTTTAACCGCCCCCTCCCCACCCCCCAAAAAAACTGTAAAGATGCAAAAACGTAATATCCAT
GAAGATCCTATTACCTAGGAAGATTTTGATGTTTTGCTGCGAATGCGGTGTTGGGATTTATTT
GTTCTTGGAGTGTTCTGCGTGGCTGGCAAAGAATAATGTTCCAAAATCGGTCCATCTCCCAAG
GGGTCCAATTTTTCTTCTGGGTGTCAGCGAGCCCTGACTCACTACAGTGCAGCTGACAGGGG
CTGTCATGCAACTGGCCCCCTAAGCCAAAGCAAAAGACCTAAGGACGACCTTTGAACAATACAA
AGGATGGGTTTCAATGTAATTAGGCTACTGAGCGGATCAGCTGTAGCACTGGTTATAGCCCCC
ACTGTCTTACTGACAATGCTTTCTTCTGCCGAACGAGGATGCCCTAAGGGCTGTAGGTGTGAA
GGCAAAATGGTATATTGTGAATCTCAGAAATTACAGGAGATACCCTCAAGTATATCTGCTGGT
TGCTTAGGTTTGTCCCTTCGCTATAACAGCCTTCAAAAACCTTAAGTATAATCAATTTAAAGGG
CTCAACCAGCTCACCTGGCTATACTTGACCATAACCATATCAGCAATATTGACGAAAATGCT
TTTAATGGAATACGCAGACTCAAAGAGCTGATTCTTAGTTCCAATAGAATCTCCTATTTTCTT
ACAATACCTTCAGACCTGTGACAAATTTACGGAAGTGGATCTGTCCTATAATCAGCTGCAT
TCTCTGGGATCTGAACAGTTTCGGGGCTTGCGGAAGCTGCTGAGTTTACATTTACGGTCTAAC
TCCCTGAGAACCATCCCTGTGCGAATATTCCAAGACTGCCGCAACCTGGAAGTTTTGGACCTG
GGATATAACCGGATCCGAAGTTTAGCCAGGAATGTCTTTGCTGGCATGATCAGACTCAAAGAA
CTTCACCTGGAGCACAATCAATTTTCCAAGCTCAACCTGGCCCTTTTTCCAAGGTTGGTCAGC
CTTCAGAACCTTTACTTGCAGTGGAATAAAATCAGTGTCATAGGACAGACCATGTCCTGGACC
TGGAGCTCCTTACAAAGGCTTGATTTATCAGGCAATGAGATCGAAGCTTTCAGTGGACCCAGT
GTTTTCCAGTGTGTCCCGAATCTGCAGCGCCTCAACCTGGATTCCAACAAGCTCACATTTATT
GGTCAAGAGATTTTGGATTCTTGGATATCCCTCAATGACATCAGTCTTGCTGGGAATATATGG
GAATGCAGCAGAAATATTTGCTCCCTTGTAAGTGGCTGAAAAGTTTTAAAGGTCTAAGGGAG
AATACAATTATCTGTGCCAGTCCCAAAGAGCTGCAAGGAGTAAATGTGATCGATGCAGTGAAG
AACTACAGCATCTGTGGCAAAGTACTACAGAGAGGTTTGATCTGGCCAGGGCTCTCCCAAAG
CCGACGTTTAAGCCCAAGCTCCCCAGGCCGAAGCATGAGAGCAAACCCCTTTGCCCCCGACG
GTGGGAGCCACAGAGCCCGGCCAGAGACCGATGCTGACGCCGAGCACATCTCTTTCCATAAA
ATCATCGCGGGCAGCGTGGCGCTTTTCCTGTCCGTGCTCGTCATCCTGCTGGTTATCTACGTG
TCATGGAAGCGGTACCCTGCGAGCATGAAGCAGCTGCAGCAGCGCTCCCTCATGCGAAGGCAC
AGGAAAAAGAAAAGACAGTCCCTAAAGCAAATGACTCCCAGCACCCAGGAATTTTATGTAGAT
TATAAACCCACCAACACGGAGACCAGCGAGATGCTGCTGAATGGGACGGGACCCTGCACCTAT
AACAAATCGGGCTCCAGGGAGTGTGAGGTATGAACCATTTGTGATAAAAAGAGCTCTTAAAAGC
TGGGAAATAAGTGGTGCTTTATTGAACTCTGGTGACTATCAAGGGAACGCGATGCCCCCCTC
CCCTTCCCTCTCCCTCTCACTTTGGTGGCAAGATCCTTCCTTGTCGGTTTTAGTGCATTCATA
ATACTGGTCATTTTCCTCTCATACATAATCAACCCATTGAAATTTAAATACCACAATCAATGT
GAAGCTTGAAGTCCGGTTTAATATAATACCTATTGTATAAGACCCTTTACTGATTCCATTAAT
GTCGCATTTGTTTTAAGATAAACTTCTTTCATAGGTAAAAA

124/168

FIGURE 124

MGFNVIRLLSGSAVALVIAPT VLLTMLSSAERGCPKGCRCEGKMVYCESQKLQEIPSSISAGC
LGLSLRYNSLQKLKYNQFKGLNQLTWLYLDHNNHISNIDENAFNGIRRLKELILSSNRISYFLN
NTFRPVTNLRNLDLSYNQLHSLGSEQFRGLRKLLSLHLRSNSLRTIPVRI FQDCRNLELLDLG
YNRIRSLARNVFAGMIRLKEHLEHNQFSKLNLALEPRLVSLQONLYLQWNKISVIGQTMSWTW
SSLQRLDLSGNEIEAFSGPSVFQCVPNLQRLNLD SNKLTFIGQEILDSWISLNDISLAGNIWE
CSRNICSLVNWLKSFKGLRENTIICASPKELOGVNVIDAVKNYSICGKSTTERFDLARALPKP
TFKPKLPRPKHESKPPLPPTVGATEPGPETDADAEHISFHKIIAGSVALFLSVLVILLVIYVS
WKRYPASMKQLQQRSLMRRHRKKKRQSLKQMTPTSTQEFYVDYKPTNTETSEM LLLNGTGPCTYN
KSGSRECEV

Important features of the protein:**Signal peptide:**

amino acids 1-33

Transmembrane domain:

amino acids 420-442

N-glycosylation sites.

amino acids 126-129, 357-360, 496-499, 504-507

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 465-468

Tyrosine kinase phosphorylation site.

amino acids 136-142

N-myristoylation sites.

amino acids 11-16, 33-38, 245-250, 332-337, 497-502, 507-512

125/168

FIGURE 125

CCGTTATCGTCTTGCGCTACTGCTGAATGTCCGTCCCGGAGGAGGAGGAGAGGCTTTTGCCGC
TGACCCAGAGATGGCCCCGAGCGAGCAAATTCCTACTGTCCGGCTGCGCGGCTACCGTGGCCG
AGCTAGCAACCTTTCCCCTGGATCTCACAAAACTCGACTCCAAATGCAAGGAGAAGCAGCTC
TTGCTCGGTTGGGAGACGGTGCAAGAGAATCTGCCCCCTATAGGGGAATGGTGCGCACAGCCC
TAGGGATCATTGAAGAGGAAGGCTTTCTAAAGCTTTGGCAAGGAGTGACACCCGCCATTTACA
GACACGTAGTGTATTCTGGAGGTCGAATGGTCACATATGAACATCTCCGAGAGGTTGTGTTTG
GCAAAAGTGAAGATGAGCATTATCCCCTTTGGAAATCAGTCATTGGAGGGATGATGGCTGGTG
TTATTGGCCAGTTTTTAGCCAATCCAACCTGACCTAGTGAAGGTTTCAGATGCAAATGGAAGGAA
AAAGGAACTGGAAGGAAAACCATTCGCGATTTTCGTGGTGTACATCATGCATTTGCAAAAATCT
TAGCTGAAGGAGGAATACGAGGGCTTTGGGCAGGCTGGGTACCCAATATACAAAGAGCAGCAC
TGGTGAATATGGGAGATTTAACCCTTATGATACAGTGAAACACTACTTGGTATTGAATACAC
CACTTGAGGACAATATCATGACTCACGGTTTATCAAGTTTATGTTCTGGACTGGTAGCTTCTA
TTCTGGGAACACCAGCCGATGTCATCAAAGCAGAATAATGAATCAACCACGAGATAAACAAG
GAAGGGGACTTTTGTATAAATCATCGACTGACTGCTTGATTCAGGCTGTTCAAGGTGAAGGAT
TCATGAGTCTATATAAAGGCTTTTACCATCTTGGCTGAGAATGACCCCTTGGTCAATGGTGT
TCTGGCTTACTTATGAAAAAATCAGAGAGATGAGTGGAGTCAGTCCATTTTAA

126/168

FIGURE 126

MSVPEEEERLLPLTQRWPRASKFLLSGCAATVAELATFPLDLTKTRLQMQGEAALARLGDGAR
ESAPYRGMVRTALGIIIEEGFLKLWQGVTPAIYRHVVYSGGRMVITYEHLREVVFGKSEDEHYP
LWKSIVIGGMMAGVIGQFLANPTDLVKVQMOMEGKRKLEGKPLRFRGVHHAFAKILAEGGIRGL
WAGWVPNIQRAALVNMGDLTTYDTVKHYLVNLNTPLEDNIMTHGLSSLCSGLVASILGTPADVI
KSRIMNQPRDKQGRGLLYKSSTDCLIQAVQGEFMSLYKGFLPSWLRMTPWSMVFWLTYEKIR
EMSGVSPF

Transmembrane domains:

amino acids 25-38, 130-147, 233-248

127/168

FIGURE 127

CGCGGATCGGACCCAAGCAGGTCGGCGGGCGGCGGCAGGAGAGCGGCCGGGCGTCAGCTCCTCG
ACCCCCGTGTCGGGCTAGTCCAGCGAGGCGGACGGGGCGGCGTGGGCCCATGGCCAGGCCCGGC
ATGGAGCGGTGGCGCGACCGGCTGGCGCTGGTGACGGGGGGCCTCGGGGGGCATCGGCGCGGCC
GTGGCCCCGGGGCCCTGGTCCAGCAGGGACTGAAGGTGGTGGGCTGCGCCCCGCACTGTGGGCAAC
ATCGAGGAGCTGGCTGCTGAATGTAAGAGTGCAGGCTACCCCGGGACTTTGATCCCCTACAGA
TGTGACCTATCAAATGAAGAGGACATCCTCTCCATGTTCTCAGCTATCCGTTCTCAGCACAGC
GGTGTAGACATCTGCATCAACAATGCTGGCTTGGCCCCGGCCTGACACCCTGCTCTCAGGCAGC
ACCAGTGGTTGGAAGGACATGTTCAATGTGAACGTGCTGGCCCTCAGCATCTGCACACGGGAA
GCCTACCAGTCCATGAAGGAGCGGAATGTGGACGATGGGCACATCATTAAACATCAATAGCATG
TCTGGCCACCGAGTGTTACCCCTGTCTGTGACCCACTTCTATAGTGCCACCAAGTATGCCGTC
ACTGCGCTGACAGAGGGACTGAGGCAAGAGCTTCGGGAGGCCAGACCCACATCCGAGCCACG
TGCATCTCTCCAGGTGTGGTGGAGACACAATTGCGCTTCAAACCTCCACGACAAGGACCCTGAG
AAGGCAGCTGCCACCTATGAGCAAATGAAGTGTCTCAAACCCGAGGATGTGGCCGAGGCTGTT
ATCTACGTCCTCAGCACCCCCGCACACATCCAGATTGGAGACATCCAGATGAGGCCACGGAG
CAGGTGACCTAGTGACTGTGGGAGCTCCTCCTTCCCTCCCCACCCTTCATGGCTTGCCTCCTG
CCTCTGGATTTTAGGTGTTGATTTCTGGATCACGGGATACCACTTCCTGTCCACACCCCGACC
AGGGGCTAGAAAATTTGTTTGAGATTTTATATCATCTTGTCAAATTGCTTCAGTTGTAAATG
TGAAAAATGGGCTGGGGAAAGGAGGTGGTGTCCCTAATTGTTTTACTTGTTAACTTGTTCTTG
TGCCCCCTGGGCACTTGGCCTTTGTCTGCTCTCAGTGTCTTCCCTTTGACATGGGAAAGGAGTT
GTGGCCAAAATCCCCATCTTCTTGACCTCAACGTCTGTGGCTCAGGGCTGGGGTGGCAGAGG
GAGGCCTTCACCTTATATCTGTGTTGTTATCCAGGGCTCCAGACTTCCTCCTCTGCCTGCCCC
ACTGCACCCTCTCCCCCTTATCTATCTCCTTCTCGGCTCCCCAGCCCAGTCTTGGCTTCTTGT
CCCCTCCTGGGGTCATCCCTCCACTCTGACTCTGACTATGGCAGCAGAACACCAGGGCCTGGC
CCAGTGGATTTTCATGGTGATCATTAAAAAAGAAAAATCGCAACCAAAAAAAAAAAAAA

128/168

FIGURE 128

MARPGMERWRDRLALVTGASGGIGA A VARALVQQGLKVVG CARTVGNIEELAAECKSAGYPGT
LIPYRCDLSNEEDILSMFSAIRSQHSGVDICINNAGLARPD TLLSGSTSGWKDMFNVNVLALS
ICTREAYQSMKERNVDDGHIININSMGHRVLPLSVTHFY SATKYAVTALTEGLRQELREAQT
HIRATCISPGVVETQFAFKLHDKDPEKAAATYEQMKCLKPEDVAEAVIYVLSTPAHIQIGDIQ
MRPTEQVT

Important features of the protein:

Signal peptide:

amino acids 1-17

N-myristoylation sites.

amino acids 18-24, 21-27, 22-28, 24-30, 40-46, 90-96, 109-115,
199-205

Short-chain alcohol dehydrogenase.

amino acids 30-42, 104-114

129/168

FIGURE 129

AACTTCTACATGGGCCTCCTGCTGCTGGTGCTCTTCCTCAGCCTCCTGCCGGTGGCCTACACC
ATCATGTCCCTCCCACCCTCCTTTGACTGCGGGCCGTTCAAGTGCAGAGTCTCAGTTGCCCGG
GAGCACCTCCCCTCCCGAGGCAGTCTGCTCAGAGGGCCTCGGCCCAGAATTCCAGTTCTGGTT
TCATGCCAGCCTGTAAAAGGCCATGGAACCTTGGGTGAATCACCGATGCCATTTAAGAGGGTT
TTCTGCCAGGATGGAAATGTTAGGTCGTTCTGTGTCTGCGCTGTTCAATTCAGTAGCCACCAG
CCACCTGTGGCCGTTGAGTGCTTGAAATTGAGGGAAGTGAAGAAAATTAATTTCTCATGTATTTTT
CTCATTTATTTATTAATTTTTTAAGTATAGTTGTACATATTTGGGGGTACATGTGATATTTGG
ATACATGTATACAATATATAATGATCAAATCAGGGTAACTGGGATATCCATCACATCAAACAT
TTATTTTTTTATTCTTTTTTAGACAGAGTCTCACTCTGTCACCCAGGCTGGAGTGCAGTGGTGCC
ATCTCAGCTTACTGCAACCTCTGCCTGCCAGGTTCAAGCGATTCTCATGCCTCCACCTCCCAA
GTAGCTGGGACTACAGGCATGCACCACAATGCCCAACTAATTTTTGTATTTTTAGTAGAGACG
GGGTTTTGCCATGTTGCCCAGGCTGGCCTTGAACTCCTGGCCTCAAACAATCCACTTGCCTCG
GCCTCCCAAAGTGTTATGATTACAGGCGTGAGCCACCGTGCCTGGCCTAAACATTTATCTTTT
CTTTGTGTTGGGAACCTTTGAAATTATACAATGAATTATTGTAACTGTCATCTCCCTGCTGTG
CTATGGAACACTGGGACTTCTTCCCTCTATCTAACTGTATATTTGTACCAGTTAACCAACCGT
ACTTCATCCCCACTCCTCTCTATCCTTCCCAACCTCTGATCACCTCATTCTACTCTCTACCTC
CATGAGATCCACTTTTTTAGCTCCCACATGTGAGTAAGAAAATGCAATATTTGTCTTTCTGTG
CCTGGCTTATTTCACTTAACATAATGACTTCCTGTTCCATCCATGTTGCTGCAAATGACAGGA
TTTCGTTCTTAATTTCAATTAAATAACCACACATGGCAAAAA

WO 01/16318

PCT/US00/23328

130/168

FIGURE 130

MGLLLLVLFLSLLPVAYTIMSLPPSFDCGPFRCRVSVAREHLPSRGSLLRGPRPRI PVLVSCQ
PVKGHGTLGESPMPEFKRVFCQDGNVRSFCVCAVHFSSHQPPVAVECLK

Important features of the protein:

Signal peptide:

amino acids 1-18

N-myristoylation site.

amino acids 86-92

Zinc carboxypeptidases, zinc-binding region 2 signature.

amino acids 68-79

131/168

FIGURE 131

TTCTGAAGTAACGGAAGCTACCTTGTATAAAGACCTCAACACTGCTGACCATGATCAGCGCAG
CCTGGAGCATCTTCCTCATCGGGACTAAAATTGGGCTGTTCCCTCAAGTAGCACCTCTATCAG
TTATGGCTAAATCCTGTCCATCTGTGTGTCGCTGCGATGCGGGTTTCATTTACTGTAATGATC
GCTTTCTGACATCCATTCCAACAGGAATACCAGAGGATGCTACAACCTCTCTACCTTCAGAACA
ACCAAATAAATAATGCTGGGATTCCTTCAGATTTGAAAAACTTGCTGAAAGTAGAAAGAATAT
ACCTATAACCACAACAGTTTAGATGAATTCCTACCAACCTCCCAAAGTATGTAAAAGAGTTAC
ATTTGCAAGAAAATAACATAAGGACTATCACTTATGATTCACTTTCAAAAATTCCCTATCTGG
AAGAATTACATTTAGATGACAACCTCTGTCTCTGCAGTTAGCATAGAAGAGGGAGCATTCCGAG
ACAGCAACTATCTCCGACTGCTTTTCCTGTCCCGTAATCACCTTAGCACAATTCCCTGGGGTT
TGCCCAGGACTATAGAAGAACTACGCTTGGATGATAATCGCATATCCACTATTTTCATCACCAT
CTCTTCAAGGTCTCACTAGTCTAAAACGCCTGGTTCTAGATGGAAACCTGTTGAACAATCATG
GTTTAGGTGACAAAGTTTCTTCAACCTAGTTAATTTGACAGAGCTGTCCCTGGTGCGGAATT
CCCTGACTGCTGCACCAGTAAACCTTCCAGGCACAAACCTGAGGAAGCTTTATCTTCAAGATA
ACCACATCAATCGGGTGCCCCCAAATGCTTTTTCTTATCTAAGGCAGCTCTATCGACTGGATA
TGTC CAATAATAACCTAAGTAATTTACCTCAGGGTATCTTTGATGATTTGGACAATATAACAC
AACTGATTCTTCGCAACAATCCCTGGTATTGCGGGTGCAAGATGAAATGGGTACGTGACTGGT
TACAATCACTACCTGTGAAGGTCAACGTGCGTGGGCTCATGTGCCAAGCCCCAGAAAAGGTTC
GTGGGATGGCTATTAAGGATCTCAATGCAGAACTGTTTGATTGTAAGGACAGTGGGATTGTAA
GCACCATT CAGATAACCACTGCAATACCCAACACAGTGTATCCTGCCCAAGGACAGTGGCCAG
CTCCAGTGACC AAACAGCCAGATATTAAGAACCCCAAGCTCACTAAGGATCAACAAACCACAG
GGAGTCCCTCAAGAAAAACAATTACAATTACTGTGAAGTCTGTCACCTCTGATACCATT CATA
TCTCTTGGAACTTGCTCTACCTATGACTGCTTTGAGACTCAGCTGGCTTAAACTGGGCCATA
GCCCCGGCATT TTGGATCTATAACAGAAACAATTGTAACAGGGGAACGCAGTGAGTACTTGGTCA
CAGCCCTGGAGCCTGATTCACCCTATAAAGTATGCATGGTTCCCATGGAAACCAGCAACCTCT
ACCTATTTGATGAAACTCCTGTTTGTATTGAGACTGAAACTGCACCCCTTCGAATGTACAACC
CTACAACCACCCTCAATCGAGAGCAAGAGAAAGAACCTTACAAAACCCCAATTTACCTTTGG
CTGCCATCATTGGTGGGGCTGTGGCCCTGGTTACCATTGCCCTTCTTGCTTTAGTGTGTTGGT
ATGTT CATAGGAATGGATCGCTCTTCTCAAGGAACTGTGCATATAGCAAAGGGAGGAGAAGAA
AGGATGACTATGCAGAAGCTGGCACTAAGAAGGACAACCTCTATCCTGGAAATCAGGGAAACTT
CTTTTCAGATGTTACCAATAAGCAATGAACCCATCTCGAAGGAGGAGTTTGTAATACACACCA
TATTT CCTCCTAATGGAATGAATCTGTACAAAACAATCACAGTGAAAGCAGTAGTAACCGAA
GCTACAGAGACAGTGGTATTCAGACTCAGATCACTCACACTCATGATGCTGAAGGACTCACA
GCAGACTTGTGTTTTGGGTTTTTTAAACCTAAGGGAGGTGATGGT

132/168

FIGURE 132

MISAAWSIFLIGTKIGLFLQVAPLSVMAKSCPSVCRC DAGFIYCNDRFLTSIPTGIPEDATTL
YLQNNQINNAGIPSDLKNLLKVERIYLYHNSLDEFPTNLPKYVKELHLQENNIRTITYDSLK
IPYLEELHLDDNSVSAVSIEEGA FRDSNYLRLLFLSRNHLSTIPWGLPRTIEELRLDDNRIST
ISSPSLQGLTSLKRLVLDGNLLNNHGLGDKVFFNLVNLTELSLVRNSLTAAPVNLPGTNLRKL
YLQDNHINRVPPNAFSYLRQLYRLDMSNNNLSNLPQGI FDDL DNITQLILRNNPWYCGCKMKW
VRDWLQSLPVKVNVRGLMCQAPEKVRGMAIKDLNAELFDCKDSGIVSTIQITTAIPNTVYPAQ
GQWPAPVTKQPDIKNPKLTKDQOTTGSPSRKTITITVKS VTS DTI HISWKLALPMTALRLSWL
KLGHSPAFGSITETIVTGERSEYLVTALEPDSPYKVCMPMETS NLYLFDETPVCIETETAPL
RMYNPTTTLNREQEKEPYKNPNLPLAAIIGGAVALVTIAL LALVCWYVHRNGSLFSRNCAYSK
GRRRKDDYAEAGTKKDNSILEIRETSFQMLPISNEPISKEEFVIHTIFPPNGMNLYKNNHSES
SSNRSYRDSGIPDSDHSHS

Important features of the protein:**Signal peptide:**

amino acids 1-28

Transmembrane domain:

amino acids 531-552

N-glycosylation sites.

amino acids 226-229, 282-285, 296-299, 555-558, 626-629, 633-636

Tyrosine kinase phosphorylation site.

amino acids 515-522

N-myristoylation sites.amino acids 12-17, 172-177, 208-213, 359-364, 534-539, 556-561,
640-645**Amidation site.**

amino acids 567-570

Leucine zipper pattern.

amino acids 159-180

Phospholipase A2 aspartic acid active site.

amino acids 34-44

133/168

FIGURE 133

CCGTCATCCCCCTGCAGCCACCCTTCCCAGAGTCCTTTGCCCAGGCCACCCCAGGCTTCTTGG
CAGCCCTGCCGGGGCCACTTGTCTTCATGTCTGCCAGGGGGAGGTGGGAAGGAGGTGGGAGGAG
GGCGTGCAGAGGCAGTCTGGGCTTGGCCAGAGCTCAGGGTGCTGAGCGTGTGACCAGCAGTGA
GCAGAGGCCGGCCATGGCCAGCCTGGGGCTGCTGCTCCTGCTCTTACTGACAGCACTGCCACC
GCTGTGGTCCTCCTCACTGCCTGGGCTGGACACTGCTGAAAGTAAAGCCACCATTGCAGACCT
GATCCTGTCTGCGCTGGAGAGAGCCACCGTCTTCCTAGAACAGAGGCTGCCTGAAATCAACCT
GGATGGCATGGTGGGGGTCCGAGTGCTGGAAGAGCAGCTAAAAAGTGTCGGGGAGAAGTGGGC
CCAGGAGCCCCCTGCTGCAGCCGCTGAGCCTGCGCGTGGGGATGCTGGGGGAGAAGCTGGAGGC
TGCCATCCAGAGATCCCTCCACTACCTCAAGCTGAGTGATCCCAAGTACCTAAGAGAGTTCCA
GCTGACCCTCCAGCCCCGGGTTTTGGAAGCTCCCACATGCCTGGATCCACACTGATGCCTCCTT
GGTGTACCCACGTTTCGGGCCCCAGGACTCATTCTCAGAGGAGAGAAGTGACGTGTGCCTGGT
GCAGCTGCTGGGAACCGGGACGGACAGCAGCGAGCCCTGCGGCCTCTCAGACCTCTGCAGGAG
CCTCATGACCAAGCCCCGGCTGCTCAGGCTACTGCCTGTCCCACCAACTGCTCTTCTTCCTCTG
GGCCAGAATGAGGGGATGCACACAGGGACCACTCCAACAGAGCCAGGACTATATCAACCTCTT
CTGCGCCAACATGATGGACTTGAACCGCAGAGCTGAGGCCATCGGATACGCCTACCCTACCCG
GGACATCTTCATGGAAAACATCATGTTCTGTGGAATGGGCGGCTTCTCCGACTTCTACAAGCT
CCGGTGGCTGGAGGCCATTCTCAGCTGGCAGAAACAGCAGGAAGGATGCTTCGGGGAGCCTGA
TGCTGAAGATGAAGAATTATCTAAAGCTATTCAATATCAGCAGCATTTTTTCGAGGAGAGTGAA
GAGGCGAGAAAAACAATTTCCAGATTCTCGCTCTGTTGCTCAGGCTGGAGTACAGTGGCGCAA
TCTCGGCTCACTGCAACCTTTGCCTCCTGGGTTCAAGCAATTCTCTTGCCTCATCCTCCCGAG
TAGCTGGGACTACAGGAGCGTGCCACCATACTGGCTAATTTTTTATATTTTTTTTAGTAGAGAC
AGGGTTTCATCATGTTGCTCATGCTGGTCTCGAACTCCTGATCTCAAGAGATCCGCCCACCTC
AGGCTCCCAAAGTGTGGGATTATTAGGTGTGAGCCACCGTGTCTGGCTGAAAAGCACTTTCAAA
GAGACTGTGTTGAATAAAGGGCCAAGGTTCTTGCCACCCAGCACTCATGGGGGCTCTCTCCCC
TAGATGGCTGCTCCTCCCACAACACAGCCACAGCAGTGGCAGCCCTGGGTGGCTTCCTATACA
TCCTGGCAGAATAACCCCCAGCAAACAGAGAGCCACACCCATCCACACCGCCACCACCAAGCA
GCCGCTGAGACGGACGGTTCCATGCCAGCTGCCTGGAGGAGGAACAGACCCCTTTAGTCCTCA
TCCCTTAGATCCTGGAGGGCACGGATCACATCCTGGGAAGAAGGCATCTGGAGGATAAGCAAA
GCCACCCCGACACCCAATCTTGGAAGCCCTGAGTAGGCAGGGCCAGGGTAGGTGGGGGCGGG
AGGGACCCAGGTGTGAACGGATGAATAAAGTTCAACTGCAACTGAAAAAAAAAAAA

134/168

FIGURE 134

MSARGRWEGGRRACRGSGLLARAQGAERVTSSEQRPAMASLGLLLLLLLLLTALPPLWSSSLPG
LDTAESKATIADLILSALERATVFLEQRLPEINLDGMVGVRVLEEQLKSVREKWAQEPLLQPL
SLRVGMLGEKLEAAIQRS LHYLKLSDPKYLREFQLTLQPGFWKLPHAWIHTDASLVYPTFGPQ
DSFSEERSDVCLVQLLGTGTDSEPCGLSDLCRSLMTKPGCSGYCLSHQLLFFLWARMRGCTQ
GPLQQSQDYINLFCANMMDLNRRAEAIGYAYPTRDIFMENIMFCGMGGFSDFYKLRWLEAILS
WQKQQEGCFGEPPDAEDEELSKAIQYQQHFSSRRVKRREKQFPDSRSVAQAGVQWRNLGSLQPLP
PGFKQFSC LILPSSWDYRSVPPYLANFYIFLVETGFHHVAHAGLELLISRDPPTSGSQSVGL

Important features of the protein:

Signal peptide:

amino acids 1-26

Transmembrane domain:

amino acids 39-56

Tyrosine kinase phosphorylation sites.

amino acids 149-156, 274-282

N-myristoylation sites.

amino acids 10-16, 20-26, 63-69, 208-214

Amidation site.

amino acids 10-14

Glycoprotein hormones beta chain signature 1.

amino acids 230-237

135/168

FIGURE 135

GGTCTGAGTGCAGAGCTGCTGTCATGGCGGCCGCTCTGTGGGGCTTCTTTCCCGTCCTGCTGC
TGCTGCTGCTATCGGGGGATGTCCAGAGCTCGGAGGTGCCCGGGGCTGCTGCTGAGGGATCGG
GAGGGAGTGGGGTCGGCATAGGAGATCGCTTCAAGATTGAGGGGCGTGCAAGTTGTTCCAGGGG
TGAAGCCTCAGGACTGGATCTCGGCGGCCCGAGTGCTGGTAGACGGAGAAGAGCACGTCGGTT
TCCTTAAGACAGATGGGAGTTTTGTGGTTCATGATATACCTTCTGGATCTTATGTAGTGGAAG
TTGTATCTCCAGCTTACAGATTTGATCCCGTTCGAGTGGATATCACTTCGAAAGGAAAAATGA
GAGCAAGATATGTGAATTACATCAAAACATCAGAGGTTGTCAGACTGCCCTATCCTCTCCAAA
TGAAATCTTCAGGTCCACCTTCTTACTTTATTAAAAGGGAATCGTGGGGCTGGACAGACTTTC
TAATGAACCCAATGGTTATGATGATGGTTCTTCCTTTATTGATATTTGTGCTTCTGCCTAAAG
TGGTCAACACAAGTGATCCTGACATGAGACGGGAAATGGAGCAGTCAATGAATATGCTGAATT
CCAACCATGAGTTGCCTGATGTTTCTGAGTTCATGACAAGACTCTTCTCTTCAAAATCATCTG
GCAAATCTAGCAGCGGCAGCAGTAAAACAGGCAAAAGTGGGGCTGGCAAAAGGAGGTAGTCAG
GCCGTCCAGAGCTGGCATTTCACAAACACGGCAACACTGGGTGGCATCCAAGTCTTGGA AAA
CCGTGTGAAGCAACTACTATAAACTTGAGTCATCCCGACGTTGATCTCTTACA ACTGTGTATGTT
AACTTTTTAGCACATGTTTTGTACTTGGTACACGAGAAAACCCAGCTTTCATCTTTTGTCTGT
ATGAGGTCAATATTGATGTC ACTGAATTAATTACAGTGTCCTATAGAAAATGCCATTAATAAA
TTATATGA ACTACTATACATTATGTATATTAATTA AAACATCTTAATCCAGAAATCAAAAAA
AAAAAAAAAAAAAAAAAAAAA

136/168

FIGURE 136

MAAALWGFFPVLLLLLLSGDVQSSEVPGAAAEGSGGSGVGIGDRFKIEGRAVVPGVKPQDWIS
AARVLVDGEEHVGFLKTDGSFVVHDIPSGSYVVEVVS PAYRFDPVRVDITSKGKMRARYVNYI
KTSEVVRLPYPLQMKSSGPPSYFIKRESWGWTDFLMNPMVMMVLPLLI FVLLPKVVNTSDPD
MRREMEQSMNMLNSNHELDPVSEFMTRLFSSKSSGKSSSGSSKTGKSGAGKRR

Important features of the protein:

Signal sequence:

amino acids 1-23

Transmembrane domain:

amino acids 161-182

N-glycosylation site.

amino acids 184-187

Glycosaminoglycan attachment sites.

amino acids 37-40, 236-239

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 151-154

N-myristoylation sites.

amino acids 33-38, 36-41, 38-44, 229-234

Amidation site.

amino acids 238-241

ATP/GTP-binding site motif A (P-loop).

amino acids 229-236

137/168

FIGURE 137

GATGGCGCAGCCACAGCTTCTGTGAGATTCTGATTTCTCCCCAGTTCCCCTGTGGGTCTGAGGG
GACCAGAAGGGTGAGCTACGTTGGCTTTCTGGAAGGGGAGGCTATATGCGTCAATTCCCCAAA
ACAAGTTTTGACATTTCCCCTGAAATGTCATTCTCTATCTATTCACTGCAAGTGCCTGCTGTT
CCAGGCCTTACCTGCTGGGCACTAACGGCGGAGCCAGGATGGGGACAGAATAAAGGAGCCACG
ACCTGTGCCACCAACTCGCACTCAGACTCTGAACTCAGACCTGAAATCTTCTCTTCACGGGAG
GCTTGGCAGTTTTTCTTACTCCTGTGGTCTCCAGATTTCAAGGCCTAAGATGAAAGCCTCTAGT
CTTGCCTTCAGCCTTCTCTCTGCTGCGTTTTATCTCCTATGGACTCCTTCCACTGGACTGAAG
ACACTCAATTTGGGAAGCTGTGTGATCGCCACAAACCTTCAGGAAATACGAAATGGATTTTCT
GAGATACGGGGCAGTGTGCAAGCCAAAGATGGAAACATTGACATCAGAATCTTAAGGAGGACT
GAGTCTTTGCAAGACACAAAGCCTGCGAATCGATGCTGCCTCCTGCGCCATTTGCTAAGACTC
TATCTGGACAGGGTATTTAAAACTACCAGACCCCTGACCATTATACTCTCCGGAAGATCAGC
AGCCTCGCCAATTCCTTTCTTACCATCAAGAAGGACCTCCGGCTCTCTCATGCCCACATGACA
TGCCATTGTGGGGAGGAAGCAATGAAGAAATACAGCCAGATTCTGAGTCACTTTGAAAAGCTG
GAACCTCAGGCAGCAGTTGTGAAGGCTTTGGGGGAACTAGACATTCTTCTGCAATGGATGGAG
GAGACAGAATAGGAGGAAAGTGATGCTGCTGCTAAGAATATTCGAGGTCAAGAGCTCCAGTCT
TCAATACCTGCAGAGGAGGCATGACCCCAAACCACCATCTCTTTACTGTACTAGTCTTGTGCT
GGTCACAGTGTATCTTATTTATGCATTACTTGCTTCCTTGCATGATTGTCTTTATGCATCCCC
AATCTTAATTGAGACCATACTTGTATAAGATTTTTTGTAATATCTTTCTGCTATTGGATATATT
TATTAGTTAATATATTTATTTATTTTTTTGCTATTTAATGTATTTATTTTTTTTACTTGGACATG
AACTTTAAAAAAATTCACAGATTATATTTATAACCTGACTAGAGCAGGTGATGTATTTTTTAT
ACAGTAAAAAAAACCTTGTAATTCTAGAAGAGTGGCTAGGGGGGTATTTCATTTGTAT
TCAACTAAGGACATATTTACTCATGCTGATGCTCTGTGAGATATTTGAAATTGAACCAATGAC
TACTTAGGATGGGTGTGGAATAAGTTTTGATGTGGAATTGCACATCTACCTTACAATTACTG
ACCATCCCCAGTAGACTCCCCAGTCCCATAATTGTGTATCTTCCAGCCAGGAATCCTACACGG
CCAGCATGTATTTCTACAAATAAAGTTTTCTTTGCATACCAAAAAAAAAAAAAAAAAAAAAA

138/168

FIGURE 138

MRQFPKTSFDISPMSFSIYSLQVPAVPGLTCWALTAEPGWGQNKGATTCATNSHSDSELRPE
IFSSREAWQFFLLLWSPDFRPKMKASSLAFSLLSAAFYLLWTPSTGLKTLNLGSCVIATNLQE
IRNGFSEIRGSVQAKDGNIDIRILRRTESLQDTKPANRCCLLRHLLRLYLDRVFKNYQTPDHY
TLRKISSLANSFLLTIKKDLRLSHAHMTCHCGEAMKKYSQILSHFEKLEPQAAVVKALGELDI
LLQWMEETE

Important features of the protein:

Signal peptide:

amino acids 1-42

cAMP- and cGMP-dependent protein kinase phosphorylation sites.

amino acids 192-195, 225-228

N-myristoylation sites.

amino acids 42-47, 46-51, 136-141

139/168

FIGURE 139

CCTGGAGCCGGAAGCGCGGCTGCAGCAGGGCGAGGCTCCAGGTGGGGTCGGTTCCGCATCCAG
CCTAGCGTGTCCACGATGCGGCTGGGCTCCGGGACTTTTCGCTACCTGTTGCGTAGCGATCGAG
GTGCTAGGGATCGCGGTCTTCCTTCGGGGATTCTTCCCGGCTCCCGTTCTGTTCTCTGCCAGA
GCGGAACACGGAGCGGAGCCCCAGCGCCCGAACCCTCGGCTGGAGCCAGTTCTAACTGGACC
ACGCTGCCACCACCTCTCTTCAGTAAAGTTGTTATTGTTCTGATAGATGCCTTGAGAGATGAT
TTTGTGTTTGGGTCAAAGGGTGTGAAATTTATGCCCTACACAACCTTACCTTGTGGAAAAAGGA
GCATCTCACAGTTTTGTGGCTGAAGCAAAGCCACCTACAGTTACTATGCCTCGAATCAAGGCA
TTGATGACGGGGAGCCTTCCTGGCTTTGTGCGACGTCATCAGGAACCTCAATTCTCCTGCACTG
CTGGAAGACAGTGTGATAAGACAAGCAAAAGCAGCTGGAAAAAGAATAGTCTTTTATGGAGAT
GAAACCTGGGTAAATTATTCCCAAAGCATTTTTGTGGAATATGATGGAACAACCTCATTTTTTC
GTGTCAGATTACACAGAGGTGGATAATAATGTCACGAGGCATTTGGATAAAGTATTAAAAAGA
GGAGATTGGGACATATTAATCCTCCACTACCTGGGGCTGGACCACATTGGCCACATTTTCAGGG
CCCAACAGCCCCCTGATTGGGCAGAAGCTGAGCGAGATGGACAGCGTGCTGATGAAGATCCAC
ACCTCACTGCAGTCGAAGGAGAGAGAGACGCCTTTACCCAATTTGCTGGTTCTTTGTGGTGAC
CATGGCATGTCTGAAACAGGAAGTCACGGGGCCTCCTCCACCGAGGAGGTGAATACACCTCTG
ATTTTAATCAGTTCTGCGTTTGAAAGGAAACCCGGTGATATCCGACATCCAAAGCACGTCCAA
TAGACGGATGTGGCTGCGACACTGGCGATAGCACTTGGCTTACCGATTCCAAAAGACAGTGTA
GGGAGCCTCCTATTCCCAGTTGTGGAAGGAAGACCAATGAGAGAGCAGTTGAGATTTTTTACAT
TTGAATACAGTGCAGCTTAGTAAACTGTTGCAAGAGAATGTGCCGTCATATGAAAAAGATCCT
GGGTTTGAGCAGTTTAAAATGTCAGAAAGATTGCATGGGAACTGGATCAGACTGTACTTGGAG
GAAAAGCATTGAGAAGTCCTATTCAACCTGGGCTCCAAGGTTCTCAGGCAGTACCTGGATGCT
CTGAAGACGCTGAGCTTGTCCCTGAGTGCACAAGTGGCCCAGTTCTCACCCCTGCTCCTGCTCA
GCGTCCCACAGGCACTGCACAGAAAGGCTGAGCTGGAAGTCCCAGTGTCTCCTGGGTTTTT
CTCTGCTCTTTTATTTGGTGATCCTGGTTCTTTTCGGCCGTTACGTCATTGTGTGCACCTCAG
CTGAAAGTTTCGTGCTACTTCTGTGGCCTCTCGTGGCTGGCGGCAGGCTGCCTTTTCGTTTACCA
GACTCTGGTTGAACACCTGGTGTGTGCCAAGTGCTGGCAGTGCCCTGGACAGGGGGCCTCAGG
GAAGGACGTGGAGCAGCCTTATCCCAGGCCTCTGGGTGTCCCGACACAGGTGTTACATCTGT
GCTGTCAGGTCAGATGCCTCAGTTCTTGGAAGCTAGGTTCTGCGACTGTTACCAAGGTGAT
TGTAAGAGCTGGCGGTACAGAGGAACAAGCCCCCAGCTGAGGGGGTGTGTGAATCGGACA
GCCTCCCAGCAGAGGTGTGGGAGCTGCAGCTGAGGGAAGAAGAGACAATCGGCCTGGACACTC
AGGAGGGTCAAAGGAGACTTGGTCGCACCACTCATCCTGCCACCCCCAGAATGCATCCTGCC
TCATCAGGTCCAGATTTCTTTCCAAGGCGGACGTTTTCTGTTGGAATTCTTAGTCCTTGGCCT
CGGACACCTTCATTTCGTTAGCTGGGGAGTGGTGGTGAGGCAGTGAAGAAGAGGCGGATGGTCA
CACTCAGATCCACAGAGCCCAGGATCAAGGGACCCACTGCAGTGGCAGCAGGACTGTTGGGCC
CCCACCCCAACCCTGCACAGCCCTCATCCCCCTCTTGGCTTGAGCCGTCAGAGGCCCTGTGCTG
AGTGTCTGACCGAGACACTCACAGCTTTGTTCATCAGGGCACAGGCTTCCTCGGAGCCAGGATG
ATCTGTGCCACGCTTGACCTCGGGCCCATCTGGGCTCATGCTCTCTCTCCTGCTATTGAATT
AGTACCTAGCTGCACACAGTATGTAGTTACCAAAGAATAAACGGCAATAATTGAGAAAAAAA

140/168

FIGURE 140

MRLGSGTFATCCVAIEVLGIAVFLRGFFPAPVRSSARAEGAEPPAPEPSAGASSNWTTLPPP
LFSKVVIVLIDALRDDFVFGSKGVKEMPYTTYLVEKGASHSFVAEAKPPTVTMPRIKALMTGS
LPGFVDVIRNLNSPALLEDVIRQAKAAGKRIVFYGDETWVKLFPKHFVEYDGTTSFFVSDYT
EVDNNVTRHLDKVLKRGDWDILILHYLGLDGHIGHISGPNSPLIGQKLSEMDSVLMKIHTSLQS
KERETPLPNLLVLCGDHGMSETGSHGASSTEEVNTPLILISSAFERKPGDIRHPKHVQ

Important features of the protein:**Signal peptide:**

amino acids 1-34

Transmembrane domain:

amino acids 58-76

N-glycosylation sites.

amino acids 56-60, 194-198

N-myristoylation sites.amino acids 6-12, 52-58, 100-106, 125-131, 233-239, 270-276,
275-281, 278-284**Amidation site.**

amino acids 154-158

Cell attachment sequence.

amino acids 205-208

141/168

FIGURE 141

GGCACGAGGCAAGCCTTCCAGGTTATCGTGACGCACCTTGAAAGTCTGAGAGCTACTGCCCTA
CAGAAAGTTACTAGTGCCCTAAAGCTGGCGCTGGCACTGATGTTACTGCTGCTGTTGGAGTAC
AACTTCCCTATAGAAAACAACCTGCCAGCACCTTAAGACCACTCACACCTTCAGAGTGAAGAAC
TTAAACCCGAAGAAATTCAGCATTTCATGACCAGGATCACAAAGTACTGGTCCTGGACTCTGGG
AATCTCATAGCAGTTCCAGATAAAAACTACATACGCCCAGAGATCTTCTTTGCATTAGCCTCA
TCCTTGAGCTCAGCCTCTGCGGAGAAAGGAAGTCCGATTCTCCTGGGGGTCTCTAAAGGGGAG
TTTTGTCTCTACTGTGACAAGGATAAAGGACAAAGTCATCCATCCCTTCAGCTGAAGAAGGAG
AAACTGATGAAGCTGGCTGCCCAAAGGAATCAGCACGCCGGCCCTTCATCTTTTATAGGGCT
CAGGTGGGCTCCTGGAACATGCTGGAGTCGGCGGCTCACCCCGGATGGTTCATCTGCACCTCC
TGCAATTGTAATGAGCCTGTTGGGGTGACAGATAAATTTGAGAACAGGAAACACATTGAATTT
TCATTTCAACCAGTTTGCAAAGCTGAAATGAGCCCCAGTGAGGTCAGCGATTAGGAAACTGCC
CCATTGAACGCCTTCCTCGCTAATTTGAACTAATTGTATAAAAACACCAAACCTGCTCACT

WO 01/16318

PCT/US00/23328

142/168

FIGURE 142

MLLLLEYNFPIENNCQHLKTTHTFRVKNLNPKKFSIHDQDHKVLVLD SGNLI AVPDKNYIRP
EIFFALASSLSSASAEKGSPILLGVSKGEFCLYCDKDKGQSHPSLQLKKEKLMKLAAQKESAR
RPFIFYRAQVGSWNMLESAAHPGWFICTSCNCNEPVGVTDKFENRKHIEFSFQPVCKAEMSPS
EVSD

cAMP- and cGMP-dependent protein kinase phosphorylation site.
amino acids 33-36

N-myristoylation site.
amino acids 50-55, 87-92

Interleukin-1
amino acids 37-182

[illegible]

WO 01/16318

PCT/US00/23328

144/168

FIGURE 144

MLGLPWKGGLSWALLLLLLLGSQILLIYAWHFHEQRDCDEHNVMARYLPATVEFAVHTFNQQSK
DYYAYRLGHILNSWKEQVESKTVFSMELLGRTRCGKFEDDIDNCHFQESTELNNTFTCFETI
STRPWMTQFSLLNKTCLEGFH

Important features of the protein:

Signal peptide:

amino acids 1-25

N-glycosylation sites.

amino acids 117-121, 139-143

N-myristoylation site.

amino acids 9-15

145/168

FIGURE 145

CTGTGCAGCTCGAGGCTCCAGAGGCACACTCCAGAGAGAGCCAAGGTTCTGACGCGATGAGGA
AGCACCTGAGCTGGTGGTGGCTGGCCACTGTCTGCATGCTGCTCTTCAGCCACCTCTCTGCGG
TCCAGACGAGGGGCATCAAGCACAGAATCAAGTGGAACCGGAAGGCCCTGCCCAGCACTGCCC
AGATCACTGAGGCCCAGGTGGCTGAGAACCGCCCCGGGAGCCTTCATCAAGCAAGGCCGCAAGC
TCGACATTGACTTCGGAGCCGAGGGCAACAGGTACTACGAGGCCAACTACTGGCAGTTCCCCG
ATGGCATCCACTACAACGGCTGCTCTGAGGCTAATGTGACCAAGGAGGCATTTGTCACCGGCT
GCATCAATGCCACCCAGGCGGCGAACCAGGGGGAGTTCCAGAAGCCAGACAACAAGCTCCACC
AGCAGGTGCTCTGGCGGCTGGTCCAGGAGCTCTGCTCCCTCAAGCATTGCGAGTTTTGGTTGG
AGAGGGGCGCAGGACTTCGGGTCACCATGCACCAGCCAGTGCTCCTCTGCCTTCTGGCTTTGA
TCTGGCTCATGGTGAAATAAGCTTGCCAGGAGGCTGGCAGTACAGAGCGCAGCAGCGAGCAAA
TCCTGGCAAGTGACCCAGCTCTTCTCCCCCAAACCCACGCGTGTTCTGAAGGTGCCCAGGAGC
GGCGATGCACTCGCACTGCAAATGCCGCTCCCACGTATGCGCCCTGGTATGTGCCTGCGTTCT
GATAGATGGGGGACTGTGGCTTCTCCGTCACTCCATTCTCAGCCCCCTAGCAGAGCGTCTGGCA
CACTAGATTAGTAGTAAATGCTTGATGAGAAGAACACATCAGGCACTGCGCCACCTGCTTCAC
AGTACTTCCCAACAACCTCTTAGAGGTAGGTGTATTCCCGTTTTACAGATAAGGAAACTGAGGC
CCAGAGAGCTGAAGTACTGCACCCAGCATCACCAGCTAGAAAGTGGCAGAGCCAGGATTCAAC
CCTGGCTTGTCTAACCCAGGTTTTCTGCTCTGTCCAATTCCAGAGCTGTCTGGTGATCACTT
TATGTCTCACAGGGACCCACATCCAAACATGTATCTCTAATGAAATTGTGAAAGCTCCATGTT
TAGAAATAAATGAAAACACCTGA

146/168

FIGURE 146

MRKHLSSWWLATVCMLLEFSHLSAVQTRGIKHRIKWNRKALPSTAQITEAQVAENRPGAFIKQG
RKLDIDFGAEGNRYYEANYWQFPDGIHYNGCSEANVTKEAFVTGCINATQAANQGEFQKPDNK
LHQQVLWRLVQELCSLKHCEFWLERGAGLRVTMHQPVLLCLLALIWLMLVK

Important features of the protein:

Signal peptide:

amino acids 1-26

Transmembrane domain:

amino acids 157-171

N-glycosylation sites.

amino acids 98-102, 110-114

Tyrosine kinase phosphorylation site.

amino acids 76-83

N-myristoylation sites.

amino acids 71-77, 88-94, 93-99, 107-113, 154-160

Amidation site.

amino acids 62-66

147/168

FIGURE 147

GCCTTGGCCTCCCAAAGGGCTGGGATTATAGGCGTGACCACCATGTCTGGTCCAGAGTCTCAT
TTCCTGATGATTTATAGACTCAAAGAAAACTATGTTTCAGAAGCTCTCTTCTCTTCTGGCCTC
CTCTCTGTCTTCTTTCCCTCTTTCTTCTTATTTTAATTAGTAGCATCTACTCAGAGTCATGCA
AGCTGGAAATCTTTCATTTTGCTTGTCAGTGGGGTAGGTCAGTCTTAGTTTTTATTTTT
TGAAATTTCAACTTTCAGATTCAGGGGGTACATGTGAAGGTTTGTTTTATGAGTATATTGCAT
GATGCTGAGGTTTGGGGT

WO 01/16318

PCT/US00/23328

148/168

FIGURE 148

MFRSSLLFWPPLCLLSLFLILISSIYSESKLEIFHFACQWGRSLSLSFYFLKFQLSDSGGT
CEGLFYEYIA

Important features of the protein:

Signal peptide:

amino acids 1-25

N-myristoylation site.

amino acids 62-68

149/168

FIGURE 149

GTCTCCGCGTCACAGGAAC TTCAGCACCCACAGGGCGGACAGCGCTCCCCTCTACCTGGAGAC
TTGACTCCCGCGCGCCCCAACCCTGCTTATCCCTTGACCGTCGAGTGTGAGAGATCCTGCAGC
CGCCCAGTCCCGGCCCCCTCTCCCGCCCCACACCCACCCTCCTGGCTCTTCCTGTTTTTACTCC
TCCTTTTTCATTCATAACAAAAGCTACAGCTCCAGGAGCCCAGCGCCGGGCTGTGACCCAAGCC
GAGCGTGGAAGAATGGGGTTCCTCGGGACCGGCACCTTGGATTCTGGTGTTAGTGCTCCCGATT
CAAGCTTTCCCCAAACCTGGAGGAAGCCAAGACAAATCTCTACATAATAGAGAATTAAGTGCA
GAAAGACCTTTGAATGAACAGATTGCTGAAGCAGAAGAAGACAAGATTAAAAAACATATCCT
CCAGAAAACAAGCCAGGTCAGAGCAACTATTCTTTTGTTGATAACTTGAACCTGCTAAAGGCA
ATAACAGAAAAGGAAAAAATTGAGAAAGAAAGACAATCTATAAGAAGCTCCCCACTTGATAAT
AAGTTGAATGTGGAAGATGTTGATTCAACCAAGAATCGAAAACCTGATCGATGATTATGACTCT
ACTAAGAGTGGATTGGATCATAAATTTCAAGATGATCCAGATGGTCTTCATCAACTAGACGGG
ACTCCTTTAACCGCTGAAGACATTGTCCATAAAATCGCTGCCAGGATTTATGAAGAAAATGAC
AGAGCCGTGTTTGACAAGATTGTTTCTAACTACTTAATCTCGGCCTTATCACAGAAAGCCAA
GCACATACACTGGAAGATGAAGTAGCAGAGGTTTTACAAAAATTAATCTCAAAGGAAGCCAAC
AATTATGAGGAGGATCCCAATAAGCCCACAAGCTGGACTGAGAATCAGGCTGGAAAAATACCA
GAGAAAGTGACTCCAATGGCAGCAATTCAAGATGGTCTTGCTAAGGGAGAAAACGATGAAACA
GTATCTAACACATTAACTTGACAAATGGCTTGGAAGGAGAACTAAAACCTACAGTGAAGAC
AACTTTGAGGAACTCCAATATTTCCCAAATTTCTATGCGCTACTGAAAAGTATTGATTCAGAA
AAAGAAGCAAAAGAGAAAGAAACACTGATTACTATCATGAAAACACTGATTGACTTTGTGAAG
ATGATGGTGAAATATGGAACAATATCTCCAGAAGAAGGTGTTTCCTACCTTGAAAACCTTGGAT
GAAATGATTGCTCTTCAGACCAAAAACAAGCTAGAAAAAAATGCTACTGACAATATAAGCAAG
CTTTTCCCAGCACCATCAGAGAAGAGTCATGAAGAAACAGACAGTACCAAGGAAGAAGCAGCT
AAGATGGAAAAGGAATATGGAAGCTTGAAGGATTCCACAAAAGATGATAACTCCAACCCAGGA
GGAAAGACAGATGAACCCAAAGGAAAAACAGAAGCCTATTTGGAAGCCATCAGAAAAAATATT
GAATGGTTGAAGAAACATGACAAAAGGGAAATAAAGAAGATTATGACCTTTCAAAGATGAGA
GACTTCATCAATAAACAAGCTGATGCTTATGTGGAGAAAGGCATCCTTGACAAGGAAGAAGCC
GAGGCCATCAAGCGCATTATATAGCAGCCTGTAAAAAATGGCAAAAGATCCAGGAGTCTTTCAAC
TGTTTCAGAAAACATAATATAGCTTAAACACTTCTAATTCTGTGATTAAATTTTTTGACCC
AAGGGTTATTAGAAAGTGCTGAATTTACAGTAGTTAACCTTTTACAAGTGGTTAAACATAGC
TTTCTTCCCGTAAAAACTATCTGAAAGTAAAGTTGTATGTAAGCTGAAAAAAAAAAAAAAAAAA
AAA

150/168

FIGURE 150

MGFLGTGTWILVLVLP IQAFPKPGGSQDKSLHNRELSAERPLNEQIAEAEEDKIKKTYPPENK
PGQSNYSFVDNLNLLKAITEKEKIEKERQSIRSSPLDNKLNVEDVDSTKNRKLIDDYDSTKSG
LDHKFQDDPDGLHQLDGTPLTAEDIVHKIAARIYEENDRAVEDKIVSKLLNLGLITESQAHTL
EDEVAEVLQKLISKEANNYEEDPNKPTSWTENQAGKIPEKVTPMAAIQDGLAKGENDETVSNT
LTLTNGLERRTKTYSEDNFEELQYFPNFYALLKSIDSEKEAKEKETLITIMKTLIDFVKMMVK
YGTISPEEGVSYLENLDEMIALQTKNKLEKNATDNISKLFPAPEKSHEETDSTKEEAAKMEK
EYGLKDKSTKDDNSNPGGKTDEPKGKTEAYLEAIRKNIEWLKKHDKKGNKEDYDLSKMRDFIN
KQADAYVEKGILDKEEAIEAIKRIYSSL

N-glycosylation sites:

amino acids 68-71, 346-349, 350-353

Casein kinase II phosphorylation site:

amino acids 70-73, 82-85, 97-100, 125-128, 147-150, 188-191, 217-
220, 265-268, 289-292, 305-308, 320-323, 326-329, 362-365, 368-
341, 369-372, 382-385, 386-389, 387-390

N-myristoylation sites:

amino acids 143-148, 239-244

151/168

FIGURE 151

CGGCTCGAGGCTCCCGCCAGGAGAAAGGAACATTCTGAGGGGAGTCTACACCCTGTGGAGCTC
AAGATGGTCCTGAGTGGGGCGCTGTGCTTCCGAATGAAGGACTCGGCATTGAAGGTGCTTTAT
CTGCATAATAACCAGCTTCTAGCTGGAGGGCTGCATGCAGGGAAGGTCATTAAAGGTGAAGAG
ATCAGCGTGGTCCCCAATCGGTGGCTGGATGCCAGCCTGTCCCCCGTCATCCTGGGTGTCCAG
GGTGGGAAGCCAGTGCCTGTCATGTGGGGTGGGGCAGGAGCCGACTCTAACACTAGAGCCAGTG
AACATCATGGAGCTCTATCTTGGTGCCAAGGAATCCAAGAGCTTCACCTTCTACCGGCGGGAC
ATGGGGCTCACCTCCAGCTTCGAGTCGGCTGCCTACCCGGGCTGGTTCCTGTGCACGGTGCCT
GAAGCCGATCAGCCTGTCAGACTCACCCAGCTTCCCGAGAATGGTGGCTGGAATGCCCCCATC
ACAGACTTCTACTTCCAGCAGTGTGACTAGGGCAACGTGCCCCCAGAACTCCCTGGGCAGAG
CCAGCTCGGGTGAGGGGTGAGTGGAGGAGACCCATGGCGGACAATCACTCTCTCTGCTCTCAG
GACCCCCACGTCTGACTTAGTGGGCACCTGACCACTTTGTCTTCTGGTTCCAGTTTGGATAA
ATTCTGAGATTTGGAGCTCAGTCCACGGTCCTCCCCCACTGGATGGTGCTACTGCTGTGGAAC
CTTGTA AAAACCATGTGGGGTAAACTGGGAATAACATGAAAAGATTTCTGTGGGGGTGGGGTG
GGGAGTGGTGGGAATCATTCTGCTTAATGGTAACTGACAAGTGTTACCCTGAGCCCCGCAG
GCCAACCCTATCCCCAGTTGAGCCTTATAGGGTCAGTAGCTCTCCACATGAAGTCCTGTCACTC
ACCACTGTGCAGGAGAGGGAGGTGGTCATAGAGTCAGGGATCTATGGCCCTTGGCCCAGCCCC
ACCCCTTCCCTTTAATCCTGCCACTGTCATATGCTACCTTTCCCTATCTCTTCCCTCATCATC
TTGTTGTGGGCATGAGGAGGTGGTGATGTCAGAAGAAATGGCTCGAGCTCAGAAGATAAAAGA
TAAGTAGGGTATGCTGATCCTCTTTTAAAAACCCAAGATACAATCAAAATCCCAGATGCTGGT
CTCTATTCCCATGAAAAAGTGCTCATGACATATTGAGAAGACCTACTTACAAAGTGGCATATA
TTGCAATTTATTTTAATTAAAAGATACCTATTTATATATTTCTTTATAGAAAAAGTCTGGAA
GAGTTTACTTCAATTGTAGCAATGTCAGGGTGGTGGCAGTATAGGTGATTTTTCTTTTAATTC
TGTTAATTTATCTGTATTTCTTAATTTTTCTACAATGAAGATGAATTCCTTGTATAAAAATAA
GAAAAGAAATTAATCTTGAGGTAAAGCAGAGCAGACATCATCTCTGATTGTCCTCAGCCTCCAC
TTCCCCAGAGTAAATTCAAATTGAATCGAGCTCTGCTGCTCTGGTTGGTTGTAGTAGTGATCA
GGAAACAGATCTCAGCAAAGCCACTGAGGAGGAGGCTGTGCTGAGTTTGTGTGGCTGGAATCT
CTGGGTAAAGGAACCTTAAAGAACA AAAATCATCTGGTAATTCTTTCTAGAAAGGATCACAGCCC
CTGGGATTCCAAGGCATTGGATCCAGTCTCTAAGAAGGCTGCTGTACTGGTTGAATTGTGTCC
CCCTCAAATTCACATCCTTCTTGGAATCTCAGTCTGTGAGTTTATTTGGAGATAAGGTCTCTG
CAGATGTAGTTAGTTAAGACAAGGTCATGCTGGATGAAGGTAGACCTAAATTCAATATGACTG
GTTTCCTTGTATGAAAAGGAGAGGACACAGAGACAGAGGAGACGCGGGGAAGACTATGTAAAG
ATGAAGGCAGAGATCGGAGTTTTGCAGCCACAAGCTAAGAAACACCAAGGATTGTGGCAACCA
TCAGAAGCTTGGAAGAGGCCAAAGAAGAATTCTTCCCTAGAGGCTTTAGAGGGATAACGGCTCT
GCTGAAACCTTAATCTCAGACTTCCAGCCTCCTGAACGAAGAAAGAATAAATTTCCGGCTGTTT
TAAGCCACCAAGGATAATTGGTTACAGCAGCTCTAGGAACTAATACAGCTGCTAAAATGATC
CCTGTCTCCTCGTGTTTACATTCTGTGTGTGTCCCCCTCCCACAATGTACCAAAGTTGTCTTTG
TGACCAATAGAATATGGCAGAAGTGATGGCATGCCACTTCCAAGATTAGGTTATAAAAGACAC
TGCAGCTTCTACTTGAGCCCTCTCTCTCTGCCACCCACCGCCCCCAATCTATCTTGGCTCACT
CGCTCTGGGGGAAGCTAGCTGCCATGCTATGAGCAGGCCCTATAAAGAGACTTACGTGGTAAAA
AATGAAGTCTCCTGCCACAGCCACATTAGTGAACCTAGAAGCAGAGACTCTGTGAGATAATC
GATGTTTGTGTTTAAAGTTGCTCAGTTTTGGTCTAACTTGTTATGCAGCAATAGATAAATAA
TATGCAGAGAAAGAG

152/168

FIGURE 152

MVLSGALCFRMKDSALKVLYLHNNQLLAGGLHAGKVIKGEEISVVPNRWLDASLSPVILGVQG
GSQCLSCGVGQEPTLTLEPVNIMELYLGAKESKSFTFYRRDMGLTSSFESAAYPGWFLCTVPE
ADQPVRLTQLPENGGWNAPITDFYFQQCD

N-myristoylation sites.

amino acids 29-34, 30-35, 60-65, 63-68, 73-78, 91-96, 106-111

Interleukin-1 signature.

amino acids 111-131

Interleukin-1 proteins.

amino acids 8-29, 83-120, 95-134, 64-103

153/168

FIGURE 153

CTTCAGAACAGGTTCTCCTTCCCCAGTCACCAGTTGCTCGAGTTAGAATTGTCTGCAATGGCC
GCCCTGCAGAAATCTGTGAGCTCTTTCCTTATGGGGACCCTGGCCACCAGCTGCCTCCTTCTC
TTGGCCCTCTTGGTACAGGGAGGAGCAGCTGCGCCCATCAGCTCCCAGTGCAGGCTTGACAAG
TCCAACCTCCAGCAGCCCTATATCACCAACCGCACCTTCATGCTGGCTAAGGAGGCTAGCTTG
GCTGATAACAACACAGACGTTCTGTCTCATTGGGGAGAACTGTTCCACGGAGTCAGTATGAGT
GAGCGCTGCTATCTGATGAAGCAGGTGCTGAACTTCACCCTTGAAGAAGTGCTGTTCCCTCAA
TCTGATAGGTTCCAGCCTTATATGCAGGAGGTGGTGCCCTTCCTGGCCAGGCTCAGCAACAGG
CTAAGCACATGTCATATTGAAGGTGATGACCTGCATATCCAGAGGAATGTGCAAAAGCTGAAG
GACACAGTGAAAAAGCTTGGAGAGAGTGGAGAGATCAAAGCAATTGGAGAACTGGATTTGCTG
TTTATGTCTCTGAGAAATGCCTGCATTTGACCAGAGCAAAGCTGAAAAATGAATAACTAACCC
CCTTTCCTGCTAGAAATAACAATTAGATGCCCCAAAGCGATTTTTTTTAAACCAAAGGAAGA
TGGGAAGCCAACTCCATCATGATGGGTGGATTCCAAATGAACCCCTGCGTTAGTTACAAAGG
AAACCAATGCCACTTTTGTTTATAAGACCAGAAGGTAGACTTTCTAAGCATAGATATTTATTG
ATAACATTTTCATTGTAAGTGGTGTCTATACACAGAAAACAATTTATTTTTTAAATAATTGTC
TTTTTCCATAAAAAAGATTACTTTCCATTCCTTTAGGGGAAAAAACCCCTAAATAGCTTCATG
TTTCCATAATCAGTACTTTATATTTATAAATGTATTTATTATTATTATAAGACTGCATTTTAT
TTATATCATTTTATTAATATGGATTTATTTATAGAAACATCATTCGATATTGCTACTTGAGTG
TAAGGCTAATATTGATATTTATGACAATAATTATAGAGCTATAACATGTTTATTTGACCTCAA
TAAACACTTGGATATCCC

154/168

FIGURE 154

MAALQKSVSSFLMGTLATSCLLLLALLVQGGAAAPISSHCRLDKSNFQQPYITNRTFMLAKEA
SLADNNTDVRLIGEKLFGVSMSERCYLMKQVLNFTLEEVLFPPQSDRFQPYMQEVPFLARLS
NRLSTCHIEGDDLHIQRNVQKLKDTVKKLGESGEIKAIGELDLLFMSLRNACI

Important features of the protein:

Signal peptide:

amino acids 1-33.

N-glycosylation sites.

amino acids 54-58, 68-72, 97-101

N-myristoylation sites.

amino acids 14-20, 82-88

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 10-21

155/168

FIGURE 155

GGCTTGCTGAAAATAAAATCAGGACTCCTAACCTGCTCCAGTCAGCCTGCTTCCACGAGGCCT
GTCAGTCAGTGCCCGACTTGTGACTGAGTGTGCAGTGCCCGAGCATGTACCAGGTCAGTGCAGA
GGGCTGCCTGAGGGCTGTGCTGAGAGGGAGAGGAGCAGAGATGCTGCTGAGGGTGGAGGGAGG
CCAAGCTGCCAGGTTTGGGGCTGGGGGCCAAGTGGAGTGAGAACTGGGATCCCAGGGGGAGG
GTGCAGATGAGGGAGCGACCCAGATTAGGTGAGGACAGTTCTCTCATTAGCCTTTTCCTACAG
GTGGTTGCATTCTTGGCAATGGTCATGGGAACCCACACCTACAGCCACTGGCCCAGCTGCTGC
CCGAGCAAAGGGCAGGACACCTCTGAGGAGCTGCTGAGGTGGAGCACTGTGCCTGTGCCTCCC
CTAGAGCCTGCTAGGCCCAACCGCCACCCAGAGTCCTGTAGGGCCAGTGAAGATGGACCCCTC
AACAGCAGGGCCATCTCCCCCTGGAGATATGAGTTGGACAGAGACTTGAACCGGCTCCCCCAG
GACCTGTACCACGCCCCGTTGCCTGTGCCCGCACTGCGTCAGCCTACAGACAGGCTCCCACATG
GACCCCCGGGGCAACTCGGAGCTGCTCTACCACAACCAGACTGTCTTCTACAGGCGGCCATGC
CATGGCGAGAAGGGCACCCACAAGGGCTACTGCCTGGAGCGCAGGCTGTACCGTGTTTCCTTA
GCTTGTGTGTGTGTGCGGCCCCGTGTGATGGGCTAGCCGGACCTGCTGGAGGCTGGTCCCTTT
TTGGGAAACCTGGAGCCAGGTGTACAACCACTTGCCATGAAGGGCCAGGATGCCCAGATGCTT
GGCCCCCTGTGAAGTGCTGTCTGGAGCAGCAGGATCCCGGGACAGGATGGGGGGCTTTGGGGAA
AACCTGCACTTCTGCACATTTTGAAAAGAGCAGCTGCTGCTTAGGGCCGCCGGAAGCTGGTGT
CCTGTCATTTTCTCTCAGGAAAGGTTTTCAAAGTTCTGCCCATTTCTGGAGGCCACCACTCCT
GTCTCTTCCTCTTTTCCCATCCCCTGCTACCCTGGCCCAGCACAGGCACTTTCTAGATATTTC
CCCCTTGCTGGAGAAGAAAGAGCCCCTGGTTTTATTTGTTTGTTTACTCATCACTCAGTGAGC
ATCTACTTTGGGTGCATTCTAGTGTAGTTACTAGTCTTTTGACATGGATGATTCTGAGGAGGA
AGCTGTTATTGAATGTATAGAGATTTATCCAAATAAATATCTTTATTTAAAAATGAAAAA

156/168

FIGURE 156

MRERPRLGEDSSLISLFLQVVAFLAMVMGTHYSHWPSCCPSKGQDTSEELLRWSTVPVPPE
PARPNRHPESCRASEDGPLNSRAISPWRYELDRDLNRLPQDLYHARCLCPHCVSLQTGSHMDP
RGNSELLYHNQTVFYRRPCHGEKGTKGYCLERRLYRVSLACVCVRPRVMG

Important features of the protein:

Signal peptide:

amino acids 1-32

N-glycosylation site.

amino acids 136-140

Tyrosine kinase phosphorylation site.

amino acids 127-135

N-myristoylation sites.

amino acids 44-50, 150-156

157/168

FIGURE 157

CCGGCGATGTCGCTCGTGCTGCTAAGCCTGGCCGCGCTGTGCAGGAGCGCCGTACCCCGAGAG
CCGACCGTTCAATGTGGCTCTGAAACTGGGCCATCTCCAGAGTGGATGCTACAACATGATCTA
ATCCCCGGAGACTTGAGGGACCTCCGAGTAGAACCTGTTACAAGTAGTGTTGCAACAGGGGAC
TATTCAATTTTGATGAATGTAAGCTGGGTACTCCGGGCAGATGCCAGCATCCGCTTGTTGAAG
GCCACCAAGATTTGTGTGACGGGCAAAAGCAACTTCCAGTCCTACAGCTGTGTGAGGTGCAAT
TACACAGAGGCCTTCCAGACTCAGACCAGACCCTCTGGTGGTAAATGGACATTTTCCTACATC
GGCTTCCCTGTAGAGCTGAACACAGTCTATTTTCATTGGGGCCCATAATATTCCTAATGCAAAT
ATGAATGAAGATGGCCCTTCCATGTCTGTGAATTTACCTCACCAGGCTGCCTAGACCACATA
ATGAAATATAAAAAAAGTGTGTCAAGGCCGGAAGCCTGTGGGATCCGAACATCACTGCTTGT
AAGAAGAATGAGGAGACAGTAGAAGTGAAGTTCACAACCACTCCCCTGGGAAACAGATACATG
GCTCTTATCCAACACAGCACTATCATCGGGTTTTCTCAGGTGTTTGAGCCACACCAGAAGAAA
CAAACGCGAGCTTCAGTGGTGATTCCAGTGAAGTGGGGATAGTGAAGGTGCTACGGTGCAGCTG
ACTCCATATTTTCCTACTTGTGGCAGCGACTGCATCCGACATAAAGGAACAGTTGTGCTCTGC
CCACAAACAGGCGTCCCTTTCCCTCTGGATAACAACAAAAGCAAGCCGGGAGGCTGGCTGCCT
CTCCTCCTGCTGTCTCTGCTGGTGGCCACATGGGTGCTGGTGGCAGGGATCTATCTAATGTGG
AGGCACGAAAGGATCAAGAAGACTTCCTTTTCTACCACCACACTACTGCCCCCATTAAGGTT
CTTGTGGTTTACCCATCTGAAATATGTTTCCATCACACAATTTGTTACTTCACTGAATTTCTT
CAAACCATTTGCAGAAGTGAGGTCATCCTTGAAAAGTGGCAGAAAAAGAAAATAGCAGAGATG
GGTCCAGTGCAGTGGCTTGCCACTCAAAGAAGGCAGCAGACAAAGTCGTCTTCCTTCTTTCC
AATGACGTCAACAGTGTGTGCGATGGTACCTGTGGCAAGAGCGAGGGCAGTCCCAGTGAGAAC
TCTCAAGACCTCTTCCCCCTTGCCTTTAACCTTTTCTGCAGTGATCTAAGAAGCCAGATTCAT
CTGCACAAATACGTGGTGGTCTACTTTAGAGAGATTGATACAAAAGACGATTACAATGCTCTC
AGTGTCTGCCCCAAGTACCACCTCATGAAGGATGCCACTGCTTTCTGTGCAGAACTTCTCCAT
GTCAAGCAGCAGGTGTCAGCAGGAAAAAGATCACAAGCCTGCCACGATGGCTGCTGCTCCTTG
TAG

158/168

FIGURE 158

MSLVLLSLAALCRSAVPREPTVQCGSETGPSPEWMLQHDLI PGDLRLDLRVEPVTTSVATGDYS
ILMNVS WVL RADASIRLLKATKICVTGKSNFQSYSCVRCNYTEAFQTQTRPSGGKWTFSYIGF
PVELNTVYFIGAHNIPNANMNEDGPSMSVNFTSPGCLDHIMKYKKKCVKAGSLWDPNITACKK
NEETVEVNFTTTPLGNRYMALIQHSTIIGFSQVFEPHQKKQTRASVVI PVTGDSEGATVQLTP
YFPTCGSDCIRHKGTVVLC PQTGVPFPLDNNKSKPGGWLPLLLLSLLVATWVLVAGIYLMWRH
ERIKKTSFSTTTLLPPIKVLVVYPSEICFHHTICYFTEFLQNHCRSEVILEKWQKKKIAEMGP
VQWLATQKKAADKVVFLLSNDVNSVCDGTCGKSEGSPSENSQDLFPLAFNLFCSDLRSQIHLH
KYVVVYFREIDTKDDYNALSVC PKYHLMKDATAFCAELLHV KQQVSAGKRSQACHDGCCSL

Important features of the protein:**Signal peptide:**

amino acids 1-14

Transmembrane domain:

amino acids 290-309

N-glycosylation sites.amino acids 67 - 71, 103 - 107, 156 - 160, 183 - 187, 197 - 201
and 283 - 287**cAMP- and cGMP-dependent protein kinase phosphorylation sites.**

amino acids 228 - 232 and 319 - 323

Casein kinase II phosphorylation sites.

amino acids 178 - 182, 402 - 406, 414 - 418 and 453 - 457

N-myristoylation site.

amino acids 116-122

Amidation site.

amino acids 488-452

159/168

FIGURE 159

AGCCACCAGCGCAACATGACAGTGAAGACCCTGCATGGCCCAGCCATGGTCAAGTACTTGCTG
CTGTCGATATTGGGGCTTGCCTTTCTGAGTGAGGCGGCAGCTCGGAAAATCCCCAAAGTAGGA
CATACTTTTTTCCAAAAGCCTGAGAGTTGCCCCGCCTGTGCCAGGAGGTAGTATGAAGCTTGAC
ATTGGCATCATCAATGAAAACCAGCGCGTTTCCATGTCACGTAACATCGAGAGCCGCTCCACC
TCCCCCTGGAATTACACTGTCACCTGGGACCCCAACCGGTACCCCTCGGAAGTTGTACAGGCC
CAGTGTAGGAACTTGGGCTGCATCAATGCTCAAGGAAAGGAAGACATCTCCATGAATTCCGTT
CCCATCCAGCAAGAGACCCTGGTTCGTCCGGAGGAAGCACCAAGGCTGCTCTGTTTCTTTCCAG
TTGGAGAAGGTGCTGGTGACTGTTGGCTGCACCTGCGTCACCCCTGTCATCCACCATGTGCAG
TAAGAGGTGCATATCCACTCAGCTGAAGAAG

160/168

FIGURE 160

MTVKTLHGPMVKYLLLSILGLAFLSEAAARKIPKVGHTFFQKPESCPPVPGGSMKLDIGIIN
ENQRVSMERNIESRSTSPWNYTVTWDPNRYPSSEVVQAQCRNLGCINAQGKEDISMNSVPIQQE
TLVVRKHKQGCSVSFQLEKVLVTVGCTCVTPVIHHVQ

Signal sequence:

amino acids 1-30

N-glycosylation site.

amino acids 83-87

N-myristoylation sites.

amino acids 106-111, 136-141

161/168

FIGURE 161

ACACTGGCCAAACAAAAACGAAAGCACTCCGTGCTGGAAGTAGGAGGAGAGTCAGGACTCCCA
GGACAGAGAGTGCACAAACTACCCAGCACAGCCCCCTCCGCCCCCTCTGGAGGCTGAAGAGGG
ATTCCAGCCCCCTGCCACCCACAGACACGGGCTGACTGGGGTGTCTGCCCCCCTTGGGGGGGGG
CAGCACAGGGCCTCAGGCCTGGGTGCCACCTGGCACCTAGAAGATGCTCTGTGCCCTGGTTCTT
GCTGTCCTTGGCACTGGGCCGAAGCCCAGTGGTCCTTTCTCTGGAGAGGCTTGTGGGGCCTCA
GGACGCTACCCACTGCTCTCCGGGCTCTCCTGCCGCCTCTGGGACAGTGACATACTCTGCCT
GCCTGGGGACATCGTGCCTGCTCCGGGGCCCCGTGCTGGCGCCTACGCACCTGCAGACAGAGCT
GGTGTGAGGTGCCAGAAGGAGACCGACTGTGACCTCTGTCTGCGTGTGGCTGTCCACTTGGC
CGTGCATGGGCACTGGGAAGAGCCTGAAGATGAGGAAAAGTTTGGAGGAGCAGCTGACTCAGG
GGTGGAGGAGCCTAGGAATGCCTCTCTCCAGGCCCAAGTCGTGCTCTCCTTCCAGGCCTACCC
TACTGCCCCGCTGCGTCCTGCTGGAGGTGCAAGTGCCTGCTGCCCTTGTGCAGTTTGGTCAGTC
TGTGGGCTCTGTGGTATATGACTGCTTCGAGGCTGCCCTAGGGAGTGAGGTACGAATCTGGTC
CTATACTCAGCCCAGGTACGAGAAGGAACTCAACCACACACAGCAGCTGCCTGCCCTGCCCTG
GCTCAACGTGTCAGCAGATGGTGACAACGTGCATCTGGTTCTGAATGTCTCTGAGGAGCAGCA
CTTCGGCCTCTCCCTGTACTGGAATCAGGTCCAGGGCCCCCCCCAAAACCCCGGTGGCACAAAAA
CCTGACTGGACCGCAGATCATTACCTTGAACCACACAGACCTGGTTCCCTGCCTCTGTATTCA
GGTGTGGCCTCTGGAACCTGACTCCGTTAGGACGAACATCTGCCCCCTTCAGGGAGGACCCCCG
CGCACACCAGAACCTCTGGCAAGCCGCCCCGACTGCGACTGCTGACCTTGCAGAGCTGGCTGCT
GGACGCACCGTGCTCGCTGCCCGCAGAAGCGGCCTGTGCTGGCGGGCTCCGGGTGGGGACCC
CTGCCAGCCACTGGTCCCACCGCTTTCCTGGGAGAACGTCACTGTGGACAAGGTTCTCGAGTT
CCCATTGCTGAAAGGCCACCCTAACCTCTGTGTTTCAGGTGAACAGCTCGGAGAAGCTGCAGCT
GCAGGAGTGCTTGTGGGCTGACTCCCTGGGGCCTCTCAAAGACGATGTGCTACTGTTGGAGAC
ACGAGGCCCCCAGGACAACAGATCCCTCTGTGCCTTGGAAACCAGTGGCTGTACTTCACTACC
CAGCAAAGCCTCCACGAGGGCAGCTCGCCTTGGAGAGTACTTACTACAAGACCTGCAGTCAGG
CCAGTGTCTGCAGCTATGGGACGATGACTTGGGAGCGCTATGGGCCTGCCCCATGGACAAATA
CATCCACAAGCGCTGGGCCCTCGTGTGGCTGGCCTGCCTACTCTTTGCCGCTGCGCTTTCCCT
CATCCTCCTTCTCAAAAAGGATCACGCGAAAGGGTGGCTGAGGCTCTTGAAACAGGACGTCCG
CTCGGGGGCGGGCCGCCAGGGGGCCGCGCGGCTCTGCTCCTCTACTCAGCCGATGACTCGGGTTT
CGAGCGCCTGGTGGGCGCCCTGGCGTCGGCCCTGTGCCAGCTGCCGCTGCGCGTGGCCGTAGA
CCTGTGGAGCCGTCGTGAACTGAGCGCGCAGGGGGCCCGTGGCTTGGTTTCACGCGCAGCGGCG
CCAGACCCTGCAGGAGGGCGGCGTGGTGGTCTTGCTCTTCTCTCCCGGTGCGGTGGCGCTGTG
CAGCGAGTGGCTACAGGATGGGGTGTCCGGGGCCCCGGGGCGCACGGCCCCGCACGACGCCTTCCG
CGCCTCGCTCAGCTGCGTGCTGCCCGACTTCTTGCAAGGGCCGGGGCGCCCCGGCAGCTACGTGGG
GGCCTGCTTCGACAGGCTGCTCCACCCGGACGCCGTACCCGCCCTTTTCCGCACCGTGCCCCGT
CTTCACACTGCCCTCCCAACTGCCAGACTTCTTGGGGGCCCTGCAGCAGCCTCGCGCCCCGCG
TTCCGGGGCGGCTCCAAGAGAGAGCGGAGCAAGTGTCCCGGGGCCCTTCAGCCAGCCCTGGATAG
CTACTTCCATCCCCCGGGGACTCCCGCGCCGGGACGCGGGGTGGGACCAGGGGGCGGGACCTGG
GGCGGGGGACGGGACTTAAATAAAGGCAGACGCTGTTTTTCTAAAAAA

162/168

FIGURE 162

MPVPWFLLSLALGRSPVVL\$LERLVGPQDATHCSPGLSCRLWDS\$DILCLPGDIVPAPGPVLAP
THLQTELVLR\$CQKETDCDLCLRVAVHLAVHGHWEPEDEEEKFGGAADSGVEEPRNASLQAQVV
LSFQAYPTARCVLLEVQVPAALVQFGQSVG\$SVVYDCFEAALGSEVRIWSYTQPRYEKELNHTQ
QLPALPWLNV\$ADGDNVHLVLNVSEEQHFGLSLYWNQVQ\$GPPKPRWHKNLTGPQIITLNHTDL
VPCLCIQVWPLEPDSVRTNICPFREDPRAHQNLWQAARLRLLTQ\$SWLLDAPCSLPAEAALCW
RAPGGDPCQPLVPPLSWENVTV\$DKVLEFPLLKGHPNLCVQVNSSEKLQLQECLWADSLGPLKD
DVL\$LLETRGPQDNRS\$LCAL\$PSGCTSLPSKASTRAARLGEYLLQDLQ\$GQCLQLWDDDLGALW
ACPM\$DKYIHKRWALVWLACLLFAAALS\$ILL\$LLKKDHAKGWLRL\$LLKQDVRSGAAARGRAALLY
SADDSGFERLVGALASALCQLPLRVAVDLWSRRELSAQGPV\$AWFHAQRRQTLQEGGVVLLFS
P\$AVALCSEWLQDGVSGPGAHGPHDAFRASLSCVLPDFLQGRAPGSYVGACFDRL\$LLHPDAVPA
LFRTVPVFTLPSQLPDFLGALQ\$QPRAPRSGRLQERAEQVSRALQPALDSYFHPPGTPAPGRGV
GPGAGPGAGDGT

Signal sequence:

amino acids 1-20

Transmembrane domain.

amino acids 453-475

N-glycosylation sites.amino acids 118-121, 186-189, 198-201, 211-214, 238-241, 248-251,
334-337, 357-360, 391-394**Glycosaminoglycan attachment site.**

amino acids 583-586

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 552-555

N-myristoylation sites.amino acids 107-112, 152-157, 319-324, 438-443, 516-521, 612-617,
692-697, 696-701, 700-705

163/168

FIGURE 163

GGGAGGGCTCTGTGCCAGCCCCGATGAGGACGCTGCTGACCATCTTGACTGTGGGATCCCTGG
CTGCTCACGCCCCTGAGGACCCCTCGGATCTGCTCCAGCACGTGAAATTCCAGTCCAGCAACT
TTGAAAACATCCTGACGTGGGACAGCGGGCCAGAGGGCACCCCAGACACGGTCTACAGCATCG
AGTATAAGACGTACGGAGAGAGGGACTGGGTGGCAAAGAAGGGCTGTCAGCGGATCACCCGGA
AGTCCTGCAACCTGACGGTGGAGACGGGCAACCTCACGGAGCTCTACTATGCEAGGGTCACCGCT
GTCAGTGCGGGAGGCCGGTCAGCCACCAAGATGACTGACAGGTTTCTCTCTGTCAGCACACT
ACCCTCAAGCCACCTGATGTGACCTGTATCTCCAAAGTGAGATCGATTGAGATGATTGTTTCAT
CCTACCCCCACGCCAATCCGTGCAGGCGATGGCCACCGGCTAACCCTGGAAGACATCTTCCAT
GACCTGTTCTACCACTTAGAGCTCCAGGTCAACCGCACCTACCAAATGCACCTTGGAGGGGAAG
CAGAGAGAATATGAGTTCTTCGGCCTGACCCCTGACACAGAGTTCCTTGGCACCATCATGATT
TGCGTTCCACCTGGGCCAAGGAGAGTGCCCCCTACATGTGCCGAGTGAAGACACTGCCAGAC
CGGACATGGACCTACTCCTTCTCCGGAGCCTTCCTGTTCTCCATGGGCTTCCTCGTCGCAGTA
CTCTGCTACCTGAGCTACAGATATGTCACCAAGCCGCCTGCACCTCCCAACTCCCTGAACGTC
CAGCGAGTCTGACTTTCCAGCCGCTGCGCTTCATCCAGGAGCACGTCTGATCCCTGTCTTT
GACCTCAGCGGCCCCAGCAGTCTGGCCCAGCCTGTCCAGTACTCCCAGATCAGGGTGTCTGGA
CCCAGGGAGCCCGCAGGAGCTCCACAGCGGCATAGCCTGTCCGAGATCACCTACTTAGGGCAG
CCAGACATCTCCATCCTCCAGCCCTCCAACGTGCCACCTCCCCAGATCCTCTCCCCACTGTCC
TATGCCCCAAACGCTGCCCCCTGAGGTGCGGCCCCCATCCTATGCACCTCAGGTGACCCCCGAA
GCTCAATTCCCATTTCTACGCCCCACAGGCCATCTCTAAGGTCCAGCCTTCCTCCTATGCCCCCT
CAAGCCACTCCGGACAGCTGGCCTCCCTCCTATGGGGTATGCATGGAAGGTTCTGGCAAAGAC
TCCCCCACTGGGACACTTTCTAGTCTTAAACACCTTAGGCCTAAAGGTCAGCTTCAGAAAGAG
CCACCAGCTGGAAGCTGCATGTTAGGTGGCCTTTCTCTGCAGGAGGTGACCTCCTTGGCTATG
GAGGAATCCCAAGAAGCAAAATCATTGCACCAGCCCCTGGGGATTGTCACAGACAGAACATCT
GACCCAAATGTGCTACACAGTGGGGAGGAAGGGACACCACAGTACCTAAAGGGGCCAGCTCCCC
CTCCTCTCCTCAGTCCAGATCGAGGGCCACCCCATGTCCCTCCCTTTGCAACCTCCTTCCGGT
CCATGTTCCCCCTCGGACCAAGGTCCAAGTCCCTGGGGCCTGCTGGAGTCCCTTGTGTGTCCC
AAGGATGAAGCCAAGAGCCCAGCCCCCTGAGACCTCAGACCTGGAGCAGCCCACAGAACTGGAT
TCTCTTTTCAGAGGCCTGGCCCTGACTGTGCAGTGGGAGTCCCTGAGGGGAATGGGAAAGGCTT
GGTGCTTCCTCCCTGTCCCTACCCAGTGTACATCCTTGGCTGTCAATCCCATGCCTGCCCAT
GCCACACACTCTGCGATCTGGCCTCAGACGGGTGCCCTTGAGAGAAGCAGAGGGGAGTGGCATG
CAGGGCCCCCTGCCATGGGTGCGCTCCTCACCGGAACAAAGCAGCATGATAAGGACTGCAGCGG
GGGAGCTCTGGGGAGCAGCTTGTGTAGACAAGCGCGTGCTCGCTGAGCCCTGCAAGGCAGAAA
TGACAGTGCAAGGAGGAAATGCAGGGGAACTCCCGAGGTCCAGAGCCCCACCTCCTAACACCA
TGGATTCAAAGTGCTCAGGGAATTTGCCTCTCCTTGCCCCATTCTTGCCAGTTTCACAATCT
AGCTCGACAGAGCATGAGGCCCTGCCTCTTCTGTCAATTGTTCAAAGGTGGGAAGAGAGCCTG
GAAAAGAACCAGGCCTGGAAAAGAACCAGAAGGAGGCTGGGCAGAACCAACAACCTGCACT
TCTGCCAAGGCCAGGGCCAGCAGGACGGCAGGACTCTAGGGAGGGGTGTGGCCTGCAGCTCAT
TCCCAGCCAGGGCAACTGCCTGACGTTGCACGATTTTCTAGCTTCATTCTCTGATAGAACAAG
CGAAATGCAGGTCCACCAGGGAGGGAGACACACAAGCCTTTTCTGCAGGCAGGAGTTTCAGAC
CCTATCCTGAGAATGGGGTTTGAAAGGAAGGTGAGGGCTGTGGCCCCCTGGACGGGTACAATAA
CACACTGTACTGATGTCACAACCTTTGCAAGCTCTGCCTTGGGTTCAGCCCATCTGGGCTCAA
TTCCAGCCTCACCCTCACAAGCTGTGTGACTTCAAACAATGAAATCAGTGCCCAGAACCTC
GGTTTCCTCATCTGTAATGTGGGGATCATAACACCTACCTCATGGAGTTGTGGTGAAGATGAA
ATGAAGTCATGTCTTTAAAGTGCTTAATAGTGCCTGGTACATGGGCAGTGCCCAATAAACGGT
AGCTATTTAAAAA

164/168

FIGURE 164

MRTLLTILTVGSLAAHAPEDPSDLLQHVKFQSSNFENILTWDSGPEGTPDTVYSIEYKTYGER
DWVAKKGCQRITRKSCNLTVETGNLTLEYARVTAVSAGGRSATKMTDRFSSLQHTTLKPPDV
TCISKVRSIQMIVHPTPTPIRAGDGHRLTLEDIFHDLFYHLELQVNRITYQMHLGGKQREYEFF
GLTPDTEFLGTIMICVPTWAKESAPYMCRVKTLPDRTWTYSFSGAFLFSMGFLVAVLCYLSYR
YVTKPPAPPNSLVQRVLTFQPLRFIQEHVLI PVFDLSGPSSLAQPVQYSQIRVSGPREPAGA
PQRHSLSEITYLGQPDISILQPSNVPPPQILSPLSYAPNAAPEVGPPSYAPQVTPEAQFPFYA
PQAISKVQPSSYAPQATPDSWPPSYGVCMEGSGKDSPTGTLSSPKHLRPKGQLQKEPPAGSCM
LGGLSLQEVTS LAMEESQEAKSLHQPLGICTDRTSDPNVLHSGEEGTPQYLLKGQLPLLSSVQI
EGHPMSLPLQPPSGPCSPSDQGPSPWGLLESLVCPKDEAKSPAPETSDLEQPTELDSLFRGLA
LTVQWES

Signal sequence.

amino acids 1-17

Transmembrane domain.

amino acids 233-250

N-glycosylation sites.

amino acids 80-83, 87-90, 172-175

N-myristoylation sites.

amino acids 11-16, 47-52, 102-107, 531-536, 565-570

165/168

FIGURE 165

TGGCCTACTGGAAAAAAAAAAAAAAAAAAAAAGTCACCCGGGCCCCGCGGTGGCCACAACAT
GGCTGCGGGCGCCGGGGCTGCTCTTCTGGCTGTTTCGTGCTGGGGGCGCTCTGGTGGGTCCCGGG
CCAGTCGGATCTCAGCCACGGACGGCGTTTCTCGGACCTCAAAGTGTGCGGGGACGAAGAGTG
CAGCATGTTAATGTACCGTGGGAAAGCTCTTGAAGACTTCACGGGCCCTGATTGTCGTTTTGT
GAATTTTAAAAAAGGTGACGATGTATATGTCTACTACAACTGGCAGGGGGATCCCTTGAAC
TTGGGCTGGAAGTGTTGAACACAGTTTTGGATATTTTCCAAAAGATTTGATCAAGGTACTTCA
TAAATACACGGAAGAAGAGCTACATATTCAGCAGATGAGACAGACTTTGTCTGCTTTGAAGG
AGGAAGAGATGATTTTAATAGTTATAATGTAGAAGAGCTTTTAGGATCTTTGGAAGTGGAGGA
CTCTGTACCTGAAGAGTCGAAGAAAGCTGAAGAAGTTTCTCAGCACAGAGAGAAATCTCCTGA
GGAGTCTCGGGGGCGTGAACTTGACCCTGTGCCTGAGCCCGAGGCATTCAGAGCTGATTCAGA
GGATGGAGAAGGTGCTTTCTCAGAGAGCACCGAGGGGGCTGCAGGGACAGCCCTCAGCTCAGGA
GAGCCACCCTCACACCAGCGGTCCTGCGGCTAACGCTCAGGGAGTGCAGTCTTCGTTGGACAC
TTTTGAAGAAATTCTGCACGATAAATTGAAAGTGCCGGGAAGCGAAAGCAGAACTGGCAATAG
TTCTCCTGCCTCGGTGGAGCGGGAGAAGACAGATGCTTACAAAGTCCTGAAAACAGAAATGAG
TCAGAGAGGAAGTGGACAGTGCGTTATTCATTACAGCAAAGGATTCGTTGGCATCAAATCT
AAGTTTGTTTTACAAAGATTGTTTTTTAGTACTAAGCTGCCTTGGCAGTTTGCATTTTTGAGCC
AAACAAAAATATATTATTTTCCCTTCTAAGTAAAAAAAAAAAAAAAAAAAAA

166/168

FIGURE 166

MAAAPGLLFWLFLVGLGALWWVPGQSDLSHGRRFSDLKVCGDEEC SMLMYRGKALEDFTGPDCRF
VNFKKGDDVYVYYKLAGGSLELWAGSVEHSFGYFPKDLIKVLH KYTEEELHIPADETDFVCFE
GGRDDFNSYNVEELLGSLELEDSVPEESKKAEEVSQHREKSPEESRGRELDPVPEPEAFRADS
EDGEGAFSESTEG LQGQPSAQESH PHTSGPAANAQGVQSSLDTFEEILHDKLKVPGSESRTGN
SSPASVEREKTDAYKVLKTEMSQRGSGQCVIHYSKGFRRWHQNL SLFYKDCF

Important features of the protein:**Signal peptide:**

amino acids 1-22

N-glycosylation site.

amino acids 294-298

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 30-34

Tyrosine kinase phosphorylation site.

amino acids 67-76

N-myristoylation sites.

amino acids 205-211, 225-231, 277-283

Amidation site.

amino acids 28-32

167/168

FIGURE 167

CCAGGACCAGGGCGCACCGGCTCAGCCTCTCACTTGTCAGAGGCCGGGGAAGAGAAGCAAAGC
GCAACGGTGTGGTCCAAGCCGGGGCTTCTGCTTCGCCTCTAGGACATACACGGGACCCCCTAA
CTTCAGTCCCCCAAACGCGCACCCCTCGAAGTCTTGAAGTCCAGCCCCGCACATCCACGCGCGG
CACAGGCGCGGCAGGCGGCAGGTCCCGGCCGAAGGCGATGCGCGCAGGGGGTTCGGGCAGCTGG
GCTCGGGCGGGCGGGAGTAGGGCCCCGGCAGGGAGGCAGGGAGGCTGCATATTCAGAGTCGCGGG
CTGCGCCCTGGGCAGAGGCCGCCCTCGCTCCACGCAACACCTGCTGCTGCCACCGCGCCGCGA
TGAGCCGCGTGGTCTCGCTGCTGCTGGGCGCCGCGCTGCTCTGCGGCCACGGAGCCTTCTGCC
GCCGCGTGGTCAGCGGCCAAAAGGTGTGTTTTGCTGACTTCAAGCATCCCTGCTACAAAATGG
CCTACTTCCATGAACTGTCCAGCCGAGTGAGCTTTCAGGAGGCACGCCTGGCTTGTGAGAGTG
AGGGAGGAGTCCTCCTCAGCCTTGAGAATGAAGCAGAACAGAAGTTAATAGAGAGCATGTTGC
AAAACCTGACAAAACCCGGGACAGGGATTTCTGATGGTGATTTCTGGATAGGGCTTTGGAGGA
ATGGAGATGGGCAAACATCTGGTGCCCTGCCCAGATCTCTACCAGTGGTCTGATGGAAGCAATT
CCCAGTACCGAACTGGTACACAGATGAACCTTCCTGCGGAAGTGAAAAGTGTGTTGTGATGT
ATCACCAACCAACTGCCAATCCTGGCCTTGGGGGTCCCTACCTTTACCAGTGGAATGATGACA
GGTGTAACATGAAGCACAATTATATTTGCAAGTATGAACCAGAGATTAATCCAACAGCCCCTG
TAGAAAAGCCTTATCTTACAAATCAACCAGGAGACACCCATCAGAATGTGGTTGTTACTGAAG
CAGGTATAATTCCCAATCTAATTTATGTTGTTATACCAACAATAACCCCTGCTCTTACTGATAC
TGGTTGCTTTTGGAACTGTGTTTCCAGATGCTGCATAAAAGTAAAGGAAGAACAAAACCTA
GTCCAAACCAGTCTACACTGTGGATTTCAAAGAGTACCAGAAAAGAAAGTGGCATGGAAGTAT
AATAACTCATTGACTTGGTTCCAGAATTTTGTAATTCTGGATCTGTATAAGGAATGGCATCAG
AACAAATAGCTTGGAAATGGCTTGAAATCACAAAGGATCTGCAAGATGAACTGTAAGCTCCCCCT
TGAGGCAAATATTAAAGTAATTTTATATGTCTATTATTTTCAATTTAAAGAATATGCTGTGCTA
ATAATGGAGTGAGACATGCTTATTTTGCTAAAGGATGCACCCAAACTTCAAACCTTCAAGCAAA
TGAAATGGACAATGCAGATAAAGTTGTTATCAACACGTCGGGAGTATGTGTGTTAGAAGCAAT
TCCTTTTATTTCTTTTACCTTTTATAAGTTGTTATCTAGTCAATGTAATGTATATTGTATTGA
AATTTACAGTGTGCAAAAGTATTTTACCTTTGCATAAGTGTTTGATAAAAATGAACTGTTCTA
ATATTTATTTTATGGCATCTCATTTTCAATACATGCTCTTTTGATTAAAGAACTTATTAC
TGTTGTCAACTGAATTCACACACACACAAATATAGTACCATAGAAAAAGTTTGTCTTCTCGAA
ATAATTCATCTTTCAGCTTCTCTGCTTTTGGTCAATGTCTAGGAAATCTCTTCAGAAATAAGA
AGCTATTTTCAATTAAGTGTGATATAAACCTCCTCAAACATTTTACTTAGAGGCAAGGATTGTCT
AATTTCAATTGTGCAAGACATGTGCCTTATAATTATTTTATAGCTTAAAATTAAACAGATTTTG
TAATAATGTAACCTTTGTTAATAGGTGCATAAACACTAATGCAGTCAATTTGAACAAAAGAAGT
GACATACACAATATAAATCATATGTCTTCACACGTTGCCTATATAATGAGAAGCAGCTCTCTG
AGGGTTCTGAAATCAATGTGGTCCCTCTCTTGCCCACTAAACAAAGATGGTTGTTTCGGGGTTT
GGGATTGACACTGGAGGCAGATAGTTGCAAAGTTAGTCTAAGGTTTCCCTAGCTGTATTTAGC
CTCTGACTATATTAGTATACAAAGAGGTCATGTGGTTGAGACCAGGTGAATAGTCACTATCAG
TGTGGAGACAAGCACAGCACACAGACATTTTAGGAAGGAAAGGAACACGAAATCGTGTGAAA
ATGGGTTGGAACCCATCAGTGATCGCATATTCATTGATGAGGGTTTGCTTGAGATAGAAAATG
GTGGCTCCTTTCTGTCTTATCTCCTAGTTTCTTCAATGCTTACGCCTTGTTCTTCTCAAGAGA
AAGTTGTAACCTCTGCTTTCATATGTCCCTGTGCTCCTTTTAACCAAATAAAGAGTTCTTG
TTTCTGGGGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

168/168

FIGURE 168

MSRVVSLLLGAALLCGHGAFRRVVSGQKVCFADFKHPCYKMAYFHELSSRVSFQEARLACES
EGGVLLSLENEAEQKLIESMLQNLT KP GTGISDGDFWIGLWRNGDGQTSGACPDLYQWSDGSN
SQYRNWYTDEPSCGSEKCVVMYHQPTANPGLGGPYLYQWNDDRCNMKHNYICKYEPEINPTAP
VEKPYLTNQP GDTHQNVVVTEAGIIPNLIYVVIPTIPLLLLILVAFGTCCFQMLHKSKGR TKT
SPNQSTLWISKSTRKESGMEV

Important features of the protein:**Signal peptide:**

amino acids 1-21

Transmembrane domain:

amino acids 214-235

N-glycosylation sites.

amino acids 86-89, 255-258

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 266-269

N-myristoylation sites.

amino acids 27-32, 66-71, 91-96, 93-98, 102-107, 109-114, 140-145, 212-217

GGGGCTTCGGCGCCAGCGGCCAGCGCTAGTCGGTCTGGTAAGGATTTACAAAAGGTGCAGGTA
TGAGCAGGTCTGAAGACTAACATTTTGTGAAGTTGTAAAACAGAAAACCTGTTAGAAATGTGG
TGGTTTCAGCAAGGCCTCAGTTTCCTTCCTTCAGCCCTTGTAATTTGGACATCTGCTGCTTTC
ATATTTTCATACATTACTGCAGTAACACTCCACCATATAGACCCGGCTTTACCTTATATCAGT
GACACTGGTACAGTAGCTCCAGAAAAATGCTTATTTGGGGCAATGCTAAATATTGCGGCAGTT
TTATGCATTGCTACCATTTATGTTTCGTTATAAGCAAGTTCATGCTCTGAGTCCTGAAGAGAAC
GTTATCATCAAATTAAACAAGGCTGGCCTTGTACTTGGAATACTGAGTTGTTTAGGACTTTCT
ATTGTGGCAAACCTTCCAGAAAAACAACCCTTTTTTGCTGCACATGTAAGTGGAGCTGTGCTTACC
TTTGGTATGGGCTCATTATATATGTTTGTTTCAGACCATCCTTTCCTACCAAATGCAGCCCAA
ATÇCATGGCAAACAAGTCTTCTGGATCAGACTGTTGTTGGTTATCTGGTGTGGAGTAAGTGCA
CTTAGCATGCTGACTTGCTCATCAGTTTTGCACAGTGGCAATTTTGGGACTGATTTAGAACAG
AAACTCCATTGGAACCCCGAGGACAAAGGTTATGTGCTTCACATGATCACTACTGCAGCAGAA
TGGTCTATGTCATTTTCCTTCTTTGGTTTTTTTCCTGACTTACATTCGTGATTTTCAGAAAATT
TCTTTACGGGTGGAAGCCAATTTACATGGATTAAACCCTCTATGACACTGCACCTTGCCCTATT
AACAAATGAACGAACACGGCTACTTTCCAGAGATATTTGATGAAAGGATAAAAATATTTCTGTAA
TGATTATGATTCTCAGGGATTGGGGAAAGGTTCACAGAAGTTGCTTATTCTTCTCTGAAATTT
TCAACCACTTAATCAAGGCTGACAGTAACACTGATGAATGCTGATAATCAGGAAACATGAAAG
AAGCCATTTGATAGATTATTCTAAAGGATATCATCAAGAAGACTATTAAAAACACCTATGCCT
ATACTTTTTTATCTCAGAAAAATAAAGTCAAAAGACTATG