



(86) Date de dépôt PCT/PCT Filing Date: 2004/03/26

(87) Date publication PCT/PCT Publication Date: 2005/01/06

(45) Date de délivrance/Issue Date: 2012/05/08

(85) Entrée phase nationale/National Entry: 2005/09/27

(86) N° demande PCT/PCT Application No.: US 2004/009296

(87) N° publication PCT/PCT Publication No.: 2005/001065

(30) Priorité/Priority: 2003/04/01 (US60/459,734)

(51) Cl.Int./Int.Cl. *C12N 15/56* (2006.01),
C11D 7/42 (2006.01), *C12N 15/55* (2006.01),
C12N 9/42 (2006.01), *C12P 19/00* (2006.01),
C12P 19/14 (2006.01)


(72) Inventeurs/Inventors:
LARENAS, EDMUND A., US;
GOEDEGEBUUR, FRITS, NL;
GUALFETTI, PETER, US;
MITCHINSON, COLIN, US

(73) Propriétaire/Owner:
GENENCOR INTERNATIONAL, INC., US

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : CBH1.1 D'HUMICOLA GRISEA VARIANT

(54) Title: VARIANT HUMICOLA GRISEA CBH1.1



pRAX1

(57) Abrégé/Abstract:

ABSTRACT OF THE DISCLOSURE Disclosed are variants of Humicola grisea Cel7A (CBH1.1), H. jecorina CBH1 variant or S. thermophilum CBH1, nucleic acids encoding the same and methods for producing the same. The variant cellulases have the amino acid sequence of a glycosyl hydrolase of family 7A wherein one or more amino acid residues are substituted.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
6 January 2005 (06.01.2005)

PCT

(10) International Publication Number
WO 2005/001065 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number:
PCT/US2004/009296
- (22) International Filing Date: 26 March 2004 (26.03.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/459,734 1 April 2003 (01.04.2003) US
- (71) Applicant (for all designated States except US): **GENENCOR INTERNATIONAL, INC.** [US/US]; 925 Page Mill Road, Palo Alto, CA 94304 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LARENAS, Edmund, A.** [US/US]; 301 Nevada, Moss Beach, CA 94038 (US). **GOEDEGEBUUR, Frits** [NL/NL]; Rozenlaan 128, NL-3135XV Vlaardingen (NL). **GUALFETTI, Peter** [US/US]; 129 Arkansas Street #1, San Francisco, CA 94107 (US). **MITCHINSON, Colin** [GB/US]; 381 Myrtle Street, Half Moon Bay, CA 94019 (US).
- (74) Agent: **BOYD?, Victoria, L.?**; GENENCOR INTERNATIONAL, INC., 925 Page Mill Road, Palo Alto, CA 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: VARIANT HUMICOLA GRISEA CBH1.1



(57) Abstract: ABSTRACT OF THE DISCLOSURE
Disclosed are variants of Humicola grisea Cel7A (CBH1.1), H. jecorina CBH1 variant or S. thermophilum CBH1, nucleic acids encoding the same and methods for producing the same. The variant cellulases have the amino acid sequence of a glycosyl hydrolase of family 7A wherein one or more amino acid residues are substituted.

WO 2005/001065 A2

VARIANT HUMICOLA GRISEA CBH1.1

CROSS-REFERENCE TO RELATED APPLICATIONS

[01] This application claims priority to U.S. Provisional Application No. 60/459,734 filed April 1, 2003 (Attorney Docket No. GC794P).

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[02] Portions of this work were funded by Subcontract No. ZCO-0-30017-01 with the National Renewable Energy Laboratory under Prime Contract No. DE-AC36-99GO10337 with the U.S. Department of Energy. Accordingly, the United States Government may have certain rights in this invention.

FIELD OF THE INVENTION

[03] The present invention relates to polypeptides having cellobiohydrolase I (also referred to as CBH I or CBH1) activity and polynucleotides having a nucleotide sequence which encodes for the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the nucleic acid constructs as well as methods for producing and using the polypeptides.

BACKGROUND OF THE INVENTION

[04] Cellulose is an important industrial raw material and a source of renewable energy. The physical structure and morphology of native cellulose are complex and the fine details of its structure have been difficult to determine experimentally. However, the chemical composition of cellulose is simple, consisting of D-glucose residues linked by beta-1 4-glycosidic bonds to form linear chains.

[05] In order to be efficient, the digestion of cellulose requires several types of enzymes acting cooperatively. At least three categories of enzymes are necessary to convert cellulose into glucose: endo (1,4)-beta-D-glucanases (EC 3 1.4) that cut the cellulose chains at random; cellobiohydrolases (EC 3 1.91) which cleave cellobiosyl units from the cellulose chain ends and beta-glucosidases (EC 3 1.21) that convert cellobiose and

soluble cellodextrins into glucose. Among these three categories of enzymes involved in the biodegradation of cellulose, cellobiohydrolases are the key enzymes for the degradation of native crystalline cellulose.

[06] Exo-cellobiohydrolases (Cellobiohydrolase 1, or CBH 1) refer to the cellobiohydrolases which degrade cellulose by hydrolyzing the cellobiose from the non-reducing end of the cellulose polymer chains.

[07] It is an object of the present invention to provide improved polypeptides having cellobiohydrolase I activity and polynucleotides encoding the polypeptides. The improved polypeptides may have improved specific activity and/or improved stability - in particular improved thermostability.

[08] Although cellulase compositions have been previously described, there remains a need for new and improved cellulase compositions for use in household detergents, stonewashing compositions or laundry detergents, etc. Cellulases that exhibit improved performance are of particular interest.

REFERENCES

- Altschul, S. F., *et al.*, J. Mol. Biol. 215:403-410, 1990.
- Altschul, S. F., *et al.*, Nucleic Acids Res. 25:3389-3402, 1997.
- Aro, N., *et al.*, J. Biol. Chem., 10.1074/ M003624200, April 13, 2001.
- Aubert, *et al.*, Ed., p11 et seq., Academic Press, 1988.
- Ausubel G. M., *et al.* CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y., 1993.
- Baker et al., Appl. Biochem. and Biotechnol. 45/46:245 - 256, 1994.
- Bhikhabhai, R. *et al.*, J. Appl. Biochem. 6:336, 1984.
- Boel et al. EMBO J 3:1581-1585 1984.
- Brumbauer, A. *et al.*, Bioseparation 7:287-295, 1999.
- Deutscher, M.P., Methods Enzymol. 182:779-80, 1990.
- Ellouz, S. *et al.*, J. Chromatography 396:307, 1987.
- Filho, *et al.* Can. J. Microbiol. 42:1-5, 1996.
- Fliess, A., *et al.*, Eur. J. Appl. Microbiol. Biotechnol. 17:314, 1983.
- Goedegebuur et al., Curr. Genet. 41 :89 – 98, 2002.
- Goyal, A. *et al.* Bioresource Technol. 36:37, 1991.
- Hazell, B. W. et al., Lett. Appl. Microbiol. 30:282-286, 2000.
- Herr *et al.*, Appl. Microbiol. Biotechnol. 5:29-36, 1978.
- Hu et al., Mol. Cell. Biol. 11:5792-9, 1991.

- Jeeves et al., *Biotechnol. Genet. Eng. Rev.* 9:327 -369, 1991.
- Kawaguchi, T *et al.*, *Gene* 173(2):287-8, 1996.
- Kelley et al. *EMBO J.* 4 :475-479, 1985.
- Knowles, J. *et al.*, *TIBTECH* 5, 255-261, 1987.
- Krishna, S. *et al.*, *Bioresource Tech.* 77:193-196, 2001.
- Kuhls K. *et al.*, *Proc. Natl. Acad. Sci. USA* 93(15): 7755 - 7760, 1996.
- Kumar, A., *et al.*, *Textile Chemist and Colorist* 29:37-42, 1997.
- Medve, J. *et al.*, *J. Chromatography A* 808:153, 1998.
- Mohagheghi, A. et al., *Int. J. Syst. Bacteriol.* 36:435 - 443, 1986.
- Nieves et al., *Appl. Biochem. and Biotechnol.* 51/52 211 - 223, 1995.
- Nunberg et al. *Mol. Cell Biol.* 4:2306-2315 1984.
- Ohmiya *et al.*, *Biotechnol. Gen. Engineer. Rev.* 14:365-414, 1997.
- Okada, M. et al., *Appl. Environ. Microbiol.*, 64:555-563, 1988.
- Ooi *et al.*, *Nucleic Acid Res.* 18:5884, 1990
- Penttila *et al.*, *Gene* 45:253-263, 1986.
- Penttila et al., *Gene* 61: 155 – 164, 1987.
- Penttila *et al.*, *Gene* 63: 103-112, 1988.
- Pere, J., *et al.*, In *Proc. Tappi Pulping Conf.*, Nashville, TN, 27-31, pp. 693-696, 1996.-
- Saarilahti *et al.*, *Gene* 90:9-14, 1990.
- Sakamoto *et al.*, *Curr. Genet.* 27:435-439, 1995.
- Saloheimo M, *et al.*, *Gene* 63:11-22, 1988.
- Saloheimo, A. *et al.*, *Molecular Microbiology*, 13:219-228, 1994.
- Saloheimo, M. et al., *Eur. J. Biochem.*, 249:584-591, 1997.
- Sambrook *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL* (Second Edition), Cold Spring Harbor Press, Plainview, N.Y., 1989.
- Schulein, *Methods Enzymol.*, 160, 25, pages 234 et seq, 1988.
- Scopes, *Methods Enzymol.* 90 Pt E:479-90, 1982.
- Shoemaker et al., *Biochem. Biophys. Acat.* 523 :133-146 1978.
- Shoemaker, S. et al., *Bio/Technology*, 1:691-696, 1983
- Srisodsuk, M. et al. *J. Biol. Chem.* 268(28): 20756 - 20761, 1993.
- Strathern *et al.*, eds. (1981) *The Molecular Biology of the Yeast Saccharomyces*.
- Suurnakki, A. *et al.*, *Cellulose* 7:189-209, 2000.
- Teeri, T. et al., *Gene*, 51:43-52, 1987
- Tilbeurgh, H. *et al.*, *FEBS Lett.* 16:215, 1984.
- Tomaz, C. and Queiroz, J., *J. Chromatography A* 865:123-128, 1999.
- Tomme, P. *et al.*, *Eur. J. Biochem.* 170:575-581, 1988.

Van Tilbeurgh, H. *et al.*, FEBS Lett. 204:223-227, 1986.

Ward, M. *et al.*, Appl. Microbiol. Biotechnol. 39:738 – 743, 1993.

Wood, Biochem. Soc. Trans., 13, pp. 407-410, 1985.

Wood *et al.*, METHODS IN ENZYMOLOGY, 160, 25, p. 87 et seq., Academic Press, New York, 1988.

BRIEF SUMMARY OF THE INVENTION

[09] In a first aspect, the present invention relates to a polypeptide having cellobiohydrolase I activity, selected from the group consisting of

- a) *H. grisea* CBH1.1 variant derived from CBS 225.63
- b) *H. grisea* CBH1.1 variant having the sequence given in Figure 3
- c) *H. grisea* CBH1.1 variant having the sequence given in Figure 4
- d) *Hypocrea jecorina* CBH1 variant as described herein; and
- e) *Scytalidium thermophilum* CBH1 derived from CBS 671.88.

[10] In second aspect the present invention relates to a polynucleotide encoding an *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1. In one embodiment, the polynucleotide encodes an *H. grisea* CBH1.1 variant is derived from CBS 225.63. In another embodiment, the polynucleotide encodes an *H. grisea* CBH1.1 variant shown in Figure 3. In another embodiment, the polynucleotide encodes an *H. grisea* CBH1.1 variant shown in Figure 4.

[11] In one general embodiment, polynucleotide encoding an *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 has at least 90%, preferably 95%, 98%, or more sequence identity to the *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 coding sequences presented herein using a sequence alignment program.

[12] In a third aspect the present invention relates to a nucleic acid construct comprising the nucleotide sequence, which encodes for the inventive CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1, operably linked to one or more control sequences that direct the production of the CBH1.1 variant, *H. jecorina* CBH1 variant or *S. thermophilum* CBH1 in a suitable host.

[13] In a fourth aspect the present invention relates to a recombinant expression vector comprising the nucleic acid construct of the invention.

[14] In a fifth aspect the present invention relates to a recombinant host cell comprising the nucleic acid construct of the invention.

[15] In a sixth aspect the present invention relates to a method for producing a CBH1, the method comprising

- a) transforming a host cell with a nucleic acid encoding a *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 described herein;
- b) culturing the host cell under conditions to produce the polypeptide; and
- c) recovering the polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

[16] Figure 1 is the genomic DNA sequence for *H. grisea* CBH1.1 (SEQ ID NO:1). The putative intron is in bold and underlined.

[17] Figure 2 is the cDNA sequence for *H. grisea* CBH1.1 (SEQ ID NO:2). The putative intron sequence (nucleotides 413-472 in Figure 1) has been deleted.

[18] Figure 3 is the signal sequence and mature amino acid sequence for *H. grisea* CBH1.1 (SEQ ID NO:3). The signal sequence is in bold and underlined.

[19] Figure 4 is the mature amino acid sequence for *H. grisea* CBH1.1 (SEQ ID NO:4).

[20] Figure 5 shows an alignment of two public and a variant *Humicola grisea* CBH1.1 mature sequences. The two public sequences are X17258 (SEQ ID NO:5) and D63515 (SEQ ID NO:6).

[21] Figure 6 is the pRAX1 plasmid. This vector is based on the plasmid pGAPT2 except a 5259bp HindIII fragment of *Aspergillus nidulans* genomic DNA fragment AMA1 sequence (Molecular Microbiology 1996 19:565-574) was inserted. Base 1 to 1134 contains *Aspergillus niger* glucoamylase gene promoter. Base 3098 to 3356 and 4950 to 4971 contains *Aspergillus niger* glucoamylase terminator. *Aspergillus nidulans* pyrG gene was inserted from 3357 to 4949 as a marker for fungal transformation. There is a multiple cloning site (MCS) into which genes may be inserted.

[22] Figure 7 is the pRAXdes2 vector backbone. This vector is based on the plasmid vector pRAX1. A Gateway cassette has been inserted into pRAX1 vector (indicated by the arrow on the interior of the circular plasmid). This cassette contains recombination sequence attR1 and attR2 and the selection marker catH and ccdB. The vector has been made according to the manual given in Gateway™ Cloning Technology: version 1 page 34-38 and can only replicate in *E. coli* DB3.1 from Invitrogen; in other *E. coli* hosts the ccdB gene is lethal. First a PCR fragment is made with primers containing attB1/2 recombination sequences. This fragment is recombined with pDONR201 (commercially available from Invitrogen); this vector contains attP1/2 recombination sequences with catH and ccdB in between the recombination sites. The BP clonase enzymes from Invitrogen are used to recombine the PCR fragment in this so-called ENTRY vector, clones with the

PCR fragment inserted can be selected at 50µg/ml kanamycin because clones expressing *ccdB* do not survive. Now the att sequences are altered and called attL1 and attL2. The second step is to recombine this clone with the pRAXdes2 vector (containing attR1 and attR2 catH and *ccdB* in between the recombination sites). The LR clonase enzymes from Invitrogen are used to recombine the insert from the ENTRY vector in the destination vector. Only pRAXCBH1 vectors are selected using 100µg/ml ampicillin because *ccdB* is lethal and the ENTRY vector is sensitive to ampicillin. By this method the expression vector is now prepared and can be used to transform *A. niger*. The *H. grisea* CBH1.1 expression is under the *Aspergillus* glucoamylase promoter and terminator control. The transformation marker *pyrG* gene and the AMA 1 sequence are from *Aspergillus nidulans*.

[23] Figure 8 provides an illustration of the pRAXdes2cbh1 vector which was used for expression of the nucleic acids encoding the CBH1.1 variants in *Aspergillus*. A nucleic acid encoding a CBH1.1 enzyme variant was cloned into the vector by homologous recombination of the att sequences.

[24] Figure 9 illustrates the activity of various CBH1 cellulases in a cellulose conversion assay at one day, 38°C, and 15 mg of total enzyme /g cellulose using pretreated corn stover (PCS) as a substrate. This assay combines at about a 50:50 mass ratio the CBH1.1 sample to be tested with supernatant from growths of a *T. reesei* strain which has been deleted for CBH1. At 38°C the *H. grisea* CBH1.1 is unremarkable in this assay.

[25] Legend: delCBH1 is a strain that has had its endogenous CBH1 deleted; it is CBH1 deficient. rCBH1 is a purified enzyme from an *A. niger* strain that has had a *T. reesei* CBH1 gene inserted. *A. niger* is a purified CBHB from an *A. niger* strain overexpressing its endogenous cellulase CBHB; *Hschweinitzii*/An is a purified CBH1 from an *A. niger* strain expressing an inserted heterologous CBH1 gene from *Hypocrea schweinitzii*. *Tpseudokonii*/An is a purified CBH1 from an *A. niger* strain expressing an inserted heterologous CBH1 gene from *Trichoderma pseudokoningii*. *Hgrisea*/An-1 is a purified CBH1.1 from a first *A. niger* clone expressing an inserted heterologous CBH1.1 gene from *H. grisea*. *Hgrisea*/An-2 is a purified CBH1.1 from a second *A. niger* clone expressing an inserted heterologous CBH1.1 gene from *H. grisea*. *Hgrisea*/An-1 and *Hgrisea*/An-2 are two clones from the same transformation of *A. niger* with the *H. grisea* variant CBH1.1 gene.

[26] Figure 10 shows the activity CBH1 cellulases in a cellulose conversion assay at one day, 65°C, and 15 mg of total enzyme /g cellulose using pretreated corn stover as a substrate. This assay combines at about a 50:50 mass ratio the CBH1 sample to be tested with supernatant from growths of a *T. reesei* strain which has been deleted for CBH1.

- 7 -

Unlike the 38°C result both *H. grisea* CBH1.1 samples are clearly better than other samples. Legend: same as for Figure 9.

[27] Figure 11 illustrates the activity of *H. grisea* CBH1.1 and *T. reesei* CBH1 cellulases in a cellulose conversion assay at one day, 38°C, and 15 mg of total enzyme /g cellulose using pretreated corn stover (PCS) as a substrate. This assay combines at about a 50:50 mass ratio the CBH1.1 sample to be tested with supernatant from the growth of a *T. reesei* strain which has been deleted for CBH1. At 38°C the *H. grisea* CBH1.1 is unremarkable in this assay. Native CBH1 (nCBH1) was purified from a *T. reesei* whole cellulase using methods known in the art (see Methods for Purifying CBH, below)..

[28] Figure 12 shows the activity results for the *H. grisea* and *T. reesei* rCBH1 enzymes only in a cellulose conversion assay at one day, 65°C, and 15 mg of total enzyme /g cellulose using pretreated corn stover as a substrate. This assay combines at about a 50:50 mass ratio the CBH1 sample to be tested with supernatant from the growth of a *T. reesei* strain which has been deleted for CBH1. Unlike the 38°C result the *H. grisea* CBH1.1 sample are is better than *T. reesei*.

[29] Figure 13A - C shows the activity CBH1 cellulases in a cellulose conversion assay on Phosphoric acid swollen cellulose (PASC) at various time points. In panel A, the temperature is 38°C with measurements being taken over a period of 120 minutes. In panel B, measurements were taken at the temperature 65°C. In panel C, measurements were taken at the temperature 70°C. It can be seen that the variant *H. grisea* CBH1.1 releases more cellobiose than the *T. reesei* CBH1 at any time point measured.

[30] Figure 14A – 14C shows the genomic DNA (SEQ ID NO:7), the cDNA (SEQ ID NO:8) and amino acid (SEQ ID NO:9) sequences of the *Scytalidium thermophilum* CBH1. The amino acid sequence includes the signal sequence.

[31] Figure 15 shows the alignment of the mature forms (i.e., without a signal sequence) of *H. grisea* CBH 1.1, *H. jecorina* CBH1 (SEQ ID NO: 10) and *Scytalidium thermophilum* CBH1 (SEQ ID NO:11). Also shown is the consensus sequence (SEQ ID NO:12). Six residues in the *H. jecorina* CBH1 catalytic domain are bold and underlined indicating the sites may be important for enhanced stability.

[32] Figure 16A and B show the thermostability profiles for *H. grisea* CBH1.1 and *S. thermophilum* CBH1.

DETAILED DESCRIPTION

[33] The invention will now be described in detail by way of reference only using the following definitions and examples. All patents and publications, including all sequences

disclosed within such patents and publications.

[34] Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 2D ED., John Wiley and Sons, New York (1994), and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY, Harper Perennial, NY (1991) provide one of skill with a general dictionary of many of the terms used in this invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. Practitioners are particularly directed to Sambrook *et al.*, 1989, and Ausubel FM *et al.*, 1993, for definitions and terms of the art. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary.

[35] The headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole.

[36]

I. DEFINITIONS

[37] "Cellulase," "cellulolytic enzymes" or "cellulase enzymes" means bacterial, or fungal exoglucanases or exocellobiohydrolases, and/or endoglucanases, and/or β -glucosidases.

[38] The term "cellulase" refers to a category of enzymes capable of hydrolyzing cellulose polymers to shorter cello-oligosaccharide oligomers, cellobiose and/or glucose. Numerous examples of cellulases, such as exoglucanases, exocellobiohydrolases, endoglucanases, and glucosidases have been obtained from cellulolytic organisms, particularly including fungi, and bacteria. The enzymes made by these microbes are mixtures of proteins with three types of actions useful in the conversion of cellulose to glucose: endoglucanases (EG), cellobiohydrolases (CBH), and beta-glucosidase. These

three different types of cellulase enzymes act synergistically to convert cellulose and its derivatives to glucose.

[39] Many microbes make enzymes that hydrolyze cellulose, including the fungus *Trichoderma*, the compost bacteria *Thermomonospora*, *Bacillus*, and *Cellulomonas*; *Streptomyces*; and the fungi *Humicola*, *Aspergillus* and *Fusarium*.

[40] "Cellobiohydrolase 1 activity" means a cellulose 1,4-beta-cellobiosidase (also called Exo-glucanase, Exo-cellobiohydrolase, CBH1 or 1,4-beta-cellobiohydrolase) activity, as defined in the enzyme class EC3.2.1.91 which catalyzes the hydrolysis of 1,4-beta-D-glucosidic linkages in cellulose and cellotetraose, releasing cellobiose from the non-reducing ends of the cellulose chains. For the present invention, the CBH1 activity may be determined according to the procedure described in Example 4.

[41] By the term "host cell" is meant a cell that contains a vector and supports the replication, and/or transcription or transcription and translation (expression) of the expression construct. Host cells for use in the present invention can be prokaryotic cells, such as *E. coli*, or eukaryotic cells such as yeast, plant, insect, amphibian, or mammalian cells. In general, host cells are filamentous fungi.

[42] The term "recombinant" when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

[43] The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide" or "secretory signal peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger peptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway to yield the secretory signal peptide and a smaller peptide commonly referred to as the mature polypeptide.

[44] As used herein, the phrases "whole cellulase preparation" and "whole cellulase composition" are used interchangeably and refer to both naturally occurring and non-naturally occurring compositions. A "naturally occurring" composition is one produced by a naturally occurring source and which comprises one or more cellobiohydrolase-type, one or more endoglucanase-type, and one or more β -glucosidase components wherein each of these components is found at the ratio produced by the source. A naturally occurring composition is one that is produced by an organism unmodified with respect to the

cellulolytic enzymes such that the ratio of the component enzymes is unaltered from that produced by the native organism.

[45] A "non-naturally occurring" composition encompasses those compositions produced by: (1) combining component cellulolytic enzymes either in a naturally occurring ratio or non-naturally occurring, i.e., altered, ratio; or (2) modifying an organism to overexpress or underexpress one or more cellulolytic enzyme; or (3) modifying an organism such that at least one cellulolytic enzyme is deleted or (4) modifying an organism to express a heterologous component cellulolytic enzyme.

[46] As used herein, the term "promoter" refers to a nucleic acid sequence that functions to direct transcription of a downstream gene. The promoter will generally be appropriate to the host cell in which the target gene is being expressed. The promoter together with other transcriptional and translational regulatory nucleic acid sequences (also termed "control sequences") are necessary to express a given gene. In general, the transcriptional and translational regulatory sequences include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences. The promoter may be the promoter normally associated with the downstream gene or it may be heterologous, i.e., from another gene or another microorganism as long as it function to direct the gene. In one aspect the promoter is an inducible promoter. In one aspect the promoter is the *T. reesei cbh1* promoter which is deposited in GenBank under Accession Number D86235. In another aspect the promoter is a *cbh II* or xylanase promoter from *T. reesei*.

[47] Examples include the promoter from the *A. awamori* or *A. niger* glucoamylase genes (Nunberg, J. H. et al. (1984) Mol. Cell. Biol. 4, 2306-2315; Boel, E. et al. (1984) EMBO J. 3, 1581-1585), the *Mucor miehei* carboxyl protease gene herein, the *Trichoderma reesei* cellobiohydrolase I gene (Shoemaker, S. P. et al. (1984) European Patent Application No. EPO0137280A1), the *A. nidulans trpC* gene (Yelton, M. et al. (1984) Proc. Natl. Acad. Sci. USA 81, 1470-1474; Mullaney, E. J. et al. (1985) Mol. Gen. Genet. 199, 37-45) the *A. nidulans alcA* gene (Lockington, R. A. et al. (1986) Gene 33, 137-149), the *A. nidulans tpiA* gene (McKnight, G. L. et al. (1986) Cell 46, 143-147), the *A. nidulans amdS* gene (Hynes, M. J. et al. (1983) Mol. Cell Biol. 3, 1430-1439), the *T. reesei xln1* gene, the *T. reesei cbh2* gene, the *T. reesei egl1* gene, the *T. reesei egl2* gene, the *T. reesei egl3* gene, and higher eukaryotic promoters such as the SV40 early promoter (Barclay, S. L. and E. Meller (1983) Molecular and Cellular Biology 3, 2117-2130).

[48] A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA encoding a secretory leader, i.e.,

a signal peptide, is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[49] As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, that may or may not include regions preceding and following the coding region, e.g. 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

[50] The gene may encode therapeutically significant proteins or peptides, such as growth factors, cytokines, ligands, receptors and inhibitors, as well as vaccines and antibodies. The gene may encode commercially important industrial proteins or peptides, such as enzymes, e.g., cellulases such as an *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1. The gene of interest may be a naturally occurring gene, a mutated gene or a synthetic gene.

[51] The "filamentous fungi" of the present invention are eukaryotic microorganisms and include all filamentous forms of the subdivision Eumycotina (see Alexopoulos, C. J. (1962), Introductory Mycology, New York: Wiley). These fungi are characterized by a vegetative mycelium with a cell wall composed of chitin, cellulose, and other complex polysaccharides. The filamentous fungi of the present invention are morphologically, physiologically, and genetically distinct from yeasts. Vegetative growth by filamentous fungi is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *S. cerevisiae* is by budding of a unicellular thallus, and carbon catabolism may be fermentative. *S. cerevisiae* has a prominent, very stable diploid phase, whereas diploids exist only briefly prior to meiosis in filamentous fungi, e.g., *Aspergilli* and *Neurospora*. *S. cerevisiae* has 17 chromosomes as opposed to 8 and 7 for *A. nidulans* and *N. crassa* respectively. Recent illustrations of differences between *S. cerevisiae* and filamentous fungi include the inability of *S. cerevisiae* to process *Aspergillus* and *Trichoderma* introns and the inability to recognize many transcriptional regulators of filamentous fungi (Innis, M. A. et al. (1985) Science, 228, 21-26).

[52] The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not normally found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences, e.g., from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein will often refer to two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

[53] The terms "isolated" or "purified" as used herein refer to a nucleic acid or amino acid that is removed from at least one component with which it is naturally associated.

[54] In the present context, the term "substantially pure polypeptide" means a polypeptide preparation which contains at the most 10% by weight of other polypeptide material with which it is natively associated (lower percentages of other polypeptide material are preferred, e.g. at the most 8% by weight, at the most 6% by weight, at the most 5% by weight, at the most 4% at the most 3% by weight, at the most 2% by weight, at the most 1% by weight, and at the most 1/2% by weight). Thus, it is preferred that the substantially pure polypeptide is at least 92% pure, i.e. that the polypeptide constitutes at least 92% by weight of the total polypeptide material present in the preparation, and higher percentages are preferred such as at least 94% pure, at least 95% pure, at least 96% pure, at least 96% pure, at least 97% pure, at least 98% pure, at least 99%, and at the most 99.5% pure. The polypeptides disclosed herein are preferably in a substantially pure form. In particular, it is preferred that the polypeptides disclosed herein are in "essentially pure form", i.e. that the polypeptide preparation is essentially free of other polypeptide material with which it is natively associated. This can be accomplished, for example, by preparing the polypeptide by means of well-known recombinant methods. Herein, the term "substantially pure polypeptide" is synonymous with the terms "isolated polypeptide" and "polypeptide in isolated form".

[55] In general, nucleic acid molecules which encode the variant *H. grisea* CBH1.1 will hybridize, under moderate to high stringency conditions to the sequence provided herein as SEQ ID NO:1 (the variant *H. grisea* CBH1.1). However, in some cases a CBH1-encoding nucleotide sequence is employed that possesses a substantially different codon usage, while the protein encoded by the CBH1-encoding nucleotide sequence has the same or substantially the same amino acid sequence as the native protein. For example, the coding sequence may be modified to facilitate faster expression of CBH1 in a particular prokaryotic or eukaryotic expression system, in accordance with the frequency

with which a particular codon is utilized by the host. Te'o, *et al.* (2000), for example, describes the optimization of genes for expression in filamentous fungi.

[56] A nucleic acid sequence is considered to be "selectively hybridizable" to a reference nucleic acid sequence if the two sequences specifically hybridize to one another under moderate to high stringency hybridization and wash conditions. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe. For example, "maximum stringency" typically occurs at about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe); "high stringency" at about $5-10^\circ\text{C}$ below the T_m ; "moderate" or "intermediate stringency" at about $10-20^\circ\text{C}$ below the T_m of the probe; and "low stringency" at about $20-25^\circ\text{C}$ below the T_m . Functionally, maximum stringency conditions may be used to identify sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to identify sequences having about 80% or more sequence identity with the probe.

[57] Moderate and high stringency hybridization conditions are well known in the art (see, for example, Sambrook, *et al.*, 1989, Chapters 9 and 11, and in Ausubel, F.M., *et al.*, 1993). An example of high stringency conditions includes hybridization at about 42°C in 50% formamide, 5X SSC, 5X Denhardt's solution, 0.5% SDS and 100 $\mu\text{g/ml}$ denatured carrier DNA followed by washing two times in 2X SSC and 0.5% SDS at room temperature and two additional times in 0.1X SSC and 0.5% SDS at 42°C .

[58] The term "% homology" is used interchangeably herein with the term "% identity" herein and refers to the level of nucleic acid or amino acid sequence identity between the nucleic acid sequence that encodes an *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 amino acid sequence, when aligned using a sequence alignment program.

[59] For example, as used herein, 80% homology means the same thing as 80% sequence identity determined by a defined algorithm, and accordingly a homologue of a given sequence has greater than 80% sequence identity over a length of the given sequence. Exemplary levels of sequence identity include, but are not limited to, 80, 85, 90, 95, 98% or more sequence identity to a given sequence, e.g., the coding sequence for an *H. grisea* CBH1.1 variant, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1, as described herein.

[60] Exemplary computer programs which can be used to determine identity between two sequences include, but are not limited to, the suite of BLAST programs, e.g., BLASTN, BLASTX, and TBLASTX, BLASTP and TBLASTN.

See also, Altschul, *et al.*, 1990 and Altschul, *et al.*, 1997.

[61] Sequence searches are typically carried out using the BLASTN program when evaluating a given nucleic acid sequence relative to nucleic acid sequences in the GenBank DNA Sequences and other public databases. The BLASTX program is preferred for searching nucleic acid sequences that have been translated in all reading frames against amino acid sequences in the GenBank Protein Sequences and other public databases. Both BLASTN and BLASTX are run using default parameters of an open gap penalty of 11.0, and an extended gap penalty of 1.0, and utilize the BLOSUM-62 matrix. (See, *e.g.*, Altschul, *et al.*, 1997.)

[62] A preferred alignment of selected sequences in order to determine "% identity" between two or more sequences, is performed using for example, the CLUSTAL-W program in MacVector version 6.5, operated with default parameters, including an open gap penalty of 10.0, an extended gap penalty of 0.1, and a BLOSUM 30 similarity matrix.

[63] As used herein, the terms "transformed", "stably transformed" or "transgenic" with reference to a cell means the cell has a non-native (heterologous) nucleic acid sequence integrated into its genome or as an episomal plasmid that is maintained through multiple generations.

[64] As used herein, the term "expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

[65] The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

[66] It follows that the term "CBH1.1 expression" refers to transcription and translation of the variant CBH1.1 cellulase gene, the products of which include precursor RNA, mRNA, polypeptide, post-translationally processed polypeptides. By way of example, assays for CBH1.1 expression include Western blot for CBH1.1 protein, Northern blot analysis and reverse transcriptase polymerase chain reaction (RT-PCR) assays for CBH1 mRNA, and endoglucanase activity assays as described in Shoemaker S.P. and Brown R.D.Jr. (Biochim. Biophys. Acta, 1978, 523:133-146) and Schulein (1988). Similarly, as used herein, "CBH1 expression" refers to transcription and translation of a *H. jecorina*

CBH1 variant or a *S. thermophilum* CBH1 cellulase gene, the products of which include precursor RNA, mRNA, polypeptide, post-translationally processed polypeptides.

[67] As used herein, the term "surfactant" refers to any compound generally recognized in the art as having surface active qualities. Thus, for example, surfactants comprise anionic, cationic and nonionic surfactants such as those commonly found in detergents. Anionic surfactants include linear or branched alkylbenzenesulfonates; alkyl or alkenyl ether sulfates having linear or branched alkyl groups or alkenyl groups; alkyl or alkenyl sulfates; olefinsulfonates; and alkanesulfonates. Ampholytic surfactants include quaternary ammonium salt sulfonates, and betaine-type ampholytic surfactants. Such ampholytic surfactants have both the positive and negative charged groups in the same molecule. Nonionic surfactants may comprise polyoxyalkylene ethers, as well as higher fatty acid alkanolamides or alkylene oxide adduct thereof, fatty acid glycerine monoesters, and the like.

[68] As used herein, the term "cellulose containing fabric" refers to any sewn or unsewn fabrics, yarns or fibers made of cotton or non-cotton containing cellulose or cotton or non-cotton containing cellulose blends including natural cellulosics and manmade cellulosics (such as jute, flax, ramie, rayon, and lyocell).

[69] As used herein, the term "cotton-containing fabric" refers to sewn or unsewn fabrics, yarns or fibers made of pure cotton or cotton blends including cotton woven fabrics, cotton knits, cotton denims, cotton yarns, raw cotton and the like.

[70] As used herein, the term "stonewashing composition" refers to a formulation for use in stonewashing cellulose containing fabrics. Stonewashing compositions are used to modify cellulose containing fabrics prior to sale, *i.e.*, during the manufacturing process. In contrast, detergent compositions are intended for the cleaning of soiled garments and are not used during the manufacturing process.

[71] As used herein, the term "detergent composition" refers to a mixture which is intended for use in a wash medium for the laundering of soiled cellulose containing fabrics. In the context of the present invention, such compositions may include, in addition to cellulases and surfactants, additional hydrolytic enzymes, builders, bleaching agents, bleach activators, bluing agents and fluorescent dyes, caking inhibitors, masking agents, cellulase activators, antioxidants, and solubilizers.

[72] As used herein, the term "decrease or elimination in expression of the *cbh1.1* gene" means that either that the *cbh1.1* gene has been deleted from the genome and therefore cannot be expressed by the recombinant host microorganism; or that the *cbh1.1* gene has been modified such that a functional CBH1.1 enzyme is not produced by the host microorganism.

[73] The term "variant *cbh1.1 gene*" or "variant CBH1.1" means, respectively, that the nucleic acid sequence of the *cbh1.1* gene from *H. grisea* has been altered by removing, adding, and/or manipulating the coding sequence or the amino acid sequence of the expressed protein has been modified consistent with the invention described herein.

[74] The term "variant *cbh1 gene*" or "variant CBH1" means, respectively, that the nucleic acid sequence of the *cbh1* gene from *H. jecorina* has been altered by removing, adding, and/or manipulating the coding sequence or the amino acid sequence of the expressed protein has been modified consistent with the invention described herein.

[75] As used herein, the terms "active" and "biologically active" refer to a biological activity associated with a particular protein and are used interchangeably herein. For example, the enzymatic activity associated with a protease is proteolysis and, thus, an active protease has proteolytic activity. It follows that the biological activity of a given protein refers to any biological activity typically attributed to that protein by those of skill in the art.

[76] When employed in enzymatic solutions, the homolog or variant CBH1.1, *H. jecorina* CBH1 variant or *S. thermophilum* CBH1 component is generally added in an amount sufficient to allow the highest rate of release of soluble sugars from the biomass. The amount of homolog or variant CBH1.1, *H. jecorina* CBH1 variant or *S. thermophilum* CBH1 component added depends upon the type of biomass to be saccharified which can be readily determined by the skilled artisan. However, when employed, the weight percent of the variant CBH1.1, *H. jecorina* CBH1 variant or *S. thermophilum* CBH1 component relative to any EG type components present in the cellulase composition is from preferably about 1, preferably about 5, preferably about 10, preferably about 15, or preferably about 20 weight percent to preferably about 25, preferably about 30, preferably about 35, preferably about 40, preferably about 45 or preferably about 50 weight percent. Furthermore, preferred ranges may be about 0.5 to about 15 weight percent, about 0.5 to about 20 weight percent, from about 1 to about 10 weight percent, from about 1 to about 15 weight percent, from about 1 to about 20 weight percent, from about 1 to about 25 weight percent, from about 5 to about 20 weight percent, from about 5 to about 25 weight percent, from about 5 to about 30 weight percent, from about 5 to about 35 weight percent, from about 5 to about 40 weight percent, from about 5 to about 45 weight percent, from about 5 to about 50 weight percent, from about 10 to about 20 weight percent, from about 10 to about 25 weight percent, from about 10 to about 30 weight percent, from about 10 to about 35 weight percent, from about 10 to about 40 weight percent, from about 10 to about 45 weight percent, from about 10 to about 50 weight percent, from about 15 to about 20 weight percent, from about 15 to about 25 weight

percent, from about 15 to about 30 weight percent, from about 15 to about 35 weight percent, from about 15 to about 30 weight percent, from about 15 to about 45 weight percent, from about 15 to about 50 weight percent.

II. HOST ORGANISMS

[77] Filamentous fungi include all filamentous forms of the subdivision Eumycota and Oomycota. The filamentous fungi are characterized by vegetative mycelium having a cell wall composed of chitin, glucan, chitosan, mannan, and other complex polysaccharides, with vegetative growth by hyphal elongation and carbon catabolism that is obligately aerobic.

[78] In the present invention, the filamentous fungal parent cell may be a cell of a species of, but not limited to, *Trichoderma*, e.g., *Trichoderma longibrachiatum*, *Trichoderma viride*, *Trichoderma koningii*, *Trichoderma harzianum*; *Penicillium* sp.; *Humicola* sp., including *Humicola insolens* and *Humicola grisea*; *Chrysosporium* sp., including *C. lucknowense*; *Gliocladium* sp.; *Aspergillus* sp.; *Fusarium* sp., *Neurospora* sp., *Hypocrea* sp., and *Emericella* sp. As used herein, the term "*Trichoderma*" or "*Trichoderma* sp." refers to any fungal strains which have previously been classified as *Trichoderma* or are currently classified as *Trichoderma*.

[79] In one preferred embodiment, the filamentous fungal parent cell is an *Aspergillus niger*, *Aspergillus awamori*, *Aspergillus aculeatus*, or *Aspergillus nidulans* cell.

[80] In another preferred embodiment, the filamentous fungal parent cell is a *Trichoderma reesei* cell.

III. CELLULASES

[81] Cellulases are known in the art as enzymes that hydrolyze cellulose (beta-1,4-glucan or beta D-glucosidic linkages) resulting in the formation of glucose, cellobiose, celooligosaccharides, and the like. As set forth above, cellulases have been traditionally divided into three major classes: endoglucanases (EC 3.2.1.4) ("EG"), exoglucanases or cellobiohydrolases (EC 3.2.1.91) ("CBH") and beta-glucosidases (EC 3.2.1.21) ("BG"). (Knowles, *et al.*, 1987; Schulein, 1988).

[82] Certain fungi produce complete cellulase systems which include exo-cellobiohydrolases or CBH-type cellulases, endoglucanases or EG-type cellulases and beta-glucosidases or BG-type cellulases (Schulein, 1988). However, sometimes these systems lack CBH-type cellulases and bacterial cellulases also typically include little or no CBH-type cellulases. In addition, it has been shown that the EG components and CBH

components synergistically interact to more efficiently degrade cellulose. See, e.g., Wood, 1985. The different components, *i.e.*, the various endoglucanases and exocellobiohydrolases in a multi-component or complete cellulase system, generally have different properties, such as isoelectric point, molecular weight, degree of glycosylation, substrate specificity and enzymatic action patterns.

[83] Cellulase compositions have also been shown to degrade cotton-containing fabrics, resulting in reduced strength loss in the fabric (U.S. Patent No. 4,822,516), contributing to reluctance to use cellulase compositions in commercial detergent applications. Cellulase compositions comprising endoglucanase components have been suggested to exhibit reduced strength loss for cotton-containing fabrics as compared to compositions comprising a complete cellulase system.

[84] Cellulases have also been shown to be useful in degradation of cellulase biomass to ethanol (wherein the cellulase degrades cellulose to glucose and yeast or other microbes further ferment the glucose into ethanol), in the treatment of mechanical pulp (Pere *et al.*, 1996), for use as a feed additive (WO 91/04673) and in grain wet milling.

[85] Most CBHs and EGs have a multidomain structure consisting of a core domain separated from a cellulose binding domain (CBD) by a linker peptide (Suurnakki *et al.*, 2000). The core domain contains the active site whereas the CBD interacts with cellulose by binding the enzyme to it (van Tilbeurgh *et al.*, 1986; Tomme *et al.*, 1988). The CBDs are particularly important in the hydrolysis of crystalline cellulose. It has been shown that the ability of cellobiohydrolases to degrade crystalline cellulose clearly decreases when the CBD is absent (Linder and Teeri, 1997). However, the exact role and action mechanism of CBDs is still a matter of speculation. It has been suggested that the CBD enhances the enzymatic activity merely by increasing the effective enzyme concentration at the surface of cellulose (Stahlberg *et al.*, 1991), and/or by loosening single cellulose chains from the cellulose surface (Tormo *et al.*, 1996). Most studies concerning the effects of cellulase domains on different substrates have been carried out with core proteins of cellobiohydrolases, as their core proteins can easily be produced by limited proteolysis with papain (Tomme *et al.*, 1988). Numerous cellulases have been described in the scientific literature, examples of which include: from *Trichoderma reesei*:

Shoemaker, S. *et al.*, Bio/Technology, 1:691-696, 1983, which discloses CBHI; Teeri, T. *et al.*, Gene, 51:43-52, 1987, which discloses CBHII. Cellulases from species other than *Trichoderma* have also been described e.g., Ooi *et al.*, 1990, which discloses the cDNA sequence coding for endoglucanase F1-CMC produced by *Aspergillus aculeatus*; Kawaguchi T *et al.*, 1996, which discloses the cloning and sequencing of the cDNA encoding beta-glucosidase 1 from *Aspergillus aculeatus*; Sakamoto *et al.*, 1995, which

discloses the cDNA sequence encoding the endoglucanase CMCase-1 from *Aspergillus kawachii* IFO 4308; Saarilahti *et al.*, 1990 which discloses an endoglucanase from *Erwinia carotovora*; Spilliaert R, *et al.*, 1994, which discloses the cloning and sequencing of *bglA*, coding for a thermostable beta-glucanase from *Rhodothermus marinus*; and Halldorsdottir S *et al.*, 1998, which discloses the cloning, sequencing and overexpression of a *Rhodothermus marinus* gene encoding a thermostable cellulase of glycosyl hydrolase family 12. However, there remains a need for identification and characterization of novel cellulases, with improved properties, such as improved performance under conditions of thermal stress or in the presence of surfactants, increased specific activity, altered substrate cleavage pattern, and/or high level expression *in vitro*.

[86] The development of new and improved cellulase compositions that comprise varying amounts CBH-type cellulase is of interest for use: (1) in detergent compositions that exhibit enhanced cleaning ability, function as a softening agent and/or improve the feel of cotton fabrics (e.g., "stone washing" or "biopolishing"); (2) in compositions for degrading wood pulp or other biomass into sugars (e.g., for bio-ethanol production); and/or (3) in feed compositions.

IV. MOLECULAR BIOLOGY

[87] In one embodiment this invention provides for the expression of variant *H. grisea* CBH1.1 cellulase genes under the control of a promoter functional in a filamentous fungus. Therefore, this invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2nd ed. 1989); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and Ausubel *et al.*, eds., *Current Protocols in Molecular Biology* (1994)).

A. Methods for Identifying Variant CBH1.1 Genes

[88] Two publically available *H. grisea* CBH1.1 nucleic acid sequences are shown in Figure 5. The invention, in one aspect, encompasses a nucleic acid molecule encoding a variant *H. grisea* CBH1.1 described herein. The nucleic acid may be a DNA molecule.

[89] Techniques that can be used to isolate variant CBH1.1-encoding DNA sequences are well known in the art and include, but are not limited to, cDNA and/or genomic library screening with a homologous DNA probes and expression screening with activity assays or antibodies against CBH1. Any of these methods can be found in Sambrook, *et al.* or in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, F. Ausubel, *et al.*, ed. Greene Publishing and Wiley-Interscience, New York (1987) ("Ausubel").

B. Nucleic Acid Constructs/Expression Vectors.

[90] Natural or synthetic polynucleotide fragments encoding a variant *H. grisea* CBH1.1 cellulase ("variant *H. grisea* CBH1.1 cellulase-encoding nucleic acid sequences") may be incorporated into heterologous nucleic acid constructs or vectors, capable of introduction into, and replication in, a filamentous fungal or yeast cell. The vectors and methods disclosed herein are suitable for use in host cells for the expression of a variant CBH1.1 cellulase. Any vector may be used as long as it is replicable and viable in the cells into which it is introduced. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. Cloning and expression vectors are also described in Sambrook *et al.*, 1989, Ausubel FM *et al.*, 1989, and Strathern *et al.*, 1981.

Appropriate expression vectors for fungi are described in van den Hondel, C.A.M.J.J. *et al.* (1991) In: Bennett, J.W. and Lasure, L.L. (eds.) *More Gene Manipulations in Fungi*. Academic Press, pp. 396-428. The appropriate DNA sequence may be inserted into a plasmid or vector (collectively referred to herein as "vectors") by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by standard procedures. Such procedures and related sub-cloning procedures are deemed to be within the scope of knowledge of those skilled in the art.

[91] Recombinant filamentous fungi comprising the coding sequence for a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase may be produced by introducing a heterologous nucleic acid construct comprising the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase coding sequence into the cells of a selected strain of the filamentous fungi.

[92] Once the desired form of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase nucleic acid sequence is obtained, it may be modified in a variety of ways. Where the sequence involves non-coding flanking regions, the flanking regions may be subjected to resection, mutagenesis, etc. Thus, transitions, transversions, deletions, and insertions may be performed on the naturally occurring sequence.

[93] A selected variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase coding sequence may be inserted into a suitable vector according to well-known recombinant techniques and used to transform filamentous fungi capable of cellulase expression. Due to the inherent degeneracy of the genetic code, other nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence may be used to clone and express a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase. Therefore it is appreciated that such

substitutions in the coding region fall within the sequence variants covered by the present invention.

[94] The present invention also includes recombinant nucleic acid constructs comprising one or more of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase-encoding nucleic acid sequences as described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation.

[95] Heterologous nucleic acid constructs may include the coding sequence for a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase: (i) in isolation; (ii) in combination with additional coding sequences; such as fusion protein or signal peptide coding sequences, where the desired cellulase coding sequence is the dominant coding sequence; (iii) in combination with non-coding sequences, such as introns and control elements, such as promoter and terminator elements or 5' and/or 3' untranslated regions, effective for expression of the coding sequence in a suitable host; and/or (iv) in a vector or host environment in which the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase coding sequence is a heterologous gene.

[96] In one aspect of the present invention, a heterologous nucleic acid construct is employed to transfer a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase-encoding nucleic acid sequence into a cell *in vitro*, with established filamentous fungal and yeast lines preferred. For long-term, production of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase, stable expression is preferred. It follows that any method effective to generate stable transformants may be used in practicing the invention.

[97] Appropriate vectors are typically equipped with a selectable marker-encoding nucleic acid sequence, insertion sites, and suitable control elements, such as promoter and termination sequences. The vector may comprise regulatory sequences, including, for example, non-coding sequences, such as introns and control elements, *i.e.*, promoter and terminator elements or 5' and/or 3' untranslated regions, effective for expression of the coding sequence in host cells (and/or in a vector or host cell environment in which a modified soluble protein antigen coding sequence is not normally expressed), operably linked to the coding sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, many of which are commercially available and/or are described in Sambrook, *et al.*, (*supra*).

[98] Exemplary promoters include both constitutive promoters and inducible promoters, examples of which include a CMV promoter, an SV40 early promoter, an RSV promoter,

an EF-1 α promoter, a promoter containing the tet responsive element (TRE) in the tet-on or tet-off system as described (ClonTech and BASF), the beta actin promoter and the metallothionine promoter that can upregulated by addition of certain metal salts. A promoter sequence is a DNA sequence which is recognized by the particular filamentous fungus for expression purposes. It is operably linked to DNA sequence encoding a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 polypeptide. Such linkage comprises positioning of the promoter with respect to the initiation codon of the DNA sequence encoding the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 polypeptide in the disclosed expression vectors. The promoter sequence contains transcription and translation control sequence which mediate the expression of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 polypeptide. Examples include the promoters from the *Aspergillus niger*, *A. awamori* or *A. oryzae* glucoamylase, alpha-amylase, or alpha-glucosidase encoding genes; the *A. nidulans* *gpdA* or *trpC* Genes; the *Neurospora crassa* *cbh1* or *trp1* genes; the *A. niger* or *Rhizomucor miehei* aspartic proteinase encoding genes; the *H. jecorina* *cbh1*, *cbh2*, *egl1*, *egl2*, or other cellulase encoding genes.

[99] The choice of the proper selectable marker will depend on the host cell, and appropriate markers for different hosts are well known in the art. Typical selectable marker genes include *argB* from *A. nidulans* or *H. jecorina*, *amdS* from *A. nidulans*, *pyr4* from *Neurospora crassa* or *H. jecorina*, *pyrG* from *Aspergillus niger* or *A. nidulans*. Additional exemplary selectable markers include, but are not limited to *trpc*, *trp1*, *oliC31*, *niaD* or *leu2*, which are included in heterologous nucleic acid constructs used to transform a mutant strain such as *trp*-, *pyr*-, *leu*- and the like.

[100] Such selectable markers confer to transformants the ability to utilize a metabolite that is usually not metabolized by the filamentous fungi. For example, the *amdS* gene from *H. jecorina* which encodes the enzyme acetamidase that allows transformant cells to grow on acetamide as a nitrogen source. The selectable marker (e.g. *pyrG*) may restore the ability of an auxotrophic mutant strain to grow on a selective minimal medium or the selectable marker (e.g. *oliC31*) may confer to transformants the ability to grow in the presence of an inhibitory drug or antibiotic.

[101] The selectable marker coding sequence is cloned into any suitable plasmid using methods generally employed in the art. Exemplary plasmids include pUC18, pBR322, pRAX and pUC100. The pRAX plasmid contains AMA1 sequences from *A. nidulans*, which make it possible to replicate in *A. niger*.

[102] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and

immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, Sambrook *et al.*, 1989; Freshney, 1987; Ausubel, *et al.*, 1993; and Coligan *et al.*, 1991.

C. Methods for Transforming a Host Cell

[103] In the present invention, the filamentous fungal parent cell may be a cell of a species of, but not limited to, *Trichoderma*, e.g., *Trichoderma longibrachiatum* (reesei), *Trichoderma viride*, *Trichoderma koningii*, *Trichoderma harzianum*; *Penicillium* sp.; *Humicola* sp., including *Humicola insolens*; *Chrysosporium* sp., including *C. lucknowense*; *Gliocladium* sp.; *Aspergillus* sp.; *Fusarium* sp., *Neurospora* sp., *Hypocrea* sp., and *Emmericella* sp. As used herein, the term "*Trichoderma*" or "*Trichoderma* sp." refers to any fungal strains which have previously been classified as *Trichoderma* or are currently classified as *Trichoderma*.

[104] Examples of parental cell lines which may be treated and/or modified for variant *H. grisea* CBH1.1, *H. jecorina* CBH1 variant or *S. thermophilium* CBH1 expression include, but are not limited to, filamentous fungal cells. Examples of appropriate primary cell types for use in practicing the invention include, but are not limited to, *Aspergillus* and *Trichoderma*.

[105] In one embodiment, the filamentous fungal parent cell is an *Aspergillus niger*, *Aspergillus awamori*, *Aspergillus aculeatus*, or *Aspergillus nidulans* cell.

[106] In a second embodiment, the filamentous fungal parent cell is a *Hypocrea jecorina* cell. This cell was previously referred to as *T. reesei*.

[107] After DNA sequences that encode the CBH1.1 variants, *H. jecorina* CBH1 variant or *S. thermophilium* CBH1 have been cloned into DNA constructs, the DNA is used to transform microorganisms. The microorganism to be transformed for the purpose of expressing a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 according to the present invention may advantageously comprise a strain derived from *Trichoderma* sp. Thus, a preferred mode for preparing variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulases according to the present invention comprises transforming a *Trichoderma* sp. host cell with a DNA construct comprising at least a fragment of DNA encoding a portion or all of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1. The DNA construct will generally be functionally attached, i.e., operably linked, to a promoter. The transformed host cell is then grown under conditions so as to express the variant *H. grisea* CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1. Subsequently, the variant *H. grisea* CBH1.1, a

H. jecorina CBH1 variant or a *S. thermophilium* CBH1 may be isolated. It may be desirable to have the variant *H. grisea* CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 in a substantially pure form. Similarly, it may be desirable to have the variant *H. grisea* CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 in an essentially pure form.

[108] However, it may in fact be that the best expression vehicle for a given DNA encoding a variant CBH1.1 may differ from *H. jecorina* (i.e., *T. reesei*). Thus, it may be that it will be most advantageous to express a protein in a transformation host that bears phylogenetic similarity to the source organism for the variant CBH1.1. In an alternative embodiment, *Aspergillus niger* can be used as an expression vehicle. For a description of transformation techniques with *A. niger*, see WO 98/31821, the disclosure of which is incorporated by reference in its entirety.

[109] Accordingly, the present description of a *Trichoderma spp.* expression system is provided for illustrative purposes only and as one option for expressing the variant CBH1.1 of the invention. One of skill in the art, however, may be inclined to express the DNA encoding variant CBH1.1 in a different host cell if appropriate and it should be understood that the source of the variant CBH1.1 should be considered in determining the optimal expression host. Additionally, the skilled worker in the field will be capable of selecting the best expression system for a particular gene through routine techniques utilizing the tools available in the art.

D. Methods for Expressing a Variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1

[110] The methods of the invention rely on the use cells to express a variant CBH1.1 cellulase, with no particular method of expression required.

[111] The invention provides host cells that have been transduced, transformed or transfected with an expression vector comprising a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase-encoding nucleic acid sequence. The culture conditions, such as temperature, pH and the like, are those previously used for the parental host cell prior to transduction, transformation or transfection and will be apparent to those skilled in the art.

[112] In one approach, a filamentous fungal cell or yeast cell is transfected with an expression vector having a promoter or biologically active promoter fragment or one or more (e.g., a series) of enhancers which functions in the host cell line, operably linked to a DNA segment encoding a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S.*

thermophilium CBH1 cellulase, such that the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase is expressed in the cell line.

[113] Thus, the present invention provides filamentous fungi comprising cells which have been modified, selected and cultured in a manner effective to result in variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase production or expression relative to the corresponding non-transformed parental fungi.

[114] Examples of species of parental filamentous fungi that may be treated and/or modified for desired cellulase expression include, but are not limited to *Trichoderma*, *Penicillium* sp., *Humicola* sp., including *Humicola insolens*; *Aspergillus* sp., including *Aspergillus niger*, *Chrysosporium* sp., *Fusarium* sp., *Hypocrea* sp., and *Emericella* sp.

[115] Cells expressing a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase are cultured under conditions typically employed to culture the parental fungal line. Generally, cells are cultured in a standard medium containing physiological salts and nutrients, such as described in Pourquie, J. et al., *Biochemistry and Genetics of Cellulose Degradation*, eds. Aubert, J. P. et al., Academic Press, pp. 71-86, 1988 and Ilmen, M. et al., *Appl. Environ. Microbiol.* 63:1298-1306, 1997. Culture conditions are also standard, e.g., cultures are incubated at 28°C in shaker cultures or fermenters until desired levels of variant CBH1.1 cellulase expression are achieved.

[116] Preferred culture conditions for a given filamentous fungus may be found in the scientific literature and/or from the source of the fungi such as the American Type Culture Collection.

After fungal growth has been established, the cells are exposed to conditions effective to cause or permit the expression of a variant CBH1.1 cellulase.

[117] In cases where a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase coding sequence is under the control of an inducible promoter, the inducing agent, e.g., a sugar, metal salt or antibiotics, is added to the medium at a concentration effective to induce variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase expression.

[118] In one embodiment, the strain comprises *Aspergillus niger*, which is a useful strain for obtaining overexpressed protein. For example *A. niger* var *awamori* dgr246 is known to secrete elevated amounts of secreted cellulases (Goedegebuur et al, *Curr. Genet* (2002) 41: 89-98). Other strains of *Aspergillus niger* var *awamori* such as GCDAP3, GCDAP4 and GAP3-4 are known Ward et al (Ward, M, Wilson, L.J. and Kodama, K.H., 1993, *Appl. Microbiol. Biotechnol.* 39:738-743).

[119] In another embodiment, the strain comprises *Trichoderma reesei*, which is a useful strain for obtaining overexpressed protein. For example, RL-P37, described by Sheir-

Neiss, *et al.*, *Appl. Microbiol. Biotechnol.* 20:46-53 (1984) is known to secrete elevated amounts of cellulase enzymes. Functional equivalents of RL-P37 include *Trichoderma reesei* strain RUT-C30 (ATCC No. 56765) and strain QM9414 (ATCC No. 26921). It is contemplated that these strains would also be useful in overexpressing variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1.

[120] Where it is desired to obtain the desired cellulase in the absence of potentially detrimental native cellulolytic activity, it is useful to obtain a host cell strain which has had one or more cellulase genes deleted prior to introduction of a DNA construct or plasmid containing the DNA fragment encoding the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase. Such strains may be prepared by the method disclosed in U.S. Patent No. 5,246,853 and WO 92/06209.

By expressing a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase in a host microorganism that is missing one or more cellulase genes, the identification and subsequent purification procedures are simplified. Any gene from *Trichoderma* sp. which has been cloned can be deleted, for example, the *cbh1*, *cbh2*, *egl1*, and *egl2* genes as well as those encoding a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 protein (see e.g., U.S. Patent No. 5,475,101 and WO 94/28117, respectively).

[121] Gene deletion may be accomplished by inserting a form of the desired gene to be deleted or disrupted into a plasmid by methods known in the art. The deletion plasmid is then cut at an appropriate restriction enzyme site(s), internal to the desired gene coding region, and the gene coding sequence or part thereof replaced with a selectable marker. Flanking DNA sequences from the locus of the gene to be deleted or disrupted, preferably between about 0.5 to 2.0 kb, remain on either side of the selectable marker gene. An appropriate deletion plasmid will generally have unique restriction enzyme sites present therein to enable the fragment containing the deleted gene, including flanking DNA sequences, and the selectable marker gene to be removed as a single linear piece.

[122] A selectable marker must be chosen so as to enable detection of the transformed microorganism. Any selectable marker gene that is expressed in the selected microorganism will be suitable. For example, with *Aspergillus* sp., the selectable marker is chosen so that the presence of the selectable marker in the transformants will not significantly affect the properties thereof. Such a selectable marker may be a gene that encodes an assayable product. For example, a functional copy of a *Aspergillus* sp. gene may be used which if lacking in the host strain results in the host strain displaying an auxotrophic phenotype.

[123] In one embodiment, a *pyrG*⁻ derivative strain of *Aspergillus sp.* is transformed with a functional *pyrG* gene, which thus provides a selectable marker for transformation. A *pyrG*⁻ derivative strain may be obtained by selection of *Aspergillus sp.* strains that are resistant to fluoro-orotic acid (FOA). The *pyrG* gene encodes orotidine-5'-monophosphate decarboxylase, an enzyme required for the biosynthesis of uridine. Strains with an intact *pyrG* gene grow in a medium lacking uridine but are sensitive to fluoro-orotic acid. It is possible to select *pyrG*⁻ derivative strains that lack a functional orotidine monophosphate decarboxylase enzyme and require uridine for growth by selecting for FOA resistance. Using the FOA selection technique it is also possible to obtain uridine-requiring strains which lack a functional orotate pyrophosphoribosyl transferase. It is possible to transform these cells with a functional copy of the gene encoding this enzyme (Berges & Barreau, *Curr. Genet.* 19:359-365 (1991), and van Hartingsveldt et al., (1986) Development of a homologous transformation system for *Aspergillus niger* based on the *pyrG* gene. *Mol. Gen. Genet.* 206:71-75). Selection of derivative strains is easily performed using the FOA resistance technique referred to above, and thus, the *pyrG* gene is preferably employed as a selectable marker. In another embodiment, a *pyr4*⁻ derivative strain of *Trichoderma sp.* is transformed with a functional *pyr4* gene, which thus provides a selectable marker for transformation. Although the following discusses the *Aspergillus* system, similar procedures for *Trichoderma* and other fungal systems may be used as will be appreciated by one skilled in the art.

[124] To transform *pyrG*⁻ *Aspergillus sp.* so as to be lacking in the ability to express one or more cellulase genes, a single DNA fragment comprising a disrupted or deleted cellulase gene is then isolated from the deletion plasmid and used to transform an appropriate *pyr*⁻ *Aspergillus* host. Transformants are then identified and selected based on their ability to express the *pyrG* gene product and thus complement the uridine auxotrophy of the host strain. Southern blot analysis is then carried out on the resultant transformants to identify and confirm a double crossover integration event that replaces part or all of the coding region of the genomic copy of the gene to be deleted with the *pyr4* selectable markers.

[125] Although the specific plasmid vectors described above relate to preparation of *pyr*⁻ transformants, the present invention is not limited to these vectors. Various genes can be deleted and replaced in the *Aspergillus sp.* strain using the above techniques. In addition, any available selectable markers can be used, as discussed above. In fact, any

Aspergillus sp. gene that has been cloned, and thus identified, can be deleted from the genome using the above-described strategy.

[126] As stated above, the host strains used are derivatives of *Aspergillus sp.* that lack or have a nonfunctional gene or genes corresponding to the selectable marker chosen. For example, if the selectable marker of *pyrG* is chosen, then a specific *pyrG*⁻ derivative strain is used as a recipient in the transformation procedure. Similarly, selectable markers comprising *Aspergillus sp.* genes equivalent to the *Aspergillus nidulans* genes *amdS*, *argB*, *trpC*, *niaD* may be used. The corresponding recipient strain must therefore be a derivative strain such as *argB*⁻, *trpC*⁻, *niaD*⁻, respectively.

[127] DNA encoding the variant CBH1.1 cellulase is then prepared for insertion into an appropriate microorganism. According to the present invention, DNA encoding a variant CBH1.1 cellulase comprises the DNA necessary to encode for a protein that has functional cellulolytic activity. The DNA fragment encoding the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase may be functionally attached to a fungal promoter sequence, for example, the promoter of the *glaA* gene.

[128] It is also contemplated that more than one copy of DNA encoding a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase may be recombined into the strain to facilitate overexpression. The DNA encoding the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase may be prepared by the construction of an expression vector carrying the DNA encoding the cellulase. The expression vector carrying the inserted DNA fragment encoding the variant CBH1.1 cellulase may be any vector which is capable of replicating autonomously in a given host organism or of integrating into the DNA of the host, typically a plasmid. In preferred embodiments two types of expression vectors for obtaining expression of genes are contemplated. The first contains DNA sequences in which the promoter, gene-coding region, and terminator sequence all originate from the gene to be expressed. Gene truncation may be obtained where desired by deleting undesired DNA sequences (e.g., coding for unwanted domains) to leave the domain to be expressed under control of its own transcriptional and translational regulatory sequences. A selectable marker is also contained on the vector allowing the selection for integration into the host of multiple copies of the novel gene sequences.

[129] The second type of expression vector is preassembled and contains sequences required for high-level transcription and a selectable marker. It is contemplated that the coding region for a gene or part thereof can be inserted into this general-purpose expression vector such that it is under the transcriptional control of the expression

cassettes promoter and terminator sequences. For example, pRAX is such a general-purpose expression vector. Genes or part thereof can be inserted downstream of the strong *glaA* promoter. An example of an integrative expression vector is the pTrex vector. Genes or part thereof can be inserted downstream of the strong *cbh1* promoter.

[130] In the vector, the DNA sequence encoding the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase of the present invention should be operably linked to transcriptional and translational sequences, *i.e.*, a suitable promoter sequence and signal sequence in reading frame to the structural gene. The promoter may be any DNA sequence that shows transcriptional activity in the host cell and may be derived from genes encoding proteins either homologous or heterologous to the host cell. An optional signal peptide provides for extracellular production of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase. The DNA encoding the signal sequence is preferably that which is naturally associated with the gene to be expressed, however the signal sequence from any suitable source is contemplated in the present invention.

[131] The procedures used to fuse the DNA sequences coding for the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase of the present invention with the promoter into suitable vectors are well known in the art.

[132] Various methods may be employed for delivering an expression vector, DNA vector or construct described above into cells *in vitro*. Methods of introducing nucleic acids into cells for expression of heterologous nucleic acid sequences are also known to the ordinarily skilled artisan, including, but not limited to electroporation; nuclear microinjection or direct microinjection into single cells; bacterial protoplast fusion with intact cells; use of polycations, *e.g.*, polybrene or polyornithine; membrane fusion with liposomes, lipofectamine or lipofection-mediated transfection; high velocity bombardment with DNA-coated microprojectiles; incubation with calcium phosphate-DNA precipitate; DEAE-Dextran mediated transfection; infection with modified viral nucleic acids; *Agrobacterium*-mediated transfer of DNA; and the like. In addition, heterologous nucleic acid constructs comprising a variant CBH1.1-encoding nucleic acid sequence can be transcribed *in vitro*, and the resulting RNA introduced into the host cell by well-known methods, *e.g.*, by injection.

[133] The preferred method in the present invention to prepare *Aspergillus sp.* for transformation involves the preparation of protoplasts from fungal mycelium. See Campbell *et al.* Improved transformation efficiency of *A.niger* using homologous *niaD* gene for nitrate reductase. Curr. Genet. 16:53-56; 1989. The mycelium can be obtained from germinated vegetative spores. The mycelium is treated with an enzyme that digests the

cell wall resulting in protoplasts. The protoplasts are then protected by the presence of an osmotic stabilizer in the suspending medium. These stabilizers include sorbitol, mannitol, potassium chloride, magnesium sulfate and the like. Usually the concentration of these stabilizers varies between 0.8 M and 1.2 M. It is preferable to use about a 1.2 M solution of sorbitol in the suspension medium.

[134] Uptake of the DNA into the host *Aspergillus sp.* strain is dependent upon the calcium ion concentration. Generally between about 10 mM CaCl_2 and 50 mM CaCl_2 is used in an uptake solution. Besides the need for the calcium ion in the uptake solution, other items generally included are a buffering system such as TE buffer (10 Mm Tris, pH 7.4; 1 mM EDTA) or 10 mM MOPS, pH 6.0 buffer (morpholinepropanesulfonic acid) and polyethylene glycol (PEG). It is believed that the polyethylene glycol acts to fuse the cell membranes thus permitting the contents of the medium to be delivered into the cytoplasm of the *Aspergillus sp.* strain and the plasmid DNA is transferred to the nucleus. This fusion frequently leaves multiple copies of the plasmid DNA tenderly integrated into the host chromosome.

[135] Usually a suspension containing the *Aspergillus sp.* protoplasts or cells that have been subjected to a permeability treatment at a density of 10^5 to $10^6/\text{mL}$, preferably $2 \times 10^5/\text{mL}$ are used in transformation. A volume of 100 μL of these protoplasts or cells in an appropriate solution (e.g., 1.2 M sorbitol; 50 mM CaCl_2) are mixed with the desired DNA. Generally a high concentration of PEG is added to the uptake solution. From 0.1 to 1 volume of 25% PEG 4000 can be added to the protoplast suspension. However, it is preferable to add about 0.25 volumes to the protoplast suspension. Additives such as dimethyl sulfoxide, heparin, spermidine, potassium chloride and the like may also be added to the uptake solution and aid in transformation. Similar procedures are available for other fungal host cells. See, for example, U.S. Patent No. 6,268,328.

[136] Generally, the mixture is then incubated at approximately 0°C for a period of between 10 to 30 minutes. Additional PEG is then added to the mixture to further enhance the uptake of the desired gene or DNA sequence. The 25% PEG 4000 is generally added in volumes of 5 to 15 times the volume of the transformation mixture; however, greater and lesser volumes may be suitable. The 25% PEG 4000 is preferably about 10 times the volume of the transformation mixture. After the PEG is added, the transformation mixture is then incubated either at room temperature or on ice before the addition of a sorbitol and CaCl_2 solution. The protoplast suspension is then further added to molten aliquots of a growth medium. This growth medium permits the growth of transformants only. Any growth medium can be used in the present invention that is

suitable to grow the desired transformants. However, if *Pyr*⁺ transformants are being selected it is preferable to use a growth medium that contains no uridine. The subsequent colonies are transferred and purified on a growth medium depleted of uridine.

[137] At this stage, stable transformants may be distinguished from unstable transformants by their faster growth rate and the formation of circular colonies with a smooth, rather than ragged outline on solid culture medium lacking uridine. Additionally, in some cases a further test of stability may be made by growing the transformants on solid non-selective medium (*i.e.* containing uridine), harvesting spores from this culture medium and determining the percentage of these spores which will subsequently germinate and grow on selective medium lacking uridine. Alternatively, other methods known in the art may be used to select transformants.

[138] In a particular embodiment of the above method, the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase(s) are recovered in active form from the host cell after growth in liquid media either as a result of the appropriate post translational processing of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase.

E. Methods of Analysis For CBH1 Nucleic Acid Coding Sequences and/or Protein Expression.

[139] In order to evaluate the expression of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase by a cell line that has been transformed with a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase-encoding nucleic acid construct, assays can be carried out at the protein level, the RNA level or by use of functional bioassays particular to cellobiohydrolase activity and/or production.

[140] In one exemplary application of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase nucleic acid and protein sequences described herein, a genetically modified strain of filamentous fungi, *e.g.*, *Trichoderma reesei*, is engineered to produce an increased amount of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase. Such genetically modified filamentous fungi would be useful to produce a cellulase product with greater increased cellulolytic capacity. In one approach, this is accomplished by introducing the coding sequence for a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase into a suitable host, *e.g.*, a filamentous fungi such as *Aspergillus niger*.

[141] Accordingly, the invention includes methods for expressing a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase in a filamentous fungus or

other suitable host by introducing an expression vector containing the DNA sequence encoding a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase into cells of the filamentous fungus or other suitable host.

[142] In another aspect, the invention includes methods for modifying the expression of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase in a filamentous fungus or other suitable host. Such modification includes a decrease or elimination in expression of the endogenous CBH.

[143] In general, assays employed to analyze the expression of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase include, Northern blotting, dot blotting (DNA or RNA analysis), RT-PCR (reverse transcriptase polymerase chain reaction), or *in situ* hybridization, using an appropriately labeled probe (based on the nucleic acid coding sequence) and conventional Southern blotting and autoradiography.

[144] In addition, the production and/or expression of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase may be measured in a sample directly, for example, by assays for cellobiohydrolase activity, expression and/or production. Such assays are described, for example, in Becker et al., Biochem J. (2001) 356:19-30 and Mitsuishi et al., FEBS (1990) 275:135-138.

The ability of CBH1 to hydrolyze isolated soluble and insoluble substrates can be measured using assays described in Srisodsuk et al., J. Biotech. (1997) 57:49-57 and Nidetzky and Claeysens Biotech. Bioeng. (1994) 44:961-966. Substrates useful for assaying cellobiohydrolase, endoglucanase or β -glucosidase activities include crystalline cellulose, filter paper, phosphoric acid swollen cellulose, cellooligosaccharides, methylumbelliferyl lactoside, methylumbelliferyl cellobioside, orthonitrophenyl lactoside, paranitrophenyl lactoside, orthonitrophenyl cellobioside, paranitrophenyl cellobioside.

[145] In addition, protein expression, may be evaluated by immunological methods, such as immunohistochemical staining of cells, tissue sections or immunoassay of tissue culture medium, e.g., by Western blot or ELISA. Such immunoassays can be used to qualitatively and quantitatively evaluate expression of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase. The details of such methods are known to those of skill in the art and many reagents for practicing such methods are commercially available.

[146] A purified form of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase may be used to produce either monoclonal or polyclonal antibodies specific to the expressed protein for use in various immunoassays. (See, e.g., Hu et al., 1991). Exemplary assays include ELISA, competitive immunoassays,

radioimmunoassays, Western blot, indirect immunofluorescent assays and the like. In general, commercially available antibodies and/or kits may be used for the quantitative immunoassay of the expression level of cellobiohydrolase proteins.

F. Methods for Purifying a CBH1

[147] In general, a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase protein produced in cell culture is secreted into the medium and may be purified or isolated, e.g., by removing unwanted components from the cell culture medium. However, in some cases, a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase protein may be produced in a cellular form necessitating recovery from a cell lysate. In such cases the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase protein is purified from the cells in which it was produced using techniques routinely employed by those of skill in the art. Examples include, but are not limited to, affinity chromatography (Tilbeurgh *et al.*, 1984), ion-exchange chromatographic methods (Goyal *et al.*, 1991; Fliess *et al.*, 1983; Bhikhabhai *et al.*, 1984; Ellouz *et al.*, 1987), including ion-exchange using materials with high resolution power (Medve *et al.*, 1998), hydrophobic interaction chromatography (Tomaz and Queiroz, 1999), and two-phase partitioning (Brumbauer, *et al.*, 1999).

[148] Typically, the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase protein is fractionated to segregate proteins having selected properties, such as binding affinity to particular binding agents, e.g., antibodies or receptors; or which have a selected molecular weight range, or range of isoelectric points.

[149] Once expression of a given variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase protein is achieved, the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase protein thereby produced is purified from the cells or cell culture. Exemplary procedures suitable for such purification include the following: antibody-affinity column chromatography, ion exchange chromatography; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; and gel filtration using, e.g., Sephadex G-75. Various methods of protein purification may be employed and such methods are known in the art and described e.g. in Deutscher, 1990; Scopes, 1982. The purification step(s) selected will depend, e.g., on the nature of the production process used and the particular protein produced.

V. Utility of *cbh1* and CBH1

[150] It can be appreciated that the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase nucleic acids, the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase protein and compositions comprising variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase protein activity find utility in a wide variety applications, some of which are described below.

[151] New and improved cellulase compositions that comprise varying amounts of a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase find utility in detergent compositions that exhibit enhanced cleaning ability, function as a softening agent and/or improve the feel of cotton fabrics (e.g., "stone washing" or "biopolishing"), in compositions for degrading wood pulp into sugars (e.g., for bio-ethanol production), and/or in feed compositions. The isolation and characterization of cellulase of each type provides the ability to control the aspects of such compositions.

[152] Variant CBH1.1, *H. jecorina* CBH1 variant or *S. thermophilium* CBH1 cellulases with decreased thermostability find uses, for example, in areas where the enzyme activity is required to be neutralized at lower temperatures so that other enzymes that may be present are left unaffected. In addition, the enzymes may find utility in the limited conversion of cellulotics, for example, in controlling the degree of crystallinity or of cellulosic chain-length. After reaching the desired extent of conversion the saccharifying temperature can be raised above the survival temperature of the de-stabilized CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1. As the CBH1 activity is essential for hydrolysis of crystalline cellulose, conversion of crystalline cellulose will cease at the elevated temperature.

[153] In one approach, the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase of the invention finds utility in detergent compositions or in the treatment of fabrics to improve the feel and appearance.

[154] Since the rate of hydrolysis of cellulosic products may be increased by using a transformant having at least one additional copy of the variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilium* CBH1 cellulase gene, either as a replicative plasmid or inserted into the genome, products that contain cellulose or heteroglycans can be degraded at a faster rate and to a greater extent. Products made from cellulose such as paper, cotton, cellulosic diapers and the like can be degraded more efficiently in a landfill. Thus, the fermentation product obtainable from the transformants or the transformants alone may be used in compositions to help degrade by liquefaction a variety of cellulose products that add to the overcrowded landfills.

[155] Separate saccharification and fermentation is a process whereby cellulose present in biomass, e.g., corn stover, is converted to glucose and subsequently yeast strains convert glucose into ethanol. Simultaneous saccharification and fermentation is a process whereby cellulose present in biomass, e.g., corn stover, is converted to glucose and, at the same time and in the same reactor, yeast strains convert glucose into ethanol. Thus, in another approach, the desired cellulase of the invention finds utility in the degradation of biomass to ethanol. Ethanol production from readily available sources of cellulose provides a stable, renewable fuel source.

[156] Cellulose-based feedstocks are comprised of agricultural wastes, grasses and woods and other low-value biomass such as municipal waste (e.g., recycled paper, yard clippings, etc.). Ethanol may be produced from the fermentation of any of these cellulosic feedstocks. However, the cellulose must first be converted to sugars before there can be conversion to ethanol. Thus, the CBH1 cellulase described herein find use in the conversion of biomass to sugars.

[157] A large variety of feedstocks may be used with the inventive variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase(s) and the one selected for use may depend on the region where the conversion is being done. For example, in the Midwestern United States agricultural wastes such as wheat straw, corn stover and bagasse may predominate while in California rice straw may predominate. However, it should be understood that any available cellulosic biomass may be used in any region.

[158] A cellulase composition containing an enhanced amount of cellobiohydrolase finds utility in ethanol production. Ethanol from this process can be further used as an octane enhancer or directly as a fuel in lieu of gasoline which is advantageous because ethanol as a fuel source is more environmentally friendly than petroleum derived products. It is known that the use of ethanol will improve air quality and possibly reduce local ozone levels and smog. Moreover, utilization of ethanol in lieu of gasoline can be of strategic importance in buffering the impact of sudden shifts in non-renewable energy and petrochemical supplies.

[159] Ethanol can be produced via saccharification and fermentation processes from cellulosic biomass such as trees, herbaceous plants, municipal solid waste and agricultural and forestry residues. However, the ratio of individual cellulase enzymes within a naturally occurring cellulase mixture produced by a microbe may not be the most efficient for rapid conversion of cellulose in biomass to glucose. It is known that endoglucanases act to produce new cellulose chain ends which themselves are substrates for the action of cellobiohydrolases and thereby improve the efficiency of

hydrolysis of the entire cellulase system. Therefore, the use of increased or optimized cellobiohydrolase activity may greatly enhance the production of ethanol.

[160] Thus, the inventive cellobiohydrolase(s) finds use in the hydrolysis of cellulose to its sugar components. In one embodiment, a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase is added to the biomass prior to the addition of a fermentative organism. In a second embodiment, a variant CBH1.1, a *H. jecorina* CBH1 variant or a *S. thermophilum* CBH1 cellulase is added to the biomass at the same time as a fermentative organism. Optionally, there may be other cellulase components present in either embodiment. Enhanced cellulose conversion may be achieved at higher temperatures using the CBH1 polypeptides described herein.

[161] In another embodiment the cellulosic feedstock may be pretreated. Pretreatment may be by elevated temperature and the addition of either of dilute acid, concentrated acid or dilute alkali solution. The pretreatment solution is added for a time sufficient to at least partially hydrolyze the hemicellulose components and then neutralized.

[162] The detergent compositions of this invention may employ besides the cellulase composition (irrespective of the cellobiohydrolase content, i.e., cellobiohydrolase-free, substantially cellobiohydrolase-free, or cellobiohydrolase enhanced), a surfactant, including anionic, non-ionic and ampholytic surfactants, a hydrolase, building agents, bleaching agents, bluing agents and fluorescent dyes, caking inhibitors, solubilizers, cationic surfactants and the like. All of these components are known in the detergent art. The cellulase composition as described above can be added to the detergent composition either in a liquid diluent, in granules, in emulsions, in gels, in pastes, and the like. Such forms are well known to the skilled artisan. When a solid detergent composition is employed, the cellulase composition is preferably formulated as granules. Preferably, the granules can be formulated so as to contain a cellulase protecting agent. For a more thorough discussion, see US Patent Number 6,162,782 entitled "Detergent compositions containing cellulase compositions deficient in CBH1 type components".

[163] Preferably the cellulase compositions are employed from about 0.00005 weight percent to about 5 weight percent relative to the total detergent composition. More preferably, the cellulase compositions are employed from about 0.0002 weight percent to about 2 weight percent relative to the total detergent composition.

[164] In addition the desired cellulase nucleic acid sequence finds utility in the identification and characterization of related nucleic acid sequences. A number of techniques useful for determining (predicting or confirming) the function of related genes or gene products include, but are not limited to, (A) DNA/RNA analysis, such as (1)

overexpression, ectopic expression, and expression in other species; (2) gene knock-out (reverse genetics, targeted knock-out, viral induced gene silencing (VIGS, see Baulcombe, 1999); (3) analysis of the methylation status of the gene, especially flanking regulatory regions; and (4) in situ hybridization; (B) gene product analysis such as (1) recombinant protein expression; (2) antisera production, (3) immunolocalization; (4) biochemical assays for catalytic or other activity; (5) phosphorylation status; and (6) interaction with other proteins via yeast two-hybrid analysis; (C) pathway analysis, such as placing a gene or gene product within a particular biochemical or signaling pathway based on its overexpression phenotype or by sequence homology with related genes; and (D) other analyses which may also be performed to determine or confirm the participation of the isolated gene and its product in a particular metabolic or signaling pathway, and help determine gene function.

EXAMPLES

[165] The present invention is described in further detail in the following examples which are not in any way intended to limit the scope of the invention as claimed. The attached Figures are meant to be considered as integral parts of the specification and description of the invention. All references cited are herein specifically incorporated by reference for all that is described therein.

Example 1 Identification of CBH1.1 variants

[166] This example illustrates the isolation and characterization of the nucleic acid encoding a variant *H. grisea* CBH1.1.

Isolation of Genomic DNA

[167] Genomic DNA may be isolated using any method known in the art. In this set of experiments a sample of *Humicola grisea* var *thermoidea* (CBS 225.63) was obtained. However, the following protocol may be used:

[168] Cells are grown at 45°C in 20 ml Potato Dextrose Broth (PDB) for 24 hours. The cells are diluted 1:20 in fresh PDB medium and grown overnight. Two milliliters of cells are centrifuged and the pellet washed in 1 ml KC (60g KCl, 2g citric acid per liter, pH adjusted to 6.2 with 1M KOH). The cell pellet is resuspended in 900 µl KC. 100 µl (20 mg/ml) Novozyme® is added, mixed gently and the protoplastation followed microscopically at 37°C until greater than 95% protoplasts are formed for a maximum of 2 hours. The cells are centrifuged at 1500 rpm (460g) for 10 minutes. 200 µl TES/SDS

- 38 -

(10mM Tris, 50mM EDTA, 150mM NaCl, 1% SDS) is added, mixed and incubated at room temperature for 5 minutes. DNA is isolated using a Qiagen mini-prep isolation kit (Qiagen). The column is eluted with 100 µl milli-Q water and the DNA collected.

[169] This is the method used for the isolation of genomic DNA of *H. grisea var thermoidea* from PDA plates grown at 45°C. An alternative method using the FastPrep® method may be desirable. The system consists of the FastPrep® Instrument as well as FastPrep® kits for nucleic acid isolation. FastPrep® is available from Qbiogene, Inc., Qbiogene, Inc., 2251 Rutherford Road, Carlsbad, CA 92008.

Construction of primers

[170] PCR was performed on a standard PCR machine such as the PTC-200 Peltier Thermal Cycler from MJ Research Inc. under the following conditions:

- 1) 1 minute at 96°C for 1 cycle
- 2) 30 seconds at 94°C
60 seconds at 55°C
2 minutes at 72°C
- 3) Repeat step 2 for 30 cycles
- 4) 7 minutes at 72°C for 1 cycle, and
- 5) lower temperature to 15°C for storage and further analysis.

[171] The following DNA primers were constructed for use in amplification of homologous CBH1 genes from genomic DNA's isolated from various microorganisms. All symbols used herein for protein and DNA sequences correspond to IUPAC IUB Biochemical Nomenclature Commission codes.

[172] Homologous 5' (PVS203) and 3' (PVS204) primers were developed based on the sequence of CBH1 from *Humicola grisea var thermoidea* (IFO9854) This strain expresses the sequence given in the alignment figure as D63515 (Figure 5). Both primers contained Gateway cloning sequences from Invitrogen® at the 5' of the primer. Primer PVS203 contained attB1 sequence and primer PVS204 contained attB2 sequence.

Sequence of PVS203 without the attB1:

5' ATGCGTACCGCCAAGTTCGC 3' (signal sequence of CBH1)

Sequence of PVS204 without the attB2:

5' TTACAGGCACTGAGAGTACCAG 3' (cellulose binding module of CBH1)

[173] PCR conditions were as follows: 20µL of 5X reaction buffer (5X reaction buffer comprising 50mM Tris HCl, pH 8.5; 87.5 mM (NH₄)₂SO₄; 6.25 mM MgCl₂; 2.5% Teen20 (v/v) 7.5% DMSO (v/v)); 0.2 mM each of dATP, dTTP, dGTP, dCTP (final concentration), 1 µL of 100 ng/µL genomic DNA, 1 µL of Tgo polymerase (Roche diagnostics GmbH, Cat#3186199) at 1 unit per µL, 0.2µM of each primer, PVS203 and PVS204, (final concentration), and water to 100 µL.

Isolation of variant *H. grisea* CBH1.1 gene sequence

[174] The full length sequence was obtained directly by using the N terminal (PVS203) and C terminal (PVS204) primers. The full length DNA sequence was translated into three open reading frames using Vector NTI software. Comparison of DNA and protein sequences to two known sequences for *H. grisea* Cel7A (X17258 and D63515) were performed to identify any putative intron sequences. Translation of the genomic DNA sequence without the intron sequences revealed the protein sequence of the variant *H. grisea* CBH1.1. The full length gene has been obtained and is provided in Figure 1 (genomic DNA). The putative cDNA is presented in Figure 2.

Example 2 Expression of CBH1.1 variants

[175] The following example details how the expression of the variant *H. grisea* CBH1.1 gene was performed.

[176] The full-length gene from Example 1 were transferred to the *A. niger* Gateway compatible destination vector, which was developed by Genencor. This vector was built by using the pRAX1 as a backbone, shown in Figure 6, according to the manual given in Gateway™ Cloning Technology: version 1 page 34-38.

[177] The newly developed expression vector is shown in Figure 7; this is a product of transferring the new gene into the destination vector pRAXdes2. This resulted in the final expression vectors called pRAXdesCBH1.1 (see Figure 8).

[178] The construct has been transformed into *A. niger* var. *awamori* according to the method described by Cao *et al* (Cao Q-N, Stubbs M, Ngo KQP, Ward M, Cunningham A, Pai EF, Tu G-C and Hofmann T (2000) Penicillopepsin-JT2 a recombinant enzyme from *Penicillium janthinellum* and contribution of a hydrogen bond in subsite S3 to *kcat* *Protein Science* 9:991-1001).

[179] Transformants were streaked on minimal medium plates (Ballance DJ, Buxton FP, and Turner G (1983) Transformation of *Aspergillus nidulans* by the orotidine-5'-phosphate decarboxylase gene of *Neurospora crassa* *Biochem Biophys Res Commun* 112:284-289) and grown for 4 days at 30°C. Spores were collected using methods well known in the art.

A. nidulans conidia are harvested in water

(by rubbing the surface of a conidiating culture with a sterile bent glass rod to dislodge the spores) and can be stored for weeks to months at 4°C without a serious loss of viability. However, freshly harvested spores germinate more reproducibly. For long-term storage, spores can be stored in 50% glycerol at -20°C, or in 15-20% glycerol at -80°C. Glycerol is more easily pipetted as an 80% solution in water. 800µl of aqueous conidial suspension

- 40 -

(as made for 4°C storage) added to 200 µl 80% glycerol is used for a -80°C stock; 400 µl suspension added to 600 µl 80% glycerol is used for a -20°C stock. Vortex before freezing. For mutant collections, small pieces of conidiating cultures can be excised and placed in 20% glycerol, vortexed, and frozen as -80°C stocks. In our case we store them in 50% glycerol at -80°C.

[180] *A. niger* var *awamori* transformants were grown on minimal medium lacking uridine (Ballance et al. 1983). Transformants were screened for cellulase activity by inoculating 1 cm² of spore suspension from the sporulated grown agar plate into 100 ml shake flasks for 3 days at 37°C as described by Cao et al. (2000).

[181] The CBHI activity assay is based on the hydrolysis of the nonfluorescent 4-methylumbelliferyl-β-lactoside to the products lactose and 7-hydroxy-4-methylcoumarin, the latter product is responsible for the fluorescent signal. Pipette 170 µl 50 mM NaAc buffer pH 4.5 in a 96-well microtiter plate (MTP) (Greiner, Fluotrac 200, art. nr. 655076) suitable for fluorescence. Add 10 µl of supernatant and then add 10 µl of MUL (1 mM 4-methylumbelliferyl-β-lactoside (MUL) in milliQ water) and put the MTP in the Fluostar Galaxy (BMG Labtechnologies; D-77656 Offenburg). Measure the kinetics for 16 min. (8 cycles of 120s each) using λ_{320 nm} (excitation) and λ_{460 nm} (emission) at 50°C. Supernatants having CBH1 activity were then subjected to Hydrophobic Interaction Chromatography as described in Example 3 below.

[182] The amino acid sequences were deduced as stated above in Example 1. The amino acid sequence of the variant CBH1.1 is shown in Figure 3 with a signal sequence, and in Figure 4 without a signal sequence. The signal sequence is shown in Figure 3 in bold and underlined font.

Example 3 Thermostability of CBH1 variants

[183] The following example details how the variant *H. grisea* CBH1.1 differs in thermostability from a *T. reesei* CBH1 cellulase enzyme.

[184] CBH I cellulase variants were cloned and expressed as above (see Example 2). Cel7A wild type and variants were then purified from cell-free supernatants of these cultures by column chromatography. Proteins were purified using hydrophobic interaction chromatography (HIC). Columns were run on a BioCAD® Sprint Perfusion Chromatography System using Poros® 20 HP2 resin both made by Applied Biosystems.

[185] HIC columns were equilibrated with 5 column volumes of 0.020 M sodium phosphate, 0.5 M ammonium sulfate at pH 6.8. Ammonium sulfate was added to the supernatants to a final concentration of approximately 0.5 M and the pH was adjusted to

- 41 -

6.8. After filtration, the supernatant was loaded onto the column. After loading, the column was washed with 10 column volumes of equilibration buffer and then eluted with a 10 column volume gradient from 0.5 M ammonium sulfate to zero ammonium sulfate in 0.02 M sodium phosphate pH 6.8. Cel7A was eluted approximately mid-gradient. Fractions were collected and pooled on the basis of reduced, SDS-PAGE gel analysis.

[186] The melting points were determined according to the methods of Luo, *et al.*, *Biochemistry* 34:10669 and Gloss, *et al.*, *Biochemistry* 36:5612.

[187] Data was collected on the Aviv 215 circular dichroism spectrophotometer. Spectra of the variants between 210 and 260 nanometers were taken at 25° C. Buffer conditions were 50 mM Bis Tris Propane/50 mM ammonium acetate/glacial acetic acid at pH 5.5. The protein concentration was kept between 0.25 and 0.5 mgs/mL. After determining the optimal wavelength to monitor unfolding, the samples were thermally denatured by ramping the temperature from 25° C to 75° C under the same buffer conditions. Data was collected for 5 seconds every 2 degrees. Partially reversible unfolding was monitored at 230 nanometers in an 0.1 centimeter path length cell.

[188] The variant *H. grisea* CBH 1.1 cellulase has an enhanced thermostability profile as compared wild type *T. reesei* CBH1. The variant *H. grisea* CBH1.1 has a T_m of 72.5°C; *T. reesei* has a T_m of 62.3°C.

Example 4 **Activity of CBH1 variants**

[189] The following example details how the activity of the *H. grisea* CBH1.1 variant was evaluated.

[190] Cellulose conversion was evaluated by techniques known in the art. See, for example, Baker et al, Appl Biochem Biotechnol 1998 Spring; 70-72():395-403.

[191] A standard cellulosic conversion assay was used in the experiments. In this assay enzyme and buffered substrate were placed in containers and incubated at a temperature over time. The reaction was quenched with enough 100 mM Glycine, pH 11.0 to bring the pH of the reaction mixture to at least pH10. Once the reaction was quenched, an aliquot of the reaction mixture was filtered through a 0.2 micron membrane to remove solids. The filtered solution was then assayed for soluble sugars by HPLC according to the methods described in Baker et al., Appl. Biochem. Biotechnol. 70-72:395 – 403 (1998).

[192] Pretreated corn stover (PCS) – Corn stover was pretreated with 2% w/w H₂SO₄ as described in Schell, D. et al., J. Appl. Biochem. Biotechnol. 105:69 – 86 (2003) and followed by multiple washes with deionized water to obtain a pH of 4.5. Sodium acetate

was added to make a final concentration of 50mM and this was titrated to pH 5.0. The cellulose concentration in the reaction mixture was approximately 7%.

[193] Phosphoric acid swollen cellulose (PASC) - PASC was prepared from Avicel according to the method described in Walseth (1971) Tappi 35: 228 (1971) and Wood Biochem J. 121:353 (1971). This material was diluted with buffer and water to achieve a 0.63 % w/v mixture such that the final concentration of sodium acetate was 50mM, pH 5.0. The enzymes were dosed at 1.6 mg of total protein per gram of cellulose.

[194] The first set of experiments examined the percent conversion at 38°C for 1 day on 12.66% pretreated corn stover (PCS) (see Schell, D. et al., J. Appl. Biochem. Biotechnol. (2003) 105:69 – 86) using 15.5 mg enzyme/gm cellulose. Samples were agitated at 700rpm. Comparisons were made between: 1) a cellulase from a CBH1 deleted *Trichoderma* strain (delCBH1); 2) an *A. niger* strain that had a *T. reesei* CBH1 gene inserted (rCBH1); 3) an *A. niger* strain overexpressing its native CBHB (Aniger); 4) an *A. niger* strain that had a *H. schweinitzii* CBH1 gene inserted (Hschweinitzii/An); 5) an *A. niger* strain that had a *T. pseudokoningii* CBH1 gene inserted (Tpseudokoni/An); 6) an *A. niger* strain that had a *H. grisea* variant CBH1.1 gene inserted (Hgrisea/An-1); 7) an *A. niger* strain that had a *H. grisea* variant CBH1.1 gene inserted (Hgrisea/An-2).

Hgrisea/An-1 and Hgrisea/An-2 are two clones from the same transformation of *A. niger* with the *H. grisea* variant CBH1.1 gene. The results from the first set of experiments are presented in Figure 9. As can be seen, the *H. grisea* variant CBH1.1 does not out perform any of the other CBH's tested. Figure 11 is a comparison of the *H. grisea* variant CBH1.1 against the *T. reesei* nCBH1 results from the same experimental conditions.

[195] The second set of experiments examined the percent conversion under similar conditions as the first set except the incubation temperature was 65°C, not 38°C. Results for this set of experiments are presented in Figure 10. As can be seen, the *H. grisea* variant CBH1.1 out performed the other CBH's tested. Figure 12 is a comparison of the *H. grisea* variant CBH1.1 against the *T. reesei* rCBH1 results from the same experimental conditions.

[196] The third set of experiments examined the rate of cellobiose generation from PASC under similar test conditions as the previous experiments; the temperatures used were 38°C, 65°C and 70°C. Results are presented in Figure 13. As can be seen, the *H. grisea* variant CBH1.1 out performed the *T. reesei* CBH1 at all temperatures tested.

Example 5 Isolation of *Scytalidium thermophilum* CBH1

[197] This example illustrates the isolation and characterization of the nucleic acid encoding a *S. thermophilum* CBH1.

Isolation of Genomic DNA and Construction of Primers

[198] Methods for cloning, expressing and purifying the *Scytalidium thermophilum* CBH1 were all as described for *Humicola grisea* var. *thermoidea* CBH1.1 (see Example 1). The CBH1 gene from *Scytalidium thermophilum* was amplified from genomic DNA from public strain collection entry CBS 671.88, using the same PCR primers, PVS203 and PVS204, used to amplify the *Humicola grisea* var. *thermoidea* CBH1.1.

Isolation of *S. thermophilum* CBH1 gene sequence

[199] The full length sequence was obtained directly by using the N terminal (PVS203) and C terminal (PVS204) primers. The full length DNA sequence was translated into three open reading frames using Vector NTI software. Translation of the genomic DNA sequence without the intron sequences revealed the protein sequence of the *S. thermophilum* CBH1. The full length gene has been obtained and is provided in Figure 14A (genomic DNA). The putative cDNA is presented in Figure 14B. The amino acid sequence with and without signal sequence is provided in Figures 14C and 15, respectively.

[200] The *S. thermophilum* CBH1, *Humicola grisea* var. *thermoidea* CBH1.1 and *H. jecorina* CBH1 protein sequences were aligned. See Figure 15.

[201] By identifying sites of difference between *Humicola grisea* var. *thermoidea* CBH1.1 and the significantly more stable *Scytalidium thermophilum* CBH1, and then using the alignment in Figure 15 to identify the corresponding site in *Hypocrea jecorina* CBH1, which is much less stable than either *Humicola grisea* var. *thermoidea* CBH1.1 or *Scytalidium thermophilum* CBH1.) The following sites in *Hypocrea jecorina* CBH1 (mature protein residue numbering) are identified as being important for stability:

Thr(T)55, preferably Thr55Glu (T55E) & Thr55Lys (T55K)

Ser(S)58, preferably Ser58Thr (S58T)

Gln(Q)101, preferably Gln101Tyr (Q101Y) & Gln101His (Q101H)

Asn(N)250, preferably Asn250Asp (N250D) & Asn250Glu (N250E)

Pro(P)265, preferably Pro265Ala (P265A) & Pro265Ser (P265S)

Leu(L)288, preferably Leu288Ile (L288I).

Example 6
Thermostability of *Scytalidium thermophilum* CBH1

[202] This example describes the thermostability measurement of *S. thermophilum* CBH1 and *H. grisea* var. *thermoidea* CBH1.1.

[203] *S. thermophilum* CBH1 and *H. grisea* var. *thermoidea* CBH1.1 were expressed using the methods as in Example 2, purified as in Example 3 and analyzed by differential scanning calorimetry using a Microcal VP-DSC. Samples were scanned from 30 to 95 degrees C at 90 degrees C/hour. The purified proteins were desalted into 50 mM ammonium acetate and 50 mM bistris propane at pH 5.5. The final protein concentration was between 0.05 and 0.25 mg/mL for all samples.

[204] The thermal transition observed for *S. thermophilum* CBH1 had a midpoint of 78.3 degrees C. *H. grisea* CBH1.1 has a transition with a midpoint of 76.0 degrees C. See Figure 16.

[205] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

SEQUENCE LISTING

<110> Genencor International, Inc.

<120> Variant Humicola grisea CBH1.1

<130> 11816-106

<140> CA 2,520,636

<141> 2004-03-26

<150> US 60/459,734

<151> 2003-04-01

<160> 15

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1638

<212> DNA

<213> Humicola grisea

<400> 1

atgcgtaccg	ccaagtctgc	caccctcgcc	gcccttgtgg	cctcggccgc	cgcccagcag	60
gcgtgcagtc	tcaccaccga	gaggcaccct	tccctctctt	ggaagaagtg	caccgcccgc	120
ggccagtgcc	agaccgtcca	ggcttccatc	actctcgact	ccaactggcg	ctggactcac	180
caggtgtctg	gctccaccaa	ctgctacacg	ggcaacaagt	gggatactag	catctgcact	240
gatgccaaagt	cgtgcgctca	gaactgctgc	gtcgatgggtg	ccgactacac	cagcacctat	300
ggcatcacca	ccaacgggtga	ttccctgagc	ctcaagttcg	tcaccaaggg	ccagcactcg	360
accaacgtcg	gctcgcgtac	ctacctgatg	gacggcgagg	acaagtatca	gagtacgttc	420
tatcttcagc	cttctcgcgc	cttgaatcct	ggctaacgtt	tacacttcac	agccttcgag	480
ctcctcggca	acgagttcac	cttcgatgtc	gatgtctcca	acatcggctg	cggctctcaac	540
ggcgccctgt	acttcgtctc	catggacgcc	gatgggtggc	tcagccgcta	tcctggcaac	600
aaggctgggtg	ccaagtacgg	taccggctac	tgcgatgtct	agtgcccccg	tgacatcaag	660
ttcatcaacg	gcgaggccaa	cattgagggc	tggaccggct	ccaccaacga	ccccaacgcc	720
ggcgcgggcc	gctatggtac	ctgctgctct	gagatggata	tctgggaagc	caacaacatg	780
gctactgcct	tcactcctca	cccttgcacc	atcattggcc	agagccgctg	cgagggcgac	840
tcgtgcgggtg	gcacctacag	caacgagcgc	tacgccggcg	tctgcgaccc	cgatggctgc	900
gacttcaact	cgtaccgcca	gggcaacaag	accttctacg	gcaagggcat	gaccgtcgac	960
accaccaaga	agatcactgt	cgtcaccacg	ttcctcaagg	atgccaacgg	cgatctcggc	1020
gagatcaagc	gcttctacgt	ccaggatggc	aagatcatcc	ccaactccga	gtccaccatc	1080
cccggcgctc	agggcaattc	catcaccacg	gactgggtgc	accgccagaa	ggttgccttt	1140
ggcgacattg	acgacttcaa	ccgcaagggc	ggcatgaagc	agatgggcaa	ggccctcgcc	1200
ggccccatgg	tcctgggtcat	gtccatctgg	gatgaccacg	cctccaacat	gctctggctc	1260
gactcgacct	tccctgtcga	tgccgctggc	aagcccggcg	ccgagcgcg	tgcttgccc	1320
accacctcgg	gtgtccctgc	tgaggttgag	gccgaggccc	ccaacagcaa	cgctgtcttc	1380
tccaacatcc	gcttcggccc	catcggtcgc	accgttgctg	gtctccccgg	cgcgggcaac	1440
ggcggaaca	acggcggcaa	ccccccgccc	cccaccacca	ccacctctc	ggctccggcc	1500
accaccacca	ccgccagcgc	tgcccccaag	gctggccgct	ggcagcagtg	cgcgggcatc	1560
ggcttctactg	gcccgaacca	gtgcgaggag	ccctacactt	gcaccaagct	caacgactgg	1620
tactctcagt	gcctgtaa					1638

<210> 2

<211> 1323

<212> DNA

<213> Humicola grisea

<400> 2

```

gctcagaact gctgcgtcga tgggtgccgac tacaccagca cctatggcat caccaccaac      60
ggtgattccc tgagcctcaa gttcgtcacc aagggccagc actcgaccaa cgtcgggctcg      120
cgtacctacc tgatggacgg cgaggacaag tatcagacct tcgagctcct cggcaacgag      180
ttcaccttcg atgtcgaatg ctccaacatc ggctgcggtc tcaacggcgc cctgtacttc      240
gtctccatgg acgccgatgg tgggtctcagc cgctatcctg gcaacaaggc tggtgccaag      300
tacgggtaccg gctactgcga tgctcagtgc ccccgtgaca tcaagtccat caacggcgag      360
gccaacattg agggctggac cggctccacc aacgacccca acgccggcgc gggccgctat      420
ggtacctgct gctctgagat ggatatctgg gaagccaaca acatgggtac tgccttcact      480
cctcaccctt gcaccatcat tggccagagc cgctgcgagg gcgactcgtg cgggtggcacc      540
tacagcaacg agcgctacgc cggcgctctg gaccccgatg gctgcgactt caactcgtac      600
cgccagggca acaagacctt ctacggcaag ggcattgacc tcgacaccac caagaagatc      660
actgtcgtca cccagttcct caaggatgcc aacggcgatc tcggcgagat caagcgcttc      720
tacgtccagg atggcaagat catccccaac tccgagtcca ccatccccgg cgtcgagggc      780
aattccatca cccaggactg gtgcgaccgc cagaagggtg cctttggcga cattgacgac      840
ttcaaccgca agggcgggcat gaagcagatg ggcaaggccc tcgccggccc catggtcctg      900
gtcatgtcca tctgggatga ccacgcctcc aacatgctct ggctcgactc gaccttcct      960
gtcgatgccg ctggcaagcc cggcgccgag cgcggtgcct gcccgaccac ctcgggtgtc     1020
cctgctgagg ttgaggccga ggcccccaac agcaacgtcg tcttctccaa catccgcttc     1080
ggccccatcg gctcgaccgt tgctgggtctc cccggcgcgg gcaacggcgg caacaacggc     1140
ggcaaccccc cgccccccac caccaccacc tcctcggtc cggccaccac caccaccgcc     1200
agcgctggcc ccaaggctgg ccgctggcag cagtgcggcg gcatcggtt cactggcccg     1260
acccagtgcg aggagcccta cacttgacc aagctcaacg actggtactc tcagtgcctg     1320
taa                                                                1323

```

<210> 3

<211> 525

<212> PRT

<213> Humicola grisea

<400> 3

```

Met Arg Thr Ala Lys Phe Ala Thr Leu Ala Ala Leu Val Ala Ser Ala
 1          5          10          15
Ala Ala Gln Gln Ala Cys Ser Leu Thr Thr Glu Arg His Pro Ser Leu
          20          25          30
Ser Trp Lys Lys Cys Thr Ala Gly Gly Gln Cys Gln Thr Val Gln Ala
          35          40          45
Ser Ile Thr Leu Asp Ser Asn Trp Arg Trp Thr His Gln Val Ser Gly
          50          55          60
Ser Thr Asn Cys Tyr Thr Gly Asn Lys Trp Asp Thr Ser Ile Cys Thr
          65          70          75          80
Asp Ala Lys Ser Cys Ala Gln Asn Cys Cys Val Asp Gly Ala Asp Tyr
          85          90          95
Thr Ser Thr Tyr Gly Ile Thr Thr Asn Gly Asp Ser Leu Ser Leu Lys
          100          105          110
Phe Val Thr Lys Gly Gln His Ser Thr Asn Val Gly Ser Arg Thr Tyr
          115          120          125
Leu Met Asp Gly Glu Asp Lys Tyr Gln Thr Phe Glu Leu Leu Gly Asn
          130          135          140
Glu Phe Thr Phe Asp Val Asp Val Ser Asn Ile Gly Cys Gly Leu Asn
          145          150          155          160
Gly Ala Leu Tyr Phe Val Ser Met Asp Ala Asp Gly Gly Leu Ser Arg
          165          170          175
Tyr Pro Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp
          180          185          190
Ala Gln Cys Pro Arg Asp Ile Lys Phe Ile Asn Gly Glu Ala Asn Ile
          195          200          205
Glu Gly Trp Thr Gly Ser Thr Asn Asp Pro Asn Ala Gly Ala Gly Arg

```

210	215	220
Tyr Gly Thr Cys Cys Ser Glu Met Asp Ile Trp Glu Ala Asn Asn Met		
225	230	235
Ala Thr Ala Phe Thr Pro His Pro Cys Thr Ile Ile Gly Gln Ser Arg		240
	245	250
Cys Glu Gly Asp Ser Cys Gly Gly Thr Tyr Ser Asn Glu Arg Tyr Ala		255
	260	265
Gly Val Cys Asp Pro Asp Gly Cys Asp Phe Asn Ser Tyr Arg Gln Gly		270
	275	280
Asn Lys Thr Phe Tyr Gly Lys Gly Met Thr Val Asp Thr Thr Lys Lys		285
	290	295
Ile Thr Val Val Thr Gln Phe Leu Lys Asp Ala Asn Gly Asp Leu Gly		300
305	310	315
Glu Ile Lys Arg Phe Tyr Val Gln Asp Gly Lys Ile Ile Pro Asn Ser		320
	325	330
Glu Ser Thr Ile Pro Gly Val Glu Gly Asn Ser Ile Thr Gln Asp Trp		335
	340	345
Cys Asp Arg Gln Lys Val Ala Phe Gly Asp Ile Asp Asp Phe Asn Arg		350
	355	360
Lys Gly Gly Met Lys Gln Met Gly Lys Ala Leu Ala Gly Pro Met Val		365
370	375	380
Leu Val Met Ser Ile Trp Asp Asp His Ala Ser Asn Met Leu Trp Leu		385
385	390	395
Asp Ser Thr Phe Pro Val Asp Ala Ala Gly Lys Pro Gly Ala Glu Arg		400
	405	410
Gly Ala Cys Pro Thr Thr Ser Gly Val Pro Ala Glu Val Glu Ala Glu		415
	420	425
Ala Pro Asn Ser Asn Val Val Phe Ser Asn Ile Arg Phe Gly Pro Ile		430
	435	440
Gly Ser Thr Val Ala Gly Leu Pro Gly Ala Gly Asn Gly Gly Asn Asn		445
450	455	460
Gly Gly Asn Pro Pro Pro Pro Thr Thr Thr Thr Ser Ser Ala Pro Ala		465
465	470	475
Thr Thr Thr Thr Ala Ser Ala Gly Pro Lys Ala Gly Arg Trp Gln Gln		480
	485	490
Cys Gly Gly Ile Gly Phe Thr Gly Pro Thr Gln Cys Glu Glu Pro Tyr		495
	500	505
Thr Cys Thr Lys Leu Asn Asp Trp Tyr Ser Gln Cys Leu		510
	515	520
		525

<210> 4
 <211> 507
 <212> PRT
 <213> Humicola grisea

<400> 4
 Gln Gln Ala Cys Ser Leu Thr Thr Glu Arg His Pro Ser Leu Ser Trp
 1 5 10 15
 Lys Lys Cys Thr Ala Gly Gly Gln Cys Gln Thr Val Gln Ala Ser Ile
 20 25 30
 Thr Leu Asp Ser Asn Trp Arg Trp Thr His Gln Val Ser Gly Ser Thr
 35 40 45
 Asn Cys Tyr Thr Gly Asn Lys Trp Asp Thr Ser Ile Cys Thr Asp Ala
 50 55 60
 Lys Ser Cys Ala Gln Asn Cys Cys Val Asp Gly Ala Asp Tyr Thr Ser
 65 70 75 80
 Thr Tyr Gly Ile Thr Thr Asn Gly Asp Ser Leu Ser Leu Lys Phe Val
 85 90 95
 Thr Lys Gly Gln His Ser Thr Asn Val Gly Ser Arg Thr Tyr Leu Met

			100					105					110				
Asp	Gly	Glu	Asp	Lys	Tyr	Gln	Thr	Phe	Glu	Leu	Leu	Gly	Asn	Glu	Phe		
		115					120					125					
Thr	Phe	Asp	Val	Asp	Val	Ser	Asn	Ile	Gly	Cys	Gly	Leu	Asn	Gly	Ala		
	130					135					140						
Leu	Tyr	Phe	Val	Ser	Met	Asp	Ala	Asp	Gly	Gly	Leu	Ser	Arg	Tyr	Pro		
145					150					155					160		
Gly	Asn	Lys	Ala	Gly	Ala	Lys	Tyr	Gly	Thr	Gly	Tyr	Cys	Asp	Ala	Gln		
			165					170						175			
Cys	Pro	Arg	Asp	Ile	Lys	Phe	Ile	Asn	Gly	Glu	Ala	Asn	Ile	Glu	Gly		
			180					185					190				
Trp	Thr	Gly	Ser	Thr	Asn	Asp	Pro	Asn	Ala	Gly	Ala	Gly	Arg	Tyr	Gly		
	195					200						205					
Thr	Cys	Cys	Ser	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Asn	Met	Ala	Thr		
	210					215					220						
Ala	Phe	Thr	Pro	His	Pro	Cys	Thr	Ile	Ile	Gly	Gln	Ser	Arg	Cys	Glu		
225					230					235					240		
Gly	Asp	Ser	Cys	Gly	Gly	Thr	Tyr	Ser	Asn	Glu	Arg	Tyr	Ala	Gly	Val		
			245					250						255			
Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Ser	Tyr	Arg	Gln	Gly	Asn	Lys		
			260					265					270				
Thr	Phe	Tyr	Gly	Lys	Gly	Met	Thr	Val	Asp	Thr	Thr	Lys	Lys	Ile	Thr		
	275					280						285					
Val	Val	Thr	Gln	Phe	Leu	Lys	Asp	Ala	Asn	Gly	Asp	Leu	Gly	Glu	Ile		
	290					295					300						
Lys	Arg	Phe	Tyr	Val	Gln	Asp	Gly	Lys	Ile	Ile	Pro	Asn	Ser	Glu	Ser		
305					310					315					320		
Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	Asp		
			325					330					335				
Arg	Gln	Lys	Val	Ala	Phe	Gly	Asp	Ile	Asp	Asp	Phe	Asn	Arg	Lys	Gly		
			340					345					350				
Gly	Met	Lys	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	Val		
	355					360					365						
Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Ser	Asn	Met	Leu	Trp	Leu	Asp	Ser		
	370					375					380						
Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala		
385					390					395					400		
Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro		
			405					410						415			
Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser		
			420					425					430				
Thr	Val	Ala	Gly	Leu	Pro	Gly	Ala	Gly	Asn	Gly	Gly	Asn	Asn	Gly	Gly		
	435					440						445					
Asn	Pro	Pro	Pro	Pro	Thr	Thr	Thr	Thr	Ser	Ser	Ala	Pro	Ala	Thr	Thr		
	450					455					460						
Thr	Thr	Ala	Ser	Ala	Gly	Pro	Lys	Ala	Gly	Arg	Trp	Gln	Gln	Cys	Gly		
465					470					475					480		
Gly	Ile	Gly	Phe	Thr	Gly	Pro	Thr	Gln	Cys	Glu	Glu	Pro	Tyr	Thr	Cys		
			485					490					495				
Thr	Lys	Leu	Asn	Asp	Trp	Tyr	Ser	Gln	Cys	Leu							
			500					505									

<210> 5

<211> 507

<212> PRT

<213> Humicola grisea

<400> 5

Gln Gln Ala Cys Ser Leu Thr Thr Glu Arg His Pro Ser Leu Ser Trp

1				5				10				15				
Asn	Lys	Cys	Thr	Ala	Gly	Gly	Gln	Cys	Gln	Thr	Val	Gln	Ala	Ser	Ile	
			20					25					30			
Thr	Leu	Asp	Ser	Asn	Trp	Arg	Trp	Thr	His	Gln	Val	Ser	Gly	Ser	Thr	
		35					40					45				
Asn	Cys	Tyr	Thr	Gly	Asn	Lys	Trp	Asp	Thr	Ser	Ile	Cys	Thr	Asp	Ala	
	50					55					60					
Lys	Ser	Cys	Ala	Gln	Asn	Cys	Cys	Val	Asp	Gly	Ala	Asp	Tyr	Thr	Ser	
65					70					75					80	
Thr	Tyr	Gly	Ile	Thr	Thr	Asn	Gly	Asp	Ser	Leu	Ser	Leu	Lys	Phe	Val	
				85					90					95		
Thr	Lys	Gly	Gln	His	Ser	Thr	Asn	Val	Gly	Ser	Arg	Thr	Tyr	Leu	Met	
			100					105					110			
Asp	Gly	Glu	Asp	Lys	Tyr	Gln	Thr	Phe	Glu	Leu	Leu	Gly	Asn	Glu	Phe	
		115					120					125				
Thr	Phe	Asp	Val	Asp	Val	Ser	Asn	Ile	Gly	Cys	Gly	Leu	Asn	Gly	Ala	
	130					135					140					
Leu	Tyr	Phe	Val	Ser	Met	Asp	Ala	Asp	Gly	Gly	Leu	Ser	Arg	Tyr	Pro	
145					150					155					160	
Gly	Asn	Lys	Ala	Gly	Ala	Lys	Tyr	Gly	Thr	Gly	Tyr	Cys	Asp	Ala	Gln	
				165				170						175		
Cys	Pro	Arg	Asp	Ile	Lys	Phe	Ile	Asn	Gly	Glu	Ala	Asn	Ile	Glu	Gly	
			180					185					190			
Trp	Thr	Gly	Ser	Thr	Asn	Asp	Pro	Asn	Ala	Gly	Ala	Gly	Arg	Tyr	Gly	
		195				200						205				
Thr	Cys	Cys	Ser	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Asn	Met	Ala	Thr	
	210					215					220					
Ala	Phe	Thr	Pro	His	Pro	Cys	Thr	Ile	Ile	Gly	Gln	Ser	Arg	Cys	Glu	
225					230					235					240	
Gly	Asp	Ser	Cys	Gly	Gly	Thr	Tyr	Ser	Asn	Glu	Arg	Tyr	Ala	Gly	Val	
				245					250					255		
Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Ser	Tyr	Arg	Gln	Gly	Asn	Lys	
			260					265					270			
Thr	Phe	Tyr	Gly	Lys	Gly	Met	Thr	Val	Asp	Thr	Thr	Lys	Lys	Ile	Thr	
		275					280					285				
Val	Val	Thr	Gln	Phe	Leu	Lys	Asp	Ala	Asn	Gly	Asp	Leu	Gly	Glu	Ile	
	290					295					300					
Lys	Arg	Phe	Tyr	Val	Gln	Asp	Gly	Lys	Ile	Ile	Pro	Asn	Ser	Glu	Ser	
305					310					315					320	
Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	Asp	
				325					330					335		
Arg	Gln	Lys	Val	Ala	Phe	Gly	Asp	Ile	Asp	Asp	Phe	Asn	Arg	Lys	Gly	
			340					345					350			
Gly	Met	Lys	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	Val	
		355					360					365				
Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Ser	Asn	Met	Leu	Trp	Leu	Asp	Ser	
	370					375					380					
Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala	
385					390					395					400	
Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro	
				405					410					415		
Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser	
			420					425					430			
Thr	Val	Ala	Gly	Leu	Pro	Gly	Ala	Gly	Asn	Gly	Gly	Asn	Asn	Gly	Gly	
		435					440					445				
Asn	Pro	Pro	Pro	Pro	Thr	Thr	Thr	Thr	Ser	Ser	Ala	Pro	Ala	Thr	Thr	
	450					455					460					
Thr	Thr	Ala	Ser	Ala	Gly	Pro	Lys	Ala	Gly	Arg	Trp	Gln	Gln	Cys	Gly	
465					470					475					480	

Gly Ile Gly Phe Thr Gly Pro Thr Gln Cys Glu Glu Pro Tyr Ile Cys
 485 490 495
 Thr Lys Leu Asn Asp Trp Tyr Ser Gln Cys Leu
 500 505

<210> 6

<211> 507

<212> PRT

<213> Humicola grisea

<400> 6

Gln Gln Ala Cys Ser Leu Thr Thr Glu Arg His Pro Ser Leu Ser Trp
 1 5 10 15
 Lys Lys Cys Thr Ala Gly Gly Gln Cys Gln Thr Val Gln Ala Ser Ile
 20 25 30
 Thr Leu Asp Ser Asn Trp Arg Trp Thr His Gln Val Ser Gly Ser Thr
 35 40 45
 Asn Cys Tyr Thr Gly Asn Lys Trp Asp Thr Ser Ile Cys Thr Asp Ala
 50 55 60
 Lys Ser Cys Ala Gln Asn Cys Cys Val Asp Gly Ala Asp Tyr Thr Ser
 65 70 75 80
 Thr Tyr Gly Ile Thr Thr Asn Gly Asp Ser Leu Ser Leu Lys Phe Val
 85 90 95
 Thr Lys Gly Gln Tyr Ser Thr Asn Val Gly Ser Arg Thr Tyr Leu Met
 100 105 110
 Asp Gly Glu Asp Lys Tyr Gln Thr Phe Glu Leu Leu Gly Asn Glu Phe
 115 120 125
 Thr Phe Asp Val Asp Val Ser Asn Ile Gly Cys Gly Leu Asn Gly Ala
 130 135 140
 Leu Tyr Phe Val Ser Met Asp Ala Asp Gly Gly Leu Ser Arg Tyr Pro
 145 150 155 160
 Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp Ala Gln
 165 170 175
 Cys Pro Arg Asp Ile Lys Phe Ile Asn Gly Glu Ala Asn Ile Glu Gly
 180 185 190
 Trp Thr Gly Ser Thr Asn Asp Pro Asn Ala Gly Ala Gly Arg Tyr Gly
 195 200 205
 Thr Cys Cys Ser Glu Met Asp Ile Trp Glu Ala Asn Asn Met Ala Thr
 210 215 220
 Ala Phe Thr Pro His Pro Cys Thr Ile Ile Gly Gln Ser Arg Cys Glu
 225 230 235 240
 Gly Asp Ser Cys Gly Gly Thr Tyr Ser Asn Glu Arg Tyr Ala Gly Val
 245 250 255
 Cys Asp Pro Asp Gly Cys Asp Phe Asn Ser Tyr Arg Gln Gly Asn Lys
 260 265 270
 Thr Phe Tyr Gly Lys Gly Met Thr Val Asp Thr Thr Lys Lys Ile Thr
 275 280 285
 Val Val Thr Gln Phe Leu Lys Asp Ala Asn Gly Asp Leu Gly Glu Ile
 290 295 300
 Lys Arg Phe Tyr Val Gln Asp Gly Lys Ile Ile Pro Asn Ser Glu Ser
 305 310 315 320
 Thr Ile Pro Gly Val Glu Gly Asn Ser Ile Thr Gln Asp Trp Cys Asp
 325 330 335
 Arg Gln Lys Val Ala Phe Gly Asp Ile Asp Asp Phe Asn Arg Lys Gly
 340 345 350
 Gly Met Lys Gln Met Gly Lys Ala Leu Ala Gly Pro Met Val Leu Val
 355 360 365
 Met Ser Ile Trp Asp Asp His Ala Ser Asn Met Leu Trp Leu Asp Ser
 370 375 380

Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala	
385					390					395					400	
Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro	
				405					410					415		
Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser	
			420					425					430			
Thr	Val	Ala	Gly	Leu	Pro	Gly	Ala	Gly	Asn	Gly	Gly	Asn	Asn	Gly	Gly	
		435					440					445				
Asn	Pro	Pro	Pro	Pro	Thr	Thr	Thr	Thr	Ser	Ser	Ala	Pro	Ala	Thr	Thr	
	450					455					460					
Thr	Thr	Ala	Ser	Ala	Gly	Pro	Lys	Ala	Gly	Arg	Trp	Gln	Gln	Cys	Gly	
465					470					475					480	
Gly	Ile	Gly	Phe	Thr	Gly	Pro	Thr	Gln	Cys	Glu	Glu	Pro	Tyr	Thr	Cys	
				485					490						495	
Thr	Lys	Leu	Asn	Asp	Trp	Tyr	Ser	Gln	Cys	Leu						
			500					505								

<210> 7

<211> 1662

<212> DNA

<213> Scytalidium thermophilum

<400> 7

atgcgtaccg	ccaagttcgc	caccctcgcc	gcccttgtgg	cctcggccgc	cgcccagcag	60
gcgtgcagcc	tcaccaccga	gaggcaccct	tccctctcct	ggaagaagtg	caccgcccgc	120
ggccagtgcc	agaccgtcca	ggcttccatc	actctcgact	ccaactggcg	ctggactcac	180
caggtgtctg	gctccaccaa	ctgctacacg	ggcaacgagt	gggattctag	catctgcact	240
gatgccaagt	cgtgcgctca	gaactgctgc	gtcgatgggtg	ctgactacac	cagcacctat	300
ggcatcacca	ccaacggtga	ttccctgagc	ctcaagttcg	tcaccaaggg	ccagtactcg	360
accaacgtcg	gctcgcgtac	ctacctgatg	gacggcgagg	acaagtatca	gagtaggttc	420
tatcttcagc	cttctcgcgc	cttgaatcct	ggctaacttt	tacacttcac	agccttcgag	480
ctcctcggca	acgagttcac	cttcgatgtc	gatgtctcca	acatcggctg	cggtctcaac	540
ggcgccctgt	acttcgtctc	catggacgcc	gatggtggtc	tcagccgcta	tcctggcaac	600
aaggctggtg	ccaagtacgg	taccggctac	tgcgatgctc	agtgcctccg	tgacatcaag	660
ttcatcaacg	gcgaggccaa	cattgagggc	tggaccggct	ccaccaacga	ccccaacgcc	720
ggcgcgggcc	gctatggtac	ctgctgctct	gagatggata	tctgggaggc	caacaacatg	780
gctactgcct	tcactcctca	cccttgcact	atcattggcc	agagccgctg	cgagggcgac	840
tcgtgcggtg	gcacctacag	caacgaccgc	tacgccggcg	tctgcgacce	cgatggctgc	900
gacttcaacg	cgtatcgcca	gggcaacaag	accttctacg	gcaagggcac	gaccgtcgac	960
accaccaaga	agctcaccgt	cgtcacccag	ttcctcaagg	acgccaacgg	cgatctcggc	1020
gagatcaagc	gcttctacgt	ccaggatggg	aagatcatcc	ccaactccga	gtccaccatc	1080
cccggcgctc	agggcaactc	catcacccag	gattggtgcg	accgccagaa	ggttgccttt	1140
ggcgacattg	acgacttcaa	ccgcaagggc	ggcatgaagc	agatgggcaa	ggccctcgcc	1200
ggccccatgg	tcctgggtcat	gtccatctgg	gatgaccacg	cctccaacat	gctctggctc	1260
gactcgacct	tccctgtcga	tgccgctggc	aagcccggcg	ccgagcgcg	tgccctgccc	1320
accacctcgg	gtgtccctgc	tgaggttgag	gccgaggccc	ccaacagcaa	cgctcgtctc	1380
tccaacatcc	gcttcggccc	catcggtcgc	accgttgccg	gccttcccag	cgatggcggc	1440
aacaacggcg	gcaacaccac	cgtccagccc	ccgcccagca	ccaccaccac	ctctgccagc	1500
agcagcacca	cctcggctcc	tgccaccacc	accaccgcca	gcgctggccc	caaggctggc	1560
cgctggcagc	agtgcggcgg	catcggtctc	actggcccga	cccagtgcga	ggagccctac	1620
acttgcacca	agctcaacga	ctggtactct	cagtgcctgt	aa		1662

<210> 8

<211> 1602

<212> DNA

<213> Scytalidium thermophilum

<400> 8

atgcgtaccg	ccaagttcgc	caccctcgcc	gcccttgtgg	cctcggccgc	cgcccagcag	60
------------	------------	------------	------------	------------	------------	----

```

gcgtgcagcc tcaccaccga gaggcaccct tccctctcct ggaagaagtg caccgccggc 120
ggccagtgcc agaccgtcca ggcttccatc actctcgact ccaactggcg ctggactcac 180
caggtgtctg gctccaccaa ctgctacacg ggcaacgagt gggattctag catctgcact 240
gatgccaaagt cgtgcgctca gaactgctgc gtcgatgggtg ctgactacac cagcacctat 300
ggcatcacca ccaacgggtga ttccctgagc ctcaagtctg tcaccaaggg ccagtactcg 360
accaacgtcg gctcgcgtac ctacctgatg gacggcgagg acaagtatca gaccttcgag 420
ctcctcggca acgagttcac cttcgatgtc gatgtctcca acatcggctg cggctctaac 480
ggcgccctgt acttcgtctc catggacgcc gatgggtggc tcagccgcta tcctggcaac 540
aaggctgggtg ccaagtacgg taccggctac tgcgatgtc agtgcccccg tgacatcaag 600
ttcatcaacg gcgaggccaa cattgagggc tggaccggct ccaccaacga ccccaacgcc 660
ggcgcgggcc gctatggtac ctgctgctct gagatggata tctgggagge caacaacatg 720
gctactgcct tcactcctca cccttgcact atcattggcc agagccgctg cgagggcgac 780
tcgtgcggtg gcacctacag caacgaccgc tacgccggcg tctgcgacce cgatggctgc 840
gacttcaacg cgtatcgcca gggcaacaag accttctacg gcaagggcat gaccgtcgac 900
accaccaaga agctcaccgt cgtcaccag ttctcaagg acgccaacgg cgatctcggc 960
gagatcaagc gcttctacgt ccaggatggg aagatcatcc ccaactccga gtccaccatc 1020
cccggcgtcg agggcaactc catcaccag gattgggtgc accgccagaa gggtgccttt 1080
ggcgacattg acgacttcaa ccgcaagggc ggcatgaagc agatgggcaa ggccctcgcc 1140
ggcccatgg tcctgggtcat gtccatctgg gatgaccacg cctccaacat gctctggctc 1200
gactcgacct tccctgtcga tgccgctggc aagcccggcg ccgagcgcgg tgccctgccc 1260
accacctcgg gtgtccctgc tgaggttgag gccgaggccc ccaacagcaa cgtcgtcttc 1320
tccaacatcc gcttcggccc catcggtcgc accgttgccg gccttcccag cgatggcggc 1380
aacaacggcg gcaacaccac cgtccagccc ccgcccagca ccaccaccac ctctgccagc 1440
agcagcacca cctcgggtcc tgccaccacc accaccgcca gcgctggccc caaggctggc 1500
cgctggcagc agtgcgggcg catcggtctc actggcccga cccagtgcga ggagccctac 1560
acttgacca agctcaacga ctggtactct cagtgcctgt aa 1602

```

<210> 9

<211> 533

<212> PRT

<213> *Scytalidium thermophilum*

<400> 9

```

Met Arg Thr Ala Lys Phe Ala Thr Leu Ala Ala Leu Val Ala Ser Ala
 1          5          10          15
Ala Ala Gln Gln Ala Cys Ser Leu Thr Thr Glu Arg His Pro Ser Leu
 20          25          30
Ser Trp Lys Lys Cys Thr Ala Gly Gly Gln Cys Gln Thr Val Gln Ala
 35          40          45
Ser Ile Thr Leu Asp Ser Asn Trp Arg Trp Thr His Gln Val Ser Gly
 50          55          60
Ser Thr Asn Cys Tyr Thr Gly Asn Glu Trp Asp Ser Ser Ile Cys Thr
 65          70          75          80
Asp Ala Lys Ser Cys Ala Gln Asn Cys Cys Val Asp Gly Ala Asp Tyr
 85          90          95
Thr Ser Thr Tyr Gly Ile Thr Thr Asn Gly Asp Ser Leu Ser Leu Lys
100          105          110
Phe Val Thr Lys Gly Gln Tyr Ser Thr Asn Val Gly Ser Arg Thr Tyr
115          120          125
Leu Met Asp Gly Glu Asp Lys Tyr Gln Thr Phe Glu Leu Leu Gly Asn
130          135          140
Glu Phe Thr Phe Asp Val Asp Val Ser Asn Ile Gly Cys Gly Leu Asn
145          150          155          160
Gly Ala Leu Tyr Phe Val Ser Met Asp Ala Asp Gly Gly Leu Ser Arg
165          170          175
Tyr Pro Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp
180          185          190
Ala Gln Cys Pro Arg Asp Ile Lys Phe Ile Asn Gly Glu Ala Asn Ile
195          200          205

```

Glu	Gly	Trp	Thr	Gly	Ser	Thr	Asn	Asp	Pro	Asn	Ala	Gly	Ala	Gly	Arg
	210					215					220				
Tyr	Gly	Thr	Cys	Cys	Ser	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Asn	Met
225					230					235					240
Ala	Thr	Ala	Phe	Thr	Pro	His	Pro	Cys	Thr	Ile	Ile	Gly	Gln	Ser	Arg
				245					250						255
Cys	Glu	Gly	Asp	Ser	Cys	Gly	Gly	Thr	Tyr	Ser	Asn	Asp	Arg	Tyr	Ala
			260					265					270		
Gly	Val	Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Ala	Tyr	Arg	Gln	Gly
		275					280					285			
Asn	Lys	Thr	Phe	Tyr	Gly	Lys	Gly	Met	Thr	Val	Asp	Thr	Thr	Lys	Lys
	290					295					300				
Leu	Thr	Val	Val	Thr	Gln	Phe	Leu	Lys	Asp	Ala	Asn	Gly	Asp	Leu	Gly
305					310					315					320
Glu	Ile	Lys	Arg	Phe	Tyr	Val	Gln	Asp	Gly	Lys	Ile	Ile	Pro	Asn	Ser
				325					330						335
Glu	Ser	Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp
			340					345						350	
Cys	Asp	Arg	Gln	Lys	Val	Ala	Phe	Gly	Asp	Ile	Asp	Asp	Phe	Asn	Arg
		355					360					365			
Lys	Gly	Gly	Met	Lys	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val
	370					375					380				
Leu	Val	Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Ser	Asn	Met	Leu	Trp	Leu
385					390					395					400
Asp	Ser	Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg
				405					410						415
Gly	Ala	Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu
			420					425					430		
Ala	Pro	Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile
		435					440					445			
Gly	Ser	Thr	Val	Ala	Gly	Leu	Pro	Ser	Asp	Gly	Gly	Asn	Asn	Gly	Gly
	450					455					460				
Asn	Thr	Thr	Val	Gln	Pro	Pro	Pro	Ser	Thr	Thr	Thr	Thr	Ser	Ala	Ser
465					470					475					480
Ser	Ser	Thr	Thr	Ser	Ala	Pro	Ala	Thr	Thr	Thr	Thr	Ala	Ser	Ala	Gly
				485					490						495
Pro	Lys	Ala	Gly	Arg	Trp	Gln	Gln	Cys	Gly	Gly	Ile	Gly	Phe	Thr	Gly
			500					505					510		
Pro	Thr	Gln	Cys	Glu	Glu	Pro	Tyr	Thr	Cys	Thr	Lys	Leu	Asn	Asp	Trp
		515					520					525			
Tyr	Ser	Gln	Cys	Leu											
	530														

<210> 10

<211> 497

<212> PRT

<213> Hypocrea jecorina

<400> 10

Gln	Ser	Ala	Cys	Thr	Leu	Gln	Ser	Glu	Thr	His	Pro	Pro	Leu	Thr	Trp
1				5					10					15	
Gln	Lys	Cys	Ser	Ser	Gly	Gly	Thr	Cys	Thr	Gln	Gln	Thr	Gly	Ser	Val
			20					25					30		
Val	Ile	Asp	Ala	Asn	Trp	Arg	Trp	Thr	His	Ala	Thr	Asn	Ser	Ser	Thr
		35					40					45			
Asn	Cys	Tyr	Asp	Gly	Asn	Thr	Trp	Ser	Ser	Thr	Leu	Cys	Pro	Asp	Asn
	50					55					60				
Glu	Thr	Cys	Ala	Lys	Asn	Cys	Cys	Leu	Asp	Gly	Ala	Ala	Tyr	Ala	Ser
65					70					75					80

10

<400> 11

Gln	Gln	Ala	Cys	Ser	Leu	Thr	Thr	Glu	Arg	His	Pro	Ser	Leu	Ser	Trp
1				5					10					15	
Lys	Lys	Cys	Thr	Ala	Gly	Gly	Gln	Cys	Gln	Thr	Val	Gln	Ala	Ser	Ile
			20					25					30		
Thr	Leu	Asp	Ser	Asn	Trp	Arg	Trp	Thr	His	Gln	Val	Ser	Gly	Ser	Thr
		35					40					45			
Asn	Cys	Tyr	Thr	Gly	Asn	Glu	Trp	Asp	Ser	Ser	Ile	Cys	Thr	Asp	Ala
	50					55					60				
Lys	Ser	Cys	Ala	Gln	Asn	Cys	Cys	Val	Asp	Gly	Ala	Asp	Tyr	Thr	Ser
65					70					75					80
Thr	Tyr	Gly	Ile	Thr	Thr	Asn	Gly	Asp	Ser	Leu	Ser	Leu	Lys	Phe	Val
			85						90					95	
Thr	Lys	Gly	Gln	Tyr	Ser	Thr	Asn	Val	Gly	Ser	Arg	Thr	Tyr	Leu	Met
			100					105					110		
Asp	Gly	Glu	Asp	Lys	Tyr	Gln	Thr	Phe	Glu	Leu	Leu	Gly	Asn	Glu	Phe
		115					120					125			
Thr	Phe	Asp	Val	Asp	Val	Ser	Asn	Ile	Gly	Cys	Gly	Leu	Asn	Gly	Ala
	130					135					140				
Leu	Tyr	Phe	Val	Ser	Met	Asp	Ala	Asp	Gly	Gly	Leu	Ser	Arg	Tyr	Pro
145					150					155					160
Gly	Asn	Lys	Ala	Gly	Ala	Lys	Tyr	Gly	Thr	Gly	Tyr	Cys	Asp	Ala	Gln
			165					170						175	
Cys	Pro	Arg	Asp	Ile	Lys	Phe	Ile	Asn	Gly	Glu	Ala	Asn	Ile	Glu	Gly
			180					185					190		
Trp	Thr	Gly	Ser	Thr	Asn	Asp	Pro	Asn	Ala	Gly	Ala	Gly	Arg	Tyr	Gly
		195					200					205			
Thr	Cys	Cys	Ser	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Asn	Met	Ala	Thr
	210					215					220				
Ala	Phe	Thr	Pro	His	Pro	Cys	Thr	Ile	Ile	Gly	Gln	Ser	Arg	Cys	Glu
225					230					235					240
Gly	Asp	Ser	Cys	Gly	Gly	Thr	Tyr	Ser	Asn	Asp	Arg	Tyr	Ala	Gly	Val
			245						250					255	
Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Ala	Tyr	Arg	Gln	Gly	Asn	Lys
			260				265						270		
Thr	Phe	Tyr	Gly	Lys	Gly	Met	Thr	Val	Asp	Thr	Thr	Lys	Lys	Leu	Thr
		275					280					285			
Val	Val	Thr	Gln	Phe	Leu	Lys	Asp	Ala	Asn	Gly	Asp	Leu	Gly	Glu	Ile
	290					295					300				
Lys	Arg	Phe	Tyr	Val	Gln	Asp	Gly	Lys	Ile	Ile	Pro	Asn	Ser	Glu	Ser
305					310					315					320
Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	Asp
			325						330					335	
Arg	Gln	Lys	Val	Ala	Phe	Gly	Asp	Ile	Asp	Asp	Phe	Asn	Arg	Lys	Gly
			340				345						350		
Gly	Met	Lys	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	Val
		355					360					365			
Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Ser	Asn	Met	Leu	Trp	Leu	Asp	Ser
	370					375					380				
Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala
385					390					395					400
Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro
			405						410					415	
Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser
			420					425					430		
Thr	Val	Ala	Gly	Leu	Pro	Ser	Asp	Gly	Gly	Asn	Asn	Gly	Gly	Asn	Thr
		435					440					445			
Thr	Val	Gln	Pro	Pro	Pro	Ser	Thr	Thr	Thr	Thr	Ser	Ala	Ser	Ser	Ser

450		455		460
Thr Thr Ser Ala Pro Ala Thr Thr Thr Thr Ala Ser Ala Gly Pro Lys				
465		470		475
Ala Gly Arg Trp Gln Gln Cys Gly Gly Ile Gly Phe Thr Gly Pro Thr				
		485		490
Gln Cys Glu Glu Pro Tyr Thr Cys Thr Lys Leu Asn Asp Trp Tyr Ser				
		500		505
				510
Gln Cys Leu				
515				

<210> 12

<211> 507

<212> PRT

<213> Artificial Sequence

<220>

<223> consensus sequence

<400> 12

Gln Gln Ala Cys Ser Leu Thr Thr Glu Arg His Pro Ser Leu Ser Trp				
1		5		10
Lys Lys Cys Thr Ala Gly Gly Gln Cys Gln Thr Val Gln Ala Ser Ile				
		20		25
Thr Leu Asp Ser Asn Trp Arg Trp Thr His Gln Val Ser Gly Ser Thr				
		35		40
Asn Cys Tyr Thr Gly Asn Lys Trp Asp Ser Ser Ile Cys Thr Asp Ala				
		50		55
Lys Ser Cys Ala Gln Asn Cys Cys Val Asp Gly Ala Asp Tyr Thr Ser				
65		70		75
Thr Tyr Gly Ile Thr Thr Asn Gly Asp Ser Leu Ser Leu Lys Phe Val				
		85		90
Thr Lys Gly Gln His Ser Thr Asn Val Gly Ser Arg Thr Tyr Leu Met				
		100		105
Asp Gly Glu Asp Lys Tyr Gln Thr Phe Glu Leu Leu Gly Asn Glu Phe				
		115		120
Thr Phe Asp Val Asp Val Ser Asn Ile Gly Cys Gly Leu Asn Gly Ala				
		130		135
Leu Tyr Phe Val Ser Met Asp Ala Asp Gly Gly Leu Ser Arg Tyr Pro				
145		150		155
Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp Ala Gln				
		165		170
Cys Pro Arg Asp Ile Lys Phe Ile Asn Gly Glu Ala Asn Ile Glu Gly				
		180		185
Trp Thr Gly Ser Thr Asn Asp Pro Asn Ala Gly Ala Gly Arg Tyr Gly				
		195		200
Thr Cys Cys Ser Glu Met Asp Ile Trp Glu Ala Asn Asn Met Ala Thr				
		210		215
Ala Phe Thr Pro His Pro Cys Thr Ile Ile Gly Gln Ser Arg Cys Glu				
225		230		235
Gly Asp Ser Cys Gly Gly Thr Tyr Ser Asn Glu Arg Tyr Ala Gly Val				
		245		250
Cys Asp Pro Asp Gly Cys Asp Phe Asn Ser Tyr Arg Gln Gly Asn Lys				
		260		265
Thr Phe Tyr Gly Lys Gly Met Thr Val Asp Thr Thr Lys Lys Ile Thr				
		275		280
Val Val Thr Gln Phe Leu Lys Asp Ala Asn Gly Asp Leu Gly Glu Ile				
		290		295
Lys Arg Phe Tyr Val Gln Asp Gly Lys Ile Ile Pro Asn Ser Glu Ser				
305		310		315
				320

Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	Asp	
				325					330					335		
Arg	Gln	Lys	Val	Ala	Phe	Gly	Asp	Ile	Asp	Asp	Phe	Asn	Arg	Lys	Gly	
			340					345					350			
Gly	Met	Lys	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	Val	
		355					360					365				
Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Ser	Asn	Met	Leu	Trp	Leu	Asp	Ser	
	370					375					380					
Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala	
385					390					395					400	
Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro	
				405					410					415		
Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser	
			420					425					430			
Thr	Val	Ala	Gly	Leu	Pro	Gly	Ala	Gly	Asn	Gly	Gly	Asn	Asn	Gly	Gly	
		435					440					445				
Asn	Pro	Pro	Pro	Pro	Thr	Thr	Thr	Thr	Ser	Ser	Ala	Pro	Ala	Thr	Thr	
	450					455					460					
Thr	Thr	Ala	Ser	Ala	Gly	Pro	Lys	Ala	Gly	Arg	Trp	Gln	Gln	Cys	Gly	
465					470					475					480	
Gly	Ile	Gly	Phe	Thr	Gly	Pro	Thr	Gln	Cys	Glu	Glu	Pro	Tyr	Thr	Cys	
				485					490					495		
Thr	Lys	Leu	Asn	Asp	Trp	Tyr	Ser	Gln	Cys	Leu						
			500					505								

<210> 13

<211> 507

<212> PRT

<213> Artificial Sequence

<220>

<223> consensus sequence

<400> 13

Gln	Gln	Ala	Cys	Ser	Leu	Thr	Thr	Glu	Arg	His	Pro	Ser	Leu	Ser	Trp	
1				5					10					15		
Lys	Lys	Cys	Thr	Ala	Gly	Gly	Gln	Cys	Gln	Thr	Val	Gln	Ala	Ser	Ile	
			20					25					30			
Thr	Leu	Asp	Ser	Asn	Trp	Arg	Trp	Thr	His	Gln	Val	Ser	Gly	Ser	Thr	
		35					40					45				
Asn	Cys	Tyr	Thr	Gly	Asn	Lys	Trp	Asp	Thr	Ser	Ile	Cys	Thr	Asp	Ala	
	50					55					60					
Lys	Ser	Cys	Ala	Gln	Asn	Cys	Cys	Val	Asp	Gly	Ala	Asp	Tyr	Thr	Ser	
65					70					75					80	
Thr	Tyr	Gly	Ile	Thr	Thr	Asn	Gly	Asp	Ser	Leu	Ser	Leu	Lys	Phe	Val	
				85					90					95		
Thr	Lys	Gly	Gln	His	Ser	Thr	Asn	Val	Gly	Ser	Arg	Thr	Tyr	Leu	Met	
			100					105						110		
Asp	Gly	Glu	Asp	Lys	Tyr	Gln	Thr	Phe	Glu	Leu	Leu	Gly	Asn	Glu	Phe	
		115					120						125			
Thr	Phe	Asp	Val	Asp	Val	Ser	Asn	Ile	Gly	Cys	Gly	Leu	Asn	Gly	Ala	
	130						135					140				
Leu	Tyr	Phe	Val	Ser	Met	Asp	Ala	Asp	Gly	Gly	Leu	Ser	Arg	Tyr	Pro	
145					150					155					160	
Gly	Asn	Lys	Ala	Gly	Ala	Lys	Tyr	Gly	Thr	Gly	Tyr	Cys	Asp	Ala	Gln	
				165					170					175		
Cys	Pro	Arg	Asp	Ile	Lys	Phe	Ile	Asn	Gly	Glu	Ala	Asn	Ile	Glu	Gly	
			180					185					190			
Trp	Thr	Gly	Ser	Thr	Asn	Asp	Pro	Asn	Ala	Gly	Ala	Gly	Arg	Tyr	Gly	

	195		200		205										
Thr	Cys	Cys	Ser	Glu	Met	Asp	Ile	Trp	Glu	Ala	Asn	Asn	Met	Ala	Thr
	210					215					220				
Ala	Phe	Thr	Pro	His	Pro	Cys	Thr	Ile	Ile	Gly	Gln	Ser	Arg	Cys	Glu
225					230					235					240
Gly	Asp	Ser	Cys	Gly	Gly	Thr	Tyr	Ser	Asn	Glu	Arg	Tyr	Ala	Gly	Val
				245					250						255
Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Ser	Tyr	Arg	Gln	Gly	Asn	Lys
			260				265						270		
Thr	Phe	Tyr	Gly	Lys	Gly	Met	Thr	Val	Asp	Thr	Thr	Lys	Lys	Ile	Thr
		275					280					285			
Val	Val	Thr	Gln	Phe	Leu	Lys	Asp	Ala	Asn	Gly	Asp	Leu	Gly	Glu	Ile
	290					295				300					
Lys	Arg	Phe	Tyr	Val	Gln	Asp	Gly	Lys	Ile	Ile	Pro	Asn	Ser	Glu	Ser
305					310					315					320
Thr	Ile	Pro	Gly	Val	Glu	Gly	Asn	Ser	Ile	Thr	Gln	Asp	Trp	Cys	Asp
				325					330					335	
Arg	Gln	Lys	Val	Ala	Phe	Gly	Asp	Ile	Asp	Asp	Phe	Asn	Arg	Lys	Gly
			340				345						350		
Gly	Met	Lys	Gln	Met	Gly	Lys	Ala	Leu	Ala	Gly	Pro	Met	Val	Leu	Val
	355						360					365			
Met	Ser	Ile	Trp	Asp	Asp	His	Ala	Ser	Asn	Met	Leu	Trp	Leu	Asp	Ser
	370					375					380				
Thr	Phe	Pro	Val	Asp	Ala	Ala	Gly	Lys	Pro	Gly	Ala	Glu	Arg	Gly	Ala
385					390					395					400
Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Glu	Val	Glu	Ala	Glu	Ala	Pro
				405					410					415	
Asn	Ser	Asn	Val	Val	Phe	Ser	Asn	Ile	Arg	Phe	Gly	Pro	Ile	Gly	Ser
			420				425						430		
Thr	Val	Ala	Gly	Leu	Pro	Gly	Ala	Gly	Asn	Gly	Gly	Asn	Asn	Gly	Gly
	435						440					445			
Asn	Pro	Pro	Pro	Pro	Thr	Thr	Thr	Thr	Ser	Ser	Ala	Pro	Ala	Thr	Thr
	450					455					460				
Thr	Thr	Ala	Ser	Ala	Gly	Pro	Lys	Ala	Gly	Arg	Trp	Gln	Gln	Cys	Gly
465					470					475					480
Gly	Ile	Gly	Phe	Thr	Gly	Pro	Thr	Gln	Cys	Glu	Glu	Pro	Tyr	Thr	Cys
				485					490					495	
Thr	Lys	Leu	Asn	Asp	Trp	Tyr	Ser	Gln	Cys	Leu					
			500					505							

<210> 14
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 14
 atgcgtaccg ccaagttcgc

20

<210> 15
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 15
ttacaggcac tgagagtacc ag

22

- 45 -

CLAIMS:

1. An isolated polypeptide comprising the amino acid sequence of SEQ ID NO: 3.
2. The polypeptide of claim 1 further comprising a signal sequence.
3. An isolated polypeptide comprising the amino acid sequence of SEQ ID NO: 4.
4. The polypeptide of claim 3 further comprising a signal sequence.
5. A polynucleotide encoding the polypeptide of SEQ ID NO: 3.
6. A polynucleotide encoding the polypeptide of SEQ ID NO: 4.
7. A nucleic acid construct comprising a polynucleotide according to claim 5 or 6, operably linked to one or more control sequences.
8. A recombinant expression vector comprising the nucleic acid construct of claim 7.
9. A host cell transformed with the nucleic acid construct of claim 8.
10. A method for producing a CBH1 polypeptide, said method comprising:
 - a) transforming a host cell with a nucleic acid comprising a polynucleotide according to claim 5 or 6;
 - b) culturing the host cell under conditions to produce the polypeptide; and
 - c) recovering the polypeptide.

- 46 -

11. A detergent composition comprising a polypeptide having cellobiohydrolase I activity according to any one of claims 1 to 4 and a surfactant.

12. A method of converting biomass to sugars comprising contacting the biomass with a polypeptide having cellobiohydrolase I activity according to any one of claims 1 to 4 wherein said polypeptide has enhanced activity relative to *T. reesei* in either a pretreated corn stover (PCS) conversion assay at 65°C or a phosphoric acid swollen cellulase (PASC) assay at 38°C.

13. A method of converting biomass to sugars comprising contacting said biomass with a polypeptide having cellobiohydrolase I activity according to any one of claims 1 to 4.

H. grisea CBH1.1

Figure 1: Total Genomic Sequence (1638 nucleotides):

1	ATGCGTACCG	CCAAGTTCGC	CACCCTCGCC	GCCCTTGTGG	CCTCGGCCGC	50
51	CGCCCAGCAG	GCGTGCAGTC	TCACCACCGA	GAGGCACCCT	TCCCTCTCTT	100
101	GGAAGAAGTG	CACCGCCGGC	GGCCAGTGCC	AGACCGTCCA	GGCTTCCATC	150
151	ACTCTCGACT	CCAAGTGGCG	CTGGACTCAC	CAGGTGTCTG	GCTCCACCAA	200
201	CTGCTACACG	GGCAACAAGT	GGGATACTAG	CATCTGCACT	GATGCCAAGT	250
251	CGTGCGCTCA	GAAGTGTGCT	GTCGATGGTG	CCGACTACAC	CAGCACCTAT	300
301	GGCATCACCA	CCAACGGTGA	TTCCCTGAGC	CTCAAGTTCT	TCACCAAGGG	350
351	CCAGCACTCG	ACCAACGTCG	GCTCGCGTAC	CTACCTGATG	GACGGCGAGG	400
401	ACAAGTATCA	GAGTACGTTT	TATCTTCAGC	CTTCTCGCGC	CTTGAATCCT	450
451	GGCTAACGTT	TACACTTCAC	AGCCTTCGAG	CTCCTCGGCA	ACGAGTTCAC	500
501	CTTCGATGTC	GATGTCTCCA	ACATCGGCTG	CGGTCTCAAC	GGCGCCCTGT	550
551	ACTTCGTCTC	CATGGACGCC	GATGGTGGTC	TCAGCCGCTA	TCCTGGCAAC	600
601	AAGGCTGGTG	CCAAGTACGG	TACCGGCTAC	TGCGATGCTC	AGTGCCCCCG	650
651	TGACATCAAG	TTCATCAACG	GCGAGGCCAA	CATTGAGGGC	TGGACCGGCT	700
701	CCACCAACGA	CCCCAACGCC	GGCGCGGGCC	GCTATGGTAC	CTGCTGCTCT	750
751	GAGATGGATA	TCTGGGAAGC	CAACAACATG	GCTACTGCCT	TCACTCCTCA	800
801	CCCTTGACAC	ATCATTTGCC	AGAGCCGCTG	CGAGGGCGAC	TCGTGCGGTG	850
851	GCACCTACAG	CAACGAGCGC	TACGCCGGCG	TCTGCGACCC	CGATGGCTGC	900
901	GACTTCAACT	CGTACCGCCA	GGGCAACAAG	ACCTTCTACG	GCAAGGGCAT	950
951	GACCGTCGAC	ACCACCAAGA	AGATCACTGT	CGTCACCCAG	TTCTTCAAGG	1000
1001	ATGCCAACGG	CGATCTCGGC	GAGATCAAGC	GCTTCTACGT	CCAGGATGGC	1050
1051	AAGATCATCC	CCAAGTCCGA	GTCCACCATC	CCCGGCGTCG	AGGGCAATTC	1100
1101	CATCACCCAG	GACTGGTGCG	ACCGCCAGAA	GGTTGCCTTT	GGCGACATTG	1150
1151	ACGACTTCAA	CCGCAAGGGC	GGCATGAAGC	AGATGGGCAA	GGCCCTCGCC	1200
1201	GGCCCCATGG	TCCTGGTTCAT	GTCCATCTGG	GATGACCACG	CCTCCAACAT	1250
1251	GCTCTGGCTC	GACTCGACCT	TCCCTGTCTG	TGCCGCTGGC	AAGCCCGGCG	1300
1301	CCGAGCGCGG	TGCCTGCCCC	ACCACCTCGG	GTGTCCCTGC	TGAGGTTGAG	1350
1351	GCCGAGGCCC	CCAACAGCAA	CGTCGTCTTC	TCCAACATCC	GCTTCGGCCC	1400
1401	CATCGGCTCG	ACCGTTGCTG	GTCTCCCCGG	CGCGGGCAAC	GGCGGCAACA	1450
1451	ACGGCGGCAA	CCCCCCGCCC	CCCACCACCA	CCACCTCCTC	GGCTCCGGCC	1500
1501	ACCACCACCA	CCGCCAGCGC	TGGCCCCAAG	GCTGGCCGCT	GGCAGCAGTG	1550
1551	CGGCGGCATC	GGCTTCACTG	GCCCGACCCA	GTGCGAGGAG	CCCTACACTT	1600
1601	GCACCAAGCT	CAACGACTGG	TACTCTCAGT	GCCTGTAA		1638

Figure 2: Putative intron sequence deleted (GTACGTT...CAG = 413-472) Gives the cDNA sequence (1578 nucleotides):

1	ATGCGTACCG	CCAAGTTCGC	CACCCTCGCC	GCCCTTGTGG	CCTCGGCCGC	50
51	CGCCCAGCAG	GCGTGCAATC	TCACCACCGA	GAGGCACCCT	TCCCTCTCTT	100
101	GGAAGAAGTG	CACCGCCGGC	GGCCAGTGCC	AGACCGTCCA	GGCTTCCATC	150
151	ACTCTCGACT	CCAAGTGGCG	CTGGACTCAC	CAGGTGTCTG	GCTCCACCAA	200
201	CTGCTACACG	GGCAACAAGT	GGGATACTAG	CATCTGCACT	GATGCCAAGT	250
251	CGTGCCTCA	GAAGTGTGTC	GTCGATGGTG	CCGACTACAC	CAGCACCTAT	300
301	GGCATCACCA	CCAACGGTGA	TTCCCTGAGC	CTCAAGTTCG	TCACCAAGGG	350
351	CCAGCACTCG	ACCAACGTCG	GCTCGCGTAC	CTACCTGATG	GACGGCGAGG	400
401	ACAAGTATCA	GACCTTCGAG	CTCCTCGGCA	ACGAGTTCAC	CTTCGATGTC	450
451	GATGTCTCCA	ACATCGGCTG	CGGTCTCAAC	GGCGCCCTGT	ACTTCGTCTC	500
501	CATGGACGCC	GATGGTGGTC	TCAGCCGCTA	TCCTGGCAAC	AAGGCTGGTG	550
551	CCAAGTACGG	TACCGGCTAC	TGCGATGCTC	AGTGGCCCCG	TGACATCAAG	600
601	TTCATCAACG	GCGAGGCCAA	CATTGAGGGC	TGGACCGGCT	CCACCAACGA	650
651	CCCCAACGCC	GGCGCGGGCC	GCTATGGTAC	CTGCTGCTCT	GAGATGGATA	700
701	TCTGGGAAGC	CAACAACATG	GCTACTGCCT	TCACTCCTCA	CCCTTGCACC	750
751	ATCATTTGGC	AGAGCCGCTG	CGAGGGCGAC	TCGTGCGGTG	GCACCTACAG	800
801	CAACGAGCGC	TACGCCGGCG	TCTGCGACCC	CGATGGCTGC	GAATTCAACT	850
851	CGTACCGCCA	GGGCAACAAG	ACCTTCTACG	GCAAGGGCAT	GACCGTCGAC	900
901	ACCACCAAGA	AGATCACTGT	CGTCACCCAG	TTCTTCAAGG	ATGCCAACGG	950
951	CGATCTCGGC	GAGATCAAGC	GCTTCTACGT	CCAGGATGGC	AAGATCATCC	1000
1001	CCAAGTCCGA	GTCCACCATC	CCCGGCGTCG	AGGGCAATTC	CATCACCAG	1050
1051	GACTGGTGCG	ACCGCCAGAA	GGTTGCCTTT	GGCGACATTG	ACGACTTCAA	1100
1101	CCGCAAGGGC	GGCATGAAGC	AGATGGGCAA	GGCCCTCGCC	GGCCCCATGG	1150
1151	TCCTGGTCAT	GTCCATCTGG	GATGACCACG	CCTCCAACAT	GCTCTGGCTC	1200
1201	GACTCGACCT	TCCCTGTCGA	TGCCGCTGGC	AAGCCCGGCG	CCGAGCGCGG	1250
1251	TGCCTGCCCC	ACCACCTCGG	GTGTCCCTGC	TGAGGTTGAG	GCCGAGGCCC	1300
1301	CCAACAGCAA	CGTCGTCTTC	TCCAACATCC	GCTTCGGCCC	CATCGGCTCG	1350
1351	ACCGTTGCTG	GTCTCCCCGG	CGCGGGCAAC	GGCGGCAACA	ACGGCGGCAA	1400
1401	CCCCCGCCCC	CCCACCAACCA	CCACCTCCTC	GGCTCCGGCC	ACCACCAACCA	1450
1451	CCGCCAGCGC	TGGCCCCAAG	GCTGGCCGCT	GGCAGCAGTG	CGGCGGCATC	1500
1501	GGCTTCACTG	GCCCGACCCA	GTGCGAGGAG	CCCTACACTT	GCACCAAGCT	1550
1551	CAACGACTGG	TACTCTCAGT	GCCTGTAA			1578

Figure 3: Translation of the cDNA sequence gives the *H. grisea* var *thermoidea* CBH1 precursor (i.e.: with signal sequence) protein sequence (525 amino acids):

1	<u>MRTAKFATLA</u>	<u>ALVASAAAQQ</u>	ACSLTTERHP	SLSWKKCTAG	GQCQTVQASI	50
51	TLDSNWRWITH	QVSGSTNCYT	GNKWDTSICT	DAKSCAQNCC	VDGADYTSTY	100
101	GITTNGDSLS	LKFVTKGQHS	TNVGSRTYLM	DGEDKYQTFE	LLGNEFTFDV	150
151	DVSNIGCGLN	GALYFVSMDA	DGGLSRYPGN	KAGAKYGTGY	CDAQCPRIK	200
201	FINGEANIEG	WTGSTNDPNA	GAGRYGTCCS	EMDIWEANNM	ATAFTPHPCT	250
251	IIGQSRCEGD	SCGGTYSNER	YAGVCDPDGC	DFNSYRQGNK	TFYGKGMTVD	300
301	TTKKITVVTQ	FLKDANGDLG	EIKRFYVQDG	KIIPNSESTI	PGVEGNSITQ	350
351	DWCDRQKVAF	GDIDDFNRKG	GMKQMGKALA	GPMVLVMSIW	DDHASNMLWL	400
401	DSTFPVDAAG	KPGAERGACP	TTSGVPAEVE	AEAPNSNVVF	SNIRFGPIGS	450
451	TVAGLPGAGN	GGNNGGNPPP	PTTTTSSAPA	TTTTASAGPK	AGRWQQCGGI	500
501	GFTGPTQCEE	PYTCTKLNDW	YSQCL			525

Figure 4: Mature (i.e.: expressed protein with the putative signal sequence removed) protein sequence (507 amino acids):

1	QQACSLTTER	HPSLSWKKCT	AGGQCQTVQA	SITLDSNWRW	THQVSGSTNC	50
51	YTGNKWDTSI	CTDAKSCAQN	CCVDGADYTS	TYGITTNGDS	LSLKFVTKGQ	100
101	HSTNVGSRTY	LMDGEDKYQT	FELLGNEFTF	DVDVSNIGCG	LNGALYFVSM	150
151	DADGGLSRYP	GNKAGAKYGT	GYCDAQCPRD	IKFINGEANI	EGWTGSTNDP	200
201	NAGAGRYGTC	CSEMDIWEAN	NMATAFTPHP	CTIIGQSRCE	GDSCGGTYSN	250
251	ERYAGVCDPD	GCDFNSYRQG	NKTFYGKGMT	VDTTKKITVV	TQFLKDANGD	300
301	LGEIKRFYVQ	DGKIIPNSES	TIPGVEGNSI	TQDWCDRQKV	AFGDIDDFNR	350
351	KGGMKQMGKA	LAGPMVLVMS	IWDDHASNML	WLDSTFPVDA	AGKPGAERGA	400
401	CPTTSGVPAE	VEAEAPNSNV	VFSNIRFGPI	GSTVAGLPGA	GNGGNNGGNP	450
451	PPPTTTTSSA	PATTTTASAG	PKAGRWQQCG	GIGFTGPTQC	EEPYTCTKLN	500
501	DWYSQCL					507

Figure 5. Sequence alignment of two public sequences and variant *H. grisea* CBH1.1




Figure 6: pRAX1



Figure 7: Destination vector pRAXdes2 for expression in *A. niger*

Figure 8: Replicative expression pRAXdesCBH1 vector of CBH1 genes under the control of the glucoamylase promotor.







Figure 11: CBHI Conversion Assay:
38C, 700 rpm agitation, 1 day, 12.66% PCS, 15.5 mg enzyme/g cellulose




Figure 12: CBHI Conversion Assay:
65C, 700 rpm agitation, 1 day, 12.66% PCS, 15.5 mg enzyme/g cellulose




Figure 13A
3 fold rate difference b/n *H. grisea* and rCBHI




Figure 13B
4.8 fold rate difference





Figure 13C



```
1  ATGCGTACCG CCAAGTTCGC CACCCTCGCC GCCCTTGTGG CCTCGGCCGC
51  CGCCCAGCAG GCGTGCAGCC TCACCACCGA GAGGCACCCT TCCCTCTCCT
101 GGAAGAAGTG CACCGCCGGC GGCCAGTGCC AGACCGTCCA GGCTTCCATC
151 ACTCTCGACT CCAACTGGCG CTGGACTCAC CAGGTGTCTG GCTCCACCAA
201 CTGCTACACG GGCAACGAGT GGGATTCTAG CATCTGCACT GATGCCAAGT
251 CGTGCGCTCA GAACTGCTGC GTCGATGGTG CTGACTACAC CAGCACCTAT
301 GGCATCACCA CCAACGGTGA TTCCCTGAGC CTCAAGTTCG TCACCAAGGG
351 CCAGTACTCG ACCAACGTCG GCTCGCGTAC CTACCTGATG GACGGCGAGG
401 ACAAGTATCA GAGTAGGTTC TATCTTCAGC CTTCTCGCGC CTTGAATCCT
451 GGCTAACTTT TACACTTCAC AGCCTTCGAG CTCCTCGGCA ACGAGTTCAC
501 CTTGATGTC GATGTCTCCA ACATCGGCTG CGGTCTCAAC GGCGCCCTGT
551 ACTTCGTCTC CATGGACGCC GATGGTGGTC TCAGCCGCTA TCCTGGCAAC
601 AAGGCTGGTG CCAAGTACGG TACCGGCTAC TGCGATGCTC AGTGCCCCCG
651 TGACATCAAG TTCATCAACG GCGAGGCCAA CATTGAGGGC TGGACCGGCT
701 CCACCAACGA CCCCAACGCC GCGCGGGGCC GCTATGGTAC CTGCTGCTCT
751 GAGATGGATA TCTGGGAGGC CAACAACATG GCTACTGCCT TCACTCCTCA
801 CCCTTGCACT ATCATTGGCC AGAGCCGCTG CGAGGGCGAC TCGTGCGGTG
851 GCACCTACAG CAACGACCGC TACGCCGGCG TCTGCGACCC CGATGGCTGC
901 GACTTCAACG CGTATCGCCA GGGCAACAAG ACCTTCTACG GCAAGGGCAT
951 GACCGTCGAC ACCACCAAGA AGCTCACCGT CGTCACCCAG TTCCTCAAGG
1001 ACGCCAACGG CGATCTCGGC GAGATCAAGC GCTTCTACGT CCAGGATGGG
1051 AAGATCATCC CCAACTCCGA GTCCACCATC CCCGGCGTCG AGGGCAACTC
1101 CATCACCCAG GATTGGTGCG ACCGCCAGAA GGTTCCTTTT GGCGACATTG
1151 ACGACTTCAA CCGCAAGGGC GGCATGAAGC AGATGGGCAA GGCCCTCGCC
1201 GGCCCCATGG TCCTGGTCAT GTCCATCTGG GATGACCACG CCTCCAACAT
1251 GCTCTGGCTC GACTCGACCT TCCCTGTCGA TGCCGCTGGC AAGCCCGGCG
1301 CCGAGCGCGG TGCTTGCCCG ACCACCTCGG GTGTCCCTGC TGAGGTTGAG
1351 GCCGAGGCCC CCAACAGCAA CGTCGTCTTC TCCAACATCC GCTTCGGCCC
1401 CATCGGCTCG ACCGTTGCCG GCCTTCCCAG CGATGGCGGC AACAACGGCG
1451 GCAACACCAC CGTCCAGCCC CCGCCCAGCA CCACCACCAC CTCTGCCAGC
1501 AGCAGCACCA CCTCGGCTCC TGCCACCACC ACCACCGCCA GCGCTGGCCC
1551 CAAGGCTGGC CGCTGGCAGC AGTGCGGCGG CATCGGCTTC ACTGGCCCGA
1601 CCCAGTGCGA GGAGCCCTAC ACTTGACCA AGCTCAACGA CTGGTACTCT
1651 CAGTGCCTGT AA
```

Figure 14A
Scytalidium thermophilum CBH1 Genomic DNA

```

1   ATGCGTACCG CCAAGTTCGC CACCCTCGCC GCCCTTGTGG CCTCGGCCGC
    CGCCCAGCAG GCGTGCAGCC TCACCACCGA GAGGCACCCT TCCCTCTCCT
101  GGAAGAAGTG CACCGCCGGC GGCCAGTGCC AGACCGTCCA GGCTTCCATC
    ACTCTCGACT CCAACTGGCG CTGGACTCAC CAGGTGTCTG GCTCCACCAA
201  CTGCTACACG GGCAACGAGT GGGATTCTAG CATCTGCACT GATGCCAAGT
    CGTGCGCTCA GAACTGCTGC GTCGATGGTG CTGACTACAC CAGCACCTAT
301  GGCATCACCA CCAACGGTGA TTCCCTGAGC CTCAAGTTCG TCACCAAGGG
    CCAGTACTCG ACCAACGTCG GCTCGCGTAC CTACCTGATG GACGGCGAGG
401  ACAAGTATCA GACCTTCGAG CTCCTCGGCA ACGAGTTCAC CTTGATGTC
    GATGTCTCCA ACATCGGCTG CGGTCTCAAC GCGGCCCTGT ACTTCGTCTC
501  CATGGACGCC GATGGTGGTC TCAGCCGCTA TCCTGGCAAC AAGGCTGGTG
    CCAAGTACGG TACCGGCTAC TGCGATGCTC AGTGCCCCCG TGACATCAAG
601  TTCATCAACG GCGAGGCCAA CATTGAGGGC TGGACCGGCT CCACCAACGA
    CCCCACGCC GCGCGGGGCC GCTATGGTAC CTGCTGCTCT GAGATGGATA
701  TCTGGGAGGC CAACAACATG GCTACTGCCT TCACTCCTCA CCCTTGCACT
    ATCATTGGCC AGAGCCGCTG CGAGGGCGAC TCGTGCGGTG GCACCTACAG
801  CAACGACCGC TACGCCGGCG TCTGCGACCC CGATGGCTGC GACTTCAACG
    CGTATCGCCA GGGCAACAAG ACCTTCTACG GCAAGGGCAT GACCGTCGAC
901  ACCACCAAGA AGCTCACCGT CGTCACCCAG TTCCTCAAGG ACGCCAACGG
    CGATCTCGGC GAGATCAAGC GCTTCTACGT CCAGGATGGG AAGATCATCC
1001 CCAACTCCGA GTCCACCATC CCCGGCGTCG AGGGCAACTC CATCACCAG
    GATTGGTGCG ACCGCCAGAA GGTTCGCTTT GCGGACATTG ACGACTTCAA
1101 CCGCAAGGGC GGCATGAAGC AGATGGGCAA GGCCCTCGCC GGCCCCATGG
    TCCTGGTCAT GTCCATCTGG GATGACCACG CCTCCAACAT GCTCTGGCTC
1201 GACTCGACCT TCCCTGTCGA TGCCGCTGGC AAGCCCGGCG CCGAGCGCGG
    TGCTTGCCCG ACCACCTCGG GTGTCCCTGC TGAGGTTGAG GCCGAGGCCC
1301 CCAACAGCAA CGTCGTCTTC TCCAACATCC GCTTCGGCCC CATCGGCTCG
    ACCGTTGCCG GCCTTCCCAG CGATGGCGGC AACAACGGCG GCAACACCAC
1401 CGTCCAGCCC CCGCCCAGCA CCACCACCAC CTCTGCCAGC AGCAGCACCA
    CCTCGGCTCC TGCCACCACC ACCACCGCCA GCGCTGGCCC CAAGGCTGGC
1501 CGCTGGCAGC AGTGCGGCGG CATCGGCTTC ACTGGCCCGA CCCAGTGCGA
    GGAGCCCTAC ACTTGCACCA AGCTCAACGA CTGGTACTCT CAGTGCCTGT
1601 AA

```

Figure 14B
Scytalidium thermophilum CBH1 cDNA

1	MRTAKFATLAALVASAAAQQACSLTTERHPSLSWKKCTAGGQCQTVQASI	50
51	TLDSNWRWTHQVSGSTNCYTGNEWDSSICTDAKSCAQNCCVDGADYTSTY	100
101	GITTNGDSLKLVTKGQYSTNVGSRTYLMDGEDKYQTFELLGNEFTFDV	150
151	DVSNIGCGLNGALYFVSMDADGGLSRYPGNKAGAKYGTGYCDAQCPRIK	200
201	FINGEANIEGWTGSTNDPNAGAGRYGTCCSEMDIWEANNMATAFTPHPCT	250
251	IIGQSRCEGDSCGGTYSNDRYAGVCDPDGCDFNAYRQGNKTFYKGKMTVD	300
301	TTKKLTVVTQFLKDANGDLGEIKRFYVQDGKIIIPNSESTIPGVEGNSITQ	350
351	DWCDRQKVAFGDIDDFNRKGGMKQMGKALAGPMVLVMSIWDDHASNMLWL	400
401	DSTFPVDAAGKPGAERGACPTTSGVPAEVEAEAPNSNVVFSNIRFGPIGS	450
451	TVAGLPSDGGNNGGNTTVQPPPSTTTTSASSSTTSAPATTTTASAGPKAG	500
501	RWQQCGGIGFTGPTQCEEPYTCTKLNDWYSQCL-	534

Figure 14C

Scytalidium thermophilum CBHI, including signal sequence

Figure 15: Alignment of the mature protein sequences for *Hypocrea jecorina* CBH1, *Humicola grisea* var. *thermoidea* CBH1.1, and *Scytalidium thermophilum* CBH

Genencor <i>Hypocrea jecorina</i> Cel7A	1	75
<i>Humicola grisea</i> CBH1.1	(1)	QSACTLQSETHPPLTWQKSSGGTCTQQTGTVVIDANWRWTHATNSSTNCYDGN TW STLCPDNETCAKNCCLDG
<i>Scytalidium thermophilum</i> 69	(1)	QOACSLTTERHPSLSWNKCTAGGQCQTQVQASITLDSNRWTHQVSGSTNCYTGKWDTSICTDAKSCAQNCCVDG
Consensus	(1)	QOACSLTTERHPSLSWKKCTAGGQCQTQVQASITLDSNRWTHQVSGSTNCYTGNEWDSSICTDAKSCAQNCCVDG
Genencor <i>Hypocrea jecorina</i> Cel7A	76	150
<i>Humicola grisea</i> CBH1.1	(76)	AAYASTYGVTTSGNSLSIGFVTSQAQKN-VGARLYLMASDTTYQEFTLLGNEFSFDVDVSQPCGLNGALYFVSM
<i>Scytalidium thermophilum</i> 69	(76)	ADYTSTYGITTTNGDSLSLKFEVTKGQHSSTNVGSRTYLMDGEDKYQTFELLGNEFTFDVDVSNIGCGLNGALYFVSM
Consensus	(76)	ADYTSTYGITTTNGDSLSLKFEVTKGQYSTNVGSRTYLMDGEDKYQTFELLGNEFTFDVDVSNIGCGLNGALYFVSM
Genencor <i>Hypocrea jecorina</i> Cel7A	151	225
<i>Humicola grisea</i> CBH1.1	(150)	DADGGVSKYPTNTAGAKYGTGYCDSCQCPRLKFFINGQANVEGWEPSSNNANTGIGGHGCCSEMEDIWEANSISEA
<i>Scytalidium thermophilum</i> 69	(151)	DADGGLSRYPGNKAGAKYGTGYCDAQCPRIKFFINGEANIEGWTGSTNDPNAGAGRYGTCCSEMEDIWEANNMATA
Consensus	(151)	DADGGLSRYPGNKAGAKYGTGYCDAQCPRIKFFINGEANIEGWTGSTNDPNAGAGRYGTCCSEMEDIWEANNMATA
Genencor <i>Hypocrea jecorina</i> Cel7A	226	300
<i>Humicola grisea</i> CBH1.1	(225)	LTPHPCTTVGQEI ~~CE~~ GDGCGGTYSNDNRYGGTCDPDGCDWNPYRLGNTSFYGPSSFTLDTTKKLTVVVTQFETSG-
<i>Scytalidium thermophilum</i> 69	(226)	FTPHPCTTIIGSRCEGSDSCGGTYSNERYAGVCDPDGCDENSYRQGNKTFYGKG--MTVDTTKKITVVTQFLKDN
Consensus	(226)	FTPHPCTTIIGSRCEGSDSCGGTYSNDRYAGVCDPDGCDENSYRQGNKTFYGKG--MTVDTTKKLTVVTQFLKDN
Genencor <i>Hypocrea jecorina</i> Cel7A	301	375
<i>Humicola grisea</i> CBH1.1	(299)	----AINRYYVQNGVTFQQPNAEELGSYSGNELNDDYCTAEAEAEFGSS-FSDKGGLTQFKKATSGGMVLVMSLWD
<i>Scytalidium thermophilum</i> 69	(299)	GDLGEIKRFYVQDGKIIIPNSESTIPGVEGNSITQDWCDRQKVAFGDIIDFNKGGMKQMGKALAGPMVLVMSIWD
Consensus	(301)	GDLGEIKRFYVQDGKIIIPNSESTIPGVEGNSITQDWCDRQKVAFGDIIDFNKGGMKQMGKALAGPMVLVMSIWD
Genencor <i>Hypocrea jecorina</i> Cel7A	376	450
<i>Humicola grisea</i> CBH1.1	(369)	DYANMLWLDSTYPTNETSSTPGAVRGSCSTSSGVPQAQVESQSPNAKVTFFSNIKFGPIGSTGNPESGGNP-----
<i>Scytalidium thermophilum</i> 69	(374)	DHASNMLWLDSTFPVDAAG-KPGAERGACPTTSGVPAEVEAEAPNSNVVFSNIRFGPIGSTVAGLPGAG--NGGN
Consensus	(376)	DHASNMLWLDSTFPVDAAG-KPGAERGACPTTSGVPAEVEAEAPNSNVVFSNIRFGPIGSTVAGLPGAG NGGN
Genencor <i>Hypocrea jecorina</i> Cel7A	451	518
<i>Humicola grisea</i> CBH1.1	(438)	PGGNPPG-----TTTTRRPATTTGSSPGPTQSHYGQCGGIGYSGPTVCASGTTCCQVLNPPYYSQCL
<i>Scytalidium thermophilum</i> 69	(446)	NGGNPPP-----PTTTSSAPATTTTASAGPKAGRWQCCGGIGFTGPTQCEEPTCTCKLNDWYSQCL
Consensus	(448)	TTVQPPPPSTTTTSASSSTTSAPATTTTASAGPKAGRWQCCGGIGFTGPTQCEEPTCTCKLNDWYSQCL
	(451)	NGGNPPP PTTTSSAPATTTTASAGPKAGRWQCCGGIGFTGPTQCEEPTCTCKLNDWYSQCL





Figure 16

