



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2567749 C 2013/07/23

(11)(21) **2 567 749**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2005/05/26
(87) Date publication PCT/PCT Publication Date: 2005/12/08
(45) Date de délivrance/Issue Date: 2013/07/23
(85) Entrée phase nationale/National Entry: 2006/11/22
(86) N° demande PCT/PCT Application No.: IL 2005/000540
(87) N° publication PCT/PCT Publication No.: 2005/116252
(30) Priorité/Priority: 2004/05/26 (US60/574,210)

(51) Cl.Int./Int.Cl. *C12P 21/06* (2006.01),
G01N 33/00 (2006.01), *G06F 7/00* (2006.01)
(72) Inventeur/Inventor:
SMILANSKY, ZE'EV, IL
(73) Propriétaire/Owner:
ANIMA CELL METROLOGY, IL
(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : PROCEDE D'EVALUATION DE SEQUENCES DE RIBONUCLEOTIDES
(54) Title: METHODS FOR EVALUATING RIBONUCLEOTIDE SEQUENCES

(57) **Abrégé/Abstract:**

The present invention relates to methods for identifying ribonucleotide sequences, in vitro, using the ribosome-mediated translation.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 December 2005 (08.12.2005)

PCT

(10) International Publication Number
WO 2005/116252 A3

(51) International Patent Classification⁷: **C12P 21/06**,
G01N 33/00, G06F 7/00

(21) International Application Number:
PCT/IL2005/000540

(22) International Filing Date: 26 May 2005 (26.05.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/574,210 26 May 2004 (26.05.2004) US

(71) Applicant (for all designated States except US): **ANIMA CELL METROLOGY** [IL/IL]; 41 Moshav Meishar, D.N. Emek Sorek 76850 (IL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **SMILANSKY, Ze'ev** [IL/IL]; 41 Moshav Meishar, 76850 D.N. Emek Sorek (IL).

(74) Agent: **WEBB, Cynthia**; Webb & Associates, P.O. Box 2189, 76121 Rehovot (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,

GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
2 March 2006

(48) Date of publication of this corrected version:
6 July 2006

(15) Information about Correction:
see PCT Gazette No. 27/2006 of 6 July 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR EVALUATING RIBONUCLEOTIDE SEQUENCES

(57) Abstract: The present invention relates to methods for identifying ribonucleotide sequences, *in vitro*, using the ribosome-mediated translation.



WO 2005/116252 A3

METHODS FOR EVALUATING RIBONUCLEOTIDE SEQUENCES

FIELD OF THE INVENTION

5 The present invention relates to methods for identifying ribonucleotide sequences by monitoring ribosomal translation, *in vitro*.

BACKGROUND OF THE INVENTION

RNA reading has an important value in biological and pharmaceutical industries.
10 Identifying expression levels of multiple genes between various biological samples enables to perform genotyping, study disease pathways and obtain improved diagnosis and prognosis of diseases among other applications.

Since the early 1990s, the simultaneous measure of the expression of thousands of
15 different RNA gene products in a biological sample, such as a cell lysate, became feasible by the introduction of DNA microarrays (DNA chips). A DNA chip consists of numerous addressable locations. In each location numerous copies of a specific single strand DNA molecule (probes) are attached. When a sample containing a DNA strand that is complementary to one or more of the DNA molecules on the chip, hybridization
20 takes place. With appropriate sample labeling strategies, a pattern indicating the identity of the DNA strands and their amounts is obtained,. The chip with its large number of probes can identify, quantitate and compare the RNA sequences expressed in a set of samples (e.g. Nature Genetics, January 1999 Supplement).

25 The technology of DNA chips has several major drawbacks: design and production of chips is lengthy and expensive; assay performance takes several days and may include biased intermediate stages such as amplification; inaccurate quantitation and insensitivity to mRNA isoforms. The most prominent drawback of these methods is that they allow only partial analysis of gene products. In the case of oligo chips, only a
30 predetermined oligo sequence designed for that chip can be detected. In the case of cDNA chips, the content of the chip is obtained by 'trial and error' and hence gene coverage is not guaranteed. Moreover, any attempt to analyze hundreds of thousands of

RNA isoforms would result in impractical chip density which would not enable to distinguish between RNA variants and isoforms. In addition, commercial off the shelf chips usually encompass well-known, recognized genes and thus analysis is limited to identification of such already well-known genes. Non-adequate evaluation of expression
5 magnitude is another disadvantage of the commercial chips as it is common to have several spots that putatively cover the same gene, and show gross differences in expression estimates, sometimes of a factor of 3 or more.

A protein synthesis monitoring (also termed hereinafter "PSM") system and
10 methods of using same is disclosed by the inventor of the present invention in International Patent Application No. PCT/IL03/01011, Publication No. WO2004/050825. PSM includes a plurality of markers, each marker encompasses a pair of interacting labeling moieties, the first moiety being attached to a ribosome or a fragment thereof and the second moiety being
15 attached to one of the following entities: the ribosome or the fragment thereof, tRNA or amino acid. Protein synthesis in PSM is carried out by monitoring the signal sequences generated upon excitation of the markers. WO2004/050825 discloses that using the PSM system enables real-time monitoring of proteins synthesis *in vivo* and further allow identifying the amino-acid sequences of the protein being synthesized through database
20 interrogation process.

U.S. Patent No. 5,706,498 discloses a gene database retrieval system for retrieving gene sequences having a sequence similar to a sequence data from the gene database. The system is capable of storing the sequence data of genes whose structures or
25 sequences were analyzed and identified. The system includes a dynamic programming operation unit for determining the degree of similarity between target data and key data by utilizing the sequence data of the bases of the gene from the gene database as the target data and the sequence data of the bases as the key for retrieval, and further contains a central processing device unit for allowing access to the gene database in parallel to the
30 operation process for determining the degree of similarity. U.S. Patent No. 5,706,498 merely provides a database retrieval tool *in silico* but does not teach or even suggest identification of mRNA molecules in cellular systems.

U.S. Patent No. 5,856,928 discloses a system for characterizing and interpreting nucleotide and amino acid sequences. Natural numbers are assigned to represent DNA and mRNA nucleotide bases (n-numbers 0, 1, 2, 3), base pairing numbers in RNA (p-numbers 0, 1, 2, 3), and amino acids in protein (z-numbers with seventeen prime numbers and odd numbers 1, 25, 45; all smaller than 64). Gene and protein sequences may be represented, characterized and interpreted by their specific n-sums and z-sums. The system disclosed in US 5,856,928 is in fact a representational scheme facilitating computation and characterization of nucleotide and amino acid sequences *in silico*. This system cannot provide mRNA identification in cellular systems.

Nowhere in the background art is it taught or suggested that mRNA may be identified by utilizing the putative transcription activity. Moreover, there is an unmet need to measure RNA through its natural role, namely as a template for protein production, rather than through reverse transcription followed and/or hybridization techniques.

SUMMARY OF THE INVENTION

The present invention provides methods for identifying the ribonucleotide sequences of mRNA molecules. The method of the present invention is essentially different from any other method known in the art for mRNA identification as it uses the cellular translation mechanism which is carried out by ribosomes for identifying the RNA molecules being used by this mechanism as a template for protein translation. Thus, the method of the invention is devoid of the drawbacks characterizing the methods known in the art. Particularly, the methods of the invention do not require use of DNA chips and thus the entire stage of chip design is avoided. Moreover, RNA identification according to the methods of the present invention does not require reverse transcription, amplification or fluorescent labeling. Another advantage of RNA identification according to the methods of the present invention over methods known in the art is that the fundamental components of the PSM system, namely a solid substrate such as a microscope slide with ribosomes immobilized thereto and a cell-free translation system, are readily accessible and can be adjusted for any assay and biologic source.

In the disclosed method, RNA is fed into a cell-free translation system, also termed hereinafter PSM system, where ribosomes are immobilized on a solid substrate, for example, a microscope slide. PSM systems, disclosed in WO2004/050825, enables to perform a PSM assay which includes monitoring protein synthesis through detection of the signal sequences produced by fluorescent markers being attached to ribosomes or fragments thereof, amino acids and tRNAs. The signal sequences are generated in the PSM system upon excitation of the fluorescent markers. mRNA molecules that correspond to signal sequences obtained in the PSM assay may be identified by performing database interrogation in a database, termed hereinafter "PSM database". The PSM database is specifically designed for the method of the invention and comprises a plurality of signal sequences or the corresponding data streams wherein each signal sequence is assigned to a particular RNA molecule.

The method of the present invention is particularly advantageous over other methods known in the art for mRNA identification since mRNA is monitored substantially throughout its length, thus facilitating to distinguish between variants and isoforms of the same generic RNA molecule.

Moreover, using the methods of the present invention RNA molecules are detected at a resolution of a single RNA molecule, thereby the number of RNA molecules processed in the PSM assay can be calculated. Thus, the methods of the invention provide analysis on single molecule basis, hence providing an ultimate signal to noise ratios. Additionally, RNA identification according to the method of the present invention is carried out in real time substantially during elongation of the RNA molecules. Thus, using the method of the invention the RNA sequence enables discrimination between isoforms. In addition, the entire identification assay carried out by the method of the present invention is completed within hours rather than days.

According to one aspect, the present invention provides a method for RNA identification comprising:

- (a) providing a PSM system, wherein the PSM system comprises:
 - (i) at least one marker detectable through detection of electromagnetic radiation, the at least one marker

- comprising a pair of interacting labeling moieties, wherein the first moiety being bound to a ribosome or a fragment thereof, the ribosome or the fragment thereof being attached to a solid substrate, and the second moiety being bound to an entity selected from the group consisting of: the ribosome or the labeled fragment thereof, tRNA and amino acid, wherein the marker is capable of emitting electromagnetic radiation in response to translation activity;
- (ii) at least one translation component selected from the group consisting of: aminoacyl-tRNA synthetases, initiation factors, elongation factors, termination factors, energy sources and energy regenerating molecules; and
- (iii) detection means adapted to measure emitted radiation from the PSM system;
- (b) introducing at least one RNA molecule into the PSM system: and
- (c) detecting electromagnetic radiation signals obtained in response to translation activity.

According to one embodiment, the PSM system comprises a plurality of markers. According to another embodiment, the method comprises introducing a plurality of RNA molecules into the PSM system.

According to another embodiment, the solid substrate is selected from the group consisting of: glass, glass slide adapted for microscope means and a solid substrate having a mica surface.

According to yet another embodiment, the method further comprises identifying the ribonucleotide sequence of the at least one RNA molecule.

According to another embodiment, the step of identifying the ribonucleotide sequence of the at least one RNA molecule comprises:

performing PSM database interrogation, thereby assigning said at least one signal sequence to at least one particular RNA molecule.

According to yet another embodiment, identifying the ribonucleotide sequence of at least one RNA molecule further comprises storing the at least one signal sequence in a PSM database.

According to yet another embodiment, the step of performing PSM database interrogation comprises:

determining the probability of an RNA molecule in the PSM database to produce said at least one signal sequence ; and
selecting one or more RNA molecules having the highest probabilities thereby assigning the one or more RNA molecules to said at least one signal sequence.

According to yet another embodiment, the at least one signal sequence is composed of one or more values selected from the group consisting of: time, spatial coordinates, signal type and signal intensity.

According to yet another embodiment, signals are obtained by energy transfer between the pair of interacting labeling moieties. According to yet another embodiment, the signals are selected from the group consisting of: FRET signals, quenching signals and a fluorescent signals.

According to yet another embodiment, the marker comprises a label selected from the group consisting of: a fluorescent dye, a fluorescent amino acid, a fluorescent peptide or protein, a fluorescent nucleotide, a quantum dot, a luminescent substance, a donor-quencher pair and a fluorescent donor-acceptor pair. According to yet another embodiment, the second labeling moiety is a fluorescent amino acid.

According to yet another embodiment, the ribosomal fragment is selected from the group consisting of: ribosomal RNA, a ribosomal protein, ribosomal protein L1, ribosomal protein L11, ribosomal protein S1 and fragments thereof. According to an alternative embodiment, the ribosomal fragment is located near a ribosomal site selected

from the group consisting of: ribosomal A site, ribosomal P site, ribosomal E site, peptide exit channel site, L1 arm, and L7/L12 arm.

BRIEF DESCRIPTION OF THE DRAWINGS

5 The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of providing what is believed to be the most useful and readily understood
10 description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the invention, the description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

15

In the drawings:

FIGURE 1 describes a general embodiment of the system with TIR illumination and optical setup for double labeling.

FIGURE 2 describes a cartoon of FRET signal outputs from a doubly labeled
20 synthesis system.

FIGURE 3 describes the processing of an amino acid sequence into a PSM sequence for the case of double labeling.

DETAILED DESCRIPTION

25 The present invention provides a method for RNA identification, preferably by feeding the RNA into an *in-vitro* protein synthesis monitoring system, also termed hereinafter “PSM system”, and determining the identity of the RNA by monitoring translation activity at the PSM system.

30 The terms “RNA” and “mRNA” are interchangeably used herein to describe a ribonucleotide sequence that transfers genetic information to ribosomes, where it serves as a template for protein synthesis. A ribonucleotide sequence is a polymer of ribonucleic

acids, and is a constituent of all living cells and many viruses. It consists of a long, usually single-stranded chain of alternating phosphate and ribose units with the bases adenine, guanine, cytosine, and uracil bonded to the ribose. The structure and base sequence of RNA are determinants of protein synthesis and the transmission of genetic
5 information.

The term “*in-vitro*” as used herein, refers to cellular systems rather than *in silico* systems. In a particular embodiment, *in vitro* is used to define a cell-free system comprising ribosome(s) and components required for carrying out translation in the
10 ribosome(s) of the system. However, *in vitro* as used herein may also refer to a cellular system with the proviso that such system would enable RNA identification by PSM assay in accordance with the teaching of the present invention.

The term “translation activity” as used herein refers to any step during the
15 transition from mRNA to an amino acid or an amino acid sequence, including, but not limited to, mRNA-tRNA recognition and pairing (also known as codon-anti codon paring), amino acid activation (or tRNA aminoacylation), attachment of an amino acid to the tRNA and addition of the amino acid to a growing peptide chain.

20 The background art merely teaches how to identify RNA molecules by using its propensity for hybridization, and does not consider or suggest using its natural role as a template for protein synthesis. The background art suffers from severe disadvantages such as the need for elaborate chip design, ambiguous results, complex and lengthy result analysis, insensitivity to RNA isoforms, inaccurate quantitation and lengthy assay
25 preparation. A comprehensive review on DNA chips is provided in the supplemental issue of Nature published January 1999. Additional publications are available from commercial suppliers such as Affymetrix, Agilent, and Amersham Pharmacia.

In principle, the probes contained within DNA chips (or microarrays) are broadly
30 divided into two categories: oligonucleotides and cDNAs. In the first category, the strands are relatively short (20-70 base pairs). They are synthesized either directly on the chip or in a solution and subsequently printed. cDNA strands are usually longer, often

between 200-1000 base pairs, and they are produced by techniques of genetic engineering (or cloning) from living cells.

5 In oligo microarrays, the chip is carefully designed and the content of each spot is well defined. Because of the relatively short sequence length, several oligos are required to properly identify a gene. Identification of hybridized mRNAs is complicated in cases where one gene gives rise to several different mRNAs (as in the event of alternative splicing). Further, hybridization signals tend to fluctuate, yielding relatively low signal to noise. Due to these obstacles, the designing of an oligo microarray is a difficult, costly
10 and time-consuming process.

In cDNA microarrays, chip content commonly arises from a gene library. There is no assurance that the cDNAs on the chip cover all genes of interest, neither is there any assurance of uniqueness. Repetitive sequences, as well as sequences common to several gene families, can lead to cross-hybridization. In fact chip content is often unknown until
15 after the assay is analyzed, as cDNA sequencing is expensive and thus performed only for spots of interest. There are serious quality control problems involved with cDNA chips, such as variation in the concentration of the probes on a slide, incorrect identity of some probes due to mis-annotation, cross-hybridization of splice-variants, differences in the probe length, pin-to-pin variation and spot size variation.

20

Preparation of a DNA chip may last several weeks and even several months, depending on the type of chip, number of genes and the required precision. In the case of oligo chips, off-the-shelf chips or oligo libraries ready for printing can be purchased. Performance of the analysis usually takes a few days. The stages typically involve
25 harvesting total RNA from a single cell or from a plurality of cells; applying reverse transcription thereby obtaining cDNA; optionally amplifying the resulting products, either through bacterial growth or through PCR; fluorescent labeling and hybridization to the chip and finally the chip is scanned and the results analyzed.

30 The present invention overcomes the disadvantages of background art by providing a method for RNA identification, preferably by feeding the RNA into an in-vitro protein synthesis system, and determining the identity of the mRNA being translated by monitoring the synthesis process.

In order to better describe the disclosure, the technology of protein synthesis monitoring (PSM) will be now briefly explained. This technology with its applications is described in detail in WO2004/050825.

5 In PSM, use is made of the physical phenomenon of FRET (fluorescence resonance energy transfer). A donor fluorophore and a matching acceptor fluorophore are selected so that the emission spectrum of the donor overlaps the excitation spectrum of the acceptor. When donor and acceptor are in close proximity (usually less than 10 nm), excitation of the donor will cause emission from the acceptor, as some of the energy
10 passes from donor to acceptor via a quantum coupling effect. Thus, donor and acceptor serve as a proximity gauge: when they are near, a FRET signal can be generated, and when they are not, a FRET signal cannot be generated. In practice, a ribosome is engineered to carry a donor fluorophore, and tRNA and/or amino acids and/or some other part of the ribosome are either engineered to carry acceptor fluorophores or else their
15 natural fluorescent properties are utilized as acceptors. The PSM assay involves translation of a pool of RNA molecules *in vitro* or *in vivo* using ribosomes tRNA and amino acid molecules. In order to monitor the translation, a light source illuminates the ribosomes thus exciting the donor fluorophores and thereby the acceptor fluorophores whenever these components are in sufficient proximity to each other. The resulting
20 signals are detected and optionally stored. Using data base interrogation, the protein being synthesized is identified.

 According to WO2004/050825, PSM uses the FRET effect to monitor the synthesis process performed by a ribosome, and identify the protein being synthesized,
25 substantially in real time. Various labeling strategies can be used for PSM. In one preferred embodiment, the ribosome is labeled with donor and one or more amino acid species are labeled with acceptor fluorophore. According to certain embodiments, all amino acid species are labeled. When a labeled amino acid is processed by the ribosome, a FRET signal is generated. The resulting signal sequence corresponds to the loci of
30 labeled amino acids within the protein sequence, and therefore is a characteristic of this protein and enables its identification.

Other labeling strategies involve labeling tRNA rather than amino acids; labeling both donor and acceptor on selected parts of the ribosome; labeling with more than one type of FRET pair; labeling all amino acids and /or all tRNAs; and many other variants of the basic idea. Some of these variants are discussed below while many others will be
5 evident to anyone skilled in the appropriate art.

Thus, according to one aspect the present invention provides a method for RNA identification the method comprising:

- (a) providing a PSM system;
- 10 (b) introducing at least one RNA molecule into the PSM system; and
- (c) detecting electromagnetic radiation signals obtained in response to translation activity; and optionally
- (d) identifying the ribonucleotide sequence of the at least one RNA molecule.

15 One exemplary flow of operation according to the aspect of the invention is now described. An optical apparatus monitors the protein synthesis system (PSM), described hereinabove and in WO2004/050825, optionally by directing electromagnetic radiation of the required wavelength and energy onto the marked system, thereby exciting the donor fluorophores. The acceptor fluorophores on the tRNAs and/or amino acids and/or
20 on the ribosome respond to this energy with the FRET signal whenever a donor and acceptor pair are in sufficient proximity, indicative of particular steps of translation activity, for example, indicating the incorporation of a particular tRNA or amino acid by said synthesis system. Fluorescence radiation emitted from acceptor fluorophores is detected by the optical apparatus and the event is recorded by the image acquisition
25 device connected to a computerized analysis unit. The acquired image sequence is then analyzed by the software for the purpose of identifying the mRNA(s) associated with the recorded translation activity.

Analysis includes separation of the initial signals (for example as a stream of
30 video frames) into a set of signal sequences, wherein each signal sequence is emitted from a single ribosome. Each ribosomal signal sequence is separately analyzed to identify the mRNA that served as a template for its synthesis. Alternatively, the mRNA that served as a template for the recorded synthesis is analyzed directly by interrogating

the signal sequence in a PSM databases which includes signal sequence data associated with RNA rather than with proteins.

Optionally, a signal sequence data obtained from the PSM assay is stored in the PSM database. The correspondence between the data in the PSM database and the RNA molecules assigned thereto depends on the labeling strategy used in the PSM assay. Several labeling strategies are detailed below. The resulting sequence of detection events is a characteristic of the RNA used as a template for the protein synthesis. This signal sequence may be described as a bit stream with zeroes and ones (as commonly used to describe data in computer science terminology). This stream may contain some uncertainty as to the number of bits in each field, as well as other elements of uncertainty.

The signal is subsequently used to interrogate a database of signals computed *in silico* from a relevant database of protein or nucleotide sequences. The method disclosed herein uses this signal to identify the mRNA that most likely produced the signal. In the event that for a particular RNA molecule only part of the elongation and/or translation event is recorded, then the resulting signal sequence does not correspond directly to the RNA. However, the PSM database includes information relating to the labeling strategy and to various translation events and therefore can assign any signal sequence to an RNA molecule even if the signal sequence does not include record of the entire translation event.

In the present invention, ribosomes are immobilized on a solid substrate. Numerous solid substrates are suitable for carrying out the method of the invention providing that the surface enables to immobilize ribosomes thereto and allows the immobilized ribosome to execute translation activity in the cell-free translation system. Solid substrates that may be utilized for immobilizing ribosomes are known in the art (see for example Ha, T, *Single-molecule fluorescence resonance energy transfer*, METHODS 25, 78-86 (2001)).

According to certain embodiments, the solid surface is adapted for use under a microscope, thereby enabling to monitor the translation activity through the microscope.

Preferably, the microscope as well as any monitoring means that is utilized in the context of the present invention is adapted for detecting single molecule fluorescence. According to some embodiment, the detected activity is stored.

5 Preferably, ribosomes are immobilized at a density that permits optical resolution at the level of a single ribosome. The ribosomes are labeled with one or more types of donor fluorophores, and either some tRNA species, or some amino acid species, or another part of that ribosome, or a combination thereof, labeled with acceptor fluorophores that match the donor(s). The donor-acceptor pair is also referred herein as
10 the first and second labeling moieties of the marker used in the PSM assay. In other preferred embodiments assignment of donors and acceptors to molecules could be switched. Next, the sample containing the mixture of mRNA molecules to be assayed is introduced with the other components required for performing translation in a cell-free system including tRNAs, labeled and unlabeled; synthetases; initiation, elongation and
15 termination factors energy sources (ATP, GTP), energy regenerating systems (creatine phosphate and creatine phosphokinase for eukaryotic systems (Shimizu *et al.*, *Cell-free translation reconstituted with purified components*. Nat Biotechnol. 2001, 19(8):751-5). The entire translation system is placed under a microscope equipped for single molecule detection, such as instruments available from Zeiss (Oberkochen, Germany) and Leica
20 (Wetzlar, Germany), with an image acquisition device operable at a sufficient rate (10-100 frames per second), and computational units that can acquire and analyze the resulting images and data.

 The marker optionally comprises at least one photoactive component. The emitted
25 electromagnetic radiation is detected and can be analyzed to identify and measure RNA molecules in a sample mixture. The procedure can optionally be performed simultaneously for hundreds, thousands and even millions of single ribosomes.

 The method of the invention may be carried out in accordance with the following
30 alternatives:

tRNA labeling. In this embodiment the acceptor fluorophore or fluorophores are attached to one or more species of tRNA. Methods of

labeling tRNA are discussed in detail in WO2004/050825.

Amino acid labeling. Amino acid labeling is preferable for measuring
5 RNA *in vitro* (more preferable than it is for *in-vivo* applications of PSM)
since proper structure, folding and functionality of the synthesized protein is
not crucial. In addition, amino acid labeling is relatively straightforward.

Double labeling. According to an alternative embodiment, more than
10 one FRET pair is utilized thereby distinct tRNAs or amino acids being
attached to distinct labeling moieties. For example, two donors with distinct
emission spectra are placed on one ribosome, and matching fluorescent
acceptors can be attached to tRNAs or to amino acids. For example, an
arginine tRNA is labeled with an acceptor of one color, and a lysine tRNA is
15 labeled with an acceptor of a different color. The appropriate pair of donors is
attached to, for example, ribosomal proteins L1 and S1, respectively. This
scheme is beneficial even if only two out of the 20 amino acids are labeled,
since the identification of the mRNA being translated in this setting depends
on the interspersed profile of one amino acid relative to the other, rather than
20 on the timing of the FRET signals. This removes the dependence of the
identification process on temporal aspects of the synthesis, such as codon
bias, variable tRNA species abundance, effects of RNA secondary structure
(Pelletier J, Sonenberg N. The involvement of mRNA secondary structure in
protein synthesis. *Biochem Cell Biol.* 1987, 65:576-581), ribosome pausing
25 (Wolin SL and Walter P., *EMBO J.* 1988, 7:3559-3569), among other aspects
of synthesis. In another preferred embodiment, the two spectrally distinct
acceptors are selected to respond to a single donor.

Full sequence labeling. This scheme calls for labeling the ribosome
30 with one, two or more donors with distinct emission spectra, and labeling the
entire set of amino acids and/or tRNAs with acceptors of corresponding
excitation spectra. This strategy is in contrast to labeling with only one FRET
pair, where one subset of amino acids is labeled and the complementary

subset remains unlabeled, in order to produce an informative signal sequence. Such full sequence labeling allows the sequence of amino acids to be read in its entirety, obviating the need for estimating the numbers of consecutive unlabeled events (see WO2004/050825 for section entitled “Data
5 interpretation simulation”). As in the strategy of double labeling described above, full sequence labeling is independent from temporal aspects of the synthesis. Full sequence labeling also offers important additional advantages, such as the possibility of identifying mutations, and that of identifying mRNAs from a given organism based on an mRNA database of another
10 (similar) organism, based on sequence matching.

Identifying the ribonucleotide sequence of the at least one RNA molecule, using PSM assay according to the principles of the present invention, comprises:
performing PSM database interrogation, thereby retrieving one or more PSM signal
15 sequences corresponding to one or more ribonucleotide sequences from the PSM database that conforms with said at least one signal sequence, thereby assigning said at least one signal sequence to at least one particular RNA molecule.

The identification process may further includes, prior to database interrogation any
20 one of the following steps:

- a) transferring the signals to a computerized analysis station; and/or
- b) clustering said signals into a list of signal sequences, the list comprises at least one signal sequence wherein the at least one signal sequence corresponds to signals obtained from a single ribosome; and/or
- 25 c) transforming the at least one signal sequence into at least one data stream.

In order to identify a proteins or RNA molecules from PSM signal sequence, the signal sequence data obtained during PSM assay must be processed and compiled to enable such identification. In order to allow efficient and accurate identification, the
30 following general three steps must be taken: (i) modeling of the data obtained from the PSM process; (ii) construction of a scoring function; and (iii) performance of efficient and accurate classification of the modeled data. The term “scoring function” as used herein refers to an estimate of the probability that a given signal sequence obtained from a

PSM assay corresponds to a particular ribonucleotide sequence or subsequence. This probability is in effect the probability that a specific ribonucleotide sequence or subsequence, the details of which are stored in the PSM database, would yield a signal sequence that is substantially identical to said signal sequence obtained from the PSM assay. A value obtained from the scoring function is not necessarily identical to the probability value. In fact, it may be difficult or even impossible to compute said probability value with sufficient precision. However, for the purpose of prioritizing and/or determining the best RNA candidates, stored in the database, that are the plausible generators of the measured signal, the value yielded by the scoring function may provide sufficient information.

I. Modeling the data obtained from the PSM process

In the modeling step, the physical and chemical events leading to the production of the PSM signal sequence are modeled, as precisely as possible, by computerized algorithms. Such an algorithm produces, from a ribonucleotide sequence (or a protein sequence), an expected PSM signal sequence that is stored in the PSM database.

In modeling the PSM process, as much as possible information about the factors effecting the production of the PSM signals needs to be known. The most important factors are labeling strategy and translation mechanism characteristics. Accordingly, a signal sequence is commonly of the form $S = (t_1, x_1, y_1, s_1), (t_2, x_2, y_2, s_2), \dots, (t_n, x_n, y_n, s_n), \dots$. Where t_i denotes a timing value, x_i and y_i denote image coordinates, and s_i denotes signal type or intensity (of both donor and acceptor). A signal sequence obtained during PSM assay is recorded and forwarded to a software module (also termed a "sequence analyzer") which transforms the sequence s into one or more data-stream of FRET on/off signals, as described in detail in WO2004/050825.

Preferably, every stretch of length K of a protein or an RNA sequence is modeled, where the number K depends on the labeling strategy used in the PSM assay and additional factors. If, for example, $K=100$ then an mRNA with 300 codons will have 200 entries in the database, one for each subsequence of length 100. The reason for this indexing is that it is impossible to determine that the synthesis process is recorded right

from its initiation. However, the modeling algorithm is directed to identify an mRNA even if the ribosome is monitored only during part of the translation period. The following labeling strategies are used in a PSM assay of the present invention:

- 5 1. Labeling of all amino acids with 20 distinct colors.

Though this may introduce technical complexity it is relatively easy to analyze the resulting signal *in silico* using bioinformatic techniques. The mRNA sequence is identical to the labeled sequence, and no modeling is required. The scoring function can be simply based upon the familiar BLAST algorithm.

10

2. Labeling of all amino acids or tRNAs with 2 distinct colors

This case is very similar to the previous one, except that instead of 20 or more colors we obtain a binary sequence with only two types of colors. In the modelling, the same process is repeated, transforming the sequence into a binary sequence. Again, the
15 BLAST algorithm can be used for scoring.

3. Double labeling of two amino acids or tRNAs with 2 distinct colors

This case is very similar to the previous two, the difference being that most amino acids (or tRNAs) are disregarded, leaving a labeled binary subsequence of the original,
20 full sequence. In the modelling, the same process is repeated, transforming the sequence into a binary sequence with only the labeled tRNAs accounted for. Again, the BLAST algorithm can be used for scoring.

4. Single labeling of several amino acids or tRNAs with a single color

25 In this scenario, only a single label is used. Thus, the amino acid (or tRNA) sequence is partitioned into two types – labeled versus unlabeled. The signals obtained are interspersed with “no signals”. Thus, the data stream of PSM signals measured from a single ribosome consists of a list of temporal data, indicating the precise time measurement at signal detection: t1, t2, t3, and so on. We can call this time sequence the
30 timing fingerprint of an mRNA. In the database, we need to be able to compute the timing fingerprint for each mRNA in the database. For this, we need to measure the timing fingerprints for a sufficient number of mRNA species, and from these

measurements deduce the correct model. For example, we can measure the timing fingerprints for mRNAs which are polymers of a single codon, or a single tRNA, and thereby deduce the average and standard deviation of timing for incorporation of that particular tRNA or amino acid. Once the model is available, we can compile a database
5 of timing fingerprints for each of the mRNAs in the PSM mRNA database .

In this setting, the scoring function that should be used is based on the description above, where the comparison between a measured timing fingerprint and a candidate timing fingerprint from the database is based on evaluation of the probability that the
10 candidate mRNA actually yielded the measured timing fingerprint, based on the statistical measurements made while constructing the database.

II. Construction of a scoring function

Upon production of a set of expected sequences corresponding to a set of mRNA
15 molecules of interest and storage of this set, a scoring function is applied. The scoring function provides the probability (p -value) for a particular signal sequence stored in the PSM database to yield the signal sequence measure by the PSM assay. Thus, for each entry in the PSM database, the probability of matching to the measured PSM signal sequence obtained during PSM assay is computed. As a result of such computation, the
20 most probable matches are selected. The signal sequence obtained during PSM assay is assigned to the RNA molecules corresponding to the most probable matches.

For the purpose of scoring, the degree of similarity between the signals produced during the PSM assay and the signals stored in the PSM database can be calculated using the system disclosed in US Patent No. 5,706,498. This patent discloses a database
25 retrieval system adapted for determining the degree of similarity between target data and key data.

III. Performance of efficient and accurate classification of the modeled data.

The database and scoring function described above, need to be efficiently arranged in order to enable quick and efficient analysis. Database arrangements are
30 discussed in detail in numerous prior art documents, such as for example Donald E. Knuth, The Art of Computer Programming, Addison-Wesley. US Patent No. 6,189,013

discloses a relational database system for storing biomolecular sequence information in a manner that allows sequences to be catalogued and searched according to association with one or more projects for obtaining full-length biomolecular sequences from shorter sequences.

5

Numerous applications of RNA measurement are currently practiced while others are attempted or will become practicable in the next few years. Reviews of current and foreseen applications for RNA reading devices can be found in the following market research reports: DNA Microarrays and Their Materials, published in January 2004 by
10 Business Communications Company Inc of Norwalk, CT; DNA Probes-based Diagnostics, published in January 2002 by Global Industry Analysts, Inc, of San Jose, Ca. ; Outlook for DNA Microarrays: Emerging Applications and Insights on Optimizing Microarray Studies, published in January 2002 by the Cambridge Healthtec Institute of Newton Upper Falls, MA . In addition, discussion of major applications can be found in
15 the scientific literature. For example, use of DNA chips for diagnosis and prognosis is discussed in Ken Garber, Science, 303:1754-1755 , 2004; use of DNA chips for SNP analysis is discussed in Erdogan F, *et al.*, Nucleic Acids Res. 2001, 29:E36. The term “SNP” refers to a single nucleotide polymorphism, a mutation involving a single base.

20 Illustrative, optional methods for monitoring protein synthesis are now described in detail below. The method will be described in selected variants and applications by way of an example, and it should be recognized that the illustrated embodiments should not be taken as a limitation on the scope of the disclosure. Some, but not all of the variants of the invention not mentioned here are: using a different type of microscope or
25 illumination; using a different fluorescent labeling scheme or strategy; Attaching fluorophores to other ribosomal proteins or locations, or using alternative labeling protocols; using other types of fluorophores, such as organic dyes, fluorescent proteins such as GFP or any of its variants; using a different data analysis method and system; using a different substrate for ribosome immobilization or a different strategy for viewing
30 functioning ribosomes.

EXAMPLES

Example 1: RNA identification with two donors and amino-acid labeling

Figure 1 describes one optional but preferred embodiment for an exemplary
5 apparatus for data acquisition, based on a wide-field microscope equipped with TIR
(Total internal reflection, a microscopy illumination method that illuminates a volume at
the interface of two materials with different refractive indices) illumination and
intensified CCD camera. This setup is useful for *in-vitro* single-molecule protein
synthesis monitoring application, where the ribosomes are immobilized on the
10 microscope slide.

In this preferred embodiment the ribosome is labeled with one or two quantum
dots as fluorescent donors. A ribosomal protein, such as optionally and preferably L1, is
first biotinylated using FluoReporter Biotin-XX protein labeling kit (Molecular Probes,
15 cat# F-2610) according to manufacturer protocol. Then, the biotinylated protein is linked
to Qdot™ 525 Streptavidin Conjugate (QuantumDot) according to manufacturer
protocol. In a similar way, a second ribosomal protein, such as optionally and preferably
S1, is labeled with Qdot™ 605 Streptavidin Conjugate. Then, a fluorescent acceptor with
excitation maximum near 525nm is attached to one amino acid, in this example lysine,
20 and a fluorescent acceptor with excitation maximum near 605nm is attached to the other
amino acid, in this example arginine (see below for techniques of amino acid labeling).
For the convenience of the ensuing discussion the appearance of the emission from the
lysine acceptor is termed “green” and the emission of the arginine acceptor is termed
“red”.

25

The two-color labeling scheme has an advantage over one-color labeling, even if
only two amino acids are labeled as the identification of the mRNA being translated
depends on the interspersed profile of one amino acid relative to the other, and not solely
on the temporal sequence of events. This is explained in more detail below. Double
30 labeling also removes the dependence on temporal aspects of synthesis that affect the
synthesis rate, as discussed above.

Referring to Figure 1, laser 100 is a diode-pumped doubled YAG laser (Crystalaser, Reno, NV) that can excite a wide range of dyes. Laser illumination 102 travels through a dichroic mirror 104 (Chroma Technology, Brattleboro, Vermont) and into a 'dove prism 106 such as a small Pellin Broca prism (CVI laser, <http://www.cvilaser.com/>) where the illumination undergoes TIR. The prism is optically coupled to the fused silica bottom of the sample chamber 108, so that evanescent waves illuminate up to 150 nm above the surface of the fused silica, thereby illuminating the immobilized ribosomes. The emitted fluorescence signals (both donors and acceptors) pass through objective 110 (Olympus, DPLanApo 100UV 1.3oil, or PLAPO60XO, Plan APO 60X oil immersion, NA=1.4 working distance=0.15 mm), through a dichroic splitting filter 111 (Chroma Technology, Brattleboro, Vermont) which splits the image into green (112,114,118) and red (113, 115, 119) channels. In each of the channels the emitted signals pass through fluorescent filters (112 and 113), through imaging lenses (114 and 115) into intensified CCD (ICCD) cameras 118 and 119 such as Cascade:512B available from Roper Scientific Photometrics. The readout from both cameras is transferred digitally to computer system 120 for image analysis, signal processing and subsequent identification of the mRNA being translated.

The ribosomes are immobilized on the microscope slide. The slide may have a mica surface (Novascan Techniques™) which is transparent and flat on a molecular size scale. Ribosomes, either labeled or unlabeled, undergo binding to mica in a few seconds, allowing the detection of single fluorescence images in aqueous buffer. A large excess of ribosomes and a short incubation period are employed for single molecule detection. Mica-bound ribosomes retain their activities, as shown in Sytnik et al., J. Mol. Biol. (1999), 285, 49-54, where detailed protocols are provided. Ribosomes can also be immobilized on surface treated glass slides. Ribosomes should be immobilized at a density that permits optical resolution of single ribosomes. For example, a minimal distance of 1-2 microns could be kept between ribosomes.

As a ribosome processes the mRNA template, electromagnetic radiation of the appropriate wavelength is used to simultaneously excite the two types of donors. This is possible since quantum dots have a wide excitation range, but narrow and specific emission. Emission from the two types of acceptors is detected separately using double

detection channels, each with its own spectral filter. The resulting signal sequence consists of green signals, red signals, and no signals. In applications of PSM that involve only one type of acceptor fluorophore, the exact timing of signal detection is important to obtain the identification of the mRNA. Specific algorithms are used in that case to
 5 estimate the numbers of unlabeled tRNAs or amino acids that were processed between two consecutive labeled events (see WO2004/050825).

In a preferred embodiment mRNA identification is based on the interspersion profile of red and green signals. In Figure 2 an example is shown, where a dark disc **210**
 10 corresponds to a green signal (lysine) and an bright disc **220** to a red signal (arginine). The events are shown as they appear on temporal axis **230**. This signal is translated into the code **grggrrrgrrgg** indicating the appearance order of green and red events. This code is used to search in an mRNA database precompiled for such searches. In Figure 3, amino acid sequences **301** and **302** of two distinct mRNA templates are shown, with
 15 lysine (K) and arginine (R) underlined. The corresponding subsequences of lysines and arginines are shown in **303** and **304**. It is clear that the code **grggrrrgrrgg** is compatible with sequence **303** and not with sequence **304**, and therefore matches mRNA **301** but not **302**. This approach serves as a basis for construction of an algorithm for mRNA identification.

20

Example 2: tRNA labeling and usage.

In this preferred embodiment the acceptor fluorophore is attached to one or more species of tRNA. Methods of labeling tRNA are discussed in detail in WO2004/050825. The labeled tRNAs are known to be processed normally both by
 25 ribosome and by the cognate synthetases.

Ribosome labeling is performed using one of several preferred methods. These include labeling with naturally fluorescent proteins, with organic dyes, and with semiconductor quantum dots. Labeling strategies included labeling ribosomal proteins
 30 such as ribosomal proteins L1, S1, S21 and others; In addition, 3' and 5' ends of 5S, 16S and 23S rRNA have been labeled (Robbins and Hardesty, Biochemistry. 1983, 22; 22(24):5675-5679).

For *in-vitro* labeling, there are several strategies. Organic dyes can be used to label ribosomal proteins using standard protein labeling techniques. Suppliers of these dyes publish detailed protocols describing their use. General procedures label proteins
5 through their amino groups (lysine). Other procedures target cysteines which are sometimes available for precisely located labeling. In this way, ribosomal proteins S1 and S8 were labeled by coumarin (Bakin *et al.*, J Mol Biol., 1991 221:441-453), and ribosomal proteins were tagged with fluorescein attached to a cysteine residue (Odom *et al.*, Biochemistry, 1990, 29:10734-10744).

10

A novel labeling strategy uses quantum dots, which are commercially available pre-conjugated with biotin or streptavidin. In such cases, proteins can be labeled with biotin, so the streptavidin conjugated Qdots (commercially available as QdotTM525/ QdotTM 565/ QdotTM 605/ QdotTM 655 streptavidin conjugated quantum dots, from
15 Quantum Dot Corporation, Hayward, Ca, USA; the numbers relate to maximal emission wavelength, nm) bind specifically to them. One useful method of generating a biotin-labeled protein involves creating a fusion protein between the protein of choice and biotin carboxyl carrier protein (BCCP). In the fusion protein, the original protein sequence is fused optionally to the last 87 (or 110) codons of the E. coli BCCP. When the
20 fusion protein is translated, it has the biotin tag attached to it and binds specifically to streptavidin (cf. Surrey *et al.*, Proc. Natl. Acad. Sci. USA, 95: 4293-4298, 1998).

Another method that is useful for labeling of ribosomal proteins is the well known strategy of fusing the protein of choice with a naturally fluorescent protein, such
25 as green fluorescent protein, yellow/cyan/blue fluorescent proteins or any other naturally fluorescent protein. An example where L1 was labeled by fusing it with a naturally fluorescent protein is described in (Mascarenhas *et al.*, EMBO Rep. 2001, 2:685-689). Numerous additional strategies for ribosome labeling were tested and others are clearly suitable for use with the present invention.

30

For the cell-free translation system, it is also possible to use cross systems. For example, an E. Coli translation system can be used for producing mammalian proteins. Alternatively, a mammalian translation system can be used. Sometimes it is beneficial to

switch some system elements. For example, using some E. Coli tRNAs in a mammalian system can allow specific labeling of tRNAs that have appropriately modified bases. Several E. Coli tRNAs have the uridine in position 8 modified to thiouridine. This makes such tRNA types suitable for attaching a fluorescent label. A complete database of tRNA
5 sequences can be found online . A
database of known RNA modifications can be found online

10 **Example 3: Amino acid labeling.**

In contrast with *in-vivo* applications of PSM, for RNA identification amino-acid labeling could be the method of choice since the functionality of the protein is not crucial, and since amino acids can be labeled relatively easily.

15 There are several ways of fluorescent labeling for amino acids in a cell free translation system. One way is to introduce labeled amino acids into the system. In this case one has to verify that the cognate synthetases function normally for charging the tRNAs with the labeled amino acid. For this step, it may be required to optimize the structure of the synthetases. Methods for high-throughput adaptation of synthetases for
20 unnatural or labeled amino acids are discussed for example in Santoro *et al.*, Nat. Biotechnol., 20:1044-1048 2002. In such cases it is important that no unlabeled amino acid of this type exist in the system, to avoid confounding the PSM signal sequence. To achieve this, a synthetic construction of the cell free translation system is preferably prepared as described in (Shimizu *et al.*, Nat Biotechnol. 2001, 19:751-755), since the
25 components of the translation system are introduced in a controlled manner.

A second way of introducing labeled amino acids is by fluorescent labeling of acylated tRNA. In other words, what is being labeled is the charged tRNA with its amino acid. Fluorescent labeled acylated tRNA can be purchased, for example the
30 FluoroTect™ GreenLys in vitro Translation Labeling System available from Promega Corporation (Madison, WI). It is important in this system also that no unlabeled amino acids exist to avoid confounding the PSM signal sequence. One way to ensure this is by

removing the cognate synthetases of labeled amino acids. This will ensure that recharging of tRNA does not occur, and only the labeled tRNA that was introduced will be used.

5 **Example 4: Additional PSM strategies**

In one preferred embodiment, a quenching strategy is used instead of FRET, as noted previously. Accordingly, instead of a fluorescent donor and acceptor, there is a fluorescent donor and an acceptor quencher that captures the donor energy without emission. Donor fluorescence is detected as long as the quencher is not sufficiently near
10 to the donor. Use of this strategy in PSM, generates a signal of donor fluorescence intermitted by periods of quenching.

In another preferred embodiment, a combination of methods is used. For example, some tRNAs and some amino acids may be labeled as acceptors, either with distinct or
15 with indistinguishable optical characteristics. Preferably, the amino acids are labeled with fluorescent labels that are distinct from tRNA labels. The signal analysis system accepts both signals and uses both in order to identify more confidently the mRNA being translated. It is obvious to anyone skilled in the art of single molecule detection and analysis that this is just one example of a wide variety of methods that can be derived
20 from this particular example.

In view of the large number of possible applications and embodiments of the present disclosure it should be recognized that the illustrated embodiments are only particular examples and should not be taken as a limitation on the scope of the disclosure.
25 Some of the possible additional applications which are clearly enabled by the present invention are clinical applications, diagnostic applications, production of food, cosmetics, and other bioproducts, military applications concerning biological warfare, and many more.

30 Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such

WO 2005/116252

PCT/IL2005/000540

alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims.

5

Citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention.

10

WE CLAIM:

1. A method for RNA identification the method comprising: (a) providing a protein synthesis monitoring (PSM) system, wherein the PSM system comprises: (i) at least one marker detectable through detection of electromagnetic radiation, the at least one marker comprising a pair of interacting labeling moieties, wherein the first moiety being bound to a ribosome or a fragment thereof, the ribosome or the fragment thereof being attached to a solid substrate, and the second moiety being bound to an entity selected from the group consisting of: the ribosome or the labeled fragment thereof, tRNA and amino acid, wherein the marker is capable of emitting electromagnetic radiation in response to translation activity; (ii) at least one translation component selected from the group consisting of: aminoacyl-tRNA synthetases, initiation factors, elongation factors, termination factors, energy sources and energy regenerating molecules; and (iii) detection means adapted to measure emitted radiation from the PSM system; (b) introducing at least one RNA molecule into the PSM system; (c) detecting electromagnetic radiation signals obtained in response to translation activity; and identifying the ribonucleotide sequence of the at least one RNA molecule by performing PSM database interrogation, thereby assigning at least one signal sequence to at least one particular RNA.
2. The method of claim 1, comprising introducing a plurality of RNA molecules into the PSM system.
3. The method according to claim 1, wherein the solid substrate is selected from the group consisting of: glass, glass slide adapted for microscope means and a solid substrate having a mica surface.
4. The method according to claim 1, wherein identifying the ribonucleotide sequence of at least one RNA molecule further comprises storing the at least one signal sequence in a PSM database.
5. The method according to claim 1, wherein the step of performing PSM database interrogation comprises: determining the probability of an RNA molecule in the PSM database to generate said at least one signal sequence; and selecting one or more RNA molecules having the highest scoring function value, thereby assigning the one or more RNA molecules to said at least one signal sequence.
6. The method according to claim 1, wherein the at least one signal sequence is composed of one or more values selected from the group consisting of: time, spatial coordinates, signal type and signal intensity.
7. The method according to claim 1, wherein the signals are obtained by energy transfer between the pair of interacting labeling moieties.
8. The method according to claim 7, wherein the signals are selected from the group consisting of: FRET signals, quenching signals and a fluorescent signal.
9. The method according to claim 1, wherein the marker comprises a label selected from the group consisting of: a fluorescent dye, a fluorescent amino acid, a fluorescent

peptide or protein, a fluorescent nucleotide, a quantum dot, a luminescent substance, a donor-quencher pair and a fluorescent donor- acceptor pair.

10. The method of claim 1, wherein the second labeling moiety is a fluorescent amino acid.
11. The method according to claim 1, wherein the ribosomal fragment is selected from the group consisting of: ribosomal RNA, a ribosomal protein, ribosomal protein L1, ribosomal protein L11, ribosomal protein S1 and fragments thereof.
12. The method according to claim 11, wherein According to an alternative embodiment, the ribosomal fragment is located near a ribosomal site selected from the group consisting of: ribosomal A site, ribosomal P site, ribosomal E site, peptide exit channel site, L1 arm, and L7/L12 arm.
13. The method according to claim 1, wherein the PSM system comprises: a plurality of markers detectable through detection of electromagnetic radiation, wherein each marker comprises a pair of interacting labeling moieties, wherein the first moiety being bound to a ribosome or a labeled fragment thereof, the ribosome or the fragment thereof being attached to a solid substrate, and the second moiety being bound to an entity selected from the group consisting of: the ribosome or the fragment thereof, tRNA and amino acid, wherein the marker is capable of emitting electromagnetic radiation in response to translation activity.
14. The method according to claim 13, wherein distinct tRNAs or amino acids are bound to distinct markers.
15. The method of claim 13, the marker being one of two distinct colors.

1/2




FIGURE 1

2/2




FIGURE 2

301. VGSDKCTTIHYNMCNRSSCMGQLKKWVDSTPKPPGT
RVRKAMAIYQSRRQHMTEVSVTKCTYSPALNVQLVK
 ESGGGRLVQPGGSLR

302. SCAASGYTFTNYGMNWVRQQDQPMREEEEVETFAFQA
 EIAQLMSLIINTFYSNKEIFLRENSSDALDKTIAKSG
 TKAFME

303. KRKKKRRKRRKK

304. RKRRKKK

FIGURE 3