

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 June 2010 (17.06.2010)

(10) International Publication Number
WO 2010/068452 A1

PCT

(51) International Patent Classification:

C07D 401/12 (2006.01) **A61K 31/47** (2006.01)
C07D 471/04 (2006.01) **A61P 29/00** (2006.01)

(21) International Application Number:

PCT/US2009/065752

(22) International Filing Date:

24 November 2009 (24.11.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/117,880 25 November 2008 (25.11.2008) US

(71) Applicant (for all designated States except US):
JANSSEN PHARMACEUTICA NV [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BREITENBUCHER, J., Guy** [US/US]; 2433 North Summit Circle Glen, Escondido, CA 92026 (US). **KEITH, John, M.** [US/US]; 16724 Alondra Drive, San Diego, CA 92128 (US). **TICHENOR, Mark, S.** [US/US]; 4065 Porte La Paz 156, San Diego, CA 92122 (US). **CHAMBERS, Alison, L.** [US/US]; 5235 Fiore Terrace C312, San Diego, CA 92122 (US). **JONES, William, M.** [US/US]; 8520 Costa Verde Blvd 3325, San Diego, CA 92122 (US). **HAWRYLUK, Natalie, A.** [CA/US]; 4379 Mistral Place, San Diego, CA 92130 (US). **TIMMONS, Amy, K.** [US/US]; 5370 Toscana Way H116, San Diego, CA 92122 (US). **MERIT, Jeffrey, E.** [US/US]; 344 Olmsted Road, Stanford, CA 94305 (US). **SEIERSTAD, Mark, J.** [US/US]; 2230 Shadyridge Avenue, Escondido, CA 92029 (US).

(74) Agents: **JOHNSON, Philip, S.** et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))
- with amended claims and statement (Art. 19(1))

(54) Title: HETEROARYL-SUBSTITUTED UREA MODULATORS OF FATTY ACID AMIDE HYDROLASE

(57) Abstract: Certain heteroaryl-substituted piperidiny and piperaziny urea compounds are described, which are useful as FAAH inhibitors. Such compounds may be used in pharmaceutical compositions and methods for the treatment of disease states, disorders, and conditions mediated by fatty acid amide hydrolase (FAAH) activity, such as anxiety, pain, inflammation, sleep disorders, eating disorders, insulin resistance, diabetes, osteoporosis, and movement disorders (e.g., multiple sclerosis).



WO 2010/068452 A1

5

HETEROARYL-SUBSTITUTED UREA MODULATORS OF FATTY ACID AMIDE HYDROLASE

Cross Reference to Related Application

This application claims the benefit of US provisional patent application serial
10 number 61/117,880, filed November 25, 2008.

Field of the Invention

The present invention relates to certain heteroaryl-substituted piperidinyl and
piperazinyl urea compounds, pharmaceutical compositions containing them, and
methods of using them for the treatment of disease states, disorders, and conditions
15 mediated by fatty acid amide hydrolase (FAAH) activity.

Background of the Invention


Medicinal benefits have been attributed to the cannabis plant for centuries.
The primary bioactive constituent of cannabis is Δ^9 -tetrahydro-cannabinol (THC).
20 The discovery of THC eventually led to the identification of two endogenous
cannabinoid receptors responsible for its pharmacological actions, namely CB₁ and
CB₂ (Goya, *Exp. Opin. Ther. Patents* 2000, 10, 1529). These discoveries not only
established the site of action of THC, but also inspired inquiries into the endogenous
agonists of these receptors, or "endocannabinoids". The first endocannabinoid
25 identified was the fatty acid amide anandamide (AEA). AEA itself elicits many of the
pharmacological effects of exogenous cannabinoids (Piomelli, *Nat. Rev. Neurosci.*
2003, 4(11), 873).

The catabolism of AEA is primarily attributable to the integral membrane
bound protein fatty acid amide hydrolase (FAAH), which hydrolyzes AEA to
30 arachidonic acid. FAAH was characterized in 1996 by Cravatt and co-workers
(Cravatt, *Nature* 1996, 384, 83). It was subsequently determined that FAAH is
additionally responsible for the catabolism of a large number of important lipid
signaling fatty acid amides including: another major endocannabinoid, 2-
arachidonoylglycerol (2-AG) (*Science* 1992, 258, 1946-1949); the sleep-inducing
35 substance, oleamide (OEA) (*Science* 1995, 268, 1506); the appetite-suppressing


- 5 agent, N-oleoylethanolamine (Rodriguez de Fonesca, *Nature* 2001, 414, 209); and the anti-inflammatory agent, palmitoylethanolamide (PEA) (Lambert, *Curr. Med. Chem.* 2002, 9(6), 663).

Small-molecule inhibitors of FAAH should elevate the concentrations of these endogenous signaling lipids and thereby produce their associated beneficial
10 pharmacological effects. There have been some reports of the effects of various FAAH inhibitors in pre-clinical models.

In particular, two carbamate-based inhibitors of FAAH were reported to have analgesic properties in animal models. In rats, BMS-1 (see WO 02/087569), which has the structure shown below, was reported to have an analgesic effect in the
15 Chung spinal nerve ligation model of neuropathic pain, and the Hargraves test of acute thermal nociception. URB-597 was reported to have efficacy in the zero plus maze model of anxiety in rats, as well as analgesic efficacy in the rat hot plate and formalin tests (Kathuria, *Nat. Med.* 2003, 9(1), 76). The sulfonylfluoride AM374 was also shown to significantly reduce spasticity in chronic relapsing experimental
20 autoimmune encephalomyelitis (CREAE) mice, an animal model of multiple sclerosis (Baker, *FASEB J.* 2001, 15(2), 300).



In addition, the oxazolopyridine ketone OL-135 is reported to be a potent inhibitor of FAAH, and has been reported to have analgesic activity in both the hot
25 plate and tail emersion tests of thermal nociception in rats (WO 04/033652).



5 Results of research on the effects of certain exogenous cannabinoids has elucidated that a FAAH inhibitor may be useful for treating various conditions, diseases, disorders, or symptoms. These include pain, nausea/emesis, anorexia, spasticity, movement disorders, epilepsy and glaucoma. To date, approved therapeutic uses for cannabinoids include the relief of chemotherapy-induced nausea
10 and emesis among patients with cancer and appetite enhancement in patients with HIV/AIDs who experience anorexia as a result of wasting syndrome. Two products are commercially available in some countries for these indications, namely, dronabinol (Marinol[®]) and nabilone.

 Apart from the approved indications, a therapeutic field that has received
15 much attention for cannabinoid use is analgesia, i.e., the treatment of pain. Five small randomized controlled trials showed that THC is superior to placebo, producing dose-related analgesia (Robson, *Br. J. Psychiatry* 2001, 178, 107-115). Atlantic Pharmaceuticals is reported to be developing a synthetic cannabinoid, CT-3, a 1,1-dimethyl heptyl derivative of the carboxylic metabolite of tetrahydrocannabinol, as an
20 orally active analgesic and anti-inflammatory agent. A pilot phase II trial in chronic neuropathic pain with CT-3 was reportedly initiated in Germany in May 2002.

 A number of individuals with locomotor activity-related diseases, such as multiple sclerosis have claimed a benefit from cannabis for both disease-related pain and spasticity, with support from small controlled trials (Croxford et al., *J. Neuroimmunol*, 2008, 193, 120-9; Svendsen, *Br. Med. J.* 2004, 329, 253). Likewise,
25 various victims of spinal cord injuries, such as paraplegia, have reported that their painful spasms are alleviated after smoking marijuana. A report showing that cannabinoids appear to control spasticity and tremor in the CREA model of multiple sclerosis demonstrated that these effects are mediated by CB₁ and CB₂ receptors
30 (Baker, *Nature* 2000, 404, 84-87). Phase 3 clinical trials have been undertaken in multiple sclerosis and spinal cord injury patients with a narrow ratio mixture of tetrahydrocannabinol/cannabidiol (THC/CBD). It has been reported that FAAH knockout mice consistently recover to a better clinical score than wild type controls, and this improvement is not a result of anti-inflammatory activity, but rather may
35 reflect some neuroprotection or remyelination promoting effect of lack of the enzyme (Webb et al, *Neurosci Lett.*, 2008, vol. 439, 106-110).

5 Reports of small-scale controlled trials to investigate other potential commercial uses of cannabinoids have been made. Trials in volunteers have been reported to have confirmed that oral, injected, and smoked cannabinoids produced dose-related reductions in intraocular pressure (IOP) and therefore may relieve glaucoma symptoms. Ophthalmologists have prescribed cannabis for patients with
10 glaucoma in whom other drugs have failed to adequately control intraocular pressure (Robson, 2001, *supra*).

Inhibition of FAAH using a small-molecule inhibitor may be advantageous compared to treatment with a direct-acting CB₁ agonist. Administration of exogenous CB₁ agonists may produce a range of responses, including reduced
15 nociception, catalepsy, hypothermia, and increased feeding behavior. These four in particular are termed the “cannabinoid tetrad.” Experiments with FAAH *-/-* mice show reduced responses in tests of nociception, but did not show catalepsy, hypothermia, or increased feeding behavior (Cravatt, *Proc. Natl. Acad. Sci. USA* 2001, 98(16), 9371). Fasting caused levels of AEA to increase in rat limbic
20 forebrain, but not in other brain areas, providing evidence that stimulation of AEA biosynthesis may be anatomically regionalized to targeted CNS pathways (Kirkham, *Br. J. Pharmacol.* 2002, 136, 550). The finding that AEA increases are localized within the brain, rather than systemic, suggests that FAAH inhibition with a small molecule could enhance the actions of AEA and other fatty acid amides in tissue
25 regions where synthesis and release of these signaling molecules is occurring in a given pathophysiological condition (Piomelli, 2003, *supra*).

In addition to the effects of a FAAH inhibitor on AEA and other endocannabinoids, inhibitors of FAAH's catabolism of other lipid mediators may be used in treating certain other therapeutic indications. For example, PEA has
30 demonstrated biological effects in animal models of inflammation (Holt, et al. *Br. J. Pharmacol.* 2005, 146, 467-476), immunosuppression, analgesia, and neuroprotection (Ueda, *J. Biol. Chem.* 2001, 276(38), 35552). Oleamide, another substrate of FAAH, induces sleep (Boger, *Proc. Natl. Acad. Sci. USA* 2000, 97(10), 5044; Mendelson, *Neuropsychopharmacology* 2001, 25, S36). Inhibition of FAAH
35 has also been implicated in cognition (Varvel et al., *J. Pharmacol. Exp. Ther.* 2006,

- 5 317(1), 251-257) and depression (Gobbi et al., *Proc. Natl. Acad. Sci. USA* 2005, 102(51), 18620-18625).

Two additional indications for FAAH are supported by recent data indicating that FAAH substrate activated receptors are important in energy metabolism, and in bone homeostasis (Overton et al., *Br. J. Pharmacol.* 2008, in press; and Plutzky, 10 *Diab. Vasc. Dis. Res.* 2007, 4 Suppl 3, S12-4). It has been shown that the previously mentioned lipid signaling fatty acid amides catabolized by FAAH, oleoylethanolamide (OEA), is one of the most active agonists of the recently orphanised GPCR 119 (GPR119) (also termed glucose dependent insulinotropic receptor). This receptor is expressed predominantly in the pancreas in humans and 15 activation improves glucose homeostasis via glucose-dependent insulin release in pancreatic beta-cells. GPR119 agonists can suppress glucose excursions when administered during oral glucose tolerance tests, and OEA has also been shown independently to regulate food intake and body weight gain when administered to rodents, indicating a probable benefit in energy metabolism disorders, such as 20 insulin resistance and diabetes. The FAAH substrate palmitoylethanolamide (PEA) is an agonist at the PPAR α receptor. Evidence from surrogate markers in human studies with the PPAR α agonist fenofibrate is supportive of the concept that PPAR α agonism offers the potential for inducing a coordinated PPAR α response that may improve dyslipidaemia, repress inflammation and limit atherosclerosis in patients with 25 the metabolic syndrome or type 2 diabetes. The FAAH substrate anandamide (AEA) is an agonist at the PPAR γ receptor. Anandamide treatment induces 3T3-L1 differentiation into adipocytes, as well as triglyceride droplet accumulation and expression of adiponectin (Bouaboula et al., *E. J. Pharmacol.* 2005, 517, 174-181). Low dose cannabinoid therapy has been shown to reduce atherosclerosis in mice, 30 further suggesting a therapeutic benefit of FAAH inhibition in dyslipidemia, liver steatosis, steatohepatitis, obesity, and metabolic syndrome (Steffens et al., *Nature*, 2005, 434, 782-6).

Osteoporosis is one of the most common degenerative diseases. It is characterized by reduced bone mineral density (BMD) with an increased risk for 35 bone fractures. CB₂-deficient mice have a markedly accelerated age-related trabecular bone loss and cortical expansion. A CB₂-selective agonism enhances

5 endocortical osteoblast number and activity and restrains trabecular
osteoclastogenesis and attenuates ovariectomy-induced bone loss (Ofek et al., *Proc.*
Natl. Acad. Sci. U.S.A. 2006, 103, 696-701). There is a substantial genetic
contribution to BMD, although the genetic factors involved in the pathogenesis of
human osteoporosis are largely unknown. The applicability to human BMD is
10 suggested by genetic studies in which a significant association of single
polymorphisms and haplotypes was found encompassing the CNR2 gene on human
chromosome 1p36, demonstrating a role for the peripherally expressed CB₂ receptor
in the etiology of osteoporosis (Karsak et al., *Hum. Mol. Genet.*, 2005, 14, 3389-96).

Thus, small-molecule FAAH inhibitors should be useful in treating pain of
15 various etiologies, anxiety, multiple sclerosis and other movement disorders,
nausea/emesis, eating disorders, epilepsy, glaucoma, inflammation,
immunosuppression, neuroprotection, depression, cognition enhancement, and
sleep disorders, and potentially with fewer side effects than treatment with an
exogenous cannabinoid.


20 A number of heteroaryl-substituted ureas have been reported in various
publications. Certain substituted thiophene ureas are described in US Patent No.
6,881,741. Certain ureido-pyrazoles are described in US Patent No. 6,387,900.
Certain benzothiazole amide derivatives are described in US Patent Publication US
2003/149036. Certain ureas are reported as prenyltransferase inhibitors in WO
25 2003/047569. Piperidinyl ureas are described as histamine H₃ receptor antagonists
in US Patent No. 6,100,279. Piperazinyl ureas are disclosed as calcitonin mimetics
in US Patent Nos. 6,124,299 and 6,395,740. Various ureas are reported as small-
molecule FAAH modulators in US Patent Publication Nos. US 2006/173184 and US
2007/0004741, in Intl. Patent Appl. Nos. WO 2008/023720, WO 2008/047229, and
30 WO 2008/024139, and by Cravatt et al. (*Biochemistry* 2007, 46(45), 13019. Ureas
are described as modulators of other targets in U.S. Pat. Appl. Publ. US
2007/270433, and in Intl. Pat. Appl. Publ. Nos. WO 2007/096251 and WO
2006/085108. Certain piperidinyl ureas and piperazinyl ureas have been previously
described as FAAH modulators in U.S. Pat. Appl. No. 60/931,920, filed May 25,
35 2007. However, there remains a desire for potent FAAH modulators with suitable
pharmaceutical properties.

5

Summary of the Invention

Certain heteroaryl-substituted piperidinyl urea derivatives have been found to have FAAH-modulating activity. Thus, the invention is directed to the general and preferred embodiments defined, respectively, by the independent and dependent claims appended hereto, which are incorporated by reference herein.

In one general aspect, the invention is directed to a chemical entity of Formula (I):



Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

(i) phenyl unsubstituted or substituted with one or two R^a moieties;

where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃,

-OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c,

-SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;

or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -

OCF₂O-;

where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b

and R^c taken together with the atoms of attachment form a 4-8 membered ring;

(ii) phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties, wherein:

5 L is a linker selected from the group consisting of $-(CH_2)_{1-6}-$, $-CH=CH-$, $-O-$, $-C_{1-6}alkyl-O-C_{1-6}alkyl-$, $-C_{1-6}alkyl-N(C_{1-6}alkyl)-C_{1-6}alkyl-$, $-C_{1-6}alkyl-S(O)_{0-2}C_{1-6}alkyl-$, $-C\equiv C-$, $-C(O)-$, or a covalent bond;

Ar³ is:

- (a) phenyl unsubstituted or substituted with one or two R^a moieties;
- 10 (b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom
- 15 selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;

and pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, and

20 pharmaceutically active metabolites of such compounds. In especially preferred embodiments, the invention is directed to compounds described or exemplified in the detailed description below and their pharmaceutically acceptable salts.

In a further general aspect, the invention relates to pharmaceutical compositions each comprising an effective amount of at least one chemical entity

25 selected from compounds of Formula (I), pharmaceutically acceptable salts of compounds of Formula (I), pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I); and (b) a pharmaceutically acceptable excipient.

In another general aspect, the invention is directed to a method of treating a

30 subject suffering from or diagnosed with a disease, disorder, or medical condition mediated by FAAH activity, comprising administering to the subject in need of such treatment an effective amount of at least one chemical entity selected from compounds of Formula (I) and their pharmaceutically acceptable salts, pharmaceutically active prodrugs, and pharmaceutically active metabolites. In

35 preferred embodiments of the inventive method, the disease, disorder, or medical condition is selected from: anxiety, depression, pain, sleep disorders, eating

5 disorders, inflammation, multiple sclerosis and other movement disorders, HIV
wasting syndrome, closed head injury, stroke, learning and memory disorders,
Alzheimer's disease, epilepsy, Tourette's syndrome, Niemann-Pick disease,
Parkinson's disease, Huntington's chorea, optic neuritis, autoimmune uveitis,
10 symptoms of drug withdrawal, nausea, emesis, sexual dysfunction, post-traumatic
stress disorder, cerebral vasospasm, glaucoma, irritable bowel syndrome,
inflammatory bowel disease, immunosuppression, gastroesophageal reflux disease,
paralytic ileus, secretory diarrhea, gastric ulcer, rheumatoid arthritis, unwanted
pregnancy, hypertension, cancer, hepatitis, allergic airway disease, auto-immune
diabetes, intractable pruritis, and neuroinflammation.

15 Additional embodiments, features, and advantages of the invention will be
apparent from the following detailed description and through practice of the
invention.

Detailed Description of Invention and Its Preferred Embodiments

The invention may be more fully appreciated by reference to the following
20 detailed description, including the following glossary of terms and the concluding
examples. For the sake of brevity, the disclosures of the publications, including
patents, cited in this specification are herein incorporated by reference.


As used herein, the terms "including", "containing" and "comprising" are used
in their open, non-limiting sense.

25 The term "alkyl" refers to a straight- or branched-chain alkyl group having from
1 to 12 carbon atoms in the chain. Examples of alkyl groups include methyl (Me,
which also may be structurally depicted by / symbol), ethyl (Et), n-propyl, isopropyl,
butyl, isobutyl, sec-butyl, tert-butyl (tBu), pentyl, isopentyl, tert-pentyl, hexyl, isohexyl,
and so on.

30 The term "alkenyl" refers to a straight- or branched-chain alkenyl group having
from 2 to 12 carbon atoms in the chain. (The double bond of the alkenyl group is
formed by two sp^2 hybridized carbon atoms.) Illustrative alkenyl groups include prop-
2-enyl, but-2-enyl, but-3-enyl, 2-methylprop-2-enyl, hex-2-enyl, and so on.


The term "cycloalkyl" refers to a saturated or partially saturated, monocyclic,
35 fused polycyclic, or spiro polycyclic carbocycle having from 3 to 12 ring atoms per

5 carbocycle. Illustrative examples of cycloalkyl groups include the following entities, in the form of properly bonded moieties:



10 A “heterocycloalkyl” refers to a monocyclic, or fused, bridged, or spiro polycyclic ring structure that is saturated or partially saturated and has from 3 to 12 ring atoms per ring structure selected from carbon atoms and up to three heteroatoms selected from nitrogen, oxygen, and sulfur. The ring structure may optionally contain up to two oxo groups on carbon or sulfur ring members.

15 Illustrative examples of heterocycloalkyl groups include the following entities, in the form of properly bonded moieties:




20

The term “heteroaryl” refers to a monocyclic, fused bicyclic, or fused polycyclic aromatic heterocycle (ring structure having ring atoms selected from carbon atoms and up to four heteroatoms selected from nitrogen, oxygen, and sulfur) having from 3 to 12 ring atoms per heterocycle. Illustrative examples of heteroaryl

25

groups include the following entities, in the form of properly bonded moieties:

5



10 The term "halogen" represents chlorine, fluorine, bromine or iodine. The term "halo" represents chloro, fluoro, bromo or iodo.

The term "substituted" means that the specified group or moiety bears one or more substituents. The term "unsubstituted" means that the specified group bears no substituents. The term "optionally substituted" means that the specified group is unsubstituted or substituted by one or more substituents. Where the term

15 "substituted" is used to describe a structural system, the substitution is meant to occur at any valency-allowed position on the system. In cases where a specified moiety or group is not expressly noted as being optionally substituted or substituted with any specified substituent, it is understood that such a moiety or group is intended to be unsubstituted.

20 A structural formula given herein is intended to represent compounds having structures depicted by the formula as well as equivalent variations or forms. For example, compounds encompassed by Formula (I) may have asymmetric centers and therefore exist in different enantiomeric forms. All optical isomers and stereoisomers of the compounds of the general formula, and mixtures thereof, are

25 considered within the scope of the formula. Thus, a general formula given herein is intended to represent a racemate, one or more enantiomeric forms, one or more diastereomeric forms, one or more atropisomeric forms, and mixtures thereof. Furthermore, certain structures may exist as geometric isomers (i.e., *cis* and *trans* isomers), as tautomers (e.g. pyrazole, benzimidazole, tetrazole, or benzotriazole

30 tautomers), or as atropisomers, which are intended to be represented by the

5 structural formula. Additionally, a formula given herein is intended to embrace hydrates, solvates, and polymorphs of such compounds, and mixtures thereof.

A structural formula given herein is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or
10 more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, and chlorine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{32}P , ^{33}P , ^{35}S , ^{18}F , ^{36}Cl , and ^{125}I , respectively. Such isotopically labeled compounds are useful in
15 metabolic studies (preferably with ^{14}C), reaction kinetic studies (with, for example ^2H or ^3H), detection or imaging techniques [such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT)], including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ^{18}F - or ^{11}C -labeled compound may be preferred for PET or SPECT
20 studies. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and
25 preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

When referring to any formula given herein, the selection of a particular moiety from a list of possible species for a specified variable is not intended to define the moiety for the variable appearing elsewhere. In other words, where a formula
30 variable appears more than once, the choice of the species from a specified list is independent of the choice of the species for the same variable elsewhere in the formula.

In certain embodiments, Ar^1 is isoxazolo[5,4-c]pyridin-3-yl, isoxazolo[4,5-c]pyridin-3-yl, isoxazolo[4,5-b]pyridin-3-yl, isoxazolo[5,4-b]pyridin-3-yl, imidazo[1,2-a]pyridin-8-yl, imidazo[1,2-a]pyridin-7-yl, imidazo[1,2-a]pyridin-6-yl, imidazo[1,2-a]pyridin-5-yl, imidazo[1,2-b]pyridazin-3-yl, imidazo[1,2-a]pyridin-3-yl, imidazo[1,2-
35 a]pyridin-5-yl, imidazo[1,2-b]pyridazin-3-yl, imidazo[1,2-a]pyridin-3-yl, imidazo[1,2-

5 a]pyridin-2-yl, imidazo[1,2-a]pyrimidin-7-yl, imidazo[1,2-a]pyrimidin-5-yl, imidazo[1,2-c]pyrimidin-7-yl, benzooxazol-6-yl, 1H-indazol-7-yl, quinolin-8-yl, isoquinolin-5-yl, isoquinolin-4-yl, 1,1-dioxo-1H-1 λ ⁶-benzo[d]isothiazol-3-yl, 1H-pyrrolo[2,3-b]pyridin-5-yl, 1H-pyrrolo[2,3-b]pyridin-4-yl, 1H-pyrrolo[3,2-b]pyridin-6-yl, 2-methyl-benzothiazol-6-yl, 1-isopropyl-1H-pyrazolo[3,4-b]pyridin-5-yl, 2-methyl-benzooxazol-5-yl, 1H-
 10 indazol-7-yl, 1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl, 2-methyl-2H-indazol-4-yl, benzooxazole-2-yl group, 6-[1,2,3]triazol-1-yl-pyridin-3-yl, 6-[1,2,4]triazol-4-yl-pyridin-3-yl, 6-[1,2,4]triazol-1-yl-pyridin-3-yl, 6-[1,2,3]triazol-2-yl-pyridin-3-yl, 4-[1,2,3]triazol-2-yl-phenyl, 4-[1,2,3]triazol-1-yl-phenyl, 5-methyl-[1,3,4]oxadiazol-2-yl, 2-phenyl-pyrimidin-5-yl, 3-pyridyl, benzisoxazol-3-yl, pyrimidin-4-yl, isoxazol-3-yl, 6-
 15 fluorobenzo[d]isoxazol-3-yl, 3-phenyl-[1,2,4]thiadiazol-5-yl, 1H-tetrazol-5-yl, benzo[1,2,5]thiadiazol-4-yl, benzo[1,2,5]oxadiazol-4-yl, thiophen-2-yl, thiophen-3-yl, 6-chloro-pyridazin-3-yl, pyrazin-2-yl, 1H-benzotriazol-5-yl, [1,5]naphthyridin-2-yl, benzothiazol-6-yl, or 1H-pyrazol-3-yl. In certain embodiments, Ar¹ is pyrimidine, isoxazole, benzisoxazole, imidazo[1,2-b]pyridazine, or pyridine, wherein each group
 20 is optionally substituted with one or two R^a moieties.

In certain embodiments, Ar² is phenyl substituted with one or two R^a moieties. In certain embodiments, Ar² is phenyl, substituted with one or two R^a moieties wherein each wherein each R^a moiety is independently selected from the group consisting of: chloro, cyano, isobutyl, methylsulfanyl, methanesulfonyl,
 25 trifluoromethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, fluoro, methyl, methoxy, tert-butyl, bromo, methoxycarbonyl, cyanomethyl, methoxycarbonylmethyl, trifluoromethanesulfonyl, trifluoromethanesulfanyl, and butyl; or two adjacent R^a moieties taken together form -OCH₂O- or -OCF₂O-. In further embodiments, each R^a moiety is independently selected from the group consisting of: F, Cl, Br, CF₃, or
 30 two adjacent R^a moieties taken together form -OCF₂O-.

In certain embodiments, Ar² is phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties. In further embodiments, Ar² is phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties and wherein L is -CH₂CH₂-,
 35 -O-, -OCH₂-, or -C≡C-. In further embodiments, L is -O-. In further embodiments, L is phenyl, unsubstituted or substituted with one or two R^a moieties and each R^a

5 moiety is independently selected from the group consisting of: chloro, cyano, isobutyl, methylsulfanyl, methanesulfonyl, trifluoromethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, fluoro, methyl, methoxy, tert-butyl, bromo, methoxycarbonyl, cyanomethyl, methoxycarbonylmethyl, trifluoromethanesulfonyl, trifluoromethanesulfanyl, and butyl; or two adjacent R^a moieties taken together form
10 -OCH₂O- or -OCF₂O-. In certain preferred embodiments, each R^a moiety is independently fluoro.

In a further embodiment, Ar² is a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said
15 fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties. In one embodiment, Ar² is a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom selected from the group consisting of N, O, and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said
20 fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties. In still further preferred embodiments, Ar² is a benzimidazolyl, indazolyl, benzothiophenyl, quinolinyl, indolyl, or benzofuranyl group. In a further embodiment Ar² is quinolinyl.

The invention also relates to pharmaceutically acceptable salts of the free
25 acids or bases represented by compounds in Table 1, preferably of the preferred embodiments described above and of the specific compounds exemplified herein. The therapeutic compositions and methods of the invention may employ pharmaceutically acceptable salts of the free acids or bases represented by compounds in Table 1, preferably of the preferred embodiments described above
30 and of the specific compounds exemplified herein. A "pharmaceutically acceptable salt" is intended to mean a salt of a free acid or base of a compound represented by Formula (I) that is non-toxic, biologically tolerable, or otherwise biologically suitable for administration to the subject. See, generally, S.M. Berge, et al., "Pharmaceutical Salts", J. Pharm. Sci., 1977, 66:1-19, and *Handbook of Pharmaceutical Salts, Properties, Selection, and Use*, Stahl and Wermuth, Eds., Wiley-VCH and VHCA,
35 Zurich, 2002.

5 Preferred pharmaceutically acceptable salts are those that are pharmacologically effective and suitable for contact with the tissues of patients without undue toxicity, irritation, or allergic response. A compound from Table 1 may possess a sufficiently acidic group, a sufficiently basic group, or both types of functional groups, and accordingly react with a number of inorganic or organic
10 bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt. Examples of pharmaceutically acceptable salts include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrogen-phosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates,
15 isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates, γ -hydroxybutyrates,
20 glycolates, tartrates, methane-sulfonates, propanesulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, and mandelates.

If the compound in Table 1 contains a basic nitrogen, the desired pharmaceutically acceptable salt may be prepared by any suitable method available in the art, for example, by treatment of the free base with an inorganic acid, such as
25 hydrochloric acid, hydrobromic acid, sulfuric acid, sulfamic acid, nitric acid, boric acid, phosphoric acid, and the like; or with an organic acid, such as acetic acid, phenylacetic acid, propionic acid, stearic acid, lactic acid, ascorbic acid, maleic acid, hydroxymaleic acid, isethionic acid, succinic acid, valeric acid, fumaric acid, malonic acid, pyruvic acid, oxalic acid, glycolic acid, salicylic acid, oleic acid, palmitic acid,
30 lauric acid, a pyranosidyl acid, such as glucuronic acid or galacturonic acid, an α -hydroxy acid, such as mandelic acid, citric acid, or tartaric acid; an amino acid, such as aspartic acid or glutamic acid; an aromatic acid, such as benzoic acid, 2-acetoxybenzoic acid, naphthoic acid, or cinnamic acid; a sulfonic acid, such as laurylsulfonic acid, p-toluenesulfonic acid, methanesulfonic acid, or ethanesulfonic
35 acid; or any compatible mixture of acids such as those given as examples herein.

5 If the compound in Table 1 is an acid such as a carboxylic acid or sulfonic acid, the desired pharmaceutically acceptable salt may be prepared by any suitable method, for example, by treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary or tertiary), an alkali metal hydroxide, alkaline earth metal hydroxide, or any compatible mixture of bases such as those given as
10 examples herein. Illustrative examples of suitable salts include organic salts derived from amino acids, such as glycine and arginine, ammonia, carbonates, bicarbonates, primary, secondary, and tertiary amines, and cyclic amines, such as benzylamines, pyrrolidines, piperidine, morpholine, and piperazine, and inorganic salts derived from sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum,
15 and lithium.

The invention also relates to pharmaceutically acceptable prodrugs of the compounds of Table 1. The term "prodrug" means a precursor of a designated compound that, following administration to a subject, yields the compound *in vivo* via a chemical or physiological process such as solvolysis or enzymatic cleavage, or
20 under physiological conditions (e.g., a prodrug on being brought to physiological pH is converted to the compound of Table 1). A "pharmaceutically acceptable prodrug" is a prodrug that is non-toxic, biologically tolerable, and otherwise biologically suitable for administration to the subject. Illustrative procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of
25 Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

Examples of prodrugs include compounds having an amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid residues, covalently joined through an amide or ester bond to a free amino, hydroxy, or carboxylic acid group of a compound of Formula (I). Examples of amino acid
30 residues include the twenty naturally occurring amino acids, commonly designated by three letter symbols, as well as 4-hydroxyproline, hydroxylysine, demosine, isodemosine, 3-methylhistidine, norvalin, beta-alanine, gamma-aminobutyric acid, citrulline homocysteine, homoserine, ornithine and methionine sulfone.

Additional types of prodrugs may be produced, for instance, by derivatizing
35 free carboxyl groups of structures of compounds of Table 1 as amides or alkyl esters. Examples of amides include those derived from ammonia, primary C₁₋₆alkyl

5 amines and secondary di(C₁₋₆alkyl) amines. Secondary amines include 5- or 6-membered heterocycloalkyl or heteroaryl ring moieties. Examples of amides include those that are derived from ammonia, C₁₋₃alkyl primary amines, and di(C₁₋₂alkyl)amines. Examples of esters of the invention include C₁₋₇alkyl, C₅₋₇cycloalkyl, phenyl, and phenyl(C₁₋₆alkyl) esters. Preferred esters include methyl esters.

10 Prodrugs may also be prepared by derivatizing free hydroxy groups using groups including hemisuccinates, phosphate esters, dimethylaminoacetates, and phosphoryloxymethyloxycarbonyls, following procedures such as those outlined in Fleisher et al., *Adv. Drug Delivery Rev.* 1996, 19, 115-130. Carbamate derivatives of hydroxy and amino groups may also yield prodrugs. Carbonate derivatives,

15 sulfonate esters, and sulfate esters of hydroxy groups may also provide prodrugs. Derivatization of hydroxy groups as (acyloxy)methyl and (acyloxy)ethyl ethers, wherein the acyl group may be an alkyl ester, optionally substituted with one or more ether, amine, or carboxylic acid functionalities, or where the acyl group is an amino acid ester as described above, is also useful to yield prodrugs. Prodrugs of this type

20 may be prepared as described in Robinson et al., *J. Med. Chem.* 1996, 39, 10-18. Free amines can also be derivatized as amides, sulfonamides or phosphoramidates. All of these prodrug moieties may incorporate groups including ether, amine, and carboxylic acid functionalities.

The present invention also relates to pharmaceutically active metabolites of

25 compounds of Table 1. A "pharmaceutically active metabolite" means a pharmacologically active product of metabolism in the body of a compound of Table 1 or salt thereof. Prodrugs and active metabolites of a compound may be determined using routine techniques known or available in the art. See, e.g., Bertolini et al., *J. Med. Chem.* 1997, 40, 2011-2016; Shan et al., *J. Pharm. Sci.* 1997,

30 86 (7), 765-767; Bagshawe, *Drug Dev. Res.* 1995, 34, 220-230; Bodor, *Adv. Drug Res.* 1984, 13, 255-331; Bundgaard, *Design of Prodrugs* (Elsevier Press, 1985); and Larsen, *Design and Application of Prodrugs*, Drug Design and Development (Krogsgaard-Larsen et al., eds., Harwood Academic Publishers, 1991).

The compounds of Table 1, and their pharmaceutically acceptable salts,

35 pharmaceutically acceptable prodrugs, and pharmaceutically active metabolites (collectively, "active agents") of the present invention are useful as FAAH inhibitors in

5 the methods of the invention. The active agents may be used in the inventive methods for the treatment of medical conditions, diseases, or disorders mediated through inhibition or modulation of FAAH, such as those described herein. Active agents according to the invention may therefore be used as an analgesic, anti-depressant, cognition enhancer, neuroprotectant, sedative, appetite stimulant, or
10 contraceptive.

Exemplary medical conditions, diseases, and disorders mediated by FAAH activity include anxiety, depression, pain, sleep disorders, eating disorders, inflammation, multiple sclerosis and other movement disorders, HIV wasting syndrome, closed head injury, stroke, learning and memory disorders, Alzheimer's
15 disease, epilepsy, Tourette's syndrome, epilepsy, Niemann-Pick disease, Parkinson's disease, Huntington's chorea, optic neuritis, autoimmune uveitis, symptoms of drug withdrawal, nausea, emesis, sexual dysfunction, post-traumatic stress disorder, cerebral vasospasm, diabetes, metabolic syndrome and osteoporosis.

20 Thus, the active agents may be used to treat subjects diagnosed with or suffering from such a disease, disorder, or condition. The term "treat" or "treating" as used herein is intended to refer to administration of an agent or composition of the invention to a subject for the purpose of effecting a therapeutic benefit through modulation of FAAH activity. Treating includes reversing, ameliorating, alleviating,
25 inhibiting the progress of, lessening the severity of, reducing the incidence of, or preventing a disease, disorder, or condition, or one or more symptoms of such disease, disorder or condition mediated through modulation of FAAH activity. The term "subject" refers to a mammalian patient in need of such treatment, such as a human. "Modulators" include both inhibitors and activators, where "inhibitors" refer
30 to compounds that decrease, prevent, inactivate, desensitize or down-regulate FAAH expression or activity, and "activators" are compounds that increase, activate, facilitate, sensitize, or up-regulate FAAH expression or activity.

Accordingly, the invention relates to methods of using the active agents described herein to treat subjects diagnosed with or suffering from a disease,
35 disorder, or condition mediated through FAAH activity, such as: anxiety, pain, sleep disorders, eating disorders, inflammation, movement disorders (e.g., multiple

5 sclerosis), energy metabolism (e.g. insulin resistance, diabetes, dyslipidemia, liver steatosis, steatohepatitis, obesity, and metabolic syndrome) and bone homeostasis (e.g. osteoporosis).

Symptoms or disease states are intended to be included within the scope of "medical conditions, disorders, or diseases." For example, pain may be associated
10 with various diseases, disorders, or conditions, and may include various etiologies. Illustrative types of pain treatable with a FAAH-modulating agent, in one example herein a FAAH-inhibiting agent, according to the invention include cancer pain, postoperative pain, GI tract pain, spinal cord injury pain, visceral hyperalgesia, thalamic pain, headache (including stress headache and migraine), low back pain,
15 neck pain, musculoskeletal pain, peripheral neuropathic pain, central neuropathic pain, neurogenerative disorder related pain, and menstrual pain. HIV wasting syndrome includes associated symptoms such as appetite loss and nausea. Parkinson's disease includes, for example, levodopa-induced dyskinesia. Treatment of multiple sclerosis may include treatment of symptoms such as spasticity,
20 neurogenic pain, central pain, or bladder dysfunction. Symptoms of drug withdrawal may be caused by, for example, addiction to opiates or nicotine. Nausea or emesis may be due to chemotherapy, postoperative, or opioid related causes. Treatment of sexual dysfunction may include improving libido or delaying ejaculation. Treatment of cancer may include treatment of glioma. Sleep disorders include, for example,
25 sleep apnea, insomnia, and disorders calling for treatment with an agent having a sedative or narcotic-type effect. Eating disorders include, for example, anorexia or appetite loss associated with a disease such as cancer or HIV infection/AIDS.

In treatment methods according to the invention, an effective amount of at least one active agent according to the invention is administered to a subject
30 suffering from or diagnosed as having such a disease, disorder, or condition. A "therapeutically effective amount" or "effective amount" means an amount or dose of a FAAH-modulating agent sufficient to generally bring about a therapeutic benefit in patients in need of treatment for a disease, disorder, or condition mediated by FAAH activity. Effective amounts or doses of the active agents of the present invention
35 may be ascertained by routine methods such as modeling, dose escalation studies or clinical trials, and by taking into consideration routine factors, e.g., the mode or

5 route of administration or drug delivery, the pharmacokinetics of the agent, the severity and course of the disease, disorder, or condition, the subject's previous or ongoing therapy, the subject's health status and response to drugs, and the judgment of the treating physician. An exemplary dose is in the range of from about 0.0001 to about 200 mg of active agent per kg of subject's body weight per day, preferably about 0.001 to 100 mg/kg/day, or about 0.01 to 35 mg/kg/day, or about 0.1 to 10 mg/kg daily in single or divided dosage units (e.g., BID, TID, QID). For a 10 70-kg human, an illustrative range for a suitable dosage amount is from about 0.05 to about 7 g/day, or about 0.2 to about 5 g/day. Once improvement of the patient's disease, disorder, or condition has occurred, the dose may be adjusted for maintenance treatment. For example, the dosage or the frequency of administration, or both, may be reduced as a function of the symptoms, to a level at which the desired therapeutic effect is maintained. Of course, if symptoms have been alleviated to an appropriate level, treatment may cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of 20 symptoms.

In addition, the active agents of the invention may be used in combination with additional active ingredients in the treatment of the above conditions. The additional active ingredients may be coadministered separately with an active agent of compounds of Table 1 or included with such an agent in a pharmaceutical 25 composition according to the invention. In an exemplary embodiment, additional active ingredients are those that are known or discovered to be effective in the treatment of conditions, disorders, or diseases mediated by FAAH activity, such as another FAAH modulator or a compound active against another target associated with the particular condition, disorder, or disease. The combination may serve to increase efficacy (e.g., by including in the combination a compound potentiating the 30 potency or effectiveness of an active agent according to the invention), decrease one or more side effects, or decrease the required dose of the active agent according to the invention. In one illustrative embodiment, a composition according to the invention may contain one or more additional active ingredients selected from 35 opioids, NSAIDs (e.g., ibuprofen, cyclooxygenase-2 (COX-2) inhibitors, and naproxen), gabapentin, pregabalin, tramadol, acetaminophen, and aspirin.

5 The active agents of the invention are used, alone or in combination with one or more additional active ingredients, to formulate pharmaceutical compositions of the invention. A pharmaceutical composition of the invention comprises: (a) an effective amount of at least one active agent in accordance with the invention; and (b) a pharmaceutically acceptable excipient.

10 A "pharmaceutically acceptable excipient" refers to a substance that is non-toxic, biologically tolerable, and otherwise biologically suitable for administration to a subject, such as an inert substance, added to a pharmacological composition or otherwise used as a vehicle, carrier, or diluent to facilitate administration of a agent and that is compatible therewith. Examples of excipients include calcium carbonate, 15 calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils, and polyethylene glycols.

 Delivery forms of the pharmaceutical compositions containing one or more dosage units of the active agents may be prepared using suitable pharmaceutical excipients and compounding techniques known or that become available to those 20 skilled in the art. The compositions may be administered in the inventive methods by a suitable route of delivery, e.g., oral, parenteral, rectal, topical, or ocular routes, or by inhalation.

 The preparation may be in the form of tablets, capsules, sachets, dragees, powders, granules, lozenges, powders for reconstitution, liquid preparations, or 25 suppositories. Preferably, the compositions are formulated for intravenous infusion, topical administration, or oral administration.

 For oral administration, the active agents of the invention can be provided in the form of tablets or capsules, or as a solution, emulsion, or suspension. To prepare the oral compositions, the active agents may be formulated to yield a 30 dosage of, e.g., from about 5 mg to 5 g daily, or from about 50 mg to 5 g daily, in single or divided doses. For example, a total daily dosage of about 5 mg to 5 g daily may be accomplished by dosing once, twice, three, or four times per day.

 Oral tablets may include the active ingredient(s) mixed with compatible pharmaceutically acceptable excipients such as diluents, disintegrating agents, 35 binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservative agents. Suitable inert fillers include sodium and calcium

5 carbonate, sodium and calcium phosphate, lactose, starch, sugar, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol, and the like. Exemplary liquid oral excipients include ethanol, glycerol, water, and the like. Starch, polyvinyl-pyrrolidone (PVP), sodium starch glycolate, microcrystalline cellulose, and alginic acid are
10 The lubricating agent, if present, may be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or glyceryl distearate to delay absorption in the gastrointestinal tract, or may be coated with an enteric coating.

Capsules for oral administration include hard and soft gelatin capsules. To
15 prepare hard gelatin capsules, active ingredient(s) may be mixed with a solid, semi-solid, or liquid diluent. Soft gelatin capsules may be prepared by mixing the active ingredient with water, an oil such as peanut oil or olive oil, liquid paraffin, a mixture of mono and di-glycerides of short chain fatty acids, polyethylene glycol 400, or propylene glycol.

20 Liquids for oral administration may be in the form of suspensions, solutions, emulsions or syrups or may be lyophilized or presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid compositions may optionally contain: pharmaceutically-acceptable excipients such as suspending agents (for example, sorbitol, methyl cellulose, sodium alginate,
25 gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminum stearate gel and the like); non-aqueous vehicles, e.g., oil (for example, almond oil or fractionated coconut oil), propylene glycol, ethyl alcohol, or water; preservatives (for example, methyl or propyl p-hydroxybenzoate or sorbic acid); wetting agents such as lecithin; and, if desired, flavoring or coloring agents.

30 The active agents of this invention may also be administered by non-oral routes. For example, compositions may be formulated for rectal administration as a suppository. For parenteral use, including intravenous, intramuscular, intraperitoneal, or subcutaneous routes, the agents of the invention may be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and
35 isotonicity or in parenterally acceptable oil. Suitable aqueous vehicles include Ringer's solution and isotonic sodium chloride. Such forms may be presented in


5 unit-dose form such as ampules or disposable injection devices, in multi-dose forms such as vials from which the appropriate dose may be withdrawn, or in a solid form or pre-concentrate that can be used to prepare an injectable formulation. Illustrative infusion doses range from about 1 to 1000 µg/kg/minute of agent admixed with a pharmaceutical carrier over a period ranging from several minutes to several days.

10 For topical administration, the agents may be mixed with a pharmaceutical carrier at a concentration of about 0.1% to about 10% of drug to vehicle. Another mode of administering the agents of the invention may utilize a patch formulation to affect transdermal delivery.

Active agents may alternatively be administered in methods of this invention
15 by inhalation, via the nasal or oral routes, e.g., in a spray formulation also containing a suitable carrier.

Exemplary active agents useful in methods of the invention will now be described by reference to illustrative synthetic schemes for their general preparation below and the specific examples that follow. Artisans will recognize that, to obtain
20 the various compounds herein, starting materials may be suitably selected so that the ultimately desired substituents will be carried through the reaction scheme with or without protection as appropriate to yield the desired product. Alternatively, it may be necessary or desirable to employ, in the place of the ultimately desired substituent, a suitable group that may be carried through the reaction scheme and
25 replaced as appropriate with the desired substituent. Unless otherwise specified, the variables are as defined above in reference to Formula (I).


SCHEME A



30 Referring to Scheme A, a carbamate of formula (IV) may be obtained by reacting a compound of formula (II) with a compound of formula (III), in which Q¹ represents an aryl group, under chloroformate condensation conditions. Preferably, Q¹ is substituted or unsubstituted phenyl, and the reaction occurs with or without a base, in a solvent such as acetonitrile, at a temperature from about 0 °C to about 80


- 5 °C. More preferably, Q¹ is phenyl, and the reaction occurs in pyridine at room temperature ("rt"), or in acetonitrile at 50 °C without added base.

SCHEME B



- 10 Referring to Scheme B, a compound of formula (VI) is obtained by reacting a compound of formula (V) with a compound of formula (IV) under aryl carbamate condensation conditions. The reaction may preferably take place in a solvent at a temperature from about rt to about 120 °C. Preferably, Q¹ is phenyl, and the reaction is performed in dimethylsulfoxide (DMSO) in a microwave reactor at about 100 °C or
- 15 by conventional heating from about rt to about 50 °C.

SCHEME C



- Intermediate compounds of formula (IX) are prepared according to Scheme C. Hydrometallation of an alkenyl compound (VII) gives an activated species, which is
- 20 subsequently reacted with a suitable reagent Ar²-HAL (where HAL is chloride, bromide, or iodide) to provide a compound (IX). Preferably, hydrometallation is accomplished by hydroboration using a suitable dialkylborane reagent such as 9-borabicyclo[3.3.1]nonane (9-BBN) or diisopinocampheylborane, in a solvent such as THF. The resulting boron adduct is preferably reacted with compounds of formula
- 25 (VIII) in the presence of a suitable palladium catalyst, a base such as K₂CO₃ or Cs₂CO₃, in a solvent such as N,N-dimethylformamide (DMF) or an aqueous mixture thereof. When Q³ is a suitable protecting group, then compounds of formula (IX) can be converted into compounds of formula (V) using suitable deprotection conditions.

Compounds of Table 1 may be converted to their corresponding salts by

5 applying general techniques described in the art. For example, a compound of Table 1 may be treated with trifluoroacetic acid, HCl, or citric acid in a solvent such as Et₂O, 1,4-dioxane, dichloromethane (DCM), tetrahydrofuran (THF), or MeOH to provide the corresponding salt forms.

10 Compounds prepared according to the schemes described above may be obtained as single enantiomers, diastereomers, or regioisomers, by enantio-, diastereo-, or regio-specific synthesis, or by resolution. Compounds prepared according to the schemes above may alternatively be obtained as racemic (1:1) or non-racemic (not 1:1) mixtures or as mixtures of diastereomers or regioisomers. Where racemic and non-racemic mixtures of enantiomers are obtained, single
15 enantiomers may be isolated using conventional separation methods, such as chiral chromatography, recrystallization, diastereomeric salt formation, derivatization into diastereomeric adducts, biotransformation, or enzymatic transformation. Where regioisomeric or diastereomeric mixtures are obtained, single isomers may be separated using conventional methods such as chromatography or crystallization.

20 The following specific examples are provided to further illustrate the invention and various preferred embodiments.

EXAMPLES

Chemistry:

25 In preparing the examples listed below, the following general experimental and analytical methods were used.

Reaction mixtures were stirred under a nitrogen atmosphere at rt unless otherwise noted. Where solutions or mixtures are concentrated, they are typically concentrated under reduced pressure using a rotary evaporator. Where solutions are dried, they are typically dried over a drying agent such as MgSO₄ or Na₂SO₄,
30 unless otherwise noted.

Microwave reactions were carried out in either a CEM Discover® or a Biotage Initiator™ Microwave at specified temperatures.

Normal phase flash column chromatography (FCC) was performed on silica gel columns using ethyl acetate (EtOAc)/hexanes as eluent, unless otherwise
35 indicated.


5 Reversed-Phase High Performance Liquid Chromatography (HPLC) was performed using: Shimadzu instrument with a Phenomenex Gemini column 5 μ m C18 (150 x 21.2 mm) or Waters Xterra RP18 OBD column 5 μ m (100 x 30 mm), a gradient of 95:5 to 0:100 water (0.05% TFA)/CH₃CN (0.05% TFA), a flow rate of 80 mL/min, and detection at 254 nm.

10 Mass spectra were obtained on an Agilent series 1100 MSD using electrospray ionization (ESI) in positive mode unless otherwise indicated.

 NMR spectra were obtained on either a Bruker model DPX400 (400 MHz), DPX500 (500 MHz) or DRX600 (600 MHz) spectrometer. The format of the ¹H NMR data below is: chemical shift in ppm down field of the tetramethylsilane reference
15 (multiplicity, coupling constant *J* in Hz, integration).


 Chemical names were generated using ChemDraw Ultra 6.0.2 (CambridgeSoft Corp., Cambridge, MA) or ACD/Name Version 9 (Advanced Chemistry Development, Toronto, Ontario, Canada).

20 Intermediate 1: Isoxazolo[5,4-c]pyridin-3-ylamine.




 To a solution of 3-chloro-isonicotinitrile (1.13 g, 8.36 mmol) in DMF (6.0 mL) were added potassium carbonate (1.69 g, 12.2 mmol) and acetohydroxamic acid (0.91 g, 12.2 mmol). The reaction mixture was stirred at rt overnight, diluted with
25 EtOAc (200 mL) and extracted with saturated aqueous NaHCO₃ (200 mL) then saturated aqueous NaCl (100 mL). The aqueous layers were back extracted with EtOAc (200 mL) and the combined organic layers were dried (MgSO₄) and concentrated. The crude residue was purified (FCC, 2 N NH₃ in MeOH/DCM) to give isoxazolo[5,4-c]pyridin-3-ylamine (0.447 g, 41%). MS (ESI⁺): calcd for C₆H₅N₃O m/z
30 135.04, found 136.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.93 (d, *J* = 0.8, 1H), 8.45 (d, *J* = 5.2, 1H), 7.88-7.86 (dd, *J* = 5.2, 1.2, 1H), 6.72 (br s, 2H).

 Intermediates 2 and 3 were prepared using methods analogous to those described for intermediate 1 with the stated changes to reagents described below.


5 Intermediate 2: Isoxazolo[4,5-c]pyridin-3-ylamine.

The title compound was prepared using methods analogous to those described for intermediate 1 except 4-chloro-3-cyanopyridine was the starting material. MS (ESI⁺): calcd for C₆H₅N₃O m/z 135.04, found 136.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 9.09 (d, *J* = 1.0, 1H), 8.57 (d, *J* = 5.9, 1H), 7.53 (dd, *J* = 5.9, 1.0, 1H), 6.77 (br s, 2H).

10 Intermediate 3: Isoxazolo[4,5-b]pyridin-3-ylamine.

15 The title compound was prepared using methods analogous to those described for intermediate 1 except 3-chloro-2-cyanopyridine was the starting material. MS (ESI⁺): calcd for C₆H₅N₃O m/z 135.04, 136.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.58 (dd, *J* = 4.5, 1.1, 1H), 7.98 (dd, *J* = 8.5, 1.1, 1H), 7.57 (dd, *J* = 8.5, 4.5, 1H), 6.55 (br s, 2H).


20

Intermediate 4: Imidazo[1,2-a]pyridin-8-ylamine.


To a solution of 50% aqueous chloroacetaldehyde (4.14 mL, 32.6 mmol) were added sodium bicarbonate (3.19 g, 40.0 mmol) and 2,3-diaminopyridine (3.56 g, 32.6 mmol). The reaction mixture was stirred at rt overnight, then diluted with saturated aqueous NaCl (100 mL) and extracted with n-butanol (3x200 mL). The organic layers were combined, dried (MgSO₄), and concentrated. The crude residue was purified (FCC, 2 N NH₃ in MeOH/DCM) to give imidazo[1,2-a]pyridin-8-ylamine (2.18 g, 51%). MS (ESI⁺): calcd for C₇H₇N₃ m/z 133.06, found 134.1 (M+H)⁺. ¹H NMR (d₆-DMSO): 7.79-7.75 (m, 2H), 7.40 (d, *J* = 1.1, 1H), 6.60 (s, 1H), 6.21 (dd, *J* = 7.3, 1.1, 1H), 5.53 (br s, 2H).

30

5

Intermediate 5: Imidazo[1,2-a]pyridin-7-ylamine.


Intermediate 5 was prepared using methods analogous to those described for intermediate 4 except 2,4-diaminopyridine was the starting material. MS (ESI⁺): calcd for C₇H₇N₃ m/z 133.06, found 134.1 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.13-8.10 (m, 1H), 7.48 (s, 1H), 7.15 (s, 1H), 6.36-6.33 (m, 2H), 5.52 (br s, 2H).

Intermediate 6: Imidazo[1,2-a]pyridin-5-ylamine.

To a solution of 2,6-diaminopyridine (5.00 g, 45.8 mmol) in 1:1 H₂O:EtOH (10 mL) were added sodium bicarbonate (5.00 g, 59.5 mmol) and chloroacetaldehyde (4.31 g, 27.5 mmol). The reaction mixture was heated to 70 °C overnight, then cooled to rt, diluted with saturated aqueous NaCl (100 mL), and extracted with n-butanol (3x200 mL). The organic layers were combined, dried (MgSO₄), and concentrated. The crude residue was purified (FCC, 2 N NH₃ in MeOH/DCM) to give imidazo[1,2-a]pyridin-5-ylamine (1.32 g, 22%). MS (ESI⁺): calcd for C₇H₇N₃ m/z 133.06, found 134.1 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.23-8.21 (m, 1H), 7.91 (d, J = 1.3, 1H), 7.50 (dd, J = 8.8, 7.4, 1H), 7.22 (d, J = 8.8, 1H), 6.97 (br s, 2H), 6.37 (dd, J = 7.4, 0.9, 1H).

25


Intermediates 7 to 9 were prepared using methods analogous to those described for intermediate 6 with the stated changes to reagents described below.

Intermediate 7: Imidazo[1,2-a]pyrimidin-7-ylamine.

30


- 5 Intermediate 7 was prepared using methods analogous to those described for Intermediate 6 except 2,4-diaminopyrimidine was the starting material. MS (ESI⁺): calcd for C₆H₆N₄ m/z 134.06, found 135.1 (M+H)⁺. ¹H NMR (d₄-methanol): 8.27 (d, *J* = 7.2, 1H), 7.31 (d, *J* = 1.8, 1H), 7.18 (d, *J* = 1.8, 1H), 6.35 (d, *J* = 7.2, 1H).

10 Intermediate 8: Imidazo[1,2-a]pyrimidin-5-ylamine.



- Intermediate 8 was prepared using methods analogous to those described for intermediate 6 except 2,4-diaminopyrimidine was the starting material. MS (ESI⁺): calcd for C₆H₆N₄ m/z 134.06, found 135.1 (M+H)⁺. ¹H NMR (d₄-methanol): 8.41 (d, *J* = 5.4, 1H), 7.72 (d, *J* = 1.8, 1H), 7.55 (d, *J* = 1.8, 1H), 6.17 (d, *J* = 5.4, 1H).
- 15


Intermediate 9: Imidazo[1,2-c]pyrimidin-7-ylamine.



- Intermediate 9 was prepared using methods analogous to those described for intermediate 6 except 4,6-diaminopyrimidine was the starting material. MS (ESI⁺): calcd for C₆H₆N₄ m/z 134.06, found 135.1 (M+H)⁺. ¹H NMR (d₆-DMSO): 9.80 (s, 1H), 8.39 (s, 1H), 8.03 (s, 1H), 7.01 (s, 1H), 6.90 (s, 2H).
- 20


- Intermediates 10 to 18 were prepared by treatment of the appropriate aminoheterocycle intermediate with 1 equivalent of phenylchloroformate in dry CH₃CN and 1 equivalent of pyridine. The reagents were stirred at rt for 4 h, diluted with EtOAc, washed with NaHCO₃, and dried over MgSO₄. The material was purified by FCC to provide pure phenyl carbamate intermediates.
- 25

30 Intermediate 10: Imidazo[1,2-a]pyridin-8-yl-carbamic acid phenyl ester.



5 MS (ESI⁺): calcd for C₁₄H₁₁N₃O₂ m/z 253.09, found 254.1 (M+H)⁺.


Intermediate 11: Imidazo[1,2-a]pyridin-3-yl-carbamic acid phenyl ester.



MS (ESI⁺): calcd for C₁₄H₁₁N₃O₂ m/z 253.09, found 254.1 (M+H)⁺.


10

Intermediate 12: Imidazo[1,2-a]pyridin-2-yl-carbamic acid phenyl ester.




MS (ESI⁺): calcd for C₁₄H₁₁N₃O₂ m/z 253.09, found 254.1 (M+H)⁺.

15 Intermediate 13: (1H-Indazol-7-yl)-carbamic acid phenyl ester.



MS (ESI⁺): calcd for C₁₄H₁₁N₃O₂ m/z 253.09, found 254.3 (M+H)⁺.


Intermediate 14: Quinolin-8-yl-carbamic acid phenyl ester.



20


MS (ESI⁺): calcd for C₁₆H₁₂N₂O₂ m/z 264.09, found 265.1 (M+H)⁺.

Intermediate 15: Isoquinolin-5-yl-carbamic acid phenyl ester.



5 MS (ESI⁺): calcd for C₁₆H₁₂N₂O₂ m/z 264.09, found 265.1 (M+H)⁺.


Intermediate 16: Isoquinolin-4-yl-carbamic acid phenyl ester.



MS (ESI⁺): calcd for C₁₆H₁₂N₂O₂ m/z 264.09, found 265.1 (M+H)⁺.


10

Intermediate 17: (1H-Pyrrolo[2,3-b]pyridin-4-yl)-carbamic acid phenyl ester.




MS (ESI⁺): calcd for C₁₄H₁₁N₃O₂ m/z 253.09, found 254.1 (M+H)⁺.

15 Intermediate 18: (1H-Pyrrolo[3,2-b]pyridin-6-yl)-carbamic acid phenyl ester.



MS (ESI⁺): calcd for C₁₄H₁₁N₃O₂ m/z 253.09, found 254.1 (M+H)⁺.


Intermediate 19: Imidazo[1,2-b]pyridazin-3-yl-carbamic acid phenyl ester.



20


To a solution of imidazo[1,2-b]pyridazin-3-ylamine (209 mg, 1.56 mmol) in dry DMF (5.0 mL) was added pyridine (382 μL, 4.68 mmol) and phenyl chloroformate (244 mg, 1.56 mmol). After 3 h at rt, the mixture was diluted with EtOAc (100 mL) and washed with saturated aqueous NaHCO₃ (50 mL). The organic layer was dried (MgSO₄) and concentrated. The residue was purified (FCC) to give imidazo[1,2-b]pyridazin-3-yl-carbamic acid phenyl ester (285 mg, 72%). MS (ESI⁺): calcd for C₁₃H₁₀N₄O₂ m/z 254.08, found 255.3 (M+H)⁺.

25


5 Intermediate 20: Isoxazolo[5,4-c]pyridin-3-yl-carbamic acid phenyl ester.

To a suspension of isoxazolo[5,4-c]pyridin-3-ylamine (1.49 g, 11.0 mmol) in CH₃CN (50 mL) was added phenyl chloroformate (0.696 mL, 5.50 mmol). The reaction mixture was heated at 70 °C overnight, then cooled to rt, diluted with H₂O (100 mL) and saturated aqueous NaHCO₃ (10 mL). The resulting yellow precipitate was filtered and dried under vacuum to give isoxazolo[5,4-c]pyridin-3-yl-carbamic acid phenyl ester (1.099 g, 39%). MS (ESI⁺): calcd for C₁₃H₉N₃O₃ m/z 255.06, found 256.1 (M+H)⁺.

15 Intermediates 21 to 23 were prepared using methods analogous to those described for Intermediate 20.


Intermediate 21: Isoxazolo[5,4-b]pyridin-3-yl-carbamic acid phenyl ester.

20 MS (ESI⁺): calcd for C₁₃H₉N₃O₃ m/z 255.06, found 256.1 (M+H)⁺.

Intermediate 22: Isoxazolo[4,5-c]pyridin-3-yl-carbamic acid phenyl ester.


MS (ESI⁺): calcd for C₁₃H₉N₃O₃ m/z 255.06, found 256.1 (M+H)⁺.

25

Intermediate 23: Isoxazolo[4,5-b]pyridin-3-yl-carbamic acid phenyl ester.

MS (ESI⁺): calcd for C₁₃H₉N₃O₃ m/z 255.06, found 256.1 (M+H)⁺.

5 Example 1: 3-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide.



Step A: 3-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid tert-butyl ester. 1-Boc-3-methylene piperidine (842 mg, 4.27 mmol) was
10 degassed (neat) for 15 minutes and then treated with a THF solution of 9-BBN (0.5 M in THF, 8.6 mL, 4.3 mmol). The reaction mixture was refluxed for 2 h, then cooled to rt. The reaction mixture was then added, via cannula, to a preformed solution of 5-bromo-2,2-difluoro-1,3-benzodioxole (1.00 g, 4.22 mmol), Pd(dppf)Cl₂•CH₂Cl₂ (94 mg, 0.128 mmol), and potassium carbonate (746 mg, 5.40 mmol) in DMF/H₂O (10
15 mL/1 mL). The resultant mixture was heated at 60 °C for 18 h, cooled to rt, poured into water, basified to pH 11 with 1 N NaOH, and extracted with EtOAc (3x). The organic layers were combined, dried (Na₂SO₄), and concentrated. The crude residue was purified (FCC) to give 3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid tert-butyl ester (1.00 g, 67%).

20 Step B: 3-[(2,2-Difluoro-1,3-benzodioxol-5-yl)methyl]-piperidine hydrochloride.
To a solution of 3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid tert-butyl ester (1.0 g, 2.8 mmol) in CH₂Cl₂ (11 mL) was added HCl (4 M in dioxane, 4.2 mL, 16.9 mmol). The reaction mixture was stirred at r.t. overnight and then concentrated to give 3-[(2,2-difluoro-1,3-benzodioxol-5-yl)methyl]-piperidine as
25 the hydrochloride salt (791 mg, 96%). The crude product was used without further purification.


Step C: 3-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide. To a solution of 3-[(2,2-difluoro-1,3-benzodioxol-5-yl)methyl]-piperidine hydrochloride (100 mg, 0.34 mmol) in CH₃CN (2 mL) was added Et₃N
30 (0.07 mL, 0.52 mmol). To this reaction mixture, pyridin-3-yl-carbamic acid phenyl ester (77.0 mg, 1.05 mmol) was added and the reaction mixture was heated overnight at 35 °C. The reaction mixture was then concentrated and the crude residue was purified (FCC, 2 N NH₃ in MeOH/DCM) to give 3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide (120 mg,

5 93%). MS (ESI⁺): calcd for C₁₉H₁₉F₂N₃O₃ m/z 375.14, found 376.2 (M+H)⁺. ¹H NMR (d₄-methanol): 8.54 (d, *J* = 2.3, 1H), 8.17-8.13 (m, 1H), 7.86 (ddd, *J* = 8.4, 2.6, 1.4, 1H), 7.34-7.30 (m, 1H), 7.10-7.06 (m, 2H), 6.99 (dd, *J* = 8.2, 1.5, 1H), 4.05-3.95 (m, 2H), 3.04-2.98 (m, 1H), 2.74-2.64 (m, 2H), 2.58-2.53 (m, 1H), 1.86-1.74 (m, 3H), 1.56-1.47 (m, 1H), 1.30-1.22 (m, 1H).

10

The compounds in Examples 2 to 9 were prepared using methods analogous to those described in Example 1.


Example 2: N-Pyrimidin-4-yl-3-[4-(trifluoromethyl)phenyl]methyl}piperidine-1-
15 carboxamide.



20

¹H NMR (d₄-methanol): 8.76 (d, *J* = 1.0, 1H), 8.53 (d, *J* = 6.0, 1H), 8.00 (dd, *J* = 1.5, 6.0, 1H), 7.57 (d, *J* = 8.0, 2H), 7.37 (br s, 1H), 7.28 (d, *J* = 8.0, 2H), 4.04 (d, *J* = 12.5, 1H), 3.90 (d, *J* = 12.5, 1H), 3.75-3.02 (m, 1H), 2.74-2.70 (m, 2H), 2.56 (dd, *J* = 8, 13.5, 1H), 1.96-1.75 (m, 3H), 1.57-1.47 (m, 1H), 1.30-1.18 (m, 1H).

Example 3: N-Pyrimidin-4-yl-3-[3-(trifluoromethyl)phenyl]methyl}piperidine-1-
carboxamide.




25

MS (ESI⁺): calcd for C₁₈H₁₉F₃N₄O m/z 364.15, found 365.2 (M+H)⁺. ¹H NMR (CDCl₃): 8.76 (d, *J* = 1.0, 1H), 8.53 (d, *J* = 5.5, 1H), 8.00 (dd, *J* = 1.5, 6.0, 1H), 7.50 (d, *J* = 7.5, 1H), 7.43-7.42 (m, 2H), 7.38-7.35 (m, 2H), 4.04 (d, *J* = 12.5, 1H), 3.90 (d, *J* = 12.5, 1H), 3.07-3.01 (m, 1H), 2.76-2.70 (m, 2H), 2.56 (dd, *J* = 8.0, 13.5, 1H), 1.97-1.75 (m, 3H), 1.57-1.46 (m, 1H), 1.29-1.19 (m, 1H).


30

Example 4: 3-[(4-Fluorophenyl)methyl]-N-pyrimidin-4-ylpiperidine-1-carboxamide.




¹H NMR (CDCl₃): 8.76 (d, *J* = 1.0, 1H), 8.52 (d, *J* = 6.0, 1H), 8.00 (dd, *J* = 1.0, 6.0, 1H), 7.40 (br s, 1H), 7.11-7.10 (m, 2H), 7.01-6.97 (m, 2H), 3.99 (d, *J* = 12.5, 1H), 3.91 (d, *J* = 12.5, 1H), 3.04-2.99 (m, 1H), 2.69-2.63 (m, 1H), 2.62 (dd, *J* = 6.5, 14.0, 1H), 2.49 (dd, *J* = 7.5, 14.0, 1H), 1.87-1.71 (m, 3H), 1.56-1.44 (m, 1H), 1.29-1.14 (m, 1H).

Example 5: N-1,2-Benzisoxazol-3-yl-3-([4-(trifluoromethyl)phenyl]methyl)-piperidine-1-carboxamide.




¹H NMR (CDCl₃): 9.04 (s, 1H), 8.05 (d, *J* = 8.5, 1H), 7.58-7.54 (m, 3H), 7.45 (d, *J* = 8.5, 1H), 7.33-7.28 (m, 3H), 4.21-4.14 (m, 2H), 3.09-3.05 (m, 1H), 2.84-2.75 (m, 2H), 2.56 (dd, *J* = 8.0, 13.5, 1H), 1.99-1.91 (m, 1H), 1.88-1.76 (m, 2H), 1.64-1.50 (m, 1H), 1.30-1.17 (m, 1H).

Example 6: N-1,2-Benzisoxazol-3-yl-3-([3-(trifluoromethyl)phenyl]methyl)-piperidine-1-carboxamide.



MS (ESI⁺): calcd for C₂₁H₂₀F₃N₃O₂ *m/z* 403.15, found 404.2 (M+H)⁺. ¹H NMR (CDCl₃): 9.02 (s, 1H), 8.06 (d, *J* = 8.5, 1H), 7.57-7.54 (m, 1H), 7.48-7.45 (m, 3H), 7.41-7.36 (m, 2H), 7.31-7.28 (m, 1H), 4.20 (d, *J* = 10.5, 1H), 4.15 (d, *J* = 11.5, 1H), 3.10-3.05 (m, 1H), 2.83-2.79 (m, 2H), 2.55 (dd, *J* = 8.5, 13.5, 1H), 1.99-1.90 (m, 1H), 1.87-1.77 (m, 2H), 1.64-1.51 (m, 1H), 1.29-1.17 (m, 1H).

Example 7: N-1,2-Benzisoxazol-3-yl-3-[(4-fluorophenyl)methyl]piperidine-1-carboxamide.




5

MS (ESI⁺): calcd for C₂₀H₂₀FN₃O₂ m/z 353.15, found 354.2 (M+H)⁺. ¹H NMR (CDCl₃): 8.50 (s, 1H), 8.06 (d, *J* = 8.0, 1H), 7.55-7.52 (m, 1H), 7.45 (d, *J* = 8.5, 1H), 7.29-7.28 (m, 1H), 7.14-7.12 (m, 2H), 7.00-6.96 (m, 2H), 4.17-4.10 (m, 2H), 3.10-3.04 (m, 1H), 2.79-2.76 (m, 1H), 2.70 (dd, *J* = 6.5, 13.5, 1H), 1.94-1.75 (m, 3H), 1.68-1.52 (m, 1H), 1.28-1.15 (m, 1H).

10

Example 8: N-Isoxazol-3-yl-3-{[4-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide.




15

¹H NMR (CDCl₃): 8.93 (s, 1H), 8.21 (d, *J* = 2.0, 1H), 7.55 (d, *J* = 8.0, 2H), 7.28 (d, *J* = 8.0, 2H), 7.05 (d, *J* = 2.0, 1H), 4.12 (d, *J* = 12.0, 1H), 4.03 (d, *J* = 13.0, 1H), 3.04-2.99 (m, 1H), 2.77-2.72 (m, 2H), 2.54 (dd, *J* = 8.5, 13.5, 1H), 1.94 (br s, 1H), 1.86-1.73 (m, 2H), 1.59-1.45 (m, 1H), 1.29-1.15 (m, 1H).

20


Example 9: N-Isoxazol-3-yl-3-{[3-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide.



25

MS (ESI⁺): calcd for C₁₇H₁₈F₃N₃O₃ m/z 353.14, found 354.2 (M+H)⁺. ¹H NMR (CDCl₃): 9.01 (s, 1H), 8.20 (d, *J* = 2.0, 1H), 7.48 (d, *J* = 7.5, 1H), 7.44-7.39 (m, 2H), 7.34 (d, *J* = 7.5, 1H), 7.03 (d, *J* = 2.0, 1H), 4.14 (d, *J* = 13.0, 1H), 4.05 (d, *J* = 13.0, 1H), 3.04-2.99 (m, 1H), 2.78-2.70 (m, 2H), 2.55-2.50 (m, 1H), 1.90-1.86 (m, 1H), 1.82-1.75 (m, 2H), 1.61-1.43 (m, 1H), 1.28-1.16 (m, 1H).

Example 10: 3-[3-(4-Fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide.



5

Step A: 3-(4-Fluoro-phenoxy)-bromobenzene. To a solution of 1-bromo-3-iodo-benzene (1.00 g, 3.55 mmol), 4-fluorophenol (462 mg, 3.59 mmol), N,N-dimethylglycine hydrochloride (149 mg, 1.06 mmol), and cesium carbonate (2.28 g, 7.00 mmol) in DMA was added CuI (66.8 mg, 0.351 mmol). The reaction mixture was heated at 90 °C for 48 h, cooled to rt, and then poured into water and extracted with EtOAc (3x). The organic layers were combined, dried (Na₂SO₄), and concentrated. The crude residue was purified (FCC) to give the title compound as a colorless oil (319 mg, 32%).

Step B: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid tert-butyl ester. The title compound was prepared using methods analogous to those described in Example 1, Step A.


Step C: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine hydrochloride. The title compound was prepared using methods analogous to those described in Example 1, Step B.

Step D: 3-[3-(4-Fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide. The title compound was prepared using methods analogous to those described in Example 1, Step C. MS (ESI⁺): calcd for C₂₄H₂₄FN₃O₂ m/z 405.18, found 406.2 (M+H)⁺. ¹H NMR (d₄-methanol): 8.54-8.53 (m, 1H), 8.15 (dd, J = 4.8, 1.4, 1H), 7.87 (ddd, J = 8.4, 2.6, 1.4, 1H), 7.34-7.31 (m, 1H), 7.25 (t, J = 7.9, 1H), 7.08-7.03 (m, 2H), 7.00-6.94 (m, 3H), 6.83-6.81 (m, 1H), 6.79-6.77 (m, 1H), 4.05-4.01 (m, 1H), 4.00-3.95 (m, 1H), 3.03-2.95 (m, 1H), 2.69 (dd, J = 13.2, 10.1, 1H), 2.64-2.60 (m, 1H), 2.53-2.48 (m, 1H), 1.85-1.71 (m, 3H), 1.55-1.46 (m, 1H), 1.29-1.21 (m, 1H).

The compounds in Examples 11 to 15 were prepared using methods analogous to those described in Example 10.

Example 11: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide.


5



MS (ESI⁺): calcd for C₂₄H₂₃ClFN₃O₂ m/z 439.14, found 440.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.58 (s, 1H), 8.34 (s, 1H), 8.28 (d, *J* = 5.3, 1H), 7.56 (d, *J* = 5.3, 1H), 7.29 (t, *J* = 7.8, 1H), 7.22-7.14 (m, 2H), 7.07-7.01 (m, 2H), 6.97 (d, *J* = 7.6, 1H), 6.86-6.82 (m, 1H), 6.82-6.76 (m, 1H), 3.93 (d, *J* = 13.2, 2H), 2.95-2.87 (m, 1H), 2.68-2.43 (m, 3H), 1.78-1.59 (m, 3H), 1.47-1.29 (m, 1H), 1.23-1.07 (m, 1H).

Example 12: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide.

15




MS (ESI⁺): calcd for C₂₅H₂₄FN₅O₂ m/z 445.19, found 446.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.80 (s, 1H), 8.64-8.61 (m, 1H), 8.16 (dd, *J* = 9.2, 1.4, 1H), 7.78 (s, 1H), 7.34 (dd, *J* = 9.2, 4.5, 1H), 7.29 (t, *J* = 7.8, 1H), 7.21-7.14 (m, 2H), 7.07-7.01 (m, 2H), 6.98 (d, *J* = 7.6, 1H), 6.87-6.83 (m, 1H), 6.82-6.77 (m, 1H), 3.97 (d, *J* = 12.9, 2H), 2.91 (s, 1H), 2.70-2.44 (m, 3H), 1.81-1.59 (m, 3H), 1.41 (m, 1H), 1.24-1.12 (m, 1H).

Examples 13 and 14 were resolved using chiral column chromatography:

Jasco Preparative SFC System using a Phenomenex Lux II column (5 μm 250 x 21 mm, L x I.D.) @ 40 °C, 150 bar, 8.3 mL/min ethanol with 0.2% triethylamine and 33.3 mL/min CO₂, detection at 220 nm. Retention times: Example 13, (+)-enantiomer: 33-38 minutes; Example 14, (–)-enantiomer: 42-48 minutes.


Example 13: (+)-3-[3-(4-Chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide.

5




MS (ESI⁺): calcd for C₂₄H₂₄ClN₃O₂ m/z 421.16, found 422.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.64-8.58 (m, 2H), 8.13 (dd, *J* = 4.6, 1.5, 1H), 7.86-7.81 (m, 1H), 7.40 (d, *J* = 9.0, 2H), 7.32 (t, *J* = 7.8, 1H), 7.24 (dd, *J* = 8.3, 4.7, 1H), 7.05-6.98 (m, 3H), 6.91-6.88 (m, 1H), 6.88-6.84 (m, 1H), 4.01-3.92 (m, 2H), 2.90-2.82 (m, 1H), 2.63-2.43 (m, 3H), 1.75-1.61 (m, 3H), 1.45-1.31 (m, 1H), 1.22-1.07 (m, 1H); [α]²³_D + 22.4 (c 0.15, MeOH).

Example 14: (-)-3-[3-(4-Chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide.



MS (ESI⁺): calcd for C₂₄H₂₄ClN₃O₂ m/z 421.16, found 422.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.64-8.58 (m, 2H), 8.13 (dd, *J* = 4.6, 1.5, 1H), 7.86-7.81 (m, 1H), 7.40 (d, *J* = 9.0, 2H), 7.32 (t, *J* = 7.8, 1H), 7.24 (dd, *J* = 8.3, 4.7, 1H), 7.05-6.98 (m, 3H), 6.91-6.88 (m, 1H), 6.88-6.84 (m, 1H), 4.01-3.92 (m, 2H), 2.90-2.82 (m, 1H), 2.63-2.43 (m, 3H), 1.75-1.61 (m, 3H), 1.45-1.31 (m, 1H), 1.22-1.07 (m, 1H); [α]²³_D - 22.0 (c 0.15, MeOH).

Example 15: 3-[3-(4-Trifluoromethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide.




MS (ESI⁺): calcd for C₂₅H₂₄F₃N₃O₃ m/z 471.18, found 472.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.35 (br s, 1H), 8.27 (br s, 1H), 7.99-7.95 (m, 1H), 7.31-7.27 (m, 1H), 7.24-7.20 (m, 1H), 7.19-7.15 (m, 2H), 6.99 (d, *J* = 9.1, 2H), 6.95 (d, *J* = 7.5, 1H),

- 5 6.88-6.84 (m, 2H), 6.26 (s, 1H), 3.94-3.88 (m, 1H), 3.87-3.82 (m, 1H), 3.09-3.01 (m, 1H), 2.69 (dd, $J = 13.2, 9.8$, 1H), 2.62 (dd, $J = 13.8, 7.2$, 1H), 2.54 (dd, $J = 13.8, 7.0$, 1H), 1.92-1.83 (m, 2H), 1.81-1.74 (m, 1H), 1.59-1.50 (m, 1H), 1.29-1.20 (m, 1H).


The compounds in Examples 16 to 19 were prepared using methods
10 analogous to those described in Example 1.

Example 16: 3-Quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide.



- 15 MS (ESI⁺): calcd for C₂₁H₂₂N₄O m/z 346.18, found 347.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.77 (d, $J = 2.2$, 1H), 8.36 (d, $J = 2.5$, 1H), 8.26 (dd, $J = 4.7, 1.3$, 1H), 8.09 (d, $J = 8.5$, 1H), 7.96-7.91 (m, 2H), 7.76 (dd, $J = 8.2, 1.0$, 1H), 7.71-7.66 (m, 1H), 7.56-7.51 (m, 1H), 7.21 (dd, $J = 8.3, 4.7$, 1H), 6.33 (s, 1H), 4.08-4.01 (m, 1H), 3.89-3.81 (m, 1H), 3.11-3.02 (m, 1H), 2.87 (dd, $J = 13.9, 6.7$, 1H), 2.81-2.67 (m, 2H),
20 2.06-1.95 (m, 1H), 1.92-1.84 (m, 1H), 1.84-1.75 (m, 1H), 1.61-1.49 (m, 1H), 1.35-1.24 (m, 1H).


Example 17: 3-(3,4-Dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide.



- 25 MS (ESI⁺): calcd for C₁₈H₁₉Cl₂N₃O m/z 363.09, found 364.1 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.39 (d, $J = 2.4$, 1H), 8.27 (dd, $J = 4.7, 1.4$, 1H), 7.95 (ddd, $J = 8.4, 2.6, 1.5$, 1H), 7.36 (d, $J = 8.2$, 1H), 7.27 (s, 1H), 7.22 (dd, $J = 8.3, 4.7$, 1H), 7.01 (dd, $J = 8.2, 2.0$, 1H), 6.28 (s, 1H), 4.00-3.93 (m, 1H), 3.87-3.80 (m, 1H), 3.08-3.00 (m, 1H), 2.67 (dd, $J = 13.2, 10.0$, 1H), 2.64-2.57 (m, 1H), 2.48 (dd, $J = 13.8, 7.4$, 1H), 1.90-1.73 (m, 3H), 1.60-1.48 (m, 1H), 1.27-1.16 (m, 1H).
- 30

5


Example 18: 3-(3-Trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide.



MS (ESI⁺): calcd for C₁₉H₂₀F₃N₃O₂ m/z 379.15, found 380.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.36 (d, *J* = 2.5, 1H), 8.26 (dd, *J* = 4.7, 1.3, 1H), 7.95 (ddd, *J* = 8.3, 2.5, 1.4, 1H), 7.32 (t, *J* = 7.9, 1H), 7.22 (dd, *J* = 8.3, 4.7, 1H), 7.12-7.07 (m, 2H), 7.02 (s, 1H), 6.34 (s, 1H), 3.98-3.92 (m, 1H), 3.91-3.84 (m, 1H), 3.06-2.98 (m, 1H), 2.72-2.64 (m, 2H), 2.54 (dd, *J* = 13.7, 7.5, 1H), 1.92-1.81 (m, 2H), 1.80-1.74 (m, 1H), 1.59-1.49 (m, 1H), 1.28-1.18 (m, 1H).

15

Example 19: 3-(4-Trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide.




MS (ESI⁺): calcd for C₁₉H₂₀F₃N₃O₂ m/z 379.15, found 380.2 (M+H)⁺. ¹H NMR (d₆-DMSO): 8.37 (d, *J* = 2.4, 1H), 8.26 (dd, *J* = 4.7, 1.4, 1H), 7.94 (ddd, *J* = 8.4, 2.6, 1.5, 1H), 7.24-7.12 (m, 5H), 6.32 (s, 1H), 3.98-3.91 (m, 1H), 3.89-3.81 (m, 1H), 3.08-2.99 (m, 1H), 2.73-2.62 (m, 2H), 2.52 (dd, *J* = 13.8, 7.3, 1H), 1.91-1.73 (m, 3H), 1.60-1.48 (m, 1H), 1.29-1.16 (m, 1H).

The compounds in Examples 20 to 23 can be synthesized using the synthetic schemes given in Example 10.

Example 20: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide.


30

5




Example 21: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide.


10



Example 22: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide.




Example 23: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-ylamide.



Example 24: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide.

20




To a solution consisting of 1 equivalent of imidazo[1,2-a]pyridin-7-ylamine in CH_2Cl_2 is added 1 equivalent of N,N'-disuccinimidyl carbonate and 1 equivalent of 4-dimethylaminopyridine (DMAP). The reaction mixture is stirred at rt for 6 h, then
treated with 1 equivalent of 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine hydrochloride

- 5 and 2 equivalents of diisopropylethylamine and stirred for an additional 16 h at rt. The reaction mixture is diluted with EtOAc and extracted with H₂O then saturated aqueous NaCl. The organic layer is dried (MgSO₄) and concentrated. The crude residue is purified (FCC, 2 N NH₃ in MeOH/DCM) to give 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide.


10

The compounds in Examples 25 to 28 can be synthesized using the synthetic scheme given in Example 24.


Example 25: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid
15 imidazo[1,2-a]pyridin-5-ylamide.



Example 26: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid
20 imidazo[1,2-a]pyrimidin-7-ylamide.




Example 27: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid
25 imidazo[1,2-a]pyrimidin-5-ylamide.




Example 28: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid
imidazo[1,2-c]pyrimidin-7-ylamide.

5




The compounds in Examples 29 to 32 can be synthesized using the synthetic schemes given in Example 10.

10 Example 29: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-amide.




15 Example 30: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide.




Example 31: 3-[3-(4-Fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide.

20



Example 32: 3-[3-(4-Trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide.


25



5

The compound in Example 33 can be synthesized using the synthetic schemes given in Example 1.

10 Example 33: 3-(4-Chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide.



15 Biological Testing:

- Assay Method 1

A. Transfection of Cells with Human FAAH

A 10-cm tissue culture dish with a confluent monolayer of SK-N-MC cells was split 2 days (d) prior to transfection. Using sterile technique, the media was removed and the cells were detached from the dish by the addition of trypsin. One fifth of the cells were then placed onto a new 10-cm dish. Cells were grown in a 37 °C incubator with 5% CO₂ in Minimal Essential Media Eagle with 10% Fetal Bovine Serum. After 2 d, cells were approximately 80% confluent. These cells were removed from the dish with trypsin and pelleted in a clinical centrifuge. The pellet was re-suspended in 400 µL complete media and transferred to an electroporation cuvette with a 0.4 cm gap between the electrodes. Supercoiled human FAAH cDNA (1 µg) was added to the cells and mixed. The voltage for the electroporation was set at 0.25 kV, and the capacitance was set at 960 µF. After electroporation, the cells were diluted into complete media (10 mL) and plated onto four 10-cm dishes. Because of the variability in the efficiency of electroporation, four different concentrations of cells were plated. The ratios used were 1:20, 1:10, and 1:5, with

25

30

5 the remainder of the cells being added to the fourth dish. The cells were allowed to recover for 24 h before adding the selection media (complete media with 600 $\mu\text{g/mL}$ G418). After 10 d, dishes were analyzed for surviving colonies of cells. Dishes with well-isolated colonies were used. Cells from individual colonies were isolated and tested. The clones that showed the most FAAH activity, as measured by
10 anandamide hydrolysis, were used for further study.

B. FAAH Assay

T84 frozen cell pellets or transfected SK-N-MC cells (contents of 1 x 15 cm culture dishes) were homogenized in 50 mL of FAAH assay buffer (125 mM Tris, 1mM EDTA, 0.2% Glycerol, 0.02% Triton X-100, 0.4 mM Hepes, pH 9). The assay
15 mixture consisted of 50 μL of the cell homogenate, 10 μL of the test compound, and 40 μL of anandamide [$1\text{-}^3\text{H}$ -ethanolamine] (^3H -AEA, Perkin-Elmer, 10.3 Ci/mmol), which was added last, for a final tracer concentration of 80 nM. The reaction mixture was incubated at rt for 1 h. During the incubation, 96-well Multiscreen filter plates (catalog number MAFCNOB50; Millipore, Bedford, MA, USA) were loaded with 25 μL
20 of activated charcoal (Multiscreen column loader, catalog number MACL09625, Millipore) and washed once with 100 μL of MeOH. Also during the incubation, 96-well DYNEX MicroLite plates (catalog number NL510410) were loaded with 100 μL of MicroScint40 (catalog number 6013641, Packard Bioscience, Meriden, CT, USA). After the 1 h incubation, 60 μL of the reaction mixture were transferred to the
25 charcoal plates, which were then assembled on top of the DYNEX plates using Centrifuge Alignment Frames (catalog number MACF09604, Millipore). The unbound labeled ethanolamine was centrifuged through to the bottom plate (5 min at 2000 rpm), which was preloaded with the scintillant, as described above. The plates were sealed and left at rt for 1 h before counting on a Hewlett Packard TopCount.

30 • Assay Method 2

A. Transfection of Cells with Rat FAAH

A 10-cm tissue culture dish with a confluent monolayer of SK-N-MC cells was split 2 days (d) prior to transfection. Using sterile technique, the media was removed and the cells were detached from the dish by the addition of trypsin. One fifth of the
35 cells were then placed onto a new 10-cm dish. Cells were grown in a 37 °C incubator with 5% CO_2 in Minimal Essential Media Eagle with 10% Fetal Bovine

5 Serum. After 2 d, cells were approximately 80% confluent. These cells were removed from the dish with trypsin and pelleted in a clinical centrifuge. The pellet was re-suspended in 400 μ L complete media and transferred to an electroporation cuvette with a 0.4 cm gap between the electrodes. Supercoiled rat FAAH cDNA (1 μ g) was added to the cells and mixed. The voltage for the electroporation was set at
10 0.25 kV, and the capacitance was set at 960 μ F. After electroporation, the cells were diluted into complete media (10 mL) and plated onto four 10-cm dishes. Because of the variability in the efficiency of electroporation, four different concentrations of cells were plated. The ratios used were 1:20, 1:10, and 1:5, with the remainder of the cells being added to the fourth dish. The cells were allowed to recover for 24 h
15 before adding the selection media (complete media with 600 μ g/mL G418). After 10 d, dishes were analyzed for surviving colonies of cells. Dishes with well-isolated colonies were used. Cells from individual colonies were isolated and tested. The clones that showed the most FAAH activity, as measured by anandamide hydrolysis, were used for further study.

20 B. FAAH Assay

T84 frozen cell pellets or transfected SK-N-MC cells (contents of 1 x 15 cm culture dishes) were homogenized in 50 mL of FAAH assay buffer (125 mM Tris, 1 mM EDTA, 0.2% Glycerol, 0.02% Triton X-100, 0.4 mM Hepes, pH 9). The assay mixture consisted of 50 μ L of the cell homogenate, 10 μ L of the test compound, and
25 40 μ L of anandamide [1-³H-ethanolamine] (³H-AEA, Perkin-Elmer, 10.3 Ci/mmol), which was added last, for a final tracer concentration of 80 nM. The reaction mixture was incubated at rt for 1 h. During the incubation, 96-well Multiscreen filter plates (catalog number MAFCNOB50; Millipore, Bedford, MA, USA) were loaded with 25 μ L of activated charcoal (Multiscreen column loader, catalog number MACL09625,
30 Millipore) and washed once with 100 μ L of MeOH. Also during the incubation, 96-well DYNEX MicroLite plates (catalog number NL510410) were loaded with 100 μ L of MicroScint40 (catalog number 6013641, Packard Bioscience, Meriden, CT, USA). After the 1 h incubation, 60 μ L of the reaction mixture were transferred to the charcoal plates, which were then assembled on top of the DYNEX plates using
35 Centrifuge Alignment Frames (catalog number MACF09604, Millipore). The unbound labeled ethanolamine was centrifuged through to the bottom plate (5 min at

5 2000 rpm), which was preloaded with the scintillant, as described above. The plates were sealed and left at rt for 1 h before counting on a Hewlett Packard TopCount.

Results for compounds tested in these assays are summarized in Table 1, as an average of results obtained. Compounds were tested in free base, hydrochloride
 10 salt, and/or trifluoroacetic acid salt forms. Where activity is shown as greater than (>) a particular value, the value is the solubility limit of the compound in the assay medium or the highest concentration tested in the assay.


Table 1

Ex.	Assay 1 IC₅₀ (μM)	Assay 2 IC₅₀ (μM)		Ex.	Assay 1 IC₅₀ (μM)	Assay 2 IC₅₀ (μM)
1	0.410	0.604		11	0.087	0.008
2	8.000	>10		12	0.056	0.007
3	8.000	10.000		13	0.019	0.031
4	10.000	>10		14	0.073	0.027
5	2.000	0.183		15	0.011	0.017
6	0.645	0.050		16	1.000	>10
7	2.000	0.152		17	0.048	0.180
8	10.000	>10		18	0.190	0.450
9	7.000	8.999		19	0.430	1.600
10	0.018	0.019				

15 While the invention has been illustrated by reference to exemplary and preferred embodiments, it will be understood that the invention is intended not to be limited to the foregoing detailed description, but to be defined by the appended claims as properly construed under principles of patent law.

5 What is claimed is:

1. A chemical entity selected from compounds of Formula (I):



wherein:

10 Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently
 15 unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

(i) phenyl unsubstituted or substituted with one or two R^a moieties;

20 where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃, -OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c, -SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;
 or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -OCF₂O-;

25 where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b and R^c taken together with the atoms of attachment form a 4-8 membered ring;

(ii) phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties, wherein:

30 L is a linker selected from the group consisting of -(CH₂)₁₋₆-, -CH=CH-, -O-, -C₁₋₆alkyl-O-C₁₋₆alkyl-, -C₁₋₆alkyl-N(C₁₋₆alkyl)-C₁₋₆alkyl-, -C₁₋₆alkyl-S(O)₀₋₂C₁₋₆alkyl-, -C≡C-, -C(O)-, or a covalent bond;

Ar³ is:

(a) phenyl unsubstituted or substituted with one or two R^a moieties;

- 5 (b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom
10 selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;
- and pharmaceutically acceptable salts of compounds of Formula (I),
- 15 pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I).

2. A chemical entity as in claim 1, wherein Ar¹ is isoxazolo[5,4-c]pyridin-3-yl, isoxazolo[4,5-c]pyridin-3-yl, isoxazolo[4,5-b]pyridin-3-yl, isoxazolo[5,4-b]pyridin-3-yl,
20 imidazo[1,2-a]pyridin-8-yl, imidazo[1,2-a]pyridin-7-yl, imidazo[1,2-a]pyridin-6-yl, imidazo[1,2-a]pyridin-5-yl, imidazo[1,2-b]pyridazin-3-yl, imidazo[1,2-a]pyridin-3-yl, imidazo[1,2-a]pyridin-2-yl, imidazo[1,2-a]pyrimidin-7-yl, imidazo[1,2-a]pyrimidin-5-yl, imidazo[1,2-c]pyrimidin-7-yl, benzooxazol-6-yl, 1H-indazol-7-yl, quinolin-8-yl, isoquinolin-5-yl, isoquinolin-4-yl, 1,1-dioxo-1H-1λ⁶-benzo[d]isothiazol-3-yl, 1H-
25 pyrrolo[2,3-b]pyridin-5-yl, 1H-pyrrolo[2,3-b]pyridin-4-yl, 1H-pyrrolo[3,2-b]pyridin-6-yl, 2-methyl-benzothiazol-6-yl, 1-isopropyl-1H-pyrazolo[3,4-b]pyridin-5-yl, 2-methyl-benzooxazol-5-yl, 1H-indazol-7-yl, 1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl, 2-methyl-2H-indazol-4-yl, benzooxazole-2-yl group, 6-[1,2,3]triazol-1-yl-pyridin-3-yl, 6-[1,2,4]triazol-4-yl-pyridin-3-yl, 6-[1,2,4]triazol-1-yl-pyridin-3-yl, 6-[1,2,3]triazol-2-yl-
30 pyridin-3-yl, 4-[1,2,3]triazol-2-yl-phenyl, 4-[1,2,3]triazol-1-yl-phenyl, 5-methyl-[1,3,4]oxadiazol-2-yl, 2-phenyl-pyrimidin-5-yl, 3-pyridyl, benzisoxazol-3-yl, pyrimidin-4-yl, isoxazol-3-yl, 6-fluorobenzo[d]isoxazol-3-yl, 3-phenyl-[1,2,4]thiadiazol-5-yl, 1H-tetrazol-5-yl, benzo[1,2,5]thiadiazol-4-yl, benzo[1,2,5]oxadiazol-4-yl, thiophen-2-yl, thiophen-3-yl, 6-chloro-pyridazin-3-yl, pyrazin-2-yl, 1H-benzotriazol-5-yl,
35 [1,5]naphthyridin-2-yl, benzothiazol-6-yl, or 1H-pyrazol-3-yl.


- 5 3. A chemical entity as in claim 1, wherein Ar¹ is pyrimidine, isoxazole, benzisoxazole, or pyridine, each group optionally substituted with one or two R^a.
4. A chemical entity as in claim 1, wherein Ar² is phenyl substituted with one or two R^a moieties.
- 10 5. A chemical entity as in claim 4, wherein each R^a moiety is independently selected from the group consisting of: chloro, cyano, isobutyl, methylsulfanyl, methanesulfonyl, trifluoromethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, fluoro, methyl, methoxy, tert-butyl, bromo, methoxycarbonyl, cyanomethyl, methoxycarbonylmethyl, trifluoromethanesulfonyl, trifluoromethanesulfanyl, and butyl; or two adjacent R^a moieties taken together form -OCH₂O- or -OCF₂O-.
- 15 6. A chemical entity as in claim 4, wherein each R^a moiety is independently selected from the group consisting of: F, Cl, Br, CF₃, or two adjacent R^a moieties taken together form -OCF₂O-.
- 20 7. A chemical entity as in claim 1, wherein Ar² is phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties.
- 25 8. A chemical entity as in claim 7, wherein L is -CH₂CH₂-, -O-, -OCH₂-, or -C≡C-.
9. A chemical entity as in claim 7, wherein L is -O-.
- 30 10. A chemical entity as in claim 7, wherein Ar³ is phenyl, unsubstituted or substituted with one or two R^a moieties.
11. A chemical entity as in claim 10, wherein each R^a moiety is independently selected from the group consisting of: chloro, cyano, isobutyl, methylsulfanyl, methanesulfonyl, trifluoromethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, fluoro, methyl, methoxy, tert-butyl, bromo, methoxycarbonyl, cyanomethyl,
- 35

- 5 methoxycarbonylmethyl, trifluoromethanesulfonyl, trifluoromethanesulfanyl, and butyl; or two adjacent R^a moieties taken together form –OCH₂O– or –OCF₂O–.
12. A chemical entity as in claim 10, wherein each R^a moiety is independently fluoro.
- 10 13. A chemical entity selected from the group consisting of:
3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
N-pyrimidin-4-yl-3-[[4-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
15 N-pyrimidin-4-yl-3-[[3-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
3-[(4-fluorophenyl)methyl]-N-pyrimidin-4-ylpiperidine-1-carboxamide;
N-1,2-benzisoxazol-3-yl-3-[[4-(trifluoromethyl)-phenyl]-methyl]-piperidine-1-carboxamide;
N-1,2-benzisoxazol-3-yl-3-[[3-(trifluoromethyl)-phenyl]-methyl]-piperidine-1-
20 carboxamide;
N-1,2-benzisoxazol-3-yl-3-[(4-fluorophenyl)methyl]piperidine-1-carboxamide;
N-isoxazol-3-yl-3-[[4-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
N-isoxazol-3-yl-3-[[3-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
3-[3-(4-fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide;
25 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide;
(+)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
30 (–)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-trifluoromethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide;
3-quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3,4-dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
35 3-(3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;

- 5 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide;
10 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide;
15 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-5-ylamide;
20 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-c]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-amide;
25 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide;
3-[3-(4-trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
30 and pharmaceutically acceptable salts thereof.

14. A pharmaceutical composition for treating a disease, disorder, or medical condition mediated by FAAH activity, comprising:

- (a) an effective amount of at least one chemical entity selected from
35 compounds of Formula (I):



wherein:

Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

(i) phenyl unsubstituted or substituted with one or two R^a moieties;

where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃, -OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c, -SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;

or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -

OCF₂O-;

where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b and R^c taken together with the atoms of attachment form a 4-8 membered ring;

(ii) phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties, wherein:

L is a linker selected from the group consisting of -(CH₂)₁₋₆-, -CH=CH-, -O-, -C₁₋₆alkyl-O-C₁₋₆alkyl-, -C₁₋₆alkyl-N(C₁₋₆alkyl)-C₁₋₆alkyl-, -C₁₋₆alkyl-S(O)₀₋₂C₁₋₆alkyl-, -C≡C-, -C(O)-, or a covalent bond;

Ar³ is:

(a) phenyl unsubstituted or substituted with one or two R^a moieties;

(b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or

(c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or

- 5 (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;
- 10 and pharmaceutically acceptable salts of compounds of Formula (I), pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I) ; and
- (b) a pharmaceutically acceptable excipient.
- 15 15. A pharmaceutical composition according to claim 14, wherein said at least one chemical entity is selected from the group consisting of:
- 3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
- N-pyrimidin-4-yl-3-{{4-(trifluoromethyl)phenyl}methyl}piperidine-1-carboxamide;
- 20 N-pyrimidin-4-yl-3-{{3-(trifluoromethyl)phenyl}methyl}piperidine-1-carboxamide;
- 3-{{(4-fluorophenyl)methyl}-N-pyrimidin-4-yl}piperidine-1-carboxamide;
- N-1,2-benzisoxazol-3-yl-3-{{4-(trifluoromethyl)-phenyl}-methyl}-piperidine-1-carboxamide;
- N-1,2-benzisoxazol-3-yl-3-{{3-(trifluoromethyl)-phenyl}-methyl}-piperidine-1-
- 25 carboxamide;
- N-1,2-benzisoxazol-3-yl-3-{{(4-fluorophenyl)methyl}piperidine-1-carboxamide;
- N-isoxazol-3-yl-3-{{4-(trifluoromethyl)phenyl}methyl}piperidine-1-carboxamide;
- N-isoxazol-3-yl-3-{{3-(trifluoromethyl)phenyl}methyl}piperidine-1-carboxamide;
- 3-{{3-(4-fluorophenoxy)benzyl}-N-pyridin-3-yl}piperidine-1-carboxamide;
- 30 3-{{3-(4-fluoro-phenoxy)-benzyl}-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide;
- 3-{{3-(4-fluoro-phenoxy)-benzyl}-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide;
- (+)-3-{{3-(4-chloro-phenoxy)-benzyl}-piperidine-1-carboxylic acid pyridin-3-ylamide;
- 35 (−)-3-{{3-(4-chloro-phenoxy)-benzyl}-piperidine-1-carboxylic acid pyridin-3-ylamide;


- 5 3-[3-(4-trifluoromethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide;
3-quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3,4-dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
- 10 3-(4-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide;
- 15 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-
- 20 ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-7-ylamide;
- 25 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-c]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-
- 30 amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide;
3-[3-(4-trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
- 35 3-(4-chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
and pharmaceutically acceptable salts thereof.

5

16. A pharmaceutical composition according to claim 14, further comprising: an analgesic selected from the group consisting of opioids and non-steroidal anti-inflammatory drugs.

10 17. A pharmaceutical composition according to claim 14, further comprising: an additional active ingredient selected from the group consisting of aspirin, acetaminophen, opioids, ibuprofen, naproxen, COX-2 inhibitors, gabapentin, pregabalin, and tramadol.

15 18. A method of treating a subject suffering from or diagnosed with a disease, disorder, or medical condition mediated by FAAH activity, comprising administering to the subject in need of such treatment an effective amount of at least one chemical entity selected from compounds of Formula (I):



20 wherein:

Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

(i) phenyl unsubstituted or substituted with one or two R^a moieties;

30 where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃, -OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c, -SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;
or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -OCF₂O-;

- 5 where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b and R^c taken together with the atoms of attachment form a 4-8 membered ring;
- (ii) phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties, wherein:
- 10 L is a linker selected from the group consisting of -(CH₂)₁₋₆-, -CH=CH-, -O-, -C₁₋₆alkyl-O-C₁₋₆alkyl-, -C₁₋₆alkyl-N(C₁₋₆alkyl)-C₁₋₆alkyl-, -C₁₋₆alkyl-S(O)₀₋₂C₁₋₆alkyl-, -C≡C-, -C(O)-, or a covalent bond;
- Ar³ is:
- (a) phenyl unsubstituted or substituted with one or two R^a moieties;
- 15 (b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom
- 20 selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;
- and pharmaceutically acceptable salts of compounds of Formula (I),
- 25 pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I).

19. A method according to claim 18, wherein said at least one chemical entity is selected from the group consisting of:
- 30 3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
- N-pyrimidin-4-yl-3-[[4-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
- N-pyrimidin-4-yl-3-[[3-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
- 3-[(4-fluorophenyl)methyl]-N-pyrimidin-4-ylpiperidine-1-carboxamide;
- 35 N-1,2-benzisoxazol-3-yl-3-[[4-(trifluoromethyl)-phenyl]-methyl]-piperidine-1-carboxamide;

- 5 N-1,2-benzisoxazol-3-yl-3-[[3-(trifluoromethyl)-phenyl]-methyl]-piperidine-1-carboxamide;
N-1,2-benzisoxazol-3-yl-3-[(4-fluorophenyl)methyl]piperidine-1-carboxamide;
N-isoxazol-3-yl-3-[[4-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
N-isoxazol-3-yl-3-[[3-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;
- 10 3-[3-(4-fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide;
- 15 (+)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
(-)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-trifluoromethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide;
3-quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide;
- 20 3-(3,4-dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide;
- 25 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-
- 30 ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-5-ylamide;
- 35 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-7-ylamide;

- 5 3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-c]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-
10 amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide;
3-[3-(4-trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
15 3-(4-chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
and pharmaceutically acceptable salts thereof.

20. A method according to claim 18, wherein the disease, disorder, or medical condition is selected from the group consisting of: anxiety, depression, pain, sleep
20 disorders, eating disorders, inflammation, movement disorders, HIV wasting syndrome, closed head injury, stroke, learning and memory disorders, Alzheimer's disease, epilepsy, Tourette's syndrome, Niemann-Pick disease, Parkinson's disease, Huntington's chorea, optic neuritis, autoimmune uveitis, drug withdrawal, nausea, emesis, sexual dysfunction, post-traumatic stress disorder, cerebral vasospasm,
25 glaucoma, irritable bowel syndrome, inflammatory bowel disease, immunosuppression, gastroesophageal reflux disease, paralytic ileus, secretory diarrhea, gastric ulcer, rheumatoid arthritis, unwanted pregnancy, hypertension, cancer, hepatitis, allergic airway disease, autoimmune diabetes, intractable pruritis, and neuroinflammation.

30

21. A method according to claim 18, wherein the disease, disorder, or medical condition is pain or inflammation.

22. A method according to claim 18, wherein the disease, disorder, or medical
35 condition is anxiety, a sleep disorder, an eating disorder, or a movement disorder.

5 23. A method according to claim 18, wherein the disease, disorder, or medical condition is multiple sclerosis.

24. A method according to claim 18, wherein the disease, disorder, or medical condition is energy metabolism or bone homeostasis.


10

AMENDED CLAIMS

received by the International Bureau on 20 April 2010 (20.04.2010)

What is claimed is:

1. A chemical entity selected from compounds of Formula (I):



wherein:

Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

- (i) phenyl unsubstituted or substituted with one or two R^a moieties;

where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃, -OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c, -SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;

or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -OCF₂O-;

where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b and R^c taken together with the atoms of attachment form a 4-8 membered ring;

- (ii) phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties, wherein:

L is a linker selected from the group consisting of -(CH₂)₁₋₆-, -CH=CH-, -O-, -C₁₋₆alkyl-O-C₁₋₆alkyl-, -C₁₋₆alkyl-N(C₁₋₆alkyl)-C₁₋₆alkyl-, -C₁₋₆alkyl-S(O)₀₋₂C₁₋₆alkyl-, -C≡C-, -C(O)-, or a covalent bond;

Ar³ is:

- (a) phenyl unsubstituted or substituted with one or two R^a moieties;

- (b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or

- (c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;

where when Ar² is 3-trifluoromethyl-phenyl, Ar¹ is not thiazolo[5,4-d]pyrimidine; and pharmaceutically acceptable salts of compounds of Formula (I), pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I).

2. A chemical entity as in claim 1, wherein Ar¹ is isoxazolo[5,4-c]pyridin-3-yl, isoxazolo[4,5-c]pyridin-3-yl, isoxazolo[4,5-b]pyridin-3-yl, isoxazolo[5,4-b]pyridin-3-yl, imidazo[1,2-a]pyridin-8-yl, imidazo[1,2-a]pyridin-7-yl, imidazo[1,2-a]pyridin-6-yl, imidazo[1,2-a]pyridin-5-yl, imidazo[1,2-b]pyridazin-3-yl, imidazo[1,2-a]pyridin-3-yl, imidazo[1,2-a]pyridin-2-yl, imidazo[1,2-a]pyrimidin-7-yl, imidazo[1,2-a]pyrimidin-5-yl, imidazo[1,2-c]pyrimidin-7-yl, benzooxazol-6-yl, 1H-indazol-7-yl, quinolin-8-yl, isoquinolin-5-yl, isoquinolin-4-yl, 1,1-dioxo-1H-1λ⁶-benzo[d]isothiazol-3-yl, 1H-pyrrolo[2,3-b]pyridin-5-yl, 1H-pyrrolo[2,3-b]pyridin-4-yl, 1H-pyrrolo[3,2-b]pyridin-6-yl, 2-methyl-benzothiazol-6-yl, 1-isopropyl-1H-pyrazolo[3,4-b]pyridin-5-yl, 2-methyl-benzooxazol-5-yl, 1H-indazol-7-yl, 1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl, 2-methyl-2H-indazol-4-yl, benzooxazole-2-yl group, 6-[1,2,3]triazol-1-yl-pyridin-3-yl, 6-[1,2,4]triazol-4-yl-pyridin-3-yl, 6-[1,2,4]triazol-1-yl-pyridin-3-yl, 6-[1,2,3]triazol-2-yl-pyridin-3-yl, 4-[1,2,3]triazol-2-yl-phenyl, 4-[1,2,3]triazol-1-yl-phenyl, 5-methyl-[1,3,4]oxadiazol-2-yl, 2-phenyl-pyrimidin-5-yl, 3-pyridyl, benzisoxazol-3-yl, pyrimidin-4-yl, isoxazol-3-yl, 6-fluorobenzo[d]isoxazol-3-yl, 3-phenyl-[1,2,4]thiadiazol-5-yl, 1H-tetrazol-5-yl, benzo[1,2,5]thiadiazol-4-yl, benzo[1,2,5]oxadiazol-4-yl, thiophen-2-yl, thiophen-3-yl, 6-chloro-pyridazin-3-yl, pyrazin-2-yl, 1H-benzotriazol-5-yl, [1,5]naphthyridin-2-yl, benzothiazol-6-yl, or 1H-pyrazol-3-yl.

3. A chemical entity as in claim 1, wherein Ar¹ is pyrimidine, isoxazole, benzisoxazole, or pyridine, each group optionally substituted with one or two R^a.
4. A chemical entity as in claim 1, wherein Ar² is phenyl substituted with one or two R^a moieties.
5. A chemical entity as in claim 4, wherein each R^a moiety is independently selected from the group consisting of: chloro, cyano, isobutyl, methylsulfanyl, methanesulfonyl, trifluoromethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, fluoro, methyl, methoxy, tert-butyl, bromo, methoxycarbonyl, cyanomethyl, methoxycarbonylmethyl, trifluoromethanesulfonyl, trifluoromethanesulfanyl, and butyl; or two adjacent R^a moieties taken together form -OCH₂O- or -OCF₂O-.
6. A chemical entity as in claim 4, wherein each R^a moiety is independently selected from the group consisting of: F, Cl, Br, CF₃, or two adjacent R^a moieties taken together form -OCF₂O-.
7. A chemical entity as in claim 1, wherein Ar² is phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties.
8. A chemical entity as in claim 7, wherein L is -CH₂CH₂-, -O-, -OCH₂-, or -C≡C-.
9. A chemical entity as in claim 7, wherein L is -O-.
10. A chemical entity as in claim 7, wherein Ar³ is phenyl, unsubstituted or substituted with one or two R^a moieties.
11. A chemical entity as in claim 10, wherein each R^a moiety is independently selected from the group consisting of: chloro, cyano, isobutyl, methylsulfanyl, methanesulfonyl, trifluoromethyl, trifluoromethoxy, 2,2,2-trifluoroethoxy, fluoro, methyl, methoxy, tert-butyl, bromo, methoxycarbonyl, cyanomethyl, methoxycarbonylmethyl,

trifluoromethanesulfonyl, trifluoromethanesulfanyl, and butyl; or two adjacent R^a moieties taken together form –OCH₂O– or –OCF₂O–.

12. A chemical entity as in claim 10, wherein each R^a moiety is independently fluoro.

13. A chemical entity selected from the group consisting of:

3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;

N-pyrimidin-4-yl-3-[[4-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;

N-pyrimidin-4-yl-3-[[3-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;

3-[(4-fluorophenyl)methyl]-N-pyrimidin-4-ylpiperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-[[4-(trifluoromethyl)-phenyl]-methyl]-piperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-[[3-(trifluoromethyl)-phenyl]-methyl]-piperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-[(4-fluorophenyl)methyl]piperidine-1-carboxamide;

N-isoxazol-3-yl-3-[[4-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;

N-isoxazol-3-yl-3-[[3-(trifluoromethyl)phenyl]methyl]piperidine-1-carboxamide;

3-[3-(4-fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide;

3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide;

3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide;

(+)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;

(–)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;

3-[3-(4-trifluoromethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide;

3-quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide;

3-(3,4-dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;


3-(3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;

3-(4-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;

3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-c]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide;
3-[3-(4-trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
and pharmaceutically acceptable salts thereof.

14. A pharmaceutical composition for treating a disease, disorder, or medical condition mediated by FAAH activity, comprising:

(a) an effective amount of at least one chemical entity selected from compounds of Formula (I):



wherein:

Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

(i) phenyl unsubstituted or substituted with one or two R^a moieties;

where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃, -OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c, -SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;

or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -OCF₂O-; where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b and R^c taken together with the atoms of attachment form a 4-8 membered ring;

(ii) phenyl substituted at the 3- or 4-position with -L-Ar³, unsubstituted or substituted with one or two R^a moieties, wherein:

L is a linker selected from the group consisting of -(CH₂)₁₋₆-, -CH=CH-, -O-, -C₁₋₆alkyl-O-C₁₋₆alkyl-, -C₁₋₆alkyl-N(C₁₋₆alkyl)-C₁₋₆alkyl-, -C₁₋₆alkyl-S(O)₀₋₂C₁₋₆alkyl-, -C≡C-, -C(O)-, or a covalent bond;

Ar³ is:

- (a) phenyl unsubstituted or substituted with one or two R^a moieties;
- (b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom selected from the group consisting of N, O, and S, and optionally having up to four

additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;

and pharmaceutically acceptable salts of compounds of Formula (I), pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I) ; and

(b) a pharmaceutically acceptable excipient.

15. A pharmaceutical composition according to claim 14, wherein said at least one chemical entity is selected from the group consisting of:

3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;

N-pyrimidin-4-yl-3-{[4-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;

N-pyrimidin-4-yl-3-{[3-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;

3-[(4-fluorophenyl)methyl]-N-pyrimidin-4-ylpiperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-{[4-(trifluoromethyl)-phenyl]-methyl}-piperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-{[3-(trifluoromethyl)-phenyl]-methyl}-piperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-[(4-fluorophenyl)methyl]piperidine-1-carboxamide;

N-isoxazol-3-yl-3-{[4-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;

N-isoxazol-3-yl-3-{[3-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;

3-[3-(4-fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide;

3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide;

3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide;

(+)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;

(-)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;


3-[3-(4-trifluormethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide;

3-quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3,4-dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-c]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide;
3-[3-(4-trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
and pharmaceutically acceptable salts thereof.

16. A pharmaceutical composition according to claim 14, further comprising: an analgesic selected from the group consisting of opioids and non-steroidal anti-inflammatory drugs.

17. A pharmaceutical composition according to claim 14, further comprising: an additional active ingredient selected from the group consisting of aspirin, acetaminophen, opioids, ibuprofen, naproxen, COX-2 inhibitors, gabapentin, pregabalin, and tramadol.

18. A method of treating a subject suffering from or diagnosed with a disease, disorder, or medical condition mediated by FAAH activity, comprising administering to the subject in need of such treatment an effective amount of at least one chemical entity selected from compounds of Formula (I):



wherein:

Ar¹ is mono- or fused bicyclic heteroaryl group consisting of 5-10 ring atoms and having one heteroatom selected from the group consisting of N, O and S, with a carbon atom as point of ring attachment, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said heteroaryl group having not more than five heteroatoms, each heteroaryl group independently unsubstituted or substituted with halo, -C₁₋₄alkyl, -OC₁₋₄alkyl, -OCF₃, -CN, or -CF₃;

Ar² is

(i) phenyl unsubstituted or substituted with one or two R^a moieties;

where each R^a moiety is independently -C₁₋₈alkyl, -OC₁₋₈alkyl, halo, -CF₃, -OCF₃, -OCH₂CF₃, -S(O)₀₋₂C₁₋₈alkyl, -S(O)₀₋₂CF₃, -CO₂H, -N(R^b)R^c, -SO₂NR^bR^c, -NR^bSO₂R^c, -C(O)NR^bR^c, or -NO₂;

or two adjacent R^a moieties taken together form -O(CH₂)₁₋₂O- or -OCF₂O-;

where R^b and R^c are each independently -H or -C₁₋₈alkyl, or optionally R^b and R^c taken together with the atoms of attachment form a 4-8 membered ring;

(ii) phenyl substituted at the 3- or 4-position with $-L-Ar^3$, unsubstituted or substituted with one or two R^a moieties, wherein:

L is a linker selected from the group consisting of $-(CH_2)_{1-6}-$, $-CH=CH-$, $-O-$, $-C_{1-6}alkyl-O-C_{1-6}alkyl-$, $-C_{1-6}alkyl-N(C_{1-6}alkyl)-C_{1-6}alkyl-$, $-C_{1-6}alkyl-S(O)_{0-2}C_{1-6}alkyl-$, $-C\equiv C-$, $-C(O)-$, or a covalent bond;

Ar^3 is:

- (a) phenyl unsubstituted or substituted with one or two R^a moieties;
 - (b) a monocyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
 - (c) a 9- or 10-membered fused bicyclic heteroaryl group unsubstituted or substituted with one or two R^a moieties; or
- (iii) a 9- or 10-membered fused bicyclic heteroaryl having one heteroatom selected from the group consisting of N, O, and S, and optionally having up to four additional carbon ring atoms replaced with nitrogen, said fused bicyclic heteroaryl having not more than five heteroatoms, and unsubstituted or substituted with one, two or three R^a moieties;

and pharmaceutically acceptable salts of compounds of Formula (I), pharmaceutically acceptable prodrugs of compounds of Formula (I), and pharmaceutically active metabolites of compounds of Formula (I).

19. A method according to claim 18, wherein said at least one chemical entity is selected from the group consisting of:

3-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;

N-pyrimidin-4-yl-3-{[4-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;

N-pyrimidin-4-yl-3-{[3-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;

3-[(4-fluorophenyl)methyl]-N-pyrimidin-4-ylpiperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-{[4-(trifluoromethyl)-phenyl]-methyl}-piperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-{[3-(trifluoromethyl)-phenyl]-methyl}-piperidine-1-carboxamide;

N-1,2-benzisoxazol-3-yl-3-[(4-fluorophenyl)methyl]piperidine-1-carboxamide;
N-isoxazol-3-yl-3-{[4-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;
N-isoxazol-3-yl-3-{[3-(trifluoromethyl)phenyl]methyl}piperidine-1-carboxamide;
3-[3-(4-fluorophenoxy)benzyl]-N-pyridin-3-ylpiperidine-1-carboxamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-chloro-pyridin-3-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-b]pyridazin-3-ylamide;
(+)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
(-)-3-[3-(4-chloro-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-trifluoromethoxy-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridine-3-ylamide;
3-quinolin-3-ylmethyl-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3,4-dichloro-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (1H-pyrrolo[2,3-b]pyridin-4-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[4,5-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoxazolo[5,4-c]pyridin-3-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyridin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-7-ylamide;

3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-a]pyrimidin-5-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid imidazo[1,2-c]pyrimidin-7-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid (4-cyano-pyridin-3-yl)-amide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid quinolin-8-ylamide;
3-[3-(4-fluoro-phenoxy)-benzyl]-piperidine-1-carboxylic acid isoquinolin-5-ylamide;
3-[3-(4-trifluoromethyl-phenoxy)-benzyl]-piperidine-1-carboxylic acid pyridin-3-ylamide;
3-(4-chloro-3-trifluoromethoxy-benzyl)-piperidine-1-carboxylic acid pyridin-3-ylamide;
and pharmaceutically acceptable salts thereof.

20. A method according to claim 18, wherein the disease, disorder, or medical condition is selected from the group consisting of: anxiety, depression, pain, sleep disorders, eating disorders, inflammation, movement disorders, HIV wasting syndrome, closed head injury, stroke, learning and memory disorders, Alzheimer's disease, epilepsy, Tourette's syndrome, Niemann-Pick disease, Parkinson's disease, Huntington's chorea, optic neuritis, autoimmune uveitis, drug withdrawal, nausea, emesis, sexual dysfunction, post-traumatic stress disorder, cerebral vasospasm, glaucoma, irritable bowel syndrome, inflammatory bowel disease, immunosuppression, gastroesophageal reflux disease, paralytic ileus, secretory diarrhea, gastric ulcer, rheumatoid arthritis, unwanted pregnancy, hypertension, cancer, hepatitis, allergic airway disease, autoimmune diabetes, intractable pruritis, and neuroinflammation.

21. A method according to claim 18, wherein the disease, disorder, or medical condition is pain or inflammation.

22. A method according to claim 18, wherein the disease, disorder, or medical condition is anxiety, a sleep disorder, an eating disorder, or a movement disorder.

23. A method according to claim 18, wherein the disease, disorder, or medical condition is multiple sclerosis.

24. A method according to claim 18, wherein the disease, disorder, or medical condition is energy metabolism or bone homeostasis.

Statement Under Article 19(1):

In replacement claim 1: thiazolo [5,4-d] pyrimidine has been deleted from original claim 1 as a choice for the bicyclic heteroaryl group recited in the definition of the formula variable Ar¹ when Ar² is 3-trifluoromethyl-phenyl.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/065752

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D401/12 C07D471/04 A61K31/47 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/134958 A (HOFFMANN LA ROCHE [CH]; BRINKMAN JOHN A [US]; CHEUNG ADRIAN WAI-HING [] 29 November 2007 (2007-11-29) example 68	1,4-6, 14,16, 18,20-24
A	WO 2006/014136 A1 (ASTRAZENECA AB [SE]; FOLMER JAMES [US]; HUNT SIMON FRASER [GB]; HAMLEY) 9 February 2006 (2006-02-09) the whole document	1-24



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

11 February 2010

Date of mailing of the international search report

22/02/2010

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Hacking, Michiel

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2009/065752

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007134958 A	29-11-2007	AR 060957 A1	23-07-2008
		AU 2007253485 A1	29-11-2007
		CA 2651769 A1	29-11-2007
		CN 101448844 A	03-06-2009
		EP 2027132 A1	25-02-2009
		JP 2009537473 T	29-10-2009
		KR 20080111141 A	22-12-2008
		US 2007270433 A1	22-11-2007
WO 2006014136 A1	09-02-2006	AU 2005267932 A1	09-02-2006
		BR PI0514035 A	27-05-2008
		CA 2576112 A1	09-02-2006
		CN 1993325 A	04-07-2007
		EP 1781613 A1	09-05-2007
		JP 2008508353 T	21-03-2008
		KR 20070043998 A	26-04-2007
		US 2008064706 A1	13-03-2008
		ZA 200700683 A	27-08-2008