



(43) International Publication Date
24 September 2015 (24.09.2015)

WIPO | PCT

(10) International Publication Number
WO 2015/140071 A1

(51) International Patent Classification:

A01N 43/80 (2006.01) *A01N 57/12* (2006.01)
A01N 37/34 (2006.01) *A01N 37/36* (2006.01)
A01N 47/34 (2006.01) *A01P 3/00* (2006.01)

(21) International Application Number:

PCT/EP2015/055317

(22) International Filing Date:

13 March 2015 (13.03.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

14160404.1 18 March 2014 (18.03.2014) EP

(71) Applicant: **BAYER CROPSCIENCE AG** [DE/DE]; Alfred-Nobel-Str. 50, 40789 Monheim (DE).

(72) Inventors: **HILLEBRAND, Stefan**; Lothringer Str. 22, 41462 Neuss (DE). **OLENIK, Britta**; Birkenstr. 30, 46242 Bottrop (DE). **WEISS, Martin**; Galäerweg 22, 53332 Bornheim/Sechtem (DE). **WACHENDORFF-NEUMANN, Ulrike**; Oberer Markweg 85, 56566 Neuwied (DE).

(74) Agent: **BIP PATENTS**; c/o Bayer Intellectual Property GmbH, Alfred-Nobel-Str. 10, 40789 Monheim am Rhein (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

(54) Title: ACTIVE COMPOUND COMBINATIONS

(57) Abstract: The present invention relates to active compound combinations, in particular within a fungicide composition, which comprises (A) crystalline thiazolylisoxazoline of formula (I) in its Form B and a further fungicidally active compound (B). Moreover, the invention relates to a method for curatively or preventively controlling the phytopathogenic fungi of plants or crops, to the use of a combination according to the invention for the treatment of seed, to a method for protecting a seed and not at least to the treated seed.



WO 2015/140071 A1

Active compound combinations

The present invention relates to active compound combinations, in particular within a fungicide composition, which comprises (A) crystalline thiazolyloxazoline of formula (I) in its Form B and a further fungicidally active compound (B). Moreover, the invention relates to a method for curatively or preventively
 5 controlling the phytopathogenic fungi of plants or crops, to the use of a combination according to the invention for the treatment of seed, to a method for protecting a seed and not at least to the treated seed.

It is already known that compound of formula (I) can be used as fungicide (WO 2008/013925).


Crystalline thiazolyloxazoline of formula (I) in its Form B is known from WO 2010/123791 (see page 4, line 5 to 10) and also some mixtures of compound of formula (I) and (B) are known (WO 2009/055514).

10 Since the ecological and economic demands made on modern active ingredients, for example fungicides, are increasing constantly, for example with respect to activity spectrum, toxicity, selectivity, application rate, formation of residues and favourable manufacture, and there can also be problems, for example, with resistances, there is a constant need to develop novel fungicidal compositions which have advantages over the known compositions at least in some areas.

15 The present invention provides synergistic active compound combinations/compositions which in some aspects at least achieve the stated objective.

The present invention provides a synergistic combination comprising:

(A) Crystalline thiazolyloxazoline of formula (I)



(I)

20 in its Form B (see WO 2010/123791, page 4, line 5 to 10) characterized by powder X-ray diffraction pattern having at least the 2Θ reflection positions

2Θ
14.902
18.123
18.87

20.204
20.883
21.79
24.186
26.947

and

(B) at least one further active compound selected from the following groups

- (1) inhibitors of the ergosterol synthesis,
- 5 (2) inhibitors of the respiratory chain at complex I or II,
- (3) inhibitors of the respiratory chain at complex III,
- (4) inhibitors of the mitosis and cell division,
- (5) compounds capable of having a multisite action,
- (6) compounds capable of inducing a host defense,
- 10 (7) inhibitors of the amino acid and/or protein biosynthesis,
- (8) inhibitors of the ATP production,
- (9) inhibitors of the cell wall synthesis,
- (10) inhibitors of the lipid and membrane synthesis,
- (11) inhibitors of the melanine biosynthesis,
- 15 (12) inhibitors of the nucleic acid synthesis,
- (13) inhibitors of the signal transduction,
- (14) compounds capable of acting as uncoupler,
- (15) other fungicides.

In the description below numbers in parenthesis behind a compound name represent the CAS Registry No. of said compound.

The combinations according to the invention not only bring about the additive enhancement of the spectrum of action with respect to the phytopathogen to be controlled that was in principle to be expected but achieves
5 a synergistic effect which is higher than for the mixture with amorphous form of compound of formula (I).

In addition to the fungicidal synergistic activity, the active compound combinations according to the invention have further surprising properties which, in a wider sense, may also be called synergistic, such as, for example: broadening of the activity spectrum to other phytopathogens, for example to resistant strains of plant diseases; lower application rates of the active compounds; sufficient control of pests with the aid of the
10 active compound combinations according to the invention even at application rates where the individual compounds show no or virtually no activity; advantageous behaviour during formulation or during use, for example during grinding, sieving, emulsifying, dissolving or dispensing; improved storage stability and light stability; advantageous residue formation; improved toxicological or ecotoxicological behaviour; improved properties of the plant, for example better growth, increased harvest yields, a better developed root system, a
15 larger leaf area, greener leaves, stronger shoots, less seed required, lower phytotoxicity, mobilization of the defence system of the plant, good compatibility with plants. Thus, the use of the active compound combinations or compositions according to the invention contributes considerably to keeping young cereal stands healthy, which increases, for example, the winter survival of the cereal seed treated, and also safeguards quality and yield. Moreover, the active compound combinations according to the invention may contribute to enhanced systemic action. Even if the individual compounds of the combination have no sufficient systemic
20 properties, the active compound combinations according to the invention may still have this property. In a similar manner, the active compound combinations according to the invention may result in higher persistency of the fungicidal action.

Preference is given to combinations comprising at least one further active compound (B) selected from the
25 following groups:

(1) Inhibitors of the ergosterol biosynthesis, for example (1.1) aldimorph (1704-28-5), (1.2) azaconazole (60207-31-0), (1.3) bitertanol (55179-31-2), (1.4) bromuconazole (116255-48-2), (1.5) cyproconazole (113096-99-4), (1.6) diclobutrazole (75736-33-3), (1.7) difenoconazole (119446-68-3), (1.8) diniconazole (83657-24-3), (1.9) diniconazole-M (83657-18-5), (1.10) dodemorph (1593-77-7), (1.11) dodemorph acetate (31717-87-0), (1.12) epoxiconazole (106325-08-0), (1.13) etaconazole (60207-93-4), (1.14) fenarimol (60168-88-9), (1.15) fenbuconazole (114369-43-6), (1.16) fenhexamid (126833-17-8), (1.17) fenpropidin (67306-00-7), (1.18) fenpropimorph (67306-03-0), (1.19) fluquinconazole (136426-54-5), (1.20) flurprimidol (56425-91-3), (1.21) flusilazole (85509-19-9), (1.22) flutriafol (76674-21-0), (1.23) furconazole (112839-33-5), (1.24) furconazole-cis (112839-32-4), (1.25) hexaconazole (79983-71-4), (1.26) imazalil (60534-80-7), (1.27) imazalil sulfate (58594-72-2), (1.28) imibenconazole (86598-92-7), (1.29) ipconazole (125225-28-7), (1.30) metconazole (125116-23-6), (1.31) myclobutanil (88671-89-0), (1.32) naftifine (65472-88-0), (1.33) nuarimol (63284-71-9), (1.34) oxpoconazole (174212-12-5), (1.35) paclobutrazol

(76738-62-0), (1.36) pefurazoate (101903-30-4), (1.37) penconazole (66246-88-6), (1.38) piperalin (3478-94-2), (1.39) prochloraz (67747-09-5), (1.40) propiconazole (60207-90-1), (1.41) prothioconazole (178928-70-6), (1.42) pyributicarb (88678-67-5), (1.43) pyrifenoxy (88283-41-4), (1.44) quinconazole (103970-75-8), (1.45) simeconazole (149508-90-7), (1.46) spiroxamine (118134-30-8), (1.47) tebuconazole (107534-96-3), (1.48) terbinafine (91161-71-6), (1.49) tetraconazole (112281-77-3), (1.50) triadimefon (43121-43-3), (1.51) triadimenol (89482-17-7), (1.52) tridemorph (81412-43-3), (1.53) triflumizole (68694-11-1), (1.54) triforine (26644-46-2), (1.55) triticonazole (131983-72-7), (1.56) uniconazole (83657-22-1), (1.57) uniconazole-p (83657-17-4), (1.58) viniconazole (77174-66-4), (1.59) voriconazole (137234-62-9), (1.60) 1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol (129586-32-9), (1.61) methyl 1-(2,2-dimethyl-2,3-dihydro-1H-inden-1-yl)-1H-imidazole-5-carboxylate (110323-95-0), (1.62) N'-{5-(difluoromethyl)-2-methyl-4-[3-(trimethylsilyl)propoxy]phenyl}-N-ethyl-N-methylimidoforamide, (1.63) N-ethyl-N-methyl-N'-{2-methyl-5-(trifluoromethyl)-4-[3-(trimethylsilyl)propoxy]phenyl}imidoforamide, (1.64) O-[1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl] 1H-imidazole-1-carbothioate (111226-71-2).

15

(2) inhibitors of the respiratory chain at complex I or II, for example (2.1) bixafen (581809-46-3), (2.2) boscalid (188425-85-6), (2.3) carboxin (5234-68-4), (2.4) diflometorim (130339-07-0), (2.5) fenfuram (24691-80-3), (2.6) fluopyram (658066-35-4), (2.7) flutolanil (66332-96-5), (2.8) fluxapyroxad (907204-31-3), (2.9) furametpyr (123572-88-3), (2.10) furmecyclox (60568-05-0), (2.11) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR) (881685-58-1), (2.12) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.13) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.14) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.15) isopyrazam (syn epimeric racemate 1RS,4SR,9RS), (2.16) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.17) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (2.18) mepronil (55814-41-0), (2.19) oxycarboxin (5259-88-1), (2.20) penflufen (494793-67-8), (2.21) penthiopyrad (183675-82-3), (2.22) sedaxane (874967-67-6), (2.23) thifluzamide (130000-40-7), (2.24) 1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (2.25) 3-(difluoromethyl)-1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-1H-pyrazole-4-carboxamide, (2.26) 3-(difluoromethyl)-N-[4-fluoro-2-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-1-methyl-1H-pyrazole-4-carboxamide, (2.27) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide (1092400-95-7), (2.28) 5,8-difluoro-N-[2-(2-fluoro-4-{[4-(trifluoromethyl)pyridin-2-yl]oxy}phenyl)ethyl]quinazolin-4-amine (1210070-84-0), (2.29) benzovindiflupyr, (2.30) N-[(1S,4R)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.31) N-[(1R,4S)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.32) 3-(Difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (2.33) 1,3,5-Trimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (2.34) 1-Methyl-3-(trifluoromethyl)-N-(1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazol-4-carboxamid, (2.35) 1-Methyl-3-(trifluoromethyl)-N-[(1S)-1,3,3-trimethyl-

2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (2.36) 1-Methyl-3-(trifluormethyl)-N-[(1R)-1,3,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (2.37) 3-(Difluormethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (2.38) 3-(Difluormethyl)-1-methyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (2.39) 1,3,5-Trimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid, (2.40) 1,3,5-Trimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazol-4-carboxamid.

(3) inhibitors of the respiratory chain at complex III, for example (3.1) ametocetradin (865318-97-4), (3.2) amisulbrom (348635-87-0), (3.3) azoxystrobin (131860-33-8), (3.4) cyazofamid (120116-88-3), (3.5) coumethoxystrobin (850881-30-0), (3.6) coumoxystrobin (850881-70-8), (3.7) dimoxystrobin (141600-52-4), (3.8) enestroburin (238410-11-2), (3.9) famoxadone (131807-57-3), (3.10) fenamidone (161326-34-7), (3.11) fenoxystrobin (918162-02-4), (3.12) fluoxastrobin (361377-29-9), (3.13) kresoxim-methyl (143390-89-0), (3.14) metominostrobin (133408-50-1), (3.15) orysastrobin (189892-69-1), (3.16) picoxystrobin (117428-22-5), (3.17) pyraclostrobin (175013-18-0), (3.18) pyrametostrobin (915410-70-7), (3.19) pyraoxystrobin (862588-11-2), (3.20) pyribencarb (799247-52-2), (3.21) triclopyricarb (902760-40-1), (3.22) trifloxystrobin (141517-21-7), (3.23) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl}oxy}phenyl)-2-(methoxyimino)-N-methylethanamide, (3.24) (2E)-2-(methoxyimino)-N-methyl-2-(2-{{((1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)ethanamide, (3.25) (2E)-2-(methoxyimino)-N-methyl-2-{2-[(E)-({1-[3-(trifluoromethyl)phenyl]ethoxy}imino)methyl]phenyl}ethanamide (158169-73-4), (3.26) (2E)-2-{2-[[({(1E)-1-(3-{{(E)-1-fluoro-2-phenylethenyl}oxy}phenyl)ethylidene}amino}oxy)methyl]phenyl}-2-(methoxyimino)-N-methylethanamide (326896-28-0), (3.27) (2E)-2-{2-[[{(2E,3E)-4-(2,6-dichlorophenyl)but-3-en-2-ylidene}amino}oxy)methyl]phenyl}-2-(methoxyimino)-N-methylethanamide, (3.28) 2-chloro-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)pyridine-3-carboxamide (119899-14-8), (3.29) 5-methoxy-2-methyl-4-(2-{{((1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, (3.30) methyl (2E)-2-{2-[(cyclopropyl[(4-methoxyphenyl)imino]methyl}sulfanyl)methyl]phenyl}-3-methoxyprop-2-enoate (149601-03-6), (3.31) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-(formylamino)-2-hydroxybenzamide (226551-21-9), (3.32) 2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide (173662-97-0), (3.33) (2R)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide (394657-24-0).

(4) Inhibitors of the mitosis and cell division, for example (4.1) benomyl (17804-35-2), (4.2) carbendazim (10605-21-7), (4.3) chlorfenazole (3574-96-7), (4.4) diethofencarb (87130-20-9), (4.5) ethaboxam (162650-77-3), (4.6) fluopicolide (239110-15-7), (4.7) fuberidazole (3878-19-1), (4.8) pencycuron (66063-05-6), (4.9) thiabendazole (148-79-8), (4.10) thiophanate-methyl (23564-05-8), (4.11) thiophanate (23564-06-9), (4.12) zoxamide (156052-68-5), (4.13) 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine (214706-53-3), (4.14) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine (1002756-87-7).

- (5) Compounds capable to have a multisite action, like for example (5.1) bordeaux mixture (8011-63-0), (5.2) captafol (2425-06-1), (5.3) captan (133-06-2), (5.4) chlorothalonil (1897-45-6), (5.5) copper hydroxide (20427-59-2), (5.6) copper naphthenate (1338-02-9), (5.7) copper oxide (1317-39-1), (5.8) copper oxychloride (1332-40-7), (5.9) copper(2+) sulfate (7758-98-7), (5.10) dichlofluanid (1085-98-9), (5.11) dithianon (3347-22-6), (5.12) dodine (2439-10-3), (5.13) dodine free base, (5.14) ferbam (14484-64-1), (5.15) fluorofolpet (719-96-0), (5.16) folpet (133-07-3), (5.17) guazatine (108173-90-6), (5.18) guazatine acetate, (5.19) iminoctadine (13516-27-3), (5.20) iminoctadine albesilate (169202-06-6), (5.21) iminoctadine triacetate (57520-17-9), (5.22) mancooper (53988-93-5), (5.23) mancozeb (8018-01-7), (5.24) maneb (12427-38-2), (5.25) metiram (9006-42-2), (5.26) metiram zinc (9006-42-2), (5.27) oxine-copper (10380-28-6), (5.28) propamidine (104-32-5), (5.29) propineb (12071-83-9), (5.30) sulphur and sulphur preparations including calcium polysulphide (7704-34-9), (5.31) thiram (137-26-8), (5.32) tolylfluanid (731-27-1), (5.33) zineb (12122-67-7), (5.34) ziram (137-30-4).
- (6) Compounds capable to induce a host defence, like for example (6.1) acibenzolar-S-methyl (135158-54-2), (6.2) isotianil (224049-04-1), (6.3) probenazole (27605-76-1), (6.4) tiadinil (223580-51-6).
- (7) Inhibitors of the amino acid and/or protein biosynthesis, for example (7.1) andoprim (23951-85-1), (7.2) blasticidin-S (2079-00-7), (7.3) cyprodinil (121552-61-2), (7.4) kasugamycin (6980-18-3), (7.5) kasugamycin hydrochloride hydrate (19408-46-9), (7.6) mepanipyrin (110235-47-7), (7.7) pyrimethanil (53112-28-0), (7.8) 3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinoline (861647-32-7).
- (8) Inhibitors of the ATP production, for example (8.1) fentin acetate (900-95-8), (8.2) fentin chloride (639-58-7), (8.3) fentin hydroxide (76-87-9), (8.4) silthiofam (175217-20-6).
- (9) Inhibitors of the cell wall synthesis, for example (9.1) benthiavalicarb (177406-68-7), (9.2) dimethomorph (110488-70-5), (9.3) flumorph (211867-47-9), (9.4) iprovalicarb (140923-17-7), (9.5) mandipropamid (374726-62-2), (9.6) polyoxins (11113-80-7), (9.7) polyoxorim (22976-86-9), (9.8) validamycin A (37248-47-8), (9.9) valifenalate (283159-94-4; 283159-90-0).
- (10) Inhibitors of the lipid and membrane synthesis, for example (10.1) biphenyl (92-52-4), (10.2) chloroneb (2675-77-6), (10.3) dicloran (99-30-9), (10.4) edifenphos (17109-49-8), (10.5) etridiazole (2593-15-9), (10.6) iodocarb (55406-53-6), (10.7) iprobenfos (26087-47-8), (10.8) isoprothiolane (50512-35-1), (10.9) propamocarb (25606-41-1), (10.10) propamocarb hydrochloride (25606-41-1), (10.11) prothiocarb (19622-08-3), (10.12) pyrazophos (13457-18-6), (10.13) quintozone (82-68-8), (10.14) tecnazene (117-18-0), (10.15) tolclofos-methyl (57018-04-9).
- (11) Inhibitors of the melanine biosynthesis, for example (11.1) carpropamid (104030-54-8), (11.2) diclocymet (139920-32-4), (11.3) fenoxanil (115852-48-7), (11.4) phthalide (27355-22-2), (11.5) pyroquilon (57369-32-1), (11.6) tricyclazole (41814-78-2), (11.7) 2,2,2-trifluoroethyl {3-methyl-1-[(4-methylbenzoyl)amino]butan-2-yl} carbamate (851524-22-6).

- (12) Inhibitors of the nucleic acid synthesis, for example (12.1) benalaxyl (71626-11-4), (12.2) benalaxyl-M (kiralaxyl) (98243-83-5), (12.3) bupirimate (41483-43-6), (12.4) clozylacon (67932-85-8), (12.5) dime-thirimol (5221-53-4), (12.6) ethirimol (23947-60-6), (12.7) furalaxyl (57646-30-7), (12.8) hymexazol (10004-44-1), (12.9) metalaxyl (57837-19-1), (12.10) metalaxyl-M (mefenoxam) (70630-17-0), (12.11) ofurace (58810-48-3), (12.12) oxadixyl (77732-09-3), (12.13) oxolinic acid (14698-29-4).
- (13) Inhibitors of the signal transduction, for example (13.1) chlozolate (84332-86-5), (13.2) fenpiclonil (74738-17-3), (13.3) fludioxonil (131341-86-1), (13.4) iprodione (36734-19-7), (13.5) procymidone (32809-16-8), (13.6) quinoxifen (124495-18-7), (13.7) vinclozolin (50471-44-8).
- (14) Compounds capable to act as an uncoupler, like for example (14.1) binapacryl (485-31-4), (14.2) dinocap (131-72-6), (14.3) ferimzone (89269-64-7), (14.4) fluazinam (79622-59-6), (14.5) meptyldinocap (131-72-6).
- (15) Further compounds, like for example (15.1) benthiazole (21564-17-0), (15.2) bethoxazin (163269-30-5), (15.3) capsimycin (70694-08-5), (15.4) carvone (99-49-0), (15.5) chinomethionat (2439-01-2), (15.6) pyriofenone (chlazafenone) (688046-61-9), (15.7) cufraneb (11096-18-7), (15.8) cyflufenamid (180409-60-3), (15.9) cymoxanil (57966-95-7), (15.10) cyprosulfamide (221667-31-8), (15.11) dazomet (533-74-4), (15.12) debacarb (62732-91-6), (15.13) dichlorophen (97-23-4), (15.14) diclomezine (62865-36-5), (15.15) difenzoquat (49866-87-7), (15.16) difenzoquat methylsulphate (43222-48-6), (15.17) diphenylamine (122-39-4), (15.18) ecomate, (15.19) fenpyrazamine (473798-59-3), (15.20) flumetover (154025-04-4), (15.21) fluoroimide (41205-21-4), (15.22) flusulfamide (106917-52-6), (15.23) flutianil (304900-25-2), (15.24) fosetyl-aluminium (39148-24-8), (15.25) fosetyl-calcium, (15.26) fosetyl-sodium (39148-16-8), (15.27) hexachlorobenzene (118-74-1), (15.28) irumamycin (81604-73-1), (15.29) methasulfocarb (66952-49-6), (15.30) methyl isothiocyanate (556-61-6), (15.31) metrafenone (220899-03-6), (15.32) mildiomyacin (67527-71-3), (15.33) natamycin (7681-93-8), (15.34) nickel dimethyldithiocarbamate (15521-65-0), (15.35) nitrothal-isopropyl (10552-74-6), (15.36) octhilinone (26530-20-1), (15.37) oxamocarb (917242-12-7), (15.38) oxyfenthiin (34407-87-9), (15.39) pentachlorophenol and salts (87-86-5), (15.40) phenothrin, (15.41) phosphorous acid and its salts (13598-36-2), (15.42) propamocarb-fosetilate, (15.43) propanosine-sodium (88498-02-6), (15.44) proquinazid (189278-12-4), (15.45) pyrimorph (868390-90-3), (15.45e) (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one (1231776-28-5), (15.45z) (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one (1231776-29-6), (15.46) pyrrolnitrine (1018-71-9), (15.47) tebufloquin (376645-78-2), (15.48) tecloftalam (76280-91-6), (15.49) tolifenide (304911-98-6), (15.50) triazoxide (72459-58-6), (15.51) trichlamide (70193-21-4), (15.52) zarilamid (84527-51-5), (15.53) (3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl)carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate (517875-34-2), (15.54) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003319-79-6), (15.55) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003319-80-9), (15.56) 1-

(4-{4-[5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003318-67-9), (15.57) 1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl 1H-imidazole-1-carboxylate (111227-17-9), (15.58) 2,3,5,6-tetrachloro-4-(methylsulfonyl)pyridine (13108-52-6), (15.59) 2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)-one (221451-58-7), (15.60) 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, (15.61) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5R)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone (1003316-53-7), (15.62) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5S)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone (1003316-54-8), (15.63) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-{4-[4-(5-phenyl-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl}ethanone (1003316-51-5), (15.64) 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, (15.65) 2-chloro-5-[2-chloro-1-(2,6-difluoro-4-methoxyphenyl)-4-methyl-1H-imidazol-5-yl]pyridine, (15.66) 2-phenylphenol and salts (90-43-7), (15.67) 3-(4,4,5-trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline (861647-85-0), (15.68) 3,4,5-trichloropyridine-2,6-dicarbonitrile (17824-85-0), (15.69) 3-[5-(4-chlorophenyl)-2,3-dimethyl-1,2-oxazolidin-3-yl]pyridine, (15.70) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (15.71) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (15.72) 5-amino-1,3,4-thiadiazole-2-thiol, (15.73) 5-chloro-N'-phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide (134-31-6), (15.74) 5-fluoro-2-[(4-fluorobenzyl)oxy]pyrimidin-4-amine (1174376-11-4), (15.75) 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine (1174376-25-0), (15.76) 5-methyl-6-octyl[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, (15.77) ethyl (2Z)-3-amino-2-cyano-3-phenylprop-2-enoate, (15.78) N'-(4-{[3-(4-chlorobenzyl)-1,2,4-thiadiazol-5-yl]oxy}-2,5-dimethylphenyl)-N-ethyl-N-methylimidoformamide, (15.79) N-(4-chlorobenzyl)-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (15.80) N-[(4-chlorophenyl)(cyano)methyl]-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (15.81) N-[(5-bromo-3-chloropyridin-2-yl)methyl]-2,4-dichloropyridine-3-carboxamide, (15.82) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloropyridine-3-carboxamide, (15.83) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-iodopyridine-3-carboxamide, (15.84) N-{(E)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide (221201-92-9), (15.85) N-{(Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide (221201-92-9), (15.86) N'-{4-[(3-tert-butyl-4-cyano-1,2-thiazol-5-yl)oxy]-2-chloro-5-methylphenyl}-N-ethyl-N-methylimidoformamide, (15.87) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-1,3-thiazole-4-carboxamide (922514-49-6), (15.88) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide (922514-07-6), (15.89) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1S)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide (922514-48-5), (15.90) pentyl {6-[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylidene]amino}oxy)methylpyridin-2-yl}carbamate, (15.91) phenazine-1-carboxylic acid, (15.92) quinolin-8-ol (134-31-6), (15.93) quinolin-8-ol sulfate (2:1) (134-31-6), (15.94) tert-butyl {6-[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methylpyridin-2-yl}carbamate.

- (16) Further compounds, like for example (16.1) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (16.2) N-(4'-chlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (16.3) N-(2',4'-dichlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (16.4) 3-(difluoromethyl)-1-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (16.5) N-(2',5'-difluorobiphenyl-2-yl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (16.6) 3-(difluoromethyl)-1-methyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (16.7) 5-fluoro-1,3-dimethyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (16.8) 2-chloro-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (16.9) 3-(difluoromethyl)-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, (16.10) N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, (16.11) 3-(difluoromethyl)-N-(4'-ethynylbiphenyl-2-yl)-1-methyl-1H-pyrazole-4-carboxamide, (16.12) N-(4'-ethynylbiphenyl-2-yl)-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, (16.13) 2-chloro-N-(4'-ethynylbiphenyl-2-yl)pyridine-3-carboxamide, (16.14) 2-chloro-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (16.15) 4-(difluoromethyl)-2-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1,3-thiazole-5-carboxamide, (16.16) 5-fluoro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, (16.17) 2-chloro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (16.18) 3-(difluoromethyl)-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, (16.19) 5-fluoro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, (16.20) 2-chloro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide, (16.21) (5-bromo-2-methoxy-4-methylpyridin-3-yl)(2,3,4-trimethoxy-6-methylphenyl)methanone, (16.22) N-[2-(4-{[3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-methoxyphenyl)ethyl]-N2-(methylsulfonyl)valinamide (220706-93-4), (16.23) 4-oxo-4-[(2-phenylethyl)amino]butanoic acid, (16.24) but-3-yn-1-yl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl} carbamate, (16.25) 4-Amino-5-fluoropyrimidin-2-ol (mesomere Form: 6-Amino-5-fluoropyrimidin-2(1H)-on), (16.26) propyl 3,4,5-trihydroxybenzoate.

All named mixing partners of the classes (1) to (16) can, if their functional groups enable this, optionally form salts with suitable bases or acids.

- Particular preference is further given to combinations comprising at least one further active compound (B) selected from the following groups:

- (2.1) bixafen, (2.2) boscalid, (2.6) fluopyram, (2.8) fluxapyroxad, (2.11) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR), (2.12) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.13) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.14) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.15) isopyrazam (syn epimeric racemate 1RS,4SR,9RS), (2.16) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.17) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (2.27) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.29) N-[9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-

yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (3.1) ametocradin, (3.2) amisulbrom, (3.3) azoxystrobin, (3.4) cyazofamid, (3.9) famoxadone, (3.10) fenamidone, (3.12) fluoxastrobin, (3.16) picoxystrobin, (3.17) pyraclostrobin, (3.22) trifloxystrobin, (4.6) fluopicolide, (5.1) bordeaux mixture, (5.4) chlorothalonil, (5.5) copper hydroxide, (5.7) copper oxide, (5.8) copper oxychloride, (5.9) copper(2+) sulfate, (5.16) folpet, (5.23) mancozeb, (5.25) metiram, (5.26) metiram zinc, (5.29) propineb, (5.30) sulphur and sulphur preparations including calcium polysulphide, (7.7) pyrimethanil, (9.2) dimethomorph, (9.4) iprovalicarb, (9.5) mandipropamid, (10.9) propamocarb, (10.10) propamocarb hydrochloride, (12.9) metalaxyl, (12.10) metalaxyl-M (mefenoxam), (14.4) fluazinam, (15.9) cymoxanil, (15.24) fosetyl-aluminium, (15.25) fosetyl-calcium, (15.26) fosetyl-sodium, (15.41) phosphorous acid and its salts, (15.42) propamocarb-fosetate, (15.54) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.55) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.56) 1-(4-{4-[5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.60) 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, (15.90) pentyl {6-[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylidene]amino}oxy)methylpyridin-2-yl} carbamate.

All named mixing partners of the classes (1) to (15) can, if their functional groups enable this, optionally form salts with suitable bases or acids.

If the active compounds in the active compound combinations according to the invention are present in certain weight ratios, the synergistic effect is particularly pronounced. However, the weight ratios of the active compounds in the active compound combinations can be varied within a relatively wide range.

In the combinations according to the invention the compounds (A) and (B) are present in a synergistically effective weight ratio of A:B in a range of 50:1 to 1:5000, preferably in a weight ratio of 1:5 to 1:5000, most preferably in a weight ratio of 1:50 to 1:5000. Further ratios of A:B which can be used according to the present invention are: 1:5 to 1:50; 1:50 to 1:500; 1:500 to 1:5000.

Further preferred ratios of A:B which can be used according to the present invention are: 1:1 to 1:5000, 1:1 to 1:2000, 1:1 to 1:1000, 1:10 to 1:5000, 1:10 to 1:2000, 1:10 to 1:1000, 1:20 to 1:5000, 1:20 to 1:2000, 1:20 to 1:1000, 1:100 to 1:5000, 1:100 to 1:2000, 1:100 to 1:1000, 1:200 to 1:5000, 1:200 to 1:2000, 1:200 to 1:1000, 1:500 to 1:5000, 1:500 to 1:2000, 1:500 to 1:1000.

Where a compound (A) or a compound (B) can be present as mixtures of various possible isomeric forms, in particular of stereoisomers, such as, for example, E and Z, threo and erythro, and also optical isomers, and, if appropriate, also of tautomers. What is claimed are both the E and the Z isomers and the threo and erythro and also the optical isomers (R and S), any mixtures of these isomers, and also the possible tautomeric forms.

- Compound (A) or compound (B) having at least one basic centre are capable of forming, for example, acid addition salts, e.g. with strong inorganic acids, such as mineral acids, e.g. perchloric acid, sulfuric acid, nitric acid, nitrous acid, a phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as unsubstituted substituted, e.g. halo-substituted, C₁-C₄ alkanecarboxylic acids, e.g. acetic acid, saturated or
- 5 unsaturated dicarboxylic acids, e.g. oxalic, malonic, succinic, maleic, fumaric and phthalic acid, hydroxycarboxylic acids, e.g. ascorbic, lactic, malic, tartaric and citric acid, or benzoic acid, or with organic sulfonic acids, such as unsubstituted or substituted, e.g. halo-substituted, C₁-C₄alkane- or aryl-sulfonic acids, e.g. methane- or p-toluene-sulfonic acid. Compounds (A) or compounds (B) having at least one acid
- 10 group are capable of forming, for example, salts with bases, e.g. metal salts, such as alkali metal or alkaline earth metal salts, e.g. sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkylamine, e.g. ethyl-, diethyl-, triethyl- or dimethyl-propyl-amine, or a mono-, di- or tri-hydroxy-lower alkylamine, e.g. mono-, di- or tri-ethanolamine. In addition, corresponding internal salts may optionally be formed. In the context of the
- 15 invention, preference is given to agrochemically advantageous salts. In view of the close relationship between the compounds (A) or the compounds (B) in free form and in the form of their salts, hereinabove and herein below any reference to the free compounds (A) or free compounds (B) or to their salts should be understood as including also the corresponding salts or the free compounds (A) or free compounds (B), respectively, where appropriate and expedient. The equivalent also applies to tautomers of compounds (A) or compounds (B) and to their salts.
- 20 According to the invention the expression "combination" stands for the various combinations of compounds (A) and (B), for example in a single "ready-mix" form, in a combined spray mixture composed from separate formulations of the single active compounds, such as a "tank-mix", and in a combined use of the single active ingredients when applied in a sequential manner, i.e. one after the other with a reasonably short period, such as a few hours or days. Preferably the order of applying the compounds (A) and (B)
- 25 is not essential for working the present invention.

Composition / Formulation

- The present invention furthermore relates to compositions for combating/controlling undesirable microorganisms comprising the active compound combinations according to the invention. Preferably, the compositions are fungicidal compositions comprising agriculturally suitable auxiliaries, solvents, carriers, surfac-
- 30 tants or extenders.

- In the context of the present invention, "control of harmful microorganisms" means a reduction in infestation by harmful microorganisms, compared with the untreated plant measured as fungicidal efficacy, preferably a reduction by 25-50 %, compared with the untreated plant (100 %), more preferably a reduction by 40-79 %, compared with the untreated plant (100 %); even more preferably, the infection by harmful microorganisms is entirely suppressed (by 70-100 %). The control may be curative, i.e. for treatment of already
- 35 infected plants, or protective, for protection of plants which have not yet been infected.

An "effective but non-phytotoxic amount" means an amount of the inventive composition which is sufficient to control the fungal disease of the plant in a satisfactory manner or to eradicate the fungal disease completely, and which, at the same time, does not cause any significant symptoms of phytotoxicity. In general, this application rate may vary within a relatively wide range. It depends on several factors, for example on the
5 fungus to be controlled, the plant, the climatic conditions and the ingredients of the inventive compositions.

Suitable organic solvents include all polar and non-polar organic solvents usually employed for formulation purposes. Preferable the solvents are selected from ketones, e.g. methyl-isobutyl-ketone and cyclohexanone, amides, e.g. dimethyl formamide and alkanecarboxylic acid amides, e.g. N,N-dimethyl decaneamide and N,N-dimethyl octanamide, furthermore cyclic solvents, e.g. N-methyl-pyrrolidone, N-octyl-pyrrolidone, N-
10 dodecyl-pyrrolidone, N-octyl-caprolactame, N-dodecyl-caprolactame and butyrolactone, furthermore strong polar solvents, e.g. dimethylsulfoxide, and aromatic hydrocarbons, e.g. xylol, Solvesso™, mineral oils, e.g. white spirit, petroleum, alkyl benzenes and spindle oil, also esters, e.g. propyleneglycol-monomethylether acetate, adipic acid dibutylester, acetic acid hexylester, acetic acid heptylester, citric acid tri-*n*-butylester and phthalic acid di-*n*-butylester, and also alcohols, e.g. benzyl alcohol and 1-methoxy-2-propanol.

15 Furthermore the invention relates to a method of combating undesirable microorganisms, characterized in that the active compound combinations according to the invention are applied to the phytopathogenic fungi and/or their habitat.

According to the invention, carrier is to be understood as meaning a natural or synthetic, organic or inorganic substance which is mixed or combined with the active compounds for better applicability, in particular
20 for application to plants or plant parts or seeds. The carrier, which may be solid or liquid, is generally inert and should be suitable for use in agriculture.

According to the invention, a carrier is a natural or synthetic, organic or inorganic substance with which the active ingredients are mixed or combined for better applicability, in particular for application to plants or plant parts or seed. The carrier, which may be solid or liquid, is generally inert and should be suitable
25 for use in agriculture.

Useful solid or liquid carriers include: for example ammonium salts and natural rock dusts, such as kaolins, clays, talc, chalk, quartz, attapulgit, montmorillonite or diatomaceous earth, and synthetic rock dusts, such as finely divided silica, alumina and natural or synthetic silicates, resins, waxes, solid fertilizers, water, alcohols, especially butanol, organic solvents, mineral and vegetable oils, and derivatives thereof. Mix-
30 tures of such carriers can likewise be used.

Suitable solid filler and carrier include inorganic particles, e.g. carbonates, silicates, sulphates and oxides with an average particle size of between 0.005 and 20 µm, preferably of between 0.02 to 10 µm, for example ammonium sulphate, ammonium phosphate, urea, calcium carbonate, calcium sulphate, magnesium sulphate, magnesium oxide, aluminium oxide, silicium dioxide, so-called fine-particle silica, silica gels,

natural or synthetic silicates, and aluminosilicates and plant products like cereal flour, wood powder/sawdust and cellulose powder.

Useful solid carriers for granules include: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite, dolomite, and synthetic granules of inorganic and organic meals, and also
5 granules of organic material such as sawdust, coconut shells, maize cobs and tobacco stalks.

Useful liquefied gaseous extenders or carriers are those liquids which are gaseous at standard temperature and under standard pressure, for example aerosol propellants such as halohydrocarbons, and also butane, propane, nitrogen and carbon dioxide.

In the formulations, it is possible to use tackifiers such as carboxymethylcellulose, and natural and synthetic
10 polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids, such as cephalins and lecithins, and synthetic phospholipids. Further additives may be mineral and vegetable oils.

If the extender used is water, it is also possible for example, to use organic solvents as auxiliary solvents. Suitable liquid solvents are essentially: aromatic compounds, such as xylene, toluene or alkylnaphthalenes,
15 chlorinated aromatic compounds or chlorinated aliphatic hydrocarbons, such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons, such as cyclohexane or paraffins, for example mineral oil fractions, mineral and vegetable oils, alcohols, such as butanol or glycol, and also ethers and esters thereof, ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents, such as dimethylformamide and dimethyl sulfoxide, and also water.

20 The compositions according to the invention may comprise additional further components, such as, for example, surfactants. Suitable surfactants are emulsifiers, dispersants or wetting agents having ionic or nonionic properties, or mixtures of these surfactants. Examples of these are salts of polyacrylic acid, salts of lignosulphonic acid, salts of phenolsulphonic acid or naphthalenesulphonic acid, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (preferably
25 alkylphenols or arylphenols), salts of sulphosuccinic esters, taurine derivatives (preferably alkyl taurates), phosphoric esters of polyethoxylated alcohols or phenols, fatty esters of polyols, and derivatives of the compounds containing sulphates, sulphonates and phosphates. The presence of a surfactant is required if one of the active compounds and/or one of the inert carriers is insoluble in water and when the application takes place in water. The proportion of surfactants is between 5 and 40 per cent by weight of the composition according to the invention.
30

Suitable surfactants (adjuvants, emulsifiers, dispersants, protective colloids, wetting agent and adhesive) include all common ionic and non-ionic substances, for example ethoxylated nonylphenols, polyalkylene glycolether of linear or branched alcohols, reaction products of alkyl phenols with ethylene oxide and/or propylene oxide, reaction products of fatty acid amines with ethylene oxide and/or propylene oxide, furthermore fatty acid esters, alkyl sulfonates, alkyl sulphates, alkyl ethersulphates, alkyl etherphosphates,
35

arylsulphate, ethoxylated arylalkylphenols, e.g. tristyryl-phenol-ethoxylates, furthermore ethoxylated and propoxylated arylalkylphenols like sulphated or phosphated arylalkylphenol-ethoxylates and -ethoxy- and -propoxylates. Further examples are natural and synthetic, water soluble polymers, e.g. lignosulphonates, gelatine, gum arabic, phospholipides, starch, hydrophobic modified starch and cellulose derivatives, in particular cellulose ester and cellulose ether, further polyvinyl alcohol, polyvinyl acetate, polyvinyl pyrrolidone, polyacrylic acid, polymethacrylic acid and co-polymerisates of (meth)acrylic acid and (meth)acrylic acid esters, and further co-polymerisates of methacrylic acid and methacrylic acid esters which are neutralized with alkalimetal hydroxide and also condensation products of optionally substituted naphthalene sulfonic acid salts with formaldehyde.

10 It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide, Prussian blue, and organic dyes, such as alizarin dyes, azo dyes and metal phthalocyanine dyes, and trace nutrients, such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Antifoams which may be present in the formulations include e.g. silicone emulsions, longchain alcohols, fatty acids and their salts as well as fluoroorganic substances and mixtures thereof.

15 Examples of thickeners are polysaccharides, e.g. xanthan gum or veegum, silicates, e.g. attapulgite, bentonite as well as fine-particle silica.

If appropriate, other additional components may also be present, for example protective colloids, binders, adhesives, thickeners, thixotropic substances, penetrants, stabilizers, sequestering agents, complex formers. In general, the active compounds can be combined with any solid or liquid additive customarily used for formulation purposes.

The inventive compositions can be used as such or, depending on their particular physical and/or chemical properties, in the form of their formulations or the use forms prepared therefrom, such as aerosols, capsule suspensions, cold-fogging concentrates, warm-fogging concentrates, encapsulated granules, fine granules, flowable concentrates for the treatment of seed, ready-to-use solutions, dustable powders, emulsifiable concentrates, oil-in-water emulsions, water-in-oil emulsions, macrogranules, microgranules, oil-dispersible powders, oil-miscible flowable concentrates, oil-miscible liquids, gas (under pressure), gas generating product, foams, pastes, pesticide coated seed, suspension concentrates, suspoemulsion concentrates, soluble concentrates, suspensions, wettable powders, soluble powders, dusts and granules, water-soluble and water-dispersible granules or tablets, water-soluble and water-dispersible powders for the treatment of seed, wettable powders, natural products and synthetic substances impregnated with active ingredient, and also microencapsulations in polymeric substances and in coating materials for seed, and also ULV cold-fogging and warm-fogging formulations.

The inventive compositions include not only formulations which are already ready for use and can be applied with a suitable apparatus to the plant or the seed, but also commercial concentrates which have to be diluted with water prior to use. Customary applications are for example dilution in water and subsequent

spraying of the resulting spray liquor, application after dilution in oil, direct application without dilution, seed treatment or soil application of granules.

In general, the compositions according to the invention comprise between 0.05 and 99 per cent by weight, 0.01 and 98 per cent by weight, preferable between 0.1 and 95 per cent by weight, particularly preferred
5 between 0.5 and 90 per cent by weight of the active compound combination according to the invention, very particularly preferable between 10 and 70 per cent by weight.

The contents of active ingredient in the application forms prepared from the commercial formulations may vary in a broad range. The concentration of the active ingredients in the application forms is generally between 0.000001 to 95 % by weight, preferably between 0.0001 and 2 % by weight.

10 The formulations mentioned can be prepared in a manner known per se, for example by mixing the active ingredients with at least one customary extender, solvent or diluent, adjuvant, emulsifier, dispersant, and/or binder or fixative, wetting agent, water repellent, if appropriate desiccants and UV stabilizers and, if appropriate, dyes and pigments, antifoams, preservatives, inorganic and organic thickeners, adhesives, gibberellins and also further processing auxiliaries and also water. Depending on the formulation type to be prepared further
15 processing steps are necessary, e.g. wet grinding, dry grinding and granulation.

The inventive composition may be present as such or in their (commercial) formulations and in the use forms prepared from these formulations as a mixture with other (known) active ingredients, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators, herbicides, fertilizers, safeners and/or semiochemicals.

20 The compositions according to the invention do not only comprise ready-to-use compositions which can be applied with suitable apparatus to the plant or the seed, but also commercial concentrates which have to be diluted with water prior to use.

The treatment according to the invention of the plants and plant parts with the active compounds or compositions is carried out directly or by action on their surroundings, habitat or storage space using customary
25 treatment methods, for example by dipping, spraying, atomizing, irrigating, evaporating, dusting, fogging, broadcasting, foaming, painting, spreading-on, watering (drenching), drip irrigating and, in the case of propagation material, in particular in the case of seeds, furthermore as a powder for dry seed treatment, a solution for seed treatment, a water-soluble powder for slurry treatment, by incrusting, by coating with one or more layers, etc. It is furthermore possible to apply the active compounds by the ultra-low volume
30 method, or to inject the active compound preparation or the active compound itself into the soil.

Plant/Crop Protection

The inventive active ingredients or compositions have potent microbicidal activity and can be used for control of unwanted microorganisms, such as fungi and bacteria, in crop protection and in the protection of materials.

The invention also relates to a method for controlling unwanted microorganisms, characterized in that the inventive active ingredients are applied to the phytopathogenic fungi, phytopathogenic bacteria and/or their habitat.

5 Fungicides can be used in crop protection for control of phytopathogenic fungi. They are characterized by an outstanding efficacy against a broad spectrum of phytopathogenic fungi, including soilborne pathogens, which are in particular members of the classes *Plasmodiophoromycetes*, *Peronosporomycetes* (Syn. *Oomycetes*), *Chytridiomycetes*, *Zygomycetes*, *Ascomycetes*, *Basidiomycetes* and *Deuteromycetes* (Syn. *Fungi imperfecti*). Some fungicides are systemically active and can be used in plant protection as foliar, seed dressing or soil fungicide. Furthermore, they are suitable for combating fungi, which inter alia infest wood
10 or roots of plant.

Bactericides can be used in crop protection for control of *Pseudomonadaceae*, *Rhizobiaceae*, *Enterobacteriaceae*, *Corynebacteriaceae* and *Streptomycetaceae*.

Non-limiting examples of pathogens of fungal diseases which can be treated in accordance with the invention include:

15 diseases caused by powdery mildew pathogens, for example *Blumeria* species, for example *Blumeria graminis*; *Podosphaera* species, for example *Podosphaera leucotricha*; *Sphaerotheca* species, for example *Sphaerotheca fuliginea*; *Uncinula* species, for example *Uncinula necator*;

diseases caused by rust disease pathogens, for example *Gymnosporangium* species, for example *Gymnosporangium sabinae*; *Hemileia* species, for example *Hemileia vastatrix*; *Phakopsora* species, for example
20 *Phakopsora pachyrhizi* and *Phakopsora meibomiae*; *Puccinia* species, for example *Puccinia recondite*, *P. trititica*, *P. graminis* or *P. striiformis*; *Uromyces* species, for example *Uromyces appendiculatus*;

diseases caused by pathogens from the group of the *Oomycetes*, for example *Albugo* species, for example *Albugo candida*; *Bremia* species, for example *Bremia lactucae*; *Peronospora* species, for example *Peronospora pisi* or *P. brassicae*; *Phytophthora* species, for example *Phytophthora infestans*; *Plasmopara*
25 species, for example *Plasmopara viticola*; *Pseudoperonospora* species, for example *Pseudoperonospora humuli* or *Pseudoperonospora cubensis*; *Pythium* species, for example *Pythium ultimum*;

leaf blotch diseases and leaf wilt diseases caused, for example, by *Alternaria* species, for example *Alternaria solani*; *Cercospora* species, for example *Cercospora beticola*; *Cladosporium* species, for example *Cladosporium cucumerinum*; *Cochliobolus* species, for example *Cochliobolus sativus* (conidia form: Drechslera, Syn: Helminthosporium), *Cochliobolus miyabeanus*; *Colletotrichum* species, for example
30 *Colletotrichum lindemuthianum*; *Cycloconium* species, for example *Cycloconium oleaginum*; *Diaporthe* species, for example *Diaporthe citri*; *Elsinoe* species, for example *Elsinoe fawcettii*; *Gloeosporium* species, for example *Gloeosporium laeticolor*; *Glomerella* species, for example *Glomerella cingulata*; *Guignardia* species, for example *Guignardia bidwelli*; *Leptosphaeria* species, for example *Leptosphaeria maculans*, *Leptosphaeria nodorum*; *Magnaporthe* species, for example *Magnaporthe grisea*; *Microdochi-*
35

um species, for example *Microdochium nivale*; *Mycosphaerella* species, for example *Mycosphaerella graminicola*, *M. arachidicola* and *M. fijiensis*; *Phaeosphaeria* species, for example *Phaeosphaeria nodorum*; *Pyrenophora* species, for example *Pyrenophora teres*, *Pyrenophora tritici repentis*; *Ramularia* species, for example *Ramularia collo-cygni*, *Ramularia areola*; *Rhynchosporium* species, for example
 5 *Rhynchosporium secalis*; *Septoria* species, for example *Septoria apii*, *Septoria lycopersii*; *Typhula* species, for example *Typhula incarnata*; *Venturia* species, for example *Venturia inaequalis*;

root and stem diseases caused, for example, by *Corticium* species, for example *Corticium graminearum*; *Fusarium* species, for example *Fusarium oxysporum*; *Gaeumannomyces* species, for example *Gaeumannomyces graminis*; *Rhizoctonia* species, such as, for example *Rhizoctonia solani*; *Sarocladium* diseases
 10 caused for example by *Sarocladium oryzae*; *Sclerotium* diseases caused for example by *Sclerotium oryzae*; *Tapesia* species, for example *Tapesia acuformis*; *Thielaviopsis* species, for example *Thielaviopsis basicola*;

ear and panicle diseases (including corn cobs) caused, for example, by *Alternaria* species, for example *Alternaria* spp.; *Aspergillus* species, for example *Aspergillus flavus*; *Cladosporium* species, for example
 15 *Cladosporium cladosporioides*; *Claviceps* species, for example *Claviceps purpurea*; *Fusarium* species, for example *Fusarium culmorum*; *Gibberella* species, for example *Gibberella zeae*; *Monographella* species, for example *Monographella nivalis*; *Septoria* species, for example *Septoria nodorum*;

diseases caused by smut fungi, for example *Sphacelotheca* species, for example *Sphacelotheca reiliana*; *Tilletia* species, for example *Tilletia caries*, *T. controversa*; *Urocystis* species, for example *Urocystis occulta*;
 20 *Ustilago* species, for example *Ustilago nuda*, *U. nuda tritici*;

fruit rot caused, for example, by *Aspergillus* species, for example *Aspergillus flavus*; *Botrytis* species, for example *Botrytis cinerea*; *Penicillium* species, for example *Penicillium expansum* and *P. purpurogenum*; *Sclerotinia* species, for example *Sclerotinia sclerotiorum*; *Verticillium* species, for example *Verticillium albo-*
tratum;

seed and soilborne decay, mould, wilt, rot and damping-off diseases caused, for example, by *Alternaria* species, caused for example by *Alternaria brassicicola*; *Aphanomyces* species, caused for example by *Aphanomyces euteiches*; *Ascochyta* species, caused for example by *Ascochyta lentis*; *Aspergillus* species, caused for example by *Aspergillus flavus*; *Cladosporium* species, caused for example by *Cladosporium herbarum*; *Cochliobolus* species, caused for example by *Cochliobolus sativus*; (Conidiaform: Drechslera,
 25 Bipolaris Syn: Helminthosporium); *Colletotrichum* species, caused for example by *Colletotrichum cocco-*
 30 *codes*; *Fusarium* species, caused for example by *Fusarium culmorum*; *Gibberella* species, caused for example by *Gibberella zeae*; *Macrophomina* species, caused for example by *Macrophomina phaseolina*; *Monographella* species, caused for example by *Monographella nivalis*; *Penicillium* species, caused for example by *Penicillium expansum*; *Phoma* species, caused for example by *Phoma lingam*; *Phomopsis* species, caused for example by *Phomopsis sojae*; *Phytophthora* species, caused for example by *Phytophthora cactorum*; *Pyrenophora* species, caused for example by *Pyrenophora graminea*; *Pyricularia* species,
 35

caused for example by *Pyricularia oryzae*; *Pythium* species, caused for example by *Pythium ultimum*; *Rhizoctonia* species, caused for example by *Rhizoctonia solani*; *Rhizopus* species, caused for example by *Rhizopus oryzae*; *Sclerotium* species, caused for example by *Sclerotium rolfsii*; *Septoria* species, caused for example by *Septoria nodorum*; *Typhula* species, caused for example by *Typhula incarnata*; *Verticillium* species, caused for example by *Verticillium dahliae*;

cancers, galls and witches' broom caused, for example, by *Nectria* species, for example *Nectria galligena*;

wilt diseases caused, for example, by *Monilinia* species, for example *Monilinia laxa*;

leaf blister or leaf curl diseases caused, for example, by *Exobasidium* species, for example *Exobasidium vexans*;

10 *Taphrina* species, for example *Taphrina deformans*;

decline diseases of wooden plants caused, for example, by Esca disease, caused for example by *Phaeo-niella clamydospora*, *Phaeoacremonium aleophilum* and *Fomitiporia mediterranea*; Eutypa dyebark, caused for example by *Eutypa lata* ; Ganoderma diseases caused for example by *Ganoderma boninense*; Rigidoporus diseases caused for example by *Rigidoporus lignosus*;

15 diseases of flowers and seeds caused, for example, by *Botrytis* species, for example *Botrytis cinerea*;

diseases of plant tubers caused, for example, by *Rhizoctonia* species, for example *Rhizoctonia solani*; *Helminthosporium* species, for example *Helminthosporium solani*;

Club root caused, for example, by *Plasmodiophora* species, for example *Plasmodiophora brassicae*;

diseases caused by bacterial pathogens, for example *Xanthomonas* species, for example *Xanthomonas campestris* pv. *oryzae*; *Pseudomonas* species, for example *Pseudomonas syringae* pv. *lachrymans*; *Erwinia* species, for example *Erwinia amylovora*.

The following diseases of soya beans can be controlled with preference:

Fungal diseases on leaves, stems, pods and seeds caused, for example, by *Alternaria* leaf spot (*Alternaria spec. atrans tenuissima*), Anthracnose (*Colletotrichum gloeosporoides dematium* var. *truncatum*), brown spot (*Septoria glycines*), cercospora leaf spot and blight (*Cercospora kikuchii*), choanephora leaf blight (*Choanephora infundibulifera trisporea* (Syn.)), dactuliophora leaf spot (*Dactuliophora glycines*), downy mildew (*Peronospora manshurica*), drechslera blight (*Drechslera glycini*), frog-eye leaf spot (*Cercospora sojae*), leptosphaerulina leaf spot (*Leptosphaerulina trifolii*), phyllosticta leaf spot (*Phyllosticta sojaecola*), pod and stem blight (*Phomopsis sojae*), powdery mildew (*Microsphaera diffusa*), pyrenochaeta leaf spot (*Pyrenochaeta glycines*), rhizoctonia aerial, foliage, and web blight (*Rhizoctonia solani*), rust (*Phakopsora pachyrhizi*, *Phakopsora meibomia*), scab (*Sphaceloma glycines*), stemphylium leaf blight (*Stemphylium botryosum*), target spot (*Corynespora cassiicola*).

Fungal diseases on roots and the stem base caused, for example, by black root rot (*Calonectria crotalariae*), charcoal rot (*Macrophomina phaseolina*), fusarium blight or wilt, root rot, and pod and collar rot (*Fusarium oxysporum*, *Fusarium orthoceras*, *Fusarium semitectum*, *Fusarium equiseti*), mycoleptodiscus root rot (*Mycoleptodiscus terrestris*), neocosmospora (*Neocosmospora vasinfecta*), pod and stem blight
 5 (*Diaporthe phaseolorum*), stem canker (*Diaporthe phaseolorum* var. *caulivora*), phytophthora rot (*Phytophthora megasperma*), brown stem rot (*Phialophora gregata*), pythium rot (*Pythium aphanidermatum*, *Pythium irregulare*, *Pythium debaryanum*, *Pythium myriotylum*, *Pythium ultimum*), rhizoctonia root rot, stem decay, and damping-off (*Rhizoctonia solani*), sclerotinia stem decay (*Sclerotinia sclerotiorum*), sclerotinia southern blight (*Sclerotinia rolfsii*), thielaviopsis root rot (*Thielaviopsis basicola*).

10 The inventive fungicidal compositions can be used for curative or protective/preventive control of phytopathogenic fungi. The invention therefore also relates to curative and protective methods for controlling phytopathogenic fungi by the use of the inventive active ingredients or compositions, which are applied to the seed, the plant or plant parts, the fruit or the soil in which the plants grow.

The fact that the active ingredients are well tolerated by plants at the concentrations required for controlling plant diseases allows the treatment of above-ground parts of plants, of propagation stock and seeds,
 15 and of the soil.

According to the invention all plants and plant parts can be treated. By plants is meant all plants and plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by
 20 conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering methods. By plant parts is meant all above ground and below ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed
 25 as well as roots, corms and rhizomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners and seeds also belong to plant parts.

The inventive active compositions, when they are well tolerated by plants, have favourable homeotherm toxicity and are well tolerated by the environment, are suitable for protecting plants and plant organs, for enhancing harvest yields, for improving the quality of the harvested material. They can preferably be used
 30 as crop protection compositions. They are active against normally sensitive and resistant species and against all or some stages of development.

Plants which can be treated in accordance with the invention include the following main crop plants: maize, soya bean, alfalfa, cotton, sunflower, *Brassica* oil seeds such as *Brassica napus* (e.g. canola, rapeseed), *Brassica rapa*, *B. juncea* (e.g. (field) mustard) and *Brassica carinata*, *Arecaceae* sp. (e.g. oilpalm,
 35 coconut), rice, wheat, sugar beet, sugar cane, oats, rye, barley, millet and sorghum, triticale, flax, nuts, grapes and vine and various fruit and vegetables from various botanic taxa, e.g. *Rosaceae* sp. (e.g. pome

fruits such as apples and pears, but also stone fruits such as apricots, cherries, almonds, plums and peaches, and berry fruits such as strawberries, raspberries, red and black currant and gooseberry), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp. (e.g. olive tree), *Actinidaceae* sp., *Lauraceae* sp. (e.g. avocado, cinnamon, camphor), *Musaceae* sp. (e.g. banana trees and plantations), *Rubiaceae* sp. (e.g. coffee), *Theaceae* sp. (e.g. tea), *Sterculiaceae* sp., *Rutaceae* sp. (e.g. lemons, oranges, mandarins and grapefruit); *Solanaceae* sp. (e.g. tomatoes, potatoes, peppers, capsicum, aubergines, tobacco), *Liliaceae* sp., *Compositae* sp. (e.g. lettuce, artichokes and chicory – including root chicory, endive or common chicory), *Umbelliferae* sp. (e.g. carrots, parsley, celery and celeriac), *Cucurbitaceae* sp. (e.g. cucumbers – including gherkins, pumpkins, watermelons, calabashes and melons), *Alliaceae* sp. (e.g. leeks and onions), *Cruciferae* sp. (e.g. white cabbage, red cabbage, broccoli, cauliflower, Brussels sprouts, pak choi, kohlrabi, radishes, horseradish, cress and chinese cabbage), *Leguminosae* sp. (e.g. peanuts, peas, lentils and beans – e.g. common beans and broad beans), *Chenopodiaceae* sp. (e.g. Swiss chard, fodder beet, spinach, beetroot), *Linaceae* sp. (e.g. hemp), *Cannabaceae* sp. (e.g. cannabis), *Malvaceae* sp. (e.g. okra, cocoa), *Papaveraceae* (e.g. poppy), *Asparagaceae* (e.g. asparagus); useful plants and ornamental plants in the garden and woods including turf, lawn, grass and *Stevia rebaudiana*; and in each case genetically modified types of these plants.

In particular, the mixtures and compositions according to the invention are suitable for controlling the following plant diseases:

Albugo spp. (white rust) on ornamental plants, vegetable crops (e.g. *A. candida*) and sunflowers (e.g. *A. tragopogonis*); *Alternaria* spp. (black spot disease, black blotch) on vegetables, oilseed rape (e.g. *A. brassicola* or *A. brassicae*), sugar beet (e.g. *A. tenuis*), fruit, rice, soybeans and also on potatoes (e.g. *A. solani* or *A. alternata*) and tomatoes (e.g. *A. solani* or *A. alternata*) and *Alternaria* spp. (black head) on wheat; *Aphanomyces* spp. on sugar beet and vegetables; *Ascochyta* spp. on cereals and vegetables, e.g. *A. tritici* (*Ascochyta* leaf blight) on wheat and *A. hordei* on barley; *Bipolaris* and *Drechslera* spp. (teleomorph: *Cochliobolus* spp.), e.g. leaf spot diseases (*D. maydis* and *B. zeicola*) on corn, e.g. glume blotch (*B. sorokiniana*) on cereals and e.g. *B. oryzae* on rice and on lawn; *Blumeria* (old name: *Erysiphe*) *graminis* (powdery mildew) on cereals (e.g. wheat or barley); *Botryosphaeria* spp. ('Slack Dead Arm Disease') on grapevines (e.g. *B. obtusa*); *Botrytis cinerea* (teleomorph: *Botryotinia fuckeliana*: gray mold, gray rot) on soft fruit and pomaceous fruit (inter alia strawberries), vegetables (inter alia lettuce, carrots, celeriac and cabbage), oilseed rape, flowers, grapevines, forest crops and wheat (ear mold); *Bremia lactucae* (downy mildew) on lettuce; *Ceratocystis* (syn. *Ophiostoma*) spp. (blue stain fungus) on deciduous trees and coniferous trees, e.g. *C. ulmi* (Dutch elm disease) on elms; *Cercospora* spp. (*Cercospora* leaf spot) on corn (e.g. *C. zea-maydis*), rice, sugar beet (e.g. *C. beticola*), sugar cane, vegetables, coffee, soybeans (e.g. *C. sojae* or *C. kikuchii*) and rice; *Cladosporium* spp. on tomato (e.g. *C. fulvum*: tomato leaf mold) and cereals, e.g. *C. herbarum* (ear rot) on wheat; *Claviceps purpurea* (ergot) on cereals; *Cochliobolus* (anamorph: *Helminthosporium* or *Bipolaris*) spp. (leaf spot) on corn (e.g. *C. carbonum*), cereals (e.g. *C. sativus*, anamorph: *B. sorokiniana*: glume blotch) and rice (for example *C. miyabeanus*, anamorph: *H. oryzae*); *Colletotrichum* (teleomorph: *Glomerella*) spp. (anthracnose) on cotton (e.g. *C. gossypii*), corn (e.g. *C. graminicola*).

la: stem rot and anthracnosis), soft fruit, potatoes (e.g. *C. coccodes*: wilt disease), beans (e.g. *C. lindemuthianum*) and soybeans (e.g. *C. truncatum*); *Corticium* spp., e.g. *C. sasakii* (sheath blight) on rice; *Corynespora cassiicola* (leaf spot) on soybeans and ornamental plants; *Cyloconium* spp., e.g. *C. oleaginum* on olives; *Cylindrocarpon* spp. (e.g. fruit tree cancer or black foot disease of grapevine, teleomorph: *Nectria* or *Neonectria* spp.) on fruit trees, grapevines (e.g. *C. liriodendri*; teleomorph: *Neonectria liriodendri*, black foot disease) and many ornamental trees; *Dematophora* (teleomorph: *Rosellinia*) *necatrix* (root/stem rot) on soybeans; *Diaporthe* spp. e.g. *D. phaseolorum* (stem disease) on soybeans; *Drechslera* (syn. *Helminthosporium*, teleomorph: *Pyrenophora*) spp. on corn, cereals, such as barley (e.g. *D. teres*, net blotch) and on wheat (e.g. *D. tritici-repentis*: DTR leaf spot), rice and lawn; Esca disease (dieback of grapevine, apoplexia) on grapevines, caused by *Formitiporia* (syn. *Phellinus*) *punctata*, *F. mediterranea*. *Phaeomoniella chlamydospora* (old name *Phaeoacremonium chlamydosporum*), *Phaeoacremonium aleophilum* and/or *Botryosphaeria obtusa*; *Elsinoe* spp. on pome fruit (*E. pyri*) and soft fruit (*E. veneta*: anthracnosis) and also grapevines (*E. ampelina*: anthracnosis); *Entyloma oryzae* (leaf smut) on rice; *Epicoccum* spp. (black head) on wheat; *Erysiphe* spp. (powdery mildew) on sugar beet (*E. betae*), vegetables (e.g. *E. pisi*), such as cucumber species (e.g. *E. cichoracearum*) and cabbage species, such as oilseed rape (e.g. *E. cruciferarum*); *Eutypa fata* (*Eutypa* cancer or dieback, anamorph: *Cytosporina lata*, syn. *Libertella blepharis*) on fruit trees, grapevines and many ornamental trees; *Exserohilum* (syn. *Helminthosporium*) spp. on corn (e.g. *E. turcicum*); *Fusarium* (teleomorph: *Gibberella*) spp. (wilt disease, root and stem rot) on various plants, such as e.g. *F. graminearum* or *F. culmorum* (root rot and silver-top) on cereals (e.g. wheat or barley), *F. oxysporum* on tomatoes, *F. solani* on soybeans and *F. verticillioides* on corn; *Gaeumannomyces graminis* (takeall) on cereals (e.g. wheat or barley) and corn; *Gibberella* spp. on cereals (e.g. *G. zeae*) and rice (e.g. *G. fujikuroi*: bakanae disease); *Glomerella cingulata* on grapevines, pomeaceous fruit and other plants and *G. gossypii* on cotton; grainstaining complex on rice; *Guignardia bidwellii* (black rot) on grapevines; *Gymnosporangium* spp. on Rosaceae and juniper, e.g. *G. sabinae* (pear rust) on pears; *Helminthosporium* spp. (syn. *Drechslera*, teleomorph: *Cochliobolus*) on corn, cereals and rice; *Hemileia* spp., e.g. *H. vastatrix* (coffee leaf rust) on coffee; *Isariopsis clavispora* (syn. *Cladosporium vitis*) on grapevines; *Macrophomina phaseolina* (syn. *phaseoli*) (root/stem rot) on soybeans and cotton; *Microdochium* (syn. *Fusarium*) *nivale* (pink snow mold) on cereals (e.g. wheat or barley); *Microsphaera diffusa* (powdery mildew) on soybeans; *Monilinia* spp., e.g. *M. laxa*, *M. fructicola* and *M. fructigena* (blossom and twig blight) on stone fruit and other Rosaceae; *Mycosphaerella* spp. on cereals, bananas, soft fruit and peanuts, such as e.g. *M. graminicola* (anamorph: *Septoria tritici*, *Septoria* leaf blotch) on wheat or *M. fijiensis* (Sigatoka disease) on bananas; *Peronospora* spp. (downy mildew) on cabbage (e.g. *P. brassicae*), oilseed rape (e.g. *P. parasitica*), bulbous plants (e.g. *P. destructor*), tobacco (*P. tabacina*) and soybeans (e.g. *P. manshurica*); *Phakopsora pachyrhizi* and *P. meibomia* (soybean rust) on soybeans; *Phialophora* spp. e.g. on grapevines (e.g. *P. tracheiphila* and *P. tetraspora*) and soybeans (e.g. *P. gregata*: stem disease); *Phoma lingam* (root and stem rot) on oilseed rape and cabbage and *P. betae* (leaf spot) on sugar beet; *Phomopsis* spp. on sunflowers, grapevines (e.g. *P. viticola*: dead-arm disease) and soybeans (e.g. stem canker/stem blight: *P. phaseoli*, teleomorph: *Diaporthe phaseolorum*); *Physoderma maydis* (brown spot) on corn; *Phytophthora* spp. (wilt disease, root, leaf, stem and fruit rot) on various plants, such as on

bell peppers and cucumber species (e.g. *P. capsici*), soybeans (e.g. *P. megasperma*, syn. *P. sojae*), potatoes and tomatoes (e.g. *P. infestans*, late blight and brown rot) and deciduous trees (e.g. *P. ramorum* sudden oak death); *Plasmodiophora brassicae* (club-root) on cabbage, oilseed rape, radish and other plants; *Plasmopara* spp., e.g. *P. viticola* (peronospora of grapevines, downy mildew) on grapevines and *P. halstedii* on sunflowers; *Podosphaera* spp. (powdery mildew) on Rosaceae, hops, pomaceous fruit and soft fruit, e.g. *P. leucotricha* on apple; *Polymyxa* spp., e.g. on cereals, such as barley and wheat (*P. graminis*) and sugar beet (*P. betae*) and the viral diseases transmitted thereby; *Pseudocercospora herpotrichoides* (eyespot/stem break, teleomorph: *Tapesia yallundae*) on cereals, e.g. wheat or barley; *Pseudoperonospora* (downy mildew) on various plants, e.g. *P. cubensis* on cucumber species or *P. humili* on hops; *Pseudopezizicola tracheiphila* (angular leaf scorch, anamorph *Phialophora*) on grapevines; *Puccinia* spp. (rust disease) on various plants, e.g. *P. tritici* (brown rust of wheat), *P. striiformis* (yellow rust), *P. hordei* (dwarf leaf rust), *P. graminis* (black rust) or *P. recondita* (brown rust of rye) on cereals, such as e.g. wheat, barley or rye, *P. kuehnii* on sugar cane and, e.g., on asparagus (e.g. *P. asparagi*); *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (speckled leaf blotch) on wheat or *P. teres* (net blotch) on barley; *Pyricularia* spp., e.g. *P. oryzae* (teleomorph: *Magnaporthe grisea*, rice blast) on rice and *P. grisea* on lawn and cereals; *Pythium* spp. (damping-off disease) on lawn, rice, corn, wheat, cotton, oilseed rape, sunflowers, sugar beet, vegetables and other plants (e.g. *P. ultimum* or *P. aphanidermatum*); *Ramularia* spp., e.g. *R. collo-cygni* (*Ramularia* leaf and lawn spot/physiological leaf spot) on barley and *R. beticola* on sugar beet; *Rhizoctonia* spp. on cotton, rice, potatoes, lawn, corn, oilseed rape, potatoes, sugar beet, vegetables and on various other plants, for example *R. solani* (root and stem rot) on soybeans, *R. solani* (sheath blight) on rice or *R. cerealis* (sharp eyespot) on wheat or barley; *Rhizopus stolonifer* (soft rot) on strawberries, carrots, cabbage, grapevines and tomato; *Rhynchosporium secalis* (leaf spot) on barley, rye and triticale; *Sarocladium oryzae* and *S. attenuatum* (sheath rot) on rice; *Sclerotinia* spp. (stem or white rot) on vegetable and field crops, such as oilseed rape, sunflowers (e.g. *Sclerotinia sclerotiorum*) and soybeans (e.g. *S. rolfsii*); *Septoria* spp. on various plants, e.g. *S. glycines* (leaf spot) on soybeans, *S. tritici* (*Septoria* leaf blotch) on wheat and *S.* (syn. *Stagonospora*) *nodorum* (leaf blotch and glume blotch) on cereals; *Uncinula* (syn. *Erysiphe*) *necator* (powdery mildew, anamorph: *Oidium tuckeri*) on grapevines; *Setosphaeria* spp. (leaf spot) on corn (e.g. *S. turcicum*, syn. *Helminthosporium turcicum*) and lawn; *Sphacelotheca* spp. (head smut) on corn, (e.g. *S. reiliana*: kernel smut), millet and sugar cane; *Sphaerotheca fuliginea* (powdery mildew) on cucumber species; *Spongospora subterranea* (powdery scab) on potatoes and the viral diseases transmitted thereby; *Stagonospora* spp. on cereals, e.g. *S. nodorum* (leaf blotch and glume blotch, teleomorph: *Leptosphaeria* [syn. *Phaeosphaeria*] *nodorum*) on wheat; *Synchytrium endobioticum* on potatoes (potato wart disease); *Taphrina* spp., e.g. *T. deformans* (curly-leaf disease) on peach and *T. pruni* (plum-pocket disease) on plums; *Thielaviopsis* spp. (black root rot) on tobacco, pome fruit, vegetable crops, soybeans and cotton, e.g. *T. basicola* (syn. *Chalara elegans*); *Tilletia* spp. (bunt or stinking smut) on cereals, such as e.g. *T. tritici* (syn. *T. caries*, wheat bunt) and *T. controversa* (dwarf bunt) on wheat; *Typhula incarnata* (gray snow mold) on barley or wheat; *Urocystis* spp., e.g. *U. occulta* (flag smut) on rye; *Uromyces* spp. (rust) on vegetable plants, such as beans (e.g. *U. appendiculatus*, syn. *U. phaseoli*) and sugar beet (e.g. *U. betae*); *Ustilago* spp. (loose smut) on cereals (e.g. *U. nuda* and *U. avenae*), corn (e.g. *U. maydis*:

corn smut) and sugar cane; *Venturia* spp. (scab) on apples (e.g. *V. inaequalis*) and pears and *Verticillium* spp. (leaf and shoot wilt) on various plants, such as fruit trees and ornamental trees, grapevines, soft fruit, vegetable and field crops, such as e.g. *V. dahliae* on strawberries, oilseed rape, potatoes and tomatoes.

Plant Growth Regulation

- 5 In some cases, the inventive compositions can, at particular concentrations or application rates, also be used as herbicides, safeners, growth regulators or agents to improve plant properties, or as microbicides, for example as fungicides, antimycotics, bactericides, viricides (including compositions against viroids) or as compositions against MLO (Mycoplasma-like organisms) and RLO (Rickettsia-like organisms). If appropriate, they can also be used as intermediates or precursors for the synthesis of other active ingredients.
- 10 The inventive active combinations intervene in the metabolism of the plants and can therefore also be used as growth regulators.

Plant growth regulators may exert various effects on plants. The effect of the substances depends essentially on the time of application in relation to the developmental stage of the plant, and also on the amounts of active ingredient applied to the plants or their environment and on the type of application. In each case,
15 growth regulators should have a particular desired effect on the crop plants.

Plant growth-regulating compounds can be used, for example, to inhibit the vegetative growth of the plants. Such inhibition of growth is of economic interest, for example, in the case of grasses, since it is thus possible to reduce the frequency of grass cutting in ornamental gardens, parks and sport facilities, on roadsides, at airports or in fruit crops. Also of significance is the inhibition of the growth of herbaceous
20 and woody plants on roadsides and in the vicinity of pipelines or overhead cables, or quite generally in areas where vigorous plant growth is unwanted.

Also important is the use of growth regulators for inhibition of the longitudinal growth of cereal. This reduces or completely eliminates the risk of lodging of the plants prior to harvest. In addition, growth regulators in the case of cereals can strengthen the culm, which also counteracts lodging. The employment of
25 growth regulators for shortening and strengthening culms allows the deployment of higher fertilizer volumes to increase the yield, without any risk of lodging of the cereal crop.

In many crop plants, inhibition of vegetative growth allows denser planting, and it is thus possible to achieve higher yields based on the soil surface. Another advantage of the smaller plants obtained in this way is that the crop is easier to cultivate and harvest.

- 30 Inhibition of the vegetative plant growth may also lead to enhanced yields because the nutrients and assimilates are of more benefit to flower and fruit formation than to the vegetative parts of the plants.

Frequently, growth regulators can also be used to promote vegetative growth. This is of great benefit when harvesting the vegetative plant parts. However, promoting vegetative growth may also promote generative growth in that more assimilates are formed, resulting in more or larger fruits.

5 In some cases, yield increases may be achieved by manipulating the metabolism of the plant, without any detectable changes in vegetative growth. In addition, growth regulators can be used to alter the composition of the plants, which in turn may result in an improvement in quality of the harvested products. For example, it is possible to increase the sugar content in sugar beet, sugar cane, pineapples and in citrus fruit, or to increase the protein content in soya or cereals. It is also possible, for example, to use growth regulators to inhibit the degradation of desirable ingredients, for example sugar in sugar beet or sugar cane, before or after harvest. It is also possible to positively influence the production or the elimination of secondary plant ingredients. One example is the stimulation of the flow of latex in rubber trees.

Under the influence of growth regulators, parthenocarpic fruits may be formed. In addition, it is possible to influence the sex of the flowers. It is also possible to produce sterile pollen, which is of great importance in the breeding and production of hybrid seed.

15 Use of growth regulators can control the branching of the plants. On the one hand, by breaking apical dominance, it is possible to promote the development of side shoots, which may be highly desirable particularly in the cultivation of ornamental plants, also in combination with an inhibition of growth. On the other hand, however, it is also possible to inhibit the growth of the side shoots. This effect is of particular interest, for example, in the cultivation of tobacco or in the cultivation of tomatoes.

20 Under the influence of growth regulators, the amount of leaves on the plants can be controlled such that defoliation of the plants is achieved at a desired time. Such defoliation plays a major role in the mechanical harvesting of cotton, but is also of interest for facilitating harvesting in other crops, for example in viticulture. Defoliation of the plants can also be undertaken to lower the transpiration of the plants before they are transplanted.

25 Growth regulators can likewise be used to regulate fruit dehiscence. On the one hand, it is possible to prevent premature fruit dehiscence. On the other hand, it is also possible to promote fruit dehiscence or even flower abortion to achieve a desired mass ("thinning"), in order to eliminate alternation. Alternation is understood to mean the characteristic of some fruit species, for endogenous reasons, to deliver very different yields from year to year. Finally, it is possible to use growth regulators at the time of harvest to reduce the forces required to detach the fruits, in order to allow mechanical harvesting or to facilitate manual harvesting.

35 Growth regulators can also be used to achieve faster or else delayed ripening of the harvested material before or after harvest. This is particularly advantageous as it allows optimal adjustment to the requirements of the market. Moreover, growth regulators in some cases can improve the fruit colour. In addition, growth regulators can also be used to concentrate maturation within a certain period of time. This estab-

lishes the prerequisites for complete mechanical or manual harvesting in a single operation, for example in the case of tobacco, tomatoes or coffee.

By using growth regulators, it is additionally possible to influence the resting of seed or buds of the plants, such that plants such as pineapple or ornamental plants in nurseries, for example, germinate, sprout or
5 flower at a time when they are normally not inclined to do so. In areas where there is a risk of frost, it may be desirable to delay budding or germination of seeds with the aid of growth regulators, in order to avoid damage resulting from late frosts.

Finally, growth regulators can induce resistance of the plants to frost, drought or high salinity of the soil. This allows the cultivation of plants in regions which are normally unsuitable for this purpose.

10 Resistance Induction / Plant Health and other effects

The active combinations according to the invention also exhibit a potent strengthening effect in plants. Accordingly, they can be used for mobilizing the defences of the plant against attack by undesirable microorganisms.

Plant-strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances which are capable of stimulating the defence system of plants in such a way that the
15 treated plants, when subsequently inoculated with undesirable microorganisms, develop a high degree of resistance to these microorganisms.

The active combinations according to the invention are also suitable for increasing the yield of crops. In addition, they show reduced toxicity and are well tolerated by plants.

20 Further, in context with the present invention plant physiology effects comprise the following:

Abiotic stress tolerance, comprising temperature tolerance, drought tolerance and recovery after drought stress, water use efficiency (correlating to reduced water consumption), flood tolerance, ozone stress and UV tolerance, tolerance towards chemicals like heavy metals, salts, pesticides (safener) etc..

Biotic stress tolerance, comprising increased fungal resistance and increased resistance against nematodes,
25 viruses and bacteria. In context with the present invention, biotic stress tolerance preferably comprises increased fungal resistance and increased resistance against nematodes

Increased plant vigor, comprising plant health / plant quality and seed vigor, reduced stand failure, improved appearance, increased recovery, improved greening effect and improved photosynthetic efficiency.

Effects on plant hormones and/or functional enzymes.

30 Effects on growth regulators (promoters), comprising earlier germination, better emergence, more developed root system and/or improved root growth, increased ability of tillering, more productive tillers, earli-

er flowering, increased plant height and/or biomass, shorting of stems, improvements in shoot growth, number of kernels/ear, number of ears/m², number of stolons and/or number of flowers, enhanced harvest index, bigger leaves, less dead basal leaves, improved phyllotaxy, earlier maturation / earlier fruit finish, homogenous riping, increased duration of grain filling, better fruit finish, bigger fruit/vegetable size, sprouting resistance and reduced lodging.

Increased yield, referring to total biomass per hectare, yield per hectare, kernel/fruit weight, seed size and/or hectolitre weight as well as to increased product quality, comprising:

improved processability relating to size distribution (kernel, fruit, etc.), homogenous riping, grain moisture, better milling, better vinification, better brewing, increased juice yield, harvestability, digestibility, sedimentation value, falling number, pod stability, storage stability, improved fiber length/strength/uniformity, increase of milk and/or meat quality of silage fed animals, adaption to cooking and frying;

further comprising improved marketability relating to improved fruit/grain quality, size distribution (kernel, fruit, etc.), increased storage / shelf-life, firmness / softness, taste (aroma, texture, etc.), grade (size, shape, number of berries, etc.), number of berries/fruits per bunch, crispness, freshness, coverage with wax, frequency of physiological disorders, colour, etc.;

further comprising increased desired ingredients such as e.g. protein content, fatty acids, oil content, oil quality, aminoacid composition, sugar content, acid content (pH), sugar/acid ratio (Brix), polyphenols, starch content, nutritional quality, gluten content/index, energy content, taste, etc.;

and further comprising decreased undesired ingredients such as e.g. less mycotoxines, less aflatoxines, geosmin level, phenolic aromas, lacchase, polyphenol oxidases and peroxidases, nitrate content etc.

Sustainable agriculture, comprising nutrient use efficiency, especially nitrogen (N)-use efficiency, phosphours (P)-use efficiency, water use efficiency, improved transpiration, respiration and/or CO₂ assimilation rate, better nodulation, improved Ca-metabolism etc..

Delayed senescence, comprising improvement of plant physiology which is manifested, for example, in a longer grain filling phase, leading to higher yield, a longer duration of green leaf colouration of the plant and thus comprising colour (greening), water content, dryness etc.. Accordingly, in the context of the present invention, it has been found that the specific inventive application of the active compound combination makes it possible to prolong the green leaf area duration, which delays the maturation (senescence) of the plant. The main advantage to the farmer is a longer grain filling phase leading to higher yield. There is also an advantage to the farmer on the basis of greater flexibility in the harvesting time.

Therein "sedimentation value" is a measure for protein quality and describes according to Zeleny (Zeleny value) the degree of sedimentation of flour suspended in a lactic acid solution during a standard time interval. This is taken as a measure of the baking quality. Swelling of the gluten fraction of flour in lactic acid

solution affects the rate of sedimentation of a flour suspension. Both a higher gluten content and a better gluten quality give rise to slower sedimentation and higher Zeleny test values. The sedimentation value of flour depends on the wheat protein composition and is mostly correlated to the protein content, the wheat hardness, and the volume of pan and hearth loaves. A stronger correlation between loaf volume and Zeleny sedimentation volume compared to SDS sedimentation volume could be due to the protein content influencing both the volume and Zeleny value (*Czech J. Food Sci. Vol. 21, No. 3: 91–96, 2000*).

Further the “falling number” as mentioned herein is a measure for the baking quality of cereals, especially of wheat. The falling number test indicates that sprout damage may have occurred. It means that changes to the physical properties of the starch portion of the wheat kernel has already happened. Therein, the falling number instrument analyzes viscosity by measuring the resistance of a flour and water paste to a falling plunger. The time (in seconds) for this to happen is known as the falling number. The falling number results are recorded as an index of enzyme activity in a wheat or flour sample and results are expressed in time as seconds. A high falling number (for example, above 300 seconds) indicates minimal enzyme activity and sound quality wheat or flour. A low falling number (for example, below 250 seconds) indicates substantial enzyme activity and sprout-damaged wheat or flour.

The term “more developed root system” / “improved root growth” refers to longer root system, deeper root growth, faster root growth, higher root dry/fresh weight, higher root volume, larger root surface area, bigger root diameter, higher root stability, more root branching, higher number of root hairs, and/or more root tips and can be measured by analyzing the root architecture with suitable methodologies and Image analysis programmes (e.g. WinRhizo).

The term “crop water use efficiency” refers technically to the mass of agriculture produce per unit water consumed and economically to the value of product(s) produced per unit water volume consumed and can e.g. be measured in terms of yield per ha, biomass of the plants, thousand-kernel mass, and the number of ears per m².

The term “nitrogen-use efficiency” refers technically to the mass of agriculture produce per unit nitrogen consumed and economically to the value of product(s) produced per unit nitrogen consumed, reflecting uptake and utilization efficiency.

Improvement in greening / improved colour and improved photosynthetic efficiency as well as the delay of senescence can be measured with well-known techniques such as a HandyPea system (Hansatech). Fv/Fm is a parameter widely used to indicate the maximum quantum efficiency of photosystem II (PSII). This parameter is widely considered to be a selective indication of plant photosynthetic performance with healthy samples typically achieving a maximum Fv/Fm value of approx. 0.85. Values lower than this will be observed if a sample has been exposed to some type of biotic or abiotic stress factor which has reduced the capacity for photochemical quenching of energy within PSII. Fv/Fm is presented as a ratio of variable fluorescence (Fv) over the maximum fluorescence value (Fm). The Performance Index is essentially an

indicator of sample vitality. (See e.g. *Advanced Techniques in Soil Microbiology*, **2007**, 11, 319-341; *Applied Soil Ecology*, **2000**, 15, 169-182.)

5 The improvement in greening / improved colour and improved photosynthetic efficiency as well as the delay of senescence can also be assessed by measurement of the net photosynthetic rate (Pn), measurement of the chlorophyll content, e.g. by the pigment extraction method of Ziegler and Ehle, measurement of the photochemical efficiency (Fv/Fm ratio), determination of shoot growth and final root and/or canopy biomass, determination of tiller density as well as of root mortality.

10 Within the context of the present invention preference is given to improving plant physiology effects which are selected from the group comprising: enhanced root growth / more developed root system, improved greening, improved water use efficiency (correlating to reduced water consumption), improved nutrient use efficiency, comprising especially improved nitrogen (N)-use efficiency, delayed senescence and enhanced yield.

15 Within the enhancement of yield preference is given as to an improvement in the sedimentation value and the falling number as well as to the improvement of the protein and sugar content – especially with plants selected from the group of cereals (preferably wheat).

20 Preferably the novel use of the fungicidal compositions of the present invention relates to a combined use of a) preventively and/or curatively controlling pathogenic fungi and/or nematodes, with or without resistance management, and b) at least one of enhanced root growth, improved greening, improved water use efficiency, delayed senescence and enhanced yield. From group b) enhancement of root system, water use efficiency and N-use efficiency is particularly preferred.

Seed Treatment

The invention further comprises a method for treating seed.

25 The invention further relates to seed which has been treated by one of the methods described in the previous paragraph. The inventive seeds are employed in methods for the protection of seed from unwanted microorganisms. In these methods, seed treated with at least one inventive active ingredient is used.

30 The inventive compositions are also suitable for treating seed. A large part of the damage to crop plants caused by harmful organisms is triggered by the infection of the seed during storage or after sowing, and also during and after germination of the plant. This phase is particularly critical since the roots and shoots of the growing plant are particularly sensitive, and even minor damage may result in the death of the plant. There is therefore a great interest in protecting the seed and the germinating plant by using appropriate compositions.

The control of phytopathogenic fungi by treating the seed of plants has been known for a long time and is the subject of constant improvements. However, the treatment of seed entails a series of problems which

cannot always be solved in a satisfactory manner. For instance, it is desirable to develop methods for protecting the seed and the germinating plant, which dispense with, or at least significantly reduce, the additional deployment of crop protection compositions after planting or after emergence of the plants. It is also desirable to optimize the amount of the active ingredient used so as to provide the best possible protection for the seed and the germinating plant from attack by phytopathogenic fungi, but without damaging the plant itself by the active ingredient employed. In particular, methods for the treatment of seed should also take account of the intrinsic fungicidal properties of transgenic plants in order to achieve optimal protection of the seed and the germinating plant with a minimum expenditure of crop protection compositions.

The present invention therefore also relates to a method for protection of seed and germinating plants from attack by phytopathogenic fungi, by treating the seed with an inventive composition. The invention likewise relates to the use of the inventive compositions for treatment of seed to protect the seed and the germinating plant from phytopathogenic fungi. The invention further relates to seed which has been treated with an inventive composition for protection from phytopathogenic fungi.

The control of phytopathogenic fungi which damage plants post-emergence is effected primarily by treating the soil and the above-ground parts of plants with crop protection compositions. Owing to the concerns regarding a possible influence of the crop protection compositions on the environment and the health of humans and animals, there are efforts to reduce the amount of active ingredients deployed.

One of the advantages of the present invention is that the particular systemic properties of the inventive active ingredients and compositions mean that treatment of the seed with these active ingredients and compositions not only protects the seed itself, but also the resulting plants after emergence, from phytopathogenic fungi. In this way, the immediate treatment of the crop at the time of sowing or shortly thereafter can be dispensed with.

It is likewise considered to be advantageous that the inventive compositions can especially also be used with transgenic seed, in which case the plant growing from this seed is capable of expressing a protein which acts against pests. By virtue of the treatment of such seed with the inventive active ingredients or compositions, merely the expression of the protein, for example an insecticidal protein, can control certain pests. Surprisingly, a further synergistic effect can be observed in this case, which additionally increases the effectiveness for protection against attack by pests.

The inventive compositions are suitable for protecting seed of any plant variety which is used in agriculture, in greenhouses, in forests or in horticulture and viticulture. In particular, this is the seed of cereals (such as wheat, barley, rye, triticale, sorghum/millet and oats), maize, cotton, soya beans, rice, potatoes, sunflower, bean, coffee, beet (for example sugar beet and fodder beet), peanut, oilseed rape, poppy, olive, coconut, cocoa, sugar cane, tobacco, vegetables (such as tomato, cucumbers, onions and lettuce), turf and ornamentals (see also below). The treatment of the seed of cereals (such as wheat, barley, rye, triticale and oats), maize and rice is of particular significance.

As also described below, the treatment of transgenic seed with the inventive active ingredients or compositions is of particular significance. This relates to the seed of plants containing at least one heterologous gene which enables the expression of a polypeptide or protein having insecticidal properties. The heterologous gene in transgenic seed can originate, for example, from microorganisms of the species *Bacillus*, *Rhizobium*, *Pseudomonas*, *Serratia*, *Trichoderma*, *Clavibacter*, *Glomus* or *Gliocladium*. This heterologous gene preferably originates from *Bacillus* sp., in which case the gene product is effective against the European maize borer and/or the Western maize rootworm. The heterologous gene more preferably originates from *Bacillus thuringiensis*.

In the context of the present invention, the inventive composition is applied to the seed alone or in a suitable formulation. Preferably, the seed is treated in a state in which it is sufficiently stable for no damage to occur in the course of treatment. In general, the seed can be treated at any time between harvest and sowing. It is customary to use seed which has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh of the fruits. For example, it is possible to use seed which has been harvested, cleaned and dried down to a moisture content of less than 15 % by weight. Alternatively, it is also possible to use seed which, after drying, for example, has been treated with water and then dried again.

When treating the seed, care must generally be taken that the amount of the inventive composition applied to the seed and/or the amount of further additives is selected such that the germination of the seed is not impaired, or that the resulting plant is not damaged. This has to be borne in mind in particular in the case of active ingredients which can have phytotoxic effects at certain application rates.

The inventive compositions can be applied directly, i.e. without containing any other components and without having been diluted. In general, it is preferable to apply the compositions to the seed in the form of a suitable formulation. Suitable formulations and methods for seed treatment are known to those skilled in the art and are described, for example, in the following documents: US 4,272,417, US 4,245,432, US 4,808,430, US 5,876,739, US 2003/0176428 A1, WO 2002/080675, WO 2002/028186.

The active ingredients usable in accordance with the invention can be converted to the customary seed dressing formulations, such as solutions, emulsions, suspensions, powders, foams, slurries or other coating compositions for seed, and also ULV formulations.

These formulations are prepared in a known manner, by mixing the active ingredients with customary additives, for example customary extenders and also solvents or diluents, dyes, wetting agents, dispersants, emulsifiers, antifoams, preservatives, secondary thickeners, adhesives, gibberellins and also water.

Useful dyes which may be present in the seed dressing formulations usable in accordance with the invention are all dyes which are customary for such purposes. It is possible to use either pigments, which are sparingly soluble in water, or dyes, which are soluble in water. Examples include the dyes known by the names Rhodamine B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

Useful wetting agents which may be present in the seed dressing formulations usable in accordance with the invention are all substances which promote wetting and which are conventionally used for the formulation of active agrochemical ingredients. Preference is given to using alkyl naphthalenesulphonates, such as diisopropyl or diisobutyl naphthalenesulphonates.

- 5 Useful dispersants and/or emulsifiers which may be present in the seed dressing formulations usable in accordance with the invention are all nonionic, anionic and cationic dispersants conventionally used for the formulation of active agrochemical ingredients. Usable with preference are nonionic or anionic dispersants or mixtures of nonionic or anionic dispersants. Suitable nonionic dispersants include especially ethylene oxide/propylene oxide block polymers, alkylphenol polyglycol ethers and tristyrylphenol polyglycol ether, and the phosphated or sulphated derivatives thereof. Suitable anionic dispersants are especially lignosulphonates, polyacrylic acid salts and arylsulphonate/formaldehyde condensates.
- 10

Antifoams which may be present in the seed dressing formulations usable in accordance with the invention are all foam-inhibiting substances conventionally used for the formulation of active agrochemical ingredients. Silicone antifoams and magnesium stearate can be used with preference.

- 15 Preservatives which may be present in the seed dressing formulations usable in accordance with the invention are all substances usable for such purposes in agrochemical compositions. Examples include dichlorophene and benzyl alcohol hemiformal.

- Secondary thickeners which may be present in the seed dressing formulations usable in accordance with the invention are all substances usable for such purposes in agrochemical compositions. Preferred examples include cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and finely divided silica.
- 20

Adhesives which may be present in the seed dressing formulations usable in accordance with the invention are all customary binders usable in seed dressing products. Preferred examples include polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose.

- 25 The gibberellins which may be present in the seed dressing formulations usable in accordance with the invention may preferably be gibberellins A1, A3 (= gibberellic acid), A4 and A7; particular preference is given to using gibberellic acid. The gibberellins are known (cf. R. Wegler "Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel" [Chemistry of the Crop Protection Compositions and Pesticides], vol. 2, Springer Verlag, 1970, p. 401-412).

- 30 The seed dressing formulations usable in accordance with the invention can be used, either directly or after previously having been diluted with water, for the treatment of a wide range of different seed, including the seed of transgenic plants. In this case, additional synergistic effects may also occur in interaction with the substances formed by expression.

For treatment of seed with the seed dressing formulations usable in accordance with the invention, or the preparations prepared therefrom by adding water, all mixing units usable customarily for the seed dressing are useful. Specifically, the procedure in the seed dressing is to place the seed into a mixer, to add the particular desired amount of seed dressing formulations, either as such or after prior dilution with water, and
5 to mix everything until the formulation is distributed homogeneously on the seed. If appropriate, this is followed by a drying process.

Mycotoxins

In addition, the inventive treatment can reduce the mycotoxin content in the harvested material and the foods and feeds prepared therefrom. Mycotoxins include particularly, but not exclusively, the following:
10 deoxynivalenol (DON), nivalenol, 15-Ac-DON, 3-Ac-DON, T2- and HT2-toxin, fumonisins, zearalenon, moniliformin, fusarin, diacetoxyscirpenol (DAS), beauvericin, enniatin, fusaroproliferin, fusarenol, ochratoxins, patulin, ergot alkaloids and aflatoxins which can be produced, for example, by the following fungi: *Fusarium* spec., such as *F. acuminatum*, *F. asiaticum*, *F. avenaceum*, *F. crookwellense*,
15 *F. culmorum*, *F. graminearum* (*Gibberella zeae*), *F. equiseti*, *F. fujikoro*i, *F. musarum*, *F. oxysporum*, *F. proliferatum*, *F. poae*, *F. pseudograminearum*, *F. sambucinum*, *F. scirpi*, *F. semitectum*, *F. solani*, *F. sporotrichoides*, *F. langsethiae*, *F. subglutinans*, *F. tricinctum*, *F. verticillioides* etc., and also by *Aspergillus* spec., such as *A. flavus*, *A. parasiticus*, *A. nomius*, *A. ochraceus*, *A. clavatus*, *A. terreus*, *A. versicolor*, *Penicillium* spec., such as *P. verrucosum*, *P. viridicatum*, *P. citrinum*, *P. expansum*, *P. claviforme*,
20 *P. roqueforti*, *Claviceps* spec., such as *C. purpurea*, *C. fusiformis*, *C. paspali*, *C. africana*, *Stachybotrys* spec. and others.

Material Protection

The inventive compositions can also be used in the protection of materials, for protection of industrial materials against attack and destruction by unwanted microorganisms, for example fungi and insects.

In addition, the inventive combinations can be used as antifouling compositions, alone or in combinations
25 with other active ingredients.

Industrial materials in the present context are understood to mean inanimate materials which have been prepared for use in industry. For example, industrial materials which are to be protected by inventive active ingredients from microbial alteration or destruction may be adhesives, glues, paper, wallpaper and board/cardboard, textiles, carpets, leather, wood, fibers and tissues, paints and plastic articles, cooling lubricants and other materials which can be infected with or destroyed by microorganisms. Parts of production plants and buildings, for example cooling-water circuits, cooling and heating systems and ventilation and air-conditioning units, which may be impaired by the proliferation of microorganisms may also be mentioned within the scope of the materials to be protected. Industrial materials within the scope of the present invention preferably include adhesives, sizes, paper and card, leather, wood, paints, cooling lubricants and heat transfer fluids, more preferably wood.
35

The inventive active combinations may prevent adverse effects, such as rotting, decay, discoloration, decoloration or formation of mould.

In the case of treatment of wood the compositions according to the invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting one or more compounds according to the invention or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

In addition, the inventive combinations can be used to protect objects which come into contact with salt-water or brackish water, especially hulls, screens, nets, buildings, moorings and signalling systems, from fouling.

The inventive method for controlling unwanted fungi can also be employed for protecting storage goods. Storage goods are understood to mean natural substances of vegetable or animal origin or processed products thereof which are of natural origin, and for which long-term protection is desired. Storage goods of vegetable origin, for example plants or plant parts, such as stems, leaves, tubers, seeds, fruits, grains, can be protected freshly harvested or after processing by (pre)drying, moistening, comminuting, grinding, pressing or roasting. Storage goods also include timber, both unprocessed, such as construction timber, electricity poles and barriers, or in the form of finished products, such as furniture. Storage goods of animal origin are, for example, hides, leather, furs and hairs. The inventive active ingredients may prevent adverse effects, such as rotting, decay, discoloration, decoloration or formation of mould.

Microorganisms capable of degrading or altering the industrial materials include, for example, bacteria, fungi, yeasts, algae and slime organisms. The inventive active ingredients preferably act against fungi, especially moulds, wood-discoloring and wood-destroying fungi (*Ascomycetes*, *Basidiomycetes*, *Deuteromycetes* and *Zygomycetes*), and against slime organisms and algae. Examples include microorganisms of the following genera: *Alternaria*, such as *Alternaria tenuis*; *Aspergillus*, such as *Aspergillus niger*; *Chaetomium*, such as *Chaetomium globosum*; *Coniophora*, such as *Coniophora puetana*; *Lentinus*, such as *Lentinus tigrinus*; *Penicillium*, such as *Penicillium glaucum*; *Polyporus*, such as *Polyporus versicolor*; *Aureobasidium*, such as *Aureobasidium pullulans*; *Sclerophoma*, such as *Sclerophoma pityophila*; *Trichoderma*, such as *Trichoderma viride*; *Ophiostoma* spp., *Ceratocystis* spp., *Humicola* spp., *Petriella* spp., *Trichurus* spp., *Coriolus* spp., *Gloeophyllum* spp., *Pleurotus* spp., *Poria* spp., *Serpula* spp. and *Tyromyces* spp., *Cladosporium* spp., *Paecilomyces* spp. *Mucor* spp., *Escherichia*, such as *Escherichia coli*; *Pseudomonas*, such as *Pseudomonas aeruginosa*; *Staphylococcus*, such as *Staphylococcus aureus*, *Candida* spp. and *Saccharomyces* spp., such as *Saccharomyces cerevisiae*.

Antimycotic Activity

In addition, the inventive combinations also have very good antimycotic activity. They have a very broad antimycotic activity spectrum, especially against dermatophytes and yeasts, moulds and diphasic fungi (for example against *Candida* species, such as *C. albicans*, *C. glabrata*), and *Epidermophyton floccosum*, *Aspergillus* species, such as *A. niger* and *A. fumigatus*, *Trichophyton* species, such as *T. mentagrophytes*, *Microsporon* species such as *M. canis* and *M. audouinii*. The list of these fungi by no means constitutes a restriction of the mycotic spectrum covered, and is merely of illustrative character.

The inventive combinations can therefore be used both in medical and in non-medical applications.

Application Rates and Timing

When using the inventive combinations as fungicides, the application rates can be varied within a relatively wide range, depending on the kind of application. The application rate of the inventive active ingredients is

- in the case of treatment of plant parts, for example leaves: from 0.1 to 10 000 g/ha, preferably from 10 to 1000 g/ha, more preferably from 10 to 800 g/ha, even more preferably from 50 to 300 g/ha (in the case of application by watering or dripping, it is even possible to reduce the application rate, especially when inert substrates such as rockwool or perlite are used);
- in the case of seed treatment: from 2 to 200 g per 100 kg of seed, preferably from 3 to 150 g per 100 kg of seed, more preferably from 2.5 to 25 g per 100 kg of seed, even more preferably from 2.5 to 12.5 g per 100 kg of seed;
- in the case of soil treatment: from 0.1 to 10 000 g/ha, preferably from 1 to 5000 g/ha.

These application rates are merely by way of example and are not limiting for the purposes of the invention.

The inventive compositions can thus be used to protect plants from attack by the pathogens mentioned for a certain period of time after treatment. The period for which protection is provided extends generally for 1 to 28 days, preferably for 1 to 14 days, more preferably for 1 to 10 days, most preferably for 1 to 7 days, after the treatment of the plants with the active ingredients, or for up to 200 days after a seed treatment.

The method of treatment according to the invention also provides the use or application of compounds (A) and (B) and/or (C) in a simultaneous, separate or sequential manner. If the single active ingredients are applied in a sequential manner, i.e. at different times, they are applied one after the other within a reasonably short period, such as a few hours or days. Preferably the order of applying the compounds (A) and (B) and/or (C) is not essential for working the present invention.

The plants listed can particularly advantageously be treated in accordance with the invention with the compounds of the general formula (I) and the inventive compositions. The preferred ranges stated above for the active ingredients or compositions also apply to the treatment of these plants. Particular emphasis is

given to the treatment of plants with the compounds or compositions specifically mentioned in the present text.

5

The advanced fungicidal activity of the active compound combinations according to the invention is evident from the example below. While the individual active compounds exhibit weaknesses with regard to the fungicidal activity, the combinations have an activity which exceeds a simple addition of activities.

10 A synergistic effect of fungicides is always present when the fungicidal activity of the active compound combinations exceeds the total of the activities of the active compounds when applied individually. The expected activity for a given combination of two active compounds can be calculated as follows (cf. Colby, S.R., "Calculating Synergistic and Antagonistic Responses of Herbicide Combinations", *Weeds* **1967**, 15, 20-22):

15 If

X is the efficacy when active compound A is applied at an application rate of m ppm (or g/ha),

Y is the efficacy when active compound B is applied at an application rate of n ppm (or g/ha),

E is the efficacy when the active compounds A and B are applied at application rates of m and n ppm (or g/ha), respectively, and

20 then

$$E = X + Y - \frac{X \cdot Y}{100}$$

The degree of efficacy, expressed in % is denoted. 0 % means an efficacy which corresponds to that of the control while an efficacy of 100 % means that no disease is observed.

25 If the actual fungicidal activity exceeds the calculated value, then the activity of the combination is super-additive, i.e. a synergistic effect exists. In this case, the efficacy which was actually observed must be greater than the value for the expected efficacy (E) calculated from the abovementioned formula.

A further way of demonstrating a synergistic effect is the method of Tammes (cf. "Isoboles, a graphic representation of synergism in pesticides" in *Neth. J. Plant Path.*, **1964**, 70, 73-80).

The invention is illustrated by the following example. However the invention is not limited to the example.

Example**Phytophthora test (tomatoes) / preventive**

Compounds used for the test are Chlorothalonil, Cymoxanil, Fosetyl-Al, Mandipropamid and Metalaxyl as active compounds (1 part by weight), solved in acetone/dimethylacetamide (24.5/24.5 part by weight) and
5 alkylaryl polyglycol ether (1 part by weight). Compound of formula (I) (Oxathiapiprolin) was used in a 50g a.i./l SC-Formulation as amorphous active compound ("AM") and as crystalline active compound ("CRYST") which corresponds to Form B in WO 2010/123791. Solved active compounds, suspended SC-formulations and combinations thereof were diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound or
10 compound combination at the stated rate of application. After the spray coating has dried on, the plants are inoculated with an aqueous spore suspension of *Phytophthora infestans*. The plants are then placed in an incubation cabinet at approximately 20 °C and a relative atmospheric humidity of 100%.

The test is evaluated 3 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

15 The table below clearly shows that the observed activity of the active compound combination according to the invention is greater than the calculated activity, i.e. a synergistic effect is present.

Table

Phytophthora test (tomatoes) / protective

Active compounds	Application rate of active compound in ppm a.i.	Efficacy in %		
		found*	calc.**	diff.***
("AM") Oxathiapiprolin amorphous	0.005	30		
("CRYST") Oxathiapiprolin crystalline, Form B (see WO 2010/123791)	0.005	30		
Chlorothalonil	10	0		
Cymoxanil	25	0		
Fosetyl-Al	25	8		
Mandipropamid	0.25	60		
Metalaxyl	5	0		
("AM") + Chlorothalonil 1:2000	0.005 + 10	40	30	10
("CRYST") + Chlorothalonil 1:2000	0.005 + 10	84	30	54
("AM") + Cymoxanil 1:5000	0.005 + 25	30	30	0
("CRYST") + Cymoxanil 1:5000	0.005 + 25	60	30	30
("AM") + Fosetyl-Al 1:5000	0.005 + 25	45	36	9
("CRYST") + Fosetyl-Al 1:5000	0.005 + 25	50	36	14
("AM") + Mandipropamid 1:50	0.005 + 0.25	58	72	-14
("CRYST") + Mandipropamid 1:50	0.005 + 0.25	83	72	11
("AM") + Metalaxyl 1:1000	0.005 + 5	35	30	5
("CRYST") + Metalaxyl 1:1000	0.005 + 5	50	30	20

* found = activity found


** calc. = activity calculated using Colby's formula

*** diff. = difference between activity found and activity calculated

Patent Claims

1. Synergistic combination comprising:

(A) Crystalline thiazolyloxazoline of formula (I)



(I)

5 in its Form B (see WO 2010/123791, page 4, line 5 to 10) characterized by powder X-ray diffraction pattern having at least the 2Θ reflection positions

2Θ
14.902
18.123
18.87
20.204
20.883
21.79
24.186
26.947

and

(B) at least one further active compound selected from the following groups

- (1) inhibitors of the ergosterol synthesis,
- (2) inhibitors of the respiratory chain at complex I or II,
- (3) inhibitors of the respiratory chain at complex III,
- (4) inhibitors of the mitosis and cell division,

- (5) compounds capable of having a multisite action,
 - (6) compounds capable of inducing a host defense,
 - (7) inhibitors of the amino acid and/or protein biosynthesis,
 - (8) inhibitors of the ATP production,
 - 5 (9) inhibitors of the cell wall synthesis,
 - (10) inhibitors of the lipid and membrane synthesis,
 - (11) inhibitors of the melanine biosynthesis,
 - (12) inhibitors of the nucleic acid synthesis,
 - (13) inhibitors of the signal transduction,
 - 10 (14) compounds capable of acting as uncoupler,
 - (15) other fungicides.
2. Method for controlling phytopathogenic harmful fungi, characterized in that mixtures according to Claim 1 are applied to the phytopathogenic harmful fungi and/or their habitat.
- 15 3. Composition for controlling phytopathogenic harmful fungi, characterized by a content of at least one mixture according to Claim 1, in addition to extenders and/or surfactants.
4. Composition according to Claim 1 comprising at least one further active ingredient selected from the group of the insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators, herbicides, fertilizers, safeners and semiochemicals.
- 20 5. Use of mixtures according to Claim 1 for control of phytopathogenic harmful fungi.
6. Use of mixtures according to Claim 1 as plant growth regulators.
- 25 7. Process for producing compositions for controlling phytopathogenic harmful fungi, characterized in that mixtures according to Claim 1 are mixed with extenders and/or surfactants.

8. Use of compounds of the mixtures according to Claim 1 for treatment of transgenic plants.
9. Use of compounds of mixtures according to Claim 1 for treatment of seed and of seed of transgenic plants.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2015/055317

A. CLASSIFICATION OF SUBJECT MATTER

INV. A01N43/80 A01N37/34 A01N47/34 A01N57/12 A01N37/36
A01P3/00

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/123791 A1 (DU PONT [US]; HANAGAN MARY ANN [US]; OBERHOLZER MATTHEW RICHARD [US];) 28 October 2010 (2010-10-28) cited in the application page 71, line 35 - page 72, line 6 table 7 claims 15,16	1-9
X	WO 2009/055514 A2 (DU PONT [US]; GREGORY VANN [US]; PASTERIS ROBERT JAMES [US]) 30 April 2009 (2009-04-30) cited in the application pag391: compound 214 tables M, N, O, P	1-9



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

4 May 2015

Date of mailing of the international search report

12/05/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Götz, Gerhard

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2015/055317

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010123791	A1	28-10-2010	AR 076384 A1 08-06-2011
		AU 2010239456 A1 13-10-2011	
		CA 2757055 A1 28-10-2010	
		CN 102459256 A 16-05-2012	
		CO 6400143 A2 15-03-2012	
		DK 2421861 T3 15-04-2013	
		EP 2421861 A1 29-02-2012	
		ES 2402113 T3 29-04-2013	
		JP 2012524784 A 18-10-2012	
		KR 20120007056 A 19-01-2012	
		MY 152264 A 15-09-2014	
		NZ 595278 A 22-02-2013	
		PE 08792012 A1 16-08-2012	
		PT 2421861 E 18-03-2013	
		RU 2011147184 A 27-05-2013	
		TW 201038568 A 01-11-2010	
		US 2012034315 A1 09-02-2012	
		US 2014127322 A1 08-05-2014	
		UY 32567 A 30-11-2010	
		WO 2010123791 A1 28-10-2010	
WO 2009055514	A2	30-04-2009	AR 069368 A1 20-01-2010
		AU 2008316880 A1 30-04-2009	
		CA 2699933 A1 30-04-2009	
		CN 101969781 A 09-02-2011	
		CO 6382166 A2 15-02-2012	
		EP 2187748 A2 26-05-2010	
		GE P20135758 B 25-02-2013	
		GT 201000114 A 16-04-2012	
		JP 5364712 B2 11-12-2013	
		JP 2011500826 A 06-01-2011	
		KR 20100083179 A 21-07-2010	
		MY 150537 A 30-01-2014	
		PE 03152013 A1 26-03-2013	
		PE 09742009 A1 13-08-2009	
		RU 2010120684 A 27-11-2011	
		TW 200917963 A 01-05-2009	
		US 2010240619 A1 23-09-2010	
		UY 31416 A1 29-05-2009	
		VN 24622 A1 27-12-2010	
		WO 2009055514 A2 30-04-2009	