

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2008343923 C1**

(54) Title
Use of tetramic acid derivatives for controlling nematodes

(51) International Patent Classification(s)
A01N 43/38 (2006.01) **A01P 5/00** (2006.01)
A01N 47/06 (2006.01)

(21) Application No: **2008343923** (22) Date of Filing: **2008.12.18**

(87) WIPO No: **WO9/085176**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/008,507	2007.12.20	US

(43) Publication Date: **2009.07.09**

(44) Accepted Journal Date: **2014.02.06**

(44) Amended Journal Date: **2014.04.10**

(71) Applicant(s)
Bayer Intellectual Property GmbH

(72) Inventor(s)
Perry, John L.;Jimenez, Manuel;Steffens, Robert;Fischer, Reiner;Krueger, Stephen;Bell, John W.

(74) Agent / Attorney
Davies Collison Cave, Level 14 255 Elizabeth Street, Sydney, NSW, 2000

(56) Related Art
CA 2 492 096 C
EP 0 076 580 B1
WO 2007/126691 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 July 2009 (09.07.2009)

PCT

(10) International Publication Number
WO 2009/085176 A3

(51) International Patent Classification:

A01N 43/38 (2006.01) A01P 5/00 (2006.01)
A01N 47/06 (2006.01)

(21) International Application Number:

PCT/US2008/013829

(22) International Filing Date:

18 December 2008 (18.12.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/008,507 20 December 2007 (20.12.2007) US

(71) Applicant (for all designated States except US): **BAYER CROPSCIENCE LP** [US/US]; Patent Department, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FISCHER, Reiner** [DE/DE]; Nelly-Sachs-Str. 23, Monheim Am Rhein (DE). **STEFFENS, Robert** [US/US]; 312 Hogan Valley Way, Cary, NC 27513 (US). **BELL, John, W.** [US/US]; 344 Wilkinson Creek Lane, Chapel Hill, NC 27516 (US). **JIMENEZ, Manuel** [US/US]; 323 Old Line Avenue, Exeter, CA 93322 (US). **KRUEGER, Stephen** [US/US]; 604 Compton Road, Raleigh, NC 27609 (US). **PERRY, John, L.** [US/US]; 2321 21st Avenue, Kingsburg, CA 93631 (US).

(74) Agent: **HARMUTH, Raymond, J.**; Bayer CropScience LP, Patent Department, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available):

AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

3 September 2009



WO 2009/085176 A3

(54) Title: USE OF TETRAMIC ACID DERIVATIVES FOR CONTROLLING NEMATODES

(57) Abstract: The present application relates to the use of tetramic acid derivatives for controlling soil-dwelling nematodes.

Use of tetramic acid derivatives for controlling nematodes

The present invention relates to the use of known tetramic acid derivatives for controlling soil-dwelling nematodes.

It is already known that certain cyclic ketoenols have herbicidal, insecticidal and acaricidal properties. The activity- of these substances is good; however, it is sometimes unsatisfactory at low application rates.

Known to have insecticidal and/or acaricidal action are 1H-3-aryl-pyrrolidine-2,4-dione derivatives (WO 98/05638) and their cis-isomers (WO 04/007448).


Also known are mixtures of compounds from WO 98/05638 with other insecticides and/or acaricides: WO 01/89300, WO 02/00025, WO 02/05648, WO 02/17715, WO 02/19824, WO 02/30199, WO 02/37963, WO 05/004603, WO 05/053405, WO 06/089665, DE-A-10342673, WO 2008/006516, WO 2008/006514, WO 2008/006513, WO 2008/006515, WO 2008/006512, WO 2008/009379. Also known are mixtures with nematicides: file number EP07112279.

Moreover, they are known to be active against plant bugs (family: Miridae) (WO 2007/131681)


Also, WO 2007/126691 discloses an activity against leaf nematodes following dip treatment of flower bulbs.

Summary of the Invention

Surprisingly, it has now been found that the compounds of the formulae (I-1) and (I-2)



(I-1)




(I-2)


known from WO 04/007448 reduce the population density of soil-dwelling plant-damaging nematodes in annual and perennial crops after foliar treatment.

2008343923 23 Dec 2013

In a first aspect, the present invention provides use of a compound of the formulae (I-1) and/or (I-2)




(I-1)




(I-2)

in a foliar treatment for controlling soil-dwelling plant-damaging nematodes in annual and perennial crops.

In a second aspect, the present invention provides a method for controlling soil nematodes in annual and perennial crops, the method comprising foliar treatment of the annual and perennial crops with a compound of the formulae (I-1) and/or (I-2)



(I-1)



(I-2)

Preference is given to using the compound of the formula (I-1) for controlling soil-dwelling plant-damaging nematodes in perennial crops.

Preference is given to using the compound of the formula (I-1) for controlling soil-dwelling plant-damaging nematodes in annual crops.

- 5 Preference is also given to using the compound of the formula (I-2) for controlling soil-dwelling plant-damaging nematodes in perennial crops.

Preference is also given to using the compound of the formula (I-2) for controlling soil-dwelling plant-damaging nematodes in annual crops.

- The phytoparasitic nematodes include, for example, *Pratylenchus* spp., *Radopholus similis*,
10 *Ditylenchus dipsaci*, *Tylenchulus semipenetrans*, *Heterodera* spp., *Globodera* spp., *Meloidogyne* spp.,
Aphelenchoides spp., *Longidorus* spp., *Xiphinema* spp., *Trichodorus* spp., *Bursaphelenchus* spp.,
Belonolaimus longicaudatus, *Mesocriconema xenoplax*, *Tylenchorhynchus* spp., *Rotylenchulus* spp.,
Helicotylenchus multicinctus, *Paratrichodorus* spp., *Paratylenchus* spp., *Criconemella* spp.,
Hoplolaimus spp., *Scutellonema* spp., and *Dolichodorus* spp..

- 15 Perennial crops are to be understood as meaning citrus fruit, pomme fruit, stone fruit, grapevine, tea, almonds, nuts, coffee, tropical fruit, soft fruit, ornamental plants, lawn and olives.

Annual crops are to be understood as meaning vegetable, tobacco, melons, beet, sugar beet, cereals, corn, cotton, soya and potatoes.

- Growth-regulating insecticides such as the compounds of the formulae (I-1) and (I-2) generally act
20 slowly and have no killing effect on adult animals. Owing to the slow onset of action and a short half-life in the soil, a nematode-controlling application against soil-dwelling nematodes was not expected to be feasible. It is highly surprising that the compounds of the formulae (I-1) and (I-2) are, after foliar application, suitable for controlling nematodes in spite of the slow onset of action.

- The crops to be protected, which have only been described in a general manner, are specified in a
25 differentiated and more in-depth manner below. Thus, with a view to application, citrus fruit is to be understood as meaning, for example, oranges, clementines, satsumas, lemons, grapefruits, cumquats, mandarines,

furthermore pomme fruit, such as apples, pears, but also stone fruit, such as peaches, nectarines, cherries, apricots,

furthermore grapevine, olives, tea, and tropical crops, such as, for example, mangoes, papayas, figs, pineapples, dates, bananas, durians, passion fruit, kakis, coconuts, cacao, coffee, avocados, lychees, maracujas, guavas, sugar cane,

5 moreover almonds and nuts, such as, for example, hazelnuts, walnuts, pistachios, cashew nuts, brazil nuts, pecan nuts, butter nuts, chestnuts, hickory nuts, macadamia nuts, peanuts,

additionally also soft fruit, such as, for example, blackcurrants, gooseberries, raspberries, blackberries, blueberries, strawberries, red bilberries, kiwis, cranberries.

10 With respect to the use, ornamental plants are to be understood as meaning, for example, cut flowers, such as, for example, roses, carnations, gerbera, lilies, marguerites, chrysanthemums, tulips, daffodils, anemones, poppies, amaryllis, dahlias, azaleas, malves, gardenias, euphobias,

furthermore, for example, bedding plants, potted plants and shrubs, such as, for example, roses, hibiscus, chrysanthemums,

15 furthermore, for example, bushes and conifers, such as, for example, fig trees, rhododendron, spruce trees, fir trees, pine trees, yew trees, juniper trees, but also lawn, such as, for example, golf lawn, garden lawn.

With respect to the use, vegetables are understood as meaning for example fruiting vegetables and inflorescences as vegetables, for example bell peppers, chillies, tomatoes, aubergines, cucumbers, pumpkins, courgettes, broad beans, climbing and dwarf beans, peas, artichokes, maize;

20 but also leafy vegetables, for example head-forming lettuce, chicory, endives, various types of cress, of rocket, lamb's lettuce, iceberg lettuce, leeks, spinach, Swiss chard;

furthermore tuber vegetables, root vegetables and stem vegetables, for example celeriac/celery, beetroot, carrots, radish, horseradish, scorzonera, asparagus, beet for human consumption, palm hearts, bamboo shoots, furthermore bulb vegetables, for example onions, leeks, Florence fennel, garlic;

25 furthermore Brassica vegetables such as cauliflower, broccoli, kohlrabi, red cabbage, white cabbage, curly kale, Savoy cabbage, Brussels sprouts, Chinese cabbage.

With respect to the use in cereal crops, cereal is to be understood as meaning, for example, wheat, barley, rye, oats, triticale but also maize and millet.

Particularly preferred are, from the family of the

Pratylenchidae,

- | | | |
|----|---------------------------|--|
| | Radopholus brevicaudatus | in citrus fruit, tropical fruit, e.g. bananas, coffee, |
| | Radopholus capitatus | coconuts, avocado; tea, ornamental plants, lawn |
| 5 | Radopholus cavenessi | |
| | Radopholus clarus | |
| | Radopholus citrophilus | |
| | Radopholus crenatus | |
| | Radopholus inaequalis | |
| 10 | Radopholus inanis | |
| | Radopholus intermedius | |
| | Radopholus laevis | |
| | Radopholus litoralis | |
| | Radopholus magniglans | |
| 15 | Radopholus megadorus | |
| | Radopholus nativus | |
| | Radopholus neosimilis | |
| | Radopholus nigeriensis | |
| | Radopholus rectus | |
| 20 | Radopholus rotundisemenus | |
| | Radopholus serratus | |
| | Radopholus similis | |
| | Radopholus trilineatus | |
| | Radopholus triversus | |
| 25 | Radopholus vacuus | |
| | Radopholus vangundyi | |
| | Radopholus vertexplanus | |
| | Radopholus williamsi | |
| | Pratylenchus coffeae | in tropical fruit, e.g. bananas, coffee, pineapples, |
| 30 | Pratylenchus brachyurus | nuts, e.g. walnuts, ornamental plants, e.g. roses; |
| | Pratylenchus fallax | strawberries |
| | Pratylenchus goodeyi | |
| | Pratylenchus vulnus | |

Pratylenchus penetrans

Particular preference is furthermore given to

5 *Xiphinema americanum* in crops such as grapevines, soft fruit, e.g. strawberries, conifers, e.g. pines, ornamental plants, e.g. roses, stone fruit

Xiphinema diversicaudatum

Xiphinema index

Longidorus elongatus in crops such as soft fruit, e.g. strawberries, shrubs, perennial crops

Meloidogyne incognita in crops such as grapevines, peanuts, sugar cane, tomatoes

10 *Meloidogyne hapla*

Meloidogyne arenaria

Meloidogyne javanica

15 *Tylenchulus semipenetrans* in crops such as citrus fruit, e.g. oranges, grapefruits, (Family: Tylenchulidae) lemons, mandarines, grapevines, olives, tropical fruit, e.g. persimmon

Belonolaimus longicaudatus in crops such as citrus fruit, e.g. oranges, grapefruits, (Family: Belonolaimidae) lemons, mandarines, soft fruit, e.g. strawberries, lawn, conifers, e.g. spruce

20 *Mesocriconema xenoplax* in crops such as grapevines, nuts, e.g. almonds, walnuts
Rotylenchulus reniformis in crops such as tropical fruit, e.g. bananas, pineapples, papayas, melons, passion fruit, coffee, in citrus fruit, e.g. oranges, grapefruits, ornamental plants, e.g. gardenias, euphorbias

Helicotylenchus multicinctus in crops such as tropical fruit, e.g. bananas

25 All plants and plant parts can be treated in accordance with the invention. In this context, plants are understood as meaning all plants and plant populations such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants can be plants which can be obtained by traditional breeding and optimization methods or by biotechnological and recombinant

methods, or combinations of these methods, including the transgenic plants and including the plant varieties which are capable or not capable of being protected by Plant Breeders' Rights. Plant parts are understood as meaning all aerial and subterranean parts and organs of the plants such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stalks, stems, flowers, fruit bodies, fruits and seeds, but also roots, tubers and rhizomes. The plant parts also include crop material and vegetative and generative propagation material, for example cuttings, tubers, rhizomes, slips and seeds.

The treatment according to the invention with the active compound, of the plants and plant parts, is effected directly or by treating their environment, habitat or store using conventional treatment methods, for example by dipping, spraying, fumigating, fogging, scattering, brushing on, injecting, and, in the case of propagation material, in particular seeds, furthermore by coating with one or more coats.

As already mentioned above, all plants and their parts can be treated in accordance with the invention. In a preferred embodiment, plant species and plant varieties which are found in the wild or which are obtained by traditional biological breeding methods, such as hybridization or protoplast fusion, and parts of these species and varieties are treated. In a further preferred embodiment, transgenic plants and plant varieties which were obtained by recombinant methods, if appropriate in combination with traditional methods (genetically modified organisms) and their parts are treated. The terms "parts", "parts of plants" or "plant parts" were described above.

Plants which are especially preferably treated in accordance with the invention are those of the varieties which are in each case commercially available or in use. Plant varieties are understood as meaning plants with novel traits which have been bred both by conventional breeding, by mutagenesis or by recombinant DNA techniques. They may take the form of varieties, biotypes or genotypes.

Depending on the plant species or plant varieties, their location and growth conditions (soils, climate, vegetation period, nutrition), superadditive ("synergistic") effects may also occur as a result of the treatment according to the invention. Effects which exceed the effects actually to be expected are, for example, reduced application rates and/or widened activity spectrum and/or an enhancement of the activity of the substances and compositions which can be used in accordance with the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salinity, increased flowering performance, facilitated harvest, speedier maturation, higher yields, higher quality and/or higher nutritional value of the crop products, better storability and/or processibility of the crop products.

The preferred transgenic plants or plant varieties (plants or plant varieties obtained by means of genetic engineering) which are to be treated in accordance with the invention include all plants which, by means of the recombinant modification, have received genetic material which confers particularly advantageous valuable traits to these plants. Examples of such traits are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salinity, increased flowering performance, facilitated harvest, speedier maturation, higher yields, higher quality and/or higher nutritional value of the crop products, better storability and/or processibility of the crop products. Other examples of such traits which are particularly emphasized are an improved defence of the plants against animal and microbial pests such as insects, mites, phytopathogenic fungi, bacteria and/or viruses, and an increased tolerance of the plants to specific herbicidal active compounds. Examples of transgenic plants which are mentioned are the important crop plants such as cereals (wheat, rice), maize, soybean, potato, cotton, tobacco, oilseed rape and fruit plants (with the fruits apples, pears, citrus fruits and grapes), with particular emphasis on maize, soybean, potatoes, cotton, tobacco and oilseed rape. Traits which are particularly emphasized are the increased defence of the plants against insects, arachnids, nematodes and slugs and snails as the result of toxins formed in the plants, in particular toxins which are produced in the plants by the genetic material of *Bacillus Thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and their combinations) (hereinbelow "Bt plants"). Traits which are also particularly emphasized are the increased defence of plants against fungi, bacteria and viruses by systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and resistance genes and correspondingly expressed proteins and toxins. Traits which are furthermore especially emphasized are the increased tolerance of the plants to specific herbicidal active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinothricin (for example "PAT" gene). The specific genes which confer the desired traits can also occur in combinations with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are maize varieties, cotton varieties, soybean varieties and potato varieties sold under the trade names YIELD GARD® (for example maize, cotton, soybean), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucoatn® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soybean varieties which are sold under the trade names Roundup Ready® (glyphosate tolerance, for example maize, cotton, soybean), Liberty Link® (phosphinothricin tolerance, for example oilseed rape), IMI® (imidazolinone tolerance) and STS® (sulphonylurea tolerance, for example maize). Herbicide-resistant plants (bred conventionally for herbicide tolerance) which may also be mentioned are the varieties sold under the name Clearfield® (for example maize). Naturally, what has been said also applies to plant varieties which will be developed, or marketed, in the future and which have these genetic traits or traits to be developed in the future.

The active compounds can be converted into the customary formulations, such as solutions, emulsions, wettable powders, suspensions, powders, dusts, pastes, soluble powders, granules, suspoemulsion concentrates, natural and synthetic materials impregnated with active compound, and ultrafine encapsulations in polymeric materials.

- 5 These formulations are produced in the known manner, for example by mixing the active compound with extenders, that is, liquid solvents and/or solid carriers, optionally with the use of surfactants, that is, emulsifiers and/or dispersants and/or foam formers.

Suitable extenders are, for example, water, polar and unpolar organic chemical liquids, for example from the classes of the aromatic and nonaromatic hydrocarbons (such as paraffins, alkylbenzenes, 10 alkylnaphthalenes, chlorobenzenes), of the alcohols and polyols (which can optionally also be substituted, etherified and/or esterified), of the ketones (such as acetone, cyclohexanone), esters (including fats and oils) and (poly)ethers, of the unsubstituted and substituted amines, amides, lactams (such as N-alkylpyrrolidones) and lactones, the sulphones and sulphoxides (such as dimethyl sulphoxide).

- 15 In the case of the use of water as an extender, organic solvents can, for example, also be used as cosolvents. Liquid solvents which are suitable are mainly: aromatics, such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons, such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons, such as cyclohexane or paraffins, for example mineral oil fractions, mineral oils and vegetable oils, alcohols, such as 20 butanol or glycol as well as their ethers and esters, ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents, such as dimethylformamide and dimethyl sulphoxide, and water.

Solid carriers which are suitable are:

- for example ammonium salts and ground natural minerals, such as kaolins, clays, talc, chalk, quartz, 25 attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as highly-disperse silica, alumina and silicates; suitable solid carriers for granules are: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, and synthetic granules of inorganic and organic meals, and granules of organic material such as sawdust, coconut shells, maize cobs and tobacco stalks; suitable emulsifiers and/or foam formers are: for example non- 30 ionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates as well as protein hydrolysates; suitable dispersants are: for example lignin-sulphite waste liquors and methylcellulose.

Adhesives such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, and natural phospholipids, such as cephalins and lecithins, and synthetic phospholipids, can be used in the formulations. Other additives can be mineral and vegetable oils.

- 5 It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs, such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

- 10 The formulations in general comprise between 0.1 and 95% by weight of active compound, preferably between 0.5 and 90%, and additionally preferably extenders and/or surfactants.

The active compound content of the use forms prepared from the commercially available formulations can vary within wide ranges. The active compound concentration of the use forms can be in the range of from 0.0000001 up to 95% by weight of active compound, preferably between 0.0001 and 1% by weight.

- 15 Application is in a customary manner which is appropriate for the use forms.

Use examples**Example 1**

In 5 replications, in each case 15 vines of the cultivar „Thompson Seedless 1948“ are treated against soil-dwelling nematodes using a fogger. Here, a tank mix of the active compound (I-2) (SC 240) is applied after foliar application at the stated application rates with 0.25% of the adjuvant Acitvator 90, in comparison to imidacloprid (SC 550) after drench application by micro-irrigation injection. The water application rate is 75 gallons per acre = 605 plants. Two applications of active compound (I-2) are carried out at an interval of 32 days.

Evaluation is carried out prior to the treatment and 30 days and 60 days after the first treatment by counting the number of nematodes in 500 g of soil.

a)

Active compounds	Application rate per 605 plants	Activity against <i>Meloidogyne incognita</i>		
		0 d	30 d	60 d
Imidacloprid	14 fluid ounces	267	171	188
Active compound (I-2)	12.5 fluid ounces	312	140.8	178.8
Control	0	342	336	308

b)

Active compounds	Application rate per 605 plants	Activity against <i>Mesocriconema xenoplax</i>		
		0 d	30 d	60 d
Imidacloprid	14 fluid ounces	1863	808.4	958.4
Active compound (I-2)	12.5 fluid ounces	1817	861.2	942.8
Control	0	1630	1397.6	2083.2

c)

Active compounds	Application rate per 605 plants	Activity against <i>Pratylenchus</i> sp.		
		0 d	30 d	60 d
Imidacloprid	14 fluid ounces	406	320	319.6
Active compound (I-2)	12.5 fluid ounces	826	371.2	454
Control	0	848	540	646

Example 2

Foliar Application of (I-2) to Citrus in Pots for Control of *Tylenchulus semipenetrans*

Treatments: 15 trees sprayed with (I-2 = Movento), 15 trees sprayed with water (presample each pot); 25 mL/tree

5 *Protocol:*

Add *Tylenchulus semipenetrans* larvae to all pots. The following day take trees out of greenhouse. Cover top of pot to exclude spray from soil. (I-2) (SC 240) is sprayed in a concentration of 2.6 ml plus 2.5 ml adjuvant MSO (methylated seed oil), (100% active w/v, added to the spray solution at 0.25% v/v) per liter sprac solution. Spray canopy from two sides of trees to runoff with a 2 gallon
 10 “Sure Spray” pump sprayer (Chapin Mfg.) using a nozzle that delivers a cone shape spray pattern. Allow material to dry before returning plants to greenhouse. Repeat in 14 days.

For this citrus nematode trial, 5-year old Volkamer lemon seedlings are infested with *T. semipenetrans* obtained from the field and maintained in the greenhouse (25-32 °C) for use in pesticide trials. Trees used in the experiment are not treated with nematicidal compounds. The
 15 seedlings are growing in a 50:50 mixture of Astatula sand (97% sand) and Pro-mix potting mixture in rectangular pots (10x10x30 cm). Trees are irrigated normally, but receive no fertilizer or pesticide treatments other than (I-2) during the experiment. Effects on citrus nematode are evaluated by soil sampling 30 days and 60 days after treatment.

ANCOVA: pf versus treat (First nematode evaluation)

20

Factor	Levels	Values
treat	2	0 1

Analysis of Covariance for pf

25

Source	DF	Adj SS	MS	F	P
Covariates	1	12890661	12890661	15.12	0.001
treat	1	5665625	5665625	6.64	0.016
Error	27	23026066	852817		
Total	29	41567938			

30

Covariate	Coef	SE Coef	T	P
pi	0.2656	0.0683	3.888	0.001

35

ANCOVA: ln pf versus treat

Factor	Levels	Values
treat	2	0 1

5 Analysis of Covariance for ln pf

Source	DF	Adj SS	MS	F	P
Covariates	1	46.987	46.987	69.98	0.000
treat	1	5.563	5.563	8.29	0.008
Error	27	18.129	0.671		
Total	29	68.114			

Covariate	Coef	SE Coef	T	P
ln pi	0.8075	0.0965	8.365	0.000

15

ANCOVA: pf2 versus treat (Second nematode evaluation)

Factor	Levels	Values
treat	2	0 1

20 Analysis of Covariance for pf2

Source	DF	Adj SS	MS	F	P
Covariates	1	37890104	37890104	12.06	0.002
treat	1	5907691	5907691	1.88	0.182
Error	27	84807883	3141033		
Total	29	128580451			

Covariate	Coef	SE Coef	T	P
pi	0.4553	0.131	3.473	0.002

30

ANCOVA: ln pf2 versus treat


Factor	Levels	Values
treat	2	0 1

35 Analysis of Covariance for ln pf2

Source	DF	Adj SS	MS	F	P
Covariates	1	32.587	32.587	32.61	0.000
treat	1	2.982	2.982	2.98	0.096
Error	27	26.980	0.999		
Total	29	61.015			

Covariate	Coef	SE Coef	T	P
ln pi	0.6724	0.118	5.711	0.000

45



(Movento = I-2)

Tylenchulus semipenetrans raw data (juvenile and male nematodes per 100 cm² soil. Pi is initial population before treatment, Pf is final population at one and two months post treatment).

Treat Pi	Treat Pf	Treat Pf2	No treat Pi	No treat Pf	No treat Pf2
1625.0	638.16	4122.22	1387.5	3272.73	660.71
125.0	87.21	175.44	2237.5	733.77	6031.91
2190.0	1803.37	1409.84	2980.0	1850.57	4348.21
4250.0	1402.78	3205.36	7650.0	2308.22	5977.78
210.0	60.81	92.59	400.0	43.48	44.44
2490.0	1448.86	1481.13	100.0	269.23	428.57
250.0	86.21	551.02	530.0	1776.12	3084.91
125.0	54.35	531.25	7000.0	1664.63	4812.50
2412.5	168.83	181.82	787.5	1306.67	2336.73
2100.0	519.74	380.95	3760.0	3080.25	2314.81
1700.0	534.25	338.71	6675.0	4981.71	3564.81
9237.5	1440.00	1207.55	50.0	58.82	106.38
4180.0	2753.33	7807.69	1050.0	1954.55	1442.31
400.0	336.84	347.46	25.0	23.81	90.91
4250.0	708.33	1908.33	850.0	1739.13	1780.70

Example 3

The experiment is carried out in a shade house at a temperature of around 78 F (76-80 F). The experiment is conducted in a complete randomize design with seven treatments and six replications. 6 inch pots are filled with 1,500 g of steam pasteurized (194 F for 2:30 h) field soil per pot and a 4 weeks old tomato cv. Rutgers transplant is added. The soil is an Arredondo fine sand with ca. 92% sand and 1% om. The soil is taken from a site with no history of root-knot nematode infestation.

The first preventive treatments of Vydate CLV and (I-2) (SC 240; adjuvant: Dyne-Amic (a blend of nonionic and organosilicone surfactants and MSO), 99% active w/v, added 0.25% v/v) are applied as foliar spray. Fosthiazate 900 is applied as a soil drench in 40 gallons water per acre. The second preventive application of Vydate and (I-2) are sprayed over foliage 7 days after 1st treatment. The soil is inoculated by adding 5,000 eggs/pot of *Meloidogyne arenaria* one day after the 2nd treatment. The curative treatments of Vydate and (I-2) are applied as a foliar spray on 21 and 28 days after inoculation with *Meloidogyne arenaria*.

All (I-2) and Vydate treatments are applied in 40 gallons of water/acre by a CO₂ pressurized sprayer with a single 8003 VS TeeJet nozzle. The tomato plants for each treatment are lined up in a row with the spray directed over their foliage until noticeable drip.

78 days after the 1st treatment the plants are harvested. Each root system is evaluated for the percentage of root-knot nematode galling and eggs per root system. A 1 to 6 scale is used for estimating the number galls per root system as follows: 1 = no galls on the root system; 2 = 10%; 3 = 11 to 30%; 4 = 31 to 70%; 5 = 71 to 90%; 6= 100% root system galled. Following the determination of percentage of galling the roots are processed in a bleach solution to dissolve away the gelatinous matrix surrounding the eggs. Individual eggs are collection on a 500 mesh sieve and poured into a counting dish for determining their numbers.

RESULTS

The nontreated uninoculated control plants have no galling and the roots are white and healthy in appearance (Table 1). Nontreated inoculated plants have darkened, rotting roots that showed visual signs of decay. Galls and egg masses are numerous and there are many large coalesced galls.

- 5 Plants treated with fosthiazate, except for one plant out of six, have darkened decaying roots with numerous galls and egg masses. Large portions of the root system have coalesced galls. Their general appearance is similar to that of the nontreated inoculated plants. The root system of plants treated with (I-2) via a foliar spray as a preventive treatment have numerous egg masses and coalesced galls; roots are dark in appearance but the number of egg masses appear less than that in the
- 10 nontreated inoculated control plants. Roots of plants treated with Vydate foliar preventative treatment have numerous coalesced galls, but roots are whiter in appearance and rotting is much less apparent. There are less egg masses.

- 15 Roots of plants treated with (I-2) as a curative treatment have little darkening or rotting appearance. They have more fibrous roots and overall their appear to be a larger root system. Three of the six plants have little gall coalescing and there is an obvious appearance of fewer egg masses. Plants treated with Vydate as a curative treatment have more coalesced galls than the (I-2) treated plants. There is little appearance of rot and there are fewer egg masses.

Because there appeared to be fewer egg masses on tomato roots of some treatments the decision is made to count individual eggs to make a more quantitative determination among the treatments.

Table 1. Average root-knot nematode galling indices and number of eggs per tomato root system following treatments of fosthiazate, (I-2), and Vydate.

Treatment ¹	Rate	Galling index ¹	Number of eggs/root system 78 d after first treatment
1. Untreated (without nematode)		0.0	0
2. Untreated (with nematode)		6.0	727,820
3. Fosthiazate – preventive	6.0 lb ai/A	5.3	453,500
4. Movento (I-2) – preventive	0.0785 lb ai/A	6.0	421,600
5. Vydate – preventive	4.0 lb ai/A	5.2	333,500
6. Movento (I-2) – curative	0.0785 lb ai/A	4.5	141,000
7. Vydate – curative	4.0 lb ai/A	4.3	60,250

5

17

10

15

¹Subjective root-knot nematode gall rating scale 1 to 6 where 1 = no galls on root system;

2 = 10%; 3 = 11 to 30%; 4 = 31 to 70%; 5 = 71 to 90%; 6 = 100% of root system galled.

Data are means of four replications. Means within a column followed by a common letter are not different according to Duncan’s multiple-range test (P. 0.05).

2008343923 23 Dec 2013


Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.


2008343923 23 Dec 2013

The claims defining the invention are as follows:

1. Use of a compound of the formulae (I-1) and/or (I-2)



(I-1)




(I-2)


in a foliar treatment for controlling soil-dwelling plant-damaging nematodes in annual and perennial crops.

2. A use according to claim 1 for controlling soil dwelling plant-damaging nematodes in grapevines, wherein the compound is of formula (I-1).
3. A use according to claim 1 for controlling soil dwelling plant-damaging nematodes in vegetables, wherein the compound is of formula (I-1).
4. A use according to claim 1 for controlling soil dwelling plant-damaging nematodes in citrus, wherein the compound is of formula (I-1).
5. A use according to claim 1 for controlling soil dwelling plant-damaging nematodes in grapevines, wherein the compound is of formula (I-2).
6. A use according to claim 1 for controlling soil dwelling plant-damaging nematodes in vegetables, wherein the compound is of formula (I-2).
7. A use according to claim 1 for controlling soil dwelling plant-damaging nematodes in citrus, wherein the compound is of formula (I-2).
8. A method for controlling soil nematodes in annual and perennial crops, the method comprising foliar treatment of the annual and perennial crops with a compound of the formulae (I-1) and/or (I-2)

2008343923 23 Dec 2013



(I-1)



(I-2)

9. A method according to claim 7, wherein the annual and perennial crops are grapevines and the compound is of formula (I-1).
10. A method according to claim 7 for controlling soil dwelling plant-damaging nematodes in vegetables, wherein the compound is of formula (I-1).
11. A method according to claim 7 for controlling soil dwelling plant-damaging nematodes in citrus, wherein the compound is of formula (I-1).
12. A method according to claim 7 for controlling soil dwelling plant-damaging nematodes in grapevines, wherein the compound is of formula (I-2).
13. A method according to claim 7 for controlling soil dwelling plant-damaging nematodes in vegetables, wherein the compound is of formula (I-2).
14. A method according to claim 7 for controlling soil dwelling plant-damaging nematodes in citrus, wherein the compound is of formula (I-2).
15. Use according to claim 1, or a method according to claim 8, substantially as herein described with reference to the examples..